

$B_J(5970)^+$

 $I(J^P) = \frac{1}{2}(?)$
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_J(5970)^+$ MASS

 OUR FIT uses m_{B^0} and $m_{B_J(5970)^+} - m_{B^0}$ to determine $m_{B_J(5970)^+}$.

VALUE (MeV)

DOCUMENT ID

5964 ± 5 OUR FIT

$m_{B_J(5970)^+} - m_{B^0}$

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

685 ± 5 OUR FIT
685 ± 5 OUR AVERAGE

685.3 ± 4.1 ± 2.5

2K

¹ AAIJ

15AB LHCb

 pp at 7, 8 TeV

681 ± 5 ± 12

1.4k

² AALTONEN

14l CDF

 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.8 ± 4.5 ± 2.5

2K

³ AAIJ

15AB LHCb

 pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 545.8 \pm 4.1 \pm 2.5$ MeV which we adjust by

 the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AALTONEN 14l reports $m_{B_J(5970)^+} - m_{B^0} - m_{\pi^+} = 541 \pm 5 \pm 12$ MeV which we adjusted by the π^+ mass.

³ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 547 \pm 5 \pm 3$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$m_{B_J(5970)^+} - m_{B^{*0}}$

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.0 ± 4.0 ± 2.5

2k

¹ AAIJ

15AB LHCb

 pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 547 \pm 4 \pm 3$ MeV which

 we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5970)^+$ WIDTH

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

62 ± 20 OUR AVERAGE

63 ± 15 ± 17

2K

¹ AAIJ

15AB LHCb

 pp at 7, 8 TeV

 60⁺³⁰₋₂₀ ± 40

1.4k

AALTONEN

14l CDF

 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

61 ± 14 ± 17

2K

² AAIJ

15AB LHCb

 pp at 7, 8 TeV

61 ± 15 ± 17

2K

³ AAIJ

15AB LHCb

 pp at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5970)^+$ DECAY MODES

NODE=M220215;NODE=M220

Mode		Fraction (Γ_i/Γ)
Γ_1	$B^0 \pi^+$	possibly seen
Γ_2	$B^{*0} \pi^+$	seen

DESIG=1
DESIG=2

$B_J(5970)^+$ BRANCHING RATIOS

NODE=M220220

$\Gamma(B^0 \pi^+)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
possibly seen	2K	¹ AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV	
possibly seen	1.4k	AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV	

NODE=M220R01
NODE=M220R01

¹ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.

NODE=M220R01;LINKAGE=A

$\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	2k	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV	
seen	1.4k	AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV	

NODE=M220R02
NODE=M220R02

$B_J(5970)^+$ REFERENCES

NODE=M220

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14l PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)

REFID=56628
REFID=56029