

$B_J(5840)^+$

$$I(J^P) = \frac{1}{2}(?)$$

I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M224

NODE=M224

 $B_J(5840)^+$ MASSOUR FIT uses m_{B^0} and $m_{B_J(5840)^+} - m_{B^0}$ to determine $m_{B_J(5840)^+}$.

NODE=M224M

NODE=M224M

NODE=M224M

VALUE (MeV)

DOCUMENT ID

5851 ± 19 OUR FIT **$m_{B_J(5840)^+} - m_{B^0}$**

NODE=M224DM

NODE=M224DM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

571 ± 19 OUR FIT**571 ± 13 ± 14**

7k

1 AAIJ

15AB LHCB

 pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

595 ± 26 ± 14

7k

2 AAIJ

15AB LHCB

 pp at 7, 8 TeV

OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$ MeV which we adjust by

the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$ MeV which we adjust by

the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224DM;LINKAGE=A

NODE=M224DM;LINKAGE=B

 $m_{B_J(5840)^+} - m_{B^{*0}}$

NODE=M224DM2

NODE=M224DM2

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

565 ± 15 ± 14

7k

1 AAIJ

15AB LHCB

 pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$

MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224DM2;LINKAGE=A

 $B_J(5840)^+$ WIDTH

NODE=M224W

NODE=M224W

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

224 ± 24 ± 80

7k

1 AAIJ

15AB LHCB

 pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

215 ± 27 ± 80

7k

2 AAIJ

15AB LHCB

 pp at 7, 8 TeV

OCCUR=2

229 ± 27 ± 80

7k

3 AAIJ

15AB LHCB

 pp at 7, 8 TeV

OCCUR=3

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224W;LINKAGE=A

NODE=M224W;LINKAGE=B

NODE=M224W;LINKAGE=C

 $B_J(5840)^+$ DECAY MODES

NODE=M224215;NODE=M224

	Mode	Fraction (Γ_i/Γ)
Γ_1	$B^{*0} \pi^+$	seen
Γ_2	$B^0 \pi^+$	possibly seen

DESIG=1

DESIG=2

$B_J(5840)^+$ BRANCHING RATIOS

$\Gamma(B^{*0}\pi^+)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
seen	7k	AAIJ	15AB LHCB	pp at 7, 8 TeV	

$\Gamma(B^0\pi^+)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
possibly seen	7k	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV	

¹ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.

$B_J(5840)^+$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------

NODE=M224220

NODE=M224R01
NODE=M224R01

NODE=M224R02
NODE=M224R02

NODE=M224R02;LINKAGE=A

NODE=M224

REFID=56628