

$B_J(5840)^0$

$$I(J^P) = \frac{1}{2}(?)$$

I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

 $B_J(5840)^0$ MASSOUR FIT uses m_{B^+} and $m_{B_J(5840)^0} - m_{B^+}$ to determine $m_{B_J(5840)^0}$.

VALUE (MeV)

DOCUMENT ID

5863±9 OUR FIT **$m_{B_J(5840)^0} - m_{B^+}$**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

584± 9 OUR FIT**584± 5±7**

12k

¹ AAIJ

15AB LHCB

 pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

610±22±7

12k

² AAIJ

15AB LHCB

 pp at 7, 8 TeV¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust bythe π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.² AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$ MeV which we adjust bythe π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$m_{B_J(5840)^0} - m_{B^{*+}}$**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

584±5±7

12k

¹ AAIJ

15AB LHCB

 pp at 7, 8 TeV¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$ MeVwhich we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$B_J(5840)^0$ WIDTH**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

127±17±34

12k

¹ AAIJ

15AB LHCB

 pp at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

107±20±34

12k

² AAIJ

15AB LHCB

 pp at 7, 8 TeV

119±17±34

12k

³ AAIJ

15AB LHCB

 pp at 7, 8 TeV¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference. **$B_J(5840)^0$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*+} \pi^-$	seen
Γ_2 $B^+ \pi^-$	possibly seen

NODE=M225

NODE=M225

NODE=M225M

NODE=M225M

NODE=M225M

NODE=M225DM

NODE=M225DM

OCCUR=2

NODE=M225DM;LINKAGE=A

NODE=M225DM;LINKAGE=B

NODE=M225DM2

NODE=M225DM2

NODE=M225DM2;LINKAGE=A

NODE=M225W

NODE=M225W

OCCUR=2

OCCUR=3

NODE=M225W;LINKAGE=A

NODE=M225W;LINKAGE=B

NODE=M225W;LINKAGE=C

NODE=M225215;NODE=M225

DESIG=1

DESIG=2

$B_J(5840)^0$ BRANCHING RATIOS

$\Gamma(B^{*+}\pi^-)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	12k	AAIJ	15AB LHCB	pp at 7, 8 TeV	

$\Gamma(B^+\pi^-)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
possibly seen		¹ AAIJ	15AB LHCB	pp at 7, 8 TeV	

¹ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.

$B_J(5840)^0$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------

NODE=M225220

NODE=M225R01
NODE=M225R01

NODE=M225R02
NODE=M225R02

NODE=M225R02;LINKAGE=A

NODE=M225

REFID=56628