

$T_{b\bar{s}}(5568)^+$

$$I(J^P) = 1(?^?)$$

OMITTED FROM SUMMARY TABLE
was $X(5568)^\pm$

Seen as a peak in the $B_s \pi^\pm$ mass spectrum with a significance of more than 3σ by ABAZOV 16E and ABAZOV 18A in inclusive $p\bar{p}$ collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

NODE=M232

NODE=M232

$T_{b\bar{s}}(5568)^+$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$5566.9^{+3.2+0.6}_{-3.1-1.2}$	278	¹ ABAZOV	18A D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5567.8 \pm 2.9^{+0.9}_{-1.9}$	133	² ABAZOV	16E D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$
¹ From the combined analysis of $B_s^0 \rightarrow J/\psi \phi$ and $B_s^0 \rightarrow D_s^\pm \mu^\mp X$ decays.				
² Assumes $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ decay. If $T_{b\bar{s}}(5568)^\pm \rightarrow B_s^* \pi^\pm$ decay is assumed, the mass shifts upward by 49 MeV.				

NODE=M232M

NODE=M232M

NODE=M232M;LINKAGE=B

NODE=M232M;LINKAGE=A

$T_{b\bar{s}}(5568)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$18.6^{+7.9+3.5}_{-6.1-3.8}$	278	¹ ABAZOV	18A D0	$p\bar{p} \rightarrow B_s \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$21.9 \pm 6.4^{+5.0}_{-2.5}$	133	ABAZOV	16E D0	$p\bar{p} \rightarrow B_s \pi^\pm X$
¹ From the combined analysis of $B_s^0 \rightarrow J/\psi \phi$ and $B_s^0 \rightarrow D_s^\pm \mu^\mp X$ decays.				

NODE=M232W

NODE=M232W

NODE=M232W;LINKAGE=B

$T_{b\bar{s}}(5568)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B_s \pi^\pm$	seen

NODE=M232215;NODE=M232

DESIG=1

$T_{b\bar{s}}(5568)^+$ BRANCHING RATIOS

$\Gamma(B_s \pi^\pm)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	145	¹ ABAZOV	18A D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$	
seen	133	² ABAZOV	16E D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen		³ AABOUD	18L ATLS	$pp \rightarrow B_s^0 \pi^\pm X$	
not seen		⁴ AALTONEN	18A CDF	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$	
not seen		⁵ SIRUNYAN	18J CMS	$pp \rightarrow B_s^0 \pi^\pm X$	
not seen		⁶ AAIJ	16AI LHCB	$pp \rightarrow B_s^0 \pi^\pm X$	

NODE=M232220

NODE=M232R01
NODE=M232R01

OCCUR=2

NODE=M232R01;LINKAGE=F
NODE=M232R01;LINKAGE=A

NODE=M232R01;LINKAGE=E

NODE=M232R01;LINKAGE=D

NODE=M232R01;LINKAGE=C

¹ With B_s mesons reconstructed in decays to $D_s^\pm \mu^\mp X$.

² Seen in $p\bar{p}$ collisions at 1.96 TeV at a rate of $(8.6 \pm 1.9 \pm 1.4)\%$ relative to inclusive B_s production in the kinematic region $10 < p_T(B_s) < 30$ GeV/c, with B_s mesons reconstructed in decays to $J/\psi \phi$. An alternative possibility, $T_{b\bar{s}}(5568)^\pm \rightarrow B_s^* \pi^\pm$ with a missing γ , could not be ruled out.

³ Not seen in 24.4 fb^{-1} of pp collision data at $\sqrt{s} = 7$ and 8 TeV with B_s mesons reconstructed in decays to $J/\psi \phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 1.5% at $p_T(B_s) > 10$ GeV/c and less than 1.6% at $p_T(B_s) > 15$ GeV/c at 95% CL.

⁴ Not seen in 9.6 fb^{-1} of $p\bar{p}$ collision data at $\sqrt{s} = 1.96$ TeV with B_s mesons reconstructed in decays to $J/\psi \phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 6.7% at 95% CL.

⁵ Not seen in 19.7 fb^{-1} of pp collisions data at $\sqrt{s} = 8$ TeV with B_s mesons reconstructed in decays to $J/\psi \phi$. An upper limit on the production rate times branching fraction for

$T_{b\bar{s}}(5568)^{\pm} \rightarrow B_s \pi^{\pm}$ relative to inclusive B_s production is less than 1.1% at $p_T(B_s) > 10$ GeV/c and less than 1.0% at $p_T(B_s) > 15$ GeV/c at 95%CL.

⁶ Not seen in 3 fb⁻¹ of pp collision data at $\sqrt{s} = 7$ and 8 TeV in a scan over the $T_{b\bar{s}}(5568)$ mass and width, with B_s mesons reconstructed in decays to $D_s^- \pi^+$ or $J/\psi \phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^{\pm} \rightarrow B_s \pi^{\pm}$ relative to inclusive B_s production is less than 2.1% at $p_T(B_s) > 10$ GeV/c at 90% CL.

NODE=M232R01;LINKAGE=B

$T_{b\bar{s}}(5568)^+$ REFERENCES

NODE=M232

AABOUD	18L	PRL 120 202007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AALTONEN	18A	PRL 120 202006	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	18A	PR D97 092004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
SIRUNYAN	18J	PRL 120 202005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	16AI	PRL 117 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	16E	PRL 117 022003	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=58829
REFID=58828
REFID=58937
REFID=58827
REFID=57549
REFID=57453