

LIGHT UNFLAVORED MESONS ($S = C = B = 0$)

For $I = 1$ (π, b, ρ, a): $u\bar{d}, (u\bar{u} - d\bar{d})/\sqrt{2}, d\bar{u}$;
for $I = 0$ ($\eta, \eta', h, h', \omega, \phi, f, f'$): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

NODE=MXXX005

NODE=MXXX005

NODE=M014

 $f_0(500)$ $I^G(J^{PC}) = 0^+(0^{++})$ also known as σ ; was $f_0(600)$

See the related review(s):

[Scalar Mesons below 2 GeV](#)

$f_0(500)$ T-MATRIX POLE \sqrt{s}

Note that $\Gamma \approx 2 \text{ Im}(\sqrt{s}_{\text{pole}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–550)–i(200–350) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
(512 ± 15)–i(188 ± 12)	1 ABLIKIM 17	BES3	$J/\psi \rightarrow \gamma 3\pi$
(440 ± 10)–i(238 ± 10)	2 ALBALADEJO 12	RVUE	Compilation
(445 ± 25)–i(278 ⁺²² _{−18})	3,4 GARCIA-MAR..11	RVUE	Compilation
(457 ⁺¹⁴ _{−13})–i(279 ⁺¹¹ _{−7})	3,5 GARCIA-MAR..11	RVUE	Compilation
(442 ⁺⁵ _{−8})–i(274 ⁺⁶ _{−5})	6 MOUSSALLAM11	RVUE	Compilation
(452 ± 13)–i(259 ± 16)	7 MENNESSIER 10	RVUE	Compilation
(448 ± 43)–i(266 ± 43)	8 MENNESSIER 10	RVUE	Compilation
(455 ± 6 ⁺³¹ _{−13})–i(278 ± 6 ⁺³⁴ _{−43})	9 CAPRINI 08	RVUE	Compilation
(463 ± 6 ⁺³¹ _{−17})–i(259 ± 6 ⁺³³ _{−34})	10 CAPRINI 08	RVUE	Compilation
(552 ⁺⁸⁴ _{−106})–i(232 ⁺⁸¹ _{−72})	11 ABLIKIM 07A	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
(466 ± 18)–i(223 ± 28)	12 BONVICINI 07	CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
(472 ± 30)–i(271 ± 30)	13 BUGG 07A	RVUE	Compilation
(484 ± 17)–i(255 ± 10)	GARCIA-MAR..07	RVUE	Compilation
(430)–i(325)	14 ANISOVICH 06	RVUE	Compilation
(441 ⁺¹⁶ _{−8})–i(272 ⁺⁹ _{−12.5})	15 CAPRINI 06	RVUE	$\pi\pi \rightarrow \pi\pi$
(470 ± 50)–i(285 ± 25)	16 ZHOU 05	RVUE	
(541 ± 39)–i(252 ± 42)	17 ABLIKIM 04A	BES2	$J/\psi \rightarrow \omega \pi^+ \pi^-$
(528 ± 32)–i(207 ± 23)	18 GALLEGOS 04	RVUE	Compilation
(440 ± 8)–i(212 ± 15)	19 PELAEZ 04A	RVUE	$\pi\pi \rightarrow \pi\pi$
(533 ± 25)–i(249 ± 25)	20 BUGG 03	RVUE	
517 – i240	BLACK 01	RVUE	$\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$
(470 ± 30)–i(295 ± 20)	15 COLANGELO 01	RVUE	$\pi\pi \rightarrow \pi\pi$
(535 ⁺⁴⁸ _{−36})–i(155 ⁺⁷⁶ _{−53})	21 ISHIDA 01		$\Upsilon(3S) \rightarrow \Upsilon \pi\pi$
610 ± 14 – i620 ± 26	22 SUROVTSEV 01	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
(540 ⁺³⁶ _{−29})–i(193 ⁺³² _{−40})	ISHIDA 00B		$p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$
445 – i235	HANNAH 99	RVUE	π scalar form factor
(523 ± 12)–i(259 ± 7)	KAMINSKI 99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
442 – i 227	OLLER 99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
469 – i203	OLLER 99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
445 – i221	OLLER 99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
(1530 ⁺⁹⁰ _{−250})–i(560 ± 40)	ANISOVICH 98B	RVUE	Compilation
420 – i 212	LOCHER 98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
440 – i245	23 DOBADO 97	RVUE	Compilation
(602 ± 26)–i(196 ± 27)	24 ISHIDA 97		$\pi\pi \rightarrow \pi\pi$
(537 ± 20)–i(250 ± 17)	25 KAMINSKI 97B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, 4\pi$
470 – i250	26,27 TORNQVIST 96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
387 – i305	27,28 JANSEN 95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
420 – i370	29 ACHASOV 94	RVUE	$\pi\pi \rightarrow \pi\pi$
(506 ± 10)–i(247 ± 3)	KAMINSKI 94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$

370 – i356	³⁰ ZOU	94B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
408 – i342	^{27,30} ZOU	93	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
470 – i208	³¹ VANBEVEREN	86	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta,$
(750 ± 50) – i(450 ± 50)	³² ESTABROOKS	79	RVUE	...
(660 ± 100) – i(320 ± 70)	PROTOPOPESCU	73	HBC	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
650 – i370	³³ BASDEVANT	72	RVUE	$\pi\pi \rightarrow \pi\pi$
1 S-matrix pole; 8595 events.				
2 Applying the chiral unitary approach at NLO to the K_{e4} data of BATLEY 10 and $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.				
3 Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.				
4 Analytic continuation using Roy equations.				
5 Analytic continuation using GKPY equations.				
6 Using Roy equations.				
7 Average of three variants of the analytic K-matrix model. Uses the K_{e4} data of BATLEY 08A and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73 and GRAYER 74.				
8 Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.				
9 From the K_{e4} data of BATLEY 08A and $\pi N \rightarrow \pi\pi N$ data of HYAMS 73.				
10 From the K_{e4} data of BATLEY 08A and $\pi N \rightarrow \pi\pi N$ data of PROTOPOPESCU 73, GRAYER 74, and ESTABROOKS 74.				
11 From a mean of three different $f_0(500)$ parametrizations. Uses 40k events.				
12 From an isobar model using 2.6k events.				
13 Reanalysis of ABLIKIM 04A, PISLAK 01, and HYAMS 73 data.				
14 Using the N/D method.				
15 From the solution of the Roy equation (ROY 71) for the isoscalar S-wave and using a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.				
16 Reanalysis of the data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, ROSSELET 77, PISLAK 03, and AKHMETSHIN 04.				
17 From a mean of six different analyses and $f_0(500)$ parameterizations.				
18 Using data on $\psi(2S) \rightarrow J/\psi\pi\pi$ from BAI 00E and on $\gamma(nS) \rightarrow \gamma(mS)\pi\pi$ from BUTLER 94B and ALEXANDER 98.				
19 Reanalysis of data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.				
20 From a combined analysis of HYAMS 73, AUGUSTIN 89, AITALA 01B, and PISLAK 01.				
21 A similar analysis (KOMADA 01) finds $(580^{+79}_{-30}) - i(190^{+107}_{-49})$ MeV.				
22 Coupled channel reanalysis of BATON 70, BENSINGER 71, BAILLON 72, HYAMS 73, HYAMS 75, ROSSELET 77, COHEN 80, and ETKIN 82B using the uniformizing variable.				
23 Using the inverse amplitude method and data of ESTABROOKS 73, GRAYER 74, and PROTOPOPESCU 73.				
24 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.				
25 Average and spread of 4 variants ("up" and "down") of KAMINSKI 97B 3-channel model.				
26 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.				
27 Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.				
28 Analysis of data from FALVARD 88.				
29 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.				
30 Analysis of data from OCHS 73, GRAYER 74, and ROSSELET 77.				
31 Coupled-channel analysis using data from PROTOPOPESCU 73, HYAMS 73, HYAMS 75, GRAYER 74, ESTABROOKS 74, ESTABROOKS 75, FROGGATT 77, CORDEN 79, BISWAS 81.				
32 Analysis of data from APEL 72C, GRAYER 74, CASON 76, PAWLICKI 77. Includes spread and errors of 4 solutions.				
33 Analysis of data from BATON 70, BENSINGER 71, COLTON 71, BAILLON 72, PROTOPOPESCU 73, and WALKER 67.				

$f_0(500)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETERS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–550) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
513 ± 32	³⁴ MURAMATSU 02	CLEO	$e^+ e^- \approx 10$ GeV
$478^{+24}_{-23} \pm 17$	AITALA	01B	E791 $D^+ \rightarrow \pi^- \pi^+ \pi^+$
563^{+58}_{-29}	³⁵ ISHIDA	01	$\gamma(3S) \rightarrow \gamma\pi\pi$
555	³⁶ ASNER	00	CLE2 $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
540 ± 36	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$
750 ± 4	ALEKSEEV	99	SPEC 1.78 π^- $p_{\text{polar}} \rightarrow \pi^- \pi^+ n$
744 ± 5	ALEKSEEV	98	SPEC 1.78 π^- $p_{\text{polar}} \rightarrow \pi^- \pi^+ n$

NODE=M014PP;LINKAGE=I
 NODE=M014PP;LINKAGE=AD
 NODE=M014PP;LINKAGE=GM
 NODE=M014PP;LINKAGE=GR
 NODE=M014PP;LINKAGE=GC
 NODE=M014PP;LINKAGE=MO
 NODE=M014PP;LINKAGE=ME
 NODE=M014PP;LINKAGE=MN
 NODE=M014PP;LINKAGE=CA
 NODE=M014PP;LINKAGE=AP
 NODE=M014PP;LINKAGE=AL
 NODE=M014PP;LINKAGE=BO
 NODE=M014PP;LINKAGE=BU
 NODE=M014PP;LINKAGE=AN
 NODE=M014PP;LINKAGE=CL
 NODE=M014PP;LINKAGE=ZH
 NODE=M014PP;LINKAGE=AB
 NODE=M014PP;LINKAGE=GA
 NODE=M014PP;LINKAGE=PE
 NODE=M014PP;LINKAGE=PS
 NODE=M014PP;LINKAGE=KI
 NODE=M014PP;LINKAGE=SU
 NODE=M014PP;LINKAGE=DO
 NODE=M014PP;LINKAGE=AA
 NODE=M014PP;LINKAGE=E
 NODE=M014PP;LINKAGE=B

 NODE=M014PP;LINKAGE=G
 NODE=M014PP;LINKAGE=C
 NODE=M014PP;LINKAGE=D
 NODE=M014PP;LINKAGE=F
 NODE=M014PP;LINKAGE=BV

 NODE=M014PP;LINKAGE=A
 NODE=M014PP;LINKAGE=J

 NODE=M014M
 NODE=M014M
 → UNCHECKED ←

759 ± 5	³⁷ TROYAN	98	5.2 $np \rightarrow np\pi^+\pi^-$
780 ± 30	ALDE	97	GAM2 450 $pp \rightarrow pp\pi^0\pi^0$
585 ± 20	³⁸ ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
761 ± 12	³⁹ SVEC	96	RVUE 6–17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 860	^{40,41} TORNQVIST	96	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1165 ± 50	^{42,43} ANISOVICH	95	RVUE $\pi^- p \rightarrow \pi^0\pi^0 n, \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
~ 1000	⁴⁴ ACHASOV	94	RVUE $\pi\pi \rightarrow \pi\pi$
414 ± 20	³⁹ AUGUSTIN	89	DM2

34 Statistical uncertainty only.

35 A similar analysis (KOMADA 01) finds 526^{+48}_{-37} MeV.

36 From the best fit of the Dalitz plot.

37 6σ effect, no PWA.

38 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

39 Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+ N$ on polarized targets. The fit does not include $f_0(980)$.

40 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

41 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.

42 Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.

43 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

44 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.

$f_0(500)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400–700) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
335 ± 67	⁴⁵ MURAMATSU	02 CLEO	$e^+e^- \approx 10$ GeV
$324^{+42}_{-40} \pm 21$	AITALA	01B E791	$D^+ \rightarrow \pi^-\pi^+\pi^+$
372^{+229}_{-95}	⁴⁶ ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
540	⁴⁷ ASNER	00 CLE2	$\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
372 ± 80	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
119 ± 13	ALEKSEEV	99 SPEC	1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
77 ± 22	ALEKSEEV	98 SPEC	1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
35 ± 12	⁴⁸ TROYAN	98	5.2 $np \rightarrow np\pi^+\pi^-$
780 ± 60	ALDE	97 GAM2	450 $pp \rightarrow pp\pi^0\pi^0$
385 ± 70	⁴⁹ ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
290 ± 54	⁵⁰ SVEC	96 RVUE	6–17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 880	^{51,52} TORNQVIST	96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
460 ± 40	^{53,54} ANISOVICH	95 RVUE	$\pi^- p \rightarrow \pi^0\pi^0 n, \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
~ 3200	⁵⁵ ACHASOV	94 RVUE	$\pi\pi \rightarrow \pi\pi$
494 ± 58	⁵⁰ AUGUSTIN	89 DM2	

45 Statistical uncertainty only.

46 A similar analysis (KOMADA 01) finds 301^{+145}_{-100} MeV.

47 From the best fit of the Dalitz plot.

48 6σ effect, no PWA.

49 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

50 Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+ N$ on polarized targets. The fit does not include $f_0(980)$.

51 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

52 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.

53 Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.

54 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

55 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.

NODE=M014M;LINKAGE=UT

NODE=M014M;LINKAGE=KI

NODE=M014M;LINKAGE=KK

NODE=M014M;LINKAGE=TN

NODE=M014M;LINKAGE=AA

NODE=M014M;LINKAGE=E

NODE=M014M;LINKAGE=B

NODE=M014M;LINKAGE=GG

NODE=M014M;LINKAGE=F

NODE=M014M;LINKAGE=G

NODE=M014M;LINKAGE=D

NODE=M014W

NODE=M014W

→ UNCHECKED ←

NODE=M014W;LINKAGE=UT

NODE=M014W;LINKAGE=KI

NODE=M014W;LINKAGE=KK

NODE=M014W;LINKAGE=TN

NODE=M014W;LINKAGE=AA

NODE=M014W;LINKAGE=E

NODE=M014W;LINKAGE=B

NODE=M014W;LINKAGE=GG

NODE=M014W;LINKAGE=F

NODE=M014W;LINKAGE=G

NODE=M014W;LINKAGE=D1

$f_0(500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 \gamma\gamma$	seen

 $f_0(500)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2
<u>VALUE (keV)</u>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.05±0.21	56 DAI	14A	RVUE Compilation	
1.7 ± 0.4	57 HOFERICHTER11	RVUE	Compilation	
3.08±0.82	58 MENNESSIER 11	RVUE	Compilation	
2.08±0.2 $^{+0.07}_{-0.04}$	59 MOUSSALLAM11	RVUE	Compilation	
2.08	60 MAO	09	RVUE Compilation	
1.2 ± 0.4	61 BERNABEU	08	RVUE	
3.9 ± 0.6	58 MENNESSIER 08	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
1.8 ± 0.4	62 OLLER	08	RVUE Compilation	
1.68±0.15	62,63 OLLER	08A	RVUE Compilation	
3.1 ± 0.5	64,65 PENNINGTON 08	RVUE	Compilation	
2.4 ± 0.4	65,66 PENNINGTON 08	RVUE	Compilation	
4.1 ± 0.3	67 PENNINGTON 06	RVUE	$\gamma\gamma \rightarrow \pi^0\pi^0$	
3.8 ± 1.5	68,69 BOGLIONE	99	RVUE $\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
5.4 ± 2.3	68 MORGAN	90	RVUE $\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
10 ± 6	COURAU	86	$e^+e^- \rightarrow \pi^+\pi^- e^+e^-$	
56 Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.				
57 Using Roy-Steiner equations with $\pi\pi$ phase shifts from an update of COLANGELO 01 and from GARCIA-MARTIN 11A.				
58 Using an analytic K-matrix model.				
59 Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.				
60 Used dispersion theory. The value quoted used the $f_0(500)$ pole position of 457 – i276 MeV.				
61 Using p, n polarizabilities from PDG 06 and fitting to $\pi\pi$ phase motion from GARCIA-MARTIN 07 and σ -poles from GARCIA-MARTIN 07 and CAPRINI 06.				
62 Using twice-subtracted dispersion integrals.				
63 Supersedes OLLER 08.				
64 Solution A (preferred solution based on χ^2 -analysis).				
65 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.				
66 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).				
67 Using unitarity and the σ pole position from CAPRINI 06.				
68 This width could equally well be assigned to the $f_0(1370)$. The authors analyse data from BOYER 90 and MARSISKE 90 and report strong correlation with $\gamma\gamma$ width of $f_2(1270)$.				
69 Supersedes MORGAN 90.				

 $f_0(500)$ REFERENCES

ABLIKIM 17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DAI 14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
ALBALADEJO 12	PR D86 034003	M. Albaladejo, J.A. Oller	(MURC)
GARCIA-MAR...11	PR D74 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
GARCIA-MAR...11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
HOFERICHTER 11	EPJ C71 1743	M. Hoferichter, D.R. Phillips, C. Schat	(BONN+)
MENNESSIER 11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang	
MOUSSALLAM 11	EPJ C71 1814	B. Moussallam	
BATLEY 10	PL B686 101	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY 10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
MENNESSIER 10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang	
MAO 09	PR D79 116008	Y. Mao <i>et al.</i>	
BATLEY 08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BERNABEU 08	PRL 100 241804	J. Bernabeu, J. Prades	(IFIC, GRAN)
CAPRINI 08	PR D77 114019	I. Caprini	
MENNESSIER 08	PL B665 205	G. Mennessier, S. Narison, W. Ochs	
OLLER 08	PL B659 201	J.A. Oller, L. Roca, C. Schat	(MURC, UBA)
OLLER 08A	EPJ A37 15	J.A. Oller, L. Roca	(MURC)
PENNINGTON 08	EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA 08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM 07A	PL B645 19	M. Ablikim <i>et al.</i>	(BES Collab.)
BONVICINI 07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG 07A	JP G34 151	D.V. Bugg <i>et al.</i>	
GARCIA-MAR...07	PR D76 074034	R. Garcia-Martin, J.R. Pelaez, F.J. Yndurain	
MORI 07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)

NODE=M014215;NODE=M014

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M014220

NODE=M014W2
NODE=M014W2

OCCUR=2

NODE=M014W2;LINKAGE=B

NODE=M014W2;LINKAGE=HO

NODE=M014W2;LINKAGE=ME

NODE=M014W2;LINKAGE=MO

NODE=M014W2;LINKAGE=MA

NODE=M014W2;LINKAGE=BE

NODE=M014W2;LINKAGE=OL

NODE=M014W2;LINKAGE=LL

NODE=M014W2;LINKAGE=P1

NODE=M014W2;LINKAGE=P3

NODE=M014W2;LINKAGE=P2

NODE=M014W2;LINKAGE=PE

NODE=M014W2;LINKAGE=A

NODE=M014W2;LINKAGE=BL

NODE=M014

REFID=57650

REFID=55923

REFID=54277

REFID=16761

REFID=54121

REFID=53966

REFID=53637

REFID=53975

REFID=53285

REFID=53567

REFID=53657

REFID=52992

REFID=52487

REFID=52271

REFID=52273

REFID=52299

REFID=53976

REFID=53977

REFID=52303

REFID=52309

REFID=51614

REFID=51721

REFID=53252

REFID=51949

REFID=51652

ANISOVICH	06	IJMP A21 3615	V.V. Anisovich	REFID=51137
CAPRINI	06	PRL 96 132001	I. Caprini, G. Colangelo, H. Leutwyler W.-M. Yao <i>et al.</i>	(BCIP+) REFID=51076
PDG	06	JP G33 1	M.R. Pennington	(PDG Collab.) REFID=51004
PENNINGTON	06	PR D97 011601	Z.Y. Zhou <i>et al.</i>	REFID=51184
ZHOU	05	JHEP 0502 043	M. Ablikim <i>et al.</i>	(BES Collab.) REFID=50823
ABLIKIM	04A	PL B598 149	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.) REFID=49740
AKHMETSHIN	04	PL B578 285	P. Buettiker, S. Descotes-Genon, B. Moussallam	REFID=49609
BUETTIKER	04	EPJ C33 409	A. Gallegos <i>et al.</i>	REFID=56428;ERROR=1
GALLEGOS	04	PR D69 074033	J.R. Pelaez	REFID=49769
PELAEZ	04A	MPL A19 2879	D.V. Bugg	REFID=50347
BUGG	03	PL B572 1	S. Pislak <i>et al.</i>	REFID=49586
PISLAK	03	PR D67 072004	(BNL E865 Collab.)	REFID=49344
Also		PR D81 119903E	S. Pislak <i>et al.</i>	REFID=53337
MURAMATSU	02	PRL 89 251802	H. Muramatsu <i>et al.</i>	(CLEO Collab.) REFID=49081
Also		PRL 90 059901 (errat.)	H. Muramatsu <i>et al.</i>	(CLEO Collab.) REFID=49385
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.) REFID=48005
BLACK	01	PR D64 014031	D. Black <i>et al.</i>	REFID=48314
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler	REFID=49180
ISHIDA	01	PL B518 47	M. Ishida <i>et al.</i>	REFID=48354
KOMADA	01	PL B508 31	T. Komada <i>et al.</i>	REFID=48541
PISLAK	01	PRL 87 221801	S. Pislak <i>et al.</i>	(BNL E865 Collab.) REFID=48433
Also		PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.) REFID=49344
Also		PRL 105 019901E	S. Pislak <i>et al.</i>	(BNL E865 Collab.) REFID=53338
SUROVTSEV	01	PR D63 054024	Y.S. Surovtsev, D. Krupa, M. Nagy	REFID=48310
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.) REFID=47339
BAI	00E	PR D62 032002	J. Bai <i>et al.</i>	(BES Collab.) REFID=47955
ISHIDA	00B	PTP 104 203	M. Ishida <i>et al.</i>	REFID=48358
ALEKSEEV	99	NP B541 3	I.G. Alekseev <i>et al.</i>	REFID=46614
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington	REFID=46931
HANNAH	99	PR D60 017502	T. Hannah	REFID=46935
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN) REFID=46927
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>	(CLEO Collab.) REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset	REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset	REFID=47386
ALEKSEEV	98	PAN 61 174	I.G. Alekseev <i>et al.</i>	REFID=46328
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.) REFID=46329
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	REFID=46331
		Translated from UFN 168 481.		
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI) REFID=46372
TROYAN	98	JINRRC 5-91 33	Yu. Troyan <i>et al.</i>	REFID=46615
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.) REFID=45392
DOBADO	97	PR D56 3057	A. Dobado, J.R. Pelaez	REFID=53964
ISHIDA	97	PTP 98 1005	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK) REFID=45998
KAMINSKI	97B	PL B413 130	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, IPN) REFID=45778
Also		PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK) REFID=45770
SVEC	96	PR D53 2343	M. Svec	(MCGI) REFID=44509
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS) REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.) REFID=44375
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP) REFID=44442
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADDL, JULI) REFID=44508
ACHASOV	94	PR D49 5779	N.N. Achasov, G.N. Shestakov	(NOVM) REFID=44087
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) REFID=43659
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.) REFID=43799
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+) REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM) REFID=44072
ZOU	93	PR D48 3948	B.S. Zou, D.V. Bugg	(LOQM) REFID=43672
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.) REFID=43172
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) REFID=41862
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.) REFID=41362
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.) REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH) REFID=41583
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.) REFID=41004
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) REFID=40262
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+) REFID=40576
COURAU	86	NP B271 1	A. Courau <i>et al.</i>	(CLER, LALO) REFID=44510
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i>	(NIJM, BIEL) REFID=45769
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL) REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND) REFID=20390
BISWAS	81	PRL 47 1378	N.N. Biswas <i>et al.</i>	(NDAM, ANL) REFID=21106
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP REFID=20381
MUKHIN	80	JETPL 32 601	K.N. Mukhin <i>et al.</i>	(KIAE) REFID=44528
		Translated from ZETFP 32 616.		
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP REFID=20374
ESTABROOKS	79	PR D19 2678	P. Estabrooks	(CARL) REFID=20375
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD) REFID=21072
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJ REFID=20367
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL) REFID=11004
CASON	76	PR D16 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL) IJ REFID=21064
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH) REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM) REFID=20355
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL) REFID=21062
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH) REFID=20111
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM) REFID=20113
ESTABROOKS	73	Tallahassee	P.G. Estabrooks <i>et al.</i>	(CERN, MPIM) REFID=20345
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM) REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI) REFID=20349
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL) REFID=20108
APEL	72C	PL 41B 542	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA) REFID=21013
BAILLON	72	PL 38B 555	P.H. Baillon <i>et al.</i>	(SLAC) REFID=20093
BASDEVANT	72	PL 41B 178	J.L. Basdevant, C.D. Froggatt, J.L. Petersen	(CERN) REFID=20095
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN) REFID=44530
BENSINGER	71	PL 36B 134	J.R. Bensinger <i>et al.</i>	(WISC) REFID=21006
COLTON	71	PR D3 2028	E.P. Colton <i>et al.</i>	(LBL, FNAL, UCLA+) REFID=44533
ROY	71	PL 36B 353	S.M. Roy	REFID=51107
BATON	70	PL 33B 528	J.P. Baton, G. Laurens, J. Reignier	(SACL) REFID=20086
WALKER	67	RMP 39 695	W.D. Walker	(WISC) REFID=20960

$\rho(770)$ $I^G(J^{PC}) = 1^+(1^{--})$

See the related review(s):

 $\rho(770)$ **$\rho(770)$ MASS**

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NEUTRAL ONLY, e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
775.26±0.25 OUR AVERAGE				
775.02±0.35		1 LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
775.97±0.46±0.70	900k	2 AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
774.6 ±0.4 ±0.5	800k	3,4 ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
775.65±0.64±0.50	114k	5,6 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
775.9 ±0.5 ±0.5	1.98M	7 ALOISIO 03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.8 ±0.9 ±2.0	500k	7 ACHASOV 02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ±1.1		8 BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
763.49±0.53		9 BARTOS 17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
758.23±0.46		10 BARTOS 17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
775.8 ±0.5 ±0.3	1.98M	11 ALOISIO 03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ±0.6 ±0.5	1.98M	12 ALOISIO 03	KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.0 ±0.6 ±1.1	500k	13 ACHASOV 02	SND	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±0.7 ±5.3		14 BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-,\mu^+\mu^-$
770.5 ±1.9 ±5.1		15 GARDNER 98	RVUE	0.28–0.92 $e^+e^- \rightarrow \pi^+\pi^-$
764.1 ±0.7		16 O'CONNELL 97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
757.5 ±1.5		17 BERNICHA 94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
768 ±1		18 GESHKEN... 89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$

¹ Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

² A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

³ Supersedes ACHASOV 05A.

⁴ A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

⁵ Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.

⁶ Update of AKHMETSHIN 02.

⁷ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁸ From the GOUNARIS 68 parametrization of the pion form factor.

⁹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

¹⁰ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

¹¹ Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.

¹² Without limitations on masses and widths.

¹³ Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0\pi\pi} = g_{\rho^\pm\pi\pi}$.

¹⁴ Using the data of BARKOV 85 in the hidden local symmetry model.

¹⁵ From the fit to $e^+e^- \rightarrow \pi^+\pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.

¹⁶ A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.

¹⁷ Applying the S-matrix formalism to the BARKOV 85 data.

¹⁸ Includes BARKOV 85 data. Model-dependent width definition.

CHARGED ONLY, τ DECAYS and e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
775.11±0.34 OUR AVERAGE					
774.6 ±0.2 ±0.5	5.4M	1,2 FUJIKAWA 08	BELL	±	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
775.5 ±0.7		2,3 SCHABEL 05C	ALEP		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
775.5 ±0.5 ±0.4	1.98M	4 ALOISIO 03	KLOE		1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±1.1 ±0.5	87k	5,6 ANDERSON 00A	CLE2		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

NODE=M009

NODE=M009205

NODE=M009205

NODE=M009M0
NODE=M009M0

OCCUR=2

OCCUR=3

OCCUR=3

NODE=M009M0;LINKAGE=LE

NODE=M009M0;LINKAGE=AK
NODE=M009M0;LINKAGE=AC
NODE=M009M0;LINKAGE=SNNODE=M009M0;LINKAGE=GS
NODE=M009M0;LINKAGE=PTNODE=M009M0;LINKAGE=CH
NODE=M009M0;LINKAGE=K
NODE=M009M0;LINKAGE=A

NODE=M009M0;LINKAGE=B

NODE=M009M0;LINKAGE=DF
NODE=M009M0;LINKAGE=WONODE=M009M0;LINKAGE=HC
NODE=M009M0;LINKAGE=K2
NODE=M009M0;LINKAGE=G8NODE=M009M0;LINKAGE=AB
NODE=M009M0;LINKAGE=AA
NODE=M009M0;LINKAGE=FNODE=M009M5
NODE=M009M5

• • • We do not use the following data for averages, fits, limits, etc. • • •

761.60 ± 0.95	⁷ BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
$774.8 \pm 0.6 \pm 0.4$ 1.98M	⁸ ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	OCCUR=2
$776.3 \pm 0.6 \pm 0.7$ 1.98M	⁸ ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	OCCUR=3
$773.9 \pm 2.0 \pm 0.3$ 1.0	⁹ SANZ-CILLERO03		RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
$774.5 \pm 0.7 \pm 1.5$ 500k	⁴ ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	OCCUR=2
775.1 ± 0.5	¹⁰ PICH	01	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	

1 $|F_\pi(0)|^2$ fixed to 1.

2 From the GOUNARIS 68 parametrization of the pion form factor.

3 The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.

4 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

5 $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.

6 From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.

7 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.

8 Without limitations on masses and widths.

9 Using the data of BARATE 97M and the effective chiral Lagrangian.

10 From a fit of the model-independent parameterization of the pion form factor to the data of BARATE 97M.

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
763.0±0.3±1.2	600k	¹ ABELE	99E	CBAR	$0 \pm 0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$

1 Assuming the equality of ρ^+ and ρ^- masses and widths.

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
766.5±1.1 OUR AVERAGE					

763.7 ± 3.2		ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
768 ± 9		AGUILAR-...	91	EHS	$400 pp$
767 ± 3	2935	¹ CAPRARO	87	SPEC	$200 \pi^- Cu \rightarrow \pi^- \pi^0 Cu$
761 ± 5	967	¹ CAPRARO	87	SPEC	$200 \pi^- Pb \rightarrow \pi^- \pi^0 Pb$
771 ± 4		HUSTON	86	SPEC	$202 \pi^+ A \rightarrow \pi^+ \pi^0 A$
766 ± 7	6500	² BYERLY	73	OSPK	$5 \pi^- p$
766.8 ± 1.5	9650	³ PISUT	68	RVUE	$1.7-3.2 \pi^- p, t < 10$
767 ± 6	900	¹ EISNER	67	HBC	$4.2 \pi^- p, t < 10$

1 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

2 Phase shift analysis. Systematic errors added corresponding to spread of different fits.

3 From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
769.0± 1.0 OUR AVERAGE				

$771 \pm 2 \pm 2$	63.5k	¹ ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
$770 \pm 2 \pm 1$	79k	² BREITWEG	98B	ZEUS $50-100 \gamma p$
767.6 ± 2.7		BARTALUCCI	78	CNTR $\gamma p \rightarrow e^+ e^- p$
775 ± 5		GLADDING	73	CNTR $2.9-4.7 \gamma p$
767 ± 4	1930	BALLAM	72	HBC $2.8 \gamma p$
770 ± 4	2430	BALLAM	72	HBC $4.7 \gamma p$
765 ± 10		ALVENSLEB...	70	CNTR $\gamma A, t < 0.01$
767.7 ± 1.9	140k	BIGGS	70	CNTR $<4.1 \gamma C \rightarrow \pi^+ \pi^- C$
765 ± 5	4000	ASBURY	67B	CNTR $\gamma + Pb$

• • • We do not use the following data for averages, fits, limits, etc. • • •

771 ± 2	79k	³ BREITWEG	98B	ZEUS $50-100 \gamma p$
-------------	-----	-----------------------	-----	------------------------

1 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.

2 From the parametrization according to SOEDING 66.

3 From the parametrization according to ROSS 66.

NODE=M009M5;LINKAGE=FU
NODE=M009M5;LINKAGE=GO
NODE=M009M5;LINKAGE=SC
NODE=M009M5;LINKAGE=CH
NODE=M009M5;LINKAGE=A6
NODE=M009M5;LINKAGE=K1
NODE=M009M5;LINKAGE=A
NODE=M009M5;LINKAGE=WO
NODE=M009M5;LINKAGE=Z
NODE=M009M5;LINKAGE=PC

NODE=M009M7
NODE=M009M7

NODE=M009M7;LINKAGE=LB

NODE=M009M2
NODE=M009M2

OCCUR=2

NODE=M009M2;LINKAGE=Z
NODE=M009M2;LINKAGE=X
NODE=M009M2;LINKAGE=A

NODE=M009M0P
NODE=M009M0P

OCCUR=2

OCCUR=2

NODE=M009M0P;LINKAGE=AB
NODE=M009M0P;LINKAGE=B5
NODE=M009M0P;LINKAGE=B6

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
769.0 ±0.9 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
765 ±6		BERTIN 97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
773 ±1.6		WEIDENAUER 93	ASTE	$\bar{p}p \rightarrow \pi^+ \pi^- \omega$
762.6 ±2.6		AGUILAR-...	EHS	400 $p p$
770 ±2	1	HEYN 81	RVUE	Pion form factor
768 ±4	2,3	BOHACIK 80	RVUE	
769 ±3		WICKLUND 78	ASPK	3,4,6 $\pi^\pm N$
768 ±1	76k	DEUTSCH...	HBC	16 $\pi^+ p$
767 ±4	4100	ENGLER 74	DBC	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
775 ±4	32k	PROTOPOP...	HBC	7.1 $\pi^+ p, t < 0.4$
764 ±3	6.8k	RATCLIFF 72	ASPK	15 $\pi^- p, t < 0.3$
774 ±3	1.7k	REYNOLDS 69	HBC	2.26 $\pi^- p$
769.2 ±1.5	13.3k	PISUT 68	RVUE	1.7–3.2 $\pi^- p, t < 10$

• • • We do not use the following data for averages, fits, limits, etc. • • •

774.34 ±0.18 ±0.35	970k	7 ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma \pi^+ \pi^-$
772.93 ±0.18 ±0.34	970k	8 ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma \pi^+ \pi^-$
773.5 ±2.5		9 COLANGELO 01	RVUE	$\pi\pi \rightarrow \pi\pi$
762.3 ±0.5 ±1.2	600k	10 ABELE 99E	CBAR	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
777 ±2	4.9k	11 ADAMS 97	E665	470 $\mu p \rightarrow \mu XB$
770 ±2		12 BOGOLYUB...	MIRA	32 $\bar{p}p \rightarrow \pi^+ \pi^- X$
768 ±8		12 BOGOLYUB...	MIRA	32 $p p \rightarrow \pi^+ \pi^- X$
761.1 ±2.9		DUBNICKA 89	RVUE	Pion form factor
777.4 ±2.0		13 CHABAUD 83	ASPK	17 $\pi^- p$ polarized
769.5 ±0.7		2,3 LANG 79	RVUE	
770 ±9		3 ESTABROOKS 74	RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
773.5 ±1.7	11.2k	14 JACOBS 72	HBC	2.8 $\pi^- p$
775 ±3	2.2k	15 HYAMS 68	OSPK	11.2 $\pi^- p$

1 HEYN 81 includes all spacelike and timelike F_π values until 1978.

2 From pole extrapolation.

3 From phase shift analysis of GRAYER 74 data.

4 Phase shift analysis. Systematic errors added corresponding to spread of different fits.

5 Published values contain misprints. Corrected by private communication RATCLIFF 74.

6 Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.

7 From a fit to $\pi^+ \pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.

8 From a fit to $\pi^+ \pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.

9 Breit-Wigner mass from a phase-shift analysis of HYAMS 73 and PROTOPOPOV 73 data.

10 Using relativistic Breit-Wigner and taking into account $\rho\omega$ interference.

11 Systematic errors not evaluated.

12 Systematic effects not studied.

13 From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P-wave intensity. CHABAUD 83 includes data of GRAYER 74.

14 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

15 Of HYAMS 68 six parametrizations, this is theoretically soundest. MR

NODE=M009M0R

NODE=M009M0R

OCCUR=2

OCCUR=2

NODE=M009M0R;LINKAGE=B0
NODE=M009M0R;LINKAGE=C0
NODE=M009M0R;LINKAGE=H
NODE=M009M0R;LINKAGE=X
NODE=M009M0R;LINKAGE=03
NODE=M009M0R;LINKAGE=R

NODE=M009M0R;LINKAGE=B


NODE=M009M0R;LINKAGE=C

NODE=M009M0R;LINKAGE=CL

NODE=M009M0R;LINKAGE=BL
NODE=M009M0R;LINKAGE=A1
NODE=M009M0R;LINKAGE=QQ
NODE=M009M0R;LINKAGE=G

NODE=M009M0R;LINKAGE=Z

NODE=M009M0R;LINKAGE=02



$m_{\rho(770)^0} - m_{\rho(770)^{\pm}}$

NODE=M009D

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.7 ±0.8 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
-2.4 ±0.8		1 SCHAEL	05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
0.4 ±0.7 ±0.6	1.98M	2 ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.3 ±1.1 ±2.0	500k	2 ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.6 ±0.6 ±1.7	600k	ABELE	99E	CBAR	$\pm 0 \quad 0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
-4 ±4	3000	3 REYNOLDS	69	HBC	-0 $2.26 \pi^- p$
-5 ±5	3600	3 FOSTER	68	HBC	$\pm 0 \quad 0.0 \bar{p}p$
2.4 ±2.1	22950	4 PISUT	68	RVUE	$\pi N \rightarrow \rho N$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
-3.37±1.06		5 BARTOS	17A	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

NODE=M009D;LINKAGE=SC

¹ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEL 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

² Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

³ From quoted masses of charged and neutral modes.

⁴ Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDHABER 64, ABOLINS 63.


⁵ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M009D;LINKAGE=CH

NODE=M009D;LINKAGE=A

NODE=M009D;LINKAGE=R

NODE=M009D;LINKAGE=B



VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.5 \pm 0.8 \pm 0.7$	1.98M	¹ ALOISIO	03	KLOE $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
¹ Without limitations on masses and widths.				

$\rho(770)$ RANGE PARAMETER

The range parameter R enters an energy-dependent correction to the width, of the form $(1 + q_f^2 R^2) / (1 + q^2 R^2)$, where q is the momentum of one of the pions in the $\pi\pi$ rest system. At resonance, $q = q_f$.

VALUE (GeV $^{-1}$)	DOCUMENT ID	TECN	CHG	COMMENT
$5.3^{+0.9}_{-0.7}$	¹ CHABAUD	83	ASPK	0 $\pi^- \rho$ polarized

¹ The old PISUT 68 value, properly corrected, was 3.2 ± 0.6 .

$\rho(770)$ WIDTH

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NEUTRAL ONLY, $e^+ e^-$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
147.8 ± 0.9 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.				
149.59 ± 0.67		¹ LEES	12G	BABR $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
145.98 ± 0.75 ± 0.50	900k	² AKHMETSHIN	07	$e^+ e^- \rightarrow \pi^+ \pi^-$
146.1 ± 0.8 ± 1.5	800k	^{3,4} ACHASOV	06	$e^+ e^- \rightarrow \pi^+ \pi^-$
143.85 ± 1.33 ± 0.80	114k	^{5,6} AKHMETSHIN	04	CMD2 $e^+ e^- \rightarrow \pi^+ \pi^-$
147.3 ± 1.5 ± 0.7	1.98M	⁷ ALOISIO	03	KLOE $1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
151.1 ± 2.6 ± 3.0	500k	⁷ ACHASOV	02	SND $1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.5 ± 3.0		⁸ BARKOV	85	OLYA $e^+ e^- \rightarrow \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
144.06 ± 0.85		⁹ BARTOS	17	RVUE $e^+ e^- \rightarrow \pi^+ \pi^-$
144.56 ± 0.80		¹⁰ BARTOS	17A	RVUE $e^+ e^- \rightarrow \pi^+ \pi^-$
143.9 ± 1.3 ± 1.1	1.98M	¹¹ ALOISIO	03	KLOE $1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
147.4 ± 1.5 ± 0.7	1.98M	¹² ALOISIO	03	KLOE $1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M009D1

NODE=M009D1

NODE=M009D;LINKAGE=WO

NODE=M009R

NODE=M009R

NODE=M009R

NODE=M009R;LINKAGE=01

NODE=M009220

NODE=M009220

NODE=M009W0
NODE=M009W0

OCCUR=2

OCCUR=3

149.8 \pm 2.2 \pm 2.0	500k	13 ACHASOV	02 SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
147.9 \pm 1.5 \pm 7.5		14 BENAYOUN	98 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
153.5 \pm 1.3 \pm 4.6		15 GARDNER	98 RVUE	$0.28-0.92 e^+ e^- \rightarrow \pi^+ \pi^-$
145.0 \pm 1.7		16 O'CONNELL	97 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
142.5 \pm 3.5		17 BERNICHA	94 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
138 \pm 1		18 GEHKEN...	89 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$

1 Using the GOUNARIS 68 parametrization with the complex phase of the $\rho-\omega$ interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

2 A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

3 Supersedes ACHASOV 05A.

4 A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

5 Using the GOUNARIS 68 parametrization with the complex phase of the $\rho-\omega$ interference.

6 From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02.

7 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

8 From the GOUNARIS 68 parametrization of the pion form factor.

9 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

10 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

11 Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.

12 Without limitations on masses and widths.

13 Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0 \pi\pi} = g_{\rho^\pm \pi\pi}$.

14 Using the data of BARKOV 85 in the hidden local symmetry model.

15 From the fit to $e^+ e^- \rightarrow \pi^+ \pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.

16 A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.

17 Applying the S-matrix formalism to the BARKOV 85 data.

18 Includes BARKOV 85 data. Model-dependent width definition.

OCCUR=3

NODE=M009W0;LINKAGE=LE

NODE=M009W;LINKAGE=AK
NODE=M009W0;LINKAGE=AC
NODE=M009W0;LINKAGE=SN

NODE=M009W5;LINKAGE=GS
NODE=M009W5;LINKAGE=P2


NODE=M009W;LINKAGE=CH
NODE=M009W;LINKAGE=K
NODE=M009W0;LINKAGE=A

NODE=M009W0;LINKAGE=B

NODE=M009W;LINKAGE=DF
NODE=M009W;LINKAGE=WO

NODE=M009W;LINKAGE=HC
NODE=M009W;LINKAGE=K2
NODE=M009W;LINKAGE=G8

NODE=M009W;LINKAGE=AB
NODE=M009W;LINKAGE=AA
NODE=M009W;LINKAGE=F



CHARGED ONLY, τ DECAYS and $e^+ e^-$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
149.1 \pm 0.8 OUR FIT					
149.1 \pm 0.8 OUR AVERAGE					
148.1 \pm 0.4 \pm 1.7 5.4M	1,2 FUJIKAWA	08 BELL	\pm	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
149.0 \pm 1.2	2,3 SCHABEL	05C ALEP		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
149.9 \pm 2.3 \pm 2.0 500k	4 ACHASOV	02 SND	\pm	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
150.4 \pm 1.4 \pm 1.4 87k	5,6 ANDERSON	00A CLE2		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	

NODE=M009W5
NODE=M009W5

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

139.90 ± 0.46	⁷ BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
$143.7 \pm 1.3 \pm 1.2$ 1.98M	⁴ ALOISIO	03	KLOE	$\pm 1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
$142.9 \pm 1.3 \pm 1.4$ 1.98M	⁸ ALOISIO	03	KLOE	$- 1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	OCCUR=2
$144.7 \pm 1.4 \pm 1.2$ 1.98M	⁸ ALOISIO	03	KLOE	$+ 1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	OCCUR=3
$150.2 \pm 2.0 \pm 0.7$ $- 1.6$	⁹ SANZ-CILLERO03		RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	OCCUR=4
$150.9 \pm 2.2 \pm 2.0$ 500k	¹⁰ ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	

¹ $|F_\pi(0)|^2$ fixed to 1.

² From the GOUNARIS 68 parametrization of the pion form factor.

³ The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.

⁴ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁵ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.

⁶ From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.

⁷ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.

⁸ Without limitations on masses and widths.

⁹ Using the data of BARATE 97M and the effective chiral Lagrangian.

¹⁰ Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0 \pi \pi} = g_{\rho^\pm \pi \pi}$.

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
149.5 ± 1.3	600k	¹ ABELE	99E	CBAR	$0 \pm 0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$

¹ Assuming the equality of ρ^+ and ρ^- masses and widths.

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150.2 ± 2.4 OUR FIT					

150.2 ± 2.4 OUR AVERAGE

152.8 ± 4.3		ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
155 ± 11	2.9k	¹ CAPRARO	87	SPEC	$- 200 \pi^- Cu \rightarrow \pi^- \pi^0 Cu$
154 ± 20	967	¹ CAPRARO	87	SPEC	$- 200 \pi^- Pb \rightarrow \pi^- \pi^0 Pb$
150 ± 5		HUSTON	86	SPEC	$+ 202 \pi^+ A \rightarrow \pi^+ \pi^0 A$
146 ± 12	6.5k	² BYERLY	73	OSPK	$- 5 \pi^- p$
148.2 ± 4.1	9.6k	³ PISUT	68	RVUE	$- 1.7-3.2 \pi^- p, t < 10$
146 ± 13	900	EISNER	67	HBC	$- 4.2 \pi^- p, t < 10$

• • • We do not use the following data for averages, fits, limits, etc. • • •

137.0 ± 0.4		⁴ ABLIKIM	17	BES3	$J/\psi \rightarrow \gamma 3\pi$
-------------	--	----------------------	----	------	----------------------------------

¹ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

² Phase shift analysis. Systematic errors added corresponding to spread of different fits.

³ From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

⁴ S-matrix pole at a fixed ρ meson mass of 775.49 MeV.

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
151.7 ± 2.6 OUR AVERAGE				

$155 \pm 5 \pm 2$	63.5k	¹ ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
$146 \pm 3 \pm 13$	79k	² BREITWEG	98B	ZEUS $50-100 \gamma p$
150.9 ± 3.0		BARTALUCCI	78	CNTR $\gamma p \rightarrow e^+ e^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

138 ± 3	79k	³ BREITWEG	98B	ZEUS $50-100 \gamma p$
147 ± 11		GLADDING	73	CNTR $2.9-4.7 \gamma p$
155 ± 12	2430	BALLAM	72	HBC $4.7 \gamma p$
145 ± 13	1930	BALLAM	72	HBC $2.8 \gamma p$
140 ± 5		ALVENSLEB...	70	CNTR $\gamma A, t < 0.01$
146.1 ± 2.9	140k	BIGGS	70	CNTR $<4.1 \gamma C \rightarrow \pi^+ \pi^- C$
160 ± 10		LANZEROTTI	68	CNTR γp
130 ± 5	4000	ASBURY	67B	CNTR $\gamma + Pb$

¹ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.

² From the parametrization according to SOEDING 66.

³ From the parametrization according to ROSS 66.

NODE=M009W5;LINKAGE=FU
NODE=M009W5;LINKAGE=GO
NODE=M009W5;LINKAGE=SC

NODE=M009W5;LINKAGE=CH
NODE=M009W5;LINKAGE=A6
NODE=M009W5;LINKAGE=K1

NODE=M009W5;LINKAGE=A
NODE=M009W5;LINKAGE=WO
NODE=M009W5;LINKAGE=Z
NODE=M009W5;LINKAGE=HC

NODE=M009W7
NODE=M009W7

NODE=M009W;LINKAGE=LB

NODE=M009W2
NODE=M009W2

OCCUR=2

NODE=M009W2;LINKAGE=Z
NODE=M009W2;LINKAGE=X
NODE=M009W2;LINKAGE=A1

NODE=M009W2;LINKAGE=A

NODE=M009W0P
NODE=M009W0P

OCCUR=2

OCCUR=2

NODE=M009W0P;LINKAGE=AB
NODE=M009W;LINKAGE=B5
NODE=M009W;LINKAGE=B6

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
150.9 ± 1.7 OUR AVERAGE				Error includes scale factor of 1.1.
122 ± 20		BERTIN 97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
145.7 ± 5.3		WEIDENAUER 93	ASTE	$\bar{p}p \rightarrow \pi^+ \pi^- \omega$
144.9 ± 3.7		DUBNICKA 89	RVUE	π form factor
148 ± 6	1,2	BOHACIK 80	RVUE	
152 ± 9	3	WICKLUND 78	ASPK	3,4,6 $\pi^\pm p N$
154 ± 2	76k	DEUTSCH... 76	HBC	16 $\pi^+ p$
157 ± 8	6.8k	RATCLIFF 72	ASPK	15 $\pi^- p$, $t < 0.3$
143 ± 8	1.7k	REYNOLDS 69	HBC	2.26 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
150.85 ± 0.55 ± 0.67	970k	5 ABLIKIM	18C BES3	$\eta'(958) \rightarrow \gamma \pi^+ \pi^-$
150.18 ± 0.55 ± 0.65	970k	6 ABLIKIM	18C BES3	$\eta'(958) \rightarrow \gamma \pi^+ \pi^-$
147.0 ± 2.5	600k	7 ABELE	99E CBAR	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
146 ± 3	4.9k	8 ADAMS	97 E665	470 $\mu p \rightarrow \mu XB$
160.0 + 4.1 - 4.0		9 CHABAUD	83 ASPK	17 $\pi^- p$ polarized
155 ± 1	10 HEYN	81 RVUE		π form factor
148.0 ± 1.3	1,2 LANG	79 RVUE		
146 ± 14	4.1k	ENGLER 74	DBC	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
143 ± 13	2 ESTABROOKS 74	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
160 ± 10	32k	1 PROTOPOP... 73	HBC	7.1 $\pi^+ p$, $t < 0.4$
145 ± 12	2.2k	3,11 HYAMS 68	OSPK	11.2 $\pi^- p$
163 ± 15	13.3k	12 PISUT	68 RVUE	1.7–3.2 $\pi^- p$, $t < 10$

1 From pole extrapolation.

2 From phase shift analysis of GRAYER 74 data.

3 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

4 Published values contain misprints. Corrected by private communication RATCLIFF 74.

5 From a fit to $\pi^+ \pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.6 From a fit to $\pi^+ \pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.7 Using relativistic Breit-Wigner and taking into account ρ - ω interference.

8 Systematic errors not evaluated.

9 From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P -wave intensity. CHABAUD 83 includes data of GRAYER 74.10 HEYN 81 includes all spacelike and timelike F_π values until 1978.

11 Of HYAMS 68 six parametrizations this is theoretically soundest. MR

12 Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.

$\Gamma_{\rho(770)^0} - \Gamma_{\rho(770)^\pm}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.3 ± 1.3 OUR AVERAGE				Error includes scale factor of 1.4.
-0.2 ± 1.0	1 SCHAEL 05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
3.6 ± 1.8 ± 1.7	1.98M	2 ALOISIO 03	KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.66 ± 0.85	3 BARTOS	17A RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	

1 From the combined fit of the τ^- data from ANDERSON 00A and SCHAEL 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.2 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

3 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

$\Gamma_{\rho(770)^+} - \Gamma_{\rho(770)^-}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.8 ± 2.0 ± 0.5	1.98M	1 ALOISIO	03 KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

1 Without limitations on masses and widths.

NODE=M009W0R

NODE=M009W0R

OCCUR=2

NODE=M009W;LINKAGE=C

NODE=M009W;LINKAGE=H

NODE=M009W;LINKAGE=Z

NODE=M009W;LINKAGE=03

NODE=M009W0R;LINKAGE=B

NODE=M009W0R;LINKAGE=C

NODE=M009W;LINKAGE=BL

NODE=M009W;LINKAGE=A1

NODE=M009W;LINKAGE=G

NODE=M009W;LINKAGE=B

NODE=M009W;LINKAGE=02

NODE=M009W;LINKAGE=R

NODE=M009W6

NODE=M009W6

NODE=M009W6;LINKAGE=SC

NODE=M009W6;LINKAGE=CH

NODE=M009W6;LINKAGE=A

NODE=M009W16

NODE=M009W16

NODE=M009W16;LINKAGE=WO

$\rho(770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \pi\pi$	~ 100	%
$\rho(770)^{\pm}$ decays		
$\Gamma_2 \pi^{\pm}\pi^0$	~ 100	%
$\Gamma_3 \pi^{\pm}\gamma$	(4.5 ± 0.5)	$\times 10^{-4}$ S=2.2
$\Gamma_4 \pi^{\pm}\eta$	< 6	$\times 10^{-3}$ CL=84%
$\Gamma_5 \pi^{\pm}\pi^+\pi^-\pi^0$	< 2.0	$\times 10^{-3}$ CL=84%
$\rho(770)^0$ decays		
$\Gamma_6 \pi^+\pi^-$	~ 100	%
$\Gamma_7 \pi^+\pi^-\gamma$	(9.9 ± 1.6)	$\times 10^{-3}$
$\Gamma_8 \pi^0\gamma$	(4.7 ± 0.6)	$\times 10^{-4}$ S=1.4
$\Gamma_9 \eta\gamma$	(3.00 ± 0.21)	$\times 10^{-4}$
$\Gamma_{10} \pi^0\pi^0\gamma$	(4.5 ± 0.8)	$\times 10^{-5}$
$\Gamma_{11} \mu^+\mu^-$	[a] (4.55 ± 0.28)	$\times 10^{-5}$
$\Gamma_{12} e^+e^-$	[a] (4.72 ± 0.05)	$\times 10^{-5}$
$\Gamma_{13} \pi^+\pi^-\pi^0$	$(1.01^{+0.54}_{-0.36} \pm 0.34)$	$\times 10^{-4}$
$\Gamma_{14} \pi^+\pi^-\pi^+\pi^-$	(1.8 ± 0.9)	$\times 10^{-5}$
$\Gamma_{15} \pi^+\pi^-\pi^0\pi^0$	(1.6 ± 0.8)	$\times 10^{-5}$
$\Gamma_{16} \pi^0e^+e^-$	< 1.2	$\times 10^{-5}$ CL=90%
$\Gamma_{17} \eta e^+e^-$		

[a] The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+\mu^-) = \Gamma(\rho^0 \rightarrow e^+e^-) \times 0.99785$.

NODE=M009225;NODE=M009

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M009;CLUMP=A
 DESIG=11;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=3
 DESIG=5
 DESIG=21

NODE=M009;CLUMP=B
 DESIG=12;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=60
 DESIG=40
 DESIG=8
 DESIG=80
 DESIG=6
 DESIG=4
 DESIG=7;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=22
 DESIG=30
 DESIG=9
 DESIG=10

LINKAGE=MD2

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 10 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.7$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_3 & -100 & \\ \hline \Gamma & 15 & -15 \\ & x_2 & x_3 \end{array}$$

Mode	Rate (MeV)	Scale factor
$\Gamma_2 \pi^{\pm}\pi^0$	150.2 ± 2.4	
$\Gamma_3 \pi^{\pm}\gamma$	0.068 ± 0.007	2.3

DESIG=11
 DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 7 branching ratios uses 22 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 9.5$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

	x_7	x_8	x_9	x_{10}	x_{11}	x_{12}	x_{14}	Γ
	x_6	x_7	x_8	x_9	x_{10}	x_{11}	x_{12}	x_{14}
x_7	-100							
x_8	-4	0						
x_9	-1	0	1					
x_{10}	-1	0	0	0				
x_{11}	2	-3	0	0	0			
x_{12}	0	0	-8	-9	0	0		
x_{14}	-1	0	0	0	0	0	0	
Γ	0	0	4	5	0	0	-54	0

Mode		Rate (MeV)	Scale factor
Γ_6	$\pi^+ \pi^-$	147.5 ± 0.9	DESIG=12
Γ_7	$\pi^+ \pi^- \gamma$	1.48 ± 0.24	DESIG=60
Γ_8	$\pi^0 \gamma$	0.070 ± 0.009	DESIG=40
Γ_9	$\eta \gamma$	0.0447 ± 0.0032	DESIG=8
Γ_{10}	$\pi^0 \pi^0 \gamma$	0.0066 ± 0.0012	DESIG=80
Γ_{11}	$\mu^+ \mu^-$	[a] 0.0068 ± 0.0004	DESIG=6
Γ_{12}	$e^+ e^-$	[a] 0.00704 ± 0.00006	DESIG=4
Γ_{14}	$\pi^+ \pi^- \pi^+ \pi^-$	0.0027 ± 0.0014	DESIG=22

$\rho(770)$ PARTIAL WIDTHS

$\Gamma(\pi^\pm \gamma)$

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
68 ± 7 OUR FIT	Error includes scale factor of 2.3.			
68 ± 7 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.			
81 ± 4 ± 4	CAPRARO	87	SPEC	-
59.8 ± 4.0	HUSTON	86	SPEC	+
71 ± 7	JENSEN	83	SPEC	-
				200 $\pi^- A \rightarrow \pi^- \pi^0 A$
				202 $\pi^+ A \rightarrow \pi^+ \pi^0 A$
				156–260 $\pi^- A \rightarrow \pi^- \pi^0 A$


WEIGHTED AVERAGE
68 ± 7 (Error scaled by 2.2)

Γ_3

NODE=M009230

NODE=M009W3
NODE=M009W3

Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.



$\Gamma(\pi^0\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
77±17±11	36500	¹ ACHASOV 03	SND	0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
121±31		DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

¹ Using $\Gamma_{\text{total}} = 147.9 \pm 1.3$ MeV and $B(\rho \rightarrow \pi^0\gamma)$ from ACHASOV 03.

 Γ_8

NODE=M009W31
NODE=M009W31

 $\Gamma(\eta\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
62±17		¹ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$

¹ Solution corresponding to constructive ω - ρ interference.

 Γ_9

NODE=M009W31;LINKAGE=AV
NODE=M009W32
NODE=M009W32

 $\Gamma(e^+e^-)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.04 ± 0.06 OUR FIT				
7.04 ± 0.06 OUR AVERAGE				
7.048±0.057±0.050	900k	¹ AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
7.06 ± 0.11 ± 0.05	114k	^{2,3} AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
6.77 ± 0.10 ± 0.30		BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.12 ± 0.02 ± 0.11	800k	⁴ ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
6.3 ± 0.1		⁵ BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$

 Γ_{12}

NODE=M009W32;LINKAGE=L
NODE=M009W4
NODE=M009W4

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.8±1.4±0.5	153	AKHMETSHIN 00	CMD2	$0.6\text{--}0.97 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

 Γ_{14}

NODE=M009W4;LINKAGE=AK
NODE=M009W4;LINKAGE=GS
NODE=M009W4;LINKAGE=P2
NODE=M009W4;LINKAGE=AC
NODE=M009W4;LINKAGE=K2

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

2.8±1.4±0.5 153 AKHMETSHIN 00 CMD2 $0.6\text{--}0.97 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M009W33
NODE=M009W33

NODE=M009233

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.876±0.023±0.064	800k	^{1,2} ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.72 ± 0.02		³ BENAYOUN 10	RVUE	$0.4\text{--}1.05 e^+e^-$
1 Supersedes ACHASOV 05A.				
2 A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.				
3 A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.				

NODE=M009G4
NODE=M009G4

NODE=M009G4;LINKAGE=AC
NODE=M009G4;LINKAGE=SN

NODE=M009G4;LINKAGE=BE

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_6/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
1.42±0.10 OUR FIT				

1.45±0.12 OUR AVERAGE

1.32±0.14±0.08	33k	¹ ACHASOV 07B	SND	$0.6\text{--}1.38 e^+e^- \rightarrow \eta\gamma$
1.50±0.65±0.09	17.4k	² AKHMETSHIN 05	CMD2	$0.60\text{--}1.38 e^+e^- \rightarrow \eta\gamma$
1.61±0.20±0.11	23k	^{3,4} AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.85±0.49		⁵ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.05±0.02		⁶ BENAYOUN 10	RVUE	$0.4\text{--}1.05 e^+e^-$
-----------	--	--------------------------	------	---------------------------

NODE=M009G1
NODE=M009G1

NODE=M009G1;LINKAGE=AH

NODE=M009G;LINKAGE=AH
NODE=M009G;LINKAGE=AK
NODE=M009G;LINKAGE=BQ

NODE=M009G;LINKAGE=LP
NODE=M009G1;LINKAGE=BE

¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
⁵ Recalculated by us from the cross section in the peak.
⁶ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
2.22 ± 0.29 OUR FIT				Error includes scale factor of 1.4.
2.22 ± 0.26 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
1.98 ± 0.22 ± 0.10		1 ACHASOV	16A SND	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$
2.90 ± 0.60 ± 0.18	18k	AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$
2.37 ± 0.53 ± 0.33	36k	2 ACHASOV	03 SND	$0.60-0.97 e^+e^- \rightarrow \pi^0\gamma$
3.61 ± 0.74 ± 0.49	10k	3 DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.875 ± 0.026		4 BENAYOUN	10 RVUE	$0.4-1.05 e^+e^-$


¹ From the VMD model with the rho(770), omega(782), phi(1020) resonances, and an additional resonance describing the total contribution of the rho(1450) and omega(1420) states. Supersedes ACHASOV 03.

² Using $\sigma_{\phi \rightarrow \pi^0\gamma}$ from ACHASOV 00 and $m_\rho = 775.97$ MeV in the model with the energy-independent phase of rho-omega interference equal to $(-10.2 \pm 7.0)^\circ$.

³ Recalculated by us from the cross section in the peak.

⁴ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

WEIGHTED AVERAGE
2.22 ± 0.26 (Error scaled by 1.3)



$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ $\Gamma_{12}/\Gamma \times \Gamma_{13}/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.903 ± 0.076		1 BENAYOUN	10 RVUE	$0.4-1.05 e^+e^-$
4.58 ± 2.46 ± 1.56	1.2M	2 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

² Statistical significance is less than 3 σ.

$\rho(770)$ BRANCHING RATIOS

Γ(π±η)/Γ(ππ)					Γ ₄ /Γ ₁
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<60	84	FERBEL	66 HBC	±	$\pi^\pm p$ above 2.5

Γ(π±π+π-π0)/Γ(ππ)					Γ ₅ /Γ ₁
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<20	84	FERBEL	66 HBC	±	$\pi^\pm p$ above 2.5

• • • We do not use the following data for averages, fits, limits, etc. • • •

35 ± 40 JAMES 66 HBC + 2.1 $\pi^+ p$

NODE=M009G2
NODE=M009G2

NODE=M009G2;LINKAGE=B

NODE=M009G;LINKAGE=SH

NODE=M009G2;LINKAGE=LP

NODE=M009G2;LINKAGE=BE

NODE=M009G3
NODE=M009G3

NODE=M009G3;LINKAGE=BE
NODE=M009G3;LINKAGE=AC

NODE=M009235

NODE=M009R4
NODE=M009R4

NODE=M009R1
NODE=M009R1

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.0099±0.0016 OUR FIT					NODE=M009R12 NODE=M009R12
0.0099±0.0016		1 DOLINSKY 91 ND	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0111±0.0014		2 VASSERMAN 88 ND	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		OCCUR=2
<0.005	90	3 VASSERMAN 88 ND	$e^+e^- \rightarrow \pi^+\pi^-\gamma$		NODE=M009R12;LINKAGE=J NODE=M009R12;LINKAGE=I NODE=M009R12;LINKAGE=N
1 Bremsstrahlung from a decay pion and for photon energy above 50 MeV.					
2 Superseded by DOLINSKY 91.					
3 Structure radiation due to quark rearrangement in the decay.					

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.20±0.52		1 ACHASOV 16A SND	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$		
6.21 ^{+1.28} _{-1.18} ±0.39	18k	2,3 AKHMETSHIN 05 CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$		
5.22±1.17±0.75	36k	3,4 ACHASOV 03 SND	$0.60-0.97 e^+e^- \rightarrow \pi^0\gamma$		
6.8 ±1.7		5 BENAYOUN 96 RVUE	$0.54-1.04 e^+e^- \rightarrow \pi^0\gamma$		
7.9 ±2.0		3 DOLINSKY 89 ND	$e^+e^- \rightarrow \pi^0\gamma$		
1 Using $B(\rho \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 03.					
2 Using $B(\rho \rightarrow e^+e^-) = (4.67 \pm 0.09) \times 10^{-5}$.					
3 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.					
4 Using $B(\rho \rightarrow e^+e^-) = (4.54 \pm 0.10) \times 10^{-5}$.					
5 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.					

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	CHG COMMENT	
3.00±0.21 OUR FIT					NODE=M009R7 NODE=M009R7
2.90±0.32 OUR AVERAGE					
2.79±0.34±0.03	33k	1 ACHASOV 07B SND	$0.6-1.38 e^+e^- \rightarrow \eta\gamma$		
3.6 ±0.9		2 ANDREWS 77 CNTR 0	$6.7-10 \gamma\text{Cu}$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.21±1.39±0.20	17.4k	3,4 AKHMETSHIN 05 CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$		
3.39±0.42±0.23		2,5,6 AKHMETSHIN 01B CMD2	$e^+e^- \rightarrow \eta\gamma$		
1.9 ^{+0.6} _{-0.8}		7 BENAYOUN 96 RVUE	$0.54-1.04 e^+e^- \rightarrow \eta\gamma$		
4.0 ±1.1		2,4 DOLINSKY 89 ND	$e^+e^- \rightarrow \eta\gamma$		
1 ACHASOV 07B reports $[\Gamma(\rho(770) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\rho(770) \rightarrow e^+e^-)] = (1.32 \pm 0.14 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\rho(770) \rightarrow e^+e^-) = (4.72 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.					NODE=M009R7;LINKAGE=AO
2 Solution corresponding to constructive $\omega\rho$ interference.					
3 Using $B(\rho \rightarrow e^+e^-) = (4.67 \pm 0.09) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.					
4 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.					
5 The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).					
6 Using $B(\rho \rightarrow e^+e^-) = (4.75 \pm 0.10) \times 10^{-5}$ from AKHMETSHIN 02 and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.					
7 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution. Constructive $\rho\omega$ interference solution.					

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.5±0.8 OUR FIT					
4.5^{+0.9}_{-0.8} OUR AVERAGE					
5.2 ^{+1.5} _{-1.3} ±0.6	190	1 AKHMETSHIN 04B CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$		OCCUR=2
4.1 ^{+1.0} _{-0.9} ±0.3	295	2 ACHASOV 02F SND	$0.36-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.8 ^{+3.4} _{-1.8} ±0.5	63	3 ACHASOV 00G SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$		

¹This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$, and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0\pi^0$ with a branching ratio $(2.0^{+1.1}_{-0.9} \pm 0.3) \times 10^{-5}$ differing from zero by 2.0 standard deviations.

²This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$ and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0\pi^0$ with a branching ratio $(1.9^{+0.9}_{-0.8} \pm 0.4) \times 10^{-5}$ differing from zero by 2.4 standard deviations. Supersedes ACHASOV 00G.

³Superseded by ACHASOV 02F.

$\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-)$

VALUE (units 10^{-5})

4.60 ± 0.28 OUR FIT

4.6 ± 0.2 ± 0.2

ANTIPOV 89 SIGM π^- Cu $\rightarrow \mu^+\mu^-\pi^-$ Cu

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.2 $^{+1.6}_{-3.6}$ ¹ ROTHWELL 69 CNTR Photoproduction

5.6 ± 1.5 ² WEHMANN 69 OSPK 12 π^- C, Fe

9.7 $^{+3.1}_{-3.3}$ ^{3,4} HYAMS 67 OSPK 11 π^- Li, H

Γ_{11}/Γ_6

NODE=M009R14;LINKAGE=AH

¹Possibly large $\rho\omega$ interference leads us to increase the minus error.

²Result contains $11 \pm 11\%$ correction using SU(3) for central value. The error on the correction takes account of possible $\rho\omega$ interference and the upper limit agrees with the upper limit of $\omega \rightarrow \mu^+\mu^-$ from this experiment.

³But he even enlarges his error to take residual ω contamination into account. Since his value is high, seems the other experiments also can't have too many ω 's. But maybe Hyams has additional μ 's from $\rho \rightarrow \pi\pi$, decaying π 's.

⁴HYAMS 67's mass resolution is 20 MeV. The ω region was excluded.

$\Gamma(e^+e^-)/\Gamma(\pi\pi)$

VALUE (units 10^{-4})

0.40 ± 0.05

• • • We do not use the following data for averages, fits, limits, etc. • • •

1,2 BENAKSAS 72 OSPK $e^+e^- \rightarrow \pi^+\pi^-$

¹The ρ' contribution is not taken into account.

²Barkov excludes Auslender and Benaksas for large statistical and systematic errors.

Γ_{12}/Γ_1

NODE=M009R5;LINKAGE=R
NODE=M009R5;LINKAGE=W

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})

1.01 ± 0.54 ± 0.34

<1.2

90

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2M 1 ACHASOV 03D RVUE $0.44^{+2.00}_{-0.36} e^+e^- \rightarrow \pi^+\pi^-\pi^0$

VASSERMAN 88B ND $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹Statistical significance is less than 3σ .

Γ_{13}/Γ

NODE=M009R5;LINKAGE=01

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi\pi)$

VALUE

~0.01

<0.01

84

• • • We do not use the following data for averages, fits, limits, etc. • • •

BRAMON 86 RVUE 0 $J/\psi \rightarrow \omega\pi^0$

¹ABRAMS 71 HBC 0 $3.7\pi^+p$

¹Model dependent, assumes $I = 1, 2$, or 3 for the 3π system.

Γ_{13}/Γ_1

NODE=M009R;LINKAGE=KS
NODE=M009R3;LINKAGE=01

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})

1.8 ± 0.9 OUR FIT

1.8 ± 0.9 ± 0.3

153 AKHMETSHIN 00 CMD2 $0.6-0.97 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20

90

KURDADZE 88 OLYA $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

Γ_{14}/Γ

NODE=M009R6;LINKAGE=NS

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$

CL%

1.8 ± 0.9 ± 0.3

153 AKHMETSHIN 00 CMD2 $0.6-0.97 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<20

90

CHUNG 68 HBC 0 $3.2, 4.2\pi^-p$

<20

90

HUSON 68 HLBC 0 $16.0\pi^-p$

<80

JAMES 66 HBC 0 $2.1\pi^+p$

Γ_{14}/Γ_1

NODE=M009R13
NODE=M009R13

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma(\pi\pi)$

CL%

<15

90

ERBE 69 HBC 0 $2.5-5.8\gamma p$

<20

CHUNG 68 HBC 0 $3.2, 4.2\pi^-p$

<20

90

HUSON 68 HLBC 0 $16.0\pi^-p$

<80

JAMES 66 HBC 0 $2.1\pi^+p$

NODE=M009R11
NODE=M009R11

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$				Γ_{15}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.60±0.74±0.18		1 ACHASOV	09A SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 4	90	AULCHENKO	87C ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
<20	90	KURDADZE	86 OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
1 Assuming no interference between the ρ and ω contributions.				
$\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$				Γ_{16}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	ACHASOV	08 SND	$0.36-0.97 e^+e^- \rightarrow \pi^0e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.6		AKHMETSHIN	05A CMD2	0.72-0.84 e^+e^-
$\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$				Γ_{17}/Γ
VALUE (units 10^{-5})		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7		AKHMETSHIN	05A CMD2	0.72-0.84 e^+e^-

$\rho(770)$ REFERENCES

ABLIKIM	18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>	
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>	
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova	
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 136 442.		
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)
		Translated from ZETF 134 80.		
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 130 437.		
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
		Translated from ZETFP 84 491.		
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETF 128 1201.		
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ALOISIO	05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
		Translated from ZETFP 82 841.		
SCHAEL	05C	PRPL 421 191	S. Schael <i>et al.</i>	(ALEPH Collab.)
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)
SANZ-CILLERO	03	EPJ C27 587	J.J. Sanz-Cillero, A. Pich	
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler	
PICH	01	PR D63 093005	A. Pich, J. Portoles	
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 72 411.		
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
		Translated from ZETFP 71 519.		
AKHMETSHIN	00	PL B475 190	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)
ABELE	99E	PL B469 270	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+) (ZEUS Collab.)
BREITWEG	98B	EPJ C2 247	J. Breitweg <i>et al.</i>	
GARDNER	98	PR D57 2716	S. Gardner, H.B. O'Connell	
Also		PR D62 019903 (errat.)	S. Gardner, H.B. O'Connell	
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ADAMS	97	ZPHY C74 237	M.R. Adams <i>et al.</i>	(E665 Collab.)
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BOGOLYUB...	97	PAN 60 46	M.Y. Bogolyubsky <i>et al.</i>	(MOSU, SERP)
		Translated from YAF 60 53.		

NODE=M009R8
NODE=M009R8NODE=M009R8;LINKAGE=AC
NODE=M009R15
NODE=M009R15

NODE=M009

REFID=58843
REFID=57650
REFID=58325
REFID=58367
REFID=57138
REFID=57513
REFID=56977;ERROR=2
REFID=54274
REFID=54299
REFID=16683
REFID=53212
REFID=58797
REFID=53101
REFID=53136
REFID=52258
REFID=52536
REFID=51942
REFID=51615
REFID=51113
REFID=51133
REFID=51513
REFID=51045
REFID=50330
REFID=50508
REFID=51112
REFID=51060
REFID=50845
REFID=49609
REFID=49610
REFID=49187
REFID=49577
REFID=49404
REFID=49399
REFID=48549
REFID=48816
REFID=48565
REFID=48167
REFID=49180
REFID=48313
REFID=47417
REFID=47882
REFID=47929
REFID=47421
REFID=47468
REFID=47414
REFID=45859
REFID=46354
REFID=46366
REFID=47981
REFID=45415
REFID=45533
REFID=45622
REFID=45701
REFID=45393

O'CONNELL	97	NP A623 559	H.B. O'Connell <i>et al.</i>	(ADLD)	REFID=45860
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
BERNICA	94	PR D50 4454	A. Bernicha, G. Lopez Castro, J. Pestieau	(LOUV+)	REFID=44097
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
AGUILAR... DOLINSKY	91	ZPHY C50 405 PRPL 202 99	M. Aguilar-Benitez <i>et al.</i> S.I. Dolinsky <i>et al.</i>	(LEBC-EHS Collab.) (NOVO)	REFID=41637 REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ANTIPOV	89	ZPHY C42 185	Y.M. Antipov <i>et al.</i>	(SERP, JINR, BGNA+)	REFID=40739
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN... KURDADZE	89 88	ZPHY C45 351 JETPL 47 512	B.V. Geshkenbein L.M. Kurdadze <i>et al.</i>	(ITEP) (NOVO)	REFID=41017 REFID=41121
VASSERMAN	88	Translated from ZETFP 47 432. SJNP 47 1035	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=41019
VASSERMAN	88B	Translated from YAF 47 1635. SJNP 48 480	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=41020
AULCHENKO	87C	Translated from YAF 48 753. IYF 87-90 Preprint	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=41370
CAPRARO	87	NP B288 659	L. Capraro <i>et al.</i>	(CLER, FRAS, MILA+)	REFID=40003
BRAMON	86	PL B173 97	A. Bramon, J. Casulleras	(BARC)	REFID=22102
HUSTON	86	PR D33 3199	J. Huston <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=20137
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
JENSEN	83	PR D27 26	T. Jensen <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=20132
HEYN	81	ZPHY C7 169	M.F. Heyn, C.B. Lang	(GRAZ)	REFID=20129
BOHACIK	80	PR D21 1342	J. Bohacik, H. Kuhnel	(SLOV, WIEN)	REFID=20128
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)	REFID=20126
BARTALUCCI	78	NC 44A 587	S. Bartalucci <i>et al.</i>	(DESY, FRAS)	REFID=20122
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
DEUTSCH... ENGLER	76 74	NP B103 426 PR D10 2070	M. Deutschmann <i>et al.</i> A. Engler <i>et al.</i>	(AACH3, BERL, BONN+) (CMU, CASE)	REFID=20119 REFID=20110
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
RATCLIFF	74	Private Comm.			REFID=40128; ERROR=3
BYERLY	73	PR D7 637	W.L. Byerly <i>et al.</i>	(MICH)	REFID=20104
GLADDING	73	PR D8 3721	G.E. Gladding <i>et al.</i>	(HARV)	REFID=20106
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
PROTOPOP... BALLAM	73 72	PR D7 1279 PR D5 545	S.D. Protopopescu <i>et al.</i> J. Ballam <i>et al.</i>	(LBL)	REFID=20108
BENAKSAS	72	PL 39B 289	D. Benakasas <i>et al.</i>	(SLAC, LBL, TUFTS)	REFID=20094
JACOBS	72	PR D6 1291	L.D. Jacobs	(ORSAY)	REFID=20096
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SACL)	REFID=20101
ABRAMS	71	PR D4 653	G.S. Abrams <i>et al.</i>	(SLAC)	REFID=20102
ALVENSLEB... BIGGS	70 70	PRL 24 786 PRL 24 1197	H. Alvensleben <i>et al.</i> P.J. Biggs <i>et al.</i>	(LBL) (DESY)	REFID=20090 REFID=20085
ERBE	69	PR 188 2060	R. Erbe <i>et al.</i>	(DARE)	REFID=20087
MALAMUD	69	Argonne Conf. 93	R. Erbe <i>et al.</i> (German Bubble Chamber Collab.)	(UCLA)	REFID=20074
REYNOLDS	69	PR 184 1424	E.I. Malamud, P.E. Schlein	(FSU)	REFID=20077
ROTHWELL	69	PRL 23 1521	B.G. Reynolds <i>et al.</i>	(NEAS)	REFID=20080
WEHMANN	69	PR 178 2095	P.L. Rothwell <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20082
ARMENISE	68	NC 54A 999	A.A. Wehmann <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20084
BATON	68	PR 176 1574	N. Armenise <i>et al.</i>	(SACL)	REFID=20054
CHUNG	68	PR 165 1491	J.P. Baton, G. Laurens	(SACL)	REFID=20056
FOSTER	68	NP B6 107	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
GOUNARIS	68	PRL 21 244	M. Foster <i>et al.</i>	(CERN, CDEF)	REFID=20061
HUSON	68	PL 28B 208	G.J. Gounaris, J.J. Sakurai	(CERN, MILA, UCLA)	REFID=48054
HYAMS	68	NP B7 1	R. Huson <i>et al.</i>	(CERN, MPIM)	REFID=20062
LANZEROTTI	68	PR 166 1365	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20063
PISUT	68	NP B6 325	L.J. Lanzerotti <i>et al.</i>	(HARV)	REFID=20068
ASBURY	67B	PRL 19 865	J. Pisut, M. Roos	(CERN)	REFID=20070
BACON	67	PR 157 1263	J.G. Asbury <i>et al.</i>	(DESY, COLU)	REFID=20038
EISNER	67	PR 164 1699	T.C. Bacon <i>et al.</i>	(BNL)	REFID=20039
HUWE	67	PL 24B 252	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
HYAMS	67	PL 24B 634	D.O. Huwe <i>et al.</i>	(COLU)	REFID=20049
MILLER	67B	PR 153 1423	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20050
ALFF.... FERBEL	66 66	PR 145 1072 PL 21 111	D.H. Miller <i>et al.</i> C. Alff-Steinberger <i>et al.</i>	(PURD)	REFID=20051
HAGOPIAN	66	PR 145 1128	T. Ferbel	(COLU, RUTG)	REFID=10762
HAGOPIAN	66B	PR 152 1183	V. Hagopian <i>et al.</i>	(ROCH)	REFID=20028
JACOBS	66B	UCRL 16877	V. Hagopian, Y.L. Pan	(PENN, SACL)	REFID=20030
JAMES	66	PR 142 896	L.D. Jacobs	(PENN, LRL)	REFID=20031
ROSS	66	PR 149 1172	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=20033
SOEDING	66	PL B19 702	M. Ross, L. Stodolsky	(LRL)	REFID=10770
WEST	66	PR 149 1089	P. Soeding	(WISC)	REFID=46380
BLIEDEN	65	PL 19 444	E. West <i>et al.</i>	(CERN MMS Collab.)	REFID=46385
CARMONY	64	PRL 12 254	H.R. Blieden <i>et al.</i>	(UCB)	REFID=20035
GOLDHABER	64	PRL 12 336	D.D. Carmony <i>et al.</i>	(LRL, UCB)	REFID=20578
ABOLINS	63	PRL 11 381	G. Goldhaber <i>et al.</i>	(UCSD)	REFID=20013
			M.A. Abolins <i>et al.</i>	(UCSD)	REFID=20006

$\omega(782)$ $I^G(J^{PC}) = 0^-(1^{--})$

NODE=M001

 $\omega(782)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
782.65±0.12 OUR AVERAGE		Error includes scale factor of 1.9.		See the ideogram below.
783.20±0.13±0.16	18680	AKHMETSHIN 05	CMD2	0.60–1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
782.68±0.09±0.04	11200	¹ AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.79±0.08±0.09	1.2M	² ACHASOV 03D	RVUE	$0.44–2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.7 ± 0.1 ± 1.5	19500	WURZINGER 95	SPEC	1.33 $p d \rightarrow {}^3\text{He} \omega$
781.96±0.17±0.80	11k	³ AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
782.08±0.36±0.82	3463	⁴ AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
781.96±0.13±0.17	15k	AMSLER 93B	CBAR	0.0 $\bar{p} p \rightarrow \omega \pi^0 \pi^0$
782.4 ± 0.2	270k	WEIDENAUER 93	ASTE	$\bar{p} p \rightarrow 2\pi^+ 2\pi^- \pi^0$
782.2 ± 0.4	1488	KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.4 ± 0.5	7000	⁵ KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
781.91±0.24		⁶ LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
781.78±0.10		⁷ BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
783.3 ± 0.4	433	CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.5 ± 0.8	33260	ROOS 80	RVUE	0.0–3.6 $\bar{p} p$
782.6 ± 0.8	3000	BENKHEIRI 79	OMEG	9–12 $\pi^\pm p$
781.8 ± 0.6	1430	COOPER 78B	HBC	0.7–0.8 $\bar{p} p \rightarrow 5\pi$
782.7 ± 0.9	535	VANAPEL...	HBC	7.2 $\bar{p} p \rightarrow \bar{p} p \omega$
783.5 ± 0.8	2100	GESSAROLI 77	HBC	11 $\pi^- p \rightarrow \omega n$
782.5 ± 0.8	418	AGUILAR-...	HBC	3.9, 4.6 $K^- p$
783.4 ± 1.0	248	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K^+ K^- \omega$
781.0 ± 0.6	510	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K_1 K_1 \omega$
783.7 ± 1.0	3583	⁸ COYNE 71	HBC	3.7 $\pi^+ p \rightarrow p \pi^+ \pi^+ \pi^- \pi^0$
784.1 ± 1.2	750	ABRAMOVI...	HBC	3.9 $\pi^- p$
783.2 ± 1.6		⁹ BIGGS 70B	CNTR	<4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
782.4 ± 0.5	2400	BIZZARRI 69	HBC	0.0 $\bar{p} p$

¹ Update of AKHMETSHIN 00C.² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.³ From the $\eta \rightarrow \gamma \gamma$ decay.⁴ From the $\eta \rightarrow 3\pi^0$ decay.⁵ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.⁶ From the $\rho - \omega$ interference in the $\pi^+ \pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.⁷ Systematic uncertainties underestimated.⁸ From best-resolution sample of COYNE 71.⁹ From ω - ρ interference in the $\pi^+ \pi^-$ mass spectrum assuming ω width 12.6 MeV.

NODE=M001M

NODE=M001M

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M001M;LINKAGE=PT

NODE=M001M;LINKAGE=VH

NODE=M001M;LINKAGE=S1

NODE=M001M;LINKAGE=S2


NODE=M001M;LINKAGE=B

NODE=M001M;LINKAGE=LE

NODE=M001M;LINKAGE=KB

NODE=M001M;LINKAGE=D

NODE=M001M;LINKAGE=F



$\omega(782)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.49±0.08 OUR AVERAGE				
8.68±0.23±0.10	11200	1 AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
8.68±0.04±0.15	1.2M	2 ACHASOV 03D	RVUE	$0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
8.2 ± 0.3	19500	WURZINGER 95	SPEC	$1.33 pd \rightarrow {}^3He\omega$
8.4 ± 0.1		3 AULCHENKO 87	ND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
8.30±0.40		BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
9.8 ± 0.9	1488	KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
9.0 ± 0.8	433	CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
9.1 ± 0.8	451	BENAKSAS 72B	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.13±0.45		4 LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
12 ± 2	1430	COOPER 78B	HBC	$0.7-0.8 \bar{p}p \rightarrow 5\pi$
9.4 ± 2.5	2100	GEASSAROLI 77	HBC	$11 \pi^- p \rightarrow \omega n$
10.22±0.43	20000	5 KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
13.3 ± 2	418	AGUILAR-...	72B	HBC $3.9, 4.6 K^- p$
10.5 ± 1.5		BORENSTEIN 72	HBC	$2.18 K^- p$
7.70±0.9 ±1.15	940	BROWN 72	MMS	$2.5 \pi^- p \rightarrow n MM$
10.3 ± 1.4	510	BIZZARRI 71	HBC	$0.0 p\bar{p} \rightarrow K_1 K_1 \omega$
12.8 ± 3.0	248	BIZZARRI 71	HBC	$0.0 p\bar{p} \rightarrow K^+ K^- \omega$
9.5 ± 1.0	3583	COYNE 71	HBC	$3.7 \pi^+ p \rightarrow p\pi^+ \pi^+ \pi^- \pi^0$

1 Update of AKHMETSHIN 00C.

2 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

3 Relativistic Breit-Wigner includes radiative corrections.

4 From the $\rho - \omega$ interference in the $\pi^+ \pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.

5 Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

NODE=M001W

NODE=M001W

OCCUR=2
NODE=M001W;LINKAGE=PT
NODE=M001W;LINKAGE=VH

NODE=M001W;LINKAGE=D
NODE=M001W;LINKAGE=LE

NODE=M001W;LINKAGE=B

NODE=M001215;NODE=M001

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \pi^+ \pi^- \pi^0$	(89.3 ± 0.6) %	
$\Gamma_2 \pi^0 \gamma$	(8.40 ± 0.22) %	S=1.8
$\Gamma_3 \pi^+ \pi^-$	(1.53 ± 0.06) %	
Γ_4 neutrals (excluding $\pi^0 \gamma$)	(7 ± 4) × 10 ⁻³	S=1.1
$\Gamma_5 \eta \gamma$	(4.5 ± 0.4) × 10 ⁻⁴	S=1.1

DESIG=1

DESIG=3

DESIG=2

DESIG=13

DESIG=6

Γ_6	$\pi^0 e^+ e^-$	$(7.7 \pm 0.6) \times 10^{-4}$		DESIG=14
Γ_7	$\pi^0 \mu^+ \mu^-$	$(1.34 \pm 0.18) \times 10^{-4}$	S=1.5	DESIG=11
Γ_8	$\eta e^+ e^-$			DESIG=18
Γ_9	$e^+ e^-$	$(7.36 \pm 0.15) \times 10^{-5}$	S=1.5	DESIG=7
Γ_{10}	$\pi^+ \pi^- \pi^0 \pi^0$	$< 2 \times 10^{-4}$	CL=90%	DESIG=12
Γ_{11}	$\pi^+ \pi^- \gamma$	$< 3.6 \times 10^{-3}$	CL=95%	DESIG=4
Γ_{12}	$\pi^+ \pi^- \pi^+ \pi^-$	$< 1 \times 10^{-3}$	CL=90%	DESIG=15
Γ_{13}	$\pi^0 \pi^0 \gamma$	$(6.7 \pm 1.1) \times 10^{-5}$		DESIG=5
Γ_{14}	$\eta \pi^0 \gamma$	$< 3.3 \times 10^{-5}$	CL=90%	DESIG=17
Γ_{15}	$\mu^+ \mu^-$	$(7.4 \pm 1.8) \times 10^{-5}$		DESIG=8
Γ_{16}	3γ	$< 1.9 \times 10^{-4}$	CL=95%	DESIG=10
Charge conjugation (C) violating modes				
Γ_{17}	$\eta \pi^0$	C $< 2.2 \times 10^{-4}$	CL=90%	NODE=M001;CLUMP=A
Γ_{18}	$2\pi^0$	C $< 2.2 \times 10^{-4}$	CL=90%	DESIG=9
Γ_{19}	$3\pi^0$	C $< 2.3 \times 10^{-4}$	CL=90%	DESIG=193
Γ_{20}	invisible	C $< 7 \times 10^{-5}$	CL=90%	DESIG=16
				DESIG=194

CONSTRAINED FIT INFORMATION

An overall fit to 15 branching ratios uses 55 measurements and one constraint to determine 10 parameters. The overall fit has a $\chi^2 = 57.0$ for 46 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	28											
x_3	-9	-3										
x_4	-95	-55	0									
x_5	7	15	-1	-12								
x_6	-1	0	0	0	0							
x_7	0	0	0	0	0	0						
x_9	-35	-70	3	52	-22	0	0					
x_{13}	1	3	0	-2	0	0	0	0	-2			
x_{15}	0	0	0	0	0	0	0	0	0	0	0	
	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_9	x_{13}			

$\omega(782)$ PARTIAL WIDTHS

$\Gamma(\pi^0\gamma)$	Γ_2			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
880±50	7815	¹ ACHASOV	13	SND 1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
788±12±27	36500	² ACHASOV	03	SND 0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
764±51	10625	DOLINSKY	89	ND $e^+e^- \rightarrow \pi^0\gamma$

¹ Systematic uncertainty not estimated.

² Using $\Gamma_{\gamma\gamma} \equiv 8.44 \pm 0.09$ MeV and $B(\omega \rightarrow \pi^0 \gamma)$ from ACHASOV 03.

$\Gamma(\eta\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_5
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.1 ± 2.5		1 DOLINSKY	89	ND	$e^+e^- \rightarrow \eta\gamma$
¹ Using $\Gamma = 8.4 \pm 0.1$ MeV and $B(\omega \rightarrow \eta\gamma)$ from DOLINSKY 89					

- Using $\Gamma_\omega = 0.4 \pm 0.1$ MeV and $B(\omega \rightarrow \eta\gamma)$ from BULINSKY 89.

NODE=M001218

NODE=M001W1
NODE=M001W1

NODE=M001W1;LINKAGE=AC
NODE=M001W1;LINKAGE=AD

NODE=M001W2
NODE=M001W2

NODE=M001W2;LINKAGE=DA

$\Gamma(e^+e^-)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_9
0.60 ±0.02 OUR EVALUATION					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.591±0.015	11200	1,2 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
0.653±0.003±0.021	1.2M	3 ACHASOV 03D	RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
0.600±0.031	10625	DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$	
1 Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.891 \pm 0.007$ and $\Gamma_{\text{total}} = 8.44 \pm 0.09$ MeV.					
2 Update of AKHMETSHIN 00C.					
3 Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.					

NODE=M001W7

NODE=M001W7

→ UNCHECKED ←

NODE=M001W7;LINKAGE=3P

NODE=M001W7;LINKAGE=PT

NODE=M001W;LINKAGE=VF

NODE=M001225

NODE=M001G2

NODE=M001G2

 $\omega(782) \Gamma(e^+e^-)\Gamma(i)/\Gamma^2(\text{total})$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma \times \Gamma_1/\Gamma$
6.56±0.12 OUR FIT Error includes scale factor of 1.6.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.24±0.11±0.08	11.2k	1 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
6.70±0.06±0.27		AUBERT,B 04N	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$	
6.74±0.04±0.24	1.2M	2,3 ACHASOV 03D	RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
6.37±0.35		2 DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
6.45±0.24		2 BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
5.79±0.42	1488	2 KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
5.89±0.54	433	2 CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
7.54±0.84	451	2 BENAKSAS 72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.20±0.13		4 BENAYOUN 10	RVUE	$0.4-1.05 e^+e^-$	

1 Update of AKHMETSHIN 00C.

2 Recalculated by us from the cross section in the peak.

3 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.4 A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M001G;LINKAGE=PT

NODE=M001G;LINKAGE=LP

NODE=M001G;LINKAGE=VH

NODE=M001G2;LINKAGE=BE

NODE=M001G4

NODE=M001G4

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma \times \Gamma_2/\Gamma$
6.18 ±0.11 OUR FIT Error includes scale factor of 1.6.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.336±0.056±0.089		1 ACHASOV 16A	SND	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$	
6.47 ±0.14 ±0.39	18k	AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$	
6.50 ±0.11 ±0.20	36k	2 ACHASOV 03	SND	$0.60-0.97 e^+e^- \rightarrow \pi^0\gamma$	
6.34 ±0.21 ±0.21	10k	3 DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.80 ±0.13		4 BENAYOUN 10	RVUE	$0.4-1.05 e^+e^-$	

1 From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 03.2 Using $\sigma_{\phi \rightarrow \pi^0\gamma}$ from ACHASOV 00 and $m_\omega = 782.57$ MeV in the model with the energy-independent phase of $\rho\omega$ interference equal to $(-10.2 \pm 7.0)^\circ$.

3 Recalculated by us from the cross section in the peak.

4 A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M001G4;LINKAGE=A

NODE=M001G;LINKAGE=SH

NODE=M001G4;LINKAGE=LP

NODE=M001G4;LINKAGE=BE

NODE=M001G5

NODE=M001G5

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma \times \Gamma_3/\Gamma$
1.225±0.058±0.041					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.166±0.036		2 BENAYOUN 13	RVUE	$0.4-1.05 e^+e^-$	
1.05 ±0.08		3 DAVIER 13	RVUE	$e^+e^- \rightarrow \pi^+\pi^-(\gamma)$	
1 Supersedes ACHASOV 05A.					
2 A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data. Supersedes BENAYOUN 10.					
3 From $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ data of LEES 12G.					

NODE=M001G5;LINKAGE=AC

NODE=M001G5;LINKAGE=B

NODE=M001G5;LINKAGE=A

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$
 $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
3.32±0.28 OUR FIT		Error includes scale factor of 1.1.		
3.18±0.28 OUR AVERAGE				
3.10±0.31±0.11	33k	1 ACHASOV	07B SND	$0.6-1.38 e^+e^- \rightarrow \eta\gamma$
3.17 ^{+1.85} _{-1.31} ±0.21	17.4k	2 AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$
3.41±0.52±0.21	23k	3,4 AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.50±0.10		5 BENAYOUN	10 RVUE	$0.4-1.05 e^+e^-$
1 From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.				
2 From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.				
3 From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.				
4 The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).				
5 A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.				

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$
 $\Gamma_9/\Gamma \times \Gamma_{15}/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±1.8±2.2	4.5M	1 ANASTASI	17 KLOE	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
1 From a fit of the real part of the vacuum polarization by a sum of the leptonic and hadronic contributions, where the hadronic contribution is parametrized as a sum of Breit-Wigner resonances $\omega(782)$, $\phi(1020)$ and using a GOUNARIS 68 parametrization for the $\rho(770)$, and a non-resonant term.				

$\omega(782)$ BRANCHING RATIOS

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$
 Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.9024±0.0019		1 AMBROSINO 08G	KLOE	$1.0-1.03 e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
0.8965±0.0016±0.0048	1.2M	2,3 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.880 ± 0.020 ± 0.032	11200	3,4 AKHMETSHIN 00C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.8942±0.0062		3 DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

1 Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

2 Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

3 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0) / \Gamma_{\text{total}}^2$.

4 Using $\Gamma(e^+e^-) = 0.60 \pm 0.02$ keV.

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$
 Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.88±0.18		1 ACHASOV	16A SND	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$
8.09±0.14		2 AMBROSINO	08G KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
9.06±0.20±0.57	18k	3,4 AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$
9.34±0.15±0.31	36k	4 ACHASOV	03 SND	$0.60-0.97 e^+e^- \rightarrow \pi^0\gamma$
8.65±0.16±0.42	1.2M	5,6 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.39±0.24	9k	7 BENAYOUN	96 RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.88±0.62	10k	4 DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\gamma$

1 Using $B(\omega \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 03.

2 Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

3 Using $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$.

4 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma) / \Gamma_{\text{total}}^2$.

5 Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

6 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0) / \Gamma_{\text{total}}^2$.

7 Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001G3

NODE=M001G3

NODE=M001G3;LINKAGE=AH

NODE=M001G;LINKAGE=AH

NODE=M001G;LINKAGE=AK

NODE=M001G;LINKAGE=BQ

NODE=M001G3;LINKAGE=BE

NODE=M001G01

NODE=M001G01

NODE=M001G01;LINKAGE=A

NODE=M001220

NODE=M001R21

NODE=M001R21

NODE=M001R21

NODE=M001R21;LINKAGE=AM

NODE=M001R;LINKAGE=VF

NODE=M001R;LINKAGE=ZL

NODE=M001R;LINKAGE=KH

NODE=M001R28

NODE=M001R28

NODE=M001R28;LINKAGE=A

NODE=M001R28;LINKAGE=AM

NODE=M001R;LINKAGE=AH

NODE=M001R;LINKAGE=VL

NODE=M001R28;LINKAGE=VF

NODE=M001R28;LINKAGE=ZL

NODE=M001R28;LINKAGE=A1

$\Gamma(\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
9.41±0.23 OUR FIT	Error includes scale factor of 2.0.			
9.05±0.27 OUR AVERAGE	Error includes scale factor of 1.8.			
8.97±0.16	AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$	
9.94±0.36±0.38	1 AULCHENKO 00A	SND	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$	
8.4 ±1.3	KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$	
10.9 ±2.5	BENAKSAS 72C	OSPK	$e^+e^- \rightarrow \pi^0\gamma$	
8.1 ±2.0	BALDIN 71	HLBC	$2.9\pi^+p$	
13 ±4	JACQUET 69B	HLBC	$2.05\pi^+p \rightarrow \pi^+\rho\omega$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.7 ±0.2 ±0.5	2,3 ACHASOV 03D	RVUE	$0.44-2.00\pi^+\pi^-\pi^0$	
9.9 ±0.7	2 DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$	

1 From $\sigma_0^{\omega\pi^0 \rightarrow \pi^0\pi^0\gamma}(m_\phi)/\sigma_0^{\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0\pi^0}(m_\phi)$ with a phase-space correction factor of 1/1.023.

2 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

3 Using ACHASOV 03. Based on 1.2M events.

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
1.53±0.06 OUR FIT					
1.51±0.07 OUR AVERAGE	Error includes scale factor of 1.1.				
1.52±0.08	1 HANHART 18	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$		
1.46±0.12±0.02	900k 2 AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$		
1.30±0.24±0.05	11.2k 3 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$		
2.38 ^{+1.77} _{-0.90} ±0.18	5.4k 4 ACHASOV 02E	SND	$1.1-1.38\pi^+\pi^-\pi^0 \rightarrow \pi^+\pi^-\pi^0$		
2.3 ±0.5	BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$		
1.6 ^{+0.9} _{-0.7}	QUENZER 78	DM1	$e^+e^- \rightarrow \pi^+\pi^-$		
3.6 ±1.9	BENAKSAS 72	OSPK	$e^+e^- \rightarrow \pi^+\pi^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.29±0.22±0.03	970k 5,6 ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$		
1.28±0.22±0.03	970k 7,8 ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$		
1.75±0.11	4.5M 9 ACHASOV 05A	SND	$e^+e^- \rightarrow \pi^+\pi^-$		
2.01±0.29	10 BENAYOUN 03	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$		
1.9 ±0.3	11 GARDNER 99	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$		
2.3 ±0.4	12 BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$		
1.0 ±0.11	13 WICKLUND 78	ASPK	$3,4,6\pi^\pm N$		
1.22±0.30	ALVENSLEB... 71C	CNTR	Photoproduction		
1.3 ^{+1.2} _{-0.9}	MOFFEIT 71	HBC	$2.8,4.7\gamma p$		
0.80 ^{+0.28} _{-0.20}	14 BIGGS 70B	CNTR	$4.2\gamma C \rightarrow \pi^+\pi^- C$		

1 Dispersive analysis. Value extracted from average of data from AUBERT 09AS, AKHMETSHIN 07, ACHASOV 06, AMBROSINO 11A, BABUSCI 13D, ABLIKIM 16B normalised by PDG evaluation for $\Gamma(\omega \rightarrow e^+e^-)$.

2 A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

3 Update of AKHMETSHIN 02.

4 From the $m_{\pi^+\pi^-}$ spectrum taking into account the interference of the $\rho\pi$ and $\omega\pi$ amplitudes.

5 From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.

6 ABLIKIM 18C reports $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.25 \pm 0.21 \pm 0.52) \times 10^{-4}$ which we divide by our best value $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

7 From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.

8 ABLIKIM 18C reports $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.22 \pm 0.21 \pm 0.52) \times 10^{-4}$ which we divide by our best value $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

9 Using $\Gamma(\omega \rightarrow e^+e^-)$ from the 2004 Edition of this Review (PDG 04).

10 Using the data of AKHMETSHIN 02 in the hidden local symmetry model.

11 Using the data of BARKOV 85.

NODE=M001R3
NODE=M001R3

OCCUR=2

OCCUR=3

NODE=M001R15;LINKAGE=D

NODE=M001R15;LINKAGE=AK

NODE=M001R15;LINKAGE=PT

NODE=M001R;LINKAGE=VE

NODE=M001R15;LINKAGE=E

NODE=M001R15;LINKAGE=H

NODE=M001R15;LINKAGE=J

NODE=M001R15;LINKAGE=K

NODE=M001R;LINKAGE=SN

NODE=M001R;LINKAGE=BY

NODE=M001R15;LINKAGE=H4

12 Using the data of BARKOV 85 in the hidden local symmetry model.

13 From a model-dependent analysis assuming complete coherence.

14 Re-evaluated under $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ by BEHREND 71 using more accurate $\omega \rightarrow \rho$ photoproduction cross-section ratio.

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$

See also $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$.

VALUE		DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.0171 ± 0.0007 OUR FIT					
0.026 ± 0.005 OUR AVERAGE					
0.021 +0.028 -0.009	1,2 RATCLIFF	72 ASPK	15 $\pi^- p \rightarrow n 2\pi$		
0.028 ± 0.006	1 BEHREND	71 ASPK	Photoproduction		
0.022 +0.009 -0.01	3 ROOS	70 RVUE			

¹ The fitted width of these data is 160 MeV in agreement with present average, thus the ω contribution is overestimated. Assuming ρ width 145 MeV.

² Significant interference effect observed. NB of $\omega \rightarrow 3\pi$ comes from an extrapolation.

³ ROOS 70 combines ABRAMOVICH 70 and BIZZARRI 70.

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\gamma)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
0.20 ± 0.04	1.98M	1 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	

¹ Using the data of ALOISIO 02D.

$\Gamma(\text{ neutrals})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$(\Gamma_2 + \Gamma_4)/\Gamma$
0.091 ± 0.006 OUR FIT					
0.081 ± 0.011 OUR AVERAGE					
0.075 ± 0.025	BIZZARRI	71 HBC	0.0 $p\bar{p}$		
0.079 ± 0.019	DEINET	69B OSPK	1.5 $\pi^- p$		
0.084 ± 0.015	BOLLINI	68C CNTR	2.1 $\pi^- p$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.073 ± 0.018	42 BASILE	72B CNTR	1.67 $\pi^- p$		

$\Gamma(\text{ neutrals})/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$(\Gamma_2 + \Gamma_4)/\Gamma_1$
0.102 ± 0.008 OUR FIT					
0.103 +0.011 -0.010 OUR AVERAGE					
0.15 ± 0.04	46 AGUILAR-...	72B HBC	3.9,4.6 $K^- p$		
0.10 ± 0.03	19 BARASH	67B HBC	0.0 $\bar{p}p$		
0.134 ± 0.026	850 DIGIUGNO	66B CNTR	1.4 $\pi^- p$		
0.097 ± 0.016	348 FLATTE	66 HBC	1.4 – 1.7 $K^- p \rightarrow \Lambda M\bar{M}$		
0.06 +0.05 -0.02	JAMES	66 HBC	2.1 $\pi^+ p$		
0.08 ± 0.03	35 KRAEMER	64 DBC	1.2 $\pi^+ d$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ± 0.02	20 BUSCHBECK	63 HBC	1.5 $K^- p$		

$\Gamma(\pi^0\gamma)/\Gamma(\text{ neutrals})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/(\Gamma_2 + \Gamma_4)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.78 ± 0.07 ¹ DAKIN 72 OSPK 1.4 $\pi^- p \rightarrow n M\bar{M}$

>0.81 90 DEINET 69B OSPK

¹ Error statistical only. Authors obtain good fit also assuming $\pi^0\gamma$ as the only neutral decay.

$\Gamma(\text{ neutrals})/\Gamma(\text{ charged particles})$

VALUE		DOCUMENT ID	TECN	COMMENT	$(\Gamma_2 + \Gamma_4)/(\Gamma_1 + \Gamma_3)$
0.100 ± 0.008 OUR FIT					
0.124 ± 0.021	FELDMAN	67C OSPK	1.2 $\pi^- p$		

NODE=M001R15;LINKAGE=Q

NODE=M001R15;LINKAGE=F

NODE=M001R15;LINKAGE=B

NODE=M001R2

NODE=M001R2

NODE=M001R2

NODE=M001R2;LINKAGE=A

NODE=M001R2;LINKAGE=S

NODE=M001R2;LINKAGE=R

NODE=M001R33

NODE=M001R33

NODE=M001R;LINKAGE=KL

NODE=M001R14

NODE=M001R14

NODE=M001R1

NODE=M001R1

NODE=M001R18

NODE=M001R18

NODE=M001R18;LINKAGE=D

NODE=M001R9

NODE=M001R9

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$	Γ_5/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
4.5 ± 0.4 OUR FIT	Error includes scale factor of 1.1.
6.3 ± 1.3 OUR AVERAGE	Error includes scale factor of 1.2.
6.6 ± 1.7	¹ ABELE 97E CBAR 0.0 $\bar{p}p \rightarrow 5\gamma$
8.3 ± 2.1	ALDE 93 GAM2 $38\pi^- p \rightarrow \omega n$
3.0 ± 2.5	² ANDREWS 77 CNTR 6.7–10 γCu
• • • We do not use the following data for averages, fits, limits, etc. • • •	
4.2 ± 0.4 ± 0.1	33k ³ ACHASOV 07B SND 0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
4.44 ± 2.59 ± 0.28	17.4k ^{4,5} AKHMETSHIN 05 CMD2 0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
5.10 ± 0.72 ± 0.34	23k ⁶ AKHMETSHIN 01B CMD2 $e^+e^- \rightarrow \eta\gamma$
0.7 to 5.5	⁷ CASE 00 CBAR 0.0 $p\bar{p} \rightarrow \eta\eta\gamma$
6.56 ± 2.41 ± 2.55	3525 ^{2,8} BENAYOUN 96 RVUE $e^+e^- \rightarrow \eta\gamma$
7.3 ± 2.9	^{2,4} DOLINSKY 89 ND $e^+e^- \rightarrow \eta\gamma$
1 No flat $\eta\eta\gamma$ background assumed.	
2 Solution corresponding to constructive ω - ρ interference.	
3 ACHASOV 07B reports $[\Gamma(\omega(782) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow e^+e^-)] = (3.10 \pm 0.31 \pm 0.11) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow e^+e^-) = (7.36 \pm 0.15) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.	
4 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.	
5 Using $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.	
6 Using $B(\omega \rightarrow e^+e^-) = (7.07 \pm 0.19) \times 10^{-5}$ and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$. Solution corresponding to constructive ω - ρ interference. The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively). Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.	
7 Depending on the degree of coherence with the flat $\eta\eta\gamma$ background and using $B(\omega \rightarrow \pi^0\gamma) = (8.5 \pm 0.5) \times 10^{-2}$.	
8 Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.	

NODE=M001R19
NODE=M001R19

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$	Γ_5/Γ_2
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.0098 ± 0.0024	¹ ALDE 93 GAM2 $38\pi^- p \rightarrow \omega n$
0.0082 ± 0.0033	² DOLINSKY 89 ND $e^+e^- \rightarrow \eta\gamma$
0.010 ± 0.045	APEL 72B OSPK 4–8 $\pi^- p \rightarrow n3\gamma$


NODE=M001R;LINKAGE=EA
NODE=M001R19;LINKAGE=A
NODE=M001R19;LINKAGE=AO

$\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
7.7 ± 0.6 OUR FIT	
7.7 ± 0.6 OUR AVERAGE	
7.61 ± 0.53 ± 0.64	ACHASOV 08 SND 0.36–0.97 $e^+e^- \rightarrow \pi^0e^+e^-$
8.19 ± 0.71 ± 0.62	AKHMETSHIN 05A CMD2 0.72–0.84 e^+e^-
5.9 ± 1.9	⁴³ DOLINSKY 88 ND $e^+e^- \rightarrow \pi^0e^+e^-$

NODE=M001R11;LINKAGE=A
NODE=M001R11;LINKAGE=K

$\Gamma(\pi^0\mu^+\mu^-)/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.34 ± 0.18 OUR FIT	Error includes scale factor of 1.5.
1.34 ± 0.19 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.
1.41 ± 0.09 ± 0.15	ARNALDI 16 NA60 400 GeV (p - A) collisions
1.72 ± 0.25 ± 0.14	3k ARNALDI 09 NA60 158A In–In collisions
0.96 ± 0.23	DZHELYADIN 81B CNTR 25–33 $\pi^- p \rightarrow \omega n$

NODE=M001R12
NODE=M001R12



$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$	Γ_8/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
<1.1	AKHMETSHIN 05A CMD2 0.72-0.84 $e^+ e^-$

NODE=M001R34
NODE=M001R34

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$	Γ_9/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.736±0.015 OUR FIT Error includes scale factor of 1.5.	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
0.700±0.016	11200 1,2 AKHMETSHIN 04 CMD2 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.752±0.004±0.024	1.2M 2,3 ACHASOV 03D RVUE $0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.714±0.036	2 DOLINSKY 89 ND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.72 ± 0.03	2 BARKOV 87 CMD $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.64 ± 0.04	2 KURDADZE 83B OLYA $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.675±0.069	433 2 CORDIER 80 DM1 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.83 ± 0.10	451 2 BENAKSAS 72B OSPK $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.77 ± 0.06	4 AUGUSTIN 69D OSPK $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.65 ± 0.13	33 5 ASTVACAT... 68 OSPK Assume SU(3)+mixing

NODE=M001R13
NODE=M001R13

- 1 Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = 0.891 \pm 0.007$. Update of AKHMETSHIN 00C.
- 2 Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}^2$.
- 3 Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+ \pi^-) = (1.70 \pm 0.28)\%$.
- 4 Rescaled by us to correspond to ω width 8.4 MeV. Systematic errors underestimated.
- 5 Not resolved from ρ decay. Error statistical only.

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$	Γ_{10}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
< 2 90 ACHASOV 09A SND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
<200 90 KURDADZE 86 OLYA $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	

NODE=M001R13;LINKAGE=4P
NODE=M001R13;LINKAGE=ZL
NODE=M001R13;LINKAGE=VF
NODE=M001R13;LINKAGE=E
NODE=M001R13;LINKAGE=A

$\Gamma(\pi^+ \pi^- \gamma)/\Gamma_{\text{total}}$	Γ_{11}/Γ
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<0.0036 95 WEIDENAUER 90 ASTE $p\bar{p} \rightarrow \pi^+ \pi^- \pi^+ \pi^- \gamma$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
<0.004 95 BITYUKOV 88B SPEC $32 \pi^- p \rightarrow \pi^+ \pi^- \gamma X$	

NODE=M001R22
NODE=M001R22

$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{11}/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.066	90	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$	NODE=M001R4
<0.05	90	FLATTE	66	$1.2 - 1.7 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$	NODE=M001R4

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{12}/Γ
$<1 \times 10^{-3}$	90	KURDADZE	88	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$	NODE=M001R24 NODE=M001R24

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ
6.7 ± 1.1 OUR FIT					NODE=M001R29 NODE=M001R29
6.5 ± 1.2 OUR AVERAGE					

$6.4^{+2.4}_{-2.0} \pm 0.8$	190	1 AKHMETSHIN 04B	CMD2	$0.6 - 0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$6.6^{+1.4}_{-1.3} \pm 0.6$	295	ACHASOV 02F	SND	$0.36 - 0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$11.8^{+2.1}_{-1.9} \pm 1.4$	190	2 AKHMETSHIN 04B	CMD2	$0.6 - 0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2
$7.8 \pm 2.7 \pm 2.0$	63	1,3 ACHASOV 00G	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2
$12.7 \pm 2.3 \pm 2.5$	63	2,3 ACHASOV 00G	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	

1 In the model assuming the $\rho \rightarrow \pi^0\pi^0\gamma$ decay via the $\omega\pi$ and $f_0(500)\gamma$ mechanisms.

2 In the model assuming the $\rho \rightarrow \pi^0\pi^0\gamma$ decay via the $\omega\pi$ mechanism only.

3 Superseded by ACHASOV 02F.

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ_1
<0.00045	90	DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	NODE=M001R10 NODE=M001R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.08	95	JACQUET 69B	HLBC	$2.05 \pi^+ p \rightarrow \pi^+ p\omega$
-------	----	-------------	------	--

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^0\gamma)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ_2
7.9 ± 1.3 OUR FIT						NODE=M001R7 NODE=M001R7

8.5 ± 2.9 40 ± 14 ALDE 94B GAM2 $38\pi^- p \rightarrow \pi^0\pi^0\gamma n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 50	90	DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
<1800	95	KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
<1500	90	BENAKSAS 72C	OSPK	e^+e^-
<1400		BALDIN 71	HLBC	$2.9 \pi^+ p$
<1000	90	BARMIN 64	HLBC	$1.3 - 2.8 \pi^- p$

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\text{ neutrals})$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{13}/(\Gamma_2 + \Gamma_4)$
--------------	------------	--------------------	-------------	----------------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22 \pm 0.07		1 DAKIN 72	OSPK	$1.4 \pi^- p \rightarrow n\text{MM}$
<0.19	90	DEINET 69B	OSPK	

1 See $\Gamma(\pi^0\gamma)/\Gamma(\text{ neutrals})$.

 $\Gamma(\eta\pi^0\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{14}/Γ
<3.3	90	AKHMETSHIN 04B	CMD2	$0.6 - 0.97 e^+e^- \rightarrow \eta\pi^0\gamma$	NODE=M001R32 NODE=M001R32

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{15}/Γ
7.4 ± 1.8 OUR FIT					NODE=M001R30 NODE=M001R30
7.4 ± 1.8 OUR AVERAGE					

6.6 \pm 1.4 \pm 1.7	4.5M	1 ANASTASI 17	KLOE	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
9.0 \pm 2.9 \pm 1.1	18	HEISTER 02C	ALEP	$Z \rightarrow \mu^+\mu^- + X$

1 Assuming lepton universality in the decay $\omega \rightarrow \ell^+\ell^-$ and correcting for different phase space between electron and muon final states.

$\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{15}/Γ_1
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.2	90	WILSON 69	OSPK	$12 \pi^- C \rightarrow Fe$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.7	74	FLATTE 66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \mu^+ \mu^-$	
<1.2		BARBARO-...	65	HBC	$2.7 K^- p$

$\Gamma(\pi^0\mu^+\mu^-)/\Gamma(\mu^+\mu^-)$					Γ_7/Γ_{15}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.2 ± 0.6	30	¹ DZHELYADIN 79	CNTR	$25-33 \pi^- p$	

¹ Superseded by DZHELYADIN 81B result above.

$\Gamma(3\gamma)/\Gamma_{\text{total}}$					Γ_{16}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.9	95	¹ ABELE 97E	CBAR	$0.0 \bar{p}p \rightarrow 5\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2	90	¹ PROKOSHKIN 95	GAM2	$38 \pi^- p \rightarrow 3\gamma n$	

¹ From direct 3γ decay search.

$\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{17}/Γ
Violates C conservation.					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.001	90	ALDE 94B	GAM2	$38\pi^- p \rightarrow \eta\pi^0 n$	

$[\Gamma(\eta\gamma) + \Gamma(\eta\pi^0)]/\Gamma(\pi^+\pi^-\pi^0)$					$(\Gamma_5 + \Gamma_{17})/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.016	90	¹ FLATTE 66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \pi^+ \pi^- MM$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

<0.045 95 JACQUET 69B HLBC $2.05 \pi^+ p \rightarrow \pi^+ p\omega$

¹ Restated by us using $B(\eta \rightarrow \text{charged modes}) = 29.2\%$.

$\Gamma(\eta\pi^0)/\Gamma(\pi^0\gamma)$					Γ_{17}/Γ_2
Violates C conservation.					
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.6	90	¹ STAROSTIN 09	CRYM	$\gamma p \rightarrow \eta\pi^0 p$	

1 STAROSTIN 09 reports $[\Gamma(\omega(782) \rightarrow \eta\pi^0)/\Gamma(\omega(782) \rightarrow \pi^0\gamma)] \times [B(\eta \rightarrow 2\gamma)] < 1.01 \times 10^{-3}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

$\Gamma(2\pi^0)/\Gamma(\pi^0\gamma)$					Γ_{18}/Γ_2
Violates C conservation and Bose-Einstein statistics.					
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.59	90	STAROSTIN 09	CRYM	$\gamma p \rightarrow 2\pi^0 p$	

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$					Γ_{19}/Γ
Violates C conservation.					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3 $\times 10^{-4}$	90	PROKOSHKIN 95	GAM2	$38 \pi^- p \rightarrow 3\pi^0 n$	

$\Gamma(3\pi^0)/\Gamma(\pi^0\gamma)$					Γ_{19}/Γ_2
Violates C conservation.					
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.72	90	STAROSTIN 09	CRYM	$\gamma p \rightarrow 3\pi^0 p$	

$\Gamma(3\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{19}/Γ_1
Violates C conservation.					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.009	90	BARBERIS 01	450 pp	$\rightarrow p_f 3\pi^0 p_s$	

NODE=M001R6
NODE=M001R6

NODE=M001R20
NODE=M001R20

NODE=M001R27;LINKAGE=S
NODE=M001R27
NODE=M001R27

NODE=M001R8
NODE=M001R8

NODE=M001R35;LINKAGE=ST
NODE=M001R35
NODE=M001R35
NODE=M001R35

NODE=M001R36
NODE=M001R36
NODE=M001R36

NODE=M001R26
NODE=M001R26
NODE=M001R26

NODE=M001R37
NODE=M001R37
NODE=M001R37

NODE=M001R31
NODE=M001R31
NODE=M001R31

$\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ_1
$<8.1 \times 10^{-5}$	90	ABLIKIM	18S	BES3	$J/\psi \rightarrow \omega\eta \rightarrow \omega\pi^+\pi^-\pi^0$

PARAMETER Λ IN $\omega \rightarrow \pi^0\ell^+\ell^-$ DECAY

In the pole approximation the electromagnetic transition form factor for a resonance of mass M is given by the expression:

$$|F|^2 = (1 - M^2/\Lambda^2)^{-2},$$

where for the parameter Λ vector dominance predicts $\Lambda = M_p \approx 0.770$ GeV. The ARNALDI 09 measurement is in obvious conflict with this expectation. Note that for $\eta \rightarrow \gamma\mu^+\mu^-$ decay ARNALDI 09 and DZHELYADIN 80 obtain the value of Λ consistent with vector dominance.

PARAMETER Λ IN $\omega \rightarrow \pi^0\mu^+\mu^-$ DECAY

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ_1
0.670 ± 0.006 OUR AVERAGE					
$0.6707 \pm 0.0039 \pm 0.0056$	1	ARNALDI	16	NA60 400 GeV (p -A) collisions	
$0.668 \pm 0.009 \pm 0.003$	3k	ARNALDI	09	NA60 158A In-In collisions	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.65 ± 0.03		DZHELYADIN	81B	CNTR 25–33 $\pi^- p \rightarrow \omega n$	
1 ARNALDI 16 reports $\Lambda^{-2}(\omega) = 2.223 \pm 0.026 \pm 0.037$ GeV $^{-2}$ which we converted to the quoted Λ value.					NODE=M001LAM;LINKAGE=A
2 ARNALDI 09 reports $\Lambda^{-2}(\omega) = 2.24 \pm 0.06 \pm 0.02$ GeV $^{-2}$ which we converted to the quoted Λ value.					NODE=M001LAM;LINKAGE=B

PARAMETER Λ IN $\omega \rightarrow \pi^0e^+e^-$ DECAY

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ_1
0.709 ± 0.037	1.1k	1 ADLARSON	17B	A2MM $\gamma p \rightarrow \omega p$	
1 ADLARSON 17B reports $\Lambda^{-2}(\omega\pi^0) = 1.99 \pm 0.21$ GeV $^{-2}$ that we converted to the quoted Λ value.					NODE=M001A02;LINKAGE=A

ENERGY DEPENDENCE OF $\omega \rightarrow \pi^+\pi^-\pi^0$ DALITZ PLOT

The following experiments fit to one or more of the coefficients α , β , γ for $|\text{matrix element}|^2 \propto P(1 + 2\alpha Z + 2\beta Z^{3/2} \sin(3\phi) + 2\gamma Z^2 + O(Z^{5/2}))$ where P is the P -wave phase-space factor and Z , ϕ are kinematical variables as defined in ADLARSON 17.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ_1
0.133 ± 0.008 OUR AVERAGE					
$0.1321 \pm 0.0067 \pm 0.0046$	260k	1 ABLIKIM	18AD BES3	$J/\psi \rightarrow \omega\eta$	
0.147 ± 0.036	44k	ADLARSON	17 WASA	α in $pd \rightarrow {}^3\text{He} \omega$, $pp \rightarrow pp\omega$	

1 Keeping a term linear in Z only. A fit with the terms proportional to Z and $Z^{3/2}$ gives $\alpha = 0.133 \pm 0.041$ and $\beta = 0.037 \pm 0.054$.

 $\omega(782)$ REFERENCES

ABLIKIM	18AD	PR D98 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59498
ABLIKIM	18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58971
HANHART	18	EPJ C78 450	C. Hanhart <i>et al.</i>		REFID=59186
ADLARSON	17	PL B770 418	P. Adlarson <i>et al.</i>	(WASA-at-COSY Collab.)	REFID=57907
ADLARSON	17B	PR C95 035208	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)	REFID=58155
ANASTASI	17	PL B767 485	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)	REFID=57887
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57126
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57513
ARNALDI	16	PL B757 437	R. Arnaldi <i>et al.</i>	(NA60 Collab.)	REFID=57220
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=4
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
BABUSCI	13D	PL B720 336	D. Babusci <i>et al.</i>	(CATA, CALB, BARI)	REFID=55337
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono (PARIN, BERLIN+)		REFID=55357
DAVIER	13	EPJ C73 2597	M. Davier <i>et al.</i>		REFID=55499
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54305
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53101
		Translated from ZETF 136, 442.			
ARNALDI	09	PL B677 260	R. Arnaldi <i>et al.</i>	(NA60 Collab.)	REFID=52720
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
STAROSTIN	09	PR C79 065201	A. Starostin <i>et al.</i>	(Crystal Ball Collab. at MAMI)	REFID=53001
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52258
		Translated from ZETF 134, 80.			

AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52573
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51513
		Translated from ZETFP 84 491.			
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51045
		Translated from ZETF 128 1201.			
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
		Translated from ZETFP 82 841.			
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49610
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49187
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
BENAYOUN	03	EPJ C29 397	M. Benayoun <i>et al.</i>		REFID=49477
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48816
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48565
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48564
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48324
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
		Translated from ZETFP 72 411.			
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47929
		Translated from ZETFP 71 519.			
AKHMETSHIN	00C	PL B476 33	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47423
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
		Translated from ZETFP 117 1067.			
CASE	00	PR D61 032002	T. Case <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47409
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
GARDNER	99	PR D59 076002	S. Gardner, H.B. O'Connell		REFID=46919
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)	REFID=45859
ABELE	97E	PL B411 361	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45755
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
PROKOSHKIN	95	PD 40 273	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=44616
		Translated from DANS 342 610.			
WURZINGER	95	PR C51 443	R. Würzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=45209
ALDE	94B	PL B340 122	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=44100
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
ALDE	93	PAN 56 1229	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43603
		Translated from YAF 56 137.			
Also		ZPHY C61 35	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43790
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43602
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
WEIDENAUER	90	ZPHY C47 353	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=41368
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
BITYUKOV	88B	SJNP 47 800	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=41021
		Translated from YAF 47 1258.			
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41022
		Translated from YAF 48 442.			
KURDADZE	88	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=41121
		Translated from ZETFP 47 432.			
AULCHENKO	87	PL B186 432	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=40007
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
		Translated from ZETFP 43 497.			
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
KURDADZE	83B	JETPL 36 274	A.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20244
		Translated from ZETFP 36 221.			
DZHELYADIN	81B	PL 102B 296	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20242
CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
DZHELYADIN	80	PL 94B 548	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10831
ROOS	80	LNC 27 321	M. Roos, A. Pellinen	(HELS)	REFID=20241
BENKHIERI	79	NP B150 268	P. Benkhieri <i>et al.</i>	(EPOL, CERN, CDEF+)	REFID=20238
DZHELYADIN	79	PL 84B 143	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20239
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
QUENZER	78	PL 76B 512	A. Quenzer <i>et al.</i>	(LALO)	REFID=20123
VANAPEL...	78	NP B133 245	G.W. van Apeldoorn <i>et al.</i>	(ZEEM)	REFID=20234
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
KEYNE	76	PR D14 28	J. Keyne <i>et al.</i>	(LOIC, SHMP)	REFID=20226
		PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72B	PL 41B 234	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20206
BASILE	72B	Phil. Conf. 153	M. Basile <i>et al.</i>	(CERN)	REFID=20207
BENAKSAS	72	PL 39B 289	D. Benakas <i>et al.</i>	(ORSAY)	REFID=20096
BENAKSAS	72B	PL 42B 507	D. Benakas <i>et al.</i>	(ORSAY)	REFID=20209
BENAKSAS	72C	PL 42B 511	D. Benakas <i>et al.</i>	(ORSAY)	REFID=20517
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
BROWN	72	PL 42B 117	R.M. Brown <i>et al.</i>	(ILL, ILLC)	REFID=20211
DAKIN	72	PR D6 2321	J.T. Dakin <i>et al.</i>	(PRIN)	REFID=20212
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)	REFID=20102
ALVENSLEB...	71C	PRL 27 888	H. Alvensleben <i>et al.</i>	(DESY)	REFID=20193
BALDIN	71	SJNP 13 758	A.B. Baldin <i>et al.</i>	(ITEP)	REFID=20195
		Translated from YAF 13 1318.			

BEHREND	71	PRL 27 61	H.J. Behrend <i>et al.</i>	(ROCH, CORN, FNAL)	REFID=20197
BIZZARRI	71	NP B27 140	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20198
COYNE	71	NP B32 333	D.G. Coyne <i>et al.</i>	(LRL)	REFID=20201
MOFFEIT	71	NP B29 349	K.C. Moffeit <i>et al.</i>	(LRL, UCB, SLAC+)	REFID=20204
ABRAMOVI...	70	NP B20 209	M. Abramovich <i>et al.</i>	(CERN)	REFID=20180
BIGGS	70B	PRL 24 1201	P.J. Biggs <i>et al.</i>	(DARE)	REFID=20184
BIZZARRI	70	PRL 25 1385	R. Bizzarri <i>et al.</i>	(ROMA, SYRA)	REFID=20181
ROOS	70	DNPL/R7 173	M. Roos	(CERN)	REFID=20191
Proc. Daresbury Study Weekend No. 1.					
AUGUSTIN	69D	PL 28B 513	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=20169
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20171
DEINET	69B	PL 30B 426	W. Deinet <i>et al.</i>	(KARL, CERN)	REFID=20173
JACQUET	69B	NC 63A 743	F. Jacquet <i>et al.</i>	(EPOL, BERG)	REFID=20176
WILSON	69	Private Comm.	R. Wilson	(HARV)	REFID=20179
Also		PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20084
ASTVACAT...	68	PL 27B 45	R.G. Astvatsaturov <i>et al.</i>	(JINR, MOSU)	REFID=20055
BOLLINI	68C	NC 56A 531	D. Bollini <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20164
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>		REFID=20160
FELDMAN	67C	PR 159 1219	M. Feldman <i>et al.</i>	(PENN)	REFID=20161
DIGIUGNO	66B	NC 44A 1272	G. Di Giugno <i>et al.</i>	(NAPL, FRAS, TRST)	REFID=20156
FLATTE	66	PR 145 1050	S.M. Flatte <i>et al.</i>	(LRL)	REFID=20157
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
BARBARO...	65	PRL 14 279	A. Barbaro-Galtieri, R.D. Tripp	(LRL)	REFID=20152
BARMIN	64	JETP 18 1289	V.V. Barmin <i>et al.</i>	(ITEP)	REFID=20149
		Translated from ZETF 45 1879.			
KRAEMER	64	PR 136 B496	R.W. Kraemer <i>et al.</i>	(JHU, NWES, WOOD)	REFID=10755
BUSCHBECK	63	Siena Conf. 1 166	B. Buschbeck <i>et al.</i>	(VIEN, CERN, ANIK)	REFID=20146

 $\eta'(958)$

$I^G(J^{PC}) = 0^+(0^-+)$

 $\eta'(958)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
957.78 ±0.06 OUR AVERAGE					
957.793 ±0.054 ±0.036	3.9k	LIBBY	08	CLEO $J/\psi \rightarrow \gamma\eta'$	
957.9 ±0.2 ±0.6	4800	WURZINGER	96	SPEC $1.68 pd \rightarrow {}^3He\eta'$	
957.46 ±0.33		DUANE	74	MMS $\pi^- p \rightarrow nMM$	
958.2 ±0.5	1414	DANBURG	73	HBC $2.2 K^- p \rightarrow \Lambda\eta'$	
958 ±1	400	JACOBS	73	HBC $2.9 K^- p \rightarrow \Lambda\eta'$	
956.1 ±1.1	3415	¹ BASILE	71	CNTR $1.6 \pi^- p \rightarrow nn'$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
957.5 ±0.2		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
959 ±1	630	² BELADIDZE	92c	VES $36 \pi^- Be \rightarrow \pi^-\eta'{}^3He$	
958 ±1	340	² ARMSTRONG	91B	OMEG $300 pp \rightarrow pp\eta\pi^+\pi^-$	
958.2 ±0.4	622	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
957.8 ±0.2	2420	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
956.3 ±1.0	143	² GIDAL	87	MRK2 $e^+ e^- \rightarrow e^+ e^-\eta\pi^+\pi^-$	OCCUR=2
957.4 ±1.4	535	³ BASILE	71	CNTR $1.6 \pi^- p \rightarrow nn'$	OCCUR=2
957 ±1		RITTENBERG	69	HBC $1.7-2.7 K^- p$	

1 Using all η' decays.

2 Systematic uncertainty not estimated.

3 Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement. **$\eta'(958)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.188±0.006 OUR FIT					
[0.196 ± 0.009 MeV OUR 2019 FIT]					
0.230±0.021 OUR AVERAGE					
0.226 ±0.017 ±0.014	2300	CZERWINSKI	10	MMS $pp \rightarrow pp\eta'$	
0.40 ±0.22	4800	WURZINGER	96	SPEC $1.68 pd \rightarrow {}^3He\eta'$	
0.28 ±0.10	1000	BINNIE	79	MMS 0 $\pi^- p \rightarrow nMM$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.20 ±0.04		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	

NODE=M002

NODE=M002M

NODE=M002M

OCCUR=2

OCCUR=2

NODE=M002M;LINKAGE=BS

NODE=M002M;LINKAGE=NS

NODE=M002M;LINKAGE=BA

NODE=M002W

NODE=M002W

NEW

η' (958) DECAY MODES

NODE=M002215;NODE=M002

Mode	Fraction (Γ_i/Γ)	Confidence level	
$\Gamma_1 \pi^+ \pi^- \eta$	(42.5 \pm 0.5) %		DESIG=1
$\Gamma_2 \rho^0 \gamma$ (including non-resonant $\pi^+ \pi^- \gamma$)	(29.5 \pm 0.4) %		DESIG=9
$\Gamma_3 \rho^0 \gamma$			DESIG=213
$\Gamma_4 \pi^0 \pi^0 \eta$	(22.4 \pm 0.5) %		DESIG=2
$\Gamma_5 \omega \gamma$	(2.52 \pm 0.07) %		DESIG=7
$\Gamma_6 \omega e^+ e^-$	(2.0 \pm 0.4) $\times 10^{-4}$		DESIG=205
$\Gamma_7 \gamma \gamma$	(2.307 \pm 0.033) %		DESIG=6
$\Gamma_8 3\pi^0$	(2.50 \pm 0.17) $\times 10^{-3}$		DESIG=8
$\Gamma_9 \mu^+ \mu^- \gamma$	(1.13 \pm 0.28) $\times 10^{-4}$		DESIG=20
$\Gamma_{10} \pi^+ \pi^- \mu^+ \mu^-$	< 2.9 $\times 10^{-5}$	90%	DESIG=201
$\Gamma_{11} \pi^+ \pi^- \pi^0$	(3.61 \pm 0.17) $\times 10^{-3}$		DESIG=121
$\Gamma_{12} (\pi^+ \pi^- \pi^0)$ S-wave	(3.8 \pm 0.5) $\times 10^{-3}$		DESIG=211
$\Gamma_{13} \pi^\mp \rho^\pm$	(7.4 \pm 2.3) $\times 10^{-4}$		DESIG=210
$\Gamma_{14} \pi^0 \rho^0$	< 4 %	90%	DESIG=18
$\Gamma_{15} 2(\pi^+ \pi^-)$	(8.4 \pm 0.9) $\times 10^{-5}$		DESIG=131
$\Gamma_{16} \pi^+ \pi^- 2\pi^0$	(1.8 \pm 0.4) $\times 10^{-4}$		DESIG=202
$\Gamma_{17} 2(\pi^+ \pi^-)$ neutrals	< 1 %	95%	DESIG=132
$\Gamma_{18} 2(\pi^+ \pi^-) \pi^0$	< 1.8 $\times 10^{-3}$	90%	DESIG=141
$\Gamma_{19} 2(\pi^+ \pi^-) 2\pi^0$	< 1 %	95%	DESIG=15
$\Gamma_{20} 3(\pi^+ \pi^-)$	< 3.1 $\times 10^{-5}$	90%	DESIG=203
$\Gamma_{21} K^\pm \pi^\mp$	< 4 $\times 10^{-5}$	90%	DESIG=207
$\Gamma_{22} \pi^+ \pi^- e^+ e^-$	(2.4 \pm 1.3) $\times 10^{-3}$		DESIG=10
$\Gamma_{23} \pi^+ e^- \nu_e + \text{c.c.}$	< 2.1 $\times 10^{-4}$	90%	DESIG=204
$\Gamma_{24} \gamma e^+ e^-$	(4.91 \pm 0.27) $\times 10^{-4}$		DESIG=28
$\Gamma_{25} \pi^0 \gamma \gamma$	(3.20 \pm 0.24) $\times 10^{-3}$		DESIG=24
$\Gamma_{26} \pi^0 \gamma \gamma$ (non resonant)	(6.2 \pm 0.9) $\times 10^{-4}$		DESIG=212
$\Gamma_{27} \eta \gamma \gamma$	< 1.33 $\times 10^{-4}$	90%	DESIG=214
$\Gamma_{28} 4\pi^0$	< 3.2 $\times 10^{-4}$	90%	DESIG=26
$\Gamma_{29} e^+ e^-$	< 5.6 $\times 10^{-9}$	90%	DESIG=150
Γ_{30} invisible	< 6 $\times 10^{-4}$	90%	DESIG=200

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes

NODE=M002;CLUMP=B

$\Gamma_{31} \pi^+ \pi^-$	P,CP	< 1.8	$\times 10^{-5}$	90%	DESIG=111
$\Gamma_{32} \pi^0 \pi^0$	P,CP	< 4	$\times 10^{-4}$	90%	DESIG=25
$\Gamma_{33} \pi^0 e^+ e^-$	C	[a] < 1.4	$\times 10^{-3}$	90%	DESIG=16
$\Gamma_{34} \eta e^+ e^-$	C	[a] < 2.4	$\times 10^{-3}$	90%	DESIG=17
$\Gamma_{35} 3\gamma$	C	< 1.0	$\times 10^{-4}$	90%	DESIG=23
$\Gamma_{36} \mu^+ \mu^- \pi^0$	C	[a] < 6.0	$\times 10^{-5}$	90%	DESIG=22
$\Gamma_{37} \mu^+ \mu^- \eta$	C	[a] < 1.5	$\times 10^{-5}$	90%	DESIG=21
$\Gamma_{38} e \mu$	LF	< 4.7	$\times 10^{-4}$	90%	DESIG=27

[a] C parity forbids this to occur as a single-photon process.

LINKAGE=CS

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 19 branching ratios uses 51 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 69.4$ for 43 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

	x_2	x_4	x_5	x_7	x_8	x_{11}	x_{22}	
	x_1	x_2	x_4	x_5	x_7	x_8	x_{11}	x_{22}
x_2	-24							
x_4	-74	-42						
x_5	-7	-6	-2					
x_7	-11	-7	9	-1				
x_8	-17	-9	19	0	2			
x_{11}	-1	-1	-1	0	0	0		
x_{22}	-9	-7	-7	-1	-2	-2	0	
Γ	11	-10	-1	1	-40	0	0	2

Mode	Rate (MeV)	
$\Gamma_1 \pi^+ \pi^- \eta$	0.0799 ± 0.0029	DESIG=1
$\Gamma_2 \rho^0 \gamma$ (including non-resonant $\pi^+ \pi^- \gamma$)	0.0554 ± 0.0019	DESIG=9
$\Gamma_4 \pi^0 \pi^0 \eta$	0.0421 ± 0.0017	DESIG=2
$\Gamma_5 \omega \gamma$	0.00474 ± 0.00020	DESIG=7
$\Gamma_7 \gamma \gamma$	0.00434 ± 0.00013	DESIG=6
$\Gamma_8 3\pi^0$	(4.7 ± 0.4) $\times 10^{-4}$	DESIG=8
$\Gamma_{11} \pi^+ \pi^- \pi^0$	(6.8 ± 0.4) $\times 10^{-4}$	DESIG=121
$\Gamma_{22} \pi^+ \pi^- e^+ e^-$	(4.4 ± 2.3) $\times 10^{-4}$	DESIG=10

$\eta'(958)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$		Γ_7
VALUE (keV)	EVTS	DOCUMENT ID
4.34 ± 0.14 OUR FIT		
[4.36 ± 0.14 keV OUR 2019 FIT]		
4.28 ± 0.19 OUR AVERAGE		
4.17 ± 0.10 ± 0.27	2000	¹ ACCIARRI
4.53 ± 0.29 ± 0.51	266	KARCH
3.61 ± 0.13 ± 0.48		² BEHREND
4.6 ± 1.1 ± 0.6	23	BARU
4.57 ± 0.25 ± 0.44		BUTLER
5.08 ± 0.24 ± 0.71	547	³ ROE
3.8 ± 0.7 ± 0.6	34	AIHARA
4.9 ± 0.5 ± 0.5	136	⁴ WILLIAMS
• • • We do not use the following data for averages, fits, limits, etc. • • •		
4.7 ± 0.6 ± 0.9	143	⁵ GIDAL
4.0 ± 0.9		⁶ BARTEL
• • • We do not use the following data for averages, fits, limits, etc. • • •		

¹ No non-resonant $\pi^+ \pi^-$ contribution found.

² Reevaluated by us using $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$.

³ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁴ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁵ Superseded by BUTLER 90.

⁶ Systematic error not evaluated.

NODE=M002220

NODE=M002W4

NODE=M002W4

NEW

NODE=M002W4;LINKAGE=AC

NODE=M002W4;LINKAGE=K1

NODE=M002W4;LINKAGE=K2

NODE=M002W4;LINKAGE=K3

NODE=M002W4;LINKAGE=C

NODE=M002W4;LINKAGE=A

$\Gamma(e^+e^-)$					Γ_{29}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.1 \times 10^{-3}$	90	1,2 ACHASOV	15	SND 0.958 $e^+e^- \rightarrow \pi\pi\eta$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<2.0 \times 10^{-3}$	90	2 ACHASOV	15	SND 0.958 $e^+e^- \rightarrow \pi\pi\eta$	
$<2.4 \times 10^{-3}$	90	2 AKHMETSHIN	15	CMD3 0.958 $e^+e^- \rightarrow \pi^+\pi^-\eta$	
1 Combining data of ACHASOV 15 and AKHMETSHIN 15.					
2 Using η and η' branching fractions from PDG 14.					

$\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

$\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_2/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.28±0.04 OUR FIT					
[1.26 ± 0.04 keV OUR 2019 FIT]					
1.26±0.07 OUR AVERAGE				Error includes scale factor of 1.2.	
$1.09 \pm 0.04 \pm 0.13$		BEHREND 91	CELL	$e^+e^- \rightarrow e^+e^-\rho(770)^0\gamma$	
$1.35 \pm 0.09 \pm 0.21$		AIHARA 87	TPC	$e^+e^- \rightarrow e^+e^-\rho\gamma$	
$1.13 \pm 0.04 \pm 0.13$	867	ALBRECHT 87B	ARG	$e^+e^- \rightarrow e^+e^-\rho\gamma$	
$1.53 \pm 0.09 \pm 0.21$		ALTHOFF 84E	TASS	$e^+e^- \rightarrow e^+e^-\rho\gamma$	
$1.14 \pm 0.08 \pm 0.11$	243	BERGER 84B	PLUT	$e^+e^- \rightarrow e^+e^-\rho\gamma$	
$1.73 \pm 0.34 \pm 0.35$	95	JENNI 83	MRK2	$e^+e^- \rightarrow e^+e^-\rho\gamma$	
$1.49 \pm 0.13 \pm 0.027$	213	BARTEL 82B	JADE	$e^+e^- \rightarrow e^+e^-\rho\gamma$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$1.85 \pm 0.31 \pm 0.24$	43	BEHREND 82C	CELL	$e^+e^- \rightarrow e^+e^-\rho\gamma$	

$\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_4/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
0.97±0.04 OUR FIT			Error includes scale factor of 1.1. [1.00 ± 0.05 keV OUR 2019 FIT]		
0.92±0.06±0.11	1 KARCH 92	CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$0.95 \pm 0.05 \pm 0.08$	2 KARCH 90	CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$		
$1.00 \pm 0.08 \pm 0.10$	2,3 ANTREASYAN 87	CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$		
1 Reevaluated by us using $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$. Supersedes ANTREASYAN 87 and KARCH 90.					NODE=M002G2;LINKAGE=K4
2 Superseded by KARCH 92.					NODE=M002G2;LINKAGE=A
3 Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.					NODE=M002G2;LINKAGE=D

$\eta'(958) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_{29}/\Gamma$
VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	1 AKHMETSHIN 15	CMD3	$0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$	
1 AKHMETSHIN 15 reports [$\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta) \times \Gamma(\eta'(958) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] $\times [B(\eta \rightarrow 2\gamma)] < 4.1 \times 10^{-4}$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.					

$\eta'(958) \text{ BRANCHING RATIOS}$

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
42.5 ± 0.5 OUR FIT				Error includes scale factor of 1.1. [0.426 ± 0.007 OUR 2019 FIT]	
41.24±0.08±1.24	312k	ABLIKIM 19T	BES	$J/\psi \rightarrow \gamma\eta'$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
42.4 ± 1.1 ± 0.4	1.2k	1 PEDLAR	09	CLEO $J/\psi \rightarrow \gamma\eta'$	
1 Not independent of other η' branching fractions and ratios in PEDLAR 09.					NODE=M002R47;LINKAGE=PE

NODE=M002W1
NODE=M002W1

OCCUR=2

NODE=M002W1;LINKAGE=A
NODE=M002W1;LINKAGE=B

NODE=M002223

NODE=M002223

NODE=M002G1
NODE=M002G1

NEW

NODE=M002G2
NODE=M002G2

NEW

NODE=M002G2;LINKAGE=K4

NODE=M002G2;LINKAGE=A

NODE=M002G2;LINKAGE=D

NODE=M002224

NODE=M002G01
NODE=M002G01

NODE=M002G01;LINKAGE=A

NODE=M002230

NODE=M002R47
NODE=M002R47

NEW

$\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{\text{total}}$
 $0.2810\Gamma_1/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1194±0.0015 OUR FIT		Error includes scale factor of 1.1. [0.1196 ± 0.0019 OUR 2019 FIT]		

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123 ± 0.014	107	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
0.10 ± 0.04	10	LONDON	66	HBC	2.24 $K^- p \rightarrow \Lambda 2\pi^+ 2\pi^- \pi^0$
0.07 ± 0.04	7	BADIER	65B	HBC	3 $K^- p$

 $\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{\text{total}}$
 $0.7212\Gamma_1/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.306±0.004 OUR FIT		Error includes scale factor of 1.1. [0.307 ± 0.005 OUR 2019 FIT]		

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.314±0.026	281	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
-------------	-----	------------	----	-----	-----------------

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$
 Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
29.5 ± 0.4 OUR FIT		Error includes scale factor of 1.1. [0.289 ± 0.005 OUR 2019 FIT]		

29.9 ± 0.6 OUR AVERAGE [0.319 ± 0.030 OUR 2009 AVERAGE]

29.90±0.03±0.55	913k	ABLIKIM	19T	BES	$J/\psi \rightarrow \gamma\eta'$
------------------------	------	---------	-----	-----	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

28.7 ± 0.7 ± 0.4	0.2k	¹ PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
32.9 ± 3.3	298	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
20 ± 10	20	LONDON	66	HBC	2.24 $K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$
34 ± 9	35	BADIER	65B	HBC	3 $K^- p$

¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

 $\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$
 Γ_3/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

33.34±0.06±1.60	970k	¹ ABLIKIM	18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
34.43±0.52±1.97	970k	² ABLIKIM	18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$

¹ From a fit to $\pi^+\pi^-$ mass using $\rho(770)$, $\omega(782)$, and box anomaly components.

² From a fit to $\pi^+\pi^-$ mass using $\rho(770)$, $\omega(782)$, and $\rho(1450)$ components.

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta)$
 Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.694±0.014 OUR FIT		Error includes scale factor of 1.1. [0.679 ± 0.017 OUR 2019 FIT]	

0.683±0.020 OUR AVERAGE

0.677±0.024±0.011		PEDLAR	09	CLE3	$J/\psi \rightarrow \eta'\gamma$
0.69 ± 0.03		ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta'\gamma$

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))$
 $\Gamma_2/0.714\Gamma_1$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.972±0.020 OUR FIT		Error includes scale factor of 1.1. [0.951 ± 0.024 OUR 2019 FIT]		

0.97 ± 0.09 OUR AVERAGE

0.70 ± 0.22		AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\eta$
1.07 ± 0.17		BELADIDZE	92C	VES	$36 \pi^- \text{Be} \rightarrow \pi^-\eta' \eta \text{Be}$
0.92 ± 0.14	473	DANBURG	73	HBC	$2.2 K^- p \rightarrow \Lambda X^0$
1.11 ± 0.18	192	JACOBS	73	HBC	$2.9 K^- p \rightarrow \Lambda X^0$

 $\Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$
 Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
22.4 ± 0.6 OUR FIT		Error includes scale factor of 1.1. [0.228 ± 0.008 OUR 2019 FIT]		

21.36±0.10±0.92	52k	ABLIKIM	19T	BES	$J/\psi \rightarrow \gamma\eta'$
------------------------	-----	---------	-----	-----	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

23.5 ± 1.3 ± 0.4	3.2k	¹ PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
------------------	------	---------------------	----	------	----------------------------------

¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R3

NODE=M002R3

NEW

NODE=M002R1

NODE=M002R1

NEW

NODE=M002R6

NODE=M002R6

NEW

OCCUR=2

NODE=M002R66;LINKAGE=A

NODE=M002R66;LINKAGE=B

NODE=M002R43

NODE=M002R43

NEW

NODE=M002R27

NODE=M002R27

NEW

NODE=M002R48

NODE=M002R48

NEW

NODE=M002R48;LINKAGE=PE

$\Gamma(\pi^0\pi^0\eta(3\pi^0\text{decay}))/\Gamma_{\text{total}}$
 $0.321\Gamma_4/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0718±0.0018 OUR FIT	Error includes scale factor of 1.1.	[0.0733 ± 0.0026 OUR 2019 FIT]		

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.11 ± 0.06 4 BENSINGER 70 DBC 2.2 $\pi^+ p$

 $\Gamma(\pi^0\pi^0\eta)/\Gamma(\pi^+\pi^-\eta)$
 Γ_4/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.527±0.019 OUR FIT	Error includes scale factor of 1.1.	[0.536 ± 0.026 OUR 2019 FIT]	

0.555±0.043±0.013 PEDLAR 09 CLE3 $J/\psi \rightarrow \eta'\gamma$

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi\pi\eta)$
 $\Gamma_2/(\Gamma_1+\Gamma_4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.454±0.009 OUR FIT	Error includes scale factor of 1.1.	[0.442 ± 0.012 OUR 2019 FIT]	

0.43 ±0.02 ±0.02 BARBERIS 98C OMEG 450 $p p \rightarrow p_f \eta' p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.15 DAVIS 68 HBC 5.5 $K^- p$

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$
 Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.52 ±0.07 OUR FIT	$[(2.62 \pm 0.13) \times 10^{-2}$ OUR 2019 FIT]			

2.50 ±0.07 OUR AVERAGE

$[(2.55 \pm 0.16) \times 10^{-2}$ OUR 2019 AVERAGE]

2.489 ± 0.018 ± 0.074 23k ABLIKIM 19T BES $J/\psi \rightarrow \gamma\eta'$
 2.55 ± 0.03 ± 0.16 33.2k 1 ABLIKIM 15AD BES3 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.34 ± 0.30 ± 0.04 70 2 PEDLAR 09 CLEO $J/\psi \rightarrow \gamma\eta'$

1 Using $B(J/\psi \rightarrow \eta'\gamma) = (5.15 \pm 0.16) \times 10^{-3}$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

2 Not independent of other η' branching fractions and ratios in PEDLAR 09.

 $\Gamma(\omega\gamma)/\Gamma(\pi^+\pi^-\eta)$
 Γ_5/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0593±0.0018 OUR FIT	Error includes scale factor of 1.1.	[0.0615 ± 0.0033 OUR 2019 FIT]		

0.055 ±0.007 ±0.001 PEDLAR 09 CLE3 $J/\psi \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.068 ± 0.013 68 ZANFINO 77 ASPK 8.4 $\pi^- p$

 $\Gamma(\omega\gamma)/\Gamma(\pi^0\pi^0\eta)$
 Γ_5/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.113±0.004 OUR FIT	$[0.115 \pm 0.007$ OUR 2019 FIT]		

0.147±0.016 ALDE 87B GAM2 38 $\pi^- p \rightarrow n4\gamma$

 $\Gamma(\omega e^+e^-)/\Gamma(\omega\gamma)$
 Γ_6/Γ_5

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

7.71 ± 1.34 ± 0.54 1 ABLIKIM 15AD BES3 $J/\psi \rightarrow \eta'\gamma$

1 Obtained from other ABLIKIM 15AD measurements with common systematics taken into account

 $\Gamma(\omega e^+e^-)/\Gamma_{\text{total}}$
 Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.97±0.34±0.17	66	1 ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta'\gamma$

1 Using $B(J/\psi \rightarrow \eta'\gamma) = (5.15 \pm 0.16) \times 10^{-3}$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/[\Gamma(\pi^+\pi^-\eta) + \Gamma(\pi^0\pi^0\eta) +$
 $\Gamma(\omega\gamma)]$
 $\Gamma_2/(\Gamma_1+\Gamma_4+\Gamma_5)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.437±0.008 OUR FIT	Error includes scale factor of 1.1.	[0.425 ± 0.011 OUR 2019 FIT]	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ± 0.14 DAUBER 64 HBC 1.95 $K^- p$

NODE=M002R26

NODE=M002R26

NEW

NODE=M002R45

NODE=M002R45

NEW

NODE=M002R7

NODE=M002R7

NEW

NODE=M002R49

NODE=M002R49

NEW

NODE=M002R49;LINKAGE=A

NODE=M002R49;LINKAGE=PE

NODE=M002R17

NODE=M002R17

NEW

NODE=M002R33

NODE=M002R33

NEW

NODE=M002R60

NODE=M002R60

NODE=M002R60;LINKAGE=A

NODE=M002R59

NODE=M002R59

NODE=M002R59;LINKAGE=A

NODE=M002R18

NODE=M002R18

NEW

$$\frac{[\Gamma(\pi^0\pi^0\eta(\text{charged decay})) + \Gamma(\omega(\text{charged decay})\gamma)]/\Gamma_{\text{total}}}{(0.286\Gamma_4+0.89\Gamma_5)/\Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0864±0.0017 OUR FIT	Error includes scale factor of 1.1. [0.0886 ± 0.0026 OUR 2019 FIT]			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ± 0.029 42 RITTENBERG 69 HBC 1.7–2.7 $K^- p$

$$\frac{\Gamma(\pi^+\pi^-\text{ neutrals})/\Gamma_{\text{total}}}{(0.714\Gamma_1+0.286\Gamma_4+0.89\Gamma_5)/\Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3897±0.0028 OUR FIT	Error includes scale factor of 1.1. [0.3926 ± 0.0035 OUR 2019 FIT]			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 ± 0.1 39 LONDON 66 HBC 2.24 $K^- p \rightarrow \Lambda\pi^+\pi^-\text{ neutrals}$
0.35 ± 0.06 33 BADIER 65B HBC 3 $K^- p$

$$\frac{\Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\Gamma_7/\Gamma}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.307±0.035 OUR FIT	Error includes scale factor of 1.1. [(2.22 ± 0.08) × 10 $^{-2}$ OUR 2019 FIT]			

2.31 ± 0.06 OUR AVERAGE Error includes scale factor of 1.8. [(2.00 ± 0.15) × 10 $^{-2}$ OUR 2019 AVERAGE]

2.331±0.012±0.035	71k	ABLIKIM	19T	BES	$J/\psi \rightarrow \gamma\eta'$
1.99 $^{+0.31}_{-0.27}$ ± 0.07	114	¹ WICHT	08	BELL	$B^\pm \rightarrow K^\pm\gamma\gamma$
2.00 ± 0.18		² STANTON	80	SPEC	$8.45\pi^- p \rightarrow n\pi^+\pi^- 2\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.25 ± 0.16 ± 0.03	0.3k	³ PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
1.8 ± 0.2	6000	⁴ APEL	79	NICE	$15\text{--}40\pi^- p \rightarrow n2\gamma$
2.5 ± 0.7		DUANE	74	MMS	$\pi^- p \rightarrow n\text{MM}$
1.71 ± 0.33	68	DALPIAZ	72	CNTR	$1.6\pi^- p \rightarrow nX^0$
2.0 $^{+0.8}_{-0.6}$	31	HARVEY	71	OSPK	$3.65\pi^- p \rightarrow nX^0$

¹WICHT 08 reports $[\Gamma(\eta'(958) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta'K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta'K^+) = (7.04 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Includes APEL 79 result.

³Not independent of other η' branching fractions and ratios in PEDLAR 09.

⁴Data is included in STANTON 80 evaluation.

$$\frac{\Gamma(\gamma\gamma)/\Gamma(\pi^+\pi^-\eta)}{\Gamma_7/\Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0543±0.0012 OUR FIT	Error includes scale factor of 1.1. [0.0522 ± 0.0022 OUR 2019 FIT]		

0.053 ± 0.004 ± 0.001 PEDLAR 09 CLE3 $J/\psi \rightarrow \eta'\gamma$

$$\frac{\Gamma(\gamma\gamma)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))}{\Gamma_7/\Gamma_2}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0783±0.0016 OUR FIT	Error includes scale factor of 1.1. [0.0768 ± 0.0033 OUR 2019 FIT]		

0.080 ± 0.008 ABLIKIM 06E BES2 $J/\psi \rightarrow \eta'\gamma$

$$\frac{\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)}{\Gamma_7/\Gamma_4}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.1031±0.0028 OUR FIT [0.097 ± 0.004 OUR 2019 FIT]			

0.105 ± 0.010 OUR AVERAGE Error includes scale factor of 1.9.

0.091 ± 0.009	AMSLER	93	CBAR	0.0 $\bar{p}p$
0.112 ± 0.002 ± 0.006	ALDE	87B	GAM2	$38\pi^- p \rightarrow n2\gamma$

$$\frac{\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta(\text{neutral decay}))}{\Gamma_7/0.714\Gamma_4}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.144±0.004 OUR FIT [0.136 ± 0.006 OUR 2019 FIT]				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.188±0.058 16 APEL 72 OSPK 3.8 $\pi^- p \rightarrow nX^0$

NODE=M002R4

NODE=M002R4

NEW

NODE=M002R2

NODE=M002R2

NEW

NODE=M002R19

NODE=M002R19

NEW

NODE=M002R19;LINKAGE=WI

NODE=M002R19;LINKAGE=S

NODE=M002R19;LINKAGE=PE

NODE=M002R19;LINKAGE=A

NODE=M002R46

NODE=M002R46

NEW

NODE=M002R42

NODE=M002R42

NEW

NODE=M002R38

NODE=M002R38

NEW

NODE=M002R28

NODE=M002R28

NEW

$\Gamma(\text{neutrals})/\Gamma_{\text{total}}$		$(0.714\Gamma_4 + 0.09\Gamma_5 + \Gamma_7)/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.185 ± 0.004 OUR FIT		Error includes scale factor of 1.1.	[0.188 ± 0.006 OUR 2019 FIT]	[0.188 ± 0.006 OUR 2019 FIT]	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.185 ± 0.022	535	BASILE	71	CNTR	$1.6\pi^- p \rightarrow nX^0$
0.189 ± 0.026	123	RITTENBERG	69	HBC	1.7–2.7 $K^- p$

$\Gamma(3\pi^0)/\Gamma_{\text{total}}$		Γ_8/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.50 ± 0.17 OUR FIT					
[(2.54 ± 0.18) × 10 ⁻³ OUR 2019 FIT]					
3.57 ± 0.26 OUR AVERAGE					
3.522 ± 0.082 ± 0.254	2015	ABLIKIM	17	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
4.79 ± 0.59 ± 1.14	183	¹ ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.56 ± 0.22 ± 0.34	309	² ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$

¹We have added all systematic uncertainties in quadrature to a single value.
²Superseded by ABLIKIM 17.

$\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$		Γ_8/Γ_4			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
112 ± 8 OUR FIT					
[(111 ± 8) × 10 ⁻⁴ OUR 2019 FIT]					
78 ± 10 OUR AVERAGE					
86 ± 19	235	BLIK	08	GAMS	$32\pi^- p \rightarrow \eta' n$
74 ± 15		ALDE	87B	GAM2	$38\pi^- p \rightarrow n6\gamma$
75 ± 18		BINON	84	GAM2	$30\text{--}40\pi^- p \rightarrow n6\gamma$

$\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$		Γ_9/Γ_7			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.9 ± 1.2	33	VIKTOROV	80	CNTR	$25,33\pi^- p \rightarrow 2\mu\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$		Γ_{10}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.29	90	¹ ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
<2.4	90	² NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$

1 Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

2 Not independent of measured value of Γ_{10}/Γ_1 from NAIK 09.

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$		Γ_{10}/Γ_1			
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.5	90	¹ NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
1 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.					

$\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$		Γ_{10}/Γ_2			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$		Γ_{11}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.61 ± 0.18 OUR FIT					
3.61 ± 0.18 OUR AVERAGE					
3.591 ± 0.054 ± 0.174	6067	ABLIKIM	17	BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
4.28 ± 0.49 ± 1.11	78	¹ ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
3.7 ^{+1.1} _{-0.9} ± 0.4		² NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

3.83 ± 0.15 ± 0.39 1014 ³ABLIKIM 12E BES3 $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$

1 We have added all systematic uncertainties in quadrature to a single value.

2 Not independent of measured value of Γ_{11}/Γ_1 from NAIK 09.

3 Superseded by ABLIKIM 17.

NODE=M002R5

NODE=M002R5

NEW

NODE=M002R55;LINKAGE=A
NODE=M002R55;LINKAGE=B

NODE=M002R32
NODE=M002R32

NEW

NODE=M002R50
NODE=M002R50

NODE=M002R57
NODE=M002R57

NODE=M002R21;LINKAGE=A
NODE=M002R21;LINKAGE=NA
NODE=M002R21;LINKAGE=B

$\Gamma((\pi^+\pi^-\pi^0) \text{ S-wave})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$37.63 \pm 0.77 \pm 5.00$	6580	1 ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$

¹We have added all systematic uncertainties in quadrature .

 $\Gamma(\pi^\mp\rho^\pm)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.44 \pm 0.60 \pm 2.23$	1231	1 ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^\mp\rho^\pm)$

¹We have added all systematic uncertainties in quadrature .

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5 ± 0.4 OUR FIT	Error includes scale factor of 1.1.			$[(8.5 \pm 0.4) \times 10^{-3}$ OUR 2019 FIT]

 $8.28^{+2.49}_{-2.12} \pm 0.04$ 20 1 NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

¹NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (21^{+6}_{-5} \pm 2) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^0\rho^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	RITTENBERG 65	HBC	$2.7 K^- p$

 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$8.4 \pm 0.9 \pm 0.1$	199	1 ABLIKIM	14M BES3	$J/\psi \rightarrow \gamma\eta'$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 24	90	2 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<1000	90	RITTENBERG 69	HBC	$1.7\text{--}2.7 K^- p$

¹ABLIKIM 14M reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] = (4.40 \pm 0.35 \pm 0.30) \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Not independent of measured value of Γ_{15}/Γ_1 from NAIK 09.

 $\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.6	90	1 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

¹NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.79 \pm 0.38 \pm 0.02$	84	1 ABLIKIM	14M BES3	$J/\psi \rightarrow \gamma\eta'$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<27	90	2 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
-----	----	--------	---------	----------------------------------

¹ABLIKIM 14M reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] = (9.38 \pm 1.79 \pm 0.89) \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

²Not independent of measured value of Γ_{16}/Γ_1 from NAIK 09.

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	1 NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

¹NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 15 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$.

 Γ_{12}/Γ

NODE=M002R63
NODE=M002R63

NODE=M002R63;LINKAGE=A

 Γ_{13}/Γ

NODE=M002R62
NODE=M002R62

NODE=M002R62;LINKAGE=A

 Γ_{11}/Γ_1

NODE=M002R01
NODE=M002R01

NEW

NODE=M002R01;LINKAGE=NA

 Γ_{14}/Γ

NODE=M002R10
NODE=M002R10

 Γ_{15}/Γ

NODE=M002R24
NODE=M002R24

NODE=M002R24;LINKAGE=A

NODE=M002R24;LINKAGE=NA

 Γ_{15}/Γ_1

NODE=M002R04
NODE=M002R04

NODE=M002R04;LINKAGE=NA

 Γ_{16}/Γ

NODE=M002R51
NODE=M002R51

NODE=M002R51;LINKAGE=A

NODE=M002R51;LINKAGE=NA

 Γ_{16}/Γ_1

NODE=M002R05
NODE=M002R05

NODE=M002R05;LINKAGE=NA

$\Gamma(2(\pi^+\pi^-)\text{ neutrals})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
<0.01	95	DANBURG 73	HBC	2.2 $K^- p \rightarrow \Lambda X^0$	NODE=M002R22 NODE=M002R22
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$	

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.002	90	¹ NAIK 09	CLEO	$J/\psi \rightarrow \gamma\eta'$	NODE=M002R23 NODE=M002R23
<0.01	90	RITTENBERG 69	HBC	1.7–2.7 $K^- p$	

¹ Not independent of measured value of Γ_{18}/Γ_1 from NAIK 09. $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ_1
<4	90	¹ NAIK 09	CLEO	$J/\psi \rightarrow \gamma\eta'$	NODE=M002R06 NODE=M002R06

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$. $\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{19}/Γ
<0.01	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)+\text{MM}$	NODE=M002R16 NODE=M002R16
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.01	90	LONDON 66	HBC	Compilation	

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ
< 3.1	90	¹ ABLIKIM 13U	BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$	NODE=M002R07 NODE=M002R07
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 53	90	² NAIK 09	CLEO	$J/\psi \rightarrow \gamma\eta'$	
<500	95	KALBFLEISCH 64B	HBC	$K^- p \rightarrow \Lambda 2(\pi^+\pi^-)$	

¹ Using $B(J/\psi \rightarrow \gamma\eta' (958)) = (5.16 \pm 0.15) \times 10^{-3}$.² Not independent of measured value of Γ_{20}/Γ_1 from NAIK 09. $\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ_1
<1.2	90	¹ NAIK 09	CLEO	$J/\psi \rightarrow \gamma\eta'$	NODE=M002R08 NODE=M002R08

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$. $\Gamma(K^\pm\pi^\mp)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ_2
< 1.3×10^{-4}	90	ABLIKIM 16M	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$	NODE=M002R61 NODE=M002R61

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
2.4 ± 1.3 OUR FIT						

 $[(2.4^{+1.3}_{-1.0}) \times 10^{-3} \text{ OUR 2019 FIT}]$ $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ 2.11 $\pm 0.12 \pm 0.14$ 429 ABLIKIM 130 BES3 $J/\psi \rightarrow \gamma\eta'$ 2.5 ± 1.2 ± 0.5 2 NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$ <6 90 RITTENBERG 65 HBC 2.7 $K^- p$ ¹ Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.² Not independent of measured value of Γ_{22}/Γ_1 from NAIK 09.

NODE=M002R22

NODE=M002R22

NODE=M002R23

NODE=M002R23

NODE=M002R23;LINKAGE=NA

NODE=M002R06

NODE=M002R06

NODE=M002R06;LINKAGE=NA

NODE=M002R16

NODE=M002R16

NODE=M002R07

NODE=M002R07

NODE=M002R08;LINKAGE=NA

NODE=M002R61

NODE=M002R61

NODE=M002R12

NODE=M002R12

NEW

NODE=M002R12;LINKAGE=A

NODE=M002R12;LINKAGE=NA

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ_1
---	-------------	--------------------	-------------	----------------	------------------------

5.5 ± 3.0 OUR FIT $[(5.5^{+3.0}_{-2.2}) \times 10^{-3}$ OUR 2019 FIT]**5.52 ± 3.00 ± 0.03** 8 1 NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ_2
7.2$\pm 0.4 \pm 0.5$	429	ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$

 $\Gamma(\pi^+e^-\nu_e + \text{c.c.})/\Gamma(\pi^+\pi^-\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{23}/Γ_1
<5.0	90	ABLIKIM	13G	BES3	$J/\psi \rightarrow \phi\eta'$

 $\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{24}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.9	90	BRIERE	00	CLEO	$10.6 e^+e^-$

 $\Gamma(\gamma e^+e^-)/\Gamma(\gamma\gamma)$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{24}/Γ_7
2.13$\pm 0.09 \pm 0.07$	864	ABLIKIM	150	BES3	$J/\psi \rightarrow \gamma e^+e^-$

 $\Gamma(\pi^0\gamma\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{25}/Γ
3.20$\pm 0.07 \pm 0.23$	3.4k	ABLIKIM	17T	BES3	$J/\psi \rightarrow \gamma\eta'$

 $\Gamma(\pi^0\gamma\gamma(\text{non resonant}))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{26}/Γ
6.16$\pm 0.64 \pm 0.67$	655	ABLIKIM	17T	BES3	$J/\psi \rightarrow \gamma\eta'$

 $\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{25}/Γ_4
<37	90	ALDE	87B	GAM2	$38 \pi^- p \rightarrow n4\gamma$

 $\Gamma(\eta\gamma\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{27}/Γ
<1.33	90	ABLIKIM	19AW	BES3	$J/\psi \rightarrow \gamma\eta' \rightarrow \gamma\gamma\gamma\gamma$

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{28}/Γ
<3.2 $\times 10^{-4}$	90	DONSKOV	14	GAM4	$32.5 \pi^- p \rightarrow \eta' n$

 $\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{28}/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<23	90	ALDE	87B	GAM2	$38 \pi^- p \rightarrow n8\gamma$

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{29}/Γ
< 5.6 $\times 10^{-9}$	90	1 ACHASOV	15	SND	$0.958 e^+e^- \rightarrow \pi\pi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12 $\times 10^{-9}$ 90 ² AKHMETSHIN 15 CMD3 $0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$

< 2.1 $\times 10^{-7}$ 90 VOROBIEV 88 ND $e^+e^- \rightarrow \pi^+\pi^-\eta$

¹ Combining data of ACHASOV 15 and AKHMETSHIN 15 and using $\Gamma(\eta') = 0.198 \pm 0.009$ MeV.

² Using $\Gamma_{\eta'(958)} = 198 \pm 9$ keV, $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.9 \pm 0.7)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.20)\%$.

NODE=M002R02

NODE=M002R02

NEW

NODE=M002R02;LINKAGE=NA

NODE=M002R56

NODE=M002R56

NODE=M002R54

NODE=M002R54

NODE=M002R40

NODE=M002R40

NODE=M002R00

NODE=M002R00

NODE=M002R64

NODE=M002R64

NODE=M002R65

NODE=M002R65

NODE=M002R35

NODE=M002R35

NODE=M002R67

NODE=M002R67

NODE=M002R58

NODE=M002R58

NODE=M002R37

NODE=M002R37

NODE=M002R39

NODE=M002R39

NODE=M002R39;LINKAGE=B

NODE=M002R39;LINKAGE=A

$\Gamma(\text{invisible})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{30}/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<9.5	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$	NODE=M002R52 NODE=M002R52

1 Not independent of measured value of Γ_{30}/Γ_1 from NAIK 09. $\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{30}/Γ_7
< 2.4	90	ABLIKIM	13	BES3 $J/\psi \rightarrow \phi\eta'$	NODE=M002R44 NODE=M002R44

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

<6.69	90	ABLIKIM	06Q	BES $J/\psi \rightarrow \phi\eta'$
-------	----	---------	-----	------------------------------------

 $\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\eta)$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{30}/Γ_1
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<2.1	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$	NODE=M002R09 NODE=M002R09

1 NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 5.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.41 \times 10^{-2}$. $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{31}/Γ
< 0.18	90	¹ AAIJ	17D	LHCb $D_{(s)}^+ \rightarrow \pi^+\pi^-\pi^+$	NODE=M002R20 NODE=M002R20

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

< 0.5	90	² ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-$
< 29	90	³ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+\pi^-$
< 3.3	90	⁴ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+\pi^-$
<800	95	DANBURG	73	HBC $2.2 K^- p \rightarrow \Lambda X^0$
<200	90	RITTENBERG	69	HBC $1.7-2.7 K^- p$

1 Using branching fractions of $D_{(s)}^+$ decays from PDG 15.2 ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.25 \times 10^{-3}$.3 Taking into account interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.4 Without interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum. $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{32}/Γ
< 4 x 10⁻⁴	90	¹ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$	NODE=M002R53 NODE=M002R53

1 ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.25 \times 10^{-3}$. $\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{32}/Γ_4
< 45	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n4\gamma$	NODE=M002R36 NODE=M002R36

 $\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{33}/Γ
< 1.4	90	BRIERE	00	CLEO $10.6 e^+ e^-$	NODE=M002R8 NODE=M002R8

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

<13	90	RITTENBERG	65	HBC $2.7 K^- p$
-----	----	------------	----	-----------------

 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{34}/Γ
< 2.4	90	BRIERE	00	CLEO $10.6 e^+ e^-$	NODE=M002R9 NODE=M002R9

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

<11	90	RITTENBERG	65	HBC $2.7 K^- p$
-----	----	------------	----	-----------------

 $\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{35}/Γ_4
< 4.6	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n3\gamma$	NODE=M002R34 NODE=M002R34

$\Gamma(\mu^+ \mu^- \pi^0)/\Gamma_{\text{total}}$				Γ_{36}/Γ	
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<6.0	90	DZHELYADIN	81	CNTR	$30 \pi^- p \rightarrow \eta' n$
$\Gamma(\mu^+ \mu^- \eta)/\Gamma_{\text{total}}$				Γ_{37}/Γ	
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.5	90	DZHELYADIN	81	CNTR	$30 \pi^- p \rightarrow \eta' n$
$\Gamma(e\mu)/\Gamma_{\text{total}}$				Γ_{38}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<4.7	90	BRIERE	00	CLEO	$10.6 e^+ e^-$

$\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$$

X and Y are Dalitz variables; α is complex and C, and D are real-valued. Parameters C and D are not necessarily equal to c and d, respectively, in the generalized parameterization following this one. May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

$\text{Re}(\alpha)$ decay parameter

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.034 ± 0.002 ± 0.002	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.054 ± 0.004 ± 0.001	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.033 ± 0.005 ± 0.003	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.072 ± 0.012 ± 0.006	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
-0.021 ± 0.018 ± 0.017	6.7k	³ BRIERE	00	CLEO $10.6 e^+ e^- \rightarrow \eta\pi^+\pi^- X$
-0.058 ± 0.013 ± 0.003	5.4k	⁴ ALDE	86	GAM2 $38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
-0.08 ± 0.03		^{4,5} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

1 See ABLIKIM 11 for the full correlation matrix.

2 Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

3 Assuming $\text{Im}(\alpha) = 0$, $C = 0$, and $D = 0$.

4 Assuming $C = 0$.

5 From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002R31
NODE=M002R31

NODE=M002R30
NODE=M002R30

NODE=M002R41
NODE=M002R41

NODE=M002225

NODE=M002225

NODE=M002A0
NODE=M002A0

OCCUR=2

NODE=M002A0;LINKAGE=AB
NODE=M002A0;LINKAGE=AM

NODE=M002A0;LINKAGE=BR
NODE=M002A0;LINKAGE=A
NODE=M002A0;LINKAGE=KA

NODE=M002IA0
NODE=M002IA0

OCCUR=2

NODE=M002IA0;LINKAGE=AB
NODE=M002IA0;LINKAGE=AM

NODE=M002IA0;LINKAGE=A
NODE=M002IA0;LINKAGE=KA

NODE=M002C0
NODE=M002C0

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.000 ± 0.019 ± 0.001	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
0.000 ± 0.038 ± 0.002	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
0.000 ± 0.049 ± 0.001	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.0 ± 0.1 ± 0.0	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
-0.00 ± 0.13 ± 0.00	5.4k	³ ALDE	86	GAM2 $38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
0.0 ± 0.3		^{3,4} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

1 See ABLIKIM 11 for the full correlation matrix.

2 Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

3 Assuming $C = 0$.

4 From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0027 ± 0.0024 ± 0.0015	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
0.018 ± 0.009 ± 0.003	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.020 ± 0.018 ± 0.004	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$

1 See ABLIKIM 11 for the full correlation matrix.

2 Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

NODE=M002C0;LINKAGE=AB
NODE=M002C0;LINKAGE=AM

D decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.053±0.004±0.004	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.061±0.009±0.005	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.059±0.012±0.004	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.066±0.030±0.015	7k	² AMELIN	05A	VES 28 $\pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
0.00 ±0.03 ±0.00	5.4k	³ ALDE	86	GAM2 38 $\pi^- p \rightarrow n\eta\pi^0\pi^0$
0		^{3,4} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

¹ See ABLIKIM 11 for the full correlation matrix.² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.³ Assuming C = 0.⁴ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002D0

NODE=M002D0

OCCUR=2

NODE=M002D0;LINKAGE=AB

NODE=M002D0;LINKAGE=AM

NODE=M002D0;LINKAGE=AL

NODE=M002D0;LINKAGE=KA

NODE=M002227

NODE=M002227

NODE=M002DPA

NODE=M002DPA

OCCUR=2

NODE=M002DPA;LINKAGE=A

NODE=M002DPA;LINKAGE=AB

NODE=M002DPA;LINKAGE=BL

NODE=M002DPA;LINKAGE=DO

NODE=M002DPB

NODE=M002DPB

OCCUR=2

NODE=M002DPB;LINKAGE=A

NODE=M002DPB;LINKAGE=AB

NODE=M002DPB;LINKAGE=BL

NODE=M002DPB;LINKAGE=DO

NODE=M002DPC

NODE=M002DPC

 $\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + a Y + b Y^2 + c X + d X^2$$

X and Y are Dalitz variables and a, b, c, and d are real-valued parameters.

May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays.

We do not average measurements in the section below because parameter values from each experiment are strongly correlated.

a decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.056±0.004±0.002	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.087±0.009±0.006	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.074±0.008±0.006	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta\pi^0\pi^0$
-0.072±0.007±0.008		¹ GONZALEZ-S..18A	RVUE	$\eta' \rightarrow \eta\pi^0\pi^0$
-0.047±0.011±0.003	44k	² ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.066±0.016±0.003	15k	³ BLIK	09	GAM4 32.5 $\pi^- p \rightarrow \eta' n$
-0.127±0.016±0.008	20k	⁴ DOROFEEV	07	VES 27 $\pi^- p \rightarrow \eta' n$, $\pi^- A \rightarrow \eta'\pi^- A^*$

¹ Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.² See ABLIKIM 11 for the full correlation matrix.³ From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.⁴ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.**b decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.049±0.006±0.006	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.073±0.014±0.005	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.063±0.014±0.005	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta\pi^0\pi^0$
-0.052±0.001±0.002		¹ GONZALEZ-S..18A	RVUE	$\eta' \rightarrow \eta\pi^0\pi^0$
-0.069±0.019±0.009	44k	² ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.063±0.028±0.004	15k	³ BLIK	09	GAM4 32.5 $\pi^- p \rightarrow \eta' n$
-0.106±0.028±0.014	20k	⁴ DOROFEEV	07	VES 27 $\pi^- p \rightarrow \eta' n$, $\pi^- A \rightarrow \eta'\pi^- A^*$

¹ Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.² See ABLIKIM 11 for the full correlation matrix.³ From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.⁴ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.**c decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0027±0.0024±0.0018	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
0.019 ±0.011 ±0.003	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.107 ±0.096 ±0.003	15k	² BLIK	09	GAM4 32.5 $\pi^- p \rightarrow \eta' n$
0.015 ±0.011 ±0.014	20k	³ DOROFEEV	07	VES 27 $\pi^- p \rightarrow \eta' n$, $\pi^- A \rightarrow \eta'\pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.² From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.³ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

NODE=M002D0

NODE=M002D0

OCCUR=2

NODE=M002D0;LINKAGE=AB

NODE=M002D0;LINKAGE=AM

NODE=M002D0;LINKAGE=AL

NODE=M002D0;LINKAGE=KA

NODE=M002227

NODE=M002227

NODE=M002DPA

NODE=M002DPA

OCCUR=2

NODE=M002DPA;LINKAGE=A

NODE=M002DPA;LINKAGE=AB

NODE=M002DPA;LINKAGE=BL

NODE=M002DPA;LINKAGE=DO

NODE=M002DPB

NODE=M002DPB

OCCUR=2

NODE=M002DPB;LINKAGE=A

NODE=M002DPB;LINKAGE=AB

NODE=M002DPB;LINKAGE=BL

NODE=M002DPB;LINKAGE=DO

NODE=M002DPC

NODE=M002DPC

NODE=M002DPC;LINKAGE=AB

NODE=M002DPC;LINKAGE=BL

NODE=M002DPC;LINKAGE=DO

d decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.063±0.004±0.003	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.074±0.009±0.004	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.050±0.009±0.005	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta\pi^0\pi^0$
-0.051±0.008±0.006		¹ GONZALEZ-S..18A	RVUE	$\eta' \rightarrow \eta\pi^0\pi^0$
-0.073±0.012±0.003	44k	² ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.018±0.078±0.006	15k	³ BLIK	09	GAM4 $32.5\pi^- p \rightarrow \eta' n$
-0.082±0.017±0.008	20k	⁴ DOROFEEV	07	VES $27\pi^- p \rightarrow \eta' n, \pi^- A \rightarrow \eta'\pi^- A^*$

¹Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

²See ABLIKIM 11 for the full correlation matrix.

³From $\eta' \rightarrow \eta\pi^0\pi^0$ decay. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$.

⁴From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

NODE=M002DPD
NODE=M002DPD

OCCUR=2

NODE=M002DPD;LINKAGE=A

NODE=M002DPD;LINKAGE=AB

NODE=M002DPD;LINKAGE=BL

NODE=M002DPD;LINKAGE=DO

NODE=M002226

NODE=M002226

NODE=M002B0
NODE=M002B0

NODE=M002235

NODE=M002235

NODE=M002A
NODE=M002A

NODE=M002FFL
NODE=M002FFL

NODE=M002FFL

NODE=M002FFL;LINKAGE=A

NODE=M002

REFID=60000
REFID=59773
REFID=58711
REFID=58843
REFID=58957
REFID=59316
REFID=57702
REFID=57650
REFID=58013
REFID=57511
REFID=56983
REFID=56602
REFID=56780
REFID=56781

$\eta'(958)$ β PARAMETER $|\text{MATRIX ELEMENT}|^2 = (1 + 2\beta Z)$

See the "Note on η Decay Parameters" in our 1994 edition Physical Review D50 1173 (1994), p. 1454.

 β decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.61 ±0.08 OUR AVERAGE				Error includes scale factor of 1.2.
-0.640±0.046±0.047	1.8k	ABLIKIM	15G	BES3 $J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$
-0.59 ±0.18	235	BLIK	08	GAMS $32\pi^- p \rightarrow \eta' n$
-0.1 ±0.3		ALDE	87B	GAM2 $38\pi^- p \rightarrow n3\pi^0$

 $\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

DECAY ASYMMETRY PARAMETER FOR $\pi^+\pi^-\gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.03 ±0.04 OUR AVERAGE				
-0.019±0.056		AIHARA	87	TPC $2\gamma \rightarrow \pi^+\pi^-\gamma$
-0.069±0.078	295	GRIGORIAN	75	STRC $2.1\pi^- p$
0.00 ±0.10	103	KALBFLEISCH	75	HBC $2.18K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.07 ±0.08	152	RITTENBERG	65	HBC $2.1-2.7K^- p$

 $\eta'(958) \rightarrow \gamma\ell^+\ell^-$ TRANSITION FORM FACTOR SLOPE

Related to the effective virtual meson mass Λ , via slope $\approx \Lambda^{-2}$. See e.g. LANDSBERG 85, eq. (3.8), for a detailed definition.

VALUE (GeV ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
1.62±0.17 OUR AVERAGE				
1.60±0.17±0.08	864	¹ ABLIKIM	150	BES3 $J/\psi \rightarrow \gamma e^+e^-$
1.7 ±0.4	33	¹ VIKTOROV	80	$25,33\pi^- p \rightarrow 2\mu\gamma$

¹In the single-pole Ansatz where slope = $1/(\Lambda^2 + \gamma^2)$ with Λ , γ being a Breit-Wigner mass, width for the effective contributing vector meson.

 $\eta'(958)$ REFERENCES

ABLIKIM	19AW	PR D100 052015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18	PR D97 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18C	PR C120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADLARSON	18A	PR D98 012001	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)
GONZALEZ-S...18A		EPJ C78 758	S. Gonzalez-Solis, E. Passemar	(BEIJ, IND+)
AAIJ	17D	PL B764 233	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17T	PR D96 012005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AD	PR D92 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15G	PR D92 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15O	PR D92 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ACHASOV	15	PR D91 092010	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=56788
AKHMETSHIN	15	PL B740 273	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=56386
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=5
ABLIKIM	14M	PRL 112 251801	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55904
DONSKOV	14	MPL A29 1450213	S. Donskov <i>et al.</i>	(GAMS-4π Collab.)	REFID=56321
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ABLIKIM	13	PR D87 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54829
ABLIKIM	13G	PR D87 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54952
ABLIKIM	13O	PR D87 092011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55388
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM	11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53711
CZERWINSKI	10	PRL 105 122001	E. Czerwinski <i>et al.</i>	(COSY-11 Collab.)	REFID=53364
BLIK	09	PAN 72 231	A.M. Blik <i>et al.</i>	(IHEP (Protvino))	REFID=52727
		Translated from YAF 72 258.			
NAIK	09	PRL 102 061801	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52678
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
BLIK	08	PAN 71 2124	A. Blik <i>et al.</i>	(GAMS-4π Collab.)	REFID=52663
		Translated from YAF 71 2161.			
LIBBY	08	PRL 101 182002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=52591
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
DOROFEEV	07	PL B651 22	V. Dorofeev <i>et al.</i>	(VES Collab.)	REFID=51711
MORI	07A	JPSJ 76 074102	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51691
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06Q	PR D97 202002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51487
AMELIN	05A	PAN 68 372	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=50766
		Translated from YAF 68 401.			
AMSLER	04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51079
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
BRIERE	00	PRL 84 26	R. Briere <i>et al.</i>	(CLEO Collab.)	REFID=47410
ACCIARRI	98Q	PL B418 399	M. Acciari <i>et al.</i>	(L3 Collab.)	REFID=46316
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
WURZINGER	96	PL B374 283	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=44992
PDG	94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)	REFID=43653
AMSLER	93	ZPHY C58 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43311
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)	REFID=43175
		Translated from YAF 55 2748.			
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=42170
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRMP+)	REFID=41862
BEHREND	91	ZPHY C49 401	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41497
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
KARCH	90	PL B249 353	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=41377
ROE	90	PR D41 17	N.A. Roe <i>et al.</i>	(ASP Collab.)	REFID=41014
AIHARA	88C	PR D38 1	H. Aihsara <i>et al.</i>	(TPC-2γ Collab.)	REFID=40564
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
WILLIAMS	88	PR D38 1365	D.A. Williams <i>et al.</i>	(Crystal Ball Collab.)	REFID=40567
AIHARA	87	PR D35 2650	H. Aihsara <i>et al.</i>	(TPC-2γ Collab.)	REFID=40009
ALBRECHT	87B	PL B199 457	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40265
ALDE	87B	ZPHY C36 603	D.M. Alde <i>et al.</i>	(LANL, BELG, SERP, LAPP)	REFID=40236
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ALDE	86	PL B177 115	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=20310
BARTEL	85E	PL 160B 421	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10843
LANDSBERG	85	PRPL 128 301	L.G. Landsberg	(SERP)	REFID=10844
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=20305
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)	REFID=20306
BINON	84	PL 140B 264	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP+)	REFID=20307
JENNI	83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)	REFID=20304
BARTEL	82B	PL 113B 190	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=20300
BEHREND	82C	PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
DZHELYADIN	81	PL 105B 239	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10836
STANTON	80	PL B92 353	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+)	REFID=40294
VIKTOROV	80	SJNP 32 520	V.A. Viktorov <i>et al.</i>	(SERP)	REFID=20298
		Translated from YAF 32 1005.			
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)	REFID=20295
BINNIE	79	PL 83B 141	D.M. Binnie <i>et al.</i>	(LOIC)	REFID=20296
ZANFINO	77	PRL 38 930	C. Zanfino <i>et al.</i>	(CARL, MCGI, OHIO+)	REFID=20293
GRIGORIAN	75	NP B91 232	A. Grigorian <i>et al.</i>	(+)	REFID=20287
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
DUANE	74	PRL 32 425	A. Duane <i>et al.</i>	(LOI, SHMP)	REFID=20284
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)	REFID=20286
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH)	REFID=20280
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+)	REFID=20281
AGUILAR...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20275
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)	REFID=20278
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20270
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)	REFID=20272
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=20268
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL)	REFID=20266
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)	REFID=20263
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJP	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=20253
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)	REFID=10761
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP	REFID=20247
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP	REFID=20252

$f_0(980)$ $I^G(J^{PC}) = 0^+(0^{++})$

See also the minireview on scalar mesons under $f_0(500)$. (See the index for the page number.)

NODE=M003

NODE=M003

NODE=M003M1

NODE=M003M1
→ UNCHECKED ←

 $f_0(980)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
990 ±20 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
992.0 \pm 8.5 \pm 8.6		1 AAIJ	19H LHCb $p\bar{p} \rightarrow D^\pm X$	
989.4 \pm 1.3	424	ABLIKIM	15P BES3 $J/\psi \rightarrow K^+ K^- 3\pi$	
989.9 \pm 0.4	706	ABLIKIM	12E BES3 $J/\psi \rightarrow \gamma 3\pi$	
1003 \pm 5 -27		2,3 GARCIA-MAR..11	RVUE Compilation	
996 \pm 7		2,4 GARCIA-MAR..11	RVUE Compilation	OCCUR=2
996 \pm 4 -14		5 MOUSSALLAM11	RVUE Compilation	
981 \pm 43		6 MENNESSIER 10	RVUE Compilation	
1030 \pm 30 -10		7 ANISOVICH 09	RVUE $0.0 \bar{p}p, \pi N$	
977 \pm 11 -9 \pm 1	44	8 ECKLUND 09	CLEO $4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + c.c.$	
982.2 \pm 1.0 \pm 8.1 -8.0		9 UEHARA 08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
976.8 \pm 0.3 \pm 10.1 -0.6	64k	10 AMBROSINO 07 KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
984.7 \pm 0.4 \pm 2.4 -3.7	64k	11 AMBROSINO 07 KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
973 \pm 3	262 \pm 30	12 AUBERT 07AKBABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$	
970 \pm 7	54 \pm 9	12 AUBERT 07AKBABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \pi^0 \gamma$	OCCUR=2
953 \pm 20	2.6k	13 BONVICINI 07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$	
985.6 \pm 1.2 \pm 1.1 -1.5 -1.6		14 MORI 07 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
983.0 \pm 0.6 \pm 4.0 -3.0		15 AMBROSINO 06B KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
977.3 \pm 0.9 \pm 3.7 -4.3		16 AMBROSINO 06B KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	OCCUR=2
950 \pm 9	4286	17 GARMASH 06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$	
965 \pm 10		18 ABLIKIM 05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-, \phi K^+ K^-$	
1031 \pm 8		19 ANISOVICH 03 RVUE		
1037 \pm 31		TIKHOMIROV 03 SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
973 \pm 1	2438	20 ALOISIO 02D KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
977 \pm 3 \pm 2	848	21 AITALA 01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$	
969.8 \pm 4.5	419	22 ACHASOV 00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
985 \pm 16 -12	419	23,24 ACHASOV 00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
976 \pm 5 \pm 6		25 AKHMETSHIN 99B CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
977 \pm 3 \pm 6	268	25 AKHMETSHIN 99C CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
975 \pm 4 \pm 6		26 AKHMETSHIN 99C CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=3
975 \pm 4 \pm 6		27 AKHMETSHIN 99C CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma, \pi^0 \pi^0 \gamma$	
985 \pm 10		BARBERIS 99 OMEG 450	$p\bar{p} \rightarrow p_s p_f K^+ K^-$	
		BARBERIS 99B OMEG 450	$p\bar{p} \rightarrow p_s p_f \pi^+ \pi^-$	
		BARBERIS 99C OMEG 450	$p\bar{p} \rightarrow p_s p_f \pi^0 \pi^0$	
982 \pm 3		28 BARBERIS 99D OMEG 450	$p\bar{p} \rightarrow K^+ K^-, \pi^+ \pi^-$	
982 \pm 3		BELLAZZINI 99 GAM4 450	$p\bar{p} \rightarrow p\bar{p} \pi^0 \pi^0$	
987 \pm 6 \pm 6		29 KAMINSKI 99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$	
989 \pm 15		29 OLLER 99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
991 \pm 3		OLLER 99B RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~ 980				
~ 993.5				

~ 987	²⁹ OLLER	99c	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
957 ± 6	³⁰ ACKERSTAFF	98Q	OPAL	$Z \rightarrow f_0 X$
960 ± 10	ALDE	98	GAM4	
1015 ± 15	²⁹ ANISOVICH	98B	RVUE	Compilation
1008	³¹ LOCHER	98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
955 ± 10	³⁰ ALDE	97	GAM2	$450 pp \rightarrow pp\pi^0\pi^0$
994 ± 9	³² BERTIN	97c	OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
$993.2 \pm 6.5 \pm 6.9$	³³ ISHIDA	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
1006	TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
997 ± 5	3k	³⁴ ALDE	95B	GAM2 $38 \pi^- p \rightarrow \pi^0\pi^0 n$
960 ± 10	10k	³⁵ ALDE	95B	GAM2 $38 \pi^- p \rightarrow \pi^0\pi^0 n$
994 ± 5	AMSLER	95B	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
~ 996	³⁶ AMSLER	95D	CBAR	$0.0 \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$
987 ± 6	³⁷ ANISOVICH	95	RVUE	
1015	JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
983	³⁸ BUGG	94	RVUE	$\bar{p}p \rightarrow \eta_2\pi^0$
973 ± 2	³⁹ KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
988	⁴⁰ ZOU	94B	RVUE	
988 ± 10	⁴¹ MORGAN	93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_s \rightarrow \pi(\pi\pi)$
971.1 ± 4.0	³⁰ AGUILAR-...	91	EHS	$400 pp$
979 ± 4	⁴² ARMSTRONG	91	OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
956 ± 12	BREAKSTONE	90	SFM	$pp \rightarrow pp\pi^+\pi^-$
959.4 ± 6.5	³⁰ AUGUSTIN	89	DM2	$J/\psi \rightarrow \omega\pi^+\pi^-$
978 ± 9	³⁰ ABACHI	86B	HRS	$e^+ e^- \rightarrow \pi^+\pi^- X$
985.0 ± 9.0	ETKIN	82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$
974 ± 4	⁴² GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+\pi^- X$
975	⁴³ ACHASOV	80	RVUE	
986 ± 10	⁴² AGUILAR-...	78	HBC	$0.7 \bar{p}p \rightarrow K_S^0 K_S^0$
969 ± 5	⁴² LEEPER	77	ASPK	$2-2.4 \pi^- p \rightarrow \pi^+\pi^- n, K^+K^-n$
987 ± 7	⁴² BINNIE	73	CNTR	$\pi^- p \rightarrow nMM$
1012 ± 6	⁴⁴ GRAYER	73	ASPK	$17 \pi^- p \rightarrow \pi^+\pi^- n$
1007 ± 20	⁴⁴ HYAMS	73	ASPK	$17 \pi^- p \rightarrow \pi^+\pi^- n$
997 ± 6	⁴⁴ PROTOPOP...	73	HBC	$7 \pi^+ p \rightarrow \pi^+ p\pi^+\pi^-$

1 From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

2 Quoted number refers to real part of pole position.

3 Analytic continuation using Roy equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.

4 Analytic continuation using GKPY equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.

5 Pole position. Used Roy equations.

6 Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.

7 On sheet II in a 2-pole solution. The other pole is found on sheet III at $(850-100i)$ MeV

8 Using a relativistic Breit-Wigner function and taking into account the finite D_s mass.

9 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K\bar{K}/g_{f_0}\pi\pi = 0$.

10 In the kaon-loop fit.

11 In the no-structure fit.

12 Systematic errors not estimated.

13 FLATTE 76 parameterization. $g_{f_0}\pi\pi = 329 \pm 96$ MeV/c² assuming $g_{f_0} K\bar{K}/g_{f_0}\pi\pi = 2$.

14 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K\bar{K}/g_{f_0}\pi\pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.

15 In the kaon-loop fit following formalism of ACHASOV 89.

16 In the no-structure fit assuming a direct coupling of ϕ to $f_0\gamma$.

17 FLATTE 76 parameterization. Supersedes GARMASH 05.

18 FLATTE 76 parameterization, $g_{f_0} K\bar{K}/g_{f_0}\pi\pi = 4.21 \pm 0.25 \pm 0.21$.

19 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0\pi^0 n, \pi^- p \rightarrow K\bar{K} n, \pi^+\pi^- \rightarrow \pi^+\pi^-, \bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta, \pi^+\pi^-\pi^0, K^+K^-\pi^0, K_S^0 K_S^0\pi^0$,

$K^+K_S^0\pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+, K_S^0 K^- \pi^0, K_S^0 K_S^0\pi^-$ at rest.

OCCUR=2

NODE=M003M1;LINKAGE=F

NODE=M003M1;LINKAGE=GC

NODE=M003M1;LINKAGE=GM

NODE=M003M1;LINKAGE=GI

NODE=M003M1;LINKAGE=MU

NODE=M003M1;LINKAGE=ME

NODE=M003M1;LINKAGE=AO

NODE=M003M1;LINKAGE=EC

NODE=M003M1;LINKAGE=UE

NODE=M003M1;LINKAGE=AK

NODE=M003M1;LINKAGE=AS

NODE=M003M1;LINKAGE=NS

NODE=M003M1;LINKAGE=BO

NODE=M003M1;LINKAGE=MO

NODE=M003M1;LINKAGE=AB

NODE=M003M1;LINKAGE=AM

NODE=M003M1;LINKAGE=GR

NODE=M003M1;LINKAGE=AL

NODE=M003M;LINKAGE=KM

- 20 From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution.
- 21 Coupled-channel Breit-Wigner, couplings $g_\pi=0.09\pm 0.01\pm 0.01$, $g_K=0.02\pm 0.04\pm 0.03$.
- 22 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
- 23 Supersedes ACHASOV 98I.
- 24 In the “narrow resonance” approximation.
- 25 Assuming $\Gamma(f_0)=40$ MeV.
- 26 From a narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.
- 27 From the combined fit of the photon spectra in the reactions $e^+e^- \rightarrow \pi^+\pi^-\gamma$, $\pi^0\pi^0\gamma$.
- 28 Supersedes BARBERIS 99 and BARBERIS 99B
- 29 T-matrix pole.
- 30 From invariant mass fit.
- 31 On sheet II in a 2 pole solution. The other pole is found on sheet III at $(1039-93i)$ MeV.
- 32 On sheet II in a 2 pole solution. The other pole is found on sheet III at $(963-29i)$ MeV.
- 33 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 34 At high $|t|$.
- 35 At low $|t|$.
- 36 On sheet II in a 4-pole solution, the other poles are found on sheet III at $(953-55i)$ MeV and on sheet IV at $(938-35i)$ MeV.
- 37 Combined fit of ALDE 95B, ANISOVICH 94, AMSLER 94D.
- 38 On sheet II in a 2 pole solution. The other pole is found on sheet III at $(996-103i)$ MeV.
- 39 From sheet II pole position.
- 40 On sheet II in a 2 pole solution. The other pole is found on sheet III at $(797-185i)$ MeV and can be interpreted as a shadow pole.
- 41 On sheet II in a 2 pole solution. The other pole is found on sheet III at $(978-28i)$ MeV.
- 42 From coupled channel analysis.
- 43 Coupled channel analysis with finite width corrections.
- 44 Included in AGUILAR-BENITEZ 78 fit.

$f_0(980)$ WIDTH

Width determination very model dependent. Peak width in $\pi\pi$ is about 50 MeV, but decay width can be much larger.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10 to 100 OUR ESTIMATE					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
15.3 ± 4.7	424	ABLIKIM	15P	BES3 $J/\psi \rightarrow K^+K^-3\pi$	
9.5 ± 1.1	706	ABLIKIM	12E	BES3 $J/\psi \rightarrow \gamma 3\pi$	
42 + 20 - 16		1,2 GARCIA-MAR..11	RVUE	Compilation	
50 + 20 - 12		2,3 GARCIA-MAR..11	RVUE	Compilation	
48 + 22 - 6		4 MOUSSALLAM11	RVUE	Compilation	
36 ± 22		5 MENNESSIER 10	RVUE	Compilation	
70 + 20 - 32		6 ANISOVICH 09	RVUE	0.0 $\bar{p}p, \pi N$	
91 + 30 - 22 ± 3	44	7 ECKLUND	09	CLEO $4.17 e^+e^- \rightarrow D_s^- D_s^{*+} + c.c.$	I
66.9 ± 2.2 + 17.6 - 12.5		8 UEHARA	08A	BELL $10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	
65 ± 13	262 ± 30	9 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$	
81 ± 21	54 ± 9	9 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\pi^0\gamma$	OCCUR=2
51.3 + 20.8 + 13.2 - 17.7 - 3.8		10 MORI	07	BELL $10.6 e^+e^- \rightarrow e^+e^-\pi^+\pi^-$	
61 ± 9 + 14 - 8	2584	11 GARMASH	05	BELL $B^+ \rightarrow K^+\pi^+\pi^-$	
64 ± 16		12 ANISOVICH	03	RVUE	
121 ± 23		TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
~ 70		13 BRAMON	02	RVUE $1.02 e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2
44 ± 2 ± 2	848	14 AITALA	01A	E791 $D_s^+ \rightarrow \pi^-\pi^+\pi^+$	
201 ± 28	419	15 ACHASOV	00H	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$	

NODE=M003M1;LINKAGE=KD

NODE=M003M;LINKAGE=TL

NODE=M003M;LINKAGE=V9

NODE=M003M;LINKAGE=V8

NODE=M003M1;LINKAGE=AI

NODE=M003M;LINKAGE=SM

NODE=M003M;LINKAGE=ST

NODE=M003M;LINKAGE=SL

NODE=M003M1;LINKAGE=BD

NODE=M003M1;LINKAGE=AN

NODE=M003M1;LINKAGE=A

NODE=M003M1;LINKAGE=LC

NODE=M003M1;LINKAGE=X

NODE=M003M1;LINKAGE=AA

NODE=M003M1;LINKAGE=LA

NODE=M003M1;LINKAGE=LB

NODE=M003M1;LINKAGE=KL

NODE=M003M1;LINKAGE=CF

NODE=M003M1;LINKAGE=C2

NODE=M003M1;LINKAGE=KM

NODE=M003M1;LINKAGE=L

NODE=M003M1;LINKAGE=K

NODE=M003M1;LINKAGE=B

NODE=M003M;LINKAGE=B

NODE=M003M;LINKAGE=R

NODE=M003W1

NODE=M003W1

NODE=M003W1

→ UNCHECKED ←

OCCUR=2

OCCUR=2

122 ± 13	419	16,17	ACHASOV	00H	SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
56 ± 20		18	AKHMETSHIN	99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
65 ± 20			BARBERIS	99	OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$	
80 ± 10			BARBERIS	99B	OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$	
80 ± 10			BARBERIS	99C	OMEG	$450 pp \rightarrow p_s p_f \pi^0 \pi^0$	
48 ± 12 ± 8		19	BARBERIS	99D	OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$	
65 ± 25			BELLAZZINI	99	GAM4	$450 pp \rightarrow pp\pi^0\pi^0$	
71 ± 14		20	KAMINSKI	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$	
~28		20	OLLER	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~25			OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~14		20	OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$	
70 ± 20			ALDE	98	GAM4		
86 ± 16		20	ANISOVICH	98B	RVUE	Compilation	
54		21	LOCHER	98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
69 ± 15		22	ALDE	97	GAM2	$450 pp \rightarrow pp\pi^0\pi^0$	
38 ± 20		23	BERTIN	97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
~100		24	ISHIDA	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
34			TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$	
48 ± 10	3k	25	ALDE	95B	GAM2	$38 \pi^- p \rightarrow \pi^0 \pi^0 n$	
95 ± 20	10k	26	ALDE	95B	GAM2	$38 \pi^- p \rightarrow \pi^0 \pi^0 n$	OCCUR=2
26 ± 10			AMSLER	95B	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$	
~112		27	AMSLER	95D	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$	
80 ± 12		28	ANISOVICH	95	RVUE		
30			JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
74		29	BUGG	94	RVUE	$\bar{p}p \rightarrow \eta_2 \pi^0$	
29 ± 2		30	KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
46		31	ZOU	94B	RVUE		
48 ± 12		32	MORGAN	93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_s \rightarrow \pi(\pi\pi)$	
37.4 ± 10.6		22	AGUILAR-...	91	EHS	$400 pp$	
72 ± 8		33	ARMSTRONG	91	OMEG	$300 pp \rightarrow pp\pi\pi, ppKK$	
110 ± 30			BREAKSTONE	90	SFM	$pp \rightarrow pp\pi^+\pi^-$	
29 ± 13		22	ABACHI	86B	HRS	$e^+ e^- \rightarrow \pi^+ \pi^- X$	
120 ± 281 ± 20			ETKIN	82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$	
28 ± 10		33	GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+ \pi^- X$	
70 to 300		34	ACHASOV	80	RVUE		
100 ± 80		35	AGUILAR-...	78	HBC	$0.7 \bar{p}p \rightarrow K_S^0 K_S^0$	
30 ± 8		33	LEEPER	77	ASPK	$2-2.4 \pi^- p \rightarrow \pi^+ \pi^- n, K^+ K^- n$	
48 ± 14		33	BINNIE	73	CNTR	$\pi^- p \rightarrow nMM$	
32 ± 10		36	GRAYER	73	ASPK	$17 \pi^- p \rightarrow \pi^+ \pi^- n$	
30 ± 10		36	HYAMS	73	ASPK	$17 \pi^- p \rightarrow \pi^+ \pi^- n$	
54 ± 16		36	PROTOPOPESCU	73	HBC	$7 \pi^+ p \rightarrow \pi^+ p\pi^+ \pi^-$	

1 Analytic continuation using Roy equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPOVESCU 73.

2 Quoted number refers to twice imaginary part of pole position.

3 Analytic continuation using GKPY equations. Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPOVESCU 73.

4 Pole position. Used Roy equations.

5 Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.

6 On sheet II in a 2-pole solution. The other pole is found on sheet III at (850–100) MeV

7 Using a relativistic Breit-Wigner function and taking into account the finite D_s mass.

8 Breit-Wigner $\pi\pi$ width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K_K / g_{f_0} \pi\pi = 0$.

9 Systematic errors not estimated.

10 Breit-Wigner $\pi\pi$ width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K_K / g_{f_0} \pi\pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.

NODE=M003W1;LINKAGE=GC

NODE=M003W1;LINKAGE=GI

NODE=M003W1;LINKAGE=GM

NODE=M003W1;LINKAGE=MU

NODE=M003W1;LINKAGE=ME

NODE=M003W1;LINKAGE=AO

NODE=M003W1;LINKAGE=EC

NODE=M003W1;LINKAGE=UE

NODE=M003W1;LINKAGE=NS

NODE=M003W1;LINKAGE=MO

- 11 Breit-Wigner, solution 1, PWA ambiguous.
 12 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.
 13 Using the data of AKHMETSHIN 99c, ACHASOV 00H, and ALOISIO 02D.
 14 Breit-Wigner width.
 15 Supersedes ACHASOV 98l. Using the model of ACHASOV 89.
 16 Supersedes ACHASOV 98l.
 17 In the "narrow resonance" approximation.
 18 From the combined fit of the photon spectra in the reactions $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$.
 19 Supersedes BARBERIS 99 and BARBERIS 99B
 20 T-matrix pole.
 21 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039–93*i*) MeV.
 22 From invariant mass fit.
 23 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963–29*i*) MeV.
 24 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
 25 At high $|t|$.
 26 At low $|t|$.
 27 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953–55*i*) MeV and on sheet IV at (938–35*i*) MeV.
 28 Combined fit of ALDE 95B, ANISOVICH 94,
 29 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996–103*i*) MeV.
 30 From sheet II pole position.
 31 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797–185*i*) MeV and can be interpreted as a shadow pole.
 32 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978–28*i*) MeV.
 33 From coupled channel analysis.
 34 Coupled channel analysis with finite width corrections.
 35 From coupled channel fit to the HYAMS 73 and PROTOPOPESCU 73 data. With a simultaneous fit to the $\pi\pi$ phase-shifts, inelasticity and to the $K_S^0 K_S^0$ invariant mass.
 36 Included in AGUILAR-BENITEZ 78 fit.

$f_0(980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 K\bar{K}$	seen
$\Gamma_3 \gamma\gamma$	seen
$\Gamma_4 e^+ e^-$	

$f_0(980)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	DOCUMENT ID	TECN	COMMENT	Γ_3
VALUE (keV)				

0.31 $^{+0.05}_{-0.04}$ OUR AVERAGE

0.32 ± 0.05	1 DAI	14A RVUE	Compilation	
0.286 ± 0.017 $^{+0.211}_{-0.070}$	2 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
0.205 $^{+0.095}_{-0.083}$ $^{+0.147}_{-0.117}$	3 MORI	07 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
0.42 ± 0.06 ± 0.18	4 OEST	90 JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.16 ± 0.01	5 MENNESSIER	11 RVUE		
0.29 ± 0.21 $^{+0.02}_{-0.07}$	6 MOUSSALLAM	11 RVUE	Compilation	
0.42	7,8 PENNINGTON	08 RVUE	Compilation	
0.10	8,9 PENNINGTON	08 RVUE	Compilation	
0.28 $^{+0.09}_{-0.13}$	10 BOGLIONE	99 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$	
0.29 ± 0.07 ± 0.12	11,12 BOYER	90 MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
0.31 ± 0.14 ± 0.09	11,12 MARSISKE	90 CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
0.63 ± 0.14	13 MORGAN	90 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$	

NODE=M003W1;LINKAGE=GA
 NODE=M003W;LINKAGE=KM

NODE=M003W;LINKAGE=BR
 NODE=M003W;LINKAGE=TL
 NODE=M003W;LINKAGE=V9
 NODE=M003W;LINKAGE=V8
 NODE=M003W1;LINKAGE=AI
 NODE=M003W;LINKAGE=SL

NODE=M003W1;LINKAGE=BD
 NODE=M003W1;LINKAGE=AN
 NODE=M003W1;LINKAGE=LO
 NODE=M003W1;LINKAGE=A
 NODE=M003W1;LINKAGE=X
 NODE=M003W1;LINKAGE=AA

NODE=M003W1;LINKAGE=LA
 NODE=M003W1;LINKAGE=LB
 NODE=M003W1;LINKAGE=KL

NODE=M003W1;LINKAGE=CF
 NODE=M003W1;LINKAGE=C2
 NODE=M003W1;LINKAGE=KM
 NODE=M003W1;LINKAGE=L

NODE=M003W1;LINKAGE=K
 NODE=M003W1;LINKAGE=B
 NODE=M003W;LINKAGE=B
 NODE=M003W;LINKAGE=C

NODE=M003W;LINKAGE=R

NODE=M003215;NODE=M003

DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=5;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=4

NODE=M003220

NODE=M003W4
 NODE=M003W4

OCCUR=2

- 1 Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.
 2 Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0 \pi \pi} = 0$.
 3 Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0 \pi \pi} = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.
 4 OEST 90 quote systematic errors ± 0.08 . We use ± 0.18 . Observed 60 events.
 5 Uses an analytic K-matrix model. Compilation.
 6 Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.
 7 Solution A (preferred solution based on χ^2 -analysis).
 8 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.
 9 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).
 10 Supersedes MORGAN 90.
 11 From analysis allowing arbitrary background unconstrained by unitarity.
 12 Data included in MORGAN 90, BOGLIONE 99 analyses.
 13 From amplitude analysis of BOYER 90 and MARSISKE 90, data corresponds to resonance parameters $m = 989$ MeV, $\Gamma = 61$ MeV.

 $\Gamma(e^+ e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4
<8.4	90	VOROBIEV	88	ND	$e^+ e^- \rightarrow \pi^0 \pi^0$

 $f_0(980)$ BRANCHING RATIOS **$\Gamma(\pi\pi)/[\Gamma(\pi\pi) + \Gamma(K\bar{K})]$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/(\Gamma_1 + \Gamma_2)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.52 \pm 0.12	9.9k	1 AUBERT	060 BABR	$B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$	
0.75 \pm 0.11 -0.13		2 ABLIKIM	05Q BES2	$\chi_{c0} \rightarrow 2\pi^+ 2\pi^-$, $\pi^+ \pi^- K^+ K^-$	
0.84 \pm 0.02 \sim 0.68		3 ANISOVICH OLLER	02D SPEC 99B RVUE	Combined fit $\pi\pi \rightarrow \pi\pi, K\bar{K}$	
0.67 \pm 0.09		4 LOVERRE	80 HBC	$4\pi^- p \rightarrow n2K_S^0$	
0.81 \pm 0.09 -0.04		4 CASON	78 STRC	$7\pi^- p \rightarrow n2K_S^0$	
0.78 \pm 0.03		4 WETZEL	76 OSPK	$8.9\pi^- p \rightarrow n2K_S^0$	

1 Recalculated by us using $\Gamma(K^+ K^-) / \Gamma(\pi^+ \pi^-) = 0.69 \pm 0.32$ from AUBERT 060 and isospin relations.

2 Using data from ABLIKIM 04G.

3 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta \eta n$, $\eta \eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

4 Measure $\pi\pi$ elasticity assuming two resonances coupled to the $\pi\pi$ and $K\bar{K}$ channels only.

 $f_0(980)$ REFERENCES

AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GARCIA-MAR... 11		PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
GARCIA-MAR... 11A		PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
MENNESSIER	11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang	
MOUSSALLAM	11	EPJ C71 1814	B. Moussallam	
BATLEY	10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
MENNESSIER	10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang	
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
ECKLUND	09	PR D80 052009	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)
BATLEY	08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)
AMBROSINO	06B	PL B634 148	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AUBERT	06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)
GARMASH	06	PR D 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ACHASOV	05	PR D72 013006	N.N. Achasov, G.N. Shestakov	
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
BUETTIKER	04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam	
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	

Translated from YAF 66 860.

NODE=M003W4;LINKAGE=D

NODE=M003W4;LINKAGE=UE

NODE=M003W4;LINKAGE=MO

NODE=M003W4;LINKAGE=H

NODE=M003W4;LINKAGE=ME

NODE=M003W4;LINKAGE=MU

NODE=M003W4;LINKAGE=P1

NODE=M003W4;LINKAGE=P3

NODE=M003W4;LINKAGE=P2

NODE=M003W4;LINKAGE=BL

NODE=M003W4;LINKAGE=B

NODE=M003W4;LINKAGE=C

NODE=M003W4;LINKAGE=A

NODE=M003W3

NODE=M003W3

NODE=M003225

NODE=M003R1

NODE=M003R1

NODE=M003R1;LINKAGE=AU

NODE=M003R1;LINKAGE=AB

NODE=M003R;LINKAGE=CH

NODE=M003R1;LINKAGE=B

NODE=M003

REFID=59670

REFID=59987

REFID=56781

REFID=55923

REFID=54270

REFID=16761

REFID=54121

REFID=53637

REFID=53975

REFID=53567

REFID=53657

REFID=52719

REFID=53041

REFID=52487

REFID=52303

REFID=52309

REFID=51616

REFID=51908

REFID=51721

REFID=51652

REFID=51043

REFID=51141

REFID=51162

REFID=50450

REFID=50958

REFID=50762

REFID=50641

REFID=50187

REFID=56428;ERROR=6

REFID=49401

REFID=49423

ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			
BRAMON	02	EPJ C26 253	A. Bramon <i>et al.</i>		REFID=49178
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)	REFID=48312
ITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
ITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99C	PL B453 325	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46923
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>		REFID=46600
ACKERSTAFF	98Q	EPJ C4 19	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46145
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)	REFID=46372
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ISHIDA	96	PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45770
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=44375
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)	REFID=44442
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)	REFID=44072
MORGAN	93	PR D48 1185	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=43614
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivchenko		REFID=48021
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ACHASOV	80	SJNP 32 566	N.N. Achasov, S.A. Devyanin, G.N. Shestakov	(NOVM)	REFID=20458
		Translated from YAF 32 1098.			
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP	REFID=20381
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+) IJP	REFID=20382
AGUILAR-...	78	NP B140 73	M. Aguilar-Benitez <i>et al.</i>	(MADR, BOMB+)	REFID=20368
CASON	78	PRL 41 271	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20370
LEEPER	77	PR D16 2054	R.J. Leeper <i>et al.</i>	(ISU)	REFID=20365
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)	REFID=20446
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)	REFID=20362
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)	REFID=21062
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
BINNIE	73	PRL 31 1534	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20343
GRAYER	73	Tallahassee	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20347
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108

$a_0(980)$ $I^G(J^P C) = 1^-(0^{++})$

See our minireview on scalar mesons under $f_0(500)$. (See the index for the page number.)

 $a_0(980)$ MASSVALUE (MeV)DOCUMENT ID **980 ± 20 OUR ESTIMATE**

Mass determination very model dependent

 $\eta\pi$ FINAL STATE ONLYVALUE (MeV)EVTSDOCUMENT IDTECNCHGCOMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

982.5	± 1.6	± 1.1	16.9k	1	AMBROSINO	09F	KLOE	$1.02 e^+ e^- \rightarrow \eta\pi^0\gamma$
986	± 4				ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$
982.3	$+ 0.6$	$+ 3.1$		2	UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
987.4	± 1.0	± 3.0		3,4	BUGG	08A	RVUE 0	$\bar{p}p \rightarrow \pi^0\pi^0\eta$
989.1	± 1.0	± 3.0		4,5	BUGG	08A	RVUE 0	$\bar{p}p \rightarrow \pi^0\pi^0\eta$
985	± 4	± 6	318		ACHARD	02B	L3	$183-209 e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
995	$+ 52$		36	6	ACHASOV	00F	SND	$e^+ e^- \rightarrow \eta\pi^0\gamma$
994	$+ 33$		36	7	ACHASOV	00F	SND	$e^+ e^- \rightarrow \eta\pi^0\gamma$
975	± 7				BARBERIS	00H		$450 pp \rightarrow p_f \eta\pi^0 p_s$
988	± 8				BARBERIS	00H		$450 pp \rightarrow \Delta_f^{++} \eta\pi^- p_s$
~ 1055				8	OLLER	99	RVUE	$\eta\pi, K\bar{K}$
~ 1009.2				8	OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
993.1	± 2.1			9	TEIGE	99	B852	$18.3 \pi^- p \rightarrow \eta\pi^+\pi^- n$
988	± 6			8	ANISOVICH	98B	RVUE	Compilation
987					TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
991					JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi, \eta\pi$
984.45	$\pm 1.23 \pm 0.34$				AMSLER	94C	CBAR	$0.0 \bar{p}p \rightarrow \omega\eta\pi^0$
982	± 2			10	AMSLER	92	CBAR	$0.0 \bar{p}p \rightarrow \eta\eta\pi^0$
984	± 4		1040	10	ARMSTRONG	91B	OMEG ±	$300 pp \rightarrow pp\eta\pi^+\pi^-$
976	± 6				ATKINSON	84E	OMEG ±	$25-55 \gamma p \rightarrow \eta\pi n$
986	± 3		500	11	EVANGELIS...	81	OMEG ±	$12 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
990	± 7			11	GURTU	79	HBC	$4.2 K^- p \rightarrow \Lambda\eta 2\pi$
980	± 11				CONFORTO	78	OSPK	$4.5 \pi^- p \rightarrow pX^-$
978	± 16				CORDEN	78	OMEG ±	$12-15 \pi^- p \rightarrow n\eta 2\pi$
977	± 7				GRASSLER	77	HBC	$16 \pi^\mp p \rightarrow p\eta 3\pi$
989	± 4		70		WELLS	75	HBC	$3.1-6 K^- p \rightarrow \Lambda\eta 2\pi$
972	± 10		150		DEFOIX	72	HBC	$0.7 \bar{p}p \rightarrow 7\pi$
970	± 15		20		BARNES	69C	HBC	$4-5 K^- p \rightarrow \Lambda\eta 2\pi$
980	± 10				CAMPBELL	69	DBC	$2.7 \pi^+ d$
980	± 10		15		MILLER	69B	HBC	$4.5 K^- N \rightarrow \eta\pi\Lambda$
980	± 10		30		AMMAR	68	HBC	$5.5 K^- p \rightarrow \Lambda\eta 2\pi$

1 Using the model of ACHASOV 89 and ACHASOV 03B.

2 From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

3 Parameterizes couplings to $\bar{K}K$, $\pi\eta$, and $\pi\eta'$.

4 Using AMSLER 94D and ABELE 98.

5 From the T-matrix pole on sheet II.

6 Using the model of ACHASOV 89. Supersedes ACHASOV 98B.

7 Using the model of JAFFE 77. Supersedes ACHASOV 98B.

8 T-matrix pole.

9 Breit-Wigner fit, average between a_0^\pm and a_0^0 . The fit favors a slightly heavier a_0^\pm .

10 From a single Breit-Wigner fit.

11 From $f_1(1285)$ decay.

NODE=M036

NODE=M036

NODE=M036205

NODE=M036MX

→ UNCHECKED ←

NODE=M036M1

NODE=M036M1

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M036M1;LINKAGE=AM

NODE=M036M1;LINKAGE=UE

NODE=M036M1;LINKAGE=BP

NODE=M036M1;LINKAGE=BU

NODE=M036M1;LINKAGE=BT

NODE=M036M1;LINKAGE=V1

NODE=M036M1;LINKAGE=M2

NODE=M036M1;LINKAGE=AN

NODE=M036M1;LINKAGE=BF

NODE=M036M1;LINKAGE=A

NODE=M036M1;LINKAGE=R

$K\bar{K}$ ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
947.7 \pm 5.5 5.0	1 AAIJ	19H LHCb	$p\bar{p} \rightarrow D^\pm X$	
925 \pm 5 \pm 8 \sim 1053	190k	2 AAIJ 3 OLLER	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$ $\pi\pi \rightarrow \pi\pi, K\bar{K}$	
982 \pm 3 975 \pm 15		4 ABELE	$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$	
976 \pm 6 1016 \pm 10	316 100	BERTIN DEBILLY 5 ASTIER	98B OBLX 80 HBC 67 HBC	$0.0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp$ $1.2-2 \bar{p}p \rightarrow f_1(1285)\omega$
1003.3 \pm 7.0	143	6 ROSENFELD	65 RVUE	

1 From the $D^\pm \rightarrow K^\pm K^\pm K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

2 Using a two-channel resonance parametrization with couplings fixed to ABELE 98.

3 T-matrix pole.

4 T-matrix pole on sheet II, the pole on sheet III is at 1006-i49 MeV.

5 ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.

6 Plus systematic errors.

NODE=M036M2
NODE=M036M2

 $a_0(980)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
50 to 100 OUR ESTIMATE Width determination very model dependent. Peak width in $\eta\pi$ is about 60 MeV, but decay width can be much larger.					

• • • We do not use the following data for averages, fits, limits, etc. • • •

75.6 \pm 1.6 \pm 17.4 80.2 \pm 3.8 \pm 5.4 50 \pm 13 \pm 4 72 \pm 16 61 \pm 19 \sim 42 \sim 112 71 \pm 7 92 \pm 20 65 \pm 10 \sim 100 202 54.12 \pm 0.34 \pm 0.12 54 \pm 10 95 \pm 14 62 \pm 15 60 \pm 20 60 \pm 50 86.0 \pm 60.0 44 \pm 22 80 to 300 16.0 \pm 25.0 30 \pm 5 40 \pm 15 60 \pm 30 80 \pm 30	1 UEHARA 2 BUGG 318 ACHARD BARBERIS BARBERIS 3 OLLER 3 OLLER TEIGE 3 ANISOVICH 4 BERTIN TORNQVIST JANSSEN AMSLER 5 AMSLER 5 ARMSTRONG 6 EVANGELIS... 6 GURTU 47 CONFORTO 50 CORDEN GRASSLER 7 FLATTE WELLS DEFOIX CAMPBELL MILLER AMMAR	09A BELL 08A RVUE 0 02B L3 00H 00H 99 RVUE 99B RVUE 99 B852 98B RVUE 98B OBLX \pm 96 RVUE 95 RVUE 94C CBAR 92 CBAR 91B OMEG \pm 500 81 OMEG \pm 145 47 50 77 76 70 150 69 15 30 1	0.0 $\bar{p}p \rightarrow \omega\eta\pi^0$ 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$ 300 $p\bar{p} \rightarrow p\bar{p}\eta\pi^+\pi^-$ 12 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-$ 4.2 $K^- p \rightarrow \Lambda\eta 2\pi$ 4.5 $\pi^- p \rightarrow pX^-$ 12-15 $\pi^- p \rightarrow n\eta 2\pi$ 16 $\pi^\mp p \rightarrow p\eta 3\pi$ 4.2 $K^- p \rightarrow \Lambda\eta 2\pi$ 3.1-6 $K^- p \rightarrow \Lambda\eta 2\pi$ 0.7 $\bar{p}p \rightarrow 7\pi$ 2.7 $\pi^+ d$ 4.5 $K^- N \rightarrow \eta\pi\Lambda$ 5.5 $K^- p \rightarrow \Lambda\eta 2\pi$		
---	---	---	--	--	--

NODE=M036M2;LINKAGE=C

NODE=M036M2;LINKAGE=B

NODE=M036M2;LINKAGE=AN

NODE=M036M2;LINKAGE=Q

NODE=M036M2;LINKAGE=A

NODE=M036M2;LINKAGE=S

NODE=M036210

NODE=M036W1

→ UNCHECKED ←

OCCUR=2

NODE=M036W1;LINKAGE=UE

NODE=M036W1;LINKAGE=BU

NODE=M036W1;LINKAGE=AN

NODE=M036W1;LINKAGE=BE

NODE=M036W1;LINKAGE=A

NODE=M036W1;LINKAGE=R

NODE=M036W1;LINKAGE=F

1 From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

2 From the T-matrix pole on sheet II, using AMSLER 94D and ABELE 98.

3 T-matrix pole.

4 The $\eta\pi$ width.

5 From a single Breit-Wigner fit.

6 From $f_1(1285)$ decay.

7 Using a two-channel resonance parametrization of GAY 76B data.

$K\bar{K}$ ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
92± 8		1 ABELE	98	CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 24		2 OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
~ 25	100	3 ASTIER	67	HBC	±
57±13	143	4 ROSENFELD	65	RVUE	±
1 T-matrix pole on sheet II, the pole on sheet III is at 1006-i49 MeV.					
2 T-matrix pole.					
3 ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.					
4 Plus systematic errors.					

NODE=M036W2

NODE=M036W2

 $a_0(980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta\pi$	seen
$\Gamma_2 K\bar{K}$	seen
$\Gamma_3 \rho\pi$	not seen
$\Gamma_4 \gamma\gamma$	seen
$\Gamma_5 e^+e^-$	

 $a_0(980)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	Γ_4
VALUE (keV)	DOCUMENT ID TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.30±0.10	1 AMSLER 98 RVUE
1 Using $\Gamma_{\gamma\gamma} B(a_0(980) \rightarrow \eta\pi) = 0.24 \pm 0.08$ keV.	

DESIG=1;OUR EST; \rightarrow UNCHECKED ←
DESIG=3;OUR EST; \rightarrow UNCHECKED ←
DESIG=2;OUR EST; \rightarrow UNCHECKED ←
DESIG=5;OUR EST; \rightarrow UNCHECKED ←
DESIG=6

NODE=M036215;NODE=M036

 $a_0(980)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_4/\Gamma$
VALUE (keV)	DOCUMENT ID TECN COMMENT

0.21 +0.08 -0.04 OUR AVERAGE

0.128 +0.003 +0.502 -0.002 -0.043	1 UEHARA 09A BELL $\gamma\gamma \rightarrow \pi^0\eta$
0.28 ± 0.04 ± 0.10	44 OEST 90 JADE $e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.19 ± 0.07 +0.10 -0.07	ANTREASYAN 86 CBAL $e^+e^- \rightarrow e^+e^-\pi^0\eta$

1 From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

NODE=M036217

NODE=M036W4
NODE=M036W4

NODE=M036W4;LINKAGE=A

NODE=M036220

NODE=M036G1
NODE=M036G1

NODE=M036G1;LINKAGE=UE

NODE=M036G2
NODE=M036G2

NODE=M036225

NODE=M036R2
NODE=M036R2 **$a_0(980)$ BRANCHING RATIOS**

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$	Γ_2/Γ_1
VALUE	DOCUMENT ID TECN CHG COMMENT

0.183±0.024 OUR AVERAGE Error includes scale factor of 1.2.

0.57 ± 0.16	1 BARGIOTTI 03 OBLX $\bar{p}p$
0.23 ± 0.05	2 ABELE 98 CBAR $0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
0.166±0.01 ±0.02	3 BARBERIS 98C OMEG $450 pp \rightarrow p_f f_1(1285)p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
1.20 ± 0.15	4 ANISOVICH 09 RVUE $0.0 \bar{p}p, \pi N$
1.05 ± 0.07 ± 0.05	5 BUGG 08A RVUE 0 $\bar{p}p \rightarrow \pi^0\pi^0\eta$
~ 0.60	OLLER 99B RVUE $\pi\pi \rightarrow \eta\pi, K\bar{K}$
0.7 ± 0.3	3 CORDEN 78 OMEG $12-15 \pi^- p \rightarrow n\eta/2\pi$
0.25 ± 0.08	3 DEFOIX 72 HBC ± $0.7 \bar{p} \rightarrow 7\pi$

$\Gamma(\rho\pi)/\Gamma(\eta\pi)$ $\rho\pi$ forbidden.

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.25	70	AMMAR	70	HBC	\pm 4.1, 5.5 $K^- p \rightarrow \Lambda\eta 2\pi$
1 Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.					
2 Using $\pi^0 \pi^0 \eta$ from AMSLER 94D.					
3 From the decay of $f_1(1285)$.					
4 This is a ratio of couplings.					
5 A ratio of couplings, using AMSLER 94D and ABELE 98. Supersedes BUGG 94.					

 Γ_3/Γ_1

NODE=M036R1

NODE=M036R1

NODE=M036R1

NODE=M036R;LINKAGE=BG

NODE=M036R2;LINKAGE=Q

NODE=M036R2;LINKAGE=L

NODE=M036R2;LINKAGE=AN

NODE=M036R2;LINKAGE=BU

a₀(980) REFERENCES

AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
BUGG	08A	PR D78 074023	D.V. Bugg	(LOQM)	REFID=52578
ACHASOV	03B	PR D68 014006	N.N. Achasov, A.V. Kiselev		REFID=49476
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48574
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
TEIGE	99	PR D59 012001	S. Teige <i>et al.</i>	(BNL E852 Collab.)	REFID=47386
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46613
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=45863
AMSLER	98	RMP 70 1293	C. Amsler		REFID=46317
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46601
Translated from UFN 168 481.					
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46331
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46346
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=46351
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44507
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44508
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
VOROB'YEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
Translated from YAF 48 436.					
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20624
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
DEBILLY	80	NP B176 1	L. de Billy <i>et al.</i>	(CURIN, LAUS, NEUC+)	REFID=20461
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
CONFORTO	78	LNC 23 419	B. Conforto <i>et al.</i>	(RHEL, TNTO, CHIC+)	REFID=20451
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20452
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
JAFFE	77	PR D15 267,281	R. Jaffe	(MIT)	REFID=43673
FLATTE	76	PL 63B 224	S.M. Flatté	(CERN)	REFID=20446
GAY	76B	PL 63B 220	J.B. Gay <i>et al.</i>	(CERN, AMST, NIJM) JP	REFID=20445
WELLS	75	NP B101 333	J. Wells <i>et al.</i>	(OXF)	REFID=20444
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
AMMAR	70	PR D2 430	R. Ammar <i>et al.</i>	(KANS, NWES, ANL, WISC)	REFID=20428
BARNES	69C	PRL 23 610	V.E. Barnes <i>et al.</i>	(BNL, SYRA)	REFID=20418
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)	REFID=20419
MILLER	69B	PL 29B 255	D.H. Miller <i>et al.</i>	(PURD)	REFID=20424
Also		PR 188 2011	W.L. Yen <i>et al.</i>	(PURD)	REFID=20425
AMMAR	68	PRL 21 1832	R. Ammar <i>et al.</i>	(NWES, ANL)	REFID=20412
ASTIER	67	PL 25B 294	A. Astier <i>et al.</i>	(CDEF, CERN, IRAD)	REFID=20405
Includes data of BARLOW 67, CONFORTO 67, and ARMENTEROS 65.					
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
CONFORTO	67	NP B3 469	G. Conforto <i>et al.</i>	(CERN, CDEF, IPNP+)	REFID=20411
ARMENTEROS	65	PL 17 344	R. Armenteros <i>et al.</i>	(CERN, CDEF)	REFID=20396
ROSENFELD	65	Oxford Conf. 58	A.H. Rosenfeld	(LRL)	REFID=20399

NODE=M036

NODE=M004

 $\phi(1020)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\phi(1020)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	NODE
1019.461 ± 0.016 OUR AVERAGE					NODE=M004M
1019.463 ± 0.061	2.3M	1 KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$, $K_S^0 K_L^0$	OCCUR=2
1019.462 $\pm 0.042 \pm 0.056$	28k	2 LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$	
1019.51 $\pm 0.02 \pm 0.05$		3 LEES	13Q	BABR $e^+ e^- \rightarrow K^+ K^- \gamma$	
1019.30 $\pm 0.02 \pm 0.10$	105k	AKHMETSHIN 06	CMD2	0.98-1.06 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
1019.52 $\pm 0.05 \pm 0.05$	17.4k	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+ e^- \rightarrow \eta \gamma$	
1019.483 $\pm 0.011 \pm 0.025$	272k	4 AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$	
1019.42 ± 0.05	1900k	5 ACHASOV	01E	SND $e^+ e^- \rightarrow K^+ K^-$, $K_S^0 K_L^0, \pi^+ \pi^- \pi^0$	
1019.40 $\pm 0.04 \pm 0.05$	23k	AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta \gamma$	
1019.36 ± 0.12		6 ACHASOV	00B	SND $e^+ e^- \rightarrow \eta \gamma$	
1019.38 $\pm 0.07 \pm 0.08$	2200	7 AKHMETSHIN 99F	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \geq 2\gamma$	
1019.51 $\pm 0.07 \pm 0.10$	11169	AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
1019.5 ± 0.4		BARBERIS	98	OMEG 450 $p p \rightarrow p p 2K^+ 2K^-$	
1019.42 ± 0.06	55600	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow$ hadrons	
1019.7 ± 0.3	2012	DAVENPORT	86	MPSF 400 $pA \rightarrow 4KX$	
1019.7 $\pm 0.1 \pm 0.1$	5079	ALBRECHT	85D	ARG 10 $e^+ e^- \rightarrow K^+ K^- X$	
1019.3 ± 0.1	1500	ARENTON	82	AEMS 11.8 polar. $p p \rightarrow KK$	
1019.67 ± 0.17	25080	8 PELLINEN	82	RVUE	
1019.52 ± 0.13	3681	BUKIN	78C	OLYA $e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1019.54 $\pm 0.10 \pm 0.51$		9 AAIJ	19H	LHCb $p p \rightarrow D^\pm X$	OCCUR=2
1019.469 ± 0.061	1.7M	KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$	
1019.457 ± 0.061	610k	KOZYREV	16	CMD3 $e^+ e^- \rightarrow K_S^0 K_L^0$	
1019.48 ± 0.01		LEES	13F	BABR $D^+ \rightarrow K^+ K^- \pi^+$	
1019.441 $\pm 0.008 \pm 0.080$	542k	10 AKHMETSHIN 08	CMD2	1.02 $e^+ e^- \rightarrow K^+ K^-$	
1019.63 ± 0.07	12540	11 AUBERT,B	05J	BABR $D^0 \rightarrow \bar{K}^0 K^+ K^-$	
1019.8 ± 0.7		ARMSTRONG	86	OMEG 85 $\pi^+ / p p \rightarrow \pi^+ / p 4K p$	
1020.1 ± 0.11	5526	11 ATKINSON	86	OMEG 20-70 γp	
1019.7 ± 1.0		BEBEK	86	CLEO $e^+ e^- \rightarrow \Upsilon(4S)$	
1019.411 ± 0.008	642k	12 DIJKSTRA	86	SPEC 100-200 $\pi^\pm, \bar{p}, p, K^\pm$, on Be	
1020.9 ± 0.2		11 FRAME	86	OMEG 13 $K^+ p \rightarrow \phi K^+ p$	
1021.0 ± 0.2		11 ARMSTRONG	83B	OMEG 18.5 $K^- p \rightarrow K^- K^+ \Lambda$	
1020.0 ± 0.5		11 ARMSTRONG	83B	OMEG 18.5 $K^- p \rightarrow K^- K^+ \Lambda$	OCCUR=2
1019.7 ± 0.3		11 BARATE	83	GOLI 190 $\pi^- Be \rightarrow 2\mu X$	
1019.8 $\pm 0.2 \pm 0.5$	766	IVANOV	81	OLYA 1-1.4 $e^+ e^- \rightarrow K^+ K^-$	
1019.4 ± 0.5	337	COOPER	78B	HBC 0.7-0.8 $\bar{p} p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$	
1020 ± 1	383	11 BALDI	77	CNTR 10 $\pi^- p \rightarrow \pi^- \phi p$	OCCUR=2
1018.9 ± 0.6	800	COHEN	77	ASPK 6 $\pi^\pm N \rightarrow K^+ K^- N$	
1019.7 ± 0.5	454	KALBFLEISCH	76	HBC 2.18 $K^- p \rightarrow \Lambda K \bar{K}$	
1019.4 ± 0.8	984	BESCH	74	CNTR 2 $\gamma p \rightarrow p K^+ K^-$	
1020.3 ± 0.4	100	BALLAM	73	HBC 2.8-9.3 γp	
1019.4 ± 0.7		BINNIE	73B	CNTR $\pi^- p \rightarrow \phi n$	
1019.6 ± 0.5	120	13 AGUILAR...	72B	HBC 3.9,4.6 $K^- p \rightarrow \Lambda K^+ K^-$	
1019.9 ± 0.5	100	13 AGUILAR...	72B	HBC 3.9,4.6 $K^- p \rightarrow K^- p K^+ K^-$	OCCUR=2
1020.4 ± 0.5	131	COLLEY	72	HBC 10 $K^- p \rightarrow K^+ p \phi$	
1019.9 ± 0.3	410	STOTTLE...	71	HBC 2.9 $K^- p \rightarrow \Sigma/\Lambda K \bar{K}$	

- 1 Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.
- 2 Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.
- 3 Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.
- 4 Update of AKHMETSHIN 99D
- 5 From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
- 6 Using a total width of 4.43 ± 0.05 MeV. Systematic uncertainty included.
- 7 Using a total width of 4.43 ± 0.05 MeV.
- 8 PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DEGROOT 74.
- 9 From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.
- 10 Strongly correlated with AKHMETSHIN 04.
- 11 Systematic errors not evaluated.
- 12 Weighted and scaled average of 12 measurements of DIJKSTRA 86.
- 13 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

NODE=M004M;LINKAGE=G
 NODE=M004M;LINKAGE=E
 NODE=M004M;LINKAGE=C
 NODE=M004M;LINKAGE=GS
 NODE=M004M;LINKAGE=AE
 NODE=M004M;LINKAGE=G2
 NODE=M004M;LINKAGE=F2
 NODE=M004M;LINKAGE=R
 NODE=M004M;LINKAGE=I
 NODE=M004M;LINKAGE=AH
 NODE=M004M;LINKAGE=A
 NODE=M004M;LINKAGE=B
 NODE=M004M;LINKAGE=D

$\phi(1020)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.249±0.013 OUR AVERAGE				Error includes scale factor of 1.1.
4.245±0.013	2.3M	1 KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+K^-$, $K_S^0 K_L^0$
4.205±0.103±0.067	28k	2 LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
4.29 ± 0.04 ± 0.07		3 LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$
4.30 ± 0.06 ± 0.17	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.280±0.033±0.025	272k	4 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
4.21 ± 0.04	1900k	5 ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-$, $K_S K_L$, $\pi^+\pi^-\pi^0$
4.44 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
4.5 ± 0.7	1500	ARENTON	82	AEMS 11.8 polar. $p p \rightarrow K K$
4.2 ± 0.6	766	6 IVANOV	81	OLYA 1–1.4 $e^+e^- \rightarrow K^+K^-$
4.3 ± 0.6		6 CORDIER	80	DM1 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.36 ± 0.29	3681	6 BUKIN	78C	OLYA $e^+e^- \rightarrow$ hadrons
4.4 ± 0.6	984	6 BESCH	74	CNTR $2\gamma p \rightarrow p K^+K^-$
4.67 ± 0.72	681	6 BALAKIN	71	OSPK $e^+e^- \rightarrow$ hadrons
4.09 ± 0.29		BIZOT	70	OSPK $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.249±0.015	1.7M	KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+K^-$
4.240±0.017	610k	KOZYREV	16	CMD3 $e^+e^- \rightarrow K_S^0 K_L^0$
4.37 ± 0.02		LEES	13F	BABR $D^+ \rightarrow K^+K^-\pi^+$
4.24 ± 0.02 ± 0.03	542k	7 AKHMETSHIN 08	CMD2	1.02 $e^+e^- \rightarrow K^+K^-$
4.28 ± 0.13	12540	8 AUBERT,B	05J	BABR $D^0 \rightarrow \bar{K}^0 K^+K^-$
4.45 ± 0.06	271k	DIJKSTRA	86	SPEC 100 π^- Be
3.6 ± 0.8	337	6 COOPER	78B	HBC 0.7–0.8 $\bar{p}p \rightarrow$ $K_S^0 K_L^0 \pi^+\pi^-$
4.5 ± 0.50	1300	6,8 AKERLOF	77	SPEC 400 $pA \rightarrow K^+K^-X$
4.5 ± 0.8	500	6,8 AYRES	74	ASPK 3–6 $\pi^-p \rightarrow$ K^+K^-n , $K^-p \rightarrow$ $K^+K^-\Lambda/\Sigma^0$
3.81 ± 0.37		COSME	74B	OSPK $e^+e^- \rightarrow K_L^0 K_S^0$
3.8 ± 0.7	454	6 BORENSTEIN	72	HBC 2.18 $K^-p \rightarrow K\bar{K}n$

- 1 Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.
- 2 Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.
- 3 Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.
- 4 Update of AKHMETSHIN 99D
- 5 From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
- 6 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- 7 Strongly correlated with AKHMETSHIN 04.
- 8 Systematic errors not evaluated.

NODE=M004W;LINKAGE=G
 NODE=M004W;LINKAGE=E
 NODE=M004W;LINKAGE=C
 NODE=M004W;LINKAGE=GS
 NODE=M004W;LINKAGE=AE
 NODE=M004W;LINKAGE=D
 NODE=M004W;LINKAGE=AH
 NODE=M004W;LINKAGE=A

$\phi(1020)$ DECAY MODES

NODE=M004215;NODE=M004

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
$\Gamma_1 K^+ K^-$	(49.2 \pm 0.5) %	S=1.3	
$\Gamma_2 K_L^0 K_S^0$	(34.0 \pm 0.4) %	S=1.3	
$\Gamma_3 \rho\pi + \pi^+\pi^-\pi^0$	(15.24 \pm 0.33) %	S=1.2	
$\Gamma_4 \rho\pi$			DESIG=16
$\Gamma_5 \pi^+\pi^-\pi^0$			DESIG=3
$\Gamma_6 \eta\gamma$	(1.303 \pm 0.025) %	S=1.2	DESIG=4
$\Gamma_7 \pi^0\gamma$	(1.30 \pm 0.05) $\times 10^{-3}$		DESIG=7
$\Gamma_8 \ell^+\ell^-$	—		DESIG=256;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_9 e^+e^-$	(2.973 \pm 0.034) $\times 10^{-4}$	S=1.3	DESIG=5
$\Gamma_{10} \mu^+\mu^-$	(2.86 \pm 0.19) $\times 10^{-4}$		DESIG=6
$\Gamma_{11} \eta e^+e^-$	(1.08 \pm 0.04) $\times 10^{-4}$		DESIG=17
$\Gamma_{12} \pi^+\pi^-$	(7.3 \pm 1.3) $\times 10^{-5}$		DESIG=8
$\Gamma_{13} \omega\pi^0$	(4.7 \pm 0.5) $\times 10^{-5}$		DESIG=25
$\Gamma_{14} \omega\gamma$	< 5 %	CL=84%	DESIG=10
$\Gamma_{15} \rho\gamma$	< 1.2 $\times 10^{-5}$	CL=90%	DESIG=12
$\Gamma_{16} \pi^+\pi^-\gamma$	(4.1 \pm 1.3) $\times 10^{-5}$		DESIG=9
$\Gamma_{17} f_0(980)\gamma$	(3.22 \pm 0.19) $\times 10^{-4}$	S=1.1	DESIG=20
$\Gamma_{18} \pi^0\pi^0\gamma$	(1.12 \pm 0.06) $\times 10^{-4}$		DESIG=19
$\Gamma_{19} \pi^+\pi^-\pi^+\pi^-$	(3.9 \pm 2.8) $\times 10^{-6}$		DESIG=15
$\Gamma_{20} \pi^+\pi^+\pi^-\pi^-\pi^0$	< 4.6 $\times 10^{-6}$	CL=90%	DESIG=14
$\Gamma_{21} \pi^0e^+e^-$	(1.33 \pm 0.07) $\times 10^{-5}$		DESIG=21
$\Gamma_{22} \pi^0\eta\gamma$	(7.27 \pm 0.30) $\times 10^{-5}$	S=1.5	DESIG=22
$\Gamma_{23} a_0(980)\gamma$	(7.6 \pm 0.6) $\times 10^{-5}$		DESIG=23
$\Gamma_{24} K^0\bar{K}^0\gamma$	< 1.9 $\times 10^{-8}$	CL=90%	DESIG=257
$\Gamma_{25} \eta'(958)\gamma$	(6.22 \pm 0.21) $\times 10^{-5}$		DESIG=194
$\Gamma_{26} \eta\pi^0\pi^0\gamma$	< 2 $\times 10^{-5}$	CL=90%	DESIG=195
$\Gamma_{27} \mu^+\mu^-\gamma$	(1.4 \pm 0.5) $\times 10^{-5}$		DESIG=196
$\Gamma_{28} \rho\gamma\gamma$	< 1.2 $\times 10^{-4}$	CL=90%	DESIG=250
$\Gamma_{29} \eta\pi^+\pi^-$	< 1.8 $\times 10^{-5}$	CL=90%	DESIG=255
$\Gamma_{30} \eta\mu^+\mu^-$	< 9.4 $\times 10^{-6}$	CL=90%	DESIG=26
$\Gamma_{31} \eta U \rightarrow \eta e^+e^-$	< 1 $\times 10^{-6}$	CL=90%	DESIG=259
Γ_{32} invisible	< 1.7 $\times 10^{-4}$	CL=90%	DESIG=260
Lepton Family number (LF) violating modes			
$\Gamma_{33} e^\pm\mu^\mp$	$LF < 2$	$\times 10^{-6}$	CL=90% NODE=M004;CLUMP=A DESIG=258

CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 82 measurements and one constraint to determine 14 parameters. The overall fit has a $\chi^2 = 63.7$ for 69 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-78									
x_3	-59 -4									
x_6	-23 19 6									
x_7	-15 14 4 10									
x_9	54 -52 -17 -38 -27									
x_{10}	-7 7 2 5 3 -13									
x_{12}	-3 3 1 2 2 -6 1									
x_{13}	-5 4 1 3 2 -8 1 1									
x_{17}	0 0 0 0 0 0 0 0 0									
x_{18}	-11 10 3 19 5 -20 2 1 2 0									
x_{19}	-1 1 0 1 0 -2 0 0 0 0									
x_{23}	0 0 0 0 0 0 0 0 0 0									
x_{25}	-8 6 2 33 3 -12 2 1 1 0									
	x_1	x_2	x_3	x_6	x_7	x_9	x_{10}	x_{12}	x_{13}	x_{17}
x_{19}	0									
x_{23}	0 0									
x_{25}	6 0 0									
	x_{18}	x_{19}	x_{23}							

$\phi(1020)$ PARTIAL WIDTHS

$\Gamma(\eta\gamma)$

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

Γ_6

• • • We do not use the following data for averages, fits, limits, etc. • • •

$58.9 \pm 0.5 \pm 2.4$

ACHASOV

00

SND

$e^+ e^- \rightarrow \eta\gamma$

$\Gamma(\pi^0\gamma)$

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

Γ_7

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.40 \pm 0.16^{+0.43}_{-0.40}$

ACHASOV

00

SND

$e^+ e^- \rightarrow \pi^0\gamma$

$\Gamma(\ell^+\ell^-)$

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

Γ_8

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.320 \pm 0.017 \pm 0.015$

¹ AMBROSINO

05

KLOE

$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$

¹ Weighted average of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ from AMBROSINO 05 assuming lepton universality.

$\Gamma(e^+e^-)$

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

Γ_9

1.27 ± 0.04 OUR EVALUATION

1.251 ± 0.021 OUR AVERAGE

Error includes scale factor of 1.1.

$1.235 \pm 0.006 \pm 0.022$

¹ AKHMETSHIN

11

CMD2

$1.02 e^+ e^- \rightarrow \phi$

$1.32 \pm 0.05 \pm 0.03$

² AMBROSINO

05

KLOE

$1.02 e^+ e^- \rightarrow e^+ e^-$

1.28 ± 0.05

AKHMETSHIN

95

CMD2

$1.02 e^+ e^- \rightarrow \phi$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$, $\eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² From forward-backward asymmetry and using $\Gamma_{\text{total}} = 4.26 \pm 0.05$ MeV from the 2004 edition of this Review.

NODE=M004218

NODE=M004W6

NODE=M004W6

NODE=M004W7

NODE=M004W7

NODE=M004W5

NODE=M004W5

NODE=M004W5;LINKAGE=AM

NODE=M004W8

NODE=M004W8

→ UNCHECKED ←

NODE=M004W8;LINKAGE=AK

NODE=M004W8;LINKAGE=AM

$(\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-))^{1/2}$	$(\Gamma_9 \Gamma_{10})^{1/2}$			
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
$1.320 \pm 0.018 \pm 0.017$	AMBROSINO 05	KLOE	$1.02 e^+ e^- \rightarrow \mu^+ \mu^-$	

NODE=M004W9
NODE=M004W9

$\phi(1020) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$				
$\Gamma(K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_1 \Gamma_9/\Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.6340 \pm 0.0070 \pm 0.0039$	1 LEES	13Q BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.669 \pm 0.001 \pm 0.023$	1.7M	KOZYREV	18 CMD3	$e^+ e^- \rightarrow K^+ K^-$

¹ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.

$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_9/\Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.4200 \pm 0.0033 \pm 0.0123$	28k	1 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$

¹ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

NODE=M004G01;LINKAGE=A

$\phi(1020) \Gamma(i) \Gamma(e^+ e^-)/\Gamma^2(\text{total})$				
$\Gamma(K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma \times \Gamma_9/\Gamma$			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
14.63 ± 0.29 OUR FIT				Error includes scale factor of 1.5.
14.6 ± 0.5 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.

15.789 ± 0.541	1.7M	KOZYREV	18 CMD3	$e^+ e^- \rightarrow K^+ K^-$
14.27 ± 0.05	± 0.31	542k	AKHMETSHIN 08	CMD2 $1.02 e^+ e^- \rightarrow K^+ K^-$
13.93 ± 0.14	± 0.99	1000k	¹ ACHASOV 01E SND	$e^+ e^- \rightarrow K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.


NODE=M004GXX
NODE=M004GXX

NODE=M004GXX;LINKAGE=A

NODE=M004224

NODE=M004G10
NODE=M004G10

NODE=M004G10;LINKAGE=AE



$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma \times \Gamma_9/\Gamma$			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
10.10 ± 0.12 OUR FIT	Error includes scale factor of 1.1.			
10.07 ± 0.13 OUR AVERAGE				
10.078 ± 0.223	610k	1 KOZYREV	16 CMD3	$e^+ e^- \rightarrow K_S^0 K_L^0$
10.01 ± 0.04 ± 0.17	272k	2 AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
10.27 ± 0.07 ± 0.34	500k	3 ACHASOV	01E SND	$e^+ e^- \rightarrow K^+ K^-$, $K_S K_L, \pi^+ \pi^- \pi^0$

1 KOZYREV 16 also reports $\Gamma(e^+ e^-) B(\phi \rightarrow K_S^0 K_L^0) = (0.428 \pm 0.001 \pm 0.009)$ keV.

2 Update of AKHMETSHIN 99D

3 From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_9/\Gamma$			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT

4.53 ± 0.10 OUR FIT Error includes scale factor of 1.1.

4.46 ± 0.12 OUR AVERAGE

4.51 ± 0.16 ± 0.11	105k	AKHMETSHIN 06	CMD2	$0.98-1.06 \pi^+ \pi^- \pi^0$
4.30 ± 0.08 ± 0.21	AUBERT,B	04N BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
4.665 ± 0.042 ± 0.261	400k	1 ACHASOV	01E SND	$e^+ e^- \rightarrow K^+ K^-$, $K_S K_L, \pi^+ \pi^- \pi^0$
4.35 ± 0.27 ± 0.08	11169	2 AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.38 ± 0.12	BENAYOUN	10 RVUE	0.4-1.05	$e^+ e^-$
-----------------	----------	---------	----------	-----------

1 From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

2 Recalculated by us from the cross section in the peak.

$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma \times \Gamma_9/\Gamma$			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT

3.87 ± 0.07 OUR FIT Error includes scale factor of 1.2.

3.93 ± 0.09 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

4.050 ± 0.067 ± 0.118	33k	1 ACHASOV	07B SND	$0.6-1.38 e^+ e^- \rightarrow \eta\gamma$
4.093 ± 0.040 ± 0.247	17.4k	2 AKHMETSHIN 05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta\gamma$
3.850 ± 0.041 ± 0.159	23k	3,4 AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
4.00 ± 0.04 ± 0.11	5 ACHASOV	00 SND	$e^+ e^- \rightarrow \eta\gamma$	
3.53 ± 0.08 ± 0.17	2200	6,7 AKHMETSHIN 99F	CMD2	$e^+ e^- \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.19 ± 0.06	8 BENAYOUN	10 RVUE	0.4-1.05	$e^+ e^-$
-----------------	------------	---------	----------	-----------

1 From a combined fit of $\sigma(e^+ e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.

2 From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

3 From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

4 The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

5 From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$.

6 Recalculated by the authors from the cross section in the peak.

7 From the $\eta \rightarrow \pi^+ \pi^- \pi^0$ decay and using $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (23.1 \pm 0.5) \times 10^{-2}$.

8 A simultaneous fit of $e^+ e^- \rightarrow \pi^+ \pi^-$, $\pi^+ \pi^- \pi^0$, $\pi^0 \gamma$, $\eta\gamma$ data.

NODE=M004G6
NODE=M004G6

NODE=M004G6;LINKAGE=A
NODE=M004G;LINKAGE=GS
NODE=M004G6;LINKAGE=AE

NODE=M004G7
NODE=M004G7

NODE=M004G7;LINKAGE=AE

NODE=M004G;LINKAGE=B


NODE=M004G2
NODE=M004G2

NODE=M004G2;LINKAGE=AH

NODE=M004G2;LINKAGE=AK
NODE=M004G;LINKAGE=AK
NODE=M004G;LINKAGE=BQ

NODE=M004G2;LINKAGE=A
NODE=M004G;LINKAGE=A
NODE=M004G2;LINKAGE=C
NODE=M004G7;LINKAGE=BE

WEIGHTED AVERAGE
3.93±0.09 (Error scaled by 1.3)



$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma \times \Gamma_9/\Gamma$$

$$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_7/\Gamma \times \Gamma_9/\Gamma$$

3.88±0.14 OUR FIT

NODE=M004G3
NODE=M004G3

3.87±0.15 OUR AVERAGE

4.04±0.09±0.19

¹ ACHASOV 16A SND 0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$

3.75±0.11±0.29

AKHMETSHIN 05 CMD2 0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$

3.67±0.10^{+0.27}
-0.25

² ACHASOV 00 SND $e^+e^- \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.29±0.11

³ BENAYOUN 10 RVUE 0.4–1.05 e^+e^-

1 From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$ resonances, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 00.

NODE=M004G3;LINKAGE=B

2 From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\pi^0 \rightarrow 2\gamma) = (98.798 \pm 0.032) \times 10^{-2}$.

NODE=M004G3;LINKAGE=A

3 A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.

NODE=M004G3;LINKAGE=BE

$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$$

**8.5^{+0.5}
-0.6 OUR FIT**

NODE=M004G5

8.8 ±0.9 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

NODE=M004G5

8.36±0.59±0.37

ACHASOV 01G SND $e^+e^- \rightarrow \mu^+\mu^-$

9.9 ±1.4 ±0.9

¹ ACHASOV 99C SND $e^+e^- \rightarrow \mu^+\mu^-$

14.4 ±3.0

² VASSERMAN 81 OLYA $e^+e^- \rightarrow \mu^+\mu^-$

8.6 ±5.9


² AUGUSTIN 73 OSPK $e^+e^- \rightarrow \mu^+\mu^-$

1 Recalculated by the authors from the cross section in the peak.

NODE=M004G5;LINKAGE=A

2 Recalculated by us from the cross section in the peak.

NODE=M004G5;LINKAGE=B



$$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

2.2 ±0.4 OUR FIT

2.2 ±0.4 OUR AVERAGE

$2.1 \pm 0.3 \pm 0.3$ ¹ ACHASOV 00C SND $e^+ e^- \rightarrow \pi^+ \pi^-$

$1.95^{+1.15}_{-0.87}$ ² GOLUBEV 86 ND $e^+ e^- \rightarrow \pi^+ \pi^-$

$6.01^{+3.19}_{-2.51}$ ² VASSERMAN 81 OLYA $e^+ e^- \rightarrow \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.31 ± 0.99 ³ BENAYOUN 13 RVUE $0.4-1.05 e^+ e^-$

¹ Recalculated by the authors from the cross section in the peak.

² Recalculated by us from the cross section in the peak.

³ A simultaneous fit to $e^+ e^- \rightarrow \pi^+ \pi^-$, $\pi^+ \pi^- \pi^0$, $\pi^0 \gamma$, $\eta \gamma$, $K\bar{K}$, and $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$ data.

$$\Gamma(\omega \pi^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

1.40±0.15 OUR FIT

1.37±0.17±0.01

^{1,2} AMBROSINO 08G KLOE $e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0, 2\pi^0 \gamma$

¹ Recalculated by the authors from the cross section at the peak.

² AMBROSINO 08G reports $[\Gamma(\phi(1020) \rightarrow \omega \pi^0)/\Gamma_{\text{total}} \times \Gamma(\phi(1020) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\pi^0 \pi^0 \gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

3.34±0.17 OUR FIT

$3.33^{+0.04+0.19}_{-0.09-0.20}$

¹ AMBROSINO 07 KLOE $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

¹ Calculated by the authors from the cross section at the peak.

$$\Gamma(\pi^+ \pi^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-9}) EVTS DOCUMENT ID TECN COMMENT

1.2 +0.8 -0.7 OUR FIT

1.17±0.52±0.64 3285 ¹ AKHMETSHIN 00E CMD2 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$

¹ Recalculated by the authors from the cross section in the peak.

NODE=M004G4
NODE=M004G4

NODE=M004G4;LINKAGE=AC
NODE=M004G4;LINKAGE=B
NODE=M004G4;LINKAGE=A

NODE=M004G11
NODE=M004G11

NODE=M004G11;LINKAGE=AB
NODE=M004G11;LINKAGE=AM

NODE=M004G9
NODE=M004G9

NODE=M004G9;LINKAGE=AM

NODE=M004G8
NODE=M004G8

NODE=M004G8;LINKAGE=A

φ(1020) BRANCHING RATIOS

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.492±0.005 OUR FIT Error includes scale factor of 1.3.					NODE=M004R220
0.493±0.010 OUR AVERAGE					NODE=M004R1 NODE=M004R1
0.492±0.012	2913	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow K^+ K^-$	
0.44 ± 0.05	321	KALBFLEISCH 76	HBC	$2.18 K^- p \rightarrow \Lambda K^+ K^-$	
0.49 ± 0.06	270	DEGROOT 74	HBC	$4.2 K^- p \rightarrow \Lambda \phi$	
0.540±0.034	565	BALAKIN 71	OSPK	$e^+ e^- \rightarrow K^+ K^-$	
0.48 ± 0.04	252	LINDSEY 66	HBC	$2.1-2.7 K^- p \rightarrow \Lambda K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.493±0.003±0.007	1	AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow K^+ K^-$	
0.476±0.017	1000k	ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$	

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$, $\eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.340±0.004 OUR FIT Error includes scale factor of 1.3.					NODE=M004R1;LINKAGE=AK
0.331±0.009 OUR AVERAGE					NODE=M004R1;LINKAGE=B2
0.335±0.010	40644	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$	
0.326±0.035	DOLINSKY 91	ND		$e^+ e^- \rightarrow K_L^0 K_S^0$	
0.310±0.024	DRUZHININ 84	ND		$e^+ e^- \rightarrow K_L^0 K_S^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.336±0.002±0.006	1	AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow K_S^0 K_L^0$	
0.351±0.013	500k	ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$	
0.27 ± 0.03	133	KALBFLEISCH 76	HBC	$2.18 K^- p \rightarrow \Lambda K_L^0 K_S^0$	
0.257±0.030	95	BALAKIN 71	OSPK	$e^+ e^- \rightarrow K_L^0 K_S^0$	
0.40 ± 0.04	167	LINDSEY 66	HBC	$2.1-2.7 K^- p \rightarrow \Lambda K_L^0 K_S^0$	

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-$, $K_S^0 K_L^0$, $\pi^+ \pi^- \pi^0$, $\eta \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Balakin error increased by Paul.

$\Gamma(K_L^0 K_S^0)/\Gamma(K^+ K^-)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.690±0.015 OUR FIT Error includes scale factor of 1.3.					NODE=M004R19 NODE=M004R19
0.740±0.031 OUR AVERAGE					
0.70 ± 0.06	2732	BUKIN	78C OLYA	$e^+ e^- \rightarrow K_L^0 K_S^0$	
0.82 ± 0.08	LOSTY	78	HBC	$4.2 K^- p \rightarrow \phi \text{ hyperon}$	
0.71 ± 0.05	LAVEN	77	HBC	$10 K^- p \rightarrow K^+ K^- \Lambda$	
0.71 ± 0.08	LYONS	77	HBC	$3-4 K^- p \rightarrow \Lambda \phi$	
0.89 ± 0.10	144	AGUILAR...	72B HBC	$3.9, 4.6 K^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.638±0.022	2.3M	1 KOZYREV	18	CMD3 $e^+ e^- \rightarrow K_L^0 K_S^0, K^+ K^-$	
0.68 ± 0.03	2 AKHMETSHIN 95	CMD2		$e^+ e^- \rightarrow K_L^0 K_S^0, K^+ K^-$	

¹ The prediction taking into account phase-space difference, radiative corrections, isospin breaking, and the Sommerfeld-Gamow-Sakharov factor gives 0.630.

² Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNSKIY 07. BENAYOUN 12 obtains 0.71 ± 0.01 in the HLS model.

NODE=M004R220

NODE=M004R1
NODE=M004R1

NODE=M004R1;LINKAGE=AK

NODE=M004R1;LINKAGE=B2

NODE=M004R2
NODE=M004R2

NODE=M004R2;LINKAGE=AK

NODE=M004R2;LINKAGE=B2

NODE=M004R2;LINKAGE=01

NODE=M004R19
NODE=M004R19

NODE=M004R19;LINKAGE=A

NODE=M004R19;LINKAGE=KH

$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.408±0.005 OUR FIT		Error includes scale factor of 1.3.		
0.45 ±0.04 OUR AVERAGE				
0.44 ±0.07		1 LONDON	66 HBC	$2.24 K^- p \rightarrow \Lambda K\bar{K}$

0.48 ±0.07 52 BADIER 65B HBC $3 K^- p$

0.40 ±0.10 34 SCHLEIN 63 HBC $1.95 K^- p \rightarrow \Lambda K\bar{K}$

1 This is probably not affected by their controversial background subtraction; the value is from their numbers of $K_1 K_2$ vs $K^+ K^-$ events.

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1524±0.0033 OUR FIT		Error includes scale factor of 1.2.		
0.151 ±0.009 OUR AVERAGE				Error includes scale factor of 1.7.
0.161 ±0.008	11761	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

0.143 ±0.007 DOLINSKY 91 ND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.155 ±0.002 ±0.005	1 AKHMETSHIN 11	CMD2	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.159 ±0.008	400k	2 ACHASOV 01E	SND $e^+ e^- \rightarrow K^+ K^-$, $K_S K_L, \pi^+ \pi^- \pi^0$
0.145 ±0.009 ±0.003	11169	3 AKHMETSHIN 98	CMD2 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.139 ±0.007		4 PARROUR 76B	OSPK $e^+ e^-$

1 Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

2 Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

3 Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

4 Using $\Gamma(\phi) = 4.1$ MeV. If interference between the $\rho\pi$ and 3π modes is neglected, the fraction of the $\rho\pi$ is more than 80% at the 90% confidence level.

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K^+ K^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.310±0.009 OUR FIT		Error includes scale factor of 1.2.		
0.28 ±0.09	34	AGUILAR---	72B HBC	$3.9, 4.6 K^- p$

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K\bar{K})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.183±0.005 OUR FIT		Error includes scale factor of 1.2.		
0.24 ±0.04 OUR AVERAGE				

0.237±0.039 CERRADA 77B HBC $4.2 K^- p \rightarrow \Lambda 3\pi$

0.30 ±0.15 LONDON 66 HBC $2.24 K^- p \rightarrow \Lambda \pi^+ \pi^- \pi^0$

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K_L^0 K_S^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.448±0.011 OUR FIT		Error includes scale factor of 1.1.		
0.51 ±0.05 OUR AVERAGE				

0.56 ±0.07 3681 BUKIN 78C OLYA $e^+ e^- \rightarrow K_L^0 K_S^0, \pi^+ \pi^- \pi^0$

0.47 ±0.06 516 COSME 74 OSPK $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					

≈ 0.0087 1.98M 1,2 ALOISIO 03 KLOE $1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

<0.0006 90 3 ACHASOV 02 SND $1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

<0.23 90 3 CORDIER 80 DM1 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

<0.20 90 3 PARROUR 76B OSPK $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

1 From a fit without limitations on charged and neutral ρ masses and widths.

2 Adding the direct and $\omega\pi$ contributions and considering the interference between the $\rho\pi$ and $\pi^+ \pi^- \pi^0$.

3 Neglecting the interference between the $\rho\pi$ and $\pi^+ \pi^- \pi^0$.

 $\Gamma_2/(\Gamma_1 + \Gamma_2)$

NODE=M004R5
NODE=M004R5

NODE=M004R5;LINKAGE=01

NODE=M004R3
NODE=M004R3

NODE=M004R3;LINKAGE=AK

NODE=M004R3;LINKAGE=B2

NODE=M004R;LINKAGE=8D

NODE=M004R3;LINKAGE=E

NODE=M004R20
NODE=M004R20

NODE=M004R6
NODE=M004R6

NODE=M004R7
NODE=M004R7

NODE=M004R46
NODE=M004R46

NODE=M004R;LINKAGE=L1

NODE=M004R;LINKAGE=L2

NODE=M004R;LINKAGE=46

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.303 ± 0.025 OUR FIT Error includes scale factor of 1.2.

1.26 ± 0.04 OUR AVERAGE

1.246 ± 0.025 ± 0.057	10k	1 ACHASOV	98F SND	$e^+ e^- \rightarrow 7\gamma$
1.18 ± 0.11	279	2 AKHMETSHIN	95 CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
1.30 ± 0.06		3 DRUZHININ	84 ND	$e^+ e^- \rightarrow 3\gamma$
1.4 ± 0.2		4 DRUZHININ	84 ND	$e^+ e^- \rightarrow 6\gamma$
0.88 ± 0.20	290	KURDADZE	83C OLYA	$e^+ e^- \rightarrow 3\gamma$
1.35 ± 0.29		ANDREWS	77 CNTR	6.7–10 γ Cu
1.5 ± 0.4	54	3 COSME	76 OSPK	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.38 ± 0.02 ± 0.02		5 AKHMETSHIN	11 CMD2	$1.02 e^+ e^- \rightarrow \eta\gamma$
1.36 ± 0.05 ± 0.02	33k	6 ACHASOV	07B SND	$0.6\text{--}1.38 e^+ e^- \rightarrow \eta\gamma$
1.373 ± 0.014 ± 0.085	17.4k	7,8 AKHMETSHIN	05 CMD2	$0.60\text{--}1.38 e^+ e^- \rightarrow \eta\gamma$
1.287 ± 0.013 ± 0.063		9,10 AKHMETSHIN	01B CMD2	$e^+ e^- \rightarrow \eta\gamma$
1.338 ± 0.012 ± 0.052		11 ACHASOV	00 SND	$e^+ e^- \rightarrow \eta\gamma$
1.18 ± 0.03 ± 0.06	2200	12 AKHMETSHIN	99F CMD2	$e^+ e^- \rightarrow \eta\gamma$
1.21 ± 0.07		13 BENAYOUN	96 RVUE	$0.54\text{--}1.04 e^+ e^- \rightarrow \eta\gamma$

1 Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$.

2 From $\pi^+ \pi^- \pi^0$ decay mode of η .

3 From 2γ decay mode of η .

4 From $3\pi^0$ decay mode of η .

5 Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

6 ACHASOV 07B reports $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+ e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$ which we divide by our best value $B(\phi(1020) \rightarrow e^+ e^-) = (2.973 \pm 0.034) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

7 Using $B(\phi \rightarrow e^+ e^-) = (2.98 \pm 0.04) \times 10^{-4}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

8 Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

9 Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

10 The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

11 From the $\eta \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

12 From $\pi^+ \pi^- \pi^0$ decay mode of η and using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

13 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

Γ_6/Γ

NODE=M004R11

NODE=M004R11

OCCUR=2

NODE=M004R11;LINKAGE=AC

NODE=M004R11;LINKAGE=Z3

NODE=M004R11;LINKAGE=A

NODE=M004R11;LINKAGE=C

NODE=M004R11;LINKAGE=AN

NODE=M004R11;LINKAGE=AO

NODE=M004R11;LINKAGE=AH

NODE=M004R11;LINKAGE=AK

NODE=M004R;LINKAGE=AK

NODE=M004R;LINKAGE=BQ

NODE=M004R;LINKAGE=GA

NODE=M004R;LINKAGE=FF

NODE=M004R;LINKAGE=TS

Γ_7/Γ

NODE=M004R17

NODE=M004R17

OCCUR=3

NODE=M004R17;LINKAGE=D

NODE=M004R17;LINKAGE=AH

NODE=M004R17;LINKAGE=AK

NODE=M004R;LINKAGE=3G

NODE=M004R17;LINKAGE=TS

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.30 ± 0.05 OUR FIT

1.31 ± 0.13 OUR AVERAGE

1.30 ± 0.13		DRUZHININ	84 ND	$e^+ e^- \rightarrow 3\gamma$
1.4 ± 0.5	32	COSME	76 OSPK	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.367 ± 0.072		1 ACHASOV	16A SND	$0.60\text{--}1.38 e^+ e^- \rightarrow \pi^0\gamma$
1.258 ± 0.037 ± 0.077	18k	2,3 AKHMETSHIN	05 CMD2	$0.60\text{--}1.38 e^+ e^- \rightarrow \pi^0\gamma$
1.226 ± 0.036 ± 0.096		4 ACHASOV	00 SND	$e^+ e^- \rightarrow \pi^0\gamma$
1.26 ± 0.17		5 BENAYOUN	96 RVUE	$0.54\text{--}1.04 e^+ e^- \rightarrow \pi^0\gamma$

1 Using $B(\phi \rightarrow e^+ e^-)$ from PDG 15. Supersedes ACHASOV 00.

2 Using $B(\phi \rightarrow e^+ e^-) = (2.98 \pm 0.04) \times 10^{-4}$.

3 Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

4 From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

5 Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.9 ± 0.3 ± 0.7	ACHASOV	00 SND	$e^+ e^- \rightarrow \eta\gamma, \pi^0\gamma$
------------------	---------	--------	---

Γ_6/Γ_7

NODE=M004R42

NODE=M004R42

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
2.973 ± 0.034 OUR FIT				Error includes scale factor of 1.3.	
2.98 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.1.	
2.93 ± 0.14	1900k	1 ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-$, $K_S K_L, \pi^+\pi^-\pi^0$	
2.88 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \text{hadrons}$	
3.00 ± 0.21	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow \text{hadrons}$	
3.10 ± 0.14		2 PARROUR	76 OSPK	e^+e^-	
3.3 ± 0.3		COSME	74 OSPK	$e^+e^- \rightarrow \text{hadrons}$	
2.81 ± 0.25	681	BALAKIN	71 OSPK	$e^+e^- \rightarrow \text{hadrons}$	
3.50 ± 0.27		CHATELUS	71 OSPK	e^+e^-	

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

² Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
2.86 ± 0.19 OUR FIT				
2.5 ± 0.4 OUR AVERAGE				
2.69 ± 0.46	1 HAYES	71 CNTR	$8.3, 9.8 \gamma C \rightarrow \mu^+\mu^-X$	
2.17 ± 0.60	1 EARLES	70 CNTR	$6.0 \gamma C \rightarrow \mu^+\mu^-X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.87 $\pm 0.20 \pm 0.14$	2 ACHASOV	01G SND	$e^+e^- \rightarrow \mu^+\mu^-$	
3.30 $\pm 0.45 \pm 0.32$	3 ACHASOV	99C SND	$e^+e^- \rightarrow \mu^+\mu^-$	
4.83 ± 1.02	4 VASSERMAN	81 OLYA	$e^+e^- \rightarrow \mu^+\mu^-$	
2.87 ± 1.98	4 AUGUSTIN	73 OSPK	$e^+e^- \rightarrow \mu^+\mu^-$	

1 Neglecting interference between resonance and continuum.

2 Using $B(\phi \rightarrow e^+e^-) = (2.91 \pm 0.07) \times 10^{-4}$.

3 Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

4 Recalculated by us using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
1.08 ± 0.04 OUR AVERAGE					
1.075 $\pm 0.007 \pm 0.038$	30k	1 BABUSCI	15 KLOE	$1.02 e^+e^- \rightarrow \eta e^+e^-$	I
1.19 $\pm 0.19 \pm 0.12$	213	2 ACHASOV	01B SND	$e^+e^- \rightarrow \eta e^+e^-$	
1.14 $\pm 0.10 \pm 0.06$	355	3 AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.13 $\pm 0.14 \pm 0.07$	183	4 AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$	
1.21 $\pm 0.14 \pm 0.09$	130	5 AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$	
1.04 $\pm 0.20 \pm 0.08$	42	6 AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$	
1.3 ± 0.8	7	GOLUBEV	85 ND	$e^+e^- \rightarrow \eta e^+e^-$	

1 Using $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$ from PDG 12.

2 Using $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$, $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$, and $B(\phi \rightarrow e^+e^-) = (3.00 \pm 0.06) \times 10^{-4}$.

3 The average of the branching ratios separately obtained from the $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$ decays.

4 From $\eta \rightarrow \gamma\gamma$ decays and using $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

5 From $\eta \rightarrow 3\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

6 From $\eta \rightarrow \pi^+\pi^-\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+e^-\gamma) = (1.198 \pm 0.032) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.0 \pm 0.4) \times 10^{-2}$, $B(\phi \rightarrow \pi^+\pi^-\pi^0) = (15.5 \pm 0.6) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

NODE=M004R16

NODE=M004R16

NODE=M004R16;LINKAGE=AE

NODE=M004R16;LINKAGE=E

NODE=M004R10

NODE=M004R10

NODE=M004R10;LINKAGE=A

NODE=M004R;LINKAGE=GZ

NODE=M004R10;LINKAGE=8D

NODE=M004R;LINKAGE=VA

NODE=M004R24

NODE=M004R24

OCCUR=2

OCCUR=3

OCCUR=4

NODE=M004R24;LINKAGE=A

NODE=M004R;LINKAGE=VM

NODE=M004R;LINKAGE=H1

NODE=M004R;LINKAGE=H2

NODE=M004R;LINKAGE=H3

NODE=M004R;LINKAGE=H4

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{12}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.71 \pm 0.11 \pm 0.09		1 ACHASOV 00C	SND	$e^+e^- \rightarrow \pi^+\pi^-$	
0.65 \pm 0.38 - 0.29		1 GOLUBEV 86	ND	$e^+e^- \rightarrow \pi^+\pi^-$	
2.01 \pm 1.07 - 0.84		1 VASSERMAN 81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$	
<6.6	95	BUKIN 78B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$	
<2.7	95	ALVENSLEB... 72	CNTR	$6.7 \gamma C \rightarrow C\pi^+\pi^-$	

¹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ
4.7 \pm 0.5 OUR FIT				

**5.2 \pm 1.3
- 1.1**

1,2 AULCHENKO 00A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
-------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.4 \pm 0.6	3 AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
\sim 5.4	4 ACHASOV 00E	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
5.5 \pm 1.6 - 1.4 \pm 0.3	2,5 AULCHENKO 00A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
4.8 \pm 1.9 - 1.7 \pm 0.8	4 ACHASOV 99	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

¹ Using the 1996 and 1998 data.

² (2.3 ± 0.3)% correction for other decay modes of the $\omega(782)$ applied.

³ Not independent of the corresponding $\Gamma(\omega\pi^0) \times \Gamma(e^+e^-) / \Gamma^2(\text{total})$.

⁴ Using the 1996 data.

⁵ Using the 1998 data.

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{14}/Γ
<0.05	84	LINDSEY	66	HBC	2.1–2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals

 $\Gamma(\rho\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{15}/Γ
< 0.12	90	1 AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7	90	AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<200	84	LINDSEY	66	HBC

¹ Supersedes AKHMETSHIN 97C.

 $\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{16}/Γ
0.41 \pm 0.12 \pm 0.04	30175	1	AKHMETSHIN 99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.3	90	2 AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<600	90	KALBFLEISCH 75	HBC	2.18 $K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$
< 70	90	COSME	74	OSPK
<400	90	LINDSEY	65	HBC

$2.1-2.7 K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals

¹ For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible. Supersedes AKHMETSHIN 97C.

² For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible.

 $\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{17}/Γ
3.22 \pm 0.19 OUR FIT Error includes scale factor of 1.1.						
3.21 \pm 0.19 OUR AVERAGE						

3.21 \pm 0.03 - 0.09	18	AMBROSINO 07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
2.90 \pm 0.21 \pm 1.54		2 AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma, \pi^0\pi^0\gamma$

NODE=M004R18
NODE=M004R18

NODE=M004R18;LINKAGE=8D

NODE=M004R28
NODE=M004R28

OCCUR=2

NODE=M004R28;LINKAGE=K2
NODE=M004R28;LINKAGE=K3
NODE=M004R28;LINKAGE=AM
NODE=M004R28;LINKAGE=VS
NODE=M004R28;LINKAGE=K1

NODE=M004R14
NODE=M004R14

NODE=M004R15
NODE=M004R15

NODE=M004R;LINKAGE=1N

NODE=M004R12
NODE=M004R12

NODE=M004R;LINKAGE=2N

NODE=M004R12;LINKAGE=Z3

NODE=M004R30
NODE=M004R30

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.47 ± 0.21	2438	3 ALOISIO	02D KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
3.5 ± 0.3 $^{+1.3}_{-0.5}$	419	4,5 ACHASOV	00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
1.93 $\pm 0.46 \pm 0.50$	27188	6 AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
3.05 $\pm 0.25 \pm 0.72$	268	7 AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
1.5 ± 0.5	268	8 AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	OCCUR=2
3.42 $\pm 0.30 \pm 0.36$	164	4 ACHASOV	98I SND	$e^+ e^- \rightarrow 5\gamma$	
< 1	90	9 AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	OCCUR=2
< 7	90	10 AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
< 20	90	DRUZHININ	87 ND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	

¹ Obtained by the authors taking into account the $\pi^+ \pi^-$ decay mode. Includes a component due to $\pi\pi$ production via the $f_0(500)$ meson. Supersedes ALOISIO 02D.

² From the combined fit of the photon spectra in the reactions $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$.

³ From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution. Superseded by AMBROSINO 07.

⁴ Assuming that the $\pi^0 \pi^0 \gamma$ final state is completely determined by the $f_0 \gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+ \pi^-) = 2B(f_0 \rightarrow \pi^0 \pi^0)$.

⁵ Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

⁶ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

⁷ Neglecting other intermediate mechanisms ($\rho\pi$, $\sigma\gamma$).

⁸ A narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.

⁹ For destructive interference with the Bremsstrahlung process

¹⁰ For constructive interference with the Bremsstrahlung process

$\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.47 ± 0.15 OUR FIT				Error includes scale factor of 1.1.

2.6 ± 0.2 $^{+0.8}_{-0.3}$	419	1 ACHASOV	00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
---	-----	-----------	---------	--

¹ Assuming that the $\pi^0 \pi^0 \gamma$ final state is completely determined by the $f_0 \gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+ \pi^-) = 2B(f_0 \rightarrow \pi^0 \pi^0)$.

Γ_{17}/Γ_6

- NODE=M004R30;LINKAGE=MB
- NODE=M004R;LINKAGE=SL
- NODE=M004R;LINKAGE=KD
- NODE=M004R;LINKAGE=AI
- NODE=M004R;LINKAGE=U8
- NODE=M004R;LINKAGE=3N
- NODE=M004R;LINKAGE=SM
- NODE=M004R;LINKAGE=ST
- NODE=M004R30;LINKAGE=A
- NODE=M004R30;LINKAGE=B
- NODE=M004R44
- NODE=M004R44

$\Gamma(\pi^0 \pi^0 \gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.06 OUR AVERAGE					

1.07 ± 0.01 $^{+0.06}_{-0.03}$		1 AMBROSINO 07	KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
1.08 ± 0.17 ± 0.09	268	AKHMETSHIN 99C	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.09 ± 0.03 ± 0.05	2438	ALOISIO	02D KLOE	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
1.158 $\pm 0.093 \pm 0.052$	419	2,3 ACHASOV	00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$	
< 10	90	DRUZHININ	87 ND	$e^+ e^- \rightarrow 5\gamma$	

¹ Supersedes ALOISIO 02D.

² Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

³ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

Γ_{18}/Γ

- NODE=M004R44;LINKAGE=AI
- NODE=M004R26
- NODE=M004R26
- NODE=M004R26;LINKAGE=MB
- NODE=M004R26;LINKAGE=U8
- NODE=M004R26;LINKAGE=V8
- NODE=M004R39
- NODE=M004R39

$\Gamma(\pi^0 \pi^0 \gamma)/\Gamma(\eta\gamma)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.04 OUR FIT				

0.865 $\pm 0.070 \pm 0.017$	419	1 ACHASOV	00H SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
---	-----	-----------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90 ± 0.08 ± 0.07	164	ACHASOV	98I SND	$e^+ e^- \rightarrow 5\gamma$
----------------------------	-----	---------	---------	-------------------------------

¹ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

Γ_{18}/Γ_6

- NODE=M004R39;LINKAGE=V8
- NODE=M004R22
- NODE=M004R22
- NODE=M004R22;LINKAGE=A

$\Gamma(\pi^+ \pi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.5 ± 2.7 ± 1.6		6.8k	1 AKHMETSHIN 17	CMD3	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.93 $\pm 1.74 \pm 2.14$	3.3k	AKHMETSHIN 00E	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$
< 870	90	CORDIER	79 WIRE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$

¹ Using the cross section at the ϕ meson peak $\sigma(\phi) = 4172 \pm 42$ nb, the nonresonant cross section $\sigma(0) = 1.263 \pm 0.027$ nb and $\text{Re}(Z) = 0.146 \pm 0.030$, $\text{Im}(Z) = -0.002 \pm 0.024$ for the complex amplitude of the $\phi \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ transition.

$\Gamma(\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<150	95	BARKOV	88	CMD $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

 Γ_{20}/Γ

NODE=M004R27
NODE=M004R27

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.33 $^{+0.07}_{-0.10}$ OUR AVERAGE

1.35 ± 0.05	9.5k	1 ANASTASI	16B	KLOE	$e^+e^- \rightarrow \pi^0 e^+ e^-$
1.01 ± 0.28	52	2 ACHASOV	02D	SND	$e^+e^- \rightarrow \pi^0 e^+ e^-$
1.22 ± 0.34	46	3 AKHMETSHIN 01C	CMD2		$e^+e^- \rightarrow \pi^0 e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	DOLINSKY	88	ND	$e^+e^- \rightarrow \pi^0 e^+ e^-$
-----	----	----------	----	----	------------------------------------

1 Using $B(\pi^0 \rightarrow \gamma\gamma)$ from the 2014 Edition of this Review (PDG 14).

2 Using various branching ratios from the 2000 Edition of this Review (PDG 00).

3 Using $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$, $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$, and $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$.

 $\Gamma(\pi^0\eta\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

7.27 ± 0.30 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

7.06 ± 0.22	16.9k	1 AMBROSINO	09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
8.51 ± 0.51	607	2 ALOISIO	02C	KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
7.96 ± 0.60	197	3 ALOISIO	02C	KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
8.8 ± 1.4 ± 0.9	36	4 ACHASOV	00F	SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
9.0 ± 2.4 ± 1.0	80	AKHMETSHIN 99C	CMD2		$e^+e^- \rightarrow \eta\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.01 ± 0.10	13.3k	2,5 AMBROSINO	09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
7.12 ± 0.13	3.6k	3,6 AMBROSINO	09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
8.3 ± 2.3 ± 1.2	20	ACHASOV	98B	SND	$e^+e^- \rightarrow 5\gamma$
<250	90	DOLINSKY	91	ND	$e^+e^- \rightarrow \pi^0\eta\gamma$

1 Combined results of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decay modes measurements.

2 From the decay mode $\eta \rightarrow \gamma\gamma$.

3 From the decay mode $\eta \rightarrow \pi^+\pi^-\pi^0$.

4 Supersedes ACHASOV 98B.


5 Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$.

6 Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.73 \pm 0.28)\%$.

 Γ_{21}/Γ

NODE=M004R31
NODE=M004R31

WEIGHTED AVERAGE
7.27 ± 0.30 (Error scaled by 1.5)



NODE=M004R31;LINKAGE=A

NODE=M004R;LINKAGE=DS

NODE=M004R;LINKAGE=5H

NODE=M004R32
NODE=M004R32

OCCUR=2

OCCUR=2

OCCUR=3

NODE=M004R32;LINKAGE=AM

NODE=M004R;LINKAGE=C1

NODE=M004R;LINKAGE=C2

NODE=M004R32;LINKAGE=AF

NODE=M004R32;LINKAGE=AB

NODE=M004R32;LINKAGE=AR

$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$				Γ_{23}/Γ		
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
7.6±0.6 OUR FIT						
7.6±0.6 OUR AVERAGE						
7.4±0.7			1 ALOISIO	02C KLOE	$e^+ e^- \rightarrow \eta \pi^0 \gamma$	
8.8±1.7		36	2 ACHASOV	00F SND	$e^+ e^- \rightarrow \eta \pi^0 \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
11 ± 2			3 GOKALP	02 RVUE	$e^+ e^- \rightarrow \eta \pi^0 \gamma$	
<500	90		DOLINSKY	91 ND	$e^+ e^- \rightarrow \pi^0 \eta \gamma$	

¹ Using $M_{a_0(980)} = 984.8$ MeV and assuming $a_0(980)\gamma$ dominance.

² Assuming $a_0(980)\gamma$ dominance in the $\eta \pi^0 \gamma$ final state.

³ Using data of ACHASOV 00F.

$\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$				Γ_{17}/Γ_{23}		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
6.1±0.6		1 ALOISIO	02C KLOE	$e^+ e^- \rightarrow \eta \pi^0 \gamma$		
1 Using results of ALOISIO 02D and assuming that $f_0(980)$ decays into $\pi\pi$ only and $a_0(980)$ into $\eta\pi$ only.						
$\Gamma(K^0 \bar{K}^0 \gamma)/\Gamma_{\text{total}}$				Γ_{24}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.9 × 10⁻⁸	90	AMBROSINO	09C KLOE	$e^+ e^- \rightarrow K_S^0 \bar{K}_S^0 \gamma$		

$\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$				Γ_{25}/Γ		
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
6.22±0.21 OUR FIT						
6.22±0.30 OUR AVERAGE						
6.22±0.27±0.12	3407	1 AMBROSINO	07A KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- 7\gamma$		
6.7 ± 2.8 ± 0.8	12	2 AULCHENKO	03B SND	$e^+ e^- \rightarrow \eta' \gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
6.7 ± 5.0 ± 1.5	7	AULCHENKO	03B SND	$e^+ e^- \rightarrow 7\gamma$		
6.10±0.61±0.43	120	3 ALOISIO	02E KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$		
8.2 ± 2.1 ± 1.1	21	4 AKHMETSHIN	00B CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$		
4.9 ± 2.2 ± 0.6	9	5 AKHMETSHIN	00F CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \geq 2\gamma$		OCCUR=2
6.4 ± 1.6	30	6 AKHMETSHIN	00F CMD2	$e^+ e^- \rightarrow \eta'(958)\gamma$		OCCUR=2
6.7 ± 3.4 ± 1.0	5	7 AULCHENKO	99 SND	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$		
<11	90	AULCHENKO	98 SND	$e^+ e^- \rightarrow 7\gamma$		
12 ± 7 ± 2	6	4 AKHMETSHIN	97B CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$		
<41	90	DRUZHININ	87 ND	$e^+ e^- \rightarrow \gamma \eta \pi^+ \pi^-$		

¹ AMBROSINO 07A reports $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma)/\Gamma_{\text{total}}] / [B(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$ which we multiply by our best value $B(\phi(1020) \rightarrow \eta\gamma) = (1.303 \pm 0.025) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Averaging AULCHENKO 03B with AULCHENKO 99.

³ Using $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$.

⁴ Using the value $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$.

⁵ Using $B(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$.

⁶ Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.

⁷ Using the value $B(\eta' \rightarrow \eta\pi^+\pi^-) = (43.7 \pm 1.5) \times 10^{-2}$ and $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$.

$\Gamma(\eta'(958)\gamma)/\Gamma(K_L^0 K_S^0)$				Γ_{25}/Γ_2		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.83±0.06 OUR FIT						
1.46±0.64±0.18	9	1 AKHMETSHIN	00F CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \geq 2\gamma$		

¹ Using various branching ratios of K_S^0 , K_L^0 , η , η' from the 2000 edition (The European Physical Journal C15 1 (2000)) of this Review.

NODE=M004R33
NODE=M004R33

NODE=M004R;LINKAGE=C3
NODE=M004R33;LINKAGE=AF
NODE=M004R;LINKAGE=GK

NODE=M004R47
NODE=M004R47

NODE=M004R;LINKAGE=C4

NODE=M004R48
NODE=M004R48

NODE=M004R25
NODE=M004R25

OCCUR=2

NODE=M004R25;LINKAGE=AM

NODE=M004R25;LINKAGE=BK
NODE=M004R;LINKAGE=E2
NODE=M004R25;LINKAGE=Q
NODE=M004R;LINKAGE=T2
NODE=M004R;LINKAGE=T3
NODE=M004R25;LINKAGE=AU

NODE=M004R43
NODE=M004R43

NODE=M004R;LINKAGE=T1

$\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$				Γ_{25}/Γ_6
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.77±0.15 OUR FIT				
4.78±0.20 OUR AVERAGE				
4.77±0.09±0.19	3407	AMBROSINO 07A	KLOE 1.02	$e^+ e^- \rightarrow \pi^+ \pi^- 7\gamma$
4.70±0.47±0.31	120	ALOISIO 02E	KLOE 1.02	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
6.5 $^{+1.7}_{-1.5}$ ±0.8	21	AKHMETSHIN 00B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.5 $^{+5.2}_{-4.0}$ ±1.4	6	AKHMETSHIN 97B	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- 3\gamma$

¹ From the decay mode $\eta' \rightarrow \eta \pi^+ \pi^-$, $\eta \rightarrow \gamma\gamma$.

² Superseded by AKHMETSHIN 00B.

$\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$				Γ_{26}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	AULCHENKO 98	SND	$e^+ e^- \rightarrow 7\gamma$

$\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$				Γ_{27}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.43±0.45±0.14	27188	¹ AKHMETSHIN 99B	CMD2	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.3 ±1.0	824 ± 33	² AKHMETSHIN 97C	CMD2	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
1 For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.				
2 For $E_\gamma > 20$ MeV.				

$\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$				Γ_{28}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.2	90	AULCHENKO 08	CMD2	$\phi \rightarrow \pi^+ \pi^- \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5	90	AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma\gamma$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{29}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.8	90	AKHMETSHIN 00E	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.1	90	AULCHENKO 08	CMD2	$\phi \rightarrow \eta \pi^+ \pi^-$
<30	90	AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma\gamma$

$\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$				Γ_{30}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<9.4	90	AKHMETSHIN 01	CMD2	$e^+ e^- \rightarrow \eta e^+ e^-$

$\Gamma(\eta U \rightarrow \eta e^+ e^-)/\Gamma_{\text{total}}$				Γ_{31}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1 × 10 ⁻⁶	90	¹ BABUSCI 13B	KLOE	$1.02 e^+ e^- \rightarrow \eta e^+ e^-$
1 For a narrow vector U with mass between 5 and 470 MeV, from the combined analysis of $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow \pi^0 \pi^0 \pi^0$ from ARCHILLI 12. Measured 90% CL limits as a function of m_U range from 2.2×10^{-8} to 10^{-6} .				

$\Gamma(\text{invisible})/\Gamma(K^+K^-)$				Γ_{32}/Γ_1
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.4 × 10 ⁻⁴	90	ABLIKIM 18s	BES3	$J/\psi \rightarrow \phi \eta \rightarrow \phi \pi^+ \pi^- \pi^0$

Lepton Family number (LF) violating modes

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$				Γ_{33}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2 × 10 ⁻⁶	90	ACHASOV 10A	SND	$e^+ e^- \rightarrow e^\pm \mu^\mp$

$$\pi^+ \pi^- \pi^0 / \rho\pi \text{ AMPLITUDE RATIO } a_1 \text{ IN DECAY OF } \phi \rightarrow \pi^+ \pi^- \pi^0$$

NODE=M004R34
NODE=M004R34

NODE=M004R;LINKAGE=E1
NODE=M004R;LINKAGE=KS

NODE=M004R36
NODE=M004R36

NODE=M004R35;LINKAGE=3N
NODE=M004R35;LINKAGE=A

NODE=M004R37
NODE=M004R37

NODE=M004R38
NODE=M004R38

NODE=M004R45
NODE=M004R45

NODE=M004R01
NODE=M004R01

NODE=M00422A

NODE=M004R29
NODE=M004R29

NODE=M004D1

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.2 OUR AVERAGE					
10.1±4.4±1.7	80k	1 AKHMETSHIN 06	CMD2	$1.017\text{--}1.021 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
9.0±1.1±0.6	1.98M	2,3 ALOISIO 03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$-6 < a_1 < 6$	500k	3 ACHASOV 02	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$	
$-16 < a_1 < 11$	90	9.8k 1,4 AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma\gamma$	
1 Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.					
2 From a fit without limitations on charged and neutral ρ masses and widths.					
3 Recalculated by us to match the notations of AKHMETSHIN 98.					
4 Assuming zero phase for the contact term.					

NODE=M004D1

NODE=M004D1

PARAMETER β IN $\phi \rightarrow Pe^+e^-$ DECAYS

In the one-pole approximation the electromagnetic transition form factor for $\phi \rightarrow Pe^+e^-$ ($P = \pi, \eta$) is given as a function of the e^+e^- invariant mass squared, q^2 , by the expression:

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter $\Lambda \approx 0.770$ GeV ($\Lambda^{-2} \approx 1.687$ GeV $^{-2}$). The slope of this form factor, $\beta = dF/dq^2(q^2=0)$, equals Λ^{-2} in this approximation.

The measurements below obtain β in the one-pole approximation.

PARAMETER β IN $\phi \rightarrow \pi^0 e^+e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
2.02±0.11	9.5k	1 ANASTASI 16B	KLOE	$1.02 e^+ e^- \rightarrow \pi^0 e^+ e^-$

1 The error combines statistical and systematic uncertainties.

NODE=M004D1;LINKAGE=AK

NODE=M004D;LINKAGE=L1

NODE=M004D;LINKAGE=L3

NODE=M004D1;LINKAGE=KL

NODE=M004230

NODE=M004230

PARAMETER β IN $\phi \rightarrow \eta e^+e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29±0.13 OUR AVERAGE				

1.28±0.10 $^{+0.09}_{-0.08}$	30k	BABUSCI 15	KLOE	$1.02 e^+ e^- \rightarrow \eta e^+ e^-$
3.8 ± 1.8	213	1 ACHASOV 01B	SND	$1.02 e^+ e^- \rightarrow \eta e^+ e^-$

1 The uncertainty is statistical only. The systematic one is negligible, in comparison.

NODE=M004A00

NODE=M004A00

NODE=M004A00;LINKAGE=A

NODE=M004BFP

NODE=M004BFP

NODE=M004BFP;LINKAGE=A

$\phi(1020)$ REFERENCES

AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	
KOZYREV	18	PL B779 64	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
AKHMETSHIN	17	PL B768 345	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)
ANASTASI	16B	PL B757 362	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)
KOZYREV	16	PL B760 314	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)
BABUSCI	15	PL B742 1	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
BABUSCI	13B	PL B720 111	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono (PARIN, BERLIN+)	
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>	
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)
AKHMETSHIN	11	PL B695 412	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>	
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)

Translated from ZETFP 88 93.

NODE=M004

REFID=59670

REFID=58971

REFID=59987

REFID=58794

REFID=57893

REFID=57513

REFID=57399

REFID=57514

REFID=56374

REFID=56977;ERROR=7

REFID=55940

REFID=55687

REFID=55068

REFID=55357

REFID=55127

REFID=55404

REFID=53951

REFID=54281

REFID=54305

REFID=54066

REFID=53645

REFID=53352

REFID=53212

REFID=52969

REFID=53105

REFID=52572

REFID=52573

REFID=52268

FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro	(SND Collab.)	REFID=52584
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(KLOE Collab.)	REFID=51942
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51616
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51646
DUBYNISKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>		REFID=51719
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=51465
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=50453
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50824
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49613
		Translated from ZETFP 124 28.			
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48814
		Translated from ZETFP 75 539.			
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48823
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48825
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl		REFID=48575
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>		REFID=49167
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48111
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Zubin	(Novosibirsk SND Collab.)	REFID=48312
ACHASOV	01G	PRL 86 1698	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48315
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48110
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48323
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell		REFID=48570
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47425
		Translated from ZETFP 117 22.			
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47431
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
		Translated from ZETFP 72 411.			
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47927
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47422
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47936
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47937
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
		Translated from ZETFP 117 1067.			
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>		REFID=47969
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>		REFID=46896
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>		REFID=46939
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47397
AULCHENKO	99A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
		Translated from ZETFP 117 1067.			
ACHASOV	99B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46317
ACHASOV	99F	JETPL 68 573	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46321
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>		REFID=46600
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=46325
AULCHENKO	98	PL B436 199	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46336
BARBERIS	98	PL B432 436	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46344
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin <i>et al.</i>	(NOVO, BOST, PIT+) (Novosibirsk CMD-2 Collab.)	REFID=45801
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=45802
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=44617
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
BARKOV	88	SJNP 47 248	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=41024
		Translated from YAF 47 393.			
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41022
		Translated from YAF 48 442.			
DRUZHININ	87	ZPHY C37 1	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=40448
ARMSTRONG	86	PL 166B 245	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=20563
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=11540
DAVENPORT	86	PR D33 2519	T.F. Davenport	(TUFTS, ARIZ, FNAL, FSU, NDAM+)	REFID=20567
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra <i>et al.</i>	(ANIK, BRIS, CERN+)	REFID=20568
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)	REFID=20569
GOLUBEV	86	SJNP 44 409	V.B. Golubev <i>et al.</i>	(NOVO)	REFID=40449
		Translated from YAF 44 633.			
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=20562
GOLUBEV	85	SJNP 41 756	V.B. Golubev <i>et al.</i>	(NOVO)	REFID=40450
		Translated from YAF 41 1183.			
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)	REFID=12177
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20560
		Translated from ZETFP 38 306.			
ARENTON	82	PR D25 2241	M.W. Arenton <i>et al.</i>	(ANL, ILL)	REFID=20556
PELLINEN	82	PS 25 599	A. Pellinen, M. Roos	(HELS)	REFID=20557
DAUM	81	PL 100B 439	C. Daum <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)	REFID=20552
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
Also		Private Comm.	S.I. Eidelman	(NOVO)	REFID=20554
VASSERMAN	81	PL 99B 62	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=20555
Also		SJNP 35 240	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=47475
		Translated from YAF 35 352.			

CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
CORDIER	79	PL 81B 389	A. Cordier <i>et al.</i>	(LALO)	REFID=20549
BUKIN	78B	SJNP 27 521	A.D. Bokin <i>et al.</i>	(NOVO)	REFID=20545
		Translated from YAF 27 985.			
BUKIN	78C	SJNP 27 516	A.D. Bokin <i>et al.</i>	(NOVO)	REFID=20544
		Translated from YAF 27 976.			
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
LOSTY	78	NP B133 38	M.J. Losty <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=20547
AKERLOF	77	PRL 39 861	C.W. Akerlof <i>et al.</i>	(FNAL, MICH, PURD)	REFID=20534
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
BALDI	77	PL 68B 381	R. Baldi <i>et al.</i>	(GEVA)	REFID=20536
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+)	REFID=20537
COHEN	77	PRL 38 269	D. Cohen <i>et al.</i>	(ANL)	REFID=20538
LAVEN	77	NP B127 43	H. Laven <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)	REFID=20541
LYONS	77	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)	REFID=20232
COSME	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20529
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20531
PARROUR	76	PL 63B 357	G. Parrou <i>et al.</i>	(ORSAY)	REFID=20532
PARROUR	76B	PL 63B 362	G. Parrou <i>et al.</i>	(ORSAY)	REFID=20533
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AYRES	74	PRL 32 1463	D.S. Ayres <i>et al.</i>	(ANL)	REFID=20522
BESCH	74	NP B70 257	H.J. Besch <i>et al.</i>	(BONN)	REFID=20523
COSME	74	PL 48B 155	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20525
COSME	74B	PL 48B 159	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20526
DEGROOT	74	NP B74 77	A.J. de Groot <i>et al.</i>	(AMST, NIJM)	REFID=20527
AUGUSTIN	73	PRL 30 462	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=47515
BALLAM	73	PR D7 3150	J. Ballam <i>et al.</i>	(SLAC, LBL)	REFID=20520
BINNIE	73B	PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
AGUILAR...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
ALVENSELB...	72	PRL 28 66	H. Alvensleben <i>et al.</i>	(MIT, DESY)	REFID=20514
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
COLLEY	72	NP B50 1	D.C. Colley <i>et al.</i>	(BIRM, GLAS)	REFID=20519
BALAKIN	71	PL 34B 328	V.E. Balakin <i>et al.</i>	(NOVO)	REFID=20507
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)	REFID=20508
Also		PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)	REFID=20501
HAYES	71	PR D4 899	S. Hayes <i>et al.</i>	(CORN)	REFID=20511
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemyer	(UMD)	REFID=20512
BIZOT	70	PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)	REFID=20501
Also		Liverpool Sym. 69	J.P. Perez-y-Jorba		REFID=20502
EARLES	70	PRL 25 1312	D.R. Earles <i>et al.</i>	(NEAS)	REFID=20504
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)	REFID=20481
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IGJPC	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=20253
LINDSEY	65	PRL 15 221	J.S. Lindsey, G.A. Smith	(LRL)	REFID=20478
LINDSEY	65 data included in LINDSEY 66.				
SCHLEIN	63	PRL 10 368	P.E. Schlein <i>et al.</i>	(UCLA) IGJP	REFID=20474

 $h_1(1170)$ $I^G(J^{PC}) = 0^-(1^{+-})$ **$h_1(1170)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

 1170 ± 20 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

1168 ± 4	ANDO	92	SPEC	$8\pi^- p \rightarrow \pi^+\pi^-\pi^0 n$
1166 ± 5 ± 3	1 ANDO	92	SPEC	$8\pi^- p \rightarrow \pi^+\pi^-\pi^0 n$
1190 ± 60	2 DANKOWY...	81	SPEC 0	$8\pi p \rightarrow 3\pi n$

1 Average and spread of values using 2 variants of the model of BOWLER 75.

2 Uses the model of BOWLER 75.

NODE=M030M

NODE=M030M

→ UNCHECKED ←

OCCUR=2

NODE=M030M;LINKAGE=B

NODE=M030M;LINKAGE=C

NODE=M030W

NODE=M030W

→ UNCHECKED ←

OCCUR=2

NODE=M030W;LINKAGE=B

NODE=M030W;LINKAGE=C

 $h_1(1170)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

 360 ± 40 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

345 ± 6	ANDO	92	SPEC	$8\pi^- p \rightarrow \pi^+\pi^-\pi^0 n$
375 ± 6 ± 34	3 ANDO	92	SPEC	$8\pi^- p \rightarrow \pi^+\pi^-\pi^0 n$
320 ± 50	4 DANKOWY...	81	SPEC 0	$8\pi p \rightarrow 3\pi n$

3 Average and spread of values using 2 variants of the model of BOWLER 75.

4 Uses the model of BOWLER 75.

h₁(1170) DECAY MODES

NODE=M030215;NODE=M030

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi$	seen

h₁(1170) BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ/Γ
<u>VALUE</u>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	ANDO 92	SPEC	$8\pi^- p \rightarrow \pi^+\pi^-\pi^0 n$	
seen	ATKINSON 84	OMEG	$20\text{--}70\gamma p \rightarrow \pi^+\pi^-\pi^0 p$	
seen	DANKOWY... 81	SPEC	$8\pi p \rightarrow 3\pi n$	

h₁(1170) REFERENCES

ANDO 92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
ATKINSON 84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
DANKOWY... 81	PRL 46 580	J.A. Dankowich <i>et al.</i>	(TNTD, BNL, CARL+)
BOWLER 75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)


b₁(1235)

$I^G(J^{PC}) = 1^+(1^{+-})$

b₁(1235) MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
1229.5 ± 3.2 OUR AVERAGE					
		Error includes scale factor of 1.6. See the ideogram below.			
1225 ± 5		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
1235 ± 15		ALDE	92C GAM2		$38,100\pi^- p \rightarrow \omega\pi^0 n$
1236 ± 16		FUKUI	91 SPEC		$8.95\pi^- p \rightarrow \omega\pi^0 n$
1222 ± 6		ATKINSON	84E OMEG ±		$25\text{--}55\gamma p \rightarrow \omega\pi X$
1237 ± 7		ATKINSON	84E OMEG 0		$25\text{--}55\gamma p \rightarrow \omega\pi X$
1239 ± 5		EVANGELIS...	81 OMEG -		$12\pi^- p \rightarrow \omega\pi\pi$
1251 ± 8	450	GESELLER	77 HBC -		$11\pi^- p \rightarrow \pi^-\omega p$
1245 ± 11	890	FLATTE	76C HBC -		$4.2 K^- p \rightarrow \pi^-\omega\Sigma^+$
1222 ± 4	1400	CHALOUPKA	74 HBC -		$3.9\pi^- p$
1220 ± 7	600	KARSHON	74B HBC +		$4.9\pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1190 ± 10		AUGUSTIN	89 DM2 ±		$e^+ e^- \rightarrow 5\pi$
1213 ± 5		ATKINSON	84C OMEG 0		$20\text{--}70\gamma p$
1271 ± 11		COLLICK	84 SPEC +		$200\pi^+ Z \rightarrow Z\pi\omega$

WEIGHTED AVERAGE
1229.5±3.2 (Error scaled by 1.6)



DESIG=1;OUR EST;→ UNCHECKED ←

NODE=M030220

NODE=M030R1
NODE=M030R1

NODE=M030

REFID=43171
REFID=20574
REFID=20572
REFID=20571

NODE=M011

NODE=M011M

NODE=M011M

OCCUR=2

***b₁(1235)* WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
142± 9 OUR AVERAGE		Error includes scale factor of 1.2.			
113±12		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
160±30		ALDE	92C GAM2		$38,100 \pi^- p \rightarrow \omega \pi^0 n$
151±31		FUKUI	91 SPEC		$8.95 \pi^- p \rightarrow \omega \pi^0 n$
170±15		EVANGELIS...	81 OMEG	-	$12 \pi^- p \rightarrow \omega \pi p$
170±50	225	BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p 4\pi$
155±32	450	GESELLER	77 HBC	-	$11 \pi^- p \rightarrow \pi^- \omega p$
182±45	890	FLATTE	76C HBC	-	$4.2 K^- p \rightarrow \pi^- \omega \Sigma^+$
135±20	1400	CHALOUPKA	74 HBC	-	$3.9 \pi^- p$
156±22	600	KARSHON	74B HBC	+	$4.9 \pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
210±19		AUGUSTIN	89 DM2	±	$e^+ e^- \rightarrow 5\pi$
231±14		ATKINSON	84C OMEG	0	$20-70 \gamma p$
232±29		COLLICK	84 SPEC	+	$200 \pi^+ Z \rightarrow Z \pi \omega$

NODE=M011W

NODE=M011W

***b₁(1235)* DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \omega \pi$ [D/S amplitude ratio = 0.277 ± 0.027]	seen	
$\Gamma_2 \pi^\pm \gamma$	$(1.6 \pm 0.4) \times 10^{-3}$	
$\Gamma_3 \eta \rho$	seen	
$\Gamma_4 \pi^+ \pi^+ \pi^- \pi^0$	< 50 %	84%
$\Gamma_5 K^*(892)^\pm K^\mp$	seen	
$\Gamma_6 (K\bar{K})^\pm \pi^0$	< 8 %	90%
$\Gamma_7 K_S^0 K_L^0 \pi^\pm$	< 6 %	90%
$\Gamma_8 K_S^0 K_S^0 \pi^\pm$	< 2 %	90%
$\Gamma_9 \phi \pi$	< 1.5 %	84%

NODE=M011215;NODE=M011

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=74

DESIG=71;OUR EST;→ UNCHECKED ←

DESIG=73;OUR EST;→ UNCHECKED ←

DESIG=72;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

***b₁(1235)* PARTIAL WIDTHS**

$\Gamma(\pi^\pm \gamma)$	Γ_2
VALUE (keV)	
230±60	COLLICK 84 SPEC + $200 \pi^+ Z \rightarrow Z \pi \omega$


NODE=M011220

NODE=M011W3
NODE=M011W3***b₁(1235)* D-wave/S-wave AMPLITUDE RATIO
IN DECAY OF $b_1(1235) \rightarrow \omega \pi$**

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.277±0.027 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.			
0.269±0.009±0.010		NOZAR	02 MPS	-	$18 \pi^- p \rightarrow \omega \pi^- p$
0.23 ±0.03		AMSLER	94C CBAR		$0.0 \bar{p}p \rightarrow \omega \eta \pi^0$
0.45 ±0.04		AMSLER	93B CBAR		$0.0 \bar{p}p \rightarrow \omega \pi^0 \pi^0$
0.235±0.047		ATKINSON	84C OMEG		$20-70 \gamma p$
0.4 +0.1 -0.1		GESELLER	77 HBC	-	$11 \pi^- p \rightarrow \pi^- \omega p$
0.21 ±0.08		CHUNG	75B HBC	+	$7.1 \pi^+ p$
0.3 ±0.1		CHALOUPKA	74 HBC	-	$3.9-7.5 \pi^- p$
0.35 ±0.25	600	KARSHON	74B HBC	+	$4.9 \pi^+ p$

NODE=M011DS

NODE=M011DS



$b_1(1235)$ D-wave/S-wave amplitude ratio in decay of $b_1(1235) \rightarrow \omega\pi$

**$b_1(1235)$ D-wave/S-wave AMPLITUDE PHASE DIFFERENCE
IN DECAY OF $b_1(1235) \rightarrow \omega\pi$**

NODE=M011PH

VALUE (°)	DOCUMENT ID	TECN	CHG	COMMENT
10.5±2.4±3.9	NOZAR	02	MPS	- 18 $\pi^- p \rightarrow \omega\pi^- p$

NODE=M011PH

$b_1(1235)$ BRANCHING RATIOS

NODE=M011230

$\Gamma(\eta\rho)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_3/Γ_1	
<0.10	ATKINSON	84D	OMEG	20–70 γp

NODE=M011R9

NODE=M011R9

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_4/Γ_1	
<0.5	ABOLINS	63	HBC	+ 3.5 $\pi^+ p$

NODE=M011R1

NODE=M011R1

$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	Γ_5/Γ_1	
seen	¹ ABLIKIM	10E	BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M011R10

NODE=M011R10

¹ From a fit including ten additional resonances and energy-independent Breit-Wigner width.

NODE=M011R10;LINKAGE=AB

$\Gamma((K\bar{K})^\pm\pi^0)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_6/Γ_1	
<0.08	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R6

NODE=M011R6

$\Gamma(K_S^0 K_L^0 \pi^\pm)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_7/Γ_1	
<0.06	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R8

NODE=M011R8

$\Gamma(K_S^0 K_S^0 \pi^\pm)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_8/Γ_1	
<0.02	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R7

NODE=M011R7

$\Gamma(\phi\pi)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_9/Γ_1	
<0.004	VIKTOROV	96	SPEC	0 32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$

NODE=M011R4

NODE=M011R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	95	BIZZARRI	69	HBC	± 0.0 $\bar{p}p$
<0.015		DAHL	67	HBC	1.6–4.2 $\pi^- p$

b₁(1235) REFERENCES

ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
NOZAR	02	PL B541 35	M. Nozar <i>et al.</i>		REFID=48850
VIKTOROV	96	PAN 59 1184	V.A. Viktorov <i>et al.</i>	(SERP)	REFID=45203
		Translated from YAF 59 1239.			
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43602
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41859
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41581
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
ATKINSON	84C	NP B243 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+) JP	REFID=20625
ATKINSON	84D	NP B242 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20623
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20624
COLLICK	84	PRL 53 2374	B. Collick <i>et al.</i>	(MINN, ROCH, FNAL)	REFID=20626
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
GESAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+) JP	REFID=20230
FLATTE	76C	PL 64B 225	S.M. Flatte <i>et al.</i>	(CERN, AMST, NIJM+) JP	REFID=20615
CHUNG	75B	PR D11 2426	S.U. Chung <i>et al.</i>	(BNL, LBL, UCSC) JP	REFID=20613
CHALOUPKA	74	PL 51B 407	V. Chaloupka <i>et al.</i>	(CERN) JP	REFID=20611
KARSHON	74B	PR D10 3608	U. Karshon <i>et al.</i>	(REHO) JP	REFID=20612
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20171
BALTAY	67	PRL 18 93	C. Baltay <i>et al.</i>	(COLU)	REFID=20159
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL)	REFID=20321
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)	REFID=20006

a₁(1260)

$$I^G(J^{PC}) = 1^-(1^{++})$$

See also our review under the $a_1(1260)$ in PDG 06, Journal of Physics **G33** 1 (2006).

a₁(1260) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1230 ±40 OUR ESTIMATE					
1299 +12 -28	46M	¹ AGHASYAN	18B COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1195.05 ± 1.05 ± 6.33	894k	AAIJ	18AI LHCb	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$	
1209 ± 4 +12 -9		² MIKHASENKO	18 RVUE	$\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$	
1225 ± 9 ± 20	7k	³ DARGENT	17 RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$	
1255 ± 6 +7 -17	420k	⁴ ALEKSEEV	10 COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
1243 ± 12 ± 20		⁵ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \rho^0 \rho^\pm \pi^\mp \gamma$	
1230–1270	6360	⁶ LINK	07A FOCS	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$	
1203 ± 3		⁷ GOMEZ-DUM..04	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$	
1330 ± 24	90k	SALVINI	04 OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$	
1331 ± 10 ± 3	37k	⁸ ASNER	00 CLE2	$10.6 e^+ e^- \rightarrow \tau^+ \tau^-, \tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$	
1255 ± 7 ± 6	5904	⁹ ABREU	98G DLPH	$e^+ e^-$	
1207 ± 5 ± 8	5904	¹⁰ ABREU	98G DLPH	$e^+ e^-$	OCCUR=2
1196 ± 4 ± 5	5904	^{11,12} ABREU	98G DLPH	$e^+ e^-$	OCCUR=3
1240 ± 10		BARBERIS	98B	$450 pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$	
1262 ± 9 ± 7		^{9,13} ACKERSTAFF	97R OPAL	$E_{cm}^{ee} = 88\text{--}94, \tau^- \rightarrow 3\pi\nu$	
1210 ± 7 ± 2		^{10,13} ACKERSTAFF	97R OPAL	$E_{cm}^{ee} = 88\text{--}94, \tau^- \rightarrow 3\pi\nu$	OCCUR=2
1211 ± 7 +50 -0		¹⁰ ALBRECHT	93C ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1121 ± 8		¹⁴ ANDO	92 SPEC	$8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
1242 ± 37		¹⁵ IVANOV	91 RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1260 ± 14		¹⁶ IVANOV	91 RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=2
1250 ± 9		¹⁷ IVANOV	91 RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
1208 ± 15		ARMSTRONG	90 OMEG	$300.0 pp \rightarrow pp \pi^+ \pi^- \pi^0$	
1220 ± 15		¹⁸ ISGUR	89 RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1260 ± 25		¹⁹ BOWLER	88 RVUE		
1166 ± 18 ± 11		BAND	87 MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	

NODE=M011

REFID=53361
REFID=48850
REFID=45203

REFID=44091
REFID=43602
REFID=43585

REFID=41859
REFID=41581
REFID=41004

REFID=20625
REFID=20623
REFID=20624

REFID=20626
REFID=20462
REFID=21265

REFID=20230
REFID=20615
REFID=20613

REFID=20611
REFID=20612
REFID=20171

REFID=20159
REFID=20321
REFID=20006

NODE=M010

NODE=M010M

NODE=M010M
→ UNCHECKED ←

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

1164	± 41	± 23	BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^- \nu$
1250	± 40		18 TORNQVIST	87	RVUE	
1046	± 11		ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1056	± 20	± 15	RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1194	± 14	± 10	SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1255	± 23		BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
1240	± 80		20 DANKOWY...	81	SPEC	$8.45 \pi^- p \rightarrow n3\pi$
1280	± 30		20 DAUM	81B	CNTR	$63.94 \pi^- p \rightarrow p3\pi$
1041	± 13		21 GAVILLET	77	HBC	$4.2 K^- p \rightarrow \Sigma 3\pi$

OCCUR=2

1 Statistical error negligible.

2 From the pole position. Using an amplitude analysis based on approximate three-body unitary of τ data from SCHAEFEL 05C.

3 Reanalysis of CLEO data using Breit-Wigner parameterization.

4 Superseded by AGHASYAN 2018B.

5 The $\rho^\pm \pi^\mp$ state can be also due to the $\pi(1300)$.

6 Using the Breit-Wigner parameterization; strong correlation between mass and width.

7 Using the data of BARATE 98R.

8 From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.

9 Uses the model of KUHN 90.

10 Uses the model of ISGUR 89.

11 Includes the effect of a possible a'_1 state.

12 Uses the model of FEINDT 90.

13 Supersedes AKERS 95P.

14 Average and spread of values using 2 variants of the model of BOWLER 75.

15 Reanalysis of RUCKSTUHL 86.

16 Reanalysis of SCHMIDKE 86.

17 Reanalysis of ALBRECHT 86B.

18 From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

19 From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

20 Uses the model of BOWLER 75.

21 Produced in K^- backward scattering.

NODE=M010M;LINKAGE=Q

NODE=M010M;LINKAGE=S

NODE=M010M;LINKAGE=V

NODE=M010M;LINKAGE=R

NODE=M010M;LINKAGE=AU

NODE=M010M;LINKAGE=LI

NODE=M010M;LINKAGE=GO

NODE=M010M;LINKAGE=B6

NODE=M010M;LINKAGE=KS

NODE=M010M;LINKAGE=IM

NODE=M010M;LINKAGE=A1

NODE=M010M;LINKAGE=F1

NODE=M010M;LINKAGE=X

NODE=M010M;LINKAGE=P

NODE=M010M;LINKAGE=I

NODE=M010M;LINKAGE=L

NODE=M010M;LINKAGE=M

NODE=M010M;LINKAGE=K

NODE=M010M;LINKAGE=G

NODE=M010M;LINKAGE=D

NODE=M010M;LINKAGE=F

 $a_1(1260)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
250 to 600 OUR ESTIMATE				
420 ± 35 OUR AVERAGE				
380 ± 80	46M	1 AGHASYAN	18B COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
430 ± 24 ± 31		DARGENT	17 RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
422.01 ± 2.10 ± 12.72 894k		AAIJ	18AI LHCb	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
576 ± 11 ± 89		2 MIKHAENKO	18 RVUE	$\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$
367 ± 9 ± 28	420k	3 ALEKSEEV	10 COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
410 ± 31 ± 30		4 AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \rho^0 \rho^\pm \pi^\mp \gamma$
520–680	6360	5 LINK	07A FOCS	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
480 ± 20		6 GOMEZ-DUM..04	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$
580 ± 41	90k	7 SALVINI	04 OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
460 ± 85	205	7 DRUTSKOY	02 BELL	$B \rightarrow D^{(*)} K^- K^{*0}$
814 ± 36 ± 13	37k	8 ASNER	00 CLE2	$10.6 e^+ e^- \rightarrow \tau^+ \tau^-, \tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
450 ± 50	22k	9 AKHMETSHIN	99E CMD2	$1.05-1.38 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
570 ± 10		10 BONDAR	99 RVUE	$e^+ e^- \rightarrow 4\pi, \tau \rightarrow 3\pi \nu_\tau$
587 ± 27 ± 21	5904	11 ABREU	98G DLPH	$e^+ e^-$
478 ± 3 ± 15	5904	12 ABREU	98G DLPH	$e^+ e^-$
425 ± 14 ± 8	5904	13,14 ABREU	98G DLPH	$e^+ e^-$
400 ± 35		BARBERIS	99B	$450 pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
621 ± 32 ± 58		11,15 ACKERSTAFF	97R OPAL	$E_{cm}^{ee} = 88-94, \tau \rightarrow 3\pi \nu$

NODE=M010W

NODE=M010W

→ UNCHECKED ←

OCCUR=2

OCCUR=3

457	± 15	± 17	12,15	ACKERSTAFF	97R	OPAL	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94, \tau \rightarrow 3\pi\nu$	OCCUR=2
446	± 21	$+140$ -0	12	ALBRECHT	93C	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
239	± 11		ANDO		92	SPEC	$8\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
266	± 13	± 4	16	ANDO	92	SPEC	$8\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	OCCUR=3
465	$+228$ -143		17	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	
298	$+40$ -34		18	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=2
488	± 32		19	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
430	± 50		ARMSTRONG	90	OMEG	300.0 $p p \rightarrow p p \pi^+ \pi^- \pi^0$		
420	± 40		20	ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
396	± 43		21	BOWLER	88	RVUE		
405	± 75	± 25	BAND		87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
419	± 108	± 57	BAND		87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$	
521	± 27		ALBRECHT		86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=2
476	$+132$ -120	± 54	RUCKSTUHL		86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
462	± 56	± 30	SCHMIDKE		86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
292	± 40		BELLINI		85	SPEC	$40\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
380	± 100		22	DANKOWY...	81	SPEC	$8.45\pi^- p \rightarrow n3\pi$	
300	± 50		22	DAUM	81B	CNTR	$63.94\pi^- p \rightarrow p3\pi$	
230	± 50		23	GAVILLET	77	HBC	$4.2 K^- p \rightarrow \Sigma 3\pi$	

1 Statistical error negligible.

2 From the pole position. Using an amplitude analysis based on approximate three-body unitary of τ data from SCHAEL 05C.

3 Superseded by AGHASYAN 2018B.

4 The $\rho^\pm \pi^\mp$ state can be also due to the $\pi(1300)$.

5 Using the Breit-Wigner parameterization; strong correlation between mass and width.

6 Using the data of BARATE 98R.

7 From a fit of the $K^- K^{*0}$ distribution assuming $m_{a_1} = 1230$ MeV and purely resonant production of the $K^- K^{*0}$ system.

8 From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.

9 Using the $a_1(1260)$ mass of 1230 MeV.

10 From AKHMETSHIN 99E and ASNER 00 data using the $a_1(1260)$ mass of 1230 MeV.

11 Uses the model of KUHN 90.

12 Uses the model of ISGUR 89.

13 Includes the effect of a possible a'_1 state.

14 Uses the model of FEINDT 90.

15 Supersedes AKERS 95P.

16 Average and spread of values using 2 variants of the model of BOWLER 75.

17 Reanalysis of RUCKSTUHL 86.

18 Reanalysis of SCHMIDKE 86.

19 Reanalysis of ALBRECHT 86B.

20 From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

21 From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

22 Uses the model of BOWLER 75.

23 Produced in K^- backward scattering.

$a_1(1260)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $(\rho\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi$	seen
Γ_3 $(\rho\pi)_{D\text{-wave}}, \rho \rightarrow \pi\pi$	seen
Γ_4 $(\rho(1450)\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi$	seen
Γ_5 $(\rho(1450)\pi)_{D\text{-wave}}, \rho \rightarrow \pi\pi$	seen
Γ_6 $f_0(500)\pi, f_0 \rightarrow \pi\pi$	seen
Γ_7 $f_0(980)\pi, f_0 \rightarrow \pi\pi$	not seen
Γ_8 $f_0(1370)\pi, f_0 \rightarrow \pi\pi$	seen
Γ_9 $f_2(1270)\pi, f_2 \rightarrow \pi\pi$	seen
Γ_{10} $\pi^+ \pi^- \pi^0$	seen
Γ_{11} $\pi^0 \pi^0 \pi^0$	not seen
Γ_{12} $K\bar{K}\pi$	seen
Γ_{13} $K^*(892)K$	seen
Γ_{14} $\pi\gamma$	seen

NODE=M010W;LINKAGE=Q
NODE=M010W;LINKAGE=S

NODE=M010W;LINKAGE=R
NODE=M010W;LINKAGE=AU
NODE=M010W;LINKAGE=LI
NODE=M010W;LINKAGE=GO
NODE=M010W;LINKAGE=DR

NODE=M010W;LINKAGE=B6
NODE=M010W;LINKAGE=WE
NODE=M010W;LINKAGE=WB
NODE=M010W;LINKAGE=KS
NODE=M010W;LINKAGE=IM
NODE=M010W;LINKAGE=A1
NODE=M010W;LINKAGE=F1
NODE=M010W;LINKAGE=X
NODE=M010W;LINKAGE=P
NODE=M010W;LINKAGE=I
NODE=M010W;LINKAGE=L
NODE=M010W;LINKAGE=M
NODE=M010W;LINKAGE=K
NODE=M010W;LINKAGE=G
NODE=M010W;LINKAGE=D
NODE=M010W;LINKAGE=F

NODE=M010215;NODE=M010

DESIG=17;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=7;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=8;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=9;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=10;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=16;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=11;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=12;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=13;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=22
DESIG=23;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=18;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=14;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow

$a_1(1260)$ PARTIAL WIDTHS

$\Gamma(\pi\gamma)$		DOCUMENT ID	TECN	COMMENT	Γ_{14}
VALUE (keV)					
640±246		ZIELINSKI	84C SPEC	200 $\pi^+ Z \rightarrow Z 3\pi^-$	


D-wave/S-wave AMPLITUDE RATIO IN DECAY OF $a_1(1260) \rightarrow \rho\pi$

VALUE	DOCUMENT ID	TECN	COMMENT	
-0.062±0.020 OUR AVERAGE			Error includes scale factor of 2.3. See the ideogram below.	
-0.043±0.009±0.005	LINK	07A FOCS	$D^0 \rightarrow \pi^-\pi^+\pi^-\pi^+$	
-0.14 ± 0.04 ± 0.07	1 CHUNG	02 B852	$18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$	
-0.10 ± 0.02 ± 0.02	2,3 ACKERSTAFF	97R OPAL	$E_{cm}^{ee} = 88-94, \tau \rightarrow 3\pi\nu$	
-0.11 ± 0.02	2 ALBRECHT	93C ARG	$\tau^+ \rightarrow \pi^+\pi^+\pi^- \nu$	

1 Deck-type background not subtracted.

2 Uses the model of ISGUR 89.

3 Supersedes AKERS 95P.

**D-wave/S-wave AMPLITUDE RATIO IN DECAY OF $a_1(1260) \rightarrow \rho\pi$** **$a_1(1260)$ BRANCHING RATIOS**

$\Gamma((\rho\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi)/\Gamma_{\text{total}}$		Γ_2/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID
60.19	37k	1 ASNER 00 CLE2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.6 e^+ e^- \rightarrow \tau^+\tau^-, \tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

$\Gamma((\rho\pi)_D\text{-wave}, \rho \rightarrow \pi\pi)/\Gamma_{\text{total}}$		Γ_3/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID
1.30±0.60±0.22	37k	1 ASNER 00 CLE2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.6 e^+ e^- \rightarrow \tau^+\tau^-, \tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

$\Gamma((\rho(1450)\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi)/\Gamma_{\text{total}}$		Γ_4/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID
0.56±0.84±0.32	37k	1,2 ASNER 00 CLE2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.6 e^+ e^- \rightarrow \tau^+\tau^-, \tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

NODE=M010220

NODE=M010W4
NODE=M010W4

NODE=M010DS

NODE=M010DS

NODE=M010DS;LINKAGE=C
NODE=M010DS;LINKAGE=IM
NODE=M010DS;LINKAGE=X

NODE=M010225

NODE=M010R5
NODE=M010R5NODE=M010R6
NODE=M010R6NODE=M010R7
NODE=M010R7

$\Gamma((\rho(1450)\pi)_{D\text{-wave}}, \rho \rightarrow \pi\pi)/\Gamma_{\text{total}}$					Γ_5/Γ	NODE=M010R8 NODE=M010R8
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
$2.04 \pm 1.20 \pm 0.28$	37k	1, ² ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
$\Gamma(f_0(500)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$					Γ_6/Γ	NODE=M010R9 NODE=M010R9
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
seen		CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$		
$18.76 \pm 4.29 \pm 1.48$	37k	1, ³ ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$		
$\Gamma(f_0(500)\pi, f_0 \rightarrow \pi\pi)/\Gamma((\rho\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi)$					Γ_6/Γ_2	NODE=M010R4 NODE=M010R4
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
0.06 ± 0.05	90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$		
~ 0.3	28k	AKHMETSHIN	99E	CMD2 $1.05-1.38 e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$		
0.003 ± 0.003		⁴ LONGACRE	82	RVUE		
$\Gamma(f_0(980)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$					Γ_7/Γ	NODE=M010R10 NODE=M010R10
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
not seen	37k	ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$		
$\Gamma(f_0(1370)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$					Γ_8/Γ	NODE=M010R11 NODE=M010R11
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
$7.40 \pm 2.71 \pm 1.26$	37k	1, ⁵ ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$		
$\Gamma(f_2(1270)\pi, f_2 \rightarrow \pi\pi)/\Gamma_{\text{total}}$					Γ_9/Γ	NODE=M010R12 NODE=M010R12
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
$1.19 \pm 0.49 \pm 0.17$	37k	1, ⁶ ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$		
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{10}/Γ	NODE=M010R00 NODE=M010R00
VALUE		DOCUMENT ID		COMMENT		
seen		BARBERIS	98B	$450 pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$		
$\Gamma(\pi^0 \pi^0 \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$					Γ_{11}/Γ_{10}	NODE=M010R15 NODE=M010R15
VALUE	CL%	DOCUMENT ID		COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
<0.008	90	⁷ BARBERIS	01	$450 pp \rightarrow p_f 3\pi^0 p_s$		
$\Gamma(K^*(892)K)/\Gamma_{\text{total}}$					Γ_{13}/Γ	NODE=M010R13 NODE=M010R13
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
2.2 ± 0.5	2255	⁸ COAN	04	CLEO $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$		
8 to 15	205	⁹ DRUTSKOY	02	BELL $B \rightarrow D^{(*)} K^- K^{*0}$		
$3.3 \pm 0.5 \pm 0.1$	37k	¹⁰ ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$		
2.6 ± 0.3		¹¹ BARATE	99R	ALEP $\tau \rightarrow K\bar{K}\pi\nu_\tau$		

- 1 From a fit to the Dalitz plot.
 2 Assuming for $\rho(1450)$ mass and width of 1370 and 386 MeV respectively.
 3 Assuming for $f_0(500)$ (σ) mass and width of 860 and 880 MeV respectively.
 4 Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from GAVILET 77, DAUM 80, and DANKOWYCH 81.
 5 Assuming for $f_0(1370)$ mass and width of 1186 and 350 MeV respectively.
 6 Assuming for $f_2(1270)$ mass and width of 1275 and 185 MeV respectively.
 7 Inconsistent with observations of $\sigma\pi$, $f_0(1370)\pi$, and $f_2(1270)\pi$ decay modes.
 8 Using structure functions from KUHN 92 and DECKER 93A and $B(\tau^- \rightarrow K^-\pi^-K^+\nu_\tau) = (0.155 \pm 0.006 \pm 0.009)\%$ from BRIERE 03.
 9 From a comparison to ALAM 94 assuming purely resonant production of the K^-K^{*0} system.
 10 From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.
 11 Assuming $a_1(1260)$ dominance and taking $B(\tau \rightarrow a_1(1260)\nu_\tau)$ from BUSKULIC 96.

NODE=M010R;LINKAGE=B1
 NODE=M010R;LINKAGE=B2
 NODE=M010R;LINKAGE=B3
 NODE=M010R4;LINKAGE=E

 NODE=M010R;LINKAGE=B4
 NODE=M010R;LINKAGE=B5
 NODE=M010R;LINKAGE=RB
 NODE=M010R13;LINKAGE=CO

 NODE=M010R;LINKAGE=DR

 NODE=M010R;LINKAGE=B6
 NODE=M010R13;LINKAGE=BA

NODE=M010

$a_1(1260)$ REFERENCES			
AAIJ	18AI	EPJ C78 443	R. Aaij <i>et al.</i> (LHCb Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i> (COMPASS Collab.)
MIKHASENKO	18	PR D98 096021	M. Mikhasenko <i>et al.</i> (JPAC Collab.)
DARGENT	17	JHEP 1705 143	P. d'Argent <i>et al.</i> (HEID, BRIS)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i> (COMPASS Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i> (BABAR Collab.)
LINK	07A	PR D75 052003	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i> (PDG Collab.)
SCHAEL	05C	PRPL 421 191	S. Schael <i>et al.</i> (ALEPH Collab.)
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i> (CLEO Collab.)
GOMEZ-DUM...	04	PR D69 073002	D. Gomez Dunn, A. Pich, J. Portoles
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i> (OBELIX Collab.)
BRIERE	03	PRL 90 181802	R. A. Briere <i>et al.</i> (CLEO Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i> (BNL E852 Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i> (BELLE Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i> (CLEO Collab.)
AKHMETSHIN	99E	PL B466 392	R.R. Akhmetshin <i>et al.</i> (Novosibirsk CMD-2 Collab.)
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i> (ALEPH Collab.)
BONDAR	99	PL B466 403	A.E. Bondar <i>et al.</i> (Novosibirsk CMD-2 Collab.)
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i> (DELPHI Collab.)
BARATE	98R	EPJ C4 409	R. Barate <i>et al.</i> (ALEPH Collab.)
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i> (WA 102 Collab.)
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff <i>et al.</i> (OPAL Collab.)
BUSKULIC	96	ZPHY C70 579	D. Buskulic <i>et al.</i> (ALEPH Collab.)
AKERS	95P	ZPHY C67 45	R. Akers <i>et al.</i> (OPAL Collab.)
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i> (CLEO Collab.)
ALBRECHT	93C	ZPHY C58 61	H. Albrecht <i>et al.</i> (ARGUS Collab.)
DECKER	93A	ZPHY C58 445	R. Decker <i>et al.</i>
ANDO	92	PL B291 496	A. Ando <i>et al.</i> (KEK, KYOT, NIRS, SAGA+)
KUHN	92	ZPHY C56 661	J.H. Kuhn, E. Mirkes
IVANOV	91	ZPHY C49 563	Y.P. Ivanov, A.A. Osipov, M.K. Volkov (JINR)
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch (WA76 Coll.)
FEINDT	90	ZPHY C48 681	M. Feindt (HAMB)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i> (MPIM)
ISGUR	89	PR D39 1357	N. Isgur, C. Morningstar, C. Reader (TNTO)
BOWLER	88	PL B209 99	M.G. Bowler (OXF)
BAND	87	PL B198 297	H.R. Band <i>et al.</i> (MAC Collab.)
TORNQVIST	87	ZPHY C36 695	N.A. Tornqvist (HELS)
ALBRECHT	86B	ZPHY C33 7	H. Albrecht <i>et al.</i> (ARGUS Collab.)
RUCKSTUHL	86	PRl 56 2132	W. Ruckstuhl <i>et al.</i> (DELCO Collab.)
SCHMIDKE	86	PRL 57 527	W.B. Schmidke <i>et al.</i> (Mark II Collab.)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>
		Translated from YAF 41 1223.	
ZIELINSKI	84C	PRL 52 1195	M. Zielinski <i>et al.</i> (ROCH, MINN, FNAL)
LONGACRE	82	PR D26 82	R.S. Longacre (BNL)
DANKOWY...	81	PRL 46 580	J.A. Dankowycz <i>et al.</i> (TNTO, BNL, CARL+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i> (AMST, CERN, CRAC, MPIM+)
DAUM	80	PL 89B 281	C. Daum <i>et al.</i> (AMST, CERN, CRAC, MPIM+) JP
GAVILLET	77	PL 69B 119	P. Gavillet <i>et al.</i> (AMST, CERN, NIJM+) JP
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i> (OXFTP, DARE)

REFID=59187
 REFID=59471
 REFID=59487
 REFID=58121
 REFID=53356
 REFID=52049
 REFID=51713
 REFID=51004
 REFID=50845
 REFID=49945
 REFID=49771
 REFID=53226
 REFID=49360
 REFID=48837
 REFID=48780
 REFID=48324
 REFID=47339
 REFID=47411
 REFID=47366
 REFID=47358
 REFID=45909
 REFID=46147
 REFID=46345
 REFID=45616
 REFID=44588
 REFID=44366
 REFID=43738
 REFID=43310
 REFID=51577
 REFID=43171
 REFID=51576
 REFID=41750
 REFID=41375
 REFID=45912
 REFID=45862
 REFID=40730
 REFID=40578
 REFID=40263
 REFID=40030
 REFID=20884
 REFID=10349
 REFID=10350
 REFID=47490

 REFID=20882
 REFID=20878
 REFID=20572
 REFID=20872
 REFID=20868
 REFID=20852
 REFID=20571

$f_2(1270)$ $I^G(J^{PC}) = 0^+(2^{++})$

NODE=M005

 $f_2(1270)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1275.5 ± 0.8 OUR AVERAGE					
1275.8 ± 1.0 ± 0.4		¹ BOGOLYUB...	13	SPEC $7\pi^+(K^+, p)A \rightarrow n\gamma + X$	
1262 $\pm \frac{1}{2} \pm 8$		² ABLIKIM	06v	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
1275 ± 15		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$	
1283 ± 5		ALDE	98	GAM4 $100\pi^-p \rightarrow \pi^0\pi^0n$	
1278 ± 5		³ BERTIN	97c	OBLX $0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
1272 ± 8	200k	PROKOSHKIN	94	GAM2 $38\pi^-p \rightarrow \pi^0\pi^0n$	
1269.7 ± 5.2	5730	AUGUSTIN	89	DM2 $e^+e^- \rightarrow 5\pi$	
1283 ± 8	400	⁴ ALDE	87	GAM4 $100\pi^-p \rightarrow 4\pi^0n$	
1274 ± 5		⁴ AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$	
1283 ± 6		⁵ LONGACRE	86	MPS $22\pi^-p \rightarrow n2K_S^0$	
1276 ± 7		COURAU	84	DLCO $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$	
1273.3 ± 2.3		⁶ CHABAUD	83	ASPK $17\pi^-p$ polarized	
1280 ± 4		⁷ CASON	82	STRC $8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$	
1281 ± 7	11600	GIDAL	81	MRK2 J/ψ decay	
1282 ± 5		⁸ CORDEN	79	OMEG $12\text{--}15\pi^-p \rightarrow n2\pi$	
1269 ± 4	10k	APEL	75	NICE $40\pi^-p \rightarrow n2\pi^0$	
1272 ± 4	4600	ENGLER	74	DBC $6\pi^+n \rightarrow \pi^+\pi^-p$	
1277 ± 4	5300	FLATTE	71	HBC $7.0\pi^+p$	
1273 ± 8		⁴ STUNTEBECK	70	HBC $8\pi^-p, 5.4\pi^+d$	
1265 ± 8		BOESEBECK	68	HBC $8\pi^+p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1259 $\pm 4 \pm 4$	1.7k	^{9,10} DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
1267 $\pm 4 \pm 3$	1.5k	^{9,10} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$	
1270 ± 8		¹¹ ANISOVICH	09	RVUE $0.0\bar{p}p, \pi N$	
1277 ± 6	870	¹² SCHEGELSKY	06a	RVUE $\gamma\gamma \rightarrow K_S^0K_S^0$	
1251 ± 10		TIKHOMIROV	03	SPEC $40.0\pi^-C \rightarrow K_S^0K_S^0K_L^0X$	
1260 ± 10		¹³ ALDE	97	GAM2 $450pp \rightarrow pp\pi^0\pi^0$	
1278 ± 6		¹³ GRYGOREV	96	SPEC $40\pi^-N \rightarrow K_S^0K_S^0X$	
1262 ± 11		AGUILAR-...	91	EHS $400pp$	
1275 ± 10		AKER	91	CBAR $0.0\bar{p}p \rightarrow 3\pi^0$	
1220 ± 10		BREAKSTONE	90	SFM $pp \rightarrow pp\pi^+\pi^-$	
1288 ± 12		ABACHI	86b	HRS $e^+e^- \rightarrow \pi^+\pi^-X$	
1284 ± 30	3k	BINON	83	GAM2 $38\pi^-p \rightarrow n2\eta$	
1280 ± 20	3k	APEL	82	CNTR $25\pi^-p \rightarrow n2\pi^0$	
1284 ± 10	16000	DEUTSCH...	76	HBC $16\pi^+p$	
1258 ± 10	600	TAKAHASHI	72	HBC $8\pi^-p \rightarrow n2\pi$	
1275 ± 13		ARMENISE	70	HBC $9\pi^+n \rightarrow p\pi^+\pi^-$	
1261 ± 5	1960	⁴ ARMENISE	68	DBC $5.1\pi^+n \rightarrow p\pi^+MM^-$	
1270 ± 10	360	⁴ ARMENISE	68	DBC $5.1\pi^+n \rightarrow p\pi^0MM$	
1268 ± 6		¹⁴ JOHNSON	68	HBC $3.7\text{--}4.2\pi^-p$	

1 Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.

2 Breit-Wigner mass.

3 T-matrix pole.

4 Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

5 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

6 From an energy-independent partial-wave analysis.

7 From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$.

8 From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.

9 Using CLEO-c data but not authored by the CLEO Collaboration.

10 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 185$ MeV.

11 4-poles, 5-channel K matrix fit.

12 From analysis of L3 data at 91 and 183–209 GeV.

13 Systematic uncertainties not estimated.

14 JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

NODE=M005M

NODE=M005M

OCCUR=2

NODE=M005M;LINKAGE=B

NODE=M005M;LINKAGE=K

NODE=M005M;LINKAGE=A

NODE=M005M;LINKAGE=T

NODE=M005M;LINKAGE=L

NODE=M005M;LINKAGE=O

NODE=M005M;LINKAGE=P

NODE=M005M;LINKAGE=S

NODE=M005M;LINKAGE=C

NODE=M005M;LINKAGE=D

NODE=M005M;LINKAGE=AN

NODE=M005M;LINKAGE=SC

NODE=M005M;LINKAGE=QQ

NODE=M005M;LINKAGE=J

$f_2(1270)$ WIDTH

NODE=M005W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

186.7 \pm 2.2 OUR FIT Error includes scale factor of 1.4.

185.9 \pm 2.8 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

190.3 \pm 1.9 \pm 1.8	1 BOGOLYUB...	13	SPEC	$7\pi^+(K^+, p)A \rightarrow n\gamma + X$	
175 \pm 6 \pm 10	2 ABLIKIM	06v	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
190 \pm 20	ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$	
171 \pm 10	ALDE	98	GAM4	$100\pi^-p \rightarrow \pi^0\pi^0n$	
204 \pm 20	3 BERTIN	97c	OBLX	$0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
192 \pm 5	PROKOSHKIN	94	GAM2	$38\pi^-p \rightarrow \pi^0\pi^0n$	
180 \pm 24	AGUILAR....	91	EHS	$400pp$	
169 \pm 9	4 AUGUSTIN	89	DM2	$e^+e^- \rightarrow 5\pi$	
150 \pm 30	4 ALDE	87	GAM4	$100\pi^-p \rightarrow 4\pi^0n$	
186 \pm 9	5 LONGACRE	86	MPS	$22\pi^-p \rightarrow n2K_S^0$	
179.2 \pm 6.9	6 CHABAUD	83	ASPK	$17\pi^-p$ polarized	
160 \pm 11	DENNEY	83	LASS	$10\pi^+N$	
196 \pm 10	3k APEL	82	CNTR	$25\pi^-p \rightarrow n2\pi^0$	
152 \pm 9	7 CASON	82	STRC	$8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$	
186 \pm 27	11600 GIDAL	81	MRK2	J/ψ decay	
216 \pm 13	8 CORDEN	79	OMEG	$12\text{--}15\pi^-p \rightarrow n2\pi$	
190 \pm 10	APEL	75	NICE	$40\pi^-p \rightarrow n2\pi^0$	
192 \pm 16	4600 ENGLER	74	DBC	$6\pi^+n \rightarrow \pi^+\pi^-p$	
183 \pm 15	5300 FLATTE	71	HBC	$7\pi^+p \rightarrow \Delta^{++}f_2$	
196 \pm 30	4 STUNTEBECK	70	HBC	$8\pi^-p, 5.4\pi^+d$	
216 \pm 20	4 ARMENISE	68	DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$	OCCUR=2
128 \pm 27	4 BOESEBECK	68	HBC	$8\pi^+p$	
176 \pm 21	4.9 JOHNSON	68	HBC	$3.7\text{--}4.2\pi^-p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
194 \pm 36	10 ANISOVICH	09	RVUE	$0.0\bar{p}p, \pi N$	
195 \pm 15	870 SCHEGELSKY	06a	RVUE	$\gamma\gamma \rightarrow K_S^0K_S^0$	
121 \pm 26	TIKHOMIROV	03	SPEC	$40.0\pi^-C \rightarrow K_S^0K_S^0K_L^0X$	
187 \pm 20	12 ALDE	97	GAM2	$450pp \rightarrow pp\pi^0\pi^0$	
184 \pm 10	12 GRYGOREV	96	SPEC	$40\pi^-N \rightarrow K_S^0K_S^0X$	
200 \pm 10	AKER	91	CBAR	$0.0\bar{p}p \rightarrow 3\pi^0$	
240 \pm 40	3k BINON	83	GAM2	$38\pi^-p \rightarrow n2\eta$	
187 \pm 30	650 ANTIPOV	77	CIBS	$25\pi^-p \rightarrow p3\pi$	
225 \pm 38	16000 DEUTSCH...	76	HBC	$16\pi^+p$	
166 \pm 28	600 TAKAHASHI	72	HBC	$8\pi^-p \rightarrow n2\pi$	
173 \pm 53	4 ARMENISE	70	HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$	OCCUR=2

¹ Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.

² Breit-Wigner width

³ T-matrix pole.

⁴ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁵ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

⁶ From an energy-independent partial-wave analysis.

⁷ From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$.

⁸ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.

⁹ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

¹⁰ 4-poles, 5-channel K matrix fit.

¹¹ From analysis of L3 data at 91 and 183–209 GeV.

¹² Systematic uncertainties not estimated.

NODE=M005W;LINKAGE=C

NODE=M005W;LINKAGE=D

NODE=M005W;LINKAGE=QA

NODE=M005W;LINKAGE=T

NODE=M005W;LINKAGE=L

NODE=M005W;LINKAGE=R

NODE=M005W;LINKAGE=Q


NODE=M005W;LINKAGE=U

NODE=M005W;LINKAGE=J

NODE=M005W;LINKAGE=AN

NODE=M005W;LINKAGE=SC

NODE=M005W;LINKAGE=QQ

 **$f_2(1270)$ width (MeV)** **$f_2(1270)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
$\Gamma_1 \pi\pi$	(84.2 $^{+2.9}_{-0.9}$) %	S=1.1	DESIG=1
$\Gamma_2 \pi^+ \pi^- 2\pi^0$	(7.7 $^{+1.1}_{-3.2}$) %	S=1.2	DESIG=3
$\Gamma_3 K\bar{K}$	(4.6 $^{+0.5}_{-0.4}$) %	S=2.7	DESIG=4
$\Gamma_4 2\pi^+ 2\pi^-$	(2.8 ± 0.4) %	S=1.2	DESIG=2
$\Gamma_5 \eta\eta$	(4.0 ± 0.8) $\times 10^{-3}$	S=2.1	DESIG=7
$\Gamma_6 4\pi^0$	(3.0 ± 1.0) $\times 10^{-3}$		DESIG=9
$\Gamma_7 \gamma\gamma$	(1.42 ± 0.24) $\times 10^{-5}$	S=1.4	DESIG=8
$\Gamma_8 \eta\pi\pi$	< 8 $\times 10^{-3}$	CL=95%	DESIG=6
$\Gamma_9 K^0 K^- \pi^+ + \text{c.c.}$	< 3.4 $\times 10^{-3}$	CL=95%	DESIG=5
$\Gamma_{10} e^+ e^-$	< 6 $\times 10^{-10}$	CL=90%	DESIG=10

NODE=M005215;NODE=M005

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 45 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 83.0$ for 38 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-90						
x_3	10	-39					
x_4	10	-38	1				
x_5	1	-6	0	0			
x_6	0	-7	0	0	0		
x_7	3	1	-15	0	0	0	
Γ	-71	65	-10	-7	-1	0	-6
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode		Rate (MeV)	Scale factor	
Γ_1	$\pi\pi$	157.2	$+4.0$ -1.1	DESIG=1
Γ_2	$\pi^+ \pi^- 2\pi^0$	14.4	$+2.1$ -6.0	1.2 DESIG=3
Γ_3	$K\bar{K}$	8.5	± 0.8	2.8 DESIG=4
Γ_4	$2\pi^+ 2\pi^-$	5.2	± 0.7	1.2 DESIG=2
Γ_5	$\eta\eta$	0.75	± 0.14	2.1 DESIG=7
Γ_6	$4\pi^0$	0.56	± 0.19	DESIG=9
Γ_7	$\gamma\gamma$	0.0026 ± 0.0005	1.4 DESIG=8	

$f_2(1270)$ PARTIAL WIDTHS

$\Gamma(\pi\pi)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

157.2 $^{+4.0}_{-1.1}$ OUR FIT

157.0 $^{+6.0}_{-1.0}$ 1 LONGACRE 86 MPS $22\pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

152 ± 8 870 2 SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

Γ_1

NODE=M005220

NODE=M005W1
NODE=M005W1

$\Gamma(K\bar{K})$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

8.5 ± 0.8 OUR FIT Error includes scale factor of 2.8.

9.0 $^{+0.7}_{-0.3}$ 1 LONGACRE 86 MPS $22\pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5 ± 2.0 870 2 SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

Γ_3

NODE=M005W4
NODE=M005W4

$\Gamma(\eta\eta)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.75 ± 0.14 OUR FIT Error includes scale factor of 2.1.

1.0 ± 0.1 1 LONGACRE 86 MPS $22\pi^- p \rightarrow n2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.4 870 2 SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

Γ_5

NODE=M005W7
NODE=M005W7

$\Gamma(\gamma\gamma)$

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an S-wave contribution, values are systematically higher (typically around 3 keV).

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR FIT				Error includes scale factor of 1.4.
2.93± 0.40		3 DAI	14A	RVUE Compilation

Γ_7

NODE=M005W8
NODE=M005W8

NODE=M005W8

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.14±0.20	4.5 PENNINGTON 08	RVUE	Compilation	OCCUR=2
3.82±0.30	5.6 PENNINGTON 08	RVUE	Compilation	OCCUR=3
2.55±0.15	870 2 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
2.84±0.35	BOGLIONE 99	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$	
2.93±0.23±0.32	7 YABUKI 95	VNS		
2.58±0.13 ^{+0.36} _{-0.27}	8 BEHREND 92	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
3.10±0.35±0.35	9 BLINOV 92	MD1	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
2.27±0.47±0.11	ADACHI 90D	TOPZ	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
3.15±0.04±0.39	BOYER 90	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
3.19±0.16 ^{+0.29} _{-0.28}	MARSISKE 90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
2.35±0.65	10 MORGAN 90	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$	
3.19±0.09 ^{+0.22} _{-0.38}	2177 OEST 90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
3.2 ±0.1 ±0.4	11 AIHARA 86B	TPC	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
2.5 ±0.1 ±0.5	BEHREND 84B	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
2.85±0.25±0.5	12 BERGER 84	PLUT	$e^+ e^- \rightarrow e^+ e^- 2\pi$	
2.70±0.05±0.20	COURAU 84	DLCO	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
2.52±0.13±0.38	13 SMITH 84C	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
2.7 ±0.2 ±0.6	EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$	
2.9 ^{+0.6} _{-0.4} ±0.6	14 EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$	OCCUR=2
3.2 ±0.2 ±0.6	BRANDELIK 81B	TASS	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
3.6 ±0.3 ±0.5	ROUSSARIE 81	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$	
2.3 ±0.8	15 BERGER 80B	PLUT	$e^+ e^-$	

$\Gamma(e^+ e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}
<0.11	90	ACHASOV 00K	SND	$e^+ e^- \rightarrow \pi^0 \pi^0$	NODE=M005W9 NODE=M005W9
<1.7	90	VOROBYEV 88	ND	$e^+ e^- \rightarrow \pi^0 \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 1 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- 2 From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.
- 3 Based on a K-matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.
- 4 Solution A (preferred solution based on χ^2 -analysis).
- 5 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.
- 6 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).
- 7 With a narrow scalar state around 1220 MeV.
- 8 Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.
- 9 Using the unitarized model of LYTH 85.
- 10 Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1370)$: $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$ KeV.
- 11 Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.
- 12 Using the MENNESSIER 83 model.
- 13 Superseded by BOYER 90.
- 14 If helicity = 2 assumption is not made.
- 15 Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.

$f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_3\Gamma_7/\Gamma$
0.121±0.020 OUR FIT	Error includes scale factor of 1.3.			
0.091±0.007±0.027	¹ ALBRECHT 90G	ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	NODE=M005G1 NODE=M005G1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.104±0.007±0.072	² ALBRECHT 90G	ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	OCCUR=2

1 Using an incoherent background.

2 Using a coherent background.

NODE=M005PW;LINKAGE=L
NODE=M005W1;LINKAGE=SC
NODE=M005W8;LINKAGE=A

NODE=M005W8;LINKAGE=P1
NODE=M005W8;LINKAGE=P3

NODE=M005W8;LINKAGE=P2

NODE=M005W8;LINKAGE=YA
NODE=M005W;LINKAGE=B
NODE=M005W;LINKAGE=A
NODE=M005PW;LINKAGE=C

NODE=M005PW;LINKAGE=B

NODE=M005PW;LINKAGE=X
NODE=M005PW;LINKAGE=V
NODE=M005PW;LINKAGE=H
NODE=M005PW;LINKAGE=A

NODE=M005223

NODE=M005G1
NODE=M005G1

OCCUR=2

NODE=M005G1;LINKAGE=A
NODE=M005G1;LINKAGE=K

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_5\Gamma_7/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
11.5^{+1.8+4.5}_{-2.0-3.7}	1 UEHARA	10A BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$		NODE=M005G02 NODE=M005G02

1 Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_0(Y)$.

Helicity-0/Helicity-2 RATIO IN $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$					
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT		
3.7^{+15.9}_{-2.9}	UEHARA	08A BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5 ± 1.8	¹ DAI	14A RVUE	Compilation
13	^{2,3} PENNINGTON 08	RVUE	Compilation
26	^{3,4} PENNINGTON 08	RVUE	Compilation

1 Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.

2 Solution A (preferred solution based on χ^2 -analysis).

3 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

4 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

$f_2(1270)$ BRANCHING RATIOS					
$\Gamma(\pi\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.842^{+0.029}_{-0.009} OUR FIT		Error includes scale factor of 1.1.			

0.837 ± 0.020 OUR AVERAGE

0.849 ± 0.025	CHABAUD	83 ASPK	17 $\pi^- p$ polarized
0.85 ± 0.05	BEAUPRE	71 HBC	8 $\pi^+ p \rightarrow \Delta^{++} f_2$
0.8 ± 0.04	OH	70 HBC	1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(\pi\pi)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
Should be twice $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$ if decay is $\rho\rho$. (See ASCOLI 68D.)					

0.091^{+0.014}_{-0.040} OUR FIT Error includes scale factor of 1.2.

0.15 ± 0.06 600 EISENBERG 74 HBC $4.9 \pi^+ p \rightarrow \Delta^{++} f_2$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 EMMS 75D DBC $4 \pi^+ n \rightarrow p f_2$

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$					Γ_3/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.					

0.054^{+0.005}_{-0.006} OUR FIT Error includes scale factor of 2.7.

0.041 ± 0.004 OUR AVERAGE

0.045 ± 0.01	¹ BARGIOTTI	03 OBLX	$\bar{p}p$
0.037 ± 0.008 _{-0.021}	ETKIN	82B MPS	$23 \pi^- p \rightarrow n 2K_S^0$
0.045 ± 0.009	CHABAUD	81 ASPK	17 $\pi^- p$ polarized
0.039 ± 0.008	LOVERRE	80 HBC	$4 \pi^- p \rightarrow K\bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.052 ± 0.025	ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
0.036 ± 0.005	² COSTA	80 OMEG	1-2.2 $\pi^- p \rightarrow K^+ K^- n$
0.030 ± 0.005	³ MARTIN	79 RVUE	
0.027 ± 0.009	⁴ POLYCHRO...	79 STRC	$7 \pi^- p \rightarrow n 2K_S^0$
0.025 ± 0.015	EMMS	75D DBC	$4 \pi^+ n \rightarrow p f_2$
0.031 ± 0.012	20 ADERHOLZ	69 HBC	$8 \pi^+ p \rightarrow K^+ K^- \pi^+ p$

NODE=M005R2
NODE=M005R2

NODE=M005G02;LINKAGE=UE

NODE=M005HR0
NODE=M005HR0

OCCUR=2
OCCUR=3
NODE=M005HR0;LINKAGE=A

NODE=M005HR0;LINKAGE=P1
NODE=M005HR0;LINKAGE=P3
NODE=M005HR0;LINKAGE=P2

NODE=M005225

NODE=M005R10
NODE=M005R10

NODE=M005R2
NODE=M005R2
NODE=M005R2

NODE=M005R3
NODE=M005R3
NODE=M005R3

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
0.033±0.005 OUR FIT		Error includes scale factor of 1.2.			NODE=M005R1 NODE=M005R1
0.033±0.004 OUR AVERAGE		Error includes scale factor of 1.1.			
0.024±0.006	160	EMMS	75D DBC	$4\pi^+ n \rightarrow p f_2$	
0.051±0.025	70	EISENBERG	74 HBC	$4.9\pi^+ p \rightarrow \Delta^{++} f_2$	
$0.043^{+0.007}_{-0.011}$	285	LOUIE	74 HBC	$3.9\pi^- p \rightarrow n f_2$	
0.037±0.007	154	ANDERSON	73 DBC	$6\pi^+ n \rightarrow p f_2$	
0.047±0.013	OH		70 HBC	$1.26\pi^- p \rightarrow \pi^+\pi^- n$	

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
4.0±0.8 OUR FIT	Error includes scale factor of 2.1.			NODE=M005R7 NODE=M005R7

2.9±0.5 OUR AVERAGE

2.7±0.7	BINON	05 GAMS	$33\pi^- p \rightarrow \eta\eta n$	
2.8±0.7	ALDE	86D GAM4	$100\pi^- p \rightarrow 2\eta n$	
5.2±1.7	BINON	83 GAM2	$38\pi^- p \rightarrow 2\eta n$	

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_1
0.003±0.001		BARBERIS	00E	$450pp \rightarrow p_f\eta\eta p_s$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.05	95	EDWARDS	82F CBAL	$e^+e^- \rightarrow e^+e^- 2\eta$
<0.016	95	EMMS	75D DBC	$4\pi^+ n \rightarrow p f_2$
<0.09	95	EISENBERG	74 HBC	$4.9\pi^+ p \rightarrow \Delta^{++} f_2$

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
0.0030±0.0010 OUR FIT					NODE=M005R11 NODE=M005R11

0.003 ± 0.001

400 ± 50 ALDE 87 GAM4 $100\pi^- p \rightarrow 4\pi^0 n$

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				NODE=M005R13 NODE=M005R13

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.57±0.01 $^{+1.39}_{-0.14}$	UEHARA	08A BELL	$10.6e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	
------------------------------	--------	----------	---	--

 $\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_1
<0.010	95	EMMS	75D DBC	$4\pi^+ n \rightarrow p f_2$	NODE=M005R5 NODE=M005R5

 $\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_1
<0.004	95	EMMS	75D DBC	$4\pi^+ n \rightarrow p f_2$	NODE=M005R4 NODE=M005R4

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-10})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
<6	90	ACHASOV	00K SND	$e^+e^- \rightarrow \pi^0\pi^0$	

1 Coupled channel analysis of $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, and $K^\pm K_S^0\pi^\mp$.

2 Re-evaluated by CHABAUD 83.

3 Includes PAWLICKI 77 data.

4 Takes into account the $f_2(1270)-f'_2(1525)$ interference.

 $f_2(1270)$ REFERENCES

DOBBS 15 PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
DAI 14A PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
BOGOLYUB... 13 PAN 76 1324	M.Yu. Bogolyubsky <i>et al.</i>	(HYPERON-M Collab.)
	Translated from YAF 76 1389.	
UEHARA 13 PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
UEHARA 10A PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH 09 IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
UEHARA 09 PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
PDG 08 PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
PENNINGTON 08 EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA 08A PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
MORI 07 PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)
ABLIKIM 06V PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)
SCHEGELSKY 06A EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
ABLIKIM 05 PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
BINON 05 PAN 68 960	F. Binon <i>et al.</i>	
	Translated from YAF 68 998.	

NODE=M005

REFID=56805

REFID=55923

REFID=55585

REFID=55592

REFID=53641

REFID=52719

REFID=52761

REFID=52166

REFID=52303

REFID=52309

REFID=51652

REFID=51507

REFID=51185

REFID=50450

REFID=50780

NODE=M005R;LINKAGE=BG

NODE=M005R3;LINKAGE=D

NODE=M005R3;LINKAGE=F

NODE=M005R3;LINKAGE=M

ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
GRYGOREV	96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov	(ITEP)	REFID=45566
		Translated from YAF 59 2187.			
YABUKI	95	JPSJ 64 435	F. Yabuki <i>et al.</i>	(VENUS Collab.)	REFID=46384
PROKOSHKIN	94	PD 39 420	Y.D. Prokoshkin, A.A. Kondashov	(SERP)	REFID=44094
		Translated from DANS 336 613.			
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
BLINOV	92	ZPHY C53 33	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41858
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	REFID=41587
ADACHI	90D	PL B234 185	I. Adachi <i>et al.</i>	(TOPAZ Collab.)	REFID=41345
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LAJO, CLER, FRAS+)	REFID=40268
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=20764
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LANDRO	86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen	(UTRO)	REFID=20767
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
LYTH	85	JP G11 459	D.H. Lyth		REFID=42169
BEHREND	84B	ZPHY C23 223	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20757
BERGER	84	ZPHY C26 199	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20760
COURAU	84	PL 147B 227	A. Courau <i>et al.</i>	(CIT, SLAC)	REFID=20758
SMITH	84C	PR D30 851	J.R. Smith <i>et al.</i>	(SLAC, LBL, HARV)	REFID=20759
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20751
		Translated from YAF 38 934.			
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)	REFID=20393
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20745
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)	REFID=20741
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)	REFID=20388
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20736
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)	REFID=20382
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
PDG	78	PL 75B 1	C. Bricman <i>et al.</i>		REFID=40124
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)	REFID=20367
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)	REFID=20721
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=20715
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)	REFID=20719
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)	REFID=20710
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)	REFID=20103
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)	REFID=20698
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)	REFID=20700
ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP	REFID=20335
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)	REFID=20681
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)	REFID=20065
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
DERADO	65	PR 14 872	I. Derado <i>et al.</i>	(NDAM)	REFID=20668
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)	REFID=20663
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)	REFID=20657

$f_1(1285)$ $I^G(J^{PC}) = 0^+(1^{++})$

NODE=M008

 $f_1(1285)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1281.9 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.			
1281.0 ± 0.8		DICKSON	16	$\gamma p \rightarrow \eta \pi^+ \pi^- p$	
1287.4 ± 3.0	87	ABLIKIM	15P	$J/\psi \rightarrow K^+ K^- 3\pi$	
1281.16 ± 0.39 ± 0.45		¹ LEES	12X	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$	
1285.1 ± 1.0 + 1.6 - 0.3		² ABLIKIM	11J	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$	
1281 ± 2 ± 1		AUBERT	07AU	$BABR 10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$	
1276.1 ± 8.1 ± 8.0	203	BAI	04J	$BES2 J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$	
1274 ± 6	237	ABDALLAH	03H	$DLPH 91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$	
1280 ± 4		ACCIARRI	01G	L3	
1288 ± 4 ± 5	20k	ADAMS	01B	$B852 18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$	
1284 ± 6	1400	ALDE	97B	$GAM4 100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$	
1281 ± 1		BARBERIS	97B	$OMEG 450 pp \rightarrow pp2(\pi^+ \pi^-)$	
1281 ± 1		BARBERIS	97C	$OMEG 450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$	
1280 ± 2		³ ANTINORI	95	$OMEG 300,450 pp \rightarrow pp2(\pi^+ \pi^-)$	
1282.2 ± 1.5		LEE	94	$MPS2 18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$	
1279 ± 5		FUKUI	91C	$SPEC 8.95 \pi^- p \rightarrow \eta \pi^+ \pi^- n$	
1278 ± 2	140	ARMSTRONG	89	$OMEG 300 pp \rightarrow K \bar{K} \pi pp$	
1278 ± 2		ARMSTRONG	89G	$OMEG 85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$	
1280.1 ± 2.1	60	RATH	89	$MPS 21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$	
1285 ± 1	4750	⁴ BIRMAN	88	$MPS 8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$	
1280 ± 1	504	BITYUKOV	88	$SPEC 32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$	
1280 ± 4		ANDO	86	$SPEC 8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$	
1277 ± 2	420	REEVES	86	$SPEC 6.6 p\bar{p} \rightarrow K K \pi X$	
1285 ± 2		CHUNG	85	$SPEC 8 \pi^- p \rightarrow N K \bar{K} \pi$	
1279 ± 2	604	ARMSTRONG	84	$OMEG 85 \pi^+ p \rightarrow K \bar{K} \pi \pi p, pp \rightarrow K \bar{K} \pi pp$	
1286 ± 1		CHAUVAT	84	$SPEC ISR 31.5 pp$	
1278 ± 4		EVANGELIS...	81	$OMEG 12 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$	
1283 ± 3	103	DIONISI	80	$HBC 4 \pi^- p \rightarrow K \bar{K} \pi n$	
1282 ± 2	320	NACASCH	78	$HBC 0.7, 0.76 \bar{p}p \rightarrow K \bar{K} 3\pi$	
1279 ± 5	210	GRASSLER	77	$HBC 16 \pi^\mp p$	
1286 ± 3	180	DUBOC	72	$HBC 1.2 \bar{p}p \rightarrow 2K 4\pi$	
1283 ± 5		DAHL	67	$HBC 1.6-4.2 \pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1289.3 ± 2.8	234	ABLIKIM	19BA	$BES3 e^+ e^- \rightarrow \psi(2S)$	
1284.2 ± 2.2		⁵ AAIJ	14Y	$LHCb \bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$	
1281.9 ± 0.5		⁵ SOSA	99	$SPEC pp \rightarrow p_{slow} (K_S^0 K^+ \pi^-) p_{fast}$	
1282.8 ± 0.6		⁵ SOSA	99	$SPEC pp \rightarrow p_{slow} (K_S^0 K^- \pi^+) p_{fast}$	OCCUR=2
1270 ± 10		AMELIN	95	$VES 37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$	
1280 ± 2		ABATZIS	94	$OMEG 450 pp \rightarrow pp2(\pi^+ \pi^-)$	
1282 ± 4		ARMSTRONG	93C	$E760 \bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
1270 ± 6 ± 10		ARMSTRONG	92C	$OMEG 300 pp \rightarrow pp\pi^+ \pi^- \gamma$	
1281 ± 1		ARMSTRONG	89E	$OMEG 300 pp \rightarrow pp2(\pi^+ \pi^-)$	
1279 ± 6 ± 10	16	BECKER	87	$MRK3 e^+ e^- \rightarrow \phi K \bar{K} \pi$	
1286 ± 9		GIDAL	87	$MRK2 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$	
1287 ± 5	353	BITYUKOV	84B	$SPEC 32 \pi^- p \rightarrow K^+ K^- \pi^0 n$	
~ 1279		⁶ TORNQVIST	82B	$RVUE$	

NODE=M008M

NODE=M008M

1275	\pm 6	31	BROMBERG	80	SPEC	100	$\pi^- p \rightarrow K\bar{K}\pi X$
1288	\pm 9	200	GURTU	79	HBC	4.2	$K^- p \rightarrow n\eta 2\pi$
~ 1275.0		46	⁷ STANTON	79	CNTR	8.5	$\pi^- p \rightarrow n2\gamma 2\pi$
1271	\pm 10	34	CORDEN	78	OMEG	12–15	$\pi^- p \rightarrow K^+ K^- \pi n$
1295	\pm 12	85	CORDEN	78	OMEG	12–15	$\pi^- p \rightarrow n5\pi$
1292	\pm 10	150	DEFOIX	72	HBC	0.7	$\bar{p}p \rightarrow 7\pi$
1280	\pm 3	500	⁸ THUN	72	MMS	13.4	$\pi^- p$
1303	\pm 8		BARDADIN...	71	HBC	8	$\pi^+ p \rightarrow p6\pi$
1283	\pm 6		BOESEBECK	71	HBC	16.0	$\pi p \rightarrow p5\pi$
1270	\pm 10		CAMPBELL	69	DBC	2.7	$\pi^+ d$
1285	\pm 7		LORSTAD	69	HBC	0.7	$\bar{p}p$, 4,5-body
1290	\pm 7		D'ANDLAU	68	HBC	1.2	$\bar{p}p$, 5–6 body

1 Using the $2\pi^+ 2\pi^-$ and $\pi^+ \pi^- \eta$ modes of $f_1(1285)$ decay.

2 The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

3 Supersedes ABATZIS 94, ARMSTRONG 89E.

4 From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

5 No systematic error given.

6 From a unitarized quark-model calculation.

7 From phase shift analysis of $\eta \pi^+ \pi^-$ system.

8 Seen in the missing mass spectrum.

OCCUR=2

NODE=M008M;LINKAGE=LE

NODE=M008M;LINKAGE=BL

NODE=M008M;LINKAGE=B


NODE=M008M;LINKAGE=A

NODE=M008M;LINKAGE=N1

NODE=M008M;LINKAGE=T

NODE=M008M;LINKAGE=P

NODE=M008M;LINKAGE=S



$f_1(1285)$ mass (MeV)

$f_1(1285)$ WIDTH

NODE=M008W

Only experiments giving width error less than 20 MeV are kept for averaging.

NODE=M008W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
22.7\pm 1.1 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.	
18.4 \pm 1.4		DICKSON	16	CLAS	2.55 $\gamma p \rightarrow \eta \pi^+ \pi^- p$

NODE=M008W

18.3 ± 6.3	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$	
22.0 ± 3.1 ^{+ 2.0} _{- 1.5}	1	ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$	
35 ± 6 ± 4		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$	
40.0 ± 8.6 ± 9.3	203	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
29 ± 12	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$	
45 ± 9 ± 7	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$	
55 ± 18	1400	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta\pi^0\pi^0 n$	
24 ± 3		BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$	
20 ± 2		BARBERIS	97C OMEG	$450 pp \rightarrow ppK_S^0 K^\pm \pi^\mp$	
36 ± 5		² ANTINORI	95 OMEG	$300,450 pp \rightarrow pp2(\pi^+\pi^-)$	
29.0 ± 4.1		LEE	94 MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$	
25 ± 4	140	ARMSTRONG	89 OMEG	$300 pp \rightarrow K\bar{K}\pi pp$	
22 ± 2	4750	³ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$	
25 ± 4	504	BITYUKOV	88 SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$	
19 ± 5		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta\pi^+\pi^- n$	
32 ± 8	420	REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow KK\pi X$	
22 ± 2		CHUNG	85 SPEC	$8 \pi^- p \rightarrow NK\bar{K}\pi$	
32 ± 3	604	ARMSTRONG	84 OMEG	$85 \pi^+ p \rightarrow K\bar{K}\pi\pi p, pp \rightarrow K\bar{K}\pi pp$	
24 ± 3		CHAUVAT	84 SPEC	ISR 31.5 pp	
29 ± 10	103	DIONISI	80 HBC	$4 \pi^- p \rightarrow K\bar{K}\pi n$	
28.3 ± 6.7	320	NACASCH	78 HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.1 ± 3.4	234	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$	
32.4 ± 5.8		⁴ AAIJ	14Y LHCb	$\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+\pi^-)$	
18.2 ± 1.2		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$	
19.4 ± 1.5		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$	OCCUR=2
40 ± 5		ABATZIS	94 OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$	
31 ± 5		ARMSTRONG	89E OMEG	$300 pp \rightarrow pp2(\pi^+\pi^-)$	
41 ± 12		ARMSTRONG	89G OMEG	$85 \pi^+ p \rightarrow 4\pi\pi p, pp \rightarrow 4\pi pp$	
17.9 ± 10.9	60	RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$	
14 ⁺²⁰ ₋₁₄ ± 10	16	BECKER	87 MRK3	$e^+ e^- \rightarrow \phi K\bar{K}\pi$	
26 ± 12		EVANGELIS...	81 OMEG	$12 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$	
25 ± 15	200	GURTU	79 HBC	$4.2 K^- p \rightarrow n\eta 2\pi$	
~ 10		⁵ STANTON	79 CNTR	$8.5 \pi^- p \rightarrow n2\gamma 2\pi$	
24 ± 18	210	GRASSLER	77 HBC	$16 \pi^\mp p$	
28 ± 5	150	⁶ DEFOIX	72 HBC	$0.7 \bar{p}p \rightarrow 7\pi$	
46 ± 9	180	⁶ DUBOC	72 HBC	$1.2 \bar{p}p \rightarrow 2K4\pi$	
37 ± 5	500	⁷ THUN	72 MMS	$13.4 \pi^- p$	
10 ± 10		BOESEBECK	71 HBC	$16.0 \pi p \rightarrow p5\pi$	
30 ± 15		CAMPBELL	69 DBC	$2.7 \pi^+ d$	
60 ± 15		⁶ LORSTAD	69 HBC	$0.7 \bar{p}p, 4,5\text{-body}$	
35 ± 10		⁶ DAHL	67 HBC	$1.6\text{--}4.2 \pi^- p$	

1 The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

2 Supersedes ABATZIS 94, ARMSTRONG 89E.

3 From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

4 No systematic error given.

5 From phase shift analysis of $\eta\pi^+\pi^-$ system.

6 Resolution is not unfolded.

7 Seen in the missing mass spectrum.

NODE=M008W;LINKAGE=BL

NODE=M008W;LINKAGE=B

NODE=M008W;LINKAGE=A


NODE=M008W;LINKAGE=N1

NODE=M008W;LINKAGE=P

NODE=M008W;LINKAGE=R

NODE=M008W;LINKAGE=S

WEIGHTED AVERAGE
22.7±1.1 (Error scaled by 1.5)



$f_1(1285)$ width (MeV)

$f_1(1285)$ DECAY MODES

NODE=M008215;NODE=M008

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
$\Gamma_1 4\pi$	(32.7± 1.9) %	S=1.2	DESIG=21
$\Gamma_2 \pi^0 \pi^0 \pi^+ \pi^-$	(21.8± 1.3) %	S=1.2	DESIG=22
$\Gamma_3 2\pi^+ 2\pi^-$	(10.9± 0.6) %	S=1.2	DESIG=20
$\Gamma_4 \rho^0 \pi^+ \pi^-$	(10.9± 0.6) %	S=1.2	DESIG=191
$\Gamma_5 \rho^0 \rho^0$	seen		DESIG=23
$\Gamma_6 4\pi^0$	< 7 $\times 10^{-4}$	CL=90%	DESIG=7
$\Gamma_7 \eta \pi^+ \pi^-$	(35 ± 15) %		DESIG=198
$\Gamma_8 \eta \pi \pi$	(52.2± 2.0) %	S=1.2	DESIG=3
$\Gamma_9 a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$]	(38 ± 4) %		DESIG=4
$\Gamma_{10} \eta \pi \pi$ [excluding $a_0(980)\pi$]	(14 ± 4) %		DESIG=5
$\Gamma_{11} K\bar{K}\pi$	(9.0± 0.4) %	S=1.1	DESIG=1
$\Gamma_{12} K\bar{K}^*(892)$	not seen		DESIG=6
$\Gamma_{13} \pi^+ \pi^- \pi^0$	(3.0± 0.9) $\times 10^{-3}$		DESIG=197
$\Gamma_{14} \rho^\pm \pi^\mp$	< 3.1 $\times 10^{-3}$	CL=95%	DESIG=199
$\Gamma_{15} \gamma \rho^0$	(6.1± 1.0) %	S=1.7	DESIG=13
$\Gamma_{16} \phi \gamma$	(7.4± 2.6) $\times 10^{-4}$		DESIG=10
$\Gamma_{17} e^+ e^-$	< 9.4 $\times 10^{-9}$	CL=90%	DESIG=200
$\Gamma_{18} \gamma \gamma^*$			DESIG=9
$\Gamma_{19} \gamma \gamma$			DESIG=8

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 18 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 24.0$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	-30				
x_{10}	-12	-88			
x_{11}	22	-10	-4		
x_{15}	-25	-7	-3	-27	
	x_1	x_9	x_{10}	x_{11}	

$f_1(1285) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{19} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{19} / \Gamma$			
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.62	95	GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{18} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{18} / \Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.4 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.4.	
1.18 ± 0.25 ± 0.20	26	1,2 AIHARA	88B TPC	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$	
2.30 ± 0.61 ± 0.42		1,3 GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.8 ± 0.3 ± 0.3	420	4 ACHARD	02B L3	$183-209 e^+ e^- \rightarrow e^+ e^- \eta\pi^+\pi^-$	

1 Assuming a ρ -pole form factor.

2 Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.

3 Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49.

4 Published value multiplied by the $\eta\pi\pi$ branching ratio 0.52.

$f_1(1285)$ BRANCHING RATIOS

$\Gamma(K\bar{K}\pi) / \Gamma(4\pi)$		Γ_{11} / Γ_1			
VALUE	DOCUMENT ID	TECN	COMMENT		
0.274 ± 0.017 OUR FIT	Error includes scale factor of 1.4. [0.272 ± 0.016 OUR 2019 FIT Scale factor = 1.3]				
0.271 ± 0.016 OUR AVERAGE	Error includes scale factor of 1.2.				

0.265 ± 0.014	¹ BARBERIS 97C OMEG 450 $p p \rightarrow p p K_S^0 K^\pm \pi^\mp$
0.28 ± 0.05	² ARMSTRONG 89E OMEG 300 $p p \rightarrow p p f_1(1285)$
0.37 ± 0.03 ± 0.05	³ ARMSTRONG 89G OMEG 85 $\pi p \rightarrow 4\pi X$

1 Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.

2 Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.

3 4π consistent with being entirely $\rho\pi\pi$.

$\Gamma(\pi^0\pi^0\pi^+\pi^-) / \Gamma_{\text{total}}$		$\Gamma_2 / \Gamma = \frac{2}{3} \Gamma_1 / \Gamma$			
VALUE	DOCUMENT ID	TECN	COMMENT		
0.218 ± 0.013 OUR FIT	Error includes scale factor of 1.2. [0.223 ± 0.013 OUR 2019 FIT Scale factor = 1.3]				

$\Gamma(2\pi^+ 2\pi^-) / \Gamma_{\text{total}}$		$\Gamma_3 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$			
VALUE	DOCUMENT ID	TECN	COMMENT		
0.109 ± 0.006 OUR FIT	Error includes scale factor of 1.2. [0.112 ± 0.007 OUR 2019 FIT Scale factor = 1.3]				

$\Gamma(\rho^0\pi^+\pi^-) / \Gamma_{\text{total}}$		$\Gamma_4 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$			
VALUE	DOCUMENT ID	TECN	COMMENT		
0.109 ± 0.006 OUR FIT	Error includes scale factor of 1.2. [0.112 ± 0.007 OUR 2019 FIT Scale factor = 1.3]				

NODE=M008217

NODE=M008G2
NODE=M008G2

NODE=M008G3
NODE=M008G3

NODE=M008G3;LINKAGE=A
NODE=M008G3;LINKAGE=F
NODE=M008G3;LINKAGE=B
NODE=M008G3;LINKAGE=AC

NODE=M008220

NODE=M008R1
NODE=M008R1
NEW

NODE=M008R1;LINKAGE=B
NODE=M008R1;LINKAGE=M
NODE=M008R1;LINKAGE=A

NODE=M008R18
NODE=M008R18

NEW

NODE=M008R17
NODE=M008R17

NEW

NODE=M008R19
NODE=M008R19

NEW

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2\pi^+ 2\pi^-)$				Γ_4/Γ_3	NODE=M008R6 NODE=M008R6			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>					
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$								
1.0 ± 0.4	GRASSLER	77	HBC	$16 \text{ GeV } \pi^\pm p$				
$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$								
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>			Γ_5/Γ			
seen	BARBERIS	00C	$450 \text{ pp} \rightarrow p_f 4\pi p_s$					
$\Gamma(4\pi^0)/\Gamma_{\text{total}}$								
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ			
<7	90	ALDE	87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$			
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma(\eta \pi^+ \pi^-)$								
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ_7			
0.86 ± 0.16 ± 0.20	2.3k	¹ DOROFEEV	11	VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$			
1 Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+ \pi^-$ mass spectrum.								
$\Gamma(\eta \pi \pi)/\Gamma_{\text{total}}$								
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>			$\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$			
0.522 ± 0.020 OUR FIT	Error includes scale factor of 1.2. [0.520 ^{+0.018} _{-0.021} OUR 2019 FIT				NODE=M008R02 NODE=M008R02			
Scale factor = 1.2]								
$\Gamma(4\pi)/\Gamma(\eta \pi \pi)$								
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9 + \Gamma_{10})$	NODE=M008R22 NODE=M008R22			
0.63 ± 0.06 OUR FIT	Error includes scale factor of 1.3. [0.64 ^{+0.06} _{-0.05} OUR 2019 FIT				NEW			
Scale factor = 1.2]								
0.41 ± 0.14 OUR AVERAGE								
0.37 ± 0.11 ± 0.11	BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$				
0.64 ± 0.40	GURTU	79	HBC	$4.2 K^- p$				
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$								
0.93 ± 0.30	¹ GRASSLER	77	HBC	$16 \pi^\mp p$				
1 Assuming $\rho \pi \pi$ and $a_0(980) \pi$ intermediate states.								
$\Gamma(2\pi^+ 2\pi^-)/\Gamma(\eta \pi \pi)$								
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_8	NODE=M008R4 NODE=M008R4			
0.28 ± 0.02 ± 0.02	¹ LEES	12X	BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$	NEW			
1 Assuming $B(f_1(1285) \rightarrow \pi \pi \eta) = 3/2 B(f_1(1285) \rightarrow \pi^+ \pi^- \eta)$.								
$\Gamma(a_0(980)\pi [\text{ignoring } a_0(980) \rightarrow K\bar{K}])/ \Gamma(\eta \pi \pi)$								
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_9/\Gamma_8 = \Gamma_9/(\Gamma_9 + \Gamma_{10})$			
0.72 ± 0.08 OUR FIT					NODE=M008R04;LINKAGE=M			
0.72 ± 0.07 OUR AVERAGE					NODE=M008R04;LINKAGE=LE			
0.74 ± 0.02 ± 0.09	DICKSON	16	CLAS	$\gamma p \rightarrow f_1(1285) p$				
0.72 ± 0.15	GURTU	79	HBC	$4.2 K^- p$				
0.6 ^{+0.3} _{-0.2}	CORDEN	78	OMEG	12–15 $\pi^- p$				
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$								
>0.69	ACHARD	02B	L3	$183\text{--}209 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$				
0.28 ± 0.07	ALDE	97B	GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$				
1.0 ± 0.3	GRASSLER	77	HBC	$16 \pi^\mp p$				
$\Gamma(K\bar{K}\pi)/\Gamma(\eta \pi \pi)$								
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9 + \Gamma_{10})$	NODE=M008R2 NODE=M008R2			
0.172 ± 0.012 OUR FIT	Error includes scale factor of 1.1. [0.176 ± 0.012 OUR 2019 FIT				NEW			
Scale factor = 1.1]								
0.176 ± 0.012 OUR AVERAGE								
0.216 ± 0.010 ± 0.031	DICKSON	16	CLAS	$\gamma p \rightarrow f_1(1285) p$	OCCUR=2			
0.166 ± 0.01 ± 0.008	BARBERIS	98C	OMEG	$450 \text{ pp} \rightarrow p_f f_1(1285) p_s$				
0.42 ± 0.15	GURTU	79	HBC	$4.2 K^- p$				
0.5 ± 0.2	¹ CORDEN	78	OMEG	12–15 $\pi^- p$				
0.20 ± 0.08	² DEFIX	72	HBC	$0.7 \bar{p} p \rightarrow 7\pi$				
0.16 ± 0.08	CAMPBELL	69	DBC	$2.7 \pi^+ d$				
1 CORDEN 78 assumes low-mass $\eta \pi \pi$ region is dominantly 1^{++} . See BARBERIS 98C and MANAK 00A for discussion.								
2 $K\bar{K}$ system characterized by the $I = 1$ threshold enhancement. (See under $a_0(980)$).								

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
not seen	NACASCH 78	HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				

seen 1 ACHARD 07 L3 $183-209 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm\pi^\mp$

1 A clear signal of 19.8 ± 4.4 events observed at high Q^2 .

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
0.30±0.055±0.074	2.3k	1 DOROFEEV	11 VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$	

1 Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.

 $\Gamma(\rho^\pm\pi^\mp)/\Gamma_{\text{total}}$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
<0.31	95	DOROFEEV	11 VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$	

 $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
6.1±1.0 OUR FIT	Error includes scale factor of 1.7. [(5.3 ± 1.2) $\times 10^{-2}$ OUR 2019 FIT Scale factor = 2.9]				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.8 \pm 0.7 \pm 0.6$	1 AMELIN	95 VES	$37 \pi^- N \rightarrow \pi^-\pi^+\pi^-\gamma N$	
<5	95	BITYUKOV	91B SPEC	$32 \pi^- p \rightarrow \pi^+\pi^-\gamma n$

1 Not an independent measurement.

 $\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$
0.55±0.10 OUR FIT	Error includes scale factor of 1.5. [0.48 ± 0.13 OUR 2019 FIT Scale factor = 2.5]			

0.45±0.18

1 COFFMAN 90 MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

1 Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.

 $\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$


VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma_{15} = (\Gamma_9 + \Gamma_{10})/\Gamma_{15}$
8.6±1.6 OUR FIT	Error includes scale factor of 1.9. [9.7 ± 1.9 OUR 2019 FIT Scale factor = 2.4]			

8.5±2.0 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

21.3±4.4	DICKSON	16 CLAS	$\gamma p \rightarrow f_1(1285)p$	
10.0±1.0±2.0	BARBERIS	98C OMEG	$450 pp \rightarrow p_f f_1(1285)p_s$	
7.5±1.0	1 ARMSTRONG	92C OMEG	$300 pp \rightarrow pp\pi^+\pi^-\gamma, pp\eta\pi^+\pi^-$	

1 Published value multiplied by 1.5.

WEIGHTED AVERAGE
8.5±2.0 (Error scaled by 2.2)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

NODE=M008R5
NODE=M008R5

NODE=M008R5;LINKAGE=CH

NODE=M008R01
NODE=M008R01

NODE=M008R01;LINKAGE=DO

NODE=M008R03
NODE=M008R03

NODE=M008R15
NODE=M008R15

NEW

NODE=M008R15;LINKAGE=A

NODE=M008R13
NODE=M008R13
NEW

NODE=M008R13;LINKAGE=E

NODE=M008R16
NODE=M008R16
NEW

NODE=M008R16;LINKAGE=B

$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ_{11}
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.035	90	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	NODE=M008R12 NODE=M008R12
1 Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$.					

 $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ_{11}
0.82±0.21±0.20		19	BITYUKOV	88	SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.50	95		BARBERIS	98C	OMEG	$450 pp \rightarrow p_f f_1(1285) p_s$
<0.93	95		AMELIN	95	VES	$37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
<9.4 × 10⁻⁹	90	¹ ACHASOV	20	SND	$e^+ e^- \rightarrow \eta\pi^0\pi^0$

1 Two candidate events are found corresponding to a significance of 2.5σ and the branching fraction of $(5.1^{+3.7}_{-2.7}) \times 10^{-9}$.

f₁(1285) REFERENCES

ACHASOV	20	PL B800 135074	ACHASOV 2020 <i>et al.</i>	(SND)	
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60144;ERROR=8
DICKSON	16	PR C93 065202	R. Dickson <i>et al.</i>	(JLab CLAS Collab.)	REFID=60024
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57487
AAIJ	14Y	PRL 112 091802	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56781
LEES	12X	PR D86 092010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55837
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54714
DOROFEEV	11	EPJ A47 68	V. Dorofeev <i>et al.</i>	(SERP, MIPT)	REFID=53931
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)	REFID=16755
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=53229
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51698
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=52049
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=50167
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49548
ACCIARRI	01G	PL B501 1	M. Acciari <i>et al.</i>	(L3 Collab.)	REFID=48574
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=48319
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=49649
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47959
SOSA	99	PRC 83 913	M. Sosa <i>et al.</i>	(WA 102 Collab.)	REFID=47989
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(GAMS Collab.)	REFID=46937
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46346
		Translated from YAF 60 458.			REFID=45396
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
AMELIN	95	ZPHY C66 71	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=44376
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44437
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44090
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=44092
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=43587
ARMSTRONG	92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(Mark III Collab.)	REFID=42097
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BITYUKOV	91B	SNJNP 54, 318	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=41864
		Translated from YAF 54 529.			
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
ARMSTRONG	89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+, JPC)	REFID=40729
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
ARMSTRONG	89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)	REFID=40930
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNLU, CUNY+)	REFID=40924
AIHARA	88B	PL B209 107	H. Aibara <i>et al.</i>	(TPC-2γ Collab.)	REFID=40572
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
BITYUKOV	88	PL B203 327	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=40569
MIR	88	Photon-Photon 88, 126	R. Mir	(Mark III Collab.)	REFID=41574
Conference					
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
BECKER	87	PRL 59 186	J.Y. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
GIDAL	87	PR 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) JP	REFID=20891
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP	REFID=20936
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP	REFID=20934
ARMSTRONG	84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=20929
BITYUKOV	84B	PL 144B 133	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=20468
CHAUVAT	84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCL+)	REFID=20932
TORNQVIST	82B	PL B203 268	N.A. Tornqvist	(HELS)	REFID=20573
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)	REFID=20922
DIONISI	80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+)	REFID=20924
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP	REFID=20887
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHET, TELA+) JP	REFID=20452
NACASCH	78	NP B135 203	R. Nacash <i>et al.</i>	(PARIS, MADR, CERN)	REFID=20919
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
DUBOC	72	NP B46 429	J. Duboc <i>et al.</i>	(PARIS, LIVP)	REFID=20339
THUN	72	PRL 28 1733	R. Thun <i>et al.</i>	(STON, NEAS)	REFID=20911
BARDADIN...	71	PR D4 2711	M. Bardadin-Otwinowska <i>et al.</i>	(WARS)	REFID=20196
BOESEBECK	71	PL 34B 659	K. Boesebeck (AACH, BERL, BONN, CERN, CRAC+)	(PURD)	REFID=20905
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>		REFID=20419
LORSTAD	69	NP B14 63	B. Lorstad <i>et al.</i>	(CDEF, CERN) JP	REFID=20901
D'ANDLAU	68	NP B5 693	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+) JP	REFID=20897
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP	REFID=20321

$\eta(1295)$ $I^G(J^{PC}) = 0^+(0^-+)$ See also the mini-review under $\eta(1405)$

NODE=M037

NODE=M037

NODE=M037M


NODE=M037M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1294±4 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.				
1302±9±8	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1282±5	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1299±4	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1295±4		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1264±8	¹ AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~1275	STANTON	79	CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$

WEIGHTED AVERAGE
1294±4 (Error scaled by 1.6)

 $\eta(1295)$ mass (MeV)

¹ PWA analysis of AUGUSTIN 92 assigns 0^-+ quantum numbers to this state rather than 1^{++} as before.

NODE=M037M;LINKAGE=AG

 $\eta(1295)$ WIDTH

NODE=M037W

NODE=M037W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
55± 5 OUR AVERAGE				
57±23±21	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
66±13	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
53± 6		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<40	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
44±20		² AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~70		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$

² PWA analysis of AUGUSTIN 92 assigns 0^-+ quantum numbers to this state rather than 1^{++} as before.

NODE=M037W;LINKAGE=AG

$\eta(1295)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta\pi^+\pi^-$	seen
$\Gamma_2 a_0(980)\pi$	seen
$\Gamma_3 \gamma\gamma$	
$\Gamma_4 \eta\pi^0\pi^0$	seen
$\Gamma_5 \eta(\pi\pi)S\text{-wave}$	seen
$\Gamma_6 \sigma\eta$	
$\Gamma_7 K\bar{K}\pi$	

$\eta(1295) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_3/\Gamma$
<u>VALUE (keV)</u>	<u>CL%</u>
<0.066	95
AIHARA	ACCIARRI
88C	01G
$e^+e^- \rightarrow e^+e^- \eta\pi^+\pi^-$	L3
ANTREASYAN 87	CBAL
$e^+e^- \rightarrow e^+e^-\eta\pi\pi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_7\Gamma_3/\Gamma$
<u>VALUE (keV)</u>	<u>CL%</u>
<0.014	90
AHOHE	3,4
90	AHOHE
$e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$	05
CLE2	

3 Using $\eta(1295)$ mass and width 1294 MeV and 55 MeV, respectively.

4 Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

$\eta(1295)$ BRANCHING RATIOS

$\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
not seen	BERTIN
	97
	OBLX
	$0.0 \bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
seen	BIRMAN
	88
	MPS
	$8 \pi^- p \rightarrow K^+\bar{K}^0\pi^-\pi^-n$
large	ANDO
large	STANTON
	86
	SPEC
	$8 \pi^- p \rightarrow \eta\pi^+\pi^-n$
	79
	CNTR
	$8.4 \pi^- p \rightarrow n\eta\pi^-2\pi$

$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi^0\pi^0)$	Γ_2/Γ_4
<u>VALUE</u>	<u>DOCUMENT ID</u>
0.65±0.10	5 ALDE
	97B
	GAM4
	$100 \pi^- p \rightarrow \eta\pi^0\pi^0n$

5 Assuming that $a_0(980)$ decays only to $\eta\pi$.

$\Gamma(\eta\pi\pi)S\text{-wave})/\Gamma(\eta\pi^0\pi^0)$	Γ_5/Γ_4
<u>VALUE</u>	<u>DOCUMENT ID</u>
0.35±0.10	ALDE
	97B
	GAM4
	$100 \pi^- p \rightarrow \eta\pi^0\pi^0n$

$\Gamma(a_0(980)\pi)/\Gamma(\sigma\eta)$	Γ_2/Γ_6
<u>VALUE</u>	<u>EVTS</u>
0.48±0.22	9082
	MANAK
	00A
	MPS
	$18 \pi^- p \rightarrow \eta\pi^+\pi^-n$

$\eta(1295)$ REFERENCES

AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 60 458.		
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+) (SUGI, NAGO, KEK, KYOT+)
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIIRS, SAGA+) IJP
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP

NODE=M037215;NODE=M037

DESIG=2;OUR EST; \rightarrow UNCHECKED ←
 DESIG=1;OUR EST; \rightarrow UNCHECKED ←
 DESIG=3
 DESIG=4;OUR EST; \rightarrow UNCHECKED ←
 DESIG=5;OUR EST; \rightarrow UNCHECKED ←
 DESIG=6
 DESIG=7

NODE=M037220

NODE=M037G2
NODE=M037G2

NODE=M037G3
NODE=M037G3

NODE=M037G3;LINKAGE=AH
NODE=M037G3;LINKAGE=B3

NODE=M037225

NODE=M037R1
NODE=M037R1

NODE=M037R2
NODE=M037R2

NODE=M037R2;LINKAGE=A

NODE=M037R4
NODE=M037R4

NODE=M037R5
NODE=M037R5

NODE=M037

REFID=50764
REFID=48319
REFID=49649
REFID=47989
REFID=45396
REFID=45417
REFID=41584
REFID=41748
REFID=41352
REFID=40564
REFID=40568
REFID=40008
REFID=20891
REFID=20887

$\pi(1300)$ $I^G(J^{PC}) = 1^-(0^-+)$ **$\pi(1300)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1300±100 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1128± 26±70		DARGENT 17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1345± 8±10	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
1200± 40	90k	SALVINI 04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1343± 15±24		CHUNG 02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1375± 40		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1275± 15		BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
~ 1114		ABELE 96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
1190± 30		ZIELINSKI 84	SPEC	$200 \pi^+ Z \rightarrow Z 3\pi$
1240± 30		BELLINI 82	SPEC	$40 \pi^- A \rightarrow A 3\pi$
1273± 50		² AARON 81	RVUE	
1342± 20		BONESINI 81	OMEG	$12 \pi^- p \rightarrow p 3\pi$
~ 1400		DAUM 81B	SPEC	$63.94 \pi^- p$

¹ From analysis of L3 data at 183–209 GeV.² Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058M

NODE=M058M
→ UNCHECKED ← **$\pi(1300)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
200 to 600 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
314± 39±66		DARGENT 17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
260± 20±30	18k	³ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
470±120	90k	SALVINI 04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
449± 39±47		CHUNG 02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
268± 50		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
218±100		BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
~ 340		ABELE 96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
440± 80		ZIELINSKI 84	SPEC	$200 \pi^+ Z \rightarrow Z 3\pi$
360±120		BELLINI 82	SPEC	$40 \pi^- A \rightarrow A 3\pi$
580±100		⁴ AARON 81	RVUE	
220± 70		BONESINI 81	OMEG	$12 \pi^- p \rightarrow p 3\pi$
~ 600		DAUM 81B	SPEC	$63.94 \pi^- p$

³ From analysis of L3 data at 183–209 GeV.⁴ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.NODE=M058M;LINKAGE=SC
NODE=M058M;LINKAGE=E

NODE=M058W

NODE=M058W
→ UNCHECKED ← **$\pi(1300)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi$	seen
$\Gamma_2 \pi(\pi\pi)_{S\text{-wave}}$	seen
$\Gamma_3 \gamma\gamma$	

NODE=M058W;LINKAGE=SC
NODE=M058W;LINKAGE=E

NODE=M058215;NODE=M058

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4

NODE=M058218

NODE=M058G1
NODE=M058G1

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1 \Gamma_3 / \Gamma$
<0.085	90	ACCIARRI 97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.8	95	⁵ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	
<0.54	90	ALBRECHT 97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	

⁵ From analysis of L3 data at 183–209 GeV.

NODE=M058G1;LINKAGE=SC

$\pi(1300)$ BRANCHING RATIOS

$\Gamma(\pi(\pi\pi)S\text{-wave})/\Gamma(\rho\pi)$		Γ_2/Γ_1			
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.2 ± 0.4		90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
seen			CHUNG	02	B852 18.3 $\pi^- p \rightarrow \pi^+ 2\pi^- p$
<0.15		90	ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
2.12		6	AARON	81	RVUE

⁶ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058220

NODE=M058R1
NODE=M058R1

NODE=M058R1;LINKAGE=E

$\pi(1300)$ REFERENCES

DARGENT	17	JHEP 1705 143	P. d'Argent <i>et al.</i>	(HEID, BRIS)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACCIARRI	97T	PL B413 147	M. Acciari <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERTIN	97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ZIELINSKI	84	PR D30 1855	M. Zielinski <i>et al.</i>	(ROCH, MINN, FNAL)
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)
AARON	81	PR D24 1207	R.A. Aaron, R.S. Longacre	(NEAS, BNL)
BONESINI	81	PL 103B 75	M. Bonesini <i>et al.</i>	(MILA, LIVP, DARE+)
DANKOWY...	81	PR D46 580	J.A. Dankowycz <i>et al.</i>	(TNTO, BNL, CARL+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
DAUM	80	PL 89B 281	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

NODE=M058

REFID=58121
REFID=51186
REFID=53226
REFID=48837
REFID=48334
REFID=45761
REFID=45418
REFID=45763
REFID=45011
REFID=20881
REFID=21134
REFID=20870
REFID=21130
REFID=20572
REFID=20872
REFID=20868
REFID=20571

NODE=M012

 $a_2(1320)$


$$I^G(J^{PC}) = 1^-(2^{++})$$

$a_2(1320)$ MASS

VALUE (MeV)	DOCUMENT ID
1316.9 ± 0.9 OUR AVERAGE	Includes data from the 4 datablocks that follow this one. Error includes scale factor of 1.9. See the ideogram below.

NODE=M012M0

NODE=M012M0



$a_2(1320)$ MASS (MeV)

3 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

NODE=M012M1

NODE=M012M1

1318.6± 1.3 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

1314.5 ± 4.0	46M	1 AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1326 ± 2 ± 2		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1317 ± 3		BARBERIS	98B		450 $p p \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
1323 ± 4 ± 3		ACCIARRI	97T	L3	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^- \pi^0$
1320 ± 7		ALBRECHT	97B	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^- \pi^0$
1311.3 ± 1.6 ± 3.0	72.4k	AMELIN	96	VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1310 ± 5		ARMSTRONG	90	OMEG 0	300.0 $p p \rightarrow p p \pi^+ \pi^- \pi^0$
1323.8 ± 2.3	4022	AUGUSTIN	89	DM2 ±	$J/\psi \rightarrow \rho^\pm a_2^\mp$
1320.6 ± 3.1	3562	AUGUSTIN	89	DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$
1317 ± 2	25k	2 DAUM	80C	SPEC -	63,94 $\pi^- p \rightarrow 3\pi p$
1320 ± 10	1097	2 BALTAY	78B	HBC +0	15 $\pi^+ p \rightarrow p 4\pi$
1306 ± 8		FERRERSORIA	78	OMEG -	9 $\pi^- p \rightarrow p 3\pi$
1318 ± 7	1.6k	2 EMMS	75	DBC 0	4 $\pi^+ n \rightarrow p (3\pi)^0$
1315 ± 5		2 ANTIPOV	73C	CNTR -	25,40 $\pi^- p \rightarrow p \eta \pi^-$
1306 ± 9	1580	CHALOUPKA	73	HBC -	3.9 $\pi^- p$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1321	± 1	$^{+0}_{-7}$	420k	³ ALEKSEEV	10	COMP	190	$\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1300	± 2	± 4	18k	⁴ SCHEGELSKY	06	RVUE	0	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
1305	± 14			CONDOR	93	SHF		$\gamma p \rightarrow n \pi^+ \pi^+ \pi^-$
1310	± 2			² EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow 3\pi^- p$
1343	± 11		490	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow \Delta 3\pi$
1309	± 5		5k	BINNIE	71	MMS	-	$\pi^- p$ near a_2 threshold
1299	± 6		28k	BOWEN	71	MMS	-	5 $\pi^- p$
1300	± 6		24k	BOWEN	71	MMS	+	5 $\pi^+ p$
1309	± 4		17k	BOWEN	71	MMS	-	7 $\pi^- p$
1306	± 4		941	ALSTON...	70	HBC	+	7.0 $\pi^+ p \rightarrow 3\pi^- p$

1 Statistical error negligible.

2 From a fit to $J^P = 2^+$ $\rho\pi$ partial wave.

3 Superseded by AGHASYAN 2018B.

4 From analysis of L3 data at 183–209 GeV.

OCCUR=2

OCCUR=2


OCCUR=3

NODE=M012M1;LINKAGE=D

NODE=M012M1;LINKAGE=P

NODE=M012M1;LINKAGE=C

NODE=M012M1;LINKAGE=SC



KK MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M012M2

NODE=M012M2

The data in this block is included in the average printed for a previous datablock.

1318.1± 0.7 OUR AVERAGE

1319	± 5	4700	^{1,2} CLELAND	82B	SPEC	+	50 $\pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
1324	± 6	5200	^{1,2} CLELAND	82B	SPEC	-	50 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
1320	± 2	4000	CHABAUD	80	SPEC	-	17 $\pi^- A \rightarrow K_S^0 K^- A$	
1312	± 4	11000	CHABAUD	78	SPEC	-	9.8 $\pi^- p \rightarrow K^- K_S^0 p$	
1316	± 2	4730	CHABAUD	78	SPEC	-	18.8 $\pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
1318	± 1		^{1,3} MARTIN	78D	SPEC	-	10 $\pi^- p \rightarrow K_S^0 K^- p$	
1320	± 2	2724	MARGULIE	76	SPEC	-	23 $\pi^- p \rightarrow K^- K_S^0 p$	
1313	± 4	730	FOLEY	72	CNTR	-	20.3 $\pi^- p \rightarrow K^- K_S^0 p$	
1319	± 3	1500	³ GRAYER	71	ASPK	-	17.2 $\pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1304	± 10	870	⁴ SCHEGELSKY	06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$
1330	± 11	1000	^{1,2} CLELAND	82B	SPEC	+	30 $\pi^+ p \rightarrow K_S^0 K^+ p$
1324	± 5	350	HYAMS	78	ASPK	+	12.7 $\pi^+ p \rightarrow K^+ K_S^0 p$

NODE=M012M2;LINKAGE=P

NODE=M012M2;LINKAGE=W

NODE=M012M2;LINKAGE=S

NODE=M012M2;LINKAGE=SC

1 From a fit to $J^P = 2^+$ partial wave.

2 Number of events evaluated by us.

3 Systematic error in mass scale subtracted.

4 From analysis of L3 data at 91 and 183–209 GeV.

$\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

1312.2± 2.8 OUR AVERAGE Error includes scale factor of 2.6. See the ideogram below.

1306.0± 0.8±1.3	1 RODAS	19 JPAC	191 $\pi^- p \rightarrow \eta(\prime)\pi^- p$	
1308 ± 9	BARBERIS	00H	450 $p p \rightarrow p_f \eta \pi^0 p_s$	
1316 ± 9	BARBERIS	00H	450 $p p \rightarrow \Delta_f^{++} \eta \pi^- p_s$	OCCUR=2
1317 ± 1 ±2	THOMPSON	97 MPS	18 $\pi^- p \rightarrow \eta \pi^- p$	
1315 ± 5 ±2	AMSLER	94D CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$	
1325.1± 5.1	AOYAGI	93 BKEI	$\pi^- p \rightarrow \eta \pi^- p$	
1317.7± 1.4±2.0	BELADIDZE	93 VES	37 $\pi^- N \rightarrow \eta \pi^- N$	
1323 ± 8 1000	3 KEY	73 OSPK -	6 $\pi^- p \rightarrow p \pi^- \eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1307 ± 1 ±6	4 JACKURA	18 JPAC	$\pi^- p \rightarrow \eta \pi^- p$	
1315 ±12	5 ADOLPH	15 COMP	191 $\pi^- p \rightarrow \eta(\prime)\pi^- p$	
1309 ± 4	ANISOVICH	09 RVUE	$\bar{p} p, \pi N$	
1324 ± 5	ARMSTRONG	93C E760 0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
1336.2± 1.7 2561	DELFOSSÉ	81 SPEC +	$\pi^\pm p \rightarrow p \pi^\pm \eta$	
1330.7± 2.4 1653	DELFOSSÉ	81 SPEC -	$\pi^\pm p \rightarrow p \pi^\pm \eta$	OCCUR=2
1324 ± 8 6200	3,6 CONFORTO	73 OSPK -	6 $\pi^- p \rightarrow p MM^-$	

1 The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data.
The mass is extracted from the T-matrix pole.

2 The systematic error of 2 MeV corresponds to the spread of solutions.

3 Error includes 5 MeV systematic mass-scale error.


4 Superseded by RODAS 19.

5 ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the $\eta\pi$ and $\rho\pi$ channels into account.

6 Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.

NODE=M012M3

NODE=M012M3



NODE=M012M3;LINKAGE=C

NODE=M012M3;LINKAGE=DD

NODE=M012M3;LINKAGE=E

NODE=M012M3;LINKAGE=B

NODE=M012M3;LINKAGE=A

NODE=M012M3;LINKAGE=M

 $\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

1322 ± 7 OUR AVERAGE

1318 ± 8 ±3	IVANOV	01 B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
1327.0±10.7	BELADIDZE	93 VES	37 $\pi^- N \rightarrow \eta' \pi^- N$

NODE=M012M4

NODE=M012M4

$a_2(1320)$ WIDTH

NODE=M012210

 3π MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

CHG

COMMENT

NODE=M012W1

NODE=M012W1

 105.0 ± 1.7 OUR AVERAGE

106.6 \pm 3.4 7.0	46M	1 AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	I
108 \pm 3 \pm 15		CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
120 \pm 10		BARBERIS	98B	450 $p p \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$	
105 \pm 10 \pm 11		ACCIARRI	97T L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
120 \pm 10		ALBRECHT	97B ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
103.0 \pm 6.0 \pm 3.3 72.4k		AMELIN	96 VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
120 \pm 10		ARMSTRONG	90 OMEG 0	300.0 $p p \rightarrow p p \pi^+ \pi^- \pi^0$	
107.0 \pm 9.7	4022	AUGUSTIN	89 DM2 \pm	$J/\psi \rightarrow \rho^\pm a_2^\mp$	
118.5 \pm 12.5	3562	AUGUSTIN	89 DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$	OCCUR=2
97 \pm 5		2 EVANGELIS...	81 OMEG -	12 $\pi^- p \rightarrow 3\pi p$	
96 \pm 9	25k	2 DAUM	80C SPEC -	63,94 $\pi^- p \rightarrow 3\pi p$	
110 \pm 15	1097	2 BALTAY	78B HBC +0	15 $\pi^+ p \rightarrow p 4\pi$	
112 \pm 18	1.6k	2 EMMS	75 DBC 0	4 $\pi^+ n \rightarrow p (3\pi)^0$	
122 \pm 14	1.2k	2,3 WAGNER	75 HBC 0	7 $\pi^+ p \rightarrow \Delta^{++} (3\pi)^0$	
115 \pm 15		2 ANTIPOV	73C CNTR -	25,40 $\pi^- p \rightarrow p \eta \pi^-$	
99 \pm 15	1580	CHALOUPKA	73 HBC -	3.9 $\pi^- p$	
105 \pm 5	28k	BOWEN	71 MMS -	5 $\pi^- p$	OCCUR=2
99 \pm 5	24k	BOWEN	71 MMS +	5 $\pi^+ p$	OCCUR=3
103 \pm 5	17k	BOWEN	71 MMS -	7 $\pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

110 \pm 2 \pm 2 -15	420k	4 ALEKSEEV	10 COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	I
117 \pm 6 \pm 20	18k	5 SCHEGELSKY	06 RVUE 0	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	
120 \pm 40		CONDО	93 SHF	$\gamma p \rightarrow n \pi^+ \pi^+ \pi^-$	
115 \pm 14	490	BALTAY	78B HBC 0	15 $\pi^+ p \rightarrow \Delta 3\pi$	OCCUR=2
72 \pm 16	5k	BINNIE	71 MMS -	$\pi^- p$ near a_2 thresh- old	OCCUR=2
79 \pm 12	941	ALSTON-...	70 HBC +	7.0 $\pi^+ p \rightarrow 3\pi p$	

1 Statistical error negligible.

2 From a fit to $J^P = 2^+ \rho \pi$ partial wave.3 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

4 Superseded by AGHASYAN 2018B.

5 From analysis of L3 data at 183–209 GeV.

 $K\bar{K}$ AND $\eta\pi$ MODES

VALUE (MeV)

DOCUMENT ID

107 \pm 5 OUR ESTIMATE**112.5 \pm 1.2 OUR AVERAGE** Includes data from the 2 datablocks that follow this one. **$K\bar{K}$ MODE**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

CHG

COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M012W1;LINKAGE=C

NODE=M012W1;LINKAGE=P

NODE=M012W1;LINKAGE=S

NODE=M012W1;LINKAGE=E

NODE=M012W1;LINKAGE=SC

NODE=M012W0

NODE=M012W0

→ UNCHECKED ←

NODE=M012W2

NODE=M012W2

109.8 \pm 2.4 OUR AVERAGE

112 \pm 20	4700	1,2 CLELAND	82B SPEC +	50 $\pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
120 \pm 25	5200	1,2 CLELAND	82B SPEC -	50 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
106 \pm 4	4000	CHABAUD	80 SPEC -	17 $\pi^- A \rightarrow K_S^0 K^- A$	
126 \pm 11	11000	CHABAUD	78 SPEC -	9.8 $\pi^- p \rightarrow K^- K_S^0 p$	
101 \pm 8	4730	CHABAUD	78 SPEC -	18.8 $\pi^- p \rightarrow K^- K_S^0 p$	
113 \pm 4		1,3 MARTIN	78D SPEC -	10 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=2
105 \pm 8	2724	3 MARGULIE	76 SPEC -	23 $\pi^- p \rightarrow K^- K_S^0 p$	
113 \pm 19	730	FOLEY	72 CNTR -	20.3 $\pi^- p \rightarrow K^- K_S^0 p$	
123 \pm 13	1500	3 GRAYER	71 ASPK -	17.2 $\pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

120	± 15	870	⁴ SCHEGELSKY	06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$
121	± 51	1000	^{1,2} CLELAND	82B	SPEC	+	$30\pi^+ p \rightarrow K_S^0 K^+ p$
110	± 18	350	HYAMS	78	ASPK	+	$12.7\pi^+ p \rightarrow K^+ K_S^0 p$

1 From a fit to $J^P = 2^+$ partial wave.

2 Number of events evaluated by us.

3 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

4 From analysis of L3 data at 91 and 183–209 GeV.

NODE=M012W2;LINKAGE=P
NODE=M012W2;LINKAGE=W
NODE=M012W2;LINKAGE=S
NODE=M012W2;LINKAGE=SC

$\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

113.4 \pm 1.3 OUR AVERAGE

114.4 \pm 1.6 \pm 0.0		¹ RODAS	19	JPAC	$191\pi^- p \rightarrow \eta(\prime)\pi^- p$		
115	± 20	BARBERIS	00H		$450pp \rightarrow p_f\eta\pi^0 p_s$		
112	± 14	BARBERIS	00H		$450pp \rightarrow \Delta_f^{++}\eta\pi^- p_s$	OCCUR=2	
112	± 3 ± 2	² AMSLER	94D	CBAR	$0.0\bar{p}p \rightarrow \pi^0\pi^0\eta$		
103	± 6 ± 3	BELADIDZE	93	VES	$37\pi^- N \rightarrow \eta\pi^- N$		
112.2 \pm 5.7	2561	DELFOSSE	81	SPEC	$+\pi^\pm p \rightarrow p\pi^\pm\eta$		
116.6 \pm 7.7	1653	DELFOSSE	81	SPEC	$-\pi^\pm p \rightarrow p\pi^\pm\eta$		
108	± 9	KEY	73	OSPK	$-6\pi^- p \rightarrow p\pi^- \eta$		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
112	± 1 ± 8	³ JACKURA	18	JPAC	$\pi^- p \rightarrow \eta\pi^- p$		
119	± 14	⁴ ADOLPH	15	COMP	$191\pi^- p \rightarrow \eta(\prime)\pi^- p$		
110	± 4	ANISOVICH	09	RVUE	$\bar{p}p, \pi N$		
127	± 2 ± 2	⁵ THOMPSON	97	MPS	$18\pi^- p \rightarrow \eta\pi^- p$		
118	± 10	ARMSTRONG	93c	E760	$0\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$		
104	± 9	6200	⁶ CONFORTO	73	OSPK	$-6\pi^- p \rightarrow pMM^-$	

1 The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data. The width is extracted from the T-matrix pole.

2 The systematic error of 2 MeV corresponds to the spread of solutions.

3 Superseded by RODAS 19.

4 ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the $\eta\pi$ and $\rho\pi$ channels into account.

5 Resolution is not unfolded.

6 Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.

NODE=M012W3;LINKAGE=E
NODE=M012W3;LINKAGE=DD
NODE=M012W3;LINKAGE=C
NODE=M012W3;LINKAGE=B
NODE=M012W3;LINKAGE=A
NODE=M012W3;LINKAGE=M

$\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
119\pm 25 OUR AVERAGE			

140 \pm 35 \pm 20	IVANOV	01	B852	$18\pi^- p \rightarrow \eta'\pi^- p$
106 \pm 32	BELADIDZE	93	VES	$37\pi^- N \rightarrow \eta'\pi^- N$

NODE=M012W4
NODE=M012W4

$a_2(1320)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 3π	(70.1 \pm 2.7) %	S=1.2
Γ_2 $\rho(770)\pi$		
Γ_3 $f_2(1270)\pi$		
Γ_4 $\rho(1450)\pi$		
Γ_5 $\eta\pi$	(14.5 \pm 1.2) %	
Γ_6 $\omega\pi\pi$	(10.6 \pm 3.2) %	S=1.3
Γ_7 $K\bar{K}$	(4.9 \pm 0.8) %	
Γ_8 $\eta'(958)\pi$	(5.5 \pm 0.9) $\times 10^{-3}$	
Γ_9 $\pi^\pm\gamma$	(2.91 \pm 0.27) $\times 10^{-3}$	
Γ_{10} $\gamma\gamma$	(9.4 \pm 0.7) $\times 10^{-6}$	
Γ_{11} e^+e^-	< 5 $\times 10^{-9}$	CL=90%

NODE=M012215;NODE=M012

DESIG=1
DESIG=11
DESIG=12
DESIG=13
DESIG=3
DESIG=4
DESIG=2
DESIG=8
DESIG=7
DESIG=9
DESIG=10

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 18 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 9.3$ for 15 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	10				
x_6	-89	-46			
x_7	-1	-2	-24		
	x_1	x_5	x_6		

$a_2(1320)$ PARTIAL WIDTHS

$\Gamma(\eta\pi)$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.5 ± 3.0 870 ¹ SCHEGELSKY 06A RVUE 0 $\gamma\gamma \rightarrow K_S^0 K_S^0$

1 From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$ keV and SU(3) relations.

Γ_5

NODE=M012220

NODE=M012W6
NODE=M012W6

$\Gamma(K\bar{K})$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.0^{+2.0}_{-1.5}$ 870 ¹ SCHEGELSKY 06A RVUE 0 $\gamma\gamma \rightarrow K_S^0 K_S^0$

1 From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$ keV and SU(3) relations.

Γ_7

NODE=M012W5
NODE=M012W5

$\Gamma(\pi^\pm\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

311± 25 OUR AVERAGE

$358 \pm 6 \pm 42$ 1 ADOLPH 14 COMP – $190 \pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$

$284 \pm 25 \pm 25$ 7.1k MOLCHANOV 01 SELX $600 \pi^- \text{A} \rightarrow \pi^+ \pi^- \pi^- \text{A}$

295 ± 60 CIHANGIR 82 SPEC + $200 \pi^+ \text{A}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

461 ± 110 ² MAY 77 SPEC ± 9.7 γA

1 Primakoff reaction using $a_2(1320) \rightarrow 3\pi$ branching ratio of 70.1%.

2 Assuming one-pion exchange.

Γ_9

NODE=M012W5;LINKAGE=SC

$\Gamma(\gamma\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

1.00±0.06 OUR AVERAGE

$0.98 \pm 0.05 \pm 0.09$ ACCIARRI 97T L3 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

$0.96 \pm 0.03 \pm 0.13$ ALBRECHT 97B ARG $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

$1.26 \pm 0.26 \pm 0.18$ 36 BARU 90 MD1 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

$1.00 \pm 0.07 \pm 0.15$ 415 BEHREND 90C CELL 0 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

$1.03 \pm 0.13 \pm 0.21$ BUTLER 90 MRK2 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

$1.01 \pm 0.14 \pm 0.22$ 85 OEST 90 JADE $e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$

$0.90 \pm 0.27 \pm 0.15$ 56 ¹ ALTHOFF 86 TASS 0 $e^+ e^- \rightarrow e^+ e^- 3\pi$

$1.14 \pm 0.20 \pm 0.26$ ² ANTREASYAN 86 CBAL 0 $e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$

$1.06 \pm 0.18 \pm 0.19$ BERGER 84C PLUT 0 $e^+ e^- \rightarrow e^+ e^- 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.81 \pm 0.19^{+0.42}_{-0.11}$ 35 ¹ BEHREND 82C CELL 0 $e^+ e^- \rightarrow e^+ e^- 3\pi$

$0.77 \pm 0.18 \pm 0.27$ 22 ² EDWARDS 82F CBAL 0 $e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$

Γ_{10}

NODE=M012W7;LINKAGE=AD
NODE=M012W;LINKAGE=M2

NODE=M012W9
NODE=M012W9

1 From $\rho\pi$ decay mode.

2 From $\eta\pi^0$ decay mode.

NODE=M012W;LINKAGE=F
NODE=M012W;LINKAGE=G

$\Gamma(e^+e^-)$						Γ_{11}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		
< 0.56	90	ACHASOV 00K	SND	$e^+e^- \rightarrow \pi^0\pi^0$		NODE=M012W10 NODE=M012W10
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<25	90	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^0\eta$		

a₂(1320) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(3\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$						$\Gamma_1\Gamma_{10}/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.65 ± 0.02 ± 0.02	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$		

1 From analysis of L3 data at 183–209 GeV.

$\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$						$\Gamma_5\Gamma_{10}/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT			
• • • We do not use the following data for averages, fits, limits, etc. • • •						

0.145^{+0.097}_{-0.034} ¹ UEHARA 09A BELL $e^+e^- \rightarrow e^+e^-\eta\pi^0$ 1 From the D_2 -wave. The fraction of the D_0 -wave is 3.4^{+2.3%}_{-1.1%}.

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$						$\Gamma_7\Gamma_{10}/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT			
0.126 ± 0.007 ± 0.028 ¹ ALBRECHT 90G ARG $e^+e^- \rightarrow e^+e^-K^+K^-$						

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.081 ± 0.006 ± 0.027 ² ALBRECHT 90G ARG $e^+e^- \rightarrow e^+e^-K^+K^-$

1 Using an incoherent background.

2 Using a coherent background.

a₂(1320) BRANCHING RATIOS

$[\Gamma(f_2(1270)\pi) + \Gamma(\rho(1450)\pi)]/\Gamma(\rho(770)\pi)$						$(\Gamma_3 + \Gamma_4)/\Gamma_2$
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.12	90	ABRAMOVI...	70B	HBC	—	3.93 $\pi^- p$

$\Gamma(\rho(770)\pi)/\Gamma(f_2(1270)\pi)$						Γ_2/Γ_3
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
16.5 ^{+1.2} _{-2.4}	46M	¹ AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^-\pi^+\pi^- p$		

1 Statistical error negligible.

$\Gamma(\eta\pi)/\Gamma(3\pi)$						Γ_5/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
0.207 ± 0.018 OUR FIT						

0.213 ± 0.020 OUR AVERAGE

0.18 ± 0.05		FORINO	76	HBC	11 $\pi^- p$	
0.22 ± 0.05	52	ANTIPOV	73	CNTR	40 $\pi^- p$	
0.211 ± 0.044	149	CHALOUPKA	73	HBC	— 3.9 $\pi^- p$	
0.246 ± 0.042	167	ALSTON-...	71	HBC	+ 7.0 $\pi^+ p$	
0.25 ± 0.09	15	BOECKMANN	70	HBC	+ 5.0 $\pi^+ p$	
0.23 ± 0.08	22	ASCOLI	68	HBC	— 5 $\pi^- p$	
0.12 ± 0.08		CHUNG	68	HBC	— 3.2 $\pi^- p$	
0.22 ± 0.09		CONTE	67	HBC	— 11.0 $\pi^- p$	


$\Gamma(\omega\pi\pi)/\Gamma(3\pi)$						Γ_6/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
0.15 ± 0.05 OUR FIT Error includes scale factor of 1.3.						

0.15 ± 0.05 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

0.28 ± 0.09	60	DIAZ	74	DBC	0 6 $\pi^+ n$	
0.18 ± 0.08		¹ KARSHON	74	HBC	Avg. of above two	OCCUR=3
0.10 ± 0.05	279	² CHALOUPKA	73	HBC	— 3.9 $\pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.29 ± 0.08	140	¹ KARSHON	74	HBC	0 4.9 $\pi^+ p$	
0.10 ± 0.04	60	¹ KARSHON	74	HBC	+ 4.9 $\pi^+ p$	OCCUR=2
0.19 ± 0.08		DEFOIX	73	HBC	0 0.7 $\bar{p}p$	

¹KARSHON 74 suggest an additional $I = 0$ state strongly coupled to $\omega\pi\pi$ which could explain discrepancies in branching ratios and masses. We use a central value and a systematic spread.

²Decays to $b_1(1040)\pi$, $b_1 \rightarrow \omega\pi$. Error increased to account for possible systematic errors of complicated analysis.



$\Gamma(K\bar{K})/\Gamma(3\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.070±0.012 OUR FIT					
0.078±0.017		CHABAUD 78	RVUE		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.011±0.003	1	BERTIN 98B	OBLX	0.0	$\bar{p}p \rightarrow K^\pm K_s \pi^\mp$
0.056±0.014	50	2 CHALOUPKA 73	HBC	—	3.9 $\pi^- p$
0.097±0.018	113	2 ALSTON-...	71	HBC	+ 7.0 $\pi^+ p$
0.06 ± 0.03		2 ABRAMOVIC 70B	HBC	—	3.93 $\pi^- p$
0.054±0.022		2 CHUNG 68	HBC	—	3.2 $\pi^- p$

¹ Using 4π data from BERTIN 97D.

² Included in CHABAUD 78 review.

Γ_7/Γ_1

NODE=M012R1
NODE=M012R1

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.08±0.02	1	BERTIN 98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_s \pi^\mp$

¹ Using $\eta\pi\pi$ data from AMSLER 94D.

Γ_7/Γ_5

NODE=M012R14
NODE=M012R14

$\Gamma(\eta\pi)/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.162±0.012 OUR FIT					
0.140±0.028 OUR AVERAGE					
0.13 ± 0.04		ESPIGAT 72	HBC	±	0.0 $\bar{p}p$
0.15 ± 0.04	34	BARNHAM 71	HBC	+	3.7 $\pi^+ p$

NODE=M012R14;LINKAGE=BE

$\Gamma(K\bar{K})/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.054±0.009 OUR FIT					
0.048±0.012 OUR AVERAGE					
0.05 ± 0.02		TOET 73	HBC	+	5 $\pi^+ p$
0.09 ± 0.04		TOET 73	HBC	0	5 $\pi^+ p$
0.03 ± 0.02	8	¹ DAMERI 72	HBC	—	11 $\pi^- p$
0.06 ± 0.03	17	BARNHAM 71	HBC	+	3.7 $\pi^+ p$

NODE=M012R8
NODE=M012R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.020±0.004 ² ESPIGAT 72 HBC ± 0.0 $\bar{p}p$

OCCUR=2

¹ Montanet agrees. Vlada.

² Not averaged because of discrepancy between masses from $K\bar{K}$ and $\rho\pi$ modes.

NODE=M012R8;LINKAGE=01
NODE=M012R8;LINKAGE=A

$\Gamma(\eta'(958)\pi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_8/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.006	95	ALDE	92B	GAM2	38,100 $\pi^- p \rightarrow \eta' \pi^0 n$	NODE=M012R4
<0.02	97	BARNHAM	71	HBC	+ 3.7 $\pi^+ p$	NODE=M012R4
0.004±0.004	1	BOESEBECK	68	HBC	+ 8 $\pi^+ p$	

¹ No longer valid since $\Gamma(K\bar{K})/\Gamma(3\pi)$ value has changed (MORRISON 71).

 $\Gamma(\eta'(958)\pi)/\Gamma(3\pi)$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_8/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.011	90	EISENSTEIN	73	HBC	- 5 $\pi^- p$	NODE=M012R5
<0.04		ALSTON---	71	HBC	+ 7.0 $\pi^+ p$	NODE=M012R5
0.04 +0.03 -0.04		BOECKMANN	70	HBC	0 5.0 $\pi^+ p$	

 $\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_5
0.038±0.005 OUR AVERAGE				
0.05 ± 0.02	ADOLPH	15	COMP 191 $\pi^- p \rightarrow \eta' \pi^- p$	I
0.032±0.009	ABELE	97C	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta'$	
0.047±0.010±0.004	¹ BELADIDZE	93	VES 37 $\pi^- N \rightarrow a_2^- N$	
0.034±0.008±0.005	BELADIDZE	92	VES 36 $\pi^- C \rightarrow a_2^- C$	

¹ Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = 0.441$, $B(\eta \rightarrow \gamma\gamma) = 0.389$ and $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 0.236$.

 $\Gamma(\pi^\pm\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.005 +0.005 -0.003	¹ EISENBERG	72	HBC 4.3,5.25,7.5 γp	NODE=M012R11

¹ Pion-exchange model used in this estimation.

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6	90	ACHASOV	00K	SND $e^+ e^- \rightarrow \pi^0 \pi^0$	NODE=M012R15

 $a_2(1320)$ REFERENCES

RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=59554
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)	REFID=59003
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=55911
ALEKSEEV	10	PRI 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(BELLE Collab.)	REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	(BNL E852 Collab.)	REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	(BNL E852 Collab.)	REFID=51185
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
IVANOV	01	PRI 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
MOLCHANOV	01	PL B521 171	V.V. Molchanov <i>et al.</i>	(FNAL SELEX Collab.)	REFID=48559
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
ABELE	97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45531
ACCIARRI	97T	PL B413 147	M. Acciari <i>et al.</i>	(L3 Collab.)	REFID=45761
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=45418
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=45584
AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44649
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BELADIDZE	93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43598
CONDO	93	PR D48 3045	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=43600
ALDE	92B	ZPHY C54 549	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41852
BELADIDZE	92	ZPHY C54 235	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42171
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch (WA76 Coll.)	REFID=41375	
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41356
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBIEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023

Translated from YAF 48 436.

ALTHOFF	86	ZPHY C31 537	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21287
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
BERGER	84C	PL 149B 427	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=21286
BEHREND	82C	PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
CIHANGIR	82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BÖNN, CERN+)	REFID=20462
CHABAUD	80	NP B175 189	V. Chabaud <i>et al.</i>	(CERN, MPIM, AMST)	REFID=21274
DAUM	80C	PL 89B 276	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP	REFID=21275
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
CHABAUD	78	NP B145 349	V. Chabaud <i>et al.</i>	(CERN, MPIM)	REFID=21267
FERRERSORIA	78	PL 74B 287	A. Ferrer Soria <i>et al.</i>	(ORSAY, CERN, CDEF+)	REFID=21270
HYAMS	78	NP B146 303	B.D. Hyams <i>et al.</i>	(CERN, MPIM, ATEN)	REFID=21271
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA) JP	REFID=21272
MAY	77	PR D16 1983	E.N. May <i>et al.</i>	(ROCH, CORN)	REFID=20450
FORINO	76	NC 35A 465	A. Forino <i>et al.</i>	(BGNA, FIRZ, GENO, MILA+)	REFID=21259
MARGULIE	76	PR D14 667	M. Margulies <i>et al.</i>	(BNL, CUNY)	REFID=21261
EMMS	75	PL 58B 117	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL) JP	REFID=21254
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP	REFID=20843
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)	REFID=21248
KARSHON	74	PRL 32 852	U. Karshon <i>et al.</i>	(REHO)	REFID=21249
ANTIPOV	73	NP B63 175	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=21238
ANTIPOV	73C	NP B63 153	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=20817
CHALOUPKA	73	PL 44B 211	V. Chaloupka <i>et al.</i>	(CERN)	REFID=21242
CONFORTO	73	PL 45B 154	G. Conforto <i>et al.</i>	(EFI, FNAL, TNOT+) JP	REFID=21243
DEFOIX	73	PL 43B 141	C. Defoix <i>et al.</i>	(CDEF)	REFID=21244
EISENSTEIN	73	PR D7 278	L. Eisenstein <i>et al.</i>	(ILL)	REFID=21245
KEY	73	PRL 30 503	A.W. Key <i>et al.</i>	(TNOT, EFI, FNAL, WISC)	REFID=21246
TOET	73	NP B63 248	D.Z. Toet <i>et al.</i>	(NIJM, BONN, DURH, TORI)	REFID=20714
DAMERI	72	NC 9A 1	M. Dameri <i>et al.</i>	(GENO, MILA, SACL)	REFID=20338
EISENBERG	72	PR D5 15	Y. Eisenberg <i>et al.</i>	(REHO, SLAC, TELA)	REFID=20098
ESPIGAT	72	NP B36 93	P. Espigat <i>et al.</i>	(CERN, CDEF)	REFID=21232
FOLEY	72	PR D6 747	K.J. Foley <i>et al.</i>	(BNL, CUNY)	REFID=21233
ALSTON-...	71	PL 34B 156	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21214
BARNHAM	71	PRL 26 1494	K.W.J. Barnham <i>et al.</i>	(LBL)	REFID=21215
BINNIE	71	PL 36B 257	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=21217
BOWEN	71	PRL 26 1663	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21219
GRAYER	71	PL 34B 333	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=21223
ABRAMOVICH	70B	NP B23 466	M. Abramovich <i>et al.</i>	(CERN) JP	REFID=21195
ALSTON-...	70	PL 33B 607	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21196
BOECKMANN	70	NP B16 221	K. Boeckmann <i>et al.</i>	(BONN, DURH, NIJM+)	REFID=21202
ASCOLI	68	PRL 20 1321	G. Ascoli <i>et al.</i>	(ILL) JP	REFID=21171
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
CONTE	67	NC 51A 175	F. Conte <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=21166

NODE=M147

 $f_0(1370)$ $I^G(J^{PC}) = 0^+(0^{++})$

See also the mini-reviews on scalar mesons under $f_0(500)$ (see the index for the page number) and on non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).

$f_0(1370)$ T-MATRIX POLE POSITION

Note that $\Gamma \approx 2 \operatorname{Im}(\sqrt{s_{\text{pole}}})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1200–1500)–i(150–250) OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
(1290 ± 50)– i (170 ⁺²⁰ _{−40})	¹ ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$
(1373 ± 15)– i (137 ± 10)	² BARGIOTTI	03 OBLX	$\bar{p}p$
(1302 ± 17)– i (166 ± 18)	³ BARBERIS	00C	450 $p p \rightarrow p_f 4\pi p_s$
(1312 ± 25 ± 10)– i (109 ± 22 ± 15)	BARBERIS	99D OMEG	450 $p p \rightarrow K^+ K^-, \pi^+ \pi^-$
(1406 ± 19)– i (80 ± 6)	⁴ KAMINSKI	99 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
(1300 ± 20)– i (120 ± 20)	ANISOVICH	98B RVUE	Compilation
(1290 ± 15)– i (145 ± 15)	BARBERIS	97B OMEG	450 $p p \rightarrow \bar{p}p 2(\pi^+ \pi^-)$
(1548 ± 40)– i (560 ± 40)	BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
(1380 ± 40)– i (180 ± 25)	ABELE	96B CBAR	0.0 $\bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
(1300 ± 15)– i (115 ± 8)	BUGG	96 RVUE	
(1330 ± 50)– i (150 ± 40)	⁵ AMSLER	95B CBAR	$\bar{p}p \rightarrow 3\pi^0$
(1360 ± 35)– i (150–300)	⁵ AMSLER	95C CBAR	$\bar{p}p \rightarrow \pi^0 \eta\eta$
(1390 ± 30)– i (190 ± 40)	⁶ AMSLER	95D CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
1346 – i 249	^{7,8} JANSEN	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
1214 – i 168	^{8,9} TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1364 – i 139	AMSLER	94D CBAR	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
(1365 ⁺²⁰ _{−55})– i (134 ± 35)	ANISOVICH	94 CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$
(1340 ± 40)– i (127 ⁺³⁰ _{−20})	¹⁰ BUGG	94 RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$
(1430 ± 5)– i (73 ± 13)	¹¹ KAMINSKI	94 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
1420 – i 220	¹² AU	87 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$

¹ Another pole is found at $(1510 \pm 130) - i(800 \pm 100)$ MeV.

² Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

³ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

⁴ T-matrix pole on sheet ——.

⁵ Supersedes ANISOVICH 94.

⁶ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$, and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁷ Analysis of data from FALVARD 88.

⁸ The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

⁹ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

¹⁰ Reanalysis of ANISOVICH 94 data.

¹¹ T-matrix pole on sheet III.

¹² Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

$f_0(1370)$ BREIT-WIGNER MASS OR K-MATRIX POLE PARAMETER

VALUE (MeV)	DOCUMENT ID
1200 to 1500 OUR ESTIMATE	

NODE=M147

NODE=M147PP

NODE=M147PP

NODE=M147PP

→ UNCHECKED ←

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M147PP;LINKAGE=AO

NODE=M147PP;LINKAGE=BG

NODE=M147PP;LINKAGE=PC

NODE=M147PP;LINKAGE=TK

NODE=M147PP;LINKAGE=K

NODE=M147PP;LINKAGE=A

NODE=M147PP;LINKAGE=C

NODE=M147PP;LINKAGE=DD

NODE=M147PP;LINKAGE=BB

NODE=M147PP;LINKAGE=C1

NODE=M147PP;LINKAGE=KM

NODE=M147PP;LINKAGE=H

NODE=M147205

NODE=M147M

→ UNCHECKED ←

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1400±40		¹ AUBERT	09L	BABR $B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 ^{+ 6 +} _{- 7 -} 72		² UEHARA	08A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1259±55	2.6k	BONVICINI	07	CLEO $D^+ \rightarrow \pi^- \pi^+ \pi^+$
1309± 1± 15		³ BUGG	07A	RVUE $0.0 p\bar{p} \rightarrow 3\pi^0$
1449±13	4.3k	⁴ GARMASH	06	BELL $B^+ \rightarrow K^+ \pi^+ \pi^-$
1350±50		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$
1265±30 ^{+ 20} _{- 35}		ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1434±18± 9	848	AITALA	01A	E791 $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
1308±10		BARBERIS	99B	OMEG $450 pp \rightarrow p_s p_f \pi^+ \pi^-$
1315±50		BELLAZZINI	99	GAM4 $450 pp \rightarrow pp \pi^0 \pi^0$
1315±30		ALDE	98	GAM4 $100 \pi^- p \rightarrow \pi^0 \pi^0 n$
1280±55		BERTIN	98	OBLX $0.05-0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1186		^{5,6} TORNQVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472±12		ARMSTRONG	91	OMEG $300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275±20		BREAKSTONE	90	SFM $62 pp \rightarrow pp\pi^+ \pi^-$
1420±20		AKESSON	86	SPEC $63 pp \rightarrow pp\pi^+ \pi^-$
1256		FROGGATT	77	RVUE $\pi^+ \pi^-$ channel
1 Breit-Wigner mass.				
2 Breit-Wigner mass. May also be the $f_0(1500)$.				
3 Reanalysis of ABELE 96C data.				
4 Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.				
5 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.				
6 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$ decays				

NODE=M147M1

NODE=M147M1

 $K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1422±15±28		¹ AAIJ	19H	LHCb $pp \rightarrow D^\pm X$
1360±31±28	430	^{2,3} DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
1350±48±15	168	^{2,3} DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
1440± 6		VLADIMIRSK...	06	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1391±10		TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440±50		BOLONKIN	88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1463± 9		ETKIN	82B	MPS $23 \pi^- p \rightarrow n2K_S^0$
1425±15		WICKLUND	80	SPEC $6 \pi N \rightarrow K^+ K^- N$
~1300		POLYCHRO...	79	STRC $7 \pi^- p \rightarrow n2K_S^0$

NODE=M147M1;LINKAGE=BW

NODE=M147M1;LINKAGE=UE

NODE=M147M1;LINKAGE=BU

NODE=M147M1;LINKAGE=GR

NODE=M147M1;LINKAGE=BB

NODE=M147M1;LINKAGE=FF

NODE=M147M2

NODE=M147M2

OCCUR=4

OCCUR=2

NODE=M147M2;LINKAGE=F

NODE=M147M2;LINKAGE=A

NODE=M147M2;LINKAGE=B

NODE=M147M3

NODE=M147M3

 4π MODE 2($\pi\pi$) $s+\rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1395±40		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1374±38		AMSLER	94	CBAR $0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
1345±12		ADAMO	93	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
1386±30		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~1410	5751	¹ BETTINI	66	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
1 $\rho\rho$ dominant.				

NODE=M147M3;LINKAGE=BE

NODE=M147M4

NODE=M147M4

 $\eta\eta$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1262 ^{+51 +} _{-78 -} 82		¹ UEHARA	10A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1430		AMSLER	92	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
1220±40		ALDE	86D	GAM4 $100 \pi^- p \rightarrow n2\eta$
1 Breit-Wigner mass. May also be the $f_0(1500)$.				

NODE=M147M4;LINKAGE=UE

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1330.2 \pm 5.9	¹ AAIJ 19H LHCb $p\bar{p} \rightarrow D^\pm X$		
1306 \pm 20	² ANISOVICH 03 RVUE		
¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18. ² K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.			

NODE=M147M5

NODE=M147M5

 $f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID
200 to 500 OUR ESTIMATE	

NODE=M147210

NODE=M147W

→ UNCHECKED ←

ππ MODE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
300 \pm 80	¹ AUBERT 09L BABR $B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$		
90+ 90- 2+ 1- 50 22	² UEHARA 08A BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$		
298 \pm 21	³ BONVICINI 07 CLEO $D^+ \rightarrow \pi^- \pi^+ \pi^+$		
126 \pm 25	³ GARMASH 06 BELL $B^+ \rightarrow K^+ \pi^+ \pi^-$		
265 \pm 40	ABLIKIM 05 BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$		
350 \pm 100+ 105 60	ABLIKIM 05Q BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$		
173 \pm 32 \pm 6	⁸⁴⁸ AITALA 01A E791 $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$		
222 \pm 20	BARBERIS 99B OMEG 450 $p\bar{p} \rightarrow p_s p_f \pi^+ \pi^-$		
255 \pm 60	BELLAZZINI 99 GAM4 450 $p\bar{p} \rightarrow p p \pi^0 \pi^0$		
190 \pm 50	ALDE 98 GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$		
323 \pm 13	BERTIN 98 OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$		
350	^{4,5} TORNQVIST 95 RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$		
195 \pm 33	ARMSTRONG 91 OMEG 300 $p\bar{p} \rightarrow p p \pi\pi, p p K\bar{K}$		
285 \pm 60	BREAKSTONE 90 SFM 62 $p\bar{p} \rightarrow p p \pi^+ \pi^-$		
460 \pm 50	AKESSON 86 SPEC 63 $p\bar{p} \rightarrow p p \pi^+ \pi^-$		
~400	⁶ FROGGATT 77 RVUE $\pi^+ \pi^-$ channel		

NODE=M147W1
NODE=M147W1

1 The systematic errors are not reported.

2 Breit-Wigner width. May also be the $f_0(1500)$.3 Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays. Supersedes GARMASH 05.

4 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

5 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^- \pi^- \pi^0 \nu_\tau$ decays

6 Width defined as distance between 45 and 135° phase shift.

NODE=M147W1;LINKAGE=NS

NODE=M147W1;LINKAGE=UE

NODE=M147W1;LINKAGE=GR

NODE=M147W1;LINKAGE=BB

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
324 \pm 38 \pm 42	¹ AAIJ 19H LHCb $p\bar{p} \rightarrow D^\pm X$		
121 \pm 15	VLADIMIRSK..06 SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$		
55 \pm 26	TIKHOMIROV 03 SPEC 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$		
250 \pm 80	BOLONKN 88 SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$		
118+ 16	ETKIN 82B MPS 23 $\pi^- p \rightarrow n2K_S^0$		
160 \pm 30	WICKLUND 80 SPEC 6 $\pi N \rightarrow K^+ K^- N$		
~150	POLYCHRO... 79 STRC 7 $\pi^- p \rightarrow n2K_S^0$		

NODE=M147W2
NODE=M147W2

OCCUR=3

1 From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the isobar model A.

NODE=M147W2;LINKAGE=C

4π MODE $2(\pi\pi)_S + \rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
275 \pm 55		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 \pm 61		AMSLER 94	CBAR	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
398 \pm 26		ADAMO 93	OBLX	$\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 \pm 50		GASPERO 93	DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
~ 90	5751	¹ BETTINI 66	DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
$1 \rho\rho$ dominant.				

 $\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$484^{+246}_{-170}{}^{+246}_{-263}$	¹ UEHARA 10A	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
250	AMSLER 92	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
320 ± 40	ALDE 86D	GAM4	$100 \pi^- p \rightarrow n 2\eta$

¹ Breit-Wigner width. May also be the $f_0(1500)$.

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
147^{+30}_{-50}	¹ ANISOVICH 03	RVUE
¹ K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.		

 $f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 4\pi$	seen
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 2\pi^+ 2\pi^-$	seen
$\Gamma_5 \pi^+ \pi^- 2\pi^0$	seen
$\Gamma_6 \rho\rho$	seen
$\Gamma_7 2(\pi\pi)_S$ -wave	seen
$\Gamma_8 \pi(1300)\pi$	seen
$\Gamma_9 a_1(1260)\pi$	seen
$\Gamma_{10} \eta\eta$	seen
$\Gamma_{11} K\bar{K}$	seen
$\Gamma_{12} K\bar{K}n\pi$	not seen
$\Gamma_{13} 6\pi$	not seen
$\Gamma_{14} \omega\omega$	not seen
$\Gamma_{15} \gamma\gamma$	seen
$\Gamma_{16} e^+ e^-$	not seen

 $f_0(1370)$ PARTIAL WIDTHS **$\Gamma(\gamma\gamma)$**

See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90.

 Γ_{15}

NODE=M147W3
NODE=M147W3

 $\Gamma(e^+ e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBYEV 88	ND	$e^+ e^- \rightarrow \pi^0 \pi^0$

 Γ_{16}

NODE=M147W3;LINKAGE=BE

NODE=M147W4
NODE=M147W4

NODE=M147W4;LINKAGE=UE

NODE=M147W5
NODE=M147W5

NODE=M147W;LINKAGE=KM

NODE=M147215;NODE=M147

DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=10;OUR EST; \rightarrow UNCHECKED
 DESIG=4;OUR EST; \rightarrow UNCHECKED
 DESIG=5;OUR EST; \rightarrow UNCHECKED
 DESIG=6;OUR EST; \rightarrow UNCHECKED
 DESIG=14;OUR EST; \rightarrow UNCHECKED
 DESIG=15;OUR EST; \rightarrow UNCHECKED
 DESIG=16;OUR EVAL; \rightarrow UNCHECKED
 DESIG=17;OUR EVAL; \rightarrow UNCHECKED
 DESIG=2;OUR EST; \rightarrow UNCHECKED
 DESIG=11;OUR EST; \rightarrow UNCHECKED
 DESIG=18;OUR EVAL; \rightarrow UNCHECKED
 DESIG=19;OUR EVAL; \rightarrow UNCHECKED
 DESIG=20;OUR EVAL; \rightarrow UNCHECKED
 DESIG=12;OUR EST; \rightarrow UNCHECKED
 DESIG=13;OUR EST; \rightarrow UNCHECKED

NODE=M147217

NODE=M147W11
NODE=M147W11
NODE=M147W11

NODE=M147W12
NODE=M147W12

$f_0(1370) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	$\Gamma_{10}\Gamma_{15}/\Gamma$
VALUE (eV)			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
121 $^{+133}_{-53}$ $^{+169}_{-106}$	¹ UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1 Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.			

 $f_0(1370) \text{ BRANCHING RATIOS}$

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	Γ_1/Γ
VALUE	CL%		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
<0.10	95	OCHS	13 RVUE
0.26 \pm 0.09		BUGG	96 RVUE
<0.15	¹ AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
<0.06		GASPERO	93 DBC 0.0 $\bar{p}n \rightarrow$ hadrons
1 Using AMSLER 95B ($3\pi^0$).			

$\Gamma(4\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	$\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
>0.72		GASPERO	93 DBC 0.0 $\bar{p}n \rightarrow$ hadrons

$\Gamma(4\pi^0)/\Gamma(4\pi)$	DOCUMENT ID	TECN	Γ_3/Γ_2
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
seen	ABELE	96	CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$
0.068 \pm 0.005	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

 1 Model-dependent evaluation.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(4\pi)$	DOCUMENT ID	TECN	$\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.420 \pm 0.014	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$

 1 Model-dependent evaluation.

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(4\pi)$	DOCUMENT ID	TECN	$\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.512 \pm 0.019	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

 1 Model-dependent evaluation.

$\Gamma(\rho\rho)/\Gamma(4\pi)$	DOCUMENT ID	TECN	Γ_6/Γ_2
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.26 \pm 0.07	ABELE	01B CBAR	0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(2(\pi\pi)s\text{-wave})/\Gamma(\pi\pi)$	DOCUMENT ID	TECN	Γ_7/Γ_1
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
5.6 \pm 2.6	¹ ABELE	01 CBAR	0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$

 1 From the combined data of ABELE 96 and ABELE 96C.

$\Gamma(2(\pi\pi)s\text{-wave})/\Gamma(4\pi)$	DOCUMENT ID	TECN	Γ_7/Γ_2
VALUE			
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.51 \pm 0.09	ABELE	01B CBAR	0.0 $\bar{p}d \rightarrow 5\pi p$

NODE=M147225

NODE=M147G01
NODE=M147G01

NODE=M147G01;LINKAGE=UE

NODE=M147220

NODE=M147R3
NODE=M147R3

NODE=M147R3;LINKAGE=B

NODE=M147R4
NODE=M147R4NODE=M147R12
NODE=M147R12

NODE=M147R12;LINKAGE=GA

NODE=M147R5
NODE=M147R5

NODE=M147R5;LINKAGE=A

NODE=M147R6
NODE=M147R6

NODE=M147R6;LINKAGE=A

NODE=M147R17
NODE=M147R17NODE=M147R15
NODE=M147R15

NODE=M147R;LINKAGE=KZ

NODE=M147R16
NODE=M147R16

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)s\text{-wave})$					Γ_6/Γ_7
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R10 NODE=M147R10
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
large	BARBERIS 00C		$450 \text{ } pp \rightarrow p_f 4\pi p_s$		
1.6 ± 0.2	AMSLER 94	CBAR	$\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$		
~ 0.65	GASPERO 93	DBC	$0.0 \text{ } \bar{p}n \rightarrow \text{hadrons}$		OCCUR=2
$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$					Γ_8/Γ_2
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R18 NODE=M147R18
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.17 ± 0.06	ABELE 01B	CBAR	$0.0 \text{ } \bar{p}d \rightarrow 5\pi p$		
$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$					Γ_9/Γ_2
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R19 NODE=M147R19
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.06 ± 0.02	ABELE 01B	CBAR	$0.0 \text{ } \bar{p}d \rightarrow 5\pi p$		
$\Gamma(\eta\eta)/\Gamma(4\pi)$					$\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R14 NODE=M147R14
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$(28 \pm 11) \times 10^{-3}$	¹ ANISOVICH 02D	SPEC	Combined fit		
$(4.7 \pm 2.0) \times 10^{-3}$	BARBERIS 00E		$450 \text{ } pp \rightarrow p_f \eta\eta p_s$		
1 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.					
$\Gamma(K\bar{K})/\Gamma_{\text{total}}$					Γ_{11}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R11 NODE=M147R11
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.35 ± 0.13	BUGG 96	RVUE			
$\Gamma(K\bar{K})/\Gamma(\pi\pi)$					Γ_{11}/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R13 NODE=M147R13
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.08 ± 0.08	ABLIKIM 05	BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$, $\phi K^+ K^-$		
0.91 ± 0.20	¹ BARGIOTTI 03	OBLX	$\bar{p}p$		
0.12 ± 0.06	² ANISOVICH 02D	SPEC	Combined fit		
$0.46 \pm 0.15 \pm 0.11$	BARBERIS 99D	OMEG	$450 \text{ } pp \rightarrow K^+ K^-$, $\pi^+ \pi^-$		
1 Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.					
2 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.					
$\Gamma(K\bar{K}\eta\eta)/\Gamma_{\text{total}}$					Γ_{12}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R20 NODE=M147R20
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.03	GASPERO 93	DBC	$0.0 \text{ } \bar{p}n \rightarrow \text{hadrons}$		
$\Gamma(6\pi)/\Gamma_{\text{total}}$					Γ_{13}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R21 NODE=M147R21
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.22	GASPERO 93	DBC	$0.0 \text{ } \bar{p}n \rightarrow \text{hadrons}$		
$\Gamma(\omega\omega)/\Gamma_{\text{total}}$					Γ_{14}/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M147R22 NODE=M147R22
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.13	GASPERO 93	DBC	$0.0 \text{ } \bar{p}n \rightarrow \text{hadrons}$		

f₀(1370) REFERENCES

NODE=M147

AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	(NWES)	REFID=59987
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>		REFID=56805
OCHS	13	JP G40 043001	W. Ochs		REFID=55367
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52723
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=51721
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>		REFID=53252
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51594
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51162
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45011
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45038
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)	REFID=45094
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist	(HELS)	REFID=44529
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC	REFID=43660
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC	REFID=43657
GASPERO	93	NP A562 407	M. Gaspero	(ROMAI) JPC	REFID=43658
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)	REFID=40064
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)	REFID=21123
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20383
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)	REFID=20349
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)	REFID=44530
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)	REFID=21361

$\pi_1(1400)$

$I^G(J^{PC}) = 1^-(1^-+)$

See also the mini-review under non- $q\bar{q}$ candidates in PDG 06, Journal of Physics **G33** 1 (2006).


$\pi_1(1400)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1354 ± 25 OUR AVERAGE					Error includes scale factor of 1.8. See the ideogram below.
1257 ± 20	± 25	23.5k	ADAMS	07B B852	18 $\pi^- p \rightarrow \eta \pi^0 n$
1384 ± 20	± 35	90k	SALVINI	04 OBLX	$\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
1360 ± 25			ABELE	99 CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$
1400 ± 20	± 20		ABELE	98B CBAR	0.0 $\bar{p} n \rightarrow \pi^- \pi^0 \eta$
1370 ± 16	+ 50 - 30		1 THOMPSON	97 MPS	18 $\pi^- p \rightarrow \eta \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1323.1 ± 4.6		2 AOYAGI	93 BKEI		$\pi^- p \rightarrow \eta \pi^- p$
1406 ± 20		3 ALDE	88B GAM4 0		100 $\pi^- p \rightarrow \eta \pi^0 n$

1 Natural parity exchange, questioned by DZIERBA 03.

2 Unnatural parity exchange.

3 Seen in the P_0 -wave intensity of the $\eta \pi^0$ system, unnatural parity exchange.



$\pi_1(1400)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
330 ± 35 OUR AVERAGE					
354 ± 64	± 58	23.5k	ADAMS	07B B852	18 $\pi^- p \rightarrow \eta \pi^0 n$
378 ± 50	± 50	90k	SALVINI	04 OBLX	$\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
220 ± 90			ABELE	99 CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$
310 ± 50	+ 50 - 30		ABELE	98B CBAR	0.0 $\bar{p} n \rightarrow \pi^- \pi^0 \eta$
385 ± 40	+ 65 - 105		4 THOMPSON	97 MPS	18 $\pi^- p \rightarrow \eta \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
143.2 ± 12.5		5 AOYAGI	93 BKEI		$\pi^- p \rightarrow \eta \pi^- p$
180 ± 20		6 ALDE	88B GAM4 0		100 $\pi^- p \rightarrow \eta \pi^0 n$

4 Resolution is not unfolded, natural parity exchange, questioned by DZIERBA 03.

5 Unnatural parity exchange.

6 Seen in the P_0 -wave intensity of the $\eta \pi^0$ system, unnatural parity exchange.

NODE=M111M

NODE=M111M

NODE=M111M;LINKAGE=B

NODE=M111M;LINKAGE=C

NODE=M111M;LINKAGE=A

NODE=M111W

NODE=M111W

NODE=M111W;LINKAGE=QQ

NODE=M111W;LINKAGE=C

NODE=M111W;LINKAGE=A

$\pi_1(1400)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta\pi^0$	seen
$\Gamma_2 \eta\pi^-$	seen
$\Gamma_3 \eta'\pi$	
$\Gamma_4 \rho(770)\pi$	not seen

 $\pi_1(1400)$ BRANCHING RATIOS **$\Gamma(\eta\pi^0)/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen	PROKOSHKIN 95B	GAM4		$100 \pi^- p \rightarrow \eta\pi^0 n$	
not seen	7 BUGG	94	RVUE	$\bar{p}p \rightarrow \eta 2\pi^0$	
not seen	8 APEL	81	NICE 0	$40 \pi^- p \rightarrow \eta\pi^0 n$	

7 Using Crystal Barrel data.

8 A general fit allowing S , D , and P waves (including $m=0$) is not done because of limited statistics. **$\Gamma(\eta\pi^-)/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
possibly seen	BELADIDZE 93	VES	$37\pi^- N \rightarrow \eta\pi^- N$	

 $\Gamma(\eta'\pi)/\Gamma(\eta\pi^0)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.80	95	BOUTEMEUR 90	GAM4	$100 \pi^- p \rightarrow 4\gamma n$	

 $\Gamma(\rho(770)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
not seen	AGHASYAN 18B	18B COMP	$190 \pi^- p \rightarrow \pi^-\pi^+\pi^- p$	

 $\pi_1(1400)$ REFERENCES

AGHASYAN 18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ADAMS 07B	PL B657 27	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
PDG 06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
SALVINI 04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)
DZIERBA 03	PR D67 094015	A.R. Dzierba <i>et al.</i>	
ABELE 99	PL B446 349	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE 98B	PL B423 175	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
THOMPSON 97	PRC 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)
PROKOSHKIN 95B	PAN 58 606	Y.D. Prokoshkin, S.A. Sadovsky	(SERP)
	Translated from YAF 58 662.		
BUGG 94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
AOYAGI 93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)
BELADIDZE 93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)
BOUTEMEUR 90	Hadron 89 Conf. p 119	M. Boutemeur, M. Poulet	(SERP, BELG, LANL, LAPP) IGJPC
ALDE 88B	PL B205 397	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP) IGJPC
APEL 81	NP B193 269	W.D. Apel <i>et al.</i>	(SERP, CERN)

NODE=M111215;NODE=M111

DESIG=1;OUR EST; \rightarrow UNCHECKED ←
DESIG=4;OUR EST; \rightarrow UNCHECKED ←
DESIG=3
DESIG=5

NODE=M111220

NODE=M111R1
NODE=M111R1NODE=M111R1;LINKAGE=C
NODE=M111R1;LINKAGE=BNODE=M111R4
NODE=M111R4NODE=M111R3
NODE=M111R3NODE=M111R00
NODE=M111R00

NODE=M111

REFID=59471
REFID=52048
REFID=51004
REFID=53226
REFID=49412
REFID=46602
REFID=45864
REFID=45584
REFID=44619

REFID=44078
REFID=43599
REFID=43598
REFID=41751
REFID=40558
REFID=22913

$\eta(1405)$

$I^G(J^{PC}) = 0^+(0^-+)$

See also the $\eta(1475)$.

See the related review(s):

Pseudoscalar and Pseudovector Mesons in the 1400 MeV Region

NODE=M027

NODE=M027


$\eta(1405)$ MASS

NODE=M027205

VALUE (MeV)

DOCUMENT ID

1408.8±2.0 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.



$\eta\pi\pi$ MODE

NODE=M027M1

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M027M1

1405.8± 2.6 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.

1409.0± 1.7	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
1407.0± 3.5	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$
1399.8± 2.2 ^{+2.8} _{-0.1}	¹	ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
1392 ± 14	900 ± 375	AMSLER	04B CBAR	$0\bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$
1394 ± 8	6.6 ± 2.0k	AMSLER	04B CBAR	$0\bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$
1404 ± 6	9082	MANAK	00A MPS	$18\pi^-p \rightarrow \eta\pi^+\pi^-n$
1424 ± 6	2200	ALDE	97B GAM4	$100\pi^-p \rightarrow \eta\pi^0\pi^0n$
1400 ± 6	²	BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
1388 ± 4		FUKUI	91C SPEC	$8.95\pi^-p \rightarrow \eta\pi^+\pi^-n$
1398 ± 6	261	³ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
1420 ± 5		ANDO	86 SPEC	$8\pi^-p \rightarrow \eta\pi^+\pi^-n$

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1404.0±11.0	195	ABLIKIM	19BABES3	$e^+e^- \rightarrow \psi(2S)$
1385 ± 7		BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
1409 ± 3		⁴ AMSLER	95F CBAR	$0\bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$

¹ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.

² From fit to the $a_0(980)\pi^- + \pi^+$ partial wave.

³ Best fit with a single Breit Wigner.


⁴ Superseded by AMSLER 04B.

NODE=M027M1;LINKAGE=BL

NODE=M027M1;LINKAGE=J1

NODE=M027M1;LINKAGE=A1

NODE=M027M1;LINKAGE=A



K $\bar{K}\pi$ MODE ($a_0(980)\pi$ or direct K $\bar{K}\pi$)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M027M4
NODE=M027M4

1413.9± 1.7 OUR AVERAGE Error includes scale factor of 1.1.

1413 ± 14	3651	1 NICHITU	02 OBLX	$0 \bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1416 ± 4 ± 2	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
1405 ± 5		2 CICALO	99 OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
1407 ± 5		2 BERTIN	97 OBLX	$0 \bar{p}p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
1416 ± 2		2 BERTIN	95 OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
1416 ± 8	+7 -5	700	3 BAI	90C MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1413 ± 5		3 RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K^0_S \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1459 ± 5		4 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
----------	--	------------	--------	---

1 Decaying dominantly directly to $K^+ K^- \pi^0$.

2 Decaying into $(K\bar{K})_S \pi$, $(K\pi)_S \bar{K}$, and $a_0(980)\pi$.

3 From fit to the $a_0(980)\pi$ $0^- +$ partial wave. Cannot rule out a $a_0(980)\pi$ 1^{++} partial wave.

4 Excluded from averaging because averaging would be meaningless.

OCCUR=2

OCCUR=3

NODE=M027M;LINKAGE=NC

NODE=M027M4;LINKAGE=FX

NODE=M027M4;LINKAGE=C2

NODE=M027M4;LINKAGE=AA

NODE=M027M2

NODE=M027M2

$\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1403±17 OUR AVERAGE Error includes scale factor of 1.8.

1390±12	235 ± 91	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$
1424±10±11	547	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
• • •	We do not use the following data for averages, fits, limits, etc. • • •			
1401±18		1,2 AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+ \pi^- \gamma\gamma$
1432±8		2 COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+ \pi^- 2\gamma$

OCCUR=4

NODE=M027M2;LINKAGE=E

NODE=M027M2;LINKAGE=X

4 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1420±20		BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+ \pi^- \pi^+ \pi^-$
1489±12	3270	1 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

NODE=M027M3

NODE=M027M3

1 Estimated by us from various fits.

NODE=M027M3;LINKAGE=E

$K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1452.7 ± 3.3	191	1,2 ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$	
1437.6 ± 3.2	249 ± 35	1,2 ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$	
1445.9 ± 5.7	62 ± 18	1,2 ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$	OCCUR=2
1442 ± 10	410	1 BAI	98C BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	
1445 ± 8	693	1 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
1433 ± 8	296	1 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	OCCUR=2
1413 ± 8	500	1 DUCH	89 ASTE	$\bar{p}p \rightarrow \pi^+ \pi^- K^\pm \pi^\mp K^0$	
1453 ± 7	170	1 RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$	OCCUR=2
1419 ± 1	8800	1 BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$	
1424 ± 3	620	1 REEVES	86 SPEC	$6.6 p \bar{p} \rightarrow K\bar{K}\pi X$	
1421 ± 2		1 CHUNG	85 SPEC	$8 \pi^- p \rightarrow K\bar{K}\pi n$	
1440 ⁺²⁰ ₋₁₅	174	1 EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	
1440 ⁺¹⁰ ₋₁₅		1 SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
1425 ± 7	800	1,3 BAUILLON	67 HBC	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$	

1 These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

2 Systematic uncertainty not evaluated.

3 From best fit of $0^- +$ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.

NODE=M027M6
NODE=M027M6

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M027M;LINKAGE=NP

NODE=M027M;LINKAGE=NS


NODE=M027M6;LINKAGE=H

NODE=M027210

NODE=M027WX

 $\eta(1405)$ WIDTH

VALUE (MeV)	DOCUMENT ID	
50.1 ± 2.6 OUR AVERAGE	Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.7. See the ideogram below.	



NODE=M027W1
NODE=M027W1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
The data in this block is included in the average printed for a previous datablock.					

52.6 ± 3.2 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

48.3 ± 5.2	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^0)$	
55.0 ± 11.0	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^0 \pi^0 \pi^0)$	OCCUR=2
52.8 ± 7.6 ^{+0.1} _{-7.6}		1 ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+ \pi^-)$	
55 ± 11	900	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$	

55	± 12	6.6k	AMSLER	04B	CBAR	0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \gamma$
80	± 21	9.0k	MANAK	00A	MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
85	± 18	2.2k	ALDE	97B	GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
47	± 13		² BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
59	± 4		FUKUI	91C	SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
53	± 11		³ AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
31	± 7		ANDO	86	SPEC	8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

79.0 ± 16.0	195	ABLIKIM	19BA	BES3	$e^+ e^- \rightarrow \psi(2S)$
86 ± 10		⁴ AMSLER	95F	CBAR	0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta$


1 The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.

2 From fit to the $a_0(980) \pi^- \pi^+$ partial wave.

3 From $\eta \pi^+ \pi^-$ mass distribution - mainly $a_0(980) \pi^-$ - no spin-parity determination available.

4 Superseded by AMSLER 04B.

OCCUR=2



K̄K̄π MODE ($a_0(980)\pi$ or direct $K\bar{K}\pi$)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

NODE=M027W4
NODE=M027W4

48 \pm 4 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.
51 \pm 6	3651
42 \pm 10 \pm 9	20k
50 \pm 4	
48 \pm 5	
50 \pm 4	
75 \pm 9	
91 \pm 67 \pm 15	
91 \pm 31 \pm 38	
19 \pm 7	
1	NICHITIU
2	ADAMS
3	CICALO
2	BERTIN
2	BERTIN
	AUGUSTIN
3	BAI
3	RATH
02	OBLX
01B	B852
99	OBLX
97	OBLX
95	OBLX
92	DM2
90C	MRK3
89	MPS
0	$\bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
18	GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
0	$\bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
0	$\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
$K\bar{K}$	$\rightarrow J/\psi \rightarrow \gamma K\bar{K}\pi$
$K\bar{K}$	$\rightarrow J/\psi \rightarrow \gamma K^0 S \bar{K}^\pm \pi^\mp$
$K\bar{K}$	$\rightarrow J/\psi \rightarrow \gamma K^0 S K^\pm \pi^\mp$
21.4	$\pi^- p \rightarrow n K^0 S K^\pm \pi^\mp$

OCCUR=2


OCCUR=3

NODE=M027W;LINKAGE=NC
NODE=M027W4;LINKAGE=FX
NODE=M027W4;LINKAGE=C

1 Decaying dominantly directly to $K^+ K^- \pi^0$.

2 Decaying into $(K\bar{K})_S \pi$, $(K\pi)_S \bar{K}$, and $a_0(980)\pi$.

3 From fit to the $a_0(980)\pi^- \pi^+$ partial wave, but $a_0(980)\pi^- \pi^+$ cannot be excluded.



$\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
89 ±17 OUR AVERAGE				Error includes scale factor of 1.7.
64 ±18	235 ± 91	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \gamma$
101.0 ± 8.8 ± 8.8	547	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
174 ±44		AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+ \pi^- \gamma \gamma$
90 ±26	1 COFFMAN		90 MRK3	$J/\psi \rightarrow \pi^+ \pi^- 2\gamma$

1 This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

NODE=M027W2
NODE=M027W2

4π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
160 ± 30		BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
144 ± 13	3270	1 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

1 Estimated by us from various fits.

OCCUR=2

$K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
45.9 ± 8.2	191	1,2 ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K\bar{K}\pi$
48.9 ± 9.0	249 ± 35	1,2 ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
34.2 ± 18.5	62 ± 18	1,2 ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
93 ± 14	296	1 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
105 ± 10	693	1 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
62 ± 16	500	1 DUCH	89 ASTE	$\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
100 ± 11	170	1 RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
66 ± 2	8800	1 BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
60 ± 10	620	1 REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K\bar{K}\pi X$
60 ± 10		1 CHUNG	85 SPEC	$8 \pi^- p \rightarrow K\bar{K}\pi n$
55 ± 20	174	1 EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
50 ± 30		1 SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
80 ± 10	800	1,3 BAILLON	67 HBC	$0.0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

NODE=M027W2;LINKAGE=X

NODE=M027W3
NODE=M027W3

1 These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

NODE=M027W3;LINKAGE=F2

2 Systematic uncertainty not evaluated.

NODE=M027W6
NODE=M027W6

3 From best fit to 0^-+ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M027W;LINKAGE=NP

NODE=M027W;LINKAGE=NS

NODE=M027W6;LINKAGE=H1

$\eta(1405)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 K\bar{K}\pi$	seen	
$\Gamma_2 \eta\pi\pi$	seen	
$\Gamma_3 a_0(980)\pi$	seen	
$\Gamma_4 \eta(\pi\pi)_S$ -wave	seen	
$\Gamma_5 f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$	not seen	
$\Gamma_6 f_0(980)\eta$	seen	
$\Gamma_7 4\pi$	seen	
$\Gamma_8 \rho\rho$	<58 %	99.85%
$\Gamma_9 \gamma\gamma$		
$\Gamma_{10} \rho^0\gamma$	seen	
$\Gamma_{11} \phi\gamma$		
$\Gamma_{12} K^*(892)K$	seen	

 $\eta(1405) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_9/\Gamma$
$\text{VALUE (keV)} \quad \text{CL\%}$	$\text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.035 90 1,2 AHOHE 05 CLE2 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$

1 Using $\eta(1405)$ mass and width 1410 MeV and 51 MeV, respectively.

2 Assuming three-body phase-space decay to $K_S^0K^\pm\pi^\mp$.

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_9/\Gamma$
$\text{VALUE (keV)} \quad \text{CL\%}$	$\text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.095 95 ACCIARRI 01G L3 183–202 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

$\Gamma(\rho^0\gamma) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{10}\Gamma_9/\Gamma$
$\text{VALUE (keV)} \quad \text{CL\%}$	$\text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5 95 ALTHOFF 84E TASS $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$

 $\eta(1405)$ BRANCHING RATIOS

$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$	Γ_2/Γ_1
$\text{VALUE} \quad \text{CL\%}$	$\text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

1.09±0.48 1 AMSLER 04B CBAR 0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.5 90 EDWARDS 83B CBAL $J/\psi \rightarrow \eta\pi\pi\gamma$

<1.1 90 SCHARRÉ 80 MRK2 $J/\psi \rightarrow \eta\pi\pi\gamma$

<1.5 95 FOSTER 68B HBC 0.0 $\bar{p}p$

1 Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.

$\Gamma(\rho^0\gamma)/\Gamma(\eta\pi\pi)$	Γ_{10}/Γ_2
VALUE	$\text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

0.111±0.064 AMSLER 04B CBAR 0 $\bar{p}p$

$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$	Γ_3/Γ_1
VALUE	$\text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~0.15 1 BERTIN 95 OBLX 0 $\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

~0.8 500 1 DUCH 89 ASTE $\bar{p}p \rightarrow \pi^+\pi^-K^\pm\pi^\mp K^0$

~0.75 1 REEVES 86 SPEC 6.6 $p\bar{p} \rightarrow K\bar{K}\pi X$

1 Assuming that the $a_0(980)$ decays only into $K\bar{K}$.

$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$	Γ_3/Γ_2
VALUE	$\text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29±0.10 ABELE 98E CBAR 0 $p\bar{p} \rightarrow \eta\pi^0\pi^0\pi^0$

0.19±0.04 2200 1 ALDE 97B GAM4 100 $\pi^-\rho \rightarrow \eta\pi^0\pi^0\eta$

0.56±0.04±0.03 1 AMSLER 95F CBAR 0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$

1 Assuming that the $a_0(980)$ decays only into $\eta\pi$.

NODE=M027215;NODE=M027

DESIG=2;OUR EST; \rightarrow UNCHECKED ←
 DESIG=5;OUR EST; \rightarrow UNCHECKED ←
 DESIG=4;OUR EST; \rightarrow UNCHECKED ←
 DESIG=9;OUR EST; \rightarrow UNCHECKED ←
 DESIG=15
 DESIG=10;OUR EST; \rightarrow UNCHECKED ←
 DESIG=6;OUR EST; \rightarrow UNCHECKED ←
 DESIG=12
 DESIG=7
 DESIG=8;OUR EST; \rightarrow UNCHECKED ←
 DESIG=13
 DESIG=11;OUR EST; \rightarrow UNCHECKED ←

NODE=M027220

NODE=M027G3
NODE=M027G3

NODE=M027G3;LINKAGE=AH
NODE=M027G3;LINKAGE=B3

NODE=M027G5
NODE=M027G5

NODE=M027G8
NODE=M027G8

NODE=M027225

NODE=M027R3
NODE=M027R3

NODE=M027R3;LINKAGE=AM

NODE=M027R12
NODE=M027R12

NODE=M027R4
NODE=M027R4

NODE=M027R4;LINKAGE=C

NODE=M027R2
NODE=M027R2

NODE=M027R2;LINKAGE=A

$\Gamma(a_0(980)\pi)/\Gamma(\eta(\pi\pi)S\text{-wave})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.91±0.12		ANISOVICH 01	SPEC	0.0 $\bar{p}p \rightarrow \eta\pi^+\pi^-\pi^+\pi^-$
0.15±0.04	9082	¹ MANAK 00A	MPS	18 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
0.70±0.12±0.20		² BAI 99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Statistical error only.² Assuming that the $a_0(980)$ decays only into $\eta\pi$. Γ_3/Γ_4 NODE=M027R9
NODE=M027R9 $\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0152±0.0038	¹ COFFMAN 90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
¹ Using $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi) = 4.2 \times 10^{-3}$ and $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\gamma\rho^0) = 6.4 \times 10^{-5}$.			

 Γ_{10}/Γ_1

NODE=M027R;LINKAGE=K3

NODE=M027R9;LINKAGE=BK

 $\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.78 × 10⁻³	90	¹ ABLIKIM 180	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
¹ Using results from BAI 00D.				

 Γ_9/Γ_1 NODE=M027R02
NODE=M027R02 $\Gamma(\eta(\pi\pi)S\text{-wave})/\Gamma(\eta\pi\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.81±0.04	2200	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta\pi^0\pi^0 n$

 Γ_4/Γ_2 NODE=M027R8
NODE=M027R8 $\Gamma(f_0(980)\eta)/\Gamma(\eta\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.32±0.07	¹ ANISOVICH 00	SPEC	0.9–1.2 $\bar{p}p \rightarrow \eta\pi^0$
¹ Using preliminary Crystal Barrel data.			

 Γ_6/Γ_2 NODE=M027R10
NODE=M027R10 $\Gamma(f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM 17AJ	BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$
¹ ABLIKIM 17AJ reports $B(\psi(2S) \rightarrow \gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0) < 5.0 \times 10^{-7}$.			

 Γ_5/Γ NODE=M027R00
NODE=M027R00 $\Gamma(\rho\rho)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.58	99.85	1,2 AMSLER	04B CBAR	0 $\bar{p}p$

¹ Assuming that the $\eta(1405)$ decays are saturated by the $\pi\pi\eta$, $K\bar{K}\pi$ and $\rho\rho$ modes.² Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$. Γ_8/Γ NODE=M027R13
NODE=M027R13 $\Gamma(K^*(892)K)/\Gamma(a_0(980)\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

 Γ_{12}/Γ_3 NODE=M027R13;LINKAGE=AM
NODE=M027R13;LINKAGE=AS

0.084±0.024	¹ ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
-------------	--------------------	----------	--

¹ Statistical error only. $\Gamma(\phi\gamma)/\Gamma(\rho^0\gamma)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

 Γ_{11}/Γ_{10} NODE=M027R14
NODE=M027R14

0.09±0.03	¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
0.13±0.04	² ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$

OCCUR=2

<0.77	95	³ BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma K^+ K^-$
-------	----	------------------	----------	---

¹ Constructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18I reports the inverse as 11.10 ± 3.5 .² Destructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18I reports the inverse as 7.53 ± 2.49 .³ Calculated by us from $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \phi\gamma\gamma) < 0.82 \times 10^{-4}$ and $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \rho^0\gamma\gamma) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$.

NODE=M027R14;LINKAGE=A

NODE=M027R14;LINKAGE=B

NODE=M027R14;LINKAGE=BA


$\eta(1405)$ REFERENCES

NODE=M027

ABLIKIM 18I	19BA PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60024
ABLIKIM 18O	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM 18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ABLIKIM 17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58322
ABLIKIM 13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55386
ABLIKIM 12E	PRl 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
ABLIKIM 11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53931
ABLIKIM 08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
AHOHE 05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)	REFID=50764
AMSLER 04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51079
BAI 04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
NICHITIU 02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
ACCIARRI 01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS 01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
ANISOVICH 01	NP A690 567	A.V. Anisovich <i>et al.</i>		REFID=48308
ANISOVICH 00	PL B472 168	A.V. Anisovich <i>et al.</i>		REFID=47429
BAI 00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
MANAK 00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
BAI 99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
CICALO 99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)	REFID=47394
ABELE 98E	NP B514 45	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46314
BAI 98C	PL B440 217	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46337
ALDE 97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
Translated from YAF 60 458.				
BERTIN 97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
AMSLER 95F	PL B358 389	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44613
BERTIN 95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=44614
BUGG 95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
AUGUSTIN 92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON 92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
FUKUI 91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
AUGUSTIN 90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BAI 90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
COFFMAN 90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
BISELLO 89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
DUCH 89	ZPHY C45 223	K.D. Duch <i>et al.</i>	(ASTERIX Collab.)	REFID=41016
RATH 89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)	REFID=40924
BIRMAN 88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD)	REFID=40568
ANDO 86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)	REFID=20891
REEVES 86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+)	REFID=20936
CHUNG 85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+)	REFID=20934
ALTHOFF 84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=20305
EDWARDS 83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
EDWARDS 82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
Also	PRL 50 219	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21315
SCHARRE 80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
FOSTER 68B	NP B8 174	M. Foster <i>et al.</i>	(CERN, CDEF)	REFID=21179
BAILLON 67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)	REFID=20407

$h_1(1415)$ $I^G(J^{PC}) = 0^-(1^{+-})$ was $h_1(1380)$ **$h_1(1415)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1416± 8 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
1423± 2.1±7.3	2.2k	1 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$
1412± 4 ±8		1 ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1,2} \rightarrow$ $\gamma \phi(h_1 \rightarrow K^* \bar{K})$
1440±60		ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$
1380±20		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

1 Final states $K^+ K^- \pi^0$ and $K_S^0 K^\pm \pi^\mp$. $h_1(1415)$ MASS (MeV) **$h_1(1415)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
90 ±15 OUR AVERAGE				
90.3± 9.8±17.5	2.2k	1 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$
84 ±12 ±40		1 ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1,2} \rightarrow$ $\gamma \phi(h_1 \rightarrow K^* \bar{K})$
170 ±80		ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$
80 ±30		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

1 Final states $K^+ K^- \pi^0$ and $K_S^0 K^\pm \pi^\mp$.

NODE=M109M

NODE=M109M

NODE=M109M;LINKAGE=A

 $h_1(1415)$ DECAY MODES

Mode
$\Gamma_1 K \bar{K}^*(892) + \text{c.c.}$

NODE=M109W

NODE=M109W

NODE=M109W;LINKAGE=A

NODE=M109215;NODE=M109

DESIG=1

NODE=M109

REFID=59456
REFID=56778
REFID=45765
REFID=40282 **$h_1(1415)$ REFERENCES**

ABLIKIM	18AB PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABELE	97H PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ASTON	88C PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

$a_1(1420)$ $I^G(J^{PC}) = 1^-(1^{++})$

OMITTED FROM SUMMARY TABLE

 $a_1(1420)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1411$^{+4}_{-5}$	46M	¹ AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1414 $^{+15}_{-13}$	2,3 ADOLPH	15C	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1 Statistical error negligible. 2 Using the isobar model and partial-wave analysis with 88 waves. 3 Superseded by AGHASYAN 2018B.				

NODE=M230M

NODE=M230M

NODE=M230M;LINKAGE=C
NODE=M230M;LINKAGE=A
NODE=M230M;LINKAGE=B **$a_1(1420)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
161$^{+11}_{-14}$	46M	¹ AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
153 $^{+8}_{-23}$	2,3 ADOLPH	15C	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1 Statistical error negligible. 2 Using the isobar model and partial-wave analysis with 88 waves. 3 Superseded by AGHASYAN 2018B.				

NODE=M230W

NODE=M230W

NODE=M230W;LINKAGE=D
NODE=M230W;LINKAGE=A
NODE=M230W;LINKAGE=C **$a_1(1420)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 f_0(980)\pi$	seen

NODE=M230215;NODE=M230

DESIG=1

 $a_1(1420)$ BRANCHING RATIOS

$\Gamma(f_0(980)\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	¹ ADOLPH	15C	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1 Using the isobar model and partial-wave analysis with 88 waves.				

NODE=M230220

NODE=M230R01
NODE=M230R01

NODE=M230R01;LINKAGE=A

 $a_1(1420)$ REFERENCES

AGHASYAN ADOLPH	18B 15C	PR D98 092003 PRL 115 082001	M. Aghasyan <i>et al.</i> C. Adolph <i>et al.</i>	(COMPASS Collab.) (COMPASS Collab.)
--------------------	------------	---------------------------------	--	--

NODE=M230

REFID=59471
REFID=56790

NODE=M006

 $f_1(1420)$ $I^G(J^{PC}) = 0^+(1^{++})$ See the minireview under $\eta(1405)$. **$f_1(1420)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1426.3 ± 0.9 OUR AVERAGE				
1434 ± 5 ± 5	133	1 ACHARD	07 L3	Error includes scale factor of 1.1. 183–209 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1426 ± 6	711	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
1420 ± 14	3651	NICHITIU	02 OBLX	0 $\bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$
1428 ± 4 ± 2	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+K^-\pi^0 n$
1426 ± 1		BARBERIS	97C OMEG	450 $p\bar{p} \rightarrow ppK_S^0 K^\pm \pi^\mp$
1425 ± 8		BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
1430 ± 4		2 ARMSTRONG	92E OMEG	85,300 $\pi^+ p, p\bar{p} \rightarrow \pi^+ p, pp(K\bar{K}\pi)$
1462 ± 20		3 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
+ 7 + 3 - 6 - 2	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1425 ± 10	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
1442 ± 5 + 10 - 17	111	BECKER	87 MRK3	$e^+e^-, \omega K\bar{K}\pi$
1423 ± 4		GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
1417 ± 13	13	AIHARA	86C TPC	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
1422 ± 3		CHAUVAT	84 SPEC	ISR 31.5 $p\bar{p}$
1440 ± 10		4 BROMBERG	80 SPEC	100 $\pi^- p \rightarrow K\bar{K}\pi X$
1426 ± 6	221	DIONISI	80 HBC	4 $\pi^- p \rightarrow K\bar{K}\pi n$
1420 ± 20		DAHL	67 HBC	1.6–4.2 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1430.8 ± 0.9		5 SOSA	99 SPEC	$p\bar{p} \rightarrow p_{slow} (K_S^0 K^+\pi^-) p_{fast}$
1433.4 ± 0.8		5 SOSA	99 SPEC	$p\bar{p} \rightarrow p_{slow} (K_S^0 K^-\pi^+) p_{fast}$
1435 ± 9		PROKOSHKIN	97B GAM4	100 $\pi^- p \rightarrow \eta\pi^0\pi^0 n$
1429 ± 3	389	ARMSTRONG	89 OMEG	300 $p\bar{p} \rightarrow K\bar{K}\pi p\bar{p}$
1425 ± 2	1520	ARMSTRONG	84 OMEG	85 $\pi^+ p, p\bar{p} \rightarrow (\pi^+, p)(K\bar{K}\pi)p$
~ 1420		BITYUKOV	84 SPEC	32 $K^-\bar{p} \rightarrow K^+K^-\pi^0 Y$

¹ From a fit with a width fixed at 55 MeV.² This result supersedes ARMSTRONG 84, ARMSTRONG 89.³ From fit to the $K^*(892)K 1^{++}$ partial wave.⁴ Mass error increased to account for $a_0(980)$ mass cut uncertainties.⁵ No systematic error given.

NODE=M006M2;LINKAGE=CH

NODE=M006M2;LINKAGE=C

NODE=M006M2;LINKAGE=B

NODE=M006M2;LINKAGE=A

NODE=M006M2;LINKAGE=N1

 $f_1(1420)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
54.5 ± 2.6 OUR AVERAGE				
51 ± 14	711	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
61 ± 8	3651	NICHITIU	02 OBLX	0 $\bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$
38 ± 9 ± 6	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+K^-\pi^0 n$
58 ± 4		BARBERIS	97C OMEG	450 $p\bar{p} \rightarrow ppK_S^0 K^\pm \pi^\mp$
45 ± 10		BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
58 ± 10		6 ARMSTRONG	92E OMEG	85,300 $\pi^+ p, p\bar{p} \rightarrow \pi^+ p, pp(K\bar{K}\pi)$

NODE=M006W

NODE=M006W

129	± 41		7 AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
68	$+29$	$+8$	-18	1100	BAI	90C MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
42	± 22			17	BEHREND	89 CELL $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
40	$+17$	± 5		111	BECKER	87 MRK3 $e^+ e^- \rightarrow \omega K\bar{K}\pi$
35	$+47$		-20	13	AIHARA	86C TPC $e^+ e^- \rightarrow e^+ e^- K\bar{K}\pi$
47	± 10				CHAUVAT	84 SPEC ISR 31.5 pp
62	± 14				BROMBERG	80 SPEC $100 \pi^- p \rightarrow K\bar{K}\pi X$
40	± 15			221	DIONISI	80 HBC $4 \pi^- p \rightarrow K\bar{K}\pi n$
60	± 20				DAHL	67 HBC $1.6-4.2 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
68.7 \pm 2.9				8 SOSA	99 SPEC	$pp \rightarrow p_{slow}$ $(K_S^0 K^+ \pi^-) p_{fast}$
58.8 \pm 3.3				8 SOSA	99 SPEC	$pp \rightarrow p_{slow}$ $(K_S^0 K^- \pi^+) p_{fast}$
90 ± 25				PROKOSHKIN 97B	GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
58 ± 8			389	ARMSTRONG 89	OMEG	$300 pp \rightarrow K\bar{K}\pi pp$
62 ± 5			1520	ARMSTRONG 84	OMEG	$85 \pi^+ p, pp \rightarrow (\pi^+, p)(K\bar{K}\pi)p$
~ 50				BITYUKOV 84	SPEC	$32 K^- p \rightarrow K^+ K^- \pi^0 Y$

⁶This result supersedes ARMSTRONG 84, ARMSTRONG 89.

⁷From fit to the $K^*(892)K^- \pi^+$ partial wave.

⁸No systematic error given.

OCCUR=2

NODE=M006W;LINKAGE=C

NODE=M006W;LINKAGE=B

NODE=M006W;LINKAGE=N1

NODE=M006215;NODE=M006

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}\pi$	seen
$\Gamma_2 K\bar{K}^*(892) + c.c.$	seen
$\Gamma_3 \eta \pi \pi$	possibly seen
$\Gamma_4 a_0(980)\pi$	
$\Gamma_5 \pi \pi \rho$	
$\Gamma_6 4\pi$	
$\Gamma_7 \rho^0 \gamma$	
$\Gamma_8 \phi \gamma$	seen

$f_1(1420) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.9 \pm 0.4 OUR AVERAGE					
3.2 $\pm 0.6 \pm 0.7$		133	9,10 ACHARD	07 L3	$183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$
3.0 $\pm 0.9 \pm 0.7$			11,12 BEHREND	89 CELL	$e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm$
$2.3^{+1.0}_{-0.9} \pm 0.8$			HILL	89 JADE	$e^+ e^- \rightarrow e^+ e^- K^\pm K_S^0 \pi^\mp$
1.3 $\pm 0.5 \pm 0.3$			AIHARA	88B TPC	$e^+ e^- \rightarrow e^+ e^- K^\pm K_S^0 \pi^\mp$
1.6 $\pm 0.7 \pm 0.3$			11,13 GIDAL	87B MRK2	$e^+ e^- \rightarrow e^+ e^- K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.0 95 JENNI 83 MRK2 $e^+ e^- \rightarrow e^+ e^- K\bar{K}\pi$

⁹From a fit with a width fixed at 55 MeV.

¹⁰The form factor parameter from the fit is 926 ± 78 MeV.

¹¹Assume a ρ -pole form factor.

¹²A ϕ - pole form factor gives considerably smaller widths.

¹³Published value divided by 2.

DESIG=2;OUR EST; \rightarrow UNCHECKED
 DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=5;OUR EST; \rightarrow UNCHECKED
 DESIG=4
 DESIG=3
 DESIG=6
 DESIG=8
 DESIG=9;OUR EST; \rightarrow UNCHECKED

NODE=M006220

NODE=M006G2
 NODE=M006G2

NODE=M006G2;LINKAGE=CH
 NODE=M006G2;LINKAGE=CR
 NODE=M006G2;LINKAGE=A
 NODE=M006G2;LINKAGE=D
 NODE=M006G2;LINKAGE=B

$f_1(1420)$ BRANCHING RATIOS **$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(K\bar{K}\pi)$**

<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.76 ± 0.06		BROMBERG	80	SPEC 100 $\pi^- p \rightarrow K\bar{K}\pi X$	
0.86 ± 0.12		DIONISI	80	HBC 4 $\pi^- p \rightarrow K\bar{K}\pi n$	

 $\Gamma(\pi\pi\rho)/\Gamma(K\bar{K}\pi)$

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.3	95	CORDEN	78	OMEG 12–15 $\pi^- p$		
<2.0		DAHL	67	HBC 1.6–4.2 $\pi^- p$		

 $\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_1
<0.1						
<0.1	95	ARMSTRONG	91B	OMEG 300 $pp \rightarrow pp\eta\pi^+\pi^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.35 ± 0.75		KOPKE	89	MRK3 $J/\psi \rightarrow \omega\eta\pi\pi(K\bar{K}\pi)$		
<0.6	90	GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$		
<0.5	95	CORDEN	78	OMEG 12–15 $\pi^- p$		
1.5 ± 0.8		DEFOIX	72	HBC 0.7 $\bar{p}p \rightarrow 7\pi$		

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_3
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>0.1	90	PROKOSHKIN	97B	GAM4 100 $\pi^- p \rightarrow \eta\pi^0\pi^0n$		
not seen in either mode		ANDO	86	SPEC 8 $\pi^- p$		
not seen in either mode		CORDEN	78	OMEG 12–15 $\pi^- p$		
0.4 ± 0.2		DEFOIX	72	HBC 0.7 $\bar{p}p \rightarrow 7\pi$		

 $\Gamma(4\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.90	95	DIONISI	80	HBC 4 $\pi^- p$		

 $\Gamma(K\bar{K}\pi)/[\Gamma(K\bar{K}^*(892)+c.c.) + \Gamma(a_0(980)\pi)]$

<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_1/(\Gamma_2+\Gamma_4)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.65 ± 0.27	14	DIONISI	80	HBC 4 $\pi^- p$	

14 Calculated using $\Gamma(K\bar{K})/\Gamma(\eta\pi) = 0.24 \pm 0.07$ for $a_0(980)$ fractions. **$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$**

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_2
0.04 ± 0.01 ± 0.01						
		BARBERIS	98C	OMEG 450 $pp \rightarrow p_f f_1(1420) p_s$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	68	ARMSTRONG	84	OMEG 85 $\pi^+ p$
-------	----	-----------	----	-------------------

 $\Gamma(4\pi)/\Gamma(K\bar{K}\pi)$

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_6/Γ_1
<0.62						
<0.62	95	ARMSTRONG	89G	OMEG 85 $\pi p \rightarrow 4\pi X$		

 $\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ
<0.08						
<0.08	95	15 ARMSTRONG	92C	SPEC 300 $pp \rightarrow pp\pi^+\pi^-\gamma$		

15 Using the data on the $\bar{K}K\pi$ mode from ARMSTRONG 89. **$\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$**

<u>VALUE</u>	<u>CL%</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ_1
<0.02						
<0.02	95	BARBERIS	98C	OMEG 450 $pp \rightarrow p_f f_1(1420) p_s$		

NODE=M006225

NODE=M006R1

NODE=M006R1

NODE=M006R2

NODE=M006R2

NODE=M006R3

NODE=M006R3

NODE=M006R4

NODE=M006R4

NODE=M006R5

NODE=M006R5

NODE=M006R6

NODE=M006R6

NODE=M006R6;LINKAGE=C

NODE=M006R7

NODE=M006R7

NODE=M006R8

NODE=M006R8

NODE=M006R9

NODE=M006R9

NODE=M006R10

NODE=M006R10

$\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$ VALUE
0.003±0.001±0.001

DOCUMENT ID	TECN	COMMENT
BARBERIS 98C	OMEG	450 $p p \rightarrow p_f f_1(1420) p_s$

 Γ_8/Γ_1 NODE=M006R11
NODE=M006R11 $f_1(1420)$ REFERENCES

ACHARD 07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
ABDALLAH 03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
NICHITIU 02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ADAMS 01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
SOSA 99	PRL 83 913	M. Sosa <i>et al.</i>	
BARBERIS 98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN 97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
PROKOSHKIN 97B	PD 42 298	Yu.D. Prokoshkin, S.A. Sadovsky	
Translated from DANS 354 751.			
ARMSTRONG 92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ARMSTRONG 92E	ZPHY C56 29	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JPC
AUGUSTIN 92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
ARMSTRONG 91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BAI 90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
ARMSTRONG 89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+) JPC
ARMSTRONG 89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)
BEHREND 89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
HILL 89	ZPHY C42 355	P. Hill <i>et al.</i>	(JADE Collab.) JP
KOPKE 89	PRPL 174 67	L. Kopke <i>et al.</i>	(CERN)
AIHARA 88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
BECKER 87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.) JP
GIDAL 87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
GIDAL 87B	PRL 59 2016	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
AIHARA 86C	PRL 57 2500	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.) JP
ANDO 86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
ARMSTRONG 84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP
BITYUKOV 84	SJNP 39 735	S. Bityukov <i>et al.</i>	(SERP)
Translated from YAF 39 1165.			
CHAUVAT 84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)
JENNI 83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)
BROMBERG 80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)
DIONISI 80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+) IJP
CORDEN 78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
DEFOIX 72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)
DAHL 67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP
Also	PRL 14 1074	D.H. Miller <i>et al.</i>	(LRL, UCB)

NODE=M006

REFID=51698
REFID=49548
REFID=48848
REFID=49649
REFID=46937
REFID=46346
REFID=45759
REFID=45417
REFID=45549

REFID=42097
REFID=43173
REFID=41584
REFID=41862
REFID=41578
REFID=40729
REFID=40930
REFID=40732
REFID=40741
REFID=41863
REFID=40572
REFID=40015
REFID=40223
REFID=40224
REFID=21326
REFID=20891
REFID=20929
REFID=45856

REFID=20932
REFID=20304
REFID=20922
REFID=20924
REFID=20452
REFID=20435
REFID=20321
REFID=21291

 $\omega(1420)$

$I^G(J^{PC}) = 0^-(1^- -)$

 $\omega(1420)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
(1400–1450) OUR ESTIMATE				

• • • We do not use the following data for averages, fits, limits, etc. • • •

1418± 30± 10	824	¹ AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+ e^- \rightarrow \omega \eta$
1470± 50	13.1k	² AULCHENKO 15A	SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1382± 23± 70		AUBERT 07AU	BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
1350± 20± 20		AUBERT,B 04N	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
1400± 50± 130	1.2M	³ ACHASOV 03D	RVUE	0.44–2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1450± 10		⁴ HENNER 02	RVUE	1.2–2.0 $e^+ e^- \rightarrow \rho \pi, \omega \pi \pi$
1373± 70	177	⁵ AKHMETSHIN 00D	CMD2	1.2–1.38 $e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1370± 25	5095	ANISOVICH 00H	SPEC	0.0 $p\bar{p} \rightarrow \omega \pi^0 \pi^0 \pi^0$
1400 ⁺¹⁰⁰ ₋₂₀₀		⁶ ACHASOV 98H	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
~ 1400		⁷ ACHASOV 98H	RVUE	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
~ 1460		⁸ ACHASOV 98H	RVUE	$e^+ e^- \rightarrow K^+ K^-$
1440± 70		⁹ CLEGG 94	RVUE	
1419± 31	315	¹⁰ ANTONELLI 92	DM2	1.34–2.4 $e^+ e^- \rightarrow \rho \pi$

NODE=M125M

NODE=M125M
→ UNCHECKED ←OCCUR=2
OCCUR=3

- 1 From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.
- 2 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.
- 3 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- 4 Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.
- 5 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
- 6 Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.
- 7 Using the data from ANTONELLI 92.
- 8 Using the data from IVANOV 81 and BISELLO 88B.
- 9 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- 10 From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed $(+, -, +)$ phases.

$\omega(1420)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
(180-250) OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
104 ± 35 ± 10	824	1 AKHMETSHIN 17A	CMD3	1.4–2.0 $e^+ e^- \rightarrow \omega \eta$
880 ± 170	13.1k	2 AULCHENKO 15A	SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
480 ± 180		3 ACHASOV 10D	SND	1.075–2.0 $e^+ e^- \rightarrow \pi^0 \gamma$
130 ± 50 ± 100		AUBERT 07AU BABR	10.6	$e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
450 ± 70 ± 70		AUBERT,B 04N BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
870 ± 500 ± 450	1.2M	4 ACHASOV 03D	RVUE	0.44–2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
199 ± 15		5 HENNER 02	RVUE	1.2–2.0 $e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$
188 ± 45	177	6 AKHMETSHIN 00D	CMD2	1.2–1.38 $e^+ e^- \rightarrow \omega\pi^+ \pi^-$
360 ± 100	5095	ANISOVICH 00H	SPEC	0.0 $p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
240 ± 70		7 CLEGG 94	RVUE	
174 ± 59	315	8 ANTONELLI 92	DM2	1.34–2.4 $e^+ e^- \rightarrow \rho\pi$

- 1 From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.
- 2 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.
- 3 From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.
- 4 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- 5 Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.
- 6 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
- 7 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- 8 From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed $(+, -, +)$ phases.

$\omega(1420)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi$	seen
$\Gamma_2 \omega\pi\pi$	seen
$\Gamma_3 \omega\eta$	
$\Gamma_4 b_1(1235)\pi$	seen
$\Gamma_5 e^+ e^-$	seen
$\Gamma_6 \pi^0\gamma$	

NODE=M125M;LINKAGE=F

NODE=M125M;LINKAGE=E

NODE=M125M;LINKAGE=VH

NODE=M125M;LINKAGE=AB

NODE=M125M;LINKAGE=KL

NODE=M125M;LINKAGE=L1

NODE=M125M;LINKAGE=L2

NODE=M125M;LINKAGE=L3

NODE=M125M;LINKAGE=AD

NODE=M125M;LINKAGE=B

NODE=M125W

NODE=M125W

→ UNCHECKED ←

NODE=M125W;LINKAGE=F

NODE=M125W;LINKAGE=E

NODE=M125W;LINKAGE=G

NODE=M125W;LINKAGE=VH

NODE=M125W;LINKAGE=AB

NODE=M125W;LINKAGE=KL

NODE=M125W;LINKAGE=AD

NODE=M125W;LINKAGE=B

NODE=M125215;NODE=M125

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=7

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=6

$\omega(1420) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma \times \Gamma_5/\Gamma$			
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.73 ± 0.08	13.1k	1 AULCHENKO	15A SND	$1.05-1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.82 ± 0.05 ± 0.06		AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
0.65 ± 0.13 ± 0.21	1.2M	2,3 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.625 ± 0.160		4,5 CLEGG	94 RVUE	
0.466 ± 0.178		6,7 ANTONELLI	92 DM2	$1.34-2.4 e^+e^- \rightarrow \rho\pi$

1 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

2 Calculated by us from the cross section at the peak.

3 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

4 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

5 From the partial and leptonic width given by the authors.

6 From a fit to two Breit-Wigner functions interfering between them and with the ω,ϕ tails with fixed $(+, -, +)$ phases.

7 From the product of the leptonic width and partial branching ratio given by the authors.

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_5/\Gamma$

<u>VALUE (units 10^{-8})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.7 ± 5.7	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1.9 ± 1.9	1 AKHMETSHIN 00D	CMD2	$1.2-2.4 e^+e^- \rightarrow \omega\pi^+\pi^-$

1 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

 $\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_5/\Gamma$

<u>VALUE (units 10^{-8})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ± 1.0		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
5.0 $\pm 2.6 \pm 0.3$	824	1 AKHMETSHIN 17A	CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$
1.6 ± 0.9	898	2 ACHASOV	16B SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$

1 From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1420) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1420) \rightarrow e^+e^-) = 5.3 \pm 1.6$ eV.

2 From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass and the width of $\omega(1420)$ are fixed to the 2014 edition (PDG 14) of this review.

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_5/\Gamma$

<u>VALUE (units 10^{-8})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.23 ± 0.14		1 ACHASOV	10D SND	$1.075-2.0 e^+e^- \rightarrow \pi^0\gamma$
2.03 ± 0.70		2 AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$

1 From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.

2 Using 1420 MeV and 220 MeV for the $\omega(1420)$ mass and width.

 $\omega(1420)$ BRANCHING RATIOS

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$	Γ_2/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.301 ± 0.029		1 HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
possibly seen		AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$

NODE=M125230

NODE=M125G3

NODE=M125G3

NODE=M125G3;LINKAGE=A

NODE=M125G;LINKAGE=AW

NODE=M125G;LINKAGE=VH

NODE=M125G;LINKAGE=AD

NODE=M125G;LINKAGE=SE

NODE=M125G;LINKAGE=A

NODE=M125G;LINKAGE=ES

NODE=M125G4

NODE=M125G4

NODE=M125G;LINKAGE=KL

NODE=M125G6

NODE=M125G6

NODE=M125G6;LINKAGE=B

NODE=M125G6;LINKAGE=A

NODE=M125G5

NODE=M125G5

NODE=M125G5;LINKAGE=A

NODE=M125G5;LINKAGE=AK

NODE=M125225

NODE=M125R2

NODE=M125R2

$\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_4
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.60 \pm 0.16	5095	ANISOVICH	00H SPEC	0.0 $p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$	

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.699 \pm 0.029	¹ HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$	

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
~ 6.6	1.2M	^{2,3} ACHASOV	03D RVUE	$0.44\text{--}2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
23 \pm 1		¹ HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$	

1 Assuming that the $\omega(1420)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

2 Calculated by us from the cross section at the peak.

3 Assuming that the $\omega(1420)$ decays into $\rho\pi$ only.

$\omega(1420)$ REFERENCES

ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
		Translated from ZETF 148 34.		
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>	
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORsay)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)

NODE=M125R1
NODE=M125R1

NODE=M125R3
NODE=M125R3

NODE=M125R4
NODE=M125R4

NODE=M125R;LINKAGE=AC
NODE=M125R;LINKAGE=AW
NODE=M125R;LINKAGE=GS

NODE=M125

REFID=59887
REFID=58239
REFID=57537
REFID=56843

REFID=55687
REFID=59494
REFID=52049
REFID=50330
REFID=50184
REFID=49577
REFID=48815
REFID=49177
REFID=48311
REFID=47935
REFID=47948
REFID=47391
REFID=46323
REFID=44081
REFID=43168
REFID=41369
REFID=40581
REFID=40280

REFID=21586
REFID=20553

$f_2(1430)$ $I^G(J^{PC}) = 0^+(2^{++})$

OMITTED FROM SUMMARY TABLE

This entry lists nearby peaks observed in the D wave of the $K\bar{K}$ and $\pi^+\pi^-$ systems. Needs confirmation.

 $f_2(1430)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1430 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1453 ± 4	1 VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1421 ± 5	AUGUSTIN 87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1480 ± 50	AKESSON 86	SPEC	$p p \rightarrow p p \pi^+ \pi^-$
1436 $^{+26}_{-16}$	DAUM 84	CNTR	17–18 $\pi^- p \rightarrow K^+ K^- n$
1412 ± 3	DAUM 84	CNTR	63 $\pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
1439 $^{+5}_{-6}$	2 BEUSCH 67	OSPK	5,7,12 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1 $J^{PC} = 0^{++}$ or 2^{++} .			
2 Not seen by WETZEL 76.			

NODE=M066

NODE=M066M1

NODE=M066M1
→ UNCHECKED ←

OCCUR=2

NODE=M066M;LINKAGE=AC
NODE=M066M;LINKAGE=C

NODE=M066W1

NODE=M066W1

OCCUR=2

NODE=M066W;LINKAGE=AC
NODE=M066W;LINKAGE=C

NODE=M066215;NODE=M066

DESIG=1

DESIG=2

NODE=M066

REFID=48571

REFID=40268

REFID=21123

REFID=21372

REFID=20362

REFID=20320

 $f_2(1430)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
13 ± 5	3 VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
30 ± 9	AUGUSTIN 87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
150 ± 50	AKESSON 86	SPEC	$p p \rightarrow p p \pi^+ \pi^-$
81 $^{+56}_{-29}$	DAUM 84	CNTR	17–18 $\pi^- p \rightarrow K^+ K^- n$
14 ± 6	DAUM 84	CNTR	63 $\pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
43 $^{+17}_{-18}$	4 BEUSCH 67	OSPK	5,7,12 $\pi^- p \rightarrow K_S^0 K_S^0 n$
3 $J^{PC} = 0^{++}$ or 2^{++} .			
4 Not seen by WETZEL 76.			

 $f_2(1430)$ DECAY MODES

Mode
$\Gamma_1 \quad K\bar{K}$
$\Gamma_2 \quad \pi\pi$

 $f_2(1430)$ REFERENCES

VLADIMIRSK... 01	PAN 64 1895	V.V. Vladmirsky <i>et al.</i>
	Translated from YAF 64 1979.	
AUGUSTIN 87	ZPHY C36 369	J.E. Augustin <i>et al.</i>
AKESSON 86	NP B264 154	T. Akesson <i>et al.</i>
DAUM 84	ZPHY C23 339	C. Daum <i>et al.</i>
WETZEL 76	NP B115 208	W. Wetzel <i>et al.</i>
BEUSCH 67	PL 25B 357	W. Beusch <i>et al.</i>

(LALO, CLER, FRAS+)

(Axial Field Spec. Collab.)

(AMST, CERN, CRAC, MPIM+) JP

(ETH, CERN, LOIC)

(ETH, CERN)

NODE=M149

 $a_0(1450)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

See minireview on scalar mesons under $f_0(500)$. **$a_0(1450)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1474 ±19 OUR AVERAGE				
1480 ±30		ABELE 98	CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
1470 ±25		1 AMSLER 95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1458 ±14	±15	190k 2 AAIJ	LHCb	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
1515 ±30		3 ANISOVICH 09	RVUE	0.0 $\bar{p}p$, πN
1316.8 ±0.7	+24.7	4 UEHARA 09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
1432 ±13	±25	5 BUGG 08A	RVUE	$\bar{p}p$
1477 ±10		6 UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
1441 ±40	-15	3 BAKER 03	SPEC	$\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
1303 ±16		7 BARGIOTTI 03	OBLX	$\bar{p}p$
1296 ±10		8 AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1565 ±30		8 ANISOVICH 98B	RVUE	Compilation
1290 ±10		9 BERTIN 98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
1450 ±40		AMSLER 94D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1410 ±25		ETKIN 82C	MPS	23 $\pi^- p \rightarrow n2K_S^0$
~ 1300		MARTIN 78	SPEC	10 $K^\pm p \rightarrow K_S^0 \pi p$
1255 ±5		10 CASON 76		

1 Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

2 Using a model with Gaussian constraints to the PDG averaged values .

3 From the pole position.

4 May be a different state.

5 Using data from AMSLER 94D, ABELE 98, and BAKER 03. Supersedes BUGG 94.

6 Statistical error only.

7 Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

8 T-matrix pole.

9 Not confirmed by BUGG 08A.

10 Isospin 0 not excluded.

NODE=M149

NODE=M149M

NODE=M149M

NODE=M149M;LINKAGE=AB

NODE=M149M;LINKAGE=A

NODE=M149M;LINKAGE=PP

NODE=M149M;LINKAGE=UE

NODE=M149M;LINKAGE=BU

NODE=M149M;LINKAGE=ST

NODE=M149M;LINKAGE=BG

NODE=M149M;LINKAGE=AN

NODE=M149M;LINKAGE=BE

NODE=M149M;LINKAGE=CC

NODE=M149W

NODE=M149W

 $a_0(1450)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
265 ±13 OUR AVERAGE				
265 ±15		ABELE 98	CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
265 ±30		1 AMSLER 95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
282 ±12	±13	190k 2 AAIJ	LHCb	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
230 ±36		3 ANISOVICH 09	RVUE	0.0 $\bar{p}p$, πN
65.0 ±2.1	+99.1	4 UEHARA 09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
- 5.4	-32.6	5 BUGG 08A	RVUE	$\bar{p}p$
196 ±10	±10	6 UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
267 ±11		35280 3 BAKER 03	SPEC	$\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
110 ±14		7 BARGIOTTI 03	OBLX	$\bar{p}p$
92 ±16		8 AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
81 ±21		8 ANISOVICH 98B	RVUE	Compilation
292 ±40		9 BERTIN 98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
80 ±5		AMSLER 94D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
270 ±40		ETKIN 82C	MPS	23 $\pi^- p \rightarrow n2K_S^0$
230 ±30		MARTIN 78	SPEC	10 $K^\pm p \rightarrow K_S^0 \pi p$
~ 250		10 CASON 76		
79 ±10				

NODE=M149W

- 1 Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.
 2 Using a model with Gaussian constraints to the PDG averaged values .
 3 From the pole position.
 4 May be a different state.
 5 Using data from AMSLER 94D, ABELE 98, and BAKER 03. Supersedes BUGG 94.
 6 Statistical error only.
 7 Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_L^0 \pi^\mp$.
 8 T-matrix pole.
 9 Not confirmed by BUGG 08A.
 10 Isospin 0 not excluded.

$a_0(1450)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi \eta$	0.093 ± 0.020
$\Gamma_2 \pi \eta'(958)$	0.033 ± 0.017
$\Gamma_3 K\bar{K}$	0.082 ± 0.028
$\Gamma_4 \omega \pi \pi$	DEFINED AS 1
$\Gamma_5 a_0(980) \pi \pi$	seen
$\Gamma_6 \gamma \gamma$	seen

$a_0(1450) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1 \Gamma_6/\Gamma$
<u>VALUE</u> (eV)	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
$432 \pm 6^{+1073}_{-256}$	¹ UEHARA 09A BELL $\gamma\gamma \rightarrow \pi^0 \eta$
$\bullet \bullet \bullet$ May be a different state.	

$a_0(1450)$ BRANCHING RATIOS

$\Gamma(\pi\eta'(958))/\Gamma(\pi\eta)$	Γ_2/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.35±0.16	
¹ ABELE 98 CBAR	$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
0.43 ± 0.19	ABELE 97C CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \eta'$
$\bullet \bullet \bullet$ Using $\pi^0 \eta$ from AMSLER 94D.	

$\Gamma(K\bar{K})/\Gamma(\pi\eta)$	Γ_3/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.88±0.23	
¹ ABELE 98 CBAR	$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
$\bullet \bullet \bullet$ Using $\pi^0 \eta$ from AMSLER 94D.	

$\Gamma(\omega\pi\pi)/\Gamma(\pi\eta)$	Γ_4/Γ_1
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
10.7±2.3	
35280	¹ BAKER 03 SPEC $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
$\bullet \bullet \bullet$ Using results on $\bar{p}p \rightarrow a_0(1450) \pi^0$, $a_0(1450) \rightarrow \eta \pi^0$ from ABELE 96C and assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.	

$\Gamma(a_0(980)\pi\pi)/\Gamma_{\text{total}}$	Γ_5/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	BUGG 08A RVUE $\bar{p}p$

$\Gamma(a_0(980)\pi\pi)/\Gamma(\pi\eta)$	Γ_5/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
≤ 4.3	ANISOVICH 01 RVUE 0 $\bar{p}p \rightarrow \eta 2\pi^+ 2\pi^-$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	¹ UEHARA 09A BELL $\gamma\gamma \rightarrow \pi^0 \eta$
$\bullet \bullet \bullet$ May be a different state.	

NODE=M149W;LINKAGE=AB
 NODE=M149W;LINKAGE=A
 NODE=M149W;LINKAGE=PP
 NODE=M149W;LINKAGE=UE
 NODE=M149W;LINKAGE=BU
 NODE=M149W;LINKAGE=ST
 NODE=M149W;LINKAGE=BG
 NODE=M149W;LINKAGE=AN
 NODE=M149W;LINKAGE=BE
 NODE=M149W;LINKAGE=CC

NODE=M149215;NODE=M149

DESIG=1
 DESIG=2
 DESIG=3
 DESIG=4
 DESIG=5
 DESIG=6

NODE=M149225

NODE=M149G01
 NODE=M149G01

NODE=M149G01;LINKAGE=UE

NODE=M149220
 NODE=M149R1
 NODE=M149R1

NODE=M149R1;LINKAGE=A

NODE=M149R2
 NODE=M149R2

NODE=M149R2;LINKAGE=A

NODE=M149R3
 NODE=M149R3

NODE=M149R;LINKAGE=PP

NODE=M149R01
 NODE=M149R01

NODE=M149R02
 NODE=M149R02

NODE=M149R03
 NODE=M149R03

NODE=M149R03;LINKAGE=UE

a₀(1450) REFERENCES

AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)
BUGG	08A	PR D78 074023	D.V. Bugg	(LOQM)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	01	NP A690 567	A.V. Anisovich <i>et al.</i>	
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
Translated from UFN 168 481.				
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(IGJPC)
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)
CASON	76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL)

NODE=M149

REFID=57273
REFID=52719
REFID=53002
REFID=52578
REFID=51063
REFID=49414
REFID=49217
REFID=48580
REFID=48308
REFID=45863
REFID=46331

REFID=46351
REFID=45531
REFID=45076
REFID=44377
REFID=44440
REFID=44441
REFID=44093
REFID=44078
REFID=20391
REFID=22446
REFID=21064

NODE=M105

 $\rho(1450)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

See the related review(s):

 $\rho(1450)$ and $\rho(1700)$ **$\rho(1450)$ MASS** **$\rho(1450)$ MASS**

VALUE (MeV) DOCUMENT ID

1465±25 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values. **$\eta\rho^0$ MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1500±10	7.4k	1 ACHASOV	18 SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1497±14		2 AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1421±15		3 AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1470±20		ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1446±10		FUKUI	88 SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

1 From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

2 Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$.

3 Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

 $\omega\pi$ MODE

VALUE (MeV) DOCUMENT ID

• • • We do not use the following data for averages, fits, limits, etc. • • •

1510± 7	10.2k	1 ACHASOV	16D SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1544±22 ⁺¹¹ ₋₄₆	821	2 MATVIENKO	15 BELL	$\bar{B}^0 \rightarrow D^*+\omega\pi^-$
1491±19	7815	3 ACHASOV	13 SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1582±17±25	2382	4 AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1349±25 ⁺¹⁰ ₋₅	341	5 ALEXANDER	01B CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
1523±10		6 EDWARDS	00A CLE2	$\tau^- \rightarrow \omega\pi^-\nu_\tau$
1463±25		7 CLEGG	94 RVUE	
1250		8 ASTON	80C OMEG	20–70 $\gamma p \rightarrow \omega\pi^0 p$
1290±40		8 BARBER	80C SPEC	3–5 $\gamma p \rightarrow \omega\pi^0 p$

NODE=M105205

NODE=M105M0
NODE=M105M0

→ UNCHECKED ←

NODE=M105M1
NODE=M105M1

NODE=M105M1;LINKAGE=A

NODE=M105M;LINKAGE=SW

NODE=M105M1;LINKAGE=KL

NODE=M105M3
NODE=M105M3

OCCUR=2

- ¹ From a phenomenological model based on vector meson dominance with interfering $\rho(770)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.
- ² Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.
- ³ From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.
- ⁴ Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the $\omega\pi^0$ and $\pi^+\pi^-$ mass dependence of the total width. $\rho(1700)$ mass and width fixed at 1700 MeV and 240 MeV, respectively.
- ⁵ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming the $\omega\pi^-$ mass dependence for the total width.
- ⁶ Mass-independent width parameterization. $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
- ⁷ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.
- ⁸ Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.

4π MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1435 ± 40	ABELE 01B	CBAR	$0.0 \bar{p}n \rightarrow 2\pi^- 2\pi^0 \pi^+$
1350 ± 50	ACHASOV 97	RVUE	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1449 ± 4	¹ ARMSTRONG 89E	OMEG	$300 pp \rightarrow pp 2(\pi^+ \pi^-)$

¹ Not clear whether this observation has $I=1$ or 0.

ππ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1326.35 ± 3.46	¹ BARTOS 17	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$	
1342.31 ± 46.62	² BARTOS 17A	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$	
1373.83 ± 11.37	³ BARTOS 17A	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
1429 ± 41	20K ⁴ LEES 17C	BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$	
1350 ± 20	+20 -30 63.5k	⁵ ABRAMOWICZ 12	ZEUS	$ep \rightarrow e\pi^+ \pi^- p$
1493 ± 15	⁶ LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$	
1446 ± 7	±28 5.4M ^{7,8} FUJIKAWA 08	BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
1328 ± 15	⁹ SCHABEL 05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
1406 ± 15	87k ^{7,10} ANDERSON 00A	CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$	
~1368	¹¹ ABELE 99C	CBAR	$0.0 \bar{p}d \rightarrow \pi^+ \pi^- \pi^- p$	
1348 ± 33	BERTIN 98	OBLX	$0.05-0.405 \bar{n}p \rightarrow 2\pi^+ \pi^-$	
1411 ± 14	¹² ABELE 97	CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$	
1370 +90 -70	ACHASOV 97	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$	
1359 ± 40	¹⁰ BERTIN 97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
1282 ± 37	BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$	
1424 ± 25	BISELLO 89	DM2	$e^+ e^- \rightarrow \pi^+ \pi^-$	
1265.5 ± 75.3	DUBNICKA 89	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$	
1292 ± 17	¹³ KURDADZE 83	OLYA	$0.64-1.4 e^+ e^- \rightarrow \pi^+ \pi^-$	

- ¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
- ² Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
- ³ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.
- ⁴ From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.
- ⁵ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- ⁶ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
- ⁷ From the GOUNARIS 68 parametrization of the pion form factor.
- ⁸ $|F_\pi(0)|^2$ fixed to 1.
- ⁹ From the combined fit of the τ^- data from ANDERSON 00A and SCHABEL 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. $\rho(1700)$ mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
- ¹⁰ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV, respectively.
- ¹¹ $\rho(1700)$ mass and width fixed at 1780 MeV and 275 MeV respectively.
- ¹² T-matrix pole.
- ¹³ Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

NODE=M105M3;LINKAGE=D

NODE=M105M3;LINKAGE=C

NODE=M105M3;LINKAGE=AC

NODE=M105M3;LINKAGE=HK

NODE=M105M3;LINKAGE=3Z

NODE=M105M;LINKAGE=E1

NODE=M105M3;LINKAGE=B

NODE=M105M3;LINKAGE=A

NODE=M105M6

NODE=M105M6

NODE=M105M6;LINKAGE=A

NODE=M105M5

NODE=M105M5

OCCUR=2

OCCUR=2

NODE=M105M5;LINKAGE=D

NODE=M105M5;LINKAGE=E

NODE=M105M5;LINKAGE=F

NODE=M105M5;LINKAGE=B

NODE=M105M5;LINKAGE=AB

NODE=M105M5;LINKAGE=LE

NODE=M105M;LINKAGE=1K

NODE=M105M5;LINKAGE=FU

NODE=M105M5;LINKAGE=SC

NODE=M105M5;LINKAGE=A

NODE=M105M5;LINKAGE=C5

NODE=M105M5;LINKAGE=QQ

NODE=M105M5;LINKAGE=KD

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1208 \pm 8 \pm 9	190k	1 AAIJ	16N LHCb	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$	
1422.8 \pm 6.5	27k	2 ABELE	99D CBAR	$\pm 0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$	

1 Using the GOUNARIS 68 parameterization with fixed width.
2 K-matrix pole. Isospin not determined, could be $\omega(1420)$.

NODE=M105M7
NODE=M105M7

 $K\bar{K}^*(892) + c.c.$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1505 \pm 19 \pm 7	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K\bar{K}^*(892)\gamma$

NODE=M105M7;LINKAGE=A
NODE=M105M7;LINKAGE=AN

NODE=M105M8
NODE=M105M8

 $m_{\rho(1450)^0} = m_{\rho(1450)^\pm}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-31.53 \pm 47.99	1 BARTOS	17A RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M105DM
NODE=M105DM

 $\rho(1450)$ WIDTH **$\rho(1450)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
400 \pm 60 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.			

• • • We do not use the following data for averages, fits, limits, etc. • • •

480 \pm 180	1 ACHASOV	10D SND	$1.075-2.0 e^+ e^- \rightarrow \pi^0 \gamma$
1 From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.			

NODE=M105DM;LINKAGE=A

NODE=M105210

NODE=M105W0
NODE=M105W0
→ UNCHECKED ←

NODE=M105W0;LINKAGE=A

 $\eta\rho^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

280 \pm 20	7.4k	1 ACHASOV	18 SND	$1.22-2.00 e^+ e^- \rightarrow \eta\pi^+\pi^-$
226 \pm 44		2 AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
211 \pm 31		3 AKHMETSHIN 00D	CMD2	$e^+ e^- \rightarrow \eta\pi^+\pi^-$
230 \pm 30		ANTONELLI	88 DM2	$e^+ e^- \rightarrow \eta\pi^+\pi^-$
60 \pm 15		FUKUI	88 SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$

1 From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.
2 Using the data of AKHMETSHIN 01B on $e^+ e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+ e^- \rightarrow \eta\pi^+\pi^-$.
3 Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

NODE=M105W1;LINKAGE=A

NODE=M105W;LINKAGE=SW

NODE=M105W1;LINKAGE=KL

 $\omega\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

440 \pm 40	10.2k	1 ACHASOV	16D SND	$1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
303 \pm 31 \pm 69	821	2 MATVIENKO	15 BELL	$\bar{B}^0 \rightarrow D^*+\omega\pi^-$
429 \pm 42 \pm 10	2382	3 AKHMETSHIN 03B	CMD2	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
547 \pm 86 \pm 46	341	4 ALEXANDER	01B CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
400 \pm 35		5 EDWARDS	00A CLE2	$\tau^- \rightarrow \omega\pi^-\nu_\tau$
311 \pm 62		6 CLEGG	94 RVUE	
300		7 ASTON	80C OMEG	$20-70 \gamma p \rightarrow \omega\pi^0 p$
320 \pm 100		7 BARBER	80C SPEC	$3-5 \gamma p \rightarrow \omega\pi^0 p$

NODE=M105W3
NODE=M105W3

OCCUR=3

- ¹ From a phenomenological model based on vector meson dominance with interfering $\rho(770)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.
- ² Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.
- ³ Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the $\omega\pi^0$ and $\pi^+\pi^-$ mass dependence of the total width. $\rho(1700)$ mass and width fixed at 1700 MeV and 240 MeV, respectively.
- ⁴ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming the $\omega\pi^-$ mass dependence for the total width.
- ⁵ Mass-independent width parameterization. $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.
- ⁶ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.
- ⁷ Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.

4 π MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
325 \pm 100	ABELE	01B CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0 \pi^+$

 $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
324.13 \pm 12.01		¹ BARTOS	17 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
492.17 \pm 138.38		² BARTOS	17A RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
340.87 \pm 23.84		³ BARTOS	17A RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
576 \pm 29	20K	⁴ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
460 \pm 30	$^{+40}_{-45}$ 63.5k	⁵ ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
427 \pm 31		⁶ LEES	12G BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
434 \pm 16	\pm 60 5.4M	^{7,8} FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
468 \pm 41		⁹ SCHael	05C ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
455 \pm 41	87k	^{7,10} ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
\sim 374		¹¹ ABELE	99C CBAR	$0.0 \bar{p}d \rightarrow \pi^+ \pi^- \pi^- p$
275 \pm 10		BERTIN	98 OBLX	$0.05-0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
343 \pm 20		¹² ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
310 \pm 40		¹⁰ BERTIN	97C OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
236 \pm 36		BERTIN	97D OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
269 \pm 31		BISELLO	89 DM2	$e^+ e^- \rightarrow \pi^+ \pi^-$
391 \pm 70		DUBNICKA	89 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
218 \pm 46		¹³ KURDADZE	83 OLYA	$0.64-1.4 e^+ e^- \rightarrow \pi^+ \pi^-$

- ¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
- ² Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
- ³ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.
- ⁴ From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.
- ⁵ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- ⁶ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
- ⁷ From the GOUNARIS 68 parametrization of the pion form factor.
- ⁸ $|F_\pi(0)|^2$ fixed to 1.
- ⁹ From the combined fit of the τ^- data from ANDERSON 00A and SCHael 05C and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. $\rho(1700)$ mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
- ¹⁰ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV, respectively.
- ¹¹ $\rho(1700)$ mass and width fixed at 1780 MeV and 275 MeV respectively.
- ¹² T-matrix pole.
- ¹³ Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

NODE=M105W3;LINKAGE=D

NODE=M105W3;LINKAGE=C

NODE=M105W3;LINKAGE=HK

NODE=M105W3;LINKAGE=3Z

NODE=M105W;LINKAGE=E1

NODE=M105W3;LINKAGE=B

NODE=M105W3;LINKAGE=A

NODE=M105W66

NODE=M105W66

NODE=M105W5

NODE=M105W5

OCCUR=2

NODE=M105W5;LINKAGE=C

NODE=M105W5;LINKAGE=D

NODE=M105W5;LINKAGE=E

NODE=M105W5;LINKAGE=B

NODE=M105W5;LINKAGE=AB

NODE=M105W5;LINKAGE=LE

NODE=M105W;LINKAGE=1K

NODE=M105W5;LINKAGE=FU

NODE=M105W5;LINKAGE=SC

NODE=M105W5;LINKAGE=A

NODE=M105W5;LINKAGE=C5

NODE=M105W5;LINKAGE=QQ

NODE=M105W5;LINKAGE=KD

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
410 ± 19 ± 35	190k	1 AAIJ	16N LHCb	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$	
146.5 ± 10.5	27k	2 ABELE	99D CBAR	$\pm 0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$	

1 Using the GOUNARIS 68 parameterization with fixed mass.
2 K-matrix pole. Isospin not determined, could be $\omega(1420)$.

 $K\bar{K}^*(892) + c.c.$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
418 $\pm 25 \pm 4$	AUBERT	08s BABR	$10.6 e^+ e^- \rightarrow K\bar{K}^*(892)\gamma$

$$\Gamma_{\rho(1450)^0} = \Gamma_{\rho(1450)^\pm}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
151.30 ± 140.42	1 BARTOS	17A RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

 $\rho(1450)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 \pi^+ \pi^-$	seen
$\Gamma_3 4\pi$	seen
$\Gamma_4 \omega\pi$	
$\Gamma_5 a_1(1260)\pi$	
$\Gamma_6 h_1(1170)\pi$	
$\Gamma_7 \pi(1300)\pi$	
$\Gamma_8 \rho\rho$	
$\Gamma_9 \rho(\pi\pi)s\text{-wave}$	
$\Gamma_{10} e^+ e^-$	seen
$\Gamma_{11} \eta\rho$	seen
$\Gamma_{12} a_2(1320)\pi$	not seen
$\Gamma_{13} K\bar{K}$	seen
$\Gamma_{14} K^+ K^-$	seen
$\Gamma_{15} K\bar{K}^*(892) + c.c.$	possibly seen
$\Gamma_{16} \pi^0\gamma$	
$\Gamma_{17} \eta\gamma$	seen
$\Gamma_{18} f_0(500)\gamma$	not seen
$\Gamma_{19} f_0(980)\gamma$	not seen
$\Gamma_{20} f_0(1370)\gamma$	not seen
$\Gamma_{21} f_2(1270)\gamma$	not seen

 $\rho(1450) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.12	1 DIEKMAN	88 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$	
$0.027^{+0.015}_{-0.010}$	2 KURDADZE	83 OLYA	$0.64-1.4 e^+ e^- \rightarrow \pi^+ \pi^-$	

1 Using total width = 235 MeV.

2 Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

NODE=M105W7

NODE=M105W7

NODE=M105W7;LINKAGE=A

NODE=M105W7;LINKAGE=AN

NODE=M105W8

NODE=M105W8

NODE=M105DW

NODE=M105DW

NODE=M105DW;LINKAGE=A

NODE=M105215;NODE=M105

DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=20;OUR EVAL; \rightarrow UNCHECKED
 DESIG=2;OUR EST; \rightarrow UNCHECKED
 DESIG=6
 DESIG=10
 DESIG=11
 DESIG=12
 DESIG=13
 DESIG=14
 DESIG=4;OUR EST; \rightarrow UNCHECKED
 DESIG=3
 DESIG=8;OUR EST; \rightarrow UNCHECKED
 DESIG=7;OUR EVAL; \rightarrow UNCHECKED
 DESIG=21;OUR EVAL; \rightarrow UNCHECKED
 DESIG=15;OUR EST; \rightarrow UNCHECKED
 DESIG=23
 DESIG=9
 DESIG=16;OUR EST; \rightarrow UNCHECKED
 DESIG=17;OUR EST; \rightarrow UNCHECKED
 DESIG=18;OUR EST; \rightarrow UNCHECKED
 DESIG=19;OUR EST; \rightarrow UNCHECKED

NODE=M105220

NODE=M105G3

NODE=M105G3

NODE=M105G3;LINKAGE=B

NODE=M105G3;LINKAGE=KD

$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{11}\Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
210±24±10	¹ LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-$	
74±20	² AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$	
91±19	ANTONELLI	88	DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$	

¹ Includes non-resonant contribution. The selected fit model includes three ρ excited states. Model uncertainty is 20%.

² Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

 $\Gamma(\eta\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{17}\Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

<16.4	¹ AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
2.2±0.5±0.3	² AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$

¹ From 2γ decay mode of η using 1465 MeV and 310 MeV for the $\rho(1450)$ mass and width. Recalculated by us.

² Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$. Recalculated by us using width of 226 MeV.

 $\Gamma(K\bar{K}^*(892)+\text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{15}\Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

127±15±6	AUBERT	085	BABR 10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma$
----------	--------	-----	--

 $\rho(1450) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$ $\Gamma(\omega\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_{10}/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma \times \Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

2.1±0.4	10.2k	¹ ACHASOV	16D	SND 1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$	
5.3±0.4	7815	² ACHASOV	13	SND 1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$	

¹ From a phenomenological model based on vector meson dominance with interfering $\rho(770)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

² From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

 $\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{11}/\Gamma \times \Gamma_{10}/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{11}/\Gamma \times \Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

7.3±0.3	7.4k	¹ ACHASOV	18	SND 1.22-2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$	
4.3 ^{+1.1} _{-0.9} ±0.2	4.9k	² AULCHENKO	15	SND 1.22-2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$	

¹ From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

² From a fit to the $e^+e^- \rightarrow \eta\pi^+\pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

 $\Gamma(f_0(500)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}/\Gamma \times \Gamma_{10}/\Gamma$

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{18}/\Gamma \times \Gamma_{10}/\Gamma$
<4.0	90	ACHASOV	11	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$	

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{16}/\Gamma \times \Gamma_{10}/\Gamma$

VALUE (units 10^{-9})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{16}/\Gamma \times \Gamma_{10}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

2.3±1.4	¹ ACHASOV	10D	SND 1.075-2.0 $e^+e^- \rightarrow \pi^0\gamma$	
---------	----------------------	-----	--	--

¹ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.

NODE=M105G4

NODE=M105G4

NODE=M105G4;LINKAGE=A

NODE=M105G4;LINKAGE=KL

NODE=M105G6

NODE=M105G6

NODE=M105G6;LINKAGE=AK

NODE=M105G;LINKAGE=SW

NODE=M105G8

NODE=M105G8

NODE=M105230

NODE=M105R05

NODE=M105R05

OCCUR=3

NODE=M105R05;LINKAGE=A

NODE=M105R00

NODE=M105R00

NODE=M105R00;LINKAGE=B

NODE=M105R00;LINKAGE=A

NODE=M105R01

NODE=M105R01

NODE=M105R17

NODE=M105R17

NODE=M105R17;LINKAGE=A

$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_{19}/\Gamma \times \Gamma_{10}/\Gamma$		
<u>VALUE</u> (units 10^{-9})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.6	90	ACHASOV	11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$\Gamma(f_0(1370)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_{20}/\Gamma \times \Gamma_{10}/\Gamma$		
<u>VALUE</u> (units 10^{-9})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.5	90	ACHASOV	11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$\Gamma(f_2(1270)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_{21}/\Gamma \times \Gamma_{10}/\Gamma$		
<u>VALUE</u> (units 10^{-9})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.8	90	¹ ACHASOV	11	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Using Breit-Wigner parametrization of the $\rho(1450)$ with mass and width of 1465 MeV and 400 MeV, respectively.

$\rho(1450)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma(4\pi)$			Γ_1/Γ_3		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.37 ± 0.10	^{1,2} ABELE	01B	CBAR	0.0	$\bar{p}n \rightarrow 5\pi$
¹ $\omega\pi$ not included. ² Using ABELE 97.					

$\Gamma(K^+K^-)/\Gamma(\pi^+\pi^-)$			Γ_{14}/Γ_2		
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
30.7 ± 8.4 ± 8.2	20K	¹ LEES	17C	BABR	$J/\psi \rightarrow h^+h^-\pi^0$

¹ From Dalitz plot analyses in isobar models.

$\Gamma(\omega\pi)/\Gamma_{\text{total}}$			Γ_4/Γ		
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
seen	821	¹ MATVIENKO	15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
seen	1.6k	ACHASOV	12	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
~0.21		CLEGG	94	RVUE	

¹ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.

$\Gamma(\pi\pi)/\Gamma(\omega\pi)$			Γ_1/Γ_4		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
~0.32	CLEGG	94	RVUE		

$\Gamma(\omega\pi)/\Gamma(4\pi)$			Γ_4/Γ_3		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.14	CLEGG	88	RVUE		

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$			Γ_5/Γ_3		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.27 ± 0.08	¹ ABELE	01B	CBAR	0.0	$\bar{p}n \rightarrow 5\pi$

¹ $\omega\pi$ not included.

$\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$			Γ_6/Γ_3		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.08 ± 0.04	¹ ABELE	01B	CBAR	0.0	$\bar{p}n \rightarrow 5\pi$

¹ $\omega\pi$ not included.

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$			Γ_7/Γ_3		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.37 ± 0.13	¹ ABELE	01B	CBAR	0.0	$\bar{p}n \rightarrow 5\pi$

¹ $\omega\pi$ not included.

NODE=M105R02
NODE=M105R02

NODE=M105R03
NODE=M105R03

NODE=M105R04
NODE=M105R04

NODE=M105225

NODE=M105R15
NODE=M105R15

NODE=M105R;LINKAGE=BL
NODE=M105R;LINKAGE=LK

NODE=M105R08
NODE=M105R08

NODE=M105R08;LINKAGE=A

NODE=M105R5
NODE=M105R5

NODE=M105R6
NODE=M105R6

NODE=M105R3
NODE=M105R3

NODE=M105R10
NODE=M105R10

NODE=M105R11
NODE=M105R11

NODE=M105R11;LINKAGE=BL

NODE=M105R12
NODE=M105R12

NODE=M105R12;LINKAGE=BL

$\Gamma(\rho\rho)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_3
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.11 ± 0.05	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$	NODE=M105R13 NODE=M105R13
$1 \omega\pi$ not included.				

 $\Gamma(\rho(\pi\pi)s\text{-wave})/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_3
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17 ± 0.09	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$	NODE=M105R14 NODE=M105R14
$1 \omega\pi$ not included.				

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
seen	35	¹ ACHASOV	14	SND 1.15–2.00 $e^+e^- \rightarrow \eta\gamma$	NODE=M105R2 NODE=M105R2
• • • We do not use the following data for averages, fits, limits, etc. • • •					

 <0.04

DONNACHIE 87B RVUE

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

 $\Gamma(\eta\rho)/\Gamma(\omega\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.081 ± 0.020	^{1,2} AULCHENKO 15	SND	$1.22\text{--}2.00 e^+e^- \rightarrow \eta\pi^+\pi^-$	NODE=M105R4 NODE=M105R4
~ 0.24	³ DONNACHIE 91	RVUE		
>2	FUKUI 91	SPEC	$8.95 \pi^- p \rightarrow \omega\pi^0 n$	
¹ From a fit to the $e^+e^- \rightarrow \eta\pi^+\pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.				
² Reports the inverse of the quoted value as 12.3 ± 3.1 .				
³ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.				

 $\Gamma(\pi\pi)/\Gamma(\eta\rho)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_{11}
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.3 ± 0.4	¹ AULCHENKO 15	SND	$1.22\text{--}2.00 e^+e^- \rightarrow \eta\pi^+\pi^-$	NODE=M105R07 NODE=M105R07
¹ From a fit to the $e^+e^- \rightarrow \eta\pi^+\pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.				

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AMELIN 00	VES	$37 \pi^- p \rightarrow \eta\pi^+\pi^- n$	NODE=M105R9 NODE=M105R9

 $\Gamma(K\bar{K})/\Gamma(\omega\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.08	¹ DONNACHIE 91	RVUE		NODE=M105R8 NODE=M105R8

¹ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

 $\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
possibly seen	COAN 04	CLEO	$\tau^- \rightarrow K^-\pi^-K^+\nu_\tau$	NODE=M105R16 NODE=M105R16

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
seen	35	¹ ACHASOV 14	SND	$1.15\text{--}2.00 e^+e^- \rightarrow \eta\gamma$	NODE=M105R06 NODE=M105R06
¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.					

$\rho(1450)$ REFERENCES

NODE=M105

ACHASOV	18	PR D97 012008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58785
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16D	PR D94 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57618
AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIEJKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55912
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
		Translated from ZETFP 94 796.			
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ACHASOV	11	JETP 113 75	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=16721
		Translated from ZETF 140 87.			
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
ALOISIO	05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
SCHAEL	05C	PRPL 421 191	S. Schael <i>et al.</i>	(ALEPH Collab.)	REFID=50845
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
ALEXANDER	01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48391
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)	REFID=47468
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE	99C	PL B450 275	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46916
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45622
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
BERTIN	97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45763
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=41752
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41632
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41581
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
CLEGG	88	ZPHY C40 313	A.B. Clegg, A. Donnachie	(MCHS, LANC)	REFID=40922
DIEMAN	88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40418
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
		Translated from ZETFP 37 613.			
ASTON	80C	PL 92B 211	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=20652
BARBER	80C	ZPHY C4 169	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)	REFID=20653
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

$\eta(1475)$

$$I^G(J^{PC}) = 0^+(0^-+)$$


See the $\eta(1405)$ and the related review on "Pseudoscalar and Pseudovector Mesons in the 1400 MeV Region."

$\eta(1475)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1475± 4 OUR AVERAGE				
1469±14±13	74	ACHARD	07 L3	Error includes scale factor of 1.4. See the ideogram below.
1460±12	3651	NICHITIU	02 OBLX	$e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
1485± 8± 5	20k	ADAMS	01B B852	$0 \bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0 n$
1500±10		CICALO	99 OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+\pi^-$
1464±10		BERTIN	97 OBLX	$0 \bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
1460±10		BERTIN	95 OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
1490 ^{+14 + 3} _{-8 - 16}	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1475± 4		RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1477± 7±13		¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
1565± 8 ^{+ 0} ₋₆₃		² ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
1421±14		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$

¹ From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded.

² Could also be the $\eta(1405)$.



$\eta(1475)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
90± 9 OUR AVERAGE				
67±18± 7	74	ACHARD	07 L3	Error includes scale factor of 1.6. See the ideogram below.
120±15	3651	NICHITIU	02 OBLX	$e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
98±18± 3	20k	ADAMS	01B B852	$0 \bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0 n$
100±20		CICALO	99 OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+\pi^-$
105±15		BERTIN	97 OBLX	$0 \bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
105±15		BERTIN	95 OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
54 ^{+37 + 13} _{-21 - 24}		BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
51±13		RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$

NODE=M175

NODE=M175

NODE=M175M5

NODE=M175M5

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M175M5;LINKAGE=B

NODE=M175M5;LINKAGE=A

NODE=M175W5

NODE=M175W5

OCCUR=2


OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$118 \pm 22 \pm 17$	¹ ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
$45^{+14}_{-13}{}^{+21}_{-28}$	² ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
63 ± 18	AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$

¹ From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded.

² Could also be the $\eta(1405)$.



$\eta(1475)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}\pi$	seen
$\Gamma_2 K\bar{K}^*(892) + \text{c.c.}$	seen
$\Gamma_3 a_0(980)\pi$	seen
$\Gamma_4 \gamma\gamma$	seen
$\Gamma_5 K_S^0 K_S^0 \eta$	possibly seen
$\Gamma_6 \gamma\phi(1020)$	possibly seen

$\eta(1475) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_4/\Gamma$				
VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.23 \pm 0.05 \pm 0.05$	74	¹ ACHARD	07	L3	$183\text{--}209 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.089	90	^{2,3} AHOHE	05	CLE2	$10.6 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
---------	----	----------------------	----	------	--

¹ Supersedes ACCIARRI 01G. Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895$.

² Using $\eta(1475)$ mass of 1481 MeV and width of 48 MeV. The upper limit increases to 0.140 keV if the world average value, 87 MeV, of the width is used.

³ Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

$\eta(1475)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K\bar{K}\pi)$	Γ_2/Γ_1			
VALUE	DOCUMENT ID	TECN	COMMENT	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.50 ± 0.10	¹ BAILLON	67	HBC	$0.0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
-------------	----------------------	----	-----	--

¹ Data could also refer to $\eta(1405)$.

OCCUR=2

NODE=M175W5;LINKAGE=B

NODE=M175W5;LINKAGE=A

NODE=M175215;NODE=M175

DESIG=2;OUR EST; \rightarrow UNCHECKED
DESIG=1;OUR EST; \rightarrow UNCHECKED
DESIG=4;OUR EST; \rightarrow UNCHECKED
DESIG=7;OUR EST; \rightarrow UNCHECKED
DESIG=8;OUR EVAL; \rightarrow UNCHECKED
DESIG=9

NODE=M175220

NODE=M175G2

NODE=M175G2

NODE=M175G2;LINKAGE=CH

NODE=M175G2;LINKAGE=AH

NODE=M175G2;LINKAGE=B3

NODE=M175225

NODE=M175R1

NODE=M175R1

NODE=M175R;LINKAGE=BL

$$\Gamma(K\bar{K}^*(892)+c.c.)/[\Gamma(K\bar{K}^*(892)+c.c.) + \Gamma(a_0(980)\pi)] \quad \Gamma_2/(\Gamma_2+\Gamma_3)$$

VALUE CL% DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.25 90 EDWARDS 82E CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$

$$\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$$

VALUE CL% DOCUMENT ID TECN COMMENT

$<1.27 \times 10^{-3}$ 90 ¹ ABLIKIM 180 BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$

¹ Using results from BAI 00D.

$$\Gamma(\gamma\phi(1020))/\Gamma_{\text{total}}$$

VALUE DOCUMENT ID TECN COMMENT

possibly seen ¹ ABLIKIM 18I BES3 $J/\psi \rightarrow \gamma \gamma \phi(1020)$

¹ Seen as a peak in $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded. Also see $\eta(1405)$.

$$\Gamma_4/\Gamma_1$$

NODE=M175R6

NODE=M175R6

NODE=M175R01
NODE=M175R01

NODE=M175R01;LINKAGE=A

NODE=M175R00
NODE=M175R00

NODE=M175R00;LINKAGE=A

NODE=M175

REFID=58893
REFID=58925
REFID=56785
REFID=51698
REFID=50764
REFID=48848
REFID=48319
REFID=49649
REFID=47954
REFID=47394
REFID=45417
REFID=44614
REFID=41584
REFID=41578
REFID=40924
REFID=21314
REFID=20407

NODE=M152

$\eta(1475)$ REFERENCES

ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PR D115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)

$f_0(1500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See also the reviews on "Scalar mesons below 2 GeV" and on "Non- $q\bar{q}$ mesons".

$f_0(1500)$ MASS

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

1506 ± 6 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

1515 ± 12	1	BARBERIS	00A	450 $p p \rightarrow p f \eta \eta p_s$
1511 ± 9	1,2	BARBERIS	00C	450 $p p \rightarrow p f 4\pi p_s$
1510 ± 8	1	BARBERIS	00E	450 $p p \rightarrow p f \eta \eta p_s$
1522 ± 25	1	BERTIN	98	OBLX 0.05–0.405 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
1449 ± 20	1	BERTIN	97C	OBLX 0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
1500 ± 10	3	AMSLER	95D	CBAR 0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1465 ± 18	4	ROPERTZ	18	RVUE $\bar{B}_s^0 \rightarrow J/\psi (\pi^+ \pi^- / K^+ K^-)$
1447 ± 16 ± 13	5.6	DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1442 ± 9 ± 4	5.6	DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
1460.9 ± 2.9	7	AAIJ	14BR	LHCb $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$
1468 ± 14 ± 23	5.5k	ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1486 ± 10	1	ANISOVICH	09	RVUE 0.0 $\bar{p} p, \pi N$
1470 ± 60	568	KLEMPF	08	E791 $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
1470 ± 6 ± 72	10	UEHARA	08A	BELL 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0$
1466 ± 6 ± 20	11	ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1495 ± 4	AMSLER	06	CBAR	0.9 $\bar{p} p \rightarrow K^+ K^- \pi^0$
1539 ± 20	9.9k	AUBERT	060	BABR $B^+ \rightarrow K^+ K^+ K^-$
1473 ± 5	80k	UMAN	11,12	E835 5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$

OCCUR=2

NODE=M152M

NODE=M152

1478	± 6		VLADIMIRSK...	06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1493	± 7		11 BINON	05	GAMS	$33 \pi^- p \rightarrow \eta \eta n$
1524	± 14	1400	13 GARMASH	05	BELL	$B^+ \rightarrow K^+ K^+ K^-$
1489	± 8		14 ANISOVICH	03	RVUE	
1490	± 30		11 ABELE	01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1497	± 10		11 BARBERIS	99	OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$
1502	± 10		11 BARBERIS	99B	OMEG	$450 pp \rightarrow p_s p_f \pi^+ \pi^-$
1502	$\pm 12 \pm 10$		15 BARBERIS	99D	OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$
1530	± 45		11 BELLAZZINI	99	GAM4	$450 pp \rightarrow pp\pi^0\pi^0$
1505	± 18		11 FRENCH	99		$300 pp \rightarrow p_f(K^+ K^-)p_s$
1447	± 27		16 KAMINSKI	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
1580	± 80		11 ALDE	98	GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
1499	± 8		1 ANISOVICH	98B	RVUE	Compilation
~ 1520			REYES	98	SPEC	$800 pp \rightarrow p_s p_f K_S^0 K_S^0$
1510	± 20		1 BARBERIS	97B	OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
~ 1475			FRAZETTI	97D	E687	$D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 1505			ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
1515	± 20		ABELE	96B	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
1500	± 8		1 ABELE	96C	RVUE	Compilation
1460	± 20	120	11 AMELIN	96B	VES	$37 \pi^- A \rightarrow \eta \eta \pi^- A$
1500	± 8		BUGG	96	RVUE	
1500	± 15		17 AMSLER	95B	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
1505	± 15		18 AMSLER	95C	CBAR	$0.0 \bar{p}p \rightarrow \eta \eta \pi^0$
1445	± 5		19 ANTINORI	95	OMEG	$300, 450 pp \rightarrow pp2(\pi^+ \pi^-)$
1497	± 30		11 ANTINORI	95	OMEG	$300, 450 pp \rightarrow pp\pi^+\pi^-$
~ 1505			BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1446	± 5		11 ABATZIS	94	OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
1545	± 25		11 AMSLER	94E	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta \eta'$
1520	± 25		1,20 ANISOVICH	94	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$
1505	± 20		1,21 BUGG	94	RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0, \eta \pi^0 \pi^0$
1560	± 25		11 AMSLER	92	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta \eta$
1550	$\pm 45 \pm 30$		11 BELADIDZE	92C	VES	$36 \pi^- Be \rightarrow \pi^- \eta' \eta Be$
1449	± 4		11 ARMSTRONG	89E	OMEG	$300 pp \rightarrow pp2(\pi^+ \pi^-)$
1610	± 20		11 ALDE	88	GAM4	$300 \pi^- N \rightarrow \pi^- N 2\eta$
~ 1525			ASTON	88D	LASS	$11 K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1570	± 20	600	11 ALDE	87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
1575	± 45		22 ALDE	86D	GAM4	$100 \pi^- p \rightarrow 2\eta n$
1568	± 33		11 BINON	84C	GAM2	$38 \pi^- p \rightarrow \eta \eta' n$
1592	± 25		11 BINON	83	GAM2	$38 \pi^- p \rightarrow 2\eta n$
1525	± 5		11 GRAY	83	DBC	$0.0 \bar{p}N \rightarrow 3\pi$

1 T-matrix pole.

2 Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

3 T-matrix pole. Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

4 T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.

5 Using CLEO-c data but not authored by the CLEO Collaboration.

6 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 109$ MeV.

7 Solution I, statistical error only.

8 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.9 Reanalysis of AITALA 01A data. This state could also be $f_0(1370)$.10 Breit-Wigner mass. May also be the $f_0(1370)$.

11 Breit-Wigner mass.

12 Statistical error only.

13 Breit-Wigner, solution 1, PWA ambiguous.

14 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

15 Supersedes BARBERIS 99 and BARBERIS 99B.

16 T-matrix pole on sheet $- - +$.

17 T-matrix pole, supersedes ANISOVICH 94.

18 T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

19 Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

NODE=M152M;LINKAGE=PP

NODE=M152M;LINKAGE=PC

NODE=M152M;LINKAGE=AB

NODE=M152M;LINKAGE=H

NODE=M152M;LINKAGE=F

NODE=M152M;LINKAGE=G

NODE=M152M;LINKAGE=I

NODE=M152M;LINKAGE=C

NODE=M152M;LINKAGE=KL

NODE=M152M;LINKAGE=UE

NODE=M152M;LINKAGE=E

NODE=M152M;LINKAGE=ST

NODE=M152M;LINKAGE=GA

NODE=M152M;LINKAGE=KM

NODE=M152M;LINKAGE=BD

NODE=M152M;LINKAGE=TK

NODE=M152M;LINKAGE=D

NODE=M152M;LINKAGE=D1

NODE=M152M;LINKAGE=B

20 From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$.


21 Reanalysis of ANISOVICH 94 data.

22 From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152M;LINKAGE=A

NODE=M152M;LINKAGE=C1

NODE=M152M;LINKAGE=AZ



f₀(1500) WIDTH

NODE=M152W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
112± 9 OUR AVERAGE				
110± 24	1 BARBERIS	00A	450 $\bar{p}p \rightarrow p_f \eta \eta p_s$	
102± 18	1,2 BARBERIS	00C	450 $\bar{p}p \rightarrow p_f 4\pi p_s$	
110± 16	1 BARBERIS	00E	450 $\bar{p}p \rightarrow p_f \eta \eta p_s$	
108± 33	1 BERTIN	98 OBLX	0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$	
114± 30	1 BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
154± 30	3 AMSLER	95D CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

100± 18	4 ROPERTZ	18 RVUE	$\bar{B}_s^0 \rightarrow J/\psi(\pi^+ \pi^- / K^+ K^-)$	
124± 7	5 AAIJ	14BR LHCb	$\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$	
136± 41± 28 – 26 – 100	5.5k	6 ABLIKIM	13N BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$	
114± 10	1 ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$	
90± 2± 50 – 1 – 22	7 UEHARA	08A BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
108± 14± 25 – 11 –	8 ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$	
121± 8	AMSLER	06 CBAR	0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$	
257± 33	9.9k	AUBERT	060 BABR $B^+ \rightarrow K^+ K^+ K^-$	
108± 9	80k	8,9 UMAN	06 E835 5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$	
119± 10		VLADIMIRSK...06	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$	
90± 15	8 BINON	05 GAMS	33 $\pi^- p \rightarrow \eta \eta n$	
136± 23	1400	10 GARMASH	05 BELL $B^+ \rightarrow K^+ K^+ K^-$	
102± 10	11 ANISOVICH	03 RVUE		
140± 40	8 ABELE	01 CBAR	0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$	
104± 25	8 BARBERIS	99 OMEG	450 $\bar{p}p \rightarrow p_s p_f K^+ K^-$	
131± 15	8 BARBERIS	99B OMEG	450 $\bar{p}p \rightarrow p_s p_f \pi^+ \pi^-$	
98± 18± 16	12 BARBERIS	99D OMEG	450 $\bar{p}p \rightarrow K^+ K^-, \pi^+ \pi^-$	
160± 50	8 BELLAZZINI	99 GAM4	450 $\bar{p}p \rightarrow p p \pi^0 \pi^0$	
100± 33	8 FRENCH	99	300 $\bar{p}p \rightarrow p_f (K^+ K^-) p_s$	
108± 46	13 KAMINSKI	99 RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, \sigma \sigma$	
280± 100	8 ALDE	98 GAM4	100 $\pi^- p \rightarrow \pi^0 \pi^0 n$	
130± 20	1 ANISOVICH	98B RVUE	Compilation	
120± 35	1 BARBERIS	97B OMEG	450 $\bar{p}p \rightarrow p p 2(\pi^+ \pi^-)$	

~ 100	FRABETTI	97D E687	$D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 169	ABELE	96 CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
105 ± 15	ABELE	96B CBAR	$0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
100 ± 30	120 8 AMELIN	96B VES	$37 \pi^- A \rightarrow \eta \eta \pi^- A$
132 ± 15	BUGG	96 RVUE	
120 ± 25	14 AMSLER	95B CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
120 ± 30	15 AMSLER	95C CBAR	$0.0 \bar{p}p \rightarrow \eta \eta \pi^0$
65 ± 10	16 ANTINORI	95 OMEG	$300,450 pp \rightarrow pp2(\pi^+ \pi^-)$
199 ± 30	8 ANTINORI	95 OMEG	$300,450 pp \rightarrow pp\pi^+\pi^-$
56 ± 12	8 ABATZIS	94 OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
100 ± 40	8 AMSLER	94E CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta \eta'$
148 ± 20	1,17 ANISOVICH	94 CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$
150 ± 20	1,18 BUGG	94 RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0, \eta \pi^0 \pi^0$
245 ± 50	8 AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta \eta$
$153 \pm 67 \pm 50$	8 BELADIDZE	92C VES	$36 \pi^- Be \rightarrow \pi^- \eta' \eta Be$
78 ± 18	8 ARMSTRONG	89E OMEG	$300 pp \rightarrow pp2(\pi^+ \pi^-)$
170 ± 40	8 ALDE	88 GAM4	$300 \pi^- N \rightarrow \pi^- N2\eta$
150 ± 20	600 8 ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
265 ± 65	19 ALDE	86D GAM4	$100 \pi^- p \rightarrow 2\eta n$
260 ± 60	8 BINON	84C GAM2	$38 \pi^- p \rightarrow \eta \eta' n$
210 ± 40	8 BINON	83 GAM2	$38 \pi^- p \rightarrow 2\eta n$
101 ± 13	8 GRAY	83 DBC	$0.0 \bar{p}N \rightarrow 3\pi$

1 T-matrix pole.

2 Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

3 T-matrix pole. Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

4 T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.

5 Solution I, statistical error only.

6 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.7 Breit-Wigner width. May also be the $f_0(1370)$.

8 Breit-Wigner width.

9 Statistical error only.

10 Breit-Wigner, solution 1, PWA ambiguous.

11 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \eta \eta'$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

12 Supersedes BARBERIS 99 and BARBERIS 99B.

13 T-matrix pole on sheet $-- +$.

14 T-matrix pole, supersedes ANISOVICH 94.

15 T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

16 Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

17 From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$.

18 Reanalysis of ANISOVICH 94 data.

19 From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152W;LINKAGE=PP

NODE=M152W;LINKAGE=PC

NODE=M152W;LINKAGE=AB

NODE=M152W;LINKAGE=F

NODE=M152W;LINKAGE=G

NODE=M152W;LINKAGE=C

NODE=M152W;LINKAGE=UE

NODE=M152W;LINKAGE=E

NODE=M152W;LINKAGE=ST

NODE=M152W;LINKAGE=GA

NODE=M152W;LINKAGE=KM

NODE=M152W;LINKAGE=BD

NODE=M152W;LINKAGE=TK

NODE=M152W;LINKAGE=D

NODE=M152W;LINKAGE=D1

NODE=M152W;LINKAGE=B

NODE=M152W;LINKAGE=A

NODE=M152W;LINKAGE=C1

NODE=M152W;LINKAGE=AZ

NODE=M152215;NODE=M152

DESIG=8

DESIG=9

DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow

DESIG=7

DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=6;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=11;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=12;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=13;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=14;OUR EST; \rightarrow UNCHECKED \leftarrow

DESIG=1

DESIG=2

DESIG=4

DESIG=10;OUR EST; \rightarrow UNCHECKED \leftarrow

$f_0(1500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
$\Gamma_1 \pi\pi$	(34.5 ± 2.2) %	1.2
$\Gamma_2 \pi^+ \pi^-$	seen	
$\Gamma_3 2\pi^0$	seen	
$\Gamma_4 4\pi$	(48.9 ± 3.3) %	1.2
$\Gamma_5 4\pi^0$	seen	
$\Gamma_6 2\pi^+ 2\pi^-$	seen	
$\Gamma_7 2(\pi\pi)_S$ -wave	seen	
$\Gamma_8 \rho\rho$	seen	
$\Gamma_9 \pi(1300)\pi$	seen	
$\Gamma_{10} a_1(1260)\pi$	seen	
$\Gamma_{11} \eta\eta$	(6.0 ± 0.9) %	1.1
$\Gamma_{12} \eta\eta'(958)$	(2.2 ± 0.8) %	1.4
$\Gamma_{13} K\bar{K}$	(8.5 ± 1.0) %	1.1
$\Gamma_{14} \gamma\gamma$	not seen	

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 10 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 5.6$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-88			
x_{11}	27	-56		
x_{12}	3	-32	26	
x_{13}	43	-64	20	2
	x_1	x_4	x_{11}	x_{12}

$f_0(1500) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$	$\Gamma_1 \Gamma_{14} / \Gamma$			
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
33 ^{+12 +1809} _{-6 -21}	1	UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
not seen		ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}} = 91, 183-209 \text{ GeV}$
<460	95	BARATE	00E ALEP	$\gamma\gamma \rightarrow \pi^+ \pi^-$
1 May also be the $f_0(1370)$. Multiplied by us by 3 to obtain the $\pi\pi$ value.				

NODE=M152217

NODE=M152G1
NODE=M152G1

NODE=M152G1;LINKAGE=UE

NODE=M152220

NODE=M152R8
NODE=M152R8NODE=M152R10
NODE=M152R10NODE=M152R6
NODE=M152R6NODE=M152R6;LINKAGE=C
NODE=M152R6;LINKAGE=CHNODE=M152R14
NODE=M152R14

NODE=M152R;LINKAGE=KZ

NODE=M152R15
NODE=M152R15

$f_0(1500)$ BRANCHING RATIOS

$\Gamma(\pi\pi) / \Gamma_{\text{total}}$	Γ_1 / Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.454 ± 0.104	BUGG	96	RVUE	
$\Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}}$	Γ_2 / Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	BERTIN	98	OBLX	$0.05-0.405 \bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
possibly seen	FRABETTI	97D E687	$D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$	

$\Gamma(4\pi) / \Gamma(\pi\pi)$	Γ_4 / Γ_1			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.42 ± 0.18 OUR FIT Error includes scale factor of 1.2.				
1.42 ± 0.18 OUR AVERAGE Error includes scale factor of 1.2.				
1.37 ± 0.16	BARBERIS	00D		$450 p\bar{p} \rightarrow p_f 4\pi p_s$
2.1 ± 0.6	1 AMSLER	98	RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.1 ± 0.2	2 ANISOVICH	02D SPEC	Combined fit	
3.4 ± 0.8	1 ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$

1 Excluding $p\bar{p}$ contribution to 4π .2 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

$\Gamma(2(\pi\pi)s\text{-wave}) / \Gamma(\pi\pi)$	Γ_7 / Γ_1			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.42 ± 0.26	¹ ABELE	01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0$
1 From the combined data of ABELE 96 and ABELE 96C.				

NODE=M152R6;LINKAGE=C
NODE=M152R6;LINKAGE=CH

$\Gamma(2(\pi\pi)s\text{-wave}) / \Gamma(4\pi)$	Γ_7 / Γ_4			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.26 ± 0.07	ABELE	01B	CBAR	$0.0 \bar{p}d \rightarrow 5\pi^0$

NODE=M152R;LINKAGE=KZ

NODE=M152R15
NODE=M152R15

$\Gamma(\rho\rho)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.13 \pm 0.08	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$	

 $\Gamma(\rho\rho)/\Gamma(2(\pi\pi)s\text{-wave})$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_7
2.87 \pm 0.34 OUR AVERAGE Error includes scale factor of 1.1.				
3.3 \pm 0.5	BARBERIS	00C	450 $p p \rightarrow p_f \pi^+ \pi^- 2\pi^0 p_s$	
2.6 \pm 0.4	BARBERIS	00C	450 $p p \rightarrow p_f 2(\pi^+ \pi^-) p_s$	

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.50 \pm 0.25	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$	

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.12 \pm 0.05	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$	

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
large	ALDE	88	GAM4 300 $\pi^- N \rightarrow \eta\eta\pi^- N$	
large	BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$	

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_1
0.173 \pm 0.024 OUR FIT Error includes scale factor of 1.1.				
0.175 \pm 0.027 OUR AVERAGE				
0.18 \pm 0.03	BARBERIS	00E	450 $p p \rightarrow p_f \eta\eta p_s$	
0.157 \pm 0.060	¹ AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.080 \pm 0.033	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$	
0.11 \pm 0.03	² ANISOVICH	02D	SPEC Combined fit	
0.078 \pm 0.013	³ ABELE	96C	RVUE Compilation	
0.230 \pm 0.097	⁴ AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$	
¹ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.				
² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.				
³ 2 π width determined to be 60 ± 12 MeV.				
⁴ Using AMSLER 95B ($3\pi^0$).				

 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_{11}
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.8 \pm 0.3	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$	

 $\Gamma(\eta\eta'(958))/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ_1
0.064 \pm 0.022 OUR FIT Error includes scale factor of 1.4.				
0.095 \pm 0.026				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.005 \pm 0.003	¹ ANISOVICH	02D	SPEC Combined fit	
¹ From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.				

NODE=M152R16
NODE=M152R16NODE=M152R11
NODE=M152R11OCCUR=2

NODE=M152R17
NODE=M152R17NODE=M152R18
NODE=M152R18

NODE=M152R1
NODE=M152R1NODE=M152R13
NODE=M152R13NODE=M152R3;LINKAGE=AB
NODE=M152R;LINKAGE=CHNODE=M152R3;LINKAGE=CM
NODE=M152R3;LINKAGE=ANODE=M152R5
NODE=M152R5NODE=M152R12
NODE=M152R12

NODE=M152R12;LINKAGE=CH

$\Gamma(\eta\eta'(958))/\Gamma(\eta\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ_{11}
0.37±0.13 OUR FIT	Error includes scale factor of 1.5.			
0.29±0.10	¹ AMSLER 95C CBAR 0.0 $p\bar{p} \rightarrow \eta\eta\pi^0$			NODE=M152R2 NODE=M152R2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.05±0.03	² ANISOVICH 02D SPEC Combined fit			
0.84±0.23	ABELE 96C RVUE Compilation			
2.7 ± 0.8	BINON 84C GAM2 38 $\pi^- p \rightarrow \eta\eta' n$			
1 Using AMSLER 94E ($\eta\eta'\pi^0$).				
2 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.			NODE=M152R2;LINKAGE=A NODE=M152R2;LINKAGE=CH	

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	Γ_{13}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.044±0.021	BUGG 96 RVUE		NODE=M152R9 NODE=M152R9

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_1
0.246±0.025 OUR FIT				NODE=M152R7 NODE=M152R7
0.236±0.026 OUR AVERAGE				
0.25 ± 0.03	¹ BARGIOTTI 03 OBLX $\bar{p}p$			
0.19 ± 0.07	² ABELE 98 CBAR 0.0 $\bar{p}p \rightarrow K^0 K^\pm \pi^\mp$			
0.20 ± 0.08	³ ABELE 96B CBAR 0.0 $\bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.16 ± 0.05	⁴ ANISOVICH 02D SPEC Combined fit			
0.33 ± 0.03 ± 0.07	BARBERIS 99D OMEG 450 $p\bar{p} \rightarrow K^+ K^-, \pi^+ \pi^-$			
1 Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.				
2 Using $\pi^0\pi^0$ from AMSLER 95B.				NODE=M152R;LINKAGE=BG
3 Using AMSLER 95B ($3\pi^0$), AMSLER 94C ($2\pi^0\eta$) and SU(3).				NODE=M152R7;LINKAGE=A
4 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.				NODE=M152R7;LINKAGE=D NODE=M152R7;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma(\eta\eta)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_{11}
1.43±0.24 OUR FIT		Error includes scale factor of 1.1.			
1.85±0.41		BARBERIS 00E 450 $p\bar{p} \rightarrow p_f \eta\eta p_s$			NODE=M152R4 NODE=M152R4
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.5 ± 0.6		¹ ANISOVICH 02D SPEC Combined fit			
<0.4	90	² PROKOSHKIN 91 GAM4 300 $\pi^- p \rightarrow \pi^- p\eta\eta$			
<0.6		³ BINON 83 GAM2 38 $\pi^- p \rightarrow 2\eta n$			
1 From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n$, $\eta\eta n$, $\eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.					NODE=M152R4;LINKAGE=CH
2 Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production.					
3 Using ETKIN 82B and COHEN 80.					

 $f_0(1500)$ REFERENCES

ROPERTZ 18 EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis (BONN, JULI) (LHCb Collab.)	REFID=59332
AAIJ 17V JHEP 1708 037	R. Aaij <i>et al.</i> (NWES)	REFID=57828
DOBBS 15 PR D91 052006	S. Dobbs <i>et al.</i> (LHCb Collab.)	REFID=56805
AAIJ 14BR PR D89 092006	R. Aaij <i>et al.</i> (BESIII Collab.)	REFID=56035
ABLIKIM 13N PR D87 092009	Ablikim M. <i>et al.</i> (BESIII Collab.)	REFID=55387
ANISOVICH 09 IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev (BONN+)	REFID=52719
KLEMPT 08 EPJ C55 39	E. Klemp, M. Matveev, A.V. Sarantsev (BONN+)	REFID=52286
UEHARA 08A PR D78 052004	S. Uehara <i>et al.</i> (BELLE Collab.)	REFID=52309
ABLIKIM 06V PL B642 441	M. Ablikim <i>et al.</i> (BES Collab.)	REFID=51507
AMSLER 06 PL B639 165	C. Amsler <i>et al.</i> (CBAR Collab.)	REFID=51136
AUBERT 06O PR D74 032003	B. Aubert <i>et al.</i> (BABAR Collab.)	REFID=51141
UMAN 06 PR D73 052009	I. Uman <i>et al.</i> (FNAL E835)	REFID=51063
VLADIMIRSKY 06 PAN 69 493	V.V. Vladimirska <i>et al.</i> (ITEP, Moscow)	REFID=51191
	Translated from YAF 69 515.	
BINON 05 PAN 68 960	F. Binon <i>et al.</i>	REFID=50780
	Translated from YAF 68 998.	
GARMASH 05 PR D71 092003	A. Garmash <i>et al.</i> (BELLE Collab.)	REFID=50641
ANISOVICH 03 EPJ A16 229	V.V. Anisovich <i>et al.</i> (OBELIX Collab.)	REFID=49401
BARGIOTTI 03 EPJ C26 371	M. Bargiotti <i>et al.</i>	REFID=49217
AMSLER 02 EPJ C23 29	C. Amsler <i>et al.</i>	REFID=48580
ANISOVICH 02D PAN 65 1545	V.V. Anisovich <i>et al.</i>	REFID=48831
	Translated from YAF 65 1583.	

ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
AITALA	01A	PRD 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00D	PL B474 423	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47960
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
AMSLER	98	RMP 70 1293	C. Amsler		REFID=46601
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
REYES	98	PRD 81 4079	M.A. Reyes <i>et al.</i>		REFID=46378
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=45554
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45011
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45038
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
AMELIN	96B	PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44725
		Translated from YAF 59 1021.			
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)	REFID=45094
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44437
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44090
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
AMSLER	94E	PL B340 259	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44080
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)	REFID=43175
		Translated from YAF 55 2748.			
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
ALDE	88	PL B201 160	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=40283
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40330
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=21418
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20751
		Translated from YAF 38 934.			
GRAY	83	PR D27 307	L. Gray <i>et al.</i>	(SYRA)	REFID=21370
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL)	REFID=20381

$f_1(1510)$ $I^G(J^{PC}) = 0^+(1^{++})$

OMMITTED FROM SUMMARY TABLE

See the minireview under $\eta(1405)$.

NODE=M084

 $f_1(1510)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1518± 5 OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.
1530±10		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
1512± 4	600	¹ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1526± 6	271	GAVILLET	82 HBC	$4.2 K^- p \rightarrow \Lambda K K \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~1525		² BAUER	93B	$\gamma\gamma^* \rightarrow \pi^+ \pi^- \pi^0 \pi^0$


¹ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.² Not seen by AIHARA 88C in the $K_S^0 K^\pm \pi^\mp$ final state.

NODE=M084

NODE=M084M

NODE=M084M

WEIGHTED AVERAGE
1518±5 (Error scaled by 1.7)



NODE=M084M;LINKAGE=A

NODE=M084M;LINKAGE=C

 $f_1(1510)$ WIDTH

NODE=M084W


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
73±25 OUR AVERAGE				Error includes scale factor of 2.5. See the ideogram below.
100±40		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
35±15	600	³ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
107±15	271	GAVILLET	82 HBC	$4.2 K^- p \rightarrow \Lambda K K \pi$

³ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.

NODE=M084W

NODE=M084W

NODE=M084W;LINKAGE=A



$f_1(1510)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad K\bar{K}^*(892) + \text{c.c.}$	seen
$\Gamma_2 \quad \pi^+\pi^-\eta'$	seen

$f_1(1510)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$	EVTS	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	Γ_2/Γ
seen	230	ABLIKIM	11C	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

$f_1(1510)$ REFERENCES

ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BAUER	93B	PR D48 3976	D.A. Bauer <i>et al.</i>	(SLAC)
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
GAVILLET	82	ZPHY C16 119	P. Gavillet <i>et al.</i>	(CERN, CDEF, PADO+)

NODE=M084215;NODE=M084

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=2

NODE=M084225

NODE=M084R01
NODE=M084R01

NODE=M084

REFID=53684
REFID=43678
REFID=40564
REFID=40282
REFID=40568
REFID=20877

$f'_2(1525)$ $I^G(J^{PC}) = 0^+(2^{++})$

NODE=M013

 $f'_2(1525)$ MASS

VALUE (MeV)

DOCUMENT ID

1525±5 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.**PRODUCED BY PION BEAM**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1521±13		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1547 ⁺¹⁰ ₋₂	1	LONGACRE 86	MPS	$22 \pi^- p \rightarrow K_S^0 K_S^0 n$
1496 ⁺⁹ ₋₈	2	CHABAUD 81	ASPK	$6 \pi^- p \rightarrow K^+ K^- n$
1497 ⁺⁸ ₋₉		CHABAUD 81	ASPK	$18.4 \pi^- p \rightarrow K^+ K^- n$
1492±29		GORLICH 80	ASPK	$17 \pi^- p$ polarized $\rightarrow K^+ K^- n$
1502±25	3	CORDEN 79	OMEG	$12-15 \pi^- p \rightarrow \pi^+ \pi^- n$
1480	14	CRENNELL 66	HBC	$6.0 \pi^- p \rightarrow K_S^0 K_S^0 n$

NODE=M013205

NODE=M013MX


→ UNCHECKED ←

PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1518.1± 2.8 OUR AVERAGE Includes data from the datablock that follows this one.				
Error includes scale factor of 3.0. See the ideogram below.				
1526.8± 4.3		ASTON 88D	LASS	$11 K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1504 ± 12		BOLONKIN 86	SPEC	$40 K^- p \rightarrow K_S^0 K_S^0 Y$
1529 ± 3		ARMSTRONG 83B	OMEG	$18.5 K^- p \rightarrow K^- K^+ \Lambda$
1521 ± 6	650	AGUILAR...	81B	HBC $4.2 K^- p \rightarrow \Lambda K^+ K^-$
1521 ± 3	572	ALHARRAN 81	HBC	$8.25 K^- p \rightarrow \Lambda K \bar{K}$
1522 ± 6	123	BARREIRO 77	HBC	$4.15 K^- p \rightarrow \Lambda K_S^0 K_S^0$
1528 ± 7	166	EVANGELIS...	77	OMEG $10 K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1527 ± 3	120	BRANDENB...	76C	ASPK $13 K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1519 ± 7	100	AGUILAR...	72B	HBC $3.9, 4.6 K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1514 ± 8	61	BINON 07	GAMS	$32.5 K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
1513 ± 10		⁴ BARKOV 99	SPEC	$40 K^- p \rightarrow K_S^0 K_S^0 y$

NODE=M013M2

NODE=M013M2

WEIGHTED AVERAGE
1518.1±2.8 (Error scaled by 3.0)PRODUCED BY K^\pm BEAM (MeV)

PRODUCED IN $e^+ e^-$ ANNIHILATION AND PARTICLE DECAYS


VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

1514 ± 5 OUR AVERAGE Error includes scale factor of 3.8. See the ideogram below.

1522.2 \pm 2.8 \pm 5.3	AAIJ	13AN LHCb	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$	
1513 \pm 5 \pm 4 5.5k	5 ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$	
1525.3 \pm 1.2 \pm 3.7	UEHARA	13 BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$	
1521 \pm 5	ABLIKIM	05 BES2	$J/\psi \rightarrow \phi K^+ K^-$	
1518 \pm 1 \pm 3	ABE	04 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
1519 \pm 2 \pm 15	BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$	
1523 \pm 6 331	6 ACCIARRI	01H L3	91, 183–209 $e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
1535 \pm 5 \pm 4	ABREU	96C DLPH	$Z^0 \rightarrow K^+ K^- + X$	
1516 \pm 5 \pm 9	BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$	
1531.6 \pm 10.0	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$	
1496 \pm 2	7 FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$	OCCUR=2
1525 \pm 10 \pm 10	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1532 \pm 3 \pm 6 644 8.9 DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$	
1557 \pm 9 \pm 3 113 8.9 DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$	
1526 \pm 7 29 10 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$		OCCUR=2
1523 \pm 5 870 11 SCHEGELSKY 06A RVUE			$\gamma \gamma \rightarrow K_S^0 K_S^0$	
1515 \pm 5	12 FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$	

WEIGHTED AVERAGE
1514+5-4 (Error scaled by 3.8)



$f_2'(1525)$ mass, $e^+ e^-$ annihilation and particle decays (MeV)

PRODUCED IN $\bar{p}p$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1512 \pm 4 OUR AVERAGE

1513 \pm 4	AMSLER	06 CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
1508 \pm 9	13 AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1530 \pm 12	14 ANISOVICH	09 RVUE	$0.0 \bar{p}p, \pi N$
---------------	--------------	---------	-----------------------

CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1515 \pm 15	BARBERIS	99 OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$
---------------------------------	----------	---------	--------------------------------------

NODE=M013M3

NODE=M013M3

NODE=M013M9

NODE=M013M9

NODE=M013M4

NODE=M013M4

PRODUCED IN $e p$ COLLISIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$1512 \pm 3^{+1.4}_{-0.5}$	15	CHEKANOV	08	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1537^{+9}_{-8}	84	16 CHEKANOV	04	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
1 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. 2 CHABAUD 81 is a reanalysis of PAWLICKI 77 data. 3 From an amplitude analysis where the $f'_2(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious. 4 Systematic errors not estimated. 5 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances. 6 Supersedes ACCIARRI 95J. 7 From an analysis including interference with $f_0(1710)$. 8 Using CLEO-c data but not authored by the CLEO Collaboration. 9 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 73$ MeV. 10 From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated. 11 From analysis of L3 data at 91 and 183–209 GeV. 12 From an analysis ignoring interference with $f_0(1710)$. 13 T-matrix pole. 14 4-poles, 5-channel K matrix fit. 15 In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background. 16 Systematic errors not estimated.				

NODE=M013M10

NODE=M013M10

 $f'_2(1525)$ WIDTH

VALUE (MeV)		DOCUMENT ID		COMMENT
73^{+6}_{-5} OUR FIT				
76 ± 10		PDG	90	For fitting

NODE=M013210

NODE=M013WX

PRODUCED BY PION BEAM

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
102 ± 42		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
108^{+5}_{-2}	17	LONGACRE	86	$MPS 22 \pi^- p \rightarrow K_S^0 K_S^0 n$
69^{+22}_{-16}	18	CHABAUD	81	$ASPK 6 \pi^- p \rightarrow K^+ K^- n$
137^{+23}_{-21}	CHABAUD	81	ASPK	$18.4 \pi^- p \rightarrow K^+ K^- n$
150^{+83}_{-50}	GORLICH	80	ASPK	$17 \pi^- p$ polarized $\rightarrow K^+ K^- n$
165 ± 42	19	CORDEN	79	$OMEG 12\text{--}15 \pi^- p \rightarrow \pi^+ \pi^- n$
92^{+39}_{-22}	20	POLYCHRO...	79	$STRC 7 \pi^- p \rightarrow n K_S^0 K_S^0$

NODE=M013W1

NODE=M013W1

OCCUR=2

PRODUCED BY K^\pm BEAM


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$88.5^{+2.5}_{-2.2}$ OUR AVERAGE				Includes data from the datablock that follows this one. Error includes scale factor of 1.4. See the ideogram below.
90 ± 12		ASTON 88D	LASS	$11 K^- p \rightarrow K_S^0 K_S^0 \Lambda$
73 ± 18		BOLONKIN 86	SPEC	$40 K^- p \rightarrow K_S^0 K_S^0 Y$
83 ± 15		ARMSTRONG 83B	OMEG	$18.5 K^- p \rightarrow K^- K^+ \Lambda$
85 ± 16	650	AGUILAR-...	HBC	$4.2 K^- p \rightarrow \Lambda K^+ K^-$
80^{+14}_{-11}	572	ALHARRAN 81	HBC	$8.25 K^- p \rightarrow \Lambda K\bar{K}$
72 ± 25	166	EVANGELIS...	77	$OMEG 10 K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
69 ± 22	100	AGUILAR-...	72B	$HBC 3.9, 4.6 K^- p \rightarrow K\bar{K} (\Lambda, \Sigma)$

NODE=M013W2

NODE=M013W2

• • • We do not use the following data for averages, fits, limits, etc. • • •

92 ± 25	61	BINON	07	GAMS	32.5 $K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
75 ± 20	21	BARKOV	99	SPEC	40 $K^- p \rightarrow K_S^0 K_S^0 \gamma$
62 ± 19	123	BARREIRO	77	HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
61 ± 8	120	BRANDENB...	76C	ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$



PRODUCED IN $e^+ e^-$ ANNIHILATION AND PARTICLE DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M013W3
NODE=M013W3


89.2 \pm 3.4 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.

84 ± 6 ± 10	AAIJ	13AN LHCb	$B_s^0 \rightarrow J/\psi K^+ K^-$
75 ± 12 ± 16	5.5k	22 ABLIKIM	13N BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
82.9 ± 2.1 ± 3.3		UEHARA	13 BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
77 ± 15		ABLIKIM	05 BES2 $J/\psi \rightarrow \phi K^+ K^-$
82 ± 2 ± 3		ABE	04 BELL $10.6 \frac{e^+ e^-}{e^+ e^- K^+ K^-} \rightarrow$
75 ± 4 ± 15		BAI	03G BES $J/\psi \rightarrow \gamma K\bar{K}$
100 ± 15	331	23 ACCIARRI	91, 183–209 $e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$
60 ± 20 ± 19		ABREU	96C DLPH $Z^0 \rightarrow K^+ K^- + X$
60 ± 23 ± 13		BAI	96C BES $J/\psi \rightarrow \gamma K^+ K^-$
103 ± 30		AUGUSTIN	88 DM2 $J/\psi \rightarrow \gamma K^+ K^-$
100 ± 3	24 FALVAR	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$
85 ± 35		BALTRUSAIT...87	MRK3 $J/\psi \rightarrow \gamma K^+ K^-$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

37 ± 12	29	25 LEES	14H BABR $e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
104 ± 10	870	26 SCHEGELSKY	06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
62 ± 10		27 FALVAR	88 DM2 $J/\psi \rightarrow \phi K^+ K^-$



PRODUCED IN $\bar{p}p$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
77± 5 OUR AVERAGE			
76± 6	AMSLER 06	CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
79± 8	28 AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
128±20	29 ANISOVICH 09	RVUE	$0.0 \bar{p}p, \pi N$

NODE=M013W9
NODE=M013W9

CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70±25			
	BARBERIS 99	OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$

NODE=M013W4
NODE=M013W4

PRODUCED IN ep COLLISIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83± 9⁺⁵₋₄	30	CHEKANOV 08	ZEUS	$ep \rightarrow K_S^0 K_S^0 X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
50 ⁺³⁴ ₋₂₂	84	31 CHEKANOV 04	ZEUS	$ep \rightarrow K_S^0 K_S^0 X$

NODE=M013W10
NODE=M013W10

- 17 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- 18 CHABAUD 81 is a reanalysis of PAWLICKI 77 data.
- 19 From an amplitude analysis where the $f'_2(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.
- 20 From a fit to the D with $f_2(1270)-f'_2(1525)$ interference. Mass fixed at 1516 MeV.
- 21 Systematic errors not estimated.
- 22 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.
- 23 Supersedes ACCIARRI 95J.
- 24 From an analysis including interference with $f_0(1710)$.
- 25 From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated.
- 26 From analysis of L3 data at 91 and 183–209 GeV.
- 27 From an analysis ignoring interference with $f_0(1710)$.
- 28 T-matrix pole.
- 29 4-poles, 5-channel K matrix fit.
- 30 In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.
- 31 Systematic errors not estimated.

NODE=M013W;LINKAGE=L
NODE=M013W;LINKAGE=D
NODE=M013W;LINKAGE=N

NODE=M013W;LINKAGE=M
NODE=M013W2;LINKAGE=SK
NODE=M013W3;LINKAGE=A

NODE=M013W;LINKAGE=HA
NODE=M013W;LINKAGE=F2
NODE=M013W3;LINKAGE=B

NODE=M013W3;LINKAGE=SC
NODE=M013W;LINKAGE=F1
NODE=M013W;LINKAGE=TT
NODE=M013W9;LINKAGE=AN
NODE=M013W10;LINKAGE=HE

NODE=M013W10;LINKAGE=CH

$f'_2(1525)$ DECAY MODES

NODE=M013215;NODE=M013

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 K\bar{K}$	(88.7 \pm 2.2) %	DESIG=2
$\Gamma_2 \eta\eta$	(10.4 \pm 2.2) %	DESIG=4
$\Gamma_3 \pi\pi$	(8.2 \pm 1.5) $\times 10^{-3}$	DESIG=1
$\Gamma_4 K\bar{K}^*(892) + \text{c.c.}$		DESIG=3
$\Gamma_5 \pi K\bar{K}$		DESIG=6
$\Gamma_6 \pi\pi\eta$		DESIG=5
$\Gamma_7 \pi^+\pi^-\pi^-\pi^-$		DESIG=7
$\Gamma_8 \gamma\gamma$	(1.10 \pm 0.14) $\times 10^{-6}$	DESIG=8

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 2 partial widths, a combination of partial widths obtained from integrated cross sections, and 3 branching ratios uses 17 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 14.3$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100			
x_3	-6	-1		
x_8	-6	6	1	
Γ	-23	23	-1	-56
	x_1	x_2	x_3	x_8

Mode	Rate (MeV)	
$\Gamma_1 K\bar{K}$	65 \pm 5	DESIG=2
$\Gamma_2 \eta\eta$	7.6 \pm 1.8	DESIG=4
$\Gamma_3 \pi\pi$	0.60 \pm 0.12	DESIG=1
$\Gamma_8 \gamma\gamma$	(8.1 \pm 0.9) $\times 10^{-5}$	DESIG=8

 $f'_2(1525)$ PARTIAL WIDTHS

$\Gamma(K\bar{K})$	Γ_1
VALUE (MeV)	DOCUMENT ID TECN COMMENT

 65^{+5}_{-4} OUR FIT **63^{+6}_{-5}** $^{32}\text{LONGACRE}$ 86 MPS $22\pi^- p \rightarrow K_S^0 K_S^0 n$

$\Gamma(\eta\eta)$	Γ_2
VALUE (MeV)	DOCUMENT ID TECN COMMENT

 7.6 ± 1.8 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.0 \pm 0.8	870	$^{33}\text{SCHEGELSKY}$ 06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
24 \pm 3		$^{32}\text{LONGACRE}$	86 MPS $22\pi^- p \rightarrow K_S^0 K_S^0 n$

$\Gamma(\pi\pi)$	Γ_3
VALUE (MeV)	DOCUMENT ID TECN COMMENT

 0.60 ± 0.12 OUR FIT **$1.4^{+1.0}_{-0.5}$** $^{32}\text{LONGACRE}$ 86 MPS $22\pi^- p \rightarrow K_S^0 K_S^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2 \pm 1.0	870	$^{33}\text{SCHEGELSKY}$ 06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
-0.2			

NODE=M013220

NODE=M013W6
NODE=M013W6NODE=M013W7
NODE=M013W7NODE=M013W5
NODE=M013W5

$\Gamma(\gamma\gamma)$

<u>VALUE</u> (keV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_8
0.081±0.009 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.13 ± 0.03	870	33 SCHEGELSKY 06A RVUE	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
32 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.					NODE=M013W8
33 From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f'_2(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.					NODE=M013W8;LINKAGE=L

 $f'_2(1525) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

<u>VALUE</u> (keV)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	$\Gamma_1\Gamma_8/\Gamma$
0.072 ± 0.007 OUR FIT					
0.072 ± 0.007 OUR AVERAGE					
0.048 +0.067 -0.008	+0.108 -0.012	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
0.0564 ± 0.0048 ± 0.0116		ABE	04 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
0.076 ± 0.006 ± 0.011	331	34 ACCIARRI	01H L3	$e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
0.067 ± 0.008 ± 0.015		35 ALBRECHT	90G ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
0.11 +0.03 -0.02	± 0.02	BEHREND	89C CELL	$e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
0.10 +0.04 -0.03	± 0.03	BERGER	88 PLUT	$e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
0.12 ± 0.07 ± 0.04		35 AIHARA	86B TPC	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
0.11 ± 0.02 ± 0.04		35 ALTHOFF	83 TASS	$e^+ e^- \rightarrow e^+ e^- K\bar{K}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0314 ± 0.0050 ± 0.0077		36 ALBRECHT	90G ARG	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$	OCCUR=2
34 Supersedes ACCIARRI 95J. From analysis of L3 data at 91 and 183–209 GeV,					NODE=M013G;LINKAGE=HA
35 Using an incoherent background.					NODE=M013G1;LINKAGE=A
36 Using a coherent background.					NODE=M013G1;LINKAGE=B

 $f'_2(1525) \text{ BRANCHING RATIOS}$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
seen		UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$		
0.10 ± 0.03		37 PROKOSHKIN 91	GAM4	$300 \pi^- p \rightarrow \pi^- p \eta\eta$		
37 Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.						

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ_1
0.118±0.028 OUR FIT						
0.115±0.028 OUR AVERAGE						
0.119 ± 0.015 ± 0.036		61 38 BINON	07 GAMS	$32.5 K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$		
0.11 ± 0.04		39 PROKOSHKIN 91	GAM4	$300 \pi^- p \rightarrow \pi^- p \eta\eta$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 0.14	90	BARBERIS	00E	$450 pp \rightarrow p_f \eta\eta p_s$		
< 0.50		BARNES	67 HBC	$4.6, 5.0 K^- p$		
38 Using the compilation of the cross sections for $f'_2(1525)$ production in $K^- p$ collisions from ASTON 88D.						
39 Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.						

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ
0.0082±0.0016 OUR FIT						
0.0075±0.0016 OUR AVERAGE						
0.007 ± 0.002		COSTA	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$		
0.027 +0.071 -0.013		40 GORLICH	80 ASPK	$17, 18 \pi^- p$		
0.0075 ± 0.0025		40, 41 MARTIN	79 RVUE			

NODE=M013R8
NODE=M013R8NODE=M013PW;LINKAGE=L
NODE=M013W8;LINKAGE=SC

NODE=M013223

NODE=M013G1
NODE=M013G1

OCCUR=2

NODE=M013G;LINKAGE=HA
NODE=M013G1;LINKAGE=A
NODE=M013G1;LINKAGE=B

NODE=M013225

NODE=M013R8
NODE=M013R8

NODE=M013R8;LINKAGE=B

NODE=M013R3
NODE=M013R3

NODE=M013R3;LINKAGE=BI

NODE=M013R3;LINKAGE=B

NODE=M013R1
NODE=M013R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.06	95	AGUILAR-...	81B	HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
0.19 ± 0.03		CORDEN	79	OMEG	12–15 $\pi^- p \rightarrow \pi^+ \pi^- n$
<0.045	95	BARREIRO	77	HBC	4.15 $K^- p \rightarrow \Lambda K^0 S K^0 S$
0.012 ± 0.004	40	PAWICKI	77	SPEC	6 $\pi N \rightarrow K^+ K^- N$
<0.063	90	BRANDENB...	76C	ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
<0.0086	40	BEUSCH	75B	OSPK	8.9 $\pi^- p \rightarrow K^0 \bar{K}^0 n$

40 Assuming that the $f'_2(1525)$ is produced by an one-pion exchange production mechanism.

41 MARTIN 79 uses the PAWICKI 77 data with different input value of the $f'_2(1525) \rightarrow K\bar{K}$ branching ratio.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$

VALUE		DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.0092±0.0018 OUR FIT					
0.075 ± 0.035		AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

NODE=M013R1;LINKAGE=C

NODE=M013R1;LINKAGE=D

$[\Gamma(K\bar{K}^*(892)+c.c.) + \Gamma(\pi K\bar{K})]/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$(\Gamma_4+\Gamma_5)/\Gamma_1$
-------	-----	-------------	------	---------	--------------------------------

NODE=M013R5

NODE=M013R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.35	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
<0.4	67	AMMAR	67	HBC	

$\Gamma(\pi\pi\eta)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_1
-------	-----	-------------	------	---------	---------------------

NODE=M013R4

NODE=M013R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.41	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
<0.3	67	AMMAR	67	HBC	

$\Gamma(\pi^+ \pi^+ \pi^- \pi^-)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_1
-------	-----	-------------	------	---------	---------------------

NODE=M013R6

NODE=M013R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.32	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
-------	----	-------------	-----	-----	-----------------

$f'_2(1525)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
AAIJ	13AN	PR D87 072004	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55137
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		REFID=52719
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
BINON	07	PAN 70 1713	F. Binon <i>et al.</i>	(GAMS Collab.)	REFID=52057
AMSLER	06	PL B639 165	Translated from YAF 70 1758. C. Amsler <i>et al.</i>	(CBAR Collab.)	REFID=51136
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=49672
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
AMSLER	02	EPJ C23 29	Translated from YAF 66 860. C. Amsler <i>et al.</i>		REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciari <i>et al.</i>	(L3 Collab.)	REFID=48321
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARKOV	99	JETPL 70 248	B.P. Barkov <i>et al.</i>		REFID=47379
ABREU	96C	PL B379 309	Translated from ZETFP 70 242. P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
ACCIARRI	95J	PL B363 118	M. Acciari <i>et al.</i>	(L3 Collab.)	REFID=44615
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
ALBRECHT	90G	ZPHY C48 183	Translated from DANS 316 900. H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
PDG	90	PL B239 1	J.J. Hernandez <i>et al.</i>	(IFIC, BOST, CIT+)	REFID=40744
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40330
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BERGER	88	ZPHY C37 329	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=40566
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=20764
BOLONKIN	86	SJNP 43 776	B.V. Bolonkin <i>et al.</i>	(ITEP) JP	REFID=44646
		Translated from YAF 43 1211.			

LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	83	PL 121B 216	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21408
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
AGUILAR-...	81B	ZPHY C8 313	M. Aguilar-Benitez <i>et al.</i>	(CERN, CDEF+)	REFID=21104
ALHARRAN	81	NP B191 26	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=21403
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlitch <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
BARREIRO	77	NP B121 237	F. Barreiro <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=21392
EVANGELIS...	77	NP B127 384	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20540
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJP	REFID=20367
BRANDENB...	76C	NP B104 413	G.W. Brandenburg <i>et al.</i>	(SLAC)	REFID=20225
BEUSCH	75B	PL 60B 101	W. Beusch <i>et al.</i>	(CERN, ETH)	REFID=21390
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
AMMAR	67	PRL 19 1071	R. Ammar <i>et al.</i>	(NWES, ANL) JP	REFID=21382
BARNES	67	PRL 19 964	V.E. Barnes <i>et al.</i>	(BNL, SYRA) IJPC	REFID=21383
CRENNELL	66	PRL 16 1025	D.J. Crennell <i>et al.</i>	(BNL) I	REFID=20317

 $f_2(1565)$

$I^G(J^{PC}) = 0^+(2^{++})$

OMMITTED FROM SUMMARY TABLE

Seen mostly in antinucleon-nucleon annihilation. Needs confirmation
in other channels. **$f_2(1565)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
1542±19 OUR AVERAGE			Error includes scale factor of 2.2. See the ideogram below.	
1552±13	1 AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$	
1575±18	1 BERTIN	98	OBLX $0.05-0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$	
1507±15	1 BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1560±15	2 ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$	
1590±10	3,4 AMELIN	06	VES $36 \pi^- p \rightarrow \omega\omega n$	
1550±10±20	4 AMELIN	00	VES $37 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^-$	
1598±11± 9	BAKER	99B	SPEC $0 \bar{p}p \rightarrow \omega\omega\pi^0$	
1534±20	5 ABELE	96C	RVUE Compilation	
~ 1552	6 AMSLER	95D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$	
1598±72	BALOSHIN	95	SPEC $40 \pi^- C \rightarrow K_S^0 K_S^0 X$	
1566 ⁺⁸⁰ ₋₅₀	7 ANISOVICH	94	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0$	
1502± 9	ADAMO	93	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$	
1488±10	8 ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$	
1508±10	8 ARMSTRONG	93D	E760 $\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$	
1525±10	8 ARMSTRONG	93D	E760 $\bar{p}p \rightarrow \eta\pi^0 \pi^0 \rightarrow 6\gamma$	
~ 1504	9 WEIDENAUER	93	ASTE $0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$	
1540±15	8 ADAMO	92	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$	
1515±10	10 AKER	91	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$	
1565±20	MAY	90	ASTE $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$	
1477± 5	BRIDGES	86C	DBC $0.0 \bar{p}N \rightarrow 3\pi^- 2\pi^+$	

1 T-matrix pole.

2 On sheet II in a two-pole solution.

3 Supersedes the $\omega\omega$ state of BELADIDZE 92B earlier assigned to the $f_2(1640)$.

4 Breit-Wigner width.

5 T-matrix pole, large coupling to $\rho\rho$ and $\omega\omega$, could be $f_2(1640)$.

6 Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

7 From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$ including AKER 91 data.8 J^P not determined, could be partly $f_0(1500)$.9 J^P not determined.

10 Superseded by AMSLER 95B.

REFID=20768
 REFID=21408
 REFID=20558
 REFID=21104
 REFID=21403
 REFID=20742
 REFID=20737
 REFID=20738
 REFID=20374
 REFID=20377
 REFID=20378
 REFID=21392
 REFID=20540
 REFID=20367
 REFID=20225
 REFID=21390
 REFID=20205
 REFID=21382
 REFID=21383
 REFID=20317

NODE=M123

NODE=M123


NODE=M123M

NODE=M123M

OCCUR=2

NODE=M123M;LINKAGE=G
 NODE=M123M;LINKAGE=AN
 NODE=M123M;LINKAGE=AM
 NODE=M123M;LINKAGE=D
 NODE=M123M;LINKAGE=AA
 NODE=M123M;LINKAGE=AB
 NODE=M123M;LINKAGE=C

NODE=M123M;LINKAGE=E
 NODE=M123M;LINKAGE=F
 NODE=M123M;LINKAGE=BA



f₂(1565) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
122± 13 OUR AVERAGE			
113± 23	11 AMSLER	02 CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
119± 24	11 BERTIN	98 OBLX	0.05–0.405 $\bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
130± 20	11 BERTIN	97C OBLX	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
280± 40	12 ANISOVICH	09 RVUE	0.0 $\bar{p}p, \pi N$
140± 11	13,14 AMELIN	06 VES	36 $\pi^- p \rightarrow \omega \omega n$
130± 20±40	14 AMELIN	00 VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
180± 60	15 ABELE	96C RVUE	Compilation
~142	16 AMSLER	95D CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$
263±101	BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
166 ₋ 80 ₊ 20	17 ANISOVICH	94 CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0$
130± 10	18 ADAMO	93 OBLX	$\bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
148± 27	19 ARMSTRONG	93C E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
103± 15	19 ARMSTRONG	93D E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111± 10	19 ARMSTRONG	93D E760	$\bar{p}p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~206	20 WEIDENAUER	93 ASTE	0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$
132± 37	19 ADAMO	92 OBLX	$\bar{p}p \rightarrow \pi^+ \pi^+ \pi^-$
120± 10	21 AKER	91 CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$
170± 40	MAY	90 ASTE	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
116± 9	BRIDGES	86C DBC	0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$

11 T-matrix pole.

12 On sheet II in a two-pole solution.

13 Supersedes the $\omega \omega$ state of BELADIDZE 92B earlier assigned to the f₂(1640).

14 Breit-Wigner width.

15 T-matrix pole, large coupling to $\rho \rho$ and $\omega \omega$, could be f₂(1640).

16 Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

17 From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$ including AKER 91 data.

18 Supersedes ADAMO 92.

19 J^P not determined, could be partly f₀(1500).

20 J^P not determined.

21 Superseded by AMSLER 95B.

NODE=M123W

NODE=M123W

OCCUR=2

NODE=M123W;LINKAGE=G

NODE=M123W;LINKAGE=AN

NODE=M123W;LINKAGE=AM

NODE=M123W;LINKAGE=H

NODE=M123W;LINKAGE=CC

NODE=M123W;LINKAGE=AB

NODE=M123W;LINKAGE=D

NODE=M123W;LINKAGE=C

NODE=M123W;LINKAGE=E

NODE=M123W;LINKAGE=F

NODE=M123W;LINKAGE=BA

$f_2(1565)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 \pi^+\pi^-$	seen
$\Gamma_3 \pi^0\pi^0$	seen
$\Gamma_4 \rho^0\rho^0$	seen
$\Gamma_5 2\pi^+2\pi^-$	seen
$\Gamma_6 \eta\eta$	seen
$\Gamma_7 \omega\omega$	seen
$\Gamma_8 K\bar{K}$	seen
$\Gamma_9 \gamma\gamma$	seen

 $f_2(1565)$ PARTIAL WIDTHS

$\Gamma(\eta\eta)$	Γ_6
<u>VALUE (MeV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
1.2 ± 0.3	870 22 SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
$\Gamma(K\bar{K})$	Γ_8
<u>VALUE (MeV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
2.0 ± 1.0	870 22 SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
$\Gamma(\gamma\gamma)$	Γ_9
<u>VALUE (keV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.70 ± 0.14	870 22 SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
22 From analysis of L3 data at 91 and 183–209 GeV, using $f_2(1565)$ mass of 1570 MeV, width of 160 MeV, $\Gamma(\pi\pi) = 25$ MeV, and SU(3) relations.	

 $f_2(1565)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	BAKER 99B SPEC $0 \bar{p}p \rightarrow \omega\omega\pi^0$
$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	BERTIN 98 OBLX $0.05\text{--}0.405 \bar{n}p \rightarrow \pi^+\pi^+\pi^-$
not seen	23 ANISOVICH 94B RVUE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
seen	MAY 89 ASTE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
23 ANISOVICH 94B is from a reanalysis of MAY 90.	
$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	AMSLER 95B CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$
$\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$	Γ_2/Γ_4
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.042 ± 0.013	BRIDGES 86B DBC $\bar{p}N \rightarrow 3\pi^-2\pi^+$
$\Gamma(\eta\eta)/\Gamma(\pi^0\pi^0)$	Γ_6/Γ_3
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$0.024 \pm 0.005 \pm 0.012$	24 ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
24 J^P not determined, could be partly $f_0(1500)$.	

NODE=M123215;NODE=M123

DESIG=6;OUR EST; \rightarrow UNCHECKED ←
DESIG=1;OUR EST; \rightarrow UNCHECKED ←
DESIG=3;OUR EST; \rightarrow UNCHECKED ←
DESIG=2;OUR EST; \rightarrow UNCHECKED ←
DESIG=5;OUR EST; \rightarrow UNCHECKED ←
DESIG=4;OUR EST; \rightarrow UNCHECKED ←
DESIG=7;OUR EST; \rightarrow UNCHECKED ←
DESIG=9;OUR EST; \rightarrow UNCHECKED ←
DESIG=10;OUR EST; \rightarrow UNCHECKED ←

NODE=M123225

NODE=M123W3
NODE=M123W3NODE=M123W1
NODE=M123W1NODE=M123W2
NODE=M123W2

NODE=M123W1;LINKAGE=SC

NODE=M123220

NODE=M123R5
NODE=M123R5NODE=M123R1
NODE=M123R1

NODE=M123R1;LINKAGE=A

NODE=M123R3
NODE=M123R3NODE=M123R2
NODE=M123R2NODE=M123R4
NODE=M123R4

NODE=M123R4;LINKAGE=E

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
seen	BAKER 99B	SPEC	$0 \bar{p} p \rightarrow \omega\omega\pi^0$	

 $f_2(1565)$ REFERENCES

ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev		
AMELIN 06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	
SCHEGELSKY 06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>		
AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	
BAKER 99B	PL B467 147	C.A. Baker <i>et al.</i>		
BERTIN 98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	
BERTIN 97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	
ABELE 96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	
AMSLER 95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	
AMSLER 95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	
AMSLER 95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	
BALOSHIN 95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	
Translated from YAF 69 715.				
AMSLER 94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	
ANISOVICH 94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	
ANISOVICH 94B	PR D50 1972	V.V. Anisovich <i>et al.</i>	(LOQM)	
ADAMO 93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.)	
ARMSTRONG 93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	
ARMSTRONG 93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	
WEIDENAUER 93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	
ADAMO 92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)	
BELADIDZE 92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)	
AKER 91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	
MAY 90	ZPHY C46 203	B. May <i>et al.</i>	(ASTERIX Collab.)	
MAY 89	PL B225 450	B. May <i>et al.</i>	(ASTERIX Collab.) IJP	
BRIDGES 86B	PRL 56 215	D.L. Bridges <i>et al.</i>	(SYRA, CASE)	
BRIDGES 86C	PRL 57 1534	D.L. Bridges <i>et al.</i>	(SYRA)	
Translated from YAF 58 50.				

 $\rho(1570)$

$$\Gamma^G(J^{PC}) = 1^+(1^-^-)$$

OMITTED FROM SUMMARY TABLE

May be an OZI-violating decay mode of $\rho(1700)$. See our mini-review under the $\rho(1700)$.

 $\rho(1570)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1570±36±62	54	¹ AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi\pi^0\gamma$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
1480±40		² BITYUKOV	87	SPEC	$32.5 \pi^- p \rightarrow \phi\pi^0 n$

¹ From the fit with two resonances.

² Systematic errors not estimated.

 $\rho(1570)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
144±75±43	54	³ AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi\pi^0\gamma$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
130±60		⁴ BITYUKOV	87	SPEC	$32.5 \pi^- p \rightarrow \phi\pi^0 n$

³ From the fit with two resonances.

⁴ Systematic errors not estimated.

 $\rho(1570)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	
$\Gamma_2 \phi\pi$	not seen
$\Gamma_3 \omega\pi$	

NODE=M123R6
NODE=M123R6

NODE=M123

REFID=52719
REFID=51574

REFID=51185
REFID=48580
REFID=47432
REFID=47398
REFID=45782
REFID=45701
REFID=45076
REFID=44377
REFID=44440
REFID=44441
REFID=44621

REFID=44093
REFID=43659
REFID=44071
REFID=43657
REFID=43587
REFID=43596
REFID=43585
REFID=42177
REFID=42172
REFID=41587
REFID=41365
REFID=40921
REFID=21376
REFID=21377

NODE=M188

NODE=M188

NODE=M188M

NODE=M188M

NODE=M188M;LINKAGE=AU
NODE=M188M;LINKAGE=BI

NODE=M188W

NODE=M188W

NODE=M188W;LINKAGE=AU
NODE=M188W;LINKAGE=BI

NODE=M188215;NODE=M188

DESIG=1
DESIG=2
DESIG=3

$\rho(1570) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\phi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$			$\Gamma_2\Gamma_1/\Gamma$		
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±0.9±0.3		54	5 AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<70	90	6 AULCHENKO	87B ND	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$	

5 From the fit with two resonances.

6 Using mass and width of BITYUKOV 87.

 $\rho(1570)$ BRANCHING RATIOS

$\Gamma(\phi\pi)/\Gamma_{\text{total}}$			Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.01	7 DONNACHIE	91 RVUE			
7 Using data from BISELLO 91B, DOLINSKY 86, and ALBRECHT 87L.					
$\Gamma(\phi\pi)/\Gamma(\omega\pi)$			Γ_2/Γ_3		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.5	95	BITYUKOV 87	SPEC	$32.5 \pi^- p \rightarrow \phi\pi^0 n$	

 $\rho(1570)$ REFERENCES

AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AULCHENKO	87B	JETPL 45 145 Translated from ZETFP 45 118.	V.M. Aulchenko <i>et al.</i>	(NOVO)
BITYUKOV	87	PL B188 383	S.I. Bityukov <i>et al.</i>	(SERP)
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)

 $h_1(1595)$

$I^G(J^{PC}) = 0^-(1^{+-})$

OMITTED FROM SUMMARY TABLE

Seen in a partial-wave analysis of the $\omega\eta$ system produced in the reaction $\pi^- p \rightarrow \omega\eta n$ at 18 GeV/c.

 $h_1(1595)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1594±15⁺¹⁰₋₆₀	EUGENIO 01	SPEC	$18 \pi^- p \rightarrow \omega\eta n$

 $h_1(1595)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
384±60⁺⁷⁰₋₁₀₀	EUGENIO 01	SPEC	$18 \pi^- p \rightarrow \omega\eta n$

 $h_1(1595)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \omega\eta$	seen

 $h_1(1595)$ REFERENCES

EUGENIO 01	PL B497 190	P. Eugenio <i>et al.</i>
------------	-------------	--------------------------

NODE=M188225

NODE=M188G01

NODE=M188G01

NODE=M188G01;LINKAGE=AU

NODE=M188G01;LINKAGE=AL

NODE=M188220

NODE=M188R01

NODE=M188R01

NODE=M188R01;LINKAGE=DO

NODE=M188R02

NODE=M188R02

NODE=M188

REFID=52242

REFID=45765

REFID=41752

REFID=41632

REFID=40418

REFID=41373

REFID=40011

REFID=20246

NODE=M166

NODE=M166

NODE=M166M

NODE=M166M

NODE=M166W

NODE=M166W

NODE=M166215;NODE=M166

DESIG=1;OUR EST;→ UNCHECKED ←

NODE=M166

REFID=48010

NODE=M164

 $\pi_1(1600)$ $I^G(J^{PC}) = 1^-(1^-+)$ **$\pi_1(1600)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

 1660 ± 15 OUR AVERAGE

Error includes scale factor of 1.2.

$1564 \pm 24 \pm 86$	19 ± 86	¹ RODAS	19 JPAC	$191 \pi^- p \rightarrow \eta(\prime) \pi^- p$
1600 ± 110	$46M$	² AGHASYAN	18B COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$1664 \pm 8 \pm 10$	$145k$	³ LU	05 B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$1709 \pm 24 \pm 41$	$69k$	⁴ KUHN	04 B852	$18 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
$1597 \pm 10 \pm 45$		⁴ IVANOV	01 B852	$18 \pi^- p \rightarrow \eta' \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1660 \pm 10 \pm 0$	$420k$	⁵ ALEKSEEV	10 COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$1593 \pm 8 \pm 29$	4.6 ADAMS		98B B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

1 The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15data.
The mass is extracted from the T-matrix pole.

2 Statistical error negligible.

3 May be a different state: natural and unnatural parity exchanges.

4 Natural parity exchange.

5 Superseded by AGHASYAN 2018B.

6 Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with
2.6 M events of $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852
data.

NODE=M164M

NODE=M164M

NODE=M164M;LINKAGE=D

NODE=M164M;LINKAGE=B

NODE=M164M;LINKAGE=LU

NODE=M164M;LINKAGE=A

NODE=M164M;LINKAGE=C

NODE=M164M;LINKAGE=DZ

 $\pi_1(1600)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

 257 ± 60 OUR AVERAGE

Error includes scale factor of 1.9. See the ideogram below.

$492 \pm 54 \pm 102$		¹ RODAS	19 JPAC	$191 \pi^- p \rightarrow \eta(\prime) \pi^- p$
580 ± 100	$46M$	² AGHASYAN	18B COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$185 \pm 25 \pm 28$	$145k$	³ LU	05 B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$403 \pm 80 \pm 115$	$69k$	⁴ KUHN	04 B852	$18 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
$340 \pm 40 \pm 50$		⁴ IVANOV	01 B852	$18 \pi^- p \rightarrow \eta' \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$269 \pm 21 \pm 42$	$420k$	⁵ ALEKSEEV	10 COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$168 \pm 20 \pm 150$		^{4,6} ADAMS	98B B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

1 The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15data.
The width is extracted from the T-matrix pole.

2 Statistical error negligible.

3 May be a different state: natural and unnatural parity exchanges.

4 Natural parity exchange.

5 Superseded by AGHASYAN 2018B.

6 Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with
2.6 M events of $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852
data.

NODE=M164W

NODE=M164W

NODE=M164W;LINKAGE=D


NODE=M164W;LINKAGE=B

NODE=M164W;LINKAGE=LU

NODE=M164W;LINKAGE=A

NODE=M164W;LINKAGE=C

NODE=M164W;LINKAGE=DZ



$\pi_1(1600)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi\pi$	seen
$\Gamma_2 \rho^0\pi^-$	seen
$\Gamma_3 f_2(1270)\pi^-$	not seen
$\Gamma_4 b_1(1235)\pi$	seen
$\Gamma_5 \eta'(958)\pi^-$	seen
$\Gamma_6 f_1(1285)\pi$	seen

$\pi_1(1600)$ BRANCHING RATIOS

$\Gamma(\rho^0\pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	ALEKSEEV 10 COMP 190 $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p b'$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
not seen	NOZAR 09 CLAS $\gamma p \rightarrow 2\pi^+ \pi^- n$
not seen	¹ DZIERBA 06 B852 18 $\pi^- p$

¹ From the PWA analysis of 2.6 M $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data. Supersedes ADAMS 98B.

$\Gamma(f_2(1270)\pi^-)/\Gamma_{\text{total}}$	Γ_3/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
not seen	¹ DZIERBA 06 B852 18 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
1 From the PWA analysis of 2.6 M $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data. Supersedes CHUNG 02.	

$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$	Γ_4/Γ
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	35280 1 BAKER 03 SPEC $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	145k LU 05 B852 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
	1 B(($b_1\pi$) _{D-wave})/B(($b_1\pi$) _{S-wave})=0.3 ± 0.1.

$\Gamma(\eta'(958)\pi^-)/\Gamma_{\text{total}}$	Γ_5/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	IVANOV 01 B852 18 $\pi^- p \rightarrow \eta' \pi^- p$

NODE=M164215;NODE=M164

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2
DESIG=4
DESIG=5
DESIG=3
DESIG=6;OUR EST;→ UNCHECKED ←

NODE=M164220

NODE=M164R1
NODE=M164R1

NODE=M164R1;LINKAGE=DZ

NODE=M164R3
NODE=M164R3

NODE=M164R3;LINKAGE=DZ

NODE=M164R4
NODE=M164R4

NODE=M164R;LINKAGE=RB

NODE=M164R2
NODE=M164R2

$\Gamma(f_1(1285)\pi)/\Gamma(\eta'(958)\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_5
3.80±0.78	69k	1 KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$	

1 Using $\eta'(958)\pi$ data from IVANOV 01. $\pi_1(1600)$ REFERENCES

RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
NOZAR	09	PRL 102 102002	M. Nozar <i>et al.</i>	(JLab CLAS Collab.)
DZIERBA	06	PR D73 072001	A.R. Dzierba <i>et al.</i>	(BNL E852 Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)
ADAMS	98B	PRL 81 5760	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)

 $a_1(1640)$

$I^G(J^{PC}) = 1^-(1^{++})$

Possibly seen in the study of the hadronic structure in decay $\tau \rightarrow 3\pi\nu_\tau$ (ABREU 98G and ASNER 00). $a_1(1640)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1655± 16 OUR AVERAGE				Error includes scale factor of 1.2.
1700 ⁺³⁵ -130	46M	1 AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^-\pi^+\pi^-\pi^- p$
1691± 18±30		DARGENT	17 RVUE	$D^0 \rightarrow \pi^-\pi^+\pi^-\pi^+$
1630± 20	35k	2 BAKER	03 SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
1714± 9±36		CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
1640± 12±30		BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1670± 90		BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^-\pi^+\pi^- A$

1 Statistical error negligible.

2 Using the $a_1(1260)$ mass and width results of BOWLER 88. $a_1(1640)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
254± 40 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
510 ⁺¹⁷⁰ -90	46M	1 AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^-\pi^+\pi^-\pi^- p$
171± 33±40		DARGENT	17 RVUE	$D^0 \rightarrow \pi^-\pi^+\pi^-\pi^+$
225± 30	35k	2 BAKER	03 SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$
308± 37±62		CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
300± 22±40		BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
300±100		BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^-\pi^+\pi^- A$

1 Statistical error negligible.

2 Using the $a_1(1260)$ mass and width results of BOWLER 88.NODE=M164R5
NODE=M164R5

NODE=M164R;LINKAGE=KU

NODE=M164

REFID=59554
REFID=59471
REFID=56385
REFID=53356
REFID=52758
REFID=51077
REFID=50459
REFID=49773
REFID=49414
REFID=48837
REFID=48317
REFID=46610

NODE=M161

NODE=M161

NODE=M161M


NODE=M161M

NODE=M161M;LINKAGE=A
NODE=M161M;LINKAGE=KB

NODE=M161W

NODE=M161W

NODE=M161W;LINKAGE=A
NODE=M161W;LINKAGE=KB



a₁(1640) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi\pi$	seen
$\Gamma_2 f_2(1270)\pi$	seen
$\Gamma_3 \sigma\pi$	seen
$\Gamma_4 \rho\pi S-wave$	seen
$\Gamma_5 \rho\pi D-wave$	seen
$\Gamma_6 \omega\pi\pi$	seen
$\Gamma_7 f_1(1285)\pi$	seen
$\Gamma_8 a_1(1260)\eta$	not seen

a₁(1640) BRANCHING RATIOS

$\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$	Γ_2/Γ_3
VALUE 0.24±0.07	DOCUMENT ID BAKER TECN COMMENT 1.94 $\bar{p}p \rightarrow 4\pi^0$

$\Gamma(\rho\pi D-wave)/\Gamma_{total}$	Γ_5/Γ
VALUE • • • We do not use the following data for averages, fits, limits, etc. • • •	DOCUMENT ID TECN COMMENT

seen CHUNG 02 B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
seen AMELIN 95B VES 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$

$\Gamma(\omega\pi\pi)/\Gamma_{total}$	Γ_6/Γ
VALUE • • • We do not use the following data for averages, fits, limits, etc. • • •	EVTS DOCUMENT ID TECN COMMENT

seen 35280 1 BAKER 03 SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

1 Assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.

$\Gamma(f_1(1285)\pi)/\Gamma_{total}$	Γ_7/Γ
VALUE • • • We do not use the following data for averages, fits, limits, etc. • • •	DOCUMENT ID TECN COMMENT

not seen KUHN 04 B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
seen LEE 94 MPS2 18 $\pi^- p \rightarrow K^+\bar{K}^0\pi^-\pi^- p$

$\Gamma(a_1(1260)\eta)/\Gamma_{total}$	Γ_8/Γ
VALUE not seen	DOCUMENT ID TECN COMMENT

KUHN 04 B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M161215;NODE=M161

DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=7;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=6;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=8

NODE=M161220

NODE=M161R1
NODE=M161R1

NODE=M161R2
NODE=M161R2

NODE=M161R3
NODE=M161R3

NODE=M161R;LINKAGE=KB

NODE=M161R4
NODE=M161R4

NODE=M161R5
NODE=M161R5

a₁(1640) REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
DARGENT	17	JHEP 1705 143	P. d'Argent <i>et al.</i>	(HEID, BRIS)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(Serp, TBIL)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
		Translated from YAF 41 1223.		

f₂(1640)

$$f_2(1640) = 0^+(2^{++})$$

OMMITTED FROM SUMMARY TABLE

f₂(1640) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1639± 6 OUR AVERAGE	Error includes scale factor of 1.2.		
1620± 16	BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1647± 7	ADAMO	92	OBLX $\bar{n}p \rightarrow 3\pi^+2\pi^-$
1635± 7	ALDE	90	GAM2 $38\pi^-p \rightarrow \omega\omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1640± 5	AMSLER	06	CBAR $0.9\bar{p}p \rightarrow K^+K^-\pi^0$
1659± 6	VLADIMIRSK...	06	SPEC $40\pi^-p \rightarrow K_S^0K_S^0n$
1643± 7	¹ ALDE	89B	GAM2 $38\pi^-p \rightarrow \omega\omega n$

¹ Superseded by ALDE 90.

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
99⁺⁶⁰₋₄₀ OUR AVERAGE	Error includes scale factor of 2.9.			
140 ⁺⁶⁰ ₋₂₀		BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
58±20		ADAMO	92	OBLX $\bar{n}p \rightarrow 3\pi^+2\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
44± 9		AMSLER	06	CBAR $0.9\bar{p}p \rightarrow K^+K^-\pi^0$
152±18		VLADIMIRSK...	06	SPEC $40\pi^-p \rightarrow K_S^0K_S^0n$
< 70	90	ALDE	90	GAM2 $38\pi^-p \rightarrow \omega\omega n$

f₂(1640) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega\omega$	seen
Γ_2 4π	seen
Γ_3 $K\bar{K}$	seen

f₂(1640) BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
seen	AMSLER	06	CBAR $0.9\bar{p}p \rightarrow K^+K^-\pi^0$

f₂(1640) REFERENCES

AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH) JP
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	89B	PL B216 451	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+) IGJPC

NODE=M161

REFID=59471
 REFID=58121
 REFID=49773
 REFID=49414
 REFID=48837
 REFID=47339
 REFID=46888
 REFID=45909
 REFID=44433
 REFID=44092
 REFID=40578
 REFID=47490

NODE=M117

NODE=M117M

NODE=M117M

NODE=M117M;LINKAGE=BB

NODE=M117W

NODE=M117W

NODE=M117215;NODE=M117

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=3

NODE=M117220

NODE=M117R2
 NODE=M117R2

NODE=M117

REFID=51136
 REFID=51191
 REFID=44438
 REFID=42177
 REFID=40935
 REFID=40735

$\eta_2(1645)$ $I^G(J^{PC}) = 0^+(2^-+)$

NODE=M154

 $\eta_2(1645)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1617± 5 OUR AVERAGE				
1613± 8	BARBERIS 00B		450	$p p \rightarrow p_f \eta \pi^+ \pi^- p_s$
1617± 8	BARBERIS 00C		450	$p p \rightarrow p_f 4\pi p_s$
1620±20	BARBERIS 97B OMEG		450	$p p \rightarrow p p 2(\pi^+ \pi^-)$
1645±14±15	ADOMEIT 96 CBAR 0	1.94	$\bar{p} p \rightarrow \eta 3\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1645± 6±20	ANISOVICH 00E SPEC		0.9–1.94	$\bar{p} p \rightarrow \eta 3\pi^0$

NODE=M154M

NODE=M154M

 $\eta_2(1645)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
181±11 OUR AVERAGE				
185±17	BARBERIS 00B		450	$p p \rightarrow p_f \eta \pi^+ \pi^- p_s$
177±18	BARBERIS 00C		450	$p p \rightarrow p_f 4\pi p_s$
180±25	BARBERIS 97B OMEG		450	$p p \rightarrow p p 2(\pi^+ \pi^-)$
180 ⁺⁴⁰ ₋₂₁ ±25	ADOMEIT 96 CBAR 0	1.94	$\bar{p} p \rightarrow \eta 3\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200±25	ANISOVICH 00E SPEC		0.9–1.94	$\bar{p} p \rightarrow \eta 3\pi^0$

NODE=M154W

NODE=M154W

 $\eta_2(1645)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 a_2(1320)\pi$	seen
$\Gamma_2 K\bar{K}\pi$	seen
$\Gamma_3 K^*\bar{K}$	seen
$\Gamma_4 \eta\pi^+\pi^-$	seen
$\Gamma_5 a_0(980)\pi$	seen
$\Gamma_6 f_2(1270)\eta$	not seen

NODE=M154215;NODE=M154

DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=2;OUR EST; \rightarrow UNCHECKED
 DESIG=3;OUR EST; \rightarrow UNCHECKED
 DESIG=4;OUR EST; \rightarrow UNCHECKED
 DESIG=5;OUR EST; \rightarrow UNCHECKED
 DESIG=6;OUR EST; \rightarrow UNCHECKED

 $\eta_2(1645)$ BRANCHING RATIOS

$\Gamma(K\bar{K}\pi)/\Gamma(a_2(1320)\pi)$	Γ_2/Γ_1
0.07±0.03	1 BARBERIS 97C OMEG 450 $p p \rightarrow p p K\bar{K}\pi$

¹ Using $2(\pi^+\pi^-)$ data from BARBERIS 97B.

NODE=M154220

NODE=M154R1
NODE=M154R1

$\Gamma(a_2(1320)\pi)/\Gamma(a_0(980)\pi)$	Γ_1/Γ_5
13.1±2.3 OUR AVERAGE	1 ANISOVICH 11 SPEC 0.9–1.94 $p\bar{p}$ 2 BARBERIS 00B 450 $p p \rightarrow p_f \eta \pi^+ \pi^- p_s$

NODE=M154R1;LINKAGE=A

NODE=M154R3
NODE=M154R3

$\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$	Γ_6/Γ
not seen	BARBERIS 00B 450 $p p \rightarrow p_f \eta \pi^+ \pi^- p_s$

NODE=M154R3;LINKAGE=AN

NODE=M154R4
NODE=M154R4 **$\eta_2(1645)$ REFERENCES**

ANISOVICH 11 EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
ANISOVICH 00E PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS 00B PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 00C PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 97B PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 97C PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ADOMEIT 96 ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)

NODE=M154

REFID=53631
 REFID=47945
 REFID=47958
 REFID=47959
 REFID=45758
 REFID=45759
 REFID=45202

$\omega(1650)$ $I^G(J^{PC}) = 0^-(1^- -)$

NODE=M126

 $\omega(1650)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1670 ± 30 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1651 \pm 3 \pm 16	183k	1 ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
1673 \pm 6		ACHASOV	19 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$
1671 \pm 6 \pm 10	824	2 AKHMETSHIN	17A CMD3	1.4–2.0 $e^+ e^- \rightarrow \omega \eta$
1660 \pm 10	898	3 ACHASOV	16B SND	1.34–2.00 $e^+ e^- \rightarrow \omega \eta$
1680 \pm 10	13.1k	4 AULCHENKO	15A SND	1.05–1.80 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1667 \pm 13 \pm 6		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
1645 \pm 8	13	AUBERT	06D BABR	10.6 $e^+ e^- \rightarrow \omega \eta \gamma$
1660 \pm 10 \pm 2		AUBERT,B	04N BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
1770 \pm 50 \pm 60	1.2M	5 ACHASOV	03D RVUE	0.44–2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1619 \pm 5		6 HENNER	02 RVUE	1.2–2.0 $e^+ e^- \rightarrow \rho \pi,$ $\omega \pi \pi$
1700 \pm 20		EUGENIO	01 SPEC	18 $\pi^- p \rightarrow \omega \eta \eta$
1705 \pm 26	612	7 AKHMETSHIN	00D CMD2	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1820 \pm 190		8 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1840 \pm 100		9 ACHASOV	98H RVUE	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$
1780 \pm 170		10 ACHASOV	98H RVUE	$e^+ e^- \rightarrow K^+ K^-$
~ 2100		11 ACHASOV	98H RVUE	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606 \pm 9		12 CLEGG	94 RVUE	
1662 \pm 13	750	13 ANTONELLI	92 DM2	1.34–2.4 $e^+ e^- \rightarrow \rho \pi,$ $\omega \pi \pi$
1670 \pm 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
1657 \pm 13		CORDIER	81 DM1	$e^+ e^- \rightarrow \omega 2\pi$
1679 \pm 34	21	ESPOSITO	80 FRAM	$e^+ e^- \rightarrow 3\pi$
1652 \pm 17		COSME	79 OSPK	$e^+ e^- \rightarrow 3\pi$

¹ Could also be $\rho(1700)$. Branching ratio $J/\psi \rightarrow X \pi^0 \rightarrow K^+ K^- \pi^0 = (5.3 \pm 0.3)^{+0.6}_{-0.5) \times 10^{-5}$.

² From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

³ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

⁴ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

⁵ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁶ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁷ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho \pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

⁸ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

⁹ Using the data from ANTONELLI 92.

¹⁰ Using the data from IVANOV 81 and BISELLO 88B.

¹¹ Using the data from BISELLO 91C.

¹² From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹³ From the combined fit of the $\rho \pi$ and $\omega \pi \pi$ final states.

NODE=M126M

NODE=M126M

→ UNCHECKED ←

 $\omega(1650)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
315 ± 35 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
194 \pm 8 \pm 15	183k	1 ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
95 \pm 11		ACHASOV	19 SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \eta$
113 \pm 9 \pm 10	824	2 AKHMETSHIN	17A CMD3	1.4–2.0 $e^+ e^- \rightarrow \omega \eta$
110 \pm 20	898	3 ACHASOV	16B SND	1.34–2.00 $e^+ e^- \rightarrow \omega \eta$

NODE=M126W

NODE=M126W

→ UNCHECKED ←

310 ± 30	13.1k	⁴ AULCHENKO	15A SND	$1.05\pi^+\pi^-\pi^0 \rightarrow$
222 ± 25 ± 20		AUBERT	07AU BABR	$10.6\pi^+\pi^- \rightarrow \omega\pi^+\pi^-\gamma$
114 ± 14	13	AUBERT	06D BABR	$10.6\pi^+\pi^- \rightarrow \omega\eta\gamma$
230 ± 30 ± 20		AUBERT,B	04N BABR	$10.6\pi^+\pi^- \rightarrow \pi^+\pi^-\pi^0\gamma$
$490^{+200}_{-150} \pm 130$	1.2M	⁵ ACHASOV	03D RVUE	$0.44\pi^+\pi^-\pi^0 \rightarrow$
250 ± 14		⁶ HENNER	02 RVUE	$1.2\pi^+\pi^- \rightarrow \rho\pi, \omega\pi\pi$
250 ± 50		EUGENIO	01 SPEC	$18\pi^-p \rightarrow \omega\eta n$
370 ± 25	612	⁷ AKHMETSHIN	00D CMD2	$\pi^+\pi^- \rightarrow \omega\pi^+\pi^-$
113 ± 20		⁸ CLEGG	94 RVUE	
280 ± 24	750	⁹ ANTONELLI	92 DM2	$1.34\pi^+\pi^- \rightarrow \rho\pi, \omega\pi\pi$
160 ± 20		ATKINSON	83B OMEG	$20\pi^-p \rightarrow 3\pi X$
136 ± 46		CORDIER	81 DM1	$\pi^+\pi^- \rightarrow \omega 2\pi$
99 ± 49	21	ESPOSITO	80 FRAM	$\pi^+\pi^- \rightarrow 3\pi$
42 ± 17		COSME	79 OSPK	$\pi^+\pi^- \rightarrow 3\pi$

1 Could also be $\rho(1700)$. Branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (5.3 \pm 0.3)^{+0.6}_{-0.5} \times 10^{-5}$.

2 From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

3 From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

4 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

5 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

6 Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

7 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

8 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

9 From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

OCCUR=2

OCCUR=5

OCCUR=4

NODE=M126W;LINKAGE=G

NODE=M126W;LINKAGE=F

NODE=M126W;LINKAGE=E

NODE=M126W;LINKAGE=A

NODE=M126W;LINKAGE=VH

NODE=M126W;LINKAGE=AB

NODE=M126W;LINKAGE=KI

NODE=M126W;LINKAGE=AD

NODE=M126W;LINKAGE=AE

NODE=M126215;NODE=M126

$\omega(1650)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi$	seen
$\Gamma_2 \omega\pi\pi$	seen
$\Gamma_3 \omega\eta$	seen
$\Gamma_4 e^+e^-$	seen
$\Gamma_5 \pi^0\gamma$	not seen

$\omega(1650) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.56 ± 0.23	13.1k	¹ AULCHENKO	15A SND	$1.05\pi^+\pi^-\pi^0 \rightarrow$
1.3 ± 0.1 ± 0.1		AUBERT,B	04N BABR	$10.6\pi^+\pi^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1.2 ± 0.4 ± 0.8	1.2M	^{2,3} ACHASOV	03D RVUE	$0.44\pi^+\pi^-\pi^0 \rightarrow$
0.921 ± 0.230		^{4,5} CLEGG	94 RVUE	
0.479 ± 0.050	750	^{6,7} ANTONELLI	92 DM2	$1.34\pi^+\pi^- \rightarrow \rho\pi, \omega\pi\pi$

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=5

NODE=M126230

NODE=M126G3

NODE=M126G3

1 From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

2 Calculated by us from the cross section at the peak.

3 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

4 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

5 From the partial and leptonic width given by the authors.

6 From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

7 From the product of the leptonic width and partial branching ratio given by the authors.

NODE=M126G3;LINKAGE=A
 NODE=M126G;LINKAGE=AW
 NODE=M126G;LINKAGE=VH

NODE=M126G;LINKAGE=AD

NODE=M126G;LINKAGE=SE

NODE=M126G;LINKAGE=AE

NODE=M126G;LINKAGE=ES

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.0 \pm 0.5		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
4.1 \pm 0.9 \pm 1.3	1.2M	1,2 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 \pm 0.95		3 AKHMETSHIN 00D	CMD2	$1.2-1.38 e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 \pm 0.80		4,5 CLEGG	94	RVUE
6.07 \pm 0.61	750	6,7 ANTONELLI	92	DM2 $1.34-2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

1 Calculated by us from the cross section at the peak.

2 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.3 Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

4 From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

5 From the partial and leptonic width given by the authors.

6 From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

7 From the product of the leptonic width and partial branching ratio given by the authors.

 $\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_4/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

$5.62^{+0.45}_{-0.42}$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
$4.5 \pm 0.3 \pm 0.3$	824	1 AKHMETSHIN 17A	CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$
4.4 ± 0.5	898	2 ACHASOV	16B SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$
5.7 ± 0.6	13	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
< 60 at 90% CL		3 AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$

1 From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1650) \rightarrow \omega\eta) / \Gamma_{\text{total}} \times \Gamma(\omega(1650) \rightarrow e^+e^-) = 51 \pm 3$ eV.2 From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.3 $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively. **$\omega(1650)$ BRANCHING RATIOS**
 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

~ 0.65	1.2M	1 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.380 ± 0.014		2 HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

1 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.2 Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.
 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

~ 0.35	1.2M	1 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.620 ± 0.014		2 HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

1 From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.2 Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.
 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

~ 18	1.2M	1,2 ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
32 ± 1		2 HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

1 Calculated by us from the cross section at the peak.

2 Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

NODE=M126G4

NODE=M126G4

NODE=M126G4;LINKAGE=AW

NODE=M126G4;LINKAGE=VH

NODE=M126G4;LINKAGE=KL

NODE=M126G4;LINKAGE=AD

NODE=M126G4;LINKAGE=SE

NODE=M126G4;LINKAGE=AE

NODE=M126G4;LINKAGE=ES

NODE=M126G5

NODE=M126G5

NODE=M126G5;LINKAGE=B

NODE=M126G5;LINKAGE=A

NODE=M126G5;LINKAGE=KH

NODE=M126225

NODE=M126R3

NODE=M126R3

NODE=M126R;LINKAGE=VH

NODE=M126R;LINKAGE=AC

NODE=M126R2

NODE=M126R2

NODE=M126R2;LINKAGE=VH

NODE=M126R2;LINKAGE=AC

NODE=M126R4

NODE=M126R4

NODE=M126R;LINKAGE=AW

NODE=M126R4;LINKAGE=AC

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
not seen	1 ACHASOV	10D	SND 1.075–2.0 e ⁺ e ⁻ → $\pi^0\gamma$	

¹ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. The width of the highest mass effective resonance is fixed at 315 MeV.

 $\omega(1650)$ REFERENCES

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)
		Translated from ZETF 148 34.		
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>	(NOVO)
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov	(NOVO)
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)
		Translated from ZETFP 46 132.		
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORsay)
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)

 $\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^- -)$$

 $\omega_3(1670)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1667 ± 4 OUR AVERAGE				
1665.3 ± 5.2 ± 4.5	23400	AMELIN	96	VES $36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1685 ± 20	60	BAUBILLIER	79	HBC 8.2 $K^- p$ backward
1673 ± 12	430	1,2 BALTAY	78E	HBC 15 $\pi^+ p \rightarrow \Delta 3\pi$
1650 ± 12		CORDEN	78B	OMEG 8–12 $\pi^- p \rightarrow N 3\pi$
1669 ± 11	600	2 WAGNER	75	HBC 7 $\pi^+ p \rightarrow \Delta^{++} 3\pi$
1678 ± 14	500	DIAZ	74	DBC 6 $\pi^+ n \rightarrow p 3\pi^0$
1660 ± 13	200	DIAZ	74	DBC 6 $\pi^+ n \rightarrow p \omega \pi^0 \pi^0$
1679 ± 17	200	MATTHEWS	71D	DBC 7.0 $\pi^+ n \rightarrow p 3\pi^0$
1670 ± 20		KENYON	69	DBC 8 $\pi^+ n \rightarrow p 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1700	110	¹ CERRADA	77B	HBC 4.2 $K^- p \rightarrow \Lambda 3\pi$
1695 ± 20		BARNES	69B	HBC 4.6 $K^- p \rightarrow \omega 2\pi X$
1636 ± 20		ARMENISE	68B	DBC 5.1 $\pi^+ n \rightarrow p 3\pi^0$

¹ Phase rotation seen for $J^P = 3^-$ $\rho\pi$ wave.

² From a fit to $I(J^P) = 0(3^-)$ $\rho\pi$ partial wave.

 $\omega_3(1670)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
168 ± 10 OUR AVERAGE				
149 ± 19 ± 7	23400	AMELIN	96	VES $36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
160 ± 80	60	³ BAUBILLIER	79	HBC 8.2 $K^- p$ backward
173 ± 16	430	^{4,5} BALTAY	78E	HBC 15 $\pi^+ p \rightarrow \Delta 3\pi$
253 ± 39		CORDEN	78B	OMEG 8–12 $\pi^- p \rightarrow N 3\pi$
173 ± 28	600	^{3,5} WAGNER	75	HBC 7 $\pi^+ p \rightarrow \Delta^{++} 3\pi$
167 ± 40	500	DIAZ	74	DBC 6 $\pi^+ n \rightarrow p 3\pi^0$
122 ± 39	200	DIAZ	74	DBC 6 $\pi^+ n \rightarrow p \omega \pi^0 \pi^0$
155 ± 40	200	³ MATTHEWS	71D	DBC 7.0 $\pi^+ n \rightarrow p 3\pi^0$

NODE=M126R00
NODE=M126R00

NODE=M126R00;LINKAGE=A

NODE=M126

REFID=59909
REFID=59887
REFID=58239
REFID=57537
REFID=56843

REFID=59494
REFID=52049
REFID=51047
REFID=50184
REFID=49577
REFID=49406
REFID=48815
REFID=49177
REFID=48311
REFID=48010
REFID=47935
REFID=47391
REFID=46323
REFID=44081
REFID=43168
REFID=41867
REFID=41369
REFID=40581
REFID=40280

NODE=M045

NODE=M045M

NODE=M045M

OCCUR=2

NODE=M045M;LINKAGE=E
NODE=M045M;LINKAGE=P

NODE=M045W

NODE=M045W

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

90 ± 20	BARNES	69B	HBC	$4.6 K^- p \rightarrow \omega 2\pi$
100 ± 40	KENYON	69	DBC	$8 \pi^+ n \rightarrow p 3\pi^0$
$112 + 60$	ARMENISE	68B	DBC	$5.1 \pi^+ n \rightarrow p 3\pi^0$

³Width errors enlarged by us to $4\Gamma/\sqrt{N}$: see the note with the $K^*(892)$ mass.

⁴ Phase rotation seen for $J^P = 3^-$ $\rho\pi$ wave

⁵ From a fit to $I(P) = 0(3^-)$ $\sigma\pi$ partial wave.

NODE=M045W;LINKAGE=S
NODE=M045W;LINKAGE=E
NODE=M045W;LINKAGE=P

$\omega_3(1670)$ DECAY MODES

Mode	Fraction (Γ_j/Γ)
$\Gamma_1 \quad \rho\pi$	seen
$\Gamma_2 \quad \omega\pi\pi$	seen
$\Gamma_3 \quad b_1(1235)\pi$	possibly seen

$\omega_3(1670)$ BRANCHING RATIOS

$\Gamma(\omega\pi\pi)/\Gamma(\rho\pi)$	$EVTS$	$DOCUMENT_ID$	$TECN$	$COMMENT$	Γ_2/Γ_1
VALUE					

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.71 ± 0.27 100 DIAZ 74 DBC $6\pi^+\eta \rightarrow p5\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←

NODE=M045220

NODE=M045R3
NODE=M045R3

possibly seen DIAZ 74 DBC 0 π · II -

$\Gamma(b_1(1235)\pi)/\Gamma(\omega\pi\pi)$	Γ_3/Γ_2
<i>value</i>	<i>CL%</i>

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.75 68 BAUBILLIER 79 HBC 8.2 $K^- p$ backward

$\omega_3(1670)$ REFERENCES

AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BAUBILLIER	79	PL 89B 131	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BALTAY	78E	PRL 40 87	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
CORDEN	78B	NP B138 235	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+) JP
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)
MATTHEWS	71D	PR D3 2561	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC)
BARNES	69B	PRL 23 142	V.E. Barnes <i>et al.</i>	(BNL)
KENYON	69	PRL 23 146	I.R. Kenyon <i>et al.</i>	(BNL, UCND, ORNL)
ARMENISE	68B	PL 26B 336	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)

NODE=M045R4
NODE=M045R4

NODE=M045R5
NODE=M045R5

NODE=M045

REFID=44649
REFID=21522
REFID=21520
REFID=21269
REFID=20537
REFID=20843
REFID=21248
REFID=21515
REFID=21512
REFID=20800
REFID=20783

NODE=M034

 $\pi_2(1670)$ $I^G(J^{PC}) = 1^-(2^-+)$ **$\pi_2(1670)$ MASS**

NODE=M034M

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

CHG

COMMENT

NODE=M034M

1670.6 \pm 2.9 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

1642	± 12	46M	1 AGHASYAN	18B COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	
1749	± 10	± 100	145k	LU 05 B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$	
1676	± 3	± 8	2 CHUNG	02 B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
1685	± 10	± 30	BARBERIS	01	$450 pp \rightarrow p_f 3\pi^0 p_s$	
1687	± 9	± 15	AMELIN	99 VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$	
1669	± 4		BARBERIS	98B	$450 pp \rightarrow p_f \rho \pi p_s$	
1670	± 4		BARBERIS	98B	$450 pp \rightarrow p_f f_2(1270)\pi p_s$	OCCUR=2
1690	± 14		3 BERDNIKOV	94 VES	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$	
1710	± 20	700	ANTIPOV	87 SIGM -	$50 \pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$	
1676	± 6		3 EVANGELIS...	81 OMEG -	$12 \pi^- p \rightarrow 3\pi p$	
1657	± 14		3,4 DAUM	80D SPEC -	$63-94 \pi p \rightarrow 3\pi X$	
1662	± 10	2000	3 BALTAY	77 HBC +	$15 \pi^+ p \rightarrow p 3\pi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1658	± 3	± 24	420k	5 ALEKSEEV	10 COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
1730	± 20			6 AMELIN	95B VES	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
1742	± 31	± 49		ANTREASYAN	90 CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$	
1624	± 21			2 BELLINI	85 SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1622	± 35			7 BELLINI	85 SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	OCCUR=2
1693	± 28			8 BELLINI	85 SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	OCCUR=3
1710	± 20			9 DAUM	81B SPEC -	$63,94 \pi^- p$	
1660	± 10			3 ASCOLI	73 HBC -	$5-25 \pi^- p \rightarrow p \pi_2$	

1 Statistical error negligible.

2 From $f_2(1270)\pi$ decay.3 From a fit to $J^P = 2^- S$ -wave $f_2(1270)\pi$ partial wave.4 Clear phase rotation seen in $2^- S$, $2^- P$, $2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.

5 Superseded by AGHASYAN 2018B.

6 J^{PC} ambiguous.7 From $\rho \pi$ decay.8 From $\sigma \pi$ decay.9 From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.

NODE=M034M;LINKAGE=A

NODE=M034M;LINKAGE=F2

NODE=M034M;LINKAGE=P

NODE=M034M;LINKAGE=D


NODE=M034M;LINKAGE=B

NODE=M034M;LINKAGE=AX

NODE=M034M;LINKAGE=R2

NODE=M034M;LINKAGE=S2

NODE=M034M;LINKAGE=L

 **$\pi_2(1670)$ WIDTH**

NODE=M034W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	NODE
258^{+ 8}_{- 9} OUR AVERAGE					Error includes scale factor of 1.2.	

311 ^{+ 12} _{- 23}	46M	10 AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
408 ^{± 60 ± 250}	145k	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
254 ^{± 3 ± 31}		11 CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
265 ^{± 30 ± 40}		BARBERIS	01	450 $p p \rightarrow p_f 3\pi^0 p_s$
168 ^{± 43 ± 53}		AMELIN	99 VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
268 ^{± 15}		BARBERIS	98B	450 $p p \rightarrow p_f \rho \pi p_s$
256 ^{± 15}		BARBERIS	98B	450 $p p \rightarrow p_f f_2(1270) \pi p_s$
190 ^{± 50}		12 BERDNIKOV	94 VES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
170 ^{± 80}	700	ANTIPOV	87 SIGM	50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
260 ^{± 20}		12 EVANGELIS...	81 OMEG	12 $\pi^- p \rightarrow 3\pi p$
219 ^{± 20}		12,13 DAUM	80D SPEC	63–94 $\pi p \rightarrow 3\pi X$
285 ^{± 60}	2000	12 BALTAY	77 HBC	15 $\pi^+ p \rightarrow p 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

271 ^{± 9_{- 24}}	420k	14 ALEKSEEV	10 COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
310 ^{± 20}		15 AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
236 ^{± 49 ± 36}		ANTREASYAN	90 CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
304 ^{± 22}		11 BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
404 ^{± 108}		16 BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
330 ^{± 90}		17 BELLINI	85 SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
312 ^{± 50}		18 DAUM	81B SPEC	63,94 $\pi^- p$
270 ^{± 60}		12 ASCOLI	73 HBC	5–25 $\pi^- p \rightarrow p \pi_2$

10 Statistical error negligible.

11 From $f_2(1270)\pi$ decay.

12 From a fit to $J^P = 2^- f_2(1270)\pi$ partial wave.

13 Clear phase rotation seen in $2^- S$, $2^- P$, $2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.

14 Superseded by AGHASYAN 2018B.

15 J^P ambiguous.

16 From $\rho\pi$ decay.

NODE=M034W;LINKAGE=A

NODE=M034W;LINKAGE=F2

NODE=M034W;LINKAGE=P

NODE=M034W;LINKAGE=D

NODE=M034W;LINKAGE=B

NODE=M034W;LINKAGE=AX

NODE=M034W;LINKAGE=R2

17 From $\sigma\pi$ decay.18 From a two-resonance fit to four 2^-0^+ waves. This should not be averaged with all the single resonance fits. **$\pi_2(1670)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 3\pi$	$(95.8 \pm 1.4) \%$	
$\Gamma_2 \pi^+ \pi^- \pi^0$		
$\Gamma_3 \pi^0 \pi^0 \pi^0$		
$\Gamma_4 f_2(1270)\pi$	$(56.3 \pm 3.2) \%$	
$\Gamma_5 \rho\pi$	$(31 \pm 4) \%$	
$\Gamma_6 \sigma\pi$	$(10 \pm 4) \%$	
$\Gamma_7 \pi(\pi\pi)_{S\text{-wave}}$	$(8.7 \pm 3.4) \%$	
$\Gamma_8 \pi^\pm \pi^+ \pi^-$	$(53 \pm 4) \%$	
$\Gamma_9 K\bar{K}^*(892) + \text{c.c.}$	$(4.2 \pm 1.4) \%$	
$\Gamma_{10} \omega\rho$	$(2.7 \pm 1.1) \%$	
$\Gamma_{11} \pi^\pm \gamma$	$(7.0 \pm 1.2) \times 10^{-4}$	
$\Gamma_{12} \gamma\gamma$	$< 2.8 \times 10^{-7}$	90%
$\Gamma_{13} \eta\pi$	$< 5 \%$	
$\Gamma_{14} \pi^\pm 2\pi^+ 2\pi^-$	$< 5 \%$	
$\Gamma_{15} \rho(1450)\pi$	$< 3.6 \times 10^{-3}$	97.7%
$\Gamma_{16} b_1(1235)\pi$	$< 1.9 \times 10^{-3}$	97.7%
$\Gamma_{17} \eta 3\pi$		
$\Gamma_{18} f_1(1285)\pi$	possibly seen	
$\Gamma_{19} a_2(1320)\pi$	not seen	

NODE=M034W;LINKAGE=S2

NODE=M034W;LINKAGE=L

NODE=M034215;NODE=M034

DESIG=20

DESIG=22

DESIG=23

DESIG=8

DESIG=2

DESIG=13

DESIG=11

DESIG=10

DESIG=5

DESIG=14

DESIG=27

DESIG=12

DESIG=3

DESIG=4

DESIG=15

DESIG=16

DESIG=24

DESIG=25

DESIG=26

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.9$ for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|ccc} & & & \\ x_5 & -53 & & \\ x_7 & -29 & -59 & \\ x_9 & -8 & -21 & -9 \\ & x_4 & x_5 & x_7 \end{array}$$

 $\pi_2(1670)$ PARTIAL WIDTHS

$\Gamma(\pi^\pm\gamma)$				Γ_{11}
VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
$181 \pm 11 \pm 27$	19 ADOLPH	14 COMP	-	190 π^- Pb $\rightarrow \pi^+ \pi^- \pi^-$ Pb'

19 Primakoff reaction. Assumes incoherent $f_2(1270)\pi$ contribution to 3π final state and uses $B(\pi_2(1670) \rightarrow f_2\pi) = 56\%$.

NODE=M034217

NODE=M034W2

NODE=M034W2

NODE=M034W2;LINKAGE=AD

$\Gamma(\gamma\gamma)$				Γ_{12}
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.072	90	20 ACCIARRI	97T L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc.				• • •
<0.19	90	20 ALBRECHT	97B ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
1.41 $\pm 0.23 \pm 0.28$		ANTREASYAN 90	CBAL 0	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
0.8 $\pm 0.3 \pm 0.12$		21 BEHREND	90C CELL 0	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
1.3 $\pm 0.3 \pm 0.2$		22 BEHREND	90C CELL 0	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

NODE=M034W1

NODE=M034W1

OCCUR=2

NODE=M034W1;LINKAGE=QQ

NODE=M034W1;LINKAGE=C

NODE=M034W1;LINKAGE=G

20 Decaying into $f_2(1270)\pi$ and $\rho\pi$.21 Constructive interference between $f_2(1270)\pi, \rho\pi$ and background.

22 Incoherent Ansatz.

$\pi_2(1670) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_{12}/\Gamma$
VALUE (keV) CL% DOCUMENT ID TECN COMMENT <0.1 95 23 SCHEGELSKY 06 RVUE $\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	

23 From analysis of L3 data at 183–209 GeV.

 $\pi_2(1670)$ BRANCHING RATIOS

$\Gamma(3\pi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma = (\Gamma_4 + \Gamma_5 + \Gamma_7)/\Gamma$
VALUE DOCUMENT ID 0.958±0.014 OUR FIT	

$\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$	Γ_3/Γ_2
VALUE DOCUMENT ID COMMENT 0.29±0.03±0.05 BARBERIS 01 450 $p p \rightarrow p_f 3\pi^0 p_s$	

$\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi)$	$\Gamma_5/0.565\Gamma_4$
(With $f_2(1270) \rightarrow \pi^+\pi^-$)	
VALUE DOCUMENT ID TECN COMMENT 0.97±0.09 OUR AVERAGE	Error includes scale factor of 1.9.
0.76±0.07±0.10 CHUNG 02 B852 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$	
1.01±0.05 BARBERIS 98B 450 $p p \rightarrow p_f \pi^+\pi^-\pi^0 p_s$	

$\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi)$	Γ_6/Γ_4
VALUE DOCUMENT ID TECN COMMENT 0.17±0.02±0.07 CHUNG 02 B852 18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	

0.24±0.10 24,25 BAKER 99 SPEC 1.94 $\bar{p} p \rightarrow 4\pi^0$

$\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$	$\frac{1}{2}\Gamma_5/\Gamma_8 = \frac{1}{2}\Gamma_5/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$
VALUE DOCUMENT ID TECN CHG COMMENT 0.29±0.04 OUR FIT	
0.29±0.05 26 DAUM 81B SPEC 63,94 $\pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<0.3 BARTSCH 68 HBC + 8 $\pi^+ p \rightarrow 3\pi p$	

$0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$	
(With $f_2(1270) \rightarrow \pi^+\pi^-$)	$0.565\Gamma_4/\Gamma_8 = 0.565\Gamma_4/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$
VALUE DOCUMENT ID TECN CHG COMMENT 0.604±0.035 OUR FIT	Error includes scale factor of 1.3.
0.60 ± 0.05 OUR AVERAGE	
0.61 ± 0.04 26 DAUM 81B SPEC 63,94 $\pi^- p$	
0.76 +0.24 -0.34 ARMENISE 69 DBC + 5.1 $\pi^+ d \rightarrow d 3\pi$	
0.35 ± 0.20 BALTAY 68 HBC + 7–8.5 $\pi^+ p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.59 BARTSCH 68 HBC + 8 $\pi^+ p \rightarrow 3\pi p$	

$0.624\Gamma(\pi(\pi\pi)_S\text{-wave})/\Gamma(\pi^\pm\pi^+\pi^-)$	
(With $(\pi\pi)_S\text{-wave} \rightarrow \pi^+\pi^-$)	$0.624\Gamma_7/\Gamma_8 = 0.624\Gamma_7/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$
VALUE DOCUMENT ID TECN COMMENT 0.10±0.04 OUR FIT	
0.10±0.05 26 DAUM 81B SPEC 63,94 $\pi^- p$	

$\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(f_2(1270)\pi)$	Γ_9/Γ_4
VALUE DOCUMENT ID TECN CHG COMMENT 0.075±0.025 OUR FIT	

0.075±0.025 27 ARMSTRONG 82B OMEG – 16 $\pi^- p \rightarrow K^+ K^- \pi^- p$

$\Gamma(\omega\rho)/\Gamma_{\text{total}}$	Γ_{10}/Γ
VALUE DOCUMENT ID TECN COMMENT 0.027±0.004±0.010 28 AMELIN 99 VES 37 $\pi^- A \xrightarrow{\omega\pi^-} \pi^0 A^*$	

NODE=M034230

NODE=M034G01
NODE=M034G01

NODE=M034G01;LINKAGE=SC

NODE=M034220

NODE=M034R20
NODE=M034R20NODE=M034R21
NODE=M034R21NODE=M034R16
NODE=M034R16
NODE=M034R16NODE=M034R15
NODE=M034R15NODE=M034R2
NODE=M034R2NODE=M034R3
NODE=M034R3
NODE=M034R3NODE=M034R11
NODE=M034R11
NODE=M034R11NODE=M034R13
NODE=M034R13NODE=M034R17
NODE=M034R17

$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$ (All η decays.)	$\Gamma_{13}/\Gamma_8 = \Gamma_{13}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$	NODE=M034R5 NODE=M034R5 NODE=M034R5		
VALUE	<u>DOCUMENT ID</u>	<u>TECN</u> <u>CHG</u> <u>COMMENT</u>		
<0.09	BALTAY	68 HBC + 7-8.5 $\pi^+ p$		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.10	CRENNELL	70 HBC - 6 $\pi^- p \rightarrow f_2 \pi^- N$		
$\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$	$\Gamma_{14}/\Gamma_8 = \Gamma_{14}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$	NODE=M034R6 NODE=M034R6		
VALUE	<u>DOCUMENT ID</u>	<u>TECN</u> <u>CHG</u> <u>COMMENT</u>		
<0.10	CRENNELL	70 HBC - 6 $\pi^- p \rightarrow f_2 \pi^- N$		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.1	BALTAY	68 HBC + 7,8.5 $\pi^+ p$		
$\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$		Γ_{15}/Γ		
VALUE	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0036	97.7	AMELIN	99 VES	37 $\pi^- A \xrightarrow{\omega\pi^-} \pi^0 A^*$
$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$		Γ_{16}/Γ		
VALUE	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.0019	97.7	AMELIN	99 VES	37 $\pi^- A \xrightarrow{\omega\pi^-} \pi^0 A^*$
$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$		Γ_{18}/Γ		
VALUE	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
possibly seen	69k	KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$		Γ_{19}/Γ		
VALUE	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
not seen	69k	KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
D-wave/S-wave RATIO FOR $\pi_2(1670) \rightarrow f_2(1270)\pi$		NODE=M034R14 NODE=M034R14		
VALUE		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.18±0.06	24 BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.22±0.10	26 DAUM	81B SPEC	63.94 $\pi^- p$	
F-wave/P-wave RATIO FOR $\pi_2(1670) \rightarrow \rho\pi$		NODE=M034R22 NODE=M034R22		
VALUE		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-0.72±0.07±0.14	CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$	
24 Using preliminary CBAR data.				NODE=M034R;LII
25 With the $\sigma\pi$ in $L=2$ and the $f_2(1270)\pi$ in $L=0$.				NODE=M034R15;LII
26 From a two-resonance fit to four 2^-0^+ waves.				NODE=M034R;LII
27 From a partial-wave analysis of $K^+K^-\pi^-$ system.				NODE=M034R13;LII
28 Normalized to the $B(\pi_2(1670) \rightarrow f_2\pi)$.				NODE=M034R;LII
$\pi_2(1670)$ REFERENCES				
AGHASYAN 18B PR D98 092003 M. Aghasyan et al. (COMPASS Collab.)				
ADOLPH 14 EPJ A50 79 C. Adolph et al. (COMPASS Collab.)				
ALEKSEEV 10 PRL 104 241803 M.G. Alekseev et al. (COMPASS Collab.)				
SCHEGELSKY 06 EPJ A27 199 V.A. Schegelsky et al. (BNL E852 Collab.)				
LU 05 PRL 94 032002 M. Lu et al. (BNL E852 Collab.)				
KUHN 04 PL B595 109 J. Kuhn et al. (BNL E852 Collab.)				
CHUNG 02 PR D65 072001 S.U. Chung et al. (BNL E852 Collab.)				
BARBERIS 01 PL B507 14 D. Barberis et al. (BNL E852 Collab.)				
AMELIN 99 PAN 62 445 D.V. Amelin et al. (VES Collab.)				
BAKER 99 PL B449 114 C.A. Baker et al. (WA 102 Collab.)				
BARBERIS 98B PL B422 399 D. Barberis et al. (L3 Collab.)				
ACCIARRI 97T PL B413 147 M. Acciari et al. (ARGUS Collab.)				
ALBRECHT 97B ZPHY C74 469 H. Albrecht et al. (SERP, TBIL)				
AMELIN 95B PL B356 595 D.V. Amelin et al. (SERP, TBIL)				
BERDNIKOV 94 PL B337 219 E.B. Berdnikov et al. (Crystal Ball Collab.)				
ANTREASYAN 90 ZPHY C48 561 D. Antreasyan et al. (CELLO Collab.)				
BEHREND 90C ZPHY C46 583 H.J. Behrend et al. (CELLO Collab.)				
ANTIPOV 87 EPL 4 403 Y.M. Antipov et al. (SERP, JINR, INRM+)				
BELLINI 85 SJNP 41 781 D. Bellini et al. (SERP, JINR, INRM+)				
ARMSTRONG 82B NP B202 1 T.A. Armstrong, B. Baccari (AACH3, BARI, BONN+)				
DAUM 81B NP B182 269 C. Daum et al. (AMST, CERN, CRAC, MPIM+)				
EVANGELIS... 81 NP B178 197 C. Evangelista et al. (BARI, BONN, CERN+)				
Also NP B186 594 C. Evangelista				
DAUM 80D PL 89B 285 C. Daum et al. (AMST, CERN, CRAC, MPIM+)				
BALTY 77 PRL 39 591 C. Baltay, C.V. Cautis, M. Kalelkar (COLU) JP				
ASCOLI 73 PR D7 669 G. Ascoli (ILL, TNTO, GENO, HAMB, MILA+) JP				
CRENNELL 70 PRL 24 781 D.J. Crennell et al. (BNL)				
ARMENISE 69 LNC 2 501 N. Armenise et al. (BARI, BGNA, FIRZ)				
BALTY 68 PRL 20 887 C. Baltay et al. (COLU, ROCH, RUTG, YALE) I				
BARTSCH 68 NP B7 345 J. Bartsch et al. (AACH, BERL, CERN) JP				

NODE=M067

 $\phi(1680)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\phi(1680)$ MASS** **e^+e^- PRODUCTION**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1680 ± 20 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1641 ⁺²⁴ ₋₁₈		ACHASOV	19	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
1667 \pm 5 \pm 11	3k	1 IVANOV	19A	CMD3 $1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$
1700 \pm 23	2k	2 ACHASOV	18A	SND $1.3-2.0 e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$
1674 \pm 12 \pm 6	6.2k	3 LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1733 \pm 10 \pm 10		4 LEES	12F	BABR $10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
1689 \pm 7 \pm 10	4.8k	5 SHEN	09	BELL $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
1709 \pm 20 \pm 43		6 AUBERT	08S	BABR $10.6 e^+e^- \rightarrow \text{hadrons}$
1623 \pm 20	948	7 AKHMETSHIN 03	CMD2	$1.05-1.38 e^+e^- \rightarrow K_L^0 K_S^0$
~ 1500		8 ACHASOV	98H	RVUE $e^+e^- \rightarrow \pi^+\pi^-\pi^0, \omega\pi^+\pi^-$, K^+K^-
~ 1900		9 ACHASOV	98H	RVUE $e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$
1700 \pm 20		10 CLEGG	94	RVUE $e^+e^- \rightarrow K^+K^-, K_S^0 K\pi$
1657 \pm 27	367	11 BISELLO	91C	DM2 $e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$
1655 \pm 17		12 BISELLO	88B	DM2 $e^+e^- \rightarrow K^+K^-$
1680 \pm 10		12 BUON	82	DM1 $e^+e^- \rightarrow \text{hadrons}$
1677 \pm 12		13 MANE	82	DM1 $e^+e^- \rightarrow K_S^0 K\pi$

- 1 From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.
 2 Assuming the $K\bar{K}^*(892) + \text{c.c.}$ dynamics. Systematic uncertainties not estimated.
 3 Using a vector meson dominance model with contribution from $\phi(1020)$, $\phi(1680)$, and higher mass excitations of $\rho(770)$ and $\omega(782)$.
 4 Using events with $\pi\pi$ invariant mass less than 0.85 GeV.
 5 From a fit with two incoherent Breit-Wigners.
 6 From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.
 7 From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.
 8 Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.
 9 Using the data from BISELLO 91C.
 10 Using BISELLO 88B and MANE 82 data.
 11 From global fit including ρ , ω , ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.
 12 From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm\pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.
 13 Fit to one channel only, neglecting interference with ω , $\rho(1700)$.

PHOTOPRODUCTION

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1753 \pm 3		1 LINK	02K	FOCS $20-160 \gamma p \rightarrow K^+K^-p$
1726 \pm 22		1 BUSENITZ	89	TPS $\gamma p \rightarrow K^+K^-X$
1760 \pm 20		1 ATKINSON	85C	OMEG $20-70 \gamma p \rightarrow K\bar{K}X$
1690 \pm 10		1 ASTON	81F	OMEG $25-70 \gamma p \rightarrow K^+K^-X$

¹ We list here a state decaying into K^+K^- possibly different from $\phi(1680)$.

 $p\bar{p}$ ANNIHILATION

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1700 \pm 8		1 AMSLER	06	CBAR $0.9 \bar{p}p \rightarrow K^+K^-\pi^0$

¹ Could also be $\rho(1700)$.

NODE=M067205

NODE=M067M1

NODE=M067M1

→ UNCHECKED ←

OCCUR=4

NODE=M067M1;LINKAGE=D
NODE=M067M1;LINKAGE=C
NODE=M067M1;LINKAGE=BNODE=M067M1;LINKAGE=A
NODE=M067M1;LINKAGE=SH
NODE=M067M1;LINKAGE=AU

NODE=M067M;LINKAGE=HK

NODE=M067M1;LINKAGE=L1

NODE=M067M1;LINKAGE=L4
NODE=M067M;LINKAGE=A
NODE=M067M;LINKAGE=E

NODE=M067M;LINKAGE=C

NODE=M067M;LINKAGE=D

NODE=M067M2

NODE=M067M2

NODE=M067M2;LINKAGE=LK

NODE=M067M3

NODE=M067M3

NODE=M067M3;LINKAGE=AM

$\phi(1680)$ WIDTH

e^+e^- PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
150±50 OUR ESTIMATE				This is only an educated guess; the error given is larger than the error on the average of the published values.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
103 ⁺²⁶ ₋₂₄		ACHASOV 19	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
176±23± 38	3k	¹ IVANOV 19A	CMD3	1.59-2.007 $e^+e^- \rightarrow K^+K^-\eta$
300±50	2k	2 ACHASOV 18A	SND	1.3-2.0 $e^+e^- \rightarrow K_S^0K_L^0\pi^0$
165±38± 70	6.2k	3 LEES 14H	BABR	$e^+e^- \rightarrow K_S^0K_L^0\gamma$
300±15± 37		4 LEES 12F	BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
211±14± 19	4.8k	5 SHEN 09	BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
322±77±160		6 AUBERT 08S	BABR	10.6 $e^+e^- \rightarrow$ hadrons
139±60	948	7 AKHMETSHIN 03	CMD2	1.05-1.38 $e^+e^- \rightarrow K_L^0K_S^0$
300±60		8 CLEGG 94	RVUE	$e^+e^- \rightarrow K^+K^-, K_S^0K\pi$
146±55	367	9 BISELLO 91C	DM2	$e^+e^- \rightarrow K_S^0K^\pm\pi^\mp$
207±45		10 BUON 88B	DM2	$e^+e^- \rightarrow K^+K^-$
185±22		11 MANE 82	DM1	$e^+e^- \rightarrow$ hadrons
102±36		11 MANE 82	DM1	$e^+e^- \rightarrow K_S^0K\pi$

- 1 From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.
 2 Assuming the $K\bar{K}^*(892) + c.c.$ dynamics. Systematic uncertainties not estimated.
 3 Using a vector meson dominance model with contribution from $\phi(1020)$, $\phi(1680)$, and higher mass excitations of $\rho(770)$ and $\omega(782)$.
 4 Using events with $\pi\pi$ invariant mass less than 0.85 GeV.
 5 From a fit with two incoherent Breit-Wigners.
 6 From the simultaneous fit to the $K\bar{K}^*(892) + c.c.$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.
 7 From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.
 8 Using BISELLO 88B and MANE 82 data.
 9 From global fit including ρ , ω , ϕ and $\rho(1700)$.
 10 From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_L^0K_S^0$, $K_S^0K^\pm\pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.
 11 Fit to one channel only, neglecting interference with ω , $\rho(1700)$.

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
122±63	¹ LINK 02K	FOCS	$20-160 \gamma p \rightarrow K^+K^-p$
121±47	¹ BUSENITZ 89	TPS	$\gamma p \rightarrow K^+K^-X$
80±40	¹ ATKINSON 85C	OMEG	$20-70 \gamma p \rightarrow K\bar{K}X$
100±40	¹ ASTON 81F	OMEG	$25-70 \gamma p \rightarrow K^+K^-X$

1 We list here a state decaying into K^+K^- possibly different from $\phi(1680)$.

$p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
143±24	¹ AMSLER 06	CBAR	$0.9 \bar{p}p \rightarrow K^+K^-\pi^0$

1 Could also be $\rho(1700)$.

$\phi(1680)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}^*(892) + c.c.$	seen
$\Gamma_2 K_S^0K\pi$	seen
$\Gamma_3 K\bar{K}$	seen
$\Gamma_4 K_L^0K_S^0$	seen
$\Gamma_5 e^+e^-$	seen
$\Gamma_6 \omega\pi\pi$	not seen
$\Gamma_7 \phi\pi\pi$	
$\Gamma_8 K^+K^-\pi^+\pi^-$	seen
$\Gamma_9 \eta\phi$	seen
$\Gamma_{10} K^+K^-\eta$	
$\Gamma_{11} \eta\gamma$	
$\Gamma_{12} K^+K^-\pi^0$	seen

NODE=M067210

NODE=M067W1

NODE=M067W1

→ UNCHECKED ←

NODE=M067W1;LINKAGE=D

NODE=M067W1;LINKAGE=C

NODE=M067W1;LINKAGE=B

NODE=M067W1;LINKAGE=A

NODE=M067W1;LINKAGE=SH

NODE=M067W1;LINKAGE=AU

NODE=M067W;LINKAGE=HK

NODE=M067W;LINKAGE=A

NODE=M067W;LINKAGE=E

NODE=M067W;LINKAGE=C

NODE=M067W;LINKAGE=D

NODE=M067W2

NODE=M067W2

NODE=M067W2;LINKAGE=LK

NODE=M067W3

NODE=M067W3

NODE=M067W3;LINKAGE=AM

NODE=M067215;NODE=M067

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=11

DESIG=12;OUR EVAL;→ UNCHECKED ←

DESIG=10

DESIG=14

DESIG=13

DESIG=2

$\phi(1680) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the integrated cross section into channel (i) in $e^+ e^-$ annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.3 \pm 2.4 \pm 6.2 6.2k ¹ LEES 14H BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$

¹ Using a vector meson dominance model with contribution from $\phi(1020)$, $\phi(1680)$, and higher mass excitations of $\rho(770)$ and $\omega(782)$.

$\Gamma_4 \Gamma_5 / \Gamma$

NODE=M067A00
NODE=M067A00

$\Gamma(\phi \pi \pi) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 \pm 0.2 \pm 0.3 LEES 12F BABR 10.6 $e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$

$\Gamma_7 \Gamma_5 / \Gamma$

NODE=M067G02
NODE=M067G02

$\Gamma(\eta \phi) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

94 \pm 13 \pm 15 3k ¹ IVANOV 19A CMD3 $1.59-2.007 e^+ e^- \rightarrow K^+ K^- \eta$

¹ From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.

$\Gamma_9 \Gamma_5 / \Gamma$

NODE=M067R00
NODE=M067R00

$\phi(1680) \Gamma(i) \Gamma(e^+ e^-)/\Gamma^2(\text{total})$

This combination of a branching ratio into channel (i) and branching ratio into $e^+ e^-$ is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (i) or $e^+ e^-$.

$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma \times \Gamma_5/\Gamma$

NODE=M067G5
NODE=M067G5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.131 \pm 0.059 948 ¹ AKHMETSHIN 03 CMD2 $1.05-1.38 e^+ e^- \rightarrow K_L^0 K_S^0$

¹ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known. Recalculated by us.

$\Gamma(K \bar{K}^*(892)+\text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma \times \Gamma_5/\Gamma$

NODE=M067G;LINKAGE=GK

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.15 \pm 0.16 \pm 0.01 1 AUBERT 08S BABR $10.6 e^+ e^- \rightarrow K \bar{K}^*(892) \gamma +$

3.29 \pm 1.57 367 ² BISELLO 91C DM2 $1.35-2.40 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From the simultaneous fit to the $K \bar{K}^*(892)+\text{c.c.}$ and $\phi \eta$ data from AUBERT 08S using the results of AUBERT 07AK.

² Recalculated by us with the published value of $B(K \bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+ e^-)$.

NODE=M067G6
NODE=M067G6

$\Gamma(\phi \pi \pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

$\Gamma_7/\Gamma \times \Gamma_5/\Gamma$

NODE=M067G6;LINKAGE=AU

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.86 \pm 0.14 \pm 0.21 4.8k ¹ SHEN 09 BELL $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ Multiplied by 3/2 to take into account the $\phi \pi^0 \pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

NODE=M067G01
NODE=M067G01

NODE=M067G01;LINKAGE=SH

$\Gamma(\eta\phi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma \times \Gamma_5/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.64 ^{+1.74} _{-1.80}		ACHASOV	19	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$	
5.3 ± 0.6 ± 0.9	3k	¹ IVANOV	19A	CMD3 $1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$	
4.3 ± 1.0 ± 0.9		² AUBERT	08S	BABR $10.6 e^+e^- \rightarrow \phi\eta\gamma$	

¹ From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.

² From the simultaneous fit to the $K\bar{K}^*(892) + c.c.$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

 $\phi(1680)$ BRANCHING RATIOS
 $\Gamma(K\bar{K}^*(892) + c.c.)/\Gamma(K_S^0 K\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
dominant	MANE	82	DM1 $e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$	

 $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892) + c.c.)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.07 ± 0.01 BUON 82 DM1 e^+e^-

 $\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892) + c.c.)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_1
<0.10	BUON	82	DM1 e^+e^-	

 $\Gamma(\eta\phi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
seen	35	¹ ACHASOV	14	SND $1.15-2.00 e^+e^- \rightarrow \eta\gamma$	

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

 $\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892) + c.c.)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				

≈ 0.37 ¹ AUBERT 08S BABR $10.6 e^+e^- \rightarrow$ hadrons

¹ From the fit including data from AUBERT 07AK.

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
seen	35	¹ ACHASOV	14	SND $1.15-2.00 e^+e^- \rightarrow \eta\gamma$	

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

 $\phi(1680)$ REFERENCES

ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	NODE=M067G7
IVANOV	19A	PL B798 134946	V.L. Ivanov <i>et al.</i>	(CMD-3 Collab.)	NODE=M067G7
ACHASOV	18A	PR D97 032011	M.N. Achasov <i>et al.</i>	(SND Collab.)	NODE=M067G7
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	NODE=M067G7
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	NODE=M067R2
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	NODE=M067R2
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	NODE=M067R2
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	NODE=M067R2
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	NODE=M067R2
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	NODE=M067R2
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)	NODE=M067R2
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	NODE=M067R2
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin		NODE=M067R2
		Translated from YAF 65 1255.			
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	NODE=M067R2
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		NODE=M067R2
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	NODE=M067R2
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	NODE=M067R2
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)	NODE=M067R2
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	NODE=M067R2
BUSENITZ	89	PR D40 1	J.K. Busenitz <i>et al.</i>	(ILL, FNAL)	NODE=M067R2
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLÉR, FRAS+)	NODE=M067R2
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	NODE=M067R2
		Translated from ZETFP 46 132.			
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=59887
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)	REFID=60133
MANE	82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)	REFID=58883
ASTON	81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=55912
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=55940
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ØRSAY)	REFID=54298
		Translated from ZETFP 46 132.			REFID=54066
					REFID=53000
					REFID=52242
					REFID=51908
					REFID=51136
					REFID=49172
					REFID=48827
					REFID=48845
					REFID=46323
					REFID=44081
					REFID=43168
					REFID=41867
					REFID=41369
					REFID=40927
					REFID=40581
					REFID=40280
					REFID=21596
					REFID=21494
					REFID=21590
					REFID=21585
					REFID=20553
					REFID=21588

$\rho_3(1690)$ $I^G(J^{PC}) = 1^+(3^{--})$

NODE=M015

 $\rho_3(1690)$ MASSVALUE (MeV)DOCUMENT ID**1688.8±2.1 OUR AVERAGE**

Includes data from the 5 datablocks that follow this one.

2 π MODEVALUE (MeV)EVTSDOCUMENT IDTECNCHGCOMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015205

NODE=M015M

NODE=M015M1

NODE=M015M1

1686± 4 OUR AVERAGE

1677±14		EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow 2\pi p$
1679±11	476	BALTAJ	78B	HBC	0	15 $\pi^+ p \rightarrow \pi^+ \pi^- n$
1678±12	175	¹ ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
1690± 7	600	¹ ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
1693± 8		² GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1678±12		MATTHEWS	71c	DBC	0	7 $\pi^+ N$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1734±10		³ CORDEN	79	OMEG		12–15 $\pi^- p \rightarrow n2\pi$
1692±12		^{2,4} ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1737±23		ARMENISE	70	DBC	0	9 $\pi^+ N$
1650±35	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
1687±21		STUNTEBECK	70	HDBC	0	8 $\pi^- p, 5.4 \pi^+ d$
1683±13		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
1670±30		GOLDBERG	65	HBC	0	6 $\pi^+ d, 8 \pi^- p$

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.² Uses same data as HYAMS 75.³ From a phase shift solution containing a $f'_2(1525)$ width two times larger than the $K\bar{K}$ result.⁴ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.

NODE=M015M1;LINKAGE=E

NODE=M015M1;LINKAGE=G

NODE=M015M1;LINKAGE=M

NODE=M015M1;LINKAGE=I

NODE=M015M2

NODE=M015M2

 $K\bar{K}$ AND $K\bar{K}\pi$ MODESVALUE (MeV)EVTSDOCUMENT IDTECNCHGCOMMENT

The data in this block is included in the average printed for a previous datablock.

1696± 4 OUR AVERAGE

1699± 5		ALPER	80	CNTR	0	62 $\pi^- p \rightarrow K^+ K^- n$
1698±12	6k	^{5,6} MARTIN	78D	SPEC		10 $\pi p \rightarrow K_S^0 K^- p$
1692± 6		BLUM	75	ASPK	0	18.4 $\pi^- p \rightarrow nK^+ K^-$
1690±16		ADERHOLZ	69	HBC	+	8 $\pi^+ p \rightarrow K\bar{K}\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1694± 8		⁷ COSTA	80	OMEG		10 $\pi^- p \rightarrow K^+ K^- n$

⁵ From a fit to $J^P = 3^-$ partial wave.⁶ Systematic error on mass scale subtracted.⁷ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

NODE=M015M2;LINKAGE=P

NODE=M015M2;LINKAGE=S

NODE=M015M2;LINKAGE=L

NODE=M015M3

NODE=M015M3

(4 π) \pm MODEVALUE (MeV)EVTSDOCUMENT IDTECNCHGCOMMENT

The data in this block is included in the average printed for a previous datablock.

1686± 5 OUR AVERAGE Error includes scale factor of 1.1.

1694± 6		⁸ EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p4\pi$
1665±15	177	BALTAJ	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
1670±10		THOMPSON	74	HBC	+	13 $\pi^+ p$
1687±20		CASON	73	HBC	—	8,18.5 $\pi^- p$
1685±14		⁹ CASON	73	HBC	—	8,18.5 $\pi^- p$
1680±40	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N4\pi$
1689±20	102	⁹ BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\rho$
1705±21		CASO	70	HBC	—	11.2 $\pi^- p \rightarrow n\rho2\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1718±10		¹⁰ EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p4\pi$
1673± 9		¹¹ EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p4\pi$
1733± 9	66	⁹ KLIGER	74	HBC	—	4.5 $\pi^- p \rightarrow p4\pi$
1630±15		HOLMES	72	HBC	+	10–12 $K^+ p$
1720±15		BALTAJ	68	HBC	+	7, 8.5 $\pi^+ p$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

- 8 From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.
 9 From $\rho^\pm \rho^0$ mode.
 10 From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.
 11 From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

 $\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

1681± 7 OUR AVERAGE

1670±25	12 ALDE	95 GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
1690±15	EVANGELIS...	81 OMEG	12 $\pi^- p \rightarrow \omega\pi p$
1666±14	GESELLAROLI	77 HBC	11 $\pi^- p \rightarrow \omega\pi p$
1686± 9	THOMPSON	74 HBC	13 $\pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1654±24	BARNHAM	70 HBC	10 $K^+ p \rightarrow \omega\pi X$

12 Supersedes ALDE 92c.

 $\eta\pi^+\pi^-$ MODE(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.				

1682±12 OUR AVERAGE


1685±10±20	AMELIN	00 VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
1680±15	FUKUI	88 SPEC	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1700±47	13 ANDERSON	69 MMS	16 $\pi^- p$ backward
1632±15	13,14 FOCACCI	66 MMS	7-12 $\pi^- p \rightarrow p\text{MM}$
1700±15	13,14 FOCACCI	66 MMS	7-12 $\pi^- p \rightarrow p\text{MM}$
1748±15	13,14 FOCACCI	66 MMS	7-12 $\pi^- p \rightarrow p\text{MM}$

13 Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+ 2\pi^-$, with 0, 1, 2 $\pi^+ \pi^-$ pairs in ρ band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1976)

14 Not seen by BOWEN 72.

 $\rho_3(1690)$ WIDTH **2π , $K\bar{K}$, AND $K\bar{K}\pi$ MODES**

VALUE (MeV)	DOCUMENT ID
161±10 OUR AVERAGE Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.	

WEIGHTED AVERAGE
161±10 (Error scaled by 1.5)

NODE=M015M3;LINKAGE=A

NODE=M015M3;LINKAGE=F

NODE=M015M3;LINKAGE=B

NODE=M015M3;LINKAGE=C

NODE=M015M5

NODE=M015M5

NODE=M015M5;LINKAGE=A

NODE=M015M6

NODE=M015M6

NODE=M015M6

OCCUR=2

OCCUR=3

NODE=M015M6;LINKAGE=R

NODE=M015M6;LINKAGE=N

NODE=M015210

NODE=M015W

NODE=M015W

2π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

186±14 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

220±29	DENNEY	83	LASS	10	$\pi^+ N$
246±37	EVANGELIS...	81	OMEG	—	$12 \pi^- p \rightarrow 2\pi p$
116±30	BALTAY	78B	HBC	0	$15 \pi^+ p \rightarrow \pi^+ \pi^- n$
162±50	ANTIPOV	77	CIBS	0	$25 \pi^- p \rightarrow p 3\pi$
167±40	ENGLER	74	DBC	0	$6 \pi^+ n \rightarrow \pi^+ \pi^- p$
200±18	GRAYER	74	ASPK	0	$17 \pi^- p \rightarrow \pi^+ \pi^- n$
156±36	MATTHEWS	71C	DBC	0	$7 \pi^+ N$
171±65	ARMENISE	70	DBC	0	$9 \pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
322±35	CORDEN	79	OMEG	—	$12-15 \pi^- p \rightarrow n 2\pi$
240±30	ESTABROOKS	75	RVUE	—	$17 \pi^- p \rightarrow \pi^+ \pi^- n$
180±30	BARTSCH	70B	HBC	+	$8 \pi^+ p \rightarrow N 2\pi$
267 ⁺⁷² -46	STUNTEBECK	70	HDBC	0	$8 \pi^- p, 5.4 \pi^+ d$
188±49	ARMENISE	68	DBC	0	$5.1 \pi^+ d$
180±40	GOLDBERG	65	HBC	0	$6 \pi^+ d, 8 \pi^- p$

15 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

16 Uses same data as HYAMS 75 and BECKER 79.

17 From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K\bar{K}$ result.

18 From phase-shift analysis. Error takes account of spread of different phase-shift solutions.

NODE=M015W1

NODE=M015W1


NODE=M015W1;LINKAGE=T

NODE=M015W1;LINKAGE=G

NODE=M015W1;LINKAGE=M

NODE=M015W1;LINKAGE=I

WEIGHTED AVERAGE
186±14 (Error scaled by 1.3)



$\rho_3(1690)$ width, 2π mode (MeV)

 $K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

204±18 OUR AVERAGE

199±40	6000	19 MARTIN	78D	SPEC	$10 \pi p \rightarrow K_S^0 K^- p$
205±20		BLUM	75	ASPK	$18.4 \pi^- p \rightarrow n K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
219±4		ALPER	80	CNTR	$62 \pi^- p \rightarrow K^+ K^- n$
186±11		20 COSTA	80	OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
112±60		ADERHOLZ	69	HBC	$8 \pi^+ p \rightarrow K\bar{K}\pi$

19 From a fit to $J^P = 3^-$ partial wave.

20 They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

NODE=M015W2

NODE=M015W2

NODE=M015W2;LINKAGE=P

NODE=M015W2;LINKAGE=L

(4 π) \pm MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

129 \pm 10 OUR AVERAGE

123 \pm 13	21	EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p4\pi$
105 \pm 30	177	BALTAJ	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
169 $^{+70}_{-48}$		CASON	73	HBC	—	8,18.5 $\pi^- p$
135 \pm 30	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N4\pi$
160 \pm 30	102	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
230 \pm 28	22	EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p4\pi$
184 \pm 33	23	EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow p4\pi$
150	66	KLIGER	74	HBC	—	4.5 $\pi^- p \rightarrow p4\pi$
106 \pm 25		THOMPSON	74	HBC	+	13 $\pi^+ p$
125 $^{+83}_{-35}$	24	CASON	73	HBC	—	8,18.5 $\pi^- p$
130 \pm 30		HOLMES	72	HBC	+	10-12 $K^+ p$
180 \pm 30	90	24 BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N_{a_2}\pi$
100 \pm 35		BALTAJ	68	HBC	+	7, 8.5 $\pi^+ p$

21 From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.

22 From $a_2(1320)^-\pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.

23 From $a_2(1320)^0\pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

24 From $\rho^\pm \rho^0$ mode.

NODE=M015W3

NODE=M015W3

 $\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

190 \pm 40 OUR AVERAGE

230 \pm 65	25	ALDE	95	GAM2	—	38 $\pi^- p \rightarrow \omega\pi^0 n$
190 \pm 65		EVANGELIS...	81	OMEG	—	12 $\pi^- p \rightarrow \omega\pi p$
160 \pm 56		GESSAROLI	77	HBC	—	11 $\pi^- p \rightarrow \omega\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
89 \pm 25		THOMPSON	74	HBC	+	13 $\pi^+ p$
130 $^{+73}_{-43}$		BARNHAM	70	HBC	+	10 $K^+ p \rightarrow \omega\pi X$

25 Supersedes ALDE 92C.

NODE=M015W3;LINKAGE=A

NODE=M015W3;LINKAGE=B

NODE=M015W3;LINKAGE=C

NODE=M015W3;LINKAGE=F

NODE=M015W5

NODE=M015W5

 $\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015W5;LINKAGE=A

NODE=M015W6

NODE=M015W6

NODE=M015W6

126 \pm 40 OUR AVERAGE Error includes scale factor of 1.8.

220 \pm 30 \pm 50	AMELIN	00	VES	—	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
106 \pm 27	FUKUI	88	SPEC	0	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

195	26 ANDERSON	69	MMS	—	16 $\pi^- p$ backward
< 21	26,27 FOCACCI	66	MMS	—	7-12 $\pi^- p \rightarrow p\text{MM}$
< 30	26,27 FOCACCI	66	MMS	—	7-12 $\pi^- p \rightarrow p\text{MM}$
< 38	26,27 FOCACCI	66	MMS	—	7-12 $\pi^- p \rightarrow p\text{MM}$

OCCUR=2

OCCUR=3

NODE=M015W6;LINKAGE=R

NODE=M015W6;LINKAGE=N

26 Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+ 2\pi^-$, with 0, 1, 2 $\pi^+ \pi^-$ pairs in ρ^0 band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1979)

27 Not seen by BOWEN 72.

$\rho_3(1690)$ DECAY MODES

NODE=M015215;NODE=M015

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 4π	(71.1 \pm 1.9) %	
Γ_2 $\pi^\pm \pi^+ \pi^- \pi^0$	(67 \pm 22) %	
Γ_3 $\omega \pi$	(16 \pm 6) %	
Γ_4 $\pi \pi$	(23.6 \pm 1.3) %	
Γ_5 $K \bar{K} \pi$	(3.8 \pm 1.2) %	
Γ_6 $K \bar{K}$	(1.58 \pm 0.26) %	1.2
Γ_7 $\eta \pi^+ \pi^-$	seen	
Γ_8 $\rho(770) \eta$	seen	
Γ_9 $\pi \pi \rho$	seen	
Γ_{10} $a_2(1320) \pi$	seen	
Γ_{11} $\rho \rho$	seen	
Γ_{12} $\phi \pi$		
Γ_{13} $\eta \pi$		
Γ_{14} $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 14.7$ for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-77			
x_5	-74	17		
x_6	-15	2	0	
	x_1	x_4	x_5	

 $\rho_3(1690)$ BRANCHING RATIOS

NODE=M015220

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ
0.236 \pm 0.013 OUR FIT					
0.243 \pm 0.013 OUR AVERAGE					
0.259 \pm 0.018	BECKER 79	ASPK 0		17 $\pi^- p$ polarized	
0.23 \pm 0.02	CORDEN 79	OMEG		12–15 $\pi^- p \rightarrow n 2\pi$	
0.22 \pm 0.04	28 MATTHEWS 71C	HDBC 0		7 $\pi^+ n \rightarrow \pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.245 \pm 0.006	29 ESTABROOKS 75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$	

NODE=M015220

NODE=M015R1

NODE=M015R1

$\Gamma(\pi\pi)/\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ_2
0.35 \pm 0.11	CASON 73	HBC	—	8,18.5 $\pi^- p$	

NODE=M015R1;LINKAGE=P

NODE=M015R1;LINKAGE=G

NODE=M015R2

NODE=M015R2

$\Gamma(\pi\pi)/\Gamma(4\pi)$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ_1
0.332 \pm 0.026 OUR FIT	Error includes scale factor of 1.1.				
0.30 \pm 0.10	BALTAY 78B	HBC 0		15 $\pi^+ p \rightarrow p 4\pi$	


NODE=M015R3

NODE=M015R3

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ_4
0.067±0.011 OUR FIT				Error includes scale factor of 1.2.	NODE=M015R4 NODE=M015R4
0.118^{+0.040}_{-0.032} OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.	
0.191 ^{+0.040} _{-0.037}	GORLICH	80	ASPK	0	17,18 $\pi^- p$ polarized
0.08 ± 0.03	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.08 ^{+0.08} _{-0.03}	CRENNELL	68B	HBC		6.0 $\pi^- p$

WEIGHTED AVERAGE
0.118+0.040-0.032 (Error scaled by 1.7)

 $\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_4
0.16±0.05 OUR FIT					NODE=M015R5 NODE=M015R5
0.16±0.05	30 BARTSCH	70B	HBC	+	8 $\pi^+ p$
30 Increased by us to correspond to $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.					NODE=M015R5;LINKAGE=A
[$\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$					NODE=M015R6 NODE=M015R6
0.94±0.09 OUR AVERAGE					
0.96±0.21	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.88±0.15	BALLAM	71B	HBC	-	16 $\pi^- p$
1 ± 0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
consistent with 1	CASO	68	HBC	-	11 $\pi^- p$

 $\Gamma(\rho\rho)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{11}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.12±0.11		BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.56	66	KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$
0.13±0.09	31	THOMPSON	74	HBC	+	13 $\pi^+ p$
0.7 ± 0.15		BARTSCH	70B	HBC	+	8 $\pi^+ p$

³¹ $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

NODE=M015R7
NODE=M015R7

 $\Gamma(\rho\rho)/[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_{11}/(\Gamma_9+\Gamma_{10}+\Gamma_{11})$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.48±0.16	CASO	68	HBC	-	11 $\pi^- p$

NODE=M015R8
NODE=M015R8

$\Gamma(a_2(1320)\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE		DOCUMENT ID	TECN	CHG	COMMENT	Γ_{10}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.66 ± 0.08	BALTAY	78B	HBC	+	$15 \pi^+ p \rightarrow p4\pi$	
0.36 ± 0.14	32 THOMPSON	74	HBC	+	$13 \pi^+ p$	
not seen	CASON	73	HBC	-	$8,18.5 \pi^- p$	
0.6 ± 0.15	BARTSCH	70B	HBC	+	$8 \pi^+ p$	
0.6	BALTAY	68	HBC	+	$7.8.5 \pi^+ p$	

32 $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

NODE=M015R9

NODE=M015R9

 $\Gamma(\omega\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_2
0.23 ± 0.05 OUR AVERAGE						
					Error includes scale factor of 1.2.	
0.33 ± 0.07		THOMPSON	74	HBC	+	$13 \pi^+ p$
0.12 ± 0.07		BALLAM	71B	HBC	-	$16 \pi^- p$
0.25 ± 0.10		BALTAY	68	HBC	+	$7.8.5 \pi^+ p$
0.25 ± 0.10		JOHNSTON	68	HBC	-	$7.0 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	95	BALTAY	78B	HBC	+	$15 \pi^+ p \rightarrow p4\pi$
<0.09		KLIGER	74	HBC	-	$4.5 \pi^- p \rightarrow p4\pi$

NODE=M015R9;LINKAGE=T

NODE=M015R10

NODE=M015R10

 $\Gamma(\phi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE		DOCUMENT ID	TECN	CHG	COMMENT	Γ_{12}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						

<0.11 BALTAY 68 HBC + $7.8.5 \pi^+ p$

NODE=M015R11

NODE=M015R11

 $\Gamma(\pi^\pm 2\pi^+ 2\pi^- \pi^0)/\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)$

VALUE		DOCUMENT ID	TECN	CHG	COMMENT	Γ_{14}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						

<0.15 BALTAY 68 HBC + $7.8.5 \pi^+ p$

NODE=M015R12

NODE=M015R12

 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

VALUE		DOCUMENT ID	TECN	CHG	COMMENT	Γ_{13}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •						

<0.02 THOMPSON 74 HBC + $13 \pi^+ p$

NODE=M015R13

NODE=M015R13

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$

VALUE		DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ
0.0158 ± 0.0026 OUR FIT Error includes scale factor of 1.2.						

0.0130 ± 0.0024 OUR AVERAGE

0.013 ± 0.003	COSTA	80	OMEG 0		$10 \pi^- p \rightarrow K^+ K^- n$
0.013 ± 0.004	33 MARTIN	78B	SPEC	-	$10 \pi p \rightarrow K_S^0 K^- p$

33 From $(\Gamma_4\Gamma_6)^{1/2} = 0.056 \pm 0.034$ assuming $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.

NODE=M015R14

NODE=M015R14

 $\Gamma(\omega\pi)/[\Gamma(\omega\pi) + \Gamma(\rho\rho)]$

VALUE		DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_3/(\Gamma_3 + \Gamma_{11})$
• • • We do not use the following data for averages, fits, limits, etc. • • •						

0.22 ± 0.08 CASON 73 HBC - $8,18.5 \pi^- p$

NODE=M015R14;LINKAGE=B

NODE=M015R16

NODE=M015R16

 $\Gamma(\eta\pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE		DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
seen					

NODE=M015R17

NODE=M015R17

 $\Gamma(a_2(1320)\pi)/\Gamma(\rho(770)\eta)$

VALUE		DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ_8
5.5 ± 2.0					

NODE=M015R18

NODE=M015R18

$\rho_3(1690)$ REFERENCES

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP	REFID=44371
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41859
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21665
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlitch <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
MARTIN	78B	NP B140 158	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=21273
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=21272
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP	REFID=21651
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
KLIGER	74	SJNP 19 428	G.K. Kliger <i>et al.</i>	(ITEP)	REFID=21648
Translated from YAF 19 839.					
OREN	74	NP B71 189	Y. Oren <i>et al.</i>	(ANL, OXF)	REFID=20221
THOMPSON	74	NP B69 220	G. Thompson <i>et al.</i>	(PURD)	REFID=21650
CASON	73	PR D7 1971	N.M. Cason <i>et al.</i>	(NDAM)	REFID=20606
BOWEN	72	PRL 29 890	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21711
HOLMES	72	PR D6 3336	R. Holmes <i>et al.</i>	(ROCH)	REFID=21639
BALLAM	71B	PR D3 2606	J. Ballam <i>et al.</i>	(SLAC)	REFID=21630
MATTHEWS	71C	NP B33 1	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC) JP	REFID=21633
ARMENISE	70	LNC 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
BARNHAM	70	PRL 24 1083	K.W.J. Barnham <i>et al.</i>	(BIRM)	REFID=21624
BARTSCH	70B	NP B22 109	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN)	REFID=21625
CASO	70	LNC 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20590
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ANDERSON	69	PRL 22 1390	E.W. Anderson <i>et al.</i>	(BNL, CMU)	REFID=20795
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE)	REFID=21531
CASO	68	NC 54A 983	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20586
CRENNELL	68B	PL 28B 136	D.J. Crennell <i>et al.</i>	(BNL)	REFID=21616
JOHNSTON	68	PRL 20 1414	T.F. Johnston <i>et al.</i>	(TNTO, WISC) IJP	REFID=21617
FOCACCI	66	PRL 17 890	M.N. Focacci <i>et al.</i>	(CERN)	REFID=20402
GOLDBERG	65	PL 17 354	M. Goldberg <i>et al.</i>	(CERN, EPOL, ORSAY+)	REFID=21601

 $\rho(1700)$

$I^G(J^{PC}) = 1^+(1^{--})$

See the related review(s):

[ρ\(1450\)](#) and [ρ\(1700\)](#) **$\rho(1700)$ MASS** **$\eta\rho^0$ AND $\pi^+\pi^-$ MODES**

VALUE (MeV)

DOCUMENT ID

1720±20 OUR ESTIMATE

 $\eta\rho^0$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

1840±10	7.4k	¹ ACHASOV	18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1740±20		ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1701±15		² FUKUI	88	SPEC	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

¹ From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

² Assuming $\rho_0^+(1370)$ decay mode interferes with $a_1(1260)^+\pi^-$ background. From a two Breit-Wigner fit.

NODE=M015

REFID=47432
REFID=44371
REFID=41859
REFID=40273
REFID=20754
REFID=20462
REFID=21665
REFID=20737
REFID=20738
REFID=21084
REFID=20374
REFID=21265
REFID=21273
REFID=21272
REFID=20728
REFID=20230
REFID=21651
REFID=20642
REFID=20355
REFID=20110
REFID=20113
REFID=21648
REFID=20221
REFID=21650
REFID=20606
REFID=21711
REFID=21639
REFID=21630
REFID=21633
REFID=20693
REFID=21624
REFID=21625
REFID=20590
REFID=20696
REFID=20687
REFID=20795
REFID=20054
REFID=21531
REFID=20586
REFID=21616
REFID=21617
REFID=20402
REFID=21601

NODE=M065

NODE=M065205
NODE=M065M0
NODE=M065M0
→ UNCHECKED ←

NODE=M065M6
NODE=M065M6

NODE=M065M6;LINKAGE=A

NODE=M065M;LINKAGE=B

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

• • • We do not use the following data for averages, fits, limits, etc. • • •

1770.54 ± 5.49		1 BARTOS	17 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1718.50 ± 65.44		2 BARTOS	17A RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1766.80 ± 52.36		3 BARTOS	17A RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
1644 ± 36	20K	4 LEES	17c BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
1780 ± 20	+15 -20	5 ABRAMOWICZ12	ZEUS	$e p \rightarrow e \pi^+ \pi^- p$
1861 ± 17		6 LEES	12G BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
1728 ± 17	±89	7,8 FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
1780 ± 37 -29		9 ABELE	97 CBAR	$\bar{p} n \rightarrow \pi^- \pi^0 \pi^0$
1719 ± 15		9 BERTIN	97C OBLX	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
1730 ± 30		CLEGG	94 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1768 ± 21		BISELLO	89 DM2	$e^+ e^- \rightarrow \pi^+ \pi^-$
1745.7 ± 91.9		DUBNICKA	89 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
1546 ± 26		GESHKEN...	89 RVUE	
1650		10 ERKAL	85 RVUE	20–70 $\gamma p \rightarrow \gamma \pi$
1550 ± 70		ABE	84B HYBR	20 $\gamma p \rightarrow \pi^+ \pi^- p$
1590 ± 20		11 ASTON	80 OMEG	20–70 $\gamma p \rightarrow p 2\pi$
1600 ± 10		12 ATIYA	79B SPEC	50 $\gamma C \rightarrow C 2\pi$
1598 ± 24 -22		BECKER	79 ASPK	17 $\pi^- p$ polarized
1659 ± 25		10 LANG	79 RVUE	
1575		10 MARTIN	78C RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1610 ± 30		10 FROGGATT	77 RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1590 ± 20		13 HYAMS	73 ASPK	17 $\pi^- p \rightarrow \pi^+ \pi^- n$

1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

2 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

3 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.

4 From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.

5 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.

6 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

7 $|F_\pi(0)|^2$ fixed to 1.

8 From the GOUNARIS 68 parametrization of the pion form factor.

9 T-matrix pole.

10 From phase shift analysis of HYAMS 73 data.

11 Simple relativistic Breit-Wigner fit with constant width.

12 An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

13 Included in BECKER 79 analysis.

NODE=M065M1

NODE=M065M1

OCCUR=2

NODE=M065M1;LINKAGE=C

NODE=M065M1;LINKAGE=D

NODE=M065M1;LINKAGE=E

NODE=M065M1;LINKAGE=A

NODE=M065M1;LINKAGE=AB

NODE=M065M1;LINKAGE=LE

NODE=M065M1;LINKAGE=FU

NODE=M065M1;LINKAGE=GO

NODE=M065M;LINKAGE=QQ

NODE=M065M;LINKAGE=P

NODE=M065M;LINKAGE=M

NODE=M065M;LINKAGE=R

NODE=M065M;LINKAGE=H

NODE=M065M8

NODE=M065M8

OCCUR=2

NODE=M065M8;LINKAGE=AC

NODE=M065M;LINKAGE=I1

NODE=M065M;LINKAGE=I2

 $\pi\omega$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

• • • We do not use the following data for averages, fits, limits, etc. • • •

1708 ± 41	7815	1 ACHASOV	13 SND	$1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
1550 to 1620		2 ACHASOV	00I SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
1580 to 1710		3 ACHASOV	00I SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
1710 ± 90		ACHASOV	97 RVUE	$e^+ e^- \rightarrow \omega \pi^0$

1 From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

2 Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00I on $e^+ e^- \rightarrow \omega \pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega \pi^- \nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.

3 Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00I on $e^+ e^- \rightarrow \omega \pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega \pi^- \nu_\tau$.

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1541 \pm 12	\pm 33	190k	¹ AAIJ	16N LHCb	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
1740.8 \pm 22.2	27k		² ABELE	99D CBAR	\pm $0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$
1582 \pm 36	1600		CLELAND	82B SPEC	\pm $50 \pi p \rightarrow K_S^0 K^\pm p$

¹ Using the GOUNARIS 68 parameterization with a fixed width. Value is average using different $K\pi$ S-wave parametrizations in fit.

² K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

2 ($\pi^+\pi^-$) MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1851 \pm 27		ACHASOV	97 RVUE	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1570 \pm 20		¹ CORDIER	82 DM1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1520 \pm 30		² ASTON	81E OMEG	$20-70 \gamma p \rightarrow p4\pi$
1654 \pm 25		³ DIBIANCA	81 DBC	$\pi^+ d \rightarrow pp2(\pi^+ \pi^-)$
1666 \pm 39		¹ BACCI	80 FRAG	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1780	34	KILLIAN	80 SPEC	$11 e^- p \rightarrow 2(\pi^+ \pi^-)$
1500		⁴ ATIYA	79B SPEC	$50 \gamma C \rightarrow C4\pi^\pm$
1570 \pm 60	65	⁵ ALEXANDER	75 HBC	$7.5 \gamma p \rightarrow p4\pi$
1550 \pm 60		² CONVERSI	74 OSPK	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
1550 \pm 50	160	SCHACHT	74 STRC	$5.5-9 \gamma p \rightarrow p4\pi$
1450 \pm 100	340	SCHACHT	74 STRC	$9-18 \gamma p \rightarrow p4\pi$
1430 \pm 50	400	BINGHAM	72B HBC	$9.3 \gamma p \rightarrow p4\pi$

¹ Simple relativistic Breit-Wigner fit with model dependent width.

² Simple relativistic Breit-Wigner fit with constant width.

³ One peak fit result.

⁴ Parameters roughly estimated, not from a fit.

⁵ Skew mass distribution compensated by Ross-Stodolsky factor.

 $\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1660 \pm 30	ATKINSON	85B OMEG	$20-70 \gamma p$

3($\pi^+\pi^-$) AND 2($\pi^+\pi^-\pi^0$) MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1730 \pm 34	¹ FRABETTI	04 E687	$\gamma p \rightarrow 3\pi^+ 3\pi^- p$
1783 \pm 15	CLEGG	90 RVUE	$e^+ e^- \rightarrow 3(\pi^+ \pi^-) 2(\pi^+ \pi^- \pi^0)$

¹ From a fit with two resonances with the JACOB 72 continuum.

$$m_{\rho(1700)^0} = m_{\rho(1700)^\pm}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-48.30 \pm 83.81	¹ BARTOS	17A RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

 $\rho(1700)$ WIDTH **$\eta\rho^0$ AND $\pi^+\pi^-$ MODES**

VALUE (MeV)	DOCUMENT ID
250 \pm 100	OUR ESTIMATE

NODE=M065M2

NODE=M065M2

NODE=M065M2;LINKAGE=A

NODE=M065M2;LINKAGE=AN

NODE=M065M4

NODE=M065M4

OCCUR=2

NODE=M065M;LINKAGE=A

NODE=M065M4;LINKAGE=M

NODE=M065M;LINKAGE=O

NODE=M065M;LINKAGE=C

NODE=M065M;LINKAGE=D

NODE=M065M5

NODE=M065M5

NODE=M065M7

NODE=M065M7

NODE=M065M;LINKAGE=PI

NODE=M065DM

NODE=M065DM

NODE=M065DM;LINKAGE=A

NODE=M065210

NODE=M065W0

NODE=M065W0

→ UNCHECKED ←

$\eta\rho^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

• • • We do not use the following data for averages, fits, limits, etc. • • •

132±40	7.4k	¹ ACHASOV	18	SND $e^+e^- \rightarrow \eta\pi^+\pi^-$
150±30		ANTONELLI	88	DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$
282±44		² FUKUI	88	SPEC $8.95\pi^-p \rightarrow \eta\pi^+\pi^-n$

- ¹ From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.
² Assuming $\rho^+f_0(1370)$ decay mode interferes with $a_1(1260)^+\pi^-$ background. From a two Breit-Wigner fit.

NODE=M065W6
NODE=M065W6

 $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

• • • We do not use the following data for averages, fits, limits, etc. • • •

268.98± 11.40		¹ BARTOS	17	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
489.58± 16.95		² BARTOS	17A	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
414.71±119.48		³ BARTOS	17A	RVUE $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
109 ± 19	20K	⁴ LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
310 ± 30	+25 -35	63.5k	⁵ ABRAMOWICZ12	ZEUS $e p \rightarrow e \pi^+\pi^-p$
316 ± 26		⁶ LEES	12G	BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$
164 ± 21	+89 -26	5.4M	^{7,8} FUJIKAWA	08 BELL $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
275 ± 45		⁹ ABELE	97	CBAR $\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
310 ± 40		⁹ BERTIN	97C	OBLX $0.0\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
400 ± 100		CLEGG	94	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
224 ± 22		BISELLLO	89	DM2 $e^+e^- \rightarrow \pi^+\pi^-$
242.5 ± 163.0		DUBNICKA	89	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
620 ± 60		GESHKEN...	89	RVUE
<315		¹⁰ ERKAL	85	RVUE 20-70 $\gamma p \rightarrow \gamma\pi$
280 ± 30	-80	ABE	84B	HYBR 20 $\gamma p \rightarrow \pi^+\pi^-p$
230 ± 80		¹¹ ASTON	80	OMEG 20-70 $\gamma p \rightarrow p2\pi$
283 ± 14		¹² ATIYA	79B	SPEC 50 $\gamma C \rightarrow C2\pi$
175 ± 98	-53	BECKER	79	ASPK 17 π^-p polarized
232 ± 34		¹⁰ LANG	79	RVUE
340		¹⁰ MARTIN	78C	RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$
300 ± 100		¹⁰ FROGGATT	77	RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$
180 ± 50		¹³ HYAMS	73	ASPK 17 $\pi^-p \rightarrow \pi^+\pi^-n$

NODE=M065W1
NODE=M065W1

- ¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
² Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
³ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.
⁴ From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.
⁵ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
⁶ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
⁷ $|F_\pi(0)|^2$ fixed to 1.
⁸ From the GOUNARIS 68 parametrization of the pion form factor.
⁹ T-matrix pole.
¹⁰ From phase shift analysis of HYAMS 73 data.
¹¹ Simple relativistic Breit-Wigner fit with constant width.
¹² An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.
¹³ Included in BECKER 79 analysis.

NODE=M065W1;LINKAGE=C

NODE=M065W1;LINKAGE=D

NODE=M065W1;LINKAGE=E

NODE=M065W1;LINKAGE=A

NODE=M065W1;LINKAGE=AB

NODE=M065W1;LINKAGE=LE

NODE=M065W1;LINKAGE=FU

NODE=M065W1;LINKAGE=GO

NODE=M065W;LINKAGE=QQ

NODE=M065W;LINKAGE=P

NODE=M065W;LINKAGE=M

NODE=M065W;LINKAGE=R

NODE=M065W;LINKAGE=H

KK MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
187.2 ± 26.7	27k	¹ ABELE	99D	CBAR	± 0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$
265 ± 120	1600	CLELAND	82B	SPEC	± 50 $\pi p \rightarrow K_S^0 K^\pm p$

¹ K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

NODE=M065W2

NODE=M065W2

2 ($\pi^+ \pi^-$) MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
510 ± 40		¹ CORDIER	82	DM1 $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
400 ± 50		² ASTON	81E	OMEG 20–70 $\gamma p \rightarrow p4\pi$
400 ± 146		³ DIBIANCA	81	DBC $\pi^+ d \rightarrow pp2(\pi^+ \pi^-)$
700 ± 160		¹ BACCI	80	FRAG $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
100	34	KILLIAN	80	SPEC 11 $e^- p \rightarrow 2(\pi^+ \pi^-)$
600		⁴ ATIYA	79B	SPEC 50 $\gamma C \rightarrow C4\pi^\pm$
340 ± 160	65	⁵ ALEXANDER	75	HBC 7.5 $\gamma p \rightarrow p4\pi$
360 ± 100		² CONVERSI	74	OSPK $e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
400 ± 120	160	⁶ SCHACHT	74	STRC 5.5–9 $\gamma p \rightarrow p4\pi$
850 ± 200	340	⁶ SCHACHT	74	STRC 9–18 $\gamma p \rightarrow p4\pi$
650 ± 100	400	BINGHAM	72B	HBC 9.3 $\gamma p \rightarrow p4\pi$

¹ Simple relativistic Breit-Wigner fit with model-dependent width.

² Simple relativistic Breit-Wigner fit with constant width.

³ One peak fit result.

⁴ Parameters roughly estimated, not from a fit.

⁵ Skew mass distribution compensated by Ross-Stodolsky factor.

⁶ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M065W2;LINKAGE=AN

NODE=M065W4

NODE=M065W4

OCCUR=2

NODE=M065W;LINKAGE=A

NODE=M065W4;LINKAGE=M

NODE=M065W;LINKAGE=O

NODE=M065W;LINKAGE=C

NODE=M065W;LINKAGE=D

NODE=M065W;LINKAGE=E

NODE=M065W5

NODE=M065W5

 $\pi^+ \pi^- \pi^0 \pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
300 ± 50	ATKINSON	85B	OMEG 20–70 γp

 $\omega \pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 350 to 580 ¹ ACHASOV 00I SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
- 490 to 1040 ² ACHASOV 00I SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
- ¹ Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00I on $e^+ e^- \rightarrow \omega \pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega \pi^- \nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.
- ² Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00I on $e^+ e^- \rightarrow \omega \pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega \pi^- \nu_\tau$.

NODE=M065W9

NODE=M065W9

OCCUR=2

NODE=M065W;LINKAGE=I1

NODE=M065W;LINKAGE=I2

NODE=M065W7

NODE=M065W7

NODE=M065W;LINKAGE=PI

NODE=M065DW

NODE=M065DW

3($\pi^+ \pi^-$) AND 2($\pi^+ \pi^- \pi^0$) MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
315 ± 100	¹ FRABETTI 04	E687	$\gamma p \rightarrow 3\pi^+ 3\pi^- p$
285 ± 20	CLEGG	90	RVUE $e^+ e^- \rightarrow 3(\pi^+ \pi^-) 2(\pi^+ \pi^- \pi^0)$

¹ From a fit with two resonances with the JACOB 72 continuum.

$$\Gamma_{\rho(1700)^0} - \Gamma_{\rho(1700)^\pm}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 74.87 ± 120.67 ¹ BARTOS 17A RVUE $e^+ e^- \rightarrow \pi^+ \pi^-, \tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
- ¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M065DW;LINKAGE=A

$\rho(1700)$ DECAY MODES

NODE=M065215;NODE=M065

Mode	Fraction (Γ_i/Γ)
Γ_1 4π	
Γ_2 $2(\pi^+ \pi^-)$	seen
Γ_3 $\rho \pi \pi$	seen
Γ_4 $\rho^0 \pi^+ \pi^-$	seen
Γ_5 $\rho^0 \pi^0 \pi^0$	seen
Γ_6 $\rho^\pm \pi^\mp \pi^0$	seen
Γ_7 $a_1(1260)\pi$	seen
Γ_8 $h_1(1170)\pi$	seen
Γ_9 $\pi(1300)\pi$	seen
Γ_{10} $\rho\rho$	seen
Γ_{11} $\pi^+ \pi^-$	seen
Γ_{12} $\pi\pi$	seen
Γ_{13} $K\bar{K}^*(892) + \text{c.c.}$	seen
Γ_{14} $\eta\rho$	seen
Γ_{15} $a_2(1320)\pi$	not seen
Γ_{16} KK	seen
Γ_{17} $e^+ e^-$	seen
Γ_{18} $\pi^0 \omega$	seen
Γ_{19} $\pi^0 \gamma$	not seen

 $\rho(1700) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the cross-section into channel i in $e^+ e^-$ annihilation.

$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_{17}/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.6 ± 0.2	DELCOURT	81B DM1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$
2.83 ± 0.42	BACCI	80 FRAG	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)$

NODE=M065225

NODE=M065225

$\Gamma(\pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{11} \Gamma_{17}/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.13	¹ DIEKMAN	88 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
0.029 ^{+0.016} _{-0.012}	KURDADZE	83 OLYA	0.64–1.4 $e^+ e^- \rightarrow \pi^+ \pi^-$

NODE=M065G4
NODE=M065G4

$\Gamma(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{13} \Gamma_{17}/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.305 ± 0.071	¹ BIZOT	80 DM1	$e^+ e^-$

NODE=M065G4;LINKAGE=B

NODE=M065G10
NODE=M065G10

$\Gamma(\eta\rho) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{14} \Gamma_{17}/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
84±26±4	¹ LEES	18 BABR	$e^+ e^- \rightarrow \eta \pi^+ \pi^-$
7 ± 3	ANTONELLI	88 DM2	$e^+ e^- \rightarrow \eta \pi^+ \pi^-$

NODE=M065G;LINKAGE=M

NODE=M065G11
NODE=M065G11

$\Gamma(K\bar{K}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{16} \Gamma_{17}/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.035 ± 0.029	¹ BIZOT	80 DM1	$e^+ e^-$

NODE=M065G11;LINKAGE=A

NODE=M065G5
NODE=M065G5

NODE=M065G5;LINKAGE=M

¹ Includes non-resonant contribution. The selected fit model includes three ρ excited states.

Model uncertainty is 80%.

$\Gamma(\rho\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_{17}/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
3.510 ± 0.090	¹ BIZOT	80	DM1 e^+e^-
¹ Model dependent.			

$\rho(1700) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$	$\Gamma_{18}/\Gamma \times \Gamma_{17}/\Gamma$			
$\Gamma(\pi^0\omega)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{18}/\Gamma \times \Gamma_{17}/\Gamma$			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
0.09 ± 0.05	10.2k	¹ ACHASOV	16D	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.7 ± 0.4	7815	² ACHASOV	13	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$

- ¹ From a phenomenological model based on vector meson dominance with interfering $\rho(700)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainty not estimated. Supersedes ACHASOV 13.
- ² From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{14}/\Gamma \times \Gamma_{17}/\Gamma$			
VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$8.3^{+3.8}_{-3.1}$	7.4k	¹ ACHASOV	18	SND $e^+e^- \rightarrow \eta\pi^+\pi^-$
¹ From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.				

$\rho(1700)$ BRANCHING RATIOS

$\Gamma(\rho\pi\pi)/\Gamma(4\pi)$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.28 ± 0.06	¹ ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$

¹ $\omega\pi$ not included.

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$	Γ_4/Γ_2			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
~ 1.0		DELCOURT	81B	DM1 $e^+e^- \rightarrow 2(\pi^+\pi^-)$
0.7 ± 0.1	500	SCHACHT	74	STRC $5.5\text{--}18 \gamma p \rightarrow p4\pi$
0.80		¹ BINGHAM	72B	HBC $9.3 \gamma p \rightarrow p4\pi$

¹ The $\pi\pi$ system is in S-wave.

$\Gamma(\rho^0\pi^0\pi^0)/\Gamma(\rho^\pm\pi^\mp\pi^0)$	Γ_5/Γ_6			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.10		ATKINSON	85B	OMEG $20\text{--}70 \gamma p$
<0.15		ATKINSON	82	OMEG 0 $20\text{--}70 \gamma p \rightarrow p4\pi$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$	Γ_7/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.16 ± 0.05	¹ ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$

¹ $\omega\pi$ not included.

$\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$	Γ_8/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
0.17 ± 0.06	¹ ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$

¹ $\omega\pi$ not included.

NODE=M065G12
NODE=M065G12

NODE=M065G12;LINKAGE=M

NODE=M065240

NODE=M065R01
NODE=M065R01

OCCUR=4

NODE=M065R01;LINKAGE=B

NODE=M065R01;LINKAGE=AC

NODE=M065R00
NODE=M065R00

NODE=M065R00;LINKAGE=A

NODE=M065230

NODE=M065R19
NODE=M065R19

NODE=M065R;LINKAGE=BL

NODE=M065R1
NODE=M065R1

NODE=M065R1;LINKAGE=S

NODE=M065R6
NODE=M065R6

NODE=M065R15
NODE=M065R15

NODE=M065R15;LINKAGE=BL

NODE=M065R16
NODE=M065R16

NODE=M065R16;LINKAGE=BL

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.30±0.10	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$	
$1\omega\pi$ not included.				

NODE=M065R17
NODE=M065R17

NODE=M065R17;LINKAGE=BL

 $\Gamma(\rho\rho)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.09±0.03	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$	
$1\omega\pi$ not included.				

NODE=M065R18
NODE=M065R18

NODE=M065R18;LINKAGE=BL

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.287 ^{+0.043} _{-0.042}	BECKER	79	ASPK 17 $\pi^- p$ polarized	
0.15 to 0.30	¹ MARTIN	78C	RVUE 17 $\pi^- p \rightarrow \pi^+ \pi^- n$	
<0.20	² COSTA...	77B	RVUE $e^+ e^- \rightarrow 2\pi, 4\pi$	
0.30 ± 0.05	¹ FROGGATT	77	RVUE 17 $\pi^- p \rightarrow \pi^+ \pi^- n$	
<0.15	³ EISENBERG	73	HBC 5 $\pi^+ p \rightarrow \Delta^{++} 2\pi$	
0.25 ± 0.05	⁴ HYAMS	73	ASPK 17 $\pi^- p \rightarrow \pi^+ \pi^- n$	

NODE=M065R5
NODE=M065R5¹ From phase shift analysis of HYAMS 73 data.² Estimate using unitarity, time reversal invariance, Breit-Wigner.³ Estimated using one-pion-exchange model.⁴ Included in BECKER 79 analysis. $\Gamma(\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.13±0.05	ASTON	80	OMEG 20–70 $\gamma p \rightarrow p 2\pi$	
<0.14	¹ DAVIER	73	STRC 6–18 $\gamma p \rightarrow p 4\pi$	
<0.2	² BINGHAM	72B	HBC 9.3 $\gamma p \rightarrow p 2\pi$	

NODE=M065R5;LINKAGE=P

NODE=M065R5;LINKAGE=C

NODE=M065R5;LINKAGE=E

NODE=M065R5;LINKAGE=H

 $\Gamma(\pi\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.16±0.04	^{1,2} ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$	
¹ Using ABELE 97. ² $\omega\pi$ not included.				

NODE=M065R3

NODE=M065R3

NODE=M065R3;LINKAGE=E

NODE=M065R3;LINKAGE=S

 $\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
possibly seen	COAN	04	CLEO $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$	

NODE=M065R;LINKAGE=LK

NODE=M065R20;LINKAGE=BL

 $\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.15±0.03	¹ DELCOURT	81B	DM1 $e^+ e^- \rightarrow \bar{K}K\pi$	
¹ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.				

NODE=M065R21

NODE=M065R21

NODE=M065R9

NODE=M065R9

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
possibly seen		AKHMETSHIN 00D	CMD2	$e^+ e^- \rightarrow \eta\pi^+\pi^-$	
<0.04		DONNACHIE	87B	RVUE	
<0.02	58	ATKINSON	86B	OMEG 20–70 γp	

NODE=M065R12

NODE=M065R12

 $\Gamma(\eta\rho)/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.123±0.027	¹ DELCOURT	82	DM1 $e^+ e^- \rightarrow \pi^+\pi^- \text{MM}$	
~0.1	ASTON	80	OMEG 20–70 γp	

NODE=M065R8

NODE=M065R8

NODE=M065R8

$\Gamma(\pi^+\pi^- \text{ neutrals})/\Gamma(2(\pi^+\pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.6±0.4	¹ BALLAM	74	HBC 9.3 γp
1 Upper limit. Background not subtracted.			

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

 $\Gamma(K\bar{K})/\Gamma(2(\pi^+\pi^-))$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.015±0.010	¹ DELCOURT	81B	DM1		$e^+e^- \rightarrow K\bar{K}$
<0.04	95	BINGHAM	72B	HBC 0	9.3 γp

¹ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

 $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+\text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.052±0.026	BUON	82	DM1 $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma(\pi^0\omega)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^+\omega\pi^-$
seen	1.6k	ACHASOV 12	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
not seen	2382	AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
seen		ACHASOV 97	RVUE	$e^+e^- \rightarrow \omega\pi^0$

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ACHASOV	10D	SND 1.075–2.0 $e^+e^- \rightarrow \pi^0\gamma$

¹ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. The width of the highest mass effective resonance is fixed at 315 MeV.

 $\rho(1700)$ REFERENCES

ACHASOV	18	PR D97 012008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58785
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16D	PR D94 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57618
AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
Translated from ZETFP 94 796.					
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
DUBICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETPL 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
Translated from ZETFP 130 437.					
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
FRABETTI	04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=49614
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACHASOV	00I	PL B486 29	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47931
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=41355
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
BISELLA	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740

DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN... ¹	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
DIEMAN	88	PRPL 159 99	D. Diekmann	(BONN)	REFID=40272
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
ATKINSON	86B	ZPHY C30 531	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21508
ATKINSON	85B	ZPHY C26 499	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21506
ERKAL	85	ZPHY C29 485	C. Erkal, M.G. Olsson	(WISC)	REFID=20136
ABE	84B	PRL 53 751	K. Abe <i>et al.</i>	(SLAC HFP Collab.)	REFID=21503
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
		Translated from ZETFP 37 613.			
ATKINSON	82	PL 108B 55	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21493
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)	REFID=21494
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
CORDIER	82	PL 109B 129	A. Cordier <i>et al.</i>	(LALO)	REFID=21495
DELCOURT	82	PL 113B 93	B. Delcourt <i>et al.</i>	(LALO)	REFID=21496
ASTON	81E	NP B189 15	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21487
DELCOURT	81B	Bonn Conf. 205	B. Delcourt	(ORSAY)	REFID=21490
Also		PL 109B 129	A. Cordier <i>et al.</i>	(LALO)	REFID=21495
DIBIANCA	81	PR D23 595	F.A. di Bianca <i>et al.</i>	(CASE, CMU)	REFID=21492
ASTON	80	PL 92B 215	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21478
BACCI	80	PL 95B 139	C. Bacci <i>et al.</i>	(ROMA, FRAS)	REFID=21481
BIROT	80	Madison Conf. 546	J.C. Bizot <i>et al.</i>	(LALO, MONP)	REFID=21482
KILLIAN	80	PR D21 3005	T.J. Killian <i>et al.</i>	(CORN)	REFID=21484
ATIYA	79B	PRL 43 1691	M.S. Atiya <i>et al.</i>	(COLU, ILL, FNAL)	REFID=21470
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIIM, CERN, ZEEM, CRAC)	REFID=21084
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)	REFID=20126
MARTIN	78C	ANP 114 1	A.D. Martin, M.R. Pennington	(CERN)	REFID=21661
COSTA...	77B	PL 71B 345	B. Costa de Beauregard, B. Pire, T.N. Truong	(EPOL)	REFID=21465
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
ALEXANDER	75	PL 57B 487	G. Alexander <i>et al.</i>	(TEL)	REFID=21450
BALLAM	74	NP B76 375	J. Ballam <i>et al.</i>	(SLAC, LBL, MPIM)	REFID=20610
CONVERSI	74	PL 52B 493	M. Conversi <i>et al.</i>	(ROMA, FRAS)	REFID=20637
SCHACHT	74	NP B81 205	P. Schacht <i>et al.</i>	(MPIM)	REFID=21449
DAVIER	73	NP B58 31	M. Davier <i>et al.</i>	(SLAC)	REFID=21434
EISENBERG	73	PL 43B 149	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=21435
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
BINGHAM	72B	PL 41B 635	H.H. Bingham <i>et al.</i>	(LBL, UCB, SLAC) IGJP	REFID=21426
JACOB	72	PR D5 1847	M. Jacob, R. Slansky		REFID=49668
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

 $a_2(1700)$ $I^G(J^PC) = 1^-(2^{++})$ **$a_2(1700)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1705±40 OUR AVERAGE					
1722±15±67		¹ RODAS	19	JPAC $191 \pi^- p \rightarrow \eta(\prime) \pi^- p$	
1698±44		² AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta\eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1681^{+22}_{-35}	46M	3,4 AGHASYAN	18B	COMP $190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	
$1720 \pm 10 \pm 60$		⁵ JACKURA	18	JPAC $\pi^- p \rightarrow \eta \pi^- p$	
$1726 \pm 12 \pm 25$		⁴ ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	
1675 ± 25		ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$	
$1722 \pm 9 \pm 15$	18k	⁶ SCHEGELSKY	06	RVUE $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	
1702 ± 7	80k	⁷ UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$	
$1721 \pm 13 \pm 44$	145k	LU	05	B852 $18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$	
$1737 \pm 5 \pm 7$		ABE	04	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
1767 ± 14	221	⁸ ACCIARRI	01H	L3 $\gamma\gamma \rightarrow K^0_S K^0_S, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$	
1660 ± 40		⁴ ABELE	99B	CBAR $1.94 \bar{p}p \rightarrow \pi^0 \eta\eta$	
~ 1775		⁹ GRYGOREV	99	SPEC $40 \pi^- p \rightarrow K^0_S K^0_S n$	
$1752 \pm 21 \pm 4$		ACCIARRI	97T	L3 $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	

¹ The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15data. The mass is extracted from the T-matrix pole.² T-matrix pole.³ Statistical error negligible.⁴ Breit-Wigner mass.⁵ Superseded by RODAS 19.⁶ From analysis of L3 data at 183–209 GeV.⁷ Statistical error only.⁸ Spin 2 dominant, isospin not determined, could also be $I=1$.⁹ Possibly two $J^P = 2^+$ resonances with isospins 0 and 1.

NODE=M162

NODE=M162M

NODE=M162M

NODE=M162M;LINKAGE=C

NODE=M162M;LINKAGE=TT

NODE=M162M;LINKAGE=B

NODE=M162M;LINKAGE=E

NODE=M162M;LINKAGE=D

NODE=M162M;LINKAGE=SC

NODE=M162M;LINKAGE=ST

NODE=M162M;LINKAGE=HA

NODE=M162M;LINKAGE=GR

$a_2(1700)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
258± 40 OUR AVERAGE				
247± 17±63		1 RODAS	19 JPAC	$191 \pi^- p \rightarrow \eta(\prime) \pi^- p$
265± 55		2 AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
436± 20 16	46M	3,4 AGHASYAN	18B COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
280± 10±70		5 JACKURA	18 JPAC	$\pi^- p \rightarrow \eta \pi^- p$
190± 18±30		4 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
270± 50 20		ANISOVICH	09 RVUE	$0.0 \bar{p}p, \pi N$
336± 20±20	18k	6 SCHEGELSKY	06 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
417± 19	80k	7 UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta \pi^0$
279± 49±66	145k	LU	05 B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
151± 22±24		ABE	04 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
187± 60	221	8 ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
280± 70		4 ABELE	99B CBAR	$1.94 \bar{p}p \rightarrow \pi^0 \eta\eta$
150± 110±34		ACCIARRI	97T L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

1 The coupled-channel analysis of both the $\eta\pi$ and $\eta' \pi$ systems using ADOLPH 15data.
The width is extracted from the T-matrix pole.

2 T-matrix pole.

3 Statistical error negligible.

4 Breit-Wigner width.

5 Superseded by RODAS 19.

6 From analysis of L3 data at 183–209 GeV.

7 Statistical error only.

8 Spin 2 dominant, isospin not determined, could also be $J=1$.

NODE=M162W

NODE=M162W

NODE=M162W;LINKAGE=C

NODE=M162W;LINKAGE=TT

NODE=M162W;LINKAGE=B

NODE=M162W;LINKAGE=E

NODE=M162W;LINKAGE=D

NODE=M162W;LINKAGE=SC

NODE=M162W;LINKAGE=ST

NODE=M162W;LINKAGE=HA

NODE=M162215;NODE=M162

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta\pi$	(3.7 ± 1.0) %
$\Gamma_2 \gamma\gamma$	$(1.16 \pm 0.27) \times 10^{-6}$
$\Gamma_3 \rho\pi$	seen
$\Gamma_4 f_2(1270)\pi$	seen
$\Gamma_5 K\bar{K}$	(1.9 ± 1.2) %
$\Gamma_6 \omega\pi^-\pi^0$	seen
$\Gamma_7 \omega\rho$	seen

 $a_2(1700)$ PARTIAL WIDTHS

$\Gamma(\eta\pi)$				Γ_1
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.5±2.0	870	1 SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

DESIG=4

DESIG=1

DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow

DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow

DESIG=5

DESIG=6;OUR EVAL; \rightarrow UNCHECKED \leftarrow

DESIG=7;OUR EVAL; \rightarrow UNCHECKED \leftarrow

$\Gamma(\gamma\gamma)$				Γ_2
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.05	870	1 SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M162220

NODE=M162W3

NODE=M162W3

$\Gamma(K\bar{K})$				Γ_5
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.0±3.0	870	1 SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M162W2

NODE=M162W2

1 From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W1

NODE=M162W1

$a_2(1700) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$				
$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		$(\Gamma_3 + \Gamma_4)\Gamma_2/\Gamma$		
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.04±0.02	18k	1 SCHEGELSKY	06 RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

NODE=M162W1;LINKAGE=SC

NODE=M162225

NODE=M162G1

NODE=M162G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.37^{+0.12}_{-0.08}±0.10 18k 1 SCHEGELSKY 06 RVUE $\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_5\Gamma_2/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
20.6 ± 4.2 ± 4.6	² ABE	04	BELL	$e^+ e^- \rightarrow e^+ e^- K^+ K^-$
49 ± 11 ± 13	³ ACCIARRI	01H L3		$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183\text{--}209 \text{ GeV}$

1 From analysis of L3 data at 183–209 GeV.
 2 Assuming spin 2.
 3 Spin 2 dominant, isospin not determined, could also be $I=1$.

$a_2(1700)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$			Γ_3/Γ_4	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.4 ± 0.4 ± 0.1	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

1 From analysis of L3 data at 183–209 GeV.

$a_2(1700)$ REFERENCES

RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>	
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ACCIARRI	01H	PL B501 173	M. Acciari <i>et al.</i>	(L3 Collab.)
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>	
Translated from YAF 62 513.				
ACCIARRI	97T	PL B413 147	M. Acciari <i>et al.</i>	(L3 Collab.)

$f_0(1710)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See our mini-review in the 2004 edition of this Review, Physics Letters **B592** 1 (2004). See also the mini-review on scalar mesons under $f_0(500)$ (see the index for the page number).

$f_0(1710)$ MASS

OUR EVALUATION below is based on T-matrix poles from BARBERIS 00E and BARBERIS 99D.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1704 ± 12 OUR EVALUATION				
1732 + 9 - 7	OUR AVERAGE			Error includes scale factor of 1.6. See the ideogram below.
1759 ± 6 + 14 - 25	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1750 + 6 + 29 - 7 - 18		² UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
1701 ± 5 + 9 - 2	4k	³ CHEKANOV	08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
1765 + 4 - 3	± 13	⁴ ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
1738 ± 30		ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
1740 ± 4 + 10 - 25		BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
1740 + 30 - 25		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1710 ± 25		⁵ FRENCH	99	$300 p p \rightarrow p_f(K^+ K^-)p_s$

NODE=M162G2
NODE=M162G2

NODE=M162G1;LINKAGE=SC
NODE=M162G2;LINKAGE=AB
NODE=M162G;LINKAGE=HA

NODE=M162235

NODE=M162R01
NODE=M162R01

NODE=M162R01;LINKAGE=SC

NODE=M162

REFID=59554
REFID=59471
REFID=59003
REFID=57953
REFID=56385
REFID=52719
REFID=51186
REFID=51185
REFID=51063
REFID=50459
REFID=49650
REFID=48580
REFID=48321
REFID=46904
REFID=46909
REFID=45761

NODE=M068

NODE=M068

NODE=M068M

NODE=M068M

NODE=M068M
→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

1744± 7	± 5	381	6,7 DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$		OCCUR=2
1705±11	± 5	237	6,7 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$		OCCUR=3
1706± 4	± 5	1.0k	6,7 DOBBS	15	$J/\psi \rightarrow \gamma K^+K^-$		OCCUR=4
1690± 8	± 3	349	6,7 DOBBS	15	$\psi(2S) \rightarrow \gamma K^+K^-$		
1750±13			AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+K^-\pi^0$		
1747± 5		80k	4,8 UMAN	06	E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$		
1776±15			VLADIMIRSK...	06	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$		
1790 ⁺⁴⁰ -30			9 ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$		
1760±15	⁺¹⁵ -10		9 ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^- K^+K^-$		
1670±20			4 BINON	05	GAMS 33 $\pi^- p \rightarrow \eta\eta n$		
1732±15			10 ANISOVICH	03	RVUE		
1682±16			TIKHOMIROV	03	SPEC 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_S^0 X$		
1670±26		3.6k	11 NICHITIU	02	OBLX 0 $\bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$		
1698±18			12 BARBERIS	00E	450 $p p \rightarrow p_f \eta\eta p_s$		
1770±12			13 ANISOVICH	99B	SPEC 0.6–1.2 $p\bar{p} \rightarrow \eta\eta\pi^0$		
1730±15			BARBERIS	99	OMEG 450 $p p \rightarrow p_s p_f K^+K^-$		
1750±20			BARBERIS	99B	OMEG 450 $p p \rightarrow p_s p_f \pi^+\pi^-$		
1710±12	± 11		14 BARBERIS	99D	OMEG 450 $p p \rightarrow K^+K^-, \pi^+\pi^-$		
1750±30			15 ANISOVICH	98B	RVUE Compilation		
1720±39			BAI	98H	BES $J/\psi \rightarrow \gamma\pi^0\pi^0$		
1775± 1.5		57	16 BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$		
1690±11			17 ABREU	96C	DLPH $Z^0 \rightarrow K^+K^- + X$		
1696± 5	^{+ 9} -34		18 BAI	96C	BES $J/\psi \rightarrow \gamma K^+K^-$		OCCUR=2
1781± 8	⁺¹⁰ -31		BAI	96C	BES $J/\psi \rightarrow \gamma K^+K^-$		
1768±14			BALOSHIN	95	SPEC 40 $\pi^- C \rightarrow K_S^0 K_S^0 X$		
1750±15			19 BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$		OCCUR=2
1620±16			18 BUGG	95	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$		
1748±10			20 ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$		
~1750			BREAKSTONE	93	SFM $p p \rightarrow p p \pi^+\pi^-\pi^+\pi^-$		
1744±15			21 ALDE	92D	GAM2 38 $\pi^- p \rightarrow \eta\eta n$		
1713±10			22 ARMSTRONG	89D	OMEG 300 $p p \rightarrow p p K^+K^-$		
1706±10			22 ARMSTRONG	89D	OMEG 300 $p p \rightarrow p p K_S^0 K_S^0$		OCCUR=2
1707±10			20 AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+K^-, K_S^0 K_S^0$		
1700±15			18 BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$		
1720±60			18 BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$		OCCUR=2
1638±10			23 FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+K^-, K_S^0 K_S^0$		
1690± 4			24 FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+K^-, K_S^0 K_S^0$		OCCUR=2
1698±15			20 AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$		
1720±10	± 10		18 BALTRUSAIT...	87	MRK3 $J/\psi \rightarrow \gamma K^+K^-$		
1755± 8			25 ALDE	86C	GAM2 38 $\pi^- p \rightarrow n2\eta$		
1730 ^{+ 2} -10			26 LONGACRE	86	RVUE 22 $\pi^- p \rightarrow n2K_S^0$		
1742±15			20 WILLIAMS	84	MPSF 200 $\pi^- N \rightarrow 2K_S^0 X$		
1670±50			BLOOM	83	CBAL $J/\psi \rightarrow \gamma 2\eta$		
1650±50			BURKE	82	MRK2 $J/\psi \rightarrow \gamma 2\rho$		
1640±50		27,28	EDWARDS	82D	CBAL $J/\psi \rightarrow \gamma 2\eta$		
1730±10	± 20		29 ETKIN	82C	MPS 23 $\pi^- p \rightarrow n2K_S^0$		

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² Spin 0 favored over spin 2.

³ In the SU(3) based model with a specific interference pattern of the $f_0(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

⁴ Breit-Wigner mass.

⁵ $J^P = 0^+$, supersedes by ARMSTRONG 89D.

⁶ Using CLEO-c data but not authored by the CLEO Collaboration.

⁷ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 135$ MeV.

⁸ Systematic errors not estimated.

⁹ This state may be different from $f_0(1710)$, see CLOSE 05.

¹⁰ K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0\pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+\pi^- \rightarrow \pi^+\pi^-$, $\bar{p}p \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$, $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

NODE=M068M;LINKAGE=D

NODE=M068M;LINKAGE=H

NODE=M068M;LINKAGE=HE

NODE=M068M;LINKAGE=BW

NODE=M068M;LINKAGE=C3

NODE=M068M;LINKAGE=F


NODE=M068M;LINKAGE=G

NODE=M068M;LINKAGE=CH

NODE=M068M;LINKAGE=AB

NODE=M068M;LINKAGE=KM

- 11 Decaying to $f_0(1370)\pi\pi$.
 12 T-matrix pole.
 13 Not seen by AMSLER 02.
 14 Supersedes BARBERIS 99 and BARBERIS 99B.
 15 T-matrix pole, assuming $J^P = 0^+$
 16 No J^{PC} determination.
 17 No J^{PC} determination, width not determined.
 18 $J^P = 2^+$.
 19 From a fit to the 0^+ partial wave.
 20 No J^{PC} determination.
 21 ALDE 92D combines all the GAMS-2000 data.
 22 $J^P = 2^+$, superseded by FRENCH 99.
 23 From an analysis ignoring interference with $f'_2(1525)$.
 24 From an analysis including interference with $f'_2(1525)$.
 25 Superseded by ALDE 92D.
 26 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
 27 $J^P = 2^+$ preferred.
 28 From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
 29 Superseded by LONGACRE 86.
- NODE=M068M;LINKAGE=NC
 NODE=M068M;LINKAGE=TP
 NODE=M068M;LINKAGE=NS
 NODE=M068M;LINKAGE=BD
 NODE=M068M;LINKAGE=AN
 NODE=M068M;LINKAGE=4A
 NODE=M068M;LINKAGE=A4
 NODE=M068M;LINKAGE=A3
 NODE=M068M;LINKAGE=Q0
 NODE=M068M;LINKAGE=A1
 NODE=M068M;LINKAGE=AA
 NODE=M068M;LINKAGE=C
 NODE=M068M;LINKAGE=A
 NODE=M068M;LINKAGE=B
 NODE=M068M;LINKAGE=BB
 NODE=M068M;LINKAGE=A9
 NODE=M068M;LINKAGE=B2
 NODE=M068M;LINKAGE=E
 NODE=M068M;LINKAGE=B1



$f_0(1710)$ WIDTH

OUR EVALUATION below is based on T-matrix poles from BARBERIS 00E and BARBERIS 99D.

NODE=M068W

NODE=M068W

NODE=M068W

→ UNCHECKED ←

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
123 ± 18 OUR EVALUATION				
147 ± 12 OUR AVERAGE	Error includes scale factor of 1.2.			
172 ± 10	$\frac{+32}{-16}$	5.5k	1 ABLIKIM	13N BES3 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
139 ± 11	$\frac{+96}{-50}$		2 UEHARA	13 BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
100 ± 24	$\frac{+7}{-22}$	4k	3 CHEKANOV	08 ZEUS $e p \rightarrow K_S^0 K_S^0 X$
145 ± 8	$\frac{\pm 69}{-}$		4 ABLIKIM	06V BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
125 ± 20			ABLIKIM	04E BES2 $J/\psi \rightarrow \omega K^+ K^-$
166 ± 5	$\frac{+15}{-8}$		BAI	03G BES $J/\psi \rightarrow \gamma K\bar{K}$
120 ± 50			BAI	00A BES $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
105 ± 34			5 FRENCH	99 300 $p p \rightarrow p_f(K^+K^-)p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

148	± 40	AMSLER	06	CBAR	1.64 $\bar{p}p \rightarrow K^+ K^- \pi^0$	
188	± 13	80k	4,6	UMAN	06	E835 5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
250	± 30			VLADIMIRSK...	06	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
270	± 60		7	ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$
125	± 25	± 10	4	ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
260	± 50		4	BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
144	± 30		8,9	ANISOVICH	03	RVUE
320	± 50		9,10	ANISOVICH	03	RVUE
102	± 26			TIKHOMIROV	03	SPEC 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_S^0 X$
267	± 44	3651	11	NICHITIU	02	OBLX 0 $\bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
120	± 26		12	BARBERIS	00E	450 $p p \rightarrow p_f \eta \eta p_s$
220	± 40		13,14	ANISOVICH	99B	SPEC 0.6–1.2 $p\bar{p} \rightarrow \eta \eta \pi^0$
100	± 25			BARBERIS	99	OMEG 450 $p p \rightarrow p_s p_f K^+ K^-$
160	± 30			BARBERIS	99B	OMEG 450 $p p \rightarrow p_s p_f \pi^+ \pi^-$
126	± 16	± 18	12,15	BARBERIS	99D	OMEG 450 $p p \rightarrow K^+ K^-, \pi^+ \pi^-$
250	± 140		16	ANISOVICH	98B	RVUE Compilation
30	± 7	57	17	BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$
103	± 18	± 30	18	BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$
85	± 24	± 22		BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$
56	± 19			BALOSHIN	95	SPEC 40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
160	± 40		19	BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
160	± 60	± 20	18	BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
264	± 25		20	ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
200	to 300			BREAKSTONE	93	SFM $p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
< 80	90% CL		21	ALDE	92D	GAM2 38 $\pi^- p \rightarrow \eta \eta N^*$
181	± 30		22	ARMSTRONG	89D	OMEG 300 $p p \rightarrow p p K^+ K^-$
104	± 30		22	ARMSTRONG	89D	OMEG 300 $p p \rightarrow p p K_S^0 K_S^0$
166.4	± 33.2		20	AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$
30	± 20		18	BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
350	± 150			BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
148	± 17		23	FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
184	± 6		24	FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
136	± 28		20	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
130	± 20		18	BALTRUSAIT..	87	MRK3 $J/\psi \rightarrow \gamma K^+ K^-$
122	± 74	± 15	25	LONGACRE	86	RVUE 22 $\pi^- p \rightarrow n 2 K_S^0$
57	± 38		26	WILLIAMS	84	MPSF 200 $\pi^- N \rightarrow 2 K_S^0 X$
160	± 80			BLOOM	83	CBAL $J/\psi \rightarrow \gamma 2\eta$
200	± 100			BURKE	82	MRK2 $J/\psi \rightarrow \gamma 2\rho$
220	± 100	± 70	27,28	EDWARDS	82D	CBAL $J/\psi \rightarrow \gamma 2\eta$
200	± 156	± 9	29	ETKIN	82B	MPS 23 $\pi^- p \rightarrow n 2 K_S^0$

1 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

2 Spin 0 favored over spin 2.

3 In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f'_2(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

4 Breit-Wigner width.

5 $J^P = 0^+$, supersedes by ARMSTRONG 89D.

6 Systematic errors not estimated.

7 This state may be different from $f_0(1710)$, see CLOSE 05.

8 (Solution I)

9 K-matrix pole, assuming $J^P = 0^+$, from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K}n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

10 (Solution I)

11 Decaying to $f_0(1370)\pi\pi$.

12 T-matrix pole.

13 $J^P = 0^+$.

NODE=M068W;LINKAGE=F

NODE=M068W;LINKAGE=G

NODE=M068W;LINKAGE=HE

NODE=M068W;LINKAGE=BW

NODE=M068W;LINKAGE=C3

NODE=M068W;LINKAGE=CH

NODE=M068W;LINKAGE=AB

NODE=M068W;LINKAGE=K1

NODE=M068W;LINKAGE=KM

NODE=M068W;LINKAGE=K2

NODE=M068W;LINKAGE=NC

NODE=M068W;LINKAGE=TP

NODE=M068W;LINKAGE=AV

- 14 Not seen by AMSLER 02.
 15 Supersedes BARBERIS 99 and BARBERIS 99B.
 16 T-matrix pole, assuming $J^P = 0^+$
 17 No JPC determination.
 18 $J^P = 2^+$.
 19 From a fit to the 0^+ partial wave.
 20 No JPC determination.
 21 ALDE 92D combines all the GAMS-2000 data.
 22 $J^P = 2^+$, (0^+ excluded).
 23 From an analysis ignoring interference with $f'_2(1525)$.
 24 From an analysis including interference with $f'_2(1525)$.
 25 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
 26 No JPC determination.
 27 $J^P = 2^+$ preferred.
 28 From fit neglecting nearby $f'_2(1525)$. Replaced by BLOOM 83.
 29 From an amplitude analysis of the $K_S^0 K_S^0$ system, superseded by LONGACRE 86.

$f_0(1710)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}$	seen
$\Gamma_2 \eta\eta$	seen
$\Gamma_3 \pi\pi$	seen
$\Gamma_4 \gamma\gamma$	seen
$\Gamma_5 \omega\omega$	seen

$f_0(1710) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_4/\Gamma$
<u>VALUE (eV)</u>	<u>CL%</u>
$12^{+3}_{-2}{}^{+227}_{-8}$	
UEHARA	13
BELL	
$\gamma\gamma \rightarrow K_S^0 K_S^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<480	95
ALBRECHT	90G
$\gamma\gamma \rightarrow K^+ K^-$	
<110	95
BEHREND	89C
$\gamma\gamma \rightarrow K_S^0 K_S^0$	
<280	95
ALTHOFF	85B
$\gamma\gamma \rightarrow K\bar{K}\pi$	

1 Assuming helicity 2.

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_4/\Gamma$
<u>VALUE (keV)</u>	<u>CL%</u>
<0.82	
BARATE	95
00E	
ALEP	
$\gamma\gamma \rightarrow \pi^+ \pi^-$	

1 Assuming spin 0.

$f_0(1710)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	1004
seen	349
0.36 ± 0.12	1 DOBBS
$0.38^{+0.09}_{-0.19}$	1 DOBBS
	15
	15
	RVUE
	ALBALADEJO 08
	2 LONGACRE
	86
	MPS
	$22 \pi^- p \rightarrow n2K_S^0$

1 Using CLEO-c data but not authored by the CLEO Collaboration.

2 From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.22 ± 0.12	ALBALADEJO 08 RVUE
0.18 $^{+0.03}_{-0.13}$	1 LONGACRE 86 RVUE

1 From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

NODE=M068W;LINKAGE=NS
 NODE=M068W;LINKAGE=BD
 NODE=M068W;LINKAGE=AN
 NODE=M068W;LINKAGE=4A
 NODE=M068W;LINKAGE=A3
 NODE=M068W;LINKAGE=Q0
 NODE=M068W;LINKAGE=A1
 NODE=M068W;LINKAGE=AA
 NODE=M068W;LINKAGE=B
 NODE=M068W;LINKAGE=C
 NODE=M068W;LINKAGE=D
 NODE=M068W;LINKAGE=A9
 NODE=M068M;LINKAGE=WI
 NODE=M068W;LINKAGE=B2
 NODE=M068W;LINKAGE=E
 NODE=M068W;LINKAGE=A

NODE=M068215;NODE=M068

DESIG=2;OUR EST; \rightarrow UNCHECKED ←
 DESIG=1;OUR EST; \rightarrow UNCHECKED ←
 DESIG=5;OUR EST; \rightarrow UNCHECKED ←
 DESIG=6;OUR EST; \rightarrow UNCHECKED ←
 DESIG=4

NODE=M068220

NODE=M068G2
 NODE=M068G2

NODE=M068G2;LINKAGE=F

NODE=M068G3
 NODE=M068G3

NODE=M068G;LINKAGE=Z

NODE=M068225

NODE=M068R2
 NODE=M068R2

OCCUR=2

NODE=M068R2;LINKAGE=A
 NODE=M068R;LINKAGE=L

NODE=M068R1
 NODE=M068R1

NODE=M068R1;LINKAGE=L

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	381	¹ DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
seen	237	¹ DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
not seen		AMSLER	02	CBAR 0.9 $p\bar{p} \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
0.039 ^{+0.002} -0.024		² LONGACRE	86	RVUE

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity. Γ_3/Γ

NODE=M068R5

NODE=M068R5

OCCUR=2

NODE=M068R5;LINKAGE=A

NODE=M068R5;LINKAGE=L

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.23±0.05 OUR AVERAGE				
0.64±0.27	±0.18	LEES	18A	BABR $\Gamma(1S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.41 ^{+0.11} -0.17		ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
0.2 ± 0.024 ± 0.036		BARBERIS	99D	OMEG 450 $p\bar{p} \rightarrow K^+K^-, \pi^+\pi^-$
0.39±0.14		ARMSTRONG	91	OMEG 300 $p\bar{p} \rightarrow pp\pi\pi, ppK\bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.32±0.14		ALBALADEJO	08	RVUE
<0.11	95	¹ ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+K^-$
5.8 ^{+9.1} -5.5		² ANISOVICH	02D	SPEC Combined fit

¹ Using data from ABLIKIM 04A.² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data. Γ_3/Γ_1

NODE=M068R6

NODE=M068R6

 $\Gamma(\eta\eta)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.48±0.15				
		BARBERIS	00E	450 $p\bar{p} \rightarrow p_f\eta\eta p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.46 ^{+0.70} -0.38		¹ ANISOVICH	02D	SPEC Combined fit
<0.02	90	² PROKOSHKIN	91	GA24 300 $\pi^-p \rightarrow \pi^-p\eta\eta$

¹ From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K}n$) data.² Combining results of GAM4 with those of ARMSTRONG 89D. Γ_2/Γ_1

NODE=M068R7

NODE=M068R7

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	180	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$

NODE=M068

LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)	
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BESIII Collab.)	
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	
ALBALADEJO	08	PR D101 252002	M. Albaladejo, J.A. Oller		
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)	
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	
Vladimirsk...	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)	
		Translated from YAF 69 515.			
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		
		Translated from YAF 68 998.			
CLOSE	05	PR D71 094022	F.E. Close, Q. Zhao		
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	
Tikhomirov	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		
		Translated from YAF 66 860.			
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>		
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		
		Translated from YAF 65 1583.			

REFID=58950

REFID=56805

REFID=55387

REFID=55592

REFID=52656

REFID=52275

REFID=51125

REFID=51507

REFID=51136

REFID=51063

REFID=51191

REFID=50450

REFID=50958

REFID=50780

REFID=50788

REFID=49740

REFID=50174

REFID=49653

REFID=49401

REFID=49580

REFID=49423

REFID=48580

REFID=48831

NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>		REFID=46886
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>		REFID=46616
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=44621
		Translated from YAF 58 50.			
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)	REFID=43312
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=41591
Also		SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=44696
		Translated from YAF 54 745.			
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41010
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=21694
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)	REFID=21693
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20391

 $\eta(1760)$ $I^G(J^{PC}) = 0^+(0^- +)$

OMMITTED FROM SUMMARY TABLE

Seen by DM2 in the $\rho\rho$ system (BISELLO 89B). Structure in this region has been reported before in the same system (BALTRUSAITIS 86B) and in the $\omega\omega$ system (BALTRUSAITIS 85C, BISELLO 87).

 $\eta(1760)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1751±15 OUR AVERAGE					
1768 ⁺²⁴ ₋₂₅ ± 10	465	1 ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	
1744 ^{±10} ± 15	1045	2 ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1703 ⁺¹² ₋₁₁ ± 2		3 ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	OCCUR=2
1760 ± 11	320	4 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

1 From a single-resonance fit.

2 From a partial wave analysis including $\eta(1760)$, $f_0(1710)$, $f_2(1640)$, and $f_2(1910)$.

3 From a two-resonance fit.

4 Estimated by us from various fits. Systematic uncertainties not estimated.

NODE=M114

NODE=M114

NODE=M114M

NODE=M114M

OCCUR=2

NODE=M114M;LINKAGE=ZA

NODE=M114M;LINKAGE=MA

NODE=M114M;LINKAGE=ZH

NODE=M114M;LINKAGE=A

NODE=M114W

NODE=M114W

OCCUR=2

NODE=M114W;LINKAGE=ZA

NODE=M114W;LINKAGE=MA

NODE=M114W;LINKAGE=ZH

NODE=M114W;LINKAGE=B

 $\eta(1760)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
240±30 OUR AVERAGE					
224 ⁺⁶² ₋₅₆ ± 25	465	5 ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	
244 ⁺²⁴ ₋₂₁ ± 25	1045	6 ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
42 ⁺³⁶ ₋₂₂ ± 15		7 ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	
60 ± 16	320	8 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

5 From a single-resonance fit.

6 From a partial wave analysis including $\eta(1760)$, $f_0(1710)$, $f_2(1640)$, and $f_2(1910)$.

7 From a two-resonance fit.

8 Estimated by us from various fits. Systematic uncertainties not estimated.

$\eta(1760)$ DECAY MODES

NODE=M114215;NODE=M114

Mode	Fraction (Γ_i/Γ)
Γ_1 4π	
Γ_2 $2\pi^+ 2\pi^-$	seen
Γ_3 $\pi^+ \pi^- 2\pi^0$	seen
Γ_4 $\rho^0 \rho^0$	seen
Γ_5 $\rho^+ \rho^-$	seen
Γ_6 $\omega \omega$	seen
Γ_7 $\eta' \pi^+ \pi^-$	seen
Γ_8 $\gamma \gamma$	seen

 $\eta(1760) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta' \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_7 \Gamma_8/\Gamma$
VALUE (eV)	EVTS DOCUMENT ID TECN COMMENT
$28.2^{+7.9}_{-7.5} \pm 3.7$	
465	9 ZHANG 12A BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$3.0^{+2.0}_{-1.2} \pm 0.8$	52 10 ZHANG 12A BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
$18^{+13}_{-10} \pm 5$	315 11 ZHANG 12A BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
9 From a single-resonance fit.	
10 From a two-resonance fit. For constructive interference with the $X(1835)$.	
11 From a two-resonance fit. For destructive interference with the $X(1835)$.	

 $\eta(1760)$ BRANCHING RATIOS

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	
	BISELLO 89B DM2 $J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$
$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	
	BISELLO 89B DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^- 2\pi^0$
$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$	Γ_4/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	BISELLO 89B DM2 $J/\psi \rightarrow \gamma \rho^0 \rho^0$
seen	BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \gamma \rho^0 \rho^0$
$\Gamma(\rho^+ \rho^-)/\Gamma_{\text{total}}$	Γ_5/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	BISELLO 89B DM2 $J/\psi \rightarrow \gamma \rho^+ \rho^-$
seen	BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \gamma \rho^+ \rho^-$
$\Gamma(\omega \omega)/\Gamma_{\text{total}}$	Γ_6/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	BISELLO 87 DM2 $J/\psi \rightarrow \omega \omega$
seen	BALTRUSAIT..85c MRK3 $J/\psi \rightarrow \gamma \omega \omega$
$\Gamma(\gamma\gamma)/\Gamma(\omega \omega)$	Γ_8/Γ_6
VALUE	CL% DOCUMENT ID TECN COMMENT
$<2.48 \times 10^{-3}$	90 12 ABLIKIM 180 BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$

12 Using results from ABLIKIM 06H.

 $\eta(1760)$ REFERENCES

ABLIKIM	180	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+) (Mark III Collab.)
BALTRUSAIT...86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	(Mark III Collab.)
BALTRUSAIT...86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	(CIT, UCSC+)
BALTRUSAIT...85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>		

DESIG=1
DESIG=2
DESIG=3
DESIG=4
DESIG=5
DESIG=7
DESIG=8;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=9;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M114225

NODE=M114G01
NODE=M114G01

OCCUR=2

OCCUR=3

NODE=M114G01;LINKAGE=ZH
NODE=M114G01;LINKAGE=ZA
NODE=M114G01;LINKAGE=ZN

NODE=M114210

NODE=M114R01
NODE=M114R01NODE=M114R02
NODE=M114R02NODE=M114R03
NODE=M114R03NODE=M114R04
NODE=M114R04NODE=M114R06
NODE=M114R06NODE=M114R08
NODE=M114R08

NODE=M114R08;LINKAGE=A

NODE=M114

REFID=58925
REFID=54763
REFID=51125
REFID=40575
REFID=40012
REFID=22009
REFID=22100
REFID=22095

NODE=M075

 $\pi(1800)$

$$I^G(J^{PC}) = 1^-(0^{-+})$$

See the related review on "Non- $q\bar{q}$ mesons." **$\pi(1800)$ MASS**

NODE=M075

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

1810 $^{+9}_{-11}$ OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

1804 $^{+6}_{-9}$	46M	1 AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	I
1876 $\pm 18 \pm 16$	4k	2 EUGENIO	08 B852	18 $\pi^- p \rightarrow \eta \eta \pi^- p$	
1774 $\pm 18 \pm 20$		3 CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
1863 $\pm 9 \pm 10$		4 CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	OCCUR=2
1840 $\pm 10 \pm 10$	1.2k	AMELIN	96B VES	37 $\pi^- A \rightarrow \eta \eta \pi^- A$	
1775 $\pm 7 \pm 10$		5 AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
1790 ± 14		6 BERDNIKOV	94 VES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$	
1873 $\pm 33 \pm 20$		BELADIDZE	92C VES	36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$	
1814 $\pm 10 \pm 23$	426	BITYUKOV	91 VES	36 $\pi^- C \rightarrow \pi^- \eta \eta C$	
1770 ± 30	1.1k	BELLINI	82 SPEC	40 $\pi^- A \rightarrow 3\pi A$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1785 $\pm 9^{+12}_{-6}$	420k	7 ALEKSEEV	10 COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	I
1737 $\pm 5 \pm 15$		AMELIN	99 VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$	

1 Statistical error negligible.

2 From a single-pole fit.

3 In the $f_0(980)\pi$ wave.

4 In the $f_0(500)\pi$ wave.


5 From a fit to $J^{PC} = 0^{-+}$ $f_0(980)\pi$, $f_0(1370)\pi$ waves.

6 From a fit to $J^{PC} = 0^{-+}$ $K_0^*(1430)K^-$ and $f_0(980)\pi^-$ waves.

7 Superseded by AGHASYAN 2018B.

NODE=M075M;LINKAGE=B
NODE=M075M;LINKAGE=SP
NODE=M075M;LINKAGE=C1
NODE=M075M;LINKAGE=C2
NODE=M075M;LINKAGE=AX
NODE=M075M;LINKAGE=A
NODE=M075M;LINKAGE=C

WEIGHTED AVERAGE
1810-9-11 (Error scaled by 2.2)

 **$\pi(1800)$ WIDTH**

NODE=M075W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
215₋₈⁺⁷ OUR AVERAGE						NODE=M075W
220 ₋₁₁ ⁺⁸	46M	8 AGHASYAN	18B COMP		190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	
221 ₋₃₈ ⁺²⁶	4k	9 EUGENIO	08 B852	-	18 $\pi^- p \rightarrow \eta \eta \pi^- p$	
223 ₋₅₀ ⁺⁴⁸		10 CHUNG	02 B852		18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
191 ₋₂₀ ⁺²¹		11 CHUNG	02 B852		18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	OCCUR=2
210 ₋₃₀ ⁺³⁰	1.2k	AMELIN	96B VES	-	37 $\pi^- A \rightarrow \eta \eta \pi^- A$	
190 ₋₁₅ ⁺¹⁵		12 AMELIN	95B VES	-	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
210 ₋₇₀		13 BERDNIKOV	94 VES	-	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$	
225 ₋₂₀ ⁺³⁵		BELADIDZE	92C VES	-	36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$	
205 ₋₃₂ ⁺¹⁸	426	BITYUKOV	91 VES	-	36 $\pi^- C \rightarrow \pi^- \eta \eta C$	
310 ₋₅₀	1.1k	BELLINI	82 SPEC	-	40 $\pi^- A \rightarrow 3\pi A$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
208 ₋₃₇ ⁺²¹	420k	14 ALEKSEEV	10 COMP		190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
259 ₋₆ ⁺¹⁹		AMELIN	99 VES		37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$	
8 Statistical error negligible.						
9 From a single-pole fit.						
10 In the $f_0(980)\pi^-$ wave.						
11 In the $f_0(500)\pi^-$ wave.						
12 From a fit to $J^{PC} = 0^- + f_0(980)\pi, f_0(1370)\pi$ waves.						
13 From a fit to $J^{PC} = 0^- + K_0^*(1430)K^-$ and $f_0(980)\pi^-$ waves.						
14 Superseded by AGHASYAN 2018B.						

$\pi(1800)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+ \pi^- \pi^-$	seen
$\Gamma_2 f_0(500)\pi^-$	seen
$\Gamma_3 f_0(980)\pi^-$	seen
$\Gamma_4 f_0(1370)\pi^-$	seen
$\Gamma_5 f_0(1500)\pi^-$	not seen
$\Gamma_6 \rho \pi^-$	not seen
$\Gamma_7 \eta \eta \pi^-$	seen
$\Gamma_8 a_0(980)\eta$	seen
$\Gamma_9 a_2(1320)\eta$	not seen
$\Gamma_{10} f_2(1270)\pi$	not seen
$\Gamma_{11} f_0(1370)\pi^-$	not seen
$\Gamma_{12} f_0(1500)\pi^-$	seen
$\Gamma_{13} \eta \eta'(958)\pi^-$	seen
$\Gamma_{14} K_0^*(1430)K^-$	seen
$\Gamma_{15} K^*(892)K^-$	not seen

$\pi(1800)$ BRANCHING RATIOS

$\Gamma(f_0(980)\pi^-)/\Gamma(f_0(500)\pi^-)$		Γ_3/Γ_2
0.44 _{-0.38} ^{+0.08}	15 CHUNG	02 B852
		18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
$\Gamma(f_0(980)\pi^-)/\Gamma(f_0(1370)\pi^-)$		Γ_3/Γ_4
1.7 _{-1.3} ^{+1.3}	16 AMELIN	95B VES
		36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
$\Gamma(f_0(1370)\pi^-)/\Gamma_{\text{total}}$		Γ_4/Γ
seen	BELLINI	82 SPEC
		40 $\pi^- A \rightarrow 3\pi A$

NODE=M075W;LINKAGE=B
 NODE=M075W;LINKAGE=SP
 NODE=M075W;LINKAGE=C1
 NODE=M075W;LINKAGE=C2
 NODE=M075W;LINKAGE=AX
 NODE=M075W;LINKAGE=A
 NODE=M075W;LINKAGE=C

NODE=M075215;NODE=M075

DESIG=10;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=11;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=1
 DESIG=12
 DESIG=2
 DESIG=7;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=13
 DESIG=14
 DESIG=15
 DESIG=6;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=8;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=4
 DESIG=9

NODE=M075220

NODE=M075R11
 NODE=M075R11

NODE=M075R5
 NODE=M075R5

NODE=M075R1
 NODE=M075R1

$\Gamma(f_0(1500)\pi^-)/\Gamma_{\text{total}}$						Γ_5/Γ	NODE=M075R12 NODE=M075R12
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
not seen		CHUNG	02	B852	18.3	$\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
$\Gamma(\rho\pi^-)/\Gamma_{\text{total}}$						Γ_6/Γ	NODE=M075R2 NODE=M075R2
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen		BELLINI	82	SPEC	—	$40 \pi^- A \rightarrow 3\pi A$	
$\Gamma(\rho\pi^-)/\Gamma(f_0(980)\pi^-)$						Γ_6/Γ_3	NODE=M075R6 NODE=M075R6
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<0.25		CHUNG	02	B852	18.3	$\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
<0.14	90	AMELIN	95B	VES	—	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
$\Gamma(\eta\eta\pi^-)/\Gamma(\pi^+\pi^-\pi^-)$						Γ_7/Γ_1	NODE=M075R8 NODE=M075R8
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
0.5 ± 0.1	1200	¹⁶ AMELIN	96B	VES	—	$37 \pi^- A \rightarrow \eta\eta\pi^- A$	
$\Gamma(a_2(1320)\eta)/\Gamma_{\text{total}}$						Γ_9/Γ	NODE=M075R13 NODE=M075R13
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
not seen		EUGENIO	08	B852	18	$\pi^- p \rightarrow \eta\eta\pi^- p$	
$\Gamma(f_2(1270)\pi)/\Gamma_{\text{total}}$						Γ_{10}/Γ	NODE=M075R14 NODE=M075R14
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
not seen		EUGENIO	08	B852	18	$\pi^- p \rightarrow \eta\eta\pi^- p$	
$\Gamma(f_0(1370)\pi^-)/\Gamma_{\text{total}}$						Γ_{11}/Γ	NODE=M075R15 NODE=M075R15
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
not seen		EUGENIO	08	B852	18	$\pi^- p \rightarrow \eta\eta\pi^- p$	
$\Gamma(f_0(1500)\pi^-)/\Gamma(a_0(980)\eta)$						Γ_{12}/Γ_8	NODE=M075R7 NODE=M075R7
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
0.48 ± 0.17	4k ^{16,17}	EUGENIO	08	B852	—	$18 \pi^- p \rightarrow \eta\eta\pi^- p$	
0.030 ^{+0.014} —0.011	16	ANISOVICH	01B	SPEC	0	$0.6-1.94 p\bar{p} \rightarrow \eta\eta\pi^0\pi^0$	
0.08 ± 0.03	1200 ^{16,18}	AMELIN	96B	VES	—	$37 \pi^- A \rightarrow \eta\eta\pi^- A$	
$\Gamma(\eta\eta'(958)\pi^-)/\Gamma(\eta\eta\pi^-)$						Γ_{13}/Γ_7	NODE=M075R10 NODE=M075R10
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
0.29 ± 0.07		¹⁶ BELADIDZE	92C	VES	—	$36 \pi^- Be \rightarrow \pi^- \eta' \eta Be$	
0.3 ± 0.1	426 ± 57	¹⁶ BITYUKOV	91	VES	—	$36 \pi^- C \rightarrow \pi^- \eta \eta C$	
$\Gamma(K_0^*(1430)K^-)/\Gamma_{\text{total}}$						Γ_{14}/Γ	NODE=M075R4 NODE=M075R4
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
seen		BERDNIKOV	94	VES	—	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$	
$\Gamma(K^*(892)K^-)/\Gamma_{\text{total}}$						Γ_{15}/Γ	NODE=M075R9 NODE=M075R9
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen		BERDNIKOV	94	VES	—	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$	
15 Assuming that $f_0(980)$ decays only to $\pi\pi$.							
16 Systematic errors not estimated.							
17 From a single-pole fit.							
18 Assuming that $f_0(1500)$ decays only to $\eta\eta$ and $a_0(980)$ decays only to $\eta\pi$.							

(1800) REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>	
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)
AMELIN	96B	Translated from YAF 62 487	D.V. Amelin <i>et al.</i>	(SERP, TBIL) IGJPC
		Translated from YAF 59 1021		
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)
		Translated from YAF 55 2748.		
BITYUKOV	91	PL B268 137	S.I. Bityukov <i>et al.</i>	(SERP, TBIL)
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)

NODE=M075

REFID=59471
REFID=53356
REFID=52160
REFID=48837
REFID=48318
REFID=46910
REFID=44725
REFID=44433
REFID=44073
REFID=43175
REFID=41749
REFID=21134

$f_2(1810)$ $I^G(J^{PC}) = 0^+(2^{++})$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M038

 $f_2(1810)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1815±12 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.				
1822 ⁺²⁹ ₋₂₄ ⁺⁶⁶ ₋₅₇	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1737 ^{± 9} _{- 65} ⁺¹⁹⁸		² UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta \eta$
1800 ^{± 30}	40	ALDE	88D GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$
1806 ^{± 10}	1600	ALDE	87 GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
1870 ^{± 40}		³ ALDE	86D GAM4	$100 \pi^- p \rightarrow \eta \eta n$
1857 ⁺³⁵ ₋₂₄		⁴ COSTA	80 OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1858 ⁺¹⁸ ₋₇₁		⁵ LONGACRE	86 RVUE	Compilation
1799 ^{± 15}		⁶ CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$

1 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.2 Breit-Wigner mass. Could also be the $f_2(1910)$.

3 Seen in only one solution.

4 Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.


5 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

6 From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHIN 97.

NODE=M038

NODE=M038M

NODE=M038M



NODE=M038M;LINKAGE=B

NODE=M038M;LINKAGE=UE

NODE=M038M;LINKAGE=F

NODE=M038M;LINKAGE=A

NODE=M038M;LINKAGE=L

NODE=M038M;LINKAGE=P1

 $f_2(1810)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197± 22 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
229 ^{+ 52} _{- 42} ^{+ 88} _{- 155}	5.5k	⁷ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
228 ^{+ 21} _{- 20} ^{+ 234} _{- 153}		⁸ UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta \eta$
160 ^{± 30}	40	ALDE	88D GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$

NODE=M038W

NODE=M038W

190 ± 20	1600	ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$
250 ± 30	9	ALDE	86D	GAM4	100	$\pi^- p \rightarrow \eta\eta n$
185^{+102}_{-139}	10	COSTA	80	OMEG	10	$\pi^- p \rightarrow K^+ K^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

388^{+15}_{-21}	11	LONGACRE	86	RVUE	Compilation
280^{+42}_{-35}	12	CASON	82	STRC	$8\pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$

7 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

8 Breit-Wigner width. Could also be the $f_2(1910)$.

9 Seen in only one solution.

10 Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

11 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

12 From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHIN 97.

NODE=M038W;LINKAGE=B

NODE=M038W;LINKAGE=UE


NODE=M038W;LINKAGE=F

NODE=M038W;LINKAGE=A

NODE=M038W;LINKAGE=L

NODE=M038W;LINKAGE=P1

WEIGHTED AVERAGE
197 \pm 22 (Error scaled by 1.5)



f₂(1810) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	
$\Gamma_2 \eta\eta$	seen
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 K^+ K^-$	
$\Gamma_5 \gamma\gamma$	seen

NODE=M038215;NODE=M038

DESIG=2

DESIG=3

DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow

DESIG=1

DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M038225

NODE=M038G01

NODE=M038G01

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID TECN COMMENT
$5.2^{+0.9+37.3}_{-0.8-4.5}$	13 UEHARA 10A BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$

13 Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

NODE=M038G01;LINKAGE=UE

$f_2(1810)$ BRANCHING RATIOS **$\Gamma(\pi\pi)/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$	
not seen	PROKOSHKIN 97	GAM2	38 $\pi^- p \rightarrow \pi^0\pi^0n$	
$0.21^{+0.02}_{-0.03}$	14 LONGACRE 86	RVUE	Compilation	
0.44 ± 0.03	15 CASON 82	STRC	8 $\pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$	

14 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

15 Included in LONGACRE 86 global analysis.

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	ABLIKIM 13N	BES3	PWA of $J/\psi \rightarrow \gamma\eta\eta$	I
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$0.008^{+0.028}_{-0.003}$ 16 LONGACRE 86 RVUE Compilation

16 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

 $\Gamma(\pi\pi)/\Gamma(4\pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_3
• • • We do not use the following data for averages, fits, limits, etc. • • •				

<0.75 ALDE 87 GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$

 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.8 ± 0.3 ALDE 87 GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				

$0.003^{+0.019}_{-0.002}$ 17 LONGACRE 86 RVUE Compilation

seen COSTA 80 OMEG 10 $\pi^- p \rightarrow K^+K^- n$

17 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038220

NODE=M038R2

NODE=M038R2

NODE=M038R2;LINKAGE=L

NODE=M038R;LINKAGE=C

NODE=M038R3

NODE=M038R3

NODE=M038R3;LINKAGE=L

NODE=M038R4

NODE=M038R4

NODE=M038R5

NODE=M038R5

NODE=M038R1

NODE=M038R1

NODE=M038R1;LINKAGE=L

NODE=M038

REFID=55387

REFID=53641

REFID=52166

REFID=48580

REFID=45386

REFID=44652

REFID=40221

REFID=20765

REFID=20768

REFID=20746

REFID=20737

 $f_2(1810)$ REFERENCES

ABLIKIM 13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BESIII Collab.)
UEHARA 10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
PDG 08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>	
PROKOSHKIN 97	PD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)
		Translated from DANS 353 323.	
ALDE 88D	SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 47 1273.	
ALDE 87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE 86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE 86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
CASON 82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
COSTA 80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)

NODE=M085

X(1835)

$I^G(J^{PC}) = ?^?(0^-+)$

OMITTED FROM SUMMARY TABLE

Could be a superposition of two states, one with small width appearing as threshold enhancement in $p\bar{p}$, the other one with a larger width. For the former ABLIKIM 12D determine $J^{PC} = 0^-+$.

NODE=M085

X(1835) MASS

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	NODE
1826.5^{+13.0}_{-3.4} OUR AVERAGE						NODE=M085M
1825.3 \pm 2.4 ^{+17.3} _{-2.4}			1 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	OCCUR=2
1844 \pm 9 ⁺¹⁶ ₋₂₅			ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1839 \pm 26 \pm 26			2 ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$	
1909.5 \pm 15.9 ^{+9.4} _{-27.5}			3 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	
1842.2 \pm 4.2 ^{+7.1} _{-2.6}	0.6k		ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$	
1832 \pm 5 ⁺¹⁹ ₋₅ \pm 26			4 ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$	
1836.5 \pm 3.0 ^{+5.6} _{-2.1}	4265		5 ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	
1877.3 \pm 6.3 ^{+3.4} _{-7.4}			6 ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$	
1837 \pm 12 ⁺⁹ ₋₇	231	7,8 ALEXANDER	10 CLEO		$J/\psi \rightarrow \gamma p\bar{p}$	
1833.7 \pm 6.1 \pm 2.7	264		ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	
1831 \pm 7		8,9 ABLIKIM		05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$	OCCUR=2
1859 \pm 10 ⁺³ ₋₁₀ \pm 5		8 BAI		03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$	
1 From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors.						NODE=M085M;LINKAGE=B
2 From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded.						NODE=M085M;LINKAGE=C
3 Pole mass from a fit of the measured $\pi^+\pi^-\eta'$ lineshape to a Flatté formula that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold; the fit also includes known backgrounds and contributors, as well as an <i>ad hoc</i> Breit-Wigner function ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit.						NODE=M085M;LINKAGE=A
4 From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.						NODE=M085M;LINKAGE=AK
5 From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-$.						NODE=M085M;LINKAGE=AI
6 The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$.						NODE=M085M;LINKAGE=BL
7 From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.						NODE=M085M;LINKAGE=AE
8 Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, DEL-AMO-SANCHEZ 12 in $B \rightarrow D(D^*)p\bar{p}(\pi)$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$.						NODE=M085M;LINKAGE=HF
9 From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A. Systematic errors not estimated.						NODE=M085M;LINKAGE=AB

X(1835) WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	NODE
242⁺¹⁴₋₁₅ OUR AVERAGE						NODE=M085W
245.2 \pm 13.1 ^{+4.6} _{-9.6}			1 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	OCCUR=2
192 ⁺²⁰ ₋₁₇ \pm 62 ₋₄₃			ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

175 ± 57 ± 25		2 ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
273.5 ± 21.4 ± 6.1		3 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
83 ± 14 ± 11	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
< 76	90	4 ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
190 ± 9 ± 38	4265	5 ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
57 ± 12 ± 19		6 ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
0 ± 44	231	7,8 ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
67.7 ± 20.3 ± 7.7	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
< 153	90	8,9 ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$
< 30		8 BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors.

² From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded.

³ Pole width from a fit of the measured $\pi^+\pi^-\eta'$ lineshape to a Flatté formula that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold; the fit also includes known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit.

⁴ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.

⁵ From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-$.

⁶ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$.

⁷ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

⁸ Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, DEL-AMO-SANCHEZ 12 in $B \rightarrow D(D^*)p\bar{p}(\pi)$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$.

⁹ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Systematic errors not estimated.

OCCUR=2

NODE=M085W;LINKAGE=B

NODE=M085W;LINKAGE=C

NODE=M085W;LINKAGE=A

NODE=M085W;LINKAGE=AK

NODE=M085W;LINKAGE=AI

NODE=M085W;LINKAGE=BL

NODE=M085W;LINKAGE=AE

NODE=M085W;LINKAGE=HF

NODE=M085W;LINKAGE=AB

NODE=M085215;NODE=M085

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=2
 DESIG=4
 DESIG=5;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=6
 DESIG=7

NODE=M085225

NODE=M085G01
 NODE=M085G01

OCCUR=2

NODE=M085G01;LINKAGE=ZH

NODE=M085G01;LINKAGE=ZA

NODE=M085220

NODE=M085R01
 NODE=M085R01

X(1835) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 p\bar{p}$	seen
$\Gamma_2 \eta'\pi^+\pi^-$	seen
$\Gamma_3 \gamma\gamma$	
$\Gamma_4 K_S^0 K_S^0\eta$	seen
$\Gamma_5 \gamma\phi(1020)$	possibly seen
$\Gamma_6 3(\pi^+\pi^-)$	seen

X(1835) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_3/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 35.6	90	1 ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
< 83	90	2 ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

¹ From a two-resonance fit and constructive interference of the $\eta(1760)$ and $X(1835)$, a significance of 2.8 σ .

² From a two-resonance fit and destructive interference of the $\eta(1760)$ and $X(1835)$, a significance of 2.8 σ .

X(1835) BRANCHING RATIOS

$\Gamma(p\bar{p})/\Gamma(\eta'\pi^+\pi^-)$	Γ_1/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
0.333	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\eta'\pi^+\pi^-)/\Gamma(K_S^0 K_S^0 \eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.7±1.8	¹ ABLIKIM	15T	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$	

¹ Using results from ABLIKIM 05R. $\Gamma(\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen				
¹ ABLIKIM 16J quotes $B(J/\psi \rightarrow \gamma X(1835)) \times B(X(1835) \rightarrow \pi^+ \pi^- \eta') = (3.93 \pm 0.38^{+0.31}_{-0.84}) \times 10^{-4}$ from a fit of the measured $\pi^+ \pi^- \eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatté formula in addition to known backgrounds and contributors, as well as an <i>ad hoc</i> Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.	¹ ABLIKIM	16J	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$	

 $\Gamma(\gamma\phi(1020))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
possibly seen				
¹ ABLIKIM 18I	¹ ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma \gamma \phi(1020)$	

¹ Seen as a peak in $\gamma\phi$ invariant mass. Angular analysis consistent with $JPC = 0-+$. Other JPC not excluded. $\Gamma(\gamma\gamma)/\Gamma(\eta'\pi^+\pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
$<9.80 \times 10^{-3}$	90	¹ ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$	

¹ Using results from ABLIKIM 16J. $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
seen	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$	

X(1835) REFERENCES

ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	12	PR D85 092017	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10G	CP C34 421	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT,B	05L	PR D72 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M085R00
NODE=M085R00

NODE=M085R00;LINKAGE=A

NODE=M085R03
NODE=M085R03

NODE=M085R03;LINKAGE=A

NODE=M085R04
NODE=M085R04

NODE=M085R04;LINKAGE=A

NODE=M085R05
NODE=M085R05

NODE=M085R05;LINKAGE=A

NODE=M085R06
NODE=M085R06

NODE=M085

REFID=58893
REFID=58925
REFID=57454
REFID=56785
REFID=55582
REFID=54269
REFID=54286
REFID=54763
REFID=53684
REFID=53931
REFID=55685
REFID=53525
REFID=52086
REFID=50993
REFID=50985
REFID=50827
REFID=51038
REFID=50651
REFID=49473
REFID=48690
REFID=48980

$\phi_3(1850)$

$I^G(J^{PC}) = 0^-(3^- -)$

NODE=M054

$\phi_3(1850)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1854± 7 OUR AVERAGE				
1855±10		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda, K_S^0 K^\pm \pi^\mp \Lambda$
1870 ⁺³⁰ ₋₂₀	430	ARMSTRONG 82	OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$
1850±10	123	ALHARRAN 81B	HBC	8.25 $K^- p \rightarrow K\bar{K}\Lambda$

NODE=M054M

NODE=M054M

$\phi_3(1850)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
87⁺²⁸₋₂₃ OUR AVERAGE Error includes scale factor of 1.2.				
64±31		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda, K_S^0 K^\pm \pi^\mp \Lambda$
160 ⁺⁹⁰ ₋₅₀	430	ARMSTRONG 82	OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$
80 ⁺⁴⁰ ₋₃₀	123	ALHARRAN 81B	HBC	8.25 $K^- p \rightarrow K\bar{K}\Lambda$

NODE=M054W

NODE=M054W

$\phi_3(1850)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}$	seen
$\Gamma_2 K\bar{K}^*(892) + \text{c.c.}$	seen

NODE=M054215;NODE=M054

$\phi_3(1850)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.55^{+0.85}_{-0.45}	ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda, K_S^0 K^\pm \pi^\mp \Lambda$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.8 ± 0.4	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K\bar{K}\pi\Lambda$	

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M054220

NODE=M054R1
NODE=M054R1

$\phi_3(1850)$ REFERENCES

ASTON 88E PL B208 324	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) IGJPC
ARMSTRONG 82 PL 110B 77	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+) JP
ALHARRAN 81B PL 101B 357	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)

NODE=M054

REFID=40577
REFID=21405
REFID=21702

$\eta_2(1870)$

$I^G(J^{PC}) = 0^+(2^-)$

Needs confirmation.

$\eta_2(1870)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1842± 8 OUR AVERAGE				
1835±12		BARBERIS	00B	$450 \text{ } pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1844±13		BARBERIS	00C	$450 \text{ } pp \rightarrow p_f 4\pi p_s$
1840±25		BARBERIS	97B OMEG	$450 \text{ } pp \rightarrow pp2(\pi^+ \pi^-)$
1875±20±35		ADOMEIT	96 CBAR	$1.94 \bar{p}p \rightarrow \eta 3\pi^0$
1881±32±40	26	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1860± 5±15		ANISOVICH	00E SPEC	$0.9-1.94 \bar{p}p \rightarrow \eta 3\pi^0$
1840±15		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\eta_2(1870)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
225±14 OUR AVERAGE				
235±22		BARBERIS	00B	$450 \text{ } pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
228±23		BARBERIS	00C	$450 \text{ } pp \rightarrow p_f 4\pi p_s$
200±40		BARBERIS	97B OMEG	$450 \text{ } pp \rightarrow pp2(\pi^+ \pi^-)$
200±25±45		ADOMEIT	96 CBAR	$1.94 \bar{p}p \rightarrow \eta 3\pi^0$
221±92±44	26	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
250±25 ⁺⁵⁰ ₋₃₅		ANISOVICH	00E SPEC	$0.9-1.94 \bar{p}p \rightarrow \eta 3\pi^0$
170±40		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

$\eta_2(1870)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta \pi \pi$	
$\Gamma_2 a_2(1320) \pi$	
$\Gamma_3 f_2(1270) \eta$	
$\Gamma_4 a_0(980) \pi$	
$\Gamma_5 \gamma \gamma$	seen

$\eta_2(1870)$ BRANCHING RATIOS

$\Gamma(a_2(1320)\pi)/\Gamma(f_2(1270)\eta)$	Γ_2/Γ_3
1.7 ±0.4 OUR AVERAGE	
1.60±0.40	¹ ANISOVICH 11 SPEC $0.9-1.94 p\bar{p}$
20.4 ±6.6	BARBERIS 00B $450 \text{ } pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
4.1 ±2.3	ADOMEIT 96 CBAR $1.94 \bar{p}p \rightarrow \eta 3\pi^0$

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

$\Gamma(a_2(1320)\pi)/\Gamma(a_0(980)\pi)$	Γ_2/Γ_4
32.6±12.6	
	BARBERIS 00B $450 \text{ } pp \rightarrow p_f \eta \pi^+ \pi^- p_s$

$\Gamma(a_0(980)\pi)/\Gamma(f_2(1270)\eta)$	Γ_4/Γ_3
0.48±0.45	
	¹ ANISOVICH 11 SPEC $0.9-1.94 p\bar{p}$

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_5/Γ
seen	
	KARCH 92 CBAL $e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$

NODE=M101

NODE=M101

NODE=M101M

NODE=M101M

NODE=M101W

NODE=M101W

NODE=M101225;NODE=M101

DESIG=1

DESIG=4

DESIG=8

DESIG=2

DESIG=9

NODE=M101230

NODE=M101R2

NODE=M101R2

NODE=M101R2;LINKAGE=AN

NODE=M101R4

NODE=M101R4

NODE=M101R01

NODE=M101R01

NODE=M101R01;LINKAGE=AN

NODE=M101R02

NODE=M101R02

$\eta_2(1870)$ REFERENCES

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)

NODE=M101


REFID=53631
 REFID=47945
 REFID=47958
 REFID=47959
 REFID=46606
 REFID=45758
 REFID=45202
 REFID=42170

 $\pi_2(1880)$

$$\mathcal{I}^G(J^{PC}) = 1^-(2^-+)$$

 $\pi_2(1880)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG COMMENT
1874⁺²⁶₋₅ OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.			
1847 ⁺²⁰ ₋₃	46M	¹ AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1929 \pm 24 \pm 18	4k	EUGENIO	08 B852	— 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
1876 \pm 11 \pm 67	145k	LU	05 B852	— 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
2003 \pm 88 \pm 148	69k	KUHN	04 B852	— 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1880 \pm 20		ANISOVICH	01B SPEC 0	0.6–1.94 $\bar{p}p \rightarrow \eta \eta \pi^0 \pi^0$

¹ Statistical error negligible.

NODE=M185M

NODE=M185M

NODE=M185M;LINKAGE=A

 $\pi_2(1880)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG COMMENT
237^{+ 33}₋₃₀ OUR AVERAGE	Error includes scale factor of 1.2.			
246 ^{+ 33} ₋₂₈	46M	² AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
323 \pm 87 \pm 43	4k	EUGENIO	08 B852	— 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
146 \pm 17 \pm 62	145k	LU	05 B852	— 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
306 \pm 132 \pm 121	69k	KUHN	04 B852	— 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
255 \pm 45		ANISOVICH	01B SPEC 0	0.6–1.94 $\bar{p}p \rightarrow \eta \eta \pi^0 \pi^0$

² Statistical error negligible.

NODE=M185W

NODE=M185W

NODE=M185W;LINKAGE=A

$\pi_2(1880)$ DECAY MODES

NODE=M185215;NODE=M185

Mode
$\Gamma_1 \eta\eta\pi^-$
$\Gamma_2 a_0(980)\eta$
$\Gamma_3 a_2(1320)\eta$
$\Gamma_4 f_0(1500)\pi$
$\Gamma_5 f_1(1285)\pi$
$\Gamma_6 \omega\pi^-\pi^0$

 $\Gamma(a_2(1320)\eta)/\Gamma(f_1(1285)\pi)$ Γ_3/Γ_5

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

22.7 ± 7.3	69k	KUHN	04	B852	-	$18 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
------------	-----	------	----	------	---	--

 $\Gamma(f_0(1500)\pi)/\Gamma(a_0(980)\eta)$ Γ_4/Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.28^{+0.20}_{-0.15}$	3 ANISOVICH	01B	SPEC	0	$0.6-1.94 \bar{p}p \rightarrow \eta\eta\pi^0\pi^0$
------------------------	-------------	-----	------	---	--

³ Systematic errors not estimated.

 $\pi_2(1880)$ REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>	

 $\rho(1900)$

$$\rho(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

See our mini-review under the $\rho(1700)$.

 $\rho(1900)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1909 \pm 17 \pm 25$	54	¹ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
1880 ± 30		AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
1860 ± 20		AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
1910 ± 10	^{2,3}	FRABETTI	04 E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
1870 ± 10		ANTONELLI	96 SPEC	$e^+e^- \rightarrow \text{hadrons}$

¹ From the fit with two resonances.

² From a fit with two resonances with the JACOB 72 continuum.

³ Supersedes FRABETTI 01.

DESIG=1
DESIG=2
DESIG=3
DESIG=4
DESIG=5
DESIG=6

NODE=M185R01
NODE=M185R01

NODE=M185R02
NODE=M185R02

NODE=M185R02;LINKAGE=NS

NODE=M185

REFID=59471
REFID=52160
REFID=50459
REFID=49773
REFID=48318

NODE=M170

NODE=M170

NODE=M170M

NODE=M170M

OCCUR=2

NODE=M170M;LINKAGE=AU
NODE=M170M;LINKAGE=PI
NODE=M170M;LINKAGE=RS

NODE=M170W

NODE=M170W

OCCUR=2

NODE=M170W;LINKAGE=AU
NODE=M170W;LINKAGE=PI
NODE=M170W;LINKAGE=RS

 $\rho(1900)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$48 \pm 17 \pm 2$	54	⁴ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
130 ± 30		AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
160 ± 20		AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
37 ± 13	^{5,6}	FRABETTI	04 E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
10 ± 5		ANTONELLI	96 SPEC	$e^+e^- \rightarrow \text{hadrons}$

⁴ From the fit with two resonances.

⁵ From a fit with two resonances with the JACOB 72 continuum.

⁶ Supersedes FRABETTI 01.

$\rho(1900) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(\phi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$		$\Gamma_4/\Gamma \times \Gamma_6/\Gamma$		
VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$4.2 \pm 1.2 \pm 0.8$	54	⁷ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
⁷ From the fit with two resonances.				

 $\rho(1900)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 6π	seen
Γ_2 $3\pi^+3\pi^-$	seen
Γ_3 $2\pi^+2\pi^-2\pi^0$	
Γ_4 $\phi\pi$	
Γ_5 hadrons	seen
Γ_6 e^+e^-	seen
Γ_7 NN	not seen

 $\rho(1900)$ BRANCHING RATIOS

$\Gamma(6\pi)/\Gamma_{\text{total}}$		Γ_1/Γ		
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	8k	AKHMETSHIN 13	CMD3	$e^+e^- \rightarrow 3\pi^+3\pi^-$
not seen		AGNELLO 02	OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-\pi^0$
seen		FRABETTI 01	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
seen		ANTONELLI 96	SPEC	$e^+e^- \rightarrow \text{hadrons}$

 $\rho(1900)$ REFERENCES

AKHMETSHIN 13	PL B723 82	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
AUBERT 085	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
FRABETTI 04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AGNELLO 02	PL B527 39	M. Agnello <i>et al.</i>	(OBELIX Collab.)
FRABETTI 01	PL B514 240	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANTONELLI 96	PL B365 427	A. Antonelli <i>et al.</i>	(FENICE Collab.)
JACOB 72	PR D5 1847	M. Jacob, R. Slansky	

 $f_2(1910)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

We list here three different peaks with close masses and widths seen in the mass distributions of $\omega\omega$, $\eta\eta'$, and K^+K^- final states. ALDE 91B argues that they are of different nature.

 $f_2(1910)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 \pm 9 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.			
1890 \pm 10	¹ AMELIN 06	VES	$36\pi^-p \rightarrow \omega\omega n$
1897 \pm 11	BARBERIS 00F		$450pp \rightarrow p_f\omega\omega p_s$
1924 \pm 14	ALDE 90	GAM2	$38\pi^-p \rightarrow \omega\omega n$

NODE=M170215

NODE=M170B01

NODE=M170B01

NODE=M170B01;LINKAGE=AU

NODE=M170225;NODE=M170

DESIG=5;OUR EST; \rightarrow UNCHECKED
 DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=6
 DESIG=7
 DESIG=2;OUR EST; \rightarrow UNCHECKED
 DESIG=3;OUR EST; \rightarrow UNCHECKED
 DESIG=4;OUR EST; \rightarrow UNCHECKED

NODE=M170230

NODE=M170R1
 NODE=M170R1

NODE=M170

REFID=55370
 REFID=52242
 REFID=51047
 REFID=49614
 REFID=48576
 REFID=48350
 REFID=44633
 REFID=49668

NODE=M142

NODE=M142


NODE=M142205

NODE=M142MX

NODE=M142M2
 NODE=M142M2

¹ Supersedes BELADIDZE 92B.

NODE=M142M2;LINKAGE=AM

 **$f_2(1910) \eta\eta'$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1934±16	1 BARBERIS	00A	$450 \bar{p}p \rightarrow p_f \eta\eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1934±20	2 ANISOVICH	00J	SPEC
1911±10	ALDE	91B	$GAM2 \ 38 \pi^- p \rightarrow \eta\eta' n$

¹ Also compatible with $J^{PC}=1-+$.
² Combined fit with $\eta\eta$, $\pi\pi$, and $\eta\pi\pi$.

NODE=M142M3
NODE=M142M3 **$f_2(1910) K^+ K^-$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1941±18	1 AMSLER	06	$CBAR \ 1.64 \bar{p}p \rightarrow K^+ K^- \pi^0$

¹ Tentative, could be $f_2(1950)$.

NODE=M142M3;LINKAGE=KS

NODE=M142M3;LINKAGE=AN

NODE=M142M4
NODE=M142M4

NODE=M142M4;LINKAGE=A

NODE=M142210

NODE=M142WX


NODE=M142W2
NODE=M142W2

NODE=M142W2;LINKAGE=AM

 $f_2(1910) \omega\omega$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
167±21 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
165±19	1 AMELIN	06	$36 \pi^- p \rightarrow \omega\omega n$
202±32	BARBERIS	00F	$450 \bar{p}p \rightarrow p_f \omega\omega p_s$

¹ Supersedes BELADIDZE 92B.



f₂(1910) $\eta\eta'$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
141±41	1 BARBERIS	00A	450 $\bar{p}p \rightarrow p_f \eta\eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
271±25	2 ANISOVICH	00J	SPEC
90±35	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

1 Also compatible with $JPC=1-+$.
2 Combined fit with $\eta\eta$, $\pi\pi$, and $\eta\pi\pi$.

NODE=M142W3
NODE=M142W3

f₂(1910) $K^+ K^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
120±40	AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+ K^- \pi^0$

NODE=M142W3;LINKAGE=KS
NODE=M142W3;LINKAGE=AN

NODE=M142W4
NODE=M142W4

NODE=M142215;NODE=M142

f₂(1910) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^0 \pi^0$	
$\Gamma_2 K^+ K^-$	seen
$\Gamma_3 K_S^0 K_S^0$	
$\Gamma_4 \eta\eta$	seen
$\Gamma_5 \omega\omega$	seen
$\Gamma_6 \eta\eta'$	seen
$\Gamma_7 \eta'\eta'$	
$\Gamma_8 \rho\rho$	seen
$\Gamma_9 a_2(1320)\pi$	seen
$\Gamma_{10} f_2(1270)\eta$	seen

DESIG=6
DESIG=11
DESIG=8
DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=9
DESIG=10;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=12;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=13;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M142225

f₂(1910) BRANCHING RATIOS

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	Γ_2/Γ
seen	1 AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+ K^- \pi^0$

NODE=M142R11
NODE=M142R11

NODE=M142R11;LINKAGE=A

1 Tentative, could be $f_2(1950)$.

$\Gamma(\pi^0 \pi^0)/\Gamma(\eta\eta')$	DOCUMENT ID	TECN	Γ_1/Γ_6
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.1	ALDE	89	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

NODE=M142R4
NODE=M142R4

$\Gamma(K_S^0 K_S^0)/\Gamma(\eta\eta')$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_3/Γ_6
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.066	90	BALOSHIN	86	SPEC $40\pi p \rightarrow K_S^0 K_S^0 n$	NODE=M142R7 NODE=M142R7

 $\Gamma(\eta\eta)/\Gamma(\eta\eta')$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_4/Γ_6
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<0.05	90	ALDE	91B GAM2	$38\pi^- p \rightarrow \eta\eta' n$	NODE=M142R6 NODE=M142R6

 $\Gamma(\omega\omega)/\Gamma(\eta\eta')$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ_6
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
2.6 ± 0.6	BARBERIS	00F	$450 pp \rightarrow p_f \omega\omega p_s$	NODE=M142R10 NODE=M142R10

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
probably not seen	BARBERIS	00A	$450 pp \rightarrow p_f \eta' \eta' p_s$	NODE=M142R8 NODE=M142R8
possibly seen	BELADIDZE	92D VES	$37\pi^- p \rightarrow \eta' \eta' n$	

 $\Gamma(\rho\rho)/\Gamma(\omega\omega)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_8/Γ_5
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
2.6 ± 0.4	BARBERIS	00F	$450 pp \rightarrow p_f \omega\omega p_s$	NODE=M142R9 NODE=M142R9

 $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{10}/Γ_9
0.09 ± 0.05	¹ ANISOVICH	11	SPEC $0.9\text{--}1.94 p\bar{p}$	NODE=M142R12 NODE=M142R12

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

f₂(1910) REFERENCES

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 69 715.		
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(CBAR Collab.)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(Crystal Barrel Collab.)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(VES Collab.)
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)
BELADIDZE	92D	ZPHY C57 13	G.M. Beladidze <i>et al.</i>	(VES Collab.)
ALDE	91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
		Translated from YAF 54 751.		
Also		PL B276 375	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	89	PL B216 447	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
Also		SJNP 48 1035	D.M. Alde <i>et al.</i>	(BELG, SERP, LANL, LAPP)
		Translated from YAF 48 1724.		
BALOSHIN	86	SJNP 43 959	O.N. Baloshin <i>et al.</i>	(ITEP)
		Translated from YAF 43 1487.		

REFID=41911
REFID=40935
REFID=40727
REFID=44697

REFID=40734

REFID=53631
REFID=51574

REFID=51136
REFID=47945
REFID=47950
REFID=47957
REFID=47962
REFID=45202
REFID=42172
REFID=43309
REFID=41844

REFID=40727
REFID=44697

REFID=40734

REFID=40734

REFID=40734

NODE=M227

 $a_0(1950)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. Seen in $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$ by LEES 16A
with significance 2.5σ in $K_S^0 K^\pm \pi^\mp$ and 4.2σ in $K^+ K^- \pi^0$.

NODE=M227

 $a_0(1950)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$1931 \pm 14 \pm 22$	12k	1,2 LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1949 $\pm 32 \pm 76$	8k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
1927 $\pm 15 \pm 23$	4k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$

¹ From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating width.

² Weighted average of the $K_S^0 K^\pm$ and $K^+ K^-$ decay modes.

NODE=M227M

NODE=M227M

OCCUR=3

OCCUR=2

NODE=M227M;LINKAGE=A

NODE=M227M;LINKAGE=B

NODE=M227W

NODE=M227W

OCCUR=3

OCCUR=2

NODE=M227W;LINKAGE=A

NODE=M227W;LINKAGE=B

NODE=M227215;NODE=M227

 $a_0(1950)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$271 \pm 22 \pm 29$	12k	1,2 LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

265 $\pm 36 \pm 110$	8k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
274 $\pm 28 \pm 30$	4k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$

¹ From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating mass.

² Weighted average of the $K_S^0 K^\pm$ and $K^+ K^-$ decay modes.

DESIG=1

NODE=M227225

NODE=M227R01

NODE=M227R01

NODE=M227R01;LINKAGE=A

NODE=M227

REFID=57125

 $a_0(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}$	seen

 $a_0(1950)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_1/Γ
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

¹ From a model-independent partial wave analysis.

NODE=M227R01

NODE=M227R01

NODE=M227R01

NODE=M227R01;LINKAGE=A

NODE=M227

REFID=57125

 $a_0(1950)$ REFERENCES

LEES	16A PR D93 012005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
------	-------------------	-------------------------	-----------------

$f_2(1950)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$f_2(1950)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1936±12 OUR AVERAGE			Error includes scale factor of 1.3. See the ideogram below.
1940±50	BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
1980±22	1 BARBERIS	00C	450 $p p \rightarrow p p 4\pi$
1940±22	2 BARBERIS	00C	450 $p p \rightarrow p p 2\pi 2\pi^0$
1960±30	BARBERIS	97B OMEG	450 $p p \rightarrow p p 2(\pi^+ \pi^-)$
1918±12	ANTINORI	95 OMEG	300,450 $p p \rightarrow p p 2(\pi^+ \pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2038 ⁺¹³⁺¹² ₋₁₁₋₇₃	3 UEHARA	09 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1930±25	4 BINON	05 GAMS	$33 \pi^- p \rightarrow \eta \eta n$
1980± 2±14	ABE	04 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1867±46	5 AMSLER	02 CBAR	$0.9 \bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
2010±25	ANISOVICH	00J SPEC	
1980±50	ANISOVICH	99B SPEC	$1.35-1.94 p \bar{p} \rightarrow \eta \eta \pi^0$
~1990	6 OAKDEN	94 RVUE	$0.36-1.55 \bar{p} p \rightarrow \pi \pi$
1950±15	7 ASTON	91 LASS	$11 K^- p \rightarrow \Lambda K \bar{K} \pi \pi$

1 Decaying into $\pi^+ \pi^- \pi^0$.2 Decaying into $2(\pi^+ \pi^-)$.3 Taking into account $f_4(2050)$.

4 First solution, PWA is ambiguous.

5 T-matrix pole.

6 From solution B of amplitude analysis of data on $\bar{p} p \rightarrow \pi \pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

7 Cannot determine spin to be 2.

NODE=M135M

NODE=M135M

OCCUR=2

NODE=M135M;LINKAGE=A4

NODE=M135M;LINKAGE=B4


NODE=M135M;LINKAGE=UE

NODE=M135M;LINKAGE=BI

NODE=M135M;LINKAGE=TT

NODE=M135M;LINKAGE=BB

NODE=M135M;LINKAGE=A

 **$f_2(1950)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
464± 24 OUR AVERAGE			
380 ⁺¹²⁰ ₋₉₀	BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
520± 50	8 BARBERIS	00C	450 $p p \rightarrow p p 4\pi$
485± 55	9 BARBERIS	00C	450 $p p \rightarrow p p 4\pi$
460± 40	BARBERIS	97B OMEG	450 $p p \rightarrow p p 2(\pi^+ \pi^-)$
390± 60	ANTINORI	95 OMEG	300,450 $p p \rightarrow p p 2(\pi^+ \pi^-)$

NODE=M135W

NODE=M135W

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$441^{+27+28}_{-25-192}$	10 UEHARA	09 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
450 ± 50	11 BINON	05 GAMS	$33 \pi^- p \rightarrow \eta \eta n$
$297 \pm 12 \pm 6$	ABE	04 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
385 ± 58	12 AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
495 ± 35	ANISOVICH	00J SPEC	
500 ± 100	ANISOVICH	99B SPEC	$1.35-1.94 p\bar{p} \rightarrow \eta \eta \pi^0$
~ 100	13 OAKDEN	94 RVUE	$0.36-1.55 \bar{p}p \rightarrow \pi\pi$
250 ± 50	14 ASTON	91 LASS	$11 K^- p \rightarrow \Lambda K\bar{K} \pi\pi$

8 Decaying into $\pi^+ \pi^- 2\pi^0$.

9 Decaying into $2(\pi^+ \pi^-)$.

10 Taking into account $f_4(2050)$.

11 First solution, PWA is ambiguous.

12 T-matrix pole.

13 From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

14 Cannot determine spin to be 2.

NODE=M135W;LINKAGE=A4

NODE=M135W;LINKAGE=B4

NODE=M135W;LINKAGE=UE

NODE=M135W;LINKAGE=BI

NODE=M135W;LINKAGE=TT

NODE=M135W;LINKAGE=BB

NODE=M135W;LINKAGE=A

NODE=M135215;NODE=M135

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892) \bar{K}^*(892)$	seen
$\Gamma_2 \pi\pi$	
$\Gamma_3 \pi^+ \pi^-$	seen
$\Gamma_4 \pi^0 \pi^0$	seen
$\Gamma_5 4\pi$	seen
$\Gamma_6 \pi^+ \pi^- \pi^+ \pi^-$	
$\Gamma_7 a_2(1320)\pi$	
$\Gamma_8 f_2(1270)\pi\pi$	
$\Gamma_9 \eta\eta$	seen
$\Gamma_{10} K\bar{K}$	seen
$\Gamma_{11} \gamma\gamma$	seen
$\Gamma_{12} p\bar{p}$	seen

$f_2(1950) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$122 \pm 4 \pm 26$ 15 ABE 04 BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$

15 Assuming spin 2.

$\Gamma_{10}\Gamma_{11}/\Gamma$

NODE=M135225

NODE=M135G1

NODE=M135G1

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$162^{+69+1137}_{-42-204}$ 16 UEHARA 09 BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$

16 Taking into account $f_4(2050)$.

$\Gamma_2\Gamma_{11}/\Gamma$

NODE=M135G1;LINKAGE=AB

NODE=M135G2

NODE=M135G2

$f_2(1950) \text{BRANCHING RATIOS}$

$\Gamma(K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

seen ASTON 91 LASS 0 $11 K^- p \rightarrow \Lambda K\bar{K} \pi\pi$

Γ_1/Γ

NODE=M135G2;LINKAGE=UE

NODE=M135220

NODE=M135R1

NODE=M135R1

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen BARBERIS 00B $450 p\bar{p} \rightarrow p_f \eta \pi^+ \pi^- p_s$

not seen BARBERIS 00C $450 p\bar{p} \rightarrow p_f 4\pi p_s$

possibly seen BARBERIS 97B OMEG $450 p\bar{p} \rightarrow p p 2(\pi^+ \pi^-)$

Γ_7/Γ

NODE=M135R3

NODE=M135R3

$\Gamma(\eta\eta)/\Gamma(4\pi)$				Γ_9/Γ_5
VALUE	CL%	DOCUMENT ID	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<5.0 \times 10^{-3}$	90	BARBERIS 00E	$450 \text{ pp} \rightarrow p_f \eta\eta p_s$	
$\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$				
VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_3
0.14±0.05	AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta\eta, \pi^0 \pi^0 \pi^0$	NODE=M135R6 NODE=M135R6
$\Gamma(p\bar{p})/\Gamma_{\text{total}}$				
VALUE	EVTS	DOCUMENT ID	TECN	Γ_{12}/Γ
seen	111	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

$f_2(1950)$ REFERENCES

ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)

$a_4(1970)$

$I^G(J^{PC}) = 1^-(4^{++})$

was $a_4(2040)$

$a_4(1970)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1967±16 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.					
1935 ⁺¹¹ ₋₁₃	46M	¹ AGHASYAN	18B	COMP	$190 \pi^- p \rightarrow \pi^- \pi^- \pi^- \pi^- p$
1900 ⁺⁸⁰ ₋₂₀		ADOLPH	15	COMP	$191 \pi^- p \rightarrow \eta(\prime) \pi^- p$
1985±10±13	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
1996±25±43		CHUNG	02	B852	$18.3 \pi^- p \rightarrow 3\pi p$
2000±40 ⁺⁶⁰ ₋₂₀		IVANOV	01	B852	$18 \pi^- p \rightarrow \eta' \pi^- p$
2010±20	² DONSKOV	96	GAM2 0		$38 \pi^- p \rightarrow \eta \pi^0 n$
2040±30	³ CLELAND	82B	SPEC	±	$50 \pi^- p \rightarrow K_S^0 K^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1885±13 ⁺⁵⁰ ₋₂	420k	⁴ ALEKSEEV	10	COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
2004± 6	80k	⁵ UMAN	06	E835	$5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
2005 ⁺²⁵ ₋₄₅		⁶ ANISOVICH	01F	SPEC	$2.0 \bar{p}p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
1944± 8±50		⁷ AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1903±10		⁸ BALDI	78	SPEC	—
2030±50		⁹ CORDEN	78C	OMEG 0	$10 \pi^- p \rightarrow p K_S^0 K^-$
					$15 \pi^- p \rightarrow 3\pi n$

1 Statistical error negligible.

2 From a simultaneous fit to the G_+ and G_0 wave intensities.

3 From an amplitude analysis.

4 Superseded by AGHASYAN 2018B.

5 Statistical error only.

6 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

7 May be a different state.

8 From a fit to the Y_8^0 moment. Limited by phase space.

9 $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.

NODE=M135R5
NODE=M135R5

NODE=M135R6
NODE=M135R6

NODE=M135R07
NODE=M135R07

NODE=M135

REFID=53525
REFID=52761
REFID=50780


REFID=49650
REFID=48580
REFID=47950
REFID=47426
REFID=47958
REFID=47959
REFID=47961
REFID=46886
REFID=45758
REFID=45212
REFID=44437
REFID=45210
REFID=41746

NODE=M017

NODE=M017M

NODE=M017M

NODE=M017M;LINKAGE=B
NODE=M017M;LINKAGE=A
NODE=M017M;LINKAGE=C
NODE=M017M;LINKAGE=D
NODE=M017M;LINKAGE=ST
NODE=M017M;LINKAGE=AN
NODE=M017M;LINKAGE=DM
NODE=M017M;LINKAGE=Y
NODE=M017M;LINKAGE=M

 $a_4(1970)$ MASS (MeV) $a_4(1970)$ WIDTH

NODE=M017W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

324 \pm 15 OUR AVERAGE

333 \pm 16	46M	1 AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
300 \pm 80		ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
231 \pm 30 \pm 46	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
298 \pm 81 \pm 85		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi^- p$
350 \pm 100 \pm 70		IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
370 \pm 80		2 DONSKOV	96	GAM2 0	38 $\pi^- p \rightarrow \eta \pi^0 n$
380 \pm 150		3 CLELAND	82B	SPEC \pm	50 $\pi^- p \rightarrow K_S^0 K^\pm p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

294 \pm 25 \pm 46	420k	4 ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
401 \pm 16	80k	5 UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
180 \pm 30		6 ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
324 \pm 26 \pm 75		7 AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
166 \pm 43		8 BALDI	78	SPEC -	10 $\pi^- p \rightarrow p K_S^0 K^-$
510 \pm 200		9 CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi^- n$

1 Statistical error negligible.

2 From a simultaneous fit to the G_+ and G_0 wave intensities.

3 From an amplitude analysis.

4 Superseded by AGHASYAN 2018B.

5 Statistical error only.

6 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

7 May be a different state.

8 From a fit to the Y_8^0 moment. Limited by phase space.9 $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.

NODE=M017W;LINKAGE=B

NODE=M017W;LINKAGE=A

NODE=M017W;LINKAGE=C

NODE=M017W;LINKAGE=D

NODE=M017W;LINKAGE=ST

NODE=M017W;LINKAGE=AN

NODE=M017W;LINKAGE=DM

NODE=M017W;LINKAGE=Y

NODE=M017W;LINKAGE=M

 $a_4(1970)$ DECAY MODES

NODE=M017215;NODE=M017

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\bar{K}$	seen
$\Gamma_2 \pi^+ \pi^- \pi^0$	seen
$\Gamma_3 \rho\pi$	seen
$\Gamma_4 f_2(1270)\pi$	seen

DESIG=1

DESIG=2

DESIG=5;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=6;OUR EST; \rightarrow UNCHECKED \leftarrow

Γ_5	$\omega\pi^-\pi^0$	seen
Γ_6	$\omega\rho$	seen
Γ_7	$\eta\pi$	seen
Γ_8	$\eta'(958)\pi$	seen

DESIG=7;OUR EST; \rightarrow UNCHECKED \leftarrow
 DESIG=8
 DESIG=3
 DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow

a4(1970) BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ				
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
seen	BALDI	78	SPEC	\pm	$10 \pi^- p \rightarrow K_S^0 K^- p$
$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_2/Γ				
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
seen	CORDEN	78C	OMEG	0	$15 \pi^- p \rightarrow 3\pi n$
$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$	Γ_3/Γ_4				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.7^{+0.9}_{-0.8}$ OUR AVERAGE				Error includes scale factor of 3.7.	
$2.9^{+0.6}_{-0.4}$	46M	1 AGHASYAN	18B	COMP	$190 \pi^- p \rightarrow \pi^-\pi^+\pi^- p$
$1.1 \pm 0.2 \pm 0.2$		CHUNG	02	B852	$18.3 \pi^- p \rightarrow 3\pi p$
1 Statistical error negligible.					
$\Gamma(\eta\pi)/\Gamma_{\text{total}}$	Γ_7/Γ				
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
seen	DONSKOV	96	GAM2	0	$38 \pi^- p \rightarrow \eta\pi^0 n$
$\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$	Γ_8/Γ_7				
VALUE	DOCUMENT ID	TECN	COMMENT		
0.23 ± 0.07	ADOLPH	15	COMP	$191 \pi^- p \rightarrow \eta'(\gamma)\pi^- p$	
$\Gamma(\omega\rho)/\Gamma_{\text{total}}$	Γ_6/Γ				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega\pi^-\pi^0 p$

a4(1970) REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>	
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 62 487.		
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>	
DONSKOV	96	PAN 59 982	S.V. Donskov <i>et al.</i>	(GAMS Collab.) IGPJC
		Translated from YAF 59 1027.		
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
BALDI	78	PL 74B 413	R. Baldi <i>et al.</i>	(GEVA) JP
CORDEN	78C	NP B136 77	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP

NODE=M017220
 NODE=M017R1
 NODE=M017R1
 NODE=M017R2
 NODE=M017R2
 NODE=M017R4
 NODE=M017R4
 NODE=M017R4;LINKAGE=A
 NODE=M017R3
 NODE=M017R3
 NODE=M017R01
 NODE=M017R01
 NODE=M017R5
 NODE=M017R5
 NODE=M017

REFID=59471
 REFID=56385
 REFID=53356
 REFID=51063
 REFID=50459
 REFID=48837
 REFID=48352
 REFID=48317
 REFID=46910
 REFID=46903
 REFID=46902
 REFID=45207
 REFID=21281
 REFID=21783
 REFID=20859

NODE=M167

 $\rho_3(1990)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

OMITTED FROM SUMMARY TABLE

 $\rho_3(1990)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1982±14	1 ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~2007	HASAN 94	RVUE	$\bar{p}p \rightarrow \pi\pi$

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167M

NODE=M167M

NODE=M167M;LINKAGE=AY

 $\rho_3(1990)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
188±24	2 ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~287	HASAN 94	RVUE	$\bar{p}p \rightarrow \pi\pi$

² From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167W

NODE=M167W

NODE=M167W;LINKAGE=AY

 $\rho_3(1990)$ REFERENCES

ANISOVICH 02	PL B542 8	A.V. Anisovich <i>et al.</i>
ANISOVICH 01D	PL B508 6	A.V. Anisovich <i>et al.</i>
ANISOVICH 01E	PL B513 281	A.V. Anisovich <i>et al.</i>
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg (RAL, LOQM, PNPI+) (LOQM)

 $\pi_2(2005)$

$$I^G(J^{PC}) = 1^-(2^{+-})$$

OMITTED FROM SUMMARY TABLE

 $\pi_2(2005)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1963⁺¹⁷₋₂₇ OUR AVERAGE				
1962 ⁺¹⁷ ₋₂₉	46M	1 AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^-\pi^+\pi^- p$
1974±14±83	145k	LU 05 B852	18	$\pi^- p \rightarrow \omega\pi^-\pi^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2005±15		ANISOVICH 01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

¹ Statistical uncertainty negligible.

NODE=M167

REFID=48828
REFID=48327
REFID=48349
REFID=47950
REFID=44103

NODE=M239

NODE=M239M

NODE=M239M

NODE=M239M;LINKAGE=A

NODE=M239W

NODE=M239W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
370^{+ 16}_{- 90} OUR AVERAGE				
371 ^{+ 16} _{- 120}	46M	1 AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^-\pi^+\pi^- p$
341± 61±139	145k	LU 05 B852	18	$\pi^- p \rightarrow \omega\pi^-\pi^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200± 40		ANISOVICH 01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

¹ Statistical uncertainty negligible.

NODE=M239W;LINKAGE=A

$\pi_2(2005)$ DECAY MODES

NODE=M239215;NODE=M239

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^- \pi^+ \pi^-$	seen
$\Gamma_2 \omega \pi^0 \pi^-$	seen

 $\pi_2(2005)$ BRANCHING RATIOS

$\Gamma(\pi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE seen	DOCUMENT ID AGHASYAN 18B TECN COMP COMMENT 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

$\Gamma(\omega \pi^0 \pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE seen	DOCUMENT ID LU 05 TECN B852 COMMENT 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$

 $\pi_2(2005)$ REFERENCES

AGHASYAN LU ANISOVICH	18B 05 01F	PR D98 092003 PRL 94 032002 PL B517 261	M. Aghasyan <i>et al.</i> M. Lu <i>et al.</i> A.V. Anisovich <i>et al.</i>	(COMPASS Collab.) (BNL E852 Collab.)
-----------------------------	------------------	---	--	---

 $f_2(2010)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

 $f_2(2010)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2011^{+ 62}_{- 76}	¹ ETKIN 88 MPS		$22 \pi^- p \rightarrow \phi \phi n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2005 \pm 12	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1980 \pm 20	² BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
2050 $^{+ 90}_{- 50}$	ETKIN 85	MPS	$22 \pi^- p \rightarrow 2\phi n$
2120 $^{+ 20}_{- 120}$	LINDENBAUM 84	RVUE	
2160 \pm 50	ETKIN 82	MPS	$22 \pi^- p \rightarrow 2\phi n$

¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi \phi 2^{++} S_2$, D_2 , and D_0 is 98^{+1}_{-3} , 0^{+1}_{-0} , and 2^{+2}_{-1} , respectively.

² Statistically very weak, only 1.4 s.d.

 $f_2(2010)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
202^{+ 67}_{- 62}	³ ETKIN 88 MPS		$22 \pi^- p \rightarrow \phi \phi n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

209 \pm 32	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
145 \pm 50	⁴ BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
200 $^{+ 160}_{- 50}$	ETKIN 85	MPS	$22 \pi^- p \rightarrow 2\phi n$
300 $^{+ 150}_{- 50}$	LINDENBAUM 84	RVUE	
310 \pm 70	ETKIN 82	MPS	$22 \pi^- p \rightarrow 2\phi n$

³ Includes data of ETKIN 85.

⁴ Statistically very weak, only 1.4 s.d.

 $f_2(2010)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \phi \phi$	seen
$\Gamma_2 K \bar{K}$	seen

DESIG=1

DESIG=2

NODE=M239220

NODE=M239R00
NODE=M239R00NODE=M239R01
NODE=M239R01

NODE=M239

REFID=59471
REFID=50459
REFID=48352

NODE=M106

NODE=M106M

NODE=M106M

NODE=M106M;LINKAGE=C

NODE=M106M;LINKAGE=E

NODE=M106W

NODE=M106W

NODE=M106W;LINKAGE=C

NODE=M106W;LINKAGE=E

NODE=M106215;NODE=M106

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2

$f_2(2010)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$		DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	

 $f_2(2010)$ REFERENCES

VLADIMIRSK... 06	PAN 69 493 Translated from YAF 69 515.	V.V. Vladimirsy <i>et al.</i>	(ITEP, Moscow)
BOLONKIN 88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ETKIN 88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)
ETKIN 85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)
LINDENBAUM 84	CNPP 13 285	S.J. Lindenbaum	(CUNY)
ETKIN 82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)
Also	Brighton Conf. 351	S.J. Lindenbaum	(BNL, CUNY)

 $f_0(2020)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

 $f_0(2020)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1992±16	1,2	BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1910±50	3 ROPERTZ	18 RVUE	$\bar{B}_s^0 \rightarrow J/\psi(\pi^+\pi^-/K^+K^-)$	1
2037± 8	80k 4 UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$	
2040±38	ANISOVICH	00J SPEC		
2010±60	ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0\pi^0 n$	
2020±35	BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$	
1 Average between $\pi^+\pi^-2\pi^0$ and $2(\pi^+\pi^-)$. 2 T-matrix pole. 3 T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants. 4 Statistical error only.				

NODE=M106

REFID=51191

REFID=40580

REFID=40285

REFID=21871

REFID=21869

REFID=21866

REFID=21867

NODE=M156

NODE=M156

NODE=M156M

NODE=M156M

NODE=M156M;LINKAGE=PC

NODE=M156M;LINKAGE=PP

NODE=M156M;LINKAGE=A

NODE=M156M;LINKAGE=ST

NODE=M156W

NODE=M156W

NODE=M156W;LINKAGE=PC

NODE=M156W;LINKAGE=PP

NODE=M156W;LINKAGE=A

NODE=M156W;LINKAGE=ST

NODE=M156215;NODE=M156

 $f_0(2020)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
442± 60	1,2 BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_s$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
400± 80	3 ROPERTZ	18 RVUE	$\bar{B}_s^0 \rightarrow J/\psi(\pi^+\pi^-/K^+K^-)$	1
296± 17	80k 4 UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$	
405± 40	ANISOVICH	00J SPEC		
240±100	ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0\pi^0 n$	
410± 50	BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+\pi^-)$	
1 Average between $\pi^+\pi^-2\pi^0$ and $2(\pi^+\pi^-)$. 2 T-matrix pole. 3 T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants. 4 Statistical error only.				

 $f_0(2020)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\pi\pi$	seen
$\Gamma_2 \pi^0\pi^0$	seen
$\Gamma_3 \rho\rho$	seen
$\Gamma_4 \omega\omega$	seen
$\Gamma_5 \eta\eta$	seen

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=5

$f_0(2020)$ BRANCHING RATIOS **$\Gamma(\rho\rho)/\Gamma(\omega\omega)$**

VALUE	DOCUMENT ID	COMMENT	Γ_3/Γ_4
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 3	BARBERIS 00F	450 $p p \rightarrow p_f \omega\omega p_s$	

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
seen	UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$	

 $f_0(2020)$ REFERENCES

ROPERTZ 18	EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis (BONN, JULI)
AAIJ 17V	JHEP 1708 037	R. Aaij <i>et al.</i> (LHCb Collab.)
AAIJ 14BR	PR D89 092006	R. Aaij <i>et al.</i> (LHCb Collab.)
UMAN 06	PR D73 052009	I. Uman <i>et al.</i> (FNAL E835)
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i> (RAL, LOQM, PNPI+)
BARBERIS 00C	PL B471 440	D. Barberis <i>et al.</i> (WA 102 Collab.)
BARBERIS 00F	PL B484 198	D. Barberis <i>et al.</i> (WA 102 Collab.)
ALDE 98	EPJ A3 361	D. Alde <i>et al.</i> (GAM4 Collab.)
Also	PAN 62 405	D. Alde <i>et al.</i> (GAMS Collab.)
	Translated from YAF 62 446.	
BARBERIS 97B	PL B413 217	D. Barberis <i>et al.</i> (WA 102 Collab.)

 $f_4(2050)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

 $f_4(2050)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2018±11 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.	
1960±15		AMELIN	06 VES	$36 \pi^- p \rightarrow \omega\omega n$	
2005±10		¹ BINON	05 GAMS	$33 \pi^- p \rightarrow \eta\eta n$	
1998±15		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$	
2060±20		ALDE	90 GAM2	$38 \pi^- p \rightarrow \omega\omega n$	
2038±30		AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
2086±15		BALTRUSAIT..	87 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
2000±60		ALDE	86D GAM4	$100 \pi^- p \rightarrow n2\eta$	
2020±20	40k	² BINON	84B GAM2	$38 \pi^- p \rightarrow n2\pi^0$	
2015±28		³ CASON	82 STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$	
2031^{+25}_{-36}		ETKIN	82B MPS	$23 \pi^- p \rightarrow n2K_S^0$	
2020±30	700	APEL	75 NICE	$40 \pi^- p \rightarrow n2\pi^0$	
2050±25		BLUM	75 ASPK	$18.4 \pi^- p \rightarrow nK^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1966±25		⁴ ANISOVICH	09 RVUE	$0.0 \bar{p}p, \pi N$	
$1885^{+14}_{-13}{}^{+218}_{-25}$		⁵ UEHARA	09 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	
2018± 6		ANISOVICH	00J SPEC	$2.0 \bar{p}p \rightarrow \eta\pi^0 \pi^0, \pi^0 \pi^0,$ $\eta\eta, \eta\eta', \pi\pi$	
~ 2000		⁶ MARTIN	98 RVUE	$\bar{N}\bar{N} \rightarrow \pi\pi$	
~ 2010		⁷ MARTIN	97 RVUE	$\bar{N}\bar{N} \rightarrow \pi\pi$	
~ 2040		⁸ OAKDEN	94 RVUE	$0.36\text{--}1.55 \bar{p}p \rightarrow \pi\pi$	
~ 1990		⁹ OAKDEN	94 RVUE	$0.36\text{--}1.55 \bar{p}p \rightarrow \pi\pi$	
1978± 5		¹⁰ ALPER	80 CNTR	$62 \pi^- p \rightarrow K^+ K^- n$	
2040±10		¹⁰ ROZANSKA	80 SPRK	$18 \pi^- p \rightarrow p\bar{p}n$	
1935±13		¹⁰ CORDEN	79 OMEG	$12\text{--}15 \pi^- p \rightarrow n2\pi$	
1988± 7		EVANGELIS...	79B OMEG	$10 \pi^- p \rightarrow K^+ K^- n$	
1922±14		¹¹ ANTIPOV	77 CIBS	$25 \pi^- p \rightarrow p3\pi$	

¹ From the first PWA solution.

² From a partial-wave analysis of the data.

³ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.

⁴ K matrix pole.

⁵ Taking into account the $f_2(1950)$. Helicity-2 production favored.

⁶ Energy-dependent analysis.

⁷ Single energy analysis.

NODE=M156220

NODE=M156R1

NODE=M156R01
NODE=M156R01

NODE=M156

REFID=59332
REFID=57828
REFID=56035
REFID=51063
REFID=47950
REFID=47959
REFID=47962
REFID=46605
REFID=46914
REFID=45758

NODE=M016


NODE=M016M

NODE=M016M

OCCUR=2

NODE=M016M;LINKAGE=BI
NODE=M016M;LINKAGE=N
NODE=M016M;LINKAGE=NN
NODE=M016M;LINKAGE=KM
NODE=M016M;LINKAGE=UE
NODE=M016M;LINKAGE=RB
NODE=M016M;LINKAGE=BR

- 8 From solution A of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
- 9 From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
- 10 $I(JP) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.
- 11 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.



f4(2050) WIDTH				
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
237± 18 OUR AVERAGE				Error includes scale factor of 1.9. See the ideogram below.
290± 20		AMELIN	06	$36 \pi^- p \rightarrow \omega\omega n$
340± 80		12 BINON	05	$33 \pi^- p \rightarrow \eta\eta n$
395± 40		ALDE	98	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
170± 60		ALDE	90	$38 \pi^- p \rightarrow \omega\omega n$
304± 60		AUGUSTIN	87	$J/\psi \rightarrow \gamma\pi^+\pi^-$
210± 63		BALTRUSAIT...	87	$J/\psi \rightarrow \gamma\pi^+\pi^-$
400±100		ALDE	86D	$100 \pi^- p \rightarrow n2\eta$
240± 40	40k	13 BINON	84B	$38 \pi^- p \rightarrow n2\pi^0$
190± 14		DENNEY	83	$10 \pi^+ n/\pi^+ p$
186 ⁺¹⁰³ ₋₅₈		14 CASON	82	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
305 ⁺³⁶ ₋₁₁₉		ETKIN	82B	$23 \pi^- p \rightarrow n2K_S^0$
180± 60	700	APEL	75	$40 \pi^- p \rightarrow n2\pi^0$
225 ⁺¹²⁰ ₋₇₀		BLUM	75	$18.4 \pi^- p \rightarrow nK^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
260± 40		15 ANISOVICH	09	$0.0 \bar{p}p, \pi N$
453± 20 ⁺³¹ ₋₁₂₉		16 UEHARA	09	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
182± 7		ANISOVICH	00J	$2.0 \bar{p}p \rightarrow \eta\pi^0 \pi^0, \pi^0 \pi^0,$ $\eta\eta, \eta\eta', \pi\pi$
~ 170		17 MARTIN	98	$NN \rightarrow \pi\pi$
~ 200		18 MARTIN	97	$\bar{N}N \rightarrow \pi\pi$
~ 60		19 OAKDEN	94	$RVUE 0.36-1.55 \bar{p}p \rightarrow \pi\pi$
~ 80		20 OAKDEN	94	$RVUE 0.36-1.55 \bar{p}p \rightarrow \pi\pi$
243± 16		21 ALPER	80	$CNTR 62 \pi^- p \rightarrow K^+ K^- n$
140± 15		21 ROZANSKA	80	$SPRK 18 \pi^- p \rightarrow p\bar{p}n$
263± 57		21 CORDEN	79	$OMEG 12-15 \pi^- p \rightarrow n2\pi$
100± 28		EVANGELIS...	79B	$OMEG 10 \pi^- p \rightarrow K^+ K^- n$
107± 56		22 ANTIPOV	77	$CIBS 25 \pi^- p \rightarrow p3\pi$

NODE=M016M;LINKAGE=B

NODE=M016M;LINKAGE=BB

NODE=M016M;LINKAGE=M


NODE=M016M;LINKAGE=T

NODE=M016W

NODE=M016W

OCCUR=2

- 12 From the first PWA solution.
 13 From a partial-wave analysis of the data.
 14 From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.
 15 K matrix pole.
 16 Taking into account the $f_2(1950)$. Helicity-2 production favored.
 17 Energy-dependent analysis.
 18 Single energy analysis.
 19 From solution A of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
 20 From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
 21 $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.
 22 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.



f₄(2050) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \omega\omega$	seen
$\Gamma_2 \pi\pi$	$(17.0 \pm 1.5)\%$
$\Gamma_3 K\bar{K}$	$(6.8^{+3.4}_{-1.8}) \times 10^{-3}$
$\Gamma_4 \eta\eta$	$(2.1 \pm 0.8) \times 10^{-3}$
$\Gamma_5 4\pi^0$	< 1.2 %
$\Gamma_6 \gamma\gamma$	
$\Gamma_7 a_2(1320)\pi$	seen

f₄(2050) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_6/\Gamma$
VALUE (keV)	CL%
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
<0.29	95 ALTHOFF 85B TASS $\gamma\gamma \rightarrow K\bar{K}\pi$

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_6/\Gamma$
VALUE (eV)	CL% EVTS DOCUMENT ID TECHN COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	

$23.1^{+3.6+70.5}_{-3.3-15.6}$	23 UEHARA 09 BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
<1100	95 13 ± 4 OEST 90 JADE $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$\bullet \bullet \bullet$ Taking into account the $f_2(1950)$. Helicity-2 production favored.	

NODE=M016W;LINKAGE=BI
 NODE=M016W;LINKAGE=N
 NODE=M016W;LINKAGE=NN
 NODE=M016W;LINKAGE=KM
 NODE=M016W;LINKAGE=UE
 NODE=M016W;LINKAGE=RB
 NODE=M016W;LINKAGE=BR
 NODE=M016W;LINKAGE=BW

NODE=M016W;LINKAGE=BB

NODE=M016W;LINKAGE=M
 NODE=M016W;LINKAGE=T

NODE=M016215;NODE=M016

DESIG=6
 DESIG=1
 DESIG=2
 DESIG=3
 DESIG=5
 DESIG=4
 DESIG=7

NODE=M016220

NODE=M016G2
 NODE=M016G2

NODE=M016G3
 NODE=M016G3

NODE=M016G3;LINKAGE=UE

$f_4(2050)$ BRANCHING RATIOS **$\Gamma(\omega\omega)/\Gamma_{\text{total}}$** VALUE**seen**

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

DOCUMENT IDTECNCOMMENT

AMELIN

06

VES

36 $\pi^- p \rightarrow \omega\omega n$ **Γ_1/Γ**

NODE=M016225

NODE=M016R7

NODE=M016R7

 $\Gamma(\omega\omega)/\Gamma(\pi\pi)$ VALUE **1.5 ± 0.3** DOCUMENT IDTECNCOMMENT

ALDE

90

GAM2

38 $\pi^- p \rightarrow \omega\omega n$ **Γ_1/Γ_2**

NODE=M016R5

NODE=M016R5

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ VALUE **0.170 ± 0.015 OUR AVERAGE**0.18 ± 0.03 0.16 ± 0.03 0.17 ± 0.02 DOCUMENT IDTECNCOMMENT

BINON

83C

GAM2

38 $\pi^- p \rightarrow n4\gamma$

CASON

82

STRC

8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$

CORDEN

79

OMEG

12–15 $\pi^- p \rightarrow n2\pi$ **Γ_2/Γ**

NODE=M016R1

NODE=M016R1

24 Assuming one pion exchange.

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$ VALUE **$0.04^{+0.02}_{-0.01}$** DOCUMENT IDTECNCOMMENT

ETKIN

82B

MPS

23 $\pi^- p \rightarrow n2K_S^0$ **Γ_3/Γ_2**

NODE=M016R2

NODE=M016R2

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) **2.1 ± 0.8** DOCUMENT IDTECNCOMMENT

ALDE

86D

GAM4

100 $\pi^- p \rightarrow n4\gamma$ **Γ_4/Γ**

NODE=M016R3

NODE=M016R3

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ VALUE**<0.012**DOCUMENT IDTECNCOMMENT

ALDE

87

GAM4

100 $\pi^- p \rightarrow 4\pi^0 n$ **Γ_5/Γ**

NODE=M016R4

NODE=M016R4

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ VALUE**seen**DOCUMENT IDTECNCOMMENT

AMELIN

00

VES

37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$ **Γ_7/Γ**

NODE=M016R6

NODE=M016R6

 $f_4(2050)$ REFERENCES

ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
TRANSLATED FROM YAF 69 715.				
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
ALDE	Also	PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
MARTIN	98	PR C57 3492	B.R. Martin <i>et al.</i>	
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP)
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+)
TRANSLATED FROM YAF 62 446.				
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)
CASON	82	PR C48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+) JP
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP
TRANSLATED FROM YAF 38 1199.				

NODE=M016

REFID=52719

REFID=52761

REFID=51574

REFID=50780

REFID=47432

REFID=47950

REFID=47962

REFID=46605

REFID=46914

REFID=46373

REFID=45685

REFID=45212

REFID=45210

REFID=40935

REFID=41358

REFID=40221

REFID=40268

REFID=40010

REFID=20765

REFID=21349

REFID=21780

REFID=40288

REFID=20754

REFID=20746

REFID=20390

REFID=21665

REFID=21774

REFID=20374

REFID=21967

REFID=20728

REFID=20720

REFID=21651

$\pi_2(2100)$ $I^G(J^{PC}) = 1^-(2^-+)$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

 $\pi_2(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2090± 29 OUR AVERAGE			
2090± 30	1 AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
2100±150	2 DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$

¹ From a fit to $J^{PC} = 2^-+$ $f_2(1270)\pi$, $(\pi\pi)_S\pi$ waves.
² From a two-resonance fit to four 2^-0^+ waves.

 $\pi_2(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
625± 50 OUR AVERAGE Error includes scale factor of 1.2.			
520±100	3 AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
651± 50	4 DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$

³ From a fit to $J^{PC} = 2^-+$ $f_2(1270)\pi$, $(\pi\pi)_S\pi$ waves.
⁴ From a two-resonance fit to four 2^-0^+ waves.

 $\pi_2(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $\rho\pi$	seen
Γ_3 $f_2(1270)\pi$	seen
Γ_4 $(\pi\pi)_S\pi$	seen

 $\pi_2(2100)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(3\pi)$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.19±0.05	5 DAUM	81B CNTR	63,94 $\pi^- p$

$\Gamma(f_2(1270)\pi)/\Gamma(3\pi)$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.36±0.09	5 DAUM	81B CNTR	63,94 $\pi^- p$

$\Gamma((\pi\pi)_S\pi)/\Gamma(3\pi)$	Γ_4/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.45±0.07	5 DAUM	81B CNTR	63,94 $\pi^- p$

D-wave/S-wave RATIO FOR $\pi_2(2100) \rightarrow f_2(1270)\pi$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.39±0.23	5 DAUM	81B CNTR	63,94 $\pi^- p$

5 From a two-resonance fit to four 2^-0^+ waves. **$\pi_2(2100)$ REFERENCES**

AMELIN DAUM	95B 81B	PL B356 595 NP B182 269	D.V. Amelin <i>et al.</i> C. Daum <i>et al.</i>	(SERP, TBIL) (AMST, CERN, CRAC, MPIM+)
----------------	------------	----------------------------	--	---

NODE=M020

NODE=M020

NODE=M020M

NODE=M020M

NODE=M020M;LINKAGE=AX
NODE=M020M;LINKAGE=L

NODE=M020W

NODE=M020W

NODE=M020W;LINKAGE=AX
NODE=M020W;LINKAGE=L

NODE=M020215;NODE=M020

DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M020220

NODE=M020R1
NODE=M020R1NODE=M020R2
NODE=M020R2NODE=M020R3
NODE=M020R3NODE=M020R4
NODE=M020R4

NODE=M020R;LINKAGE=L

NODE=M020

REFID=44433

REFID=20872

NODE=M168

 $f_0(2100)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

 $f_0(2100)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

 2086^{+20}_{-24} OUR AVERAGE

$2081 \pm 13^{+24}_{-36}$	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
2090 ± 30		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2090 \pm 10 \pm 6$	529	^{2,3} DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
$2099 \pm 17 \pm 8$	283	^{2,3} DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
2105 ± 8	80k	⁴ UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
2102 ± 13		⁵ ANISOVICH	00J SPEC	$2.0 \bar{p}p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$ $\eta \eta, \eta \eta', \pi^+ \pi^-$
2105 ± 10		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p}p \rightarrow \eta \eta, \eta \eta'$
~ 2104		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
~ 2122		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi \pi$

1 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

2 Using CLEO-c data but not authored by the CLEO Collaboration.

3 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 209$ MeV.

4 Statistical error only.

5 Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

NODE=M168

NODE=M168M

NODE=M168M

OCCUR=2

NODE=M168M;LINKAGE=A

NODE=M168M;LINKAGE=B

NODE=M168M;LINKAGE=C

NODE=M168M;LINKAGE=ST

NODE=M168M;LINKAGE=AN

NODE=M168W

NODE=M168W

 $f_0(2100)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

 284^{+60}_{-32} OUR AVERAGE

273^{+27+70}_{-24-23}	5.5k	⁶ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
330 ± 100		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
236 ± 14	80k	⁷ UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
211 ± 29		⁸ ANISOVICH	00J SPEC	$2.0 \bar{p}p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$ $\eta \eta, \eta \eta', \pi^+ \pi^-$
200 ± 25		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p}p \rightarrow \eta \eta, \eta \eta'$
~ 203		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
~ 273		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi \pi$

NODE=M168W;LINKAGE=A

NODE=M168W;LINKAGE=ST

NODE=M168W;LINKAGE=AN

NODE=M168

 $f_0(2100)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
ABLIKIM	13N	PR D87 092009	Ablkim M. <i>et al.</i>	(BESIII Collab.)
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>	
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)

REFID=56805

REFID=55387

REFID=51063

REFID=47942

REFID=47950

REFID=47426

REFID=47472

REFID=44438

REFID=44103

$f_2(2150)$ $I^G(J^P)$ = $0^+(2^{++})$

OMITTED FROM SUMMARY TABLE

This entry was previously called T_0 . **$f_2(2150)$ MASS** **$f_2(2150)$ MASS, COMBINED MODES (MeV)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2157±12 OUR AVERAGE Includes data from the datablock that follows this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2170± 6 80k 1 UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$

1 Statistical error only.

 $\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

2157±12 OUR AVERAGE2151±16 BARBERIS 00E 450 $p p \rightarrow p_f \eta\eta p_s$
2175±20 PROKOSHIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$,
450 $p p \rightarrow p p 2\eta$ 2130±35 SINGOVSKI 94 GAM4 450 $p p \rightarrow p p 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2140±30 ² ABELE 99B CBAR 1.94 $\bar{p}p \rightarrow \pi^0 \eta\eta$
2104±20 ³ ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$

2 Spin not determined.

3 No J^P determination. **$\eta\pi\pi$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135±20±45 ⁴ ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$ 4 ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance. **$\bar{p}p \rightarrow \pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~2090 ⁵ OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~2120 ⁶ OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~2170 ⁷ MARTIN 80B RVUE
~2150 ⁷ MARTIN 80C RVUE
~2150 ⁸ DULUDE 78B OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$ 5 OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.6 From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.7 $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.8 $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.**S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2139⁺₋ 8 ⁹ EVANGELIS... 97 SPEC 0.6–2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
~2190 ⁹ CUTTS 78B CNTR 0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155₋⁺¹⁵ COUPLAND 77 CNTR 0 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193₋² ALSPECTOR 73 CNTR $\bar{p}p$ S channel

9 Isospins 0 and 1 not separated.

10 From a fit to the total elastic cross section.

11 Referred to as T or T region by ALSPECTOR 73.

NODE=M042

NODE=M042

NODE=M042205

NODE=M042M

NODE=M042M

NODE=M042M;LINKAGE=ST

NODE=M042M3

NODE=M042M3

NODE=M042M3;LINKAGE=K3
NODE=M042M3;LINKAGE=A

NODE=M042M4

NODE=M042M4

NODE=M042M4;LINKAGE=AD

NODE=M042M1
NODE=M042M1

OCCUR=2

NODE=M042M1;LINKAGE=B

NODE=M042M1;LINKAGE=BB
NODE=M042M1;LINKAGE=P
NODE=M042M1;LINKAGE=L

NODE=M042M2

NODE=M042M2

NODE=M042M2;LINKAGE=I

NODE=M042M2;LINKAGE=E

NODE=M042M2;LINKAGE=M

$K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2200±13	VLADIMIRSK...06	SPEC 40	$\pi^- p \rightarrow K_S^0 K_S^0 n$
2150±20	ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
2130±35	BARBERIS 99	OMEG 450	$p p \rightarrow p_s p_f K^+ K^-$

NODE=M042M5
NODE=M042M5

 $f_2(2150)$ WIDTH **$f_2(2150)$ WIDTH, COMBINED MODES (MeV)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
152±30 OUR AVERAGE Includes data from the datablock that follows this one. Error includes scale factor of 1.4. See the ideogram below.				

• • • We do not use the following data for averages, fits, limits, etc. • • •

182±11	80k	12 UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
--------	-----	---------	---------	--


12 Statistical error only.

NODE=M042210

NODE=M042W
NODE=M042W

NODE=M042W;LINKAGE=ST

WEIGHTED AVERAGE
152±30 (Error scaled by 1.4)

 **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

NODE=M042W3
NODE=M042W3

152±30 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

280±70	BARBERIS 00E	450 $p p \rightarrow p_f \eta\eta p_s$
150±35	PROKOSHKIN 95D	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $\bar{p}p \rightarrow pp 2\eta$
130±30	SINGOVSKI 94	GAM4 450 $p p \rightarrow pp 2\eta$


• • • We do not use the following data for averages, fits, limits, etc. • • •

310±50	13 ABELE 99B	CBAR 1.94 $\bar{p}p \rightarrow \pi^0 \eta\eta$
203±10	14 ARMSTRONG 93C	E760 $\bar{p}p \rightarrow \pi^0 \eta\eta \rightarrow 6\gamma$

13 Spin not determined.

14 No JPC determination.

NODE=M042W3;LINKAGE=K3
NODE=M042W3;LINKAGE=A



$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
250±25±45	15 ADOMEIT	96 CBAR 0	1.94	$\bar{p}p \rightarrow \eta 3\pi^0$
15 ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.				

NODE=M042W4
NODE=M042W4

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
250 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 70	16 OAKDEN	94 RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	17 MARTIN	80B RVUE	
~ 250	17 MARTIN	80C RVUE	
~ 250	18 DULUDE	78B OSPK	1–2 $\bar{p}p \rightarrow \pi^0\pi^0$

NODE=M042W4;LINKAGE=AD

16 See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

17 $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

18 $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

NODE=M042W1
NODE=M042W1
→ UNCHECKED ←

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
56+31 -16	19 EVANGELIS...	97 SPEC	0.6–2.4	$\bar{p}p \rightarrow K_S^0 K_S^0$
135±75	20,21 COUPLAND	77 CNTR 0	0.7–2.4	$\bar{p}p \rightarrow \bar{p}p$
98± 8	21 ALSPECTOR	73 CNTR		$\bar{p}p$ S channel

NODE=M042W1;LINKAGE=CC

NODE=M042W1;LINKAGE=P
NODE=M042W1;LINKAGE=L

NODE=M042W2
NODE=M042W2

19 Isospin 0 and 2 not separated.

20 From a fit to the total elastic cross section.

21 Isospins 0 and 1 not separated.

NODE=M042W2;LINKAGE=F
NODE=M042W2;LINKAGE=E
NODE=M042W2;LINKAGE=I

$\bar{K}K$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
91±62	VLADIMIRSK...	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
150±30	ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
270±50	BARBERIS	99 OMEG	$450 pp \rightarrow p_s p_f K^+ K^-$

NODE=M042W5
NODE=M042W5

f₂(2150) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	
$\Gamma_2 \eta\eta$	seen
$\Gamma_3 K\bar{K}$	seen
$\Gamma_4 f_2(1270)\eta$	seen
$\Gamma_5 a_2(1320)\pi$	seen
$\Gamma_6 p\bar{p}$	seen

f₂(2150) BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\eta\eta)$	Γ_3/Γ_2
1.28±0.23	BARBERIS 00E 450 pp → $p_f \eta\eta p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	95	22 PROKOSHIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 pp → pp2η
------	----	--

22 Using data from ARMSTRONG 89d.

$\Gamma(\pi\pi)/\Gamma(\eta\eta)$	Γ_1/Γ_2
0.33	PROKOSHIN 95D GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 pp → pp2η

23 Derived from a $\pi^0 \pi^0/\eta\eta$ limit.

$\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$	Γ_4/Γ_5
0.79±0.11	ADOMEIT 96 CBAR 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

24 Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$	Γ_6/Γ
seen	ALEXANDER 10 CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

f₂(2150) REFERENCES

ALEXANDER 10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
VLADIMIRSKY 06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.	
ABLIKIM 04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)
ANISOVICH 00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS 00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ABELE 99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BARBERIS 99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)
EVANGELIS... 97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
MARTIN 97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)
ADOMEIT 96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
PROKOSHIN 95D	PD 40 495	Y.D. Prokoshkin	(SERP) IGJPC
		Translated from DANS 344 469.	
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
SINGOVSKI 94	NC A107 1911	A.V. Singovsky	(SERP)
ARMSTRONG 93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG 89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE 78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)

NODE=M042215;NODE=M042

DESIG=1
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6

NODE=M042220

NODE=M042R1
NODE=M042R1

NODE=M042R1;LINKAGE=A

NODE=M042R2
NODE=M042R2

NODE=M042R2;LINKAGE=A

NODE=M042R3
NODE=M042R3

NODE=M042R3;LINKAGE=A

NODE=M042R04
NODE=M042R04

NODE=M042

REFID=53525
REFID=51063
REFID=51191
REFID=50174
REFID=47945
REFID=47961
REFID=46904
REFID=46921
REFID=45687
REFID=45685
REFID=45202
REFID=45212
REFID=44647
REFID=44103
REFID=45210
REFID=44648
REFID=43587
REFID=41010
REFID=21838
REFID=21837
REFID=21733
REFID=21850
REFID=21830
REFID=21813

$\rho(2150)$ $I^G(J^{PC}) = 1^+(1^-^-)$

OMITTED FROM SUMMARY TABLE

This entry was previously called $T_1(2190)$. See our mini-review under the $\rho(1700)$.

 $\rho(2150)$ MASS **e^+e^- PRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2201 ± 19	1 LEES	20 BABR	$e^+e^- \rightarrow K^+K^-\gamma$	
2227 $\pm 9 \pm 9$	2 LEES	20 RVUE	$e^+e^- \rightarrow K^+K^-$	
2039 ± 8 $^{+36}_{-18}$ 13k	3 ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+K^-\pi^0$	
2239.2 $\pm 7.1 \pm 11.3$	4 ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$	
2254 ± 22	5 LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
2150 $\pm 40 \pm 50$	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$	
1990 ± 80	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$	
2153 ± 37	BIAGINI	91 RVUE	$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-$	
2110 ± 50	6 CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-), 2(\pi^+\pi^-\pi^0)$	

1 From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is 3.5σ .

2 From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.

3 Could also be another state. Seen in J/ψ decay with branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (6.7 \pm 1.1^{+2.2}_{-1.8}) \times 10^{-6}$.

4 The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 2191	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2070	7 OAKDEN	94 RVUE	$0.36-1.55 \bar{p}p \rightarrow \pi\pi$
~ 2170	8 MARTIN	80B RVUE	
~ 2100	8 MARTIN	80C RVUE	

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2110 ± 35	9 ANISOVICH	02 SPEC	$0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2190	10 CUTTS	78B CNTR	$0.97-3 \bar{p}p \rightarrow \bar{N}N$
2155 ± 15	10,11 COUPLAND	77 CNTR	$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
2193 ± 2	10,12 ALSPECTOR	73 CNTR	$\bar{p}p$ S channel
2190 ± 10	13 ABRAMS	70 CNTR	$\bar{p}N$

 $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2140 ± 30	ALDE	95 GAM2	$38 \pi^- p \rightarrow \omega\pi^0 n$
2170 ± 30	ALDE	92C GAM4	$100 \pi^- p \rightarrow \omega\pi^0 n$

5 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

6 Includes ATKINSON 85.

7 See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

8 $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

9 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

10 Isospins 0 and 1 not separated.

11 From a fit to the total elastic cross section.

12 Referred to as T or T region by ALSPECTOR 73.

13 Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M032

NODE=M032205

NODE=M032M3
NODE=M032M3

OCCUR=2

ERROR=9

OCCUR=2

NODE=M032M3;LINKAGE=C

NODE=M032M3;LINKAGE=D

NODE=M032M3;LINKAGE=B

NODE=M032M3;LINKAGE=E

NODE=M032M1

NODE=M032M1

NODE=M032M2

NODE=M032M2

NODE=M032M4

NODE=M032M4

NODE=M032M3;LINKAGE=LE

NODE=M032M3;LINKAGE=A

NODE=M032M1;LINKAGE=CC

NODE=M032M;LINKAGE=P

NODE=M032M;LINKAGE=AY

NODE=M032M;LINKAGE=I

NODE=M032M;LINKAGE=E

NODE=M032M;LINKAGE=M

NODE=M032M;LINKAGE=B

$\rho(2150)$ WIDTH

e^+e^- PRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
70 \pm 38	14	LEES	20	BABR $e^+e^- \rightarrow K^+K^-\gamma$
127 \pm 14 \pm 4	15	LEES	20	RVUE $e^+e^- \rightarrow K^+K^-$
196 \pm 23 \pm 25	16	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+K^-\pi^0$
139.8 \pm 12.3 \pm 20.6	17	ABLIKIM	19L	BES3 $e^+e^- \rightarrow K^+K^-$
109 \pm 76	18	LEES	12G	BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$
350 \pm 40 \pm 50		AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
310 \pm 140		AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$
389 \pm 79		BIAGINI	91	RVUE $e^+e^- \rightarrow \pi^+\pi^-, K^+K^-$
410 \pm 100	19	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-), 2(\pi^+\pi^-\pi^0)$

- 14 From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is 3.5.
- 15 From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.
- 16 Could also be another state. Seen in J/ψ decay with branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (6.7 \pm 1.1^{+2.2}_{-1.8}) \times 10^{-6}$.
- 17 The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 296	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 40	20 OAKDEN	94	RVUE $0.36-1.55 \bar{p}p \rightarrow \pi\pi$
~ 250	21 MARTIN	80B	RVUE
~ 200	21 MARTIN	80C	RVUE

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
230 \pm 50	22 ANISOVICH	02	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
135 \pm 75	23,24 COUPLAND	77	CNTR $0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
98 \pm 8	24 ALSPECTOR	73	CNTR $\bar{p}p$ S channel
~ 85	25 ABRAMS	70	CNTR S channel $\bar{p}N$

$\pi^-p \rightarrow \omega\pi^0n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
320 \pm 70	ALDE	95 GAM2	$38 \pi^-p \rightarrow \omega\pi^0n$
~ 300	ALDE	92C GAM4	$100 \pi^-p \rightarrow \omega\pi^0n$
18 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.			
19 Includes ATKINSON 85.			
20 See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.			
21 $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.			
22 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.			
23 From a fit to the total elastic cross section.			
24 Isospins 0 and 1 not separated.			
25 Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.			

NODE=M032210

NODE=M032W3

NODE=M032W3

OCCUR=2

ERROR=10

OCCUR=2

NODE=M032W3;LINKAGE=C

NODE=M032W3;LINKAGE=D

NODE=M032W3;LINKAGE=B

NODE=M032W3;LINKAGE=E

NODE=M032W1

NODE=M032W1

NODE=M032W2

NODE=M032W2

NODE=M032W4

NODE=M032W4

NODE=M032W3;LINKAGE=LE

NODE=M032W3;LINKAGE=A

NODE=M032W1;LINKAGE=CC

NODE=M032W;LINKAGE=P

NODE=M032W;LINKAGE=AY

NODE=M032W;LINKAGE=E

NODE=M032W;LINKAGE=I

NODE=M032W;LINKAGE=B

$\rho(2150)$ DECAY MODES

NODE=M032215;NODE=M032

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	
$\Gamma_2 \pi^+ \pi^-$	seen
$\Gamma_3 K^+ K^-$	seen
$\Gamma_4 3(\pi^+ \pi^-)$	seen
$\Gamma_5 2(\pi^+ \pi^- \pi^0)$	seen
$\Gamma_6 \eta' \pi^+ \pi^-$	seen
$\Gamma_7 f_1(1285) \pi^+ \pi^-$	seen
$\Gamma_8 \omega \pi^0$	seen
$\Gamma_9 \omega \pi^0 \eta$	seen
$\Gamma_{10} p\bar{p}$	

 $\rho(2150) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(f_1(1285)\pi^+\pi^-)/\Gamma_{\text{total}}$	$\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-7})	DOCUMENT ID	TECN COMMENT

3.1±0.6±0.5 26 AUBERT 07AU BABR $10.6 e^+ e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$

26 Calculated by us from the reported value of cross section at the peak.

$\Gamma(\eta' \pi^+ \pi^-)/\Gamma_{\text{total}}$	$\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-8})	DOCUMENT ID	TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9±1.9 27 AUBERT 07AU BABR $10.6 e^+ e^- \rightarrow \eta' \pi^+ \pi^-\gamma$

27 Calculated by us from the reported value of cross section at the peak.

 $\rho(2150)$ REFERENCES

LEES	20	PR D101 012011	ABLIKIM <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab., JP)
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
BIAGINI	91	NC 104A 363	M.E. Biagini <i>et al.</i>	(FRAS, PRAG)
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	

DESIG=1

DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=4;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=5;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=6;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=7;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=8;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=9;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=10

NODE=M032230

NODE=M032G01
NODE=M032G01

NODE=M032G01;LINKAGE=AU

NODE=M032G02
NODE=M032G02

NODE=M032G02;LINKAGE=AU

NODE=M032

REFID=60211;ERROR=11
 REFID=59909
 REFID=59612
 REFID=55404
 REFID=54299
 REFID=52049
 REFID=48828
 REFID=48327
 REFID=48349
 REFID=47950
 REFID=45212
 REFID=44371
 REFID=44103
 REFID=45210
 REFID=41859
 REFID=41894
 REFID=41355
 REFID=22000
 REFID=21838
 REFID=21837
 REFID=21733
 REFID=21830
 REFID=21824
 REFID=21813
 REFID=21807
 REFID=21805
 REFID=48054

NODE=M103

 $\phi(2170)$ $I^G(J^{PC}) = 0^-(1^{--})$

Observed by AUBERT,BE 06D in the initial-state radiation process
 $e^+ e^- \rightarrow \phi f_0(980) \gamma$.

 $\phi(2170)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2135 \pm 8 \pm 9	95	ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta \phi f_0(980)$
2239.2 \pm 7.1 \pm 11.3	1	ABLIKIM	19L BES3	$e^+ e^- \rightarrow K^+ K^-$
2200 \pm 6 \pm 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta \phi \pi^+ \pi^-$
2180 \pm 8 \pm 8	2,3 LEES		12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
2079 \pm 13 \pm 79	4.8k	4 SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
2186 \pm 10 \pm 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$
2125 \pm 22 \pm 10	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$
2192 \pm 14	116	5 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
2169 \pm 20	149	5 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
2175 \pm 10 \pm 15	201	3,6 AUBERT,BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$

1 The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.2 Fit includes interference with the $\phi(1680)$.3 From the $\phi f_0(980)$ component.

4 From a fit with two incoherent Breit-Wigners.

5 From the $K^+ K^- f_0(980)$ component.

6 Superseded by LEES 12F.

NODE=M103

NODE=M103M

NODE=M103M

OCCUR=2

NODE=M103M;LINKAGE=C

NODE=M103M;LINKAGE=A

NODE=M103M;LINKAGE=AB

NODE=M103M;LINKAGE=SH

NODE=M103M;LINKAGE=AU

NODE=M103M;LINKAGE=B

 $\phi(2170)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
104 \pm 24 \pm 12	95	ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta \phi f_0(980)$
139.8 \pm 12.3 \pm 20.6	7	ABLIKIM	19L BES3	$e^+ e^- \rightarrow K^+ K^-$
104 \pm 15 \pm 15	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta \phi \pi^+ \pi^-$
77 \pm 15 \pm 10	8,9 LEES		12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
192 \pm 23 \pm 25	4.8k	10 SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
65 \pm 23 \pm 17	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$
61 \pm 50 \pm 13	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$
71 \pm 21	116	11 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
102 \pm 27	149	11 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
58 \pm 16 \pm 20	201	9,12 AUBERT,BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$

7 The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.8 Fit includes interference with the $\phi(1680)$.9 From the $\phi f_0(980)$ component.

10 From a fit with two incoherent Breit-Wigners.

11 From the $K^+ K^- f_0(980)$ component.

12 Superseded by LEES 12F.

NODE=M103W

NODE=M103W

OCCUR=2

NODE=M103W;LINKAGE=C

NODE=M103W;LINKAGE=A

NODE=M103W;LINKAGE=AB

NODE=M103W;LINKAGE=SH

NODE=M103W;LINKAGE=AU

NODE=M103W;LINKAGE=B

$\phi(2170)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	seen
$\Gamma_2 \phi\eta$	
$\Gamma_3 \phi\pi\pi$	
$\Gamma_4 \phi f_0(980)$	seen
$\Gamma_5 K^+ K^- \pi^+ \pi^-$	
$\Gamma_6 K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_7 K^+ K^- \pi^0 \pi^0$	
$\Gamma_8 K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
$\Gamma_9 K^{*0} K^\pm \pi^\mp$	not seen
$\Gamma_{10} K^*(892)^0 \bar{K}^*(892)^0$	not seen

$\phi(2170) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_2 \Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1.7 \pm 0.7 \pm 1.3$	483	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi\eta\gamma$

$\Gamma(\phi f_0(980)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_4 \Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 0.3 \pm 0.3$	13,14	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \phi\pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

$2.5 \pm 0.8 \pm 0.4$ 201 14,¹⁵ AUBERT,BE 06D BABR $10.6 e^+ e^- \rightarrow K^+ K^- \pi\pi\gamma$

13 From a fit with constructive interference with the $\phi(1680)$. In a fit with destructive interference, the value is larger by a factor of 12.

14 From the $\phi f_0(980)$ component.

15 Superseded by LEES 12F.

$\phi(2170) \Gamma(i) \Gamma(e^+ e^-)/\Gamma^2(\text{total})$

$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

$1.65 \pm 0.15 \pm 0.18$ 4.8k 16 SHEN 09 BELL $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

16 Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

$\phi(2170)$ BRANCHING RATIOS

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

$\Gamma(K^{*0} K^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	AUBERT	07AK BABR	$10.6 \text{ GeV } e^+ e^-$

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

$\phi(2170)$ REFERENCES

ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M103215;NODE=M103

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=5
 DESIG=9
 DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=3
 DESIG=6
 DESIG=4
 DESIG=7
 DESIG=8
 DESIG=10

NODE=M103230

NODE=M103G2
NODE=M103G2

NODE=M103G1
NODE=M103G1

NODE=M103G1;LINKAGE=A

NODE=M103G1;LINKAGE=AB
NODE=M103G1;LINKAGE=B

NODE=M103220

NODE=M103G01
NODE=M103G01

NODE=M103G01;LINKAGE=SH

NODE=M103225

NODE=M103R01
NODE=M103R01

NODE=M103R02
NODE=M103R02

NODE=M103R03
NODE=M103R03

NODE=M103R04
NODE=M103R04

NODE=M103

REFID=59605
REFID=59612
REFID=56773
REFID=54298
REFID=53349
REFID=53000
REFID=52154
REFID=52242
REFID=51908
REFID=51511

$f_0(2200)$ $I^G(J^{PC}) = 0^+(0^{++})$

OMITTED FROM SUMMARY TABLE

Seen in $K_S^0 K_S^0$ (AUGUSTIN 88), $K^+ K^-$ (ABLIKIM 05Q) and $\eta\eta$ (BINON 05) system. Not seen in $\Upsilon(1S)$ radiative decays (BARU 89).

NODE=M112

 $f_0(2200)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2187±14 OUR AVERAGE				
2170±20 ⁺¹⁰ ₋₁₅		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
2197±17	1 AUGUSTIN	88 DM2		$J/\psi \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2206±12± 8	381	2,3 DOBBS	15	$J/\psi \rightarrow \gamma K^+K^-$
2188±17±16	203	2,3 DOBBS	15	$\psi(2S) \rightarrow \gamma K^+K^-$
2210±50		4 BINON	05 GAMS	33 $\pi^- p \rightarrow \eta\eta n$
~ 2122		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2321		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$

1 Cannot determine spin to be 0.

2 Using CLEO-c data but not authored by the CLEO Collaboration.

3 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 238$ MeV.

4 First solution, PWA is ambiguous.

NODE=M112M

NODE=M112M

OCCUR=2

OCCUR=2

NODE=M112M;LINKAGE=A

NODE=M112M;LINKAGE=B

NODE=M112M;LINKAGE=C

NODE=M112M;LINKAGE=BI

 $f_0(2200)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
207±40 OUR AVERAGE			
220±60 ⁺⁴⁰ ₋₄₅	ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
201±51	5 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
380±90	6 BINON	05 GAMS	33 $\pi^- p \rightarrow \eta\eta n$
~ 273	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 223	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
5 Cannot determine spin to be 0.			
6 First solution, PWA is ambiguous.			

NODE=M112W

NODE=M112W

OCCUR=2

NODE=M112W;LINKAGE=A

NODE=M112W;LINKAGE=BI

 $f_0(2200)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)

NODE=M112

REFID=56805

REFID=50958

REFID=50780

REFID=44103

REFID=40917

REFID=40574

$f_J(2220)$ $I^G(J^{PC}) = 0^+(2^{++} \text{ or } 4^{++})$

OMITTED FROM SUMMARY TABLE

Needs confirmation. See our mini-review in the 2004 edition of this
Review, PDG 04.

$f_J(2220)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2231.1 ± 3.5 OUR AVERAGE				

2235	± 4	± 6	74	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
2230	± 6	± 16	46	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=2
2232	± 8	± 15	23	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$	OCCUR=3
2235	± 4	± 5	32	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$	OCCUR=4
2209	± 17	± 10		ASTON	88F	LASS	$11 K^- p \rightarrow K^+ K^- \Lambda$	
2230	± 20			BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
2220	± 10		41	¹ ALDE	86B	GA24	$38-100 \pi p \rightarrow \eta\eta\eta'$	
2230	± 6	± 14	93	BALTRUSAIT..86D	MRK3	$e^+ e^- \rightarrow \gamma K^+ K^-$		
2232	± 7	± 7	23	BALTRUSAIT..86D	MRK3	$e^+ e^- \rightarrow \gamma K_S^0 K_S^0$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

2223.9 ± 2.5 ² VLADIMIRSK..08 SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$

2246 ± 36 BAI 98H BES $J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ ALDE 86B uses data from both the GAMS-2000 and GAMS-4000 detectors.
² $J^{PC} = 2^{++}$. Systematic uncertainties not evaluated

NODE=M082

NODE=M082M

NODE=M082M

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

NODE=M082M;LINKAGE=A
NODE=M082M;LINKAGE=VL

$f_J(2220)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
23 ± 8 OUR AVERAGE					

19 ± 13	± 12	74	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
20 ± 20	± 17	46	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=2
20 ± 25	± 14	23	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$	OCCUR=3
15 ± 12	± 9	32	BAI	96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$	OCCUR=4
60 ± 107			ASTON	88F	LASS	$11 K^- p \rightarrow K^+ K^- \Lambda$	
80 ± 30			BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
26 ± 20	± 17	93	BALTRUSAIT..86D	MRK3	$e^+ e^- \rightarrow \gamma K^+ K^-$		
18 ± 23	± 10	23	BALTRUSAIT..86D	MRK3	$e^+ e^- \rightarrow \gamma K_S^0 K_S^0$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.6 ± 2.5 ¹ VLADIMIRSK..08 SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$

<80 90 ALDE 87C GAM2 $38 \pi^- p \rightarrow \eta'\eta n$

¹ $J^{PC} = 2^{++}$. Systematic uncertainties not evaluated

NODE=M082W

NODE=M082W

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

NODE=M082W;LINKAGE=VL

NODE=M082215;NODE=M082

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	not seen
$\Gamma_2 \pi^+ \pi^-$	not seen
$\Gamma_3 K\bar{K}$	not seen
$\Gamma_4 p\bar{p}$	not seen
$\Gamma_5 \gamma\gamma$	not seen
$\Gamma_6 \eta\eta'(958)$	seen
$\Gamma_7 \phi\phi$	not seen
$\Gamma_8 \eta\eta$	not seen

DESIG=5

DESIG=6;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=1

DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=7;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=8;OUR EST; \rightarrow UNCHECKED \leftarrow

$f_J(2220) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_5/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
< 1.4	95	1 ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}} = 91, 183-209 \text{ GeV}$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 5.6	95	1 GODANG	97 CLE2	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
< 86	95	1 ALBRECHT	90G ARG	$\gamma\gamma \rightarrow K^+ K^-$	
<1000	95	2 ALTHOFF	85B TASS	$\gamma\gamma, K\bar{K}\pi$	

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_5/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.5	95	ALAM	98C CLE2	$\gamma\gamma \rightarrow \pi^+ \pi^-$	
1 Assuming $J^P = 2^+$.					
2 True for $J^P = 0^+$ and $J^P = 2^+$.					

 $f_J(2220) \Gamma(i)\Gamma(p\bar{p})/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<18	95	1 AMSLER	01 CBAR	$1.4-1.5 \text{ p}\bar{p} \rightarrow \pi^0 \pi^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<(11-42)	99	2 HASAN	96 SPEC	$1.35-1.55 \text{ p}\bar{p} \rightarrow \pi^+ \pi^-$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_7/\Gamma$
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<6	95	3 EVANGELIS...	98 SPEC	$1.1-2.0 \text{ p}\bar{p} \rightarrow \phi\phi$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_8/\Gamma$
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<4	95	1 AMSLER	01 CBAR	$1.4-1.5 \text{ p}\bar{p} \rightarrow \eta\eta$	
1 For $J^P = 2^+$ in the mass range 2222–2240 MeV and the total width between 10 and 20 MeV.					
2 For $J^P = 2^+$ and $J^P = 4^+$ in the mass range 2220–2245 MeV and the total width of 15 MeV.					
3 For $J^P = 2^+$, the mass of 2235 MeV and the total width of 15 MeV.					

 $f_J(2220) \text{ BRANCHING RATIOS}$

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		1 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$	
not seen		1 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$	

1 Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		1 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$	
not seen		1 DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$	

1 Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$					Γ_1/Γ_3
VALUE		DOCUMENT ID	TECN	COMMENT	
1.0±0.5		BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma 2\pi, K\bar{K}$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
not seen		1 AUBERT	07AV BABR	$B \rightarrow p\bar{p}K(*)$	
not seen		WANG	05A BELL	$B^+ \rightarrow \bar{p}pK^+$	
<3.0	95	2 EVANGELIS...	97 SPEC	$1.96-2.40 \bar{p}p \rightarrow K_S^0 K_S^0$	
<1.1	99.7	3 BARNES	93 SPEC	$1.3-1.57 \bar{p}p \rightarrow K_S^0 K_S^0$	
<2.6	99.7	3 BARDIN	87 CNTR	$1.3-1.5 \bar{p}p \rightarrow K^+ K^-$	
<3.6	99.7	3 SCULLI	87 CNTR	$1.29-1.55 \bar{p}p \rightarrow K^+ K^-$	

NODE=M082220

NODE=M082G1

NODE=M082G1

NODE=M082G3

NODE=M082G3

NODE=M082223

NODE=M082GG1

NODE=M082GG1

NODE=M082GG2

NODE=M082GG2

NODE=M082GG3

NODE=M082GG3

NODE=M082GG;LINKAGE=A

NODE=M082GG;LINKAGE=B

NODE=M082GG;LINKAGE=C

NODE=M082225

NODE=M082R00

NODE=M082R00

OCCUR=2

NODE=M082R00;LINKAGE=A

NODE=M082R01

NODE=M082R01

OCCUR=2

NODE=M082R01;LINKAGE=A

NODE=M082R2

NODE=M082R2

NODE=M082R1

NODE=M082R1

¹ Assuming $\Gamma < 30$ MeV.

² Assuming $\Gamma \sim 20$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.

³ Assuming $\Gamma = 30\text{-}35$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.

$\Gamma(p\bar{p})/\Gamma(K\bar{K})$					Γ_4/Γ_3
VALUE	DOCUMENT ID	TECN	COMMENT		
0.17±0.09	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}, K\bar{K}$		

$f_J(2220)$ REFERENCES

DOBBS 15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
VЛАДИМИРСК... 08	PAN 71 2129 Translated from YAF 71 2166.	V.V. Vladimirska <i>et al.</i>	(ITEP)
AUBERT 07AV	PR D76 092004	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG 05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
PDG 04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ACCIARRI 01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
AMSLER 01	PL B520 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ALAM 98C	PRl 81 3328	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BAI 98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
EVANGELIS... 98	PR D57 5370	C. Evangelista <i>et al.</i>	(JETSET Collab.)
EVANGELIS... 97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
GODANG 97	PRL 79 3829	R. Godang <i>et al.</i>	(CLEO Collab.)
BAI 96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
HASAN 96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BARNES 93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
ALBRECHT 90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASTON 88F	PL B215 199	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BOLONKIN 88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ALDE 87C	SJNP 45 255 Translated from YAF 45 405	D. Alde <i>et al.</i>	
BARDIN 87	PL B195 292	G. Bardin <i>et al.</i>	(SACL, FERR, CERN, PADO+)
SCULLI 87	PRL 58 1715	J. Sculli <i>et al.</i>	(NYU, BNL)
ALDE 86B	PL B177 120	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BALTRUSAIT... 86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
ALTHOFF 85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)

NODE=M082R1;LINKAGE=AU

NODE=M082R1;LINKAGE=C

NODE=M082R1;LINKAGE=B

NODE=M082R3

NODE=M082R3

NODE=M082

REFID=56805

REFID=52681

REFID=51990

REFID=50651

REFID=49653

REFID=48321

REFID=48558

REFID=46326

REFID=46342

REFID=46365

REFID=45687

REFID=45760

REFID=44736

REFID=45197

REFID=43601

REFID=41374

REFID=40585

REFID=40580

REFID=47474

REFID=40235

REFID=40023

REFID=21864

REFID=21865

REFID=21349

OTHER RELATED PAPERS

DEL-AMO-SA... 100 PRL 105 172001 P. del Amo Sanchez *et al.* (BABAR Collab.)

REFID=53533

NODE=M115

$\eta(2225)$

$I^G(J^{PC}) = 0^+(0^-+)$

OMMITTED FROM SUMMARY TABLE

Seen in $J/\psi \rightarrow \gamma\phi\phi$. Possibly seen in $B \rightarrow \phi\phi K$ by LEES 11A.

NODE=M115

$\eta(2225)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2221⁺¹³₋₁₀ OUR AVERAGE				

2216^{+ 4 +21}_{- 5 -11} ¹ ABLIKIM 16N BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

2240^{+30 +30}_{-20 -20} 196 ± 19 ABLIKIM 08I BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

2230 ± 25 ± 15 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

2214 ± 20 ± 13 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2220 BISELLO 86B DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

OCCUR=2

NODE=M115M;LINKAGE=B

NODE=M115W

NODE=M115W

$\eta(2225)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
185^{+ 40}_{- 20} OUR AVERAGE				

185^{+ 12 +43}_{- 14 -17} ¹ ABLIKIM 16N BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

190^{+ 30 +60}_{- 40 -40} 196 ± 19 ABLIKIM 08I BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

150^{+ 300}_{- 60} ± 60 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 80 BISELLO 86B DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

1 From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

$\eta(2225)$ REFERENCES

ABLIKIM LEES ABLIKIM BAI BISELLO	16N 11A 08I 90B 86B	PR D93 112011 PR D84 012001 PL B662 330 PRL 65 1309 PL B179 294	M. Ablikim J.P. Lees <i>et al.</i> M. Ablikim <i>et al.</i> Z. Bai <i>et al.</i> D. Bisello <i>et al.</i>	(BESIII Collab.) (BABAR Collab.) (BES Collab.) (Mark III Collab.) (DM2 Collab.)
--	---------------------------------	---	---	---

$\rho_3(2250)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

OMMITTED FROM SUMMARY TABLE

Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $f_4(2300)$, $\rho_5(2350)$.

$\rho_3(2250)$ MASS

$\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~2232	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~2090	¹ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~2250	² MARTIN	80B	RVUE	
~2300	² MARTIN	80C	RVUE	
~2140	³ CARTER	78B	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~2150	⁴ CARTER	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

¹ See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

³ $I = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.

⁴ $I(J^P) = 1(3^-)$ from amplitude analysis.

S-CHANNEL $\bar{p}p$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2260±20	⁵ ANISOVICH	02	SPEC	$0.6-1.9 p\bar{p} \rightarrow \omega\pi^0$, $\omega\eta\pi^0, \pi^+\pi^-$
~2190	⁶ CUTTS	78B	CNTR	0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155±15	^{6,7} COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193± 2	^{6,8} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2190±10	⁹ ABRAMS	70	CNTR	S channel $\bar{p}N$

⁵ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

⁶ Isospins 0 and 1 not separated.

⁷ From a fit to the total elastic cross section.

⁸ Referred to as T or T' region by ALSPECTOR 73.

⁹ Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

$\pi^- p \rightarrow \eta\pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2290±20±30	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

$\rho_3(2250)$ WIDTH

NODE=M115W;LINKAGE=A

NODE=M115

REFID=57512
REFID=16595
REFID=52255
REFID=41354
REFID=22101

NODE=M044

NODE=M044

NODE=M044205

NODE=M044M1
NODE=M044M1

NODE=M044M1;LINKAGE=CC

NODE=M044M1;LINKAGE=P
NODE=M044M1;LINKAGE=K
NODE=M044M1;LINKAGE=J

NODE=M044M2
NODE=M044M2

NODE=M044M;LINKAGE=AY

NODE=M044M2;LINKAGE=I
NODE=M044M2;LINKAGE=E
NODE=M044M2;LINKAGE=M
NODE=M044M2;LINKAGE=B

NODE=M044M3
NODE=M044M3

NODE=M044210

$\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 220	HASAN 94	RVUE		$\bar{p}p \rightarrow \pi\pi$
~ 60	10 OAKDEN 94	RVUE		0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	11 MARTIN 80B	RVUE		
~ 200	11 MARTIN 80C	RVUE		
~ 150	12 CARTER 78B	CNTR 0		0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 200	13 CARTER 77	CNTR 0		0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

10 See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

11 $I(JP) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

12 $I = 0, 1$. $JP = 3^-$ from Barrelet-zero analysis.

13 $I(JP) = 1(3^-)$ from amplitude analysis.

NODE=M044W1
NODE=M044W1

S-CHANNEL $\bar{p}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
160±25	14 ANISOVICH 02	SPEC		$0.6-1.9 p\bar{p} \rightarrow \omega\pi^0,$ $\omega\eta\pi^0, \pi^+\pi^-$
135±75	15,16 COUPLAND 77	CNTR 0		0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98±8	16 ALSPECTOR 73	CNTR		$\bar{p}p$ S channel
~ 85	17 ABRAMS 70	CNTR		S channel $\bar{p}N$

14 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

15 From a fit to the total elastic cross section.

16 Isospins 0 and 1 not separated.

17 Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M044W1;LINKAGE=CC

NODE=M044W1;LINKAGE=P

NODE=M044W1;LINKAGE=K

NODE=M044W1;LINKAGE=J

NODE=M044W2
NODE=M044W2

 $\pi^-p \rightarrow \eta\pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
230±50±80	AMELIN 00	VES	37 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

NODE=M044W3
NODE=M044W3

 $\rho_3(2250)$ REFERENCES

ANISOVICH 02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH 01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER 78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
CARTER 77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE 75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS 70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER 68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)

NODE=M044

REFID=48828
REFID=48327
REFID=48349
REFID=47432
REFID=47950
REFID=45212
REFID=44103
REFID=45210
REFID=21838
REFID=21837
REFID=21964
REFID=21733
REFID=21963
REFID=21830
REFID=21824
REFID=21813
REFID=21807
REFID=21805

NODE=M107

 $f_2(2300)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$f_2(2300)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2297±28	¹ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2243 $^{+7+3}_{-6-29}$	² UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
2270 ± 12	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2327 $\pm 9 \pm 6$	ABE	04	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
2231 ± 10	BOOTH	86	OMEG 85 $\pi^- Be \rightarrow 2\phi Be$
2220 $^{+90}_{-20}$	LINDENBAUM	84	RVUE
2320 ± 40	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi 2^{++} S_2$, D_2 , and D_0 is 6^{+15}_{-5} , 25^{+18}_{-14} , and 69^{+16}_{-27} , respectively.

² Spin 2 preferred, tentatively assigned to $f_2(2300)$.

NODE=M107M

NODE=M107M

NODE=M107M;LINKAGE=C

NODE=M107M;LINKAGE=A

NODE=M107W

NODE=M107W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
149±41	³ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
145 $\pm 12^{+27}_{-34}$	⁴ UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
90 ± 29	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
275 $\pm 36 \pm 20$	ABE	04	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
133 ± 50	BOOTH	86	OMEG 85 $\pi^- Be \rightarrow 2\phi Be$
200 ± 50	LINDENBAUM	84	RVUE
220 ± 70	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

³ Includes data of ETKIN 85.

⁴ Spin 2 preferred, tentatively assigned to $f_2(2300)$.

NODE=M107W;LINKAGE=C
NODE=M107W;LINKAGE=A

NODE=M107215;NODE=M107

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \phi\phi$	seen
$\Gamma_2 K\bar{K}$	seen
$\Gamma_3 \gamma\gamma$	seen

 $f_2(2300) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_3/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.2 $^{+0.5+1.3}_{-0.4-2.2}$	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$	
44 $\pm 6 \pm 12$	⁵ ABE	04	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$	
5 Assuming spin 2.				

DESIG=1;OUR EST; \rightarrow UNCHECKED
DESIG=2;OUR EST; \rightarrow UNCHECKED
DESIG=3;OUR EST; \rightarrow UNCHECKED

NODE=M107225

NODE=M107G1
NODE=M107G1

NODE=M107G1;LINKAGE=AB

NODE=M107

REFID=55592
REFID=51191REFID=49650
REFID=40285
REFID=21870
REFID=21871
REFID=21869
REFID=21866

UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirska <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIPV, GLAS, CERN)
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)
ETKIN	82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)

NODE=M041

 $f_4(2300)$ $I^G(J^{PC}) = 0^+(4^{++})$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_0(2350)$. Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $\rho_5(2350)$.

NODE=M041

 $f_4(2300)$ MASS

NODE=M041205

NODE=M041M

NODE=M041M1

NODE=M041M1

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 2314	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2300	1 MARTIN	80B	RVUE
~ 2300	1 MARTIN	80C	RVUE
~ 2340	2 CARTER	78B	CNTR 0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2330	DULUDE	78B	OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$
~ 2310	3 CARTER	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

1 $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.2 $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.3 $I(J^P) = 0(4^+)$ from amplitude analysis.

NODE=M041M1;LINKAGE=P

NODE=M041M1;LINKAGE=K

NODE=M041M1;LINKAGE=J

S-CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2283±17	4 ANISOVICH	00J	SPEC
~ 2380	5 CUTTS	78B	CNTR 0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2345±15	5,6 COUPLAND	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2359± 2	5,7 ALSPECTOR	73	CNTR $\bar{p}p$ S channel
2375±10	ABRAMS	70	CNTR S channel $\bar{N}N$

4 From the combined analysis of ANISOVICH 99c and ANISOVICH 99f on $\bar{p}p \rightarrow \eta\pi^0\pi^0$, $\pi^0\pi^0$, $\eta\eta$, $\eta\eta'$, $\pi^+\pi^-$.

5 Isospins 0 and 1 not separated.

6 From a fit to the total elastic cross section.

7 Referred to as U or U region by ALSPECTOR 73.

NODE=M041M2;LINKAGE=AN

NODE=M041M2;LINKAGE=I

NODE=M041M2;LINKAGE=EE

NODE=M041M2;LINKAGE=M

 $\pi^- p \rightarrow \eta\pi\pi n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2330±20±40	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M041M3

NODE=M041M3

NODE=M041M4

NODE=M041M4

→ UNCHECKED ←

 pp CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	COMMENT
2320±60 OUR ESTIMATE		
2332±15	BARBERIS	00F 450 $pp \rightarrow p_f \omega \omega p_s$

NODE=M041210

NODE=M041W1

NODE=M041W1

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 278	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 200	8 MARTIN	80C	RVUE
~ 150	9 CARTER	78B	CNTR 0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 210	10 CARTER	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

8 $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.9 $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.10 $I(J^P) = 0(4^+)$ from amplitude analysis.

NODE=M041W1;LINKAGE=P

NODE=M041W1;LINKAGE=K

NODE=M041W1;LINKAGE=J

S CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
310 ± 25	11 ANISOVICH 00J	SPEC	
135 ± 150 - 65	12,13 COUPLAND 77	CNTR	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
165 ± 18 - 8	13 ALSPECTOR 73	CNTR	$\bar{p}p$ S channel
~190	ABRAMS 70	CNTR	S channel $\bar{N}N$
11 From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on $\bar{p}p \rightarrow \eta\pi^0\pi^0$, $\pi^0\pi^0$, $\eta\eta$, $\eta\eta'$, $\pi^+\pi^-$.			
12 From a fit to the total elastic cross section.			
13 Isospins 0 and 1 not separated.			

NODE=M041W2
NODE=M041W2 **$\pi^- p \rightarrow \eta\pi\pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
235 ± 50 ± 40	AMELIN 00	VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M041W2;LINKAGE=AN

NODE=M041W2;LINKAGE=E
NODE=M041W2;LINKAGE=INODE=M041W3
NODE=M041W3 **$p\bar{p}$ CENTRAL PRODUCTION**

VALUE (MeV)	DOCUMENT ID	COMMENT
250 ± 80 OUR ESTIMATE		
• • • We do not use the following data for averages, fits, limits, etc. • • •		
260 ± 57	BARBERIS 00F	450 $p\bar{p} \rightarrow p_f\omega\omega p_s$

NODE=M041W4
NODE=M041W4
→ UNCHECKED ← **$f_4(2300)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho\rho$	seen
$\Gamma_2 \omega\omega$	seen
$\Gamma_3 \eta\pi\pi$	seen
$\Gamma_4 \pi\pi$	seen
$\Gamma_5 K\bar{K}$	seen
$\Gamma_6 N\bar{N}$	seen

NODE=M041215;NODE=M041

 $f_4(2300)$ BRANCHING RATIOS

VALUE	DOCUMENT ID	COMMENT	Γ_1/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.8 ± 0.5	BARBERIS 00F	450 $p\bar{p} \rightarrow p_f\omega\omega p_s$	

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←

NODE=M041220

NODE=M041R1
NODE=M041R1 **$f_4(2300)$ REFERENCES**

AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARBERIS 00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH 99C	PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH 99F	NP A651 253	A.V. Anisovich <i>et al.</i>	
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER 78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE 78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARII) JP
CARTER 77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS 70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)

NODE=M041

REFID=47432
REFID=47950
REFID=47962
REFID=46903
REFID=46926
REFID=44103
REFID=21838
REFID=21837
REFID=21964
REFID=21733
REFID=21850
REFID=21963
REFID=21830
REFID=21813
REFID=21807

NODE=M169

f₀(2330)

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

f₀(2330) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2314±25	1 BUGG	04A RVUE	
2337±14	ANISOVICH	00J SPEC	2.0 $\bar{p}p \rightarrow \pi\pi, \eta\eta$
~2321	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
1 Partial wave analysis of the data on $p\bar{p} \rightarrow \Lambda\bar{\Lambda}$ from BARNES 00.			

NODE=M169M

NODE=M169M

NODE=M169M;LINKAGE=BU

NODE=M169W

NODE=M169W

NODE=M169W;LINKAGE=BU

NODE=M169

REFID=50158
 REFID=47950
 REFID=47965
 REFID=44103

NODE=M108

f₂(2340)

$$I^G(J^{PC}) = 0^+(2^{++})$$

f₂(2340) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2345⁺⁵⁰₋₄₀ OUR AVERAGE				
2362 ⁺³¹ ₋₃₀ ⁺¹⁴⁰ ₋₆₃	5.5k	1 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
2339±55		2 ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2350±7	80k	3 UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
2392±10		BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
2360±20		LINDENBAUM	84 RVUE	

NODE=M108M

NODE=M108M

NODE=M108M;LINKAGE=A

NODE=M108M;LINKAGE=C

NODE=M108M;LINKAGE=ST

NODE=M108W

NODE=M108W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
322⁺⁷⁰₋₆₀ OUR AVERAGE				
334 ⁺⁶² ₋₅₄ ⁺¹⁶⁵ ₋₁₀₀	5.5k	4 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
319 ⁺⁸¹ ₋₆₉		5 ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi\phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
218±16	80k	6 UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
198±50		BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
150 ⁺¹⁵⁰ ₋₅₀		LINDENBAUM	84 RVUE	

4 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

5 Includes data of ETKIN 85.

6 Statistical error only.

$f_2(2340)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \phi\phi$	seen
$\Gamma_2 \eta\eta$	seen

$f_2(2340)$ BRANCHING RATIOS

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_2/Γ
seen	UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$

$f_2(2340)$ REFERENCES

ABLIKIM UMAN ETKIN BOOTH ETKIN LINDENBAUM	13N 06 88 86 85 84	PR D87 092009 PR D73 052009 PL B201 568 NP B273 677 PL 165B 217 CNPP 13 285	Ablikim M. <i>et al.</i> I. Uman <i>et al.</i> A. Etkin <i>et al.</i> P.S.L. Booth <i>et al.</i> A. Etkin <i>et al.</i> S.J. Lindenbaum	(BESIII Collab.) (FNAL E835) (BNL, CUNY) (LIVP, GLAS, CERN) (BNL, CUNY) (CUNY)
--	-----------------------------------	--	--	---

$\rho_5(2350)$

$$\rho^G(J^{PC}) = 1^+(5^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_1(2400)$. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $f_4(2300)$.

$\rho_5(2350)$ MASS

$\pi^- p \rightarrow \omega\pi^0 n$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2330±35	ALDE	95	GAM2 38 $\pi^- p \rightarrow \omega\pi^0 n$

$\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 2303	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2300	¹ MARTIN	80B	RVUE	
~ 2250	¹ MARTIN	80C	RVUE	
~ 2500	² CARTER	78B	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2480	³ CARTER	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

S-CHANNEL $\bar{N}N$

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2300±45	⁴ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0$, $\omega\eta\pi^0$, $\pi^+\pi^-$
2295±30	ANISOVICH	00J	SPEC	
~ 2380	⁵ CUTTS	78B	CNTR	0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2345±15	^{5,6} COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2359± 2	^{5,7} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2350±10	⁸ ABRAMS	70	CNTR	S channel $\bar{N}N$
2360±25	⁹ OH	70B	HDBC –0	$\bar{p}(pn)$, $K^* K 2\pi$

NODE=M108W;LINKAGE=A

NODE=M108W;LINKAGE=C

NODE=M108W;LINKAGE=ST

NODE=M108215;NODE=M108

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2

NODE=M108220

NODE=M108R01
NODE=M108R01

NODE=M108

REFID=55387
REFID=51063
REFID=40285
REFID=21870
REFID=21871
REFID=21869

NODE=M033

NODE=M033

NODE=M033205

NODE=M033M

NODE=M033M3
NODE=M033M3

NODE=M033M1
NODE=M033M1

NODE=M033M2
NODE=M033M2

$\pi^- p \rightarrow K^+ K^- n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2307±6	ALPER	80	CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$
1	$I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.			
2	$I = 0(1); J^P = 5^-$ from Barrelet-zero analysis.			
3	$I(J^P) = 1(5^-)$ from amplitude analysis.			
4	From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.			
5	Isospins 0 and 1 not separated.			
6	From a fit to the total elastic cross section.			
7	Referred to as U or U region by ALSPECTOR 73.			
8	For $I = 1 \bar{N}N$.			
9	No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.			

NODE=M033M4
NODE=M033M4NODE=M033M1;LINKAGE=P
NODE=M033M1;LINKAGE=K
NODE=M033M1;LINKAGE=J
NODE=M033M2;LINKAGE=AYNODE=M033M2;LINKAGE=I
NODE=M033M2;LINKAGE=E
NODE=M033M2;LINKAGE=M
NODE=M033M2;LINKAGE=A
NODE=M033M2;LINKAGE=N

NODE=M033210

NODE=M033W3
NODE=M033W3NODE=M033W1
NODE=M033W1 $\rho_5(2350)$ WIDTH $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
400±100	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~169	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~250	10 MARTIN	80B	RVUE	
~300	10 MARTIN	80C	RVUE	
~150	11 CARTER	78B	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~210	12 CARTER	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
260±75	13 ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
235^{+65}_{-40}	ANISOVICH	00J	SPEC	
135^{+150}_{-65}	14,15 COUPLAND	77	CNTR 0	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
165^{+18}_{-8}	15 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
< 60	16 OH	70B	HDBC –0	$\bar{p}(pn), K^* K 2\pi$
~140	ABRAMS	67C	CNTR	S channel $\bar{p}N$

NODE=M033W2
NODE=M033W2 $\pi^- p \rightarrow K^+ K^- n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
245±20	ALPER	80	CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$
10	$I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.			
11	$I = 0(1); J^P = 5^-$ from Barrelet-zero analysis.			
12	$I(J^P) = 1(5^-)$ from amplitude analysis.			
13	From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.			
14	From a fit to the total elastic cross section.			
15	Isospins 0 and 1 not separated.			
16	No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.			

NODE=M033W4
NODE=M033W4

OCCUR=2

NODE=M033W1;LINKAGE=P
NODE=M033W1;LINKAGE=K
NODE=M033W1;LINKAGE=J
NODE=M033W2;LINKAGE=AYNODE=M033W2;LINKAGE=E
NODE=M033W2;LINKAGE=I
NODE=M033W2;LINKAGE=N

$f_5(2350)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington
CARTER	78B	NP B141 467	A.A. Carter
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>
OH	73	NP B51 57	B.Y. Oh <i>et al.</i>
CHAPMAN	71B	PR D4 1275	J.W. Chapman <i>et al.</i>
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>
OH	70B	PRL 24 1257	B.Y. Oh <i>et al.</i>
ABRAMS	67C	PRL 18 1209	R.J. Abrams <i>et al.</i>
			(RAL, LOQM, PNPI+) (GAMS Collab.) JP (LOQM) (AMST, CERN, CRAC, MPIM+) (LOUC, RHEL) JP (DURH) JP (LOQM) (STON, WISC) (LOQM, RHEL) JP (LOQM, RHEL) (RUTG, UPNJ) (MSU) (MICH) (BNL) (MSU) (BNL)

 $f_6(2510)$

$$I^G(J^{PC}) = 0^+(6^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

 $f_6(2510)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2465±50 OUR AVERAGE			Error includes scale factor of 2.1.
2420±30	ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
2510±30	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2485±40	¹ ANISOVICH	00J	SPEC 1.92–2.41 $p\bar{p}$

¹ From the combined analysis of ANISOVICH 99c, ANISOVICH 99f, ANISOVICH 99j, ANISOVICH 99k, and ANISOVICH 00b.

 $f_6(2510)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
255±40 OUR AVERAGE			
270±60	ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
240±60	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
410±90	² ANISOVICH	00J	SPEC 1.92–2.41 $p\bar{p}$

² From the combined analysis of ANISOVICH 99c, ANISOVICH 99f, ANISOVICH 99j, ANISOVICH 99k, and ANISOVICH 00b.

 $f_6(2510)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	(6.0±1.0) %

 $f_6(2510)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.06±0.01	³ BINON	83C	GAM2 38 $\pi^- p \rightarrow n 4\gamma$

³ Assuming one pion exchange and using data of BOLOTOV 74.

 $f_6(2510)$ REFERENCES

ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>
Also		PAN 62 405	D. Alde <i>et al.</i>
		Translated from YAF 62 446.	(GAM4 Collab.) (GAMS Collab.)
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>
BOLOTOV	74	Translated from YAF 38 1199.	V.N. Bolotov <i>et al.</i>
			(SERP, BELG, LAPP) JP (SERP, BRUX+) (SERP)

NODE=M033

REFID=48828
REFID=48327
REFID=48349
REFID=47950
REFID=44371
REFID=44103
REFID=21665
REFID=21838
REFID=21837
REFID=21964
REFID=21733
REFID=21963
REFID=21830
REFID=21813
REFID=21931
REFID=21926
REFID=21807
REFID=21925
REFID=21804

NODE=M089

NODE=M089M

NODE=M089M

NODE=M089M;LINKAGE=AN

NODE=M089W

NODE=M089W

NODE=M089W;LINKAGE=AN

NODE=M089215;NODE=M089

DESIG=1

NODE=M089220

NODE=M089R1
NODE=M089R1

NODE=M089R1;LINKAGE=A

NODE=M089

REFID=47942
REFID=47950
REFID=46903
REFID=46926
REFID=47416
REFID=47472
REFID=46605
REFID=46914
REFID=21780
REFID=40288
REFID=44705

OTHER LIGHT MESONS

Further States

OMITTED FROM SUMMARY TABLE

This section contains states observed by a single group or states poorly established that thus need confirmation.

QUANTUM NUMBERS, MASSES, WIDTHS, AND BRANCHING RATIOS

X(360)	$I^G(J^{PC}) = ?^?(?^+)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$360 \pm 7 \pm 9$	64 ± 18	2.3k	1 ABRAAMYAN 09	CNTR	$2.75 d C \rightarrow \gamma\gamma X$		

¹ Not seen in $pC \rightarrow \gamma\gamma X$ at 5.5 GeV/c.

NODE=MXXX015

X(1070)	$I^G(J^{PC}) = ?^?(0^++)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
1072 ± 1	3.5 ± 0.5	2 VLADIMIRSK...08		$40 \pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$	

² Supersedes GRIGOR'EV 05.

NODE=M300

NODE=M300

X(1110)	$I^G(J^{PC}) = 0^+(even++)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1107 ± 4	$111 \pm 8 \pm 15$	DAFTARI	87	DBC	$0. \bar{p}n \rightarrow \rho^- \pi^+ \pi^-$	

NODE=M300K08

NODE=M300K08

NODE=M300K08;LINKAGE=AB

NODE=M300J07

NODE=M300J07

NODE=M300J07;LINKAGE=VL

NODE=M300J30

NODE=M300J30

f₀(1200-1600)	$I^G(J^{PC}) = 0^+(0^++)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1323 ± 8	237 ± 20	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$		
1480^{+100}_{-150}	1030^{+80}_{-170}	3 ANISOVICH 03	SPEC			
1530^{+90}_{-250}	560 ± 40	4 ANISOVICH 03	SPEC			
3 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest. 4 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K} n$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$ at rest.						

NODE=M300J98

NODE=M300J98

OCCUR=2

NODE=M300;LINKAGE=KM

NODE=M300;LINKAGE=MK

X(1420)	$I^G(J^{PC}) = 2^+(0^++)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1420 ± 20	160 ± 10	FILIPPI	00	OBLX	$0 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$	

NODE=M300J61

NODE=M300J61

X(1545)	$I^G(J^{PC}) = ?^?(?^++)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
1545 ± 3	6.0 ± 2.5	5 VLADIMIRSK...08	$40 \pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$		

⁵ Supersedes VLADIMIRSKII 00.

NODE=M300K07

NODE=M300K07

NODE=M300K07;LINKAGE=VL

X(1575)	$I^G(J^{PC}) = ?^?(1^- -)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1576^{+49+98}_{-55-91}	$818^{+22+64}_{-23-133}$	6 ABLIKIM	06s	BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

NODE=M300J08

NODE=M300J08

⁶A broad peak observed at $K^+ K^-$ invariant mass. Mass and width above are its pole position. The observed branching ratio is $B(J/\psi \rightarrow X\pi^0) B(X \rightarrow K^+ K^-) = (8.5 \pm 0.6^{+2.7}_{-3.6}) \times 10^{-4}$.

X(1600)	$I^G(J^{PC}) = 2^+(2^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1600 \pm 100	400 \pm 200	7 ALBRECHT	91F ARG	10.2 $e^+ e^- \rightarrow e^+ e^- 2(\pi^+ \pi^-)$
7 Our estimate.				

X(1650)	$I^G(J^{PC}) = 0^-(?^-)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1652 \pm 7	<50	100	PROKOSHKIN 96	GAM2	32,38 $\pi p \rightarrow \omega \eta n$

X(1730)	$I^G(J^{PC}) = ??(?^?)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1731.0 \pm 1.2 \pm 2.0	3.2 \pm 0.8 \pm 1.3	58	VLADIMIRSK..07	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 X$

X(1750)	$I^G(J^{PC}) = ?^?(1^{--})$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1753.5 \pm 1.5 \pm 2.3	122.2 \pm 6.2 \pm 8.0	LINK	02K FOCS	20-160 $\gamma p \rightarrow K^+ K^- p$	

$B(X(1750) \rightarrow K^*(892)^0 K^0 \rightarrow K^\pm \pi^\mp K_S^0)/B(X(1750) \rightarrow K^+ K^-)$					
VALUE	CL%	DOCUMENT ID	TECN		
<0.065	90	LINK	02K FOCS		

$B(X(1750) \rightarrow K^*(892)^\pm K^\mp \rightarrow K_S^0 \pi^\pm K^\mp)/B(X(1750) \rightarrow K^+ K^-)$					
VALUE	CL%	DOCUMENT ID	TECN		
<0.183	90	LINK	02K FOCS		

f₂(1750)	$I^G(J^{PC}) = 0^+(2^{++})$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1755 \pm 10	67 \pm 12	870	8 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(K\bar{K})$	$I^G(J^{PC}) = 0^+(2^{++})$			
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
17 \pm 5	870	9 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(\gamma\gamma)$	$I^G(J^{PC}) = 0^+(2^{++})$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.13 \pm 0.04	870	9 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(\pi\pi)$	$I^G(J^{PC}) = 0^+(2^{++})$			
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 \pm 1.0	870	9 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

$\Gamma(\eta\eta)$	$I^G(J^{PC}) = 0^+(2^{++})$			
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.0 \pm 0.5	870	9 SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

⁸ From analysis of L3 data at 91 and 183–209 GeV.

⁹ From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

X(1775)	$I^G(J^{PC}) = 1^-(?^-)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1763 \pm 20	192 \pm 60	CONDO	91	SHF $\gamma p \rightarrow (p\pi^+)(\pi^+\pi^-\pi^-)$
1787 \pm 18	118 \pm 60	CONDO	91	SHF $\gamma p \rightarrow n\pi^+ \pi^+ \pi^-$

f₀(1800)	$I^G(J^{PC}) = 0^+(0^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1795 \pm 7 ⁺²³ ₋₂₀	95 \pm 10 ⁺⁷⁸ ₋₈₂	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma\omega\phi$
1812 \pm 18 ⁺¹⁹ ₋₂₆	105 \pm 20 \pm 28	10 ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma\omega\phi$

NODE=M300J08;LINKAGE=AB

NODE=M300J99
NODE=M300J99

NODE=M300J62
NODE=M300J62

NODE=M300K06
NODE=M300K06

NODE=M300J94
NODE=M300J94

NODE=M300B5
NODE=M300B5

NODE=M300B6
NODE=M300B6

NODE=M300JAM
NODE=M300JAM

NODE=M300JA1
NODE=M300JA1

NODE=M300JA2
NODE=M300JA2

NODE=M300JA3
NODE=M300JA3

NODE=M300JA4
NODE=M300JA4

NODE=M300JAM;LINKAGE=SC
NODE=M300JA;LINKAGE=SC

NODE=M300J60
NODE=M300J60

OCCUR=2

NODE=M300K29
NODE=M300K29

¹⁰ Not seen by LIU 09 in $B^\pm \rightarrow K^\pm \omega\phi$.

X(1850 - 3100)		$I^G(J^{PC}) = ?^?(1^- -)$		
$\Gamma(e^+ e^-) \cdot B(X \rightarrow \text{hadrons})$ (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<120	90	11 ANASHIN	11 KEDR	$e^+ e^- \rightarrow \text{hadrons}$

¹¹ This limit is center-of-mass energy dependent. We quote the most stringent one.

X(1855)		$I^G(J^{PC}) = ?^?(???)$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1856.6 ± 5	20 ± 5	BRIDGES	86D SPEC	0. $\bar{p}d \rightarrow \pi\pi N$

X(1870)		$I^G(J^{PC}) = ?^?(2??)$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1870 ± 40	250 ± 30	ALDE	86D GAM4	100 $\pi^- p \rightarrow 2\eta X$

a₃(1875)		$I^G(J^{PC}) = 1^-(3^{++})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1874 ± 43 ± 96	385 ± 121 ± 114	CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

B(a₃(1875) → f₂(1270)π)/B(a₃(1875) → ρπ)				
VALUE	DOCUMENT ID	TECN	COMMENT	
0.8 ± 0.2	12 CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

¹² Using the observable fractions of 50.0% $\rho\pi$, 56.5% $f_2\pi$, and 11.8% $\rho_3\pi$.

B(a₃(1875) → ρ₃(1690)π)/B(a₃(1875) → ρπ)				
VALUE	DOCUMENT ID	TECN	COMMENT	
0.9 ± 0.3	13 CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

¹³ Using the observable fractions of 50.0% $\rho\pi$, 56.5% $f_2\pi$, and 11.8% $\rho_3\pi$.

a₁(1930)		$I^G(J^{PC}) = 1^-(1^{++})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1930 ⁺³⁰ ₋₇₀	155 ± 45	ANISOVICH	01F SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

X(1935)		$I^G(J^{PC}) = 1^+(1^-?)$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1935 ± 20	215 ± 30	EVANGELIS... 79	OMEG	10,16 $\pi^- p \rightarrow \bar{p}pn$

ρ₂(1940)		$I^G(J^{PC}) = 1^+(2^{--})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1940 ± 40	155 ± 40	14 ANISOVICH	02 SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

¹⁴ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02Z.

ω₃(1945)		$I^G(J^{PC}) = 0^-(3^{--})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1945 ± 20	115 ± 22	15 ANISOVICH	02B SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

¹⁵ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

a₂(1950)		$I^G(J^{PC}) = 1^-(2^{++})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1950 ⁺³⁰ ₋₇₀	180 ⁺³⁰ ₋₇₀	16 ANISOVICH	01F SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K29;LINKAGE=AB

NODE=M300K28
NODE=M300K28

NODE=M300K28;LINKAGE=AN

NODE=M300J31
NODE=M300J31

NODE=M300J45
NODE=M300J45

NODE=M300J95
NODE=M300J95

NODE=M300B7
NODE=M300B7

NODE=M300B;LINKAGE=C1

NODE=M300B8
NODE=M300B8

NODE=M300B8;LINKAGE=C1

NODE=M300J92
NODE=M300J92

NODE=M300J33
NODE=M300J33

NODE=M300J85
NODE=M300J85

NODE=M300J85;LINKAGE=AY

NODE=M300J65
NODE=M300J65

NODE=M300J65;LINKAGE=AZ

NODE=M300K24
NODE=M300K24

16 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

$\omega(1960)$	$I^G(J^{PC}) = 0^-(1^{--})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1960 \pm 25	195 \pm 60	17	ANISOVICH	02B	SPEC	$0.6\text{--}1.9 \ p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

17 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

$b_1(1960)$	$I^G(J^{PC}) = 1^+(1^{+-})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1960 \pm 35	230 \pm 50	18	ANISOVICH	02	SPEC	$0.6\text{--}1.9 \ p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

18 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

$h_1(1965)$	$I^G(J^{PC}) = 0^-(1^{+-})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1965 \pm 45	345 \pm 75	19	ANISOVICH	02B	SPEC	$0.6\text{--}1.9 \ p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

19 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

$f_1(1970)$	$I^G(J^{PC}) = 0^+(1^{++})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1971 \pm 15	240 \pm 45			ANISOVICH	00J	SPEC

$X(1970)$	$I^G(J^{PC}) = ??(???)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1970 \pm 10	40 \pm 20			CHLIAPNIK...	80	HBC $32 \ K^+ p \rightarrow 2K_S^0 2\pi X$

$X(1975)$	$I^G(J^{PC}) = ??(???)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1973 \pm 15	80			30	CASO	70	HBC $11.2 \ \pi^- p \rightarrow \rho 2\pi$

$\omega_2(1975)$	$I^G(J^{PC}) = 0^-(2^{--})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1975 \pm 20	175 \pm 25	20	ANISOVICH	02B	SPEC	$0.6\text{--}1.9 \ p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

20 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

$a_2(1990)$	$I^G(J^{PC}) = 1^-(2^{++})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2050 \pm 10 \pm 40	190 \pm 22 \pm 100	18k	21	SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
2003 \pm 10 \pm 19	249 \pm 23 \pm 32			LU	05	B852	$18 \ \pi^- p \rightarrow \omega\pi^-\pi^0 p$

21 From analysis of L3 data at 183–209 GeV.

$\Gamma(\gamma\gamma) \Gamma(\pi^+\pi^-\pi^0) / \Gamma(\text{total})$						
<u>VALUE (keV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.11 \pm 0.04 \pm 0.05	18k	22 SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$		

22 From analysis of L3 data at 183–209 GeV.

$\rho(2000)$	$I^G(J^{PC}) = 1^+(1^{--})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2000 \pm 30	260 \pm 45	23 BUGG		04C	RVUE	Compilation
~ 1988	~ 244	HASAN		94	RVUE	$\bar{p}p \rightarrow \pi\pi$

NODE=M300K24;LINKAGE=AN

NODE=M300J79
NODE=M300J79

NODE=M300J67
NODE=M300J67

NODE=M300J67;LINKAGE=AY

NODE=M300J64
NODE=M300J64

NODE=M300J1
NODE=M300J1

NODE=M300J46
NODE=M300J46

NODE=M300J47
NODE=M300J47

NODE=M300J81
NODE=M300J81

NODE=M300J81;LINKAGE=AZ

NODE=M300J2
NODE=M300J2

NODE=M300J2;LINKAGE=SC

NODE=M300J2G
NODE=M300J2G

NODE=M300J2G;LINKAGE=SC

NODE=M300J77
NODE=M300J77

²³ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

f₂(2000)	$I^G(JPC) = 0^+(2^{++})$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2001 \pm 10	312 \pm 32	ANISOVICH 00J	SPEC		
\sim 1996	\sim 134	HASAN 94	RVUE	$\bar{p}p \rightarrow \pi\pi$	

NODE=M300;LINKAGE=AY

X(2000)	$I^G(JPC) = 1^-(?^+)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
1964 \pm 35	225 \pm 50	24 ARMSTRONG 93D	E760		$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
\sim 2100	\sim 500	24 ANTIPOV 77	CIBS	-	$25\pi^- p \rightarrow p\pi^-\rho_3$
2214 \pm 15	355 \pm 21	25 BALTAY 77	HBC	0	$15\pi^- p \rightarrow \Delta^{++}3\pi$
2080 \pm 40	340 \pm 80	KALELKAR 75	HBC	+	$15\pi^+ p \rightarrow p\pi^+\rho_3$

²⁴ Cannot determine spin to be 3.²⁵ BALTAY 77 favors $J^P = ,3^+$.NODE=M300J25
NODE=M300J25

X(2000)	$I^G(JPC) = ?(4^{++})$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1998 \pm 3 \pm 5	<15	VLADIMIRSK..03	SPEC	$\pi^- p \rightarrow K_S^0 K_S^0 MM$	

NODE=M300K01
NODE=M300K01

η(2010)	$I^G(JPC) = 0^+(0^-+)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2010 $^{+35}_{-60}$	270 \pm 60	ANISOVICH 00J	SPEC		

NODE=M300J97
NODE=M300J97

π_1(2015)	$I^G(JPC) = 1^-(1^-+)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2014 \pm 20 \pm 16	230 \pm 32 \pm 73	145k	LU 05	B852	$18\pi^- p \rightarrow \omega\pi^-\pi^0 p$
2001 \pm 30 \pm 92	333 \pm 52 \pm 49	69k	KUHN 04	B852	$18\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M300J5
NODE=M300J5

a₀(2020)	$I^G(JPC) = 1^-(0^{++})$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2025 \pm 30	330 \pm 75	ANISOVICH 99C	SPEC		

NODE=M300J6
NODE=M300J6

X(2020)	$I^G(JPC) = ??(???)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2015 \pm 3	10 \pm 4	FERRER 99	RVUE	$\pi p \rightarrow p\bar{p}\pi(\pi)$	

NODE=M300J34
NODE=M300J34

h_3(2025)	$I^G(JPC) = 0^-(3^{+-})$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2025 \pm 20	145 \pm 30	26 ANISOVICH 02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$	

²⁶ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.NODE=M300J78
NODE=M300J78

b_3(2030)	$I^G(JPC) = 1^+(3^{+-})$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2032 \pm 12	117 \pm 11	27 ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$	

²⁷ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.NODE=M300J69
NODE=M300J69

a_2(2030)	$I^G(JPC) = 1^-(2^{++})$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2030 \pm 20	205 \pm 30	28 ANISOVICH 01F	SPEC	1.96–2.41 $\bar{p}p$	

NODE=M300K23
NODE=M300K23

28 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

$a_3(2030)$	$I^G(J^{PC}) = 1^-(3^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2031 \pm 12	150 \pm 18	29 ANISOVICH	01F SPEC	1.96–2.41 $p\bar{p}$
29 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.				

$\eta_2(2030)$	$I^G(J^{PC}) = 0^+(2^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	
2030 \pm 5 \pm 15	205 \pm 10 \pm 15	ANISOVICH	00E SPEC	

$B(a_2\pi)_{L=0}/B(a_2\pi)_{L=2}$				
VALUE	DOCUMENT ID	TECN	COMMENT	
0.05 \pm 0.03	30 ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$

30 Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

$B(a_0\pi)/B(a_2\pi)_{L=2}$				
VALUE	DOCUMENT ID	TECN	COMMENT	
0.10 \pm 0.08	31 ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$

31 Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

$B(f_2\eta)/B(a_2\pi)_{L=2}$				
VALUE	DOCUMENT ID	TECN	COMMENT	
0.13 \pm 0.06	32 ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$

32 Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

$f_3(2050)$	$I^G(J^{PC}) = 0^+(3^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2048 \pm 8	213 \pm 34	ANISOVICH	00J SPEC	2.0 $p\bar{p} \rightarrow \eta\pi^0\pi^0$

$f_0(2060)$	$I^G(J^{PC}) = 0^+(0^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
\sim 2050	\sim 120	33 OAKDEN	94 RVUE	0.36 – 1.55 $p\bar{p} \rightarrow \pi\pi$
\sim 2060	\sim 50	33 OAKDEN	94 RVUE	0.36 – 1.55 $p\bar{p} \rightarrow \pi\pi$

33 See SEMENOV 99 and KLOET 96.

$\pi(2070)$	$I^G(J^{PC}) = 1^-(0^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2070 \pm 35	310^{+100}_{-50}	ANISOVICH	01F SPEC	$2.0 p\bar{p} \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

$X(2075)$	$I^G(J^{PC}) = ?(?)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2075 \pm 12 \pm 5 $90 \pm 35 \pm 9$ 34 ABLIKIM 04J BES2 $J/\psi \rightarrow K^-\rho\Lambda$				
34 From a fit in the region $M_{p\bar{\Lambda}} - M_p - M_\Lambda < 150$ MeV. S-wave in the $p\bar{\Lambda}$ system preferred.				

A similar near-threshold enhancement in the $p\bar{\Lambda}$ system is observed in $B^+ \rightarrow p\bar{\Lambda}\bar{D}^0$ by CHEN 11F.

$X(2080)$	$I^G(J^{PC}) = ?(?)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2080 \pm 10	110 \pm 20	KREYMER	80 STRC	$13 \pi^- d \rightarrow p\bar{p}n(n_s)$

$X(2080)$	$I^G(J^{PC}) = ?(3^{-?})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2080 \pm 10	190 \pm 15	ROZANSKA	80 SPRK	$18 \pi^- p \rightarrow p\bar{p}n$

$a_1(2095)$	$I^G(J^{PC}) = 1^-(1^{++})$			
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN COMMENT
2096 \pm 17 \pm 121	451 \pm 41 \pm 81	69k	KUHN	04 B852 $18 \pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M300K23;LINKAGE=AN

NODE=M300K20
NODE=M300K20

NODE=M300J8
NODE=M300J8

NODE=M300B1
NODE=M300B1

NODE=M300B1;LINKAGE=AN

NODE=M300B2
NODE=M300B2

NODE=M300B2;LINKAGE=AN

NODE=M300B3
NODE=M300B3

NODE=M300B3;LINKAGE=AN

NODE=M300J7
NODE=M300J7

NODE=M300J59
NODE=M300J59

OCCUR=2
NODE=M300J;LINKAGE=A

NODE=M300J91
NODE=M300J91

NODE=M300J01
NODE=M300J01

NODE=M300J01;LINKAGE=AB

NODE=M300J35
NODE=M300J35

NODE=M300J37
NODE=M300J37

NODE=M300J04
NODE=M300J04

B(a₁(2095) → f₁(1285)π) / B(a₁(2095) → a₁(1260))				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
3.18±0.64	69k	KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M300B03
NODE=M300B03

$\eta(2100)$	$I^G(J^{PC}) = 0^+(0^-+)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2050 _{-24 -26} ^{+30 +75}	250 _{-30 -164} ^{+36 +181}	35	ABLIKIM	16N	BES3 $J/\psi \rightarrow \gamma K^+$ $K^- K^+ K^-$
2103±50	187 ± 75	586	36 BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
35	From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$, for which the primary signal is $\eta(2225) \rightarrow \phi\phi$, and that also finds significant signals for for 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).				
36	ASTON 81B sees no peak, has 850 events in Ajinenko+Barth bins. ARESTOV 80 sees no peak.				

NODE=M300J48
NODE=M300J48

$X(2100)$	$I^G(J^{PC}) = ?^?(0??)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2100±40	250 ± 40		ALDE	86D	GAM4 100 $\pi^- p \rightarrow 2\eta X$

NODE=M300J49
NODE=M300J49

$X(2110)$	$I^G(J^{PC}) = 1^+(3-?)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2110±10	330 ± 20		EVANGELIS...	79	OMEG 10,16 $\pi^- p \rightarrow \bar{p}pn$

NODE=M300J36
NODE=M300J36

$f_2(2140)$	$I^G(J^{PC}) = 0^+(2++)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2141±12	49 ± 28	389	GREEN	86	MPSF 400 pA → 4KX

NODE=M300J50
NODE=M300J50

$X(2150)$	$I^G(J^{PC}) = ?^?(2+?)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2150±10	260 ± 10		ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J38
NODE=M300J38

$a_2(2175)$	$I^G(J^{PC}) = 1^-(2++)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2175±40	310 ₋₄₅ ⁺⁹⁰		ANISOVICH	01F	SPEC 2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J88
NODE=M300J88

$\eta(2190)$	$I^G(J^{PC}) = 0^+(0^-+)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2190±50	850 ± 100		BUGG	99	BES

NODE=M300J13
NODE=M300J13

$\omega_2(2195)$	$I^G(J^{PC}) = 0^-(2--)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2195±30	225 ± 40	37	ANISOVICH	02B	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$
37	From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.				

NODE=M300J82
NODE=M300J82

NODE=M300J82;LINKAGE=AZ

$\omega(2205)$	$I^G(J^{PC}) = 0^-(1--)$				
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2205±30	350 ± 90	38	ANISOVICH	02B	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J80
NODE=M300J80

38 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

X(2210)	$I^G(J^{PC}) = ?^?(???)$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2210 $^{+79}_{-21}$	203 $^{+437}_{-87}$	EVANGELIS...	79B OMEG	$10 \pi^- p \rightarrow K^+ K^- n$

X(2210)	$I^G(J^{PC}) = ?^?(???)$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2207 ± 22	130	CASO	70 HBC	$11.2 \pi^- p$

h₁(2215)	$I^G(J^{PC}) = 0^-(1^{+-})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2215 ± 40	325 ± 55	39 ANISOVICH	02B SPEC	$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

39 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

$\rho_2(2225)$	$I^G(J^{PC}) = 1^+(2^{--})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2225 ± 35	335^{+100}_{-50}	40 ANISOVICH	02 SPEC	$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

40 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

$\rho_4(2230)$	$I^G(J^{PC}) = 1^+(4^{--})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2230 ± 25	210 ± 30	41 ANISOVICH	02 SPEC	$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

41 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

$b_1(2240)$	$I^G(J^{PC}) = 1^+(1^{+-})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2240 ± 35	320 ± 85	42 ANISOVICH	02 SPEC	$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

42 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

$f_2(2240)$	$I^G(J^{PC}) = 0^+(2^{++})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2240 ± 15	241 ± 30	43 ANISOVICH	00J SPEC	$1.92\text{--}2.41 p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 2226 \quad \sim 226 \quad \text{HASAN} \quad 94 \quad \text{RVUE} \quad p\bar{p} \rightarrow \pi\pi$

43 From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

$b_3(2245)$	$I^G(J^{PC}) = 1^+(3^{+-})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2245 ± 50	320 ± 70	44 BUGG	04C RVUE	

44 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

$\eta_2(2250)$	$I^G(J^{PC}) = 0^+(2^{-+})$	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>			
2248 ± 20	280 ± 20	ANISOVICH	00I SPEC	
2267 ± 14	290 ± 50	ANISOVICH	00J SPEC	

NODE=M300J80;LINKAGE=AZ

NODE=M300J39
NODE=M300J39

NODE=M300J51
NODE=M300J51

NODE=M300J27
NODE=M300J27

NODE=M300J27;LINKAGE=AZ

NODE=M300J70
NODE=M300J70

NODE=M300J74
NODE=M300J74

NODE=M300J74;LINKAGE=AY

NODE=M300K26
NODE=M300K26

NODE=M300K26;LINKAGE=AN

NODE=M300K10
NODE=M300K10

NODE=M300K10;LINKAGE=AY

NODE=M300J17
NODE=M300J17

$\pi_4(2250)$	$I^G(J^{PC}) = 1^-(4 - +)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 15	215 ± 25	ANISOVICH	01F	SPEC 2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J73
NODE=M300J73

$\omega_4(2250)$	$I^G(J^{PC}) = 0^-(4 --)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 30	150 ± 50	45	ANISOVICH	02B SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$
45 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.				

NODE=M300J84
NODE=M300J84

NODE=M300J84;LINKAGE=AZ

$\omega_5(2250)$	$I^G(J^{PC}) = 0^-(5 --)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 70	320 ± 95	46	BUGG	04 RVUE
46 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.				

NODE=M300K11
NODE=M300K11

NODE=M300;LINKAGE=AZ

$\omega_3(2255)$	$I^G(J^{PC}) = 0^-(3 --)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255 ± 15	175 ± 30	47	ANISOVICH	02B SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$
47 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.				

NODE=M300J66
NODE=M300J66

NODE=M300J66;LINKAGE=AZ

$a_4(2255)$	$I^G(J^{PC}) = 1^-(4 ++)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2237 ± 5	291 ± 12	UMAN	06	E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
2255 ± 40	330 ± 110	48	ANISOVICH	01F SPEC 1.96–2.41 $\bar{p}p$
48 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.				

NODE=M300K21
NODE=M300K21

NODE=M300K21;LINKAGE=AN

$a_2(2255)$	$I^G(J^{PC}) = 1^-(2 ++)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255 ± 20	230 ± 15	49	ANISOVICH	01G SPEC 1.96–2.41 $\bar{p}p$
49 From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.				

NODE=M300K22
NODE=M300K22

NODE=M300K22;LINKAGE=AN

$X(2260)$	$I^G(J^{PC}) = 0^+(4+?)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2260 ± 20	400 ± 100	EVANGELIS...	79	OMEG 10,16 $\pi^- p \rightarrow \bar{p}pn$

NODE=M300J40
NODE=M300J40

$\rho(2270)$	$I^G(J^{PC}) = 1^+(1 --)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2265 ± 40	325 ± 80	50	ANISOVICH	02 SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
2280 ± 50	440 ± 110	ATKINSON	85	OMEG 20–70 $\gamma p \rightarrow p\omega\pi^+\pi^-\pi^0$
50 From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.				

NODE=M300J86
NODE=M300J86

NODE=M300J86;LINKAGE=AY

$a_1(2270)$	$I^G(J^{PC}) = 1^-(1 ++)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2270 ± 55	305 ± 70	ANISOVICH	01F	SPEC 2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J72
NODE=M300J72

$h_3(2275)$	$I^G(J^{PC}) = 0^-(3 +-)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2275 ± 25	190 ± 45	51	ANISOVICH	02B SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J28
NODE=M300J28

NODE=M300J28;LINKAGE=AZ

$a_3(2275)$	$I^G(J^{PC}) = 1^-(3 ++)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2275 ± 35	350 ± 100	52	ANISOVICH	01G SPEC 1.96–2.41 $\bar{p}p$

NODE=M300K19
NODE=M300K19

52 From the combined analysis of ANISOVICH 99c, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

$\pi_2(2285)$	$I^G(JPC) = 1^-(2^-+)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2285 \pm 20 \pm 25$	$250 \pm 20 \pm 25$	53	ANISOVICH	11	SPEC	$0.9\text{--}1.94 p\bar{p}$

53 Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

$\omega_3(2285)$	$I^G(JPC) = 0^-(3^- -)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2278 ± 28	224 ± 50	54	BUGG	04A	RVUE	
2285 ± 60	230 ± 40	55	ANISOVICH	02B	SPEC	$0.6\text{--}1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

54 Partial wave analysis of the data on $p\bar{p} \rightarrow \Lambda\Lambda$ from BARNES 00.

55 From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

$\omega(2290)$	$I^G(JPC) = 0^-(1^- -)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
2290 ± 20	275 ± 35	56	BUGG	04A	RVUE

56 Partial wave analysis of the data on $p\bar{p} \rightarrow \Lambda\Lambda$ from BARNES 00.

$f_2(2295)$	$I^G(JPC) = 0^+(2^++)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2293 ± 13	216 ± 37	57	ANISOVICH	00J	SPEC	$1.92\text{--}2.41 p\bar{p}$

57 From the combined analysis of ANISOVICH 99c, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

$f_3(2300)$	$I^G(JPC) = 0^+(3^{++})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
2334 ± 25	200 ± 20	58	BUGG	04A	RVUE

58 Partial wave analysis of the data on $p\bar{p} \rightarrow \Lambda\Lambda$ from BARNES 00.

$f_1(2310)$	$I^G(JPC) = 0^+(1^{++})$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
2310 ± 60	255 ± 70	ANISOVICH	00J	SPEC	

$\eta(2320)$	$I^G(JPC) = 0^+(0^- +)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
2320 ± 15	230 ± 35	59	ANISOVICH	00M	SPEC

59 From the combined analysis of $\bar{p}p \rightarrow \eta\eta\eta$ from ANISOVICH 00M and $\bar{p}p \rightarrow \eta\pi^0\pi^0$ from ANISOVICH 00J.

$\eta_4(2330)$	$I^G(JPC) = 0^+(4^- +)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2328 ± 38	240 ± 90	ANISOVICH	00J	SPEC		$2.0 p\bar{p} \rightarrow \eta\pi^0\pi^0$

$\omega(2330)$	$I^G(JPC) = 0^-(1^- -)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2330 ± 30	435 ± 75	ATKINSON	88	OMEG		$25\text{--}50 \gamma p \rightarrow \rho^\pm \rho^0 \pi^\mp$

$X(2340)$	$I^G(JPC) = ??(???)$	<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2340 ± 20	180 ± 60	126	60	BALTAY	75	HBC	$15 \pi^+ p \rightarrow p\bar{p}$

NODE=M300K19;LINKAGE=AN

NODE=M300K25
NODE=M300K25

NODE=M300K25;LINKAGE=AN

NODE=M300J83
NODE=M300J83

NODE=M300J83;LINKAGE=BU
NODE=M300J83;LINKAGE=AZ

NODE=M300J02
NODE=M300J02

NODE=M300J02;LINKAGE=BU

NODE=M300K27
NODE=M300K27

NODE=M300K27;LINKAGE=AN

NODE=M300J19
NODE=M300J19

NODE=M300J19;LINKAGE=BU

NODE=M300J23
NODE=M300J23

NODE=M300J18
NODE=M300J18

NODE=M300;LINKAGE=B

NODE=M300J22
NODE=M300J22

NODE=M300J53
NODE=M300J53

NODE=M300J54
NODE=M300J54

60 Dominant decay into $\rho^0 \rho^0 \pi^+$. BALTAY 78 finds confirmation in $2\pi^+ \pi^- 2\pi^0$ events which contain $\rho^+ \rho^0 \pi^0$ and $2\rho^+ \pi^-$.

$\pi(2360)$	$I^G(J^{PC}) = 1^-(0 - +)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2360 ± 25	300^{+100}_{-50}	ANISOVICH	01F	SPEC	$2.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J;LINKAGE=B1

$X(2360)$	$I^G(J^{PC}) = ??(4+?)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2360 ± 10	430 ± 30	ROZANSKA	80	SPRK	$18 \pi^- p \rightarrow p\bar{p}n$

NODE=M300J42
NODE=M300J42

$X(2440)$	$I^G(J^{PC}) = ??(5-?)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2440 ± 10	310 ± 20	ROZANSKA	80	SPRK	$18 \pi^- p \rightarrow p\bar{p}n$

NODE=M300J43
NODE=M300J43

$a_6(2450)$	$I^G(J^{PC}) = 1^-(6 + +)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2450 ± 130	400 ± 250	CLELAND	82B	SPEC	$50 \pi p \rightarrow K_S^0 K^\pm p$

NODE=M300K12
NODE=M300K12

$X(2540)$	$I^G(J^{PC}) = 0^+(0 + +)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2539 \pm 14^{+38}_{-14}$	$274^{+77+126}_{-61-163}$	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300K30
NODE=M300K30

$\Gamma(\gamma\gamma) \times B(K\bar{K})$					
<u>VALUE (eV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
40^{+9+17}_{-7-40}		UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300K3G
NODE=M300K3G

$X(2632)$	$I^G(J^{PC}) = ??(???)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2635.2 ± 3.3		61 EVDOKIMOV	04	SELX	$X(2632) \rightarrow D_s^+ \eta$
2631.6 ± 2.1	< 17	62 EVDOKIMOV	04	SELX	$X(2632) \rightarrow D^0 K^+$

NODE=M300J03
NODE=M300J03

$B(X(2632) \rightarrow D^0 K^+)/B(X(2632) \rightarrow D_s^+ \eta)$				
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	
0.14 ± 0.06		63 EVDOKIMOV	04	SELX

OCCUR=2

NODE=M300J03;LINKAGE=EV
NODE=M300J03;LINKAGE=ED

$X(2680)$	$I^G(J^{PC}) = ??(???)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2676 ± 27	150	CASO	70	HBC	$11.2 \pi^- p \rightarrow \rho^- \pi^+ \pi^- p$

NODE=M300J55
NODE=M300J55

$X(2710)$	$I^G(J^{PC}) = ??(6+?)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2710 ± 20	170 ± 40	ROZANSKA	80	SPRK	$18 \pi^- p \rightarrow p\bar{p}n$

NODE=M300J44
NODE=M300J44

$X(2750)$	$I^G(J^{PC}) = ??(7-?)$				
<u>MASS (MeV)</u>	<u>WIDTH (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2747 ± 32	195 ± 75	DENNEY	83	LASS	$10 \pi^+ p \rightarrow K^+ K^- \pi^+ p$

NODE=M300J56
NODE=M300J56

f₆(3100)	<i>I^G(JPC) = 0⁺(6⁺⁺)</i>	<i>MASS (MeV)</i>	<i>WIDTH (MeV)</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	NODE=M300J06 NODE=M300J06	
3100±100	700 ± 130	BINON	05	GAMS	33	$\pi^- p \rightarrow \eta\eta n$		
X(3250)	<i>I^G(JPC) = ??(???) 3-Body Decays</i>	<i>MASS (MeV)</i>	<i>WIDTH (MeV)</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	NODE=M300J57 NODE=M300J57	
3250±8±20	45 ± 18	ALEEV	93	BIS2	$X(3250) \rightarrow \Lambda\bar{p}K^+$			
3265±7±20	40 ± 18	ALEEV	93	BIS2	$X(3250) \rightarrow \bar{\Lambda}pK^-$	OCCUR=2		
X(3250)	<i>I^G(JPC) = ??(???) 4-Body Decays</i>	<i>MASS (MeV)</i>	<i>WIDTH (MeV)</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	NODE=M300J58 NODE=M300J58	
3245±8±20	25 ± 11	ALEEV	93	BIS2	$X(3250) \rightarrow \Lambda\bar{p}K^+\pi^\pm$			
3250±9±20	50 ± 20	ALEEV	93	BIS2	$X(3250) \rightarrow \bar{\Lambda}pK^-\pi^\mp$	OCCUR=2		
3270±8±20	25 ± 11	ALEEV	93	BIS2	$X(3250) \rightarrow K_S^0 p\bar{p}K^\pm$	OCCUR=3		
X(3350)	<i>I^G(JPC) = ??(???)</i>	<i>MASS (MeV)</i>	<i>WIDTH (MeV)</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	NODE=M300J09 NODE=M300J09
3350 ⁺¹⁰ ₋₂₀ ±20	70 ⁺⁴⁰ ₋₃₀ ± 40	50 ± 10	64	GABYSHEV	06A	BELL	$B^- \rightarrow \Lambda_c^+ \bar{p}\pi^-$	
64 A similar enhancement in the $\Lambda_c^+ \bar{p}$ final state is also reported by BABAR collaboration in AUBERT 10H.							NODE=M300J09;LINKAGE=AU	
REFERENCES for Further States							NODE=M300	
ABLIKIM 16N	PR D93 112011	M. Ablikim				(BESIII Collab.)	REFID=57512	
ABLIKIM 13J	PR D87 032008	M. Ablikim <i>et al.</i>				(BESIII Collab.)	REFID=54955	
UEHARA 13	PTEP 2013 123C01	S. Uehara <i>et al.</i>				(BELLE Collab.)	REFID=55592	
ANISOVICH 12	PR D85 014001	A.V. Anisovich <i>et al.</i>					REFID=53961	
ANASHIN 11	PL B703 543	V.V. Anashin <i>et al.</i>				(KEDR Collab.)	REFID=53932	
ANISOVICH 11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>				(LOQM, RAL, PNPI)	REFID=53631	
CHEN 11F	PR D84 071501	P. Chen <i>et al.</i>				(BELLE Collab.)	REFID=53814	
AUBERT 10H	PR D82 031102	B. Aubert <i>et al.</i>				(BABAR Collab.)	REFID=53363	
ABRAMANYAN 09	PR C80 034001	Kh.U. Abraamyan <i>et al.</i>					REFID=53100	
LIU 09	PR D79 071102	C. Liu <i>et al.</i>				(BELLE Collab.)	REFID=52752	
VLADIMIRSK... 08	PAN 71 2129	V.V. Vladimirska <i>et al.</i>				(ITEP)	REFID=52681	
VLADIMIRSK... 07	PAN 70 1706	V. Vladimirska <i>et al.</i>					REFID=52058	
VLADIMIRSK... 07	Translated from YAF 71 2166.							
YASUI 07	PR D76 034009	S. Yasui, M. Oka					REFID=51907	
ABLIKIM 06J	PRL 96 162002	M. Ablikim <i>et al.</i>				(BES Collab.)	REFID=51127	
ABLIKIM 06S	PRL 97 142002	M. Ablikim <i>et al.</i>				(BES Collab.)	REFID=51451	
GABYSHEV 06A	PRL 97 242001	N. Gabyshev <i>et al.</i>				(BELLE Collab.)	REFID=51565	
SCHEGELSKY 06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>					REFID=51186	
SCHEGELSKY 06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>					REFID=51185	
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>				(FNAL E835)	REFID=51063	
VLADIMIRSK... 06	PAN 69 493	V.V. Vladimirska <i>et al.</i>				(ITEP, Moscow)	REFID=51191	
BINON 05	PAN 68 960	F. Binon <i>et al.</i>					REFID=50780	
GRIGOR'EV 05	PAN 68 1271	V.K. Grigor'ev <i>et al.</i>				(ITEP)	REFID=50844	
GRIGOR'EV 05	Translated from YAF 68 998.							
LU 05	PRL 94 032002	M. Lu <i>et al.</i>				(BNL E852 Collab.)	REFID=50459	
ABLIKIM 04J	PRL 93 112002	M. Ablikim <i>et al.</i>				(BES Collab.)	REFID=50196	
BUGG 04	PL B595 556 (errat.)	D.V. Bugg					REFID=49763	
BUGG 04A	EPJ C36 161	D.V. Bugg					REFID=50158	
BUGG 04C	PRPL 397 257	D.V. Bugg					REFID=50203	
EVDOKIMOV 04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>				(SELEX Collab.)	REFID=50337	
KUHN 04	PL B595 109	J. Kuhn <i>et al.</i>				(BNL E852 Collab.)	REFID=49773	
ANISOVICH 03	EPJ A16 229	V.V. Anisovich <i>et al.</i>					REFID=49401	
VLADIMIRSK... 03	PAN 66 700	V.V. Vladimirska <i>et al.</i>					REFID=49419	
ANISOVICH 02	PL B542 8	A.V. Anisovich <i>et al.</i>					REFID=48828	
ANISOVICH 02B	PL B542 19	A.V. Anisovich <i>et al.</i>					REFID=48829	
CHUNG 02	PR D65 072001	S.U. Chung <i>et al.</i>					REFID=48837	
LINK 02K	PL B545 50	J.M. Link <i>et al.</i>					REFID=48845	
ANISOVICH 01C	PL B507 23	A.V. Anisovich <i>et al.</i>					REFID=48325	
ANISOVICH 01D	PL B508 6	A.V. Anisovich <i>et al.</i>					REFID=48327	
ANISOVICH 01E	PL B513 281	A.V. Anisovich <i>et al.</i>					REFID=48349	
ANISOVICH 01F	PL B517 261	A.V. Anisovich <i>et al.</i>					REFID=48352	
ANISOVICH 01G	PL B517 273	A.V. Anisovich <i>et al.</i>					REFID=48353	
ANISOVICH 00B	NP A662 319	A.V. Anisovich <i>et al.</i>					REFID=47942	
ANISOVICH 00D	PL B476 15	A.V. Anisovich <i>et al.</i>					REFID=47944	
ANISOVICH 00E	PL B477 19	A.V. Anisovich <i>et al.</i>					REFID=47945	
ANISOVICH 00I	PL B491 40	A.V. Anisovich <i>et al.</i>					REFID=47949	
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>					REFID=47950	
ANISOVICH 00M	PL B496 145	A.V. Anisovich <i>et al.</i>					REFID=48009	
BARNES 00	PR C62 055203	P.D. Barnes <i>et al.</i>					REFID=47965	
FILIPPI 00	PL B495 284	A. Filippi <i>et al.</i>					REFID=48006	
VLADIMIRSKII 00	JETPL 72 486	V.V. Vladimirska <i>et al.</i>					REFID=47997	
	Translated from ZETFP 72 698.							

ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>	REFID=46903	
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>	REFID=46902	
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>	REFID=46926	
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>	REFID=47416	
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>	REFID=47472	
BUGG	99	PL B458 511	D.V. Bugg <i>et al.</i>	REFID=46938	
FERRER	99	EPJ C10 249	A. Ferrer <i>et al.</i>	REFID=47404	
SEmenov	99	SPU 42 847	S.V. Semenov	REFID=47363	
		Translated from UFN 42 937.			
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
PROKOSHIN	96	PD 41 247	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=45182
		Translated from DANS 348 481.			
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALEEV	93	PAN 56 1358	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)	REFID=43668
		Translated from YAF 56 100.			
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43596
ALBRECHT	91F	ZPHY C50 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41658
CONDO	91	PR D43 2787	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=41588
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
ATKINSON	88	ZPHY C38 535	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=40556
DAFTARI	87	PRU 58 859	I.K. Daftari <i>et al.</i>	(SYRA)	REFID=40412
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
BRIDGES	86D	PL B180 313	D.L. Bridges <i>et al.</i>	(SYRA, BNL, CASE+)	REFID=21984
GREEN	86	PRL 56 1639	D.R. Green <i>et al.</i>	(FNAL, ARIZ, FSU+)	REFID=21872
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=22000
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
ASTON	81B	NP B189 205	D. Aston <i>et al.</i>	(BONN, CERN, EPOL, GLAS+)	REFID=11553
ARESTOV	80	IHEP 80-165	Y.I. Arestov <i>et al.</i>	(SERP)	REFID=22312
CHLIAPNIK...	80	ZPHY C3 285	P.V. Chliapnikov <i>et al.</i>	(SERP, BRUX, MONS)	REFID=21996
KREYMER	80	PR D22 36	A.E. Kreymer <i>et al.</i>	(IND, PURD, SLAC+)	REFID=21970
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)	REFID=21774
EVANGELIS...	79	NP B153 253	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21966
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21967
BALTAY	78	PR D17 52	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21569
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU)	REFID=20847
BALTAY	75	PRL 35 891	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21994
KALELKAR	75	Thesis Nevis 207	M.S. Kalelkar	(COLU)	REFID=21564
CASO	70	LNC 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20590

STRANGE MESONS ($S = \pm 1, C = B = 0$)

$K^+ = u\bar{s}$, $K^0 = d\bar{s}$, $\bar{K}^0 = \bar{d}s$, $K^- = \bar{u}s$, similarly for K^* 's

$K_0^*(700)$

$I(J^P) = \frac{1}{2}(0^+)$

also known as κ ; was $K_0^*(800)$

Needs confirmation. See the mini-review on scalar mesons under $f_0(500)$ (see the index for the page number).

NODE=MXXX020

NODE=MXXX020

NODE=M174

NODE=M174

NODE=M174TMP

NODE=M174TMP

→ UNCHECKED ←

$K_0^*(700)$ T-Matrix Pole \sqrt{s}

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(630–730) – i (260–340) OUR EVALUATION			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
(670 ± 18) – i (295 ± 28)	1 PELAEZ	17 RVUE	
(764 ± 63 ⁺⁷¹ ₋₅₄) – i (306 ± 149 ⁺¹⁴³ ₋₈₅)	2 ABLIKIM	11B BES2	1.3k $J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
(665 ± 9) – i (268 ⁺²¹ ₋₆)	3 GUO	11B RVUE	
(849 ± 77 ⁺¹⁸ ₋₁₄) – i (256 ± 40 ⁺⁴⁶ ₋₂₂)	2 ABLIKIM	10E BES2	1.4k $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
(663 ± 8 ± 34) – i (329 ± 5 ± 22)	4 BUGG	10 RVUE	S-matrix pole
(706.0 ± 1.8 ± 22.8) – i (319.4 ± 2.2 ± 20.2)	5 BONVICINI	08A CLEO	141k $D^+ \rightarrow K^- \pi^+ \pi^+$
(841 ± 30 ⁺⁸¹ ₋₇₃) – i (309 ± 45 ⁺⁴⁸ ₋₇₂)	2 ABLIKIM	06C BES2	25k $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
(750 ⁺³⁰ ₋₅₅) – i (342 ± 60)	6 BUGG	06 RVUE	
(658 ± 13) – i (279 ± 12)	7 DESCOTES-G..06	RVUE	$\pi K \rightarrow \pi K$

(757 ± 33) – <i>i</i> (279 ± 41)	8 GUO	06 RVUE	
(694 ± 53) – <i>i</i> (303 ± 30)	9 ZHOU	06 RVUE	$K p \rightarrow K^- \pi^+ n$
(754 ± 22) – <i>i</i> (230 ± 27)	10 PELAEZ	04A RVUE	$K \pi \rightarrow K \pi$
(594 ± 79) – <i>i</i> (362 ± 166)	9 ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
(722 ± 60) – <i>i</i> (386 ± 50)	9 BUGG	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
(875 ± 75) – <i>i</i> (335 ± 110)	11 ISHIDA	97B RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
727 – <i>i</i> 263	12 VANBEVEREN	86 RVUE	

- 1 Extracted from Forward Dispersion Relations using sequences of Pade approximants .
 2 Extracted from Breit-Wigner parameters.
 3 Fit to scattering phase shifts using UChPT amplitudes with explicit resonances.
 4 Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an *s*-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.
 5 From a complex pole included in the fit. Using parameters from the model that fits data best.
 6 Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C using for the κ an *s*-dependent width with an Adler zero near threshold.
 7 Using Roy-Steiner equations (ROY 71) consistent with unitarity, analyticity and crossing symmetry constraints.
 8 From UChPT fitted to MERCER 71, BINGHAM 72 and ESTABROOKS 78. Amplitude shown to be consistent with data of ABLIKIM 06C.
 9 Reanalysis of ASTON 88 data.
 10 Reanalysis of data from LINGLIN 73, ESTABROOKS 78, and ASTON 88 using the Inverse Amplitude Method.
 11 Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes. Extracted from Breit-Wigner parameters.
 12 Unitarized Quark Model.

NODE=M174TMP;LINKAGE=N
 NODE=M174TMP;LINKAGE=A
 NODE=M174TMP;LINKAGE=D
 NODE=M174TMP;LINKAGE=H

NODE=M174TMP;LINKAGE=B
 NODE=M174TMP;LINKAGE=G
 NODE=M174TMP;LINKAGE=I
 NODE=M174TMP;LINKAGE=O
 NODE=M174TMP;LINKAGE=F
 NODE=M174TMP;LINKAGE=K
 NODE=M174TMP;LINKAGE=M
 NODE=M174TMP;LINKAGE=E

$K_0^*(700)$ Breit-Wigner Mass				
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
824 ± 30 OUR AVERAGE				
826 ± 49	1.3k	1 ABLIKIM	11B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
810 ± 68	1.4k	2 ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
878 ± 23	25k	3 ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
797 ± 19	15k	4,5 AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
888.0 ± 1.9	141k	6 BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
856 ± 17	54k	7 LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
855 ± 15	0.6k	8 CAWLFIELD	06A CLEO	$D^0 \rightarrow K^+ K^- \pi^0$
905 ± 65		9 ISHIDA	97B RVUE	$11 K^- p \rightarrow K^- \pi^+ n$

NODE=M174M
 NODE=M174M

OCCUR=2
 OCCUR=2

OCCUR=2

NODE=M174M;LINKAGE=LI

NODE=M174M;LINKAGE=BL

NODE=M174M;LINKAGE=EP

NODE=M174M;LINKAGE=A

NODE=M174M;LINKAGE=AU

NODE=M174M;LINKAGE=C
 NODE=M174M;LINKAGE=BW
 NODE=M174M;LINKAGE=CA

NODE=M174M;LINKAGE=IS

NODE=M174W

$K_0^*(700)$ Breit-Wigner Width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
478 ± 50 OUR AVERAGE					NODE=M174W
449 ± 156	+144 -81	1.3k	1 ABLIKIM	11B BES2 $J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	
536 ± 87	+106 -47	1.4k	2 ABLIKIM	10E BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$	OCCUR=2
499 ± 52	+ 55 - 87	25k	3 ABLIKIM	06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$	OCCUR=2
410 ± 43	± 87	15k	4,5 AITALA	02 E791 $D^+ \rightarrow K^- \pi^+ \pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
550.4 ± 11.8	141k	6 BONVICINI	08A CLEO $D^+ \rightarrow K^- \pi^+ \pi^+$		OCCUR=2
464 ± 28	± 22	54k	7 LINK	07B FOCS $D^+ \rightarrow K^- \pi^+ \pi^+$	
251 ± 48	0.6k	8 CAWLFIELD	06A CLEO $D^0 \rightarrow K^+ K^- \pi^0$		
545 +235 -110		9 ISHIDA	97B RVUE $11 K^- p \rightarrow K^- \pi^+ n$		
1 The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$ MeV.					NODE=M174W;LINKAGE=LI
2 From a fit including ten additional resonances and energy-independent Breit-Wigner width.					NODE=M174W;LINKAGE=BL
3 A fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model with mass and width of the $K_0^*(700)$ from ABLIKIM 06C well describes the left slope of the $K_S^0 \pi^-$ invariant mass spectrum in $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$ decay studied by EPIFANOV 07. Averaged value from different parameterizations.					NODE=M174W;LINKAGE=EP
4 Not seen by KOPP 01 using 7070 events of $D^0 \rightarrow K^- \pi^+ \pi^0$. LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than $K_0^*(700)$ in their high statistics analysis of $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$.					NODE=M174W;LINKAGE=A
5 AUBERT 07T does not find evidence for the charged $K_0^*(700)$ using 11k events of $D^0 \rightarrow K^- K^+ \pi^0$.					NODE=M174W;LINKAGE=AU
6 Using parameters from the model that fits data best.					
7 A Breit-Wigner mass and width.					
8 Statistical error only. A fit to the Dalitz plot including the $K_0^*(700)^\pm$, $K^*(892)^\pm$, and ϕ resonances modeled as Breit-Wigners. A significant S-wave can be also modeled as a non-resonant contribution.					NODE=M174W;LINKAGE=CA
9 Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.					NODE=M174W;LINKAGE=IS

$K_0^*(700)$ REFERENCES

PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.Ruiz de Elvira		REFID=57836
ABLIKIM	11B	PL B698 183	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53683
GUO	11B	PR D84 034005	Z.-H. Guo, J.A. Oller		REFID=58808
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)	REFID=53213
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=53056
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
AUBERT	07T	PR D76 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51726
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)	REFID=51929
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51875
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51051
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51458
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)	REFID=50996
CAWLFIELD	06A	PR D74 031108	C. Cawfield <i>et al.</i>	(CLEO Collab.)	REFID=51153
DESCOTES-G...	06	EPJ C48 553	S. Descotes-Genon, B. Moussallam		REFID=51518
GUO	06	NP A773 78	F.K. Guo <i>et al.</i>		REFID=51164
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng		REFID=51198
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=50679
PELAEZ	04A	MPL A19 2879	J.R. Pelaez		REFID=50347
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>		REFID=50165
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
ITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48807
LINK	02E	PL B535 43	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=48728
KOPP	01	PR D63 092001	S. Kopp <i>et al.</i>	(CLEO Collab.)	REFID=48134
ISHIDA	97B	PTP 98 621	S. Ishida <i>et al.</i>		REFID=48655
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i>	(NIJM, BIEL)	REFID=45769
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
BINGHAM	72	NP B41 1	H.H. Bingham <i>et al.</i>	(International K^+ Collab.)	REFID=22415
MERCER	71	NP B32 381	R. Mercer <i>et al.</i>	(JHU)	REFID=22412
ROY	71	PL 36B 353	S.M. Roy		REFID=51107

NODE=M018

K^{*}(892) $I(J^P) = \frac{1}{2}(1^-)$ **K^{*}(892) MASS****CHARGED ONLY, HADROPRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
891.66±0.26 OUR AVERAGE					
892.6 ± 0.5	5840	BAUBILLIER	84B	HBC	—
888 ± 3		NAPIER	84	SPEC	+
891 ± 1		NAPIER	84	SPEC	—
891.7 ± 2.1	3700	BARTH	83	HBC	+
891 ± 1	4100	TOAFF	81	HBC	—
892.8 ± 1.6		AJINENKO	80	HBC	+
890.7 ± 0.9	1800	AGUILAR-...	78B	HBC	±
886.6 ± 2.4	1225	BALAND	78	HBC	±
891.7 ± 0.6	6706	COOPER	78	HBC	±
891.9 ± 0.7	9000	¹ PALER	75	HBC	—
892.2 ± 1.5	4404	AGUILAR-...	71B	HBC	—
891 ± 2	1000	CRENNELL	69D	DBC	—
890 ± 3.0	720	BARLOW	67	HBC	±
889 ± 3.0	600	BARLOW	67	HBC	±
891 ± 2.3	620	² DEBAERE	67B	HBC	+
891.0 ± 1.2	1700	³ WOJCICKI	64	HBC	—

• • • We do not use the following data for averages, fits, limits, etc. • • •

893.6 ± 0.1	^{+0.2} _{-0.3}	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
895.6 ± 0.8	4K	⁴ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$		
893.2 ± 0.1	± 1.0	190k	⁵ AAIJ	16N LHCb	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$	
893.5 ± 1.1	27k	⁶ ABELE	99D CBAR	±	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$	
890.4 ± 0.2	± 0.5	80k	⁷ BIRD	89 LASS	—	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
890.0 ± 2.3	800	^{2,3} CLELAND	82	SPEC	+	$30 K^+ p \rightarrow K_S^0 \pi^+ p$
896.0 ± 1.1	3200	^{2,3} CLELAND	82	SPEC	+	$50 K^+ p \rightarrow K_S^0 \pi^+ p$
893 ± 1	3600	^{2,3} CLELAND	82	SPEC	—	$50 K^+ p \rightarrow K_S^0 \pi^- p$
896.0 ± 1.9	380	DELFOSSE	81	SPEC	+	$50 K^\pm p \rightarrow K^\pm \pi^0 p$
886.0 ± 2.3	187	DELFOSSE	81	SPEC	—	$50 K^\pm p \rightarrow K^\pm \pi^0 p$
894.2 ± 2.0	765	² CLARK	73	HBC	—	$3.13 K^- p \rightarrow \bar{K}^0 \pi^- p$
894.3 ± 1.5	1150	^{2,3} CLARK	73	HBC	—	$3.3 K^- p \rightarrow \bar{K}^0 \pi^- p$
892.0 ± 2.6	341	² SCHWEING...68		HBC	—	$5.5 K^- p \rightarrow \bar{K}^0 \pi^- p$

1 Inclusive reaction. Complicated background and phase-space effects.

2 Mass errors enlarged by us to Γ/\sqrt{N} . See note.

3 Number of events in peak reevaluated by us.

4 From a Dalitz plot analysis in an isobar model with charged and neutral K^{*}(892) masses and widths floating.5 Average of fit results with different parametrizations for the K π S-wave.

6 K-matrix pole.

7 From a partial wave amplitude analysis.

CHARGED ONLY, PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
895.47±0.20±0.74	53k	¹ EPIFANOV	07	BELL $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
892.0 ± 0.5		² BOITO	10	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
892.0 ± 0.9		^{3,4} BOITO	09	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
895.3 ± 0.2		^{4,5} JAMIN	08	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
896.4 ± 0.9	12k	⁶ BONVICINI	02	CLEO $\tau^- \rightarrow K^- \pi^0 \nu_\tau$
895 ± 2		⁷ BARATE	99R ALEP	$\tau^- \rightarrow K^- \pi^0 \nu_\tau$

NODE=M018205

NODE=M018M1
NODE=M018M1

OCCUR=2

OCCUR=2

OCCUR=2
OCCUR=3

OCCUR=2

OCCUR=2
OCCUR=2NODE=M018M;LINKAGE=I
NODE=M018M;LINKAGE=D
NODE=M018M;LINKAGE=W
NODE=M018M1;LINKAGE=BNODE=M018M1;LINKAGE=A
NODE=M018M1;LINKAGE=AN
NODE=M018M1;LINKAGE=FNODE=M018MCT
NODE=M018MCT

¹ From a fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model.

² From the pole position of the $K\pi$ vector form factor using EPIFANOV 07 and constraints from K_{l3} decays in ANTONELLI 10.

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

⁴ Systematic uncertainties not estimated.

⁵ Reanalysis of EPIFANOV 07 using resonance chiral theory.

⁶ Calculated by us from the shift by 4.7 ± 0.9 MeV (statistical uncertainty only) reported in BONVICINI 02 with respect to the world average value from PDG 00.

⁷ With mass and width of the $K^*(1410)$ fixed at 1412 MeV and 227 MeV, respectively.

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
895.55±0.20 OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.
894.68±0.25±0.05		1 ABLIKIM	16F BES3	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.4 ± 0.2 ± 0.2	243k	2 DEL-AMO-SA..11I	BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.7 ± 0.2 ± 0.3	141k	3 BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
895.41±0.32 ^{+0.35} _{-0.43}	18k	4 LINK	05I FOCS	$D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
896 ± 2		BARBERIS	98E OMEG	$450 pp \rightarrow p_f p_s K^* \bar{K}^*$
895.9 ± 0.5 ± 0.2		ASTON	88 LASS	$11 K^- p \rightarrow K^- \pi^+ n$
894.52±0.63	25k	5 ATKINSON	86 OMEG	20–70 γp
894.63±0.76	20k	5 ATKINSON	86 OMEG	20–70 γp
897 ± 1	28k	EVANGELIS...	80 OMEG	$10 \pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$
898.4 ± 1.4	1180	AGUILAR...	78B HBC	$0.76 \bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
894.9 ± 1.6		WICKLUND	78 ASPK	$3,4,6 K^\pm N \rightarrow (K\pi)^0 N$
897.6 ± 0.9		BOWLER	77 DBC	$5.4 K^+ d \rightarrow K^+ \pi^- pp$
895.5 ± 1.0	3600	MCCUBBIN	75 HBC	$3.6 K^- p \rightarrow K^- \pi^+ n$
897.1 ± 0.7	22k	5 PALER	75 HBC	$14.3 K^- p \rightarrow (K\pi)^0 X$
896.0 ± 0.6	10k	FOX	74 RVUE	$2 K^- p \rightarrow K^- \pi^+ n$
896.0 ± 0.6		FOX	74 RVUE	$2 K^+ n \rightarrow K^+ \pi^- p$
896 ± 2		6 MATISON	74 HBC	$12 K^+ p \rightarrow K^+ \pi^- \Delta$
896 ± 1	3186	LEWIS	73 HBC	$2.1\text{--}2.7 K^+ p \rightarrow K\pi\pi p$
894.0 ± 1.3		6 LINGLIN	73 HBC	$2\text{--}13 K^+ p \rightarrow K^+ \pi^- \pi^+ p$
898.4 ± 1.3	1700	7 BUCHNER	72 DBC	$4.6 K^+ n \rightarrow K^+ \pi^- p$
897.9 ± 1.1	2934	7 AGUILAR...	71B HBC	$3.9,4.6 K^- p \rightarrow K^- \pi^+ n$
898.0 ± 0.7	5362	7 AGUILAR...	71B HBC	$3.9,4.6 K^- p \rightarrow K^- \pi^+ \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
898.1 ± 1.0	4K	9 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
895.53±0.17		LEES	13F BABR	$D^+ \rightarrow K^+ K^- \pi^+$
894.9 ± 0.5 ± 0.7	14.4k	10 MITCHELL	09A CLEO	$D_s^+ \rightarrow K^+ K^- \pi^+$
896.2 ± 0.3	20k	11 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
900.7 ± 1.1	5900	BARTH	83 HBC	$70 K^+ p \rightarrow K^+ \pi^- X$

¹ Taking also into account the $K_0^*(1430)^0$ and $K_2^*(1430)^0$.

² Taking into account the $K^*(892)^0$, S-wave and P-wave ($K^*(1410)^0$).

³ From the isobar model with a complex pole for the κ .

⁴ Fit to $K\pi$ mass spectrum includes a non-resonant scalar component.

⁵ Inclusive reaction. Complicated background and phase-space effects.

⁶ From pole extrapolation.

⁷ Mass errors enlarged by us to Γ/\sqrt{N} . See note.

⁸ Number of events in peak reevaluated by us.

⁹ From a Dalitz plot analysis in an isobar model with charged and neutral $K^*(892)$ masses and widths floating.

¹⁰ This value comes from a fit with χ^2 of 178/117.

¹¹ Systematic uncertainties not estimated.

NODE=M018MCT;LINKAGE=EF

NODE=M018MCT;LINKAGE=BT

NODE=M018MCT;LINKAGE=BI

NODE=M018MCT;LINKAGE=NS

NODE=M018MCT;LINKAGE=JA

NODE=M018MCT;LINKAGE=BO

NODE=M018MCT;LINKAGE=BA

NODE=M018M2

NODE=M018M2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M018M2;LINKAGE=BB

NODE=M018M2;LINKAGE=DE

NODE=M018M2;LINKAGE=BO

NODE=M018M2;LINKAGE=LI

NODE=M018M2;LINKAGE=I

NODE=M018M;LINKAGE=C


NODE=M018M2;LINKAGE=D

NODE=M018M2;LINKAGE=W

NODE=M018M2;LINKAGE=A

NODE=M018M2;LINKAGE=MI

NODE=M018M2;LINKAGE=NS



$K^*(892)^0$ mass (MeV)

NODE=M018209

$K^*(892)$ MASSES AND MASS DIFFERENCES

Unrealistically small errors have been reported by some experiments. We use simple “realistic” tests for the minimum errors on the determination of a mass and width from a sample of N events:

$$\delta_{\min}(m) = \frac{\Gamma}{\sqrt{N}}, \quad \delta_{\min}(\Gamma) = 4 \frac{\Gamma}{\sqrt{N}}. \quad (1)$$

We consistently increase unrealistic errors before averaging. For a detailed discussion, see the 1971 edition of this Note.

$m_{K^*(892)^0} - m_{K^*(892)^{\pm}}$

NODE=M018D

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
6.7 ± 1.2 OUR AVERAGE					
7.7 ± 1.7	2980	AGUILAR...	78B	HBC	±0 0.76 $\bar{p}p \rightarrow K^{\mp} K_S^0 \pi^{\pm}$
5.7 ± 1.7	7338	AGUILAR...	71B	HBC	-0 3.9, 4.6 $K^- p$
6.3 ± 4.1	283	¹ BARASH	67B	HBC	0.0 $\bar{p}p$

¹ Number of events in peak reevaluated by us.

NODE=M018D

NODE=M018D

NODE=M018D;LINKAGE=W

$K^*(892)$ RANGE PARAMETER

NODE=M018R

NODE=M018R

NODE=M018R

VALUE (GeV^{-1})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.1 ± 0.5 ± 0.5	243k	¹ DEL-AMO-SA.11I	BABR	0	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
3.96 ± 0.54 ^{+1.31} _{-0.90}	18k	² LINK	05I	FOCS	0 $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
3.4 ± 0.7		ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
12.1 ± 3.2 ± 3.0		BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

¹ Taking into account the $K^*(892)^0$, S -wave and P -wave ($K^*(1410)^0$).

² Fit to $K\pi$ mass spectrum includes a non-resonant scalar component.

$K^*(892)$ WIDTH

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

50.8±0.9 OUR FIT

50.8±0.9 OUR AVERAGE

49 ± 2	5840	BAUBILLIER	84B	HBC	—	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
56 ± 4		NAPIER	84	SPEC	—	200 $\pi^- p \rightarrow 2K_S^0 X$
51 ± 2	4100	TOAFF	81	HBC	—	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
50.5±5.6		AJINENKO	80	HBC	+	32 $K^+ p \rightarrow K^0 \pi^+ X$
45.8±3.6	1800	AGUILAR-...	78B	HBC	±	0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
52.0±2.5	6706	¹ COOPER	78	HBC	±	0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$
52.1±2.2	9000	² PALER	75	HBC	—	14.3 $K^- p \rightarrow (K\pi)^- X$
46.3±6.7	765	¹ CLARK	73	HBC	—	3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$
48.2±5.7	1150	^{1,3} CLARK	73	HBC	—	3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$
54.3±3.3	4404	¹ AGUILAR-...	71B	HBC	—	3.9, 4.6 $K^- p \rightarrow (K\pi)^- p$
46 ± 5	1700	^{1,3} WOJCICKI	64	HBC	—	1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.7±0.2 ^{+0.1} _{-0.2}	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
43.6±1.3	4K	⁴ LEES	17C BABR		$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
47.2±0.3±2.3	190k	⁵ AAIJ	16N LHCb		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
54.8±1.7	27k	⁶ ABELE	99D CBAR	±	0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$
45.2±1 ± 2	80k	⁷ BIRD	89 LASS	—	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
42.8±7.1	3700	BARTH	83 HBC	+	70 $K^+ p \rightarrow K^0 \pi^+ X$
64.0±9.2	800	^{1,3} CLELAND	82 SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
62.0±4.4	3200	^{1,3} CLELAND	82 SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
55 ± 4	3600	^{1,3} CLELAND	82 SPEC	—	50 $K^+ p \rightarrow K_S^0 \pi^- p$
62.6±3.8	380	DELFOSSE	81 SPEC	+	50 $K^\pm p \rightarrow K^\pm \pi^0 p$
50.5±3.9	187	DELFOSSE	81 SPEC	—	50 $K^\pm p \rightarrow K^\pm \pi^0 p$

¹ Width errors enlarged by us to $4 \times \Gamma / \sqrt{N}$; see note.

² Inclusive reaction. Complicated background and phase-space effects.

³ Number of events in peak reevaluated by us.

⁴ From a Dalitz plot analysis in an isobar model with charged and neutral $K^*(892)$ masses and widths floating.

⁵ Average of fit results with different parametrizations for the $K\pi$ S -wave.

⁶ K -matrix pole.

⁷ From a partial wave amplitude analysis.

CHARGED ONLY, PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

46.2±0.6±1.2 53k ¹ EPIFANOV 07 BELL $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.5±1.1		² BOITO	10	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
46.2±0.4		^{3,4} BOITO	09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
47.5±0.4		^{4,5} JAMIN	08	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
55 ± 8		⁶ BARATE	99R	ALEP	$\tau^- \rightarrow K^- \pi^0 \nu_\tau$

¹ From a fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model.

² From the pole position of the $K\pi$ vector form factor using EPIFANOV 07 and constraints from $K_{3/2}$ decays in ANTONELLI 10.

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

⁴ Systematic uncertainties not estimated.

⁵ Reanalysis of EPIFANOV 07 using resonance chiral theory.

⁶ With mass and width of the $K^*(1410)$ fixed at 1412 MeV and 227 MeV, respectively.

NODE=M018R;LINKAGE=DE

NODE=M018R;LINKAGE=LI

NODE=M018215

NODE=M018W1

NODE=M018W1

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M018W;LINKAGE=D

NODE=M018W;LINKAGE=I

NODE=M018W;LINKAGE=W

NODE=M018W1;LINKAGE=B

NODE=M018W1;LINKAGE=A

NODE=M018W1;LINKAGE=AN

NODE=M018W1;LINKAGE=F

NODE=M018W5

NODE=M018W5

NODE=M018W5;LINKAGE=EF

NODE=M018W5;LINKAGE=BT

NODE=M018W5;LINKAGE=BI

NODE=M018W5;LINKAGE=NS

NODE=M018W5;LINKAGE=JA

NODE=M018W5;LINKAGE=BA

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47.3 ±0.5 OUR FIT		Error includes scale factor of 1.9.		
47.3 ±0.5 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.		
46.53±0.56±0.31		¹ ABLIKIM 16F BES3 $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$		
46.5 ±0.3 ±0.2	243k	² DEL-AMO-SA..11I BABR $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$		
45.3 ±0.5 ±0.6	141k	³ BONVICINI 08A CLEO $D^+ \rightarrow K^- \pi^+ \pi^+$		
47.79±0.86 ^{+1.32} _{-1.06}	18k	⁴ LINK 05I FOCS $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$		
54 ±3		BARBERIS 98E OMEG 450 $p p \rightarrow p_f p_s K^* \bar{K}^*$		
50.8 ±0.8 ±0.9		ASTON 88 LASS 11 $K^- p \rightarrow K^- \pi^+ n$		
46.5 ±4.3	5900	BARTH 83 HBC 70 $K^+ p \rightarrow K^+ \pi^- X$		
54 ±2	28k	EVANGELIS... 80 OMEG 10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$		
45.9 ±4.8	1180	AGUILAR-... 78B HBC 0.76 $\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$		
51.2 ±1.7		WICKLUND 78 ASPK 3,4,6 $K^\pm N \rightarrow (K\pi)^0 N$		
48.9 ±2.5		BOWLER 77 DBC 5.4 $K^+ d \rightarrow K^+ \pi^- p p$		
48 ±3	3600	MCCUBBIN 75 HBC 3.6 $K^- p \rightarrow K^- \pi^+ n$		
50.6 ±2.5	22k	⁵ PALER 75 HBC 14.3 $K^- p \rightarrow (K\pi)^0 X$		
47 ±2	10k	FOX 74 RVUE 2 $K^- p \rightarrow K^- \pi^+ n$		
51 ±2		FOX 74 RVUE 2 $K^+ n \rightarrow K^+ \pi^- p$		OCCUR=2
46.0 ±3.3	3186	⁶ LEWIS 73 HBC 2.1-2.7 $K^+ p \rightarrow K\pi\pi p$		
51.4 ±5.0	1700	⁶ BUCHNER 72 DBC 4.6 $K^+ n \rightarrow K^+ \pi^- p$		
55.8 ±4.2 _{-3.4}	2934	⁶ AGUILAR-... 71B HBC 3.9,4.6 $K^- p \rightarrow K^- \pi^+ n$		
48.5 ±2.7	5362	AGUILAR-... 71B HBC 3.9,4.6 $K^- p \rightarrow K^- \pi^+ \pi^- p$		OCCUR=2
54.0 ±3.3	4300	^{6,7} HABER 70 DBC 3 $K^- N \rightarrow K^- \pi^+ X$		
53.2 ±2.1	10k	⁶ DAVIS 69 HBC 12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$		
44 ±5.5	1040	⁶ DAUBER 67B HBC 2.0 $K^- p \rightarrow K^- \pi^+ \pi^- p$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

52.6 ±1.7	4K	⁸ LEES 17C BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	
44.90±0.30		LEES 13F BABR $D^+ \rightarrow K^+ K^- \pi^+$	
45.7 ±1.1 ±0.5	14.4k	⁹ MITCHELL 09A CLEO $D_s^+ \rightarrow K^+ K^- \pi^+$	
50.6 ±0.9	20k	¹⁰ AUBERT 07AK BABR $10.6 \frac{e^+ e^-}{K^{*0} K^\pm \pi^\mp \gamma}$	

1 Taking also into account the $K_0^*(1430)^0$ and $K_2^*(1430)^0$.

2 Taking into account the $K^*(892)^0$, S-wave and P-wave ($K^*(1410)^0$).

3 From the isobar model with a complex pole for the κ .

4 Fit to $K\pi$ mass spectrum includes a non-resonant scalar component.

5 Inclusive reaction. Complicated background and phase-space effects.

6 Width errors enlarged by us to $4 \times \Gamma/\sqrt{N}$; see note.

7 Number of events in peak reevaluated by us.

8 From a Dalitz plot analysis in an isobar model with charged and neutral $K^*(892)$ masses and widths floating.

9 This value comes from a fit with χ^2 of 178/117.


10 Systematic uncertainties not estimated.

NODE=M018W2
NODE=M018W2

NODE=M018W2;LINKAGE=B
NODE=M018W2;LINKAGE=DE
NODE=M018W2;LINKAGE=BO
NODE=M018W2;LINKAGE=LN
NODE=M018W2;LINKAGE=I
NODE=M018W2;LINKAGE=D
NODE=M018W2;LINKAGE=W
NODE=M018W2;LINKAGE=A

NODE=M018W2;LINKAGE=MI
NODE=M018W2;LINKAGE=NS

WEIGHTED AVERAGE
47.3±0.5 (Error scaled by 2.0)



NEUTRAL ONLY (MeV)

K*(892) DECAY MODES

NODE=M018220;NODE=M018

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \ K\pi$	~ 100	%
$\Gamma_2 \ (K\pi)^{\pm}$	(99.901 ± 0.009)	%
$\Gamma_3 \ (K\pi)^0$	(99.754 ± 0.021)	%
$\Gamma_4 \ K^0\gamma$	(2.46 ± 0.21) × 10 ⁻³	
$\Gamma_5 \ K^{\pm}\gamma$	(9.9 ± 0.9) × 10 ⁻⁴	
$\Gamma_6 \ K\pi\pi$	< 7	× 10 ⁻⁴ 95%

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=11
 DESIG=12
 DESIG=4
 DESIG=3
 DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 7.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_5 & -100 \\ \Gamma & 19 & -19 \\ \hline x_2 & x_5 \end{array}$$

Mode	Rate (MeV)
$\Gamma_2 \ (K\pi)^{\pm}$	50.7 ± 0.9
$\Gamma_5 \ K^{\pm}\gamma$	0.050 ± 0.005

DESIG=11
 DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 23 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 68.4$ for 21 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-100			
Γ	12	-12		
x_3		x_4		

Mode	Rate (MeV)	Scale factor
$\Gamma_3 (K\pi)^0$	47.2 ± 0.5	1.9
$\Gamma_4 K^0 \gamma$	0.117 ± 0.010	

$K^*(892)$ PARTIAL WIDTHS

$\Gamma(K^0 \gamma)$	Γ_4
<u>VALUE (keV)</u>	<u>EVTS</u>
116 ± 10 OUR FIT	

116.5 ± 9.9 584 CARLSMITH 86 SPEC 0 $K_L^0 A \rightarrow K_S^0 \pi^0 A$

$\Gamma(K^\pm \gamma)$	Γ_5
<u>VALUE (keV)</u>	<u>DOCUMENT ID</u>
50 ± 5 OUR FIT	
50 ± 5 OUR AVERAGE	
48 ± 11	BERG 83 SPEC - 156 $K^- A \rightarrow \bar{K} \pi A$
51 ± 5	CHANDLEE 83 SPEC + 200 $K^+ A \rightarrow K \pi A$

$K^*(892)$ BRANCHING RATIOS

$\Gamma(K^0 \gamma) / \Gamma_{\text{total}}$	Γ_4 / Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>

2.46 ± 0.21 OUR FIT • • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ± 0.7 CARITHERS 75B CNTR 0 8–16 $\bar{K}^0 A$

$\Gamma(K^\pm \gamma) / \Gamma_{\text{total}}$	Γ_5 / Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>

0.99 ± 0.09 OUR FIT • • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6 95 BEMPORAD 73 CNTR + 10–16 $K^+ A$

$\Gamma(K\pi\pi) / \Gamma((K\pi)^\pm)$	Γ_6 / Γ_2
<u>VALUE</u>	<u>CL%</u>

< 7×10^{-4} 95 JONGEJANS 78 HBC 4 $K^- p \rightarrow p \bar{K}^0 2\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 20×10^{-4} WOJCICKI 64 HBC – 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$

$K^*(892)$ REFERENCES

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	16F	PR D94 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA... ANTONELLI	11I 10	PR D83 072001 EPJ C69 399	P. del Amo Sanchez <i>et al.</i> M. Antonelli <i>et al.</i>	(FlaviaNet Working Group)
BOITO	10	JHEP 1009 031	D.R. Boito, R. Escrivano, M. Jamin	(BARC)
BOITO	09	EPJ C59 821	D.R. Boito, R. Escrivano, M. Jamin	(CLEO Collab.)
MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
JAMIN	08	PL B664 78	M. Jamin, A. Pich, J. Portoles	(BABAR Collab.)
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BELLE Collab.)
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)

NODE=M018

REFID=59909
REFID=57981
REFID=57273
REFID=57307
REFID=55127
REFID=16493
REFID=53448
REFID=53632
REFID=52728
REFID=52756
REFID=52426
REFID=52285
REFID=51908
REFID=51929

LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=50679
BONVICINI	02	PRL 88 111803	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=48701
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ABELE	99D	PL B468 178	D. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47366
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46348
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
CARLSMITH	86	PRL 56 18	D. Carlsmith <i>et al.</i>	(EFI, SACL)	REFID=22461
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
NAPIER	84	PL 149B 514	A. Napier <i>et al.</i>	(TUFTS, ARIZ, FNAL, FLOR+)	REFID=22460
BARTH	83	NP B223 296	M. Barth <i>et al.</i>	(BRUX, CERN, GENO, MONS+)	REFID=22456
BERG	83	Thesis UMI 83-21652	D.M. Berg	(ROCH)	REFID=22457
CHANDLEE	83	PRL 51 168	C. Chandlee <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=22458
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
AJINENKO	80	ZPHY C5 177	I.V. Ajinenko <i>et al.</i>	(SERP, BRUX, MONS+)	REFID=22449
EVANGELIS...	80	NP B165 383	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=22450
AGUILAR-...	78B	NP B141 101	M. Aguilar-Benitez <i>et al.</i>	(MADR, TATA+)	REFID=22438
BALAND	78	NP B140 220	J.F. Baland <i>et al.</i>	(MONS, BELG, CERN+)	REFID=20369
COOPER	78	NP B136 365	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=22441
JONGEJANS	78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
BOWLER	77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
CARITHERS	75B	PRL 35 349	W.C.J. Carithers <i>et al.</i>	(ROCH, MCGI)	REFID=22433
MCCUBBIN	75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
PALER	75	NP B96 1	K. Paler <i>et al.</i>	(RHEL, SACL, EPOL)	REFID=22435
FOX	74	NP B80 403	G.C. Fox, M.L. Griss	(CIT)	REFID=22430
MATISON	74	PR D9 1872	M.J. Matison <i>et al.</i>	(LBL)	REFID=22431
BEMPORAD	73	NP B51 1	C. Bemporad <i>et al.</i>	(CERN, ETH, LOIC)	REFID=22416
CLARK	73	NP B54 432	A.G. Clark, L. Lyons, D. Radojicic	(OXF)	REFID=22426
LEWIS	73	NP B60 283	P.H. Lewis <i>et al.</i>	(LOWC, LOIC, CDEF)	REFID=22427
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
BUCHNER	72	NP B45 333	K. Buchner <i>et al.</i>	(MPIIM, CERN, BRUX)	REFID=22418
AGUILAR-...	71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
HABER	70	NP B17 289	B. Haber <i>et al.</i>	(REHO, SACL, BGNA, EPOL)	REFID=22406
CRENNELL	69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS	69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
SCHWEING...	68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
DAUBER	67B	PR 153 1403	P.M. Dauber <i>et al.</i>	(UCLA)	REFID=22389
DEBAERE	67B	NC 51A 401	W. de Baere <i>et al.</i>	(BRUX, CERN)	REFID=22390
WOJCICKI	64	PR 135 B484	S.G. Wojcicki	(LRL)	REFID=22379

$K_1(1270)$ $I(J^P) = \frac{1}{2}(1^+)$ **$K_1(1270)$ MASS**VALUE (MeV) DOCUMENT ID**1272±7 OUR AVERAGE** Includes data from the 2 datablocks that follow this one.**PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE**VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

1275±10 700 GAVILLET 78 HBC + 4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$ **PRODUCED BY K BEAMS**VALUE (MeV) DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

1270±10 ¹ DAUM 81C CNTR - 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~1276	² TORNQVIST	82B	RVUE	
~1300	VERGEEST	79	HBC	-
1289±25	³ CARNEGIE	77	ASPK	±
~1300	BRANDENB...	76	ASPK	±
~1270	OTTER	76	HBC	-
1260	DAVIS	72	HBC	+
1234±12	FIRESTONE	72B	DBC	+

1 Well described in the chiral unitary approach of GEN 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

2 From a unitarized quark-model calculation.

3 From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

PRODUCED BY BEAMS OTHER THAN K MESONSVALUE (MeV) EVTS DOCUMENT ID TECN COMMENT**1248.1 ± 3.3 ± 1.4** GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1289.81 ± 0.56 ± 1.66	894k	AAIJ	18AI	LHCb	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
1279 ± 10	25k	¹ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
1294 ± 10	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
1300	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
1242 ⁺⁹ ₋₁₀		² ASTIER	69	HBC	$\bar{p}p$
1300	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

1 Systematic errors not estimated.

2 This was called the C meson.**PRODUCED IN τ LEPTON DECAYS**VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT**1254±33±34** 7k ASNER 00B CLEO ± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ **$K_1(1270)$ WIDTH**VALUE (MeV) DOCUMENT ID**90±20 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.**87± 7 OUR AVERAGE** Includes data from the 2 datablocks that follow this one.**PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE**VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

75±15 700 GAVILLET 78 HBC + 4.2 $K^- p \rightarrow \Xi^- K\pi\pi$

NODE=M028205

NODE=M028MX

NODE=M028M2

NODE=M028M2

NODE=M028M3

NODE=M028M3

NODE=M028M3;LINKAGE=DA

NODE=M028M3;LINKAGE=T

NODE=M028M3;LINKAGE=E

NODE=M028M1

NODE=M028M1

NODE=M028M1;LINKAGE=AB

NODE=M028M1;LINKAGE=A

NODE=M028MT

NODE=M028MT

NODE=M028210

NODE=M028WX

→ UNCHECKED ←

NODE=M028W2

NODE=M028W2

PRODUCED BY K BEAMS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

90± 8 ¹ DAUM 81C CNTR — 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 150	VERGEEST	79	HBC	—	4.2 $K^- p \rightarrow (\bar{K}\pi\pi)^- p$
150±71	² CARNEGIE	77	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
~ 200	BRANDENB...	76	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
120	DAVIS	72	HBC	+	12 $K^+ p$
188±21	FIRESTONE	72B	DBC	+	12 $K^+ d$

¹ Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

² From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

119.5 ± 5.2 ±6.7 GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

116.11± 1.65±2.96	894k	AAIJ	18AI LHCb	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
131 ± 21	25k	¹ ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
66 ± 15	310	RODEBACK	81 HBC	4 $\pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72 HBC	4.5 $\pi^- p \rightarrow \Lambda K 2\pi$
127 + 7 - 25		ASTIER	69 HBC	$\bar{p} p$
60	45	CRENNELL	67 HBC	6 $\pi^- p \rightarrow \Lambda K 2\pi$

¹ Systematic errors not estimated.

PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**260 +90
-70 ±80** 7k ASNER 00B CLEO ± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$

 $K_1(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\rho$	(42 ± 6) %
$\Gamma_2 K_0^*(1430)\pi$	(28 ± 4) %
$\Gamma_3 K^*(892)\pi$	(16 ± 5) %
$\Gamma_4 K\omega$	(11.0 ± 2.0) %
$\Gamma_5 K f_0(1370)$	(3.0 ± 2.0) %
$\Gamma_6 \gamma K^0$	seen

 $K_1(1270)$ PARTIAL WIDTHS

$\Gamma(K\rho)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1
-----------------	-------------	------	-----	---------	------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

57±5	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^-(K\pi\pi)^+$
75±6	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$\Gamma(K_0^*(1430)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2
--------------------------	-------------	------	-----	---------	------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

26±6	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
------	----------	-----	------	---	--

$\Gamma(K^*(892)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3
-----------------------	-------------	------	-----	---------	------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14±11	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^-(K\pi\pi)^+$
2± 2	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W3

NODE=M028W3

NODE=M028W3;LINKAGE=DA

NODE=M028W3;LINKAGE=E

NODE=M028W1

NODE=M028W1

NODE=M028W1;LINKAGE=AB

NODE=M028WT

NODE=M028WT

DESIG=2

DESIG=7

DESIG=1

DESIG=5

DESIG=8

DESIG=9;OUR EST;→ UNCHECKED ←

NODE=M028220

NODE=M028W5

NODE=M028W5

NODE=M028W7

NODE=M028W7

NODE=M028W4

NODE=M028W4

$\Gamma(K\omega)$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4±4	MAZZUCATO 79	HBC	+	4.2 $K^- p \rightarrow \Xi^-(K\pi\pi)^+$
24±3	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_4

NODE=M028W6
NODE=M028W6

 $\Gamma(K\eta(1370))$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22±5	CARNEGIE 77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

 Γ_5

NODE=M028W8
NODE=M028W8

 $\Gamma(\gamma K^0)$

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
73.2±6.1±28.3	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$	

 Γ_6

NODE=M028W9
NODE=M028W9

 $K_1(1270)$ BRANCHING RATIOS $\Gamma(K\rho)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.42 ±0.06	¹ DAUM 81C	CNTR	$63 K^- p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.584±0.043 dominant	² GULER 11 RODEBACK 81	BELL HBC	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

 Γ_1/Γ

NODE=M028225

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.28 ±0.04	¹ DAUM 81C	CNTR	$63 K^- p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0201±0.0064	² GULER 11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

 Γ_2/Γ

NODE=M028R4
NODE=M028R4

 $\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.16 ±0.05	¹ DAUM 81C	CNTR	$63 K^- p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.171±0.023	² GULER 11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

 Γ_3/Γ

NODE=M028R1
NODE=M028R1

 $\Gamma(K\omega)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.11 ±0.02	¹ DAUM 81C	CNTR	$63 K^- p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.225±0.052	² GULER 11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

 Γ_4/Γ

NODE=M028R3
NODE=M028R3

 $\Gamma(K\omega)/\Gamma(K\rho)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.30	95	RODEBACK 81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$

 Γ_4/Γ_1

NODE=M028R6
NODE=M028R6

 $\Gamma(K\eta(1370))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.03±0.02	¹ DAUM 81C	CNTR	$63 K^- p \rightarrow K^- 2\pi p$

 Γ_5/Γ

NODE=M028R5
NODE=M028R5

D-wave/S-wave RATIO FOR $K_1(1270) \rightarrow K^*(892)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
1.0±0.7	¹ DAUM 81C	CNTR	$63 K^- p \rightarrow K^- 2\pi p$

 Γ_5/Γ

NODE=M028R9
NODE=M028R9

¹ Average from low and high t data.

² Assuming that decays are saturated by the $K\rho$, $K_0^*(1430)\pi$, $K^*(892)\pi$, $K\omega$ decay modes and neglecting interference between them. The values $B(\omega \rightarrow \pi^+ \pi^-) = (1.53^{+0.11}_{-0.13})\%$ and $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10)\%$ are used. Systematic uncertainties not estimated.

NODE=M028R;LINKAGE=F
NODE=M028R1;LINKAGE=GU

K₁(1270) REFERENCES

AAIJ	18AI	EPJ C78	443	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83	032005	H. Guler <i>et al.</i>	(BELLE Collab.)
GENG	07	PR D75	014017	L.S. Geng <i>et al.</i>	
ABLIKIM	06C	PL B633	681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL	89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62	072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
TORNQVIST	82B	NP B203	268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187	1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
RODEBACK	81	ZPHY C9	9	S. Rodeback <i>et al.</i>	(CERN, CDEF, MADR+)
MAZZUCATO	79	NP B156	532	M. Mazzucato <i>et al.</i>	(CERN, ZEEM, NIJM+)
VERGEEST	79	NP B158	265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
GAVILLET	78	PL 76B	517	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+) JP
CARNEGIE	77	NP B127	509	R.K. Carnegie <i>et al.</i>	(SLAC)
CARNEGIE	77B	PL 68B	287	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL	36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
OTTER	76	NP B106	77	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
CRENNELL	72	PR D6	1220	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS	72	PR D5	2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5	505	A. Firestone <i>et al.</i>	(LBL)
ASTIER	69	NP B10	65	A. Astier <i>et al.</i>	(CDEF, CERN, IPNP, LIVP) IJP
CRENNELL	67	PRL	19 44	D.J. Crennell <i>et al.</i>	(BNL) I

NODE=M028

REFID=59187
REFID=53668
REFID=51623
REFID=51037
REFID=48822
REFID=47766
REFID=20573
REFID=22548
REFID=22550
REFID=20867
REFID=22542
REFID=22538
REFID=22535
REFID=22536
REFID=22532
REFID=22533
REFID=22419
REFID=22505
REFID=22506
REFID=22482
REFID=22473

K₁(1400)

$$I(J^P) = \frac{1}{2}(1^+)$$

K₁(1400) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1403± 7 OUR AVERAGE					
1463±64±68	7k	ASNER	00B	CLEO	\pm $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
1373±14±18	1	ASTON	87	LASS	0 $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1392±18		BAUBILLIER	82B	HBC	0 $8.25 K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
1410±25		DAUM	81C	CNTR	— $63 K^- p \rightarrow K^- 2\pi p$
1415±15		ETKIN	80	MPS	0 $6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1404±10	2	CARNEGIE	77	ASPK	\pm $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1418± 8	25k	3 ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 1350		4 TORNQVIST	82B	RVUE	
~ 1400		VERGEEST	79	HBC	— $4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 1400		BRANDENB...	76	ASPK	\pm $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
1420		DAVIS	72	HBC	+
1368±18		FIRESTONE	72B	DBC	+

1 From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.

2 From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

3 Systematic errors not estimated.

4 From a unitarized quark-model calculation.

NODE=M064

NODE=M064M

NODE=M064M

NODE=M064M;LINKAGE=P

NODE=M064M;LINKAGE=E

NODE=M064M;LINKAGE=AB

NODE=M064M;LINKAGE=T

K₁(1400) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
174± 13 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.					
300 ⁺³⁷⁰ ₋₁₁₀ ±140	7k	ASNER	00B	CLEO	\pm $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
188± 54± 60	5	ASTON	87	LASS	0 $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
276± 65		BAUBILLIER	82B	HBC	0 $8.25 K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
195± 25		DAUM	81C	CNTR	— $63 K^- p \rightarrow K^- 2\pi p$
180± 10		ETKIN	80	MPS	0 $6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
142± 16	6	CARNEGIE	77	ASPK	\pm $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
152± 16	25k	7 ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 200		VERGEEST	79	HBC	— $4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 160		BRANDENB...	76	ASPK	\pm $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
80		DAVIS	72	HBC	+
241± 30		FIRESTONE	72B	DBC	+

NODE=M064W

NODE=M064W

5 From partial-wave analysis of $K^0\pi^+\pi^-$ system.


6 From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

7 Systematic errors not estimated.

NODE=M064W;LINKAGE=P

NODE=M064W;LINKAGE=E

NODE=M064W;LINKAGE=AB



K₁(1400) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892)\pi$	(94 ± 6) %
$\Gamma_2 K\rho$	(3.0 ± 3.0) %
$\Gamma_3 Kf_0(1370)$	(2.0 ± 2.0) %
$\Gamma_4 K\omega$	(1.0 ± 1.0) %
$\Gamma_5 K_0^*(1430)\pi$	not seen
$\Gamma_6 \gamma K^0$	seen

K₁(1400) PARTIAL WIDTHS

$\Gamma(K^*(892)\pi)$	Γ_1
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
117±10	CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
$\Gamma(K\rho)$	Γ_2
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
2±1	CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
$\Gamma(K\omega)$	Γ_4
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
23±12	CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
$\Gamma(\gamma K^0)$	Γ_6
VALUE (keV)	DOCUMENT ID TECN COMMENT
280.8±23.2±40.4	ALAVI-HARATI02B KTEV $K + A \rightarrow K^* + A$

K₁(1400) BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	DOCUMENT ID TECN COMMENT
0.94±0.06	8 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$
$\Gamma(K\rho)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE	DOCUMENT ID TECN COMMENT
0.03±0.03	8 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064215;NODE=M064

DESIG=1

DESIG=2

DESIG=8

DESIG=5

DESIG=7;OUR EST;→ UNCHECKED ←
DESIG=9;OUR EST;→ UNCHECKED ←

NODE=M064220

NODE=M064W1

NODE=M064W1

NODE=M064W2

NODE=M064W2

NODE=M064W5

NODE=M064W5

NODE=M064W6

NODE=M064W6

NODE=M064225

NODE=M064R1

NODE=M064R1

NODE=M064R2

NODE=M064R2

$\Gamma(K f_0(1370))/\Gamma_{\text{total}}$					Γ_3/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.02±0.02	8 DAUM	81C CNTR	63	$K^- p \rightarrow K^- 2\pi p$	
$\Gamma(K\omega)/\Gamma_{\text{total}}$					Γ_4/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.01±0.01	8 DAUM	81C CNTR	63	$K^- p \rightarrow K^- 2\pi p$	
$\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$					Γ_5/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
not seen	8 DAUM	81C CNTR	63	$K^- p \rightarrow K^- 2\pi p$	
D-wave/S-wave RATIO FOR $K_1(1400) \rightarrow K^*(892)\pi$					
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.04±0.01	8 DAUM	81C CNTR	63	$K^- p \rightarrow K^- 2\pi p$	

8 Average from low and high t data.

K₁(1400) REFERENCES

ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENBURG	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)

K^{*}(1410)

$$I(J^P) = \frac{1}{2}(1^-)$$

K^{*}(1410) MASS

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
1414±15 OUR AVERAGE					Error includes scale factor of 1.3.
1380±21±19		ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
1420± 7±10		ASTON	87	LASS 0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1437± 8±16	190k	¹ AAIJ	16N	LHCb	$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
1426± 8±24	190k	² AAIJ	16N	LHCb	$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
1276 ⁺⁷² ₋₇₇		^{3,4} BOITO	09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
1367±54		BIRD	89	LASS	—
1474±25		BAUBILLIER	82B	HBC 0	$8.25 K^- p \rightarrow \bar{K}^0 2\pi n$
1500±30		ETKIN	80	MPS 0	$6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

1 Using a parametrization for the $K\pi$ S-wave similar to ASTON 88 with fixed resonance width.

2 Using a $K\pi$ S-wave parametrization with resonant and non-resonant contributions.

3 From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

4 Systematic uncertainties not estimated.

NODE=M064R5
NODE=M064R5

NODE=M064R3
NODE=M064R3

NODE=M064R4
NODE=M064R4

NODE=M064R9
NODE=M064R9

NODE=M064R;LINKAGE=F

NODE=M064

REFID=51037
REFID=48822
REFID=47766
REFID=40234
REFID=22551
REFID=20573
REFID=22548
REFID=22545
REFID=22542
REFID=22535
REFID=22532
REFID=22505
REFID=22506

NODE=M094

NODE=M094M

NODE=M094M

OCCUR=2

NODE=M094M;LINKAGE=A

NODE=M094M;LINKAGE=C

NODE=M094M;LINKAGE=BI

NODE=M094M;LINKAGE=NS

NODE=M094W

NODE=M094W

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
232± 21 OUR AVERAGE					Error includes scale factor of 1.1.
176± 52±22		ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
240± 18±12		ASTON	87	LASS 0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M064R5

NODE=M064R3

NODE=M064R4

NODE=M064R9

NODE=M064R;LINKAGE=F

NODE=M064

REFID=51037
REFID=48822
REFID=47766
REFID=40234
REFID=22551
REFID=20573
REFID=22548
REFID=22545
REFID=22542
REFID=22535
REFID=22532
REFID=22505
REFID=22506

NODE=M094

NODE=M094M

NODE=M094M

OCCUR=2

NODE=M094M;LINKAGE=A

NODE=M094M;LINKAGE=C

NODE=M094M;LINKAGE=BI

NODE=M094M;LINKAGE=NS

NODE=M094W

NODE=M094W

• • • We do not use the following data for averages, fits, limits, etc. • • •

210 ± 20 ± 60	190k	1 AAIJ	16N LHCb	$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
270 ± 20 ± 40	190k	1 AAIJ	16N LHCb	$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
198 ± 61		2,3 BOITO	09 RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
114 ± 101		BIRD	89 LASS	—
275 ± 65		BAUBILLIER	82B HBC	0 8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
500 ± 100		ETKIN	80 MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

1 Using a $K\pi$ S-wave parametrization with resonant and non-resonant contributions.

2 From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

3 Systematic uncertainties not estimated.

OCCUR=2

NODE=M094W;LINKAGE=A

NODE=M094W;LINKAGE=BI

NODE=M094W;LINKAGE=NS

NODE=M094215;NODE=M094

$K^*(1410)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 K^*(892)\pi$	> 40 %	95%
$\Gamma_2 K\pi$	(6.6 ± 1.3) %	
$\Gamma_3 K\rho$	< 7 %	95%
$\Gamma_4 \gamma K^0$	< 2.3 $\times 10^{-4}$	90%

$K^*(1410)$ PARTIAL WIDTHS

$\Gamma(\gamma K^0)$	Γ_4
VALUE (keV) CL%	DOCUMENT ID TECN COMMENT
<52.9 90	ALAVI-HARATI02B KTeV $K + A \rightarrow K^* + A$

DESIG=2

DESIG=1

DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=4

NODE=M094217

NODE=M094W1

NODE=M094W1

$K^*(1410)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	Γ_3/Γ_1
VALUE CL% DOCUMENT ID TECN CHG COMMENT	
<0.17 95 ASTON 84 LASS 0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$	

NODE=M094220

NODE=M094R1

NODE=M094R1

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$	Γ_2/Γ_1
VALUE CL% DOCUMENT ID TECN CHG COMMENT	
<0.16 95 ASTON 84 LASS 0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$	

NODE=M094R2

NODE=M094R2

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE DOCUMENT ID TECN CHG COMMENT	
0.066 ± 0.010 ± 0.008 ASTON 88 LASS 0 11 $K^- p \rightarrow K^- \pi^+ n$	

NODE=M094R3

NODE=M094R3

$K^*(1410)$ REFERENCES

AAIJ 16N PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
BOITO 09 EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin	
EPIFANOV 07 PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)
ALAVI-HARATI 02B PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
BIRD 89 SLAC-332	P.F. Bird	(SLAC)
ASTON 88 NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON 87 NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON 84 PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
BAUBILLIER 82B NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ETKIN 80 PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP

NODE=M094

REFID=57273

REFID=52728

REFID=51929

REFID=48822

REFID=41002

REFID=40262

REFID=40234

REFID=22689

REFID=22551

REFID=22545

NODE=M019

 $K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

See our minireview in the 1994 edition and in this edition under the $f_0(500)$.

 $K_0^*(1430)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1425 ± 50 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1438 ± 8 ± 4 5.4k		1 LEES	14E BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
1427 ± 4 ± 13		2 BUGG	10 RVUE	S-matrix pole
1466.6 ± 0.7 ± 3.4 141k		3 BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1412		4 LINK	07 FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
1461.0 ± 4.0 ± 2.1 54k		5 LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
1406 ± 29		6 BUGG	06 RVUE	
1435 ± 6		7 ZHOU	06 RVUE	$Kp \rightarrow K^- \pi^+ n$
1455 ± 20 ± 15		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1456 ± 8		8 ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
~ 1419		9 BUGG	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
~ 1440		10 LI	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
1459 ± 9	15k	11 AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1440		12 JAMIN	00 RVUE	$Kp \rightarrow Kp$
1436 ± 8		13 BARBERIS	98E OMEG	$450 pp \rightarrow p_f p_s K^+ K^- \pi^+ \pi^-$
1415 ± 25		9 ANISOVICH	97C RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
~ 1450		14 TORNQVIST	96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi$
1412 ± 6		15 ASTON	88 LASS	$11 K^- p \rightarrow K^- \pi^+ n$
~ 1430		BAUBILLIER	84B HBC	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 1425		16 ESTABROOKS	78 ASPK	$13 K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$
~ 1450.0		MARTIN	78 SPEC	$10 K_S^\pm p \rightarrow K_S^0 \pi p$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed width $\Gamma(K_0^*(1430)) = 210$ MeV.

² S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.

³ From the isobar model with a complex pole for the κ .

⁴ From a non-parametric analysis.

⁵ A Breit-Wigner mass and width.

⁶ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s-dependent width and an Adler zero near threshold.

⁷ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1950)$.

⁸ Using ASTON 88 and assuming $K_0^*(700)$.

⁹ T-matrix pole. Reanalysis of ASTON 88 data.

¹⁰ Breit-Wigner fit. Using ASTON 88.

¹¹ Assuming a low-mass scalar $K\pi$ resonance, $\kappa(700)$.

¹² T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

¹³ J^P not determined, could be $K_2^*(1430)$.

¹⁴ T-matrix pole.

¹⁵ Uses a model for the background, without this background they get a mass 1340 MeV, where the phase shift passes 90°.

¹⁶ Mass defined by pole position. From elastic $K\pi$ partial-wave analysis.

NODE=M019

NODE=M019M

NODE=M019M
→ UNCHECKED ←

NODE=M019M;LINKAGE=LE

NODE=M019M;LINKAGE=BG

NODE=M019M;LINKAGE=BO

NODE=M019M;LINKAGE=LI

NODE=M019M;LINKAGE=BW

NODE=M019M;LINKAGE=BU

NODE=M019M;LINKAGE=ZU

NODE=M019M;LINKAGE=ZH

NODE=M019M;LINKAGE=A1

NODE=M019M;LINKAGE=E

NODE=M019M;LINKAGE=A0

NODE=M019M;LINKAGE=JM

NODE=M019M;LINKAGE=JP

NODE=M019M;LINKAGE=TT

NODE=M019M;LINKAGE=D

NODE=M019M;LINKAGE=A

NODE=M019W

NODE=M019W
→ UNCHECKED ← **$K_0^*(1430)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
270 ± 80 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				

210	± 20	± 12	5.4k	¹ LEES	14E BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
270	± 10	± 40		² BUGG	10 RVUE	S-matrix pole
174.2 \pm 1.9 \pm 3.2	141k			³ BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 500				⁴ LINK	07 FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
177.0 \pm 8.0 \pm 3.4	54k			⁵ LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
350	± 40			⁶ BUGG	06 RVUE	
288	± 22			⁷ ZHOU	06 RVUE	$K p \rightarrow K^- \pi^+ n$
270	± 45	$+30$ -35		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
217	± 31			⁸ ZHENG	04 RVUE	$K^- p \rightarrow K^- \pi^+ n$
~ 316				⁹ BUGG	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
~ 350				¹⁰ LI	03 RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
175 ± 17	15k			¹¹ AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 300				¹² JAMIN	00 RVUE	$K p \rightarrow K p$
196 ± 45				¹³ BARBERIS	98E OMEG	$450 pp \rightarrow p_f p_s K^+ K^- \pi^+ \pi^-$
330 ± 50				⁹ ANISOVICH	97C RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
~ 320				¹⁴ TORNQVIST	96 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi$
294 ± 23				ASTON	88 LASS	$11 K^- p \rightarrow K^- \pi^+ n$
~ 200				BAUBILLIER	84B HBC	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
200 to 300				¹⁵ ESTABROOKS	78 ASPK	$13 K^\pm p \rightarrow K^\pm \pi^\pm(n, \Delta)$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed mass $M(K_0^*(1430)) = 1435$ MeV.

² S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s -dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.

³ From the isobar model with a complex pole for the κ .

⁴ From a non-parametric analysis.

⁵ A Breit-Wigner mass and width.

⁶ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s -dependent width and an Adler zero near threshold.

⁷ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700), K_0^*(1950)$.

⁸ Using ASTON 88 and assuming $K_0^*(700)$.

⁹ T-matrix pole. Reanalysis of ASTON 88 data.

10 Breit-Wigner fit. Using ASTON 88.

11 Assuming a low-mass scalar $K\pi$ resonance, $\kappa(700)$.

12 T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

13 J^P not determined, could be $K_2^*(1430)$.

14 T-matrix pole.

15 From elastic $K\pi$ partial-wave analysis.

NODE=M019W;LINKAGE=LE

NODE=M019W;LINKAGE=BG

NODE=M019W;LINKAGE=BO

NODE=M019W;LINKAGE=LI

NODE=M019W;LINKAGE=BW

NODE=M019W;LINKAGE=BU

NODE=M019W;LINKAGE=ZU

NODE=M019W;LINKAGE=ZH

NODE=M019W;LINKAGE=A1

NODE=M019W;LINKAGE=E

NODE=M019W;LINKAGE=A0

NODE=M019W;LINKAGE=JM

NODE=M019W;LINKAGE=JP

NODE=M019W;LINKAGE=TT

NODE=M019W;LINKAGE=C

NODE=M019215;NODE=M019

DESIG=1

DESIG=2

DESIG=3

NODE=M019220

NODE=M019R1

NODE=M019R1

NODE=M019R01

NODE=M019R01

NODE=M019R01;LINKAGE=LE

NODE=M019R00

NODE=M019R00

$K_0^*(1430)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\pi$	(93 ± 10 %)
$\Gamma_2 K\eta$	(8.6 \pm 2.7 3.4 %)
$\Gamma_3 K\eta'(958)$	seen

$K_0^*(1430)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	DOCUMENT ID
0.93\pm0.04\pm0.09	ASTON 88 TECN 0 COMMENT $11 K^- p \rightarrow K^- \pi^+ n$

$\Gamma(K\eta)/\Gamma(K\pi)$	Γ_2/Γ_1
VALUE (%)	EVTS DOCUMENT ID TECN COMMENT
9.2\pm2.5\pm1.0	5.4k ¹ LEES 14E BABR $\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a Dalitz analysis in the presence of several interfering scalar-meson resonances.

$\Gamma(K\eta'/(958))/\Gamma_{\text{total}}$	Γ_3/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	ABLIKIM 14J BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

K₀^{*}(1430) REFERENCES

ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55901
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)	REFID=53213
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=53056
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
LINK	07	PL B648 156	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51702
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51875
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51051
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=51458
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)	REFID=50996
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng		REFID=51198
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>		REFID=50165
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
LI	03	PR D67 034025	L. Li, B. Zou, G. Li		REFID=49192
AITALA	02	PRC 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48807
JAMIN	00	NP B587 331	M. Jamin <i>et al.</i>		REFID=47983
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46348
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev		REFID=45815
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=22446

NODE=M019

K₂^{*}(1430)

$$I(J^P) = \frac{1}{2}(2^+)$$

We consider that phase-shift analyses provide more reliable determinations of the mass and width.

NODE=M022

K₂^{*}(1430) MASS**CHARGED ONLY, WITH FINAL STATE K_π**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1427.3 ± 1.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below. [1425.6 ± 1.5 MeV OUR 2019 AVERAGE Scale factor = 1.1]			

NODE=M022205

NODE=M022M1
NODE=M022M1

NEW

1432.7 ± 0.7 ^{+2.2} _{-2.3}	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
1420 ± 4	1587	BAUBILLIER	84B HBC	—	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
1436 ± 5.5	400	1,2 CLELAND	82 SPEC	+	$30 K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1500	1,2 CLELAND	82 SPEC	+	$50 K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1200	1,2 CLELAND	82 SPEC	—	$50 K^+ p \rightarrow K_S^0 \pi^- p$
1423 ± 5	935	TOAFF	81 HBC	—	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
1428.0 ± 4.6		³ MARTIN	78 SPEC	+	$10 K^\pm p \rightarrow K_S^0 \pi p$
1423.8 ± 4.6		³ MARTIN	78 SPEC	—	$10 K^\pm p \rightarrow K_S^0 \pi p$
1420.0 ± 3.1	1400	AGUILAR-...	71B HBC	—	$3.9, 4.6 K^- p$
1425 ± 8.0	225	1,2 BARNHAM	71C HBC	+	$K^+ p \rightarrow K^0 \pi^+ p$
1416 ± 10	220	CRENNELL	69D DBC	—	$3.9 K^- N \rightarrow \bar{K}^0 \pi^- N$
1414 ± 13.0	60	¹ LIND	69 HBC	+	$9 K^+ p \rightarrow K^0 \pi^+ p$
1427 ± 12	63	¹ SCHWEING...	68 HBC	—	$5.5 K^- p \rightarrow \bar{K} \pi N$
1423 ± 11.0	39	¹ BASSANO	67 HBC	—	$4.6-5.0 K^- p \rightarrow \bar{K}^0 \pi^- p$

OCCUR=2


OCCUR=3

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1423.4 ± 2	± 3	24809 ± 820	⁴ BIRD	89 LASS	—	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
------------	-----	-------------	-------------------	---------	---	--

**NEUTRAL ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1432.4± 1.3 OUR AVERAGE				
1431.2± 1.8± 0.7	5	ASTON	88	LASS $11 K^- p \rightarrow K^- \pi^+ n$
1434 ± 4 ± 6	5	ASTON	87	LASS $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1433 ± 6 ± 10	5	ASTON	84B	LASS $11 K^- p \rightarrow \bar{K}^0 2\pi n$
1471 ± 12	5	BAUBILLIER	82B	HBC $8.25 K^- p \rightarrow N K_S^0 \pi \pi$
1428 ± 3	5	ASTON	81C	LASS $11 K^- p \rightarrow K^- \pi^+ n$
1434 ± 2	5	ESTABROOKS	78	ASPK $13 K^\pm p \rightarrow p K \pi$
1440 ± 10	5	BOWLER	77	DBC $5.5 K^+ d \rightarrow K \pi pp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1428.5± 3.9	1786± 127	AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
1420 ± 7	300	HENDRICK	76	DBC $8.25 K^+ N \rightarrow K^+ \pi N$
1421.6± 4.2	800	MCCUBBIN	75	HBC $3.6 K^- p \rightarrow K^- \pi^+ n$
1420.1± 4.3	7	LINGLIN	73	HBC $2-13 K^+ p \rightarrow K^+ \pi^- X$
1419.1± 3.7	1800	AGUILAR-...	71B	HBC $3.9, 4.6 K^- p$
1416 ± 6	600	CORDS	71	DBC $9 K^+ n \rightarrow K^+ \pi^- p$
1421.1± 2.6	2200	DAVIS	69	HBC $12 K^+ p \rightarrow K^+ \pi^- X$

NODE=M022M4
NODE=M022M4

¹ Errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

² Number of events in peak re-evaluated by us.

³ Systematic error added by us.

⁴ From a partial wave amplitude analysis.

⁵ From phase shift or partial-wave analysis.

⁶ Systematic errors not estimated.

⁷ From pole extrapolation, using world $K^+ p$ data summary tape.

NODE=M022M;LINKAGE=D
NODE=M022M;LINKAGE=W
NODE=M022M;LINKAGE=B
NODE=M022M;LINKAGE=F
NODE=M022M;LINKAGE=P
NODE=M022M4;LINKAGE=NS
NODE=M022M;LINKAGE=C

NODE=M022210

NODE=M022W1
NODE=M022W1
NEW

NEW

CHARGED ONLY, WITH FINAL STATE $K\pi$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
100.0± 2.1 OUR FIT					
[98.5 ± 2.7 MeV OUR 2019 FIT Scale factor = 1.1]					
100.0± 2.2 OUR AVERAGE					Error includes scale factor of 1.1. [98.5 ± 2.9 MeV OUR 2019 AVERAGE Scale factor = 1.1]
102.5± 1.6 ^{+3.1} _{-2.8}	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
109 ± 22	400	8, ⁹ CLELAND	82	SPEC +	$30 K^+ p \rightarrow K_S^0 \pi^+ p$
124 ± 12.8	1500	8, ⁹ CLELAND	82	SPEC +	$50 K^+ p \rightarrow K_S^0 \pi^+ p$
113 ± 12.8	1200	8, ⁹ CLELAND	82	SPEC -	$50 K^+ p \rightarrow K_S^0 \pi^- p$
85 ± 16	935	TOAFF	81	HBC -	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
96.5± 3.8		MARTIN	78	SPEC +	$10 K^\pm p \rightarrow K_S^0 \pi p$

OCCUR=2


OCCUR=3

97.7 ± 4.0	MARTIN	78	SPEC	—	$10 K^\pm p \rightarrow K_S^0 \pi p$	OCCUR=2
$94.7^{+15.1}_{-12.5}$	1400	AGUILAR-...	71B	HBC	—	$3.9, 4.6 K^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$98 \pm 4 \pm 4$	25k	10 BIRD	89	LASS	—	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
109 ± 5 OUR AVERAGE				Error includes scale factor of 1.9. See the ideogram below.	
$116.5 \pm 3.6 \pm 1.7$	11 ASTON	88	LASS	$11 K^- p \rightarrow K^- \pi^+ n$	
$129 \pm 15 \pm 15$	11 ASTON	87	LASS	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
$131 \pm 24 \pm 20$	11 ASTON	84B	LASS	$11 K^- p \rightarrow \bar{K}^0 2\pi n$	
143 ± 34	11 BAUBILLIER	82B	HBC	$8.25 K^- p \rightarrow N K_S^0 \pi \pi$	
98 ± 8	11 ASTON	81C	LASS	$11 K^- p \rightarrow K^- \pi^+ n$	
140 ± 30	11 ETKIN	80	SPEC	$6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
98 ± 5	11 ESTABROOKS 78	ASPK		$13 K^\pm p \rightarrow p K \pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
113.7 ± 9.2	1786 ± 127	12 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^* 0 K^\pm \pi^\mp \gamma$	
125 ± 29	300	8 HENDRICK	76	DBC	$8.25 K^+ N \rightarrow K^+ \pi N$
116 ± 18	800	MCCUBBIN	75	HBC	$3.6 K^- p \rightarrow K^- \pi^+ n$
61 ± 14	13 LINGLIN	73	HBC	$2-13 K^+ p \rightarrow K^+ \pi^- X$	
$116.6^{+10.3}_{-15.5}$	1800	AGUILAR-...	71B	HBC	$3.9, 4.6 K^- p$
144 ± 24.0	600	8 CORDS	71	DBC	$9 K^+ n \rightarrow K^+ \pi^- p$
101 ± 10	2200	DAVIS	69	HBC	$12 K^+ p \rightarrow K^+ \pi^- \pi^+ p$

WEIGHTED AVERAGE
109±5 (Error scaled by 1.9)



$K_2^*(1430)^0$ width (MeV)

⁸ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁹ Number of events in peak re-evaluated by us.

¹⁰ From a partial wave amplitude analysis.

¹¹ From phase shift or partial-wave analysis.

¹² Systematic errors not estimated.

¹³ From pole extrapolation, using world $K^+ p$ data summary tape.

NODE=M022W;LINKAGE=D
NODE=M022W;LINKAGE=W
NODE=M022W;LINKAGE=F
NODE=M022W;LINKAGE=P
NODE=M022W4;LINKAGE=NS
NODE=M022W;LINKAGE=C

NODE=M022215;NODE=M022

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 K \pi$	$(49.9 \pm 1.2) \%$	
$\Gamma_2 K^*(892) \pi$	$(24.7 \pm 1.5) \%$	
$\Gamma_3 K^*(892) \pi \pi$	$(13.4 \pm 2.2) \%$	
$\Gamma_4 K \rho$	$(8.7 \pm 0.8) \%$	S=1.2

DESIG=1
DESIG=2
DESIG=6
DESIG=3

Γ_5	$K\omega$	(2.9 ± 0.8) %		DESIG=4
Γ_6	$K^+\gamma$	(2.4 ± 0.5) $\times 10^{-3}$	S=1.1	DESIG=8
Γ_7	$K\eta$	($1.5^{+3.4}_{-1.0}$) $\times 10^{-3}$	S=1.3	DESIG=5
Γ_8	$K\omega\pi$	< 7.2 $\times 10^{-4}$	CL=95%	DESIG=7
Γ_9	$K^0\gamma$	< 9 $\times 10^{-4}$	CL=90%	DESIG=10; OUR EVAL; \rightarrow UNCHECKED \leftarrow

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 10 branching ratios uses 32 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 21.1$ for 25 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

	x_2	x_3	x_4	x_5	x_6	x_7
x_2	-9					
x_3	-40	-73				
x_4	-8	36	-52			
x_5	-11	-3	-26	-7		
x_6	-1	-1	-1	-1	0	
x_7	-4	-7	-5	-5	-2	0
Γ	0	0	0	0	0	-10
	x_1	x_2	x_3	x_4	x_5	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $K\pi$	49.9 ± 1.6	
Γ_2 $K^*(892)\pi$	24.7 ± 1.6	
Γ_3 $K^*(892)\pi\pi$	13.5 ± 2.3	
Γ_4 $K\rho$	8.7 ± 0.8	1.2
Γ_5 $K\omega$	2.9 ± 0.8	
Γ_6 $K^+\gamma$	0.24 ± 0.05	1.1
Γ_7 $K\eta$	$0.15^{+0.34}_{-0.10}$	1.3

$K_2^*(1430)$ PARTIAL WIDTHS

$\Gamma(K^+\gamma)$	Γ_6
VALUE (keV)	DOCUMENT ID
241 ± 50 OUR FIT	Error includes scale factor of 1.1.
240 ± 45	CIHANGIR 82 SPEC + 200 $K^+ Z \rightarrow Z K^+ \pi^0, Z K_S^0 \pi^+$

NODE=M022220

NODE=M022W8
NODE=M022W8

$\Gamma(K^0\gamma)$	Γ_9
VALUE (keV)	DOCUMENT ID
< 5.4	ALAVI-HARATI02B KTEV $K + A \rightarrow K^* + A$

* * * We do not use the following data for averages, fits, limits, etc. * * *

< 84	90 CARLSMITH 87 SPEC 0 $60-200 K_L^0 A \rightarrow K_S^0 \pi^0 A$
--------	---

NODE=M022W9
NODE=M022W9

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
0.499 ± 0.012 OUR FIT	
0.488 ± 0.014 OUR AVERAGE	
$0.485 \pm 0.006 \pm 0.020$ 14 ASTON 88 LASS 0 $11 K^- p \rightarrow K^- \pi^+ n$	
0.49 ± 0.02 14 ESTABROOKS 78 ASPK $\pm 13 K^\pm p \rightarrow p K\pi$	

NODE=M022225

NODE=M022R1
NODE=M022R1

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_2/Γ_1
0.496±0.034 OUR FIT					
0.47 ±0.04 OUR AVERAGE					
0.44 ±0.09	ASTON	84B	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
0.62 ±0.19	LAUSCHER	75	HBC	0	10,16 $K^- p \rightarrow K^- \pi^+ n$
0.54 ±0.16	DEHM	74	DBC	0	4.6 $K^+ N$
0.47 ±0.08	AGUILAR-...	71B	HBC		3.9,4.6 $K^- p$
0.47 ±0.10	BASSANO	67	HBC	-0	4.6,5.0 $K^- p$
0.45 ±0.13	BADIER	65C	HBC	-	3 $K^- p$

NODE=M022R4
NODE=M022R4

 $\Gamma(K\omega)/\Gamma(K\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_5/Γ_1
0.059±0.017 OUR FIT					
0.070±0.035 OUR AVERAGE					
0.05 ±0.04	AGUILAR-...	71B	HBC		3.9,4.6 $K^- p$
0.13 ±0.07	BASSOMPIE...	69	HBC	0	5 $K^+ p$

NODE=M022R5
NODE=M022R5

 $\Gamma(K\rho)/\Gamma(K\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ_1
0.174±0.017 OUR FIT	Error includes scale factor of 1.2.				

NODE=M022R6
NODE=M022R6

 $0.150^{+0.029}_{-0.017}$ OUR AVERAGE

0.18 ±0.05	ASTON	84B	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
0.02 +0.10 -0.02	DEHM	74	DBC	0	4.6 $K^+ N$
0.16 ±0.05	AGUILAR-...	71B	HBC		3.9,4.6 $K^- p$
0.14 ±0.10	BASSANO	67	HBC	-0	4.6,5.0 $K^- p$
0.14 ±0.07	BADIER	65C	HBC	-	3 $K^- p$

 $\Gamma(K\rho)/\Gamma(K^*(892)\pi)$


<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_4/Γ_2
0.350±0.031 OUR FIT	Error includes scale factor of 1.4.				

NODE=M022R7
NODE=M022R7

 $0.354±0.033$ OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

0.293±0.032±0.020	ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
0.38 ±0.09	BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow NK_S^0 \pi\pi$
0.39 ±0.03	DAUM	81C	CNTR		63 $K^- p \rightarrow K^- 2\pi p$

WEIGHTED AVERAGE
0.354±0.033 (Error scaled by 1.4)


 $\Gamma(K\omega)/\Gamma(K^*(892)\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	Γ_5/Γ_2
0.118±0.034 OUR FIT					
0.10 ±0.04 OUR AVERAGE	FIELD	67	HBC	-	3.8 $K^- p$

NODE=M022R8
NODE=M022R8

$\Gamma(K\eta)/\Gamma(K^*(892)\pi)$						Γ_7/Γ_2		
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			NODE=M022R9	NODE=M022R9
0.006^{+0.014}_{-0.004} OUR FIT				Error includes scale factor of 1.2.				
0.07 ±0.04	FIELD	67	HBC	—	3.8 $K^- p$			
$\Gamma(K\eta)/\Gamma(K\pi)$						Γ_7/Γ_1		
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT		NODE=M022R10	NODE=M022R10
0.0030^{+0.0070}_{-0.0020} OUR FIT				Error includes scale factor of 1.3.				
0 ±0.0056	15 ASTON	88B	LASS	—	11 $K^- p \rightarrow K^- \eta p$			
• • • We do not use the following data for averages, fits, limits, etc. • • •								
<0.04	95 AGUILAR-...	71B	HBC		3.9, 4.6 $K^- p$			
<0.065	16 BASSOMPIE...	69	HBC		5.0 $K^+ p$			
<0.02	BISHOP	69	HBC		3.5 $K^+ p$			
$\Gamma(K^*(892)\pi\pi)/\Gamma_{\text{total}}$						Γ_3/Γ		
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			NODE=M022R11	NODE=M022R11
0.134±0.022 OUR FIT								
0.12 ±0.04	17 GOLDBERG	76	HBC	—	3 $K^- p \rightarrow p\bar{K}^0 \pi\pi\pi$			
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$						Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			NODE=M022R12	NODE=M022R12
0.27±0.05 OUR FIT								
0.21±0.08	16,17 JONGEJANS	78	HBC	—	4 $K^- p \rightarrow p\bar{K}^0 \pi\pi\pi$			
$\Gamma(K\omega\pi)/\Gamma_{\text{total}}$						Γ_8/Γ		
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M022R13	NODE=M022R13
<0.72	95	0	JONGEJANS	78	HBC	4 $K^- p \rightarrow p\bar{K}^0 4\pi$		
14 From phase shift analysis.								
15 ASTON 888 quote < 0.0092 at CL=95%. We convert this to a central value and 1 sigma error in order to be able to use it in our constrained fit.								
16 Restated by us.								
17 Assuming $\pi\pi$ system has isospin 1, which is supported by the data.								
$K_2^*(1430)$ REFERENCES								
ABLIKIM	19AQ	PR D100	032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)		REFID=59909	
AUBERT	07AK	PR D76	012008	B. Aubert <i>et al.</i>	(BABAR Collab.)		REFID=51908	
ALAVI-HARATI	02B	PRL	89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)		REFID=48822	
BIRD	89	SLAC-332		P.F. Bird	(SLAC)		REFID=41002	
ASTON	88	NP B296	493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)		REFID=40262	
ASTON	88B	PL B201	169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)		REFID=40281	
ASTON	87	NP B292	693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)		REFID=40234	
CARLSMITH	87	PR D36	3502	D. Carlsmith <i>et al.</i>	(IFI, SACL)		REFID=40557	
ASTON	84B	NP B247	261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)		REFID=22763	
BAUBILLIER	84B	ZPHY C26	37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)		REFID=22459	
BAUBILLIER	82B	NP B202	21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)		REFID=22551	
CIHANGIR	82	PL 117B	123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)		REFID=21280	
CLELAND	82	NP B208	189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)		REFID=22455	
ASTON	81C	PL 106B	235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP		REFID=22821	
DAUM	81C	NP B187	1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)		REFID=22548	
TOAFF	81	PR D23	1500	S. Toaff <i>et al.</i>	(ANL, KANS)		REFID=22454	
ETKIN	80	PR D22	42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP		REFID=22545	
ESTABROOKS	78	NP B133	490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)		REFID=22443	
Also		PR D17	658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)		REFID=22444	
JONGEJANS	78	NP B139	383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)		REFID=22445	
MARTIN	78	NP B134	392	A.D. Martin <i>et al.</i>	(DURH, GEVA)		REFID=22446	
BOWLER	77	NP B126	31	M.G. Bowler <i>et al.</i>	(OXF)		REFID=22437	
GOLDBERG	76	LNC	17 253	J. Goldberg	(HAIF)		REFID=22742	
HENDRICK	76	NP B112	189	K. Hendrickx <i>et al.</i>	(MONS, SACL, PARIS+)		REFID=22743	
LAUSCHER	75	NP B86	189	P. Lauscher <i>et al.</i>	(ABCLV Collab.) JP		REFID=22582	
MCCUBBIN	75	NP B86	13	N.A. McCubbin, L. Lyons	(OXF)		REFID=22434	
DEHM	74	NP B75	47	G. Dehm <i>et al.</i>	(MPIM, BRUX, MONS, CERN)		REFID=22736	
LINGLIN	73	NP B55	408	D. Linglin	(CERN)		REFID=22428	
AGUILAR-...	71B	PR D4	2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)		REFID=22408	
BARNHAM	71C	NP B28	171	K.W.J. Barnham <i>et al.</i>	(BIRM, GLAS)		REFID=22409	
CORDS	71	PR D4	1974	D. Cords <i>et al.</i>	(PURD, UCD, IUPU)		REFID=22411	
BASSOMPIE...	69	NP B13	189	G. Bassompierre <i>et al.</i>	(CERN, BRUX) JP		REFID=22710	
BISHOP	69	NP B9	403	J.M. Bishop <i>et al.</i>	(WISC)		REFID=22485	
CRENNELL	69D	PRL	22 487	D.J. Crennell <i>et al.</i>	(BNL)		REFID=22399	
DAVIS	69	PRL	23 1071	P.J. Davis <i>et al.</i>	(LRL)		REFID=22400	
LIND	69	NP B14	1	V.G. Lind <i>et al.</i>	(LRL) JP		REFID=22404	
SCHWEING...	68	PR 166	1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)		REFID=22398	
Also		Thesis		F.L. Schweingruber	(NWES, NWES)		REFID=22709	
BASSANO	67	PRL	19 968	D. Bassano <i>et al.</i>	(BNL, SYRA)		REFID=22695	
FIELD	67	PL 24B	638	J.H. Field <i>et al.</i>	(UCSD)		REFID=22701	
BADIER	65C	PL 19	612	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)		REFID=22690	

K(1460)

$$I(J^P) = \frac{1}{2}(0^-)$$

OMMITTED FROM SUMMARY TABLE
Observed in $K\pi\pi$ partial-wave analysis.

K(1460) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1482.40 \pm 3.58 \pm 15.22	894k	AAIJ	18AI	LHCb	$D^0 \rightarrow K^\mp 2\pi^\pm \pi^\mp$
~ 1460	63	DAUM	81C	CNTR	$K^- p \rightarrow K^- 2\pi p$
~ 1400	13	¹ BRANDENB...	76B	ASPK	$K^\pm p \rightarrow K^\pm 2\pi p$
¹ Coupled mainly to $Kf_0(1370)$. Decay into $K^*(892)\pi$ seen.					

NODE=M021

NODE=M021M

NODE=M021M

NODE=M021M;LINKAGE=A

NODE=M021W

NODE=M021W

NODE=M021W;LINKAGE=A

NODE=M021215;NODE=M021

DESIG=1;OUR EST; \rightarrow UNCHECKED
DESIG=2;OUR EST; \rightarrow UNCHECKED
DESIG=3;OUR EST; \rightarrow UNCHECKED

NODE=M021220

NODE=M021W1
NODE=M021W1NODE=M021W2
NODE=M021W2NODE=M021W3
NODE=M021W3

NODE=M021

REFID=59187
REFID=22548
REFID=22767***K(1460) WIDTH***

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
335.60 \pm 6.20 \pm 8.65	894k	AAIJ	18AI	LHCb	$D^0 \rightarrow K^\mp 2\pi^\pm \pi^\mp$
~ 260	63	DAUM	81C	CNTR	$K^- p \rightarrow K^- 2\pi p$
~ 250	15	¹ BRANDENB...	76B	ASPK	$K^\pm p \rightarrow K^\pm 2\pi p$
¹ Coupled mainly to $Kf_0(1370)$. Decay into $K^*(892)\pi$ seen.					

K(1460) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K\rho$	seen
Γ_3 $K_0^*(1430)\pi$	seen

K(1460) PARTIAL WIDTHS

$\Gamma(K^*(892)\pi)$	Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •	
~ 109	DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K\rho)$	Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •	
~ 34	DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K_0^*(1430)\pi)$	Γ_3
• • • We do not use the following data for averages, fits, limits, etc. • • •	
~ 117	DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

K(1460) REFERENCES

AAIJ	18AI	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BRANDENB...	76B	PRL 36 1239	G.W. Brandenburg <i>et al.</i>	(SLAC) JP

NODE=M021

$K_2(1580)$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+ \pi^-$ system. Needs confirmation.

 $K_2(1580)$ MASS

<u>VALUE</u> (MeV)	<u>DOCUMENT ID</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1580	OTTER	79	— 10,14,16 $K^- p$

NODE=M039

 $K_2(1580)$ WIDTH

<u>VALUE</u> (MeV)	<u>DOCUMENT ID</u>	<u>CHG</u>	<u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 110	OTTER	79	— 10,14,16 $K^- p$

NODE=M039M

NODE=M039M

 $K_2(1580)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892)\pi$	seen
$\Gamma_2 K_2^*(1430)\pi$	possibly seen

NODE=M039W

NODE=M039W

 $K_2(1580)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
seen OTTER 79 HBC — 10,14,16 $K^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
possibly seen	GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
$\Gamma(K_2^*(1430)\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
possibly seen OTTER 79 HBC — 10,14,16 $K^- p$	

NODE=M039215; NODE=M039

DESIG=1

DESIG=2

NODE=M039220

NODE=M039R1

NODE=M039R1

NODE=M039R2

NODE=M039R2

 $K_2(1580)$ REFERENCES

GULER 11 PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
OTTER 79 NP B147 1	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP

NODE=M039

REFID=53668

REFID=22772

K(1630)

$$I(J^P) = \frac{1}{2}(?)$$

OMITTED FROM SUMMARY TABLE

Seen as a narrow peak, compatible with the experimental resolution, in the invariant mass of the $K_S^0\pi^+\pi^-$ system produced in π^-p interactions at high momentum transfers.

NODE=M160

K(1630) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1629±7	~ 75	KARNAUKHOV98	BC	$16.0 \pi^- p \rightarrow (K_S^0\pi^+\pi^-) X^+\pi^-X^0$

NODE=M160M

NODE=M160M

K(1630) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
16⁺¹⁹₋₁₆	~ 75	¹ KARNAUKHOV98	BC	$16.0 \pi^- p \rightarrow (K_S^0\pi^+\pi^-) X^+\pi^-X^0$

NODE=M160W

NODE=M160W

¹ Compatible with an experimental resolution of 14 ± 1 MeV.

NODE=M160W;LINKAGE=A

K(1630) DECAY MODES

Mode
$\Gamma_1 K_S^0\pi^+\pi^-$

DESIG=1

NODE=M160

REFID=46371

NODE=M099

K₁(1650)

$$I(J^P) = \frac{1}{2}(1^+)$$

OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems ($K^+\phi$, $K\pi\pi$) reported in partial-wave analysis in the 1600–1900 mass region.

NODE=M099

K₁(1650) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1672±50 OUR AVERAGE					Error includes scale factor of 1.1.
1793±59 ⁺¹⁵³ ₋₁₀₁	4289	¹ AAIJ	17C LHCb	B ⁺ → $J/\psi\phi K^+$	
1650±50		FRAME	86 OMEG +	13 K ⁺ p → $\phi K^+ p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 1840		ARMSTRONG	83 OMEG -	18.5 K ⁻ p → 3Kp	
~ 1800		DAUM	81C CNTR -	63 K ⁻ p → K ⁻ 2πp	

NODE=M099M

NODE=M099M

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 7.6 σ.

NODE=M099M;LINKAGE=A

K₁(1650) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
158± 50 OUR AVERAGE					
365±157 ⁺¹³⁸ ₋₂₁₅	4289	² AAIJ	17C LHCb	B ⁺ → $J/\psi\phi K^+$	
150± 50		FRAME	86 OMEG +	13 K ⁺ p → $\phi K^+ p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 250		DAUM	81C CNTR -	63 K ⁻ p → K ⁻ 2πp	

NODE=M099W

NODE=M099W

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 7.6 σ.

NODE=M099W;LINKAGE=A

K₁(1650) DECAY MODES

NODE=M099215;NODE=M099

Mode	
Γ_1	$K\pi\pi$
Γ_2	$K\phi$

K₁(1650) REFERENCES

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

K^{*}(1680)

$$I(J^P) = \frac{1}{2}(1^-)$$

K^{*}(1680) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1718±18 OUR AVERAGE					
1722±20 ⁺³³ ₋₁₀₉	4289	1 AAIJ	17C LHCb		$B^+ \rightarrow J/\psi\phi K^+$
1677±10± 32		ASTON	88 LASS 0	11	$K^- p \rightarrow K^- \pi^+ n$
1735±10± 20		ASTON	87 LASS 0	11	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1678±64		BIRD	89 LASS -	11	$K^- p \rightarrow \bar{K}^0 \pi^- p$
1800±70		ETKIN	80 MPS 0	6	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
~ 1650		ESTABROOKS	78 ASPK 0	13	$K^\pm p \rightarrow K^\pm \pi^\pm n$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 8.5 σ .

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
322±110 OUR AVERAGE					
Error includes scale factor of 4.2.					
354± 75 ⁺¹⁴⁰ ₋₁₈₁	4289	2 AAIJ	17C LHCb		$B^+ \rightarrow J/\psi\phi K^+$
205± 16± 34		ASTON	88 LASS 0	11	$K^- p \rightarrow K^- \pi^+ n$
423± 18± 30		ASTON	87 LASS 0	11	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
454±270		BIRD	89 LASS -	11	$K^- p \rightarrow \bar{K}^0 \pi^- p$
170± 30		ETKIN	80 MPS 0	6	$K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
250 to 300		ESTABROOKS	78 ASPK 0	13	$K^\pm p \rightarrow K^\pm \pi^\pm n$

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 8.5 σ .

K^{*}(1680) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(38.7±2.5) %
Γ_2 $K\rho$	(31.4 ^{+5.0} _{-2.1}) %
Γ_3 $K^*(892)\pi$	(29.9 ^{+2.2} _{-5.0}) %
Γ_4 $K\phi$	seen

DESIG=1

DESIG=2

NODE=M099

REFID=57657

REFID=57636

REFID=20569

REFID=22801

REFID=22548

NODE=M095

NODE=M095M

NODE=M095M

NODE=M095M;LINKAGE=A

NODE=M095W

NODE=M095W

NODE=M095W;LINKAGE=A

NODE=M095215;NODE=M095

DESIG=1

DESIG=3

DESIG=2

DESIG=4

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 2.9$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	−36			
x_3	−39	−72		
	x_1	x_2		

K*(1680) BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$

VALUE

0.387 ± 0.026 OUR FIT

0.388 ± 0.014 ± 0.022

DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-----	---------

ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
-------	----	------	---

Γ_1/Γ

NODE=M095220

NODE=M095R4

NODE=M095R4

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$

VALUE

1.30 ± 0.23 OUR FIT

2.8 ± 1.1

DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-----	---------

ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
-------	----	------	--

Γ_1/Γ_3

NODE=M095R2

NODE=M095R2

$\Gamma(K\rho)/\Gamma(K\pi)$

VALUE

0.81 ± 0.14 OUR FIT

1.2 ± 0.4

DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-----	---------

ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
-------	----	------	--

Γ_2/Γ_1

NODE=M095R3

NODE=M095R3

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$

VALUE

1.05 ± 0.27 OUR FIT

0.97 ± 0.09 ± 0.30

DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-----	---------

ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
-------	----	------	---

Γ_2/Γ_3

NODE=M095R1

NODE=M095R1

$\Gamma(K\phi)/\Gamma_{\text{total}}$

VALUE

seen

EVTS	DOCUMENT ID	TECN	COMMENT
------	-------------	------	---------

4289	3 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
------	--------	----------	-----------------------------------

Γ_4/Γ

NODE=M095R00

NODE=M095R00

3 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5 σ .

NODE=M095R00;LINKAGE=A

K*(1680) REFERENCES

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP

REFID=57657

REFID=57636

REFID=41002

REFID=40262

REFID=40234

REFID=22689

REFID=22545

REFID=22443

NODE=M095

NODE=M023

K₂(1770)

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this *Review*, PDG 04.***K₂(1770) MASS***

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1773± 8 OUR AVERAGE					
1777±35 ⁺¹²² ₋₇₇	4289	1 AAIJ	17C LHCb		$B^+ \rightarrow J/\psi \phi K^+$
1773± 8		2 ASTON	93 LASS		$11K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1743±15		TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1810±20		FRAME 86	OMEG +		$13 K^+ p \rightarrow \phi K^+ p$
~1730		ARMSTRONG 83	OMEG -		$18.5 K^- p \rightarrow 3K p$
~1780		3 DAUM 81C	CNTR -		$63 K^- p \rightarrow K^- 2\pi p$
1710±15	60	CHUNG 74	HBC -		$7.3 K^- p \rightarrow K^- \omega p$
1767± 6		BLIEDEN 72	MMS -		$11-16 K^- p$
1730±20	306	4 FIRESTONE 72B	DBC +		$12 K^+ d$
1765±40		5 COLLEY 71	HBC +		$10 K^+ p \rightarrow K 2\pi N$
1740		DENEGR 71	DBC -		$12.6 K^- d \rightarrow \bar{K} 2\pi d$
1745±20		AGUILAR-...	70C HBC	-	$4.6 K^- p$
1780±15		BARTSCH 70C	HBC -		$10.1 K^- p$
1760±15		LUDLAM 70	HBC -		$12.6 K^- p$

1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.0 σ .2 From a partial wave analysis of the $K^- \omega$ system.3 From a partial wave analysis of the $K^- 2\pi$ system.

4 Produced in conjunction with excited deuteron.

5 Systematic errors added correspond to spread of different fits.

NODE=M023

NODE=M023M

NODE=M023M

NODE=M023M;LINKAGE=C

NODE=M023M;LINKAGE=A

NODE=M023M;LINKAGE=B

NODE=M023M;LINKAGE=P

NODE=M023M;LINKAGE=X

K₂(1770) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
186± 14 OUR AVERAGE					
217±116 ⁺²²¹ ₋₁₅₄	4289	6 AAIJ	17C LHCb		$B^+ \rightarrow J/\psi \phi K^+$
186± 14		7 ASTON	93 LASS		$11K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
147± 70		TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
140± 40		FRAME 86	OMEG +		$13 K^+ p \rightarrow \phi K^+ p$
~220		ARMSTRONG 83	OMEG -		$18.5 K^- p \rightarrow 3K p$
~210		8 DAUM 81C	CNTR -		$63 K^- p \rightarrow K^- 2\pi p$
110± 50	60	CHUNG 74	HBC -		$7.3 K^- p \rightarrow K^- \omega p$
100± 26		BLIEDEN 72	MMS -		$11-16 K^- p$
210± 30	306	9 FIRESTONE 72B	DBC +		$12 K^+ d$
90± 70		10 COLLEY 71	HBC +		$10 K^+ p \rightarrow K 2\pi N$
130		DENEGR 71	DBC -		$12.6 K^- d \rightarrow \bar{K} 2\pi d$
100± 50		AGUILAR-...	70C HBC	-	$4.6 K^- p$
138± 40		BARTSCH 70C	HBC -		$10.1 K^- p$
50 ^{+ 40} _{- 20}		LUDLAM 70	HBC -		$12.6 K^- p$

6 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.0 σ .7 From a partial wave analysis of the $K^- \omega$ system.8 From a partial wave analysis of the $K^- 2\pi$ system.

9 Produced in conjunction with excited deuteron.

10 Systematic errors added correspond to spread of different fits.

NODE=M023W

NODE=M023W

NODE=M023W;LINKAGE=A

NODE=M023W;LINKAGE=B

NODE=M023W;LINKAGE=C

NODE=M023W;LINKAGE=P

NODE=M023W;LINKAGE=X

K₂(1770) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\pi\pi$	
$\Gamma_2 K_2^*(1430)\pi$	seen
$\Gamma_3 K^*(892)\pi$	seen
$\Gamma_4 Kf_2(1270)$	seen
$\Gamma_5 Kf_0(980)$	seen
$\Gamma_6 K\phi$	seen
$\Gamma_7 K\omega$	seen

K₂(1770) BRANCHING RATIOS

$$\frac{\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)}{(\Gamma_2/K_2^*(1430) \rightarrow K\pi)}$$

$$\frac{\Gamma_2/\Gamma_1}{\Gamma_1}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 0.03	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
~ 1.0	11 FIRESTONE	72B	DBC	+ 12 $K^+ d$
< 1.0	COLLEY	71	HBC	10 $K^+ p$
0.2 ± 0.2	AGUILAR-...	70C	HBC	- 4.6 $K^- p$
< 1.0	BARTSCH	70C	HBC	- 10.1 $K^- p$
1.0	BARBARO-...	69	HBC	+ 12.0 $K^+ p$

11 Produced in conjunction with excited deuteron.

$$\frac{\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)}{(\Gamma_3/K^*(892)\pi \rightarrow \pi\pi)}$$

$$\frac{\Gamma_3/\Gamma_1}{\Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.23	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$$\frac{\Gamma(Kf_2(1270))/\Gamma(K\pi\pi)}{(\Gamma_4/Kf_2(1270) \rightarrow \pi\pi)}$$

$$\frac{\Gamma_4/\Gamma_1}{\Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.74	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$$\frac{\Gamma(Kf_0(980))/\Gamma_{\text{total}}}{\text{VALUE}}$$

$$\frac{\Gamma_5/\Gamma}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
possibly seen	TIKHOMIROV 03	SPEC	40.0 $\pi^- C \xrightarrow{K_S^0 K_S^0 K_L^0 X}$

$$\frac{\Gamma(K\phi)/\Gamma_{\text{total}}}{\text{VALUE}}$$

$$\frac{\Gamma_6/\Gamma}{\Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	4289	12 AAIJ	17C	LHCb	$B^+ \rightarrow J/\psi\phi K^+$
seen		ARMSTRONG 83	OMEG	-	18.5 $K^- p \rightarrow K^- \phi N$

12 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.0 σ .

$$\frac{\Gamma(K\omega)/\Gamma_{\text{total}}}{\text{VALUE}}$$

$$\frac{\Gamma_7/\Gamma}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	OTTER	81	HBC	± 8.25, 10, 16 $K^\pm p$
seen	CHUNG	74	HBC	- 7.3 $K^- p \rightarrow K^- \omega p$

K₂(1770) REFERENCES

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860		
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
		NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+, CRAC, MPIM+)
ARMSTRONG	83	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
DAUM	81C	NP B181 1	G. Otter <i>et al.</i>	(AACH3, BERL, LOIC, VIEN, BIRM+)
OTTER	81	NP B181 1	H.R. Bliden <i>et al.</i>	(STON, NEAS)
CHUNG	74	PL 51B 413	S.U. Chung <i>et al.</i>	(BNL)
BLIEDEN	72	PL 39B 668	D.C. Colley <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(BIRM, GLAS)
COLLEY	71	NP B26 71	D. Denegri <i>et al.</i>	(JHU) JP
DENEGR	71	NP B28 13	M. Aguilar-Benitez <i>et al.</i>	(BNL)
AGUILAR-...	70C	PRL 25 54	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN+)
BARTSCH	70C	PL 33B 186	T. Ludlam, J. Sandweiss, A.J. Slaughter	(YALE)
LUDLAM	70	PR D2 1234	A. Barbaro-Galtieri <i>et al.</i>	(LRL)
BARBARO-...	69	PRL 22 1207		

NODE=M023215;NODE=M023

DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=2;OUR EST; \rightarrow UNCHECKED
 DESIG=4;OUR EST; \rightarrow UNCHECKED
 DESIG=9;OUR EST; \rightarrow UNCHECKED
 DESIG=11
 DESIG=10
 DESIG=8

NODE=M023220

NODE=M023R1
 NODE=M023R1
 NODE=M023R1

NODE=M023R1;LINKAGE=P

NODE=M023R3
 NODE=M023R3

NODE=M023R4
 NODE=M023R4
 NODE=M023R4

NODE=M023R6
 NODE=M023R6

NODE=M023R5
 NODE=M023R5

NODE=M023R5;LINKAGE=A

NODE=M023R2
 NODE=M023R2

NODE=M023

REFID=57657
 REFID=57636
 REFID=49653
 REFID=49423
 REFID=43597
 REFID=20569
 REFID=22801
 REFID=22548
 REFID=22549
 REFID=22735
 REFID=22788
 REFID=22506
 REFID=22785
 REFID=22497
 REFID=22782
 REFID=22783
 REFID=22784
 REFID=22483

NODE=M060

 $K_3^*(1780)$ $I(J^P) = \frac{1}{2}(3^-)$ **$K_3^*(1780)$ MASS**

NODE=M060M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1776 ± 7 OUR AVERAGE		Error includes scale factor of 1.1.			
$1781 \pm 8 \pm 4$		1 ASTON	88 LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
$1740 \pm 14 \pm 15$		1 ASTON	87 LASS	0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1779 ± 11		2 BALDI	76 SPEC	+	$10 K^+ p \rightarrow K^0 \pi^+ p$
1776 ± 26		3 BRANDENB...	76D ASPK	0	$13 K^\pm p \rightarrow K^\pm \pi^\mp N$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1720 \pm 10 \pm 15$	6111	4 BIRD	89 LASS	-	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
1749 ± 10		ASTON	88B LASS	-	$11 K^- p \rightarrow K^- \eta p$
1780 ± 9	300	BAUBILLIER	84B HBC	-	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
1790 ± 15		BAUBILLIER	82B HBC	0	$8.25 K^- p \rightarrow K_S^0 2\pi N$
1784 ± 9	2060	CLELAND	82 SPEC	\pm	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
1786 ± 15		5 ASTON	81D LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
1762 ± 9	190	TOAFF	81 HBC	-	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
1850 ± 50		ETKIN	80 MPS	0	$6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
1812 ± 28		BEUSCH	78 OMEG		$10 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1786 ± 8		CHUNG	78 MPS	0	$6 K^- p \rightarrow K^- \pi^+ n$

1 From energy-independent partial-wave analysis.

2 From a fit to Y_6^2 moment. $J^P = 3^-$ found.3 Confirmed by phase shift analysis of ESTABROOKS 78, yields $J^P = 3^-$.

4 From a partial wave amplitude analysis.

5 From a fit to the Y_6^0 moment.NODE=M060M;LINKAGE=K
NODE=M060M;LINKAGE=M
NODE=M060M;LINKAGE=A
NODE=M060M;LINKAGE=F
NODE=M060M;LINKAGE=J **$K_3^*(1780)$ WIDTH**

NODE=M060W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
159 ± 21 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
$203 \pm 30 \pm 8$		6 ASTON	88 LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
$171 \pm 42 \pm 20$		6 ASTON	87 LASS	0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
135 ± 22		7 BALDI	76 SPEC	+	$10 K^+ p \rightarrow K^0 \pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$187 \pm 31 \pm 20$	6111	8 BIRD	89 LASS	-	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
193^{+51}_{-37}		ASTON	88B LASS	-	$11 K^- p \rightarrow K^- \eta p$
99 ± 30	300	BAUBILLIER	84B HBC	-	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 130		BAUBILLIER	82B HBC	0	$8.25 K^- p \rightarrow K_S^0 2\pi N$
191 ± 24	2060	CLELAND	82 SPEC	\pm	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
225 ± 60		9 ASTON	81D LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
~ 80	190	TOAFF	81 HBC	-	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
240 ± 50		ETKIN	80 MPS	0	$6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
181 ± 44		10 BEUSCH	78 OMEG		$10 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
96 ± 31		CHUNG	78 MPS	0	$6 K^- p \rightarrow K^- \pi^+ n$
270 ± 70		11 BRANDENB...	76D ASPK	0	$13 K^\pm p \rightarrow K^\pm \pi^\mp N$

NODE=M060W

⁶ From energy-independent partial-wave analysis.

⁷ From a fit to Y_6^2 moment. $J^P = 3^-$ found.


⁸ From a partial wave amplitude analysis.

⁹ From a fit to Y_6^0 moment.

¹⁰ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹¹ ESTABROOKS 78 find that BRANDENBURG 76D data are consistent with 175 MeV width. Not averaged.

NODE=M060W;LINKAGE=K
NODE=M060W;LINKAGE=M
NODE=M060W;LINKAGE=F
NODE=M060W;LINKAGE=J
NODE=M060W;LINKAGE=D
NODE=M060W;LINKAGE=E



$K_3^*(1780)$ DECAY MODES

NODE=M060215;NODE=M060

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 K\rho$	(31 ± 9) %	
$\Gamma_2 K^*(892)\pi$	(20 ± 5) %	
$\Gamma_3 K\pi$	(18.8 ± 1.0) %	
$\Gamma_4 K\eta$	(30 ± 13) %	
$\Gamma_5 K_2^*(1430)\pi$	< 16 %	95%

DESIG=3
DESIG=2
DESIG=1
DESIG=6
DESIG=4

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 4 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|ccc} & x_2 & & \\ x_2 & 85 & & \\ & 18 & 21 & \\ x_3 & -98 & -94 & -27 \\ x_4 & & x_1 & x_2 & x_3 \end{array}$$

$K_3^*(1780)$ BRANCHING RATIOS

NODE=M060220

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	Γ_1/Γ_2
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
1.52 ± 0.23 OUR FIT	ASTON 87 LASS 0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M060R5
NODE=M060R5

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$						Γ_2/Γ_3
<u>VALUE</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
1.09±0.26 OUR FIT						
1.09±0.26	ASTON	84B	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$	
$\Gamma(K\pi)/\Gamma_{\text{total}}$						Γ_3/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
0.188±0.010 OUR FIT						
0.188±0.010 OUR AVERAGE						
0.187±0.008±0.008	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	
0.19 ± 0.02	ESTABROOKS	78	ASPK	0	13 $K^\pm p \rightarrow K\pi N$	
$\Gamma(K\eta)/\Gamma(K\pi)$						Γ_4/Γ_3
<u>VALUE</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	
1.6 ±0.7 OUR FIT						
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.41±0.050	12 BIRD	89	LASS	—	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
0.50±0.18	ASTON	88B	LASS	—	11 $K^- p \rightarrow K^- \eta p$	

12 This result supersedes ASTON 88B.

$\Gamma(K_2^*(1430)\pi)/\Gamma(K^*(892)\pi)$						Γ_5/Γ_2
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<0.78	95	ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$K_3^*(1780)$ REFERENCES

BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
ASTON	81D	PL 99B 502	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
BEUSCH	78	PL 74B 282	W. Beusch <i>et al.</i>	(CERN, AACB3, ETH) JP
CHUNG	78	PRL 40 355	S.U. Chung <i>et al.</i>	(BNL, BRAN, CUNY+) JP
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH-) JP
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH-) JP
BALDI	76	PL 63B 344	R. Baldi <i>et al.</i>	(GEVA) JP
BRANDENB...	76D	PL 60B 478	G.W. Brandenburg <i>et al.</i>	(SLAC) JP

NODE=M060R7
NODE=M060R7

NODE=M060R4
NODE=M060R4

NODE=M060R8
NODE=M060R8

NODE=M060R8;LINKAGE=H

NODE=M060R6
NODE=M060R6

NODE=M060

REFID=41002
REFID=40262
REFID=40281
REFID=40234
REFID=22763
REFID=22459
REFID=22551
REFID=22455
REFID=22820
REFID=22454
REFID=22545
REFID=22537
REFID=22814
REFID=22443
REFID=22444
REFID=22807
REFID=22808

$K_2(1820)$

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this Review (PDG 04)
under $K_2(1770)$.

 $K_2(1820)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1819 ± 12 OUR AVERAGE				
$1853 \pm 27^{+18}_{-35}$	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
1816 ± 13		² ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1840		³ DAUM	81C CNTR	$63K^- p \rightarrow K^- 2\pi p$
1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 σ .				
2 From a partial wave analysis of the $K^- \omega$ system.				
3 From a partial wave analysis of the $K^- 2\pi$ system.				

NODE=M146

NODE=M146M

NODE=M146M

NODE=M146M;LINKAGE=B
NODE=M146M;LINKAGE=A
NODE=M146M;LINKAGE=C

 $K_2(1820)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
264 ± 34 OUR AVERAGE				
$167 \pm 58^{+82}_{-72}$	4289	⁴ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
276 ± 35		⁵ ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 230		⁶ DAUM	81C CNTR	$63K^- p \rightarrow K^- 2\pi p$
4 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 σ .				
5 From a partial wave analysis of the $K^- \omega$ system.				
6 From a partial wave analysis of the $K^- 2\pi$ system.				

NODE=M146W

NODE=M146W

NODE=M146W;LINKAGE=A
NODE=M146W;LINKAGE=B
NODE=M146W;LINKAGE=C

NODE=M146215;NODE=M146

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi\pi$	
Γ_2 $K_2^*(1430)\pi$	seen
Γ_3 $K^*(892)\pi$	seen
Γ_4 $Kf_2(1270)$	seen
Γ_5 $K\omega$	seen
Γ_6 $K\phi$	seen

DESIG=5

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=6;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=7

NODE=M146220

NODE=M146R1
NODE=M146R1

NODE=M146R2
NODE=M146R2

NODE=M146R3
NODE=M146R3

NODE=M146R00
NODE=M146R00

NODE=M146R00;LINKAGE=A

$\Gamma(K^*(1430)\pi)/\Gamma(K\pi\pi)$	DOCUMENT ID	TECN	Γ_2/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.77	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K}2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$	DOCUMENT ID	TECN	Γ_3/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.05	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K}2\pi p$
$\Gamma(Kf_2(1270))/\Gamma(K\pi\pi)$	DOCUMENT ID	TECN	Γ_4/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.18	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K}2\pi p$
$\Gamma(K\phi)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	Γ_6/Γ
seen	4289	⁷ AAIJ	17C LHCb $B^+ \rightarrow J/\psi \phi K^+$
7 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 σ .			

K₂(1820) REFERENCES

AAIJ Also	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	04	PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
ASTON	93	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
DAUM	81C	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
		NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

NODE=M146

REFID=57657
 REFID=57636
 REFID=49653
 REFID=43597
 REFID=22548

K(1830)

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of $K\phi$ system. Needs confirmation.***K(1830) MASS***

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1874±43⁺⁵⁹₋₁₁₅	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1830 ARMSTRONG 83 OMEG – 18.5 $K^- p \rightarrow 3K p$ ² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.5 σ .

NODE=M088

NODE=M088

NODE=M088M

NODE=M088M

NODE=M088M;LINKAGE=A

NODE=M088W

NODE=M088W

NODE=M088W;LINKAGE=A

NODE=M088215;NODE=M088

DESIG=1

NODE=M088

REFID=57657
 REFID=57636
 REFID=22801

K(1830) DECAY MODES

Mode
$\Gamma_1 \quad K\phi$

K(1830) REFERENCES

AAIJ Also	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
ARMSTRONG	83	PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
		NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+) JP

$K_0^*(1950)$

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+$ system. Needs confirmation.

 $K_0^*(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1945±10±20	1 ASTON 88 LASS 0			11 $K^- p \rightarrow K^- \pi^+ n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1917±12	2 ZHOU 06 RVUE			$K^- p \rightarrow K^- \pi^+ n$
1820±40	3 ANISOVICH 97C RVUE			11 $K^- p \rightarrow K^- \pi^+ n$

1 We take the central value of the two solutions and the larger error given.

2 S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.

3 T-matrix pole. Reanalysis of ASTON 88 data.

 $K_0^*(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
201± 34±79	4 ASTON 88 LASS 0			11 $K^- p \rightarrow K^- \pi^+ n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

145± 38	5 ZHOU 06 RVUE			$K^- p \rightarrow K^- \pi^+ n$
250±100	6 ANISOVICH 97C RVUE			11 $K^- p \rightarrow K^- \pi^+ n$

4 We take the central value of the two solutions and the larger error given.

5 S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.

6 T-matrix pole. Reanalysis of ASTON 88 data.

 $K_0^*(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad K^- \pi^+$	(52±14) %

 $K_0^*(1950)$ BRANCHING RATIOS

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
0.52±0.08±0.12	7 ASTON 88 LASS 0			11 $K^- p \rightarrow K^- \pi^+ n$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

~0.60	8 ZHOU 06 RVUE			$K^- p \rightarrow K^- \pi^+ n$
-------	----------------	--	--	---------------------------------

7 We take the central value of the two solutions and the larger error given.

8 S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.

 $K_0^*(1950)$ REFERENCES

ZHOU 06 NP A775 212	Z.Y. Zhou, H.Q. Zheng
ANISOVICH 97C PL B413 137	A.V. Anisovich, A.V. Sarantsev
ASTON 88 NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)

NODE=M134

NODE=M134

NODE=M134M

NODE=M134M

NODE=M134M;LINKAGE=A

NODE=M134M;LINKAGE=ZU

NODE=M134M;LINKAGE=A1

NODE=M134W

NODE=M134W

NODE=M134W;LINKAGE=A

NODE=M134W;LINKAGE=ZU

NODE=M134W;LINKAGE=A1

NODE=M134215;NODE=M134

DESIG=1

NODE=M134220

NODE=M134R1

NODE=M134R1

NODE=M134R1;LINKAGE=A

NODE=M134R1;LINKAGE=ZU

NODE=M134

REFID=51198

REFID=45815

REFID=40262

$K_2^*(1980)$ $I(J^P) = \frac{1}{2}(2^+)$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M104

NODE=M104

NODE=M104M

NODE=M104M

NEW

 $K_2^*(1980)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1943±50 OUR AVERAGE					Error includes scale factor of 2.2. [1974 ± 26 MeV OUR 2019 AVERAGE]
1868± 8± 40 57	183k	ABLIKIM	19AQBES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
2073±94±245 240	4289	1AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$	
1973± 8± 25		ASTON	87 LASS	0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2020±20		TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1978±40	241	BIRD	89 LASS	-	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$

1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 σ .

NODE=M104M;LINKAGE=B

 $K_2^*(1980)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
307± 50 OUR AVERAGE					Error includes scale factor of 1.2. [376 ± 70 MeV OUR 2019 AVERAGE]
272± 24± 50 15	183k	ABLIKIM	19AQBES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
678±311±1153 559	4289	2AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$	
373± 33± 60		ASTON	87 LASS	0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
180± 70		TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
398± 47	241	BIRD	89 LASS	-	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$

2 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 σ .

NODE=M104W;LINKAGE=B

 $K_2^*(1980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892)\pi$	possibly seen
$\Gamma_2 K\rho$	possibly seen
$\Gamma_3 K f_2(1270)$	possibly seen
$\Gamma_4 K\phi$	seen

DESIG=2

DESIG=3

DESIG=4

DESIG=5

NODE=M104220

NODE=M104R01
NODE=M104R01NODE=M104R02
NODE=M104R02NODE=M104R1
NODE=M104R1NODE=M104R3
NODE=M104R3 **$K_2^*(1980)$ BRANCHING RATIOS**

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
$\Gamma(K\rho)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	DOCUMENT ID	TECN	CHG	Γ_2/Γ_1
1.49±0.24±0.09	ASTON	87 LASS	0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
$\Gamma(K f_2(1270))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
possibly seen	TIKHOMIROV 03	SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$

$\Gamma(K\phi)/\Gamma_{\text{total}}$				Γ_4/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	4289	3 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$	
3 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 σ .					

K₂^{*}(1980) REFERENCES

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
TIKHOMIROV	03	PAN 66 828 Translated from YAF 66 860.	G.D. Tikhomirov <i>et al.</i>	
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

K₄^{*}(2045)

$$I(J^P) = \frac{1}{2}(4^+)$$

K₄^{*}(2045) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2048⁺ 8₋₉ OUR AVERAGE		Error includes scale factor of 1.1. [2045 ± 9 MeV OUR 2019 AVERAGE Scale factor = 1.1]			
2090± 9 ⁺¹¹ ₋₂₉	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
2062± 14±13		¹ ASTON	86 LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
2039± 10	400	2,3 CLELAND	82 SPEC	±	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$
2070 ⁺¹⁰⁰ ₋₄₀		⁴ ASTON	81C LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2079± 7	431	TORRES	86 MPSF	400 pA → 4KX
2088± 20	650	BAUBILLIER	82 HBC	— 8.25 $K^- p \rightarrow K_S^0 \pi^- p$
2115± 46	488	CARMONY	77 HBC	0 9 $K^+ d \rightarrow K^+ \pi^s X$

1 From a fit to all moments.

2 From a fit to 8 moments.

3 Number of events evaluated by us.

4 From energy-independent partial-wave analysis.

NODE=M104R00

NODE=M104R00

NODE=M104R00;LINKAGE=A

NODE=M104

REFID=59909

REFID=57657

REFID=57636

REFID=53668

REFID=49423

REFID=41002

REFID=40234

NODE=M035

NODE=M035M

NODE=M035M

NEW

NODE=M035M;LINKAGE=E

NODE=M035M;LINKAGE=B

NODE=M035M;LINKAGE=W

NODE=M035M;LINKAGE=D

NODE=M035W

NODE=M035W

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
199⁺ 27 OUR AVERAGE		[198 ± 30 MeV OUR 2019 AVERAGE]			
201± 19 ⁺⁵⁷ ₋₁₇	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
221± 48±27		⁵ ASTON	86 LASS	0	$11 K^- p \rightarrow K^- \pi^+ n$
189± 35	400	6,7 CLELAND	82 SPEC	±	$50 K^+ p \rightarrow K_S^0 \pi^\pm p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

61± 58	431	TORRES	86 MPSF	400 pA → 4KX
170 ⁺¹⁰⁰ ₋₅₀	650	BAUBILLIER	82 HBC	— 8.25 $K^- p \rightarrow K_S^0 \pi^- p$
240 ⁺⁵⁰⁰ ₋₁₀₀		⁸ ASTON	81C LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
300± 200		CARMONY	77 HBC	0 9 $K^+ d \rightarrow K^+ \pi^s X$

5 From a fit to all moments.

6 From a fit to 8 moments.

7 Number of events evaluated by us.

8 From energy-independent partial-wave analysis.

NODE=M035W;LINKAGE=E

NODE=M035W;LINKAGE=B

NODE=M035W;LINKAGE=W

NODE=M035W;LINKAGE=D

***K₄**(2045) DECAY MODES**

NODE=M035215;NODE=M035

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K\pi$	(9.9±1.2) %
$\Gamma_2 K^*(892)\pi\pi$	(9 ± 5) %
$\Gamma_3 K^*(892)\pi\pi\pi$	(7 ± 5) %
$\Gamma_4 \rho K\pi$	(5.7±3.2) %
$\Gamma_5 \omega K\pi$	(5.0±3.0) %
$\Gamma_6 \phi K\pi$	(2.8±1.4) %
$\Gamma_7 \phi K^*(892)$	(1.4±0.7) %

***K₄**(2045) BRANCHING RATIOS**

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.099±0.012	ASTON 88 LASS 0 11 $K^- p \rightarrow K^- \pi^+ n$
$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$	Γ_2/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.89±0.53	BAUBILLIER 82 HBC — 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$	Γ_3/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.75±0.49	BAUBILLIER 82 HBC — 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(\rho K\pi)/\Gamma(K\pi)$	Γ_4/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.58±0.32	BAUBILLIER 82 HBC — 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(\omega K\pi)/\Gamma(K\pi)$	Γ_5/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
0.50±0.30	BAUBILLIER 82 HBC — 8.25 $K^- p \rightarrow p K_S^0 3\pi$
$\Gamma(\phi K\pi)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.028±0.014	9 TORRES 86 MPSF 400 pA → 4KX
$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.014±0.007	9 TORRES 86 MPSF 400 pA → 4KX

9 Error determination is model dependent.

***K₄**(2045) REFERENCES**

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
TORRES	86	PR D34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)

DESIG=1
DESIG=2
DESIG=5
DESIG=3
DESIG=4
DESIG=6
DESIG=7

NODE=M035220

NODE=M035R1
NODE=M035R1NODE=M035R2
NODE=M035R2NODE=M035R5
NODE=M035R5NODE=M035R3
NODE=M035R3NODE=M035R4
NODE=M035R4NODE=M035R6
NODE=M035R6NODE=M035R7
NODE=M035R7

NODE=M035R;LINKAGE=A

NODE=M035

REFID=59909
REFID=40262
REFID=22462
REFID=22845
REFID=22842
REFID=22455
REFID=22821
REFID=22811

$K_2(2250)$

$I(J^P) = \frac{1}{2}(2^-)$

OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems reported in the 2150–2260 MeV region, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the $J^P = 2^-$ wave.

$K_2(2250)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2247±17 OUR AVERAGE					
2200±40		¹ ARMSTRONG 83C	OMEG	—	18 $K^- p \rightarrow \Lambda \bar{p} X$
2235±50		¹ BAUBILLIER 81	HBC	—	8 $K^- p \rightarrow \Lambda \bar{p} X$
2260±20		¹ CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2280±20		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
2147± 4	37	CHLIAPNIK...	79	HBC	+ 32 $K^+ p \rightarrow \bar{\Lambda} p X$
2240±20	20	LISSAUER	70	HBC	9 $K^+ p$
$1 J^P = 2^-$ from moments analysis.					

NODE=M040

NODE=M040M

NODE=M040M

$K_2(2250)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
180±30 OUR AVERAGE					
150±30		² ARMSTRONG 83C	OMEG	—	18 $K^- p \rightarrow \Lambda \bar{p} X$
210±30		² CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
180±60		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
~ 200		² BAUBILLIER 81	HBC	—	8 $K^- p \rightarrow \Lambda \bar{p} X$
~ 40	37	CHLIAPNIK...	79	HBC	+ 32 $K^+ p \rightarrow \bar{\Lambda} p X$
80±20	20	LISSAUER	70	HBC	9 $K^+ p$
$2 J^P = 2^-$ from moments analysis.					

NODE=M040M;LINKAGE=Q

NODE=M040W

NODE=M040W

$K_2(2250)$ DECAY MODES

Mode
$\Gamma_1 K \pi \pi$
$\Gamma_2 K f_2(1270)$
$\Gamma_3 K^*(892) f_0(980)$
$\Gamma_4 p \bar{\Lambda}$

NODE=M040W;LINKAGE=Q

NODE=M040215;NODE=M040

DESIG=1

DESIG=3

DESIG=4

DESIG=2

NODE=M040

REFID=49423

REFID=22852

REFID=22850

REFID=22851

REFID=22849

REFID=22847

$K_2(2250)$ REFERENCES

TIKHOMIROV 03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>
	Translated from YAF 66 860.	
ARMSTRONG 83C	NP B227 365	T.A. Armstrong <i>et al.</i>
BAUBILLIER 81	NP B183 1	M. Baubillier <i>et al.</i>
CLELAND 81	NP B184 1	W.E. Cleland <i>et al.</i>
CHLIAPNIK...	NP B158 253	P.V. Chliapnikov <i>et al.</i>
LISSAUER 70	NP B18 491	D. Lissauer <i>et al.</i>

(BARI, BIRM, CERN+)
(BIRM, CERN, GLAS+) JP
(PITT, GEVA, LAUS+) JP
(CERN, BELG, MONS)
(LBL)

$K_3(2320)$

$I(J^P) = \frac{1}{2}(3^+)$

OMMITTED FROM SUMMARY TABLE

Seen in the $J^P = 3^+$ wave of the antihyperon-nucleon system.
Needs confirmation. **$K_3(2320)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2324±24 OUR AVERAGE				
2330±40	¹ ARMSTRONG 83C	OMEG	—	18 $K^- p \rightarrow \Lambda \bar{p} X$
2320±30	¹ CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$

 $1 J^P = 3^+$ from moments analysis.

NODE=M090

NODE=M090M

NODE=M090M

NODE=M090M;LINKAGE=P

NODE=M090W

NODE=M090W

NODE=M090W;LINKAGE=P

NODE=M090215;NODE=M090

 $K_3(2320)$ DECAY MODES

Mode
$\Gamma_1 p \bar{\Lambda}$

 $K_3(2320)$ REFERENCES

ARMSTRONG 83C	NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
CLELAND 81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+)

 $K_5^*(2380)$

$I(J^P) = \frac{1}{2}(5^-)$

OMMITTED FROM SUMMARY TABLE

Needs confirmation.

DESIG=1

NODE=M090

REFID=22852

REFID=22851

NODE=M098

NODE=M098

NODE=M098M

NODE=M098M

NODE=M098M;LINKAGE=E

NODE=M098W

NODE=M098W

NODE=M098W;LINKAGE=E

NODE=M098215;NODE=M098

 $K_5^*(2380)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2382±14±19	¹ ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

1 From a fit to all the moments.

 $K_5^*(2380)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
178±37±32	² ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

2 From a fit to all the moments.

 $K_5^*(2380)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K \pi$	(6.1±1.2) %

 $K_5^*(2380)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT
0.061±0.012	ASTON 88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

DESIG=1

NODE=M098220

NODE=M098R1

NODE=M098R1

K₅^{*(2380)} REFERENCES

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

K₄(2500)

$I(J^P) = \frac{1}{2}(4^-)$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

K₄(2500) MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2490±20	¹ CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda\bar{p}$

 $1 J^P = 4^-$ from moments analysis.**K₄(2500) WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~250	² CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda\bar{p}$

 $2 J^P = 4^-$ from moments analysis.**K₄(2500) DECAY MODES**

Mode
$\Gamma_1 p\bar{\Lambda}$

K₄(2500) REFERENCES

CLELAND	81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+)
---------	----	-----------	----------------------------	---------------------

K(3100)

$I^G(J^P) = ??(??)$

OMITTED FROM SUMMARY TABLE

Narrow peak observed in several ($\Lambda\bar{p}$ + pions) and ($\bar{\Lambda}p$ + pions) states in Σ^- Be reactions by BOURQUIN 86 and in np and nA reactions by ALEEV 93. Not seen by BOEHNLEIN 91. If due to strong decays, this state has exotic quantum numbers ($B=0, Q=+1, S=-1$ for $\Lambda\bar{p}\pi^+\pi^+$ and $I \geq 3/2$ for $\Lambda\bar{p}\pi^-$). Needs confirmation.

K(3100) MASS

VALUE (MeV)	DOCUMENT ID
≈ 3100 OUR ESTIMATE	

3-BODY DECAYS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3054±11 OUR AVERAGE			
3060± 7±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+$
3056± 7±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-$
3055± 8±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^-$
3045± 8±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^+$

4-BODY DECAYS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3059±11 OUR AVERAGE			
3067± 6±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
3060± 8±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$
3055± 7±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$
3052± 8±20	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3105±30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
3115±30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

NODE=M098

REFID=40262

REFID=22462

NODE=M091

NODE=M091

NODE=M091M

NODE=M091M

NODE=M091M;LINKAGE=R

NODE=M091W

NODE=M091W

NODE=M091W;LINKAGE=R

NODE=M091215;NODE=M091

DESIG=1

NODE=M091

REFID=22851

NODE=M129

NODE=M129

NODE=M129205

NODE=M129M
→ UNCHECKED ←NODE=M129M1
NODE=M129M1

OCCUR=2

OCCUR=3

OCCUR=4

NODE=M129M2
NODE=M129M2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

5-BODY DECAYS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3095±30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$
¹ Supersedes ALEEV 90.			

NODE=M129M3
NODE=M129M3

NODE=M129M;LINKAGE=A

NODE=M129210

NODE=M129W1
NODE=M129W1OCCUR=2
OCCUR=3
OCCUR=4NODE=M129W2
NODE=M129W2OCCUR=2
OCCUR=3
OCCUR=4

OCCUR=2

NODE=M129W3
NODE=M129W3

NODE=M129W;LINKAGE=A

NODE=M129215;NODE=M129

3-BODY DECAYS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
42±16	2 ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+$
36±15	2 ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-$
50±18	2 ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^-$
30±15	2 ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^+$

4-BODY DECAYS

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22±8	2 ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$	
28±12	2 ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$	
32±15	2 ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$	
30±15	2 ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$	
<30	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
<80	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

5-BODY DECAYS

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<30	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$
² Supersedes ALEEV 90.				

K(3100) DECAY MODES

Mode
$\Gamma_1 K(3100)^0 \rightarrow \Lambda\bar{p}\pi^+$
$\Gamma_2 K(3100)^{--} \rightarrow \Lambda\bar{p}\pi^-$
$\Gamma_3 K(3100)^- \rightarrow \Lambda\bar{p}\pi^+\pi^-$
$\Gamma_4 K(3100)^+ \rightarrow \Lambda\bar{p}\pi^+\pi^+$
$\Gamma_5 K(3100)^0 \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$
$\Gamma_6 K(3100)^0 \rightarrow \Sigma(1385)^+\bar{p}$

DESIG=1
DESIG=2
DESIG=3
DESIG=4
DESIG=5
DESIG=6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_1
<0.04	90	ALEEV 93	BIS2	$K(3100)^0 \rightarrow \Sigma(1385)^+\bar{p}$	

NODE=M129R1
NODE=M129R1**K(3100) REFERENCES**

ALEEV 93	PAN 56 1358	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
BOEHNLEIN 91	NPBPS B21 174	A. Boehnlein <i>et al.</i>	(FLOR, BNL, IND+)
ALEEV 90	ZPHY C47 533	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
BOURQUIN 86	PL B172 113	M.H. Bourquin <i>et al.</i>	(GEVA, RAL, HEIDP+)

NODE=M129
REFID=43668
REFID=41743
REFID=42173
REFID=22928

CHARMED MESONS ($C = \pm 1$)

$D^+ = c\bar{d}$, $D^0 = c\bar{u}$, $\bar{D}^0 = \bar{c}u$, $D^- = \bar{c}d$, similarly for D^* 's

$D^*(2007)^0$

$I(J^P) = \frac{1}{2}(1^-)$
 I , J , P need confirmation.

J consistent with 1, value 0 ruled out (NGUYEN 77).

$D^*(2007)^0$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2006.85±0.05 OUR FIT	Error includes scale factor of 1.1.		
• • • We do not use the following data for averages, fits, limits, etc. • • •			

2006 ± 1.5 ¹ GOLDHABER 77 MRK1 e^+e^-

¹ From simultaneous fit to $D^*(2010)^+$, $D^*(2007)^0$, D^+ , and D^0 .

$m_{D^*(2007)^0} - m_{D^0}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
142.014±0.030 OUR FIT	Error includes scale factor of 1.5.			
142.016±0.030 OUR AVERAGE	Error includes scale factor of 1.5.			
142.007 $\pm 0.015 \pm 0.014$	10K	² TOMARADZE 15 CLEO $e^+e^- \rightarrow$ hadrons		
142.2 ± 0.3	± 0.2 145	ALBRECHT 95F ARG $e^+e^- \rightarrow$ hadrons		
142.12 ± 0.05	± 0.05 1176	BORTOLETTO92B CLE2 $e^+e^- \rightarrow$ hadrons		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
142.2 ± 2.0		SADROZINSKI 80 CBAL $D^{*0} \rightarrow D^0\pi^0$		
142.7 ± 1.7		³ GOLDHABER 77 MRK1 e^+e^-		
2 Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. This value comes from the average of the results for two decay modes, $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$.				
3 From simultaneous fit to $D^*(2010)^+$, $D^*(2007)^0$, D^+ , and D^0 .				

$D^*(2007)^0$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	⁴ ABACHI 88B HRS		$D^{*0} \rightarrow D^+\pi^-$

⁴ Assuming $m_{D^{*0}} = 2007.2 \pm 2.1$ MeV/ c^2 .

$D^*(2007)^0$ DECAY MODES

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0\pi^0$	(64.7 \pm 0.9) %
$\Gamma_2 D^0\gamma$	(35.3 \pm 0.9) %

NODE=MXXX035

NODE=MXXX035

NODE=M061

NODE=M061

NODE=M061M

NODE=M061M

NODE=M061M

NODE=M061M;LINKAGE=G

NODE=M061DM

NODE=M061DM

NODE=M061DM

NODE=M061DM;LINKAGE=A

NODE=M061DM;LINKAGE=G

NODE=M061W

NODE=M061W

NODE=M061W;LINKAGE=A

NODE=M061220;NODE=M061

NODE=M061

DESIG=1

DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 5 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 2.5$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100
x_1	

$D^*(2007)^0$ BRANCHING RATIOS

$\Gamma(D^0\pi^0)/\Gamma(D^0\gamma)$	Γ_1/Γ_2			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.83 ± 0.07 OUR FIT		Error includes scale factor of 1.1.		
1.85 ± 0.07 OUR AVERAGE				
$1.90 \pm 0.07 \pm 0.05$	4.9k	ABLIKIM	15B BES3	$10.6 e^+ e^- \rightarrow \text{hadrons}$
$1.74 \pm 0.02 \pm 0.13$		AUBERT,BE	05G BABR	$10.6 e^+ e^- \rightarrow \text{hadrons}$

NODE=M061225

$\Gamma(D^0\pi^0)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.647 ± 0.009 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.655 \pm 0.008 \pm 0.005$	3.2k	5 ABLIKIM	15B BES3	$e^+ e^- \rightarrow \text{hadrons}$
$0.635 \pm 0.003 \pm 0.017$	69k	5 AUBERT,BE	05G BABR	$10.6 e^+ e^- \rightarrow \text{hadrons}$
$0.596 \pm 0.035 \pm 0.028$	858	6 ALBRECHT	95F ARG	$e^+ e^- \rightarrow \text{hadrons}$
$0.636 \pm 0.023 \pm 0.033$	1097	6 BUTLER	92 CLE2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M061R3
NODE=M061R3

$\Gamma(D^0\gamma)/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.353 ± 0.009 OUR FIT				
0.381 ± 0.029 OUR AVERAGE				
$0.404 \pm 0.035 \pm 0.028$	456	6 ALBRECHT	95F ARG	$e^+ e^- \rightarrow \text{hadrons}$
$0.364 \pm 0.023 \pm 0.033$	621	6 BUTLER	92 CLE2	$e^+ e^- \rightarrow \text{hadrons}$
$0.37 \pm 0.08 \pm 0.08$		ADLER	88D MRK3	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.345 \pm 0.008 \pm 0.005$	1.8k	5 ABLIKIM	15B BES3	$e^+ e^- \rightarrow \text{hadrons}$
$0.365 \pm 0.003 \pm 0.017$	68k	5 AUBERT,BE	05G BABR	$10.6 e^+ e^- \rightarrow \text{hadrons}$
0.47 ± 0.23		LOW	87 HRS	29 GeV $e^+ e^-$
0.53 ± 0.13		BARTEL	85G JADE	$e^+ e^-$, hadrons
0.47 ± 0.12		COLES	82 MRK2	$e^+ e^-$
0.45 ± 0.15		GOLDHABER	77 MRK1	$e^+ e^-$

NODE=M061R1
NODE=M061R1

5 Derived from the ratio $\Gamma(D^0\pi^0) / \Gamma(D^0\gamma)$ assuming that the branching fractions of $D^{*0} \rightarrow D^0\pi^0$ and $D^{*0} \rightarrow D^0\gamma$ decays sum to 100%

NODE=M061R;LINKAGE=AU

6 The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

NODE=M061R;LINKAGE=A

$D^*(2007)^0$ REFERENCES

ABLIKIM	15B	PR D91 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
TOMARADZE	15	PR D91 011102	A. Tomaradze <i>et al.</i>	(NWES)
AUBERT,BE	05G	PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92B	PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)
ABACHI	88B	PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)
ADLER	88D	PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)
LOW	87	PL B183 232	E.H. Low <i>et al.</i>	(HRS Collab.)
BARTEL	85G	PL 161B 197	W. Bartel <i>et al.</i>	(JADE Collab.)
COLES	82	PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)
SADROZINSKI	80	Madison Conf. 681	H.F.W. Sdrozinski <i>et al.</i>	(PRIN, CIT+)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
NGUYEN	77	PRL 39 262	H.K. Nguyen <i>et al.</i>	(LBL, SLAC) J

NODE=M061

REFID=56375
REFID=57142
REFID=50942
REFID=44374
REFID=43116
REFID=43170
REFID=40584
REFID=40579
REFID=40017
REFID=22880
REFID=22866
REFID=22877
REFID=11434
REFID=11543

NODE=M062

 $D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

 $D^*(2010)^\pm$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^{*(2460)}{}^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

2010.26±0.05 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2008 ±3	¹ GOLDHABER 77	MRK1	±	e^+e^-
2008.6 ±1.0	² PERUZZI 77	LGW	±	e^+e^-

¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 ; not independent of FELDMAN 77B mass difference below.

² PERUZZI 77 mass not independent of FELDMAN 77B mass difference below and PERUZZI 77 D^0 mass value.

NODE=M062M

NODE=M062M

NODE=M062M

 $m_{D^*(2010)^+} - m_{D^+}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^{*(2460)}{}^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
140.603 ±0.015 OUR FIT				
140.602 ±0.014 OUR AVERAGE				
140.6010 ±0.0068 ±0.0129	151k	LEES	17F	BABR $e^+e^- \rightarrow$ hadrons
140.64 ±0.08 ±0.06	620	BORTOLETTO92B	CLE2	$e^+e^- \rightarrow$ hadrons

NODE=M062MD

NODE=M062MD

NODE=M062MD

 $m_{D^*(2010)^+} - m_{D^0}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^{*(2460)}{}^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
145.4257 ±0.0017 OUR FIT				
145.4258 ±0.0020 OUR AVERAGE				Error includes scale factor of 1.2.
145.4259 ±0.0004 ±0.0017	312.8k	LEES	13X	BABR $D^{*\pm} \rightarrow D^0\pi^\pm$ $(K\pi, K3\pi)\pi^\pm$
145.412 ±0.002 ±0.012		ANASTASSOV	02	CLE2 $D^{*\pm} \rightarrow D^0\pi^\pm$ $(K\pi)\pi^\pm$
145.54 ±0.08	611	³ ADINOLFI	99	BEAT $D^{*\pm} \rightarrow D^0\pi^\pm$
145.45 ±0.02		³ BREITWEG	99	ZEUS $D^{*\pm} \rightarrow D^0\pi^\pm$ $(K\pi)\pi^\pm$
145.42 ±0.05		³ BREITWEG	99	ZEUS $D^{*\pm} \rightarrow D^0\pi^\pm$ $(K^-3\pi)\pi^\pm$
145.5 ±0.15	103	⁴ ADLOFF	97B	H1 $D^{*\pm} \rightarrow D^0\pi^\pm$
145.44 ±0.08	152	⁴ BREITWEG	97	ZEUS $D^{*\pm} \rightarrow D^0\pi^\pm$,
145.42 ±0.11	199	⁴ BREITWEG	97	ZEUS $D^{*\pm} \rightarrow D^0\pi^\pm$ $D^0 \rightarrow K^-3\pi$ $D^0 \rightarrow K^-\pi^+$
145.4 ±0.2	48	⁴ DERRICK	95	ZEUS $D^{*\pm} \rightarrow D^0\pi^\pm$
145.39 ±0.06 ±0.03		BARLAG	92B	ACCM π^- 230 GeV
145.5 ±0.2	115	⁴ ALEXANDER	91B	OPAL $D^{*\pm} \rightarrow D^0\pi^\pm$
145.30 ±0.06		⁴ DECAMP	91J	ALEP $D^{*\pm} \rightarrow D^0\pi^\pm$
145.40 ±0.05 ±0.10		ABACHI	88B	HRS $D^{*\pm} \rightarrow D^0\pi^\pm$
145.46 ±0.07 ±0.03		ALBRECHT	85F	ARG $D^{*\pm} \rightarrow D^0\pi^+$
145.5 ±0.3	28	BAILEY	83	SPEC $D^{*\pm} \rightarrow D^0\pi^\pm$
145.5 ±0.3	60	FITCH	81	SPEC $\pi^- A$
145.3 ±0.5	30	FELDMAN	77B	MRK1 $D^{*+} \rightarrow D^0\pi^+$

NODE=M062DM

NODE=M062DM

NODE=M062DM

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

145.4256 $\pm 0.0006 \pm 0.0017$	138.5k	LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- \pi^+) \pi^\pm$
145.4266 $\pm 0.0005 \pm 0.0019$	174.3k	LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- 2\pi^+ \pi^-) \pi^\pm$
145.44 ± 0.09	122	⁴ BREITWEG	97B ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm, D^0 \rightarrow K^- \pi^+$
145.8 ± 1.5	16	AHLEN	83 HRS	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.1 ± 1.8	12	BAILEY	83 SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.1 ± 0.5	14	BAILEY	83 SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.5 ± 0.5	14	YELTON	82 MRK2	$29 e^+ e^- \rightarrow K^- \pi^+$
~ 145.5		AVERY	80 SPEC	γA
145.2 ± 0.6	2	BLIETSCHAU	79 BEBC	νp

³ Statistical errors only.

⁴ Systematic error not evaluated.

OCCUR=2

OCCUR=3

NODE=M062DM;LINKAGE=AV
NODE=M062DM;LINKAGE=A

$m_{D^*(2010)^+} - m_{D^*(2007)^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.6 ± 1.8	⁵ PERUZZI	77 LGW	$e^+ e^-$
5 Not independent of FELDMAN 77B mass difference above, PERUZZI 77 D^0 mass, and GOLDHABER 77 $D^*(2007)^0$ mass.			

NODE=M062EM

NODE=M062EM

NODE=M062EM;LINKAGE=P

$D^*(2010)^{\pm}$ WIDTH

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
83.4 ± 1.8 OUR AVERAGE					
83.3 $\pm 1.2 \pm 1.4$	312.8k	⁶ LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K\pi, K3\pi) \pi^\pm$	
96 $\pm 4 \pm 22$		⁶ ANASTASSOV	02 CLE2	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K\pi) \pi^\pm$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
83.4 $\pm 1.7 \pm 1.5$	138.5k	⁶ LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- \pi^+) \pi^\pm$	
83.2 $\pm 1.5 \pm 2.6$	174.3k	⁶ LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- 2\pi^+ \pi^-) \pi^\pm$	
<131	90	110	BARLAG	92B ACCM	$\pi^- 230 \text{ GeV}$
6 Ignoring the electromagnetic contribution from $D^{*\pm} \rightarrow D^\pm \gamma$.					

NODE=M062W

NODE=M062W

OCCUR=3

OCCUR=2

NODE=M062W;LINKAGE=LE

NODE=M062225;NODE=M062

NODE=M062

DESIG=1

DESIG=3

DESIG=2

$D^*(2010)^{\pm}$ DECAY MODES

$D^*(2010)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0 \pi^+$	(67.7 ± 0.5) %
$\Gamma_2 D^+ \pi^0$	(30.7 ± 0.5) %
$\Gamma_3 D^+ \gamma$	(1.6 ± 0.4) %

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 6 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.3$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{cc|cc} & & & \\ x_2 & & -62 & \\ & & -43 & -44 \\ & & \hline x_3 & & x_1 & x_2 \end{array}$$

$D^*(2010)^+$ BRANCHING RATIOS **$\Gamma(D^0\pi^+)/\Gamma_{\text{total}}$**

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.677 ± 0.005 OUR FIT				
0.677 ± 0.006 OUR AVERAGE				
0.6759 ± 0.0029 ± 0.0064	7,8,9 BARTELT	98 CLE2	$e^+ e^-$	
0.688 ± 0.024 ± 0.013	ALBRECHT	95F ARG	$e^+ e^- \rightarrow \text{hadrons}$	
0.681 ± 0.010 ± 0.013	7 BUTLER	92 CLE2	$e^+ e^- \rightarrow \text{hadrons}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.57 ± 0.04 ± 0.04	ADLER	88D MRK3	$e^+ e^-$	
0.44 ± 0.10	COLES	82 MRK2	$e^+ e^-$	
0.6 ± 0.15	9 GOLDHABER	77 MRK1	$e^+ e^-$	

NODE=M062230

NODE=M062R1

NODE=M062R1

 $\Gamma(D^+\pi^0)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.307 ± 0.005 OUR FIT					
0.3073 ± 0.0013 ± 0.0062					
0.312 ± 0.011 ± 0.008	1404 ALBRECHT	95F ARG	$e^+ e^- \rightarrow \text{hadrons}$		
0.308 ± 0.004 ± 0.008	410 7 BUTLER	92 CLE2	$e^+ e^- \rightarrow \text{hadrons}$		
0.26 ± 0.02 ± 0.02	ADLER	88D MRK3	$e^+ e^-$		
0.34 ± 0.07	COLES	82 MRK2	$e^+ e^-$		

NODE=M062R3

NODE=M062R3

 $\Gamma(D^+\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.016 ± 0.004 OUR FIT						
0.016 ± 0.005 OUR AVERAGE						
0.0168 ± 0.0042 ± 0.0029			7,8 BARTELT	98 CLE2	$e^+ e^-$	
0.011 ± 0.014 ± 0.016		12	7 BUTLER	92 CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

NODE=M062R2

NODE=M062R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.052	90	ALBRECHT	95F ARG	$e^+ e^- \rightarrow \text{hadrons}$
0.17 ± 0.05 ± 0.05		ADLER	88D MRK3	$e^+ e^-$
0.22 ± 0.12		10 COLES	82 MRK2	$e^+ e^-$

7 The branching ratios are not independent, they have been constrained by the authors to sum to 100%.

8 Systematic error includes theoretical error on the prediction of the ratio of hadronic modes.

9 Assuming that isospin is conserved in the decay.

10 Not independent of $\Gamma(D^0\pi^+)/\Gamma_{\text{total}}$ and $\Gamma(D^+\pi^0)/\Gamma_{\text{total}}$ measurement.

NODE=M062R;LINKAGE=A

NODE=M062R;LINKAGE=B

NODE=M062R;LINKAGE=G

NODE=M062R;LINKAGE=C

 $D^*(2010)^\pm$ REFERENCES

LEES	17F	PRL 119 202003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58277
LEES	13X	PRL 111 111801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55564
Also		PR D88 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55547
Also		PR D88 079902 (errat.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55695
ANASTASSOV	02	PR D65 032003	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=48550
ADINOLFI	99	NP B547 3	M. Adinolfi <i>et al.</i>	(Beatrice Collab.)	REFID=46925
BREITWEG	99	EPJ C6 67	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=46604
BARTEL	98	PRL 80 3919	J. Bartelt <i>et al.</i>	(CLEO Collab.)	REFID=46349
ADLOFF	97B	ZPHY C72 593	C. Adloff <i>et al.</i>	(H1 Collab.)	REFID=45421
BREITWEG	97	PL B401 192	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45520
BREITWEG	97B	PL B407 402	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45699
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44374
DERRICK	95	PL B349 225	M. Derrick <i>et al.</i>	(ZEUS Collab.)	REFID=44373
BARLAG	92B	PL B278 480	S. Barlag <i>et al.</i>	(ACCMOR Collab.)	REFID=42174
BORTOLETTO	92B	PRL 69 2046	D. Bortolotto <i>et al.</i>	(CLEO Collab.)	REFID=43116
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43170
ALEXANDER	91B	PL B262 341	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=41553
DECAMP	91J	PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)	REFID=41614
ABACHI	88B	PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)	REFID=40584
ADLER	88D	PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40579
ALBRECHT	85F	PL 150B 235	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=11527
AHLEN	83	PRL 51 1147	S.P. Ahlen <i>et al.</i>	(ANL, IND, LBL+)	REFID=22868
BAILEY	83	PL 132B 230	R. Bailey <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)	REFID=22870
COLES	82	PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)	REFID=22866
YELTON	82	PRL 49 430	J.M. Yelton <i>et al.</i>	(SLAC, LBL, UCB+)	REFID=22867
FITCH	81	PRL 46 761	V.L. Fitch <i>et al.</i>	(PRIN, SACL, TORI+)	REFID=22863
AVERY	80	PRL 44 1309	P. Avery <i>et al.</i>	(ILL, FNAL, COLU)	REFID=11498
BLIETSCHAU	79	PL 86B 108	J. Blietschau <i>et al.</i>	(AACH3, BONN, CERN+)	REFID=22861
FELDMAN	77B	PRL 38 1313	G.J. Feldman <i>et al.</i>	(Mark I Collab.)	REFID=22858
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)	REFID=11434
PERUZZI	77	PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)	REFID=11435

NODE=M062

$D_0^*(2300)^0$

$$I(J^P) = \frac{1}{2}(0^+)$$

was $D_0^*(2400)^0$

$J^P = 0^+$ assignment favored (ABE 04D).

$D_0^*(2300)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2300±19 OUR AVERAGE				
2297± 8±20	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
2308±17±32		ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2407±21±35	9.8k	1 LINK	04A FOCS	γ A

¹ Possibly the feed-down from another state.

NODE=M178

NODE=M178M

NODE=M178M

NODE=M178M;LINKAGE=A

NODE=M178W

NODE=M178W

NODE=M178W;LINKAGE=A

NODE=M178215;NODE=M178

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M178

REFID=52941
REFID=50011
REFID=49775

NODE=M179

$D_0^*(2300)^{\pm}$

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

was $D_0^*(2400)^{\pm}$

J, P need confirmation.

NODE=M179

NODE=M179M

NODE=M179M

OCCUR=2

$D_0^*(2300)^{\pm}$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2349± 7 OUR AVERAGE				
2360±15±30		1 AAIJ	15X LHCb	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2349± 6± 4		2 AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2354± 7±11		3 AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2403±14±35	18.8k	4 LINK	04A FOCS	γ A

- 1 From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S-wave and the $D\pi$ S- and P-waves.
 2 Modeling the $\pi^+\pi^-$ S-wave with the Isobar formalism.
 3 Modeling the $\pi^+\pi^-$ S-wave with the K-matrix formalism.
 4 Possibly the feed-down from another state.

NODE=M179M;LINKAGE=A

NODE=M179M;LINKAGE=B

NODE=M179M;LINKAGE=C

NODE=M179M;LINKAGE=D

$D_0^*(2300)^\pm$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
221±18 OUR AVERAGE				
255±26±51	1	AAIJ	15X LHCb	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
217±13±13	2	AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
230±15±21	3	AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
283±24±34	18.8k	4 LINK	04A FOCUS	γA
1 From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S-wave and the $D\pi$ S- and P-waves. 2 Modeling the $\pi^+\pi^-$ S-wave with the Isobar formalism. 3 Modeling the $\pi^+\pi^-$ S-wave with the K-matrix formalism. 4 Possibly the feed-down from another state.				

NODE=M179W

NODE=M179W

OCCUR=2

NODE=M179W;LINKAGE=A

NODE=M179W;LINKAGE=B

NODE=M179W;LINKAGE=C

NODE=M179W;LINKAGE=D

$D_0^*(2300)^\pm$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0 \pi^+$	seen

NODE=M179215;NODE=M179

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M179

REFID=56588

REFID=56609

REFID=49775

NODE=M097

 $D_1(2420)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

$D_1(2420)^0$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^{*}(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.


NODE=M097M

NODE=M097M

NODE=M097M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2420.8±0.5 OUR FIT Error includes scale factor of 1.3.				
2420.5±0.6 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
2419.6±0.1±0.7	210k	AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$
2423.1±1.5±0.4	2.7k	1 ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2420.1±0.1±0.8	103k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2426 ± 3 ± 1	151	ABE	05A BELL	$B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
2421.4±1.5±0.9	2 ABE		04D BELL	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2421 $\frac{+1}{-2}$ ± 2	286	AVERY	94C CLE2	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2422 ± 2 ± 2	51	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} \pi^- X$
2428 ± 3 ± 2	279	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2414 ± 2 ± 5	171	ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2428 ± 8 ± 5	171	ANJOS	89C TPS	$\gamma N \rightarrow D^{*+} \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2420.5±2.1±0.9	3110±340	3 CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
2421.7±0.7±0.6	7.5k	ABULENCIA	06A CDF	$1900 p\bar{p} \rightarrow D^{*+} \pi^- X$
2425 ± 3	235	4 ABREU	98M DLPH	$e^+ e^-$

- ¹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions, and A_{D_2} fixed to the theoretical prediction of -1 .
- ² Fit includes the contribution from $D_1^*(2430)^0$.
- ³ Calculated using the mass difference $m(D_1^0) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of 0.9 MeV.
- ⁴ No systematic error given.



VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
410.6\pm0.5 OUR FIT	Error includes scale factor of 1.3.			
411.5\pm0.8 OUR AVERAGE				
410.2 \pm 2.1 \pm 0.9	3110 \pm 340	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+}\pi^- X$
411.7 \pm 0.7 \pm 0.4	7.5k	ABULENCIA 06A	CDF	$1900 p\bar{p} \rightarrow D^{*+}\pi^- X$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7\pm 2.5 OUR AVERAGE	Error includes scale factor of 3.5. See the ideogram below.			
35.2 \pm 0.4 \pm 0.9	210k	AAIJ	13CC LHCb	$p\bar{p} \rightarrow D^{*+}\pi^- X$
38.8 \pm 5.0 \pm 1.9	2.7k	¹ ABRAMOWICZ 13	ZEUS	$e^\pm p \rightarrow D^{(*)+}\pi^- X$
31.4 \pm 0.5 \pm 1.3	103k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+}\pi^- X$
20.0 \pm 1.7 \pm 1.3	7.5k	ABULENCIA 06A	CDF	$1900 p\bar{p} \rightarrow D^0\pi^+\pi^-\pi^-$
24 \pm 7 \pm 8	151	ABE 05A	BELL	$B^- \rightarrow D^0\pi^+\pi^-\pi^-$
23.7 \pm 2.7 \pm 4.0		² ABE 04D	BELL	$B^- \rightarrow D^{*+}\pi^-\pi^-$
20 \pm 6 \pm 3	286	AVERY 94C	CLE2	$e^+ e^- \rightarrow D^{*+}\pi^- X$
15 \pm 8 \pm 4	51	FRABETTI 94B	E687	$\gamma Be \rightarrow D^{*+}\pi^- X$
23 \pm 8 \pm 10	279	AVERY 90	CLEO	$e^+ e^- \rightarrow D^{*+}\pi^- X$
13 \pm 6 \pm 10	171	ALBRECHT 89H	ARG	$e^+ e^- \rightarrow D^{*+}\pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
53.2 \pm 7.2 \pm 3.3	3110 \pm 340	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+}\pi^- X$
58 \pm 14 \pm 10	171	ANJOS 89C	TPS	$\gamma N \rightarrow D^{*+}\pi^- X$

NODE=M097M;LINKAGE=AR

NODE=M097M;LINKAGE=AB

NODE=M097M;LINKAGE=CH

NODE=M097M;LINKAGE=K

NODE=M097DM

NODE=M097DM

NODE=M097DM

NODE=M097W


NODE=M097W

¹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions, and A_{D_2} fixed to the theoretical prediction of -1 .

² Fit includes the contribution from $D_1^*(2430)^0$.

NODE=M097W;LINKAGE=AR

NODE=M097W;LINKAGE=AB



$D_1(2420)^0$ DECAY MODES

$\overline{D}_1(2420)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^*(2010)^+ \pi^-$	seen
$\Gamma_2 D^0 \pi^+ \pi^-$	seen
$\Gamma_3 D^0 \rho^0$	
$\Gamma_4 D^0 f_0(500)$	
$\Gamma_5 D_0^*(2300)^+ \pi^-$	
$\Gamma_6 D^+ \pi^-$	not seen
$\Gamma_7 D^{*0} \pi^+ \pi^-$	not seen

$D_1(2420)^0$ BRANCHING RATIOS

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen	ACKERSTAFF 97W OPAL $e^+ e^- \rightarrow D^{*+} \pi^- X$
seen	AVERY 90 CLEO $e^+ e^- \rightarrow D^{*+} \pi^- X$
seen	ALBRECHT 89H ARG $e^+ e^- \rightarrow D^* \pi^- X$
seen	ANJOS 89C TPS $\gamma N \rightarrow D^{*+} \pi^- X$

$\Gamma(D^+ \pi^-)/\Gamma(D^*(2010)^+ \pi^-)$	Γ_6/Γ_1
VALUE CL%	DOCUMENT ID TECN COMMENT
<0.24	90 AVERY 90 CLEO $e^+ e^- \rightarrow D^+ \pi^- X$

$D_1(2420)^0$ POLARIZATION AMPLITUDE A_{D_1}

A polarization amplitude A_{D_1} is a parameter that depends on the initial polarization of the D_1 and is sensitive to a possible S -wave contribution to its decay. For D_1 decays the helicity angle, θ_h , distribution varies like $1 + A_{D_1} \cos^2 \theta_h$, where θ_h is the angle in the D^* rest frame between the two pions emitted by the $D_1 \rightarrow D^* \pi$ and the $D^* \rightarrow D \pi$.

Unpolarized D_1 decaying purely via D -wave is predicted to give $A_{D_1} = 3$.

NODE=M097215;NODE=M097

NODE=M097

DESIG=1

DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow

DESIG=4

DESIG=5

DESIG=6

DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow

DESIG=7;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M097220

NODE=M097R1

NODE=M097R1

NODE=M097R2

NODE=M097R2

NODE=M097PAH

NODE=M097PAH

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
5.73±0.25 OUR AVERAGE				
7.8 ± 6.7 ± 4.6	2.7k	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)} + \pi^- X$
-2.7 ± 1.8				
5.72±0.25	103k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$
5.9 ± 3.0 ± 2.4		CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
-1.7 ± 1.0				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.30±0.48	210k	² AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$
3.8 ± 0.6 ± 0.8		³ AUBERT	09Y BABR	$B^+ \rightarrow D_1^0 \ell^+ \nu_\ell$
2.74 ± 1.40		⁴ AVERY	94C CLE2	$e^+ e^- \rightarrow D^{*+} \pi^- X$
-0.93				
¹ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1. A pure D -wave not excluded although some S -wave mixing possible. ² Systematic uncertainty not estimated. Resonance parameters fixed. ³ Assuming $\Gamma(\gamma(4S) \rightarrow B^+ B^-) / \Gamma(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths and helicity angle distributions for charged and neutral D_1 mesons. ⁴ Systematic uncertainties not estimated.				

NODE=M097PAH

$D_1(2420)^0$ REFERENCES

AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABRAMOWICZ	13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
DEL-AMO-SA..	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE	05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)

NODE=M097PAH;LINKAGE=AR

NODE=M097PAH;LINKAGE=A

NODE=M097PAH;LINKAGE=AU

NODE=M097PAH;LINKAGE=AV

NODE=M097

REFID=55581
 REFID=54743
 REFID=53534
 REFID=52929
 REFID=52733
 REFID=51054
 REFID=50755
 REFID=50011
 REFID=46315
 REFID=45788
 REFID=44096
 REFID=43687
 REFID=41013
 REFID=41001
 REFID=40737

NODE=M120

 $D_1(2420)^\pm$

$$I(J^P) = \frac{1}{2}(??)$$

I needs confirmation.

OMMITTED FROM SUMMARY TABLE
 Seen in $D^*(2007)^0 \pi^+$. $J^P = 0^+$ ruled out.

NODE=M120

NODE=M120M


NODE=M120M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2423.2±2.4 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
2421.9 ± 4.7 ± 3.4	759	¹ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)} 0 \pi^+ X$
-1.2				
2421 ± 2 ± 1	124	ABE	05A BELL	$\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
2425 ± 2 ± 2	146	BERGFELD	94B CLE2	$e^+ e^- \rightarrow D^{*0} \pi^+ X$
2443 ± 7 ± 5	190	ANJOS	89C TPS	$\gamma N \rightarrow D^0 \pi^+ X^0$

NODE=M120M;LINKAGE=AB

¹ From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1, respectively.

NODE=M120

 **$D_1(2420)^{\pm}$ MASS (MeV)**

$$m_{D_1^*(2420)^{\pm}} - m_{D_1^*(2420)^0}$$

NODE=M120DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4⁺²₋₃^{±3}	BERGFELD	94B	CLE2 $e^+ e^- \rightarrow$ hadrons

NODE=M120DM

 $D_1(2420)^{\pm}$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
25± 6 OUR AVERAGE				
21± 5±8	124	ABE	05A	$\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
26 ⁺⁸ ₋₇ ^{±4}	146	BERGFELD	94B	$e^+ e^- \rightarrow D^{*0} \pi^+ X$
41±19±8	190	ANJOS	89C	$\gamma N \rightarrow D^0 \pi^+ X^0$

NODE=M120W

NODE=M120W

 $D_1(2420)^{\pm}$ DECAY MODES

NODE=M120215; NODE=M120

NODE=M120

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^*(2007)^0 \pi^+$	seen
$\Gamma_2 D^+ \pi^+ \pi^-$	seen
$\Gamma_3 D^+ \rho^0$	
$\Gamma_4 D^+ f_0(500)$	
$\Gamma_5 D_0^*(2300)^0 \pi^+$	
$\Gamma_6 D^0 \pi^+$	not seen
$\Gamma_7 D^{*+} \pi^+ \pi^-$	not seen

DESIG=1

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

DESIG=5

DESIG=6

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

 $D_1(2420)^{\pm}$ BRANCHING RATIOS

NODE=M120220

$\Gamma(D^*(2007)^0 \pi^+)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	Γ_1/Γ
seen	ANJOS	89C	TPS $\gamma N \rightarrow D^0 \pi^+ X^0$

NODE=M120R1

NODE=M120R1

$\Gamma(D^0\pi^+)/\Gamma(D^*(2007)^0\pi^+)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.18	90	BERGFELD 94B	CLE2	$e^+e^- \rightarrow \text{hadrons}$	

 $D_1(2420)^\pm$ POLARIZATION AMPLITUDE A_{D_1}

A polarization amplitude A_{D_1} is a parameter that depends on the initial polarization of the D_1 and is sensitive to a possible S -wave contribution to its decay. For D_1 decays the helicity angle, θ_h , distribution varies like $1 + A_{D_1} \cos^2 \theta_h$, where θ_h is the angle in the D^* rest frame between the two pions emitted by the $D_1 \rightarrow D^* \pi$ and the $D^* \rightarrow D \pi$.

Unpolarized D_1 decaying purely via D -wave is predicted to give $A_{D_1} = 3$.

VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3.8 \pm 0.6 \pm 0.8$	² AUBERT	09Y BABR	$B^0 \rightarrow D_1^- \ell^+ \nu_\ell$	
² Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+ B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths and helicity angle distributions for charged and neutral D_1 mesons.				

 $D_1(2420)^\pm$ REFERENCES

ABRAMOWICZ 13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
AUBERT 09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE 05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)
BERGFELD 94B	PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
ANJOS 89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)

 $D_1(2430)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

OMITTED FROM SUMMARY TABLE
 $J = 1^+$ assignment favored (ABE 04D).

 $D_1(2430)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
2427 $\pm 26 \pm 25$	ABE	04D BELL	$B^- \rightarrow D^{*+} \pi^- \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2477 ± 28	¹ AUBERT	06L BABR	$\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$	

¹ Systematic errors not estimated.

 $D_1(2430)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
384 $\pm 107 \pm 74$	ABE	04D BELL	$B^- \rightarrow D^{*+} \pi^- \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
266 ± 97	² AUBERT	06L BABR	$\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$	

² Systematic errors not estimated.

 $D_1(2430)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 D^*(2010)^+ \pi^-$	seen	

 $D_1(2430)^0$ REFERENCES

AUBERT 06L PR D74 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE 04D PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M120R2
NODE=M120R2

NODE=M120PAH

NODE=M120PAH

NODE=M120PAH

NODE=M120PAH;LINKAGE=AU

NODE=M120

REFID=54743
REFID=52929
REFID=50755
REFID=44099
REFID=40737

NODE=M180

NODE=M180

NODE=M180M

NODE=M180M

NODE=M180M;LINKAGE=AU

NODE=M180W

NODE=M180W

NODE=M180W;LINKAGE=AU

NODE=M180215;NODE=M180

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M180

REFID=51140
REFID=50011

NODE=M119

 $D_2^*(2460)^0$ $I(J^P) = \frac{1}{2}(2^+)$

$J^P = 2^+$ assignment strongly favored (ALBRECHT 89B, ALBRECHT 89H), natural parity confirmed by the helicity analysis (DEL-AMO-SANCHEZ 10P). AAIJ 13CC confirms $J^P = 2^+$ and natural parity.

NODE=M119

 $D_2^*(2460)^0$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2460.7 ±0.4 OUR FIT Error includes scale factor of 3.1.

2460.56±0.35 OUR AVERAGE Error includes scale factor of 2.6. See the ideogram below.

2463.7 ±0.4 ±0.7	28k	¹ AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$	
2460.4 ±0.4 ±1.2	82k	AAIJ	13CC LHCb	$p\bar{p} \rightarrow D^{*+} \pi^- X$	
2460.4 ±0.1 ±0.1	675k	AAIJ	13CC LHCb	$p\bar{p} \rightarrow D^+ \pi^- X$	
2462.5 ±2.4 ±1.3	2.3k	² ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$	
2462.2 ±0.1 ±0.8	243k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$	
2460.4 ±1.2 ±2.2	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$	
2461.6 ±2.1 ±3.3		³ ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$	
2464.5 ±1.1 ±1.9	5.8k	³ LINK	04A FOCS	γA	
2465 ±3 ±3	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$	
2453 ±3 ±2	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$	
2461 ±3 ±1	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
2455 ±3 ±5	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$	
2459 ±3 ±2	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

2469.1 ±3.7 ±1.2	1.5k	⁴ CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$	
2463.3 ±0.6 ±0.8	20k	ABULENCIA	06A CDF	$1900 p\bar{p} \rightarrow D^+ \pi^- X$	
2461 ±6	126	⁵ ABREU	98M DLPH	$e^+ e^-$	
2466 ±7	1	ASRATYAN	95 BEBC	$53,40 \nu(\bar{\nu}) \rightarrow pX, dX$	

¹From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_2^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

²From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

³Fit includes the contribution from $D_2^*(2400)^0$.

⁴Calculated using the mass difference $m(D_2^{*0}) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of $+1.2$ MeV.

⁵No systematic error given.

NODE=M119M

NODE=M119M

NODE=M119M

OCCUR=2

OCCUR=2

NODE=M119M;LINKAGE=B


NODE=M119M;LINKAGE=AR

NODE=M119M;LINKAGE=LI

NODE=M119M;LINKAGE=CH

NODE=M119M;LINKAGE=K

WEIGHTED AVERAGE
2460.56±0.35 (Error scaled by 2.6)



$D_2^*(2460)^0$ mass (MeV)

$m_{D_2^{*0}} - m_{D^+}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
591.0±0.4 OUR FIT	Error includes scale factor of 2.9.			NODE=M119DM
593.9±0.6±0.5	20k	ABULENCIA	06A CDF	$1900 p\bar{p} \rightarrow D^+ \pi^- X$

$m_{D_2^{*0}} - m_{D^{*+}}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
450.4±0.4 OUR FIT	Error includes scale factor of 2.9.			NODE=M119DM2
458.8±3.7^{+1.2}_{-1.3}	1560 ± 230	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)} + \pi^- X$

$D_2^*(2460)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47.5± 1.1 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			NODE=M119W
47.0± 0.8± 1.0	28k	⁶ AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$
43.2± 1.2± 3.0	82k	AAIJ	13CC LHCb	$p\bar{p} \rightarrow D^{*+} \pi^- X$
45.6± 0.4± 1.1	675k	AAIJ	13CC LHCb	$p\bar{p} \rightarrow D^+ \pi^- X$
46.6± 8.1 ^{+5.9} _{-3.8}	2.3k	⁷ ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)} + \pi^- X$
50.5± 0.6± 0.7	243k	DEL-AMO-SA... 10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$
41.8± 2.5± 2.9	3.4k	AUBERT	09AB BABR	$B^- \rightarrow D^+ \pi^- \pi^-$
49.2± 2.3± 1.3	20k	ABULENCIA	06A CDF	$1900 p\bar{p} \rightarrow D^+ \pi^- X$
45.6± 4.4± 6.7	⁸ ABE	04D BELL	$B^- \rightarrow D^+ \pi^- \pi^-$	
38.7± 5.3± 2.9	5.8k	⁸ LINK	04A FOCS	γA
28 ⁺⁸ ₋₇ ± 6	486	AVERY	94C CLE2	$e^+ e^- \rightarrow D^+ \pi^- X$
25 ⁺¹⁰ ₋₁₀ ± 5	128	FRABETTI	94B E687	$\gamma Be \rightarrow D^+ \pi^- X$
20 ⁺⁹ ₋₁₂ ± 9	440	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
15 ⁺¹³ ₋₁₀ ± 5	337	ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- X$
20 ⁺¹⁰ ₋₁₀ ± 5	153	ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- X$

NODE=M119DM2

NODE=M119DM2

NODE=M119DM2

NODE=M119DM2

NODE=M119DM2

NODE=M119W

NODE=M119W

OCCUR=3


OCCUR=2

- 6 From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.
- 7 From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 .
- 8 Fit includes the contribution from $D_0^*(2400)^0$.

NODE=M119W;LINKAGE=D

NODE=M119W;LINKAGE=AR

NODE=M119W;LINKAGE=LI



$D_2^*(2460)^0$ DECAY MODES

$\bar{D}_2^*(2460)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^+ \pi^-$	seen
Γ_2 $D^*(2010)^+ \pi^-$	seen
Γ_3 $D^0 \pi^+ \pi^-$	not seen
Γ_4 $D^{*0} \pi^+ \pi^-$	not seen

NODE=M119215;NODE=M119

NODE=M119

CLUMP=A;DESIG=1

DESIG=2

DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M119220

NODE=M119R1

NODE=M119R1

NODE=M119R2

NODE=M119R2

NODE=M119R3

NODE=M119R3

$\Gamma(D^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	3.4k AUBERT 09AB BABR $B^- \rightarrow D^+ \pi^- \pi^-$
seen	337 ALBRECHT 89B ARG $e^+ e^- \rightarrow D^+ \pi^- X$
seen	ANJOS 89C TPS $\gamma N \rightarrow D^+ \pi^- X$

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	ACKERSTAFF 97W OPAL $e^+ e^- \rightarrow D^{*+} \pi^- X$
seen	AVERY 90 CLEO $e^+ e^- \rightarrow D^{*+} \pi^- X$
seen	ALBRECHT 89H ARG $e^+ e^- \rightarrow D^* \pi^- X$

$\Gamma(D^+ \pi^-)/\Gamma(D^*(2010)^+ \pi^-)$	Γ_1/Γ_2
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.54±0.15 OUR AVERAGE	
$1.4 \pm 0.3 \pm 0.3$	2.3k ABRAMOWICZ13 ZEUS $e^\pm p \rightarrow D^{(*)+} \pi^- X$
$1.47 \pm 0.03 \pm 0.16$	379k DEL-AMO-SA..10P BABR $e^+ e^- \rightarrow D^{(*)+} \pi^- X$
$2.8 \pm 0.8 \pm 0.5$	1560 ± 230 CHEKANOV 09 ZEUS $e^\pm p \rightarrow D^{(*)+} \pi^- X$

$2.2 \pm 0.7 \pm 0.6$	AVERY	94C	CLE2	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2.3 ± 0.8	AVERY	90	CLEO	$e^+ e^-$
$3.0 \pm 1.1 \pm 1.5$	ALBRECHT	89H	ARG	$e^+ e^- \rightarrow D^* \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.9 ± 0.5	ABE	04D	BELL	$B^- \rightarrow D^{(*)+} \pi^- \pi^-$

9 From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

$\Gamma(D^+ \pi^-) / [\Gamma(D^+ \pi^-) + \Gamma(D^*(2010)^+ \pi^-)]$	$\Gamma_1 / (\Gamma_1 + \Gamma_2)$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.62 \pm 0.03 \pm 0.02$	8414	10	AUBERT	$09Y$ BABR $B^+ \rightarrow D_2^{*0} \ell^+ \nu_\ell$
10 Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+ B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral D_2^* mesons.				

$D_2^*(2460)^0$ POLARIZATION AMPLITUDE A_{D_2}

A polarization amplitude A_{D_2} is a parameter that depends on the initial polarization of the D_2 . For D_2 decays the helicity angle, θ_H , distribution varies like $1 + A_{D_2} \cos^2(\theta_H)$, where θ_H is the angle in the D^* rest frame between the two pions emitted by the $D_2 \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-1.16 ± 0.35	2.3k	11	ABRAMOWICZ13	ZEUS $e^\pm p \rightarrow D^{(*)+} \pi^- X$
consistent with -1	243k		DEL-AMO-SA..10P	BABR $e^+ e^- \rightarrow D^+ \pi^- X$
$-0.74^{+0.49}_{-0.38}$		12	AVERY	94C CLE2 $e^+ e^- \rightarrow D^{*+} \pi^- X$

11 From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions.

12 Systematic uncertainties not estimated.

$D_2^*(2460)^0$ REFERENCES

AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABRAMOWICZ	13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
DEL-AMO-SA..	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)
LINK	04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ASRATYAN	95	ZPHY C68 43	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+) (CLEO Collab.)
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(FNAL E687 Collab.)
FRAEBETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(CLEO Collab.)
AVERY	90	PR D41 774	P. Avery, D. Besson	(FNAL E687 Collab.)
ALBRECHT	89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)

NODE=M119R3;LINKAGE=AR

NODE=M119R01
NODE=M119R01

NODE=M119R01;LINKAGE=AU

NODE=M119PAM

NODE=M119PAM

NODE=M119PAM

NODE=M119PAM;LINKAGE=AB
NODE=M119PAM;LINKAGE=AV

NODE=M119

REFID=57518
REFID=55581
REFID=54743
REFID=53534
REFID=52941
REFID=52929
REFID=52733
REFID=51054
REFID=50011
REFID=49775
REFID=46315
REFID=45788
REFID=44439
REFID=44096
REFID=43687
REFID=41013
REFID=40736
REFID=41001
REFID=40737

$D_2^*(2460)^\pm$

$I(J^P) = \frac{1}{2}(2^+)$

$J^P = 2^+$ assignment strongly favored(ALBRECHT 89B).

$D_2^*(2460)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2465.4±1.3 OUR AVERAGE				Error includes scale factor of 3.1. See the ideogram below.

2465.6±1.8±1.3	1	AAIJ	15X LHCb	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2468.6±0.6±0.3	2	AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2463.1±0.2±0.6	342k	AAIJ	13CC LHCb	$p p \rightarrow D^0 \pi^+ X$
2460.6±4.4 ^{+3.6} _{-0.8}	1371	ABRAMOWICZ13	ZEUS	$e^\pm p \rightarrow D^{(*)0} \pi^+ X$
2465.4±0.2±1.1	111k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^0 \pi^+ X$
2465.7±1.8 ^{+1.4} _{-4.8}	2909	KUZMIN	07 BELL	$e^+ e^- \rightarrow \text{hadrons}$
2463 ± 3 ± 3	310	BERGFELD	94B CLE2	$e^+ e^- \rightarrow D^0 \pi^+ X$
2453 ± 3 ± 2	185	FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^0 \pi^+ X$
2469 ± 4 ± 6		ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2468.1±0.6±0.5	5	AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2467.6±1.5±0.8	3.5k	LINK	04A FOCS	γA

¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S-wave and the $\bar{D}\pi$ S- and P-waves.


² Modeling the $\pi^+ \pi^-$ S-wave with the Isobar formalism.

³ From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1, respectively.

⁴ At a fixed width of 50.5 MeV.

⁵ Modeling the $\pi^+ \pi^-$ S-wave with the K-matrix formalism.

⁶ Fit includes the contribution from $D_0^*(2400)^\pm$. Not independent of the corresponding mass difference measurement, $(m_{D_2^*(2460)^\pm}) - (m_{D_2^*(2460)^0})$.



$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2.4±1.7 OUR AVERAGE			

3.1±1.9±0.9	LINK	04A FOCS	γA
- 2 ± 4 ± 4	BERGFELD	94B CLE2	$e^+ e^- \rightarrow \text{hadrons}$
0 ± 4	FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D\pi X$
14 ± 5 ± 8	ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ X$

NODE=M150

NODE=M150

NODE=M150M

NODE=M150M

OCCUR=2

NODE=M150M;LINKAGE=A

NODE=M150M;LINKAGE=B

NODE=M150M;LINKAGE=AB

NODE=M150M;LINKAGE=DE

NODE=M150M;LINKAGE=C

NODE=M150M;LINKAGE=LI

NODE=M150DM

NODE=M150DM

$D_2^*(2460)^{\pm}$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
46.7± 1.2 OUR AVERAGE				
46.0± 3.4±3.2		1 AAIJ	15X LHCb	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
47.3± 1.5±0.7		2 AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
48.6± 1.3±1.9	342k	AAIJ	13CC LHCb	$p p \rightarrow D^0 \pi^+ X$
49.7± 3.8±6.4	2909	KUZMIN	07 BELL	$e^+ e^- \rightarrow \text{hadrons}$
34.1± 6.5±4.2	3.5k	3 LINK	04A FOCS	γA
27 ± 11 ± 5	310	BERGFELD	94B CLE2	$e^+ e^- \rightarrow D^0 \pi^+ X$
23 ± 9 ± 5	185	FRABETTI	94B E687	$\gamma Be \rightarrow D^0 \pi^+ X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
46.0± 1.4±1.8		4 AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$

1 From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S -wave and the $D\pi$ S - and P -waves.

2 Modeling the $\pi^+ \pi^-$ S -wave with the Isobar formalism.

3 Fit includes the contribution from $D_0^*(2400)^\pm$.

4 Modeling the $\pi^+ \pi^-$ S -wave with the K-matrix formalism.

NODE=M150W

NODE=M150W

OCCUR=2

NODE=M150W;LINKAGE=A

NODE=M150W;LINKAGE=B

NODE=M150W;LINKAGE=LI

NODE=M150W;LINKAGE=C

 $D_2^*(2460)^{\pm}$ DECAY MODES

$D_2^*(2460)^-$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0 \pi^+$	seen
$\Gamma_2 D^{*0} \pi^+$	seen
$\Gamma_3 D^+ \pi^+ \pi^-$	not seen
$\Gamma_4 D^{*+} \pi^+ \pi^-$	not seen

 $D_2^*(2460)^{\pm}$ BRANCHING RATIOS

$\Gamma(D^0 \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	ALBRECHT 89F ARG $e^+ e^- \rightarrow D^0 \pi^+ X$

$\Gamma(D^0 \pi^+)/\Gamma(D^{*0} \pi^+)$	Γ_1/Γ_2
1.2±0.4 OUR AVERAGE	ALBRECHT 89F ARG $e^+ e^- \rightarrow D^0 \pi^+ X$

1.1±0.4±0.3	1371	1 ABRAMOWICZ13 ZEUS $e^\pm p \rightarrow D^{(*)0} \pi^+ X$
1.9±1.1±0.3		BERGFELD 94B CLE2 $e^+ e^- \rightarrow \text{hadrons}$

1 From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1, respectively.

$\Gamma(D^0 \pi^+)/[\Gamma(D^0 \pi^+) + \Gamma(D^{*0} \pi^+)]$	$\Gamma_1/(\Gamma_1+\Gamma_2)$
seen	ALBRECHT 89F ARG $e^+ e^- \rightarrow D^0 \pi^+ X$

• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.62±0.03±0.02	3361 1 AUBERT 09Y BABR $\bar{B}^0 \rightarrow D_2^{*+} \ell^- \nu_\ell$

1 Assuming $\Gamma(\gamma(4S) \rightarrow B^+ B^-) / \Gamma(\gamma(4S) \rightarrow B^0 \bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral D_2^* mesons.

DESIG=1

DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow

NODE=M150220

NODE=M150R1

NODE=M150R1

NODE=M150R2

NODE=M150R2

NODE=M150R2;LINKAGE=AB

NODE=M150R01

NODE=M150R01

NODE=M150R01;LINKAGE=AU

NODE=M150

REFID=56588

REFID=56609

REFID=55581

REFID=54743

REFID=53534

REFID=52929

REFID=51854

REFID=49775

REFID=44099

REFID=43687

REFID=40736

REFID=40931

 $D_2^*(2460)^{\pm}$ REFERENCES

AAIJ 15X PR D92 012012	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 15Y PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABRAMOWICZ 13 NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)
DEL-AMO-SA... 10P PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT 09Y PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)
KUZMIN 07 PR D76 012006	A. Kuzmin <i>et al.</i>	(BELLE Collab.)
LINK 04A PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)
BERGFELD 94B PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)
FRABETTI 94B PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ALBRECHT 89B PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ALBRECHT 89F PL B231 208	H. Albrecht <i>et al.</i>	(ARGUS Collab.)

NODE=M198

D(2550)⁰

$$I(J^P) = \frac{1}{2}(?)$$

OMITTED FROM SUMMARY TABLE

Unnatural parity according to the helicity analysis of DEL-AMO-SANCHEZ 10P and AAIJ 13CC. DEL-AMO-SANCHEZ 10P suggests $J^P = 0^-$.

NODE=M198

D(2550)⁰ MASS

NODE=M198M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2564 ± 20 OUR AVERAGE				Error includes scale factor of 3.9.
2579.5 ± 3.4 ± 5.5	60k	AAIJ 13CC LHCb	$p p \rightarrow D^* + \pi^- X$	
2539.4 ± 4.5 ± 6.8	34k	DEL-AMO-SA..10P BABR	$e^+ e^- \rightarrow D^* + \pi^- X$	

NODE=M198M

D(2550)⁰ WIDTH

NODE=M198W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
135 ± 17 OUR AVERAGE				
177.5 ± 17.8 ± 46.0	60k	AAIJ 13CC LHCb	$p p \rightarrow D^* + \pi^- X$	
130 ± 12 ± 13	34k	DEL-AMO-SA..10P BABR	$e^+ e^- \rightarrow D^* + \pi^- X$	

NODE=M198W

D(2550)⁰ DECAY MODES

NODE=M198215;NODE=M198

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^* + \pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

D(2550)⁰ POLARIZATION AMPLITUDE A_{D_J}

NODE=M198PAM

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$ decays.

NODE=M198PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.2 ± 1.3	60k	¹ AAIJ 13CC LHCb	$p p \rightarrow D^* + \pi^- X$	

NODE=M198PAM

¹ Systematic uncertainty not estimated.

NODE=M198PAM;LINKAGE=A

D(2550)⁰ REFERENCES

NODE=M198

AAIJ 13CC JHEP 1309 145 DEL-AMO-SA... 10P PR D82 111101	R. Aaij <i>et al.</i> P. del Amo Sanchez <i>et al.</i>	(LHCb Collab.) (BABAR Collab.)
--	---	-----------------------------------

REFID=55581

REFID=53534

NODE=M199

 $D_J^*(2600)$ $I(J^P) = \frac{1}{2}(?)$

OMITTED FROM SUMMARY TABLE

was $D(2600)$ J^P consistent with natural parity (DEL-AMO-SANCHEZ 10P,
AAIJ 13CC).

NODE=M199


 $D_J^*(2600)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2623 ±12 OUR AVERAGE					Error includes scale factor of 4.8. See the ideogram below.
2681.1 ± 5.6 ± 14.0	28k	1 AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$	
2649.2 ± 3.5 ± 3.5	51k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$	
2608.7 ± 2.4 ± 2.5	26k	DEL-AMO-SA..10P	BABR 0	$e^+ e^- \rightarrow D^+ \pi^- X$	
2621.3 ± 3.7 ± 4.2	13k	2 DEL-AMO-SA..10P	BABR +	$e^+ e^- \rightarrow D^0 \pi^+ X$	

¹ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

² At a fixed width of 93 MeV.

WEIGHTED AVERAGE
2623±12 (Error scaled by 4.8)

 $D_J^*(2600)$ MASS (MeV)

NODE=M199M

NODE=M199M

OCCUR=2

NODE=M199M;LINKAGE=A

NODE=M199M;LINKAGE=DE

 $D_J^*(2600)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
139 ±31 OUR AVERAGE				Error includes scale factor of 3.2. See the ideogram below.
186.7 ± 8.5 ± 11.9	28k	3 AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$
140.2 ± 17.1 ± 18.6	51k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$


93 ± 6 ± 13 26k DEL-AMO-SA..10P BABR $e^+ e^- \rightarrow D^+ \pi^- X$

³ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

NODE=M199W

NODE=M199W

NODE=M199W;LINKAGE=A



$D_s^*(2600)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D\pi$	seen
$\Gamma_2 D^+\pi^-$	seen
$\Gamma_3 D^0\pi^\pm$	seen
$\Gamma_4 D^*\pi$	seen
$\Gamma_5 D^{*+}\pi^-$	seen

$D_s^*(2600)$ BRANCHING RATIOS

$\Gamma(D^+\pi^-)/\Gamma(D^{*+}\pi^-)$	Γ_2/Γ_5
VALUE $0.32 \pm 0.02 \pm 0.09$	EVTS 76k DOCUMENT ID DEL-AMO-SA...10P TECN BABR COMMENT $e^+e^- \rightarrow D^{(*)+}\pi^-X$

$D_s^*(2600)$ REFERENCES

AAIJ 16AH PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA...10P PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

NODE=M199215;NODE=M199

DESIG=1;OUR EVAL; \rightarrow UNCHECKED
 DESIG=2;OUR EVAL; \rightarrow UNCHECKED
 DESIG=3;OUR EVAL; \rightarrow UNCHECKED
 DESIG=4;OUR EVAL; \rightarrow UNCHECKED
 DESIG=5;OUR EVAL; \rightarrow UNCHECKED

NODE=M199220

NODE=M199R01
 NODE=M199R01

NODE=M199

REFID=57518
 REFID=55581
 REFID=53534

$D^*(2640)^\pm$

$$I(J^P) = \frac{1}{2}(??)$$

OMITTED FROM SUMMARY TABLE

Seen in Z decays by ABREU 98M. Not seen by ABBIENDI 01N and CHEKANOV 09. Needs confirmation.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2637±2±6	66 ± 14	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

 $D^*(2640)^\pm$ MASS

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<15	95	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

 $D^*(2640)^\pm$ WIDTH

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^*(2010)^+ \pi^+ \pi^-$	seen

 $D^*(2640)^\pm$ REFERENCES

CHEKANOV 09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABBIENDI 01N	EPJ C20 445	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU 98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)

 $D(2740)^0$

$$I(J^P) = \frac{1}{2}(??)$$

OMITTED FROM SUMMARY TABLE
 J^P consistent with unnatural parity (AAIJ 13CC).

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2737.0±3.5±11.2	7.7k	AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$

 $D(2740)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
73.2±13.4±25.0	7.7k	AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$

 $D(2740)^0$ WIDTH

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^{*+} \pi^-$	seen

 $D(2740)^0$ DECAY MODES

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

3.1±2.2 7.7k ¹ AAIJ 13CC LHCb $p p \rightarrow D^{*+} \pi^- X$

1 Systematic uncertainty not estimated.

NODE=M158

NODE=M158

NODE=M158M

NODE=M158M

NODE=M158W

NODE=M158W

NODE=M158215;NODE=M158

NODE=M158

DESIG=1;OUR EST;→ UNCHECKED ←

NODE=M158

REFID=52733

REFID=48296

REFID=46315

NODE=M228

NODE=M228

NODE=M228M

NODE=M228M

NODE=M228W

NODE=M228W

NODE=M228215;NODE=M228

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M228PAM

NODE=M228PAM

NODE=M228PAM

NODE=M228PAM;LINKAGE=A

D(2740)⁰ REFERENCES

AAIJ

13CC JHEP 1309 145

R. Aaij *et al.*

(LHCb Collab.)

 $D_3^*(2750)$ $I(J^P) = \frac{1}{2}(3^-)$

OMITTED FROM SUMMARY TABLE

J^P determined by AAIJ 15Y from the Dalitz plot analysis of $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$ decays. J^P consistent with natural parity (AAIJ 13CC).

NODE=M228

REFID=55581

NODE=M203

NODE=M203


NODE=M203M

NODE=M203M

OCCUR=2
OCCUR=3OCCUR=2
OCCUR=3

OCCUR=2

NODE=M203M;LINKAGE=C

NODE=M203M;LINKAGE=A
NODE=M203M;LINKAGE=DE
NODE=M203M;LINKAGE=DA
NODE=M203M;LINKAGE=BWEIGHTED AVERAGE
 2763.5 ± 3.4 (Error scaled by 2.2) **$D_3^*(2750)$ WIDTH**

NODE=M203W

NODE=M203W

OCCUR=2

 66 ± 5 OUR AVERAGE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
66 ± 5 OUR AVERAGE					
95.3 ± 9.6 ± 34.0	28k	6 AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$	
105 ± 18 ± 24		7 AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$	
74.4 ± 3.4 ± 37.0	14k	AAIJ	13CC LHCb 0	$p p \rightarrow D^{*+} \pi^- X$	
74.4 ± 3.4 ± 19.1	56k	AAIJ	13CC LHCb 0	$p p \rightarrow D^+ \pi^- X$	

$66.7 \pm 6.6 \pm 10.5$	20k	AAIJ	13CC LHCb + 8 DEL-AMO-SA..10P BABR	$p p \rightarrow D^0 \pi^+ X$ $e^+ e^- \rightarrow D^{*+} \pi^- X$ $e^+ e^- \rightarrow D^+ \pi^- X$
$71 \pm 6 \pm 11$	23.5k			

• • • We do not use the following data for averages, fits, limits, etc. • • •

154 $\pm 27 \pm 16$	9 AAIJ	15Y LHCb	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
---------------------	--------	----------	---

⁶ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

⁷ Modeling the $\pi^+ \pi^-$ S-wave with the Isobar formalism.

⁸ The states observed in the $D^* \pi$ and $D \pi$ final states are not necessarily the same.

⁹ Modeling the $\pi^+ \pi^-$ S-wave with the K-matrix formalism.

OCCUR=4

OCCUR=2

OCCUR=2

NODE=M203W;LINKAGE=C

NODE=M203W;LINKAGE=A

NODE=M203W;LINKAGE=DE

NODE=M203W;LINKAGE=B

NODE=M203215;NODE=M203

$D_3^*(2750)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D \pi$	seen
$\Gamma_2 D^+ \pi^-$	seen
$\Gamma_3 D^0 \pi^\pm$	seen
$\Gamma_4 D^* \pi$	seen
$\Gamma_5 D^{*+} \pi^-$	seen

$D_3^*(2750)$ BRANCHING RATIOS

$\Gamma(D^+ \pi^-)/\Gamma(D^{*+} \pi^-)$	Γ_2/Γ_5			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.42 $\pm 0.05 \pm 0.11$	34.8k	10 DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{(*)+} \pi^- X$

¹⁰ The states observed in the $D^* \pi$ and $D \pi$ final states are not necessarily the same.

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=4;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=5;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M203220

NODE=M203R01
NODE=M203R01

NODE=M203R01;LINKAGE=DE

NODE=M203PAM

NODE=M203PAM

NODE=M203PAM

NODE=M203PAM;LINKAGE=DE

NODE=M203

REFID=57518
REFID=56609
REFID=55581
REFID=53534

$D_3^*(2750)$ REFERENCES

AAIJ	16AH PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15Y PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA... 10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

NODE=M229

D(3000)⁰

$I(J^P) = \frac{1}{2}(?)$

OMITTED FROM SUMMARY TABLE

Both natural- and unnatural-parity components observed depending on the decay mode (AAIJ 13CC).

NODE=M229

D(3000)⁰ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3214 ±29 ±49	28k	1 AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2971.8 ± 8.7	9.5k	2,3 AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$
3008.1 ± 4.0	17.6k	2,4 AAIJ	13CC LHCb	$p p \rightarrow D^+ \pi^- X$
1 From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.				
2 Systematic uncertainty not estimated.				
3 Unnatural parity preferred.				
4 Natural parity state. A state $D(3000)^+$ is possibly seen in $D^0 \pi^+$ final state.				

NODE=M229M

NODE=M229M

OCCUR=2

NODE=M229M;LINKAGE=D

NODE=M229M;LINKAGE=A
NODE=M229M;LINKAGE=B
NODE=M229M;LINKAGE=C

D(3000)⁰ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
186 ±38 ±72	28k	5 AAIJ	16AH LHCb	$B^- \rightarrow D^+ \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
188.1 ± 44.8	9.5k	6,7 AAIJ	13CC LHCb	$p p \rightarrow D^{*+} \pi^- X$
110.5 ± 11.5	17.6k	6,8 AAIJ	13CC LHCb	$p p \rightarrow D^+ \pi^- X$
5 From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.				
6 Systematic uncertainty not estimated.				
7 Unnatural parity preferred.				
8 Natural parity state. A state $D(3000)^+$ is possibly seen in $D^0 \pi^+$ final state.				

NODE=M229W

NODE=M229W

OCCUR=2

NODE=M229W;LINKAGE=D

NODE=M229W;LINKAGE=A
NODE=M229W;LINKAGE=B
NODE=M229W;LINKAGE=C

NODE=M229215;NODE=M229

D(3000)⁰ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^{*+} \pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

NODE=M229PAM

NODE=M229PAM

NODE=M229PAM

NODE=M229PAM;LINKAGE=A

NODE=M229

REFID=57518

REFID=55581

D(3000)⁰ REFERENCES

AAIJ	16AH PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)

CHARMED, STRANGE MESONS ($C = S = \pm 1$)

$D_s^+ = c\bar{s}$, $D_s^- = \bar{c}s$, similarly for $D_s^{*\pm}$

$D_s^{*\pm}$

$I(J^P) = 0(?)$

J^P is natural, width and decay modes consistent with 1^- .

$D_s^{*\pm}$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^{*}(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2112.2±0.4 OUR FIT			
2106.6±2.1±2.7	¹ BLAYLOCK 87 MRK3 $e^+ e^- \rightarrow D_s^\pm \gamma X$		

¹ Assuming D_s^\pm mass = 1968.7 ± 0.9 MeV.

$m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^{*}(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
143.8 ± 0.4 OUR FIT				
143.9 ± 0.4 OUR AVERAGE				
143.76 ± 0.39 ± 0.40		GRONBERG 95 CLE2 $e^+ e^-$		
144.22 ± 0.47 ± 0.37		BROWN 94 CLE2 $e^+ e^-$		
142.5 ± 0.8 ± 1.5		² ALBRECHT 88 ARG $e^+ e^- \rightarrow D_s^\pm \gamma X$		
139.5 ± 8.3 ± 9.7	60	AIHARA 84D TPC $e^+ e^- \rightarrow$ hadrons		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
143.0 ± 18.0	8	ASRATYAN 85 HLBC FNAL 15-ft, $\nu^2 H$		
110 ± 46		BRANDELIK 79 DASP $e^+ e^- \rightarrow D_s^\pm \gamma X$		

² Result includes data of ALBRECHT 84B.

$D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.9	90	GRONBERG 95 CLE2 $e^+ e^-$		
< 4.5	90	ALBRECHT 88 ARG $E_{cm}^{ee} = 10.2$ GeV		

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.9	90	BROWN 94 CLE2 $e^+ e^-$
<22	90	BLAYLOCK 87 MRK3 $e^+ e^- \rightarrow D_s^\pm \gamma X$

D_s^{*+} DECAY MODES

D_s^{*-} modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D_s^+ \gamma$	(93.5 ± 0.7) %
$\Gamma_2 D_s^+ \pi^0$	(5.8 ± 0.7) %
$\Gamma_3 D_s^+ e^+ e^-$	(6.7 ± 1.6) × 10 ⁻³

NODE=MXXX040

NODE=MXXX040

NODE=S074

NODE=S074

NODE=S074M

NODE=S074M

NODE=S074M

NODE=S074M;LINKAGE=E

NODE=S074DM

NODE=S074DM

NODE=S074DM

NODE=S074DM;LINKAGE=A

NODE=S074W

NODE=S074W

NODE=S074215;NODE=S074

NODE=S074

DESIG=1

DESIG=2

DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 3 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-97	
x_3	-19	-4
	x_1	x_2

D_s^{*+} BRANCHING RATIOS

$\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.935 ± 0.007 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ASRATYAN	91	HLBC	$\bar{\nu}_\mu$ Ne
seen	ALBRECHT	88	ARG	$e^+ e^- \rightarrow D_s^+ \gamma X$
seen	AIHARA	84D		
seen	ALBRECHT	84B		
seen	BRANDELIK	79		

$\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.062 ± 0.008 OUR FIT			
0.062 ± 0.008 OUR AVERAGE			

$0.062 \pm 0.005 \pm 0.006$	AUBERT,BE	05G	BABR	$10.6 e^+ e^- \rightarrow \text{hadrons}$
$0.062^{+0.020}_{-0.018} \pm 0.022$	GRONBERG	95	CLE2	$e^+ e^-$

$\Gamma(D_s^+ e^+ e^-) / \Gamma(D_s^+ \gamma)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 1.7 OUR FIT				

$7.2^{+1.5}_{-1.3} \pm 1.0$	38	CRONIN-HEN..12	CLEO	$4.17 e^+ e^- \rightarrow \text{hadrons}$
---	----	----------------	------	---

D_s^{*+} REFERENCES

CRONIN-HEN... 12	PR D86 072005	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=54627
AUBERT,BE 05G	PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50942
GRONBERG 95	PRL 75 3232	J. Gronberg <i>et al.</i>	(CLEO Collab.)	REFID=44568
BROWN 94	PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)	REFID=43868
ASRATYAN 91	PL B257 525	A.E. Asratyan <i>et al.</i>	(ITEP, BELG, SACL+)	REFID=41582
ALBRECHT 88	PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40269
BLAYLOCK 87	PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)	REFID=40005
ASRATYAN 85	PL 156B 441	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=22887
AIHARA 84D	PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=11561
ALBRECHT 84B	PL 146B 111	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22886
BRANDELIK 79	PL 80B 412	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=11442

NODE=S074220

NODE=S074R1
NODE=S074R1NODE=S074R2
NODE=S074R2NODE=S074R01
NODE=S074R01

NODE=S074

NODE=M172

 $D_{s0}^*(2317)^\pm$
 $I(J^P) = 0(0^+)$
 J, P need confirmation.

AUBERT 06P and CHOI 15A do not observe neutral and doubly charged partners of the $D_{s0}^*(2317)^\pm$.

NODE=M172

 $D_{s0}^*(2317)^\pm$ MASS
The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^{*}(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2317.8±0.5 OUR FIT				
2318.0±0.7 OUR AVERAGE				
2318.3±1.2±1.2	115	1 ABLIKIM	18J BES3	4.6 $e^+ e^- \rightarrow D_s^{*\pm} D_{s0}^*(2317)^\mp$
2319.6±0.2±1.4	3.1k	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^0 X$
2317.3±0.4±0.8	1.0k	2 AUBERT	04E BABR	10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2317.2±1.3	88	3 AUBERT,B	04S BABR	$B \rightarrow D_{s0}^{(*)}(2317)^+ \bar{D}^{(*)}$
2317.2±0.5±0.9	761	4 MIKAMI	04 BELL	10.6 $e^+ e^-$
2316.8±0.4±3.0	1.2k	4.5 AUBERT	03G BABR	10.6 $e^+ e^-$
2317.6±1.3	273	4.6 AUBERT	03G BABR	10.6 $e^+ e^-$
2319.8±2.1±2.0	24	4 KROKOVNY	03B BELL	10.6 $e^+ e^-$

¹ From a fit of the D_s^* recoil mass where the $D_{s0}^*(2317)$ signal is described with a Crystal Ball function convolved with a Gaussian function.

² Supersedes AUBERT 03G.

³ Systematic errors not evaluated.

⁴ Not independent of the corresponding $m_{D_{s0}^*(2317)} - m_{D_s}$.

⁵ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.

⁶ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.

NODE=M172M

NODE=M172M

NODE=M172M

OCCUR=2

NODE=M172M;LINKAGE=A

NODE=M172M;LINKAGE=AU

NODE=M172M;LINKAGE=AB

NODE=M172M;LINKAGE=B1

NODE=M172M;LINKAGE=A1

NODE=M172M;LINKAGE=A2

NODE=M172DM

NODE=M172DM

NODE=M172DM

OCCUR=2

NODE=M172DM;LINKAGE=K3

NODE=M172DM;LINKAGE=A1

NODE=M172DM;LINKAGE=C1

NODE=M172DM;LINKAGE=A2

NODE=M172DM;LINKAGE=C2

NODE=M172W

NODE=M172W

 $D_{s0}^*(2317)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.8	95	3180	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^0 X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 4.6	90	761	MIKAMI	04 BELL	10.6 $e^+ e^-$
< 10			AUBERT	03G BABR	10.6 $e^+ e^-$
< 7	90	135	BESSION	03 CLE2	10.6 $e^+ e^-$

$D_{s0}^*(2317)^\pm$ DECAY MODES $D_{s0}^*(2317)^-$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 D_s^+ \pi^0$	$(100^{+0}_{-20})\%$	DESIG=1
$\Gamma_2 D_s^+ \gamma$	< 5 %	90%
$\Gamma_3 D_s^*(2112)^+ \gamma$	< 6 %	90%
$\Gamma_4 D_s^+ \gamma\gamma$	< 18 %	95%
$\Gamma_5 D_s^*(2112)^+ \pi^0$	< 11 %	90%
$\Gamma_6 D_s^+ \pi^+ \pi^-$	< 4 $\times 10^{-3}$	90%
$\Gamma_7 D_s^+ \pi^0 \pi^0$	not seen	DESIG=7;OUR EVAL; \rightarrow UNCHECKED \leftarrow

 $D_{s0}^*(2317)^\pm$ BRANCHING RATIOS

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$	Γ_1/Γ
1.00$^{+0.00}_{-0.14}$$^{+0.00}_{-0.14}$	47 ABLIKIM 18J BES3 4.6 $e^+ e^- \rightarrow D_s^* D_{s0}^*(2317)^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 1.5k AUBERT 03G BABR 10.6 $e^+ e^-$

$\Gamma(D_s^+ \gamma)/\Gamma(D_s^+ \pi^0)$	Γ_2/Γ_1
<0.05 90 MIKAMI 04 BELL 10.6 $e^+ e^-$	NODE=M172R5 NODE=M172R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.14 95 AUBERT 06P BABR 10.6 $e^+ e^-$
<0.052 90 BESSON 03 CLE2 10.6 $e^+ e^-$

$\Gamma(D_s^*(2112)^+ \gamma)/\Gamma(D_s^+ \pi^0)$	Γ_3/Γ_1
<0.059 90 BESSON 03 CLE2 10.6 $e^+ e^-$	NODE=M172R6 NODE=M172R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.16 95 AUBERT 06P BABR 10.6 $e^+ e^-$
<0.18 90 MIKAMI 04 BELL 10.6 $e^+ e^-$

$\Gamma(D_s^+ \gamma\gamma)/\Gamma(D_s^+ \pi^0)$	Γ_4/Γ_1
<0.18 95 AUBERT 06P BABR 10.6 $e^+ e^-$	NODE=M172R7 NODE=M172R7

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen AUBERT 03G BABR 10.6 $e^+ e^-$

$\Gamma(D_s^*(2112)^+ \pi^0)/\Gamma(D_s^+ \pi^0)$	Γ_5/Γ_1
<0.11 90 BESSON 03 CLE2 10.6 $e^+ e^-$	NODE=M172R8 NODE=M172R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma(D_s^+ \pi^0)$	Γ_6/Γ_1
<0.004 90 MIKAMI 04 BELL 10.6 $e^+ e^-$	NODE=M172R9 NODE=M172R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.005 95 AUBERT 06P BABR 10.6 $e^+ e^-$
<0.019 90 BESSON 03 CLE2 10.6 $e^+ e^-$

$\Gamma(D_s^+ \pi^0 \pi^0)/\Gamma(D_s^+ \pi^0)$	Γ_7/Γ_1
<0.25 95 AUBERT 06P BABR 10.6 $e^+ e^-$	NODE=M172R10 NODE=M172R10

NODE=M172215;NODE=M172

NODE=M172

$D_{s0}^*(2317)^\pm$ REFERENCES

ABLIKIM	18J	PR D97 051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHOI	15A	PR D91 092011	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)
AUBERT	03G	PRL 90 242001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSON	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)

NODE=M172

REFID=58895
 REFID=56577
 REFID=51144
 REFID=49747
 REFID=50195
 REFID=49629
 REFID=49417
 REFID=49583
 REFID=49615

$D_{s1}(2460)^\pm$

$$I(J^P) = 0(1^+)$$

$D_{s1}(2460)^\pm$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2459.5±0.6 OUR FIT Error includes scale factor of 1.1.

2459.6±0.9 OUR AVERAGE Error includes scale factor of 1.3.

2460.1±0.2±0.8 1 AUBERT 06P BABR 10.6 $e^+ e^-$

2458.0±1.0±1.0 195 AUBERT 04E BABR 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2459.5±1.2±3.7	920	AUBERT	06P BABR	$10.6 \quad e^+ e^- \rightarrow D_s^+ \gamma X$
2458.6±1.0±2.5	560	AUBERT	06P BABR	$10.6 \quad e^+ e^- \rightarrow D_s^+ \pi^0 \gamma X$
2460.2±0.2±0.8	123	AUBERT	06P BABR	$10.6 \quad e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$
2458.9±1.5	112	2 AUBERT,B	04S BABR	$B \rightarrow D_{s1}(2460)^+ \bar{D}^(*)$
2461.1±1.6	139	3 AUBERT,B	04S BABR	$B \rightarrow D_{s1}(2460)^+ \bar{D}^(*)$
2456.5±1.3±1.3	126	4,5 MIKAMI	04 BELL	$10.6 \quad e^+ e^-$
2459.5±1.3±2.0	152	6,7 MIKAMI	04 BELL	$10.6 \quad e^+ e^-$
2459.9±0.9±1.6	60	6,7 MIKAMI	04 BELL	$10.6 \quad e^+ e^-$
2459.2±1.6±2.0	57	KROKOVNY	03B BELL	$10.6 \quad e^+ e^-$

1 The average of the values obtained from the $D_s^+ \gamma$, $D_s^+ \pi^0 \gamma$, $D_s^+ \pi^+ \pi^-$ final state.

2 Systematic errors not evaluated. From the decay to $D_s^{*+} \pi^0$.

3 Systematic errors not evaluated. From the decay to $D_s^+ \gamma$.

4 Not independent of the corresponding $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$.

5 Using $m_{D_s^{*+}} = 2112.4 \pm 0.7$ MeV.

6 Not independent of the corresponding $m_{D_{s1}(2460)^\pm} - m_{D_s^\pm}$.

7 Using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV.

NODE=M173M

NODE=M173M

NODE=M173M

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

OCCUR=2

OCCUR=3

NODE=M173M;LINKAGE=UB

NODE=M173M;LINKAGE=AU

NODE=M173M;LINKAGE=AB

NODE=M173M;LINKAGE=B1

NODE=M173M;LINKAGE=B2

NODE=M173M;LINKAGE=B3

NODE=M173M;LINKAGE=B4

NODE=M173MD

NODE=M173MD

NODE=M173MD

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------


347.3±0.7 OUR FIT Error includes scale factor of 1.2.

347.1±2.2 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

344.1±1.3±1.1	126	MIKAMI	04 BELL	$10.6 \quad e^+ e^-$
351.2±1.7±1.0	41	BESSON	03 CLE2	$10.6 \quad e^+ e^-$
346.8±1.6±1.9	57	8 KROKOVNY	03B BELL	$10.6 \quad e^+ e^-$

⁸ Recalculated by us using $m_{D_s^{*+}} = 2112.4 \pm 0.7$ MeV.

NODE=M173MD;LINKAGE=K3



$m_{D_{s1}(2460)^{\pm}} - m_{D_s^{*\pm}}$

The fit includes D^{\pm} , D^0 , D_s^{\pm} , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^{\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

491.2 \pm 0.6 OUR FIT Error includes scale factor of 1.1.

491.3 \pm 1.4 OUR AVERAGE

491.0 \pm 1.3 \pm 1.9	152	9 MIKAMI	04 BELL	10.6 $e^+ e^-$
491.4 \pm 0.9 \pm 1.5	60	10 MIKAMI	04 BELL	10.6 $e^+ e^-$

9 From the decay to $D_s^{\pm} \gamma$.

10 From the decay to $D_s^{\pm} \pi^+ \pi^-$.

NODE=M173DM

NODE=M173DM

NODE=M173DM

OCCUR=2

NODE=M173DM;LINKAGE=M1

NODE=M173DM;LINKAGE=M2

NODE=M173W

NODE=M173W

OCCUR=2

NODE=M173215;NODE=M173

NODE=M173

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.5	95	123	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6.3	95	560	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^0 \gamma X$
< 10		195	AUBERT	04E BABR	10.6 $e^+ e^-$
< 5.5	90	126	MIKAMI	04 BELL	10.6 $e^+ e^-$
< 7	90	41	BESSON	03 CLE2	10.6 $e^+ e^-$

$D_{s1}(2460)^+$ DECAY MODES

$D_{s1}(2460)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 D_s^{*+} \pi^0$	(48 \pm 11) %	
$\Gamma_2 D_s^+ \gamma$	(18 \pm 4) %	
$\Gamma_3 D_s^+ \pi^+ \pi^-$	(4.3 \pm 1.3) %	S=1.1
$\Gamma_4 D_s^{*+} \gamma$	< 8 %	CL=90%

DESIG=1

DESIG=2

DESIG=3

DESIG=4

Γ_5	$D_{s0}^*(2317)^+ \gamma$	(3.7 $^{+ 5.0}_{- 2.4}$) %	DESIG=5
Γ_6	$D_s^+ \pi^0$		DESIG=7
Γ_7	$D_s^+ \pi^0 \pi^0$		DESIG=8
Γ_8	$D_s^+ \gamma \gamma$		DESIG=9

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 8 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 3.4$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	80			
x_3	68	62		
x_5	-3	25	26	
	x_1	x_2	x_3	

$D_{s1}(2460)^{\pm}$ BRANCHING RATIOS

$\Gamma(D_s^{*+} \pi^0) / \Gamma_{\text{total}}$	Γ_1 / Γ				
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

0.48 \pm 0.11 OUR FIT

0.56 \pm 0.13 \pm 0.09 11 AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^- \bar{D}^(*)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 41 BESSON 03 CLE2 10.6 $e^+ e^-$

11 Evaluated in AUBERT 06N including measurements from AUBERT,B 04S.

$\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$	Γ_2 / Γ			
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

0.18 \pm 0.04 OUR FIT

0.16 \pm 0.04 \pm 0.03 12 AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^- \bar{D}^(*)$

12 Evaluated in AUBERT 06N including measurements from AUBERT,B 04S.

$\Gamma(D_s^+ \gamma) / \Gamma(D_s^{*+} \pi^0)$	Γ_2 / Γ_1					
<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

0.38 \pm 0.05 OUR FIT
0.44 \pm 0.09 OUR AVERAGE

0.55 \pm 0.13 \pm 0.08 152 MIKAMI 04 BELL 10.6 $e^+ e^-$

0.38 \pm 0.11 \pm 0.04 38 KROKOVNY 03B BELL 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.274 \pm 0.045 \pm 0.020 251 13 AUBERT,B 04S BABR $B \rightarrow D_{s1}(2460)^+ \bar{D}^(*)$

<0.49 90 BESSON 03 CLE2 10.6 $e^+ e^-$

13 Used by AUBERT 06N in their measurement of $B(D_s^{*-} \pi^0)$ and $B(D_s^- \gamma)$.

$\Gamma(D_s^+ \pi^+ \pi^-) / \Gamma(D_s^{*+} \pi^0)$	Γ_3 / Γ_1					
<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

0.090 \pm 0.020 OUR FIT Error includes scale factor of 1.2.

0.14 \pm 0.04 \pm 0.02 60 MIKAMI 04 BELL 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.08 90 BESSON 03 CLE2 10.6 $e^+ e^-$

$\Gamma(D_s^{*+} \gamma) / \Gamma(D_s^{*+} \pi^0)$	Γ_4 / Γ_1				
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	

<0.16 90 BESSON 03 CLE2 10.6 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.31 90 MIKAMI 04 BELL 10.6 $e^+ e^-$

NODE=M173220

NODE=M173R1

NODE=M173R1

NODE=M173R1;LINKAGE=AU

NODE=M173R6

NODE=M173R6

NODE=M173R6;LINKAGE=AU

NODE=M173R2

NODE=M173R2

NODE=M173R2;LINKAGE=AU

NODE=M173R3

NODE=M173R3

NODE=M173R4

NODE=M173R4

$\Gamma(D_{s0}^*(2317)^+\gamma)/\Gamma(D_s^{*+}\pi^0)$				Γ_5/Γ_1	NODE=M173R5 NODE=M173R5
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.22	95	AUBERT	04E	BABR 10.6 e ⁺ e ⁻	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.58	90	BESSON	03	CLE2 10.6 e ⁺ e ⁻	
$\Gamma(D_s^{*+}\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_1/(\Gamma_1+\Gamma_5)$	NODE=M173R7 NODE=M173R7
VALUE	DOCUMENT ID	TECN	COMMENT		
0.93±0.09 OUR FIT					
0.97±0.09±0.05	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_s^+\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_2/(\Gamma_1+\Gamma_5)$	NODE=M173R8 NODE=M173R8
VALUE	DOCUMENT ID	TECN	COMMENT		
0.35 ±0.04 OUR FIT					
0.337±0.036±0.038	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_s^+\pi^+\pi^-)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_3/(\Gamma_1+\Gamma_5)$	NODE=M173R9 NODE=M173R9
VALUE	DOCUMENT ID	TECN	COMMENT		
0.083±0.017 OUR FIT	Error includes scale factor of 1.2.				
0.077±0.013±0.008	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	
$\Gamma(D_s^{*+}\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_4/(\Gamma_1+\Gamma_5)$	NODE=M173R10 NODE=M173R10
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.24	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
$\Gamma(D_{s0}^*(2317)^+\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_5/(\Gamma_1+\Gamma_5)$	NODE=M173R11 NODE=M173R11
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.25	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
$\Gamma(D_s^+\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_6/(\Gamma_1+\Gamma_5)$	NODE=M173R12 NODE=M173R12
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.042	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
$\Gamma(D_s^+\pi^0\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_7/(\Gamma_1+\Gamma_5)$	NODE=M173R13 NODE=M173R13
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.68	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	
$\Gamma(D_s^+\gamma\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$				$\Gamma_8/(\Gamma_1+\Gamma_5)$	NODE=M173R14 NODE=M173R14
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.33	95	AUBERT	06P	BABR 10.6 e ⁺ e ⁻	

$D_{s1}(2460)^{\pm}$ REFERENCES

AUBERT	06N	PR D74 031103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51142
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
BESSON	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

NODE=M173

$D_{s1}(2536)^{\pm}$

$I(J^P) = 0(1^+)$
 J, P need confirmation.

Seen in $D^*(2010)^+ K^0$, $D^*(2007)^0 K^+$, and $D_s^+ \pi^+ \pi^-$. Not seen
in $D^+ K^0$ or $D^0 K^+$. $J^P = 1^+$ assignment strongly favored.

NODE=M121

NODE=M121

NODE=M121M

NODE=M121M

NODE=M121M

$D_{s1}(2536)^{\pm}$ MASS

The fit includes $D^{\pm}, D^0, D_s^{\pm}, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^{\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2535.11±0.06 OUR FIT				
2535.21±0.28 OUR AVERAGE				
2537.7 ± 0.5 ± 3.1	24	1 ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
2535.7 ± 0.6 ± 0.5	46	2 ABAZOV	09G D0	$B_s^0 \rightarrow D_s^- \mu^+ \nu_\mu X$
2534.78 ± 0.31 ± 0.40	182	AUBERT	08B BABR	$B \rightarrow \bar{D}^{(*)} D^* K$
2534.6 ± 0.3 ± 0.7	193	AUBERT	06P BABR	$10.6 e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$
2535.3 ± 0.7	92	3 HEISTER	02B ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X, D^{*0} K^+ X$
2534.2 ± 1.2	9	ASRATYAN	94 BEBC	$\nu N \rightarrow D^* K^0 X, D^{*0} K^\pm X$
2535 ± 0.6 ± 1	75	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X, D^{*0} K^+ X$
2535.2 ± 0.5 ± 1.5	28	ALBRECHT	92R ARG	$10.4 e^+ e^- \rightarrow D^{*0} K^+ X$
2536.6 ± 0.7 ± 0.4		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.9 ± 0.6 ± 2.0		ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2534.1 ± 0.6	116	4 AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2535.08 ± 0.01 ± 0.15	8038	5 LEES	11B BABR	$10.6 e^+ e^- \rightarrow D^{*+} K_S^0 X$
2535.57 $^{+0.44}_{-0.41}$ ± 0.10	236	6 CHEKANOV	09 ZEUS	$e^{\pm} p \rightarrow D^{*+} K_S^0 X, D^{*0} K^+ X$
2535.3 ± 0.2 ± 0.5	134	7 ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*0} K^+ X$
2534.8 ± 0.6 ± 0.6	44	8 ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535 ± 28		9 ASRATYAN	88 HLBC	$\nu N \rightarrow D_s \gamma\gamma X$

1 From a fit of the D_s^+ recoil mass distribution with an incoherent sum of the S-wave and D-wave Breit-Wigner line shapes.

2 Using the $D^*(2010)^{\pm}$ mass of 2010.0 ± 0.4 MeV from PDG 06.

3 Calculated using $m(D^*(2010)^{\pm}) = 2010.0 \pm 0.5$ MeV, $m(D^*(2007)^0) = 2006.7 \pm 0.5$ MeV, and the mass difference below.

4 Systematic uncertainties not evaluated.

5 Calculated using the mass difference $m(D_{s1}^+) - m(D^{*+})_{PDG}$ below and $m(D^{*+})_{PDG} = 2010.25 \pm 0.14$ MeV. Assuming S-wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$, using a Breit-Wigner line shape corresponding to $L=0$.

6 Calculated using the mass difference $m(D_{s1}^+) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV.

7 Calculated using $m(D^*(2007)^0) = 2006.6 \pm 0.5$ MeV and the mass difference below.

8 Calculated using $m(D^*(2010)^{\pm}) = 2010.1 \pm 0.6$ MeV and the mass difference below.

9 Not seen in $D^* K$.

OCCUR=2

NODE=M121M;LINKAGE=D

NODE=M121M;LINKAGE=AB

NODE=M121M;LINKAGE=HI

NODE=M121M;LINKAGE=AU

NODE=M121M;LINKAGE=LE

NODE=M121M;LINKAGE=CH

NODE=M121M;LINKAGE=A

NODE=M121M;LINKAGE=C

NODE=M121M;LINKAGE=B

NODE=M121DM

NODE=M121DM

NODE=M121DM

$m_{D_{s1}(2536)^{\pm}} - m_{D_s^*(2111)}$

The fit includes $D^{\pm}, D^0, D_s^{\pm}, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^{\pm}$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
422.9± 0.4 OUR FIT			
424 ± 28	ASRATYAN 88 HLBC	$D_s^{*\pm} \gamma$	

NODE=M121DN

$m_{D_{s1}(2536)^{\pm}} - m_{D^*(2010)^{\pm}}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DN

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

524.85±0.04 OUR FIT**524.84±0.04 OUR AVERAGE**

524.83±0.01±0.04	8038	10 LEES	11B BABR	$10.6 e^+ e^- \rightarrow D^{*+} K_S^0 X$
525.30 ^{+0.44} _{-0.41} ± 0.10	236 ± 30	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X$
525.3 ± 0.6 ± 0.1	41	HEISTER 02B	ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X$
524.7 ± 0.6 ± 0.2	44	ALEXANDER93	CLE2	$e^+ e^- \rightarrow D^{*+} K_S^0 X$

10 Assuming S -wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$, using a Breit-Wigner line shape corresponding to $L=0$.

NODE=M121DN

OCCUR=2

NODE=M121DN;LINKAGE=LE

 $m_{D_{s1}(2536)^\pm} = m_{D^*(2007)^0}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DP

NODE=M121DP

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
528.26±0.05 OUR FIT		Error includes scale factor of 1.2.		
528.68±0.28 OUR AVERAGE				
528.7 ± 1.9 ± 0.5	51	HEISTER 02B	ALEP	$e^+ e^- \rightarrow D^{*0} K^+ X$
527.3 ± 2.2	29	ACKERSTAFF 97W	OPAL	$e^+ e^- \rightarrow D^{*0} K^+ X$
528.7 ± 0.2 ± 0.2	134	ALEXANDER 93	CLE2	$e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121DP

OCCUR=2

NODE=M121W

NODE=M121W

 $D_{s1}(2536)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.92±0.05 OUR AVERAGE					
1.7 ± 1.2 ± 0.6	24	11 ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$	
0.92±0.03±0.04	8038	12 LEES	11B BABR	$10.6 e^+ e^- \rightarrow D^{*+} K_S^0 X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.75±0.23	116	13 AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ + D^{(*)}$	
< 2.5	95	193 AUBERT	06P BABR	$10.6 e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$	
< 3.2	90	75 FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X$, $D^{*0} K^+ X$	
< 2.3	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$	
< 3.9	90	ALBRECHT 92R	ARG	$10.4 e^+ e^- \rightarrow D^{*0} K^+ X$	
< 5.44	90	AVERY 90	CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$	
< 4.6	90	ALBRECHT 89E	ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$	

NODE=M121W;LINKAGE=A

11 From a fit of the D_s^+ recoil mass distribution with an incoherent sum of the S -wave and S -wave Breit-Wigner line shapes.

12 Assuming S -wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$, using a Breit-Wigner line shape corresponding to $L=0$.

13 Systematic uncertainties not evaluated.

NODE=M121W;LINKAGE=LE

NODE=M121W;LINKAGE=AU

NODE=M121215;NODE=M121

NODE=M121

 $D_{s1}(2536)^+$ DECAY MODES

$D_{s1}(2536)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 D^*(2010)^+ K^0$	0.85 ± 0.12	
$\Gamma_2 (D^*(2010)^+ K^0)_{S-wave}$	0.61 ± 0.09	
$\Gamma_3 (D^*(2010)^+ K^0)_{D-wave}$		
$\Gamma_4 D^+ \pi^- K^+$	0.028±0.005	
$\Gamma_5 D^*(2007)^0 K^+$	DEFINED AS 1	
$\Gamma_6 D^+ K^0$	<0.34	90%
$\Gamma_7 D^0 K^+$	<0.12	90%
$\Gamma_8 D_s^{*+} \gamma$	possibly seen	
$\Gamma_9 D_s^+ \pi^+ \pi^-$	seen	

DESIG=1

DESIG=7

DESIG=9

DESIG=8

DESIG=4

DESIG=2

DESIG=5

DESIG=3

DESIG=6

$D_{s1}(2536)^+$ BRANCHING RATIOS

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^*(2010)^+ K^0)$					Γ_5/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
1.18±0.16 OUR AVERAGE					
0.88±0.24±0.08	116	AUSHEV	11	BELL $B \rightarrow D_{s1}(2536)^+ D^0$	
2.3 ± 0.6 ± 0.3	236 ± 30	CHEKANOV	09	ZEUS $e^\pm p \rightarrow D^{*+} K_S^0 X$	
1.32±0.47±0.23	92	14 HEISTER	02B	ALEP $e^+ e^- \rightarrow D^{*0} K^+ X$	
1.9 $^{+1.1}_{-0.9}$ ± 0.4	35	14 ACKERSTAFF	97W	OPAL $e^+ e^- \rightarrow D^{*0} K^+ X$	
1.1 ± 0.3		ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*+} K^0 X$	
1.4 ± 0.3 ± 0.2		15 ALBRECHT	92R	ARG $10.4 e^+ e^- \rightarrow D^{*0} K^+ X, D^{*+} K^0 X$	

14 Ratio of the production rates measured in Z^0 decays.

15 Evaluated by us from published inclusive cross-sections.

$\Gamma((D^*(2010)^+ K^0)_{S-wave})/\Gamma(D^*(2010)^+ K^0)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.72±0.05±0.01	5485	BALAGURA	08	BELL $10.6 e^+ e^- \rightarrow D^{*+} K^0 X$	

$\Gamma(D^+ \pi^- K^+)/\Gamma(D^*(2010)^+ K^0)$					Γ_4/Γ_1
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.27±0.18±0.37	1264	BALAGURA	08	BELL $10.6 e^+ e^- \rightarrow D^+ \pi^- K^+ X$	

$\Gamma(D^+ K^0)/\Gamma(D^*(2010)^+ K^0)$					Γ_6/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.40	90	ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*+} K^0 X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.43	90	ALBRECHT	89E	ARG $D_{s1}^* \rightarrow D^*(2010) K^0$	

$\Gamma(D^0 K^+)/\Gamma(D^*(2007)^0 K^+)$					Γ_7/Γ_5
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.12	90	ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*0} K^+ X$	

$\Gamma(D_s^{*+} \gamma)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
possibly seen	ASRATYAN	88	HLBC $\nu N \rightarrow D_s \gamma \gamma X$		

$\Gamma(D_s^{*+} \gamma)/\Gamma(D^*(2007)^0 K^+)$					Γ_8/Γ_5
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.42	90	ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*0} K^+ X$	

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	AUBERT	06P	BABR $10.6 e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$		

 $D_{s1}(2536)^{\pm}$ REFERENCES

ABLIKIM	19P	CP C43 031001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59617
AUSHEV	11	PR D83 051102	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=16505
LEES	11B	PR D83 072003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16773
ABAZOV	09G	PRL 102 051801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52652
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BALAGURA	08	PR D77 032001	V. Balagura <i>et al.</i>	(BELLE Collab.)	REFID=52133
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN	94	ZPHY C61 563	A.E. Asratyan <i>et al.</i>	(BIRN, BELG, CERN+) (FNAL E687 Collab.)	REFID=43667
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(CLEO Collab.)	REFID=43687
ALEXANDER	93	PL B303 377	J. Alexander <i>et al.</i>	(ARGUS Collab.)	REFID=43316
ALBRECHT	92R	PL B297 425	H. Albrecht <i>et al.</i>	(CLEO Collab.)	REFID=43179
AVERY	90	PR D41 774	P. Avery, D. Besson	(ARGUS Collab.)	REFID=41013
ALBRECHT	89E	PL B230 162	H. Albrecht <i>et al.</i>	(ITEP, SERP)	REFID=40914
ASRATYAN	88	ZPHY C40 483	A.E. Asratyan <i>et al.</i>		REFID=40916

NODE=M121220

NODE=M121R6

NODE=M121R6

NODE=M121R6;LINKAGE=6A

NODE=M121R6;LINKAGE=A

NODE=M121R8

NODE=M121R8

NODE=M121R9

NODE=M121R9

NODE=M121R1

NODE=M121R1

NODE=M121R4

NODE=M121R4

NODE=M121R3

NODE=M121R3

NODE=M121R5

NODE=M121R5

NODE=M121

NODE=M148

 $D_{s2}^*(2573)$

$I(J^P) = 0(2^+)$

J^P is natural, width and decay modes consistent with 2^+ .
 AAIJ 14AW confirms $J^P = 2^+$.

NODE=M148

 $D_{s2}^*(2573)$ MASS

NODE=M148M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2569.1 ±0.8 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
2570.7 ±2.0 ±1.7	62	¹ ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
2568.39±0.29±0.26		AAIJ	14AWLHCb	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
2569.4 ±1.6 ±0.5	82	AAIJ	11A LHCb	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$
2572.2 ±0.3 ±1.0		AUBERT,BE	06E BABR	$e^+ e^- \rightarrow D K X$
2574.5 ±3.3 ±1.6		ALBRECHT	96 ARG	$e^+ e^- \rightarrow D^0 K^+ X$
2573.2 $\begin{array}{l} +1.7 \\ -1.6 \end{array}$ ±0.9	217	KUBOTA	94 CLE2	$e^+ e^- \sim 10.5 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2570.0 ±4.3	25	² EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$
2568.6 ±3.2	64	³ HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$

1 From a fit of the D_s^+ recoil mass distribution .


2 Not independent of the mass difference below.

3 Calculated using $m_{D^0} = 1864.5 \pm 0.5 \text{ MeV}$ and the mass difference below.

NODE=M148M

NODE=M148M

NODE=M148M



NODE=M148M;LINKAGE=A

NODE=M148M;LINKAGE=EV

NODE=M148M;LINKAGE=HI

$m_{D_{s2}^*(2573)} = m_{D^0}$

NODE=M148DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
704 ±3 ±1	64	HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
705.4±4.3	25	¹ EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$

NODE=M148DM

NODE=M148DM

1 Systematic errors not estimated.

NODE=M148DM;LINKAGE=EV

 $D_{s2}^*(2573)$ WIDTH

NODE=M148W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.9±0.7 OUR AVERAGE				
17.2±3.6±1.1	62	¹ ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$

NODE=M148W

NODE=M148W

16.9±0.5±0.6		AAIJ	14AWLHCb	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
12.1±4.5±1.6	82	AAIJ	11A LHCb	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$
27.1±0.6±5.6		AUBERT,BE	06E BABR	$e^+ e^- \rightarrow D K X$
10.4±8.3±3.0		ALBRECHT	96 ARG	$e^+ e^- \rightarrow D^0 K^+ X$
16 ± 5 ± 3	217	KUBOTA	94 CLE2	$e^+ e^- \sim 10.5 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 9 -6	25	² EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$
------------------	----	------------------------	---------	--

¹ From a fit of the D_s^+ recoil mass distribution .

² Systematic errors not estimated.

$D_{s2}^*(2573)^+$ DECAY MODES

$D_{s2}^*(2573)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0 K^+$	seen
$\Gamma_2 D^*(2007)^0 K^+$	not seen

$D_{s2}^*(2573)^+$ BRANCHING RATIOS

$\Gamma(D^0 K^+)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
seen	217 KUBOTA 94 CLE2 \pm $e^+ e^- \sim 10.5 \text{ GeV}$
$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^0 K^+)$	Γ_2/Γ_1
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>CHG</u> <u>COMMENT</u>
<0.33	90 KUBOTA 94 CLE2 + $e^+ e^- \sim 10.5 \text{ GeV}$

$D_{s2}^*(2573)$ REFERENCES

ABLIKIM	19P	CP C43 031001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	14AW	PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	11A	PL B698 14	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT,BE	06E	PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)
EVDOKEKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)
ALBRECHT	96	ZPHY C69 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
KUBOTA	94	PRL 72 1972	Y. Kubota <i>et al.</i>	(CLEO Collab.)

NODE=M148W;LINKAGE=A
NODE=M148W;LINKAGE=EV

NODE=M148215;NODE=M148

NODE=M148

DESIG=1
DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M148220

NODE=M148R2
NODE=M148R2

NODE=M148R1
NODE=M148R1

NODE=M148

REFID=59617
REFID=56105
REFID=16665
REFID=51512
REFID=50337
REFID=48562
REFID=44631
REFID=43781

NODE=M182

 $D_{s1}^*(2700)^\pm$ $I(J^P) = 0(1^-)$ **$D_{s1}^*(2700)^+ \text{ MASS}$**

NODE=M182M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2708.3 \pm 4.0 OUR AVERAGE				

2699 \pm 14 2709.2 \pm 1.9 \pm 4.5 2710 \pm 2 \pm 12 2708 \pm 9 \pm 11	7 52k 10.4k 182	1 LEES 2 AAIJ 3 AUBERT BRODZICKA	15C BABR 12AU LHCb 09AR BABR 08 BELL	$B \rightarrow DD^0 K^+$ $pp \rightarrow (DK)^+ X$ at 7 TeV $e^+ e^- \rightarrow D^{(*)} K X$ $B^+ \rightarrow D^0 \bar{D}^0 K^+$
---	--------------------------	---	---	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

2694 \pm 8 \pm 13 2707 \pm 8 \pm 8 2688 \pm 4 \pm 3	LEES LEES 4 AUBERT,BE	15C BABR 15C BABR 06E BABR	$B^0 \rightarrow D^- D^0 K^+$ $B^+ \rightarrow \bar{D}^0 D^0 K^+$ $10.6 e^+ e^- \rightarrow DK X$
---	-----------------------------	----------------------------------	---

1 From a combined analysis of $B^0 \rightarrow D^- D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 D^0 K^+$.

2 From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{sJ}^*(2860)^+$.

3 From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.

4 Superseded by AUBERT 09AR.

OCCUR=2

OCCUR=3

NODE=M182M;LINKAGE=B

NODE=M182M;LINKAGE=AA

NODE=M182M;LINKAGE=AB

NODE=M182M;LINKAGE=AU

 $D_{s1}^*(2700)^+ \text{ WIDTH}$

NODE=M182W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
120 \pm 11 OUR AVERAGE				

127 \pm 24 115.8 \pm 7.3 \pm 12.1 149 \pm 7 \pm 39 108 \pm 23 \pm 36	52k 10.4k 182	5 LEES 7 AUBERT BRODZICKA	15C BABR 09AR BABR 08 BELL	$B \rightarrow DD^0 K^+$ $e^+ e^- \rightarrow D^{(*)} K X$ $B^+ \rightarrow D^0 \bar{D}^0 K^+$
---	---------------------	---------------------------------	----------------------------------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

145 \pm 24 \pm 22 113 \pm 21 \pm 20 112 \pm 7 \pm 36	LEES LEES 8 AUBERT,BE	15C BABR 15C BABR 06E BABR	$B^0 \rightarrow D^- D^0 K^+$ $B^+ \rightarrow \bar{D}^0 D^0 K^+$ $10.6 e^+ e^- \rightarrow DK X$
--	-----------------------------	----------------------------------	---

OCCUR=2

OCCUR=3

NODE=M182W;LINKAGE=A

NODE=M182W;LINKAGE=AA

NODE=M182W;LINKAGE=AB

NODE=M182W;LINKAGE=AU

5 From a combined analysis of $B^0 \rightarrow D^- D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 D^0 K^+$.

6 From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{sJ}^*(2860)^+$.

7 From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.

8 Superseded by AUBERT 09AR.

NODE=M182215;NODE=M182

 $D_{s1}^*(2700)^\pm \text{ DECAY MODES}$

Mode

Γ_1	DK
Γ_2	$D^0 K^+$
Γ_3	$D^+ K_S^0$
Γ_4	$D^* K$
Γ_5	$D^{*0} K^+$
Γ_6	$D^{*+} K_S^0$

DESIG=2

DESIG=1

DESIG=3

DESIG=4

DESIG=5

DESIG=6

$D_{s1}^*(2700)^{\pm}$ BRANCHING RATIOS

$\Gamma(D^* K)/\Gamma(DK)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
0.91±0.13±0.12	10.4k	9 AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$ 9 From the average of the corresponding ratios with $D^{(*)}0 K^+$ and $D^{(*)}+ K_S^0$.	

$\Gamma(D^{*0} K^+)/\Gamma(D^0 K^+)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.88±0.14±0.14	7716	10 AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$ 10 From the $D^{*0} K^+$ and $D^0 K^+$, where $D^{*0} \rightarrow D^0 \pi^0$.	

$\Gamma(D^{*+} K_S^0)/\Gamma(D^+ K_S^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.14±0.39±0.23	2700	11 AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$ 11 From the $D^{*+} K_S^0$ and $D^+ K_S^0$, where $D^{*+} \rightarrow D^+ \pi^0$.	

$D_{s1}^*(2700)^{\pm}$ REFERENCES

LEES	15C	PR D91 052002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	12AU	JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR	PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
AUBERT,BE	06E	PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

$D_{s1}^*(2860)^{\pm}$

$$I(J^P) = 0(1^-)$$

OMITTED FROM SUMMARY TABLE

J^P consistent with 1^- from angular analysis of AAIJ 14AW. Observed by AUBERT,BE 06E and AUBERT 09AR in inclusive production of DK and $D^* K$ in $e^+ e^-$ annihilation.

$D_{s1}^*(2860)^{+}$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2859 ±12 ±24		1 AAIJ	14AW LHCb	$B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

2866.1± 1.0± 6.3	36k	2,3 AAIJ	12AU LHCb	$p p \rightarrow (DK)^+ X$ at 7 TeV
2862 ± 2 + 5 - 2	3122	3,4 AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$
2856.6± 1.5± 5.0		5 AUBERT,BE	06E BABR	$e^+ e^- \rightarrow DK X$

1 Separated from the spin-3 component $D_{s3}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .
2 From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{sJ}^*(2860)^+$.
3 Possible contribution from the $D_{s3}^*(2860)$ state.
4 From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.
5 Superseded by AUBERT 09AR.

NODE=M182225

NODE=M182R01
NODE=M182R01

NODE=M182R01;LINKAGE=AU

NODE=M182R02
NODE=M182R02

NODE=M182R02;LINKAGE=AU

NODE=M182R03
NODE=M182R03

NODE=M182R03;LINKAGE=AU

NODE=M182

REFID=56412
REFID=54735
REFID=53135
REFID=52144
REFID=51512

NODE=M196

NODE=M196

NODE=M196M

NODE=M196M

NODE=M196M;LINKAGE=A

NODE=M196M;LINKAGE=AA

NODE=M196M;LINKAGE=B
NODE=M196M;LINKAGE=AB

NODE=M196M;LINKAGE=AU

NODE=M196W

NODE=M196W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
159 ±23 ±77		1 AAIJ	14AW LHCb	$B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

69.9± 3.2± 6.6	36k	2,3 AAIJ	12AU LHCb	$p p \rightarrow (DK)^+ X$ at 7 TeV
48 ± 3 ± 6	3122	3,4 AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$
47 ± 7 ± 10		5 AUBERT,BE	06E BABR	$e^+ e^- \rightarrow DK X$

- 1 Separated from the spin-3 component $D_{s3}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .
- 2 From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{sJ}^*(2860)^+$.
- 3 Possible contribution from the $D_{s3}^*(2860)$ state.
- 4 From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.
- 5 Superseded by AUBERT 09AR.

$D_{s1}^*(2860)^\pm$ DECAY MODES

Mode
$\Gamma_1 D K$
$\Gamma_2 D^0 K^+$
$\Gamma_3 D^+ K_S^0$
$\Gamma_4 D^* K$
$\Gamma_5 D^{*0} K^+$
$\Gamma_6 D^{*+} K_S^0$

$D_{s1}^*(2860)^\pm$ BRANCHING RATIOS

$\Gamma(D^* K)/\Gamma(DK)$	Γ_4/Γ_1
$1.10 \pm 0.15 \pm 0.19$	3122 1 AUBERT 09AR BABR $e^+ e^- \rightarrow D^{(*)} K X$

¹ From the average of the corresponding ratios with $D^{(*)0} K^+$ and $D^{(*)+} K_S^0$.

$\Gamma(D^{*0} K^+)/\Gamma(D^0 K^+)$	Γ_5/Γ_2
$1.04 \pm 0.17 \pm 0.20$	2241 1 AUBERT 09AR BABR $e^+ e^- \rightarrow D^{(*)} K X$

¹ From the $D^{*0} K^+$ and $D^0 K^+$, where $D^{*0} \rightarrow D^0 \pi^0$.

$\Gamma(D^{*+} K_S^0)/\Gamma(D^+ K_S^0)$	Γ_6/Γ_3
$1.38 \pm 0.35 \pm 0.49$	881 1 AUBERT 09AR BABR $e^+ e^- \rightarrow D^{(*)} K X$

¹ From the $D^{*+} K_S^0$ and $D^+ K_S^0$, where $D^{*+} \rightarrow D^+ \pi^0$.

$D_{s1}^*(2860)^\pm$ REFERENCES

AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	12AU JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06E PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M196W;LINKAGE=A

NODE=M196W;LINKAGE=AA

NODE=M196W;LINKAGE=B

NODE=M196W;LINKAGE=AB

NODE=M196W;LINKAGE=AU

NODE=M196215;NODE=M196

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

NODE=M196225

NODE=M196R01

NODE=M196R01

NODE=M196R01;LINKAGE=AU

NODE=M196R02

NODE=M196R02

NODE=M196R02;LINKAGE=AU

NODE=M196R03

NODE=M196R03

NODE=M196R03;LINKAGE=AU

NODE=M196

REFID=56105

REFID=54735

REFID=53135

REFID=51512

NODE=M226

 $D_{s3}^*(2860)^\pm$ $I(J^P) = 0(3^-)$

OMITTED FROM SUMMARY TABLE

 J^P consistent with 3^- from angular analysis of AAIJ 14AW.

VALUE (MeV)

 $2860.5 \pm 2.6 \pm 6.5$ **$D_{s3}^*(2860)^+$ MASS**

DOCUMENT ID

1 AAIJ

TECN

14AWLHCb

COMMENT

 $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$

¹ Separated from the spin-1 component $D_{s1}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

VALUE (MeV)

 $53 \pm 7 \pm 7$ **$D_{s3}^*(2860)^+$ WIDTH**

DOCUMENT ID

1 AAIJ

TECN

14AWLHCb

COMMENT

 $B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$

¹ Separated from the spin-1 component $D_{s1}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

 $D_{s3}^*(2860)^\pm$ REFERENCES

AAIJ

14AW PRL 113 162001

R. Aaij *et al.*

(LHCb Collab.) JP

 $D_{sJ}(3040)^\pm$ $I(J^P) = 0(?)$

OMITTED FROM SUMMARY TABLE

Observed by AUBERT 09AR in inclusive production of $D^* K$ in $e^+ e^-$ annihilation.

VALUE (MeV)

 $3044 \pm 8^{+30}_{-5}$ **$D_{sJ}(3040)^+$ MASS**

DOCUMENT ID

AUBERT

TECN

09AR BABR $e^+ e^- \rightarrow D^* K X$

COMMENT

VALUE (MeV)

 $239 \pm 35^{+46}_{-42}$ **$D_{sJ}(3040)^+$ WIDTH**

DOCUMENT ID

AUBERT

TECN

09AR BABR $e^+ e^- \rightarrow D^* K X$

COMMENT

 $D_{sJ}(3040)^\pm$ DECAY MODES

Mode

Γ_1	$D^* K$
Γ_2	$D^{*0} K^+$
Γ_3	$D^{*+} K_S^0$

NODE=M197

NODE=M197M

NODE=M197W

NODE=M197215;NODE=M197

DESIG=1

DESIG=2

DESIG=3

AUBERT 09AR PR D80 092003

B. Aubert *et al.*

(BABAR Collab.)

NODE=M197

REFID=53135

BOTTOM MESONS ($B = \pm 1$)

$B^+ = u\bar{b}$, $B^0 = d\bar{b}$, $\bar{B}^0 = \bar{d}b$, $B^- = \bar{u}b$, similarly for B^* 's

$B_1(5721)^+$

$I(J^P) = \frac{1}{2}(1^+)$
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_1(5721)^+$ MASS

OUR FIT uses $m_{B^{*0}}$ and $m_{B_1^+} - m_{B^{*0}}$ to determine $m_{B_1(5721)^+}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

5725.9^{+2.5}_{-2.7} OUR FIT

$m_{B_1^+} - m_{B^{*0}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

401.2^{+2.4}_{-2.7} OUR FIT

401.2^{+2.4}_{-2.7} OUR AVERAGE

400.5 ± 1.8 ± 3.1	8K	¹ AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV
402 ± 3 ± 3		² AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_1^+} - m_{B^0}] - (m_{B^{*0}} - m_{B^0}) - m_{\pi^+} = 260.9 \pm 1.8 \pm 3.1$ MeV which we adjust by the π^+ mass and assume $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_1(5721)^+} - m_{B^{*0}} - m_{\pi^+} = 262 \pm 3^{+1}_{-3}$ MeV which we adjusted by the π^+ mass.

$B_1(5721)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31 ± 6 OUR AVERAGE				Error includes scale factor of 1.1.
29.1 ± 3.6 ± 4.3	8K	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV
49 ± 12 ± 2		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
-10 -13				

$B_1(5721)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 B^{*0} \pi^+$	seen

$B_1(5721)^+$ BRANCHING RATIOS

$\Gamma(B^{*0}\pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	8K AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV
seen	AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

$B_1(5721)^+$ REFERENCES

AAIJ AALTONEN	15AB 14I	JHEP 1504 024 PR D90 012013	R. Aaij <i>et al.</i> T. Aaltonen <i>et al.</i>	(LHCb Collab.) (CDF Collab.)
------------------	-------------	--------------------------------	--	---------------------------------

NODE=MXXX045

NODE=MXXX045

NODE=M218

NODE=M218

NODE=M218M

NODE=M218M

NODE=M218M

NODE=M218DM

NODE=M218DM

NODE=M218DM;LINKAGE=A

NODE=M218DM;LINKAGE=AA

NODE=M218W

NODE=M218W

NODE=M218215;NODE=M218

DESIG=1

NODE=M218220

NODE=M218R01
NODE=M218R01

NODE=M218

REFID=56628
REFID=56029

$B_1(5721)^0$
 $I(J^P) = \frac{1}{2}(1^+)$
I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_1(5721)^0$ MASSOUR FIT uses mass differences measurements listed below to determine the mass $m_{B_1(5721)^0}$.

VALUE (MeV)	DOCUMENT ID
5726.1±1.3 OUR FIT	Error includes scale factor of 1.2.

$m_{B_1^0} - m_{B^+}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
446.7±1.3 OUR FIT	Error includes scale factor of 1.2.		

441.5±2.4±1.3 ¹ ABAZOV 07T D0 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

446.2^{+1.9+1.0}_{-2.1-1.2} ¹ AALTONEN 09D CDF Repl. by AALTONEN 14I

¹ Observed in $B_1^0 \rightarrow B^{*+} \pi^-$.

$m_{B_1^0} - m_{B^{*+}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
401.4±1.2 OUR FIT	Error includes scale factor of 1.2.			

402.8±1.1 OUR AVERAGE

403.4 $\pm 0.7 \pm 1.5$ 35K ¹ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV

402.3 $\pm 0.9 \pm 1.1$ 2 AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_1^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 263.9 \pm 0.7 \pm 1.4$

MeV which we adjust by the π^- mass and $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_1(5721)^0} - m_{B^{*+}} - m_{\pi^-} = 262.7 \pm 0.9 \pm 1.1$ MeV which we adjusted by the π^- mass.

 $B_1(5721)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.5±3.4 OUR AVERAGE	Error includes scale factor of 1.1.			
30.1 $\pm 1.5 \pm 3.5$	35k	AAIJ 15AB LHCb	$p\bar{p}$ at 7, 8 TeV	
23 $\pm 3 \pm 4$	AALTONEN 14I	CDF	$p\bar{p}$ at 1.96 TeV	

 $B_1(5721)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B^{*+} \pi^-$	seen

 $B_1(5721)^0$ BRANCHING RATIOS

$\Gamma(B^{*+} \pi^-)/\Gamma_{\text{total}}$		Γ_1/Γ
seen	35K	AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV
seen		AALTONEN 09D CDF $p\bar{p}$ at 1.96 TeV
seen	¹ ABAZOV 07T D0	$p\bar{p}$ at 1.96 TeV

¹ Observed in $B_1^0 \rightarrow B^{*+} \pi^-$ with $B^{*+} \rightarrow B^+ \gamma$ and $B^+ \rightarrow J/\psi \pi^+$.

 $B_1(5721)^0$ REFERENCES

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D	PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T	PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)

NODE=M183

NODE=M183

NODE=M183M

NODE=M183M

NODE=M183M

NODE=M183DM

NODE=M183DM

NODE=M183DM;LINKAGE=AA

NODE=M183DM2

NODE=M183DM2

NODE=M183DM2;LINKAGE=B

NODE=M183DM2;LINKAGE=AA

NODE=M183W

NODE=M183W

NODE=M183215;NODE=M183

DESIG=1

NODE=M183220

NODE=M183R01

NODE=M183R01

NODE=M183R01;LINKAGE=AB

NODE=M183

REFID=56628

REFID=56029

REFID=52700

REFID=52014

NODE=M151

 $B_J^*(5732)$
 $I(J^P) = ?(?)$
I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

also known as B^{**}

Quantum numbers shown are quark-model predictions. Signal can be interpreted as stemming from several narrow and broad resonances.

NODE=M151

 $B_J^*(5732)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5698± 8 OUR AVERAGE				Error includes scale factor of 1.2.
5710±20		1 AFFOLDER	01F CDF	$p\bar{p}$ at 1.8 TeV
5695 ⁺¹⁷ ₋₁₉		2 BARATE	98L ALEP	$e^+ e^- \rightarrow Z$
5704± 4±10	1944	3 BUSKULIC	96D ALEP	$E_{cm}^{ee} = 88\text{--}94$ GeV
5732± 5±20	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88\text{--}94$ GeV
5681±11	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88\text{--}94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5713± 2		4 ACCIARRI	99N L3	$e^+ e^- \rightarrow Z$

¹AFFOLDER 01F uses the reconstructed B meson through semileptonic decay channels. The fraction of light B mesons that are produced at $L=1$ B^{**} states is measured to be $0.28 \pm 0.06 \pm 0.03$.

²BARATE 98L uses fully reconstructed B mesons to search for B^{**} production in the $B\pi^\pm$ system. In the framework of heavy quark symmetry (HQS), they also measured the mass of B_2^* to be 5739^{+8+6}_{-11-4} MeV/ c^2 and the relative production rate of $B(b \rightarrow B_2^* \rightarrow B^{(*)}\pi)/B(b \rightarrow B_{u,d}) = (31 \pm 9^{+6}_{-5})\%$.

³Using $m_{B\pi} - m_B = 424 \pm 4 \pm 10$ MeV.

⁴ACCIARRI 99N uses inclusive reconstructed B mesons to search for B^{**} production in the $B^{(*)}\pi^\pm$ system. In the framework of HQET, they measured the mass of B_1^* and B_2^* to be $5670 \pm 10 \pm 13$ MeV and $5768 \pm 5 \pm 6$ with the $B(b \rightarrow B^{**}) = (32 \pm 3 \pm 6) \times 10^{-2}$. They also reported the evidence for the existence of an excited B -meson state or mixture of states in the region 5.9–6.0 GeV.

NODE=M151M

NODE=M151M

NODE=M151M;LINKAGE=MF

NODE=M151M;LINKAGE=B

NODE=M151M;LINKAGE=A

NODE=M151M;LINKAGE=N

 $B_J^*(5732)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
128±18 OUR AVERAGE				
145±28	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88\text{--}94$ GeV
116±24	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88\text{--}94$ GeV

NODE=M151W

NODE=M151W

NODE=M151215;NODE=M151

 $B_J^*(5732)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B^*\pi + B\pi$	seen
$\Gamma_2 \quad B^*\pi(X)$	[a] (85±29) %

[a] X refers to decay modes with or without additional accompanying decay particles.

DESIG=1;OUR EST;→ UNCHECKED ← DESIG=2

LINKAGE=151

NODE=M151220

NODE=M151220

NODE=M151R1
NODE=M151R1 **$B_J^*(5732)$ BRANCHING RATIOS**

X refers to decay modes with or without additional accompanying decay particles.

$\Gamma(B^*\pi(X))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.85^{+0.26}_{-0.27}±0.12	ABBIENDI	02E OPAL	$e^+ e^- \rightarrow Z$	

$B_s^*(5732)$ REFERENCES

ABBIENDI	02E	EPJ C23 437	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	01F	PR D64 072002	T. Affolder <i>et al.</i>	(CDF Collab.)
ACCIARRI	99N	PL B465 323	M. Acciari <i>et al.</i>	(L3 Collab.)
BARATE	98L	PL B425 215	R. Barate <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96D	ZPHY C69 393	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	95B	PL B345 598	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	95E	ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)

 $B_2^*(5747)^+$

$$I(J^P) = \frac{1}{2}(2^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_2^*(5747)^+$ MASSOUR FIT uses $m_{B_s^0}$ and $m_{B_2^{*+}} - m_{B_s^0}$ to determine $m_{B_2^*(5747)^+}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5737.2 ± 0.7 OUR FIT				

$m_{B_2^{*+}} = m_{B^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
457.5 ± 0.7 OUR FIT				

457.5 ± 0.7 OUR AVERAGE

457.62 ± 0.72 ± 0.40	4K	1 AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV
457.3 ± 1.3 ± 0.9		2 AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

¹AAIJ 15AB reports $[m_{B_2^{*+}} - m_{B_s^0}] - m_{\pi^+} = 318.1 \pm 0.7 \pm 0.4$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event.

²AALTONEN 14I reports $m_{B_2^*(5747)^+} - m_{B_s^0} - m_{\pi^+} = 317.7 \pm 1.2^{+0.3}_{-0.9}$ MeV which we adjusted by the π^+ mass.

 $B_2^*(5747)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 ± 5 OUR AVERAGE				Error includes scale factor of 2.2.
23.6 ± 2.0 ± 2.1	4K	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV
11 +4 +3 -3 -4		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

 $B_2^*(5747)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 B^0 \pi^+$	seen
$\Gamma_2 B^{*0} \pi^+$	seen

 $B_2^*(5747)^+$ BRANCHING RATIOS

$\Gamma(B^0 \pi^+)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_1/Γ
seen	4K	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV
seen		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

$\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_2/Γ
seen	4k	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV

$\Gamma(B^{*0} \pi^+)/\Gamma(B^0 \pi^+)$	EVTS	DOCUMENT ID	TECN	Γ_2/Γ_1
1.0 ± 0.5 ± 0.8	4k	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV

NODE=M151

REFID=48742
REFID=48369
REFID=47247
REFID=46082
REFID=44677
REFID=44131
REFID=44182

NODE=M219

NODE=M219M

NODE=M219M

NODE=M219M

NODE=M219DM

NODE=M219DM

NODE=M219DM;LINKAGE=B

NODE=M219DM;LINKAGE=A

NODE=M219W

NODE=M219W

NODE=M219215;NODE=M219

DESIG=1
DESIG=2

NODE=M219220

NODE=M219R01
NODE=M219R01NODE=M219R02
NODE=M219R02NODE=M219R03
NODE=M219R03

$B_2^*(5747)^+$ REFERENCES

AAIJ 15AB JHEP 1504 024
AALTONEN 14I PR D90 012013

R. Aaij *et al.*
T. Aaltonen *et al.*

(LHCb Collab.)
(CDF Collab.)

$B_2^*(5747)^0$

$I(J^P) = \frac{1}{2}(2^+)$
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_2^*(5747)^0$ MASS

OUR FIT uses m_{B^+} , $m_{B_1^0} - m_{B^+}$, and $m_{B_2^{*0}} - m_{B_1^0}$ to determine $m_{B_2^*(5747)^0}$. The -0.659 correlation between statistical uncertainties of $m_{B_1^0} - m_{B^+}$ and $m_{B_2^{*0}} - m_{B_1^0}$ measurements reported by ABAZOV 07T is taken into account.

VALUE (MeV)	DOCUMENT ID
5739.5±0.7 OUR FIT	Error includes scale factor of 1.4.

$$m_{B_2^{*0}} - m_{B_1^0}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.4±1.4 OUR FIT	Error includes scale factor of 1.3.		
26.2±3.1±0.9	¹ ABAZOV 07T D0 $p\bar{p}$ at 1.96 TeV		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
14.9 ^{+2.2} _{-2.5} ^{+1.2} _{-1.4}	¹ AALTONEN 09D CDF Repl. by AALTONEN 14I		
1 Observed in $B_2^{*0} \rightarrow B^+ \pi^-$ and $B_2^{*0} \rightarrow B^+ \pi^-$.			
	$m_{B_2^{*0}} - m_{B^+}$		

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
460.2 ± 0.6 OUR FIT	Error includes scale factor of 1.4.			
459.9 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.8.			
460.18±0.37±0.33	17K ¹ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV			
457.5 ± 1.2 ^{+0.8} _{-0.9}	² AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV			
1 AAIJ 15AB reports $[m_{B_2^{*0}} - m_{B^+}] - m_{\pi^-} = 320.6 \pm 0.4 \pm 0.3$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event.				
2 AALTONEN 14I reports $m_{B_2^*(5747)^0} - m_{B^+} - m_{\pi^-} = 317.9 \pm 1.2 \pm 0.8$ MeV which we adjusted by the π^- mass.				

$B_2^*(5747)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
24.2±1.7 OUR AVERAGE				
24.5±1.0± 1.5	17K AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV			
22 ⁺³ ₋₂ ⁺⁴ ₋₅	AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22.7 ^{+3.8} _{-3.2} ^{+3.2} _{-10.2}	AALTONEN 09D CDF Repl. by AALTONEN 14I			

$B_2^*(5747)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 B^+ \pi^-$	seen
$\Gamma_2 B^{*+} \pi^-$	seen

NODE=M219

REFID=56628

REFID=56029

NODE=M184

NODE=M184

NODE=M184M

NODE=M184M

NODE=M184M

NODE=M184DM

NODE=M184DM

NODE=M184DM;LINKAGE=AB

NODE=M184DM2

NODE=M184DM2

NODE=M184DM2;LINKAGE=A

NODE=M184DM2;LINKAGE=AA

NODE=M184W

NODE=M184W

NODE=M184215;NODE=M184

DESIG=1

DESIG=2

$B_2^*(5747)^0$ BRANCHING RATIOS **$\Gamma(B^+\pi^-)/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	17K	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV	
seen		AALTONEN	09D CDF	$p\bar{p}$ at 1.96 TeV	
seen		ABAZOV	07T D0	$p\bar{p}$ at 1.96 TeV	

 Γ_1/Γ

NODE=M184220

NODE=M184R01

NODE=M184R01

 $\Gamma(B^{*+}\pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	17K	AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV	
seen		AALTONEN	09D CDF	$p\bar{p}$ at 1.96 TeV	
seen		ABAZOV	07T D0	$p\bar{p}$ at 1.96 TeV	

 Γ_2/Γ

NODE=M184R02

NODE=M184R02

 $\Gamma(B^{*+}\pi^-)/\Gamma(B^+\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.82 ± 0.28 OUR AVERAGE					

NODE=M184R03

NODE=M184R03

0.71 ± 0.14 ± 0.30

1.10 ± 0.42 ± 0.31

17K AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV1 ABAZOV 07T D0 $p\bar{p}$ at 1.96 TeV1 Converted from measured ratio of $R = B(B_2^{*0} \rightarrow B^{*+}\pi^-) / B(B_2^{*0} \rightarrow B^{(*)+}\pi^-)$
= 0.475 ± 0.095 ± 0.069. **$B_2^*(5747)^0$ REFERENCES**

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D	PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T	PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)

 $B_J(5840)^+$ $I(J^P) = \frac{1}{2}(??)$
 I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

 $B_J(5840)^+$ MASSOUR FIT uses m_{B^0} and $m_{B_J(5840)^+} - m_{B^0}$ to determine $m_{B_J(5840)^+}$.

VALUE (MeV)	DOCUMENT ID
5851 ± 19 OUR FIT	

NODE=M184

REFID=56628

REFID=56029

REFID=52700

REFID=52014

NODE=M224

NODE=M224

NODE=M224M

NODE=M224M

NODE=M224M

NODE=M224DM

NODE=M224DM

OCCUR=2

NODE=M224DM;LINKAGE=A

NODE=M224DM;LINKAGE=B

NODE=M224DM2

NODE=M224DM2

 $m_{B_J(5840)^+} - m_{B^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
571 ± 19 OUR FIT				

571 ± 13 ± 147k ¹ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

595 ± 26 ± 14

7k ² AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV1 AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.2 AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$m_{B_J(5840)^+} = m_{B^0}$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
565 ± 15 ± 14	7k ¹ AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

565 ± 15 ± 14

7k ¹ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$

MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M224DM2;LINKAGE=A

$B_J(5840)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
224±24±80	7k	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
215±27±80	7k	² AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
229±27±80	7k	³ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference. ² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference. ³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.				

NODE=M224W

NODE=M224W

OCCUR=2
OCCUR=3

NODE=M224W;LINKAGE=A

NODE=M224W;LINKAGE=B

NODE=M224W;LINKAGE=C

$B_J(5840)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B^{*0} \pi^+$	seen
$\Gamma_2 \quad B^0 \pi^+$	possibly seen

NODE=M224215;NODE=M224

DESIG=1
DESIG=2

NODE=M224220

NODE=M224R01
NODE=M224R01

$B_J(5840)^+$ BRANCHING RATIOS

$\Gamma(B^{*0}\pi^+)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	7k	AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
$\Gamma(B^0\pi^+)/\Gamma_{\text{total}}$				
VALUE	EVTS	DOCUMENT ID	TECN	Γ_2/Γ
possibly seen	7k	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV
¹ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.				

NODE=M224220

NODE=M224R02
NODE=M224R02

NODE=M224R02;LINKAGE=A

NODE=M224

REFID=56628

$B_J(5840)^+$ REFERENCES

AAIJ 15AB JHEP 1504 024 R. Aaij *et al.* (LHCb Collab.)

$B_J(5840)^0$
 $I(J^P) = \frac{1}{2}(?)$
I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

 $B_J(5840)^0$ MASSOUR FIT uses m_{B^+} and $m_{B_J(5840)^0} - m_{B^+}$ to determine $m_{B_J(5840)^0}$.

VALUE (MeV)	DOCUMENT ID
5863±9 OUR FIT	

 $m_{B_J(5840)^0} - m_{B^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
584± 9 OUR FIT				
584± 5±7	12k	¹ AAIJ	15AB LHCb	pp at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
610±22±7	12k	² AAIJ	15AB LHCb	pp at 7, 8 TeV
1 AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.				
2 AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.				

 $m_{B_J(5840)^0} - m_{B^{*+}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
584±5±7	12k	¹ AAIJ	15AB LHCb	pp at 7, 8 TeV
1 AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.				

 $B_J(5840)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
127±17±34	12k	¹ AAIJ	15AB LHCb	pp at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
107±20±34	12k	² AAIJ	15AB LHCb	pp at 7, 8 TeV
119±17±34	12k	³ AAIJ	15AB LHCb	pp at 7, 8 TeV
1 Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.				
2 Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.				
3 Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.				

 $B_J(5840)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 B^{*+} \pi^-$	seen
$\Gamma_2 B^+ \pi^-$	possibly seen

NODE=M225

NODE=M225

NODE=M225M

NODE=M225M

NODE=M225M

NODE=M225DM

NODE=M225DM

OCCUR=2

NODE=M225DM;LINKAGE=A

NODE=M225DM;LINKAGE=B

NODE=M225DM2

NODE=M225DM2

NODE=M225DM2;LINKAGE=A

NODE=M225W

NODE=M225W

OCCUR=2

OCCUR=3

NODE=M225W;LINKAGE=A

NODE=M225W;LINKAGE=B

NODE=M225W;LINKAGE=C

NODE=M225215;NODE=M225

DESIG=1

DESIG=2

$B_J(5840)^0$ BRANCHING RATIOS

$\Gamma(B^{*+}\pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	12k AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV
$\Gamma(B^+\pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
possibly seen	1 AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV
¹ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.	

$B_J(5840)^0$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------

$B_J(5970)^+$

$$I(J^P) = \frac{1}{2}(?)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_J(5970)^+$ MASS

OUR FIT uses m_{B^0} and $m_{B_J(5970)^+} - m_{B^0}$ to determine $m_{B_J(5970)^+}$.

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
5964±5 OUR FIT	

$$m_{B_J(5970)^+} = m_{B^0}$$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
685 ± 5 OUR FIT				
685 ± 5 OUR AVERAGE				
685.3±4.1± 2.5	2K	¹ AAIJ	15AB LHCb	$p\bar{p}$ at 7, 8 TeV

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

686.8±4.5± 2.5 2K ³ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV

¹AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 545.8 \pm 4.1 \pm 2.5$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

²AALTOMEN 14I reports $m_{B_J(5970)^+} - m_{B^0} - m_{\pi^+} = 541 \pm 5 \pm 12$ MeV which we adjusted by the π^+ mass.

³AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 547 \pm 5 \pm 3$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$$m_{B_J(5970)^+} - m_{B^{*0}}$$

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------------	-------------	--------------------	-------------	----------------

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

686.0±4.0±2.5 2k ¹ AAIJ 15AB LHCb $p\bar{p}$ at 7, 8 TeV

¹AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 547 \pm 4 \pm 3$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5970)^+$ WIDTH

NODE=M225220

NODE=M225R01

NODE=M225R01

NODE=M225R02

NODE=M225R02

NODE=M225R02;LINKAGE=A

NODE=M225

REFID=56628

NODE=M220

NODE=M220

NODE=M220M

NODE=M220M

NODE=M220M

NODE=M220DM

NODE=M220DM

OCCUR=2

NODE=M220DM;LINKAGE=B

NODE=M220DM;LINKAGE=A

NODE=M220DM;LINKAGE=C

NODE=M220DM2

NODE=M220DM2

NODE=M220DM2;LINKAGE=A

NODE=M220W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M220W
62±20 OUR AVERAGE						
63±15±17	2K	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV		
60 ⁺³⁰ ₋₂₀ ±40	1.4k	AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
61±14±17	2K	² AAIJ	15AB LHCb	$p p$ at 7, 8 TeV		OCCUR=2
61±15±17	2K	³ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV		OCCUR=3
1 Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.						
2 Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.						
3 Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.						

$B_J(5970)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 B^0 \pi^+$	possibly seen	
$\Gamma_2 B^{*0} \pi^+$	seen	

$B_J(5970)^+$ BRANCHING RATIOS

$\Gamma(B^0 \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ
possibly seen	2K ¹ AAIJ 15AB LHCb $p p$ at 7, 8 TeV
possibly seen	1.4k AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV
1 A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.	

$\Gamma(B^{*0} \pi^+)/\Gamma_{\text{total}}$	Γ_2/Γ
seen	2k AAIJ 15AB LHCb $p p$ at 7, 8 TeV
seen	1.4k AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

$B_J(5970)^+$ REFERENCES

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)

$B_J(5970)^0$

$I(J^P) = \frac{1}{2}(?)$
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_J(5970)^0$ MASS

OUR FIT uses m_{B^+} and $m_{B_J(5970)^0} - m_{B^+}$ to determine $m_{B_J(5970)^0}$.

VALUE (MeV)	DOCUMENT ID			
5971±5 OUR FIT				

$m_{B_J(5970)^0} - m_{B^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M221DM
691 ±5 OUR FIT						
691 ±5 OUR AVERAGE						
689.9±2.9± 5.1	10K	¹ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV		
698 ±5 ±12	2.6k	² AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
714.3±6.4± 5.1	10K	³ AAIJ	15AB LHCb	$p p$ at 7, 8 TeV		OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 550.4 \pm 2.9 \pm 5.1$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AALTONEN 14I reports $m_{B_J(5970)^0} - m_{B^+} - m_{\pi^-} = 558 \pm 5 \pm 12$ MeV which we adjusted by the π^- mass.

³ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 575 \pm 6 \pm 5$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$$m_{B_J(5970)^0} = m_{B^{*+}}$$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$691.6 \pm 3.7 \pm 5.1$ 10k ¹ AAIJ 15AB LHCb $p p$ at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 552 \pm 4 \pm 5$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5970)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

81±12 OUR AVERAGE

$82 \pm 8 \pm 9$ 10K ¹ AAIJ 15AB LHCb $p p$ at 7, 8 TeV

$70^{+30}_{-20} \pm 30$ 2.6k AALTONEN 14I CDF $p \bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$56 \pm 7 \pm 9$ 10K ² AAIJ 15AB LHCb $p p$ at 7, 8 TeV

$82 \pm 10 \pm 9$ 10K ³ AAIJ 15AB LHCb $p p$ at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

$B_J(5970)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B^+ \pi^-$	possibly seen
$\Gamma_2 \quad B^{*+} \pi^-$	seen

$B_J(5970)^0$ BRANCHING RATIOS

$\Gamma(B^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

possibly seen 10K ¹ AAIJ 15AB LHCb $p p$ at 7, 8 TeV

possibly seen 2.6k AALTONEN 14I CDF $p \bar{p}$ at 1.96 TeV

¹ A $B \pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^* \pi$ is allowed.

Γ_1/Γ

NODE=M221DM;LINKAGE=B

NODE=M221DM;LINKAGE=A

NODE=M221DM;LINKAGE=C

NODE=M221DM2

NODE=M221DM2

NODE=M221DM2;LINKAGE=A

NODE=M221W

NODE=M221W

OCCUR=2

OCCUR=3

NODE=M221W;LINKAGE=A

NODE=M221W;LINKAGE=B

NODE=M221W;LINKAGE=C

NODE=M221215;NODE=M221

DESIG=1

DESIG=2

NODE=M221220

NODE=M221R01

NODE=M221R01

OCCUR=2

NODE=M221R01;LINKAGE=A

NODE=M221R02

NODE=M221R02

OCCUR=2

NODE=M221

REFID=56628

REFID=56029

$\Gamma(B^{*+} \pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 10K AAIJ 15AB LHCb $p p$ at 7, 8 TeV

seen 2.6k AALTONEN 14I CDF $p \bar{p}$ at 1.96 TeV

Γ_2/Γ

$B_J(5970)^0$ REFERENCES

AAIJ 15AB JHEP 1504 024
AALTONEN 14I PR D90 012013

R. Aaij *et al.*
T. Aaltonen *et al.*

(LHCb Collab.)
(CDF Collab.)

BOTTOM, STRANGE MESONS ($B = \pm 1$, $S = \mp 1$)

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \text{ similarly for } B_s^{*+}'s$$

$X(5568)^\pm$

$I(J^P) = ?(?)$

OMITTED FROM SUMMARY TABLE

Seen as a peak in the $B_s\pi^\pm$ mass spectrum with a significance of more than 3σ by ABAZOV 16E and ABAZOV 18A in inclusive $p\bar{p}$ collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

$X(5568)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5566.9 +3.2 -3.1 +0.6 -1.2	278	¹ ABAZOV	18A D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5567.8 ± 2.9 $+0.9$ -1.9	133	² ABAZOV	16E D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$
1 From the combined analysis of $B_s^0 \rightarrow J/\psi\phi$ and $B_s^0 \rightarrow D_s^\pm \mu^\mp X$ decays.				
2 Assumes $X(5568)^\pm \rightarrow B_s\pi^\pm$ decay. If $X(5568)^\pm \rightarrow B_s^*\pi^\pm$ decay is assumed, the mass shifts upward by 49 MeV.				

$X(5568)^\pm$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
18.6 +7.9 -6.1 +3.5 -3.8	278	¹ ABAZOV	18A D0	$p\bar{p} \rightarrow B_s\pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
21.9 ± 6.4 $+5.0$ -2.5	133	ABAZOV	16E D0	$p\bar{p} \rightarrow B_s\pi^\pm X$
1 From the combined analysis of $B_s^0 \rightarrow J/\psi\phi$ and $B_s^0 \rightarrow D_s^\pm \mu^\mp X$ decays.				

$X(5568)^\pm$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 B_s\pi^\pm$	seen

$X(5568)^\pm$ BRANCHING RATIOS

$\Gamma(B_s\pi^\pm)/\Gamma_{\text{total}}$			Γ_1/Γ
seen	145	¹ ABAZOV	18A D0
seen	133	² ABAZOV	16E D0
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen		³ AABOUD	18L ATLS
not seen		⁴ AALTONEN	18A CDF
not seen		⁵ SIRUNYAN	18J CMS
not seen		⁶ AAIJ	16AI LHCb

NODE=MXXX046

NODE=MXXX046

NODE=M232

NODE=M232

NODE=M232M

NODE=M232M

NODE=M232M;LINKAGE=B

NODE=M232M;LINKAGE=A

NODE=M232W

NODE=M232W

NODE=M232W;LINKAGE=B

NODE=M232215;NODE=M232

DESIG=1

NODE=M232220

NODE=M232R01
NODE=M232R01

OCCUR=2

¹With B_s mesons reconstructed in decays to $D_s^\pm \mu^\mp X$.

²Seen in $p\bar{p}$ collisions at 1.96 TeV at a rate of $(8.6 \pm 1.9 \pm 1.4)\%$ relative to inclusive B_s production in the kinematic region $10 < p_T(B_s) < 30$ GeV/c, with B_s mesons reconstructed in decays to $J/\psi \phi$. An alternative possibility, $X(5568)^\pm \rightarrow B_s^* \pi^\pm$ with a missing γ , could not be ruled out.

³Not seen in 24.4 fb^{-1} of $p\bar{p}$ collision data at $\sqrt{s} = 7$ and 8 TeV with B_s mesons reconstructed in decays to $J/\psi \phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 1.5% at $p_T(B_s) > 10$ GeV/c and less than 1.6% at $p_T(B_s) > 15$ GeV/c at 95% CL.

⁴Not seen in 9.6 fb^{-1} of $p\bar{p}$ collision data at $\sqrt{s} = 1.96$ TeV with B_s mesons reconstructed in decays to $J/\psi \phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 6.7% at 95% CL.

⁵Not seen in 19.7 fb^{-1} of $p\bar{p}$ collisions data at $\sqrt{s} = 8$ TeV with B_s mesons reconstructed in decays to $J/\psi \phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 1.1% at $p_T(B_s) > 10$ GeV/c and less than 1.0% at $p_T(B_s) > 15$ GeV/c at 95% CL.

⁶Not seen in 3 fb^{-1} of $p\bar{p}$ collision data at $\sqrt{s} = 7$ and 8 TeV in a scan over the $X(5568)$ mass and width, with B_s mesons reconstructed in decays to $D_s^- \pi^+$ or $J/\psi \phi$. An upper limit on the production rate times branching fraction for $X(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 2.1% at $p_T(B_s) > 10$ GeV/c at 90% CL.

$X(5568)^\pm$ REFERENCES

AABOUD	18L	PRL 120 202007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AALTONEN	18A	PRL 120 202006	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	18A	PR D97 092004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
SIRUNYAN	18J	PRL 120 202005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	16A1	PRL 117 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	16E	PRL 117 022003	V.M. Abazov <i>et al.</i>	(D0 Collab.)

$B_{s1}(5830)^0$

$I(J^P) = 0(1^+)$
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_{s1}(5830)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

5828.70 ± 0.20 OUR FIT

5828.65 ± 0.24 OUR AVERAGE

5828.78 ± 0.09 ± 0.29 SIRUNYAN 18DF CMS $p\bar{p}$ at 8 TeV

5828.40 ± 0.04 ± 0.41 ¹AAIJ 130 LHCb $p\bar{p}$ at 7 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5829.4 ± 0.7 ²AALTOMEN 08K CDF Repl. by AALTOMEN 14I

¹ Uses $B_{s1}(5830)^0 \rightarrow B^{*+} K^-$ decay.

² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.

$$m_{B_{s1}^0} - m_{B^{*+}}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

504.00 ± 0.17 OUR FIT

504.03 ± 0.12 ± 0.15

¹ AALTOMEN 14I CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

504.41 ± 0.21 ± 0.14 ²AALTOMEN 08K CDF Repl. by AALTOMEN 14I

¹ AALTOMEN 14I reports $m_{B_{s1}(5830)^0} - m_{B^{*+}} - m_{K^-} = 10.35 \pm 0.12 \pm 0.15$ MeV which we adjusted by the K^- mass.

² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.

$B_{s1}(5830)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.5 ± 0.3 ± 0.3

NODE=M232R01;LINKAGE=F

NODE=M232R01;LINKAGE=A

NODE=M232R01;LINKAGE=E

NODE=M232R01;LINKAGE=D

NODE=M232R01;LINKAGE=C

NODE=M232R01;LINKAGE=B

NODE=M232

REFID=58829

REFID=58828

REFID=58937

REFID=58827

REFID=57549

REFID=57453

NODE=M187

NODE=M187

NODE=M187M

NODE=M187M

NODE=M187M;LINKAGE=AI

NODE=M187M;LINKAGE=AA

NODE=M187DM

NODE=M187DM

NODE=M187DM;LINKAGE=AL

NODE=M187DM;LINKAGE=AA

NODE=M187W

NODE=M187W

$B_{s1}(5830)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B^{*+} K^-$	seen
$\Gamma_2 \quad B^{*0} K_S^0$	

 $B_{s1}(5830)^0$ BRANCHING RATIOS

$\Gamma(B^{*+} K^-)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
seen	AALTONEN 08K CDF $p\bar{p}$ at 1.96 TeV
$\Gamma(B^{*0} K_S^0)/\Gamma(B^{*+} K^-)$	Γ_2/Γ_1
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.49 \pm 0.12 \pm 0.07$	¹ SIRUNYAN 18DF CMS $p\bar{p}$ at 8 TeV

¹ With the branching fractions $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$ and $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$.

 $B_{s1}(5830)^0$ REFERENCES

SIRUNYAN 18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AALTONEN 14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ 13O PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN 08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)

 $B_{s2}^*(5840)^0$

$$I(J^P) = 0(2^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

 $B_{s2}^*(5840)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

 5839.86 ± 0.12 OUR FIT[5839.85 ± 0.12 MeV OUR 2019 FIT] **5839.92 ± 0.14 OUR AVERAGE**

$5839.86 \pm 0.09 \pm 0.17$	SIRUNYAN 18DF CMS	$p\bar{p}$ at 8 TeV
$5839.99 \pm 0.05 \pm 0.20$	AAIJ 13O LHCb	$p\bar{p}$ at 7 TeV
$5839.6 \pm 1.1 \pm 0.7$	¹ ABAZOV 08E D0	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5839.7 ± 0.7 ²AALTONEN 08K CDF Repl. by AALTONEN 14I

1 Observed in $B_{s2}^{*0} \rightarrow B^+ K^-$. Measured production rate of B_{s2}^{*0} relative to B^+ to be $(1.15 \pm 0.23 \pm 0.13)\%$.

2 Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+, \bar{D}^0 \rightarrow K^+ \pi^-$.

$$m_{B_{s2}^{*0}} - m_{B_{s1}^0}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.5 ± 0.6 ¹AALTONEN 08K CDF Repl. by AALTONEN 14I

1 Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+, \bar{D}^0 \rightarrow K^+ \pi^-$.

$$m_{B_{s2}^{*0}} - m_{B^+}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

 560.52 ± 0.14 OUR FIT[560.53 ± 0.14 MeV OUR 2019 FIT] **$560.41 \pm 0.13 \pm 0.14$**

¹AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

NODE=M187215;NODE=M187

DESIG=1

DESIG=2

NODE=M187220

NODE=M187R01

NODE=M187R01

NODE=M187R00

NODE=M187R00

NODE=M187R00;LINKAGE=A

NODE=M187

REFID=59328

REFID=56029

REFID=54968

REFID=52235

NODE=M186

NODE=M186

NODE=M186M

NODE=M186M

NEW

NODE=M186M;LINKAGE=AB

NODE=M186M;LINKAGE=AA

NODE=M186DM

NODE=M186DM

NODE=M186DM;LINKAGE=AA

NODE=M186DM2

NODE=M186DM2

NEW

¹ AALTONEN 14I reports $m_{B_{s2}(5840)^0} - m_{B^+} - m_{K^-} = 66.73 \pm 0.13 \pm 0.14$ MeV which we adjusted by the K^- mass.

NODE=M186DM2;LINKAGE=AL

$B_{s2}^*(5840)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1.49±0.27 OUR AVERAGE			
1.52±0.34±0.30	SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV
1.4 ± 0.4 ± 0.2	AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
1.56±0.13±0.47	¹ AAIJ	130 LHCb	$p\bar{p}$ at 7 TeV
¹ Uses $B_{s2}^*(5840)^0 \rightarrow B^{*+} K^-$ decays.			

NODE=M186W

NODE=M186W

NODE=M186W;LINKAGE=AI

NODE=M186215;NODE=M186

$B_{s2}^*(5840)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	DEFINITION	DESIGNATION
$\Gamma_1 B^+ K^-$	DEFINED AS 1		
$\Gamma_2 B^{*+} K^-$	0.093±0.018		DESIG=2
$\Gamma_3 B^0 K_S^0$	0.43 ± 0.11		DESIG=4
$\Gamma_4 B^{*0} K_S^0$	0.04 ± 0.04		DESIG=3

NODE=M186220

$B_{s2}^*(5840)^0$ BRANCHING RATIOS

$\Gamma(B^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	AALTONEN	08K CDF	$p\bar{p}$ at 1.96 TeV	NODE=M186R01
seen	¹ ABAZOV	08E D0	$p\bar{p}$ at 1.96 TeV	NODE=M186R01

NODE=M186R01

NODE=M186R01

NODE=M186R01;LINKAGE=AB

$\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.093±0.013±0.012	AAIJ	130 LHCb	$p\bar{p}$ at 7 TeV	NODE=M186R02 NODE=M186R02

NODE=M186R02

NODE=M186R02

$\Gamma(B^{*0} K_S^0)/\Gamma(B^0 K_S^0)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3
0.093±0.086±0.014	¹ SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV	NODE=M186R00 NODE=M186R00

NODE=M186R00

NODE=M186R00

NODE=M186R00;LINKAGE=A

$\Gamma(B^0 K_S^0)/\Gamma(B^+ K^-)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.432±0.077±0.078	¹ SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV	NODE=M186R03 NODE=M186R03

NODE=M186R03

NODE=M186R03

NODE=M186R03;LINKAGE=A

$\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.081±0.021±0.015	¹ SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV	NODE=M186R04 NODE=M186R04

NODE=M186R04

NODE=M186R04

NODE=M186R04;LINKAGE=A

$B_{s2}^*(5840)^0$ REFERENCES

SIRUNYAN	18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	130 PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	08E PRL 100 082002	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=59328

REFID=56029

REFID=54968

REFID=52235

REFID=52232

NODE=M186

NODE=M153

 $B_{sJ}^*(5850)$
 $I(J^P) = ?(?)$
I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Signal can be interpreted as coming from $\bar{b}s$ states. Needs confirmation.

NODE=M153

 $B_{sJ}^*(5850)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5853±15	141	AKERS	95E	OPAL $E_{cm}^{ee} = 88-94$ GeV

NODE=M153M

NODE=M153M

 $B_{sJ}^*(5850)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47±22	141	AKERS	95E	OPAL $E_{cm}^{ee} = 88-94$ GeV

NODE=M153W

NODE=M153W

 $B_{sJ}^*(5850)$ REFERENCESAKERS 95E ZPHY C66 19 R. Akers *et al.* (OPAL Collab.)

NODE=M153

REFID=44182

**BOTTOM, CHARMED MESONS
($B = C = \pm 1$)**

$$B_c^+ = c\bar{b}, B_c^- = \bar{c}b, \text{ similarly for } B_c^* \text{'s}$$

 $B_c(2S)^\pm$

$$I(J^P) = 0(0^-)$$

OMITTED FROM SUMMARY TABLE

Quantum numbers neither measured nor confirmed.

NODE=MXXX049

NODE=MXXX049

NODE=M217

 $B_c(2S)^\pm$

$$I(J^P) = 0(0^-)$$

NODE=M217

NODE=M217M

NODE=M217M

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
6871.6±1.1 OUR AVERAGE				

[6871.0 ± 1.6 MeV OUR 2019 AVERAGE]

6872.1±1.3±0.8 24 1,2 AAIJ 19Y LHCb pp at 7, 8, 13 TeV6871.0±1.4±0.8 51 3,4 SIRUNYAN 19M CMS pp at 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen 5 AAIJ 18AL LHCb pp at 8 TeV6842 ± 4 ± 5 57 6,7 AAD 14AQ ATLAS pp at 7, 8 TeV

¹AAIJ 19Y observed $B_c(2S)^+$ in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 2.2 (3.2) global (local) standard deviations significance.

²AAIJ 19Y reports mass difference measurement of $M(B_c(2S)^+) - M(B_c^+) = 597.2 \pm 1.3 \pm 0.1$ MeV. We have adjusted this measurement with our best value of $M(B_c^+) = 6274.9 \pm 0.8$ MeV. The first uncertainty of the $M(B_c(2S)^+)$ value is a total of uncertainties reported by the experiment and the second one comes from our best value of $M(B_c^+)$.

³SIRUNYAN 19M observed $B_c(2S)^+$ in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 6.5 standard deviations significance.

⁴SIRUNYAN 19M reports mass difference measurement of $M(B_c(2S)^+) - M(B_c^+) = 596.1 \pm 1.2 \pm 0.8$ MeV. We have adjusted this measurement with our best value of $M(B_c^+) = 6274.9 \pm 0.8$ MeV. The first uncertainty of the $M(B_c(2S)^+)$ value is a total of uncertainties reported by the experiment and the second one comes from our best value of $M(B_c^+)$.

NODE=M217M;LINKAGE=E

NODE=M217M;LINKAGE=F

NODE=M217M;LINKAGE=B

NODE=M217M;LINKAGE=D

⁵ AAIJ 18AL reports an upper limit on the ratio of production cross sections for $[\sigma(B_c(2S)^+)/\sigma(B_c^+)] \cdot B(B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-) < 0.04-0.09$ at 95% CL for the mass value reported by AAD 14AQ.

⁶ Observed in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 5.2 standard deviations significance.

⁷ Might be the $B_c^*(2S)$.

NODE=M217M;LINKAGE=A

NODE=M217M;LINKAGE=AA

NODE=M217M;LINKAGE=C

NODE=M217215;NODE=M217

DESIG=1

NODE=M217225

NODE=M217R01
NODE=M217R01

NODE=M217R01;LINKAGE=AA
NODE=M217R01;LINKAGE=A

NODE=M217

REFID=59795
REFID=59639
REFID=59246
REFID=56117

$B_c(2S)^{\pm}$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad B_c^+ \pi^+ \pi^-$	seen

$B_c(2S)^{\pm}$ BRANCHING RATIOS

$\Gamma(B_c^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	EVTS DOCUMENT ID TECN COMMENT
seen	57 1 AAD 14AQ ATLS $p p$ at 7, 8 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •	
not seen	2 AAIJ 18AL LHCb $p p$ at 8 TeV
1 Observed with 5.2 standard deviations significance.	
2 AAIJ 18AL reports an upper limit on the ratio of production cross sections for $[\sigma(B_c(2S)^+)/\sigma(B_c^+)] \cdot B(B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-) < 0.04-0.09$ at 95% CL for the mass value reported by AAD 14AQ.	

$B_c(2S)^{\pm}$ REFERENCES

AAIJ	19Y	PRL 122 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	19M	PRL 122 132001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	18AL	JHEP 1801 138	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	14AQ	PRL 113 212004	G. Aad <i>et al.</i>	(ATLAS Collab.)

See the related review(s):

[Spectroscopy of Mesons Containing Two Heavy Quarks](#)

$c\bar{c}$ MESONS (including possibly non- $q\bar{q}$ states)

NODE=MXXX025

See the related review(s):

[Charmonium System](#)

NODE=M026

 $\eta_c(1S)$ $I^G(J^{PC}) = 0^+(0^-+)$ $\eta_c(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2983.9 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below. [2983.9 ± 0.5 MeV OUR 2019 AVERAGE Scale factor = 1.3]		
2985.9 ± 0.7 ± 2.1	1705	ABLIKIM	19AV BES3	$J/\psi \rightarrow \gamma\omega\omega$
2984.6 ± 0.7 ± 2.2	2673	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
2986.7 ± 0.5 ± 0.9	11K	¹ AAIJ	17AD LHCb	$p\bar{p} \rightarrow B^+X \rightarrow p\bar{p}K^+X$
2982.8 ± 1.0 ± 0.5	6.4k	² AAIJ	17BB LHCb	$p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
2982.2 ± 1.5 ± 0.1	2.0k	³ AAIJ	15BI LHCb	$p\bar{p} \rightarrow \eta_c(1S)X$
2983.5 ± 1.4 ± 1.6		⁴ ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
2979.8 ± 0.8 ± 3.5	4.5k	^{5,6} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
2984.1 ± 1.1 ± 2.1	900	^{5,6,7} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
2984.3 ± 0.6 ± 0.6		^{8,9} ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
2984.49 ± 1.16 ± 0.52	832	⁵ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons
2982.7 ± 1.8 ± 2.2	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
2984.5 ± 0.8 ± 3.1	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
2985.4 ± 1.5 ± 0.5	920	⁹ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0K^\pm\pi^\mp)$
2982.2 ± 0.4 ± 1.6	14k	¹⁰ LEES	10 BABR	$10.6 \frac{e^+e^-}{e^+e^-K_S^0} \rightarrow K^\pm\pi^\mp$
2985.8 ± 1.5 ± 3.1	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K^{(*)} \rightarrow K\bar{K}\pi K^{(*)}$
2986.1 ± 1.0 ± 2.5	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow \text{hadrons}$
2970 ± 5 ± 6	501	¹¹ ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
2971 ± 3 ± 2	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
2974 ± 7 ± 2	20	WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
2981.8 ± 1.3 ± 1.5	592	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0K^\pm\pi^\mp$
2984.1 ± 2.1 ± 1.0	190	¹² AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2982.5 ± 0.4 ± 1.4	12k	¹³ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
2982.2 ± 0.6		¹⁴ MITCHELL	09 CLEO	$e^+e^- \rightarrow \gamma X$
2982 ± 5	270	¹⁵ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
2982.5 ± 1.1 ± 0.9	2.5k	¹⁶ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
2977.5 ± 1.0 ± 1.2		^{14,17} BAI	03 BES	$J/\psi \rightarrow \gamma\eta_c$
2979.6 ± 2.3 ± 1.6	180	¹⁸ FANG	03 BELL	$B \rightarrow \eta_c K$
2976.3 ± 2.3 ± 1.2		^{14,19} BAI	00F BES	$J/\psi, \psi(2S) \rightarrow \gamma\eta_c$
2976.6 ± 2.9 ± 1.3	140	^{14,20} BAI	00F BES	$J/\psi \rightarrow \gamma\eta_c$
2980.4 ± 2.3 ± 0.6		²¹ BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$
2975.8 ± 3.9 ± 1.2		²⁰ BAI	99B BES	Sup. by BAI 00F
2999 ± 8	25	ABREU	980 DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
2988.3 ± 3.3		ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
2974.4 ± 1.9		^{14,22} BISELLLO	91 DM2	$J/\psi \rightarrow \eta_c\gamma$
2969 ± 4 ± 4	80	¹⁴ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K^+K^-$
2956 ± 12 ± 12		¹⁴ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K^0_K L$
2982.6 ± 2.7	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
2980.2 ± 1.6		^{14,22} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$
2984 ± 2.3 ± 4.0		¹⁴ GAISER	86 CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
2976 ± 8		^{14,23} BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$
2982 ± 8	18	²⁴ HIMEL	80B MRK2	e^+e^-
2980 ± 9		²⁴ PARTRIDGE	80B CBAL	e^+e^-

NODE=M026M

NODE=M026M
NEW

OCCUR=2


OCCUR=2

OCCUR=2

OCCUR=3

- 1 AAIJ 17AD report $m_{J/\psi} - m_{\eta_c(1S)} = 110.2 \pm 0.5 \pm 0.9$ MeV. We use the current value $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to obtain the quoted mass.
- 2 From a fit of the $\phi\phi$ invariant mass with the mass and width of $\eta_c(1S)$ as free parameters.
- 3 AAIJ 15BI reports $m_{J/\psi} - m_{\eta_c(1S)} = 114.7 \pm 1.5 \pm 0.1$ MeV from a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. We have used current value of $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to arrive at the quoted $m_{\eta_c(1S)}$ result.
- 4 Taking into account an asymmetric photon lineshape.
- 5 With floating width.
- 6 Ignoring possible interference with the non-resonant 0^- amplitude.
- 7 Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.
- 8 From a simultaneous fit to six decay modes of the η_c .
- 9 Accounts for interference with non-resonant continuum.
- 10 Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.
- 11 From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.
- 12 Using mass of $\psi(2S) = 3686.00$ MeV.
- 13 Not independent from the measurements reported by LEES 10.
- 14 MITCHELL 09 observes a significant asymmetry in the lineshapes of $\psi(2S) \rightarrow \gamma\eta_c$ and $J/\psi \rightarrow \gamma\eta_c$ transitions. If ignored, this asymmetry could lead to significant bias whenever the mass and width are measured in $\psi(2S)$ or J/ψ radiative decays.
- 15 From the fit of the kaon momentum spectrum. Systematic errors not evaluated.
- 16 Superseded by LEES 10.
- 17 From a simultaneous fit of five decay modes of the η_c .
- 18 Superseded by VINOKUROVA 11.
- 19 Weighted average of the $\psi(2S)$ and $J/\psi(1S)$ samples. Using an η_c width of 13.2 MeV.
- 20 Average of several decay modes. Using an η_c width of 13.2 MeV.
- 21 Superseded by ASNER 04.
- 22 Average of several decay modes.
- 23 $\eta_c \rightarrow \phi\phi$.
- 24 Mass adjusted by us to correspond to $J/\psi(1S)$ mass = 3097 MeV.

WEIGHTED AVERAGE
2983.9±0.5 (Error scaled by 1.3)



$\eta_c(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
32.0± 0.7 OUR FIT				
[31.9 ± 0.7 MeV OUR 2019 FIT]				
32.1± 0.8 OUR AVERAGE	Error includes scale factor of 1.1. [32.1 ± 0.8 MeV OUR 2019 AVERAGE Scale factor = 1.1]			
33.8± 1.6±4.1	1705	ABLIKIM	19AV BES3	$J/\psi \rightarrow \gamma\omega\omega$

NODE=M026M;LINKAGE=F

NODE=M026M;LINKAGE=G

NODE=M026M;LINKAGE=D

NODE=M026M;LINKAGE=E

NODE=M026M;LINKAGE=AL

NODE=M026M;LINKAGE=LS

NODE=M026M;LINKAGE=EL

NODE=M026M;LINKAGE=BL

NODE=M026M;LINKAGE=VA

NODE=M026M;LINKAGE=LE

NODE=M026M;LINKAGE=EB

NODE=M026M;LINKAGE=BG

NODE=M026M;LINKAGE=DE

NODE=M026M;LINKAGE=MI

NODE=M026M;LINKAGE=AU

NODE=M026M;LINKAGE=UB

NODE=M026M;LINKAGE=AK

NODE=M026M;LINKAGE=FA

NODE=M026M;LINKAGE=KZ

NODE=M026M;LINKAGE=C1

NODE=M026M;LINKAGE=NN

NODE=M026M;LINKAGE=A

NODE=M026M;LINKAGE=B

NODE=M026M;LINKAGE=M

NODE=M026W

NODE=M026W

NEW

NEW

$30.8^{+2.3}_{-2.2} \pm 2.9$	2673	XU	18	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$	
$34.0 \pm 1.9 \pm 1.3$	11K	AAIJ	17AD	LHCb	$p p \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$	
$31.4 \pm 3.5 \pm 2.0$	6.4k	¹ AAIJ	17BB	LHCb	$p p \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-)X$	
$27.2 \pm 3.1^{+5.4}_{-2.6}$		² ANASHIN	14	KEDR	$J/\psi \rightarrow \gamma \eta_c$	
$25.2 \pm 2.6 \pm 2.4$	4.5k	^{3,4} LEES	14E	BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$	
$34.8 \pm 3.1 \pm 4.0$	900	^{3,4,5} LEES	14E	BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$	OCCUR=2
$32.0 \pm 1.2 \pm 1.0$		^{6,7} ABLIKIM	12F	BES3	$\psi(2S) \rightarrow \gamma \eta_c$	
$36.4 \pm 3.2 \pm 1.7$	832	³ ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons	
$37.8^{+5.8}_{-5.3} \pm 3.1$	486	ZHANG	12A	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$	
$36.2 \pm 2.8 \pm 3.0$	11k	DEL-AMO-SA..11M	BABR		$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
$35.1 \pm 3.1^{+1.0}_{-1.6}$	920	⁷ VINOKUROVA	11	BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$	
$31.7 \pm 1.2 \pm 0.8$	14k	⁸ LEES	10	BABR	$10.6 \frac{e^+ e^-}{e^+ e^- K_S^0 K^\pm \pi^\mp} \rightarrow$	
$36.3^{+3.7}_{-3.6} \pm 4.4$	0.9k	AUBERT	08AB	BABR	$B \rightarrow \eta_c(1S) K^{(*)} \rightarrow K \bar{K} \pi K^{(*)}$	
$28.1 \pm 3.2 \pm 2.2$	7.5k	UEHARA	08	BELL	$\gamma \gamma \rightarrow \eta_c \rightarrow$ hadrons	
$48^{+8}_{-7} \pm 5$	195	WU	06	BELL	$B^+ \rightarrow p \bar{p} K^+$	
$40 \pm 19 \pm 5$	20	WU	06	BELL	$B^+ \rightarrow \Lambda \bar{\Lambda} K^+$	OCCUR=2
$24.8 \pm 3.4 \pm 3.5$	592	ASNER	04	CLEO	$\gamma \gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$	
$20.4^{+7.7}_{-6.7} \pm 2.0$	190	AMBROGIANI	03	E835	$\bar{p} p \rightarrow \eta_c \rightarrow \gamma \gamma$	
$23.9^{+12.6}_{-7.1}$		ARMSTRONG	95F	E760	$\bar{p} p \rightarrow \gamma \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$32.1 \pm 1.1 \pm 1.3$	12k	⁹ DEL-AMO-SA..11M	BABR		$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$	OCCUR=2
$34.3 \pm 2.3 \pm 0.9$	2.5k	¹⁰ AUBERT	04D	BABR	$\gamma \gamma \rightarrow \eta_c(1S) \rightarrow K \bar{K} \pi$	
$17.0 \pm 3.7 \pm 7.4$		¹¹ BAI	03	BES	$J/\psi \rightarrow \gamma \eta_c$	
$29 \pm 8 \pm 6$	180	¹² FANG	03	BELL	$B \rightarrow \eta_c K$	
$11.0 \pm 8.1 \pm 4.1$		¹³ BAI	00F	BES	$J/\psi \rightarrow \gamma \eta_c$ and $\psi(2S) \rightarrow \gamma \eta_c$	
$27.0 \pm 5.8 \pm 1.4$		¹⁴ BRANDENB...	00B	CLE2	$\gamma \gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$	
$7.0^{+7.5}_{-7.0}$	12	BAGLIN	87B	SPEC	$\bar{p} p \rightarrow \gamma \gamma$	
$10.1^{+33.0}_{-8.2}$	23	¹⁵ BALTRUSAIT..86	MRK3		$J/\psi \rightarrow \gamma p \bar{p}$	
11.5 ± 4.5		GAISER	86	CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$	
< 40 90% CL	18	HIMEI	80B	MRK2	$e^+ e^-$	
< 20 90% CL		PARTRIDGE	80B	CBAL	$e^+ e^-$	

¹ From a fit of the $\phi\phi$ invariant mass with the mass and width of $\eta_c(1S)$ as free parameters.

² Taking into account an asymmetric photon lineshape.

³ With floating mass.

⁴ Ignoring possible interference with the non-resonant 0^- amplitude.

⁵ Using both, $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶ From a simultaneous fit to six decay modes of the η_c .

⁷ Accounts for interference with non-resonant continuum.

⁸ Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.

⁹ Not independent from the measurements reported by LEES 10.

¹⁰ Superseded by LEES 10.

¹¹ From a simultaneous fit of five decay modes of the η_c .

¹² Superseded by VINOKUROVA 11.

¹³ From a fit to the 4-prong invariant mass in $\psi(2S) \rightarrow \gamma \eta_c$ and $J/\psi(1S) \rightarrow \gamma \eta_c$ decays.

¹⁴ Superseded by ASNER 04.

¹⁵ Positive and negative errors correspond to 90% confidence level.

NODE=M026W;LINKAGE=B
 NODE=M026W;LINKAGE=A
 NODE=M026W;LINKAGE=AL
 NODE=M026W;LINKAGE=LS
 NODE=M026W;LINKAGE=EL
 NODE=M026W;LINKAGE=BL
 NODE=M026W;LINKAGE=VA
 NODE=M026W;LINKAGE=LE
 NODE=M026W;LINKAGE=DE
 NODE=M026W;LINKAGE=UB
 NODE=M026W;LINKAGE=AK
 NODE=M026W;LINKAGE=FA
 NODE=M026W;LINKAGE=KZ
 NODE=M026W;LINKAGE>NN
 NODE=M026W;LINKAGE=L

NODE=M026215;NODE=M026

$\eta_c(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
------	--------------------------------	------------------

Decays involving hadronic resonances

Γ_1	$\eta'(958)\pi\pi$	(4.1 ± 1.7) %	NODE=M026;CLUMP=A
Γ_2	$\rho\rho$	(1.8 ± 0.5) %	DESIG=24
Γ_3	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(2.0 ± 0.7) %	DESIG=19
Γ_4	$K^*(892) \bar{K}^*(892)$	(7.0 ± 1.3) $\times 10^{-3}$	DESIG=26
Γ_5	$K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-$	(1.1 ± 0.5) %	DESIG=18
Γ_6	$\phi K^+ K^-$	(2.9 ± 1.4) $\times 10^{-3}$	DESIG=57
Γ_7	$\phi\phi$	(1.77 ± 0.19) $\times 10^{-3}$	DESIG=28
Γ_8	$\phi 2(\pi^+ \pi^-)$	< 4 $\times 10^{-3}$	DESIG=17
Γ_9	$a_0(980)\pi$	< 2 %	90% DESIG=58
Γ_{10}	$a_2(1320)\pi$	< 2 %	90% DESIG=21
Γ_{11}	$K^*(892) \bar{K}^+ + \text{c.c.}$	< 1.28 %	90% DESIG=22
Γ_{12}	$f_2(1270)\eta$	< 1.1 %	90% DESIG=40
Γ_{13}	$\omega\omega$	< 3.1 $\times 10^{-3}$	DESIG=23
Γ_{14}	$\omega\phi$	< 2.5 $\times 10^{-4}$	90% DESIG=20
Γ_{15}	$f_2(1270) f_2(1270)$	(9.8 ± 2.5) $\times 10^{-3}$	DESIG=47
Γ_{16}	$f_2(1270) f'_2(1525)$	(9.7 ± 3.2) $\times 10^{-3}$	DESIG=46
Γ_{17}	$f_0(980)\eta$	seen	DESIG=59
Γ_{18}	$f_0(1500)\eta$	seen	DESIG=70
Γ_{19}	$f_0(2200)\eta$	seen	DESIG=71
Γ_{20}	$a_0(980)\pi$	seen	DESIG=72
Γ_{21}	$a_0(1320)\pi$	seen	DESIG=73
Γ_{22}	$a_0(1450)\pi$	seen	DESIG=74
Γ_{23}	$a_0(1950)\pi$	seen	DESIG=75
Γ_{24}	$K_0^*(1430) \bar{K}$	seen	DESIG=79
Γ_{25}	$K_2^*(1430) \bar{K}$	seen	DESIG=76
Γ_{26}	$K_0^*(1950) \bar{K}$	seen	DESIG=77
			DESIG=78

Decays into stable hadrons

Γ_{27}	$K \bar{K} \pi$	(7.3 ± 0.4) %	NODE=M026;CLUMP=B
Γ_{28}	$K \bar{K} \eta$	(1.36 ± 0.15) %	DESIG=14
Γ_{29}	$\eta \pi^+ \pi^-$	(1.7 ± 0.5) %	DESIG=25
Γ_{30}	$\eta 2(\pi^+ \pi^-)$	(4.4 ± 1.3) %	DESIG=16
Γ_{31}	$K^+ K^- \pi^+ \pi^-$	(6.9 ± 1.0) $\times 10^{-3}$	DESIG=61
Γ_{32}	$K^+ K^- \pi^+ \pi^- \pi^0$	(3.5 ± 0.6) %	DESIG=15
Γ_{33}	$K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}$	(5.6 ± 1.5) %	DESIG=60
Γ_{34}	$K^+ K^- 2(\pi^+ \pi^-)$	(7.5 ± 2.4) $\times 10^{-3}$	DESIG=62
Γ_{35}	$2(K^+ K^-)$	(1.46 ± 0.30) $\times 10^{-3}$	DESIG=55
Γ_{36}	$\pi^+ \pi^- \pi^0$	< 5 $\times 10^{-4}$	90% DESIG=27
Γ_{37}	$\pi^+ \pi^- \pi^0 \pi^0$	(4.7 ± 1.0) %	DESIG=81
Γ_{38}	$2(\pi^+ \pi^-)$	(9.7 ± 1.2) $\times 10^{-3}$	DESIG=63
Γ_{39}	$2(\pi^+ \pi^- \pi^0)$	(16.1 ± 2.0) %	DESIG=11
Γ_{40}	$3(\pi^+ \pi^-)$	(1.8 ± 0.4) %	DESIG=64
Γ_{41}	$p \bar{p}$	(1.45 ± 0.14) $\times 10^{-3}$	DESIG=56
Γ_{42}	$p \bar{p} \pi^0$	(3.6 ± 1.3) $\times 10^{-3}$	DESIG=12
Γ_{43}	$\Lambda \bar{\Lambda}$	(1.07 ± 0.24) $\times 10^{-3}$	DESIG=65
Γ_{44}	$K^+ \bar{p} \Lambda + \text{c.c.}$	(2.6 ± 0.4) $\times 10^{-3}$	DESIG=45
Γ_{45}	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	(3.1 ± 1.4) $\times 10^{-3}$	DESIG=82
Γ_{46}	$\Sigma^+ \bar{\Sigma}^-$	(2.1 ± 0.6) $\times 10^{-3}$	DESIG=83
Γ_{47}	$\Xi^- \bar{\Xi}^+$	(9.0 ± 2.6) $\times 10^{-4}$	DESIG=66
Γ_{48}	$\pi^+ \pi^- p \bar{p}$	(5.3 ± 1.8) $\times 10^{-3}$	DESIG=67
			DESIG=13

Radiative decays

Γ_{49}	$\gamma\gamma$	(1.58 ± 0.11) $\times 10^{-4}$	NODE=M026;CLUMP=C
			DESIG=31

Charge conjugation (C), Parity (P), Lepton family number (LF) violating modes

Γ_{50}	$\pi^+ \pi^-$	$P, CP < 1.1 \times 10^{-4}$	90% DESIG=51
Γ_{51}	$\pi^0 \pi^0$	$P, CP < 4 \times 10^{-5}$	90% DESIG=52
Γ_{52}	$K^+ K^-$	$P, CP < 6 \times 10^{-4}$	90% DESIG=53
Γ_{53}	$K_S^0 K_S^0$	$P, CP < 3.1 \times 10^{-4}$	90% DESIG=54

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 8 combinations of partial widths obtained from integrated cross section, and 19 branching ratios uses 93 measurements and one constraint to determine 13 parameters. The overall fit has a $\chi^2 = 121.6$ for 81 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	15									
x_{15}	3	5								
x_{27}	17	33	6							
x_{28}	7	15	3	45						
x_{31}	9	17	3	19	9					
x_{35}	7	12	2	20	9	8				
x_{38}	11	21	4	24	11	13	9			
x_{41}	10	19	3	25	11	11	9	14		
x_{43}	2	4	1	6	3	3	2	3	23	
x_{49}	-26	-49	-9	-56	-25	-30	-22	-37	-36	-8
Γ	-1	-2	0	-3	-1	-2	-1	-2	6	1
	x_4	x_7	x_{15}	x_{27}	x_{28}	x_{31}	x_{35}	x_{38}	x_{41}	x_{43}
Γ										-29
										x_{49}

Mode	Rate (MeV)	
$\Gamma_4 K^*(892) \bar{K}^*(892)$	0.22 \pm 0.04	DESIG=18
$\Gamma_7 \phi\phi$	0.057 \pm 0.006	DESIG=17
$\Gamma_{15} f_2(1270) f_2(1270)$	0.31 \pm 0.08	DESIG=46
$\Gamma_{27} K\bar{K}\pi$	2.33 \pm 0.13	DESIG=14
$\Gamma_{28} K\bar{K}\eta$	0.43 \pm 0.05	DESIG=25
$\Gamma_{31} K^+ K^- \pi^+ \pi^-$	0.220 \pm 0.034	DESIG=15
$\Gamma_{35} 2(K^+ K^-)$	0.047 \pm 0.010	DESIG=27
$\Gamma_{38} 2(\pi^+ \pi^-)$	0.31 \pm 0.04	DESIG=11
$\Gamma_{41} p\bar{p}$	0.046 \pm 0.005	DESIG=12
$\Gamma_{43} \Lambda\bar{\Lambda}$	0.034 \pm 0.008	DESIG=45
$\Gamma_{49} \gamma\gamma$	0.00506 \pm 0.00034	DESIG=31

$\eta_c(1S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$				Γ_{49}
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.06 \pm 0.34 OUR FIT				
[5.0 ± 0.4 keV OUR 2019 FIT]				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.8 \pm 1.1	486	1 ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
5.2 \pm 1.2	273 \pm 43	2,3 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
5.5 \pm 1.2 \pm 1.8	157 \pm 33	4 KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$
7.4 \pm 0.4 \pm 2.3		5 ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
13.9 \pm 2.0 \pm 3.0	41	6 ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \eta_c$
3.8 \pm 1.1 \pm 1.9	190	7 AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
7.6 \pm 0.8 \pm 2.3		5,8 BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
6.9 \pm 1.7 \pm 2.1	76	9 ACCIARRI	99T L3	$e^+ e^- \rightarrow e^+ e^- \eta_c$
27 \pm 16 \pm 10	5	5 SHIRAI	98 AMY	58 $e^+ e^-$

NODE=M026217

NODE=M026W1

NODE=M026W1

NEW

6.7	± 2.4	± 2.3	4	ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
11.3	± 4.2		10	ALBRECHT	94H ARG	$e^+e^- \rightarrow e^+e^-\eta_c$
8.0	± 2.3	± 2.4	17	ADRIANI	93N L3	$e^+e^- \rightarrow e^+e^-\eta_c$
5.9	± 2.1	± 1.9	7	CHEN	90B CLEO	$e^+e^- \rightarrow e^+e^-\eta_c$
6.4	± 5.0		12	AIHARA	88D TPC	$e^+e^- \rightarrow e^+e^-X$
4.3	± 3.4	± 2.4	4	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
28	± 15		5,13	BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

1 Assuming there is no interference with the non-resonant background.

2 Calculated by us using $\Gamma(\eta_c \rightarrow K\bar{K}\pi) \times \Gamma(\eta_c \rightarrow \gamma\gamma) / \Gamma = 0.44 \pm 0.05$ keV from PDG 06 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.

3 Systematic errors not evaluated.

4 Normalized to $B(\eta_c \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$.

5 Normalized to $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$.

6 Average of $K_S^0 K^\pm \pi^\mp$, $\pi^+ \pi^- K^+ K^-$, and $2(K^+ K^-)$ decay modes.

7 Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

8 Superseded by ASNER 04.

9 Normalized to the sum of 9 branching ratios.

10 Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi\phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

11 Superseded by ACCIARRI 99T.

12 Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow 2K^+ 2K^-)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.

13 Re-evaluated by AIHARA 88D.

NODE=M026W1;LINKAGE=ZH

NODE=M026W1;LINKAGE=AU

NODE=M026W1;LINKAGE=NS

NODE=M026W1;LINKAGE=N3

NODE=M026W1;LINKAGE=N2

NODE=M026W1;LINKAGE=FF

NODE=M026W1;LINKAGE=N6

NODE=M026W1;LINKAGE=NN

NODE=M026W1;LINKAGE=N1

NODE=M026W1;LINKAGE=N5

NODE=M026W1;LINKAGE=WD

NODE=M026W1;LINKAGE=N4

NODE=M026W1;LINKAGE=A

NODE=M026220

NODE=M026G10

NODE=M026G10

NODE=M026G10;LINKAGE=A

NODE=M026G09

NODE=M026G09

NODE=M026G08

NODE=M026G08

NEW

NODE=M026G07

NODE=M026G07

NODE=M026G07;LINKAGE=LI

NODE=M026G03

NODE=M026G03

NODE=M026G03;LINKAGE=LI

$\eta_c(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$$\Gamma(\eta'(958)\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_1 \Gamma_{49}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
98.1±3.9±11.7	2673	XU	18	BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

75.8	± 6.3	± 8.4	486	¹ ZHANG	12A BELL $e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
------	-----------	-----------	-----	--------------------	---

1 Superseded by XU 18.

$$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_2 \Gamma_{49}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<39	90	< 1556	UEHARA	08	BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<39	90	< 1556	UEHARA	08	BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$
-----	----	--------	--------	----	---

$$\Gamma(K^*(892)\bar{K}^*(892)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_4 \Gamma_{49}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
36 ± 6 OUR FIT				[35 ± 6 eV OUR 2019 FIT]

32.4±4.2±5.8	882 ± 115	UEHARA	08	BELL $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
---------------------	-----------	--------	----	--

$$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_7 \Gamma_{49}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.0 ± 0.8 OUR FIT				

7.75±0.66±0.62	386 ± 31	¹ LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$
-----------------------	----------	------------------	----------	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.8	± 1.2	± 1.3	132 ± 23	UEHARA	08 BELL $\gamma\gamma \rightarrow 2(K^+K^-)$
-----	-----------	-----------	----------	--------	--

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.

$$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{13} \Gamma_{49}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.67±2.86±0.96	85 ± 29	¹ LIU	12B BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{14}\Gamma_{49}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.49	90	¹ LIU	12B BELL	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	
1 Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.					

 $\Gamma(f_2(1270)f_2(1270)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{15}\Gamma_{49}/\Gamma$
50±13 OUR FIT [49 ± 13 eV OUR 2019 FIT]					
69±17±12	3182 ± 766	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$	

 $\Gamma(f_2(1270)f'_2(1525)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{16}\Gamma_{49}/\Gamma$
49±9±13	1128 ± 206	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$	

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{27}\Gamma_{49}/\Gamma$
0.369±0.020 OUR FIT [0.367 ± 0.021 keV OUR 2019 FIT]						

0.407±0.027 OUR AVERAGE Error includes scale factor of 1.2.

$0.374 \pm 0.009 \pm 0.031$	14k	¹ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$	
$0.407 \pm 0.022 \pm 0.028$		2,3 ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0K^\pm\pi^\mp$	
$0.60 \pm 0.12 \pm 0.09$	41	3,4 ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$	
$1.47 \pm 0.87 \pm 0.27$		3 SHIRAI	98 AMY	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$	
0.84 ± 0.21		3 ALBRECHT	94H ARG	$\gamma\gamma \rightarrow K^\pm K_S^0\pi^\mp$	
0.60 ± 0.23		3 CHEN	90B CLEO	$\gamma\gamma \rightarrow \eta_c K^\pm K_S^0\pi^\mp$	
$1.06 \pm 0.41 \pm 0.27$	11	3 BRAUNSCH...	89 TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$	
1.5 ± 0.60	7	3 BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.386 \pm 0.008 \pm 0.021$	12k	5 DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$	
$0.418 \pm 0.044 \pm 0.022$		3,6 BRANDENB... 00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0\pi^\mp$	
<0.63	95	3 BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
<4.4	95	ALTHOFF	85B TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

1 From the corrected and unfolded mass spectrum.

2 Calculated by us from the value reported in ASNER 04 that assumes $B(\eta_c \rightarrow K\bar{K}\pi) = 5.5 \pm 1.7\%$ 3 We have multiplied $K^\pm K_S^0\pi^\mp$ measurement by 3 to obtain $K\bar{K}\pi$.4 Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow K_S^0K^\pm\pi^\mp) = (1.5 \pm 0.4)\%$.

5 Not independent from the measurements reported by LEES 10.

6 Superseded by ASNER 04.

 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{31}\Gamma_{49}/\Gamma$
35 ± 5 OUR FIT					
27 ± 6 OUR AVERAGE					

$25.7 \pm 3.2 \pm 4.9$	2019 ± 248	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$	
$280 \pm 100 \pm 60$	42	¹ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$	
$170 \pm 80 \pm 20$	13.9 ± 6.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$	

1 Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow \pi^+\pi^-K^+K^-) = (2.0 \pm 0.7)\%$. $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{32}\Gamma_{49}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

$0.190 \pm 0.006 \pm 0.028$	11k	¹ DEL-AMO-SA..11M BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	
-----------------------------	-----	-----------------------------------	--	--

1 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M026G04

NODE=M026G04

NODE=M026G04;LINKAGE=LI

NODE=M026G19

NODE=M026G19

NEW

NODE=M026G20

NODE=M026G20

NODE=M026G14

NODE=M026G14

NEW

NODE=M026G14;LINKAGE=LE

NODE=M026G14;LINKAGE=AA

NODE=M026G14;LINKAGE=C

NODE=M026G;LINKAGE=BB

NODE=M026G14;LINKAGE=DE

NODE=M026G14;LINKAGE=NN

NODE=M026G15

NODE=M026G15

NODE=M026G;LINKAGE=CC

NODE=M026G02

NODE=M026G02

NODE=M026G02;LINKAGE=DE

$\Gamma(2(K^+K^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{35}\Gamma_{49}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
7.4 ± 1.5 OUR FIT					NODE=M026G27 NODE=M026G27
5.8 ± 1.9 OUR AVERAGE					
5.6 ± 1.1 ± 1.6	216 ± 42	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$	
350 ± 90 ± 60	46	1 ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow 2(K^+K^-)$	
231 ± 90 ± 23	9.1 ± 3.3	2 ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(K^+K^-)$	
1 Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow K^+K^-) = (2.1 \pm 1.2)\%$.					
2 Includes all topological modes except $\eta_c \rightarrow \phi\phi$.					
$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{38}\Gamma_{49}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
49 ± 6 OUR FIT					NODE=M026G11 NODE=M026G11
42 ± 6 OUR AVERAGE					
40.7 ± 3.7 ± 5.3	5381 ± 492	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$	
180 ± 70 ± 20	21.4 ± 8.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$	
$\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{41}\Gamma_{49}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
7.4 ± 0.7 OUR FIT [7.6 ± 0.7 eV OUR 2019 FIT]					NODE=M026G01 NODE=M026G01 NEW
7.20 ± 1.53 ± 0.67	157 ± 33	1 KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.6 ± 1.3 ± 0.4	190	1 AMBROGIANI	03 E835	$\bar{p}p \rightarrow \gamma\gamma$	
8.1 ± 2.9 ± 2.0		1 ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$	
1 Not independent from the $\Gamma_{\gamma\gamma}$ reported by the same experiment.					
$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{53}\Gamma_{49}/\Gamma$	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.6	90	1 UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	NODE=M026G05 NODE=M026G05
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.29	90	2 UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	OCCUR=2
1 Taking into account interference with the non-resonant continuum.					
2 Neglecting interference with the non-resonant continuum.					
$\eta_c(1S)$ BRANCHING RATIOS					
———— HADRONIC DECAYS ——					
$\Gamma(\eta'(958)\pi\pi)/\Gamma_{\text{total}}$				Γ_1/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.041 ± 0.017	14	1 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	NODE=M026225
1 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.					
$\Gamma(\rho\rho)/\Gamma_{\text{total}}$				Γ_2/Γ	NODE=M026305
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
18 ± 5 OUR AVERAGE					NODE=M026R14 NODE=M026R14
12.6 ± 3.8 ± 5.1	72	1 ABLIKIM	05L BES2	$J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$	NODE=M026R14;LINKAGE=E
26.0 ± 2.4 ± 8.8	113	1 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^0\rho^0$	
23.6 ± 10.6 ± 8.2	32	1 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^+\rho^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<14	90	1 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	OCCUR=2
1 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.					
$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_3/Γ	NODE=M026R9;LINKAGE=E
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	NODE=M026R9 NODE=M026R9
0.02 ± 0.007	63	1,2 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
1 BALTRUSAITIS 86 has an error according to Partridge.					
2 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.					

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
70±13 OUR FIT [(71 ± 13) × 10 ⁻⁴ OUR 2019 FIT]					
91±26 OUR AVERAGE					
108±25±44 60 1 ABLIKIM 05L BES2 $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \gamma$ 82±28±27 14 1 BISELLO 91 DM2 $e^+ e^- \rightarrow \gamma K^+ K^- \pi^+ \pi^-$ 90±50 9 1 BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$					

¹ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

 $\Gamma(K^*(892)^0\bar{K}^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
113±47±24	45	1 ABLIKIM	06A BES2	$J/\psi \rightarrow K^* K^0 \pi^+ \pi^- \gamma$	
1 ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0\bar{K}^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.91 \pm 0.64 \pm 0.48) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
2.9^{+0.9}_{-0.8}±1.1	$14.1^{+4.4}_{-3.7}$	1 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$	

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
17.7± 1.9 OUR FIT [(17.9 ± 2.0) × 10 ⁻⁴ OUR 2019 FIT]					

28 ± 4 OUR AVERAGE

26 \pm 4 — 8 \pm 5 25.3 \pm 5.1 \pm 9.1 26 \pm 9 31 \pm 7 \pm 10 30 \pm 18 \pm 10 74 \pm 18 \pm 24 67 \pm 21 \pm 24	1.2k	1 ABLIKIM 2 ABLIKIM 2 BAI 2 BISELLO 2 BISELLO 2 BAI 2 BAI	17P BES3 05L BES2 04 BES 91 DM2 91 DM2 90B MRK3 90B MRK3	$J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$ $J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$ $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$ $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	I I I I I I I
--	------	---	--	---	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

18 \pm 8 — 6 \pm 7	7	3 HUANG	03 BELL	$B^+ \rightarrow (\phi\phi) K^+$	
---------------------------	---	---------	---------	----------------------------------	--

¹ ABLIKIM 17P reports $[\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (4.3 \pm 0.5^{+0.5}_{-1.2}) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

 $\Gamma(\phi\phi)/\Gamma(K \bar{K} \pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_{27}
0.0243±0.0025 OUR FIT [0.0245 ± 0.0025 OUR 2019 FIT]					

0.044^{+0.012}_{-0.010} OUR AVERAGE

0.055 \pm 0.014 \pm 0.005	AUBERT,B	04B BABR	$B^\pm \rightarrow K^\pm \eta_c$		
0.032 \pm 0.014 \pm 0.009	7	1 HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi\phi$	

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R8

NODE=M026R8

NEW

NODE=M026R8;LINKAGE=E

NODE=M026R25

NODE=M026R25

NODE=M026R25;LINKAGE=AB

NODE=M026R21

NODE=M026R21

NODE=M026R;LINKAGE=BB

NODE=M026R7

NODE=M026R7

NEW

OCCUR=2

OCCUR=3

NODE=M026R7;LINKAGE=A

NODE=M026R;LINKAGE=E

NODE=M026R7;LINKAGE=BB

NODE=M026R39

NODE=M026R39

NEW

NODE=M026R39;LINKAGE=BB

$\Gamma(\phi\phi)/\Gamma(p\bar{p})$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ_{41}
$1.79 \pm 0.14 \pm 0.32$	6.4k	1 AAIJ	17BB LHCb	$p\bar{p} \rightarrow b\bar{b} X \rightarrow 2(K^+K^-)X$	
1 Using inputs from AAIJ 15AS and AAIJ 15BI and $\Gamma(b \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}} = (1.16 \pm 0.10)\%$ and $\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} = (2.120 \pm 0.029) \times 10^{-3}$ from PDG 16.					

 $\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_8/Γ
<40	90	1 ABLIKIM	06A	$BES2 \quad J/\psi \rightarrow \phi 2(\pi^+\pi^-)\gamma$	
1 ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow \phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.603 \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.					

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_9/Γ
<0.02	90	1,2 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
1 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. 2 We are assuming $B(a_0(980) \rightarrow \eta\pi) > 0.5$.					

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{10}/Γ
<0.02	90	1 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
1 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.					

 $\Gamma(K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{11}/Γ
<0.0128	90	BISELLO	91	$DM2 \quad J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
<0.0132	90	1 BISELLO	91	$DM2 \quad J/\psi \rightarrow \gamma K^+ K^- \pi^0$	
1 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.					

 $\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{12}/Γ
<0.011	90	1 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
1 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.					

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ
$2.9 \pm 0.5 \pm 0.6$	1705	1 ABLIKIM	19AV	BES3	$J/\psi \rightarrow \gamma\omega\omega$	
<3.1	90	2 BALTRUSAIT..86	MRK3		$J/\psi \rightarrow \eta_c\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						

<6.3	90	2 ABLIKIM	05L	BES2	$J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$	
<6.3	90	2 BISELLO	91	DM2	$J/\psi \rightarrow \gamma\omega\omega$	
1 ABLIKIM 19AV reports $[\Gamma(\eta_c(1S) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (4.90 \pm 0.17 \pm 0.77) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.,						

2 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.	Γ_{14}/Γ
---	----------------------

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{14}/Γ
$< 2.5 \times 10^{-4}$	90	1 ABLIKIM	17P	BES3	$J/\psi \rightarrow \pi^+\pi^-\pi^0K^+K^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

<17 $\times 10^{-4}$	90	2 ABLIKIM	05L	BES2	$J/\psi \rightarrow \pi^+\pi^-\pi^0K^+K^-\gamma$	
----------------------	----	-----------	-----	------	--	--

1 Using $B(J/\psi \rightarrow \gamma\eta_c) = 0.017 \pm 0.004$.

2 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R52
NODE=M026R52

NODE=M026R52;LINKAGE=A

NODE=M026R26
NODE=M026R26

NODE=M026R26;LINKAGE=AB

NODE=M026R11
NODE=M026R11

NODE=M026R11;LINKAGE=E
NODE=M026R11;LINKAGE=F

NODE=M026R12
NODE=M026R12

NODE=M026R12;LINKAGE=E

NODE=M026R17
NODE=M026R17

OCCUR=2

NODE=M026R17;LINKAGE=E

NODE=M026R13
NODE=M026R13

NODE=M026R13;LINKAGE=E

NODE=M026R10
NODE=M026R10

NODE=M026R10;LINKAGE=C

NODE=M026R10;LINKAGE=E

NODE=M026R22
NODE=M026R22

NODE=M026R22;LINKAGE=A

NODE=M026R22;LINKAGE=E

$\Gamma(f_2(1270)f_2(1270))/\Gamma_{\text{total}}$				Γ_{15}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.98 ± 0.25 OUR FIT					NODE=M026R19 NODE=M026R19
0.77^{+0.25}_{-0.30} ± 0.17	91.2 ± 19.8	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$	
		1 ABLIKIM 04M reports $[\Gamma(\eta_c(1S) \rightarrow f_2(1270)f_2(1270))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.3 \pm 0.3^{+0.3}_{-0.4}) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.			
$\Gamma(f_0(980)\eta)/\Gamma_{\text{total}}$				Γ_{17}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R41 NODE=M026R41
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$	
$\Gamma(f_0(1500)\eta)/\Gamma_{\text{total}}$				Γ_{18}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R42 NODE=M026R42
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$	
$\Gamma(f_0(2200)\eta)/\Gamma_{\text{total}}$				Γ_{19}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R43 NODE=M026R43
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta$	
$\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$				Γ_{20}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R44 NODE=M026R44
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$	
$\Gamma(a_0(1320)\pi)/\Gamma_{\text{total}}$				Γ_{21}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R45 NODE=M026R45
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$	
$\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$				Γ_{22}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R46 NODE=M026R46
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$	
$\Gamma(a_0(1950)\pi)/\Gamma_{\text{total}}$				Γ_{23}/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M026R00 NODE=M026R00
seen	12k	¹ LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$	
		1 From a model-independant partial wave analysis.			
$\Gamma(K_0^*(1430)\bar{K})/\Gamma_{\text{total}}$				Γ_{24}/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M026R47 NODE=M026R47
seen	12k	¹ LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$	
seen		LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta/\pi^0$	
		1 From a model-independant partial wave analysis.			
$\Gamma(K_2^*(1430)\bar{K})/\Gamma_{\text{total}}$				Γ_{25}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M026R48 NODE=M026R48
seen	LEES	14E	BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$	
$\Gamma(K_0^*(1950)\bar{K})/\Gamma_{\text{total}}$				Γ_{26}/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M026R49 NODE=M026R49
seen	12K	¹ LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$	
seen		LEES	14E	BABR Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta/\pi^0$	
		1 From a Dalitz plot analysis using an isobar model.			
$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$				Γ_{27}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
7.3 ± 0.4 OUR FIT					NODE=M026R4 NODE=M026R4
		[(7.3 ± 0.5) × 10 ⁻² OUR 2019 FIT]			
6.9 ± 0.4 OUR AVERAGE					NEW
		[(6.5 ± 0.6) × 10 ⁻² OUR 2019 AVERAGE]			
6.9 ± 0.7 ± 0.6	146	¹ ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$	
7.8 ± 0.6 ± 0.6	267	² ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$	
6.3 ± 1.3 ± 0.6	55	3,4 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^0$	OCCUR=2

7.9 ± 1.4 ± 0.7	107	5,6 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\pm$	
8.5 ± 1.8		7 AUBERT	06E BABR	$B^\pm \rightarrow K^\pm \chi_{c\bar{c}}$	OCCUR=2
5.1 ± 2.1	0.6k	8 BAI	04 BES	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$	
6.90 ± 1.42 ± 1.32	33	8 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	
5.43 ± 0.94 ± 0.94	68	8 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$	OCCUR=2
4.8 ± 1.7	95	8,9 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$	
16.1 +9.2 -7.3		10,11 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 10.7 90% CL 8,12 PARTRIDGE 80B CBAL $J/\psi \rightarrow \eta_c \gamma$

1 ABLIKIM 19AP quotes $B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.15 \pm 0.12 \pm 0.10) \times 10^{-2}$ which we multiply by 6 to account for isospin symmetry.

2 ABLIKIM 19AP quotes $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (2.60 \pm 0.21 \pm 0.20) \times 10^{-2}$ which we multiply by 3 to account for isospin symmetry.

3 ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (4.54 \pm 0.76 \pm 0.48) \times 10^{-6}$ which we multiply by 6 to account for isospin symmetry.

4 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (27.24 \pm 4.56 \pm 2.88) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

5 ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (11.35 \pm 1.25 \pm 1.50) \times 10^{-6}$ which we multiply by 3 to account for isospin symmetry.

6 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (34.05 \pm 3.75 \pm 4.50) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

7 Determined from the ratio of $B(B^\pm \rightarrow K^\pm \eta_c) B(\eta_c \rightarrow K\bar{K}\pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT,B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E.

8 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

9 Average from $K^+ K^- \pi^0$ and $K^\pm K_S^0 \pi^\mp$ decay channels.

10 $K^\pm K_S^0 \pi^\mp$ corrected to $K\bar{K}\pi$ by factor 3. KS, MR.

11 Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

12 $K^+ K^- \pi^0$ corrected to $K\bar{K}\pi$ by factor 6. KS, MR

$\Gamma(\phi K^+ K^-)/\Gamma(K\bar{K}\pi)$						Γ_6/Γ_{27}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
0.052 +0.016 -0.014 ± 0.014	7	1 HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi \phi$		

1 Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 + 0.10 - 0.12) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(K\bar{K}\eta)/\Gamma_{\text{total}}$						Γ_{28}/Γ
VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
1.36 ± 0.15 OUR FIT						

$[(1.36 \pm 0.16) \times 10^{-2}$ OUR 2019 FIT]

1.0 ± 0.5 ± 0.2 7 1,2 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma \eta K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.1 90 3 BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$

1 ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (2.11 \pm 1.01 \pm 0.32) \times 10^{-6}$ which we multiply by 2 to account for isospin symmetry.

2 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (4.22 \pm 2.02 \pm 0.64) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (51 \pm 6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

3 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

OCCUR=2

OCCUR=2

NODE=M026R4;LINKAGE=C

NODE=M026R4;LINKAGE=F

NODE=M026R4;LINKAGE=BK

NODE=M026R4;LINKAGE=CK

NODE=M026R4;LINKAGE=BL

NODE=M026R4;LINKAGE=CL

NODE=M026R4;LINKAGE=AB

NODE=M026R4;LINKAGE=E

NODE=M026R4;LINKAGE=D

NODE=M026R4;LINKAGE=01

NODE=M026R4;LINKAGE=A

NODE=M026R4;LINKAGE=02

NODE=M026R02

NODE=M026R02

NODE=M026R02;LINKAGE=BB

NODE=M026R15

NODE=M026R15

NEW

OCCUR=2

NODE=M026R15;LINKAGE=AK

NODE=M026R15;LINKAGE=AM

NODE=M026R15;LINKAGE=E

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$	Γ_{28}/Γ_{27}	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
0.186±0.018 OUR FIT		
0.190±0.008±0.017	5.4k	1 LEES 14E BABR $\gamma\gamma \rightarrow K^+ K^- \eta/\pi^0$
1 LEES 14E reports $B(\eta_c(1S) \rightarrow K^+ K^- \eta)/B(\eta_c(1S) \rightarrow K^+ K^- \pi^0) = 0.571 \pm 0.025 \pm 0.051$, which we divide by 3 to account for isospin symmetry. It uses both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.		
$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{29}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
1.7±0.4±0.1	33	1 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma \eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
5.4±2.0	75	2 BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$
3.7±1.3±2.0	18	2 PARTRIDGE 80B CBAL $J/\psi \rightarrow \eta\pi^+\pi^-\gamma$
1 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (7.22 \pm 1.47 \pm 1.11) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M026R6;LINKAGE=AB
2 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.		NODE=M026R6;LINKAGE=E
$\Gamma(\eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_{30}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
4.4±1.2±0.4	39	1 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma \eta 2(\pi^+\pi^-)$
1 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (19.17 \pm 3.77 \pm 3.72) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M026R05;LINKAGE=AB
$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_{31}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
6.9±1.0 OUR FIT		
$[(6.9 \pm 1.1) \times 10^{-3}$ OUR 2019 FIT]		
11.2±1.9 OUR AVERAGE		
9.7±2.2±0.9	38	1 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^+ \pi^-$
12±4	0.4k	2 BAI 04 BES $J/\psi \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
21±7	110	2 BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c \gamma$
14 ⁺²² ₋₉	3 HIMEL 80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$
1 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (4.16 \pm 0.76 \pm 0.59) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M026R5;LINKAGE=AB
2 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.		NODE=M026R5;LINKAGE=E
3 Estimated using $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = 0.0028 \pm 0.0006$.		NODE=M026R5;LINKAGE=A
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$	Γ_{32}/Γ_{27}	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
0.477±0.017±0.070	11k	1 DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1 We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.		NODE=M026R01;LINKAGE=DE
$\Gamma(K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{33}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
5.6±1.4±0.5	43	1,2 ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\mp 2\pi^\pm$
		NODE=M026R06

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^- \pi^- 2\pi^+) = (12.01 \pm 2.22 \pm 2.04) \times 10^{-6}$ which we multiply by 2 to take c.c. into account.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + c.c.) / \Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}}] = (24.02 \pm 4.44 \pm 4.08) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.5 ± 2.4 OUR AVERAGE				
8 ± 4 ± 1	10	¹ ABLIKIM 12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- 2(\pi^+ \pi^-)$	
$7.2 \pm 2.4 \pm 1.5$	100	² ABLIKIM 06A BES2	$J/\psi \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$	
¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}}] = (3.60 \pm 1.71 \pm 0.64) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.21 \pm 0.32 \pm 0.24) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.46 ± 0.30 OUR FIT				
$[(1.47 \pm 0.31) \times 10^{-3}$ OUR 2019 FIT]				
$2.2 \pm 0.9 \pm 0.2$	7	¹ ABLIKIM 12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(K^+ K^-)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.4 ± 0.5 ± 0.6	$14.5^{+4.6}_{-3.0}$	² HUANG 03 BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$	
21 ± 10 ± 6		³ ALBRECHT 94H ARG	$\gamma \gamma \rightarrow K^+ K^- K^+ K^-$	
¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-)) / \Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}}] = (0.94 \pm 0.37 \pm 0.14) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.				
³ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi \phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.				

$\Gamma(2(K^+ K^-)) / \Gamma(K \bar{K} \pi)$ Γ_{35}/Γ_{27}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.020 ± 0.004 OUR FIT				
0.024 ± 0.007 OUR AVERAGE				
0.023 $\pm 0.007 \pm 0.006$		AUBERT,B 04B BABR	$B^\pm \rightarrow K^\pm \eta_c$	
$0.026^{+0.009}_{-0.007} \pm 0.007$	15	¹ HUANG 03 BELL	$B^\pm \rightarrow K^\pm (2K^+ 2K^-)$	
¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.				

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5 \times 10^{-4}$	90	¹ ABLIKIM 17AJ BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$	
¹ ABLIKIM 17AJ reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \eta_c(1S))] < 1.6 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 3.4 \times 10^{-3}$.				

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 0.9 \pm 0.4$				
¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}}] = (20.31 \pm 2.20 \pm 3.33) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P)) / \Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M026R06;LINKAGE=AA

NODE=M026R06;LINKAGE=AB

NODE=M026R23
NODE=M026R23

NODE=M026R23;LINKAGE=AL

NODE=M026R23;LINKAGE=AB

NODE=M026R20
NODE=M026R20

NEW

NODE=M026R20;LINKAGE=AB

NODE=M026R20;LINKAGE=BB

NODE=M026R20;LINKAGE=AL

NODE=M026R38
NODE=M026R38

NODE=M026R38;LINKAGE=BB

NODE=M026R51
NODE=M026R51

NODE=M026R51;LINKAGE=A

NODE=M026R07
NODE=M026R07

NODE=M026R07;LINKAGE=AB

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{38}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.97 ± 0.12 OUR FIT					NODE=M026R1 NODE=M026R1
1.35 ± 0.21 OUR AVERAGE					
1.74 ± 0.32 ± 0.15	100	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^-)$	
1.0 ± 0.5	542 ± 75	2 BAI	04 BES	$J/\psi \rightarrow \gamma 2(\pi^+ \pi^-)$	
1.05 ± 0.17 ± 0.34	137	2 BISELLO	91 DM2	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$	
1.3 ± 0.6	25	2 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$	
2.0 ± 1.5		3 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$	

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (7.51 \pm 0.85 \pm 1.11) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 0.0028 \pm 0.0006$.

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$					Γ_{39}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
16.1 ± 2.0 OUR AVERAGE					
$[(17.4 \pm 3.3) \times 10^{-2}$ OUR 2019 AVERAGE]					
15.3 ± 1.8 ± 1.8	333	ABLIKIM	19AP BES3	$h_c \rightarrow \gamma \eta_c$	
17.4 ± 2.9 ± 1.5	175	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^- \pi^0)$	

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (75.13 \pm 7.42 \pm 9.99) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{40}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
18 ± 4 OUR AVERAGE					
20 ± 5 ± 2	51	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 3(\pi^+ \pi^-)$	
15.4 ± 3.4 ± 3.3	479	2 ABLIKIM	06A BES2	$J/\psi \rightarrow 3(\pi^+ \pi^-) \gamma$	

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (8.82 \pm 1.57 \pm 1.59) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (2.59 \pm 0.32 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$					Γ_{41}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
14.5 ± 1.4 OUR FIT					
$[(15.1 \pm 1.6) \times 10^{-4}$ OUR 2019 FIT]					
12.7 ± 2.0 OUR AVERAGE					
$[(13.2 \pm 2.7) \times 10^{-4}$ OUR 2019 AVERAGE]					

12.0 ± 2.6 ± 1.5	34	ABLIKIM	19AP BES3	$h_c \rightarrow \gamma \eta_c$
15 ± 5 ± 1	15	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}$
15 ± 6	213 ± 33	2 BAI	04 BES	$J/\psi \rightarrow \gamma p\bar{p}$
10 ± 3 ± 4	18	2 BISELLO	91 DM2	$J/\psi \rightarrow \gamma p\bar{p}$
11 ± 6	23	2 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
29 ± 29		3 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$
29 ± 15				

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.4 ± 1.8 ± 1.1	195	4 WU	06 BELL	$B^+ \rightarrow p\bar{p} K^+$
------------------	-----	------	---------	--------------------------------

NODE=M026R1
NODE=M026R1

NODE=M026R1;LINKAGE=AB

NODE=M026R1;LINKAGE=E

NODE=M026R1;LINKAGE=A

NODE=M026R08
NODE=M026R08
NEW

NODE=M026R08;LINKAGE=AB

NODE=M026R24;LINKAGE=AL

NODE=M026R24;LINKAGE=AB

NODE=M026R2
NODE=M026R2

NEW

NEW

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (0.65 \pm 0.19 \pm 0.10) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

³ Estimated using $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = 0.0028 \pm 0.0006$.

⁴ WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (1.42 \pm 0.11 \pm 0.16) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.06 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p})/\Gamma(K\bar{K}\pi)$					Γ_{41}/Γ_{27}
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0199 ± 0.0019 OUR FIT					
[0.0207 ± 0.0021 OUR 2019 FIT]					
0.021 ± 0.002	+0.004	195	¹ WU	06	BELL $B^\pm \rightarrow K^\pm p\bar{p}$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \pm 0.10) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$					$\Gamma_{41}/\Gamma \times \Gamma_7/\Gamma$
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.26 ± 0.04 OUR FIT					
[$(0.27 \pm 0.05) \times 10^{-5}$ OUR 2019 FIT]					
4.0	+3.5		BAGLIN	89	SPEC $\bar{p}p \rightarrow K^+ K^- K^+ K^-$

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$					Γ_{42}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.36 ± 0.13 ± 0.03	14	¹ ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}\pi^0$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (1.53 \pm 0.49 \pm 0.23) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{43}/Γ
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.7 ± 2.4 OUR FIT					
[$(10.9 \pm 2.4) \times 10^{-4}$ OUR 2019 FIT]					
11.8 ± 2.3 ± 2.5		¹ ABLIKIM	12B	BES3	

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.9 ± 2.5	± 0.7	20	² WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda} K^+$
<20		90	³ BISELLO	91	DM2 $e^+ e^- \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ ABLIKIM 12B reports $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (0.198 \pm 0.021 \pm 0.032) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (0.95 \pm 0.25 \pm 0.08) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.06 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$					Γ_{43}/Γ_{41}
VALUE	DOCUMENT ID	TECN	COMMENT		
0.73 ± 0.16 OUR FIT					
[0.72 ± 0.16 OUR 2019 FIT]					
0.67 ± 0.19 ± 0.12		¹ WU	06	BELL $B^+ \rightarrow p\bar{p} K^+, \Lambda\bar{\Lambda} K^+$	

¹ Not independent from other $\eta_c \rightarrow \Lambda\bar{\Lambda}$, $p\bar{p}$ branching ratios reported by WU 06.

NODE=M026R2;LINKAGE=AB

NODE=M026R2;LINKAGE=E

NODE=M026R2;LINKAGE=A

NODE=M026R2;LINKAGE=WU

NODE=M026R03
NODE=M026R03
NEW

NODE=M026R03;LINKAGE=BB

NODE=M026R33
NODE=M026R33
NEW

NODE=M026R09
NODE=M026R09

NODE=M026R09;LINKAGE=AB

NODE=M026R18
NODE=M026R18
NEW

NODE=M026R18;LINKAGE=AB

NODE=M026R18;LINKAGE=WU

NODE=M026R18;LINKAGE=E

NODE=M026R27
NODE=M026R27
NEW

NODE=M026R27;LINKAGE=WU

$\Gamma(K^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.6 ± 0.4 OUR AVERAGE				$[(2.5 \pm 0.4) \times 10^{-3}$ OUR 2019 AVERAGE]
2.56^{+0.35}_{-0.33}±0.21	157	1 LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$

¹ LU 19 reports $(2.83^{+0.36}_{-0.34} \pm 0.35) \times 10^{-3}$ from a measurement of $[\Gamma(\eta_c(1S) \rightarrow K^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)]$ assuming $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \eta_c K^+) = (1.06 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{44}/Γ

NODE=M026R53

NODE=M026R53

NEW

NODE=M026R53;LINKAGE=A

 $\Gamma(\bar{\Lambda}(1520)\Lambda+c.c.)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.1±1.4 OUR AVERAGE				$[(3.1 \pm 1.3) \times 10^{-3}$ OUR 2019 AVERAGE]
3.1±1.4±0.3	43	1 LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$

¹ LU 19 reports $(3.48 \pm 1.48 \pm 0.46) \times 10^{-3}$ from a measurement of $[\Gamma(\eta_c(1S) \rightarrow \bar{\Lambda}(1520)\Lambda+c.c.)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)]$ assuming $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \eta_c K^+) = (1.06 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{45}/Γ

NODE=M026R54

NODE=M026R54

NEW

NODE=M026R54;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.1±0.3±0.5	112	1 ABLIKIM	13C BES3	$J/\psi \rightarrow \gamma p\bar{p}\pi^0\pi^0$

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.60 \pm 0.48 \pm 0.31) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{46}/Γ

NODE=M026R28

NODE=M026R28

NODE=M026R28;LINKAGE=AB

 $\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.90±0.18±0.19	78	1 ABLIKIM	13C BES3	$J/\psi \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.51 \pm 0.27 \pm 0.14) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{47}/Γ

NODE=M026R29

NODE=M026R29

NODE=M026R29;LINKAGE=AB

 $\Gamma(\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.3±1.7±0.5	19	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma p\bar{p}\pi^+\pi^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12 90 HIMEL 80B MRK2 $\psi(2S) \rightarrow \eta_c\gamma$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}] \times [\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}] = (2.30 \pm 0.65 \pm 0.36) \times 10^{-6}$ which we divide by our best value $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}} = (4.3 \pm 0.4) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{48}/Γ

NODE=M026R3

NODE=M026R3

NODE=M026R3;LINKAGE=AB

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.58±0.11 OUR FIT					$[(1.57 \pm 0.12) \times 10^{-4}$ OUR 2019 FIT]

1.9^{+0.7}_{-0.6} OUR AVERAGE

2.7 ± 0.8 ± 0.6	1 ABLIKIM	13I BES3
1.4 ^{+0.7} _{-0.5} ± 0.3	1.2 ^{+2.8} _{-1.1}	2 ADAMS 08 CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 Γ_{49}/Γ

NODE=M026R31

NODE=M026R31

NEW

NODE=M026310

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 $^{+0.9}_{-0.7}$	± 0.2	13	³ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
2.80 $^{+0.67}_{-0.58}$	± 1.0		⁴ ARMSTRONG	95F	E760	$\bar{p}p \rightarrow \gamma\gamma$
< 9		90	⁵ BISELLO	91	DM2	$J/\psi \rightarrow \gamma\gamma\gamma$
6 $^{+4}_{-3}$	± 4		⁴ BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma$
< 18		90	⁶ BLOOM	83	CBAL	$J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 13I reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (4.5 \pm 1.2 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ADAMS 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (2.4^{+1.1}_{-0.8} \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ WICHT 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (2.2^{+0.9+0.4}_{-0.7-0.2}) \times 10^{-7}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.06 \pm 0.09) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Not independent from the values of the total and two-photon width quoted by the same experiment.

⁵ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

⁶ Using $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{49}/Γ_{27}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.17 ± 0.23 OUR FIT				
$[(2.15 \pm 0.28) \times 10^{-3}$ OUR 2019 FIT]				

3.2 $^{+1.3}_{-1.0} \pm 0.8$ 13 ¹ WICHT 08 BELL $B^\pm \rightarrow K^\pm \gamma\gamma$

¹ Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{41}/\Gamma \times \Gamma_{49}/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
0.230 ± 0.022 OUR FIT				
$[(0.238 \pm 0.023) \times 10^{-6}$ OUR 2019 FIT]				

0.26 ± 0.05 OUR AVERAGE Error includes scale factor of 1.4.

0.224 $^{+0.038}_{-0.037}$	± 0.020	190	AMBROGIANI 03	E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
0.336 $^{+0.080}_{-0.070}$			ARMSTRONG 95F	E760	$\bar{p}p \rightarrow \gamma\gamma$
0.68 $^{+0.42}_{-0.31}$		12	BAGLIN	87B	SPEC $\bar{p}p \rightarrow \gamma\gamma$

— Charge conjugation (C), Parity (P), — — Lepton family number (LF) violating modes —

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	¹ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 70 90 ² ABLIKIM 06B BES2 $J/\psi \rightarrow \pi^+\pi^-\gamma$

¹ ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.82 \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.

² ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.1 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$.

$\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4	90	¹ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 40 90 ² ABLIKIM 06B BES2 $J/\psi \rightarrow \pi^0\pi^0\gamma$

NODE=M026R31;LINKAGE=AL

NODE=M026R31;LINKAGE=AD

NODE=M026R31;LINKAGE=WI

NODE=M026R31;LINKAGE=AB

NODE=M026R31;LINKAGE=E

NODE=M026R31;LINKAGE=C

NODE=M026R04

NODE=M026R04

NEW

NODE=M026R04;LINKAGE=BB

NODE=M026R32

NODE=M026R32

NEW

NODE=M026320

NODE=M026R34

NODE=M026R34

NODE=M026R34;LINKAGE=AL

NODE=M026R34;LINKAGE=AB

NODE=M026R35

NODE=M026R35

¹ ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

² ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.71 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{52}/Γ
<60	90	¹ ABLIKIM 06B	BES2	$J/\psi \rightarrow K^+ K^- \gamma$	

¹ ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.96 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{53}/Γ
<31	90	¹ ABLIKIM 06B	BES2	$J/\psi \rightarrow K_S^0 K_S^0 \gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	² UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
< 5.6	90	³ UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.53 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$.

² Taking into account interference with the non-resonant continuum.

³ Neglecting interference with the non-resonant continuum.

NODE=M026R35;LINKAGE=AL

NODE=M026R35;LINKAGE=AB

NODE=M026R36
NODE=M026R36

NODE=M026R36;LINKAGE=AB

NODE=M026R37
NODE=M026R37

OCCUR=2

NODE=M026R37;LINKAGE=AB

NODE=M026R37;LINKAGE=U1
NODE=M026R37;LINKAGE=U2

NODE=M026

REFID=59901

REFID=59998;ERROR=12

REFID=59614

REFID=59453

REFID=57896

REFID=58191

REFID=58322

REFID=57989

REFID=57125

REFID=57140

REFID=56857

REFID=57147

REFID=56130

REFID=55937

REFID=54878

REFID=54954

REFID=55592

REFID=54267

REFID=54271

REFID=54741

REFID=54303

REFID=54763

REFID=53711

REFID=16751

REFID=53927

REFID=53236

REFID=52676

REFID=52261

REFID=52267

REFID=52064

REFID=52204

REFID=51627

REFID=50987

REFID=50988

REFID=51059

REFID=51004

REFID=51472

REFID=50837

REFID=50801

REFID=50182

REFID=50329

REFID=49745

REFID=49746

REFID=50043

REFID=49620

REFID=49625

REFID=49465

REFID=49185

REFID=49206

REFID=49621

REFID=49188

REFID=48546

REFID=48553

$\eta_c(1S)$ REFERENCES

ABLIKIM	19AP	PR D100 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AV	PR D100 052012	ABLIKIM <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17P	PR D95 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	16A	PR D93 012005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
AAIJ	15AS	JHEP 1510 053	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13C	PR D87 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12F	PRL 108 222002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LIU	12B	PR 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
LEES	10	PR D81 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PRL D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BES Collab.)
ABLIKIM	06A	PL B633 19	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	06B	EPJ C45 337	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	05L	PR D72 072005	M. Ablikim <i>et al.</i>	(BES Collab.)
KUO	05	PL B621 41	C.C. Kuo <i>et al.</i>	(BELLE Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,B	04B	PR D70 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABDALLAH	03J	EPJ C31 481	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
AMBROGIANI	03	PL B566 45	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
BAI	03	PL B555 174	J.Z. Bai <i>et al.</i>	(BES Collab.)
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
BAI	00F	PR D62 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)
BRANDENB...	00B	PRL 85 3095	G. Brandenburg <i>et al.</i>	(CLEO Collab.)

ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47476
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ABREU	980	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46553
SHIRAI	98	PL B424 405	M. Shirai <i>et al.</i>	(AMY Collab.)	REFID=46381
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERM, GENO+)	REFID=44623
ALBRECHT	94H	PL B338 390	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44098
ADRIANI	93N	PL B318 575	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=43670
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
CHEN	90B	PL B243 169	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=41360
BAGLIN	89	PL B231 557	C. Baglin, S. Baird, G. Bassompierre	(R704 Collab.)	REFID=41015
BEHREND	89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40732
BRAUNSCH...	89	ZPHY C41 533	W. Braunschweig <i>et al.</i>	(TASSO Collab.)	REFID=40728
AIHARA	88D	PRL 60 2355	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=40588
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BERGER	86	PL 167B 120	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22010
GAISER	86	PR D34 711	J. Gaisser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BALTRUSAIT...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+) JP	REFID=22006
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
HIMEL	80B	PRL 45 1146	T.M. Himmel <i>et al.</i>	(SLAC, LBL, UCB)	REFID=22003
PARTRIDGE	80B	PRL 45 1150	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22004

 $J/\psi(1S)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$J/\psi(1S)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
3096.900 ± 0.006 OUR AVERAGE					
3096.900 $\pm 0.002 \pm 0.006$		1 ANASHIN	15	KEDR $e^+ e^- \rightarrow$ hadrons	
3096.89 ± 0.09	502	2 ARTAMONOV	00	OLYA $e^+ e^- \rightarrow$ hadrons	
3096.91 $\pm 0.03 \pm 0.01$		3 ARMSTRONG	93B	E760 $\bar{p}p \rightarrow e^+ e^-$	
3096.95 $\pm 0.1 \pm 0.3$	193	BAGLIN	87	SPEC $\bar{p}p \rightarrow e^+ e^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3096.66 $\pm 0.19 \pm 0.02$	6.1k	4 AAIJ	15BI	LHCb $p\bar{p} \rightarrow J/\psi X$	
3096.917 $\pm 0.010 \pm 0.007$		AULCHENKO	03	KEDR $e^+ e^- \rightarrow$ hadrons	
3097.5 ± 0.3		GRIBUSHIN	96	FMPS 515 $\pi^- Be \rightarrow 2\mu X$	
3098.4 ± 2.0	38k	LEMOIGNE	82	GOLI 185 $\pi^- Be \rightarrow \gamma\mu^+\mu^- A$	
3096.93 ± 0.09	502	5 ZHOLENTZ	80	REDE $e^+ e^-$	
3097.0 ± 1		6 BRANDELIK	79C	DASP $e^+ e^-$	

¹ Supersedes AULCHENKO 03.² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴ From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵ Superseded by ARTAMONOV 00.⁶ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$ and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$. **$J/\psi(1S)$ WIDTH**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
92.9 ± 2.8 OUR AVERAGE Error includes scale factor of 1.1.					
96.1 ± 3.2	13k	1 ADAMS	06A	CLEO $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	
84.4 ± 8.9		BAI	95B	BES $e^+ e^-$	
91 $\pm 11 \pm 6$		2 ARMSTRONG	93B	E760 $\bar{p}p \rightarrow e^+ e^-$	
85.5 ± 6.1		3 HSUEH	92	RVUE See γ mini-review	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
92.94 ± 1.83		4 ANASHIN	18A	KEDR $e^+ e^-$	
94.1 ± 2.7		5 ANASHIN	10	KEDR 3.097 $e^+ e^- \rightarrow e^+ e^-$, $\mu^+ \mu^-$	
93.7 ± 3.5	7.8k	1 AUBERT	04	BABR $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	

¹ Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(e^+ e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].³ Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.⁴ Using $\Gamma(e^+ e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ from PDG 16.⁵ Assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$ and using $\Gamma(e^+ e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

NODE=M070

REFID=47476

REFID=47385

REFID=46553

REFID=46381

REFID=44623

REFID=44098

REFID=43670

REFID=41668

REFID=41354

REFID=41360

REFID=41015

REFID=40732

REFID=40728

REFID=40588

REFID=40018

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

REFID=22008

REFID=22009

REFID=22010

REFID=22012

REFID=21349

REFID=22006

REFID=21682

REFID=22003

REFID=22004

REFID=22005

REFID=22006

REFID=22007

J/ψ(1S) DECAY MODES

NODE=M070215;NODE=M070

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 \pm 0.5) %	DESIG=3
Γ_2 virtual $\gamma \rightarrow$ hadrons	(13.50 \pm 0.30) %	DESIG=4
Γ_3 ggg	(64.1 \pm 1.0) %	DESIG=249
Γ_4 γgg	(8.8 \pm 1.1) %	DESIG=250
Γ_5 $e^+ e^-$	(5.971 \pm 0.032) %	DESIG=1
Γ_6 $e^+ e^- \gamma$	[a] (8.8 \pm 1.4) $\times 10^{-3}$	DESIG=5
Γ_7 $\mu^+ \mu^-$	(5.961 \pm 0.033) %	DESIG=2
Decays involving hadronic resonances		
Γ_8 $\rho\pi$	(1.69 \pm 0.15) %	S=2.4
Γ_9 $\rho^0\pi^0$	(5.6 \pm 0.7) $\times 10^{-3}$	DESIG=20
Γ_{10} $\rho(770)^{\mp} K^{\pm} K_S^0$	(1.9 \pm 0.4) $\times 10^{-3}$	DESIG=21
Γ_{11} $\rho(1450)\pi$		DESIG=342
Γ_{12} $\rho(1450)\pi \rightarrow \pi^+ \pi^- \pi^0$	(2.3 \pm 0.7) $\times 10^{-3}$	DESIG=310
Γ_{13} $\rho(1450)^{\pm} \pi^{\mp} \rightarrow K_S^0 K^{\pm} \pi^{\mp}$	(3.5 \pm 0.6) $\times 10^{-4}$	DESIG=328
Γ_{14} $\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0$	(2.7 \pm 0.6) $\times 10^{-4}$	DESIG=329
Γ_{15} $\rho(1450)\eta'(958) \rightarrow \pi^+ \pi^- \eta'(958)$	(3.3 \pm 0.7) $\times 10^{-6}$	DESIG=312
Γ_{16} $\rho(1700)\pi$		DESIG=345
Γ_{17} $\rho(1700)\pi \rightarrow \pi^+ \pi^- \pi^0$	(1.7 \pm 1.1) $\times 10^{-4}$	DESIG=325
Γ_{18} $\rho(2150)\pi$		DESIG=313
Γ_{19} $\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$	(8 \pm 40) $\times 10^{-6}$	DESIG=326
Γ_{20} $\rho_3(1690)\pi \rightarrow \pi^+ \pi^- \pi^0$		DESIG=314
Γ_{21} $a_2(1320)\rho$	(1.09 \pm 0.22) %	DESIG=316
Γ_{22} $\omega\pi^+\pi^+\pi^-\pi^-$	(8.5 \pm 3.4) $\times 10^{-3}$	DESIG=43
Γ_{23} $\omega\pi^+\pi^-\pi^0$	(4.0 \pm 0.7) $\times 10^{-3}$	DESIG=26
Γ_{24} $\omega\pi^+\pi^-$	(7.2 \pm 1.0) $\times 10^{-3}$	DESIG=211
Γ_{25} $\omega f_2(1270)$	(4.3 \pm 0.6) $\times 10^{-3}$	DESIG=24
Γ_{26} $K^*(892)^0 \bar{K}^*(892)^0$	(2.3 \pm 0.6) $\times 10^{-4}$	DESIG=28
Γ_{27} $K^*(892)^{\pm} K^*(892)^{\mp}$	(1.00 \pm 0.22) $\times 10^{-3}$	DESIG=46
Γ_{28} $K^*(892)^{\pm} K^*(700)^{\mp}$	(1.1 \pm 1.0) $\times 10^{-3}$	DESIG=256
Γ_{29} $K_S^0\pi^- K^*(892)^+ + \text{c.c.}$	(2.0 \pm 0.5) $\times 10^{-3}$	DESIG=257
Γ_{30} $K_S^0\pi^- K^*(892)^+ + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	(6.7 \pm 2.2) $\times 10^{-4}$	DESIG=299
Γ_{31} $K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0$	(6.3 \pm 0.5) $\times 10^{-6}$	DESIG=300
Γ_{32} $K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	(2.69 \pm 0.25) $\times 10^{-4}$	DESIG=376
Γ_{33} $K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	(1.10 \pm 0.60) $\times 10^{-5}$	DESIG=381
Γ_{34} $K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	(6.2 \pm 2.9) $\times 10^{-6}$	DESIG=382
Γ_{35} $\eta K^*(892)^0 \bar{K}^*(892)^0$	(1.15 \pm 0.26) $\times 10^{-3}$	DESIG=383
Γ_{36} $\eta' K^{\pm} K^{\mp}$	(1.48 \pm 0.13) $\times 10^{-3}$	DESIG=252
Γ_{37} $\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	(1.66 \pm 0.21) $\times 10^{-3}$	DESIG=355
Γ_{38} $\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	(2.16 \pm 0.31) $\times 10^{-4}$	DESIG=357
Γ_{39} $\eta' h_1(1415) \rightarrow \eta' K^{\pm} K^{\mp}$	(1.51 \pm 0.23) $\times 10^{-4}$	DESIG=353
Γ_{40} $K^*(1410) \bar{K} + \text{c.c.}$		DESIG=354
Γ_{41} $K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^{\pm} K^{\mp} \pi^0$	(7 \pm 4) $\times 10^{-5}$	DESIG=317
Γ_{42} $K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^{\pm} \pi^{\mp}$	(8 \pm 6) $\times 10^{-5}$	DESIG=330
Γ_{43} $K_2^*(1430) \bar{K} + \text{c.c.}$		DESIG=318
Γ_{44} $K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^{\pm} K^{\mp} \pi^0$	(1.0 \pm 0.5) $\times 10^{-4}$	DESIG=321

Γ_{45}	$K_2^*(1430)\bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(4.0 \pm 1.0) \times 10^{-4}$	DESIG=320
Γ_{46}	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	$(4.66 \pm 0.31) \times 10^{-3}$	DESIG=48
Γ_{47}	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$	DESIG=303
Γ_{48}	$K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(4 \pm 4) \times 10^{-4}$	DESIG=304
Γ_{49}	$K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$	DESIG=235
Γ_{50}	$\omega K^*(892) \bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$	DESIG=102
Γ_{51}	$\bar{K} K^*(892) + \text{c.c.}$		DESIG=331
Γ_{52}	$\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(5.0 \pm 0.5) \times 10^{-3}$	DESIG=332
Γ_{53}	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm 0.8) \times 10^{-3}$	S=2.9 DESIG=121
Γ_{54}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.13) \times 10^{-3}$	DESIG=231
Γ_{55}	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$	DESIG=232
Γ_{56}	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$	DESIG=122
Γ_{57}	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$	DESIG=233
Γ_{58}	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$	DESIG=132
Γ_{59}	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(7.7 \pm 1.6) \times 10^{-3}$	DESIG=214
Γ_{60}	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$	DESIG=343
Γ_{61}	$K^*(892)^0 K_S^0 \pi^0$	$(6 \pm 4) \times 10^{-4}$	DESIG=344
Γ_{62}	$\omega \pi^0 \pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$	DESIG=140
Γ_{63}	$\omega \pi^0 \eta$	$(3.4 \pm 1.7) \times 10^{-4}$	DESIG=360
Γ_{64}	$b_1(1235)^\pm \pi^\mp$	[b] $(3.0 \pm 0.5) \times 10^{-3}$	DESIG=49
Γ_{65}	$\omega K^\pm K_S^0 \pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$	DESIG=101
Γ_{66}	$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$	DESIG=160
Γ_{67}	$\eta K^\pm K_S^0 \pi^\mp$	[b] $(2.2 \pm 0.4) \times 10^{-3}$	DESIG=230
Γ_{68}	$\phi K^*(892) \bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$	DESIG=104
Γ_{69}	$\omega K \bar{K}$	$(1.9 \pm 0.4) \times 10^{-3}$	DESIG=27
Γ_{70}	$\omega f_0(1710) \rightarrow \omega K \bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$	DESIG=130
Γ_{71}	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$	DESIG=35
Γ_{72}	$\Delta(1232)^{++} \bar{p} \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$	DESIG=70
Γ_{73}	$\omega \eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6 DESIG=30
Γ_{74}	$\omega \eta' \pi^+ \pi^-$	$(1.12 \pm 0.13) \times 10^{-3}$	DESIG=385
Γ_{75}	$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3 DESIG=36
Γ_{76}	$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$	DESIG=305
Γ_{77}	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$	DESIG=129
Γ_{78}	$\phi K^+ K^-$	$(8.3 \pm 1.2) \times 10^{-4}$	DESIG=295
Γ_{79}	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$	DESIG=39
Γ_{80}	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$	DESIG=66
Γ_{81}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.})$	[b] $(1.16 \pm 0.05) \times 10^{-3}$	DESIG=67
Γ_{82}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(1.07 \pm 0.08) \times 10^{-3}$	DESIG=309
Γ_{83}	$K^+ K^- f'_2(1525)$	$(1.04 \pm 0.35) \times 10^{-3}$	DESIG=308
Γ_{84}	$\phi f'_2(1525)$	$(8 \pm 4) \times 10^{-4}$	S=2.7 DESIG=40
Γ_{85}	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7 DESIG=34
Γ_{86}	$\phi \pi^0 \pi^0$	$(5.0 \pm 1.0) \times 10^{-4}$	DESIG=76
Γ_{87}	$\phi K^\pm K_S^0 \pi^\mp$	[b] $(7.2 \pm 0.8) \times 10^{-4}$	DESIG=103
Γ_{88}	$\omega f_1(1420)$	$(6.8 \pm 2.4) \times 10^{-4}$	DESIG=105
Γ_{89}	$\phi \eta$	$(7.4 \pm 0.8) \times 10^{-4}$	S=1.5 DESIG=37
Γ_{90}	$\Xi^0 \bar{\Xi}^0$	$(1.17 \pm 0.04) \times 10^{-3}$	DESIG=248
Γ_{91}	$\Xi(1530)^- \bar{\Xi}^+ + \text{c.c.}$	$(3.18 \pm 0.08) \times 10^{-4}$	DESIG=107
Γ_{92}	$p K^- \bar{\Sigma}(1385)^0$	$(5.1 \pm 3.2) \times 10^{-4}$	DESIG=74
Γ_{93}	$\omega \pi^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4 DESIG=32
Γ_{94}	$\omega \pi^0 \rightarrow \pi^+ \pi^- \pi^0$	$(1.7 \pm 0.8) \times 10^{-5}$	DESIG=327

Γ_{95}	$\phi\eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2	DESIG=38
Γ_{96}	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9	DESIG=41
Γ_{97}	$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$	$(2.59 \pm 0.34) \times 10^{-4}$		DESIG=236
Γ_{98}	$\phi f_0(980) \rightarrow \phi\pi^0\pi^0$	$(1.8 \pm 0.5) \times 10^{-4}$		DESIG=237
Γ_{99}	$\phi\eta\eta'$	$(2.32 \pm 0.17) \times 10^{-4}$		DESIG=387
Γ_{100}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$		DESIG=278
Γ_{101}	$\phi\pi^0 f_0(980) \rightarrow \phi\pi^0 p^0\pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$		DESIG=279
Γ_{102}	$\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$		DESIG=229
Γ_{103}	$\phi a_0(980)^0 \rightarrow \phi\eta\pi^0$	$(4.4 \pm 1.4) \times 10^{-6}$		DESIG=258
Γ_{104}	$\Xi(1530)^0 \bar{\Xi}^0$	$(3.2 \pm 1.4) \times 10^{-4}$		DESIG=108
Γ_{105}	$\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.})$	[b] $(3.1 \pm 0.5) \times 10^{-4}$		DESIG=68
Γ_{106}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$		DESIG=106
Γ_{107}	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$		DESIG=280
Γ_{108}	$\phi f_1(1285) \rightarrow \phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^0\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$		DESIG=281
Γ_{109}	$\eta\pi^+\pi^-$	$(3.8 \pm 0.7) \times 10^{-4}$		DESIG=239
Γ_{110}	$\eta\rho$	$(1.93 \pm 0.23) \times 10^{-4}$		DESIG=22
Γ_{111}	$\omega\eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$		DESIG=31
Γ_{112}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$		DESIG=150
Γ_{113}	$\rho\eta'(958)$	$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6	DESIG=23
Γ_{114}	$a_2(1320)^{\pm}\pi^{\mp}$	[b] $< 4.3 \times 10^{-3}$	CL=90%	DESIG=42
Γ_{115}	$K\bar{K}_2^*(1430)^+ \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%	DESIG=45
Γ_{116}	$K_1(1270)^{\pm} K^{\mp}$	$< 3.0 \times 10^{-3}$	CL=90%	DESIG=131
Γ_{117}	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$		DESIG=377
Γ_{118}	$K_S^0\pi^- K_2^*(1430)^+ + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$		DESIG=301
Γ_{119}	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3}$	CL=90%	DESIG=47
Γ_{120}	$\phi\pi^0$	$3 \times 10^{-6} \text{ or } 1 \times 10^{-7}$		DESIG=33; OUR EVAL; → UNCHECKED ←
Γ_{121}	$\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$		DESIG=128
Γ_{122}	$\omega f_2'(1525)$	$< 2.2 \times 10^{-4}$	CL=90%	DESIG=29
Γ_{123}	$\omega X(1835) \rightarrow \omega p\bar{p}$	$< 3.9 \times 10^{-6}$	CL=95%	DESIG=263
Γ_{124}	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	$< 6.2 \times 10^{-5}$		DESIG=386
Γ_{125}	$\phi X(1835) \rightarrow \phi p\bar{p}$	$< 2.1 \times 10^{-7}$	CL=90%	DESIG=291
Γ_{126}	$\phi X(1835) \rightarrow \phi\eta\pi^+\pi^-$	$< 2.8 \times 10^{-4}$	CL=90%	DESIG=288
Γ_{127}	$\phi X(1870) \rightarrow \phi\eta\pi^+\pi^-$	$< 6.13 \times 10^{-5}$	CL=90%	DESIG=289
Γ_{128}	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$		DESIG=287
Γ_{129}	$\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0$	$< 2.52 \times 10^{-4}$	CL=90%	DESIG=253
Γ_{130}	$\Sigma(1385)^0 \bar{\Lambda}^+ \text{c.c.}$	$< 8.2 \times 10^{-6}$	CL=90%	DESIG=111
Γ_{131}	$\Delta(1232)^+\bar{p}$	$< 1 \times 10^{-4}$	CL=90%	DESIG=112
Γ_{132}	$\Lambda(1520)\bar{\Lambda}^+ \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda}$	$< 4.1 \times 10^{-6}$	CL=90%	DESIG=260
Γ_{133}	$\bar{\Lambda}(1520)\Lambda^+ \text{c.c.}$	$< 1.80 \times 10^{-3}$	CL=90%	DESIG=364
Γ_{134}	$\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	$< 1.1 \times 10^{-5}$	CL=90%	DESIG=205
Γ_{135}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$< 2.1 \times 10^{-5}$	CL=90%	DESIG=206
Γ_{136}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$< 1.6 \times 10^{-5}$	CL=90%	DESIG=207
Γ_{137}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$< 5.6 \times 10^{-5}$	CL=90%	DESIG=208
Γ_{138}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$< 1.1 \times 10^{-5}$	CL=90%	DESIG=209

Decays into stable hadrons

			NODE=M070;CLUMP=B
Γ_{139}	$2(\pi^+\pi^-)\pi^0$	(3.73 \pm 0.32) %	S=1.4 DESIG=9
Γ_{140}	$3(\pi^+\pi^-)\pi^0$	(2.9 \pm 0.6) %	DESIG=11
Γ_{141}	$\pi^+\pi^-\pi^0$	(2.10 \pm 0.08) %	DESIG=7
Γ_{142}	$\pi^+\pi^-\pi^0\pi^0\pi^0$	(2.71 \pm 0.29) %	DESIG=358
Γ_{143}	$\rho^\pm\pi^\mp\pi^0\pi^0$	(1.41 \pm 0.22) %	DESIG=362
Γ_{144}	$\rho^+\rho^-\pi^0$	(6.0 \pm 1.1) $\times 10^{-3}$	DESIG=363
Γ_{145}	$\pi^+\pi^-\pi^0K^+K^-$	(1.20 \pm 0.30) %	DESIG=18
Γ_{146}	$4(\pi^+\pi^-)\pi^0$	(9.0 \pm 3.0) $\times 10^{-3}$	DESIG=12
Γ_{147}	$\pi^+\pi^-K^+K^-$	(6.84 \pm 0.32) $\times 10^{-3}$	DESIG=16
Γ_{148}	$\pi^+\pi^-K_S^0K_L^0$	(3.8 \pm 0.6) $\times 10^{-3}$	DESIG=296
Γ_{149}	$\pi^+\pi^-K_S^0K_S^0$	(1.68 \pm 0.19) $\times 10^{-3}$	DESIG=297
Γ_{150}	$\pi^\pm\pi^0K^\mp K_S^0$	(5.7 \pm 0.5) $\times 10^{-3}$	DESIG=341
Γ_{151}	$K^+K^-K_S^0K_S^0$	(4.1 \pm 0.8) $\times 10^{-4}$	DESIG=298
Γ_{152}	$\pi^+\pi^-K^+K^-\eta$	(4.7 \pm 0.7) $\times 10^{-3}$	DESIG=238
Γ_{153}	$\pi^0\pi^0K^+K^-$	(2.12 \pm 0.23) $\times 10^{-3}$	DESIG=234
Γ_{154}	$\pi^0\pi^0K_S^0K_L^0$	(1.9 \pm 0.4) $\times 10^{-3}$	DESIG=337
Γ_{155}	$K\bar{K}\pi$	(6.1 \pm 1.0) $\times 10^{-3}$	DESIG=15
Γ_{156}	$K^+K^-\pi^0$	(2.88 \pm 0.12) $\times 10^{-3}$	DESIG=334
Γ_{157}	$K_S^0K^\pm\pi^\mp$	(5.6 \pm 0.5) $\times 10^{-3}$	DESIG=335
Γ_{158}	$K_S^0K_L^0\pi^0$	(2.06 \pm 0.27) $\times 10^{-3}$	DESIG=336
Γ_{159}	$K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0$	(1.21 \pm 0.18) $\times 10^{-3}$	DESIG=339
Γ_{160}	$K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0$	(4.3 \pm 1.3) $\times 10^{-4}$	DESIG=338
Γ_{161}	$K_S^0K_L^0\eta$	(1.44 \pm 0.34) $\times 10^{-3}$	DESIG=340
Γ_{162}	$2(\pi^+\pi^-)$	(3.57 \pm 0.30) $\times 10^{-3}$	DESIG=8
Γ_{163}	$3(\pi^+\pi^-)$	(4.3 \pm 0.4) $\times 10^{-3}$	DESIG=10
Γ_{164}	$2(\pi^+\pi^-\pi^0)$	(1.61 \pm 0.21) %	DESIG=210
Γ_{165}	$2(\pi^+\pi^-)\eta$	(2.26 \pm 0.28) $\times 10^{-3}$	DESIG=201
Γ_{166}	$3(\pi^+\pi^-)\eta$	(7.2 \pm 1.5) $\times 10^{-4}$	DESIG=202
Γ_{167}	$\pi^+\pi^-\pi^0\pi^0\eta$	(2.3 \pm 0.5) $\times 10^{-3}$	DESIG=359
Γ_{168}	$\rho^\pm\pi^\mp\pi^0\eta$	(1.9 \pm 0.8) $\times 10^{-3}$	DESIG=361
Γ_{169}	$p\bar{p}$	(2.121 \pm 0.029) $\times 10^{-3}$	DESIG=50
Γ_{170}	$p\bar{p}\pi^0$	(1.19 \pm 0.08) $\times 10^{-3}$	S=1.1 DESIG=52
Γ_{171}	$p\bar{p}\pi^+\pi^-$	(6.0 \pm 0.5) $\times 10^{-3}$	DESIG=54
Γ_{172}	$p\bar{p}\pi^+\pi^-\pi^0$	[c] (2.3 \pm 0.9) $\times 10^{-3}$	S=1.9 DESIG=55
Γ_{173}	$p\bar{p}\eta$	(2.00 \pm 0.12) $\times 10^{-3}$	DESIG=56
Γ_{174}	$p\bar{p}\rho$	< 3.1 $\times 10^{-4}$	CL=90% DESIG=57
Γ_{175}	$p\bar{p}\omega$	(9.8 \pm 1.0) $\times 10^{-4}$	S=1.3 DESIG=58
Γ_{176}	$p\bar{p}\eta'(958)$	(1.29 \pm 0.14) $\times 10^{-4}$	S=2.0 DESIG=59
Γ_{177}	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	(6.8 \pm 1.8) $\times 10^{-5}$	DESIG=276
Γ_{178}	$p\bar{p}\phi$	(5.19 \pm 0.33) $\times 10^{-5}$	DESIG=127
Γ_{179}	$n\bar{n}$	(2.09 \pm 0.16) $\times 10^{-3}$	DESIG=64
Γ_{180}	$n\bar{n}\pi^+\pi^-$	(4 \pm 4) $\times 10^{-3}$	DESIG=65
Γ_{181}	$\Sigma^+\bar{\Sigma}^-$	(1.50 \pm 0.24) $\times 10^{-3}$	DESIG=247
Γ_{182}	$\Sigma^0\bar{\Sigma}^0$	(1.172 \pm 0.032) $\times 10^{-3}$	S=1.4 DESIG=63
Γ_{183}	$2(\pi^+\pi^-)K^+K^-$	(3.1 \pm 1.3) $\times 10^{-3}$	DESIG=17
Γ_{184}	$p\bar{n}\pi^-$	(2.12 \pm 0.09) $\times 10^{-3}$	DESIG=53
Γ_{185}	$nN(1440)$	seen	DESIG=215;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{186}	$nN(1520)$	seen	DESIG=216;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{187}	$nN(1535)$	seen	DESIG=217;OUR EST; \rightarrow UNCHECKED \leftarrow
Γ_{188}	$\Xi^-\bar{\Xi}^+$	(9.7 \pm 0.8) $\times 10^{-4}$	S=1.4 DESIG=62
Γ_{189}	$\Lambda\bar{\Lambda}$	(1.89 \pm 0.09) $\times 10^{-3}$	S=2.8 DESIG=60
Γ_{190}	$\Lambda\bar{\Sigma}^-\pi^+ (\text{or c.c.})$	[b] (8.3 \pm 0.7) $\times 10^{-4}$	S=1.2 DESIG=71
Γ_{191}	$pK^-\bar{\Lambda}+\text{c.c.}$	(8.7 \pm 1.1) $\times 10^{-4}$	DESIG=72
Γ_{192}	$2(K^+K^-)$	(7.2 \pm 0.8) $\times 10^{-4}$	DESIG=19
Γ_{193}	$pK^-\bar{\Sigma}^0$	(2.9 \pm 0.8) $\times 10^{-4}$	DESIG=73

Γ_{194}	$K^+ K^-$	$(2.86 \pm 0.21) \times 10^{-4}$	DESIG=13
Γ_{195}	$K_S^0 K_L^0$	$(1.95 \pm 0.11) \times 10^{-4}$	S=2.4 DESIG=75
Γ_{196}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$	DESIG=261
Γ_{197}	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$	DESIG=228
Γ_{198}	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$	DESIG=109
Γ_{199}	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$	DESIG=225
Γ_{200}	$\pi^+\pi^-$	$(1.47 \pm 0.14) \times 10^{-4}$	DESIG=6
Γ_{201}	$\Lambda\bar{\Sigma} + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$	DESIG=61
Γ_{202}	$K_S^0 K_S^0$	$< 1.4 \times 10^{-8}$	CL=95% DESIG=14

Radiative decays

Γ_{203}	3γ	$(1.16 \pm 0.22) \times 10^{-5}$	NODE=M070;CLUMP=C DESIG=81
Γ_{204}	4γ	$< 9 \times 10^{-6}$	CL=90% DESIG=244
Γ_{205}	5γ	$< 1.5 \times 10^{-5}$	CL=90% DESIG=245
Γ_{206}	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$	DESIG=283
Γ_{207}	$\gamma\eta\pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$	DESIG=292
Γ_{208}	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$	$< 2.5 \times 10^{-6}$	CL=95% DESIG=293
Γ_{209}	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$	$< 6.6 \times 10^{-6}$	CL=95% DESIG=294
Γ_{210}	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$	DESIG=378
Γ_{211}	$\gamma\eta_c(1S)$	$(1.7 \pm 0.4) \%$	S=1.5 DESIG=85
Γ_{212}	$\gamma\eta_c(1S) \rightarrow 3\gamma$	$(3.8 \pm 1.3) \times 10^{-6}$	S=1.1 DESIG=246
Γ_{213}	$\gamma\pi^+\pi^-2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	DESIG=99
Γ_{214}	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	DESIG=96
Γ_{215}	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$	DESIG=142
Γ_{216}	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	[d] $(2.8 \pm 0.6) \times 10^{-3}$	S=1.6 DESIG=89
Γ_{217}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8 DESIG=171
Γ_{218}	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$	DESIG=170
Γ_{219}	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95% DESIG=212
Γ_{220}	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	$< 2.63 \times 10^{-6}$	CL=90% DESIG=348
Γ_{221}	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	$< 1.86 \times 10^{-6}$	CL=90% DESIG=349
Γ_{222}	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$	DESIG=94
Γ_{223}	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	CL=90% DESIG=226
Γ_{224}	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90% DESIG=227
Γ_{225}	$\gamma\eta'(958)$	$(5.25 \pm 0.07) \times 10^{-3}$	S=1.3 DESIG=84
Γ_{226}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9 DESIG=95
Γ_{227}	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$	DESIG=203
Γ_{228}	$\gamma f_2(1270) f_2(1270) (\text{non resonant})$	$(8.2 \pm 1.9) \times 10^{-4}$	DESIG=204
Γ_{229}	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$	DESIG=143
Γ_{230}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$	DESIG=100
Γ_{231}	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$	DESIG=97
Γ_{232}	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3 DESIG=124
Γ_{233}	$\gamma f_2(1270)$	$(1.64 \pm 0.12) \times 10^{-3}$	S=1.3 DESIG=86
Γ_{234}	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm 0.60) \times 10^{-5}$	DESIG=373
Γ_{235}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$	DESIG=284
Γ_{236}	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$	DESIG=368
Γ_{237}	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm 0.24) \times 10^{-5}$	DESIG=369
Γ_{238}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm 1.0) \times 10^{-4}$	S=1.5 DESIG=91
Γ_{239}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$	DESIG=135
Γ_{240}	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$	DESIG=221
Γ_{241}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm 1.2) \times 10^{-4}$	DESIG=266
Γ_{242}	$\gamma\eta$	$(1.108 \pm 0.027) \times 10^{-3}$	DESIG=83
Γ_{243}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$	DESIG=175
Γ_{244}	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$	DESIG=88
Γ_{245}	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$	DESIG=141

Γ_{246}	$\gamma f'_2(1525)$	(5.7 \pm 0.8) $\times 10^{-4}$	S=1.5	DESIG=87
Γ_{247}	$\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0$	(8.0 \pm 0.7) $\times 10^{-5}$		DESIG=374
Γ_{248}	$\gamma f'_2(1525) \rightarrow \gamma \eta \eta$	(3.4 \pm 1.4) $\times 10^{-5}$		DESIG=268
Γ_{249}	$\gamma f_2(1640) \rightarrow \gamma \omega \omega$	(2.8 \pm 1.8) $\times 10^{-4}$		DESIG=222
Γ_{250}	$\gamma f_2(1910) \rightarrow \gamma \omega \omega$	(2.0 \pm 1.4) $\times 10^{-4}$		DESIG=223
Γ_{251}	$\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0$	(1.11 \pm 0.20) $\times 10^{-5}$		DESIG=370
Γ_{252}	$\gamma f_0(1800) \rightarrow \gamma \omega \phi$	(2.5 \pm 0.6) $\times 10^{-4}$		DESIG=262
Γ_{253}	$\gamma f_2(1810) \rightarrow \gamma \eta \eta$	(5.4 \pm 3.5) $\times 10^{-5}$		DESIG=269
Γ_{254}	$\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)$	(7.0 \pm 2.2) $\times 10^{-4}$		DESIG=144
Γ_{255}	$\gamma K^*(892) \bar{K}^*(892)$	(4.0 \pm 1.3) $\times 10^{-3}$		DESIG=145
Γ_{256}	$\gamma \phi \phi$	(4.0 \pm 1.2) $\times 10^{-4}$	S=2.1	DESIG=98
Γ_{257}	$\gamma p \bar{p}$	(3.8 \pm 1.0) $\times 10^{-4}$		DESIG=90
Γ_{258}	$\gamma \eta(2225)$	(3.14 \pm 0.50) $\times 10^{-4}$		DESIG=126
Γ_{259}	$\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0$	(1.3 \pm 0.9) $\times 10^{-4}$		DESIG=125
Γ_{260}	$\gamma \eta(1760) \rightarrow \gamma \omega \omega$	(1.98 \pm 0.33) $\times 10^{-3}$		DESIG=224
Γ_{261}	$\gamma \eta(1760) \rightarrow \gamma \gamma \gamma$	< 4.80 $\times 10^{-6}$	CL=90%	DESIG=347
Γ_{262}	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	(2.77 \pm 0.34) $\times 10^{-4}$	S=1.1	DESIG=213
Γ_{263}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	(7.7 \pm 1.5) $\times 10^{-5}$		DESIG=254
Γ_{264}	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	(3.3 \pm 2.0) $\times 10^{-5}$		DESIG=282
Γ_{265}	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$			DESIG=346
Γ_{266}	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	< 3.56 $\times 10^{-6}$	CL=90%	DESIG=350
Γ_{267}	$\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-)$	(2.4 \pm 0.7) $\times 10^{-5}$		DESIG=264
Γ_{268}	$\gamma(K \bar{K} \pi) [J^{PC} = 0^- +]$	(7 \pm 4) $\times 10^{-4}$	S=2.1	DESIG=176
Γ_{269}	$\gamma \pi^0$	(3.56 \pm 0.17) $\times 10^{-5}$		DESIG=82
Γ_{270}	$\gamma p \bar{p} \pi^+ \pi^-$	< 7.9 $\times 10^{-4}$	CL=90%	DESIG=93
Γ_{271}	$\gamma \Lambda \bar{\Lambda}$	< 1.3 $\times 10^{-4}$	CL=90%	DESIG=200
Γ_{272}	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	(1.13 \pm 0.60) $\times 10^{-4}$		DESIG=267
Γ_{273}	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	(6.2 \pm 1.0) $\times 10^{-4}$		DESIG=286
Γ_{274}	$\gamma f_0(2200)$			DESIG=123
Γ_{275}	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	(5.9 \pm 1.3) $\times 10^{-4}$		DESIG=285
Γ_{276}	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	(2.72 \pm 0.19) $\times 10^{-4}$		DESIG=371
Γ_{277}	$\gamma f_J(2220)$			DESIG=92
Γ_{278}	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	< 3.9 $\times 10^{-5}$	CL=90%	DESIG=136
Γ_{279}	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	< 4.1 $\times 10^{-5}$	CL=90%	DESIG=137
Γ_{280}	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	(1.5 \pm 0.8) $\times 10^{-5}$		DESIG=138
Γ_{281}	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	(4.9 \pm 0.7) $\times 10^{-5}$		DESIG=372
Γ_{282}	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	(5.6 \pm 2.4) $\times 10^{-5}$		DESIG=270
Γ_{283}	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	(5.5 \pm 4.0) $\times 10^{-5}$		DESIG=375
Γ_{284}	$\gamma f_0(1500) \rightarrow \gamma \pi \pi$	(1.09 \pm 0.24) $\times 10^{-4}$		DESIG=172
Γ_{285}	$\gamma f_0(1500) \rightarrow \gamma \eta \eta$	(1.7 \pm 0.6) $\times 10^{-5}$		DESIG=265
Γ_{286}	$\gamma A \rightarrow \gamma \text{invisible}$	[e] < 6.3 $\times 10^{-6}$	CL=90%	DESIG=251
Γ_{287}	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[f] < 5 $\times 10^{-6}$	CL=90%	DESIG=259

Dalitz decays

Γ_{288}	$\pi^0 e^+ e^-$	(7.6 \pm 1.4) $\times 10^{-7}$	NODE=M070;CLUMP=G	
Γ_{289}	$\eta e^+ e^-$	(1.43 \pm 0.07) $\times 10^{-5}$	DESIG=271	
Γ_{290}	$\eta'(958) e^+ e^-$	(6.59 \pm 0.18) $\times 10^{-5}$	DESIG=272	
Γ_{291}	$\eta U \rightarrow \eta e^+ e^-$	< 9.11 $\times 10^{-7}$	CL=90%	DESIG=273
Γ_{292}	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	< 2.0 $\times 10^{-7}$	CL=90%	DESIG=352
Γ_{293}	$\phi e^+ e^-$	< 1.2 $\times 10^{-7}$	CL=90%	DESIG=366
				DESIG=384

Weak decays

Γ_{294}	$D^- e^+ \nu_e + c.c.$	< 1.2	$\times 10^{-5}$	CL=90%
Γ_{295}	$\bar{D}^0 e^+ e^- + c.c.$	< 8.5	$\times 10^{-8}$	CL=90%
Γ_{296}	$D^- e^+ \nu_e + c.c.$	< 1.3	$\times 10^{-6}$	CL=90%
Γ_{297}	$D_s^* e^+ \nu_e + c.c.$	< 1.8	$\times 10^{-6}$	CL=90%
Γ_{298}	$D^- \pi^+ + c.c.$	< 7.5	$\times 10^{-5}$	CL=90%
Γ_{299}	$\bar{D}^0 \bar{K}^0 + c.c.$	< 1.7	$\times 10^{-4}$	CL=90%
Γ_{300}	$\bar{D}^0 \bar{K}^{*0} + c.c.$	< 2.5	$\times 10^{-6}$	CL=90%
Γ_{301}	$D_s^- \pi^+ + c.c.$	< 1.3	$\times 10^{-4}$	CL=90%
Γ_{302}	$D_s^- \rho^+ + c.c.$	< 1.3	$\times 10^{-5}$	CL=90%

NODE=M070;CLUMP=E

DESIG=218

DESIG=219

DESIG=220

DESIG=290

DESIG=241

DESIG=242

DESIG=275

DESIG=243

DESIG=274

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

Γ_{303}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%
Γ_{304}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%
Γ_{305}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%
Γ_{306}	$e^\pm \tau^\mp$	LF	< 8.3	$\times 10^{-6}$	CL=90%
Γ_{307}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%
Γ_{308}	$\Lambda_c^+ e^- + c.c.$		< 6.9	$\times 10^{-8}$	CL=90%

NODE=M070;CLUMP=D

DESIG=80

DESIG=277

DESIG=177

DESIG=178

DESIG=179

DESIG=379

Other decays

Γ_{309}	invisible	< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	-----	------------------	--------

NODE=M070;CLUMP=F

DESIG=240

[a] For $E_\gamma > 100$ MeV.

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

[c] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.[d] See the "Note on the $\eta(1405)$ " in the $\eta(1405)$ Particle Listings.[e] For a narrow state A with mass less than 960 MeV.[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

LINKAGE=EGM

LINKAGE=SG

LINKAGE=MF

LINKAGE=MG

LINKAGE=NSA

LINKAGE=NA0

J/ ψ (1S) PARTIAL WIDTHS **Γ (hadrons)**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
74.1 ± 8.1	BAI	95B	BES $e^+ e^-$	
59 ± 24	BALDINI...	75	FRAG $e^+ e^-$	
59 ± 14	BOYARSKI	75	MRK1 $e^+ e^-$	
50 ± 25	ESPOSITO	75B	FRAM $e^+ e^-$	

NODE=M070220

NODE=M070W3

NODE=M070W3

 $\Gamma(e^+ e^-)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5
5.53 ± 0.10 OUR AVERAGE					
5.550 ± 0.056 ± 0.089		1 ANASHIN	18A	KEDR $e^+ e^-$	
5.36 ± 0.29		2 HSUEH	92	RVUE See γ mini-review	

NODE=M070W1

NODE=M070W1

OCCUR=4

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.58 ± 0.05 ± 0.08		3 ABLIKIM	16Q	BES3 $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	
5.71 ± 0.16	13k	4 ADAMS	06A	CLEO $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	
5.57 ± 0.19	7.8k	4 AUBERT	04	BABR $e^+ e^- \rightarrow \mu^+ \mu^- \gamma$	
5.14 ± 0.39		BAI	95B	BES $e^+ e^-$	
4.72 ± 0.35		ALEXANDER	89	RVUE See γ mini-review	
4.4 ± 0.6		2 BRANDELIK	79C	DASP $e^+ e^-$	
4.6 ± 0.8		5 BALDINI...	75	FRAG $e^+ e^-$	
4.8 ± 0.6		BOYARSKI	75	MRK1 $e^+ e^-$	
4.6 ± 1.0		ESPOSITO	75B	FRAM $e^+ e^-$	

NODE=M070W1;LINKAGE=D

NODE=M070W1;LINKAGE=F

NODE=M070W1;LINKAGE=A

NODE=M070W1;LINKAGE=AA

NODE=M070W1;LINKAGE=B

1 From the cross sections of $e^+ e^- \rightarrow e^+ e^-$ and $e^+ e^- \rightarrow$ hadrons near the J/ ψ (1S) peak.2 From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.3 Using $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.4 Calculated by us from the reported values of $\Gamma(e^+ e^-) \times B(\mu^+ \mu^-)$ using $B(\mu^+ \mu^-) = (5.93 \pm 0.06)\%$.5 Assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$.

$\Gamma(\mu^+\mu^-)$				Γ_7
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.13 \pm 0.52	BAI	95B	BES e^+e^-	
4.8 \pm 0.6	BOYARSKI	75	MRK1 e^+e^-	
5 \pm 1	ESPOSITO	75B	FRAM e^+e^-	

$\Gamma(\gamma\gamma)$				Γ_{303}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79c	DASP e^+e^-

$J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel i in the e^+e^- annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_1\Gamma_5/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
4.884 \pm 0.048 \pm 0.078	¹ ANASHIN	18A	KEDR e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4 \pm 0.8	² BALDINI-...	75	FRAG e^+e^-	
3.9 \pm 0.8	² ESPOSITO	75B	FRAM e^+e^-	

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Data redundant with branching ratios or partial widths above.

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_5\Gamma_5/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
333.1 \pm 6.6 \pm 4.0	¹ ANASHIN	18A	KEDR e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
332.3 \pm 6.4 \pm 4.8	ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow e^+e^-$	
350 \pm 20	BRANDELIK	79c	DASP e^+e^-	
320 \pm 70	² BALDINI-...	75	FRAG e^+e^-	
340 \pm 90	² ESPOSITO	75B	FRAM e^+e^-	
360 \pm 100	² FORD	75	SPEC e^+e^-	

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Data redundant with branching ratios or partial widths above.

$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_7\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 \pm 4 OUR AVERAGE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
333.4 \pm 2.5 \pm 4.4	ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$	
331.8 \pm 5.2 \pm 6.3	ANASHIN	10	KEDR $3.097 e^+e^- \rightarrow \mu^+\mu^-$	
338.4 \pm 5.8 \pm 7.1	13k	ADAMS	06A CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$	
330.1 \pm 7.7 \pm 7.3	7.8k	AUBERT	04 BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
510 \pm 90	DASP	75	DASP e^+e^-	
380 \pm 50	¹ ESPOSITO	75B	FRAM e^+e^-	

¹ Data redundant with branching ratios or partial widths above.

$\Gamma(\rho(770)^{\mp} K^{\pm} K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{10}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4 \pm 1.0 \pm 1.9	130	LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^{\pm} \pi^{\mp} \pi^0 \gamma$

$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{23}\Gamma_5/\Gamma$
VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 \pm 0.3 \pm 0.2	170	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

NODE=M070W2
NODE=M070W2

NODE=M070W70
NODE=M070W70

NODE=M070225

NODE=M070225

NODE=M070G3
NODE=M070G3

NODE=M070G1
NODE=M070G1

OCCUR=2

NODE=M070G1;LINKAGE=A

NODE=M070G1;LINKAGE=S

NODE=M070G2
NODE=M070G2

NODE=M070G2;LINKAGE=S

NODE=M070G31
NODE=M070G31

NODE=M070G8
NODE=M070G8

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{24}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6±5.0±0.4	788	1 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 47.8 \pm 3.1 \pm 3.2 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{62}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.8±3.5±0.2	398	1 LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 24.8 \pm 1.8 \pm 2.5 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{26}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28±0.34±0.07	47±12	1 LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.28 \pm 0.40 \pm 0.11 \quad 25 \pm 8 \quad ^{1,2} \text{ AUBERT} \quad 07AK \text{ BABR} \quad 10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹ Dividing by $(2/3)^2$ to take twice into account that $B(K^{*0} \rightarrow K^+\pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

² Superseded by LEES 12F.

$\Gamma(K^*(892)^{\pm}K^*(892)^{\mp}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{27}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80±0.48±0.32	1 ± 5	1 LEES	14H BABR	$e^+e^- \rightarrow \pi^+\pi^-K_S^0K_S^0\gamma$

¹ Dividing by $(1/4)^2$ to take twice into account $B(K^*(892) \rightarrow K_S^0\pi) = 1/4$.

$\Gamma(K_S^0\pi^-K^*(892)^++\text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{29}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0±2.8 OUR AVERAGE				

9.2±1.2±3.2	64	1 LEES	17D BABR	$e^+e^- \rightarrow K_S^0K^{\pm}\pi^{\mp}\pi^0\gamma$
14.8±4.8±1.2	53	2 LEES	14H BABR	$e^+e^- \rightarrow \pi^+\pi^-K_S^0K_S^0\gamma$

¹ Dividing by 1/2 to take into account $B(K^*(892)^{\pm} \rightarrow K^{\pm}\pi^{\mp}) = 1/2$.

² Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0\pi) = 1/4$.

$\Gamma(K_S^0\pi^-K^*(892)^++\text{c.c.} \rightarrow K_S^0K_S^0\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{30}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7±1.2±0.3	53	LEES	14H BABR	$e^+e^- \rightarrow \pi^+\pi^-K_S^0K_S^0\gamma$

$\Gamma(K^*(892)^0\bar{K}_2^*(1430)^0+\text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{46}\Gamma_5/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.8±1.4±0.6	710	1,2,3 LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$33 \pm 4 \pm 1 \quad 317 \quad 2,4 \text{ AUBERT} \quad 07AK \text{ BABR} \quad 10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 12.89 \pm 0.54 \pm 0.41 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+\pi^-) = 2/3 B(K^{*0} \rightarrow K\pi)$.

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^0\bar{K}_2^*(1430)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K\pi)] = 16.4 \pm 1.1 \pm 1.4 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K\pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G24
NODE=M070G24

NODE=M070G24;LINKAGE=AU

NODE=M070P54
NODE=M070P54

NODE=M070P54;LINKAGE=A

NODE=M070G01
NODE=M070G01

NODE=M070G01;LINKAGE=AE

NODE=M070G01;LINKAGE=A

NODE=M070GY8
NODE=M070GY8

NODE=M070GY8;LINKAGE=A

NODE=M070GY4
NODE=M070GY4

NODE=M070GY4;LINKAGE=B

NODE=M070GY4;LINKAGE=A

NODE=M070GY5
NODE=M070GY5

NODE=M070G02

NODE=M070G02

NODE=M070G02;LINKAGE=A

NODE=M070G02;LINKAGE=AE

NODE=M070G02;LINKAGE=B

NODE=M070G02;LINKAGE=UB

$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{47} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070GY9 NODE=M070GY9
18.6 ± 16.1 ± 0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$		
¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$ $B(K^*(1430) \rightarrow K \pi)$.						
² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K^*(892)^+ K_2^*(1430)^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						
$\Gamma(K^*(892)^+ K_2^*(1430)^- + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{48} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070GZ0 NODE=M070GZ0
2.32 ± 2.00 ± 0.08	8 ± 8	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$		
¹ Dividing by $1/4$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.						
$\Gamma(K^*(892)^0 \bar{K}_2(1770)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{49} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G03 NODE=M070G03
3.8 ± 0.4 ± 0.3	110 ± 14	1 AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$		
¹ Dividing by $2/3$ to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.						
$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{53} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G18 NODE=M070G18
29.0 ± 1.7 ± 1.3		AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K^+ K^*(892)^- \gamma$		
$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{54} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G20 NODE=M070G20
10.96 ± 0.85 ± 0.70	155	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$		
$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}}$					Γ_{54} / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070P79 NODE=M070P79
2.69 ± 0.01 ± 0.13	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$		
$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{55} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G21 NODE=M070G21
16.76 ± 1.70 ± 1.00	89	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$		
$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{56} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G19 NODE=M070G19
26.6 ± 2.5 ± 1.5		AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$		
$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{57} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G22 NODE=M070G22
17.70 ± 1.70 ± 1.00	94	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$		
$\Gamma(K_S^0 K^*(892)^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$					Γ_{31} / Γ	
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070P71 NODE=M070P71
6.28 ± 0.16 ± 0.59		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$		
¹ Dividing by $1/6$ to account for $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$.						
$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{59} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G39 NODE=M070G39
42.6 ± 4.8 ± 7.2	99	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$		
¹ Dividing by $1/6$ to account for $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$.						
$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$					$\Gamma_{60} \Gamma_5 / \Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G32 NODE=M070G32
22.8 ± 2.8 ± 6.8	80	1 LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$		
¹ Dividing by $1/4$ to account for $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$.						

$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_{32}/Γ	NODE=M070P76 NODE=M070P76
$\text{VALUE (units } 10^{-4}\text{)}$	EVTS	DOCUMENT ID
$2.69 \pm 0.04 \pm 0.25$	183k	ABLIKIM
		19AQ BES
		$J/\psi \rightarrow K^+ K^- \pi^0$
$\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_{33}/Γ	NODE=M070P77 NODE=M070P77
$\text{VALUE (units } 10^{-5}\text{)}$	EVTS	DOCUMENT ID
$1.1 \pm 0.1 \pm 0.6$	183k	ABLIKIM
		19AQ BES
		$J/\psi \rightarrow K^+ K^- \pi^0$
$\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_{34}/Γ	NODE=M070P78 NODE=M070P78
$\text{VALUE (units } 10^{-6}\text{)}$	EVTS	DOCUMENT ID
$6.2 \pm 0.7 \pm 2.8$	183k	ABLIKIM
		19AQ BES
		$J/\psi \rightarrow K^+ K^- \pi^0$
$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{61}\Gamma_5/\Gamma$	NODE=M070G33 NODE=M070G33
VALUE (eV)	EVTS	DOCUMENT ID
$3.60 \pm 0.75 \pm 2.25$	34	¹ LEES
		17D BABR
		$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
¹ Dividing by 2/3 to account for $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$.		
$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{67}\Gamma_5/\Gamma$	NODE=M070G38 NODE=M070G38
VALUE (eV)	EVTS	DOCUMENT ID
$7.3 \pm 1.4 \pm 0.4$	44	LEES
		17D BABR
		$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
$\Gamma(\omega K\bar{K}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{69}\Gamma_5/\Gamma$	NODE=M070G29 NODE=M070G29
VALUE (eV)	EVTS	DOCUMENT ID
$3.70 \pm 1.98 \pm 0.03$	24	¹ AUBERT
		07AU BABR
		$10.6 e^+ e^- \rightarrow \omega K^+ K^- \gamma$
¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 3.3 \pm 1.3 \pm 1.2 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{71}\Gamma_5/\Gamma$	NODE=M070G10 NODE=M070G10
$\text{VALUE (10}^{-2} \text{ keV)}$	EVTS	DOCUMENT ID
$0.95 \pm 0.19 \pm 0.01$	35	¹ AUBERT
		06D BABR
		$10.6 e^+ e^- \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+ \pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2} \text{ keV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{76}\Gamma_5/\Gamma$	NODE=M070GZ1 NODE=M070GZ1
VALUE (eV)	EVTS	DOCUMENT ID
$3.25 \pm 0.84 \pm 0.03$	29	¹ LEES
		14H BABR
		$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
¹ LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{78}\Gamma_5/\Gamma$	NODE=M070G09 NODE=M070G09
VALUE (eV)	EVTS	DOCUMENT ID
$4.59 \pm 0.62 \pm 0.05$	163	¹ LEES
		12F BABR
		$10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{79}\Gamma_5/\Gamma$	NODE=M070G07 NODE=M070G07
VALUE (eV)	EVTS	DOCUMENT ID
$1.79 \pm 0.32 \pm 0.02$	61	^{1,2,3} LEES
		12F BABR
		$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
$4.08 \pm 0.73 \pm 0.04$	44	^{2,4} AUBERT
		07AK BABR
		$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

- ¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 1.51 \pm 0.25 \pm 0.10$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2 \pm 2.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.
- ³ Using $\pi^+ \pi^-$ invariant mass between 1.1 and 1.5 GeV. May include other sources such as $f_0(1370)$.
- ⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = 3.44 \pm 0.55 \pm 0.28$ eV which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2 \pm 2.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- f'_2(1525)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{83}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

- ¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$.
- ² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{84}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.1±3.2±0.2	11	1,2 LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

- ¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$ and using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.
- ² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 7.2 \pm 2.8 \pm 0.3$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{85}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.47±0.35 OUR AVERAGE				

4.45±0.49±0.05	181	¹ LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
4.50±0.48±0.05	254 ± 23	² SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.3 ± 0.7 ± 0.1	103	³ AUBERT,BE 06D BABR	10.6	$e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
-----------------	-----	---------------------------------	------	--

- ¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)]$ assuming $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

- ³ Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_{86}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.76±0.57±0.03	45	¹ LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.13±0.88±0.03	23	² AUBERT,BE 06D BABR	10.6	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
----------------	----	---------------------------------	------	--

NODE=M070G07;LINKAGE=A

NODE=M070G07;LINKAGE=AE

NODE=M070G07;LINKAGE=B

NODE=M070G07;LINKAGE=UB

NODE=M070GZ4

NODE=M070GZ4

NODE=M070GZ4;LINKAGE=A

NODE=M070GZ4;LINKAGE=B

NODE=M070GZ2

NODE=M070GZ2

NODE=M070GZ2;LINKAGE=A

NODE=M070GZ2;LINKAGE=B

NODE=M070G14

NODE=M070G14

NODE=M070G14;LINKAGE=B

NODE=M070G14;LINKAGE=SH

NODE=M070G14;LINKAGE=AU

NODE=M070G15

NODE=M070G15

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{89}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
6.1±2.7±0.4	6	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$	
¹ AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05 \text{ eV}$.					

NODE=M070G15;LINKAGE=A

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{97}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.44±0.19 OUR AVERAGE					
1.40±0.25±0.02 57±9 ¹ LEES 12F BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$					
1.48±0.27±0.09 60±11 ² SHEN 09 BELL $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.02±0.24±0.01 20±5 ³ AUBERT 07AK BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$					
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² Multiplied by 2/3 to take into account the $\phi\pi^+\pi^-$ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.					
³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M070G28

NODE=M070G28

NODE=M070G28;LINKAGE=AU

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{98}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.98±0.26±0.01	16±4	¹ LEES 12F BABR	$10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.95±0.40±0.01 7.0±2.8 ² AUBERT 07AK BABR $10.6 e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$					
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.48 \pm 0.12 \pm 0.05 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.47 \pm 0.19 \pm 0.05 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M070G06

NODE=M070G06

NODE=M070G06;LINKAGE=A

$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{109}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.3 ±0.4 OUR AVERAGE					
2.34±0.43±0.16 49 LEES 18 BABR $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$					
2.23±0.97±0.03 9 ¹ AUBERT 07AU BABR $10.6 e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$					
¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.92 \pm 0.28) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M070G25

NODE=M070G25

NODE=M070G25;LINKAGE=AU

$\Gamma(K_S^0\pi^- K_2^*(1430)^+ + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{118}\Gamma_5/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
20.1±9.8±0.5	35	^{1,2} LEES 14H BABR	$e^+e^- \rightarrow \pi^+\pi^-K_S^0K_2^0\gamma$		

NODE=M070GY6

NODE=M070GY6

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$.

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_S^0 \pi^- K_2^*(1430)^+ + c.c.) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)] / [\Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8 \text{ eV}$ which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{139} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303 ± 5 ± 18	4990	AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$

NODE=M070G23
NODE=M070G23

$\Gamma(\pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{141} \Gamma_5 / \Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.122 ± 0.005 ± 0.008		AUBERT,B	04N BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$

NODE=M070G5
NODE=M070G5

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{142} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
150.0 ± 4.0 ± 15.0	2.3k	LEES	18E BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

NODE=M070P53
NODE=M070P53

$\Gamma(\rho^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{143} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0 ± 9.0 ± 8.0	1.2k	LEES	18E BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

NODE=M070P55
NODE=M070P55

$\Gamma(\rho^+ \rho^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{144} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0 ± 5.0 ± 3.3	529	LEES	18E BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

NODE=M070P56
NODE=M070P56

$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{145} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0 ± 4.3 ± 6.4	768	AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$

NODE=M070G27
NODE=M070G27

$\Gamma(\pi^+ \pi^- K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{147} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
37.94 ± 0.81 ± 1.10	3.1k	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ± 1.3 ± 2.1	1.5k	¹ AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
33.6 ± 2.7 ± 2.7	233	² AUBERT	05D BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

¹ Superseded by LEES 12F.

² Superseded by AUBERT 07AK.

NODE=M070G12;LINKAGE=B

NODE=M070G12;LINKAGE=AU

$\Gamma(\pi^+ \pi^- K_S^0 K_L^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{148} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.8 ± 2.3 ± 2.1	248	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$

NODE=M070G12

NODE=M070G12

OCCUR=2

$\Gamma(\pi^+ \pi^- K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{149} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3 ± 0.9 ± 0.5	133	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GY2

NODE=M070GY2

$\Gamma(\pi^\pm \pi^0 K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{150} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7 ± 1.9 ± 1.8	393	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 \pi^\pm \pi^0 \gamma$

NODE=M070G34

NODE=M070G34

$\Gamma(K^+ K^- K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{151} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 ± 0.1	29	LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

NODE=M070GY3

NODE=M070GY3

$\Gamma(\pi^+ \pi^- K^+ K^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{152} \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9 ± 3.9 ± 0.1	73	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

NODE=M070G30


NODE=M070G30

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- K^+ K^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-)] / [\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G30;LINKAGE=AU

$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{153}\Gamma_5/\Gamma$	NODE=M070G04 NODE=M070G04
11.75±0.81±0.90 388 LEES 12F BABR 10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$		
13.6 ± 1.1 ± 1.3 203 ¹ AUBERT 07AK BABR 10.6 $e^+e^- \rightarrow \pi^0\pi^0K^+K^-\gamma$		
¹ Superseded by LEES 12F.		
$\Gamma(\pi^0\pi^0K_S^0K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{154}\Gamma_5/\Gamma$	NODE=M070G40 NODE=M070G40
10.3±2.3±0.5 47 LEES 17A BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\pi^0\gamma$		
$\Gamma(K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{158}\Gamma_5/\Gamma$	NODE=M070G41 NODE=M070G41
11.4±1.3±0.6 182 LEES 17A BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$		
$\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{159}\Gamma_5/\Gamma$	NODE=M070G42 NODE=M070G42
6.7±0.9±0.4 106 LEES 17A BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$		
$\Gamma(K_2^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{160}\Gamma_5/\Gamma$	NODE=M070G43 NODE=M070G43
2.4±0.7±0.1 37 LEES 17A BABR $e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$		
$\Gamma(K_S^0K_L^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{161}\Gamma_5/\Gamma$	NODE=M070G35 NODE=M070G35
8.0±1.8±0.4 45 LEES 17A BABR $e^+e^- \rightarrow K_S^0K_L^0\eta\gamma$		
$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{162}\Gamma_5/\Gamma$	NODE=M070G11 NODE=M070G11
20.4±0.9±0.4 LEES 12E BABR 10.6 $e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$		
19.5 ± 1.4 ± 1.3 270 ¹ AUBERT 05D BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$		
¹ Superseded by LEES 12E.		
$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{163}\Gamma_5/\Gamma$	NODE=M070G6 NODE=M070G6
2.37±0.16±0.14 496 AUBERT 06D BABR 10.6 $e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$		
$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{164}\Gamma_5/\Gamma$	NODE=M070G7 NODE=M070G7
8.9±0.5±1.0 761 AUBERT 06D BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$		
$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{165}\Gamma_5/\Gamma$	NODE=M070G26 NODE=M070G26
13.1±2.4±0.1 85 ¹ AUBERT 07AU BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$		
1 AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 5.16 \pm 0.85 \pm 0.39$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{167}\Gamma_5/\Gamma$	NODE=M070P57 NODE=M070P57
12.8±1.8±2.0 203 LEES 18E BABR $\pi^+\pi^-\pi^0\pi^0\eta\gamma$		
$\Gamma(\omega\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{63}\Gamma_5/\Gamma$	NODE=M070P58 NODE=M070P58
1.90±0.96±0.01 27 ¹ LEES 18E BABR $\pi^+\pi^-\pi^0\pi^0\eta\gamma$		
1 LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		

$\Gamma(\rho^\pm \pi^\mp \pi^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{168} \Gamma_5/\Gamma$	NODE=M070P59 NODE=M070P59			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$10.5 \pm 4.1 \pm 1.6$	168	LEES	18E	BABR	$10.6 \frac{e^+ e^-}{\pi^+ \pi^-} \rightarrow \pi^0 \pi^0 \eta \gamma$
$\Gamma(p\bar{p}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{169} \Gamma_5/\Gamma$	NODE=M070G4 NODE=M070G4			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
11.9 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.				
$11.3 \pm 0.4 \pm 0.3$	821	¹ LEES	130	BABR	$e^+ e^- \rightarrow p\bar{p}\gamma$
$12.9 \pm 0.4 \pm 0.4$	918	² LEES	13Y	BABR	$e^+ e^- \rightarrow p\bar{p}\gamma$
9.7 ± 1.7		³ ARMSTRONG	93B	E760	$\bar{p}p \rightarrow e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$12.0 \pm 0.6 \pm 0.5$	438	⁴ AUBERT	06B	BABR	$e^+ e^- \rightarrow p\bar{p}\gamma$
¹ ISR photon reconstructed in the detector					
² ISR photon undetected					
³ Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.					
⁴ Superseded by LEES 130					



$\Gamma(\Sigma^0 \bar{\Sigma}^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{182}\Gamma_5/\Gamma$	NODE=M070G17		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	NODE=M070G17
6.4±1.2±0.6	AUBERT	07BD BABR	$10.6 \text{ e}^+ \text{ e}^- \rightarrow \Sigma^0 \bar{\Sigma}^0 \gamma$	

$\Gamma(2(\pi^+\pi^-)K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{183}\Gamma_5/\Gamma$	NODE=M070G9			
VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.75±0.23±0.17	205	AUBERT	06D	BABR	$10.6 \frac{e^+e^-}{K^+K^- 2(\pi^+\pi^-)\gamma}$

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{189}\Gamma_5/\Gamma$	NODE=M070G16		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	NODE=M070G16
10.7±0.9±0.7	AUBERT	07BD BABR	$10.6 \text{ e}^+\text{e}^- \rightarrow \Lambda\bar{\Lambda}\gamma$	

$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_{192}\Gamma_5/\Gamma$	NODE=M070G13				
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070G13
4.00±0.33±0.29	287 ± 24	LEES	12F	BABR	$10.6 e^+e^- \rightarrow 2(K^+K^-)\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
4.11±0.39±0.30	156 ± 15	1 AUBERT	07AK	BABR	$10.6 e^+e^- \rightarrow 2(K^+K^-)\gamma$	
4.0 ± 0.7 ± 0.6	38	2 AUBERT	05D	BABR	$10.6 e^+e^- \rightarrow 2(K^+K^-)\gamma$	

¹Superseded by LEFS 12E

² Superseded by ALBERT 07AK

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{194}\Gamma_5/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.78 \pm 0.11 \pm 0.05	462	¹ LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$	
1.94 \pm 0.11 \pm 0.05	462	² LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$	
1.42 \pm 0.23 \pm 0.08	51	³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$	
1 $\sin\phi > 0$.					
2 $\sin\phi < 0$.					
3 Interference with non-resonant K^+K^- production not taken into account.					

 $J/\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ above.

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.877 \pm 0.005 OUR AVERAGE				
0.878 \pm 0.005	BAI	95B	BES e^+e^-	
0.86 \pm 0.02	BOYARSKI	75	MRK1 e^+e^-	

 $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.135 \pm 0.003				
1,2 SETH	04	RVUE	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17 \pm 0.02	¹ BOYARSKI	75	MRK1 e^+e^-	

1 Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.2 Using $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. $\Gamma(ggg)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
64.1 \pm 1.0					
6 M	¹ BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^-\gamma + \text{hadrons}$	
1 Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+\ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma \eta_C)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.					

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
8.79 \pm 1.05					
200 k	¹ BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^-\gamma + \text{hadrons}$	
1 Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 08.					

 $\Gamma(ggg)/\Gamma(ggg)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3
13.7 \pm 0.1 \pm 0.7					
6 M	BESSON	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
5.971 \pm 0.032 OUR AVERAGE					
5.983 \pm 0.007 \pm 0.037	720k	ABLIKIM	13R	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	
5.945 \pm 0.067 \pm 0.042	15k	LI	05C	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	
5.90 \pm 0.05 \pm 0.10		BAI	98D	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	
6.09 \pm 0.33		BAI	95B	e^+e^-	
5.92 \pm 0.15 \pm 0.20		COFFMAN	92	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	
6.9 \pm 0.9		BOYARSKI	75	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$	

 $\Gamma(e^+e^-\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
8.8 \pm 1.3 \pm 0.4					
¹ ARMSTRONG	96	E760		$\bar{p}p \rightarrow e^+e^-\gamma$	

1 For $E_\gamma > 100$ MeV.

NODE=M070G08

NODE=M070G08

OCCUR=2

NODE=M070230

NODE=M070300

NODE=M070R3

NODE=M070R3

NODE=M070R4

NODE=M070R4

NODE=M070R4;LINKAGE=C

NODE=M070R4;LINKAGE=SE

NODE=M070S65

NODE=M070S65

NODE=M070S65;LINKAGE=BE

NODE=M070S66

NODE=M070S66

NODE=M070S66;LINKAGE=BE

NODE=M070S67

NODE=M070S67

NODE=M070R1

NODE=M070R1

NODE=M070S33

NODE=M070S33

NODE=M070S33;LINKAGE=A

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_7/Γ
5.961±0.033 OUR AVERAGE					
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
5.960±0.065±0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
5.84 ± 0.06 ± 0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
6.08 ± 0.33		BAI	95B	BES $e^+ e^-$	
5.90 ± 0.15 ± 0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
6.9 ± 0.9		BOYARSKI	75	MRK1 $e^+ e^-$	

 $\Gamma(e^+e^-)/\Gamma(\mu^+\mu^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_5/Γ_7
1.0016±0.0031 OUR AVERAGE				
1.0022±0.0044±0.0048	¹ AULCHENKO 14	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$	
1.0017±0.0017±0.0033	² ABLIKIM 13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
1.002 ± 0.021 ± 0.013	³ ANASHIN 10	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$	
0.997 ± 0.012 ± 0.006	LI 05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.011 ± 0.013 ± 0.016	BAI 98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
1.00 ± 0.07	BAI 95B	BES	$e^+ e^-$	
1.00 ± 0.05	BOYARSKI 75	MRK1	$e^+ e^-$	
0.91 ± 0.15	ESPOSITO 75B	FRAM	$e^+ e^-$	
0.93 ± 0.10	FORD 75	SPEC	$e^+ e^-$	

¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.² Not independent of the corresponding measurements of $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$.³ Not independent of the corresponding measurements of $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$.**HADRONIC DECAYS** $\Gamma(\rho\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_8/Γ
1.69 ± 0.15 OUR AVERAGE					
2.18 ± 0.19		1,2 AUBERT,B 04N	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$	
2.184±0.005±0.201	220k	2,3 BAI	04H	BES $e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$	
2.091±0.021±0.116		2,4 BAI	04H	BES $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
1.21 ± 0.20		BAI 96D	BES	$e^+ e^- \rightarrow \rho \pi$	
1.42 ± 0.01 ± 0.19		COFFMAN 88	MRK3	$e^+ e^-$	
1.3 ± 0.3	150	FRANKLIN 83	MRK2	$e^+ e^-$	
1.6 ± 0.4	183	ALEXANDER 78	PLUT	$e^+ e^-$	
1.33 ± 0.21		BRANDELIK 78B	DASP	$e^+ e^-$	
1.0 ± 0.2	543	BARTEL 76	CNTR	$e^+ e^-$	
1.3 ± 0.3	153	JEAN-MARIE 76	MRK1	$e^+ e^-$	

¹ From the ratio of $\Gamma(e^+ e^-) B(\pi^+ \pi^- \pi^0)$ and $\Gamma(e^+ e^-) B(\mu^+ \mu^-)$ (AUBERT 04).² Not independent of their $B(\pi^+ \pi^- \pi^0)$.³ From $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ events directly.⁴ Obtained comparing the rates for $\pi^+ \pi^- \pi^0$ and $\mu^+ \mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$ and with $B(J/\psi \rightarrow \mu^+ \mu^-) = 5.88 \pm 0.10\%$.NODE=M070R2
NODE=M070R2NODE=M070R5
NODE=M070R5NODE=M070R5;LINKAGE=A
NODE=M070R5;LINKAGE=AB


NODE=M070R5;LINKAGE=AN

NODE=M070305

NODE=M070R20
NODE=M070R20

OCCUR=2

NODE=M070R20;LINKAGE=AU
NODE=M070R20;LINKAGE=BU
NODE=M070R20;LINKAGE=BA
NODE=M070R20;LINKAGE=BI

 $\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_{141}
1.142±0.011±0.026	20K	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

1.331±0.033 20K 2 LEES 17C BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

1 From a Dalitz plot analysis in an isobar model.

2 From a Dalitz plot analysis in a Veneziano model.

NODE=M070P18
NODE=M070P18

OCCUR=2

NODE=M070P18;LINKAGE=A
NODE=M070P18;LINKAGE=B $\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_8
0.328±0.005±0.027	COFFMAN 88	MRK3	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.35 ± 0.08 ALEXANDER 78 PLUT e^+e^-
0.32 ± 0.08 BRANDELIK 78B DASP e^+e^-
0.39 ± 0.11 BARTEL 76 CNTR e^+e^-
0.37 ± 0.09 JEAN-MARIE 76 MRK1 e^+e^- NODE=M070R21
NODE=M070R21 $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ_{141}
10.9 ± 1.7 ± 2.7	20K	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

0.80±0.27 20K 2 LEES 17C BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

1 From a Dalitz plot analysis in an isobar model.

2 From a Dalitz plot analysis in a Veneziano model.

NODE=M070P25
NODE=M070P25

OCCUR=2

NODE=M070P25;LINKAGE=A
NODE=M070P25;LINKAGE=B $\Gamma(\rho(1450)^{\pm}\pi^{\mp} \rightarrow K_S^0 K^{\pm}\pi^{\mp})/\Gamma(K_S^0 K^{\pm}\pi^{\mp})$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_{157}
6.3±0.8±0.6	4K	1 LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^{\pm}\pi^{\mp}$	

1 From a Dalitz plot analysis in an isobar model.

NODE=M070P31
NODE=M070P31

NODE=M070P31;LINKAGE=A

 $\Gamma(\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0)/\Gamma(K^+K^-\pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ_{156}
9.3±2.0±0.6	2K	1 LEES	17C BABR	$J/\psi \rightarrow K^+K^-\pi^0$	

1 From a Dalitz plot analysis in an isobar model.

NODE=M070P27
NODE=M070P27

NODE=M070P27;LINKAGE=A

 $\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+\pi^-\eta'(958))/\Gamma_{\text{total}}$


VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
3.28±0.55±0.44	119	1 ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+\pi^-\eta'$	

1 From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.NODE=M070P36
NODE=M070P36

NODE=M070P36;LINKAGE=A

$\Gamma(\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{17}/Γ_{141}	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
8±2±5	20K	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
22±6	20K	2 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$		
1 From a Dalitz plot analysis in an isobar model. 2 From a Dalitz plot analysis in a Veneziano model.						
$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{19}/Γ_{141}	
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
4± 1±20	20K	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
600±250	20K	2 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$		
1 From a Dalitz plot analysis in an isobar model. 2 From a Dalitz plot analysis in a Veneziano model.						
$\Gamma(\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{20}/Γ_{141}	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
4.0±0.8	20K	1 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$		
1 From a Dalitz plot analysis in a Veneziano model.						
$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$					Γ_{21}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
10.9±2.2 OUR AVERAGE						
11.7±0.7±2.5	7584	AUGUSTIN	89 DM2	$J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$		
8.4±4.5	36	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$		
$\Gamma(\omega\pi^+\pi^-\pi^-\pi^+)/\Gamma_{\text{total}}$					Γ_{22}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
85±34	140	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$		
$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{24}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
7.2±1.0 OUR AVERAGE						
7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$		
7.8±1.6	215	BURMESTER	77D PLUT	e^+e^-		
6.8±1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$		
$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{74}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.12±0.02±0.13	14k	1 ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$		
1 Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.						
$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$					Γ_{25}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
4.3±0.6 OUR AVERAGE						
4.3±0.2±0.6	5860	AUGUSTIN	89 DM2	e^+e^-		
4.0±1.6	70	BURMESTER	77D PLUT	e^+e^-		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.9±0.8	81	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$		
$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}}$					Γ_{26}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<5	90	VANNUCCI	77 MRK1	$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$		
$\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$					Γ_{27}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.00±0.19±0.11	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0\pi^\mp\pi^0$		

$\Gamma(K^*(892)^{\pm} K^*(700)^{\mp})/\Gamma_{\text{total}}$	Γ_{28}/Γ	NODE=M070S74 NODE=M070S74
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$1.09 \pm 0.18 \pm 0.94$	655	ABLIKIM
		10E BES2
		$J/\psi \rightarrow K^{\pm} K_S^0 \pi^{\mp} \pi^0$
$\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$	Γ_{35}/Γ	NODE=M070S69 NODE=M070S69
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$1.15 \pm 0.13 \pm 0.22$	209	ABLIKIM
		10C BES2
		$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$
$\Gamma(K^*(1410) \bar{K} + c.c. \rightarrow K^{\pm} K^{\mp} \pi^0)/\Gamma(K^+ K^- \pi^0)$	Γ_{41}/Γ_{156}	NODE=M070P28 NODE=M070P28
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$2.3 \pm 1.1 \pm 0.7$	2K	¹ LEES
		17C BABR
		$J/\psi \rightarrow K^+ K^- \pi^0$
1 From a Dalitz plot analysis in an isobar model.		
$\Gamma(K^*(1410) \bar{K} + c.c. \rightarrow K_S^0 K^{\pm} \pi^{\mp})/\Gamma(K_S^0 K^{\pm} \pi^{\mp})$	Γ_{42}/Γ_{157}	NODE=M070P32 NODE=M070P32
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$1.5 \pm 0.5 \pm 0.9$	4K	¹ LEES
		17C BABR
		$J/\psi \rightarrow K_S^0 K^{\pm} \pi^{\mp}$
1 From a Dalitz plot analysis in an isobar model.		
$\Gamma(K_2^*(1430) \bar{K} + c.c. \rightarrow K^{\pm} K^{\mp} \pi^0)/\Gamma(K^+ K^- \pi^0)$	Γ_{44}/Γ_{156}	NODE=M070P29 NODE=M070P29
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$3.5 \pm 1.3 \pm 0.9$	2K	¹ LEES
		17C BABR
		$J/\psi \rightarrow K^+ K^- \pi^0$
1 From a Dalitz plot analysis in an isobar model.		
$\Gamma(K_2^*(1430) \bar{K} + c.c. \rightarrow K_S^0 K^{\pm} \pi^{\mp})/\Gamma(K_S^0 K^{\pm} \pi^{\mp})$	Γ_{45}/Γ_{157}	NODE=M070P33 NODE=M070P33
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$7.1 \pm 1.3 \pm 1.2$	4K	¹ LEES
		17C BABR
		$J/\psi \rightarrow K_S^0 K^{\pm} \pi^{\mp}$
1 From a Dalitz plot analysis in an isobar model.		
$\Gamma(K^*(892)^0 \bar{K}^*(1430)^0 + c.c.)/\Gamma_{\text{total}}$	Γ_{46}/Γ	NODE=M070R48 NODE=M070R48
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
6.7 ± 2.6	40	VANNUCCI 77
		MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
$\Gamma(\omega K^*(892) \bar{K} + c.c.)/\Gamma_{\text{total}}$	Γ_{50}/Γ	NODE=M070S2 NODE=M070S2
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
61 ± 9 OUR AVERAGE		
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM
53 ± 14 ± 14	530 ± 140	BECKER
		87 MRK3 $e^+ e^- \rightarrow \text{hadrons}$
$\Gamma(\bar{K} K^*(892) + c.c. \rightarrow K_S^0 K^{\pm} \pi^{\mp})/\Gamma(K_S^0 K^{\pm} \pi^{\mp})$	Γ_{52}/Γ_{157}	NODE=M070P30 NODE=M070P30
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$90.5 \pm 0.9 \pm 3.8$	4K	¹ LEES
		17C BABR
		$J/\psi \rightarrow K_S^0 K^{\pm} \pi^{\mp}$
1 From a Dalitz plot analysis in an isobar model.		
$\Gamma(K^+ K^*(892)^- + c.c.)/\Gamma_{\text{total}}$	Γ_{53}/Γ	NODE=M070S15 NODE=M070S15
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$6.0^{+0.8}_{-1.0}$ OUR AVERAGE	Error includes scale factor of 2.9. See the ideogram below. [(5.0 ± 0.4) × 10^{-3} OUR 2019 AVERAGE]	
8.07 ± 0.04 ± 0.38	183k	ABLIKIM
4.57 ± 0.17 ± 0.70	2285	JOUSSET
5.26 ± 0.13 ± 0.53		COFFMAN
		88 MRK3 $J/\psi \rightarrow K^{\pm} K_S^0 \pi^{\mp}, K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
2.6 ± 0.6	24	FRANKLIN
3.2 ± 0.6	48	VANNUCCI
4.1 ± 1.2	39	BRAUNSCH...
		76 DASP $J/\psi \rightarrow K^{\pm} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
OCCUR=2		
NEW		



$\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{54}/Γ_{156}
VALUE (%) $92.4 \pm 1.5 \pm 3.4$	EVTS 2K DOCUMENT ID 1 LEES TECN BABR COMMENT $J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{56}/Γ
VALUE (units 10^{-3}) 4.2 ± 0.4 OUR AVERAGE	EVTS DOCUMENT ID TECN COMMENT

3.96±0.15±0.60 1192 JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$
4.33±0.12±0.45 COFFMAN 88 MRK3 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6 45 VANNUCCI 77 MRK1 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$	Γ_{58}/Γ
VALUE (units 10^{-3}) 3.8 ± 0.8 ± 1.2	DOCUMENT ID TECN COMMENT

3.8±0.8±1.2 1 BAI 99C BES $e^+ e^-$

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{59}/Γ
VALUE seen	DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1 ABLIKIM 06C BES2 $J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

1 A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

$\Gamma(\omega \pi^0 \pi^0)/\Gamma_{\text{total}}$	Γ_{62}/Γ
VALUE (units 10^{-3}) 3.4 ± 0.3 ± 0.7	EVTS DOCUMENT ID TECN COMMENT

3.4±0.3±0.7 509 AUGUSTIN 89 DM2 $J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$	Γ_{64}/Γ
VALUE (units 10^{-4}) 30 ± 5 OUR AVERAGE	EVTS DOCUMENT ID TECN COMMENT

31±6 4600 AUGUSTIN 89 DM2 $J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
29±7 87 BURMESTER 77D PLUT $e^+ e^-$

$\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$	Γ_{65}/Γ
VALUE (units 10^{-4}) 34 ± 5 OUR AVERAGE	EVTS DOCUMENT ID TECN COMMENT

37.7±0.8±5.8 1972 ± 41 ABLIKIM 08E BES2 $e^+ e^- \rightarrow J/\psi$
29.5±1.4±7.0 879 ± 41 BECKER 87 MRK3 $e^+ e^- \rightarrow \text{hadrons}$

NODE=M070P26
NODE=M070P26

NODE=M070P26;LINKAGE=A

NODE=M070S16
NODE=M070S16

NODE=M070S35
NODE=M070S35

NODE=M070S35;LINKAGE=M3

NODE=M070S52
NODE=M070S52

NODE=M070S52;LINKAGE=AB

NODE=M070S26
NODE=M070S26

NODE=M070R49
NODE=M070R49

NODE=M070S1
NODE=M070S1

$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$				Γ_{66}/Γ	NODE=M070S28 NODE=M070S28
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
23±3±5	229	AUGUSTIN	89	DM2 $e^+ e^-$	
$\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$				Γ_{67}/Γ	NODE=M070S57 NODE=M070S57
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
21.8±2.2±3.4	232 ± 23	ABLIKIM	08E	BES2 $e^+ e^- \rightarrow J/\psi$	
$\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{37}/Γ	NODE=M070P49 NODE=M070P49
<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.66±0.03±0.21		1 ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$	
¹ From $\eta' K_S^0 K^\pm \pi^\mp$.					NODE=M070P49;LINKAGE=A
$\Gamma(\eta' K^{*\pm} \bar{K}^\mp)/\Gamma_{\text{total}}$				Γ_{36}/Γ	NODE=M070P48 NODE=M070P48
<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.48±0.13 OUR AVERAGE		1 ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$	
1.50±0.02±0.19		2 ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' K^* \bar{K}$	
1.47±0.03±0.17					OCCUR=2
¹ From $\eta' K^+ K^- \pi^0$.					NODE=M070P48;LINKAGE=A
² From $\eta' K_S^0 K^\pm \pi^\mp$.					NODE=M070P48;LINKAGE=B
$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} \bar{K}^\mp)/\Gamma_{\text{total}}$				Γ_{39}/Γ	NODE=M070P44 NODE=M070P44
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.51±0.09±0.21	1.0k	1 ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$	
¹ From $\eta' K^+ K^- \pi^0$.					NODE=M070P44;LINKAGE=A
$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{38}/Γ	NODE=M070P52 NODE=M070P52
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.16±0.12±0.29	1.1k	1 ABLIKIM	18AB	BES3 $J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$	
¹ From $\eta' K_S^0 K^\pm \pi^\mp$.					NODE=M070P52;LINKAGE=A
$\Gamma(\phi K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{68}/Γ	NODE=M070S4 NODE=M070S4
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
21.8±2.3 OUR AVERAGE		ABLIKIM	08E	BES2 $J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$	
20.8±2.7±3.9	195 ± 25	ABLIKIM	08E	BES2 $J/\psi \rightarrow \phi K^+ K^- \pi^0$	
29.6±3.7±4.7	238 ± 30	FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$	
20.7±2.4±3.0		BECKER	87	MRK3 $e^+ e^- \rightarrow \text{hadrons}$	
20 ± 3 ± 3	155 ± 20				OCCUR=2
$\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$				Γ_{69}/Γ	NODE=M070R27 NODE=M070R27
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
19 ± 4 OUR AVERAGE		1 FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$	
19.8±2.1±3.9		2 FELDMAN	77	MRK1 $e^+ e^-$	
16 ± 10	22				
¹ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.					NODE=M070R27;LINKAGE=B
$\Gamma(\omega f_0(1710) \rightarrow \omega K \bar{K})/\Gamma_{\text{total}}$				Γ_{70}/Γ	NODE=M070S25 NODE=M070S25
<u>VALUE (units 10^{-4})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
4.8±1.1±0.3	1,2	FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$	
¹ Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.					NODE=M070S25;LINKAGE=F
² Addition of $f_0(1710) \rightarrow K^+ K^-$ and $f_0(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.					NODE=M070S25;LINKAGE=G
$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$				Γ_{71}/Γ	NODE=M070R35 NODE=M070R35
<u>VALUE (units 10^{-4})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
16.0±1.0±3.0		FALVARD	88	DM2 $J/\psi \rightarrow \text{hadrons}$	
$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$				Γ_{72}/Γ	NODE=M070R70 NODE=M070R70
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.58±0.23±0.40	332	EATON	84	MRK2 $e^+ e^-$	

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ± 0.20 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
2.352 ± 0.273	5k	1 ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta$
1.44 ± 0.40 ± 0.14	13	2 AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \omega\eta\gamma$
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi\eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+ e^-) = 5.52 \pm 0.14 \pm 0.04 \text{ keV}$.

 Γ_{73}/Γ

NODE=M070R30
NODE=M070R30

 $\Gamma(\phi K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.7 ± 1.6 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
16.6 ± 1.9 ± 1.2	163 ± 19	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
21.4 ± 0.4 ± 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
48 ± 20 ± 6	9.0 ± 3.7	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 ± 0.8 ± 2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	$e^+ e^-$

¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K\bar{K}$.


² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

 Γ_{75}/Γ

NODE=M070R36
NODE=M070R36

NODE=M070R36;LINKAGE=AA
NODE=M070R36;LINKAGE=CC
NODE=M070R36;LINKAGE=A

 $\Gamma(\phi f_0(1710) \rightarrow \phi K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6±0.2±0.6	1,2 FALVARD	88	$J/\psi \rightarrow \text{hadrons}$

¹ Including interference with $f'_2(1525)$.² Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$. Γ_{77}/Γ NODE=M070S24
NODE=M070S24 $\Gamma(\phi f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.45	90	FALVARD	88	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

 Γ_{79}/Γ NODE=M070R39
NODE=M070R39 $\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10±0.09±0.28	233	EATON	84	MRK2 $e^+ e^-$

 Γ_{80}/Γ NODE=M070R66
NODE=M070R66 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.}))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.16 ± 0.05 OUR AVERAGE				
1.096±0.012±0.071	43K	ABLIKIM	16L	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
1.258±0.014±0.078	53k	ABLIKIM	16L	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
1.23 ± 0.07 ± 0.30	0.8k	ABLIKIM	12P	$J/\psi \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$
1.50 ± 0.08 ± 0.38	1k	ABLIKIM	12P	$J/\psi \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
1.00 ± 0.04 ± 0.21	0.6k	HENRARD	87	$e^+ e^- \rightarrow \Sigma^{*-}$
1.19 ± 0.04 ± 0.25	0.7k	HENRARD	87	$e^+ e^- \rightarrow \Sigma^{*+}$
0.86 ± 0.18 ± 0.22	56	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*-}$
1.03 ± 0.24 ± 0.25	68	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^{*+}$

 Γ_{81}/Γ NODE=M070R67
NODE=M070R67 $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.071±0.009±0.082	103k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

 Γ_{82}/Γ NODE=M070P17
NODE=M070P17
OCCUR=2 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE Error includes scale factor of 2.7.				
12.3±0.6±2.0	1,2	FALVARD	88	$J/\psi \rightarrow \text{hadrons}$
4.8±1.8	46	GIDAL	81	MRK2 $J/\psi \rightarrow K^+ K^- K^+ K^-$

 Γ_{84}/Γ NODE=M070R40
NODE=M070R40¹ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.² Including interference with $f_0(1710)$.NODE=M070R40;LINKAGE=B
NODE=M070R40;LINKAGE=C

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.94±0.15 OUR AVERAGE Error includes scale factor of 1.7.				
1.09±0.02±0.13		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
0.78±0.03±0.12		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ± 0.9	23	FELDMAN 77	MRK1	e^+e^-

 Γ_{85}/Γ

NODE=M070R34
NODE=M070R34

 $\Gamma(\phi K^\pm K_S^0\pi^\mp)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.2±0.8 OUR AVERAGE				
7.4±0.6±1.4	227 ± 19	ABLIKIM 08E	BES2	$e^+e^- \rightarrow J/\psi$
7.4±0.9±1.1		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
7 ± 0.6±1.0	163 ± 15	BECKER 87	MRK3	$e^+e^- \rightarrow \text{hadrons}$

 Γ_{87}/Γ

NODE=M070S3
NODE=M070S3

 $\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.8^{+1.9}_{-1.6}±1.7	111^{+31}_{-26}	BECKER 87	MRK3	$e^+e^- \rightarrow \text{hadrons}$

 Γ_{88}/Γ

NODE=M070S5
NODE=M070S5


 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.74 ± 0.08 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
0.898±0.024±0.089		ABLIKIM 05B	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadr}$
0.64 ± 0.04 ± 0.11	346	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
0.661±0.045±0.078		COFFMAN 88	MRK3	$e^+e^- \rightarrow K^+K^-\eta$

 Γ_{89}/Γ

NODE=M070R37
NODE=M070R37

WEIGHTED AVERAGE
0.74±0.08 (Error scaled by 1.5)



$\Gamma(\phi\eta)/\Gamma_{\text{total}}$ (units 10^{-3})

 $\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.32±0.06±0.16	2.2k	1 ABLIKIM	19AN BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

 Γ_{99}/Γ

NODE=M070P85
NODE=M070P85

¹ Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

NODE=M070P85;LINKAGE=A

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.17 ± 0.04 OUR AVERAGE				
1.165±0.004±0.043	135K	ABLIKIM 17E	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
1.20 ± 0.12 ± 0.21	206	ABLIKIM 080	BES2	$e^+e^- \rightarrow J/\psi$

 Γ_{90}/Γ

NODE=M070S64
NODE=M070S64

$\Gamma(\Xi(1530)^-\bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{91}/Γ
0.318±0.008 OUR AVERAGE					
[(0.59 ± 0.15) $\times 10^{-3}$ OUR 2019 AVERAGE]					
0.317±0.002±0.008	70k	ABLIKIM	20	BES3 $e^+e^- \rightarrow J/\psi$	
0.59 ± 0.09 ± 0.12	75	HENRARD	87	DM2 e^+e^-	


 $\Gamma(pK^-\bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{92}/Γ
0.51±0.26±0.18	89	EATON	84	MRK2 e^+e^-	

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{93}/Γ
0.45 ± 0.05 OUR AVERAGE					
Error includes scale factor of 1.4. See the ideogram below.					
0.538±0.012±0.065	2090	1 ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\pi^0$	
0.360±0.028±0.054	222	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$	
0.482±0.019±0.064		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^0\pi^+\pi^-\pi^0$	

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

 $\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{94}/Γ_{141}
8±3±2	20K	1 LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$	

¹ From a Dalitz plot analysis in an isobar model and significance 4.9 σ.

 $\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{95}/Γ
4.6 ± 0.5 OUR AVERAGE						

Error includes scale factor of 2.2. See the ideogram below.

5.10±0.03±0.32	31k	ABLIKIM	16M	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
5.46±0.31±0.56		ABLIKIM	05B	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
4.1 ± 0.3 ± 0.8	167	JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
3.08±0.34±0.36		COFFMAN	88	MRK3 $e^+e^- \rightarrow K^+K^-\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 13 90 VANNUCCI 77 MRK1 e^+e^-

NODE=M070S9

NODE=M070S9

NEW

NODE=M070R74

NODE=M070R74

NODE=M070R32

NODE=M070R32

NODE=M070R32;LINKAGE=BL


NODE=M070P23

NODE=M070P23

NODE=M070P23;LINKAGE=A

NODE=M070R38

NODE=M070R38

 $\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2±0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
4.6±0.4±0.8		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6±0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

 Γ_{96}/Γ

NODE=M070R41
NODE=M070R41

 $\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.50±0.80±0.61	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

 Γ_{100}/Γ

NODE=M070S97
NODE=M070S97

 $\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67±0.50±0.24	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

 Γ_{101}/Γ

NODE=M070S98
NODE=M070S98

 $\Gamma(\eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.23±0.75±0.73	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$

 Γ_{102}/Γ

NODE=M070R08
NODE=M070R08

 $\Gamma(\phi a_0(980)^0 \rightarrow \phi\eta\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.37±1.35	¹ ABLIKIM	18D BES3	$J/\psi \rightarrow \phi\eta\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.0 ± 2.7 ± 2.5	2 ABLIKIM	11D BES3	$J/\psi \rightarrow \phi\eta\pi^0$
-----------------	-----------	----------	------------------------------------

 Γ_{103}/Γ

NODE=M070S75
NODE=M070S75

¹ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.

² Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

NODE=M070S75;LINKAGE=A

 $\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.32±0.12±0.07	24 ± 9	HENRARD	87 DM2	$e^+ e^-$

 Γ_{104}/Γ

NODE=M070S10
NODE=M070S10

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.}))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31±0.05 OUR AVERAGE				
0.30±0.03±0.07	74 ± 8	HENRARD	87 DM2	$e^+ e^- \rightarrow \Sigma^{*-}$
0.34±0.04±0.07	77 ± 9	HENRARD	87 DM2	$e^+ e^- \rightarrow \Sigma^{*+}$
0.29±0.11±0.10	26	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^{*-}$
0.31±0.11±0.11	28	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^{*+}$

 Γ_{105}/Γ

NODE=M070R68
NODE=M070R68

OCCUR=2

OCCUR=2

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{106}/Γ</u>
2.6 ± 0.5 OUR AVERAGE					
$3.4 \pm 1.8 \pm 1.5$	1.1k	1 ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	
$3.2 \pm 0.6 \pm 0.4$		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$	
$2.1 \pm 0.5 \pm 0.4$	25	2 JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$0.6 \pm 0.2 \pm 0.1$	16	BECKER	87 MRK3	$J/\psi \rightarrow \phi K \bar{K} \pi$	
¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+ \pi^- \eta$ invariant mass distribution at 1297 MeV.					

NODE=M070S6
NODE=M070S6

OCCUR=2

NODE=M070S6;LINKAGE=A

NODE=M070S6;LINKAGE=Q

NODE=M070S99
NODE=M070S99NODE=M070S00
NODE=M070S00NODE=M070P81
NODE=M070P81

NODE=M070P81;LINKAGE=A

NODE=M070R22
NODE=M070R22NODE=M070R31
NODE=M070R31NODE=M070R31;LINKAGE=A
NODE=M070R31;LINKAGE=BLNODE=M070S27
NODE=M070S27

NODE=M070S27;LINKAGE=K

NODE=M070R23
NODE=M070R23

NODE=M070R23;LINKAGE=A

NODE=M070R42
NODE=M070R42 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{107}/Γ

<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{107}/Γ</u>
$9.36 \pm 2.31 \pm 1.54$	78	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$	

 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

<u>VALUE (units 10^{-7})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{108}/Γ</u>
$2.08 \pm 1.63 \pm 1.47$	9	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$	

 $\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{109}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{109}/Γ</u>
3.78 ± 0.68	471	1 ABLIKIM	19Q BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \eta \pi^+ \pi^-$	

¹ From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow \eta \pi^+ \pi^-$ assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$. $\Gamma(\eta \rho)/\Gamma_{\text{total}}$ Γ_{110}/Γ

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{110}/Γ</u>
0.193 ± 0.023 OUR AVERAGE					
0.194 $\pm 0.017 \pm 0.029$	299	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$	
0.193 $\pm 0.013 \pm 0.029$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^+ \pi^- \eta$	

 $\Gamma(\omega \eta'(958))/\Gamma_{\text{total}}$ Γ_{111}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{111}/Γ</u>
1.89 ± 0.18 OUR AVERAGE					
2.08 $\pm 0.30 \pm 0.14$	137	1 ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$	
2.26 ± 0.43	218	2 ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \eta'$	
$1.8 \pm 1.0 \pm 0.3$	6	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$	
1.66 $\pm 0.17 \pm 0.19$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow 3\pi \eta'$	

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.² Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+ \pi^- \gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$. $\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ Γ_{112}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{112}/Γ</u>
$1.41 \pm 0.27 \pm 0.47$		1 AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$	

Assuming $B(f_0(980) \rightarrow \pi \pi) = 0.78$. $\Gamma(\rho \eta'(958))/\Gamma_{\text{total}}$ Γ_{113}/Γ

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{113}/Γ</u>
8.1 ± 0.8 OUR AVERAGE Error includes scale factor of 1.6.					
7.90 $\pm 0.19 \pm 0.49$	3476	1 ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$	
8.3 $\pm 3.0 \pm 1.2$	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$	
11.4 $\pm 1.4 \pm 1.6$		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$	

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$. $\Gamma(a_2(1320)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$ Γ_{114}/Γ

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{114}/Γ</u>
$< 43 \times 10^{-4}$	90	BRAUNSCH...	76 DASP	$e^+ e^-$	

$\Gamma(K\bar{K}_2^*(1430) + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{115}/Γ
$<40 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$	NODE=M070R45 NODE=M070R45
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<66 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$	

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{116}/Γ
$<3.0 \times 10^{-3}$	90	1 BAI	99C	BES	$e^+ e^-$

¹ Assuming $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$ $\Gamma(K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT	Γ_{117}/Γ
$8.54^{+1.07+2.35}_{-1.20-2.13}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	I

 $\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{119}/Γ
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$	NODE=M070R47 NODE=M070R47

 $\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{120}/Γ
$2.94 \pm 0.16 \pm 0.16$	0.8k	1 ABLIKIM	15K	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$	I
$0.124 \pm 0.033 \pm 0.030$	35 ± 9	2 ABLIKIM	15K	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma\gamma$	I

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{120}/Γ
<6.4	90	3 ABLIKIM	05B	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\gamma\gamma$	I
<6.8	90	COFFMAN 88	MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$	OCCUR=2	

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi\pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

3 Superseded by ABLIKIM 15K.

 $\Gamma(\phi\eta(1405) \rightarrow \phi\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{121}/Γ
$2.01 \pm 0.58 \pm 0.82$	172	1 ABLIKIM	15H	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$	I

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{121}/Γ
<17	90	2 FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$	I

¹ With 3.6σ significance.² Includes unknown branching fraction $\eta(1405) \rightarrow \eta\pi\pi$. $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{122}/Γ
$<2.2 \times 10^{-4}$	90	1 VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-$	NODE=M070R29 NODE=M070R29

 $\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{122}/Γ
$<2.8 \times 10^{-4}$	90	1 FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$	I

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$. $\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{123}/Γ
$<3.9 \times 10^{-6}$	95	ABLIKIM 13P	BES3	$J/\psi \rightarrow \gamma\pi^0 p\bar{p}$	NODE=M070S81 NODE=M070S81

 $\Gamma(\omega X(1835), X \rightarrow \eta'\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{124}/Γ
$<6.2 \times 10^{-5}$	1 ABLIKIM 19AC	BES3		$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$	NODE=M070P84 NODE=M070P84

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.NODE=M070P72
NODE=M070P72NODE=M070R47
NODE=M070R47NODE=M070R33
NODE=M070R33

OCCUR=2

NODE=M070R33;LINKAGE=A

NODE=M070R33;LINKAGE=C

NODE=M070R33;LINKAGE=B

NODE=M070S23
NODE=M070S23NODE=M070R29
NODE=M070R29

NODE=M070R29;LINKAGE=C

NODE=M070S81
NODE=M070S81

NODE=M070P84;LINKAGE=A

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-7}$	90	1 ABLIKIM	16K BES3	$J/\psi \rightarrow p\bar{p} K_S^0 K_L^0$ $p\bar{p} K^+ K^-$

¹ Upper limit applies to any $p\bar{p}$ mass enhancement near threshold.

 Γ_{125}/Γ

NODE=M070P00
NODE=M070P00
OCCUR=2

 $\Gamma(\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-4}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

 Γ_{126}/Γ

NODE=M070B10
NODE=M070B10

 $\Gamma(\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.13 \times 10^{-5}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

 Γ_{127}/Γ

NODE=M070B11
NODE=M070B11

 $\Gamma(\eta \phi(2170) \rightarrow \eta \phi f_0(980) \rightarrow \eta \phi \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.14 \pm 0.37$	471	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

 Γ_{128}/Γ

NODE=M070B12
NODE=M070B12

 $\Gamma(\eta \phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.52 \times 10^{-4}$	90	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

 Γ_{129}/Γ

NODE=M070S70
NODE=M070S70

 $\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.82 \times 10^{-5}$	90	ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p} \pi^+ \pi^- \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.2 \times 10^{-3}$	90	HENRARD	87 DM2	$e^+ e^-$

 Γ_{130}/Γ

NODE=M070S13
NODE=M070S13

 $\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.1 \times 10^{-3}$	90	HENRARD	87 DM2	$e^+ e^-$

 Γ_{131}/Γ

NODE=M070S14
NODE=M070S14

 $\Gamma(\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.1 \times 10^{-6}$	90	ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda \bar{\Lambda} \gamma$

 Γ_{132}/Γ

NODE=M070S77
NODE=M070S77

 $\Gamma(\bar{\Lambda}(1520) \Lambda + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.80 \times 10^{-3}$	90	LU	19 BELL	$B^+ \rightarrow \bar{p} \Lambda K^+ K^+$

 Γ_{133}/Γ

NODE=M070P60
NODE=M070P60

 $\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 Γ_{134}/Γ

NODE=M070S47
NODE=M070S47

 $\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 Γ_{135}/Γ

NODE=M070S48
NODE=M070S48

 $\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 Γ_{136}/Γ

NODE=M070S49
NODE=M070S49

 $\Gamma(\Theta(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 Γ_{137}/Γ

NODE=M070S50
NODE=M070S50

 $\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^- \bar{n})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90	BAI	04G BES2	$e^+ e^-$

 Γ_{138}/Γ

NODE=M070S51
NODE=M070S51

STABLE HADRONS **$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$**


VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.73 ± 0.32 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.

$[(3.37 \pm 0.26) \times 10^{-2}$ OUR 2019 AVERAGE]

4.73 ± 0.44	228K	1 ABLIKIM	19Q BES3	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$
3.64 ± 0.52	1500	BURMESTER	77D PLUT	e^+e^-
4 ± 1	675	JEAN-MARIE	76 MRK1	e^+e^-

¹ From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$, assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

WEIGHTED AVERAGE
 3.73 ± 0.32 (Error scaled by 1.4)

 **$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.029 ± 0.006 OUR AVERAGE				

0.028 ± 0.009	11	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$
0.029 ± 0.007	181	JEAN-MARIE	76 MRK1	e^+e^-

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
21.0 ± 0.8 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.

$21.37 \pm 0.04^{+0.64}_{-0.62}$	1.8M	1,2 ABLIKIM	12H BES3	$e^+e^- \rightarrow J/\psi$
$23.0 \pm 2.0 \pm 0.4$	256	3 AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
$21.84 \pm 0.05 \pm 2.01$	220k	1,4 BAI	04H BES	e^+e^-
$20.91 \pm 0.21 \pm 1.16$		4,5 BAI	04H BES	e^+e^-
15 ± 2	168	FRANKLIN	83 MRK2	e^+e^-

¹ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.

² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.

³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.013 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ Mostly $\rho\pi$, see also $\rho\pi$ subsection.

⁵ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

NODE=M070307

NODE=M070R9

NODE=M070R9

NEW

NODE=M070R9;LINKAGE=A

NODE=M070R11

NODE=M070R11

NODE=M070R7

NODE=M070R7

OCCUR=2


NODE=M070R;LINKAGE=BA

NODE=M070R7;LINKAGE=AB

NODE=M070R7;LINKAGE=AU

NODE=M070R;LINKAGE=BU

NODE=M070R;LINKAGE=BI

 $\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.3	309	VANNUCCI	77	MRK1 $e^+ e^-$

 Γ_{145}/Γ NODE=M070R18
NODE=M070R18 $\Gamma(4(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
90±30	13	JEAN-MARIE	76	MRK1 $e^+ e^-$

 Γ_{146}/Γ NODE=M070R12
NODE=M070R12 $\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

 Γ_{147}/Γ NODE=M070R16
NODE=M070R16 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ±10 OUR AVERAGE				

 Γ_{155}/Γ NODE=M070R15
NODE=M070R15

55.2±12.0

78.0±21.0

25

126

FRANKLIN

VANNUCCI

83

77

MRK2

MRK1

 $e^+ e^- \rightarrow K^+ K^- \pi^0$ $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$ $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.88±0.01±0.12	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

 Γ_{156}/Γ NODE=M070P80
NODE=M070P80 $\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0±0.3±0.9	23K	LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$

 $\Gamma_{156}/\Gamma_{141}$ NODE=M070P34
NODE=M070P34 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
26.5±0.5±2.1	24K	LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$

 $\Gamma_{157}/\Gamma_{141}$ NODE=M070P35
NODE=M070P35 $\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.57±0.30 OUR AVERAGE	1107	¹ ABLIKIM	05H BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+ \pi^-, J/\psi \rightarrow 2(\pi^+ \pi^-)$

 Γ_{162}/Γ NODE=M070R8
NODE=M070R8

3.53±0.12±0.29

4.0 ±1.0

1107

76

ABLIKIM

JEAN-MARIE

05H

76

BES2

MRK1

 $e^+ e^-$ ¹ Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

NODE=M070R8;LINKAGE=AB

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

40±20 32 JEAN-MARIE 76 MRK1 e^+e^-

 Γ_{163}/Γ

NODE=M070R10
NODE=M070R10

 $\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.26 \pm 0.08 \pm 0.27$	4.8k	ABLIKIM	05C	$e^+e^- \rightarrow 2(\pi^+\pi^-)\eta$

 Γ_{165}/Γ

NODE=M070S42
NODE=M070S42

 $\Gamma(3(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$7.24 \pm 0.96 \pm 1.11$	616	ABLIKIM	05C	$e^+e^- \rightarrow 3(\pi^+\pi^-)\eta$

 Γ_{166}/Γ

NODE=M070S43
NODE=M070S43

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

 2.121 ± 0.029 OUR AVERAGE

2.112 $\pm 0.004 \pm 0.031$	314k	ABLIKIM	12C	e^+e^-
2.20 $\pm 0.16 \pm 0.06$	317	WU	06	$B^+ \rightarrow p\bar{p}K^+$
2.26 $\pm 0.01 \pm 0.14$	63316	BAI	04E	$e^+e^- \rightarrow J/\psi$
1.97 ± 0.22	99	BALDINI	98	FENI e^+e^-
1.91 $\pm 0.04 \pm 0.30$		PALLIN	87	DM2 e^+e^-
2.16 $\pm 0.07 \pm 0.15$	1420	EATON	84	MRK2 e^+e^-
2.5 ± 0.4	133	BRANDELIK	79C	DASP e^+e^-
2.0 ± 0.5		BESCH	78	BONA e^+e^-
2.2 ± 0.2	331	PERUZZI	78	MRK1 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3 48 ANTONELLI 93 SPEC e^+e^-

 Γ_{169}/Γ

NODE=M070R50
NODE=M070R50

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.006 \pm 0.027) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Assuming angular distribution $(1+\cos^2\theta)$.

NODE=M070R50;LINKAGE=WU

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

 1.19 ± 0.08 OUR AVERAGE Error includes scale factor of 1.1.

1.33 $\pm 0.02 \pm 0.11$	11k	ABLIKIM	09B	e^+e^-
1.13 $\pm 0.09 \pm 0.09$	685	EATON	84	MRK2 e^+e^-
1.4 ± 0.4		BRANDELIK	79C	DASP e^+e^-
1.00 ± 0.15	109	PERUZZI	78	MRK1 e^+e^-

 Γ_{170}/Γ

NODE=M070R52
NODE=M070R52

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------


 6.0 ± 0.5 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

6.46 $\pm 0.17 \pm 0.43$	1435	EATON	84	MRK2 e^+e^-
3.8 ± 1.6	48	BESCH	81	BONA e^+e^-
5.5 ± 0.6	533	PERUZZI	78	MRK1 e^+e^-

 Γ_{171}/Γ

NODE=M070R54
NODE=M070R54

WEIGHTED AVERAGE
6.0±0.5 (Error scaled by 1.3)



$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ±0.9 OUR AVERAGE Error includes scale factor of 1.9.				
$3.36 \pm 0.65 \pm 0.28$	364	EATON	84	MRK2 e^+e^-
1.6 ± 0.6	39	PERUZZI	78	MRK1 e^+e^-

Γ_{172}/Γ

NODE=M070R55
NODE=M070R55
NODE=M070R55

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$

Γ_{173}/Γ

NODE=M070R56
NODE=M070R56

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00±0.12 OUR AVERAGE				

$1.91 \pm 0.02 \pm 0.17$	13k	¹ ABLIKIM	09	BES2 e^+e^-
$2.03 \pm 0.13 \pm 0.15$	826	EATON	84	MRK2 e^+e^-
2.5 ± 1.2		BRANDELIK	79c	DASP e^+e^-
2.3 ± 0.4	197	PERUZZI	78	MRK1 e^+e^-

¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

$\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$

Γ_{174}/Γ

NODE=M070R57
NODE=M070R57

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.31 \times 10^{-3}$	90	EATON	84	MRK2 $e^+e^- \rightarrow \text{hadrons}\gamma$


$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

Γ_{175}/Γ

NODE=M070R58
NODE=M070R58

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98±0.10 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				

$0.90 \pm 0.02 \pm 0.09$	2670	ABLIKIM	13P	BES3 e^+e^-
$0.98 \pm 0.03 \pm 0.14$	2449	ABLIKIM	08	BES2 e^+e^-
$1.10 \pm 0.17 \pm 0.18$	486	EATON	84	MRK2 e^+e^-
1.6 ± 0.3	77	PERUZZI	78	MRK1 e^+e^-

 $\Gamma(p\bar{p}\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.129±0.014 OUR AVERAGE				Error includes scale factor of 2.0.
0.126±0.002±0.007	16K	1 ABLIKIM	19N BES3	$e^+ e^-$
0.200±0.023±0.028	265 ± 31	2 ABLIKIM	09 BES2	$e^+ e^-$
0.68 ± 0.23 ± 0.17	19	EATON	84 MRK2	$e^+ e^-$
1.8 ± 0.6	19	PERUZZI	78 MRK1	$e^+ e^-$

1 From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.2 From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\gamma\rho^0$ channels. Γ_{176}/Γ NODE=M070R59
NODE=M070R59 $\Gamma(p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{177}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8±1.2±1.3		ABLIKIM	14N BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070R59;LINKAGE=A
NODE=M070R59;LINKAGE=AB $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{178}/Γ NODE=M070S22
NODE=M070S22**0.519±0.033 OUR AVERAGE**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.523±0.006±0.033	14K	ABLIKIM	16K BES3	$J/\psi \rightarrow p\bar{p}K_S^0 K_L^0$
0.45 ± 0.13 ± 0.07		FALVARD	88 DM2	$p\bar{p}K^+ K^-$ $J/\psi \rightarrow \text{hadrons}$

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$ Γ_{179}/Γ NODE=M070R64
NODE=M070R64**2.09±0.16 OUR AVERAGE**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.09±0.16 OUR AVERAGE				
2.07±0.01±0.17	36k	ABLIKIM	12C BES3	$e^+ e^-$
2.31±0.49	79	BALDINI	98 FENI	$e^+ e^-$
1.8 ± 0.9		BESCH	78 BONA	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.90±0.55	40	ANTONELLI	93 SPEC	$e^+ e^-$

 $\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{180}/Γ NODE=M070R65
NODE=M070R65

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8±3.6	5	BESCH	81 BONA	$e^+ e^-$

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{181}/Γ NODE=M070S09
NODE=M070S09

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.50±0.10±0.22	399	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$

<i>VALUE (units 10^{-3})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	Γ_{182}/Γ
1.172 ± 0.032 OUR AVERAGE					
1.164 $\pm 0.004 \pm 0.023$	111k	ABLIKIM	17L	BES3 $J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0$	
1.33 $\pm 0.04 \pm 0.11$	1.7k	ABLIKIM	06	BES2 $J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0$	
1.06 $\pm 0.04 \pm 0.23$	884	PALLIN	87	DM2 $e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$	
1.58 $\pm 0.16 \pm 0.25$	90	EATON	84	MRK2 $e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$	
1.3 ± 0.4	52	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
2.4 ± 2.6	3	BESCH	81	BONA $e^+ e^- \rightarrow \Sigma^+ \bar{\Sigma}^-$	

 $\Gamma(2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$

<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	Γ_{183}/Γ
31 ± 13	30	VANNUCCI	77	MRK1 $e^+ e^-$	


 $\Gamma(p \bar{n} \pi^-)/\Gamma_{\text{total}}$

<i>VALUE (units 10^{-3})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	Γ_{184}/Γ
1.12 ± 0.09 OUR AVERAGE					
2.36 $\pm 0.02 \pm 0.21$	59k	ABLIKIM	06K	BES2 $J/\psi \rightarrow p \pi^- \bar{n}$	
2.47 $\pm 0.02 \pm 0.24$	55k	ABLIKIM	06K	BES2 $J/\psi \rightarrow \bar{p} \pi^+ n$	OCCUR=2
2.02 $\pm 0.07 \pm 0.16$	1288	EATON	84	MRK2 $e^+ e^- \rightarrow p \pi^-$	OCCUR=2
1.93 $\pm 0.07 \pm 0.16$	1191	EATON	84	MRK2 $e^+ e^- \rightarrow \bar{p} \pi^+$	OCCUR=2
1.7 ± 0.7	32	BESCH	81	BONA $e^+ e^- \rightarrow p \pi^-$	OCCUR=2
1.6 ± 1.2	5	BESCH	81	BONA $e^+ e^- \rightarrow \bar{p} \pi^+$	OCCUR=2
2.16 ± 0.29	194	PERUZZI	78	MRK1 $e^+ e^- \rightarrow p \pi^-$	OCCUR=2
2.04 ± 0.27	204	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \bar{p} \pi^+$	OCCUR=2

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

<i>VALUE (units 10^{-3})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	Γ_{188}/Γ
0.97 ± 0.08 OUR AVERAGE					
1.040 $\pm 0.006 \pm 0.074$	43k	ABLIKIM	16L	BES3 $J/\psi \rightarrow \Xi^- \bar{\Xi}^+$	
0.90 $\pm 0.03 \pm 0.18$	961	ABLIKIM	12P	BES2 $J/\psi \rightarrow \Xi^- \bar{\Xi}^+$	
0.70 $\pm 0.06 \pm 0.12$	132	HENRARD	87	DM2 $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$	
1.14 $\pm 0.08 \pm 0.20$	194	EATON	84	MRK2 $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$	
1.4 ± 0.5	51	PERUZZI	78	MRK1 $e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$	

WEIGHTED AVERAGE
 0.97 ± 0.08 (Error scaled by 1.4)



NODE=M070R63
NODE=M070R63

NODE=M070R17
NODE=M070R17

NODE=M070R53
NODE=M070R53

NODE=M070R62
NODE=M070R62

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89 ± 0.09 OUR AVERAGE				Error includes scale factor of 2.8. See the ideogram below.
1.943 ± 0.003 ± 0.033	441k	ABLIKIM	17L	BES3 $e^+ e^-$
2.03 ± 0.03 ± 0.15	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$
2.0 ± 0.5 ± 0.1	46	¹ WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ± 0.06 ± 0.24	631	BAI	98G	BES $e^+ e^-$
1.38 ± 0.05 ± 0.20	1847	PALLIN	87	DM2 $e^+ e^-$
1.58 ± 0.08 ± 0.19	365	EATON	84	MRK2 $e^+ e^-$
2.6 ± 1.6	5	BESCH	81	BONA $e^+ e^-$
1.1 ± 0.2	196	PERUZZI	78	MRK1 $e^+ e^-$

¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.006 \pm 0.027) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R60
NODE=M070R60

 Γ_{189}/Γ

NODE=M070R60;LINKAGE=WU

 $\Gamma(\Lambda\bar{\Sigma}^-\pi^+ (\text{or c.c.}))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.2.
0.770 ± 0.051 ± 0.083	335	¹ ABLIKIM	07H BES2	$e^+ e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
0.747 ± 0.056 ± 0.076	254	¹ ABLIKIM	07H BES2	$e^+ e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
0.90 ± 0.06 ± 0.16	225 ± 15	HENRARD	87 DM2	$e^+ e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
1.11 ± 0.06 ± 0.20	342 ± 18	HENRARD	87 DM2	$e^+ e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
1.53 ± 0.17 ± 0.38	135	EATON	84 MRK2	$e^+ e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
1.38 ± 0.21 ± 0.35	118	EATON	84 MRK2	$e^+ e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

NODE=M070R71
NODE=M070R71

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M070R71;LINKAGE=AB

 Γ_{190}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.87 ± 0.11 OUR AVERAGE				
0.85 ± 0.17 ± 0.02	45	¹ LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
0.89 ± 0.07 ± 0.14	307	EATON	84 MRK2	$e^+ e^-$

¹ LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda}+\text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.006 \pm 0.027) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R72
NODE=M070R72

NODE=M070R72;LINKAGE=A

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{192}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	1 HUANG 03	BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$	
0.7 ± 0.3		VANNUCCI 77	MRK1	$e^+ e^-$	
1 Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.					

 $\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{193}/Γ
$0.29 \pm 0.06 \pm 0.05$	90	EATON	84	MRK2 $e^+ e^-$	

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{194}/Γ
$2.86 \pm 0.09 \pm 0.19$	1k	1 METREVELI 12		$\psi(2S) \rightarrow \pi^+ \pi^- K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.39 \pm 0.24 \pm 0.22$	107	2 BALTRUSAIT..85D	MRK3	$e^+ e^-$
2.2 ± 0.9	6	2 BRANDELIK	79C DASP	$e^+ e^-$

1 Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.**2 Interference with non-resonant $K^+ K^-$ production not taken into account.** $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{195}/Γ
1.95 ± 0.11 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.	


$1.93 \pm 0.01 \pm 0.05$	110K	ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$
$2.62 \pm 0.15 \pm 0.14$	0.3k	1 METREVELI	12	$\psi(2S) \rightarrow \pi^+ \pi^- K_S^0 K_L^0$
$1.82 \pm 0.04 \pm 0.13$	2.1k	2 BAI	04A BES2	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.18 \pm 0.12 \pm 0.18$		JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$1.01 \pm 0.16 \pm 0.09$	74	BALTRUSAIT..85D	MRK3	$e^+ e^-$

1 Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.**2 Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6868 \pm 0.0027$.**

WEIGHTED AVERAGE
1.95 ± 0.11 (Error scaled by 2.4)

 Γ_{195}/Γ $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{196}/Γ
4.30 $\pm 0.13 \pm 0.99$	2.4k	ABLIKIM	12P BES2	J/ψ	

NODE=M070R19
NODE=M070R19

NODE=M070R19;LINKAGE=CC

NODE=M070R73
NODE=M070R73NODE=M070R13
NODE=M070R13NODE=M070R13;LINKAGE=ME
NODE=M070R13;LINKAGE=BANODE=M070R75
NODE=M070R75NODE=M070R75;LINKAGE=ME
NODE=M070R;LINKAGE=HZNODE=M070S78
NODE=M070S78

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				
15.7±0.80±1.54	454	1 ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2±6.0 ±4.4	44	2 ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

 Γ_{197}/Γ

NODE=M070R07
NODE=M070R07

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	1 ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	2 ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$
23 ± 7 ± 8	11	BAI	98G BES	e^+e^-
22 ± 5 ± 5	19	HENRARD	87 DM2	e^+e^-

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

 Γ_{198}/Γ

NODE=M070S11
NODE=M070S11

 $\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46±0.20±1.07	1058	1 ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				

1.47±0.13±0.13	140	1 METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-
1.0 ± 0.5	5	BRANDELIK	78B DASP	e^+e^-
1.6 ± 1.6	1	VANNUCCI	77 MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

 Γ_{199}/Γ

NODE=M070S56
NODE=M070S56

 $\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83±0.23 OUR AVERAGE					

2.74±0.24±0.22	234 ± 21	1 ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92±0.22±0.24	308 ± 24	2 ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18		2 HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15	90	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

 Γ_{201}/Γ

NODE=M070R61
NODE=M070R61

OCCUR=2

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁸				

95	1 ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$
----	-----------	-----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1 × 10 ⁻⁶	95	1 BAI	04D BES	e^+e^-
<5.2 × 10 ⁻⁶	90	1 BALTRUSAIT..85C	MRK3	e^+e^-

¹ Forbidden by CP.

 Γ_{202}/Γ

NODE=M070R14
NODE=M070R14

NODE=M070R14;LINKAGE=C

NODE=M070310

NODE=M070R81
NODE=M070R81

 $\Gamma(3\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.6±2.2 OUR AVERAGE					

11.3±1.8±2.0	113 ± 18	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ± 3 ± 2	24.2 ^{+7.2} _{-6.0}	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	PARTRIDGE	80 CBAL	e^+e^-
-----	----	-----------	---------	----------

 Γ_{203}/Γ

$\Gamma(4\gamma)/\Gamma_{\text{total}}$					Γ_{204}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S06 NODE=M070S06
$<9 \times 10^{-6}$	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$		
$\Gamma(5\gamma)/\Gamma_{\text{total}}$					Γ_{205}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S07 NODE=M070S07
$<15 \times 10^{-6}$	90	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$		
$\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_{206}/Γ	
<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070B00 NODE=M070B00
1.15 ± 0.05		¹ ABLIKIM 15AE BES3		$J/\psi \rightarrow \gamma\pi^0\pi^0$		
1 The uncertainty is systematic as statistical is negligible.						
$\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{207}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P01 NODE=M070P01
$21.4 \pm 1.8 \pm 2.5$	596	ABLIKIM 16P	BES3	$J/\psi \rightarrow 5\gamma$		
$\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{208}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P02 NODE=M070P02
$<2.5 \times 10^{-6}$	95	ABLIKIM 16P	BES3	$J/\psi \rightarrow 5\gamma$		
$\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$					Γ_{209}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P03 NODE=M070P03
$<6.6 \times 10^{-6}$	95	ABLIKIM 16P	BES3	$J/\psi \rightarrow 5\gamma$		
$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{210}/Γ	
<u>VALUE (units 10^{-4})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P73 NODE=M070P73
8.1 ± 0.4		ABLIKIM 18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$		
$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					Γ_{211}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070R85 NODE=M070R85
1.7 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.5.				
2.00 \pm 0.31 \pm 0.02		¹ MITCHELL 09	CLEO	$e^+ e^- \rightarrow \gamma X$		
1.27 \pm 0.36		GAISER 86	CBAL	$J/\psi \rightarrow \gamma X$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
seen		ANASHIN 14	KEDR	$J/\psi \rightarrow \gamma\eta_c$		
0.79 \pm 0.20	273 \pm 43	² AUBERT 06E	BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$		
seen	16	BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$		
1 MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						NODE=M070R85;LINKAGE=MI
2 Calculated by the authors using an average of $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$ from BALTRUSAITIS 86, BISELLO 91, BAI 04 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.						NODE=M070R85;LINKAGE=AU
$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{\text{total}}$					Γ_{212}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S08 NODE=M070S08
$3.8^{+1.3}_{-1.0}$ OUR AVERAGE		Error includes scale factor of 1.1.				
4.5 \pm 1.2 \pm 0.6	33 \pm 9	ABLIKIM 13I	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$		
1.2 \pm 2.7 \pm 0.3	1.2 \pm 2.8 \pm 1.1	ADAMS 08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$		
$\Gamma(\gamma\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$					Γ_{213}/Γ	
<u>VALUE (units 10^{-3})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070R99 NODE=M070R99
$8.3 \pm 0.2 \pm 3.1$		¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$		
1 4π mass less than 2.0 GeV.						NODE=M070R99;LINKAGE=M

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
6.1 ± 1.0 OUR AVERAGE			

 $5.85 \pm 0.3 \pm 1.05$ $7.8 \pm 1.2 \pm 2.4$

1 Broad enhancement at 1700 MeV.

 $\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.2 ± 2.2 ± 0.9			

6.2 ± 2.2 ± 0.9

BAI

DOCUMENT ID	TECN	COMMENT
BAI	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

 Γ_{214}/Γ NODE=M070R96
NODE=M070R96

OCCUR=2

NODE=M070R96;LINKAGE=M


NODE=M070S37
NODE=M070S37 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE			

2.8 ± 0.6 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

 $1.66 \pm 0.1 \pm 0.58$ $3.8 \pm 0.3 \pm 0.6$ $4.0 \pm 0.7 \pm 1.0$ 4.3 ± 1.7 $1.78 \pm 0.21 \pm 0.33$ $0.83 \pm 0.13 \pm 0.18$ $0.66^{+0.17}_{-0.16} {}^{+0.24}_{-0.15}$ $1.03^{+0.21}_{-0.18} {}^{+0.26}_{-0.19}$ 1,2 BAI 00D BES $J/\psi \rightarrow \gamma K^{\pm} K_S^0 \pi^{\mp}$ 3 AUGUSTIN 90 DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$ 3 EDWARDS 82E CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$ 3,4 SCHARRE 80 MRK2 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3,5,6 AUGUSTIN 92 DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$ 3,7,8 AUGUSTIN 92 DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$ 3,6,9 BAI 90C MRK3 $J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$ 3,8,10 BAI 90C MRK3 $J/\psi \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$ 1 Interference with the $J/\psi(1S)$ radiative transition to the broad $K\bar{K}\pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.2 Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.3 Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.4 Corrected for spin-zero hypothesis for $\eta(1405)$.5 From fit to the $a_0(980)\pi^-$ partial wave.6 $a_0(980)\pi^-$ mode.7 From fit to the $K^*(892)K^-$ partial wave.8 K^*K^- mode.9 From $a_0(980)\pi^-$ final state.10 From $K^*(890)K^-$ final state.WEIGHTED AVERAGE
 2.8 ± 0.6 (Error scaled by 1.6) $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.20 OUR AVERAGE			

0.78 ± 0.20 OUR AVERAGE Error includes scale factor of 1.8.


 $1.07 \pm 0.17 \pm 0.11$ $0.64 \pm 0.12 \pm 0.07$ 1 Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$. Γ_{217}/Γ NODE=M070S30
NODE=M070S30

NODE=M070S30;LINKAGE=C

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{218}/Γ	NODE=M070S29 NODE=M070S29
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
3.0 ± 0.5 OUR AVERAGE						
2.6 ± 0.7 ± 0.4		BAI	99	BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$		
3.38 ± 0.33 ± 0.64		1 BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
7.0 ± 0.6 ± 1.1	261	2 AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$		
1 Via $a_0(980)\pi$.						
2 Includes unknown branching fraction to $\eta\pi^+\pi^-$.						
$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$					Γ_{219}/Γ	
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<82	95		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
7.03 ± 0.92 ± 0.91	1.3k	1 ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$		
10.36 ± 1.51 ± 1.54	1.9k	2 ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$		
1 Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.						
2 Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.						
$\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{220}/Γ	
VALUE	CL%		DOCUMENT ID	TECN	COMMENT	
<2.63 × 10⁻⁶	90		ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$	
$\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{221}/Γ	
VALUE	CL%		DOCUMENT ID	TECN	COMMENT	
<1.86 × 10⁻⁶	90		ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$	
$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$					Γ_{222}/Γ	
VALUE (units 10^{-3})	CL%		DOCUMENT ID	TECN	COMMENT	
4.5 ± 0.8 OUR AVERAGE						
4.7 ± 0.3 ± 0.9			1 BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$	
3.75 ± 1.05 ± 1.20			2 BURKE	82	MRK2 $J/\psi \rightarrow 4\pi\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.09	90	3 BISELLO	89B		$J/\psi \rightarrow 4\pi\gamma$	
1 4π mass less than 2.0 GeV.						
2 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .						
3 4π mass in the range 2.0–25 GeV.						
$\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$					Γ_{223}/Γ	
VALUE	CL%		DOCUMENT ID	TECN	COMMENT	
<5.4 × 10⁻⁴	90		ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$	
$\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$					Γ_{224}/Γ	
VALUE	CL%		DOCUMENT ID	TECN	COMMENT	
<8.8 × 10⁻⁵	90		ABLIKIM	08A	BES2 $e^+e^- \rightarrow J/\psi$	
$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$					Γ_{225}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
5.25 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.		
[(5.21 ± 0.17) × 10 ⁻³ OUR 2019 AVERAGE Scale factor = 1.4]						
5.27 ± 0.03 ± 0.05	36k	ABLIKIM	19T	BES $J/\psi \rightarrow \gamma\eta'$		
5.43 ± 0.23 ± 0.09	5.0k	1 ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$		
4.77 ± 0.22 ± 0.06		2 ABLIKIM	11	BES3 $J/\psi \rightarrow \eta'\gamma$		
5.24 ± 0.12 ± 0.11		PEDLAR	09	CLE3 $J/\psi \rightarrow \eta'\gamma$		
5.55 ± 0.44	35k	ABLIKIM	06E	BES2 $J/\psi \rightarrow \eta'\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
4.50 ± 0.14 ± 0.53		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$		
4.30 ± 0.31 ± 0.71		BOLTON	92B	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$		OCCUR=2
4.04 ± 0.16 ± 0.85	622	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$		
4.39 ± 0.09 ± 0.66	2420	AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$		OCCUR=2
4.1 ± 0.3 ± 0.6		BLOOM	83	CBAL $e^+e^- \rightarrow 3\gamma + \text{hadrons}$		
2.9 ± 1.1	6	BRANDELIK	79C	DASP $e^+e^- \rightarrow 3\gamma$		
2.4 ± 0.7	57	BARTEL	76	CNTR $e^+e^- \rightarrow 2\gamma\rho$		

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.



$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})

	DOCUMENT ID	TECN	COMMENT
2.8 ±0.5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		
4.32±0.14±0.73	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
2.08±0.13±0.35	² BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$
3.05±0.08±0.45	² BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
4.85±0.45±1.20	³ BURKE	82 MRK2	$e^+ e^-$

¹ 4π mass less than 3.0 GeV.

² 4π mass less than 2.0 GeV.

³ 4π mass less than 2.5 GeV.

Γ_{226}/Γ


NODE=M070R95
NODE=M070R95

OCCUR=2

NODE=M070R95;LINKAGE=A
NODE=M070R95;LINKAGE=B
NODE=M070R95;LINKAGE=M

NODE=M070R84;LINKAGE=A

NODE=M070R84;LINKAGE=AB

 $\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5±0.7±1.6	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

 Γ_{227}/Γ NODE=M070S45
NODE=M070S45 $\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.2±0.8±1.7	¹ ABLIKIM	04M BES		$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

 Γ_{228}/Γ NODE=M070S46
NODE=M070S46¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays. $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.1±0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

 Γ_{229}/Γ NODE=M070B05
NODE=M070B05 $\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.7±0.5±0.5	¹ BALTRUSAIT..87		MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

 Γ_{230}/Γ NODE=M070S7
NODE=M070S7¹ Assuming branching fraction $f_4(2050) \rightarrow \pi\pi/\text{total} = 0.167$. $\Gamma(\gamma \omega \omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61±0.33 OUR AVERAGE				
6.0 ± 4.8 ± 1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma \omega \pi^+ \pi^-$
1.41 ± 0.2 ± 0.42	120 ± 17	BISELLLO	87 SPEC	$e^+ e^-$, hadrons γ
1.76 ± 0.09 ± 0.45		BALTRUSAIT..85C	MRK3	$e^+ e^- \rightarrow$ hadrons γ

 Γ_{231}/Γ NODE=M070R97
NODE=M070R97 $\Gamma(\gamma \eta(1405/1475) \rightarrow \gamma \rho^0 \rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.3.
2.1 ± 0.4		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1.36 ± 0.38		^{1,2} BISSELLLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

 Γ_{232}/Γ NODE=M070S19
NODE=M070S19¹ Estimated by us from various fits.² Includes unknown branching fraction to $\rho^0 \rho^0$. $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.64±0.12 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
$2.07 \pm 0.16^{+0.02}_{-0.07}$	2.4k	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi\pi$
$1.63 \pm 0.26^{+0.02}_{-0.06}$		³ ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

 Γ_{233}/Γ NODE=M070R86
NODE=M070R86NODE=M070S19;LINKAGE=A
NODE=M070S19;LINKAGE=B

1.42 ± 0.21	$+0.01$	-0.05	4	ABLIKIM	06v	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
1.33 ± 0.05	± 0.20		5	AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.36 ± 0.09	± 0.23		5	BALTRUSAIT..	87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1.48 ± 0.25	± 0.30	178		EDWARDS	82B	CBAL	$e^+ e^- \rightarrow 2\pi^0 \gamma$
2.0	± 0.7	35		ALEXANDER	78	PLUT	$e^+ e^-$
1.2	± 0.6	30	6	BRANDELIK	78B	DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270)) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 06v reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270)) / \Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270)) / \Gamma_{\text{total}}] \times [\mathcal{B}(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $\mathcal{B}(f_2(1270) \rightarrow \pi\pi) = (84.2^{+2.9}_{-0.9}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

5 Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁶ Restated by us to take account of spread of E1, M2, E3 transitions.

OCCUR=2


NODE=M070R86;LINKAGE=A
NODE=M070R86;LINKAGE=DO

NODE=M070R86;LINKAGE=AI

NODE=M070R86;LINKAGE=AL

NODE=M070R86;LINKAGE=X

NODE=M070R86;LINKAGE=T



$$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$$

<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.58 +0.08 -0.09	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_\xi^0 K_\xi^0$

NODE=M070P68
NODE=M070P68

$$\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
4.19+0.73+1.34	478	1 DOBBS	$U^{1/2} \rightarrow \gamma K\bar{K}$

NODE=M070R00
NODE=M070R00

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$$\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 \bar{K}_S^0)/\Gamma_{\text{total}}$$

<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.07 +0.08 -0.07	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE-M070B63

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-5}) $1.59 \pm 0.16^{+0.18}_{-0.56}$

DOCUMENT ID

ABLIKIM

TECN

18AA BES3

COMMENT

 Γ_{237}/Γ

NODE=M070P64

NODE=M070P64

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

CL% EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{238}/Γ

NODE=M070R91

NODE=M070R91

9.5 \pm 1.0 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.00 \pm 0.12 \pm 1.24 0.08 \pm 0.40	1 ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
11.76 \pm 0.54 \pm 0.94 1.2k	2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
9.62 \pm 0.29 \pm 3.51 -1.86	3 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
5.0 \pm 0.8 \pm 1.8 -0.4	1,4 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
9.2 \pm 1.4 \pm 1.4 10.4 \pm 1.2 \pm 1.6	1 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
9.6 \pm 1.2 \pm 1.8	1 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
	1 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.6 \pm 0.2 \pm 0.6 < 0.8 90	1,5 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
1.6 \pm 0.4 \pm 0.3	6 BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
3.8 \pm 1.6	7 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$

¹ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.

⁴ Assuming $J^P = 2^+$ for $f_0(1710)$.

⁵ Assuming $J^P = 0^+$ for $f_0(1710)$.

⁶ Includes unknown branching fraction to $\rho^0 \rho^0$.

⁷ Includes unknown branching fraction to $\pi^+ \pi^-$.

⁸ Includes unknown branching fraction to $\eta\eta$.

OCCUR=2

OCCUR=2

NODE=M070R91;LINKAGE=B

NODE=M070R91;LINKAGE=D

NODE=M070R91;LINKAGE=K9

NODE=M070R91;LINKAGE=A1


NODE=M070R91;LINKAGE=A2

NODE=M070R91;LINKAGE=C

NODE=M070R91;LINKAGE=Z

NODE=M070R91;LINKAGE=A

WEIGHTED AVERAGE
9.5+1.0-0.5 (Error scaled by 1.5)

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**3.8 \pm 0.5 OUR AVERAGE**3.72 \pm 0.30 \pm 0.43 4833.96 \pm 0.06 \pm 1.123.99 \pm 0.15 \pm 2.64

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.5 \pm 1.6 \pm 0.8 BAI 98H BES $J/\psi \rightarrow \gamma\pi^0\pi^0$ Γ_{239}/Γ

NODE=M070B01

NODE=M070B01

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi\pi$.

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{240}/Γ
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H	BES	$J/\psi \rightarrow \gamma\omega\omega$

$\Gamma(\gamma f_0(1710) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{241}/Γ
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N	BES3	$J/\psi \rightarrow \gamma\eta\eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{242}/Γ
1.108 ± 0.027 OUR AVERAGE					

1.12 $\pm 0.05 \pm 0.01$	18.6k	¹ ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$
1.101 $\pm 0.029 \pm 0.022$		PEDLAR	09	CLE3	$J/\psi \rightarrow \eta\eta$
1.123 ± 0.089	11k	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 $\pm 0.08 \pm 0.11$		BLOOM	83	CBAL	$e^+ e^-$
0.82 ± 0.10		BRANDELIK	79C	DASP	$e^+ e^-$
1.3 ± 0.4	21	BARTEL	77	CNTR	$e^+ e^-$

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_{243}/Γ
0.79 ± 0.13 OUR AVERAGE				

0.68 $\pm 0.04 \pm 0.24$	BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.76 $\pm 0.15 \pm 0.21$	^{1,2} AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
0.87 $\pm 0.14^{+0.14}_{-0.11}$	¹ BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Included unknown branching fraction $f_1(1420) \rightarrow K\bar{K}\pi$.

² From fit to the $K^*(892)K$ 1^{++} partial wave.

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_{244}/Γ
0.61 ± 0.08 OUR AVERAGE				

0.69 $\pm 0.16 \pm 0.20$	¹ BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\rho^0$
0.61 $\pm 0.04 \pm 0.21$	² BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 $\pm 0.09 \pm 0.17$	³ BAI	99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.625 $\pm 0.063 \pm 0.103$	⁴ BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.70 $\pm 0.08 \pm 0.16$	⁵ BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Assuming $B(f_1(1285) \rightarrow \rho^0\gamma) = 0.055 \pm 0.013$.

² Assuming $\Gamma(f_1(1285) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$.

³ Assuming $\Gamma(f_1(1285) \rightarrow \eta\pi\pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.

⁴ Obtained summing the sequential decay channels

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi\pi\pi\pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow \eta\pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980)\pi, a_0(980) \rightarrow K\bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;

$B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma\rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.

⁵ Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.

$\Gamma(\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{245}/Γ
$4.5 \pm 1.0 \pm 0.7$	BAI	99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M070B01;LINKAGE=A
NODE=M070B01;LINKAGE=AB

NODE=M070R01
NODE=M070R01

NODE=M070S84
NODE=M070S84

NODE=M070R83
NODE=M070R83

NODE=M070R83;LINKAGE=A

NODE=M070S31;LINKAGE=A
NODE=M070S31;LINKAGE=D

NODE=M070R88
NODE=M070R88

NODE=M070R88;LINKAGE=BI
NODE=M070R88;LINKAGE=BD
NODE=M070R88;LINKAGE=BA
NODE=M070R88;LINKAGE=B

NODE=M070R88;LINKAGE=A

NODE=M070S36
NODE=M070S36

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.7 $^{+0.8}_{-0.5}$ OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.0 \pm 0.9 \pm 0.2$	750	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		3 BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		3 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		3 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	4	4 BRANDELIK	79c DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78 PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$


1 Using CLEO-c data but not authored by the CLEO Collaboration.

2 DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.888$.

4 Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

Γ_{246}/Γ
NODE=M070R87
NODE=M070R87

 $\Gamma(\gamma f'_2(1525) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

7.99 $^{+0.03}_{-0.04} \pm 0.69$ -0.50

 $\Gamma(\gamma f'_2(1525) \rightarrow \gamma \eta\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.42 $^{+0.43}_{-0.51} \pm 1.37$ -1.30

1 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

 Γ_{247}/Γ

NODE=M070P69
NODE=M070P69

 $\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.28 $\pm 0.05 \pm 0.17$

 $\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.20 $\pm 0.04 \pm 0.13$

 Γ_{249}/Γ

NODE=M070R02
NODE=M070R02

 Γ_{250}/Γ

NODE=M070R03
NODE=M070R03

$\Gamma(\gamma f_0(1750) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{251}/Γ			
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	
$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$	NODE=M070P65 NODE=M070P65

$\Gamma(\gamma f_0(1800) \rightarrow \gamma \omega \phi)/\Gamma_{\text{total}}$	Γ_{252}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.6 OUR AVERAGE				
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma \omega \phi$
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma \omega \phi$

$\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$	Γ_{253}/Γ			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N	$J/\psi \rightarrow \gamma \eta \eta$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.


$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_{254}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.7 \pm 0.1 \pm 0.2$		BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$	Γ_{255}/Γ			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.0 \pm 0.3 \pm 1.3$	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

¹ Summed over all charges.

$\Gamma(\gamma \phi \phi)/\Gamma_{\text{total}}$	Γ_{256}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
$7.5 \pm 0.6 \pm 1.2$	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
$3.4 \pm 0.8 \pm 0.6$	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$3.1 \pm 0.7 \pm 0.4$		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ $\phi \phi$ mass less than 2.9 GeV, η_c excluded.



$\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}}$	Γ_{257}/Γ			
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	COMMENT
$0.38 \pm 0.07 \pm 0.07$		49	EATON	84 MRK2 $e^+ e^-$
<0.11		90	PERUZZI	78 MRK1 $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

$\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{258}/Γ
$3.14^{+0.50}_{-0.19}$ OUR AVERAGE					
$2.40 \pm 0.10^{+2.47}_{-0.18}$		1,2 ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
$4.4 \pm 0.4 \pm 0.8$	196	² ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$3.3 \pm 0.8 \pm 0.5$		² BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
$2.7 \pm 0.6 \pm 0.6$		² BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	
$2.4^{+1.5}_{-1.0}$		^{3,4} BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

² Includes unknown branching fraction to $\phi\phi$.

³ Estimated by us from various fits.

⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{259}/Γ
0.13 ± 0.09		1,2 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{260}/Γ
$1.98 \pm 0.08 \pm 0.32$	1045	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$	

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{261}/Γ
$<4.80 \times 10^{-6}$	90	ABLIKIM	180 BES3		$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$	

 $\Gamma(\gamma X(1835) \rightarrow \gamma\pi^+\pi^-\eta')/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{262}/Γ
$2.77^{+0.34}_{-0.40}$ OUR AVERAGE				Error includes scale factor of 1.1.	

$3.93 \pm 0.38^{+0.31}_{-0.84}$		1 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	
$2.87 \pm 0.09^{+0.49}_{-0.52}$	4265	² ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	
$2.2 \pm 0.4 \pm 0.4$	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	

¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatté formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two unconfirmed states $\gamma X(2120)$, and $\gamma X(2370)$, for $M(p\bar{p}) < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$.

 $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{263}/Γ
$0.77^{+0.15}_{-0.09}$ OUR AVERAGE					

$0.90^{+0.04+0.27}_{-0.11-0.55}$		1 ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$	
$1.14^{+0.43+0.42}_{-0.30-0.26}$	231	² ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$	
$0.70 \pm 0.04^{+0.19}_{-0.08}$		BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$	

¹ From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A.

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

NODE=M070S21
NODE=M070S21

NODE=M070S21;LINKAGE=C

OCCUR=2

NODE=M070S21;LINKAGE=U
NODE=M070S21;LINKAGE=A
NODE=M070S21;LINKAGE=B

NODE=M070S20
NODE=M070S20

NODE=M070S20;LINKAGE=A
NODE=M070S20;LINKAGE=B

NODE=M070R04
NODE=M070R04

NODE=M070P40
NODE=M070P40

NODE=M070R78
NODE=M070R78

NODE=M070R78;LINKAGE=A

NODE=M070R78;LINKAGE=AI

NODE=M070S71
NODE=M070S71

NODE=M070S71;LINKAGE=AK

NODE=M070S71;LINKAGE=AL

$\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$					Γ_{264}/Γ	
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>		<u>TECN</u>	<u>COMMENT</u>		NODE=M070S96 NODE=M070S96
$3.31^{+0.33+1.96}_{-0.30-1.29}$	ABLIKIM	15T	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$		
$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\phi(1020))/\Gamma_{\text{total}}$					Γ_{265}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P37 NODE=M070P37
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.77 \pm 0.35 \pm 0.25	305	¹ ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$	
8.09 \pm 1.99 \pm 1.36	1.3k	² ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$	
1 Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.						
2 Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.						
$\Gamma(\gamma X(1835) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{266}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070P41 NODE=M070P41
$<3.56 \times 10^{-6}$	90	ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$	
$\Gamma(\gamma X(1840) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$					Γ_{267}/Γ	
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S82 NODE=M070S82
$2.44^{+0.36+0.60}_{-0.74}$	0.6k	ABLIKIM	13U	BES3	$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$	
$\Gamma(\gamma(K\bar{K}\pi)[J^PC=0-+])/ \Gamma_{\text{total}}$					Γ_{268}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M070S38 NODE=M070S38
0.7 \pm 0.4 OUR AVERAGE	Error includes scale factor of 2.1.					
0.58 \pm 0.03 \pm 0.20	¹ BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$		
2.1 \pm 0.1 \pm 0.7	² BAI	00D	BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$		
1 For a broad structure around 1800 MeV.						
2 For a broad structure around 2040 MeV.						
$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$					Γ_{269}/Γ	
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070R82 NODE=M070R82
3.56 ± 0.17 OUR AVERAGE						
3.59 \pm 0.20 \pm 0.03	1.6k	¹ ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$	
3.63 \pm 0.36 \pm 0.13		PEDLAR	09	CLE3	$J/\psi \rightarrow \pi^0 \gamma$	
$3.13^{+0.65}_{-0.47}$	586	ABLIKIM	06E	BES2	$J/\psi \rightarrow \pi^0 \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
3.6 \pm 1.1 \pm 0.7		BLOOM	83	CBAL	$e^+ e^-$	
7.3 \pm 4.7	10	BRANDELIK	79C	DASP	$e^+ e^-$	
1 ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.						NODE=M070R82;LINKAGE=A
$\Gamma(\gamma p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{270}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070R93 NODE=M070R93
$<0.79 \times 10^{-3}$	90	EATON	84	MRK2	$e^+ e^-$	
$\Gamma(\gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{271}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S8 NODE=M070S8
$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<0.16 \times 10^{-3}$	90	BAI	98G	BES	$e^+ e^-$	
$\Gamma(\gamma f_0(2100) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$					Γ_{272}/Γ	
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M070S85 NODE=M070S85
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N	BES3	$J/\psi \rightarrow \gamma\eta\eta$	
1 From partial wave analysis including all possible combinations of 0^{++}, 2^{++}, and 4^{++} resonances.						NODE=M070S85;LINKAGE=A

$\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	COMMENT
6.24±0.48±0.87	744	1 DOBBS	15 $J/\psi \rightarrow \gamma\pi\pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 Γ_{273}/Γ

NODE=M070B08
NODE=M070B08

 $\Gamma(\gamma f_0(2200))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	1 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	------------	--------	---

¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

 Γ_{274}/Γ

NODE=M070S18
NODE=M070S18

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	COMMENT
5.86±0.49±1.20	490	1 DOBBS	15 $J/\psi \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 Γ_{275}/Γ

NODE=M070B09
NODE=M070B09

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.72±0.08±0.17	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300	1 BAI	96B BES	$e^+ e^- \rightarrow \gamma \bar{p}p, K\bar{K}$
>250	2 HASAN	96 SPEC	$\bar{p}p \rightarrow \pi^+ \pi^-$
< 2.3	3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	3 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$	23	3 BALTRUSAIT..86D MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$	93	3 BALTRUSAIT..86D MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

 Γ_{277}/Γ

NODE=M070R92
NODE=M070R92

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.9	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 8 ± 4	BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$
$8.4 \pm 2.6 \pm 3.0$	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

 Γ_{278}/Γ

NODE=M070B02
NODE=M070B02

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	1,2 DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6	3 DEL-AMO-SA..100 BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$	
< 2.9	3 DEL-AMO-SA..100 BABR	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$	
$6.6 \pm 2.9 \pm 2.4$	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

 Γ_{279}/Γ

NODE=M070B03
NODE=M070B03

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.5±0.6±0.5	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$

 Γ_{280}/Γ

NODE=M070B04
NODE=M070B04

$\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{281}/Γ	
<u>VALUE</u> (units 10^{-5})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
4.95±0.21^{+0.66}_{-0.72}		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$		NODE=M070P67 NODE=M070P67
$\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$					Γ_{282}/Γ	
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
5.60^{+0.62}_{-0.65}^{+2.37}_{-2.07}	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$		NODE=M070S88 NODE=M070S88
¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.						NODE=M070S88;LINKAGE=A
$\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{283}/Γ	
<u>VALUE</u> (units 10^{-5})		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
5.54^{+0.34}_{-0.40}^{+3.82}_{-1.49}		ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$		NODE=M070P70 NODE=M070P70
$\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$					Γ_{284}/Γ	
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.09±0.24 OUR AVERAGE						NODE=M070S32 NODE=M070S32
1.21±0.29±0.24	174	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$		
1.00±0.03±0.45		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$		
1.02±0.09±0.45		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$		OCCUR=2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
5.7 ± 0.8		3.4 BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$		
¹ Using CLEO-c data but not authored by the CLEO Collaboration.						
² Including unknown branching fraction to $\pi \pi$.						
³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.						
⁴ Assuming that $f_0(1500)$ decays only to two S -wave dipions.						
$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$					Γ_{285}/Γ	
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.65^{+0.26}_{-0.31}^{+0.51}_{-1.40}	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$		NODE=M070S83 NODE=M070S83
¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.						NODE=M070S83;LINKAGE=A
$\Gamma(\gamma A \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ (narrow state A with $m_A < 960$ MeV)					Γ_{286}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<6.3 × 10⁻⁶	90	¹ INSLER	10	CLEO $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$		NODE=M070S68 NODE=M070S68
¹ The limit varies with mass m_A of a narrow state A and is 4.3×10^{-6} for $m_A = 0$ MeV, reaches its largest value of 6.3×10^{-6} at $m_A = 500$ MeV, and is 3.6×10^{-6} at $m_A = 960$ MeV.						NODE=M070S68;LINKAGE=IN
$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ (narrow state A^0 with 0.2 GeV $< m_{A^0} < 3$ GeV)					Γ_{287}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.5 × 10⁻⁵	90	¹ ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$		NODE=M070S76 NODE=M070S76
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<2.1 × 10 ⁻⁵	90	² ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$		
¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.						NODE=M070S76;LINKAGE=A
² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .						NODE=M070S76;LINKAGE=AB
DALITZ DECAYS						
$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$					Γ_{288}/Γ	
<u>VALUE</u> (units 10^{-7})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
7.56±1.32±0.50	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$		NODE=M070S89 NODE=M070S89

$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$					Γ_{289}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070S90 NODE=M070S90
$1.43 \pm 0.04 \pm 0.06$	2.47k	1,2 ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM	14I BES3	$J/\psi \rightarrow \eta e^+ e^-$		
1 Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.						NODE=M070S90;LINKAGE=A
2 Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.84 \pm 0.11 \pm 0.08$ GeV. Supersedes ABLIKIM 14I.						NODE=M070S90;LINKAGE=C
$\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$					Γ_{290}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M070S91 NODE=M070S91
$6.59 \pm 0.07 \pm 0.17$	8.9k	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$5.81 \pm 0.16 \pm 0.31$	1.4k	1,2 ABLIKIM	14I BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$		
1 Using both $\eta' \rightarrow \gamma\pi^+\pi^-$ and $\eta' \rightarrow \pi^+\pi^-\eta$ decays.						NODE=M070S91;LINKAGE=A
2 Superseded by ABLIKIM 19H.						NODE=M070S91;LINKAGE=B
$\Gamma(\eta U \rightarrow \eta e^+ e^-)/\Gamma_{\text{total}}$					Γ_{291}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070P42 NODE=M070P42
$<9.11 \times 10^{-7}$	90	¹ ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$		
1 For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .						NODE=M070P42;LINKAGE=A
$\Gamma(\eta'(958) U \rightarrow \eta'(958) e^+ e^-)/\Gamma_{\text{total}}$					Γ_{292}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070P61 NODE=M070P61
$<2.0 \times 10^{-7}$	90	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$		
1 For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .						NODE=M070P61;LINKAGE=A
$\Gamma(\phi e^+ e^-)/\Gamma_{\text{total}}$					Γ_{293}/Γ	
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070P82 NODE=M070P82
<1.2	90	¹ ABLIKIM	19AB BES3	$J/\psi \rightarrow \phi e^+ e^-$		
1 Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) = (34.49 \pm 0.30)\%$.						NODE=M070P82;LINKAGE=A
WEAK DECAYS						NODE=M070320
$\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{294}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070S53 NODE=M070S53
$<1.2 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$		
$\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{295}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070S54 NODE=M070S54
$<8.5 \times 10^{-8}$	90	¹ ABLIKIM	17AF BES3	$e^+ e^- \rightarrow J/\psi$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<1.1 \times 10^{-5}$	90	ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$		
1 Using D^0 decays to $K^-\pi^+$, $K^-\pi^+\pi^0$, and $K^-\pi^+\pi^+\pi^-$.						NODE=M070S54;LINKAGE=A
$\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{296}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070S55 NODE=M070S55
$<1.3 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
$<3.6 \times 10^{-5}$	90	¹ ABLIKIM	06M BES2	$e^+ e^- \rightarrow J/\psi$		
1 Using $B(D_s^- \rightarrow \phi\pi^-) = 4.4 \pm 0.5\%$.						NODE=M070S55;LINKAGE=AB
$\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{297}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M070B13 NODE=M070B13
$<1.8 \times 10^{-6}$	90	ABLIKIM	14R BES3	$e^+ e^- \rightarrow J/\psi$		

$\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$<7.5 \times 10^{-5}$	90

 $\Gamma(\bar{D}^0 K^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$<1.7 \times 10^{-4}$	90

 $\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$<2.5 \times 10^{-6}$	90

 $\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$<1.3 \times 10^{-4}$	90

 $\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$<1.3 \times 10^{-5}$	90

 Γ_{298}/Γ NODE=M070S61
NODE=M070S61 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	CL%
$< 2.7 \times 10^{-7}$	90

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 0.5 \times 10^{-5}$	90	ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$< 1.6 \times 10^{-4}$	90	¹ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
$< 2.2 \times 10^{-5}$	90	ABLIKIM	07J	BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$< 50 \times 10^{-5}$	90	BARTEL	77	CNTR	$e^+ e^-$

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.006 \times 10^{-3}$.

 Γ_{303}/Γ NODE=M070R80
NODE=M070R80 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.4 \times 10^{-6}$	90

 Γ_{304}/Γ NODE=M070S95
NODE=M070S95 $\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$

VALUE	CL%
$< 1.6 \times 10^{-7}$	90

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	CL%
$< 1.1 \times 10^{-6}$	90

 Γ_{305}/Γ NODE=M070S39
NODE=M070S39 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$

VALUE	CL%
$< 8.3 \times 10^{-6}$	90

 Γ_{306}/Γ NODE=M070S40
NODE=M070S40 $\Gamma(A_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%
$< 6.9 \times 10^{-8}$	90

 Γ_{308}/Γ NODE=M070P74
NODE=M070P74 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$

VALUE	CL%
$< 2.0 \times 10^{-6}$	90

 Γ_{307}/Γ NODE=M070S41
NODE=M070S41 $\Gamma(\text{invisible})/\Gamma(e^+ e^-)$

VALUE	CL%
$< 6.6 \times 10^{-2}$	90

 Γ_{309}/Γ_5 NODE=M070S80
NODE=M070S80**OTHER DECAYS****DOCUMENT ID****TECN****COMMENT****LEES****BABR** $B \rightarrow K^{(*)} J/\psi$

$\Gamma(\text{invisible})/\Gamma(\mu^+\mu^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{309}/Γ_7
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G	BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$

 $J/\psi(1S)$ REFERENCES

ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60048
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59517
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59846
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59850
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59857
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59890
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59604
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59615
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59747
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59773
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)	REFID=59614
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59456
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58849
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=59102
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58315
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58317
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58324
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57903
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57967
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57966
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57990
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57265
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57454
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57509
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57510
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57511
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)	REFID=57512
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57522
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57566
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56773
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56776
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
ABLIKIM	15T	PR D115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56785
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55900
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55902
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55905
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56238
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56388
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=55655
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54920
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54954
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54955
ABLIKIM	13L	PR D87 112007	Ablikim M. <i>et al.</i>	(BESIII Collab.)	REFID=55300
ABLIKIM	13N	PR D87 092009	Ablikim M. <i>et al.</i>	(BESIII Collab.)	REFID=55387
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55392
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55402
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54265
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54267
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54268
ABLIKIM	12D	PR D108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54269
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54273
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54863
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16715
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53220
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53533
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)	REFID=53359
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52718
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53099
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52047

ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52128
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52253
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52255
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52256
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(CLEO Collab.)	REFID=52571
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(BABAR Collab.)	REFID=52261
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(CLEO Collab.)	REFID=52242
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(PDG Collab.)	REFID=52685
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(BELLE Collab.)	REFID=52166
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BES Collab.)	REFID=52204
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52072
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50986
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51058
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51128
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51130
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51036
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50496
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50507
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)	REFID=50802
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer		REFID=51038
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49739
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49611
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49620
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49607
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49751
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49754
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49403
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47427
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98E	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46341
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)	REFID=46608
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45146
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44736
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45198
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)	REFID=45197
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44434
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)	REFID=43314
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)	REFID=43601
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41866
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)	REFID=41899
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41359
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350

JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)	REFID=41349
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=40346
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAZO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LAZO, CLER, FRAS+)	REFID=40268
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)	REFID=40002
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LAZO+)	REFID=40261
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LAZO, PADO)	REFID=40243
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BALTRUSAIT...	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)	REFID=21865
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=22101
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
BALTRUSAIT...	85C	PR D55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095
BALTRUSAIT...	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22097
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
BALTRUSAIT...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22006
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)	REFID=22092
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22080
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)	REFID=22077
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22073
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
Translated from YAF 34 1471.					
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22065
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MANZ)	REFID=22066
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22067
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)	REFID=22068
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22058
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22060
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)	REFID=22063
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22054
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG	REFID=22056
BALDINI...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)	REFID=22026
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC	REFID=22030
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22036
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=22038
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)	REFID=22039

See the related review(s):

Branching Ratios of $\psi(2S)$ and $\chi_{c,1,2}^0$

NODE=M056

 $\chi_{c0}(1P)$ $I^G(J^{PC}) = 0^+(0^{++})$ $\chi_{c0}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3414.71 ± 0.30 OUR AVERAGE				
3413.0 ± 1.9 ± 0.6	933	¹ AAIJ	17BB LHCb	$p\bar{p} \rightarrow b\bar{b} X \rightarrow 2(K^+K^-)X$
3414.2 ± 0.5 ± 2.3	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
3406 ± 7 ± 6	230	² ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
3414.21 ± 0.39 ± 0.27		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
3414.7 ± 0.7 ± 0.2		³ ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
3415.5 ± 0.4 ± 0.4	392	⁴ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
3417.4 ± 1.8 ± 0.2		³ AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
3414.1 ± 0.6 ± 0.8		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3417.8 ± 0.4 ± 4		³ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3416 ± 3 ± 4		⁵ TANENBAUM	78 MRK1	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3414.6 ± 1.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
3416.5 ± 3.0		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c0}$
3422 ± 10		⁵ BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3415 ± 9		⁵ BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$

¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c0}(1P)$ fixed to the PDG 16 value.

² From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

³ Using mass of $\psi(2S) = 3686.0$ MeV.

⁴ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

⁵ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

NODE=M056M

NODE=M056M

 $\chi_{c0}(1P)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.8 ± 0.6 OUR FIT				
10.5 ± 0.8 OUR AVERAGE Error includes scale factor of 1.1.				
10.6 ± 1.9 ± 2.6	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
12.6 ^{+1.5} _{-1.6} ^{+0.9} _{-1.1}		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
8.6 ^{+1.7} _{-1.3} ^{+0.1}		ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
9.7 ± 1.0	392	¹ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
16.6 ^{+5.2} _{-3.7} ^{+0.1}		AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
14.3 ± 2.0 ± 3.0		BAI	98I BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
13.5 ± 3.3 ± 4.2		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X, \gamma\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13.2 ± 2.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ Recalculated by ANDREOTTI 05A.

NODE=M056M;LINKAGE=A

NODE=M056M;LINKAGE=EB

NODE=M056M;LINKAGE=C

NODE=M056M;LINKAGE=NW

NODE=M056M;LINKAGE=D

NODE=M056W

NODE=M056W

 $\chi_{c0}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	(2.34 ± 0.18) %	
Γ_2 $\rho^0\pi^+\pi^-$	(9.1 ± 2.9) × 10 ⁻³	
Γ_3 $\rho^0\rho^0$		
Γ_4 $f_0(980)f_0(980)$	(6.6 ± 2.1) × 10 ⁻⁴	
Γ_5 $\pi^+\pi^-\pi^0\pi^0$	(3.3 ± 0.4) %	
Γ_6 $\rho^+\pi^-\pi^0 + \text{c.c.}$	(2.9 ± 0.4) %	
Γ_7 $4\pi^0$	(3.3 ± 0.4) × 10 ⁻³	
Γ_8 $\pi^+\pi^-K^+K^-$	(1.81 ± 0.14) %	

NODE=M056;CLUMP=A

DESIG=3

DESIG=9

DESIG=54

DESIG=20

DESIG=61

DESIG=62

DESIG=70

DESIG=5

NODE=M056W;LINKAGE=AN

NODE=M056215;NODE=M056

Γ_9	$K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-$	$(9.8 \pm 4.0) \times 10^{-4}$		DESIG=31
Γ_{10}	$K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(8.0 \pm 2.0) \times 10^{-4}$		DESIG=32
Γ_{11}	$K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(6.3 \pm 1.9) \times 10^{-3}$		DESIG=33
Γ_{12}	$K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$< 2.7 \times 10^{-3}$	CL=90%	DESIG=34
Γ_{13}	$f_0(980) f_0(980)$	$(1.6 \pm 1.0) \times 10^{-4}$		DESIG=23
Γ_{14}	$f_0(980) f_0(2200)$	$(7.9 \pm 2.0) \times 10^{-4}$		DESIG=24
Γ_{15}	$f_0(1370) f_0(1370)$	$< 2.7 \times 10^{-4}$	CL=90%	DESIG=25
Γ_{16}	$f_0(1370) f_0(1500)$	$< 1.7 \times 10^{-4}$	CL=90%	DESIG=26
Γ_{17}	$f_0(1370) f_0(1710)$	$(6.7 \pm 3.5) \times 10^{-4}$		DESIG=27
Γ_{18}	$f_0(1500) f_0(1370)$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=28
Γ_{19}	$f_0(1500) f_0(1500)$	$< 5 \times 10^{-5}$	CL=90%	DESIG=29
Γ_{20}	$f_0(1500) f_0(1710)$	$< 7 \times 10^{-5}$	CL=90%	DESIG=30
Γ_{21}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(8.6 \pm 0.9) \times 10^{-3}$		DESIG=75
Γ_{22}	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(4.2 \pm 0.4) \times 10^{-3}$		DESIG=87
Γ_{23}	$K^+ K^- \pi^0 \pi^0$	$(5.6 \pm 0.9) \times 10^{-3}$		DESIG=63
Γ_{24}	$K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(2.49 \pm 0.33) \%$		DESIG=65
Γ_{25}	$\rho^+ K^- K^0 + \text{c.c.}$	$(1.21 \pm 0.21) \%$		DESIG=66
Γ_{26}	$K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(4.6 \pm 1.2) \times 10^{-3}$		DESIG=67
Γ_{27}	$K_S^0 K_S^0 \pi^+ \pi^-$	$(5.7 \pm 1.1) \times 10^{-3}$		DESIG=41
Γ_{28}	$K^+ K^- \eta \pi^0$	$(3.0 \pm 0.7) \times 10^{-3}$		DESIG=68
Γ_{29}	$3(\pi^+ \pi^-)$	$(1.20 \pm 0.18) \%$		DESIG=4
Γ_{30}	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(7.5 \pm 1.6) \times 10^{-3}$		DESIG=10
Γ_{31}	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.7 \pm 0.6) \times 10^{-3}$		DESIG=21
Γ_{32}	$\pi \pi$	$(8.51 \pm 0.33) \times 10^{-3}$		DESIG=18
Γ_{33}	$\pi^0 \eta$	$< 1.8 \times 10^{-4}$		DESIG=35
Γ_{34}	$\pi^0 \eta'$	$< 1.1 \times 10^{-3}$		DESIG=36
Γ_{35}	$\pi^0 \eta_c$	$< 1.6 \times 10^{-3}$	CL=90%	DESIG=86
Γ_{36}	$\eta \eta$	$(3.01 \pm 0.19) \times 10^{-3}$		DESIG=13
Γ_{37}	$\eta \eta'$	$(9.1 \pm 1.1) \times 10^{-5}$		DESIG=37
Γ_{38}	$\eta' \eta'$	$(2.17 \pm 0.12) \times 10^{-3}$		DESIG=46
Γ_{39}	$\omega \omega$	$(9.7 \pm 1.1) \times 10^{-4}$		DESIG=22
Γ_{40}	$\omega \phi$	$(1.41 \pm 0.13) \times 10^{-4}$		DESIG=76
Γ_{41}	$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$		DESIG=88
Γ_{42}	$K^+ K^-$	$(6.05 \pm 0.31) \times 10^{-3}$		DESIG=2
Γ_{43}	$K_S^0 K_S^0$	$(3.16 \pm 0.17) \times 10^{-3}$		DESIG=15
Γ_{44}	$\pi^+ \pi^- \eta$	$< 2.0 \times 10^{-4}$	CL=90%	DESIG=50
Γ_{45}	$\pi^+ \pi^- \eta'$	$< 4 \times 10^{-4}$	CL=90%	DESIG=53
Γ_{46}	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$< 9 \times 10^{-5}$	CL=90%	DESIG=17
Γ_{47}	$K^+ K^- \pi^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=47
Γ_{48}	$K^+ K^- \eta$	$< 2.3 \times 10^{-4}$	CL=90%	DESIG=51
Γ_{49}	$K^+ K^- K_S^0 K_S^0$	$(1.4 \pm 0.5) \times 10^{-3}$		DESIG=42
Γ_{50}	$K_S^0 K_S^0 K_S^0 K_S^0$	$(5.8 \pm 0.5) \times 10^{-4}$		DESIG=94
Γ_{51}	$K^+ K^- K^+ K^-$	$(2.82 \pm 0.29) \times 10^{-3}$		DESIG=14
Γ_{52}	$K^+ K^- \phi$	$(9.7 \pm 2.5) \times 10^{-4}$		DESIG=44
Γ_{53}	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.7 \pm 0.6) \times 10^{-3}$		DESIG=91
Γ_{54}	$K^+ K^- \pi^0 \phi$	$(1.90 \pm 0.35) \times 10^{-3}$		DESIG=92
Γ_{55}	$\phi \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.15) \times 10^{-3}$		DESIG=89
Γ_{56}	$\phi \phi$	$(8.0 \pm 0.7) \times 10^{-4}$		DESIG=16
Γ_{57}	$\phi \phi \eta$	$(8.4 \pm 1.0) \times 10^{-4}$		DESIG=96
Γ_{58}	$p \bar{p}$	$(2.21 \pm 0.08) \times 10^{-4}$		DESIG=11
Γ_{59}	$p \bar{p} \pi^0$	$(7.0 \pm 0.7) \times 10^{-4}$	S=1.3	DESIG=48

Γ_{60}	$p\bar{p}\eta$	$(3.5 \pm 0.4) \times 10^{-4}$	DESIG=52
Γ_{61}	$p\bar{p}\omega$	$(5.2 \pm 0.6) \times 10^{-4}$	DESIG=69
Γ_{62}	$p\bar{p}\phi$	$(6.0 \pm 1.4) \times 10^{-5}$	DESIG=74
Γ_{63}	$p\bar{p}\pi^+\pi^-$	$(2.1 \pm 0.7) \times 10^{-3}$	S=1.4
Γ_{64}	$p\bar{p}\pi^0\pi^0$	$(1.04 \pm 0.28) \times 10^{-3}$	DESIG=8
Γ_{65}	$p\bar{p}K^+K^-$ (non-resonant)	$(1.22 \pm 0.26) \times 10^{-4}$	DESIG=64
Γ_{66}	$p\bar{p}K_S^0K_S^0$	$< 8.8 \times 10^{-4}$	DESIG=71
Γ_{67}	$p\bar{n}\pi^-$	$(1.27 \pm 0.11) \times 10^{-3}$	DESIG=40
Γ_{68}	$\bar{p}n\pi^+$	$(1.37 \pm 0.12) \times 10^{-3}$	DESIG=43
Γ_{69}	$p\bar{n}\pi^-\pi^0$	$(2.34 \pm 0.21) \times 10^{-3}$	DESIG=82
Γ_{70}	$\bar{p}n\pi^+\pi^0$	$(2.21 \pm 0.18) \times 10^{-3}$	DESIG=83
Γ_{71}	$\Lambda\bar{\Lambda}$	$(3.27 \pm 0.24) \times 10^{-4}$	DESIG=84
Γ_{72}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(1.18 \pm 0.13) \times 10^{-3}$	DESIG=19
Γ_{73}	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$< 5 \times 10^{-4}$	DESIG=38
Γ_{74}	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	$< 5 \times 10^{-4}$	DESIG=77
Γ_{75}	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 5 \times 10^{-4}$	DESIG=78
Γ_{76}	$K^+\bar{p}\Lambda + \text{c.c.}$	$(1.25 \pm 0.12) \times 10^{-3}$	DESIG=79
Γ_{77}	$K^*(892)^+\bar{p}\Lambda + \text{c.c.}$	$(4.8 \pm 0.9) \times 10^{-4}$	DESIG=49
Γ_{78}	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	$(2.9 \pm 0.7) \times 10^{-4}$	DESIG=98
Γ_{79}	$\Lambda(1520)\bar{\Lambda}(1520)$	$(3.1 \pm 1.2) \times 10^{-4}$	DESIG=72
Γ_{80}	$\Sigma^0\bar{\Sigma}^0$	$(4.68 \pm 0.32) \times 10^{-4}$	DESIG=73
Γ_{81}	$\Sigma^+\bar{p}K_S^0 + \text{c.c.}$	$(3.52 \pm 0.27) \times 10^{-4}$	DESIG=58
Γ_{82}	$\Sigma^+\bar{\Sigma}^-$	$(4.6 \pm 0.8) \times 10^{-4}$	DESIG=97
Γ_{83}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(1.6 \pm 0.6) \times 10^{-4}$	S=2.6
Γ_{84}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$(2.3 \pm 0.7) \times 10^{-4}$	DESIG=80
Γ_{85}	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.94 \pm 0.35) \times 10^{-4}$	DESIG=81
Γ_{86}	$\Xi^0\bar{\Xi}^0$	$(3.1 \pm 0.8) \times 10^{-4}$	DESIG=85
Γ_{87}	$\Xi^-\bar{\Xi}^+$	$(4.8 \pm 0.7) \times 10^{-4}$	DESIG=60
Γ_{88}	$\eta_c\pi^+\pi^-$	$< 7 \times 10^{-4}$	DESIG=39
	Radiative decays		
Γ_{89}	$\gamma J/\psi(1S)$	$(1.40 \pm 0.05) \%$	DESIG=90%
Γ_{90}	$\gamma\rho^0$	$< 9 \times 10^{-6}$	DESIG=55
Γ_{91}	$\gamma\omega$	$< 8 \times 10^{-6}$	DESIG=56
Γ_{92}	$\gamma\phi$	$< 6 \times 10^{-6}$	DESIG=57
Γ_{93}	$\gamma\gamma$	$(2.04 \pm 0.09) \times 10^{-4}$	DESIG=7
Γ_{94}	$e^+e^-J/\psi(1S)$	$(1.33 \pm 0.29) \times 10^{-4}$	DESIG=93
Γ_{95}	$\mu^+\mu^-J/\psi(1S)$	$< 1.9 \times 10^{-5}$	DESIG=95

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 378.1$ for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_2	24								
x_8	9	2							
x_{30}	5	1	28						
x_{32}	8	2	10	3					
x_{36}	4	1	5	1	14				
x_{42}	8	2	8	3	18	11			
x_{43}	7	2	8	2	18	10	14		
x_{51}	5	1	5	2	9	5	7	7	
x_{56}	7	2	6	2	9	5	7	7	4
x_{58}	3	1	4	1	3	-1	7	7	3
x_{71}	4	1	5	1	13	8	10	10	5
x_{89}	5	1	6	2	17	11	13	12	6
x_{93}	-8	-2	-2	-3	14	9	10	10	1
Γ	-26	-6	-19	-10	-15	-7	-14	-12	-10
		x_1	x_2	x_8	x_{30}	x_{32}	x_{36}	x_{42}	x_{43}
									x_{51}
									x_{56}
x_{71}		5							
x_{89}		-19	9						
x_{93}		6	9	13					
Γ		-4	-7	-9	-38				
		x_{58}	x_{71}	x_{89}	x_{93}				

$\chi_{c0}(1P)$ PARTIAL WIDTHS

— $\chi_{c0}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$ —

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}}$	$\Gamma_{58} \Gamma_{89} / \Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.6 ± 2.3 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

26.6 ± 2.6 ± 1.4	392	1,2 BAGNASCO	02	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$
48.7 ^{+11.3} _{-8.9} ± 2.4		1,2 AMBROGIANI	99B	E835	$\bar{p}p \rightarrow \gamma J/\psi$

1 Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

2 Values in $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$ and $(\Gamma(p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$ are not independent. The latter is used in the fit since it is less correlated to the total width.

— $\chi_{c0}(1P) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$ —

$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$	$\Gamma_1 \Gamma_{93} / \Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
52 ± 4 OUR FIT				

49 ± 10 OUR AVERAGE Error includes scale factor of 1.8.

44.7 ± 3.6 ± 4.9	3.6k	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$
75 ± 13 ± 8		EISENSTEIN	01	CLE2	$e^+ e^- \rightarrow e^+ e^- \chi_{c0}$

$\Gamma(\rho^0 \rho^0) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$	$\Gamma_3 \Gamma_{93} / \Gamma$				
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<12	90	<252	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$

NODE=M056217

NODE=M056223

NODE=M056G1

NODE=M056G1

NODE=M056G;LINKAGE=7A

NODE=M056G;LINKAGE=KS

NODE=M056224

NODE=M056G2

NODE=M056G2

NODE=M056G07

NODE=M056G07

$\Gamma(\pi^+\pi^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G08 NODE=M056G08
40.0±3.5 OUR FIT						
38.8±3.7±4.7	1.7k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^-$		
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{21}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G01 NODE=M056G01
26±4±4	1094	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{30}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G09 NODE=M056G09
16 ±4 OUR FIT						
16.7±6.1±3.0	495 ± 182	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^-$		
$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{31}\Gamma_{93}/\Gamma$	
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	NODE=M056G10 NODE=M056G10
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<6	90	<148	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{32}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G3 NODE=M056G3
18.8± 1.3 OUR FIT						
23 ± 5 OUR AVERAGE						
29.7 ^{+17.4} _{-12.0} ± 4.8	103 ⁺⁶⁰ ₋₄₂	1 UEHARA	09	BELL $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$		
22.7 ± 3.2 ± 3.5	129 ± 18	2 NAKAZAWA	05	BELL $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$		
1 We multiplied the measurement by 3 to convert from $\pi^0 \pi^0$ to $\pi\pi$. Interference with the continuum included.						
2 We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.						
$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{36}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G06 NODE=M056G06
9.4±2.3±1.2	22	1 UEHARA	10A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$		
1 Interference with the continuum not included.						
$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{39}\Gamma_{93}/\Gamma$	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M056G02 NODE=M056G02
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<3.9	90	1 LIU	12B	BELL $\gamma\gamma \rightarrow 2(\pi^+ \pi^- \pi^0)$		
1 Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.						
$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{40}\Gamma_{93}/\Gamma$	
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M056G03 NODE=M056G03
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.34	90	1 LIU	12B	BELL $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
1 Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.						
$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{42}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G4 NODE=M056G4
13.4±1.0 OUR FIT						
14.3±1.6±2.3	153 ± 17	NAKAZAWA	05	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$		
$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{43}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G5 NODE=M056G5
7.0 ±0.5 OUR FIT						
8.7 ±1.7 ±0.9	266	1 UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
7.00±0.65±0.71	134 ± 12	CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$		
1 Supersedes CHEN 07B.						
$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{51}\Gamma_{93}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M056G11 NODE=M056G11
6.2±0.7 OUR FIT						
7.9±1.3±1.1	215 ± 36	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+ K^-)$		

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{56}\Gamma_{93}/\Gamma$				
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.76±0.18 OUR FIT					
1.72±0.33±0.14	56 ± 11	1 LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.3 ± 0.9 ± 0.4	23.6 ± 9.6	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+K^-)$	
1 Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.					

$\chi_{c0}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>		
0.0234±0.0018 OUR FIT			

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$	Γ_2/Γ_1		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.39±0.12 OUR FIT			
0.39±0.12	TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>		
0.0091±0.0029 OUR FIT			

$\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}$	Γ_4/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
6.6±2.1±0.1					
36 ± 9	1 ABLIKIM	04G BES		$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	
1 ABLIKIM 04G reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (6.5 \pm 1.6 \pm 1.3) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_5/Γ				
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.3±0.4±0.1					
1751.4	1 HE	08B CLEO		$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$	
1 HE 08B reports $3.54 \pm 0.10 \pm 0.43 \pm 0.18\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}$	Γ_6/Γ				
<u>VALUE (%)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.9±0.4±0.1					
1358.5	1,2 HE	08B CLEO		$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$	
1 HE 08B reports $3.04 \pm 0.18 \pm 0.42 \pm 0.16\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$	Γ_7/Γ				
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.3±0.4±0.1					
3296	1 ABLIKIM	11A BES3		$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	
1 ABLIKIM 11A reports $(3.34 \pm 0.06 \pm 0.44) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$	Γ_8/Γ		
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>		
18.1±1.4 OUR FIT			

NODE=M056G12
NODE=M056G12

NODE=M056G12;LINKAGE=LI

NODE=M056220

NODE=M056305

NODE=M056R2
NODE=M056R2

NODE=M056R54
NODE=M056R54

NODE=M056R24
NODE=M056R24

NODE=M056R62;LINKAGE=HE

NODE=M056R63
NODE=M056R63

NODE=M056R63;LINKAGE=HE

NODE=M056R71
NODE=M056R71

NODE=M056R71;LINKAGE=AB

NODE=M056R3
NODE=M056R3

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(\pi^+ \pi^- K^+ K^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{30}/Γ
0.41±0.09 OUR FIT				NODE=M056R55 NODE=M056R55
0.41±0.10	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$	

 $\Gamma(K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
9.8^{+3.6}_{-2.8}±0.2	83	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $(10.44 \pm 2.37^{+3.05}_{-1.90}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_0^*(1430)^0 \bar{K}_0^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
8.0^{+2.0}_{-2.4}±0.2	62	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $(8.49 \pm 1.66^{+1.32}_{-1.99}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0 \bar{K}_0^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
6.3^{+1.9}_{-1.0}±0.1	68	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $(6.66 \pm 1.31^{+1.60}_{-1.51}) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The measurement assumes $B(K_1(1270) \rightarrow K_P(770)) = 42 \pm 6\%$.

 $\Gamma(K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
<2.7	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $< 2.85 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$. The measurement assumes $B(K_1(1400) \rightarrow K^*(892)\pi) = 94 \pm 6\%$.

 $\Gamma(f_0(980) f_0(980))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
16.2^{+10.4}_{-9.0}±0.3	28	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980) f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.59 \pm 0.50^{+0.89}_{-0.72}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. One of the $f_0(980)$ mesons is identified via decay to $\pi^+ \pi^-$ while the other via $K^+ K^-$ decay.

 $\Gamma(f_0(980) f_0(2200))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
7.9^{+2.0}_{-2.5}±0.2	77	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	

NODE=M056R55

NODE=M056R55

NODE=M056R36

NODE=M056R36

NODE=M056R36;LINKAGE=AB

NODE=M056R37

NODE=M056R37

NODE=M056R38

NODE=M056R38

NODE=M056R38;LINKAGE=AB

NODE=M056R39

NODE=M056R39

NODE=M056R39;LINKAGE=AB

NODE=M056R28

NODE=M056R28

NODE=M056R28;LINKAGE=AB

NODE=M056R29

NODE=M056R29

¹ ABLIKIM 05Q reports $(8.42 \pm 1.42^{+1.65}_{-2.29}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(2200)) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(980) \rightarrow \pi^+\pi^-$ and $f_0(2200) \rightarrow K^+K^-$ decays.

$\Gamma(f_0(1370)f_0(1370))/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	1 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
¹ ABLIKIM 05Q reports $< 2.9 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1370)) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$. One of the $f_0(1370)$ mesons is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

$\Gamma(f_0(1370)f_0(1500))/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	1 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
¹ ABLIKIM 05Q reports $< 1.8 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1500)) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+\pi^-$ and $f_0(1500) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

$\Gamma(f_0(1370)f_0(1710))/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.7^{+3.5}_{-2.3} \pm 0.1$	61	1 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
¹ ABLIKIM 05Q reports $(7.12 \pm 1.85^{+3.28}_{-1.68}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1710)) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+\pi^-$ and $f_0(1710) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

$\Gamma(f_0(1500)f_0(1370))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	1 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
¹ ABLIKIM 05Q reports $< 1.4 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1370)) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+\pi^-$ and $f_0(1370) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

$\Gamma(f_0(1500)f_0(1500))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	1 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
¹ ABLIKIM 05Q reports $< 0.55 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1500)) / \Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$. One of the $f_0(1500)$ is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

NODE=M056R29;LINKAGE=AB

NODE=M056R30
NODE=M056R30

NODE=M056R30;LINKAGE=AB

NODE=M056R31
NODE=M056R31

NODE=M056R31;LINKAGE=AB

NODE=M056R32
NODE=M056R32

NODE=M056R32;LINKAGE=AB

NODE=M056R33
NODE=M056R33

NODE=M056R33;LINKAGE=AB

NODE=M056R34
NODE=M056R34

NODE=M056R34;LINKAGE=AB

$\Gamma(f_0(1500)f_0(1710))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	1 ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
1 ABLIKIM 05Q reports $< 0.73 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+\pi^-$ and $f_0(1710) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

 Γ_{20}/Γ NODE=M056R35
NODE=M056R35 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.61±0.13±0.94	9.0k	1 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$.				

 Γ_{21}/Γ NODE=M056R85
NODE=M056R85 $\Gamma(K_S^0K^\pm\pi^\mp\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.22±0.10±0.43	2.7k	1 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$.				

 Γ_{22}/Γ NODE=M056R86
NODE=M056R86 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.56±0.09±0.01	213.5	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

 Γ_{23}/Γ NODE=M056R64
NODE=M056R64

1 HE 08B reports $0.59 \pm 0.05 \pm 0.08 \pm 0.03\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.49±0.33±0.05	401.7	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

 Γ_{24}/Γ NODE=M056R66
NODE=M056R66

1 HE 08B reports $2.64 \pm 0.15 \pm 0.31 \pm 0.14\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\rho^+K^-K^0+c.c.)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.21±0.21±0.02	179.7	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

 Γ_{25}/Γ NODE=M056R67
NODE=M056R67

1 HE 08B reports $1.28 \pm 0.16 \pm 0.15 \pm 0.07\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+K^-K^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^-\bar{K}^+\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.46±0.12±0.01	64.1	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

 Γ_{26}/Γ NODE=M056R68
NODE=M056R68

1 HE 08B reports $0.49 \pm 0.10 \pm 0.07 \pm 0.03\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^-\bar{K}^+\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_S^0K_S^0\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.7±1.0±0.1	152 ± 14	1 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$

 Γ_{27}/Γ NODE=M056R47
NODE=M056R47

1 ABLIKIM 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0K_S^0\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.558 \pm 0.051 \pm 0.089) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R47;LINKAGE=AB

$\Gamma(K^+ K^- \eta\pi^0)/\Gamma_{\text{total}}$	Γ_{28}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.07±0.01	56.4	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.32 \pm 0.05 \pm 0.05 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$	Γ_{29}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
12.0±1.8 OUR EVALUATION	Treating systematic error as correlated.		

$\Gamma(12.0\pm1.7\text{ OUR AVERAGE})$	Γ_{29}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
11.7±1.0±1.9	1 BAI 99B BES	$\psi(2S) \rightarrow \gamma\chi_{c0}$	
12.5±2.9±0.5	1 TANENBAUM 78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$	

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{30}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.0075±0.0016 OUR FIT			

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$	Γ_{31}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.72±0.60±0.04	64 1 ABLIKIM 05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+ \pi^- K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.56±0.40±0.03 30 ± 6 2,3 ABLIKIM 04H BES Repl. by ABLIKIM 05Q

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.168 \pm 0.035^{+0.047}_{-0.040}) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

3 ABLIKIM 04H reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.53 \pm 0.29 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_{32}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.51±0.33 OUR FIT			

$\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$	Γ_{35}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻³	90	1 ABLIKIM 15N BES3	$\psi(2s) e^+ e^- \rightarrow \gamma\pi^0 \eta_c$	

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma\gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	Γ_{36}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.01±0.19 OUR FIT			

$\Gamma(\eta\eta)/\Gamma(\pi\pi)$	Γ_{36}/Γ_{32}		
VALUE	DOCUMENT ID	TECN	COMMENT
0.353±0.025 OUR FIT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.09 ± 0.03	1 ANDREOTTI 05C E835	$\bar{p}p \rightarrow 2 \text{ mesons}$
0.24 ± 0.10 ± 0.08	1 BAI 03C BES	$\psi(2S) \rightarrow 5\gamma$

¹ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi\pi$.

NODE=M056R69
NODE=M056R69

NODE=M056R69;LINKAGE=HE

NODE=M056R4
NODE=M056R4
→ UNCHECKED ←

NODE=M056R;LINKAGE=X1

NODE=M056R10
NODE=M056R10

NODE=M056R26
NODE=M056R26

NODE=M056R26;LINKAGE=AI
NODE=M056R26;LINKAGE=AB

NODE=M056R22
NODE=M056R22

NODE=M056R00
NODE=M056R00

NODE=M056R00;LINKAGE=A

NODE=M056R13
NODE=M056R13

NODE=M056R20
NODE=M056R20

NODE=M056R;LINKAGE=D1

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.1±0.2		85	1 ABLIKIM	17AI BES3	$\psi(2S) \rightarrow \gamma\eta'\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<24	90	35 ± 13	2 ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta'\eta$
<50	90		3 ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ ABLIKIM 17AI reports $(8.92 \pm 0.84 \pm 0.65) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $< 0.25 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

³ Superseded by ASNER 09. ADAMS 07 reports $< 0.5 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

NODE=M056R03
NODE=M056R03

NODE=M056R03;LINKAGE=A

NODE=M056R03;LINKAGE=AS

NODE=M056R03;LINKAGE=AD

NODE=M056R04
NODE=M056R04

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.17±0.12 OUR AVERAGE				
2.23±0.13±0.05	2.5k	1 ABLIKIM	17AI BES3	$\psi(2S) \rightarrow \gamma\eta'\eta'$
2.00±0.21±0.04	0.4k	2 ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta'\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.60±0.41±0.03	23	3 ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
1 ABLIKIM 17AI reports $(2.19 \pm 0.03 \pm 0.14) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
2 ASNER 09 reports $(2.12 \pm 0.13 \pm 0.21) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
3 Superseded by ASNER 09. ADAMS 07 reports $(1.7 \pm 0.4 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 0.0922 \pm 0.0011 \pm 0.0046$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{38}/Γ

NODE=M056R04
NODE=M056R04

NODE=M056R04;LINKAGE=A

NODE=M056R04;LINKAGE=AS

NODE=M056R04;LINKAGE=AD

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.11 OUR AVERAGE				
0.93±0.11±0.02	991	1 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$
2.16±0.66±0.04	38.1±9.6	2 ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma 6\pi$
1 ABLIKIM 11K reports $(0.95 \pm 0.03 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
2 ABLIKIM 05N reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.212 \pm 0.053 \pm 0.037) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{39}/Γ

NODE=M056R27
NODE=M056R27

NODE=M056R27;LINKAGE=AL

NODE=M056R27;LINKAGE=AB

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.41±0.13±0.03	486	1 ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.18±0.22±0.02	76	2,3 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$

 Γ_{40}/Γ

NODE=M056R76
NODE=M056R76

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (13.83 \pm 0.70 \pm 1.01) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 11K reports $(1.2 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Superseded by ABLIKIM 19J.

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{41}/Γ
$1.94 \pm 0.06 \pm 0.20$	1.4k	¹ ABLIKIM	13B	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$.

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	Γ_{42}/Γ
6.05 ± 0.31 OUR FIT		

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	Γ_{43}/Γ
3.16 ± 0.17 OUR FIT		

$\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{43}/Γ_{32}
0.371 ± 0.023 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.31 \pm 0.05 \pm 0.05$ ^{1,2} CHEN 07B BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$

¹ Using $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from the $\pi^+ \pi^-$ measurement of NAKAZAWA 05 rescaled by 3/2 to convert to $\pi\pi$.

² Not independent from other measurements.

$\Gamma(K_S^0 K_S^0)/\Gamma(K^+ K^-)$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{43}/Γ_{42}
0.52 ± 0.04 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.49 \pm 0.07 \pm 0.08$ ^{1,2} CHEN 07B BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c0}$

¹ Using $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from NAKAZAWA 05.

² Not independent from other measurements.

$\Gamma(\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{44}/Γ
<0.20	90	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.0 90 ² ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ ATHAR 07 reports $< 0.21 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

² ABLIKIM 06R reports $< 1.1 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{45}/Γ
<0.4	90	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$	

¹ ATHAR 07 reports $< 0.38 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

NODE=M056R76;LINKAGE=A

NODE=M056R76;LINKAGE=AL

NODE=M056R76;LINKAGE=B

NODE=M056R87
NODE=M056R87

NODE=M056R87;LINKAGE=A

NODE=M056R6
NODE=M056R6

NODE=M056R15
NODE=M056R15

NODE=M056R53
NODE=M056R53

NODE=M056R53;LINKAGE=CH

NODE=M056R53;LINKAGE=NI

NODE=M056R52
NODE=M056R52

NODE=M056R52;LINKAGE=CH
NODE=M056R52;LINKAGE=NI

NODE=M056R08
NODE=M056R08

NODE=M056R08;LINKAGE=AT

NODE=M056R08;LINKAGE=AB

NODE=M056R51
NODE=M056R51

NODE=M056R51;LINKAGE=AT

$\Gamma(K^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.09	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.7	90	2,3 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
<0.7	90	3,4 BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ATHAR 07 reports $< 0.10 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

² ABLIKIM 06R reports $< 0.70 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

³ We have multiplied the $K_S^0 K^+ \pi^-$ measurement by a factor of 2 to convert to $K^0 K^+ \pi^-$.

⁴ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
1 ATHAR 07 reports $< 0.06 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.				

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.23	90	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
1 ATHAR 07 reports $< 0.24 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.				

$\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.47 ± 0.03	16.8 ± 4.8	1 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
1 ABLIKIM 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.138 \pm 0.039 \pm 0.025) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.8 ± 0.5 ± 0.1	319	1 ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$
1 Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (5.64 \pm 0.33 \pm 0.37) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..				

$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID
2.82 ± 0.29 OUR FIT	

$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.25 ± 0.02	38	1 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
1 ABLIKIM 06T reports $(1.03 \pm 0.22 \pm 0.15) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

Γ_{46}/Γ

NODE=M056R17
NODE=M056R17

NODE=M056R17;LINKAGE=AT

NODE=M056R17;LINKAGE=AB

NODE=M056R17;LINKAGE=BA

NODE=M056R17;LINKAGE=X1

NODE=M056R05
NODE=M056R05

NODE=M056R05;LINKAGE=AT

NODE=M056R09
NODE=M056R09

NODE=M056R09;LINKAGE=AT

NODE=M056R48
NODE=M056R48

NODE=M056R48;LINKAGE=AB

NODE=M056R95
NODE=M056R95

NODE=M056R95;LINKAGE=A

NODE=M056R14
NODE=M056R14

NODE=M056R01
NODE=M056R01

NODE=M056R01;LINKAGE=AB

$\Gamma(K^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{53}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M056R90 NODE=M056R90
$3.68 \pm 0.30 \pm 0.50$	ABLIKIM	15M	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$	
$\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$	Γ_{54}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M056R91 NODE=M056R91
$1.90 \pm 0.14 \pm 0.32$	ABLIKIM	15M	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$	
$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$	Γ_{55}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M056R88 NODE=M056R88
$1.18 \pm 0.07 \pm 0.13$	538	1 ABLIKIM	13B	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$	
1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.					
$\Gamma(\phi \phi)/\Gamma_{\text{total}}$	Γ_{56}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>				NODE=M056R16 NODE=M056R16
$0.80 \pm 0.07 \text{ OUR FIT}$					
$\Gamma(\phi \phi \eta)/\Gamma_{\text{total}}$	Γ_{57}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M056R98 NODE=M056R98
$8.4 \pm 0.7 \pm 0.6$	186.6	1 ABLIKIM	20B	$B(\psi(2S) \rightarrow \gamma \phi \phi \eta)$	
1 ABLIKIM 20B reports $(8.41 \pm 0.74 \pm 0.62) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$.					
$\Gamma(p\bar{p})/\Gamma_{\text{total}}$	Γ_{58}/Γ				
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>				NODE=M056R11 NODE=M056R11
$2.21 \pm 0.08 \text{ OUR FIT}$					
$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$	Γ_{59}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M056R06 NODE=M056R06
$0.70 \pm 0.07 \text{ OUR AVERAGE}$	Error includes scale factor of 1.3.				
0.73 $\pm 0.06 \pm 0.01$	1 ONYISI	10	CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$	
0.56 $\pm 0.12 \pm 0.01$	2 ATHAR	07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$	
1 ONYISI 10 reports $(7.76 \pm 0.37 \pm 0.51 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 ATHAR 07 reports $(0.59 \pm 0.10 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$	Γ_{60}/Γ				
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M056R50 NODE=M056R50
$0.35 \pm 0.04 \text{ OUR AVERAGE}$					
0.35 $\pm 0.04 \pm 0.01$	1 ONYISI	10	CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$	
0.37 $\pm 0.11 \pm 0.01$	2 ATHAR	07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$	
1 ONYISI 10 reports $(3.73 \pm 0.38 \pm 0.28 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 ATHAR 07 reports $(0.39 \pm 0.11 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.52±0.06±0.01	1 ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$

¹ ONYISI 10 reports $(5.57 \pm 0.48 \pm 0.42 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{61}/Γ

NODE=M056R70
NODE=M056R70

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0±1.4±0.1	42 ± 8	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $(6.12 \pm 1.18 \pm 0.86) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{62}/Γ

NODE=M056R75
NODE=M056R75

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.7 OUR EVALUATION	Error includes scale factor of 1.4. Treating systematic error as correlated.		

2.1 ± 1.0 OUR AVERAGE Error includes scale factor of 2.0.

1.57±0.21±0.53	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c0}$
4.20±1.15±0.18	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

 Γ_{63}/Γ

NODE=M056R7
NODE=M056R7
→ UNCHECKED ←

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.104±0.028±0.002	39.5	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹ HE 08B reports $0.11 \pm 0.02 \pm 0.02 \pm 0.01\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{64}/Γ

NODE=M056R65
NODE=M056R65

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.22±0.26±0.02	48 ± 8	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $(1.24 \pm 0.20 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{65}/Γ

NODE=M056R72
NODE=M056R72

 $\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	¹ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

¹ Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

 Γ_{66}/Γ

NODE=M056R46
NODE=M056R46

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.7±1.1 OUR AVERAGE				

12.9±1.1±0.3	5150	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
11.2±3.1±0.2		² ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^-X$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.26 \pm 0.02 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R;LINKAGE=AB

² ABLIKIM 06I reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.10 \pm 0.24 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{67}/Γ

NODE=M056R49
NODE=M056R49

NODE=M056R49;LINKAGE=AL

NODE=M056R49;LINKAGE=AB

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{68}/Γ
$13.7 \pm 1.2 \pm 0.3$	5808	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.34 \pm 0.03 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(p\bar{n}\pi^- \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{69}/Γ
$23.4 \pm 2.0 \pm 0.5$	2480	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^- \pi^0$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (2.29 \pm 0.08 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\bar{p}n\pi^+ \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{70}/Γ
$22.1 \pm 1.8 \pm 0.5$	2757	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+ \pi^0$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+ \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (2.16 \pm 0.07 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	Γ_{71}/Γ
3.27 ± 0.24 OUR FIT		

 $\Gamma(\Lambda\bar{\Lambda}\pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{72}/Γ
$118 \pm 12 \pm 2$	426	¹ ABLIKIM	12I	BES3	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}\pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<400 90 ² ABLIKIM 06D BES2 $\psi(2S) \rightarrow \chi_{c0}\gamma$						
1 ABLIKIM 12I reports $(119.0 \pm 6.4 \pm 11.4) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						
2 Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$						

 $\Gamma(\Lambda\bar{\Lambda}\pi^+ \pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{73}/Γ
<50	90	¹ ABLIKIM	12I	BES3	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}\pi^+ \pi^-$
1 ABLIKIM 12I reports $< 54 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+ \pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{74}/Γ
<50	90	¹ ABLIKIM	12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+\bar{\Lambda}\pi^-$
1 ABLIKIM 12I reports $< 55 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{75}/Γ
<50	90	¹ ABLIKIM	12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^-\bar{\Lambda}\pi^+$
1 ABLIKIM 12I reports $< 50 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					

NODE=M056R82

NODE=M056R82

NODE=M056R83

NODE=M056R83

NODE=M056R84

NODE=M056R84

NODE=M056R44

NODE=M056R44

NODE=M056R77

NODE=M056R77

NODE=M056R78

NODE=M056R78

NODE=M056R79

NODE=M056R79

$\Gamma(K^+\bar{\Lambda}+c.c.)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{76}/Γ
1.25±0.12 OUR AVERAGE Error includes scale factor of 1.3.					
1.30±0.09±0.03	9k	1,2 ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} K^+$	
1.01±0.19±0.02		3 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$	
1 ABLIKIM 13D reports $(1.32 \pm 0.03 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{\Lambda} + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \approx (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 Using $B(\Lambda \rightarrow p \pi^-) = 63.9\%$.					
3 ATHAR 07 reports $(1.07 \pm 0.17 \pm 0.12) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{\Lambda} + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \approx (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(K^*(892)^+\bar{\Lambda}+c.c.)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{77}/Γ
4.8±0.9±0.1					
254	1 ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^* \bar{\Lambda}$		
1 ABLIKIM 19AU reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^+ \bar{\Lambda} + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.7 \pm 0.7 \pm 0.5) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(K^+\bar{\Lambda}(1520)+c.c.)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{78}/Γ
2.9±0.7±0.1					
62 ± 12	1 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$		
1 ABLIKIM 11F reports $(3.00 \pm 0.58 \pm 0.50) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{\Lambda}(1520) + c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \approx (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{79}/Γ
3.1±1.2±0.1					
28 ± 10	1 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$		
1 ABLIKIM 11F reports $(3.18 \pm 1.11 \pm 0.53) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \approx (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{80}/Γ
4.68±0.32 OUR AVERAGE					
4.82±0.34±0.10	1046	1 ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$	
4.2 ± 0.7 ± 0.1	78 ± 10	2 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4.7 ± 0.5 ± 0.1	243	3,4 ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$	
1 ABLIKIM 18V reports $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.72 \pm 0.18 \pm 0.28) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 NAIK 08 reports $(4.41 \pm 0.56 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \approx (9.62 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
3 ABLIKIM 13H reports $(4.78 \pm 0.34 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \approx (9.62 \pm 0.31 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

4 Superseded by ABLIKIM 18V

NODE=M056R07
NODE=M056R07NODE=M056R07;LINKAGE=AB
NODE=M056R07;LINKAGE=ATNODE=M056R99
NODE=M056R99

NODE=M056R99;LINKAGE=A

NODE=M056R73
NODE=M056R73

NODE=M056R73;LINKAGE=AB

NODE=M056R74
NODE=M056R74

NODE=M056R74;LINKAGE=AB

NODE=M056R59
NODE=M056R59

NODE=M056R59;LINKAGE=NA

NODE=M056R59;LINKAGE=AB

NODE=M056R59;LINKAGE=B

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{82}/Γ	
4.6 ± 0.8 OUR AVERAGE				Error includes scale factor of 2.6.		
5.10 ± 0.35 ± 0.10	747	1 ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$		
3.1 ± 0.7 ± 0.1	39 ± 7	2 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
4.5 ± 0.5 ± 0.1	148	3.4 ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$		
1 ABLIKIM 18V reports $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.99 \pm 0.24 \pm 0.24) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R60;LINKAGE=A	
2 NAIK 08 reports $(3.25 \pm 0.57 \pm 0.43) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R60;LINKAGE=NA	
3 ABLIKIM 13H reports $(4.54 \pm 0.42 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R60;LINKAGE=AB	
4 Superseded by ABLIKIM 18V						NODE=M056R60;LINKAGE=B

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{83}/Γ
16.2 ± 5.8 ± 0.3	27	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$	
1 ABLIKIM 12I reports $(16.4 \pm 5.7 \pm 1.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R80

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{84}/Γ
23.2 ± 6.5 ± 0.5	33	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$	
1 ABLIKIM 12I reports $(23.5 \pm 6.2 \pm 2.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R81

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{85}/Γ
1.94 ± 0.35 ± 0.04	57	1 ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	
1 ABLIKIM 15I reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (1.90 \pm 0.30 \pm 0.16) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R92

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{86}/Γ
3.1 ± 0.8 ± 0.1	23.3 ± 4.9	1 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$	
1 NAIK 08 reports $(3.34 \pm 0.70 \pm 0.48) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M056R61

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{87}/Γ
4.8 ± 0.7 ± 0.1	95 ± 11	1 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<10.3	90	2 ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0} \gamma$		

NODE=M056R60
NODE=M056R60

NODE=M056R60;LINKAGE=A

NODE=M056R60;LINKAGE=NA

NODE=M056R60;LINKAGE=AB

NODE=M056R60;LINKAGE=B

NODE=M056R80
NODE=M056R80

NODE=M056R80;LINKAGE=AL

NODE=M056R81
NODE=M056R81

NODE=M056R81;LINKAGE=AL

NODE=M056R92
NODE=M056R92

NODE=M056R92;LINKAGE=A

NODE=M056R61
NODE=M056R61

NODE=M056R61;LINKAGE=NA

NODE=M056R45
NODE=M056R45

¹ NAIK 08 reports $(5.14 \pm 0.60 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{88}/Γ
$< 7 \times 10^{-4}$	90	1,2 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$< 41 \times 10^{-4}$	90	1,3 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	
1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$.					
2 From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.					
3 From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.					

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{58}/\Gamma \times \Gamma_{32}/\Gamma$
18.8 ± 1.0 OUR FIT				
$15.3 \pm 2.4 \pm 0.8$	¹ ANDREOTTI 03	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$	

¹ We have multiplied $B(p\bar{p}) \cdot B(\pi^0\pi^0)$ measurement by 3 to obtain $B(p\bar{p}) \cdot B(\pi\pi)$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{58}/\Gamma \times \Gamma_{33}/\Gamma$
<0.4	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0\eta$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{58}/\Gamma \times \Gamma_{34}/\Gamma$
<2.5	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0\eta'$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{58}/\Gamma \times \Gamma_{36}/\Gamma$
6.7 ± 0.5 OUR FIT				
$4.0 \pm 1.2 \pm 0.5$	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \eta\eta$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT	$\Gamma_{58}/\Gamma \times \Gamma_{37}/\Gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$2.1^{+2.3}_{-1.5}$	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0\eta'$	

RADIATIVE DECAYS

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{89}/Γ
1.40 ± 0.05 OUR FIT					

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

0.25 $\pm 0.16 \pm 2.15$	12k	¹ ABLIKIM	17U BES3	$e^+e^- \rightarrow \gamma X$	
2.0 $\pm 0.2 \pm 0.2$		² ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	

¹ Not independent from $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))$ and the product $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))$ also measured in ABLIKIM 17U.

² Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c0})$ from ATHAR 04.

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{90}/Γ
< 9	90	1.2 ± 4.5	¹ BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$	

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

< 10	90	6 ± 12	² ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$	
--------	----	------------	----------------------	----------	---	--

¹ BENNETT 08A reports $< 9.6 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}]}$ $\times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

² ABLIKIM 11E reports $< 10.5 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}]}$ $\times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.

NODE=M056R45;LINKAGE=NA

NODE=M056R45;LINKAGE=AB

NODE=M056R89
NODE=M056R89

OCCUR=2

NODE=M056R89;LINKAGE=A
NODE=M056R89;LINKAGE=B
NODE=M056R89;LINKAGE=C

NODE=M056R21
NODE=M056R21

NODE=M056R;LINKAGE=AD

NODE=M056R41
NODE=M056R41

NODE=M056R42
NODE=M056R42

NODE=M056R40
NODE=M056R40

NODE=M056R43
NODE=M056R43

NODE=M056R8
NODE=M056R8

NODE=M056R8;LINKAGE=A

NODE=M056R8;LINKAGE=AD

NODE=M056R56
NODE=M056R56

NODE=M056R56;LINKAGE=BE

NODE=M056R56;LINKAGE=AB

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$					Γ_{91}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 8	90	0.0 ± 2.8	1 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13	90	5 ± 11	2 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
1 BENNETT 08A reports $< 8.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					
2 ABLIKIM 11E reports $< 12.9 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$					Γ_{92}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 6	90	0.1 ± 1.6	1 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	15 ± 7	2 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
1 BENNETT 08A reports $< 6.4 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					
2 ABLIKIM 11E reports $< 16.2 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$.					

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{93}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.04 ± 0.09 OUR FIT					

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	1 WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
1 WICHT 08 reports $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^\pm \rightarrow \chi_{c0} K^\pm)] < 0.11 \times 10^{-6}$ which we divide by our best value $B(B^\pm \rightarrow \chi_{c0} K^\pm) = 1.50 \times 10^{-4}$.				

$\Gamma(e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}$					Γ_{94}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

1.54 ± 0.33 ± 0.03	56	1,2 ABLIKIM	17I BES3	$\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$
1 ABLIKIM 17I reports $(1.51 \pm 0.30 \pm 0.13) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

2 Not independent from other measurements reported by ABLIKIM 17I

$\Gamma(e^+ e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$					Γ_{94}/Γ_{89}
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
9.5 ± 1.9 ± 0.7					

1 Uses $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(\mu^+ \mu^- J/\psi(1S))/\Gamma(e^+ e^- J/\psi(1S))$					Γ_{95}/Γ_{94}
<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.14					

$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$					Γ_{93}/Γ_{89}
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.45 ± 0.08 OUR FIT					

2.0 ± 0.4 OUR AVERAGE

2.2 ± 0.4	+0.1 -0.2		1 ANDREOTTI	04 E835	$p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$
2 AMBROGIANI 00B E835 $\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$					

1 The values of $B(p\bar{p})B(\gamma\gamma)$ and $B(\gamma\gamma)B(\gamma J/\psi)$ measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

2 Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

NODE=M056R57
NODE=M056R57

NODE=M056R58
NODE=M056R58

NODE=M056R1
NODE=M056R1

NODE=M056R93
NODE=M056R93

NODE=M056R94
NODE=M056R94

NODE=M056R97
NODE=M056R97

NODE=M056R18
NODE=M056R18

NODE=M056R;LINKAGE=AN

NODE=M056R;LINKAGE=7A

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{89}/\Gamma$
VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
31.1±1.5 OUR FIT				
28.2±2.1 OUR AVERAGE				
28.0±1.9±1.3	392	1,2,3 BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
29.3 $^{+5.7}_{-4.7}$ ±1.5	89	1,2 AMBROGIANI	99B	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$

¹ Values in ($\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$) and ($\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$) are not independent. The latter is used in the fit since it is less correlated to the total width.

² Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

³ Recalculated by ANDREOTTI 05A.

NODE=M056R19
NODE=M056R19

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{93}/\Gamma$
VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT

4.52±0.27 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.52±1.18 $^{+0.48}_{-0.72}$		¹ ANDREOTTI	04 E835	$p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$
------------------------------	--	------------------------	---------	---

¹ The values of $B(p\bar{p})B(\gamma\gamma)$ and $B(\gamma\gamma)B(\gamma J/\psi)$ measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

NODE=M056R;LINKAGE=KS

NODE=M056R19;LINKAGE=7A
NODE=M056R19;LINKAGE=AN

NODE=M056R25
NODE=M056R25

$\chi_{c0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma^{\psi(2S)}$
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT

21.7±0.9 OUR FIT

23.7±1.0 OUR AVERAGE

23.7±0.8±0.9	1222	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
23.7±1.4±1.4	383±22	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
23.6 $^{+3.7}_{-3.4}$ ±3.4	89.5 $^{+14}_{-13}$	BAI	04F BES	$\psi(2S) \rightarrow \gamma\chi_{c0}(1P) \rightarrow \gamma p\bar{p}$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow p\bar{p}) = (25.7 \pm 1.5 \pm 1.5 \pm 1.3) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056R25;LINKAGE=AN

NODE=M056230

NODE=M056B6
NODE=M056B6

$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$				$\Gamma_{58}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT

6.25±0.26 OUR FIT

4.6 ±1.9

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B6;LINKAGE=NA

NODE=M056B1
NODE=M056B1

NODE=M056B;LINKAGE=B1

$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$				$\Gamma_{71}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT

32.0±2.3 OUR FIT

31.7±2.3 OUR AVERAGE

32.0±1.9±2.2	369	¹ ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
31.2±3.3±2.0	131±12	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

NODE=M056B20
NODE=M056B20

¹ Calculated by us. ABLIKIM 13H reports $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.3 \pm 2.0 \pm 2.6) \times 10^{-5}$ from a measurement of $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c0})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.62 \pm 0.31)\%$.

NODE=M056B20;LINKAGE=AB

² Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.8 \pm 3.6 \pm 2.2 \pm 1.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B20;LINKAGE=NA

$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$				$\Gamma_{71}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT

9.2±0.7 OUR FIT

13.0 $^{+3.6}_{-3.5}$ ±2.5

¹ BAI

NODE=M056B21
NODE=M056B21

¹ BAI 03E reports [$B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c0}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$] \times [$B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})$] = $(2.45^{+0.68}_{-0.65} \pm 0.46)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

NODE=M056B21;LINKAGE=BA

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{89}/\Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.138±0.005 OUR FIT**0.147±0.029 OUR AVERAGE**

0.158±0.003±0.006	4.8k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$
0.024±0.015±0.205	12k	ABLIKIM	17U BES3	$e^+e^- \rightarrow \gamma X$
0.069±0.018		² OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c0}$
0.4 ± 0.3		³ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma\chi_{c0}$
0.16 ± 0.11		³ BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma\chi_{c0}$
3.3 ± 1.7		⁴ BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.151±0.003±0.010	4.3k	⁵ ABLIKIM	120 BES3	$\psi(2S) \rightarrow \gamma\chi_{c0}$
0.125±0.007±0.013	560	⁶ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
0.18 ± 0.01 ± 0.02	172	⁷ ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

NODE=M056B2
NODE=M056B2

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{anything})$$

$$\Gamma_{89}/\Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma_9^{\psi(2S)} = \Gamma_{89}/\Gamma \times \Gamma_{153}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + 0.343\Gamma_{154}^{\psi(2S)} + 0.190\Gamma_{155}^{\psi(2S)})$$

$$\Gamma_{89}/\Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma_9^{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.224±0.009 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.201±0.011±0.021	560	¹ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
0.31 ± 0.02 ± 0.03	172	ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M056B2;LINKAGE=A

NODE=M056B;LINKAGE=3Q

NODE=M056B;LINKAGE=2Q

NODE=M056B;LINKAGE=EA

NODE=M056B2;LINKAGE=B

NODE=M056B2;LINKAGE=ME

NODE=M056B2;LINKAGE=AD

NODE=M056B7

NODE=M056B7

NODE=M056B7

NODE=M056B7;LINKAGE=ME

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{89}/\Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.397±0.015 OUR FIT

0.358±0.020±0.037	560	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
--------------------------	-----	--------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.55 ± 0.04 ± 0.06	172	¹ ADAM	05A CLEO	Repl. by MENDEZ 08
--------------------	-----	-------------------	----------	--------------------

¹ Not independent from other values reported by ADAM 05A.

NODE=M056B8

NODE=M056B8

NODE=M056B;LINKAGE=AD

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{93}/\Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.00±0.08 OUR FIT**1.95±0.09 OUR AVERAGE**

1.93±0.08±0.05	3.5k	ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
2.17±0.32±0.10	0.2k	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
3.7 ± 1.8 ± 1.0		LEE	85 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.17±0.17±0.12	0.8k	¹ ABLIKIM	12A BES3	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
----------------	------	----------------------	----------	--

¹ Superseded by ABLIKIM 17AE.

NODE=M056B3

NODE=M056B3

NODE=M056B3;LINKAGE=A

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{32}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.34±0.29 OUR FIT**8.80±0.34 OUR AVERAGE**

9.11±0.08±0.65	17k	¹ ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
8.81±0.11±0.43	8.9k	² ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
8.13±0.19±0.89	2.8k	³ ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (3.23 \pm 0.03 \pm 0.23 \pm 0.14) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^+\pi^-) = (6.37 \pm 0.08 \pm 0.31 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

³ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (2.94 \pm 0.07 \pm 0.32 \pm 0.15) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{32}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

24.0±0.8 OUR FIT**20.7±1.7 OUR AVERAGE**

23.9±2.7±4.1	97 ± 11	¹ BAI	03C BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^0\pi^0$
20.2±1.1±1.5	720 ± 32	² BAI	98I BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^+\pi^-$

¹ We have multiplied $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. The value for $B(\chi_{c0} \rightarrow \pi^+\pi^-)$ reported in BAI 98I is derived using $B(\psi' \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi' \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{36}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.95±0.18 OUR FIT**3.12±0.19 OUR AVERAGE**

3.23±0.09±0.23	2132	¹ ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
2.93±0.12±0.29	0.9k	² ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.86±0.46±0.37	48	³ ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
----------------	----	--------------------	---------	--

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \eta\eta) = (3.44 \pm 0.10 \pm 0.24 \pm 0.13) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$.

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \eta\eta) = (3.18 \pm 0.13 \pm 0.31 \pm 0.16) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

³ Superseded by ASNER 09. Calculated by us. The value of $B(\chi_{c0}(1P) \rightarrow \eta\eta)$ reported by ADAMS 07 was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46)\%$ (ATHAR 04).

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{36}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.85 ± 0.05 OUR FIT**0.578±0.241±0.158**

BAI	03C BES	$\psi(2S) \rightarrow \gamma\eta\eta$
-----	---------	---------------------------------------

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{42}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.92±0.28 OUR FIT**5.97±0.07±0.32**

8.1k	¹ ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma K^+K^-$
------	--------------------	---------	--------------------------------------

¹ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow K^+K^-) = (6.47 \pm 0.08 \pm 0.35 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B22

NODE=M056B22

OCCUR=2

NODE=M056B22;LINKAGE=AB

NODE=M056B22;LINKAGE=AS

NODE=M056B22;LINKAGE=AN

NODE=M056B5

NODE=M056B5

NODE=M056B;LINKAGE=D1

NODE=M056B;LINKAGE=D2

NODE=M056B11

NODE=M056B11

NODE=M056B11;LINKAGE=AB

NODE=M056B11;LINKAGE=AS

NODE=M056B11;LINKAGE=AD

NODE=M056B10

NODE=M056B10

NODE=M056B23

NODE=M056B23

NODE=M056B23;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{42}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

1.71±0.08 OUR FIT

1.63±0.10±0.15 774 ± 38 1 BAI $98I$ BES $\psi(2S) \rightarrow \gamma K^+ K^-$

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{43}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

3.10±0.16 OUR FIT

3.18±0.17 OUR AVERAGE

$3.22 \pm 0.07 \pm 0.17$	$2.1k$	1	ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
$3.02 \pm 0.19 \pm 0.33$	322	1	ABLIKIM	050	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow K_S^0 K_S^0) = (3.49 \pm 0.08 \pm 0.18 \pm 0.17) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{43}/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

8.9±0.5 OUR FIT

5.6±0.8±1.3 1 BAI $99B$ BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_1/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

6.6±0.5 OUR FIT

6.9±2.4 OUR AVERAGE Error includes scale factor of 3.8.

$4.4 \pm 0.1 \pm 0.9$	1	BAI	$99B$	BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
9.3 ± 0.9	2	TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

² The value $B(\psi(1S) \rightarrow \gamma \chi_{c0}) \times B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_8/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

1.78±0.14 OUR FIT

1.64±0.05±0.2 1 BAI $05Q$ BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_8/\Gamma \times \Gamma_{153}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

5.1 ±0.4 OUR FIT

5.8 ±1.6 OUR AVERAGE Error includes scale factor of 2.3.

$4.22 \pm 0.20 \pm 0.97$	1	BAI	$99B$	BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
7.4 ± 1.0	1	TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+\pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

NODE=M056B9

NODE=M056B9

NODE=M056B9;LINKAGE=BA

NODE=M056B12

NODE=M056B12

NODE=M056B12;LINKAGE=AS

NODE=M056B13

NODE=M056B13

NODE=M056B13;LINKAGE=BA

NODE=M056B4

NODE=M056B4

NODE=M056B;LINKAGE=B2

NODE=M056B;LINKAGE=J1

NODE=M056B18

NODE=M056B18

NODE=M056B19

NODE=M056B19

NODE=M056B19;LINKAGE=TA

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{51} / \Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.76±0.28 OUR FIT				
3.20±0.11±0.41	278	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

$$\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}$$

$$\Gamma_{51} / \Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±0.8 OUR FIT				
6.1±0.8±0.9	1 BAI	99B BES		$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{56} / \Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.78±0.07 OUR FIT				
0.78±0.08 OUR AVERAGE				

0.77±0.03±0.08	612	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
0.86±0.19±0.12	26	² ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31)\%$.

² Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

$$\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{56} / \Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.25±0.21 OUR FIT				
2.6 ±1.0 ±1.1	1 BAI	99B BES		$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{81} / \Gamma \times \Gamma_{153}^{\psi(2S)} / \Gamma^{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.45±0.17±0.19	493±26	¹ ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c. c.}$

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_c^0 \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (3.52 \pm 0.19 \pm 0.21) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_c^0) = (9.79 \pm 0.20)\%$ and other branching fractions from PDG 18.

$\chi_{c0}(1P)$ REFERENCES

ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60212
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59844
ABLIKIM	19AU	PR D100 052010	Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996;ERROR=13
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60026
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59606
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59837
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58990
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58310
ABLIKIM	17AI	PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58321
ABLIKIM	17I	PR D118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56778
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56779
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583

UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
ANDREOTTI	05C	PR D72 112002	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50991
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50187
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ANDREOTTI	04	PL B584 16	M. Andreotti <i>et al.</i>	(E835 Collab.)	REFID=49744
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
ANDREOTTI	03	PRL 91 091801	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=49578
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAGNASCO	02	PL B533 237	S. Bagnasco <i>et al.</i>	(FNAL E835 Collab.)	REFID=48833
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
AMBROGIANI	99B	PRL 83 2902	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47389
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
GAISER	86	PR D34 711	J. Gaisser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
O REGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059

$\chi_{c1}(1P)$

$I^G(J^{PC}) = 0^+(1^{++})$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

NODE=M055

$\chi_{c1}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3510.67 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.2.		
3508.4 ± 1.9 ± 0.7	460	1 AAIJ	17BB LHCb	$p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3510.71 ± 0.04 ± 0.09	4.8k	2 AAIJ	17BI LHCb	$\chi_{c1} \rightarrow J/\psi\mu^+\mu^-$
3510.30 ± 0.14 ± 0.16		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
3510.719 ± 0.051 ± 0.019		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3509.4 ± 0.9		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3510.60 ± 0.087 ± 0.019	513	3 ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3511.3 ± 0.4 ± 0.4	30	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
3512.3 ± 0.3 ± 4.0		4 GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7	91	5 LEMOIGNE	82 GOLI	$185\pi^-Be \rightarrow \gamma\mu^+\mu^-A$
3510.4 ± 0.6		OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1	254	6 HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3509 ± 11	21	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3507 ± 3		6 BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4		6,7 TANENBAUM	78 MRK1	e^+e^-
3513 ± 7	367	6 BIDDICK	77 CNTR	$\psi(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3500 ± 10 40 TANENBAUM 75 MRK1 Hadrons γ

1 From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c1}(1P)$ fixed to the PDG 16 value.

2 AAIJ 17BI reports also $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$ MeV.

3 Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

4 Using mass of $\psi(2S) = 3686.0$ MeV.

5 $J/\psi(1S)$ mass constrained to 3097 MeV.

6 Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

7 From a simultaneous fit to radiative and hadronic decay channels.

NODE=M055M

NODE=M055M

OCCUR=2

NODE=M055M;LINKAGE=A

NODE=M055M;LINKAGE=B

NODE=M055M;LINKAGE=NW

NODE=M055M;LINKAGE=C

NODE=M055M;LINKAGE=P

NODE=M055M;LINKAGE=D

NODE=M055M;LINKAGE=M

NODE=M055W

NODE=M055W

$\chi_{c1}(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 ± 0.04 OUR FIT					
0.88 ± 0.05 OUR AVERAGE					
1.39 + 0.40 + 0.26 - 0.38 - 0.77			ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
0.876 ± 0.045 ± 0.026			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
0.87 ± 0.11 ± 0.08	513	1 ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3 95 BAGLIN 86B SPEC $\bar{p}p \rightarrow e^+e^-X$
<3.8 90 GAISER 86 CBAL $\psi(2S) \rightarrow \gamma X$

1 Recalculated by ANDREOTTI 05A.

NODE=M055W;LINKAGE=AN

NODE=M055215;NODE=M055

$\chi_{c1}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
------	--------------------------------	-----------------------------------

Hadronic decays

Γ_1	$3(\pi^+\pi^-)$	$(5.8 \pm 1.4) \times 10^{-3}$	S=1.2	NODE=M055;CLUMP=A DESIG=6
Γ_2	$2(\pi^+\pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$		DESIG=5
Γ_3	$\pi^+\pi^-\pi^0\pi^0$	$(1.19 \pm 0.15)\%$		DESIG=51
Γ_4	$\rho^+\pi^-\pi^0 + \text{c.c.}$	$(1.45 \pm 0.24)\%$		DESIG=52
Γ_5	$\rho^0\pi^+\pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$		DESIG=9
Γ_6	$4\pi^0$	$(5.4 \pm 0.8) \times 10^{-4}$		DESIG=60
Γ_7	$\pi^+\pi^-K^+K^-$	$(4.5 \pm 1.0) \times 10^{-3}$		DESIG=7
Γ_8	$K^+K^-\pi^0\pi^0$	$(1.12 \pm 0.27) \times 10^{-3}$		DESIG=53
Γ_9	$K^+K^-\pi^+\pi^-\pi^0$	$(1.15 \pm 0.13)\%$		DESIG=79
Γ_{10}	$K_S^0K^\pm\pi^\mp\pi^+\pi^-$	$(7.5 \pm 0.8) \times 10^{-3}$		DESIG=84
Γ_{11}	$K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(8.6 \pm 1.4) \times 10^{-3}$		DESIG=55
Γ_{12}	$\rho^-\bar{K}^+\bar{K}^0 + \text{c.c.}$	$(5.0 \pm 1.2) \times 10^{-3}$		DESIG=56
Γ_{13}	$K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(2.3 \pm 0.6) \times 10^{-3}$		DESIG=57
Γ_{14}	$K^+K^-\eta\pi^0$	$(1.12 \pm 0.34) \times 10^{-3}$		DESIG=58
Γ_{15}	$\pi^+\pi^-K_S^0K_S^0$	$(6.9 \pm 2.9) \times 10^{-4}$		DESIG=28
Γ_{16}	$K^+K^-\eta$	$(3.2 \pm 1.0) \times 10^{-4}$		DESIG=42
Γ_{17}	$\bar{K}^0K^+\pi^- + \text{c.c.}$	$(7.0 \pm 0.6) \times 10^{-3}$		DESIG=17
Γ_{18}	$K^*(892)^0\bar{K}^0 + \text{c.c.}$	$(10 \pm 4) \times 10^{-4}$		DESIG=32
Γ_{19}	$K^*(892)^+K^- + \text{c.c.}$	$(1.4 \pm 0.6) \times 10^{-3}$		DESIG=33
Γ_{20}	$K_J^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow K_S^0K^+\pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$	CL=90%	DESIG=34
Γ_{21}	$K_J^*(1430)^+K^- + \text{c.c.} \rightarrow K_S^0K^+\pi^- + \text{c.c.}$	$< 2.1 \times 10^{-3}$	CL=90%	DESIG=35
Γ_{22}	$K^+K^-\pi^0$	$(1.81 \pm 0.24) \times 10^{-3}$		DESIG=38
Γ_{23}	$\eta\pi^+\pi^-$	$(4.62 \pm 0.23) \times 10^{-3}$		DESIG=31
Γ_{24}	$a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(3.2 \pm 0.4) \times 10^{-3}$	S=2.2	DESIG=36
Γ_{25}	$a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(1.76 \pm 0.24) \times 10^{-4}$		DESIG=93
Γ_{26}	$a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(4.6 \pm 0.7) \times 10^{-5}$		DESIG=96
Γ_{27}	$f_2(1270)\eta \rightarrow \eta\pi^+\pi^-$	$(3.5 \pm 0.6) \times 10^{-4}$		DESIG=94
Γ_{28}	$f_4(2050)\eta \rightarrow \eta\pi^+\pi^-$	$(2.5 \pm 0.9) \times 10^{-5}$		DESIG=95
Γ_{29}	$\pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 5 \times 10^{-5}$	CL=90%	DESIG=97
Γ_{30}	$\pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 1.5 \times 10^{-5}$	CL=90%	DESIG=98
Γ_{31}	$\pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 8 \times 10^{-6}$	CL=90%	DESIG=99
Γ_{32}	$f_2(1270)\eta$	$(6.7 \pm 1.1) \times 10^{-4}$		DESIG=37
Γ_{33}	$\pi^+\pi^-\eta'$	$(2.2 \pm 0.4) \times 10^{-3}$		DESIG=44
Γ_{34}	$K^+K^-\eta'(958)$	$(8.8 \pm 0.9) \times 10^{-4}$		DESIG=85
Γ_{35}	$K_0^*(1430)^+K^- + \text{c.c.}$	$(6.4^{+2.2}_{-2.8}) \times 10^{-4}$		DESIG=86
Γ_{36}	$f_0(980)\eta'(958)$	$(1.6^{+1.4}_{-0.7}) \times 10^{-4}$		DESIG=87
Γ_{37}	$f_0(1710)\eta'(958)$	$(7^{+7}_{-5}) \times 10^{-5}$		DESIG=88
Γ_{38}	$f'_2(1525)\eta'(958)$	$(9 \pm 6) \times 10^{-5}$		DESIG=89
Γ_{39}	$\pi^0f_0(980) \rightarrow \pi^0\pi^+\pi^-$	$(3.5 \pm 0.9) \times 10^{-7}$		DESIG=61
Γ_{40}	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(3.2 \pm 2.1) \times 10^{-3}$		DESIG=10
Γ_{41}	$K^*(892)^0\bar{K}^*(892)^0$	$(1.4 \pm 0.4) \times 10^{-3}$		DESIG=21
Γ_{42}	$K^+K^-K_S^0K_S^0$	$< 4 \times 10^{-4}$	CL=90%	DESIG=29
Γ_{43}	$K_S^0K_S^0K_S^0K_S^0$	$(3.5 \pm 1.0) \times 10^{-5}$		DESIG=102
Γ_{44}	$K^+K^-K^+K^-$	$(5.4 \pm 1.1) \times 10^{-4}$		DESIG=14
Γ_{45}	$K^+K^-\phi$	$(4.1 \pm 1.5) \times 10^{-4}$		DESIG=30
Γ_{46}	$\bar{K}^0K^+\pi^-\phi + \text{c.c.}$	$(3.3 \pm 0.5) \times 10^{-3}$		DESIG=90
Γ_{47}	$K^+K^-\pi^0\phi$	$(1.62 \pm 0.30) \times 10^{-3}$		DESIG=91
Γ_{48}	$\phi\pi^+\pi^-\pi^0$	$(7.5 \pm 1.0) \times 10^{-4}$		DESIG=82
Γ_{49}	$\omega\omega$	$(5.7 \pm 0.7) \times 10^{-4}$		DESIG=66
Γ_{50}	ωK^+K^-	$(7.8 \pm 0.9) \times 10^{-4}$		DESIG=81
Γ_{51}	$\omega\phi$	$(2.7 \pm 0.4) \times 10^{-5}$		DESIG=67
Γ_{52}	$\phi\phi$	$(4.2 \pm 0.5) \times 10^{-4}$		DESIG=68

Γ_{53}	$\phi\phi\eta$	$(3.0 \pm 0.5) \times 10^{-4}$	DESIG=104
Γ_{54}	$p\bar{p}$	$(7.60 \pm 0.34) \times 10^{-5}$	DESIG=11
Γ_{55}	$p\bar{p}\pi^0$	$(1.55 \pm 0.18) \times 10^{-4}$	DESIG=39
Γ_{56}	$p\bar{p}\eta$	$(1.45 \pm 0.25) \times 10^{-4}$	DESIG=43
Γ_{57}	$p\bar{p}\omega$	$(2.12 \pm 0.31) \times 10^{-4}$	DESIG=59
Γ_{58}	$p\bar{p}\phi$	$< 1.7 \times 10^{-5}$	CL=90% DESIG=65
Γ_{59}	$p\bar{p}\pi^+\pi^-$	$(5.0 \pm 1.9) \times 10^{-4}$	DESIG=8
Γ_{60}	$p\bar{p}\pi^0\pi^0$	$< 5 \times 10^{-4}$	CL=90% DESIG=54
Γ_{61}	$p\bar{p}K^+K^-$ (non-resonant)	$(1.27 \pm 0.22) \times 10^{-4}$	DESIG=62
Γ_{62}	$p\bar{p}K_S^0K_S^0$	$< 4.5 \times 10^{-4}$	CL=90% DESIG=25
Γ_{63}	$p\bar{n}\pi^-$	$(3.8 \pm 0.5) \times 10^{-4}$	DESIG=74
Γ_{64}	$\bar{p}n\pi^+$	$(3.9 \pm 0.5) \times 10^{-4}$	DESIG=75
Γ_{65}	$p\bar{n}\pi^-\pi^0$	$(1.03 \pm 0.12) \times 10^{-3}$	DESIG=76
Γ_{66}	$\bar{p}n\pi^+\pi^0$	$(1.01 \pm 0.12) \times 10^{-3}$	DESIG=77
Γ_{67}	$\Lambda\bar{\Lambda}$	$(1.14 \pm 0.11) \times 10^{-4}$	DESIG=19
Γ_{68}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(2.9 \pm 0.5) \times 10^{-4}$	DESIG=24
Γ_{69}	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$(2.5 \pm 0.6) \times 10^{-4}$	DESIG=69
Γ_{70}	$\Sigma(1385)^+\Lambda\pi^- + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90% DESIG=70
Γ_{71}	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 1.3 \times 10^{-4}$	CL=90% DESIG=71
Γ_{72}	$K^+\bar{p}\Lambda + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-4}$	S=1.2 DESIG=40
Γ_{73}	$K^*(892)^+\bar{p}\Lambda + \text{c.c.}$	$(4.9 \pm 0.7) \times 10^{-4}$	DESIG=106
Γ_{74}	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	$(1.7 \pm 0.4) \times 10^{-4}$	DESIG=63
Γ_{75}	$\Lambda(1520)\bar{\Lambda}(1520)$	$< 9 \times 10^{-5}$	CL=90% DESIG=64
Γ_{76}	$\Sigma^0\bar{\Sigma}^0$	$(4.2 \pm 0.6) \times 10^{-5}$	DESIG=48
Γ_{77}	$\Sigma^+\bar{p}K_S^0 + \text{c.c.}$	$(1.53 \pm 0.12) \times 10^{-4}$	DESIG=105
Γ_{78}	$\Sigma^+\bar{\Sigma}^-$	$(3.6 \pm 0.7) \times 10^{-5}$	DESIG=49
Γ_{79}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$< 9 \times 10^{-5}$	CL=90% DESIG=72
Γ_{80}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$< 5 \times 10^{-5}$	CL=90% DESIG=73
Γ_{81}	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.35 \pm 0.24) \times 10^{-4}$	DESIG=92
Γ_{82}	$\Xi^0\bar{\Xi}^0$	$< 6 \times 10^{-5}$	CL=90% DESIG=50
Γ_{83}	$\Xi^-\bar{\Xi}^+$	$(8.0 \pm 2.1) \times 10^{-5}$	DESIG=26
Γ_{84}	$\pi^+\pi^- + K^+K^-$	$< 2.1 \times 10^{-3}$	DESIG=23
Γ_{85}	$K_S^0K_S^0$	$< 6 \times 10^{-5}$	CL=90% DESIG=27
Γ_{86}	$\eta_c\pi^+\pi^-$	$< 3.2 \times 10^{-3}$	CL=90% DESIG=83

Radiative decays

Γ_{87}	$\gamma J/\psi(1S)$	$(34.3 \pm 1.0) \%$	NODE=M055;CLUMP=B
Γ_{88}	$\gamma\rho^0$	$(2.16 \pm 0.17) \times 10^{-4}$	DESIG=1
Γ_{89}	$\gamma\omega$	$(6.8 \pm 0.8) \times 10^{-5}$	DESIG=45
Γ_{90}	$\gamma\phi$	$(2.4 \pm 0.5) \times 10^{-5}$	DESIG=46
Γ_{91}	$\gamma\gamma$	$< 6.3 \times 10^{-6}$	DESIG=47
Γ_{92}	$e^+e^- J/\psi(1S)$	$(3.46 \pm 0.22) \times 10^{-3}$	DESIG=4
Γ_{93}	$\mu^+\mu^- J/\psi(1S)$	$(2.33 \pm 0.29) \times 10^{-4}$	DESIG=100
			DESIG=103

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 378.1$ for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{44}	3				
x_{54}	4	2			
x_{67}	7	3	4		
x_{87}	23	9	2	20	
Γ	-12	-5	-63	-10	-41
	x_{17}	x_{44}	x_{54}	x_{67}	x_{87}

$\chi_{c1}(1P)$ PARTIAL WIDTHS **$\chi_{c1}(1P) \Gamma(i)\Gamma(\gamma J/\psi(1S))/\Gamma(\text{total})$**

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$	$\Gamma_{54}\Gamma_{87}/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT

 21.9 ± 0.8 OUR FIT **21.4 ± 0.9 OUR AVERAGE**21.5 \pm 0.5 \pm 0.821.4 \pm 1.5 \pm 2.219.9 $^{+4.4}_{-4.0}$ 1 ANDREOTTI 05A E835 $p\bar{p} \rightarrow e^+ e^- \gamma$ 1,2 ARMSTRONG 92 E760 $\bar{p}p \rightarrow e^+ e^- \gamma$ 1 BAGLIN 86B SPEC $\bar{p}p \rightarrow e^+ e^- X$ 1 Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

2 Recalculated by ANDREOTTI 05A.

NODE=M055220

NODE=M055223

NODE=M055G1

NODE=M055G1

NODE=M055G;LINKAGE=7A

NODE=M055G;LINKAGE=AN

NODE=M055225

NODE=M055305

NODE=M055R6

NODE=M055R6

→ UNCHECKED ←

 $\chi_{c1}(1P)$ BRANCHING RATIOS**HADRONIC DECAYS**

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT

 5.8 ± 1.4 OUR EVALUATION Error includes scale factor of 1.2. Treating systematic error as correlated. **5.8 ± 1.1 OUR AVERAGE**5.4 \pm 0.7 \pm 0.916.0 \pm 5.9 \pm 0.81 BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c1}$ 1 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$ 1 Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055225

NODE=M055305

NODE=M055R6

NODE=M055R6

→ UNCHECKED ←

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT

 7.6 ± 2.6 OUR EVALUATION Treating systematic error as correlated. **8 ± 4 OUR AVERAGE** Error includes scale factor of 1.5.4.6 \pm 2.1 \pm 2.612.5 \pm 4.2 \pm 0.61 BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c1}$ 1 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$ 1 Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R;LINKAGE=X2

NODE=M055R4

NODE=M055R4

→ UNCHECKED ←

NODE=M055R4;LINKAGE=X2

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT

 $1.19 \pm 0.15 \pm 0.03$ 604.7 1 HE 08B CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ 1 HE 08B reports $1.28 \pm 0.06 \pm 0.15 \pm 0.08\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R35

NODE=M055R35

NODE=M055R35;LINKAGE=HE

$\Gamma(\rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}$	Γ_4/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT

 $1.45 \pm 0.24 \pm 0.04$ 712.3 1,2 HE 08B CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$ 1 HE 08B reports $1.56 \pm 0.13 \pm 0.22 \pm 0.10\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R36

NODE=M055R36

NODE=M055R36;LINKAGE=HE

2 Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M055R36

NODE=M055R36

NODE=M055R36;LINKAGE=OC

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_5/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT

 3.9 ± 3.5 1 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$ 1 Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R8

NODE=M055R8

NODE=M055R;LINKAGE=T

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.4±0.8±0.1	608	1 ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$	

1 ABLIKIM 11A reports $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT		
4.5±1.0 OUR EVALUATION	Treating systematic error as correlated.				

4.5±0.9 OUR AVERAGE

4.2±0.4±0.9	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c1}$		
7.3±3.0±0.4	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$		
1 Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.					

$\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.12±0.27±0.03	45.1	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

1 HE 08B reports $(0.12 \pm 0.02 \pm 0.02 \pm 0.01) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
11.46±0.12±1.29	12k	1 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$	

1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

$\Gamma(K_s^0K^\pm\pi^\mp\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
7.52±0.11±0.79	5.1k	1 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$	

1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

$\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.86±0.13±0.02	141.3	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

1 HE 08B reports $0.92 \pm 0.09 \pm 0.11 \pm 0.06\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^-\bar{K}^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.50±0.12±0.01	141.3	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

1 HE 08B reports $0.54 \pm 0.11 \pm 0.07 \pm 0.03\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^-\bar{K}^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{13}/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.23±0.06±0.01	141.3	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

1 HE 08B reports $0.25 \pm 0.06 \pm 0.03 \pm 0.02\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R44
NODE=M055R44

NODE=M055R44;LINKAGE=AB

NODE=M055R5
NODE=M055R5
→ UNCHECKED ←

NODE=M055R5;LINKAGE=X2

NODE=M055R37
NODE=M055R37

NODE=M055R37;LINKAGE=HE

NODE=M055R00
NODE=M055R00

NODE=M055R00;LINKAGE=A

NODE=M055R60
NODE=M055R60

NODE=M055R60;LINKAGE=A

NODE=M055R39
NODE=M055R39

NODE=M055R39;LINKAGE=HE

NODE=M055R40
NODE=M055R40

NODE=M055R40;LINKAGE=HE

NODE=M055R41
NODE=M055R41

NODE=M055R41;LINKAGE=HE

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$	Γ_{14}/Γ
<u>VALUE (%)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.112±0.034±0.003	141.3 1 HE 08B CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.12 \pm 0.03 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{15}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
6.9±2.9±0.2	19.8 ± 7.7 1 ABLIKIM 050 BES2 $\psi(2S) \rightarrow \chi_{c1} \gamma$

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$	Γ_{16}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
3.2±1.0±0.1	1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{17}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
7.0±0.6 OUR FIT	

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{18}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
0.98±0.37±0.02	22 1 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 06R reports $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{19}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
1.43±0.65±0.03	27 1 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 06R reports $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{20}/Γ
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<8 × 10⁻⁴	90 1 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 06R reports $< 0.9 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{21}/Γ
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<2.1 × 10⁻³	90 1 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 06R reports $< 2.4 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R42
NODE=M055R42

NODE=M055R42;LINKAGE=HE

NODE=M055R05
NODE=M055R05

NODE=M055R05;LINKAGE=AB

NODE=M055R25
NODE=M055R25

NODE=M055R25;LINKAGE=AT

NODE=M055R09
NODE=M055R09

NODE=M055R09;LINKAGE=AB

NODE=M055R10
NODE=M055R10

NODE=M055R10;LINKAGE=AB

NODE=M055R12
NODE=M055R12

NODE=M055R12;LINKAGE=AB

NODE=M055R13
NODE=M055R13

NODE=M055R13;LINKAGE=AB

$\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) **$1.81 \pm 0.24 \pm 0.04$**

DOCUMENT ID

1 ATHAR

TECN

CLEO

 Γ_{22}/Γ

NODE=M055R20

NODE=M055R20

¹ ATHAR 07 reports $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) **4.62 ± 0.23 OUR AVERAGE** $4.58 \pm 0.23 \pm 0.11$ $4.7 \pm 0.5 \pm 0.1$ $5.3 \pm 0.9 \pm 0.1$

DOCUMENT ID

1,2 ABLIKIM

TECN

 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ Γ_{23}/Γ

NODE=M055R08

NODE=M055R08

¹ From an amplitude analysis using an isobar model.
² ABLIKIM 17K reports $(4.67 \pm 0.03 \pm 0.23 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
³ ATHAR 07 reports $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
⁴ ABLIKIM 06R reports $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) **3.2 ± 0.4 OUR AVERAGE**

Error includes scale factor of 2.2.

 $3.33 \pm 0.19 \pm 0.08$ $1.79 \pm 0.63 \pm 0.04$

DOCUMENT ID

1,2 ABLIKIM

TECN

 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ Γ_{24}/Γ

NODE=M055R15

NODE=M055R15

¹ From an amplitude analysis using an isobar model.
² ABLIKIM 17K reports $(3.40 \pm 0.03 \pm 0.19 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
³ ABLIKIM 06R reports $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) **$0.176 \pm 0.023 \pm 0.004$**

DOCUMENT ID

1,2 ABLIKIM

TECN

 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ Γ_{25}/Γ

NODE=M055R72

NODE=M055R72

¹ From an amplitude analysis using an isobar model.
² ABLIKIM 17K reports $(0.18 \pm 0.01 \pm 0.02 \pm 0.01) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-5}) **$4.6 \pm 0.7 \pm 0.1$**

DOCUMENT ID

1,2 ABLIKIM

TECN

 $\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$ Γ_{26}/Γ

NODE=M055R75

NODE=M055R75

NODE=M055R20;LINKAGE=AT

NODE=M055R08

NODE=M055R08

NODE=M055R08;LINKAGE=AB

NODE=M055R08;LINKAGE=AB

NODE=M055R15;LINKAGE=A

NODE=M055R15;LINKAGE=B

NODE=M055R15;LINKAGE=AB

NODE=M055R72

NODE=M055R72

NODE=M055R72;LINKAGE=A

NODE=M055R72;LINKAGE=B

NODE=M055R75

NODE=M055R75

1 From an amplitude analysis using an isobar model.

2 ABLIKIM 17K reports $(4.7 \pm 0.4 \pm 0.6 \pm 0.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f_2(1270)\eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{27}/Γ
$3.5 \pm 0.6 \pm 0.1$	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	

1 From an amplitude analysis using an isobar model.

2 ABLIKIM 17K reports $(0.36 \pm 0.01 \pm 0.06 \pm 0.01) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(f_4(2050)\eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_{28}/Γ
$2.5 \pm 0.9 \pm 0.1$	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	

1 From an amplitude analysis using an isobar model.

2 ABLIKIM 17K reports $(2.6 \pm 0.4 \pm 0.8 \pm 0.1) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_4(2050)\eta \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{29}/Γ
$< 5 \times 10^{-5}$	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	

1 From an amplitude analysis using an isobar model.

2 ABLIKIM 17K reports $< 4.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{30}/Γ
$< 1.5 \times 10^{-5}$	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	

1 From an amplitude analysis using an isobar model.

2 ABLIKIM 17K reports $< 1.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(\pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{31}/Γ
$< 8 \times 10^{-6}$	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	

1 From an amplitude analysis using an isobar model.

2 ABLIKIM 17K reports $< 8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\mathcal{B}(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{32}/Γ
$0.67 \pm 0.11 \pm 0.02$ OUR AVERAGE		1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$	
$0.63 \pm 0.11 \pm 0.02$		53	³ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R75;LINKAGE=A

NODE=M055R75;LINKAGE=B

NODE=M055R73

NODE=M055R73

OCCUR=2

NODE=M055R73;LINKAGE=A

NODE=M055R73;LINKAGE=D

NODE=M055R74

NODE=M055R74

NODE=M055R74;LINKAGE=A

NODE=M055R74;LINKAGE=B

NODE=M055R76

NODE=M055R76

NODE=M055R76;LINKAGE=A

NODE=M055R76;LINKAGE=B

NODE=M055R77

NODE=M055R77

NODE=M055R77;LINKAGE=A

NODE=M055R77;LINKAGE=B

NODE=M055R78

NODE=M055R78

NODE=M055R78;LINKAGE=A

NODE=M055R78;LINKAGE=B

NODE=M055R16

NODE=M055R16

¹ ABLIKIM 17K reports $(6.4 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From an amplitude analysis using an isobar model.

³ ABLIKIM 06R reports $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	Γ_{33}/Γ
$2.2 \pm 0.4 \pm 0.1$	1 ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$	

¹ ATHAR 07 reports $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{34}/Γ
8.75 ± 0.87	310	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$	

¹ Derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.2 \pm 0.4)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

$\Gamma(K_0^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{35}/Γ
$6.41 \pm 0.57 \pm 2.09$	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$	

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$\Gamma(f_0(980)\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ
$1.65 \pm 0.47 \pm 1.32$	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$	

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$\Gamma(f_0(1710)\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{37}/Γ
$0.71 \pm 0.22 \pm 0.68$	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$	

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$\Gamma(f'_2(1525)\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{38}/Γ
$0.92 \pm 0.23 \pm 0.55$	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$	

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

$\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{39}/Γ
0.35 ± 0.09		ABLIKIM 18D	BES3	$\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 1 ABLIKIM 11D BES3 $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

¹ ABLIKIM 11D reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(K^+\bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ
32 ± 21	1 TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$	

¹ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R16;LINKAGE=B

NODE=M055R16;LINKAGE=D
NODE=M055R16;LINKAGE=C

NODE=M055R28
NODE=M055R28

NODE=M055R28;LINKAGE=AT

NODE=M055R64
NODE=M055R64

NODE=M055R64;LINKAGE=A

NODE=M055R65
NODE=M055R65

NODE=M055R65;LINKAGE=A

NODE=M055R66
NODE=M055R66

NODE=M055R66;LINKAGE=A

NODE=M055R67
NODE=M055R67

NODE=M055R67;LINKAGE=A

NODE=M055R68
NODE=M055R68

NODE=M055R18;LINKAGE=BR

NODE=M055R9
NODE=M055R9

NODE=M055R9;LINKAGE=T

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$					Γ_{41}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.44 \pm 0.36 \pm 0.03$	28.4 ± 5.5	1,2 ABLIKIM	04H BES	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$	
1 ABLIKIM 04H reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.					

$\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{42}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-4}$	90	3.2 ± 2.4	1 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$
1 ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

$\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{43}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.35 \pm 0.10 \pm 0.01$	22	1 ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4 K_S^0$	
1 Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (3.4 \pm 0.9 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..					

$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{44}/Γ
VALUE (units 10^{-3})	DOCUMENT ID				
0.54 ± 0.11 OUR FIT					

$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$					Γ_{45}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.41 \pm 0.15 \pm 0.01$	17	1 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2 K^+ 2 K^-$	
1 ABLIKIM 06T reports $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(K^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{46}/Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT		
$3.27 \pm 0.28 \pm 0.46$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$		

$\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$					Γ_{47}/Γ
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT		
$1.62 \pm 0.12 \pm 0.28$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$		

$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{48}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.75 \pm 0.06 \pm 0.08$	373	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$	
1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.					

$\Gamma(\omega \omega)/\Gamma_{\text{total}}$					Γ_{49}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$5.7 \pm 0.7 \pm 0.1$	597	1 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{ hadrons}$	
1 ABLIKIM 11K reports $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M055R26
NODE=M055R26

NODE=M055R26;LINKAGE=AB

NODE=M055R26;LINKAGE=AL

NODE=M055R06
NODE=M055R06NODE=M055R82
NODE=M055R82NODE=M055R14
NODE=M055R14NODE=M055R07
NODE=M055R07

NODE=M055R07;LINKAGE=AB

NODE=M055R69
NODE=M055R69NODE=M055R70
NODE=M055R70

NODE=M055R62;LINKAGE=A

NODE=M055R49
NODE=M055R49

NODE=M055R49;LINKAGE=AL

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{50}/Γ</u>
$0.78 \pm 0.04 \pm 0.08$	628	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$	

1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

 $\Gamma(\omega \phi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{51}/Γ</u>
$0.27 \pm 0.04 \pm 0.01$	105	1 ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<u>0.21 $\pm 0.06 \pm 0.01$</u>	<u>15</u>	<u>2,3 ABLIKIM</u>	<u>11K BES3</u>	<u>$\psi(2S) \rightarrow \gamma$ hadrons</u>
--	-----------	--------------------	-----------------	---

1 ABLIKIM 19J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (2.67 \pm 0.31 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 ABLIKIM 11K reports $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 Superseded by ABLIKIM 19J.

 $\Gamma(\phi \phi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{52}/Γ</u>
$4.2 \pm 0.5 \pm 0.1$	366	1 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons	

1 ABLIKIM 11K reports $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\phi \phi \eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{53}/Γ</u>
$3.0 \pm 0.4 \pm 0.2$	83.6	1 ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma \phi \phi \eta$	

1 ABLIKIM 20B reports $(2.96 \pm 0.43 \pm 0.22) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi \phi \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$.

 $\Gamma(p \bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
0.760 ± 0.034 OUR FIT	

 $\Gamma(p \bar{p} \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{55}/Γ</u>
0.155 ± 0.018 OUR AVERAGE				
0.163 $\pm 0.019 \pm 0.004$	1 ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p \bar{p} X$	

0.112 $\pm 0.047 \pm 0.003$	2 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------------------	---------	---------	---

1 ONYISI 10 reports $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p \bar{p} \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 ATHAR 07 reports $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p \bar{p} \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(p \bar{p} \eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	<u>Γ_{56}/Γ</u>
$0.145 \pm 0.024 \pm 0.004$	1 ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p \bar{p} X$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.15	90	2 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-------	----	---------	---------	---

NODE=M055R61
NODE=M055R61

NODE=M055R61;LINKAGE=A

NODE=M055R50
NODE=M055R50

NODE=M055R50;LINKAGE=A

NODE=M055R50;LINKAGE=B

NODE=M055R51
NODE=M055R51

NODE=M055R51;LINKAGE=AL

NODE=M055R85
NODE=M055R85

NODE=M055R85;LINKAGE=A

NODE=M055R11
NODE=M055R11

NODE=M055R21
NODE=M055R21

NODE=M055R21;LINKAGE=ON

NODE=M055R21;LINKAGE=AT

NODE=M055R27
NODE=M055R27

¹ ONYISI 10 reports $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATTHAR 07 reports $< 0.16 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$		Γ_{57}/Γ		
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT
0.212 ± 0.030 ± 0.005		¹ ONYISI	10	CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

¹ ONYISI 10 reports $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$		Γ_{58}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 × 10⁻⁵	90	¹ ABLIKIM	11F	BES3 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $< 1.82 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$		Γ_{59}/Γ		
VALUE (units 10^{-3})		DOCUMENT ID	TECN	COMMENT
0.50 ± 0.19 OUR EVALUATION		Treating systematic error as correlated.		
0.50 ± 0.19 OUR AVERAGE				

0.46 ± 0.12 ± 0.15 ¹BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c1}$
 1.08 ± 0.77 ± 0.05 ¹TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$		Γ_{60}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5 × 10⁻⁴	90	¹ HE	08B	CLEO $e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

¹ HE 08B reports $< 0.05 \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

$\Gamma(p\bar{p}K^+K^- (\text{non-resonant}))/\Gamma_{\text{total}}$		Γ_{61}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.27 ± 0.22 ± 0.03	82 ± 9	¹ ABLIKIM	11F	BES3 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

¹ ABLIKIM 11F reports $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}K^+K^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$		Γ_{62}/Γ		
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	¹ ABLIKIM	06D	BES2 $\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.1 \pm 0.6)\%$.

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$		Γ_{63}/Γ		
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5 ± 0.1	1412	¹ ABLIKIM	12J	BES3 $\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.37 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R27;LINKAGE=ON

NODE=M055R27;LINKAGE=AT

NODE=M055R43
NODE=M055R43NODE=M055R48
NODE=M055R48

NODE=M055R48;LINKAGE=AB

NODE=M055R7
NODE=M055R7
→ UNCHECKED ←

NODE=M055R7;LINKAGE=X2

NODE=M055R38
NODE=M055R38

NODE=M055R38;LINKAGE=HE

NODE=M055R45
NODE=M055R45

NODE=M055R45;LINKAGE=AB

NODE=M055R02
NODE=M055R02

NODE=M055R;LINKAGE=AB

NODE=M055R56
NODE=M055R56

NODE=M055R56;LINKAGE=AL

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.9 \pm 0.5 \pm 0.1$	1625	1 ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+$
¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.38 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{64}/Γ NODE=M055R57
NODE=M055R57 $\Gamma(p\bar{n}\pi^- \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$10.3 \pm 1.1 \pm 0.2$	1082	1 ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^- \pi^0$
¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.00 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{65}/Γ NODE=M055R58
NODE=M055R58 $\Gamma(\bar{p}n\pi^+ \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$10.1 \pm 1.1 \pm 0.2$	1261	1 ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma \bar{p}n\pi^+ \pi^0$
¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+ \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.98 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{66}/Γ NODE=M055R59
NODE=M055R59 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
1.14 ± 0.11 OUR FIT	

 Γ_{67}/Γ NODE=M055R23
NODE=M055R23 $\Gamma(\Lambda\bar{\Lambda}\pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$29 \pm 5 \pm 1$	105	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}\pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<150	90	2 ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$	
¹ ABLIKIM 12I reports $(31.1 \pm 3.4 \pm 3.9) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 Using $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.1 \pm 0.6)\%$.					

 Γ_{68}/Γ NODE=M055R01
NODE=M055R01 $\Gamma(\Lambda\bar{\Lambda}\pi^+ \pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$25 \pm 6 \pm 1$	13	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}\pi^+ \pi^-$	
¹ ABLIKIM 12I reports $(26.2 \pm 5.5 \pm 3.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+ \pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 Γ_{69}/Γ

NODE=M055R01;LINKAGE=AB

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<1.3 \times 10^{-4}$	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+\bar{\Lambda}\pi^-$	
¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

 Γ_{70}/Γ NODE=M055R52
NODE=M055R52 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<13	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^-\bar{\Lambda}\pi^+$	
¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

 Γ_{71}/Γ NODE=M055R53
NODE=M055R53

NODE=M055R53;LINKAGE=AL

$\Gamma(K^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{72}/Γ
4.2±0.4 OUR AVERAGE Error includes scale factor of 1.2.					
9.0 $^{+2.7}_{-2.3}\pm 0.6$	24	1 LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$	
4.2 $\pm 0.4\pm 0.1$	3k	2,3 ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$	
3.1 $\pm 0.9\pm 0.1$		4 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$	
				1 LU 19 reports $(9.15^{+2.63}_{-2.25} \pm 0.86) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(1P)K^+)]$ assuming $B(B^+ \rightarrow \chi_{c1}(1P)K^+) = (4.79 \pm 0.23) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \chi_{c1}(1P)K^+) = (4.85 \pm 0.33) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
				2 ABLIKIM 13D reports $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
				3 Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.	
				4 ATHAR 07 reports $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	

NODE=M055R22
NODE=M055R22

NODE=M055R22;LINKAGE=A

NODE=M055R22;LINKAGE=AB

NODE=M055R22;LINKAGE=LB
NODE=M055R22;LINKAGE=AT $\Gamma(K^*(892)^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{73}/Γ
4.9±0.7±0.1	328	1 ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^*+\bar{p}\Lambda$	
1 ABLIKIM 19AU reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (4.8 \pm 0.5 \pm 0.4) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M055R86
NODE=M055R86

NODE=M055R86;LINKAGE=F

 $\Gamma(K^+\bar{\Lambda}(1520)+c.c.)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{74}/Γ
1.71±0.44±0.04	48 ± 10	1 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$	
1 ABLIKIM 11F reports $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{\Lambda}(1520)+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M055R46
NODE=M055R46

NODE=M055R46;LINKAGE=AB

 $\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{75}/Γ
<9 × 10⁻⁵	90	1 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$	
1 ABLIKIM 11F reports $< 1.00 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

NODE=M055R47
NODE=M055R47

NODE=M055R47;LINKAGE=AB

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{76}/Γ
4.2±0.6±0.1	103	1 ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{\Sigma}^0$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<6	90	2 ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{\Sigma}^0$		
<4	90	3.8 ± 2.5	3 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Sigma^0\bar{\Sigma}^0$	
1 ABLIKIM 18V reports $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.41 \pm 0.05 \pm 0.03) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						
2 ABLIKIM 13H reports $< 0.62 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.						
3 NAIK 08 reports $< 0.44 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \approx (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.						

NODE=M055R32
NODE=M055R32

NODE=M055R32;LINKAGE=B

NODE=M055R32;LINKAGE=AB

NODE=M055R32;LINKAGE=NA

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{78}/Γ
3.6±0.6±0.1		59	1 ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<8	90	2 ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$		
<6	90	4.3 ± 2.3	3 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$	

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.35 \pm 0.06 \pm 0.02) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 13H reports $< 0.87 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

³ NAIK 08 reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{79}/Γ
<9 × 10⁻⁵	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$	
1 ABLIKIM 12I reports $< 10 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{80}/Γ
<5 × 10⁻⁵	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$	
1 ABLIKIM 12I reports $< 5.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{81}/Γ
1.35±0.24±0.03	49	1 ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	
1 ABLIKIM 15I reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.32 \pm 0.20 \pm 0.12) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{82}/Γ
<6 × 10⁻⁵	90	1.7 ± 2.4	1 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$	
1 NAIK 08 reports $< 0.60 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.						

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{83}/Γ
0.80±0.21±0.02	16.4 ± 4.3	1 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						

<3.4 90 2 ABLIKIM 06D BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ NAIK 08 reports $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.1 \pm 0.6)\%$.

NODE=M055R33
NODE=M055R33

NODE=M055R33;LINKAGE=B

NODE=M055R33;LINKAGE=AB

NODE=M055R33;LINKAGE=NA

NODE=M055R54
NODE=M055R54

NODE=M055R54;LINKAGE=AL

NODE=M055R55
NODE=M055R55

NODE=M055R55;LINKAGE=AL

NODE=M055R71
NODE=M055R71

NODE=M055R71;LINKAGE=A

NODE=M055R34
NODE=M055R34

NODE=M055R34;LINKAGE=NA

NODE=M055R03
NODE=M055R03

NODE=M055R03;LINKAGE=NA

NODE=M055R03;LINKAGE=AB

$\Gamma(\pi^+\pi^-) + \Gamma(K^+K^-)]/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{84}/Γ
$<21 \times 10^{-4}$		1 FELDMAN	77	MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	NODE=M055R2 NODE=M055R2
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<38 \times 10^{-4}$	90	1 BRANDELIK	79B	DASP $\psi(2S) \rightarrow \gamma\chi_{c1}$	

¹ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{85}/Γ
$<6 \times 10^{-5}$	90	1 ABLIKIM	050	BES2 $\psi(2S) \rightarrow \chi_{c1}\gamma$	NODE=M055R04 NODE=M055R04
¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] < 0.6 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					NODE=M055R04;LINKAGE=AB

 $\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{86}/Γ
$<3.2 \times 10^{-3}$	90	1,2 ABLIKIM	13B	BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$	NODE=M055R63 NODE=M055R63
• • • We do not use the following data for averages, fits, limits, etc. • • •					OCCUR=2
$<4.4 \times 10^{-3}$	90	1,3 ABLIKIM	13B	BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$	
¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.					NODE=M055R63;LINKAGE=A
² Using the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.					NODE=M055R63;LINKAGE=B
³ Using the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.					NODE=M055R63;LINKAGE=C

RADIATIVE DECAYS **$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{87}/Γ
34.3 ± 1.0 OUR FIT					NODE=M055R1 NODE=M055R1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$34.75 \pm 0.11 \pm 1.70$	1.9M	1 ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$	
$37.9 \pm 0.8 \pm 2.1$		2 ADAM	05A	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$	
¹ Not independent from $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))$ and the product $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))$ also measured in ABLIKIM 17U.					NODE=M055R1;LINKAGE=A
² Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c1})$ from ATHAR 04.					NODE=M055R1;LINKAGE=AD

 $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{88}/Γ
216 ± 17 OUR AVERAGE					NODE=M055R29 NODE=M055R29
$215 \pm 22 \pm 5$	432 ± 25	1 ABLIKIM	11E	BES3 $\psi(2S) \rightarrow \gamma\gamma\rho^0$	
$217 \pm 24 \pm 5$	186 ± 15	2 BENNETT	08A	CLEO $\psi(2S) \rightarrow \gamma\gamma\rho^0$	
¹ ABLIKIM 11E reports $(228 \pm 13 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\chi_{c1}(1P) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				NODE=M055R29;LINKAGE=AB	
² BENNETT 08A reports $(243 \pm 19 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\chi_{c1}(1P) \rightarrow \gamma\gamma\rho^0)/\Gamma_{\text{total}}]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M055R29;LINKAGE=BE

 $\Gamma(\gamma\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{89}/Γ
68 ± 8 OUR AVERAGE					NODE=M055R30 NODE=M055R30
$66 \pm 9 \pm 2$	136 ± 14	1 ABLIKIM	11E	BES3 $\psi(2S) \rightarrow \gamma\gamma\omega$	
$74 \pm 17 \pm 2$	39 ± 7	2 BENNETT	08A	CLEO $\psi(2S) \rightarrow \gamma\gamma\omega$	
¹ ABLIKIM 11E reports $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\chi_{c1}(1P) \rightarrow \gamma\gamma\omega)/\Gamma_{\text{total}}]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				NODE=M055R30;LINKAGE=AB	
² BENNETT 08A reports $(83 \pm 15 \pm 12) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] \times [B(\chi_{c1}(1P) \rightarrow \gamma\gamma\omega)/\Gamma_{\text{total}}]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				NODE=M055R30;LINKAGE=BE	

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$					Γ_{90}/Γ
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
24±5±1		43 ± 9	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<23	90	5.2 ± 3.1	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
1 ABLIKIM 11E reports $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 BENNETT 08A reports $< 26 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.					

NODE=M055R31
NODE=M055R31

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{91}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 6.3×10^{-6}	90	ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.5×10^{-5}	90	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$	
< 150×10^{-5}	90	¹ YAMADA	77 DASP	$e^+e^- \rightarrow 3\gamma$	
1 Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.					

NODE=M055R31
NODE=M055R3

$\Gamma(e^+e^- J/\psi(1S))/\Gamma_{\text{total}}$					Γ_{92}/Γ
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.65±0.23±0.09	1.9k	^{1,2} ABLIKIM	17I BES3	$\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$	
1 ABLIKIM 17I reports $(3.73 \pm 0.09 \pm 0.25) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 Not independent from other measurements reported by ABLIKIM 17I					

NODE=M055R;LINKAGE=T1
NODE=M055R79
NODE=M055R79

$\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$					Γ_{92}/Γ_{87}
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
10.1±0.3±0.5	1.9k	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$	
1 Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.					
$\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S))$					Γ_{93}/Γ_{92}
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
6.73±0.51±0.50	222	ABLIKIM	19Z BES3	$\psi(2S) \rightarrow \gamma\chi_c \rightarrow \gamma(\mu^+\mu^- J/\psi)$	

NODE=M055R79;LINKAGE=B
NODE=M055R79;LINKAGE=C
NODE=M055R80
NODE=M055R80

$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$					$\Gamma_{54}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
2.14±0.10 OUR FIT					
1.1 ± 1.0	¹ BAI	98I BES	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma p\bar{p}$		
1 Calculated by us. The value for $B(\chi_{c1} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].					

NODE=M055230
NODE=M055B1
NODE=M055B1

$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$					$\Gamma_{67}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
11.1±1.1 OUR FIT	136	¹ ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$	
10.9±1.1 OUR AVERAGE	46 ± 7	² NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$	

NODE=M055B10
NODE=M055B10

¹ Calculated by us. ABLIKIM 13H reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (12.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ from a measurement of $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c1})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.2 \pm 0.4)\%$.

² Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$		$\Gamma_{67}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN COMMENT
3.20 ± 0.30 OUR FIT			
7.1 $\begin{array}{l} +2.8 \\ -2.4 \end{array}$ ± 1.3	$9.0 \begin{array}{l} +3.5 \\ -3.1 \end{array}$	1 BAI	03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ BAI 03E reports [$B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c1}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$] $\times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33 \begin{array}{l} +0.52 \\ -0.46 \end{array} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$		$\Gamma_{67}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN COMMENT
3.34 ± 0.06 OUR FIT			

3.24 ± 0.16 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.			
3.518 ± 0.010 ± 0.120	143k	1 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$	
3.442 ± 0.010 ± 0.132	1.9M	ABLIKIM	17U BES3	$e^+e^- \rightarrow \gamma X$	
2.81 ± 0.05 ± 0.23	13k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$	
2.56 ± 0.12 ± 0.20		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$	
2.78 ± 0.30		OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c1}$	
2.2 ± 0.5		BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma\chi_{c1}$	
2.9 ± 0.5		BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma\chi_{c1}$	
5.0 ± 1.5		BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$	
2.8 ± 0.9		WHITAKER	76 MRK1	e^+e^-	

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.377 ± 0.009 ± 0.183	142k	5 ABLIKIM	120 BES3	$\psi(2S) \rightarrow \gamma\chi_{c1}$
3.56 ± 0.03 ± 0.12	24.9k	6 MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c1}$
3.44 ± 0.06 ± 0.13	3.7k	7 ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

NODE=M055B10;LINKAGE=AB

NODE=M055B10;LINKAGE=NA

NODE=M055B11
NODE=M055B11

NODE=M055B11;LINKAGE=BA

NODE=M055B2
NODE=M055B2

NODE=M055B2;LINKAGE=A

NODE=M055B;LINKAGE=3Q


NODE=M055B;LINKAGE=2Q

NODE=M055B;LINKAGE=EA

NODE=M055B2;LINKAGE=B

NODE=M055B2;LINKAGE=ME

NODE=M055B;LINKAGE=AD



$10^{-2})$

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{anything}) = \frac{\Gamma_{87} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_9^{\psi(2S)}}{\Gamma_{87} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_9^{\psi(2S)} + (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + 0.343 \Gamma_{154}^{\psi(2S)} + 0.190 \Gamma_{155}^{\psi(2S)})}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.43±0.10 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.70 \pm 0.04 \pm 0.15$	24.9k	¹ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
$5.77 \pm 0.10 \pm 0.12$	3.7k	ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = \frac{\Gamma_{87} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}{\Gamma_{87} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.63±0.17 OUR FIT**10.15±0.28 OUR AVERAGE**

$10.17 \pm 0.07 \pm 0.27$	24.9k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
$12.6 \pm 0.3 \pm 3.8$	3k	¹ ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
8.5 ± 2.1		² HIMEL	80 MRK2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$10.24 \pm 0.17 \pm 0.23$	3.7k	³ ADAM	05A CLEO	Repl. by MENDEZ 08
---------------------------	------	-------------------	----------	--------------------

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$ quoted in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³ Not independent from other values reported by ADAM 05A.

$$\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} = \frac{\Gamma_{17} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}{\Gamma_{17} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.8±0.5 OUR FIT**7.2±0.6 OUR AVERAGE**

$7.3 \pm 0.5 \pm 0.5$	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
$7.0 \pm 0.5 \pm 0.9$	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K^0 K^+ \pi^- + \text{c.c.})$ reported by ATHAR 07 was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$. We use $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$.

$$\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) / \Gamma_{\text{total}} = \frac{\Gamma_{17} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}{\Gamma_{17} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

19.6±1.6 OUR FIT**13.2±2.4±3.2**

¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
------------------	---------	---

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} = \frac{\Gamma_{44} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}{\Gamma_{44} / \Gamma \times \Gamma_{154}^{\psi(2S)} / \Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.53±0.11 OUR FIT**0.61±0.11±0.08**

¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$
----------------------	----------	---

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$.

NODE=M055B7

NODE=M055B7

NODE=M055B7

NODE=M055B7;LINKAGE=ME

NODE=M055B3

NODE=M055B3

NODE=M055B;LINKAGE=AB

NODE=M055B;LINKAGE=J3

NODE=M055B3;LINKAGE=AD

NODE=M055B16

NODE=M055B16

NODE=M055B16;LINKAGE=AT

NODE=M055B16;LINKAGE=AB

NODE=M055B17

NODE=M055B17

NODE=M055B17;LINKAGE=BA

NODE=M055B14

NODE=M055B14

NODE=M055B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} / \frac{\Gamma_{44}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

1.52 ± 0.31 OUR FIT

1.13 ± 0.40 ± 0.29

1 BAI

99B

BES

$\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{54}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT


7.41 ± 0.35 OUR FIT

7.8 ± 0.6 OUR AVERAGE

Error includes scale factor of 1.4. See the ideogram below.

7.9 ± 0.4	± 0.3	453	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
8.2 ± 0.7	± 0.4	141 ± 13	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
4.8 ± 1.4	± 0.6	18.2 ± 5.5	BAI	04F	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}(1P) \rightarrow \gamma \bar{p}p$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow p\bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.



$$\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}} (\text{units } 10^{-6})$$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{77}/\Gamma \times \Gamma_{154}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

1.49 ± 0.09 ± 0.07	258 ± 17	¹ ABLIKIM	19BB	BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$
---------------------------	----------	----------------------	------	------	--

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_{c1} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (1.53 \pm 0.10 \pm 0.08) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.75 \pm 0.24)\%$ and other branching fractions from PDG 18.

MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

$$a_2 = M2/\sqrt{E1^2 + M2^2} \text{ Magnetic quadrupole fractional transition amplitude}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

-6.7 ± 0.9 OUR AVERAGE Error includes scale factor of 2.6. See the ideogram below.

-7.40 ± 0.33 ± 0.34	164k	¹ ABLIKIM	17N	BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
-6.26 ± 0.63 ± 0.24	39k	ARTUSO	09	CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.2 ± 3.2 ± 0.4	2090	AMBROGIANI	02	E835	$p\bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi \gamma$
-0.2 ± 0.8	921	OREGLIA	82	CBAL	$\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$

NODE=M055B15

NODE=M055B15

NODE=M055B15;LINKAGE=BA

NODE=M055B6

NODE=M055B6

NODE=M055B6;LINKAGE=NA

NODE=M055B01

NODE=M055B01

NODE=M055B01;LINKAGE=A


NODE=M055240

NODE=M055A1

NODE=M055A1

¹ Correlated with b_2 with correlation coefficient $\rho_{a_2 b_2} = 0.133$.

NODE=M055A1;LINKAGE=A



MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma \chi_{c1}(1S)$ RADIATIVE DECAY

$b_2 = M^2 / \sqrt{E^2 + M^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.4 OUR AVERAGE				
2.29 ± 0.39 ± 0.27	164k	¹ ABLIKIM	17N	BES3 $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
2.76 ± 0.73 ± 0.23	39k	ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
7.7 $^{+5.0}_{-4.5}$	921	OREGLIA	82	CBAL $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Correlated with a_2 with correlation coefficient $\rho_{a_2 b_2} = 0.133$.

NODE=M055250

NODE=M055QB2
NODE=M055QB2

NODE=M055QB2;LINKAGE=A

NODE=M055260

NODE=M055QAR
NODE=M055QAR

NODE=M055QAR;LINKAGE=AR

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS $\psi(2S) \rightarrow \gamma \chi_{c1}(1S)$ and $\chi_{c1} \rightarrow \gamma J/\psi(1S)$

a_2/b_2 Magnetic quadrupole transition amplitude ratio

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-2.27 $^{+0.57}_{-0.99}$	39k	¹ ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ Statistical and systematic errors combined. Not independent of $a_2(\chi_{c1})$ and $b_2(\chi_{c1})$ values from ARTUSO 09.

NODE=M055

REFID=60212
REFID=59844
REFID=59996;ERROR=14
REFID=60026
REFID=59606
REFID=59837
REFID=59614
REFID=58849
REFID=58990
REFID=59304
REFID=58191
REFID=58278
REFID=58310
REFID=57931
REFID=57953
REFID=57978
REFID=58026
REFID=57140
REFID=56774
REFID=56778
REFID=55901
REFID=54877
REFID=54879
REFID=54953
REFID=55583
REFID=54736

$\chi_{c1}(1P)$ REFERENCES

ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU	PR D100 052010	Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17B1	PR 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I	PR 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16715
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05A	PRC 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48552
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41865
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41907
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)	REFID=22145
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
Also		Private Comm.	M.J. Oreglia	(EF) (LBL, SLAC)	REFID=22143
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, UCB)	REFID=22119
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)	REFID=22064
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)	REFID=22106

$h_c(1P)$ $I^G(J^{PC}) = 0^-(1^{+-})$

Quantum numbers are quark model prediction, $C = -$ established by $\eta_c \gamma$ decay.

 $h_c(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3525.38±0.11 OUR AVERAGE				
3525.31±0.11±0.14	832	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
3525.40±0.13±0.18	3679	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
3525.20±0.18±0.12	1282	2 DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3525.8 ± 0.2 ± 0.2	13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3525.6 ± 0.5	92+23 -22	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2(\pi^+ \pi^- \pi^0)$
3524.4 ± 0.6 ± 0.4	168 ± 40	3 ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3527 ± 8	42	ANTONIAZZI	94 E705	300 π^\pm , $p\text{Li} \rightarrow J/\psi \pi^0 X$
3526.28±0.18±0.19	59	4 ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$
3525.4 ± 0.8 ± 0.4	5	BAGLIN	86 SPEC	$\bar{p}p \rightarrow J/\psi X$

¹ With floating width.² Combination of exclusive and inclusive analyses for the reaction $\psi(2S) \rightarrow \pi^0 h_c \rightarrow \pi^0 \eta_c \gamma$. This result is the average of DOBBS 08A and ROSNER 05.³ Superseded by DOBBS 08A.⁴ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.

NODE=M144

NODE=M144M

NODE=M144M

NODE=M144M;LINKAGE=AB

NODE=M144M;LINKAGE=DO

NODE=M144M;LINKAGE=RO

NODE=M144M;LINKAGE=NW

 $h_c(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.28±0.22		832	1 ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.44	90	3679	2 ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
< 1		13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
< 1.1	90	59	ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$

¹ With floating mass.² The central value is $\Gamma = 0.73 \pm 0.45 \pm 0.28$ MeV.

NODE=M144W

NODE=M144W

NODE=M144W;LINKAGE=AL

NODE=M144W;LINKAGE=AB

NODE=M144215;NODE=M144

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 J/\psi(1S) \pi^0$		
$\Gamma_2 J/\psi(1S) \pi \pi$	not seen	
$\Gamma_3 J/\psi(1S) \pi^+ \pi^-$	< 2.3 $\times 10^{-3}$	90%
$\Gamma_4 p\bar{p}$	< 1.5 $\times 10^{-4}$	90%
$\Gamma_5 p\bar{p} \pi^+ \pi^-$	(2.9±0.6) $\times 10^{-3}$	
$\Gamma_6 \pi^+ \pi^- \pi^0$	(1.6±0.5) $\times 10^{-3}$	
$\Gamma_7 2\pi^+ 2\pi^- \pi^0$	(8.1±1.8) $\times 10^{-3}$	
$\Gamma_8 3\pi^+ 3\pi^- \pi^0$	< 9 $\times 10^{-3}$	90%
$\Gamma_9 K^+ K^- \pi^+ \pi^-$	< 6 $\times 10^{-4}$	90%
Radiative decays		
$\Gamma_{10} \gamma \eta$	(4.7±2.1) $\times 10^{-4}$	
$\Gamma_{11} \gamma \eta'(958)$	(1.5±0.4) $\times 10^{-3}$	
$\Gamma_{12} \gamma \eta_c(1S)$	(51 ± 6) %	

DESIG=1

DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow

DESIG=10

DESIG=3

DESIG=11

DESIG=5

DESIG=6

DESIG=7

DESIG=12

NODE=M144;CLUMP=R

DESIG=9

DESIG=8

DESIG=4

NODE=M144220

 $h_c(1P)$ PARTIAL WIDTHS

$h_c(1P) \Gamma(i)\Gamma(\bar{p}p)/\Gamma(\text{total})$

$\Gamma(\gamma\eta_c(1S)) \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$	$\Gamma_{12}\Gamma_4/\Gamma$			
<u>VALUE (eV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
12.0 \pm 4.5	13	1 ANDREOTTI 05B E835	$\bar{p}p \rightarrow \eta_c \gamma$	
1 Assuming $\Gamma = 1$ MeV.				

 $h_c(1P)$ BRANCHING RATIOS

$\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$	Γ_2/Γ_1			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.18	90	ARMSTRONG 92D	E760	$\bar{p}p \rightarrow J/\psi\pi^0$

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_3/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<2.3 \times 10^{-3}$	90	1 ABLIKIM 18M	BES3	$\psi(2S) \rightarrow \pi^0\pi^+\pi^- J/\psi$

1 ABLIKIM 18M reports $[\Gamma(h_c(1P) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 2.0 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ				
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.6 \pm 0.5 \pm 0.2$	101	1 ABLIKIM	19AG	BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<2.2	90	2 ADAMS	09	CLEO	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

1 ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (1.38 \pm 0.35 \pm 0.17) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 0.19 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.

$\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_7/Γ			
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.81 ± 0.18 OUR AVERAGE				
$[(2.2 \pm 0.8) \times 10^{-2}$ OUR 2019 AVERAGE]				
0.74 \pm 0.14 \pm 0.11	254	1 ABLIKIM	19AG	BES3
$2.2 \begin{array}{l} +0.8 \\ -0.6 \end{array} \pm 0.3$	92	2 ADAMS	09	CLEO

1 ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow 2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (6.40 \pm 0.81 \pm 0.87) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (1.88 \pm 0.48 \pm 0.47) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_8/Γ			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<9 \times 10^{-3}$ (CL = 90%)		[$<2.9 \times 10^{-2}$ OUR 2019 BEST LIMIT]		
$<9 \times 10^{-3}$	90	1 ABLIKIM	19AG	BES3
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<0.029	90	2 ADAMS	09	CLEO

1 ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow 3\pi^+3\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 7.5 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.

2 ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 3\pi^+3\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 2.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.

NODE=M144223

NODE=M144G1
NODE=M144G1

NODE=M144G1;LINKAGE=AN

NODE=M144225

NODE=M144R1
NODE=M144R1NODE=M144R07
NODE=M144R07
OCCUR=2

NODE=M144R07;LINKAGE=B

NODE=M144R01
NODE=M144R01

NODE=M144R01;LINKAGE=A

NODE=M144R02
NODE=M144R02
NEW

NODE=M144R02;LINKAGE=A

NODE=M144R02;LINKAGE=AD

NODE=M144R03
NODE=M144R03

NODE=M144R03;LINKAGE=A

NODE=M144R03;LINKAGE=AD

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$2.9 \pm 0.5 \pm 0.4$	230	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$	
1 ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (2.49 \pm 0.27 \pm 0.28) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6 \times 10^{-4}$	90	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$	
1 ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] < 0.5 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = 8.6 \times 10^{-4}$.					

RADIATIVE DECAYS

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
$4.7 \pm 1.5 \pm 1.4$	18	ABLIKIM	16I	BES3	$\psi(2S) \rightarrow \pi^0 \gamma\eta$

$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.52 \pm 0.27 \pm 0.29$	44	ABLIKIM	16I	BES3	$\psi(2S) \rightarrow \pi^0 \gamma\eta'(958)$

$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
51 ± 6 OUR AVERAGE					
54.3 \pm 6.7 \pm 5.2	3679	ABLIKIM	10B	BES3	$\psi(2S) \rightarrow \pi^0 \gamma\eta_c$
48 \pm 6 \pm 7		¹ DOBBS	08A	CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
48 \pm 6 \pm 7	1282	² DOBBS	08A	CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
46 \pm 12 \pm 7	168	³ ROSNER	05	CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

1 Average of DOBBS 08A and ROSNER 05. DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (4.16 \pm 0.30 \pm 0.37) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
2 DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (4.19 \pm 0.32 \pm 0.45) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
3 ROSNER 05 reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \pi^0 h_c(1P))] = (4.0 \pm 0.8 \pm 0.7) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \pi^0 h_c(1P)) = (8.6 \pm 1.3) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(h_c(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_{15}^{\psi(2S)}/\Gamma^{\psi(2S)}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-7}$	90	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

$\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \pi^0 h_c(1P))/\Gamma_{\text{total}}$					$\Gamma_{12}/\Gamma \times \Gamma_{15}^{\psi(2S)}/\Gamma^{\psi(2S)}$
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.3 ± 0.4 OUR AVERAGE					
4.58 \pm 0.40 \pm 0.50	3679	¹ ABLIKIM	10B	BES3	$\psi(2S) \rightarrow \pi^0 \gamma X$
4.16 \pm 0.30 \pm 0.37	1430	² DOBBS	08A	CLEO	$\psi(2S) \rightarrow \pi^0 \gamma\eta_c$

¹ Not independent of other branching fractions in ABLIKIM 10B.

² Not independent of other branching fractions in DOBBS 08A.

NODE=M144R08
NODE=M144R08

NODE=M144R08;LINKAGE=A

NODE=M144R09
NODE=M144R09

NODE=M144R09;LINKAGE=A

NODE=M144230

NODE=M144R06
NODE=M144R06

NODE=M144R00
NODE=M144R00

NODE=M144R2
NODE=M144R2

OCCUR=2

NODE=M144R2;LINKAGE=DB

NODE=M144R2;LINKAGE=DO

NODE=M144R2;LINKAGE=RO

NODE=M144235

NODE=M144R05
NODE=M144R05

NODE=M144R04
NODE=M144R04

NODE=M144R04;LINKAGE=AB

NODE=M144R04;LINKAGE=DO

$\chi_c(1P)$ REFERENCES

ABLIKIM	19AG	PR D99 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18M	PR D97 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16I	PRL 116 251802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12N	PR D85 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)
ANDREOTTI	05B	PR D72 032001	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)
ANTONIAZZI	94	PR D50 4258	L. Antoniazz <i>et al.</i>	(E705 Collab.)
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)
ARMSTRONG	92D	PRL 69 2337	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
BAGLIN	86	PL B171 135	C. Baglin <i>et al.</i>	(LAPP, CERN, TORI, STRB+)

NODE=M144

REFID=59858
REFID=58901
REFID=57450
REFID=55583
REFID=54741
REFID=53348
REFID=53103
REFID=52579
REFID=50768
REFID=50812
REFID=49579
REFID=44074
REFID=43307
REFID=43174
REFID=43180

NODE=M057

 $\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\chi_{c2}(1P)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
3556.17 ± 0.07 OUR AVERAGE				
3557.3 ± 1.7 ± 0.7	611	1 AAIJ	17BB LHCb	$p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3556.10 ± 0.06 ± 0.11	4.0k	2 AAIJ	17BI LHCb	$\chi_{c2} \rightarrow J/\psi\mu^+\mu^-$
3555.3 ± 0.6 ± 2.2	2.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \text{hadrons}$
3555.70 ± 0.59 ± 0.39		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3556.173 ± 0.123 ± 0.020		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3559.9 ± 2.9		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
3556.4 ± 0.7		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3556.22 ± 0.131 ± 0.020	585	3 ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3556.9 ± 0.4 ± 0.5	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
3557.8 ± 0.2 ± 4		4 GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3553.4 ± 2.2	66	5 LEMOIGNE	82 GOLI	$185\pi^-Be \rightarrow \gamma\mu^+\mu^-A$
3555.9 ± 0.7		6 OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3557 ± 1.5	69	7 HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3551 ± 11	15	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4		7 BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4 ± 4		7,8 TANENBAUM	78 MRK1	e^+e^-
3563 ± 7	360	7 BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3555.4 ± 1.3	53	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
3543 ± 10	4	WHITAKER	76 MRK1	$e^+e^- \rightarrow J/\psi 2\gamma$

1 From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c2}(1P)$ fixed to the PDG 16 value.

2 AAIJ 17BI reports also $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$ MeV.

3 Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

4 Using mass of $\psi(2S) = 3686.0$ MeV.

5 $J/\psi(1S)$ mass constrained to 3097 MeV.

6 Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

7 Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

8 From a simultaneous fit to radiative and hadronic decay channels.

NODE=M057

NODE=M057M

NODE=M057M

NODE=M057M;LINKAGE=A

NODE=M057M;LINKAGE=B

NODE=M057M;LINKAGE=NW

NODE=M057M;LINKAGE=C

NODE=M057M;LINKAGE=P

NODE=M057M;LINKAGE=E

NODE=M057M;LINKAGE=D

NODE=M057M;LINKAGE=M

NODE=M057W

NODE=M057W

$\chi_{c2}(1P)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1.97 ± 0.09 OUR FIT				
2.00 ± 0.11 OUR AVERAGE				
2.10 ± 0.20 ± 0.02	4.0k	AAIJ	17BI LHCb	$\chi_{c2} \rightarrow J/\psi\mu^+\mu^-$
1.915 ± 0.188 ± 0.013		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
1.96 ± 0.17 ± 0.07	585	1 ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-$
2.6 +1.4 -1.0	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
2.8 +2.1 -2.0		2 GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

NODE=M057W

¹ Recalculated by ANDREOTTI 05A.² Errors correspond to 90% confidence level; authors give only width range. **$\chi_{c2}(1P)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Confidence level	
Hadronic decays			
Γ_1 $2(\pi^+ \pi^-)$	(1.02 ± 0.09) %		NODE=M057;CLUMP=A DESIG=3
Γ_2 $\rho\rho$			DESIG=43
Γ_3 $\pi^+ \pi^- \pi^0 \pi^0$	(1.83 ± 0.23) %		DESIG=50
Γ_4 $\rho^+ \pi^- \pi^0 + \text{c.c.}$	(2.19 ± 0.34) %		DESIG=51
Γ_5 $4\pi^0$	(1.11 ± 0.15) $\times 10^{-3}$		DESIG=62
Γ_6 $K^+ K^- \pi^0 \pi^0$	(2.1 ± 0.4) $\times 10^{-3}$		DESIG=52
Γ_7 $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(1.38 ± 0.20) %		DESIG=54
Γ_8 $\rho^- K^+ \bar{K}^0 + \text{c.c.}$	(4.1 ± 1.2) $\times 10^{-3}$		DESIG=55
Γ_9 $K^*(892)^0 K^- \pi^+ \rightarrow$ $K^- \pi^+ K^0 \pi^0 + \text{c.c.}$	(2.9 ± 0.8) $\times 10^{-3}$		DESIG=60
Γ_{10} $K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(3.8 ± 0.9) $\times 10^{-3}$		DESIG=56
Γ_{11} $K^*(892)^- K^+ \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(3.7 ± 0.8) $\times 10^{-3}$		DESIG=57
Γ_{12} $K^*(892)^+ \bar{K}^0 \pi^- \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(2.9 ± 0.8) $\times 10^{-3}$		DESIG=58
Γ_{13} $K^+ K^- \eta \pi^0$	(1.3 ± 0.4) $\times 10^{-3}$		DESIG=59
Γ_{14} $K^+ K^- \pi^+ \pi^-$	(8.4 ± 0.9) $\times 10^{-3}$		DESIG=5
Γ_{15} $K^+ K^- \pi^+ \pi^- \pi^0$	(1.17 ± 0.13) %		DESIG=67
Γ_{16} $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	(7.3 ± 0.8) $\times 10^{-3}$		DESIG=78
Γ_{17} $K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	(2.1 ± 1.1) $\times 10^{-3}$		DESIG=10
Γ_{18} $K^*(892)^0 \bar{K}^* (892)^0$	(2.3 ± 0.4) $\times 10^{-3}$		DESIG=21
Γ_{19} $3(\pi^+ \pi^-)$	(8.6 ± 1.8) $\times 10^{-3}$		DESIG=4
Γ_{20} $\phi\phi$	(1.06 ± 0.09) $\times 10^{-3}$		DESIG=16
Γ_{21} $\phi\phi\eta$	(5.3 ± 0.6) $\times 10^{-4}$		DESIG=99
Γ_{22} $\omega\omega$	(8.4 ± 1.0) $\times 10^{-4}$		DESIG=25
Γ_{23} $\omega K^+ K^-$	(7.3 ± 0.9) $\times 10^{-4}$		DESIG=79
Γ_{24} $\omega\phi$	(9.6 ± 2.7) $\times 10^{-6}$		DESIG=68
Γ_{25} $\pi\pi$	(2.23 ± 0.09) $\times 10^{-3}$		DESIG=22
Γ_{26} $\rho^0 \pi^+ \pi^-$	(3.7 ± 1.6) $\times 10^{-3}$		DESIG=9
Γ_{27} $\pi^+ \pi^- \pi^0 (\text{non-resonant})$	(2.0 ± 0.4) $\times 10^{-5}$		DESIG=95
Γ_{28} $\rho(770)^\pm \pi^\mp$	(6 ± 4) $\times 10^{-6}$		DESIG=96
Γ_{29} $\pi^+ \pi^- \eta$	(4.8 ± 1.3) $\times 10^{-4}$		DESIG=39
Γ_{30} $\pi^+ \pi^- \eta'$	(5.0 ± 1.8) $\times 10^{-4}$		DESIG=42
Γ_{31} $\eta\eta$	(5.4 ± 0.4) $\times 10^{-4}$		DESIG=14
Γ_{32} $K^+ K^-$	(1.01 ± 0.06) $\times 10^{-3}$		DESIG=2
Γ_{33} $K_S^0 K_S^0$	(5.2 ± 0.4) $\times 10^{-4}$		DESIG=15
Γ_{34} $K^*(892)^\pm K^\mp$	(1.44 ± 0.21) $\times 10^{-4}$		DESIG=87
Γ_{35} $K^*(892)^0 \bar{K}^0 + \text{c.c.}$	(1.24 ± 0.27) $\times 10^{-4}$		DESIG=88
Γ_{36} $K_2^*(1430)^\pm K^\mp$	(1.48 ± 0.12) $\times 10^{-3}$		DESIG=89
Γ_{37} $K_2^*(1430)^0 \bar{K}^0 + \text{c.c.}$	(1.24 ± 0.17) $\times 10^{-3}$		DESIG=90
Γ_{38} $K_3^*(1780)^\pm K^\mp$	(5.2 ± 0.8) $\times 10^{-4}$		DESIG=91
Γ_{39} $K_3^*(1780)^0 \bar{K}^0 + \text{c.c.}$	(5.6 ± 2.1) $\times 10^{-4}$		DESIG=92
Γ_{40} $a_2(1320)^0 \pi^0$	(1.29 ± 0.34) $\times 10^{-3}$		DESIG=93
Γ_{41} $a_2(1320)^\pm \pi^\mp$	(1.8 ± 0.6) $\times 10^{-3}$		DESIG=94
Γ_{42} $\bar{K}^0 K^+ \pi^- + \text{c.c.}$	(1.28 ± 0.18) $\times 10^{-3}$		DESIG=17
Γ_{43} $K^+ K^- \pi^0$	(3.0 ± 0.8) $\times 10^{-4}$		DESIG=36
Γ_{44} $K^+ K^- \eta$	< 3.2×10^{-4}	90%	DESIG=40
Γ_{45} $K^+ K^- \eta'(958)$	(1.94 ± 0.34) $\times 10^{-4}$		DESIG=82
Γ_{46} $\eta\eta'$	(2.2 ± 0.5) $\times 10^{-5}$		DESIG=34
Γ_{47} $\eta'\eta'$	(4.6 ± 0.6) $\times 10^{-5}$		DESIG=35

Γ_{48}	$\pi^+ \pi^- K_S^0 K_S^0$	$(2.2 \pm 0.5) \times 10^{-3}$		DESIG=29
Γ_{49}	$K^+ K^- K_S^0 K_S^0$	$< 4 \times 10^{-4}$	90%	DESIG=30
Γ_{50}	$K_S^0 K_S^0 K_S^0 K_S^0$	$(1.13 \pm 0.18) \times 10^{-4}$		DESIG=97
Γ_{51}	$K^+ K^- K^+ K^-$	$(1.65 \pm 0.20) \times 10^{-3}$		DESIG=24
Γ_{52}	$K^+ K^- \phi$	$(1.42 \pm 0.29) \times 10^{-3}$		DESIG=32
Γ_{53}	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(4.8 \pm 0.7) \times 10^{-3}$		DESIG=83
Γ_{54}	$K^+ K^- \pi^0 \phi$	$(2.7 \pm 0.5) \times 10^{-3}$		DESIG=84
Γ_{55}	$\phi \pi^+ \pi^- \pi^0$	$(9.3 \pm 1.2) \times 10^{-4}$		DESIG=80
Γ_{56}	$p\bar{p}$	$(7.33 \pm 0.33) \times 10^{-5}$		DESIG=11
Γ_{57}	$p\bar{p}\pi^0$	$(4.7 \pm 0.4) \times 10^{-4}$		DESIG=37
Γ_{58}	$p\bar{p}\eta$	$(1.74 \pm 0.25) \times 10^{-4}$		DESIG=41
Γ_{59}	$p\bar{p}\omega$	$(3.6 \pm 0.4) \times 10^{-4}$		DESIG=61
Γ_{60}	$p\bar{p}\phi$	$(2.8 \pm 0.9) \times 10^{-5}$		DESIG=66
Γ_{61}	$p\bar{p}\pi^+\pi^-$	$(1.32 \pm 0.34) \times 10^{-3}$		DESIG=8
Γ_{62}	$p\bar{p}\pi^0\pi^0$	$(7.8 \pm 2.3) \times 10^{-4}$		DESIG=53
Γ_{63}	$p\bar{p}K^+K^- \text{(non-resonant)}$	$(1.91 \pm 0.32) \times 10^{-4}$		DESIG=63
Γ_{64}	$p\bar{p}K_S^0 K_S^0$	$< 7.9 \times 10^{-4}$	90%	DESIG=28
Γ_{65}	$p\bar{n}\pi^-$	$(8.5 \pm 0.9) \times 10^{-4}$		DESIG=31
Γ_{66}	$\bar{p}n\pi^+$	$(8.9 \pm 0.8) \times 10^{-4}$		DESIG=75
Γ_{67}	$p\bar{n}\pi^-\pi^0$	$(2.17 \pm 0.18) \times 10^{-3}$		DESIG=76
Γ_{68}	$\bar{p}n\pi^+\pi^0$	$(2.11 \pm 0.18) \times 10^{-3}$		DESIG=77
Γ_{69}	$\Lambda\bar{\Lambda}$	$(1.84 \pm 0.15) \times 10^{-4}$		DESIG=19
Γ_{70}	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(1.25 \pm 0.15) \times 10^{-3}$		DESIG=27
Γ_{71}	$\Lambda\bar{\Lambda}\pi^+\pi^- \text{(non-resonant)}$	$(6.6 \pm 1.5) \times 10^{-4}$		DESIG=70
Γ_{72}	$\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.}$	$< 4 \times 10^{-4}$	90%	DESIG=71
Γ_{73}	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.}$	$< 6 \times 10^{-4}$	90%	DESIG=72
Γ_{74}	$K^+\bar{p}\Lambda + \text{c.c.}$	$(7.8 \pm 0.5) \times 10^{-4}$		DESIG=38
Γ_{75}	$K^*(892)^+\bar{p}\Lambda + \text{c.c.}$	$(8.2 \pm 1.1) \times 10^{-4}$		DESIG=101
Γ_{76}	$K^+\bar{p}\Lambda(1520) + \text{c.c.}$	$(2.8 \pm 0.7) \times 10^{-4}$		DESIG=64
Γ_{77}	$\Lambda(1520)\bar{\Lambda}(1520)$	$(4.6 \pm 1.5) \times 10^{-4}$		DESIG=65
Γ_{78}	$\Sigma^0\bar{\Sigma}^0$	$(3.7 \pm 0.6) \times 10^{-5}$		DESIG=47
Γ_{79}	$\Sigma^+\bar{p}K_S^0 + \text{c.c.}$	$(8.2 \pm 0.9) \times 10^{-5}$		DESIG=100
Γ_{80}	$\Sigma^+\bar{\Sigma}^-$	$(3.4 \pm 0.7) \times 10^{-5}$		DESIG=48
Γ_{81}	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$< 1.6 \times 10^{-4}$	90%	DESIG=73
Γ_{82}	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$< 8 \times 10^{-5}$	90%	DESIG=74
Γ_{83}	$K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.76 \pm 0.32) \times 10^{-4}$		DESIG=85
Γ_{84}	$\Xi^0\bar{\Xi}^0$	$< 1.0 \times 10^{-4}$	90%	DESIG=49
Γ_{85}	$\Xi^-\bar{\Xi}^+$	$(1.42 \pm 0.32) \times 10^{-4}$		DESIG=26
Γ_{86}	$J/\psi(1S)\pi^+\pi^-\pi^0$	$< 1.5 \%$	90%	DESIG=12
Γ_{87}	$\pi^0\eta_c$	$< 3.2 \times 10^{-3}$	90%	DESIG=81
Γ_{88}	$\eta_c(1S)\pi^+\pi^-$	$< 5.4 \times 10^{-3}$	90%	DESIG=69

Radiative decays

Γ_{89}	$\gamma J/\psi(1S)$	$(19.0 \pm 0.5) \%$		NODE=M057;CLUMP=B
Γ_{90}	$\gamma\rho^0$	$< 1.9 \times 10^{-5}$	90%	DESIG=6
Γ_{91}	$\gamma\omega$	$< 6 \times 10^{-6}$	90%	DESIG=44
Γ_{92}	$\gamma\phi$	$< 7 \times 10^{-6}$	90%	DESIG=45
Γ_{93}	$\gamma\gamma$	$(2.85 \pm 0.10) \times 10^{-4}$		DESIG=46
Γ_{94}	$e^+e^-J/\psi(1S)$	$(2.15 \pm 0.14) \times 10^{-3}$		DESIG=7
Γ_{95}	$\mu^+\mu^-J/\psi(1S)$	$(2.02 \pm 0.33) \times 10^{-4}$		DESIG=86
				DESIG=98

$242 \pm 65 \pm 51$	$1,4$	ACKER..K...	98	OPAL	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
$150 \pm 42 \pm 36$	$1,5$	DOMINICK	94	CLE2	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
$470 \pm 240 \pm 120$	$1,6$	BAUER	93	TPC	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

¹ Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1187 \pm 0.0008$.

² All systematic errors added in quadrature.

³ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACCIARRI 99E is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.0162 \pm 0.0014$.

⁴ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACKERSTAFF,K 98 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1203 \pm 0.0038$.

⁵ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in DOMINICK 94 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0597 \pm 0.0025$.

⁶ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in BAUER 93 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0597 \pm 0.0025$.

$\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

5.7 ± 0.5 OUR FIT

5.2 ± 0.7 OUR AVERAGE

$5.01 \pm 0.44 \pm 0.55$	1597 ± 138	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
$6.4 \pm 1.8 \pm 0.8$		EISENSTEIN	01	CLE2	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>CL%</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7.8	90	<598	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
--------	----	--------	--------	----	------	--

$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{14}\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

4.7 ± 0.5 OUR FIT

$4.42 \pm 0.42 \pm 0.53$	780 ± 74	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^-$
--------------------------	--------------	--------	----	------	--

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{15}\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

$6.5 \pm 0.9 \pm 1.5$	1250	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-----------------------	--------	-----------------	------	--

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{18}\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

1.26 ± 0.24 OUR FIT

$0.8 \pm 0.17 \pm 0.27$	151 ± 30	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+ K^- \pi^+ \pi^-$
-------------------------	--------------	--------	----	------	--

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{20}\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

0.60 ± 0.05 OUR FIT

$0.62 \pm 0.07 \pm 0.05$	89 ± 11	¹ LIU	12B	BELL	$\gamma\gamma \rightarrow 2(K^+ K^-)$
--------------------------	-------------	------------------	-----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.58 \pm 0.18 \pm 0.16$	26.5 ± 8.1	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$
--------------------------	----------------	--------	----	------	---

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{22}\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>CL%</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.64	90	¹ LIU	12B	BELL	$\gamma\gamma \rightarrow 2(\pi^+ \pi^- \pi^0)$
---------	----	------------------	-----	------	---

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{24}\Gamma_{93}/\Gamma$
<u>VALUE (eV)</u> <u>CL%</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	90	¹ LIU	12B	BELL	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
---------	----	------------------	-----	------	--

¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

NODE=M057G;LINKAGE=LL

NODE=M057G;LINKAGE=GT

NODE=M057G;LINKAGE=J4

NODE=M057G;LINKAGE=J5

NODE=M057G;LINKAGE=J6

NODE=M057G;LINKAGE=J7

NODE=M057224

NODE=M057G3

NODE=M057G3

NODE=M057G08

NODE=M057G08

NODE=M057G09

NODE=M057G09

NODE=M057G02

NODE=M057G02

NODE=M057G10

NODE=M057G10

NODE=M057G12

NODE=M057G12

NODE=M057G03

NODE=M057G03

NODE=M057G03;LINKAGE=LI

NODE=M057G04

NODE=M057G04

NODE=M057G04;LINKAGE=LI

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{25}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.25±0.07 OUR FIT				
1.18±0.25 OUR AVERAGE				
1.44±0.54±0.47	34 ± 13	1 UEHARA	09 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1.14±0.21±0.17	54 ± 10	2 NAKAZAWA	05 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
1 We multiplied the measurement by 3 to convert from $\pi^0 \pi^0$ to $\pi\pi$. Interference with the continuum included.				
2 We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.				
$\Gamma(\rho^0 \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{26}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.9 OUR FIT				
3.2±1.9±0.5	986 ± 578	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+ \pi^-)$
$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{31}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.53±0.22±0.09	8	1 UEHARA	10A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1 Interference with the continuum not included.				
$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{32}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.56±0.04 OUR FIT				
0.44±0.11±0.07	33 ± 8	NAKAZAWA	05	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{33}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.294±0.025 OUR FIT				
0.27 ±0.07 ±0.03	53	1 UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.31 ± 0.05 ± 0.03	38 ± 7	CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
1 Supersedes CHEN 07B.				
$\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{42}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.72±0.11 OUR FIT				
1.20±0.33±0.13	126	1 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
1 We have multiplied $\bar{K}K\pi$ by 2/3 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$				
$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{51}\Gamma_{93}/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.93±0.11 OUR FIT				
1.10±0.21±0.15	126 ± 24	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$
$\Gamma(\eta_c(1S)\pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_{88}\Gamma_{93}/\Gamma$			
VALUE (eV)	CL\%	DOCUMENT ID	TECN	COMMENT
<15.7	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

$\chi_{c2}(1P)$ BRANCHING RATIOS

— HADRONIC DECAYS —

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	DOCUMENT ID
0.0102±0.0009 OUR FIT	
0.36±0.15 OUR FIT	
0.31±0.17	TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c2}$
$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-))$	Γ_{26}/Γ_1
VALUE	DOCUMENT ID

NODE=M057G4
NODE=M057G4NODE=M057G4;LINKAGE=UE
NODE=M057G;LINKAGE=NANODE=M057G07
NODE=M057G07NODE=M057G13
NODE=M057G13NODE=M057G13;LINKAGE=UE
NODE=M057G5
NODE=M057G5NODE=M057G6
NODE=M057G6NODE=M057G6;LINKAGE=UE
NODE=M057G01
NODE=M057G01

NODE=M057G01;LINKAGE=DE

NODE=M057G11
NODE=M057G11NODE=M057G05
NODE=M057G05

NODE=M057225

NODE=M057305

NODE=M057R2
NODE=M057R2NODE=M057R38
NODE=M057R38

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.83±0.23±0.04	903.5	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

¹ HE 08B reports $1.87 \pm 0.07 \pm 0.22 \pm 0.13$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.19±0.34±0.05	1031.9	1,2 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

¹ HE 08B reports $2.23 \pm 0.11 \pm 0.32 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.11±0.15±0.02	1164	1 ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$	

¹ ABLIKIM 11A reports $(1.21 \pm 0.05 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.206±0.040±0.004	76.9	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

¹ HE 08B reports $0.21 \pm 0.03 \pm 0.03 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.38±0.19±0.03	211.6	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

¹ HE 08B reports $1.41 \pm 0.11 \pm 0.16 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\rho^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.41±0.12±0.01	62.9	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

¹ HE 08B reports $0.42 \pm 0.11 \pm 0.06 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^-\bar{K}^0\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^0K^-\pi^+ \rightarrow K^-\pi^+K^0\pi^0+c.c.)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.29±0.08±0.01	38.7	1 HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$	

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0K^-\pi^+ \rightarrow K^-\pi^+K^0\pi^0+c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R46
NODE=M057R46

NODE=M057R46;LINKAGE=HE

NODE=M057R47
NODE=M057R47

NODE=M057R47;LINKAGE=HE

NODE=M057R47;LINKAGE=OC

NODE=M057R58
NODE=M057R58

NODE=M057R58;LINKAGE=AB

NODE=M057R50
NODE=M057R50

NODE=M057R50;LINKAGE=HE

NODE=M057R51
NODE=M057R51

NODE=M057R51;LINKAGE=HE

NODE=M057R57
NODE=M057R57

NODE=M057R57;LINKAGE=HE

$\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
0.38±0.09±0.01	63.0	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	

¹ HE 08B reports $0.39 \pm 0.07 \pm 0.05 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
0.37±0.08±0.01	51.1	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	

¹ HE 08B reports $0.38 \pm 0.07 \pm 0.04 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
0.29±0.08±0.01	39.3	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
0.127±0.044±0.003	22.9	1 HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	

¹ HE 08B reports $0.13 \pm 0.04 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	Γ_{14}/Γ
8.4±0.9 OUR FIT		

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
11.69±0.13±1.31	11k	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$	

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
7.30±0.11±0.75	4.5k	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$	

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(K^+ K^- \pi^+ \pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ_{14}
0.25±0.13 OUR FIT				
0.25±0.13	TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c2}$	

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	Γ_{17}/Γ
21±11 OUR FIT		

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	Γ_{18}/Γ
2.3±0.4 OUR FIT		

NODE=M057R52
NODE=M057R52

NODE=M057R52;LINKAGE=HE

NODE=M057R53
NODE=M057R53

NODE=M057R53;LINKAGE=HE

NODE=M057R54
NODE=M057R54

NODE=M057R54;LINKAGE=HE

NODE=M057R55
NODE=M057R55

NODE=M057R55;LINKAGE=HE

NODE=M057R3
NODE=M057R3

NODE=M057R00
NODE=M057R00

NODE=M057R00;LINKAGE=A

NODE=M057R73
NODE=M057R73

NODE=M057R73;LINKAGE=A

NODE=M057R39
NODE=M057R39

NODE=M057R9
NODE=M057R9

NODE=M057R26
NODE=M057R26

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**8.6±1.8 OUR EVALUATION****8.6±1.8 OUR AVERAGE** $8.6 \pm 0.9 \pm 1.6$ $8.7 \pm 5.9 \pm 0.4$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

DOCUMENT ID

TECN

COMMENT

 Γ_{19}/Γ

NODE=M057R4

NODE=M057R4

→ UNCHECKED ←

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**1.06±0.09 OUR FIT** $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**5.3±0.5±0.4**

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{20}/Γ

NODE=M057R20

NODE=M057R20

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**0.84±0.10 OUR AVERAGE** $0.82 \pm 0.10 \pm 0.02$ $1.73 \pm 0.57 \pm 0.04$ $762 \quad 27.7 \pm 7.4$

¹ ABLIKIM 20B reports $(5.33 \pm 0.52 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$.

² ABLIKIM 20B reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**0.73±0.04±0.08**

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{23}/Γ

NODE=M057R74

NODE=M057R74

 $512 \quad 1 \text{ ABLIKIM} \quad 13\text{B} \text{ BES3} \quad e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ 1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$. $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ VALUE (units 10^{-6})**9.6±2.7±0.2**

CL% EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{24}/Γ

NODE=M057R74;LINKAGE=A

 $33 \quad 1 \text{ ABLIKIM} \quad 19\text{J} \text{ BES3} \quad \psi(2S) \rightarrow \gamma \text{ hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $<18 \quad 90 \quad 2,3 \text{ ABLIKIM} \quad 11\text{K} \text{ BES3} \quad \psi(2S) \rightarrow \gamma \text{ hadrons}$

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.91 \pm 0.23 \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 11K reports $< 2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

3 Superseded by ABLIKIM 19J.

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**2.23±0.09 OUR FIT**

DOCUMENT ID

 Γ_{25}/Γ

NODE=M057R27

NODE=M057R27

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**37±16 OUR FIT**

DOCUMENT ID

 Γ_{26}/Γ

NODE=M057R8

NODE=M057R8

$\Gamma(\pi^+\pi^-\pi^0(\text{non-resonant}))/\Gamma_{\text{total}}$	Γ_{27}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$2.01 \pm 0.42 \pm 0.04$	64 1 ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$
1 ABLIKIM 17AG reports $(2.1 \pm 0.4 \pm 0.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
$\Gamma(\rho(770)\pm\pi^\mp)/\Gamma_{\text{total}}$	Γ_{28}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.61 \pm 0.38 \pm 0.01$	15 1 ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$
1 ABLIKIM 17AG reports $(0.64 \pm 0.39 \pm 0.07) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho(770)\pm\pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$	Γ_{29}/Γ
<u>VALUE (units 10^{-3})</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.48 \pm 0.13 \pm 0.01$	1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<1.4 90 2 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
1 ATHAR 07 reports $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
2 ABLIKIM 06R reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.	
$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$	Γ_{30}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.50 \pm 0.18 \pm 0.01$	1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
1 ATHAR 07 reports $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	
$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	Γ_{31}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
$5.4 \pm 0.4 \text{ OUR FIT}$	
$\Gamma(K^+K^-)/\Gamma_{\text{total}}$	Γ_{32}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
$1.01 \pm 0.06 \text{ OUR FIT}$	
$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{33}/Γ
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>
$0.52 \pm 0.04 \text{ OUR FIT}$	
$\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$	Γ_{33}/Γ_{25}
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
$0.235 \pm 0.019 \text{ OUR FIT}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.27 $\pm 0.07 \pm 0.04$	1,2 CHEN 07B BELL $e^+e^- \rightarrow e^+e^-\chi_{c2}$
1 Using $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from the $\pi^+\pi^-$ measurement of NAKAZAWA 05 rescaled by 3/2 to convert to $\pi\pi$.	
2 Not independent from other measurements.	

$\Gamma(K_S^0 K_S^0)/\Gamma(K^+ K^-)$

VALUE

DOCUMENT ID

TECN COMMENT

0.52±0.05 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.70 \pm 0.21 \pm 0.12$ 1,2 CHEN 07B BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

1 Using $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ from NAKAZAWA 05.

2 Not independent from other measurements.

 Γ_{33}/Γ_{32}

NODE=M057R37

NODE=M057R37

 $\Gamma(K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{34}/Γ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

1.44±0.21±0.03

1 ABLIKIM

17AG BES3

 $\psi(2S) \rightarrow \gamma K \bar{K} \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.72 \pm 0.26 \pm 0.04$ 2 ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

$1.34 \pm 0.27 \pm 0.03$ 3 ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

1 ABLIKIM 17AG reports $(1.5 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 ABLIKIM 17AG reports $(1.8 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 ABLIKIM 17AG reports $(1.4 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{35}/Γ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

1.24±0.27±0.03

1 ABLIKIM

17AG BES3

 $\psi(2S) \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

1 ABLIKIM 17AG reports $(1.3 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}$ Γ_{36}/Γ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

14.8±1.2±0.3

1 ABLIKIM

17AG BES3

 $\psi(2S) \rightarrow \gamma K \bar{K} \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$17.4 \pm 1.6 \pm 0.4$ 2 ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

$13.0 \pm 1.5 \pm 0.3$ 3 ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^{\pm} \pi^{\mp}$

1 ABLIKIM 17AG reports $(15.5 \pm 0.6 \pm 1.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

2 ABLIKIM 17AG reports $(18.2 \pm 0.8 \pm 1.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 ABLIKIM 17AG reports $(13.6 \pm 0.8 \pm 1.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^{\pm} K^{\mp})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R37

NODE=M057R37

NODE=M057R37;LINKAGE=CH

NODE=M057R37;LINKAGE=NI

NODE=M057R86

NODE=M057R86

OCCUR=2

OCCUR=3

NODE=M057R86;LINKAGE=A

NODE=M057R86;LINKAGE=B

NODE=M057R86;LINKAGE=C

NODE=M057R87

NODE=M057R87

NODE=M057R87;LINKAGE=A

NODE=M057R88

NODE=M057R88

OCCUR=2

OCCUR=3

NODE=M057R88;LINKAGE=A

NODE=M057R88;LINKAGE=B

NODE=M057R88;LINKAGE=C

$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{37}/Γ	
<i>VALUE (units 10^{-4})</i>	<i>DOCUMENT ID</i>	<i>TECN</i> <i>COMMENT</i>
$12.4 \pm 1.7 \pm 0.3$	¹ ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
¹ ABLIKIM 17AG reports $(13.0 \pm 1.0 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M057R89 NODE=M057R89
$\Gamma(K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}$	Γ_{38}/Γ	
<i>VALUE (units 10^{-4})</i>	<i>DOCUMENT ID</i>	<i>TECN</i> <i>COMMENT</i>
$5.2 \pm 0.8 \pm 0.1$	¹ ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K \bar{K} \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
$5.1 \pm 1.0 \pm 0.1$	² ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
$5.6 \pm 1.8 \pm 0.1$	³ ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
¹ ABLIKIM 17AG reports $(5.4 \pm 0.5 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		OCCUR=2 OCCUR=3
² ABLIKIM 17AG reports $(5.3 \pm 0.5 \pm 0.9) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M057R90;LINKAGE=A
³ ABLIKIM 17AG reports $(5.9 \pm 1.1 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M057R90;LINKAGE=B
$\Gamma(K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{39}/Γ	
<i>VALUE (units 10^{-4})</i>	<i>DOCUMENT ID</i>	<i>TECN</i> <i>COMMENT</i>
$5.6 \pm 2.1 \pm 0.1$	¹ ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
¹ ABLIKIM 17AG reports $(5.9 \pm 1.6 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M057R91 NODE=M057R91
$\Gamma(a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}$	Γ_{40}/Γ	
<i>VALUE (units 10^{-4})</i>	<i>DOCUMENT ID</i>	<i>TECN</i> <i>COMMENT</i>
$12.9 \pm 3.4 \pm 0.3$	¹ ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
¹ ABLIKIM 17AG reports $(13.5 \pm 1.6 \pm 3.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M057R92 NODE=M057R92
$\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$	Γ_{41}/Γ	
<i>VALUE (units 10^{-4})</i>	<i>DOCUMENT ID</i>	<i>TECN</i> <i>COMMENT</i>
$17.6 \pm 6.1 \pm 0.4$	¹ ABLIKIM	17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
¹ ABLIKIM 17AG reports $(18.4 \pm 3.3 \pm 5.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } \text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		NODE=M057R93 NODE=M057R93

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) **$0.30 \pm 0.08 \pm 0.01$**

DOCUMENT ID

1 ATHAR

TECN

CLEO

 Γ_{43}/Γ $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3})**<0.32**

CL%

DOCUMENT ID

1 ATHAR

TECN

CLEO

 Γ_{44}/Γ $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $< 0.33 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

 $\Gamma(K^+ K^- \eta'(958))/\Gamma_{\text{total}}$ VALUE (units 10^{-4})**1.94 ± 0.34**

EVTS

DOCUMENT ID

1 ABLIKIM

TECN

BES3

 Γ_{45}/Γ $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

¹ Derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.72 \pm 0.34)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

 $\Gamma(\eta \eta')/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**2.17 $\pm 0.47 \pm 0.05$**

CL%

EVTS

DOCUMENT ID

1 ABLIKIM

TECN

BES3

 Γ_{46}/Γ $\psi(2S) \rightarrow \gamma \eta \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6

<23

90

3.3 \pm 8.0

2 ASNER

09

CLEO

 $\psi(2S) \rightarrow \gamma \eta \eta'$

< 23

90

3 ADAMS

07

CLEO

 $\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ ABLIKIM 17AI reports $(2.27 \pm 0.43 \pm 0.25) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $< 0.6 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

³ Superseded by ASNER 09. ADAMS 07 reports $< 2.3 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

 $\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**4.6 $\pm 0.6 \pm 0.1$**

CL%

EVTS

DOCUMENT ID

1 ABLIKIM

TECN

BES3

 Γ_{47}/Γ $\psi(2S) \rightarrow \gamma \eta' \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 10

< 30

90

12 \pm 7

2 ASNER

09

CLEO

 $\psi(2S) \rightarrow \gamma \eta' \eta'$

< 30

90

3 ADAMS

07

CLEO

 $\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ ABLIKIM 17AI reports $(4.76 \pm 0.56 \pm 0.38) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ASNER 09 reports $< 1.0 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

³ Superseded by ASNER 09. ADAMS 07 reports $< 3.1 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

NODE=M057R05

NODE=M057R05

NODE=M057R05;LINKAGE=AT

NODE=M057R09

NODE=M057R09

NODE=M057R78

NODE=M057R78

NODE=M057R03

NODE=M057R03

NODE=M057R03;LINKAGE=A

NODE=M057R03;LINKAGE=AS

NODE=M057R03;LINKAGE=AD

NODE=M057R04

NODE=M057R04

NODE=M057R04;LINKAGE=A

NODE=M057R04;LINKAGE=AS

NODE=M057R04;LINKAGE=AD

$\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.17 \pm 0.54 \pm 0.05$	57 ± 11	1 ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{48}/Γ

NODE=M057R31
NODE=M057R31

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4	90	2.3 ± 2.2	1 ABLIKIM	050 BES2	$e^+ e^- \rightarrow \chi_{c2} \gamma$

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] < 3.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

 Γ_{49}/Γ

NODE=M057R32
NODE=M057R32

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.13 \pm 0.18 \pm 0.02$	68	1 ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$

¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (10.8 \pm 1.5 \pm 0.8) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

 Γ_{50}/Γ

NODE=M057R94
NODE=M057R94

 $\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID
1.65 ± 0.20 OUR FIT	

 Γ_{51}/Γ

NODE=M057R18
NODE=M057R18

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.29 \pm 0.03$	52	1 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ ABLIKIM 06T reports $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{52}/Γ

NODE=M057R01
NODE=M057R01

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$4.83 \pm 0.32 \pm 0.66$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c2}$

 Γ_{53}/Γ

NODE=M057R79
NODE=M057R79

 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.74 \pm 0.16 \pm 0.44$	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c2}$

 Γ_{54}/Γ

NODE=M057R80
NODE=M057R80

 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.93 \pm 0.06 \pm 0.10$	408	1 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

 Γ_{55}/Γ

NODE=M057R75
NODE=M057R75

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID
0.733 ± 0.033 OUR FIT	

 Γ_{56}/Γ

NODE=M057R12
NODE=M057R12

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.47 ± 0.04 OUR AVERAGE	1 ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
0.47 $\pm 0.04 \pm 0.01$	2 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

 Γ_{57}/Γ

NODE=M057R06
NODE=M057R06

¹ ONYISI 10 reports $(4.83 \pm 0.25 \pm 0.35 \pm 0.31) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$	Γ_{58}/Γ			
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	

0.174±0.025 OUR AVERAGE

¹ ONYISI 10 reports $(1.76 \pm 0.23 \pm 0.14 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$	Γ_{59}/Γ			
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	

0.36±0.04±0.01

¹ ONYISI 10 reports $(3.68 \pm 0.35 \pm 0.26 \pm 0.24) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$	Γ_{60}/Γ			
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT

2.8±0.9±0.1

¹ ABLIKIM 11F reports $(3.04 \pm 0.85 \pm 0.43) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{61}/Γ			
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	

1.32±0.34 OUR EVALUATION Treating systematic error as correlated.

1.3 ±0.4 OUR AVERAGE Error includes scale factor of 1.3.

¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$
¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+ \pi^-$ to $K^0 K^+ \pi^-$ decay.

$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$	Γ_{62}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT

0.078±0.023±0.002

¹ HE 08B reports $0.08 \pm 0.02 \pm 0.01 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=ON

NODE=M057R06;LINKAGE=AT

NODE=M057R34
NODE=M057R34

NODE=M057R34;LINKAGE=ON

NODE=M057R34;LINKAGE=AT

NODE=M057R56
NODE=M057R56

NODE=M057R56;LINKAGE=ON

NODE=M057R62
NODE=M057R62

NODE=M057R62;LINKAGE=AB

NODE=M057R6
NODE=M057R6
→ UNCHECKED ←

NODE=M057R6;LINKAGE=X3

NODE=M057R49
NODE=M057R49

NODE=M057R49;LINKAGE=HE

$\Gamma(p\bar{p}K^+K^-(\text{non-resonant}))/\Gamma_{\text{total}}$					Γ_{63}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.91 \pm 0.32 \pm 0.04$	131 ± 12	¹ ABLIKIM	11F	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$
1 ABLIKIM 11F reports $(2.08 \pm 0.19 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}K^+K^-(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \approx (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M057R59
NODE=M057R59

NODE=M057R59;LINKAGE=AB

$\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$					Γ_{64}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<7.9	90	¹ ABLIKIM	06D	BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$
1 Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.					

NODE=M057R30
NODE=M057R30

NODE=M057R;LINKAGE=AB

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$					Γ_{65}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.5 ± 0.9 OUR AVERAGE					
8.4 $\pm 1.0 \pm 0.2$	3309	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
10.2 $\pm 3.4 \pm 0.2$		² ABLIKIM	06I	BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.80 \pm 0.02 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 ABLIKIM 06I reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M057R33
NODE=M057R33

NODE=M057R33;LINKAGE=AL

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$					Γ_{66}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
8.9 $\pm 0.8 \pm 0.2$	3732	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.85 \pm 0.02 \pm 0.07) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M057R70
NODE=M057R70

NODE=M057R70;LINKAGE=AL

$\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{67}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
21.7 $\pm 1.7 \pm 0.5$	2128	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.07 \pm 0.06 \pm 0.15) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M057R71
NODE=M057R71

NODE=M057R71;LINKAGE=AL

$\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$					Γ_{68}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
21.1 $\pm 1.8 \pm 0.4$	2352	¹ ABLIKIM	12J	BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$
1 ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.01 \pm 0.06 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M057R72
NODE=M057R72

NODE=M057R72;LINKAGE=AL

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{69}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>			
1.84 ± 0.15 OUR FIT					

NODE=M057R25
NODE=M057R25

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{70}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
125 $\pm 15 \pm 3$	371	¹ ABLIKIM	12I	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<350	90	² ABLIKIM	06D	BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$

NODE=M057R29
NODE=M057R29

¹ ABLIKIM 12I reports $(137.0 \pm 7.6 \pm 15.7) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
66±15±1	36	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

¹ ABLIKIM 12I reports $(71.8 \pm 14.5 \pm 8.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<40	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

¹ ABLIKIM 12I reports $< 42 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

$\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<60	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

¹ ABLIKIM 12I reports $< 61 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

$\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.8±0.5 OUR AVERAGE				

7.7±0.5±0.2	5k	1,2 ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
8.3±1.6±0.2		3 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ABLIKIM 13D reports $(8.4 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$.

³ ATHAR 07 reports $(8.5 \pm 1.4 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.2±1.1±0.2	476	1 ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+}\bar{p}\Lambda$

¹ ABLIKIM 19AU reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (7.8 \pm 0.9 \pm 0.6) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8±0.7±0.1	79 ± 13	1 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M057R29;LINKAGE=AL

NODE=M057R29;LINKAGE=AB

NODE=M057R65
NODE=M057R65

NODE=M057R65;LINKAGE=AL

NODE=M057R66
NODE=M057R66

NODE=M057R66;LINKAGE=AL

NODE=M057R67
NODE=M057R67

NODE=M057R67;LINKAGE=AL

NODE=M057R07
NODE=M057R07

NODE=M057R07;LINKAGE=AB

NODE=M057R07;LINKAGE=AT

NODE=M057R98
NODE=M057R98

NODE=M057R98;LINKAGE=F

NODE=M057R60
NODE=M057R60

¹ ABLIKIM 11F reports $(3.06 \pm 0.50 \pm 0.54) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.6±1.4±0.1	29 ± 7	1 ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$
¹ ABLIKIM 11F reports $(5.05 \pm 1.29 \pm 0.93) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.7±0.6±0.1	91	1 ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6	90	2 ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$	
<7	90	7.5 ± 3.4	3 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
¹ ABLIKIM 18V reports $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ = $(0.35 \pm 0.05 \pm 0.02) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² ABLIKIM 13H reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.					
³ NAIK 08 reports $< 0.75 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.					

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.4±0.7±0.1	55	1 ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<8	90	2 ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$	
<7	90	4.0 ± 3.5	3 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
¹ ABLIKIM 18V reports $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ = $(0.32 \pm 0.06 \pm 0.03) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² ABLIKIM 13H reports $< 0.88 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.					
³ NAIK 08 reports $< 0.67 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.					

$\Gamma(\Sigma(1385)^+\bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
¹ ABLIKIM 12I reports $< 17 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.				

$\Gamma(\Sigma(1385)^-\bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	1 ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Sigma}(1385)^+$
¹ ABLIKIM 12I reports $< 8.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$.				

NODE=M057R60;LINKAGE=AB

NODE=M057R61
NODE=M057R61

NODE=M057R61;LINKAGE=AB

NODE=M057R43
NODE=M057R43

NODE=M057R43;LINKAGE=A

NODE=M057R43;LINKAGE=AB

NODE=M057R44
NODE=M057R44

NODE=M057R44;LINKAGE=A

NODE=M057R44;LINKAGE=AB

NODE=M057R44;LINKAGE=NA

NODE=M057R68
NODE=M057R68

NODE=M057R68;LINKAGE=AL

NODE=M057R69
NODE=M057R69

NODE=M057R69;LINKAGE=AL

$\Gamma(K^-\Lambda\Xi^++c.c.)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.76 \pm 0.32 \pm 0.04$	51	1 ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^-\Lambda\Xi^++c.c.$
1 ABLIKIM 15I reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^-\Lambda\Xi^++c.c.)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (1.68 \pm 0.26 \pm 0.15) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{83}/Γ NODE=M057R81
NODE=M057R81 $\Gamma(\Xi^0\Xi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.0	90	2.9 ± 1.7	1 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Xi^0\Xi^0$
1 NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0\Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.					

 Γ_{84}/Γ NODE=M057R45
NODE=M057R45 $\Gamma(\Xi^-\Xi^+)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.42 \pm 0.31 \pm 0.03$	29 ± 5	1 NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma\Xi^+\Xi^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.7	90	2 ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$
1 NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^-\Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
2 Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.				

 Γ_{85}/Γ NODE=M057R17
NODE=M057R17 $\Gamma(J/\psi(1S)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.015	90	BARATE	81 SPEC	$190 \text{ GeV } \pi^- \text{ Be} \rightarrow 2\pi 2\mu$

 Γ_{86}/Γ NODE=M057R13
NODE=M057R13 $\Gamma(\pi^0\eta_c)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<3.2 \times 10^{-3}$	90	1 ABLIKIM	15N BES3	$\psi(2S) e^+e^- \rightarrow \gamma\pi^0\eta_c$
1 Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma\gamma) = (1.66 \pm 0.11) \times 10^{-2}$.				

 Γ_{87}/Γ NODE=M057R77
NODE=M057R77 $\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$<0.54 \times 10^{-2}$	90	1,2 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

 Γ_{88}/Γ NODE=M057R76
NODE=M057R76

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.2×10^{-2}	90	1,3 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
1 Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.				
2 From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.				
3 From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.				

OCCUR=2

NODE=M057R76;LINKAGE=A

NODE=M057R76;LINKAGE=B

NODE=M057R76;LINKAGE=C

 $\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma(\bar{K}^0 K^+ \pi^- + c.c.)$ Γ_{88}/Γ_{42}

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<16.4	90	1 LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

1 We divided the reported limit by 2 to take into account the $K_L^0 K^+ \pi^-$ mode.**RADIATIVE DECAYS** $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{89}/Γ NODE=M057R7
NODE=M057R7**19.0 ± 0.5 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.64 $\pm 0.08 \pm 1.69$	1.0M	1 ABLIKIM	17U BES3	$e^+e^- \rightarrow \gamma X$
19.9 $\pm 0.5 \pm 1.2$		2 ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

1 Not independent from $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))$ and the product $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))$ also measured in ABLIKIM 17U.2 Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi)$ from ADAM 05A and $B(\psi(2S) \rightarrow \gamma\chi_{c2})$ from ATHAR 04.

NODE=M057R7;LINKAGE=A

NODE=M057R7;LINKAGE=AD

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$	Γ_{90}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<19	90	13 ± 11	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<40	90	17.2 ± 6.8	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$

¹ ABLIKIM 11E reports $< 20.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

² BENNETT 08A reports $< 50 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$	Γ_{91}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<6	90	1 ± 6	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<6	90	0.0 ± 1.8	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

¹ ABLIKIM 11E reports $< 6.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

² BENNETT 08A reports $< 7.0 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$	Γ_{92}/Γ				
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 7	90	5 ± 5	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
<11	90	1.3 ± 2.5	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$

¹ ABLIKIM 11E reports $< 8.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

² BENNETT 08A reports $< 13 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{93}/Γ
<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>
2.85 \pm 0.10 OUR FIT	

$\Gamma(e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}$	Γ_{94}/Γ			
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
2.37 \pm 0.15 \pm 0.05	1.3k	^{1,2} ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- J/\psi$
¹ ABLIKIM 17I reports $(2.48 \pm 0.08 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
2 Not independent from other measurements reported by ABLIKIM 17I				

$\Gamma(e^+ e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$	Γ_{94}/Γ_{89}			
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.3 \pm 0.4 \pm 0.5	1.3k	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$
¹ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.				

$\Gamma(\mu^+ \mu^- J/\psi(1S))/\Gamma(e^+ e^- J/\psi(1S))$	Γ_{95}/Γ_{94}			
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.40 \pm 0.79 \pm 1.15	219	ABLIKIM	19z BES3	$\psi(2S) \rightarrow \gamma\chi_c \rightarrow \gamma(\mu^+ \mu^- J/\psi)$

NODE=M057R40
NODE=M057R40

NODE=M057R40;LINKAGE=AB
NODE=M057R40;LINKAGE=BE

NODE=M057R41
NODE=M057R41

NODE=M057R41;LINKAGE=AB
NODE=M057R41;LINKAGE=BE

NODE=M057R42
NODE=M057R42

NODE=M057R42;LINKAGE=AB
NODE=M057R42;LINKAGE=BE

NODE=M057R82;LINKAGE=C

NODE=M057R83
NODE=M057R83

NODE=M057R83;LINKAGE=A

NODE=M057R96
NODE=M057R96

$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$	Γ_{93}/Γ_{89}		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.50±0.05 OUR FIT			
0.99±0.18	¹ AMBROGIANI 00B	E835	$\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$
¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.			
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$	$\Gamma_{93}/\Gamma \times \Gamma_{56}/\Gamma$		
VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
2.09±0.13 OUR FIT			
1.7 ±0.4 OUR AVERAGE	ARMSTRONG 93	E760	$\bar{p}p \rightarrow \gamma\gamma X$
1.60±0.42	BAGLIN 87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma X$
9.9 ±4.5			

$\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$	$\Gamma_{14}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.31±0.26 OUR FIT			

2.5 ±0.9 OUR AVERAGE Error includes scale factor of 2.3.

1.90±0.14±0.44	BAI	99B	BES	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3.8 ±0.67	¹ TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c2}$
¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) \times B(J/\psi \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.				

$\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$	$\Gamma_{18}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma^{\psi(2S)}$		
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.1 ±0.4 OUR FIT			
3.11±0.36±0.48	ABLIKIM	04H	BES2

$\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$	$\Gamma_{56}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}$		
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.01±0.09 OUR FIT			
1.4 ±1.1	¹ BAI	98I	BES

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$	$\Gamma_{56}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma^{\psi(2S)}$			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
6.98±0.32 OUR FIT				

7.1 ±0.5 OUR AVERAGE Error includes scale factor of 1.2.

7.3 ±0.4 ±0.3	405	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
7.2 ±0.7 ±0.4	121 ± 12	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
4.4 ±1.6 ±0.6	14.3 ± 5.2	BAI	04F	BES	$\psi(2S) \rightarrow \gamma\chi_{c2}(1P) \rightarrow \gamma p\bar{p}$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow p\bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$	$\Gamma_{69}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma^{\psi(2S)}$			
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
17.5±1.3 OUR FIT				
17.4±1.4 OUR AVERAGE				

18.2±1.4±0.9	207	¹ ABLIKIM	13H	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
15.9±2.1±1.0	71 ± 9	² NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ Calculated by us. ABLIKIM 13H reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (20.8 \pm 1.6 \pm 2.3) \times 10^{-5}$ from a measurement of $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c2})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.74 \pm 0.35)\%$.

² Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

NODE=M057R23
NODE=M057R23

NODE=M057R;LINKAGE=7A

NODE=M057R24
NODE=M057R24

NODE=M057230

NODE=M057B18
NODE=M057B18

NODE=M057B18;LINKAGE=TA

NODE=M057B19
NODE=M057B19

NODE=M057B1
NODE=M057B1

NODE=M057B;LINKAGE=J8

NODE=M057B6
NODE=M057B6

NODE=M057B10
NODE=M057B10

NODE=M057B10;LINKAGE=AB

NODE=M057B10;LINKAGE=NA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{69}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

5.1±0.4 OUR FIT

7.1^{+3.1}_{-2.9}±1.3 8.3^{+3.7}_{-3.4} 1 BAI 03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ BAI 03E reports [$B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c2}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$] \times $[B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.59}_{-0.55} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{25}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

2.12±0.08 OUR FIT

2.17±0.09 OUR AVERAGE

2.19±0.05±0.15	4.5k	¹ ABLIKIM	10A	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
2.23±0.06±0.10	2.5k	² ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1.90±0.08±0.20	0.8k	³ ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.88 \pm 0.02 \pm 0.06 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^+\pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

³ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{25}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

0.612±0.023 OUR FIT

0.54 ± 0.06 OUR AVERAGE

0.66 ± 0.18 ± 0.37	21 ± 6	¹ BAI	03C	BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$
0.54 ± 0.05 ± 0.04	185 ± 16	² BAI	98I	BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

¹ We have multiplied $\pi^0\pi^0$ measurement by 3 to obtain $\pi\pi$.

² Calculated by us. The value for $B(\chi_{c2} \rightarrow \pi^+\pi^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{31}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) CL% EVTS DOCUMENT ID TECN COMMENT

0.52±0.04 OUR FIT

0.52±0.04 OUR AVERAGE

0.54±0.03±0.04	386	¹ ABLIKIM	10A	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
0.47±0.05±0.05	156	ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.44	90	² ADAMS	07	CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2}$
< 3	90	BAI	03C	BES	$\psi(2S) \rightarrow \gamma\eta\eta \rightarrow 5\gamma$
0.62±0.31±0.19		LEE	85	CBAL	$\psi(2S) \rightarrow \text{photons}$

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \eta\eta) = (0.65 \pm 0.04 \pm 0.05 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$.

² Superseded by ASNER 09.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{32}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

9.6±0.6 OUR FIT

10.5±0.3±0.6 1.6k ¹ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K^+K^-$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K^+K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

NODE=M057B11

NODE=M057B11

NODE=M057B11;LINKAGE=BA

NODE=M057B02

NODE=M057B02

OCCUR=2

NODE=M057B02;LINKAGE=AB

NODE=M057B02;LINKAGE=AS

NODE=M057B02;LINKAGE=AN

NODE=M057B9

NODE=M057B9

NODE=M057B;LINKAGE=BM

NODE=M057B;LINKAGE=BA

NODE=M057B04

NODE=M057B04

NODE=M057B04;LINKAGE=AB

NODE=M057B04;LINKAGE=AD

NODE=M057B03

NODE=M057B03

NODE=M057B03;LINKAGE=AS

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{32}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

0.276±0.017 OUR FIT

0.190±0.034±0.019 115 ± 13 1 BAI 98I BES $\psi(2S) \rightarrow \gamma K^+ K^-$

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{33}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

5.0 ±0.4 OUR FIT

5.0 ±0.4 OUR AVERAGE

4.9 ±0.3 ±0.3 373 ± 20 1 ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
5.72±0.76±0.63 65 ABLIKIM 050 BES2 $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{33}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

14.4±1.1 OUR FIT

14.7±4.1±3.3

1 BAI 99B BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{42}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

1.22±0.17 OUR FIT

1.15±0.18 OUR AVERAGE

1.21±0.19±0.09 37 1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
0.97±0.32±0.13 28 2 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Calculated by us. ATHAR 07 reports $B(\chi_{c2} \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (1.3 \pm 0.2 \pm 0.1 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp) = (0.6 \pm 0.2 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.1 \pm 0.6)\%$. We have multiplied by 2 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$ from $K_S^0 K^\pm \pi^\mp$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_1/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

2.79±0.26 OUR FIT

3.1 ±1.0 OUR AVERAGE Error includes scale factor of 2.5.

2.3 ±0.1 ±0.5 1 BAI 99B BES $\psi(2S) \rightarrow \gamma \chi_{c2}$
4.3 ±0.6 2 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ \pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S)\ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{51}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

1.57±0.19 OUR FIT

1.76±0.16±0.24 160 1 ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

NODE=M057B8

NODE=M057B8

NODE=M057B;LINKAGE=BI

NODE=M057B12

NODE=M057B12

NODE=M057B12;LINKAGE=AS

NODE=M057B13

NODE=M057B13

NODE=M057B13;LINKAGE=BA

NODE=M057B05

NODE=M057B05

NODE=M057B05;LINKAGE=AT

NODE=M057B05;LINKAGE=AB

NODE=M057B5

NODE=M057B5

NODE=M057B;LINKAGE=K1

NODE=M057B;LINKAGE=K2

NODE=M057B14

NODE=M057B14

NODE=M057B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} / \frac{\Gamma_{51}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

4.5±0.5 OUR FIT

3.6±0.6±0.6

¹ BAI

99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{20}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

1.01±0.08 OUR FIT

0.98±0.13 OUR AVERAGE Error includes scale factor of 1.3.

0.94±0.03±0.10	849	¹ ABLIKIM	11K	BES3	$\psi(2S) \rightarrow \gamma$ hadrons
1.38±0.24±0.23	41	² ABLIKIM	06T	BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35)\%$.

² Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{20}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

2.92±0.24 OUR FIT

4.8 ±1.3 ±1.3

¹ BAI

99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{79}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT

7.85±0.77±0.44

^{129±13}

¹ ABLIKIM

19BB BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c. c.}$

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_{c2} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (8.25 \pm 0.83 \pm 0.49) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$ and other branching fractions from PDG 18.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{89}/\Gamma \times \Gamma_{155}^{\psi(2S)}/\Gamma_{11}^{\psi(2S)}}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

1.81 ±0.04 OUR FIT

1.69 ±0.16 OUR AVERAGE

Error includes scale factor of 3.4. See the ideogram below.

1.996±0.008±0.070	81k	¹ ABLIKIM	17N	BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
1.793±0.008±0.163	1.0M	ABLIKIM	17U	BES3	$e^+ e^- \rightarrow \gamma X$
1.62 ± 0.04 ± 0.12	5.8k	BAI	04I	BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
0.99 ± 0.10 ± 0.08		GAISER	86	CBAL	$\psi(2S) \rightarrow \gamma X$
1.47 ± 0.17		² OREGLIA	82	CBAL	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.8 ± 0.5		³ BRANDELIK	79B	DASP	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.2 ± 0.2		³ BARTEL	78B	CNTR	$\psi(2S) \rightarrow \gamma \chi_{c2}$
2.2 ± 1.2		⁴ BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$
1.2 ± 0.7		² WHITAKER	76	MRK1	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.874±0.007±0.102	76k	⁵ ABLIKIM	120	BES3	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.95 ± 0.02 ± 0.07	12.4k	⁶ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.85 ± 0.04 ± 0.07	1.9k	⁷ ADAM	05A	CLEO	Repl. by MENDEZ 08

NODE=M057B15

NODE=M057B15

NODE=M057B15;LINKAGE=BA

NODE=M057B16

NODE=M057B16

NODE=M057B16;LINKAGE=AL

NODE=M057B16;LINKAGE=AB

NODE=M057B17

NODE=M057B17

NODE=M057B17;LINKAGE=BA

NODE=M057B07

NODE=M057B07


NODE=M057B07;LINKAGE=A

NODE=M057B2

NODE=M057B2

- 1 Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.
 2 Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.
 3 Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.
 4 Assumes isotropic gamma distribution.
 5 Superseded by ABLIKIM 17N.
 6 Not independent from other measurements of MENDEZ 08.
 7 Not independent from other values reported by ADAM 05A.

NODE=M057B2;LINKAGE=A
 NODE=M057B;LINKAGE=3Q
 NODE=M057B;LINKAGE=2Q
 NODE=M057B;LINKAGE=EA
 NODE=M057B2;LINKAGE=B
 NODE=M057B2;LINKAGE=ME
 NODE=M057B;LINKAGE=AD



$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}} \text{ (units } 10^{-2})$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{89} / \Gamma \times \Gamma_{155}^{\psi(2S)} / \Gamma_9^{\psi(2S)} = \Gamma_{89} / \Gamma \times \Gamma_{155}^{\psi(2S)} / (\Gamma_{11}^{\psi(2S)} + \Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + 0.343 \Gamma_{154}^{\psi(2S)} + 0.190 \Gamma_{155}^{\psi(2S)})$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.95 ± 0.06 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.12 \pm 0.03 \pm 0.09$	12.4k	¹ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
$3.11 \pm 0.07 \pm 0.07$	1.9k	ADAM	05A CLEO	Repl. by MENDEZ 08

¹ Not independent from other measurements of MENDEZ 08.

NODE=M057B7
 NODE=M057B7

NODE=M057B7

NODE=M057B7;LINKAGE=ME

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.22 ± 0.11 OUR FIT				

NODE=M057B3
 NODE=M057B3

5.53 ± 0.17 OUR AVERAGE

$5.56 \pm 0.05 \pm 0.16$	12.4k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
6.0 ± 2.8	1.3k	¹ ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
3.9 ± 1.2		² HIMEL	80 MRK2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.52 \pm 0.13 \pm 0.13$	1.9k	³ ADAM	05A CLEO	Repl. by MENDEZ 08
--------------------------	------	-------------------	----------	--------------------

NODE=M057B;LINKAGE=AB
 NODE=M057B;LINKAGE=H8

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

NODE=M057B3;LINKAGE=AD

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}} = \frac{\Gamma_{93}/\Gamma \times \Gamma_{155}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.71±0.08 OUR FIT**2.82±0.10 OUR AVERAGE**

2.83±0.08±0.06	5k	¹ ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
2.68±0.28±0.15	0.3k	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
7.0 ± 2.1 ± 2.0		LEE	85 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.81±0.17±0.15	1.1k	² ABLIKIM	12A BES3	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
----------------	------	----------------------	----------	--

¹ ABLIKIM 17AE measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0} / \Gamma_{\gamma\gamma}^{\lambda=2} = 0.000 \pm 0.006 \pm 0.012$.

² ABLIKIM 12A measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0} / \Gamma_{\gamma\gamma}^{\lambda=2} = 0.00 \pm 0.02 \pm 0.02$. Superseded by ABLIKIM 17AE.

NODE=M057B4

NODE=M057B4

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)/\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma) = \frac{\Gamma_{93}/\Gamma_{93}^{\chi_{c0}(1P)}}{\Gamma_{93}^{\chi_{c0}(1P)}}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.292±0.028 OUR AVERAGE

0.295±0.014±0.028	8k	¹ ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
0.278±0.050±0.036	0.5k	¹ ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.271±0.029±0.030	1.9k	^{1,2} ABLIKIM	12A BES3	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
-------------------	------	------------------------	----------	--

¹ Not independent from the values of $\Gamma(\chi_{c0}, \chi_{c2})$ and $B(\psi(2S) \rightarrow \chi_{c0}, \chi_{c2})$.² Superseded by ABLIKIM 17AE.

NODE=M057B06

NODE=M057B06

MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

$$a_2 = M^2/\sqrt{E1^2 + M^2 + E3^2} \text{ Magnetic quadrupole fractional transition amplitude}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

-11.0± 1.0 OUR AVERAGE

-12.0± 1.3±0.4	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 9.3± 1.6±0.3	19.8k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 9.3 ^{+ 3.9} _{- 4.1} ±0.6	5.9k	³ AMBROGIANI	02 E835	$p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
-14 ± 6	1.9k	³ ARMSTRONG	93E E760	$p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
-33.3 ^{+11.6} _{-29.2}	441	³ OREGLIA	82 CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 7.9± 1.9±0.3	19.8k	⁴ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
----------------	-------	---------------------	---------	---

¹ Correlated with a_3 , b_2 , and b_3 with correlation coefficients $\rho_{a_2 a_3} = 0.733$, $\rho_{a_2 b_2} = -0.605$, and $\rho_{a_2 b_3} = -0.095$.

² From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.

³ Assuming $a_3=0$.

⁴ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057A1

NODE=M057A1

OCCUR=2

NODE=M057A1;LINKAGE=B

NODE=M057A1;LINKAGE=AR

NODE=M057A1;LINKAGE=A

NODE=M057A1;LINKAGE=AT

$$a_3 = E3/\sqrt{E1^2 + M^2 + E3^2} \text{ Electric octupole fractional transition amplitude}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

-0.3±1.0 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

-1.3±0.9±0.4	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.7±1.4±0.3	19.8k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.0 ^{+5.5} _{-4.4} ±0.9	5908	AMBROGIANI	02 E835	$p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
0 ⁺⁶ ₋₅	1904	ARMSTRONG	93E E760	$p\bar{p} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$

¹ Correlated with a_2 , b_2 , and b_3 with correlation coefficients $\rho_{a_2 a_3} = 0.733$, $\rho_{a_3 b_2} = -0.422$, and $\rho_{a_3 b_3} = -0.024$.


² From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057A2

NODE=M057A2

NODE=M057A2;LINKAGE=A

NODE=M057A2;LINKAGE=AR



$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude (units 10^{-2})

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ RADIATIVE DECAY

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.9 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
1.7±0.8±0.2	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
4.6±1.0±1.3	13.8k	² ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.2±1.5±0.4	19.8k	³ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 5.1 ^{+5.4} _{-3.6}	721	² ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
13.2 ^{+9.8} _{-7.5}	441	⁴ OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.0±1.3±0.3 19.8k ⁴ ARTUSO 09 CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Correlated with a_2 , a_3 , and b_3 with correlation coefficients $\rho_{a_2} b_2 = -0.605$, $\rho_{a_3} b_2 = -0.422$, and $\rho_{b_2} b_3 = 0.384$.

² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

³ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

⁴ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.

NODE=M057250

NODE=M057QB2
NODE=M057QB2


OCCUR=2

NODE=M057QB2;LINKAGE=A

NODE=M057QB2;LINKAGE=AB

NODE=M057QB2;LINKAGE=AT

NODE=M057QB2;LINKAGE=AR



$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude (units 10^{-2})

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-1.0±0.6 OUR AVERAGE				
-1.4±0.7±0.4	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.5±0.8±1.8	13.8k	² ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
-0.8±1.2±0.2	19.8k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-2.7 ^{+4.3} _{-2.9}	721	² ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$

¹ Correlated with a_2 , a_3 , and b_2 with correlation coefficients $\rho_{a_2} b_3 = -0.095$, $\rho_{a_3} b_3 = -0.024$, and $\rho_{b_2} b_3 = 0.384$.

² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

NODE=M057QB3
NODE=M057QB3

NODE=M057QB3;LINKAGE=A

NODE=M057QB3;LINKAGE=AB

NODE=M057260

NODE=M057QAR
NODE=M057QAR

NODE=M057QAR;LINKAGE=AR

NODE=M057

REFID=60212
REFID=59844
REFID=59996;ERROR=15
REFID=60026
REFID=59606
REFID=59837
REFID=58990
REFID=59304
REFID=58191
REFID=58278
REFID=58310
REFID=58316
REFID=58321
REFID=57931
REFID=57978
REFID=58026
REFID=57140
REFID=56774
REFID=56778
REFID=56779
REFID=55901
REFID=54877
REFID=54879

$\chi_{c2}(1P)$ REFERENCES

ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AU	PR D100 052010	Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BI	PRL 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AG	PR D96 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17AI	PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17I	PR 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)

ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11I	PR D84 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53930
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
DOBBS	06	PR D73 071101	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51062
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50189
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE	02T	PL B540 33	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48813
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48552
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
ACCIARRI	99E	PL B453 73	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46943
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ACKER...K...	98	PL B439 197	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46324
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
DOMINICK	94	PR D50 4265	J. Dominick <i>et al.</i>	(CLEO Collab.)	REFID=44077
ARMSTRONG	93	PRL 70 2988	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43306
ARMSTRONG	93E	PR D48 3037	T.A. Armstrong <i>et al.</i>	(FNAL-E760 Collab.)	REFID=48616
BAUER	93	PL B302 345	D.A. Bauer <i>et al.</i>	(TPC Collab.)	REFID=43315
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41865
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41907
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)	REFID=22145
GAISER	86	PR D34 711	J. Gaisser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
Also		Private Comm.	M.J. Oreglia	(EFI)	REFID=22143
BARATE	81	PR D24 2994	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, CERN+)	REFID=22164
HIMEL	80	PRL 44 920	T. Himmel <i>et al.</i>	(LBL, SLAC)	REFID=22119
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151

NODE=M059

 $\eta_c(2S)$ $I^G(J^{PC}) = 0^+(0^-+)$

Quantum numbers are quark model predictions.

 $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3637.5±1.1 OUR AVERAGE				Error includes scale factor of 1.2.
3635.1±3.7±2.9	106	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
3633.6±1.7±0.6	106	¹ AAIJ	17ADLHCb	$pp \rightarrow B^+X \rightarrow p\bar{p}K^+X$
3636.4±4.1±0.7	365	² AAIJ	17BBLHCb	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3637.0±5.7±3.4	178	^{3,4} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
3635.1±5.8±2.1	47	^{3,5} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
3646.9±1.6±3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6±2.9±1.6	127 ± 18	⁶ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, KK\pi^0$
3638.5±1.5±0.8	624	³ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5±3.2±2.5	1201	³ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
3636.1 ^{+3.9} _{-4.2} ^{+0.7} _{-2.0}	128	⁷ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	⁸ ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
3645.0±5.5 ^{+4.9} _{-7.8}	121 ± 27	AUBERT	05C BABR	$e^+e^- \rightarrow J/\psi c\bar{c}$
3642.9±3.1±1.5	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3639 ± 7	98 ± 52	⁹ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
3630.8±3.4±1.0	112 ± 24	¹⁰ AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
3654 ± 6 ± 8	39 ± 11	¹¹ CHOI	02 BELL	$B \rightarrow KK_S K^-\pi^+$
3594 ± 5		¹² EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

¹AAIJ 17AD report $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$ MeV. We use the current value $m_{\psi(2S)} = 3686.097 \pm 0.025$ MeV to obtain the quoted mass.

²From a fit of the $\phi\phi$ invariant mass with the width of $\eta_c(2S)$ fixed to the PDG 16 value.

³Ignoring possible interference with continuum.

⁴With a width fixed to 11.3 MeV.

⁵With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.

⁶From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+K^-\pi^0$ decay modes.

⁷Accounts for interference with non-resonant continuum.

⁸From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁹From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁰Superseded by DEL-AMO-SANCHEZ 11M.

¹¹Superseded by VINOKUROVA 11.

¹²Assuming mass of $\psi(2S) = 3686$ MeV.

NODE=M059

NODE=M059M

NODE=M059M

OCCUR=2

OCCUR=2

NODE=M059M;LINKAGE=B

NODE=M059M;LINKAGE=C

NODE=M059M;LINKAGE=DE

NODE=M059M;LINKAGE=LE

NODE=M059M;LINKAGE=LS

NODE=M059M;LINKAGE=AB

NODE=M059M;LINKAGE=VA

NODE=M059M;LINKAGE=EB

NODE=M059M;LINKAGE=AU

NODE=M059M;LINKAGE=AR

NODE=M059M;LINKAGE=CH

NODE=M059M;LINKAGE=A

NODE=M059W

NODE=M059W

 $\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.3^{+ 3.2}_{- 2.9} OUR AVERAGE					
9.9 ± 4.8 ± 2.9	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	
16.9 ± 6.4 ± 4.8	127 ± 18	¹³ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, KK\pi^0$	
13.4 ± 4.6 ± 3.2	624	¹⁴ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	
6.6 ^{+ 8.4} _{- 5.1} ^{+ 2.6} _{- 0.9}	128	¹⁵ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm \pi^\mp)$	
6.3 ± 12.4 ± 4.0	61	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 23	90	98 ± 52	¹⁶ AUBERT	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$	
22 ± 14		121 ± 27	AUBERT	$e^+e^- \rightarrow J/\psi c\bar{c}$	
17.0 ± 8.3 ± 2.5		112 ± 24	¹⁷ AUBERT	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$	
< 55	90	39 ± 11	¹⁸ CHOI	$B \rightarrow KK_S K^-\pi^+$	
< 8.0	95		¹⁹ EDWARDS	$e^+e^- \rightarrow \gamma X$	

- 13 From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.
 14 Ignoring possible interference with continuum.
 15 Accounts for interference with non-resonant continuum.
 16 From the fit of the kaon momentum spectrum. Systematic errors not evaluated.
 17 Superseded by DEL-AMO-SANCHEZ 11M.
 18 For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.
 19 For a mass value of 3594 ± 5 MeV

$\eta_c(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	not seen	
Γ_2 $K\bar{K}\pi$	(1.9 ± 1.2) %	
Γ_3 $K\bar{K}\eta$	(5 ± 4) $\times 10^{-3}$	
Γ_4 $2\pi^+ 2\pi^-$	not seen	
Γ_5 $\rho^0 \rho^0$	not seen	
Γ_6 $3\pi^+ 3\pi^-$	not seen	
Γ_7 $K^+ K^- \pi^+ \pi^-$	not seen	
Γ_8 $K^{*0} \bar{K}^{*0}$	not seen	
Γ_9 $K^+ K^- \pi^+ \pi^- \pi^0$	(1.4 ± 1.0) %	
Γ_{10} $K^+ K^- 2\pi^+ 2\pi^-$	not seen	
Γ_{11} $K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$	seen	
Γ_{12} $2K^+ 2K^-$	not seen	
Γ_{13} $\phi\phi$	not seen	
Γ_{14} $p\bar{p}$	seen	
Γ_{15} $p\bar{p}\pi^+ \pi^-$	seen	
Γ_{16} $\gamma\gamma$	(1.9 ± 1.3) $\times 10^{-4}$	
Γ_{17} $\gamma J/\psi(1S)$	< 1.4 %	90%
Γ_{18} $\pi^+ \pi^- \eta$	not seen	
Γ_{19} $\pi^+ \pi^- \eta'$	not seen	
Γ_{20} $\pi^+ \pi^- \eta_c(1S)$	< 25 %	90%

$\eta_c(2S)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	Γ_{16}
<u>VALUE (keV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.44 \pm 0.14	106 20 XU 18 BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
1.3 \pm 0.6	21 ASNER 04 CLEO $\gamma\gamma \rightarrow \eta_c \rightarrow K_S^0 K^\pm \pi^\mp$
20 Assuming that the branching fraction into $\eta' \pi^+ \pi^-$ is the same as for $\eta_c(1S)$.	
21 They measure $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K\pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.	
$\Gamma(\gamma\gamma) \times \Gamma(\pi^+ \pi^- \eta') / \Gamma_{\text{total}}$	$\Gamma_{16} \Gamma_{19} / \Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
5.6 \pm 1.2 \pm 1.1	106 XU 18 BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(2\pi^+ 2\pi^-) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$	$\Gamma_4 \Gamma_{16} / \Gamma$
<u>VALUE (eV)</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
<6.5	
41 \pm 4 \pm 6	90 UEHARA 08 BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+ \pi^-)$
$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$	$\Gamma_2 \Gamma_{16} / \Gamma$
<u>VALUE (eV)</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
41 \pm 4 \pm 6	624 22 DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

22 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059W;LINKAGE=AB
 NODE=M059W;LINKAGE=DE
 NODE=M059W;LINKAGE=VA
 NODE=M059W;LINKAGE=AU
 NODE=M059W;LINKAGE=AR
 NODE=M059W;LINKAGE=W2
 NODE=M059W;LINKAGE=W
 NODE=M059215;NODE=M059
 DESIG=1
 DESIG=4
 DESIG=20
 DESIG=5
 DESIG=16
 DESIG=8;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=6
 DESIG=17
 DESIG=9
 DESIG=10;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=11
 DESIG=7
 DESIG=18
 DESIG=3
 DESIG=22
 DESIG=2
 DESIG=21
 DESIG=12;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=13;OUR EVAL; \rightarrow UNCHECKED \leftarrow
 DESIG=15

NODE=M059216
 NODE=M059W1
 NODE=M059W1
 NODE=M059W1;LINKAGE=A
 NODE=M059W1;LINKAGE=AS

NODE=M059R29
 NODE=M059R29
 NODE=M059218
 NODE=M059G01
 NODE=M059G01
 NODE=M059G04
 NODE=M059G04
 NODE=M059G04;LINKAGE=DE

$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_{16}/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<5.0	90	UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$
$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_9\Gamma_{16}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
30±6±5	1201	23 DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	23 Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$
VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT	
< 5.6	90 ^{24,25,26}	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 8.0	90 ^{24,25,27}	AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
<12.0	90	25,27 AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$	
24 Including the measurements of ARMSTRONG 95F in the AMBROGIANI 01 analysis. 25 For a total width $\Gamma=5$ MeV. 26 For the resonance mass region 3589–3599 MeV/ c^2 . 27 For the resonance mass region 3575–3660 MeV/ c^2 .					

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		ABREU	980	DLPH	$e^+e^- \rightarrow e^+e^- + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	28	EDWARDS	82C	CBAL	$e^+e^- \rightarrow \gamma X$
28 For a mass value of 3594 ± 5 MeV					

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.9±0.4±1.1	59 ± 12	29 AUBERT	08AB	BABR	$B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	127 ± 18	ABLIKIM	13K	BES3	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$
seen	39 ± 11	30 CHOI	02	BELL	$B \rightarrow KK_S K^-\pi^+$
29 Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.					
30 For a mass value of 3654 ± 6 MeV					

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$					Γ_3/Γ_2
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
27.3±7.0±9.0	225	31 LEES	14E	BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
31 LEES 14E reports $B(\eta_c(2S) \rightarrow K^+K^-\eta)/B(\eta_c(2S) \rightarrow K^+K^-\pi^0) = 0.82 \pm 0.21 \pm 0.27$, which we divide by 3 to account for isospin symmetry.					

$\Gamma(2\pi^+2\pi^-)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
not seen		UEHARA	08	BELL	$\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059G02
NODE=M059G02NODE=M059G05
NODE=M059G05

NODE=M059G05;LINKAGE=DE

NODE=M059G03
NODE=M059G03NODE=M059G06
NODE=M059G06

NODE=M059217

NODE=M059G1
NODE=M059G1OCCUR=2
OCCUR=3NODE=M059G1;LINKAGE=A
NODE=M059G1;LINKAGE=B
NODE=M059G1;LINKAGE=C1
NODE=M059G1;LINKAGE=C2

NODE=M059220

NODE=M059R1
NODE=M059R1

NODE=M059R;LINKAGE=W

NODE=M059R3
NODE=M059R3

NODE=M059R3;LINKAGE=AU

NODE=M059R;LINKAGE=W2

NODE=M059R26
NODE=M059R26

NODE=M059R26;LINKAGE=LE

NODE=M059R01
NODE=M059R01

$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$	Γ_5/Γ	NODE=M059R15 NODE=M059R15	
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
not seen	ABLIKIM 11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	
$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_7/Γ	NODE=M059R02 NODE=M059R02	
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
not seen	UEHARA 08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$	
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$	Γ_9/Γ_2	NODE=M059R21 NODE=M059R21	
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
0.73±0.17±0.17	1201 32 DEL-AMO-SA..11M	BABR $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
32 We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.			
$\Gamma(K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$	Γ_8/Γ	NODE=M059R16 NODE=M059R16	
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
not seen	ABLIKIM 11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$	
$\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{11}/Γ	NODE=M059R22 NODE=M059R22	
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
seen	57±17	ABLIKIM 13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
$\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$	Γ_{12}/Γ	NODE=M059R03 NODE=M059R03	
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
not seen	UEHARA 08 BELL	$\gamma\gamma \rightarrow \eta_c(2S)$	
$\Gamma(\phi\phi)/\Gamma_{\text{total}}$	Γ_{13}/Γ	NODE=M059R17 NODE=M059R17	
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
not seen	ABLIKIM 11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$	
$\Gamma(p\bar{p})/\Gamma_{\text{total}}$	Γ_{14}/Γ	NODE=M059R04 NODE=M059R04 OCCUR=2	
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
seen	106 33 AAIJ	17AD LHCb $p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$	
33 AAIJ 17AD report a 6.4 standard deviation signal, with $B(B^+ \rightarrow \eta_c(2S) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$.			
$\Gamma(p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_{15}/Γ	NODE=M059R30 NODE=M059R30	
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
seen	110 34 CHILIKIN	19 BELL $e^+ e^- \rightarrow \gamma(4S)$	
34 CHILIKIN 19 reports signals in $B^+ \rightarrow \eta_c(2S) K^+$ and $B^0 \rightarrow \eta_c(2S) K_S^0$ with 12.3 and 5.9 standard deviations, respectively.			
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_{16}/Γ	NODE=M059R2 NODE=M059R2	
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$<4 \times 10^{-4}$	90 35 WICHT	08 BELL $B^\pm \rightarrow K^\pm \gamma\gamma$	
not seen		AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$	
<0.01	90 LEE	85 CBAL $\psi' \rightarrow \text{photons}$	
35 WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S) K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S) K^+) = 4.4 \times 10^{-4}$.			
$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K\bar{K}\pi)$	Γ_{20}/Γ_2	NODE=M059R2;LINKAGE=WI	
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>		
<3.33	90 36 LEES	12AE BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$	
36 We divided the reported limit by 3 to take into account isospin relations.			

$\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$	NODE=M059R23 NODE=M059R23
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •		
$<11.8 \times 10^{-6}$	90 37 CRONIN-HEN..10	CLEO $\psi(2S) \rightarrow \gamma K^+ K^- \eta$

³⁷ CRONIN-HENNESSY 10 reports a limit of $< 5.9 \times 10^{-6}$ for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.6 \times 10^{-6}$	90	38 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

³⁸ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<12.7 \times 10^{-7}$	90	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<13.2 \times 10^{-6}$	90	39 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$

³⁹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.6 \times 10^{-6}$	90	40 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

⁴⁰ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<19.6 \times 10^{-7}$	90	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	41 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

⁴¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	42 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

⁴² Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma \times \frac{\Gamma_{157}^{\psi(2S)}}{\Gamma^{\psi(2S)}}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60	ABLIKIM	13K	BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.2	90	43 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
----------	----	-------------------	------	--

⁴³ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R25;LINKAGE=CR

NODE=M059R05
NODE=M059R05

NODE=M059R05;LINKAGE=CR

NODE=M059R18
NODE=M059R18

NODE=M059R06
NODE=M059R06

NODE=M059R06;LINKAGE=CR

NODE=M059R07
NODE=M059R07

NODE=M059R07;LINKAGE=CR

NODE=M059R19
NODE=M059R19

NODE=M059R08
NODE=M059R08

NODE=M059R08;LINKAGE=CR

NODE=M059R09
NODE=M059R09

NODE=M059R09;LINKAGE=CR

NODE=M059R10
NODE=M059R10

NODE=M059R10;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	$\Gamma_{13}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M059R20
NODE=M059R20

$$\Gamma(\eta_c(2S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	$\Gamma_{14}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M059R24
NODE=M059R24

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	$\Gamma_{17}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
$<9.7 \times 10^{-6}$	90	33	44 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$
44	Uses B($J/\psi \rightarrow e^+ e^-$) = ($5.971 \pm 0.032\%$) and B($J/\psi \rightarrow \mu^+ \mu^-$) = ($5.961 \pm 0.033\%$).				

NODE=M059R27
NODE=M059R27

NODE=M059R27;LINKAGE=A

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	$\Gamma_{18}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
$<4.3 \times 10^{-6}$	90	45 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+ \pi^- \eta$

NODE=M059R11
NODE=M059R11

NODE=M059R11;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	$\Gamma_{19}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
$<14.2 \times 10^{-6}$	90	46 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+ \pi^- \eta'$

NODE=M059R12
NODE=M059R12

NODE=M059R12;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	$\Gamma_{20}/\Gamma \times \Gamma_{157}^{\psi(2S)}/\Gamma^{\psi(2S)}$
$<1.7 \times 10^{-4}$	90	47 CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma\pi^+ \pi^- \eta_c(1S)$

NODE=M059R14
NODE=M059R14

NODE=M059R14;LINKAGE=CR

$\eta_c(2S)$ REFERENCES

CHILIKIN	19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PR L 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	980	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERM, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

NODE=M059

REFID=59899
REFID=59453
REFID=57896
REFID=58191
REFID=57978
REFID=57140
REFID=55937
REFID=54956
REFID=55583
REFID=54272
REFID=54752
REFID=53929
REFID=16751
REFID=53927
REFID=53233
REFID=52267
REFID=52064
REFID=52204
REFID=51627
REFID=51059
REFID=50773
REFID=50182
REFID=49745
REFID=49746
REFID=49188
REFID=48760
REFID=48340
REFID=46553
REFID=44623
REFID=40589
REFID=22173

$\psi(2S)$

$I^G(J^{PC}) = 0^-(1^- -)$

See the Review on “ $\psi(2S)$ and χ_c branching ratios” before the $\chi_{c0}(1P)$ Listings.

$\psi(2S)$ MASS

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.10 ± 0.06 OUR FIT				Error includes scale factor of 5.9. [3686.097 ± 0.025 MeV OUR 2019 FIT Scale factor = 2.6]

3686.097±0.010 OUR AVERAGE

3686.099 ± 0.004 ± 0.009	1	ANASHIN	15	KEDR $e^+ e^- \rightarrow$ hadrons
3686.12 ± 0.06 ± 0.10	4k	AAIJ	12H	LHCb $p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3685.95 ± 0.10	413	ARTAMONOV 00	OLYA	$e^+ e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3686.114 ± 0.007 ^{+0.011} _{-0.016}	4	ANASHIN	12	KEDR $e^+ e^- \rightarrow$ hadrons
3686.111 ± 0.025 ± 0.009		AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
3686.00 ± 0.10	413	ZHOLENTZ 80	OLYA	$e^+ e^-$

1 Supersedes AULCHENKO 03 and ANASHIN 12.

2 Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

3 Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

4 From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011$ ^{+0.002} _{-0.012} MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

5 Superseded by ARTAMONOV 00.

NODE=M071

NODE=M071M

NODE=M071M

NODE=M071M

NEW

$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

589.188±0.028 OUR AVERAGE

589.194 ± 0.027 ± 0.011	¹ AULCHENKO 03	KEDR	$e^+ e^- \rightarrow$ hadrons
589.7 ± 1.2	LEMOIGNE 82	GOLI	$185 \pi^- Be \rightarrow \gamma \mu^+ \mu^- A$
589.07 ± 0.13	¹ ZHOLENTZ 80	OLYA	$e^+ e^-$
588.7 ± 0.8	LUTH 75	MRK1	

• • • We do not use the following data for averages, fits, limits, etc. • • •

588 ± 1	² BAI	98E BES	$e^+ e^-$
---------	------------------	---------	-----------

1 Redundant with data in mass above.

2 Systematic errors not evaluated.

NODE=M071M;LINKAGE=A

NODE=M071M;LINKAGE=AR

NODE=M071M;LINKAGE=NW

NODE=M071M;LINKAGE=AN

NODE=M071M;LINKAGE=RZ

NODE=M071DM

NODE=M071DM

NODE=M071DM;LINKAGE=R

NODE=M071DM;LINKAGE=BD

NODE=M071W

NODE=M071W

$\psi(2S)$ WIDTH

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

294± 8 OUR FIT

286±16 OUR AVERAGE

358 ± 88 ± 4		ABLIKIM	08B BES2	$e^+ e^- \rightarrow$ hadrons
290 ± 25 ± 4	2.7k	ANDREOTTI 07	E835	$p\bar{p} \rightarrow e^+ e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM	06L BES2	$e^+ e^- \rightarrow$ hadrons
264 ± 27		¹ BAI	02B BES2	$e^+ e^-$
287 ± 37 ± 16		² ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+ e^-$

1 From a simultaneous fit to the hadronic and $\mu^+ \mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.

2 The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

NODE=M071W;LINKAGE=BC

NODE=M071W;LINKAGE=AN

$\psi(2S)$ DECAY MODES

NODE=M071220;NODE=M071

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(97.85 \pm 0.13) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(1.73 \pm 0.14) %	S=1.5
Γ_3 ggg	(10.6 \pm 1.6) %	DESIG=255
Γ_4 γgg	(1.03 \pm 0.29) %	DESIG=256
Γ_5 light hadrons	(15.4 \pm 1.5) %	DESIG=226
Γ_6 $e^+ e^-$	(7.93 \pm 0.17) $\times 10^{-3}$	DESIG=1
Γ_7 $\mu^+ \mu^-$	(8.0 \pm 0.6) $\times 10^{-3}$	DESIG=2
Γ_8 $\tau^+ \tau^-$	(3.1 \pm 0.4) $\times 10^{-3}$	DESIG=68
Decays into $J/\psi(1S)$ and anything		
Γ_9 $J/\psi(1S)$ anything	(61.4 \pm 0.6) %	
Γ_{10} $J/\psi(1S)$ neutrals	(25.38 \pm 0.32) %	
Γ_{11} $J/\psi(1S) \pi^+ \pi^-$	(34.68 \pm 0.30) %	DESIG=13
Γ_{12} $J/\psi(1S) \pi^0 \pi^0$	(18.24 \pm 0.31) %	DESIG=14
Γ_{13} $J/\psi(1S) \eta$	(3.37 \pm 0.05) %	DESIG=15
Γ_{14} $J/\psi(1S) \pi^0$	(1.268 \pm 0.032) $\times 10^{-3}$	DESIG=18
Hadronic decays		
Γ_{15} $\pi^0 h_c(1P)$	(8.6 \pm 1.3) $\times 10^{-4}$	
Γ_{16} $3(\pi^+ \pi^-) \pi^0$	(3.5 \pm 1.6) $\times 10^{-3}$	
Γ_{17} $2(\pi^+ \pi^-) \pi^0$	(2.9 \pm 1.0) $\times 10^{-3}$	
Γ_{18} $\rho a_2(1320)$	(2.6 \pm 0.9) $\times 10^{-4}$	
Γ_{19} $\pi^+ \pi^- \pi^0 \pi^0 \pi^0$	(5.3 \pm 0.9) $\times 10^{-3}$	
Γ_{20} $\rho^\pm \pi^\mp \pi^0 \pi^0$	< 2.7 $\times 10^{-3}$	CL=90%
Γ_{21} $p\bar{p}$	(2.94 \pm 0.08) $\times 10^{-4}$	
Γ_{22} $n\bar{n}$	(3.06 \pm 0.15) $\times 10^{-4}$	
Γ_{23} $\Delta^{++} \bar{\Delta}^{--}$	(1.28 \pm 0.35) $\times 10^{-4}$	
Γ_{24} $\Lambda \bar{\Lambda} \pi^0$	< 2.9 $\times 10^{-6}$	CL=90%
Γ_{25} $\Lambda \bar{\Lambda} \eta$	(2.5 \pm 0.4) $\times 10^{-5}$	
Γ_{26} $\Lambda \bar{p} K^+$	(1.00 \pm 0.14) $\times 10^{-4}$	
Γ_{27} $K^*(892)^+ \bar{p} \Lambda +$ c.c.	(6.3 \pm 0.7) $\times 10^{-5}$	
Γ_{28} $\Lambda \bar{p} K^+ \pi^+ \pi^-$	(1.8 \pm 0.4) $\times 10^{-4}$	
Γ_{29} $\Lambda \bar{\Lambda} \pi^+ \pi^-$	(2.8 \pm 0.6) $\times 10^{-4}$	
Γ_{30} $\Lambda \bar{\Lambda}$	(3.81 \pm 0.13) $\times 10^{-4}$	S=1.4
Γ_{31} $\Lambda \bar{\Sigma}^+ \pi^- +$ c.c.	(1.40 \pm 0.13) $\times 10^{-4}$	
Γ_{32} $\Lambda \bar{\Sigma}^- \pi^+ +$ c.c.	(1.54 \pm 0.14) $\times 10^{-4}$	
Γ_{33} $\Lambda \bar{\Sigma}^0$	(1.23 \pm 0.24) $\times 10^{-5}$	
Γ_{34} $\Sigma^0 \bar{p} K^+ +$ c.c.	(1.67 \pm 0.18) $\times 10^{-5}$	
Γ_{35} $\Sigma^+ \bar{\Sigma}^-$	(2.32 \pm 0.12) $\times 10^{-4}$	
Γ_{36} $\Sigma^0 \bar{\Sigma}^0$	(2.35 \pm 0.09) $\times 10^{-4}$	S=1.1
Γ_{37} $\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	(8.5 \pm 0.7) $\times 10^{-5}$	
Γ_{38} $\Sigma(1385)^- \bar{\Sigma}(1385)^+$	(8.5 \pm 0.8) $\times 10^{-5}$	
Γ_{39} $\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	(6.9 \pm 0.7) $\times 10^{-5}$	
Γ_{40} $\Xi^- \bar{\Xi}^+$	(2.87 \pm 0.11) $\times 10^{-4}$	S=1.1
Γ_{41} $\Xi^0 \bar{\Xi}^0$	(2.3 \pm 0.4) $\times 10^{-4}$	S=4.2
Γ_{42} $\Xi(1530)^0 \bar{\Xi}(1530)^0$	(5.2 \pm 3.2) $\times 10^{-5}$	
Γ_{43} $K^- \Lambda \bar{\Xi}^+ +$ c.c.	(3.9 \pm 0.4) $\times 10^{-5}$	
Γ_{44} $\Xi(1530)^- \bar{\Xi}(1530)^+$	(1.15 \pm 0.07) $\times 10^{-4}$	
Γ_{45} $\Xi(1530)^- \bar{\Xi}^+$	(7.0 \pm 1.2) $\times 10^{-6}$	
Γ_{46} $\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ c.c.	(5.2 \pm 1.6) $\times 10^{-6}$	
Γ_{47} $\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ +$ c.c.	(1.20 \pm 0.32) $\times 10^{-5}$	DESIG=295
Γ_{48} $K^- \sum_0^0 \bar{\Xi}^+ +$ c.c.	(3.7 \pm 0.4) $\times 10^{-5}$	DESIG=296
Γ_{49} $\Omega^- \bar{\Omega}^+$	(5.2 \pm 0.4) $\times 10^{-5}$	DESIG=74
Γ_{50} $\pi^0 p\bar{p}$	(1.53 \pm 0.07) $\times 10^{-4}$	DESIG=35
Γ_{51} $N(940) \bar{p} +$ c.c. $\rightarrow \pi^0 p\bar{p}$	(6.4 \pm 1.8) $\times 10^{-5}$	DESIG=267

Γ_{52}	$N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(7.3 ± 1.7) $\times 10^{-5}$	S=2.5	DESIG=261
Γ_{53}	$N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(6.4 ± 2.3) $\times 10^{-6}$		DESIG=268
Γ_{54}	$N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(2.5 ± 1.0) $\times 10^{-5}$		DESIG=269
Γ_{55}	$N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(3.8 ± 1.4) $\times 10^{-5}$		DESIG=270
Γ_{56}	$N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(1.79 ± 0.26) $\times 10^{-5}$		DESIG=271
Γ_{57}	$N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(2.6 ± 1.2) $\times 10^{-5}$		DESIG=272
Γ_{58}	$N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p}$	(2.13 ± 0.40) $\times 10^{-5}$		DESIG=273
Γ_{59}	$\pi^0 f_0(2100) \rightarrow \pi^0 p\bar{p}$	(1.1 ± 0.4) $\times 10^{-5}$		DESIG=262
Γ_{60}	$\eta p\bar{p}$	(6.0 ± 0.4) $\times 10^{-5}$		DESIG=200
Γ_{61}	$\eta f_0(2100) \rightarrow \eta p\bar{p}$	(1.2 ± 0.4) $\times 10^{-5}$		DESIG=263
Γ_{62}	$N(1535)\bar{p} \rightarrow \eta p\bar{p}$	(4.4 ± 0.7) $\times 10^{-5}$		DESIG=264
Γ_{63}	$\omega p\bar{p}$	(6.9 ± 2.1) $\times 10^{-5}$		DESIG=77
Γ_{64}	$\eta' p\bar{p}$	(1.10 ± 0.13) $\times 10^{-5}$		DESIG=317
Γ_{65}	$\phi p\bar{p}$	(6.1 ± 0.6) $\times 10^{-6}$		DESIG=80
Γ_{66}	$\phi X(1835) \rightarrow \phi p\bar{p}$	< 1.82 $\times 10^{-7}$	CL=90%	DESIG=318
Γ_{67}	$\pi^+ \pi^- p\bar{p}$	(6.0 ± 0.4) $\times 10^{-4}$		DESIG=31
Γ_{68}	$p\bar{n}\pi^-$ or c.c.	(2.48 ± 0.17) $\times 10^{-4}$		DESIG=227
Γ_{69}	$p\bar{n}\pi^- \pi^0$	(3.2 ± 0.7) $\times 10^{-4}$		DESIG=228
Γ_{70}	$2(\pi^+ \pi^- \pi^0)$	(4.8 ± 1.5) $\times 10^{-3}$		DESIG=221
Γ_{71}	$\eta \pi^+ \pi^-$	< 1.6 $\times 10^{-4}$	CL=90%	DESIG=202
Γ_{72}	$\eta \pi^+ \pi^- \pi^0$	(9.5 ± 1.7) $\times 10^{-4}$		DESIG=203
Γ_{73}	$2(\pi^+ \pi^-) \eta$	(1.2 ± 0.6) $\times 10^{-3}$		DESIG=251
Γ_{74}	$\pi^+ \pi^- \pi^0 \pi^0 \eta$	< 4 $\times 10^{-4}$	CL=90%	DESIG=313
Γ_{75}	$\eta' \pi^+ \pi^- \pi^0$	(4.5 ± 2.1) $\times 10^{-4}$		DESIG=204
Γ_{76}	$\omega \pi^+ \pi^-$	(7.3 ± 1.2) $\times 10^{-4}$	S=2.1	DESIG=75
Γ_{77}	$b_1^\pm \pi^\mp$	(4.0 ± 0.6) $\times 10^{-4}$	S=1.1	DESIG=40
Γ_{78}	$b_1^0 \pi^0$	(2.4 ± 0.6) $\times 10^{-4}$		DESIG=193
Γ_{79}	$\omega f_2(1270)$	(2.2 ± 0.4) $\times 10^{-4}$		DESIG=64
Γ_{80}	$\omega \pi^0 \pi^0$	(1.11 ± 0.35) $\times 10^{-3}$		DESIG=314
Γ_{81}	$\pi^0 \pi^0 K^+ K^-$	(2.6 ± 1.3) $\times 10^{-4}$		DESIG=298
Γ_{82}	$\pi^+ \pi^- K^+ K^-$	(7.3 ± 0.5) $\times 10^{-4}$		DESIG=26
Γ_{83}	$\pi^0 \pi^0 K_S^0 K_L^0$	(1.3 ± 0.6) $\times 10^{-3}$		DESIG=304
Γ_{84}	$\rho^0 K^+ K^-$	(2.2 ± 0.4) $\times 10^{-4}$		DESIG=205
Γ_{85}	$K^*(892)^0 \bar{K}_2^*(1430)^0$	(1.9 ± 0.5) $\times 10^{-4}$		DESIG=66
Γ_{86}	$K^+ K^- \pi^+ \pi^- \eta$	(1.3 ± 0.7) $\times 10^{-3}$		DESIG=252
Γ_{87}	$K^+ K^- 2(\pi^+ \pi^-) \pi^0$	(1.00 ± 0.31) $\times 10^{-3}$		DESIG=240
Γ_{88}	$K^+ K^- 2(\pi^+ \pi^-)$	(1.9 ± 0.9) $\times 10^{-3}$		DESIG=222
Γ_{89}	$K_1(1270)^\pm K^\mp$	(1.00 ± 0.28) $\times 10^{-3}$		DESIG=41
Γ_{90}	$K_S^0 K_S^0 \pi^+ \pi^-$	(2.2 ± 0.4) $\times 10^{-4}$		DESIG=225
Γ_{91}	$\rho^0 p\bar{p}$	(5.0 ± 2.2) $\times 10^{-5}$		DESIG=210
Γ_{92}	$K^+ \bar{K}^*(892)^0 \pi^-$ + c.c.	(6.7 ± 2.5) $\times 10^{-4}$		DESIG=34
Γ_{93}	$2(\pi^+ \pi^-)$	(2.4 ± 0.6) $\times 10^{-4}$	S=2.2	DESIG=24
Γ_{94}	$\rho^0 \pi^+ \pi^-$	(2.2 ± 0.6) $\times 10^{-4}$	S=1.4	DESIG=33
Γ_{95}	$K^+ K^- \pi^+ \pi^- \pi^0$	(1.26 ± 0.09) $\times 10^{-3}$		DESIG=206
Γ_{96}	$\omega f_0(1710) \rightarrow \omega K^+ K^-$	(5.9 ± 2.2) $\times 10^{-5}$		DESIG=216
Γ_{97}	$K^*(892)^0 K^- \pi^+ \pi^0$ + c.c.	(8.6 ± 2.2) $\times 10^{-4}$		DESIG=217
Γ_{98}	$K^*(892)^+ K^- \pi^+ \pi^-$ + c.c.	(9.6 ± 2.8) $\times 10^{-4}$		DESIG=218
Γ_{99}	$K^*(892)^+ K^- \rho^0$ + c.c.	(7.3 ± 2.6) $\times 10^{-4}$		DESIG=219
Γ_{100}	$K^*(892)^0 K^- \rho^+$ + c.c.	(6.1 ± 1.8) $\times 10^{-4}$		DESIG=220
Γ_{101}	$\eta K^+ K^-$, no $\eta \phi$	(3.1 ± 0.4) $\times 10^{-5}$		DESIG=207
Γ_{102}	$\omega K^+ K^-$	(1.62 ± 0.11) $\times 10^{-4}$	S=1.1	DESIG=76
Γ_{103}	$\omega K^*(892)^+ K^-$ + c.c.	(2.07 ± 0.26) $\times 10^{-4}$		DESIG=276
Γ_{104}	$\omega K_2^*(1430)^+ K^-$ + c.c.	(6.1 ± 1.2) $\times 10^{-5}$		DESIG=277
Γ_{105}	$\omega \bar{K}^*(892)^0 K^0$	(1.68 ± 0.30) $\times 10^{-4}$		DESIG=278

Γ_{106}	$\omega \bar{K}_2^*(1430)^0 K^0$	(5.8 \pm 2.2) $\times 10^{-5}$	DESIG=279
Γ_{107}	$\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c.	(1.6 \pm 0.4) $\times 10^{-5}$	DESIG=282
Γ_{108}	$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	(1.09 \pm 0.26) $\times 10^{-5}$	DESIG=283
Γ_{109}	$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c.	(3.0 \pm 1.0) $\times 10^{-6}$	DESIG=284
Γ_{110}	$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	(1.2 \pm 0.7) $\times 10^{-6}$	DESIG=285
Γ_{111}	$3(\pi^+ \pi^-)$	(3.5 \pm 2.0) $\times 10^{-4}$	S=2.8 DESIG=32
Γ_{112}	$p\bar{p} \pi^+ \pi^- \pi^0$	(7.3 \pm 0.7) $\times 10^{-4}$	DESIG=211
Γ_{113}	$K^+ K^-$	(7.5 \pm 0.5) $\times 10^{-5}$	DESIG=23
Γ_{114}	$K_S^0 K_L^0$	(5.34 \pm 0.33) $\times 10^{-5}$	DESIG=85
Γ_{115}	$\pi^+ \pi^- \pi^0$	(2.01 \pm 0.17) $\times 10^{-4}$	S=1.7 DESIG=36
Γ_{116}	$\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0$	(1.9 \pm 1.2) $\times 10^{-4}$	DESIG=201
Γ_{117}	$\rho(770)\pi \rightarrow \pi^+ \pi^- \pi^0$	(3.2 \pm 1.2) $\times 10^{-5}$	S=1.8 DESIG=22
Γ_{118}	$\pi^+ \pi^-$	(7.8 \pm 2.6) $\times 10^{-6}$	DESIG=21
Γ_{119}	$K_1(1400)^\pm K^\mp$	< 3.1 $\times 10^{-4}$	CL=90% DESIG=42
Γ_{120}	$K_2^*(1430)^\pm K^\mp$	(7.1 \pm 1.3) $\times 10^{-5}$	DESIG=265
Γ_{121}	$K^+ K^- \pi^0$	(4.07 \pm 0.31) $\times 10^{-5}$	DESIG=38
Γ_{122}	$K_S^0 K_L^0 \pi^0$	< 3.0 $\times 10^{-4}$	CL=90% DESIG=303
Γ_{123}	$K_S^0 K_L^0 \eta$	(1.3 \pm 0.5) $\times 10^{-3}$	DESIG=305
Γ_{124}	$K^+ K^*(892)^- +$ c.c.	(2.9 \pm 0.4) $\times 10^{-5}$	S=1.2 DESIG=39
Γ_{125}	$K^*(892)^0 \bar{K}^0 +$ c.c.	(1.09 \pm 0.20) $\times 10^{-4}$	DESIG=194
Γ_{126}	$\phi \pi^+ \pi^-$	(1.18 \pm 0.26) $\times 10^{-4}$	S=1.5 DESIG=78
Γ_{127}	$\phi f_0(980) \rightarrow \pi^+ \pi^-$	(7.5 \pm 3.3) $\times 10^{-5}$	S=1.6 DESIG=81
Γ_{128}	$2(K^+ K^-)$	(6.3 \pm 1.3) $\times 10^{-5}$	DESIG=208
Γ_{129}	$\phi K^+ K^-$	(7.0 \pm 1.6) $\times 10^{-5}$	DESIG=79
Γ_{130}	$2(K^+ K^-) \pi^0$	(1.10 \pm 0.28) $\times 10^{-4}$	DESIG=209
Γ_{131}	$\phi \eta$	(3.10 \pm 0.31) $\times 10^{-5}$	DESIG=89
Γ_{132}	$\eta \phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-$	< 2.2 $\times 10^{-6}$	CL=90% DESIG=316
Γ_{133}	$\phi \eta'$	(1.54 \pm 0.20) $\times 10^{-5}$	DESIG=90
Γ_{134}	$\phi f_1(1285)$	(3.0 \pm 1.3) $\times 10^{-5}$	DESIG=319
Γ_{135}	$\phi \eta(1405) \rightarrow \phi \pi^+ \pi^- \eta$	(8.5 \pm 1.7) $\times 10^{-6}$	DESIG=320
Γ_{136}	$\omega \eta'$	(3.2 \pm 2.5) $\times 10^{-5}$	DESIG=91
Γ_{137}	$\omega \pi^0$	(2.1 \pm 0.6) $\times 10^{-5}$	DESIG=92
Γ_{138}	$\rho \eta'$	(1.9 \pm 1.7) $\times 10^{-5}$	DESIG=93
Γ_{139}	$\rho \eta$	(2.2 \pm 0.6) $\times 10^{-5}$	S=1.1 DESIG=94
Γ_{140}	$\omega \eta$	< 1.1 $\times 10^{-5}$	CL=90% DESIG=95
Γ_{141}	$\phi \pi^0$	< 4 $\times 10^{-7}$	CL=90% DESIG=96
Γ_{142}	$\eta_c \pi^+ \pi^- \pi^0$	< 1.0 $\times 10^{-3}$	CL=90% DESIG=229
Γ_{143}	$p\bar{p} K^+ K^-$	(2.7 \pm 0.7) $\times 10^{-5}$	DESIG=212
Γ_{144}	$\bar{\Lambda} n K_S^0 +$ c.c.	(8.1 \pm 1.8) $\times 10^{-5}$	DESIG=237
Γ_{145}	$\phi f'_2(1525)$	(4.4 \pm 1.6) $\times 10^{-5}$	DESIG=67
Γ_{146}	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} +$ c.c.	< 8.8 $\times 10^{-6}$	CL=90% DESIG=195
Γ_{147}	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	< 1.0 $\times 10^{-5}$	CL=90% DESIG=196
Γ_{148}	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	< 7.0 $\times 10^{-6}$	CL=90% DESIG=197
Γ_{149}	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	< 2.6 $\times 10^{-5}$	CL=90% DESIG=198
Γ_{150}	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	< 6.0 $\times 10^{-6}$	CL=90% DESIG=199
Γ_{151}	$K_S^0 K_S^0$	< 4.6 $\times 10^{-6}$	DESIG=86
Γ_{152}	$\Lambda_c^+ \bar{p} e^+ e^- +$ c.c.	< 1.7 $\times 10^{-6}$	CL=90% DESIG=310

Radiative decays

Γ_{153}	$\gamma\chi_{c0}(1P)$	(9.79 \pm 0.20) %	NODE=M071;CLUMP=C	
Γ_{154}	$\gamma\chi_{c1}(1P)$	(9.75 \pm 0.24) %	DESIG=56	
Γ_{155}	$\gamma\chi_{c2}(1P)$	(9.52 \pm 0.20) %	DESIG=58	
Γ_{156}	$\gamma\eta_c(1S)$	(3.4 \pm 0.5) \times 10 ⁻³	DESIG=59	
Γ_{157}	$\gamma\eta_c(2S)$	(7 \pm 5) \times 10 ⁻⁴	DESIG=61	
Γ_{158}	$\gamma\pi^0$	(1.04 \pm 0.22) \times 10 ⁻⁶	DESIG=63	
Γ_{159}	$\gamma\eta'(958)$	(1.24 \pm 0.04) \times 10 ⁻⁴	DESIG=52	
Γ_{160}	$\gamma f_2(1270)$	(2.73 \pm 0.29) \times 10 ⁻⁴	DESIG=54	
Γ_{161}	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	(3.1 \pm 1.7) \times 10 ⁻⁵	DESIG=82	
Γ_{162}	$\gamma f_0(1500)$	(9.3 \pm 1.9) \times 10 ⁻⁵	DESIG=286	
Γ_{163}	$\gamma f'_2(1525)$	(3.3 \pm 0.8) \times 10 ⁻⁵	DESIG=287	
Γ_{164}	$\gamma f_0(1710)$		DESIG=288	
Γ_{165}	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	(3.5 \pm 0.6) \times 10 ⁻⁵	DESIG=236	
Γ_{166}	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	(6.6 \pm 0.7) \times 10 ⁻⁵	DESIG=83	
Γ_{167}	$\gamma f_0(2100) \rightarrow \gamma\pi\pi$	(4.8 \pm 1.0) \times 10 ⁻⁶	DESIG=84	
Γ_{168}	$\gamma f_0(2200) \rightarrow \gamma K\bar{K}$	(3.2 \pm 1.0) \times 10 ⁻⁶	DESIG=289	
Γ_{169}	$\gamma f_J(2220) \rightarrow \gamma\pi\pi$	< 5.8 \times 10 ⁻⁶	DESIG=290	
Γ_{170}	$\gamma f_J(2220) \rightarrow \gamma K\bar{K}$	< 9.5 \times 10 ⁻⁶	DESIG=291	
Γ_{171}	$\gamma\gamma$	< 1.5 \times 10 ⁻⁴	DESIG=292	
Γ_{172}	$\gamma\eta$	(9.2 \pm 1.8) \times 10 ⁻⁷	DESIG=51	
Γ_{173}	$\gamma\eta\pi^+\pi^-$	(8.7 \pm 2.1) \times 10 ⁻⁴	DESIG=53	
Γ_{174}	$\gamma\eta(1405)$		DESIG=230	
Γ_{175}	$\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi$	< 9 \times 10 ⁻⁵	DESIG=231	
Γ_{176}	$\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-$	(3.6 \pm 2.5) \times 10 ⁻⁵	DESIG=62	
Γ_{177}	$\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0$	< 5.0 \times 10 ⁻⁷	DESIG=232	
Γ_{178}	$\gamma\eta(1475)$		DESIG=308	
Γ_{179}	$\gamma\eta(1475) \rightarrow K\bar{K}\pi$	< 1.4 \times 10 ⁻⁴	DESIG=233	
Γ_{180}	$\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-$	< 8.8 \times 10 ⁻⁵	DESIG=234	
Γ_{181}	$\gamma 2(\pi^+\pi^-)$	(4.0 \pm 0.6) \times 10 ⁻⁴	DESIG=235	
Γ_{182}	$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	(3.7 \pm 0.9) \times 10 ⁻⁴	DESIG=241	
Γ_{183}	$\gamma K^{*0} \bar{K}^{*0}$	(2.4 \pm 0.7) \times 10 ⁻⁴	DESIG=242	
Γ_{184}	$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	(2.6 \pm 0.5) \times 10 ⁻⁴	DESIG=243	
Γ_{185}	$\gamma K^+ K^- \pi^+ \pi^-$	(1.9 \pm 0.5) \times 10 ⁻⁴	DESIG=244	
Γ_{186}	$\gamma p\bar{p}$	(3.9 \pm 0.5) \times 10 ⁻⁵	DESIG=245	
Γ_{187}	$\gamma f_2(1950) \rightarrow \gamma p\bar{p}$	< 1.20 \pm 0.22 \times 10 ⁻⁵	S=2.0	DESIG=246
Γ_{188}	$\gamma f_2(2150) \rightarrow \gamma p\bar{p}$	(7.2 \pm 1.8) \times 10 ⁻⁶	DESIG=257	
Γ_{189}	$\gamma X(1835) \rightarrow \gamma p\bar{p}$	(4.6 \pm 1.8) \times 10 ⁻⁶	DESIG=258	
Γ_{190}	$\gamma X \rightarrow \gamma p\bar{p}$	[a] < 2 \times 10 ⁻⁶	DESIG=259	
Γ_{191}	$\gamma\pi^+\pi^- p\bar{p}$	(2.8 \pm 1.4) \times 10 ⁻⁵	DESIG=260	
Γ_{192}	$\gamma 2(\pi^+\pi^-) K^+ K^-$	< 2.2 \times 10 ⁻⁴	DESIG=247	
Γ_{193}	$\gamma 3(\pi^+\pi^-)$	< 1.7 \times 10 ⁻⁴	DESIG=248	
Γ_{194}	$\gamma K^+ K^- K^+ K^-$	< 4 \times 10 ⁻⁵	DESIG=249	
Γ_{195}	$\gamma\gamma J/\psi$	(3.1 \pm 1.0) \times 10 ⁻⁴	DESIG=250	
Γ_{196}	$e^+ e^- \eta'$	(1.90 \pm 0.26) \times 10 ⁻⁶	DESIG=266	
Γ_{197}	$e^+ e^- \chi_{c0}(1P)$	(1.06 \pm 0.24) \times 10 ⁻³	DESIG=311	
Γ_{198}	$e^+ e^- \chi_{c1}(1P)$	(8.5 \pm 0.6) \times 10 ⁻⁴	DESIG=300	
Γ_{199}	$e^+ e^- \chi_{c2}(1P)$	(7.0 \pm 0.8) \times 10 ⁻⁴	DESIG=301	
Γ_{200}	$D^0 e^+ e^- + \text{c.c.}$	< 1.4 \times 10 ⁻⁷	DESIG=302	

Weak decays

Γ_{200}	$D^0 e^+ e^- + \text{c.c.}$	< 1.4 \times 10 ⁻⁷	CL=90%	NODE=M071;CLUMP=E
Γ_{201}	invisible	< 1.6	%	DESIG=306

Other decays

Γ_{201}	invisible	< 1.6	%	CL=90%	DESIG=275
----------------	-----------	-------	---	--------	-----------

[a] For a narrow resonance in the range 2.2 < M(X) < 2.8 GeV.

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 378.1$ for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

	x_7	3								
	x_8	1	0							
	x_{11}	29	11	2						
	x_{12}	28	6	1	48					
	x_{13}	13	4	1	36	15				
	x_{21}	0	0	0	4	3	2			
	x_{153}	1	0	0	2	1	1	0		
	x_{154}	1	0	0	2	1	1	0	0	
	x_{155}	1	0	0	3	1	1	0	0	0
Γ		-81	-4	-1	-38	-34	-16	-7	-1	-1
	x_6	x_7	x_8	x_{11}	x_{12}	x_{13}	x_{21}	x_{153}	x_{154}	x_{155}

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

258±26	BAI	02B	BES2 $e^+ e^-$
224±56	LUTH	75	MRK1 $e^+ e^-$

Γ_1

NODE=M071225

NODE=M071W3
NODE=M071W3

$\Gamma(e^+ e^-)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

2.33 ±0.04 OUR FIT**2.29 ±0.06 OUR AVERAGE**

2.23 ± 0.10 ± 0.02	¹ ABLIKIM	15V	BES3	4.0–4.4 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
2.338±0.037±0.096	ABLIKIM	08B	BES2	$e^+ e^- \rightarrow \text{hadrons}$
2.330±0.036±0.110	ABLIKIM	06L	BES2	$e^+ e^- \rightarrow \text{hadrons}$
2.44 ± 0.21	² BAI	02B	BES2	$e^+ e^-$
2.14 ± 0.21	ALEXANDER	89	RVUE	See γ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.279±0.015±0.042	³ ANASHIN	18	KEDR	$e^+ e^-$
2.282±0.015±0.042	⁴ ANASHIN	18	KEDR	$e^+ e^-$
2.0 ± 0.3	BRANDELIK	79C	DASP	$e^+ e^-$
2.1 ± 0.3	⁵ LUTH	75	MRK1	$e^+ e^-$

Γ_6

NODE=M071W1
NODE=M071W1

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau / 0.38847$.

³ Combining $\Gamma_{e^+ e^-} \cdot B(\mu^+ \mu^-)$ from ANASHIN 18 with $\Gamma_{e^+ e^-} \cdot B(\text{hadrons})$ from ANASHIN 12 and assuming lepton universality.

⁴ From the sum of $\Gamma_{e^+ e^-} \cdot B(\text{hadrons})$ from ANASHIN 12, $\Gamma_{e^+ e^-} \cdot B(e^+ e^-)$ and $\Gamma_{e^+ e^-} \cdot B(\mu^+ \mu^-)$ from ANASHIN 18, and $\Gamma_{e^+ e^-} \cdot B(\tau^+ \tau^-)$ from ANASHIN 07.

⁵ From a simultaneous fit to $e^+ e^-$, $\mu^+ \mu^-$, and hadronic channels assuming $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$.

OCCUR=2

NODE=M071W1;LINKAGE=A

NODE=M071W;LINKAGE=BB

NODE=M071W1;LINKAGE=B

NODE=M071W1;LINKAGE=C

NODE=M071W1;LINKAGE=F

$\Gamma(\gamma\gamma)$					Γ_{171}
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<43	90	BRANDELIK	79c	DASP	$e^+ e^-$

$\psi(2S) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $e^+ e^-$ and with the total width is obtained from the integrated cross section into channel(i) in the $e^+ e^-$ annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_1 \Gamma_6/\Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
2.233 ± 0.015 ± 0.042	1 ANASHIN	12	KEDR	$e^+ e^- \rightarrow \text{hadrons}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

2.2 ± 0.4 ABRAMS 75 MRK1 $e^+ e^-$

¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

$\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_6 \Gamma_6/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
21.2 ± 0.7 ± 1.2	1 ANASHIN	18	KEDR	$e^+ e^-$	

¹ From the average of nine scans of the $\psi(2S)$.

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_7 \Gamma_6/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
19.3 ± 0.3 ± 0.5	1 ANASHIN	18	KEDR	$\psi(2S) \rightarrow \mu^+ \mu^-$	

¹ From the average of nine scans of the $\psi(2S)$.

$\Gamma(\tau^+ \tau^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_8 \Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

9.0 ± 2.6 79 ¹ ANASHIN 07 KEDR $e^+ e^- \rightarrow \psi(2S) \rightarrow \tau^+ \tau^-$

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{11} \Gamma_6/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.808 ± 0.013 OUR FIT				Error includes scale factor of 1.3. See the ideogram below.	

0.837 ± 0.025 OUR AVERAGE	1 LEES 12E BABR 10.6 $e^+ e^- \rightarrow 2\pi^+ 2\pi^- \gamma$
	0.852 ± 0.010 ± 0.026 19.5k ADAM 06 CLEO 3.773 $e^+ e^- \rightarrow \gamma\psi(2S)$
	0.68 ± 0.09 2 BAI 98E BES $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.03 256 ³ AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
0.755 ± 0.048 ± 0.004 544 ⁴ AUBERT 05D BABR 10.6 $e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$

¹ LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+ \mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² The value of $\Gamma(e^+ e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

³ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+ \mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

Superseded by LEES 12E.

NODE=M071W51
NODE=M071W51

NODE=M071230

NODE=M071230

NODE=M071G3
NODE=M071G3

NODE=M071G3;LINKAGE=AN

NODE=M071P14
NODE=M071P14

NODE=M071P14;LINKAGE=A

NODE=M071P13
NODE=M071P13

NODE=M071P13;LINKAGE=A

NODE=M071G9
NODE=M071G9

NODE=M071G9;LINKAGE=AN


NODE=M071G1
NODE=M071G1

NODE=M071G1;LINKAGE=LE

NODE=M071G1;LINKAGE=A

NODE=M071G1;LINKAGE=UB

NODE=M071G1;LINKAGE=AU



$$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{12}\Gamma_6/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.425±0.009 OUR FIT				
0.411±0.008±0.018	3.6k	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.48 ± 0.09 ± 0.02	142	¹ LEES	18E	BABR $10.6 e^+e^- \rightarrow J/\psi\pi^0\pi^0\gamma$

¹ LEES 18E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0101 \pm 0.0015 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.6± 1.6 OUR FIT				
87 ± 9 OUR AVERAGE				
83 ± 25 ± 5	14	¹ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ± 7	291 ± 24	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{(\psi(2S))} \cdot B(\psi(2S) \rightarrow J/\psi\eta) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV.

$$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_6/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8	90	<37	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$

$$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{21}\Gamma_6/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.686±0.019 OUR FIT				

0.63 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.2.
0.67 ± 0.12 ± 0.02	43 ¹ LEES 130 BABR $e^+e^- \rightarrow p\bar{p}\gamma$
0.74 ± 0.07 ± 0.04	142 ² LEES 13Y BABR $e^+e^- \rightarrow p\bar{p}\gamma$
0.579±0.038±0.036	2.7k ANDREOTTI 07 E835 $p\bar{p} \rightarrow e^+e^-$, $J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.70 ± 0.17 ± 0.03	22 ³ AUBERT 06B BABR $e^+e^- \rightarrow p\bar{p}\gamma$

¹ ISR photon reconstructed in the detector

² ISR photon undetected

³ Superseded by LEES 130

NODE=M071G6
NODE=M071G6

NODE=M071G6;LINKAGE=A

NODE=M071G7
NODE=M071G7

NODE=M071G7;LINKAGE=UB

NODE=M071G8
NODE=M071G8

NODE=M071G2
NODE=M071G2

NODE=M071G2;LINKAGE=C
NODE=M071G2;LINKAGE=B
NODE=M071G2;LINKAGE=A

NODE=M071G11
NODE=M071G11

$$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_6/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.5±0.4±0.1	AUBERT	07BD	BABR $10.6 e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{70}\Gamma_6/\Gamma$	NODE=M071G4 NODE=M071G4
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
11.2±3.3±1.3	43	AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$	
$\Gamma(\pi^0\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{81}\Gamma_6/\Gamma$	NODE=M071G08 NODE=M071G08
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.60±0.31±0.03	17	LEES	12F	BABR	$10.6 \pi^0\pi^0K^+K^- \rightarrow \pi^0\pi^0K^+K^- \gamma$	
$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{88}\Gamma_6/\Gamma$	NODE=M071G5 NODE=M071G5
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
4.4±2.1±0.3	26	AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$	
$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{82}\Gamma_6/\Gamma$	NODE=M071G12 NODE=M071G12
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
1.92±0.30±0.06	133	LEES	12F	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2.56±0.42±0.16	85	¹ AUBERT	07AK	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
1 Superseded by LEES 12F.						
$\Gamma(\pi^0\pi^0K_S^0K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{83}\Gamma_6/\Gamma$	NODE=M071G14 NODE=M071G14
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
2.92±1.27±0.15	14	LEES	17A	BABR	$e^+e^- \rightarrow K_S^0K_L^0\pi^0\pi^0\gamma$	
$\Gamma(K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{122}\Gamma_6/\Gamma$	NODE=M071G15 NODE=M071G15
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<0.7	90	8	LEES	17A	BABR	$e^+e^- \rightarrow K_S^0K_L^0\pi^0\gamma$
$\Gamma(K_S^0K_L^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{123}\Gamma_6/\Gamma$	NODE=M071G16 NODE=M071G16
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
3.14±1.08±0.16	16	LEES	17A	BABR	$e^+e^- \rightarrow K_S^0K_L^0\eta\gamma$	
$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{127}\Gamma_6/\Gamma$	NODE=M071G13 NODE=M071G13
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.345±0.128±0.004	12	¹ LEES	12F	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.345±0.168±0.004	6 ± 3	² AUBERT	07AK	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$	
1 LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+K^-$) = $0.17 \pm 0.06 \pm 0.02$ eV which we divide by our best value B($\phi(1020) \rightarrow K^+K^-$) = $(49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						
2 Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+K^-$) = $0.17 \pm 0.08 \pm 0.02$ eV which we divide by our best value B($\phi(1020) \rightarrow K^+K^-$) = $(49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						
$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{128}\Gamma_6/\Gamma$	NODE=M071G07 NODE=M071G07
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.22±0.10±0.02	13	LEES	12F	BABR	$10.6 e^+e^- \rightarrow K^+K^-K^+K^-\gamma$	
$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{126}\Gamma_6/\Gamma$	NODE=M071G10 NODE=M071G10
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
0.55±0.19±0.01	19	¹ LEES	12F	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.57±0.23±0.01	10	² AUBERT,BE	06D	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$	

¹ LEES 12F reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.27 \pm 0.09 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT,BE 06D reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.28 \pm 0.11 \pm 0.02 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{17}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.7±2.2±1.8	410	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$

$\Gamma(\pi^+\pi^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{19}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±1.8±1.2	177	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

$\Gamma(\rho^\pm\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{20}\Gamma_6/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.2	90	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{76}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.01±0.84±0.02	37	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.69 \pm 0.73 \pm 0.16 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{80}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.58±0.82±0.02	33	¹ LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

¹ LEES 18E reports $[\Gamma(\psi(2S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.3 \pm 0.7 \pm 0.2 \text{ eV}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.3 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{73}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.87±1.41±0.01	16	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\pi^+\pi^-\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{74}\Gamma_6/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.85	90	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{95}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±1.3±0.3	32	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$

$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{86}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.04±1.79±0.02	7	¹ AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G10;LINKAGE=A

NODE=M071G10;LINKAGE=AU

NODE=M071G01
NODE=M071G01NODE=M071P16
NODE=M071P16NODE=M071G02
NODE=M071G02NODE=M071P17
NODE=M071P17

NODE=M071P17;LINKAGE=A

NODE=M071G03
NODE=M071G03

NODE=M071G03;LINKAGE=UB

NODE=M071P15
NODE=M071P15NODE=M071G04
NODE=M071G04NODE=M071G05
NODE=M071G05

NODE=M071G05;LINKAGE=UB

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{113}\Gamma_6/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.147 $\pm 0.035 \pm 0.005$	66	¹ LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$	
0.197 $\pm 0.035 \pm 0.005$	66	² LEES	15J BABR	$e^+e^- \rightarrow K^+K^-\gamma$	
0.35 $\pm 0.14 \pm 0.03$	11	³ LEES	13Q BABR	$e^+e^- \rightarrow K^+K^-\gamma$	
1 $\sin\phi > 0$. 2 $\sin\phi < 0$.					
3 Interference with non-resonant K^+K^- production not taken into account.					

$\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.9785 ± 0.0013 OUR AVERAGE					
0.9779 ± 0.0015	¹ BAI	02B BES2	e^+e^-		
0.981 ± 0.003	¹ LUTH	75 MRK1	e^+e^-		
1 Includes cascade decay into $J/\psi(1S)$.					
$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.0173 ± 0.0014 OUR AVERAGE Error includes scale factor of 1.5.					
0.0166 ± 0.0010	^{1,2} SETH	04 RVUE	e^+e^-		
0.0199 ± 0.0019	¹ BAI	02B BES2	e^+e^-		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
0.029 ± 0.004	¹ LUTH	75 MRK1	e^+e^-		
1 Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.					
2 Using $B(\psi(2S) \rightarrow \ell^+\ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02c.					

$\Gamma(ggg)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
10.58 ± 1.62					
10.58 ± 1.62	2.9 M	¹ LIBBY	09 CLEO	$\psi(2S) \rightarrow \text{hadrons}$	
1 Calculated using $\Gamma(\gamma gg)/\Gamma(ggg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow X J/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma \eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+\ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.					

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.025 ± 0.288					
1.025 ± 0.288	200 k	¹ LIBBY	09 CLEO	$\psi(2S) \rightarrow \gamma + \text{hadrons}$	
1 Calculated using $\Gamma(\gamma gg)/\Gamma(ggg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.					

$\Gamma(\gamma gg)/\Gamma(ggg)$					Γ_4/Γ_3
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
9.7 $\pm 2.6 \pm 1.6$					
9.7 $\pm 2.6 \pm 1.6$	2.9 M	LIBBY	09 CLEO	$\psi(2S) \rightarrow (\gamma +) \text{hadrons}$	

$\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.154 ± 0.015					
0.154 ± 0.015	¹ MENDEZ	08 CLEO	$e^+e^- \rightarrow \psi(2S)$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					

0.169 ± 0.026	² ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S)$		
1 Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.					
2 Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{cJ}\gamma)$, $B(\eta_c\gamma)$ from ATHAR 04 and $B(\ell^+\ell^-)$ from PDG 04. Superseded by MENDEZ 08.					

NODE=M071G06

NODE=M071G06

OCCUR=2

NODE=M071G06;LINKAGE=A
NODE=M071G06;LINKAGE=B
NODE=M071G06;LINKAGE=BA

NODE=M071235

NODE=M071R3
NODE=M071R3NODE=M071R;LINKAGE=P
NODE=M071R5
NODE=M071R5NODE=M071S43
NODE=M071S43NODE=M071S43;LINKAGE=LI
NODE=M071S44;LINKAGE=LINODE=M071S44
NODE=M071S44NODE=M071S27
NODE=M071S27NODE=M071S27;LINKAGE=ME
NODE=M071S27;LINKAGE=AD

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

 Γ_6/Γ

NODE=M071R1

NODE=M071R1

 79.3 ± 1.7 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

88 ± 13

1 FELDMAN 77 RVUE e^+e^-

¹ From an overall fit assuming equal partial widths for e^+e^- and $\mu^+\mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

NODE=M071R;LINKAGE=L

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

DOCUMENT ID

 Γ_7/Γ

NODE=M071R2

NODE=M071R2

 80 ± 6 OUR FIT $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$

VALUE

DOCUMENT ID

TECN

COMMENT

 Γ_7/Γ_6

NODE=M071R4

NODE=M071R4

 1.00 ± 0.08 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 ± 0.16

BOYARSKI 75C MRK1 e^+e^- $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

 Γ_8/Γ

NODE=M071R75

NODE=M071R75

 31 ± 4 OUR FIT **$30.8 \pm 2.1 \pm 3.8$** 1 ABLIKIM 06W BES $e^+e^- \rightarrow \psi(2S)$

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

NODE=M071R75;LINKAGE=AB

— DECAYS INTO $J/\psi(1S)$ AND ANYTHING — $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_9/Γ

NODE=M071R10

NODE=M071R10

 0.614 ± 0.006 OUR FIT **0.55 ± 0.07 OUR AVERAGE**

0.51 ± 0.12

BRANDELIK 79C DASP $e^+e^- \rightarrow \mu^+\mu^-X$

0.57 ± 0.08

ABRAMS 75B MRK1 $e^+e^- \rightarrow \mu^+\mu^-X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.6254 \pm 0.0016 \pm 0.0155$ 1.1M 1 MENDEZ 08 CLEO $\psi(2S) \rightarrow \ell^+\ell^-X$
 $0.5950 \pm 0.0015 \pm 0.0190$ 151k ADAM 05A CLEO Repl. by MENDEZ 08

1 Not independent from other measurements of MENDEZ 08.

NODE=M071R10;LINKAGE=ME

 $\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_6/\Gamma_9 = \Gamma_6 / (\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})$$

VALUE (units 10^{-2})

EVTS

DOCUMENT ID

TECN

COMMENT


NODE=M071R72

NODE=M071R72

 1.291 ± 0.026 OUR FIT **1.28 ± 0.04 OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.1.22 ± 0.02 ± 0.05 5097 ± 73 1 ANDREOTTI 05 E835 $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$ 1.28 ± 0.03 ± 0.02 1 AMBROGIANI 00A E835 $p\bar{p} \rightarrow \psi(2S)$ 1.44 ± 0.08 ± 0.02 1 ARMSTRONG 97 E760 $\bar{p}p \rightarrow \psi(2S)$

1 Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

NODE=M071R;LINKAGE=7A



$\Gamma(\mu^+ \mu^-)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_7/\Gamma_9 = \Gamma_7/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0130±0.0010 OUR FIT			
0.014 ± 0.003	HILGER	75	SPEC $e^+ e^-$

$\Gamma(J/\psi(1S)\text{ neutrals})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID
0.2538±0.0032 OUR FIT	

Γ_{10}/Γ

NODE=M071R74
NODE=M071R74

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3468±0.0030 OUR FIT				

Γ_{11}/Γ

NODE=M071R18
NODE=M071R18

0.348 ± 0.005 OUR AVERAGE

Error includes scale factor of 1.3. See the ideogram below.


0.3498±0.0002±0.0045	20M	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$
0.3504±0.0007±0.0077	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.323 ± 0.014		BAI	02B	BES2 $e^+ e^-$
0.32 ± 0.04		ABRAMS	75B	MRK1 $e^+ e^- \rightarrow J/\psi\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3354±0.0014±0.0110 60k ¹ADAM 05A CLEO Repl. by MENDEZ 08

¹ Not independent from other values reported by ADAM 05A.

NODE=M071R;LINKAGE=AD

 $\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0229±0.0005 OUR FIT			
0.0252±0.0028±0.0011	¹ AUBERT	02B BABR	e^+e^-

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

 Γ_6/Γ_{11}

NODE=M071R73
NODE=M071R73

 $\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0230±0.0017 OUR FIT			
0.0228±0.0018 OUR AVERAGE			
0.0230±0.0020±0.0012	¹ AAIJ	16Y LHCb	$\Lambda_b^0 \rightarrow \psi(2S)X$
0.0216±0.0026±0.0014	² AUBERT	02B BABR	e^+e^-
0.0327±0.0077±0.0072	² GRIBUSHIN	96 FMPS	$515 \pi^- Be \rightarrow 2\mu X$

¹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$.

² Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$.

 Γ_7/Γ_{11}

NODE=M071R63
NODE=M071R63

 $\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.8 ± 1.1 OUR FIT			
8.73±1.39±1.57	BAI	02 BES	e^+e^-

NODE=M071R76
NODE=M071R76

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.5645±0.0026 OUR FIT				
0.554 ± 0.008 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.5604±0.0009±0.0062	565k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.525 ± 0.009 ± 0.022	4k	ANDREOTTI	05 E835	$\psi(2S) \rightarrow J/\psi X$
0.536 ± 0.007 ± 0.016	20k	^{1,2} ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
0.496 ± 0.037		ARMSTRONG	97 E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


0.5637±0.0027±0.0046 60k ADAM 05A CLEO Repl. by MENDEZ 08

NODE=M071R70
NODE=M071R70

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$.

NODE=M071R;LINKAGE=AB
NODE=M071R;LINKAGE=AL



$$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{10}/\Gamma_{11} = (0.9761\Gamma_{12} + 0.719\Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})/\Gamma_{11}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.732±0.008 OUR FIT				
0.73 ±0.09		TANENBAUM 76	MRK1	$e^+ e^-$

NODE=M071R11
NODE=M071R11

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_{12}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1824±0.0031 OUR FIT				

NODE=M071R17
NODE=M071R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1769±0.0008±0.0053	61k	1 MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.1652±0.0014±0.0058	13.4k	2 ADAM	05A	CLEO	Repl. by MENDEZ 08

1 Not independent from other measurements of MENDEZ 08.

2 Not independent from other values reported by ADAM 05A.

NODE=M071R17;LINKAGE=ME
NODE=M071R17;LINKAGE=AD

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$$

$$\Gamma_{12}/\Gamma_9$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2968±0.0031 OUR FIT				
0.320 ±0.012 OUR AVERAGE				

NODE=M071R69
NODE=M071R69

0.300 ± 0.008 ± 0.022	1655 ± 44	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.328 ± 0.013 ± 0.008		AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2829±0.0012±0.0056	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.2776±0.0025±0.0043	13.4k	ADAM	05A	CLEO	Repl. by MENDEZ 08

$$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{12}/\Gamma_{11}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.526 ±0.008 OUR FIT				
0.513 ±0.022 OUR AVERAGE				Error includes scale factor of 2.2.

NODE=M071R14
NODE=M071R14

0.5047±0.0022±0.0102	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.570 ± 0.009 ± 0.026	14k	1 ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4924±0.0047±0.0086	73k	2,3 ADAM	05A	CLEO	Repl. by MENDEZ 08
0.571 ± 0.018 ± 0.044		4 ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.53 ± 0.06		TANENBAUM	76	MRK1	$e^+ e^-$
0.64 ± 0.15		5 HILGER	75	SPEC	$e^+ e^-$

1 From a fit to the J/ψ recoil mass spectra.

2 Not independent from other values reported by ADAM 05A.

3 Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

4 Not independent from other values reported by ANDREOTTI 05.

5 Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

NODE=M071R14;LINKAGE=AB
NODE=M071R14;LINKAGE=AD
NODE=M071R14;LINKAGE=AM
NODE=M071R;LINKAGE=AN
NODE=M071R;LINKAGE=I

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_{13}/Γ
<u>VALUE (units 10^{-3})</u>				
33.7 ± 0.5 OUR FIT				
32.9 ± 1.7 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
33.75 ± 0.17 ± 0.86	68.2k	ABLIKIM	12M BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
29.8 ± 0.9 ± 2.3	5.7k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
25.5 ± 2.9	386	OREGLIA	80 CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
45 ± 12	17	BRANDELIK	79B DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
42 ± 6	164	BARTEL	78B CNTR	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
34.3 ± 0.4 ± 0.9	18.4k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
32.5 ± 0.6 ± 1.1	2.8k	ADAM	05A CLEO	Repl. by MENDEZ 08
43 ± 8	44	TANENBAUM	76 MRK1	$e^+ e^-$


¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

³ Not independent from other measurements of MENDEZ 08.

⁴ Not independent from other values reported by ADAM 05A.

NODE=M071R15
NODE=M071R15




$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$	EVTS	DOCUMENT ID	TECN	Γ_{13}/Γ
<u>VALUE</u>				
0.0549 ± 0.0008 OUR FIT				
0.058 ± 0.007 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.050 ± 0.006 ± 0.003	298 ± 20	ANDREOTTI 05 E835	$\psi(2S) \rightarrow J/\psi X$	
0.072 ± 0.009		AMBROGIANI 00A E835	$p\bar{p} \rightarrow \psi(2S)$	OCCUR=2
0.061 ± 0.015		ARMSTRONG 97 E760	$\bar{p}p \rightarrow \psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0549 ± 0.0006 ± 0.0009	18.4k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.0546 ± 0.0010 ± 0.0007	2.8k	ADAM	05A CLEO	Repl. by MENDEZ 08

NODE=M071R68
NODE=M071R68

¹ Not independent from other measurements of MENDEZ 08.

NODE=M071R68;LINKAGE=ME



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.0972±0.0014 OUR FIT

0.0979±0.0018 OUR AVERAGE

0.0979±0.0010±0.0015	18.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.098 ± 0.005 ± 0.010	2k	1 ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ± 0.021		2 HIMEL	80	MRK2	$e^+ e^- \rightarrow \psi(2S)X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0968±0.0019±0.0013	2.8k	3 ADAM	05A	CLEO	Repl. by MENDEZ 08
0.095 ± 0.007 ± 0.007		4 ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$

1 From a fit to the J/ψ recoil mass spectra.

2 The value for $B(\psi(2S) \rightarrow J/\psi(1s)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S)) \rightarrow J/\psi(1S)\pi^+\pi^- = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (0.1181 \pm 0.0020)$.

3 Not independent from other values reported by ADAM 05A.

4 Not independent from other values reported by ANDREOTTI 05.

Γ_{13}/Γ_{11}

NODE=M071R71
NODE=M071R71

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

12.68±0.32 OUR AVERAGE

12.6 ± 0.2 ± 0.3	4.1k	ABLIKIM	12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
13.3 ± 0.8 ± 0.3	530	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
14.3 ± 1.4 ± 1.2	280	BAI	04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
14 ± 6	7	HIMEL	80	MRK2	$e^+ e^-$
9 ± 2 ± 1	23	1 OREGLIA	80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

13 ± 1 ± 1 88 ADAM 05A CLEO Repl. by MENDEZ 08

1 Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

NODE=M071R71;LINKAGE=AB
NODE=M071R;LINKAGE=8H

NODE=M071R71;LINKAGE=AD
NODE=M071R71;LINKAGE=AN

NODE=M071R16
NODE=M071R16

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$

$$\Gamma_{14}/\Gamma_9 = \Gamma_{14}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13} + 0.343\Gamma_{154} + 0.190\Gamma_{155})$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.213±0.012±0.003	527	1 MENDEZ	08	CLEO	$e^+ e^- \rightarrow J/\psi\gamma\gamma$
0.22 ± 0.02 ± 0.01		2 ADAM	05A	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$

1 Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

2 Not independent from other values reported by ADAM 05A.

NODE=M071R16;LINKAGE=3Q

NODE=M071S25
NODE=M071S25

NODE=M071S25;LINKAGE=ME
NODE=M071S25;LINKAGE=AD

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ_{11}
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.380 \pm 0.022 \pm 0.005	527	1 MENDEZ	08 CLEO	$e^+ e^- \rightarrow J/\psi\gamma\gamma$	
0.39 \pm 0.04 \pm 0.01		2 ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$	

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

— HADRONIC DECAYS —

$\Gamma(\pi^0 h_c(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
8.6 \pm 1.3 OUR AVERAGE					
9.0 \pm 1.5 \pm 1.3	3k	1 GE	11 CLEO	$\psi(2S) \rightarrow \pi^0$ anything	
8.4 \pm 1.3 \pm 1.0	11k	ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 h_c$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	92^{+23}_{-22}	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$	
seen	1282	DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$	
seen	168 ± 40	ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$	

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86$ MeV $\equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P)/\Gamma_0) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
35 \pm 16	6	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons	

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
29 \pm 10 OUR AVERAGE				Error includes scale factor of 4.7. See the ideogram below.	
24.9 \pm 0.7 \pm 3.6	2173	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$	
127 \pm 12 \pm 2	410	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+\pi^-)\pi^0 \gamma$	
26.1 \pm 0.7 \pm 3.0	1703	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$	
30 \pm 8	42	FRANKLIN	83 MRK2	$e^+ e^-$	

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
29 \pm 10 (Error scaled by 4.7)



NODE=M071S26
NODE=M071S26

NODE=M071S26;LINKAGE=ME
NODE=M071S26;LINKAGE=AD

NODE=M071310

NODE=M071S42
NODE=M071S42

NODE=M071S42;LINKAGE=GE

NODE=M071R37
NODE=M071R37

NODE=M071R22
NODE=M071R22

NODE=M071R22;LINKAGE=UB

$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{18}/Γ
$2.55 \pm 0.73 \pm 0.47$		112 ± 31	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
<2.3		90	BAI	98J BES	$e^+ e^-$	

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{21}/Γ
2.94 ± 0.08 OUR FIT					
3.02 ± 0.08 OUR AVERAGE					
3.05 $\pm 0.02 \pm 0.12$	19k	ABLIKIM	18T BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
3.08 $\pm 0.05 \pm 0.18$	4.5k	¹ DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
3.36 $\pm 0.09 \pm 0.25$	1.6k	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
2.87 $\pm 0.12 \pm 0.15$	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
1.4 ± 0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	
2.3 ± 0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 $\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{21}/Γ_{11}
8.49 ± 0.23 OUR FIT					
$6.98 \pm 0.49 \pm 0.97$		BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$	

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{22}/Γ
$3.06 \pm 0.06 \pm 0.14$	6k	ABLIKIM	18T BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$	

 $\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{23}/Γ
$12.8 \pm 1.0 \pm 3.4$	157	¹ BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons	

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{24}/Γ
< 0.29	90	¹ ABLIKIM	13F BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$		
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$						
<12	90	² ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$		

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{25}/Γ
$2.48 \pm 0.34 \pm 0.19$	60	¹ ABLIKIM	13F BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$		

$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$

<4.9	90	² ABLIKIM	07H BES2	$e^+ e^- \rightarrow \psi(2S)$
------	----	----------------------	----------	--------------------------------

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

 $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{26}/Γ
$1.0 \pm 0.1 \pm 0.1$	74.0	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^-$	

 $\Gamma(K^*(892)^+\bar{p}\Lambda+c.c.)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$6.3 \pm 0.5 \pm 0.5$	1011	ABLIKIM	19AU BES3

 Γ_{27}/Γ

NODE=M071P25
NODE=M071P25

 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{28}/Γ
$1.8 \pm 0.3 \pm 0.3$	45.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$	

NODE=M071S19
NODE=M071S19

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 0.4 \pm 0.5$	73.4	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$

 Γ_{29}/Γ

NODE=M071S17
NODE=M071S17

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.81 ± 0.13 OUR AVERAGE					Error includes scale factor of 1.4. See the ideogram below.

$3.97 \pm 0.02 \pm 0.12$	31k	ABLIKIM	17L	BES3	$e^+ e^- \rightarrow \Lambda\bar{\Lambda}$
$3.71 \pm 0.05 \pm 0.15$	6.5k	¹ DOBBS	17		$e^+ e^- \rightarrow \Lambda\bar{\Lambda}$
$3.39 \pm 0.20 \pm 0.32$	337	ABLIKIM	07C	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$6.4 \pm 1.8 \pm 0.1$		² AUBERT	07BD	BABR	$10.6 e^+ e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
$3.28 \pm 0.23 \pm 0.25$	208	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.75 \pm 0.09 \pm 0.23$	1.9k	^{1,3} DOBBS	14		$e^+ e^- \rightarrow \Lambda\bar{\Lambda}$
$1.81 \pm 0.20 \pm 0.27$	80	⁴ BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 4	90	FELDMAN	77	MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$


¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 Superseded by DOBBS 17.

4 Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.

WEIGHTED AVERAGE
 3.81 ± 0.13 (Error scaled by 1.4)

 $\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.40 \pm 0.03 \pm 0.13$	2.8k	ABLIKIM	13W	BES3 $\psi(2S) \rightarrow \text{hadrons}$

 Γ_{31}/Γ

NODE=M071S65
NODE=M071S65

 $\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.54 \pm 0.04 \pm 0.13$	2.8k	ABLIKIM	13W	BES3 $\psi(2S) \rightarrow \text{hadrons}$

 Γ_{32}/Γ

NODE=M071S66
NODE=M071S66

 $\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
$1.23 \pm 0.23 \pm 0.08$	30	¹ DOBBS	17 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{33}/Γ

NODE=M071P08
NODE=M071P08

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071P08;LINKAGE=A

$\Gamma(\Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.67 \pm 0.13 \pm 0.12$	276	¹ ABLIKIM	13D	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$

1 Using $B(\Lambda \rightarrow p \pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda \gamma) = 100\%$.

 $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.32 ± 0.12 OUR AVERAGE				

$2.31 \pm 0.06 \pm 0.10$	1.9k	¹ DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$2.57 \pm 0.44 \pm 0.68$	35	PEDLAR	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.51 \pm 0.15 \pm 0.16$	281	^{1,2} DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
--------------------------	-----	----------------------	----	---

1 Using CLEO-c data but not authored by the CLEO Collaboration.

2 Superseded by DOBBS 17.

 Γ_{34}/Γ

NODE=M071S63
NODE=M071S63

NODE=M071S63;LINKAGE=AB

 $\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.1.

$2.44 \pm 0.03 \pm 0.11$	7k	ABLIKIM	17L	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$2.22 \pm 0.05 \pm 0.11$	2.6k	¹ DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

$2.35 \pm 0.36 \pm 0.32$	59	ABLIKIM	07C	BES $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$2.63 \pm 0.35 \pm 0.21$	58	PEDLAR	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.25 \pm 0.11 \pm 0.16$	439	^{1,2} DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$1.2 \pm 0.4 \pm 0.4$	8	³ BAI	01	BES $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

1 Using CLEO-c data but not authored by the CLEO Collaboration.

2 Superseded by DOBBS 17.

3 Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

NODE=M071R51
NODE=M071R51

NODE=M071R47;LINKAGE=A
NODE=M071R47;LINKAGE=B

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5 ± 0.7 OUR AVERAGE				

$8.4 \pm 0.5 \pm 0.5$	1.5k	ABLIKIM	16L	BES3 $\psi(2S) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
$11 \pm 3 \pm 3$	14	¹ BAI	01	BES $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

1 Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

NODE=M071R52
NODE=M071R52

OCCUR=2

NODE=M071R52;LINKAGE=PP

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.5 \pm 0.6 \pm 0.6$	1.4K	ABLIKIM	16L	BES3 $\psi(2S) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$

NODE=M071R00
NODE=M071R00

 $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.69 \pm 0.05 \pm 0.05$	2.2k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071P00
NODE=M071P00

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.87 ± 0.11 OUR AVERAGE					Error includes scale factor of 1.1.

$3.03 \pm 0.05 \pm 0.14$	3.6k	¹ DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$2.78 \pm 0.05 \pm 0.14$	5k	ABLIKIM	16L	BES3 $\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$

$3.03 \pm 0.40 \pm 0.32$	67	ABLIKIM	07C	BES $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$2.38 \pm 0.30 \pm 0.21$	63	PEDLAR	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.66 \pm 0.12 \pm 0.20$	548	^{1,2} DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
$0.94 \pm 0.27 \pm 0.15$	12	³ BAI	01	BES $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

<2	90	FELDMAN	77	MRK1 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
------	----	---------	----	--

1 Using CLEO-c data but not authored by the CLEO Collaboration.

2 Superseded by DOBBS 17.

3 Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

NODE=M071R29
NODE=M071R29

NODE=M071R29;LINKAGE=A
NODE=M071R29;LINKAGE=B

NODE=M071R29;LINKAGE=PP

$\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{41}/Γ
2.3 ± 0.4 OUR AVERAGE Error includes scale factor of 4.2.					
2.73 ± 0.03 ± 0.13	11k	ABLIKIM	17E	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
1.97 ± 0.06 ± 0.11	1.2k	1 DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
2.75 ± 0.64 ± 0.61	19	PEDLAR	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.02 ± 0.19 ± 0.15	112	1.2 DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17. Γ_{41}/Γ NODE=M071R48
NODE=M071R48 $\Gamma(\Xi(1530)^0 \bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{42}/Γ
5.2 ± 0.3 ± 3.2		527	1 ABLIKIM	13S	BES3 $\psi(2S) \rightarrow \eta p\bar{p}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<32	90	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
< 8.1	90	2 BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

¹ With $N(1535)$ decaying to $p\eta$.² Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. Γ_{42}/Γ NODE=M071R48;LINKAGE=A
NODE=M071R48;LINKAGE=B $\Gamma(\Xi(1530)^- \bar{\Xi}(1530)^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{44}/Γ
11.45 ± 0.40 ± 0.59	5067	ABLIKIM	19AT	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

NODE=M071R53;LINKAGE=A

NODE=M071R53;LINKAGE=PP

 $\Gamma(\Xi(1530)^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{45}/Γ
0.70 ± 0.11 ± 0.04	399	ABLIKIM	19AT	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

NODE=M071P27
NODE=M071P27 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{43}/Γ
3.86 ± 0.27 ± 0.32	236	ABLIKIM	15I	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	

NODE=M071S82
NODE=M071S82 $\Gamma(\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{46}/Γ
5.21 ± 1.48 ± 0.57	74	ABLIKIM	15I	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	

NODE=M071S83
NODE=M071S83 $\Gamma(\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{47}/Γ
12.03 ± 2.94 ± 1.22	136	ABLIKIM	15I	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	

NODE=M071S84
NODE=M071S84 $\Gamma(K^- \Sigma^0 \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{48}/Γ
3.67 ± 0.33 ± 0.28	142	ABLIKIM	15I	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \bar{\Xi}^+ + \text{c.c.}$	

NODE=M071S85
NODE=M071S85 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{49}/Γ
0.52 ± 0.03 ± 0.03		326	1 DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

NODE=M071R54
NODE=M071R54

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.47 ± 0.09 ± 0.05	27	1,2 DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
<1.5	90	ABLIKIM	12Q	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
<1.6	90	PEDLAR	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	
<0.73	90	3 BAI	01	BES $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.
² Superseded by DOBBS 17.
³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

$\Gamma(\pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.53±0.07 OUR AVERAGE				
1.65±0.03±0.15	4.5k	ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$
1.54±0.06±0.06	948	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \pi^0 p\bar{p}$
1.32±0.10±0.15	256	¹ ABLIKIM	05E	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
1.4 ±0.5	9	FRANKLIN	83	MRK2 $e^+ e^-$

¹ Computed using $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$.

 Γ_{50}/Γ

NODE=M071R35
NODE=M071R35

 $\Gamma(N(940)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.42±0.20^{+1.78}_{-1.28}	1.9k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{51}/Γ

NODE=M071S56
NODE=M071S56

 $\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.3^{+1.7}_{-1.5} OUR AVERAGE Error includes scale factor of 2.5.

3.58±0.25 ^{+1.59} _{-0.84}	1.1k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$
8.1 ±0.7 ±0.3	474	² ALEXANDER	10	CLEO $\psi(2S) \rightarrow \pi^0 p\bar{p}$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

² From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

 Γ_{52}/Γ

NODE=M071S50
NODE=M071S50

 $\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.64±0.05^{+0.22}_{-0.17}	0.2k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{53}/Γ

NODE=M071S57
NODE=M071S57

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.47±0.28^{+0.99}_{-0.97}	0.7k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{54}/Γ

NODE=M071S58
NODE=M071S58

 $\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.76±0.28^{+1.37}_{-1.66}	1.1k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{55}/Γ

NODE=M071S59
NODE=M071S59

 $\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.79±0.10^{+0.24}_{-0.71}	0.5k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{56}/Γ

NODE=M071S60
NODE=M071S60

 $\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.62±0.28^{+1.12}_{-0.64}	0.9k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{57}/Γ

NODE=M071S61
NODE=M071S61

 $\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.13±0.08^{+0.40}_{-0.30}	0.8k	¹ ABLIKIM	13A	BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{58}/Γ

NODE=M071S62
NODE=M071S62

NODE=M071R35;LINKAGE=AB

NODE=M071S56

NODE=M071S56;LINKAGE=AB

NODE=M071S50

NODE=M071S59;LINKAGE=AB

NODE=M071S60

NODE=M071S60;LINKAGE=AB

NODE=M071S61

NODE=M071S62

NODE=M071S62;LINKAGE=AB

$\Gamma(\pi^0 f_0(2100) \rightarrow \pi^0 p\bar{p})/\Gamma_{\text{total}}$	Γ_{59}/Γ
<i>VALUE (units 10^{-5})</i>	<i>EVTS</i>
$1.1 \pm 0.4 \pm 0.1$	76

¹ From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N_1^*(1440)\bar{p}$, $\pi^0 f_0(2100)$, and two other broad, unestablished resonances.

NODE=M071S51
NODE=M071S51

NODE=M071S51;LINKAGE=AL

$\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$	Γ_{60}/Γ
<i>VALUE (units 10^{-5})</i>	<i>EVTS</i>
6.0 ± 0.4 OUR AVERAGE	
6.4 $\pm 0.2 \pm 0.6$	679
5.6 $\pm 0.6 \pm 0.3$	154
5.8 $\pm 1.1 \pm 0.7$	44.8 ± 8.5
8 $\pm 3 \pm 3$	9.8

¹ With $N(1535)$ decaying to $p\eta$.
² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.

NODE=M071R56
NODE=M071R56NODE=M071R56;LINKAGE=A
NODE=M071R56;LINKAGE=AB

$\Gamma(\eta f_0(2100) \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$	Γ_{61}/Γ
<i>VALUE (units 10^{-5})</i>	<i>EVTS</i>
$1.2 \pm 0.4 \pm 0.1$	31

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

NODE=M071S52
NODE=M071S52

NODE=M071S52;LINKAGE=AL

$\Gamma(N(1535)\bar{p} \rightarrow \eta p\bar{p})/\Gamma_{\text{total}}$	Γ_{62}/Γ
<i>VALUE (units 10^{-5})</i>	<i>EVTS</i>
$4.4 \pm 0.6 \pm 0.3$	123

¹ From a fit of the $p\bar{p}$ and $p\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and $\eta f_0(2100)$.

NODE=M071S53
NODE=M071S53

NODE=M071S53;LINKAGE=AL

$\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$	Γ_{63}/Γ
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>
0.69 ± 0.21 OUR AVERAGE	
0.6 $\pm 0.2 \pm 0.2$	21.2
0.8 $\pm 0.3 \pm 0.1$	14.9 ± 0.1

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R79
NODE=M071R79

NODE=M071R;LINKAGE=B3

$\Gamma(\eta' p\bar{p})/\Gamma_{\text{total}}$	Γ_{64}/Γ
<i>VALUE (units 10^{-5})</i>	<i>EVTS</i>
$1.10 \pm 0.10 \pm 0.08$	491

¹ From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

NODE=M071P20
NODE=M071P20

NODE=M071P20;LINKAGE=A

$\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$	Γ_{65}/Γ
<i>VALUE (units 10^{-6})</i>	<i>CL%</i>
$6.06 \pm 0.38 \pm 0.48$	753

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24	90	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
<26	90	¹ BAI	03B	BES	$\psi(2S) \rightarrow K^+K^-p\bar{p}$

NODE=M071R82
NODE=M071R82

NODE=M071R82;LINKAGE=B3

$\Gamma(\phi X(1835) \rightarrow \phi p\bar{p})/\Gamma_{\text{total}}$	Γ_{66}/Γ
<i>VALUE</i>	<i>CL%</i>
$<1.82 \times 10^{-7}$	90

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071P21
NODE=M071P21

$\Gamma(\pi^+ \pi^- p\bar{p})/\Gamma_{\text{total}}$	Γ_{67}/Γ
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>
6.0 ± 0.4 OUR AVERAGE	
5.9 $\pm 0.2 \pm 0.4$	904.5
8 ± 2	¹ TANENBAUM 78

¹ Assuming entirely strong decay.

NODE=M071R31
NODE=M071R31

NODE=M071R;LINKAGE=K

$\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**2.48±0.17 OUR AVERAGE** $2.45 \pm 0.11 \pm 0.21$ $2.52 \pm 0.12 \pm 0.22$

DOCUMENT ID TECN COMMENT

ABLIKIM 06I BES2 $e^+ e^- \rightarrow p\pi^- X$ ABLIKIM 06I BES2 $e^+ e^- \rightarrow \bar{p}\pi^+ X$ Γ_{68}/Γ

NODE=M071R01

NODE=M071R01

OCCUR=2

 $\Gamma(p\bar{n}\pi^- \pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**3.18±0.50±0.50** 135 ± 21

DOCUMENT ID TECN COMMENT

ABLIKIM 06I BES2 $e^+ e^- \rightarrow p\pi^- \pi^0 X$ Γ_{69}/Γ

NODE=M071R02

NODE=M071R02

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) CL%**<1.6**

90

DOCUMENT ID TECN COMMENT

BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ Γ_{71}/Γ

NODE=M071S06

NODE=M071S06

 $\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**9.5±0.7±1.5**1 BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.3±0.8±1.4 201.7 2 BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi (\eta \rightarrow \gamma\gamma)$ 8.1±1.4±1.6 50.0 2 BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi (\eta \rightarrow 3\pi)$ Γ_{72}/Γ

NODE=M071S07

NODE=M071S07

OCCUR=2

OCCUR=3

1 Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.

2 Not independent from other values reported by BRIERE 05.

 $\Gamma(2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS**1.2±0.6±0.1**

16

DOCUMENT ID TECN COMMENT

1 AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow 2(\pi^+\pi^-)\eta\gamma$ 1 AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+\pi^-)\eta) \cdot B(\eta \rightarrow \gamma\gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$. Γ_{73}/Γ

NODE=M071S38

NODE=M071S38

NODE=M071S38;LINKAGE=BR

NODE=M071S07;LINKAGE=BI

 $\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**4.5±1.6±1.3**

12.8

DOCUMENT ID TECN COMMENT

BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$ Γ_{75}/Γ

NODE=M071S08

NODE=M071S08

 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**7.3±1.2 OUR AVERAGE**

Error includes scale factor of 2.1. See the ideogram below.


8.4±0.5±1.2 386 ABLIKIM 07D BES2 $e^+ e^- \rightarrow \psi(2S)$ 12.2±2.2±0.7 37 1 AUBERT 07AU BABR 10.6 $e^+ e^- \rightarrow \omega\pi^+\pi^-\gamma$ 8.2±0.5±0.7 391 BRIERE 05 CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ 4.8±0.6±0.7 100 ± 22 2 BAI 03B BES $\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$ 1 AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16 \text{ eV}$.2 Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. Γ_{76}/Γ

NODE=M071R77

NODE=M071R77

NODE=M071R77;LINKAGE=UB

NODE=M071R77;LINKAGE=B3

 $\Gamma(b_1^\pm\pi^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.1.
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
4.18 ^{+0.43} _{-0.42} ± 0.92	170	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	1,2 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.2 ± 0.8 ± 1.0	1 BAI		99C BES	Repl. by BAI 03B

¹ Assuming $B(b_1 \rightarrow \omega\pi) = 1$.² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. Γ_{77}/Γ NODE=M071R40
NODE=M071R40 $\Gamma(b_1^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35^{+0.47}_{-0.42} ± 0.40	45	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

 Γ_{78}/Γ

NODE=M071R;LINKAGE=M1

NODE=M071R40;LINKAGE=B3

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{79}/Γ NODE=M071R21
NODE=M071R21

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.4 OUR AVERAGE					
2.3 ± 0.5 ± 0.4	57	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$	
2.05 ± 0.41 ± 0.38	62 ± 12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.5	90	1 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$	
<1.7	90	BAI	98J BES	Repl. by BAI 03B	

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$. $\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{82}/Γ NODE=M071R24
NODE=M071R24

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±0.5 OUR AVERAGE				
8.1±1.3±0.3	133	LEES	12F BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^- K^+K^-\gamma$
7.1±0.3±0.4	817.2	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
16 ± 4	1 TANENBAUM	78 MRK1	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
11.0±1.9±0.2	85	² AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^- K^+K^-\gamma$

¹ Assuming entirely strong decay.² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R24;LINKAGE=K

NODE=M071R24;LINKAGE=BE

$\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$					Γ_{84}/Γ	NODE=M071S09 NODE=M071S09
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
2.2±0.2±0.4	223.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$		
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$					Γ_{85}/Γ	NODE=M071R66 NODE=M071R66
<i>VALUE (units 10^{-4})</i>	<i>CL%</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>	
1.86±0.32±0.43	93 ± 16	BAI	04C	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.2	90	BAI	98J	BES $e^+ e^-$		
$\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$					Γ_{86}/Γ	NODE=M071S39 NODE=M071S39
<i>VALUE (units 10^{-3})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
1.3±0.7±0.1	7	1 AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$		
1 AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^- \eta)) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV.}$						
$\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$					Γ_{87}/Γ	NODE=M071R09 NODE=M071R09
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
10.0±2.5±1.8	65	ABLIKIM	07D	BES2 $e^+ e^- \rightarrow \psi(2S)$		
$\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$					Γ_{88}/Γ	NODE=M071R41 NODE=M071R41
<i>VALUE (units 10^{-4})</i>		<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
10.0±1.8±2.1	1	BAI	99C	BES $e^+ e^-$		
1 Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$						NODE=M071R;LINKAGE=M2
$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{90}/Γ	NODE=M071R49 NODE=M071R49
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
2.20±0.25±0.37	83 ± 9	ABLIKIM	050	BES2 $e^+ e^- \rightarrow \psi(2S)$		
$\Gamma(\rho^0 p \bar{p})/\Gamma_{\text{total}}$					Γ_{91}/Γ	NODE=M071S14 NODE=M071S14
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
0.5±0.1±0.2	61.1	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p \bar{p} \pi^+ \pi^-$		
$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{92}/Γ	NODE=M071R34 NODE=M071R34
<i>VALUE (units 10^{-4})</i>		<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
6.7±2.5		TANENBAUM	78	MRK1 $e^+ e^-$		
$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$					Γ_{93}/Γ	NODE=M071R27 NODE=M071R27
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
2.4±0.6 OUR AVERAGE	Error includes scale factor of 2.2.					
2.2±0.2±0.2	308	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$		
4.5±1.0		TANENBAUM	78	MRK1 $e^+ e^-$		
$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{94}/Γ	NODE=M071R33 NODE=M071R33
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
2.2±0.6 OUR AVERAGE	Error includes scale factor of 1.4.					
2.0±0.2±0.4	285.5	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-)$		
4.2±1.5		TANENBAUM	78	MRK1 $e^+ e^-$		
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					Γ_{95}/Γ	NODE=M071S10 NODE=M071S10
<i>VALUE (units 10^{-4})</i>	<i>EVTS</i>	<i>DOCUMENT ID</i>	<i>TECN</i>	<i>COMMENT</i>		
12.6±0.9 OUR AVERAGE	32	1 AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$		
18.9±5.7±0.3						
11.7±1.0±1.5	597	ABLIKIM	06G	BES2 $\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
12.7±0.5±1.0	711.6	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
1 AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.						NODE=M071S10;LINKAGE=UB

$\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$					Γ_{96}/Γ	NODE=M071S20 NODE=M071S20
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
5.9±2.0±0.9	19	ABLIKIM	06G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
$\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{97}/Γ	NODE=M071S21 NODE=M071S21
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
8.6±1.3±1.8	238	ABLIKIM	06G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
$\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{98}/Γ	NODE=M071S22 NODE=M071S22
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
9.6±2.2±1.7	133	ABLIKIM	06G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{99}/Γ	NODE=M071S23 NODE=M071S23
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
7.3±2.2±1.4	78	ABLIKIM	06G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
$\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{100}/Γ	NODE=M071S24 NODE=M071S24
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
6.1±1.3±1.2	125	ABLIKIM	06G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
$\Gamma(\eta K^+ K^-, \text{no } \eta\phi)/\Gamma_{\text{total}}$					Γ_{101}/Γ	NODE=M071S11 NODE=M071S11
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.08±0.29±0.25	0.3k	1	ABLIKIM	12L	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<13	90	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
1 Excluding $\eta\phi$.						
$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$					Γ_{102}/Γ	NODE=M071S11;LINKAGE=AB
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.62±0.11 OUR AVERAGE	Error includes scale factor of 1.1.					
1.56±0.04±0.11	2.8k	ABLIKIM	14G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
2.38±0.37±0.29	78	ABLIKIM	06G	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$		
1.9 ± 0.3 ± 0.3	76.8	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
1.5 ± 0.3 ± 0.2	23	1 BAI	03B	BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
1 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.						
$\Gamma(\omega K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{103}/Γ	NODE=M071R78 NODE=M071R78
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
20.7±2.6 OUR AVERAGE						
18.9±2.9±2.2	396	ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	
22.6±3.0±2.4	535	ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$	OCCUR=2
$\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{104}/Γ	NODE=M071S68 NODE=M071S68
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
6.1 ± 1.2 OUR AVERAGE						
6.39±1.50±0.78	128	ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	
5.86±1.61±0.83	143	ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$	OCCUR=2
$\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$					Γ_{105}/Γ	NODE=M071S69 NODE=M071S69
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
16.8±2.5±1.6	356	ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	
$\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$					Γ_{106}/Γ	NODE=M071S70 NODE=M071S70
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
5.82±2.08±0.72	116	ABLIKIM	13M	BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	

$\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{107}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.60±0.27±0.24	109	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	
1 $X(1440)$ compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.					

NODE=M071S71
NODE=M071S71

$\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{108}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.09±0.20±0.16	82	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$	
1 $X(1440)$ compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.					

NODE=M071S71;LINKAGE=AB

$\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{109}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.302±0.098±0.027	22	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	
1 Statistical significance 4.5 σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.					

NODE=M071S73
NODE=M071S73

$\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{110}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.125±0.070±0.013	10	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$	
1 Statistical significance 3.2 σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.					

NODE=M071S73;LINKAGE=AB

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$					Γ_{111}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.5 ± 2.0 OUR AVERAGE				Error includes scale factor of 2.8.	
5.45±0.42±0.87 671 ABLIKIM 05H BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+ \pi^-)$					
1.5 ± 1.0		¹ TANENBAUM 78 MRK1		$e^+ e^-$	

NODE=M071R32
NODE=M071R32

1 Assuming entirely strong decay.

$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{112}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
7.3±0.4±0.6	434.9	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$	

NODE=M071R32;LINKAGE=K

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$					Γ_{113}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL %</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.48±0.23±0.39	1.3k	¹ METREVELI	12	$\psi(2S) \rightarrow K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

NODE=M071R23
NODE=M071R23

6.2 ± 1.5 ± 0.2	66	2,3 LEES	15J BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$	
8.3 ± 1.5 ± 0.2	66	3,4 LEES	15J BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$	
6.3 ± 0.6 ± 0.3		5 DOBBS	06A CLEO	$e^+ e^-$	
10 ± 7		5 BRANDELIK	79C DASP	$e^+ e^-$	
< 5	90	FELDMAN	77 MRK1	$e^+ e^-$	

1 Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
2 $\sin\phi > 0$.
3 Using $\Gamma(\psi(2S) \rightarrow e^+ e^-) = (2.37 \pm 0.04)$ keV.
4 $\sin\phi < 0$.
5 Interference with non-resonant $K^+ K^-$ production not taken into account.

OCCUR=2

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$					Γ_{114}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
5.34±0.33 OUR AVERAGE		¹ METREVELI	12	$\psi(2S) \rightarrow K_S^0 K_L^0$	
5.28±0.25±0.34 478 ± 23 DOBBS 06A CLEO $e^+ e^-$					
5.8 ± 0.8 ± 0.4		² BAI	04B BES2	$\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$	
5.24±0.47±0.48 156 ± 14					

NODE=M071R87
NODE=M071R87

1 Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

2 Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.NODE=M071R87;LINKAGE=ME
NODE=M071R;LINKAGE=KZ

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.01±0.17 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.			
2.14±0.03±0.12	7k	1 ABLIKIM	12H BES3	$e^+e^- \rightarrow \psi(2S)$
1.81±0.18±0.19	260 ± 19	2 ABLIKIM	05J BES2	$e^+e^- \rightarrow \psi(2S)$
1.88±0.16±0.28	194	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
0.85±0.46	4	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.

² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

 Γ_{115}/Γ

NODE=M071R36
NODE=M071R36

 $\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.94±0.25±1.15	1 ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

NODE=M071R57
NODE=M071R57

 $\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.32±0.12 OUR AVERAGE	Error includes scale factor of 1.8.				
0.51±0.07±0.11		1 ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$	

0.24±0.08±0.02	22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
----------------	----	------	---------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1 FRANKLIN	83 MRK2	e^+e^-
<10	90	BARTEL	76 CNTR	e^+e^-
<10	90	2 ABRAMS	75 MRK1	e^+e^-

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

² Final state $\rho^0\pi^0$.

NODE=M071R57;LINKAGE=AK

NODE=M071R26
NODE=M071R26

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.78±0.26 OUR AVERAGE	Error includes scale factor of 1.8.				
0.76±0.25±0.06	30	1 METREVELI	12 BRANDELIK	$\psi(2S) \rightarrow \pi^+\pi^-$	

¹ METREVELI 12 BRANDELIK 79C DASP e^+e^-

8 ± 5

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.1	90	DOBBS	06A CLEO	$e^+e^- \rightarrow \psi(2S)$
<5	90	FELDMAN	77 MRK1	e^+e^-

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+\pi^-$ for continuum subtraction.

NODE=M071R26;LINKAGE=AK
NODE=M071R;LINKAGE=N

NODE=M071R20
NODE=M071R20

NODE=M071R20;LINKAGE=ME

$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	1 BAI	99C BES	$e^+ e^-$

1 Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$ Γ_{119}/Γ NODE=M071R45
NODE=M071R45 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.12 \pm 0.62^{+1.13}_{-0.61}$	251 ± 22	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{120}/Γ NODE=M071S54
NODE=M071S54 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.07 \pm 0.16 \pm 0.26$	0.9k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{121}/Γ NODE=M071R38
NODE=M071R38 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR AVERAGE					Error includes scale factor of 1.2.
$3.18 \pm 0.30^{+0.26}_{-0.31}$	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$	
$2.9^{+1.3}_{-1.7} \pm 0.4$	9.6 ± 4.2	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$	
$1.3^{+1.0}_{-0.7} \pm 0.3$	7	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{124}/Γ NODE=M071R39
NODE=M071R39 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
10.9 ± 2.0 OUR AVERAGE				
$13.3^{+2.4}_{-2.8} \pm 1.7$	65.6 ± 9.0	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
$9.2^{+2.7}_{-2.2} \pm 0.9$	25	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{125}/Γ NODE=M071R30
NODE=M071R30 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.06 OUR AVERAGE			
0.22 ± 0.10	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
0.14 ± 0.08	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$


 $\Gamma_{124}/\Gamma_{125}$ NODE=M071R46
NODE=M071R46 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.18 ± 0.26 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
2.3 $\pm 0.8 \pm 0.1$	19 ± 6	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.9 $\pm 0.2 \pm 0.1$	47.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
1.5 $\pm 0.2 \pm 0.2$	51.5 ± 8.3	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Γ_{126}/Γ NODE=M071R80
NODE=M071R801 Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.2 Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.3 Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.NODE=M071R80;LINKAGE=B3
NODE=M071R80;LINKAGE=BE

NODE=M071R80;LINKAGE=UB



$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75±0.33 OUR AVERAGE				Error includes scale factor of 1.6.
1.5 ± 0.5 ± 0.1	12 ± 4	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.6 ± 0.2 ± 0.1	18.4 ± 6.4	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.46±0.71±0.02	6 ± 3	2,3 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3} \text{ keV}$ which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04 \text{ keV}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

Γ_{127}/Γ

NODE=M071R83
NODE=M071R83

$\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.63±0.13 OUR AVERAGE				
0.9 ± 0.4 ± 0.1	13	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
0.6 ± 0.1 ± 0.1	59.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

Γ_{128}/Γ

NODE=M071S12
NODE=M071S12

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.16 OUR AVERAGE				
0.8 ± 0.2 ± 0.1	36.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ± 0.2 ± 0.1	16.1 ± 5.0	¹ BAI	03B BES	$\psi(2S) \rightarrow 2(K^+ K^-)$

Γ_{129}/Γ

NODE=M071R81
NODE=M071R81

$\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.2±0.2	44.7	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)\pi^0$

Γ_{130}/Γ

NODE=M071R81;LINKAGE=B3

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.10±0.31 OUR AVERAGE				
3.14±0.23±0.23	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.0 $^{+1.5}_{-1.1}$ ± 0.4	6	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.3 ± 1.1 ± 0.5	17	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

Γ_{131}/Γ

NODE=M071R89
NODE=M071R89

$\Gamma(\eta\phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{132}/Γ	NODE=M071P19 NODE=M071P19
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.2 \times 10^{-6}$	90	ABLIKIM	19I	BES3	$e^+ e^- \rightarrow \eta\phi f_0(980)$	
$\Gamma(\phi\eta')/\Gamma_{\text{total}}$					Γ_{133}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M071R90 NODE=M071R90
1.54 ± 0.20 OUR AVERAGE						NEW
$[(3.1 \pm 1.6) \times 10^{-5}$ OUR 2019 AVERAGE]						
1.51 $\pm 0.16 \pm 0.12$	201	ABLIKIM	19BA	BES3	$e^+ e^- \rightarrow \psi(2S)$	
3.1 $\pm 1.4 \pm 0.7$	8	¹ ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$	
1 Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.						
$\Gamma(\omega\eta')/\Gamma_{\text{total}}$					Γ_{136}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M071R91 NODE=M071R91
$3.2^{+2.4}_{-2.0} \pm 0.7$	4	¹ ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$	
1 Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.						
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					Γ_{137}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M071R92 NODE=M071R92
2.1 ± 0.6 OUR AVERAGE						
2.5 $\pm 1.2 \pm 0.2$	14	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$	
1.87 $\pm 0.68 \pm 0.28$	14	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$	
$\Gamma(\rho\eta')/\Gamma_{\text{total}}$					Γ_{138}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M071R93 NODE=M071R93
$1.87^{+1.64}_{-1.11} \pm 0.33$	2	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.02 $\pm 0.11 \pm 0.24$	143	¹ ABLIKIM	17AK	BES3	$e^+ e^- \rightarrow \psi(2S)$	
0.569 $\pm 0.128 \pm 0.236$	80	² ABLIKIM	17AK	BES3	$e^+ e^- \rightarrow \psi(2S)$	
1 Destructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.						
2 Constructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.						
$\Gamma(\rho\eta)/\Gamma_{\text{total}}$					Γ_{139}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M071R94 NODE=M071R94
2.2 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.1.		
3.0 $\pm 1.1 \pm 0.2$	18	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$	
1.78 $\pm 0.67 \pm 0.17$	13	ABLIKIM	04L	BES	$e^+ e^- \rightarrow \psi(2S)$	
$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					Γ_{140}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M071R95 NODE=M071R95
<1.1	90	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<3.1	90	ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$	
$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$					Γ_{141}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M071R96 NODE=M071R96
<0.04	90	ABLIKIM	12L	BES3	$e^+ e^- \rightarrow \psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.7	90	ADAM	05	CLEO	$e^+ e^- \rightarrow \psi(2S)$	
<0.4	90	ABLIKIM	04K	BES	$e^+ e^- \rightarrow \psi(2S)$	
$\Gamma(\eta_c\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{142}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M071R03 NODE=M071R03
<1.0	90	PEDLAR	07	CLEO	$e^+ e^- \rightarrow \psi(2S)$	

$\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$				Γ_{143}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{144}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.81 \pm 0.11 \pm 0.14$	50	1 ABLIKIM	08C	BES2 $e^+e^- \rightarrow J/\psi$
¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%.$				
$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$				Γ_{145}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$0.44 \pm 0.12 \pm 0.11$	20 ± 6	BAI	04C	$\psi(2S) \rightarrow 2(K^+K^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.45	90	BAI	98J	BES $e^+e^- \rightarrow 2(K^+K^-)$
$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$				Γ_{134}/Γ
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.0 \pm 0.4 \pm 1.3$	234	1 ABLIKIM	19BA	BES3 $e^+e^- \rightarrow \psi(2S)$
¹ ABLIKIM 19BA reports $[\Gamma(\psi(2S) \rightarrow \phi f_1(1285))/\Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta\pi^+\pi^-)] = (1.03 \pm 0.10 \pm 0.09) \times 10^{-5}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta\pi^+\pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
$\Gamma(\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta)/\Gamma_{\text{total}}$				Γ_{135}/Γ
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$8.46 \pm 1.37 \pm 0.92$	195	ABLIKIM	19BA	BES3 $e^+e^- \rightarrow \psi(2S)$
$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{146}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.88	90	BAI	04G	BES2 e^+e^-
$\Gamma(\Theta(1540)K^-\bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$				Γ_{147}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.0	90	BAI	04G	BES2 e^+e^-
$\Gamma(\Theta(1540)K_S^0\bar{p} \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$				Γ_{148}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.70	90	BAI	04G	BES2 e^+e^-
$\Gamma(\bar{\Theta}(1540)K^+n \rightarrow K_S^0\bar{p}K^+n)/\Gamma_{\text{total}}$				Γ_{149}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.6	90	BAI	04G	BES2 e^+e^-
$\Gamma(\bar{\Theta}(1540)K_S^0p \rightarrow K_S^0pK^-\bar{n})/\Gamma_{\text{total}}$				Γ_{150}/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.60	90	BAI	04G	BES2 e^+e^-
$\Gamma(K_S^0K_S^0)/\Gamma_{\text{total}}$				Γ_{151}/Γ
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.046	1	BAI	04D	BES e^+e^-
1 Forbidden by CP.				
$\Gamma(\Lambda_c^+\bar{p}e^+e^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{152}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$<1.7 \times 10^{-6}$	90	450M	ABLIKIM	18Q
				BES3 $e^+e^- \rightarrow \psi(2S)$

RADIATIVE DECAYS **$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.79 ± 0.20 OUR FIT**9.33 ± 0.26 OUR AVERAGE**

9.389 ± 0.014 ± 0.332	4.7M	ABLIKIM	17U	BES3	$e^+ e^- \rightarrow \gamma X$
9.22 ± 0.11 ± 0.46	72k	ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
9.9 ± 0.5 ± 0.8		¹ GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.2 ± 2.3		¹ BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$
7.5 ± 2.6		¹ WHITAKER	76	MRK1	$e^+ e^- \rightarrow \gamma X$

¹ Angular distribution ($1+\cos^2\theta$) assumed. **Γ_{153}/Γ**

NODE=M071315

NODE=M071R55

NODE=M071R55

 $\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.75 ± 0.24 OUR FIT**9.54 ± 0.29 OUR AVERAGE**

9.905 ± 0.011 ± 0.353	5.0M	ABLIKIM	17U	BES3	$e^+ e^- \rightarrow \gamma X$
9.07 ± 0.11 ± 0.54	76k	ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
9.0 ± 0.5 ± 0.7		¹ GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.1 ± 1.9		² BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$

¹ Angular distribution ($1-0.189 \cos^2\theta$) assumed.² Valid for isotropic distribution of the photon. **Γ_{154}/Γ**

NODE=M071R58

NODE=M071R58

 $\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------


9.52 ± 0.20 OUR FIT**9.42 ± 0.31 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

9.621 ± 0.013 ± 0.272	4.2M	ABLIKIM	17U	BES3	$e^+ e^- \rightarrow \gamma X$
9.33 ± 0.14 ± 0.61	79k	ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
8.0 ± 0.5 ± 0.7		¹ GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.0 ± 2.0		² BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$

¹ Angular distribution ($1-0.052 \cos^2\theta$) assumed.² Valid for isotropic distribution of the photon. **Γ_{155}/Γ**

NODE=M071R59

NODE=M071R59



VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •27.6 ± 0.3 ± 2.0 ¹ ATHAR 04 CLEO $e^+ e^- \rightarrow \gamma X$ ¹ Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.

NODE=M071R19

NODE=M071R19

NODE=M071R;LINKAGE=AH

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.02 \pm 0.01 \pm 0.07$	¹ ATHAR 04	CLEO	$e^+ e^- \rightarrow \gamma X$
1 Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.			

 $\Gamma_{153}/\Gamma_{154}$

NODE=M071R97
NODE=M071R97

 $\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$1.03 \pm 0.02 \pm 0.03$	¹ ATHAR 04	CLEO	$e^+ e^- \rightarrow \gamma X$
1 Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.			

 $\Gamma_{155}/\Gamma_{154}$

NODE=M071R97;LINKAGE=AH
NODE=M071R98
NODE=M071R98

 $\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.99 \pm 0.02 \pm 0.08$	¹ ATHAR 04	CLEO	$e^+ e^- \rightarrow \gamma X$
1 Not independent from ATHAR 04 measurements of $B(\gamma\chi_{cJ})$.			

 $\Gamma_{153}/\Gamma_{155}$

NODE=M071R99
NODE=M071R99


 $\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.

$0.432 \pm 0.016 \pm 0.060$		MITCHELL	09	CLEO	$e^+ e^- \rightarrow \gamma X$
$0.32 \pm 0.04 \pm 0.06$	2.5k	¹ ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
0.28 ± 0.06		² GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$

¹ ATHAR 04 used $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$ MeV to obtain this result.

² GAISER 86 used $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$ MeV to obtain this result.

 $\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{157}/Γ

NODE=M071R62
NODE=M071R62

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
7±2±4		¹ ABLIKIM 12G	BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi, KK\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

< 8	90	² CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K\bar{K}\pi$
< 20	90	ATHAR	04	$e^+ e^- \rightarrow \gamma X$

20–130 95 EDWARDS 82C CBAL $e^+ e^- \rightarrow \gamma X$

¹ ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M071R62;LINKAGE=AB

NODE=M071R62;LINKAGE=CR

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.04 ± 0.22 OUR AVERAGE	Error includes scale factor of 1.4.				
0.95 ± 0.16 ± 0.05		423	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^0$
1.58 ± 0.40 ± 0.13		37	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 5		90	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
< 5400		95	1 LIBERMAN	75 SPEC	$e^+ e^-$
$< 1 \times 10^4$		90	WIIK	75 DASP	$e^+ e^-$

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$.

 Γ_{158}/Γ

NODE=M071R42
NODE=M071R42

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.04 OUR AVERAGE					
1.251 ± 0.022 ± 0.062		56K	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta$, $\gamma\pi^0\pi^0\eta$
1.26 ± 0.03 ± 0.08		2226	1 ABLIKIM	10F BES3	$\psi(2S) \rightarrow 3\gamma\pi^+\pi^-$, $2\gamma\pi^+\pi^-$
1.19 ± 0.08 ± 0.03			PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
1.24 ± 0.27 ± 0.15		23	ABLIKIM	06R BES2	$e^+ e^- \rightarrow \psi(2S)$
1.54 ± 0.31 ± 0.20		~ 43	BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^-2\gamma$, $\pi^+\pi^-3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60		90	2 BRAUNSCH...	77 DASP	$e^+ e^-$
< 11		90	3 BARTEL	76 CNTR	$e^+ e^-$

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

 Γ_{159}/Γ

NODE=M071R44
NODE=M071R44

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.73 ± 0.29 OUR AVERAGE	Error includes scale factor of 1.8.				
2.84 ± 0.15 ± 0.03	1.9k	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$	
2.12 ± 0.19 ± 0.32		3,4 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.08 ± 0.19 ± 0.33	200.6 ± 18.8	3 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$	
2.90 ± 1.08 ± 1.07	29.9 ± 11.1	3 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.2 \pm 2.9) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

 Γ_{160}/Γ

NODE=M071R84
NODE=M071R84

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
3.1 ± 1.0 ± 1.4	175	1 DOBBS	15 $\psi(2S) \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

OCCUR=2
OCCUR=3

NODE=M071R84;LINKAGE=A
NODE=M071R84;LINKAGE=B

NODE=M071R;LINKAGE=3B
NODE=M071R;LINKAGE=B9

NODE=M071S75
NODE=M071S75

NODE=M071S75;LINKAGE=A

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
9.3 ± 1.8 ± 0.6	274	1,2 DOBBS	15 $\psi(2S) \rightarrow \gamma\pi\pi$

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi\pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi\pi) = (34.5 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using CLEO-c data but not authored by the CLEO Collaboration.

 Γ_{161}/Γ

NODE=M071S76
NODE=M071S76

NODE=M071S76;LINKAGE=A

NODE=M071S76;LINKAGE=B

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$					Γ_{163}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.3 \pm 0.8 \pm 0.1$	136	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$	NODE=M071S77 NODE=M071S77
¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² Using CLEO-c data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$					Γ_{165}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.5 ± 0.6 OUR AVERAGE					NODE=M071R85 NODE=M071R85
$3.6 \pm 0.4 \pm 0.5$	290	1 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$	
$3.01 \pm 0.41 \pm 1.24$	35.6 ± 4.8	² BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$	
¹ Using CLEO-c data but not authored by the CLEO Collaboration.					
² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.					
$\Gamma(\gamma f_0(1710) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$					Γ_{166}/Γ
VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.6 ± 0.7 OUR AVERAGE					NODE=M071R86 NODE=M071R86
$6.7 \pm 0.6 \pm 0.6$		375	1 DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$
$6.04 \pm 0.90 \pm 1.32$		39.6 ± 5.9	^{2,3} BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 15.6	90	6.8 ± 3.1	^{2,3} BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
¹ Using CLEO-c data but not authored by the CLEO Collaboration.					
² Includes unknown branching fractions to K^+K^- or $K_S^0 K_S^0$. We have multiplied the K^+K^- result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K\bar{K}$ result.					
³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.					
$\Gamma(\gamma f_0(2100) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$					Γ_{167}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$4.8 \pm 0.5 \pm 0.9$	373	1 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$	NODE=M071S78 NODE=M071S78
¹ Using CLEO-c data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma f_0(2200) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$					Γ_{168}/Γ
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	
$3.2 \pm 0.6 \pm 0.8$	207	1 DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$	NODE=M071S79 NODE=M071S79
¹ Using CLEO-c data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma f_J(2220) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$					Γ_{169}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 5.8 \times 10^{-6}$	90	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$	NODE=M071S80 NODE=M071S80
¹ Using CLEO-c data but not authored by the CLEO Collaboration.					
² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+\pi^-$ and $\pi^0\pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.					
$\Gamma(\gamma f_J(2220) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$					Γ_{170}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 9.5 \times 10^{-6}$	90	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$	NODE=M071S81 NODE=M071S81
¹ Using CLEO-c data but not authored by the CLEO Collaboration.					
² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for K^+K^- and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.					
$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$					Γ_{172}/Γ
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.18 OUR AVERAGE					NODE=M071R43 NODE=M071R43
$0.85 \pm 0.18 \pm 0.04$	382	1 ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$, $\gamma 3\pi^0$	
$1.38 \pm 0.48 \pm 0.09$	13	1 ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$, $\gamma 3\pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 90	90	BAI	98F	BES	$\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$
<200	90	YAMADA	77	DASP	$e^+ e^- \rightarrow 3\gamma$

¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

$\Gamma(\gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{173}/Γ
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$8.71 \pm 1.25 \pm 1.64$	418	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$					Γ_{175}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.9	90	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	90	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
<1.2	90	¹ SCHARRE	80	MRK1	$e^+ e^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K\bar{K}\pi$.

$\Gamma(\gamma\eta(1405) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{176}/Γ
<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.36 $\pm 0.25 \pm 0.05$	10	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{177}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$<5.0 \times 10^{-7}$	90	ABLIKIM	17AJ	BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

$\Gamma(\gamma\eta(1475) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}}$					Γ_{179}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.4	90	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	90	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$
------	----	---------	-----	------	---

$\Gamma(\gamma\eta(1475) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{180}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.88	90	ABLIKIM	06R	BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

$\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_{181}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
39.6 $\pm 2.8 \pm 5.0$	583	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^{*0} K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{182}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
37.0 $\pm 6.1 \pm 7.2$	237	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}}$					Γ_{183}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
24.0 $\pm 4.5 \pm 5.0$	41	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{184}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
25.6 $\pm 3.6 \pm 3.6$	115	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{185}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
19.1 $\pm 2.7 \pm 4.3$	132	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

$\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{186}/Γ
<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
3.9 ± 0.5 OUR AVERAGE				Error includes scale factor of 2.0.	

4.18 $\pm 0.26 \pm 0.18$ 348 ¹ ALEXANDER 10 CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$

2.9 $\pm 0.4 \pm 0.4$ 142 ABLIKIM 07D BES2 $e^+ e^- \rightarrow \psi(2S)$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

NODE=M071R43;LINKAGE=AB

NODE=M071R04
NODE=M071R04

NODE=M071R61
NODE=M071R61

OCCUR=2

NODE=M071R;LINKAGE=E

NODE=M071R05
NODE=M071R05

NODE=M071P09
NODE=M071P09

NODE=M071P09
NODE=M071P09

NODE=M071R06
NODE=M071R06

OCCUR=2

NODE=M071R07
NODE=M071R07

NODE=M071R07
NODE=M071R07

NODE=M071S28
NODE=M071S28

NODE=M071S28
NODE=M071S28

NODE=M071S29
NODE=M071S29

NODE=M071S29
NODE=M071S29

NODE=M071S30
NODE=M071S30

NODE=M071S30
NODE=M071S30

NODE=M071S31
NODE=M071S31

NODE=M071S31
NODE=M071S31

NODE=M071S32
NODE=M071S32

NODE=M071S32
NODE=M071S32

NODE=M071S33
NODE=M071S33

NODE=M071S33
NODE=M071S33

NODE=M071S33;LINKAGE=AL

$\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.2 \pm 0.2 \pm 0.1$	111	1 ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 Γ_{187}/Γ

NODE=M071S46
NODE=M071S46

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$0.72 \pm 0.18 \pm 0.03$	73	1 ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.

 Γ_{188}/Γ

NODE=M071S47
NODE=M071S47

 $\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$4.57 \pm 0.36 \pm 1.77$		ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
<5.4	90	ABLIKIM	07D BES	$\psi(2S) \rightarrow \gamma p\bar{p}$

 Γ_{189}/Γ

NODE=M071S48
NODE=M071S48

 $\Gamma(\gamma X \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

 Γ_{190}/Γ

NODE=M071S49
NODE=M071S49
NODE=M071S49

 $\Gamma(\gamma \pi^+ \pi^- p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$2.8 \pm 1.2 \pm 0.7$	17	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{191}/Γ

NODE=M071S34
NODE=M071S34

 $\Gamma(\gamma 2(\pi^+ \pi^-) K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<22	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{192}/Γ

NODE=M071S35
NODE=M071S35

 $\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<17	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{193}/Γ

NODE=M071S36
NODE=M071S36

 $\Gamma(\gamma K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{194}/Γ

NODE=M071S37
NODE=M071S37

 $\Gamma(\gamma \gamma J/\psi)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$3.1 \pm 0.6 \pm 0.8$	1.1k	ABLIKIM	120 BES3	$e^+ e^- \rightarrow \psi(2S)$

 Γ_{195}/Γ

NODE=M071S55
NODE=M071S55

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.2±0.6	1.1k	1 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$
---------	------	-----------	----------	--

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$. No systematic error estimation.

NODE=M071S55;LINKAGE=A

 $\Gamma(e^+ e^- \eta')/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.90 ± 0.26 OUR AVERAGE	57	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$

 Γ_{196}/Γ

NODE=M071P12
NODE=M071P12

1.99±0.33±0.12	57	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$
1.79±0.38±0.11	20	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$

 Γ_{196}/Γ

OCCUR=2

$\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}$	Γ_{197}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$10.6 \pm 2.4 \pm 0.4$	48	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

1 ABLIKIM 17I reports $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.40 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}$	Γ_{198}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.5 \pm 0.6 \pm 0.2$	873	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

1 ABLIKIM 17I reports $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c2}(1P))/\Gamma_{\text{total}}$	Γ_{199}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.0 \pm 0.7 \pm 0.2$	227	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

1 ABLIKIM 17I reports $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$ assuming $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$, which we rescale to our best value $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma(\gamma \chi_{c0}(1P))$	$\Gamma_{197}/\Gamma_{153}$			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$9.4 \pm 1.9 \pm 0.6$	48	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

1 Uses $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma(\gamma \chi_{c1}(1P))$	$\Gamma_{198}/\Gamma_{154}$			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 0.3 \pm 0.4$	873	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

1 Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$\Gamma(e^+ e^- \chi_{c2}(1P))/\Gamma(\gamma \chi_{c2}(1P))$	$\Gamma_{199}/\Gamma_{155}$			
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.6 \pm 0.5 \pm 0.4$	227	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

1 Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

WEAK DECAYS

$\Gamma(D^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{200}/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-7}$	90	1 ABLIKIM	17AF BES3	$e^+ e^- \rightarrow \psi(2S)$

1 Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

OTHER DECAYS

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$	Γ_{201}/Γ_6			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.0	90	LEES	13I BABR	$B \rightarrow K^{(*)} \psi(2S)$

$\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

NODE=M071P01
NODE=M071P01

NODE=M071P01;LINKAGE=B

NODE=M071P02
NODE=M071P02

NODE=M071P02;LINKAGE=B

NODE=M071P03
NODE=M071P03

NODE=M071P04
NODE=M071P04

NODE=M071P04;LINKAGE=A

NODE=M071P05
NODE=M071P05

NODE=M071P05;LINKAGE=A

NODE=M071P06
NODE=M071P06

NODE=M071P06;LINKAGE=A

NODE=M071330

NODE=M071P07
NODE=M071P07

NODE=M071P07;LINKAGE=A

NODE=M071320

NODE=M071S64
NODE=M071S64

NODE=M071240

NODE=M071240

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

$a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
63 ± 7 OUR AVERAGE				
61.7 ± 8.3	253k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
67 $^{+19}_{-13}$	59k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined.

² Statistical and systematic errors combined. Using values from fits with floating M2 amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed E3 amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.

$b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
60±31 OUR AVERAGE				
74±40	253k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
37 $^{+53}_{-47}$	59k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined. Derived from the reported measurement of $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$.

² Statistical and systematic errors combined. Using values from fits with floating M2 amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed E3 amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

NODE=M071250

NODE=M071QAR

NODE=M071QAR

NODE=M071QAR;LINKAGE=A

NODE=M071QAR;LINKAGE=AR

NODE=M071QBR

NODE=M071QBR

NODE=M071QBR;LINKAGE=A

NODE=M071QBR;LINKAGE=AR

NODE=M071

ABLIKIM	19AO PR D99 112010	M. Ablikim et al.	(BESIII Collab.)	REFID=59892
ABLIKIM	19AT PR D100 051101	M. Ablikim et al.	(BESIII Collab.)	REFID=59989
ABLIKIM	19AU PR D100 052010	Ablikim et al.	(BESIII Collab.)	REFID=59996;ERROR=16
ABLIKIM	19BA PR D100 092003	M. Ablikim et al.	(BESIII Collab.)	REFID=60024
ABLIKIM	19I PR D99 012014	M. Ablikim et al.	(BESIII Collab.)	REFID=59605
ABLIKIM	19N PR D99 032006	M. Ablikim et al.	(BESIII Collab.)	REFID=59615
ABLIKIM	18Q PR D97 091102	M. Ablikim et al.	(BESIII Collab.)	REFID=58933
ABLIKIM	18T PR D98 032006	M. Ablikim et al.	(BESIII Collab.)	REFID=58975
ABLIKIM	18Z PL B783 452	M. Ablikim et al.	(BESIII Collab.)	REFID=59038
ANASHIN	18 PL B781 174	V.V. Anashin et al.	(KEDR Collab.)	REFID=59013
LEES	18E PR D98 112015	J.P. Lees et al.	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF PR D96 111101	M. Ablikim et al.	(BESIII Collab.)	REFID=58315
ABLIKIM	17AJ PR D96 112008	M. Ablikim et al.	(BESIII Collab.)	REFID=58322
ABLIKIM	17AK PR D96 112012	M. Ablikim et al.	(BESIII Collab.)	REFID=58324
ABLIKIM	17E PL B770 217	M. Ablikim et al.	(BESIII Collab.)	REFID=57903
ABLIKIM	17I PRL 118 221802	M. Ablikim et al.	(BESIII Collab.)	REFID=57931
ABLIKIM	17L PR D95 050203	M. Ablikim et al.	(BESIII Collab.)	REFID=57967
ABLIKIM	17N PR D95 072004	M. Ablikim et al.	(BESIII Collab.)	REFID=57978
ABLIKIM	17U PR D96 032001	M. Ablikim et al.	(BESIII Collab.)	REFID=58026
ABLIKIM	17X PR D96 050203	M. Ablikim et al.	(BESIII Collab.)	REFID=58216
DOBBS	17 PR D96 092004	S. Dobbs et al.	(NWES, WAYN)	REFID=58670
LEES	17A PR D95 052001	J.P. Lees et al.	(BABAR Collab.)	REFID=57966
AAIJ	16Y JHEP 1605 132	R. Aaij et al.	(LHCb Collab.)	REFID=57333
ABLIKIM	16L PR D93 072003	M. Ablikim et al.	(BESIII Collab.)	REFID=57510
ABLIKIM	15I PR D91 092006	M. Ablikim et al.	(BESIII Collab.)	REFID=56774
ABLIKIM	15V PL B749 414	M. Ablikim et al.	(BESIII Collab.)	REFID=56787
ANASHIN	15 PL B749 50	V.V. Anashin et al.	(KEDR Collab.)	REFID=56792
DOBBS	15 PR D91 052006	S. Dobbs et al.	(NWES)	REFID=56805
LEES	15J PR D92 072008	J.P. Lees et al.	(BABAR Collab.)	REFID=56988
ABLIKIM	14G PR D89 112006	M. Ablikim et al.	(BESIII Collab.)	REFID=55898
DOBBS	14 PL B739 90	S. Dobbs et al.	(NWES, WAYN)	REFID=56333
ABLIKIM	13A PR 110 022001	M. Ablikim et al.	(BESIII Collab.)	REFID=54834
ABLIKIM	13D PR D87 012007	M. Ablikim et al.	(BESIII Collab.)	REFID=54879
ABLIKIM	13F PR D87 052007	M. Ablikim et al.	(BESIII Collab.)	REFID=54920
ABLIKIM	13M PR D87 092006	M. Ablikim et al.	(BESIII Collab.)	REFID=55386
ABLIKIM	13R PR D88 032007	M. Ablikim et al.	(BESIII Collab.)	REFID=55402
ABLIKIM	13S PR D88 032010	M. Ablikim et al.	(BESIII Collab.)	REFID=55403
ABLIKIM	13W PR D88 112007	M. Ablikim et al.	(BESIII Collab.)	REFID=55634
LEES	13I PR D87 112005	J.P. Lees et al.	(BABAR Collab.)	REFID=55161
LEES	13O PR D87 092005	J.P. Lees et al.	(BABAR Collab.)	REFID=55293
LEES	13Q PR D88 032013	J.P. Lees et al.	(BABAR Collab.)	REFID=55404
LEES	13Y PR D88 072009	J.P. Lees et al.	(BABAR Collab.)	REFID=55589
AAIJ	12H EPJ C72 1972	R. Aaij et al.	(LHCb Collab.)	REFID=54056
ABLIKIM	12D PR 108 112003	M. Ablikim et al.	(BESIII Collab.)	REFID=54269
ABLIKIM	12G PR L09 042003	M. Ablikim et al.	(BESIII Collab.)	REFID=54272
ABLIKIM	12H PL B710 594	M. Ablikim et al.	(BESIII Collab.)	REFID=54273
ABLIKIM	12L PR D86 072011	M. Ablikim et al.	(BESIII Collab.)	REFID=54739
ABLIKIM	12M PR D86 092008	M. Ablikim et al.	(BESIII Collab.)	REFID=54740
ABLIKIM	12O PR 109 172002	M. Ablikim et al.	(BESIII Collab.)	REFID=54742
ABLIKIM	12Q CP C36 1040	M. Ablikim et al.	(BES II Collab.)	REFID=54864
ANASHIN	12 PL B711 280	V.V. Anashin et al.	(KEDR Collab.)	REFID=54038
LEES	12E PR D85 112009	J.P. Lees et al.	(BABAR Collab.)	REFID=54297
LEES	12F PR D86 012008	J.P. Lees et al.	(BABAR Collab.)	REFID=54298
METREVELI	12 PR D85 092007	Z. Metreveli et al.	(NWES, FLOR, WAYN+)	REFID=54304
GE	11 PR D84 032008	J.Y. Ge et al.	(CLEO Collab.)	REFID=53960
ABLIKIM	10B PRL 104 132002	M. Ablikim et al.	(BESIII Collab.)	REFID=53348
ABLIKIM	10F PRL 105 261801	M. Ablikim et al.	(BESIII Collab.)	REFID=53630
ALEXANDER	10 PR D82 092002	J.P. Alexander et al.	(CLEO Collab.)	REFID=53525
CRONIN-HEN...10	PR D81 052002	D. Cronin-Hennessey et al.	(CLEO Collab.)	REFID=53233

ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=53124
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51636
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51725
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=51655
		Translated from ZETFP 85, 429.			
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)	REFID=52717;ERROR=17
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=51630
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51048
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51560
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51158
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50757
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50758
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50760
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50451
ADAM	05A	PR 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50497
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50785
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=50808
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50327
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50328
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49608
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49749
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49186
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48548
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48578
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49171
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)	REFID=48632
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48003
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47939
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46340
BAI	98J	PR D81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46554
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45416
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
GAISER	86	PR D34 711	J. Gaisser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41, 733.			
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22173
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
OREGLIA	80	PR 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22207
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
		Translated from YAF 34, 1471.			
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22197
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22198
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)	REFID=22064
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG	REFID=22194
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)	REFID=22176
ABRAMS	75B	PR 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)	REFID=22177
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)	REFID=22179
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)	REFID=22186
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)	REFID=22046
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC	REFID=22188
WIJK	75	Stanford Symp. 69	B.H. Wiik	(DESY)	REFID=22050

NODE=M053

 $\psi(3770)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\psi(3770)$ MASS (MeV)**

OUR FIT includes measurements of $m_{\psi}(2S)$, $m_{\psi}(3770)$, and $m_{\psi}(3770) - m_{\psi}(2S)$.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3773.7±0.4 OUR FIT	Error includes scale factor of 1.4. [3773.13 ± 0.35 MeV OUR 2019 FIT Scale factor = 1.1]			
3778.1±0.7 OUR AVERAGE	[3778.1 ± 1.2 MeV OUR 2019 AVERAGE]			
3778.1±0.7±0.6	1	AAIJ	19M LHCb $pp \rightarrow D\bar{D} +$ anything	
3779.2 ^{+1.8} _{-1.7} ^{+0.6} _{-0.8}	2	ANASHIN	12A KEDR $e^+ e^- \rightarrow D\bar{D}$	
3775.5±2.4±0.5	57	AUBERT	08B BABR $B \rightarrow D\bar{D}K$	
3776 ± 5 ± 4	68	BRODZICKA	08 BELL $B^+ \rightarrow D^0\bar{D}^0K^+$	
3778.8±1.9±0.9		AUBERT	07BE BABR $e^+ e^- \rightarrow D\bar{D}\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3779.8±0.6	3	SHAMOV	17 RVUE $e^+ e^- \rightarrow D\bar{D}$, hadrons	
3772.0±1.9	4,5	ABLIKIM	08D BES2 $e^+ e^- \rightarrow$ hadrons	
3778.4±3.0±1.3	34	CHISTOV	04 BELL Sup. by BRODZICKA 08	

1 Measured in prompt hadroproduction.

2 Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.3 From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.4 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.5 Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

NODE=M053M

NODE=M053M

NODE=M053M

NEW

NEW

 $m_{\psi(3770)} - m_{\psi(2S)}$

OUR FIT includes measurements of $m_{\psi}(2S)$, $m_{\psi}(3770)$, and $m_{\psi}(3770) - m_{\psi}(2S)$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
87.6±0.4 OUR FIT	Error includes scale factor of 1.4. [87.04 ± 0.35 MeV OUR 2019 FIT Scale factor = 1.1]		
86.6±0.7 OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.		
86.9±0.4	1	ABLIKIM	07E BES2 $e^+ e^- \rightarrow$ hadrons
86.7±0.7		ABLIKIM	06L BES2 $e^+ e^- \rightarrow$ hadrons
80 ± 2		SCHINDLER	80 MRK2 $e^+ e^-$
86 ± 2	2	BACINO	78 DLCO $e^+ e^-$
88 ± 3		RAPIDIS	77 LGW $e^+ e^-$

1 BES-II $\psi(2S)$ mass subtracted (see ABLIKIM 06L).2 SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).

NODE=M053DM


NODE=M053DM

NODE=M053DM

NEW

NODE=M053DM;LINKAGE=AK

NODE=M053DM;LINKAGE=S



$\psi(3770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.2± 1.0 OUR FIT				NODE=M053W
27.5± 0.9 OUR AVERAGE				NODE=M053W
24.9± 4.6±0.5		1 ANASHIN	12A KEDR	$e^+ e^- \rightarrow D\bar{D}$
30.4± 8.5		2,3 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
27 ±10 ±5	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$
28.5± 1.2±0.2		3 ABLIKIM	07E BES2	$e^+ e^- \rightarrow$ hadrons
23.5± 3.7±0.9		AUBERT	07BE BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$
26.9± 2.4±0.3		3 ABLIKIM	06L BES2	$e^+ e^- \rightarrow$ hadrons
24 ± 5		3 SCHINDLER	80 MRK2	$e^+ e^-$
24 ± 5		3 BACINO	78 DLCO	$e^+ e^-$
28 ± 5		3 RAPIDIS	77 LGW	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •		4 SHAMOV	17 RVUE	$e^+ e^- \rightarrow D\bar{D}$, hadrons
25.8± 1.3				

1 Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

2 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

3 Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

4 From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.

$\psi(3770)$ DECAY MODES

In addition to the dominant decay mode to $D\bar{D}$, $\psi(3770)$ was found to decay into the final states containing the J/ψ (BAI 05, ADAM 06). ADAMS 06 and HUANG 06A searched for various decay modes with light hadrons and found a statistically significant signal for the decay to $\phi\eta$ only (ADAMS 06).

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 D\bar{D}$	(93 ± 8) %	S=2.0
$\Gamma_2 D^0 \bar{D}^0$	(52 ± 4) %	S=2.0
$\Gamma_3 D^+ D^-$	(41 ± 4) %	S=2.0
$\Gamma_4 J/\psi \pi^+ \pi^-$	(1.93 ± 0.28) $\times 10^{-3}$	DESIG=6
$\Gamma_5 J/\psi \pi^0 \pi^0$	(8.0 ± 3.0) $\times 10^{-4}$	DESIG=4
$\Gamma_6 J/\psi \eta$	(9 ± 4) $\times 10^{-4}$	DESIG=46
$\Gamma_7 J/\psi \pi^0$	< 2.8 $\times 10^{-4}$	DESIG=47
$\Gamma_8 e^+ e^-$	(9.6 ± 0.7) $\times 10^{-6}$	DESIG=48
		DESIG=1

Decays to light hadrons					NODE=M053;CLUMP=H
Γ_9	$b_1(1235)\pi$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=20
Γ_{10}	$\phi\eta'$	< 7	$\times 10^{-4}$	CL=90%	DESIG=17
Γ_{11}	$\omega\eta'$	< 4	$\times 10^{-4}$	CL=90%	DESIG=16
Γ_{12}	$\rho^0\eta'$	< 6	$\times 10^{-4}$	CL=90%	DESIG=15
Γ_{13}	$\phi\eta$	$(3.1 \pm 0.7) \times 10^{-4}$			DESIG=8
Γ_{14}	$\omega\eta$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=14
Γ_{15}	$\rho^0\eta$	< 5	$\times 10^{-4}$	CL=90%	DESIG=13
Γ_{16}	$\phi\pi^0$	< 3	$\times 10^{-5}$	CL=90%	DESIG=12
Γ_{17}	$\omega\pi^0$	< 6	$\times 10^{-4}$	CL=90%	DESIG=11
Γ_{18}	$\pi^+\pi^-\pi^0$	< 5	$\times 10^{-6}$	CL=90%	DESIG=9
Γ_{19}	$\rho\pi$	< 5	$\times 10^{-6}$	CL=90%	DESIG=10
Γ_{20}	K^+K^-				DESIG=234
Γ_{21}	$K^*(892)^+K^- + \text{c.c.}$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=19
Γ_{22}	$K^*(892)^0\bar{K}^0 + \text{c.c.}$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=18
Γ_{23}	$K_S^0 K_L^0$	< 1.2	$\times 10^{-5}$	CL=90%	DESIG=3
Γ_{24}	$2(\pi^+\pi^-)$	< 1.12	$\times 10^{-3}$	CL=90%	DESIG=21
Γ_{25}	$2(\pi^+\pi^-)\pi^0$	< 1.06	$\times 10^{-3}$	CL=90%	DESIG=22
Γ_{26}	$2(\pi^+\pi^-\pi^0)$	< 5.85	%	CL=90%	DESIG=208
Γ_{27}	$\omega\pi^+\pi^-$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=24
Γ_{28}	$3(\pi^+\pi^-)$	< 9.1	$\times 10^{-3}$	CL=90%	DESIG=52
Γ_{29}	$3(\pi^+\pi^-)\pi^0$	< 1.37	%	CL=90%	DESIG=55
Γ_{30}	$3(\pi^+\pi^-)2\pi^0$	< 11.74	%	CL=90%	DESIG=210
Γ_{31}	$\eta\pi^+\pi^-$	< 1.24	$\times 10^{-3}$	CL=90%	DESIG=23
Γ_{32}	$\pi^+\pi^-2\pi^0$	< 8.9	$\times 10^{-3}$	CL=90%	DESIG=206
Γ_{33}	$\rho^0\pi^+\pi^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=64
Γ_{34}	$\eta 3\pi$	< 1.34	$\times 10^{-3}$	CL=90%	DESIG=25
Γ_{35}	$\eta 2(\pi^+\pi^-)$	< 2.43	%	CL=90%	DESIG=53
Γ_{36}	$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%	DESIG=221
Γ_{37}	$\eta' 3\pi$	< 2.44	$\times 10^{-3}$	CL=90%	DESIG=26
Γ_{38}	$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%	DESIG=27
Γ_{39}	$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=28
Γ_{40}	$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%	DESIG=207
Γ_{41}	$4(\pi^+\pi^-)$	< 1.67	%	CL=90%	DESIG=62
Γ_{42}	$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%	DESIG=63
Γ_{43}	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%	DESIG=29
Γ_{44}	$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%	DESIG=30
Γ_{45}	$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%	DESIG=67
Γ_{46}	$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%	DESIG=68
Γ_{47}	ωK^+K^-	< 3.4	$\times 10^{-4}$	CL=90%	DESIG=32
Γ_{48}	$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%	DESIG=69
Γ_{49}	$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%	DESIG=70
Γ_{50}	$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%	DESIG=71
Γ_{51}	$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%	DESIG=209
Γ_{52}	$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%	DESIG=57
Γ_{53}	$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%	DESIG=58
Γ_{54}	ηK^+K^-	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=31
Γ_{55}	$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%	DESIG=222
Γ_{56}	$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%	DESIG=65
Γ_{57}	$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=33
Γ_{58}	ϕK^+K^-	< 7.5	$\times 10^{-4}$	CL=90%	DESIG=34
Γ_{59}	$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=35
Γ_{60}	$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=59
Γ_{61}	$K_S^0 K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=200
Γ_{62}	$K_S^0 K^-\pi^+\pi^0$	< 1.33	%	CL=90%	DESIG=201
Γ_{63}	$K_S^0 K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%	DESIG=214
Γ_{64}	$K_S^0 K^-2\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%	DESIG=202
Γ_{65}	$K_S^0 K^-\pi^+\rho^0$	< 1.6	%	CL=90%	DESIG=215

Γ_{66}	$K_S^0 K^- \pi^+ \eta$	< 1.3	%	CL=90%	DESIG=216
Γ_{67}	$K_S^0 K^- 2\pi^+ \pi^- \pi^0$	< 4.18	%	CL=90%	DESIG=203
Γ_{68}	$K_S^0 K^- 2\pi^+ \pi^- \eta$	< 4.8	%	CL=90%	DESIG=217
Γ_{69}	$K_S^0 K^- \pi^+ 2(\pi^+ \pi^-)$	< 1.22	%	CL=90%	DESIG=204
Γ_{70}	$K_S^0 K^- \pi^+ 2\pi^0$	< 2.65	%	CL=90%	DESIG=205
Γ_{71}	$K_S^0 K^- K^+ K^- \pi^+$	< 4.9	$\times 10^{-3}$	CL=90%	DESIG=218
Γ_{72}	$K_S^0 K^- K^+ K^- \pi^+ \pi^0$	< 3.0	%	CL=90%	DESIG=219
Γ_{73}	$K_S^0 K^- K^+ K^- \pi^+ \eta$	< 2.2	%	CL=90%	DESIG=220
Γ_{74}	$K^{*0} K^- \pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%	DESIG=60
Γ_{75}	$p\bar{p}$				DESIG=233
Γ_{76}	$p\bar{p}\pi^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=54
Γ_{77}	$p\bar{p}\pi^+ \pi^-$	< 5.8	$\times 10^{-4}$	CL=90%	DESIG=36
Γ_{78}	$\Lambda\bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%	DESIG=42
Γ_{79}	$p\bar{p}\pi^+ \pi^- \pi^0$	< 1.85	$\times 10^{-3}$	CL=90%	DESIG=37
Γ_{80}	$\omega p\bar{p}$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=39
Γ_{81}	$\Lambda\bar{\Lambda}\pi^0$	< 7	$\times 10^{-5}$	CL=90%	DESIG=72
Γ_{82}	$p\bar{p}2(\pi^+ \pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%	DESIG=61
Γ_{83}	$\eta p\bar{p}$	< 5.4	$\times 10^{-4}$	CL=90%	DESIG=38
Γ_{84}	$\eta_0^0 p\bar{p}\pi^+ \pi^-$	< 3.3	$\times 10^{-3}$	CL=90%	DESIG=223
Γ_{85}	$\rho^0 p\bar{p}$	< 1.7	$\times 10^{-3}$	CL=90%	DESIG=66
Γ_{86}	$p\bar{p}K^+ K^-$	< 3.2	$\times 10^{-4}$	CL=90%	DESIG=40
Γ_{87}	$\eta p\bar{p}K^+ K^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=224
Γ_{88}	$\pi^0 p\bar{p}K^+ K^-$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=225
Γ_{89}	$\phi p\bar{p}$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=41
Γ_{90}	$\Lambda\bar{\Lambda}\pi^+ \pi^-$	< 2.5	$\times 10^{-4}$	CL=90%	DESIG=43
Γ_{91}	$\Lambda\bar{p}K^+$	< 2.8	$\times 10^{-4}$	CL=90%	DESIG=44
Γ_{92}	$\Lambda\bar{p}K^+ \pi^+ \pi^-$	< 6.3	$\times 10^{-4}$	CL=90%	DESIG=45
Γ_{93}	$\Lambda\bar{\Lambda}\eta^-$	< 1.9	$\times 10^{-4}$	CL=90%	DESIG=226
Γ_{94}	$\Sigma^+ \bar{\Sigma}^-$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=227
Γ_{95}	$\Sigma^0 \bar{\Sigma}^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=228
Γ_{96}	$\Xi^+ \bar{\Xi}^-$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=229
Γ_{97}	$\Xi^0 \bar{\Xi}^0$	< 1.4	$\times 10^{-4}$	CL=90%	DESIG=230

Radiative decays

Γ_{98}	$\gamma \chi_{c2}$	< 6.4	$\times 10^{-4}$	CL=90%	NODE=M053;CLUMP=R DESIG=51
Γ_{99}	$\gamma \chi_{c1}$	(2.49 ± 0.23)	$\times 10^{-3}$		DESIG=50
Γ_{100}	$\gamma \chi_{c0}$	(6.9 ± 0.6)	$\times 10^{-3}$		DESIG=49
Γ_{101}	$\gamma \eta_c$	< 7	$\times 10^{-4}$	CL=90%	DESIG=231
Γ_{102}	$\gamma \eta_c(2S)$	< 9	$\times 10^{-4}$	CL=90%	DESIG=232
Γ_{103}	$\gamma \eta'$	< 1.8	$\times 10^{-4}$	CL=90%	DESIG=213
Γ_{104}	$\gamma \eta$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=212
Γ_{105}	$\gamma \pi^0$	< 2	$\times 10^{-4}$	CL=90%	DESIG=211

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 23 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 20.1$ for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	99			
x_8	0	0		
Γ	0	0	-44	

$x_2 \quad x_3 \quad x_8$

Mode	Rate (MeV)	Scale factor	
Γ_2 $D^0 \bar{D}^0$	14.0 ± 1.4	1.8	DESIG=5
Γ_3 $D^+ D^-$	11.2 ± 1.1	1.7	DESIG=6
Γ_8 $e^+ e^-$	(2.62 ± 0.18) $\times 10^{-4}$	1.4	DESIG=1

$\psi(3770)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$					Γ_8
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.262±0.018 OUR FIT		Error includes scale factor of 1.4.			
0.256±0.016 OUR AVERAGE		Error includes scale factor of 1.2.			
0.154 ^{+0.079 +0.021} _{-0.058 -0.027}	1,2 ANASHIN	12A KEDR	$e^+ e^- \rightarrow D\bar{D}$		
0.22 ± 0.05	3,4 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons		
0.277±0.011±0.013	4 ABLIKIM	07E BES2	$e^+ e^- \rightarrow$ hadrons		
0.203±0.003 ^{+0.041} _{-0.027}	1.4M BESSON	06 CLEO	$e^+ e^- \rightarrow$ hadrons		
0.276±0.050	4 SCHINDLER	80 MRK2	$e^+ e^-$		
0.18 ± 0.06	4 BACINO	78 DLCO	$e^+ e^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.196±0.018	6 SHAMOV	17 RVUE	$e^+ e^- \rightarrow D\bar{D}$, hadrons	OCCUR=2	
0.414 ^{+0.072 +0.093} _{-0.080 -0.028}	2,7 ANASHIN	12A KEDR	$e^+ e^- \rightarrow D\bar{D}$	OCCUR=2	
0.37 ± 0.09	8 RAPIDIS	77 LGW	$e^+ e^-$		
1 Solution I of the two solutions.					
2 Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.					
3 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.					
4 Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.					
5 BESSON 06 (as corrected in BESSON 10) measure $\sigma(e^+ e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30} \text{ nb at } \sqrt{s} = 3773 \pm 1 \text{ MeV}$, and obtain Γ_{ee} from the Born-level cross section calculated using $\psi(3770)$ mass and width from our 2004 edition, PDG 04.					
6 From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.					
7 Solution II of the two solutions.					
8 See also $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ below.					

$\psi(3770)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma = (\Gamma_2 + \Gamma_3)/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.93^{+0.08}_{-0.09} OUR FIT		Error includes scale factor of 2.0.		
0.93^{+0.08}_{-0.09} OUR AVERAGE		Error includes scale factor of 2.1.		
0.849±0.056±0.018	1 ABLIKIM	08B BES2	$e^+ e^- \rightarrow$ non- $D\bar{D}$	
1.033±0.014 ^{+0.048} _{-0.066}	1.427M BESSON	06 CLEO	$e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.836±0.049	3 SHAMOV	17 RVUE	$e^+ e^- \rightarrow D\bar{D}$, hadrons	
0.866±0.050±0.036	4,5 ABLIKIM	07K BES2	$e^+ e^- \rightarrow$ non- $D\bar{D}$	
0.836±0.073±0.042	5 ABLIKIM	06L BES2	$e^+ e^- \rightarrow D\bar{D}$	
0.855±0.017±0.058	5,6 ABLIKIM	06N BES2	$e^+ e^- \rightarrow D\bar{D}$	
1 Neglecting interference.				
2 Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of $D\bar{D}$ reported by CLEO in DOBBS 07.				
3 From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.				
4 Using $\sigma^{obs} = 7.07 \pm 0.58 \text{ nb}$ and neglecting interference.				
5 Not independent of ABLIKIM 08B.				
6 From a measurement of $\sigma(e^+ e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773 \text{ MeV}$, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.				

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.52^{+0.04}_{-0.05} OUR FIT		Error includes scale factor of 2.0.		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.467±0.047±0.023	ABLIKIM	06L BES2	$e^+ e^- \rightarrow D^0\bar{D}^0$	
0.499±0.013±0.038	1 ABLIKIM	06N BES2	$e^+ e^- \rightarrow D^0\bar{D}^0$	

¹ From a measurement of $\sigma(e^+ e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773 \text{ MeV}$, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

NODE=M053225

NODE=M053W1

NODE=M053W1

NODE=M053W1;LINKAGE=A1

NODE=M053W1;LINKAGE=AN

NODE=M053W1;LINKAGE=AB

NODE=M053W1;LINKAGE=NI

NODE=M053W1;LINKAGE=BE

NODE=M053W1;LINKAGE=BI

NODE=M053W1;LINKAGE=A2

NODE=M053W1;LINKAGE=R

NODE=M053230

NODE=M053R1

NODE=M053R1

NODE=M053R1;LINKAGE=AI

NODE=M053R1;LINKAGE=BE

NODE=M053R1;LINKAGE=A

NODE=M053R1;LINKAGE=AL

NODE=M053R1;LINKAGE=SU

NODE=M053R1;LINKAGE=AB

NODE=M053R46

NODE=M053R46

NODE=M053R46;LINKAGE=AB

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
-------	-------------	------	---------	-------------------

0.41 ±0.04 OUR FIT Error includes scale factor of 2.0.

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.369 \pm 0.037 \pm 0.028$	ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D^+ D^-$
$0.357 \pm 0.011 \pm 0.034$	¹ ABLIKIM	06N	BES2	$e^+ e^- \rightarrow D^+ D^-$

¹ From a measurement of $\sigma(e^+ e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.NODE=M053R47
NODE=M053R47 $\Gamma(D^0 \bar{D}^0)/\Gamma(D^+ D^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_3
-------	------	-------------	------	---------	---------------------

1.253±0.016 OUR FIT**1.253±0.016 OUR AVERAGE**

$1.252 \pm 0.009 \pm 0.013$	5.3M	BONVICINI	14	CLEO	$e^+ e^- \rightarrow D\bar{D}$
$1.39 \pm 0.31 \pm 0.12$		PAKHLOVA	08	BELL	$10.6 e^+ e^- \rightarrow D\bar{D}\gamma$
$1.78 \pm 0.33 \pm 0.24$		AUBERT	07BE	BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$
$1.27 \pm 0.12 \pm 0.08$		ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D\bar{D}$
$2.43 \pm 1.50 \pm 0.43$	34	¹ CHISTOV	04	BELL	$B^+ \rightarrow \psi(3770) K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.258 \pm 0.016 \pm 0.014$		² DOBBS	07	CLEO	$e^+ e^- \rightarrow D\bar{D}$
-----------------------------	--	--------------------	----	------	--------------------------------

¹ See ADLER 88C for older measurements of this quantity.² Superseded by BONVICINI 14.

NODE=M053R47;LINKAGE=AB

 $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
--------------------------	------	-------------	------	---------	-------------------

1.93±0.28 OUR AVERAGE

$1.89 \pm 0.20 \pm 0.20$	231 ± 33	ADAM	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
$3.4 \pm 1.4 \pm 0.9$	17.8 ± 4.8	BAI	05	BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R5;LINKAGE=CH
NODE=M053R5;LINKAGE=DO $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
--------------------------	------	-------------	------	---------	-------------------

$0.080 \pm 0.025 \pm 0.016$	39 ± 14	ADAM	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
-----------------------------	-------------	------	----	------	----------------------------------

NODE=M053R7
NODE=M053R7 $\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
--------------------------	------	-------------	------	---------	-------------------

$87 \pm 33 \pm 22$	22 ± 10	ADAM	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
--------------------	-------------	------	----	------	----------------------------------

NODE=M053R8
NODE=M053R8 $\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
--------------------------	-----	------	-------------	------	---------	-------------------

<28	90	<10	ADAM	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
-----	----	-----	------	----	------	----------------------------------

NODE=M053R9
NODE=M053R9 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
--------------------------	-------------	------	---------	-------------------

0.96±0.07 OUR FIT	Error includes scale factor of 1.3.			
1.3 ±0.2		RAPIDIS	77	LGW

NODE=M053R2
NODE=M053R2**DECAYS TO LIGHT HADRONS** $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
--------------------------	-----	-------------	------	---------	-------------------

<1.4	90	¹ ADAMS	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
------	----	--------------------	----	------	----------------------------------

NODE=M053250

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R82;LINKAGE=AD

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
--------------------------	-----	-------------	------	---------	----------------------

<7	90	¹ ADAMS	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
----	----	--------------------	----	------	----------------------------------

NODE=M053R82;LINKAGE=AD

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R83;LINKAGE=AD

 $\Gamma(\omega\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
--------------------------	-----	-------------	------	---------	----------------------

<4	90	¹ ADAMS	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
----	----	--------------------	----	------	----------------------------------

NODE=M053R84;LINKAGE=AD

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R84;LINKAGE=AD

$\Gamma(\rho^0 \eta')/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{12}/Γ
<6	90	1 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{13}/Γ
3.1±0.6±0.3		1 ADAMAS	06 CLEO	$3.773 e^+ e^- \rightarrow \phi\eta$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	2 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	-----------	----------	----------------------------------

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{14}/Γ
<1.4	90	1 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\rho^0 \eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{15}/Γ
<5	90	1 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{16}/Γ
< 3	90	1 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	90	2 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	-----------	----------	----------------------------------

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{17}/Γ
<6	90	1 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{18}/Γ
<5	90	1,2 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Data suggest possible destructive interference with continuum.

² Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{19}/Γ
<5	90	1,2 ADAMAS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

² Data suggest possible destructive interference with continuum.

 $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{20}/Γ
--------------	--------------------	-------------	----------------	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 10^{-5}$	¹ DRUZHININ	15 RVUE	$e^+ e^- \rightarrow \psi(3770)$
----------------	------------------------	---------	----------------------------------

¹ DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+ e^- \rightarrow K^+ K^-$ and $e^+ e^- \rightarrow K_S^0 K_L^0$.

NODE=M053R85
NODE=M053R85

NODE=M053R85;LINKAGE=AD
NODE=M053R86
NODE=M053R86

NODE=M053R86
NODE=M053R86

NODE=M053R86;LINKAGE=AD
NODE=M053R87
NODE=M053R87

NODE=M053R11
NODE=M053R11

NODE=M053R88
NODE=M053R88

NODE=M053R88;LINKAGE=AD
NODE=M053R89
NODE=M053R89

NODE=M053R89;LINKAGE=AD
NODE=M053R89;LINKAGE=AS

NODE=M053R90
NODE=M053R90

NODE=M053R90;LINKAGE=AD
NODE=M053R90;LINKAGE=AS

NODE=M053R00
NODE=M053R00

NODE=M053R00;LINKAGE=A

$\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{21}/Γ			
$\text{VALUE (units } 10^{-5}\text{)}$	CL\%			
<1.4	90	1 ADAMS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{22}/Γ			
$\text{VALUE (units } 10^{-3}\text{)}$	CL\%			
<1.2	90	1 ADAMS	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$	Γ_{23}/Γ			
$\text{VALUE (units } 10^{-5}\text{)}$	CL\%			
< 1.2	90	1 CRONIN-HEN..06	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	2 ABLIKIM	04F BES	$e^+ e^- \rightarrow \psi(3770)$
¹ Using $\sigma(e^+ e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08^{+0.41}_{-0.30})$ nb from BESSON 06 and $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6895 \pm 0.0014$.				
² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.				

$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$	Γ_{24}/Γ			
$\text{VALUE (units } 10^{-4}\text{)}$	CL\%			
<11.2	90	1 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48	90	2 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.				
² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.				

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$	Γ_{25}/Γ			
$\text{VALUE (units } 10^{-4}\text{)}$	CL\%			
<10.6	90	1 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<62	90	2 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.				
² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.				

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$	Γ_{26}/Γ			
$\text{VALUE (units } 10^{-3}\text{)}$	EVTS			
<58.5	90 305	ABLIKIM	08N BES2	$e^+ e^- \rightarrow \psi(3770)$

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{27}/Γ			
$\text{VALUE (units } 10^{-4}\text{)}$	CL\%			
< 6.0	90	1 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	2 ABLIKIM	07I BES2	$3.77 e^+ e^-$
¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.				
² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.				

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$	Γ_{28}/Γ			
$\text{VALUE (units } 10^{-4}\text{)}$	CL\%			
<91	90	1 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R91
NODE=M053R91

NODE=M053R91;LINKAGE=AS

NODE=M053R92
NODE=M053R92

NODE=M053R92;LINKAGE=AD

NODE=M053R3
NODE=M053R3

NODE=M053R3;LINKAGE=CR

NODE=M053R3;LINKAGE=AB

NODE=M053R21
NODE=M053R21

NODE=M053R21;LINKAGE=HU
NODE=M053R21;LINKAGE=AK

NODE=M053R22
NODE=M053R22

NODE=M053R22;LINKAGE=HU
NODE=M053R22;LINKAGE=AK

NODE=M053R72
NODE=M053R72

NODE=M053R24
NODE=M053R24

NODE=M053R24;LINKAGE=HU
NODE=M053R24;LINKAGE=AK

NODE=M053R07
NODE=M053R07

NODE=M053R07;LINKAGE=AK

$\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{29}/Γ
<137	90	¹ ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{30}/Γ
<117.4	90	59	ABLIKIM	08N	BES2	$e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{31}/Γ
<1.24	90	¹ HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.3	90	² ABLIKIM	10D	BES2	$e^+ e^- \rightarrow \psi(3770)$
----------------	----	----------------------	-----	------	----------------------------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{32}/Γ
<8.9	90	218	ABLIKIM	08N	BES2	$e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{33}/Γ
<6.9	90	¹ ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(\eta 3\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{34}/Γ
<13.4	90	¹ HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

 $\Gamma(\eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{35}/Γ
<243	90	¹ ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(\eta\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{36}/Γ
<1.45	90	¹ ABLIKIM	10D	BES2	$e^+ e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(\eta' 3\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{37}/Γ
<24.4	90	¹ HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

 $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{38}/Γ
< 9.0	90	¹ HUANG	06A	CLEO	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48	90	² ABLIKIM	07B	BES2	$e^+ e^- \rightarrow \psi(3770)$
---------------	----	----------------------	-----	------	----------------------------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

NODE=M053R10
NODE=M053R10

NODE=M053R10;LINKAGE=AK

NODE=M053R74
NODE=M053R74

NODE=M053R23
NODE=M053R23

NODE=M053R23;LINKAGE=HU
NODE=M053R23;LINKAGE=AK

NODE=M053R70
NODE=M053R70

NODE=M053R53
NODE=M053R53

NODE=M053R53;LINKAGE=AK

NODE=M053R25
NODE=M053R25

NODE=M053R25;LINKAGE=HU

NODE=M053R08
NODE=M053R08

NODE=M053R08;LINKAGE=AK

NODE=M053R77
NODE=M053R77

NODE=M053R77;LINKAGE=AK

NODE=M053R26
NODE=M053R26

NODE=M053R26;LINKAGE=HU

NODE=M053R27
NODE=M053R27

NODE=M053R27;LINKAGE=HU
NODE=M053R27;LINKAGE=AK

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.1	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	2 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	-----------	----------	---------------------------------

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-2\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.2	90	14	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$

 $\Gamma(4(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<16.7	90	1 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<30.6	90	1 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<4.5	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 23.6	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<111	90	2 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
------	----	-----------	----------	---------------------------------

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-\rho^0\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<8	90	1 ABLIKIM	07I BES2	3.77 e^+e^-

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-\rho^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<146	90	1 ABLIKIM	07I BES2	3.77 e^+e^-

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\omega K^+K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 3.4	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	2 ABLIKIM	07I BES2	3.77 e^+e^-
-----	----	-----------	----------	---------------

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 Γ_{39}/Γ

NODE=M053R28

NODE=M053R28

NODE=M053R71

NODE=M053R71

NODE=M053R50

NODE=M053R50

NODE=M053R50;LINKAGE=AK

NODE=M053R52

NODE=M053R52

NODE=M053R29

NODE=M053R29

NODE=M053R29;LINKAGE=HU

NODE=M053R30

NODE=M053R30

NODE=M053R30;LINKAGE=HU

NODE=M053R30;LINKAGE=AK

NODE=M053R58

NODE=M053R58

NODE=M053R58;LINKAGE=AK

NODE=M053R59

NODE=M053R59

NODE=M053R59;LINKAGE=AK

NODE=M053R32

NODE=M053R32

NODE=M053R32;LINKAGE=HU

NODE=M053R32;LINKAGE=AK

$\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<38	90	¹ ABLIKIM	07I	BES2 3.77 e^+e^-

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^{*0}K^-\pi^+\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<162	90	¹ ABLIKIM	07I	BES2 3.77 e^+e^-

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-\pi^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<323	90	¹ ABLIKIM	07I	BES2 3.77 e^+e^-

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-\pi^+\pi^- 2\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<26.7	90	24	ABLIKIM	08N	BES2 $e^+e^- \rightarrow \psi(3770)$

 $\Gamma(K^+K^-2(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<10.3	90	¹ ABLIKIM	07F	BES2 $e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<36.0	90	¹ ABLIKIM	07F	BES2 $e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta K^+K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 4.1	90	¹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<31	90	² ABLIKIM	10D	BES2 $e^+e^- \rightarrow \psi(3770)$
---------------	----	----------------------	-----	--------------------------------------

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.24	90	¹ ABLIKIM	10D	BES2 $e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\rho^0 K^+K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5.0	90	¹ ABLIKIM	07F	BES2 $e^+e^- \rightarrow \psi(3770)$

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 6.0	90	¹ HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17	90	² ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$
---------------	----	----------------------	-----	--------------------------------------

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 Γ_{48}/Γ

NODE=M053R60
NODE=M053R60

NODE=M053R60;LINKAGE=AK

NODE=M053R61
NODE=M053R61

NODE=M053R61;LINKAGE=AK

NODE=M053R62
NODE=M053R62

NODE=M053R73
NODE=M053R73

NODE=M053R57
NODE=M053R57

NODE=M053R57;LINKAGE=AK

NODE=M053R51
NODE=M053R51

NODE=M053R51;LINKAGE=AK

NODE=M053R31
NODE=M053R31

NODE=M053R31;LINKAGE=HU
NODE=M053R31;LINKAGE=AK

NODE=M053R78
NODE=M053R78

NODE=M053R78;LINKAGE=AK

NODE=M053R54
NODE=M053R54

NODE=M053R54;LINKAGE=AK

NODE=M053R33
NODE=M053R33

NODE=M053R33;LINKAGE=HU
NODE=M053R33;LINKAGE=AK

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{58}/Γ
< 7.5	90	1 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R34 NODE=M053R34
< 24	90	2 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1 Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.
 2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{59}/Γ
< 2.9	90	1 HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R34;LINKAGE=HU NODE=M053R34;LINKAGE=AK
< 46	90	2 ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	
					NODE=M053R35;LINKAGE=HU NODE=M053R35;LINKAGE=AK

• • • We do not use the following data for averages, fits, limits, etc. • • •

1 Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.
 2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(2(K^+ K^-)\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{60}/Γ
< 3.2	90	1 ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R48 NODE=M053R48

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. $\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{61}/Γ
< 3.2	90	18	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R64 NODE=M053R64

 $\Gamma(K_S^0 K^- \pi^+\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{62}/Γ
< 13.3	90	40	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R65 NODE=M053R65

 $\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{63}/Γ
< 6.6	90		ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R15 NODE=M053R15

 $\Gamma(K_S^0 K^- 2\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{64}/Γ
< 8.7	90	39	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R66 NODE=M053R66

 $\Gamma(K_S^0 K^- \pi^+\rho^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{65}/Γ
< 1.6	90		ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R16 NODE=M053R16

 $\Gamma(K_S^0 K^- \pi^+\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{66}/Γ
< 1.3	90		ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R17 NODE=M053R17

 $\Gamma(K_S^0 K^- 2\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{67}/Γ
< 41.8	90	23	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R67 NODE=M053R67

 $\Gamma(K_S^0 K^- 2\pi^+\pi^-\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{68}/Γ
< 4.8	90		ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R18 NODE=M053R18

 $\Gamma(K_S^0 K^- \pi^+ 2(\pi^+\pi^-))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{69}/Γ
< 12.2	90	4	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R68 NODE=M053R68

$\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{70}/Γ
<26.5	90	17	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R69
NODE=M053R69 $\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{71}/Γ
<4.9	90	ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R19
NODE=M053R19 $\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{72}/Γ
<3.0	90	ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R75
NODE=M053R75 $\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-2})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{73}/Γ
<2.2	90	ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R76
NODE=M053R76 $\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{74}/Γ
<9.7	90	¹ AAIJ	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R55
NODE=M053R55

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{75}/Γ
---	-------------	--------------------	-------------	----------------	----------------------

NODE=M053R98
NODE=M053R98

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		¹ AAIJ	17AD LHCb	$p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p} K^+ X$	
$7.1^{+8.6}_{-2.9}$	684	² ABLIKIM	14L BES3	$e^+ e^- \rightarrow \psi(3770)$	
310 ± 30	684	³ ABLIKIM	14L BES3	$e^+ e^- \rightarrow \psi(3770)$	

OCCUR=2

¹ AAIJ 17AD reports $B(B^+ \rightarrow \psi(3770) K^+ \rightarrow p\bar{p} K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p} K^+) < 0.09$ (0.10) at 90% (95%) CL.

2 Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

3 Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{76}/Γ
< 0.4	90	1,2 ABLIKIM	14O BES3	$e^+ e^- \rightarrow \psi(3770)$	

NODE=M053R09
NODE=M053R09

• • • We do not use the following data for averages, fits, limits, etc. • • •

$59^{+3}_{-2} \pm 5$		^{1,3} ABLIKIM	14O BES3	$e^+ e^- \rightarrow \psi(3770)$	
<12	90	⁴ ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	

OCCUR=2

¹ Calculated by the authors using $\sigma(e^+ e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30} \text{ nb}$ from BESSON 10.

2 Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

3 Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

4 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{77}/Γ
< 5.8	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	

NODE=M053R36
NODE=M053R36

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	² ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$	
-----	----	----------------------	----------	----------------------------------	--

NODE=M053R36;LINKAGE=HU
NODE=M053R36;LINKAGE=AK

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{78}/Γ
<1.2	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$	

NODE=M053R42
NODE=M053R42

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4	90	² ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$	
----	----	----------------------	----------	----------------------------------	--

NODE=M053R42;LINKAGE=HU
NODE=M053R42;LINKAGE=AK

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<18.5	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<73	90	2 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{79}/Γ NODE=M053R37
NODE=M053R37 $\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<30	90	2 ABLIKIM	07I BES2	$3.77 e^+e^-$

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.2 Using $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$ nb and neglecting interference. Γ_{80}/Γ NODE=M053R39
NODE=M053R39 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.7	90	1 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<12	90	2 ABLIKIM	07I BES2	$3.77 e^+e^-$

1 Assuming that interference effects between resonance and continuum can be neglected.

2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{81}/Γ NODE=M053R63
NODE=M053R63 $\Gamma(p\bar{p}2(\pi^+\pi^-))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	1 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{82}/Γ NODE=M053R49
NODE=M053R49 $\Gamma(\eta p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.4	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<11	90	2 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{83}/Γ NODE=M053R38
NODE=M053R38 $\Gamma(\eta p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	1 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{84}/Γ NODE=M053R79
NODE=M053R79 $\Gamma(\rho^0 p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	1 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

1 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{85}/Γ NODE=M053R56
NODE=M053R56 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$ <11 90 2 ABLIKIM 07B BES2 $e^+e^- \rightarrow \psi(3770)$ 1 Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.2 Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb. Γ_{86}/Γ NODE=M053R40
NODE=M053R40NODE=M053R37
NODE=M053R37NODE=M053R39
NODE=M053R39NODE=M053R63
NODE=M053R63NODE=M053R49
NODE=M053R49

NODE=M053R49;LINKAGE=AK

NODE=M053R38
NODE=M053R38

NODE=M053R38;LINKAGE=AK

NODE=M053R79
NODE=M053R79

NODE=M053R79;LINKAGE=AK

NODE=M053R56
NODE=M053R56

NODE=M053R56;LINKAGE=AK

NODE=M053R40
NODE=M053R40

NODE=M053R40;LINKAGE=AK

$\Gamma(p\bar{p}K^+K^-)/\Gamma_{\text{total}}$	Γ_{87}/Γ			
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	1 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

$\Gamma(\pi^0 p\bar{p}K^+K^-)/\Gamma_{\text{total}}$	Γ_{88}/Γ			
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	1 ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

$\Gamma(\phi p\bar{p})/\Gamma_{\text{total}}$	Γ_{89}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9	90	2 ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
----	----	-----------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{90}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.7	90	2 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
<39	90	3 ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected.

³ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$	Γ_{91}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{92}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6.3	90	1 HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6 \text{ nb}$ at the resonance.

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$	Γ_{93}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	1 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$	Γ_{94}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	1 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$	Γ_{95}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	1 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

$\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$	Γ_{96}/Γ			
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	1 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R80
NODE=M053R80

NODE=M053R80;LINKAGE=AK

NODE=M053R81
NODE=M053R81

NODE=M053R81;LINKAGE=AK

NODE=M053R41
NODE=M053R41

NODE=M053R41;LINKAGE=HU
NODE=M053R41;LINKAGE=AK

NODE=M053R43
NODE=M053R43

NODE=M053R43;LINKAGE=HU
NODE=M053R43;LINKAGE=A
NODE=M053R43;LINKAGE=AK

NODE=M053R44
NODE=M053R44

NODE=M053R44;LINKAGE=HU

NODE=M053R45
NODE=M053R45

NODE=M053R45;LINKAGE=HU

NODE=M053R93
NODE=M053R93

NODE=M053R93;LINKAGE=A

NODE=M053R94
NODE=M053R94

NODE=M053R94;LINKAGE=A

NODE=M053R95
NODE=M053R95

NODE=M053R95;LINKAGE=A

NODE=M053R96
NODE=M053R96

NODE=M053R96;LINKAGE=A

$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{97}/Γ
<1.4	90	¹ ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(3770)$	

¹ Assuming that interference effects between resonance and continuum can be neglected.

RADIATIVE DECAYS $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{98}/Γ
<0.64	90	¹ ABLIKIM	15J	BES3 $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma \gamma J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0	90	² BRIERE	06	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
<0.9	90	³ COAN	06A	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma \gamma J/\psi$	

¹ This limit is equivalent to $(0.25 \pm 0.21 \pm 0.18) \times 10^{-3}$ branching fraction value.

² Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

³ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+ e^- \rightarrow D \bar{D})$ from HE 05 for $\sigma(e^+ e^- \rightarrow \psi(3770))$.

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{99}/Γ
2.49 ± 0.23 OUR AVERAGE					

1.98 ± 0.78 ± 0.05	202	¹ ABLIKIM	16B	BES3 $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
2.48 ± 0.15 ± 0.23	0.6k	ABLIKIM	15J	BES3 $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma \gamma J/\psi$	
2.4 ± 0.8 ± 0.2		² ABLIKIM	14H	BES3 $e^+ e^- \rightarrow \psi(3770) \rightarrow K_S^0 K^\pm \pi^\mp$	
2.9 ± 0.5 ± 0.4		³ BRIERE	06	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma \gamma J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.9 ± 1.4 ± 0.6	54	⁴ BRIERE	06	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
2.8 ± 0.5 ± 0.4	53	⁵ COAN	06A	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma \gamma J/\psi$	OCCUR=2

¹ ABLIKIM 16B reports $(1.94 \pm 0.42 \pm 0.64) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma \chi_{c1})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma \chi_{c1})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)] = (8.51 \pm 2.39 \pm 1.42) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp) = 0.00349 \pm 0.00029$. Our first error is their experiment's error and our second error is the systematic error from using our best value. We have calculated the best value of $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/2 of $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

³ Averages the two measurements from COAN 06A and BRIERE 06.

⁴ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

⁵ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+ e^- \rightarrow D \bar{D})$ from HE 05 for $\sigma(e^+ e^- \rightarrow \psi(3770))$.

 $\Gamma(\gamma \chi_{c1})/\Gamma(J/\psi \pi^+ \pi^-)$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{99}/Γ_4
1.49 ± 0.31 ± 0.26	53 ± 10	¹ COAN	06A	CLEO $e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma \gamma J/\psi$	

¹ Using $B(\psi(3770) \rightarrow J/\psi \pi^+ \pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$ from ADAM 06.

NODE=M053R97
NODE=M053R97

NODE=M053R97;LINKAGE=A

NODE=M053240

NODE=M053R03
NODE=M053R03

NODE=M053R03;LINKAGE=A

NODE=M053R03;LINKAGE=BR

NODE=M053R03;LINKAGE=CO

NODE=M053R02
NODE=M053R02

NODE=M053R02;LINKAGE=A

NODE=M053R02;LINKAGE=AB

NODE=M053R02;LINKAGE=BI

NODE=M053R02;LINKAGE=BR

NODE=M053R02;LINKAGE=CO

NODE=M053R04
NODE=M053R04

NODE=M053R04;LINKAGE=CO

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$				Γ_{100}/Γ	
<u>VALUE</u> (units 10^{-3})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.9 ± 0.6 OUR AVERAGE					
6.7 $\pm 0.7 \pm 0.1$	2.2K	1	ABLIKIM	16B BES3	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
7.3 $\pm 0.7 \pm 0.6$	274		BRIERE	06 CLEO	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 44	90	2	COAN	06A CLEO	$e^+ e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

¹ ABLIKIM 16B reports $(6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c0})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+ e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+ e^- \rightarrow \psi(3770))$.

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$				Γ_{100}/Γ_{98}	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>8	90	1	BRIERE	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$

¹ Not independent of other results in BRIERE 06.

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$				Γ_{100}/Γ_{99}	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.5 ± 0.6	1	BRIERE	06 CLEO	$e^+ e^- \rightarrow \psi(3770)$	

¹ Not independent of other results in BRIERE 06.

$\Gamma(\gamma\eta_c)/\Gamma_{\text{total}}$				Γ_{101}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 7 \times 10^{-4}$	90	1	ABLIKIM	14H BES3	
¹ ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 16 \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = 2.43 \times 10^{-2}$. We have calculated the best value of $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = 7.3 \times 10^{-2}$.					

$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$				Γ_{102}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$< 9 \times 10^{-4}$	90	1	ABLIKIM	14H BES3	
¹ ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 5.6 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) = 6 \times 10^{-3}$. We have calculated the best value of $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$.					

$\Gamma(\gamma\eta')/\Gamma_{\text{total}}$				Γ_{103}/Γ	
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.8	90	1	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
¹ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.					

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$				Γ_{104}/Γ	
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.5	90	1	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
¹ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.					

$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$				Γ_{105}/Γ	
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2	90	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$	

NODE=M053R01
NODE=M053R01

NODE=M053R01;LINKAGE=B

NODE=M053R01;LINKAGE=CO

NODE=M053R06

NODE=M053R06;LINKAGE=BR

NODE=M053R05

NODE=M053R05

NODE=M053R99;LINKAGE=AB

NODE=M053R20

NODE=M053R20

NODE=M053R20;LINKAGE=AB

NODE=M053R14

NODE=M053R14

NODE=M053R13;LINKAGE=PE

NODE=M053R13

NODE=M053R13

NODE=M053R12

NODE=M053R12

$\psi(3770)$ REFERENCES

NODE=M053

AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59697
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
SHAMOV	17	PL B769 187	A.G. Shamov, K.Yu. Todyshev		REFID=57900
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57126
ABLIKIM	15J	PR D91 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56775
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)	REFID=56962
ABLIKIM	14H	PR D89 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55899
ABLIKIM	14L	PL B735 101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55903
ABLIKIM	14O	PR D90 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55906
BONVICINI	14	PR D89 072002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=55798
ABLIKIM	13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55393
ANASHIN	12A	PL B711 292	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54055
ABLIKIM	10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53350
BESSON	10	PRL 104 159901 (errat.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
ABLIKIM	09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=53134
PEDALAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
ABLIKIM	08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52569
ABLIKIM	08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52570
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)	REFID=52144
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
ABLIKIM	07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51704
ABLIKIM	07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51882
ABLIKIM	07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51940
ABLIKIM	07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52045
ABLIKIM	07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52073
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52075
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51131
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
ADAMS	06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50990
BESSON	06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51041
Also		PRL 104 159901 (errat.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
BRIERE	06	PR D74 031106	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=51149
COAN	06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51155
CRONIN-HEN... 06		PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=51156
HUANG	06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=50999
BAI	05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50332
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=50924
Also		PRL 96 199903 (errat.)	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51211
ABLIKIM	04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50185
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40361
SCHINDLER	80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)	REFID=22222
BACINO	78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)	REFID=11437
RAPIDIS	77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)	REFID=22220

$\psi_2(3823)$

$I^G(J^{PC}) = 0^-(2^{--})$
 I, J, P need confirmation.

was $\psi(3823)$, $X(3823)$

Seen by BHARDWAJ 13 in $B \rightarrow \chi_{c1}\gamma K$ and ABLIKIM 15S in $e^+e^- \rightarrow \pi^+\pi^-\gamma\chi_{c1}$ decays as a narrow peak in the invariant mass distribution of the $\chi_{c1}\gamma$ system. Properties consistent with the $\psi_2(1^3D_2) c\bar{c}$ state.

$\psi_2(3823)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3822.2±1.2 OUR AVERAGE				
3821.7±1.3±0.7	19 ± 5	1 ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
3823.1±1.8±0.7	33 ± 10	2 BHARDWAJ	13 BELL	$B \rightarrow \chi_{c1}\gamma K$

- 1 From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c1}\gamma$ together, with floating mass scale offset for $\psi(2S)$, floating $\psi_2(3823)$ mass, and zero $\psi_2(3823)$ width, resulting in a significance of 5.9σ when including systematic uncertainties.
- 2 From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$ and $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$ with significance 4.0σ including systematics. Corrected for the measured $\psi(2S)$ mass using $B \rightarrow \psi(2S)K \rightarrow (\gamma\chi_{c1})K$ decays.

$\psi_2(3823)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	1 ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<24	90	2 BHARDWAJ	13 BELL	$B \rightarrow \chi_{c1}\gamma K$
1 From a fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.				
2 From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$ and $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$ with significance 4.0σ including systematics.				

$\psi_2(3823)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \chi_{c1}\gamma$	seen
$\Gamma_2 \chi_{c2}\gamma$	not seen

$\psi_2(3823)$ BRANCHING RATIOS

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	$B^+ \rightarrow \chi_{c1}\gamma K^+$
1 Reported $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$ with statistical significance 3.8σ .	

$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$	Γ_2/Γ
not seen	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
not seen	$B^+ \rightarrow \chi_{c2}\gamma K^+$

- 1 From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c2}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c2}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.
- 2 Reported $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c2}) < 3.6 \times 10^{-6}$ at 90% CL.

NODE=M212

NODE=M212

NODE=M212M

NODE=M212M

NODE=M212M;LINKAGE=B

NODE=M212M;LINKAGE=A

NODE=M212W

NODE=M212W

NODE=M212W;LINKAGE=B

NODE=M212W;LINKAGE=A

NODE=M212215;NODE=M212

DESIG=1

DESIG=2

NODE=M212225

NODE=M212R01

NODE=M212R01

NODE=M212R01;LINKAGE=A

NODE=M212R02

NODE=M212R02

NODE=M212R02;LINKAGE=B

NODE=M212R02;LINKAGE=A

$\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
<0.41	90	BHARDWAJ	13	BELL $B^+ \rightarrow \chi_{c1/c2}\gamma K^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.42	90	¹ ABLIKIM	15S	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1}\gamma$	
-------	----	----------------------	-----	--	--

¹ From a simultaneous unbinned maximum likelihood fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1}(2)\gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1}(2)\gamma$ and $\psi_2(3823) \rightarrow \chi_{c1}(2)\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.

NODE=M212R03

NODE=M212R03

NODE=M212R03;LINKAGE=A

 $\psi_2(3823)$ REFERENCES

ABLIKIM BHARDWAJ	15S 13	PRL 115 011803 PRL 111 032001	M. Ablikim <i>et al.</i> V. Bhardwaj <i>et al.</i>	(BESIII Collab.) (BELLE Collab.)
---------------------	-----------	----------------------------------	---	-------------------------------------

 $\psi_3(3842)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

J, P need confirmation.

J^P has not been measured, 3^- is the quark model prediction.

 $\psi_3(3842)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
3842.71±0.16±0.12	AAIJ	19M	LHCb $p p \rightarrow D \bar{D} +$ anything	

 $\psi_3(3842)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
2.79±0.51±0.35	AAIJ	19M	LHCb $p p \rightarrow D \bar{D} +$ anything	

 $\psi_3(3842)$ DECAY MODES

Mode	Γ_1/Γ	Γ_2/Γ
Γ_1 $D^+ D^-$		
Γ_2 $D^0 \bar{D}^0$		

 $\psi_3(3842)$ BRANCHING RATIOS

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	Γ_1/Γ
---	-------------------

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$	Γ_2/Γ
---	-------------------

$\psi_3(3842)$ REFERENCES	Γ_2/Γ_1
AAIJ 19M JHEP 1907 035	R. Aaij <i>et al.</i> (LHCb Collab.)

NODE=M212

REFID=56784

REFID=55412

NODE=M241

NODE=M241

NODE=M241M

NODE=M241M

NODE=M241W

NODE=M241W

NODE=M241215;NODE=M241

DESIG=1

DESIG=2

NODE=M241225

NODE=M241R01

NODE=M241R01

NODE=M241R02

NODE=M241R02

NODE=M241

REFID=59697

NODE=M237

 $\chi_{c0}(3860)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Observed by CHILIKIN 17 using full amplitude analysis of the process
 $e^+ e^- \rightarrow J/\psi D\bar{D}$, where $D = D^0, D^+$.

NODE=M237

 $\chi_{c0}(3860)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3862 +26 +40 -32 -13	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D\bar{D}$

NODE=M237M

NODE=M237M

 $\chi_{c0}(3860)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
201 +154 +88 -67 -82	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D\bar{D}$

NODE=M237W

NODE=M237W

 $\chi_{c0}(3860)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0 \bar{D}^0$	seen
$\Gamma_2 D^+ D^-$	seen

NODE=M237215; NODE=M237

DESIG=1

DESIG=2

 $\chi_{c0}(3860)$ BRANCHING RATIOS

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D^0 \bar{D}^0$	
$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D^+ D^-$	

NODE=M237220

NODE=M237R00
NODE=M237R00NODE=M237R01
NODE=M237R01 **$\chi_{c0}(3860)$ REFERENCES**

CHILIKIN 17 PR D95 112003 K. Chilikin *et al.* (BELLE Collab.) JPC

NODE=M237

REFID=57995

NODE=M176

 $\chi_{c1}(3872)$

$I^G(J^{PC}) = 0^+(1^{++})$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to < 4% at 95% CL.

See our note on "Developments in Heavy Quarkonium Spectroscopy".

 $\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

NODE=M176M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3871.69 ± 0.17 OUR AVERAGE				
3871.9 ± 0.7 ± 0.2	20 ± 5	ABLIKIM 14	BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
3871.95 ± 0.48 ± 0.12	0.6k	AAIJ 12H	LHCb	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3871.85 ± 0.27 ± 0.19	~ 170	¹ CHOI 11	BELL	$B \rightarrow K\pi^+\pi^-J/\psi$
3873 ± 1.8 ± 1.3	27 ± 8	² DEL-AMO-SA.10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61 ± 0.16 ± 0.19	6k	^{2,3} AALTONEN 09AU	CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3871.4 ± 0.6 ± 0.1	93.4	AUBERT 08Y	BABR	$B^+ \rightarrow K^+J/\psi\pi^+\pi^-$
3868.7 ± 1.5 ± 0.4	9.4	AUBERT 08Y	BABR	$B^0 \rightarrow K_S^0J/\psi\pi^+\pi^-$
3871.8 ± 3.1 ± 3.0	522	^{2,4} ABAZOV 04F	D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3873.3 ± 1.1 ± 1.0	45	⁵ ABLIKIM 19V	BES	$e^+e^- \rightarrow \gamma\omega J/\psi$
3860.0 ± 10.4	13.6	^{2,6} AGHASYAN 18A	COMP	$\gamma^*N \rightarrow X\pi^\pm N'$
3868.6 ± 1.2 ± 0.2	8	⁷ AUBERT 06	BABR	$B^0 \rightarrow K_S^0J/\psi\pi^+\pi^-$
3871.3 ± 0.6 ± 0.1	61	⁷ AUBERT 06	BABR	$B^- \rightarrow K^-J/\psi\pi^+\pi^-$
3873.4 ± 1.4	25	⁸ AUBERT 05R	BABR	$B^+ \rightarrow K^+J/\psi\pi^+\pi^-$
3871.3 ± 0.7 ± 0.4	730	^{2,9} ACOSTA 04	CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3872.0 ± 0.6 ± 0.5	36	¹⁰ CHOI 03	BELL	$B \rightarrow K\pi^+\pi^-J/\psi$
3836 ± 13	58	^{2,11} ANTONIAZZI 94	E705	$300\pi^\pm Li \rightarrow J/\psi\pi^+\pi^-X$

¹The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.

²Width consistent with detector resolution.

³A possible equal mixture of two states with a mass difference greater than 3.6 MeV/c² is excluded at 95% CL.

⁴Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.

⁵Fit with fixed width and including two resonances, $X(3915)$ and $X(3960)$.

⁶Could be a different state.

⁷Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.

⁸Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.

⁹Superseded by AALTONEN 09AU.

¹⁰Superseded by CHOI 11.

¹¹A lower mass value can be due to an incorrect momentum scale for soft pions.

NODE=M176

NODE=M176M

NODE=M176M

OCCUR=2

OCCUR=2

NODE=M176M;LINKAGE=CO

NODE=M176M;LINKAGE=AC

NODE=M176M;LINKAGE=AA

NODE=M176M;LINKAGE=AB

NODE=M176M;LINKAGE=B

NODE=M176M;LINKAGE=A

NODE=M176M;LINKAGE=AE

NODE=M176M;LINKAGE=AU

NODE=M176M;LINKAGE=AT

NODE=M176M;LINKAGE=CH

NODE=M176M;LINKAGE=AN

$\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3872.9 ^{+0.6+0.4} _{-0.4-0.5}	50	1,2 AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
3875.1 ^{+0.7} _{-0.5} ^{±0.5}	33 ± 6	2 AUBERT	08B BABR $B \rightarrow \bar{D}^{*0} D^0 K$	
3875.2 ^{+0.7+0.9} _{-1.8}	24 ± 6	2,3 GOKHROO	06	BELL $B \rightarrow D^0 \bar{D}^0 \pi^0 K$
¹ Calculated from the measured $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{\bar{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.				
² Experiments report $D^{*0} \bar{D}^0$ invariant mass above $D^{*0} \bar{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.				
³ Superseded by AUSHEV 10.				

$m_{\chi_{c1}(3872)} - m_{J/\psi}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
774.9^{±3.1}_{±3.0}	522	ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

$m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
187.4 ± 1.4	25	1 AUBERT	05R BABR $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$	
¹ Superseded by AUBERT 06.				

$\chi_{c1}(3872)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.2	90		CHOI	11	BELL $B \rightarrow K \pi^+ \pi^- J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.4	90		ABLIKIM	14	BES3 $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
<3.3	90		AUBERT	08Y BABR $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$	
<4.1	90	69	AUBERT	06	BABR $B \rightarrow K \pi^+ \pi^- J/\psi$
<2.3	90	36	¹ CHOI	03	BELL $B \rightarrow K \pi^+ \pi^- J/\psi$

¹ Superseded by CHOI 11.

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

3.9 ^{+2.8+0.2} _{-1.4-1.1}	50	¹ AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
3.0 ^{+1.9} _{-1.4} ^{±0.9}	33 ± 6	AUBERT	08B BABR $B \rightarrow \bar{D}^{*0} D^0 K$	

¹ With a measured value of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

$\chi_{c1}(3872)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	
$\Gamma_2 \pi^+ \pi^- J/\psi(1S)$	> 3.2 %
$\Gamma_3 \rho^0 J/\psi(1S)$	
$\Gamma_4 \omega J/\psi(1S)$	> 2.3 %
$\Gamma_5 D^0 \bar{D}^0 \pi^0$	> 40 %
$\Gamma_6 \bar{D}^{*0} D^0$	> 30 %
$\Gamma_7 \gamma\gamma$	
$\Gamma_8 D^0 \bar{D}^0$	
$\Gamma_9 D^+ D^-$	

NODE=M176MD0

NODE=M176MD0

NODE=M176MD0;LINKAGE=AS
NODE=M176MD0;LINKAGE=AU

NODE=M176MD0;LINKAGE=GO

NODE=M176207

NODE=M176DM

NODE=M176DM2

NODE=M176DM2

NODE=M176DM2;LINKAGE=AU

NODE=M176W

NODE=M176W

NODE=M176W;LINKAGE=CH

NODE=M176WD0

NODE=M176WD0

NODE=M176WD0;LINKAGE=AU

NODE=M176215;NODE=M176

DESIG=1

DESIG=2

DESIG=10

DESIG=13

DESIG=8

DESIG=12

DESIG=5

DESIG=6

DESIG=7

Γ_{10}	$\gamma\chi_{c1}$					DESIG=3
Γ_{11}	$\gamma\chi_{c2}$					DESIG=15
Γ_{12}	$\pi^0\chi_{c2}$					DESIG=20
Γ_{13}	$\pi^0\chi_{c1}$					DESIG=18
Γ_{14}	$\pi^0\chi_{c0}$					DESIG=19
Γ_{15}	$\gamma J/\psi$					DESIG=9
Γ_{16}	$\gamma\psi(2S)$					DESIG=11
Γ_{17}	$\pi^+\pi^-\eta_c(1S)$					DESIG=14;OUR EVAL; \rightarrow UNCHECKED \leftarrow
Γ_{18}	$\pi^+\pi^-\chi_{c1}$					DESIG=17
Γ_{19}	$p\bar{p}$					DESIG=16
Γ_{20}	$\eta J/\psi$					NODE=M176;CLUMP=A DESIG=4

C-violating decays

Γ_{20} $\eta J/\psi$

$\chi_{c1}(3872)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_1
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 4.3	90	1 ABLIKIM	15v BES3	$4.0-4.4 e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
<280	90	2 YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
1 ABLIKIM 15v reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-)/\Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma = 3\%$.					
2 Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. Assuming that $\Gamma(\pi^+\pi^- J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).					

$\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
< 0.13					
< 0.13	90	ABLIKIM	15v BES3	$4.0-4.4 e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
< 6.2	90	1,2 AUBERT	05D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$	
< 8.3	90	2 DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
<10	90	3 YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
1 Using $B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.					
2 Assuming $\chi_{c1}(3872)$ has $JPC = 1^{--}$.					
3 Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$.					

$\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_7/\Gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					

<12.9 90 1 DOBBS 05 CLE3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi\gamma$

1 Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.

$\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_4\Gamma_7/\Gamma$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					

<1.7 90 1 LEES 12AD BABR $e^+e^- \rightarrow e^+e^-\omega J/\psi$

1 Assuming $\chi_{c1}(3872)$ has spin 2.

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{17}\Gamma_7/\Gamma$
<11.1					

NODE=M176220

NODE=M176W1
NODE=M176W1
NODE=M176W1;LINKAGE=B
NODE=M176W1;LINKAGE=A

NODE=M176230

NODE=M176G1
NODE=M176G1

NODE=M176G1;LINKAGE=AU

NODE=M176G1;LINKAGE=DO
NODE=M176G1;LINKAGE=A

NODE=M176232

NODE=M176H1
NODE=M176H1

NODE=M176H1;LINKAGE=DO

NODE=M176G01
NODE=M176G01

NODE=M176G01;LINKAGE=LE

NODE=M176G02
NODE=M176G02

$\chi_{c1}(3872)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^- J/\psi(1S))/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
>0.032	93 ± 17	1 AUBERT	08Y BABR	$B \rightarrow \chi_{c1}(3872) K$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	151	2 BALA	15 BELL	$B \rightarrow \chi_{c1}(3872) K \pi$	
>0.05	30	3 AUBERT	05R BABR	$B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$	
>0.05	36 ± 7	4 CHOI	03 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^- J/\psi$	
¹ AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					
² BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872) K^+ \pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872) K^0 \pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.					
³ Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					
⁴ CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] / [B(B^+ \rightarrow \psi(2S) K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S) K^+) = (6.19 \pm 0.22) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (34.68 \pm 0.30) \times 10^{-2}$.					

$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
>0.023	21 ± 7	1 DEL-AMO-SA..10B BABR	$B^+ \rightarrow \omega J/\psi K^+$		
¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872) K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi \omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.					

$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$					Γ_4/Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT		
1.1±0.4 OUR AVERAGE	Error includes scale factor of 1.7. [0.8 ± 0.3 OUR 2019 AVERAGE]				
¹ Fit with fixed width and including two resonances, $X(3915)$ and $X(3960)$.					
² Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.					

$\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
>0.4	17 ± 5	1 GOKHROO	06 BELL	$B^+ \rightarrow D^0 \bar{D}^0 \pi^0 K^+$	
¹ GOKHROO 06 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					

$\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$					Γ_5/Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	1 GOKHROO	06 BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	AUSHEV	10 BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$		
¹ May not necessarily be the same state as that observed in the $J/\psi \pi^+ \pi^-$ mode. Supersedes CHISTOV 04.					

NODE=M176235

NODE=M176R6

NODE=M176R6

NODE=M176R6;LINKAGE=AB

NODE=M176R6;LINKAGE=A

NODE=M176R6;LINKAGE=AE

NODE=M176R6;LINKAGE=CH

NODE=M176R14
NODE=M176R14

NODE=M176R14;LINKAGE=DE

NODE=M176R15
NODE=M176R15
NEWNODE=M176R15;LINKAGE=A
NODE=M176R15;LINKAGE=DENODE=M176R12
NODE=M176R12

NODE=M176R12;LINKAGE=GO

NODE=M176R5
NODE=M176R5

NODE=M176R5;LINKAGE=GO

$\Gamma(\bar{D}^{*0} D^0)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
>0.30	41^{+9}_{-8}	1 AUSHEV	10 BELL	$B^+ \rightarrow \bar{D}^{*0} D^0 K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.6	27 ± 6	2 AUBERT	08B BABR	$B^+ \rightarrow \bar{D}^{*0} D^0 K^+$	
1 AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					
2 AUBERT 08B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					

NODE=M176R13
NODE=M176R13

NODE=M176R13;LINKAGE=AS

NODE=M176R13;LINKAGE=AU

 $\Gamma(D^0 \bar{D}^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	CHISTOV	04 BELL	$B \rightarrow K D^0 \bar{D}^0$	

NODE=M176R3
NODE=M176R3 $\Gamma(D^+ D^-)/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	CHISTOV	04 BELL	$B \rightarrow K D^+ D^-$	

NODE=M176R4
NODE=M176R4 $\Gamma(\gamma \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ_2
not seen		1 BHARDWAJ	13 BELL	$B^+ \rightarrow \chi_{c1} \gamma K^+$	
<0.89	90	CHOI	03 BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$	
1 Reported $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1}) < 1.9 \times 10^{-6}$ at 90% CL.					

NODE=M176R1
NODE=M176R1

NODE=M176R1;LINKAGE=A

 $\Gamma(\gamma \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S))$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_2
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	1 BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2} \gamma K^\pm$	
1 Reported $B(B^\pm \rightarrow \chi_{c1}(3872) K^\pm) \times B(\chi_{c1}(3872) \rightarrow \gamma \chi_{c2}) < 6.7 \times 10^{-6}$ at 90% CL.				

NODE=M176R01
NODE=M176R01

NODE=M176R01;LINKAGE=A

 $\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
>7 $\times 10^{-3}$		1 BHARDWAJ	11 BELL	$B^\pm \rightarrow \gamma J/\psi K^\pm$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.011	20	2 AUBERT	09B BABR	$B^+ \rightarrow \gamma J/\psi K^+$	
>0.013	19	3 AUBERT,BE	06M BABR	$B^+ \rightarrow \gamma J/\psi K^+$	
1 BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					
2 AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					
3 Superseded by AUBERT 09B. AUBERT,BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					

NODE=M176R7
NODE=M176R7

NODE=M176R7;LINKAGE=BA

NODE=M176R7;LINKAGE=AB

NODE=M176R7;LINKAGE=AU

 $\Gamma(\gamma \psi(2S))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
seen	36 ± 9	1 AAIJ	14AH LHCb	$B^+ \rightarrow \gamma \psi(2S) K^+$	
>0.04	25 ± 7	2 AUBERT	09B BABR	$B^+ \rightarrow \gamma \psi(2S) K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen		3 BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma \psi(2S) K^+$	
1 From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi \gamma$ decays with a statistical significance of 4.4σ .					
2 AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) < 2.6 \times 10^{-4}$.					
3 BHARDWAJ 11 reports $B(B^+ \rightarrow K^+ \chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma \psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.					

NODE=M176R10
NODE=M176R10

NODE=M176R10;LINKAGE=A

NODE=M176R10;LINKAGE=AU

NODE=M176R10;LINKAGE=BH

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$					Γ_{16}/Γ_{15}	
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
2.6 ± 0.6 OUR AVERAGE						NODE=M176R11 NODE=M176R11
2.46 ± 0.64 ± 0.29	36 ± 9	1	AAIJ	14AH LHCb	$B^+ \rightarrow \gamma\psi(2S)K^+$	
3.4 ± 1.4		AUBERT	09B BABR		$B^+ \rightarrow \gamma c\bar{c}K'$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<2.1	90	BHARDWAJ	11	BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$	
1 From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ .						NODE=M176R11;LINKAGE=A
$\Gamma(\pi^+\pi^-\chi_{c1})/\Gamma_{\text{total}}$					Γ_{18}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		1	BHARDWAJ	16	BELL $B^+ \rightarrow \pi^+\pi^-\chi_{c1}K^+$	NODE=M176R00 NODE=M176R00
1 BHARDWAJ 16 quotes $B(B^+ \rightarrow \chi_{c1}(3872)K^+) \cdot B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\chi_{c1}) < 1.5 \times 10^{-6}$ at 90% CL.						NODE=M176R00;LINKAGE=A
$\Gamma(p\bar{p})/\Gamma_{\text{total}}$					Γ_{19}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		1	AAIJ	17AD LHCb	$p\bar{p} \rightarrow B^+X \rightarrow p\bar{p}K^+X$	NODE=M176R03 NODE=M176R03
1 AAIJ 17AD reports $B(B^+ \rightarrow \chi_{c1}(3872)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) < 2.0 (2.5) \times 10^{-3}$ at 90% (95%) CL.						NODE=M176R03;LINKAGE=A
$\Gamma(p\bar{p})/\Gamma(\pi^+\pi^-J/\psi(1S))$					Γ_{19}/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.0 × 10⁻³	95	1	AAIJ	13S LHCb	$B^+ \rightarrow p\bar{p}K^+$	NODE=M176R02 NODE=M176R02
1 AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow p\bar{p})/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$.						NODE=M176R02;LINKAGE=A
$\Gamma(\pi^0\chi_{c0})/\Gamma(\pi^+\pi^-J/\psi(1S))$					Γ_{14}/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<19	90	ABLIKIM	19U	BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$	NODE=M176R04 NODE=M176R04
$\Gamma(\pi^0\chi_{c1})/\Gamma(\pi^+\pi^-J/\psi(1S))$					Γ_{13}/Γ_2	
VALUE (units 10 ⁻²)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
88⁺³³₋₂₇ ± 10	10.8	ABLIKIM	19U	BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$	NODE=M176R05 NODE=M176R05
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<97	90	1	BHARDWAJ	19	BELL $B^\pm \rightarrow \chi_{c1}\pi^0K^\pm$	
1 BHARDWAJ 19 reports $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \pi^0\chi_{c1}) < 8.1 \times 10^{-6}$ at 90% CL which was divided by $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) = (8.63 \pm 0.97) \times 10^{-6}$ from CHOI 11.						NODE=M176R05;LINKAGE=E
$\Gamma(\pi^0\chi_{c2})/\Gamma(\pi^+\pi^-J/\psi(1S))$					Γ_{12}/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.1	90	ABLIKIM	19U	BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$	NODE=M176R06 NODE=M176R06
C-violating decays						
$\Gamma(\eta J/\psi)/\Gamma(\pi^+\pi^-J/\psi(1S))$					Γ_{20}/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.4	90	1,2 IWASHITA	14	BELL	$B \rightarrow K\eta J/\psi$	NODE=M176R2 NODE=M176R2
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.6	90	AUBERT	04Y	BABR	$B \rightarrow K\eta J/\psi$	
1 IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+, \chi_{c1} \rightarrow J/\psi\pi^+\pi^-) = 8.6 \times 10^{-6}$.						NODE=M176R2;LINKAGE=A
2 IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.						NODE=M176R2;LINKAGE=C

$\chi_{c1}(3872)$ REFERENCES

							NODE=M176
ABLIKIM	19U	PRL 122 202001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59786		
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796		
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=59884		
AGHASYAN	18A	PL B783 334	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59036		
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896		
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=57272		
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56771		
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56787		
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)	REFID=56408		
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55897		
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55647		
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)	REFID=55925		
AAIJ	13Q	PR L10 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=54985		
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55008		
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=55412		
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056		
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54751		
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752		
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=53779		
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=53934		
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=53225		
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53362		
AALTTONEN	09AU	PRL 103 152001	T. Altonen <i>et al.</i>	(CDF Collab.)	REFID=53098		
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52722		
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120		
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52265		
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017		
AUBERT,BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51454		
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)	REFID=51432		
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50498		
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509		
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50627		
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=50458		
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=50200		
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=49742		
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49997		
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002		
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653		
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=49677		
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BES Collab.)	REFID=49628		
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(E705 Collab.)	REFID=46339		
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074		

 $Z_c(3900)$

$I^G(J^{PC}) = 1^+(1^{+-})$

was $X(3900)$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

NODE=M210

Charged $Z_c(3900)$ seen as a peak in the invariant mass distribution of the $J/\psi\pi^\pm$ system by BES III (ABLIKIM 13T) in $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ at c.m. energy of 4.26 GeV and by radiative return from e^+e^- collisions at \sqrt{s} from 9.46 to 10.86 GeV at Belle (LIU 13B). Partial wave analysis of ABLIKIM 17J determines $J^P = 1^+$ with more than 7σ significance. Neutral $Z_c(3900)$ seen in the $J/\psi\pi^0$ invariant mass distribution in $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$ at c.m. energies of 4.23, 4.26, and 4.36 GeV by BES III (ABLIKIM 15U) and at 4.17 GeV by XIAO 13A. Peaks in $(D\bar{D}^*)^{0,\pm}$ reported by BES III (ABLIKIM 14A, ABLIKIM 15AB) are assumed to be related.

NODE=M210

 $Z_c(3900)$ MASS

NODE=M210M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
3888.4±2.5 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below. [3887.2 ± 2.3 MeV OUR 2019 AVERAGE Scale factor = 1.6]				NODE=M210M
3902.6 ^{+5.2} _{-5.0} ^{+3.3} _{-1.4}	1	ABAZOV	19	D0	1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$	
3895.0 ^{+5.2} _{-2.7} ^{+4.0} _{-2.7}	502	ABAZOV	18B	D0	1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$	
3885.7 ^{+4.3} _{-5.7} ^{+8.4}		ABLIKIM	15AB	BES3	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$	
3881.7 ^{+1.6} _{-1.6}	1.2k	ABLIKIM	15AC	BES3	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$	

NODE=M210M


NEW

3894.8 \pm 2.3 \pm 3.2	356	³ ABLIKIM	15U BES3	0	$e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$	
3883.9 \pm 1.5 \pm 4.2	1.2k	³ ABLIKIM	14A BES3	\pm	$e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$	
3899.0 \pm 3.6 \pm 4.9	307	³ ABLIKIM	13T BES3	\pm	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
3894.5 \pm 6.6 \pm 4.5	159	³ LIU	13B BELL	\pm	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
3886 \pm 4 \pm 2	81	^{3,4} XIAO	13A	\pm	$4.17 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	OCCUR=2
3904 \pm 9 \pm 5	25	^{3,4} XIAO	13A	0	$4.17 e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

3881.2 \pm 4.2 \pm 52.7 6k ⁵ ABLIKIM 17J BES3 \pm $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

1 Measured in weak decays of b -flavored hadrons (nonprompt).
 2 The signal of the $Z_c(3900)$ is correlated with a parent $J/\psi \pi^+ \pi^-$ system in the invariant mass range 4.2–4.7 GeV.
 3 Neglecting interference between the $Z_c(3900)$ and non-resonant continuum.
 4 For $M^2(\pi^+ \pi^-) < 0.65$ GeV 2 . Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
 5 Pole mass obtained from a fit to a Flatte-like formula.



$Z_c(3900)$ WIDTH						
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
28.3 \pm 2.5 OUR AVERAGE						NODE=M210W
[28.2 \pm 2.6 MeV OUR 2019 AVERAGE]						NODE=M210W
28.2 \pm 2.6						NEW
32 \pm 28 \pm 26		1 ABAZOV	19 D0		1.96 TeV $p\bar{p} \rightarrow \pi^+ \pi^- J/\psi X$ (non-prompt)	
-21 $-$ 7					$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
51.8 \pm 4.6 \pm 36.0	6 k	2 ABLIKIM	17J BES3	\pm	$e^+ e^- \rightarrow \pi^0 (D\bar{D}^*)^0$	
35 \pm 11 \pm 15		3 ABLIKIM	15AB BES3	0	$e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$	
26.6 \pm 2.0 \pm 2.1	1248	3 ABLIKIM	15AC BES3	\pm	$e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$	
29.6 \pm 8.2 \pm 8.2	356	3 ABLIKIM	15U BES3	0	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
24.8 \pm 3.3 \pm 11.0	1212	3 ABLIKIM	14A BES3	\pm	$e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$	
46 \pm 10 \pm 20	307	3 ABLIKIM	13T BES3	\pm	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
63 \pm 24 \pm 26	159	3 LIU	13B BELL	\pm	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
37 \pm 4 \pm 8	81	^{3,4} XIAO	13A	\pm	$4.17 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	

- 1 Measured in weak decays of b -flavored hadrons (nonprompt).
 2 Pole width obtained from a fit to a Flatte-like formula.
 3 Neglecting interference between the $Z_c(3900)$ and non-resonant continuum.
 4 For $M^2(\pi^+ \pi^-) < 0.65$ GeV 2 . Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

Z_c(3900) DECAY MODES

NODE=M210215;NODE=M210

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 J/\psi \pi$	seen
$\Gamma_2 h_c \pi^\pm$	not seen
$\Gamma_3 \eta_c \pi^+ \pi^-$	not seen
$\Gamma_4 \eta_c(1S) \rho(770)^\pm$	
$\Gamma_5 (D\bar{D}^*)^\pm$	seen
$\Gamma_6 D^0 D^{*-} + \text{c.c.}$	seen
$\Gamma_7 D^- D^{*0} + \text{c.c.}$	seen
$\Gamma_8 \omega \pi^\pm$	not seen
$\Gamma_9 J/\psi \eta$	not seen
$\Gamma_{10} D^+ D^{*-} + \text{c.c.}$	seen
$\Gamma_{11} D^0 \bar{D}^{*0} + \text{c.c.}$	seen

Z_c(3900) BRANCHING RATIOS

$\Gamma(J/\psi \pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	356	ABLIKIM	15U	BES3	0	$e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$
seen	307	ABLIKIM	13T	BES3	\pm	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
seen	25	¹ XIAO	13A		0	$4.17 e^+ e^- \rightarrow \pi^0 \pi^0 J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	² ABAZOV	19	D0	1.96 TeV $p\bar{p} \rightarrow \pi^+ \pi^- J/\psi X$ (prompt)
not seen	³ ADOLPH	15D	COMP	\pm $\gamma N \rightarrow J/\psi \pi^\pm N$

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Upper limit for the prompt production is set: $N_{\text{prompt}}/N_{\text{nonprompt}} < 0.70$, CL = 95%.

³ ADOLPH 15D measure $B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm) \sigma(\gamma N \rightarrow Z_c(3900)^\pm N)/\sigma(\gamma N \rightarrow J/\psi N) < 3.7 \times 10^{-3}$ at 90% CL.

$\Gamma(h_c \pi^\pm)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	13X	BES3	\pm	$e^+ e^- \rightarrow h_c \pi^+ \pi^-$	

$\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	¹ VINOKUROVA 15	BELL	0	$B^+ \rightarrow K^+ \eta_c \pi^+ \pi^-$		

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ Z_c(3900)^0) \times B(X \rightarrow \eta_c \pi^+ \pi^-) < 4.7 \times 10^{-5}$ at 90% CL.

$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi \pi)$						Γ_5/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
6.2±1.1±2.7	¹ ABLIKIM	14A	BES3	\pm	$e^+ e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$	

¹ Assuming the same origin of the $(D\bar{D}^*)^\pm$ and $\pi^\pm J/\psi$ decay modes.

$\Gamma(D^0 D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_6/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
seen	ABLIKIM	15AC	BES3	\pm	$e^+ e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$	
seen	ABLIKIM	14A	BES3	\pm	$e^+ e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$	

$\Gamma(D^- D^{*0} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_7/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
seen	ABLIKIM	15AC	BES3	\pm	$e^+ e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$	
seen	ABLIKIM	14A	BES3	\pm	$e^+ e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$	

$\Gamma(\omega \pi^\pm)/\Gamma_{\text{total}}$						Γ_8/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	15R	BES3	\pm	$e^+ e^- \rightarrow \omega \pi^+ \pi^-$	

$\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$						Γ_9/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	15Q	BES3	0	$4.0-4.6 e^+ e^- \rightarrow J/\psi \eta \pi^0$	

DESIG=1
DESIG=2
DESIG=10
DESIG=11
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=8
DESIG=9
DESIG=4
DESIG=5
DESIG=6
DESIG=7

NODE=M210225

NODE=M210R01
NODE=M210R01

NODE=M210R01;LINKAGE=XI
NODE=M210R01;LINKAGE=B
NODE=M210R01;LINKAGE=A

NODE=M210R02
NODE=M210R02NODE=M210R11
NODE=M210R11

NODE=M210R11;LINKAGE=VI

NODE=M210R03
NODE=M210R03

NODE=M210R03;LINKAGE=A

NODE=M210R09
NODE=M210R09NODE=M210R10
NODE=M210R10NODE=M210R00
NODE=M210R00NODE=M210R04
NODE=M210R04

$\Gamma(J/\psi\eta)/\Gamma(J/\psi\pi)$						Γ_9/Γ_1	NODE=M210R05 NODE=M210R05
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT		
<0.15	90	ABLIKIM	15Q	BES3	0	4.226 $e^+ e^- \rightarrow J/\psi\eta\pi^0$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$							
<0.65	90	ABLIKIM	15Q	BES3	0	4.257 $e^+ e^- \rightarrow J/\psi\eta\pi^0$	OCCUR=2
$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(J/\psi\pi)$						Γ_4/Γ_1	NODE=M210R12 NODE=M210R12
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
2.3±0.8	332	1 ABLIKIM	19BC	BES3	±	$e^+ e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$	
¹ Using $e^+e^- \rightarrow \pi^\mp(Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm)$ cross section at 4.23 and 4.26 GeV from ABLIKIM 17J							
$\Gamma(D^+D^{*-} + c.c.)/\Gamma_{\text{total}}$						Γ_{10}/Γ	NODE=M210R06 NODE=M210R06
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
seen	ABLIKIM	15AB	BES3	0	$e^+ e^- \rightarrow \pi^0(D\bar{D}^*)^0$		
$\Gamma(D^0\bar{D}^{*0} + c.c.)/\Gamma_{\text{total}}$						Γ_{11}/Γ	NODE=M210R07 NODE=M210R07
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
seen	ABLIKIM	15AB	BES3	0	$e^+ e^- \rightarrow \pi^0(D\bar{D}^*)^0$		
$\Gamma(D^+D^{*-} + c.c.)/\Gamma(D^0\bar{D}^{*0} + c.c.)$						Γ_{10}/Γ_{11}	NODE=M210R08 NODE=M210R08
VALUE	DOCUMENT ID	TECN	CHG	COMMENT			
0.96±0.18±0.12	ABLIKIM	15AB	BES3	0	$e^+ e^- \rightarrow \pi^0(D\bar{D}^*)^0$		

Z_c(3900) REFERENCES

ABAZOV	19	PR D100 012005	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=59903
ABLIKIM	19BC	PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60036
ABAZOV	18B	PR D98 052010	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=59442
ABLIKIM	17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57950
ABLIKIM	15AB	PRL 115 222002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56954
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=56967
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56782
ABLIKIM	15R	PR D92 032009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56783
ABLIKIM	15U	PRL 115 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56786
ADOLPH	15D	PL B742 330	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56791
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (errat.)	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=57795
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=55648
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55409
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55635
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)	REFID=55593

X(3915)

$$I^G(J^{PC}) = 0^+(0 \text{ or } 2^{++})$$

was $\chi_{c0}(3915)$

The experimental analysis prefers $J^{PC} = 0^{++}$. However, a re-analysis presented in ZHOU 15C shows that if helicity-2 dominance assumption is abandoned and a sizable helicity-0 component is allowed, a $J^{PC} = 2^{++}$ assignment is possible.

X(3915) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3918.4 ± 1.9 OUR AVERAGE				
3919.4 ± 2.2 ± 1.6	59 ± 10	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
3919.1 ± 3.8 ± 2.0		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3915 ± 3 ± 2	49 ± 15	UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
3943 ± 11 ± 13	58 ± 11	¹ CHOI	05 BELL	$B \rightarrow \omega J/\psi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3926.4 ± 2.2 ± 1.2		² ABLIKIM	19V BES	$e^+ e^- \rightarrow \gamma \omega J/\psi$
3914.6 ± 3.8 ± 2.0		¹ AUBERT	08W BABR	Superseded by DEL-AMO-SANCHEZ 10B

¹ $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.² Could also be X(3940). Significance 3.1σ . Fit with additional resonance at 3963.7 ± 5.7 MeV, significance 3.4σ .

NODE=M159M

NODE=M159M

X(3915) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 ± 5 OUR AVERAGE				
13 ± 6 ± 3	59	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
31 ± 10 ± 5		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
17 ± 10 ± 3	49	UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
87 ± 22 ± 26	58	³ CHOI	05 BELL	$B \rightarrow \omega J/\psi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.8 ± 7.5 ± 2.6		⁴ ABLIKIM	19V BES	$e^+ e^- \rightarrow \gamma \omega J/\psi$
34 ± 12 ± 5		³ AUBERT	08W BABR	Superseded by DEL-AMO-SANCHEZ 10B

³ $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.⁴ Could also be X(3940). Significance 3.1σ . Fit with additional resonance at 3963.7 ± 5.7 MeV, significance 3.4σ .

NODE=M159M;LINKAGE=CH

NODE=M159M;LINKAGE=A

NODE=M159W

NODE=M159W

X(3915) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \frac{\omega J/\psi}{D^{*0} D^0}$	seen
$\Gamma_3 \pi^+ \pi^- \eta_c(1S)$	not seen
$\Gamma_4 \eta_c \eta$	not seen
$\Gamma_5 \eta_c \pi^0$	not seen
$\Gamma_6 K \bar{K}$	not seen
$\Gamma_7 \gamma \gamma$	seen
$\Gamma_8 \pi^0 \chi_{c1}$	

NODE=M159W;LINKAGE=CH

NODE=M159W;LINKAGE=A

NODE=M159215;NODE=M159

DESIG=1;OUR EST; \rightarrow UNCHECKED
 DESIG=3
 DESIG=4;OUR EVAL; \rightarrow UNCHECKED
 DESIG=6
 DESIG=7
 DESIG=5;OUR EVAL; \rightarrow UNCHECKED
 DESIG=2
 DESIG=8

NODE=M159220

NODE=M159G01
 NODE=M159G01

OCCUR=2

NODE=M159G01;LINKAGE=UH
 NODE=M159G01;LINKAGE=UR

$$\Gamma(\omega J/\psi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} = \Gamma_1 \Gamma_7 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
54 ± 9 OUR AVERAGE				
52 ± 10 ± 3	59 ± 10	⁵ LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
61 ± 17 ± 8	49 ± 15	⁵ UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
18 ± 5 ± 2	49 ± 15	⁶ UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$

⁵ For $J^P = 0^+$.⁶ For $J^P = 2^+$, helicity-2.

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_3\Gamma_7/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$
$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_6\Gamma_7/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.96	90	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

X(3915) BRANCHING RATIOS

$\Gamma(\omega J/\psi)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	7 DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$	
seen	8 CHOI	05 BELL	$B \rightarrow \omega J/\psi K$	
7 DEL-AMO-SANCHEZ 10B reports $B(B^\pm \rightarrow X(3915)K^\pm) \times B(X(3915) \rightarrow J/\psi\omega) = (3.0^{+0.7 \pm 0.5}_{-0.6 - 0.3}) \times 10^{-5}$ and $B(B^0 \rightarrow X(3915)K^0) \times B(X(3915) \rightarrow J/\psi\omega) = (2.1 \pm 0.9 \pm 0.3) \times 10^{-5}$.				NODE=M159R03;LINKAGE=DE
8 CHOI 05 reports $B(B \rightarrow X(3915)K) \times B(X(3915) \rightarrow J/\psi\omega) = (7.1 \pm 1.3 \pm 3.1) \times 10^{-5}$.				NODE=M159R03;LINKAGE=CH

$\Gamma(\omega J/\psi)/\Gamma(\bar{D}^{*0} D^0)$				Γ_1/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.71	90	9 AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
9 By combining the upper limit $B(B \rightarrow X(3915)K) \times B(X(3915) \rightarrow D^{*0}\bar{D}^0) < 0.67 \times 10^{-4}$ from AUSHEV 10 with the average of CHOI 05 and AUBERT 08W measurements $B(B \rightarrow X(3915)K) \times B(X(3915) \rightarrow \omega J/\psi) = (0.51 \pm 0.11) \times 10^{-4}$.				NODE=M159R02;LINKAGE=AU

$\Gamma(\eta_c\eta)/\Gamma_{\text{total}}$				Γ_4/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	10 VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \eta$	
10 VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ X(3915)^0) \times B(X \rightarrow \eta_c \eta) < 3.3 \times 10^{-5}$ at 90% CL.				NODE=M159R00;LINKAGE=VI

$\Gamma(\eta_c\pi^0)/\Gamma_{\text{total}}$				Γ_5/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	11 VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \pi^0$	
11 VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ X(3915)^0) \times B(X \rightarrow \eta_c \pi^0) < 1.8 \times 10^{-5}$ at 90% CL.				NODE=M159R04;LINKAGE=VI

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				Γ_7/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	59 ± 10	LEES	12AD BABR	$e^+e^- \rightarrow e^+e^-\omega J/\psi$
seen		UEHARA	10 BELL	$10.6 e^+e^- \rightarrow e^+e^-\omega J/\psi$

$\Gamma(\pi^0\chi_{c1})/\Gamma_{\text{total}}$				Γ_8/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	42 ± 14	12 BHARDWAJ	19 BELL	$B^\pm \rightarrow \chi_{c1}\pi^0 K^\pm$
12 BHARDWAJ 19 reports $B(B^+ \rightarrow K^+ X(3915)) \times B(X(3915) \rightarrow \chi_{c1}\pi^0) < 3.8 \times 10^{-5}$ at 90% CL. A signal significance 2.3 standard deviations.				NODE=M159R05;LINKAGE=A

X(3915) REFERENCES

ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
ZHOU	15C	PRL 115 022001	Z.-Y. Zhou, Z. Xiao, H.-Q. Zhou	(BEIJT, NANJ)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA..	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
UEHARA	10	PRL 104 092001	S. Uehara <i>et al.</i>	(BELLE Collab.)
AUBERT	08W	PRL 101 082001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHOI	05	PRL 94 182002	S.-K. Choi <i>et al.</i>	(BELLE Collab.)

NODE=M159G02
NODE=M159G02NODE=M159G03
NODE=M159G03

NODE=M159225

NODE=M159R03
NODE=M159R03

NODE=M159R03;LINKAGE=DE

NODE=M159R03;LINKAGE=CH

NODE=M159R02
NODE=M159R02

NODE=M159R02;LINKAGE=AU

NODE=M159R00
NODE=M159R00

NODE=M159R00;LINKAGE=VI

NODE=M159R04
NODE=M159R04

NODE=M159R04;LINKAGE=VI

NODE=M159R01
NODE=M159R01NODE=M159R05
NODE=M159R05

NODE=M159R05;LINKAGE=A

NODE=M159

REFID=59796

REFID=59884

REFID=56706

REFID=57795

REFID=56842

REFID=55592

REFID=54751

REFID=54752

REFID=53225

REFID=53362

REFID=53232

REFID=52263

REFID=50737

NODE=M050

 $\chi_{c2}(3930)$ $I^G(J^{PC}) = 0^+(2^{++})$ **$\chi_{c2}(3930)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3922.2±1.0 OUR AVERAGE		Error includes scale factor of 1.6. [3927.2 ± 2.6 MeV OUR 2019 AVERAGE]		
3921.9±0.6±0.2	1	AAIJ	19M LHCb	$p p \rightarrow D\bar{D} +$ anything
3926.7±2.7±1.1	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
3929 ± 5 ± 2	64	UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

¹ Measured in prompt hadroproduction.

NODE=M050M

NODE=M050M

NEW

NODE=M050M;LINKAGE=A


NODE=M050W

NODE=M050W

NEW

NODE=M050W;LINKAGE=A

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.3± 2.8 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below. [24 ± 6 MeV OUR 2019 AVERAGE]		
36.6± 1.9±0.9	2	AAIJ	19M LHCb	$p p \rightarrow D\bar{D} +$ anything
21.3± 6.8±3.6	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
29 ± 10 ± 2	64	UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

² Measured in prompt hadroproduction.WEIGHTED AVERAGE
35.3±2.8 (Error scaled by 1.4) **$\chi_{c2}(3930)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \gamma\gamma$	seen
$\Gamma_2 K\bar{K}\pi$	
$\Gamma_3 K^+ K^- \pi^+ \pi^- \pi^0$	
$\Gamma_4 D\bar{D}$	seen
$\Gamma_5 D^+ D^-$	seen
$\Gamma_6 D^0 \bar{D}^0$	seen
$\Gamma_7 \pi^+ \pi^- \eta_c(1S)$	not seen
$\Gamma_8 K\bar{K}$	not seen

NODE=M050215;NODE=M050

```

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=5
DESIG=6
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=7;OUR EVAL;→ UNCHECKED ←
DESIG=8;OUR EVAL;→ UNCHECKED ←

```

$\chi_{c2}(3930)$ PARTIAL WIDTHS **$\chi_{c2}(3930) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$**

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_3\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<3.4	90	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$\Gamma(D\bar{D}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_4\Gamma_1/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.21±0.04 OUR AVERAGE				
0.24±0.05±0.04	76 ± 17	AUBERT	10G	BABR $e^+ e^- \rightarrow e^+ e^- D\bar{D}$
0.18±0.05±0.03	64	³ UEHARA	06	BELL $e^+ e^- \rightarrow e^+ e^- D\bar{D}$
3 Assuming $B(D^+ D^-) = 0.89 B(D^0 \bar{D}^0)$.				
$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_7\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<18	90	LEES	12AE	BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$
$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_8\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.256	90	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$

 $\chi_{c2}(3930)$ BRANCHING RATIOS

$\Gamma(D^+ D^-)/\Gamma(D^0 \bar{D}^0)$				Γ_5/Γ_6
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.74±0.43±0.16	64	UEHARA	06	BELL $10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

 $\chi_{c2}(3930)$ REFERENCES

AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
DEL-AMO-SA... 11M	PR D84 012004		P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
AUBERT	10G	PR D81 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	06	PRL 96 082003	S. Uehara <i>et al.</i>	(BELLE Collab.)

NODE=M050220

NODE=M050222

NODE=M050G01

NODE=M050G01

NODE=M050G02

NODE=M050G02

NODE=M050G1

NODE=M050G1

NODE=M050G1;LINKAGE=UE

NODE=M050G03

NODE=M050G03

NODE=M050G04

NODE=M050G04

NODE=M050225

NODE=M050R01

NODE=M050R01

NODE=M050

REFID=59697

REFID=55592

REFID=54752

REFID=16751

REFID=53357

REFID=51039

X(3940) $I^G(J^{PC}) = ??(??)$

OMITTED FROM SUMMARY TABLE

Reported by ABE 07, observed in $e^+ e^- \rightarrow J/\psi X$.**X(3940) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3942\pm7\pm6	52	PAKHLOV	08	BELL $e^+ e^- \rightarrow J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3943 \pm 6 \pm 6	25	¹ ABE	07	BELL $e^+ e^- \rightarrow J/\psi X$
3936 \pm 14	266	² ABE	07	BELL $e^+ e^- \rightarrow J/\psi(c\bar{c})$

¹From a fit to $D^{*+} D^-$ and $D^{*0} \bar{D}^0$ events.
²From the inclusive fit. Not independent of the exclusive measurement by ABE 07.

X(3940) WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
37\pm26\pm8		52	PAKHLOV	08	BELL $e^+ e^- \rightarrow J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<52	90	25	ABE	07	BELL $e^+ e^- \rightarrow J/\psi X$

X(3940) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D\bar{D}^* + c.c.$	seen
$\Gamma_2 D\bar{D}$	not seen
$\Gamma_3 J/\psi\omega$	not seen

X(3940) BRANCHING RATIOS

$\Gamma(D\bar{D}^* + c.c.)/\Gamma_{\text{total}}$	Γ_1/Γ				
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.45	90	25	^{1,2} ABE	07	BELL $e^+ e^- \rightarrow J/\psi X$

¹For $X(3940)$ decaying to final states with more than two tracks.
²PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c² may consist of several states.

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.41	90	^{1,2} ABE	07	BELL $e^+ e^- \rightarrow J/\psi X$

¹For $X(3940)$ decaying to final states with more than two tracks.
²PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c² may consist of several states.

$\Gamma(J/\psi\omega)/\Gamma_{\text{total}}$	Γ_3/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.26	90	^{1,2} ABE	07	BELL $e^+ e^- \rightarrow J/\psi X$

¹For $X(3940)$ decaying to final states with more than two tracks.
²PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c² may consist of several states.

X(3940) REFERENCES

PAKHLOV ABE	08 07	PRL 100 202001 PRL 98 082001	P. Pakhlov <i>et al.</i> K. Abe <i>et al.</i>	(BELLE Collab.) (BELLE Collab.)
----------------	----------	---------------------------------	--	------------------------------------

NODE=M029

NODE=M029

NODE=M029M

NODE=M029M

OCCUR=2

NODE=M029M;LINKAGE=EB

NODE=M029M;LINKAGE=EM

NODE=M029W

NODE=M029W

NODE=M029215;NODE=M029

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M029225

NODE=M029R01
NODE=M029R01NODE=M029R01;LINKAGE=AB
NODE=M029R01;LINKAGE=AENODE=M029R02
NODE=M029R02NODE=M029R02;LINKAGE=AB
NODE=M029R02;LINKAGE=AENODE=M029R03
NODE=M029R03NODE=M029R03;LINKAGE=AB
NODE=M029R03;LINKAGE=AE

NODE=M029

REFID=52302
REFID=51627

NODE=M213

X(4020)

$$I^G(J^P C) = 1^+(?^-)$$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Charged X(4020) seen by ABLIKIM 13X from $e^+ e^- \rightarrow \pi^+ \pi^- h_c(1P)$ at c.m. energy from 3.90 to 4.42 GeV as a peak in the invariant mass distribution of the $\pi^\pm h_c(1P)$ system, and by ABLIKIM 14B from $e^+ e^- \rightarrow (D^* \bar{D}^*)^\pm \pi^\mp$ events in $(D^* \bar{D}^*)^\pm$ mass. A neutral X(4020) seen by ABLIKIM 14P at three c.m. energies in the same range in $e^+ e^- \rightarrow \pi^0 \pi^0 h_c(1P)$ as a peak in the larger of the two masses recoiling against a π^0 . ABLIKIM 15AA observes a 5.9σ signal in $(D^* \bar{D}^*)^0$ in $e^+ e^- \rightarrow (D^* \bar{D}^*)^0 \pi^0$ events using collisions at two c.m. energies. Production rates and mass values support grouping neutral and charged X(4020) together as manifestations of a single $I = 1$ particle.

NODE=M213

X(4020) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
4024.1±1.9 OUR AVERAGE					
4025.5 ^{+2.0} _{-4.7} ±3.1	116	1 ABLIKIM	15AA BES3	0	$e^+ e^- \rightarrow (D^* \bar{D}^*)^0 \pi^0$
4026.3±2.6±3.7	401	1 ABLIKIM	14B BES3	±	$e^+ e^- \rightarrow (D^* \bar{D}^*)^\pm \pi^\mp$
4023.9±2.2±3.8	61	1,2 ABLIKIM	14P BES3	0	$e^+ e^- \rightarrow \pi^0 \pi^0 h_c$
4022.9±0.8±2.7	253	1 ABLIKIM	13X BES3	±	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$

1 Neglecting interference between the X(4020) and non-resonant continuum.

2 Assuming $J^P = 1^+$ and width of 7.9 ± 2.6 MeV.

NODE=M213M

NODE=M213M

X(4020) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
13 ±5 OUR AVERAGE					
23.0±6.0±1.0	116	1 ABLIKIM	15AA BES3	0	$e^+ e^- \rightarrow (D^* \bar{D}^*)^0 \pi^0$
24.8±5.6±7.7	401	1 ABLIKIM	14B BES3	±	$e^+ e^- \rightarrow (D^* \bar{D}^*)^\pm \pi^\mp$
7.9±2.7±2.6	253	1 ABLIKIM	13X BES3	±	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$

1 Neglecting interference between the X(4020) and non-resonant continuum.


NODE=M213M;LINKAGE=AB

NODE=M213M;LINKAGE=B

NODE=M213W

NODE=M213W

WEIGHTED AVERAGE
13±5 (Error scaled by 1.7)



X(4020) DECAY MODES

NODE=M213215;NODE=M213

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 h_c(1P)\pi$	seen
$\Gamma_2 D^*\bar{D}^*$	seen
$\Gamma_3 D\bar{D}^* + \text{c.c.}$	not seen
$\Gamma_4 \eta_c\pi^+\pi^-$	not seen
$\Gamma_5 \eta_c(1S)\rho(770)^\pm$	
$\Gamma_6 J/\psi(1S)\pi^\pm$	not seen

X(4020) BRANCHING RATIOS

$\Gamma(h_c(1P)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	EVTS DOCUMENT ID TECN CHG COMMENT
seen	61 ABLIKIM 14P BES3 0 $e^+e^- \rightarrow \pi^0\pi^0 h_c$
seen	253 ABLIKIM 13X BES3 \pm $e^+e^- \rightarrow \pi^+\pi^- h_c$

$\Gamma(D^*\bar{D}^*)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE	EVTS DOCUMENT ID TECN CHG COMMENT
seen	116 1 ABLIKIM 15AA BES3 0 $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
seen	401 1 ABLIKIM 14B BES3 \pm $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$

1 Neglecting interference between the $X(4020)$ and non-resonant continuum.

$\Gamma(D\bar{D}^* + \text{c.c.})/\Gamma_{\text{total}}$	Γ_3/Γ
VALUE	DOCUMENT ID TECN CHG COMMENT
not seen	ABLIKIM 15AC BES3 \pm $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_4/Γ
VALUE	DOCUMENT ID TECN COMMENT
not seen	1 VINOKUROVA 15 BELL $B^+ \rightarrow K^+\eta_c\pi^+\pi^-$

1 VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ X(4020)^0) \times B(X \rightarrow \eta_c\pi^+\pi^-) < 1.6 \times 10^{-5}$ at 90% CL.

$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(h_c(1P)\pi)$	Γ_5/Γ_1
VALUE	CL% DOCUMENT ID TECN CHG COMMENT
<1.2	90 1 ABLIKIM 19BC BES3 \pm $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$

1 Using $e^+e^- \rightarrow \pi^\mp(Z_c(4020)^\pm \rightarrow h_c(1P)\pi^\pm)$ cross section at 4.23, 4.26 and 4.36 GeV from ABLIKIM 13X

$\Gamma(J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$	Γ_6/Γ
VALUE	DOCUMENT ID TECN COMMENT
not seen	1 ABLIKIM 17J BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$

1 From Partial Wave Analysis assuming $J^P = 1^+$.**X(4020) REFERENCES**

ABLIKIM	19BC PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17J PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AA PRL 115 182002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AC PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
VINOKUROVA	15 JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also	JHEP 1702 088 (errat.)	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	14B PRL 112 132001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	14P PRL 113 212002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13X PR 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)

REFID=60036
 REFID=57950
 REFID=56951
 REFID=56967
 REFID=56706
 REFID=57795
 REFID=55654
 REFID=56118
 REFID=55635

NODE=M213225

NODE=M213R01
NODE=M213R01NODE=M213R02
NODE=M213R02

NODE=M213R02;LINKAGE=A

NODE=M213R03
NODE=M213R03NODE=M213R00
NODE=M213R00

NODE=M213R00;LINKAGE=VI

NODE=M213R05
NODE=M213R05

NODE=M213R05;LINKAGE=A

NODE=M213R04
NODE=M213R04

NODE=M213R04;LINKAGE=A

NODE=M213

NODE=M072

 $\psi(4040)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\psi(4040)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4039 ± 1 OUR ESTIMATE			
4039.6± 4.3	1 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4034 ± 6	2 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4037 ± 2	3 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4040 ± 1	4 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4040 ± 10	BRANDELIK	78C DASP	$e^+ e^- \rightarrow$ hadrons
1 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
2 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
3 From a fit to Crystal Ball (OSTERHELD 86) data.			
4 From a fit to BES (BAI 02C) data.			

 $\psi(4040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 ±10 OUR ESTIMATE			
84.5±12.3	5 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87 ±11	6 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
85 ±10	7 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
89 ± 6	8 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
52 ±10	BRANDELIK	78C DASP	$e^+ e^- \rightarrow$ hadrons
5 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
6 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
7 From a fit to Crystal Ball (OSTERHELD 86) data.			
8 From a fit to BES (BAI 02C) data.			

 $\psi(4040)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+ e^-$	$(1.07 \pm 0.16) \times 10^{-5}$	
$\Gamma_2 D\bar{D}$	seen	
$\Gamma_3 D^0\bar{D}^0$	seen	
$\Gamma_4 D^+D^-$	seen	
$\Gamma_5 D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_6 D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_7 D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_8 D^*\bar{D}^*$	seen	
$\Gamma_9 D^*(2007)^0\bar{D}^*(2007)^0$	seen	
$\Gamma_{10} D^*(2010)^+D^*(2010)^-$	seen	
$\Gamma_{11} D\bar{D}\pi (\text{excl. } D^*\bar{D})$		
$\Gamma_{12} D^0D^-\pi^+ + \text{c.c.} (\text{excl. } D^*(2007)^0\bar{D}^0 + \text{c.c.}, D^*(2010)^+D^- + \text{c.c.})$	not seen	
$\Gamma_{13} D\bar{D}^*\pi (\text{excl. } D^*\bar{D}^*)$	not seen	

NODE=M072M

NODE=M072M

→ UNCHECKED ←

OCCUR=2

NODE=M072M;LINKAGE=AB

NODE=M072M;LINKAGE=MO

NODE=M072M;LINKAGE=ST

NODE=M072M;LINKAGE=SE

NODE=M072W

NODE=M072W

→ UNCHECKED ←

OCCUR=2

NODE=M072W;LINKAGE=AB

NODE=M072W;LINKAGE=MO

NODE=M072W;LINKAGE=ST

NODE=M072W;LINKAGE=SE

NODE=M072215;NODE=M072

NODE=M072

DESIG=5

DESIG=17;OUR EST;→ UNCHECKED ←

DESIG=1

DESIG=18

DESIG=19;OUR EST;→ UNCHECKED ←

DESIG=2

DESIG=20

DESIG=21;OUR EST;→ UNCHECKED ←

DESIG=3

DESIG=22

DESIG=23

DESIG=24

DESIG=25

Γ_{14}	$D^0 \bar{D}^{*-} \pi^+ + c.c. \text{ (excl.)}$ $D^*(2010)^+ D^*(2010)^-$	seen		DESIG=26
Γ_{15}	$D_s^+ D_s^-$	seen		DESIG=27
Γ_{16}	$J/\psi(1S) \text{ hadrons}$			DESIG=4
Γ_{17}	$J/\psi \pi^+ \pi^-$	< 4	$\times 10^{-3}$	90%
Γ_{18}	$J/\psi \pi^0 \pi^0$	< 2	$\times 10^{-3}$	90%
Γ_{19}	$J/\psi \eta$	(5.2 ± 0.7)	$\times 10^{-3}$	
Γ_{20}	$J/\psi \pi^0$	< 2.8	$\times 10^{-4}$	90%
Γ_{21}	$J/\psi \pi^+ \pi^- \pi^0$	< 2	$\times 10^{-3}$	90%
Γ_{22}	$\chi_{c1} \gamma$	< 3.4	$\times 10^{-3}$	90%
Γ_{23}	$\chi_{c2} \gamma$	< 5	$\times 10^{-3}$	90%
Γ_{24}	$\chi_{c1} \pi^+ \pi^- \pi^0$	< 1.1	%	90%
Γ_{25}	$\chi_{c2} \pi^+ \pi^- \pi^0$	< 3.2	%	90%
Γ_{26}	$h_c(1P) \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%
Γ_{27}	$\phi \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%
Γ_{28}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.9	$\times 10^{-4}$	90%
Γ_{29}	$\Lambda \bar{\Lambda} \pi^0$	< 9	$\times 10^{-5}$	90%
Γ_{30}	$\Lambda \bar{\Lambda} \eta$	< 3.0	$\times 10^{-4}$	90%
Γ_{31}	$\Sigma^+ \bar{\Sigma}^-$	< 1.3	$\times 10^{-4}$	90%
Γ_{32}	$\Sigma^0 \bar{\Sigma}^0$	< 7	$\times 10^{-5}$	90%
Γ_{33}	$\Xi^+ \bar{\Xi}^-$	< 1.6	$\times 10^{-4}$	90%
Γ_{34}	$\Xi^0 \bar{\Xi}^0$	< 1.8	$\times 10^{-4}$	90%
Γ_{35}	$\mu^+ \mu^-$			DESIG=6

 $\psi(4040)$ PARTIAL WIDTHS **$\Gamma(e^+ e^-)$** VALUE (keV)
 0.86 ± 0.07 OUR ESTIMATE **0.83 ± 0.20**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 to 1.4	10 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
0.88 ± 0.11	11 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.91 ± 0.13	12 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.75 ± 0.15	BRANDELIK	78C DASP	$e^+ e^-$

⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.

¹⁰ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.

¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.

¹² From a fit to BES (BAI 02C) data.

 Γ_1

NODE=M072220

NODE=M072W5

NODE=M072W5

→ UNCHECKED ←

OCCUR=2

NODE=M072W5;LINKAGE=AB

NODE=M072W5;LINKAGE=MO

NODE=M072W5;LINKAGE=ST

NODE=M072W5;LINKAGE=SE

NODE=M072235

NODE=M072G01

NODE=M072G01

NODE=M072G01;LINKAGE=A

NODE=M072G02

NODE=M072G02

NODE=M072G02;LINKAGE=A

 $\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$ **$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{22} \Gamma_1 / \Gamma$
<2.9	90	13 HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c1} \gamma$	

¹³ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{23} \Gamma_1 / \Gamma$
<4.6	90	14 HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c2} \gamma$	

¹⁴ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma^2(\text{total})$

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma \times \Gamma_1/\Gamma$
<u>VALUE</u> (units 10^{-8})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.1±1.4±1.5	15 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$	
12.8±2.1±1.9	16 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$	
15 Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.				
16 Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.				

$\psi(4040)$ BRANCHING RATIOS

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$				Γ_1/Γ
<u>VALUE</u> (units 10^{-5})	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~1.0	FELDMAN	77	MRK1	$e^+ e^-$

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$				Γ_3/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^0\bar{D}^0$	
seen	PAKHLOVA	08 BELL	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$	

$\Gamma(D^+D^-)/\Gamma_{\text{total}}$				Γ_4/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^+D^-\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^+D^-$	
seen	PAKHLOVA	08 BELL	$e^+ e^- \rightarrow D^+D^-\gamma$	

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D} + \text{c.c.})$				Γ_2/Γ_5
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.24±0.05±0.12	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}$	

$\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$				Γ_3/Γ_6
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.05±0.03	17 GOLDHABER	77 MRK1	$e^+ e^-$	

17 Phase-space factor (p^3) explicitly removed.

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_6/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^0\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0}\bar{D}^0$	

$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_7/Γ
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
seen	18 ZHUKOVA	18 BELL	$e^+ e^- \rightarrow D^{*+}D^-\gamma$	
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*+}D^-\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+}D^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	PAKHLOVA	07 BELL	$e^+ e^- \rightarrow D^{*+}D^-\gamma$	
18 Supersedes PAKHLOVA 07.				

$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$				Γ_7/Γ_6
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.95±0.09±0.10	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{*0}\bar{D}$	

$\Gamma(D^*\bar{D}^*)/\Gamma(D^*\bar{D} + \text{c.c.})$				Γ_8/Γ_5
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.18±0.14±0.03	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	

NODE=M072230

NODE=M072R25

NODE=M072R25

OCCUR=2

NODE=M072R25;LINKAGE=A

NODE=M072R25;LINKAGE=B

NODE=M072225

NODE=M072R4

NODE=M072R4

NODE=M072R14

NODE=M072R14

NODE=M072R15

NODE=M072R15

NODE=M072R12

NODE=M072R12

NODE=M072R1

NODE=M072R1

NODE=M072R;LINKAGE=P

NODE=M072R16

NODE=M072R16

NODE=M072R17

NODE=M072R17

OCCUR=3

NODE=M072R17;LINKAGE=C

NODE=M072R11

NODE=M072R11

NODE=M072R13

NODE=M072R13

$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$				Γ_9/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$		NODE=M072R18
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$		NODE=M072R18
$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})$				Γ_9/Γ_6	
VALUE	DOCUMENT ID	TECN	COMMENT		
32.0 ± 12.0	19 GOLDHABER 77	MRK1	$e^+ e^-$		NODE=M072R2
19 Phase-space factor (p^3) explicitly removed.					NODE=M072R2;LINKAGE=P
$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$				Γ_{10}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	20 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$		NODE=M072R19
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$		NODE=M072R19
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$		OCCUR=2
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$		NODE=M072R19;LINKAGE=B
20 Supersedes PAKHLOVA 07.					
$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.})) / \Gamma_{\text{total}}$				Γ_{12}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$		NODE=M072R20
$\Gamma(D \bar{D}^* \pi (\text{excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$				Γ_{13}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$		NODE=M072R21
$\Gamma(D^0 \bar{D}^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)) / \Gamma_{\text{total}}$				Γ_{14}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 \bar{D}^{*-} \pi^+ \gamma$		NODE=M072R22
$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$				Γ_{15}/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$		NODE=M072R23
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$		NODE=M072R23
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$		
$\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{17}/Γ	
VALUE (units 10^{-3}) CL%	DOCUMENT ID	TECN	COMMENT		
<4 90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$		NODE=M072R01
$\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$				Γ_{18}/Γ	
VALUE (units 10^{-3}) CL%	DOCUMENT ID	TECN	COMMENT		
<2 90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$		NODE=M072R02
$\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$				Γ_{19}/Γ	
VALUE (units 10^{-3}) CL%	DOCUMENT ID	TECN	COMMENT		
$5.2 \pm 0.5 \pm 0.5$ 21	ABLIKIM 12K	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$		NODE=M072R03
• • • We do not use the following data for averages, fits, limits, etc. • • •					NODE=M072R03
<7 90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$		
21 ABLIKIM 12K measure $\sigma(e^+ e^- \rightarrow J/\psi \eta) = 32.1 \pm 2.8 \pm 1.3 \text{ pb}$. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.					NODE=M072R03;LINKAGE=AB
$\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$				Γ_{20}/Γ	
VALUE (units 10^{-3}) CL%	DOCUMENT ID	TECN	COMMENT		
<0.28 90 22	ABLIKIM 12K	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$		NODE=M072R04
• • • We do not use the following data for averages, fits, limits, etc. • • •					NODE=M072R04
<2 90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$		
22 ABLIKIM 12K measure $\sigma(e^+ e^- \rightarrow J/\psi \pi^0) < 1.6 \text{ pb}$. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.					NODE=M072R04;LINKAGE=AB

$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 Γ_{21}/Γ NODE=M072R05
NODE=M072R05 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 Γ_{22}/Γ NODE=M072R06
NODE=M072R06 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
<17	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 Γ_{23}/Γ NODE=M072R07
NODE=M072R07 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 Γ_{24}/Γ NODE=M072R08
NODE=M072R08 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<32	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 Γ_{25}/Γ NODE=M072R09
NODE=M072R09 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	23 PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

 Γ_{26}/Γ NODE=M072R24
NODE=M072R24

23 From several values of \sqrt{s} near the peak of the $\psi(4040)$, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 1.0 \pm 8.0 \pm 5.4 \pm 0.2$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

 Γ_{27}/Γ NODE=M072R10
NODE=M072R10 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.9	90	24 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

 Γ_{28}/Γ NODE=M072R26
NODE=M072R26

24 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.9	90	25 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

 Γ_{29}/Γ NODE=M072R27
NODE=M072R27

25 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.0	90	26 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

 Γ_{30}/Γ NODE=M072R28
NODE=M072R28

26 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.3	90	27 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

 Γ_{31}/Γ NODE=M072R29
NODE=M072R29

27 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<0.7	90	28 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

 Γ_{32}/Γ NODE=M072R30
NODE=M072R30

28 Assuming that interference effects between resonance and continuum can be neglected.

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.6	90	29 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

 Γ_{33}/Γ NODE=M072R31
NODE=M072R31

29 Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R31;LINKAGE=A

$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{34}/Γ
<1.8	90	30 ABLIKIM	13Q	BES3 $e^+ e^- \rightarrow \psi(4040)$	

30 Assuming that interference effects between resonance and continuum can be neglected.

 $\psi(4040)$ REFERENCES

ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
ABLIKIM	13Q	PR D87 112011	Ablikim M. <i>et al.</i>	(BESIII Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	12K	PR D86 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA... 10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN... 09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PR 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
Also	ZPHY C1 233		R. Brandelik <i>et al.</i>	(DASP Collab.)
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)

 $X(4050)^{\pm}$

$$I^G(J^{PC}) = 1^-(?^+)$$

I, G, C need confirmation.

OMMITTED FROM SUMMARY TABLE

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Observed by MIZUK 08 in the $\pi^+ \chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

 $X(4050)^{\pm}$ MASS

<u>VALUE</u> (MeV)	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{34}/Γ
4051 ± 14⁺²⁰₋₄₁	¹ MIZUK	08 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$	

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

 $X(4050)^{\pm}$ WIDTH

<u>VALUE</u> (MeV)	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{34}/Γ
82⁺²¹⁺⁴⁷₋₁₇₋₂₂	¹ MIZUK	08 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$	

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

 $X(4050)^{\pm}$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^+ \chi_{c1}(1P)$	seen
$\Gamma_2 \quad \pi^\pm \psi(3770)$	not seen

NODE=M072R32
NODE=M072R32

NODE=M072R32;LINKAGE=A

NODE=M072

REFID=58710
REFID=56816
REFID=55393
REFID=55377
REFID=54738
REFID=53638
REFID=16787
REFID=53532
REFID=53540
REFID=52724
REFID=53114
REFID=53143
REFID=52142
REFID=52132
REFID=52134
REFID=51628
REFID=51075
REFID=50813
REFID=50506
REFID=50503
REFID=51064
REFID=22232
REFID=22114
REFID=22062
REFID=11434

NODE=M191

NODE=M191

NODE=M191M

NODE=M191M

NODE=M191M;LINKAGE=MI

NODE=M191W

NODE=M191W

NODE=M191W;LINKAGE=MI

NODE=M191215;NODE=M191

DESIG=1
DESIG=2

X(4050) \pm BRANCHING RATIOS

$\Gamma(\pi^+ \chi_{c1}(1P))/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	1 MIZUK	08 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			

not seen 2 LEES 12B BABR $B \rightarrow K \pi \chi_{c1}(1P)$

1 With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- X(4050)^+) \times B(X(4050)^+ \rightarrow \pi^+ \chi_{c1}(1P)) = (3.0^{+1.5+3.7}_{-0.8-1.6}) \times 10^{-5}$.

2 With a product branching fraction limit of $B(\bar{B}^0 \rightarrow X(4050)^+ K^-) \times B(X(4050)^+ \rightarrow \chi_{c1} \pi^+) < 1.8 \times 10^{-5}$ at 90% CL.

$\Gamma(\pi^\pm \psi(3770))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	1 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$
1 From a measurement of $\sigma(e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.			

X(4050) \pm REFERENCES

ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12B PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08 PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

X(4055) \pm

$$I^G(J^{PC}) = 1^+ (?^-)$$

I, G, C need confirmation.

OMMITTED FROM SUMMARY TABLE

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Needs confirmation. Seen by WANG 15A in the $\psi(2S)\pi^+$ invariant mass distribution in $\psi(4360) \rightarrow \psi(2S)\pi^+\pi^-$ decay.

X(4055) \pm MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4054 ± 3 ± 1	1 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4039.3 ± 6.0	2 ABLIKIM	18K BES3	$e^+ e^- \rightarrow \pi^0 \pi^0 \psi(2S)$
4032.1 ± 2.4	3 ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$

1 Statistical significance of 3.5 σ .

2 Statistical error only, with significance of 5.9 σ (from a fit with a 19% CL). Identified as the same structure observed in ABLIKIM 17V in $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ decays.

3 Statistical error only, with significance of 9.2 σ . From an unbinned maximum likelihood fit of the $\pi^+ \pi^- \psi(2S)$ Dalitz plot from data collected at $\sqrt{s} = 4.416$ GeV for a $J^C = 1^+$ state. The fit does not match the detailed structure of the data, having a C.L. of only 8%.

X(4055) \pm WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
45 ± 11 ± 6	1 WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

31.9 ± 14.8	2 ABLIKIM	18K BES3	$e^+ e^- \rightarrow \pi^0 \pi^0 \psi(2S)$
26.1 ± 5.3	3 ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$

1 Statistical significance of 3.5 σ .

2 Statistical error only, with significance of 5.9 σ (from a fit with a 19% CL). Identified as the same structure observed in ABLIKIM 17V in $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ decays.

3 Statistical error only, with significance of 9.2 σ . From an unbinned maximum likelihood fit of the $\pi^+ \pi^- \psi(2S)$ Dalitz plot from data collected at $\sqrt{s} = 4.416$ GeV for a $J^C = 1^+$ state. The fit does not match the detailed structure of the data, having a C.L. of only 8%.

NODE=M191225

NODE=M191R01

NODE=M191R01

NODE=M191R01;LINKAGE=MI

NODE=M191R01;LINKAGE=LE

NODE=M191R00
NODE=M191R00

NODE=M191R00;LINKAGE=A

NODE=M191

REFID=59910
REFID=54042
REFID=52535

NODE=M223

NODE=M223

NODE=M223M

NODE=M223M

OCCUR=2

NODE=M223M;LINKAGE=A

NODE=M223M;LINKAGE=C

NODE=M223M;LINKAGE=B

NODE=M223W

NODE=M223W

NODE=M223W;LINKAGE=A

NODE=M223W;LINKAGE=C

NODE=M223W;LINKAGE=B

X(4055) \pm DECAY MODES

NODE=M223215;NODE=M223

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+ \psi(2S)$	seen
$\Gamma_2 \pi^\pm \psi(3770)$	not seen

X(4055) \pm BRANCHING RATIOS

$\Gamma(\pi^+ \psi(2S))/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE seen	DOCUMENT ID ¹ WANG TECHN 15A BELL COMMENT $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

¹ Statistical significance of 3.5 σ .

$\Gamma(\pi^\pm \psi(3770))/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE not seen	DOCUMENT ID ¹ ABLIKIM TECHN 19AR BES3 COMMENT $e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$

¹ From a measurement of $\sigma(e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.**X(4055) \pm REFERENCES**

ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18K PR D97 052001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17V PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also	PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
WANG	15A PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)

X(4100) \pm

$I^G(J^{PC}) = 1^-(???)$

OMITTED FROM SUMMARY TABLEProperties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.Reported by AAIJ 18AN in the $\eta_c(1S)\pi^-$ invariant mass distribution in $B^0 \rightarrow \eta_c(1S)K^+\pi^-$ decays with a significance of 3.4σ . $J^P = 0^+$ or 1^- assignment consistent with data.**X(4100) \pm MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4096\pm20⁺¹⁸₋₂₂	AAIJ	18AN LHCb	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

X(4100) \pm WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
152\pm58⁺⁶⁰₋₃₅	AAIJ	18AN LHCb	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

X(4100) \pm DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \eta_c(1S)\pi^-$	seen
$\Gamma_2 \pi^\pm \psi(3770)$	not seen

X(4100) \pm BRANCHING RATIOS

$\Gamma(\eta_c(1S)\pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE seen	DOCUMENT ID ¹ AAIJ TECHN 18AN LHCb COMMENT $B^0 \rightarrow \eta_c(1S)K^+\pi^-$

¹ AAIJ 18AN quotes a fit fraction for $B^0 \rightarrow X(4100)^- K^+ \rightarrow \eta_c(1S)\pi^- K^+$ of $(3.3 \pm 1.1^{+1.2}_{-1.1})\%$ from an amplitude analysis.

NODE=M240215;NODE=M240

DESIG=1

DESIG=2

NODE=M240225

NODE=M240R01

NODE=M240R01

NODE=M240R01;LINKAGE=A

NODE=M240R00

NODE=M240R00

NODE=M240R00;LINKAGE=A

NODE=M223

REFID=59910

REFID=58896

REFID=58029

REFID=59611

REFID=56839

NODE=M240

NODE=M240

NODE=M240

NODE=M240M

NODE=M240M

NODE=M240W

NODE=M240W

NODE=M240215;NODE=M240

DESIG=1

DESIG=2

NODE=M240225

NODE=M240R01

NODE=M240R01

NODE=M240R01;LINKAGE=A

$\Gamma(\pi^\pm \psi(3770))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
not seen	1 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$	
1 From a measurement of $\sigma(e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.				

 $X(4100)^\pm$ REFERENCES

ABLIKIM AAIJ	19AR PR D100 032005 18AN EPJ C78 1019	M. Ablikim <i>et al.</i> R. Aaij <i>et al.</i>	(BESIII Collab.) (LHCb Collab.)
-----------------	--	---	------------------------------------

 $\chi_{c1}(4140)$

$I^G(J^{PC}) = 0^+(1^{++})$

was $X(4140)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen by AALTONEN 09AH, ABAZOV 14A, CHATRCHYAN 14M,
AAIJ 17C in $B^+ \rightarrow \chi_{c1} K^+$, $\chi_{c1} \rightarrow J/\psi \phi$, and by ABAZOV 15M
separately in both prompt (4.7σ) and non-prompt (5.6σ) production
in $p\bar{p} \rightarrow J/\psi \phi +$ anything. Not seen by SHEN 10 in $\gamma\gamma \rightarrow J/\psi \phi$ and ABLIKIM 15 in $e^+ e^- \rightarrow \gamma J/\psi \phi$ at $\sqrt{s} = 4.23, 4.26,$
 4.36 GeV.

 $\chi_{c1}(4140)$ MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
4146.8 ± 2.4 OUR AVERAGE		Error includes scale factor of 1.1.			NODE=M193M
4146.5 ± 4.5 -2.8	4289	1 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193M
4143.4 ± 2.9 -3.0	19	2 AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193M
4152.5 ± 1.7 -5.4	616	3 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi \phi +$ anything	NODE=M193M
4159.0 ± 4.3 ± 6.6	52	4 ABAZOV	14A D0	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193M
4148.0 ± 2.4 ± 6.3	0.3k	5 CHATRCHYAN 14M	CMS	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193M
• • • We do not use the following data for averages, fits, limits, etc. • • •					
4143.0 ± 2.9 ± 1.2	14 6,7	AALTONEN 09AH	CDF	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193M
1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.4σ . 2 Statistical significance of more than 5σ . 3 Statistical significance of more than 6σ . 4 Statistical significance of 3.1σ . 5 From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ . 6 Statistical significance of 3.8σ . 7 Superseded by AALTONEN 17.					

 $\chi_{c1}(4140)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
22 ± 8 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			NODE=M193W
83 ± 21 -14	4289	1 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193W
15.3 ± 10.4 -6.1 ± 2.5	19	2 AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193W
16.3 ± 5.6 ± 11.4	616	3 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi \phi +$ anything	NODE=M193W
20 ± 13 -8	52	4 ABAZOV	14A D0	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193W
28 ± 15 -11	0.3k	5 CHATRCHYAN 14M	CMS	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193W
• • • We do not use the following data for averages, fits, limits, etc. • • •					
11.7 ± 8.3 -5.0 ± 3.7	14 6,7	AALTONEN 09AH	CDF	$B^+ \rightarrow J/\psi \phi K^+$	NODE=M193W

- 1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 8.4σ .
 2 Statistical significance of more than 5σ .
 3 Statistical significance of more than 6σ .
 4 Statistical significance of 3.1σ .
 5 From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ .
 6 Statistical significance of 3.8σ .
 7 Superseded by AALTONEN 17.

NODE=M193W;LINKAGE=D
 NODE=M193W;LINKAGE=E
 NODE=M193W;LINKAGE=C
 NODE=M193W;LINKAGE=A
 NODE=M193W;LINKAGE=B
 NODE=M193W;LINKAGE=AA
 NODE=M193W;LINKAGE=F



$\chi_{c1}(4140)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen
Γ_2 $\gamma\gamma$	not seen

NODE=M193215;NODE=M193

$\Gamma(\gamma\gamma) \times \Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	CL%
<41	90
¹ SHEN	
10 BELL	
10.6 $e^+ e^- \rightarrow e^+ e^- J/\psi\phi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
< 6	90
² SHEN	
10 BELL	
10.6 $e^+ e^- \rightarrow e^+ e^- J/\psi\phi$	
¹ For $J^P = 0^+$.	
² For $J^P = 2^+$.	

NODE=M193220

NODE=M193G01
NODE=M193G01

OCCUR=2

NODE=M193G01;LINKAGE=S0
NODE=M193G01;LINKAGE=S2

NODE=M193225

NODE=M193R01
NODE=M193R01

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	EVTS DOCUMENT ID TECN COMMENT
seen	4289 ¹ AAIJ 17C LHCb $B^+ \rightarrow J/\psi\phi K^+$
seen	616 ² ABAZOV 15M D0 $p\bar{p} \rightarrow J/\psi\phi + \text{anything}$
seen	52 ³ ABAZOV 14A D0 $B^+ \rightarrow J/\psi\phi K^+$
seen	0.3k ⁴ CHATRCHYAN 14M CMS $B^+ \rightarrow J/\psi\phi K^+$
seen	14 ⁵ AALTONEN 09AH CDF $B^+ \rightarrow J/\psi\phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
not seen	⁶ ABLIKIM 15 BES3 $e^+ e^- \rightarrow \gamma\phi J/\psi$
not seen	⁷ AAIJ 12AA LHCb $pp \rightarrow B^+ X \text{ at } 7 \text{ TeV}$

- 1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 8.4σ .
 2 Statistical significance of more than 6σ .
 3 ABAZOV 14A reports $B(B^+ \rightarrow \chi_{c1}(4140)K^+ \rightarrow J/\psi\phi K^+)/B(B^+ \rightarrow J/\psi\phi K^+) = (19 \pm 7 \pm 4)\%$ with 3.1σ significance.
 4 From a fit assuming an S -wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ .
 5 Statistical significance of 3.8σ .
 6 Reported $\sigma(e^+e^- \rightarrow \gamma\chi_{c1}(4140)) \cdot B(\chi_{c1}(4140) \rightarrow J/\psi\phi) < 0.35, 0.28, \text{ and } 0.33 \text{ pb}$ at $4.23, 4.26, \text{ and } 4.36 \text{ GeV}$, respectively, at $90\% \text{ CL}$.
 7 Reported $B(B^+ \rightarrow \chi_{c1}(4140)K^+) \cdot B(\chi_{c1}(4140) \rightarrow J/\psi\phi)/B(B^+ \rightarrow J/\psi\phi K^+) < 0.07$ at $90\% \text{ CL}$.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$			Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	SHEN	10 BELL	$10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$

$\chi_{c1}(4140)$ REFERENCES

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	17	MPL A32 1750139	T. Altonen <i>et al.</i>	(CDF Collab.)
ABAZOV	15M	PRL 115 232001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	15	PR D91 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABAZOV	14A	PR D89 012004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	14M	PL B734 261	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAIJ	12AA	PR D85 091103	R. Aaij <i>et al.</i>	(LHCb Collab.)
SHEN	10	PRL 104 112004	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AALTONEN	09AH	PRL 102 242002	T. Aaltonen <i>et al.</i>	(CDF Collab.)

$\psi(4160)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

$\psi(4160)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4191 \pm 5 OUR AVERAGE			
4191 \pm 9	AAIJ	13BC LHCb	$B^+ \rightarrow K^+\mu^+\mu^-$
4191.7 \pm 6.5	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4193 \pm 7	² MO	10 RVUE	$e^+e^- \rightarrow \text{hadrons}$
4151 \pm 4	³ SETH	05A RVUE	$e^+e^- \rightarrow \text{hadrons}$
4155 \pm 5	⁴ SETH	05A RVUE	$e^+e^- \rightarrow \text{hadrons}$
4159 \pm 20	BRANDELIK	78C DASP	e^+e^-
1 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.			
2 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
3 From a fit to Crystal Ball (OSTERHELD 86) data.			
4 From a fit to BES (BAI 02C) data.			

$\psi(4160)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70 \pm 10 OUR AVERAGE			
65 \pm 22	AAIJ	13BC LHCb	$B^+ \rightarrow K^+\mu^+\mu^-$
71.8 \pm 12.3	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
79 \pm 14	² MO	10 RVUE	$e^+e^- \rightarrow \text{hadrons}$
107 \pm 10	³ SETH	05A RVUE	$e^+e^- \rightarrow \text{hadrons}$
107 \pm 16	⁴ SETH	05A RVUE	$e^+e^- \rightarrow \text{hadrons}$
78 \pm 20	BRANDELIK	78C DASP	e^+e^-
1 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.			
2 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
3 From a fit to Crystal Ball (OSTERHELD 86) data.			
4 From a fit to BES (BAI 02C) data.			

NODE=M193R01;LINKAGE=E
 NODE=M193R01;LINKAGE=D
 NODE=M193R01;LINKAGE=A
 NODE=M193R01;LINKAGE=C
 NODE=M193R01;LINKAGE=AA
 NODE=M193R01;LINKAGE=B
 NODE=M193R01;LINKAGE=AI

NODE=M193R02
 NODE=M193R02

NODE=M193
 REFID=57657
 REFID=57636
 REFID=58161
 REFID=56957
 REFID=56368
 REFID=55650
 REFID=55753
 REFID=54263
 REFID=53235
 REFID=52968

NODE=M025

NODE=M025M

NODE=M025M

OCCUR=2

NODE=M025M;LINKAGE=AB

NODE=M025M;LINKAGE=MO

NODE=M025M;LINKAGE=ST

NODE=M025M;LINKAGE=SE

NODE=M025W

NODE=M025W

OCCUR=2

NODE=M025W;LINKAGE=AB

NODE=M025W;LINKAGE=MO

NODE=M025W;LINKAGE=ST

NODE=M025W;LINKAGE=SE

$\psi(4160)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode		Fraction (Γ_i/Γ)	Confidence level	
Γ_1	$e^+ e^-$	$(6.9 \pm 3.3) \times 10^{-6}$		DESIG=1
Γ_2	$\mu^+ \mu^-$	seen		DESIG=33
Γ_3	$D\bar{D}$	seen		DESIG=15;OUR EVAL; \rightarrow UNCHECKED \leftarrow
Γ_4	$D^0 \bar{D}^0$	seen		DESIG=16
Γ_5	$D^+ D^-$	seen		DESIG=17
Γ_6	$D^* \bar{D} + \text{c.c.}$	seen		DESIG=18;OUR EVAL; \rightarrow UNCHECKED \leftarrow
Γ_7	$D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen		DESIG=19
Γ_8	$D^*(2010)^+ D^- + \text{c.c.}$	seen		DESIG=20
Γ_9	$D^* \bar{D}^*$	seen		DESIG=21;OUR EVAL; \rightarrow UNCHECKED \leftarrow
Γ_{10}	$D^*(2007)^0 \bar{D}^*(2007)^0$	seen		DESIG=22
Γ_{11}	$D^*(2010)^+ D^*(2010)^-$	seen		DESIG=23
Γ_{12}	$D^0 D^- \pi^+ + \text{c.c. (excl.)}$ $D^*(2007)^0 \bar{D}^0 + \text{c.c.},$ $D^*(2010)^+ D^- + \text{c.c.})$	not seen		DESIG=24
Γ_{13}	$D\bar{D}^* \pi^+ + \text{c.c. (excl. } D^* \bar{D}^*)$	seen		DESIG=25
Γ_{14}	$D^0 D^{*-} \pi^+ + \text{c.c. (excl.)}$ $D^*(2010)^+ D^*(2010)^-$	not seen		DESIG=26
Γ_{15}	$D_s^+ D_s^-$	not seen		DESIG=27
Γ_{16}	$D_s^{*+} D_s^- + \text{c.c.}$	seen		DESIG=28
Γ_{17}	$J/\psi \pi^+ \pi^-$	$< 3 \times 10^{-3}$	90%	DESIG=2
Γ_{18}	$J/\psi \pi^0 \pi^0$	$< 3 \times 10^{-3}$	90%	DESIG=3
Γ_{19}	$J/\psi K^+ K^-$	$< 2 \times 10^{-3}$	90%	DESIG=4
Γ_{20}	$J/\psi \eta$	$< 8 \times 10^{-3}$	90%	DESIG=5
Γ_{21}	$J/\psi \pi^0$	$< 1 \times 10^{-3}$	90%	DESIG=6
Γ_{22}	$J/\psi \eta'$	$< 5 \times 10^{-3}$	90%	DESIG=7
Γ_{23}	$J/\psi \pi^+ \pi^- \pi^0$	$< 1 \times 10^{-3}$	90%	DESIG=8
Γ_{24}	$\psi(2S) \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%	DESIG=9
Γ_{25}	$\chi_{c1} \gamma$	$< 5 \times 10^{-3}$	90%	DESIG=10
Γ_{26}	$\chi_{c2} \gamma$	$< 1.3 \%$	90%	DESIG=11
Γ_{27}	$\chi_{c1} \pi^+ \pi^- \pi^0$	$< 2 \times 10^{-3}$	90%	DESIG=12
Γ_{28}	$\chi_{c2} \pi^+ \pi^- \pi^0$	$< 8 \times 10^{-3}$	90%	DESIG=13
Γ_{29}	$h_c(1P) \pi^+ \pi^-$	$< 5 \times 10^{-3}$	90%	DESIG=29
Γ_{30}	$h_c(1P) \pi^0 \pi^0$	$< 2 \times 10^{-3}$	90%	DESIG=30
Γ_{31}	$h_c(1P) \eta$	$< 2 \times 10^{-3}$	90%	DESIG=31
Γ_{32}	$h_c(1P) \pi^0$	$< 4 \times 10^{-4}$	90%	DESIG=32
Γ_{33}	$\phi \pi^+ \pi^-$	$< 2 \times 10^{-3}$	90%	DESIG=14
Γ_{34}	$\gamma \chi_{c1}(3872) \rightarrow \gamma J/\psi \pi^+ \pi^-$	$< 6.8 \times 10^{-5}$	90%	DESIG=34
Γ_{35}	$\gamma X(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-$	$< 1.36 \times 10^{-4}$	90%	DESIG=35
Γ_{36}	$\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-$	$< 1.18 \times 10^{-4}$	90%	DESIG=36
Γ_{37}	$\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-$	$< 1.47 \times 10^{-4}$	90%	DESIG=37
Γ_{38}	$\gamma \chi_{c1}(3872) \rightarrow \gamma \gamma J/\psi$	$< 1.05 \times 10^{-4}$	90%	DESIG=38
Γ_{39}	$\gamma X(3915) \rightarrow \gamma \gamma J/\psi$	$< 1.26 \times 10^{-4}$	90%	DESIG=39
Γ_{40}	$\gamma X(3930) \rightarrow \gamma \gamma J/\psi$	$< 8.8 \times 10^{-5}$	90%	DESIG=40
Γ_{41}	$\gamma X(3940) \rightarrow \gamma \gamma J/\psi$	$< 1.79 \times 10^{-4}$	90%	DESIG=41
Γ_{42}	$K^+ K^-$			DESIG=42
Γ_{43}	$K_S^0 K^\pm \pi^\mp$			DESIG=43

NODE=M025215;NODE=M025

NODE=M025

$\psi(4160)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$					Γ_1
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
0.48±0.22	1 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.4 to 1.1	2 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons		
0.83±0.08	3 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons		
0.84±0.13	4 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons		
0.77±0.23	BRANDELIK	78C DASP	$e^+ e^-$		
1 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.					
2 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.					
3 From a fit to Crystal Ball (OSTERHELD 86) data.					
4 From a fit to BES (BAI 02C) data.					

$\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$					
$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{25}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.2	90	1 HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c1}\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.					
$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{26}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.1	90	1 HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c2}\gamma$	
1 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.					
$\Gamma(J/\psi\eta') \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{22}\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.17±0.04	86	1,2 ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$	
1.07±0.09	86	1,3 ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$	
1 Based on a fit to $\sigma(e^+ e^- \rightarrow \eta' J/\psi)$ from $\sqrt{s} = 4.18$ to 4.60 GeV assuming interfering $\psi(4160)$ and $\psi(4260)$ contributions. At $\sqrt{s} = 4.18$ GeV, $\sigma(e^+ e^- \rightarrow \eta' J/\psi) = 2.4 \pm 0.3 \pm 0.2$ pb.					
2 Solution I of the fit, corresponding to a phase of -0.03 ± 0.44 rad.					
3 Solution II of the fit, corresponding to a phase of 2.54 ± 0.04 rad.					

$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		$\Gamma_{43}\Gamma_1/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.71 ± 0.13 ± 0.12	1 ABLIKIM	19AE BES3	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$		
0.0095±0.0088±0.0004	2 ABLIKIM	19AE BES3	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$		
1 Solution I of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.					
2 Solution II of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.					

$\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma^2(\text{total})$					
$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.8±0.9±0.9	1 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$		
12.8±1.7±2.0	2 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$		

NODE=M025220

NODE=M025W1

NODE=M025W1

OCCUR=2

NODE=M025W1;LINKAGE=AB

NODE=M025W1;LINKAGE=MO

NODE=M025W1;LINKAGE=ST

NODE=M025W1;LINKAGE=SE

NODE=M025235

NODE=M025G01

NODE=M025G01

NODE=M025G01;LINKAGE=A

NODE=M025G02

NODE=M025G02

NODE=M025G02;LINKAGE=A

NODE=M025R42

NODE=M025R42

OCCUR=2

NODE=M025R42;LINKAGE=A

NODE=M025R42;LINKAGE=B

NODE=M025R42;LINKAGE=C

NODE=M025R00

NODE=M025R00

OCCUR=5

NODE=M025R00;LINKAGE=A

NODE=M025R00;LINKAGE=D

NODE=M025230

NODE=M025R32

NODE=M025R32

OCCUR=2

¹Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

²Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

$\psi(4160)$ BRANCHING RATIOS

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	1 AAIJ	13BC LHCb	$B^+ \rightarrow K^+ \mu^+ \mu^-$	
			¹ AAIJ 13BC report $B(B^+ \rightarrow K^+ \psi(4160)) B(\psi(4160) \rightarrow \mu^+ \mu^-) = (3.5^{+0.9}_{-0.8}) \times 10^{-9}$.	

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_9
0.02±0.03±0.02	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^0\bar{D}^0$	
seen	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$	

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^+ D^-$	
seen	PAKHLOVA 08	BELL	$e^+ e^- \rightarrow D^+ D^- \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^+ D^- \gamma$	

$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^0\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0}\bar{D}^0$	

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
seen	1 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$	I
seen	AUBERT 09M BABR	e ⁺ e ⁻	$\rightarrow D^{*+} D^- \gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$	I

¹Supersedes PAKHLOVA 07.

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_9
0.34±0.14±0.05	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0}\bar{D}^{*0}$	

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
seen	1 ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	I
seen	AUBERT 09M BABR	e ⁺ e ⁻	$\rightarrow D^{*+} D^{*-} \gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	I

¹Supersedes PAKHLOVA 07.

NODE=M025R32;LINKAGE=A

NODE=M025R32;LINKAGE=B

NODE=M025225

NODE=M025R31
NODE=M025R31

NODE=M025R31;LINKAGE=A

NODE=M025R14
NODE=M025R14

NODE=M025R16
NODE=M025R16

NODE=M025R17
NODE=M025R17

NODE=M025R18
NODE=M025R18

NODE=M025R19
NODE=M025R19

NODE=M025R19;LINKAGE=A

NODE=M025R15
NODE=M025R15

NODE=M025R20
NODE=M025R20

NODE=M025R21
NODE=M025R21

NODE=M025R21;LINKAGE=A

$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.})) / \Gamma_{\text{total}}$				Γ_{12}/Γ	NODE=M025R22 NODE=M025R22
<u>VALUE</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
not seen				PAKHLOVA	08A BELL
$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$					
$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)) / \Gamma_{\text{total}}$				Γ_{13}/Γ	NODE=M025R23 NODE=M025R23
<u>VALUE</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
seen				CRONIN-HEN..09	CLEO
$e^+ e^- \rightarrow D \bar{D}^* \pi$					
$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)) / \Gamma_{\text{total}}$				Γ_{14}/Γ	NODE=M025R24 NODE=M025R24
<u>VALUE</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
not seen				PAKHLOVA	09 BELL
$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$					
$\Gamma(D_s^+ D_s^-) / \Gamma_{\text{total}}$				Γ_{15}/Γ	NODE=M025R25 NODE=M025R25
<u>VALUE</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
not seen				PAKHLOVA	11 BELL
$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$					
not seen				DEL-AMO-SA..10N	BABR
$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$					
not seen				CRONIN-HEN..09	CLEO
$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$					
$\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma_{\text{total}}$				Γ_{16}/Γ	NODE=M025R26 NODE=M025R26
<u>VALUE</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
seen				PAKHLOVA	11 BELL
$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$					
seen				DEL-AMO-SA..10N	BABR
$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$					
seen				CRONIN-HEN..09	CLEO
$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$					
$\Gamma(J/\psi \pi^+ \pi^-) / \Gamma_{\text{total}}$				Γ_{17}/Γ	NODE=M025R01 NODE=M025R01
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<3				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					
$\Gamma(J/\psi \pi^0 \pi^0) / \Gamma_{\text{total}}$				Γ_{18}/Γ	NODE=M025R02 NODE=M025R02
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<3				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					
$\Gamma(J/\psi K^+ K^-) / \Gamma_{\text{total}}$				Γ_{19}/Γ	NODE=M025R03 NODE=M025R03
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<2				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					
$\Gamma(J/\psi \eta) / \Gamma_{\text{total}}$				Γ_{20}/Γ	NODE=M025R04 NODE=M025R04
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<8				COAN	06 CLEO
90					
We do not use the following data for averages, fits, limits, etc. • • •					
possibly seen				¹ ABLIKIM	15L BES3
seen				WANG	13B BELL
$e^+ e^- \rightarrow J/\psi \eta$					
1 An enhancement around 4.2 GeV is observed.					
$\Gamma(J/\psi \pi^0) / \Gamma_{\text{total}}$				Γ_{21}/Γ	NODE=M025R05 NODE=M025R05
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<1				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					
$\Gamma(J/\psi \eta') / \Gamma_{\text{total}}$				Γ_{22}/Γ	NODE=M025R06 NODE=M025R06
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<5				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					
$\Gamma(J/\psi \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$				Γ_{23}/Γ	NODE=M025R07 NODE=M025R07
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<1				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					
$\Gamma(\psi(2S) \pi^+ \pi^-) / \Gamma_{\text{total}}$				Γ_{24}/Γ	NODE=M025R08 NODE=M025R08
<u>VALUE (units 10^{-3})</u>				<u>DOCUMENT ID</u>	<u>TECN</u>
<4				COAN	06 CLEO
90					
$4.12\text{--}4.2 e^+ e^- \rightarrow \text{hadrons}$					

$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7

90

COAN

06

CLEO

4.12–4.2 $e^+e^- \rightarrow$ hadrons Γ_{25}/Γ

NODE=M025R09
NODE=M025R09

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<13

90

COAN

06

CLEO

4.12–4.2 $e^+e^- \rightarrow$ hadrons Γ_{26}/Γ

NODE=M025R10
NODE=M025R10

 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2

90

COAN

06

CLEO

4.12–4.2 $e^+e^- \rightarrow$ hadrons Γ_{27}/Γ

NODE=M025R11
NODE=M025R11

 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<8

90

COAN

06

CLEO

4.12–4.2 $e^+e^- \rightarrow$ hadrons Γ_{28}/Γ

NODE=M025R12
NODE=M025R12

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<5

90

1 PEDLAR

11

CLEO

 $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

1 At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 Γ_{29}/Γ

NODE=M025R27
NODE=M025R27

 $\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2

90

1 PEDLAR

11

CLEO

 $e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

1 At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 Γ_{30}/Γ

NODE=M025R28
NODE=M025R28

 $\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	-------------	--------------------	-------------	----------------

<2

90

1 PEDLAR

11

CLEO

 $e^+e^- \rightarrow h_c(1P)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen

41

2 ABLIKIM

17R

BES3

 $e^+e^- \rightarrow h_c(1P)\eta$

1 At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

2 An enhancement around 4.2 GeV is observed.

 Γ_{31}/Γ

NODE=M025R29
NODE=M025R29

 $\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<0.4

90

1 PEDLAR

11

CLEO

 $e^+e^- \rightarrow h_c(1P)\pi^0$

1 At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

 Γ_{32}/Γ

NODE=M025R30
NODE=M025R30

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	------------	--------------------	-------------	----------------

<2

90

COAN

06

CLEO

4.12–4.2 $e^+e^- \rightarrow$ hadrons Γ_{33}/Γ

NODE=M025R13
NODE=M025R13

 $\Gamma(\gamma\chi_{c1}(3872) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
--------------	------------	--------------------	----------------

 $<0.68 \times 10^{-4}$

90

1 XIAO

13

 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 Γ_{34}/Γ

NODE=M025R34
NODE=M025R34

 $\Gamma(\gamma X(3915) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>
--------------	------------	--------------------	----------------

 $<1.36 \times 10^{-4}$

90

1 XIAO

13

 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

 Γ_{35}/Γ

NODE=M025R35
NODE=M025R35

NODE=M025R35;LINKAGE=A

$\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{36}/Γ
VALUE	CL%	DOCUMENT ID	COMMENT		
$<1.18 \times 10^{-4}$	90	1 XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$		NODE=M025R36 NODE=M025R36
1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{37}/Γ
VALUE	CL%	DOCUMENT ID	COMMENT		
$<1.47 \times 10^{-4}$	90	1 XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$		NODE=M025R37 NODE=M025R37
1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma \chi_{c1}(3872) \rightarrow \gamma \gamma J/\psi)/\Gamma_{\text{total}}$					Γ_{38}/Γ
VALUE	CL%	DOCUMENT ID	COMMENT		
$<1.05 \times 10^{-4}$	90	1 XIAO	13 $\psi(4160) \rightarrow \gamma \gamma J/\psi$		NODE=M025R38 NODE=M025R38
1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3915) \rightarrow \gamma \gamma J/\psi)/\Gamma_{\text{total}}$					Γ_{39}/Γ
VALUE	CL%	DOCUMENT ID	COMMENT		
$<1.26 \times 10^{-4}$	90	1 XIAO	13 $\psi(4160) \rightarrow \gamma \gamma J/\psi$		NODE=M025R39 NODE=M025R39
1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3930) \rightarrow \gamma \gamma J/\psi)/\Gamma_{\text{total}}$					Γ_{40}/Γ
VALUE	CL%	DOCUMENT ID	COMMENT		
$<0.88 \times 10^{-4}$	90	1 XIAO	13 $\psi(4160) \rightarrow \gamma \gamma J/\psi$		NODE=M025R40 NODE=M025R40
1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3940) \rightarrow \gamma \gamma J/\psi)/\Gamma_{\text{total}}$					Γ_{41}/Γ
VALUE	CL%	DOCUMENT ID	COMMENT		
$<1.79 \times 10^{-4}$	90	1 XIAO	13 $\psi(4160) \rightarrow \gamma \gamma J/\psi$		NODE=M025R41 NODE=M025R41
1 Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$					Γ_{42}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2 \times 10^{-5}$	90	1 DRUZHININ	15 RVUE	$e^+ e^- \rightarrow \psi(3770)$	NODE=M025R33 NODE=M025R33
1 DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+ e^- \rightarrow K^+ K^-$ and $e^+ e^- \rightarrow K_S^0 K_L^0$.					

$\psi(4160)$ REFERENCES

ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan	
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)
ABLIKIM	15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)
HAN	15	PR D92 021011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
XIAO	13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH	05A	PR D72 017501	K.K. Seth	
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)

NODE=M025

REFID=60210
 REFID=59856
 REFID=58710
 REFID=58009
 REFID=57991
 REFID=57140
 REFID=56777
 REFID=56962
 REFID=56816
 REFID=55229
 REFID=55377
 REFID=55381
 REFID=53638
 REFID=16787
 REFID=53532
 REFID=53540
 REFID=52724
 REFID=53114
 REFID=53143
 REFID=52142
 REFID=52132
 REFID=52134
 REFID=51628
 REFID=51075
 REFID=50813
 REFID=50506
 REFID=50503
 REFID=51064
 REFID=22232

NODE=M190

X(4160)

$$I^G(J^{PC}) = ??(??)$$

OMMITTED FROM SUMMARY TABLE

Seen by PAKHLOV 08 in $e^+e^- \rightarrow J/\psi X, X \rightarrow D^* \bar{D}^*$ **X(4160) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4156⁺²⁵₋₂₀^{±15}	24	PAKHLOV	08	BELL $e^+e^- \rightarrow J/\psi X$

X(4160) WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
139⁺¹¹¹₋₆₁^{±21}	24	PAKHLOV	08	BELL $e^+e^- \rightarrow J/\psi X$

X(4160) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D\bar{D}$	not seen
$\Gamma_2 D^*\bar{D} + c.c.$	not seen
$\Gamma_3 D^*\bar{D}^*$	seen

X(4160) BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$	Γ_1/Γ_3
VALUE	CL%
<0.09	90
PAKHLOV	08
BELL	$e^+e^- \rightarrow J/\psi X$
$\Gamma(D^*\bar{D} + c.c.)/\Gamma(D^*\bar{D}^*)$	Γ_2/Γ_3
VALUE	CL%
<0.22	90
PAKHLOV	08
BELL	$e^+e^- \rightarrow J/\psi X$

X(4160) REFERENCESPAKHLOV 08 PRL 100 202001 P. Pakhlov *et al.* (BELLE Collab.)**Z_c(4200)**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

I, G, C need confirmation.

OMMITTED FROM SUMMARY TABLE

was $X(4200)^{\pm}$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Reported by CHILIKIN 14 in $J/\psi\pi^+$ at a significance of 6.2σ . Assignments of $0^-, 1^-, 2^-,$ and 2^+ excluded at $6.1\sigma, 7.4\sigma, 4.4\sigma,$ and 7.0σ level, respectively. Needs confirmation.

Z_c(4200) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4196⁺³¹⁺¹⁷₋₂₉₋₁₃	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^-\pi^+$

Z_c(4200) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
370^{±70}₋₁₃₂⁺⁷⁰	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^-\pi^+$

NODE=M190

NODE=M190

NODE=M190M

NODE=M190M

NODE=M190W

NODE=M190W

NODE=M190215;NODE=M190

DESIG=1;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow

NODE=M190225

NODE=M190R01
NODE=M190R01NODE=M190R02
NODE=M190R02

NODE=M190

REFID=52302

NODE=M231

NODE=M231

NODE=M231M

NODE=M231M

NODE=M231W

NODE=M231W

Z_c(4200) DECAY MODES

NODE=M231215;NODE=M231

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad J/\psi\pi^+$	seen

Z_c(4200) BRANCHING RATIOS

$\Gamma(J/\psi\pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE	DOCUMENT ID TECN COMMENT
seen CHILIKIN 14 BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
possibly seen 1 AAIJ 19R LHCb $B^0 \rightarrow K^+ \pi^- J/\psi + \text{c.c.}$	

¹ From a model-independent analysis.**Z_c(4200) REFERENCES**

AAIJ 19R PRL 122 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)
CHILIKIN 14 PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)

 $\psi(4230)$

$$\Gamma^G(J^{PC}) = 0^-(1^{--})$$

was $X(4230)$

The recent measurement of $e^+ e^- \rightarrow J/\psi\pi\pi$ (ABLIKIM 17B) led to a downward shift in the mass of the $\psi(4260)$, also known as $Y(4260)$, such that a distinction between the $\psi(4260)$ and $\psi(4230)$ no longer appears justified. Therefore, starting from this edition, we include the data of ABLIKIM 17B in this node and have listed the $\psi(4230)$ in the summary tables instead of the $\psi(4260)$.

 $\psi(4230)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4220 ± 15 OUR ESTIMATE				
4219.9 ± 2.2 OUR AVERAGE				
[4218 ⁺⁵ ₋₄ MeV OUR 2019 AVERAGE Scale factor = 1.2]				
4218.5 ± 1.6 ± 4.0	1 ABLIKIM	19AI BES3	$e^+ e^- \rightarrow \omega\chi_{c0}$	
4228.6 ± 4.1 ± 6.3	ABLIKIM	19R BES3	$e^+ e^- \rightarrow \pi^+ D^0 D^{*-+}$ c.c.	
4222.0 ± 3.1 ± 1.4	2 ABLIKIM	17B BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$	
4218 ± 5.5 ± 0.9	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$	
4209.5 ± 7.4 ± 1.4	3 ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4200.6 ± 7.9 ± 3.0	4 ABLIKIM	19V BES3	$e^+ e^- \rightarrow \gamma\chi_{c1}(3872)$	
4230 ± 8 ± 6	180 5 ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega\chi_{c0}$	

- 1 From a fit of the measured cross section from $\sqrt{s} = 4.178\text{--}4.278$ GeV. Supersedes ABLIKIM 15C.
- 2 From a three-resonance fit.
- 3 From a fit to the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb⁻¹.
- 4 Simultaneous fit to $\chi_{c1} \rightarrow \omega J/\psi$ and $\chi_{c1} \rightarrow \pi^+ \pi^- J/\psi$.
- 5 From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$, $\chi_{c0} \rightarrow K^+ K^-$, and $\omega \rightarrow \pi^+ \pi^- \pi^0$.

 $\psi(4230)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 to 100 OUR ESTIMATE				
43 ± 9 OUR AVERAGE Error includes scale factor of 3.3. See the ideogram below.				
[59 ⁺¹² ₋₁₀ MeV OUR 2019 AVERAGE Scale factor = 1.5]				
28.2 ± 3.9 ± 1.6	1 ABLIKIM	19AI BES3	$e^+ e^- \rightarrow \omega\chi_{c0}$	

DESIG=1

NODE=M231220

NODE=M231R01
NODE=M231R01

NODE=M231R01;LINKAGE=C

NODE=M231

REFID=59776
REFID=56344

NODE=M222

NODE=M222

NODE=M222M

NODE=M222M
→ UNCHECKED ←
NEW

NODE=M222M;LINKAGE=C

NODE=M222M;LINKAGE=B

NODE=M222M;LINKAGE=F
NODE=M222M;LINKAGE=A

NODE=M222W

NODE=M222W
→ UNCHECKED ←
NEW

$77.0 \pm 6.8 \pm 6.3$	ABLIKIM	19R BES3	$e^+ e^- \rightarrow \pi^+ D^0 D^{*-} +$
$44.1 \pm 4.3 \pm 2.0$	² ABLIKIM	17B BES3	$e^+ e^- \xrightarrow{\text{c.c.}} \pi^+ \pi^- J/\psi$
$66.0 \pm 12.3 \pm 0.4$	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$
$80.1 \pm 24.6 \pm 2.9$	³ ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$			
$115 \pm 38 \pm 12$	⁴ ABLIKIM	19V BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$
$38 \pm 12 \pm 2$	⁵ ABLIKIM	15C BES3	$e^+ e^- \rightarrow \omega \chi_{c0}$

¹ From a fit of the measured cross section from $\sqrt{s} = 4.178\text{--}4.278$ GeV. Supersedes ABLIKIM 15C.

² From a three-resonance fit.

³ From a fit to the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-) \ell^+ \ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} .

⁴ Simultaneous fit to $\chi_{c1} \rightarrow \omega J/\psi$ and $\chi_{c1} \rightarrow \pi^+ \pi^- J/\psi$.

⁵ From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$, $\chi_{c0} \rightarrow K^+ K^-$, and $\omega \rightarrow \pi^+ \pi^- \pi^0$.

NODE=M222W;LINKAGE=C


NODE=M222W;LINKAGE=D

NODE=M222W;LINKAGE=B

NODE=M222W;LINKAGE=E

NODE=M222W;LINKAGE=A

WEIGHTED AVERAGE
43 \pm 9 (Error scaled by 3.3)



$\psi(4230)$ WIDTH (MeV)

$\psi(4230)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	
$\Gamma_2 \omega \chi_{c0}$	seen
$\Gamma_3 \pi^+ \pi^- h_c$	seen
$\Gamma_4 \pi^+ \pi^- \psi(2S)$	seen
$\Gamma_5 \pi^+ D^0 D^{*-} + \text{c.c.}$	seen
$\Gamma_6 \Xi^- \bar{\Xi}^+$	

NODE=M222215;NODE=M222

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

NODE=M222220

NODE=M222G01

NODE=M222G01

NEW

NODE=M222G01;LINKAGE=B

NODE=M222G01;LINKAGE=A

$\Gamma(\omega \chi_{c0}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_1/\Gamma$
VALUE (eV)	EVTS DOCUMENT ID TECN COMMENT
2.5 ± 0.4 OUR AVERAGE	[2.7 ± 0.6 eV OUR 2019 AVERAGE]
$2.5 \pm 0.2 \pm 0.3$	¹ ABLIKIM 19AI BES3 $e^+ e^- \rightarrow \omega \chi_{c0}$
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$	
$2.7 \pm 0.5 \pm 0.4$	180 ² ABLIKIM 15C BES3 $e^+ e^- \rightarrow \omega \chi_{c0}$

¹ From a fit of the measured cross section from $\sqrt{s} = 4.178\text{--}4.278$ GeV. Supersedes ABLIKIM 15C.

² From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+ \pi^-$, $\chi_{c0} \rightarrow K^+ K^-$, and $\omega \rightarrow \pi^+ \pi^- \pi^0$.

$\Gamma(\pi^+\pi^-\psi(2S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_4\Gamma_1/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		NODE=M222R03 NODE=M222R03
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.6 \pm 1.3	¹ ABLIKIM	19K	BES3 $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$		OCCUR=2
1.8 \pm 1.4	² ABLIKIM	19K	BES3 $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$		NODE=M222R03;LINKAGE=A NODE=M222R03;LINKAGE=B
1 Solution I of two equivalent solutions in a fit using two interfering resonances. 2 Solution II of two equivalent solutions in a fit using two interfering resonances.					
$\psi(4230)$ BRANCHING RATIOS					
$\Gamma(\omega\chi_{c0})/\Gamma_{\text{total}}$				Γ_2/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	180	¹ ABLIKIM	15C	BES3 $e^+e^- \rightarrow \omega\chi_{c0}$	NODE=M222225
1 From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.					
$\Gamma(\pi^+\pi^-h_c)/\Gamma_{\text{total}}$				Γ_3/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		ABLIKIM	17G	BES3 $e^+e^- \rightarrow \pi^+\pi^-h_c$	NODE=M222R00 NODE=M222R00
$\Gamma(\pi^+\pi^-\psi(2S))/\Gamma_{\text{total}}$				Γ_4/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		¹ ABLIKIM	17V	BES3 $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	NODE=M222R02 NODE=M222R02
1 From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb $^{-1}$.					
$\Gamma(\pi^+D^0D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$				Γ_5/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		ABLIKIM	19R	BES3 $e^+e^- \rightarrow \pi^+D^0D^{*-} + \text{c.c.}$	NODE=M222R04 NODE=M222R04
$\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_6\Gamma_1/\Gamma$	
VALUE (10 $^{-3}$ eV)	CL%	DOCUMENT ID	TECN	COMMENT	
0.33	90	ABLIKIM	20C	BES3 $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$	NODE=M222R05 NODE=M222R05

 $\psi(4230)$ REFERENCES

ABLIKIM	20C	PRL 124 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60215
ABLIKIM	19A1	PR D99 091103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59871
ABLIKIM	19K	PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	19R	PRL 122 102002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59765
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796
ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57755
ABLIKIM	17G	PRL 118 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57915
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58029
Also		PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	15C	PRL 114 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56401

NODE=M222

NODE=M216

R_{c0}(4240)

$I^G(J^{PC}) = 1^+(0^{--})$
I, G, C need confirmation.

OMMITTED FROM SUMMARY TABLE
 was $X(4240)^{\pm}$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Spin and parity assignment $J^P = 0^-$ is favored over $1^-, 2^-$, and 2^+ by 8σ and over 1^+ by 1σ , according to the four-dimensional amplitude analysis of AAIJ 14AG.

NODE=M216

R_{c0}(4240) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4239±18⁺⁴⁵₋₁₀	¹ AAIJ	14AG LHCb	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $Z_c(4430)$ fit, with significance 6σ including systematic variations.

NODE=M216M

NODE=M216M

NODE=M216M;LINKAGE=AA

R_{c0}(4240) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
220±47⁺¹⁰⁸₋₇₄	¹ AAIJ	14AG LHCb	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $Z_c(4430)$ fit, with significance 6σ including systematic variations.

NODE=M216W

NODE=M216W

NODE=M216W;LINKAGE=AA

R_{c0}(4240) DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^- \psi(2S)$	seen

NODE=M216215;NODE=M216

DESIG=1

NODE=M216225

NODE=M216R01
 NODE=M216R01

NODE=M216R01;LINKAGE=AA

NODE=M216

REFID=55896

R_{c0}(4240) BRANCHING RATIOS

$\Gamma(\pi^- \psi(2S))/\Gamma_{\text{total}}$			Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	14AG LHCb	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $Z_c(4430)$ fit. No partial branching fraction quoted.

R_{c0}(4240) REFERENCES

AAIJ	14AG PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	---------------------	-----------------------	----------------

X(4250) $^\pm$

$I^G(JPC) = 1^-(?^+)$
 I, G, C need confirmation.

OMITTED FROM SUMMARY TABLE

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Observed by MIZUK 08 in the $\pi^+ \chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

NODE=M192

X(4250) $^\pm$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4248 $+44 +180$ $-29 - 35$	¹ MIZUK	08 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192M

NODE=M192M

NODE=M192M;LINKAGE=MI

NODE=M192W

NODE=M192W

NODE=M192W;LINKAGE=MI

NODE=M192215;NODE=M192

X(4250) $^\pm$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
177 $+54 +316$ $-39 - 61$	¹ MIZUK	08 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192W

NODE=M192W

NODE=M192W;LINKAGE=MI

X(4250) $^\pm$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^+ \chi_{c1}(1P)$	seen

DESIG=1

X(4250) $^\pm$ BRANCHING RATIOS

$\Gamma(\pi^+ \chi_{c1}(1P))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	Γ_1/Γ
seen	¹ MIZUK	08 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}(1P)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ² LEES 12B BABR $B \rightarrow K\pi \chi_{c1}(1P)$

¹ With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- X(4250)^+) \times B(X(4250)^+ \rightarrow \pi^+ \chi_{c1}(1P)) = (4.0^{+2.3+19.7}_{-0.9-0.5}) \times 10^{-5}$.

² With a product branching fraction limit of $B(\bar{B}^0 \rightarrow X(4250)^+ K^-) \times B(X(4250)^+ \rightarrow \chi_{c1} \pi^+) < 4.0 \times 10^{-5}$ at 90% CL.

NODE=M192225

NODE=M192R01
NODE=M192R01

NODE=M192R01;LINKAGE=MI

NODE=M192R01;LINKAGE=LE

X(4250) $^\pm$ REFERENCES

LEES	12B PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08 PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M192

REFID=54042
REFID=52535

NODE=M074

 $\psi(4260)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

OMMITTED FROM SUMMARY TABLE
also known as $Y(4260)$; was $X(4260)$

The state $\psi(4260)$ received its mass label from a Breit-Wigner (BW) fit to the $J/\psi\pi\pi$ data listed below. The symmetric BW placed the mass unavoidably into the center of the distribution. The most recent measurement in the 4260 MeV mass range in the same channel (ABLIKIM 17B), however, revealed that the distribution is asymmetric and that the state has a much lower mass consistent with the entry for particle $\psi(4230)$. Thus, in this edition we merged the measurement of ABLIKIM 17B with the $\psi(4230)$ node and labeled the older measurements of this node as not used. For details see the review on "Spectroscopy of mesons containing two heavy quarks."

NODE=M074

 $\psi(4260)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4209.1 ± 6.8 ± 7.0	1 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
4223.3 ± 1.6 ± 2.5	2 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$	
4258.6 ± 8.3 ± 12.1	3 LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
4245 ± 5 ± 4	4 LEES	12AC BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
4247 ± 12 ± 17	3,5 YUAN	07 BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
4284 ± 17 ± 413.6	HE	06B CLEO	$9.4-10.6 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
4259 ± 8 ± 2	125	6 AUBERT,B	05I BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$

NODE=M074M

NODE=M074M

OCCUR=2

1 From a three-resonance fit.

2 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

3 From a two-resonance fit.

4 From a single-resonance fit. Supersedes AUBERT,B 05I.

5 Superseded by LIU 13B.

6 From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074M;LINKAGE=D

NODE=M074M;LINKAGE=C

NODE=M074M;LINKAGE=YU

NODE=M074M;LINKAGE=LE

NODE=M074M;LINKAGE=YN

NODE=M074M;LINKAGE=AU

 $\psi(4260)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
76.6 ± 14.2 ± 2.4	1 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
54.2 ± 2.6 ± 1.0	2 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$	
134.1 ± 16.4 ± 5.5	3 LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
114 ± 16 ± 7	4 LEES	12AC BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
108 ± 19 ± 10	3,5 YUAN	07 BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
73 ± 39 ± 5	13.6 HE	06B CLEO	$9.4-10.6 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
88 ± 23 ± 6	125	6 AUBERT,B	05I BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$

NODE=M074W

NODE=M074W

1 From a three-resonance fit.

2 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

3 From a two-resonance fit.

4 From a single-resonance fit. Supersedes AUBERT,B 05I.

5 Superseded by LIU 13B.

6 From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074W;LINKAGE=C

NODE=M074W;LINKAGE=B

NODE=M074W;LINKAGE=YU

NODE=M074W;LINKAGE=LE

NODE=M074W;LINKAGE=YN

NODE=M074W;LINKAGE=AU

$\psi(4260)$ DECAY MODES

NODE=M074215;NODE=M074

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 e^+ e^-$		DESIG=1
$\Gamma_2 J/\psi \pi^+ \pi^-$	seen	DESIG=2;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_3 J/\psi f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$	seen	DESIG=41;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_4 Z_c(3900)^\pm \pi^\mp, Z_c^\pm \rightarrow J/\psi \pi^\pm$	seen	DESIG=43;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_5 J/\psi \pi^0 \pi^0$	seen	DESIG=4;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_6 J/\psi K^+ K^-$	seen	DESIG=5;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_7 J/\psi K_S^0 K_S^0$	not seen	DESIG=44
$\Gamma_8 J/\psi \eta$	not seen	DESIG=6;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_9 J/\psi \pi^0$	not seen	DESIG=7;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{10} J/\psi \eta'$	not seen	DESIG=8;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{11} J/\psi \pi^+ \pi^- \pi^0$	not seen	DESIG=9;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{12} J/\psi \eta \pi^0$	not seen	DESIG=45
$\Gamma_{13} J/\psi \eta \eta$	not seen	DESIG=10;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{14} \psi(2S) \pi^+ \pi^-$	not seen	DESIG=11;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{15} \psi(2S) \eta$	not seen	DESIG=12;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{16} \chi_{c0} \omega$	not seen	DESIG=13;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{17} \chi_{c1} \pi^+ \pi^- \pi^0$	not seen	DESIG=16;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{18} \chi_{c2} \pi^+ \pi^- \pi^0$	not seen	DESIG=17;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{19} h_c(1P) \pi^+ \pi^-$	not seen	DESIG=40;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{20} \phi \pi^+ \pi^-$	not seen	DESIG=18;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{21} \phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	not seen	DESIG=22;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{22} D \bar{D}$	not seen	DESIG=19;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{23} D^0 \bar{D}^0$	not seen	DESIG=31
$\Gamma_{24} D^+ D^-$	not seen	DESIG=32
$\Gamma_{25} D^* \bar{D} + c.c.$	not seen	DESIG=23;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{26} D^*(2007)^0 \bar{D}^0 + c.c.$	not seen	DESIG=33
$\Gamma_{27} D^*(2010)^+ D^- + c.c.$	not seen	DESIG=34
$\Gamma_{28} D^* \bar{D}^*$	not seen	DESIG=24;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{29} D^*(2007)^0 \bar{D}^*(2007)^0$	not seen	DESIG=35
$\Gamma_{30} D^*(2010)^+ D^*(2010)^-$	not seen	DESIG=36
$\Gamma_{31} D \bar{D} \pi + c.c.$		DESIG=37
$\Gamma_{32} D^0 D^- \pi^+ + c.c. \text{ (excl.)}$	not seen	DESIG=38
$D^*(2007)^0 \bar{D}^{*0} + c.c.,$		
$D^*(2010)^+ D^- + c.c.)$		
$\Gamma_{33} D \bar{D}^* \pi + c.c. \text{ (excl. } D^* \bar{D}^*)$	not seen	DESIG=25
$\Gamma_{34} D^0 D^{*-} \pi^+ + c.c. \text{ (excl.}$	not seen	DESIG=39
$D^*(2010)^+ D^*(2010)^-$		
$\Gamma_{35} D^0 D^*(2010)^- \pi^+ + c.c.$	not seen	DESIG=30;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{36} D_1(2420) \bar{D} + c.c.$	not seen	DESIG=50
$\Gamma_{37} D^* \bar{D}^* \pi$	not seen	DESIG=26
$\Gamma_{38} D_s^+ D_s^-$	not seen	DESIG=27
$\Gamma_{39} D_s^{*+} D_s^- + c.c.$	not seen	DESIG=28
$\Gamma_{40} D_s^{*+} D_s^-$	not seen	DESIG=29
$\Gamma_{41} p \bar{p}$	not seen	DESIG=3;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{42} p \bar{p} \pi^0$	not seen	DESIG=46;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{43} \Xi^- \bar{\Xi}^+$		DESIG=51
$\Gamma_{44} K_S^0 K^\pm \pi^\mp$	not seen	DESIG=20;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{45} K_S^0 K^\pm \pi^\mp \pi^0$		DESIG=48
$\Gamma_{46} K_S^0 K^\pm \pi^\mp \eta$		DESIG=49
$\Gamma_{47} K^+ K^- \pi^0$	not seen	DESIG=21;OUR EVAL; \rightarrow UNCHECKED \leftarrow
Radiative decays		
$\Gamma_{48} \eta_c(1S) \gamma$	possibly seen	NODE=M074;CLUMP=C
$\Gamma_{49} \chi_{c1} \gamma$	not seen	DESIG=47
$\Gamma_{50} \chi_{c2} \gamma$	not seen	DESIG=14;OUR EVAL; \rightarrow UNCHECKED \leftarrow
$\Gamma_{51} \chi_{c1}(3872) \gamma$	seen	DESIG=15;OUR EVAL; \rightarrow UNCHECKED \leftarrow
		DESIG=42

$\psi(4260) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_1 / \Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±1.0 OUR AVERAGE				
9.2±0.8±0.7	1 LEES	12AC BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
8.9 ^{+3.9} _{-3.1} ±1.8	8.1 HE	06B CLEO	9.4–10.6 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.4±0.8±0.6	2 LIU	13B BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	OCCUR=2
20.5±1.4±2.0	3 LIU	13B BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
6.0±1.2 ^{+4.7} _{-0.5}	2,4 YUAN	07 BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
20.6±2.3 ^{+9.1} _{-1.7}	3,4 YUAN	07 BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
5.5±1.0 ^{+0.8} _{-0.7}	125 5 AUBERT,B	05I BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	

¹ From a single-resonance fit. Supersedes AUBERT,B 05I.² Solution I of two equivalent solutions in a fit using two interfering resonances.³ Solution II of two equivalent solutions in a fit using two interfering resonances.⁴ Superseded by LIU 13B.⁵ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

$\Gamma(J/\psi K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_6 \Gamma_1 / \Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	1 SHEN	14 BELL	9.4–10.9 $e^+ e^- \rightarrow \gamma K^+ K^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2	90	2 YUAN	08 BELL	$e^+ e^- \rightarrow \gamma K^+ K^- J/\psi$
------	----	--------	---------	---

¹ From a fit of the broad $K^+ K^- J/\psi$ enhancement including a coherent $\psi(4260)$ amplitude with mass and width from LIU 13B. Supersedes YUAN 08. The shape of the cross section observed by ABLIKIM 18N between 2.2 and 2.3 GeV is incompatible with that of $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ in ABLIKIM 13T and ABLIKIM 17B. They also observe a broad enhancement around 2.5 GeV.² From a fit of the broad $K^+ K^- J/\psi$ enhancement including a coherent $\psi(4260)$ amplitude with mass and width from YUAN 07.

$\Gamma(J/\psi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_7 \Gamma_1 / \Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.85	90	1 SHEN	14 BELL	9.4–10.9 $e^+ e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$

¹ From a fit of the $K_S^0 K_S^0 J/\psi$ mass range from 4.4 to 5.5 GeV including a coherent $\psi(4260)$ amplitude with mass and width from LIU 13B.

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_8 \Gamma_1 / \Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<14.2	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

$\Gamma(J/\psi \eta') \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{10} \Gamma_1 / \Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.06±0.03	46 1,2 ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$	OCCUR=2
1.38±0.11	46 1,3 ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$	

¹ Based on a fit to $\sigma(e^+ e^- \rightarrow \eta' J/\psi)$ from $\sqrt{s} = 4.18$ to 4.60 GeV assuming interfering $\psi(4160)$ and $\psi(4260)$ contributions. At $\sqrt{s} = 4.23$ GeV, $\sigma(e^+ e^- \rightarrow \eta' J/\psi) = 3.6 \pm 0.6 \pm 0.3$ pb.² Solution I of the fit, corresponding to a phase of -0.03 ± 0.44 rad.³ Solution II of the fit, corresponding to a phase of 2.54 ± 0.04 rad.

$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{14} \Gamma_1 / \Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<4.3	90	1 LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^- \gamma$
7.4 ^{+2.1} _{-1.7}	2 LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^- \gamma$	

¹ For constructive interference with the $\psi(4360)$ in a combined fit of AUBERT 07s and WANG 07D data with three resonances.² For destructive interference with the $\psi(4360)$ in a combined fit of AUBERT 07s and WANG 07D data with three resonances.

NODE=M074230

NODE=M074G1

NODE=M074G1

OCCUR=2

OCCUR=2

NODE=M074G1;LINKAGE=LE
NODE=M074G1;LINKAGE=YU
NODE=M074G1;LINKAGE=YA
NODE=M074G1;LINKAGE=YN
NODE=M074G1;LINKAGE=AUNODE=M074G3
NODE=M074G3

NODE=M074G3;LINKAGE=A

NODE=M074G3;LINKAGE=YU

NODE=M074G02
NODE=M074G02

NODE=M074G02;LINKAGE=A

NODE=M074G01
NODE=M074G01NODE=M074R34
NODE=M074R34

OCCUR=2

NODE=M074R34;LINKAGE=A

NODE=M074R34;LINKAGE=B
NODE=M074R34;LINKAGE=CNODE=M074G7
NODE=M074G7

OCCUR=2

NODE=M074G7;LINKAGE=LI

NODE=M074G7;LINKAGE=LU

$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{20}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.4	90	AUBERT,BE	06D	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{21}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.28	90	¹ AUBERT	07AK	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
1 AUBERT 07AK reports $[\Gamma(\psi(4260) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(4260) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] < 0.14 \text{ eV}$ which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = 49.2 \times 10^{-2}$.					
$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{44}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.04 ± 0.19 ± 0.09	1	ABLIKIM	19AE	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0027 ± 0.0023 ± 0.0001	2	ABLIKIM	19AE	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
< 0.5 at 90% CL		AUBERT	08S	BABR	$10.6 e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$
1 Solution I of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.					
2 Solution II of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.					
$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{45}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.05	90	ABLIKIM	19	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$
$\Gamma(K_S^0 K^\pm \pi^\mp \eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{46}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.19	90	ABLIKIM	19	BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \eta$
$\Gamma(K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{47}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.6	90	AUBERT	08S	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\gamma$
$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{49}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.4	90	¹ HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$
1 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.					
$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{50}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<4.0	90	¹ HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$
1 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.					
$\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{43}\Gamma_1/\Gamma$
VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT	
0.27	90	ABLIKIM	20C	BES3	$e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$
$\psi(4260)$ BRANCHING RATIOS					
$\Gamma(J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_3/Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.17 ± 0.13	¹ LEES	12AC	BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$	
1 Systematic uncertainties not estimated.					

$\Gamma(Z_c(3900)^{\pm} \pi^{\mp}, Z_c^{\pm} \rightarrow J/\psi \pi^{\pm}) / \Gamma(J/\psi \pi^+ \pi^-)$					Γ_4 / Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT		
0.215 ± 0.033 ± 0.075	1 ABLIKIM	13T BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$		
• • • We do not use the following data for averages, fits, limits, etc. • • •					

0.29 ± 0.08 2 LIU 13B BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

1 Assuming that the cross section of $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ is fully due to the $\psi(4260)$.

2 Systematic error not evaluated.

$\Gamma(J/\psi K_S^0 K_S^0) / \Gamma_{\text{total}}$					Γ_7 / Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	SHEN	14	BELL	$9.4 - 10.9 e^+ e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$	

$\Gamma(J/\psi \eta \pi^0) / \Gamma_{\text{total}}$					Γ_{12} / Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	ABLIKIM	15Q BES3	$4.0 - 4.6 e^+ e^- \rightarrow J/\psi \eta \pi^0$		

$\Gamma(\psi(2S) \pi^+ \pi^-) / \Gamma(J/\psi \pi^+ \pi^-)$					Γ_{14} / Γ_2
VALUE	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					

(0.11 ± 0.03 ± 0.03) to (0.55 ± 0.18 ± 0.19) 1 ZHANG 17C RVUE $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$

1 From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.

$\Gamma(h_c(1P) \pi^+ \pi^-) / \Gamma(J/\psi \pi^+ \pi^-)$					Γ_{19} / Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	1 PEDLAR	11 CLEO	$e^+ e^- \rightarrow h_c(1P) \pi^+ \pi^-$	

1 At $\sqrt{s} = 4260$ MeV, PEDLAR 11 measures $\sigma(e^+ e^- \rightarrow h_c(1P) \pi^+ \pi^-) = 32 \pm 17 \pm 6 \pm 6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

$\Gamma(D\bar{D}) / \Gamma(J/\psi \pi^+ \pi^-)$					Γ_{22} / Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0	90	1 AUBERT	07BE BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

<4.0 90 CRONIN-HEN..09 CLEO $e^+ e^-$

1 Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$.

$\Gamma(D^0 \bar{D}^0) / \Gamma_{\text{total}}$					Γ_{23} / Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen					
• • • We do not use the following data for averages, fits, limits, etc. • • •					

not seen AUBERT 09M BABR $e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$
not seen PAKHLOVA 08 BELL $e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$

$\Gamma(D^+ D^-) / \Gamma_{\text{total}}$					Γ_{24} / Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen					
• • • We do not use the following data for averages, fits, limits, etc. • • •					

not seen AUBERT 09M BABR $e^+ e^- \rightarrow D^+ D^- \gamma$
not seen PAKHLOVA 08 BELL $e^+ e^- \rightarrow D^+ D^- \gamma$

$\Gamma(D^* \bar{D} + \text{c.c.}) / \Gamma(J/\psi \pi^+ \pi^-)$					Γ_{25} / Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<34	90	AUBERT	09M BABR	$e^+ e^- \rightarrow \gamma D^* \bar{D}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

<45 90 CRONIN-HEN..09 CLEO $e^+ e^-$

$\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.}) / \Gamma_{\text{total}}$					Γ_{26} / Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen					
• • • We do not use the following data for averages, fits, limits, etc. • • •					

not seen AUBERT 09M BABR $e^+ e^- \rightarrow D^* \bar{D}^0 \gamma$

NODE=M074R01
NODE=M074R01

NODE=M074R01;LINKAGE=AB
NODE=M074R01;LINKAGE=A

NODE=M074R27
NODE=M074R27

NODE=M074R28
NODE=M074R28

NODE=M074R25
NODE=M074R25

NODE=M074R2
NODE=M074R2

NODE=M074R12
NODE=M074R12

NODE=M074R13
NODE=M074R13

NODE=M074R03
NODE=M074R03

NODE=M074R14
NODE=M074R14

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{27}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* + D^-$	
not seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^* + D^- \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT 09M BABR		$e^+ e^- \rightarrow D^* + D^- \gamma$	

NODE=M074R15
NODE=M074R15

$\Gamma(D^* \bar{D}^*)/\Gamma(J/\psi \pi^+ \pi^-)$				Γ_{28}/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	CRONIN-HEN..09	CLEO	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<40	90	AUBERT 09M BABR		$e^+ e^- \rightarrow \gamma D^* \bar{D}^*$

NODE=M074R04
NODE=M074R04

$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$				Γ_{29}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT 09M BABR		$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$	

NODE=M074R17
NODE=M074R17

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$				Γ_{30}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* + D^*-$	
not seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^* + D^* - \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	AUBERT 09M BABR		$e^+ e^- \rightarrow D^* + D^* - \gamma$	

NODE=M074R18
NODE=M074R18

$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^{*0} + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.}))/\Gamma_{\text{total}}$				Γ_{32}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	PAKHLOVA 08A	BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+$	

NODE=M074R16
NODE=M074R16

$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$				Γ_{33}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D} \pi$	

NODE=M074R22
NODE=M074R22

$\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma(J/\psi \pi^+ \pi^-)$				Γ_{33}/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R05
NODE=M074R05

$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$				Γ_{34}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$	

NODE=M074R23
NODE=M074R23

$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-)$				Γ_{35}/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R10
NODE=M074R10

$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$				$\Gamma_{35}/\Gamma \times \Gamma_1/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42 × 10 ⁻⁶	90	1 PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R11
NODE=M074R11

1 Using 4263⁺⁸₋₉ MeV for the mass of $\psi(4260)$.

$\Gamma(D^* \bar{D}^* \pi)/\Gamma_{\text{total}}$				Γ_{37}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D}^* \pi$	

NODE=M074R24
NODE=M074R24

$\Gamma(D^* \bar{D}^* \pi)/\Gamma(J/\psi \pi^+ \pi^-)$				Γ_{37}/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R06
NODE=M074R06

$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ
not seen	1 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$	
¹ Results from a measurement of $\sigma(e^+ e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.})$ between $\sqrt{s} = 4.3$ and 4.6 GeV.				

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{38}/Γ
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	PAKHLOVA	11	BELL $e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	

 $\Gamma(D_s^+ D_s^-)/\Gamma(J/\psi \pi^+ \pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{38}/Γ_2
<0.7	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.3	90	CRONIN-HEN..09	CLEO	$e^+ e^-$	

 $\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{39}/Γ
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$	
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	PAKHLOVA	11	BELL $e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$	

 $\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{39}/Γ_2
< 0.8	90	CRONIN-HEN..09	CLEO	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<44	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$	

 $\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	PAKHLOVA	11	BELL $e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$	
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$	

 $\Gamma(D_s^{*+} D_s^{*-})/\Gamma(J/\psi \pi^+ \pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ_2
< 9.5	90	CRONIN-HEN..09	CLEO	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<30	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$	

 $\Gamma(p\bar{p})/\Gamma(J/\psi \pi^+ \pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{41}/Γ_2
<0.13	90	1 AUBERT	06B	BABR $e^+ e^- \rightarrow p\bar{p}\gamma$	

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$.

 $\Gamma(p\bar{p}\pi^0)/\Gamma(J/\psi \pi^+ \pi^-)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{42}/Γ_2
<2 × 10⁻⁴	90	ABLIKIM	17F	BES3 $e^+ e^- \rightarrow \psi(4260) \rightarrow \text{hadrons}$	

Radiative decays $\Gamma(\eta_c(1S)\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{48}/Γ
possibly seen	1 ABLIKIM	17W	$e^+ e^- \rightarrow \gamma \eta_c(1S)$	

¹ Significance ranges from 4.2σ to as low as 1.5σ for a flat component plus $\psi(4260)$ spectrum. Needs confirmation.

NODE=M074R33
NODE=M074R33

NODE=M074R33;LINKAGE=A
NODE=M074R19
NODE=M074R19

NODE=M074R07
NODE=M074R07

NODE=M074R20
NODE=M074R20

NODE=M074R08
NODE=M074R08

NODE=M074R21
NODE=M074R21

NODE=M074R09
NODE=M074R09

NODE=M074R1
NODE=M074R1

NODE=M074R00
NODE=M074R00
OCCUR=2

NODE=M074310

NODE=M074R29
NODE=M074R29

NODE=M074R29;LINKAGE=A

$\Gamma(\chi_{c1}(3872)\gamma)/\Gamma_{\text{total}}$				Γ_{51}/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	20 ± 5	ABLIKIM	14	BES3 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

$\psi(4260)$ REFERENCES

ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60210
ABLIKIM	20C	PR L 124 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60215
ABLIKIM	19	PR D99 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59515
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59856
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ABLIKIM	18N	PR D97 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58914
ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57755
ABLIKIM	17F	PL B771 45	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57909
ABLIKIM	17W	PR D96 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58212
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan		REFID=57991
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang		REFID=58219
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan		REFID=58463
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56782
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55647
SHEN	14	PR D89 072015	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55944
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55409
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
LEES	12AC	PR D86 051102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54750
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDALAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA... 10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532	
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN... 09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114	
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=52296
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
YUAN	08	PR D77 011105	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=52135
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51724
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51959
YUAN	07	PRL 99 182004	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=51960
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
HE	06B	PR D74 091104	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51523
AUBERT,B	05I	PRL 95 142001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50776

NODE=M074

NODE=M074R26
NODE=M074R26

NODE=M233

 $\chi_{c1}(4274)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

was $X(4274)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

NODE=M233

Seen by AAIJ 17C in $B^+ \rightarrow \chi_{c1} K^+$, $\chi_{c1} \rightarrow J/\psi \phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ with a significance (accounting for systematic uncertainties) of 6.0 σ .

 $\chi_{c1}(4274)$ MASS

NODE=M233M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4274 ± 8 OUR AVERAGE				
4273.3 ± 8.3 $^{+17.2}_{-3.6}$	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
4274.4 ± 8.4 $^{+8.4}_{-6.7}$ ± 1.9	22	² AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M233M

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.0 σ .
² From a fit to the invariant mass spectrum with a significance of 3.1 σ .

NODE=M233M;LINKAGE=A
NODE=M233M;LINKAGE=B **$\chi_{c1}(4274)$ WIDTH**

NODE=M233W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
49 ± 12 OUR AVERAGE				
56 ± 11 $^{+8}_{-11}$	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
32.3 ± 21.9 $^{+21.9}_{-15.3}$ ± 7.6	22	² AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M233W

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.0 σ .
² From a fit to the invariant mass spectrum with a significance of 3.1 σ .

NODE=M233W;LINKAGE=A
NODE=M233W;LINKAGE=B **$\chi_{c1}(4274)$ DECAY MODES**

NODE=M233215;NODE=M233

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi \phi$	seen

DESIG=1

 $\chi_{c1}(4274)$ BRANCHING RATIOS

NODE=M233220

$\Gamma(J/\psi \phi)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_1/Γ
seen	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M233R01
NODE=M233R01

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.0 σ .

NODE=M233R01;LINKAGE=A

 $\chi_{c1}(4274)$ REFERENCES

NODE=M233

AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	17	MPL A32 1750139	T. Altonen <i>et al.</i>	(CDF Collab.)

REFID=57657
REFID=57636
REFID=58161

X(4350) $I^G(J^{PC}) = 0^+(?^+)$

OMITTED FROM SUMMARY TABLE

Seen by SHEN 10 in the $\gamma\gamma \rightarrow J/\psi\phi$. Needs confirmation.**X(4350) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4350.6$^{+4.6}_{-5.1}$$\pm 0.7$	8.8 $^{+4.2}_{-3.2}$	¹ SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

¹ Statistical significance of 3.2 σ .**X(4350) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
13$^{+18}_{-9}$$\pm 4$	8.8 $^{+4.2}_{-3.2}$	¹ SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

¹ Statistical significance of 3.2 σ .**X(4350) DECAY MODES**

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 J/\psi\phi$	seen
$\Gamma_2 \gamma\gamma$	seen

X(4350) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\gamma\gamma) \times \Gamma(J/\psi\phi)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7$^{+3.2}_{-2.4}$$\pm 1.1$	8.8 $^{+4.2}_{-3.2}$	¹ SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 $^{+0.7}_{-0.6}\pm 0.3$	8.8 $^{+4.2}_{-3.2}$	² SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$
-----------------------------	----------------------	-------------------	---------	---

¹ For $J^P = 0^+$. Statistical significance of 3.2 σ .² For $J^P = 2^+$. Statistical significance of 3.2 σ .**X(4350) BRANCHING RATIOS**

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	¹ SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	

¹ Statistical significance of 3.2 σ .

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	¹ SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	

¹ Statistical significance of 3.2 σ .**X(4350) REFERENCES**SHEN 10 PRL 104 112004 C.P. Shen *et al.* (BELLE Collab.)

3/29/2020 16:58

Page 679

NODE=M194

NODE=M194M

NODE=M194M

NODE=M194M;LINKAGE=SH

NODE=M194W

NODE=M194W

NODE=M194W;LINKAGE=SH

NODE=M194215;NODE=M194

DESIG=1

DESIG=2

NODE=M194220

NODE=M194G01

NODE=M194G01

OCCUR=2

NODE=M194G01;LINKAGE=S0

NODE=M194G01;LINKAGE=S2

NODE=M194225

NODE=M194R01

NODE=M194R01

NODE=M194R01;LINKAGE=SH

NODE=M194R02

NODE=M194R02

NODE=M194R02;LINKAGE=SH

NODE=M194

REFID=53235

NODE=M181

 $\psi(4360)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $Y(4360)$; was $X(4360)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from $e^+ e^-$ collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by AUBERT 07S, WANG 07D, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

 $\psi(4360)$ MASS

NODE=M181M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4368 ± 13 OUR AVERAGE				Error includes scale factor of 3.7. See the ideogram below.
4320.0 ± 10.4 ± 7.0		1 ABLIKIM	17B BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
4383.8 ± 4.2 ± 0.8		2 ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4347 ± 6 ± 3 279	279	3 WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4340 ± 16 ± 9 37	37	4 LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4383.7 ± 2.9 ± 6.2		5 ZHANG	17B RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4386.4 ± 2.1 ± 6.4		6 ZHANG	17C RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$
4355 ± 9 ± 9 74	74	7 LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4324 ± 24		8 AUBERT	07S BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4361 ± 9 ± 9 47	47	4 WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

1 From a three-resonance fit.

2 From a fit to the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-) \ell^+ \ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} .

3 From a two-resonance fit. Supersedes WANG 07D.

4 From a two-resonance fit.


5 From a three-resonance fit.

6 From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.

7 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

8 From a single-resonance fit. Systematic errors not estimated.

WEIGHTED AVERAGE
4368±13 (Error scaled by 3.7)



NODE=M181

 $\psi(4360)$ WIDTH

NODE=M181W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
96 ± 7 OUR AVERAGE				
101.4 ± 25.3 ± 10.2		1 ABLIKIM	17B BES3	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
101.4 ± 19.7 ± 10.2				

NODE=M181W

$84.2 \pm 12.5 \pm 2.1$	² ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
$103 \pm 9 \pm 5$	³ WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$94 \pm 32 \pm 13$	⁴ LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$94.2 \pm 7.3 \pm 2.0$	⁵ ZHANG	17B RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
$96.0 \pm 6.7 \pm 2.7$	⁶ ZHANG	17C RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi \text{ or } \psi(2S)$
$103^{+17}_{-15} \pm 11$	7 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
172 ± 33	⁸ AUBERT	07S BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$74 \pm 15 \pm 10$	⁴ WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

1 From a three-resonance fit.

2 From a fit to the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-) \ell^+ \ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} .

3 From a two-resonance fit. Supersedes WANG 07D.

4 From a two-resonance fit.

5 From a three-resonance fit.

6 From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.

7 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

8 From a single-resonance fit. Systematic errors not estimated.

NODE=M181W;LINKAGE=B

NODE=M181W;LINKAGE=C

NODE=M181W;LINKAGE=A

NODE=M181W;LINKAGE=WA

NODE=M181W;LINKAGE=E

NODE=M181W;LINKAGE=D

NODE=M181W;LINKAGE=LI

NODE=M181W;LINKAGE=AU

NODE=M181215;NODE=M181

$\psi(4360)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	
$\Gamma_2 J/\psi \pi^+ \pi^-$	
$\Gamma_3 \psi(2S) \pi^+ \pi^-$	seen
$\Gamma_4 \psi_2(3823) \pi^+ \pi^-$	possibly seen
$\Gamma_5 J/\psi \eta$	
$\Gamma_6 D^0 D^{*-} \pi^+$	
$\Gamma_7 D_1(2420) \bar{D} + \text{c.c.}$	possibly seen
$\Gamma_8 \chi_{c1} \gamma$	
$\Gamma_9 \chi_{c2} \gamma$	

$\psi(4360)$ $\Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_3 \Gamma_1/\Gamma$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
7.3 ± 2.8	¹ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
11.0 ± 3.8	² ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
$9.2 \pm 0.6 \pm 0.6$	279	³ WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$10.9 \pm 0.6 \pm 0.7$	279	⁴ WANG	15A BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$6.0 \pm 1.0 \pm 0.5$	37	¹ LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$7.2 \pm 1.0 \pm 0.6$	37	² LEES	14F BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$11.1^{+1.3}_{-1.2}$	74	⁵ LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
12.3 ± 1.2	74	⁶ LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$10.4 \pm 1.7 \pm 1.5$	47	¹ WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
$11.8 \pm 1.8 \pm 1.4$	47	² WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

1 Solution I of two equivalent solutions in a fit using two interfering resonances.

2 Solution II of two equivalent solutions in a fit using two interfering resonances.

3 Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

4 Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

5 Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

6 Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M181230

NODE=M181G1

NODE=M181G1

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M181G1;LINKAGE=WA

NODE=M181G1;LINKAGE=WN

NODE=M181G1;LINKAGE=A

NODE=M181G1;LINKAGE=B

NODE=M181G1;LINKAGE=LI

NODE=M181G1;LINKAGE=LU

NODE=M181G01

NODE=M181G01

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_5 \Gamma_1/\Gamma$			
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				

<6.8 90 WANG 13B BELL $e^+ e^- \rightarrow J/\psi \eta \gamma$

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_8\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M181G02 NODE=M181G02
<0.57	90	1 HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$	
1 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.						
$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_9\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M181G03 NODE=M181G03
<1.9	90	1 HAN	15	BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$	
1 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.						
$\psi(4360)$ BRANCHING RATIOS						
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S) \pi^+ \pi^-)$						Γ_6/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M181R01 NODE=M181R01
<8	90	PAKHLOVA 09	BELL	$e^+e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-} \pi^+$		
$\Gamma(\psi(2S) \pi^+ \pi^-)/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE		DOCUMENT ID	TECN	COMMENT		NODE=M181R00 NODE=M181R00
seen		1 ABLIKIM	17v	BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
1 From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} .						
$\Gamma(\psi(2S) \pi^+ \pi^-)/\Gamma(J/\psi \pi^+ \pi^-)$						Γ_3/Γ_2
VALUE		DOCUMENT ID	TECN	COMMENT		NODE=M181R04 NODE=M181R04
• • • We do not use the following data for averages, fits, limits, etc. • • •						
(0.81 ± 0.12 ± 0.13) to (42 ± 15 ± 15)	1 ZHANG	17C RVUE		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$		
1 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.						
$\Gamma(\psi_2(3823) \pi^+ \pi^-)/\Gamma_{\text{total}}$						Γ_4/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M181R03 NODE=M181R03
possibly seen	19	1 ABLIKIM	15S	BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$	
1 From a fit of $e^+e^- \rightarrow \pi^+\pi^-\psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1}\gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4360)$ line shape.						
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_6/\Gamma \times \Gamma_1/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		NODE=M181R02 NODE=M181R02
<0.72 × 10 ⁻⁶	90	1 PAKHLOVA 09	BELL	$e^+e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-} \pi^+$		
1 Using $4355^{+9}_{-10} \pm 9$ MeV for the mass of $\psi(4360)$.						
$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_7/Γ
VALUE		DOCUMENT ID	TECN	COMMENT		NODE=M181R05 NODE=M181R05
possibly seen		1 ABLIKIM	19AR	BES3	$e^+e^- \rightarrow \pi^+\pi^-\bar{D}D$	
1 Evidence for $e^+e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant.						
$\psi(4360)$ REFERENCES						
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>		(BESIII Collab.)	REFID=59910
ABLIKIM	19K	PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>		(BESIII Collab.)	REFID=59611
ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>		(BESIII Collab.)	REFID=57755
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>		(BESIII Collab.)	REFID=58029
Also		PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>		(BESIII Collab.)	REFID=59611
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang		(BESIII Collab.)	REFID=58219
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan		(BESIII Collab.)	REFID=58463
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>		(BESIII Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>		(BELLE Collab.)	REFID=56816
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>		(BELLE Collab.)	REFID=56839
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>		(BABAR Collab.)	REFID=55938
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>		(BELLE Collab.)	REFID=55377
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>		(BELLE Collab.)	REFID=53143
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		(BABAR Collab.)	REFID=52296
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>		(BABAR Collab.)	REFID=51724
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>		(BELLE Collab.)	REFID=51959

NODE=M236

 $\psi(4390)$

$J^P C = 0^-(1^{--})$
 / needs confirmation.

OMMITTED FROM SUMMARY TABLE
 was $X(4390)$

This state shows properties different from a conventional $q\bar{q}$ state.
 A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

 $\psi(4390)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4391.5^{+6.3}_{-6.8}± 1.0	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$

NODE=M236

NODE=M236M

NODE=M236M

 $\psi(4390)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
139.5^{+16.2}_{-20.6}± 0.6	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$

NODE=M236W

NODE=M236W

 $\psi(4390)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 \quad \pi^+ \pi^- h_c$	seen	
$\Gamma_2 \quad \pi^+ \pi^- \psi(3770)$	possibly seen	

NODE=M236215;NODE=M236

DESIG=1

DESIG=3

NODE=M236220

NODE=M236R01

NODE=M236R01

NODE=M236R02

NODE=M236R02

NODE=M236R02;LINKAGE=A

 $\psi(4390)$ BRANCHING RATIOS

$\Gamma(\pi^+ \pi^- h_c)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$	

NODE=M236220

NODE=M236R01

NODE=M236R01

$\Gamma(\pi^+ \pi^- \psi(3770))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
possibly seen	1 ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$	

NODE=M236R02

NODE=M236R02

¹ Observe $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$ at $\sqrt{s} = 4.26, 4.36$, and 4.42 GeV but cannot establish if continuum or resonant.

NODE=M236

REFID=59910

REFID=57915

ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17G PRL 118 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)

$\psi(4415)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\psi(4415)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4421 ± 4 OUR ESTIMATE			
4415.1± 7.9	¹ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	² MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
4411 ± 7	³ PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	⁴ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4429 ± 9	⁵ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C DASP	$e^+ e^-$
4414 ± 7	SIEGRIST	76 MRK1	$e^+ e^-$
¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.			
² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
³ Systematic uncertainties not estimated.			
⁴ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁵ From a fit to BES (BAI 02C) data.			

 $\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
62 ±20 OUR ESTIMATE			
71.5±19.0	⁶ ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ±32	⁷ MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
77 ±20	⁸ PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ±16	⁹ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
118 ±35	¹⁰ SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
66 ±15	BRANDELIK	78C DASP	$e^+ e^-$
33 ±10	SIEGRIST	76 MRK1	$e^+ e^-$
⁶ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.			
⁷ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁸ Systematic uncertainties not estimated.			
⁹ From a fit to Crystal Ball (OSTERHELD 86) data.			
¹⁰ From a fit to BES (BAI 02C) data.			

 $\psi(4415)$ DECAY MODES

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 D\bar{D}$	seen	
$\Gamma_2 D^0\bar{D}^0$	seen	
$\Gamma_3 D^+D^-$	seen	
$\Gamma_4 D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_5 D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_6 D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_7 D^*\bar{D}^*$	seen	
$\Gamma_8 D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen	
$\Gamma_9 D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen	

NODE=M073M

NODE=M073M

→ UNCHECKED ←

OCCUR=2

NODE=M073M;LINKAGE=AB

NODE=M073M;LINKAGE=MO

NODE=M073M;LINKAGE=NS

NODE=M073M;LINKAGE=ST

NODE=M073M;LINKAGE=SE

NODE=M073W

NODE=M073W

→ UNCHECKED ←

OCCUR=2

NODE=M073W;LINKAGE=AB

NODE=M073W;LINKAGE=MO

NODE=M073W;LINKAGE=NS

NODE=M073W;LINKAGE=ST

NODE=M073W;LINKAGE=SE

NODE=M073215;NODE=M073

NODE=M073

DESIG=7;OUR EVAL;→ UNCHECKED ←

DESIG=8

DESIG=9

DESIG=10;OUR EVAL;→ UNCHECKED ←

DESIG=11

DESIG=12

DESIG=13;OUR EVAL;→ UNCHECKED ←

DESIG=14

DESIG=15

Γ_{10}	$D^0 D^- \pi^+$ (excl. $D^*(2007)^0 \bar{D}^0$ +c.c., $D^*(2010)^+ D^-$ +c.c.)	< 2.3	%	90%	DESIG=4
Γ_{11}	$D \bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+$ +c.c.	(10 ± 4) %			DESIG=5
Γ_{12}	$D^0 D^{*-} \pi^+$ +c.c.	< 11	%	90%	DESIG=6
Γ_{13}	$D_1(2420) \bar{D} +$ c.c.	possibly seen			DESIG=25
Γ_{14}	$D_s^+ D_s^-$	not seen			DESIG=16
Γ_{15}	$\omega \chi_{c2}$	possibly seen			DESIG=20
Γ_{16}	$D_s^{*+} D_s^-$ +c.c.	seen			DESIG=17
Γ_{17}	$D_s^{*+} D_s^{*-}$	not seen			DESIG=18
Γ_{18}	$\psi_2(3823) \pi^+ \pi^-$	possibly seen			DESIG=21
Γ_{19}	$\psi(3770) \pi^+ \pi^-$	possibly seen			DESIG=24
Γ_{20}	$J/\psi \eta$	< 6 $\times 10^{-3}$		90%	DESIG=19
Γ_{21}	$\chi_{c1} \gamma$	< 8 $\times 10^{-4}$		90%	DESIG=22
Γ_{22}	$\chi_{c2} \gamma$	< 4 $\times 10^{-3}$		90%	DESIG=23
Γ_{23}	$e^+ e^-$	(9.4 ± 3.2) $\times 10^{-6}$			DESIG=1

$\psi(4415)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$	Γ_{23}
VALUE (keV)	DOCUMENT ID TECN COMMENT

0.58±0.07 OUR ESTIMATE

0.35±0.12

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 to 0.8	12 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
0.72 ± 0.11	13 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.64 ± 0.23	14 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.49 ± 0.13	BRANDELIK	78C DASP	$e^+ e^-$
0.44 ± 0.14	SIEGRIST	76 MRK1	$e^+ e^-$

11 Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

12 Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different $e^+ e^-$ partial widths. We quote only the range of values.

13 From a fit to Crystal Ball (OSTERHELD 86) data.

14 From a fit to BES (BAI 02C) data.

$\psi(4415) \Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{20} \Gamma_{23}/\Gamma$
VALUE (eV)	DOCUMENT ID TECN COMMENT

<3.6	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$
------	----	------	----------	--

$\Gamma(\chi_{c1} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{21} \Gamma_{23}/\Gamma$
VALUE (eV)	DOCUMENT ID TECN COMMENT

<0.47	90	15 HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c1} \gamma$
-------	----	--------	---------	--

15 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\Gamma(\chi_{c2} \gamma) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{22} \Gamma_{23}/\Gamma$
VALUE (eV)	DOCUMENT ID TECN COMMENT

<2.3	90	16 HAN	15 BELL	$10.58 e^+ e^- \rightarrow \chi_{c2} \gamma$
------	----	--------	---------	--

16 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

$\psi(4415)$ BRANCHING RATIOS

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE	DOCUMENT ID TECN COMMENT

seen	PAKHLOVA	08 BELL	$e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$
------	----------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$
----------	--------	----------	--

NODE=M073220

NODE=M073W1

NODE=M073W1

→ UNCHECKED ←

OCCUR=2

NODE=M073W1;LINKAGE=AB

NODE=M073W1;LINKAGE=MO

NODE=M073W1;LINKAGE=ST

NODE=M073W1;LINKAGE=SE

NODE=M073230

NODE=M073G01

NODE=M073G01

NODE=M073G02

NODE=M073G02

NODE=M073G02;LINKAGE=A

NODE=M073G03

NODE=M073G03

NODE=M073G03;LINKAGE=A

NODE=M073225

NODE=M073R04

NODE=M073R04

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	Γ_3/Γ		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
seen	PAKHLOVA 08	BELL $e^+ e^- \rightarrow D^+ D^- \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M	BABR $e^+ e^- \rightarrow D^+ D^- \gamma$	
$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$	Γ_1/Γ_7	NODE=M073R02 NODE=M073R02	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
0.14±0.12±0.03	AUBERT	09M BABR $e^+ e^- \rightarrow \gamma D^*(*) \bar{D}^(*)$	
$\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_5/Γ	NODE=M073R06 NODE=M073R06	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
seen	AUBERT 09M	BABR $e^+ e^- \rightarrow D^{*0} \bar{D}^0 \gamma$	
$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_6/Γ	NODE=M073R07 NODE=M073R07	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
seen	17 ZHUKOVA 18	BELL $e^+ e^- \rightarrow D^{*+} D^- \gamma$	
seen	AUBERT 09M	BABR $e^+ e^- \rightarrow D^{*+} D^- \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA 07	BELL $e^+ e^- \rightarrow D^{*+} D^- \gamma$	
17 Supersedes PAKHLOVA 07.			
$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$	Γ_4/Γ_7	NODE=M073R03 NODE=M073R03	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
0.17±0.25±0.03	AUBERT 09M	BABR $e^+ e^- \rightarrow \gamma D^*(*) \bar{D}^(*)$	
$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}}$	Γ_8/Γ	NODE=M073R08 NODE=M073R08	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
seen	AUBERT 09M	BABR $e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$	
$\Gamma(D^*(2010)^+ D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}}$	Γ_9/Γ	NODE=M073R09 NODE=M073R09	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
seen	18 ZHUKOVA 18	BELL $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	
seen	AUBERT 09M	BABR $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA 07	BELL $e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$	
18 Supersedes PAKHLOVA 07.			
$\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{11}/Γ	NODE=M073R3 NODE=M073R3	
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
10.5±2.4±3.8	19 PAKHLOVA 08A	BELL $10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$	
19 Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.			
$\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c.}, D^*(2010)^+ D^- + \text{c.c.}) / D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})$	Γ_{10}/Γ_{11}		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
<0.22	90	20 PAKHLOVA 08A	BELL $10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
20 Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.			
$\Gamma(D^0 D^* - \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_{12}/\Gamma \times \Gamma_{23}/\Gamma$		
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
<0.99 × 10⁻⁶	90	21 PAKHLOVA 09	BELL $e^+ e^- \rightarrow D^0 D^{*-} \pi^+$
21 Using 4421 ± 4 MeV for the mass of $\psi(4415)$.			
$\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{13}/Γ	NODE=M073R15 NODE=M073R15	
<u>VALUE</u>		<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
possibly seen	22 ABLIKIM 19AR	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$	
22 Evidence for $e^+ e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant.			
$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$	Γ_{14}/Γ	NODE=M073R10 NODE=M073R10	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>	
not seen	PAKHLOVA 11	BELL $e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	
not seen	DEL-AMO-SA..10N	BABR $e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	

$\Gamma(\omega\chi_{c2})/\Gamma_{\text{total}}$				Γ_{15}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
possibly seen	ABLIKIM	16A	BES3	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$	NODE=M073R00 NODE=M073R00
$\Gamma(D_s^{*+} D_s^- + c.c.)/\Gamma_{\text{total}}$				Γ_{16}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
seen	PAKHLOVA	11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$	NODE=M073R11 NODE=M073R11
seen	DEL-AMO-SA..10N	BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$	
$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$				Γ_{17}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
not seen	PAKHLOVA	11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$	NODE=M073R12 NODE=M073R12
not seen	DEL-AMO-SA..10N	BABR		$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$	
$\Gamma(\psi(3770)\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{19}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
possibly seen	23	ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$	NODE=M073R14 NODE=M073R14
23 Observe $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$ at $\sqrt{s} = 4.26, 4.36$, and 4.42 GeV but cannot establish if continuum or resonant.					NODE=M073R14;LINKAGE=A
$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$				Γ_{18}/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
possibly seen	19	24	ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
24 From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1} \gamma$ cross sections taken at \sqrt{s} values of $4.23, 4.26, 4.36, 4.42$, and 4.60 GeV to the $\psi(4415)$ line shape.					NODE=M073R13;LINKAGE=A

$\psi(4415)$ REFERENCES

ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
ABLIKIM	16A	PR D93 011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57122
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
DEL-AMO-SA...10N	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)	REFID=22243

$Z_c(4430)$

$I^G(J^P)$ = $1^+(1^{+-})$
 G, C need confirmation.

was $X(4430)^\pm$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

First seen by CHOI 08 in $B \rightarrow K\pi^+\psi(2S)$ decays, confirmed by AAIJ 14AG, and confirmed in a model-independent way by AAIJ 15BH. Also seen by CHILIKIN 14 in $B \rightarrow K^+\pi^+J/\psi$ decays.

J^P was determined by CHILIKIN 13 and AAIJ 14AG.

$Z_c(4430)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4478⁺¹⁵₋₁₈ OUR AVERAGE			
4475 ± 7 ⁺¹⁵ ₋₂₅	¹ AAIJ	14AG LHCb	$B^0 \rightarrow K^+\pi^-\psi(2S)$
4485 ± 22 ⁺²⁸ ₋₁₁	¹ CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4443 ⁺¹⁵⁺¹⁹ ₋₁₂₋₁₃	² MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
4433 ± 4 ± 2	³ CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
1 From a four-dimensional amplitude analysis. 2 From a Dalitz plot analysis. Superseded by CHILIKIN 13. 3 Superseded by MIZUK 09 and CHILIKIN 13.			

NODE=M195

$Z_c(4430)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
181^{±31} OUR AVERAGE			
172 ± 13 ⁺³⁷ ₋₃₄	¹ AAIJ	14AG LHCb	$B^0 \rightarrow K^+\pi^-\psi(2S)$
200 ⁺⁴¹⁺²⁶ ₋₄₆₋₃₅	¹ CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
107 ⁺⁸⁶⁺⁷⁴ ₋₄₃₋₅₆	² MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
45 ⁺¹⁸⁺³⁰ ₋₁₃₋₁₃	³ CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
1 From a four-dimensional amplitude analysis. 2 From a Dalitz plot analysis. Superseded by CHILIKIN 13. 3 Superseded by MIZUK 09 and CHILIKIN 13.			

NODE=M195M;LINKAGE=A
NODE=M195M;LINKAGE=MI
NODE=M195M;LINKAGE=CH

$Z_c(4430)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+\psi(2S)$	seen
$\Gamma_2 \pi^+J/\psi$	seen

NODE=M195W

NODE=M195W

$Z_c(4430)$ BRANCHING RATIOS

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	¹ AAIJ	14AG LHCb	$B^0 \rightarrow K^+\pi^-\psi(2S)$	
seen	² CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	³ AUBERT	09AA BABR	$B \rightarrow K\pi^+\psi(2S)$	
seen	⁴ MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$	
1 From a four-dimensional amplitude analysis. No product of branching fractions quoted. 2 From a four-dimensional amplitude analysis. Measured a product of branching fractions $B(B^0 \rightarrow Z_c(4430)^-K^+) \times B(Z_c(4430)^-\rightarrow \psi(2S)\pi^-) = (6.0^{+1.7+2.5}_{-2.0-1.4}) \times 10^{-5}$. 3 AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+\psi(2S)) < 4.7 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+\psi(2S)) < 3.1 \times 10^{-5}$ at 95% CL. 4 Measured a product of branching fractions $B(\bar{B}^0 \rightarrow K^- Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+\psi(2S)) = (3.2^{+1.8+5.3}_{-0.9-1.6}) \times 10^{-5}$. Superseded by CHILIKIN 13.				

NODE=M195215;NODE=M195

DESIG=1

DESIG=2

NODE=M195225

NODE=M195R01

NODE=M195R01

NODE=M195R01;LINKAGE=AA
NODE=M195R01;LINKAGE=A

NODE=M195R01;LINKAGE=AU

NODE=M195R01;LINKAGE=MI

$\Gamma(\pi^+ J/\psi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	1,2 CHILIKIN	14 BELL	$\bar{B}^0 \rightarrow K^- \pi^+ J/\psi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	3 AUBERT	09AA BABR	$B \rightarrow K\pi^+ J/\psi$	
1 CHILIKIN 14 reports $B(\bar{B}^0 \rightarrow Z_c(4430)^+ K^-) \times B(Z_c(4430)^+ \rightarrow J/\psi\pi^+) = (5.4^{+4.0}_{-1.0} \pm 1.1) \times 10^{-6}$.				NODE=M195R02;LINKAGE=A
2 A broad enhancement seen by AAJ 19R in the decays $B^0 \rightarrow J/\psi\pi^+ K^-$ at 4600 MeV can be due to an interplay of $Z_c(4430)$, $Z_c(4200)$ and the fitting polynomials.				NODE=M195R02;LINKAGE=E
3 AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+ J/\psi) < 1.5 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+ J/\psi) < 0.4 \times 10^{-5}$ at 95% CL.				NODE=M195R02;LINKAGE=AU

 $Z_c(4430)$ REFERENCES

AAIJ	19R PRL 122 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BH PR D92 112009	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AG PR D91 122002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
CHILIKIN	14 PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)
CHILIKIN	13 PR D88 074026	K. Chilikin <i>et al.</i>	(BELLE Collab.) JP
AUBERT	09AA PR D79 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
MIZUK	09 PR D80 031104	R. Mizuk <i>et al.</i>	(BELLE Collab.)
CHOI	08 PRL 100 142001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)

 $\chi_{c0}(4500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMMITTED FROM SUMMARY TABLE
was $X(4500)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen by AAJ 17C in $B^+ \rightarrow \chi_{c0} K^+$, $\chi_{c0} \rightarrow J/\psi\phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ with a significance (accounting for systematic uncertainties) of 6.1σ .

 $\chi_{c0}(4500)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
4506±11⁺¹²₋₁₅	4289	¹ AAJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$	

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 6.1σ .

 $\chi_{c0}(4500)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
92±21⁺²¹₋₂₀	4289	¹ AAJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$	

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 6.1σ .

 $\chi_{c0}(4500)$ DECAY MODES

Mode	Γ_1	$J/\psi\phi$	Γ_1/Γ

 $\chi_{c0}(4500)$ BRANCHING RATIOS

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	4289	¹ AAJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 6.1σ .					

 $\chi_{c0}(4500)$ REFERENCES

AAIJ Also	17C PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
	PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M195R02
NODE=M195R02

NODE=M195R02;LINKAGE=A

NODE=M195R02;LINKAGE=E

NODE=M195R02;LINKAGE=AU

NODE=M195

REFID=59776
REFID=57110
REFID=55896
REFID=56344
REFID=55551
REFID=52940
REFID=52960
REFID=52178

NODE=M234

NODE=M234

NODE=M234M

NODE=M234M

NODE=M234M;LINKAGE=A

NODE=M234W

NODE=M234W

NODE=M234W;LINKAGE=A

NODE=M234215;NODE=M234

DESIG=1

NODE=M234220

NODE=M234R01
NODE=M234R01

NODE=M234R01;LINKAGE=A

NODE=M234

REFID=57657
REFID=57636

NODE=M189

$\psi(4660)$

$J^P C = 0^-(1^{--})$
/ needs confirmation.

also known as $Y(4660)$; was $X(4660)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from $e^+ e^-$ collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

$\psi(4660)$ MASS

NODE=M189M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4633 \pm 7 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below. [4643 \pm 9 MeV OUR 2019 AVERAGE Scale factor = 1.2]

4625.9 \pm 6.2 6.0	\pm 0.4	89	1 JIA	19A BELL $e^+ e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
4652 \pm 10	\pm 11	279	2 WANG	15A BELL $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4669 \pm 21	\pm 3	37	3 LEES	14F BABR $10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4634 \pm 8 7	\pm 5 8	142	4 PAKHLOVA	08B BELL $e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4652.5 \pm 3.4 1.1	\pm 1.1	5 DAI	17 RVUE	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
4645.2 \pm 9.5 6.0	\pm 6.0	6 ZHANG	17B RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4646.4 \pm 9.7 4.8	\pm 4.8	7 ZHANG	17C RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$
4661 \pm 9 8	\pm 6	8 LIU	08H RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
4664 \pm 11 5	\pm 5	44 WANG	07D BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

1 From a fit of a Breit-Wigner convolved with a Gaussian.

2 From a two-resonance fit. Supersedes WANG 07D.

3 From a two-resonance fit.

4 The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

5 The pole parameters are extracted from the speed plot.

6 From a three-resonance fit.

7 From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.

8 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189M

NODE=M189M

NEW

NODE=M189M;LINKAGE=E

NODE=M189M;LINKAGE=A

NODE=M189M;LINKAGE=LE


NODE=M189M;LINKAGE=PA

NODE=M189M;LINKAGE=C

NODE=M189M;LINKAGE=D

NODE=M189M;LINKAGE=B

NODE=M189M;LINKAGE=LI



$\psi(4660)$ WIDTH

NODE=M189W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
64 ± 9 OUR AVERAGE				
[72 ± 11 MeV OUR 2019 AVERAGE]				
49.8 ^{+13.9} _{-11.5} ± 4.0	89	1 JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
68 ± 11 ± 5	279	2 WANG	15A BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
104 ± 48 ± 10	37	3 LEES	14F BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
92 ± 40 ± 10	142	4 PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
62.6 ± 5.6 ± 4.3	5 DAI	17 RVUE		$e^+e^- \rightarrow \Lambda_c^+\Lambda_c^-$
113.8 ± 18.1 ± 3.4	6 ZHANG	17B RVUE		$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
103.5 ± 15.6 ± 4.0	7 ZHANG	17C RVUE		$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$
42 ± 17 ± 6	44	8 LIU	08H RVUE	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
48 ± 15 ± 3	44	WANG	07D BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
1 From a fit of a Breit-Wigner convolved with a Gaussian.				
2 From a two-resonance fit. Supersedes WANG 07D.				
3 From a two-resonance fit.				
4 The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.				
5 The pole parameters are extracted from the speed plot.				
6 From a three-resonance fit.				
7 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.				
8 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.				

NODE=M189W

NEW

ψ(4660) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ ₁ e^+e^-	
Γ ₂ $\psi(2S)\pi^+\pi^-$	seen
Γ ₃ $J/\psi\eta$	
Γ ₄ $D^0D^{*-}\pi^+$	
Γ ₅ $\chi_{c1}\gamma$	
Γ ₆ $\chi_{c2}\gamma$	
Γ ₇ $\Lambda_c^+\Lambda_c^-$	
Γ ₈ $D_s^+D_{s1}(2536)^-$	

ψ(4660) Γ(i) × Γ(e⁺e⁻)/Γ(total)

Γ(ψ(2S)π ⁺ π ⁻) × Γ(e ⁺ e ⁻)/Γ _{total}	Γ ₂ Γ ₁ /Γ		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.0 ± 0.3 ± 0.2			
279	1 WANG	15A BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
8.1 ± 1.1 ± 1.0	2 WANG	15A BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
2.7 ± 1.3 ± 0.5	3 LEES	14F BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
7.5 ± 1.7 ± 0.7	4 LEES	14F BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
2.2 ^{+0.7} _{-0.6}	44	5 LIU	08H RVUE $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
5.9 ± 1.6	44	6 LIU	08H RVUE $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
3.0 ± 0.9 ± 0.3	44	3 WANG	07D BELL $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
7.6 ± 1.8 ± 0.8	44	4 WANG	07D BELL $10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=4

DESIG=3

DESIG=6

DESIG=7

DESIG=5

DESIG=8

NODE=M189230

NODE=M189G1

NODE=M189G1

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M189G1;LINKAGE=A

NODE=M189G1;LINKAGE=B

NODE=M189G1;LINKAGE=WA

NODE=M189G1;LINKAGE=WN

NODE=M189G1;LINKAGE=LI

NODE=M189G1;LINKAGE=LU

Γ(J/ψη) × Γ(e ⁺ e ⁻)/Γ _{total}	Γ ₃ Γ ₁ /Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<0.94	

NODE=M189G01

NODE=M189G01

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.94	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_5\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		
<0.45	90	1 HAN	15	BELL	10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$	NODE=M189G02 NODE=M189G02
¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.						
$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_6\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT		
<2.1	90	1 HAN	15	BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$	NODE=M189G03 NODE=M189G03
¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.						
$\psi(4660)$ BRANCHING RATIOS						
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S) \pi^+ \pi^-)$						Γ_4/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<10	90	PAKHLOVA 09	BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$		NODE=M189R01 NODE=M189R01
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_4/\Gamma \times \Gamma_1/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.37 \times 10^{-6}	90	1 PAKHLOVA 09	BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$		NODE=M189R02 NODE=M189R02
¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$.						
$\Gamma(\Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_7/\Gamma \times \Gamma_1/\Gamma$
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.68^{+0.16}_{-0.15}^{+0.29}_{-0.30}	142	1 PAKHLOVA 08B	BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$		NODE=M189R03 NODE=M189R03
¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.						
$\Gamma(D_s^+ D_{s1}(2536)^-)/\Gamma(e^+e^-)/\Gamma_{\text{total}}$						$\Gamma_8\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
14.3^{+2.8}_{-2.6}^{+1.5}	89	1 JIA	19A	BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$	NODE=M189R00 NODE=M189R00
¹ Using $D_{s1}(2536)^- \rightarrow \bar{D}^{*0} K^-$.						

$\psi(4660)$ REFERENCES

JIA	19A	PR D100 111103	S. Jia <i>et al.</i>	(BELLE Collab.)	NODE=M189
DAI	17	PR D96 116001	L.-Y. Dai, J. Haidenbauer, U.-G. Meissner	(JULI+)	REFID=60037
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang		REFID=58704
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan		REFID=58219
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=58463
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=56816
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56839
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55938
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=55377
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=53143
PAKHLOVA	08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52296
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52596
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51724
					REFID=51959

NODE=M235

 $\chi_{c0}(4700)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMMITTED FROM SUMMARY TABLE
was $X(4700)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

NODE=M235

Seen by AAIJ 17C in $B^+ \rightarrow \chi_{c0} K^+$, $\chi_{c0} \rightarrow J/\psi \phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ with a significance (accounting for systematic uncertainties) of 5.6σ .

 $\chi_{c0}(4700)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4704 \pm 10^{+14}_{-24}$	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.6σ .

NODE=M235M

NODE=M235M

NODE=M235M;LINKAGE=A

 $\chi_{c0}(4700)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$120 \pm 31^{+42}_{-33}$	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.6σ .

NODE=M235W

NODE=M235W

NODE=M235W;LINKAGE=A

 $\chi_{c0}(4700)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad J/\psi \phi$	seen

NODE=M235215;NODE=M235

DESIG=1

 $\chi_{c0}(4700)$ BRANCHING RATIOS

$\Gamma(J/\psi \phi)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	Γ_1/Γ
seen	4289	¹ AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.6σ .

NODE=M235220

NODE=M235R01
NODE=M235R01

NODE=M235R01;LINKAGE=A

NODE=M235

REFID=57657
REFID=57636

$b\bar{b}$ MESONS **(including possibly non- $q\bar{q}$ states)**

NODE=MXXX030

See the related review(s):

[Bottomonium System](#)

[Width Determination of the \$\Upsilon\$ States](#)

$\eta_b(1S)$

$$I^G(J^P C) = 0^+(0^- +)$$

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.

NODE=M171

NODE=M171

NODE=M171M


NODE=M171M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9398.7 ± 2.0 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
9394.8 ± 2.7 ± 4.5	29K	FULSOM	18	BELL $\Upsilon(2S) \rightarrow \gamma X$
9400.7 ± 1.7 ± 1.6	33.1k	TAMPONI	15	BELL $e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
9402.4 ± 1.5 ± 1.8	34k	¹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$
9391.8 ± 6.6 ± 2.0	2.3k	² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
9394.2 ± 4.8 ± 2.0	13k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
9388.9 ± 3.1 ± 2.7	19k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9393.2 ± 3.4 ± 2.3	10	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$
9300 ± 20 ± 20		HEISTER	02D ALEP	181–209 $e^+ e^-$

¹With floating width. Not independent of the corresponding mass difference measurement.

²Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.

³Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.



NODE=M171M;LINKAGE=MI

NODE=M171M;LINKAGE=AU

NODE=M171M;LINKAGE=DO


$m_{\Upsilon(1S)} - m_{\eta_b}$

NODE=M171M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3 ± 3.2 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
57.9 ± 1.5 ± 1.8	34k	¹ MIZUK	12	BELL $e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$
68.5 ± 6.6 ± 2.0	2.3 ± 0.5k	² BONVICINI	10	CLEO $\Upsilon(3S) \rightarrow \gamma X$
66.1 ± 4.8 ± 2.0	13 ± 5k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
71.4 ± 2.3 ± 2.7	19 ± 3k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1 ± 3.4 ± 2.3	10^{+5}_{-4}	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

- ¹ With floating width. Not independent of the corresponding mass measurement.
² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.
³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M2;LINKAGE=MI
 NODE=M171M2;LINKAGE=AU
 NODE=M171M2;LINKAGE=DO



γ ENERGY IN $\gamma(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
920.6^{+2.8}_{-3.2} OUR AVERAGE				
$918.6 \pm 6.0 \pm 1.9$	$2.3 \pm 0.5k$	1 BONVICINI	10 CLEO	$\gamma(3S) \rightarrow \gamma X$
$921.2 \pm 2.1 \pm 2.4$	$19 \pm 3k$	1 AUBERT	08V BABR	$\gamma(3S) \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171DM

NODE=M171DM

NODE=M171DM;LINKAGE=BO

γ ENERGY IN $\gamma(2S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
609.3^{+4.6}_{-4.5} ± 1.9				
	$13 \pm 5k$	1 AUBERT	09AQ BABR	$\gamma(2S) \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171U2S

NODE=M171U2S

NODE=M171U2S;LINKAGE=AU

$\eta_b(1S)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10⁺⁵₋₄ OUR AVERAGE				
$8 \pm 5 \pm 5$	$33.1k$	1 TAMPONI	15 BELL	$e^+ e^- \rightarrow \gamma \eta + \text{hadrons}$
$10.8 \pm 4.0 \pm 4.5$	$34k$	1 MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$

¹ With floating mass.

NODE=M171W

NODE=M171W

NODE=M171W;LINKAGE=MI

$\eta_b(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+ 3h^-$	not seen	
Γ_3 $2h^+ 2h^-$	not seen	
Γ_4 $4h^+ 4h^-$	not seen	
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+ \mu^-$	$<9 \times 10^{-3}$	90%
Γ_7 $\tau^+ \tau^-$	$<8 \%$	90%

 $\eta_b(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ **$\Gamma(3h^+ 3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<470	95	ABDALLAH	06	DLPH	161–209 $e^+ e^-$
<132	95	HEISTER	02D	ALEP	181–209 $e^+ e^-$

 $\Gamma(2h^+ 2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<190	95	ABDALLAH	06	DLPH	161–209 $e^+ e^-$
< 48	95	HEISTER	02D	ALEP	181–209 $e^+ e^-$

 $\Gamma(4h^+ 4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_4\Gamma_5/\Gamma$
------------	-----	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<660	95	ABDALLAH	06	DLPH	161–209 $e^+ e^-$
------	----	----------	----	------	-------------------

 $\eta_b(1S)$ BRANCHING RATIOS **$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
-------	------	-------------	------	---------	-------------------

seen	34k	MIZUK	12	BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
------	-----	-------	----	------	---

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
-------	-----	-------------	------	---------	-------------------

$<9 \times 10^{-3}$	90	1 AUBERT	09Z	BABR	$e^+ e^- \rightarrow \gamma(2S, 3S) \rightarrow \gamma\eta_b$
---------------------	----	----------	-----	------	---

1 Obtained using $B(\gamma(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$ and $B(\gamma(3S) \rightarrow \gamma\eta_b) = (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$. This limit is equivalent to $B(\eta_b \rightarrow \mu^+ \mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.

 $\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
-------	-----	-------------	------	---------	-------------------

$<8 \times 10^{-2}$	90	AUBERT	09P	BABR	$e^+ e^- \rightarrow \gamma\tau^+\tau^-$
---------------------	----	--------	-----	------	--

 $\eta_b(1S)$ REFERENCES

FULSAM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRC 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)

NODE=M171225;NODE=M171

DESIG=7
DESIG=1;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=2;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=4;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=3;OUR EST; \rightarrow UNCHECKED \leftarrow
DESIG=5
DESIG=6

NODE=M171230

NODE=M171G1
NODE=M171G1

NODE=M171G2
NODE=M171G2

NODE=M171G3
NODE=M171G3

NODE=M171235

NODE=M171R03
NODE=M171R03

NODE=M171R01
NODE=M171R01

NODE=M171R01;LINKAGE=AU

NODE=M171R02
NODE=M171R02

NODE=M171

REFID=59535
REFID=56996
REFID=54288
REFID=54718
REFID=53231
REFID=53106
REFID=53062
REFID=52930
REFID=52262
REFID=51042
REFID=48577

NODE=M049

 $\gamma(1S)$ $I^G(J^{PC}) = 0^-(1^-^-)$ **$\gamma(1S)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9460.30±0.26 OUR AVERAGE			Error includes scale factor of 3.3.
9460.51±0.09±0.05	1 ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
9459.97±0.11±0.07	MACKAY 84	REDE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9460.60±0.09±0.05	2,3 BARU	92B	$e^+ e^- \rightarrow$ hadrons
9460.59±0.12	BARU	86	$e^+ e^- \rightarrow$ hadrons
9460.6 ±0.4	3,4 ARTAMONOV 84	REDE	$e^+ e^- \rightarrow$ hadrons
1 Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).			NODE=M049M;LINKAGE=AR
2 Superseding BARU 86.			NODE=M049M;LINKAGE=A
3 Superseded by ARTAMONOV 00.			NODE=M049M;LINKAGE=RZ
4 Value includes data of ARTAMONOV 82.			NODE=M049M;LINKAGE=G

 $\gamma(1S)$ WIDTH

VALUE (keV)	DOCUMENT ID
54.02±1.25 OUR EVALUATION	See the Note on "Width Determinations of the γ States"

 $\gamma(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \tau^+ \tau^-$	(2.60 ± 0.10) %	
$\Gamma_2 e^+ e^-$	(2.38 ± 0.11) %	
$\Gamma_3 \mu^+ \mu^-$	(2.48 ± 0.05) %	
Hadronic decays		
$\Gamma_4 g g g$	(81.7 ± 0.7) %	
$\Gamma_5 \gamma g g$	(2.2 ± 0.6) %	
$\Gamma_6 \eta'(958)$ anything	(2.94 ± 0.24) %	
$\Gamma_7 J/\psi(1S)$ anything	(5.4 ± 0.4) × 10 ⁻⁴	S=1.4
$\Gamma_8 J/\psi(1S) \eta_c$	< 2.2 × 10 ⁻⁶	CL=90%
$\Gamma_9 J/\psi(1S) \chi_{c0}$	< 3.4 × 10 ⁻⁶	CL=90%
$\Gamma_{10} J/\psi(1S) \chi_{c1}$	(3.9 ± 1.2) × 10 ⁻⁶	
$\Gamma_{11} J/\psi(1S) \chi_{c2}$	< 1.4 × 10 ⁻⁶	CL=90%
$\Gamma_{12} J/\psi(1S) \eta_c(2S)$	< 2.2 × 10 ⁻⁶	CL=90%
$\Gamma_{13} J/\psi(1S) X(3940)$	< 5.4 × 10 ⁻⁶	CL=90%
$\Gamma_{14} J/\psi(1S) X(4160)$	< 5.4 × 10 ⁻⁶	CL=90%
$\Gamma_{15} X(4350)$ anything, $X \rightarrow J/\psi(1S) \phi$	< 8.1 × 10 ⁻⁶	CL=90%
$\Gamma_{16} Z_c(3900)^\pm$ anything, $Z_c \rightarrow J/\psi(1S) \pi^\pm$	< 1.3 × 10 ⁻⁵	CL=90%
$\Gamma_{17} Z_c(4200)^\pm$ anything, $Z_c \rightarrow J/\psi(1S) \pi^\pm$	< 6.0 × 10 ⁻⁵	CL=90%
$\Gamma_{18} Z_c(4430)^\pm$ anything, $Z_c \rightarrow J/\psi(1S) \pi^\pm$	< 4.9 × 10 ⁻⁵	CL=90%
$\Gamma_{19} X_{cs}^\pm$ anything, $X \rightarrow J/\psi K^\pm$	< 5.7 × 10 ⁻⁶	CL=90%
$\Gamma_{20} \chi_{c1}(3872)$ anything, $\chi_{c1} \rightarrow J/\psi(1S) \pi^+ \pi^-$	< 9.5 × 10 ⁻⁶	CL=90%
$\Gamma_{21} \psi(4260)$ anything, $\psi \rightarrow J/\psi(1S) \pi^+ \pi^-$	< 3.8 × 10 ⁻⁵	CL=90%
$\Gamma_{22} \psi(4260)$ anything, $\psi \rightarrow J/\psi(1S) K^+ K^-$	< 7.5 × 10 ⁻⁶	CL=90%
$\Gamma_{23} \chi_{c1}(4140)$ anything, $\chi_{c1} \rightarrow J/\psi(1S) \phi$	< 5.2 × 10 ⁻⁶	CL=90%
$\Gamma_{24} \chi_{c0}$ anything	< 4 × 10 ⁻³	CL=90%

NODE=M049M

NODE=M049M

NODE=M049M;LINKAGE=AR

NODE=M049M;LINKAGE=A

NODE=M049M;LINKAGE=RZ

NODE=M049M;LINKAGE=G

NODE=M049W

NODE=M049W

→ UNCHECKED ←

NODE=M049215;NODE=M049

DESIG=3

DESIG=2

DESIG=1

NODE=M049;CLUMP=A

DESIG=117

DESIG=118

DESIG=73

DESIG=12

DESIG=146

DESIG=147

DESIG=148

DESIG=149

DESIG=150

DESIG=151

DESIG=152

DESIG=167

DESIG=168

DESIG=169

DESIG=170

DESIG=173

DESIG=160

DESIG=161

DESIG=165

DESIG=166

DESIG=5

Γ_{25}	χ_{c1} anything	$(1.90 \pm 0.35) \times 10^{-4}$	DESIG=6
Γ_{26}	$\chi_{c1}(1P)X_{tetra}$	$< 3.78 \times 10^{-5}$	CL=90% DESIG=175
Γ_{27}	χ_{c2} anything	$(2.8 \pm 0.8) \times 10^{-4}$	DESIG=7
Γ_{28}	$\psi(2S)$ anything	$(1.23 \pm 0.20) \times 10^{-4}$	DESIG=8
Γ_{29}	$\psi(2S)\eta_c$	$< 3.6 \times 10^{-6}$	CL=90% DESIG=153
Γ_{30}	$\psi(2S)\chi_{c0}$	$< 6.5 \times 10^{-6}$	CL=90% DESIG=154
Γ_{31}	$\psi(2S)\chi_{c1}$	$< 4.5 \times 10^{-6}$	CL=90% DESIG=155
Γ_{32}	$\psi(2S)\chi_{c2}$	$< 2.1 \times 10^{-6}$	CL=90% DESIG=156
Γ_{33}	$\psi(2S)\eta_c(2S)$	$< 3.2 \times 10^{-6}$	CL=90% DESIG=157
Γ_{34}	$\psi(2S)X(3940)$	$< 2.9 \times 10^{-6}$	CL=90% DESIG=158
Γ_{35}	$\psi(2S)X(4160)$	$< 2.9 \times 10^{-6}$	CL=90% DESIG=159
Γ_{36}	$\psi(4260)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	$< 7.9 \times 10^{-5}$	CL=90% DESIG=162
Γ_{37}	$\psi(4360)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	$< 5.2 \times 10^{-5}$	CL=90% DESIG=163
Γ_{38}	$\psi(4660)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	$< 2.2 \times 10^{-5}$	CL=90% DESIG=164
Γ_{39}	$X(4050)^{\pm}$ anything, $X \rightarrow \psi(2S)\pi^{\pm}$	$< 8.8 \times 10^{-5}$	CL=90% DESIG=171
Γ_{40}	$Z_c(4430)^{\pm}$ anything, $Z_c \rightarrow \psi(2S)\pi^{\pm}$	$< 6.7 \times 10^{-5}$	CL=90% DESIG=172
Γ_{41}	$Z_c(4200)^+Z_c(4200)^-$	$< 2.23 \times 10^{-5}$	CL=90% DESIG=178
Γ_{42}	$Z_c(3900)^{\pm}Z_c(4200)^{\mp}$	$< 8.1 \times 10^{-6}$	CL=90% DESIG=179
Γ_{43}	$Z_c(3900)^+Z_c(3900)^-$	$< 1.8 \times 10^{-6}$	CL=90% DESIG=180
Γ_{44}	$X(4050)^+X(4050)^-$	$< 1.58 \times 10^{-5}$	CL=90% DESIG=181
Γ_{45}	$X(4250)^+X(4250)^-$	$< 2.66 \times 10^{-5}$	CL=90% DESIG=182
Γ_{46}	$X(4050)^{\pm}X(4250)^{\mp}$	$< 4.42 \times 10^{-5}$	CL=90% DESIG=183
Γ_{47}	$Z_c(4430)^+Z_c(4430)^-$	$< 2.03 \times 10^{-5}$	CL=90% DESIG=184
Γ_{48}	$X(4055)^{\pm}X(4055)^{\mp}$	$< 2.33 \times 10^{-5}$	CL=90% DESIG=186
Γ_{49}	$X(4055)^{\pm}Z_c(4430)^{\mp}$	$< 4.55 \times 10^{-5}$	CL=90% DESIG=189
Γ_{50}	$\rho\pi$	$< 3.68 \times 10^{-6}$	CL=90% DESIG=11
Γ_{51}	$\omega\pi^0$	$< 3.90 \times 10^{-6}$	CL=90% DESIG=131
Γ_{52}	$\pi^+\pi^-$	$< 5 \times 10^{-4}$	CL=90% DESIG=23
Γ_{53}	K^+K^-	$< 5 \times 10^{-4}$	CL=90% DESIG=24
Γ_{54}	$p\bar{p}$	$< 5 \times 10^{-4}$	CL=90% DESIG=25
Γ_{55}	$\pi^+\pi^-\pi^0$	$(2.1 \pm 0.8) \times 10^{-6}$	DESIG=72
Γ_{56}	ϕK^+K^-	$(2.4 \pm 0.5) \times 10^{-6}$	DESIG=136
Γ_{57}	$\omega\pi^+\pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$	DESIG=137
Γ_{58}	$K^*(892)^0K^-\pi^+ + c.c.$	$(4.4 \pm 0.8) \times 10^{-6}$	DESIG=138
Γ_{59}	$\phi f_2'(1525)$	$< 1.63 \times 10^{-6}$	CL=90% DESIG=139
Γ_{60}	$\omega f_2(1270)$	$< 1.79 \times 10^{-6}$	CL=90% DESIG=140
Γ_{61}	$\rho(770)a_2(1320)$	$< 2.24 \times 10^{-6}$	CL=90% DESIG=141
Γ_{62}	$K^*(892)^0\bar{K}_2^*(1430)^0 + c.c.$	$(3.0 \pm 0.8) \times 10^{-6}$	DESIG=142
Γ_{63}	$K_1(1270)^{\pm}K^{\mp}$	$< 2.41 \times 10^{-6}$	CL=90% DESIG=143
Γ_{64}	$K_1(1400)^{\pm}K^{\mp}$	$(1.0 \pm 0.4) \times 10^{-6}$	CL=90% DESIG=144
Γ_{65}	$b_1(1235)^{\pm}\pi^{\mp}$	$< 1.25 \times 10^{-6}$	CL=90% DESIG=145
Γ_{66}	$\pi^+\pi^-\pi^0\pi^0$	$(1.28 \pm 0.30) \times 10^{-5}$	DESIG=132
Γ_{67}	$K_S^0K^+\pi^- + c.c.$	$(1.6 \pm 0.4) \times 10^{-6}$	DESIG=133
Γ_{68}	$K^*(892)^0\bar{K}^0 + c.c.$	$(2.9 \pm 0.9) \times 10^{-6}$	DESIG=134
Γ_{69}	$K^*(892)^-K^+ + c.c.$	$< 1.11 \times 10^{-6}$	CL=90% DESIG=135
Γ_{70}	$f_1(1285)$ anything	$(4.6 \pm 3.1) \times 10^{-3}$	DESIG=174
Γ_{71}	$D^*(2010)^{\pm}$ anything	$(2.52 \pm 0.20) \%$	DESIG=30
Γ_{72}	$f_1(1285)X_{tetra}$	$< 6.24 \times 10^{-5}$	CL=90% DESIG=176
Γ_{73}	$\bar{2}H$ anything	$(2.85 \pm 0.25) \times 10^{-5}$	DESIG=107
Γ_{74}	Sum of 100 exclusive modes	$(1.200 \pm 0.017) \%$	DESIG=128

Radiative decays

Γ_{75}	$\gamma\pi^+\pi^-$	$(6.3 \pm 1.8) \times 10^{-5}$	NODE=M049;CLUMP=B
Γ_{76}	$\gamma\pi^0\pi^0$	$(1.7 \pm 0.7) \times 10^{-5}$	DESIG=70
Γ_{77}	$\gamma\pi\pi$ (S-wave)	$(4.6 \pm 0.7) \times 10^{-5}$	DESIG=71
Γ_{78}	$\gamma\pi^0\eta$	$< 2.4 \times 10^{-6}$	DESIG=190
Γ_{79}	γK^+K^-	[a] $(1.14 \pm 0.13) \times 10^{-5}$	DESIG=111
Γ_{80}	$\gamma p\bar{p}$	[b] $< 6 \times 10^{-6}$	DESIG=102
Γ_{81}	$\gamma 2h^+2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$	DESIG=103
Γ_{82}	$\gamma 3h^+3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$	DESIG=20
Γ_{83}	$\gamma 4h^+4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$	DESIG=21
Γ_{84}	$\gamma\pi^+\pi^-K^+K^-$	$(2.9 \pm 0.9) \times 10^{-4}$	DESIG=22
Γ_{85}	$\gamma 2\pi^+2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$	DESIG=14
Γ_{86}	$\gamma 3\pi^+3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$	DESIG=13
Γ_{87}	$\gamma 2\pi^+2\pi^-K^+K^-$	$(2.4 \pm 1.2) \times 10^{-4}$	DESIG=17
Γ_{88}	$\gamma\pi^+\pi^-p\bar{p}$	$(1.5 \pm 0.6) \times 10^{-4}$	DESIG=18
Γ_{89}	$\gamma 2\pi^+2\pi^-p\bar{p}$	$(4 \pm 6) \times 10^{-5}$	DESIG=15
Γ_{90}	$\gamma 2K^+2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$	DESIG=19
Γ_{91}	$\gamma\eta'(958)$	$< 1.9 \times 10^{-6}$	DESIG=16
Γ_{92}	$\gamma\eta$	$< 1.0 \times 10^{-6}$	DESIG=55
Γ_{93}	$\gamma f_0(980)$	$< 3 \times 10^{-5}$	DESIG=54
Γ_{94}	$\gamma f'_2(1525)$	$(2.9 \pm 0.6) \times 10^{-5}$	DESIG=105
Γ_{95}	$\gamma f_2(1270)$	$(1.01 \pm 0.06) \times 10^{-4}$	DESIG=52
Γ_{96}	$\gamma\eta(1405)$	$< 8.2 \times 10^{-5}$	DESIG=51
Γ_{97}	$\gamma f_0(1500)$	$< 1.5 \times 10^{-5}$	DESIG=65
Γ_{98}	$\gamma f_0(1500) \rightarrow \gamma K^+K^-$	$(1.0 \pm 0.4) \times 10^{-5}$	DESIG=108
Γ_{99}	$\gamma f_0(1710)$	$< 2.6 \times 10^{-4}$	DESIG=192
Γ_{100}	$\gamma f_0(1710) \rightarrow \gamma K^+K^-$	$(1.01 \pm 0.32) \times 10^{-5}$	DESIG=53
Γ_{101}	$\gamma f_0(1710) \rightarrow \gamma\pi^+\pi^-$	$(5.3 \pm 2.0) \times 10^{-6}$	DESIG=112
Γ_{102}	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$	$< 1.4 \times 10^{-6}$	DESIG=191
Γ_{103}	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$< 1.8 \times 10^{-6}$	DESIG=109
Γ_{104}	$\gamma f_4(2050)$	$< 5.3 \times 10^{-5}$	DESIG=110
Γ_{105}	$\gamma f_0(2200) \rightarrow \gamma K^+K^-$	$< 2 \times 10^{-4}$	DESIG=104
Γ_{106}	$\gamma f_J(2220) \rightarrow \gamma K^+K^-$	$< 8 \times 10^{-7}$	DESIG=69
Γ_{107}	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	$< 6 \times 10^{-7}$	DESIG=60
Γ_{108}	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	$< 1.1 \times 10^{-6}$	DESIG=61
Γ_{109}	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	$< 3 \times 10^{-5}$	DESIG=62
Γ_{110}	$\gamma\eta_c(1S)$	$< 5.7 \times 10^{-3}$	DESIG=68
Γ_{111}	$\gamma\chi_{c0}$	$< 6.5 \times 10^{-5}$	DESIG=119
Γ_{112}	$\gamma\chi_{c1}$	$< 2.3 \times 10^{-4}$	DESIG=120
Γ_{113}	$\gamma\chi_{c2}$	$< 7.6 \times 10^{-5}$	DESIG=121
Γ_{114}	$\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi$	$< 1.6 \times 10^{-6}$	DESIG=122
Γ_{115}	$\gamma\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0J/\psi$	$< 2.8 \times 10^{-6}$	DESIG=123
Γ_{116}	$\gamma X(3915) \rightarrow \omega J/\psi$	$< 3.0 \times 10^{-6}$	DESIG=124
Γ_{117}	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	$< 2.2 \times 10^{-6}$	DESIG=125
Γ_{118}	γX	[c] $< 4.5 \times 10^{-6}$	DESIG=126
Γ_{119}	$\gamma X\bar{X} (m_X < 3.1 \text{ GeV})$	[d] $< 1 \times 10^{-3}$	DESIG=66
Γ_{120}	$\gamma X\bar{X} (m_X < 4.5 \text{ GeV})$	[e] $< 2.4 \times 10^{-4}$	DESIG=67
Γ_{121}	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[f] $< 1.78 \times 10^{-4}$	DESIG=127
Γ_{122}	$\gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	[g] $< 9 \times 10^{-6}$	DESIG=113
Γ_{123}	$\gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[h] $< 1.30 \times 10^{-4}$	DESIG=114
Γ_{124}	$\gamma a_1^0 \rightarrow \gamma gg$	[h] $< 1 \% \times 10^{-3}$	DESIG=115
Γ_{125}	$\gamma a_1^0 \rightarrow \gamma s\bar{s}$	[h] $< 1 \times 10^{-3}$	DESIG=116

Lepton Family number (*LF*) violating modes

Γ_{126}	$\mu^\pm\tau^\mp$	LF	$< 6.0 \times 10^{-6}$	CL=95%	NODE=M049;CLUMP=C
					DESIG=116

Other decays					NODE=M049;CLUMP=D DESIG=106
Γ_{127} invisible	< 3.0		$\times 10^{-4}$	CL=90%	
[a] $2m_\tau < M(\tau^+ \tau^-) < 9.2$ GeV					LINKAGE=E49
[b] 2 GeV $< m_{K^+ K^-} < 3$ GeV					LINKAGE=G49
[c] $X =$ scalar with $m < 8.0$ GeV					LINKAGE=A49
[d] $X \bar{X} =$ vectors with $m < 3.1$ GeV					LINKAGE=B49
[e] X and $\bar{X} =$ zero spin with $m < 4.5$ GeV					LINKAGE=F49
[f] 1.5 GeV $< m_X < 5.0$ GeV					LINKAGE=C49
[g] 201 MeV $< M(\mu^+ \mu^-) < 3565$ MeV					LINKAGE=D49
[h] 0.5 GeV $< m_X < 9.0$ GeV, where m_X is the invariant mass of the hadronic final state.					LINKAGE=I49

 $\Gamma(1S) \Gamma(i) \Gamma(e^+ e^-)/\Gamma(\text{total})$

$\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$	$\Gamma_2 \Gamma_3/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$31.2 \pm 1.6 \pm 1.7$	KOBEL	92	$e^+ e^- \rightarrow \mu^+ \mu^-$

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_0 \Gamma_2/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.240 ± 0.016 OUR AVERAGE			

1.252 $\pm 0.004 \pm 0.019$	¹ ROSNER	06	CLEO	$9.5 e^+ e^- \rightarrow \text{hadrons}$
1.187 $\pm 0.023 \pm 0.031$	¹ BARU	92B	MD1	$e^+ e^- \rightarrow \text{hadrons}$
1.23 $\pm 0.02 \pm 0.05$	¹ JAKUBOWSKI	88	CBAL	$e^+ e^- \rightarrow \text{hadrons}$
1.37 $\pm 0.06 \pm 0.09$	² GILES	84B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$
1.23 $\pm 0.08 \pm 0.04$	² ALBRECHT	82	DASP	$e^+ e^- \rightarrow \text{hadrons}$
1.13 $\pm 0.07 \pm 0.11$	² NICZYPORUK	82	LENA	$e^+ e^- \rightarrow \text{hadrons}$
1.09 ± 0.25	² BOCK	80	CNTR	$e^+ e^- \rightarrow \text{hadrons}$
1.35 ± 0.14	³ BERGER	79	PLUT	$e^+ e^- \rightarrow \text{hadrons}$

1 Radiative corrections evaluated following KURAEV 85.

2 Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

3 Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$. **$\Gamma(1S) \text{ PARTIAL WIDTHS}$**

$\Gamma(e^+ e^-)$	Γ_2		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.340 ± 0.018 OUR EVALUATION			

 $\Gamma(1S) \text{ BRANCHING RATIOS}$

$\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN
2.60 ± 0.10 OUR AVERAGE			
2.53 $\pm 0.13 \pm 0.05$	60k	¹ BESSON	07
2.61 $\pm 0.12^{+0.09}_{-0.13}$	25k	CINABRO	94B
2.7 $\pm 0.4 \pm 0.2$		² ALBRECHT	85C
3.4 $\pm 0.4 \pm 0.4$		GILES	83
		CLEO	$e^+ e^- \rightarrow \tau^+ \tau^-$
		ARG	$\Gamma(2S) \rightarrow \pi^+ \pi^- \tau^+ \tau^-$

¹ BESSON 07 reports $[\Gamma(\Gamma(1S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}]/[B(\Gamma(1S) \rightarrow \mu^+ \mu^-)] = 1.02 \pm 0.02 \pm 0.05$ which we multiply by our best value $B(\Gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Gamma(1S) \rightarrow ee) = B(\Gamma(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN
2.38 ± 0.11 OUR AVERAGE			
2.29 $\pm 0.08 \pm 0.11$		ALEXANDER	98
2.42 $\pm 0.14 \pm 0.14$	307	ALBRECHT	87
2.8 $\pm 0.3 \pm 0.2$	826	BESSON	84
5.1 ± 3.0		BERGER	80C
		CLEO	$\Gamma(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
		ARG	$\Gamma(2S) \rightarrow \pi^+ \pi^- e^+ e^-$
		PLUT	$e^+ e^- \rightarrow e^+ e^-$

NODE=M049218

NODE=M049G1
NODE=M049G1NODE=M049G2
NODE=M049G2NODE=M049G2;LINKAGE=B
NODE=M049G2;LINKAGE=R
NODE=M049G2;LINKAGE=P

NODE=M049220

NODE=M049W2
NODE=M049W2
→ UNCHECKED ←

NODE=M049225

NODE=M049R3
NODE=M049R3

NODE=M049R3;LINKAGE=BE

NODE=M049R3;LINKAGE=A

NODE=M049R2
NODE=M049R2

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE	EVTS
0.0248±0.0005 OUR AVERAGE	

0.0249±0.0002±0.0007	345k	ADAMS	05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0249±0.0008±0.0013		ALEXANDER	98	CLE2	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.0212±0.0020±0.0010		¹ BARU	92	MD1	$e^+e^- \rightarrow \mu^+\mu^-$
0.0231±0.0012±0.0010		¹ KOBEL	92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0252±0.0007±0.0007		CHEN	89B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0261±0.0009±0.0011		KAARSBERG	89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.0230±0.0025±0.0013	86	ALBRECHT	87	ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.029 ± 0.003 ± 0.002	864	BESSON	84	CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
0.027 ± 0.003 ± 0.003		ANDREWS	83	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.032 ± 0.013 ± 0.003		ALBRECHT	82	DASP	$e^+e^- \rightarrow \mu^+\mu^-$
0.038 ± 0.015 ± 0.002		NICZYPORUK	82	LENA	$e^+e^- \rightarrow \mu^+\mu^-$
0.014 +0.034 -0.014		BOCK	80	CNTR	$e^+e^- \rightarrow \mu^+\mu^-$
0.022 ± 0.020		BERGER	79	PLUT	$e^+e^- \rightarrow \mu^+\mu^-$

¹ Taking into account interference between the resonance and continuum.

 $\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

VALUE	EVTS
1.008±0.023 OUR AVERAGE	

1.005±0.013±0.022	0.7M	¹ DEL-AMO-SA..10C	BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(1S)$
1.02 ± 0.02 ± 0.05	60k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S)$

¹ Allows any number of extra photons with total energy < 500 MeV.

 $\Gamma(ggg)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS
81.7±0.7	20M

¹ BESSON	06A	CLEO	$\Upsilon(1S) \rightarrow \text{hadrons}$
---------------------	-----	------	---

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ from BESSON 06A and PDG 08 values of $B(\mu^+\mu^-) = (2.48 \pm 0.05)\%$ and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS
2.20±0.60	400k

¹ BESSON	06A	CLEO	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
---------------------	-----	------	--

¹ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

 $\Gamma(\gamma gg)/\Gamma(ggg)$

VALUE (units 10^{-2})	EVTS
2.70±0.01±0.27	20M

BESSON	06A	CLEO	$\Upsilon(1S) \rightarrow (\gamma +) \text{hadrons}$
--------	-----	------	--

 $\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$

VALUE
0.0294±0.0024 OUR AVERAGE

0.030 ± 0.002 ± 0.002

0.028 ± 0.004 ± 0.002

DOCUMENT ID	TECN	COMMENT
-------------	------	---------

AQUINES	06A	CLE3	$\Upsilon(1S) \rightarrow \eta' \text{anything}$
ARTUSO	03	CLE2	$\Upsilon(1S) \rightarrow \eta' \text{anything}$

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS
--------------------------	-----	------

5.4 ± 0.4 OUR FIT Error includes scale factor of 1.4.

5.4 ± 0.4 OUR AVERAGE Error includes scale factor of 1.5.

5.25±0.13±0.25	3k	SHEN	16	BELL	$e^+e^- \rightarrow J/\psi X$
6.4 ± 0.4 ± 0.6	730	BRIERE	04	CLEO	$e^+e^- \rightarrow J/\psi X$
11 ± 4 ± 2		¹ FULTON	89	CLEO	$e^+e^- \rightarrow \mu^+\mu^-X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	ALBRECHT	92J	ARG	$e^+e^- \rightarrow e^+e^-X, \mu^+\mu^-X$
<17	90	MASCHMANN	90	CBAL	$e^+e^- \rightarrow \text{hadrons}$
<200	90	NICZYPORUK	83	LENA	

¹ Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.

 Γ_3/Γ

NODE=M049R1

NODE=M049R1

NODE=M049R1;LINKAGE=G

NODE=M049R43

NODE=M049R43

NODE=M049R43;LINKAGE=DE

NODE=M049R35

NODE=M049R35

NODE=M049R35;LINKAGE=BE

NODE=M049R36

NODE=M049R36

NODE=M049R36;LINKAGE=BE

NODE=M049R37

NODE=M049R37

NODE=M049R73

NODE=M049R73

NODE=M049R12

NODE=M049R12

NODE=M049R12;LINKAGE=K

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$					Γ_8/Γ	NODE=M049R85 NODE=M049R85
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.2 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$					Γ_9/Γ	NODE=M049R86 NODE=M049R86
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$					Γ_{10}/Γ	NODE=M049R87 NODE=M049R87
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT		
$3.90 \pm 1.21 \pm 0.23$	20	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$					Γ_{11}/Γ	NODE=M049R88 NODE=M049R88
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<1.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$					Γ_{12}/Γ	NODE=M049R89 NODE=M049R89
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.2 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$					Γ_{13}/Γ	NODE=M049R90 NODE=M049R90
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					Γ_{14}/Γ	NODE=M049R91 NODE=M049R91
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.4 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	
$\Gamma(X(4350) \text{ anything, } X \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$					Γ_{15}/Γ	NODE=M049P05 NODE=M049P05
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<8.1 \times 10^{-6}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi K^+ K^- X$	
$\Gamma(Z_c(3900)^{\pm} \text{ anything, } Z_c \rightarrow J/\psi(1S)\pi^{\pm})/\Gamma_{\text{total}}$					Γ_{16}/Γ	NODE=M049P06 NODE=M049P06
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<1.3 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi \pi^{\pm} X$	
$\Gamma(Z_c(4200)^{\pm} \text{ anything, } Z_c \rightarrow J/\psi(1S)\pi^{\pm})/\Gamma_{\text{total}}$					Γ_{17}/Γ	NODE=M049P07 NODE=M049P07
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<6.0 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi \pi^{\pm} X$	
$\Gamma(Z_c(4430)^{\pm} \text{ anything, } Z_c \rightarrow J/\psi(1S)\pi^{\pm})/\Gamma_{\text{total}}$					Γ_{18}/Γ	NODE=M049P08 NODE=M049P08
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<4.9 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi \pi^{\pm} X$	
$\Gamma(X_{cs}^{\pm} \text{ anything, } X \rightarrow J/\psi K^{\pm})/\Gamma_{\text{total}}$					Γ_{19}/Γ	NODE=M049P11 NODE=M049P11
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.7 \times 10^{-6}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi K^- X$	
$\Gamma(\chi_{c1}(3872) \text{ anything, } \chi_{c1} \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{20}/Γ	NODE=M049R00 NODE=M049R00
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<9.5 \times 10^{-6}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi \pi^+\pi^- X$	
$\Gamma(\psi(4260) \text{ anything, } \psi \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{21}/Γ	NODE=M049R99 NODE=M049R99
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.8 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi \pi^+\pi^- X$	
$\Gamma(\psi(4260) \text{ anything, } \psi \rightarrow J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$					Γ_{22}/Γ	NODE=M049P03 NODE=M049P03
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<7.5 \times 10^{-6}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi K^+ K^- X$	
$\Gamma(\chi_{c1}(4140) \text{ anything, } \chi_{c1} \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$					Γ_{23}/Γ	NODE=M049P04 NODE=M049P04
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.2 \times 10^{-6}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow J/\psi K^+ K^- X$	

$\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{24}/Γ	NODE=M049R25 NODE=M049R25
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<7.4	90	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$		
$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$					Γ_{25}/Γ	NODE=M049P13 NODE=M049P13
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.90±0.35 OUR FIT						
1.90±0.43±0.14	215	JIA	17	BELL $\gamma(1S) \rightarrow \gamma J/\psi(1S)$		
$\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{25}/Γ	NODE=M049R26 NODE=M049R26
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
0.35±0.07 OUR FIT						
0.35±0.08±0.06	52 ± 12	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$		
$\Gamma(\chi_{c1}(1P)X_{\text{tetra}})/\Gamma_{\text{total}}$					Γ_{26}/Γ	NODE=M049P15 NODE=M049P15
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 37.8×10^{-6}	90	¹ JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$		
1 For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 37.8×10^{-6} .						
$\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{27}/Γ	NODE=M049R27 NODE=M049R27
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
0.52±0.12±0.09	47 ± 11	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$		
$\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$					Γ_{28}/Γ	NODE=M049P12 NODE=M049P12
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.23±0.17±0.11	215	SHEN	16	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$					Γ_{28}/Γ	NODE=M049R28 NODE=M049R28
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
0.41±0.11±0.08	42 ± 11	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- X$		
$\Gamma(\psi(2S) \eta_c)/\Gamma_{\text{total}}$					Γ_{29}/Γ	NODE=M049R92 NODE=M049R92
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 3.6×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) \chi_{c0})/\Gamma_{\text{total}}$					Γ_{30}/Γ	NODE=M049R93 NODE=M049R93
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 6.5×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) \chi_{c1})/\Gamma_{\text{total}}$					Γ_{31}/Γ	NODE=M049R94 NODE=M049R94
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 4.5×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) \chi_{c2})/\Gamma_{\text{total}}$					Γ_{32}/Γ	NODE=M049R95 NODE=M049R95
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 2.1×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) \eta_c(2S))/\Gamma_{\text{total}}$					Γ_{33}/Γ	NODE=M049R96 NODE=M049R96
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 3.2×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) X(3940))/\Gamma_{\text{total}}$					Γ_{34}/Γ	NODE=M049R97 NODE=M049R97
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 2.9×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(2S) X(4160))/\Gamma_{\text{total}}$					Γ_{35}/Γ	NODE=M049R98 NODE=M049R98
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 2.9×10^{-6}	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
$\Gamma(\psi(4260) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{36}/Γ	NODE=M049P00 NODE=M049P00
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 7.9×10^{-5}	90	SHEN	16	BELL $\gamma(1S) \rightarrow \psi(2S)\pi^+\pi^- X$		

$\Gamma(\psi(4360) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{37}/Γ	NODE=M049P01 NODE=M049P01
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<5.2 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow \psi(2S)\pi^+\pi^- X$	
$\Gamma(\psi(4660) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{38}/Γ	NODE=M049P02 NODE=M049P02
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<2.2 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow \psi(2S)\pi^+\pi^- X$	
$\Gamma(X(4050)^{\pm} \text{ anything}, X \rightarrow \psi(2S)\pi^{\pm})/\Gamma_{\text{total}}$					Γ_{39}/Γ	NODE=M049P09 NODE=M049P09
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<8.8 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow \psi(2S)\pi^{\pm} X$	
$\Gamma(Z_c(4430)^{\pm} \text{ anything}, Z_c \rightarrow \psi(2S)\pi^{\pm})/\Gamma_{\text{total}}$					Γ_{40}/Γ	NODE=M049P10 NODE=M049P10
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<6.7 \times 10^{-5}$	90	SHEN	16	BELL	$\gamma(1S) \rightarrow \psi(2S)\pi^{\pm} X$	
$\Gamma(Z_c(4200)^+ Z_c(4200)^-)/\Gamma_{\text{total}}$					Γ_{41}/Γ	NODE=M049P17 NODE=M049P17
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<22.3 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow J/\psi\pi^{\pm} X$	
1 Assuming $B(Z_c(4200)^{\pm} \rightarrow J/\psi\pi^{\pm}) = 1$.						
$\Gamma(Z_c(3900)^{\pm} Z_c(4200)^{\mp})/\Gamma_{\text{total}}$					Γ_{42}/Γ	NODE=M049P18 NODE=M049P18
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<8.1 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow J/\psi\pi^{\pm} X$	
1 Assuming $B(Z_c(4200)^{\pm} \rightarrow J/\psi\pi^{\pm}) = 1 = B(Z_c(3900)^{\pm} \rightarrow J/\psi\pi^{\pm})$.						
$\Gamma(Z_c(3900)^+ Z_c(3900)^-)/\Gamma_{\text{total}}$					Γ_{43}/Γ	NODE=M049P19 NODE=M049P19
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<1.8 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow J/\psi\pi^{\pm} X$	
1 Assuming $B(Z_c(3900)^{\pm} \rightarrow J/\psi\pi^{\pm}) = 1$						
$\Gamma(X(4050)^+ X(4050)^-)/\Gamma_{\text{total}}$					Γ_{44}/Γ	NODE=M049P20 NODE=M049P20
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<15.8 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow \chi_{c1}(1P)\pi^{\pm} X$	
1 Assuming $B(X(4050)^{\pm} \rightarrow \chi_{c1}(1P)\pi^{\pm}) = 1$						
$\Gamma(X(4250)^+ X(4250)^-)/\Gamma_{\text{total}}$					Γ_{45}/Γ	NODE=M049P21 NODE=M049P21
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<26.6 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow \chi_{c1}(1P)\pi^{\pm} X$	
1 Assuming $B(X(4250)^{\pm} \rightarrow \chi_{c1}(1P)\pi^{\pm}) = 1$						
$\Gamma(X(4050)^{\pm} X(4250)^{\mp})/\Gamma_{\text{total}}$					Γ_{46}/Γ	NODE=M049P22 NODE=M049P22
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<44.2 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow \chi_{c1}(1P)\pi^{\pm} X$	
1 Assuming $B(X(4050)^{\pm} \rightarrow \chi_{c1}(1P)\pi^{\pm}) = 1 = B(X(4250)^{\pm} \rightarrow \chi_{c1}(1P)\pi^{\pm})$						
$\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$					Γ_{47}/Γ	NODE=M049P23 NODE=M049P23
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<20.3 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(2S) \rightarrow \psi(2S)\pi^{\pm} X$	
1 Assuming $B(Z_c(4430)^{\pm} \rightarrow \psi(2S)\pi^{\pm}) = 1$						
$\Gamma(X(4055)^{\pm} X(4055)^{\mp})/\Gamma_{\text{total}}$					Γ_{48}/Γ	NODE=M049P25 NODE=M049P25
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<23.3 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow \psi(2S)\pi^{\pm} X$	
1 Assuming $B(X(4055)^{\pm} \rightarrow \psi(2S)\pi^{\pm}) = 1$						
$\Gamma(X(4055)^{\pm} Z_c(4430)^{\mp})/\Gamma_{\text{total}}$					Γ_{49}/Γ	NODE=M049P26 NODE=M049P26
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<45.5 \times 10^{-6}$	90	¹ JIA	18	BELL	$\gamma(1S) \rightarrow \psi(2S)\pi^{\pm} X$	
1 Assuming $B(X(4055)^{\pm} \rightarrow \psi(2S)\pi^{\pm}) = 1 = B(Z_c(4430)^{\pm} \rightarrow \psi(2S)\pi^{\pm})$						

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.68	90	SHEN	13	BELL $\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1 $\times 10^3$	90	BLINOV	90	MD1 $\gamma(1S) \rightarrow \rho^0 \pi^0$
<2 $\times 10^2$	90	FULTON	90B	$\gamma(1S) \rightarrow \rho^0 \pi^0$
<2.1 $\times 10^3$	90	NICZYPORUK	83	LENA $\gamma(1S) \rightarrow \rho^0 \pi^0$

 Γ_{50}/Γ

NODE=M049R11
NODE=M049R11

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<3.90	90	SHEN	13	BELL $\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

 Γ_{51}/Γ

NODE=M049R05
NODE=M049R05

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	BARU	92	MD1 $\gamma(1S) \rightarrow \pi^+ \pi^-$

 Γ_{52}/Γ

NODE=M049R57
NODE=M049R57

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	BARU	92	MD1 $\gamma(1S) \rightarrow K^+ K^-$

 Γ_{53}/Γ

NODE=M049R58
NODE=M049R58

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<5	90	1 BARU	96	MD1 $\gamma(1S) \rightarrow p\bar{p}$

 Γ_{54}/Γ

NODE=M049R59
NODE=M049R59

¹ Supersedes BARU 92 in this node.

 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.14 $\pm 0.72 \pm 0.34$		26 ± 9	SHEN	13	BELL $\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0$

 Γ_{55}/Γ

NODE=M049R72
NODE=M049R72

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18.4	90	ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow$ hadrons
-------	----	------------	----	------------------------------------

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.36 $\pm 0.37 \pm 0.29$	56	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(K^+ K^-)$

 Γ_{56}/Γ

NODE=M049R75
NODE=M049R75

 $\Gamma(\omega\pi^+ \pi^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.46 $\pm 0.67 \pm 0.72$	64	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(\pi^+ \pi^-)^0$

 Γ_{57}/Γ

NODE=M049R76
NODE=M049R76

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
4.42 $\pm 0.50 \pm 0.58$	173	SHEN	12A	BELL $\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{58}/Γ

NODE=M049R77
NODE=M049R77

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.63	90	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(K^+ K^-)$

 Γ_{59}/Γ

NODE=M049R78
NODE=M049R78

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<1.79	90	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(\pi^+ \pi^-)^0$

 Γ_{60}/Γ

NODE=M049R79
NODE=M049R79

 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.24	90	SHEN	12A	BELL $\gamma(1S) \rightarrow 2(\pi^+ \pi^-)^0$

 Γ_{61}/Γ

NODE=M049R80
NODE=M049R80

 $\Gamma(K^*(892)^0 \bar{K}^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.02 $\pm 0.68 \pm 0.34$	42	SHEN	12A	BELL $\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{62}/Γ

NODE=M049R81
NODE=M049R81

 $\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<2.41	90	SHEN	12A	BELL $\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 Γ_{63}/Γ

NODE=M049R82
NODE=M049R82

$\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$					Γ_{64}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R83 NODE=M049R83
$1.02 \pm 0.35 \pm 0.22$	24	SHEN	12A	BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(b_1(1235)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$					Γ_{65}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R84 NODE=M049R84
<1.25	90	SHEN	12A	BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{66}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R06 NODE=M049R06
$12.8 \pm 2.0 \pm 2.3$	143 ± 22	SHEN	13	BELL	$\gamma(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{67}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M049R07 NODE=M049R07
$1.59 \pm 0.33 \pm 0.18$	37 ± 8	SHEN	13	BELL	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<3.4	90	¹ DOBBS	12A		$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$	
1 Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.						
$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{68}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R08 NODE=M049R08
$2.92 \pm 0.85 \pm 0.37$	16 ± 5	SHEN	13	BELL	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$	
$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{69}/Γ	
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R09 NODE=M049R09
<1.11	90	SHEN	13	BELL	$\gamma(1S) \rightarrow K_S^0 K^- \pi^+$	
$\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$					Γ_{70}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P14 NODE=M049P14
$4.6 \pm 2.8 \pm 1.3$	3.1k	JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$	
$\Gamma(D^*(2010)^{\pm} \text{ anything})/\Gamma_{\text{total}}$					Γ_{71}/Γ	
<u>VALUE (units 10^{-3})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M049R32 NODE=M049R32
$25.2 \pm 1.3 \pm 1.5$	≈ 2k	¹ AUBERT	10C	BABR	$\gamma(2S) \rightarrow \pi^+ \pi^- \gamma(1S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<19	90	² ALBRECHT	92J	ARG	$e^+ e^- \rightarrow D^0 \pi^{\pm} X$	
1 For $x_p > 0.1$.						
2 For $x_p > 0.2$.						
$\Gamma(f_1(1285) X_{\text{tetra}})/\Gamma_{\text{total}}$					Γ_{72}/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P16 NODE=M049P16
$<62.4 \times 10^{-6}$	90	¹ JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$	
1 For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.6×10^{-6} to 62.4×10^{-6} .						
$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$					Γ_{73}/Γ	
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R33 NODE=M049R33
$2.85 \pm 0.25 \text{ OUR AVERAGE}$						OCCUR=2
2.81 ± 0.49 ± 0.20		LEES	14G	BABR	$e^+ e^- \rightarrow \overline{2H} X$	
2.86 ± 0.19 ± 0.21	455	ASNER	07	CLEO	$e^+ e^- \rightarrow \overline{2H} X$	
$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$					Γ_{74}/Γ	
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>				NODE=M049R02 NODE=M049R02
1.200 ± 0.017	^{1,2} DOBBS	12A			$\gamma(1S) \rightarrow \text{hadrons}$	
1 DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.						
2 Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.						
$\Gamma(ggg, \gamma gg \rightarrow \overline{d} \text{ anything})/\Gamma(ggg, \gamma gg \rightarrow \text{anything})$						
<u>VALUE (units 10^{-5})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R34 NODE=M049R34
$3.36 \pm 0.23 \pm 0.25$	455	ASNER	07	CLEO	$e^+ e^- \rightarrow \overline{d} X$	

$\Gamma(\gamma\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**6.3±1.2±1.3**¹ For $m_{\pi\pi} > 1$ GeV. $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**1.7±0.6±0.3**¹ For $m_{\pi\pi} > 1$ GeV. $\Gamma(\gamma\pi\pi(\text{S-wave}))/\Gamma_{\text{total}}$ VALUE (units 10^{-5})**4.63±0.56±0.48** $\Gamma(\gamma\pi^0\eta)/\Gamma_{\text{total}}$ VALUE (units 10^{-6}) CL%**<2.4**

90

¹ BESSON 07A obtained this limit for $0.7 < m_{\pi^0\eta} < 3$ GeV. $\Gamma(\gamma K^+K^-)/\Gamma_{\text{total}}$ $(2 < m_{K^+K^-} < 3$ GeV)VALUE (units 10^{-5}) CL%**1.14±0.08±0.10**

90

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$ $(2 < m_{p\bar{p}} < 3$ GeV)VALUE (units 10^{-5}) CL%**<0.6**

90

 $\Gamma(\gamma 2h^+2h^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**7.0±1.1±1.0**

80 ± 12

 $\Gamma(\gamma 3h^+3h^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**5.4±1.5±1.3**

39 ± 11

 $\Gamma(\gamma 4h^+4h^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**7.4±2.5±2.5**

36 ± 12

 $\Gamma(\gamma\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**2.9±0.7±0.6**

29 ± 8

 $\Gamma(\gamma 2\pi^+2\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**2.5±0.7±0.5**

26 ± 7

 $\Gamma(\gamma 3\pi^+3\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**2.5±0.9±0.8**

17 ± 5

 $\Gamma(\gamma 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**2.4±0.9±0.8**

18 ± 7

 $\Gamma(\gamma\pi^+\pi^-p\bar{p})/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**1.5±0.5±0.3**

22 ± 6

DOCUMENT ID	TECN	COMMENT
¹ ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$

 Γ_{75}/Γ NODE=M049R70
NODE=M049R70

DOCUMENT ID	TECN	COMMENT
¹ ANASTASSOV 99	CLE2	$e^+e^- \rightarrow \text{hadrons}$

 Γ_{76}/Γ NODE=M049R71
NODE=M049R71

DOCUMENT ID	TECN	COMMENT
LEES	18A	BABR

 Γ_{77}/Γ NODE=M049P27
NODE=M049P27

DOCUMENT ID	TECN	COMMENT
¹ BESSON 07A	07A	CLEO

 Γ_{78}/Γ NODE=M049R47
NODE=M049R47

DOCUMENT ID	TECN	COMMENT
ATHAR	06	CLE3

 Γ_{79}/Γ NODE=M049R24
NODE=M049R24

DOCUMENT ID	TECN	COMMENT
ATHAR	06	CLE3

 Γ_{80}/Γ NODE=M049R29
NODE=M049R29
NODE=M049R29

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{81}/Γ NODE=M049R20
NODE=M049R20

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{82}/Γ NODE=M049R21
NODE=M049R21

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{83}/Γ NODE=M049R22
NODE=M049R22

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{84}/Γ

NODE=M049R14

NODE=M049R14

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{85}/Γ

NODE=M049R13

NODE=M049R13

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{86}/Γ

NODE=M049R17

NODE=M049R17

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{87}/Γ

NODE=M049R18

NODE=M049R18

DOCUMENT ID	TECN	COMMENT
FULTON	90B	CLEO

 Γ_{88}/Γ

NODE=M049R15

NODE=M049R15

$\Gamma(\gamma 2\pi^+ 2\pi^- p\bar{p})/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{89}/Γ
0.4±0.4±0.4	7 ± 6	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

 $\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{90}/Γ
0.2±0.2	2 ± 2	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{91}/Γ
< 1.9	90	ATHAR	07A	CLEO $\gamma(1S) \rightarrow \gamma\eta' \rightarrow \gamma\pi^+\pi^-\eta, \gamma\rho$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					

<16 90 RICHICHI 01B CLE2 $\gamma(1S) \rightarrow \gamma\eta' \rightarrow \gamma\eta\pi^+\pi^-$

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{92}/Γ
< 1.0	90	ATHAR	07A	CLEO $\gamma(1S) \rightarrow \gamma\eta \rightarrow \gamma\gamma\gamma, \gamma\pi^+\pi^-\pi^0, \gamma 3\pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21 90 MASEK 02 CLEO $\gamma(1S) \rightarrow \gamma\eta$

 $\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{93}/Γ
<3	90	¹ ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$	

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 1$.

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{94}/Γ
2.9 ±0.6 OUR AVERAGE						

2.13 ± 0.28 ± 0.72		1 LEES	18A	BABR $\gamma(1S) \rightarrow \gamma K^+ K^-$		
4.0 ± 1.4 ± 0.1		17 BESSON	11	CLEO $\gamma(1S) \rightarrow K_S^0 K_S^0$		
3.7 ± 0.9 ± 0.8		ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma K^+ K^-$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14 90 ³ FULTON 90B CLEO $\gamma(1S) \rightarrow \gamma K^+ K^-$
<19.4 90 ³ ALBRECHT 89 ARG $\gamma(1S) \rightarrow \gamma K^+ K^-$

¹ Using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.887 \pm 0.022$ and $B(K^0\bar{K}^0) = 1/2 B(K\bar{K})$.

² BESSON 11 reports $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\gamma(1S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})]$ assuming $B(f'_2(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$, which we rescale to our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (88.7 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The result also assumes $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$ and $B(f'_2(1525) \rightarrow K\bar{K}) = 4 B(f'_2(1525) \rightarrow K_S^0\bar{K}_S^0)$.

³ Assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

<u>VALUE (units 10^{-5})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{95}/Γ
10.1 ±0.6 OUR AVERAGE					

10.15 ± 0.59 ± 0.54		1 LEES	18A	BABR $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$		
10.5 ± 1.6 ± 1.9		2 BESSON	07A	CLE3 $\gamma(1S) \rightarrow \gamma\pi^0\pi^0$		
10.2 ± 0.8 ± 0.7		ATHAR	06	CLE3 $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$		
8.1 ± 2.3 ± 2.7		³ ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21 90 ³ FULTON 90B CLEO $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
<13 90 ³ ALBRECHT 89 ARG $\gamma(1S) \rightarrow \gamma\pi^+\pi^-$
<81 90 SCHMITT 88 CBAL $\gamma(1S) \rightarrow \gamma X$

¹ Using $B(f_2(1270) \rightarrow \pi^0\pi^0) = 1/3 B(f_2(1270) \rightarrow \pi\pi)$ and $B(f_2(1270) \rightarrow \pi\pi) = (84.2 \pm 2.9)\%$.

² Using $B(f_2(1270) \rightarrow \pi^0\pi^0) = B(f_2(1270) \rightarrow \pi\pi)/3$ and $B(f_2(1270) \rightarrow \pi\pi) = (84.7 \pm 2.5)\%$.

³ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.

NODE=M049R19

NODE=M049R19

NODE=M049R16

NODE=M049R16

NODE=M049R55

NODE=M049R55

NODE=M049R54

NODE=M049R54

NODE=M049R31

NODE=M049R31

NODE=M049R31;LINKAGE=AT

NODE=M049R52

NODE=M049R52

NODE=M049R52;LINKAGE=A

NODE=M049R52;LINKAGE=BE

NODE=M049R52;LINKAGE=D

NODE=M049R51

NODE=M049R51

NODE=M049R51;LINKAGE=A

NODE=M049R51;LINKAGE=BE

NODE=M049R51;LINKAGE=C

$\Gamma(\gamma\eta(1405))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{96}/Γ
<8.2	90	¹ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$	

¹ Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$.

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{97}/Γ
<1.5	90	¹ BESSON	07A CLEO	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma \pi^0 \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{97}/Γ
<6.1	90	² BESSON	07A CLEO	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma \eta \eta$	

¹ Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi\pi)/3$ and $B(f_0(1500) \rightarrow \pi\pi) = (0.349 \pm 0.023)\%$.

² Calculated by us using $B(f_0(1500) \rightarrow \eta\eta) = (5.1 \pm 0.9)\%$.

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{98}/Γ
1.04 ± 0.14 ± 0.33	1	LEES	18A BABR	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma K^+ K^-$	

¹ LEES 18A quotes $B(\gamma(1S) \rightarrow \gamma f_0(1500) \rightarrow \gamma K\bar{K}) = (2.08 \pm 0.27 \pm 0.65) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K\bar{K})$.

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
< 2.6	90	¹ ALBRECHT	89 ARG	$\gamma(1S) \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
< 6.3	90	¹ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma K^+ K^-$	

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
<19	90	¹ FULTON	90B CLEO	$\gamma(1S) \rightarrow \gamma K_S^0 K_S^0$	

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
< 8	90	² ALBRECHT	89 ARG	$\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$	

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
<24	90	³ SCHMITT	88 CBAL	$\gamma(1S) \rightarrow \gamma X$	

¹ Assuming $B(f_0(1710) \rightarrow K\bar{K}) = 0.38$.

² Assuming $B(f_0(1710) \rightarrow \pi\pi) = 0.04$.

³ Assuming $B(f_0(1710) \rightarrow \eta\eta) = 0.18$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ
1.01 ± 0.26 ± 0.18	1	LEES	18A BABR	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ
<0.7	90	ATHAR	06 CLEO	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma K^+ K^-$	

¹ LEES 18A quotes $B(\gamma(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma K\bar{K}) = (2.02 \pm 0.51 \pm 0.35) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K\bar{K})$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{101}/Γ
0.53 ± 0.17 ± 0.11	1	LEES	18A BABR	$\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$	

¹ LEES 18A quotes $B(\gamma(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi\pi) = (0.79 \pm 0.26 \pm 0.17) \times 10^{-5}$ assuming $B(\pi^0 \pi^0) = 1/3 B(\pi\pi)$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{102}/Γ
<1.4	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma \pi^0 \pi^0$	

¹ Assuming $B(f_0(1710) \rightarrow \pi\pi) = 0.17$.

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{103}/Γ
<1.8	90	BESSON	07A CLEO	$e^+ e^- \rightarrow \gamma(1S) \rightarrow \gamma \eta\eta$	

 $\Gamma(\gamma f_4(2050))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{104}/Γ
<5.3	90	¹ ATHAR	06 CLE3	$\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$	

¹ Assuming $B(f_4(2050) \rightarrow \pi\pi) = 0.17$.

NODE=M049R23

NODE=M049R23

NODE=M049R23;LINKAGE=J

NODE=M049R44

NODE=M049R44

OCCUR=2

NODE=M049R44;LINKAGE=BE

NODE=M049R44;LINKAGE=BS

NODE=M049P29

NODE=M049P29

NODE=M049R53

NODE=M049R53

OCCUR=2

OCCUR=2

NODE=M049R53;LINKAGE=E

NODE=M049R53;LINKAGE=F

NODE=M049R53;LINKAGE=A

NODE=M049R50

NODE=M049R50

NODE=M049P28

NODE=M049P28

NODE=M049R45

NODE=M049R45

NODE=M049R46

NODE=M049R46

NODE=M049R30

NODE=M049R30

NODE=M049R30;LINKAGE=AT

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$					Γ_{105}/Γ	NODE=M049R63 NODE=M049R63
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.0002	90	BARU	89	MD1	$\gamma(1S) \rightarrow \gamma K^+ K^-$	
$\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$					Γ_{106}/Γ	NODE=M049R56 NODE=M049R56
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT		
< 8	90	ATHAR	06	CLE3	$\gamma(1S) \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 160	90	MASEK	02	CLEO	$\gamma(1S) \rightarrow \gamma K^+ K^-$	
< 150	90	FULTON	90B	CLEO	$\gamma(1S) \rightarrow \gamma K^+ K^-$	
< 290	90	ALBRECHT	89	ARG	$\gamma(1S) \rightarrow \gamma K^+ K^-$	
<2000	90	BARU	89	MD1	$\gamma(1S) \rightarrow \gamma K^+ K^-$	
$\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{107}/Γ	NODE=M049R41 NODE=M049R41
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT		
< 6	90	ATHAR	06	CLE3	$\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<120	90	MASEK	02	CLEO	$\gamma(1S) \rightarrow \gamma \pi^+ \pi^-$	
$\Gamma(\gamma f_J(2220) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{108}/Γ	NODE=M049R42 NODE=M049R42
VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT		
< 11	90	ATHAR	06	CLE3	$\gamma(1S) \rightarrow \gamma p\bar{p}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<160	90	MASEK	02	CLEO	$\gamma(1S) \rightarrow \gamma p\bar{p}$	
$\Gamma(\gamma\eta(2225) \rightarrow \gamma\phi\phi)/\Gamma_{\text{total}}$					Γ_{109}/Γ	NODE=M049R62 NODE=M049R62
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.003	90	BARU	89	MD1	$\gamma(1S) \rightarrow \gamma K^+ K^- K^+ K^-$	
$\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$					Γ_{110}/Γ	NODE=M049R38 NODE=M049R38
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<5.7	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$					Γ_{111}/Γ	NODE=M049R39 NODE=M049R39
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
<6.5	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$					Γ_{112}/Γ	NODE=M049R40 NODE=M049R40
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<2.3	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$					Γ_{113}/Γ	NODE=M049R48 NODE=M049R48
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<7.6	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$					Γ_{114}/Γ	NODE=M049R49 NODE=M049R49
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<1.6	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$					Γ_{115}/Γ	NODE=M049R68 NODE=M049R68
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<2.8	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$					Γ_{116}/Γ	NODE=M049R69 NODE=M049R69
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<3.0	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	
$\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$					Γ_{117}/Γ	NODE=M049R74 NODE=M049R74
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<2.2	90	SHEN	10A	BELL	$\gamma(1S) \rightarrow \gamma X$	

$\Gamma(\gamma X)/\Gamma_{\text{total}}$ (X = scalar with $m < 8.0$ GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.5	90	1 DEL-AMO-SA..11J	BABR	$e^+ e^- \rightarrow \gamma + X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	90	2 BAEST	95	CLEO $e^+ e^- \rightarrow \gamma + X$
-----	----	---------	----	---------------------------------------

1 For a noninteracting scalar X with mass $m < 8.0$ GeV.2 For a noninteracting pseudoscalar X with mass < 7.2 GeV. Γ_{118}/Γ

NODE=M049R60

NODE=M049R60

NODE=M049R60

 $\Gamma(\gamma X\bar{X}(m_X < 3.1 \text{ GeV}))/\Gamma_{\text{total}}$ (X \bar{X} = vectors with $m < 3.1$ GeV)

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	1 BAEST	95	CLEO $e^+ e^- \rightarrow \gamma + X\bar{X}$

1 For a noninteracting vector X with mass < 3.1 GeV. Γ_{119}/Γ

NODE=M049R60;LINKAGE=DA

NODE=M049R60;LINKAGE=A

NODE=M049R61

NODE=M049R61

NODE=M049R61

 $\Gamma(\gamma X\bar{X}(m_X < 4.5 \text{ GeV}))/\Gamma_{\text{total}}$ Γ_{120}/Γ X and \bar{X} = zero spin with $m < 4.5$ GeV

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	1 DEL-AMO-SA..11J	BABR	$e^+ e^- \rightarrow \gamma + X\bar{X}$

1 For a noninteracting scalar X with mass $m < 4.5$ GeV. $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{121}/Γ (1.5 GeV $< m_X < 5.0$ GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.78	95	ROSNER	07A	CLEO $e^+ e^- \rightarrow \gamma X$

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{122}/Γ (201 $< M(\mu^+ \mu^-) < 3565$ MeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	1 LOVE	08	CLEO $e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9.7	90	2 LEES	13C	BABR $e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$
------	----	--------	-----	--

1 For a narrow scalar or pseudoscalar a_1^0 with $201 < M(\mu^+ \mu^-) < 3565$ MeV, excluding J/ψ . Measured 90% CL limits as a function of $M(\mu^+ \mu^-)$ range from $1-9 \times 10^{-6}$.2 For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9200 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from $0.28-9.7 \times 10^{-6}$. $\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ (2 $m_\tau < M(\tau^+ \tau^-) < 9.2$ GeV)

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<130	90	1 LEES	13R	BABR $\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 50	90	2 LOVE	08	CLEO $e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$
------	----	--------	----	--

1 For a narrow scalar a_1^0 with $2m_\tau < M(a_1^0) < 9.2$ GeV, which result in a 90% CL upper limits of 0.9×10^{-5} at $M(a_1^0) = 2m_\tau$, $\approx 1.5 \times 10^{-5}$ at $M(a_1^0) = 7.5$ GeV, and 13×10^{-5} at $M(a_1^0) = 9.2$ GeV.2 For a narrow scalar or pseudoscalar a_1^0 with $2m_\tau < M(a_1^0) < 7.5$ GeV, which result in a 90% CL limits ranging from 1×10^{-5} at $M(a_1^0) = 2m_\tau$ to 5×10^{-5} at $M(a_1^0) = 7.5$ GeV. $\Gamma(\gamma a_1^0 \rightarrow \gamma gg)/\Gamma_{\text{total}}$ Γ_{124}/Γ (0.5 GeV $< m < 9.0$ GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1 \times 10^{-2}$	90	1 LEES	13L	BABR $\Upsilon(1S) \rightarrow \gamma X$

1 For a narrow, CP -odd pseudoscalar a_1^0 searched for in 26 hadronic decay modes with invariant mass $0.5 \text{ GeV} < m_X < 9.0 \text{ GeV}$. Measured 90% CL limit as a function of m_X range from 10^{-6} to 10^{-2} .

NODE=M049R60

NODE=M049R60

NODE=M049R60

NODE=M049R60;LINKAGE=DA

NODE=M049R60;LINKAGE=A

NODE=M049R61

NODE=M049R61

NODE=M049R61

NODE=M049R61;LINKAGE=A

NODE=M049R01

NODE=M049R01

NODE=M049R01

NODE=M049R01;LINKAGE=DA

NODE=M049R64

NODE=M049R64

NODE=M049R64

NODE=M049R65

NODE=M049R65

NODE=M049R65

NODE=M049R65;LINKAGE=LO

NODE=M049R65;LINKAGE=LE

NODE=M049R66

NODE=M049R66

NODE=M049R66

NODE=M049R66;LINKAGE=A

NODE=M049R66;LINKAGE=LO

NODE=M049R03

NODE=M049R03

NODE=M049R03

NODE=M049R03;LINKAGE=A

$\Gamma(\gamma a_1^0 \rightarrow \gamma s\bar{s})/\Gamma_{\text{total}}$ (0.5 GeV < m < 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1 \times 10^{-3}$	90	1 LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$

¹ For a narrow, CP-odd pseudoscalar a_1^0 searched for in 14 hadronic decay modes with invariant mass $1.5 \text{ GeV} < m_X < 9.0 \text{ GeV}$. Measured 90% CL limit as a function of m_X range from 10^{-5} to 10^{-3} .

 Γ_{125}/Γ

NODE=M049R04

NODE=M049R04

NODE=M049R04

NODE=M049R04;LINKAGE=A

LEPTON FAMILY NUMBER (L_f) VIOLATING MODES $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

OTHER DECAYS $\Gamma(\text{invisible})/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.0	90	AUBERT	09AX BABR	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<39	90	RUBIN	07 CLEO	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
<25	90	TAJIMA	07 BELL	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

 $\Upsilon(1S)$ REFERENCES

JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58949
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58950
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=57635
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
SHEN	16	PR D93 112013	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=57515
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54949
LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55167
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55451
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	(BELLE Collab.)	REFID=54746
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314
BESSON	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=16737
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16495
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53211
DEL-AMO-SA...	10C	PR D104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53354
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545
AUBERT	09AX	PR L 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53201
LOVE	08	PR L 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52565
LOVE	08A	PR L 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51945
BESSON	07	PR L 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
BESSON	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51638
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)	REFID=51629
TAJIMA	07	PR L 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=51645
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51510
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50993
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PR L 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PR L 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50183
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=49395
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)	REFID=48846
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)	REFID=48345
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	(CLEO Collab.)	REFID=47424
ANASTASSOV	99	PR L 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=46609
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)	REFID=44146
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)	REFID=44102
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=42167
BARU	92	ZPHY C46 229	S.E. Baru <i>et al.</i>	(NOVO)	REFID=41860
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)	REFID=42168
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41361
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=41012
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=40918
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUELLER	88	HE e ⁺ e ⁻ Physics 412	W. Buchmuller, S. Cooper Editors: A. Ali and P. Soeding, World Scientific, Singapore	(HANN, DESY, MIT)	REFID=40034
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGPC	REFID=40742
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)	REFID=40582
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22284
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22282
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033

Translated from YAF 41 733.

ARTAMONOV	84	PL 137B	272	A.S. Artamonov <i>et al.</i>	(NOVO)
BESSON	84	PR D30	1433	D. Besson <i>et al.</i>	(CLEO Collab.)
GILES	84B	PR D29	1285	R. Giles <i>et al.</i>	(CLEO Collab.)
MACKAY	84	PR D29	2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)
ANDREWS	83	PRL	50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)
GILES	83	PRL	50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)
NICZYPORUK	83	ZPHY	C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
ALBRECHT	82	PL	116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)
ARTAMONOV	82	PL	118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)
NICZYPORUK	82	ZPHY	C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)
BERGER	80C	PL	93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)
BOCK	80	ZPHY	C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)
BERGER	79	ZPHY	C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)

REFID=22278
REFID=22279
REFID=22280
REFID=22281
REFID=22273
REFID=22274
REFID=12488
REFID=22270
REFID=22271
REFID=22272
REFID=22263
REFID=22264
REFID=22259

NODE=M076

 $\chi_{b0}(1P)$

$$\Gamma^G(JPC) = 0^+(0^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b0}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID	COMMENT		
9859.44 ± 0.42 ± 0.31 OUR EVALUATION	From average γ energy below, using $\Upsilon(2S)$ mass = 10023.26 ± 0.31 MeV			

 $m_{\chi_{b1}(1P)} - m_{\chi_{b0}(1P)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
32.49 ± 0.93	LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M076M

NODE=M076M
→ UNCHECKED ←NODE=M076M2
NODE=M076M2 **γ ENERGY IN $\Upsilon(2S)$ DECAY**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
162.5 ± 0.4 OUR AVERAGE			
162.56 ± 0.19 ± 0.42	ARTUSO	05	CLEO $\Upsilon(2S) \rightarrow \gamma X$
162.0 ± 0.8 ± 1.2	EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
162.1 ± 0.5 ± 1.4	ALBRECHT	85E	ARG $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
163.8 ± 1.6 ± 2.7	NERNST	85	CBAL $\Upsilon(2S) \rightarrow \gamma X$
158.0 ± 7 ± 1	HAAS	84	CLEO $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
149.4 ± 0.7 ± 5.0	KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma X$

NODE=M076DM

NODE=M076DM

 $\chi_{b0}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level	
$\Gamma_1 \gamma \Upsilon(1S)$	(1.94 ± 0.27) %		
$\Gamma_2 D^0 X$	< 10.4 %	90%	DESIG=1
$\Gamma_3 \pi^+ \pi^- K^+ K^- \pi^0$	< 1.6 $\times 10^{-4}$	90%	DESIG=2
$\Gamma_4 2\pi^+ \pi^- K^- K_S^0$	< 5 $\times 10^{-5}$	90%	DESIG=3
$\Gamma_5 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 5 $\times 10^{-4}$	90%	DESIG=4
$\Gamma_6 2\pi^+ 2\pi^- 2\pi^0$	< 2.1 $\times 10^{-4}$	90%	DESIG=5
$\Gamma_7 2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.6) $\times 10^{-4}$	90%	DESIG=6
$\Gamma_8 2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.7 $\times 10^{-4}$	90%	DESIG=7
$\Gamma_9 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 5 $\times 10^{-4}$	90%	DESIG=8
$\Gamma_{10} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 1.6 $\times 10^{-4}$	90%	DESIG=9
$\Gamma_{11} 3\pi^+ 3\pi^-$	< 8 $\times 10^{-5}$	90%	DESIG=10
$\Gamma_{12} 3\pi^+ 3\pi^- 2\pi^0$	< 6 $\times 10^{-4}$	90%	DESIG=11
$\Gamma_{13} 3\pi^+ 3\pi^- K^+ K^-$	(2.4 ± 1.2) $\times 10^{-4}$	90%	DESIG=12
$\Gamma_{14} 3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 1.0 $\times 10^{-3}$	90%	DESIG=13
$\Gamma_{15} 4\pi^+ 4\pi^-$	< 8 $\times 10^{-5}$	90%	DESIG=14
$\Gamma_{16} 4\pi^+ 4\pi^- 2\pi^0$	< 2.1 $\times 10^{-3}$	90%	DESIG=15
$\Gamma_{17} J/\psi J/\psi$	< 7 $\times 10^{-5}$	90%	DESIG=16
$\Gamma_{18} J/\psi \psi(2S)$	< 1.2 $\times 10^{-4}$	90%	DESIG=17
$\Gamma_{19} \psi(2S) \psi(2S)$	< 3.1 $\times 10^{-5}$	90%	DESIG=18
$\Gamma_{20} J/\psi(1S) \text{anything}$	< 2.3 $\times 10^{-3}$	90%	DESIG=19

NODE=M076215; NODE=M076

$\chi_{b0}(1P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
1.94±0.27 OUR AVERAGE						
2.07±0.24±0.21	1,2	LEES	14M	BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$	
1.76±0.30±0.18	87	KORNICER	11	CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.6	90	5 LEES	11J	BABR	$\Upsilon(2S) \rightarrow X\gamma$	
< 6	90	WALK	86	CBAL	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$	
< 11	90	PAUSS	83	CUSB	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$	

1 LEES 14M quotes $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$ = $(7.75 \pm 0.91) \times 10^{-4}$ combining the results from samples of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ with and without converted photons. Assumes $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

2 LEES 14M reports $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (7.75 \pm 0.91) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

3 Assuming $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$.

4 KORNICER 11 reports $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (6.59 \pm 0.96 \pm 0.60) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

5 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
<10.4 × 10⁻²	90	6,7 BRIERE	08	CLEO	$\Upsilon(2S) \rightarrow \gamma D^0 X$

6 For $p_{D^0} > 2.5$ GeV/c.

7 The authors also present their result as $(5.6 \pm 3.6 \pm 0.5) \times 10^{-2}$.

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
<1.6	90	8 ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$
8 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.					

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
<0.5	90	9 ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$
9 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.					

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
<5	90	10 ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$
10 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 18 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.					

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
<2.1	90	11 ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$
11 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 8 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.					

NODE=M076220

NODE=M076R1

NODE=M076R1

NODE=M076R1;LINKAGE=A

NODE=M076R1;LINKAGE=B

NODE=M076R1;LINKAGE=KA
NODE=M076R1;LINKAGE=KR

NODE=M076R1;LINKAGE=LE

NODE=M076R01
NODE=M076R01

NODE=M076R01;LINKAGE=BR
NODE=M076R01;LINKAGE=RI

NODE=M076R02
NODE=M076R02

NODE=M076R02;LINKAGE=AS

NODE=M076R03
NODE=M076R03

NODE=M076R03;LINKAGE=AS

NODE=M076R04
NODE=M076R04

NODE=M076R04;LINKAGE=AS

NODE=M076R05
NODE=M076R05

NODE=M076R05;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.6±0.1	7	12 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$
12 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] = (4 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_7/Γ

NODE=M076R06
NODE=M076R06

NODE=M076R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	13 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
13 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 10 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_8/Γ

NODE=M076R07
NODE=M076R07

NODE=M076R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	14 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
14 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 20 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_9/Γ

NODE=M076R08
NODE=M076R08

NODE=M076R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	15 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
15 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_{10}/Γ

NODE=M076R09
NODE=M076R09

NODE=M076R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	16 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
16 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_{11}/Γ

NODE=M076R10
NODE=M076R10

NODE=M076R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	17 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
17 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 22 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_{12}/Γ

NODE=M076R11
NODE=M076R11

NODE=M076R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±1.2±0.2	9	18 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
18 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] = (9 \pm 4 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{13}/Γ

NODE=M076R12
NODE=M076R12

NODE=M076R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10	90	19 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
19 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 37 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_{14}/Γ

NODE=M076R13
NODE=M076R13

NODE=M076R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	20 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
20 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

 Γ_{15}/Γ

NODE=M076R14
NODE=M076R14

NODE=M076R14;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$					Γ_{16}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<21	90	21 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$	NODE=M076R15 NODE=M076R15
21 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))]$ < 7.7×10^{-6} which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.					NODE=M076R15;LINKAGE=AS
$\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$					Γ_{17}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<7	90	22 SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$	NODE=M076R16 NODE=M076R16
22 SHEN 12 reports < 7.1×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] \text{ assuming } B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.					NODE=M076R16;LINKAGE=SH
$\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$					Γ_{18}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<12	90	23 SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$	NODE=M076R17 NODE=M076R17
23 SHEN 12 reports < 12×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] \text{ assuming } B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.					NODE=M076R17;LINKAGE=SH
$\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$					Γ_{19}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<3.1	90	24 SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$	NODE=M076R18 NODE=M076R18
24 SHEN 12 reports < 3.1×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] \text{ assuming } B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.					NODE=M076R18;LINKAGE=SH
$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$					Γ_{20}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.3 \times 10^{-3}$	90	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M076R00 NODE=M076R00

$\chi_{b0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

$\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma \times \Gamma_{61}^{(2S)}/\Gamma^{(2S)}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.7 $\times 10^{-3}$	90	25 LEES	11J BABR	$\Gamma(2S) \rightarrow X \gamma$	NODE=M076230
25 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $\Gamma(\gamma \Gamma(1S))/\Gamma_{\text{total}} < 4.6\%$ using $B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4)\%$.					NODE=M076B02 NODE=M076B02
$B(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P)) \times B(\Gamma(1S) \rightarrow \ell^+ \ell^-)$					
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.67 ± 0.28 OUR AVERAGE					NODE=M076B01 NODE=M076B01
2.9 ± 1.7 ± 0.1		26 LEES	14M BABR	$\gamma \gamma \mu^+ \mu^-$	NODE=M076B01;LINKAGE=A
1.63 ± 0.24 ± 0.15	87	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$	
26 From a sample of $\Gamma(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.					
$[B(\chi_{b0}(1P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(2S) \rightarrow \gamma \chi_{b0}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Gamma(1S)) \times B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$					
VALUE (%)		DOCUMENT ID	TECN	COMMENT	
3.28 ± 0.37		27 LEES	14M BABR	$\gamma \gamma \mu^+ \mu^-$	NODE=M076A01 NODE=M076A01
27 From a sample of $\Gamma(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ without converted photons.					NODE=M076A01;LINKAGE=A

$\chi_{b0}(1P)$ REFERENCES

JIA	17A PR D96 112002	S. Jia et al.	(BELLE Collab.)	NODE=M076
LEES	14M PR D90 112010	J.P. Lees et al.	(BABAR Collab.)	REFID=58318
SHEN	12 PR D85 071102	C.P. Shen et al.	(BELLE Collab.)	REFID=56343
KORNICER	11 PR D83 054003	M. Kornicer et al.	(CLEO Collab.)	REFID=54313
LEES	11J PR D84 072002	J.P. Lees et al.	(BABAR Collab.)	REFID=16769
ASNER	08A PR D78 091103	D.M. Asner et al.	(CLEO Collab.)	REFID=53936
BRIERE	08 PR D78 092007	R.A. Briere et al.	(CLEO Collab.)	REFID=52574
ARTUSO	05 PRL 94 032001	M. Artuso et al.	(CLEO Collab.)	REFID=52577
EDWARDS	99 PR D59 032003	K.W. Edwards et al.	(CLEO Collab.)	REFID=50454
WALK	86 PR D34 2611	W.S. Walk et al.	(CLEO Collab.)	REFID=46612
ALBRECHT	85E PL 160B 331	H. Albrecht et al.	(Crystal Ball Collab.)	REFID=22290
NERNST	85 PRL 54 2195	R. Nernst et al.	(ARGUS Collab.)	REFID=22288
HAAS	84 PRL 52 799	J. Haas et al.	(Crystal Ball Collab.)	REFID=22289
KLOPFEN... PAUSS	83 PRL 51 160 83 PL 130B 439	C. Klopfenstein et al. F. Pauss et al.	(CLEO Collab.) (CUSB Collab.) (MPIIM, COLU, CORN, LSU+)	REFID=22287 REFID=22285 REFID=22286

$\chi_{b1}(1P)$

$$I^G(JPC) = 0^+(1^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 1$ from SKWARNICKI 87.


 $\chi_{b1}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9892.78 ± 0.26 ± 0.31 OUR EVALUATION			From average γ energy below, using $\Upsilon(2S)$ mass = 10023.26 ± 0.31 MeV

 γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
129.63 ± 0.33 OUR AVERAGE			Error includes scale factor of 1.3. See the ideogram below.
129.58 ± 0.09 ± 0.29	ARTUSO 05	CLEO	$\Upsilon(2S) \rightarrow \gamma X$
128.8 ± 0.4 ± 0.6	EDWARDS 99	CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
131.7 ± 0.9 ± 1.3	WALK 86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
131.7 ± 0.3 ± 1.1	ALBRECHT 85E	ARG	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
130.6 ± 0.8 ± 2.4	NERNST 85	CBAL	$\Upsilon(2S) \rightarrow \gamma X$
129 ± 0.8 ± 1	HAAS 84	CLEO	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
128.1 ± 0.4 ± 3.0	KLOPFEN... 83	CUSB	$\Upsilon(2S) \rightarrow \gamma X$
130.6 ± 3.0	PAUSS 83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

WEIGHTED AVERAGE
129.63 ± 0.33 (Error scaled by 1.3)



$\chi_{b1}(1P)$ mass (MeV)

 $\chi_{b1}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \gamma \Upsilon(1S)$	(35.2 ± 2.0) %	
$\Gamma_2 D^0 X$	(12.6 ± 2.2) %	
$\Gamma_3 \pi^+ \pi^- K^+ K^- \pi^0$	(2.0 ± 0.6) × 10 ⁻⁴	
$\Gamma_4 2\pi^+ \pi^- K^- K_S^0$	(1.3 ± 0.5) × 10 ⁻⁴	
$\Gamma_5 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 6 × 10 ⁻⁴	90%
$\Gamma_6 2\pi^+ 2\pi^- 2\pi^0$	(8.0 ± 2.5) × 10 ⁻⁴	
$\Gamma_7 2\pi^+ 2\pi^- K^+ K^-$	(1.5 ± 0.5) × 10 ⁻⁴	
$\Gamma_8 2\pi^+ 2\pi^- K^+ K^- \pi^0$	(3.5 ± 1.2) × 10 ⁻⁴	
$\Gamma_9 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(8.6 ± 3.2) × 10 ⁻⁴	
$\Gamma_{10} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	(9.3 ± 3.3) × 10 ⁻⁴	
$\Gamma_{11} 3\pi^+ 3\pi^-$	(1.9 ± 0.6) × 10 ⁻⁴	

NODE=M077215;NODE=M077

NODE=M077

NODE=M077M

NODE=M077M
→ UNCHECKED ←

NODE=M077DM

NODE=M077DM

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

DESIG=9

DESIG=10

DESIG=11

Γ_{12}	$3\pi^+ 3\pi^- 2\pi^0$	(1.7 \pm 0.5) \times 10 $^{-3}$		DESIG=12
Γ_{13}	$3\pi^+ 3\pi^- K^+ K^-$	(2.6 \pm 0.8) \times 10 $^{-4}$		DESIG=13
Γ_{14}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	(7.5 \pm 2.6) \times 10 $^{-4}$		DESIG=14
Γ_{15}	$4\pi^+ 4\pi^-$	(2.6 \pm 0.9) \times 10 $^{-4}$		DESIG=15
Γ_{16}	$4\pi^+ 4\pi^- 2\pi^0$	(1.4 \pm 0.6) \times 10 $^{-3}$		DESIG=16
Γ_{17}	ω anything	(4.9 \pm 1.4) %		DESIG=21
Γ_{18}	ωX_{tetra}	< 4.44 \times 10 $^{-4}$	90%	DESIG=22
Γ_{19}	$J/\psi J/\psi$	< 2.7 \times 10 $^{-5}$	90%	DESIG=17
Γ_{20}	$J/\psi \psi(2S)$	< 1.7 \times 10 $^{-5}$	90%	DESIG=18
Γ_{21}	$\psi(2S) \psi(2S)$	< 6 \times 10 $^{-5}$	90%	DESIG=19
Γ_{22}	$J/\psi(1S)$ anything	< 1.1 \times 10 $^{-3}$	90%	DESIG=20
Γ_{23}	$J/\psi(1S) X_{tetra}$	< 2.27 \times 10 $^{-4}$	90%	DESIG=23

$\chi_{b1}(1P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.352 \pm 0.020 OUR AVERAGE					
0.356 \pm 0.016 - 0.022	\pm 0.019	964k	1 FULSUM	18 BELL $\Upsilon(2S) \rightarrow \gamma X$	
0.364 \pm 0.017 - 0.019		2,3,4 LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$	
0.331 \pm 0.018 - 0.017	\pm 0.017	3222 4,5 KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$	
0.350 \pm 0.023 - 0.018	\pm 0.018	13k 6 LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$	
0.34 \pm 0.07 - 0.02	\pm 0.02	53 4,7,8 WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$	
0.47 \pm 0.18		KLOPFEN...	83 CUSB	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$	
1 FULSUM 18 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (2.45 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
2 LEES 14M quotes $\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{total} = (2.51 \pm 0.12) \%$ combining the results from samples of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ with and without converted photons.					
3 LEES 14M reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (2.51 \pm 0.12) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
4 Assuming $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.					
5 KORNICER 11 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (22.8 \pm 0.4 \pm 1.2) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
6 LEES 11J reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (24.1 \pm 0.6 \pm 1.5) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
7 WALK 86 quotes $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (5.8 \pm 0.9 \pm 0.7) \%$.					
8 WALK 86 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (23.4 \pm 3.63 \pm 2.82) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

$\Gamma(D^0 X)/\Gamma_{total}$

VALUE (units 10 $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
12.6 \pm 1.9 \pm 1.1	2310	1 BRIERE	08 CLEO	$\Upsilon(2S) \rightarrow \gamma D^0 X$	

1 For $p_{D^0} > 2.5$ GeV/c.

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{total}$

VALUE (units 10 $^{-4}$)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
2.0 \pm 0.6 \pm 0.1	18	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$	
1 ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (14 \pm 3 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M077220

NODE=M077R1
NODE=M077R1

OCCUR=4

NODE=M077R1;LINKAGE=J

NODE=M077R1;LINKAGE=B

NODE=M077R1;LINKAGE=D

NODE=M077R1;LINKAGE=KA
NODE=M077R1;LINKAGE=KR

NODE=M077R1;LINKAGE=LE

NODE=M077R1;LINKAGE=A

NODE=M077R1;LINKAGE=C

NODE=M077R01
NODE=M077R01

NODE=M077R01;LINKAGE=BR

NODE=M077R02
NODE=M077R02

NODE=M077R02;LINKAGE=AS

$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$	Γ_4/Γ
$1.3 \pm 0.5 \pm 0.1$ 11 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$	NODE=M077R03 NODE=M077R03

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (9 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}$	Γ_5/Γ
<6 90 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$	NODE=M077R04 NODE=M077R04

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] < 42 \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

$\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ
$8.0 \pm 2.4 \pm 0.4$ 46 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$	NODE=M077R05 NODE=M077R05

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (55 \pm 9 \pm 14) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$	Γ_7/Γ
$1.5 \pm 0.5 \pm 0.1$ 18 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$	NODE=M077R06 NODE=M077R06

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (10 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$	Γ_8/Γ
$3.5 \pm 1.2 \pm 0.2$ 22 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$	NODE=M077R07 NODE=M077R07

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (24 \pm 6 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$	Γ_9/Γ
$8.6 \pm 3.2 \pm 0.4$ 26 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$	NODE=M077R08 NODE=M077R08

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (59 \pm 14 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$	Γ_{10}/Γ
$9.3 \pm 3.3 \pm 0.5$ 21 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$	NODE=M077R09 NODE=M077R09

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (64 \pm 16 \pm 16) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$	Γ_{11}/Γ
$1.9 \pm 0.6 \pm 0.1$ 25 1 ASNER 08A CLEO $\gamma(2S) \rightarrow \gamma 3\pi^+3\pi^-$	NODE=M077R10 NODE=M077R10

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))] = (13 \pm 3 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R03
NODE=M077R03

NODE=M077R03;LINKAGE=AS

NODE=M077R04
NODE=M077R04

NODE=M077R04;LINKAGE=AS

NODE=M077R05
NODE=M077R05

NODE=M077R05;LINKAGE=AS

NODE=M077R06
NODE=M077R06

NODE=M077R06;LINKAGE=AS

NODE=M077R07
NODE=M077R07

NODE=M077R07;LINKAGE=AS

NODE=M077R08
NODE=M077R08

NODE=M077R08;LINKAGE=AS

NODE=M077R09
NODE=M077R09

NODE=M077R09;LINKAGE=AS

NODE=M077R10
NODE=M077R10

NODE=M077R10;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
17±5±1	56	¹ ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$	

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ $= (119 \pm 18 \pm 32) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$					Γ_{13}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.6±0.8±0.1	21	¹ ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$	

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ $= (18 \pm 4 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$					Γ_{14}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
7.5±2.6±0.4	28	¹ ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$	

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ $= (52 \pm 11 \pm 14) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$					Γ_{15}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.6±0.9±0.1	24	¹ ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$	

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ $= (18 \pm 4 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$					Γ_{16}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
14±5±1	26	¹ ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$	

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ $= (96 \pm 24 \pm 29) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\omega \text{anything})/\Gamma_{\text{total}}$					Γ_{17}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.9±1.3±0.6	51k	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$	

$\Gamma(\omega X_{\text{tetra}})/\Gamma_{\text{total}}$					Γ_{18}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<44.4 × 10⁻⁵	90	¹ JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$	

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 3.3×10^{-5} to 44.4×10^{-5} .

$\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$					Γ_{19}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<2.7	90	¹ SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$	

¹ SHEN 12 reports $< 2.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.

$\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$					Γ_{20}/Γ
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.7	90	¹ SHEN	12 BELL	$\Gamma(2S) \rightarrow \gamma \psi X$	

¹ SHEN 12 reports $< 1.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Gamma(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.

NODE=M077R11
NODE=M077R11

NODE=M077R11;LINKAGE=AS

NODE=M077R12
NODE=M077R12

NODE=M077R13
NODE=M077R13

NODE=M077R14
NODE=M077R14

NODE=M077R14;LINKAGE=AS

NODE=M077R15
NODE=M077R15

NODE=M077R15;LINKAGE=AS

NODE=M077R19
NODE=M077R19

NODE=M077R23
NODE=M077R23

NODE=M077R23;LINKAGE=A

NODE=M077R16
NODE=M077R16

NODE=M077R16;LINKAGE=SH

NODE=M077R17
NODE=M077R17

NODE=M077R17;LINKAGE=SH

$\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ
<6	90	¹ SHEN	12	BELL	$\Upsilon(2S) \rightarrow \gamma\psi X$
1 SHEN 12 reports $< 6.2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] \text{ assuming } B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.					

NODE=M077R18
NODE=M077R18

NODE=M077R18;LINKAGE=SH

 $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
$<1.1 \times 10^{-3}$	90	JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$

NODE=M077R00
NODE=M077R00 $\Gamma(J/\psi(1S)X_{\text{tetra}})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
$<22.7 \times 10^{-5}$	90	¹ JIA	17A	BELL	$e^+e^- \rightarrow \text{hadrons}$

NODE=M077R22
NODE=M077R22

NODE=M077R22;LINKAGE=A

1 For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 1.8×10^{-5} to 22.7×10^{-5} . $\chi_{b1}(1P)$ Cross-Particle Branching Ratios

$$\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \frac{\Gamma_{59}^{(2S)}}{\Gamma^{(2S)}}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{24}/Γ
$24.1 \pm 0.6 \pm 1.5$	13k	LEES	11J	BABR	$\Upsilon(2S) \rightarrow X\gamma$

NODE=M077B03
NODE=M077B03


$$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{25}/Γ
5.90 ± 0.34 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.	

NODE=M077B01
NODE=M077B01

6.86 $\pm 0.47 \pm 0.44$	1 LEES	14M	BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$
6.86 $\pm 0.47 \pm 0.44$	1 LEES	14M	BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$
5.65 $\pm 0.11 \pm 0.27$	3222	KORNICER	11	CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
5.8 $\pm 0.9 \pm 0.7$	53	WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M077B01;LINKAGE=A

1 From a sample of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.

$$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{26}/Γ
1.30 ± 0.34 OUR AVERAGE					

NODE=M077B02
NODE=M077B02

1.16 $\pm 0.78 \pm 0.14$	1 LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
1.16 $\pm 0.78 \pm 0.14$	1 LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
1.33 $\pm 0.30 \pm 0.23$	50	KORNICER	11	CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M077B02;LINKAGE=A

1 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.

$B(\chi_{b2}(1P) \rightarrow pX + \bar{p}X)/B(\chi_{b1}(1P) \rightarrow pX + \bar{p}X)$

VALUE	DOCUMENT ID	TECN	COMMENT
1.068±0.010±0.040	BRIERE	07	CLEO $\gamma(2S) \rightarrow \gamma\chi_{bJ}(1P)$
1.11±0.15±0.20	BRIERE	07	CLEO $\gamma(2S) \rightarrow \gamma\chi_{bJ}(1P)$

 $\chi_{b1}(1P)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)
WALK	86	PR D34 2611	W.S. Wall <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)

 $h_b(1P)$

$I^G(J^{PC}) = 0^-(1^{+-})$

Quantum numbers are quark model predictions, $C = -$ established by $\eta_b\gamma$ decay.

 $h_b(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9899.3±0.8 OUR AVERAGE				
9899.3±0.4±1.0	112k	TAMPONI	15	BELL $e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
9899.1±0.4±1.0	70k	MIZUK	12	BELL $e^+e^- \rightarrow \pi^+\pi^- \text{ hadrons}$
9902 ± 4 ± 2	10.8k	LEES	11K	BABR $\gamma(3S) \rightarrow \eta_b\gamma\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9898.2 ^{+1.1} _{-1.0} ^{+1.0}	50.0k	¹ ADACHI	12	BELL $10.86 e^+e^- \rightarrow \pi^+\pi^- \text{ MM}$

¹ Superseded by MIZUK 12.

 $h_b(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \eta_b(1S)\gamma$	(52 ⁺⁶ ₋₅) %

 $h_b(1P)$ BRANCHING RATIOS

Γ($\eta_b(1S)\gamma$)/Γ _{total}	Γ ₁ /Γ
52 ⁺⁶ ₋₅ OUR AVERAGE	

56 ± 8 ± 4	33.1k	¹ TAMPONI	15	BELL $e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
49.2±5.7 ^{+5.6} _{-3.3}	24k	MIZUK	12	BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^- \text{ hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 10.8k LEES 11K BABR $\gamma(3S) \rightarrow \eta_b\gamma\pi^0$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.

 $h_b(1P)$ REFERENCES

TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PR D108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PR D109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)

NODE=M077R20
NODE=M077R20

NODE=M077R21
NODE=M077R21

NODE=M077

REFID=59535
REFID=58318
REFID=56343
REFID=54313
REFID=16769
REFID=53936
REFID=52574
REFID=52577
REFID=51887
REFID=50454
REFID=46612
REFID=40019
REFID=22290
REFID=22288
REFID=22289
REFID=22287
REFID=22285
REFID=22286

NODE=M204

NODE=M204

NODE=M204M

NODE=M204M

NODE=M204M;LINKAGE=AD

NODE=M204215;NODE=M204

DESIG=1

NODE=M204225

NODE=M204R01
NODE=M204R01

NODE=M204R01;LINKAGE=A

NODE=M204

REFID=56996
REFID=53962
REFID=54718
REFID=53937

$\chi_{b2}(1P)$

$I^G(JPC) = 0^+(2^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 2$ from SKWARNICKI 87.

 $\chi_{b2}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID	COMMENT	
9912.21 ± 0.26 ± 0.31 OUR EVALUATION	From average γ energy below, using $\Upsilon(2S)$ mass = 10023.26 ± 0.31 MeV		

 $m_{\chi_{b2}(1P)} - m_{\chi_{b1}(1P)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
19.10 ± 0.25 OUR AVERAGE	Error includes scale factor of 1.1.		
19.81 ± 0.65 ± 0.20	¹ AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$
19.01 ± 0.24	LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$

¹ From the $\chi_{bj}(1P) \rightarrow \Upsilon(1S)\gamma$ transition.

 γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
110.44 ± 0.29 OUR AVERAGE	Error includes scale factor of 1.1.		
110.58 ± 0.08 ± 0.30	ARTUSO	05 CLEO	$\Upsilon(2S) \rightarrow \gamma X$
110.8 ± 0.3 ± 0.6	EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma \chi(1P)$
107.0 ± 1.1 ± 1.3	WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
110.6 ± 0.3 ± 0.9	ALBRECHT	85E ARG	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
110.4 ± 0.8 ± 2.2	NERNST	85 CBAL	$\Upsilon(2S) \rightarrow \gamma X$
109.5 ± 0.7 ± 1.0	HAAS	84 CLEO	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
108.2 ± 0.3 ± 2.0	KLOPFEN...	83 CUSB	$\Upsilon(2S) \rightarrow \gamma X$
108.8 ± 4.0	PAUSS	83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

 $\chi_{b2}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level	
$\Gamma_1 \gamma \Upsilon(1S)$	(18.0 ± 1.0) %		DESIG=1
$\Gamma_2 D^0 X$	< 7.9 %	90%	DESIG=2
$\Gamma_3 \pi^+ \pi^- K^+ K^- \pi^0$	(8 ± 5) × 10 ⁻⁵		DESIG=3
$\Gamma_4 2\pi^+ \pi^- K^- K_S^0$	< 1.0 × 10 ⁻⁴	90%	DESIG=4
$\Gamma_5 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	(5.3 ± 2.4) × 10 ⁻⁴		DESIG=5
$\Gamma_6 2\pi^+ 2\pi^- 2\pi^0$	(3.5 ± 1.4) × 10 ⁻⁴		DESIG=6
$\Gamma_7 2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.4) × 10 ⁻⁴		DESIG=7
$\Gamma_8 2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.1 ± 0.9) × 10 ⁻⁴		DESIG=8
$\Gamma_9 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(3.9 ± 1.8) × 10 ⁻⁴		DESIG=9
$\Gamma_{10} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 5 × 10 ⁻⁴	90%	DESIG=10
$\Gamma_{11} 3\pi^+ 3\pi^-$	(7.0 ± 3.1) × 10 ⁻⁵		DESIG=11
$\Gamma_{12} 3\pi^+ 3\pi^- 2\pi^0$	(1.0 ± 0.4) × 10 ⁻³		DESIG=12
$\Gamma_{13} 3\pi^+ 3\pi^- K^+ K^-$	< 8 × 10 ⁻⁵	90%	DESIG=13
$\Gamma_{14} 3\pi^+ 3\pi^- K^+ K^- \pi^0$	(3.6 ± 1.5) × 10 ⁻⁴		DESIG=14
$\Gamma_{15} 4\pi^+ 4\pi^-$	(8 ± 4) × 10 ⁻⁵		DESIG=15
$\Gamma_{16} 4\pi^+ 4\pi^- 2\pi^0$	(1.8 ± 0.7) × 10 ⁻³		DESIG=16
$\Gamma_{17} J/\psi J/\psi$	< 4 × 10 ⁻⁵	90%	DESIG=17
$\Gamma_{18} J/\psi \psi(2S)$	< 5 × 10 ⁻⁵	90%	DESIG=18
$\Gamma_{19} \psi(2S) \psi(2S)$	< 1.6 × 10 ⁻⁵	90%	DESIG=19
$\Gamma_{20} J/\psi(1S) \text{anything}$	(1.5 ± 0.4) × 10 ⁻³		DESIG=20

NODE=M078M

NODE=M078M
→ UNCHECKED ←

NODE=M078DM2

NODE=M078DM2

NODE=M078DM2;LINKAGE=A

NODE=M078DM

NODE=M078DM

NODE=M078215;NODE=M078

$\chi_{b2}(1P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	Γ_1/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
0.180 ± 0.010 OUR AVERAGE		<u>TECN</u>
$0.164^{+0.009}_{-0.010} \pm 0.008$	503k	¹ FULSUM
$0.185 \pm 0.008 \pm 0.009$		^{2,3,4} LEES
$0.186 \pm 0.011 \pm 0.009$	1770	^{4,5} KORNICER
$0.194^{+0.014}_{-0.017} \pm 0.009$	8k	⁶ LEES
$0.25 \pm 0.06 \pm 0.01$	35	^{4,7,8} WALK
0.20 ± 0.05		KLOPFEN...
1		$\Upsilon(2S) \rightarrow \gamma X$
2		$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
3		$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
4		$\Upsilon(2S) \rightarrow X \gamma$
5		$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
6		$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
7		$\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P) \times B(\Upsilon(2S) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (4.4 \pm 0.9 \pm 0.5) \%$
8		$\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P) \times B(\Upsilon(2S) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (17.7 \pm 3.6 \pm 2.0) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
1		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.17 \pm 0.01 \pm 0.06) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
2		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}} = (1.32 \pm 0.06) \%$ combining the results from samples of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and without converted photons.
3		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.32 \pm 0.06) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
4		Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05) \%$.
5		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.33 \pm 0.04 \pm 0.07) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
6		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (13.9 \pm 0.5 \pm 0.9) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
7		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (4.4 \pm 0.9 \pm 0.5) \%$.
8		$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (17.7 \pm 3.6 \pm 2.0) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
$\Gamma(D^0 X)/\Gamma_{\text{total}}$	Γ_2/Γ	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>
$<7.9 \times 10^{-2}$	90	^{1,2} BRIERE
		<u>TECN</u>
		$\Upsilon(2S) \rightarrow \gamma D^0 X$
1		For $p_{D^0} > 2.5$ GeV/c.
2		The authors also present their result as $(5.4 \pm 1.9 \pm 0.5) \times 10^{-2}$.
$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$	Γ_3/Γ	
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$0.84 \pm 0.50 \pm 0.04$	8	¹ ASNER
		<u>TECN</u>
		$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$
1		ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$	Γ_4/Γ	
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>
<1.0	90	¹ ASNER
		<u>TECN</u>
		$\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$
1		ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 7 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

$\Gamma(2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.3±2.4±0.3	11	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^02\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (38 \pm 14 \pm 10) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_5/Γ NODE=M078R04
NODE=M078R04

NODE=M078R04;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±1.4±0.2	19	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (25 \pm 8 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_6/Γ NODE=M078R05
NODE=M078R05

NODE=M078R05;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.4±0.1	14	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (8 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_7/Γ NODE=M078R06
NODE=M078R06

NODE=M078R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.9±0.1	13	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (15 \pm 5 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_8/Γ NODE=M078R07
NODE=M078R07

NODE=M078R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±1.8±0.2	11	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (28 \pm 11 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_9/Γ NODE=M078R08
NODE=M078R08

NODE=M078R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] < 36 \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.				

 Γ_{10}/Γ NODE=M078R09
NODE=M078R09

NODE=M078R09;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.31±0.03	9	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+3\pi^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (5 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{11}/Γ NODE=M078R10
NODE=M078R10

NODE=M078R10;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.2±3.6±0.5	34	1 ASNER	08A CLEO	$\Gamma(2S) \rightarrow \gamma 3\pi^+3\pi^-2\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P))] = (73 \pm 16 \pm 20) \times 10^{-6}$ which we divide by our best value $B(\Gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{12}/Γ NODE=M078R11
NODE=M078R11

NODE=M078R11;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.				

 Γ_{13}/Γ NODE=M078R12
NODE=M078R12 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.6±1.5±0.2	14	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (26 \pm 8 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{14}/Γ

NODE=M078R12;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.84±0.40±0.04	7	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{15}/Γ NODE=M078R13
NODE=M078R13 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
18±7±1	29	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (132 \pm 31 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{16}/Γ NODE=M078R15
NODE=M078R15 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 4.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] \text{ assuming } B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

 Γ_{17}/Γ NODE=M078R16
NODE=M078R16 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 4.9 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] \text{ assuming } B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

 Γ_{18}/Γ NODE=M078R17
NODE=M078R17 $\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	¹ SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 1.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] \text{ assuming } B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

 Γ_{19}/Γ NODE=M078R18
NODE=M078R18 $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.50±0.34±0.22	462	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$
1 SHEN 12 reports $< 1.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] \text{ assuming } B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

 Γ_{20}/Γ NODE=M078R00
NODE=M078R00 $\chi_{b2}(1P) \text{ Cross-Particle Branching Ratios}$

$$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}} = \frac{\Gamma_1}{\Gamma} \times \frac{\Gamma_{60}}{\Gamma} / \Gamma_{\Upsilon(2S)}$$

NODE=M078230

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
13.9±0.5±0.9	8k	LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$
1 LEES 8k reports $13.9 \pm 0.5 \pm 0.9$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] \text{ assuming } B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078B03
NODE=M078B03

$B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.38 ± 0.16 OUR AVERAGE				
$3.63^{+0.36}_{-0.34} + 0.18_{-0.19}$		¹ LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$
$3.29 \pm 0.09 \pm 0.16$	1770	KORNICER	11 CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
$4.4 \pm 0.9 \pm 0.5$	35	WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ From a sample of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.

$$[B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
55.6 ± 1.6		¹ LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$

¹ From a sample of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ events without converted photons.

 $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5 OUR AVERAGE				

$4.68^{+0.99}_{-0.92} \pm 0.37$		¹ LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$3.56 \pm 0.40 \pm 0.41$	126	KORNICER	11 CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

¹ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.

 $\chi_{b2}(1P)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)
WALK	86	PR D34 2611	W.S. Wall <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFENF...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)

NODE=M078B01
NODE=M078B01

 $\eta_b(2S)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

 $\eta_b(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$9999.0 \pm 3.5^{+2.8}_{-1.9}$	26k	¹ MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

$9974.6 \pm 2.3 \pm 2.1$	11 ± 4	2,3,4 DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
--------------------------	------------	-------------	----	---

¹ Assuming $\Gamma_{\eta_b(2S)} = 4.9$ MeV. Not independent of the corresponding mass difference measurement.

² SANDILYA 13 (Belle Collaboration) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of $(157.8 \pm 3.6) \times 10^6$ $\Upsilon(2S)$ decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction $\mathcal{B}[\Upsilon(2S) \rightarrow \eta_b(2S)\gamma] \times \sum_i \mathcal{B}[\eta_b(2S) \rightarrow X_i] < 4.9 \times 10^{-6}$, summed over the exclusive hadronic final states X_i , is an order of magnitude smaller than that reported by DOBBS 12.

³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

⁴ Assuming $\Gamma_{\eta_b(2S)} = 5$ MeV. Not independent of the corresponding mass difference measurement.

NODE=M078B01

NODE=M078B01

NODE=M078B01;LINKAGE=A

NODE=M078A00

NODE=M078A00

NODE=M078A00;LINKAGE=A

NODE=M078B02

NODE=M078B02

NODE=M078B02;LINKAGE=A

NODE=M078

REFID=59535

REFID=58318

REFID=56199

REFID=56343

REFID=54313

REFID=16769

REFID=53936

REFID=52574

REFID=52577

REFID=50454

REFID=46612

REFID=40019

REFID=22290

REFID=22288

REFID=22289

REFID=22287

REFID=22285

REFID=22286

NODE=M200

NODE=M200

NODE=M200M

NODE=M200M

NODE=M200M;LINKAGE=MI

NODE=M200M;LINKAGE=A

NODE=M200M;LINKAGE=DO

NODE=M200M;LINKAGE=NI

NODE=M200DM

$$m_{\Upsilon(2S)} - m_{\eta_b(2S)}$$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
24.3±3.5^{+2.8}_{-1.9}	26k	5 MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
48.7±2.3±2.1	11 ± 4	6,7,8 DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
5 Assuming $\Gamma_{\eta_b}(2S) = 4.9$ MeV. Not independent of the corresponding mass measurement.				
6 SANDILYA 13 (Belle Collaboration) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of $(157.8 \pm 3.6) \times 10^6$ $\Upsilon(2S)$ decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction $\mathcal{B}[\Upsilon(2S) \rightarrow \eta_b(2S)\gamma] \times \sum_i \mathcal{B}[\eta_b(2S) \rightarrow X_i] < 4.9 \times 10^{-6}$, summed over the exclusive hadronic final states X_i , is an order of magnitude smaller than that reported by DOBBS 12.				
7 Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.				
8 Assuming $\Gamma_{\eta_b}(2S) = 5$ MeV. Not independent of the corresponding mass measurement.				

$\eta_b(2S)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma\pi^+\pi^-$ hadrons

$\eta_b(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	seen

$\eta_b(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	Γ_1/Γ
seen	$e^+ e^- \rightarrow \gamma\pi^+\pi^-$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	$\Upsilon(2S) \rightarrow \gamma$ hadrons
9 SANDILYA 13 (Belle Collaboration) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of $(157.8 \pm 3.6) \times 10^6$ $\Upsilon(2S)$ decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction $\mathcal{B}[\Upsilon(2S) \rightarrow \eta_b(2S)\gamma] \times \sum_i \mathcal{B}[\eta_b(2S) \rightarrow X_i] < 4.9 \times 10^{-6}$, summed over the exclusive hadronic final states X_i , is an order of magnitude smaller than that reported by DOBBS 12.	
10 Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.	

NODE=M200DM

NODE=M200DM;LINKAGE=MI

NODE=M200DM;LINKAGE=A

NODE=M200DM;LINKAGE=DO

NODE=M200DM;LINKAGE=NI

NODE=M200W

NODE=M200W

NODE=M200215;NODE=M200

DESIG=1

NODE=M200225

NODE=M200R01
NODE=M200R01

NODE=M200R01;LINKAGE=A

NODE=M200R01;LINKAGE=DO

NODE=M200

REFID=55590
REFID=54288
REFID=54718

SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

$\Upsilon(2S)$ $I^G(J^{PC}) = 0^-(1^- -)$

NODE=M052

 $\Upsilon(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10023.26 ± 0.31 OUR AVERAGE			
10023.5 \pm 0.5	¹ ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
10023.1 \pm 0.4	BARBER 84	REDE	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10023.6 \pm 0.5	^{2,3} BARU	86B	REDE $e^+ e^- \rightarrow$ hadrons
1 Reanalysis of BARU 86B using new electron mass (COHEN 87).			
2 Reanalysis of ARTAMONOV 84.			
3 Superseded by ARTAMONOV 00.			

NODE=M052M

NODE=M052M

NODE=M052M;LINKAGE=AR
 NODE=M052M;LINKAGE=C
 NODE=M052M;LINKAGE=RZ

 $m_{\Upsilon(3S)} - m_{\Upsilon(2S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$331.50 \pm 0.02 \pm 0.13$	LEES	11C	$e^+ e^- \rightarrow \pi^+ \pi^- X$

NODE=M052DM3

NODE=M052DM3

NODE=M052W

NODE=M052W
 → UNCHECKED ←

NODE=M052215;NODE=M052

 $\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID
31.98 ± 2.63 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

 $\Upsilon(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \Upsilon(1S) \pi^+ \pi^-$	(17.85 ± 0.26) %	
$\Gamma_2 \Upsilon(1S) \pi^0 \pi^0$	(8.6 ± 0.4) %	
$\Gamma_3 \tau^+ \tau^-$	(2.00 ± 0.21) %	
$\Gamma_4 \mu^+ \mu^-$	(1.93 ± 0.17) %	S=2.2
$\Gamma_5 e^+ e^-$	(1.91 ± 0.16) %	
$\Gamma_6 \Upsilon(1S) \pi^0$	< 4 $\times 10^{-5}$	CL=90%
$\Gamma_7 \Upsilon(1S) \eta$	(2.9 ± 0.4) $\times 10^{-4}$	S=2.0
$\Gamma_8 J/\psi(1S)$ anything	< 6 $\times 10^{-3}$	CL=90%
$\Gamma_9 J/\psi(1S) \eta_c$	< 5.4 $\times 10^{-6}$	CL=90%
$\Gamma_{10} J/\psi(1S) \chi_{c0}$	< 3.4 $\times 10^{-6}$	CL=90%
$\Gamma_{11} J/\psi(1S) \chi_{c1}$	< 1.2 $\times 10^{-6}$	CL=90%
$\Gamma_{12} J/\psi(1S) \chi_{c2}$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{13} J/\psi(1S) \eta_c(2S)$	< 2.5 $\times 10^{-6}$	CL=90%
$\Gamma_{14} J/\psi(1S) X(3940)$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{15} J/\psi(1S) X(4160)$	< 2.0 $\times 10^{-6}$	CL=90%
$\Gamma_{16} \chi_{c1}$ anything	(2.2 ± 0.5) $\times 10^{-4}$	
$\Gamma_{17} \chi_{c1}(1P)^0 X_{tetra}$	< 3.67 $\times 10^{-5}$	CL=90%
$\Gamma_{18} \chi_{c2}$ anything	(2.3 ± 0.8) $\times 10^{-4}$	
$\Gamma_{19} \psi(2S) \eta_c$	< 5.1 $\times 10^{-6}$	CL=90%
$\Gamma_{20} \psi(2S) \chi_{c0}$	< 4.7 $\times 10^{-6}$	CL=90%
$\Gamma_{21} \psi(2S) \chi_{c1}$	< 2.5 $\times 10^{-6}$	CL=90%
$\Gamma_{22} \psi(2S) \chi_{c2}$	< 1.9 $\times 10^{-6}$	CL=90%
$\Gamma_{23} \psi(2S) \eta_c(2S)$	< 3.3 $\times 10^{-6}$	CL=90%
$\Gamma_{24} \psi(2S) X(3940)$	< 3.9 $\times 10^{-6}$	CL=90%
$\Gamma_{25} \psi(2S) X(4160)$	< 3.9 $\times 10^{-6}$	CL=90%
$\Gamma_{26} Z_c(3900)^+ Z_c(3900)^-$	< 1.0 $\times 10^{-6}$	CL=90%
$\Gamma_{27} Z_c(4200)^+ Z_c(4200)^-$	< 1.67 $\times 10^{-5}$	CL=90%
$\Gamma_{28} Z_c(3900)^{\pm} Z_c(4200)^{\mp}$	< 7.3 $\times 10^{-6}$	CL=90%
$\Gamma_{29} X(4050)^+ X(4050)^-$	< 1.35 $\times 10^{-5}$	CL=90%

DESIG=4

DESIG=5

DESIG=3

DESIG=1

DESIG=2

DESIG=10

DESIG=6

DESIG=20

DESIG=143

DESIG=144

DESIG=145

DESIG=146

DESIG=147

DESIG=148

DESIG=149

DESIG=157

DESIG=160

DESIG=158

DESIG=150

DESIG=151

DESIG=152

DESIG=153

DESIG=154

DESIG=155

DESIG=156

DESIG=162

DESIG=163

DESIG=164

DESIG=165

Γ_{30}	$X(4250)^+ X(4250)^-$	< 2.67	$\times 10^{-5}$	CL=90%	DESIG=166
Γ_{31}	$X(4050)^{\pm} X(4250)^{\mp}$	< 2.72	$\times 10^{-5}$	CL=90%	DESIG=167
Γ_{32}	$Z_c(4430)^+ Z_c(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%	DESIG=168
Γ_{33}	$X(4055)^{\pm} X(4055)^{\mp}$	< 1.11	$\times 10^{-5}$	CL=90%	DESIG=170
Γ_{34}	$X(4055)^{\pm} Z_c(4430)^{\mp}$	< 2.11	$\times 10^{-5}$	CL=90%	DESIG=171
Γ_{35}	$\overline{^2H}$ anything	(2.78 \pm 0.30)	$\times 10^{-5}$	S=1.2	DESIG=16
Γ_{36}	hadrons	(94 \pm 11)	%		DESIG=101
Γ_{37}	ggg	(58.8 \pm 1.2)	%		DESIG=105
Γ_{38}	γgg	(1.87 \pm 0.28)	%		DESIG=106
Γ_{39}	$\phi K^+ K^-$	(1.6 \pm 0.4)	$\times 10^{-6}$		DESIG=133
Γ_{40}	$\omega \pi^+ \pi^-$	< 2.58	$\times 10^{-6}$	CL=90%	DESIG=134
Γ_{41}	$K^*(892)^0 K^- \pi^+ + c.c.$	(2.3 \pm 0.7)	$\times 10^{-6}$		DESIG=135
Γ_{42}	$\phi f_2'(1525)$	< 1.33	$\times 10^{-6}$	CL=90%	DESIG=136
Γ_{43}	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%	DESIG=137
Γ_{44}	$\rho(770) a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%	DESIG=138
Γ_{45}	$K^*(892)^0 \overline{K}_2^*(1430)^0 + c.c.$	(1.5 \pm 0.6)	$\times 10^{-6}$		DESIG=139
Γ_{46}	$K_1(1270)^{\pm} K^{\mp}$	< 3.22	$\times 10^{-6}$	CL=90%	DESIG=140
Γ_{47}	$K_1(1400)^{\pm} K^{\mp}$	< 8.3	$\times 10^{-7}$	CL=90%	DESIG=141
Γ_{48}	$b_1(1235)^{\pm} \pi^{\mp}$	< 4.0	$\times 10^{-7}$	CL=90%	DESIG=142
Γ_{49}	$\rho \pi$	< 1.16	$\times 10^{-6}$	CL=90%	DESIG=126
Γ_{50}	$\pi^+ \pi^- \pi^0$	< 8.0	$\times 10^{-7}$	CL=90%	DESIG=127
Γ_{51}	$\omega \pi^0$	< 1.63	$\times 10^{-6}$	CL=90%	DESIG=128
Γ_{52}	$\pi^+ \pi^- \pi^0 \pi^0$	(1.30 \pm 0.28)	$\times 10^{-5}$		DESIG=129
Γ_{53}	$K_S^0 K^+ \pi^- + c.c.$	(1.14 \pm 0.33)	$\times 10^{-6}$		DESIG=130
Γ_{54}	$K^*(892)^0 \overline{K}^0 + c.c.$	< 4.22	$\times 10^{-6}$	CL=90%	DESIG=131
Γ_{55}	$K^*(892)^- K^+ + c.c.$	< 1.45	$\times 10^{-6}$	CL=90%	DESIG=132
Γ_{56}	$f_1(1285)$ anything	(2.2 \pm 1.6)	$\times 10^{-3}$		DESIG=159
Γ_{57}	$f_1(1285) X_{tetra}$	< 6.47	$\times 10^{-5}$	CL=90%	DESIG=161
Γ_{58}	Sum of 100 exclusive modes	(2.90 \pm 0.30)	$\times 10^{-3}$		DESIG=121

Radiative decays

Γ_{59}	$\gamma \chi b_1(1P)$	(6.9 \pm 0.4)	%	NODE=M052;CLUMP=A DESIG=8	
Γ_{60}	$\gamma \chi b_2(1P)$	(7.15 \pm 0.35)	%	DESIG=7	
Γ_{61}	$\gamma \chi b_0(1P)$	(3.8 \pm 0.4)	%	DESIG=9	
Γ_{62}	$\gamma f_0(1710)$	< 5.9	$\times 10^{-4}$	CL=90%	DESIG=13
Γ_{63}	$\gamma f_2'(1525)$	< 5.3	$\times 10^{-4}$	CL=90%	DESIG=12
Γ_{64}	$\gamma f_2(1270)$	< 2.41	$\times 10^{-4}$	CL=90%	DESIG=11
Γ_{65}	$\gamma f_J(2220)$			DESIG=14	
Γ_{66}	$\gamma \eta_c(1S)$	< 2.7	$\times 10^{-5}$	CL=90%	DESIG=111
Γ_{67}	$\gamma \chi c_0$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=112
Γ_{68}	$\gamma \chi c_1$	< 3.6	$\times 10^{-6}$	CL=90%	DESIG=113
Γ_{69}	$\gamma \chi c_2$	< 1.5	$\times 10^{-5}$	CL=90%	DESIG=114
Γ_{70}	$\gamma \chi c_1(3872) \rightarrow \pi^+ \pi^- J/\psi$	< 8	$\times 10^{-7}$	CL=90%	DESIG=115
Γ_{71}	$\gamma \chi c_1(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	< 2.4	$\times 10^{-6}$	CL=90%	DESIG=116
Γ_{72}	$\gamma X(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%	DESIG=117
Γ_{73}	$\gamma \chi c_1(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%	DESIG=118
Γ_{74}	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%	DESIG=119
Γ_{75}	$\gamma \eta_b(1S)$	(5.5 \pm 1.1)	$\times 10^{-4}$	S=1.2	DESIG=102
Γ_{76}	$\gamma \eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7	$\times 10^{-6}$	CL=90%	DESIG=124
Γ_{77}	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9	$\times 10^{-6}$	CL=90%	DESIG=125
Γ_{78}	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%	DESIG=103
Γ_{79}	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%	DESIG=108
Γ_{80}	$\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$	< 8.3	$\times 10^{-6}$	CL=90%	DESIG=123

Lepton Family number (*LF*) violating modes

Γ_{81}	$e^\pm \tau^\mp$	<i>LF</i>	< 3.2	$\times 10^{-6}$	CL=90%
Γ_{82}	$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.3	$\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ **CONSTRAINED FIT INFORMATION**

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 11.8$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

 **$\Upsilon(2S) \Gamma(i) \Gamma(e^+ e^-) / \Gamma(\text{total})$**

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_4 \Gamma_5 / \Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
$6.5 \pm 1.5 \pm 1.0$	KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$

NODE=M052218

$\Gamma(\Upsilon(1S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_1 \Gamma_5 / \Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$105.4 \pm 1.0 \pm 4.2$	11.8K	¹ AUBERT	08BP BABR	$10.58 \text{ e}^+ \text{e}^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$

NODE=M052G1
NODE=M052G1

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{36} \Gamma_5 / \Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
0.577 ± 0.009 OUR AVERAGE				
0.581 $\pm 0.004 \pm 0.009$	¹ ROSNER	06	CLEO	$10.0 \text{ e}^+ \text{e}^- \rightarrow \text{hadrons}$
0.552 $\pm 0.031 \pm 0.017$	¹ BARU	96	MD1	$\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$
$0.54 \pm 0.04 \pm 0.02$	¹ JAKUBOWSKI	88	CBAL	$\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$
$0.58 \pm 0.03 \pm 0.04$	² GILES	84B	CLEO	$\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$
$0.60 \pm 0.12 \pm 0.07$	² ALBRECHT	82	DASP	$\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$
$0.54 \pm 0.07 \pm 0.09$	² NICZYPORUK	81C	LENA	$\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$
0.41 ± 0.18	² BOCK	80	CNTR	$\text{e}^+ \text{e}^- \rightarrow \text{hadrons}$

NODE=M052G03
NODE=M052G03

¹ Radiative corrections evaluated following KURAEV 85.
² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

NODE=M052G03;LINKAGE=AU

$\Gamma(e^+ e^-)$				Γ_5
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
0.612 ± 0.011 OUR EVALUATION				

NODE=M052G2;LINKAGE=P
NODE=M052G2;LINKAGE=R **$\Upsilon(2S) \text{ PARTIAL WIDTHS}$**

$\Gamma(e^+ e^-)$				Γ_5
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
0.612 ± 0.011 OUR EVALUATION				

NODE=M052220

 $\Upsilon(2S) \text{ BRANCHING RATIOS}$

$\Gamma(\Upsilon(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}$				Γ_1 / Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.85 ± 0.26 OUR FIT				

NODE=M052W2

NODE=M052W2

→ UNCHECKED ←

Abbreviation MM in the *COMMENT* field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.92 ± 0.26 OUR AVERAGE				

NODE=M052225

16.8 $\pm 1.1 \pm 1.3$ 906k ¹ LEES 11C BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$

17.80 $\pm 0.05 \pm 0.37$ 170k ² LEES 11L BABR $\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$

18.02 $\pm 0.02 \pm 0.61$ 851k ³ BHARI 09 CLEO $e^+ e^- \rightarrow \pi^+ \pi^- \text{MM}$

17.22 $\pm 0.17 \pm 0.75$ 11.8K ⁴ AUBERT 08BP BABR $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \ell^+ \ell^-$

19.2 $\pm 0.2 \pm 1.0$ 52.6k ⁵ ALEXANDER 98 CLE2 $\pi^+ \pi^- \ell^+ \ell^-, \pi^+ \pi^- \text{MM}$

NODE=M052R4

NODE=M052R4

NODE=M052R4

18.1 \pm 0.5 \pm 1.0	11.6k	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^-$ MM
16.9 \pm 4.0		GELPHMAN	85	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
19.1 \pm 1.2 \pm 0.6		BESSON	84	CLEO	$\pi^+ \pi^-$ MM
18.9 \pm 2.6		FONSECA	84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
21 \pm 7	7	NICZYPORUK	81B	LENA	$e^+ e^- \rightarrow \ell^+ \ell^- \pi^+ \pi^-$

¹ LEES 11C reports $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

³ A weighted average of the inclusive and exclusive results.

⁴ Using $B(\gamma(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$, $B(\gamma(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\gamma(2S)) = 0.612 \pm 0.011$ keV.

⁵ Using $B(\gamma(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$ and $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\gamma(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
8.6 \pm 0.4 OUR AVERAGE					
8.43 \pm 0.16 \pm 0.42	38k	¹ BHARI	09	CLEO	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.2 \pm 0.6 \pm 0.8	275	² ALEXANDER	98	CLE2	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
9.5 \pm 1.9 \pm 1.9	25	ALBRECHT	87	ARG	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
8.0 \pm 1.5		GELPHMAN	85	CBAL	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$
10.3 \pm 2.3		FONSECA	84	CUSB	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$

¹ Authors assume $B(\gamma(1S) \rightarrow e^+ e^-) + B(\gamma(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.

² Using $B(\gamma(1S) \rightarrow e^+ e^-) = (2.52 \pm 0.17)\%$ and $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.07)\%$.

$\Gamma(\gamma(1S)\pi^0\pi^0)/\Gamma(\gamma(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				

0.462 \pm 0.037 ¹ BHARI 09 CLEO $e^+ e^- \rightarrow \gamma(2S)$

¹ Not independent of other values reported by BHARI 09.

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
2.00 \pm 0.21 OUR AVERAGE					
2.00 \pm 0.12 \pm 0.18	22k	¹ BESSON	07	CLEO	$e^+ e^- \rightarrow \gamma(2S) \rightarrow \tau^+ \tau^-$
1.7 \pm 1.5 \pm 0.6		HAAS	84B	CLEO	$e^+ e^- \rightarrow \tau^+ \tau^-$

¹ BESSON 07 reports $[\Gamma(\gamma(2S) \rightarrow \tau^+ \tau^-)/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \mu^+ \mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\gamma(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.0193 \pm 0.0017 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.						
0.0203 \pm 0.0003 \pm 0.0008		120k	ADAMS	05	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$
0.0122 \pm 0.0028 \pm 0.0019		¹ KOBEL	92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$	
0.0138 \pm 0.0025 \pm 0.0015		KAARSBERG	89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$	
0.009 \pm 0.006 \pm 0.006		² ALBRECHT	85	ARG	$e^+ e^- \rightarrow \mu^+ \mu^-$	
0.018 \pm 0.008 \pm 0.005		HAAS	84B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.038 90 NICZYPORUK 81C LENA $e^+ e^- \rightarrow \mu^+ \mu^-$

¹ Taking into account interference between the resonance and continuum.

² Re-evaluated using $B(\gamma(1S) \rightarrow \mu^+ \mu^-) = 0.026$.

NODE=M052R4;LINKAGE=ES

NODE=M052R4;LINKAGE=LE

NODE=M052R4;LINKAGE=BH

NODE=M052R4;LINKAGE=AU

NODE=M052R4;LINKAGE=T

NODE=M052R5

NODE=M052R5

NODE=M052R5;LINKAGE=BH

NODE=M052R5;LINKAGE=T

NODE=M052R21

NODE=M052R21

NODE=M052R21;LINKAGE=BH

NODE=M052R3

NODE=M052R3


NODE=M052R3;LINKAGE=BE

NODE=M052R1

NODE=M052R1

NODE=M052R1;LINKAGE=A

NODE=M052R1;LINKAGE=R



$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$	Γ_3/Γ_4
<u>VALUE</u> $1.04 \pm 0.04 \pm 0.05$	<u>EVTS</u> 22k <u>DOCUMENT ID</u> BESSON <u>TECN</u> CLEO <u>COMMENT</u> $e^+e^- \rightarrow \gamma(2S)$

NODE=M052R17
NODE=M052R17

$\Gamma(\gamma(1S)\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ
<u>VALUE (units 10^{-5})</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

NODE=M052R10
NODE=M052R10

• • • We do not use the following data for averages, fits, limits, etc. • • •	
< 4	90 ¹ TAMPONI 13 BELL $e^+e^- \rightarrow \gamma(1S)\pi^0$
< 18	90 ² HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
< 110	90 ALEXANDER 98 CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
< 800	90 LURZ 87 CBAL $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M052R10;LINKAGE=TA

1 TAMPONI 13 reports $[\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^0)/\Gamma_{\text{total}}] / [\Gamma(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$.	Γ_6/Γ
2 Authors assume $B(\gamma(1S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.	

$\Gamma(\gamma(1S)\pi^0)/\Gamma(\gamma(1S)\pi^+\pi^-)$	Γ_6/Γ_1
<u>VALUE (units 10^{-4})</u> <2.3	<u>CL%</u> 90 <u>DOCUMENT ID</u> TAMPONI <u>TECN</u> BELL <u>COMMENT</u> $e^+e^- \rightarrow \gamma(1S)\pi^0$

NODE=M052R09
NODE=M052R09

$\Gamma(\gamma(1S)\eta)/\Gamma_{\text{total}}$	Γ_7/Γ
<u>VALUE (units 10^{-4})</u> 2.9 ± 0.4 OUR FIT Error includes scale factor of 2.0.	<u>CL%</u> <u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

NODE=M052R6
NODE=M052R6

2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.	
2.39 ± 0.31 ± 0.14	112 ¹ LEES 11L BABR $\gamma(2S) \rightarrow \ell^+\ell^-\eta$

NOTFITTED

2.1 $^{+0.7}_{-0.6} \pm 0.3$	14 ² HE 08A CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$
------------------------------	---


• • • We use the following data for averages but not for fits. • • •

3.55 ± 0.32 ± 0.05	241 ³ TAMPONI 13 BELL $e^+e^- \rightarrow \gamma(1S)\eta$
--------------------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90 1,4 AUBERT 08BP BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
< 28	90 ALEXANDER 98 CLE2 $e^+e^- \rightarrow \ell^+\ell^-\eta$
< 50	90 ALBRECHT 87 ARG $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
< 70	90 LURZ 87 CBAL $e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, 3\pi^0)$
< 100	90 BESSON 84 CLEO $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
< 20	90 FONSECA 84 CUSB $e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, \pi^+\pi^-\pi^0)$

- 1 Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.
 2 Authors assume $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$.
 3 TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
 4 Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.



$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.64±0.25 OUR FIT					Error includes scale factor of 2.0.

1.99±0.14±0.11 241 TAMPONI 13 BELL $e^+ e^- \rightarrow \Upsilon(1S)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.17±0.08	1	LEES	11L	BABR	$\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
< 5.2	90	2	AUBERT	08BP	BABR $e^+ e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

1 Not independent of other values reported by LEES 11L.

2 Not independent of other values reported by AUBERT 08BP.

Γ_7/Γ_1

NODE=M052R22
NODE=M052R22

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.13	90	TAMPONI	13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\pi^0$
-------	----	---------	----	------	---

Γ_6/Γ_7

NODE=M052R23
NODE=M052R23

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.006	90	MASCHMANN	90	CBAL $e^+ e^- \rightarrow \text{hadrons}$

Γ_8/Γ

NODE=M052R16
NODE=M052R16

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10⁻⁶	90	YANG	14	BELL $e^+ e^- \rightarrow J/\psi X$

Γ_9/Γ

NODE=M052R53
NODE=M052R53

$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 × 10⁻⁶	90	YANG	14	BELL $e^+ e^- \rightarrow J/\psi X$

Γ_{10}/Γ

NODE=M052R54
NODE=M052R54

$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10⁻⁶	90	YANG	14	BELL $e^+ e^- \rightarrow J/\psi X$

Γ_{11}/Γ

NODE=M052R55
NODE=M052R55

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$	Γ_{12}/Γ	NODE=M052R56 NODE=M052R56
<u>VALUE</u> $<2.0 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$	Γ_{13}/Γ	NODE=M052R57 NODE=M052R57
<u>VALUE</u> $<2.5 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$	Γ_{14}/Γ	NODE=M052R58 NODE=M052R58
<u>VALUE</u> $<2.0 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow J/\psi X$
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$	Γ_{15}/Γ	NODE=M052R59 NODE=M052R59
<u>VALUE</u> $<2.0 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow J/\psi X$
$\Gamma(\chi_{c1} \text{anything})/\Gamma_{\text{total}}$	Γ_{16}/Γ	NODE=M052R00 NODE=M052R00
<u>VALUE (units 10^{-4})</u> $2.24 \pm 0.44 \pm 0.20$	<u>EVTS</u> 376	<u>DOCUMENT ID</u> JIA <u>TECN</u> BELL <u>COMMENT</u> $\gamma(2S) \rightarrow \gamma J/\psi(1S)$
$\Gamma(\chi_{c1}(1P)^0 X_{\text{tetra}})/\Gamma_{\text{total}}$	Γ_{17}/Γ	NODE=M052R69 NODE=M052R69
<u>VALUE</u> $<36.7 \times 10^{-6}$	<u>CL%</u> 90	¹ <u>DOCUMENT ID</u> JIA <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \text{hadrons}$
¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 36.7×10^{-6} .		
$\Gamma(\chi_{c2} \text{anything})/\Gamma_{\text{total}}$	Γ_{18}/Γ	NODE=M052R67 NODE=M052R67
<u>VALUE (units 10^{-4})</u> $2.28 \pm 0.73 \pm 0.34$		<u>DOCUMENT ID</u> JIA <u>TECN</u> BELL <u>COMMENT</u> $\gamma(2S) \rightarrow \gamma J/\psi(1S)$
$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$	Γ_{19}/Γ	NODE=M052R60 NODE=M052R60
<u>VALUE</u> $<5.1 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$	Γ_{20}/Γ	NODE=M052R61 NODE=M052R61
<u>VALUE</u> $<4.7 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$	Γ_{21}/Γ	NODE=M052R62 NODE=M052R62
<u>VALUE</u> $<2.5 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$	Γ_{22}/Γ	NODE=M052R63 NODE=M052R63
<u>VALUE</u> $<1.9 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$	Γ_{23}/Γ	NODE=M052R64 NODE=M052R64
<u>VALUE</u> $<3.3 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$	Γ_{24}/Γ	NODE=M052R65 NODE=M052R65
<u>VALUE</u> $<3.9 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$	Γ_{25}/Γ	NODE=M052R66 NODE=M052R66
<u>VALUE</u> $<3.9 \times 10^{-6}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> YANG <u>TECN</u> BELL <u>COMMENT</u> $e^+ e^- \rightarrow \psi(2S) X$
$\Gamma(Z_c(3900)^+ Z_c(3900)^-)/\Gamma_{\text{total}}$	Γ_{26}/Γ	NODE=M052R71 NODE=M052R71
<u>VALUE</u> $<1.0 \times 10^{-6}$	<u>CL%</u> 90	¹ <u>DOCUMENT ID</u> JIA <u>TECN</u> BELL <u>COMMENT</u> $\gamma(2S) \rightarrow J/\psi \pi^\pm X$

¹ Assuming $B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$.

NODE=M052R71;LINKAGE=A

$\Gamma(Z_c(4200)^+ Z_c(4200)^-)/\Gamma_{\text{total}}$					Γ_{27}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<16.7 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(1S) \rightarrow J/\psi \pi^\pm X$	NODE=M052R72 NODE=M052R72
¹ Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$					
$\Gamma(Z_c(3900)^\pm Z_c(4200)^\mp)/\Gamma_{\text{total}}$					Γ_{28}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.3 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow J/\psi \pi^\pm X$	NODE=M052R73 NODE=M052R73
¹ Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm)$.					
$\Gamma(X(4050)^+ X(4050)^-)/\Gamma_{\text{total}}$					Γ_{29}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<13.5 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$	NODE=M052R74 NODE=M052R74
¹ Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$					
$\Gamma(X(4250)^+ X(4250)^-)/\Gamma_{\text{total}}$					Γ_{30}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<26.7 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$	NODE=M052R75 NODE=M052R75
¹ Assuming $B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$					
$\Gamma(X(4050)^\pm X(4250)^\mp)/\Gamma_{\text{total}}$					Γ_{31}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<27.2 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$	NODE=M052R76 NODE=M052R76
¹ Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$					
$\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$					Γ_{32}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<20.3 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$	NODE=M052R77 NODE=M052R77
¹ Assuming $B(Z_c(4430)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$					
$\Gamma(X(4055)^\pm X(4055)^\mp)/\Gamma_{\text{total}}$					Γ_{33}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<11.1 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$	NODE=M052R79 NODE=M052R79
¹ Assuming $B(X(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$					
$\Gamma(X(4055)^\pm Z_c(4430)^\mp)/\Gamma_{\text{total}}$					Γ_{34}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<21.1 \times 10^{-6}$	90	1 JIA	18	BELL $\gamma(2S) \rightarrow \psi(2S) \pi^\pm X$	NODE=M052R80 NODE=M052R80
¹ Assuming $B(X(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(Z_c(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$					
$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$					Γ_{35}/Γ
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.78^{+0.30}_{-0.26} OUR AVERAGE				Error includes scale factor of 1.2.	
$2.64 \pm 0.11^{+0.26}_{-0.21}$		LEES	14G BABR	$e^+ e^- \rightarrow \overline{2H} X$	
$3.37 \pm 0.50 \pm 0.25$	58	ASNER	07 CLEO	$e^+ e^- \rightarrow \overline{2H} X$	
$\Gamma(ggg)/\Gamma_{\text{total}}$					Γ_{37}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
58.8^{+1.2}_{-1.2}	6M	1 BESSON	06A CLEO	$\gamma(2S) \rightarrow \text{hadrons}$	NODE=M052R01 NODE=M052R01
¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+ \pi^- \gamma(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0 \pi^0 \gamma(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.					
$\Gamma(\gamma gg)/\Gamma(ggg)$					Γ_{38}/Γ_{37}
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.18^{+0.04}_{-0.47}	6M	BESSON	06A CLEO	$\gamma(2S) \rightarrow (\gamma +) \text{ hadrons}$	NODE=M052R03 NODE=M052R03

$\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$				Γ_{39}/Γ	NODE=M052R43 NODE=M052R43
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.58 \pm 0.33 \pm 0.18$	58	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$	
$\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_{40}/Γ	NODE=M052R44 NODE=M052R44
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<2.58	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{41}/Γ	NODE=M052R45 NODE=M052R45
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.32 \pm 0.40 \pm 0.54$	135	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$				Γ_{42}/Γ	NODE=M052R46 NODE=M052R46
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.33	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(K^+ K^-)$	
$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$				Γ_{43}/Γ	NODE=M052R47 NODE=M052R47
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.57	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$				Γ_{44}/Γ	NODE=M052R48 NODE=M052R48
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.88	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{45}/Γ	NODE=M052R49 NODE=M052R49
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$1.53 \pm 0.52 \pm 0.19$	32	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(K_1(1270)^{\pm} K^{\mp})/\Gamma_{\text{total}}$				Γ_{46}/Γ	NODE=M052R50 NODE=M052R50
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<3.22	90	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(K_1(1400)^{\pm} K^{\mp})/\Gamma_{\text{total}}$				Γ_{47}/Γ	NODE=M052R51 NODE=M052R51
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.83	90	SHEN	12A BELL	$\gamma(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	
$\Gamma(b_1(1235)^{\pm} \pi^{\mp})/\Gamma_{\text{total}}$				Γ_{48}/Γ	NODE=M052R52 NODE=M052R52
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.40	90	SHEN	12A BELL	$\gamma(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$	
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$				Γ_{49}/Γ	NODE=M052R27 NODE=M052R27
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.16	90	SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$	
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$				Γ_{50}/Γ	NODE=M052R28 NODE=M052R28
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<0.80	90	SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0$	
$\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$				Γ_{51}/Γ	NODE=M052R29 NODE=M052R29
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<1.63	90	SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	
$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$				Γ_{52}/Γ	NODE=M052R30 NODE=M052R30
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$13.0 \pm 1.9 \pm 2.1$	261 ± 37	SHEN	13 BELL	$\gamma(2S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$	
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$				Γ_{53}/Γ	NODE=M052R40 NODE=M052R40
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$1.14 \pm 0.30 \pm 0.13$	40 ± 10	SHEN	13 BELL	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.2	90	1 DOBBS	12A	$\gamma(2S) \rightarrow K_S^0 K^- \pi^+$	

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M052R40;LINKAGE=DO

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.22	90	SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 Γ_{54}/Γ NODE=M052R41
NODE=M052R41 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.45	90	SHEN	13	BELL $\gamma(2S) \rightarrow K_S^0 K^- \pi^+$

 Γ_{55}/Γ NODE=M052R42
NODE=M052R42 $\Gamma(f_1(1285) \text{anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±1.50±0.63	2.9k	JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

 Γ_{56}/Γ NODE=M052R68
NODE=M052R68 $\Gamma(f_1(1285) X_{\text{tetra}})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<64.7 × 10 ⁻⁶	90	1 JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

 Γ_{57}/Γ NODE=M052R70
NODE=M052R70

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 7.8×10^{-6} to 64.7×10^{-6} .

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
0.29±0.03	1,2 DOBBS	12A	$\gamma(2S) \rightarrow \text{hadrons}$

 Γ_{58}/Γ NODE=M052R08
NODE=M052R08

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.069 ±0.004 OUR AVERAGE				
0.0693±0.0012±0.0041	407k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.069 ±0.005 ±0.009		EDWARDS	99	CLE2 $\gamma(2S) \rightarrow \gamma \chi(1P)$
0.091 ±0.018 ±0.022		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.065 ±0.007 ±0.012		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
0.080 ±0.017 ±0.016		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.059 ±0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

 Γ_{59}/Γ NODE=M052R8
NODE=M052R8 $\Gamma(\gamma \chi_{b2}(1P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0715±0.0035 OUR AVERAGE				
0.0724±0.0011±0.0040	410k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.074 ±0.005 ±0.008		EDWARDS	99	CLE2 $\gamma(2S) \rightarrow \gamma \chi(1P)$
0.098 ±0.021 ±0.024		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.058 ±0.007 ±0.010		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
0.102 ±0.018 ±0.021		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.061 ±0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

 Γ_{60}/Γ NODE=M052R7
NODE=M052R7 $\Gamma(\gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.038 ±0.004 OUR AVERAGE				
0.0375±0.0012±0.0047	198k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.034 ±0.005 ±0.006		EDWARDS	99	CLE2 $\gamma(2S) \rightarrow \gamma \chi(1P)$
0.064 ±0.014 ±0.016		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.036 ±0.008 ±0.009		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
0.044 ±0.023 ±0.009		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.035 ±0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

 Γ_{61}/Γ NODE=M052R9
NODE=M052R9 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<59	90	¹ ALBRECHT	89	ARG $\gamma(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				

 Γ_{62}/Γ NODE=M052R13
NODE=M052R13

¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.

² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+ \pi^-$.

OCCUR=2

NODE=M052R13;LINKAGE=M
NODE=M052R13;LINKAGE=N

$\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$					Γ_{63}/Γ
<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<53	90	¹ ALBRECHT	89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$
¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$.					
$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$					Γ_{64}/Γ
<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
<24.1	90	¹ ALBRECHT	89	ARG	$\gamma(2S) \rightarrow \gamma \pi^+ \pi^-$
¹ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.					
$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$					Γ_{65}/Γ
<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.8	90	¹ ALBRECHT	89	ARG	$\gamma(2S) \rightarrow \gamma K^+ K^-$
¹ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.					
$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$					Γ_{66}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 2.7×10^{-5}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$					Γ_{67}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 1.0×10^{-4}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$					Γ_{68}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 3.6×10^{-6}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$					Γ_{69}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 1.5×10^{-5}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi)/\Gamma_{\text{total}}$					Γ_{70}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 0.8×10^{-6}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \chi_{c1}(3872) \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$					Γ_{71}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 2.4×10^{-6}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma X(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$					Γ_{72}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 2.8×10^{-6}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$					Γ_{73}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 1.2×10^{-6}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$					Γ_{74}/Γ
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
< 1.3×10^{-6}	90	WANG	11B	BELL	$\gamma(2S) \rightarrow \gamma X$
$\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$					Γ_{75}/Γ
<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.5^{+1.1}_{-0.9} OUR AVERAGE Error includes scale factor of 1.2.					
$6.1^{+0.6}_{-0.7} {}^{+0.9}_{-0.6}$	29k	FULSAM	18	BELL	$\gamma(2S) \rightarrow \gamma X$
$3.9 \pm 1.1 {}^{+1.1}_{-0.9}$	$13 \pm 5k$	¹ AUBERT	09AQ BABR		$\gamma(2S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<21	90	LEES	11J	BABR	$\gamma(2S) \rightarrow X\gamma$
< 8.4	90	¹ BONVICINI	10	CLEO	$\gamma(2S) \rightarrow \gamma X$
< 5.1	90	² ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$
¹ Assuming $\Gamma_{\eta_b}(1S) = 10$ MeV.					
² Superseded by BONVICINI 10.					

$\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{76}/Γ
$<3.7 \times 10^{-6}$	90	SANDILYA	13	BELL $\gamma(2S) \rightarrow \gamma$ hadrons	NODE=M052R25 NODE=M052R25

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{77}/Γ
< 4.9	90		SANDILYA	13	BELL $\gamma(2S) \rightarrow \gamma$ hadrons	NODE=M052R26 NODE=M052R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

$46.2^{+29.7}_{-14.2} \pm 10.6$ 10 ¹ DOBBS 12 $\gamma(2S) \rightarrow \gamma$ hadrons

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$

($1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{78}/Γ
<1.95	95	ROSNER	07A	CLEO $e^+ e^- \rightarrow \gamma X$	NODE=M052R19 NODE=M052R19 NODE=M052R19

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$

($0.3 \text{ GeV} < m_{A^0} < 7 \text{ GeV}$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{79}/Γ
$<8 \times 10^{-5}$	90	¹ LEES	11H	BABR $\gamma(2S) \rightarrow \gamma$ hadrons	NODE=M052R06 NODE=M052R06 NODE=M052R06

¹ For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{80}/Γ
<8.3	90	¹ AUBERT	09Z	BABR $e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$	NODE=M052R24 NODE=M052R24

¹ For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from 0.26×10^{-6} to 8.3×10^{-6} .

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{81}/Γ
<3.2	90	LEES	10B	BABR $e^+ e^- \rightarrow e^\pm \tau^\mp$	NODE=M052230

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{82}/Γ
< 3.3	90	LEES	10B	BABR $e^+ e^- \rightarrow \mu^\pm \tau^\mp$	NODE=M052R20 NODE=M052R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14.4 95 LOVE 08A CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

$\gamma(2S)$ Cross-Particle Branching Ratios

 $B(\gamma(2S) \rightarrow \pi^+ \pi^-) \times B(\gamma(3S) \rightarrow \gamma(2S)X)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
$1.78 \pm 0.02 \pm 0.11$	906k	LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$	NODE=M052R05 NODE=M052R05

$\gamma(2S)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)
SANDILYA	13	PR D111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PR D109 082001	S. Dobbs <i>et al.</i>	(BELLE Collab.)
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>	(BELLE Collab.)
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)

NODE=M052

REFID=59535
REFID=58949
REFID=57635
REFID=58318
REFID=55939
REFID=56345
REFID=55590
REFID=55395
REFID=54919
REFID=54288
REFID=54746
REFID=54314
REFID=16775
REFID=53877
REFID=53936
REFID=53938
REFID=53939

BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE e ⁺ e ⁻ Physics 412	W. Buchmueler, S. Cooper Editors: A. Ali and P. Soeding, World Scientific, Singapore	(HANN, DESY, MIT)	REFID=40034
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC	REFID=40742
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)	REFID=40021
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22334
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
GELPHMAN	85	PR D32 2893	D. Gelfphman <i>et al.</i>	(Crystal Ball Collab.)	REFID=22336
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>		REFID=22327; ERROR=18
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)	REFID=22329
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22332
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22319
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22318
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264

$\Upsilon_2(1D)$

$I^G(J^{PC}) = 0^-(2^- -)$

was $\Upsilon(1D)$

First observed by BONVICINI 04 in the decay to $\gamma\gamma \Upsilon(1S)$ and confirmed by DEL-AMO-SANCHEZ 10R in the decay to $\pi^+ \pi^- \Upsilon(1S)$.

Data consistent with $J^P = 2^-$. The states with $J = 1$ and 3 also possibly seen, but need confirmation.

$\Upsilon_2(1D)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10163.7±1.4 OUR AVERAGE				Error includes scale factor of 1.7.
10164.5±0.8±0.5		DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
10161.1±0.6±1.6	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Upsilon_2(1D)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma\gamma \Upsilon(1S)$	seen
$\Gamma_2 \quad \gamma\chi_{bJ}(1P)$	seen
$\Gamma_3 \quad \eta \Upsilon(1S)$	not seen
$\Gamma_4 \quad \pi^+ \pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$

$\Upsilon_2(1D)$ BRANCHING RATIOS

$\Gamma(\eta \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_3/Γ_1
VALUE	CL%
<0.25	90
	BONVICINI 04 CLE3 $\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Gamma(\pi^+ \pi^- \Upsilon(1S))/\Gamma_{\text{total}}$	Γ_4/Γ_1
VALUE (units 10^{-2})	DOCUMENT ID TECN COMMENT
0.66^{+0.15}_{-0.14}±0.06	¹ DEL-AMO-SA..10R BABR $\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$

¹ Using theoretical predictions for $B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon_2(1D))$.

$\Gamma(\pi^+ \pi^- \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_4/Γ_1
VALUE	CL% DOCUMENT ID TECN COMMENT
<1.2	90 ² BONVICINI 04 CLE3 $\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

² Assuming $J = 2$.

$\Upsilon_2(1D)$ REFERENCES

DEL-AMO-SA... 10R PR D82 111102
BONVICINI 04 PR D70 032001

P. del Amo Sanchez *et al.*
G. Bonvicini *et al.*

(BABAR Collab.)
(CLEO Collab.)

NODE=M177

NODE=M177

NODE=M177M

NODE=M177M

NODE=M177215;NODE=M177

DESIG=1;OUR EVAL; \rightarrow UNCHECKED
DESIG=2;OUR EVAL; \rightarrow UNCHECKED
DESIG=3;OUR EVAL; \rightarrow UNCHECKED
DESIG=4

NODE=M177225

NODE=M177R01
NODE=M177R01

NODE=M177R03
NODE=M177R03

NODE=M177R03;LINKAGE=DE

NODE=M177R02
NODE=M177R02

NODE=M177R02;LINKAGE=BO

NODE=M177

REFID=53634
REFID=49759

$\chi_{b0}(2P)$

$I^G(JPC) = 0^+(0^{++})$
J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b0}(2P)$ MASS

VALUE (MeV)	DOCUMENT ID
10232.5 ± 0.4 ± 0.5 OUR EVALUATION	From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

$m_{\chi_{b1}(2P)} = m_{\chi_{b0}(2P)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23.8 ± 1.7	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$


γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
121.9 ± 0.4 OUR EVALUATION				Treating systematic errors as correlated
122.2 ± 0.5 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
121.55 ± 0.16 ± 0.46		ARTUSO	05	CLEO $\Upsilon(3S) \rightarrow \gamma X$
123.0 ± 0.8	4959	¹ HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma X$
124.6 ± 1.4	17	² HEINTZ	92	CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
122.3 ± 0.3 ± 0.6	9903	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma X$

¹A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

²A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

WEIGHTED AVERAGE
122.2 ± 0.5 (Error scaled by 1.4)



$\chi_{b0}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \gamma \Upsilon(2S)$	$(1.38 \pm 0.30) \%$	
$\Gamma_2 \gamma \Upsilon(1S)$	$(3.8 \pm 1.7) \times 10^{-3}$	
$\Gamma_3 D^0 X$	$< 8.2 \%$	90%
$\Gamma_4 \pi^+ \pi^- K^+ K^- \pi^0$	$< 3.4 \times 10^{-5}$	90%
$\Gamma_5 2\pi^+ \pi^- K^- K_S^0$	$< 5 \times 10^{-5}$	90%
$\Gamma_6 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$< 2.2 \times 10^{-4}$	90%
$\Gamma_7 2\pi^+ 2\pi^- 2\pi^0$	$< 2.4 \times 10^{-4}$	90%
$\Gamma_8 2\pi^+ 2\pi^- K^+ K^-$	$< 1.5 \times 10^{-4}$	90%

NODE=M079

NODE=M079

→ UNCHECKED ←

NODE=M079M2

NODE=M079M2

NODE=M079DM

NODE=M079DM

→ UNCHECKED ←

OCCUR=2

NODE=M079DM;LINKAGE=A

NODE=M079DM;LINKAGE=B

NODE=M079215;NODE=M079

DESIG=2

DESIG=1

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

Γ_9	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.2	$\times 10^{-4}$	90%	DESIG=9
Γ_{10}	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 1.1	$\times 10^{-3}$	90%	DESIG=10
Γ_{11}	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=11
Γ_{12}	$3\pi^+ 3\pi^-$	< 7	$\times 10^{-5}$	90%	DESIG=12
Γ_{13}	$3\pi^+ 3\pi^- 2\pi^0$	< 1.2	$\times 10^{-3}$	90%	DESIG=13
Γ_{14}	$3\pi^+ 3\pi^- K^+ K^-$	< 1.5	$\times 10^{-4}$	90%	DESIG=14
Γ_{15}	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=15
Γ_{16}	$4\pi^+ 4\pi^-$	< 1.7	$\times 10^{-4}$	90%	DESIG=16
Γ_{17}	$4\pi^+ 4\pi^- 2\pi^0$	< 6	$\times 10^{-4}$	90%	DESIG=17

$\chi_{b0}(2P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
1.38±0.30 OUR AVERAGE					
$1.31 \pm 0.27^{+0.13}_{-0.12}$	3,4	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
$3.6 \pm 1.6 \pm 0.3$	3,5	HEINTZ	92	CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.8	90	6 LEES	11J	BABR $\Upsilon(3S) \rightarrow X \gamma$	
<8.9	90	7 CRAWFORD	92B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	

3 Assuming $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

4 LEES 14M reports $[\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] = (7.7 \pm 1.6) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

5 Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91.

6 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$.

7 Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) < 1.19 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P) \gamma) = 0.049$.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.38±0.17 OUR AVERAGE					
$0.36 \pm 0.17 \pm 0.03$	8,9,10	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
$0.9 \pm 0.7 \pm 0.1$	9,11	HEINTZ	92	CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.2	90	12 LEES	11J	BABR $\Upsilon(3S) \rightarrow X \gamma$	
<2.5	90	13 CRAWFORD	92B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	

8 LEES 14M quotes $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (2.1 \pm 1.0) \times 10^{-4}$ combining the results from $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ samples with and without photon conversions.

9 Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

10 LEES 14M reports $[\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] = (2.1 \pm 1.0) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

11 Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91.

12 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$.

13 Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) < 0.63 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P) \gamma) = 0.049$.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
$<8.2 \times 10^{-2}$	90	14,15 BRIERE	08	CLEO $\Upsilon(3S) \rightarrow \gamma D^0 X$	

14 For $p_{D^0} > 2.5$ GeV/c.

15 The authors also present their result as $(4.1 \pm 3.0 \pm 0.4) \times 10^{-2}$.

NODE=M079220

NODE=M079R2
NODE=M079R2

NODE=M079R2;LINKAGE=D
NODE=M079R2;LINKAGE=E

NODE=M079R2;LINKAGE=C

NODE=M079R2;LINKAGE=LE

NODE=M079R2;LINKAGE=B

NODE=M079R1
NODE=M079R1

NODE=M079R1;LINKAGE=D

NODE=M079R1;LINKAGE=E
NODE=M079R1;LINKAGE=F

NODE=M079R1;LINKAGE=C

NODE=M079R1;LINKAGE=LE

NODE=M079R1;LINKAGE=B

NODE=M079R01
NODE=M079R01

NODE=M079R01;LINKAGE=BR
NODE=M079R01;LINKAGE=RI

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.34	90	16 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$
16 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_4/Γ NODE=M079R02
NODE=M079R02 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	17 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$
17 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_5/Γ NODE=M079R03
NODE=M079R03 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	18 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$
18 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_6/Γ NODE=M079R04
NODE=M079R04 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.4	90	19 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$
19 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_7/Γ NODE=M079R05
NODE=M079R05 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	20 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$
20 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 9 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_8/Γ NODE=M079R06
NODE=M079R06 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	21 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$
21 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_9/Γ NODE=M079R07
NODE=M079R07 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	22 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$
22 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 63 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{10}/Γ NODE=M079R08
NODE=M079R08 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	23 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$
23 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 39 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{11}/Γ NODE=M079R09
NODE=M079R09 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	24 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+3\pi^-$
24 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P))] < 4 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{12}/Γ NODE=M079R10
NODE=M079R10

NODE=M079R10;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	25 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
25 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 7.2×10^{-6} which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{13}/Γ NODE=M079R11
NODE=M079R11 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	26 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
26 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 9×10^{-6} which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{14}/Γ NODE=M079R12
NODE=M079R12 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	27 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
27 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 43×10^{-6} which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{15}/Γ NODE=M079R13
NODE=M079R13 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	28 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
28 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 10×10^{-6} which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{16}/Γ NODE=M079R14
NODE=M079R14 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	29 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
29 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 38×10^{-6} which we divide by our best value $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

 Γ_{17}/Γ NODE=M079R15
NODE=M079R15 $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma \times \Gamma_{22}^{T(3S)}/\Gamma^{T(3S)}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	30 LEES	11J BABR	$\gamma(3S) \rightarrow X\gamma$
30 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \gamma(1S))/\Gamma_{\text{total}} \times \Gamma(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \gamma(1S)) < 1.2\%$ using $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.				

NODE=M079B01
NODE=M079B01 $B(\chi_{b0}(2P) \rightarrow \gamma \gamma(1S)) \times B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\gamma(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.4 ± 0.9 OUR AVERAGE			

NODE=M079B01;LINKAGE=LE

1.7 + 1.5 + 0.1 - 1.4 - 1.2 1.3 ± 1.0 ± 0.3	31 LEES	14M BABR	$\gamma(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
	32 HEINTZ	92 CSB2	$\gamma(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M079A02
NODE=M079A0231 From a sample of $\gamma(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.32 Calculated by us. HEINTZ 92 quotes $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \gamma(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$.NODE=M079A02;LINKAGE=A
NODE=M079A02;LINKAGE=K $[B(\chi_{b0}(2P) \rightarrow \gamma \gamma(1S)) \times B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \gamma(1S)) \times B(\gamma(3S) \rightarrow \gamma \chi_{b1}(2P))]$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.71 ± 0.80	33 LEES	14M BABR	$\gamma(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M079A00
NODE=M079A0033 From a sample of $\gamma(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ without converted photons.

NODE=M079A00;LINKAGE=A

 $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \gamma(2S))/\Gamma_{\text{total}} \times \Gamma(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_{22}^{T(3S)}/\Gamma^{T(3S)}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	34 LEES	11J BABR	$\gamma(3S) \rightarrow X\gamma$

NODE=M079B02
NODE=M079B0234 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \gamma(2S))/\Gamma_{\text{total}} \times \Gamma(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \gamma(2S)) < 2.8\%$ using $B(\gamma(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.

NODE=M079B02;LINKAGE=LE

$B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.4±1.6 OUR AVERAGE			
6.6 ± 4.9 ± 2.0	35 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
4.0 ± 4.0 ± 0.3	36 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
4.0 ± 1.7 ± 0.3			35 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.
			36 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$.

 $[B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
3.31±0.56			
37 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
37 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ without converted photons.			

 $\chi_{b0}(2P)$ REFERENCES

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

 $\chi_{b1}(2P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b1}(2P)$ MASS

VALUE (MeV)	DOCUMENT ID	COMMENT		
10255.46±0.22±0.50 OUR EVALUATION		From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV		

 $m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23.5±0.7±0.7	1 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$
1 From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.			

 γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
99.26±0.22 OUR EVALUATION				Treating systematic errors as correlated
99.53±0.23 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
99.15±0.07±0.25				
99 ± 1	169	ARTUSO 05	CLEO	$\Upsilon(3S) \rightarrow \gamma X$
100.1 ± 0.4	11147	CRAWFORD 92B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
100.2 ± 0.5	223	2 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
99.5 ± 0.1 ± 0.5	25759	3 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
		MORRISON 91	CLE2	$e^+ e^- \rightarrow \gamma X$

NODE=M079A03
NODE=M079A03

NODE=M079A03;LINKAGE=A
NODE=M079A03;LINKAGE=B

NODE=M079A01
NODE=M079A01

NODE=M079A01;LINKAGE=A

NODE=M079

REFID=56343
REFID=53936
REFID=52574
REFID=52577
REFID=50454
REFID=43177
REFID=43604
REFID=41580
REFID=41634
REFID=41586

NODE=M080

NODE=M080

NODE=M080M

NODE=M080M
→ UNCHECKED ←

NODE=M080M2

NODE=M080M2

NODE=M080M2;LINKAGE=A

NODE=M080DM

NODE=M080DM
→ UNCHECKED ←


OCCUR=2

²A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

³A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M080DM;LINKAGE=A

NODE=M080DM;LINKAGE=B



γ energy in $\eta(3S)$ decay (MeV)

$\chi_{b1}(2P)$ DECAY MODES

NODE=M080215;NODE=M080

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 \omega \eta(1S)$	(1.63 \pm 0.40) %	DESIG=3
$\Gamma_2 \gamma \eta(2S)$	(18.1 \pm 1.9) %	DESIG=2
$\Gamma_3 \gamma \eta(1S)$	(9.9 \pm 1.0) %	DESIG=1
$\Gamma_4 \pi\pi\chi_{b1}(1P)$	(9.1 \pm 1.3) \times 10 ⁻³	DESIG=4
$\Gamma_5 D^0 X$	(8.8 \pm 1.7) %	DESIG=5
$\Gamma_6 \pi^+\pi^-K^+K^-\pi^0$	(3.1 \pm 1.0) \times 10 ⁻⁴	DESIG=6
$\Gamma_7 2\pi^+\pi^-K^-K_S^0$	(1.1 \pm 0.5) \times 10 ⁻⁴	DESIG=7
$\Gamma_8 2\pi^+\pi^-K^-K_S^0 2\pi^0$	(7.7 \pm 3.2) \times 10 ⁻⁴	DESIG=8
$\Gamma_9 2\pi^+2\pi^-2\pi^0$	(5.9 \pm 2.0) \times 10 ⁻⁴	DESIG=9
$\Gamma_{10} 2\pi^+2\pi^-K^+K^-$	(10 \pm 4) \times 10 ⁻⁵	DESIG=10
$\Gamma_{11} 2\pi^+2\pi^-K^+K^-\pi^0$	(5.5 \pm 1.8) \times 10 ⁻⁴	DESIG=11
$\Gamma_{12} 2\pi^+2\pi^-K^+K^-2\pi^0$	(10 \pm 4) \times 10 ⁻⁴	DESIG=12
$\Gamma_{13} 3\pi^+2\pi^-K^-K_S^0\pi^0$	(6.7 \pm 2.6) \times 10 ⁻⁴	DESIG=13
$\Gamma_{14} 3\pi^+3\pi^-$	(1.2 \pm 0.4) \times 10 ⁻⁴	DESIG=14
$\Gamma_{15} 3\pi^+3\pi^-2\pi^0$	(1.2 \pm 0.4) \times 10 ⁻³	DESIG=15
$\Gamma_{16} 3\pi^+3\pi^-K^+K^-$	(2.0 \pm 0.8) \times 10 ⁻⁴	DESIG=16
$\Gamma_{17} 3\pi^+3\pi^-K^+K^-\pi^0$	(6.1 \pm 2.2) \times 10 ⁻⁴	DESIG=17
$\Gamma_{18} 4\pi^+4\pi^-$	(1.7 \pm 0.6) \times 10 ⁻⁴	DESIG=18
$\Gamma_{19} 4\pi^+4\pi^-2\pi^0$	(1.9 \pm 0.7) \times 10 ⁻³	DESIG=19

$\chi_{b1}(2P)$ BRANCHING RATIOS

NODE=M080220

$\Gamma(\omega \eta(1S))/\Gamma_{\text{total}}$	Γ_1/Γ
1.63 \pm 0.35 \pm 0.16 -0.31 -0.15	32.6 \pm 6.9 ⁴ CRONIN-HEN..04 CLE3 $\eta(3S) \rightarrow \gamma\omega\eta(1S)$

NODE=M080R3
NODE=M080R3

⁴ Using $B(\eta(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.3 \pm 0.6)\%$ and $B(\eta(1S) \rightarrow e^+e^-) = 2 B(\eta(1S) \rightarrow \mu^+\mu^-) = 2 (2.48 \pm 0.06)\%$.

NODE=M080R3;LINKAGE=CR

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$						Γ_2/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M080R2	NODE=M080R2
0.181±0.019 OUR AVERAGE							
0.211±0.017±0.019	5,6,7	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$			
0.190±0.018±0.017	4.3k	8 LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$			
0.206±0.035±0.019		5,9 CRAWFORD	92B	CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$			
0.132±0.018±0.012		5,10 HEINTZ	92	CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$			
5 Assuming $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$.							
6 LEES 14M quotes $\Gamma(\chi_b1(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P))/\Gamma_{\text{total}} = (2.66 \pm 0.22)\%$ combining the results from $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ samples with and without photon conversions.							
7 LEES 14M reports $[\Gamma(\chi_b1(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P))] = (2.66 \pm 0.22) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.							
8 LEES 11J reports $[\Gamma(\chi_b1(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.							
9 CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) \times B(\chi_b1(2P) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \ell^+\ell^-) = (10.23 \pm 1.20 \pm 1.26) 10^{-4}$.							
10 Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) \times B(\chi_b1(2P) \rightarrow \gamma \Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91.							
$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$						Γ_3/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M080R1	NODE=M080R1
0.099±0.010 OUR AVERAGE							
0.107±0.006±0.010	11,12,13	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$			
0.098±0.005±0.009	15k	14 LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$			
0.103±0.023±0.009		11,15 CRAWFORD	92B	CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$			
0.075±0.010±0.007		11,16 HEINTZ	92	CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$			
11 Assuming $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.							
12 LEES 14M quotes $\Gamma(\chi_b1(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P))/\Gamma_{\text{total}} = (13.48 \pm 0.72) \times 10^{-3}$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with and without converted photons.							
13 LEES 14M reports $[\Gamma(\chi_b1(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P))] = (13.48 \pm 0.72) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.							
14 LEES 11J reports $[\Gamma(\chi_b1(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.							
15 CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) \times B(\chi_b1(2P) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (6.47 \pm 1.12 \pm 0.82) 10^{-4}$.							
16 Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_b1(2P)) \times B(\chi_b1(2P) \rightarrow \gamma \Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91.							
$\Gamma(\pi\pi\chi_b1(1P))/\Gamma_{\text{total}}$						Γ_4/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M080R4	NODE=M080R4
9.1±1.3 OUR AVERAGE							
9.2±1.1±0.8	31k	17 LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$			
8.6±2.3±2.1		18 CAWLFIELD	06	CLE3 $\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$			
17 LEES 11C measures $B(\Upsilon(3S) \rightarrow \chi_b1(2P)X) \times B(\chi_b1(2P) \rightarrow \chi_b1(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$. We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_b1(2P)X) = B(\Upsilon(3S) \rightarrow \chi_b1(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$.							
18 CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming I-spin conservation, no D-wave contribution, $\Gamma(\chi_b1(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_b2(2P)) = 138 \pm 19$ keV.							
$\Gamma(D^0 X)/\Gamma_{\text{total}}$						Γ_5/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M080R01	NODE=M080R01
8.8±1.5±0.8	2243	19 BRIERE	08	CLEO $\Upsilon(3S) \rightarrow \gamma D^0 X$			
19 For $p_{D^0} > 2.5$ GeV/c.							

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
3.1±1.0±0.3	30	20 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$
20 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (39 \pm 8 \pm 9) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.1±0.5±0.1	10	21 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$
21 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (14 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
7.7±3.1±0.7	15	22 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$
22 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (97 \pm 30 \pm 26) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.9±2.0±0.5	36	23 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$
23 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (74 \pm 16 \pm 19) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
1.0±0.4±0.1	12	24 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$
24 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
5.5±1.7±0.5	38	25 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$
25 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (69 \pm 13 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
9.6±3.5±0.9	27	26 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$
26 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (121 \pm 29 \pm 33) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
6.7±2.5±0.6	17	27 ASNER	08A CLEO	$\Gamma(3S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$
27 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P))] = (85 \pm 23 \pm 22) \times 10^{-6}$ which we divide by our best value $B(\Gamma(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M080R02
NODE=M080R02

NODE=M080R02;LINKAGE=AS

NODE=M080R03
NODE=M080R03NODE=M080R04
NODE=M080R04NODE=M080R05
NODE=M080R05NODE=M080R05;LINKAGE=AS
NODE=M080R06
NODE=M080R06NODE=M080R07
NODE=M080R07

NODE=M080R07;LINKAGE=AS

NODE=M080R08
NODE=M080R08

NODE=M080R08;LINKAGE=AS

NODE=M080R09
NODE=M080R09

NODE=M080R09;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
1.2±0.4±0.1	18	28 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$	
28 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (15 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
12±4±1	44	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$	
29 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (150 \pm 30 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
2.0±0.7±0.2	16	30 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$	
30 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (25 \pm 7 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
6.1±2.1±0.6	25	31 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$	
31 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (77 \pm 17 \pm 21) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
1.7±0.6±0.2	16	32 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$	
32 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (22 \pm 6 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{19}/Γ
19±7±2	41	33 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$	
33 ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))] = (241 \pm 47 \pm 72) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\chi_{b1}(2P)$ Cross-Particle Branching Ratios

$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma \times \Gamma_{21}^{(3S)}/\Gamma^{(3S)}$
12.4±0.3±0.6	15k LEES 11J BABR $\Upsilon(3S) \rightarrow X \gamma$

 $B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.		
3.52 ^{+0.28} _{-0.27} ^{+0.17} _{-0.18}	34 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
3.24 ^{+0.56} _{-0.41} ^{+0.41}	58 35 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
2.34 ^{+0.28} _{-0.15} ^{+0.15}	36 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

NODE=M080R10
NODE=M080R10

NODE=M080R10;LINKAGE=AS

NODE=M080R11
NODE=M080R11NODE=M080R12
NODE=M080R12

NODE=M080R12;LINKAGE=AS

NODE=M080R13
NODE=M080R13NODE=M080R14
NODE=M080R14NODE=M080R15
NODE=M080R15

NODE=M080R15;LINKAGE=AS

NODE=M080230

NODE=M080B01
NODE=M080B01NODE=M080A00
NODE=M080A00

34 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.


35 CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.

36 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$.

NODE=M080A00;LINKAGE=A

NODE=M080A00;LINKAGE=C

NODE=M080A00;LINKAGE=B



$$\frac{\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) / \Gamma_{\text{total}}}{\Gamma_2 / \Gamma} \times \frac{\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) / \Gamma_{\text{total}}}{\Gamma_{21} / \Gamma_{21}} = \frac{\Gamma(\Upsilon(2S) \rightarrow \ell^+\ell^-)}{\Gamma(3S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 0.1 \pm 0.2$	4.3k	LEES	11J	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M080B02

NODE=M080B02

$$B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+\ell^-)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
$4.95^{+0.75+1.01}_{-0.70-0.24}$	37	LEES	14M	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$5.12 \pm 0.60 \pm 0.63$	111	38 CRAWFORD	92B	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M080A01

NODE=M080A01

37 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.


38 CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.

39 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$.

NODE=M080A01;LINKAGE=A

NODE=M080A01;LINKAGE=C

NODE=M080A01;LINKAGE=B



$$B(\chi_{b1}(2P) \rightarrow \gamma \gamma(2S)) \times B(\gamma(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\gamma(2S) \rightarrow \ell^+ \ell^-)$$

(units 10^{-4})

$$B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) \times B(\gamma(3S) \rightarrow \chi_{b1}(2P)X)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.16±0.07±0.12	31k	LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$

NODE=M080R16
NODE=M080R16

$$B(\chi_{b2}(2P) \rightarrow pX + \bar{p}X)/B(\chi_{b1}(2P) \rightarrow pX + \bar{p}X)$$

VALUE	DOCUMENT ID	TECN	COMMENT
1.109±0.007±0.040	BRIERE	07	CLEO $\gamma(3S) \rightarrow \gamma \chi_{bJ}(2P)$

NODE=M080R20
NODE=M080R20

$$B(\chi_{b0}(2P) \rightarrow pX + \bar{p}X)/B(\chi_{b1}(2P) \rightarrow pX + \bar{p}X)$$

VALUE	DOCUMENT ID	TECN	COMMENT
1.082±0.025±0.060	BRIERE	07	CLEO $\gamma(3S) \rightarrow \gamma \chi_{bJ}(2P)$

NODE=M080R21
NODE=M080R21

$\chi_{b1}(2P)$ REFERENCES

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawlfieeld <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... 04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

NODE=M080

REFID=56343
REFID=16775
REFID=53936
REFID=52574
REFID=52577
REFID=51887
REFID=50997
REFID=50454
REFID=49766
REFID=43177
REFID=43604
REFID=41580
REFID=41634
REFID=41586

NODE=M205

 $h_b(2P)$ $I^G(J^{PC}) = 0^-(1^{+-})$

OMMITTED FROM SUMMARY TABLE
 Quantum numbers are quark model predictions.

 $h_b(2P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10259.8±0.5±1.1	90k	MIZUK	12	BELL $e^+e^- \rightarrow \pi^+\pi^-$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10259.8±0.6 ^{+1.4} _{-1.0}	83.9k	¹ ADACHI	12	BELL 10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM

¹ Superseded by MIZUK 12. **$h_b(2P)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	not seen
Γ_2 $\eta_b(1S)\gamma$	(22±5) %
Γ_3 $\eta_b(2S)\gamma$	(48±13) %

 $h_b(2P)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$			Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN COMMENT
not seen			
	83.9k	ADACHI	12 BELL 10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM
$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$			
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN COMMENT
22.3±3.8^{+3.1}_{-3.3}	10k	MIZUK	12 BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons
$\Gamma(\eta_b(2S)\gamma)/\Gamma_{\text{total}}$			
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN COMMENT
47.5±10.5^{+6.8}_{-7.7}	26k	MIZUK	12 BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons

 $h_b(2P)$ REFERENCES

ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M205

NODE=M205M

NODE=M205M

NODE=M205M;LINKAGE=AD

NODE=M205215;NODE=M205

DESIG=1

DESIG=2

DESIG=3

NODE=M205225

NODE=M205R01

NODE=M205R01

NODE=M205R02

NODE=M205R02

NODE=M205R03

NODE=M205R03

NODE=M205

REFID=53962

REFID=54718

$\chi_{b2}(2P)$

$I^G(JPC) = 0^+(2^{++})$
J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b2}(2P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10268.65 ± 0.22 ± 0.50 OUR EVALUATION			From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

$m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.10 ± 0.24 OUR AVERAGE			
12.3 ± 2.6 ± 0.6	¹ AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$
13.04 ± 0.26	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
13.5 ± 0.4 ± 0.5	² HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$
1 From the $\chi_{bj}(2P) \rightarrow \Upsilon(1S)\gamma$ transition.			
2 From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.			

γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
86.19 ± 0.22 OUR EVALUATION				Treating systematic errors as correlated
86.40 ± 0.18 OUR AVERAGE				
86.04 ± 0.06 ± 0.27		ARTUSO	05 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
86 ± 1	101	CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
86.7 ± 0.4	10319	³ HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
86.9 ± 0.4	157	⁴ HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
86.4 ± 0.1 ± 0.4	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$
3 A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.				
4 A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.				

$\chi_{b2}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level	
$\Gamma_1 \omega \Upsilon(1S)$	(1.10 ± 0.30) %		DESIG=3
$\Gamma_2 \gamma \Upsilon(2S)$	(8.9 ± 1.2) %		DESIG=2
$\Gamma_3 \gamma \Upsilon(1S)$	(6.6 ± 0.8) %		DESIG=1
$\Gamma_4 \pi\pi\chi_{b2}(1P)$	(5.1 ± 0.9) × 10 ⁻³		DESIG=4
$\Gamma_5 D^0 X$	< 2.4 %	90%	DESIG=5
$\Gamma_6 \pi^+ \pi^- K^+ K^- \pi^0$	< 1.1 × 10 ⁻⁴	90%	DESIG=6
$\Gamma_7 2\pi^+ \pi^- K^- K_S^0$	< 9 × 10 ⁻⁵	90%	DESIG=7
$\Gamma_8 2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 7 × 10 ⁻⁴	90%	DESIG=8
$\Gamma_9 2\pi^+ 2\pi^- 2\pi^0$	(3.9 ± 1.6) × 10 ⁻⁴		DESIG=9
$\Gamma_{10} 2\pi^+ 2\pi^- K^+ K^-$	(9 ± 4) × 10 ⁻⁵		DESIG=10
$\Gamma_{11} 2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.4 ± 1.1) × 10 ⁻⁴		DESIG=11
$\Gamma_{12} 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(4.7 ± 2.3) × 10 ⁻⁴		DESIG=12
$\Gamma_{13} 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 4 × 10 ⁻⁴	90%	DESIG=13
$\Gamma_{14} 3\pi^+ 3\pi^-$	(9 ± 4) × 10 ⁻⁵		DESIG=14
$\Gamma_{15} 3\pi^+ 3\pi^- 2\pi^0$	(1.2 ± 0.4) × 10 ⁻³		DESIG=15
$\Gamma_{16} 3\pi^+ 3\pi^- K^+ K^-$	(1.4 ± 0.7) × 10 ⁻⁴		DESIG=16
$\Gamma_{17} 3\pi^+ 3\pi^- K^+ K^- \pi^0$	(4.2 ± 1.7) × 10 ⁻⁴		DESIG=17
$\Gamma_{18} 4\pi^+ 4\pi^-$	(9 ± 5) × 10 ⁻⁵		DESIG=18
$\Gamma_{19} 4\pi^+ 4\pi^- 2\pi^0$	(1.3 ± 0.5) × 10 ⁻³		DESIG=19

NODE=M081

NODE=M081

NODE=M081M

NODE=M081M
→ UNCHECKED ←

NODE=M081M2

NODE=M081M2

NODE=M081M2;LINKAGE=B
NODE=M081M2;LINKAGE=A

NODE=M081DM

NODE=M081DM
→ UNCHECKED ←

OCCUR=2

NODE=M081DM;LINKAGE=A

NODE=M081DM;LINKAGE=B

NODE=M081215;NODE=M081

$\chi_{b2}(2P)$ BRANCHING RATIOS

$\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
$1.10^{+0.32+0.11}_{-0.28-0.10}$	$20.1^{+5.8}_{-5.1}$	5 CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$	

⁵ Using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$.

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
-------	------	-------------	------	---------	-------------------

0.089 ± 0.012 OUR AVERAGE

$0.085 \pm 0.010 \pm 0.010$	6,7,8 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$		
$0.084 \pm 0.011 \pm 0.010$	2.5k 9 LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$		
$0.096 \pm 0.022 \pm 0.012$	7,10 CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$		
$0.106 \pm 0.016 \pm 0.013$	7,11 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$		

⁶ LEES 14M quotes $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}} = (1.12 \pm 0.13)\%$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and without converted photons.

⁷ Assuming $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

⁸ LEES 14M reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (1.12 \pm 0.13) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹ LEES 11J reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (1.1 \pm 0.1 \pm 0.1) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁰ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \ell^+ \ell^-) = (4.98 \pm 0.94 \pm 0.62) 10^{-4}$.

¹¹ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
-------	------	-------------	------	---------	-------------------

0.066 ± 0.008 OUR AVERAGE

$0.061 \pm 0.004 \pm 0.007$	12,13,14 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$		
$0.070 \pm 0.004 \pm 0.008$	11k 15 LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$		
$0.077 \pm 0.018 \pm 0.009$	13,16 CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$		
$0.061 \pm 0.009 \pm 0.007$	13,17 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$		

¹² LEES 14M quotes $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}} = (8.03 \pm 0.50) \times 10^{-3}$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and without converted photons.

¹³ Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

¹⁴ LEES 14M reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (8.03 \pm 0.50) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁵ LEES 11J reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (9.2 \pm 0.3 \pm 0.4) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

¹⁶ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (5.03 \pm 0.94 \pm 0.63) 10^{-4}$.

¹⁷ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91.

$\Gamma(\pi \pi \chi_{b2}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
--------------------------	------	-------------	------	---------	-------------------

5.1 ± 0.9 OUR AVERAGE

$4.9 \pm 0.7 \pm 0.6$	17k 18 LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$		
$6.0 \pm 1.6 \pm 1.4$	19 CAWLFIELD	06 CLE3	$\Upsilon(3S) \rightarrow 2(\gamma \pi \ell)$		

NODE=M081220

NODE=M081R3

NODE=M081R3

NODE=M081R3;LINKAGE=CR

NODE=M081R2

NODE=M081R2

NODE=M081R2;LINKAGE=D

NODE=M081R2;LINKAGE=E

NODE=M081R2;LINKAGE=F

NODE=M081R2;LINKAGE=LE

NODE=M081R2;LINKAGE=B

NODE=M081R2;LINKAGE=C

NODE=M081R1

NODE=M081R1

NODE=M081R1;LINKAGE=D

NODE=M081R1;LINKAGE=E

NODE=M081R1;LINKAGE=F

NODE=M081R1;LINKAGE=LE

NODE=M081R1;LINKAGE=B

NODE=M081R1;LINKAGE=C

NODE=M081R4

NODE=M081R4

¹⁸ $(0.64 \pm 0.05 \pm 0.08) \times 10^{-3}$. We derive the value assuming $B(\gamma(3S) \rightarrow \chi_{b2}(2P)X) = B(\gamma(3S) \rightarrow \chi_{b2}(2P)\gamma) = (13.1 \pm 1.6) \times 10^{-2}$.
¹⁹ CAWLFIELD 06 quote $\Gamma(\chi_{b2}(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming I-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

$\Gamma(D^0 X)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.4 × 10⁻²	90	20,21 BRIERE	08 CLEO	$\gamma(3S) \rightarrow \gamma D^0 X$	

20 For $p_{D^0} > 2.5$ GeV/c.

21 The authors also present their result as $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$.

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.1	90	22 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$	

22 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.9	90	23 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$	

23 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] < 12 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

$\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	
<7	90	24 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$	

24 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] < 87 \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

$\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.9±1.6±0.5	23	25 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$	

25 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.9±0.4±0.1	11	26 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$	

26 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.4±1.0±0.3	16	27 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$	

27 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] = (32 \pm 11 \pm 8) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$					Γ_{12}/Γ
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	
4.7±2.2±0.6	14	28 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$	

28 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P))] = (62 \pm 23 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\gamma(3S) \rightarrow \gamma\chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R4;LINKAGE=LE

NODE=M081R4;LINKAGE=CA

NODE=M081R01
NODE=M081R01

NODE=M081R01;LINKAGE=BR
NODE=M081R01;LINKAGE=RI

NODE=M081R02
NODE=M081R02

NODE=M081R02;LINKAGE=AS

NODE=M081R03
NODE=M081R03

NODE=M081R03;LINKAGE=AS

NODE=M081R04
NODE=M081R04

NODE=M081R04;LINKAGE=AS

NODE=M081R05
NODE=M081R05

NODE=M081R05;LINKAGE=AS

NODE=M081R06
NODE=M081R06

NODE=M081R06;LINKAGE=AS

NODE=M081R07
NODE=M081R07

NODE=M081R07;LINKAGE=AS

NODE=M081R08
NODE=M081R08

NODE=M081R08;LINKAGE=AS

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
29 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 58 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.				

 Γ_{13}/Γ NODE=M081R09
NODE=M081R09 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	14	30 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
30 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{14}/Γ NODE=M081R10
NODE=M081R10 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	45	31 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
31 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (159 \pm 33 \pm 43) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{15}/Γ NODE=M081R11
NODE=M081R11 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.4±0.7±0.2	12	32 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
32 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (19 \pm 7 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{16}/Γ NODE=M081R12
NODE=M081R12 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2±1.7±0.5	16	33 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
33 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (55 \pm 16 \pm 15) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{17}/Γ NODE=M081R13
NODE=M081R13 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	9	34 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
34 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{18}/Γ NODE=M081R14
NODE=M081R14 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
13±5±2	27	35 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
35 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (165 \pm 46 \pm 50) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

 Γ_{19}/Γ NODE=M081R15
NODE=M081R15 $\chi_{b2}(2P)$ Cross-Particle Branching Ratios

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \frac{\Gamma(3S)}{\Gamma(20)} / \Gamma(3S)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±0.3±0.4	11k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$

NODE=M081230

NODE=M081B01
NODE=M081B01

$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) / \Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) / \Gamma_{\text{total}}$					
$\frac{\Gamma_2 / \Gamma \times \Gamma^{(3S)}_{20}}{\Gamma^{(3S)}} / \Gamma^{(3S)}$					
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.1±0.1±0.1	2.5k	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X\gamma$
$B(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X)$					
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.64±0.05±0.08	17k	LEES	11C	BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
$B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$					
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.02±0.18 OUR AVERAGE					
1.95 $^{+0.22}_{-0.21}$ $^{+0.10}_{-0.16}$	36	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.52 $^{+0.47}_{-0.32}$ $^{+0.32}_{-0.28}$	48	37 CRAWFORD	92B	CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.98 $^{+0.28}_{-0.28}$ $^{+0.12}_{-0.12}$		38 HEINTZ	92	CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
36 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.					
37 CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.					
38 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma\Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$.					
$[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$					
<u>VALUE (%)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
66.6±3.0	39	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
39 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ events without converted photons.					
$B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+\ell^-)$					
<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
2.74±0.29 OUR AVERAGE					
3.22 $^{+0.58}_{-0.53}$ $^{+0.16}_{-0.71}$	40	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.49 $^{+0.47}_{-0.47}$ $^{+0.31}_{-0.31}$	53	41 CRAWFORD	92B	CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.74 $^{+0.33}_{-0.33}$ $^{+0.18}_{-0.18}$		42 HEINTZ	92	CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
40 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.					
41 CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.					
42 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma\Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$.					
$[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$					
<u>VALUE (%)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
46.9±2.0	43	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
43 From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ without converted photons.					

$\chi_{b2}(2P)$ REFERENCES

AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CAWLFIELD	06	PR D73 012003	C. Cawfield <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HEN... 04		PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)
				REFID=56199
				REFID=56343
				REFID=16775
				REFID=53936
				REFID=52574
				REFID=52577
				REFID=50997
				REFID=50454
				REFID=49766
				REFID=43177
				REFID=43604
				REFID=41580
				REFID=41634
				REFID=41586

NODE=M048

 $\Upsilon(3S)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\Upsilon(3S)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10355.2±0.5	¹ ARTAMONOV 00	MD1	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10355.3±0.5	2,3 BARU	86B REDE	$e^+ e^- \rightarrow$ hadrons
1 Reanalysis of BARU 86B using new electron mass (COHEN 87).			
2 Reanalysis of ARTAMONOV 84.			
3 Superseded by ARTAMONOV 00.			

NODE=M048M

NODE=M048M

NODE=M048M;LINKAGE=AR
 NODE=M048M;LINKAGE=C
 NODE=M048M;LINKAGE=RZ

 $m_{\Upsilon(3S)} - m_{\Upsilon(2S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$

NODE=M048DM2

NODE=M048DM2

 $\Upsilon(3S)$ WIDTH

VALUE (keV)	DOCUMENT ID
20.32±1.85 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

NODE=M048W

NODE=M048W
 → UNCHECKED ←

NODE=M048215;NODE=M048

 $\Upsilon(3S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \Upsilon(2S)$ anything	(10.6 ± 0.8) %	
$\Gamma_2 \Upsilon(2S)\pi^+\pi^-$	(2.82 ± 0.18) %	S=1.6
$\Gamma_3 \Upsilon(2S)\pi^0\pi^0$	(1.85 ± 0.14) %	
$\Gamma_4 \Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
$\Gamma_5 \Upsilon(2S)\pi^0$	< 5.1 × 10 ⁻⁴	CL=90%
$\Gamma_6 \Upsilon(1S)\pi^+\pi^-$	(4.37 ± 0.08) %	
$\Gamma_7 \Upsilon(1S)\pi^0\pi^0$	(2.20 ± 0.13) %	
$\Gamma_8 \Upsilon(1S)\eta$	< 1 × 10 ⁻⁴	CL=90%
$\Gamma_9 \Upsilon(1S)\pi^0$	< 7 × 10 ⁻⁵	CL=90%
$\Gamma_{10} h_b(1P)\pi^0$	< 1.2 × 10 ⁻³	CL=90%
$\Gamma_{11} h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 ± 1.4) × 10 ⁻⁴	
$\Gamma_{12} h_b(1P)\pi^+\pi^-$	< 1.2 × 10 ⁻⁴	CL=90%
$\Gamma_{13} \tau^+\tau^-$	(2.29 ± 0.30) %	
$\Gamma_{14} \mu^+\mu^-$	(2.18 ± 0.21) %	S=2.1
$\Gamma_{15} e^+e^-$	(2.18 ± 0.20) %	
Γ_{16} hadrons	(93 ± 12) %	
$\Gamma_{17} ggg$	(35.7 ± 2.6) %	
$\Gamma_{18} \gamma gg$	(9.7 ± 1.8) × 10 ⁻³	
$\Gamma_{19} \frac{1}{2}\bar{H}$ anything	(2.33 ± 0.33) × 10 ⁻⁵	

DESIG=8
 DESIG=4
 DESIG=10
 DESIG=12
 DESIG=107
 DESIG=3
 DESIG=11
 DESIG=9
 DESIG=106
 DESIG=112
 DESIG=113
 DESIG=114
 DESIG=16
 DESIG=1
 DESIG=2
 DESIG=101
 DESIG=109
 DESIG=110
 DESIG=117

Radiative decays

$\Gamma_{20} \gamma\chi b_2(2P)$	(13.1 ± 1.6) %	S=3.4
$\Gamma_{21} \gamma\chi b_1(2P)$	(12.6 ± 1.2) %	S=2.4
$\Gamma_{22} \gamma\chi b_0(2P)$	(5.9 ± 0.6) %	S=1.4
$\Gamma_{23} \gamma\chi b_2(1P)$	(10.0 ± 1.0) × 10 ⁻³	S=1.7
$\Gamma_{24} \gamma\chi b_1(1P)$	(9 ± 5) × 10 ⁻⁴	S=1.8
$\Gamma_{25} \gamma\chi b_0(1P)$	(2.7 ± 0.4) × 10 ⁻³	
$\Gamma_{26} \gamma\eta_b(2S)$	< 6.2 × 10 ⁻⁴	CL=90%
$\Gamma_{27} \gamma\eta_b(1S)$	(5.1 ± 0.7) × 10 ⁻⁴	
$\Gamma_{28} \gamma A^0 \rightarrow \gamma$ hadrons	< 8 × 10 ⁻⁵	CL=90%
$\Gamma_{29} \gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 × 10 ⁻⁴	CL=95%
$\Gamma_{30} \gamma a_1^0 \rightarrow \gamma\mu^+\mu^-$	< 5.5 × 10 ⁻⁶	CL=90%
$\Gamma_{31} \gamma a_1^0 \rightarrow \gamma\tau^+\tau^-$	[b] < 1.6 × 10 ⁻⁴	CL=90%

NODE=M048;CLUMP=B
 DESIG=5
 DESIG=6
 DESIG=7
 DESIG=103
 DESIG=104
 DESIG=13
 DESIG=14
 DESIG=15
 DESIG=115
 DESIG=102
 DESIG=116
 DESIG=108

Lepton Family number (*LF*) violating modes

Γ_{32}	$e^\pm \tau^\mp$	<i>LF</i>	< 4.2	$\times 10^{-6}$	CL=90%
Γ_{33}	$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.1	$\times 10^{-6}$	CL=90%

[a] $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ [b] For $m_{\tau^+ \tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. **$\Upsilon(3S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$** **$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{16}\Gamma_{15}/\Gamma$
0.414±0.007 OUR AVERAGE				

$0.413 \pm 0.004 \pm 0.006$	ROSNER	06	CLEO	$10.4 e^+ e^- \rightarrow \text{hadrons}$
$0.45 \pm 0.03 \pm 0.03$	⁴ GILES	84B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$

4 Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

 $\Gamma(\Upsilon(1S)\pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_6\Gamma_{15}/\Gamma$
18.46±0.27±0.77	6.4K	⁵ AUBERT	08BP BABR	$e^+ e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$	

5 Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$. **$\Upsilon(3S) \text{ PARTIAL WIDTHS}$** **$\Gamma(e^+ e^-)$**

VALUE (keV)	DOCUMENT ID	Γ_{15}
0.443±0.008 OUR EVALUATION		

 $\Upsilon(3S) \text{ BRANCHING RATIOS}$ **$\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}}$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.106 ± 0.008 OUR AVERAGE					

0.1023 ± 0.0105	4625	6,7,8 BUTLER	94B CLE2	$e^+ e^- \rightarrow \ell^+\ell^-X$
0.111 ± 0.012	4891	7,8,9 BROCK	91 CLEO	$e^+ e^- \rightarrow \pi^+\pi^-X, \pi^+\pi^-\ell^+\ell^-$

6 Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.7 Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.8 Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$.9 Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality. **$\Gamma(\Upsilon(2S)\pi^+ \pi^-)/\Gamma_{\text{total}}$**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
2.82±0.18 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.					

$3.00 \pm 0.02 \pm 0.14$	543k	LEES	11C BABR	$e^+ e^- \rightarrow \pi^+\pi^-X$
$2.40 \pm 0.10 \pm 0.26$	800	¹⁰ AUBERT	08BP BABR	$e^+ e^- \rightarrow \gamma\pi^+\pi^-e^+e^-$
3.12 ± 0.49	980	^{11,12} BUTLER	94B CLE2	$e^+ e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
2.13 ± 0.38	974	¹³ BROCK	91 CLEO	$e^+ e^- \rightarrow \pi^+\pi^-X, \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.82 \pm 0.65 \pm 0.53$	138	¹³ WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.1 ± 2.0	5	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

10 Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008 \text{ keV}$.

11 From the exclusive mode.

12 Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.13 Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

NODE=M048;CLUMP=C

DESIG=111

DESIG=105

LINKAGE=C48

LINKAGE=MRG

NODE=M048218

NODE=M048G2

NODE=M048G2

NODE=M048G2;LINKAGE=R

NODE=M048G01

NODE=M048G01

NODE=M048G01;LINKAGE=AU

NODE=M048220

NODE=M048W2

NODE=M048W2

→ UNCHECKED ←

NODE=M048225

NODE=M048R8

NODE=M048R8

NODE=M048R;LINKAGE=A

NODE=M048R;LINKAGE=B

NODE=M048R;LINKAGE=D

NODE=M048R;LINKAGE=C

NODE=M048R4


NODE=M048R4

NODE=M048R4;LINKAGE=AU

NODE=M048R;LINKAGE=M

NODE=M048R4;LINKAGE=A

NODE=M048R4;LINKAGE=C



$\Gamma(\gamma(2S)\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.85±0.14 OUR AVERAGE				
1.82±0.09±0.12	4391	14 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16±0.39		15,16 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.7 ± 0.5 ± 0.2	10	17 HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

14 Authors assume $B(\gamma(1S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

15 $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.

16 From the exclusive mode.

17 $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma(2S)\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0502±0.0069			
18 BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^- 2\gamma$	

18 From the exclusive mode.

$\Gamma(\gamma(2S)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.51	90	19 HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

19 Authors assume $B(\gamma(2S) \rightarrow e^+e^-) + B(\gamma(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

$\Gamma(\gamma(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.37±0.08 OUR AVERAGE				

4.32±0.07±0.13	90k	20 LEES	11L BABR	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.01±0.13	190k	21 BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^-$ MM
4.17±0.06±0.19	6.4K	22 AUBERT	08BP BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52±0.35	11830	23 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^- X, \pi^+\pi^-\ell^+\ell^-$
4.46±0.34±0.50	451	23 WU	93 CUSB	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.30	11221	23 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^- X, \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9 ± 1.0	22 GREEN	82 CLEO	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.9 ± 1.3	26 MAGERAS	82 CUSB	$\gamma(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

20 Using $B(\gamma(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

21 A weighted average of the inclusive and exclusive results.

22 Using $B(\gamma(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\gamma(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\gamma(3S)) = 0.443 \pm 0.008$ keV.

23 Using $B(\gamma(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

NODE=M048R10
NODE=M048R10

NODE=M048R10;LINKAGE=BH
NODE=M048R;LINKAGE=K
NODE=M048R10;LINKAGE=M
NODE=M048R;LINKAGE=G

NODE=M048R12
NODE=M048R12

NODE=M048R12;LINKAGE=M

NODE=M048R25
NODE=M048R25

NODE=M048R25;LINKAGE=HE

NODE=M048R3
NODE=M048R3
NODE=M048R3

NODE=M048R3;LINKAGE=LE

NODE=M048R3;LINKAGE=BH
NODE=M048R3;LINKAGE=AU

NODE=M048R3;LINKAGE=B

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.577 \pm 0.026 \pm 0.060	800	²⁴ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
24 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.				

 Γ_2/Γ_6

NODE=M048R28
NODE=M048R28

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20 \pm 0.13 OUR AVERAGE				
2.24 \pm 0.09 \pm 0.11	6584	²⁵ BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99 \pm 0.34	56	²⁶ BUTLER	94B	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 \pm 0.4 \pm 0.3	33	²⁷ HEINTZ	92	CSB2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
25 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.				
26 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.				
27 Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.				

 Γ_7/Γ

NODE=M048R11
NODE=M048R11

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.501 \pm 0.043	²⁸ BHARI	09	CLEO $e^+e^- \rightarrow \Upsilon(3S)$
28 Not independent of other values reported by BHARI 09.			

 Γ_7/Γ_6

NODE=M048R26
NODE=M048R26

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	90	²⁹ LEES	11L	BABR $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.8	90	^{29,30} AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
<0.18	90	³¹ HE	08A	CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$
<2.2	90	BROCK	91	CLEO $e^+e^- \rightarrow \ell^+\ell^-\eta$
29 Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.				
30 Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.				
31 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.				

 Γ_8/Γ

NODE=M048R9
NODE=M048R9

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<0.23	90	³² LEES	11L	BABR $\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.9	90	³³ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$
32 Not independent of other values reported by LEES 11L.				
33 Not independent of other values reported by AUBERT 08BP.				

 Γ_8/Γ_6

NODE=M048R27
NODE=M048R27

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.07	90	³⁴ HE	08A	CLEO $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
34 Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.				

 Γ_9/Γ

NODE=M048R24
NODE=M048R24

 $\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 \times 10 ⁻³	90	³⁵ GE	11	CLEO $\Upsilon(3S) \rightarrow \pi^0$ anything
35 Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.				

 Γ_{10}/Γ

NODE=M048R03
NODE=M048R03

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.3 \pm 1.1 \pm 0.9	LEES	11K	BABR $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$

 Γ_{11}/Γ

NODE=M048R33
NODE=M048R33

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{12}/Γ
<u>VALUE (units 10^{-4})</u>	<u>CL%</u>
< 1.2	90
36 LEES	36 DOCUMENT ID
11C BABR	TECN
$e^+ e^- \rightarrow \pi^+ \pi^- X$	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •	
<18	36 BUTLER
94B CLE2	$e^+ e^- \rightarrow \pi^+ \pi^- X$
<15	36 BROCK
91 CLEO	$e^+ e^- \rightarrow \pi^+ \pi^- X$
36 For $M(h_b(1P)) = 9900$ MeV.	

NODE=M048R34
NODE=M048R34

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$	Γ_{13}/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>
$2.29 \pm 0.21 \pm 0.22$	15k 37 DOCUMENT ID
BESSON 07	07 CLEO
$e^+ e^- \rightarrow \gamma(3S) \rightarrow \tau^+\tau^-$	COMMENT
37 BESSON 07 reports $[\Gamma(\gamma(3S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\gamma(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\gamma(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.	

NODE=M048R34;LINKAGE=MH

NODE=M048R18
NODE=M048R18


NODE=M048R18;LINKAGE=BE

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$	Γ_{13}/Γ_{14}
<u>VALUE</u>	<u>EVTS</u>
$1.05 \pm 0.08 \pm 0.05$	15k BESSON 07 CLEO
$e^+ e^- \rightarrow \gamma(3S)$	COMMENT

NODE=M048R19
NODE=M048R19

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$	Γ_{14}/Γ
<u>VALUE</u>	<u>EVTS</u>
0.0218 ± 0.0021 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.
0.0239 ± 0.0007 ± 0.0010	81k ADAMS 05 CLEO
0.0202 ± 0.0019 ± 0.0033	CHEN 89B CLEO
0.0173 ± 0.0015 ± 0.0011	KAARSBERG 89 CSB2
0.033 ± 0.013 ± 0.007	ANDREWS 83 CLEO
	$e^+ e^- \rightarrow \mu^+\mu^-$

NODE=M048R1
NODE=M048R1



$\Gamma(ggg)/\Gamma_{\text{total}}$	Γ_{17}/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>
35.7 ± 2.6	3M 38 DOCUMENT ID
BESSON 06A	06A CLEO
$\gamma(3S) \rightarrow \text{hadrons}$	COMMENT

NODE=M048R30
NODE=M048R30

38 Calculated using BESSON 06A value of $\Gamma(\gamma gg)/\Gamma(gg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\gamma(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+\pi^- \gamma(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0\pi^0 \gamma(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma\chi_{b2}(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma\chi_{b1}(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma\chi_{b0}(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma\chi_{b0}(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+\mu^-) = (2.18 \pm 0.21)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ BESSON 06A value.

NODE=M048R30;LINKAGE=BE

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$					Γ_{18}/Γ
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.97 ± 0.18	60k	39 BESSON	06A CLEO	$\Gamma(3S) \rightarrow \gamma + \text{hadrons}$	

39 Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(ggg)/\Gamma_{\text{total}}$ BESSON 06A value.

NODE=M048R31
NODE=M048R31

NODE=M048R31;LINKAGE=BE

$\Gamma(\gamma gg)/\Gamma(ggg)$					Γ_{18}/Γ_{17}
<u>VALUE (units 10^{-2})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$2.72 \pm 0.06 \pm 0.49$	3M	BESSON	06A CLEO	$\Gamma(3S) \rightarrow (\gamma + \text{hadrons})$	

NODE=M048R32
NODE=M048R32


$\Gamma(\overline{H} \text{ anything})/\Gamma_{\text{total}}$					Γ_{19}/Γ
<u>VALUE (units 10^{-5})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.33 \pm 0.15^{+0.31}_{-0.28}$	LEES	14G BABR	$e^+ e^- \rightarrow \overline{H} X$		I

NODE=M048R00
NODE=M048R00

$\Gamma(\gamma \chi b_2(2P))/\Gamma_{\text{total}}$					Γ_{20}/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.131 ± 0.016 OUR AVERAGE		Error includes scale factor of 3.4. See the ideogram below.			
0.1579 $\pm 0.0017 \pm 0.0073$	568k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$	
0.111 $\pm 0.005 \pm 0.004$	10319	HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$	
0.135 $\pm 0.003 \pm 0.017$	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$	

NODE=M048R5
NODE=M048R5

40 Supersedes NARAIN 91.




$\Gamma(\gamma \chi b_1(2P))/\Gamma_{\text{total}}$					Γ_{21}/Γ
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
0.126 ± 0.012 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.			
0.1454 $\pm 0.0018 \pm 0.0073$	537k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$	
0.115 $\pm 0.005 \pm 0.005$	11147	HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$	
0.105 $\pm 0.003 \pm 0.013$	25759	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$	

NODE=M048R6
NODE=M048R6

41 Supersedes NARAIN 91.

NODE=M048R6;LINKAGE=H




VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
0.059 ± 0.006 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.			
0.0677 ± 0.0020 ± 0.0065	225k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$	
0.060 ± 0.004 ± 0.006	4959	42 HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma X$	
0.049 +0.003 -0.004 ± 0.006	9903	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma X$	

42 Supersedes NARAIN 91.

NODE=M048R7
NODE=M048R7

NODE=M048R7;LINKAGE=H



VALUE (units 10^{-3})	CL %	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
10.0±1.0 OUR AVERAGE			Error includes scale factor of 1.7.			
8.0±1.3±0.4	126	43,44 KORNICER	11	CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$		
10.5±0.3 +0.7 -0.6	9.7k	LEES	11J	BABR $\gamma(3S) \rightarrow X\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<19 seen	90	45 ASNER	08A	CLEO $\gamma(3S) \rightarrow \gamma + \text{hadrons}$		
		46 HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$		

NODE=M048R21
NODE=M048R21

43 Assuming $B(\gamma(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

44 KORNICER 11 reports $[\Gamma(\gamma(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma\gamma(1S))]$
 $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma\gamma(1S)) = (18.0 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

45 ASNER 08A reports $[\Gamma(\gamma(3S) \rightarrow \gamma\chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma\chi_{b2}(1P))]$
 $< 27.1 \times 10^{-2}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

46 HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\gamma(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\gamma(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\gamma(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\gamma(1S) \rightarrow \ell^+ \ell^-$.

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$

Γ₂₄/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.9±0.5 OUR AVERAGE	Error includes scale factor of 1.8.				
1.5±0.4±0.1	50	47,48	KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$
0.5±0.3 ^{+0.2} _{-0.1}		LEES		11J	BABR $\gamma(3S) \rightarrow X\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7 seen	90	49 ASNER 50 HEINTZ	08A CLEO 92 CSB2	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$ $e^+ e^- \rightarrow \gamma\gamma\ell^+\ell^-$
--------------	----	-----------------------	---------------------	--

47 Assuming $B(\gamma(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

48 KORNICER 11 reports $[\Gamma(\gamma(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma\gamma(1S))]$
 $= (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma\gamma(1S)) = (35.2 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

49 ASNER 08A reports $[\Gamma(\gamma(3S) \rightarrow \gamma\chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P))]$
 $< 2.5 \times 10^{-2}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

50 HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\gamma(3S) \rightarrow \gamma\chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma\gamma(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0,1,2$ using inclusive $\gamma(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1,2$ using $\gamma(1S) \rightarrow \ell^+ \ell^-$.

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$

Γ₂₅/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.27±0.04 OUR AVERAGE					
0.27±0.04±0.02	2.3k	LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$
0.30±0.04±0.10	8.7k	ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90	51 ASNER	08A CLEO	$\gamma(3S) \rightarrow \gamma + \text{hadrons}$
------	----	----------	----------	--

51 ASNER 08A reports $[\Gamma(\gamma(3S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\gamma(2S) \rightarrow \gamma\chi_{b0}(1P))]$
 $< 21.9 \times 10^{-2}$ which we multiply by our best value $B(\gamma(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

$\Gamma(\gamma\eta_b(2S))/\Gamma_{\text{total}}$

Γ₂₆/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 6.2					
< 6.2	90	ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$
-----	----	------	-----	------	----------------------------------

$\Gamma(\gamma\eta_b(1S))/\Gamma_{\text{total}}$

Γ₂₇/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.1±0.7 OUR AVERAGE					
7.1±1.8±1.3	2.3±0.5k	52 BONVICINI	10	CLEO	$\gamma(3S) \rightarrow \gamma X$
4.8±0.5±0.6	19 ± 3k	52 AUBERT	09AQ	BABR	$\gamma(3S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.5	90	LEES	11J	BABR	$\gamma(3S) \rightarrow X\gamma$
4.8±0.5±1.2	19 ± 3k	52,53 AUBERT	08V	BABR	$\gamma(3S) \rightarrow \gamma X$
<4.3	90	54 ARTUSO	05	CLEO	$e^+ e^- \rightarrow \gamma X$

52 Assuming $\Gamma_{\eta_b}(1S) = 10$ MeV.

53 Systematic error re-evaluated by AUBERT 09AQ.

54 Superseded by BONVICINI 10.

NODE=M048R21;LINKAGE=KA

NODE=M048R21;LINKAGE=KR

NODE=M048R21;LINKAGE=AS

NODE=M048R21;LINKAGE=HE

NODE=M048R22

NODE=M048R22

NODE=M048R22;LINKAGE=KA

NODE=M048R22;LINKAGE=KR

NODE=M048R22;LINKAGE=AS

NODE=M048R22;LINKAGE=HE

NODE=M048R15

NODE=M048R15

NODE=M048R15;LINKAGE=AS

NODE=M048R16

NODE=M048R16

NODE=M048R17

NODE=M048R17

NODE=M048R17;LINKAGE=BO

NODE=M048R17;LINKAGE=AU

NODE=M048R17;LINKAGE=SU

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ (0.3 GeV < m_{A^0} < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-5}$	90	55 LEES	11H BABR	$\gamma(3S) \rightarrow \gamma \text{ hadrons}$
55 For a narrow scalar or pseudoscalar A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .				

 Γ_{28}/Γ

NODE=M048R02

NODE=M048R02

NODE=M048R02

NODE=M048R02;LINKAGE=LE

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ (1.5 GeV < m_X < 5.0 GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	95	ROSNER	07A CLEO	$e^+ e^- \rightarrow \gamma X$

 Γ_{29}/Γ

NODE=M048R20

NODE=M048R20

NODE=M048R20

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5.5	90	56 AUBERT	09Z BABR	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \mu^+ \mu^-$

56 For a narrow scalar or pseudoscalar a_1^0 with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of $m_{a_1^0}$ range from $0.27\text{--}5.5 \times 10^{-6}$.

NODE=M048R04

NODE=M048R04

NODE=M048R04;LINKAGE=AU

 $\Gamma(\gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.6 \times 10^{-4}$	90	57 AUBERT	09P BABR	$e^+ e^- \rightarrow \gamma a_1^0 \rightarrow \gamma \tau^+ \tau^-$

57 For a narrow scalar or pseudoscalar a_1^0 with $M(\tau^+ \tau^-)$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from $1.5\text{--}16 \times 10^{-5}$.

NODE=M048R29

NODE=M048R29

NODE=M048R29;LINKAGE=AU

LEPTON FAMILY NUMBER (LF) VIOLATING MODES $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.2	90	LEES	10B BABR	$e^+ e^- \rightarrow e^\pm \tau^\mp$

NODE=M048230

NODE=M048R01

NODE=M048R01

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	LEES	10B BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<20.3	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M048R23

NODE=M048R23

NODE=M048

LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	09AQ	PR D81 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PR D81 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
BESSON	07	PR D79 052002	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>	(CLEO Collab.)

REFID=55939

REFID=53960

REFID=16769

REFID=16775

REFID=53877

REFID=53936

REFID=53937

REFID=53938

REFID=53231

REFID=53355

REFID=53106

REFID=53062

REFID=52930

REFID=52662

REFID=52574

REFID=52660

REFID=52262

REFID=52587

REFID=52592

REFID=52166

REFID=51620

REFID=52079

REFID=51147

REFID=51035

REFID=50452

REFID=50454

REFID=47424

BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43799
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)	REFID=43313
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)	REFID=41579
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PR 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUELLER	88	HE e ⁺ e ⁻ Physics 412	W. Buchmuller, S. Cooper Editors: A. Ali and P. Soeding, World Scientific, Singapore	(HANN, DESY, MIT)	REFID=40034
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41 733			
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)	REFID=22273
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)	REFID=22321
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)	REFID=22359

$\chi_{b1}(3P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

Needs confirmation.

Observed in the radiative decay to $\Upsilon(1S, 2S, 3S)$, therefore $C = +$.
 J needs confirmation.

$\chi_{b1}(3P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
10513.42 ± 0.41 ± 0.53		1 SIRUNYAN	18N CMS	$p p \rightarrow \gamma \mu^+ \mu^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
10515.7 + 2.2 - 3.9 + 1.5 - 2.1	169	2 AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$	
10512.1 ± 2.1 ± 0.9	351	3 AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$	
10511.3 ± 1.7 ± 2.5	182	4 AAIJ	14BI LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$	
10530 ± 5 ± 9	5 AAD	12A ATLAS	$p p \rightarrow \gamma \mu^+ \mu^- X$		
10551 ± 14 ± 17	5 ABAZOV	12Q D0	$p \bar{p} \rightarrow \gamma \mu^+ \mu^- X$		

¹ Systematic error includes an additional 0.5 MeV for the uncertainty on the $\Upsilon(3S)$ mass.
Also measures $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$ MeV. A total of 372 $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ events was observed.

² From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for ±30% variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

³ The mass of the $\chi_{b1}(3P)$ state obtained by combining the results of AAIJ 14BG with that of AAIJ 14BI. The first uncertainty is experimental and the second attributable to the unknown mass splitting, assumed to be $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

⁴ From $\chi_{b1}(3P) \rightarrow \Upsilon(3S)\gamma$ transition assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

⁵ The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.

$\chi_{b1}(3P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \Upsilon(1S)\gamma$	seen
$\Gamma_2 \quad \Upsilon(2S)\gamma$	seen
$\Gamma_3 \quad \Upsilon(3S)\gamma$	seen

$\chi_{b1}(3P)$ BRANCHING RATIOS

$\Gamma(\Upsilon(1S)\gamma)/\Gamma_{\text{total}}$				Γ_1/Γ	
seen	169	1 AAIJ	14BG LHCb	$p p \rightarrow \gamma \mu^+ \mu^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen		AAD	12A ATLAS	$p p \rightarrow \gamma \mu^+ \mu^- X$	
seen		ABAZOV	12Q D0	$p \bar{p} \rightarrow \gamma \mu^+ \mu^- X$	

¹ From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for ±30% variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

REFID=43799
REFID=43313
REFID=43604
REFID=41579
REFID=41580
REFID=41634
REFID=41586
REFID=40919
REFID=40733
REFID=40034

REFID=11616
REFID=22338
REFID=40033

NODE=M206

NODE=M206

NODE=M206M

NODE=M206M

OCCUR=2

NODE=M206M;LINKAGE=D

NODE=M206M;LINKAGE=A

NODE=M206M;LINKAGE=B

NODE=M206M;LINKAGE=C

NODE=M206M;LINKAGE=AA

NODE=M206215;NODE=M206

DESIG=1

DESIG=2

DESIG=3

NODE=M206225

NODE=M206R01
NODE=M206R01

NODE=M206R01;LINKAGE=A

$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	169	1 AAIJ	14BG LHCb	$pp \rightarrow \gamma\mu^+\mu^- X$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
seen		AAD	12A ATLAS	$pp \rightarrow \gamma\mu^+\mu^- X$	
1 From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.					

 $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen		SIRUNYAN 18N CMS		$pp \rightarrow \gamma\mu^+\mu^- X$	
seen	182	AAIJ 14BI LHCb		$pp \rightarrow \gamma\mu^+\mu^- X$	

 $\chi_{b1}(3P)$ REFERENCES

SIRUNYAN 18N PRL 121 092002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ 14BG JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 14BI EPJ C74 3092	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD 12A PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV 12Q PR D86 031103	V.M. Abazov <i>et al.</i>	(D0 Collab.)

 $\chi_{b2}(3P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Needs confirmation.

Observed in the radiative decay to $\Upsilon(3S)$, therefore $C = +$. J needs confirmation. $\chi_{b2}(3P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
10524.02 \pm 0.57 \pm 0.53	1 SIRUNYAN 18N CMS		$pp \rightarrow \gamma\mu^+\mu^- X$	
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
10530 \pm 5 \pm 9	2 AAD 12A ATLAS		$pp \rightarrow \gamma\mu^+\mu^- X$	
1 Systematic error includes an additional 0.5 MeV for the uncertainty on the $\Upsilon(3S)$ mass. Also measures $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$ MeV. A total of 372 $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ events was observed. 2 The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.				

 $\chi_{b2}(3P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 \quad \Upsilon(3S)\gamma$	seen	

 $\chi_{b2}(3P)$ BRANCHING RATIOS

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	SIRUNYAN 18N CMS		$pp \rightarrow \gamma\mu^+\mu^- X$	

 $\chi_{b2}(3P)$ REFERENCES

SIRUNYAN 18N PRL 121 092002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAD 12A PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)

NODE=M206R02
NODE=M206R02

NODE=M206R02;LINKAGE=A

NODE=M206R03
NODE=M206R03

NODE=M206

REFID=58873
REFID=56199
REFID=56235
REFID=54037
REFID=54264

NODE=M238

NODE=M238

NODE=M238M

NODE=M238M

NODE=M238M;LINKAGE=A

NODE=M238M;LINKAGE=AA

NODE=M238215;NODE=M238

DESIG=1

NODE=M238225

NODE=M238R01
NODE=M238R01

NODE=M238

REFID=58873
REFID=54037

NODE=M047

 $\Upsilon(4S)$

$I^G(J^{PC}) = 0^-(1^- -)$

also known as $\Upsilon(10580)$ **$\Upsilon(4S)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10579.4±1.2 OUR AVERAGE			
10579.3±0.4±1.2	AUBERT	05Q BABR $e^+ e^- \rightarrow$ hadrons	
10580.0±3.5	¹ BEBEK	87 CLEO $e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10577.4±1.0	² LOVELOCK	85 CUSB $e^+ e^- \rightarrow$ hadrons	
1 Reanalysis of BESSON 85.			
2 No systematic error given.			

NODE=M047M

NODE=M047M

NODE=M047M;LINKAGE=C
NODE=M047M;LINKAGE=B **$\Upsilon(4S)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
20.5±2.5 OUR AVERAGE			
20.7±1.6±2.5	AUBERT	05Q BABR $e^+ e^- \rightarrow$ hadrons	
20 ± 2 ± 4	BESSON	85 CLEO $e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
25 ± 2.5	LOVELOCK	85 CUSB $e^+ e^- \rightarrow$ hadrons	

NODE=M047W

NODE=M047W

NODE=M047215;NODE=M047

 $\Upsilon(4S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level	
$\Gamma_1 B\bar{B}$	> 96 %	95%	
$\Gamma_2 B^+ B^-$	(51.4 ± 0.6) %		DESIG=8;OUR EST;→ UNCHECKED ←
$\Gamma_3 D_s^+$ anything + c.c.	(17.8 ± 2.6) %		DESIG=10
$\Gamma_4 B^0 \bar{B}^0$	(48.6 ± 0.6) %		DESIG=12
$\Gamma_5 J/\psi K_S^0 + (J/\psi, \eta_c) K_S^0$	< 4 × 10 ⁻⁷	90%	DESIG=11
Γ_6 non- $B\bar{B}$	< 4 %	95%	DESIG=15
$\Gamma_7 e^+ e^-$	(1.57 ± 0.08) × 10 ⁻⁵		DESIG=6
$\Gamma_8 \rho^+ \rho^-$	< 5.7 × 10 ⁻⁶	90%	DESIG=1
$\Gamma_9 K^*(892)^0 \bar{K}^0$	< 2.0 × 10 ⁻⁶	90%	DESIG=16
$\Gamma_{10} J/\psi(1S)$ anything	< 1.9 × 10 ⁻⁴	95%	DESIG=22
$\Gamma_{11} D^{*+}$ anything + c.c.	< 7.4 %	90%	DESIG=2
$\Gamma_{12} \phi$ anything	(7.1 ± 0.6) %		DESIG=3
$\Gamma_{13} \phi \eta$	< 1.8 × 10 ⁻⁶	90%	DESIG=4
$\Gamma_{14} \phi \eta'$	< 4.3 × 10 ⁻⁶	90%	DESIG=13
$\Gamma_{15} \rho \eta$	< 1.3 × 10 ⁻⁶	90%	DESIG=18
$\Gamma_{16} \rho \eta'$	< 2.5 × 10 ⁻⁶	90%	DESIG=19
$\Gamma_{17} \gamma(1S)$ anything	< 4 × 10 ⁻³	90%	DESIG=20
$\Gamma_{18} \gamma(1S) \pi^+ \pi^-$	(8.2 ± 0.4) × 10 ⁻⁵		DESIG=5
$\Gamma_{19} \gamma(1S) \eta$	(1.81 ± 0.18) × 10 ⁻⁴		DESIG=7
$\Gamma_{20} \gamma(1S) \eta'$	(3.4 ± 0.9) × 10 ⁻⁵		DESIG=17
$\Gamma_{21} \gamma(2S) \pi^+ \pi^-$	(8.2 ± 0.8) × 10 ⁻⁵		DESIG=26
$\Gamma_{22} h_b(1P) \pi^+ \pi^-$	not seen		DESIG=9
$\Gamma_{23} h_b(1P) \eta$	(2.18 ± 0.21) × 10 ⁻³		DESIG=21
$\Gamma_{24} {}^2H$ anything	< 1.3 × 10 ⁻⁵	90%	DESIG=23
Double Radiative Decays			
$\Gamma_{25} \gamma\gamma \Upsilon(D) \rightarrow \gamma\gamma\eta \Upsilon(1S)$	< 2.3 × 10 ⁻⁵	90%	DESIG=14

NODE=M047;CLUMP=B
DESIG=24

$\Upsilon(4S)$ PARTIAL WIDTHS

$\Gamma(e^+ e^-)$

VALUE (keV)

0.272±0.029 OUR AVERAGE

	DOCUMENT ID	TECN	COMMENT
0.321±0.017±0.029	AUBERT	05Q	BABR $e^+ e^- \rightarrow$ hadrons
0.28 ± 0.05 ± 0.01	1 ALBRECHT	95E	ARG $e^+ e^- \rightarrow$ hadrons
0.192±0.007±0.038	BESSON	85	CLEO $e^+ e^- \rightarrow$ hadrons
0.283±0.037	LOVELOCK	85	CUSB $e^+ e^- \rightarrow$ hadrons

¹ Using LEYAOUANC 77 parametrization of $\Gamma(s)$.


Γ_7

NODE=M047220

NODE=M047W1

NODE=M047W1

NODE=M047W1;LINKAGE=A



$\Upsilon(4S)$ BRANCHING RATIOS

$B\bar{B}$ DECAYS

The ratio of branching fraction to charged and neutral B mesons is often derived assuming isospin invariance in the decays, and relies on the knowledge of the B^+/B^0 lifetime ratio. "OUR EVALUATION" is obtained based on averages of rescaled data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <http://www.slac.stanford.edu/xorg/hflav/>. The averaging/rescaling procedure takes into account the common dependence of the measurement on the value of the lifetime ratio.

$\Gamma(B^+ B^-)/\Gamma_{\text{total}}$

VALUE

0.514±0.006 OUR EVALUATION

DOCUMENT ID

Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$

Γ_2/Γ

NODE=M047230

NODE=M047BBD

NODE=M047BBD

NODE=M047R11

NODE=M047R11

→ UNCHECKED ←

$\Gamma(D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE

0.178±0.021±0.016

DOCUMENT ID

1 ARTUSO

TECN CLE3 $e^+ e^- \rightarrow D_X X$

¹ ARTUSO 05B reports $[\Gamma(\Upsilon(4S) \rightarrow D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (8.0 \pm 0.2 \pm 0.9) \times 10^{-3}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

Γ_3/Γ

NODE=M047R13

NODE=M047R13

NODE=M047R13;LINKAGE=AR

$\Gamma(B^0 \bar{B}^0)/\Gamma_{\text{total}}$

VALUE

0.486±0.006 OUR EVALUATION

DOCUMENT ID

Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$

Γ_4/Γ

NODE=M047R12

NODE=M047R12

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.487±0.010±0.008 ¹ AUBERT,B 05H BABR $\Upsilon(4S) \rightarrow \bar{B}B \rightarrow D^*\ell\nu_\ell$

¹ Direct measurement. This value is averaged with the value extracted from the $\Gamma(B^+ B^-) / \Gamma(B^0 \bar{B}^0)$ measurements.

NODE=M047R12;LINKAGE=AU

$\Gamma(B^+ B^-)/\Gamma(B^0 \bar{B}^0)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_4
1.058±0.024 OUR EVALUATION				
1.006±0.036±0.031	¹ AUBERT 04F	BABR	$\gamma(4S) \rightarrow B\bar{B} \rightarrow J/\psi K$	
1.01 ± 0.03 ± 0.09	¹ HASTINGS 03	BELL	$\gamma(4S) \rightarrow B\bar{B} \rightarrow \text{dileptons}$	
1.058±0.084±0.136	² ATHAR 02	CLEO	$\gamma(4S) \rightarrow B\bar{B} \rightarrow D^* \ell\nu$	
1.10 ± 0.06 ± 0.05	³ AUBERT 02	BABR	$\gamma(4S) \rightarrow B\bar{B} \rightarrow (c\bar{c})K^*$	
1.04 ± 0.07 ± 0.04	⁴ ALEXANDER 01	CLEO	$\gamma(4S) \rightarrow B\bar{B} \rightarrow J/\psi K^*$	
1 HASTINGS 03 and AUBERT 04F assume $\tau(B^+)/\tau(B^0) = 1.083 \pm 0.017$.				
2 ATHAR 02 assumes $\tau(B^+)/\tau(B^0) = 1.074 \pm 0.028$. Supersedes BARISH 95.				
3 AUBERT 02 assumes $\tau(B^+)/\tau(B^0) = 1.062 \pm 0.029$.				
4 ALEXANDER 01 assumes $\tau(B^+)/\tau(B^0) = 1.066 \pm 0.024$.				

 $[\Gamma(J/\psi K_S^0) + \Gamma((J/\psi, \eta_c) K_S^0)]/\Gamma_{\text{total}}$ Forbidden by CP invariance.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	¹ TAJIMA	07A	BELL $\gamma(4S) \rightarrow B^0 \bar{B}^0$

1 $\gamma(4S)$ with $CP = +1$ decays to the final state with $CP = -1$. Γ_5/Γ $\Gamma(\text{non}-B\bar{B})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	95	BARISH	96B	CLEO $e^+ e^-$

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.57±0.08 OUR AVERAGE			

1.55±0.04±0.07	AUBERT	05Q	BABR $e^+ e^- \rightarrow \text{hadrons}$
2.77±0.50±0.49	¹ ALBRECHT	95E	ARG $e^+ e^- \rightarrow \text{hadrons}$

1 Using LEYAOUANC 77 parametrization of $\Gamma(s)$. $\Gamma(\rho^+ \rho^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.7×10^{-6}	90	AUBERT	08B0	BABR $e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0$

 $\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.0×10^{-6}	90	SHEN	13A	BELL $e^+ e^- \rightarrow K^*(892)^0 \bar{K}^0$

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	95	¹ ABE	02D	BELL $e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	90	¹ AUBERT	01C	BABR $e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$
------	----	---------------------	-----	---

1 Uses $B(J/\psi \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$. $\Gamma(D^{*+} \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.074	90	¹ ALEXANDER	90C	CLEO $e^+ e^-$

1 For $x > 0.473$. $\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
7.1 ± 0.1 ± 0.6		HUANG	07	CLEO $\gamma(4S) \rightarrow \phi X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23	90	¹ ALEXANDER	90C	CLEO $e^+ e^-$
-------	----	------------------------	-----	----------------

1 For $x > 0.52$. $\Gamma(\phi\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	¹ BELOUS	09	BELL $e^+ e^- \rightarrow \phi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.5	90	AUBERT,BE	06F	BABR $e^+ e^- \rightarrow \phi\eta$
------	----	-----------	-----	-------------------------------------

1 Using all intermediate branching fraction values from PDG 08.

NODE=M047R10

NODE=M047R10

→ UNCHECKED ←

NODE=M047R10;LINKAGE=F

NODE=M047R10;LINKAGE=D

NODE=M047R10;LINKAGE=E

NODE=M047R10;LINKAGE=C

NODE=M047R16

NODE=M047R16

NODE=M047R16

NODE=M047R16;LINKAGE=TA

NODE=M047NBB

NODE=M047R6

NODE=M047R6

NODE=M047R5

NODE=M047R5

NODE=M047R5;LINKAGE=A

NODE=M047R17

NODE=M047R17

NODE=M047R02

NODE=M047R02

NODE=M047R1

NODE=M047R1

NODE=M047R;LINKAGE=AC

NODE=M047R2

NODE=M047R2

NODE=M047R2;LINKAGE=A

NODE=M047R3

NODE=M047R3

NODE=M047R3;LINKAGE=A

NODE=M047R14

NODE=M047R14

NODE=M047R14;LINKAGE=BE

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{14}/Γ
<4.3	90	1 BELOUS	09	BELL $e^+ e^- \rightarrow \phi\eta'$	

¹ Using all intermediate branching fraction values from PDG 08.

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{15}/Γ
<1.3	90	1 BELOUS	09	BELL $e^+ e^- \rightarrow \rho\eta$	

¹ Using all intermediate branching fraction values from PDG 08.

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-6})	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{16}/Γ
<2.5	90	1 BELOUS	09	BELL $e^+ e^- \rightarrow \rho\eta'$	

¹ Using all intermediate branching fraction values from PDG 08.

 $\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{\text{total}}$

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{17}/Γ
<0.004	90	ALEXANDER	90C	CLEO $e^+ e^-$	

 $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{18}/Γ
8.2 ± 0.4 OUR AVERAGE			GUIDO	17	BELL $\Upsilon(4S) \rightarrow \pi^+\pi^-\mu^+\mu^-$	

8.2 $\pm 0.5 \pm 0.4$	515	GUIDO	17	BELL $\Upsilon(4S) \rightarrow \pi^+\pi^-\mu^+\mu^-$	
8.5 $\pm 1.3 \pm 0.2$	113 ± 16	1 SOKOLOV09	BELL	$e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$	
$8.00 \pm 0.64 \pm 0.27$	430	2 AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.8 $\pm 4.0 \pm 0.3$	3,4	SOKOLOV07	BELL	$e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$	
9.0 $\pm 1.5 \pm 0.2$	167 ± 19	5 AUBERT	06R BABR	$e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$	
<12	90	GLENN	99	CLE2 $e^+ e^-$	

¹ SOKOLOV 09 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (0.211 \pm 0.030 \pm 0.014) \times 10^{-5}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³ SOKOLOV 07 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (4.42 \pm 0.81 \pm 0.56) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ According to the authors, systematic errors were underestimated.

⁵ Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (2.23 \pm 0.25 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-4})	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{19}/Γ
1.81 ± 0.18 OUR AVERAGE			GUIDO	17	BELL $\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\mu^+\mu^-$	

1.70 $\pm 0.23 \pm 0.08$	49	GUIDO	17	BELL $\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\mu^+\mu^-$	
1.96 $\pm 0.26 \pm 0.09$	56	1 AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\ell^+\ell^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.7	90	2 TAMPONI	15	BELL $e^+ e^- \rightarrow \gamma\eta + \text{hadrons}$	
------	----	-----------	----	--	--

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

² Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.

 $\Gamma(\Upsilon(1S)\eta')/\Gamma_{\text{total}}$

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	Γ_{20}/Γ
$3.43 \pm 0.88 \pm 0.21$	27	GUIDO	18	BELL $\Upsilon(4S) \rightarrow (\rho^0\gamma, \pi^+\pi^-\eta)\mu^+\mu^-$	

NODE=M047R21
NODE=M047R21

NODE=M047R21;LINKAGE=BE

NODE=M047R22
NODE=M047R22

NODE=M047R22;LINKAGE=BE

NODE=M047R23
NODE=M047R23

NODE=M047R23;LINKAGE=BE

NODE=M047R4
NODE=M047R4

NODE=M047R7
NODE=M047R7

NODE=M047R7;LINKAGE=SK

NODE=M047R7;LINKAGE=UB

NODE=M047R7;LINKAGE=SO

NODE=M047R7;LINKAGE=US

NODE=M047R7;LINKAGE=AU

NODE=M047R18;LINKAGE=UB

NODE=M047R18;LINKAGE=A

NODE=M047R03
NODE=M047R03

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.41 \pm 0.40 \pm 0.12$ 56 ¹ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+ \pi^- (\pi^0) \ell^+ \ell^-$

1 Not independent of other values reported by AUBERT 08BP.

 Γ_{19}/Γ_{18}

NODE=M047R19
NODE=M047R19

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

8.2±0.8 OUR AVERAGE

$7.9 \pm 1.0 \pm 0.4$ 181 GUIDO 17 BELL $\Upsilon(4S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$

$8.6 \pm 1.1 \pm 0.7$ 220 ¹ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$8.8 \pm 1.7 \pm 0.8$ 97 ± 15 ² AUBERT 06R BABR $e^+ e^- \rightarrow \pi^+ \pi^- \mu^+ \mu^-$
 <3.9 90 GLENN 99 CLE2 $e^+ e^-$

¹ Using $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$ and $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

² Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \mu^+ \mu^-)] = (1.69 \pm 0.26 \pm 0.20) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 Γ_{21}/Γ

NODE=M047R9
NODE=M047R9

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.16 \pm 0.16 \pm 0.14$ 220 ¹ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

 Γ_{21}/Γ_{18}

NODE=M047R20
NODE=M047R20

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

not seen $(35^{+32}_{-26})k$ ¹ ADACHI 12 BELL $10.58 e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$

¹ From the upper limit on the ratio of $\sigma(e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-)$ at the $\Upsilon(4S)$ to that at the $\Upsilon(5S)$ of 0.27.

 Γ_{22}/Γ

NODE=M047R01
NODE=M047R01

 $\Gamma(h_b(1P)\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.18±0.11±0.18 112k ¹ TAMPONI 15 BELL $e^+ e^- \rightarrow h_b(1P)\eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.

 Γ_{23}/Γ

NODE=M047R00
NODE=M047R00

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.3 90 ASNER 07 CLEO $e^+ e^- \rightarrow \bar{d}X$

 Γ_{24}/Γ

NODE=M047R00;LINKAGE=AD

Double Radiative Decays $\Gamma(\gamma\gamma \Upsilon(D) \rightarrow \gamma\gamma \eta \Upsilon(1S))/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.3 × 10⁻⁵ 90 GUIDO 17 BELL $\Upsilon(4S) \rightarrow \gamma\gamma \pi^+ \pi^- \pi^0 \mu^+ \mu^-$

 Γ_{25}/Γ

NODE=M047R24
NODE=M047R24

 $\Upsilon(4S)$ REFERENCES

GUIDO	18	PRL 121 062001	E. Guido <i>et al.</i>	(BELLE Collab.)
GUIDO	17	PR D96 052005	E. Guido <i>et al.</i>	(BELLE Collab.)
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
BELOUS	09	PL B681 400	K. Belous <i>et al.</i>	(BELLE Collab.)
SOKOLOV	09	PR D79 051103	A. Sokolov <i>et al.</i>	(BELLE Collab.)
AUBERT	08BO	PR D78 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)
SOKOLOV	07	PR D75 071103	A. Sokolov <i>et al.</i>	(BELLE Collab.)
TAJIMA	07A	PRL 99 211601	O. Tajima <i>et al.</i>	(BELLE Collab.)
AUBERT	06R	PR D96 232001	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06F	PR D74 111103	B. Aubert <i>et al.</i>	(BABAR Collab.)
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)
AUBERT	05Q	PR D72 032005	B. Aubert <i>et al.</i>	(BABAR Collab.)

REFID=58860
REFID=58218
REFID=56996
REFID=55591
REFID=53962
REFID=53107
REFID=52760
REFID=52659
REFID=52660
REFID=52166
REFID=51617
REFID=51624
REFID=51715
REFID=52066
REFID=51143
REFID=51563
REFID=50992
REFID=50774

AUBERT,B	05H	PRL 95 042001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50777
AUBERT	04F	PR D69 071101	B.Aubert <i>et al.</i>		REFID=49748
HASTINGS	03	PR D67 052004	N.C. Hastings <i>et al.</i>	(BELLE Collab.)	REFID=49209
ABE	02D	PR 88 052001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48557
ATHAR	02	PR D66 052003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=48832
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48514
ALEXANDER	01	PRL 86 2737	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48316
AUBERT	01C	PRL 87 162002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48346
GLENN	99	PR D59 052003	S. Glenn <i>et al.</i>		REFID=46890
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44693
ALBRECHT	95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44372
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44139
ALEXANDER	90C	PRL 64 2226	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=41346
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=40270
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PR 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369
LEYAOUANC	77	PL B71 397	A. Le Yaouanc <i>et al.</i>	(ORSAY)	REFID=44695

Z_b(10610)

$I^G(J^{PC}) = 1^+(1^{+-})$

was X(10610)

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Observed by BONDAR 12 in $\gamma(5S)$ decays to $\gamma(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

Z_b(10610)[±] MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
10607.2±2.0	¹ BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10608.5±3.4 ^{+3.7} _{-1.4}	² GARMASH	15	BELL $e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
10608.1±1.2 ^{+1.5} _{-0.2}	² GARMASH	15	BELL $e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$	OCCUR=2
10607.4±1.5 ^{+0.8} _{-0.2}	² GARMASH	15	BELL $e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$	OCCUR=3
10611 ± 4 ± 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \gamma(1S)\pi^+\pi^-$	OCCUR=2
10609 ± 2 ± 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \gamma(2S)\pi^+\pi^-$	OCCUR=3
10608 ± 2 ± 3	³ BONDAR	12	BELL $e^+e^- \rightarrow \gamma(3S)\pi^+\pi^-$	OCCUR=4
10605 ± 2 ± 3	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$	OCCUR=5
10599 ± 6 ± 5	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$	OCCUR=6

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.**Z_b(10610)⁰ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
10609±4±4	¹ KROKOVNY	13	BELL $e^+e^- \rightarrow \gamma(2S)/\gamma(3S)\pi^0\pi^0$	

¹ From a simultaneous fit to the KROKOVNY 13 Dalitz analysis of $e^+e^- \rightarrow \gamma(2S)/\gamma(3S)\pi^0\pi^0$ decays with fixed width $\Gamma(Z_b(10610)^0) = 18.4$ MeV.

Z_b(10610)[±] WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
18.4± 2.4	¹ BONDAR	12	BELL $e^+e^- \rightarrow$ hadrons	

NODE=M207

NODE=M207

NODE=M207M

NODE=M207M

NODE=M207M;LINKAGE=BO

NODE=M207M;LINKAGE=A

NODE=M207M;LINKAGE=BN

NODE=M207M0

NODE=M207M0

NODE=M207M0;LINKAGE=A

NODE=M207W

NODE=M207W

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18.5 \pm 5.3^{+6.1}_{-2.3}$	² GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
$20.8 \pm 2.5^{+0.3}_{-2.1}$	² GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	
$18.7 \pm 3.4^{+2.5}_{-1.3}$	² GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	
$22.3 \pm 7.7^{+3.0}_{-4.0}$	³ BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
$24.2 \pm 3.1^{+2.0}_{-3.0}$	³ BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	
$17.6 \pm 3.0 \pm 3.0$	³ BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	
$11.4^{+4.5+2.1}_{-3.9-1.2}$	³ BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$	
13^{+10+9}_{-8-7}	³ BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$	

¹ Average of the BONDAR 12 measurements in separate channels.

² Correlated with the corresponding result from BONDAR 12.

³ Superseded by the average measurement of BONDAR 12.

Z_b(10610) DECAY MODES

Mode	Fraction (Γ_i/Γ)	
$\Gamma_1 \gamma(1S)\pi^+$	$(5.4^{+1.9}_{-1.5}) \times 10^{-3}$	DESIG=1
$\Gamma_2 \gamma(1S)\pi^0$	not seen	DESIG=9
$\Gamma_3 \gamma(2S)\pi^+$	$(3.6^{+1.1}_{-0.8}) \%$	DESIG=2
$\Gamma_4 \gamma(2S)\pi^0$	seen	DESIG=10
$\Gamma_5 \gamma(3S)\pi^+$	$(2.1^{+0.8}_{-0.6}) \%$	DESIG=3
$\Gamma_6 \gamma(3S)\pi^0$	seen	DESIG=11
$\Gamma_7 h_b(1P)\pi^+$	$(3.5^{+1.2}_{-0.9}) \%$	DESIG=4
$\Gamma_8 h_b(2P)\pi^+$	$(4.7^{+1.7}_{-1.3}) \%$	DESIG=5
$\Gamma_9 B^+\bar{B}^0$	not seen	DESIG=8
$\Gamma_{10} B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	$(85.6^{+2.1}_{-2.9}) \%$	DESIG=6
$\Gamma_{11} B^{*+}\bar{B}^{*0}$	not seen	DESIG=7

Z_b(10610) BRANCHING RATIOS

$\Gamma(\gamma(1S)\pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ
<u>VALUE (units 10^{-3})</u>	
$5.4^{+1.6+1.1}_{-1.3-0.8}$	¹ GARMASH 16 BELL $e^+ e^- \rightarrow \pi^-\pi^+ B^+\bar{B}^{*0}, \pi^-\bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH 15 BELL $e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$
seen	BONDAR 12 BELL $e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$

¹ Assuming the Z_b(10610) decay width is saturated by the channels $\pi^+\gamma(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^+\bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

$\Gamma(\gamma(1S)\pi^0)/\Gamma_{\text{total}}$	Γ_2/Γ
<u>VALUE</u>	
not seen	KROKOVNY 13 BELL $e^+ e^- \rightarrow \gamma(1S)\pi^0\pi^0$

$\Gamma(\gamma(2S)\pi^+)/\Gamma_{\text{total}}$	Γ_3/Γ
<u>VALUE (units 10^{-2})</u>	
$3.62^{+0.76+0.79}_{-0.59-0.53}$	¹ GARMASH 16 BELL $e^+ e^- \rightarrow \pi^-\pi^+ B^+\bar{B}^{*0}, \pi^-\bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH 15 BELL $e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$
seen	BONDAR 12 BELL $e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$

¹ Assuming the Z_b(10610) decay width is saturated by the channels $\pi^+\gamma(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^+\bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M207W;LINKAGE=BO
NODE=M207W;LINKAGE=A
NODE=M207W;LINKAGE=BN

NODE=M207215;NODE=M207

DESIG=1
DESIG=9
DESIG=2
DESIG=10
DESIG=3
DESIG=11
DESIG=4
DESIG=5
DESIG=8
DESIG=6
DESIG=7

NODE=M207225
NODE=M207R01
NODE=M207R01

NODE=M207R01;LINKAGE=A

NODE=M207R09
NODE=M207R09

NODE=M207R02
NODE=M207R02

NODE=M207R02;LINKAGE=A

$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
seen	1 KROKOVNY	13 BELL	$e^+ e^- \rightarrow \Upsilon(2S)\pi^0\pi^0$	NODE=M207R10 NODE=M207R10

¹ Combined significance in $e^+ e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$, including systematics, of 6.5 σ .

 $\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
2.15^{+0.55+0.60}_{-0.42-0.43}	1 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$	NODE=M207R03 NODE=M207R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen GARMASH 15 BELL $e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

seen BONDAR 12 BELL $e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

¹ Assuming the $Z_b(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

 $\Gamma(\Upsilon(3S)\pi^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
seen	1 KROKOVNY	13 BELL	$e^+ e^- \rightarrow \Upsilon(3S)\pi^0\pi^0$	NODE=M207R11 NODE=M207R11

¹ Combined significance in $e^+ e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$, including systematics, of 6.5 σ .

 $\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
3.45^{+0.87+0.86}_{-0.71-0.63}	1 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$	NODE=M207R04 NODE=M207R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 2 MIZUK 16 BELL $e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$

seen 3 BONDAR 12 BELL $e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$

¹ Assuming the $Z_b(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

² Using $e^+ e^-$ energies near the $\Upsilon(11020)$.

³ Using $e^+ e^-$ energies near the $\Upsilon(10860)$.

 $\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
4.67^{+1.24+1.18}_{-1.00-0.89}	1 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$	NODE=M207R05 NODE=M207R05

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 2 MIZUK 16 BELL $e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$

seen 3 BONDAR 12 BELL $e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹ Assuming the $Z_b(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

² Using $e^+ e^-$ energies near the $\Upsilon(11020)$.

³ Using $e^+ e^-$ energies near the $\Upsilon(10860)$.

 $\Gamma(B^+ \bar{B}^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
not seen	GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^0$	NODE=M207R08 NODE=M207R08

 $[\Gamma(B^+ \bar{B}^0) + \Gamma(B^{*+} \bar{B}^0)]/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
85.6^{+1.5+1.5}_{-2.0-2.1}	357	1 GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- B^{*+} \bar{B}^0$	NODE=M207R00 NODE=M207R00

¹ Assuming the $Z_b(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + B^{*+} \bar{B}^0$, and using the results from BONDAR 12 and MIZUK 16. Using the mass and width of the $Z_b(10610)$ from BONDAR 12.

 $\Gamma(B^{*+} \bar{B}^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
not seen	GARMASH	16 BELL	$e^+ e^- \rightarrow \pi^- B^{*+} \bar{B}^0$	NODE=M207R06 NODE=M207R06

$$\frac{[\Gamma(B^+ \bar{B}^{*0}) + \Gamma(B^{*+} \bar{B}^0)] / [\Gamma(\gamma(1S)\pi^+) + \Gamma(\gamma(2S)\pi^+) + \Gamma(\gamma(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)]}{\Gamma_{10}/(\Gamma_1 + \Gamma_3 + \Gamma_5 + \Gamma_7 + \Gamma_8)}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.93^{+0.99+1.01}_{-0.69-0.73}$ 357 1 GARMASH 16 BELL $e^+ e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

1 Combined with the results of BONDAR 12 and MIZUK 16. Not independent from $Z_b(10610)$ branching fractions to $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$.

NODE=M207R07
NODE=M207R07

$Z_b(10610)$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)
KROKOVNY	13	PR D88 052016	P. Krovovny <i>et al.</i>	(BELLE Collab.)
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)

$Z_b(10650)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

I, G, C need confirmation.

was $X(10650)^\pm$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Observed by BONDAR 12 in $\gamma(5S)$ decays to $\gamma(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

NODE=M207

REFID=57446
REFID=57465
REFID=56811
REFID=55588
REFID=53963

NODE=M208

NODE=M208

$Z_b(10650)$ MASS

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
10652.2 ± 1.5		1 BONDAR	12 BELL	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$10656.7 \pm 5.0^{+1.1}_{-3.1}$	2 GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
$10650.7 \pm 1.5^{+0.5}_{-0.2}$	2 GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	
$10651.2 \pm 1.0^{+0.4}_{-0.3}$	2 GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	
$10657 \pm 6 \pm 3$	3 BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
$10651 \pm 2 \pm 3$	3 BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	
$10652 \pm 1 \pm 2$	3 BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	
$10654 \pm 3 \pm 1$	3 BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$	
$10651 \pm 2 \pm 3$	3 BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$	

NODE=M208M

NODE=M208M

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

NODE=M208M;LINKAGE=BO

NODE=M208M;LINKAGE=A

NODE=M208M;LINKAGE=BN

NODE=M208W

NODE=M208W

$Z_b(10650)$ WIDTH

<u>VALUE (MeV)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
11.5 ± 2.2		4 BONDAR	12 BELL	$e^+ e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$12.1^{+11.3+2.7}_{-4.8-0.6}$	5 GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
$14.2 \pm 3.7^{+0.9}_{-0.4}$	5 GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	
$9.3 \pm 2.2^{+0.3}_{-0.5}$	5 GARMASH	15 BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	
$16.3 \pm 9.8^{+6.0}_{-2.0}$	6 BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(1S)\pi^+\pi^-$	
$13.3 \pm 3.3^{+4.0}_{-3.0}$	6 BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(2S)\pi^+\pi^-$	
$8.4 \pm 2.0 \pm 2.0$	6 BONDAR	12 BELL	$e^+ e^- \rightarrow \gamma(3S)\pi^+\pi^-$	
$20.9^{+5.4+2.1}_{-4.7-5.7}$	6 BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$	
$19 \pm 7^{+11}_{-7}$	6 BONDAR	12 BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+\pi^-$	

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

$\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$					Γ_5/Γ	
VALUE (units 10^{-2})	DOCUMENT ID		TECN	COMMENT		NODE=M208R05 NODE=M208R05
14.7^{+3.2+2.8}_{-2.8-2.3}	13	GARMASH	16	BELL	$e^+ e^- \rightarrow \pi^- B^* + \bar{B}^{*0}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
possibly seen	14	MIZUK	16	BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+ \pi^-$	
seen	15	BONDAR	12	BELL	$e^+ e^- \rightarrow h_b(2P)\pi^+ \pi^-$	
13 Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^* + \bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.						NODE=M208R05;LINKAGE=C
14 Using $e^+ e^-$ energies near the $\gamma(11020)$.						NODE=M208R05;LINKAGE=A
15 Using $e^+ e^-$ energies near the $\gamma(10860)$.						NODE=M208R05;LINKAGE=B
$\Gamma(B^+ \bar{B}^0)/\Gamma_{\text{total}}$					Γ_6/Γ	
VALUE	DOCUMENT ID		TECN	COMMENT		NODE=M208R08 NODE=M208R08
not seen	GARMASH	16	BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^0$		
$[\Gamma(B^+ \bar{B}^{*0}) + \Gamma(B^{*+} \bar{B}^0)]/\Gamma_{\text{total}}$					Γ_7/Γ	
VALUE	DOCUMENT ID		TECN	COMMENT		NODE=M208R00 NODE=M208R00
not seen	GARMASH	16	BELL	$e^+ e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$		
$\Gamma(B^{*+} \bar{B}^{*0})/\Gamma_{\text{total}}$					Γ_8/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M208R06 NODE=M208R06
73.7^{+3.4+2.7}_{-4.4-3.5}	161	16 GARMASH	16	BELL	$e^+ e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$	
16 Assuming the $Z_b(10650)$ decay width is saturated by the channels $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+} \bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.						NODE=M208R06;LINKAGE=A
Using the mass and width of the $Z_b(10650)$ from BONDAR 12.						
$\Gamma(B^{*+} \bar{B}^{*0})/[\Gamma(\gamma(1S)\pi^+) + \Gamma(\gamma(2S)\pi^+) + \Gamma(\gamma(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)]$					$\Gamma_8/(\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4 + \Gamma_5)$	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M208R07 NODE=M208R07
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2.80 ^{+0.69+0.54} _{-0.40-0.36}	161	17 GARMASH	16	BELL	$e^+ e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$	
17 Combined with the results of BONDAR 12 and MIZUK 16. Not independent from $Z_b(10650)$ branching fractions to $\pi^+ \gamma(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+} \bar{B}^{*0}$.						NODE=M208R07;LINKAGE=A

$Z_b(10650)$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)

NODE=M208

REFID=57446
REFID=57465
REFID=56811
REFID=53963

NODE=M243

 $\Upsilon(10753)$ $I^G(J^{PC}) = ?^?(1^{--})$

OMITTED FROM SUMMARY TABLE

A candidate for $\Upsilon(3D)$ state or an exotic structure.

NODE=M243

Seen by MIZUK 19 in $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n=1,2,3$) with a significance of 5.2σ .

$\Upsilon(10753)$ MASS				NODE=M243M
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	NODE=M243M
$10752.7 \pm 5.9^{+0.7}_{-1.1}$	1 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.				NODE=M243M;LINKAGE=A
$\Upsilon(10753)$ WIDTH				NODE=M243W
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	NODE=M243W
$35.5^{+17.6+3.9}_{-11.3-3.3}$	1 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.				NODE=M243W;LINKAGE=A
$\Upsilon(10753)$ DECAY MODES				NODE=M243215;NODE=M243
Mode				
Γ_1	$\Upsilon(nS)\pi^+\pi^-$ ($n=1,2,3$)			DESIG=1
Γ_2	$\Upsilon(1S)\pi^+\pi^-$			DESIG=3
Γ_3	$\Upsilon(2S)\pi^+\pi^-$			DESIG=4
Γ_4	$\Upsilon(3S)\pi^+\pi^-$			DESIG=5
Γ_5	e^+e^-			DESIG=2
$\Upsilon(10753)$ BRANCHING RATIOS				NODE=M243225
$\Gamma(\Upsilon(nS)\pi^+\pi^- (n=1,2,3)) / \Gamma_{\text{total}}$			Γ_1/Γ	NODE=M243R01
$\Gamma(e^+e^-) \times \Gamma(\Upsilon(1S)\pi^+\pi^-) / \Gamma_{\text{total}}$			$\Gamma_5\Gamma_2/\Gamma$	NODE=M243R01
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	NODE=M243R00 NODE=M243R00
$\bullet\bullet\bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet\bullet\bullet$				
0.295 \pm 0.175	1,2 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.				NODE=M243R00;LINKAGE=A
² Reported as the range 0.12–0.47 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.				NODE=M243R00;LINKAGE=B
$\Gamma(e^+e^-) \times \Gamma(\Upsilon(2S)\pi^+\pi^-) / \Gamma_{\text{total}}$			$\Gamma_5\Gamma_3/\Gamma$	NODE=M243R02 NODE=M243R02
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
$\bullet\bullet\bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet\bullet\bullet$				
0.875 \pm 0.345	1,2 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.				NODE=M243R02;LINKAGE=A
² Reported as the range 0.53–1.22 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.				NODE=M243R02;LINKAGE=B

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$	$\Gamma_5\Gamma_4/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.235 ± 0.025	^{1,2} MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

² Reported as the range 0.21–0.26 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

$\Upsilon(10753)$ REFERENCES

MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)
-------	----	---------------	------------------------	-----------------

$\Upsilon(10860)$

$$\mathcal{I}^G(J^PC) = 0^-(1^{--})$$

$\Upsilon(10860)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9723 $^{+4000}_{-2300}$ OUR AVERAGE			Error includes scale factor of 1600.9. See the ideogram below. [10889.9 $^{+3.2}_{-2.6}$ MeV OUR 2019 AVERAGE]
10885.3 \pm 1.5 $^{+2.2}_{-0.9}$	¹ MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
36.6 \pm 4.5 $^{+0.5}_{-1.1}$	² MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
10884.7 \pm 3.6 $^{+8.9}_{-1.0}$	³ MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P,2P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. **• • •**

10881.8 \pm 1.0 $^{+1.2}_{-1.1}$	^{4,5} SANTEL	16	BELL $e^+e^- \rightarrow \text{hadrons}$
10891.1 \pm 3.2 $^{+1.2}_{-2.0}$	^{6,7} SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S,2S,3S)\pi^+\pi^-$
10879 \pm 3	^{8,9} CHEN	10	BELL $e^+e^- \rightarrow \text{hadrons}$
10888.4 \pm 2.7 $^{+1.2}_{-2.6}$	¹⁰ CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S,2S,3S)\pi^+\pi^-$
10876 \pm 2	⁸ AUBERT	09E	BABR $e^+e^- \rightarrow \text{hadrons}$
10869 \pm 2	¹¹ AUBERT	09E	BABR $e^+e^- \rightarrow \text{hadrons}$
10868 \pm 6 \pm 5	¹² BESSON	85	CLEO $e^+e^- \rightarrow \text{hadrons}$
10845 \pm 20	¹³ LOVELOCK	85	CUSB $e^+e^- \rightarrow \text{hadrons}$

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

² From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

³ From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

⁴ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).

⁵ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

⁶ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

⁷ Superseded by MIZUK 19.

⁸ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

⁹ The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.

NODE=M243R03
NODE=M243R03

NODE=M243R03;LINKAGE=A

NODE=M243R03;LINKAGE=B

NODE=M243

REFID=60090

NODE=M092

NODE=M092M

NODE=M092M

NEW

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M092M;LINKAGE=E

NODE=M092M;LINKAGE=G

NODE=M092M;LINKAGE=D

NODE=M092M;LINKAGE=A

NODE=M092M;LINKAGE=B

NODE=M092M;LINKAGE=C

NODE=M092M;LINKAGE=F

NODE=M092M;LINKAGE=AU

NODE=M092M;LINKAGE=CH


- 10 In a model where a flat nonresonant $\gamma(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.
- 11 In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.
- 12 Assuming four Gaussians with radiative tails and a single step in R .
- 13 In a coupled-channel model with three resonances and a smooth step in R .

NODE=M092M;LINKAGE=CE

NODE=M092M;LINKAGE=UB

NODE=M092M;LINKAGE=BE

NODE=M092M;LINKAGE=LO

 $\gamma(10860)$ MASS (MeV) $\gamma(10860)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
41 $^{+13}_{-21}$ OUR AVERAGE [51 $^{+6}_{-7}$ MeV OUR 2019 AVERAGE]			
40.6$^{+12.7}_{-8.0}$$^{+1.1}_{-19.1}$	1 MIZUK	16 BELL	$e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
48.5 $^{+1.9}_{-1.8}$ $^{+2.0}_{-2.8}$	2,3 SANTEL	16 BELL	$e^+e^- \rightarrow$ hadrons
53.7 $^{+7.1}_{-5.6}$ $^{+1.3}_{-5.4}$	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow \gamma(1S, 2S, 3S)\pi^+\pi^-$
46 $^{+9}_{-7}$	6,7 CHEN	10 BELL	$e^+e^- \rightarrow$ hadrons
30.7 $^{+8.3}_{-7.0}$ $^{+3.1}_{-}$	8 CHEN	10 BELL	$e^+e^- \rightarrow \gamma(1S, 2S, 3S)\pi^+\pi^-$
43 $^{+4}_{-4}$	6 AUBERT	09E BABR	$e^+e^- \rightarrow$ hadrons
74 $^{+4}_{-4}$	9 AUBERT	09E BABR	$e^+e^- \rightarrow$ hadrons
112 $^{+17}_{-15}$ $^{+23}_{-}$	10 BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
110 $^{+15}_{-}$	11 LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

NODE=M092W

NODE=M092W

NEW

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M092W;LINKAGE=D

¹ From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\gamma(10860)$ and $\gamma(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

² From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\gamma(10860)$ and $\gamma(11020)$, one relative phase, and one decoherence coefficient).

³ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

⁴ From a simultaneous fit to the $\gamma(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\gamma(10860)$ and $\gamma(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M092W;LINKAGE=A

NODE=M092W;LINKAGE=B

NODE=M092W;LINKAGE=C

- 5 Superseded by MIZUK 19.
 6 In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.
 7 The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.
 8 In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.
 9 In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.
 10 Assuming four Gaussians with radiative tails and a single step in R .
 11 In a coupled-channel model with three resonances and a smooth step in R .

$\Upsilon(10860)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 B\bar{B}X$	(76.2 ± 2.7) %	
$\Gamma_2 B\bar{B}$	(5.5 ± 1.0) %	
$\Gamma_3 B\bar{B}^* + \text{c.c.}$	(13.7 ± 1.6) %	
$\Gamma_4 B^*\bar{B}^*$	(38.1 ± 3.4) %	
$\Gamma_5 B\bar{B}^{(*)}\pi$	< 19.7 %	90%
$\Gamma_6 B\bar{B}\pi$	(0.0 ± 1.2) %	
$\Gamma_7 B^*\bar{B}\pi + B\bar{B}^*\pi$	(7.3 ± 2.3) %	
$\Gamma_8 B^*\bar{B}^*\pi$	(1.0 ± 1.4) %	
$\Gamma_9 B\bar{B}\pi\pi$	< 8.9 %	90%
$\Gamma_{10} B_s^{(*)}\bar{B}_s^{(*)}$	(20.1 ± 3.1) %	
$\Gamma_{11} B_s\bar{B}_s$	(5 ± 5) $\times 10^{-3}$	
$\Gamma_{12} B_s B_s^* + \text{c.c.}$	(1.35 ± 0.32) %	
$\Gamma_{13} B_s^*\bar{B}_s^*$	(17.6 ± 2.7) %	
Γ_{14} no open-bottom	(3.8 ± 5.0) %	
$\Gamma_{15} e^+ e^-$	(7.6 ± 4.0) $\times 10^{-6}$	
$\Gamma_{16} K^*(892)^0\bar{K}^0$	< 1.0 $\times 10^{-5}$	90%
$\Gamma_{17} \Upsilon(1S)\pi^+\pi^-$	(5.3 ± 0.6) $\times 10^{-3}$	
$\Gamma_{18} \Upsilon(2S)\pi^+\pi^-$	(7.8 ± 1.3) $\times 10^{-3}$	
$\Gamma_{19} \Upsilon(3S)\pi^+\pi^-$	(4.8 ± 1.9) $\times 10^{-3}$	
$\Gamma_{20} \Upsilon(1S)K^+K^-$	(6.1 ± 1.8) $\times 10^{-4}$	
$\Gamma_{21} \eta \Upsilon_J(1D)$	(4.8 ± 1.1) $\times 10^{-3}$	
$\Gamma_{22} h_b(1P)\pi^+\pi^-$	(3.5 ± 1.0) $\times 10^{-3}$	
$\Gamma_{23} h_b(2P)\pi^+\pi^-$	(5.7 ± 1.7) $\times 10^{-3}$	
$\Gamma_{24} \chi_{bJ}(1P)\pi^+\pi^-\pi^0$	(2.5 ± 2.3) $\times 10^{-3}$	
$\Gamma_{25} \chi_{b0}(1P)\pi^+\pi^-\pi^0$	< 6.3 $\times 10^{-3}$	90%
$\Gamma_{26} \chi_{b0}(1P)\omega$	< 3.9 $\times 10^{-3}$	90%
$\Gamma_{27} \chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	< 4.8 $\times 10^{-3}$	90%
$\Gamma_{28} \chi_{b1}(1P)\pi^+\pi^-\pi^0$	(1.85 ± 0.33) $\times 10^{-3}$	
$\Gamma_{29} \chi_{b1}(1P)\omega$	(1.57 ± 0.30) $\times 10^{-3}$	
$\Gamma_{30} \chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	(5.2 ± 1.9) $\times 10^{-4}$	
$\Gamma_{31} \chi_{b2}(1P)\pi^+\pi^-\pi^0$	(1.17 ± 0.30) $\times 10^{-3}$	
$\Gamma_{32} \chi_{b2}(1P)\omega$	(6.0 ± 2.7) $\times 10^{-4}$	
$\Gamma_{33} \chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega}$	(6 ± 4) $\times 10^{-4}$	
$\Gamma_{34} \gamma X_b \rightarrow \gamma \Upsilon(1S)\omega$	< 3.8 $\times 10^{-5}$	90%

Inclusive Decays.

These decay modes are submodes of one or more of the decay modes above.

$\Gamma_{35} \phi$ anything	(13.8 ± 2.4) %
$\Gamma_{36} D^0$ anything + c.c.	(108 ± 8) %
$\Gamma_{37} D_s$ anything + c.c.	(46 ± 6) %
$\Gamma_{38} J/\psi$ anything	(2.06 ± 0.21) %
$\Gamma_{39} B^0$ anything + c.c.	(77 ± 8) %
$\Gamma_{40} B^+$ anything + c.c.	(72 ± 6) %

NODE=M092W;LINKAGE=E

NODE=M092W;LINKAGE=AU

NODE=M092W;LINKAGE=CH

NODE=M092W;LINKAGE=CE

NODE=M092W;LINKAGE=UB

NODE=M092W;LINKAGE=BE

NODE=M092W;LINKAGE=LO

NODE=M092215;NODE=M092

DESIG=9

DESIG=2

DESIG=3

DESIG=4

DESIG=10

DESIG=23

DESIG=24

DESIG=25

DESIG=11

DESIG=16

DESIG=5

DESIG=7

DESIG=8

DESIG=28

DESIG=1

DESIG=29

DESIG=17

DESIG=18

DESIG=19

DESIG=20

DESIG=40

DESIG=26

DESIG=27

DESIG=41

DESIG=30

DESIG=31

DESIG=32

DESIG=33

DESIG=34

DESIG=35

DESIG=36

DESIG=37

DESIG=38

DESIG=39

NODE=M092;CLUMP=I

NODE=M092

DESIG=12

DESIG=13

DESIG=6

DESIG=14

DESIG=21

DESIG=22

$\Upsilon(10860)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_{15}
0.31 ±0.07 OUR AVERAGE			Error includes scale factor of 1.3.	
0.22 ±0.05 ±0.07	BESSON 85	CLEO	$e^+e^- \rightarrow$ hadrons	
0.365±0.070	LOVELOCK 85	CUSB	$e^+e^- \rightarrow$ hadrons	

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{15}\Gamma_{17}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.09±0.34	1,2 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

² Reported as the range 0.75–1.43 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{15}\Gamma_{18}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.58±1.22	1,2 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

² Reported as the range 1.35–3.80 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_{15}\Gamma_{19}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.73±0.30	1,2 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$	

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

² Reported as the range 0.43–1.03 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

$\Upsilon(10860)$ BRANCHING RATIOS

"OUR EVALUATION" is obtained based on averages of rescaled data listed below. The averages and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <http://www.slac.stanford.edu/xorg/hflav/>.

$\Gamma(B\bar{B}X)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.762^{+0.027}_{-0.043} OUR EVALUATION					

0.71 ±0.06 OUR AVERAGE

0.737±0.032±0.051	1063	¹ DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$	
0.589±0.100±0.092		² HUANG	07 CLEO	$\Upsilon(5S) \rightarrow$ hadrons	

¹ Not independent of DRUTSKOY 10 values for $\Upsilon(5S) \rightarrow B^{\pm,0}$ anything.

² Using measurements or limits from AQUINES 06.

$\Gamma(B\bar{B})/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
5.5^{+1.0}_{-0.9}±0.4		¹ DRUTSKOY	10 BELL	$\Upsilon(5S) \rightarrow B^+ X, B^0 X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.8	90	² HUANG	07 CLEO	$\Upsilon(5S) \rightarrow$ hadrons
-------	----	--------------------	---------	------------------------------------

¹ Assuming isospin conservation.

² Using measurements or limits from AQUINES 06.

$\Gamma(B\bar{B})/\Gamma(B\bar{B}X)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
<0.22	90	AQUINES	06 CLE3	$\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092220

NODE=M092W1

NODE=M092W1

NODE=M092R50

NODE=M092R50

NODE=M092R50;LINKAGE=A

NODE=M092R50;LINKAGE=B

NODE=M092R51

NODE=M092R51

NODE=M092R51;LINKAGE=A

NODE=M092R51;LINKAGE=B

NODE=M092R52

NODE=M092R52

NODE=M092R52;LINKAGE=A

NODE=M092R52;LINKAGE=B

NODE=M092230

NODE=M092230

NODE=M092R13

NODE=M092R13

→ UNCHECKED ←

NODE=M092R13;LINKAGE=DR

NODE=M092R13;LINKAGE=HU

NODE=M092R16

NODE=M092R16

NODE=M092R16;LINKAGE=DR

NODE=M092R16;LINKAGE=HU

NODE=M092R05

NODE=M092R05

$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma_{\text{total}}$

VALUE
0.137±0.016 OUR AVERAGE

$0.137 \pm 0.013 \pm 0.011$
 $0.143 \pm 0.053 \pm 0.027$

¹ Assuming isospin conservation.

² Using measurements or limits from AQUINES 06.

 Γ_3/Γ

NODE=M092R15
NODE=M092R15

 $\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma(B\bar{B}X)$

VALUE
0.24±0.09±0.03

EVTS
10

DOCUMENT ID TECN COMMENT

¹ DRUTSKOY 10 BELL $\gamma(5S) \rightarrow B^+ X, B^0 X$
² HUANG 07 CLEO $\gamma(5S) \rightarrow \text{hadrons}$

 Γ_3/Γ_1

NODE=M092R15;LINKAGE=DR

NODE=M092R15;LINKAGE=HU

NODE=M092R06
NODE=M092R06

 $\Gamma(B^*\bar{B}^*)/\Gamma_{\text{total}}$

VALUE
0.381±0.034 OUR AVERAGE

$0.375^{+0.021}_{-0.019} \pm 0.030$
 $0.436 \pm 0.083 \pm 0.072$

¹ Assuming isospin conservation.

² Using measurements or limits from AQUINES 06.

 Γ_4/Γ

NODE=M092R14
NODE=M092R14

 $\Gamma(B^*\bar{B}^*)/\Gamma(B\bar{B}X)$

VALUE
0.74±0.15±0.08

EVTS
31

DOCUMENT ID TECN COMMENT

AQUINES 06 CLE3 $\gamma(5S) \rightarrow \text{hadrons}$

 Γ_4/Γ_1

NODE=M092R07
NODE=M092R07

 $\Gamma(B\bar{B}(\pi)/\Gamma_{\text{total}}$

VALUE
<0.197

CL%
90

DOCUMENT ID TECN COMMENT

¹ HUANG 07 CLEO $\gamma(5S) \rightarrow \text{hadrons}$

 Γ_5/Γ

NODE=M092R17
NODE=M092R17

¹ Using measurements or limits from AQUINES 06.

 $\Gamma(B\bar{B}(\pi)/\Gamma(B\bar{B}X)$

VALUE
<0.32

CL%
90

DOCUMENT ID TECN COMMENT

AQUINES 06 CLE3 $\gamma(5S) \rightarrow \text{hadrons}$

 Γ_5/Γ_1

NODE=M092R08
NODE=M092R08

 $\Gamma(B\bar{B}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})
0.0±1.2±0.3

EVTS
0

DOCUMENT ID TECN COMMENT

¹ DRUTSKOY 10 BELL $\gamma(5S) \rightarrow B^{+,0} \pi^- X$

 Γ_6/Γ

NODE=M092R28
NODE=M092R28

¹ Assuming isospin conservation.

 $[\Gamma(B^*\bar{B}\pi) + \Gamma(B\bar{B}^*\pi)]/\Gamma_{\text{total}}$

VALUE (units 10^{-2})
7.3±2.3±0.8

EVTS
38

DOCUMENT ID TECN COMMENT

¹ DRUTSKOY 10 BELL $\gamma(5S) \rightarrow B^{+,0} \pi^- X$

 Γ_7/Γ

NODE=M092R29
NODE=M092R29

¹ Assuming isospin conservation.

 $\Gamma(B^*\bar{B}^*\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})
1.0±1.4±0.4

EVTS
5

DOCUMENT ID TECN COMMENT

¹ DRUTSKOY 10 BELL $\gamma(5S) \rightarrow B^{+,0} \pi^- X$

 Γ_8/Γ

NODE=M092R30
NODE=M092R30

¹ Assuming isospin conservation.

 $\Gamma(B\bar{B}\pi\pi)/\Gamma_{\text{total}}$

VALUE
<0.089

CL%
90

DOCUMENT ID TECN COMMENT

¹ HUANG 07 CLEO $\gamma(5S) \rightarrow \text{hadrons}$

 Γ_9/Γ

NODE=M092R18
NODE=M092R18

¹ Using measurements or limits from AQUINES 06.

 $\Gamma(B\bar{B}\pi\pi)/\Gamma(B\bar{B}X)$

VALUE
<0.14

CL%
90

DOCUMENT ID TECN COMMENT

AQUINES 06 CLE3 $\gamma(5S) \rightarrow \text{hadrons}$

 Γ_9/Γ_1

NODE=M092R09
NODE=M092R09

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma_{\text{total}}$	$\Gamma_{10}/\Gamma = (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT

0.201^{+0.030}_{-0.031} OUR EVALUATION**0.189^{+0.027}_{-0.021} OUR AVERAGE**

0.172 \pm 0.030	1 ESEN	13 BELL	$\Gamma(5S) \rightarrow D^0 X, D_s X$
0.21 $^{+0.06}_{-0.03}$	2 HUANG	07 CLEO	$\Gamma(5S) \rightarrow D_s X$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.180 \pm 0.013 \pm 0.032	3 DRUTSKOY	07 BELL	$\Gamma(5S) \rightarrow D^0 X, D_s X$
0.160 \pm 0.026 \pm 0.058	4 ARTUSO	05B CLEO	$e^+ e^- \rightarrow D_s X$

¹ Supersedes DRUTSKOY 07.² Supersedes ARTUSO 05B. Combining inclusive ϕ , D_s , and B measurements. Using $B(D_s^+ \rightarrow \phi\pi^+) = 4.4 \pm 0.6\%$ from PDG 06.³ Using $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6)\%$ from PDG 06.⁴ Uses a model-dependent estimate $B(B_s \rightarrow D_s X) = (92 \pm 11)\%$.

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma(B\bar{B}X)$	Γ_{10}/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT

0.264^{+0.052}_{-0.045} OUR EVALUATION

$\Gamma(B_s^*\bar{B}_s^*)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$	$\Gamma_{13}/\Gamma_{10} = \Gamma_{13}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$			
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT

87.8 \pm 1.5 OUR AVERAGE

87.0 \pm 1.7	1,2 ESEN	13 BELL	$B_s^0 \rightarrow D_s^- \pi^+$
90.5 \pm 3.2 \pm 0.1	227 2,3 LI	12 BELL	$B_s^0 \rightarrow J/\psi\eta^{(')}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
90.1 $^{+3.8}_{-4.0}$ \pm 0.2	4 LOUVOT	09 BELL	$10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$
93 $^{+7}_{-9}$ \pm 1	4 DRUTSKOY	07A BELL	Superseded by LOUVOT 09

¹ Supersedes LOUVOT 09.² With $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$.³ The ratios $N(B_s^*\bar{B}_s^*) / N(B_s^{(*)}\bar{B}_s^{(*)})$ and $N(B_s^*\bar{B}_s^0) / N(B_s^{(*)}\bar{B}_s^{(*)})$ are measured with a correlation coefficient of -0.72 .⁴ From a measurement of $\sigma(e^+ e^- \rightarrow B_s^*\bar{B}_s^*) / \sigma(e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)})$ at $\sqrt{s} = 10.86$ GeV.

$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$	$\Gamma_{11}/\Gamma_{10} = \Gamma_{11}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$		
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT

2.6$^{+2.6}_{-2.5}$	LOUVOT	09 BELL	$10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$
---------------------------------------	--------	---------	--

$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^*\bar{B}_s^*)$	Γ_{11}/Γ_{13}			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT

<0.16	90	BONVICINI	06 CLE3	$e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$
-----------------	----	-----------	---------	--

$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$	$\Gamma_{12}/\Gamma_{10} = \Gamma_{12}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$			
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT

6.7\pm1.2 OUR AVERAGE	1,2 ESEN	13 BELL	$B_s^0 \rightarrow D_s^- \pi^+$
7.3 \pm 1.4	227 2,3 LI	12 BELL	$B_s^0 \rightarrow J/\psi\eta^{(')}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.3 $^{+3.3}_{-3.0}$ \pm 0.1	LOUVOT	09 BELL	$10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$
--------------------------------	--------	---------	--

¹ Supersedes LOUVOT 09.² With $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$.³ The ratios $N(B_s^*\bar{B}_s^*) / N(B_s^{(*)}\bar{B}_s^{(*)})$ and $N(B_s^*\bar{B}_s^0) / N(B_s^{(*)}\bar{B}_s^{(*)})$ are measured with a correlation coefficient of -0.72 .NODE=M092R01
NODE=M092R01

→ UNCHECKED ←

NODE=M092R01;LINKAGE=ES
NODE=M092R01;LINKAGE=HUNODE=M092R01;LINKAGE=DR
NODE=M092R01;LINKAGE=ARNODE=M092R34
NODE=M092R34

→ UNCHECKED ←

NODE=M092R19
NODE=M092R19

NODE=M092R19;LINKAGE=ES

NODE=M092R19;LINKAGE=IL

NODE=M092R19;LINKAGE=LI

NODE=M092R19;LINKAGE=DR

NODE=M092R24
NODE=M092R24NODE=M092R03
NODE=M092R03NODE=M092R25
NODE=M092R25

NODE=M092R25;LINKAGE=ES

NODE=M092R25;LINKAGE=IL

NODE=M092R25;LINKAGE=LI

$\Gamma(B_s \bar{B}_s^* + \text{c.c.})/\Gamma(B_s^* \bar{B}_s^*)$	Γ_{12}/Γ_{13}	NODE=M092R04 NODE=M092R04
<u>VALUE</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
<0.16 90	BONVICINI 06 CLE3 $e^+ e^-$	
$\Gamma(\text{no open-bottom})/\Gamma_{\text{total}}$	Γ_{14}/Γ	NODE=M092R33 NODE=M092R33
<u>VALUE</u>	<u>DOCUMENT ID</u>	
0.038 ^{+0.051} _{-0.005} OUR EVALUATION		→ UNCHECKED ←
$\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma_{\text{total}}$	Γ_{16}/Γ	NODE=M092R35 NODE=M092R35
<u>VALUE</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
<1.0 × 10 ⁻⁵ 90	SHEN 13A BELL $e^+ e^- \rightarrow K^*(892)^0 \bar{K}^0$	
$\Gamma(\eta \tau_J(1D))/\Gamma_{\text{total}}$	Γ_{21}/Γ	NODE=M092R48 NODE=M092R48
<u>VALUE (units 10⁻³)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
4.82±0.92±0.67	¹ TAMPONI 18 BELL $e^+ e^- \rightarrow \tau(5S) \rightarrow \eta X$	
1 Mainly $J = 2$, assumes no continuum contribution under $\tau(5S)$.		
$\Gamma(\tau(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{17}/Γ	NODE=M092R20 NODE=M092R20
<u>VALUE (units 10⁻³)</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
5.3±0.3±0.5 325	¹ CHEN 08 BELL 10.87 $e^+ e^- \rightarrow \tau(1S)\pi^+\pi^-$	
1 Assuming that the observed events are solely due to the $\tau(5S)$ resonance.		
$\Gamma(\tau(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{18}/Γ	NODE=M092R21 NODE=M092R21
<u>VALUE (units 10⁻³)</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
7.8±0.6±1.1 186	¹ CHEN 08 BELL 10.87 $e^+ e^- \rightarrow \tau(2S)\pi^+\pi^-$	
1 Assuming that the observed events are solely due to the $\tau(5S)$ resonance.		
$\Gamma(\tau(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$	Γ_{19}/Γ	NODE=M092R22 NODE=M092R22
<u>VALUE (units 10⁻³)</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
4.8 ^{+1.8} _{-1.5} ±0.7 10	¹ CHEN 08 BELL 10.87 $e^+ e^- \rightarrow \tau(3S)\pi^+\pi^-$	
1 Assuming that the observed events are solely due to the $\tau(5S)$ resonance.		
$\Gamma(\tau(1S)K^+K^-)/\Gamma_{\text{total}}$	Γ_{20}/Γ	NODE=M092R23 NODE=M092R23
<u>VALUE (units 10⁻⁴)</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
6.1 ^{+1.6} _{-1.4} ±1.0 20	¹ CHEN 08 BELL 10.87 $e^+ e^- \rightarrow \tau(1S)K^+K^-$	
1 Assuming that the observed events are solely due to the $\tau(5S)$ resonance.		
$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(\tau(2S)\pi^+\pi^-)$	Γ_{22}/Γ_{18}	NODE=M092R31 NODE=M092R31
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
0.45±0.08 ^{+0.07} _{-0.12}	ADACHI 12 BELL 10.86 $e^+ e^- \rightarrow \text{hadrons}$	
$\Gamma(h_b(2P)\pi^+\pi^-)/\Gamma(\tau(2S)\pi^+\pi^-)$	Γ_{23}/Γ_{18}	NODE=M092R32 NODE=M092R32
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
0.77±0.08 ^{+0.22} _{-0.17}	ADACHI 12 BELL 10.86 $e^+ e^- \rightarrow \text{hadrons}$	
$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(h_b(2P)\pi^+\pi^-)$	Γ_{22}/Γ_{23}	NODE=M092R00 NODE=M092R00
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
0.616±0.052±0.017	MIZUK 16 BELL $e^+ e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$	
$\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_{24}/Γ	NODE=M092R49 NODE=M092R49
<u>VALUE (units 10⁻³)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
2.5±0.6±2.2	YIN 18 BELL $e^+ e^- \rightarrow \text{hadrons}$	
$\Gamma(\chi_{b0}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_{25}/Γ	NODE=M092R36 NODE=M092R36
<u>VALUE</u> <u>CL%</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>	
<6.3 × 10 ⁻³ 90	¹ HE 14 BELL $\tau(5S) \rightarrow \pi^+\pi^-\pi^0\gamma \tau(1S)$	
1 Assuming that all the $b\bar{b}$ events are from $\tau(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$ from ESEN 13. Correlated with other results from HE 14.		

$\Gamma(\chi_{b0}(1P)\omega)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-3}$	90	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{26}/Γ

NODE=M092R37
NODE=M092R37

 $\Gamma(\chi_{b0}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.8 \times 10^{-3}$	90	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{27}/Γ

NODE=M092R38
NODE=M092R38

 $\Gamma(\chi_{b1}(1P)\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.85 \pm 0.23 \pm 0.23$	80	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{28}/Γ

NODE=M092R39
NODE=M092R39

 $\Gamma(\chi_{b1}(1P)\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.57 \pm 0.22 \pm 0.21$	60	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{29}/Γ

NODE=M092R40
NODE=M092R40

 $\Gamma(\chi_{b1}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.52 \pm 0.15 \pm 0.11$	24	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{30}/Γ

NODE=M092R41
NODE=M092R41

 $\Gamma(\chi_{b2}(1P)\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.17 \pm 0.27 \pm 0.14$	29	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{31}/Γ

NODE=M092R42
NODE=M092R42

 $\Gamma(\chi_{b2}(1P)\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.60 \pm 0.23 \pm 0.15$	13	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{32}/Γ

NODE=M092R43
NODE=M092R43

 $\Gamma(\chi_{b2}(1P)\omega)/\Gamma(\chi_{b1}(1P)\omega)$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$0.38 \pm 0.16 \pm 0.09$ 1 HE 14 BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Accounting for correlated systematics.

NODE=M092R44;LINKAGE=A

 $\Gamma(\chi_{b2}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.61 \pm 0.22 \pm 0.28$	16	1 HE	14	BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Assuming that all the $b\bar{b}$ events are from $\gamma(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.

 Γ_{33}/Γ

NODE=M092R45
NODE=M092R45

 $\Gamma(\chi_{b2}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma(\chi_{b1}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$1.20 \pm 0.55 \pm 0.65$ 1 HE 14 BELL $\gamma(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \gamma(1S)$

¹ Accounting for correlated systematics.

NODE=M092R46
NODE=M092R46

NODE=M092R46;LINKAGE=A

$\Gamma(\gamma X_b \rightarrow \gamma \Upsilon(1S)\omega)/\Gamma_{\text{total}}$	Γ_{34}/Γ	NODE=M092R47 NODE=M092R47
<u>VALUE</u> $<3.8 \times 10^{-5}$	<u>CL%</u> 90	<u>DOCUMENT ID</u> 1 HE
		<u>TECN</u> BELL
$\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$		
1 Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14. For a state X_b with mass between $10.55 \text{ GeV}/c^2$ and $10.65 \text{ GeV}/c^2$, the obtained 90% upper limit as a function of m_{X_b} varies from 2.6×10^{-5} to 3.8×10^{-5} .		
$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$	Γ_{35}/Γ	NODE=M092R12 NODE=M092R12
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$0.138 \pm 0.007 \pm 0.023$	HUANG	07 CLEO
$\Upsilon(5S) \rightarrow \phi X$		
$\Gamma(D^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{36}/Γ	NODE=M092R10 NODE=M092R10
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$1.076 \pm 0.040 \pm 0.068$	DRUTSKOY	07 BELL
$\Upsilon(5S) \rightarrow D^0 X$		
$\Gamma(D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{37}/Γ	NODE=M092R02 NODE=M092R02
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
0.46 ± 0.06 OUR AVERAGE		
0.472 $\pm 0.024 \pm 0.072$	1 DRUTSKOY	07 BELL
0.44 $\pm 0.09 \pm 0.04$	2 ARTUSO	05B CLE3
$\Upsilon(5S) \rightarrow D_s X$		
$e^+ e^- \rightarrow D_X X$		
1 Using $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6)\%$ from PDG 06.		
2 ARTUSO 05B reports $[\Gamma(\Upsilon(10860) \rightarrow D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = 0.0198 \pm 0.0019 \pm 0.0038$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.		
$\Gamma(J/\psi \text{ anything})/\Gamma_{\text{total}}$	Γ_{38}/Γ	NODE=M092R11 NODE=M092R11
<u>VALUE (units 10^{-2})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
$2.060 \pm 0.160 \pm 0.134$	DRUTSKOY	07 BELL
$\Upsilon(5S) \rightarrow J/\psi X$		
$\Gamma(B^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{39}/Γ	NODE=M092R26 NODE=M092R26
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$0.770 \pm 0.058 \pm 0.061$	352	DRUTSKOY
$\Upsilon(5S) \rightarrow B^0 X$		
$\Gamma(B^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$	Γ_{40}/Γ	NODE=M092R27 NODE=M092R27
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>
$0.721 \pm 0.039 \pm 0.050$	711	DRUTSKOY
$\Upsilon(5S) \rightarrow B^+ X$		

R(10860) REFERENCES

MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60090
TAMPONI	18	EPJ C78 633	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=59195
YIN	18	PR D98 091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)	REFID=59468
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
HE	14	PRL 113 142001	X.H. He <i>et al.</i>	(BELLE Collab.)	REFID=55927
ESEN	13	PR D87 031101	S. Esen <i>et al.</i>	(BELLE Collab.)	REFID=54894
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
LI	12	PRL 108 181808	J. Li <i>et al.</i>	(BELLE Collab.)	REFID=54116
CHEN	10	PR D82 091106	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53531
DRUTSKOY	10	PR D81 112003	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=53358
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661
LOUVOT	09	PRL 102 021801	R. Louvot <i>et al.</i>	(BELLE Collab.)	REFID=52646
CHEN	08	PRL 100 112001	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=52153
DRUTSKOY	07	PRL 98 052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51621
DRUTSKOY	07A	PR D76 012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51852
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
AQUINES	06	PRL 96 152001	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51106
BONVICINI	06	PRL 96 022002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=50995
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369

NODE=M093

 $\gamma(11020)$ $I^G(J^{PC}) = 0^-(1^- -)$ **$\gamma(11020)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
11000 ± 4 OUR AVERAGE			
[10992.9 ± 10.0 MeV OUR 2019 AVERAGE]			
11000.00 ± 4.0 ± 1.0	1 MIZUK	19 BELL	$e^+ e^- \rightarrow \gamma(1S, 2S, 3S)\pi^+\pi^-$
10999.0 ± 7.3 ± 16.9	2 MIZUK	16 BELL	$e^+ e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
11003.0 ± 1.1 ± 0.9	3,4 SANTEL	16 BELL	$e^+ e^- \rightarrow \text{hadrons}$
10987.5 ± 6.4 ± 9.1	5,6 SANTEL	16 BELL	$e^+ e^- \rightarrow \gamma(1S, 2S, 3S)\pi^+\pi^-$
10996 ± 2	7 AUBERT	09E BABR	$e^+ e^- \rightarrow \text{hadrons}$
11019 ± 5 ± 7	BESSON	85 CLEO	$e^+ e^- \rightarrow \text{hadrons}$
11020 ± 30	LOVELOCK	85 CUSB	$e^+ e^- \rightarrow \text{hadrons}$

¹ From a simultaneous fit to the $\gamma(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\gamma(10860)$.

² From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\gamma(10860)$ and $\gamma(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

³ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\gamma(10860)$ and $\gamma(11020)$, one relative phase, and one decoherence coefficient).

⁴ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

⁵ From a simultaneous fit to the $\gamma(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\gamma(10860)$ and $\gamma(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

⁶ Superseded by MIZUK 19.

⁷ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M093M

NODE=M093M

NEW

OCCUR=2

NODE=M093M;LINKAGE=E

NODE=M093M;LINKAGE=D

NODE=M093M;LINKAGE=A

NODE=M093M;LINKAGE=B

NODE=M093M;LINKAGE=C

NODE=M093M;LINKAGE=F
NODE=M093M;LINKAGE=AU

NODE=M093W

NODE=M093W

NEW

OCCUR=2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
24 ± 8 OUR AVERAGE			
[49 ± 15 MeV OUR 2019 AVERAGE]			
23.8 ± 8.0 ± 0.7	8 MIZUK	19 BELL	$e^+ e^- \rightarrow \gamma(nS)\pi^+\pi^-$
27 ± 27 ± 5	9 MIZUK	16 BELL	$e^+ e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
39.3 ± 1.7 ± 1.3	10,11 SANTEL	16 BELL	$e^+ e^- \rightarrow \text{hadrons}$
61 ± 9 ± 2	12,13 SANTEL	16 BELL	$e^+ e^- \rightarrow \gamma(1S, 2S, 3S)\pi^+\pi^-$
37 ± 3	14 AUBERT	09E BABR	$e^+ e^- \rightarrow \text{hadrons}$
61 ± 13 ± 22	BESSON	85 CLEO	$e^+ e^- \rightarrow \text{hadrons}$
90 ± 20	LOVELOCK	85 CUSB	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M093W

- 8 From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.
- 9 From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.
- 10 From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).
- 11 Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.
- 12 From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n=1, 2, 3$ cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.
- 13 Superseded by MIZUK 19.
- 14 In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M093W;LINKAGE=E

NODE=M093W;LINKAGE=D

NODE=M093W;LINKAGE=A

NODE=M093W;LINKAGE=B

NODE=M093W;LINKAGE=C

NODE=M093W;LINKAGE=F
NODE=M093W;LINKAGE=AU

NODE=M093215;NODE=M093

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 e^+ e^-$	$(5.4^{+1.9}_{-2.1}) \times 10^{-6}$
$\Gamma_2 \Upsilon(1S)\pi^+\pi^-$	
$\Gamma_3 \Upsilon(2S)\pi^+\pi^-$	
$\Gamma_4 \Upsilon(3S)\pi^+\pi^-$	
$\Gamma_5 \chi_{bJ}(1P)\pi^+\pi^-\pi^0$	$(9^{+9}_{-8}) \times 10^{-3}$
$\Gamma_6 \chi_{b1}(1P)\pi^+\pi^-\pi^0$	seen
$\Gamma_7 \chi_{b2}(1P)\pi^+\pi^-\pi^0$	seen

 $\Upsilon(11020)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	Γ_1
VALUE (keV)	
0.130 ± 0.030 OUR AVERAGE	
$0.095 \pm 0.03 \pm 0.035$	BESSON 85 CLEO $e^+e^- \rightarrow$ hadrons
0.156 ± 0.040	LOVELOCK 85 CUSB $e^+e^- \rightarrow$ hadrons

DESIG=1

DESIG=5

DESIG=6

DESIG=7

DESIG=2

DESIG=3

DESIG=4

NODE=M093220

NODE=M093W1

NODE=M093W1

NODE=M093R04

NODE=M093R04

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_2/\Gamma$
VALUE (eV)	
15,16 MIZUK	19 BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

NODE=M093R04;LINKAGE=A

NODE=M093R04;LINKAGE=B

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_3/\Gamma$
VALUE (eV)	
17,18 MIZUK	19 BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

NODE=M093R05

NODE=M093R05

NODE=M093R05;LINKAGE=A

NODE=M093R05;LINKAGE=B

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 15 From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.
- 16 Reported as the range 0.38–0.54 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_4/\Gamma$	NODE=M093R06 NODE=M093R06
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •		
0.33±0.16	19,20 MIZUK	19 BELL $e^+e^- \rightarrow \gamma(nS)\pi^+\pi^-$
19 From a simultaneous fit to the $\gamma(nS)\pi^+\pi^-$, $n = 1, 2, 3$ cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.		NODE=M093R06;LINKAGE=A
20 Reported as the range 0.17–0.49 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.		NODE=M093R06;LINKAGE=B
$\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_5/Γ	NODE=M093R00 NODE=M093R00
<u>VALUE (units 10^{-3})</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
8.7±4.3^{+7.6}_{-6.6}	YIN	18 BELL $e^+e^- \rightarrow \text{hadrons}$
$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_6/Γ	NODE=M093R01 NODE=M093R01
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
seen	YIN	18 BELL $e^+e^- \rightarrow \text{hadrons}$
$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	Γ_7/Γ	NODE=M093R02 NODE=M093R02
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
seen	YIN	18 BELL $e^+e^- \rightarrow \text{hadrons}$
$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)$	Γ_7/Γ_6	NODE=M093R03 NODE=M093R03
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u> <u>COMMENT</u>
0.4±0.2	YIN	18 BELL $e^+e^- \rightarrow \text{hadrons}$

$\Upsilon(11020)$ REFERENCES

MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60090
YIN	18	PR D98 091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)	REFID=59468
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369