

# LIGHT UNFLAVORED MESONS ( $S = C = B = 0$ )

For  $l = 1$  ( $\pi, b, \rho, a$ ):  $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$ ;  
for  $l = 0$  ( $\eta, \eta', h, h', \omega, \phi, f, f'$ ):  $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

NODE=MXXX005

NODE=MXXX005

NODE=M014

## $f_0(500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

also known as  $\sigma$ ; was  $f_0(600)$   
See the related review(s):

Scalar Mesons below 1 GeV

### $f_0(500)$ T-MATRIX POLE $\sqrt{s}$

NODE=M014PP

NODE=M014PP

NODE=M014PP

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(400–550)–<math>i</math>(200–350) OUR ESTIMATE</b> (see Fig. 64.3 in the review)			
(410 ± 20)– $i$ (240 ± 15)	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
(512 ± 15)– $i$ (188 ± 12)	<sup>1</sup> ABLIKIM	17	BES3 $J/\psi \rightarrow \gamma 3\pi$
(440 ± 10)– $i$ (238 ± 10)	<sup>2</sup> ALBALADEJO	12	RVUE Compilation
(445 ± 25)– $i$ (278 <sup>+22</sup> <sub>-18</sub> )	<sup>3,4</sup> GARCIA-MAR..11	RVUE	Compilation
(457 <sup>+14</sup> <sub>-13</sub> )– $i$ (279 <sup>+11</sup> <sub>-7</sub> )	<sup>3,5</sup> GARCIA-MAR..11	RVUE	Compilation
(442 <sup>+5</sup> <sub>-8</sub> )– $i$ (274 <sup>+6</sup> <sub>-5</sub> )	<sup>6</sup> MOUSSALLAM11	RVUE	Compilation
(452 ± 13)– $i$ (259 ± 16)	<sup>7</sup> MENNESSIER	10	RVUE Compilation
(448 ± 43)– $i$ (266 ± 43)	<sup>8</sup> MENNESSIER	10	RVUE Compilation
(455 ± 6 <sup>+31</sup> <sub>-13</sub> )– $i$ (278 ± 6 <sup>+34</sup> <sub>-43</sub> )	<sup>9</sup> CAPRINI	08	RVUE Compilation
(463 ± 6 <sup>+31</sup> <sub>-17</sub> )– $i$ (259 ± 6 <sup>+33</sup> <sub>-34</sub> )	<sup>10</sup> CAPRINI	08	RVUE Compilation
(552 <sup>+84</sup> <sub>-106</sub> )– $i$ (232 <sup>+81</sup> <sub>-72</sub> )	<sup>11</sup> ABLIKIM	07A	BES2 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
(466 ± 18)– $i$ (223 ± 28)	<sup>12</sup> BONVICINI	07	CLEO $D^+ \rightarrow \pi^-\pi^+\pi^+$
(472 ± 30)– $i$ (271 ± 30)	<sup>13</sup> BUGG	07A	RVUE Compilation
(484 ± 17)– $i$ (255 ± 10)	GARCIA-MAR..07	RVUE	Compilation
(430)– $i$ (325)	<sup>14</sup> ANISOVICH	06	RVUE Compilation
(441 <sup>+16</sup> <sub>-8</sub> )– $i$ (272 <sup>+9</sup> <sub>-12.5</sub> )	<sup>15</sup> CAPRINI	06	RVUE $\pi\pi \rightarrow \pi\pi$
(470 ± 50)– $i$ (285 ± 25)	<sup>16</sup> ZHOU	05	RVUE
(541 ± 39)– $i$ (252 ± 42)	<sup>17</sup> ABLIKIM	04A	BES2 $J/\psi \rightarrow \omega\pi^+\pi^-$
(528 ± 32)– $i$ (207 ± 23)	<sup>18</sup> GALLEGOS	04	RVUE Compilation
(533 ± 25)– $i$ (249 ± 25)	<sup>19</sup> BUGG	03	RVUE
517 – $i$ 240	BLACK	01	RVUE $\pi\pi \rightarrow \pi\pi$
(470 ± 30)– $i$ (295 ± 20)	<sup>15</sup> COLANGELO	01	RVUE $\pi\pi \rightarrow \pi\pi$
(535 <sup>+48</sup> <sub>-36</sub> )– $i$ (155 <sup>+76</sup> <sub>-53</sub> )	<sup>20</sup> ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
610 ± 14 – $i$ (310 ± 13)	<sup>21</sup> SUROVTSEV	01	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
(540 <sup>+36</sup> <sub>-29</sub> )– $i$ (193 <sup>+32</sup> <sub>-40</sub> )	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
445 – $i$ 235	HANNAH	99	RVUE $\pi$ scalar form factor
(523 ± 12)– $i$ (259 ± 7)	KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
442 – $i$ 227	OLLER	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
469 – $i$ 203	OLLER	99B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
445 – $i$ 221	OLLER	99C	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
420 – $i$ 212	LOCHER	98	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
440 – $i$ 245	<sup>22</sup> DOBADO	97	RVUE Compilation
(602 ± 26)– $i$ (196 ± 27)	<sup>23</sup> ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
(537 ± 20)– $i$ (250 ± 17)	<sup>24</sup> KAMINSKI	97B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, 4\pi$
470 – $i$ 250	<sup>25,26</sup> TORNQVIST	96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
387 – $i$ 305	<sup>26,27</sup> JANSSEN	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
420 – $i$ 370	<sup>28</sup> ACHASOV	94	RVUE $\pi\pi \rightarrow \pi\pi$
(506 ± 10)– $i$ (247 ± 3)	KAMINSKI	94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
370 – $i$ 356	<sup>29</sup> ZOU	94B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
408 – $i$ 342	<sup>26,29</sup> ZOU	93	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

470 – $i208$	30 VANBEVEREN 86	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta,$
(750 ± 50) – $i(450 \pm 50)$	31 ESTABROOKS 79	RVUE	$\dots$ $\pi\pi \rightarrow \pi\pi, K\bar{K}$
(660 ± 100) – $i(320 \pm 70)$	PROTOPOP... 73	HBC	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
650 – $i370$	32 BASDEVANT 72	RVUE	$\pi\pi \rightarrow \pi\pi$

- 1 S-matrix pole; 8595 events.
- 2 Applying the chiral unitary approach at NLO to the  $K_{e4}$  data of BATLEY 10 and  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.
- 3 Uses the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.
- 4 Analytic continuation using Roy equations.
- 5 Analytic continuation using GKPY equations.
- 6 Using Roy equations.
- 7 Average of three variants of the analytic K-matrix model. Uses the  $K_{e4}$  data of BATLEY 08A and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73 and GRAYER 74.
- 8 Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.
- 9 From the  $K_{e4}$  data of BATLEY 08A and  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73.
- 10 From the  $K_{e4}$  data of BATLEY 08A and  $\pi N \rightarrow \pi\pi N$  data of PROTOPOPESCU 73, GRAYER 74, and ESTABROOKS 74.
- 11 From a mean of three different  $f_0(500)$  parametrizations. Uses 40k events.
- 12 From an isobar model using 2.6k events.
- 13 Reanalysis of ABLIKIM 04A, PISLAK 01, and HYAMS 73 data.
- 14 Using the N/D method.
- 15 From the solution of the Roy equation (ROY 71) for the isoscalar S-wave and using a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.
- 16 Reanalysis of the data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, ROSSELET 77, PISLAK 03, and AKHMETSHIN 04.
- 17 From a mean of six different analyses and  $f_0(500)$  parameterizations.
- 18 Using data on  $\psi(2S) \rightarrow J/\psi\pi\pi$  from BAI 00E and on  $\mathcal{T}(nS) \rightarrow \mathcal{T}(mS)\pi\pi$  from BUTLER 94B and ALEXANDER 98.
- 19 From a combined analysis of HYAMS 73, AUGUSTIN 89, AITALA 01B, and PISLAK 01.
- 20 A similar analysis (KOMADA 01) finds  $(580_{-30}^{+79}) - i(190_{-49}^{+107})$  MeV.
- 21 Coupled channel reanalysis of BATON 70, BENSINGER 71, BAILLON 72, HYAMS 73, HYAMS 75, ROSSELET 77, COHEN 80, and ETKIN 82B using the uniformizing variable.
- 22 Using the inverse amplitude method and data of ESTABROOKS 73, GRAYER 74, and PROTOPOPESCU 73.
- 23 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 24 Average and spread of 4 variants (“up” and “down”) of KAMINSKI 97B 3-channel model.
- 25 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.
- 26 Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.
- 27 Analysis of data from FALVARD 88.
- 28 Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.
- 29 Analysis of data from OCHS 73, GRAYER 74, and ROSSELET 77.
- 30 Coupled-channel analysis using data from PROTOPOPESCU 73, HYAMS 73, HYAMS 75, GRAYER 74, ESTABROOKS 74, ESTABROOKS 75, FROGGATT 77, CORN DEN 79, BISWAS 81.
- 31 Analysis of data from APEL 72C, GRAYER 74, CASON 76, PAWLICKI 77. Includes spread and errors of 4 solutions.
- 32 Analysis of data from BATON 70, BENSINGER 71, COLTON 71, BAILLON 72, PROTOPOPESCU 73, and WALKER 67.

NODE=M014PP;LINKAGE=I  
 NODE=M014PP;LINKAGE=AD  
 NODE=M014PP;LINKAGE=GM  
 NODE=M014PP;LINKAGE=GR  
 NODE=M014PP;LINKAGE=GC  
 NODE=M014PP;LINKAGE=MO  
 NODE=M014PP;LINKAGE=ME  
 NODE=M014PP;LINKAGE=MN  
 NODE=M014PP;LINKAGE=CA  
 NODE=M014PP;LINKAGE=AP  
 NODE=M014PP;LINKAGE=AL  
 NODE=M014PP;LINKAGE=BO  
 NODE=M014PP;LINKAGE=BU  
 NODE=M014PP;LINKAGE=AN  
 NODE=M014PP;LINKAGE=CL  
 NODE=M014PP;LINKAGE=ZH  
 NODE=M014PP;LINKAGE=AB  
 NODE=M014PP;LINKAGE=GA  
 NODE=M014PP;LINKAGE=PS  
 NODE=M014PP;LINKAGE=KI  
 NODE=M014PP;LINKAGE=SU  
 NODE=M014PP;LINKAGE=DO  
 NODE=M014PP;LINKAGE=AA  
 NODE=M014PP;LINKAGE=E  
 NODE=M014PP;LINKAGE=B  
 NODE=M014PP;LINKAGE=G  
 NODE=M014PP;LINKAGE=C  
 NODE=M014PP;LINKAGE=D  
 NODE=M014PP;LINKAGE=F  
 NODE=M014PP;LINKAGE=BV  
 NODE=M014PP;LINKAGE=A  
 NODE=M014PP;LINKAGE=J

### $f_0(500)$ BREIT-WIGNER MASS

NODE=M014M  
 NODE=M014M  
 → UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>400 to 800 OUR ESTIMATE</b>			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
513 ± 32	33 MURAMATSU 02	CLEO	$e^+e^- \approx 10$ GeV
478 $_{-23}^{+24}$ ± 17	AITALA 01B	E791	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
563 $_{-29}^{+58}$	34 ISHIDA 01		$\Upsilon(3S) \rightarrow \Upsilon \pi\pi$
555	35 ASNER 00	CLE2	$\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
540 ± 36	ISHIDA 00B		$p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$
750 ± 4	ALEKSEEV 99	SPEC	$1.78 \pi^- p_{\text{polar}} \rightarrow \pi^- \pi^+ n$
744 ± 5	ALEKSEEV 98	SPEC	$1.78 \pi^- p_{\text{polar}} \rightarrow \pi^- \pi^+ n$

759 ± 5	36 TROYAN	98	5.2 $np \rightarrow np\pi^+\pi^-$
780 ± 30	ALDE	97	GAM2 450 $pp \rightarrow pp\pi^0\pi^0$
585 ± 20	37 ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
761 ± 12	38 SVEC	96	RVUE 6-17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 860	39,40 TORNQVIST	96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1165 ± 50	41,42 ANISOVICH	95	RVUE $\pi^-p \rightarrow \pi^0\pi^0n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
414 ± 20	38 AUGUSTIN	89	DM2

33 Statistical uncertainty only.

34 A similar analysis (KOMADA 01) finds  $526^{+48}_{-37}$  MeV.

35 From the best fit of the Dalitz plot.

36  $6\sigma$  effect, no PWA.

37 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

38 Breit-Wigner fit to S-wave intensity measured in  $\pi N \rightarrow \pi^-\pi^+N$  on polarized targets. The fit does not include  $f_0(980)$ .

39 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

40 Also observed by ASNER 00 in  $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$  decays.

41 Uses  $\pi^0\pi^0$  data from ANISOVICH 94, AMSLER 94D, and ALDE 95B,  $\pi^+\pi^-$  data from OCHS 73, GRAYER 74 and ROSSELET 77, and  $\eta\eta$  data from ANISOVICH 94.

42 The pole is on Sheet III. Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.

NODE=M014M;LINKAGE=UT

NODE=M014M;LINKAGE=KI

NODE=M014M;LINKAGE=KK

NODE=M014M;LINKAGE=TN

NODE=M014M;LINKAGE=AA

NODE=M014M;LINKAGE=E

NODE=M014M;LINKAGE=B

NODE=M014M;LINKAGE=GG

NODE=M014M;LINKAGE=F

NODE=M014M;LINKAGE=G

### $f_0(500)$ BREIT-WIGNER WIDTH

NODE=M014W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>100 to 800 OUR ESTIMATE</b>			

NODE=M014W

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

335 ± 67	43 MURAMATSU	02	CLEO $e^+e^- \approx 10$ GeV
324 <sup>+</sup> <sub>40</sub> ± 21	AITALA	01B	E791 $D^+ \rightarrow \pi^-\pi^+\pi^+$
372 <sup>+</sup> <sub>95</sub> ± 229	44 ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
540	45 ASNER	00	CLE2 $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
372 ± 80	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
119 ± 13	ALEKSEEV	99	SPEC 1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
77 ± 22	ALEKSEEV	98	SPEC 1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
35 ± 12	46 TROYAN	98	5.2 $np \rightarrow np\pi^+\pi^-$
780 ± 60	ALDE	97	GAM2 450 $pp \rightarrow pp\pi^0\pi^0$
385 ± 70	47 ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
290 ± 54	48 SVEC	96	RVUE 6-17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 880	49,50 TORNQVIST	96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
460 ± 40	51,52 ANISOVICH	95	RVUE $\pi^-p \rightarrow \pi^0\pi^0n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
494 ± 58	48 AUGUSTIN	89	DM2

43 Statistical uncertainty only.

44 A similar analysis (KOMADA 01) finds  $301^{+145}_{-100}$  MeV.

45 From the best fit of the Dalitz plot.

46  $6\sigma$  effect, no PWA.

47 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

48 Breit-Wigner fit to S-wave intensity measured in  $\pi N \rightarrow \pi^-\pi^+N$  on polarized targets. The fit does not include  $f_0(980)$ .

NODE=M014W;LINKAGE=UT

NODE=M014W;LINKAGE=KI

NODE=M014W;LINKAGE=KK

NODE=M014W;LINKAGE=TN

NODE=M014W;LINKAGE=AA

NODE=M014W;LINKAGE=E

NODE=M014W;LINKAGE=B

49 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

50 Also observed by ASNER 00 in  $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$  decays.

NODE=M014W;LINKAGE=GG

NODE=M014W;LINKAGE=F

51 Uses  $\pi^0\pi^0$  data from ANISOVICH 94, AMSLER 94D, and ALDE 95B,  $\pi^+\pi^-$  data from OCHS 73, GRAYER 74 and ROSSELET 77, and  $\eta\eta$  data from ANISOVICH 94.

52 The pole is on Sheet III. Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.

NODE=M014W;LINKAGE=G

$f_0(500)$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $\gamma\gamma$	seen

NODE=M014215;NODE=M014

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

 $f_0(500)$  PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_2$
------------------------	-------------	-------------	------	---------	------------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2.05±0.21	53	DAI	14A	RVUE	Compilation	
1.7 ±0.4	54	HOFERICH... 11		RVUE	Compilation	
3.08±0.82	55	MENNESSIER 11		RVUE	Compilation	
2.08±0.2 $\begin{smallmatrix} +0.07 \\ -0.04 \end{smallmatrix}$	56	MOUSSALLAM11		RVUE	Compilation	
2.08	57	MAO	09	RVUE	Compilation	
1.2 ±0.4	58	BERNABEU 08		RVUE		
3.9 ±0.6	55	MENNESSIER 08		RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
1.8 ±0.4	59	OLLER 08		RVUE	Compilation	
1.68±0.15	59,60	OLLER 08A		RVUE	Compilation	
3.1 ±0.5	61,62	PENNINGTON 08		RVUE	Compilation	
2.4 ±0.4	62,63	PENNINGTON 08		RVUE	Compilation	
4.1 ±0.3	64	PENNINGTON 06		RVUE	$\gamma\gamma \rightarrow \pi^0\pi^0$	
3.8 ±1.5	65,66	BOGLIONE 99		RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
5.4 ±2.3	65	MORGAN 90		RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
10 ±6		COURAU 86		DM1	$e^+e^- \rightarrow \pi^+\pi^-e^+e^-$	
53 Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.						NODE=M014W2;LINKAGE=B
54 Using Roy-Steiner equations with $\pi\pi$ phase shifts from an update of COLANGELO 01 and from GARCIA-MARTIN 11A.						NODE=M014W2;LINKAGE=HO
55 Using an analytic K-matrix model.						NODE=M014W2;LINKAGE=ME
56 Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.						NODE=M014W2;LINKAGE=MO
57 Used dispersion theory. The value quoted used the $f_0(500)$ pole position of 457 - i276 MeV.						NODE=M014W2;LINKAGE=MA
58 Using $p, n$ polarizabilities from PDG 06 and fitting to $\pi\pi$ phase motion from GARCIA-MARTIN 07 and $\sigma$ -poles from GARCIA-MARTIN 07 and CAPRINI 06.						NODE=M014W2;LINKAGE=BE
59 Using twice-subtracted dispersion integrals.						NODE=M014W2;LINKAGE=OL
60 Supersedes OLLER 08.						NODE=M014W2;LINKAGE=LL
61 Solution A (preferred solution based on $\chi^2$ -analysis).						NODE=M014W2;LINKAGE=P1
62 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.						NODE=M014W2;LINKAGE=P3
63 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).						NODE=M014W2;LINKAGE=P2
64 Using unitarity and the $\sigma$ pole position from CAPRINI 06.						NODE=M014W2;LINKAGE=PE
65 This width could equally well be assigned to the $f_0(1370)$ . The authors analyse data from BOYER 90 and MARSISKE 90 and report strong correlation with $\gamma\gamma$ width of $f_2(1270)$ .						NODE=M014W2;LINKAGE=A
66 Supersedes MORGAN 90.						NODE=M014W2;LINKAGE=BL

NODE=M014220

NODE=M014W2  
NODE=M014W2

OCCUR=2

NODE=M014W2;LINKAGE=B

NODE=M014W2;LINKAGE=HO

NODE=M014W2;LINKAGE=ME

NODE=M014W2;LINKAGE=MO

NODE=M014W2;LINKAGE=MA

NODE=M014W2;LINKAGE=BE

NODE=M014W2;LINKAGE=OL

NODE=M014W2;LINKAGE=LL

NODE=M014W2;LINKAGE=P1

NODE=M014W2;LINKAGE=P3

NODE=M014W2;LINKAGE=P2

NODE=M014W2;LINKAGE=PE

NODE=M014W2;LINKAGE=A

NODE=M014W2;LINKAGE=BL

 $f_0(500)$  REFERENCES

SARANTSEV 21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM 17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DAI 14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)
ALBALADEJO 12	PR D86 034003	M. Albaladejo, J.A. Oller	(MURC)
GARCIA-MAR... 11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
GARCIA-MAR... 11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)
HOFERICH... 11	EPJ C71 1743	M. Hoferichter, D.R. Phillips, C. Schat	(BONN+)
MENNESSIER 11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang	
MOUSSALLAM 11	EPJ C71 1814	B. Moussallam	
BATLEY 10	PL B686 101	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BATLEY 10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
MENNESSIER 10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang	
MAO 09	PR D79 116008	Y. Mao <i>et al.</i>	
BATLEY 08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)
BERNABEU 08	PRL 100 241804	J. Bernabeu, J. Prades	(IFIC, GRAN)
CAPRINI 08	PR D77 114019	I. Caprini	
MENNESSIER 08	PL B665 205	G. Mennessier, S. Narison, W. Ochs	
OLLER 08	PL B659 201	J.A. Oller, L. Roca, C. Schat	(MURC, UBA)
OLLER 08A	EPJ A37 15	J.A. Oller, L. Roca	(MURC)
PENNINGTON 08	EPJ C56 1	M.R. Pennington <i>et al.</i>	
UEHARA 08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
ABLIKIM 07A	PL B645 19	M. Ablikim <i>et al.</i>	(BES Collab.)
BONVICINI 07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG 07A	JP G34 151	D.V. Bugg <i>et al.</i>	
GARCIA-MAR... 07	PR D76 074034	R. Garcia-Martin, J.R. Pelaez, F.J. Yndurain	

NODE=M014

REFID=61091

REFID=57650

REFID=55923

REFID=54277

REFID=16761

REFID=54121

REFID=53966

REFID=53637

REFID=53975

REFID=53285

REFID=53567

REFID=53657

REFID=52992

REFID=52487

REFID=52271

REFID=52273

REFID=52299

REFID=53976

REFID=53977

REFID=52303

REFID=52309

REFID=51614

REFID=51721

REFID=53252

REFID=51949

MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51652
ANISOVICH	06	IJMP A21 3615	V.V. Anisovich		REFID=51137
CAPRINI	06	PRL 96 132001	I. Caprini, G. Colangelo, H. Leutwyler	(BCIP+)	REFID=51076
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
PENNINGTON	06	PRL 97 011601	M.R. Pennington		REFID=51184
ZHOU	05	JHEP 0502 043	Z.Y. Zhou <i>et al.</i>		REFID=50823
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49740
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
BUETTIKER	04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam		REFID=56428;ERROR=1
GALLEGOS	04	PR D69 074033	A. Gallegos <i>et al.</i>		REFID=49769
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
PISLAK	03	PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=49344
Also		PR D81 119903E	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=53337
MURAMATSU	02	PRL 89 251802	H. Muramatsu <i>et al.</i>	(CLEO Collab.)	REFID=49081
Also		PRL 90 059901 (err.)	H. Muramatsu <i>et al.</i>	(CLEO Collab.)	REFID=49385
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
BLACK	01	PR D64 014031	D. Black <i>et al.</i>		REFID=48314
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler		REFID=49180
ISHIDA	01	PL B518 47	M. Ishida <i>et al.</i>		REFID=48354
KOMADA	01	PL B508 31	T. Komada <i>et al.</i>		REFID=48541
PISLAK	01	PRL 87 221801	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=48433
Also		PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=49344
Also		PRL 105 019901E	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=53338
SUROVTSEV	01	PR D63 054024	Y.S. Surovtsev, D. Krupa, M. Nagy		REFID=48310
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
BAI	00E	PR D62 032002	J. Bai <i>et al.</i>	(BES Collab.)	REFID=47955
ISHIDA	00B	PTP 104 203	M. Ishida <i>et al.</i>		REFID=48358
ALEKSEEV	99	NP B541 3	I.G. Alekseev <i>et al.</i>		REFID=46614
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
HANNAH	99	PR D60 017502	T. Hannah		REFID=46935
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
ALEKSEEV	98	PAN 61 174	I.G. Alekseev <i>et al.</i>		REFID=46328
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)	REFID=46372
TROYAN	98	JINRRC 5-91 33	Yu. Troyan <i>et al.</i>		REFID=46615
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
DOBADO	97	PR D56 3057	A. Dobado, J.R. Pelaez		REFID=53964
ISHIDA	97	PTP 98 1005	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45998
KAMINSKI	97B	PL B413 130	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, IPN)	REFID=45778
Also		PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45770
SVEC	96	PR D53 2343	M. Svec	(MCGI)	REFID=44509
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=44375
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)	REFID=44442
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
ACHASOV	94	PR D49 5779	N.N. Achasov, G.N. Shestakov	(NOVM)	REFID=44087
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43799
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)	REFID=44072
ZOU	93	PR D48 3948	B.S. Zou, D.V. Bugg	(LOQM)	REFID=43672
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
COURAU	86	NP B271 1	A. Courau <i>et al.</i>	(CLER, LALO)	REFID=44510
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i>	(NIJM, BIEL)	REFID=45769
CASON	83B	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BISWAS	81	PRL 47 1378	N.N. Biswas <i>et al.</i>	(NDAM, ANL)	REFID=21106
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP	REFID=20381
MUKHIN	80	JETPL 32 601	K.N. Mukhin <i>et al.</i>	(KIAE)	REFID=44528
		Translated from ZETFP 32 616.			
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
ESTABROOKS	79	PR D19 2678	P. Estabrooks	(CARL)	REFID=20375
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJ	REFID=20367
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
CASON	76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL) IJ	REFID=21064
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)	REFID=21062
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
ESTABROOKS	73	Tallahassee	P.G. Estabrooks <i>et al.</i>	(CERN, MPIM)	REFID=20345
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)	REFID=20349
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108
APEL	72C	PL 41B 542	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=21013
BAILLON	72	PL 38B 555	P.H. Baillon <i>et al.</i>	(SLAC)	REFID=20093
BASDEVANT	72	PL 41B 178	J.L. Basdevant, C.D. Froggatt, J.L. Petersen	(CERN)	REFID=20095
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)	REFID=44530
BENSINGER	71	PL 36B 134	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=21006
COLTON	71	PR D3 2028	E.P. Colton <i>et al.</i>	(LBL, FNAL, UCLA+)	REFID=44533
ROY	71	PL 36B 353	S.M. Roy		REFID=51107
BATON	70	PL 33B 528	J.P. Baton, G. Laurens, J. Reigner	(SACL)	REFID=20086
WALKER	67	RMP 39 695	W.D. Walker	(WISC)	REFID=20960

$\rho(770)$ 

$$I^G(J^{PC}) = 1^+(1^{--})$$

 **$\rho(770)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(761–765) – <math>i</math> (71–74) OUR ESTIMATE</b>			
$(763.7^{+1.7}_{-1.5}) - i(73.2^{+1.0}_{-1.1})$	<sup>1</sup> GARCIA-MAR..11	RVUE	Compilation
$(754 \pm 18) - i(74 \pm 10)$	<sup>2</sup> PELAEZ	04A	RVUE $\pi\pi \rightarrow \pi\pi$
$(762.4 \pm 1.8) - i(72.6 \pm 1.4)$	COLANGELO	01	RVUE $\pi\pi \rightarrow \pi\pi$

<sup>1</sup> Reanalysis of the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOESCU 73 using GKPY equations.

<sup>2</sup> Reanalysis of data from PROTOPOESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.

 **$\rho(770)$  MASS**We no longer list  $S$ -wave Breit-Wigner fits, or data with high combinatorial background.**NEUTRAL ONLY,  $e^+e^-$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>775.26 ± 0.23 OUR AVERAGE</b>				
775.3 ± 0.5 ± 0.6		<sup>1</sup> ACHASOV	21	SND $e^+e^- \rightarrow \pi^+\pi^-$
775.02 ± 0.35		<sup>2</sup> LEES	12G	BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$
775.97 ± 0.46 ± 0.70	900k	<sup>3</sup> AKHMETSHIN	07	$e^+e^- \rightarrow \pi^+\pi^-$
774.6 ± 0.4 ± 0.5	800k	<sup>4,5</sup> ACHASOV	06	SND $e^+e^- \rightarrow \pi^+\pi^-$
775.65 ± 0.64 ± 0.50	114k	<sup>6,7</sup> AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow \pi^+\pi^-$
775.9 ± 0.5 ± 0.5	1.98M	<sup>8</sup> ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.8 ± 0.9 ± 2.0	500k	<sup>8</sup> ACHASOV	02	SND $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ± 1.1		<sup>9</sup> BARKOV	85	OLYA $e^+e^- \rightarrow \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
763.49 ± 0.53		<sup>10</sup> BARTOS	17	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
758.23 ± 0.46		<sup>11</sup> BARTOS	17A	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
775.8 ± 0.5 ± 0.3	1.98M	<sup>12</sup> ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ± 0.6 ± 0.5	1.98M	<sup>13</sup> ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.0 ± 0.6 ± 1.1	500k	<sup>14</sup> ACHASOV	02	SND $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ± 0.7 ± 5.3		<sup>15</sup> BENAYOUN	98	RVUE $e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
770.5 ± 1.9 ± 5.1		<sup>16</sup> GARDNER	98	RVUE $0.28-0.92 e^+e^- \rightarrow \pi^+\pi^-$
764.1 ± 0.7		<sup>17</sup> O'CONNELL	97	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
757.5 ± 1.5		<sup>18</sup> BERNICHA	94	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
768 ± 1		<sup>19</sup> GESHKEN...	89	RVUE $e^+e^- \rightarrow \pi^+\pi^-$

<sup>1</sup> From a fit of the cross section in the energy range  $0.525 < \sqrt{s} < 0.883$  GeV parameterized by the sum of the Breit-Wigner amplitudes for the  $\rho(770)$ ,  $\omega$  and  $\rho(1450)$  resonances.

<sup>2</sup> Using the GOUNARIS 68 parametrization with the complex phase of the  $\rho$ - $\omega$  interference and leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

<sup>3</sup> A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

<sup>4</sup> Supersedes ACHASOV 05A.

<sup>5</sup> A fit of the SND data from 400 to 1000 MeV using parameters of the  $\rho(1450)$  and  $\rho(1700)$  from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

<sup>6</sup> Using the GOUNARIS 68 parametrization with the complex phase of the  $\rho$ - $\omega$  interference.

<sup>7</sup> Update of AKHMETSHIN 02.

<sup>8</sup> Assuming  $m_{\rho^+} = m_{\rho^-}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-}$ .

<sup>9</sup> From the GOUNARIS 68 parametrization of the pion form factor.

<sup>10</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

<sup>11</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

<sup>12</sup> Assuming  $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$ .

<sup>13</sup> Without limitations on masses and widths.

NODE=M009

NODE=M009PP

NODE=M009PP

NODE=M009PP

→ UNCHECKED ←

OCCUR=2

OCCUR=2

NODE=M009PP;LINKAGE=C

NODE=M009PP;LINKAGE=B

NODE=M009205

NODE=M009205

NODE=M009M0

NODE=M009M0

OCCUR=2

OCCUR=3

OCCUR=3

NODE=M009M0;LINKAGE=D

NODE=M009M0;LINKAGE=LE

NODE=M009M0;LINKAGE=AK

NODE=M009M0;LINKAGE=AC

NODE=M009M0;LINKAGE=SN

NODE=M009M0;LINKAGE=GS

NODE=M009M0;LINKAGE=PT

NODE=M009M0;LINKAGE=CH

NODE=M009M0;LINKAGE=K

NODE=M009M0;LINKAGE=A

NODE=M009M0;LINKAGE=B

NODE=M009M0;LINKAGE=DF

NODE=M009M0;LINKAGE=WO

<sup>14</sup> Assuming  $m_{\rho^0} = m_{\rho^\pm}$ ,  $g_{\rho^0\pi\pi} = g_{\rho^\pm\pi\pi}$ .

<sup>15</sup> Using the data of BARKOV 85 in the hidden local symmetry model.

<sup>16</sup> From the fit to  $e^+e^- \rightarrow \pi^+\pi^-$  data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.

<sup>17</sup> A fit of BARKOV 85 data assuming the direct  $\omega\pi\pi$  coupling.

<sup>18</sup> Applying the S-matrix formalism to the BARKOV 85 data.

<sup>19</sup> Includes BARKOV 85 data. Model-dependent width definition.

NODE=M009M0;LINKAGE=HC  
 NODE=M009M0;LINKAGE=K2  
 NODE=M009M0;LINKAGE=G8

NODE=M009M0;LINKAGE=AB  
 NODE=M009M0;LINKAGE=AA  
 NODE=M009M0;LINKAGE=F

### CHARGED ONLY, $\tau$ DECAYS and $e^+e^-$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>775.11±0.34 OUR AVERAGE</b>					
774.6 ±0.2 ±0.5	5.4M	<sup>1,2</sup> FUJIKAWA	08	BELL ±	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
775.5 ±0.7		<sup>2,3</sup> SCHAEEL	05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
775.5 ±0.5 ±0.4	1.98M	<sup>4</sup> ALOISIO	03	KLOE	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±1.1 ±0.5	87k	<sup>5,6</sup> ANDERSON	00A	CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
761.60±0.95		<sup>7</sup> BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
774.8 ±0.6 ±0.4	1.98M	<sup>8</sup> ALOISIO	03	KLOE -	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
776.3 ±0.6 ±0.7	1.98M	<sup>8</sup> ALOISIO	03	KLOE +	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
773.9 ±2.0 <sup>+0.3</sup> <sub>-1.0</sub>		<sup>9</sup> SANZ-CILLERO	03	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
774.5 ±0.7 ±1.5	500k	<sup>4</sup> ACHASOV	02	SND ±	$1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ±0.5		<sup>10</sup> PICH	01	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

NODE=M009M5  
 NODE=M009M5

OCCUR=2

OCCUR=3

OCCUR=2

<sup>1</sup>  $|F_\pi(0)|^2$  fixed to 1.

<sup>2</sup> From the GOUNARIS 68 parametrization of the pion form factor.

<sup>3</sup> The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.

<sup>4</sup> Assuming  $m_{\rho^+} = m_{\rho^-}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-}$ .

<sup>5</sup>  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV respectively.

<sup>6</sup> From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.

<sup>7</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.

<sup>8</sup> Without limitations on masses and widths.

<sup>9</sup> Using the data of BARATE 97M and the effective chiral Lagrangian.

<sup>10</sup> From a fit of the model-independent parameterization of the pion form factor to the data of BARATE 97M.

NODE=M009M5;LINKAGE=FU  
 NODE=M009M5;LINKAGE=GO  
 NODE=M009M5;LINKAGE=SC

NODE=M009M5;LINKAGE=CH  
 NODE=M009M5;LINKAGE=A6  
 NODE=M009M5;LINKAGE=K1

NODE=M009M5;LINKAGE=A

NODE=M009M5;LINKAGE=WO  
 NODE=M009M5;LINKAGE=Z  
 NODE=M009M5;LINKAGE=PC

### MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>763.0±0.3±1.2</b>	600k	<sup>1</sup> ABELE	99E	CBAR 0±	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> Assuming the equality of  $\rho^+$  and  $\rho^-$  masses and widths.

NODE=M009M7  
 NODE=M009M7

NODE=M009M7;LINKAGE=LB

### CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>766.5±1.1 OUR AVERAGE</b>					
763.7±3.2		ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
768 ±9		AGUILAR...	91	EHS	400 $p p$
767 ±3	2935	<sup>1</sup> CAPRARO	87	SPEC -	200 $\pi^- \text{Cu} \rightarrow \pi^- \pi^0 \text{Cu}$
761 ±5	967	<sup>1</sup> CAPRARO	87	SPEC -	200 $\pi^- \text{Pb} \rightarrow \pi^- \pi^0 \text{Pb}$
771 ±4		HUSTON	86	SPEC +	202 $\pi^+ \text{A} \rightarrow \pi^+ \pi^0 \text{A}$
766 ±7	6500	<sup>2</sup> BYERLY	73	OSPK -	5 $\pi^- p$
766.8±1.5	9650	<sup>3</sup> PISUT	68	RVUE -	1.7-3.2 $\pi^- p$ , $t < 10$
767 ±6	900	<sup>1</sup> EISNER	67	HBC -	4.2 $\pi^- p$ , $t < 10$

NODE=M009M2  
 NODE=M009M2

OCCUR=2

<sup>1</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>2</sup> Phase shift analysis. Systematic errors added corresponding to spread of different fits.

<sup>3</sup> From fit of 3-parameter relativistic  $P$ -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

NODE=M009M2;LINKAGE=Z  
 NODE=M009M2;LINKAGE=X  
 NODE=M009M2;LINKAGE=A

**NEUTRAL ONLY, PHOTOPRODUCED**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>769.2± 0.9 OUR AVERAGE</b>				
770.8± 1.3 <sup>+2.3</sup> <sub>-2.4</sub>	900k	ANDREEV 20	H1	$ep \rightarrow e\pi^+\pi^-p$
771 ± 2 <sup>+2</sup> <sub>-1</sub>	63.5k	<sup>1</sup> ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^-p$
770 ± 2 ± 1	79k	<sup>2</sup> BREITWEG 98B	ZEUS	50–100 $\gamma p$
767.6± 2.7		BARTALUCCI 78	CNTR	$\gamma p \rightarrow e^+e^-p$
775 ± 5		GLADDING 73	CNTR	2.9–4.7 $\gamma p$
767 ± 4	1930	BALLAM 72	HBC	2.8 $\gamma p$
770 ± 4	2430	BALLAM 72	HBC	4.7 $\gamma p$
765 ± 10		ALVENSLEB... 70	CNTR	$\gamma A, t < 0.01$
767.7± 1.9	140k	BIGGS 70	CNTR	$< 4.1 \gamma C \rightarrow \pi^+\pi^-C$
765 ± 5	4000	ASBURY 67B	CNTR	$\gamma + Pb$

• • • We do not use the following data for averages, fits, limits, etc. • • •

771 ± 2	79k	<sup>3</sup> BREITWEG 98B	ZEUS	50–100 $\gamma p$
---------	-----	---------------------------	------	-------------------

<sup>1</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho$ - $\omega$  interference.  
<sup>2</sup> From the parametrization according to SOEDING 66.  
<sup>3</sup> From the parametrization according to ROSS 66.

NODE=M009M0P  
 NODE=M009M0P

OCCUR=2

OCCUR=2

NODE=M009M0P;LINKAGE=AB  
 NODE=M009M0P;LINKAGE=B5  
 NODE=M009M0P;LINKAGE=B6

**NEUTRAL ONLY, OTHER REACTIONS**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>769.0 ± 0.9 OUR AVERAGE</b>				
Error includes scale factor of 1.4. See the ideogram below.				
765 ± 6		BERTIN 97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
773 ± 1.6		WEIDENAUER 93	ASTE	$\bar{p}p \rightarrow \pi^+\pi^-\omega$
762.6 ± 2.6		AGUILAR-... 91	EHS	400 $pp$
770 ± 2		<sup>1</sup> HEYN 81	RVUE	Pion form factor
768 ± 4		<sup>2,3</sup> BOHACIK 80	RVUE	
769 ± 3		<sup>4</sup> WICKLUND 78	ASPK	3,4,6 $\pi^\pm N$
768 ± 1	76k	DEUTSCH... 76	HBC	16 $\pi^+p$
767 ± 4	4100	ENGLER 74	DBC	6 $\pi^+n \rightarrow \pi^+\pi^-p$
775 ± 4	32k	<sup>2</sup> PROTOPOP... 73	HBC	7.1 $\pi^+p, t < 0.4$
764 ± 3	6.8k	<sup>5</sup> RATCLIFF 72	ASPK	15 $\pi^-p, t < 0.3$
774 ± 3	1.7k	REYNOLDS 69	HBC	2.26 $\pi^-p$
769.2 ± 1.5	13.3k	<sup>6</sup> PISUT 68	RVUE	1.7–3.2 $\pi^-p, t < 10$

• • • We do not use the following data for averages, fits, limits, etc. • • •

774.34±0.18±0.35	970k	<sup>7</sup> ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
772.93±0.18±0.34	970k	<sup>8</sup> ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
773.5 ± 2.5		<sup>9</sup> COLANGELO 01	RVUE	$\pi\pi \rightarrow \pi\pi$
762.3 ± 0.5 ± 1.2	600k	<sup>10</sup> ABELE 99E	CBAR	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
777 ± 2	4.9k	<sup>11</sup> ADAMS 97	E665	470 $\mu p \rightarrow \mu XB$
770 ± 2		<sup>12</sup> BOGOLYUB... 97	MIRA	32 $\bar{p}p \rightarrow \pi^+\pi^-X$
768 ± 8		<sup>12</sup> BOGOLYUB... 97	MIRA	32 $pp \rightarrow \pi^+\pi^-X$
761.1 ± 2.9		DUBNICKA 89	RVUE	$\pi$ form factor
777.4 ± 2.0		<sup>13</sup> CHABAUD 83	ASPK	17 $\pi^-p$ polarized
769.5 ± 0.7		<sup>2,3</sup> LANG 79	RVUE	
770 ± 9		<sup>3</sup> ESTABROOKS 74	RVUE	17 $\pi^-p \rightarrow \pi^+\pi^-n$
773.5 ± 1.7	11.2k	<sup>14</sup> JACOBS 72	HBC	2.8 $\pi^-p$
775 ± 3	2.2k	<sup>15</sup> HYAMS 68	OSPK	11.2 $\pi^-p$

<sup>1</sup> HEYN 81 includes all spacelike and timelike  $F_\pi$  values until 1978.

<sup>2</sup> From pole extrapolation.

<sup>3</sup> From phase shift analysis of GRAYER 74 data.

<sup>4</sup> Phase shift analysis. Systematic errors added corresponding to spread of different fits.

<sup>5</sup> Published values contain misprints. Corrected by private communication RATCLIFF 74.

<sup>6</sup> Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDBERGER 64, ABOLINS 63.

<sup>7</sup> From a fit to  $\pi^+\pi^-$  mass using  $\rho(770)$  (parametrized with the Gounaris-Sakurai approach),  $\omega(782)$ , and box anomaly components.

<sup>8</sup> From a fit to  $\pi^+\pi^-$  mass using  $\rho(770)$  (parametrized with the Gounaris-Sakurai approach),  $\omega(782)$ , and  $\rho(1450)$  components.

<sup>9</sup> Breit-Wigner mass from a phase-shift analysis of HYAMS 73 and PROTOPOESCU 73 data.

<sup>10</sup> Using relativistic Breit-Wigner and taking into account  $\rho$ - $\omega$  interference.

<sup>11</sup> Systematic errors not evaluated.

<sup>12</sup> Systematic effects not studied.

<sup>13</sup> From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P-wave intensity. CHABAUD 83 includes data of GRAYER 74.

NODE=M009M0R  
 NODE=M009M0R

OCCUR=2

OCCUR=2

NODE=M009M0R;LINKAGE=B0  
 NODE=M009M0R;LINKAGE=C0  
 NODE=M009M0R;LINKAGE=H  
 NODE=M009M0R;LINKAGE=X  
 NODE=M009M0R;LINKAGE=03  
 NODE=M009M0R;LINKAGE=R

NODE=M009M0R;LINKAGE=B

NODE=M009M0R;LINKAGE=C

NODE=M009M0R;LINKAGE=CL


NODE=M009M0R;LINKAGE=BL  
 NODE=M009M0R;LINKAGE=A1  
 NODE=M009M0R;LINKAGE=QQ  
 NODE=M009M0R;LINKAGE=G



<sup>14</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>15</sup> Of HYAMS 68 six parametrizations, this is theoretically soundest. MR

NODE=M009M0R;LINKAGE=Z  
 NODE=M009M0R;LINKAGE=02



$m_{\rho(770)^0} - m_{\rho(770)^\pm}$

NODE=M009D

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
<b>-0.7 ± 0.8</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.5. See the ideogram below.			
-2.4 ± 0.8	1	SCHAEL 05C	ALEP		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
0.4 ± 0.7 ± 0.6	1.98M	2 ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.3 ± 1.1 ± 2.0	500k	2 ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.6 ± 0.6 ± 1.7	600k	ABELE	99E	CBAR ±0	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
-4 ± 4	3000	3 REYNOLDS	69	HBC -0	$2.26 \pi^- \rho$
-5 ± 5	3600	3 FOSTER	68	HBC ±0	$0.0 \bar{p} p$
2.4 ± 2.1	22950	4 PISUT	68	RVUE	$\pi N \rightarrow \rho N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
-3.37 ± 1.06		5 BARTOS	17A	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$ , $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

NODE=M009D

<sup>1</sup> From the combined fit of the  $\tau^-$  data from ANDERSON 00A and SCHAEL 05c and  $e^+ e^-$  data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

NODE=M009D;LINKAGE=SC

<sup>2</sup> Assuming  $m_{\rho^+} = m_{\rho^-}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-}$ .

NODE=M009D;LINKAGE=CH

<sup>3</sup> From quoted masses of charged and neutral modes.


NODE=M009D;LINKAGE=A

<sup>4</sup> Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDHABER 64, ABOLINS 63.

NODE=M009D;LINKAGE=R

<sup>5</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M009D;LINKAGE=B



**$m_{\rho(770)^+} - m_{\rho(770)^-}$**

NODE=M009D1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1.5 \pm 0.8 \pm 0.7$	1.98M	<sup>1</sup> ALOISIO 03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
<sup>1</sup> Without limitations on masses and widths.				

NODE=M009D1

NODE=M009D;LINKAGE=WO

**$\rho(770)$  RANGE PARAMETER**

NODE=M009R

The range parameter  $R$  enters an energy-dependent correction to the width, of the form  $(1 + q_r^2 R^2) / (1 + q^2 R^2)$ , where  $q$  is the momentum of one of the pions in the  $\pi\pi$  rest system. At resonance,  $q = q_r$ .

NODE=M009R

VALUE ( $\text{GeV}^{-1}$ )	DOCUMENT ID	TECN	CHG	COMMENT
$5.3^{+0.9}_{-0.7}$	<sup>1</sup> CHABAUD 83	ASPK	0	$17 \pi^- p$ polarized
<sup>1</sup> The old PISUT 68 value, properly corrected, was $3.2 \pm 0.6$ .				

NODE=M009R

NODE=M009R;LINKAGE=01

**$\rho(770)$  WIDTH**

NODE=M009220

We no longer list  $S$ -wave Breit-Wigner fits, or data with high combinatorial background.

NODE=M009220

**NEUTRAL ONLY,  $e^+ e^-$**


NODE=M009W0  
 NODE=M009W0

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>147.4 \pm 0.8</math></b>		<b>OUR AVERAGE</b> Error includes scale factor of 2.0. See the ideogram below.		
$145.6 \pm 0.6 \pm 0.8$		<sup>1</sup> ACHASOV 21	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$149.59 \pm 0.67$		<sup>2</sup> LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
$145.98 \pm 0.75 \pm 0.50$	900k	<sup>3</sup> AKHMETSHIN 07		$e^+ e^- \rightarrow \pi^+ \pi^-$
$146.1 \pm 0.8 \pm 1.5$	800k	<sup>4,5</sup> ACHASOV 06	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$143.85 \pm 1.33 \pm 0.80$	114k	<sup>6,7</sup> AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^-$
$147.3 \pm 1.5 \pm 0.7$	1.98M	<sup>8</sup> ALOISIO 03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$151.1 \pm 2.6 \pm 3.0$	500k	<sup>8</sup> ACHASOV 02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$150.5 \pm 3.0$		<sup>9</sup> BARKOV 85	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

144.06±0.85		10	BARTOS	17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
144.56±0.80		11	BARTOS	17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
143.9 ±1.3 ±1.1	1.98M	12	ALOISIO	03	KLOE	$1.02 \frac{e^+e^- \rightarrow \pi^+\pi^-\pi^0}{\pi^+\pi^-\pi^0}$	OCCUR=2
147.4 ±1.5 ±0.7	1.98M	13	ALOISIO	03	KLOE	$1.02 \frac{e^+e^- \rightarrow \pi^+\pi^-\pi^0}{\pi^+\pi^-\pi^0}$	OCCUR=3
149.8 ±2.2 ±2.0	500k	14	ACHASOV	02	SND	$1.02 \frac{e^+e^- \rightarrow \pi^+\pi^-\pi^0}{\pi^+\pi^-\pi^0}$	OCCUR=3
147.9 ±1.5 ±7.5		15	BENAYOUN	98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$	
153.5 ±1.3 ±4.6		16	GARDNER	98	RVUE	$0.28-0.92 \frac{e^+e^- \rightarrow \pi^+\pi^-}{\pi^+\pi^-}$	
145.0 ±1.7		17	O'CONNELL	97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
142.5 ±3.5		18	BERNICHIA	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
138 ±1		19	GESHKEN...	89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	

- 1 From a fit of the cross section in the energy range  $0.525 < \sqrt{s} < 0.883$  GeV parameterized by the sum of the Breit-Wigner amplitudes for the  $\rho(770)$ ,  $\omega$  and  $\rho(1450)$  resonances. NODE=M009W0;LINKAGE=D
- 2 Using the GOUNARIS 68 parametrization with the complex phase of the  $\rho$ - $\omega$  interference and leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit. NODE=M009W0;LINKAGE=LE
- 3 A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05. NODE=M009W;LINKAGE=AK
- 4 Supersedes ACHASOV 05A. NODE=M009W0;LINKAGE=AC  
NODE=M009W0;LINKAGE=SN
- 5 A fit of the SND data from 400 to 1000 MeV using parameters of the  $\rho(1450)$  and  $\rho(1700)$  from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A. NODE=M009W5;LINKAGE=GS  
NODE=M009W5;LINKAGE=P2
- 6 Using the GOUNARIS 68 parametrization with the complex phase of the  $\rho$ - $\omega$  interference. NODE=M009W;LINKAGE=CH  
NODE=M009W;LINKAGE=K
- 7 From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02. NODE=M009W0;LINKAGE=A  
NODE=M009W0;LINKAGE=B
- 8 Assuming  $m_{\rho^+} = m_{\rho^-}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-}$ . NODE=M009W;LINKAGE=DF  
NODE=M009W;LINKAGE=WO
- 9 From the GOUNARIS 68 parametrization of the pion form factor. NODE=M009W;LINKAGE=HC  
NODE=M009W;LINKAGE=K2
- 10 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C. NODE=M009W;LINKAGE=G8
- 11 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A. NODE=M009W;LINKAGE=AB  
NODE=M009W;LINKAGE=AA  
NODE=M009W;LINKAGE=F
- 12 Assuming  $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$ .
- 13 Without limitations on masses and widths.
- 14 Assuming  $m_{\rho^0} = m_{\rho^\pm}$ ,  $g_{\rho^0\pi\pi} = g_{\rho^\pm\pi\pi}$ .
- 15 Using the data of BARKOV 85 in the hidden local symmetry model.
- 16 From the fit to  $e^+e^- \rightarrow \pi^+\pi^-$  data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.
- 17 A fit of BARKOV 85 data assuming the direct  $\omega\pi\pi$  coupling.
- 18 Applying the S-matrix formalism to the BARKOV 85 data.
- 19 Includes BARKOV 85 data. Model-dependent width definition.



**CHARGED ONLY,  $\tau$  DECAYS and  $e^+e^-$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>149.1 <math>\pm 0.8</math> OUR FIT</b>					
<b>149.1 <math>\pm 0.8</math> OUR AVERAGE</b>					
148.1 $\pm 0.4$	$\pm 1.7$	5.4M	1,2 FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
149.0 $\pm 1.2$			2,3 SCHAEL	05c ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
149.9 $\pm 2.3$	$\pm 2.0$	500k	4 ACHASOV	02 SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.4 $\pm 1.4$	$\pm 1.4$	87k	5,6 ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
139.90 $\pm 0.46$			7 BARTOS	17A RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
143.7 $\pm 1.3$	$\pm 1.2$	1.98M	4 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
142.9 $\pm 1.3$	$\pm 1.4$	1.98M	8 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
144.7 $\pm 1.4$	$\pm 1.2$	1.98M	8 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.2 $\pm 2.0$	$+0.7$ $-1.6$		9 SANZ-CILLERO	03 RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
150.9 $\pm 2.2$	$\pm 2.0$	500k	10 ACHASOV	02 SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M009W5  
 NODE=M009W5

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

<sup>1</sup>  $|F_\pi(0)|^2$  fixed to 1.

<sup>2</sup> From the GOUNARIS 68 parametrization of the pion form factor.

<sup>3</sup> The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.

<sup>4</sup> Assuming  $m_{\rho^+} = m_{\rho^-}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-}$ .

<sup>5</sup>  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV respectively.

<sup>6</sup> From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.

<sup>7</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.

<sup>8</sup> Without limitations on masses and widths.

<sup>9</sup> Using the data of BARATE 97M and the effective chiral Lagrangian.

<sup>10</sup> Assuming  $m_{\rho^0} = m_{\rho^\pm}$ ,  $g_{\rho^0 \pi\pi} = g_{\rho^\pm \pi\pi}$ .

NODE=M009W5;LINKAGE=FU  
 NODE=M009W5;LINKAGE=GO  
 NODE=M009W5;LINKAGE=SC

NODE=M009W5;LINKAGE=CH  
 NODE=M009W5;LINKAGE=A6  
 NODE=M009W5;LINKAGE=K1

NODE=M009W5;LINKAGE=A

NODE=M009W5;LINKAGE=W0  
 NODE=M009W5;LINKAGE=Z

NODE=M009W5;LINKAGE=HC

**MIXED CHARGES, OTHER REACTIONS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>149.5 <math>\pm 1.3</math></b>	600k	1 ABELE	99E CBAR	0 $\pm$	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> Assuming the equality of  $\rho^+$  and  $\rho^-$  masses and widths.

NODE=M009W7  
 NODE=M009W7

NODE=M009W;LINKAGE=LB

**CHARGED ONLY, HADROPRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>150.2 <math>\pm 2.4</math> OUR FIT</b>					
<b>150.2 <math>\pm 2.4</math> OUR AVERAGE</b>					
152.8 $\pm 4.3$			ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
155 $\pm 11$	2.9k	1 CAPRARO	87 SPEC	-	200 $\pi^- \text{Cu} \rightarrow \pi^- \pi^0 \text{Cu}$
154 $\pm 20$	967	1 CAPRARO	87 SPEC	-	200 $\pi^- \text{Pb} \rightarrow \pi^- \pi^0 \text{Pb}$
150 $\pm 5$		HUSTON	86 SPEC	+	202 $\pi^+ \text{A} \rightarrow \pi^+ \pi^0 \text{A}$
146 $\pm 12$	6.5k	2 BYERLY	73 OSPK	-	5 $\pi^- p$
148.2 $\pm 4.1$	9.6k	3 PISUT	68 RVUE	-	1.7-3.2 $\pi^- p$ , $t < 10$
146 $\pm 13$	900	EISNER	67 HBC	-	4.2 $\pi^- p$ , $t < 10$

NODE=M009W2  
 NODE=M009W2

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

137.0  $\pm 0.4$  <sup>4</sup> ABLIKIM 17 BES3  $J/\psi \rightarrow \gamma 3\pi$

<sup>1</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>2</sup> Phase shift analysis. Systematic errors added corresponding to spread of different fits.

<sup>3</sup> From fit of 3-parameter relativistic  $P$ -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

<sup>4</sup> S-matrix pole at a fixed  $\rho$  meson mass of 775.49 MeV.

NODE=M009W2;LINKAGE=Z  
 NODE=M009W2;LINKAGE=X  
 NODE=M009W2;LINKAGE=A1

NODE=M009W2;LINKAGE=A

**NEUTRAL ONLY, PHOTOPRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>151.5 <math>^{+1.9}_{-2.1}</math> OUR AVERAGE</b>				
151.3 $\pm 2.2$	$^{+1.6}_{-2.8}$	900k	ANDREEV	20 H1 $ep \rightarrow e\pi^+\pi^-p$
155 $\pm 5$	$\pm 2$	63.5k	1 ABRAMOWICZ	12 ZEUS $ep \rightarrow e\pi^+\pi^-p$
146 $\pm 3$	$\pm 13$	79k	2 BREITWEG	98B ZEUS 50-100 $\gamma p$
150.9 $\pm 3.0$			BARTALUCCI	78 CNTR $\gamma p \rightarrow e^+e^-p$

NODE=M009W0P  
 NODE=M009W0P

• • • We do not use the following data for averages, fits, limits, etc. • • •

138 ± 3	79k	<sup>3</sup> BREITWEG	98B	ZEUS	50–100 $\gamma p$	OCCUR=2
147 ± 11		GLADDING	73	CNTR	2.9–4.7 $\gamma p$	
155 ± 12	2430	BALLAM	72	HBC	4.7 $\gamma p$	
145 ± 13	1930	BALLAM	72	HBC	2.8 $\gamma p$	OCCUR=2
140 ± 5		ALVENSLEB...	70	CNTR	$\gamma A$ , $t < 0.01$	
146.1 ± 2.9	140k	BIGGS	70	CNTR	$< 4.1 \gamma C \rightarrow \pi^+ \pi^- C$	
160 ± 10		LANZEROTTI	68	CNTR	$\gamma p$	
130 ± 5	4000	ASBURY	67B	CNTR	$\gamma + Pb$	

<sup>1</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho$ - $\omega$  interference.

<sup>2</sup> From the parametrization according to SOEDING 66.

<sup>3</sup> From the parametrization according to ROSS 66.

NODE=M009W0P;LINKAGE=AB

NODE=M009W;LINKAGE=B5

NODE=M009W;LINKAGE=B6

## NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>150.9 ± 1.7 OUR AVERAGE</b>		Error includes scale factor of 1.1.		
122 ± 20		BERTIN	97C	OBLX 0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
145.7 ± 5.3		WEIDENAUER	93	ASTE $\bar{p} p \rightarrow \pi^+ \pi^- \omega$
144.9 ± 3.7		DUBNICKA	89	RVUE $\pi$ form factor
148 ± 6		<sup>1,2</sup> BOHACIK	80	RVUE
152 ± 9		<sup>3</sup> WICKLUND	78	ASPK 3,4,6 $\pi^\pm p N$
154 ± 2	76k	DEUTSCH...	76	HBC 16 $\pi^+ p$
157 ± 8	6.8k	<sup>4</sup> RATCLIFF	72	ASPK 15 $\pi^- p$ , $t < 0.3$
143 ± 8	1.7k	REYNOLDS	69	HBC 2.26 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

150.85 ± 0.55 ± 0.67	970k	<sup>5</sup> ABLIKIM	18C	BES3 $\eta'(958) \rightarrow \gamma \pi^+ \pi^-$	
150.18 ± 0.55 ± 0.65	970k	<sup>6</sup> ABLIKIM	18C	BES3 $\eta'(958) \rightarrow \gamma \pi^+ \pi^-$	OCCUR=2
147.0 ± 2.5	600k	<sup>7</sup> ABELE	99E	CBAR 0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$	
146 ± 3	4.9k	<sup>8</sup> ADAMS	97	E665 470 $\mu p \rightarrow \mu X B$	
160.0 + 4.1 - 4.0		<sup>9</sup> CHABAUD	83	ASPK 17 $\pi^- p$ polarized	
155 ± 1		<sup>10</sup> HEYN	81	RVUE $\pi$ form factor	
148.0 ± 1.3		<sup>1,2</sup> LANG	79	RVUE	
146 ± 14	4.1k	ENGLER	74	DBC 6 $\pi^+ n \rightarrow \pi^+ \pi^- p$	
143 ± 13		<sup>2</sup> ESTABROOKS	74	RVUE 17 $\pi^- p \rightarrow \pi^+ \pi^- n$	
160 ± 10	32k	<sup>1</sup> PROTOPOP...	73	HBC 7.1 $\pi^+ p$ , $t < 0.4$	
145 ± 12	2.2k	<sup>3,11</sup> HYAMS	68	OSPK 11.2 $\pi^- p$	
163 ± 15	13.3k	<sup>12</sup> PISUT	68	RVUE 1.7–3.2 $\pi^- p$ , $t < 10$	

<sup>1</sup> From pole extrapolation.

<sup>2</sup> From phase shift analysis of GRAYER 74 data.

<sup>3</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>4</sup> Published values contain misprints. Corrected by private communication RATCLIFF 74.

<sup>5</sup> From a fit to  $\pi^+ \pi^-$  mass using  $\rho(770)$  (parametrized with the Gounaris-Sakurai approach),  $\omega(782)$ , and box anomaly components.

<sup>6</sup> From a fit to  $\pi^+ \pi^-$  mass using  $\rho(770)$  (parametrized with the Gounaris-Sakurai approach),  $\omega(782)$ , and  $\rho(1450)$  components.

<sup>7</sup> Using relativistic Breit-Wigner and taking into account  $\rho$ - $\omega$  interference.

<sup>8</sup> Systematic errors not evaluated.

<sup>9</sup> From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of  $P$ -wave intensity. CHABAUD 83 includes data of GRAYER 74.

<sup>10</sup> HEYN 81 includes all spacelike and timelike  $F_\pi$  values until 1978.

<sup>11</sup> Of HYAMS 68 six parametrizations this is theoretically soundest. MR

<sup>12</sup> Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.

NODE=M009W;LINKAGE=C

NODE=M009W;LINKAGE=H

NODE=M009W;LINKAGE=Z

NODE=M009W;LINKAGE=03

NODE=M009W0R;LINKAGE=B

NODE=M009W0R;LINKAGE=C

NODE=M009W;LINKAGE=BL

NODE=M009W;LINKAGE=A1

NODE=M009W;LINKAGE=G

NODE=M009W;LINKAGE=B

NODE=M009W;LINKAGE=02

NODE=M009W;LINKAGE=R

## $\Gamma_{\rho(770)^0} - \Gamma_{\rho(770)^\pm}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.3 ± 1.3 OUR AVERAGE</b>		Error includes scale factor of 1.4.		
-0.2 ± 1.0		<sup>1</sup> SCHAEAL	05C	ALEP $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
3.6 ± 1.8 ± 1.7	1.98M	<sup>2</sup> ALOISIO	03	KLOE 1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.66 ± 0.85		<sup>3</sup> BARTOS	17A	RVUE $e^+ e^- \rightarrow \pi^+ \pi^-, \tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
-------------	--	---------------------	-----	---

NODE=M009W6

NODE=M009W6

<sup>1</sup> From the combined fit of the  $\tau^-$  data from ANDERSON 00A and SCHAEEL 05C and  $e^+e^-$  data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

<sup>2</sup> Assuming  $m_{\rho^+} = m_{\rho^-}$ ,  $\Gamma_{\rho^+} = \Gamma_{\rho^-}$ .

<sup>3</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

$$\Gamma_{\rho(770)^+} = \Gamma_{\rho(770)^-}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.8±2.0±0.5</b>	1.98M	<sup>1</sup> ALOISIO	03	KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> Without limitations on masses and widths.

NODE=M009W6;LINKAGE=SC

NODE=M009W6;LINKAGE=CH

NODE=M009W6;LINKAGE=A

NODE=M009W16

NODE=M009W16

NODE=M009W16;LINKAGE=WO

NODE=M009225;NODE=M009

### $\rho(770)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\pi\pi$	$\sim 100$	%
$\Gamma_2$ $K\bar{K}$		
<b><math>\rho(770)^\pm</math> decays</b>		
$\Gamma_3$ $\pi^\pm\pi^0$	$\sim 100$	%
$\Gamma_4$ $\pi^\pm\gamma$	( 4.5 ± 0.5 )	$\times 10^{-4}$ S=2.2
$\Gamma_5$ $\pi^\pm\eta$	< 6	$\times 10^{-3}$ CL=84%
$\Gamma_6$ $\pi^\pm\pi^+\pi^-\pi^0$	< 2.0	$\times 10^{-3}$ CL=84%
<b><math>\rho(770)^0</math> decays</b>		
$\Gamma_7$ $\pi^+\pi^-$	$\sim 100$	%
$\Gamma_8$ $\pi^+\pi^-\gamma$	( 9.9 ± 1.6 )	$\times 10^{-3}$
$\Gamma_9$ $\pi^0\gamma$	( 4.7 ± 0.8 )	$\times 10^{-4}$ S=1.7
$\Gamma_{10}$ $\eta\gamma$	( 3.00±0.21 )	$\times 10^{-4}$
$\Gamma_{11}$ $\pi^0\pi^0\gamma$	( 4.5 ± 0.8 )	$\times 10^{-5}$
$\Gamma_{12}$ $\mu^+\mu^-$	[a] ( 4.55±0.28 )	$\times 10^{-5}$
$\Gamma_{13}$ $e^+e^-$	[a] ( 4.72±0.05 )	$\times 10^{-5}$
$\Gamma_{14}$ $\pi^+\pi^-\pi^0$	( 1.01 <sup>+0.54</sup> <sub>-0.36</sub> ± 0.34 )	$\times 10^{-4}$
$\Gamma_{15}$ $\pi^+\pi^-\pi^+\pi^-$	( 1.8 ± 0.9 )	$\times 10^{-5}$
$\Gamma_{16}$ $\pi^+\pi^-\pi^0\pi^0$	( 1.6 ± 0.8 )	$\times 10^{-5}$
$\Gamma_{17}$ $\pi^0e^+e^-$	< 1.2	$\times 10^{-5}$ CL=90%
$\Gamma_{18}$ $\eta e^+e^-$		

DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=81

NODE=M009;CLUMP=A  
DESIG=11;OUR EVAL;→ UNCHECKED ←  
DESIG=3  
DESIG=5  
DESIG=21

NODE=M009;CLUMP=B  
DESIG=12;OUR EVAL;→ UNCHECKED ←  
DESIG=60  
DESIG=40  
DESIG=8  
DESIG=80  
DESIG=6  
DESIG=4  
DESIG=7;OUR EVAL;→ UNCHECKED ←  
DESIG=22  
DESIG=30  
DESIG=9  
DESIG=10

[a] The  $\omega\rho$  interference is then due to  $\omega\rho$  mixing only, and is expected to be small. If  $e\mu$  universality holds,  $\Gamma(\rho^0 \rightarrow \mu^+\mu^-) = \Gamma(\rho^0 \rightarrow e^+e^-) \times 0.99785$ .

LINKAGE=MD2

### CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 10 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 10.7$  for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_4$		-100	
$\Gamma$		15	-15
		$x_3$	$x_4$

Mode	Rate (MeV)	Scale factor
$\Gamma_3$ $\pi^\pm\pi^0$	150.2 ± 2.4	
$\Gamma_4$ $\pi^\pm\gamma$	0.068±0.007	2.3

DESIG=11

DESIG=3

### CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 7 branching ratios uses 21 measurements and one constraint to determine 9 parameters. The overall fit has a  $\chi^2 = 9.5$  for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_8$	-100								
$x_9$	-5	0							
$x_{10}$	-1	0	1						
$x_{11}$	-1	0	0	0					
$x_{12}$	2	-3	0	0	0				
$x_{13}$	0	0	-6	-9	0	0			
$x_{15}$	-1	0	0	0	0	0	0		
$\Gamma$	0	0	3	5	0	0	-54	0	
	$x_7$	$x_8$	$x_9$	$x_{10}$	$x_{11}$	$x_{12}$	$x_{13}$	$x_{15}$	

Mode	Rate (MeV)	Scale factor
$\Gamma_7$ $\pi^+ \pi^-$	147.5 $\pm$ 0.9	
$\Gamma_8$ $\pi^+ \pi^- \gamma$	1.48 $\pm$ 0.24	
$\Gamma_9$ $\pi^0 \gamma$	0.070 $\pm$ 0.012	1.7
$\Gamma_{10}$ $\eta \gamma$	0.0447 $\pm$ 0.0032	
$\Gamma_{11}$ $\pi^0 \pi^0 \gamma$	0.0066 $\pm$ 0.0012	
$\Gamma_{12}$ $\mu^+ \mu^-$	[a] 0.0068 $\pm$ 0.0004	
$\Gamma_{13}$ $e^+ e^-$	[a] 0.00704 $\pm$ 0.00006	
$\Gamma_{15}$ $\pi^+ \pi^- \pi^+ \pi^-$	0.0027 $\pm$ 0.0014	

DESIG=12  
DESIG=60  
DESIG=40  
DESIG=8  
DESIG=80  
DESIG=6  
DESIG=4  
DESIG=22

### $\rho(770)$ PARTIAL WIDTHS

NODE=M009230


#### $\Gamma(\pi^\pm \gamma)$

$\Gamma_4$

NODE=M009W3  
NODE=M009W3

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>68 <math>\pm</math> 7 OUR FIT</b>	Error includes scale factor of 2.3.			
<b>68 <math>\pm</math> 7 OUR AVERAGE</b>	Error includes scale factor of 2.2. See the ideogram below.			
81 $\pm$ 4 $\pm$ 4	CAPRARO	87	SPEC -	200 $\pi^- A \rightarrow \pi^- \pi^0 A$
59.8 $\pm$ 4.0	HUSTON	86	SPEC +	202 $\pi^+ A \rightarrow \pi^+ \pi^0 A$
71 $\pm$ 7	JENSEN	83	SPEC -	156-260 $\pi^- A \rightarrow \pi^- \pi^0 A$

WEIGHTED AVERAGE  
68 $\pm$ 7 (Error scaled by 2.2)



$\Gamma(\pi^0\gamma)$  $\Gamma_9$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$77 \pm 17 \pm 11$	36500	<sup>1</sup> ACHASOV 03	SND	$0.60-0.97 e^+ e^- \rightarrow \pi^0 \gamma$
$121 \pm 31$		DOLINSKY 89	ND	$e^+ e^- \rightarrow \pi^0 \gamma$

<sup>1</sup> Using  $\Gamma_{\text{total}} = 147.9 \pm 1.3$  MeV and  $B(\rho \rightarrow \pi^0 \gamma)$  from ACHASOV 03.

NODE=M009W31  
NODE=M009W31

NODE=M009W31;LINKAGE=AV

 $\Gamma(\eta\gamma)$  $\Gamma_{10}$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$62 \pm 17$	<sup>1</sup> DOLINSKY 89	ND	$e^+ e^- \rightarrow \eta \gamma$
-------------	--------------------------	----	-----------------------------------

<sup>1</sup> Solution corresponding to constructive  $\omega$ - $\rho$  interference.

NODE=M009W32  
NODE=M009W32

NODE=M009W32;LINKAGE=L

 $\Gamma(e^+ e^-)$  $\Gamma_{13}$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**7.04 ± 0.06 OUR FIT**

**7.04 ± 0.06 OUR AVERAGE**

$7.048 \pm 0.057 \pm 0.050$	900k	<sup>1</sup> AKHMETSHIN 07		$e^+ e^- \rightarrow \pi^+ \pi^-$
$7.06 \pm 0.11 \pm 0.05$	114k	<sup>2,3</sup> AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^-$
$6.77 \pm 0.10 \pm 0.30$		BARKOV 85	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.12 \pm 0.02 \pm 0.11$	800k	<sup>4</sup> ACHASOV 06	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$6.3 \pm 0.1$		<sup>5</sup> BENAYOUN 98	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-, \mu^+ \mu^-$

<sup>1</sup> A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

<sup>2</sup> Using the GOUNARIS 68 parametrization with the complex phase of the  $\rho$ - $\omega$  interference.

<sup>3</sup> From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02.

<sup>4</sup> Supersedes ACHASOV 05A.

<sup>5</sup> Using the data of BARKOV 85 in the hidden local symmetry model.

NODE=M009W4  
NODE=M009W4

OCCUR=2

NODE=M009W4;LINKAGE=AK  
NODE=M009W4;LINKAGE=GS  
NODE=M009W4;LINKAGE=P2  
NODE=M009W4;LINKAGE=AC  
NODE=M009W4;LINKAGE=K2

 $\Gamma(\pi^+ \pi^- \pi^+ \pi^-)$  $\Gamma_{15}$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.8 \pm 1.4 \pm 0.5$	153	AKHMETSHIN 00	CMD2	$0.6-0.97 e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$
-----------------------	-----	---------------	------	--

NODE=M009W33  
NODE=M009W33

 $\rho(770) \Gamma(e^+ e^-) \Gamma(i) / \Gamma^2(\text{total})$ 

NODE=M009233

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}} \times \Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_{13} / \Gamma \times \Gamma_7 / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**4.89 ± 0.04 OUR AVERAGE**

$4.889 \pm 0.015 \pm 0.039$		<sup>1</sup> ACHASOV 21	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$4.876 \pm 0.023 \pm 0.064$	800k	<sup>2,3</sup> ACHASOV 06	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.72 \pm 0.02$		<sup>4</sup> BENAYOUN 10	RVUE	$0.4-1.05 e^+ e^-$
-----------------	--	--------------------------	------	--------------------

<sup>1</sup> From a fit of the cross section in the energy range  $0.525 < \sqrt{s} < 0.883$  GeV parameterized by the sum of the Breit-Wigner amplitudes for the  $\rho(770)$ ,  $\omega$  and  $\rho(1450)$  resonances.

<sup>2</sup> Supersedes ACHASOV 05A.

<sup>3</sup> A fit of the SND data from 400 to 1000 MeV using parameters of the  $\rho(1450)$  and  $\rho(1700)$  from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

<sup>4</sup> A simultaneous fit of  $e^+ e^- \rightarrow \pi^+ \pi^-, \pi^+ \pi^- \pi^0, \pi^0 \gamma, \eta \gamma$  data.

NODE=M009G4;LINKAGE=A

NODE=M009G4;LINKAGE=AC  
NODE=M009G4;LINKAGE=SN

NODE=M009G4;LINKAGE=BE

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}} \times \Gamma(\eta\gamma) / \Gamma_{\text{total}}$  $\Gamma_{13} / \Gamma \times \Gamma_{10} / \Gamma$ 

VALUE (units $10^{-8}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.42 ± 0.10 OUR FIT**

**1.45 ± 0.12 OUR AVERAGE**

$1.32 \pm 0.14 \pm 0.08$	33k	<sup>1</sup> ACHASOV 07B	SND	$0.6-1.38 e^+ e^- \rightarrow \eta \gamma$
$1.50 \pm 0.65 \pm 0.09$	17.4k	<sup>2</sup> AKHMETSHIN 05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta \gamma$
$1.61 \pm 0.20 \pm 0.11$	23k	<sup>3,4</sup> AKHMETSHIN 01B	CMD2	$e^+ e^- \rightarrow \eta \gamma$
$1.85 \pm 0.49$		<sup>5</sup> DOLINSKY 89	ND	$e^+ e^- \rightarrow \eta \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.05 \pm 0.02$		<sup>6</sup> BENAYOUN 10	RVUE	$0.4-1.05 e^+ e^-$
-----------------	--	--------------------------	------	--------------------

NODE=M009G1  
NODE=M009G1



- <sup>1</sup> From a combined fit of  $\sigma(e^+e^- \rightarrow \eta\gamma)$  with  $\eta \rightarrow 3\pi^0$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$ , and fixing  $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$ . Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- <sup>2</sup> From the  $\eta \rightarrow 2\gamma$  decay and using  $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$ .
- <sup>3</sup> From the  $\eta \rightarrow 3\pi^0$  decay and using  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ .
- <sup>4</sup> The combined fit from 600 to 1380 MeV taking into account  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ , and  $\rho(1450)$  (mass and width fixed at 1450 MeV and 310 MeV respectively).
- <sup>5</sup> Recalculated by us from the cross section in the peak.
- <sup>6</sup> A simultaneous fit of  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$  data.

NODE=M009G1;LINKAGE=AH

NODE=M009G;LINKAGE=AH  
 NODE=M009G;LINKAGE=AK  
 NODE=M009G;LINKAGE=BQ

NODE=M009G;LINKAGE=LP  
 NODE=M009G1;LINKAGE=BE

**$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\pi^0\gamma)/\Gamma_{total}$   $\Gamma_{13}/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units  $10^{-8}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**2.2 ± 0.4 OUR FIT** Error includes scale factor of 1.7.

**2.21 ± 0.34 OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.

1.98 ± 0.22 ± 0.10    <sup>1</sup> ACHASOV    16A    SND    0.60-1.38  $e^+e^- \rightarrow \pi^0\gamma$

2.90 <sup>+0.60</sup>/<sub>-0.55</sub> ± 0.18    18k    AKHMETSHIN 05    CMD2    0.60-1.38  $e^+e^- \rightarrow \pi^0\gamma$

3.61 ± 0.74 ± 0.49    10k    <sup>2</sup> DOLINSKY    89    ND     $e^+e^- \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.875 ± 0.026    <sup>3</sup> BENAYOUN    10    RVUE    0.4-1.05  $e^+e^-$

2.37 ± 0.53 ± 0.33    36k    <sup>4</sup> ACHASOV    03    SND    0.60-0.97  $e^+e^- \rightarrow \pi^0\gamma$

<sup>1</sup> From the VMD model with the  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$  resonances, and an additional resonance describing the total contribution of the  $\rho(1450)$  and  $\omega(1420)$  states. Supersedes ACHASOV 03.

<sup>2</sup> Recalculated by us from the cross section in the peak.

<sup>3</sup> A simultaneous fit of  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$  data.

<sup>4</sup> Using  $\sigma_{\phi \rightarrow \pi^0\gamma}$  from ACHASOV 00 and  $m_{\rho} = 775.97$  MeV in the model with the energy-independent phase of  $\rho$ - $\omega$  interference equal to  $(-10.2 \pm 7.0)^\circ$ .


NODE=M009G2  
 NODE=M009G2

NODE=M009G2;LINKAGE=B

NODE=M009G2;LINKAGE=LP  
 NODE=M009G2;LINKAGE=BE

NODE=M009G;LINKAGE=SH

WEIGHTED AVERAGE  
 2.21±0.34 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

**$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$   $\Gamma_{13}/\Gamma \times \Gamma_{14}/\Gamma$**

VALUE (units  $10^{-9}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.903 ± 0.076    <sup>1</sup> BENAYOUN    10    RVUE    0.4-1.05  $e^+e^-$

4.58 <sup>+2.46</sup>/<sub>-1.64</sub> ± 1.56    1.2M    <sup>2</sup> ACHASOV    03D    RVUE    0.44-2.00  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> A simultaneous fit of  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$  data.

<sup>2</sup> Statistical significance is less than  $3\sigma$ .

NODE=M009G3  
 NODE=M009G3

NODE=M009G3;LINKAGE=BE  
 NODE=M009G3;LINKAGE=AC

**$\rho(770)$  BRANCHING RATIOS**

NODE=M009235

**$\Gamma(\pi^\pm\eta)/\Gamma(\pi\pi)$   $\Gamma_5/\Gamma_1$**

VALUE (units  $10^{-4}$ )    CL%    DOCUMENT ID    TECN    CHG    COMMENT

<60    84    FERBEL    66    HBC    ±     $\pi^\pm p$  above 2.5

NODE=M009R4  
 NODE=M009R4

$\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)/\Gamma(\pi\pi)$  $\Gamma_6/\Gamma_1$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<20	84	FERBEL	66	HBC	$\pm$ $\pi^\pm p$ above 2.5
• • • We do not use the following data for averages, fits, limits, etc. • • •					
35 $\pm$ 40		JAMES	66	HBC	+ 2.1 $\pi^+ p$

NODE=M009R1  
 NODE=M009R1

 $\Gamma(\pi^+ \pi^- \gamma)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.0099 <math>\pm</math> 0.0016 OUR FIT</b>				
<b>0.0099 <math>\pm</math> 0.0016</b>		<sup>1</sup> DOLINSKY	91	ND $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0111 $\pm$ 0.0014		<sup>2</sup> VASSERMAN	88	ND $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<0.005	90	<sup>3</sup> VASSERMAN	88	ND $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

NODE=M009R12  
 NODE=M009R12

- <sup>1</sup> Bremsstrahlung from a decay pion and for photon energy above 50 MeV.  
<sup>2</sup> Superseded by DOLINSKY 91.  
<sup>3</sup> Structure radiation due to quark rearrangement in the decay.

OCCUR=2

NODE=M009R12;LINKAGE=J  
 NODE=M009R12;LINKAGE=I  
 NODE=M009R12;LINKAGE=N

 $\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.20 $\pm$ 0.52		<sup>1</sup> ACHASOV	16A	SND 0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
6.21 $^{+1.28}_{-1.18} \pm 0.39$	18k	<sup>2,3</sup> AKHMETSHIN	05	CMD2 0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
5.22 $\pm 1.17 \pm 0.75$	36k	<sup>3,4</sup> ACHASOV	03	SND 0.60-0.97 $e^+ e^- \rightarrow \pi^0 \gamma$
6.8 $\pm 1.7$		<sup>5</sup> BENAYOUN	96	RVUE 0.54-1.04 $e^+ e^- \rightarrow \pi^0 \gamma$
7.9 $\pm 2.0$		<sup>3</sup> DOLINSKY	89	ND $e^+ e^- \rightarrow \pi^0 \gamma$

NODE=M009R9  
 NODE=M009R9

- <sup>1</sup> Using  $B(\rho \rightarrow e^+ e^-)$  from PDG 15. Supersedes ACHASOV 03.  
<sup>2</sup> Using  $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$ .  
<sup>3</sup> Not independent of the corresponding  $\Gamma(e^+ e^-) \times \Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}^2$ .  
<sup>4</sup> Using  $B(\rho \rightarrow e^+ e^-) = (4.54 \pm 0.10) \times 10^{-5}$ .  
<sup>5</sup> Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

NODE=M009R9;LINKAGE=C  
 NODE=M009R9;LINKAGE=AK  
 NODE=M009R9;LINKAGE=BZ  
 NODE=M009R9;LINKAGE=AS  
 NODE=M009R9;LINKAGE=A

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>3.00 <math>\pm</math> 0.21 OUR FIT</b>					
<b>2.90 <math>\pm</math> 0.32 OUR AVERAGE</b>					
2.79 $\pm 0.34 \pm 0.03$	33k	<sup>1</sup> ACHASOV	07B	SND	0.6-1.38 $e^+ e^- \rightarrow \eta\gamma$
3.6 $\pm 0.9$		<sup>2</sup> ANDREWS	77	CNTR 0	6.7-10 $\gamma$ Cu
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.21 $\pm 1.39 \pm 0.20$	17.4k	<sup>3,4</sup> AKHMETSHIN	05	CMD2	0.60-1.38 $e^+ e^- \rightarrow \eta\gamma$
3.39 $\pm 0.42 \pm 0.23$		<sup>2,5,6</sup> AKHMETSHIN	01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
1.9 $^{+0.6}_{-0.8}$		<sup>7</sup> BENAYOUN	96	RVUE	0.54-1.04 $e^+ e^- \rightarrow \eta\gamma$
4.0 $\pm 1.1$		<sup>2,4</sup> DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$

NODE=M009R7  
 NODE=M009R7

- <sup>1</sup> ACHASOV 07B reports  $[\Gamma(\rho(770) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\rho(770) \rightarrow e^+ e^-)] = (1.32 \pm 0.14 \pm 0.08) \times 10^{-8}$  which we divide by our best value  $B(\rho(770) \rightarrow e^+ e^-) = (4.72 \pm 0.05) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.  
<sup>2</sup> Solution corresponding to constructive  $\omega$ - $\rho$  interference.  
<sup>3</sup> Using  $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$  and  $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$ .  
<sup>4</sup> Not independent of the corresponding  $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$ .  
<sup>5</sup> The combined fit from 600 to 1380 MeV taking into account  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ , and  $\rho(1450)$  (mass and width fixed at 1450 MeV and 310 MeV respectively).  
<sup>6</sup> Using  $B(\rho \rightarrow e^+ e^-) = (4.75 \pm 0.10) \times 10^{-5}$  from AKHMETSHIN 02 and  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ .  
<sup>7</sup> Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution. Constructive  $\rho$ - $\omega$  interference solution.

NODE=M009R7;LINKAGE=AO

NODE=M009R7;LINKAGE=A  
 NODE=M009R;LINKAGE=AK  
 NODE=M009R7;LINKAGE=AZ  
 NODE=M009R;LINKAGE=BQ

NODE=M009R;LINKAGE=BX

NODE=M009R7;LINKAGE=C

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{11}/\Gamma$ NODE=M009R14  
NODE=M009R14**4.5±0.8 OUR FIT****4.5<sup>+0.9</sup><sub>-0.8</sub> OUR AVERAGE**5.2<sup>+1.5</sup><sub>-1.3</sub>±0.6 190 <sup>1</sup> AKHMETSHIN 04B CMD2 0.6–0.97  $e^+e^- \rightarrow \pi^0\pi^0\gamma$ 

OCCUR=2

4.1<sup>+1.0</sup><sub>-0.9</sub>±0.3 295 <sup>2</sup> ACHASOV 02F SND 0.36–0.97  $e^+e^- \rightarrow \pi^0\pi^0\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.8<sup>+3.4</sup><sub>-1.8</sub>±0.5 63 <sup>3</sup> ACHASOV 00G SND  $e^+e^- \rightarrow \pi^0\pi^0\gamma$ 

<sup>1</sup> This branching ratio includes the conventional VMD mechanism  $\rho \rightarrow \omega\pi^0$ ,  $\omega \rightarrow \pi^0\gamma$ , and the new decay mode  $\rho \rightarrow f_0(500)\gamma$ ,  $f_0(500) \rightarrow \pi^0\pi^0$  with a branching ratio  $(2.0^{+1.1}_{-0.9} \pm 0.3) \times 10^{-5}$  differing from zero by 2.0 standard deviations.

NODE=M009R14;LINKAGE=AH

<sup>2</sup> This branching ratio includes the conventional VMD mechanism  $\rho \rightarrow \omega\pi^0$ ,  $\omega \rightarrow \pi^0\gamma$  and the new decay mode  $\rho \rightarrow f_0(500)\gamma$ ,  $f_0(500) \rightarrow \pi^0\pi^0$  with a branching ratio  $(1.9^{+0.9}_{-0.8} \pm 0.4) \times 10^{-5}$  differing from zero by 2.4 standard deviations. Supersedes ACHASOV 00G.

NODE=M009R;LINKAGE=FF

<sup>3</sup> Superseded by ACHASOV 02F.

NODE=M009R;LINKAGE=GF

 $\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-)$ VALUE (units  $10^{-5}$ )

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{12}/\Gamma_7$ NODE=M009R5  
NODE=M009R5**4.60±0.28 OUR FIT****4.6 ±0.2 ±0.2**

ANTIPOV

89

SIGM

 $\pi^- \text{Cu} \rightarrow \mu^+\mu^-\pi^- \text{Cu}$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.2<sup>+1.6</sup><sub>-3.6</sub> <sup>1</sup> ROTHWELL 69 CNTR Photoproduction5.6 ±1.5 <sup>2</sup> WEHMANN 69 OSPK 12  $\pi^- \text{C, Fe}$ 9.7<sup>+3.1</sup><sub>-3.3</sub> <sup>3,4</sup> HYAMS 67 OSPK 11  $\pi^- \text{Li, H}$ 

<sup>1</sup> Possibly large  $\rho$ - $\omega$  interference leads us to increase the minus error.

NODE=M009R5;LINKAGE=R

<sup>2</sup> Result contains  $11 \pm 11\%$  correction using SU(3) for central value. The error on the correction takes account of possible  $\rho$ - $\omega$  interference and the upper limit agrees with the upper limit of  $\omega \rightarrow \mu^+\mu^-$  from this experiment.

NODE=M009R5;LINKAGE=W

<sup>3</sup> But he even enlarges his error to take residual  $\omega$  contamination into account. Since his value is high, seems the other experiments also can't have too many  $\omega$ 's. But maybe Hyams has additional  $\mu$ 's from  $\rho \rightarrow \pi\pi$ , decaying  $\pi$ 's.

NODE=M009R5;LINKAGE=01

<sup>4</sup> HYAMS 67's mass resolution is 20 MeV. The  $\omega$  region was excluded.

NODE=M009R5;LINKAGE=H

 $\Gamma(e^+e^-)/\Gamma(\pi\pi)$ VALUE (units  $10^{-4}$ )

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{13}/\Gamma_1$ NODE=M009R3  
NODE=M009R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.40±0.05 <sup>1,2</sup> BENAJSAS 72 OSPK  $e^+e^- \rightarrow \pi^+\pi^-$ 

<sup>1</sup> The  $\rho'$  contribution is not taken into account.

NODE=M009R;LINKAGE=KS

<sup>2</sup> Barkov excludes Auslender and Benaksas for large statistical and systematic errors.

NODE=M009R3;LINKAGE=01

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )

CL% EVTS

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{14}/\Gamma$ NODE=M009R10  
NODE=M009R10**0.88±0.23±0.30**<sup>1</sup> LEES

21B

BABR

10.5  $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.01<sup>+0.54</sup><sub>-0.36</sub>±0.34 1.2M <sup>2</sup> ACHASOV 03D RVUE 0.44–2.00  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ <1.2 90 VASSERMAN 88B ND  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ 

<sup>1</sup> From the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$  with contributions from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ . Statistical evidence is more than  $6\sigma$ .

NODE=M009R10;LINKAGE=A

<sup>2</sup> Statistical significance is less than  $3\sigma$ .

NODE=M009R;LINKAGE=NS

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi\pi)$ 

VALUE

CL%

DOCUMENT ID

TECN

CHG

COMMENT

 $\Gamma_{14}/\Gamma_1$ NODE=M009R6  
NODE=M009R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

~0.01 BRAMON 86 RVUE 0  $J/\psi \rightarrow \omega\pi^0$ <0.01 84 <sup>1</sup> ABRAMS 71 HBC 0  $3.7\pi^+\pi^-$ 

<sup>1</sup> Model dependent, assumes  $l = 1, 2, \text{ or } 3$  for the  $3\pi$  system.

NODE=M009R6;LINKAGE=G

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.8±0.9 OUR FIT</b>					
<b>1.8±0.9±0.3</b>	153		AKHMETSHIN 00	CMD2	0.6–0.97 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<20	90		KURDADZE 88	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M009R13  
 NODE=M009R13

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma(\pi\pi)$   $\Gamma_{15}/\Gamma_1$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<15	90	ERBE 69	HBC	0	2.5–5.8 $\gamma p$
<20		CHUNG 68	HBC	0	3.2,4.2 $\pi^- p$
<20	90	HUSON 68	HLBC	0	16.0 $\pi^- p$
<80		JAMES 66	HBC	0	2.1 $\pi^+ p$

NODE=M009R11  
 NODE=M009R11

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>1.60±0.74±0.18</b>		<sup>1</sup> ACHASOV 09A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 4	90	AULCHENKO 87C	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
<20	90	KURDADZE 86	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

NODE=M009R8  
 NODE=M009R8

<sup>1</sup> Assuming no interference between the  $\rho$  and  $\omega$  contributions.

NODE=M009R8;LINKAGE=AC

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.2</b>	90	ACHASOV 08	SND	0.36–0.97 $e^+e^- \rightarrow \pi^0 e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<1.6		AKHMETSHIN 05A	CMD2	0.72–0.84 $e^+e^-$

NODE=M009R15  
 NODE=M009R15

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<0.7	AKHMETSHIN 05A	CMD2	0.72–0.84 $e^+e^-$

NODE=M009R16  
 NODE=M009R16

 **$\rho(770)$  REFERENCES**

ACHASOV 21	JHEP 2101 113	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61040
LEES 21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
ANDREEV 20	EPJ C80 1189	V. Andreev <i>et al.</i>	(H1 Collab.)	REFID=60773
ABLIKIM 18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
ABLIKIM 17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57650
BARTOS 17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS 17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
ABLIKIM 16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV 16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57513
PDG 15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=2
ABRAMOWICZ 12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
LEES 12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
AMBROSINO 11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
GARCIA-MAR... 11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=16761
BATLEY 10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	REFID=53567
BENAYOUN 10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
DUBNICKA 10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
ACHASOV 09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53101
	Translated from ZETF 136 442.			
AUBERT 09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
ACHASOV 08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52258
	Translated from ZETF 134 80.			
FUJIKAWA 08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
ACHASOV 07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AKHMETSHIN 07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV 06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
	Translated from ZETF 130 437.			
ACHASOV 06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AULCHENKO 06A	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51513
	Translated from ZETFP 84 491.			
ACHASOV 05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51045
	Translated from ZETF 128 1201.			
AKHMETSHIN 05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AKHMETSHIN 05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
ALOISIO 05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
AULCHENKO 05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
	Translated from ZETFP 82 841.			

NODE=M009

SCHAEL	05C	PRPL 421 191	S. Schael <i>et al.</i>	(ALEPH Collab.)	REFID=50845
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49610
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49187
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
SANZ-CILLERO	03	EPJ C27 587	J.J. Sanz-Cillero, A. Pich		REFID=49399
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48816
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48565
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler		REFID=49180
PICH	01	PR D63 093005	A. Pich, J. Portoles		REFID=48313
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
ACHASOV	00G	Translated from ZETFP 72 411. JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47929
AKHMETSHIN	00A	Translated from ZETFP 71 519. PL B475 190	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47421
ANDERSON	00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)	REFID=47468
ABELE	99E	PL B469 270	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47414
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)	REFID=45859
BREITWEG	98B	EPJ C2 247	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=46354
GARDNER	98	PR D57 2716	S. Gardner, H.B. O'Connell		REFID=46366
		PR D62 019903 (err.)	S. Gardner, H.B. O'Connell		REFID=47981
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ADAMS	97	ZPHY C74 237	M.R. Adams <i>et al.</i>	(E665 Collab.)	REFID=45533
BARATE	97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45622
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
BOGOLYUB...	97	PAN 60 46	M.Y. Bogolyubsky <i>et al.</i>	(MOSU, SERP)	REFID=45393
O'CONNELL	97	Translated from YAF 60 53. NP A623 559	H.B. O'Connell <i>et al.</i>	(ADLD)	REFID=45860
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
BERNICA	94	PR D50 4454	A. Bernica, G. Lopez Castro, J. Pestieau	(LOUV+)	REFID=44097
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
AGUILAR...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ANTIPOV	89	ZPHY C42 185	Y.M. Antipov <i>et al.</i>	(SERP, JINR, BGNA+)	REFID=40739
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
KURDADZE	88	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=41121
VASSERMAN	88	Translated from ZETFP 47 432. SJNP 47 1035	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=41019
VASSERMAN	88B	Translated from YAF 47 1035. SJNP 48 480	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=41020
AULCHENKO	87C	Translated from YAF 48 753. IYF 87-90 Preprint	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=41370
CAPRARO	87	NP B288 659	L. Capraro <i>et al.</i>	(CLER, FRAS, MILA+)	REFID=40003
BRAMON	86	PL B173 97	A. Bramon, J. Casulleras	(BARC)	REFID=22102
HUSTON	86	PR D33 3199	J. Huston <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=20137
KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
BARKOV	85	Translated from ZETFP 43 497. NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
JENSEN	83	PR D27 26	T. Jensen <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=20132
HEYN	81	ZPHY C7 169	M.F. Heyn, C.B. Lang	(GRAZ)	REFID=20129
BOHACIK	80	PR D21 1342	J. Bohacik, H. Kuhnelt	(SLOV, WIEN)	REFID=20128
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL)	REFID=20381
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)	REFID=20126
BARTALUCCI	78	NC 44A 587	S. Bartalucci <i>et al.</i>	(DESY, FRAS)	REFID=20122
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
DEUTSCH...	76	NP B103 426	M. Deuschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
RATCLIFF	74	Private Comm.			REFID=40128 ERROR=3
BYERLY	73	PR D7 637	W.L. Byerly <i>et al.</i>	(MICH)	REFID=20104
GLADDING	73	PR D8 3721	G.E. Gladding <i>et al.</i>	(HARV)	REFID=20106
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108
BALLAM	72	PR D5 545	J. Ballam <i>et al.</i>	(SLAC, LBL, TUFTS)	REFID=20094
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20096
JACOBS	72	PR D6 1291	L.D. Jacobs	(SACL)	REFID=20101
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)	REFID=20102
ABRAMS	71	PR D4 653	G.S. Abrams <i>et al.</i>	(LBL)	REFID=20090
ALVENSLEB...	70	PRL 24 786	H. Alvensleben <i>et al.</i>	(DESY)	REFID=20085
BIGGS	70	PRL 24 1197	P.J. Biggs <i>et al.</i>	(DARE)	REFID=20087
ERBE	69	PR 188 2060	R. Erbe <i>et al.</i>	(German Bubble Chamber Collab.)	REFID=20074
MALAMUD	69	Argonne Conf. 93	E.I. Malamud, P.E. Schlein	(UCLA)	REFID=20077
REYNOLDS	69	PR 184 1424	B.G. Reynolds <i>et al.</i>	(FSU)	REFID=20080
ROTHWELL	69	PRL 23 1521	P.L. Rothwell <i>et al.</i>	(NEAS)	REFID=20082
WEHMANN	69	PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20084
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
BATON	68	PR 176 1574	J.P. Baton, G. Laurens	(SACL)	REFID=20056
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
FOSTER	68	NP B6 107	M. Foster <i>et al.</i>	(CERN, CDEF)	REFID=20061
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054
HUSON	68	PL 28B 208	R. Huson <i>et al.</i>	(ORSAY, MILA, UCLA)	REFID=20062
HYAMS	68	NP B7 1	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20063
LANZEROTTI	68	PR 166 1365	L.J. Lanzerotti <i>et al.</i>	(HARV)	REFID=20068
PISUT	68	NP B6 325	J. Pisut, M. Roos	(CERN)	REFID=20070
ASBURY	67B	PRL 19 865	J.G. Asbury <i>et al.</i>	(DESY, COLU)	REFID=20038
BACON	67	PR 157 1263	T.C. Bacon <i>et al.</i>	(BNL)	REFID=20039
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
HUWE	67	PL 24B 252	D.O. Huwe <i>et al.</i>	(COLU)	REFID=20049
HYAMS	67	PL 24B 634	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20050
MILLER	67B	PR 153 1423	D.H. Miller <i>et al.</i>	(PURD)	REFID=20051
ALFF...	66	PR 145 1072	C. Alff-Steinberger <i>et al.</i>	(COLU, RUTG)	REFID=10762
FERBEL	66	PL 21 111	T. Ferbel	(ROCH)	REFID=20028
HAGOPIAN	66	PR 145 1128	V. Hagopian <i>et al.</i>	(PENN, SACL)	REFID=20030
HAGOPIAN	66B	PR 152 1183	V. Hagopian, Y.L. Pan	(PENN, LRL)	REFID=20031
JACOBS	66B	UCRL 16877	L.D. Jacobs	(LRL)	REFID=20033
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
ROSS	66	PR 149 1172	M. Ross, L. Stodolsky		REFID=46380
SOEDING	66	PL B19 702	P. Soeding		REFID=46385
WEST	66	PR 149 1089	E. West <i>et al.</i>	(WISC)	REFID=20035
BLIEDEN	65	PL 19 444	H.R. Blieden <i>et al.</i>	(CERN MMS Collab.)	REFID=20016
CARMONY	64	PRL 12 254	D.D. Carmony <i>et al.</i>	(UCB)	REFID=20578
GOLDHABER	64	PRL 12 336	G. Goldhaber <i>et al.</i>	(LRL, UCB)	REFID=20013
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)	REFID=20006

$\omega(782)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M001

 $\omega(782)$  MASS

NODE=M001M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>782.66±0.13 OUR AVERAGE</b>		Error includes scale factor of 2.0. See the ideogram below.		
777.9 ±2.2 $\begin{smallmatrix} +4.3 \\ -2.2 \end{smallmatrix}$	900k	ANDREEV	20 H1	$e p \rightarrow e \pi^+ \pi^- p$
783.20±0.13±0.16	18680	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
782.68±0.09±0.04	11200	<sup>1</sup> AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.79±0.08±0.09	1.2M	<sup>2</sup> ACHASOV 03D	RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
781.96±0.17±0.80	11k	<sup>3</sup> AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
782.08±0.36±0.82	3463	<sup>4</sup> AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
781.96±0.13±0.17	15k	AMSLER 93B	CBAR	0.0 $\bar{p} p \rightarrow \omega \pi^0 \pi^0$
782.4 ±0.2	270k	WEIDENAUER 93	ASTE	$\bar{p} p \rightarrow 2\pi^+ 2\pi^- \pi^0$
782.4 ±0.5	7000	<sup>5</sup> KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
782.58±0.03±0.01		<sup>6</sup> HOID 20	RVUE	$e^+ e^- \rightarrow \pi^0 \gamma$
781.68±0.09±0.03		<sup>7</sup> COLANGELO 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
782.63±0.03±0.01		<sup>8</sup> HOFERICHT... 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
781.91±0.24		<sup>9</sup> LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
782.7 ±0.1 ±1.5	19500	<sup>10</sup> WURZINGER 95	SPEC	1.33 $p d \rightarrow {}^3\text{He} \omega$
781.78±0.10		<sup>10</sup> BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.2 ±0.4	1488	<sup>11</sup> KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
783.3 ±0.4	433	CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.5 ±0.8	33260	ROOS 80	RVUE	0.0-3.6 $\bar{p} p$
782.6 ±0.8	3000	BENKHEIRI 79	OMEG	9-12 $\pi^\pm p$
781.8 ±0.6	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p} p \rightarrow 5\pi$
782.7 ±0.9	535	VANAPEL... 78	HBC	7.2 $\bar{p} p \rightarrow \bar{p} p \omega$
783.5 ±0.8	2100	GESSAROLI 77	HBC	11 $\pi^- p \rightarrow \omega n$
782.5 ±0.8	418	AGUILAR-... 72B	HBC	3.9,4.6 $K^- p$
783.4 ±1.0	248	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K^+ K^- \omega$
781.0 ±0.6	510	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K_1 K_1 \omega$
783.7 ±1.0	3583	<sup>12</sup> COYNE 71	HBC	3.7 $\pi^+ p \rightarrow \rho \pi^+ \pi^+ \pi^- \pi^0$
784.1 ±1.2	750	ABRAMOVI... 70	HBC	3.9 $\pi^- p$
783.2 ±1.6		<sup>13</sup> BIGGS 70B	CNTR	<4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
782.4 ±0.5	2400	BIZZARRI 69	HBC	0.0 $\bar{p} p$

NODE=M001M

OCCUR=2

OCCUR=2

OCCUR=2


<sup>1</sup> Update of AKHMETSHIN 00C.<sup>2</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+ \pi^- \pi^0$  and ANTONELLI 92 on the  $\omega \pi^+ \pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.<sup>3</sup> From the  $\eta \rightarrow \gamma \gamma$  decay.<sup>4</sup> From the  $\eta \rightarrow 3\pi^0$  decay.<sup>5</sup> Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.<sup>6</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives  $782.736 \pm 0.024$  MeV.<sup>7</sup> The  $\omega$  mass was extracted from a dispersively improved Breit-Wigner parameterization, the  $\omega$  width fixed at  $8.49 \pm 0.08$  MeV. The value does not include vacuum polarization which would shift the mass to  $781.81 \pm 0.09 \pm 0.03$  MeV. The mixing parameter is assumed real valued.<sup>8</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.<sup>9</sup> From the  $\rho - \omega$  interference in the  $\pi^+ \pi^-$  mass spectrum using the Breit-Wigner for the  $\omega$  and leaving its mass and width as free parameters of the fit.<sup>10</sup> Systematic uncertainties underestimated.<sup>11</sup> Systematic uncertainties not estimated.<sup>12</sup> From best-resolution sample of COYNE 71.<sup>13</sup> From  $\omega - \rho$  interference in the  $\pi^+ \pi^-$  mass spectrum assuming  $\omega$  width 12.6 MeV.NODE=M001M;LINKAGE=PT  
NODE=M001M;LINKAGE=VHNODE=M001M;LINKAGE=S1  
NODE=M001M;LINKAGE=S2  
NODE=M001M;LINKAGE=B  
NODE=M001M;LINKAGE=G

NODE=M001M;LINKAGE=A

NODE=M001M;LINKAGE=H

NODE=M001M;LINKAGE=LE

NODE=M001M;LINKAGE=KB  
NODE=M001M;LINKAGE=I  
NODE=M001M;LINKAGE=D  
NODE=M001M;LINKAGE=F



**ω(782) WIDTH**

NODE=M001W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.68±0.13 OUR AVERAGE</b>				
8.68±0.23±0.10	11200	1 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.68±0.04±0.15	1.2M	2 ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
8.65±0.06±0.01		3 HOID 20	RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.71±0.04±0.04		4 HOFERICH... 19	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.13±0.45		5 LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
8.2 ±0.3	19500	6 WURZINGER 95	SPEC	1.33 $p d \rightarrow {}^3\text{He}\omega$
8.4 ±0.1		7 AULCHENKO 87	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.30±0.40		6 BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.8 ±0.9	1488	8 KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0 ±0.8	433	6 CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
12 ±2	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p}p \rightarrow 5\pi$
9.4 ±2.5	2100	GESSAROLI 77	HBC	11 $\pi^-p \rightarrow \omega n$
10.22±0.43	20000	9 KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
13.3 ±2	418	AGUILAR... 72B	HBC	3.9,4.6 $K^-p$
9.1 ±0.8	451	6 BENAKSAS 72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
10.5 ±1.5		BORENSTEIN 72	HBC	2.18 $K^-p$
7.70±0.9 ±1.15	940	BROWN 72	MMS	2.5 $\pi^-p \rightarrow nMM$
10.3 ±1.4	510	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K_1^+K_1^-\omega$
12.8 ±3.0	248	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K^+K^-\omega$
9.5 ±1.0	3583	COYNE 71	HBC	3.7 $\pi^+p \rightarrow \rho\pi^+\pi^+\pi^-\pi^0$

NODE=M001W

OCCUR=2

<sup>1</sup> Update of AKHMETSHIN 00C.

NODE=M001W;LINKAGE=PT  
NODE=M001W;LINKAGE=VH

<sup>2</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

NODE=M001W;LINKAGE=E

<sup>3</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives  $8.63 \pm 0.05$  MeV.

NODE=M001W;LINKAGE=F

<sup>4</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

NODE=M001W;LINKAGE=LE

<sup>5</sup> From the  $\rho-\omega$  interference in the  $\pi^+\pi^-$  mass spectrum using the Breit-Wigner for the  $\omega$  and leaving its mass and width as free parameters of the fit.

NODE=M001W;LINKAGE=I

<sup>6</sup> Systematic uncertainties underestimated.

NODE=M001W;LINKAGE=G

<sup>7</sup> Relativistic Breit-Wigner includes radiative corrections. Systematic uncertainties not estimated.

NODE=M001W;LINKAGE=J

<sup>8</sup> Systematic uncertainties not estimated.

NODE=M001W;LINKAGE=B

<sup>9</sup> Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

$\omega(782)$  DECAY MODES

NODE=M001215;NODE=M001

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level	
$\Gamma_1$ $\pi^+\pi^-\pi^0$	(89.2 $\pm$ 0.7) %		DESIG=1
$\Gamma_2$ $\pi^0\gamma$	( 8.35 $\pm$ 0.27) %	S=2.2	DESIG=3
$\Gamma_3$ $\pi^+\pi^-$	( 1.53 $^{+0.11}_{-0.13}$ ) %	S=1.2	DESIG=2
$\Gamma_4$ neutrals (excluding $\pi^0\gamma$ )	( 7 $^{+8}_{-4}$ ) $\times 10^{-3}$	S=1.1	DESIG=13
$\Gamma_5$ $\eta\gamma$	( 4.5 $\pm$ 0.4 ) $\times 10^{-4}$	S=1.1	DESIG=6
$\Gamma_6$ $\pi^0e^+e^-$	( 7.7 $\pm$ 0.6 ) $\times 10^{-4}$		DESIG=14
$\Gamma_7$ $\pi^0\mu^+\mu^-$	( 1.34 $\pm$ 0.18) $\times 10^{-4}$	S=1.5	DESIG=11
$\Gamma_8$ $\eta e^+e^-$			DESIG=18
$\Gamma_9$ $e^+e^-$	( 7.38 $\pm$ 0.22) $\times 10^{-5}$	S=1.9	DESIG=7
$\Gamma_{10}$ $\pi^+\pi^-\pi^0\pi^0$	< 2 $\times 10^{-4}$	CL=90%	DESIG=12
$\Gamma_{11}$ $\pi^+\pi^-\gamma$	< 3.6 $\times 10^{-3}$	CL=95%	DESIG=4
$\Gamma_{12}$ $\pi^+\pi^-\pi^+\pi^-$	< 1 $\times 10^{-3}$	CL=90%	DESIG=15
$\Gamma_{13}$ $\pi^0\pi^0\gamma$	( 6.7 $\pm$ 1.1 ) $\times 10^{-5}$		DESIG=5
$\Gamma_{14}$ $\eta\pi^0\gamma$	< 3.3 $\times 10^{-5}$	CL=90%	DESIG=17
$\Gamma_{15}$ $\mu^+\mu^-$	( 7.4 $\pm$ 1.8 ) $\times 10^{-5}$		DESIG=8
$\Gamma_{16}$ $3\gamma$	< 1.9 $\times 10^{-4}$	CL=95%	DESIG=10

## Charge conjugation (C) violating modes

$\Gamma_{17}$ $\eta\pi^0$	C	< 2.1 $\times 10^{-4}$	CL=90%	NODE=M001;CLUMP=A DESIG=9
$\Gamma_{18}$ $2\pi^0$	C	< 2.2 $\times 10^{-4}$	CL=90%	DESIG=193
$\Gamma_{19}$ $3\pi^0$	C	< 2.3 $\times 10^{-4}$	CL=90%	DESIG=16
$\Gamma_{20}$ invisible		< 7 $\times 10^{-5}$	CL=90%	DESIG=194

## CONSTRAINED FIT INFORMATION

An overall fit to 15 branching ratios uses 48 measurements and one constraint to determine 10 parameters. The overall fit has a  $\chi^2 = 48.0$  for 39 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	23								
$x_3$	-18	-4							
$x_4$	-92	-55	1						
$x_5$	7	23	-1	-15					
$x_6$	-1	0	0	0	0				
$x_7$	0	0	0	0	0	0			
$x_9$	-24	-73	4	47	-31	0	0		
$x_{13}$	1	4	0	-2	1	0	0	-3	
$x_{15}$	0	0	0	0	0	0	0	0	0
	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$	$x_7$	$x_9$	$x_{13}$

 $\omega(782)$  PARTIAL WIDTHS

NODE=M001218

 $\Gamma(\pi^0\gamma)$  $\Gamma_2$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

880 $\pm$ 50	7815	<sup>1</sup> ACHASOV	13	SND	1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
788 $\pm$ 12 $\pm$ 27	36500	<sup>2</sup> ACHASOV	03	SND	0.60-0.97 $e^+e^- \rightarrow \pi^0\gamma$
764 $\pm$ 51	10625	DOLINSKY	89	ND	$e^+e^- \rightarrow \pi^0\gamma$

<sup>1</sup>Systematic uncertainty not estimated.

<sup>2</sup>Using  $\Gamma_\omega = 8.44 \pm 0.09$  MeV and  $B(\omega \rightarrow \pi^0\gamma)$  from ACHASOV 03.

NODE=M001W1  
NODE=M001W1NODE=M001W1;LINKAGE=AC  
NODE=M001W1;LINKAGE=AD



$\Gamma(\eta\gamma)$  $\Gamma_5$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1±2.5	<sup>1</sup> DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$
---------	--------------------------	----	---------------------------------

<sup>1</sup> Using  $\Gamma_\omega = 8.4 \pm 0.1$  MeV and  $B(\omega \rightarrow \eta\gamma)$  from DOLINSKY 89.

NODE=M001W2  
NODE=M001W2

NODE=M001W2;LINKAGE=DA

 $\Gamma(e^+e^-)$  $\Gamma_9$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.60 ± 0.02 OUR EVALUATION**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.591±0.015	11200	<sup>1,2</sup> AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-------------	-------	------------------------------	------	--------------------------------------

0.653±0.003±0.021	1.2M	<sup>3</sup> ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow$
-------------------	------	--------------------------	------	--------------------------------

0.600±0.031	10625	DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-------------	-------	-------------	----	--

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.891 \pm 0.007$  and  $\Gamma_{\text{total}} = 8.44 \pm 0.09$  MeV.

<sup>2</sup> Update of AKHMETSHIN 00C.

<sup>3</sup> Using ACHASOV 03, ACHASOV 03D and  $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$ .

NODE=M001W7  
NODE=M001W7

→ UNCHECKED ←

NODE=M001W7;LINKAGE=3P  
NODE=M001W7;LINKAGE=PT  
NODE=M001W;LINKAGE=VF

 $\omega(782) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M001235

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_9/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

<b>569.8±3.1±8.2</b>	<sup>1</sup> LEES 21B	BABR	10.5 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
----------------------	-----------------------	------	---

<sup>1</sup> From the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$  with contributions from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .

NODE=M001R02  
NODE=M001R02

NODE=M001R02;LINKAGE=A

 $\omega(782) \Gamma(e^+e^-)\Gamma(i)/\Gamma^2(\text{total})$ 

NODE=M001225

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma \times \Gamma_1/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**6.59±0.19 OUR FIT** Error includes scale factor of 2.1.

**6.36±0.14 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

6.24±0.11±0.08	11.2k	<sup>1</sup> AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
----------------	-------	----------------------------	------	--------------------------------------

6.74±0.04±0.24	1.2M	<sup>2,3</sup> ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow$
----------------	------	----------------------------	------	--------------------------------

6.37±0.35		<sup>2</sup> DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-----------	--	--------------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.20±0.13		<sup>4</sup> BENAYOUN 10	RVUE	0.4–1.05 $e^+e^-$
-----------	--	--------------------------	------	-------------------

6.70±0.06±0.27		<sup>5</sup> AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow$
----------------	--	---------------------------	------	---------------------------

6.45±0.24		<sup>6</sup> BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	--	------------------------	-----	--------------------------------------

5.79±0.42	1488	<sup>7</sup> KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	------	---------------------------	------	--------------------------------------

5.89±0.54	433	<sup>6</sup> CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	-----	-------------------------	-----	--------------------------------------

7.54±0.84	451	<sup>6</sup> BENAJSAS 72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	-----	---------------------------	------	--------------------------------------

<sup>1</sup> Update of AKHMETSHIN 00C.

<sup>2</sup> Recalculated by us from the cross section in the peak.

<sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

<sup>4</sup> A simultaneous fit of  $e^+e^- \rightarrow \pi^+\pi^-$ ,  $\pi^+\pi^-\pi^0$ ,  $\pi^0\gamma$ ,  $\eta\gamma$  data.

<sup>5</sup> Superseded by LEES 21B.

<sup>6</sup> Recalculated by us from the cross section in the peak. Systematic uncertainties underestimated.


<sup>7</sup> Recalculated by us from the cross section in the peak. Systematic uncertainties not estimated.

NODE=M001G2  
NODE=M001G2

NODE=M001G;LINKAGE=PT  
NODE=M001G;LINKAGE=LP  
NODE=M001G;LINKAGE=VH

NODE=M001G2;LINKAGE=BE  
NODE=M001G2;LINKAGE=C  
NODE=M001G2;LINKAGE=A

NODE=M001G2;LINKAGE=B



**$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\pi^0\gamma)/\Gamma_{total}$   $\Gamma_9/\Gamma \times \Gamma_2/\Gamma$**

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.16 ±0.14 OUR FIT</b>				Error includes scale factor of 1.8.
<b>6.34 ±0.10 OUR AVERAGE</b>				
6.336±0.056±0.089		<sup>1</sup> ACHASOV 16A SND		0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
6.47 ±0.14 ±0.39	18k	AKHMETSHIN 05 CMD2		0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
6.34 ±0.21 ±0.21	10k	<sup>2</sup> DOLINSKY 89 ND		$e^+e^- \rightarrow \pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
6.80 ±0.13		<sup>3</sup> BENAYOUN 10 RVUE		0.4–1.05 $e^+e^-$
6.50 ±0.11 ±0.20	36k	<sup>4</sup> ACHASOV 03 SND		0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$

NODE=M001G4  
NODE=M001G4

- <sup>1</sup> From the VMD model with the interfering  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ , and an additional resonance describing the total contribution of the  $\rho(1450)$  and  $\omega(1420)$  states. Supersedes ACHASOV 03.
- <sup>2</sup> Recalculated by us from the cross section in the peak.
- <sup>3</sup> A simultaneous fit of  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$  data.
- <sup>4</sup> Using  $\sigma(\phi \rightarrow \pi^0\gamma)$  from ACHASOV 00 and  $m_\omega = 782.57$  MeV in the model with the energy-independent phase of  $\rho$ - $\omega$  interference equal to  $(-10.2 \pm 7.0)^\circ$ .

NODE=M001G4;LINKAGE=A

NODE=M001G4;LINKAGE=LP  
NODE=M001G4;LINKAGE=BE  
NODE=M001G;LINKAGE=SH

**$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\pi^+\pi^-)/\Gamma_{total}$   $\Gamma_9/\Gamma \times \Gamma_3/\Gamma$**

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.28 ±0.05 OUR AVERAGE</b>				
1.318±0.051±0.021		<sup>1</sup> ACHASOV 21 SND		$e^+e^- \rightarrow \pi^+\pi^-$
1.225±0.058±0.041	800k	<sup>2</sup> ACHASOV 06 SND		$e^+e^- \rightarrow \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.166±0.036		<sup>3</sup> BENAYOUN 13 RVUE		0.4–1.05 $e^+e^-$
1.05 ±0.08		<sup>4</sup> DAVIER 13 RVUE		$e^+e^- \rightarrow \pi^+\pi^-(\gamma)$

NODE=M001G5  
NODE=M001G5

- <sup>1</sup> From a fit of the cross section in the energy range  $0.525 < \sqrt{s} < 0.883$  GeV parameterized by the sum of the Breit-Wigner amplitudes for the  $\rho(770)$ ,  $\omega$  and  $\rho(1450)$  resonances. The measured phase of the  $\rho(770)$ — $\omega$  interference is  $(110.7 \pm 1.5 \pm 1.0)^\circ$ .
- <sup>2</sup> Supersedes ACHASOV 05A.
- <sup>3</sup> A simultaneous fit to  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$ , and  $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$  data. Supersedes BENAYOUN 10.
- <sup>4</sup> From  $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$  data of LEES 12G.

NODE=M001G5;LINKAGE=C

NODE=M001G5;LINKAGE=AC  
NODE=M001G5;LINKAGE=B

NODE=M001G5;LINKAGE=A

**$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\eta\gamma)/\Gamma_{total}$   $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$**

VALUE (units $10^{-8}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.32±0.28 OUR FIT</b>				Error includes scale factor of 1.1.
<b>3.18±0.28 OUR AVERAGE</b>				
3.10±0.31±0.11	33k	<sup>1</sup> ACHASOV 07B SND		0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
3.17 <sup>+1.85</sup> <sub>-1.31</sub> ±0.21	17.4k	<sup>2</sup> AKHMETSHIN 05 CMD2		0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
3.41±0.52±0.21	23k	<sup>3,4</sup> AKHMETSHIN 01B CMD2		$e^+e^- \rightarrow \eta\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.50±0.10		<sup>5</sup> BENAYOUN 10 RVUE		0.4–1.05 $e^+e^-$

NODE=M001G3  
NODE=M001G3

- <sup>1</sup> From a combined fit of  $\sigma(e^+e^- \rightarrow \eta\gamma)$  with  $\eta \rightarrow 3\pi^0$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$ , and fixing  $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$ . Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- <sup>2</sup> From the  $\eta \rightarrow 2\gamma$  decay and using  $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$ .
- <sup>3</sup> From the  $\eta \rightarrow 3\pi^0$  decay and using  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ .
- <sup>4</sup> The combined fit from 600 to 1380 MeV taking into account  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ , and  $\rho(1450)$  (mass and width fixed at 1450 MeV and 310 MeV respectively).
- <sup>5</sup> A simultaneous fit of  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$  data.

NODE=M001G3;LINKAGE=AH

NODE=M001G;LINKAGE=AH

NODE=M001G;LINKAGE=AK

NODE=M001G;LINKAGE=BQ

NODE=M001G3;LINKAGE=BE

NODE=M001G01

NODE=M001G01

NODE=M001G01;LINKAGE=A

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \Gamma_{15}/\Gamma$

VALUE (units $10^{-9}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.3±1.8±2.2</b>	4.5M	<sup>1</sup> ANASTASI	17	KLOE $e^+e^- \rightarrow \mu^+\mu^-\gamma$

- <sup>1</sup> From a fit of the real part of the vacuum polarization by a sum of the leptonic and hadronic contributions, where the hadronic contribution is parametrized as a sum of Breit-Wigner resonances  $\omega(782)$ ,  $\phi(1020)$  and using a GOUNARIS 68 parametrization for the  $\rho(770)$ , and a non-resonant term.

NODE=M001220

## $\omega(782)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the  $\pi\pi$   $P$ -wave scattering phase shift.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9024±0.0019		<sup>1</sup> AMBROSINO 08G	KLOE	1.0–1.03 $e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
0.8965±0.0016±0.0048	1.2M	<sup>2,3</sup> ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.880 ±0.020 ±0.032	11200	<sup>3,4</sup> AKHMETSHIN 00C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.8942±0.0062		<sup>3</sup> DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

- <sup>1</sup> Not independent of  $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$  from AMBROSINO 08G.

- <sup>2</sup> Using ACHASOV 03, ACHASOV 03D and  $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$ .

- <sup>3</sup> Not independent of the corresponding  $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$ .

- <sup>4</sup> Using  $\Gamma(e^+e^-) = 0.60 \pm 0.02$  keV.

NODE=M001R21

NODE=M001R21

NODE=M001R21

NODE=M001R21;LINKAGE=AM

NODE=M001R;LINKAGE=VF

NODE=M001R;LINKAGE=ZL

NODE=M001R;LINKAGE=KH

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.88±0.18		<sup>1</sup> ACHASOV 16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
8.09±0.14		<sup>2</sup> AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
9.06±0.20±0.57	18k	<sup>3,4</sup> AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
9.34±0.15±0.31	36k	<sup>4</sup> ACHASOV 03	SND	0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
8.65±0.16±0.42	1.2M	<sup>5,6</sup> ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.39±0.24	9k	<sup>7</sup> BENAYOUN 96	RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.88±0.62	10k	<sup>4</sup> DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

- <sup>1</sup> Using  $B(\omega \rightarrow e^+e^-)$  from PDG 15. Supersedes ACHASOV 03.

- <sup>2</sup> Not independent of  $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$  from AMBROSINO 08G.

- <sup>3</sup> Using  $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$ .

- <sup>4</sup> Not independent of the corresponding  $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$ .

- <sup>5</sup> Using ACHASOV 03, ACHASOV 03D and  $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$ .

- <sup>6</sup> Not independent of the corresponding  $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$ .

- <sup>7</sup> Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001R28

NODE=M001R28

NODE=M001R28;LINKAGE=A

NODE=M001R28;LINKAGE=AM

NODE=M001R;LINKAGE=AH

NODE=M001R;LINKAGE=VL

NODE=M001R28;LINKAGE=VF

NODE=M001R28;LINKAGE=ZL

NODE=M001R28;LINKAGE=A1

$\Gamma(\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_2/\Gamma_1$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**9.35±0.30 OUR FIT** Error includes scale factor of 2.4.

**9.05±0.27 OUR AVERAGE** Error includes scale factor of 1.8.

8.97±0.16	AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
9.94±0.36±0.38	<sup>1</sup> AULCHENKO 00A	SND	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
8.4 ±1.3	KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
10.9 ±2.5	BENAKSAS 72C	OSPK	$e^+e^- \rightarrow \pi^0\gamma$
8.1 ±2.0	BALDIN 71	HLBC	$2.9 \pi^+p$
13 ±4	JACQUET 69B	HLBC	$2.05 \pi^+p \rightarrow \pi^+p\omega$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.7 ±0.2 ±0.5	<sup>2,3</sup> ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.9 ±0.7	<sup>2</sup> DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

NODE=M001R3

NODE=M001R3

<sup>1</sup> From  $\sigma_0^{\omega\pi^0 \rightarrow \pi^0\pi^0\gamma}(m_\phi)/\sigma_0^{\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0\pi^0}(m_\phi)$  with a phase-space correction factor of 1/1.023.

<sup>2</sup> Not independent of the corresponding  $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$ .

<sup>3</sup> Using ACHASOV 03. Based on 1.2M events.

NODE=M001R3;LINKAGE=AL

NODE=M001R3;LINKAGE=VL  
NODE=M001R3;LINKAGE=VW

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_3/\Gamma$

See also  $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ .

NODE=M001R15

NODE=M001R15  
NODE=M001R15

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.53<sup>+0.11</sup><sub>-0.13</sub> OUR FIT** Error includes scale factor of 1.2.

**1.49 $\pm$ 0.13 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

1.46 $\pm$ 0.12 $\pm$ 0.02	900k	<sup>1</sup> AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
1.30 $\pm$ 0.24 $\pm$ 0.05	11.2k	<sup>2</sup> AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
2.38 <sup>+1.77</sup> <sub>-0.90</sub> $\pm$ 0.18	5.4k	<sup>3</sup> ACHASOV	02E	SND 1.1-1.38 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
2.3 $\pm$ 0.5		BARKOV	85	OLYA $e^+e^- \rightarrow \pi^+\pi^-$
1.6 <sup>+0.9</sup> <sub>-0.7</sub>		QUENZER	78	DM1 $e^+e^- \rightarrow \pi^+\pi^-$
3.6 $\pm$ 1.9		BENAKSAS	72	OSPK $e^+e^- \rightarrow \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.29 $\pm$ 0.22 $\pm$ 0.03	970k	<sup>4,5</sup> ABLIKIM	18C	BES3 $\eta'(958) \rightarrow \gamma\pi^+\pi^-$
1.28 $\pm$ 0.22 $\pm$ 0.03	970k	<sup>6,7</sup> ABLIKIM	18C	BES3 $\eta'(958) \rightarrow \gamma\pi^+\pi^-$
1.52 $\pm$ 0.08		<sup>8</sup> HANHART	18	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
1.75 $\pm$ 0.11	4.5M	<sup>9</sup> ACHASOV	05A	SND $e^+e^- \rightarrow \pi^+\pi^-$
2.01 $\pm$ 0.29		<sup>10</sup> BENAYOUN	03	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
1.9 $\pm$ 0.3		<sup>11</sup> GARDNER	99	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
2.3 $\pm$ 0.4		<sup>12</sup> BENAYOUN	98	RVUE $e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
1.0 $\pm$ 0.11		<sup>13</sup> WICKLUND	78	ASPK 3,4,6 $\pi^\pm N$
1.22 $\pm$ 0.30		ALVENSLEB...	71C	CNTR Photoproduction
1.3 <sup>+1.2</sup> <sub>-0.9</sub>		MOFFEIT	71	HBC 2.8,4.7 $\gamma\rho$
0.80 <sup>+0.28</sup> <sub>-0.20</sub>		<sup>14</sup> BIGGS	70B	CNTR 4.2 $\gamma C \rightarrow \pi^+\pi^-C$

OCCUR=3  
OCCUR=2

<sup>1</sup> A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

<sup>2</sup> Update of AKHMETSHIN 02.

<sup>3</sup> From the  $m_{\pi^+\pi^-}$  spectrum taking into account the interference of the  $\rho\pi$  and  $\omega\pi$  amplitudes.

<sup>4</sup> From a fit to  $\pi^+\pi^-$  mass using  $\rho(770)$  (parametrized with the Gounaris-Sakurai approach),  $\omega(782)$ , and box anomaly components.

<sup>5</sup> ABLIKIM 18C reports  $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.25 \pm 0.21 \pm 0.52) \times 10^{-4}$  which we divide by our best value  $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>6</sup> From a fit to  $\pi^+\pi^-$  mass using  $\rho(770)$  (parametrized with the Gounaris-Sakurai approach),  $\omega(782)$ , and  $\rho(1450)$  components.

<sup>7</sup> ABLIKIM 18C reports  $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.22 \pm 0.21 \pm 0.52) \times 10^{-4}$  which we divide by our best value  $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>8</sup> Dispersive analysis. Value extracted from average of data from AUBERT 09AS, AKHMETSHIN 07, ACHASOV 06, AMBROSINO 11A, BABUSCI 13D, ABLIKIM 16B normalised by PDG 16 evaluation for  $\Gamma(\omega \rightarrow e^+e^-)$ .

<sup>9</sup> Using  $\Gamma(\omega \rightarrow e^+e^-)$  from the 2004 Edition of this Review (PDG 04).

<sup>10</sup> Using the data of AKHMETSHIN 02 in the hidden local symmetry model.

<sup>11</sup> Using the data of BARKOV 85.

<sup>12</sup> Using the data of BARKOV 85 in the hidden local symmetry model.

<sup>13</sup> From a model-dependent analysis assuming complete coherence.

<sup>14</sup> Re-evaluated under  $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$  by BEHREND 71 using more accurate  $\omega \rightarrow \rho$  photoproduction cross-section ratio.

NODE=M001R15;LINKAGE=AK  
NODE=M001R15;LINKAGE=PT  
NODE=M001R;LINKAGE=VE

NODE=M001R15;LINKAGE=E


NODE=M001R15;LINKAGE=H

NODE=M001R15;LINKAGE=J

NODE=M001R15;LINKAGE=K

NODE=M001R15;LINKAGE=D

NODE=M001R;LINKAGE=SN  
NODE=M001R;LINKAGE=BY  
NODE=M001R15;LINKAGE=H4  
NODE=M001R15;LINKAGE=Q  
NODE=M001R15;LINKAGE=F  
NODE=M001R15;LINKAGE=B



$\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$

See also  $\Gamma(\pi^+\pi^-)/\Gamma_{total}$ .

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0172±0.0014 OUR FIT</b>			Error includes scale factor of 1.2.
<b>0.026 ±0.005 OUR AVERAGE</b>			

0.021 +0.028 -0.009	1,2	RATCLIFF	72	ASPK	15 $\pi^- p \rightarrow n2\pi$
0.028 ±0.006	1	BEHREND	71	ASPK	Photoproduction
0.022 +0.009 -0.01	3	ROOS	70	RVUE	

- <sup>1</sup> The fitted width of these data is 160 MeV in agreement with present average, thus the  $\omega$  contribution is overestimated. Assuming  $\rho$  width 145 MeV.  
<sup>2</sup> Significant interference effect observed. NB of  $\omega \rightarrow 3\pi$  comes from an extrapolation.  
<sup>3</sup> ROOS 70 combines ABRAMOVICH 70 and BIZZARRI 70.

NODE=M001R2  
 NODE=M001R2  
 NODE=M001R2

NODE=M001R2;LINKAGE=A  
 NODE=M001R2;LINKAGE=S  
 NODE=M001R2;LINKAGE=R

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\gamma)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.20±0.04</b>	1.98M	<sup>1</sup> ALOISIO	03	KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

- <sup>1</sup> Using the data of ALOISIO 02D.

NODE=M001R33  
 NODE=M001R33

NODE=M001R;LINKAGE=KL

$\Gamma(\text{neutrals})/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.091±0.006 OUR FIT</b>				
<b>0.081±0.011 OUR AVERAGE</b>				

0.075±0.025		BIZZARRI	71	HBC	0.0 $p\bar{p}$
0.079±0.019		DEINET	69B	OSPK	1.5 $\pi^- p$
0.084±0.015		BOLLINI	68C	CNTR	2.1 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.073±0.018	42	BASILE	72B	CNTR	1.67 $\pi^- p$

NODE=M001R14  
 NODE=M001R14

$\Gamma(\text{neutrals})/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.102±0.008 OUR FIT</b>				
<b>0.103<sup>+0.011</sup><sub>-0.010</sub> OUR AVERAGE</b>				

0.15 ±0.04	46	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
0.10 ±0.03	19	BARASH	67B	HBC	0.0 $\bar{p}p$
0.134±0.026	850	DIGIUGNO	66B	CNTR	1.4 $\pi^- p$
0.097±0.016	348	FLATTE	66	HBC	1.4 - 1.7 $K^- p \rightarrow \Lambda MM$
0.06 +0.05 -0.02		JAMES	66	HBC	2.1 $\pi^+ p$
0.08 ±0.03	35	KRAEMER	64	DBC	1.2 $\pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ±0.02	20	BUSCHBECK	63	HBC	1.5 $K^- p$

NODE=M001R1  
 NODE=M001R1

$\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$  $\Gamma_2/(\Gamma_2+\Gamma_4)$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M001R18  
 NODE=M001R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.78 \pm 0.07$		<sup>1</sup> DAKIN	72	OSPK	$1.4 \pi^- p \rightarrow nMM$
$>0.81$	90	DEINET	69B	OSPK	

<sup>1</sup> Error statistical only. Authors obtain good fit also assuming  $\pi^0\gamma$  as the only neutral decay.

NODE=M001R18;LINKAGE=D

 $\Gamma(\text{neutrals})/\Gamma(\text{charged particles})$  $(\Gamma_2+\Gamma_4)/(\Gamma_1+\Gamma_3)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M001R9  
 NODE=M001R9

**0.100 ± 0.008 OUR FIT**

<b>0.124 ± 0.021</b>	FELDMAN	67C	OSPK	$1.2 \pi^- p$
----------------------	---------	-----	------	---------------

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M001R19  
 NODE=M001R19

**4.5 ± 0.4 OUR FIT** Error includes scale factor of 1.1.**6.3 ± 1.3 OUR AVERAGE** Error includes scale factor of 1.2.

$6.6 \pm 1.7$		<sup>1</sup> ABELE	97E	CBAR	$0.0 \bar{p} p \rightarrow 5\gamma$
$8.3 \pm 2.1$		ALDE	93	GAM2	$38\pi^- p \rightarrow \omega n$
$3.0 \begin{smallmatrix} +2.5 \\ -1.8 \end{smallmatrix}$		<sup>2</sup> ANDREWS	77	CNTR	$6.7-10 \gamma Cu$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.2 \pm 0.4 \pm 0.1$	33k	<sup>3</sup> ACHASOV	07B	SND	$0.6-1.38 e^+ e^- \rightarrow \eta\gamma$
$4.44 \begin{smallmatrix} +2.59 \\ -1.83 \end{smallmatrix} \pm 0.28$	17.4k	<sup>4,5</sup> AKHMETSHIN	05	CMD2	$0.60-1.38 e^+ e^- \rightarrow \eta\gamma$
$5.10 \pm 0.72 \pm 0.34$	23k	<sup>6</sup> AKHMETSHIN	01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
$0.7 \text{ to } 5.5$		<sup>7</sup> CASE	00	CBAR	$0.0 p\bar{p} \rightarrow \eta\eta\gamma$
$6.56 \begin{smallmatrix} +2.41 \\ -2.55 \end{smallmatrix}$	3525	<sup>2,8</sup> BENAYOUN	96	RVUE	$e^+ e^- \rightarrow \eta\gamma$
$7.3 \pm 2.9$		<sup>2,4</sup> DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$

<sup>1</sup> No flat  $\eta\eta\gamma$  background assumed.

<sup>2</sup> Solution corresponding to constructive  $\omega$ - $\rho$  interference.

<sup>3</sup> ACHASOV 07B reports  $[\Gamma(\omega(782) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow e^+ e^-)] = (3.10 \pm 0.31 \pm 0.11) \times 10^{-8}$  which we divide by our best value  $B(\omega(782) \rightarrow e^+ e^-) = (7.38 \pm 0.22) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

<sup>4</sup> Not independent of the corresponding  $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$ .

<sup>5</sup> Using  $B(\omega \rightarrow e^+ e^-) = (7.14 \pm 0.13) \times 10^{-5}$  and  $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$ .

<sup>6</sup> Using  $B(\omega \rightarrow e^+ e^-) = (7.07 \pm 0.19) \times 10^{-5}$  and using  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ . Solution corresponding to constructive  $\omega$ - $\rho$  interference. The combined fit from 600 to 1380 MeV taking into account  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ , and  $\rho(1450)$  (mass and width fixed at 1450 MeV and 310 MeV respectively). Not independent of the corresponding  $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$ .

<sup>7</sup> Depending on the degree of coherence with the flat  $\eta\eta\gamma$  background and using  $B(\omega \rightarrow \pi^0\gamma) = (8.5 \pm 0.5) \times 10^{-2}$ .

<sup>8</sup> Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001R;LINKAGE=EA  
 NODE=M001R19;LINKAGE=A  
 NODE=M001R19;LINKAGE=AO

NODE=M001R13;LINKAGE=WL  
 NODE=M001R19;LINKAGE=AK  
 NODE=M001R19;LINKAGE=TS

NODE=M001R;LINKAGE=CS

NODE=M001R19;LINKAGE=A1

 $\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$  $\Gamma_5/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M001R11  
 NODE=M001R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0098 \pm 0.0024$		<sup>1</sup> ALDE	93	GAM2	$38\pi^- p \rightarrow \omega n$
$0.0082 \pm 0.0033$		<sup>2</sup> DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$
$0.010 \pm 0.045$		APEL	72B	OSPK	$4-8 \pi^- p \rightarrow n3\gamma$

<sup>1</sup> Model independent determination.

<sup>2</sup> Solution corresponding to constructive  $\omega$ - $\rho$  interference.

NODE=M001R11;LINKAGE=A  
 NODE=M001R11;LINKAGE=K

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M001R23  
 NODE=M001R23


**7.7 ± 0.6 OUR FIT****7.7 ± 0.6 OUR AVERAGE**

$7.61 \pm 0.53 \pm 0.64$		ACHASOV	08	SND	$0.36-0.97 e^+ e^- \rightarrow \pi^0 e^+ e^-$
$8.19 \pm 0.71 \pm 0.62$		AKHMETSHIN	05A	CMD2	$0.72-0.84 e^+ e^-$
$5.9 \pm 1.9$	43	DOLINSKY	88	ND	$e^+ e^- \rightarrow \pi^0 e^+ e^-$

$\Gamma(\pi^0 \mu^+ \mu^-) / \Gamma_{\text{total}}$  $\Gamma_7 / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.34 ± 0.18 OUR FIT</b>				Error includes scale factor of 1.5.
<b>1.34 ± 0.19 OUR AVERAGE</b>				Error includes scale factor of 1.5. See the ideogram below.
1.41 ± 0.09 ± 0.15		ARNALDI	16	NA60 400 GeV ( $p$ -A) collisions
1.72 ± 0.25 ± 0.14	3k	ARNALDI	09	NA60 158A In-In collisions
0.96 ± 0.23		DZHELADIN	81B	CNTR 25-33 $\pi^- p \rightarrow \omega n$

NODE=M001R12  
 NODE=M001R12

 $\Gamma(\eta e^+ e^-) / \Gamma_{\text{total}}$  $\Gamma_8 / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
< 1.1	AKHMETSHIN 05A	CMD2	0.72-0.84 $e^+ e^-$

NODE=M001R34  
 NODE=M001R34

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}}$  $\Gamma_9 / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.738 ± 0.022 OUR FIT</b>				Error includes scale factor of 1.9.
0.700 ± 0.016	11200	1,2 AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.752 ± 0.004 ± 0.024	1.2M	2,3 ACHASOV 03D	RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.714 ± 0.036		2 DOLINSKY 89	ND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.72 ± 0.03		2 BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.64 ± 0.04	1488	2 KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.675 ± 0.069	433	2 CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.83 ± 0.10	451	2 BENAKSAS 72B	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.77 ± 0.06		4 AUGUSTIN 69D	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.65 ± 0.13	33	5 ASTVACAT...	68	OSPK Assume SU(3)+mixing

NODE=M001R13  
 NODE=M001R13

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = 0.891 \pm 0.007$ . Update of AKHMETSHIN 00C.

<sup>2</sup> Not independent of the corresponding  $\Gamma(e^+ e^-) \times \Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}^2$ .

<sup>3</sup> Using ACHASOV 03, ACHASOV 03D and  $B(\omega \rightarrow \pi^+ \pi^-) = (1.70 \pm 0.28)\%$ .

<sup>4</sup> Rescaled by us to correspond to  $\omega$  width 8.4 MeV. Systematic errors underestimated.

<sup>5</sup> Not resolved from  $\rho$  decay. Error statistical only.

NODE=M001R13;LINKAGE=4P  
 NODE=M001R13;LINKAGE=ZL  
 NODE=M001R13;LINKAGE=VF  
 NODE=M001R13;LINKAGE=E  
 NODE=M001R13;LINKAGE=A

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}$  $\Gamma_{10} / \Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 2	90	ACHASOV 09A	SND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
< 200	90	KURDADZE 86	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

NODE=M001R5  
 NODE=M001R5

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0036</b>	95	WEIDENAUER 90	ASTE	$\rho\bar{p} \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.004	95	BITYUKOV 88B	SPEC	$32 \pi^- p \rightarrow \pi^+\pi^-\gamma X$

NODE=M001R22  
NODE=M001R22

$$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{11}/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.066	90	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$
<0.05	90	FLATTE 66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$

NODE=M001R4  
NODE=M001R4

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1 x 10<sup>-3</sup></b>	90	KURDADZE 88	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M001R24  
NODE=M001R24

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (units 10 <sup>-5</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.7 ± 1.1 OUR FIT</b>				
<b>6.5 ± 1.2 OUR AVERAGE</b>				
$6.4^{+2.4}_{-2.0} \pm 0.8$	190	<sup>1</sup> AKHMETSHIN 04B	CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$
$6.6^{+1.4}_{-1.3} \pm 0.6$	295	ACHASOV 02F	SND	$0.36-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$

NODE=M001R29  
NODE=M001R29

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$11.8^{+2.1}_{-1.9} \pm 1.4$	190	<sup>2</sup> AKHMETSHIN 04B	CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2
$7.8 \pm 2.7 \pm 2.0$	63	<sup>1,3</sup> ACHASOV 00G	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
$12.7 \pm 2.3 \pm 2.5$	63	<sup>2,3</sup> ACHASOV 00G	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2

<sup>1</sup> In the model assuming the  $\rho \rightarrow \pi^0\pi^0\gamma$  decay via the  $\omega\pi$  and  $f_0(500)\gamma$  mechanisms.

<sup>2</sup> In the model assuming the  $\rho \rightarrow \pi^0\pi^0\gamma$  decay via the  $\omega\pi$  mechanism only.

<sup>3</sup> Superseded by ACHASOV 02F.

NODE=M001R29;LINKAGE=A  
NODE=M001R29;LINKAGE=B  
NODE=M001R;LINKAGE=GF

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_{13}/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.00045</b>	90	DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.08	95	JACQUET 69B	HLBC	$2.05 \pi^+ p \rightarrow \pi^+ p \omega$

NODE=M001R10  
NODE=M001R10

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^0\gamma) \quad \Gamma_{13}/\Gamma_2$$

VALUE (units 10 <sup>-4</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.0 ± 1.3 OUR FIT</b>					
<b>8.5 ± 2.9</b>					
	40 ± 14		ALDE 94B	GAM2	$38\pi^- p \rightarrow \pi^0\pi^0\gamma n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 50	90		DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
<1800	95		KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
<1500	90		BENAKSAS 72C	OSPK	$e^+e^-$
<1400			BALDIN 71	HLBC	$2.9 \pi^+ p$
<1000	90		BARMIN 64	HLBC	$1.3-2.8 \pi^- p$

NODE=M001R7  
NODE=M001R7

$$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\text{neutrals}) \quad \Gamma_{13}/(\Gamma_2+\Gamma_4)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.22 \pm 0.07$		<sup>1</sup> DAKIN 72	OSPK	$1.4 \pi^- p \rightarrow nMM$
<0.19	90	DEINET 69B	OSPK	

NODE=M001R17  
NODE=M001R17

<sup>1</sup> See  $\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$ .

NODE=M001R17;LINKAGE=D

$$\Gamma(\eta\pi^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (units 10 <sup>-5</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.3</b>	90	AKHMETSHIN 04B	CMD2	$0.6-0.97 e^+e^- \rightarrow \eta\pi^0\gamma$

NODE=M001R32  
NODE=M001R32



$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{15}/\Gamma$ NODE=M001R30  
NODE=M001R30**7.4±1.8 OUR FIT****7.4±1.8 OUR AVERAGE**

6.6±1.4±1.7	4.5M	<sup>1</sup> ANASTASI	17	KLOE	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
9.0±2.9±1.1	18	HEISTER	02C	ALEP	$Z \rightarrow \mu^+ \mu^- + X$

<sup>1</sup> Assuming lepton universality in the decay  $\omega \rightarrow \ell^+ \ell^-$  and correcting for different phase space between electron and muon final states.

NODE=M001R30;LINKAGE=A

 $\Gamma(\mu^+ \mu^-)/\Gamma(\pi^+ \pi^- \pi^0)$ VALUE (units  $10^{-3}$ )

CL%

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{15}/\Gamma_1$ NODE=M001R6  
NODE=M001R6

<0.2	90	WILSON	69	OSPK	$12 \pi^- C \rightarrow Fe$
------	----	--------	----	------	-----------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

<1.7	74	FLATTE	66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \mu^+ \mu^-$
<1.2		BARBARO-...	65	HBC	$2.7 K^- p$

 $\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma(\mu^+ \mu^-)$ 

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

 $\Gamma_7/\Gamma_{15}$ NODE=M001R20  
NODE=M001R20

••• We do not use the following data for averages, fits, limits, etc. •••

1.2±0.6	30	<sup>1</sup> DZHELYADIN	79	CNTR	$25-33 \pi^- p$
---------	----	-------------------------	----	------	-----------------

<sup>1</sup> Superseded by DZHELYADIN 81B result above.

NODE=M001R20;LINKAGE=S

 $\Gamma(3\gamma)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )

CL%

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{16}/\Gamma$ NODE=M001R27  
NODE=M001R27

<1.9	95	<sup>1</sup> ABELE	97E	CBAR	$0.0 \bar{p} p \rightarrow 5\gamma$
------	----	--------------------	-----	------	-------------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

<2	90	<sup>1</sup> PROKOSHKIN	95	GAM2	$38 \pi^- p \rightarrow 3\gamma n$
----	----	-------------------------	----	------	------------------------------------

<sup>1</sup> From direct  $3\gamma$  decay search.

NODE=M001R27;LINKAGE=A

 $\Gamma(\eta \pi^0)/\Gamma_{\text{total}}$ 

Violates C conservation.

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{17}/\Gamma$ NODE=M001R25  
NODE=M001R25  
NODE=M001R25

••• We do not use the following data for averages, fits, limits, etc. •••

<0.001	90	ALDE	94B	GAM2	$38\pi^- p \rightarrow \eta \pi^0 n$
--------	----	------	-----	------	--------------------------------------

 $[\Gamma(\eta \gamma) + \Gamma(\eta \pi^0)]/\Gamma(\pi^+ \pi^- \pi^0)$ 

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 $(\Gamma_5 + \Gamma_{17})/\Gamma_1$ NODE=M001R8  
NODE=M001R8

<0.016	90	<sup>1</sup> FLATTE	66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \pi^+ \pi^- MM$
--------	----	---------------------	----	-----	--

••• We do not use the following data for averages, fits, limits, etc. •••

<0.045	95	JACQUET	69B	HLBC	$2.05 \pi^+ p \rightarrow \pi^+ p \omega$
--------	----	---------	-----	------	---

<sup>1</sup> Restated by us using  $B(\eta \rightarrow \text{charged modes}) = 29.2\%$ .

NODE=M001R8;LINKAGE=A

 $\Gamma(\eta \pi^0)/\Gamma(\pi^0 \gamma)$ 

Violates C conservation.

VALUE (units  $10^{-3}$ )

CL%

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{17}/\Gamma_2$ NODE=M001R35  
NODE=M001R35  
NODE=M001R35

<2.6	90	<sup>1</sup> STAROSTIN	09	CRYM	$\gamma p \rightarrow \eta \pi^0 p$
------	----	------------------------	----	------	-------------------------------------

<sup>1</sup> STAROSTIN 09 reports  $[\Gamma(\omega(782) \rightarrow \eta \pi^0)/\Gamma(\omega(782) \rightarrow \pi^0 \gamma)] \times [B(\eta \rightarrow 2\gamma)] < 1.01 \times 10^{-3}$  which we divide by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M001R35;LINKAGE=ST

 $\Gamma(2\pi^0)/\Gamma(\pi^0 \gamma)$ 

Violates C conservation and Bose-Einstein statistics.

VALUE (units  $10^{-3}$ )

CL%

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{18}/\Gamma_2$ NODE=M001R36  
NODE=M001R36  
NODE=M001R36

<2.59	90	STAROSTIN	09	CRYM	$\gamma p \rightarrow 2\pi^0 p$
-------	----	-----------	----	------	---------------------------------

 $\Gamma(3\pi^0)/\Gamma_{\text{total}}$ 

Violates C conservation.

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{19}/\Gamma$ NODE=M001R26  
NODE=M001R26  
NODE=M001R26

••• We do not use the following data for averages, fits, limits, etc. •••

< $3 \times 10^{-4}$	90	PROKOSHKIN	95	GAM2	$38 \pi^- p \rightarrow 3\pi^0 n$
----------------------	----	------------	----	------	-----------------------------------

$\Gamma(3\pi^0)/\Gamma(\pi^0\gamma)$ Violates  $C$  conservation.

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.72	90	STAROSTIN 09	CRYM	$\gamma p \rightarrow 3\pi^0 p$

 $\Gamma_{19}/\Gamma_2$ 

NODE=M001R37  
 NODE=M001R37  
 NODE=M001R37

 $\Gamma(3\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Violates  $C$  conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	BARBERIS 01	450	$pp \rightarrow p_f 3\pi^0 p_s$

 $\Gamma_{19}/\Gamma_1$ 

NODE=M001R31  
 NODE=M001R31  
 NODE=M001R31

 $\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\pi^0)$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.1 $\times 10^{-5}$	90	ABLIKIM 18S	BES3	$J/\psi \rightarrow \omega\eta \rightarrow \omega\pi^+\pi^-\pi^0$

 $\Gamma_{20}/\Gamma_1$ 

NODE=M001R01  
 NODE=M001R01

**PARAMETER  $\Lambda$  IN  $\omega \rightarrow \pi^0 \ell^+ \ell^-$  DECAY**

In the pole approximation the electromagnetic transition form factor for a resonance of mass  $M$  is given by the expression:

$$|F|^2 = (1 - M^2/\Lambda^2)^{-2},$$

where for the parameter  $\Lambda$  vector dominance predicts  $\Lambda = M_p \approx 0.770$  GeV. The ARNALDI 09 measurement is in obvious conflict with this expectation. Note that for  $\eta \rightarrow \gamma\mu^+\mu^-$  decay ARNALDI 09 and DZHELYADIN 80 obtain the value of  $\Lambda$  consistent with vector dominance.

NODE=M001230

NODE=M001230

**PARAMETER  $\Lambda$  IN  $\omega \rightarrow \pi^0 \mu^+ \mu^-$  DECAY**

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.670  $\pm$  0.006 OUR AVERAGE**

0.6707 $\pm$ 0.0039 $\pm$ 0.0056		<sup>1</sup> ARNALDI	16	NA60 400 GeV ( $p$ -A) collisions
0.668 $\pm$ 0.009 $\pm$ 0.003	3k	<sup>2</sup> ARNALDI	09	NA60 158A In-In collisions

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.65 $\pm$ 0.03		DZHELYADIN 81B	CNTR	25-33 $\pi^- p \rightarrow \omega n$
-----------------	--	----------------	------	--------------------------------------

<sup>1</sup> ARNALDI 16 reports  $\Lambda^{-2}(\omega) = 2.223 \pm 0.026 \pm 0.037$  GeV<sup>-2</sup> which we converted to the quoted  $\Lambda$  value.

<sup>2</sup> ARNALDI 09 reports  $\Lambda^{-2}(\omega) = 2.24 \pm 0.06 \pm 0.02$  GeV<sup>-2</sup> which we converted to the quoted  $\Lambda$  value.

NODE=M001LAM  
 NODE=M001LAM

NODE=M001LAM;LINKAGE=A

NODE=M001LAM;LINKAGE=B

**PARAMETER  $\Lambda$  IN  $\omega \rightarrow \pi^0 e^+ e^-$  DECAY**

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.709  $\pm$  0.037**

1.1k		<sup>1</sup> ADLARSON 17B	A2MM	$\gamma p \rightarrow \omega p$
------	--	---------------------------	------	---------------------------------

<sup>1</sup> ADLARSON 17B reports  $\Lambda^{-2}(\omega\pi^0) = 1.99 \pm 0.21$  GeV<sup>-2</sup> that we converted to the quoted  $\Lambda$  value.

NODE=M001A02  
 NODE=M001A02

NODE=M001A02;LINKAGE=A

**ENERGY DEPENDENCE OF  $\omega \rightarrow \pi^+\pi^-\pi^0$  DALITZ PLOT**

The following experiments fit to one or more of the coefficients  $\alpha$ ,  $\beta$ ,  $\gamma$  for |matrix element|<sup>2</sup>  $\propto P(1 + 2\alpha Z + 2\beta Z^{3/2} \sin(3\phi) + 2\gamma Z^2 + O(Z^{5/2}))$  where  $P$  is the  $P$ -wave phase-space factor and  $Z$ ,  $\phi$  are kinematical variables as defined in ADLARSON 17.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.133  $\pm$  0.008 OUR AVERAGE**

0.1321 $\pm$ 0.0067 $\pm$ 0.0046	260k	<sup>1</sup> ABLIKIM	18AD BES3	$J/\psi \rightarrow \omega\eta$
0.147 $\pm$ 0.036	44k	ADLARSON 17	WASA	$\alpha$ in $pd \rightarrow {}^3\text{He } \omega$ , $pp \rightarrow pp\omega$

<sup>1</sup> Keeping a term linear in  $Z$  only. A fit with the terms proportional to  $Z$  and  $Z^{3/2}$  gives  $\alpha = 0.133 \pm 0.041$  and  $\beta = 0.037 \pm 0.054$ .

NODE=M001A00

NODE=M001A00

NODE=M001A00

NODE=M001A00;LINKAGE=A

**$\omega$ (782) REFERENCES**

NODE=M001

ACHASOV	21	JHEP 2101 113	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61040
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
ANDREEV	20	EPJ C80 1189	V. Andreev <i>et al.</i>	(H1 Collab.)	REFID=60773
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)	REFID=61048
COLANGELO	19	JHEP 1902 006	G. Colangelo, M. Hoferichter, P. Stoffer	(BERN+)	REFID=59585
HOFRICHT... ABLIKIM	19 18AD	JHEP 1908 137 PR D98 112007	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)	REFID=60974
ABLIKIM	18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59498
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
HANHART	18	EPJ C78 450	C. Hanhart <i>et al.</i>	(BESIII Collab.)	REFID=58971
ADLARSON	17	PL B770 418	P. Adlarson <i>et al.</i>	(WASA-at-COSY Collab.)	REFID=59186
ADLARSON	17B	PR C95 035208	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)	REFID=57907
ANASTASI	17	PL B767 485	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)	REFID=58155
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57887
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(BESIII Collab.)	REFID=57126
ARNALDI	16	PL B757 437	R. Arnaldi <i>et al.</i>	(SND Collab.)	REFID=57513
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(NA60 Collab.)	REFID=57220
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=57140
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=56977; ERROR=4
BABUSCI	13D	PL B720 336	D. Babusci <i>et al.</i>	(CATA, CALB, BARI)	REFID=55584
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)	REFID=55337
DAVIER	13	EPJ C73 2597	M. Davier <i>et al.</i>		REFID=55357
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55499
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54299
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=54305
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=16683
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53212
ARNALDI	09	Translated from ZETF 136 442. PL B677 260	R. Arnaldi <i>et al.</i>	(NA60 Collab.)	REFID=53101
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52720
STAROSTIN	09	PR C79 065201	A. Starostin <i>et al.</i>	(Crystal Ball Collab. at MAMI)	REFID=53136
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53001
AMBROSINO	08G	Translated from ZETF 134 80. PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52258
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52573
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51942
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51615
ACHASOV	06A	Translated from ZETF 130 437. PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51113
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51133
ACHASOV	05A	Translated from ZETFP 84 491. JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51513
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51045
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
AKHMETSHIN	04	Translated from ZETFP 82 841. PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49610
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=50184
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49653
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49187
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49577
BENAYOUN	03	EPJ C29 397	M. Benayoun <i>et al.</i>		REFID=49404
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49477
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48816
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48565
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48824
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48564
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48311
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48167
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48324
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00G	Translated from ZETFP 72 411. JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
AKHMETSHIN	00C	Translated from ZETFP 71 519. PL B476 33	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47929
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47423
CASE	00	Translated from ZETF 117 1067. PR D61 032002	T. Case <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47953
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47409
GARDNER	99	PR D59 076002	S. Gardner, H.B. O'Connell		REFID=47391
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)	REFID=46919
ABELE	97E	PL B411 361	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45859
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45755
PROKOSHKIN	95	PD 40 273	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=45753
WURZINGER	95	Translated from DANS 342 610. PR C51 443	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=44616
ALDE	94B	PL B340 122	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=45209
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44100
ALDE	93	PAN 56 1229	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=44091
AMSLER	93B	Translated from YAF 56 137. ZPHY C61 35	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43603
WEIDENAUER	93	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43790
ANTONELLI	92	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43602
DOLINSKY	91	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43585
WEIDENAUER	90	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=43168
DOLINSKY	89	ZPHY C47 353	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=41369
BITYUKOV	88B	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41368
DOLINSKY	88	SJNP 47 800	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=41003
KURDADZE	88	Translated from YAF 47 1258. SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41021
AULCHENKO	87	Translated from YAF 48 442. JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=41022
BARKOV	87	Translated from ZETFP 47 432. PL B186 432	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=41121
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40007
		Translated from ZETFP 46 132.		(NOVO)	REFID=40280

KURDADZE	86	JETPL 43 643 Translated from ZETFP 43 497.	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
KURDADZE	83B	JETPL 36 274 Translated from ZETFP 36 221.	A.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20244
DZHELYADIN	81B	PL 102B 296	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20242
CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
DZHELYADIN	80	PL 94B 548	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10831
ROOS	80	LNC 27 321	M. Roos, A. Pellinen	(HELS)	REFID=20241
BENKHEIRI	79	NP B150 268	P. Benkheiri <i>et al.</i>	(EPOL, CERN, CDEF+)	REFID=20238
DZHELYADIN	79	PL 84B 143	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20239
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
QUENZER	78	PL 76B 512	A. Quenzer <i>et al.</i>	(LALO)	REFID=20123
VANAPEL...	78	NP B133 245	G.W. van Apeldoorn <i>et al.</i>	(ZEEM)	REFID=20234
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
KEYNE	76	PR D14 28	J. Keyne <i>et al.</i>	(LOIC, SHMP)	REFID=20226
Also		PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72B	PL 41B 234	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20206
BASILE	72B	Phil. Conf. 153	M. Basile <i>et al.</i>	(CERN)	REFID=20207
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20096
BENAKSAS	72B	PL 42B 507	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20209
BENAKSAS	72C	PL 42B 511	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20517
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
BROWN	72	PL 42B 117	R.M. Brown <i>et al.</i>	(ILL, ILLC)	REFID=20211
DAKIN	72	PR D6 2321	J.T. Dakin <i>et al.</i>	(PRIN)	REFID=20212
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)	REFID=20102
ALVENSLEB...	71C	PRL 27 888	H. Alvensleben <i>et al.</i>	(DESY)	REFID=20193
BALDIN	71	SJNP 13 758 Translated from YAF 13 1318.	A.B. Baldin <i>et al.</i>	(ITEP)	REFID=20195
BEHREND	71	PRL 27 61	H.J. Behrend <i>et al.</i>	(ROCH, CORN, FNAL)	REFID=20197
BIZZARRI	71	NP B27 140	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20198
COYNE	71	NP B32 333	D.G. Coyne <i>et al.</i>	(LRL)	REFID=20201
MOFFFEIT	71	NP B29 349	K.C. Moffeit <i>et al.</i>	(LRL, UCB, SLAC+)	REFID=20204
ABRAMOVI...	70	NP B20 209	M. Abramovich <i>et al.</i>	(CERN)	REFID=20180
BIGGS	70B	PRL 24 1201	P.J. Biggs <i>et al.</i>	(DARE)	REFID=20184
BIZZARRI	70	PRL 25 1385	R. Bizzarri <i>et al.</i>	(ROMA, SYRA)	REFID=20181
ROOS	70	DNPL/R7 173	M. Roos	(CERN)	REFID=20191
Proc. Daresbury Study Weekend No. 1.					
AUGUSTIN	69D	PL 28B 513	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=20169
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20171
DEINET	69B	PL 30B 426	W. Deinet <i>et al.</i>	(KARL, CERN)	REFID=20173
JACQUET	69B	NC 63A 743	F. Jacquet <i>et al.</i>	(EPOL, BERG)	REFID=20176
WILSON	69	Private Comm.	R. Wilson	(HARV)	REFID=20179
Also		PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20084
ASTVACAT...	68	PL 27B 45	R.G. Astvatsaturov <i>et al.</i>	(JINR, MOSU)	REFID=20055
BOLLINI	68C	NC 56A 531	D. Bollini <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20164
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
FELDMAN	67C	PR 159 1219	M. Feldman <i>et al.</i>	(PENN)	REFID=20161
DIGIUGNO	66B	NC 44A 1272	G. Di Giugno <i>et al.</i>	(NAPL, FRAS, TRST)	REFID=20156
FLATTE	66	PR 145 1050	S.M. Flatte <i>et al.</i>	(LRL)	REFID=20157
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
BARBARO...	65	PRL 14 279	A. Barbaro-Galtieri, R.D. Tripp	(LRL)	REFID=20152
BARMIN	64	JETP 18 1289 Translated from ZETF 45 1879.	V.V. Barmin <i>et al.</i>	(ITEP)	REFID=20149
KRAEMER	64	PR 136 B496	R.W. Kraemer <i>et al.</i>	(JHU, NWES, WOOD)	REFID=10755
BUSCHBECK	63	Siena Conf. 1 166	B. Buschbeck <i>et al.</i>	(VIEN, CERN, ANIK)	REFID=20146

$\eta'(958)$ 

$$I^G(J^{PC}) = 0^+(0^{-+})$$

NODE=M002

 $\eta'(958)$  MASS

NODE=M002M

NODE=M002M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>957.78 ±0.06 OUR AVERAGE</b>				
957.793±0.054±0.036	3.9k	LIBBY	08	CLEO $J/\psi \rightarrow \gamma\eta'$
957.9 ±0.2 ±0.6	4800	WURZINGER	96	SPEC 1.68 $p d \rightarrow {}^3\text{He}\eta'$
957.46 ±0.33		DUANE	74	MMS $\pi^- p \rightarrow n\text{MM}$
958.2 ±0.5	1414	DANBURG	73	HBC 2.2 $K^- p \rightarrow \Lambda\eta'$
958 ±1	400	JACOBS	73	HBC 2.9 $K^- p \rightarrow \Lambda\eta'$
956.1 ±1.1	3415	<sup>1</sup> BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
957.5 ±0.2		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
959 ±1	630	<sup>2</sup> BELADIDZE	92C	VES 36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
958 ±1	340	<sup>2</sup> ARMSTRONG	91B	OMEG 300 $pp \rightarrow pp\eta\pi^+\pi^-$
958.2 ±0.4	622	<sup>2</sup> AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
957.8 ±0.2	2420	<sup>2</sup> AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
956.3 ±1.0	143	<sup>2</sup> GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
957.4 ±1.4	535	<sup>3</sup> BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
957 ±1		RITTENBERG	69	HBC 1.7-2.7 $K^- p$

<sup>1</sup> Using all  $\eta'$  decays.<sup>2</sup> Systematic uncertainty not estimated.<sup>3</sup> Using  $\eta'$  decays into neutrals. Not independent of the other listed BASILE 71  $\eta'$  mass measurement.

OCCUR=2

OCCUR=2

NODE=M002M;LINKAGE=BS

NODE=M002M;LINKAGE=NS

NODE=M002M;LINKAGE=BA

 $\eta'(958)$  WIDTH

NODE=M002W

NODE=M002W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.188±0.006 OUR FIT</b>					
<b>0.230±0.021 OUR AVERAGE</b>					
0.226±0.017±0.014	2300	CZERWINSKI	10	MMS	$pp \rightarrow pp\eta'$
0.40 ±0.22	4800	WURZINGER	96	SPEC	1.68 $p d \rightarrow {}^3\text{He}\eta'$
0.28 ±0.10	1000	BINNIE	79	MMS	0 $\pi^- p \rightarrow n\text{MM}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.20 ±0.04		BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

 $\eta'(958)$  DECAY MODES

NODE=M002215;NODE=M002

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\pi^+\pi^-\eta$	(42.5 ±0.5 )%	DESIG=1
$\Gamma_2$ $\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$ )	(29.5 ±0.4 )%	DESIG=9
$\Gamma_3$ $\rho^0\gamma$		DESIG=213
$\Gamma_4$ $\pi^0\pi^0\eta$	(22.4 ±0.5 )%	DESIG=2
$\Gamma_5$ $\omega\gamma$	( 2.52 ±0.07 )%	DESIG=7
$\Gamma_6$ $\omega e^+e^-$	( 2.0 ±0.4 ) × 10 <sup>-4</sup>	DESIG=205
$\Gamma_7$ $\gamma\gamma$	( 2.307±0.033 )%	DESIG=6
$\Gamma_8$ $3\pi^0$	( 2.50 ±0.17 ) × 10 <sup>-3</sup>	DESIG=8
$\Gamma_9$ $\mu^+\mu^-\gamma$	( 1.13 ±0.28 ) × 10 <sup>-4</sup>	DESIG=20
$\Gamma_{10}$ $\pi^+\pi^-\mu^+\mu^-$	( 2.0 ±0.4 ) × 10 <sup>-5</sup>	DESIG=201
$\Gamma_{11}$ $\pi^+\pi^-\pi^0$	( 3.61 ±0.17 ) × 10 <sup>-3</sup>	DESIG=121
$\Gamma_{12}$ $(\pi^+\pi^-\pi^0)$ S-wave	( 3.8 ±0.5 ) × 10 <sup>-3</sup>	DESIG=211
$\Gamma_{13}$ $\pi^\mp\rho^\pm$	( 7.4 ±2.3 ) × 10 <sup>-4</sup>	DESIG=210
$\Gamma_{14}$ $2(\pi^+\pi^-)$	( 8.4 ±0.9 ) × 10 <sup>-5</sup>	DESIG=131
$\Gamma_{15}$ $\pi^+\pi^-2\pi^0$	( 1.8 ±0.4 ) × 10 <sup>-4</sup>	DESIG=202
$\Gamma_{16}$ $2(\pi^+\pi^-)$ neutrals	< 1 %	95% DESIG=132

$\Gamma_{17}$	$2(\pi^+\pi^-\pi^0)$	$< 1.8$	$\times 10^{-3}$	90%	DESIG=141
$\Gamma_{18}$	$2(\pi^+\pi^-)2\pi^0$	$< 1$	%	95%	DESIG=15
$\Gamma_{19}$	$3(\pi^+\pi^-)$	$< 3.1$	$\times 10^{-5}$	90%	DESIG=203
$\Gamma_{20}$	$K^\pm\pi^\mp$	$< 4$	$\times 10^{-5}$	90%	DESIG=207
$\Gamma_{21}$	$\pi^+\pi^-e^+e^-$	$(2.42 \pm 0.10)$	$\times 10^{-3}$		DESIG=10
$\Gamma_{22}$	$\pi^+e^-\nu_e + \text{c.c.}$	$< 2.1$	$\times 10^{-4}$	90%	DESIG=204
$\Gamma_{23}$	$\gamma e^+e^-$	$(4.91 \pm 0.27)$	$\times 10^{-4}$		DESIG=28
$\Gamma_{24}$	$\pi^0\gamma\gamma$	$(3.20 \pm 0.24)$	$\times 10^{-3}$		DESIG=24
$\Gamma_{25}$	$\pi^0\gamma\gamma$ (non resonant)	$(6.2 \pm 0.9)$	$\times 10^{-4}$		DESIG=212
$\Gamma_{26}$	$\eta\gamma\gamma$	$< 1.33$	$\times 10^{-4}$	90%	DESIG=214
$\Gamma_{27}$	$4\pi^0$	$< 4.94$	$\times 10^{-5}$	90%	DESIG=26
$\Gamma_{28}$	$e^+e^-$	$< 5.6$	$\times 10^{-9}$	90%	DESIG=150
$\Gamma_{29}$	$e^+e^-e^+e^-$	$(4.5 \pm 1.1)$	$\times 10^{-6}$		DESIG=215
$\Gamma_{30}$	invisible	$< 6$	$\times 10^{-4}$	90%	DESIG=200

**Charge conjugation (C), Parity (P),  
Lepton family number (LF) violating modes**

NODE=M002;CLUMP=B

$\Gamma_{31}$	$\pi^+\pi^-$	$P, CP$	$< 1.8$	$\times 10^{-5}$	90%	DESIG=111
$\Gamma_{32}$	$\pi^0\pi^0$	$P, CP$	$< 4$	$\times 10^{-4}$	90%	DESIG=25
$\Gamma_{33}$	$\pi^0e^+e^-$	$C$	[a] $< 1.4$	$\times 10^{-3}$	90%	DESIG=16
$\Gamma_{34}$	$\pi^0\rho^0$	$C$	$< 4$	%	90%	DESIG=18
$\Gamma_{35}$	$\eta e^+e^-$	$C$	[a] $< 2.4$	$\times 10^{-3}$	90%	DESIG=17
$\Gamma_{36}$	$3\gamma$	$C$	$< 1.0$	$\times 10^{-4}$	90%	DESIG=23
$\Gamma_{37}$	$\mu^+\mu^-\pi^0$	$C$	[a] $< 6.0$	$\times 10^{-5}$	90%	DESIG=22
$\Gamma_{38}$	$\mu^+\mu^-\eta$	$C$	[a] $< 1.5$	$\times 10^{-5}$	90%	DESIG=21
$\Gamma_{39}$	$e\mu$	$LF$	$< 4.7$	$\times 10^{-4}$	90%	DESIG=27

[a] C parity forbids this to occur as a single-photon process.

LINKAGE=CS

**CONSTRAINED FIT INFORMATION**

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 20 branching ratios uses 52 measurements and one constraint to determine 9 parameters. The overall fit has a  $\chi^2 = 69.5$  for 44 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-25							
$x_4$	-75	-43						
$x_5$	-7	-6	-2					
$x_7$	-11	-7	9	-1				
$x_8$	-17	-10	19	0	2			
$x_{11}$	-1	-1	-1	0	0	0		
$x_{21}$	-8	30	-14	-2	-2	-3	0	
$\Gamma$	11	-10	-1	1	-40	0	0	-3
	$x_1$	$x_2$	$x_4$	$x_5$	$x_7$	$x_8$	$x_{11}$	$x_{21}$

	Mode	Rate (MeV)	
$\Gamma_1$	$\pi^+\pi^-\eta$	$0.0799 \pm 0.0029$	DESIG=1
$\Gamma_2$	$\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$ )	$0.0554 \pm 0.0019$	DESIG=9
$\Gamma_4$	$\pi^0\pi^0\eta$	$0.0421 \pm 0.0017$	DESIG=2
$\Gamma_5$	$\omega\gamma$	$0.00474 \pm 0.00020$	DESIG=7
$\Gamma_7$	$\gamma\gamma$	$0.00434 \pm 0.00013$	DESIG=6
$\Gamma_8$	$3\pi^0$	$(4.7 \pm 0.4) \times 10^{-4}$	DESIG=8
$\Gamma_{11}$	$\pi^+\pi^-\pi^0$	$(6.8 \pm 0.4) \times 10^{-4}$	DESIG=121
$\Gamma_{21}$	$\pi^+\pi^-e^+e^-$	$(4.54 \pm 0.23) \times 10^{-4}$	DESIG=10

$\eta'(958)$  PARTIAL WIDTHS

NODE=M002220

 $\Gamma(\gamma\gamma)$  $\Gamma_7$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**4.34±0.14 OUR FIT****4.28±0.19 OUR AVERAGE**

4.17±0.10±0.27	2000	1 ACCIARRI	98Q L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.53±0.29±0.51	266	KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
3.61±0.13±0.48		2 BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\eta'(958)$
4.6 ±1.1 ±0.6	23	BARU	90 MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.57±0.25±0.44		BUTLER	90 MRK2	$e^+e^- \rightarrow e^+e^-\eta'(958)$
5.08±0.24±0.71	547	3 ROE	90 ASP	$e^+e^- \rightarrow e^+e^-2\gamma$
3.8 ±0.7 ±0.6	34	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.9 ±0.5 ±0.5	136	4 WILLIAMS	88 CBAL	$e^+e^- \rightarrow e^+e^-2\gamma$
4.7 ±0.6 ±0.9	143	5 GIDAL	87 MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.0 ±0.9		6 BARTEL	85E JADE	$e^+e^- \rightarrow e^+e^-2\gamma$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup>No non-resonant  $\pi^+\pi^-$  contribution found.
- <sup>2</sup>Reevaluated by us using  $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$ .
- <sup>3</sup>Reevaluated by us using  $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$ .
- <sup>4</sup>Reevaluated by us using  $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$ .
- <sup>5</sup>Superseded by BUTLER 90.
- <sup>6</sup>Systematic error not evaluated.

NODE=M002W4  
NODE=M002W4NODE=M002W4;LINKAGE=AC  
NODE=M002W4;LINKAGE=K1  
NODE=M002W4;LINKAGE=K2  
NODE=M002W4;LINKAGE=K3  
NODE=M002W4;LINKAGE=C  
NODE=M002W4;LINKAGE=A $\Gamma(e^+e^-)$  $\Gamma_{28}$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

**<1.1 × 10<sup>-3</sup>** 90 1,2 ACHASOV 15 SND 0.958  $e^+e^- \rightarrow \pi\pi\eta$ 

- • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0 × 10<sup>-3</sup> 90 2 ACHASOV 15 SND 0.958  $e^+e^- \rightarrow \pi\pi\eta$ <2.4 × 10<sup>-3</sup> 90 2 AKHMETSHIN 15 CMD3 0.958  $e^+e^- \rightarrow \pi^+\pi^-\eta$ <sup>1</sup>Combining data of ACHASOV 15 and AKHMETSHIN 15.<sup>2</sup>Using  $\eta$  and  $\eta'$  branching fractions from PDG 14.NODE=M002W1  
NODE=M002W1

OCCUR=2

NODE=M002W1;LINKAGE=A  
NODE=M002W1;LINKAGE=B $\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

This combination of a partial width with the partial width into  $\gamma\gamma$  and with the total width is obtained from the integrated cross section into channel(i) in the  $\gamma\gamma$  annihilation.

NODE=M002223

NODE=M002223

 $\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$  $\Gamma_7\Gamma_2/\Gamma$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**1.28±0.04 OUR FIT****1.26±0.07 OUR AVERAGE** Error includes scale factor of 1.2.

1.09±0.04±0.13		BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\rho(770)^0\gamma$
1.35±0.09±0.21		AIHARA	87 TPC	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.13±0.04±0.13	867	ALBRECHT	87B ARG	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.53±0.09±0.21		ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.14±0.08±0.11	243	BERGER	84B PLUT	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.73±0.34±0.35	95	JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.49±0.13±0.027	213	BARTEL	82B JADE	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.85±0.31±0.24	43	BEHREND	82C CELL	$e^+e^- \rightarrow e^+e^-\rho\gamma$

- • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M002G1  
NODE=M002G1 $\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$  $\Gamma_7\Gamma_4/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**0.97±0.04 OUR FIT** Error includes scale factor of 1.1.**0.92±0.06±0.11** <sup>1</sup>KARCH 92 CBAL  $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$ 

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.95±0.05±0.08 <sup>2</sup>KARCH 90 CBAL  $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$ 1.00±0.08±0.10 <sup>2,3</sup>ANTREASYAN 87 CBAL  $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$ <sup>1</sup>Reevaluated by us using  $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$ . Supersedes ANTREASYAN 87 and KARCH 90.<sup>2</sup>Superseded by KARCH 92.<sup>3</sup>Using  $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$ .NODE=M002G2  
NODE=M002G2

NODE=M002G2;LINKAGE=K4

NODE=M002G2;LINKAGE=A  
NODE=M002G2;LINKAGE=D

$\eta'(958) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M002224

 $\Gamma(\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_{28}/\Gamma$ NODE=M002G01  
NODE=M002G01

VALUE ( $10^{-3}$ eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.0</b>	90	<sup>1</sup> AKHMETSHIN 15	CMD3	$0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$
<sup>1</sup> AKHMETSHIN 15 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta) \times \Gamma(\eta'(958) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 4.1 \times 10^{-4}$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .				

NODE=M002G01;LINKAGE=A

 $\eta'(958) \text{ BRANCHING RATIOS}$ 

NODE=M002230

 $\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$ NODE=M002R47  
NODE=M002R47

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>42.5 ± 0.5 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>41.24 ± 0.08 ± 1.24</b>	312k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
42.4 ± 1.1 ± 0.4	1.2k	<sup>1</sup> PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
<sup>1</sup> Not independent of other $\eta'$ branching fractions and ratios in PEDLAR 09.				

NODE=M002R47;LINKAGE=PE

 $\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{\text{total}} \quad 0.2804\Gamma_1/\Gamma$ NODE=M002R3  
NODE=M002R3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.1191 ± 0.0015 OUR FIT</b>	Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.123 ± 0.014	107	RITTENBERG	69 HBC	$1.7-2.7 K^-p$
0.10 ± 0.04	10	LONDON	66 HBC	$2.24 K^-p \rightarrow \Lambda 2\pi^+ 2\pi^- \pi^0$
0.07 ± 0.04	7	BADIER	65B HBC	$3 K^-p$

 $\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{\text{total}} \quad 0.7196\Gamma_1/\Gamma$ NODE=M002R1  
NODE=M002R1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.306 ± 0.004 OUR FIT</b>	Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.314 ± 0.026	281	RITTENBERG	69 HBC	$1.7-2.7 K^-p$

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$ NODE=M002R6  
NODE=M002R6

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>29.5 ± 0.4 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>29.90 ± 0.03 ± 0.55</b>	913k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
28.7 ± 0.7 ± 0.4	0.2k	<sup>1</sup> PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
32.9 ± 3.3	298	RITTENBERG	69 HBC	$1.7-2.7 K^-p$
20 ± 10	20	LONDON	66 HBC	$2.24 K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$
34 ± 9	35	BADIER	65B HBC	$3 K^-p$

<sup>1</sup> Not independent of other  $\eta'$  branching fractions and ratios in PEDLAR 09.

NODE=M002R6;LINKAGE=PE

 $\Gamma(\rho^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$ NODE=M002R66  
NODE=M002R66

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
33.34 ± 0.06 ± 1.60	970k	<sup>1</sup> ABLIKIM	18C BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
34.43 ± 0.52 ± 1.97	970k	<sup>2</sup> ABLIKIM	18C BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$

OCCUR=2

<sup>1</sup> From a fit to  $\pi^+\pi^-$  mass using  $\rho(770)$ ,  $\omega(782)$ , and box anomaly components.

NODE=M002R66;LINKAGE=A

<sup>2</sup> From a fit to  $\pi^+\pi^-$  mass using  $\rho(770)$ ,  $\omega(782)$ , and  $\rho(1450)$  components.

NODE=M002R66;LINKAGE=B

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta) \quad \Gamma_2/\Gamma_1$ NODE=M002R43  
NODE=M002R43

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.694 ± 0.014 OUR FIT</b>	Error includes scale factor of 1.1.		
<b>0.683 ± 0.020 OUR AVERAGE</b>			
0.677 ± 0.024 ± 0.011	PEDLAR	09 CLE3	$J/\psi \rightarrow \eta'\gamma$
0.69 ± 0.03	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta'\gamma$



$$\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi^+ \pi^- \eta (\text{neutral decay}))$$
 $\Gamma_2/0.714\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.972 ± 0.020 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>0.97 ± 0.09 OUR AVERAGE</b>				
0.70 ± 0.22		AMSLER	04B CBAR	0 $\bar{p}p \rightarrow \pi^+ \pi^- \eta$
1.07 ± 0.17		BELADIDZE	92C VES	36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
0.92 ± 0.14	473	DANBURG	73 HBC	2.2 $K^- p \rightarrow \Lambda X^0$
1.11 ± 0.18	192	JACOBS	73 HBC	2.9 $K^- p \rightarrow \Lambda X^0$

NODE=M002R27  
 NODE=M002R27

$$\Gamma(\pi^0 \pi^0 \eta) / \Gamma_{\text{total}}$$
 $\Gamma_4/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>22.4 ± 0.6 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>21.36 ± 0.10 ± 0.92</b>	52k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
23.5 ± 1.3 ± 0.4	3.2k	<sup>1</sup> PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma \eta'$

NODE=M002R48  
 NODE=M002R48

<sup>1</sup> Not independent of other  $\eta'$  branching fractions and ratios in PEDLAR 09.

NODE=M002R48;LINKAGE=PE

$$\Gamma(\pi^0 \pi^0 \eta (3\pi^0 \text{ decay})) / \Gamma_{\text{total}}$$
 $0.321\Gamma_4/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0718 ± 0.0018 OUR FIT</b>	Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.11 ± 0.06	4	BENSINGER	70 DBC	2.2 $\pi^+ d$

NODE=M002R26  
 NODE=M002R26

$$\Gamma(\pi^0 \pi^0 \eta) / \Gamma(\pi^+ \pi^- \eta)$$
 $\Gamma_4/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.527 ± 0.019 OUR FIT</b>	Error includes scale factor of 1.1.		
<b>0.555 ± 0.043 ± 0.013</b>	PEDLAR	09 CLE3	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R45  
 NODE=M002R45

$$\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi \pi \eta)$$
 $\Gamma_2/(\Gamma_1 + \Gamma_4)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.454 ± 0.009 OUR FIT</b>	Error includes scale factor of 1.1.		
<b>0.43 ± 0.02 ± 0.02</b>	BARBERIS	98C OMEG	450 $p p \rightarrow p_f \eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.31 ± 0.15	DAVIS	68 HBC	5.5 $K^- p$

NODE=M002R7  
 NODE=M002R7

$$\Gamma(\omega \gamma) / \Gamma_{\text{total}}$$
 $\Gamma_5/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.52 ± 0.07 OUR FIT</b>				
<b>2.50 ± 0.07 OUR AVERAGE</b>				
2.489 ± 0.018 ± 0.074	23k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma \eta'$
2.55 ± 0.03 ± 0.16	33.2k	<sup>1</sup> ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta' \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.34 ± 0.30 ± 0.04	70	<sup>2</sup> PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma \eta'$

NODE=M002R49  
 NODE=M002R49

<sup>1</sup> Using  $B(J/\psi \rightarrow \eta' \gamma) = (5.15 \pm 0.16) \times 10^{-3}$  and  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$ .

<sup>2</sup> Not independent of other  $\eta'$  branching fractions and ratios in PEDLAR 09.

NODE=M002R49;LINKAGE=A  
 NODE=M002R49;LINKAGE=PE

$$\Gamma(\omega \gamma) / \Gamma(\pi^+ \pi^- \eta)$$
 $\Gamma_5/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0593 ± 0.0018 OUR FIT</b>	Error includes scale factor of 1.1.			
<b>0.055 ± 0.007 ± 0.001</b>		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta' \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.068 ± 0.013	68	ZANFINO	77 ASPK	8.4 $\pi^- p$

NODE=M002R17  
 NODE=M002R17

$$\Gamma(\omega \gamma) / \Gamma(\pi^0 \pi^0 \eta)$$
 $\Gamma_5/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.113 ± 0.004 OUR FIT</b>			
<b>0.147 ± 0.016</b>	ALDE	87B GAM2	38 $\pi^- p \rightarrow n 4\gamma$

NODE=M002R33  
 NODE=M002R33

$$\Gamma(\omega e^+ e^-) / \Gamma(\omega \gamma)$$
 $\Gamma_6/\Gamma_5$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
7.71 ± 1.34 ± 0.54	<sup>1</sup> ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R60  
 NODE=M002R60

<sup>1</sup> Obtained from other ABLIKIM 15AD measurements with common systematics taken into account.

NODE=M002R60;LINKAGE=A

$$\Gamma(\omega e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>1.97 ± 0.34 ± 0.17</b>	66	<sup>1</sup> ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta' \gamma$
---------------------------	----	----------------------	-----------	-----------------------------------

<sup>1</sup> Using  $B(J/\psi \rightarrow \eta' \gamma) = (5.15 \pm 0.16) \times 10^{-3}$  and  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M002R59  
NODE=M002R59

NODE=M002R59;LINKAGE=A

$$\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / [\Gamma(\pi^+ \pi^- \eta) + \Gamma(\pi^0 \pi^0 \eta) + \Gamma(\omega \gamma)] \quad \Gamma_2/(\Gamma_1 + \Gamma_4 + \Gamma_5)$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.437 ± 0.008 OUR FIT</b>	Error includes scale factor of 1.1.		
------------------------------	-------------------------------------	--	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ± 0.14	DAUBER	64	HBC	1.95 $K^- p$
-------------	--------	----	-----	--------------

NODE=M002R18  
NODE=M002R18

$$[\Gamma(\pi^0 \pi^0 \eta (\text{charged decay})) + \Gamma(\omega (\text{charged decay}) \gamma)] / \Gamma_{\text{total}} \quad (0.286\Gamma_4 + 0.89\Gamma_5)/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

<b>0.0864 ± 0.0017 OUR FIT</b>	Error includes scale factor of 1.1.			
--------------------------------	-------------------------------------	--	--	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ± 0.029	42	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
---------------	----	------------	----	-----	-----------------

NODE=M002R4  
NODE=M002R4

$$\Gamma(\pi^+ \pi^- \text{ neutrals}) / \Gamma_{\text{total}} \quad (0.714\Gamma_1 + 0.286\Gamma_4 + 0.89\Gamma_5)/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

<b>0.3897 ± 0.0028 OUR FIT</b>	Error includes scale factor of 1.1.			
--------------------------------	-------------------------------------	--	--	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 ± 0.1	39	LONDON	66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$
0.35 ± 0.06	33	BADIER	65B	HBC	3 $K^- p$

NODE=M002R2  
NODE=M002R2

$$\Gamma(\gamma \gamma) / \Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>2.307 ± 0.035 OUR FIT</b>	Error includes scale factor of 1.1.			
------------------------------	-------------------------------------	--	--	--

<b>2.31 ± 0.06 OUR AVERAGE</b>	Error includes scale factor of 1.8.			
--------------------------------	-------------------------------------	--	--	--

2.331 ± 0.012 ± 0.035	71k	ABLIKIM	19T	BES	$J/\psi \rightarrow \gamma \eta'$
-----------------------	-----	---------	-----	-----	-----------------------------------

1.99 $^{+0.31}_{-0.27}$ ± 0.07	114	<sup>1</sup> WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
--------------------------------	-----	--------------------	----	------	---

2.00 ± 0.18		<sup>2</sup> STANTON	80	SPEC	$8.45 \pi^- p \rightarrow n \pi^+ \pi^- 2\gamma$
-------------	--	----------------------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25 ± 0.16 ± 0.03	0.3k	<sup>3</sup> PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma \eta'$
--------------------	------	---------------------	----	------	-----------------------------------

1.8 ± 0.2	6000	<sup>4</sup> APEL	79	NICE	15–40 $\pi^- p \rightarrow n 2\gamma$
-----------	------	-------------------	----	------	---------------------------------------

2.5 ± 0.7		DUANE	74	MMS	$\pi^- p \rightarrow n \text{MM}$
-----------	--	-------	----	-----	-----------------------------------

1.71 ± 0.33	68	DALPIAZ	72	CNTR	$1.6 \pi^- p \rightarrow n X^0$
-------------	----	---------	----	------	---------------------------------

2.0 $^{+0.8}_{-0.6}$	31	HARVEY	71	OSPK	$3.65 \pi^- p \rightarrow n X^0$
----------------------	----	--------	----	------	----------------------------------

<sup>1</sup> WICHT 08 reports  $[\Gamma(\eta'(958) \rightarrow \gamma \gamma) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta' K^+) = (7.04 \pm 0.25) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Includes APEL 79 result.

<sup>3</sup> Not independent of other  $\eta'$  branching fractions and ratios in PEDLAR 09.

<sup>4</sup> Data is included in STANTON 80 evaluation.

NODE=M002R19  
NODE=M002R19

NODE=M002R19;LINKAGE=WI

NODE=M002R19;LINKAGE=S  
NODE=M002R19;LINKAGE=PE  
NODE=M002R19;LINKAGE=A

$$\Gamma(\gamma \gamma) / \Gamma(\pi^+ \pi^- \eta) \quad \Gamma_7/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.0543 ± 0.0012 OUR FIT</b>	Error includes scale factor of 1.1.		
--------------------------------	-------------------------------------	--	--

<b>0.053 ± 0.004 ± 0.001</b>	PEDLAR	09	CLE3	$J/\psi \rightarrow \eta' \gamma$
------------------------------	--------	----	------	-----------------------------------

NODE=M002R46  
NODE=M002R46

$$\Gamma(\gamma \gamma) / \Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) \quad \Gamma_7/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.0783 ± 0.0016 OUR FIT</b>	Error includes scale factor of 1.1.		
--------------------------------	-------------------------------------	--	--

<b>0.080 ± 0.008</b>	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta' \gamma$
----------------------	---------	-----	------	-----------------------------------

NODE=M002R42  
NODE=M002R42

$$\Gamma(\gamma \gamma) / \Gamma(\pi^0 \pi^0 \eta) \quad \Gamma_7/\Gamma_4$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>0.1031 ± 0.0028 OUR FIT</b>			
--------------------------------	--	--	--

<b>0.105 ± 0.010 OUR AVERAGE</b>	Error includes scale factor of 1.9.		
----------------------------------	-------------------------------------	--	--

0.091 ± 0.009	AMSLER	93	CBAR	0.0 $\bar{p} p$
---------------	--------	----	------	-----------------

0.112 ± 0.002 ± 0.006	ALDE	87B	GAM2	38 $\pi^- p \rightarrow n 2\gamma$
-----------------------	------	-----	------	------------------------------------

NODE=M002R38  
NODE=M002R38

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$  (neutral decay) $\Gamma_7/0.714\Gamma_4$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M002R28  
 NODE=M002R28

**0.144 ± 0.004 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.188 ± 0.058	16	APEL	72	OSPK	$3.8 \pi^- p \rightarrow n\chi^0$
---------------	----	------	----	------	-----------------------------------

 $\Gamma(\text{neutrals})/\Gamma_{\text{total}}$  $(0.714\Gamma_4 + 0.09\Gamma_5 + \Gamma_7)/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M002R5  
 NODE=M002R5

**0.185 ± 0.004 OUR FIT** Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.185 ± 0.022	535	BASILE	71	CNTR	$1.6 \pi^- p \rightarrow n\chi^0$
0.189 ± 0.026	123	RITTENBERG	69	HBC	$1.7-2.7 K^- p$

 $\Gamma(3\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R55  
 NODE=M002R55

**2.50 ± 0.17 OUR FIT****3.57 ± 0.26 OUR AVERAGE**

3.522 ± 0.082 ± 0.254	2015	ABLIKIM	17	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
4.79 ± 0.59 ± 1.14	183	<sup>1</sup> ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.56 ± 0.22 ± 0.34	309	<sup>2</sup> ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
--------------------	-----	----------------------	-----	------	-------------------------------------

<sup>1</sup>We have added all systematic uncertainties in quadrature to a single value.<sup>2</sup>Superseded by ABLIKIM 17.

NODE=M002R55;LINKAGE=A  
 NODE=M002R55;LINKAGE=B

 $\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$  $\Gamma_8/\Gamma_4$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R32  
 NODE=M002R32

**112 ± 8 OUR FIT****78 ± 10 OUR AVERAGE**

86 ± 19	235	BLIK	08	GAMS	$32 \pi^- p \rightarrow \eta' n$
74 ± 15		ALDE	87B	GAM2	$38 \pi^- p \rightarrow n6\gamma$
75 ± 18		BINON	84	GAM2	$30-40 \pi^- p \rightarrow n6\gamma$

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$  $\Gamma_9/\Gamma_7$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R29  
 NODE=M002R29

**4.9 ± 1.2**

33	VIKTOROV	80	CNTR	$25,33 \pi^- p \rightarrow 2\mu\gamma$
----	----------	----	------	--

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M002R50  
 NODE=M002R50

**1.95 ± 0.37 ± 0.03** 53 <sup>1</sup>ABLIKIM 21I BES3  $J/\psi \rightarrow \gamma\eta'(958)$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.9	90	<sup>2</sup> ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
< 24	90	<sup>3</sup> NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$

<sup>1</sup>ABLIKIM 21I reports  $(1.97 \pm 0.33 \pm 0.19) \times 10^{-5}$  from a measurement of  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$  assuming  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.21 \pm 0.17) \times 10^{-3}$ , which we rescale to our best value  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 \pm 0.07) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup>Using  $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$  from PDG 12.<sup>3</sup>Not independent of measured value of  $\Gamma_{10}/\Gamma_1$  from NAIK 09.

NODE=M002R50;LINKAGE=E

NODE=M002R50;LINKAGE=A  
 NODE=M002R50;LINKAGE=NA

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{10}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R03  
 NODE=M002R03

**< 0.5** 90 <sup>1</sup>NAIK 09 CLEO  $J/\psi \rightarrow \gamma\eta'$ 

<sup>1</sup>NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M002R03;LINKAGE=NA

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\rho^0\gamma)$  (including non-resonant  $\pi^+\pi^-\gamma$ ) $\Gamma_{10}/\Gamma_2$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R57  
 NODE=M002R57

**< 1.0** 90 ABLIKIM 130 BES3  $J/\psi \rightarrow \gamma\eta'$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>3.61 ± 0.18 OUR FIT</b>				
<b>3.61 ± 0.18 OUR AVERAGE</b>				
3.591 ± 0.054 ± 0.174	6067	ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
4.28 ± 0.49 ± 1.11	78	<sup>1</sup> ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$
3.7 $^{+1.1}_{-0.9}$ ± 0.4		<sup>2</sup> NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.83 ± 0.15 ± 0.39 1014 <sup>3</sup> ABLIKIM 12E BES3  $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$

<sup>1</sup> We have added all systematic uncertainties in quadrature to a single value.

<sup>2</sup> Not independent of measured value of  $\Gamma_{11}/\Gamma_1$  from NAIK 09.

<sup>3</sup> Superseded by ABLIKIM 17.

NODE=M002R21  
NODE=M002R21

NODE=M002R21;LINKAGE=A  
NODE=M002R21;LINKAGE=NA  
NODE=M002R21;LINKAGE=B

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$   $\Gamma_{11}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>8.5 ± 0.4 OUR FIT</b>				Error includes scale factor of 1.1.
<b>8.27 <math>^{+2.49}_{-2.12}</math> ± 0.04</b>	20	<sup>1</sup> NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (21  $^{+6}_{-5}$  ± 2) × 10<sup>-3</sup> which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 ± 0.18) × 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.$

NODE=M002R01  
NODE=M002R01

NODE=M002R01;LINKAGE=NA

 $\Gamma((\pi^+\pi^-\pi^0) \text{ S-wave})/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>37.63 ± 0.77 ± 5.00</b>	6580	<sup>1</sup> ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$

<sup>1</sup> We have added all systematic uncertainties in quadrature .

NODE=M002R63  
NODE=M002R63

NODE=M002R63;LINKAGE=A

 $\Gamma(\pi^\mp\rho^\pm)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>7.44 ± 0.60 ± 2.23</b>	1231	<sup>1</sup> ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^\mp\rho^\pm)$

<sup>1</sup> We have added all systematic uncertainties in quadrature .

NODE=M002R62  
NODE=M002R62

NODE=M002R62;LINKAGE=A

 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>8.4 ± 0.9 ± 0.1</b>		199	<sup>1</sup> ABLIKIM	14M BES3	$J/\psi \rightarrow \gamma\eta'$
< 24	90		<sup>2</sup> NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
< 1000	90		RITTENBERG	69 HBC	1.7-2.7 $K^-p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 14M reports  $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] × [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$  =  $(4.40 ± 0.35 ± 0.30) × 10^{-7}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 ± 0.07) × 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Not independent of measured value of  $\Gamma_{14}/\Gamma_1$  from NAIK 09.

NODE=M002R24  
NODE=M002R24

NODE=M002R24;LINKAGE=A

NODE=M002R24;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$   $\Gamma_{14}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.6</b>	90	<sup>1</sup> NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$  <  $1.4 × 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 × 10^{-2}$ .

NODE=M002R04  
NODE=M002R04

NODE=M002R04;LINKAGE=NA

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1.79 ± 0.38 ± 0.02</b>		84	<sup>1</sup> ABLIKIM	14M BES3	$J/\psi \rightarrow \gamma\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 27 90 <sup>2</sup> NAIK 09 CLEO  $J/\psi \rightarrow \gamma\eta'$

<sup>1</sup> ABLIKIM 14M reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}] × [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$  =  $(9.38 ± 1.79 ± 0.89) × 10^{-7}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 ± 0.07) × 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Not independent of measured value of  $\Gamma_{15}/\Gamma_1$  from NAIK 09.

NODE=M002R51  
NODE=M002R51

NODE=M002R51;LINKAGE=A

NODE=M002R51;LINKAGE=NA

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{15}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	<sup>1</sup> NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

NODE=M002R05  
NODE=M002R05

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 15 \times 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M002R05;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-) \text{ neutrals})/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	DANBURG	73	HBC $2.2 K^-p \rightarrow \Lambda X^0$

NODE=M002R22  
NODE=M002R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01	90	RITTENBERG	69	HBC $1.7-2.7 K^-p$
-------	----	------------	----	--------------------

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M002R23  
NODE=M002R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.002	90	<sup>1</sup> NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
--------	----	-------------------	----	---------------------------------------

<0.01	90	RITTENBERG	69	HBC $1.7-2.7 K^-p$
-------	----	------------	----	--------------------

<sup>1</sup> Not independent of measured value of  $\Gamma_{17}/\Gamma_1$  from NAIK 09.

NODE=M002R23;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{17}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<4	90	<sup>1</sup> NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
----	----	-------------------	----	---------------------------------------

NODE=M002R06  
NODE=M002R06

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M002R06;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.01	95	KALBFLEISCH	64B	HBC $K^-p \rightarrow \Lambda 2(\pi^+\pi^-)+MM$
-------	----	-------------	-----	---

NODE=M002R16  
NODE=M002R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01	90	LONDON	66	HBC Compilation
-------	----	--------	----	-----------------

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 3.1	90	<sup>1</sup> ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
-------	----	----------------------	-----	--

NODE=M002R07  
NODE=M002R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 53	90	<sup>2</sup> NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
------	----	-------------------	----	---------------------------------------

<500	95	KALBFLEISCH	64B	HBC $K^-p \rightarrow \Lambda 2(\pi^+\pi^-)$
------	----	-------------	-----	--

<sup>1</sup> Using  $B(J/\psi \rightarrow \gamma\eta'(958)) = (5.16 \pm 0.15) \times 10^{-3}$ .

<sup>2</sup> Not independent of measured value of  $\Gamma_{19}/\Gamma_1$  from NAIK 09.

NODE=M002R07;LINKAGE=A  
NODE=M002R07;LINKAGE=NA $\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{19}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.2	90	<sup>1</sup> NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
------	----	-------------------	----	---------------------------------------

NODE=M002R08  
NODE=M002R08

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M002R08;LINKAGE=NA

 $\Gamma(K^\pm\pi^\mp)/\Gamma(\rho^0\gamma \text{ (including non-resonant } \pi^+\pi^-\gamma))$  $\Gamma_{20}/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.3 $\times 10^{-4}$	90	ABLIKIM	16M	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
-----------------------	----	---------	-----	---

NODE=M002R61  
NODE=M002R61 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**2.42 ± 0.10 OUR FIT**NODE=M002R12  
NODE=M002R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.11 ± 0.12 ± 0.14	429	<sup>1</sup> ABLIKIM	130	BES3 $J/\psi \rightarrow \gamma\eta'$
--------------------	-----	----------------------	-----	---------------------------------------

2.5 $\begin{smallmatrix} +1.2 \\ -0.9 \end{smallmatrix}$ ± 0.5		<sup>2</sup> NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
--	--	-------------------	----	---------------------------------------

<6	90	RITTENBERG	65	HBC $2.7 K^-p$
----	----	------------	----	----------------

<sup>1</sup> Using  $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$  from PDG 12.

<sup>2</sup> Not independent of measured value of  $\Gamma_{21}/\Gamma_1$  from NAIK 09.

NODE=M002R12;LINKAGE=A  
NODE=M002R12;LINKAGE=NA

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{21}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.69±0.25 OUR FIT</b>				

NODE=M002R02  
 NODE=M002R02

<b>5.51<sup>+3.00</sup><sub>-2.30</sub>±0.03</b>	8	<sup>1</sup> NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
--	---	-------------------	----	------	----------------------------------

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R02;LINKAGE=NA

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$  $\Gamma_{21}/\Gamma_2$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.20±0.31 OUR FIT</b>				

NODE=M002R56  
 NODE=M002R56

<b>8.20±0.16±0.27</b>	2584	ABLIKIM	21J	BES3	$J/\psi \rightarrow \gamma\eta'$
-----------------------	------	---------	-----	------	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 0.4 ± 0.5	429	<sup>1</sup> ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
-----------------	-----	----------------------	-----	------	----------------------------------

<sup>1</sup> Superseded by ABLIKIM 21J.

NODE=M002R56;LINKAGE=A

 $\Gamma(\pi^+e^-\nu_e + \text{c.c.})/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{22}/\Gamma_1$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;5.0</b>	90	ABLIKIM	13G	BES3	$J/\psi \rightarrow \phi\eta'$

NODE=M002R54  
 NODE=M002R54

 $\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.9</b>	90	BRIERE	00	CLEO	$10.6 e^+e^-$

NODE=M002R40  
 NODE=M002R40

 $\Gamma(\gamma e^+e^-)/\Gamma(\gamma\gamma)$  $\Gamma_{23}/\Gamma_7$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.13±0.09±0.07</b>	864	ABLIKIM	150	BES3	$J/\psi \rightarrow \gamma e^+e^-$

NODE=M002R00  
 NODE=M002R00

 $\Gamma(\pi^0\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.20±0.07±0.23</b>	3.4k	ABLIKIM	17T	BES3	$J/\psi \rightarrow \gamma\eta'$

NODE=M002R64  
 NODE=M002R64

 $\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$  $\Gamma_{24}/\Gamma_4$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;37</b>	90	ALDE	87B	GAM2	$38 \pi^- p \rightarrow n4\gamma$

NODE=M002R35  
 NODE=M002R35

 $\Gamma(\pi^0\gamma\gamma(\text{non resonant}))/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>6.16±0.64±0.67</b>	655	ABLIKIM	17T	BES3	$J/\psi \rightarrow \gamma\eta'$

NODE=M002R65  
 NODE=M002R65

 $\Gamma(\eta\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;1.33 × 10<sup>-4</sup></b>	90	ABLIKIM	19AW	BES3	$J/\psi \rightarrow \gamma\eta' \rightarrow \gamma\gamma\gamma 2\gamma$

NODE=M002R67  
 NODE=M002R67

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;4.94 × 10<sup>-5</sup></b>	90	ABLIKIM	20E	BES3	$J/\psi \rightarrow \eta'\gamma$

NODE=M002R58  
 NODE=M002R58

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2 × 10 <sup>-4</sup>	90	DONSKOV	14	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
-------------------------	----	---------	----	------	------------------------------------

 $\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$  $\Gamma_{27}/\Gamma_4$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;23</b>	90	ALDE	87B	GAM2	$38 \pi^- p \rightarrow n8\gamma$

NODE=M002R37  
 NODE=M002R37

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{28}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-9}$	90	<sup>1</sup> ACHASOV 15	SND	$0.958 e^+e^- \rightarrow \pi\pi\eta$
$< 12 \times 10^{-9}$	90	<sup>2</sup> AKHMETSHIN 15	CMD3	$0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$
$< 2.1 \times 10^{-7}$	90	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^+\pi^-\eta$

NODE=M002R39  
NODE=M002R39

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Combining data of ACHASOV 15 and AKHMETSHIN 15 and using  $\Gamma(\eta') = 0.198 \pm 0.009$  MeV.  
<sup>2</sup> Using  $\Gamma_{\eta'(958)} = 198 \pm 9$  keV,  $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.9 \pm 0.7)\%$ , and  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.20)\%$ .

NODE=M002R39;LINKAGE=B

NODE=M002R39;LINKAGE=A

 $\Gamma(e^+e^-e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{29}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.5$	30	<sup>1</sup> ABLIKIM 22E	BES3	$J/\psi \rightarrow \gamma\eta'$

<sup>1</sup> ABLIKIM 22E reports  $(4.5 \pm 1.0 \pm 0.5) \times 10^{-6}$  from a measurement of  $[\Gamma(\eta'(958) \rightarrow e^+e^-e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$  assuming  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 \pm 0.07) \times 10^{-3}$ .

NODE=M002R68  
NODE=M002R68

NODE=M002R68;LINKAGE=B

 $\Gamma(\text{invisible})/\Gamma_{\text{total}}$  $\Gamma_{30}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 9.5$	90	<sup>1</sup> NAIK 09	CLEO	$J/\psi \rightarrow \gamma\eta'$

<sup>1</sup> Not independent of measured value of  $\Gamma_{30}/\Gamma_1$  from NAIK 09.

NODE=M002R52  
NODE=M002R52

NODE=M002R52;LINKAGE=NA

 $\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\eta)$  $\Gamma_{30}/\Gamma_1$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1$	90	<sup>1</sup> NAIK 09	CLEO	$J/\psi \rightarrow \gamma\eta'$

<sup>1</sup> NAIK 09 reports  $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 5.4 \times 10^{-3}$  which we multiply by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M002R09  
NODE=M002R09

NODE=M002R09;LINKAGE=NA

 $\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$  $\Gamma_{30}/\Gamma_7$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.4$	90	ABLIKIM 13	BES3	$J/\psi \rightarrow \phi\eta'$
$< 6.69$	90	ABLIKIM 06Q	BES	$J/\psi \rightarrow \phi\eta'$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M002R44  
NODE=M002R44 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.18$	90	<sup>1</sup> AAIJ 17D	LHCB	$D_{(s)}^+ \rightarrow \pi^+\pi^-\pi^+$
$< 0.5$	90	<sup>2</sup> ABLIKIM 11G	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$< 29$	90	<sup>3</sup> MORI 07A	BELL	$\gamma\gamma \rightarrow \pi^+\pi^-$
$< 3.3$	90	<sup>4</sup> MORI 07A	BELL	$\gamma\gamma \rightarrow \pi^+\pi^-$
$< 800$	95	DANBURG 73	HBC	$2.2 K^-p \rightarrow \Lambda X^0$
$< 200$	90	RITTENBERG 69	HBC	$1.7-2.7 K^-p$

NODE=M002R20  
NODE=M002R20

OCCUR=2

<sup>1</sup> Using branching fractions of  $D_{(s)}^+$  decays from PDG 15.  
<sup>2</sup> ABLIKIM 11G reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.25 \times 10^{-3}$ .  
<sup>3</sup> Taking into account interference with the  $\gamma\gamma \rightarrow \pi^+\pi^-$  continuum.  
<sup>4</sup> Without interference with the  $\gamma\gamma \rightarrow \pi^+\pi^-$  continuum.

NODE=M002R20;LINKAGE=A  
NODE=M002R20;LINKAGE=ALNODE=M002R20;LINKAGE=MO  
NODE=M002R20;LINKAGE=MR $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{32}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-4}$	90	<sup>1</sup> ABLIKIM 11G	BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

<sup>1</sup> ABLIKIM 11G reports  $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.25 \times 10^{-3}$ .

NODE=M002R53  
NODE=M002R53

NODE=M002R53;LINKAGE=AL

 $\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$  $\Gamma_{32}/\Gamma_4$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 45$	90	ALDE 87B	GAM2	$38 \pi^-p \rightarrow n4\gamma$

NODE=M002R36  
NODE=M002R36

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{33}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	BRIERE 00	CLEO	10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<13	90	RITTENBERG 65	HBC	2.7 $K^- p$

NODE=M002R8  
NODE=M002R8 $\Gamma(\pi^0 \rho^0)/\Gamma_{\text{total}}$  $\Gamma_{34}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	RITTENBERG 65	HBC	2.7 $K^- p$

NODE=M002R10  
NODE=M002R10 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4	90	BRIERE 00	CLEO	10.6 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<11	90	RITTENBERG 65	HBC	2.7 $K^- p$

NODE=M002R9  
NODE=M002R9 $\Gamma(3\gamma)/\Gamma(\pi^0 \pi^0 \eta)$  $\Gamma_{36}/\Gamma_4$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.6	90	ALDE 87B	GAM2	38 $\pi^- p \rightarrow n 3\gamma$

NODE=M002R34  
NODE=M002R34 $\Gamma(\mu^+ \mu^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{37}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	90	DZHELADIN 81	CNTR	30 $\pi^- p \rightarrow \eta' n$

NODE=M002R31  
NODE=M002R31 $\Gamma(\mu^+ \mu^- \eta)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	DZHELADIN 81	CNTR	30 $\pi^- p \rightarrow \eta' n$

NODE=M002R30  
NODE=M002R30 $\Gamma(e\mu)/\Gamma_{\text{total}}$  $\Gamma_{39}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.7	90	BRIERE 00	CLEO	10.6 $e^+ e^-$

NODE=M002R41  
NODE=M002R41 $\eta'(958) \rightarrow \eta \pi \pi$  DECAY PARAMETERS

NODE=M002225

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$$

$X$  and  $Y$  are Dalitz variables;  $\alpha$  is complex and  $C$ , and  $D$  are real-valued. Parameters  $C$  and  $D$  are not necessarily equal to  $c$  and  $d$ , respectively, in the generalized parameterization following this one. May be different for  $\eta'(958) \rightarrow \eta \pi^+ \pi^-$  and  $\eta'(958) \rightarrow \eta \pi^0 \pi^0$  decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

NODE=M002225

**Re( $\alpha$ ) decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.034 \pm 0.002 \pm 0.002$	351k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.054 \pm 0.004 \pm 0.001$	56k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.033 \pm 0.005 \pm 0.003$	44k	<sup>1</sup> ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.072 \pm 0.012 \pm 0.006$	7k	<sup>2</sup> AMELIN 05A	VES	$28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$
$-0.021 \pm 0.018 \pm 0.017$	6.7k	<sup>3</sup> BRIERE 00	CLEO	$10.6 e^+ e^- \rightarrow \eta \pi^+ \pi^- X$
$-0.058 \pm 0.013 \pm 0.003$	5.4k	<sup>4</sup> ALDE 86	GAM2	$38 \pi^- p \rightarrow n \eta \pi^0 \pi^0$
$-0.08 \pm 0.03$		<sup>4,5</sup> KALBFLEISCH 74	RVUE	$\eta' \rightarrow \eta \pi^+ \pi^-$

NODE=M002A0  
NODE=M002A0

OCCUR=2

<sup>1</sup> See ABLIKIM 11 for the full correlation matrix.<sup>2</sup> Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.<sup>3</sup> Assuming  $\text{Im}(\alpha) = 0$ ,  $C = 0$ , and  $D = 0$ .<sup>4</sup> Assuming  $C = 0$ .<sup>5</sup> From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.NODE=M002A0;LINKAGE=AB  
NODE=M002A0;LINKAGE=AMNODE=M002A0;LINKAGE=BR  
NODE=M002A0;LINKAGE=A  
NODE=M002A0;LINKAGE=KA



**$Im(\alpha)$  decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.000 \pm 0.019 \pm 0.001$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$0.000 \pm 0.038 \pm 0.002$	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^0 \pi^0$
$0.000 \pm 0.049 \pm 0.001$	44k	<sup>1</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.0 \pm 0.1 \pm 0.0$	7k	<sup>2</sup> AMELIN	05A	VES $28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$
$-0.00 \pm 0.13 \pm 0.00$	5.4k	<sup>3</sup> ALDE	86	GAM2 $38 \pi^- p \rightarrow n \eta \pi^0 \pi^0$
$0.0 \pm 0.3$		<sup>3,4</sup> KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta \pi^+ \pi^-$

<sup>1</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>2</sup> Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

<sup>3</sup> Assuming  $C = 0$ .

<sup>4</sup> From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002IA0  
NODE=M002IA0

OCCUR=2

NODE=M002IA0;LINKAGE=AB  
NODE=M002IA0;LINKAGE=AM

NODE=M002IA0;LINKAGE=A  
NODE=M002IA0;LINKAGE=KA

**C decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.0027 \pm 0.0024 \pm 0.0015$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$0.018 \pm 0.009 \pm 0.003$	44k	<sup>1</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.020 \pm 0.018 \pm 0.004$	7k	<sup>2</sup> AMELIN	05A	VES $28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$

<sup>1</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>2</sup> Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

NODE=M002C0  
NODE=M002C0

NODE=M002C0;LINKAGE=AB  
NODE=M002C0;LINKAGE=AM

**D decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.053 \pm 0.004 \pm 0.004$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.061 \pm 0.009 \pm 0.005$	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.059 \pm 0.012 \pm 0.004$	44k	<sup>1</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.066 \pm 0.030 \pm 0.015$	7k	<sup>2</sup> AMELIN	05A	VES $28 \pi^- A \rightarrow \eta \pi^+ \pi^- \pi^- A^*$
$0.00 \pm 0.03 \pm 0.00$	5.4k	<sup>3</sup> ALDE	86	GAM2 $38 \pi^- p \rightarrow n \eta \pi^0 \pi^0$
0		<sup>3,4</sup> KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta \pi^+ \pi^-$

<sup>1</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>2</sup> Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

<sup>3</sup> Assuming  $C = 0$ .

<sup>4</sup> From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002D0  
NODE=M002D0

OCCUR=2

NODE=M002D0;LINKAGE=AB  
NODE=M002D0;LINKAGE=AM

NODE=M002D0;LINKAGE=AL  
NODE=M002D0;LINKAGE=KA

 **$\eta'(958) \rightarrow \eta \pi \pi$  DECAY PARAMETERS**

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + a Y + b Y^2 + c X + d X^2$$

X and Y are Dalitz variables and  $a$ ,  $b$ ,  $c$ , and  $d$  are real-valued parameters. May be different for  $\eta'(958) \rightarrow \eta \pi^+ \pi^-$  and  $\eta'(958) \rightarrow \eta \pi^0 \pi^0$  decays. We do not average measurements in the section below because parameter values from each experiment are strongly correlated.

NODE=M002227

NODE=M002227

**a decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.056 \pm 0.004 \pm 0.002$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.087 \pm 0.009 \pm 0.006$	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.074 \pm 0.008 \pm 0.006$	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.072 \pm 0.007 \pm 0.008$		<sup>1</sup> GONZALEZ-S.	18A	RVUE $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.047 \pm 0.011 \pm 0.003$	44k	<sup>2</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.066 \pm 0.016 \pm 0.003$	15k	<sup>3</sup> BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
$-0.127 \pm 0.016 \pm 0.008$	20k	<sup>4</sup> DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

<sup>1</sup> Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

<sup>2</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>3</sup> From  $\eta' \rightarrow \eta \pi^0 \pi^0$  decay.

<sup>4</sup> From  $\eta' \rightarrow \eta \pi^+ \pi^-$  decay.

NODE=M002DPA  
NODE=M002DPA

OCCUR=2

NODE=M002DPA;LINKAGE=A

NODE=M002DPA;LINKAGE=AB  
NODE=M002DPA;LINKAGE=BL  
NODE=M002DPA;LINKAGE=DO

**b decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.049 \pm 0.006 \pm 0.006$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.073 \pm 0.014 \pm 0.005$	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.063 \pm 0.014 \pm 0.005$	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.052 \pm 0.001 \pm 0.002$		<sup>1</sup> GONZALEZ-S...	18A	RVUE $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.069 \pm 0.019 \pm 0.009$	44k	<sup>2</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.063 \pm 0.028 \pm 0.004$	15k	<sup>3</sup> BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
$-0.106 \pm 0.028 \pm 0.014$	20k	<sup>4</sup> DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

<sup>1</sup> Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

<sup>2</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>3</sup> From  $\eta' \rightarrow \eta \pi^0 \pi^0$  decay.

<sup>4</sup> From  $\eta' \rightarrow \eta \pi^+ \pi^-$  decay.

NODE=M002DPB  
NODE=M002DPB

OCCUR=2

NODE=M002DPB;LINKAGE=A

NODE=M002DPB;LINKAGE=AB

NODE=M002DPB;LINKAGE=BL

NODE=M002DPB;LINKAGE=DO

**c decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.0027 \pm 0.0024 \pm 0.0018$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$0.019 \pm 0.011 \pm 0.003$	44k	<sup>1</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.107 \pm 0.096 \pm 0.003$	15k	<sup>2</sup> BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
$0.015 \pm 0.011 \pm 0.014$	20k	<sup>3</sup> DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

<sup>1</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>2</sup> From  $\eta' \rightarrow \eta \pi^0 \pi^0$  decay.

<sup>3</sup> From  $\eta' \rightarrow \eta \pi^+ \pi^-$  decay.

NODE=M002DPC  
NODE=M002DPC

NODE=M002DPC;LINKAGE=AB

NODE=M002DPC;LINKAGE=BL

NODE=M002DPC;LINKAGE=DO

**d decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.063 \pm 0.004 \pm 0.003$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.074 \pm 0.009 \pm 0.004$	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.050 \pm 0.009 \pm 0.005$	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.051 \pm 0.008 \pm 0.006$		<sup>1</sup> GONZALEZ-S...	18A	RVUE $\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.073 \pm 0.012 \pm 0.003$	44k	<sup>2</sup> ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.018 \pm 0.078 \pm 0.006$	15k	<sup>3</sup> BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
$-0.082 \pm 0.017 \pm 0.008$	20k	<sup>4</sup> DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

<sup>1</sup> Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

<sup>2</sup> See ABLIKIM 11 for the full correlation matrix.

<sup>3</sup> From  $\eta' \rightarrow \eta \pi^0 \pi^0$  decay. If  $c \equiv 0$  from Bose-Einstein symmetry,  $d = -0.067 \pm 0.020 \pm 0.003$ .

<sup>4</sup> From  $\eta' \rightarrow \eta \pi^+ \pi^-$  decay.

NODE=M002DPD  
NODE=M002DPD

OCCUR=2

NODE=M002DPD;LINKAGE=A

NODE=M002DPD;LINKAGE=AB

NODE=M002DPD;LINKAGE=BL

NODE=M002DPD;LINKAGE=DO

$$\eta'(958) \beta \text{ PARAMETER}$$

$$|\text{MATRIX ELEMENT}|^2 = (1 + 2\beta Z)$$

See the "Note on  $\eta$  Decay Parameters" in our 1994 edition Physical Review D50 1173 (1994), p. 1454.

NODE=M002226

NODE=M002226

 **$\beta$  decay parameter**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>-0.61 \pm 0.08</math> OUR AVERAGE</b>				Error includes scale factor of 1.2.
$-0.640 \pm 0.046 \pm 0.047$	1.8k	ABLIKIM	15G	BES3 $J/\psi \rightarrow \gamma (\pi^0 \pi^0 \pi^0)$
$-0.59 \pm 0.18$	235	BLIK	08	GAMS $32 \pi^- p \rightarrow \eta' n$
$-0.1 \pm 0.3$		ALDE	87B	GAM2 $38 \pi^- p \rightarrow n 3\pi^0$

NODE=M002B0  
NODE=M002B0

**$\eta'(958)$  C-NONCONSERVING DECAY PARAMETER**

NODE=M002235

See the note on  $\eta$  decay parameters in the Stable Particle Particle Listings for definition of this parameter.

NODE=M002235

**DECAY ASYMMETRY PARAMETER FOR  $\pi^+\pi^-\gamma$** 

NODE=M002A

NODE=M002A

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>-0.03 ± 0.04 OUR AVERAGE</b>				
-0.019 ± 0.056		AIHARA 87	TPC	$2\gamma \rightarrow \pi^+\pi^-\gamma$
-0.069 ± 0.078	295	GRIGORIAN 75	STRC	$2.1 \pi^- p$
0.00 ± 0.10	103	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda \pi^+\pi^-\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.07 ± 0.08	152	RITTENBERG 65	HBC	$2.1-2.7 K^- p$

 **$\eta'(958) \rightarrow \gamma \ell^+ \ell^-$  TRANSITION FORM FACTOR SLOPE**

NODE=M002FFL

NODE=M002FFL

Related to the effective virtual meson mass  $\Lambda$ , via slope  $\approx \Lambda^{-2}$ . See e.g. LANDSBERG 85, eq. (3.8), for a detailed definition.

NODE=M002FFL

VALUE (GeV <sup>-2</sup> )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1.62 ± 0.17 OUR AVERAGE</b>				
1.60 ± 0.17 ± 0.08	864	<sup>1</sup> ABLIKIM 150	BES3	$J/\psi \rightarrow \gamma e^+ e^-$
1.7 ± 0.4	33	<sup>1</sup> VIKTOROV 80		$25,33 \pi^- p \rightarrow 2\mu\gamma$

<sup>1</sup> In the single-pole Ansatz where slope =  $1/(\Lambda^2 + \gamma^2)$  with  $\Lambda$ ,  $\gamma$  being a Breit-Wigner mass, width for the effective contributing vector meson.

NODE=M002FFL;LINKAGE=A

 **$\eta'(958)$  REFERENCES**

NODE=M002

ABLIKIM 22E	PR D105 112010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61639
ABLIKIM 21I	PR D103 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61112
ABLIKIM 21J	PR D103 092005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61114
ABLIKIM 20E	PR D101 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60252
ABLIKIM 19AW	PR D100 052015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60000
ABLIKIM 19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59773
ABLIKIM 18	PR D97 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58711
ABLIKIM 18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
ADLARSON 18A	PR D98 012001	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)	REFID=58957
GONZALEZ-S... 18A	EPJ C78 758	S. Gonzalez-Solis, E. Passemar	(BEIJ, IND+)	REFID=59316
AAJ 17D	PL B764 233	R. Aajj <i>et al.</i>	(LHCb Collab.)	REFID=57702
ABLIKIM 17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57650
ABLIKIM 17T	PR D96 012005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58013
ABLIKIM 16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57511
ABLIKIM 15AD	PR D92 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56983
ABLIKIM 15G	PR D92 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56602
ABLIKIM 15O	PR D92 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56780
ABLIKIM 15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
ACHASOV 15	PR D91 092010	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=56788
AKHMETSHIN 15	PL B740 273	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=56386
PDG 15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=5
ABLIKIM 14M	PRL 112 251801	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55904
DONSKOV 14	MPL A29 1450213	S. Donskov <i>et al.</i>	(GAMS-4 $\pi$ Collab.)	REFID=56321
PDG 14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ABLIKIM 13	PR D87 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54829
ABLIKIM 13G	PR D87 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54952
ABLIKIM 13O	PR D87 092011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55388
ABLIKIM 13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
ABLIKIM 12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
PDG 12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ABLIKIM 11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM 11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53711
CZERWINSKI 10	PRL 105 122001	E. Czerwinski <i>et al.</i>	(COSY-11 Collab.)	REFID=53364
BLIK 09	PAN 72 231	A.M. Blik <i>et al.</i>	(IHEP (Protvino))	REFID=52727
	Translated from YAF 72 258.			
NAIK 09	PRL 102 061801	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52678
PEDLAR 09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
BLIK 08	PAN 71 2124	A. Blik <i>et al.</i>	(GAMS-4 $\pi$ Collab.)	REFID=52663
	Translated from YAF 71 2161.			
LIBBY 08	PRL 101 182002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=52591
WICHT 08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
DOROFEEV 07	PL B651 22	V. Dorofeev <i>et al.</i>	(VES Collab.)	REFID=51711
MORI 07A	JPSJ 76 074102	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51691
ABLIKIM 06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM 06Q	PRL 97 202002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51487
AMELIN 05A	PAN 68 372	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=50766
	Translated from YAF 68 401.			
AMSLER 04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51079
BAI 04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
BRIERE 00	PRL 84 26	R. Briere <i>et al.</i>	(CLEO Collab.)	REFID=47410
ACCIARRI 98Q	PL B418 399	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46316
BARBERIS 98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
WURZINGER 96	PL B374 283	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=44992
PDG 94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)	REFID=43653
AMSLER 93	ZPHY C58 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43311
BELADIDZE 92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)	REFID=43175
	Translated from YAF 55 2748.			

KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=42170
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BEHREND	91	ZPHY C49 401	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41497
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
KARCH	90	PL B249 353	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=41377
ROE	90	PR D41 17	N.A. Roe <i>et al.</i>	(ASP Collab.)	REFID=41014
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)	REFID=40564
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
WILLIAMS	88	Translated from YAF 48 436. PR D38 1365	D.A. Williams <i>et al.</i>	(Crystal Ball Collab.)	REFID=40567
AIHARA	87	PR D35 2650	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.) JP	REFID=40009
ALBRECHT	87B	PL B199 457	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40265
ALDE	87B	ZPHY C36 603	D.M. Alde <i>et al.</i>	(LANL, BELG, SERP, LAPP)	REFID=40236
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ALDE	86	PL B177 115	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=20310
BARTEL	85E	PL 160B 421	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10843
LANDSBERG	85	PRPL 128 301	L.G. Landsberg	(SERP)	REFID=10844
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=20305
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)	REFID=20306
BINON	84	PL 140B 264	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP+)	REFID=20307
JENNI	83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)	REFID=20304
BARTEL	82B	PL 113B 190	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=20300
BEHREND	82C	PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
DZHELYADIN	81	PL 105B 239	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10836
STANTON	80	PL B92 353	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+)	REFID=40294
VIKTOROV	80	SJNP 32 520	V.A. Viktorov <i>et al.</i>	(SERP)	REFID=20298
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)	REFID=20295
BINNIE	79	PL 83B 141	D.M. Binnie <i>et al.</i>	(LOIC)	REFID=20296
ZANFINO	77	PRL 38 930	C. Zanfino <i>et al.</i>	(CARL, MCGI, OHIO+)	REFID=20293
GRIGORIAN	75	NP B91 232	A. Grigorian <i>et al.</i>	(+)	REFID=20287
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
DUANE	74	PRL 32 425	A. Duane <i>et al.</i>	(LOIC, SHMP)	REFID=20284
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)	REFID=20286
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH) JP	REFID=20280
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+)	REFID=20281
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20275
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)	REFID=20278
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20270
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)	REFID=20272
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=20268
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I	REFID=20266
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)	REFID=20263
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJP	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=20253
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)	REFID=10761
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP	REFID=20247
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP	REFID=20252

NODE=M003

 **$f_0(980)$** 

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the related review(s):  
 Scalar Mesons below 1 GeV

 **$f_0(980)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(980–1010) – <math>i</math> (20–35) OUR ESTIMATE</b> (see Fig. 64.4 in the review)			
$(1014 \pm 8) - i(35 \pm 5)$	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(992.8 \pm 1.3) - i(30.7 \pm 2.3)$	1 ALBRECHT 20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
$(1003_{-27}^{+5}) - i(21_{-8}^{+10})$	2 GARCIA-MAR..11	RVUE	Compilation
$(996 \pm 7) - i(25_{-6}^{+10})$	3 GARCIA-MAR..11	RVUE	Compilation
$(996_{-14}^{+4}) - i(24_{-3}^{+11})$	4 MOUSSALLAM11	RVUE	Compilation
$(981 \pm 43) - i(18 \pm 11)$	5 MENNESSIER 10	RVUE	Compilation
$(1030_{-10}^{+30}) - i(35_{-16}^{+10})$	6 ANISOVICH 09	RVUE	$0.0 \bar{p}p, \pi N$
$(973_{-127}^{+39}) - i(11_{-11}^{+189})$	7 PELAEZ 04A	RVUE	$\pi\pi \rightarrow \pi\pi$

<sup>1</sup> 5 poles, 5 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ), and BINON 84C ( $\eta\eta'$ ). Based on 18.5k events. Second solution  $977.8 \pm 1.7$  MeV.

<sup>2</sup> Reanalysis of the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOESCU 73 using Roy equations.

<sup>3</sup> Reanalysis of the  $K_{e4}$  data of BATLEY 10C and the  $\pi N \rightarrow \pi\pi N$  data of HYAMS 73, GRAYER 74, and PROTOPOESCU 73 using GKPY equations.

<sup>4</sup> Uses Roy equations.

<sup>5</sup> Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.

<sup>6</sup> On sheet II in a 2-pole solution. The other pole is found on sheet III at  $(850 - i 100)$  MeV.

<sup>7</sup> Reanalysis of data from PROTOPOESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.

NODE=M003PP

NODE=M003PP

NODE=M003PP

→ UNCHECKED ←

OCCUR=2

NODE=M003PP;LINKAGE=D

NODE=M003PP;LINKAGE=B

NODE=M003PP;LINKAGE=C

NODE=M003PP;LINKAGE=F

NODE=M003PP;LINKAGE=G

NODE=M003PP;LINKAGE=H

NODE=M003PP;LINKAGE=A

 **$f_0(980)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>990 ± 20 OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$992.0_{-7.5}^{+8.5} \pm 8.6$		1 AAIJ 19H	LHCB	$pp \rightarrow D^\pm X$
$989.4 \pm 1.3$	424	ABLIKIM 15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
$989.9 \pm 0.4$	706	ABLIKIM 12E	BES3	$J/\psi \rightarrow \gamma 3\pi$
$977_{-9}^{+11} \pm 1$	44	2 ECKLUND 09	CLEO	$4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + \text{c.c.}$
$982.2 \pm 1.0_{-8.0}^{+8.1}$		3 UEHARA 08A	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$976.8 \pm 0.3_{-0.6}^{+10.1}$	64k	4 AMBROSINO 07	KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$984.7 \pm 0.4_{-3.7}^{+2.4}$	64k	5 AMBROSINO 07	KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$973 \pm 3$	262 ± 30	6 AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
$970 \pm 7$	54 ± 9	6 AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \pi^0 \gamma$
$953 \pm 20$	2.6k	7 BONVICINI 07	CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
$985.6_{-1.5}^{+1.2} \pm 1.1_{-1.6}$		8 MORI 07	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
$983.0 \pm 0.6_{-3.0}^{+4.0}$		9 AMBROSINO 06B	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

NODE=M003M1

NODE=M003M1

→ UNCHECKED ←

OCCUR=2

OCCUR=2

977.3 ± 0.9 <sup>+3.7</sup> <sub>-4.3</sub>		10	AMBROSINO	06B	KLOE	1.02 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> γ	OCCUR=2
950 ± 9	2426	11	GARMASH	06	BELL	B <sup>+</sup> → K <sup>+</sup> π <sup>+</sup> π <sup>-</sup>	
965 ± 10		12	ABLIKIM	05	BES2	J/ψ → φπ <sup>+</sup> π <sup>-</sup> , φK <sup>+</sup> K <sup>-</sup>	
1031 ± 8		13	ANISOVICH	03	RVUE		
1037 ± 31			TIKHOMIROV	03	SPEC	40.0 π <sup>-</sup> C → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> K <sub>L</sub> <sup>0</sup> X	
973 ± 1	2438	14	ALOISIO	02D	KLOE	e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ	
977 ± 3 ± 2	848	15	AITALA	01A	E791	D <sub>s</sub> <sup>+</sup> → π <sup>-</sup> π <sup>+</sup> π <sup>+</sup>	
969.8 ± 4.5	419	16	ACHASOV	00H	SND	e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ	
985 <sup>+16</sup> <sub>-12</sub>	419	17,18	ACHASOV	00H	SND	e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ	OCCUR=2
976 ± 5 ± 6		19	AKHMETSHIN	99B	CMD2	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> γ	
977 ± 3 ± 6	268	19	AKHMETSHIN	99C	CMD2	e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ	
975 ± 4 ± 6		20	AKHMETSHIN	99C	CMD2	e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ	OCCUR=2
975 ± 4 ± 6		21	AKHMETSHIN	99C	CMD2	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> γ, π <sup>0</sup> π <sup>0</sup> γ	OCCUR=3
985 ± 10			BARBERIS	99	OMEG	450 pp → p <sub>S</sub> p <sub>f</sub> K <sup>+</sup> K <sup>-</sup>	
982 ± 3			BARBERIS	99B	OMEG	450 pp → p <sub>S</sub> p <sub>f</sub> π <sup>+</sup> π <sup>-</sup>	
982 ± 3			BARBERIS	99C	OMEG	450 pp → p <sub>S</sub> p <sub>f</sub> π <sup>0</sup> π <sup>0</sup>	
987 ± 6 ± 6		22	BARBERIS	99D	OMEG	450 pp → K <sup>+</sup> K <sup>-</sup> , π <sup>+</sup> π <sup>-</sup>	
989 ± 15			BELLAZZINI	99	GAM4	450 pp → ppπ <sup>0</sup> π <sup>0</sup>	
991 ± 3		23	KAMINSKI	99	RVUE	ππ → ππ, K $\bar{K}$ , σσ	
~ 980		23	OLLER	99	RVUE	ππ → ππ, K $\bar{K}$	
~ 993.5			OLLER	99B	RVUE	ππ → ππ, K $\bar{K}$	
~ 987		23	OLLER	99C	RVUE	ππ → ππ, K $\bar{K}$ , ηη	
957 ± 6		24	ACKERSTAFF	98Q	OPAL	Z → f <sub>0</sub> X	
960 ± 10			ALDE	98	GAM4		
1015 ± 15		23	ANISOVICH	98B	RVUE	Compilation	
1008		25	LOCHER	98	RVUE	ππ → ππ, K $\bar{K}$	
955 ± 10		24	ALDE	97	GAM2	450 pp → ppπ <sup>0</sup> π <sup>0</sup>	
994 ± 9		26	BERTIN	97C	OBLX	0.0 p̄p → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup>	
993.2 ± 6.5 ± 6.9		27	ISHIDA	96	RVUE	ππ → ππ, K $\bar{K}$	
1006			TORNQVIST	96	RVUE	ππ → ππ, K $\bar{K}$ , Kπ, ηπ	
997 ± 5	3k	28	ALDE	95B	GAM2	38 π <sup>-</sup> p → π <sup>0</sup> π <sup>0</sup> n	
960 ± 10	10k	29	ALDE	95B	GAM2	38 π <sup>-</sup> p → π <sup>0</sup> π <sup>0</sup> n	OCCUR=2
994 ± 5			AMSLER	95B	CBAR	0.0 p̄p → 3π <sup>0</sup>	
~ 996		30	AMSLER	95D	CBAR	0.0 p̄p → π <sup>0</sup> π <sup>0</sup> π <sup>0</sup> , π <sup>0</sup> ηη, π <sup>0</sup> π <sup>0</sup> η	
987 ± 6		31	ANISOVICH	95	RVUE		
1015			JANSSEN	95	RVUE	ππ → ππ, K $\bar{K}$	
983		32	BUGG	94	RVUE	p̄p → η2π <sup>0</sup>	
973 ± 2		33	KAMINSKI	94	RVUE	ππ → ππ, K $\bar{K}$	
988		34	ZOU	94B	RVUE		
988 ± 10		35	MORGAN	93	RVUE	ππ(K $\bar{K}$ ) → ππ(K $\bar{K}$ ), J/ψ → φππ(K $\bar{K}$ ), D <sub>S</sub> → π(ππ)	
971.1 ± 4.0		24	AGUILAR-...	91	EHS	400 pp	
979 ± 4		36	ARMSTRONG	91	OMEG	300 pp → ppππ, ppK $\bar{K}$	
956 ± 12			BREAKSTONE	90	SFM	pp → ppπ <sup>+</sup> π <sup>-</sup>	
959.4 ± 6.5		24	AUGUSTIN	89	DM2	J/ψ → ωπ <sup>+</sup> π <sup>-</sup>	
978 ± 9		24	ABACHI	86B	HRS	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> X	
985.0 <sup>+9.0</sup> <sub>-39.0</sub>			ETKIN	82B	MPS	23 π <sup>-</sup> p → n2K <sub>S</sub> <sup>0</sup>	
974 ± 4		36	GIDAL	81	MRK2	J/ψ → π <sup>+</sup> π <sup>-</sup> X	
975		37	ACHASOV	80	RVUE		
986 ± 10		36	AGUILAR-...	78	HBC	0.7 p̄p → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup>	
969 ± 5		36	LEEPER	77	ASPK	2-2.4 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> n, K <sup>+</sup> K <sup>-</sup> n	
987 ± 7		36	BINNIE	73	CNTR	π <sup>-</sup> p → nMM	
1012 ± 6		38	GRAYER	73	ASPK	17 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> n	
1007 ± 20		38	HYAMS	73	ASPK	17 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> n	
997 ± 6		38	PROTOPOP...	73	HBC	7 π <sup>+</sup> p → π <sup>+</sup> pπ <sup>+</sup> π <sup>-</sup>	

- 1 From the  $D^\pm \rightarrow K^\pm K^+ K^-$  Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.
- 2 Using a relativistic Breit-Wigner function and taking into account the finite  $D_s$  mass.
- 3 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K K / g_{f_0} \pi \pi = 0$ .
- 4 In the kaon-loop fit.
- 5 In the no-structure fit.
- 6 Systematic errors not estimated.
- 7 FLATTE 76 parameterization.  $g_{f_0} \pi \pi = 329 \pm 96 \text{ MeV}/c^2$  assuming  $g_{f_0} K \bar{K} / g_{f_0} \pi \pi = 2$ .
- 8 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0} K K / g_{f_0} \pi \pi = 4.21 \pm 0.25 \pm 0.21$  from ABLIKIM 05.
- 9 In the kaon-loop fit following formalism of ACHASOV 89.
- 10 In the no-structure fit assuming a direct coupling of  $\phi$  to  $f_0 \gamma$ .
- 11 FLATTE 76 parameterization. Supersedes GARMASH 05.
- 12 FLATTE 76 parameterization,  $g_{f_0} K \bar{K} / g_{f_0} \pi \pi = 4.21 \pm 0.25 \pm 0.21$ .
- 13 K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.
- 14 From the negative interference with the  $f_0(500)$  meson of AITALA 01B using the ACHASOV 89 parameterization for the  $f_0(980)$ , a Breit-Wigner for the  $f_0(500)$ , and ACHASOV 01F for the  $\rho\pi$  contribution.
- 15 Coupled-channel Breit-Wigner, couplings  $g_\pi = 0.09 \pm 0.01 \pm 0.01$ ,  $g_K = 0.02 \pm 0.04 \pm 0.03$ .
- 16 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
- 17 Supersedes ACHASOV 98I.
- 18 In the "narrow resonance" approximation.
- 19 Assuming  $\Gamma(f_0) = 40 \text{ MeV}$ .
- 20 From a narrow pole fit taking into account  $f_0(980)$  and  $f_0(1200)$  intermediate mechanisms.
- 21 From the combined fit of the photon spectra in the reactions  $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$ ,  $\pi^0 \pi^0 \gamma$ .
- 22 Supersedes BARBERIS 99 and BARBERIS 99B
- 23 T-matrix pole.
- 24 From invariant mass fit.
- 25 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039–93i) MeV.
- 26 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963–29i) MeV.
- 27 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 28 At high  $|t|$ .
- 29 At low  $|t|$ .
- 30 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953–55i) MeV and on sheet IV at (938–35i) MeV.
- 31 Combined fit of ALDE 95B, ANISOVICH 94, AMSLER 94D.
- 32 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996–103i) MeV.
- 33 From sheet II pole position.
- 34 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797–185i) MeV and can be interpreted as a shadow pole.
- 35 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978–28i) MeV.
- 36 From coupled channel analysis.
- 37 Coupled channel analysis with finite width corrections.
- 38 Included in AGUILAR-BENITEZ 78 fit.

NODE=M003M1;LINKAGE=F

NODE=M003M1;LINKAGE=EC  
NODE=M003M1;LINKAGE=UE

NODE=M003M1;LINKAGE=AK  
NODE=M003M1;LINKAGE=AS  
NODE=M003M1;LINKAGE=NS

NODE=M003M1;LINKAGE=BO  
NODE=M003M1;LINKAGE=MO

NODE=M003M1;LINKAGE=AB  
NODE=M003M1;LINKAGE=AM  
NODE=M003M1;LINKAGE=GR

NODE=M003M1;LINKAGE=AL  
NODE=M003M1;LINKAGE=KM

NODE=M003M1;LINKAGE=KD

NODE=M003M1;LINKAGE=TL  
NODE=M003M1;LINKAGE=V9  
NODE=M003M1;LINKAGE=V8  
NODE=M003M1;LINKAGE=AI  
NODE=M003M1;LINKAGE=SM  
NODE=M003M1;LINKAGE=ST

NODE=M003M1;LINKAGE=SL

NODE=M003M1;LINKAGE=BD  
NODE=M003M1;LINKAGE=AN  
NODE=M003M1;LINKAGE=A  
NODE=M003M1;LINKAGE=LC  
NODE=M003M1;LINKAGE=X  
NODE=M003M1;LINKAGE=AA

NODE=M003M1;LINKAGE=LA  
NODE=M003M1;LINKAGE=LB  
NODE=M003M1;LINKAGE=KL

NODE=M003M1;LINKAGE=CF  
NODE=M003M1;LINKAGE=C2  
NODE=M003M1;LINKAGE=KM  
NODE=M003M1;LINKAGE=L

NODE=M003M1;LINKAGE=K  
NODE=M003M1;LINKAGE=B  
NODE=M003M1;LINKAGE=B  
NODE=M003M1;LINKAGE=R

## $f_0(980)$ WIDTH

Width determination very model dependent. Peak width in  $\pi\pi$  is about 50 MeV, but decay width can be much larger.

NODE=M003W1

NODE=M003W1

NODE=M003W1

→ UNCHECKED ←

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10 to 100 OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$15.3 \pm 4.7$	424	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
$9.5 \pm 1.1$	706	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma 3\pi$
$91 \begin{smallmatrix} + 30 \\ - 22 \end{smallmatrix} \pm 3$	44	1 ECKLUND	09 CLEO	$4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + \text{c.c.}$
$66.9 \pm 2.2 \begin{smallmatrix} + 17.6 \\ - 12.5 \end{smallmatrix}$		2 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$65 \pm 13$	$262 \pm 30$	3 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$

81 ± 21	54 ± 9	3	AUBERT	07AK	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\pi^0\gamma$	OCCUR=2
51.3 <sup>+</sup> <sub>-</sub> 20.8 <sup>+</sup> <sub>-</sub> 13.2 <sup>+</sup> <sub>-</sub> 17.7 <sup>-</sup> <sub>-</sub> 3.8		4	MORI	07	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$	
61 ± 9 <sup>+14</sup> <sub>-8</sub>	2584	5	GARMASH	05	BELL	$B^+ \rightarrow K^+\pi^+\pi^-$	
64 ± 16		6	ANISOVICH	03	RVUE		
121 ± 23			TIKHOMIROV	03	SPEC	40.0 $\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$	
~ 70		7	BRAMON	02	RVUE	1.02 $e^+e^- \rightarrow \pi^0\pi^0\gamma$	
44 ± 2 ± 2	848	8	AITALA	01A	E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$	
201 ± 28	419	9	ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2
122 ± 13	419	10,11	ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
56 ± 20		12	AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
65 ± 20			BARBERIS	99	OMEG	450 $pp \rightarrow p_s p_f K^+ K^-$	
80 ± 10			BARBERIS	99B	OMEG	450 $pp \rightarrow p_s p_f \pi^+ \pi^-$	
80 ± 10			BARBERIS	99C	OMEG	450 $pp \rightarrow p_s p_f \pi^0 \pi^0$	
48 ± 12 ± 8		13	BARBERIS	99D	OMEG	450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$	
65 ± 25			BELLAZZINI	99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$	
71 ± 14		14	KAMINSKI	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$	
~ 28		14	OLLER	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~ 25			OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~ 14		14	OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$	
70 ± 20			ALDE	98	GAM4		
86 ± 16		14	ANISOVICH	98B	RVUE	Compilation	
54		15	LOCHER	98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
69 ± 15		16	ALDE	97	GAM2	450 $pp \rightarrow pp\pi^0\pi^0$	
38 ± 20		17	BERTIN	97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
~ 100		18	ISHIDA	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
34			TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$	
48 ± 10	3k	19	ALDE	95B	GAM2	38 $\pi^-p \rightarrow \pi^0\pi^0n$	OCCUR=2
95 ± 20	10k	20	ALDE	95B	GAM2	38 $\pi^-p \rightarrow \pi^0\pi^0n$	
26 ± 10			AMSLER	95B	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$	
~ 112		21	AMSLER	95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$	
80 ± 12		22	ANISOVICH	95	RVUE		
30			JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
74		23	BUGG	94	RVUE	$\bar{p}p \rightarrow \eta 2\pi^0$	
29 ± 2		24	KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
46		25	ZOU	94B	RVUE		
48 ± 12		26	MORGAN	93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_s \rightarrow \pi(\pi\pi)$	
37.4 ± 10.6		16	AGUILAR...	91	EHS	400 $pp$	
72 ± 8		27	ARMSTRONG	91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$	
110 ± 30			BREAKSTONE	90	SFM	$pp \rightarrow pp\pi^+\pi^-$	
29 ± 13		16	ABACHI	86B	HRS	$e^+e^- \rightarrow \pi^+\pi^-X$	
120 ± 281 ± 20			ETKIN	82B	MPS	23 $\pi^-p \rightarrow n 2K_S^0$	
28 ± 10		27	GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+\pi^-X$	
70 to 300		28	ACHASOV	80	RVUE		
100 ± 80		29	AGUILAR...	78	HBC	0.7 $\bar{p}p \rightarrow K_S^0 K_S^0$	
30 ± 8		27	LEEPER	77	ASPK	2-2.4 $\pi^-p \rightarrow \pi^+\pi^-n, K^+K^-n$	
48 ± 14		27	BINNIE	73	CNTR	$\pi^-p \rightarrow nMM$	
32 ± 10		30	GRAYER	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$	
30 ± 10		30	HYAMS	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$	
54 ± 16		30	PROTOPOP...	73	HBC	7 $\pi^+p \rightarrow \pi^+p\pi^+\pi^-$	



- 1 Using a relativistic Breit-Wigner function and taking into account the finite  $D_S$  mass. NODE=M003W1;LINKAGE=EC  
NODE=M003W1;LINKAGE=UE
- 2 Breit-Wigner  $\pi\pi$  width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0}^2 K K / g_{f_0}^2 \pi\pi = 0$ .
- 3 Systematic errors not estimated. NODE=M003W1;LINKAGE=NS  
NODE=M003W1;LINKAGE=MO
- 4 Breit-Wigner  $\pi\pi$  width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0}^2 K K / g_{f_0}^2 \pi\pi = 4.21 \pm 0.25 \pm 0.21$  from ABLIKIM 05.
- 5 Breit-Wigner, solution 1, PWA ambiguous. NODE=M003W1;LINKAGE=GA  
NODE=M003W1;LINKAGE=KM
- 6 K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.
- 7 Using the data of AKHMETSHIN 99C, ACHASOV 00H, and ALOISIO 02D. NODE=M003W;LINKAGE=BR  
NODE=M003W;LINKAGE=TL  
NODE=M003W;LINKAGE=V9  
NODE=M003W;LINKAGE=V8  
NODE=M003W1;LINKAGE=AI  
NODE=M003W;LINKAGE=SL
- 8 Breit-Wigner width.
- 9 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
- 10 Supersedes ACHASOV 98I.
- 11 In the "narrow resonance" approximation.
- 12 From the combined fit of the photon spectra in the reactions  $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$ ,  $\pi^0 \pi^0 \gamma$ .
- 13 Supersedes BARBERIS 99 and BARBERIS 99B. NODE=M003W1;LINKAGE=BD  
NODE=M003W1;LINKAGE=AN  
NODE=M003W1;LINKAGE=LO  
NODE=M003W1;LINKAGE=A  
NODE=M003W1;LINKAGE=X  
NODE=M003W1;LINKAGE=AA
- 14 T-matrix pole.
- 15 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039–93*i*) MeV.
- 16 From invariant mass fit.
- 17 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963–29*i*) MeV.
- 18 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 19 At high  $|t|$ . NODE=M003W1;LINKAGE=LA  
NODE=M003W1;LINKAGE=LB  
NODE=M003W1;LINKAGE=KL
- 20 At low  $|t|$ .
- 21 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953–55*i*) MeV and on sheet IV at (938–35*i*) MeV.
- 22 Combined fit of ALDE 95B, ANISOVICH 94.
- 23 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996–103*i*) MeV.
- 24 From sheet II pole position.
- 25 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797–185*i*) MeV and can be interpreted as a shadow pole.
- 26 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978–28*i*) MeV. NODE=M003W1;LINKAGE=CF  
NODE=M003W1;LINKAGE=C2  
NODE=M003W1;LINKAGE=KM  
NODE=M003W1;LINKAGE=L
- 27 From coupled channel analysis. NODE=M003W1;LINKAGE=K  
NODE=M003W1;LINKAGE=B  
NODE=M003W;LINKAGE=B  
NODE=M003W;LINKAGE=C
- 28 Coupled channel analysis with finite width corrections.
- 29 From coupled channel fit to the HYAMS 73 and PROTOPOESCU 73 data. With a simultaneous fit to the  $\pi\pi$  phase-shifts, inelasticity and to the  $K_S^0 K_S^0$  invariant mass.
- 30 Included in AGUILAR-BENITEZ 78 fit. NODE=M003W;LINKAGE=R

### $f_0(980)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $K\bar{K}$	seen
$\Gamma_3$ $\gamma\gamma$	seen
$\Gamma_4$ $e^+ e^-$	

NODE=M003215;NODE=M003

DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=5;OUR EVAL;→ UNCHECKED ←  
DESIG=4

### $f_0(980)$ PARTIAL WIDTHS

NODE=M003220

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_3$
------------------------	-------------	-------------	------	---------	------------

NODE=M003W4  
NODE=M003W4

#### 0.29 $^{+0.11}_{-0.06}$ OUR AVERAGE

0.286 $\pm 0.017$ $^{+0.211}_{-0.070}$	1	UEHARA	08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
0.205 $^{+0.095+0.147}_{-0.083-0.117}$	2	MORI	07	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
0.42 $\pm 0.06 \pm 0.18$	3	OEST	90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32 ±0.05	4 DAI	14A	RVUE	Compilation
0.16 ±0.01	5 MENNESSIER	11	RVUE	
0.29 ±0.21 $\begin{smallmatrix} +0.02 \\ -0.07 \end{smallmatrix}$	6 MOUSSALLAM	11	RVUE	Compilation
0.42	7,8 PENNINGTON	08	RVUE	Compilation
0.10	8,9 PENNINGTON	08	RVUE	Compilation
0.28 $\begin{smallmatrix} +0.09 \\ -0.13 \end{smallmatrix}$	10 BOGLIONE	99	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$
0.29 ±0.07 ±0.12	11,12 BOYER	90	MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
0.31 ±0.14 ±0.09	11,12 MARSISKE	90	CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
0.63 ±0.14	13 MORGAN	90	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$

OCCUR=2

<sup>1</sup> Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0 KK}/g_{f_0 \pi\pi} = 0$ .

NODE=M003W4;LINKAGE=UE

<sup>2</sup> Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio  $g_{f_0 KK}/g_{f_0 \pi\pi} = 4.21 \pm 0.25 \pm 0.21$  from ABLIKIM 05.

NODE=M003W4;LINKAGE=MO

<sup>3</sup> OEST 90 quote systematic errors  $\begin{smallmatrix} +0.08 \\ -0.18 \end{smallmatrix}$ . We use ±0.18. Observed 60 events.

NODE=M003W4;LINKAGE=H

<sup>4</sup> Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.

NODE=M003W4;LINKAGE=D

<sup>5</sup> Uses an analytic K-matrix model. Compilation.

NODE=M003W4;LINKAGE=ME

<sup>6</sup> Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.

NODE=M003W4;LINKAGE=MU

<sup>7</sup> Solution A (preferred solution based on  $\chi^2$ -analysis).

NODE=M003W4;LINKAGE=P1

<sup>8</sup> Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

NODE=M003W4;LINKAGE=P3

<sup>9</sup> Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

NODE=M003W4;LINKAGE=P2

<sup>10</sup> Supersedes MORGAN 90.

NODE=M003W4;LINKAGE=BL

<sup>11</sup> From analysis allowing arbitrary background unconstrained by unitarity.

NODE=M003W4;LINKAGE=B

<sup>12</sup> Data included in MORGAN 90, BOGLIONE 99 analyses.

NODE=M003W4;LINKAGE=C

<sup>13</sup> From amplitude analysis of BOYER 90 and MARSISKE 90, data corresponds to resonance parameters  $m = 989$  MeV,  $\Gamma = 61$  MeV.

NODE=M003W4;LINKAGE=A

## $\Gamma(e^+e^-)$

$\Gamma_4$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;8.4</b>	90	VOROBYEV	88 ND	$e^+e^- \rightarrow \pi^0\pi^0$

NODE=M003W3  
NODE=M003W3

## $f_0(980)$ BRANCHING RATIOS

NODE=M003225

### $\Gamma(\pi\pi)/[\Gamma(\pi\pi) + \Gamma(K\bar{K})]$

$\Gamma_1/(\Gamma_1+\Gamma_2)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M003R1  
NODE=M003R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.52 ±0.12	9.9k	1 AUBERT	060	BABR	$B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$
0.75 $\begin{smallmatrix} +0.11 \\ -0.13 \end{smallmatrix}$		2 ABLIKIM	05Q	BES2	$\chi_{c0} \rightarrow 2\pi^+ 2\pi^-,$ $\pi^+\pi^- K^+ K^-$
0.84 ±0.02		3 ANISOVICH	02D	SPEC	Combined fit
~ 0.68		OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
0.67 ±0.09		4 LOVERRE	80	HBC	$4 \pi^- p \rightarrow n 2K_S^0$
0.81 $\begin{smallmatrix} +0.09 \\ -0.04 \end{smallmatrix}$		4 CASON	78	STRC	$7 \pi^- p \rightarrow n 2K_S^0$
0.78 ±0.03		4 WETZEL	76	OSPK	$8.9 \pi^- p \rightarrow n 2K_S^0$

<sup>1</sup> Recalculated by us using  $\Gamma(K^+K^-)/\Gamma(\pi^+\pi^-) = 0.69 \pm 0.32$  from AUBERT 060 and isospin relations.

NODE=M003R1;LINKAGE=AU

<sup>2</sup> Using data from ABLIKIM 04G.

NODE=M003R1;LINKAGE=AB

<sup>3</sup> From a combined K-matrix analysis of Crystal Barrel ( $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

NODE=M003R;LINKAGE=CH

<sup>4</sup> Measure  $\pi\pi$  elasticity assuming two resonances coupled to the  $\pi\pi$  and  $K\bar{K}$  channels only.

NODE=M003R1;LINKAGE=B

f<sub>0</sub>(980) REFERENCES

NODE=M003

SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
AAJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)	REFID=55923
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
GARCIA-MAR...	11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=16761
GARCIA-MAR...	11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=54121
MENNESSIER	11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang		REFID=53637
MOUSSALLAM	11	EPJ C71 1814	B. Moussallam		REFID=53975
BATLEY	10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	REFID=53567
MENNESSIER	10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang		REFID=53657
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
ECKLUND	09	PR D80 052009	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=53047
BATLEY	08A	EPJ C54 411	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	REFID=52487
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>		REFID=52303
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51616
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=51721
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51652
AMBROSINO	06B	PL B634 148	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51043
AUBERT	06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51141
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51162
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ACHASOV	05	PR D72 013006	N.N. Achasov, G.N. Shestakov		REFID=50762
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50187
BUETTIKER	04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam		REFID=56428; ERROR=6
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			
BRAMON	02	EPJ C26 253	A. Bramon <i>et al.</i>		REFID=49178
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)	REFID=48312
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
ACHASOV	00B	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	99H	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99C	PL B453 325	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46923
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>		REFID=46600
ACKERSTAFF	98Q	EPJ C4 19	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46145
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)	REFID=46372
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ISHIDA	96	PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45770
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=44375
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)	REFID=44442
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADL, JULI)	REFID=44508
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)	REFID=44072
MORGAN	93	PR D48 1185	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=43614
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
AGUILAR...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=21418
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ACHASOV	80	SJNP 32 566	N.N. Achasov, S.A. Devyanin, G.N. Shestakov	(NOVM)	REFID=20458
		Translated from YAF 32 1098.			
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP	REFID=20381
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+) IJP	REFID=20382
AGUILAR...	78	NP B140 73	M. Aguilar-Benitez <i>et al.</i>	(MADR, BOMB+)	REFID=20368
CASON	78	PRL 41 271	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20370
LEEPER	77	PR D16 2054	R.J. Leeper <i>et al.</i>	(ISU)	REFID=20365
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SAFL)	REFID=11004
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)	REFID=20446
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)	REFID=20362
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)	REFID=21062
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYR	74	NP B75 189	G. Grayr <i>et al.</i>	(CERN, MPIM)	REFID=20113
BINNIE	73	PRL 31 1534	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20343
GRAYR	73	Tallahassee	G. Grayr <i>et al.</i>	(CERN, MPIM)	REFID=20347
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108

**$a_0(980)$** 

$$I^G(J^{PC}) = 1^-(0^{++})$$

See the related review(s):  
 Scalar Mesons below 1 GeV

 **$a_0(980)$  T-MATRIX POLE  $\sqrt{s}$** 

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(960-1030) - <math>i</math> (20-70) OUR ESTIMATE</b> (see Fig. 64.2 in the review)			
$(989 \pm 5) - i(40 \pm 5)$	<sup>1</sup> BUGG	08A	RVUE $\bar{p}p$ annihilation data
$(1117_{-320}^{+24}) - i(12_{-12}^{+43})$	<sup>2</sup> PELAEZ	04A	RVUE $\pi\pi \rightarrow \pi\pi, \pi K \rightarrow \pi K$
$(982 \pm 3) - i(46 \pm 4)$	<sup>3</sup> ABELE	98	CBAR $0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
<sup>1</sup> T-matrix pole on sheet II. Parameterizes couplings to $\bar{K}K, \pi\eta,$ and $\pi\eta'$ . Uses AMSLER 94D and ABELE 98.			
<sup>2</sup> Reanalysis of data from LINGLIN 73, ESTABROOKS 78, and ASTON 88 in the unitarized ChPT model.			
<sup>3</sup> T-matrix pole on sheet II; the pole on sheet III is at $(1006 - i 49)$ MeV.			

NODE=M036PP

NODE=M036PP

NODE=M036PP

→ UNCHECKED ←

NODE=M036PP;LINKAGE=C

NODE=M036PP;LINKAGE=A

NODE=M036PP;LINKAGE=D

 **$a_0(980)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>980 ± 20 OUR ESTIMATE</b> Mass determination very model dependent			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1000.7_{-0.7}^{+12.9}$	<sup>1</sup> LU	20	RVUE $\gamma\gamma \rightarrow \pi^0\eta, K_S^0 K_S^0$
<sup>1</sup> T-matrix pole on sheet II.			

NODE=M036205

NODE=M036MX

→ UNCHECKED ←

NODE=M036MX;LINKAGE=A

 **$\eta\pi$  FINAL STATE ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$1004.1 \pm 1.5 \pm 6.5$		<sup>1</sup> ALBRECHT	20	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta,$ $\pi^0\eta\eta, \pi^0 K^+ K^-$
$982.5 \pm 1.6 \pm 1.1$	16.9k	<sup>2</sup> AMBROSINO	09F	KLOE	$1.02 e^+ e^- \rightarrow \eta\pi^0\gamma$
$986 \pm 4$		ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$
$982.3 \pm 0.6 \pm 3.1$		<sup>3</sup> UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
$985 \pm 4 \pm 6$	318	ACHARD	02B	L3	$183-209 e^+ e^- \rightarrow$ $e^+ e^- \eta\pi^+ \pi^-$
$995 \pm 52$	36	<sup>4</sup> ACHASOV	00F	SND	$e^+ e^- \rightarrow \eta\pi^0\gamma$
$994 \pm 33$	36	<sup>5</sup> ACHASOV	00F	SND	$e^+ e^- \rightarrow \eta\pi^0\gamma$
$975 \pm 7$		BARBERIS	00H		$450 p p \rightarrow p_f \eta\pi^0 p_s$
$988 \pm 8$		BARBERIS	00H		$450 p p \rightarrow$ $\Delta_f^{++} \eta\pi^- p_s$
$\sim 1055$		<sup>6</sup> OLLER	99	RVUE	$\eta\pi, K\bar{K}$
$\sim 1009.2$		<sup>6</sup> OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
$993.1 \pm 2.1$		<sup>7</sup> TEIGE	99	B852	$18.3 \pi^- p \rightarrow$ $\eta\pi^+ \pi^- n$
$988 \pm 6$		<sup>6</sup> ANISOVICH	98B	RVUE	Compilation
987		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
991		JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi,$ $\eta\pi$
$984.45 \pm 1.23 \pm 0.34$		AMSLER	94C	CBAR	$0.0 \bar{p}p \rightarrow \omega\eta\pi^0$
$982 \pm 2$		<sup>8</sup> AMSLER	92	CBAR	$0.0 \bar{p}p \rightarrow \eta\eta\pi^0$
$984 \pm 4$	1040	<sup>8</sup> ARMSTRONG	91B	OMEG ±	$300 p p \rightarrow$ $p p \eta\pi^+ \pi^-$
$976 \pm 6$		ATKINSON	84E	OMEG ±	$25-55 \gamma p \rightarrow \eta\pi n$
$986 \pm 3$	500	<sup>9</sup> EVANGELIS...	81	OMEG ±	$12 \pi^- p \rightarrow$ $\eta\pi^+ \pi^- \pi^- p$
$990 \pm 7$	145	<sup>9</sup> GURTU	79	HBC ±	$4.2 K^- p \rightarrow \Lambda\eta 2\pi$
$980 \pm 11$	47	CONFORTO	78	OSPK -	$4.5 \pi^- p \rightarrow pX^-$
$978 \pm 16$	50	CORDEN	78	OMEG ±	$12-15 \pi^- p \rightarrow n\eta 2\pi$
$977 \pm 7$		GRASSLER	77	HBC -	$16 \pi^\mp p \rightarrow p\eta 3\pi$

OCCUR=2

OCCUR=2

989 ± 4	70	WELLS	75	HBC	-	3.1-6 $K^- p \rightarrow \Lambda \eta 2\pi$
972 ± 10	150	DEFOIX	72	HBC	±	0.7 $\bar{p} p \rightarrow 7\pi$
970 ± 15	20	BARNES	69C	HBC	-	4-5 $K^- p \rightarrow \Lambda \eta 2\pi$
980 ± 10		CAMPBELL	69	DBC	±	2.7 $\pi^+ d$
980 ± 10	15	MILLER	69B	HBC	-	4.5 $K^- N \rightarrow \eta \pi \Lambda$
980 ± 10	30	AMMAR	68	HBC	±	5.5 $K^- p \rightarrow \Lambda \eta 2\pi$

OCCUR=2

<sup>1</sup> T-matrix pole with 2 poles, 2 channels, pole mass on adjacent sheet  $1002.4 \pm 1.4 \pm 6.6$  MeV.

NODE=M036M1;LINKAGE=D

<sup>2</sup> Using the model of ACHASOV 89 and ACHASOV 03B.

NODE=M036M1;LINKAGE=AM

<sup>3</sup> From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

NODE=M036M1;LINKAGE=UE

<sup>4</sup> Using the model of ACHASOV 89. Supersedes ACHASOV 98B.

NODE=M036M1;LINKAGE=V1

<sup>5</sup> Using the model of JAFFE 77. Supersedes ACHASOV 98B.

NODE=M036M1;LINKAGE=M2

<sup>6</sup> T-matrix pole.

NODE=M036M1;LINKAGE=AN

<sup>7</sup> Breit-Wigner fit, average between  $a_0^\pm$  and  $a_0^0$ . The fit favors a slightly heavier  $a_0^\pm$ .

NODE=M036M1;LINKAGE=BF

<sup>8</sup> From a single Breit-Wigner fit.

NODE=M036M1;LINKAGE=A

<sup>9</sup> From  $f_1(1285)$  decay.

NODE=M036M1;LINKAGE=R

 **$K\bar{K}$  ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M036M2

NODE=M036M2

• • • We do not use the following data for averages, fits, limits, etc. • • •

947.7 <sup>+</sup> <sub>-5.0</sub> ± 6.6		<sup>1</sup> AAIJ	19H	LHCB	$pp \rightarrow D^\pm X$
925 ± 5 ± 8	190k	<sup>2</sup> AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
~ 1053		<sup>3</sup> OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
975 ± 15		BERTIN	98B	OBLX	$0.0 \bar{p} p \rightarrow K^\pm K_S \pi^\mp$
970 ± 10	316	DEBILLY	80	HBC	$1.2-2 \bar{p} p \rightarrow f_1(1285)\omega$
1016 ± 10	100	<sup>4</sup> ASTIER	67	HBC	$0.0 \bar{p} p$
1003.3 ± 7.0	143	<sup>5,6</sup> ROSENFELD	65	RVUE	

<sup>1</sup> From the  $D^\pm \rightarrow K^\pm K^+ K^-$  Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

NODE=M036M2;LINKAGE=C

<sup>2</sup> Using a two-channel resonance parametrization with couplings fixed to ABELE 98.

NODE=M036M2;LINKAGE=B

<sup>3</sup> T-matrix pole.

NODE=M036M2;LINKAGE=AN

<sup>4</sup> ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.

NODE=M036M2;LINKAGE=A

<sup>5</sup> Note on  $J^P$ . Main argument for  $0^+$  is small  $Q$  value. Isotropy of decay distribution in  $\bar{p} p$  at rest proves nothing. See discussion by Rosenfeld (Oxford) and Butterworth (Heidelberg).

NODE=M036M2;LINKAGE=01

<sup>6</sup> Plus systematic errors.

NODE=M036M2;LINKAGE=S

 **$a_0(980)$  WIDTH**

NODE=M036210

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M036W1

**50 to 100 OUR ESTIMATE** Width determination very model dependent. Peak width in  $\eta\pi$  is about 60 MeV, but decay width can be much larger.

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

97.2 ± 1.9 ± 5.7		<sup>1</sup> ALBRECHT	20	CBAR	$0.9 \bar{p} p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
73.2 <sup>+</sup> <sub>-5.2</sub> ± 25.4		<sup>2</sup> LU	20	RVUE	$\gamma\gamma \rightarrow \pi^0 \eta, K_S^0 K_S^0$
75.6 ± 1.6 <sup>+</sup> <sub>-10.0</sub> ± 17.4		<sup>3</sup> UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
50 ± 13 ± 4	318	ACHARD	02B	L3	$183-209 e^+ e^- \rightarrow$ $e^+ e^- \eta \pi^+ \pi^-$
72 ± 16		BARBERIS	00H		$450 pp \rightarrow p_f \eta \pi^0 p_s$
61 ± 19		BARBERIS	00H		$450 pp \rightarrow$ $\Delta_f^{++} \eta \pi^- p_s$
~ 42		<sup>4</sup> OLLER	99	RVUE	$\eta\pi, K\bar{K}$
~ 112		<sup>4</sup> OLLER	99B	RVUE	$\pi\pi \rightarrow \eta\pi, K\bar{K}$
71 ± 7		TEIGE	99	B852	$18.3 \pi^- p \rightarrow$ $\eta \pi^+ \pi^- n$
92 ± 20		<sup>4</sup> ANISOVICH	98B	RVUE	Compilation
65 ± 10		<sup>5</sup> BERTIN	98B	OBLX ±	$0.0 \bar{p} p \rightarrow K^\pm K_S \pi^\mp$
~ 100		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
202		JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi,$ $\eta\pi$
54.12 ± 0.34 ± 0.12		AMSLER	94C	CBAR	$0.0 \bar{p} p \rightarrow \omega \eta \pi^0$

OCCUR=2

54 ±10		<sup>6</sup> AMSLER	92	CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
95 ±14	1040	<sup>6</sup> ARMSTRONG	91B	OMEG ±	300 $pp \rightarrow$ $pp\eta\pi^+\pi^-$
62 ±15	500	<sup>7</sup> EVANGELIS...	81	OMEG ±	12 $\pi^- p \rightarrow$ $\eta\pi^+\pi^-\pi^- p$
60 ±20	145	<sup>7</sup> GURTU	79	HBC ±	4.2 $K^- p \rightarrow \Lambda\eta 2\pi$
60 $\begin{smallmatrix} +50 \\ -30 \end{smallmatrix}$	47	CONFORTO	78	OSPK -	4.5 $\pi^- p \rightarrow pX^-$
86.0 $\begin{smallmatrix} +60.0 \\ -50.0 \end{smallmatrix}$	50	CORDEN	78	OMEG ±	12-15 $\pi^- p \rightarrow n\eta 2\pi$
44 ±22		GRASSLER	77	HBC -	16 $\pi^\mp p \rightarrow p\eta 3\pi$
80 to 300		<sup>8</sup> FLATTE	76	RVUE -	4.2 $K^- p \rightarrow \Lambda\eta 2\pi$
16.0 $\begin{smallmatrix} +25.0 \\ -16.0 \end{smallmatrix}$	70	<sup>9</sup> WELLS	75	HBC -	3.1-6 $K^- p \rightarrow \Lambda\eta 2\pi$
30 ± 5	150	<sup>10</sup> DEFOIX	72	HBC ±	0.7 $\bar{p}p \rightarrow 7\pi$
40 ±15		CAMPBELL	69	DBC ±	2.7 $\pi^+ d$
60 ±30	15	MILLER	69B	HBC -	4.5 $K^- N \rightarrow \eta\pi\Lambda$
80 ±30	30	AMMAR	68	HBC ±	5.5 $K^- p \rightarrow \Lambda\eta 2\pi$

<sup>1</sup> T-matrix pole with 2 poles, 2 channels, pole width on adjacent sheet  $127.0 \pm 2.3 \pm 6.7$  MeV.

<sup>2</sup> T-matrix pole on sheet II.

<sup>3</sup> From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

<sup>4</sup> T-matrix pole.

<sup>5</sup> The  $\eta\pi$  width.

<sup>6</sup> From a single Breit-Wigner fit.

<sup>7</sup> From  $f_1(1285)$  decay.

<sup>8</sup> Using a two-channel resonance parametrization of GAY 76B data.

<sup>9</sup> Weak evidence only for  $a_0(980)^+$  production.

<sup>10</sup> This number has very little meaning. Error is much too small. Vlada

NODE=M036W1;LINKAGE=C

NODE=M036W1;LINKAGE=D

NODE=M036W1;LINKAGE=UE

NODE=M036W1;LINKAGE=AN

NODE=M036W1;LINKAGE=BE

NODE=M036W1;LINKAGE=A

NODE=M036W1;LINKAGE=R

NODE=M036W1;LINKAGE=F

NODE=M036W1;LINKAGE=W

NODE=M036W1;LINKAGE=01

### **$K\bar{K}$ ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 48		<sup>1</sup> OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
~ 25	100	<sup>2</sup> ASTIER	67	HBC ±	
57 ± 13	143	<sup>3</sup> ROSENFELD	65	RVUE ±	

<sup>1</sup> T-matrix pole.

<sup>2</sup> ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.

<sup>3</sup> Plus systematic errors.

NODE=M036W2

NODE=M036W2

NODE=M036W2;LINKAGE=AN

NODE=M036W2;LINKAGE=A

NODE=M036W2;LINKAGE=S

### **$a_0(980)$ DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta\pi$	seen
$\Gamma_2$ $K\bar{K}$	seen
$\Gamma_3$ $\eta'\pi$	seen
$\Gamma_4$ $\rho\pi$	not seen
$\Gamma_5$ $\gamma\gamma$	seen
$\Gamma_6$ $e^+e^-$	

NODE=M036215;NODE=M036

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=8

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=6

### **$a_0(980)$ PARTIAL WIDTHS**

#### **$\Gamma(\gamma\gamma)$**

VALUE (keV)	DOCUMENT ID	TECN
-------------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.30 ± 0.10	<sup>1</sup> AMSLER	98	RVUE
-------------	---------------------	----	------

<sup>1</sup> Using  $\Gamma_{\gamma\gamma} B(a_0(980) \rightarrow \eta\pi) = 0.24 \pm 0.08$  keV.

**$\Gamma_5$**

NODE=M036217

NODE=M036W4

NODE=M036W4

NODE=M036W4;LINKAGE=A

$a_0(980) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M036220

 $\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_5/\Gamma$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.21</b>	<b>+0.08</b>	<b>OUR AVERAGE</b>		
	<b>-0.04</b>			

NODE=M036G1  
NODE=M036G1

0.128	$+0.003 + 0.502$	<sup>1</sup> UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
	$-0.002 - 0.043$				
0.28	$\pm 0.04 \pm 0.10$	44	OEST	90	JADE $e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.19	$\pm 0.07 + 0.10$		ANTREASYAN	86	CBAL $e^+e^- \rightarrow e^+e^-\pi^0\eta$
	$-0.07$				

<sup>1</sup> From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

NODE=M036G1;LINKAGE=UE

 $\Gamma(\eta\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_6/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.5</b>	90	VOROBYEV	88	ND $e^+e^- \rightarrow \pi^0\eta$

NODE=M036G2  
NODE=M036G2 $a_0(980) \text{ BRANCHING RATIOS}$ 

NODE=M036225

 $\Gamma(K\bar{K})/\Gamma(\eta\pi)$  $\Gamma_2/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

**0.172±0.019 OUR AVERAGE**

[0.177 ± 0.024 OUR 2022 AVERAGE Scale factor = 1.2]

0.137±0.036±0.042	<sup>1</sup> ABLIKIM	22AH	BES3	$D_s^+ \rightarrow K_S^0 K^+ \pi^0$
0.23 ± 0.05	<sup>2</sup> ABELE	98	CBAR	$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
0.166±0.01±0.02	<sup>3</sup> BARBERIS	98c	OMEG	$450 p p \rightarrow p_f f_1(1285) p_S$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.138±0.001±0.035	<sup>4</sup> ALBRECHT	20	CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
1.20 ± 0.15	<sup>5</sup> ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$
1.05 ± 0.07 ± 0.05	<sup>6</sup> BUGG	08A	RVUE 0	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
0.57 ± 0.16	<sup>7</sup> BARGIOTTI	03	OBLX	$\bar{p}p$
~ 0.60	OLLER	99B	RVUE	$\pi\pi \rightarrow \eta\pi, K\bar{K}$
0.7 ± 0.3	<sup>3</sup> CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow n\eta 2\pi$
0.25 ± 0.08	<sup>3</sup> DEFOIX	72	HBC ±	$0.7 \bar{p} \rightarrow 7\pi$

<sup>1</sup> Using  $D_s^+ \rightarrow a_0(980)^+ \pi^0$  from ABLIKIM 19BE.

<sup>2</sup> Using  $\pi^0 \pi^0 \eta$  from AMSLER 94D.

<sup>3</sup> From the decay of  $f_1(1285)$ .

<sup>4</sup> Residues from T-matrix pole with 2 poles, 2 channels. Solution on adjacent sheet  $0.149 \pm 0.001 \pm 0.039$ .

<sup>5</sup> This is a ratio of couplings.

<sup>6</sup> A ratio of couplings, using AMSLER 94D and ABELE 98. Supersedes BUGG 94.

<sup>7</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0,$  and  $K^\pm K_S^0 \pi^\mp$ .

NODE=M036R2  
NODE=M036R2  
NEWNODE=M036R2;LINKAGE=B  
NODE=M036R2;LINKAGE=Q  
NODE=M036R2;LINKAGE=L  
NODE=M036R2;LINKAGE=ANODE=M036R2;LINKAGE=AN  
NODE=M036R2;LINKAGE=BU  
NODE=M036R;LINKAGE=BG $\Gamma(\eta' \pi)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	116k	<sup>1</sup> CHEN	20A	BELL $D^0 \rightarrow K^- \pi^+ \eta$

NODE=M036R00  
NODE=M036R00

<sup>1</sup> From an amplitude analysis of the  $D^0 \rightarrow K^- \pi^+ \eta$  decay in a three-channel Flatte model with a 10.1  $\sigma$  significance. Earlier observed by ABLIKIM 17K in the  $\chi_{c1} \rightarrow \eta \pi^+ \pi^-$  decay with a 8.9  $\sigma$  significance.

NODE=M036R00;LINKAGE=A

 $\Gamma(\rho\pi)/\Gamma(\eta\pi)$  $\Gamma_4/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.25	70	<sup>1</sup> AMMAR	70	HBC ±	$4.1, 5.5 K^- p \rightarrow \Lambda \eta 2\pi$
-------	----	--------------------	----	-------	--

<sup>1</sup> Not clear if they really observed the  $a_0(980)$  3 standard deviations.

NODE=M036R1  
NODE=M036R1  
NODE=M036R1

NODE=M036R1;LINKAGE=01

a<sub>0</sub>(980) REFERENCES

NODE=M036

ABLIKIM	22AH	PRL 129 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61880
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)	REFID=60333
LU	20	EPJ C80 436	J. Lu, B. Moussallam		REFID=60436
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
ABLIKIM	19BE	PRL 123 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60055
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57953
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
BUGG	08A	PR D78 074023	D.V. Bugg	(LOQM)	REFID=52578
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
ACHASOV	03B	PR D68 014006	N.N. Achsaov, A.V. Kiselev		REFID=49476
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48574
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
OLLER	99	PR D60 099906 (erratum)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (erratum)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
TEIGE	99	PR D59 012001	S. Teige <i>et al.</i>	(BNL E852 Collab.)	REFID=46613
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46317
AMSLER	98	RMP 70 1293	C. Amshler		REFID=46601
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
JANSSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
AMSLER	94C	PL B327 425	C. Amshler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
AMSLER	94D	PL B333 277	C. Amshler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AMSLER	92	PL B291 347	C. Amshler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20624
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
DEBILLY	80	NP B176 1	L. de Billy <i>et al.</i>	(CURIN, LAUS, NEUC+)	REFID=20461
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
CONFORTO	78	LNC 23 419	B. Conforto <i>et al.</i>	(RHEL, TINTO, CHIC+)	REFID=20451
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20452
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
JAFFE	77	PR D15 267,281	R. Jaffe	(MIT)	REFID=43673
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)	REFID=20446
GAY	76B	PL 63B 220	J.B. Gay <i>et al.</i>	(CERN, AMST, NIJM) JP	REFID=20445
WELLS	75	NP B101 333	J. Wells <i>et al.</i>	(OXF)	REFID=20444
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
AMMAR	70	PR D2 430	R. Ammar <i>et al.</i>	(KANS, NWES, ANL, WISC)	REFID=20428
BARNES	69C	PRL 23 610	V.E. Barnes <i>et al.</i>	(BNL, SYRA)	REFID=20418
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)	REFID=20419
MILLER	69B	PL 29B 255	D.H. Miller <i>et al.</i>	(PURD)	REFID=20424
Also		PR 188 2011	W.L. Yen <i>et al.</i>	(PURD)	REFID=20425
AMMAR	68	PRL 21 1832	R. Ammar <i>et al.</i>	(NWES, ANL)	REFID=20412
ASTIER	67	PL 25B 294	A. Astier <i>et al.</i>	(CDEF, CERN, IRAD)	REFID=20405
Includes data of		BARLOW 67, CONFORTO 67, and ARMENTEROS 65.			
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
CONFORTO	67	NP B3 469	G. Conforto <i>et al.</i>	(CERN, CDEF, IPNP+)	REFID=20411
ARMENTEROS	65	PL 17 344	R. Armenteros <i>et al.</i>	(CERN, CDEF)	REFID=20396
ROSENFELD	65	Oxford Conf. 58	A.H. Rosenfeld	(LRL)	REFID=20399



$\phi(1020)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M004

 $\phi(1020)$  MASS

NODE=M004M

NODE=M004M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1019.461 ± 0.016 OUR AVERAGE</b>				
1019.463 ± 0.061	2.3M	1 KOZYREV	18 CMD3	$e^+e^- \rightarrow K^+K^-$ , $K_S^0 K_L^0$
1019.462 ± 0.042 ± 0.056	28k	2 LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1019.51 ± 0.02 ± 0.05		3 LEES	13Q BABR	$e^+e^- \rightarrow K^+K^- \gamma$
1019.30 ± 0.02 ± 0.10	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.52 ± 0.05 ± 0.05	17.4k	AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
1019.483 ± 0.011 ± 0.025	272k	4 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
1019.42 ± 0.05	1900k	5 ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-$ , $K_S K_L, \pi^+\pi^-\pi^0$
1019.40 ± 0.04 ± 0.05	23k	AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1019.36 ± 0.12		6 ACHASOV	00B SND	$e^+e^- \rightarrow \eta\gamma$
1019.38 ± 0.07 ± 0.08	2200	7 AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \pi^+\pi^- \geq 2\gamma$
1019.51 ± 0.07 ± 0.10	11169	AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.5 ± 0.4		BARBERIS 98	OMEG	450 $pp \rightarrow pp2K^+2K^-$
1019.42 ± 0.06	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
1019.7 ± 0.3	2012	DAVENPORT 86	MPSF	400 $pA \rightarrow 4KX$
1019.7 ± 0.1 ± 0.1	5079	ALBRECHT 85D	ARG	10 $e^+e^- \rightarrow K^+K^-X$
1019.3 ± 0.1	1500	ARENTO	82 AEMS	11.8 polar. $pp \rightarrow KK$
1019.67 ± 0.17	25080	8 PELLINEN	82 RVUE	
1019.52 ± 0.13	3681	BUKIN	78C OLYA	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1018.4 ± 0.5 ± 0.1		9 ALBRECHT	20 CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
1019.21 ± 0.04 ± 0.03		10 HOID	20 RVUE	$e^+e^- \rightarrow \pi^0\gamma$
1019.54 ± 0.10 ± 0.51		11 AAIJ	19H LHCB	$pp \rightarrow D^\pm X$
1019.20 ± 0.02 ± 0.01		12 HOFERICHT...	19 RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.469 ± 0.061	1.7M	KOZYREV	18 CMD3	$e^+e^- \rightarrow K^+K^-$
1019.457 ± 0.061	610k	KOZYREV	16 CMD3	$e^+e^- \rightarrow K_S^0 K_L^0$
1019.48 ± 0.01		LEES	13F BABR	$D^+ \rightarrow K^+K^-\pi^+$
1019.441 ± 0.008 ± 0.080	542k	13 AKHMETSHIN 08	CMD2	1.02 $e^+e^- \rightarrow K^+K^-$
1019.63 ± 0.07	12540	14 AUBERT,B	05J BABR	$D^0 \rightarrow \bar{K}^0 K^+ K^-$
1019.8 ± 0.7		ARMSTRONG 86	OMEG	85 $\pi^+ / pp \rightarrow \pi^+ / p4Kp$
1020.1 ± 0.11	5526	14 ATKINSON	86 OMEG	20–70 $\gamma p$
1019.7 ± 1.0		BEBEK	86 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
1019.411 ± 0.008	642k	15 DIJKSTRA	86 SPEC	100–200 $\pi^\pm, \bar{p}, p, K^\pm$ , on Be
1020.9 ± 0.2		14 FRAME	86 OMEG	13 $K^+p \rightarrow \phi K^+p$
1021.0 ± 0.2		14 ARMSTRONG 83B	OMEG	18.5 $K^-p \rightarrow K^-K^+\Lambda$
1020.0 ± 0.5		14 ARMSTRONG 83B	OMEG	18.5 $K^-p \rightarrow K^-K^+\Lambda$
1019.7 ± 0.3		14 BARATE	83 GOLI	190 $\pi^-Be \rightarrow 2\mu X$
1019.8 ± 0.2 ± 0.5	766	IVANOV	81 OLYA	1–1.4 $e^+e^- \rightarrow K^+K^-$
1019.4 ± 0.5	337	COOPER	78B HBC	0.7–0.8 $\bar{p}p \rightarrow K_S^0 K_L^0 \pi^+\pi^-$
1020 ± 1	383	14 BALDI	77 CNTR	10 $\pi^-p \rightarrow \pi^- \phi p$
1018.9 ± 0.6	800	COHEN	77 ASPK	6 $\pi^\pm N \rightarrow K^+K^-N$
1019.7 ± 0.5	454	KALBFLEISCH 76	HBC	2.18 $K^-p \rightarrow \Lambda K \bar{K}$
1019.4 ± 0.8	984	BESCH	74 CNTR	2 $\gamma p \rightarrow pK^+K^-$
1020.3 ± 0.4	100	BALLAM	73 HBC	2.8–9.3 $\gamma p$
1019.4 ± 0.7		BINNIE	73B CNTR	$\pi^-p \rightarrow \phi n$
1019.6 ± 0.5	120	16 AGUILAR-...	72B HBC	3.9, 4.6 $K^-p \rightarrow \Lambda K^+K^-$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

1019.9 ±0.5	100	<sup>16</sup> AGUILAR...	72B	HBC	3.9,4.6 $K^- p \rightarrow K^- p K^+ K^-$	OCCUR=2
1020.4 ±0.5	131	COLLEY	72	HBC	10 $K^+ p \rightarrow K^+ p \phi$	
1019.9 ±0.3	410	STOTTLE...	71	HBC	2.9 $K^- p \rightarrow \Sigma / \Lambda K \bar{K}$	
		<sup>1</sup> Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.				NODE=M004M;LINKAGE=G
		<sup>2</sup> Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$ , $\omega(782)$ , and $\phi(1020)$ .				NODE=M004M;LINKAGE=E
		<sup>3</sup> Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$ , $\omega(782)$ , $\phi(1020)$ and their higher mass excitations.				NODE=M004M;LINKAGE=C
		<sup>4</sup> Update of AKHMETSHIN 99D				NODE=M004M;LINKAGE=GS
		<sup>5</sup> From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$ , $K_S K_L$ , $\pi^+ \pi^- \pi^0$ , and $\eta \gamma$ decays modes and using ACHASOV 00B for the $\eta \gamma$ decay mode.				NODE=M004M;LINKAGE=AE
		<sup>6</sup> Using a total width of $4.43 \pm 0.05$ MeV. Systematic uncertainty included.				NODE=M004M;LINKAGE=G2
		<sup>7</sup> Using a total width of $4.43 \pm 0.05$ MeV.				NODE=M004M;LINKAGE=F2
		<sup>8</sup> PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DE-GROOT 74.				NODE=M004M;LINKAGE=R
		<sup>9</sup> Width fixed at 4.2 MeV.				NODE=M004M;LINKAGE=J
		<sup>10</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives $1019.457 \pm 0.020$ MeV.				NODE=M004M;LINKAGE=M
		<sup>11</sup> From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.				NODE=M004M;LINKAGE=I
		<sup>12</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.				NODE=M004M;LINKAGE=N
		<sup>13</sup> Strongly correlated with AKHMETSHIN 04.				NODE=M004M;LINKAGE=AH
		<sup>14</sup> Systematic errors not evaluated.				NODE=M004M;LINKAGE=A
		<sup>15</sup> Weighted and scaled average of 12 measurements of DIJKSTRA 86.				NODE=M004M;LINKAGE=B
		<sup>16</sup> Mass errors enlarged by us to $\Gamma/\sqrt{N}$ ; see the note with the $K^*(892)$ mass.				NODE=M004M;LINKAGE=D

### $\phi(1020)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>4.249±0.013 OUR AVERAGE</b>		Error includes scale factor of 1.1.				
4.245±0.013	2.3M	<sup>1</sup> KOZYREV	18	CMD3 $e^+ e^- \rightarrow K^+ K^-$ , $K_S^0 K_L^0$	OCCUR=2	
4.205±0.103±0.067	28k	<sup>2</sup> LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \gamma$		
4.29 ±0.04 ±0.07		<sup>3</sup> LEES	13Q	BABR $e^+ e^- \rightarrow K^+ K^- \gamma$		
4.30 ±0.06 ±0.17	105k	AKHMETSHIN 06	CMD2	0.98–1.06 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$		
4.280±0.033±0.025	272k	<sup>4</sup> AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$		
4.21 ±0.04	1900k	<sup>5</sup> ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-$ , $K_S K_L, \pi^+ \pi^- \pi^0$		
4.44 ±0.09	55600	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow$ hadrons		
4.5 ±0.7	1500	ARENTON 82	AEMS	11.8 polar. $pp \rightarrow KK$		
4.2 ±0.6	766	<sup>6</sup> IVANOV 81	OLYA	1–1.4 $e^+ e^- \rightarrow K^+ K^-$		
4.3 ±0.6		<sup>6</sup> CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$		
4.36 ±0.29	3681	<sup>6</sup> BUKIN 78C	OLYA	$e^+ e^- \rightarrow$ hadrons		
4.4 ±0.6	984	<sup>6</sup> BESCH 74	CNTR	$2 \gamma p \rightarrow p K^+ K^-$		
4.67 ±0.72	681	<sup>6</sup> BALAKIN 71	OSPK	$e^+ e^- \rightarrow$ hadrons		
4.09 ±0.29		BIZOT 70	OSPK	$e^+ e^- \rightarrow$ hadrons		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
4.07 ±0.13 ±0.01		<sup>7</sup> HOID 20	RVUE	$e^+ e^- \rightarrow \pi^0 \gamma$		
4.23 ±0.04 ±0.02		<sup>8</sup> HOFERICH... 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$		
4.249±0.015	1.7M	KOZYREV 18	CMD3	$e^+ e^- \rightarrow K^+ K^-$		
4.240±0.017	610k	KOZYREV 16	CMD3	$e^+ e^- \rightarrow K_S^0 K_L^0$		
4.37 ±0.02		LEES 13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$		
4.24 ±0.02 ±0.03	542k	<sup>9</sup> AKHMETSHIN 08	CMD2	1.02 $e^+ e^- \rightarrow K^+ K^-$		
4.28 ±0.13	12540	<sup>10</sup> AUBERT,B 05J	BABR	$D^0 \rightarrow \bar{K}^0 K^+ K^-$		
4.45 ±0.06	271k	DIJKSTRA 86	SPEC	100 $\pi^- Be$		
3.6 ±0.8	337	<sup>6</sup> COOPER 78B	HBC	0.7–0.8 $\bar{p} p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$		
4.5 ±0.50	1300	<sup>6,10</sup> AKERLOF 77	SPEC	400 $pA \rightarrow K^+ K^- X$		
4.5 ±0.8	500	<sup>6,10</sup> AYRES 74	ASPK	3–6 $\pi^- p \rightarrow K^+ K^- n, K^- p \rightarrow K^+ K^- \Lambda / \Sigma^0$		
3.81 ±0.37		COSME 74B	OSPK	$e^+ e^- \rightarrow K_L^0 K_S^0$		
3.8 ±0.7	454	<sup>6</sup> BORENSTEIN 72	HBC	2.18 $K^- p \rightarrow K \bar{K} n$		

- <sup>1</sup> Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.
- <sup>2</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$  and higher mass excitations of  $\rho(770)$ ,  $\omega(782)$ , and  $\phi(1020)$ .
- <sup>3</sup> Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$  and their higher mass excitations.
- <sup>4</sup> Update of AKHMETSHIN 99D
- <sup>5</sup> From the combined fit assuming that the total  $\phi(1020)$  production cross section is saturated by those of  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $\pi^+\pi^-\pi^0$ , and  $\eta\gamma$  decays modes and using ACHASOV 00B for the  $\eta\gamma$  decay mode.
- <sup>6</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.
- <sup>7</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization.
- <sup>8</sup> The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.
- <sup>9</sup> Strongly correlated with AKHMETSHIN 04.
- <sup>10</sup> Systematic errors not evaluated.

NODE=M004W;LINKAGE=G

NODE=M004W;LINKAGE=E

NODE=M004W;LINKAGE=C

NODE=M004W;LINKAGE=GS  
NODE=M004W;LINKAGE=AE

NODE=M004W;LINKAGE=D  
NODE=M004W;LINKAGE=I  
NODE=M004W;LINKAGE=K

NODE=M004W;LINKAGE=AH  
NODE=M004W;LINKAGE=A

### $\phi(1020)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $K^+K^-$	(49.1 $\pm$ 0.5 ) %	S=1.3
$\Gamma_2$ $K_L^0 K_S^0$	(33.9 $\pm$ 0.4 ) %	S=1.2
$\Gamma_3$ $\rho\pi + \pi^+\pi^-\pi^0$	(15.4 $\pm$ 0.4 ) %	S=1.2
$\Gamma_4$ $\rho\pi$		
$\Gamma_5$ $\pi^+\pi^-\pi^0$		
$\Gamma_6$ $\eta\gamma$	( 1.301 $\pm$ 0.025 ) %	S=1.2
$\Gamma_7$ $\pi^0\gamma$	( 1.32 $\pm$ 0.05 ) $\times 10^{-3}$	
$\Gamma_8$ $\ell^+\ell^-$	—	
$\Gamma_9$ $e^+e^-$	( 2.979 $\pm$ 0.033 ) $\times 10^{-4}$	S=1.3
$\Gamma_{10}$ $\mu^+\mu^-$	( 2.85 $\pm$ 0.19 ) $\times 10^{-4}$	
$\Gamma_{11}$ $\eta e^+e^-$	( 1.08 $\pm$ 0.04 ) $\times 10^{-4}$	
$\Gamma_{12}$ $\pi^+\pi^-$	( 7.3 $\pm$ 1.3 ) $\times 10^{-5}$	
$\Gamma_{13}$ $\omega\pi^0$	( 4.7 $\pm$ 0.5 ) $\times 10^{-5}$	
$\Gamma_{14}$ $\omega\gamma$	< 5 %	CL=84%
$\Gamma_{15}$ $\rho\gamma$	< 1.2 $\times 10^{-5}$	CL=90%
$\Gamma_{16}$ $\pi^+\pi^-\gamma$	( 4.1 $\pm$ 1.3 ) $\times 10^{-5}$	
$\Gamma_{17}$ $f_0(980)\gamma$	( 3.22 $\pm$ 0.19 ) $\times 10^{-4}$	S=1.1
$\Gamma_{18}$ $\pi^0\pi^0\gamma$	( 1.12 $\pm$ 0.06 ) $\times 10^{-4}$	
$\Gamma_{19}$ $\pi^+\pi^-\pi^+\pi^-$	( 3.9 $^{+2.8}_{-2.2}$ ) $\times 10^{-6}$	
$\Gamma_{20}$ $\pi^+\pi^+\pi^-\pi^-\pi^0$	< 4.6 $\times 10^{-6}$	CL=90%
$\Gamma_{21}$ $\pi^0 e^+ e^-$	( 1.33 $^{+0.07}_{-0.10}$ ) $\times 10^{-5}$	
$\Gamma_{22}$ $\pi^0\eta\gamma$	( 7.27 $\pm$ 0.30 ) $\times 10^{-5}$	S=1.5
$\Gamma_{23}$ $a_0(980)\gamma$	( 7.6 $\pm$ 0.6 ) $\times 10^{-5}$	
$\Gamma_{24}$ $K^0\bar{K}^0\gamma$	< 1.9 $\times 10^{-8}$	CL=90%
$\Gamma_{25}$ $\eta'(958)\gamma$	( 6.21 $\pm$ 0.21 ) $\times 10^{-5}$	
$\Gamma_{26}$ $\eta\pi^0\pi^0\gamma$	< 2 $\times 10^{-5}$	CL=90%
$\Gamma_{27}$ $\mu^+\mu^-\gamma$	( 1.4 $\pm$ 0.5 ) $\times 10^{-5}$	
$\Gamma_{28}$ $\rho\gamma\gamma$	< 1.2 $\times 10^{-4}$	CL=90%
$\Gamma_{29}$ $\eta\pi^+\pi^-$	< 1.8 $\times 10^{-5}$	CL=90%
$\Gamma_{30}$ $\eta\mu^+\mu^-$	< 9.4 $\times 10^{-6}$	CL=90%
$\Gamma_{31}$ $\eta U \rightarrow \eta e^+ e^-$	< 1 $\times 10^{-6}$	CL=90%
$\Gamma_{32}$ invisible	< 1.7 $\times 10^{-4}$	CL=90%
<b>Lepton Family number (LF) violating modes</b>		
$\Gamma_{33}$ $e^\pm\mu^\mp$	LF < 2 $\times 10^{-6}$	CL=90%

NODE=M004215;NODE=M004

DESIG=1

DESIG=2

DESIG=24

DESIG=16

DESIG=3

DESIG=4

DESIG=7

DESIG=256;OUR EVAL;→ UNCHECKED ←

DESIG=5

DESIG=6

DESIG=17

DESIG=8

DESIG=25

DESIG=10

DESIG=12

DESIG=9

DESIG=20

DESIG=19

DESIG=15

DESIG=14

DESIG=21

DESIG=22

DESIG=23

DESIG=257

DESIG=194

DESIG=195

DESIG=196

DESIG=250

DESIG=255

DESIG=26

DESIG=259

DESIG=260

NODE=M004;CLUMP=A  
DESIG=258

## CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 80 measurements and one constraint to determine 14 parameters. The overall fit has a  $\chi^2 = 61.8$  for 67 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-73										
$x_3$	-60	-10									
$x_6$	-21	18	2								
$x_7$	-11	11	2	8							
$x_9$	48	-51	-8	-37	-22						
$x_{10}$	-6	6	1	4	3	-12					
$x_{12}$	-3	3	0	2	1	-6	1				
$x_{13}$	-4	4	1	3	2	-8	1	0			
$x_{17}$	0	0	0	0	0	0	0	0	0		
$x_{18}$	-10	10	1	18	4	-19	2	1	2	0	
$x_{19}$	-1	1	0	1	0	-2	0	0	0	0	0
$x_{23}$	0	0	0	0	0	0	0	0	0	0	0
$x_{25}$	-7	6	1	33	3	-12	1	1	1	1	0
		$x_1$	$x_2$	$x_3$	$x_6$	$x_7$	$x_9$	$x_{10}$	$x_{12}$	$x_{13}$	$x_{17}$
$x_{19}$	0										
$x_{23}$	0	0									
$x_{25}$	6	0	0								
		$x_{18}$	$x_{19}$	$x_{23}$							

### $\phi(1020)$ PARTIAL WIDTHS

#### $\Gamma(\eta\gamma)$

VALUE (keV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  $\Gamma_6$

••• We do not use the following data for averages, fits, limits, etc. •••

58.9 ± 0.5 ± 2.4      ACHASOV    00    SND     $e^+ e^- \rightarrow \eta\gamma$

NODE=M004218

NODE=M004W6  
NODE=M004W6

#### $\Gamma(\pi^0\gamma)$

VALUE (keV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  $\Gamma_7$

••• We do not use the following data for averages, fits, limits, etc. •••

5.40 ± 0.16<sup>+0.43</sup><sub>-0.40</sub>      ACHASOV    00    SND     $e^+ e^- \rightarrow \pi^0\gamma$

NODE=M004W7  
NODE=M004W7

#### $\Gamma(\ell^+ \ell^-)$

VALUE (keV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  $\Gamma_8$

••• We do not use the following data for averages, fits, limits, etc. •••

1.320 ± 0.017 ± 0.015      <sup>1</sup> AMBROSINO    05    KLOE    1.02  $e^+ e^- \rightarrow \mu^+ \mu^-$

<sup>1</sup> Weighted average of  $\Gamma_{ee}$  and  $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$  from AMBROSINO 05 assuming lepton universality.

NODE=M004W5  
NODE=M004W5

NODE=M004W5;LINKAGE=AM

#### $\Gamma(e^+ e^-)$

VALUE (keV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  $\Gamma_9$

**1.27 ± 0.04 OUR EVALUATION**

**1.251 ± 0.021 OUR AVERAGE** Error includes scale factor of 1.1.

1.235 ± 0.006 ± 0.022      <sup>1</sup> AKHMETSHIN    11    CMD2    1.02  $e^+ e^- \rightarrow \phi$

1.32 ± 0.05 ± 0.03      <sup>2</sup> AMBROSINO    05    KLOE    1.02  $e^+ e^- \rightarrow e^+ e^-$

1.28 ± 0.05      AKHMETSHIN    95    CMD2    1.02  $e^+ e^- \rightarrow \phi$

<sup>1</sup> Combined analysis of the CMD-2 data on  $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta\gamma$  assuming that the sum of their branching fractions is 0.99741 ± 0.00007.

<sup>2</sup> From forward-backward asymmetry and using  $\Gamma_{\text{total}} = 4.26 \pm 0.05$  MeV from the 2004 edition of this Review.

NODE=M004W8  
NODE=M004W8

→ UNCHECKED ←

NODE=M004W8;LINKAGE=AK

NODE=M004W8;LINKAGE=AM

$\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-)^{1/2}$				$(\Gamma_9\Gamma_{10})^{1/2}$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
<b>1.320 ± 0.018 ± 0.017</b>	AMBROSINO	05	KLOE	1.02 $e^+e^- \rightarrow \mu^+\mu^-$

NODE=M004W9  
NODE=M004W9

### $\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M004223

$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_1\Gamma_9/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.6340 ± 0.0070 ± 0.0039</b>		<sup>1</sup> LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

NODE=M004G01  
NODE=M004G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.669 ± 0.001 ± 0.023 1.7M KOZYREV 18 CMD3  $e^+e^- \rightarrow K^+K^-$

NODE=M004G01;LINKAGE=A

<sup>1</sup> Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$  and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.

$\Gamma(K_L^0 K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_2\Gamma_9/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.4200 ± 0.0033 ± 0.0123</b>	28k	<sup>1</sup> LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$

NODE=M004GXX  
NODE=M004GXX

<sup>1</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$  and higher mass excitations of  $\rho(770)$ ,  $\omega(782)$ , and  $\phi(1020)$ .

NODE=M004GXX;LINKAGE=A

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)] \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_3\Gamma_9/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT	
<b>184.1 ± 2.1 ± 8.0</b>	<sup>1</sup> LEES	21B	BABR	10.5 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

NODE=M004R03  
NODE=M004R03

<sup>1</sup> From the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$  with contributions from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ .

NODE=M004R03;LINKAGE=A

### $\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M004224

$\Gamma(K^+K^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma \times \Gamma_9/\Gamma$
VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>14.64 ± 0.28 OUR FIT</b>				Error includes scale factor of 1.4.


NODE=M004G10  
NODE=M004G10

**14.6 ± 0.5 OUR AVERAGE** Error includes scale factor of 1.8. See the ideogram below.

15.789 ± 0.541	1.7M	KOZYREV	18	CMD3	$e^+e^- \rightarrow K^+K^-$
14.27 ± 0.05 ± 0.31	542k	AKHMETSHIN	08	CMD2	1.02 $e^+e^- \rightarrow K^+K^-$
13.93 ± 0.14 ± 0.99	1000k	<sup>1</sup> ACHASOV	01E	SND	$e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$

NODE=M004G10;LINKAGE=AE

<sup>1</sup> From the combined fit assuming that the total  $\phi(1020)$  production cross section is saturated by those of  $K^+K^-$ ,  $K_S K_L$ ,  $\pi^+\pi^-\pi^0$ , and  $\eta\gamma$  decays modes and using ACHASOV 00B for the  $\eta\gamma$  decay mode.



$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.11 ±0.12 OUR FIT</b>				
<b>10.07 ±0.13 OUR AVERAGE</b>				
10.078±0.223	610k	1 KOZYREV	16 CMD3	$e^+ e^- \rightarrow K_S^0 K_L^0$
10.01 ±0.04 ±0.17	272k	2 AKHMETSHIN	04 CMD2	$e^+ e^- \rightarrow K_L^0 K_S^0$
10.27 ±0.07 ±0.34	500k	3 ACHASOV	01E SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$

<sup>1</sup> KOZYREV 16 also reports  $\Gamma(e^+ e^-) B(\phi \rightarrow K_S^0 K_L^0) = (0.428 \pm 0.001 \pm 0.009) \text{ keV}$ .

<sup>2</sup> Update of AKHMETSHIN 99D

<sup>3</sup> From the combined fit assuming that the total  $\phi(1020)$  production cross section is saturated by those of  $K^+ K^-$ ,  $K_S K_L$ ,  $\pi^+ \pi^- \pi^0$ , and  $\eta\gamma$  decays modes and using ACHASOV 00B for the  $\eta\gamma$  decay mode.

NODE=M004G6  
NODE=M004G6

NODE=M004G6;LINKAGE=A  
NODE=M004G;LINKAGE=GS  
NODE=M004G6;LINKAGE=AE

$$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.58 ±0.11 OUR FIT</b>				Error includes scale factor of 1.1.
<b>4.51 ±0.14 OUR AVERAGE</b>				
4.51 ±0.16 ±0.11	105k	AKHMETSHIN	06 CMD2	$0.98-1.06 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
4.665±0.042±0.261	400k	<sup>1</sup> ACHASOV	01E SND	$e^+ e^- \rightarrow K^+ K^-, K_S K_L, \pi^+ \pi^- \pi^0$
4.35 ±0.27 ±0.08	11169	<sup>2</sup> AKHMETSHIN	98 CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.38 ±0.12		BENAYOUN	10 RVUE	0.4-1.05 $e^+ e^-$
4.30 ±0.08 ±0.21		<sup>3</sup> AUBERT,B	04N BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$

<sup>1</sup> From the combined fit assuming that the total  $\phi(1020)$  production cross section is saturated by those of  $K^+ K^-$ ,  $K_S K_L$ ,  $\pi^+ \pi^- \pi^0$ , and  $\eta\gamma$  decays modes and using ACHASOV 00B for the  $\eta\gamma$  decay mode.

<sup>2</sup> Recalculated by us from the cross section in the peak.

<sup>3</sup> Superseded by LEES 21B.

NODE=M004G7  
NODE=M004G7

NODE=M004G7;LINKAGE=AE

NODE=M004G;LINKAGE=B  
NODE=M004G7;LINKAGE=A

$$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.88 ±0.07 OUR FIT</b>				Error includes scale factor of 1.2.
<b>3.93 ±0.09 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
4.050±0.067±0.118	33k	<sup>1</sup> ACHASOV	07B SND	0.6-1.38 $e^+ e^- \rightarrow \eta\gamma$
4.093 <sup>+0.040</sup> <sub>-0.043</sub> ±0.247	17.4k	<sup>2</sup> AKHMETSHIN	05 CMD2	0.60-1.38 $e^+ e^- \rightarrow \eta\gamma$
3.850±0.041±0.159	23k	<sup>3,4</sup> AKHMETSHIN	01B CMD2	$e^+ e^- \rightarrow \eta\gamma$
4.00 ±0.04 ±0.11		<sup>5</sup> ACHASOV	00 SND	$e^+ e^- \rightarrow \eta\gamma$
3.53 ±0.08 ±0.17	2200	<sup>6,7</sup> AKHMETSHIN	99F CMD2	$e^+ e^- \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.19 ±0.06		<sup>8</sup> BENAYOUN	10 RVUE	0.4-1.05 $e^+ e^-$
------------	--	-----------------------	---------	--------------------

<sup>1</sup> From a combined fit of  $\sigma(e^+ e^- \rightarrow \eta\gamma)$  with  $\eta \rightarrow 3\pi^0$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$ , and fixing  $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 1.44 \pm 0.04$ . Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.

<sup>2</sup> From the  $\eta \rightarrow 2\gamma$  decay and using  $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$ .

<sup>3</sup> From the  $\eta \rightarrow 3\pi^0$  decay and using  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ .

<sup>4</sup> The combined fit from 600 to 1380 MeV taking into account  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ , and  $\rho(1450)$  (mass and width fixed at 1450 MeV and 310 MeV respectively).

<sup>5</sup> From the  $\eta \rightarrow 2\gamma$  decay and using  $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$ .

<sup>6</sup> Recalculated by the authors from the cross section in the peak.

<sup>7</sup> From the  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decay and using  $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = (23.1 \pm 0.5) \times 10^{-2}$ .


<sup>8</sup> A simultaneous fit of  $e^+ e^- \rightarrow \pi^+ \pi^-, \pi^+ \pi^- \pi^0, \pi^0 \gamma, \eta\gamma$  data.

NODE=M004G2  
NODE=M004G2

NODE=M004G2;LINKAGE=AH

NODE=M004G2;LINKAGE=AK  
NODE=M004G;LINKAGE=AK  
NODE=M004G;LINKAGE=BQ

NODE=M004G2;LINKAGE=A  
NODE=M004G;LINKAGE=A  
NODE=M004G2;LINKAGE=C  
NODE=M004G7;LINKAGE=BE



**$\Gamma(\pi^0\gamma)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_7/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units 10<sup>-7</sup>)    EVTS    DOCUMENT ID    TECN    COMMENT

**3.94±0.16 OUR FIT**

**3.95±0.17 OUR AVERAGE**

4.04±0.09±0.19    1 ACHASOV    16A    SND    0.60-1.38 e<sup>+</sup>e<sup>-</sup> → π<sup>0</sup>γ

3.75±0.11±0.29    18k    AKHMETSHIN    05    CMD2    0.60-1.38 e<sup>+</sup>e<sup>-</sup> → π<sup>0</sup>γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.29±0.11    2 BENAYOUN    10    RVUE    0.4-1.05 e<sup>+</sup>e<sup>-</sup>

3.67±0.10<sup>+0.27</sup><sub>-0.25</sub>    3 ACHASOV    00    SND    e<sup>+</sup>e<sup>-</sup> → π<sup>0</sup>γ

<sup>1</sup> From the VMD model with the interfering ρ(770), ω(782), φ(1020) resonances, and an additional resonance describing the total contribution of the ρ(1450) and ω(1420) states. Supersedes ACHASOV 00.

<sup>2</sup> A simultaneous fit of e<sup>+</sup>e<sup>-</sup> → π<sup>+</sup>π<sup>-</sup>, π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>, π<sup>0</sup>γ, ηγ data.

<sup>3</sup> From the π<sup>0</sup> → 2γ decay and using B(π<sup>0</sup> → 2γ) = (98.798 ± 0.032) × 10<sup>-2</sup>.

NODE=M004G3  
NODE=M004G3

NODE=M004G3;LINKAGE=B

NODE=M004G3;LINKAGE=BE  
NODE=M004G3;LINKAGE=A

**$\Gamma(\mu^+\mu^-)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units 10<sup>-8</sup>)    DOCUMENT ID    TECN    COMMENT

**8.5 ±0.6 OUR FIT**

**8.8 ±0.9 OUR AVERAGE** Error includes scale factor of 1.5. See the ideogram below.

8.36±0.59±0.37    ACHASOV    01G    SND    e<sup>+</sup>e<sup>-</sup> → μ<sup>+</sup>μ<sup>-</sup>

9.9 ±1.4 ±0.9    1 ACHASOV    99c    SND    e<sup>+</sup>e<sup>-</sup> → μ<sup>+</sup>μ<sup>-</sup>

14.4 ±3.0    2 VASSERMAN    81    OLYA    e<sup>+</sup>e<sup>-</sup> → μ<sup>+</sup>μ<sup>-</sup>


8.6 ±5.9    2 AUGUSTIN    73    OSPK    e<sup>+</sup>e<sup>-</sup> → μ<sup>+</sup>μ<sup>-</sup>

<sup>1</sup> Recalculated by the authors from the cross section in the peak.

<sup>2</sup> Recalculated by us from the cross section in the peak.

NODE=M004G5  
NODE=M004G5

NODE=M004G5;LINKAGE=A  
NODE=M004G5;LINKAGE=B



**$\Gamma(\pi^+\pi^-)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{12}/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units  $10^{-8}$ ) DOCUMENT ID TECN COMMENT

**2.2 ±0.4 OUR FIT**  
**2.2 ±0.4 OUR AVERAGE**

2.1 ±0.3 ±0.3	1	ACHASOV	00c	SND	$e^+e^- \rightarrow \pi^+\pi^-$
1.95 <sup>+1.15</sup> <sub>-0.87</sub>	2	GOLUBEV	86	ND	$e^+e^- \rightarrow \pi^+\pi^-$
6.01 <sup>+3.19</sup> <sub>-2.51</sub>	2	VASSERMAN	81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
3.31±0.99	3	BENAYOUN	13	RVUE	0.4-1.05 $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Recalculated by the authors from the cross section in the peak.  
<sup>2</sup> Recalculated by us from the cross section in the peak.  
<sup>3</sup> A simultaneous fit to  $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$ , and  $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$  data.

NODE=M004G4  
NODE=M004G4

NODE=M004G4;LINKAGE=AC  
NODE=M004G4;LINKAGE=B  
NODE=M004G4;LINKAGE=A

**$\Gamma(\omega\pi^0)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{13}/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units  $10^{-8}$ ) DOCUMENT ID TECN COMMENT

**1.40±0.15 OUR FIT**  
**1.37±0.17±0.01** 1,2 AMBROSINO 08G KLOE  $e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$

<sup>1</sup> Recalculated by the authors from the cross section at the peak.  
<sup>2</sup> AMBROSINO 08G reports  $[\Gamma(\phi(1020) \rightarrow \omega\pi^0)/\Gamma_{total} \times \Gamma(\phi(1020) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$  which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M004G11  
NODE=M004G11

NODE=M004G11;LINKAGE=AB  
NODE=M004G11;LINKAGE=AM

**$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{18}/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units  $10^{-8}$ ) DOCUMENT ID TECN COMMENT

**3.34±0.17 OUR FIT**  
**3.33<sup>+0.04+0.19</sup><sub>-0.09-0.20</sub>** 1 AMBROSINO 07 KLOE  $e^+e^- \rightarrow \pi^0\pi^0\gamma$

<sup>1</sup> Calculated by the authors from the cross section at the peak.

NODE=M004G9  
NODE=M004G9

NODE=M004G9;LINKAGE=AM

**$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{19}/\Gamma \times \Gamma_9/\Gamma$**

VALUE (units  $10^{-9}$ ) EVTS DOCUMENT ID TECN COMMENT

**1.2<sup>+0.8</sup><sub>-0.7</sub> OUR FIT**  
**1.17±0.52±0.64** 3285 1 AKHMETSHIN 00E CMD2  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

<sup>1</sup> Recalculated by the authors from the cross section in the peak.

NODE=M004G8  
NODE=M004G8

NODE=M004G8;LINKAGE=A



$\phi(1020)$  BRANCHING RATIOS

NODE=M004220

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ NODE=M004R1  
NODE=M004R1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.491±0.005 OUR FIT** Error includes scale factor of 1.3.**0.493±0.010 OUR AVERAGE**

0.492±0.012	2913	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K^+K^-$
0.44 ±0.05	321	KALBFLEISCH 76	HBC	2.18 $K^-p \rightarrow \Lambda K^+K^-$
0.49 ±0.06	270	DEGROOT 74	HBC	4.2 $K^-p \rightarrow \Lambda\phi$
0.540±0.034	565	BALAKIN 71	OSPK	$e^+e^- \rightarrow K^+K^-$
0.48 ±0.04	252	LINDSEY 66	HBC	2.1-2.7 $K^-p \rightarrow \Lambda K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.493±0.003±0.007		<sup>1</sup> AKHMETSHIN 11	CMD2	1.02 $e^+e^- \rightarrow K^+K^-$
0.476±0.017	1000k	<sup>2</sup> ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$

<sup>1</sup> Combined analysis of the CMD-2 data on  $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$  assuming that the sum of their branching fractions is  $0.99741 \pm 0.00007$ .<sup>2</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$ .

NODE=M004R1;LINKAGE=AK

NODE=M004R1;LINKAGE=B2

 $\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ NODE=M004R2  
NODE=M004R2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.339±0.004 OUR FIT** Error includes scale factor of 1.2.**0.331±0.009 OUR AVERAGE**

0.335±0.010	40644	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
0.326±0.035		DOLINSKY 91	ND	$e^+e^- \rightarrow K_L^0 K_S^0$
0.310±0.024		DRUZHININ 84	ND	$e^+e^- \rightarrow K_L^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.336±0.002±0.006		<sup>1</sup> AKHMETSHIN 11	CMD2	1.02 $e^+e^- \rightarrow K_S^0 K_L^0$
0.351±0.013	500k	<sup>2</sup> ACHASOV 01E	SND	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
0.27 ±0.03	133	KALBFLEISCH 76	HBC	2.18 $K^-p \rightarrow \Lambda K_L^0 K_S^0$
0.257±0.030	95	<sup>3</sup> BALAKIN 71	OSPK	$e^+e^- \rightarrow K_L^0 K_S^0$
0.40 ±0.04	167	LINDSEY 66	HBC	2.1-2.7 $K^-p \rightarrow \Lambda K_L^0 K_S^0$

<sup>1</sup> Combined analysis of the CMD-2 data on  $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$  assuming that the sum of their branching fractions is  $0.99741 \pm 0.00007$ .<sup>2</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$ .<sup>3</sup> Balakin error increased by Paul.

NODE=M004R2;LINKAGE=AK

NODE=M004R2;LINKAGE=B2

NODE=M004R2;LINKAGE=01

 $\Gamma(K_L^0 K_S^0)/\Gamma(K^+K^-)$   $\Gamma_2/\Gamma_1$ NODE=M004R19  
NODE=M004R19

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.690±0.015 OUR FIT** Error includes scale factor of 1.3.**0.740±0.031 OUR AVERAGE**

0.70 ±0.06	2732	BUKIN 78C	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0$
0.82 ±0.08		LOSTY 78	HBC	4.2 $K^-p \rightarrow \phi$ hyperon
0.71 ±0.05		LAVEN 77	HBC	10 $K^-p \rightarrow K^+K^-\Lambda$
0.71 ±0.08		LYONS 77	HBC	3-4 $K^-p \rightarrow \Lambda\phi$
0.89 ±0.10	144	AGUILAR-...	72B	HBC 3.9,4.6 $K^-p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.638±0.022	2.3M	<sup>1</sup> KOZYREV 18	CMD3	$e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$
0.68 ±0.03		<sup>2</sup> AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$

<sup>1</sup> The prediction taking into account phase-space difference, radiative corrections, isospin breaking, and the Sommerfeld-Gamow-Sakharov factor gives 0.630.<sup>2</sup> Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNSKIY 07. BENAYOUN 12 obtains  $0.71 \pm 0.01$  in the HLS model.

NODE=M004R19;LINKAGE=A

NODE=M004R19;LINKAGE=KH

$$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$$

$$\Gamma_2/(\Gamma_1+\Gamma_2)$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.408±0.005 OUR FIT</b>				Error includes scale factor of 1.3.
<b>0.45 ±0.04 OUR AVERAGE</b>				
0.44 ±0.07		<sup>1</sup> LONDON	66 HBC	2.24 $K^- p \rightarrow \Lambda K\bar{K}$
0.48 ±0.07	52	BADIER	65B HBC	3 $K^- p$
0.40 ±0.10	34	SCHLEIN	63 HBC	1.95 $K^- p \rightarrow \Lambda K\bar{K}$

NODE=M004R5  
NODE=M004R5

<sup>1</sup> This is probably not affected by their controversial background subtraction; the value is from their numbers of  $K_1 K_2$  vs  $K^+ K^-$  events.

NODE=M004R5;LINKAGE=01

$$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.154±0.004 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.151±0.009 OUR AVERAGE</b>				Error includes scale factor of 1.7.
0.161±0.008	11761	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.143±0.007		DOLINSKY 91	ND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.155±0.002±0.005		<sup>1</sup> AKHMETSHIN 11	CMD2	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.159±0.008	400k	<sup>2</sup> ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0$
0.145±0.009±0.003	11169	<sup>3</sup> AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.139±0.007		<sup>4</sup> PARROUR 76B	OSPK	$e^+ e^-$

NODE=M004R3  
NODE=M004R3

<sup>1</sup> Combined analysis of the CMD-2 data on  $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta, \gamma$  assuming that the sum of their branching fractions is  $0.99741 \pm 0.00007$ .

<sup>2</sup> Using  $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$ .

<sup>3</sup> Using  $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .

<sup>4</sup> Using  $\Gamma(\phi) = 4.1$  MeV. If interference between the  $\rho\pi$  and  $3\pi$  modes is neglected, the fraction of the  $\rho\pi$  is more than 80% at the 90% confidence level.

NODE=M004R3;LINKAGE=AK

NODE=M004R3;LINKAGE=B2

NODE=M004R;LINKAGE=8D

NODE=M004R3;LINKAGE=E

$$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma(K^+ K^-)$$

$$\Gamma_3/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.313±0.010 OUR FIT</b>				Error includes scale factor of 1.2.
<b>0.28 ±0.09</b>	34	AGUILAR-...	72B HBC	3.9,4.6 $K^- p$

NODE=M004R20  
NODE=M004R20

$$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma(K\bar{K})$$

$$\Gamma_3/(\Gamma_1+\Gamma_2)$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.185±0.005 OUR FIT</b>			Error includes scale factor of 1.2.
<b>0.24 ±0.04 OUR AVERAGE</b>			
0.237±0.039	CERRADA 77B	HBC	4.2 $K^- p \rightarrow \Lambda 3\pi$
0.30 ±0.15	LONDON 66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \pi^0$

NODE=M004R6  
NODE=M004R6

$$[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma(K_L^0 K_S^0)$$

$$\Gamma_3/\Gamma_2$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.453±0.012 OUR FIT</b>				Error includes scale factor of 1.1.
<b>0.51 ±0.05 OUR AVERAGE</b>				
0.56 ±0.07	3681	BUKIN 78C	OLYA	$e^+ e^- \rightarrow K_L^0 K_S^0, \pi^+ \pi^- \pi^0$
0.47 ±0.06	516	COSME 74	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M004R7  
NODE=M004R7

$$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
≈ 0.0087	1.98M	<sup>1,2</sup> ALOISIO 03	KLOE	1.02	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
< 0.0006	90	<sup>3</sup> ACHASOV 02	SND	1.02	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
< 0.23	90	<sup>3</sup> CORDIER 80	DM1		$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
< 0.20	90	<sup>3</sup> PARROUR 76B	OSPK		$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M004R46  
NODE=M004R46

<sup>1</sup> From a fit without limitations on charged and neutral  $\rho$  masses and widths.

<sup>2</sup> Adding the direct and  $\omega\pi$  contributions and considering the interference between the  $\rho\pi$  and  $\pi^+ \pi^- \pi^0$ .

<sup>3</sup> Neglecting the interference between the  $\rho\pi$  and  $\pi^+ \pi^- \pi^0$ .

NODE=M004R;LINKAGE=L1

NODE=M004R;LINKAGE=L2

NODE=M004R;LINKAGE=46

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.301 ± 0.025 OUR FIT** Error includes scale factor of 1.2.**1.26 ± 0.04 OUR AVERAGE**

1.246 ± 0.025 ± 0.057	10k	<sup>1</sup> ACHASOV	98F SND	$e^+e^- \rightarrow 7\gamma$
1.18 ± 0.11	279	<sup>2</sup> AKHMETSHIN	95 CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1.30 ± 0.06		<sup>3</sup> DRUZHININ	84 ND	$e^+e^- \rightarrow 3\gamma$
1.4 ± 0.2		<sup>4</sup> DRUZHININ	84 ND	$e^+e^- \rightarrow 6\gamma$
0.88 ± 0.20	290	KURDADZE	83C OLYA	$e^+e^- \rightarrow 3\gamma$
1.35 ± 0.29		ANDREWS	77 CNTR	6.7-10 $\gamma$ Cu
1.5 ± 0.4	54	<sup>3</sup> COSME	76 OSPK	$e^+e^-$

NODE=M004R11  
 NODE=M004R11

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1.38 ± 0.02 ± 0.02		<sup>5</sup> AKHMETSHIN	11 CMD2	1.02 $e^+e^- \rightarrow \eta\gamma$
1.36 ± 0.05 ± 0.02	33k	<sup>6</sup> ACHASOV	07B SND	0.6-1.38 $e^+e^- \rightarrow \eta\gamma$
1.373 ± 0.014 ± 0.085	17.4k	<sup>7,8</sup> AKHMETSHIN	05 CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
1.287 ± 0.013 ± 0.063		<sup>9,10</sup> AKHMETSHIN	01B CMD2	$e^+e^- \rightarrow \eta\gamma$
1.338 ± 0.012 ± 0.052		<sup>11</sup> ACHASOV	00 SND	$e^+e^- \rightarrow \eta\gamma$
1.18 ± 0.03 ± 0.06	2200	<sup>12</sup> AKHMETSHIN	99F CMD2	$e^+e^- \rightarrow \eta\gamma$
1.21 ± 0.07		<sup>13</sup> BENAYOUN	96 RVUE	0.54-1.04 $e^+e^- \rightarrow \eta\gamma$

<sup>1</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$  and  $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$ .<sup>2</sup> From  $\pi^+\pi^-\pi^0$  decay mode of  $\eta$ .<sup>3</sup> From  $2\gamma$  decay mode of  $\eta$ .<sup>4</sup> From  $3\pi^0$  decay mode of  $\eta$ .<sup>5</sup> Combined analysis of the CMD-2 data on  $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$  assuming that the sum of their branching fractions is  $0.99741 \pm 0.00007$ .<sup>6</sup> ACHASOV 07B reports  $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$  which we divide by our best value  $B(\phi(1020) \rightarrow e^+e^-) = (2.979 \pm 0.033) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.<sup>7</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$  and  $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$ .<sup>8</sup> Not independent of the corresponding  $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$ .<sup>9</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$  and  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ .<sup>10</sup> The combined fit from 600 to 1380 MeV taking into account  $\rho(770), \omega(782), \phi(1020)$ , and  $\rho(1450)$  (mass and width fixed at 1450 MeV and 310 MeV respectively).<sup>11</sup> From the  $\eta \rightarrow 2\gamma$  decay and using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .<sup>12</sup> From  $\pi^+\pi^-\pi^0$  decay mode of  $\eta$  and using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .<sup>13</sup> Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

NODE=M004R11;LINKAGE=AC  
 NODE=M004R11;LINKAGE=Z3  
 NODE=M004R11;LINKAGE=A  
 NODE=M004R11;LINKAGE=C  
 NODE=M004R11;LINKAGE=AN

NODE=M004R11;LINKAGE=AO

NODE=M004R11;LINKAGE=AH  
 NODE=M004R11;LINKAGE=AK  
 NODE=M004R;LINKAGE=BQ

NODE=M004R;LINKAGE=GA  
 NODE=M004R;LINKAGE=FF  
 NODE=M004R;LINKAGE=TS

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.32 ± 0.05 OUR FIT****1.31 ± 0.13 OUR AVERAGE**

1.30 ± 0.13		DRUZHININ	84 ND	$e^+e^- \rightarrow 3\gamma$
1.4 ± 0.5	32	COSME	76 OSPK	$e^+e^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1.367 ± 0.072		<sup>1</sup> ACHASOV	16A SND	0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$
1.258 ± 0.037 ± 0.077	18k	<sup>2,3</sup> AKHMETSHIN	05 CMD2	0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$
1.226 ± 0.036 <sup>+0.096</sup> <sub>-0.089</sub>		<sup>4</sup> ACHASOV	00 SND	$e^+e^- \rightarrow \pi^0\gamma$
1.26 ± 0.17		<sup>5</sup> BENAYOUN	96 RVUE	0.54-1.04 $e^+e^- \rightarrow \pi^0\gamma$

OCCUR=3

<sup>1</sup> Using  $B(\phi \rightarrow e^+e^-)$  from PDG 15. Supersedes ACHASOV 00.<sup>2</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$ .<sup>3</sup> Not independent of the corresponding  $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$ .<sup>4</sup> From the  $\pi^0 \rightarrow 2\gamma$  decay and using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .<sup>5</sup> Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

NODE=M004R17;LINKAGE=D  
 NODE=M004R17;LINKAGE=AH  
 NODE=M004R17;LINKAGE=AK  
 NODE=M004R;LINKAGE=3G  
 NODE=M004R17;LINKAGE=TS

 $\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$  $\Gamma_6/\Gamma_7$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

10.9 ± 0.3 <sup>+0.7</sup> <sub>-0.8</sub>	ACHASOV	00 SND	$e^+e^- \rightarrow \eta\gamma, \pi^0\gamma$
--	---------	--------	--

NODE=M004R42  
 NODE=M004R42

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.979 ± 0.033 OUR FIT</b>				Error includes scale factor of 1.3.
<b>2.98 ± 0.07 OUR AVERAGE</b>				Error includes scale factor of 1.1.
2.93 ± 0.14	1900k	<sup>1</sup> ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$
2.88 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow \text{hadrons}$
3.00 ± 0.21	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3.10 ± 0.14		<sup>2</sup> PARROUR 76	OSPK	$e^+e^-$
3.3 ± 0.3		COSME 74	OSPK	$e^+e^- \rightarrow \text{hadrons}$
2.81 ± 0.25	681	BALAKIN 71	OSPK	$e^+e^- \rightarrow \text{hadrons}$
3.50 ± 0.27		CHATELUS 71	OSPK	$e^+e^-$

NODE=M004R16  
 NODE=M004R16

<sup>1</sup> From the combined fit assuming that the total  $\phi(1020)$  production cross section is saturated by those of  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $\pi^+\pi^-\pi^0$ , and  $\eta\gamma$  decays modes and using ACHASOV 00B for the  $\eta\gamma$  decay mode.

NODE=M004R16;LINKAGE=A

<sup>2</sup> Using total width 4.2 MeV. They detect  $3\pi$  mode and observe significant interference with  $\omega$  tail. This is accounted for in the result quoted above.

NODE=M004R16;LINKAGE=E

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.85 ± 0.19 OUR FIT</b>			
<b>2.5 ± 0.4 OUR AVERAGE</b>			
2.69 ± 0.46	<sup>1</sup> HAYES 71	CNTR	$8.3, 9.8 \gamma C \rightarrow \mu^+\mu^- X$
2.17 ± 0.60	<sup>1</sup> EARLES 70	CNTR	$6.0 \gamma C \rightarrow \mu^+\mu^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.87 ± 0.20 ± 0.14	<sup>2</sup> ACHASOV 01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$
3.30 ± 0.45 ± 0.32	<sup>3</sup> ACHASOV 99C	SND	$e^+e^- \rightarrow \mu^+\mu^-$
4.83 ± 1.02	<sup>4</sup> VASSERMAN 81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
2.87 ± 1.98	<sup>4</sup> AUGUSTIN 73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M004R10  
 NODE=M004R10

<sup>1</sup> Neglecting interference between resonance and continuum.

NODE=M004R10;LINKAGE=A

<sup>2</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.91 \pm 0.07) \times 10^{-4}$ .

NODE=M004R;LINKAGE=GZ

<sup>3</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .

NODE=M004R10;LINKAGE=8D

<sup>4</sup> Recalculated by us using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .

NODE=M004R;LINKAGE=VA

 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.08 ± 0.04 OUR AVERAGE</b>				
1.075 ± 0.007 ± 0.038	30k	<sup>1</sup> BABUSCI 15	KLOE	$1.02 e^+e^- \rightarrow \eta e^+e^-$
1.19 ± 0.19 ± 0.12	213	<sup>2</sup> ACHASOV 01B	SND	$e^+e^- \rightarrow \eta e^+e^-$
1.14 ± 0.10 ± 0.06	355	<sup>3</sup> AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.13 ± 0.14 ± 0.07	183	<sup>4</sup> AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
1.21 ± 0.14 ± 0.09	130	<sup>5</sup> AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
1.04 ± 0.20 ± 0.08	42	<sup>6</sup> AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
1.3 $\begin{smallmatrix} +0.8 \\ -0.6 \end{smallmatrix}$	7	GOLUBEV 85	ND	$e^+e^- \rightarrow \eta e^+e^-$

NODE=M004R24  
 NODE=M004R24

OCCUR=2

OCCUR=3

OCCUR=4

<sup>1</sup> Using  $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$  from PDG 12.

NODE=M004R24;LINKAGE=A

<sup>2</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$ ,  $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$ , and  $B(\phi \rightarrow e^+e^-) = (3.00 \pm 0.06) \times 10^{-4}$ .

NODE=M004R;LINKAGE=VM

<sup>3</sup> The average of the branching ratios separately obtained from the  $\eta \rightarrow \gamma\gamma$ ,  $3\pi^0$ ,  $\pi^+\pi^-\pi^0$  decays.

NODE=M004R;LINKAGE=H1

<sup>4</sup> From  $\eta \rightarrow \gamma\gamma$  decays and using  $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$ ,  $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 11) \times 10^{-2}$ , and  $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$ .

NODE=M004R;LINKAGE=H2

<sup>5</sup> From  $\eta \rightarrow 3\pi^0$  decays and using  $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$ ,  $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$ ,  $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$ , and  $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$ .

NODE=M004R;LINKAGE=H3

<sup>6</sup> From  $\eta \rightarrow \pi^+\pi^-\pi^0$  decays and using  $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$ ,  $B(\pi^0 \rightarrow e^+e^-) = (1.198 \pm 0.032) \times 10^{-2}$ ,  $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.0 \pm 0.4) \times 10^{-2}$ ,  $B(\phi \rightarrow \pi^+\pi^-\pi^0) = (15.5 \pm 0.6) \times 10^{-2}$ , and  $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$ .

NODE=M004R;LINKAGE=H4

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M004R18  
 NODE=M004R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.71 \pm 0.11 \pm 0.09$		<sup>1</sup> ACHASOV	00C	SND	$e^+e^- \rightarrow \pi^+\pi^-$
$0.65^{+0.38}_{-0.29}$		<sup>1</sup> GOLUBEV	86	ND	$e^+e^- \rightarrow \pi^+\pi^-$
$2.01^{+1.07}_{-0.84}$		<sup>1</sup> VASSERMAN	81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
<6.6	95	BUKIN	78B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
<2.7	95	ALVENSLEB...	72	CNTR	$6.7 \gamma C \rightarrow C\pi^+\pi^-$

<sup>1</sup> Using  $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ .

NODE=M004R18;LINKAGE=8D

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M004R28  
 NODE=M004R28

**4.7 ± 0.5 OUR FIT****5.2<sup>+1.3</sup><sub>-1.1</sub>**

1,2	AULCHENKO	00A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
-----	-----------	-----	-----	---

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.4 \pm 0.6$		<sup>3</sup> AMBROSINO	08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
~5.4		<sup>4</sup> ACHASOV	00E	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$5.5^{+1.6}_{-1.4} \pm 0.3$		<sup>2,5</sup> AULCHENKO	00A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
$4.8^{+1.9}_{-1.7} \pm 0.8$		<sup>4</sup> ACHASOV	99	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

<sup>1</sup> Using the 1996 and 1998 data.  
<sup>2</sup> (2.3 ± 0.3)% correction for other decay modes of the  $\omega(782)$  applied.  
<sup>3</sup> Not independent of the corresponding  $\Gamma(\omega\pi^0) \times \Gamma(e^+e^-) / \Gamma^2(\text{total})$ .  
<sup>4</sup> Using the 1996 data.  
<sup>5</sup> Using the 1998 data.

NODE=M004R28;LINKAGE=K2  
 NODE=M004R28;LINKAGE=K3  
 NODE=M004R28;LINKAGE=AM  
 NODE=M004R28;LINKAGE=VS  
 NODE=M004R28;LINKAGE=K1

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M004R14  
 NODE=M004R14

<0.05	84	LINDSEY	66	HBC	$2.1-2.7 K^-p \rightarrow \Lambda\pi^+\pi^- \text{ neutrals}$
-------	----	---------	----	-----	---

 $\Gamma(\rho\gamma)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M004R15  
 NODE=M004R15

< 0.12	90	<sup>1</sup> AKHMETSHIN	99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
--------	----	-------------------------	-----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7	90	AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<200	84	LINDSEY	66	HBC	$2.1-2.7 K^-p \rightarrow \Lambda\pi^+\pi^- \text{ neutrals}$

<sup>1</sup> Supersedes AKHMETSHIN 97C.

NODE=M004R;LINKAGE=1N

 $\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M004R12  
 NODE=M004R12

<b>0.41 ± 0.12 ± 0.04</b>		30175	<sup>1</sup> AKHMETSHIN	99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
---------------------------	--	-------	-------------------------	-----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.3	90	<sup>2</sup> AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<600	90	KALBFLEISCH	75	HBC	$2.18 K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$
< 70	90	COSME	74	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<400	90	LINDSEY	65	HBC	$2.1-2.7 K^-p \rightarrow \Lambda\pi^+\pi^- \text{ neutrals}$

NODE=M004R;LINKAGE=2N

<sup>1</sup> For  $E_\gamma > 20$  MeV and assuming that  $B(\phi(1020) \rightarrow f_0(980)\gamma)$  is negligible. Supersedes AKHMETSHIN 97C.

<sup>2</sup> For  $E_\gamma > 20$  MeV and assuming that  $B(\phi(1020) \rightarrow f_0(980)\gamma)$  is negligible.

NODE=M004R12;LINKAGE=Z3

 $\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M004R30  
 NODE=M004R30

**3.22 ± 0.19 OUR FIT** Error includes scale factor of 1.1.**3.21 ± 0.19 OUR AVERAGE**

$3.21^{+0.03}_{-0.09} \pm 0.18$			<sup>1</sup> AMBROSINO	07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$2.90 \pm 0.21 \pm 1.54$			<sup>2</sup> AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma, \pi^0\pi^0\gamma$

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.47±0.21	2438	<sup>3</sup> ALOISIO	02D	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
3.5 ±0.3 <sup>+1.3</sup> <sub>-0.5</sub>	419	<sup>4,5</sup> ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.93±0.46±0.50	27188	<sup>6</sup> AKHMETSHIN 99B	CMD2		$e^+e^- \rightarrow \pi^+\pi^-\gamma$
3.05±0.25±0.72	268	<sup>7</sup> AKHMETSHIN 99C	CMD2		$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.5 ±0.5	268	<sup>8</sup> AKHMETSHIN 99C	CMD2		$e^+e^- \rightarrow \pi^0\pi^0\gamma$
3.42±0.30±0.36	164	<sup>4</sup> ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
< 1	90	<sup>9</sup> AKHMETSHIN 97C	CMD2		$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 7	90	<sup>10</sup> AKHMETSHIN 97C	CMD2		$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 20	90	DRUZHININ	87	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

OCCUR=2

OCCUR=2

<sup>1</sup> Obtained by the authors taking into account the  $\pi^+\pi^-$  decay mode. Includes a component due to  $\pi\pi$  production via the  $f_0(500)$  meson. Supersedes ALOISIO 02D.

NODE=M004R30;LINKAGE=MB

<sup>2</sup> From the combined fit of the photon spectra in the reactions  $e^+e^- \rightarrow \pi^+\pi^-\gamma$ ,  $\pi^0\pi^0\gamma$ .

NODE=M004R;LINKAGE=SL

<sup>3</sup> From the negative interference with the  $f_0(500)$  meson of AITALA 01B using the ACHASOV 89 parameterization for the  $f_0(980)$ , a Breit-Wigner for the  $f_0(500)$ , and ACHASOV 01F for the  $\rho\pi$  contribution. Superseded by AMBROSINO 07.

NODE=M004R;LINKAGE=KD

<sup>4</sup> Assuming that the  $\pi^0\pi^0\gamma$  final state is completely determined by the  $f_0\gamma$  mechanism, neglecting the decay  $B(\phi \rightarrow K\bar{K}\gamma)$  and using  $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$ .

NODE=M004R;LINKAGE=AI

<sup>5</sup> Using the value  $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$ .

NODE=M004R;LINKAGE=U8

<sup>6</sup> For  $E_\gamma > 20$  MeV. Supersedes AKHMETSHIN 97C.

NODE=M004R;LINKAGE=3N

<sup>7</sup> Neglecting other intermediate mechanisms ( $\rho\pi$ ,  $\sigma\gamma$ ).

NODE=M004R;LINKAGE=SM

<sup>8</sup> A narrow pole fit taking into account  $f_0(980)$  and  $f_0(1200)$  intermediate mechanisms.

NODE=M004R;LINKAGE=ST

<sup>9</sup> For destructive interference with the Bremsstrahlung process

NODE=M004R30;LINKAGE=A

<sup>10</sup> For constructive interference with the Bremsstrahlung process

NODE=M004R30;LINKAGE=B

### $\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$

 $\Gamma_{17}/\Gamma_6$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.48±0.15 OUR FIT</b>	Error includes scale factor of 1.1.			

NODE=M004R44

NODE=M004R44

<b>2.6 ±0.2 <sup>+0.8</sup><sub>-0.3</sub></b>	419	<sup>1</sup> ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
--	-----	----------------------	-----	-----	---------------------------------------

<sup>1</sup> Assuming that the  $\pi^0\pi^0\gamma$  final state is completely determined by the  $f_0\gamma$  mechanism, neglecting the decay  $B(\phi \rightarrow K\bar{K}\gamma)$  and using  $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$ .

NODE=M004R44;LINKAGE=AI

### $\Gamma(\pi^0\pi^0\gamma)/\Gamma_{total}$

 $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.07 ±0.06 OUR AVERAGE</b>					

NODE=M004R26

NODE=M004R26

1.07 <sup>+0.01</sup> <sub>-0.03</sub> <sup>+0.06</sup> <sub>-0.06</sub>		<sup>1</sup> AMBROSINO	07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
--	--	------------------------	----	------	---------------------------------------

1.08 ±0.17 ±0.09	268	AKHMETSHIN 99C	CMD2		$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------------------	-----	----------------	------	--	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.09 ±0.03 ±0.05	2438	ALOISIO	02D	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------------------	------	---------	-----	------	---------------------------------------

1.158±0.093±0.052	419	<sup>2,3</sup> ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
-------------------	-----	------------------------	-----	-----	---------------------------------------

<10	90	DRUZHININ	87	ND	$e^+e^- \rightarrow 5\gamma$
-----	----	-----------	----	----	------------------------------

<sup>1</sup> Supersedes ALOISIO 02D.

NODE=M004R26;LINKAGE=MB

<sup>2</sup> Using the value  $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$ .

NODE=M004R26;LINKAGE=U8

<sup>3</sup> Supersedes ACHASOV 98I. Excluding  $\omega\pi^0$ .

NODE=M004R26;LINKAGE=V8

### $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\eta\gamma)$

 $\Gamma_{18}/\Gamma_6$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.86 ±0.04 OUR FIT</b>				

NODE=M004R39

NODE=M004R39

<b>0.865±0.070±0.017</b>	419	<sup>1</sup> ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
--------------------------	-----	----------------------	-----	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90 ±0.08 ±0.07	164	ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
------------------	-----	---------	-----	-----	------------------------------

<sup>1</sup> Supersedes ACHASOV 98I. Excluding  $\omega\pi^0$ .

NODE=M004R39;LINKAGE=V8

### $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{total}$

 $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.5 ±2.7 ±1.6</b>	6.8k	<sup>1</sup> AKHMETSHIN 17	CMD3		$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M004R22

NODE=M004R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.93±1.74±2.14	3.3k	AKHMETSHIN 00E	CMD2		$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
----------------	------	----------------	------	--	---

< 870	90	CORDIER	79	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
-------	----	---------	----	------	---

<sup>1</sup> Using the cross section at the  $\phi$  meson peak  $\sigma(\phi) = 4172 \pm 42$  nb, the nonresonant cross section  $\sigma(0) = 1.263 \pm 0.027$  nb and  $\text{Re}(Z) = 0.146 \pm 0.030$ ,  $\text{Im}(Z) = -0.002 \pm 0.024$  for the complex amplitude of the  $\phi \rightarrow \pi^+\pi^-\pi^+\pi^-$  transition.

NODE=M004R22;LINKAGE=A

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{total}$   $\Gamma_{20}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
<150	95	BARKOV 88	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

NODE=M004R27  
NODE=M004R27

$\Gamma(\pi^0e^+e^-)/\Gamma_{total}$   $\Gamma_{21}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.33<sup>+0.07</sup><sub>-0.10</sub> OUR AVERAGE</b>					
$1.35 \pm 0.05^{+0.05}_{-0.10}$		9.5k	1 ANASTASI 16B	KLOE	$e^+e^- \rightarrow \pi^0e^+e^-$
$1.01 \pm 0.28 \pm 0.29$		52	2 ACHASOV 02D	SND	$e^+e^- \rightarrow \pi^0e^+e^-$
$1.22 \pm 0.34 \pm 0.21$		46	3 AKHMETSHIN 01C	CMD2	$e^+e^- \rightarrow \pi^0e^+e^-$
••• We do not use the following data for averages, fits, limits, etc. •••					
<12	90		DOLINSKY 88	ND	$e^+e^- \rightarrow \pi^0e^+e^-$

NODE=M004R31  
NODE=M004R31

- 1 Using  $B(\pi^0 \rightarrow \gamma\gamma)$  from the 2014 Edition of this Review (PDG 14).
- 2 Using various branching ratios from the 2000 Edition of this Review (PDG 00).
- 3 Using  $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$ ,  $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$ , and  $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$ .

NODE=M004R31;LINKAGE=A  
NODE=M004R;LINKAGE=DS  
NODE=M004R;LINKAGE=5H

$\Gamma(\pi^0\eta\gamma)/\Gamma_{total}$   $\Gamma_{22}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.27<math>\pm</math>0.30 OUR AVERAGE</b> Error includes scale factor of 1.5. See the ideogram below.					
$7.06 \pm 0.22$		16.9k	1 AMBROSINO 09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
$8.51 \pm 0.51 \pm 0.57$		607	2 ALOISIO 02C	KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
$7.96 \pm 0.60 \pm 0.40$		197	3 ALOISIO 02C	KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
$8.8 \pm 1.4 \pm 0.9$		36	4 ACHASOV 00F	SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
$9.0 \pm 2.4 \pm 1.0$		80	AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••					
$7.01 \pm 0.10 \pm 0.20$		13.3k	2,5 AMBROSINO 09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
$7.12 \pm 0.13 \pm 0.22$		3.6k	3,6 AMBROSINO 09F	KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
$8.3 \pm 2.3 \pm 1.2$		20	ACHASOV 98B	SND	$e^+e^- \rightarrow 5\gamma$
<250	90		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^0\eta\gamma$

NODE=M004R32  
NODE=M004R32

OCCUR=2

OCCUR=2


OCCUR=3

- 1 Combined results of  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$  decay modes measurements.
- 2 From the decay mode  $\eta \rightarrow \gamma\gamma$ .
- 3 From the decay mode  $\eta \rightarrow \pi^+\pi^-\pi^0$ .
- 4 Supersedes ACHASOV 98B.
- 5 Using  $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$ ,  $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$ , and  $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$ .
- 6 Using  $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$ ,  $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$ , and  $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.73 \pm 0.28)\%$ .

NODE=M004R32;LINKAGE=AM  
NODE=M004R;LINKAGE=C1  
NODE=M004R;LINKAGE=C2  
NODE=M004R32;LINKAGE=AF  
NODE=M004R32;LINKAGE=AB

NODE=M004R32;LINKAGE=AR

WEIGHTED AVERAGE  
7.27 $\pm$ 0.30 (Error scaled by 1.5)



$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ VALUE (units  $10^{-5}$ ) CL% EVTS

DOCUMENT ID TECN COMMENT

**7.6±0.6 OUR FIT**  
**7.6±0.6 OUR AVERAGE**

7.4±0.7

<sup>1</sup> ALOISIO 02C KLOE  $e^+e^- \rightarrow \eta\pi^0\gamma$ 

8.8±1.7

36

<sup>2</sup> ACHASOV 00F SND  $e^+e^- \rightarrow \eta\pi^0\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

11 ±2

<sup>3</sup> GOKALP 02 RVUE  $e^+e^- \rightarrow \eta\pi^0\gamma$ 

&lt;500 90

DOLINSKY 91 ND  $e^+e^- \rightarrow \pi^0\eta\gamma$ <sup>1</sup> Using  $M_{a_0(980)}=984.8$  MeV and assuming  $a_0(980)\gamma$  dominance.<sup>2</sup> Assuming  $a_0(980)\gamma$  dominance in the  $\eta\pi^0\gamma$  final state.<sup>3</sup> Using data of ACHASOV 00F.NODE=M004R33  
NODE=M004R33NODE=M004R;LINKAGE=C3  
NODE=M004R33;LINKAGE=AF  
NODE=M004R;LINKAGE=GK $\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$  $\Gamma_{17}/\Gamma_{23}$ 

VALUE

DOCUMENT ID TECN COMMENT

**6.1±0.6**<sup>1</sup> ALOISIO 02C KLOE  $e^+e^- \rightarrow \eta\pi^0\gamma$ <sup>1</sup> Using results of ALOISIO 02D and assuming that  $f_0(980)$  decays into  $\pi\pi$  only and  $a_0(980)$  into  $\eta\pi$  only.NODE=M004R47  
NODE=M004R47

NODE=M004R;LINKAGE=C4

 $\Gamma(K^0\bar{K}^0\gamma)/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE CL%

DOCUMENT ID TECN COMMENT

<1.9 × 10<sup>-8</sup>

90

AMBROSINO 09C KLOE  $e^+e^- \rightarrow K_S^0 K_S^0 \gamma$ NODE=M004R48  
NODE=M004R48 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ VALUE (units  $10^{-5}$ ) CL% EVTS

DOCUMENT ID TECN COMMENT

**6.21±0.21 OUR FIT****6.21±0.30 OUR AVERAGE**

6.21±0.27±0.12 3407

<sup>1</sup> AMBROSINO 07A KLOE  $1.02 e^+e^- \rightarrow \pi^+\pi^-7\gamma$ 6.7  $\begin{matrix} +2.8 \\ -2.4 \end{matrix} \pm 0.8$ 

12

<sup>2</sup> AULCHENKO 03B SND  $e^+e^- \rightarrow \eta'\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7  $\begin{matrix} +5.0 \\ -4.2 \end{matrix} \pm 1.5$ 

7

AULCHENKO 03B SND  $e^+e^- \rightarrow 7\gamma$ 

6.10±0.61±0.43

120

<sup>3</sup> ALOISIO 02E KLOE  $1.02 e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 8.2  $\begin{matrix} +2.1 \\ -1.9 \end{matrix} \pm 1.1$ 

21

<sup>4</sup> AKHMETSHIN 00B CMD2  $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 4.9  $\begin{matrix} +2.2 \\ -1.8 \end{matrix} \pm 0.6$ 

9

<sup>5</sup> AKHMETSHIN 00F CMD2  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^- \geq 2\gamma$ 

6.4 ±1.6

30

<sup>6</sup> AKHMETSHIN 00F CMD2  $e^+e^- \rightarrow \eta'(958)\gamma$ 6.7  $\begin{matrix} +3.4 \\ -2.9 \end{matrix} \pm 1.0$ 

5

<sup>7</sup> AULCHENKO 99 SND  $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 

&lt;11 90

AULCHENKO 98 SND  $e^+e^- \rightarrow 7\gamma$ 12  $\begin{matrix} +7 \\ -5 \end{matrix} \pm 2$ 

6

<sup>4</sup> AKHMETSHIN 97B CMD2  $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 

&lt;41 90

DRUZHININ 87 ND  $e^+e^- \rightarrow \gamma\eta\pi^+\pi^-$ <sup>1</sup> AMBROSINO 07A reports  $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma)/\Gamma_{\text{total}}] / [B(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$  which we multiply by our best value  $B(\phi(1020) \rightarrow \eta\gamma) = (1.301 \pm 0.025) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.<sup>2</sup> Averaging AULCHENKO 03B with AULCHENKO 99.<sup>3</sup> Using  $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$ .<sup>4</sup> Using the value  $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$ .<sup>5</sup> Using  $B(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$ .<sup>6</sup> Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.<sup>7</sup> Using the value  $B(\eta' \rightarrow \eta\pi^+\pi^-) = (43.7 \pm 1.5) \times 10^{-2}$  and  $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$ .NODE=M004R25  
NODE=M004R25

OCCUR=2

OCCUR=2

NODE=M004R25;LINKAGE=AM

NODE=M004R25;LINKAGE=BK  
NODE=M004R;LINKAGE=E2  
NODE=M004R25;LINKAGE=Q  
NODE=M004R;LINKAGE=T2  
NODE=M004R;LINKAGE=T3  
NODE=M004R25;LINKAGE=AU $\Gamma(\eta'(958)\gamma)/\Gamma(K_L^0 K_S^0)$  $\Gamma_{25}/\Gamma_2$ VALUE (units  $10^{-4}$ ) EVTS

DOCUMENT ID TECN COMMENT

**1.83±0.06 OUR FIT****1.46 $\begin{matrix} +0.64 \\ -0.54 \end{matrix} \pm 0.18$** 

9

<sup>1</sup> AKHMETSHIN 00F CMD2  $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^- \geq 2\gamma$ <sup>1</sup> Using various branching ratios of  $K_S^0$ ,  $K_L^0$ ,  $\eta$ ,  $\eta'$  from the 2000 edition (The European Physical Journal **C15** 1 (2000)) of this Review.NODE=M004R43  
NODE=M004R43

NODE=M004R;LINKAGE=T1



$\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$  $\Gamma_{25}/\Gamma_6$ VALUE (units  $10^{-3}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**4.77±0.15 OUR FIT****4.78±0.20 OUR AVERAGE**

4.77±0.09±0.19 3407

AMBROSINO 07A KLOE 1.02  $e^+e^- \rightarrow \pi^+\pi^-7\gamma$ 

4.70±0.47±0.31 120

<sup>1</sup> ALOISIO 02E KLOE 1.02  $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 6.5  $^{+1.7}_{-1.5}$  ±0.8 21AKHMETSHIN 00B CMD2  $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5  $^{+5.2}_{-4.0}$  ±1.4 6<sup>2</sup> AKHMETSHIN 97B CMD2  $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ <sup>1</sup> From the decay mode  $\eta' \rightarrow \eta\pi^+\pi^-$ ,  $\eta \rightarrow \gamma\gamma$ .<sup>2</sup> Superseded by AKHMETSHIN 00B.NODE=M004R34  
NODE=M004R34NODE=M004R;LINKAGE=E1  
NODE=M004R;LINKAGE=KS $\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ VALUE (units  $10^{-5}$ )

CL%

DOCUMENT ID

TECN

COMMENT

&lt;2

90

AULCHENKO 98

SND

 $e^+e^- \rightarrow 7\gamma$ NODE=M004R36  
NODE=M004R36 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ VALUE (units  $10^{-5}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**1.43±0.45±0.14**

27188

<sup>1</sup> AKHMETSHIN 99B CMD2  $e^+e^- \rightarrow \mu^+\mu^-\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ±1.0

824 ± 33

<sup>2</sup> AKHMETSHIN 97C CMD2  $e^+e^- \rightarrow \mu^+\mu^-\gamma$ <sup>1</sup> For  $E_\gamma > 20$  MeV. Supersedes AKHMETSHIN 97C.<sup>2</sup> For  $E_\gamma > 20$  MeV.NODE=M004R35  
NODE=M004R35NODE=M004R35;LINKAGE=3N  
NODE=M004R35;LINKAGE=A $\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{28}/\Gamma$ VALUE (units  $10^{-4}$ )

CL%

DOCUMENT ID

TECN

COMMENT

&lt;1.2

90

AULCHENKO 08

CMD2

 $\phi \rightarrow \pi^+\pi^-\gamma\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt;5

90

AKHMETSHIN 98

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ NODE=M004R37  
NODE=M004R37 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{29}/\Gamma$ VALUE (units  $10^{-5}$ )

CL%

DOCUMENT ID

TECN

COMMENT

&lt; 1.8

90

AKHMETSHIN 00E

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt; 6.1

90

AULCHENKO 08

CMD2

 $\phi \rightarrow \eta\pi^+\pi^-$ 

&lt;30

90

AKHMETSHIN 98

CMD2

 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ NODE=M004R38  
NODE=M004R38 $\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$  $\Gamma_{30}/\Gamma$ VALUE (units  $10^{-6}$ )

CL%

DOCUMENT ID

TECN

COMMENT

&lt;9.4

90

AKHMETSHIN 01

CMD2

 $e^+e^- \rightarrow \eta e^+e^-$ NODE=M004R45  
NODE=M004R45 $\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{31}/\Gamma$ 

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

<1 × 10<sup>-6</sup>

90

<sup>1</sup> BABUSCI 13B

KLOE

1.02  $e^+e^- \rightarrow \eta e^+e^-$ <sup>1</sup> For a narrow vector  $U$  with mass between 5 and 470 MeV, from the combined analysis of  $\eta \rightarrow \pi^+\pi^-\pi^0$  and  $\eta \rightarrow \pi^0\pi^0\pi^0$  from ARCHILLI 12. Measured 90% CL limits as a function of  $m_U$  range from  $2.2 \times 10^{-8}$  to  $10^{-6}$ .NODE=M004R01  
NODE=M004R01

NODE=M004R01;LINKAGE=A

 $\Gamma(\text{invisible})/\Gamma(K^+K^-)$  $\Gamma_{32}/\Gamma_1$ 

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

<3.4 × 10<sup>-4</sup>

90

ABLIKIM 18S

BES3

 $J/\psi \rightarrow \phi\eta \rightarrow \phi\pi^+\pi^-\pi^0$ NODE=M004R02  
NODE=M004R02

## ————— Lepton Family number (LF) violating modes —————

NODE=M00422A

 $\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$  $\Gamma_{33}/\Gamma$ 

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

<2 × 10<sup>-6</sup>

90

ACHASOV 10A

SND

 $e^+e^- \rightarrow e^\pm\mu^\mp$ NODE=M004R29  
NODE=M004R29 $\pi^+\pi^-\pi^0 / \rho\pi$  AMPLITUDE RATIO  $a_1$  IN DECAY OF  $\phi \rightarrow \pi^+\pi^-\pi^0$ 

NODE=M004D1

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the  $\pi\pi$   $P$ -wave scattering phase shift.

NODE=M004D1

VALUE (units $10^{-2}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.1±1.2 OUR AVERAGE</b>					
10.1±4.4±1.7		80k	<sup>1</sup> AKHMETSHIN 06	CMD2	1.017–1.021 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0±1.1±0.6		1.98M	<sup>2,3</sup> ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$-6 < a_1 < 6$		500k	<sup>3</sup> ACHASOV	02 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$-16 < a_1 < 11$	90	9.8k	<sup>1,4</sup> AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$
<sup>1</sup> Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.					
<sup>2</sup> From a fit without limitations on charged and neutral $\rho$ masses and widths.					
<sup>3</sup> Recalculated by us to match the notations of AKHMETSHIN 98.					
<sup>4</sup> Assuming zero phase for the contact term.					

NODE=M004D1

NODE=M004D1;LINKAGE=AK

NODE=M004D;LINKAGE=L1

NODE=M004D;LINKAGE=L3

NODE=M004D1;LINKAGE=KL

### PARAMETER $\beta$ IN $\phi \rightarrow P e^+ e^-$ DECAYS

In the one-pole approximation the electromagnetic transition form factor for  $\phi \rightarrow P e^+ e^-$  ( $P = \pi, \eta$ ) is given as a function of the  $e^+ e^-$  invariant mass squared,  $q^2$ , by the expression:

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter  $\Lambda \approx 0.770$  GeV ( $\Lambda^{-2} \approx 1.687$  GeV $^{-2}$ ). The slope of this form factor,  $\beta = dF/dq^2(q^2=0)$ , equals  $\Lambda^{-2}$  in this approximation.

The measurements below obtain  $\beta$  in the one-pole approximation.

NODE=M004230

NODE=M004230

### PARAMETER $\beta$ IN $\phi \rightarrow \pi^0 e^+ e^-$ DECAY

VALUE (GeV $^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.02±0.11</b>	9.5k	<sup>1</sup> ANASTASI	16B KLOE	1.02 $e^+e^- \rightarrow \pi^0 e^+e^-$

<sup>1</sup> The error combines statistical and systematic uncertainties.

NODE=M004A00  
NODE=M004A00

NODE=M004A00;LINKAGE=A

### PARAMETER $\beta$ IN $\phi \rightarrow \eta e^+ e^-$ DECAY

VALUE (GeV $^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.29±0.13 OUR AVERAGE</b>				
1.28±0.10 $^{+0.09}_{-0.08}$	30k	BABUSCI	15 KLOE	1.02 $e^+e^- \rightarrow \eta e^+e^-$
3.8 ± 1.8	213	<sup>1</sup> ACHASOV	01B SND	1.02 $e^+e^- \rightarrow \eta e^+e^-$

<sup>1</sup> The uncertainty is statistical only. The systematic one is negligible, in comparison.

NODE=M004BFP  
NODE=M004BFP

NODE=M004BFP;LINKAGE=A

### $\phi(1020)$ REFERENCES

LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)	REFID=61048
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
HOFRICHT...	19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)	REFID=60974
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58971
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
KOZYREV	18	PL B779 64	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)	REFID=58794
AKHMETSHIN	17	PL B768 345	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=57893
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57513
ANASTASI	16B	PL B757 362	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)	REFID=57399
KOZYREV	16	PL B760 314	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)	REFID=57514
BABUSCI	15	PL B742 1	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)	REFID=56374
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=7
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
BABUSCI	13B	PL B720 111	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)	REFID=55068
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)	REFID=55357
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55127
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)	REFID=53951
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>		REFID=54281
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54305
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
AKHMETSHIN	11	PL B695 412	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=53645
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=53352
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52969
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=52572
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52573
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)	REFID=52268

Translated from ZETFP 88 93.

NODE=M004

FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro		REFID=52584
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51616
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51646
DUBYSNSKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>		REFID=51719
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=51465
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=50453
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50824
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49613
		Translated from ZETF 124 28.			
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48814
		Translated from ZETFP 75 539.			
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48823
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48825
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl		REFID=48575
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>		REFID=49167
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48111
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)	REFID=48312
ACHASOV	01G	PRL 86 1698	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48315
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48110
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48323
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell		REFID=48570
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47425
		Translated from ZETF 117 22.			
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47431
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
		Translated from ZETFP 72 411.			
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47927
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47422
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47936
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47937
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
		Translated from ZETF 117 1067.			
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>		REFID=47969
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>		REFID=46896
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>		REFID=46939
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i>		REFID=47397
		Also			
		PL B508 217 (err.)	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48328
AKHMETSHIN	99F	PL B460 242	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47473
AULCHENKO	99	JETPL 69 97	V.M. Aulchenko <i>et al.</i>		REFID=46920
		Translated from ZETFP 69 87.			
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46317
ACHASOV	98F	JETPL 68 573	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46321
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>		REFID=46600
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=46325
AULCHENKO	98	PL B436 199	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46336
BARBERIS	98	PL B432 436	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46344
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin <i>et al.</i>	(NOVO, BOST, PITT+)	REFID=45801
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=45802
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=44617
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
BARKOV	88	SJNP 47 248	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=41024
		Translated from YAF 47 393.			
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41022
		Translated from YAF 48 442.			
DRUZHININ	87	ZPHY C37 1	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=40448
ARMSTRONG	86	PL 166B 245	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=20563
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=11540
DAVENPORT	86	PR D33 2519	T.F. Davenport (TUFTS, ARIZ, FNAL, FSU, NDAM+)		REFID=20567
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra <i>et al.</i>	(ANIK, BRIS, CERN+)	REFID=20568
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)	REFID=20569
GOLUBEV	86	SJNP 44 409	V.B. Golubev <i>et al.</i>	(NOVO)	REFID=40449
		Translated from YAF 44 633.			
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=20562
GOLUBEV	85	SJNP 41 756	V.B. Golubev <i>et al.</i>	(NOVO)	REFID=40450
		Translated from YAF 41 1183.			
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)	REFID=12177
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20560
		Translated from ZETFP 38 306.			
ARENTON	82	PR D25 2241	M.W. Arenton <i>et al.</i>	(ANL, ILL)	REFID=20556
PELLINEN	82	PS 25 599	A. Pellinen, M. Roos	(HELS)	REFID=20557
DAUM	81	PL 100B 439	C. Daum <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)	REFID=20552
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
		Also			
		Private Comm.	S.I. Eidelman	(NOVO)	REFID=20554
VASSERMAN	81	PL 99B 62	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=20555
		Also			
		SJNP 35 240	L.M. Kurdadze <i>et al.</i>		REFID=47475
		Translated from YAF 35 352.			

CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
CORDIER	79	PL 81B 389	A. Cordier <i>et al.</i>	(LALO)	REFID=20549
BUKIN	78B	SJNP 27 521	A.D. Bukin <i>et al.</i>	(NOVO)	REFID=20545
BUKIN	78C	Translated from YAF 27 985. SJNP 27 516	A.D. Bukin <i>et al.</i>	(NOVO)	REFID=20544
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
LOSTY	78	NP B133 38	M.J. Losty <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=20547
AKERLOF	77	PRL 39 861	C.W. Akerlof <i>et al.</i>	(FNAL, MICH, PURD)	REFID=20534
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
BALDI	77	PL 68B 381	R. Baldi <i>et al.</i>	(GEVA)	REFID=20536
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+)	REFID=20537
COHEN	77	PRL 38 269	D. Cohen <i>et al.</i>	(ANL)	REFID=20538
LAVEN	77	NP B127 43	H. Laven <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)	REFID=20541
LYONS	77	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)	REFID=20232
COSME	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20529
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20531
PARROUR	76	PL 63B 357	G. Parrou <i>et al.</i>	(ORSAY)	REFID=20532
PARROUR	76B	PL 63B 362	G. Parrou <i>et al.</i>	(ORSAY)	REFID=20533
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AYRES	74	PRL 32 1463	D.S. Ayres <i>et al.</i>	(ANL)	REFID=20522
BESCH	74	NP B70 257	H.J. Besch <i>et al.</i>	(BONN)	REFID=20523
COSME	74	PL 48B 155	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20525
COSME	74B	PL 48B 159	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20526
DEGROOT	74	NP B74 77	A.J. de Groot <i>et al.</i>	(AMST, NIJM)	REFID=20527
AUGUSTIN	73	PRL 30 462	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=47515
BALLAM	73	PR D7 3150	J. Ballam <i>et al.</i>	(SLAC, LBL)	REFID=20520
BINNIE	73B	PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
ALVENSLEB...	72	PRL 28 66	H. Alvensleben <i>et al.</i>	(MIT, DESY)	REFID=20514
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
COLLEY	72	NP B50 1	D.C. Colley <i>et al.</i>	(BIRM, GLAS)	REFID=20519
BALAKIN	71	PL 34B 328	V.E. Balakin <i>et al.</i>	(NOVO)	REFID=20507
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)	REFID=20508
Also		PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)	REFID=20501
HAYES	71	PR D4 899	S. Hayes <i>et al.</i>	(CORN)	REFID=20511
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemyer	(UMD)	REFID=20512
BIZOT	70	PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)	REFID=20501
Also		Liverpool Sym. 69	J.P. Perez-y-Jorba		REFID=20502
EARLES	70	PRL 25 1312	D.R. Earles <i>et al.</i>	(NEAS)	REFID=20504
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)	REFID=20481
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IGJPC	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=20253
LINDSEY	65	PRL 15 221	J.S. Lindsey, G.A. Smith	(LRL)	REFID=20478
LINDSEY 65 data included in LINDSEY 66.					
SCHLEIN	63	PRL 10 368	P.E. Schlein <i>et al.</i>	(UCLA) IGJP	REFID=20474

# $h_1(1170)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

## $h_1(1170)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>1166 ± 5 ± 3</b>	<sup>1</sup> ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1168 ± 4	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1190 ± 60	<sup>2</sup> DANKOWY...	81	SPEC 0	8 $\pi p \rightarrow 3\pi n$
<sup>1</sup> Average and spread of values using 2 variants of the model of BOWLER 75.				
<sup>2</sup> Uses the model of BOWLER 75.				

## $h_1(1170)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>375 ± 6 ± 34</b>	<sup>3</sup> ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
345 ± 6	ANDO	92	SPEC	8 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
320 ± 50	<sup>4</sup> DANKOWY...	81	SPEC 0	8 $\pi p \rightarrow 3\pi n$
<sup>3</sup> Average and spread of values using 2 variants of the model of BOWLER 75.				
<sup>4</sup> Uses the model of BOWLER 75.				

## $h_1(1170)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \rho\pi$	seen

NODE=M030M

NODE=M030M

OCCUR=2

NODE=M030M;LINKAGE=B  
NODE=M030M;LINKAGE=C

NODE=M030W

NODE=M030W

OCCUR=2

NODE=M030W;LINKAGE=B  
NODE=M030W;LINKAGE=C

NODE=M030215;NODE=M030

DESIG=1;OUR EST;→ UNCHECKED ←

### $h_1(1170)$ BRANCHING RATIOS

NODE=M030220

$\Gamma(\rho\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen	ANDO	92	SPEC $8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
seen	ATKINSON	84	OMEG $20-70 \gamma p \rightarrow \pi^+ \pi^- \pi^0 p$	
seen	DANKOWY...	81	SPEC $8 \pi p \rightarrow 3 \pi n$	

NODE=M030R1  
NODE=M030R1

### $h_1(1170)$ REFERENCES

NODE=M030

ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

REFID=43171  
REFID=20574  
REFID=20572  
REFID=20571

## $b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

NODE=M011


### $b_1(1235)$ MASS

NODE=M011M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1229.5 ± 3.2 OUR AVERAGE</b>		Error includes scale factor of 1.6. See the ideogram below.			
1225 ± 5		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
1235 ± 15		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
1236 ± 16		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
1222 ± 6		ATKINSON 84E	OMEG ±		25-55 $\gamma p \rightarrow \omega \pi X$
1237 ± 7		ATKINSON 84E	OMEG 0		25-55 $\gamma p \rightarrow \omega \pi X$
1239 ± 5		EVANGELIS... 81	OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
1251 ± 8	450	GESSAROLI 77	HBC -		11 $\pi^- p \rightarrow \pi^- \omega p$
1245 ± 11	890	FLATTE 76C	HBC -		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
1222 ± 4	1400	CHALOUPKA 74	HBC -		3.9 $\pi^- p$
1220 ± 7	600	KARSHON 74B	HBC +		4.9 $\pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1190 ± 10		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
1213 ± 5		ATKINSON 84C	OMEG 0		20-70 $\gamma p$
1271 ± 11		COLLICK 84	SPEC +		200 $\pi^+ Z \rightarrow Z \pi \omega$

NODE=M011M

OCCUR=2



### $b_1(1235)$ WIDTH

NODE=M011W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>142 ± 9 OUR AVERAGE</b>		Error includes scale factor of 1.2.			
113 ± 12		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
160 ± 30		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
151 ± 31		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
170 ± 15		EVANGELIS... 81	OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
170 ± 50	225	BALTAY 78B	HBC +		15 $\pi^+ p \rightarrow p 4\pi$
155 ± 32	450	GESSAROLI 77	HBC -		11 $\pi^- p \rightarrow \pi^- \omega p$
182 ± 45	890	FLATTE 76C	HBC -		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
135 ± 20	1400	CHALOUPKA 74	HBC -		3.9 $\pi^- p$
156 ± 22	600	KARSHON 74B	HBC +		4.9 $\pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
210 ± 19		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
231 ± 14		ATKINSON 84C	OMEG 0		20-70 $\gamma p$
232 ± 29		COLLICK 84	SPEC +		200 $\pi^+ Z \rightarrow Z \pi \omega$

NODE=M011W

 **$b_1(1235)$  DECAY MODES**

NODE=M011215;NODE=M011

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\omega \pi$ [D/S amplitude ratio = 0.277 ± 0.027]	seen	
$\Gamma_2$ $\pi^\pm \gamma$	(1.6 ± 0.4) × 10 <sup>-3</sup>	
$\Gamma_3$ $\eta \rho$	seen	
$\Gamma_4$ $\pi^+ \pi^+ \pi^- \pi^0$	< 50 %	84%
$\Gamma_5$ $K^*(892)^\pm K^\mp$	seen	
$\Gamma_6$ $(KK)^\pm \pi^0$	< 8 %	90%
$\Gamma_7$ $K_S^0 K_L^0 \pi^\pm$	< 6 %	90%
$\Gamma_8$ $K_S^0 K_S^0 \pi^\pm$	< 2 %	90%
$\Gamma_9$ $\phi \pi$	< 1.5 %	84%

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=74

DESIG=71;OUR EST;→ UNCHECKED ←

DESIG=73;OUR EST;→ UNCHECKED ←

DESIG=72;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

 **$b_1(1235)$  PARTIAL WIDTHS**

NODE=M011220

$\Gamma(\pi^\pm \gamma)$	VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_2$
<b>230 ± 60</b>		COLLICK 84	SPEC +		200 $\pi^+ Z \rightarrow Z \pi \omega$	

NODE=M011W3


NODE=M011W3

 **$b_1(1235)$  D-wave/S-wave AMPLITUDE RATIO  
IN DECAY OF  $b_1(1235) \rightarrow \omega \pi$** 

NODE=M011DS

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.277 ± 0.027 OUR AVERAGE</b>		Error includes scale factor of 2.4. See the ideogram below.			
0.269 ± 0.009 ± 0.010		NOZAR 02	MPS -		18 $\pi^- p \rightarrow \omega \pi^- p$
0.23 ± 0.03		AMSLER 94C	CBAR		0.0 $\bar{p}p \rightarrow \omega \eta \pi^0$
0.45 ± 0.04		AMSLER 93B	CBAR		0.0 $\bar{p}p \rightarrow \omega \pi^0 \pi^0$
0.235 ± 0.047		ATKINSON 84C	OMEG		20-70 $\gamma p$
0.4 $\begin{smallmatrix} +0.1 \\ -0.1 \end{smallmatrix}$		GESSAROLI 77	HBC -		11 $\pi^- p \rightarrow \pi^- \omega p$
0.21 ± 0.08		CHUNG 75B	HBC +		7.1 $\pi^+ p$
0.3 ± 0.1		CHALOUPKA 74	HBC -		3.9-7.5 $\pi^- p$
0.35 ± 0.25	600	KARSHON 74B	HBC +		4.9 $\pi^+ p$

NODE=M011DS



$b_1(1235)$  D-wave/S-wave amplitude ratio in decay of  $b_1(1235) \rightarrow \omega\pi$

**$b_1(1235)$  D-wave/S-wave AMPLITUDE PHASE DIFFERENCE  
IN DECAY OF  $b_1(1235) \rightarrow \omega\pi$**

NODE=M011PH

VALUE (°)	DOCUMENT ID	TECN	CHG	COMMENT
<b>10.5±2.4±3.9</b>	NOZAR	02	MPS	- 18 $\pi^- p \rightarrow \omega\pi^- p$

NODE=M011PH

**$b_1(1235)$  BRANCHING RATIOS**

NODE=M011230

**$\Gamma(\eta\rho)/\Gamma(\omega\pi)$**

**$\Gamma_3/\Gamma_1$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.10</b>	ATKINSON	84D	OMEG	20-70 $\gamma p$

NODE=M011R9  
NODE=M011R9

**$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\omega\pi)$**

**$\Gamma_4/\Gamma_1$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.5</b>	ABOLINS	63	HBC	+ 3.5 $\pi^+ p$

NODE=M011R1  
NODE=M011R1

**$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$**

**$\Gamma_5/\Gamma$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>seen</b>	<sup>1</sup> ABLIKIM	10E	BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M011R10  
NODE=M011R10

<sup>1</sup> From a fit including ten additional resonances and energy-independent Breit-Wigner width.

NODE=M011R10;LINKAGE=AB

**$\Gamma((K\bar{K})^\pm \pi^0)/\Gamma(\omega\pi)$**

**$\Gamma_6/\Gamma_1$**

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.08</b>	90	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R6  
NODE=M011R6

**$\Gamma(K_S^0 K_L^0 \pi^\pm)/\Gamma(\omega\pi)$**

**$\Gamma_7/\Gamma_1$**

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.06</b>	90	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R8  
NODE=M011R8

**$\Gamma(K_S^0 K_S^0 \pi^\pm)/\Gamma(\omega\pi)$**

**$\Gamma_8/\Gamma_1$**

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.02</b>	90	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R7  
NODE=M011R7

**$\Gamma(\phi\pi)/\Gamma(\omega\pi)$**

**$\Gamma_9/\Gamma_1$**

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.004</b>	95	VIKTOROV	96	SPEC	0 32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$

NODE=M011R4  
NODE=M011R4

••• We do not use the following data for averages, fits, limits, etc. •••

<0.04	95	BIZZARRI	69	HBC	± 0.0 $\bar{p}p$
<0.015		DAHL	67	HBC	1.6-4.2 $\pi^- p$

**$b_1(1235)$  REFERENCES**

ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
NOZAR	02	PL B541 35	M. Nozar <i>et al.</i>	
VIKTOROV	96	PAN 59 1184	V.A. Viktorov <i>et al.</i>	(SERP)
		Translated from YAF 59 1239.		
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
ATKINSON	84C	NP B243 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84D	NP B242 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
COLLICK	84	PRL 53 2374	B. Collick <i>et al.</i>	(MINN, ROCH, FNAL)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+) JP
FLATTE	76C	PL 64B 225	S.M. Flatte <i>et al.</i>	(CERN, AMST, NIJM+) JP
CHUNG	75B	PR D11 2426	S.U. Chung <i>et al.</i>	(BNL, LBL, UCSC) JP
CHALOUPKA	74	PL 51B 407	V. Chaloupka <i>et al.</i>	(CERN) JP
KARSHON	74B	PR D10 3608	U. Karshon <i>et al.</i>	(REHO) JP
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)
BALTAY	67	PRL 18 93	C. Baltay <i>et al.</i>	(COLU)
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL)
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)

NODE=M011

REFID=53361  
REFID=48850  
REFID=45203  
  
REFID=44091  
REFID=43602  
REFID=43585  
REFID=41859  
REFID=41581  
REFID=41004  
REFID=20625  
REFID=20623  
REFID=20624  
REFID=20626  
REFID=20462  
REFID=21265  
REFID=20230  
REFID=20615  
REFID=20613  
REFID=20611  
REFID=20612  
REFID=20171  
REFID=20159  
REFID=20321  
REFID=20006

**$a_1(1260)$**

$$I^G(J^{PC}) = 1^-(1^+ +)$$

See also our review under the  $a_1(1260)$  in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M010

NODE=M010

**$a_1(1260)$  T-MATRIX POLE  $\sqrt{s}$**

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M010PP

NODE=M010PP

NODE=M010PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>(1209 \pm 4^{+12}_-9) - i(288 \pm 6^{+45}_-10)</math></b>			<b>OUR ESTIMATE</b>
$(1209 \pm 4^{+12}_-9) - i(288 \pm 6^{+45}_-10)$	MIKHASENKO 18	RVUE	$\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$

→ UNCHECKED ←

**$a_1(1260)$  MASS**

NODE=M010M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1230 ± 40</b>				<b>OUR ESTIMATE</b>
<b>1299 <math>^{+12}_{-28}</math></b>	46M	1 AGHASYAN	18B	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
1195.05 ± 1.05 ± 6.33	894k	AAIJ	18A1	LHCB $D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
1225 ± 9 ± 20	7k	2 DARGENT	17	RVUE $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1255 ± 6 $^{+7}_{-17}$	420k	3 ALEKSEEV	10	COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1243 ± 12 ± 20		4 AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow \rho^0 \rho^\pm \pi^\mp \gamma$
1230–1270	6360	5 LINK	07A	FOCS $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1203 ± 3		6 GOMEZ-DUM.	04	RVUE $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$
1330 ± 24	90k	SALVINI	04	OBLX $\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
1331 ± 10 ± 3	37k	7 ASNER	00	CLE2 10.6 $e^+ e^- \rightarrow \tau^+ \tau^-$ , $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
1255 ± 7 ± 6	5904	8 ABREU	98G	DLPH $e^+ e^-$
1207 ± 5 ± 8	5904	9 ABREU	98G	DLPH $e^+ e^-$
1196 ± 4 ± 5	5904	10,11 ABREU	98G	DLPH $e^+ e^-$
1240 ± 10		BARBERIS	98B	450 $pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
1262 ± 9 ± 7		8,12 ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94$ , $\tau \rightarrow 3\pi \nu$
1210 ± 7 ± 2		9,12 ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94$ , $\tau \rightarrow 3\pi \nu$
1211 ± 7 $^{+50}_{-0}$		9 ALBRECHT	93C	ARG $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1121 ± 8		13 ANDO	92	SPEC $8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1242 ± 37		14 IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1260 ± 14		15 IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$

NODE=M010M

→ UNCHECKED ←

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2



1250 ± 9	16	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
1208 ± 15		ARMSTRONG	90	OMEG	$300.0 p p \rightarrow$ $p p \pi^+ \pi^- \pi^0$	
1220 ± 15	17	ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1260 ± 25	18	BOWLER	88	RVUE		
1166 ± 18 ± 11		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1164 ± 41 ± 23		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$	OCCUR=2
1250 ± 40	17	TORNQVIST	87	RVUE		
1046 ± 11		ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1056 ± 20 ± 15		RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1194 ± 14 ± 10		SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1255 ± 23		BELLINI	85	SPEC	$40 \pi^- A \rightarrow$ $\pi^- \pi^+ \pi^- A$	
1240 ± 80	19	DANKOWY...	81	SPEC	$8.45 \pi^- p \rightarrow n 3\pi$	
1280 ± 30	19	DAUM	81B	CNTR	$63.94 \pi^- p \rightarrow p 3\pi$	
1041 ± 13	20	GAVILLET	77	HBC	$4.2 K^- p \rightarrow \Sigma 3\pi$	

- 1 Statistical error negligible.
- 2 Reanalysis of CLEO data using Breit-Wigner parameterization.
- 3 Superseded by AGHASYAN 2018B.
- 4 The  $\rho^\pm \pi^\mp$  state can be also due to the  $\pi(1300)$ .
- 5 Using the Breit-Wigner parameterization; strong correlation between mass and width.
- 6 Using the data of BARATE 98R.
- 7 From a fit to the  $3\pi$  mass spectrum including the  $K\bar{K}^*(892)$  threshold.
- 8 Uses the model of KUHN 90.
- 9 Uses the model of ISGUR 89.
- 10 Includes the effect of a possible  $a_1'$  state.
- 11 Uses the model of FEINDT 90.
- 12 Supersedes AKERS 95P.
- 13 Average and spread of values using 2 variants of the model of BOWLER 75.
- 14 Reanalysis of RUCKSTUHL 86.
- 15 Reanalysis of SCHMIDKE 86.
- 16 Reanalysis of ALBRECHT 86B.
- 17 From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.
- 18 From a combined reanalysis of ALBRECHT 86B and DAUM 81B.
- 19 Uses the model of BOWLER 75.
- 20 Produced in  $K^-$  backward scattering.

NODE=M010M;LINKAGE=Q  
 NODE=M010M;LINKAGE=V  
 NODE=M010M;LINKAGE=R  
 NODE=M010M;LINKAGE=AU  
 NODE=M010M;LINKAGE=LI  
 NODE=M010M;LINKAGE=GO  
 NODE=M010M;LINKAGE=B6  
 NODE=M010M;LINKAGE=KS  
 NODE=M010M;LINKAGE=IM  
 NODE=M010M;LINKAGE=A1  
 NODE=M010M;LINKAGE=F1  
 NODE=M010M;LINKAGE=X  
 NODE=M010M;LINKAGE=P  
 NODE=M010M;LINKAGE=I  
 NODE=M010M;LINKAGE=L  
 NODE=M010M;LINKAGE=M  
 NODE=M010M;LINKAGE=K  
 NODE=M010M;LINKAGE=G  
 NODE=M010M;LINKAGE=D  
 NODE=M010M;LINKAGE=F

### $a_1(1260)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>250 to 600 OUR ESTIMATE</b>				
<b>420 ± 35 OUR AVERAGE</b>				
380 ± 80	46M	1 AGHASYAN	18B COMP	$190 \pi^- p \rightarrow$ $\pi^- \pi^+ \pi^- p$
430 ± 24 ± 31		DARGENT	17 RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
422.01 ± 2.10 ± 12.72	894k	AAIJ	18A LHCb	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
367 ± 9 ± 28	420k	2 ALEKSEEV	10 COMP	$190 \pi^- P b \rightarrow$ $\pi^- \pi^- \pi^+ P b'$
410 ± 31 ± 30		3 AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow$ $\rho^0 \rho^\pm \pi^\mp \gamma$
520-680	6360	4 LINK	07A FOCUS	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
480 ± 20		5 GOMEZ-DUM..04	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$
580 ± 41	90k	SALVINI	04 OBLX	$\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
460 ± 85	205	6 DRUTSKOY	02 BELL	$B \rightarrow D(*) K^- K^* 0$
814 ± 36 ± 13	37k	7 ASNER	00 CLE2	$10.6 e^+ e^- \rightarrow$ $\tau^+ \tau^-, \tau^- \rightarrow$ $\pi^- \pi^0 \pi^0 \nu_\tau$
450 ± 50	22k	8 AKHMETSHIN 99E	CMD2	$1.05-1.38 e^+ e^- \rightarrow$ $\pi^+ \pi^- \pi^0 \pi^0$
570 ± 10		9 BONDAR	99 RVUE	$e^+ e^- \rightarrow 4\pi, \tau \rightarrow$ $3\pi \nu_\tau$
587 ± 27 ± 21	5904	10 ABREU	98G DLPH	$e^+ e^-$
478 ± 3 ± 15	5904	11 ABREU	98G DLPH	$e^+ e^-$
425 ± 14 ± 8	5904	12,13 ABREU	98G DLPH	$e^+ e^-$
400 ± 35		BARBERIS	98B	$450 p p \rightarrow$ $p_f \pi^+ \pi^- \pi^0 p_s$
621 ± 32 ± 58		10,14 ACKERSTAFF 97R	OPAL	$E_{cm}^{e^+e^-} = 88-94, \tau \rightarrow$ $3\pi \nu$

NODE=M010W  
 NODE=M010W  
 → UNCHECKED ←

OCCUR=2  
 OCCUR=3

457	$\pm 15$	$\pm 17$	11,14	ACKERSTAFF	97R	OPAL	$E_{cm}^{ee} = 88-94, \tau \rightarrow 3\pi\nu$	OCCUR=2
446	$\pm 21$	$+140$ $-0$	11	ALBRECHT	93C	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
239	$\pm 11$			ANDO	92	SPEC	$8 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^0 n$	
266	$\pm 13$	$\pm 4$	15	ANDO	92	SPEC	$8 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^0 n$	OCCUR=3
465	$+228$ $-143$		16	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	
298	$+40$ $-34$		17	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=2
488	$\pm 32$		18	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
430	$\pm 50$			ARMSTRONG	90	OMEG	$300.0 p p \rightarrow p p \pi^+ \pi^- \pi^0$	
420	$\pm 40$		19	ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
396	$\pm 43$		20	BOWLER	88	RVUE		
405	$\pm 75$	$\pm 25$		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
419	$\pm 108$	$\pm 57$		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$	OCCUR=2
521	$\pm 27$			ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
476	$+132$ $-120$	$\pm 54$		RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
462	$\pm 56$	$\pm 30$		SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
292	$\pm 40$			BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
380	$\pm 100$		21	DANKOWY...	81	SPEC	$8.45 \pi^- \rho \rightarrow n 3\pi$	
300	$\pm 50$		21	DAUM	81B	CNTR	$63,94 \pi^- \rho \rightarrow p 3\pi$	
230	$\pm 50$		22	GAVILLET	77	HBC	$4.2 K^- p \rightarrow \Sigma 3\pi$	

<sup>1</sup> Statistical error negligible.

<sup>2</sup> Superseded by AGHASYAN 2018B.

<sup>3</sup> The  $\rho^\pm \pi^\mp$  state can be also due to the  $\pi(1300)$ .

<sup>4</sup> Using the Breit-Wigner parameterization; strong correlation between mass and width.

<sup>5</sup> Using the data of BARATE 98R.

<sup>6</sup> From a fit of the  $K^- K^{*0}$  distribution assuming  $m_{a_1} = 1230$  MeV and purely resonant production of the  $K^- K^{*0}$  system.

<sup>7</sup> From a fit to the  $3\pi$  mass spectrum including the  $K\bar{K}^*(892)$  threshold.

<sup>8</sup> Using the  $a_1(1260)$  mass of 1230 MeV.

<sup>9</sup> From AKHMETSHIN 99E and ASNER 00 data using the  $a_1(1260)$  mass of 1230 MeV.

<sup>10</sup> Uses the model of KUHNS 90.

<sup>11</sup> Uses the model of ISGUR 89.

<sup>12</sup> Includes the effect of a possible  $a_1'$  state.

<sup>13</sup> Uses the model of FEINDT 90.

<sup>14</sup> Supersedes AKERS 95P.

<sup>15</sup> Average and spread of values using 2 variants of the model of BOWLER 75.

<sup>16</sup> Reanalysis of RUCKSTUHL 86.

<sup>17</sup> Reanalysis of SCHMIDKE 86.

<sup>18</sup> Reanalysis of ALBRECHT 86B.

<sup>19</sup> From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

<sup>20</sup> From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

<sup>21</sup> Uses the model of BOWLER 75.

<sup>22</sup> Produced in  $K^-$  backward scattering.

NODE=M010W;LINKAGE=Q  
 NODE=M010W;LINKAGE=R  
 NODE=M010W;LINKAGE=AU  
 NODE=M010W;LINKAGE=LI  
 NODE=M010W;LINKAGE=GO  
 NODE=M010W;LINKAGE=DR

NODE=M010W;LINKAGE=B6  
 NODE=M010W;LINKAGE=WE  
 NODE=M010W;LINKAGE=WB  
 NODE=M010W;LINKAGE=KS  
 NODE=M010W;LINKAGE=IM  
 NODE=M010W;LINKAGE=A1  
 NODE=M010W;LINKAGE=F1  
 NODE=M010W;LINKAGE=X  
 NODE=M010W;LINKAGE=P  
 NODE=M010W;LINKAGE=I  
 NODE=M010W;LINKAGE=L  
 NODE=M010W;LINKAGE=M  
 NODE=M010W;LINKAGE=K  
 NODE=M010W;LINKAGE=G  
 NODE=M010W;LINKAGE=D  
 NODE=M010W;LINKAGE=F

## $a_1(1260)$ DECAY MODES

NODE=M010215;NODE=M010

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $3\pi$	seen
$\Gamma_2$ $(\rho\pi)S$ -wave, $\rho \rightarrow \pi\pi$	seen
$\Gamma_3$ $(\rho\pi)D$ -wave, $\rho \rightarrow \pi\pi$	seen
$\Gamma_4$ $(\rho(1450)\pi)S$ -wave, $\rho \rightarrow \pi\pi$	seen
$\Gamma_5$ $(\rho(1450)\pi)D$ -wave, $\rho \rightarrow \pi\pi$	seen
$\Gamma_6$ $f_0(500)\pi$ , $f_0 \rightarrow \pi\pi$	seen
$\Gamma_7$ $f_0(980)\pi$ , $f_0 \rightarrow \pi\pi$	seen
$\Gamma_8$ $f_0(1370)\pi$ , $f_0 \rightarrow \pi\pi$	seen
$\Gamma_9$ $f_2(1270)\pi$ , $f_2 \rightarrow \pi\pi$	seen
$\Gamma_{10}$ $\pi^+ \pi^- \pi^0$	seen
$\Gamma_{11}$ $\pi^0 \pi^0 \pi^0$	not seen
$\Gamma_{12}$ $K K \pi$	seen
$\Gamma_{13}$ $K^*(892) K$	seen
$\Gamma_{14}$ $\pi \gamma$	seen

DESIG=17;OUR EST;→ UNCHECKED ←  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=8;OUR EST;→ UNCHECKED ←  
 DESIG=9;OUR EST;→ UNCHECKED ←  
 DESIG=10;OUR EST;→ UNCHECKED ←  
 DESIG=16;OUR EST;→ UNCHECKED ←  
 DESIG=11  
 DESIG=12;OUR EST;→ UNCHECKED ←  
 DESIG=13;OUR EST;→ UNCHECKED ←  
 DESIG=22  
 DESIG=23;OUR EST;→ UNCHECKED ←  
 DESIG=18;OUR EST;→ UNCHECKED ←  
 DESIG=14;OUR EST;→ UNCHECKED ←  
 DESIG=4;OUR EST;→ UNCHECKED ←

**$a_1(1260)$  PARTIAL WIDTHS**

$\Gamma(\pi\gamma)$

$\Gamma_{14}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>640 ± 246</b>	ZIELINSKI	84C SPEC	200 $\pi^+ Z \rightarrow Z 3\pi$

NODE=M010220

NODE=M010W4  
NODE=M010W4

**D-wave/S-wave AMPLITUDE RATIO IN DECAY OF  $a_1(1260) \rightarrow \rho\pi$**

NODE=M010DS


VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.062 ± 0.020 OUR AVERAGE</b>	Error includes scale factor of 2.3. See the ideogram below.		

NODE=M010DS

-0.043 ± 0.009 ± 0.005	LINK	07A	FOCS $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
-0.14 ± 0.04 ± 0.07	<sup>1</sup> CHUNG	02	B852 $18.3 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^- \rho$
-0.10 ± 0.02 ± 0.02	<sup>2,3</sup> ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94, \tau \rightarrow 3\pi\nu$
-0.11 ± 0.02	<sup>2</sup> ALBRECHT	93C	ARG $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$

NODE=M010DS;LINKAGE=C  
NODE=M010DS;LINKAGE=IM  
NODE=M010DS;LINKAGE=X

- <sup>1</sup> Deck-type background not subtracted.
- <sup>2</sup> Uses the model of ISGUR 89.
- <sup>3</sup> Supersedes AKERS 95P.



**$a_1(1260)$  BRANCHING RATIOS**

NODE=M010225

$\Gamma((\rho\pi)_{S\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
60.19	37k	<sup>1</sup> ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$ , $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$

NODE=M010R5  
NODE=M010R5

- <sup>1</sup> From a fit to the Dalitz plot.

NODE=M010R5;LINKAGE=B1

$\Gamma((\rho\pi)_{D\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$

$\Gamma_3/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
1.30 ± 0.60 ± 0.22	37k	<sup>1</sup> ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$ , $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$

NODE=M010R6  
NODE=M010R6

- <sup>1</sup> From a fit to the Dalitz plot.

NODE=M010R6;LINKAGE=B1

$\Gamma((\rho(1450)\pi)_{S\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
0.56 ± 0.84 ± 0.32	37k	<sup>1,2</sup> ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$ , $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$

NODE=M010R7  
NODE=M010R7

- <sup>1</sup> From a fit to the Dalitz plot.
- <sup>2</sup> Assuming for  $\rho(1450)$  mass and width of 1370 and 386 MeV respectively.

NODE=M010R7;LINKAGE=B1  
NODE=M010R7;LINKAGE=B2

$\Gamma((\rho(1450)\pi)_{D\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.04 \pm 1.20 \pm 0.28$	37k	<sup>1,2</sup> ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$ , $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
--------------------------	-----	----------------------	----	---

<sup>1</sup> From a fit to the Dalitz plot.

<sup>2</sup> Assuming for  $\rho(1450)$  mass and width of 1370 and 386 MeV respectively.

NODE=M010R8  
NODE=M010R8

NODE=M010R8;LINKAGE=B1  
NODE=M010R8;LINKAGE=B2

 $\Gamma(f_0(500)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen		CHUNG	02	B852 18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$
$18.76 \pm 4.29 \pm 1.48$	37k	<sup>1,2</sup> ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$ , $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

<sup>1</sup> From a fit to the Dalitz plot.

<sup>2</sup> Assuming for  $f_0(500)$  ( $\sigma$ ) mass and width of 860 and 880 MeV respectively.

NODE=M010R9  
NODE=M010R9

NODE=M010R9;LINKAGE=B1  
NODE=M010R9;LINKAGE=B3

 $\Gamma(f_0(500)\pi, f_0 \rightarrow \pi\pi)/\Gamma((\rho\pi)_{S\text{-wave}, \rho \rightarrow \pi\pi})$   $\Gamma_6/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06 \pm 0.05$	90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+2\pi^-$
$\sim 0.3$	28k	AKHMETSHIN 99E	CMD2	$1.05\text{--}1.38 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
$0.003 \pm 0.003$		<sup>1</sup> LONGACRE	82	RVUE

<sup>1</sup> Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from GAVIL-LET 77, DAUM 80, and DANKOWYCH 81.

NODE=M010R4  
NODE=M010R4

NODE=M010R4;LINKAGE=E

 $\Gamma(f_0(980)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen <sup>1</sup> ALEXEEV 21 COMP  $\pi^-p \rightarrow \pi^-\pi^+\pi^-p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	37k	ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$ , $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
----------	-----	-------	----	---

<sup>1</sup> The  $a_1(1260)^- \rightarrow f_0(980)\pi^-$  decay mode via the Triangle Singularity mechanism from MIKHASENKO 15 and ACETI 16 explains the  $a_1(1420)^-$  signal observed by ADOLPH 15C.

NODE=M010R10  
NODE=M010R10

NODE=M010R10;LINKAGE=A

 $\Gamma(f_0(1370)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.40 \pm 2.71 \pm 1.26$	37k	<sup>1,2</sup> ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$ , $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
--------------------------	-----	----------------------	----	---

<sup>1</sup> From a fit to the Dalitz plot.

<sup>2</sup> Assuming for  $f_0(1370)$  mass and width of 1186 and 350 MeV respectively.

NODE=M010R11  
NODE=M010R11

NODE=M010R11;LINKAGE=B1  
NODE=M010R11;LINKAGE=B4

 $\Gamma(f_2(1270)\pi, f_2 \rightarrow \pi\pi)/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.19 \pm 0.49 \pm 0.17$	37k	<sup>1,2</sup> ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$ , $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
--------------------------	-----	----------------------	----	---

<sup>1</sup> From a fit to the Dalitz plot.

<sup>2</sup> Assuming for  $f_2(1270)$  mass and width of 1275 and 185 MeV respectively.

NODE=M010R12  
NODE=M010R12

NODE=M010R12;LINKAGE=B1  
NODE=M010R12;LINKAGE=B5

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{10}/\Gamma$ 

VALUE	DOCUMENT ID	COMMENT
-------	-------------	---------

seen BARBERIS 98B 450  $pp \rightarrow p_f\pi^+\pi^-\pi^0p_s$

NODE=M010R00  
NODE=M010R00

 $\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$   $\Gamma_{11}/\Gamma_{10}$ 

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 0.008$	90	<sup>1</sup> BARBERIS	01	450 $pp \rightarrow p_f3\pi^0p_s$
-----------	----	-----------------------	----	-----------------------------------

<sup>1</sup> Inconsistent with observations of  $\sigma\pi$ ,  $f_0(1370)\pi$ , and  $f_2(1270)\pi$  decay modes.

NODE=M010R15  
NODE=M010R15

NODE=M010R15;LINKAGE=RB

$\Gamma(K^*(892)K)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2±0.5	2255	<sup>1</sup> COAN	04	CLEO $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$
8 to 15	205	<sup>2</sup> DRUTSKOY	02	BELL $B \rightarrow D^{(*)} K^- K^{*0}$
3.3±0.5±0.1	37k	<sup>3</sup> ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$ , $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
2.6±0.3		<sup>4</sup> BARATE	99R	ALEP $\tau \rightarrow K \bar{K} \pi \nu_\tau$

<sup>1</sup> Using structure functions from KUHN 92 and DECKER 93A and  $B(\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau) = (0.155 \pm 0.006 \pm 0.009)\%$  from BRIERE 03.

<sup>2</sup> From a comparison to ALAM 94 assuming purely resonant production of the  $K^- K^{*0}$  system.

<sup>3</sup> From a fit to the  $3\pi$  mass spectrum including the  $K \bar{K}^*(892)$  threshold.

<sup>4</sup> Assuming  $a_1(1260)$  dominance and taking  $B(\tau \rightarrow a_1(1260) \nu_\tau)$  from BUSKULIC 96.

NODE=M010R13  
NODE=M010R13

NODE=M010R13;LINKAGE=CO

NODE=M010R13;LINKAGE=DR

NODE=M010R13;LINKAGE=B6

NODE=M010R13;LINKAGE=BA

### $a_1(1260)$ REFERENCES

ALEXEEV	21	PRL 127 082501	G.D. Alexeev <i>et al.</i>	(COMPASS Collab.)
AAJ	18A1	EPJ C78 443	R. Aajj <i>et al.</i>	(LHCb Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
MIKHASENKO	18	PR D98 096021	M. Mikhasenko <i>et al.</i>	(JPAC Collab.)
DARGENT	17	JHEP 1705 143	P. dArgent <i>et al.</i>	(HEID, BRIS)
ACETI	16	PR D94 096015	F. Aceti, L.R. Dai, E. Oset	(IFIC, LNUDA)
ADOLPH	15C	PRL 115 082001	C. Adolph <i>et al.</i>	(COMPASS Collab.)
MIKHASENKO	15	PR D91 094015	M. Mikhasenko, B. Ketzer, A. Sarantsev	(BONN+)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
LINK	07A	PR D75 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)
GOMEZ-DUM...	04	PR D69 073002	D. Gomez Dumm, A. Pich, J. Portoles	
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)
BRIERE	03	PRL 90 181802	R. A. Briere <i>et al.</i>	(CLEO Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>	
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AKHMETSHIN	99E	PL B466 392	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)
BONDAR	99	PL B466 403	A.E. Bondar <i>et al.</i>	(Novosibirsk CMD-2 Collab.)
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
BARATE	98R	EPJ C4 409	R. Barate <i>et al.</i>	(ALEPH Collab.)
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BUSKULIC	96	ZPHY C70 579	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
AKERS	95P	ZPHY C67 45	R. Akers <i>et al.</i>	(OPAL Collab.)
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)
ALBRECHT	93C	ZPHY C58 61	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
DECKER	93A	ZPHY C58 445	R. Decker <i>et al.</i>	
ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
KUHN	92	ZPHY C56 661	J.H. Kuhn, E. Mirkes	
IVANOV	91	ZPHY C49 563	Y.P. Ivanov, A.A. Osipov, M.K. Volkov	(JINR)
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch	(WA76 Coll.)
FEINDT	90	ZPHY C48 681	M. Feindt	(HAMB)
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)
ISGUR	89	PR D39 1357	N. Isgur, C. Morningstar, C. Reader	(TNTO)
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)
BAND	87	PL B198 297	H.R. Band <i>et al.</i>	(MAC Collab.)
TORNQVIST	87	ZPHY C36 695	N.A. Tornqvist	(HELS)
ALBRECHT	86B	ZPHY C33 7	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
RUCKSTUHL	86	PRL 56 2132	W. Ruckstuhl <i>et al.</i>	(DELCO Collab.)
SCHMIDKE	86	PRL 57 527	W.B. Schmidke <i>et al.</i>	(Mark II Collab.)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
ZIELINSKI	84C	PRL 52 1195	M. Zielinski <i>et al.</i>	(ROCH, MINN, FNAL)
LONGACRE	82	PR D26 82	R.S. Longacre	(BNL)
DANKOWY...	81	PRL 46 580	J.A. Dankowycz <i>et al.</i>	(TNTO, BNL, CARL+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
DAUM	80	PL B9B 281	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
GAVILLET	77	PL 69B 119	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+) JP
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

NODE=M010

REFID=62008

REFID=59187

REFID=59471

REFID=59487

REFID=58121

REFID=62015

REFID=56790

REFID=56825

REFID=53356

REFID=52049

REFID=51713

REFID=51004

REFID=49945

REFID=49771

REFID=53226

REFID=49360

REFID=48837

REFID=48780

REFID=48324

REFID=47339

REFID=47411

REFID=47366

REFID=47358

REFID=45909

REFID=46147

REFID=46345

REFID=45616

REFID=44588

REFID=44366

REFID=43738

REFID=43310

REFID=51577

REFID=43171

REFID=51576

REFID=41750

REFID=41375

REFID=45912

REFID=45862

REFID=40730

REFID=40578

REFID=40263

REFID=40030

REFID=20884

REFID=10349

REFID=10350

REFID=47490

REFID=20882

REFID=20878

REFID=20572

REFID=20872

REFID=20868

REFID=20852

REFID=20571

$f_2(1270)$ 

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M005

 $f_2(1270)$  T-MATRIX POLE  $\sqrt{s}$ 

NODE=M005PP

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M005PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1260–1283) – i (90–110) OUR ESTIMATE</b>			
$(1268 \pm 8) - i (101 \pm 6)$	RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1263.3 \pm 0.2 \pm 1.5) - i$ $(96.9 \pm 0.2 \pm 0.8)$	ALBRECHT	20	RVUE $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta,$ $\pi^0K^+K^-$
$(1270 \pm 8) - i (97 \pm 8)$	<sup>1</sup> ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1278 \pm 5) - i (102 \pm 10)$	<sup>1</sup> BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$

NODE=M005PP

→ UNCHECKED ←

<sup>1</sup> Amplitude did not include dispersive corrections.

NODE=M005PP;LINKAGE=A

 $f_2(1270)$  MASS

NODE=M005M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1275.4 ± 0.8 OUR AVERAGE</b>				
OUR 2022 AVERAGE] Error includes scale factor of 1.1. [1275.5 ± 0.8 MeV				
$1275.8 \pm 1.0 \pm 0.4$		<sup>1</sup> BOGOLYUB...	13	SPEC $7\pi^+(K^+,p)A \rightarrow n\gamma + X$
$1262 \pm \frac{1}{2} \pm 8$		<sup>2</sup> ABLIKIM	06v	BES2 $e^+e^- \rightarrow J/\psi \rightarrow$ $\gamma\pi^+\pi^-$
$1275 \pm 15$		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$
$1283 \pm 5$		ALDE	98	GAM4 $100 \pi^- p \rightarrow \pi^0\pi^0 n$
$1272 \pm 8$	200k	PROKOSHKIN	94	GAM2 $38 \pi^- p \rightarrow \pi^0\pi^0 n$
$1269.7 \pm 5.2$	5730	AUGUSTIN	89	DM2 $e^+e^- \rightarrow 5\pi$
$1283 \pm 8$	400	<sup>3</sup> ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$
$1274 \pm 5$		<sup>3</sup> AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
$1283 \pm 6$		<sup>4</sup> LONGACRE	86	MPS $22 \pi^- p \rightarrow n2K_S^0$
$1276 \pm 7$		COURAU	84	DLCO $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
$1273.3 \pm 2.3$		<sup>5</sup> CHABAUD	83	ASPK $17 \pi^- p$ polarized
$1280 \pm 4$		<sup>6</sup> CASON	82	STRC $8 \pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$
$1281 \pm 7$	11600	GIDAL	81	MRK2 $J/\psi$ decay
$1282 \pm 5$		<sup>7</sup> CORDEN	79	OMEG $12-15 \pi^- p \rightarrow n2\pi$
$1269 \pm 4$	10k	APEL	75	NICE $40 \pi^- p \rightarrow n2\pi^0$
$1272 \pm 4$	4600	ENGLER	74	DBC $6 \pi^+ n \rightarrow \pi^+\pi^- p$
$1277 \pm 4$	5300	FLATTE	71	HBC $7.0 \pi^+ p$
$1273 \pm 8$		<sup>3</sup> STUNTEBECK	70	HBC $8 \pi^- p, 5.4 \pi^+ d$
$1265 \pm 8$		BOESEBECK	68	HBC $8 \pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1257 \pm 6$		<sup>8</sup> KLEMP	22	RVUE $J/\psi(1S) \rightarrow \gamma\pi^0\pi^0,$ $\gamma K_S^0 K_S^0$
$1263 \pm 12$		CARVER	21	CLAS $\gamma p \rightarrow \pi^0\pi^0 p$
$1259 \pm 4 \pm 4$	1.7k	<sup>9,10</sup> DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1267 \pm 4 \pm 3$	1.5k	<sup>9,10</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
$1277 \pm 6$	870	<sup>11</sup> SCHEGELSKY	06A	RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
$1251 \pm 10$		TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
$1260 \pm 10$		<sup>12</sup> ALDE	97	GAM2 $450 pp \rightarrow pp\pi^0\pi^0$
$1278 \pm 6$		<sup>12</sup> GRYGOREV	96	SPEC $40 \pi^- N \rightarrow K_S^0 K_S^0 X$
$1262 \pm 11$		AGUILAR...	91	EHS $400 pp$
$1275 \pm 10$		AKER	91	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$
$1220 \pm 10$		BREAKSTONE	90	SFM $pp \rightarrow pp\pi^+\pi^-$
$1288 \pm 12$		ABACHI	86B	HRS $e^+e^- \rightarrow \pi^+\pi^- X$
$1284 \pm 30$	3k	BINON	83	GAM2 $38 \pi^- p \rightarrow n2\eta$
$1280 \pm 20$	3k	APEL	82	CNTR $25 \pi^- p \rightarrow n2\pi^0$
$1284 \pm 10$	16000	DEUTSCH...	76	HBC $16 \pi^+ p$
$1258 \pm 10$	600	TAKAHASHI	72	HBC $8 \pi^- p \rightarrow n2\pi$
$1275 \pm 13$		ARMENISE	70	HBC $9 \pi^+ n \rightarrow p\pi^+\pi^-$
$1261 \pm 5$	1960	<sup>3</sup> ARMENISE	68	DBC $5.1 \pi^+ n \rightarrow p\pi^+ MM^-$
$1270 \pm 10$	360	<sup>3</sup> ARMENISE	68	DBC $5.1 \pi^+ n \rightarrow p\pi^0 MM$
$1268 \pm 6$		<sup>13</sup> JOHNSON	68	HBC $3.7-4.2 \pi^- p$

NODE=M005M

NEW

OCCUR=2

OCCUR=2

OCCUR=2

- <sup>1</sup> Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
- <sup>2</sup> Breit-Wigner mass.
- <sup>3</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.
- <sup>4</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- <sup>5</sup> From an energy-independent partial-wave analysis.
- <sup>6</sup> From an amplitude analysis of the reaction  $\pi^+\pi^- \rightarrow 2\pi^0$ .
- <sup>7</sup> From an amplitude analysis of  $\pi^+\pi^- \rightarrow \pi^+\pi^-$  scattering data.
- <sup>8</sup> Fit of the tensor partial waves from BES3 in the multipole basis.
- <sup>9</sup> Using CLEO-c data but not authored by the CLEO Collaboration.
- <sup>10</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 185$  MeV.
- <sup>11</sup> From analysis of L3 data at 91 and 183–209 GeV.
- <sup>12</sup> Systematic uncertainties not estimated.
- <sup>13</sup> JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

NODE=M005M;LINKAGE=B

NODE=M005M;LINKAGE=K

NODE=M005M;LINKAGE=T

NODE=M005M;LINKAGE=L

NODE=M005M;LINKAGE=O

NODE=M005M;LINKAGE=P

NODE=M005M;LINKAGE=S

NODE=M005M;LINKAGE=Q

NODE=M005M;LINKAGE=C

NODE=M005M;LINKAGE=D

NODE=M005M;LINKAGE=SC

NODE=M005M;LINKAGE=QQ

NODE=M005M;LINKAGE=J

 **$f_2(1270)$  WIDTH**

NODE=M005W

VALUE (MeV)    EVTS    DOCUMENT ID    TECN    COMMENT  
**186.6 $\pm$  2.3 OUR FIT** Error includes scale factor of 1.5. [186.7 $^{+2.2}_{-2.5}$  MeV OUR 2022  
FIT Scale factor = 1.4]

NODE=M005W

NEW

**185.8 $^{+2.8}_{-2.1}$  OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.  
[185.9 $^{+2.8}_{-2.1}$  MeV OUR 2022 AVERAGE Scale factor = 1.6]

NEW


190.3 $\pm$ 1.9 $\pm$ 1.8		<sup>1</sup> BOGOLYUB...	13	SPEC	$7\pi^+(K^+,p)A \rightarrow n\gamma + X$	
175 $^{+6}_{-4}$ $\pm$ 10		<sup>2</sup> ABLIKIM	06v	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$	
190 $\pm$ 20		ABLIKIM	05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$	
171 $\pm$ 10		ALDE	98	GAM4	$100\pi^-p \rightarrow \pi^0\pi^0n$	
192 $\pm$ 5	200k	PROKOSHKIN	94	GAM2	$38\pi^-p \rightarrow \pi^0\pi^0n$	
180 $\pm$ 24		AGUILAR-...	91	EHS	400 $pp$	
169 $\pm$ 9	5730	<sup>3</sup> AUGUSTIN	89	DM2	$e^+e^- \rightarrow 5\pi$	
150 $\pm$ 30	400	<sup>3</sup> ALDE	87	GAM4	$100\pi^-p \rightarrow 4\pi^0n$	
186 $^{+9}_{-2}$		<sup>4</sup> LONGACRE	86	MPS	$22\pi^-p \rightarrow n2K_S^0$	
179.2 $^{+6.9}_{-6.6}$		<sup>5</sup> CHABAUD	83	ASPK	$17\pi^-p$ polarized	
160 $\pm$ 11		DENNEY	83	LASS	$10\pi^+N$	
196 $\pm$ 10	3k	APEL	82	CNTR	$25\pi^-p \rightarrow n2\pi^0$	
152 $\pm$ 9		<sup>6</sup> CASON	82	STRC	$8\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$	
186 $\pm$ 27	11600	GIDAL	81	MRK2	$J/\psi$ decay	
216 $\pm$ 13		<sup>7</sup> CORDEN	79	OMEG	$12-15\pi^-p \rightarrow n2\pi$	
190 $\pm$ 10	10k	APEL	75	NICE	$40\pi^-p \rightarrow n2\pi^0$	
192 $\pm$ 16	4600	ENGLER	74	DBC	$6\pi^+n \rightarrow \pi^+\pi^-p$	
183 $\pm$ 15	5300	FLATTE	71	HBC	$7\pi^+p \rightarrow \Delta^{++}f_2$	
196 $\pm$ 30		<sup>3</sup> STUNTEBECK	70	HBC	$8\pi^-p, 5.4\pi^+d$	
216 $\pm$ 20	1960	<sup>3</sup> ARMENISE	68	DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$	OCCUR=2
128 $\pm$ 27		<sup>3</sup> BOESEBECK	68	HBC	$8\pi^+p$	
176 $\pm$ 21		<sup>3,8</sup> JOHNSON	68	HBC	$3.7-4.2\pi^-p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

168 $\pm$ 7		<sup>9</sup> KLEMP	22	RVUE	$J/\psi(1S) \rightarrow \gamma\pi^0\pi^0,$ $\gamma K_S^0 K_S^0$	
183 $\pm$ 2		CARVER	21	CLAS	$\gamma p \rightarrow \pi^0\pi^0p$	
195 $\pm$ 15	870	<sup>10</sup> SCHEGELSKY	06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
121 $\pm$ 26		TIKHOMIROV	03	SPEC	$40.0\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$	
187 $\pm$ 20		<sup>11</sup> ALDE	97	GAM2	$450pp \rightarrow pp\pi^0\pi^0$	
184 $\pm$ 10		<sup>11</sup> GRYGOREV	96	SPEC	$40\pi^-N \rightarrow K_S^0 K_S^0 X$	
200 $\pm$ 10		AKER	91	CBAR	$0.0\bar{p}p \rightarrow 3\pi^0$	
240 $\pm$ 40	3k	BINON	83	GAM2	$38\pi^-p \rightarrow n2\eta$	
187 $\pm$ 30	650	<sup>3</sup> ANTIPOV	77	CIBS	$25\pi^-p \rightarrow p3\pi$	
225 $\pm$ 38	16000	DEUTSCH...	76	HBC	$16\pi^+p$	
166 $\pm$ 28	600	<sup>3</sup> TAKAHASHI	72	HBC	$8\pi^-p \rightarrow n2\pi$	
173 $\pm$ 53		<sup>3</sup> ARMENISE	70	HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$	OCCUR=2

- 1 Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
- 2 Breit-Wigner width
- 3 Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.
- 4 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- 5 From an energy-independent partial-wave analysis.
- 6 From an amplitude analysis of the reaction  $\pi^+\pi^- \rightarrow 2\pi^0$ .
- 7 From an amplitude analysis of  $\pi^+\pi^- \rightarrow \pi^+\pi^-$  scattering data.
- 8 JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.
- 9 Fit of the tensor partial waves from BES3 in the multipole basis.
- 10 From analysis of L3 data at 91 and 183–209 GeV.
- 11 Systematic uncertainties not estimated.

NODE=M005W;LINKAGE=C  
 NODE=M005W;LINKAGE=D  
 NODE=M005W;LINKAGE=T  
 NODE=M005W;LINKAGE=L  
 NODE=M005W;LINKAGE=R  
 NODE=M005W;LINKAGE=Q  
 NODE=M005W;LINKAGE=U  
 NODE=M005W;LINKAGE=J  
 NODE=M005W;LINKAGE=M  
 NODE=M005W;LINKAGE=SC  
 NODE=M005W;LINKAGE=QQ



**$f_2(1270)$  DECAY MODES**

NODE=M005215;NODE=M005

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\pi\pi$	(84.3 $^{+2.9}_{-0.9}$ ) %	S=1.2
$\Gamma_2$ $\pi^+\pi^-2\pi^0$	( 7.7 $^{+1.1}_{-3.2}$ ) %	S=1.2
$\Gamma_3$ $K\bar{K}$	( 4.6 $\pm 0.4$ ) %	S=2.7
$\Gamma_4$ $2\pi^+2\pi^-$	( 2.8 $\pm 0.4$ ) %	S=1.2
$\Gamma_5$ $\eta\eta$	( 4.0 $\pm 0.8$ ) $\times 10^{-3}$	S=2.1
$\Gamma_6$ $4\pi^0$	( 3.0 $\pm 1.0$ ) $\times 10^{-3}$	
$\Gamma_7$ $\gamma\gamma$	( 1.42 $\pm 0.24$ ) $\times 10^{-5}$	S=1.4
$\Gamma_8$ $\eta\pi\pi$	< 8 $\times 10^{-3}$	CL=95%
$\Gamma_9$ $K^0K^-\pi^+ + c.c.$	< 3.4 $\times 10^{-3}$	CL=95%
$\Gamma_{10}$ $e^+e^-$	< 6 $\times 10^{-10}$	CL=90%

DESIG=1  
 DESIG=3  
 DESIG=4  
 DESIG=2  
 DESIG=7  
 DESIG=9  
 DESIG=8  
 DESIG=6  
 DESIG=5  
 DESIG=10



## CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 44 measurements and one constraint to determine 8 parameters. The overall fit has a  $\chi^2 = 82.3$  for 37 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-91						
$x_3$	10	-39					
$x_4$	10	-38	1				
$x_5$	1	-6	0	0			
$x_6$	0	-7	0	0	0		
$x_7$	4	1	-15	0	0	0	
$\Gamma$	-71	65	-10	-7	-1	0	-6
	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$	$x_7$

Mode	Rate (MeV)	Scale factor
$\Gamma_1$ $\pi\pi$	157.2 $^{+4.0}_{-1.1}$	
$\Gamma_2$ $\pi^+\pi^-2\pi^0$	14.3 $^{+2.2}_{-6.0}$	1.2
$\Gamma_3$ $K\bar{K}$	8.5 $\pm 0.8$	2.8
$\Gamma_4$ $2\pi^+2\pi^-$	5.2 $\pm 0.7$	1.2
$\Gamma_5$ $\eta\eta$	0.75 $\pm 0.14$	2.1
$\Gamma_6$ $4\pi^0$	0.56 $\pm 0.19$	
$\Gamma_7$ $\gamma\gamma$	0.0026 $\pm 0.0005$	1.4

### $f_2(1270)$ PARTIAL WIDTHS

NODE=M005220

#### $\Gamma(\pi\pi)$

 $\Gamma_1$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W1  
NODE=M005W1

**157.2 $^{+4.0}_{-1.1}$  OUR FIT**

**157.0 $^{+6.0}_{-1.0}$**

<sup>1</sup> LONGACRE 86 MPS 22  $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

152  $\pm 8$  870 <sup>2</sup> SCHEGELSKY 06A RVUE  $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005W1;LINKAGE=L  
NODE=M005W1;LINKAGE=SC

#### $\Gamma(K\bar{K})$

 $\Gamma_3$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W4  
NODE=M005W4

**8.5 $\pm 0.8$  OUR FIT** Error includes scale factor of 2.8.

**9.0 $^{+0.7}_{-0.3}$**

<sup>1</sup> LONGACRE 86 MPS 22  $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5  $\pm 2.0$  870 <sup>2</sup> SCHEGELSKY 06A RVUE  $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005W4;LINKAGE=L  
NODE=M005W4;LINKAGE=SC

#### $\Gamma(\eta\eta)$

 $\Gamma_5$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W7  
NODE=M005W7

**0.75 $\pm 0.14$  OUR FIT** Error includes scale factor of 2.1.

**1.0  $\pm 0.1$**

<sup>1</sup> LONGACRE 86 MPS 22  $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8  $\pm 0.4$  870 <sup>2</sup> SCHEGELSKY 06A RVUE  $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005W7;LINKAGE=L  
NODE=M005W7;LINKAGE=SC

$\Gamma(\gamma\gamma)$  $\Gamma_7$ 

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an  $S$ -wave contribution, values are systematically higher (typically around 3 keV).

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6 ± 0.5 OUR FIT</b>	Error includes scale factor of 1.4.			
<b>2.93 ± 0.40</b>		<sup>1</sup> DAI	14A	RVUE Compilation
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3.14 ± 0.20		<sup>2,3</sup> PENNINGTON 08		RVUE Compilation
3.82 ± 0.30		<sup>3,4</sup> PENNINGTON 08		RVUE Compilation
2.55 ± 0.15	870	<sup>5</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
2.84 ± 0.35		BOGLIONE 99	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
2.93 ± 0.23 ± 0.32		<sup>6</sup> YABUKI 95	VNS	
2.58 ± 0.13 <sup>+0.36</sup> <sub>-0.27</sub>		<sup>7</sup> BEHREND 92	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.10 ± 0.35 ± 0.35		<sup>8</sup> BLINOV 92	MD1	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.27 ± 0.47 ± 0.11		ADACHI 90D	TOPZ	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.15 ± 0.04 ± 0.39		BOYER 90	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.19 ± 0.16 <sup>+0.29</sup> <sub>-0.28</sub>		MARSISKE 90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
2.35 ± 0.65		<sup>9</sup> MORGAN 90	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
3.19 ± 0.09 <sup>+0.22</sup> <sub>-0.38</sub>	2177	OEST 90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
3.2 ± 0.1 ± 0.4		<sup>10</sup> AIHARA 86B	TPC	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.5 ± 0.1 ± 0.5		BEHREND 84B	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.85 ± 0.25 ± 0.5		<sup>11</sup> BERGER 84	PLUT	$e^+ e^- \rightarrow e^+ e^- 2\pi$
2.70 ± 0.05 ± 0.20		COURAU 84	DLCO	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.52 ± 0.13 ± 0.38		<sup>12</sup> SMITH 84C	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.7 ± 0.2 ± 0.6		EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
2.9 <sup>+0.6</sup> <sub>-0.4</sub> ± 0.6		<sup>13</sup> EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
3.2 ± 0.2 ± 0.6		BRANDELIK 81B	TASS	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.6 ± 0.3 ± 0.5		ROUSSARIE 81	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.3 ± 0.8		<sup>14</sup> BERGER 80B	PLUT	$e^+ e^-$

NODE=M005W8

NODE=M005W8

NODE=M005W8

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M005W8;LINKAGE=A

NODE=M005W8;LINKAGE=P1

NODE=M005W8;LINKAGE=P3

NODE=M005W8;LINKAGE=P2

NODE=M005W8;LINKAGE=SC

NODE=M005W8;LINKAGE=YA

NODE=M005W;LINKAGE=B

NODE=M005W;LINKAGE=A

NODE=M005PW;LINKAGE=C

NODE=M005PW;LINKAGE=B

NODE=M005PW;LINKAGE=X

NODE=M005PW;LINKAGE=V

NODE=M005PW;LINKAGE=H

NODE=M005PW;LINKAGE=A

<sup>1</sup> Based on a  $K$ -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.

<sup>2</sup> Solution A (preferred solution based on  $\chi^2$ -analysis).

<sup>3</sup> Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

<sup>4</sup> Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

<sup>5</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

<sup>6</sup> With a narrow scalar state around 1220 MeV.

<sup>7</sup> Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.

<sup>8</sup> Using the unitarized model of LYTH 85.

<sup>9</sup> Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with  $\gamma\gamma$  width of  $f_0(1370)$ :  $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$  KeV.

<sup>10</sup> Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes  $2.66 \pm 0.21$  in the calculation of LANDRO 86.

<sup>11</sup> Using the MENNESSIER 83 model.

<sup>12</sup> Superseded by BOYER 90.

<sup>13</sup> If helicity = 2 assumption is not made.

<sup>14</sup> Using mass, width and  $B(f_2(1270) \rightarrow 2\pi)$  from PDG 78.

 $\Gamma(e^+ e^-)$  $\Gamma_{10}$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.11</b>	90	ACHASOV	00K	SDN $e^+ e^- \rightarrow \pi^0 \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<1.7	90	VOROBYEV	88	ND $e^+ e^- \rightarrow \pi^0 \pi^0$

NODE=M005W9

NODE=M005W9

$f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M005223

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_3\Gamma_7/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M005G1  
NODE=M005G1**0.121±0.020 OUR FIT** Error includes scale factor of 1.3.**0.091±0.007±0.027** <sup>1</sup> ALBRECHT 90G ARG  $e^+e^- \rightarrow e^+e^- K^+K^-$ 

••• We do not use the following data for averages, fits, limits, etc. •••

0.104±0.007±0.072 <sup>2</sup> ALBRECHT 90G ARG  $e^+e^- \rightarrow e^+e^- K^+K^-$ 

OCCUR=2

<sup>1</sup> Using an incoherent background.

NODE=M005G1;LINKAGE=A

<sup>2</sup> Using a coherent background.

NODE=M005G1;LINKAGE=K

 $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_5\Gamma_7/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M005G02  
NODE=M005G02**11.5<sup>+1.8+4.5</sup><sub>-2.0-3.7</sub>** <sup>1</sup> UEHARA 10A BELL 10.6  $e^+e^- \rightarrow e^+e^-\eta\eta$ <sup>1</sup> Including interference with the  $f_2'(1525)$  (parameters fixed to the values from the 2008 edition of this review, PDG 08) and  $f_0(Y)$ .

NODE=M005G02;LINKAGE=UE

**Helicity-0/Helicity-2 RATIO IN  $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$** 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M005HR0  
NODE=M005HR0**3.7±0.3<sup>+15.9</sup><sub>-2.9</sub>** UEHARA 08A BELL 10.6  $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$ 

••• We do not use the following data for averages, fits, limits, etc. •••

9.5±1.8 <sup>1</sup> DAI 14A RVUE Compilation13 <sup>2,3</sup> PENNINGTON 08 RVUE Compilation26 <sup>3,4</sup> PENNINGTON 08 RVUE Compilation

OCCUR=2

OCCUR=3

<sup>1</sup> Based on a  $K$ -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.

NODE=M005HR0;LINKAGE=A

<sup>2</sup> Solution A (preferred solution based on  $\chi^2$ -analysis).

NODE=M005HR0;LINKAGE=P1

<sup>3</sup> Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

NODE=M005HR0;LINKAGE=P3

<sup>4</sup> Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

NODE=M005HR0;LINKAGE=P2

 **$f_2(1270)$  BRANCHING RATIOS**

NODE=M005225

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R10  
NODE=M005R10**0.843<sup>+0.029</sup><sub>-0.009</sub> OUR FIT** Error includes scale factor of 1.2. [0.842<sup>+0.029</sup><sub>-0.009</sub> OUR 2022 FIT

NEW

Scale factor = 1.1]

**0.837±0.020 OUR AVERAGE**0.849±0.025 CHABAUD 83 ASPK 17  $\pi^-p$  polarized0.85 ±0.05 250 BEAUPRE 71 HBC 8  $\pi^+p \rightarrow \Delta^{++}f_2$ 0.8 ±0.04 600 OH 70 HBC 1.26  $\pi^-p \rightarrow \pi^+\pi^-n$ 

••• We do not use the following data for averages, fits, limits, etc. •••

0.856±0.001±0.05 <sup>1</sup> ALBRECHT 20 RVUE 0.9  $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$ <sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

NODE=M005R10;LINKAGE=A

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi\pi)$  $\Gamma_2/\Gamma_1$ Should be twice  $\Gamma(2\pi^+2\pi^-)/\Gamma(\pi\pi)$  if decay is  $\rho\rho$ . (See ASCOLI 68D.)

NODE=M005R2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R2  
NODE=M005R2**0.091<sup>+0.014</sup><sub>-0.040</sub> OUR FIT** Error includes scale factor of 1.2. [0.091<sup>+0.014</sup><sub>-0.040</sub> OUR 2022 FIT

NEW

Scale factor = 1.2]

**0.15 ±0.06** 600 EISENBERG 74 HBC 4.9  $\pi^+p \rightarrow \Delta^{++}f_2$ 

••• We do not use the following data for averages, fits, limits, etc. •••

0.07 EMMS 75D DBC 4  $\pi^+n \rightarrow pf_2$

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M005R00  
 NODE=M005R00

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.033±0.001±0.005	<sup>1</sup> ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
-------------------	-----------------------	----	------	--

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

NODE=M005R00;LINKAGE=A

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$  $\Gamma_3/\Gamma_1$ 

We average only experiments which either take into account  $f_2(1270)$ - $a_2(1320)$  interference explicitly or demonstrate that  $a_2(1320)$  production is negligible.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R3

NODE=M005R3

NODE=M005R3

**0.054<sup>+0.005</sup><sub>-0.006</sub> OUR FIT** Error includes scale factor of 2.7.

**0.041<sup>+0.004</sup><sub>-0.005</sub> OUR AVERAGE**

0.045±0.01	<sup>1</sup> BARGIOTTI	03	OBLX	$\bar{p}p$	
0.037 <sup>+0.008</sup> <sub>-0.021</sub>	ETKIN	82B	MPS	23 $\pi^- p \rightarrow n 2K_S^0$	
0.045±0.009	CHABAUD	81	ASPK	17 $\pi^- p$ polarized	
0.039±0.008	LOVERRE	80	HBC	4 $\pi^- p \rightarrow K\bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.052±0.025	ABLIKIM	04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$	
0.036±0.005	<sup>2</sup> COSTA	80	OMEG	1-2.2 $\pi^- p \rightarrow K^+ K^- n$	
0.030±0.005	<sup>3</sup> MARTIN	79	RVUE		
0.027±0.009	<sup>4</sup> POLYCHRO...	79	STRC	7 $\pi^- p \rightarrow n 2K_S^0$	
0.025±0.015	EMMS	75D	DBC	4 $\pi^+ n \rightarrow p f_2$	
0.031±0.012	20	ADERHOLZ	69	HBC	8 $\pi^+ p \rightarrow K^+ K^- \pi^+ p$

<sup>1</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ , and  $K^\pm K_S^0 \pi^\mp$ .

<sup>2</sup> Re-evaluated by CHABAUD 83.

<sup>3</sup> Includes PAWLICKI 77 data.

<sup>4</sup> Takes into account the  $f_2(1270)$ - $f_2'(1525)$  interference.

NODE=M005R;LINKAGE=BG

NODE=M005R3;LINKAGE=D

NODE=M005R3;LINKAGE=F

NODE=M005R3;LINKAGE=M

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\pi\pi)$  $\Gamma_4/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R1

NODE=M005R1

**0.033±0.005 OUR FIT** Error includes scale factor of 1.2.

**0.033±0.004 OUR AVERAGE** Error includes scale factor of 1.1.

0.024±0.006	160	EMMS	75D	DBC	4 $\pi^+ n \rightarrow p f_2$
0.051±0.025	70	EISENBERG	74	HBC	4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$
0.043 <sup>+0.007</sup> <sub>-0.011</sub>	285	<sup>1</sup> LOUIE	74	HBC	3.9 $\pi^- p \rightarrow n f_2$
0.037±0.007	154	ANDERSON	73	DBC	6 $\pi^+ n \rightarrow p f_2$
0.047±0.013		OH	70	HBC	1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

<sup>1</sup> LOUIE 74 was quoted as 0.065 in PDG 74. Factor 2/3 to go from  $\pi^+ \pi^- \rightarrow \pi\pi$  forgotten. Mike L.

NODE=M005R1;LINKAGE=02

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

NODE=M005R7

NODE=M005R7

**4.0±0.8 OUR FIT** Error includes scale factor of 2.1.

**2.9±0.5 OUR AVERAGE**

2.7±0.7	BINON	05	GAMS	33 $\pi^- p \rightarrow \eta\eta n$
2.8±0.7	ALDE	86D	GAM4	100 $\pi^- p \rightarrow 2\eta n$
5.2±1.7	BINON	83	GAM2	38 $\pi^- p \rightarrow 2\eta n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0±1.0±2.0	<sup>1</sup> ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
-------------	-----------------------	----	------	--

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

NODE=M005R7;LINKAGE=A

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$  $\Gamma_5/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M005R6

NODE=M005R6

**0.003±0.001** BARBERIS 00E 450  $pp \rightarrow p_f \eta \eta p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.05	95	EDWARDS	82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\eta$
<0.016	95	EMMS	75D	DBC	4 $\pi^+ n \rightarrow p f_2$
<0.09	95	EISENBERG	74	HBC	4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$					$\Gamma_6/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.0030 ± 0.0010 OUR FIT</b>					
<b>0.003 ± 0.001</b>	400 ± 50	ALDE	87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_7/\Gamma$
VALUE (units $10^{-5}$ )		DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.57 ± 0.01 <sup>+1.39</sup> <sub>-0.14</sub>		UEHARA	08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$					$\Gamma_8/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.010</b>	95	EMMS	75D	DBC	4 $\pi^+ n \rightarrow p f_2$
$\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$					$\Gamma_9/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.004</b>	95	EMMS	75D	DBC	4 $\pi^+ n \rightarrow p f_2$
$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{10}/\Gamma$
VALUE (units $10^{-10}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;6</b>	90	ACHASOV	00K	SND	$e^+ e^- \rightarrow \pi^0 \pi^0$

**f<sub>2</sub>(1270) REFERENCES**

f <sub>2</sub> (1270) REFERENCES					NODE=M005
KLEMP	22	PL B830 137171	E. Klempt <i>et al.</i>	(BONN)	REFID=61646
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
CARVER	21	PRL 126 082002	M. Carver <i>et al.</i>	(CLAS Collab.)	REFID=61097
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)	REFID=55923
BOGOLYUB...	13	PAN 76 1324	M.Yu. Bogolyubsky <i>et al.</i>	(HYPERON-M Collab.)	REFID=55585
Translated from YAF 76 1389.					
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>		REFID=52303
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51652
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
Translated from YAF 68 998.					
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
Translated from YAF 66 860.					
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BOGLIONE	99	EPJ C9 11	M. Boggione, M.R. Pennington		REFID=46931
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
Translated from YAF 62 446.					
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
GRYGOREV	96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov	(ITEP)	REFID=45566
Translated from YAF 59 2187.					
YABUKI	95	JPSJ 64 435	F. Yabuki <i>et al.</i>	(VENUS Collab.)	REFID=46384
PROKOSHKIN	94	PD 39 420	Y.D. Prokoshkin, A.A. Kondashov	(SERP)	REFID=44094
Translated from DANS 336 613.					
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
BLINOV	92	ZPHY C53 33	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41858
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	REFID=41587
ADACHI	90D	PL B234 185	I. Adachi <i>et al.</i>	(TOPAZ Collab.)	REFID=41345
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
Translated from YAF 48 436.					

ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)	REFID=20764
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LANDRO	86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen	(UTRO)	REFID=20767
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
LYTH	85	JP G11 459	D.H. Lyth		REFID=42169
BEHREND	84B	ZPHY C23 223	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20757
BERGER	84	ZPHY C26 199	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20760
COURAU	84	PL 147B 227	A. Courau <i>et al.</i>	(CIT, SLAC)	REFID=20758
SMITH	84C	PR D30 851	J.R. Smith <i>et al.</i>	(SLAC, LBL, HARV)	REFID=20759
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20751
		Translated from YAF 38 934.			
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)	REFID=20393
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20745
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)	REFID=20741
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)	REFID=20388
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20736
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)	REFID=20382
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
PDG	78	PL 75B 1	C. Bricman <i>et al.</i>		REFID=40124
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)	REFID=20367
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)	REFID=20721
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=20715
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)	REFID=20719
PDG	74	PL 50B 1	V. Chaloupka <i>et al.</i>		REFID=40125
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)	REFID=20710
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)	REFID=20103
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)	REFID=20698
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)	REFID=20700
ARMENISE	70	LCN 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP	REFID=20335
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)	REFID=20681
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)	REFID=20065
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
DERADO	65	PRL 14 872	I. Derado <i>et al.</i>	(NDAM)	REFID=20668
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)	REFID=20663
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)	REFID=20657

$f_1(1285)$ 

$$I^G(J^{PC}) = 0^+(1^{++})$$

NODE=M008

 $f_1(1285)$  MASS

NODE=M008M

NODE=M008M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1281.9 ± 0.5</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.8. See the ideogram below.		
1281.0 ± 0.8		DICKSON	16 CLAS	2.55 $\gamma p \rightarrow \eta \pi^+ \pi^- p$
1287.4 ± 3.0	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
1281.16 ± 0.39 ± 0.45		<sup>1</sup> LEES	12X BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$
1285.1 ± 1.0 $\begin{smallmatrix} + 1.6 \\ - 0.3 \end{smallmatrix}$		<sup>2</sup> ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
1281 ± 2 ± 1		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
1276.1 ± 8.1 ± 8.0	203	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
1274 ± 6	237	ABDALLAH	03H DLPH	91.2 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
1280 ± 4		ACCIARRI	01G L3	
1288 ± 4 ± 5	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1284 ± 6	1400	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1281 ± 1		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1281 ± 1		BARBERIS	97C OMEG	450 $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
1280 ± 2		<sup>3</sup> ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1282.2 ± 1.5		LEE	94 MPS2	18 $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
1279 ± 5		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1278 ± 2	140	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi pp$
1278 ± 2		ARMSTRONG	89G OMEG	85 $\pi^+ p \rightarrow 4\pi \pi p, pp \rightarrow 4\pi pp$
1280.1 ± 2.1	60	RATH	89 MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1285 ± 1	4750	<sup>4</sup> BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1280 ± 1	504	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1280 ± 4		ANDO	86 SPEC	8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1277 ± 2	420	REEVES	86 SPEC	6.6 $p\bar{p} \rightarrow K K \pi X$
1285 ± 2		CHUNG	85 SPEC	8 $\pi^- p \rightarrow N K \bar{K} \pi$
1279 ± 2	604	ARMSTRONG	84 OMEG	85 $\pi^+ p \rightarrow K \bar{K} \pi \pi p, pp \rightarrow K \bar{K} \pi pp$
1286 ± 1		CHAUVAT	84 SPEC	ISR 31.5 $pp$
1278 ± 4		EVANGELIS...	81 OMEG	12 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
1283 ± 3	103	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
1282 ± 2	320	NACASCH	78 HBC	0.7,0.76 $\bar{p}p \rightarrow K \bar{K} 3\pi$
1279 ± 5	210	GRASSLER	77 HBC	16 $\pi^\mp p$
1286 ± 3	180	DUBOC	72 HBC	1.2 $\bar{p}p \rightarrow 2K4\pi$
1283 ± 5		DAHL	67 HBC	1.6-4.2 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1289.3 ± 2.8	234	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
1284.2 ± 2.2		<sup>5</sup> AAIJ	14Y LHCB	$\bar{B}_{(s)}^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$
1281.9 ± 0.5		<sup>5</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$
1282.8 ± 0.6		<sup>5</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
1270 ± 10		AMELIN	95 VES	37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
1280 ± 2		ABATZIS	94 OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1282 ± 4		ARMSTRONG	93C E760	$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1270 ± 6 ± 10		ARMSTRONG	92C OMEG	300 $pp \rightarrow pp\pi^+ \pi^- \gamma$
1281 ± 1		ARMSTRONG	89E OMEG	300 $pp \rightarrow pp2(\pi^+ \pi^-)$
1279 ± 6 ± 10	16	BECKER	87 MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$
1286 ± 9		GIDAL	87 MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
1287 ± 5	353	BITYUKOV	84B SPEC	32 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
~ 1279		<sup>6</sup> TORNQVIST	82B RVUE	

OCCUR=2

1275 ± 6	31	BROMBERG	80	SPEC	100 $\pi^- p \rightarrow K \bar{K} \pi X$
1288 ± 9	200	GURTU	79	HBC	4.2 $K^- p \rightarrow n \eta 2\pi$
~ 1275.0	46	<sup>7</sup> STANTON	79	CNTR	8.5 $\pi^- p \rightarrow n 2\gamma 2\pi$
1271 ± 10	34	CORDEN	78	OMEG	12-15 $\pi^- p \rightarrow K^+ K^- \pi n$
1295 ± 12	85	CORDEN	78	OMEG	12-15 $\pi^- p \rightarrow n 5\pi$
1292 ± 10	150	DEFOIX	72	HBC	0.7 $\bar{p} p \rightarrow 7\pi$
1280 ± 3	500	<sup>8</sup> THUN	72	MMS	13.4 $\pi^- p$
1303 ± 8		BARDADIN...	71	HBC	8 $\pi^+ p \rightarrow p 6\pi$
1283 ± 6		BOESEBECK	71	HBC	16.0 $\pi p \rightarrow p 5\pi$
1270 ± 10		CAMPBELL	69	DBC	2.7 $\pi^+ d$
1285 ± 7		LORSTAD	69	HBC	0.7 $\bar{p} p$ , 4,5-body
1290 ± 7		D'ANDLAU	68	HBC	1.2 $\bar{p} p$ , 5-6 body

OCCUR=2

<sup>1</sup> Using the  $2\pi^+ 2\pi^-$  and  $\pi^+ \pi^- \eta$  modes of  $f_1(1285)$  decay.

<sup>2</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980) \pi$ .

<sup>3</sup> Supersedes ABATZIS 94, ARMSTRONG 89E.

<sup>4</sup> From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  system.


<sup>5</sup> No systematic error given.

<sup>6</sup> From a unitarized quark-model calculation.

<sup>7</sup> From phase shift analysis of  $\eta \pi^+ \pi^-$  system.

<sup>8</sup> Seen in the missing mass spectrum.

NODE=M008M;LINKAGE=LE  
 NODE=M008M;LINKAGE=BL  
 NODE=M008M;LINKAGE=B  
 NODE=M008M;LINKAGE=A  
 NODE=M008M;LINKAGE=N1  
 NODE=M008M;LINKAGE=T  
 NODE=M008M;LINKAGE=P  
 NODE=M008M;LINKAGE=S



**$f_1(1285)$  WIDTH**

NODE=M008W

Only experiments giving width error less than 20 MeV are kept for averaging.

NODE=M008W



VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>22.7± 1.1 OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.		
18.4± 1.4		DICKSON	16 CLAS	2.55 $\gamma p \rightarrow \eta \pi^+ \pi^- p$
18.3± 6.3	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
22.0± 3.1 <sup>+</sup> <sub>-1.5</sub>		<sup>1</sup> ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
35 ± 6 ± 4		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
40.0± 8.6± 9.3	203	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
29 ± 12	237	ABDALLAH	03H DLPH	91.2 $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
45 ± 9 ± 7	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
55 ± 18	1400	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
24 ± 3		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
20 ± 2		BARBERIS	97C OMEG	450 $pp \rightarrow ppK_S^0 K^\pm \pi^\mp$
36 ± 5		<sup>2</sup> ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+ \pi^-)$
29.0± 4.1		LEE	94 MPS2	18 $\pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
25 ± 4	140	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi pp$
22 ± 2	4750	<sup>3</sup> BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
25 ± 4	504	BITYUKOV	88 SPEC	32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$
19 ± 5		ANDO	86 SPEC	8 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
32 ± 8	420	REEVES	86 SPEC	6.6 $p \bar{p} \rightarrow K K \pi X$
22 ± 2		CHUNG	85 SPEC	8 $\pi^- p \rightarrow N K \bar{K} \pi$
32 ± 3	604	ARMSTRONG	84 OMEG	85 $\pi^+ p \rightarrow K \bar{K} \pi \pi p$ , $pp \rightarrow K \bar{K} \pi pp$
24 ± 3		CHAUVAT	84 SPEC	ISR 31.5 $pp$
29 ± 10	103	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
28.3± 6.7	320	NACASCH	78 HBC	0.7,0.76 $p \bar{p} \rightarrow K \bar{K} 3\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
17.1± 3.4	234	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
32.4± 5.8		<sup>4</sup> AAIJ	14Y LHCb	$\bar{B}_s^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$
18.2± 1.2		<sup>4</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$
19.4± 1.5		<sup>4</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$
40 ± 5		ABATZIS	94 OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
31 ± 5		ARMSTRONG	89E OMEG	300 $pp \rightarrow pp2(\pi^+ \pi^-)$
41 ± 12		ARMSTRONG	89G OMEG	85 $\pi^+ p \rightarrow 4\pi \pi p$ , $pp \rightarrow 4\pi pp$
17.9± 10.9	60	RATH	89 MPS	21.4 $\pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
14 <sup>+20</sup> <sub>-14</sub> ± 10	16	BECKER	87 MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$
26 ± 12		EVANGELIS...	81 OMEG	12 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
25 ± 15	200	GURTU	79 HBC	4.2 $K^- p \rightarrow n \eta 2\pi$
~ 10		<sup>5</sup> STANTON	79 CNTR	8.5 $\pi^- p \rightarrow n 2\gamma 2\pi$
24 ± 18	210	GRASSLER	77 HBC	16 $\pi^\mp p$
28 ± 5	150	<sup>6</sup> DEFOIX	72 HBC	0.7 $p \bar{p} \rightarrow 7\pi$
46 ± 9	180	<sup>6</sup> DUBOC	72 HBC	1.2 $p \bar{p} \rightarrow 2K 4\pi$
37 ± 5	500	<sup>7</sup> THUN	72 MMS	13.4 $\pi^- p$
10 ± 10		BOESEBECK	71 HBC	16.0 $\pi p \rightarrow p 5\pi$
30 ± 15		CAMPBELL	69 DBC	2.7 $\pi^+ d$
60 ± 15		<sup>6</sup> LORSTAD	69 HBC	0.7 $p \bar{p}$ , 4,5-body
35 ± 10		<sup>6</sup> DAHL	67 HBC	1.6-4.2 $\pi^- p$

<sup>1</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980) \pi$ .

<sup>2</sup> Supersedes ABATZIS 94, ARMSTRONG 89E.

<sup>3</sup> From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  system.

<sup>4</sup> No systematic error given.

<sup>5</sup> From phase shift analysis of  $\eta \pi^+ \pi^-$  system.

<sup>6</sup> Resolution is not unfolded.


<sup>7</sup> Seen in the missing mass spectrum.

NODE=M008W

OCCUR=2

NODE=M008W;LINKAGE=BL  
 NODE=M008W;LINKAGE=B  
 NODE=M008W;LINKAGE=A  
 NODE=M008W;LINKAGE=N1  
 NODE=M008W;LINKAGE=P  
 NODE=M008W;LINKAGE=R  
 NODE=M008W;LINKAGE=S

WEIGHTED AVERAGE  
22.7±1.1 (Error scaled by 1.5)



**f<sub>1</sub>(1285) DECAY MODES**

NODE=M008215;NODE=M008

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level	
$\Gamma_1$ $4\pi$	(32.7 ± 1.9) %	S=1.2	DESIG=21
$\Gamma_2$ $\pi^0\pi^0\pi^+\pi^-$	(21.8 ± 1.3) %	S=1.2	DESIG=22
$\Gamma_3$ $2\pi^+2\pi^-$	(10.9 ± 0.6) %	S=1.2	DESIG=20
$\Gamma_4$ $\rho^0\pi^+\pi^-$	(10.9 ± 0.6) %	S=1.2	DESIG=191
$\Gamma_5$ $\rho^0\rho^0$	seen		DESIG=23
$\Gamma_6$ $4\pi^0$	< 7 × 10 <sup>-4</sup>	CL=90%	DESIG=7
$\Gamma_7$ $\eta\pi^+\pi^-$	(35 ± 15) %		DESIG=198
$\Gamma_8$ $\eta\pi\pi$	(52.2 ± 2.0) %	S=1.2	DESIG=3
$\Gamma_9$ $a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$ ]	(38 ± 4) %		DESIG=4
$\Gamma_{10}$ $\eta\pi\pi$ [excluding $a_0(980)\pi$ ]	(14 ± 4) %		DESIG=5
$\Gamma_{11}$ $K\bar{K}\pi$	(9.0 ± 0.4) %	S=1.1	DESIG=1
$\Gamma_{12}$ $K\bar{K}^*(892)$	not seen		DESIG=6
$\Gamma_{13}$ $\pi^+\pi^-\pi^0$	(3.0 ± 0.9) × 10 <sup>-3</sup>		DESIG=197
$\Gamma_{14}$ $\rho^\pm\pi^\mp$	< 3.1 × 10 <sup>-3</sup>	CL=95%	DESIG=199
$\Gamma_{15}$ $\gamma\rho^0$	(6.1 ± 1.0) %	S=1.7	DESIG=13
$\Gamma_{16}$ $\phi\gamma$	(7.4 ± 2.6) × 10 <sup>-4</sup>		DESIG=10
$\Gamma_{17}$ $e^+e^-$	< 9.4 × 10 <sup>-9</sup>	CL=90%	DESIG=200
$\Gamma_{18}$ $\gamma\gamma^*$			DESIG=9
$\Gamma_{19}$ $\gamma\gamma$			DESIG=8

## CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 18 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 24.0$  for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_9$	-30			
$x_{10}$	-12	-88		
$x_{11}$	22	-10	-4	
$x_{15}$	-25	-7	-3	-27
	$x_1$	$x_9$	$x_{10}$	$x_{11}$

### $f_1(1285) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

NODE=M008217

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{19} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{19} / \Gamma$			
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.62</b>	95	GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

NODE=M008G2  
NODE=M008G2

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{18} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{18} / \Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.4 ± 0.4 OUR AVERAGE</b>		Error includes scale factor of 1.4.			
1.18 ± 0.25 ± 0.20	26	<sup>1,2</sup> AIHARA	88B	TPC	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
2.30 ± 0.61 ± 0.42		<sup>1,3</sup> GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

NODE=M008G3  
NODE=M008G3

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.3 ± 0.3	420	<sup>4</sup> ACHARD	02B	L3	183-209 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
-----------------	-----	---------------------	-----	----	--

<sup>1</sup> Assuming a  $\rho$ -pole form factor.

<sup>2</sup> Published value multiplied by  $\eta\pi\pi$  branching ratio 0.49.

<sup>3</sup> Published value divided by 2 and multiplied by the  $\eta\pi\pi$  branching ratio 0.49.

<sup>4</sup> Published value multiplied by the  $\eta\pi\pi$  branching ratio 0.52.

NODE=M008G3;LINKAGE=A  
NODE=M008G3;LINKAGE=F  
NODE=M008G3;LINKAGE=B  
NODE=M008G3;LINKAGE=AC

### $f_1(1285) \text{ BRANCHING RATIOS}$

NODE=M008220

$\Gamma(K\bar{K}\pi) / \Gamma(4\pi)$		$\Gamma_{11} / \Gamma_1$			
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>0.274 ± 0.017 OUR FIT</b>		Error includes scale factor of 1.4.			
<b>0.271 ± 0.016 OUR AVERAGE</b>		Error includes scale factor of 1.2.			
0.265 ± 0.014		<sup>1</sup> BARBERIS	97C	OMEG 450	$pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
0.28 ± 0.05		<sup>2</sup> ARMSTRONG	89E	OMEG 300	$pp \rightarrow pp f_1(1285)$
0.37 ± 0.03 ± 0.05		<sup>3</sup> ARMSTRONG	89G	OMEG 85	$\pi p \rightarrow 4\pi X$

NODE=M008R1  
NODE=M008R1

<sup>1</sup> Using  $2(\pi^+ \pi^-)$  data from BARBERIS 97B.

<sup>2</sup> Assuming  $\rho\pi\pi$  and  $a_0(980)\pi$  intermediate states.

<sup>3</sup>  $4\pi$  consistent with being entirely  $\rho\pi\pi$ .

NODE=M008R1;LINKAGE=B  
NODE=M008R1;LINKAGE=M  
NODE=M008R1;LINKAGE=A

$\Gamma(\pi^0 \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$		$\Gamma_2 / \Gamma = \frac{2}{3} \Gamma_1 / \Gamma$			
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>0.218 ± 0.013 OUR FIT</b>		Error includes scale factor of 1.2.			

NODE=M008R18  
NODE=M008R18

$\Gamma(2\pi^+ 2\pi^-) / \Gamma_{\text{total}}$		$\Gamma_3 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$			
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>0.109 ± 0.006 OUR FIT</b>		Error includes scale factor of 1.2.			

NODE=M008R17  
NODE=M008R17

$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$		$\Gamma_4 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$			
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>0.109 ± 0.006 OUR FIT</b>		Error includes scale factor of 1.2.			

NODE=M008R19  
NODE=M008R19

$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma(2\pi^+ 2\pi^-)$		$\Gamma_4 / \Gamma_3$			
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>1.0 ± 0.4</b>		GRASSLER	77	HBC	16 GeV $\pi^\pm p$

NODE=M008R6  
NODE=M008R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ VALUEDOCUMENT IDCOMMENTNODE=M008R21  
NODE=M008R21

seen

BARBERIS 00C 450  $p p \rightarrow p_f 4\pi p_s$  $\Gamma(4\pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ VALUE (units  $10^{-4}$ ) CL%DOCUMENT IDTECNCOMMENTNODE=M008R8  
NODE=M008R8

&lt;7

90

ALDE

87

GAM4

100  $\pi^- p \rightarrow 4\pi^0 n$  $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma(\eta \pi^+ \pi^-)$  $\Gamma_{13}/\Gamma_7$ VALUE (%) EVTsDOCUMENT IDTECNCOMMENTNODE=M008R02  
NODE=M008R02**0.86 ± 0.16 ± 0.20**

2.3k

<sup>1</sup> DOROFEEV 11 VES  $\pi^- N \rightarrow \pi^- f_1(1285) N$ <sup>1</sup> Value obtained selecting the region corresponding to  $f_0(980)$  in the  $\pi^+ \pi^-$  mass spectrum.

NODE=M008R02;LINKAGE=DO

 $\Gamma(\eta \pi \pi)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$ VALUEDOCUMENT IDNODE=M008R22  
NODE=M008R22**0.522 ± 0.020 OUR FIT** Error includes scale factor of 1.2. $\Gamma(4\pi)/\Gamma(\eta \pi \pi)$  $\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9 + \Gamma_{10})$ VALUEDOCUMENT IDTECNCOMMENTNODE=M008R4  
NODE=M008R4**0.63 ± 0.06 OUR FIT** Error includes scale factor of 1.3.**0.41 ± 0.14 OUR AVERAGE**

0.37 ± 0.11 ± 0.11

BOLTON

92

MRK3

 $J/\psi \rightarrow \gamma f_1(1285)$ 

0.64 ± 0.40

GURTU

79

HBC

4.2  $K^- p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.93 ± 0.30

<sup>1</sup> GRASSLER 77 HBC 16  $\pi^\mp p$ <sup>1</sup> Assuming  $\rho \pi \pi$  and  $a_0(980)\pi$  intermediate states.

NODE=M008R4;LINKAGE=M

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma(\eta \pi \pi)$  $\Gamma_3/\Gamma_8$ VALUEDOCUMENT IDTECNCOMMENTNODE=M008R04  
NODE=M008R04**0.28 ± 0.02 ± 0.02**<sup>1</sup> LEES 12X BABR  $\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$ <sup>1</sup> Assuming  $B(f_1(1285) \rightarrow \pi \pi \eta) = 3/2 B(f_1(1285) \rightarrow \pi^+ \pi^- \eta)$ .

NODE=M008R04;LINKAGE=LE

 $\Gamma(a_0(980)\pi [\text{ignoring } a_0(980) \rightarrow K\bar{K}])/\Gamma(\eta \pi \pi)$  $\Gamma_9/\Gamma_8 = \Gamma_9/(\Gamma_9 + \Gamma_{10})$ VALUE CL%DOCUMENT IDTECNCOMMENTNODE=M008R3  
NODE=M008R3**0.72 ± 0.08 OUR FIT****0.72 ± 0.07 OUR AVERAGE**

0.74 ± 0.02 ± 0.09

DICKSON

16

CLAS

 $\gamma p \rightarrow f_1(1285) p$ 

0.72 ± 0.15

GURTU

79

HBC

4.2  $K^- p$ 0.6 <sup>+0.3</sup><sub>-0.2</sub>

CORDEN

78

OMEG

12-15  $\pi^- p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

&gt;0.69

95

ACHARD

02B

L3

183-209  $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$ 

0.28 ± 0.07

ALDE

97B

GAM4

100  $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$ 

1.0 ± 0.3

GRASSLER

77

HBC

16  $\pi^\mp p$  $\Gamma(K\bar{K}\pi)/\Gamma(\eta \pi \pi)$  $\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9 + \Gamma_{10})$ VALUEDOCUMENT IDTECNCOMMENTNODE=M008R2  
NODE=M008R2**0.172 ± 0.012 OUR FIT** Error includes scale factor of 1.1.**0.176 ± 0.012 OUR AVERAGE**

0.216 ± 0.010 ± 0.031

DICKSON

16

CLAS

 $\gamma p \rightarrow f_1(1285) p$ 

OCCUR=2

0.166 ± 0.01 ± 0.008

BARBERIS

98C

OMEG

450  $p p \rightarrow p_f f_1(1285) p_s$ 

0.42 ± 0.15

GURTU

79

HBC

4.2  $K^- p$ 

0.5 ± 0.2

<sup>1</sup> CORDEN

78

OMEG

12-15  $\pi^- p$ 

0.20 ± 0.08

<sup>2</sup> DEFOIX

72

HBC

0.7  $\bar{p} p \rightarrow 7\pi$ 

0.16 ± 0.08

CAMPBELL

69

DBC

2.7  $\pi^+ d$ <sup>1</sup> CORDEN 78 assumes low-mass  $\eta \pi \pi$  region is dominantly  $1^{++}$ . See BARBERIS 98C and MANAK 00A for discussion.

NODE=M008R2;LINKAGE=CD

<sup>2</sup>  $K\bar{K}$  system characterized by the  $l = 1$  threshold enhancement. (See under  $a_0(980)$ ).

NODE=M008R2;LINKAGE=K

 $\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ VALUEDOCUMENT IDTECNCOMMENTNODE=M008R5  
NODE=M008R5

not seen

NACASCH 78 HBC 0.7, 0.76  $\bar{p} p \rightarrow K\bar{K} 3\pi$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

<sup>1</sup> ACHARD 07 L3 183-209  $e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$ <sup>1</sup> A clear signal of  $19.8 \pm 4.4$  events observed at high  $Q^2$ .

NODE=M008R5;LINKAGE=CH

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$   $\Gamma_{13}/\Gamma$

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.30±0.055±0.074</b>	2.3k	<sup>1</sup> DOROFEEV 11	VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$

NODE=M008R01  
NODE=M008R01

<sup>1</sup> Value obtained selecting the region corresponding to  $f_0(980)$  in the  $\pi^+\pi^-$  mass spectrum. The systematic error includes the uncertainty on the partial width  $f_1 \rightarrow \eta\pi\pi$  obtained from PDG 10 data.

NODE=M008R01;LINKAGE=DO

$\Gamma(\rho^\pm\pi^\mp)/\Gamma_{total}$   $\Gamma_{14}/\Gamma$

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.31</b>	95	DOROFEEV 11	VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$

NODE=M008R03  
NODE=M008R03

$\Gamma(\gamma\rho^0)/\Gamma_{total}$   $\Gamma_{15}/\Gamma$

VALUE (units 10 <sup>-2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<b>6.1±1.0 OUR FIT</b>				Error includes scale factor of 1.7.

NODE=M008R15  
NODE=M008R15

••• We do not use the following data for averages, fits, limits, etc. •••

2.8±0.7±0.6		<sup>1</sup> AMELIN 95	VES	37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
<5	95	BITYUKOV 91B	SPEC	32 $\pi^- \rho \rightarrow \pi^+ \pi^- \gamma n$

<sup>1</sup> Not an independent measurement.

NODE=M008R15;LINKAGE=A

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$   $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.55±0.10 OUR FIT</b>			Error includes scale factor of 1.5.
<b>0.45±0.18</b>	<sup>1</sup> COFFMAN 90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

NODE=M008R13  
NODE=M008R13

<sup>1</sup> Using  $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0)=0.25 \times 10^{-4}$  and  $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-)=0.55 \times 10^{-4}$  given by MIR 88.

NODE=M008R13;LINKAGE=E

$\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$   $\Gamma_8/\Gamma_{15} = (\Gamma_9+\Gamma_{10})/\Gamma_{15}$


VALUE	DOCUMENT ID	TECN	COMMENT
<b>8.6±1.6 OUR FIT</b>			Error includes scale factor of 1.9.
<b>8.5±2.0 OUR AVERAGE</b>			Error includes scale factor of 2.2. See the ideogram below.

NODE=M008R16  
NODE=M008R16

21.3±4.4	DICKSON 16	CLAS	$\gamma\rho \rightarrow f_1(1285)\rho$
10.0±1.0±2.0	BARBERIS 98C	OMEG 450	$\rho\rho \rightarrow \rho_f f_1(1285)\rho_S$
7.5±1.0	<sup>1</sup> ARMSTRONG 92C	OMEG 300	$\rho\rho \rightarrow \rho\rho\pi^+\pi^-\gamma, \rho\rho\eta\pi^+\pi^-$

<sup>1</sup> Published value multiplied by 1.5.

NODE=M008R16;LINKAGE=B



$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$   $\Gamma_{15}/\Gamma_{11}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt;0.035</b>	90	<sup>1</sup> COFFMAN 90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

NODE=M008R12  
NODE=M008R12

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Using  $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0)=0.25 \times 10^{-4}$  and  $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi)=<0.72 \times 10^{-3}$ .

NODE=M008R12;LINKAGE=F

$\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$  $\Gamma_{16}/\Gamma_{11}$ 

VALUE (units $10^{-2}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.82 \pm 0.21 \pm 0.20</math></b>		19	BITYUKOV	88	SPEC $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.50	95	BARBERIS	98C	OMEG	450 $pp \rightarrow p_f f_1(1285) p_s$
<0.93	95	AMELIN	95	VES	37 $\pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$

 $\Gamma(e^+e^-)/\Gamma_{total}$  $\Gamma_{17}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;9.4 \times 10^{-9}</math></b>	90	<sup>1</sup> ACHASOV	20	SND $e^+e^- \rightarrow \eta\pi^0\pi^0$

<sup>1</sup>ACHASOV 20 reports two candidate events corresponding to a significance of 2.5  $\sigma$  and the branching fraction of  $(5.1^{+3.7}_{-2.7}) \times 10^{-9}$ .

 $f_1(1285)$  REFERENCES

ACHASOV	20	PL B800 135074	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60144
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60024
DICKSON	16	PR C93 065202	R. Dickson <i>et al.</i>	(JLab CLAS Collab.)	REFID=57487
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
AAIJ	14Y	PRL 112 091802	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55837
LEES	12X	PR D86 092010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54714
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53931
DOROFEEV	11	EPJ A47 68	V. Dorofeev <i>et al.</i>	(SERP, MIPT)	REFID=16755
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)	REFID=53229
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=51698
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49548
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48574
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
SOSA	99	PRL 83 913	M. Sosa <i>et al.</i>		REFID=46937
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
AMELIN	95	ZPHY C66 71	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=44376
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44437
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=44090
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)	REFID=44092
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
ARMSTRONG	92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=42097
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BITYUKOV	91B	SJNP 54 318	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=41864
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
ARMSTRONG	89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+)	REFID=40729
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
ARMSTRONG	89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)	REFID=40930
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)	REFID=40924
AIHARA	88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)	REFID=40572
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
BITYUKOV	88	PL B203 327	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=40569
MIR	88	Photon-Photon 88, 126	R. Mir	(Mark III Collab.)	REFID=41574
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP	REFID=20891
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP	REFID=20936
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP	REFID=20934
ARMSTRONG	84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=20929
BITYUKOV	84B	PL 144B 133	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=20468
CHAUVAT	84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)	REFID=20932
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)	REFID=20573
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)	REFID=20922
DIONISI	80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+)	REFID=20924
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP	REFID=20887
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20452
NACASCH	78	NP B135 203	R. Nacasch <i>et al.</i>	(PARIS, MADR, CERN)	REFID=20919
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
DUBOC	72	NP B46 429	J. Duboc <i>et al.</i>	(PARIS, LIVP)	REFID=20339
THUN	72	PRL 28 1733	R. Thun <i>et al.</i>	(STON, NEAS)	REFID=20911
BARDADIN...	71	PR D4 2711	M. Bardadin-Otwinowska <i>et al.</i>	(WARS)	REFID=20196
BOESEBECK	71	PL 34B 659	K. Boesebeck	(AACH, BERL, BONN, CERN, CRAC+)	REFID=20905
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)	REFID=20419
LORSTAD	69	NP B14 63	B. Lorstad <i>et al.</i>	(CDEF, CERN) JP	REFID=20901
D'ANDLAU	68	NP B5 693	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+) IJP	REFID=20897
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP	REFID=20321

NODE=M008R9  
NODE=M008R9

NODE=M008R00  
NODE=M008R00

OCCUR=2

NODE=M008R00;LINKAGE=B

NODE=M008

**$\eta(1295)$** 

$$I^G(J^{PC}) = 0^+(0^-+)$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M037


NODE=M037

NODE=M037M

NODE=M037M

 **$\eta(1295)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1294±4 OUR AVERAGE</b>		Error includes scale factor of 1.6. See the ideogram below.		
1302±9±8	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1282±5	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1299±4	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1295±4		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1264±8		<sup>1</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~ 1275		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$



<sup>1</sup> PWA analysis of AUGUSTIN 92 assigns  $0^-+$  quantum numbers to this state rather than  $1^{++}$  as before.

NODE=M037M;LINKAGE=AG

 **$\eta(1295)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>55± 5 OUR AVERAGE</b>				
57±23±21	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
66±13	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
53± 6		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<40	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
44±20		<sup>1</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~ 70		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$

<sup>1</sup> PWA analysis of AUGUSTIN 92 assigns  $0^-+$  quantum numbers to this state rather than  $1^{++}$  as before.

NODE=M037W

NODE=M037W

NODE=M037W;LINKAGE=AG

$\eta(1295)$  DECAY MODES

NODE=M037215;NODE=M037

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta\pi^+\pi^-$	seen
$\Gamma_2$ $a_0(980)\pi$	seen
$\Gamma_3$ $\gamma\gamma$	
$\Gamma_4$ $\eta\pi^0\pi^0$	seen
$\Gamma_5$ $\eta(\pi\pi)S\text{-wave}$	seen
$\Gamma_6$ $\sigma\eta$	seen
$\Gamma_7$ $K\bar{K}\pi$	seen

DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=3  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←  
DESIG=7;OUR EST;→ UNCHECKED ←

 $\eta(1295)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M037220

 $\Gamma(\eta\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_1\Gamma_3/\Gamma$ NODE=M037G2  
NODE=M037G2

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.066</b>	95	ACCIARRI	01G L3	183-202 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ $e^+e^-\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6	90	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
<0.3		ANTREASYAN	87 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi\pi$

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_7\Gamma_3/\Gamma$ NODE=M037G3  
NODE=M037G3

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.014	90	1,2 AHOHE	05 CLE2	10.6 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$ $e^+e^-K_S^0K^\pm\pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Using  $\eta(1295)$  mass and width 1294 MeV and 55 MeV, respectively.<sup>2</sup> Assuming three-body phase-space decay to  $K_S^0K^\pm\pi^\mp$ .NODE=M037G3;LINKAGE=AH  
NODE=M037G3;LINKAGE=B3 $\eta(1295)$  BRANCHING RATIOS

NODE=M037225

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ NODE=M037R1  
NODE=M037R1

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
seen	BIRMAN	88 MPS	8 $\pi^-p \rightarrow K^+\bar{K}^0\pi^-n$
large	ANDO	86 SPEC	8 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
large	STANTON	79 CNTR	8.4 $\pi^-p \rightarrow n\eta 2\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi^0\pi^0)$   $\Gamma_2/\Gamma_4$ NODE=M037R2  
NODE=M037R2

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.65±0.10</b>	<sup>1</sup> ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

<sup>1</sup> Assuming that  $a_0(980)$  decays only to  $\eta\pi$ .

NODE=M037R2;LINKAGE=A

 $\Gamma(\eta(\pi\pi)S\text{-wave})/\Gamma(\eta\pi^0\pi^0)$   $\Gamma_5/\Gamma_4$ NODE=M037R4  
NODE=M037R4

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.35±0.10</b>	ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

 $\Gamma(a_0(980)\pi)/\Gamma(\sigma\eta)$   $\Gamma_2/\Gamma_6$ NODE=M037R5  
NODE=M037R5

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.48±0.22</b>	9082	MANAK	00A MPS	18 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

 $\eta(1295)$  REFERENCES

NODE=M037

AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)	REFID=50764
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
		Translated from YAF 60 458.			
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)	REFID=40564
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP	REFID=20891
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP	REFID=20887



NODE=M058

 $\pi(1300)$ 

$$I^G(J^{PC}) = 1^-(0^-+)$$

 $\pi(1300)$  MASS

NODE=M058M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**1300±100 OUR ESTIMATE**

NODE=M058M

→ UNCHECKED ←

- • • We do not use the following data for averages, fits, limits, etc. • • •

1128± 26±70		DARGENT	17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1345± 8±10	18k	<sup>1</sup> SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
1200± 40	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1343± 15±24		CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1375± 40		ABELE	01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1275± 15		BERTIN	97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
~ 1114		ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
1190± 30		ZIELINSKI	84	SPEC	$200 \pi^+ Z \rightarrow Z 3\pi$
1240± 30		BELLINI	82	SPEC	$40 \pi^- A \rightarrow A 3\pi$
1273± 50		<sup>2</sup> AARON	81	RVUE	
1342± 20		BONESINI	81	OMEG	$12 \pi^- p \rightarrow p 3\pi$
~ 1400		DAUM	81B	SPEC	$63,94 \pi^- p$

<sup>1</sup> From analysis of L3 data at 183–209 GeV.<sup>2</sup> Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.NODE=M058M;LINKAGE=SC  
NODE=M058M;LINKAGE=E $\pi(1300)$  WIDTH

NODE=M058W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**200 to 600 OUR ESTIMATE**

NODE=M058W

→ UNCHECKED ←

- • • We do not use the following data for averages, fits, limits, etc. • • •

314± 39±66		DARGENT	17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
260± 20±30	18k	<sup>3</sup> SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
470±120	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
449± 39±47		CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
268± 50		ABELE	01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
218±100		BERTIN	97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
~ 340		ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
440± 80		ZIELINSKI	84	SPEC	$200 \pi^+ Z \rightarrow Z 3\pi$
360±120		BELLINI	82	SPEC	$40 \pi^- A \rightarrow A 3\pi$
580±100		<sup>4</sup> AARON	81	RVUE	
220± 70		BONESINI	81	OMEG	$12 \pi^- p \rightarrow p 3\pi$
~ 600		DAUM	81B	SPEC	$63,94 \pi^- p$

<sup>3</sup> From analysis of L3 data at 183–209 GeV.<sup>4</sup> Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.NODE=M058W;LINKAGE=SC  
NODE=M058W;LINKAGE=E $\pi(1300)$  DECAY MODES

NODE=M058215;NODE=M058

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\pi$	seen
$\Gamma_2$ $\pi(\pi\pi)S$ -wave	seen
$\Gamma_3$ $\gamma\gamma$	

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

 $\pi(1300)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M058218

 $\Gamma(\rho\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_3/\Gamma$ 

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

<0.085	90	ACCIARRI	97T	L3 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
--------	----	----------	-----	---

- • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	95	<sup>5</sup> SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
<0.54	90	ALBRECHT	97B	ARG	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$

<sup>5</sup> From analysis of L3 data at 183–209 GeV.

NODE=M058G1

NODE=M058G1

NODE=M058G1;LINKAGE=SC

$\pi(1300)$  BRANCHING RATIOS $\Gamma(\pi\pi)_{S\text{-wave}}/\Gamma(\rho\pi)$  $\Gamma_2/\Gamma_1$ 

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.4		90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
seen			CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+ 2\pi^- p$
<0.15	90		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
2.12			<sup>6</sup> AARON	81	RVUE

<sup>6</sup> Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058220

NODE=M058R1  
NODE=M058R1

NODE=M058R1;LINKAGE=E

 $\pi(1300)$  REFERENCES

DARGENT	17	JHEP 1705 143	P. dArgent <i>et al.</i>	(HEID, BRIS)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACCIARRI	97T	PL B413 147	M. Acciari <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERTIN	97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ZIELINSKI	84	PR D30 1855	M. Zielinski <i>et al.</i>	(ROCH, MINN, FNAL)
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)
AARON	81	PR D24 1207	R.A. Aaron, R.S. Longacre	(NEAS, BNL)
BONESINI	81	PL 103B 75	M. Bonesini <i>et al.</i>	(MILA, LIVP, DARE+)
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
DAUM	80	PL 89B 281	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

NODE=M058

REFID=58121  
REFID=51186  
REFID=53226  
REFID=48837  
REFID=48334  
REFID=45761  
REFID=45418  
REFID=45763  
REFID=45011  
REFID=20881  
REFID=21134  
REFID=20870  
REFID=21130  
REFID=20572  
REFID=20872  
REFID=20868  
REFID=20571

NODE=M012

 $a_2(1320)$ 

$$I^G(J^{PC}) = 1^-(2^{++})$$

 $a_2(1320)$  T-MATRIX POLE  $\sqrt{s}$ Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1305–1321)–i(52–58) OUR ESTIMATE</b>			
$(1318.7 \pm 1.9 \pm 1.3) - i(53.8 \pm 2.3^{+1.7}_{-0.9})$	<sup>1</sup> KOPF	21	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
$(1312.5 \pm 0.7 \pm 2.6) - i(53.5 \pm 0.6 \pm 1.9)$	<sup>2</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1306.0 \pm 0.8 \pm 1.3) - i(57.2 \pm 0.8 \pm 0.0)$	<sup>3</sup> RODAS	19	RVUE $91 \pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
$(1309 \pm 4) - i(55 \pm 2)$	<sup>4</sup> ANISOVICH	09	RVUE $\bar{p}p, \pi N$

<sup>1</sup> Extraction based on a combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.

<sup>2</sup> T-matrix pole with 2 poles, 2 channels ( $\pi^0\eta$  and  $K\bar{K}$ ).

<sup>3</sup> Coupled-channel analysis of both the  $\eta\pi$  and  $\eta'\pi$  systems using ADOLPH 15 data. Supersedes JACKURA 18. Performed by JPAC.

<sup>4</sup> Amplitude did not include dispersive corrections. From analysis of  $\eta\pi$  mode.

NODE=M012PP

NODE=M012PP

NODE=M012PP

→ UNCHECKED ←

NODE=M012PP;LINKAGE=B

NODE=M012PP;LINKAGE=C

NODE=M012PP;LINKAGE=D

NODE=M012PP;LINKAGE=A

 $a_2(1320)$  MASS

VALUE (MeV)	DOCUMENT ID
<b>1318.2 ± 0.6 OUR AVERAGE</b>	Includes data from the 4 datablocks that follow this one. Error includes scale factor of 1.2.

NODE=M012M0

NODE=M012M0

**3π MODE**

NODE=M012M1  
NODE=M012M1

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

**1318.6 ± 1.3 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

1314.5 <sup>+4.0</sup> <sub>-3.3</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 π <sup>-</sup> p → π <sup>-</sup> π <sup>+</sup> π <sup>-</sup> p
1326 ± 2 ± 2		CHUNG	02	B852	18.3 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> π <sup>-</sup> p
1317 ± 3		BARBERIS	98B		450 p p → p <sub>f</sub> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> p <sub>s</sub>
1323 ± 4 ± 3		ACCIARRI	97T	L3	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup>
1320 ± 7		ALBRECHT	97B	ARG	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup>
1311.3 ± 1.6 ± 3.0	72.4k	AMELIN	96	VES	36 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> n
1310 ± 5		ARMSTRONG	90	OMEG 0	300.0 p p → p p π <sup>+</sup> π <sup>-</sup> π <sup>0</sup>
1323.8 ± 2.3	4022	AUGUSTIN	89	DM2 ±	J/ψ → ρ <sup>±</sup> a <sub>2</sub> <sup>∓</sup>
1320.6 ± 3.1	3562	AUGUSTIN	89	DM2 0	J/ψ → ρ <sup>0</sup> a <sub>2</sub> <sup>0</sup>
1317 ± 2	25k	<sup>2</sup> DAUM	80C	SPEC -	63,94 π <sup>-</sup> p → 3π p
1320 ± 10	1097	<sup>2</sup> BALTAY	78B	HBC +0	15 π <sup>+</sup> p → p 4π
1306 ± 8		FERRERSORIA	78	OMEG -	9 π <sup>-</sup> p → p 3π
1318 ± 7	1.6k	<sup>2</sup> EMMS	75	DBC 0	4 π <sup>+</sup> n → p (3π) <sup>0</sup>
1315 ± 5		<sup>2</sup> ANTIPOV	73C	CNTR -	25,40 π <sup>-</sup> p → p η π <sup>-</sup>
1306 ± 9	1580	CHALOUKKA	73	HBC -	3.9 π <sup>-</sup> p

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1321 ± 1 <sup>+0</sup> <sub>-7</sub>	420k	<sup>3</sup> ALEKSEEV	10	COMP	190 π <sup>-</sup> P b → π <sup>-</sup> π <sup>-</sup> π <sup>+</sup> P b'
1300 ± 2 ± 4	18k	<sup>4</sup> SCHEGELSKY	06	RVUE 0	γγ → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup>
1305 ± 14		CONDO	93	SHF	γγ → n π <sup>+</sup> π <sup>+</sup> π <sup>-</sup>
1310 ± 2		<sup>2</sup> EVANGELIS...	81	OMEG -	12 π <sup>-</sup> p → 3π p
1343 ± 11	490	BALTAY	78B	HBC 0	15 π <sup>+</sup> p → Δ 3π
1309 ± 5	5k	BINNIE	71	MMS -	π <sup>-</sup> p near a <sub>2</sub> thresh- old
1299 ± 6	28k	BOWEN	71	MMS -	5 π <sup>-</sup> p
1300 ± 6	24k	BOWEN	71	MMS +	5 π <sup>+</sup> p
1309 ± 4	17k	BOWEN	71	MMS -	7 π <sup>-</sup> p
1306 ± 4	941	ALSTON...	70	HBC +	7.0 π <sup>+</sup> p → 3π p

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3


<sup>1</sup> Statistical error negligible.

<sup>2</sup> From a fit to J<sup>P</sup> = 2<sup>+</sup> ρ π partial wave.

<sup>3</sup> Superseded by AGHASYAN 2018B.

<sup>4</sup> From analysis of L3 data at 183–209 GeV.

NODE=M012M1;LINKAGE=D  
NODE=M012M1;LINKAGE=P  
NODE=M012M1;LINKAGE=C  
NODE=M012M1;LINKAGE=SC



**$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M2  
NODE=M012M2**1318.1± 0.7 OUR AVERAGE**

1319 ± 5	4700	<sup>1,2</sup> CLELAND	82B	SPEC	+	50 $\pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
1324 ± 6	5200	<sup>1,2</sup> CLELAND	82B	SPEC	-	50 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
1320 ± 2	4000	CHABAUD	80	SPEC	-	17 $\pi^- A \rightarrow K_S^0 K^- A$	
1312 ± 4	11000	CHABAUD	78	SPEC	-	9.8 $\pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
1316 ± 2	4730	CHABAUD	78	SPEC	-	18.8 $\pi^- p \rightarrow K^- K_S^0 p$	
1318 ± 1		<sup>1,3</sup> MARTIN	78D	SPEC	-	10 $\pi^- p \rightarrow K_S^0 K^- p$	
1320 ± 2	2724	MARGULIE	76	SPEC	-	23 $\pi^- p \rightarrow K^- K_S^0 p$	
1313 ± 4	730	FOLEY	72	CNTR	-	20.3 $\pi^- p \rightarrow K^- K_S^0 p$	
1319 ± 3	1500	<sup>3</sup> GRAYER	71	ASPK	-	17.2 $\pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1304 ± 10	870	<sup>4</sup> SCHEGELSKY	06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
1330 ± 11	1000	<sup>1,2</sup> CLELAND	82B	SPEC	+	30 $\pi^+ p \rightarrow K_S^0 K^+ p$	
1324 ± 5	350	HYAMS	78	ASPK	+	12.7 $\pi^+ p \rightarrow K^+ K_S^0 p$	

<sup>1</sup> From a fit to  $J^P = 2^+$  partial wave.<sup>2</sup> Number of events evaluated by us.<sup>3</sup> Systematic error in mass scale subtracted.<sup>4</sup> From analysis of L3 data at 91 and 183–209 GeV.NODE=M012M2;LINKAGE=P  
NODE=M012M2;LINKAGE=W  
NODE=M012M2;LINKAGE=S  
NODE=M012M2;LINKAGE=SC **$\eta\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M3  
NODE=M012M3**1317.7± 1.4 OUR AVERAGE**

1308 ± 9		BARBERIS	00H			450 $p p \rightarrow p_f \eta \pi^0 p_s$	
1316 ± 9		BARBERIS	00H			450 $p p \rightarrow \Delta_f^{++} \eta \pi^- p_s$	OCCUR=2
1317 ± 1 ± 2		THOMPSON	97	MPS		18 $\pi^- p \rightarrow \eta \pi^- p$	
1315 ± 5 ± 2		<sup>1</sup> AMSLER	94D	CBAR		0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$	
1325.1 ± 5.1		AOYAGI	93	BKEI		$\pi^- p \rightarrow \eta \pi^- p$	
1317.7 ± 1.4 ± 2.0		BELADIDZE	93	VES		37 $\pi^- N \rightarrow \eta \pi^- N$	
1323 ± 8	1000	<sup>2</sup> KEY	73	OSPK	-	6 $\pi^- p \rightarrow p \pi^- \eta$	
1307 ± 1 ± 6		<sup>3</sup> JACKURA	18	RVUE		$\pi^- p \rightarrow \eta \pi^- p$	
1315 ± 12		<sup>4</sup> ADOLPH	15	COMP		191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$	
1324 ± 5		ARMSTRONG	93C	E760	0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
1336.2 ± 1.7	2561	DELFOSE	81	SPEC	+	$\pi^\pm p \rightarrow p \pi^\pm \eta$	OCCUR=2
1330.7 ± 2.4	1653	DELFOSE	81	SPEC	-	$\pi^\pm p \rightarrow p \pi^\pm \eta$	
1324 ± 8	6200	<sup>2,5</sup> CONFORTO	73	OSPK	-	6 $\pi^- p \rightarrow p \pi^- \eta$	

<sup>1</sup> The systematic error of 2 MeV corresponds to the spread of solutions.<sup>2</sup> Error includes 5 MeV systematic mass-scale error.<sup>3</sup> Superseded by RODAS 19.<sup>4</sup> ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the  $\eta\pi$  and  $\rho\pi$  channels into account.<sup>5</sup> Missing mass with enriched MMS =  $\eta\pi^-$ ,  $\eta = 2\gamma$ .NODE=M012M3;LINKAGE=DD  
NODE=M012M3;LINKAGE=E  
NODE=M012M3;LINKAGE=B  
NODE=M012M3;LINKAGE=A  
NODE=M012M3;LINKAGE=M **$\eta'\pi$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M4  
NODE=M012M4**1322 ± 7 OUR AVERAGE**

1318 ± 8 $^{+3}_{-5}$		IVANOV	01	B852		18 $\pi^- p \rightarrow \eta' \pi^- p$	
1327.0 ± 10.7		BELADIDZE	93	VES		37 $\pi^- N \rightarrow \eta' \pi^- N$	

**$a_2(1320)$  WIDTH**

NODE=M012210

**3 $\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

 NODE=M012W1  
 NODE=M012W1
**105.0<sup>+1.7</sup><sub>-1.9</sub> OUR AVERAGE**

106.6 <sup>+3.4</sup> <sub>-7.0</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	
108 $\pm 3$ $\pm 15$		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
120 $\pm 10$		BARBERIS	98B		450 $p p \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$	
105 $\pm 10$ $\pm 11$		ACCIARRI	97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
120 $\pm 10$		ALBRECHT	97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
103.0 $\pm 6.0$ $\pm 3.3$	72.4k	AMELIN	96	VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
120 $\pm 10$		ARMSTRONG	90	OMEG 0	300.0 $p p \rightarrow p p \pi^+ \pi^- \pi^0$	
107.0 $\pm 9.7$	4022	AUGUSTIN	89	DM2 $\pm$	$J/\psi \rightarrow \rho^\pm a_2^\mp$	
118.5 $\pm 12.5$	3562	AUGUSTIN	89	DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$	OCCUR=2
97 $\pm 5$		<sup>2</sup> EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow 3\pi p$	
96 $\pm 9$	25k	<sup>2</sup> DAUM	80C	SPEC -	63,94 $\pi^- p \rightarrow 3\pi p$	
110 $\pm 15$	1097	<sup>2</sup> BALTAY	78B	HBC +0	15 $\pi^+ p \rightarrow p 4\pi^0$	
112 $\pm 18$	1.6k	<sup>2</sup> EMMS	75	DBC 0	4 $\pi^+ n \rightarrow p(3\pi)^0$	
122 $\pm 14$	1.2k	<sup>2,3</sup> WAGNER	75	HBC 0	7 $\pi^+ p \rightarrow \Delta^{++}(3\pi)^0$	
115 $\pm 15$		<sup>2</sup> ANTIPOV	73C	CNTR -	25,40 $\pi^- p \rightarrow p \eta \pi^-$	
99 $\pm 15$	1580	CHALOUPKA	73	HBC -	3.9 $\pi^- p$	
105 $\pm 5$	28k	BOWEN	71	MMS -	5 $\pi^- p$	
99 $\pm 5$	24k	BOWEN	71	MMS +	5 $\pi^+ p$	OCCUR=2
103 $\pm 5$	17k	BOWEN	71	MMS -	7 $\pi^- p$	OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

110 $\pm 2$ $^{+2}_{-15}$	420k	<sup>4</sup> ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
117 $\pm 6$ $\pm 20$	18k	<sup>5</sup> SCHEGELSKY	06	RVUE 0	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	
120 $\pm 40$		CONDO	93	SHF	$\gamma p \rightarrow n \pi^+ \pi^+ \pi^-$	
115 $\pm 14$	490	BALTAY	78B	HBC 0	15 $\pi^+ p \rightarrow \Delta 3\pi$	OCCUR=2
72 $\pm 16$	5k	BINNIE	71	MMS -	$\pi^- p$ near $a_2$ thresh- old	OCCUR=2
79 $\pm 12$	941	ALSTON-...	70	HBC +	7.0 $\pi^+ p \rightarrow 3\pi p$	

<sup>1</sup> Statistical error negligible.

<sup>2</sup> From a fit to  $J^P = 2^+ \rho\pi$  partial wave.

<sup>3</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>4</sup> Superseded by AGHASYAN 2018B.

<sup>5</sup> From analysis of L3 data at 183–209 GeV.

 NODE=M012W1;LINKAGE=C  
 NODE=M012W1;LINKAGE=P  
 NODE=M012W1;LINKAGE=S  
 NODE=M012W1;LINKAGE=E  
 NODE=M012W1;LINKAGE=SC
 **$K\bar{K}$  AND  $\eta\pi$  MODES**

VALUE (MeV)	DOCUMENT ID
-------------	-------------

 NODE=M012W0  
 NODE=M012W0
**107  $\pm 5$  OUR ESTIMATE**

→ UNCHECKED ←

**110.4 $\pm 1.7$  OUR AVERAGE** Includes data from the 2 datablocks that follow this one. **$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

 NODE=M012W2  
 NODE=M012W2

The data in this block is included in the average printed for a previous datablock.

**109.8 $\pm 2.4$  OUR AVERAGE**

112 $\pm 20$	4700	<sup>1,2</sup> CLELAND	82B	SPEC +	50 $\pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
120 $\pm 25$	5200	<sup>1,2</sup> CLELAND	82B	SPEC -	50 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
106 $\pm 4$	4000	CHABAUD	80	SPEC -	17 $\pi^- A \rightarrow K_S^0 K^- A$	
126 $\pm 11$	11000	CHABAUD	78	SPEC -	9.8 $\pi^- p \rightarrow K^- K_S^0 p$	
101 $\pm 8$	4730	CHABAUD	78	SPEC -	18.8 $\pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
113 $\pm 4$		<sup>1,3</sup> MARTIN	78D	SPEC -	10 $\pi^- p \rightarrow K_S^0 K^- p$	
105 $\pm 8$	2724	<sup>3</sup> MARGULIE	76	SPEC -	23 $\pi^- p \rightarrow K^- K_S^0 p$	
113 $\pm 19$	730	FOLEY	72	CNTR -	20.3 $\pi^- p \rightarrow K^- K_S^0 p$	
123 $\pm 13$	1500	<sup>3</sup> GRAYER	71	ASPK -	17.2 $\pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

120 ±15	870	<sup>4</sup> SCHEGELSKY	06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$
121 ±51	1000	<sup>1,2</sup> CLELAND	82B	SPEC	+	$30 \pi^+ p \rightarrow K_S^0 K^+ p$
110 ±18	350	HYAMS	78	ASPK	+	$12.7 \pi^+ p \rightarrow K^+ K_S^0 p$

<sup>1</sup> From a fit to  $J^P = 2^+$  partial wave.

<sup>2</sup> Number of events evaluated by us.

<sup>3</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>4</sup> From analysis of L3 data at 91 and 183–209 GeV.

NODE=M012W2;LINKAGE=P  
 NODE=M012W2;LINKAGE=W  
 NODE=M012W2;LINKAGE=S  
 NODE=M012W2;LINKAGE=SC

## $\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012W3  
 NODE=M012W3

### 111.1± 2.4 OUR AVERAGE

115 ±20		BARBERIS	00H			$450 p p \rightarrow p_f \eta \pi^0 p_s$
112 ±14		BARBERIS	00H			$450 p p \rightarrow \Delta_f^{++} \eta \pi^- p_s$
112 ± 3 ±2		<sup>1</sup> AMSLER	94D	CBAR		$0.0 \bar{p} p \rightarrow \pi^0 \pi^0 \eta$
103 ± 6 ±3		BELADIDZE	93	VES		$37 \pi^- N \rightarrow \eta \pi^- N$
112.2± 5.7	2561	DELFOSSÉ	81	SPEC	+	$\pi^\pm p \rightarrow p \pi^\pm \eta$
116.6± 7.7	1653	DELFOSSÉ	81	SPEC	-	$\pi^\pm p \rightarrow p \pi^\pm \eta$
108 ± 9	1000	KEY	73	OSPK	-	$6 \pi^- p \rightarrow p \pi^- \eta$

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

112 ± 1 ±8		<sup>2</sup> JACKURA	18	RVUE		$\pi^- p \rightarrow \eta \pi^- p$
119 ±14		<sup>3</sup> ADOLPH	15	COMP		$191 \pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
127 ± 2 ±2		<sup>4</sup> THOMPSON	97	MPS		$18 \pi^- p \rightarrow \eta \pi^- p$
118 ±10		ARMSTRONG	93C	E760	0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
104 ± 9	6200	<sup>5</sup> CONFORTO	73	OSPK	-	$6 \pi^- p \rightarrow p \text{M}^-$

<sup>1</sup> The systematic error of 2 MeV corresponds to the spread of solutions.

<sup>2</sup> Superseded by RODAS 19.

<sup>3</sup> ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the  $\eta\pi$  and  $\rho\pi$  channels into account.

<sup>4</sup> Resolution is not unfolded.

<sup>5</sup> Missing mass with enriched MMS =  $\eta\pi^-$ ,  $\eta = 2\gamma$ .

NODE=M012W3;LINKAGE=DD  
 NODE=M012W3;LINKAGE=C  
 NODE=M012W3;LINKAGE=B

NODE=M012W3;LINKAGE=A  
 NODE=M012W3;LINKAGE=M

## $\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

### 119±25 OUR AVERAGE

140±35±20	IVANOV	01	B852	$18 \pi^- p \rightarrow \eta' \pi^- p$
106±32	BELADIDZE	93	VES	$37 \pi^- N \rightarrow \eta' \pi^- N$

NODE=M012W4  
 NODE=M012W4

## $a_2(1320)$ DECAY MODES

NODE=M012215;NODE=M012

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $3\pi$	(70.1 ±2.7 ) %	S=1.2
$\Gamma_2$ $\rho(770)\pi$		
$\Gamma_3$ $f_2(1270)\pi$		
$\Gamma_4$ $\rho(1450)\pi$		
$\Gamma_5$ $\eta\pi$	(14.5 ±1.2 ) %	
$\Gamma_6$ $\omega\pi\pi$	(10.6 ±3.2 ) %	S=1.3
$\Gamma_7$ $K\bar{K}$	( 4.9 ±0.8 ) %	
$\Gamma_8$ $\eta'(958)\pi$	( 5.5 ±0.9 ) × 10 <sup>-3</sup>	
$\Gamma_9$ $\pi^\pm\gamma$	( 2.91±0.27 ) × 10 <sup>-3</sup>	
$\Gamma_{10}$ $\gamma\gamma$	( 9.4 ±0.7 ) × 10 <sup>-6</sup>	
$\Gamma_{11}$ $e^+e^-$	< 5 × 10 <sup>-9</sup>	CL=90%

DESIG=1  
 DESIG=11  
 DESIG=12  
 DESIG=13  
 DESIG=3  
 DESIG=4  
 DESIG=2  
 DESIG=8  
 DESIG=7  
 DESIG=9  
 DESIG=10

## CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 18 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 9.3$  for 15 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_5$	10		
$x_6$	-89	-46	
$x_7$	-1	-2	-24
	$x_1$	$x_5$	$x_6$

### $a_2(1320)$ PARTIAL WIDTHS

#### $\Gamma(\eta\pi)$

$\Gamma_5$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18.5 \pm 3.0$	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$
----------------	-----	-----------------------------	------	---	--

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$  keV and SU(3) relations.

NODE=M012220

NODE=M012W6  
NODE=M012W6

NODE=M012W6;LINKAGE=SC

#### $\Gamma(K\bar{K})$

$\Gamma_7$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.0^{+2.0}_{-1.5}$	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$
---------------------	-----	-----------------------------	------	---	--

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$  keV and SU(3) relations.

NODE=M012W5  
NODE=M012W5

NODE=M012W5;LINKAGE=SC

#### $\Gamma(\pi^\pm\gamma)$

$\Gamma_9$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

#### **311 ± 25 OUR AVERAGE**

$358 \pm 6 \pm 42$		<sup>1</sup> ADOLPH 14	COMP	-	$190 \pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$
--------------------	--	------------------------	------	---	--

$284 \pm 25 \pm 25$	7.1k	MOLCHANOV 01	SELX		$600 \pi^- \text{A} \rightarrow \pi^+ \pi^- \pi^- \text{A}$
---------------------	------	--------------	------	--	---

$295 \pm 60$		CIHANGIR 82	SPEC	+	$200 \pi^+ \text{A}$
--------------	--	-------------	------	---	----------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$461 \pm 110$		<sup>2</sup> MAY 77	SPEC	±	$9.7 \gamma \text{A}$
---------------	--	---------------------	------	---	-----------------------

<sup>1</sup> Primakoff reaction using  $a_2(1320) \rightarrow 3\pi$  branching ratio of 70.1%.

<sup>2</sup> Assuming one-pion exchange.

NODE=M012W7  
NODE=M012W7

NODE=M012W7;LINKAGE=AD  
NODE=M012W;LINKAGE=M2

#### $\Gamma(\gamma\gamma)$

$\Gamma_{10}$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

#### **1.00 ± 0.06 OUR AVERAGE**

$0.98 \pm 0.05 \pm 0.09$		ACCIARRI 97T	L3		$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
--------------------------	--	--------------	----	--	---

$0.96 \pm 0.03 \pm 0.13$		ALBRECHT 97B	ARG		$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
--------------------------	--	--------------	-----	--	---

$1.26 \pm 0.26 \pm 0.18$	36	BARU 90	MD1		$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
--------------------------	----	---------	-----	--	---

$1.00 \pm 0.07 \pm 0.15$	415	BEHREND 90C	CELL	0	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
--------------------------	-----	-------------	------	---	---

$1.03 \pm 0.13 \pm 0.21$		BUTLER 90	MRK2		$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$
--------------------------	--	-----------	------	--	---

$1.01 \pm 0.14 \pm 0.22$	85	OEST 90	JADE		$e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$
--------------------------	----	---------	------	--	--

$0.90 \pm 0.27 \pm 0.15$	56	<sup>1</sup> ALTHOFF 86	TASS	0	$e^+ e^- \rightarrow e^+ e^- 3\pi$
--------------------------	----	-------------------------	------	---	------------------------------------

$1.14 \pm 0.20 \pm 0.26$		<sup>2</sup> ANTREASYAN 86	CBAL	0	$e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$
--------------------------	--	----------------------------	------	---	--

$1.06 \pm 0.18 \pm 0.19$		BERGER 84C	PLUT	0	$e^+ e^- \rightarrow e^+ e^- 3\pi$
--------------------------	--	------------	------	---	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.81 \pm 0.19^{+0.42}_{-0.11}$	35	<sup>1</sup> BEHREND 82C	CELL	0	$e^+ e^- \rightarrow e^+ e^- 3\pi$
---------------------------------	----	--------------------------	------	---	------------------------------------

$0.77 \pm 0.18 \pm 0.27$	22	<sup>2</sup> EDWARDS 82F	CBAL	0	$e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$
--------------------------	----	--------------------------	------	---	--

<sup>1</sup> From  $\rho\pi$  decay mode.

<sup>2</sup> From  $\eta\pi^0$  decay mode.

NODE=M012W9  
NODE=M012W9

NODE=M012W;LINKAGE=F  
NODE=M012W;LINKAGE=G

$\Gamma(e^+e^-)$  $\Gamma_{11}$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.56	90	ACHASOV	00K SND	$e^+e^- \rightarrow \pi^0\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
<25	90	VOROBYEV	88 ND	$e^+e^- \rightarrow \pi^0\eta$

NODE=M012W10  
 NODE=M012W10

 $a_2(1320) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M012223

 $\Gamma(3\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_{10}/\Gamma$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
$0.65 \pm 0.02 \pm 0.02$	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
<sup>1</sup> From analysis of L3 data at 183–209 GeV.				

NODE=M012G2  
 NODE=M012G2

NODE=M012G2;LINKAGE=SC

 $\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_5\Gamma_{10}/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
$0.145^{+0.097}_{-0.034}$	<sup>1</sup> UEHARA 09A	BELL	$e^+e^- \rightarrow e^+e^-\eta\pi^0$
<sup>1</sup> From the $D_2$ -wave. The fraction of the $D_0$ -wave is $3.4^{+2.3}_{-1.1}\%$ .			

NODE=M012G01  
 NODE=M012G01

NODE=M012G01;LINKAGE=UE

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_7\Gamma_{10}/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
$0.126 \pm 0.007 \pm 0.028$	<sup>1</sup> ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
••• We do not use the following data for averages, fits, limits, etc. •••			
$0.081 \pm 0.006 \pm 0.027$	<sup>2</sup> ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
<sup>1</sup> Using an incoherent background.			
<sup>2</sup> Using a coherent background.			

NODE=M012G1  
 NODE=M012G1

OCCUR=2

NODE=M012G1;LINKAGE=A  
 NODE=M012G1;LINKAGE=B

 $a_2(1320) \text{ BRANCHING RATIOS}$ 

NODE=M012225

 $[\Gamma(f_2(1270)\pi) + \Gamma(\rho(1450)\pi)]/\Gamma(\rho(770)\pi)$  $(\Gamma_3+\Gamma_4)/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.12	90	ABRAMOVI...	70B HBC	-	$3.93 \pi^- p$

NODE=M012R9  
 NODE=M012R9

 $\Gamma(\rho(770)\pi)/\Gamma(f_2(1270)\pi)$  $\Gamma_2/\Gamma_3$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$16.5^{+1.2}_{-2.4}$	46M	<sup>1</sup> AGHASYAN 18B	COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$

NODE=M012R00  
 NODE=M012R00

<sup>1</sup> Statistical error negligible.

NODE=M012R00;LINKAGE=A

 $\Gamma(\eta\pi)/\Gamma(3\pi)$  $\Gamma_5/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.207 ± 0.018 OUR FIT</b>					
<b>0.213 ± 0.020 OUR AVERAGE</b>					
0.18 ± 0.05		FORINO 76	HBC		11 $\pi^- p$
0.22 ± 0.05	52	ANTIPOV 73	CNTR	-	40 $\pi^- p$
0.211 ± 0.044	149	CHALOUPKA 73	HBC	-	3.9 $\pi^- p$
0.246 ± 0.042	167	ALSTON-... 71	HBC	+	7.0 $\pi^+ p$
0.25 ± 0.09	15	BOECKMANN 70	HBC	+	5.0 $\pi^+ p$
0.23 ± 0.08	22	ASCOLI 68	HBC	-	5 $\pi^- p$
0.12 ± 0.08		CHUNG 68	HBC	-	3.2 $\pi^- p$
0.22 ± 0.09		CONTE 67	HBC	-	11.0 $\pi^- p$

NODE=M012R3  
 NODE=M012R3

 $\Gamma(\omega\pi\pi)/\Gamma(3\pi)$  $\Gamma_6/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.15 ± 0.05 OUR FIT</b> Error includes scale factor of 1.3.					
<b>0.15 ± 0.05 OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below.					
0.28 ± 0.09	60	DIAZ 74	DBC	0	6 $\pi^+ n$
0.18 ± 0.08		<sup>1</sup> KARSHON 74	HBC		Avg. of above two
0.10 ± 0.05	279	<sup>2</sup> CHALOUPKA 73	HBC	-	3.9 $\pi^- p$
••• We do not use the following data for averages, fits, limits, etc. •••					
0.29 ± 0.08	140	<sup>1</sup> KARSHON 74	HBC	0	4.9 $\pi^+ p$
0.10 ± 0.04	60	<sup>1</sup> KARSHON 74	HBC	+	4.9 $\pi^+ p$
0.19 ± 0.08		DEFOIX 73	HBC	0	0.7 $\bar{p}p$

NODE=M012R12  
 NODE=M012R12

OCCUR=3

OCCUR=2




<sup>1</sup> KARSHON 74 suggest an additional  $I = 0$  state strongly coupled to  $\omega\pi\pi$  which could explain discrepancies in branching ratios and masses. We use a central value and a systematic spread.

<sup>2</sup> Decays to  $b_1(1040)\pi$ ,  $b_1 \rightarrow \omega\pi$ . Error increased to account for possible systematic errors of complicated analysis.

NODE=M012R12;LINKAGE=K

NODE=M012R12;LINKAGE=01



**$\Gamma(K\bar{K})/\Gamma(3\pi)$**

VALUE EVTS DOCUMENT ID TECN CHG COMMENT  $\Gamma_7/\Gamma_1$

**0.070±0.012 OUR FIT**

**0.078±0.017**

CHABAUD 78 RVUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.011±0.003		<sup>1</sup> BERTIN	98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
0.056±0.014	50	<sup>2</sup> CHALOUPKA	73	HBC	- 3.9 $\pi^- p$
0.097±0.018	113	<sup>2</sup> ALSTON-...	71	HBC	+ 7.0 $\pi^+ p$
0.06 ±0.03		<sup>2</sup> ABRAMOVI...	70B	HBC	- 3.93 $\pi^- p$
0.054±0.022		<sup>2</sup> CHUNG	68	HBC	- 3.2 $\pi^- p$

<sup>1</sup> Using  $4\pi$  data from BERTIN 97D.

<sup>2</sup> Included in CHABAUD 78 review.

NODE=M012R1;LINKAGE=BE  
NODE=M012R1;LINKAGE=C

**$\Gamma(K\bar{K})/\Gamma(\eta\pi)$**

VALUE DOCUMENT ID TECN COMMENT  $\Gamma_7/\Gamma_5$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ±0.22 <sup>+0.09</sup> / <sub>-0.11</sub>	<sup>1</sup> KOPF	21	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
0.352±0.011±0.175	<sup>2</sup> ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
0.08 ±0.02	<sup>3</sup> BERTIN	98B	OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$

<sup>1</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.

<sup>2</sup> Residues from T-matrix pole with 2 poles, 2 channels ( $\pi^0\eta$  and  $K\bar{K}$ ).

<sup>3</sup> Using  $\eta\pi\pi$  data from AMSLER 94D.

NODE=M012R14;LINKAGE=B

NODE=M012R14;LINKAGE=A  
NODE=M012R14;LINKAGE=BE

**$\Gamma(\eta\pi)/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$**

VALUE EVTS DOCUMENT ID TECN CHG COMMENT  $\Gamma_5/(\Gamma_1+\Gamma_5+\Gamma_7)$

**0.162±0.012 OUR FIT**

**0.140±0.028 OUR AVERAGE**

0.13 ±0.04		ESPIGAT	72	HBC	± 0.0 $\bar{p}p$
0.15 ±0.04	34	BARNHAM	71	HBC	+ 3.7 $\pi^+ p$

NODE=M012R2  
NODE=M012R2

$\Gamma(K\bar{K})/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$  $\Gamma_7/(\Gamma_1+\Gamma_5+\Gamma_7)$ 

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M012R8  
NODE=M012R8

**0.054±0.009 OUR FIT****0.048±0.012 OUR AVERAGE**

0.05 ±0.02		TOET	73	HBC	+	5 $\pi^+ p$
0.09 ±0.04		TOET	73	HBC	0	5 $\pi^+ p$
0.03 ±0.02	8	<sup>1</sup> DAMERI	72	HBC	-	11 $\pi^- p$
0.06 ±0.03	17	BARNHAM	71	HBC	+	3.7 $\pi^+ p$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.020±0.004		<sup>2</sup> ESPIGAT	72	HBC	±	0.0 $\bar{p} p$
-------------	--	----------------------	----	-----	---	-----------------

<sup>1</sup> Montanet agrees. Vlada.<sup>2</sup> Not averaged because of discrepancy between masses from  $K\bar{K}$  and  $\rho\pi$  modes.

NODE=M012R8;LINKAGE=01  
NODE=M012R8;LINKAGE=A

 $\Gamma(\eta'(958)\pi)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

NODE=M012R4  
NODE=M012R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.006	95	ALDE	92B	GAM2		38,100 $\pi^- p \rightarrow \eta' \pi^0 n$
<0.02	97	BARNHAM	71	HBC	+	3.7 $\pi^+ p$
0.004±0.004		<sup>1</sup> BOESEBECK	68	HBC	+	8 $\pi^+ p$

<sup>1</sup> No longer valid since  $\Gamma(K\bar{K})/\Gamma(3\pi)$  value has changed (MORRISON 71).

NODE=M012R4;LINKAGE=B

 $\Gamma(\eta'(958)\pi)/\Gamma(3\pi)$  $\Gamma_8/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

NODE=M012R5  
NODE=M012R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011	90	EISENSTEIN	73	HBC	-	5 $\pi^- p$
<0.04		ALSTON-...	71	HBC	+	7.0 $\pi^+ p$
0.04 $\begin{smallmatrix} +0.03 \\ -0.04 \end{smallmatrix}$		BOECKMANN	70	HBC	0	5.0 $\pi^+ p$

 $\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$  $\Gamma_8/\Gamma_5$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M012R13  
NODE=M012R13

**0.038±0.005 OUR AVERAGE**

0.05 ±0.02	ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
0.032±0.009	ABELE	97C	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta'$
0.047±0.010±0.004	<sup>1</sup> BELADIDZE	93	VES	37 $\pi^- N \rightarrow a_2^- N$
0.034±0.008±0.005	BELADIDZE	92	VES	36 $\pi^- C \rightarrow a_2^- C$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.046±0.015 $\begin{smallmatrix} +0.07 \\ -0.006 \end{smallmatrix}$	<sup>2</sup> KOPF	21	RVUE	0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
---	-------------------	----	------	--

OCCUR=2

<sup>1</sup> Using  $B(\eta' \rightarrow \pi^+ \pi^- \eta) = 0.441$ ,  $B(\eta \rightarrow \gamma \gamma) = 0.389$  and  $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 0.236$ .

NODE=M012R13;LINKAGE=A

<sup>2</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi$ ,  $\eta'\pi$  and  $K\bar{K}$  systems.

NODE=M012R13;LINKAGE=C

 $\Gamma(\pi^\pm \gamma)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M012R11  
NODE=M012R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.005 $\begin{smallmatrix} +0.005 \\ -0.003 \end{smallmatrix}$	<sup>1</sup> EISENBERG	72	HBC	4.3,5.25,7.5 $\gamma p$
--	------------------------	----	-----	-------------------------

<sup>1</sup> Pion-exchange model used in this estimation.

NODE=M012R11;LINKAGE=R

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-9}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M012R15  
NODE=M012R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90	ACHASOV	00K	SND $e^+ e^- \rightarrow \pi^0 \pi^0$
----	----	---------	-----	---------------------------------------

**a<sub>2</sub>(1320) REFERENCES**

NODE=M012

KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)	REFID=61470
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=59554
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)	REFID=59003
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=55911
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
MOLCHANOV	01	PL B521 171	V.V. Molchanov <i>et al.</i>	(FNAL SELEX Collab.)	REFID=48559
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
ABELE	97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45531
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=45418
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=45584
AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44649
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BELADIDZE	93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43598
CONDO	93	PR D48 3045	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=43600
ALDE	92B	ZPHY C54 549	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41852
BELADIDZE	92	ZPHY C54 235	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42171
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch	(WA76 Coll.)	REFID=41375
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41356
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
ALTHOFF	86	ZPHY C31 537	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21287
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
BERGER	84C	PL 149B 427	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=21286
BEHREND	82C	PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (erratum)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
CHANGIR	82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
CHABAUD	80	NP B175 189	V. Chabaud <i>et al.</i>	(CERN, MPIM, AMST)	REFID=21274
DAUM	80C	PL 89B 276	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21275
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
CHABAUD	78	NP B145 349	V. Chabaud <i>et al.</i>	(CERN, MPIM)	REFID=21267
FERRERSORIA	78	PL 74B 287	A. Ferrer Soria <i>et al.</i>	(ORSAY, CERN, CDEF+)	REFID=21270
HYAMS	78	NP B146 303	B.D. Hyams <i>et al.</i>	(CERN, MPIM, ATEN)	REFID=21271
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA) JP	REFID=21272
MAY	77	PR D16 1983	E.N. May <i>et al.</i>	(ROCH, CORN)	REFID=20450
FORINO	76	NC 35A 465	A. Forino <i>et al.</i>	(BGNA, FIRZ, GENO, MILA+)	REFID=21259
MARGULIE	76	PR D14 667	M. Margulies <i>et al.</i>	(BNL, CUNY)	REFID=21261
EMMS	75	PL 58B 117	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL) JP	REFID=21254
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP	REFID=20843
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)	REFID=21248
KARSHON	74	PRL 32 852	U. Karshon <i>et al.</i>	(REHO)	REFID=21249
ANTIPOV	73	NP B63 175	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=21238
ANTIPOV	73C	NP B63 153	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=20817
CHALOUPKA	73	PL 44B 211	V. Chaloupka <i>et al.</i>	(CERN)	REFID=21242
CONFORTO	73	PL 45B 154	G. Conforto <i>et al.</i>	(EFI, FNAL, TNTO+)	REFID=21243
DEFOIX	73	PL 43B 141	C. Defoix <i>et al.</i>	(CDEF)	REFID=21244
EISENSTEIN	73	PR D7 278	L. Eisenstein <i>et al.</i>	(ILL)	REFID=21245
KEY	73	PRL 30 503	A.W. Key <i>et al.</i>	(TNTO, EFI, FNAL, WISC)	REFID=21246
TOET	73	NP B63 248	D.Z. Toet <i>et al.</i>	(NIJM, BONN, DURH, TORI)	REFID=20714
DAMERI	72	NC 9A 1	M. Dameri <i>et al.</i>	(GENO, MILA, SACL)	REFID=20338
EISENBERG	72	PR D5 15	Y. Eisenberg <i>et al.</i>	(REHO, SLAC, TELA)	REFID=20098
ESPIGAT	72	NP B36 93	P. Espigat <i>et al.</i>	(CERN, CDEF)	REFID=21232
FOLEY	72	PR D6 747	K.J. Foley <i>et al.</i>	(BNL, CUNY)	REFID=21233
ALSTON...	71	PL 34B 156	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21214
BARNHAM	71	PRL 26 1494	K.W.J. Barnham <i>et al.</i>	(LBL)	REFID=21215
BINNIE	71	PL 36B 257	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=21217
BOWEN	71	PRL 26 1663	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21219
GRAYER	71	PL 34B 333	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=21223
ABRAMOVI...	70B	NP B23 466	M. Abramovich <i>et al.</i>	(CERN) JP	REFID=21195
ALSTON...	70	PL 33B 607	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21196
BOECKMANN	70	NP B16 221	K. Boeckmann <i>et al.</i>	(BONN, DURH, NIJM+)	REFID=21202
ASCOLI	68	PRL 20 1321	G. Ascoli <i>et al.</i>	(ILL) JP	REFID=21171
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
CONTE	67	NC 51A 175	F. Conte <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=21166

**$f_0(1370)$** 

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances" and a note on "Non- $q\bar{q}$  Candidates" in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M147

NODE=M147

 **$f_0(1370)$  T-MATRIX POLE  $\sqrt{s}$** 

NODE=M147PP

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s_{\text{pole}}})$ .

NODE=M147PP

NODE=M147PP

NEW; → UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1250–1440) –i(60–300) OUR ESTIMATE</b>			
[(1200–1500)–i(150–250) MeV OUR 2022 ESTIMATE]			
(1370 ± 40)–i(195 ± 20)	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
(1280.6 ± 1.6 ± 47.4) – i(205.2 ± 1.7 ± 20.7)	<sup>1</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
(1290 ± 50)–i(170 <sup>+20</sup> <sub>-40</sub> )	<sup>2</sup> ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
(1373 ± 15)–i(137 ± 10)	<sup>3</sup> BARGIOTTI	03	OBLX $\bar{p}p$
(1302 ± 17)–i(166 ± 18)	<sup>4</sup> BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
(1312 ± 25 ± 10)–i(109 ± 22 ± 15)	BARBERIS	99D	OMEG $450 pp \rightarrow K^+K^-, \pi^+\pi^-$
(1406 ± 19)–i(80 ± 6)	<sup>5</sup> KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
(1300 ± 20)–i(120 ± 20)	ANISOVICH	98B	RVUE Compilation
(1290 ± 15)–i(145 ± 15)	BARBERIS	97B	OMEG $450 pp \rightarrow pp2(\pi^+\pi^-)$
(1548 ± 40)–i(560 ± 40)	BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
(1380 ± 40)–i(180 ± 25)	ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
(1300 ± 15)–i(115 ± 8)	BUGG	96	RVUE
(1330 ± 50)–i(150 ± 40)	<sup>6</sup> AMSLER	95B	CBAR $\bar{p}p \rightarrow 3\pi^0$
(1360 ± 35)–i(150–300)	<sup>6</sup> AMSLER	95C	CBAR $\bar{p}p \rightarrow \pi^0\eta\eta$
(1390 ± 30)–i(190 ± 40)	<sup>7</sup> AMSLER	95D	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$
1346 – i249	<sup>8,9</sup> JANSSEN	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1214 – i168	<sup>9,10</sup> TORNQVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1364 – i139	AMSLER	94D	CBAR $\bar{p}p \rightarrow \pi^0\pi^0\eta$
(1365 <sup>+20</sup> <sub>-55</sub> )–i(134 ± 35)	ANISOVICH	94	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
(1340 ± 40)–i(127 <sup>+30</sup> <sub>-20</sub> )	<sup>11</sup> BUGG	94	RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0\pi^0$
(1430 ± 5)–i(73 ± 13)	<sup>12</sup> KAMINSKI	94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1420 – i220	<sup>13</sup> AU	87	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

<sup>1</sup> T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ), and BINON 84C ( $\eta\eta'$ ).

NODE=M147PP;LINKAGE=E

<sup>2</sup> Another pole is found at  $(1510 \pm 130) - i(800^{+100}_{-150})$  MeV.

NODE=M147PP;LINKAGE=AO

<sup>3</sup> Coupled channel analysis of  $\pi^+\pi^-\pi^0, K^+K^-\pi^0$ , and  $K^\pm K_S^0 \pi^\mp$ .

NODE=M147PP;LINKAGE=BG

<sup>4</sup> Average between  $\pi^+\pi^-\pi^0$  and  $2(\pi^+\pi^-)$ .

NODE=M147PP;LINKAGE=PC

<sup>5</sup> T-matrix pole on sheet – – –.

NODE=M147PP;LINKAGE=TK

<sup>6</sup> Supersedes ANISOVICH 94.

NODE=M147PP;LINKAGE=K

<sup>7</sup> Coupled-channel analysis of  $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ , and  $\pi^0\pi^0\eta$  on sheet IV. Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.

NODE=M147PP;LINKAGE=A

<sup>8</sup> Analysis of data from FALVARD 88.

NODE=M147PP;LINKAGE=C

<sup>9</sup> The pole is on Sheet III. Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.

NODE=M147PP;LINKAGE=DD

<sup>10</sup> Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

NODE=M147PP;LINKAGE=BB

<sup>11</sup> Reanalysis of ANISOVICH 94 data.

NODE=M147PP;LINKAGE=C1

<sup>12</sup> T-matrix pole on sheet III.

NODE=M147PP;LINKAGE=KM

<sup>13</sup> Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

NODE=M147PP;LINKAGE=H

 **$f_0(1370)$  BREIT-WIGNER MASS**

NODE=M147205

VALUE (MeV)	DOCUMENT ID
<b>1200 to 1500 OUR ESTIMATE</b>	

NODE=M147M

→ UNCHECKED ←

**$\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1400±40		<sup>1</sup> AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 <sup>+6+72</sup> <sub>-7-255</sub>		<sup>2</sup> UEHARA	08A BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1259±55	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^-\pi^+\pi^+$
1309±1±15		<sup>3</sup> BUGG	07A RVUE	0.0 $p\bar{p} \rightarrow 3\pi^0$
1449±13	4.3k	<sup>4</sup> GARMASH	06 BELL	$B^+ \rightarrow K^+\pi^+\pi^-$
1350±50		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
1265±30 <sup>+20</sup> <sub>-35</sub>		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
1434±18±9	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$
1308±10		BARBERIS	99B OMEG	450 $pp \rightarrow p_s p_f \pi^+\pi^-$
1315±50		BELLAZZINI	99 GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
1315±30		ALDE	98 GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0n$
1280±55		BERTIN	98 OBLX	0.05-0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
1186		<sup>5,6</sup> TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472±12		ARMSTRONG	91 OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275±20		BREAKSTONE	90 SFM	62 $pp \rightarrow pp\pi^+\pi^-$
1420±20		AKESSON	86 SPEC	63 $pp \rightarrow pp\pi^+\pi^-$
1256		FROGGATT	77 RVUE	$\pi^+\pi^-\text{ channel}$

<sup>1</sup> Breit-Wigner mass.<sup>2</sup> Breit-Wigner mass. May also be the  $f_0(1500)$ .<sup>3</sup> Reanalysis of ABELE 96C data.<sup>4</sup> Also observed by GARMASH 07 in  $B^0 \rightarrow K_S^0\pi^+\pi^-$  decays. Supersedes GARMASH 05.<sup>5</sup> Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.<sup>6</sup> Also observed by ASNER 00 in  $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$  decaysNODE=M147M1  
NODE=M147M1NODE=M147M1;LINKAGE=BW  
NODE=M147M1;LINKAGE=UE  
NODE=M147M1;LINKAGE=BU  
NODE=M147M1;LINKAGE=GR  
NODE=M147M1;LINKAGE=BB

NODE=M147M1;LINKAGE=FF

 **$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1422±15±28		<sup>1</sup> AAIJ	19H LHCB	$pp \rightarrow D^\pm X$
1360±31±28	430	<sup>2,3</sup> DOBBS	15	$J/\psi \rightarrow \gamma K^+K^-$
1350±48±15	168	<sup>2,3</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K^+K^-$
1440±6		VLADIMIRSK...	06 SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1391±10		TIKHOMIROV	03 SPEC	40.0 $\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440±50		BOLONKIN	88 SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1463±9		ETKIN	82B MPS	23 $\pi^-p \rightarrow n2K_S^0$
1425±15		WICKLUND	80 SPEC	6 $\pi N \rightarrow K^+K^-N$
~1300		POLYCHRO...	79 STRC	7 $\pi^-p \rightarrow n2K_S^0$

<sup>1</sup> From the  $D^\pm \rightarrow K^\pm K^+K^-$  Dalitz plot fit with the isobar model A.<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>3</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 346$  MeV.NODE=M147M2  
NODE=M147M2

OCCUR=4

OCCUR=2

NODE=M147M2;LINKAGE=F  
NODE=M147M2;LINKAGE=A  
NODE=M147M2;LINKAGE=B **$4\pi$  MODE  $2(\pi\pi)_S + \rho\rho$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1395±40		ABELE	01 CBAR	0.0 $\bar{p}d \rightarrow \pi^-4\pi^0p$
1374±38		AMSLER	94 CBAR	0.0 $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
1345±12		ADAMO	93 OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-$
1386±30		GASPERO	93 DBC	0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$
~1410	5751	<sup>1</sup> BETTINI	66 DBC	0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

<sup>1</sup>  $\rho\rho$  dominant.NODE=M147M3  
NODE=M147M3

NODE=M147M3;LINKAGE=BE

 **$\eta\eta$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1262 <sup>+51+82</sup> <sub>-78-103</sub>	<sup>1</sup> UEHARA	10A BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
1430	AMSLER	92 CBAR	0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
1220±40	ALDE	86D GAM4	100 $\pi^-p \rightarrow n2\eta$

<sup>1</sup> Breit-Wigner mass. May also be the  $f_0(1500)$ .NODE=M147M4  
NODE=M147M4

NODE=M147M4;LINKAGE=UE

**COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1330.2^{+5.9}_{-6.5} \pm 5.1$	<sup>1</sup> AAIJ	19H	LHCB $pp \rightarrow D^\pm X$
$1306 \pm 20$	<sup>2</sup> ANISOVICH	03	RVUE

<sup>1</sup> From the  $D^\pm \rightarrow K^\pm K^+ K^-$  Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

<sup>2</sup> K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.

NODE=M147M5  
NODE=M147M5

NODE=M147M5;LINKAGE=A

NODE=M147M;LINKAGE=KM

 **$f_0(1370)$  BREIT-WIGNER WIDTH**

NODE=M147210

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**200 to 500 OUR ESTIMATE**

NODE=M147W  
→ UNCHECKED ←

 **$\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$300 \pm 80$		<sup>1</sup> AUBERT	09L	BABR $B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
$90^{+2}_{-1} \begin{smallmatrix} +50 \\ -22 \end{smallmatrix}$		<sup>2</sup> UEHARA	08A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$298 \pm 21$	2.6k	BONVICINI	07	CLEO $D^+ \rightarrow \pi^- \pi^+ \pi^+$
$126 \pm 25$	4286	<sup>3</sup> GARMASH	06	BELL $B^+ \rightarrow K^+ \pi^+ \pi^-$
$265 \pm 40$		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi \pi^+ \pi^-$
$350 \pm 100 \begin{smallmatrix} +105 \\ -60 \end{smallmatrix}$		ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
$173 \pm 32 \pm 6$	848	AITALA	01A	E791 $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
$222 \pm 20$		BARBERIS	99B	OMEG $450 pp \rightarrow p_S p_f \pi^+ \pi^-$
$255 \pm 60$		BELLAZZINI	99	GAM4 $450 pp \rightarrow pp \pi^0 \pi^0$
$190 \pm 50$		ALDE	98	GAM4 $100 \pi^- p \rightarrow \pi^0 \pi^0 n$
$323 \pm 13$		BERTIN	98	OBLX $0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
$350$		<sup>4,5</sup> TORNVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
$195 \pm 33$		ARMSTRONG	91	OMEG $300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
$285 \pm 60$		BREAKSTONE	90	SFM $62 pp \rightarrow pp\pi^+ \pi^-$
$460 \pm 50$		AKESSON	86	SPEC $63 pp \rightarrow pp\pi^+ \pi^-$
$\sim 400$		<sup>6</sup> FROGGATT	77	RVUE $\pi^+ \pi^-$ channel

<sup>1</sup> The systematic errors are not reported.

<sup>2</sup> Breit-Wigner width. May also be the  $f_0(1500)$ .

<sup>3</sup> Also observed by GARMASH 07 in  $B^0 \rightarrow K_S^0 \pi^+ \pi^-$  decays. Supersedes GARMASH 05.

<sup>4</sup> Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

<sup>5</sup> Also observed by ASNER 00 in  $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$  decays

<sup>6</sup> Width defined as distance between 45 and 135° phase shift.

NODE=M147W1  
NODE=M147W1

NODE=M147W1;LINKAGE=NS  
NODE=M147W1;LINKAGE=UE  
NODE=M147W1;LINKAGE=GR  
NODE=M147W1;LINKAGE=BB

NODE=M147W1;LINKAGE=FF  
NODE=M147W1;LINKAGE=E

 **$K\bar{K}$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$324 \pm 38 \pm 42$	<sup>1</sup> AAIJ	19H	LHCB $pp \rightarrow D^\pm X$
$121 \pm 15$	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$55 \pm 26$	TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
$250 \pm 80$	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$118 \begin{smallmatrix} +138 \\ -16 \end{smallmatrix}$	ETKIN 82B	MPS	$23 \pi^- p \rightarrow n 2 K_S^0$
$160 \pm 30$	WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
$\sim 150$	POLYCHRO... 79	STRC	$7 \pi^- p \rightarrow n 2 K_S^0$

<sup>1</sup> From the  $D^\pm \rightarrow K^\pm K^+ K^-$  Dalitz plot fit with the isobar model A.

NODE=M147W2  
NODE=M147W2

OCCUR=3

NODE=M147W2;LINKAGE=C

**4 $\pi$  MODE 2( $\pi\pi$ ) $_S$ + $\rho\rho$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
275 $\pm$ 55		ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 $\pm$ 61		AMSLER	94	CBAR 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
398 $\pm$ 26		ADAMO	93	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 $\pm$ 50		GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$
$\sim$ 90	5751	<sup>1</sup> BETTINI	66	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$

<sup>1</sup>  $\rho\rho$  dominant.

NODE=M147W3  
NODE=M147W3

NODE=M147W3;LINKAGE=BE

 **$\eta\eta$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
484 $^{+246+246}_{-170-263}$	<sup>1</sup> UEHARA	10A	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$
250	AMSLER	92	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \eta\eta$
320 $\pm$ 40	ALDE	86D	GAM4 100 $\pi^- p \rightarrow n 2\eta$

<sup>1</sup> Breit-Wigner width. May also be the  $f_0(1500)$ .

NODE=M147W4  
NODE=M147W4

NODE=M147W4;LINKAGE=UE

**COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
147 $^{+30}_{-50}$	<sup>1</sup> ANISOVICH	03	RVUE

<sup>1</sup> K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta\eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.

NODE=M147W5  
NODE=M147W5

NODE=M147W;LINKAGE=KM

 **$f_0(1370)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $4\pi$	seen
$\Gamma_3$ $4\pi^0$	seen
$\Gamma_4$ $2\pi^+ 2\pi^-$	seen
$\Gamma_5$ $\pi^+ \pi^- 2\pi^0$	seen
$\Gamma_6$ $\rho\rho$	seen
$\Gamma_7$ $2(\pi\pi)_S$ -wave	seen
$\Gamma_8$ $\pi(1300)\pi$	seen
$\Gamma_9$ $a_1(1260)\pi$	seen
$\Gamma_{10}$ $\eta\eta$	seen
$\Gamma_{11}$ $K\bar{K}$	seen
$\Gamma_{12}$ $K\bar{K}n\pi$	not seen
$\Gamma_{13}$ $6\pi$	not seen
$\Gamma_{14}$ $\omega\omega$	not seen
$\Gamma_{15}$ $\gamma\gamma$	seen
$\Gamma_{16}$ $e^+ e^-$	not seen

NODE=M147215;NODE=M147

DESIG=1;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=10;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=4;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=5;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=6;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=14;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=15;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=16;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=17;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=2;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=11;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=18;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=19;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=20;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=12;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=13;OUR EST; $\rightarrow$  UNCHECKED  $\leftarrow$

 **$f_0(1370)$  PARTIAL WIDTHS** **$\Gamma(\gamma\gamma)$** 

See  $\gamma\gamma$  widths under  $f_0(500)$  and MORGAN 90.

$\Gamma_{15}$

NODE=M147217

NODE=M147W11  
NODE=M147W11  
NODE=M147W11

 **$\Gamma(e^+ e^-)$** 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBYEV	88	ND $e^+ e^- \rightarrow \pi^0 \pi^0$

$\Gamma_{16}$

NODE=M147W12  
NODE=M147W12

$f_0(1370) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M147225

 $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{10}\Gamma_{15}/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M147G01  
NODE=M147G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

$121^{+133+169}_{-53-106}$	<sup>1</sup> UEHARA	10A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
----------------------------	---------------------	-----	------	--

<sup>1</sup> Including interference with the  $f'_2(1525)$  (parameters fixed to the values from the 2008 edition of this review, PDG 08) and  $f_2(1270)$ . May also be the  $f_0(1500)$ .

NODE=M147G01;LINKAGE=UE

 $f_0(1370) \text{ BRANCHING RATIOS}$ 

NODE=M147220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M147R3  
NODE=M147R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10	95	OCHS	13	RVUE
$0.26 \pm 0.09$		BUGG	96	RVUE
<0.15		<sup>1</sup> AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
<0.06		GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

<sup>1</sup> Using AMSLER 95B ( $3\pi^0$ ).

NODE=M147R3;LINKAGE=B

 $\Gamma(4\pi)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma = (\Gamma_3+\Gamma_4+\Gamma_5)/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R4  
NODE=M147R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.72	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$
-------	---------	----	---

 $\Gamma(4\pi^0)/\Gamma(4\pi)$  $\Gamma_3/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R12  
NODE=M147R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ABELE	96	CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$
$0.068 \pm 0.005$	<sup>1</sup> GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

<sup>1</sup> Model-dependent evaluation.

NODE=M147R12;LINKAGE=GA

 $\Gamma(2\pi^+2\pi^-)/\Gamma(4\pi)$  $\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3+\Gamma_4+\Gamma_5)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R5  
NODE=M147R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.420 \pm 0.014$	<sup>1</sup> GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$
-------------------	----------------------	----	---

<sup>1</sup> Model-dependent evaluation.

NODE=M147R5;LINKAGE=A

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(4\pi)$  $\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3+\Gamma_4+\Gamma_5)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R6  
NODE=M147R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.512 \pm 0.019$	<sup>1</sup> GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$
-------------------	----------------------	----	---

<sup>1</sup> Model-dependent evaluation.

NODE=M147R6;LINKAGE=A

 $\Gamma(\rho\rho)/\Gamma(4\pi)$  $\Gamma_6/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R17  
NODE=M147R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.26 \pm 0.07$	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi\rho$
-----------------	-------	-----	--

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(\pi\pi)$  $\Gamma_7/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R15  
NODE=M147R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.6 \pm 2.6$	<sup>1</sup> ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^-4\pi^0\rho$
---------------	--------------------	----	---

<sup>1</sup> From the combined data of ABELE 96 and ABELE 96C.

NODE=M147R;LINKAGE=KZ

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$  $\Gamma_7/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R16  
NODE=M147R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.51 \pm 0.09$	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi\rho$
-----------------	-------	-----	--



$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$  $\Gamma_6/\Gamma_7$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R10  
NODE=M147R10

••• We do not use the following data for averages, fits, limits, etc. •••

large	BARBERIS	00C	450 $p\rho \rightarrow \rho_f 4\pi p_S$
$1.6 \pm 0.2$	AMSLER	94	CBAR $\bar{p}\rho \rightarrow \pi^+ \pi^- 3\pi^0$
$\sim 0.65$	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$

OCCUR=2

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$  $\Gamma_8/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R18  
NODE=M147R18

••• We do not use the following data for averages, fits, limits, etc. •••

$0.17 \pm 0.06$	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi\rho$
-----------------	-------	-----	--

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$  $\Gamma_9/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R19  
NODE=M147R19

••• We do not use the following data for averages, fits, limits, etc. •••

$0.06 \pm 0.02$	ABELE	01B	CBAR $0.0 \bar{p}d \rightarrow 5\pi\rho$
-----------------	-------	-----	--

 $\Gamma(\eta\eta)/\Gamma(4\pi)$  $\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R14  
NODE=M147R14

••• We do not use the following data for averages, fits, limits, etc. •••

$(28 \pm 11) \times 10^{-3}$	<sup>1</sup> ANISOVICH	02D	SPEC Combined fit
$(4.7 \pm 2.0) \times 10^{-3}$	BARBERIS	00E	450 $p\rho \rightarrow \rho_f \eta\eta p_S$

<sup>1</sup> From a combined K-matrix analysis of Crystal Barrel ( $0. \rho\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$ ), GAMS ( $\pi\rho \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi\rho \rightarrow K\bar{K}n$ ) data.

NODE=M147R14;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R11  
NODE=M147R11

••• We do not use the following data for averages, fits, limits, etc. •••

$0.35 \pm 0.13$	BUGG	96	RVUE
-----------------	------	----	------

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$  $\Gamma_{11}/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R13  
NODE=M147R13

••• We do not use the following data for averages, fits, limits, etc. •••

$0.08 \pm 0.08$	ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-, \phi K^+ K^-$
$0.91 \pm 0.20$	<sup>1</sup> BARGIOTTI	03	OBLX $\bar{p}\rho$
$0.12 \pm 0.06$	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit
$0.46 \pm 0.15 \pm 0.11$	BARBERIS	99D	OMEG $450 p\rho \rightarrow K^+ K^-, \pi^+ \pi^-$

<sup>1</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0$ , and  $K^\pm K_S^0 \pi^\mp$ .

<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel ( $0. \rho\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$ ), GAMS ( $\pi\rho \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi\rho \rightarrow K\bar{K}n$ ) data.

NODE=M147R;LINKAGE=BG  
NODE=M147R;LINKAGE=CH

 $\Gamma(K\bar{K}n\pi)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R20  
NODE=M147R20

••• We do not use the following data for averages, fits, limits, etc. •••

$< 0.03$	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
----------	---------	----	---

 $\Gamma(6\pi)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R21  
NODE=M147R21

••• We do not use the following data for averages, fits, limits, etc. •••

$< 0.22$	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
----------	---------	----	---

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R22  
NODE=M147R22

••• We do not use the following data for averages, fits, limits, etc. •••

$< 0.13$	GASPERO	93	DBC $0.0 \bar{p}n \rightarrow \text{hadrons}$
----------	---------	----	---

$\phi_0(1370)$  REFERENCES

NODE=M147

SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
OCHS	13	JP G40 043001	W. Ochs		REFID=55367
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52723
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=51721
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>		REFID=53252
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51594
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=51162
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05Q	PRL 87 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45011
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45038
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)	REFID=45094
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist	(HELS)	REFID=44529
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC	REFID=43660
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC	REFID=43657
GASPERO	93	NP A562 407	M. Gaspero	(ROMA1) JPC	REFID=43658
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
		Translated from YAF 48 436.			
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)	REFID=40064
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)	REFID=21123
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=21418
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20383
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SAFL)	REFID=11004
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)	REFID=20349
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)	REFID=44530
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)	REFID=21361

# π<sub>1</sub>(1400)

$$I^G(J^{PC}) = 1^-(1^-+)$$

Coupled channel analyses favor the existence of only one broad 1<sup>-</sup>+ isovector state consistent with π<sub>1</sub>(1600) in the 1400–1600 MeV region. See the review on "Spectroscopy of Light Meson Resonances." See also π<sub>1</sub>(1600).

NODE=M111

NODE=M111

## π<sub>1</sub>(1400) T-MATRIX POLE √s

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M111PP

NODE=M111PP

NODE=M111PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1405 ± 4<sup>+15</sup><sub>-18</sub>) - i (314 ± 14<sup>+18</sup><sub>-69</sub>) OUR ESTIMATE</b>			
(1405 ± 4 <sup>+15</sup> <sub>-18</sub> ) - i (314 ± 14 <sup>+18</sup> <sub>-69</sub> )	ALBRECHT	20	RVUE $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$

## π<sub>1</sub>(1400) MASS

NODE=M111M

NODE=M111M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1354 ± 25 OUR AVERAGE</b>		Error includes scale factor of 1.8. See the ideogram below.			
1257 ± 20 ± 25	23.5k	ADAMS	07B	B852	18 π <sup>-</sup> p → ηπ <sup>0</sup> n
1384 ± 20 ± 35	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1360 ± 25		ABELE	99	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1400 ± 20 ± 20		ABELE	98B	CBAR	0.0 $\bar{p}n \rightarrow \pi^- \pi^0 \eta$
1370 ± 16 <sup>+50</sup> <sub>-30</sub>		<sup>1</sup> THOMPSON	97	MPS	18 π <sup>-</sup> p → ηπ <sup>-</sup> p
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1323.1 ± 4.6		<sup>2</sup> AOYAGI	93	BKEI	π <sup>-</sup> p → ηπ <sup>-</sup> p
1406 ± 20		<sup>3</sup> ALDE	88B	GAM4 0	100 π <sup>-</sup> p → ηπ <sup>0</sup> n

<sup>1</sup> Natural parity exchange, questioned by DZIERBA 03.


<sup>2</sup> Unnatural parity exchange.

<sup>3</sup> Seen in the P<sub>0</sub>-wave intensity of the ηπ<sup>0</sup> system, unnatural parity exchange.

NODE=M111M;LINKAGE=B

NODE=M111M;LINKAGE=C

NODE=M111M;LINKAGE=A



## π<sub>1</sub>(1400) WIDTH

NODE=M111W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>330 ± 35</b>	<b>OUR AVERAGE</b>				
354 ± 64 ± 58	23.5k	ADAMS	07B	B852	18 $\pi^- p \rightarrow \eta \pi^0 n$
378 ± 50 ± 50	90k	SALVINI	04	OBLX	$\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
220 ± 90		ABELE	99	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$
310 ± 50 + 50 - 30		ABELE	98B	CBAR	0.0 $\bar{p} n \rightarrow \pi^- \pi^0 \eta$
385 ± 40 + 65 - 105		<sup>1</sup> THOMPSON	97	MPS	18 $\pi^- p \rightarrow \eta \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
143.2 ± 12.5		<sup>2</sup> AOYAGI	93	BKEI	$\pi^- p \rightarrow \eta \pi^- p$
180 ± 20		<sup>3</sup> ALDE	88B	GAM4 0	100 $\pi^- p \rightarrow \eta \pi^0 n$
<sup>1</sup> Resolution is not unfolded, natural parity exchange, questioned by DZIERBA 03.					
<sup>2</sup> Unnatural parity exchange.					
<sup>3</sup> Seen in the $P_0$ -wave intensity of the $\eta \pi^0$ system, unnatural parity exchange.					

NODE=M111W

NODE=M111W;LINKAGE=QQ  
 NODE=M111W;LINKAGE=C  
 NODE=M111W;LINKAGE=A

### $\pi_1(1400)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta \pi^0$	seen
$\Gamma_2$ $\eta \pi^-$	seen
$\Gamma_3$ $\eta' \pi$	
$\Gamma_4$ $\rho(770) \pi$	not seen

NODE=M111215;NODE=M111

DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=4;OUR EST;→ UNCHECKED ←  
 DESIG=3  
 DESIG=5

### $\pi_1(1400)$ BRANCHING RATIOS

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_1/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen	PROKOSHKIN	95B	GAM4	100 $\pi^- p \rightarrow \eta \pi^0 n$	
not seen	<sup>1</sup> BUGG	94	RVUE	$\bar{p} p \rightarrow \eta 2\pi^0$	
not seen	<sup>2</sup> APEL	81	NICE 0	40 $\pi^- p \rightarrow \eta \pi^0 n$	

NODE=M111220

NODE=M111R1  
 NODE=M111R1

<sup>1</sup> Using Crystal Barrel data.<sup>2</sup> A general fit allowing  $S$ ,  $D$ , and  $P$  waves (including  $m=0$ ) is not done because of limited statistics.

NODE=M111R1;LINKAGE=C  
 NODE=M111R1;LINKAGE=B

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
possibly seen	BELADIDZE	93	VES 37 $\pi^- N \rightarrow \eta \pi^- N$	

NODE=M111R4  
 NODE=M111R4

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma_1$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.80	95	BOUTEMEUR	90	GAM4 100 $\pi^- p \rightarrow 4\gamma n$	

NODE=M111R3  
 NODE=M111R3

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma$
not seen	AGHASYAN	18B	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	

NODE=M111R00  
 NODE=M111R00

### $\pi_1(1400)$ REFERENCES

NODE=M111

ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ADAMS	07B	PL B657 27	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=52048
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)	REFID=53226
DZIERBA	03	PR D67 094015	A.R. Dzierba <i>et al.</i>		REFID=49412
ABELE	99	PL B446 349	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46602
ABELE	98B	PL B423 175	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45864
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=45584
PROKOSHKIN	95B	PAN 58 606	Y.D. Prokoshkin, S.A. Sadovsky	(SERP)	REFID=44619
		Translated from YAF 58 662.			
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
BELADIDZE	93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43598
BOUTEMEUR	90	Hadron 89 Conf. p 119	M. Boutemeur, M. Poulet	(SERP, BELG, LANL+)	REFID=41751
ALDE	88B	PL B205 397	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP) IGJPC	REFID=40558
APEL	81	NP B193 269	W.D. Apel <i>et al.</i>	(SERP, CERN)	REFID=22913

**$\eta(1405)$**

$I^G(J^{PC}) = 0^+(0^{-+})$

See also the  $\eta(1475)$ .

NODE=M027

NODE=M027

NODE=M027205


NODE=M027MX

**$\eta(1405)$  MASS**

VALUE (MeV)

DOCUMENT ID

**1408.8±2.0 OUR AVERAGE** Includes data from the 2 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.



**$\eta\pi\pi$  MODE**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

NODE=M027M1  
NODE=M027M1

The data in this block is included in the average printed for a previous datablock.

**1405.8± 2.6 OUR AVERAGE** Error includes scale factor of 2.3. See the ideogram below.

1409.0± 1.7	743	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$	
1407.0± 3.5	198	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$	OCCUR=2
1399.8± 2.2 <sup>+2.8</sup> <sub>-0.1</sub>		1 ABLIKIM	11J	BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$	
1392 ±14	900 ± 375	AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$	
1394 ± 8	6.6 ± 2.0k	AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$	OCCUR=2
1404 ± 6	9082	MANAK	00A	MPS	$18 \pi^-\rho \rightarrow \eta\pi^+\pi^-n$	
1424 ± 6	2200	ALDE	97B	GAM4	$100 \pi^-\rho \rightarrow \eta\pi^0\pi^0n$	
1400 ± 6		2 BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
1388 ± 4		FUKUI	91C	SPEC	$8.95 \pi^-\rho \rightarrow \eta\pi^+\pi^-n$	
1398 ± 6	261	3 AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
1420 ± 5		ANDO	86	SPEC	$8 \pi^-\rho \rightarrow \eta\pi^+\pi^-n$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
1404.0±11.0	195	ABLIKIM	19B	ABES3	$e^+e^- \rightarrow \psi(2S)$	
1385 ± 7		BAI	99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
1409 ± 3		4 AMSLER	95F	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$	


<sup>1</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$ .

<sup>2</sup> From fit to the  $a_0(980)\pi 0^{-+}$  partial wave.

<sup>3</sup> Best fit with a single Breit Wigner.

<sup>4</sup> Superseded by AMSLER 04B.

NODE=M027M1;LINKAGE=BL  
NODE=M027M1;LINKAGE=J1  
NODE=M027M1;LINKAGE=A1  
NODE=M027M1;LINKAGE=A



### $K\bar{K}\pi$ MODE ( $a_0(980)\pi$ or direct $K\bar{K}\pi$ )

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

**1413.9± 1.7 OUR AVERAGE** Error includes scale factor of 1.1.

1413 ±14	3651	<sup>1</sup> NICHITIU	02 OBLX	$0 \bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
1416 ± 4 ±2	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$	
1405 ± 5		<sup>2</sup> CICALO	99 OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$	
1407 ± 5		<sup>2</sup> BERTIN	97 OBLX	$0 \bar{p}p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$	
1416 ± 2		<sup>2</sup> BERTIN	95 OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$	
1416 ± 8 $\begin{smallmatrix} +7 \\ -5 \end{smallmatrix}$	700	<sup>3</sup> BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=2
1413 ± 5		<sup>3</sup> RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$	OCCUR=3

••• We do not use the following data for averages, fits, limits, etc. •••

1459 ± 5 <sup>4</sup> AUGUSTIN 92 DM2  $J/\psi \rightarrow \gamma K\bar{K}\pi$

<sup>1</sup> Decaying dominantly directly to  $K^+ K^- \pi^0$ .

<sup>2</sup> Decaying into  $(K\bar{K})_S \pi$ ,  $(K\pi)_S \bar{K}$ , and  $a_0(980)\pi$ .

<sup>3</sup> From fit to the  $a_0(980)\pi 0^-$  partial wave. Cannot rule out a  $a_0(980)\pi 1^+$  partial wave.

<sup>4</sup> Excluded from averaging because averaging would be meaningless.

NODE=M027M4  
NODE=M027M4

OCCUR=2

OCCUR=3

NODE=M027M;LINKAGE=NC  
NODE=M027M4;LINKAGE=FX  
NODE=M027M4;LINKAGE=C2

NODE=M027M4;LINKAGE=AA

### $\pi\pi\gamma$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

**1403±17 OUR AVERAGE** Error includes scale factor of 1.8.

1390±12	235 ± 91	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$	
1424±10±11	547	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
1401±18		<sup>1,2</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+ \pi^- \gamma\gamma$	OCCUR=4
1432± 8		<sup>2</sup> COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+ \pi^- 2\gamma$	

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Best fit with a single Breit Wigner.

<sup>2</sup> This peak in the  $\gamma\rho$  channel may not be related to the  $\eta(1405)$ .

NODE=M027M2;LINKAGE=E  
NODE=M027M2;LINKAGE=X

### $4\pi$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

1420±20		BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$	
1489±12	3270	<sup>1</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$	

<sup>1</sup> Estimated by us from various fits.

NODE=M027M3  
NODE=M027M3

NODE=M027M3;LINKAGE=E

**$K\bar{K}\pi$  MODE (unresolved)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1452.7 ± 3.3	191	1,2 ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
1437.6 ± 3.2	249 ± 35	1,2 ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
1445.9 ± 5.7	62 ± 18	1,2 ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
1442 ± 10	410	1 BAI	98C BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1445 ± 8	693	1 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1433 ± 8	296	1 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1413 ± 8	500	1 DUCH	89 ASTE	$\bar{p}p \rightarrow \pi^+ \pi^- K^\pm \pi^\mp K^0$
1453 ± 7	170	1 RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1419 ± 1	8800	1 BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ K^0 \pi^- n$
1424 ± 3	620	1 REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K\bar{K}\pi X$
1421 ± 2		1 CHUNG	85 SPEC	$8 \pi^- p \rightarrow K\bar{K}\pi n$
1440 +20 -15	174	1 EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1440 +10 -15		1 SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1425 ± 7	800	1,3 BAILLON	67 HBC	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

NODE=M027M6  
NODE=M027M6

OCCUR=2

OCCUR=2

OCCUR=2

<sup>1</sup> These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to  $\eta(1475)$ .

<sup>2</sup> Systematic uncertainty not evaluated.

<sup>3</sup> From best fit of  $0^-+$  partial wave, 50%  $K^*(892)K$ , 50%  $a_0(980)\pi$ .

NODE=M027M;LINKAGE=NP

NODE=M027M;LINKAGE=NS

NODE=M027M6;LINKAGE=H

**$\eta(1405)$  WIDTH**


NODE=M027210

VALUE (MeV)	DOCUMENT ID
-------------	-------------

NODE=M027WX

**50.1 ± 2.6 OUR AVERAGE** Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.7. See the ideogram below.

WEIGHTED AVERAGE  
50.1 ± 2.6 (Error scaled by 1.7)



**$\eta\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M027W1  
NODE=M027W1

The data in this block is included in the average printed for a previous datablock.

**52.6 ± 3.2 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

48.3 ± 5.2	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^0)$
55.0 ± 11.0	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^0 \pi^0 \pi^0)$
52.8 ± 7.6 +0.1 -7.6		1 ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
55 ± 11	900	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^+ \pi^- \eta$

OCCUR=2

55 ±12	6.6k	AMSLER	04B	CBAR	0	$\bar{p}p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \gamma$
80 ±21	9.0k	MANAK	00A	MPS	18	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$
85 ±18	2.2k	ALDE	97B	GAM4	100	$\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
47 ±13		<sup>2</sup> BOLTON	92B	MRK3		$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
59 ± 4		FUKUI	91C	SPEC	8.95	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$
53 ±11		<sup>3</sup> AUGUSTIN	90	DM2		$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
31 ± 7		ANDO	86	SPEC	8	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$


OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

79.0±16.0	195	ABLIKIM	19BA	BES3		$e^+ e^- \rightarrow \psi(2S)$
86 ±10		<sup>4</sup> AMSLER	95F	CBAR	0	$\bar{p}p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta$

NODE=M027W1;LINKAGE=BL  
 NODE=M027W1;LINKAGE=A1  
 NODE=M027W1;LINKAGE=D1  
 NODE=M027W1;LINKAGE=A

- <sup>1</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$ .
- <sup>2</sup> From fit to the  $a_0(980)\pi 0^-+$  partial wave.
- <sup>3</sup> From  $\eta \pi^+ \pi^-$  mass distribution - mainly  $a_0(980)\pi$  - no spin-parity determination available.
- <sup>4</sup> Superseded by AMSLER 04B.



**$K\bar{K}\pi$  MODE ( $a_0(980)\pi$  or direct  $K\bar{K}\pi$ )**

NODE=M027W4  
 NODE=M027W4

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  
 The data in this block is included in the average printed for a previous datablock.

**48 ± 4 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

51 ± 6	3651	<sup>1</sup> NICHITIU	02	OBLX	0	$\bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
42 ±10 ± 9	20k	ADAMS	01B	B852	18	GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
50 ± 4		CICALO	99	OBLX	0	$\bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
48 ± 5		<sup>2</sup> BERTIN	97	OBLX	0.0	$\bar{p}p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
50 ± 4		<sup>2</sup> BERTIN	95	OBLX	0	$\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
75 ± 9		AUGUSTIN	92	DM2		$J/\psi \rightarrow \gamma K\bar{K}\pi$
91 +67 +15 -31 -38		<sup>3</sup> BAI	90C	MRK3		$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
19 ± 7		<sup>3</sup> RATH	89	MPS	21.4	$\pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$


OCCUR=2

OCCUR=3

- <sup>1</sup> Decaying dominantly directly to  $K^+ K^- \pi^0$ .
- <sup>2</sup> Decaying into  $(K\bar{K})_S \pi$ ,  $(K\pi)_S \bar{K}$ , and  $a_0(980)\pi$ .
- <sup>3</sup> From fit to the  $a_0(980)\pi 0^-+$  partial wave, but  $a_0(980)\pi 1^++$  cannot be excluded.

NODE=M027W;LINKAGE=NC  
 NODE=M027W4;LINKAGE=FX  
 NODE=M027W4;LINKAGE=C





**ππγ MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>89 ±17</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.7.		
64 ±18	235 ± 91	AMSLER	04B CBAR	0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
101.0± 8.8±8.8	547	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
174 ±44		AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$
90 ±26		<sup>1</sup> COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$

<sup>1</sup>This peak in the  $\gamma\rho$  channel may not be related to the  $\eta(1405)$ .

NODE=M027W2  
NODE=M027W2

OCCUR=2

NODE=M027W2;LINKAGE=X

**4π MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
160±30		BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
144±13	3270	<sup>1</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup> Estimated by us from various fits.

NODE=M027W3  
NODE=M027W3

NODE=M027W3;LINKAGE=F2

**K K-bar π MODE (unresolved)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
45.9± 8.2	191	<sup>1,2</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
48.9± 9.0	249 ± 35	<sup>1,2</sup> ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+\pi^- + c.c.$
34.2±18.5	62 ± 18	<sup>1,2</sup> ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^-\pi^0$
93 ±14	296	<sup>1</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^-\pi^0$
105 ±10	693	<sup>1</sup> AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
62 ±16	500	<sup>1</sup> DUCH	89 ASTE	$\bar{p}p \rightarrow K \bar{K} \pi \pi$
100 ±11	170	<sup>1</sup> RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
66 ± 2	8800	<sup>1</sup> BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
60 ±10	620	<sup>1</sup> REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K K \pi X$
60 ±10		<sup>1</sup> CHUNG	85 SPEC	$8 \pi^- p \rightarrow K \bar{K} \pi n$
55 <sup>+20</sup> <sub>-30</sub>	174	<sup>1</sup> EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^-\pi^0$
50 <sup>+30</sup> <sub>-20</sub>		<sup>1</sup> SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
80 ±10	800	<sup>1,3</sup> BAILLON	67 HBC	$0.0 \bar{p}p \rightarrow K \bar{K} \pi \pi$

<sup>1</sup> These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to  $\eta(1475)$ .

<sup>2</sup> Systematic uncertainty not evaluated.

<sup>3</sup> From best fit to  $0^-+$  partial wave, 50%  $K^*(892)K$ , 50%  $a_0(980)\pi$ .

NODE=M027W6  
NODE=M027W6

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M027W;LINKAGE=NP

NODE=M027W;LINKAGE=NS

NODE=M027W6;LINKAGE=H1

**$\eta(1405)$  DECAY MODES**

NODE=M027215;NODE=M027

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $K\bar{K}\pi$	seen	
$\Gamma_2$ $\eta\pi\pi$	seen	
$\Gamma_3$ $a_0(980)\pi$	seen	
$\Gamma_4$ $\eta(\pi\pi)$ S-wave	seen	
$\Gamma_5$ $f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$	not seen	
$\Gamma_6$ $f_0(980)\eta$	seen	
$\Gamma_7$ $4\pi$	seen	
$\Gamma_8$ $\rho\rho$	<58 %	99.85%
$\Gamma_9$ $\gamma\gamma$		
$\Gamma_{10}$ $\rho^0\gamma$	seen	
$\Gamma_{11}$ $\phi\gamma$		
$\Gamma_{12}$ $K^*(892)K$	seen	

DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=9;OUR EST;→ UNCHECKED ←  
DESIG=15  
DESIG=10;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←  
DESIG=12  
DESIG=7  
DESIG=8;OUR EST;→ UNCHECKED ←  
DESIG=13  
DESIG=11;OUR EST;→ UNCHECKED ←

 **$\eta(1405)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$** 

NODE=M027220

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_9/\Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027G3  
NODE=M027G3

••• We do not use the following data for averages, fits, limits, etc. •••

<0.035 90 1,2 AHOHE 05 CLE2 10.6  $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$

<sup>1</sup> Using  $\eta(1405)$  mass and width 1410 MeV and 51 MeV, respectively.

<sup>2</sup> Assuming three-body phase-space decay to  $K_S^0K^\pm\pi^\mp$ .

NODE=M027G3;LINKAGE=AH  
NODE=M027G3;LINKAGE=B3

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_9/\Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027G5  
NODE=M027G5

<0.095 95 ACCIARRI 01G L3 183–202  $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

$\Gamma(\rho^0\gamma) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{10}\Gamma_9/\Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027G8  
NODE=M027G8

••• We do not use the following data for averages, fits, limits, etc. •••

<1.5 95 ALTHOFF 84E TASS  $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$

 **$\eta(1405)$  BRANCHING RATIOS**

NODE=M027225

$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$					$\Gamma_2/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M027R3  
NODE=M027R3

1.09±0.48 <sup>1</sup> AMSLER 04B CBAR 0  $\bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.5 90 EDWARDS 83B CBAL  $J/\psi \rightarrow \eta\pi\pi\gamma$

<1.1 90 SCHARRE 80 MRK2  $J/\psi \rightarrow \eta\pi\pi\gamma$

<1.5 95 FOSTER 68B HBC 0.0  $\bar{p}p$

<sup>1</sup> Using the data of BAILLON 67 on  $\bar{p}p \rightarrow K\bar{K}\pi$ .

NODE=M027R3;LINKAGE=AM

$\Gamma(\rho^0\gamma)/\Gamma(\eta\pi\pi)$				$\Gamma_{10}/\Gamma_2$
VALUE	DOCUMENT ID	TECN	COMMENT	

NODE=M027R12  
NODE=M027R12

0.111±0.064 AMSLER 04B CBAR 0  $\bar{p}p$

$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$					$\Gamma_3/\Gamma_1$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M027R4  
NODE=M027R4

••• We do not use the following data for averages, fits, limits, etc. •••

~0.15 <sup>1</sup> BERTIN 95 OBLX 0  $\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

~0.8 500 <sup>1</sup> DUCH 89 ASTE  $\bar{p}p \rightarrow \pi^+\pi^-K^\pm\pi^\mp K^0$

~0.75 <sup>1</sup> REEVES 86 SPEC 6.6  $p\bar{p} \rightarrow KK\pi X$

<sup>1</sup> Assuming that the  $a_0(980)$  decays only into  $K\bar{K}$ .

NODE=M027R4;LINKAGE=C

$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$					$\Gamma_3/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M027R2  
NODE=M027R2

••• We do not use the following data for averages, fits, limits, etc. •••

0.29±0.10 ABELE 98E CBAR 0  $p\bar{p} \rightarrow \eta\pi^0\pi^0\pi^0$

0.19±0.04 2200 <sup>1</sup> ALDE 97B GAM4 100  $\pi^-p \rightarrow \eta\pi^0\pi^0n$

0.56±0.04±0.03 <sup>1</sup> AMSLER 95F CBAR 0  $\bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$

<sup>1</sup> Assuming that the  $a_0(980)$  decays only into  $\eta\pi$ .

NODE=M027R2;LINKAGE=A

$\Gamma(a_0(980)\pi)/\Gamma(\eta(\pi\pi)S\text{-wave})$  $\Gamma_3/\Gamma_4$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M027R9  
NODE=M027R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.91±0.12		ANISOVICH	01	SPEC	0.0 $\bar{p}p \rightarrow \eta\pi^+\pi^-\pi^+\pi^-$
0.15±0.04	9082	<sup>1</sup> MANAK	00A	MPS	18 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
0.70±0.12±0.20		<sup>2</sup> BAI	99	BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

<sup>1</sup> Statistical error only.

<sup>2</sup> Assuming that the  $a_0(980)$  decays only into  $\eta\pi$ .

NODE=M027R;LINKAGE=K3  
NODE=M027R9;LINKAGE=BK

 $\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$  $\Gamma_{10}/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M027R7  
NODE=M027R7

**0.0152±0.0038** <sup>1</sup> COFFMAN 90 MRK3  $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

<sup>1</sup> Using  $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)=4.2 \times 10^{-3}$  and  $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\gamma\rho^0)=6.4 \times 10^{-5}$ .

NODE=M027R7;LINKAGE=D

 $\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$  $\Gamma_9/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M027R02  
NODE=M027R02

**<1.78 × 10<sup>-3</sup>** 90 <sup>1</sup> ABLIKIM 180 BES3  $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma$

<sup>1</sup> Using results from BAI 00D.

NODE=M027R02;LINKAGE=A

 $\Gamma(\eta(\pi\pi)S\text{-wave})/\Gamma(\eta\pi\pi)$  $\Gamma_4/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M027R8  
NODE=M027R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.81±0.04	2200	ALDE	97B	GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$
-----------	------	------	-----	------	--

 $\Gamma(f_0(980)\eta)/\Gamma(\eta\pi\pi)$  $\Gamma_6/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M027R10  
NODE=M027R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32±0.07 <sup>1</sup> ANISOVICH 00 SPEC 0.9–1.2  $\bar{p}p \rightarrow \eta 3\pi^0$

<sup>1</sup> Using preliminary Crystal Barrel data.

NODE=M027R10;LINKAGE=D

 $\Gamma(f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M027R00  
NODE=M027R00

**not seen** <sup>1</sup> ABLIKIM 17AJ BES3  $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

<sup>1</sup> ABLIKIM 17AJ reports  $B(\psi(2S) \rightarrow \gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0) < 5.0 \times 10^{-7}$ .

NODE=M027R00;LINKAGE=A

 $\Gamma(\rho\rho)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M027R13  
NODE=M027R13

**<0.58** 99.85 <sup>1,2</sup> AMSLER 04B CBAR 0  $\bar{p}p$

<sup>1</sup> Assuming that the  $\eta(1405)$  decays are saturated by the  $\pi\pi\eta$ ,  $K\bar{K}\pi$  and  $\rho\rho$  modes.

<sup>2</sup> Using the data of BAILLON 67 on  $\bar{p}p \rightarrow K\bar{K}\pi$ .

NODE=M027R13;LINKAGE=AM  
NODE=M027R13;LINKAGE=AS

 $\Gamma(K^*(892)K)/\Gamma(a_0(980)\pi)$  $\Gamma_{12}/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M027R11  
NODE=M027R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.084±0.024	<sup>1</sup> ADAMS	01B	B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
-------------	--------------------	-----	------	--

<sup>1</sup> Statistical error only.

NODE=M027R11;LINKAGE=K3

 $\Gamma(\phi\gamma)/\Gamma(\rho^0\gamma)$  $\Gamma_{11}/\Gamma_{10}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M027R14  
NODE=M027R14

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.09±0.03	<sup>1</sup> ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
-----------	----------------------	-----	------	---

0.13±0.04	<sup>2</sup> ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
-----------	----------------------	-----	------	---

<0.77	95	<sup>3</sup> BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$
-------	----	------------------	-----	---

OCCUR=2

<sup>1</sup> Constructive interference between  $X(1835)$  and  $\eta(1405)/\eta(1475)$  decays to  $\gamma\phi$  is assumed. Also see  $\eta(1475)$ . ABLIKIM 18I reports the inverse as  $11.10 \pm 3.5$ .

NODE=M027R14;LINKAGE=A

<sup>2</sup> Destructive interference between  $X(1835)$  and  $\eta(1405)/\eta(1475)$  decays to  $\gamma\phi$  is assumed. Also see  $\eta(1475)$ . ABLIKIM 18I reports the inverse as  $7.53 \pm 2.49$ .

NODE=M027R14;LINKAGE=B

<sup>3</sup> Calculated by us from  $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \phi\gamma\gamma) < 0.82 \times 10^{-4}$  and  $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \rho^0\gamma\gamma) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$ .

NODE=M027R14;LINKAGE=BA

$\eta(1405)$  REFERENCES

NODE=M027

ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60024
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58322
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55386
ABLIKIM	12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53931
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)	REFID=50764
AMSLER	04B	EPJ C33 23	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51079
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
ANISOVICH	01	NP A690 567	A.V. Anisovich <i>et al.</i>		REFID=48308
ANISOVICH	00	PL B472 168	A.V. Anisovich <i>et al.</i>		REFID=47429
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)	REFID=47394
ABELE	98E	NP B514 45	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46314
BAI	98C	PL B440 217	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46337
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
		Translated from YAF 60 458.			
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
AMSLER	95F	PL B358 389	C. AMSler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44613
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=44614
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
DUCH	89	ZPHY C45 223	K.D. Duch <i>et al.</i>	(ASTERIX Collab.) JP	REFID=41016
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)	REFID=40924
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) JJP	REFID=20891
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP	REFID=20936
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP	REFID=20934
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=20305
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
	Also	PRL 50 219	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21315
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
FOSTER	68B	NP B8 174	M. Foster <i>et al.</i>	(CERN, CDEF)	REFID=21179
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)	REFID=20407

**$h_1(1415)$** 

$$I^G(J^{PC}) = 0^-(1^{+-})$$

was  $h_1(1380)$ 

NODE=M109

 **$h_1(1415)$  MASS**

NODE=M109M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M109M

**1409<sup>+9</sup><sub>-8</sub> OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.

NEW

[1416 ± 8 MeV OUR 2022 AVERAGE Scale factor = 1.5]


1384 ± 6 <sup>+9</sup> <sub>-0</sub>		1 ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
1423 ± 2.1 ± 7.3	2.2k	2 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$
1412 ± 4 ± 8		2 ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1,2} \rightarrow \gamma \phi(h_1 \rightarrow K^* \bar{K})$
1440 ± 60		ABELE	97H CBAR	$\bar{p} p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$
1380 ± 20		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

NODE=M109M;LINKAGE=B

<sup>1</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta' \eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma \eta'$  in the decay  $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>2</sup> Final states  $K^+ K^- \pi^0$  and  $K_S^0 K^\pm \pi^\mp$ .

NODE=M109M;LINKAGE=A

 **$h_1(1415)$  WIDTH**

NODE=M109W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M109W

**78 ± 11 OUR AVERAGE**

NEW

[90 ± 15 MeV OUR 2022 AVERAGE]

66 ± 10 <sup>+12</sup> <sub>-10</sub>		1 ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
90.3 ± 9.8 ± 17.5	2.2k	2 ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$
84 ± 12 ± 40		2 ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1,2} \rightarrow \gamma \phi(h_1 \rightarrow K^* \bar{K})$
170 ± 80		ABELE	97H CBAR	$\bar{p} p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$
80 ± 30		ASTON	88C LASS	$11 K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

NODE=M109W;LINKAGE=B

<sup>1</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta' \eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma \eta'$  in the decay  $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>2</sup> Final states  $K^+ K^- \pi^0$  and  $K_S^0 K^\pm \pi^\mp$ .

NODE=M109W;LINKAGE=A

$h_1(1415)$  DECAY MODES

NODE=M109215;NODE=M109

Mode	
$\Gamma_1$	$K\bar{K}^*(892)+c.c.$

DESIG=1

 $h_1(1415)$  REFERENCES

ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

NODE=M109

REFID=61637  
REFID=59456  
REFID=56778  
REFID=45765  
REFID=40282 $f_1(1420)$ 

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M006

NODE=M006

 $f_1(1420)$  MASS

NODE=M006M2

NODE=M006M2

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1426.3 ± 0.9 OUR AVERAGE</b>		Error includes scale factor of 1.1.		
1434 ± 5 ± 5	133	<sup>1</sup> ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
1426 ± 6	711	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0K^\pm\pi^\mp + X$
1420 ± 14	3651	NICHITIU	02 OBLX	0 $\bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$
1428 ± 4 ± 2	20k	ADAMS	01B B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
1426 ± 1		BARBERIS	97C OMEG	450 $pp \rightarrow ppK_S^0K^\pm\pi^\mp$
1425 ± 8		BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
1430 ± 4		<sup>2</sup> ARMSTRONG	92E OMEG	85,300 $\pi^+p, pp \rightarrow \pi^+p, pp(K\bar{K}\pi)$
1462 ± 20		<sup>3</sup> AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
1443 $\begin{smallmatrix} +7 \\ -6 \end{smallmatrix} \begin{smallmatrix} +3 \\ -2 \end{smallmatrix}$	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0K^\pm\pi^\mp$
1425 ± 10	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$
1442 ± 5 $\begin{smallmatrix} +10 \\ -17 \end{smallmatrix}$	111	BECKER	87 MRK3	$e^+e^-, \omega K\bar{K}\pi$
1423 ± 4		GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^-K\bar{K}\pi$
1417 ± 13	13	AIHARA	86C TPC	$e^+e^- \rightarrow e^+e^-K\bar{K}\pi$
1422 ± 3		CHAUVAT	84 SPEC	ISR 31.5 $pp$
1440 ± 10		<sup>4</sup> BROMBERG	80 SPEC	100 $\pi^-p \rightarrow K\bar{K}\pi X$
1426 ± 6	221	DIONISI	80 HBC	4 $\pi^-p \rightarrow K\bar{K}\pi n$
1420 ± 20		DAHL	67 HBC	1.6-4.2 $\pi^-p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1430.8 ± 0.9		<sup>5</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}}(K_S^0K^+\pi^-)p_{\text{fast}}$
1433.4 ± 0.8		<sup>5</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}}(K_S^0K^-\pi^+)p_{\text{fast}}$
1435 ± 9		PROKOSHKIN	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$
1429 ± 3	389	ARMSTRONG	89 OMEG	300 $pp \rightarrow K\bar{K}\pi pp$
1425 ± 2	1520	ARMSTRONG	84 OMEG	85 $\pi^+p, pp \rightarrow (\pi^+, p)(K\bar{K}\pi)p$
~ 1420		BITYUKOV	84 SPEC	32 $K^-p \rightarrow K^+K^-\pi^0\gamma$

OCCUR=2

<sup>1</sup> From a fit with a width fixed at 55 MeV.<sup>2</sup> This result supersedes ARMSTRONG 84, ARMSTRONG 89.<sup>3</sup> From fit to the  $K^*(892)K$   $1^{++}$  partial wave.<sup>4</sup> Mass error increased to account for  $a_0(980)$  mass cut uncertainties.<sup>5</sup> No systematic error given.NODE=M006M2;LINKAGE=CH  
NODE=M006M2;LINKAGE=C  
NODE=M006M2;LINKAGE=B  
NODE=M006M2;LINKAGE=A  
NODE=M006M2;LINKAGE=N1 $f_1(1420)$  WIDTH

NODE=M006W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>54.5± 2.6 OUR AVERAGE</b>				
51 ±14	711	ABDALLAH	03H DLPH	91.2 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
61 ± 8	3651	NICHITIU	02 OBLX	0 $\bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
38 ± 9 ±6	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
58 ± 4		BARBERIS	97C OMEG	450 $pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
45 ±10		BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
58 ±10		<sup>6</sup> ARMSTRONG	92E OMEG	85,300 $\pi^+ p, pp \rightarrow \pi^+ p, pp (K \bar{K} \pi)$
129 ±41		<sup>7</sup> AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
68 $\begin{smallmatrix} +29 \\ -18 \end{smallmatrix}$ $\begin{smallmatrix} +8 \\ -9 \end{smallmatrix}$	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
42 ±22	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
40 $\begin{smallmatrix} +17 \\ -13 \end{smallmatrix}$ ±5	111	BECKER	87 MRK3	$e^+e^- \rightarrow \omega K \bar{K} \pi$
35 $\begin{smallmatrix} +47 \\ -20 \end{smallmatrix}$	13	AIHARA	86C TPC	$e^+e^- \rightarrow e^+e^- K \bar{K} \pi$
47 ±10		CHAUVAT	84 SPEC	ISR 31.5 $pp$
62 ±14		BROMBERG	80 SPEC	100 $\pi^- p \rightarrow K \bar{K} \pi X$
40 ±15	221	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
60 ±20		DAHL	67 HBC	1.6-4.2 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
68.7± 2.9		<sup>8</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$
58.8± 3.3		<sup>8</sup> SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
90 ±25		PROKOSHKIN	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
58 ± 8	389	ARMSTRONG	89 OMEG	300 $pp \rightarrow K \bar{K} \pi pp$
62 ± 5	1520	ARMSTRONG	84 OMEG	85 $\pi^+ p, pp \rightarrow (\pi^+, p)(K \bar{K} \pi)p$
~ 50		BITYUKOV	84 SPEC	32 $K^- p \rightarrow K^+ K^- \pi^0 \gamma$

<sup>6</sup>This result supersedes ARMSTRONG 84, ARMSTRONG 89.

<sup>7</sup>From fit to the  $K^*(892)K 1^{++}$  partial wave.

<sup>8</sup>No systematic error given.

NODE=M006W

OCCUR=2

NODE=M006W;LINKAGE=C  
 NODE=M006W;LINKAGE=B  
 NODE=M006W;LINKAGE=N1

### $f_1(1420)$ DECAY MODES

NODE=M006215;NODE=M006

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \bar{K} \pi$	seen
$\Gamma_2$ $K \bar{K}^*(892) + \text{c.c.}$	seen
$\Gamma_3$ $\eta \pi \pi$	possibly seen
$\Gamma_4$ $a_0(980) \pi$	
$\Gamma_5$ $\pi \pi \rho$	
$\Gamma_6$ $4\pi$	
$\Gamma_7$ $\rho^0 \gamma$	
$\Gamma_8$ $\phi \gamma$	seen

DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=5;OUR EST;→ UNCHECKED ←  
 DESIG=4  
 DESIG=3  
 DESIG=6  
 DESIG=8  
 DESIG=9;OUR EST;→ UNCHECKED ←

### $f_1(1420) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M006220

$\Gamma(K \bar{K} \pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$

NODE=M006G2  
 NODE=M006G2

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.9±0.4 OUR AVERAGE</b>					
3.2±0.6±0.7		133	<sup>9,10</sup> ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
3.0±0.9±0.7			<sup>11,12</sup> BEHREND	89 CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K \pi$
2.3 $\begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix}$ ±0.8			HILL	89 JADE	$e^+e^- \rightarrow e^+e^- K^\pm K_S^0 \pi^\mp$
1.3±0.5±0.3			AIHARA	88B TPC	$e^+e^- \rightarrow e^+e^- K^\pm K_S^0 \pi^\mp$
1.6±0.7±0.3			<sup>11,13</sup> GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^- K \bar{K} \pi$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<8.0 95 JENNI 83 MRK2  $e^+e^- \rightarrow e^+e^-K\bar{K}\pi$

<sup>9</sup> From a fit with a width fixed at 55 MeV.

<sup>10</sup> The form factor parameter from the fit is  $926 \pm 78$  MeV.

<sup>11</sup> Assume a  $\rho$ -pole form factor.

<sup>12</sup> A  $\phi$ -pole form factor gives considerably smaller widths.

<sup>13</sup> Published value divided by 2.

NODE=M006G2;LINKAGE=CH  
 NODE=M006G2;LINKAGE=CR  
 NODE=M006G2;LINKAGE=A  
 NODE=M006G2;LINKAGE=D  
 NODE=M006G2;LINKAGE=B

## $f_1(1420)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(K\bar{K}\pi)$   $\Gamma_2/\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.76±0.06	BROMBERG	80	SPEC 100 $\pi^- p \rightarrow K\bar{K}\pi X$
0.86±0.12	DIONISI	80	HBC 4 $\pi^- p \rightarrow K\bar{K}\pi n$

NODE=M006225

NODE=M006R1  
 NODE=M006R1

$\Gamma(\pi\pi\rho)/\Gamma(K\bar{K}\pi)$   $\Gamma_5/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.3	95	CORDEN	78	OMEG 12-15 $\pi^- p$
<2.0		DAHL	67	HBC 1.6-4.2 $\pi^- p$

NODE=M006R2  
 NODE=M006R2

$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$   $\Gamma_3/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.1 95 ARMSTRONG 91B OMEG 300  $p\bar{p} \rightarrow p\rho\eta\pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

1.35±0.75		KOPKE	89	MRK3 $J/\psi \rightarrow \omega\eta\pi\pi(K\bar{K}\pi)$
<0.6	90	GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
<0.5	95	CORDEN	78	OMEG 12-15 $\pi^- p$
1.5 ±0.8		DEFOIX	72	HBC 0.7 $\bar{p}p$

NODE=M006R3  
 NODE=M006R3

$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$   $\Gamma_4/\Gamma_3$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

>0.1	90	PROKOSHKIN	97B	GAM4 100 $\pi^- p \rightarrow \eta\pi^0\pi^0 n$
not seen in either mode		ANDO	86	SPEC 8 $\pi^- p$
not seen in either mode		CORDEN	78	OMEG 12-15 $\pi^- p$
0.4±0.2		DEFOIX	72	HBC 0.7 $\bar{p}p \rightarrow 7\pi$

NODE=M006R4  
 NODE=M006R4

$\Gamma(4\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$   $\Gamma_6/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.90	95	DIONISI	80	HBC 4 $\pi^- p$
-------	----	---------	----	-----------------

NODE=M006R5  
 NODE=M006R5

$\Gamma(K\bar{K}\pi)/[\Gamma(K\bar{K}^*(892)+c.c.)+\Gamma(a_0(980)\pi)]$   $\Gamma_1/(\Gamma_2+\Gamma_4)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.65±0.27	<sup>14</sup> DIONISI	80	HBC 4 $\pi^- p$
-----------	-----------------------	----	-----------------

<sup>14</sup> Calculated using  $\Gamma(K\bar{K})/\Gamma(\eta\pi) = 0.24 \pm 0.07$  for  $a_0(980)$  fractions.

NODE=M006R6  
 NODE=M006R6

NODE=M006R6;LINKAGE=C

$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$   $\Gamma_4/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**0.042±0.014 OUR AVERAGE**

0.44 ±0.19	ABLIKIM	21U	BES3	$D_s^+ \rightarrow f_1(1420)\pi^+$
------------	---------	-----	------	------------------------------------

0.04 ±0.01 ±0.01	BARBERIS	98C	OMEG	450 $p\bar{p} \rightarrow p_f f_1(1420)p_s$
------------------	----------	-----	------	---

••• We do not use the following data for averages, fits, limits, etc. •••

<0.04	68	ARMSTRONG	84	OMEG 85 $\pi^+ p$
-------	----	-----------	----	-------------------

NODE=M006R7  
 NODE=M006R7

$\Gamma(4\pi)/\Gamma(K\bar{K}\pi)$   $\Gamma_6/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.62	95	ARMSTRONG	89G	OMEG 85 $\pi p \rightarrow 4\pi X$
-------	----	-----------	-----	------------------------------------

NODE=M006R8  
 NODE=M006R8



$\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma$
<0.08	95	15 ARMSTRONG 92C	SPEC	300 $p\bar{p} \rightarrow p\bar{p}\pi^+\pi^-\gamma$	

<sup>15</sup> Using the data on the  $\bar{K}K\pi$  mode from ARMSTRONG 89.

NODE=M006R9  
NODE=M006R9

NODE=M006R9;LINKAGE=A

 $\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma_1$
<0.02	95	BARBERIS 98C	OMEG	450 $p\bar{p} \rightarrow p_f f_1(1420) p_S$	

NODE=M006R10  
NODE=M006R10

 $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma_1$
<b>0.003±0.001±0.001</b>	BARBERIS 98C	OMEG	450 $p\bar{p} \rightarrow p_f f_1(1420) p_S$	

NODE=M006R11  
NODE=M006R11

### $f_1(1420)$ REFERENCES

NODE=M006

ABLIKIM 21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61155
ACHARD 07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=51698
ABDALLAH 03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49548
NICHITIU 02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
ADAMS 01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
SOSA 99	PRL 83 913	M. Sosa <i>et al.</i>		REFID=46937
BARBERIS 98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
BARBERIS 97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
BERTIN 97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
PROKOSHKIN 97B	PD 42 298	Yu.D. Prokoshkin, S.A. Sadovsky		REFID=45549
Translated from DANS 354 751.				
ARMSTRONG 92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=42097
ARMSTRONG 92E	ZPHY C56 29	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JPC	REFID=43173
AUGUSTIN 92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
ARMSTRONG 91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BAI 90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
ARMSTRONG 89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+) JPC	REFID=40729
ARMSTRONG 89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)	REFID=40930
BEHREND 89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40732
HILL 89	ZPHY C42 355	P. Hill <i>et al.</i>	(JADE Collab.) JP	REFID=40741
KOPKE 89	PRPL 174 67	L. Kopke <i>et al.</i>	(CERN)	REFID=41863
AIHARA 88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)	REFID=40572
BECKER 87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.) JP	REFID=40015
GIDAL 87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
GIDAL 87B	PRL 59 2016	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40224
AIHARA 86C	PRL 57 2500	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.) JP	REFID=21326
ANDO 86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)	REFID=20891
ARMSTRONG 84	PL 146B 273	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=20929
BITYUKOV 84	SJNP 39 735	S. Bityukov <i>et al.</i>	(SERP)	REFID=45856
Translated from YAF 39 1165.				
CHAUVAT 84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)	REFID=20932
JENNI 83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)	REFID=20304
BROMBERG 80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)	REFID=20922
DIONISI 80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+) IJP	REFID=20924
CORDEN 78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20452
DEFOIX 72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
DAHL 67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP	REFID=20321
Also	PRL 14 1074	D.H. Miller <i>et al.</i>	(LRL, UCB)	REFID=21291

$\omega(1420)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

See also the  $\omega(1650)$  particle listing.

NODE=M125

NODE=M125

NODE=M125M

NODE=M125M

→ UNCHECKED ←

 **$\omega(1420)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1410± 60 OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1418± 30± 10	824	<sup>1</sup> AKHMETSHIN 17A	CMD3	1.4-2.0 $e^+e^- \rightarrow \omega\eta$
1470± 50	13.1k	<sup>2</sup> AULCHENKO 15A	SND	1.05-1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1382± 23± 70		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1350± 20± 20		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1400± 50± 130	1.2M	<sup>3</sup> ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1450± 10		<sup>4</sup> HENNER 02	RVUE	1.2-2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1373± 70	177	<sup>5</sup> AKHMETSHIN 00D	CMD2	1.2-1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
1370± 25	5095	ANISOVICH 00H	SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
1400 <sup>+100</sup> <sub>-200</sub>		<sup>6</sup> ACHASOV 98H	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
~ 1400		<sup>7</sup> ACHASOV 98H	RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
~ 1460		<sup>8</sup> ACHASOV 98H	RVUE	$e^+e^- \rightarrow K^+K^-$
1440± 70		<sup>9</sup> CLEGG 94	RVUE	
1419± 31	315	<sup>10</sup> ANTONELLI 92	DM2	1.34-2.4 $e^+e^- \rightarrow \rho\pi$
<sup>1</sup> From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of $\pi$ and other parameters floating.				
<sup>2</sup> From a fit with contributions from $\omega(782)$ , $\phi(1020)$ , $\omega(1420)$ , and $\omega(1650)$ . See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.				
<sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.				
<sup>4</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.				
<sup>5</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.				
<sup>6</sup> Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.				
<sup>7</sup> Using the data from ANTONELLI 92.				
<sup>8</sup> Using the data from IVANOV 81 and BISELLO 88B.				
<sup>9</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.				
<sup>10</sup> From a fit to two Breit-Wigner functions interfering between them and with the $\omega, \phi$ tails with fixed (+, -, +) phases.				

NODE=M125M;LINKAGE=F

NODE=M125M;LINKAGE=E

NODE=M125M;LINKAGE=VH

NODE=M125M;LINKAGE=AB

NODE=M125M;LINKAGE=KL

NODE=M125M;LINKAGE=L1

NODE=M125M;LINKAGE=L2

NODE=M125M;LINKAGE=L3

NODE=M125M;LINKAGE=AD

NODE=M125M;LINKAGE=B

 **$\omega(1420)$  WIDTH**

NODE=M125W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>290± 190 OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
440± 125	267	<sup>1</sup> ACHASOV 20B	SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
104± 35± 10	824	<sup>2</sup> AKHMETSHIN 17A	CMD3	1.4-2.0 $e^+e^- \rightarrow \omega\eta$
880± 170	13.1k	<sup>3</sup> AULCHENKO 15A	SND	1.05-1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
480± 180		<sup>4</sup> ACHASOV 10D	SND	1.075-2.0 $e^+e^- \rightarrow \pi^0\gamma$
130± 50± 100		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
450± 70± 70		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
870 <sup>+500</sup> <sub>-300</sub> ± 450	1.2M	<sup>5</sup> ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
199± 15		<sup>6</sup> HENNER 02	RVUE	1.2-2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
188± 45	177	<sup>7</sup> AKHMETSHIN 00D	CMD2	1.2-1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
360 <sup>+100</sup> <sub>-60</sub>	5095	ANISOVICH 00H	SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
240± 70		<sup>8</sup> CLEGG 94	RVUE	
174± 59	315	<sup>9</sup> ANTONELLI 92	DM2	1.34-2.4 $e^+e^- \rightarrow \rho\pi$

NODE=M125W

→ UNCHECKED ←

- <sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass of  $\omega(1420)$  is fixed to the PDG 18 value of 1420 MeV.
- <sup>2</sup> From a fit of the interfering  $\omega(1420)$  and  $\omega(1650)$  with a relative phase of  $\pi$  and other parameters floating.
- <sup>3</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ . See ACHASOV 20A for a further analysis of the  $\pi^+\pi^-\pi^0$  data.
- <sup>4</sup> From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states  $\omega(1420)$ ,  $\rho(1450)$ ,  $\omega(1650)$ , and  $\rho(1700)$ . Systematic errors not evaluated.
- <sup>5</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- <sup>6</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.
- <sup>7</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.
- <sup>8</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- <sup>9</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega,\phi$  tails with fixed (+,-,+) phases.

NODE=M125W;LINKAGE=H

NODE=M125W;LINKAGE=F

NODE=M125W;LINKAGE=E

NODE=M125W;LINKAGE=G

NODE=M125W;LINKAGE=VH

NODE=M125W;LINKAGE=AB

NODE=M125W;LINKAGE=KL

NODE=M125W;LINKAGE=AD

NODE=M125W;LINKAGE=B

 **$\omega(1420)$  DECAY MODES**

NODE=M125215;NODE=M125

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\pi$	seen
$\Gamma_2$ $\omega\pi\pi$	seen
$\Gamma_3$ $\omega\eta$	
$\Gamma_4$ $b_1(1235)\pi$	seen
$\Gamma_5$ $e^+e^-$	seen
$\Gamma_6$ $\pi^0\gamma$	

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=7

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=6

 **$\omega(1420)$   $\Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$** 

NODE=M125230

 **$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma \times \Gamma_5/\Gamma$** NODE=M125G3  
NODE=M125G3

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.73 ± 0.08      13.1k      <sup>1</sup> AULCHENKO 15A      SND      1.05–1.80  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ 0.82 ± 0.05 ± 0.06      AUBERT,B      04N      BABR      10.6  $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$ 0.65 ± 0.13 ± 0.21      1.2M      <sup>2,3</sup> ACHASOV      03D      RVUE      0.44–2.00  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ 0.625 ± 0.160      <sup>4,5</sup> CLEGG      94      RVUE0.466 ± 0.178      <sup>6,7</sup> ANTONELLI 92      DM2      1.34–2.4  $e^+e^- \rightarrow \rho\pi$ 

<sup>1</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ . See ACHASOV 20A for a further analysis of the  $\pi^+\pi^-\pi^0$  data.

NODE=M125G3;LINKAGE=A

<sup>2</sup> Calculated by us from the cross section at the peak.

NODE=M125G;LINKAGE=AW

<sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

NODE=M125G;LINKAGE=VH

<sup>4</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

NODE=M125G;LINKAGE=AD

<sup>5</sup> From the partial and leptonic width given by the authors.

NODE=M125G;LINKAGE=SE

<sup>6</sup> From a fit to two Breit-Wigner functions interfering between them and with the  $\omega,\phi$  tails with fixed (+,-,+) phases.

NODE=M125G;LINKAGE=A

<sup>7</sup> From the product of the leptonic width and partial branching ratio given by the authors.

NODE=M125G;LINKAGE=ES

 **$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma \times \Gamma_5/\Gamma$** NODE=M125G4  
NODE=M125G4

VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

19.7 ± 5.7      AUBERT      07AU      BABR      10.6  $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$ 1.9 ± 1.9      <sup>1</sup> AKHMETSHIN 00D      CMD2      1.2–2.4  $e^+e^- \rightarrow \omega\pi^+\pi^-$ 

<sup>1</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

NODE=M125G;LINKAGE=KL

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma \times \Gamma_5/\Gamma$ 

VALUE (units $10^{-8}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2.5 \pm 0.6$	267	<sup>1</sup> ACHASOV	20B SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
$2.1^{+1.0}_{-0.8}$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
$5.0 \pm 2.6 \pm 0.3$	824	<sup>2</sup> AKHMETSHIN	17A CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$
$1.6^{+0.9}_{-0.7}$	898	<sup>3</sup> ACHASOV	16B SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$

NODE=M125G6  
 NODE=M125G6

<sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass of  $\omega(1420)$  is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of  $\omega(1420)$  to the PDG 18 value of 220 MeV results in  $(3.0 \pm 1.6) \times 10^{-8}$  measurement.

NODE=M125G6;LINKAGE=C

<sup>2</sup> From a fit of the interfering  $\omega(1420)$  and  $\omega(1650)$  with a relative phase of  $\pi$  and other parameters floating. From an alternative fit  $\Gamma(\omega(1420) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1420) \rightarrow e^+e^-) = 5.3 \pm 1.6$  eV.

NODE=M125G6;LINKAGE=B

<sup>3</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass and the width of  $\omega(1420)$  are fixed to the 2014 edition (PDG 14) of this review.

NODE=M125G6;LINKAGE=A

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma \times \Gamma_5/\Gamma$ 

VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.23 \pm 0.14$	<sup>1</sup> ACHASOV	10D SND	$1.075-2.0 e^+e^- \rightarrow \pi^0\gamma$
$2.03^{+0.70}_{-0.75}$	<sup>2</sup> AKHMETSHIN	05 CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$

NODE=M125G5  
 NODE=M125G5

<sup>1</sup> From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states  $\omega(1420)$ ,  $\rho(1450)$ ,  $\omega(1650)$ , and  $\rho(1700)$ . Systematic errors not evaluated.

NODE=M125G5;LINKAGE=A

<sup>2</sup> Using 1420 MeV and 220 MeV for the  $\omega(1420)$  mass and width.

NODE=M125G5;LINKAGE=AK

 $\omega(1420)$  BRANCHING RATIOS

NODE=M125225

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.301 \pm 0.029$ possibly seen	<sup>1</sup> HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
	AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$

NODE=M125R2  
 NODE=M125R2

 $\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$  $\Gamma_2/\Gamma_4$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.60 \pm 0.16$	5095	ANISOVICH	00H SPEC	$0.0 p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$

NODE=M125R1  
 NODE=M125R1

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	ACHASOV	20A SND	$1.15-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$0.699 \pm 0.029$	<sup>1</sup> HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M125R3  
 NODE=M125R3

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$\sim 6.6$	1.2M	<sup>2,3</sup> ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$23 \pm 1$		<sup>1</sup> HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M125R4  
 NODE=M125R4

<sup>1</sup> Assuming that the  $\omega(1420)$  decays into  $\rho\pi$  and  $\omega\pi\pi$  only.

NODE=M125R;LINKAGE=AC

<sup>2</sup> Calculated by us from the cross section at the peak.

NODE=M125R;LINKAGE=AW

<sup>3</sup> Assuming that the  $\omega(1420)$  decays into  $\rho\pi$  only.

NODE=M125R;LINKAGE=GS

$\omega(1420)$  REFERENCES

ACHASOV	20A	EPJ C80 993	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60923
ACHASOV	20B	EPJ C80 1008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60924
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59887
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=58239
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57537
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56843
		Translated from ZETF 148 34.			
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>		REFID=49177
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>		REFID=47948
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)	REFID=21586
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553

NODE=M125

 $f_2(1430)$ 

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry lists nearby peaks observed in the  $D$  wave of the  $K\bar{K}$  and  $\pi^+\pi^-$  systems. Needs confirmation.

NODE=M066

 $f_2(1430)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>\approx 1430</math> OUR ESTIMATE</b>			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1440 \pm 11 \pm 3$	LEES	21A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
$1453 \pm 4$	<sup>1</sup> VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1421 \pm 5$	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
$1480 \pm 50$	AKESSON	86	SPEC $pp \rightarrow pp \pi^+ \pi^-$
$1436^{+26}_{-16}$	DAUM	84	CNTR $17-18 \pi^- p \rightarrow K^+ K^- n$
$1412 \pm 3$	DAUM	84	CNTR $63 \pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
$1439^{+5}_{-6}$	<sup>2</sup> BEUSCH	67	OSPK $5,7,12 \pi^- p \rightarrow K_S^0 K_S^0 n$
<sup>1</sup> $J^{PC} = 0^{++}$ or $2^{++}$ .			
<sup>2</sup> Not seen by WETZEL 76.			

NODE=M066M1

NODE=M066M1

→ UNCHECKED ←

OCCUR=2

NODE=M066M;LINKAGE=AC

NODE=M066M;LINKAGE=C

 $f_2(1430)$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$46 \pm 15 \pm 5$	LEES	21A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
$13 \pm 5$	<sup>3</sup> VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$30 \pm 9$	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
$150 \pm 50$	AKESSON	86	SPEC $pp \rightarrow pp \pi^+ \pi^-$
$81^{+56}_{-29}$	DAUM	84	CNTR $17-18 \pi^- p \rightarrow K^+ K^- n$
$14 \pm 6$	DAUM	84	CNTR $63 \pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
$43^{+17}_{-18}$	<sup>4</sup> BEUSCH	67	OSPK $5,7,12 \pi^- p \rightarrow K_S^0 K_S^0 n$
<sup>3</sup> $J^{PC} = 0^{++}$ or $2^{++}$ .			
<sup>4</sup> Not seen by WETZEL 76.			

NODE=M066W1

NODE=M066W1

OCCUR=2

NODE=M066W;LINKAGE=AC

NODE=M066W;LINKAGE=C

**$f_2(1430)$  DECAY MODES**

NODE=M066215;NODE=M066

Mode	
$\Gamma_1$	$K\bar{K}$
$\Gamma_2$	$\pi\pi$

DESIG=1

DESIG=2

 **$f_2(1430)$  REFERENCES**

NODE=M066

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
VLADIMIRSK...	01	PAN 64 1895 Translated from YAF 64 1979.	V.V. Vladimisky <i>et al.</i>	
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
DAUM	84	ZPHY C23 339	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)
BEUSCH	67	PL 25B 357	W. Beusch <i>et al.</i>	(ETH, CERN)

REFID=61442  
REFID=48571REFID=40268  
REFID=21123  
REFID=21372  
REFID=20362  
REFID=20320

NODE=M149

 **$a_0(1450)$** 

$$I^G(J^{PC}) = 1^-(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M149

 **$a_0(1450)$  T-MATRIX POLE  $\sqrt{s}$** 

NODE=M149PP

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M149PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1290–1500) – <math>i</math> (30–140) OUR ESTIMATE</b>			
$(1302.1 \pm 1.1 \pm 3.9) - i$ ( $56.2 \pm 0.7 \pm 1.7$ )	<sup>1</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1515 \pm 30) - i$ ( $115 \pm 18$ )	ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1432 \pm 13 \pm 25) - i$ ( $98 \pm 5 \pm 5$ )	<sup>2</sup> BUGG	08A	RVUE $\bar{p}p$
$(1441^{+40}_{-15}) - i$ ( $55 \pm 7$ )	<sup>3</sup> BAKER	03	SPEC $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
$(1303 \pm 16) - i$ ( $46 \pm 8$ )	<sup>4</sup> BARGIOTTI	03	OBLX $\bar{p}p$
$(1296 \pm 10) - i$ ( $41 \pm 11$ )	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$(1565 \pm 30) - i$ ( $146 \pm 20$ )	ANISOVICH	98B	RVUE Compilation
$(1470 \pm 25) - i$ ( $132 \pm 15$ )	<sup>5</sup> AMSLER	95D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$

NODE=M149PP

→ UNCHECKED ←

<sup>1</sup> T-matrix pole, 2 poles, 2 channels ( $\pi\eta, K\bar{K}$ ).<sup>2</sup> Using data from AMSLER 94D, ABELE 98, and BAKER 03. Supersedes BUGG 94.<sup>3</sup> From the pole position of a fitted Breit-Wigner amplitude.<sup>4</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0,$  and  $K^\pm K_S^0 \pi^\mp$ .<sup>5</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.NODE=M149PP;LINKAGE=A  
NODE=M149PP;LINKAGE=G  
NODE=M149PP;LINKAGE=C  
NODE=M149PP;LINKAGE=D  
NODE=M149PP;LINKAGE=F **$a_0(1450)$  MASS**

NODE=M149M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1439 \pm 34</math> OUR AVERAGE</b>		Error includes scale factor of 1.8. [ $1474 \pm 19$ MeV OUR 2022 AVERAGE]		
$1480 \pm 30$		ABELE	98	CBAR $0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
$1410 \pm 25$		ETKIN	82C	MPS $23 \pi^- p \rightarrow n 2K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1458 \pm 14 \pm 15$	190k	<sup>1</sup> AAIJ	16N	LHCB $D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
$1316.8^{+0.7+24.7}_{-1.0-4.6}$		<sup>2</sup> UEHARA	09A	BELL $\gamma\gamma \rightarrow \pi^0 \eta$
$1477 \pm 10$	80k	<sup>3</sup> UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
$1290 \pm 10$		<sup>4</sup> BERTIN	98B	OBLX $0.0 \bar{p}p \rightarrow K^\pm K_S \pi^\mp$
$1450 \pm 40$		AMSLER	94D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$\sim 1300$		MARTIN	78	SPEC $10 K^\pm p \rightarrow K_S^0 \pi p$
$1255 \pm 5$		<sup>5</sup> CASON	76	

NODE=M149M

NEW

- <sup>1</sup> Using a model with Gaussian constraints to the PDG averaged values .  
<sup>2</sup> May be a different state.  
<sup>3</sup> Statistical error only.  
<sup>4</sup> Not confirmed by BUGG 08A.  
<sup>5</sup> Isospin 0 not excluded.

NODE=M149M;LINKAGE=A  
 NODE=M149M;LINKAGE=UE  
 NODE=M149M;LINKAGE=ST  
 NODE=M149M;LINKAGE=BE  
 NODE=M149M;LINKAGE=CC

### $a_0(1450)$ WIDTH

NODE=M149W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>258 ± 14 OUR AVERAGE</b>				
[265 ± 13 MeV OUR 2022 AVERAGE]				
265 ± 15		ABELE	98 CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
230 ± 30		ETKIN	82C MPS	23 $\pi^- p \rightarrow n 2K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
282 ± 12 ± 13	190k	<sup>1</sup> AAIJ	16N LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
65.0 <sup>+</sup> <sub>-5.4</sub> <sup>2.1+99.1</sup> <sub>-32.6</sub>		<sup>2</sup> UEHARA	09A BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
267 ± 11	80k	<sup>3</sup> UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
80 ± 5		<sup>4</sup> BERTIN	98B OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
270 ± 40		AMSLER	94D CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
~ 250		MARTIN	78 SPEC	10 $K^\pm p \rightarrow K_S^0 \pi p$
79 ± 10		<sup>5</sup> CASON	76	

NODE=M149W  
 NEW

- <sup>1</sup> Using a model with Gaussian constraints to the PDG averaged values .  
<sup>2</sup> May be a different state.  
<sup>3</sup> Statistical error only.  
<sup>4</sup> Not confirmed by BUGG 08A.  
<sup>5</sup> Isospin 0 not excluded.

NODE=M149W;LINKAGE=A  
 NODE=M149W;LINKAGE=UE  
 NODE=M149W;LINKAGE=ST  
 NODE=M149W;LINKAGE=BE  
 NODE=M149W;LINKAGE=CC

### $a_0(1450)$ DECAY MODES

NODE=M149215;NODE=M149

Branching fractions are given relative to the one **DEFINED AS 1**.

NODE=M149

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi \eta$	0.093 ± 0.020
$\Gamma_2$ $\pi \eta'(958)$	0.033 ± 0.017
$\Gamma_3$ $K \bar{K}$	0.082 ± 0.028
$\Gamma_4$ $\omega \pi \pi$	<b>DEFINED AS 1</b>
$\Gamma_5$ $a_0(980) \pi \pi$	seen
$\Gamma_6$ $\gamma \gamma$	seen

DESIG=1  
 DESIG=2  
 DESIG=3  
 DESIG=4  
 DESIG=5  
 DESIG=6

### $a_0(1450)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M149225

$\Gamma(\pi\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1 \Gamma_6 / \Gamma$
432 ± 6 <sup>+</sup> <sub>-256</sub> <sup>1073</sup>	<sup>1</sup> UEHARA	09A BELL	$\gamma\gamma \rightarrow \pi^0 \eta$	

NODE=M149G01  
 NODE=M149G01

- <sup>1</sup> May be a different state.

NODE=M149G01;LINKAGE=UE

### $a_0(1450)$ BRANCHING RATIOS

NODE=M149220

$\Gamma(\pi\eta'(958))/\Gamma(\pi\eta)$	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_1$
<b>0.35 ± 0.16</b>	<sup>1</sup> ABELE	98 CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$	
0.43 ± 0.19	ABELE	97C CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta'$	

NODE=M149R1  
 NODE=M149R1

- <sup>1</sup> Using  $\pi^0 \eta$  from AMSLER 94D.

NODE=M149R1;LINKAGE=A

$\Gamma(K\bar{K})/\Gamma(\pi\eta)$  $\Gamma_3/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.88 ± 0.23</b>	<sup>1</sup> ABELE 98	CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
1.887 ± 0.041 ± 0.07	<sup>2</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$

<sup>1</sup> Using  $\pi^0 \eta$  from AMSLER 94D.<sup>2</sup> Residues from T-matrix pole, 2 poles, 2 channels ( $\pi\eta$ ,  $K\bar{K}$ ).NODE=M149R2  
NODE=M149R2NODE=M149R2;LINKAGE=A  
NODE=M149R2;LINKAGE=B $\Gamma(\omega\pi\pi)/\Gamma(\pi\eta)$  $\Gamma_4/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.7 ± 2.3</b>	35280	<sup>1</sup> BAKER 03	SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

<sup>1</sup> Using results on  $\bar{p}p \rightarrow a_0(1450)^0 \pi^0$ ,  $a_0(1450) \rightarrow \eta\pi^0$  from ABELE 96C and assuming the  $\omega\rho$  mechanism for the  $\omega\pi\pi$  state.NODE=M149R3  
NODE=M149R3

NODE=M149R;LINKAGE=PP

 $\Gamma(a_0(980)\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	BUGG 08A	RVUE	$\bar{p}p$

NODE=M149R01  
NODE=M149R01 $\Gamma(a_0(980)\pi\pi)/\Gamma(\pi\eta)$  $\Gamma_5/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
$\leq 4.3$	ANISOVICH 01	RVUE	0	$\bar{p}p \rightarrow \eta 2\pi^+ 2\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M149R02  
NODE=M149R02 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> UEHARA 09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$

<sup>1</sup> May be a different state.NODE=M149R03  
NODE=M149R03

NODE=M149R03;LINKAGE=UE

 **$a_0(1450)$  REFERENCES**

ALBRECHT 20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
AAIJ 16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
ANISOVICH 09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)
UEHARA 09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)
BUGG 08A	PR D78 074023	D.V. Bugg	(LOQM)
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
BAKER 03	PL B563 140	C.A. Baker <i>et al.</i>	
BARGIOTTI 03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
AMSLER 02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH 01	NP A690 567	A.V. Anisovich <i>et al.</i>	
ABELE 98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH 98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
	Translated from UFN 168 481.		
BERTIN 98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE 97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE 96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER 94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) IGJPC
BUGG 94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
ETKIN 82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
MARTIN 78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)
CASON 76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL)

NODE=M149

REFID=60439  
REFID=57273  
REFID=52719  
REFID=53002  
REFID=52578  
REFID=51063  
REFID=49414  
REFID=49217  
REFID=48580  
REFID=48308  
REFID=45863  
REFID=46331REFID=46351  
REFID=45531  
REFID=45076  
REFID=44377  
REFID=44440  
REFID=44441  
REFID=44093  
REFID=44078  
REFID=20391  
REFID=22446  
REFID=21064



$\rho(1450)$ 

$$I^G(J^{PC}) = 1^+(1^{--})$$

NODE=M105

 **$\rho(1450)$  MASS**

NODE=M105205

 **$\rho(1450)$  MASS**

VALUE (MeV)

DOCUMENT ID

**1465±25 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.

NODE=M105M0

NODE=M105M0

→ UNCHECKED ←

 **$\eta\rho^0$  MODE**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1506±11	13.4k	<sup>1</sup> GRIBANOV 20	CMD3	1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1500±10	7.4k	<sup>2</sup> ACHASOV 18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1497±14		<sup>3</sup> AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1421±15		<sup>4</sup> AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1470±20		ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1446±10		FUKUI 88	SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

NODE=M105M1

NODE=M105M1

<sup>1</sup> Mass and width of the  $\rho(770)$  fixed at 775 and 149 MeV, respectively; solution 2 of model 2,  $\eta \rightarrow \gamma\gamma$  decays used.

NODE=M105M1;LINKAGE=B

<sup>2</sup> From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering  $\rho(1450)$ ,  $\rho(1700)$  and  $\rho(2150)$  with the parameters of the  $\rho(1450)$  and  $\rho(1700)$  floating and the mass and width of the  $\rho(2150)$  fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are  $\pi$ , 0 and  $\pi$ , respectively.

NODE=M105M1;LINKAGE=A

<sup>3</sup> Using the data of AKHMETSHIN 01B on  $e^+e^- \rightarrow \eta\gamma$ , AKHMETSHIN 00D and ANTONELLI 88 on  $e^+e^- \rightarrow \eta\pi^+\pi^-$ .

NODE=M105M;LINKAGE=SW

<sup>4</sup> Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the  $\rho(1450)$  and  $\rho(1700)$  mesons assumed.

NODE=M105M1;LINKAGE=KL

 **$\omega\pi$  MODE**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1510±7	10.2k	<sup>1</sup> ACHASOV 16D	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1544±22 <sup>+11</sup> <sub>-46</sub>	821	<sup>2</sup> MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
1491±19	7815	<sup>3</sup> ACHASOV 13	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1582±17±25	2382	<sup>4</sup> AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1349±25 <sup>+10</sup> <sub>-5</sub>	341	<sup>5</sup> ALEXANDER 01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
1523±10		<sup>6</sup> EDWARDS 00A	CLE2	$\tau^- \rightarrow \omega\pi^- \nu_\tau$
1463±25		<sup>7</sup> CLEGG 94	RVUE	
1250		<sup>8</sup> ASTON 80C	OMEG	20–70 $\gamma p \rightarrow \omega\pi^0 p$
1290±40		<sup>8</sup> BARBER 80C	SPEC	3–5 $\gamma p \rightarrow \omega\pi^0 p$

NODE=M105M3

NODE=M105M3

OCCUR=2

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$ . The  $\rho(1700)$  mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

NODE=M105M3;LINKAGE=D

<sup>2</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming equal probabilities of the  $\rho(1450) \rightarrow \pi\pi$  and  $\rho(1450) \rightarrow \omega\pi$  decays.

NODE=M105M3;LINKAGE=C

<sup>3</sup> From a phenomenological model based on vector meson dominance with the interfering  $\rho(1450)$  and  $\rho(1700)$  and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M105M3;LINKAGE=AC

<sup>4</sup> Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the  $\omega\pi^0$  and  $\pi^+\pi^-$  mass dependence of the total width.  $\rho(1700)$  mass and width fixed at 1700 MeV and 240 MeV, respectively.

NODE=M105M3;LINKAGE=HK

<sup>5</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming the  $\omega\pi^-$  mass dependence for the total width.

NODE=M105M3;LINKAGE=3Z

<sup>6</sup> Mass-independent width parameterization.  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV respectively.

NODE=M105M;LINKAGE=E1

<sup>7</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

NODE=M105M3;LINKAGE=B

<sup>8</sup> Not separated from  $b_1(1235)$ , not pure  $J^P = 1^-$  effect.

NODE=M105M3;LINKAGE=A

 **$4\pi$  MODE**

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1435±40		ABELE 01B	CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0\pi^+$
1350±50		ACHASOV 97	RVUE	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1449±4		<sup>1</sup> ARMSTRONG 89E	OMEG	300 $pp \rightarrow pp2(\pi^+\pi^-)$

NODE=M105M6

NODE=M105M6

<sup>1</sup> Not clear whether this observation has  $l=1$  or 0.

NODE=M105M6;LINKAGE=A

**$\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1326.35 ± 3.46		<sup>1</sup> BARTOS 17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1342.31 ± 46.62		<sup>2</sup> BARTOS 17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1373.83 ± 11.37		<sup>3</sup> BARTOS 17A	RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1429 ± 41	20k	<sup>4</sup> LEES 17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
1350 ± 20	$\begin{matrix} +20 \\ -30 \end{matrix}$ 63.5k	<sup>5</sup> ABRAMOWICZ12	ZEUS	$e p \rightarrow e\pi^+\pi^- p$
1493 ± 15		<sup>6</sup> LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
1446 ± 7	$\pm 28$ 5.4M	<sup>7,8</sup> FUJIKAWA 08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1328 ± 15		<sup>9</sup> SCHAELE 05C	ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1406 ± 15	87k	<sup>7,10</sup> ANDERSON 00A	CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
~ 1368		<sup>11</sup> ABELE 99C	CBAR	$0.0 \bar{p}d \rightarrow \pi^+\pi^-\pi^- p$
1348 ± 33		BERTIN 98	OBLX	$0.05-0.405 \bar{n}p \rightarrow$ $2\pi^+\pi^-$
1411 ± 14		<sup>12</sup> ABELE 97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
1370 $\begin{matrix} +90 \\ -70 \end{matrix}$		ACHASOV 97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1359 ± 40		<sup>10</sup> BERTIN 97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1282 ± 37		BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+2\pi^-$
1424 ± 25		BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
1265.5 ± 75.3		DUBNICKA 89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1292 ± 17		<sup>13</sup> KURDADZE 83	OLYA	$0.64-1.4 e^+e^- \rightarrow$ $\pi^+\pi^-$

- <sup>1</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
- <sup>2</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
- <sup>3</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.
- <sup>4</sup> From a Dalitz plot analysis in an isobar model with  $\rho(1450)$  and  $\rho(1700)$  masses and widths floating.
- <sup>5</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho-\omega$  interference.
- <sup>6</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.
- <sup>7</sup> From the GOUNARIS 68 parametrization of the pion form factor.
- <sup>8</sup>  $|F_\pi(0)|^2$  fixed to 1.
- <sup>9</sup> From the combined fit of the  $\tau^-$  data from ANDERSON 00A and SCHAELE 05C and  $e^+e^-$  data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05.  $\rho(1700)$  mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
- <sup>10</sup>  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV, respectively.
- <sup>11</sup>  $\rho(1700)$  mass and width fixed at 1780 MeV and 275 MeV respectively.
- <sup>12</sup> T-matrix pole.
- <sup>13</sup> Using for  $\rho(1700)$  mass and width  $1600 \pm 20$  and  $300 \pm 10$  MeV respectively.

NODE=M105M5  
NODE=M105M5

OCCUR=2

OCCUR=2

NODE=M105M5;LINKAGE=D

NODE=M105M5;LINKAGE=E

NODE=M105M5;LINKAGE=F

NODE=M105M5;LINKAGE=B

NODE=M105M5;LINKAGE=AB

NODE=M105M5;LINKAGE=LE

NODE=M105M5;LINKAGE=1K

NODE=M105M5;LINKAGE=FU

NODE=M105M5;LINKAGE=SC

NODE=M105M5;LINKAGE=A

NODE=M105M5;LINKAGE=C5

NODE=M105M5;LINKAGE=QQ

NODE=M105M5;LINKAGE=KD

 **$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1208 ± 8 ± 9	190k	<sup>1</sup> AAJ 16N	LHCB		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
1422.8 ± 6.5	27k	<sup>2</sup> ABELE 99D	CBAR	±	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$

- <sup>1</sup> Using the GOUNARIS 68 parameterization with fixed width.
- <sup>2</sup> K-matrix pole. Isospin not determined, could be  $\omega(1420)$ .

NODE=M105M7  
NODE=M105M7

NODE=M105M7;LINKAGE=A

NODE=M105M7;LINKAGE=AN

 **$K\bar{K}^*(892) + c.c.$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1505 ± 19 ± 7	AUBERT 08S	BABR 10.6	$e^+e^- \rightarrow K\bar{K}^*(892)\gamma$

NODE=M105M8  
NODE=M105M8 **$m_{\rho(1450)^0} - m_{\rho(1450)^\pm}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-31.53 ± 47.99	<sup>1</sup> BARTOS 17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$ , $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

NODE=M105DM  
NODE=M105DM

- <sup>1</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M105DM;LINKAGE=A

**$\rho(1450)$  WIDTH**

NODE=M105210

 **$\rho(1450)$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**400± 60 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.

• • • We do not use the following data for averages, fits, limits, etc. • • •

480±180	<sup>1</sup> ACHASOV	10D	SND	1.075–2.0 $e^+e^- \rightarrow \pi^0\gamma$
---------	----------------------	-----	-----	--

<sup>1</sup> From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states  $\omega(1420)$ ,  $\rho(1450)$ ,  $\omega(1650)$ , and  $\rho(1700)$ . Systematic errors not evaluated.

NODE=M105W0

NODE=M105W0

→ UNCHECKED ←

NODE=M105W0;LINKAGE=A

 **$\eta\rho^0$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

321±27	13.4k	<sup>1</sup> GRIBANOV	20	CMD3	1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
280±20	7.4k	<sup>2</sup> ACHASOV	18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
226±44		<sup>3</sup> AKHMETSHIN 01B	CMD2		$e^+e^- \rightarrow \eta\gamma$
211±31		<sup>4</sup> AKHMETSHIN 00D	CMD2		$e^+e^- \rightarrow \eta\pi^+\pi^-$
230±30		ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
60±15		FUKUI	88	SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

<sup>1</sup> Mass and width of the  $\rho(770)$  fixed at 775 and 149 MeV, respectively; solution 2 of model 2,  $\eta \rightarrow \gamma\gamma$  decays used.

<sup>2</sup> From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering  $\rho(1450)$ ,  $\rho(1700)$  and  $\rho(2150)$  with the parameters of the  $\rho(1450)$  and  $\rho(1700)$  floating and the mass and width of the  $\rho(2150)$  fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are  $\pi$ , 0 and  $\pi$ , respectively.

<sup>3</sup> Using the data of AKHMETSHIN 01B on  $e^+e^- \rightarrow \eta\gamma$ , AKHMETSHIN 00D and ANTONELLI 88 on  $e^+e^- \rightarrow \eta\pi^+\pi^-$ .

<sup>4</sup> Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the  $\rho(1450)$  and  $\rho(1700)$  mesons assumed.

NODE=M105W1

NODE=M105W1

NODE=M105W1;LINKAGE=B

NODE=M105W1;LINKAGE=A

NODE=M105W;LINKAGE=SW

NODE=M105W1;LINKAGE=KL

 **$\omega\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

440± 40	10.2k	<sup>1</sup> ACHASOV	16D	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
303 <sup>+</sup> <sub>–</sub> <sup>31+69</sup> <sub>52–7</sub>	821	<sup>2</sup> MATVIENKO	15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
429± 42±10	2382	<sup>3</sup> AKHMETSHIN 03B	CMD2		$e^+e^- \rightarrow \pi^0\pi^0\gamma$
547± 86 <sup>+</sup> <sub>–</sub> <sup>46</sup> <sub>45</sub>	341	<sup>4</sup> ALEXANDER	01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
400± 35		<sup>5</sup> EDWARDS	00A	CLE2	$\tau^- \rightarrow \omega\pi^-\nu_\tau$
311± 62		<sup>6</sup> CLEGG	94	RVUE	
300		<sup>7</sup> ASTON	80C	OMEG	20–70 $\gamma p \rightarrow \omega\pi^0 p$
320±100		<sup>7</sup> BARBER	80C	SPEC	3–5 $\gamma p \rightarrow \omega\pi^0 p$

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$ . The  $\rho(1700)$  mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

<sup>2</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming equal probabilities of the  $\rho(1450) \rightarrow \pi\pi$  and  $\rho(1450) \rightarrow \omega\pi$  decays.

<sup>3</sup> Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the  $\omega\pi^0$  and  $\pi^+\pi^-$  mass dependence of the total width.  $\rho(1700)$  mass and width fixed at 1700 MeV and 240 MeV, respectively.

<sup>4</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming the  $\omega\pi^-$  mass dependence for the total width.

<sup>5</sup> Mass-independent width parameterization.  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV respectively.

<sup>6</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

<sup>7</sup> Not separated from  $b_1(1235)$ , not pure  $J^P = 1^-$  effect.

NODE=M105W3

NODE=M105W3

OCCUR=3

NODE=M105W3;LINKAGE=D

NODE=M105W3;LINKAGE=C

NODE=M105W3;LINKAGE=HK

NODE=M105W3;LINKAGE=3Z

NODE=M105W;LINKAGE=E1

NODE=M105W3;LINKAGE=B

NODE=M105W3;LINKAGE=A

**4 $\pi$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

325±100	ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0\pi^+$
---------	-------	-----	------	---

NODE=M105W66

NODE=M105W66

**$\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
324.13 ± 12.01		1 BARTOS	17 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
492.17 ± 138.38		2 BARTOS	17A RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
340.87 ± 23.84		3 BARTOS	17A RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
576 ± 29	20k	4 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
460 ± 30	$\begin{matrix} +40 \\ -45 \end{matrix}$ 63.5k	5 ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^-p$
427 ± 31		6 LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
434 ± 16	±60 5.4M	7,8 FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
468 ± 41		9 SCHAEEL	05C ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
455 ± 41	87k	7,10 ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
~ 374		11 ABELE	99C CBAR	$0.0 \bar{p}d \rightarrow \pi^+\pi^-\pi^-p$
275 ± 10		BERTIN	98 OBLX	$0.05-0.405 \bar{n}p \rightarrow \pi^+\pi^+\pi^-$
343 ± 20		12 ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
310 ± 40		10 BERTIN	97C OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
236 ± 36		BERTIN	97D OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+2\pi^-$
269 ± 31		BISELLO	89 DM2	$e^+e^- \rightarrow \pi^+\pi^-$
391 ± 70		DUBNICKA	89 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
218 ± 46		13 KURDADZE	83 OLYA	$0.64-1.4 e^+e^- \rightarrow \pi^+\pi^-$

NODE=M105W5  
 NODE=M105W5

OCCUR=2

- 1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
- 2 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
- 3 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.
- 4 From a Dalitz plot analysis in an isobar model with  $\rho(1450)$  and  $\rho(1700)$  masses and widths floating.
- 5 Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho-\omega$  interference.
- 6 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.
- 7 From the GOUNARIS 68 parametrization of the pion form factor.
- 8  $|F_\pi(0)|^2$  fixed to 1.
- 9 From the combined fit of the  $\tau^-$  data from ANDERSON 00A and SCHAEEL 05C and  $e^+e^-$  data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05.  $\rho(1700)$  mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
- 10  $\rho(1700)$  mass and width fixed at 1700 MeV and 235 MeV, respectively.
- 11  $\rho(1700)$  mass and width fixed at 1780 MeV and 275 MeV respectively.
- 12 T-matrix pole.
- 13 Using for  $\rho(1700)$  mass and width  $1600 \pm 20$  and  $300 \pm 10$  MeV respectively.

NODE=M105W5;LINKAGE=C

NODE=M105W5;LINKAGE=D

NODE=M105W5;LINKAGE=E

NODE=M105W5;LINKAGE=B

NODE=M105W5;LINKAGE=AB

NODE=M105W5;LINKAGE=LE

NODE=M105W5;LINKAGE=1K

NODE=M105W5;LINKAGE=FU

NODE=M105W5;LINKAGE=SC

NODE=M105W5;LINKAGE=A

NODE=M105W5;LINKAGE=C5

NODE=M105W5;LINKAGE=QQ

NODE=M105W5;LINKAGE=KD

 **$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
410 ± 19 ± 35	190k	1 AAIJ	16N LHCb		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
146.5 ± 10.5	27k	2 ABELE	99D CBAR	±	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$

- 1 Using the GOUNARIS 68 parameterization with fixed mass.
- 2 K-matrix pole. Isospin not determined, could be  $\omega(1420)$ .

NODE=M105W7  
 NODE=M105W7

NODE=M105W7;LINKAGE=A

NODE=M105W7;LINKAGE=AN

 **$K\bar{K}^*(892) + c.c.$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
418 ± 25 ± 4	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma$

NODE=M105W8  
 NODE=M105W8

 **$\Gamma_{\rho(1450)^0} = \Gamma_{\rho(1450)^\pm}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
151.30 ± 140.42	1 BARTOS	17A RVUE	$e^+e^- \rightarrow \pi^+\pi^-$ , $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

- • • We do not use the following data for averages, fits, limits, etc. • • •

- 1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M105DW  
 NODE=M105DW

NODE=M105DW;LINKAGE=A

**$\rho(1450)$  DECAY MODES**

NODE=M105215;NODE=M105

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $\pi^+\pi^-$	seen
$\Gamma_3$ $4\pi$	seen
$\Gamma_4$ $\omega\pi$	
$\Gamma_5$ $a_1(1260)\pi$	
$\Gamma_6$ $h_1(1170)\pi$	
$\Gamma_7$ $\pi(1300)\pi$	
$\Gamma_8$ $\rho\rho$	
$\Gamma_9$ $\rho(\pi\pi)$ S-wave	
$\Gamma_{10}$ $e^+e^-$	seen
$\Gamma_{11}$ $\eta\rho$	seen
$\Gamma_{12}$ $a_2(1320)\pi$	not seen
$\Gamma_{13}$ $K\bar{K}$	seen
$\Gamma_{14}$ $K^+K^-$	seen
$\Gamma_{15}$ $K\bar{K}^*(892)+c.c.$	possibly seen
$\Gamma_{16}$ $\pi^0\gamma$	seen
$\Gamma_{17}$ $\eta\gamma$	seen
$\Gamma_{18}$ $f_0(500)\gamma$	not seen
$\Gamma_{19}$ $f_0(980)\gamma$	not seen
$\Gamma_{20}$ $f_0(1370)\gamma$	not seen
$\Gamma_{21}$ $f_2(1270)\gamma$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=20;OUR EVAL;→ UNCHECKED ←  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=6  
DESIG=10  
DESIG=11  
DESIG=12  
DESIG=13  
DESIG=14  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=3  
DESIG=8;OUR EST;→ UNCHECKED ←  
DESIG=7;OUR EVAL;→ UNCHECKED ←  
DESIG=21;OUR EVAL;→ UNCHECKED ←  
DESIG=15;OUR EST;→ UNCHECKED ←  
DESIG=23;OUR EST;→ UNCHECKED ←  
DESIG=9  
DESIG=16;OUR EST;→ UNCHECKED ←  
DESIG=17;OUR EST;→ UNCHECKED ←  
DESIG=18;OUR EST;→ UNCHECKED ←  
DESIG=19;OUR EST;→ UNCHECKED ←

 **$\rho(1450) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

NODE=M105220

 **$\Gamma(\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_1\Gamma_{10}/\Gamma$** NODE=M105G3  
NODE=M105G3

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.12	<sup>1</sup> DIEKMAN 88	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
$0.027^{+0.015}_{-0.010}$	<sup>2</sup> KURDADZE 83	OLYA	$0.64-1.4 e^+e^- \rightarrow \pi^+\pi^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<sup>1</sup> Using total width = 235 MeV.<sup>2</sup> Using for  $\rho(1700)$  mass and width  $1600 \pm 20$  and  $300 \pm 10$  MeV respectively.NODE=M105G3;LINKAGE=B  
NODE=M105G3;LINKAGE=KD **$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{11}\Gamma_{10}/\Gamma$** NODE=M105G4  
NODE=M105G4

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$335 \pm 27 \pm 20$	13.4k	<sup>1</sup> GRIBANOV 20	CMD3	$1.1-2.0 e^+e^- \rightarrow \eta\pi^+\pi^-$
$210 \pm 24 \pm 10$		<sup>2</sup> LEES 18	BABR	$e^+e^- \rightarrow \eta\pi^+\pi^-$
$74 \pm 20$		<sup>3</sup> AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
$91 \pm 19$		ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$

<sup>1</sup> Mass and width of the  $\rho(770)$  fixed at 775 and 149 MeV, respectively; solution 2 of model 2,  $\eta \rightarrow \gamma\gamma$  decays used.<sup>2</sup> Includes non-resonant contribution. The selected fit model includes three  $\rho$  excited states. Model uncertainty is 20%.<sup>3</sup> Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the  $\rho(1450)$  and  $\rho(1700)$  mesons assumed.

NODE=M105G4;LINKAGE=B

NODE=M105G4;LINKAGE=A

NODE=M105G4;LINKAGE=KL

 **$\Gamma(K\bar{K}^*(892)+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{15}\Gamma_{10}/\Gamma$** NODE=M105G8  
NODE=M105G8

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$127 \pm 15 \pm 6$	AUBERT 08S	BABR	$10.6 e^+e^- \rightarrow K\bar{K}^*(892)\gamma$

 **$\Gamma(\eta\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{17}\Gamma_{10}/\Gamma$** NODE=M105G6  
NODE=M105G6

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<16.4	<sup>1</sup> AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$
$2.2 \pm 0.5 \pm 0.3$	<sup>2</sup> AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$

<sup>1</sup> From  $2\gamma$  decay mode of  $\eta$  using 1465 MeV and 310 MeV for the  $\rho(1450)$  mass and width. Recalculated by us.<sup>2</sup> Using the data of AKHMETSHIN 01B on  $e^+e^- \rightarrow \eta\gamma$ , AKHMETSHIN 00D and ANTONELLI 88 on  $e^+e^- \rightarrow \eta\pi^+\pi^-$ . Recalculated by us using width of 226 MeV.

NODE=M105G6;LINKAGE=AK

NODE=M105G;LINKAGE=SW

$\rho(1450) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$ 

NODE=M105230

$$\Gamma(\omega\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R05  
NODE=M105R05

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.1 \pm 0.4$	10.2k	<sup>1</sup> ACHASOV	16D SND	$1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$5.3 \pm 0.4$	7815	<sup>2</sup> ACHASOV	13 SND	$1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

OCCUR=3

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$ . The  $\rho(1700)$  mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

NODE=M105R05;LINKAGE=A

<sup>2</sup> From a phenomenological model based on vector meson dominance with the interfering  $\rho(1450)$  and  $\rho(1700)$  and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M105R05;LINKAGE=AC

$$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R00  
NODE=M105R00

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.3 \pm 0.3$	7.4k	<sup>1</sup> ACHASOV	18 SND	$1.22-2.00 e^+ e^- \rightarrow \eta \pi^+ \pi^-$
$4.3^{+1.1}_{-0.9} \pm 0.2$	4.9k	<sup>2</sup> AULCHENKO	15 SND	$1.22-2.00 e^+ e^- \rightarrow \eta \pi^+ \pi^-$

<sup>1</sup> From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering  $\rho(1450)$ ,  $\rho(1700)$  and  $\rho(2150)$  with the parameters of the  $\rho(1450)$  and  $\rho(1700)$  floating and the mass and width of the  $\rho(2150)$  fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are  $\pi$ , 0 and  $\pi$ , respectively.

NODE=M105R00;LINKAGE=B

<sup>2</sup> From a fit to the  $e^+ e^- \rightarrow \eta \pi^+ \pi^-$  cross section with vector meson dominance model including  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$  decaying exclusively via  $\eta \rho(770)$ . Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

NODE=M105R00;LINKAGE=A

$$\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R17  
NODE=M105R17

VALUE (units $10^{-9}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.3 \pm 1.4$	<sup>1</sup> ACHASOV	10D SND	$1.075-2.0 e^+ e^- \rightarrow \pi^0 \gamma$
---------------	----------------------	---------	--

<sup>1</sup> From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states  $\omega(1420)$ ,  $\rho(1450)$ ,  $\omega(1650)$ , and  $\rho(1700)$ . Systematic errors not evaluated.

NODE=M105R17;LINKAGE=A

$$\Gamma(f_0(500)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R01  
NODE=M105R01

VALUE (units $10^{-9}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<4.0$	90	ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	---------	--------	--

$$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R02  
NODE=M105R02

VALUE (units $10^{-9}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<2.6$	90	ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	---------	--------	--

$$\Gamma(f_0(1370)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R03  
NODE=M105R03

VALUE (units $10^{-9}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<3.5$	90	ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	---------	--------	--

$$\Gamma(f_2(1270)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R04  
NODE=M105R04

VALUE (units $10^{-9}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<0.8$	90	<sup>1</sup> ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	----------------------	--------	--

<sup>1</sup> Using Breit-Wigner parametrization of the  $\rho(1450)$  with mass and width of 1465 MeV and 400 MeV, respectively.

NODE=M105R01;LINKAGE=AC

 $\rho(1450)$  BRANCHING RATIOS

NODE=M105225

$$\Gamma(\pi\pi)/\Gamma(4\pi) \quad \Gamma_1/\Gamma_3$$

NODE=M105R15  
NODE=M105R15

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.37 \pm 0.10$	<sup>1,2</sup> ABELE	01B CBAR	$0.0 \bar{p} n \rightarrow 5\pi$
-----------------	----------------------	----------	----------------------------------

<sup>1</sup>  $\omega\pi$  not included.

<sup>2</sup> Using ABELE 97.

NODE=M105R;LINKAGE=BL

NODE=M105R;LINKAGE=LK

$\Gamma(K^+K^-)/\Gamma(\pi^+\pi^-)$  $\Gamma_{14}/\Gamma_2$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>30.7±8.4±8.2</b>	20k	<sup>1</sup> LEES	17C	BABR $J/\psi \rightarrow h^+ h^- \pi^0$

NODE=M105R08  
 NODE=M105R08

<sup>1</sup> From Dalitz plot analyses in isobar models.

NODE=M105R08;LINKAGE=A

 $\Gamma(\omega\pi)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

seen	821	<sup>1</sup> MATVIENKO	15	BELL $\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$
seen	1.6k	ACHASOV	12	SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
~ 0.21		CLEGG	94	RVUE

NODE=M105R5  
 NODE=M105R5

<sup>1</sup> Using Breit-Wigner parameterization of the  $\rho(1450)$  and assuming equal probabilities of the  $\rho(1450) \rightarrow \pi\pi$  and  $\rho(1450) \rightarrow \omega\pi$  decays.

NODE=M105R5;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(\omega\pi)$  $\Gamma_1/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN
-------	-------------	------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 0.32	CLEGG	94	RVUE
--------	-------	----	------

NODE=M105R6  
 NODE=M105R6

 $\Gamma(\omega\pi)/\Gamma(4\pi)$  $\Gamma_4/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN
-------	-------------	------

••• We do not use the following data for averages, fits, limits, etc. •••

< 0.14	CLEGG	88	RVUE
--------	-------	----	------

NODE=M105R3  
 NODE=M105R3

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$  $\Gamma_5/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.27±0.08	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup>  $\omega\pi$  not included.

NODE=M105R10  
 NODE=M105R10

NODE=M105R10;LINKAGE=BL

 $\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$  $\Gamma_6/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.08±0.04	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup>  $\omega\pi$  not included.

NODE=M105R11  
 NODE=M105R11

NODE=M105R11;LINKAGE=BL

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$  $\Gamma_7/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.37±0.13	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup>  $\omega\pi$  not included.

NODE=M105R12  
 NODE=M105R12

NODE=M105R12;LINKAGE=BL

 $\Gamma(\rho\rho)/\Gamma(4\pi)$  $\Gamma_8/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.11±0.05	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup>  $\omega\pi$  not included.

NODE=M105R13  
 NODE=M105R13

NODE=M105R13;LINKAGE=BL

 $\Gamma(\rho(\pi\pi)_{\text{S-wave}})/\Gamma(4\pi)$  $\Gamma_9/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.17±0.09	<sup>1</sup> ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup>  $\omega\pi$  not included.

NODE=M105R14  
 NODE=M105R14

NODE=M105R14;LINKAGE=BL

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

seen	35	<sup>1</sup> ACHASOV	14	SND $1.15\text{--}2.00 e^+ e^- \rightarrow \eta\gamma$
< 0.04		DONNACHIE	87B	RVUE

NODE=M105R2  
 NODE=M105R2

<sup>1</sup> From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

NODE=M105R2;LINKAGE=A

$\Gamma(\eta\rho)/\Gamma(\omega\pi)$  $\Gamma_{11}/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R4  
 NODE=M105R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.081 \pm 0.020$	<sup>1,2</sup> AULCHENKO	15	SND	$1.22-2.00 e^+ e^- \rightarrow \eta\pi^+\pi^-$
$\sim 0.24$	<sup>3</sup> DONNACHIE	91	RVUE	
$>2$	FUKUI	91	SPEC	$8.95 \pi^- \rho \rightarrow \omega\pi^0 n$

<sup>1</sup> From a fit to the  $e^+ e^- \rightarrow \eta\pi^+\pi^-$  cross section with vector meson dominance model including  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$  decaying exclusively via  $\eta\rho(770)$ . Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

NODE=M105R4;LINKAGE=A

<sup>2</sup> Reports the inverse of the quoted value as  $12.3 \pm 3.1$ .

NODE=M105R4;LINKAGE=B  
 NODE=M105R;LINKAGE=A

<sup>3</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

 $\Gamma(\pi\pi)/\Gamma(\eta\rho)$  $\Gamma_1/\Gamma_{11}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R07  
 NODE=M105R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.3 \pm 0.4$	<sup>1</sup> AULCHENKO	15	SND	$1.22-2.00 e^+ e^- \rightarrow \eta\pi^+\pi^-$
---------------	------------------------	----	-----	--

<sup>1</sup> From a fit to the  $e^+ e^- \rightarrow \eta\pi^+\pi^-$  cross section with vector meson dominance model including  $\rho(770)$ ,  $\rho(1450)$ , and  $\rho(1700)$  decaying exclusively via  $\eta\rho(770)$ . Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

NODE=M105R07;LINKAGE=A

 $\Gamma(a_2(1320)\pi)/\Gamma_{total}$  $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R9  
 NODE=M105R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMELIN	00	VES	$37 \pi^- \rho \rightarrow \eta\pi^+\pi^- n$
----------	--------	----	-----	--

 $\Gamma(K\bar{K})/\Gamma(\omega\pi)$  $\Gamma_{13}/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R8  
 NODE=M105R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.08$	<sup>1</sup> DONNACHIE	91	RVUE	
---------	------------------------	----	------	--

<sup>1</sup> Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

NODE=M105R8;LINKAGE=A

 $\Gamma(K\bar{K}^*(892) + c.c.)/\Gamma_{total}$  $\Gamma_{15}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R16  
 NODE=M105R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	COAN	04	CLEO	$\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$
---------------	------	----	------	---

 $\Gamma(\eta\gamma)/\Gamma_{total}$  $\Gamma_{17}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M105R06  
 NODE=M105R06

seen	35	<sup>1</sup> ACHASOV	14	SND	$1.15-2.00 e^+ e^- \rightarrow \eta\gamma$
------	----	----------------------	----	-----	--

<sup>1</sup> From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

NODE=M105R06;LINKAGE=A

 $\rho(1450)$  REFERENCES

NODE=M105

GRIBANOV	20	JHEP 2001 112	S.S. Gribanov <i>et al.</i>	(CMD-3 Collab.)	REFID=60620
ACHASOV	18	PR D97 012008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58785
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16D	PR D94 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57618
AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55912
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
		Translated from ZETFP 94 796.			
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ACHASOV	11	JETP 113 75	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=16721
		Translated from ZETF 140 87.			
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			



AKHMETSHIN 05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
ALOISIO 05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
SCHAEEL 05C	PRPL 421 191	S. Schaeel <i>et al.</i>	(ALEPH Collab.)	REFID=50845
AKHMETSHIN 04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
COAN 04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
AKHMETSHIN 03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE 01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AKHMETSHIN 01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
ALEXANDER 01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48391
AKHMETSHIN 00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANDERSON 00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)	REFID=47468
EDWARDS 00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE 99C	PL B450 275	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46916
ABELE 99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BERTIN 98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
ABELE 97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV 97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BARATE 97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45622
BERTIN 97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
BERTIN 97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45763
CLEGG 94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
BISELLO 91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=41752
DOLINSKY 91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
DONNACHIE 91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41632
FUKUI 91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41581
KUHN 90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ARMSTRONG 89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
BISELLO 89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA 89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
ANTONELLI 88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
CLEGG 88	ZPHY C40 313	A.B. Clegg, A. Donnachie	(MCHS, LANC)	REFID=40922
DIEKMANN 88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI 88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
ALBRECHT 87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40418
DONNACHIE 87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
DOLINSKY 86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246
BARKOV 85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
KURDADZE 83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
ASTON 80C	Translated from ZETFP 37 613.	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=20652
BARBER 80C	PL B2B 211	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)	REFID=20653
GOUNARIS 68	ZPHY C4 169	G.J. Gounaris, J.J. Sakurai		REFID=48054
	PRL 21 244			

$\eta(1475)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See the  $\eta(1405)$  and the related review on "Spectroscopy of Light Meson Resonances."

**$\eta(1475)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1475± 4 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
1469± 14± 13	74	ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
1460± 12	3651	NICHITIU	02 OBLX	$0 \bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1485± 8± 5	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1500± 10		CICALO	99 OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
1464± 10		BERTIN	97 OBLX	$0 \bar{p}p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
1460± 10		BERTIN	95 OBLX	$0 \bar{p}p \rightarrow K \bar{K} \pi \pi$
1490 <sup>+14+3</sup> <sub>-8-16</sub>	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1475± 4		RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$
1477± 7± 13		<sup>1</sup> ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
1565± 8 <sup>+0</sup> <sub>-63</sub>		<sup>2</sup> ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
1421± 14		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M175

NODE=M175

NODE=M175M5

NODE=M175M5

OCCUR=2


OCCUR=2

OCCUR=2

<sup>1</sup> From a fit to  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded.  
<sup>2</sup> Could also be the  $\eta(1405)$ .

NODE=M175M5;LINKAGE=B

NODE=M175M5;LINKAGE=A



**$\eta(1475)$  WIDTH**

NODE=M175W5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>90± 9 OUR AVERAGE</b>		Error includes scale factor of 1.6. See the ideogram below.		
67±18± 7	74	ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$
120±15	3651	NICHITIU	02 OBLX	0 $\bar{p}p \rightarrow K^+K^-\pi^+\pi^-\pi^0$
98±18± 3	20k	ADAMS	01B B852	18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$
100±20		CICALO	99 OBLX	0 $\bar{p}p \rightarrow K^\pm K_S^0\pi^\mp\pi^+\pi^-$
105±15		BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
105±15		BERTIN	95 OBLX	0 $\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
54 <sup>+37+13</sup> <sub>-21-24</sub>		BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm\pi^\mp$
51±13		RATH	89 MPS	21.4 $\pi^-p \rightarrow nK_S^0 K_S^0\pi^0$

NODE=M175W5

• • • We do not use the following data for averages, fits, limits, etc. • • •

118±22±17		<sup>1</sup> ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
45 <sup>+14+21</sup> <sub>-13-28</sub>		<sup>2</sup> ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0\eta$
63±18		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$


OCCUR=2

<sup>1</sup> From a fit to  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded.

NODE=M175W5;LINKAGE=B

<sup>2</sup> Could also be the  $\eta(1405)$ .

NODE=M175W5;LINKAGE=A



**eta(1475) DECAY MODES**

NODE=M175215;NODE=M175

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \bar{K} \pi$	seen
$\Gamma_2$ $K \bar{K}^*(892) + c.c.$	seen
$\Gamma_3$ $a_0(980) \pi$	seen
$\Gamma_4$ $\gamma \gamma$	seen
$\Gamma_5$ $K_S^0 K_S^0 \eta$	possibly seen
$\Gamma_6$ $\gamma \phi(1020)$	possibly seen

DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=4;OUR EST;→ UNCHECKED ←  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=8;OUR EVAL;→ UNCHECKED ←  
 DESIG=9

**eta(1475)  $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$**

NODE=M175220

$\Gamma(K \bar{K} \pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$						$\Gamma_1 \Gamma_4 / \Gamma$
VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.23±0.05±0.05</b>		74	<sup>1</sup> ACHARD	07 L3	183-209 $e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$	

NODE=M175G2  
 NODE=M175G2

••• We do not use the following data for averages, fits, limits, etc. •••  
 < 0.089      90      <sup>2,3</sup> AHOHE      05 CLE2      10.6  $e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Supersedes ACCIARRI 01G. Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6895$ .

<sup>2</sup> Using  $\eta(1475)$  mass of 1481 MeV and width of 48 MeV. The upper limit increases to 0.140 keV if the world average value, 87 MeV, of the width is used.

<sup>3</sup> Assuming three-body phase-space decay to  $K_S^0 K^\pm \pi^\mp$ .

NODE=M175G2;LINKAGE=CH  
 NODE=M175G2;LINKAGE=AH

NODE=M175G2;LINKAGE=B3

**eta(1475) BRANCHING RATIOS**

NODE=M175225

$\Gamma(K \bar{K}^*(892) + c.c.) / \Gamma(K \bar{K} \pi)$					$\Gamma_2 / \Gamma_1$
VALUE	DOCUMENT ID	TECN	COMMENT		
0.50±0.10	<sup>1</sup> BAILLON	67 HBC	0.0 $\bar{p}p \rightarrow K \bar{K} \pi \pi \pi$		

NODE=M175R1  
 NODE=M175R1

<sup>1</sup> Data could also refer to  $\eta(1405)$ .

NODE=M175R;LINKAGE=BL

$\Gamma(K \bar{K}^*(892) + c.c.) / [\Gamma(K \bar{K}^*(892) + c.c.) + \Gamma(a_0(980)\pi)]$						$\Gamma_2 / (\Gamma_2 + \Gamma_3)$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.25	90	EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$		

NODE=M175R6  
 NODE=M175R6

••• We do not use the following data for averages, fits, limits, etc. •••

$\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma_1$
$<1.27 \times 10^{-3}$	90	<sup>1</sup> ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$	

<sup>1</sup> Using results from BAI 00D.NODE=M175R01  
NODE=M175R01

NODE=M175R01;LINKAGE=A

 $\Gamma(\gamma\phi(1020))/\Gamma_{total}$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma$
<b>possibly seen</b>	<sup>1</sup> ABLIKIM	181	BES3 $J/\psi \rightarrow \gamma \gamma \phi(1020)$	

<sup>1</sup> Seen as a peak in  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded. Also see  $\eta(1405)$ .NODE=M175R00  
NODE=M175R00

NODE=M175R00;LINKAGE=A

 $\eta(1475)$  REFERENCES

ABLIKIM	181	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	180	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)

NODE=M175

REFID=58893  
REFID=58925  
REFID=56785  
REFID=51698  
REFID=50764  
REFID=48848  
REFID=48319  
REFID=49649  
REFID=47954  
REFID=47394  
REFID=45417  
REFID=44614  
REFID=41584  
REFID=41578  
REFID=40924  
REFID=21314  
REFID=20407 $f_0(1500)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M152

NODE=M152

 $f_0(1500)$  T-MATRIX POLE  $\sqrt{s}$ Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M152PP

NODE=M152PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1430–1530) – <math>i</math> (40–90) OUR ESTIMATE</b>			
$(1450 \pm 10) - i (53 \pm 8)$	<sup>1</sup> RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
$(1483 \pm 15) - i (58 \pm 6)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1496 \pm 1.2^{+4.4}_{-26.4}) - i (40.4 \pm 0.3^{+10.0}_{-2.5})$	<sup>2</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
$(1465 \pm 18) - i (50 \pm 9)$	<sup>3</sup> ROPERTZ	18	RVUE $\bar{B}_s^0 \rightarrow J/\psi(\pi^+ \pi^- / K^+ K^-)$
$(1486 \pm 10) - i (57 \pm 5)$	ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1489^{+8}_{-4}) - i (51 \pm 5)$	<sup>4</sup> ANISOVICH	03	RVUE
$(1515 \pm 12) - i (55 \pm 12)$	BARBERIS	00A	$450 \bar{p}p \rightarrow p_f(\eta\eta', \eta'\eta')p_s$
$(1511 \pm 9) - i (51 \pm 9)$	<sup>5</sup> BARBERIS	00C	$450 \bar{p}p \rightarrow p_f 4\pi p_s$
$(1510 \pm 8) - i (55 \pm 8)$	BARBERIS	00E	$450 \bar{p}p \rightarrow p_f \eta\eta p_s$
$(1502 \pm 12 \pm 10) - i (49 \pm 9 \pm 8)$	<sup>6</sup> BARBERIS	99D	OMEG $450 \bar{p}p \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1447 \pm 27) - i (54 \pm 23)$	<sup>7</sup> KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$(1499 \pm 8) - i (65 \pm 10)$	ANISOVICH	98B	RVUE Compilation.
$(1510 \pm 20) - i (60 \pm 18)$	BARBERIS	97B	OMEG $450 \bar{p}p \rightarrow p p 2(\pi^+ \pi^-)$
$(1449 \pm 20) - i (57 \pm 15)$	BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1515 \pm 20) - i (53 \pm 8)$	ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
$(1500 \pm 8) - i (66 \pm 8)$	ABELE	96C	RVUE Compilation.
$(1500 \pm 10) - i (77 \pm 15)$	<sup>8</sup> AMSLER	95D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
$(1520 \pm 25) - i (74^{+10}_{-13})$	<sup>9</sup> ANISOVICH	94	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$
$(1505 \pm 20) - i (75 \pm 10)$	<sup>10</sup> BUGG	94	RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0\pi^0$

NODE=M152PP

→ UNCHECKED ←

- <sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma \pi^0 \pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).
- <sup>2</sup> T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ), and BINON 84C ( $\eta\eta'$ ).
- <sup>3</sup> T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.
- <sup>4</sup> Pole position from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K\bar{K}n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta\eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.
- <sup>5</sup> Average between  $\pi^+ \pi^- 2\pi^0$  and  $2(\pi^+ \pi^-)$ .
- <sup>6</sup> Supersedes BARBERIS 99 and BARBERIS 99B.
- <sup>7</sup> T-matrix pole on sheet  $--+$ .
- <sup>8</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.
- <sup>9</sup> From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta$ .
- <sup>10</sup> Reanalysis of ANISOVICH 94 data.

NODE=M152PP;LINKAGE=E

NODE=M152PP;LINKAGE=A

NODE=M152PP;LINKAGE=F

NODE=M152PP;LINKAGE=G

NODE=M152PP;LINKAGE=C

NODE=M152PP;LINKAGE=H

NODE=M152PP;LINKAGE=I

NODE=M152PP;LINKAGE=D

NODE=M152PP;LINKAGE=J

NODE=M152PP;LINKAGE=K

 **$f_0(1500)$  MASS**

NODE=M152M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1522 ± 25 OUR AVERAGE</b>		[1506 ± 6 MeV OUR 2022 AVERAGE		Scale factor = 1.4]
<b>1522 ± 25</b>		<sup>1</sup> BERTIN	98 OBLX	0.05-0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$

NODE=M152M

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

1492.5 ± 3.6 <sup>+</sup> <sub>-20.5</sub>		<sup>2</sup> ABLIKIM	22G BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
1447 ± 16 ± 13	163	<sup>3,4</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1442 ± 9 ± 4	261	<sup>3,4</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
1460.9 ± 2.9		<sup>5</sup> AAIJ	14BR LHCB	$\bar{B}_S^0 \rightarrow J/\psi \pi^+ \pi^-$
1468 <sup>+14</sup> <sub>-15</sub> <sup>+23</sup> <sub>-74</sub>	5.5k	<sup>6</sup> ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1470 ± 60	568	<sup>7</sup> KLEMP	08 E791	$D_S^+ \rightarrow \pi^- \pi^+ \pi^+$
1470 <sup>+6</sup> <sub>-7</sub> <sup>+72</sup> <sub>-255</sub>		<sup>8</sup> UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1466 ± 6 ± 20		<sup>9</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1495 ± 4		AMSLER	06 CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
1539 ± 20	9.9k	AUBERT	06O BABR	$B^+ \rightarrow K^+ K^+ K^-$
1473 ± 5	80k	<sup>9,10</sup> UMAN	06 E835	$5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
1478 ± 6		VLADIMIRSK...	06 SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1493 ± 7		<sup>9</sup> BINON	05 GAMS	$33 \pi^- p \rightarrow \eta \eta n$
1524 ± 14	1400	<sup>11</sup> GARMASH	05 BELL	$B^+ \rightarrow K^+ K^+ K^-$
1490 ± 30		<sup>9</sup> ABELE	01 CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1497 ± 10		<sup>9</sup> BARBERIS	99 OMEG	$450 pp \rightarrow p_S p_f K^+ K^-$
1502 ± 10		<sup>9</sup> BARBERIS	99B OMEG	$450 pp \rightarrow p_S p_f \pi^+ \pi^-$
1530 ± 45		<sup>9</sup> BELLAZZINI	99 GAM4	$450 pp \rightarrow p p \pi^0 \pi^0$
1505 ± 18		<sup>9</sup> FRENCH	99	$300 pp \rightarrow p_f (K^+ K^-) p_S$
1580 ± 80		<sup>9</sup> ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
~ 1520		REYES	98 SPEC	$800 pp \rightarrow p_S p_f K_S^0 K_S^0$
~ 1475		FRABETTI	97D E687	$D_S^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 1505		ABELE	96 CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
1460 ± 20	120	<sup>9</sup> AMELIN	96B VES	$37 \pi^- A \rightarrow \eta \eta \pi^- A$
1500 ± 8		BUGG	96 RVUE	
1500 ± 15		<sup>12</sup> AMSLER	95B CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
1505 ± 15		<sup>13</sup> AMSLER	95C CBAR	$0.0 \bar{p}p \rightarrow \eta \eta \pi^0$
1445 ± 5		<sup>14</sup> ANTINORI	95 OMEG	$300,450 pp \rightarrow p p 2(\pi^+ \pi^-)$
1497 ± 30		<sup>9</sup> ANTINORI	95 OMEG	$300,450 pp \rightarrow p p \pi^+ \pi^-$
~ 1505		BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1446 ± 5		<sup>9</sup> ABATZIS	94 OMEG	$450 pp \rightarrow p p 2(\pi^+ \pi^-)$
1545 ± 25		<sup>9</sup> AMSLER	94E CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta \eta'$
1560 ± 25		<sup>9</sup> AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta \eta$
1550 ± 45 ± 30		<sup>9</sup> BELADIDZE	92C VES	$36 \pi^- Be \rightarrow \pi^- \eta' \eta Be$
1449 ± 4		<sup>9</sup> ARMSTRONG	89E OMEG	$300 pp \rightarrow p p 2(\pi^+ \pi^-)$

OCCUR=2

OCCUR=2

OCCUR=2

1610 ± 20		<sup>9</sup> ALDE	88	GAM4	300	$\pi^- N \rightarrow \pi^- N 2\eta$
~ 1525		ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1570 ± 20	600	<sup>9</sup> ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$
1575 ± 45		<sup>15</sup> ALDE	86D	GAM4	100	$\pi^- p \rightarrow 2\eta n$
1568 ± 33		<sup>9</sup> BINON	84C	GAM2	38	$\pi^- p \rightarrow \eta\eta' n$
1592 ± 25		<sup>9</sup> BINON	83	GAM2	38	$\pi^- p \rightarrow 2\eta n$
1525 ± 5		<sup>9</sup> GRAY	83	DBC	0.0	$\bar{p} N \rightarrow 3\pi$

<sup>1</sup> Breit-Wigner mass.

<sup>2</sup> The  $\pi^+\pi^-$  mass spectrum is described by a coherent sum of two Breit-Wigner resonances,  $f_0(1500)$  and a new  $X(1540)$  with mass  $1540.2 \pm 7.0^{+36.3}_{-6.1}$  MeV and width  $157 \pm 19^{+11}_{-77}$  MeV.

<sup>3</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>4</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 109$  MeV.

<sup>5</sup> Solution I, statistical error only.

<sup>6</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>7</sup> Reanalysis of AITALA 01A data. This state could also be  $f_0(1370)$ .

<sup>8</sup> Breit-Wigner mass. May also be the  $f_0(1370)$ .

<sup>9</sup> Breit-Wigner mass.

<sup>10</sup> Statistical error only.

<sup>11</sup> Breit-Wigner, solution 1, PWA ambiguous.

<sup>12</sup> T-matrix pole, supersedes ANISOVICH 94.

<sup>13</sup> T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

<sup>14</sup> Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

<sup>15</sup> From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152M;LINKAGE=PP  
NODE=M152M;LINKAGE=M

NODE=M152M;LINKAGE=F  
NODE=M152M;LINKAGE=G  
NODE=M152M;LINKAGE=I  
NODE=M152M;LINKAGE=C

NODE=M152M;LINKAGE=KL  
NODE=M152M;LINKAGE=UE  
NODE=M152M;LINKAGE=E  
NODE=M152M;LINKAGE=ST  
NODE=M152M;LINKAGE=GA  
NODE=M152M;LINKAGE=D  
NODE=M152M;LINKAGE=D1  
NODE=M152M;LINKAGE=B  
NODE=M152M;LINKAGE=AZ

### $f_0(1500)$ WIDTH

NODE=M152W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>108 ± 33 OUR AVERAGE</b>		[112 ± 9 MeV OUR 2022 AVERAGE]		
<b>108 ± 33</b>		<sup>1</sup> BERTIN	98	OBLX 0.05–0.405 $\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
107 ± 9 <sup>+</sup> <sub>7</sub>		<sup>2</sup> ABLIKIM	22G	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
124 ± 7		<sup>3</sup> AAIJ	14BR	LHCB $\bar{B}_s^0 \rightarrow J/\psi\pi^+\pi^-$
136 <sup>+</sup> <sub>26</sub> 41 <sup>+</sup> <sub>100</sub>	5.5k	<sup>4</sup> ABLIKIM	13N	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
90 <sup>+</sup> <sub>1</sub> 2 <sup>+</sup> <sub>22</sub>		<sup>5</sup> UEHARA	08A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
108 <sup>+</sup> <sub>11</sub> ± 25		<sup>6</sup> ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
121 ± 8		AMSLER	06	CBAR 0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
257 ± 33	9.9k	AUBERT	06O	BABR $B^+ \rightarrow K^+K^+K^-$
108 ± 9	80k	<sup>6,7</sup> UMAN	06	E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
119 ± 10		VLADIMIRSK...	06	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
90 ± 15		<sup>6</sup> BINON	05	GAMS 33 $\pi^- p \rightarrow \eta\eta n$
136 ± 23	1400	<sup>8</sup> GARMASH	05	BELL $B^+ \rightarrow K^+K^+K^-$
140 ± 40		<sup>6</sup> ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
104 ± 25		<sup>6</sup> BARBERIS	99	OMEG 450 $pp \rightarrow p_s p_f K^+ K^-$
131 ± 15		<sup>6</sup> BARBERIS	99B	OMEG 450 $pp \rightarrow p_s p_f \pi^+ \pi^-$
160 ± 50		<sup>6</sup> BELLAZZINI	99	GAM4 450 $pp \rightarrow pp\pi^0\pi^0$
100 ± 33		<sup>6</sup> FRENCH	99	300 $pp \rightarrow p_f(K^+K^-)p_s$
280 ± 100		<sup>6</sup> ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0\pi^0 n$
~ 100		FRABETTI	97D	E687 $D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 169		ABELE	96	CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$
100 ± 30	120	<sup>6</sup> AMELIN	96B	VES 37 $\pi^- A \rightarrow \eta\eta\pi^- A$
132 ± 15		BUGG	96	RVUE
120 ± 25		<sup>9</sup> AMSLER	95B	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0$
120 ± 30		<sup>10</sup> AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
65 ± 10		<sup>11</sup> ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+\pi^-)$

NODE=M152W  
NEW

OCCUR=2

199 ± 30		<sup>6</sup> ANTINORI	95	OMEG	300,450	$p p \rightarrow p p \pi^+ \pi^-$	OCCUR=2
56 ± 12		<sup>6</sup> ABATZIS	94	OMEG	450	$p p \rightarrow p p 2(\pi^+ \pi^-)$	
100 ± 40		<sup>6</sup> AMSLER	94E	CBAR	0.0	$\bar{p} p \rightarrow \pi^0 \eta \eta'$	
245 ± 50		<sup>6</sup> AMSLER	92	CBAR	0.0	$\bar{p} p \rightarrow \pi^0 \eta \eta$	
153 ± 67 ± 50		<sup>6</sup> BELADIDZE	92C	VES	36	$\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$	
78 ± 18		<sup>6</sup> ARMSTRONG	89E	OMEG	300	$p p \rightarrow p p 2(\pi^+ \pi^-)$	
170 ± 40		<sup>6</sup> ALDE	88	GAM4	300	$\pi^- N \rightarrow \pi^- N 2\eta$	
150 ± 20	600	<sup>6</sup> ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$	
265 ± 65		<sup>12</sup> ALDE	86D	GAM4	100	$\pi^- p \rightarrow 2\eta n$	
260 ± 60		<sup>6</sup> BINON	84C	GAM2	38	$\pi^- p \rightarrow \eta \eta' n$	
210 ± 40		<sup>6</sup> BINON	83	GAM2	38	$\pi^- p \rightarrow 2\eta n$	
101 ± 13		<sup>6</sup> GRAY	83	DBC	0.0	$\bar{p} N \rightarrow 3\pi$	

<sup>1</sup> Breit-Wigner width.

<sup>2</sup> The  $\pi^+ \pi^-$  mass spectrum is described by a coherent sum of two Breit-Wigner resonances,  $f_0(1500)$  and a new  $X(1540)$  with mass  $1540.2 \pm 7.0^{+36.3}_{-6.1}$  MeV and width  $157 \pm 19^{+11}_{-77}$  MeV.

<sup>3</sup> Solution 1, statistical error only.

<sup>4</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>5</sup> Breit-Wigner width. May also be the  $f_0(1370)$ .

<sup>6</sup> Breit-Wigner width.

<sup>7</sup> Statistical error only.

<sup>8</sup> Breit-Wigner, solution 1, PWA ambiguous.

<sup>9</sup> T-matrix pole, supersedes ANISOVICH 94.

<sup>10</sup> T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

<sup>11</sup> Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

<sup>12</sup> From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152W;LINKAGE=PP  
NODE=M152W;LINKAGE=J

NODE=M152W;LINKAGE=G  
NODE=M152W;LINKAGE=C

NODE=M152W;LINKAGE=UE  
NODE=M152W;LINKAGE=E  
NODE=M152W;LINKAGE=ST  
NODE=M152W;LINKAGE=GA  
NODE=M152W;LINKAGE=D  
NODE=M152W;LINKAGE=D1  
NODE=M152W;LINKAGE=B  
NODE=M152W;LINKAGE=AZ

### $f_0(1500)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $\pi \pi$	(34.5 ± 2.2) %	1.2
$\Gamma_2$ $\pi^+ \pi^-$	seen	
$\Gamma_3$ $2\pi^0$	seen	
$\Gamma_4$ $4\pi$	(48.9 ± 3.3) %	1.2
$\Gamma_5$ $4\pi^0$	seen	
$\Gamma_6$ $2\pi^+ 2\pi^-$	seen	
$\Gamma_7$ $2(\pi\pi)S\text{-wave}$	seen	
$\Gamma_8$ $\rho\rho$	seen	
$\Gamma_9$ $\pi(1300)\pi$	seen	
$\Gamma_{10}$ $a_1(1260)\pi$	seen	
$\Gamma_{11}$ $\eta\eta$	( 6.0 ± 0.9) %	1.1
$\Gamma_{12}$ $\eta\eta'(958)$	( 2.2 ± 0.8) %	1.4
$\Gamma_{13}$ $K\bar{K}$	( 8.5 ± 1.0) %	1.1
$\Gamma_{14}$ $\gamma\gamma$	not seen	

NODE=M152215;NODE=M152

DESIG=8  
DESIG=9  
DESIG=3;OUR EST;→ UNCHECKED ←  
DESIG=7  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←  
DESIG=11;OUR EST;→ UNCHECKED ←  
DESIG=12;OUR EST;→ UNCHECKED ←  
DESIG=13;OUR EST;→ UNCHECKED ←  
DESIG=14;OUR EST;→ UNCHECKED ←  
DESIG=1  
DESIG=2  
DESIG=4  
DESIG=10;OUR EST;→ UNCHECKED ←

### CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 10 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 5.6$  for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_4$	-88			
$x_{11}$	27	-56		
$x_{12}$	3	-32	26	
$x_{13}$	43	-64	20	2
	$x_1$	$x_4$	$x_{11}$	$x_{12}$

$f_0(1500) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M152217

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_{14}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M152G1  
NODE=M152G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$33^{+12}_{-6} + 1809_{-21}$		<sup>1</sup> UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
not seen		ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$
<460	95	BARATE	00E ALEP	$\gamma\gamma \rightarrow \pi^+ \pi^-$

<sup>1</sup> May also be the  $f_0(1370)$ . Multiplied by us by 3 to obtain the  $\pi\pi$  value.

NODE=M152G1;LINKAGE=UE

 $f_0(1500) \text{ BRANCHING RATIOS}$ 

NODE=M152220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN
-------	-------------	------

NODE=M152R8  
NODE=M152R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.454±0.104	BUGG	96 RVUE
-------------	------	---------

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R10  
NODE=M152R10seen BERTIN 98 OBLX 0.05-0.405  $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen FRABETTI 97D E687  $D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$  $\Gamma(4\pi)/\Gamma(\pi\pi)$  $\Gamma_4/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R6  
NODE=M152R6**1.42±0.18 OUR FIT** Error includes scale factor of 1.2.**1.42±0.18 OUR AVERAGE** Error includes scale factor of 1.2.1.37±0.16 BARBERIS 00D 450  $p\bar{p} \rightarrow p_f 4\pi p_s$ 2.1 ±0.6 <sup>1</sup> AMSLER 98 RVUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ±0.2 <sup>2</sup> ANISOVICH 02D SPEC Combined fit3.4 ±0.8 <sup>1</sup> ABELE 96 CBAR 0.0  $\bar{p}p \rightarrow 5\pi^0$ <sup>1</sup> Excluding  $\rho\rho$  contribution to  $4\pi$ .<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$ ), GAMS ( $\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$ ), and BNL ( $\pi p \rightarrow K \bar{K} n$ ) data.NODE=M152R6;LINKAGE=C  
NODE=M152R6;LINKAGE=CH $\Gamma(2(\pi\pi)\text{s-wave})/\Gamma(\pi\pi)$  $\Gamma_7/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R14  
NODE=M152R14

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.42±0.26 <sup>1</sup> ABELE 01 CBAR 0.0  $\bar{p}d \rightarrow \pi^- 4\pi^0 p$ <sup>1</sup> From the combined data of ABELE 96 and ABELE 96C.

NODE=M152R;LINKAGE=KZ

 $\Gamma(2(\pi\pi)\text{s-wave})/\Gamma(4\pi)$  $\Gamma_7/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R15  
NODE=M152R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26±0.07 ABELE 01B CBAR 0.0  $\bar{p}d \rightarrow 5\pi p$  $\Gamma(\rho\rho)/\Gamma(4\pi)$  $\Gamma_8/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R16  
NODE=M152R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13±0.08 ABELE 01B CBAR 0.0  $\bar{p}d \rightarrow 5\pi p$  $\Gamma(\rho\rho)/\Gamma(2(\pi\pi)\text{s-wave})$  $\Gamma_8/\Gamma_7$ 

VALUE	DOCUMENT ID	COMMENT
-------	-------------	---------

NODE=M152R11  
NODE=M152R11**2.87±0.34 OUR AVERAGE** Error includes scale factor of 1.1.3.3 ±0.5 BARBERIS 00C 450  $p\bar{p} \rightarrow p_f \pi^+ \pi^- 2\pi^0 p_s$ 2.6 ±0.4 BARBERIS 00C 450  $p\bar{p} \rightarrow p_f 2(\pi^+ \pi^-) p_s$ 

OCCUR=2

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$  $\Gamma_9/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R17  
NODE=M152R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.50±0.25 ABELE 01B CBAR 0.0  $\bar{p}d \rightarrow 5\pi p$



$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$  $\Gamma_{10}/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R18  
 NODE=M152R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.12±0.05	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$
-----------	-------	-----	--

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R1  
 NODE=M152R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

large	ALDE	88	GAM4 300 $\pi^- N \rightarrow \eta\eta\pi^- N$
large	BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$  $\Gamma_{11}/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R13  
 NODE=M152R13

**0.173±0.024 OUR FIT** Error includes scale factor of 1.1.

**0.175±0.027 OUR AVERAGE**

0.18 ±0.03	BARBERIS	00E	450 $pp \rightarrow p_f\eta\eta p_s$
0.157±0.060	<sup>1</sup> AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.080±0.033	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
0.11 ±0.03	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit
0.078±0.013	<sup>3</sup> ABELE	96C	RVUE Compilation
0.230±0.097	<sup>4</sup> AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$

<sup>1</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

<sup>3</sup>  $2\pi$  width determined to be  $60 \pm 12$  MeV.

<sup>4</sup> Using AMSLER 95B ( $3\pi^0$ ).

NODE=M152R3;LINKAGE=AB  
 NODE=M152R;LINKAGE=CH

NODE=M152R3;LINKAGE=CM  
 NODE=M152R3;LINKAGE=A

 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$  $\Gamma_5/\Gamma_{11}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R5  
 NODE=M152R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.8±0.3	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$
---------	------	----	---

 $\Gamma(\eta\eta'(958))/\Gamma(\pi\pi)$  $\Gamma_{12}/\Gamma_1$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M152R12  
 NODE=M152R12

**6.4 ±2.2 OUR FIT** Error includes scale factor of 1.4.

<b>9.5 ±2.6</b>	BARBERIS	00A	450 $pp \rightarrow p_f\eta\eta p_s$
-----------------	----------	-----	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.96 <sup>+2.95</sup> <sub>-2.87</sub>	<sup>1</sup> ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
0.5 ±0.3	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$  P-wave. The quoted measurement will be corrected in a forthcoming erratum.

<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

NODE=M152R12;LINKAGE=A

NODE=M152R12;LINKAGE=CH

 $\Gamma(\eta\eta'(958))/\Gamma(\eta\eta)$  $\Gamma_{12}/\Gamma_{11}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R2  
 NODE=M152R2

**0.37±0.13 OUR FIT** Error includes scale factor of 1.5.

<b>0.29±0.10</b>	<sup>1</sup> AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
------------------	---------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.05±0.03	<sup>2</sup> ANISOVICH	02D	SPEC Combined fit
0.84±0.23	ABELE	96C	RVUE Compilation
2.7 ±0.8	BINON	84C	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

<sup>1</sup> Using AMSLER 94E ( $\eta\eta'\pi^0$ ).

<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

NODE=M152R2;LINKAGE=A  
 NODE=M152R2;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R9  
 NODE=M152R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.044±0.021	BUGG	96	RVUE
-------------	------	----	------

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$  $\Gamma_{13}/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.246±0.025 OUR FIT****0.236±0.026 OUR AVERAGE**

0.25 ±0.03	<sup>1</sup> BARGIOTTI	03	OBLX $\bar{p}p$
0.19 ±0.07	<sup>2</sup> ABELE	98	CBAR $0.0 \bar{p}p \rightarrow K^0 K^\pm \pi^\mp$
0.20 ±0.08	<sup>3</sup> ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K^0_L K^0_L$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.16 ±0.05	<sup>4</sup> ANISOVICH	02D	SPEC Combined fit
0.33 ±0.03 ±0.07	BARBERIS	99D	OMEG $450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$

<sup>1</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ , and  $K^\pm K^0_S \pi^\mp$ .

<sup>2</sup> Using  $\pi^0 \pi^0$  from AMSLER 95B.

<sup>3</sup> Using AMSLER 95B ( $3\pi^0$ ), AMSLER 94C ( $2\pi^0 \eta$ ) and SU(3).

<sup>4</sup> From a combined K-matrix analysis of Crystal Barrel ( $0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ), GAMS ( $\pi p \rightarrow \pi^0 \pi^0 n$ ,  $\eta \eta n$ ,  $\eta \eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

NODE=M152R7  
NODE=M152R7

NODE=M152R;LINKAGE=BG  
NODE=M152R7;LINKAGE=A  
NODE=M152R7;LINKAGE=D  
NODE=M152R7;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma(\eta\eta)$  $\Gamma_{13}/\Gamma_{11}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**1.43±0.24 OUR FIT** Error includes scale factor of 1.1.**1.85±0.41** BARBERIS 00E 450  $pp \rightarrow p_f \eta \eta p_s$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ±0.6	<sup>1</sup> ANISOVICH	02D	SPEC Combined fit
<0.4	90	<sup>2</sup> PROKOSHKIN	91 GAM4 300 $\pi^- p \rightarrow \pi^- p \eta \eta$
<0.6	<sup>3</sup> BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$

<sup>1</sup> From a combined K-matrix analysis of Crystal Barrel ( $0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ), GAMS ( $\pi p \rightarrow \pi^0 \pi^0 n$ ,  $\eta \eta n$ ,  $\eta \eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

<sup>2</sup> Combining results of GAM4 with those of WA76 on  $K\bar{K}$  central production.

<sup>3</sup> Using ETKIN 82B and COHEN 80.

NODE=M152R4  
NODE=M152R4

NODE=M152R4;LINKAGE=CH

NODE=M152R4;LINKAGE=BZ  
NODE=M152R4;LINKAGE=A

 $f_0(1500)$  REFERENCES

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
ABLIKIM	22G	PRL 129 042001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61642
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ROPERTZ	18	EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis	(BONN, JULI)	REFID=59332
AAIJ	17V	JHEP 1708 037	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57828
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
AAIJ	14BR	PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56035
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
KLEMP	08	EPJ C55 39	E. Klempt, M. Matveev, A.V. Sarantsev	(BONN+)	REFID=52286
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
AUBERT	06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51141
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00D	PL B474 423	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47960
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
AMSLER	98	RMP 70 1293	C. Amsler		REFID=46601
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			

NODE=M152

BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
REYES	98	PRL 81 4079	M.A. Reyes <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMELIN	96B	PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
Translated from YAF 59 1021.				
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94E	PL B340 259	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)
Translated from YAF 55 2748.				
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)
Translated from DANS 316 900.				
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ALDE	88	PL B201 160	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
Also				
SJNP 38 561				
Translated from YAF 38 934.				
GRAY	83	PR D27 307	L. Gray <i>et al.</i>	(SYRA)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)

REFID=45782  
 REFID=46378  
 REFID=45758  
 REFID=45701  
 REFID=45554  
 REFID=45011  
 REFID=45038  
 REFID=45076  
 REFID=44725  
 REFID=45094  
 REFID=44377  
 REFID=44440  
 REFID=44441  
 REFID=44437  
 REFID=44438  
 REFID=44090  
 REFID=44091  
 REFID=44093  
 REFID=44080  
 REFID=43659  
 REFID=44078  
 REFID=43169  
 REFID=43175  
 REFID=41719  
 REFID=41011  
 REFID=40283  
 REFID=40330  
 REFID=40221  
 REFID=20765  
 REFID=20768  
 REFID=21418  
 REFID=20750  
 REFID=20751  
 REFID=21370  
 REFID=20390  
 REFID=20381  
 REFID=20355


**$f_1(1510)$**

$$I^G(J^{PC}) = 0^+(1^{++})$$

OMITTED FROM SUMMARY TABLE  
 See the review on "Spectroscopy of Light Meson Resonances."

**$f_1(1510)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1518 ± 5 OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.		
1530 ± 10		ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
1512 ± 4	600	<sup>1</sup> BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1526 ± 6	271	GAVILLET	82 HBC	4.2 $K^- p \rightarrow \Lambda K K \pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1525		<sup>2</sup> BAUER	93B	$\gamma \gamma^* \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
<sup>1</sup> From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.				
<sup>2</sup> Not seen by AIHARA 88C in the $K_S^0 K^\pm \pi^\mp$ final state.				



NODE=M084  
 NODE=M084  
 NODE=M084M;LINKAGE=A  
 NODE=M084M;LINKAGE=C

### $f_1(1510)$ WIDTH


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>73±25 OUR AVERAGE</b>	Error includes scale factor of 2.5. See the ideogram below.			
100±40		ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
35±15	600	<sup>3</sup> BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
107±15	271	GAVILLET	82 HBC	4.2 $K^- p \rightarrow \Lambda K K \pi$

<sup>3</sup>From partial wave analysis of  $K^+ \bar{K}^0 \pi^-$  state.

NODE=M084W

NODE=M084W

NODE=M084W;LINKAGE=A



### $f_1(1510)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \bar{K}^*(892) + c.c.$	seen
$\Gamma_2$ $\pi^+ \pi^- \eta'$	seen

NODE=M084215;NODE=M084

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=2

### $f_1(1510)$ BRANCHING RATIOS

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{total}$	$\Gamma_2/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	230	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M084225

NODE=M084R01  
NODE=M084R01

### $f_1(1510)$ REFERENCES

ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BAUER	93B	PR D48 3976	D.A. Bauer <i>et al.</i>	(SLAC)
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP
GAVILLET	82	ZPHY C16 119	P. Gavillet <i>et al.</i>	(CERN, CDEF, PADO+)

NODE=M084

REFID=53684  
REFID=43678  
REFID=40564  
REFID=40282  
REFID=40568  
REFID=20877

NODE=M013

**$f'_2(1525)$**


$$I^G(J^{PC}) = 0^+(2^{++})$$

**$f'_2(1525)$  MASS**

NODE=M013MX

NODE=M013MX

VALUE (MeV)                      DOCUMENT ID  
**1517.4±2.5 OUR AVERAGE** Includes data from the 6 datablocks that follow this one.  
 Error includes scale factor of 2.8. See the ideogram below.



$f'_2(1525)$  MASS (MeV)

**PRODUCED BY PION BEAM**

NODE=M013M1  
 NODE=M013M1

VALUE (MeV)    EVTS    DOCUMENT ID    TECN    COMMENT

The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

1521±13		TIKHOMIROV	03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1547 <sup>+10</sup> <sub>-2</sub>		<sup>1</sup> LONGACRE	86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1496 <sup>+9</sup> <sub>-8</sub>		<sup>2</sup> CHABAUD	81	ASPK	6 $\pi^- p \rightarrow K^+ K^- n$
1497 <sup>+8</sup> <sub>-9</sub>		CHABAUD	81	ASPK	18.4 $\pi^- p \rightarrow K^+ K^- n$
1492±29		GORLICH	80	ASPK	17 $\pi^- p$ polarized $\rightarrow K^+ K^- n$
1502±25		<sup>3</sup> CORDEN	79	OMEG	12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
1480	14	CRENNELL	66	HBC	6.0 $\pi^- p \rightarrow K_S^0 K_S^0 n$

OCCUR=2

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

<sup>2</sup> CHABAUD 81 is a reanalysis of PAWLICKI 77 data.

<sup>3</sup> From an amplitude analysis where the  $f'_2(1525)$  width and elasticity are in complete disagreement with the values obtained from  $K\bar{K}$  channel, making the solution dubious.

NODE=M013M;LINKAGE=L  
 NODE=M013M;LINKAGE=D  
 NODE=M013M;LINKAGE=N

**PRODUCED BY  $K^\pm$  BEAM**

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  
 The data in this block is included in the average printed for a previous datablock.

NODE=M013M2  
 NODE=M013M2


**1518.1 ± 2.8 OUR AVERAGE** Includes data from the datablock that follows this one. Error includes scale factor of 3.0. See the ideogram below.

1526.8 ± 4.3		ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1504 ± 12		BOLONKIN	86	SPEC	40	$K^- p \rightarrow K_S^0 K_S^0 Y$
1529 ± 3		ARMSTRONG	83B	OMEG	18.5	$K^- p \rightarrow K^- K^+ \Lambda$
1521 ± 6	650	AGUILAR-...	81B	HBC	4.2	$K^- p \rightarrow \Lambda K^+ K^-$
1521 ± 3	572	ALHARRAN	81	HBC	8.25	$K^- p \rightarrow \Lambda K \bar{K}$
1522 ± 6	123	BARREIRO	77	HBC	4.15	$K^- p \rightarrow \Lambda K_S^0 K_S^0$
1528 ± 7	166	EVANGELIS...	77	OMEG	10	$K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1527 ± 3	120	BRANDENB...	76C	ASPK	13	$K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1519 ± 7	100	AGUILAR-...	72B	HBC	3.9,4.6	$K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$

• • • We do not use the following data for averages, fits, limits, etc. • • •  
 1514 ± 8            61        BINON        07        GAMS    32.5     $K^- p \rightarrow \eta \eta (\Lambda / \Sigma^0)$   
 1513 ± 10           1 BARKOV    99        SPEC    40        $K^- p \rightarrow K_S^0 K_S^0 y$

<sup>1</sup> Systematic errors not estimated.

NODE=M013M2;LINKAGE=SK



PRODUCED BY  $K^\pm$  BEAM (MeV)

**PRODUCED IN  $e^+ e^-$  ANNIHILATION AND PARTICLE DECAYS**

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  
 The data in this block is included in the average printed for a previous datablock.

NODE=M013M3  
 NODE=M013M3


**1514  $\pm \frac{5}{4}$  OUR AVERAGE** Error includes scale factor of 3.8. See the ideogram below.

1522.2 ± 2.8 $\frac{+}{-} \frac{5.3}{2.0}$		AAIJ	13AN	LHCB		$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
1513 ± 5 $\frac{+}{-} \frac{4}{10}$	5.5k	<sup>1</sup> ABLIKIM	13N	BES3		$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1525.3 $\frac{+}{-} \frac{1.2+3.7}{1.4-2.1}$		UEHARA	13	BELL		$\gamma \gamma \rightarrow K_S^0 K_S^0$
1521 ± 5		ABLIKIM	05	BES2		$J/\psi \rightarrow \phi K^+ K^-$
1518 ± 1 ± 3		ABE	04	BELL		$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1519 ± 2 $\frac{+15}{-5}$		BAI	03G	BES		$J/\psi \rightarrow \gamma K \bar{K}$
1523 ± 6	331	<sup>2</sup> ACCIARRI	01H	L3		$91, 183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$

1535 ± 5 ± 4	ABREU	96C	DLPH	$Z^0 \rightarrow K^+ K^- + X$	
1516 ± 5 $\begin{smallmatrix} +9 \\ -15 \end{smallmatrix}$	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$	
1531.6 ± 10.0	AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$	
1496 ± 2	<sup>3</sup> FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$	OCCUR=2
1525 ± 10 ± 10	BALTRUSAIT..87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1518 ± 3	<sup>4</sup> KLEMP	22	RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0,$ $\gamma K_S^0 K_S^0$	
1503 ± 11	<sup>5</sup> RODAS	22	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$	
1532 ± 3 ± 6	644 <sup>6,7</sup> DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$	
1557 ± 9 ± 3	113 <sup>6,7</sup> DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$	OCCUR=2
1526 ± 7	29 <sup>8</sup> LEES	14H	BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$	
1523 ± 5	870 <sup>9</sup> SCHEGELSKY	06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
1515 ± 5	<sup>10</sup> FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-$	

- <sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}, 2^{++},$  and  $4^{++}$  resonances.
- <sup>2</sup> Supersedes ACCIARRI 95J.
- <sup>3</sup> From an analysis including interference with  $f_0(1710)$ .
- <sup>4</sup> Fit of the tensor partial waves from BES3 in the multipole basis.
- <sup>5</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma \pi^0 \pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).
- <sup>6</sup> Using CLEO-c data but not authored by the CLEO Collaboration.
- <sup>7</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 73$  MeV.
- <sup>8</sup> From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated.
- <sup>9</sup> From analysis of L3 data at 91 and 183–209 GeV.
- <sup>10</sup> From an analysis ignoring interference with  $f_0(1710)$ .

NODE=M013M3;LINKAGE=A  
 NODE=M013M;LINKAGE=HA  
 NODE=M013M;LINKAGE=F2  
 NODE=M013M3;LINKAGE=F  
 NODE=M013M3;LINKAGE=E  
 NODE=M013M3;LINKAGE=B  
 NODE=M013M3;LINKAGE=C  
 NODE=M013M3;LINKAGE=D  
 NODE=M013M3;LINKAGE=SC  
 NODE=M013M;LINKAGE=F1



**PRODUCED IN  $\bar{p}p$  ANNIHILATION**

VALUE (MeV) DOCUMENT ID TECN COMMENT  
 The data in this block is included in the average printed for a previous datablock.

**1512 ± 4 OUR AVERAGE**

1513 ± 4	AMSLER	06	CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
1508 ± 9	<sup>1</sup> AMSLER	02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1495.0 ± 1.1 ± 8.1	<sup>2</sup> ALBRECHT	20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
1530 ± 12	<sup>3</sup> ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$

- <sup>1</sup> T-matrix pole.
- <sup>2</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).
- <sup>3</sup> 4-poles, 5-channel K matrix fit.

NODE=M013M;LINKAGE=TT  
 NODE=M013M9;LINKAGE=A  
 NODE=M013M9;LINKAGE=AN

**CENTRAL PRODUCTION**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

**1515±15**BARBERIS 99 OMEG 450  $pp \rightarrow p_s p_f K^+ K^-$ NODE=M013M4  
NODE=M013M4**PRODUCED IN  $e p$  COLLISIONS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M013M10  
NODE=M013M10**1512±3<sup>+1.4</sup><sub>-0.5</sub>**<sup>1</sup> CHEKANOV 08 ZEUS  $e p \rightarrow K_S^0 K_S^0 X$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

1537<sup>+9</sup><sub>-8</sub>84 <sup>2</sup> CHEKANOV 04 ZEUS  $e p \rightarrow K_S^0 K_S^0 X$ 

<sup>1</sup> In the SU(3) based model with a specific interference pattern of the  $f_2(1270)$ ,  $a_2^0(1320)$ , and  $f_2'(1525)$  mesons incoherently added to the  $f_0(1710)$  and non-resonant background.

NODE=M013M10;LINKAGE=HE

<sup>2</sup> Systematic errors not estimated.

NODE=M013M10;LINKAGE=CH

 **$f_2'(1525)$  WIDTH**

NODE=M013WX

VALUE (MeV)	DOCUMENT ID	COMMENT
-------------	-------------	---------

**86 ±5 OUR FIT** Error includes scale factor of 2.2.

NODE=M013WX

**86.9<sup>+2.3</sup><sub>-2.1</sub>**

PDG 18 Average of width measurements

**PRODUCED BY PION BEAM**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M013W1  
NODE=M013W1**86.9<sup>+2.3</sup><sub>-2.1</sub> OUR AVERAGE** Includes data from the 5 datablocks that follow this one.

Error includes scale factor of 1.4. See the ideogram below.

• • • We do not use the following data for averages, fits, limits, etc. • • •

102 ±42

TIKHOMIROV 03 SPEC 40.0  $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$ 108 <sup>+5</sup><sub>-2</sub><sup>1</sup> LONGACRE 86 MPS 22  $\pi^- p \rightarrow K_S^0 K_S^0 n$ 69 <sup>+22</sup><sub>-16</sub><sup>2</sup> CHABAUD 81 ASPK 6  $\pi^- p \rightarrow K^+ K^- n$ 137 <sup>+23</sup><sub>-21</sub>CHABAUD 81 ASPK 18.4  $\pi^- p \rightarrow K^+ K^- n$ 

OCCUR=2

150 <sup>+83</sup><sub>-50</sub>GORLICH 80 ASPK 17  $\pi^- p$  polarized  $\rightarrow K^+ K^- n$ 

165 ±42

<sup>3</sup> CORDEN 79 OMEG 12-15  $\pi^- p \rightarrow \pi^+ \pi^- n$ 92 <sup>+39</sup><sub>-22</sub><sup>4</sup> POLYCHRO... 79 STRC 7  $\pi^- p \rightarrow n K_S^0 K_S^0$ 

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

NODE=M013W;LINKAGE=L

<sup>2</sup> CHABAUD 81 is a reanalysis of PAWLICKI 77 data.

NODE=M013W;LINKAGE=D


<sup>3</sup> From an amplitude analysis where the  $f_2'(1525)$  width and elasticity are in complete disagreement with the values obtained from  $K\bar{K}$  channel, making the solution dubious.

NODE=M013W;LINKAGE=N

<sup>4</sup> From a fit to the  $D$  with  $f_2(1270)$ - $f_2'(1525)$  interference. Mass fixed at 1516 MeV.

NODE=M013W;LINKAGE=M





**PRODUCED BY  $K^\pm$  BEAM**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M013W2  
NODE=M013W2

The data in this block is included in the average printed for a previous datablock.

**82± 6 OUR AVERAGE**

90±12		ASTON	88D LASS	11 $K^- p \rightarrow K_S^0 K_S^0 \Lambda$
73±18		BOLONKIN	86 SPEC	40 $K^- p \rightarrow K_S^0 K_S^0 \gamma$
83±15		ARMSTRONG	83B OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$
85±16	650	AGUILAR-...	81B HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
80 <sup>+14</sup> <sub>-11</sub>	572	ALHARRAN	81 HBC	8.25 $K^- p \rightarrow \Lambda K \bar{K}$
72±25	166	EVANGELIS...	77 OMEG	10 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
69±22	100	AGUILAR-...	72B HBC	3.9,4.6 $K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

92 <sup>+25</sup> <sub>-16</sub>	61	BINON	07 GAMS	32.5 $K^- p \rightarrow \eta \eta (\Lambda / \Sigma^0)$
75±20		<sup>1</sup> BARKOV	99 SPEC	40 $K^- p \rightarrow K_S^0 K_S^0 \gamma$
62 <sup>+19</sup> <sub>-14</sub>	123	BARREIRO	77 HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
61± 8	120	BRANDENB...	76C ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$

<sup>1</sup> Systematic errors not estimated.

NODE=M013W2;LINKAGE=SK

**PRODUCED IN  $e^+ e^-$  ANNIHILATION AND PARTICLE DECAYS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M013W3  
NODE=M013W3

The data in this block is included in the average printed for a previous datablock.

**89.2<sup>+3.4</sup><sub>-3.0</sub> OUR AVERAGE** Error includes scale factor of 1.8. See the ideogram below.

84 ± 6 <sup>+10</sup> <sub>-5</sub>		AAIJ	13AN LHCB	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
75 <sup>+12</sup> <sub>-10</sub> <sup>+16</sup> <sub>-8</sub>	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
82.9 <sup>+2.1</sup> <sub>-2.2</sub> <sup>+3.3</sup> <sub>-2.0</sub>		UEHARA	13 BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
77 ± 15		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi K^+ K^-$
82 ± 2 ± 3		ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
75 ± 4 <sup>+15</sup> <sub>-5</sub>		BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$

100 ±15	331	<sup>2</sup> ACCIARRI	01H L3	91, 183–209 $e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
60 ±20 ±19		ABREU	96c DLPH	$Z^0 \rightarrow K^+ K^- + X$
60 ±23 <sup>+13</sup> <sub>-20</sub>		BAI	96c BES	$J/\psi \rightarrow \gamma K^+ K^-$
103 ±30		AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
100 ± 3		<sup>3</sup> FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$
85 ±35		BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
78 ± 6		<sup>4</sup> KLEMP	22 RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0, \gamma K_S^0 K_S^0$
84 ±15		<sup>5</sup> RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
37 ±12	29	<sup>6</sup> LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$
104 ±10	870	<sup>7</sup> SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
62 ±10		<sup>8</sup> FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

- <sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.
- <sup>2</sup> Supersedes ACCIARRI 95J.
- <sup>3</sup> From an analysis including interference with  $f_0(1710)$ .
- <sup>4</sup> Fit of the tensor partial waves from BES3 in the multipole basis.
- <sup>5</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma \pi^0 \pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).
- <sup>6</sup> From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated.
- <sup>7</sup> From analysis of L3 data at 91 and 183–209 GeV.
- <sup>8</sup> From an analysis ignoring interference with  $f_0(1710)$ .

NODE=M013W3;LINKAGE=A

NODE=M013W;LINKAGE=HA

NODE=M013W;LINKAGE=F2


NODE=M013W3;LINKAGE=D

NODE=M013W3;LINKAGE=C

NODE=M013W3;LINKAGE=B

NODE=M013W3;LINKAGE=SC

NODE=M013W;LINKAGE=F1



**PRODUCED IN  $\bar{p}p$  ANNIHILATION**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

**77 ± 5 OUR AVERAGE**

76 ± 6	AMSLER	06	CBAR	0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
79 ± 8	<sup>1</sup> AMSLER	02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
104.8 ± 0.9 ± 9.8	<sup>2</sup> ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
128 ± 20	<sup>3</sup> ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$

- <sup>1</sup> T-matrix pole.
- <sup>2</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).
- <sup>3</sup> K-matrix, 4-poles, 5-channel fit.

NODE=M013W9;LINKAGE=TT

NODE=M013W9;LINKAGE=A

NODE=M013W9;LINKAGE=AN

**CENTRAL PRODUCTION**

VALUE (MeV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  
 The data in this block is included in the average printed for a previous datablock.

NODE=M013W4  
 NODE=M013W4

**70±25**BARBERIS 99 OMEG 450  $p p \rightarrow p_s p_f K^+ K^-$ **PRODUCED IN  $e p$  COLLISIONS**

VALUE (MeV) \_\_\_\_\_ EVTS \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  
 The data in this block is included in the average printed for a previous datablock.

NODE=M013W10  
 NODE=M013W10

**83±<sub>-4</sub><sup>+5</sup>**<sup>1</sup> CHEKANOV 08 ZEUS  $e p \rightarrow K_S^0 K_S^0 X$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

50<sup>+34</sup><sub>-22</sub>84 <sup>2</sup> CHEKANOV 04 ZEUS  $e p \rightarrow K_S^0 K_S^0 X$ 

<sup>1</sup> In the SU(3) based model with a specific interference pattern of the  $f_2(1270)$ ,  $a_2^0(1320)$ , and  $f_2'(1525)$  mesons incoherently added to the  $f_0(1710)$  and non-resonant background.

<sup>2</sup> Systematic errors not estimated.

NODE=M013W10;LINKAGE=HE

NODE=M013W10;LINKAGE=CH

 **$f_2'(1525)$  DECAY MODES**

NODE=M013215;NODE=M013

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $K \bar{K}$	(87.6±2.2) %	1.1
$\Gamma_2$ $\eta \eta$	(11.6±2.2) %	1.1
$\Gamma_3$ $\pi \pi$	( 8.3±1.6) × 10 <sup>-3</sup>	
$\Gamma_4$ $K \bar{K}^*(892) + c.c.$		
$\Gamma_5$ $\pi K \bar{K}$		
$\Gamma_6$ $\pi \pi \eta$		
$\Gamma_7$ $\pi^+ \pi^+ \pi^- \pi^-$		
$\Gamma_8$ $\gamma \gamma$	( 9.5±1.1) × 10 <sup>-7</sup>	1.1

DESIG=2

DESIG=4

DESIG=1

DESIG=3

DESIG=6

DESIG=5

DESIG=7

DESIG=8

**CONSTRAINED FIT INFORMATION**

An overall fit to the total width, 2 partial widths, a combination of partial widths obtained from integrated cross sections, and 3 branching ratios uses 17 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 18.2$  for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-100			
$x_3$	-6	-1		
$x_8$	-19	19	1	
$\Gamma$	-4	4	0	-44
	$x_1$	$x_2$	$x_3$	$x_8$

Mode	Rate (MeV)	Scale factor
$\Gamma_1$ $K \bar{K}$	75 ±4	1.8
$\Gamma_2$ $\eta \eta$	9.9 ±1.9	1.1
$\Gamma_3$ $\pi \pi$	0.71±0.14	1.1
$\Gamma_8$ $\gamma \gamma$	( 8.2 ±0.9 ) × 10 <sup>-5</sup>	

DESIG=2

DESIG=4

DESIG=1

DESIG=8

 **$f_2'(1525)$  PARTIAL WIDTHS**

NODE=M013220

 **$\Gamma(K \bar{K})$**  **$\Gamma_1$** 

VALUE (MeV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_  
**75±4 OUR FIT** Error includes scale factor of 1.8.

NODE=M013W6  
 NODE=M013W6

**63±<sub>-5</sub><sup>+6</sup>**<sup>1</sup> LONGACRE 86 MPS 22  $\pi^- p \rightarrow K_S^0 K_S^0 n$ 

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

NODE=M013PW;LINKAGE=L

$\Gamma(\eta\eta)$  $\Gamma_2$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.9±0.19 OUR FIT** Error includes scale factor of 1.1.

••• We do not use the following data for averages, fits, limits, etc. •••

5.0±0.8	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
24 $\begin{smallmatrix} +3 \\ -1 \end{smallmatrix}$		<sup>2</sup> LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$  MeV and SU(3) relations.<sup>2</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.NODE=M013W7  
NODE=M013W7

NODE=M013W7;LINKAGE=SC

NODE=M013PW7;LINKAGE=L

 $\Gamma(\pi\pi)$  $\Gamma_3$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.71±0.14 OUR FIT** Error includes scale factor of 1.1.

1.4 $\begin{smallmatrix} +1.0 \\ -0.5 \end{smallmatrix}$		<sup>1</sup> LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
--	--	--------------------------	-----	--

••• We do not use the following data for averages, fits, limits, etc. •••

0.2 $\begin{smallmatrix} +1.0 \\ -0.2 \end{smallmatrix}$	870	<sup>2</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
--	-----	-----------------------------	------	--

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles.<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$  MeV and SU(3) relations.NODE=M013W5  
NODE=M013W5

NODE=M013PW5;LINKAGE=L

NODE=M013W5;LINKAGE=SC

 $\Gamma(\gamma\gamma)$  $\Gamma_8$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.082±0.009 OUR FIT**

••• We do not use the following data for averages, fits, limits, etc. •••

0.13 ±0.03	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
------------	-----	-----------------------------	------	--

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$  MeV and SU(3) relations.NODE=M013W8  
NODE=M013W8

NODE=M013W8;LINKAGE=SC

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.746±0.002 $\begin{smallmatrix} +0.166 \\ -0.162 \end{smallmatrix}$	<sup>1</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
--	--------------------------	------	---

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).NODE=M013R00  
NODE=M013R00

NODE=M013R00;LINKAGE=A

 $f_2'(1525) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M013223

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_8/\Gamma$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**0.072 ±0.007 OUR FIT****0.072 ±0.007 OUR AVERAGE**

0.048 $\begin{smallmatrix} +0.067 \\ -0.008 \end{smallmatrix}$ $\begin{smallmatrix} +0.108 \\ -0.012 \end{smallmatrix}$	UEHARA 13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
0.0564±0.0048±0.0116	ABE 04	BELL	10.6 $e^+e^- \rightarrow e^+e^- K^+K^-$
0.076 ±0.006 ±0.011	331 ACCIARRI 01H	L3	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.067 ±0.008 ±0.015	<sup>2</sup> ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
0.11 $\begin{smallmatrix} +0.03 \\ -0.02 \end{smallmatrix}$ ±0.02	BEHREND 89c	CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.10 $\begin{smallmatrix} +0.04 \\ -0.03 \end{smallmatrix}$ $\begin{smallmatrix} +0.03 \\ -0.02 \end{smallmatrix}$	BERGER 88	PLUT	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.12 ±0.07 ±0.04	<sup>2</sup> AIHARA 86B	TPC	$e^+e^- \rightarrow e^+e^- K^+K^-$
0.11 ±0.02 ±0.04	<sup>2</sup> ALTHOFF 83	TASS	$e^+e^- \rightarrow e^+e^- K\bar{K}$

••• We do not use the following data for averages, fits, limits, etc. •••

0.0314±0.0050±0.0077	<sup>3</sup> ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
----------------------	---------------------------	-----	------------------------------------

<sup>1</sup> Supersedes ACCIARRI 95J. From analysis of L3 data at 91 and 183–209 GeV,<sup>2</sup> Using an incoherent background.<sup>3</sup> Using a coherent background.NODE=M013G1  
NODE=M013G1

OCCUR=2

NODE=M013G;LINKAGE=HA

NODE=M013G1;LINKAGE=A

NODE=M013G1;LINKAGE=B

$f'_2(1525)$  BRANCHING RATIOS $\Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.059 ± 0.003 ± 0.026	<sup>1</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
seen	UEHARA 10A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \eta \eta$
0.10 ± 0.03	<sup>2</sup> PROKOSHKIN 91	GAM4 300	$\pi^- p \rightarrow \pi^- p \eta \eta$

<sup>1</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

<sup>2</sup> Combining results of GAM4 with those of WA76 on  $K\bar{K}$  central production and results of CBAL, MRK3 and DM2 on  $J/\psi \rightarrow \gamma\eta\eta$ .

NODE=M013225

NODE=M013R8  
NODE=M013R8

NODE=M013R8;LINKAGE=A

NODE=M013R8;LINKAGE=B

 $\Gamma(\eta\eta)/\Gamma(K\bar{K})$  $\Gamma_2/\Gamma_1$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

**0.132 ± 0.028 OUR FIT****0.115 ± 0.028 OUR AVERAGE**

0.119 ± 0.015 ± 0.036	61	<sup>1</sup> BINON	07	GAMS	32.5 $K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
0.11 ± 0.04		<sup>2</sup> PROKOSHKIN 91	GAM4 300		$\pi^- p \rightarrow \pi^- p \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.14	90	BARBERIS	00E		450 $pp \rightarrow p_f \eta \eta p_s$
< 0.50		BARNES	67	HBC	4.6, 5.0 $K^- p$

<sup>1</sup> Using the compilation of the cross sections for  $f'_2(1525)$  production in  $K^- p$  collisions from ASTON 88D.

<sup>2</sup> Combining results of GAM4 with those of WA76 on  $K\bar{K}$  central production and results of CBAL, MRK3 and DM2 on  $J/\psi \rightarrow \gamma\eta\eta$ .

NODE=M013R3  
NODE=M013R3

NODE=M013R3;LINKAGE=BI

NODE=M013R3;LINKAGE=B

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

**0.83 ± 0.16 OUR FIT****0.75 ± 0.16 OUR AVERAGE**

0.7 ± 0.2		COSTA	80	OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
2.7 <sup>+7.1</sup> <sub>-1.3</sub>		<sup>1</sup> GORLICH	80	ASPK	17, 18 $\pi^- p$
0.75 ± 0.25		<sup>1,2</sup> MARTIN	79	RVUE	
3.4 ± 1.5 ± 1.0		<sup>3</sup> ALBRECHT 20	RVUE		0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
< 6	95	AGUILAR-...	81B	HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
19 ± 3		CORDEN	79	OMEG	12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
< 4.5	95	BARREIRO	77	HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
1.2 ± 0.4		<sup>1</sup> PAWLICKI	77	SPEC	6 $\pi N \rightarrow K^+ K^- N$
< 6.3	90	BRANDENB...	76C	ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
< 0.86		<sup>1</sup> BEUSCH	75B	OSPK	8.9 $\pi^- p \rightarrow K^0 \bar{K}^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Assuming that the  $f'_2(1525)$  is produced by an one-pion exchange production mechanism.

<sup>2</sup> MARTIN 79 uses the PAWLICKI 77 data with different input value of the  $f'_2(1525) \rightarrow K\bar{K}$  branching ratio.

<sup>3</sup> Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

NODE=M013R1  
NODE=M013R1

NODE=M013R1;LINKAGE=C

NODE=M013R1;LINKAGE=D

NODE=M013R1;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$  $\Gamma_3/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.0094 ± 0.0018 OUR FIT****0.075 ± 0.035**AUGUSTIN 87 DM2  $J/\psi \rightarrow \gamma\pi^+\pi^-$ NODE=M013R7  
NODE=M013R7 $[\Gamma(K\bar{K}^*(892) + \text{c.c.}) + \Gamma(\pi K\bar{K})]/\Gamma(K\bar{K})$  $(\Gamma_4 + \Gamma_5)/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.35	95	AGUILAR-...	72B	HBC	3.9, 4.6 $K^- p$
< 0.4	67	AMMAR	67	HBC	

NODE=M013R5  
NODE=M013R5

$\Gamma(\pi\pi\eta)/\Gamma(K\bar{K})$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma_1$
-------	-----	-------------	------	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.41	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
<0.3	67	AMMAR	67	HBC	

NODE=M013R4  
NODE=M013R4

 $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K\bar{K})$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma_1$
-------	-----	-------------	------	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.32	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
-------	----	-------------	-----	-----	-----------------

NODE=M013R6  
NODE=M013R6

 $f'_2(1525)$  REFERENCES

NODE=M013

KLEMP	22	PL B830 137171	E. Klempt <i>et al.</i>	(BONN)	REFID=61646
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
AAIJ	13AN	PR D87 072004	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55137
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
BINON	07	PAN 70 1713	F. Binon <i>et al.</i>	(GAMS Collab.)	REFID=52057
		Translated from YAF 70 1758.			
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=49672
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARKOV	99	JETPL 70 248	B.P. Barkov <i>et al.</i>		REFID=47379
		Translated from ZETFP 70 242.			
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
ACCIARRI	95J	PL B363 118	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44615
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40330
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BERGER	88	ZPHY C37 329	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=40566
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 $\gamma$ Collab.)	REFID=20764
BOLONKIN	86	SJNP 43 776	B.V. Bolonkin <i>et al.</i>	(ITEP) JP	REFID=44646
		Translated from YAF 43 1211.			
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	83	PL 121B 216	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21408
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
AGUILAR-...	81B	ZPHY C8 313	M. Aguilar-Benitez <i>et al.</i>	(CERN, CDEF+)	REFID=21104
ALHARRAN	81	NP B191 26	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=21403
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
BARREIRO	77	NP B121 237	F. Barreiro <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=21392
EVANGELIS...	77	NP B127 384	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20540
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJP	REFID=20367
BRANDENB...	76C	NP B104 413	G.W. Brandenburg <i>et al.</i>	(SLAC)	REFID=20225
BEUSCH	75B	PL 60B 101	W. Beusch <i>et al.</i>	(CERN, ETH)	REFID=21390
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
AMMAR	67	PRL 19 1071	R. Ammar <i>et al.</i>	(NWES, ANL) JP	REFID=21382
BARNES	67	PRL 19 964	V.E. Barnes <i>et al.</i>	(BNL, SYRA) IJPC	REFID=21383
CRENNELL	66	PRL 16 1025	D.J. Crennell <i>et al.</i>	(BNL) I	REFID=20317

**$f_2(1565)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M123

OMITTED FROM SUMMARY TABLE

Seen mostly in antinucleon-nucleon annihilation. Needs confirmation in other channels. See the review on "Spectroscopy of Light Meson Resonances."

NODE=M123

 **$f_2(1565)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M123PP

NODE=M123PP

NODE=M123PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1495–1560) – <math>i</math> (40–110) OUR ESTIMATE</b>			
$(1560 \pm 15) - i (140 \pm 20)$	<sup>1</sup> ANISOVICH	09	RVUE 0.0 $\bar{p}p, \pi N$
$(1552 \pm 13) - i (57 \pm 12)$	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta,$ $\pi^0 \pi^0 \pi^0$
$(1507 \pm 15) - i (65 \pm 10)$	BERTIN	97C	OBLX 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1534 \pm 20) - i (90 \pm 30)$	<sup>2</sup> ABELE	96C	RVUE Compilation
$(\sim 1552) - i (\sim 71)$	<sup>3</sup> AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$

<sup>1</sup> On sheet II in a two-pole solution.<sup>2</sup> T-matrix pole, large coupling to  $\rho\rho$  and  $\omega\omega$ , could be  $f_2(1640)$ .<sup>3</sup> Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

NODE=M123PP;LINKAGE=A

NODE=M123PP;LINKAGE=D

NODE=M123PP;LINKAGE=E

 **$f_2(1565)$  MASS**

NODE=M123M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>1575 \pm 18</math> OUR AVERAGE</b>	[1542 $\pm$ 19 MeV OUR 2022 AVERAGE Scale factor = 2.2]		
<b><math>1575 \pm 18</math></b>	<sup>4</sup> BERTIN	98	OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1590 \pm 10$	<sup>5,6</sup> AMELIN	06	VES 36 $\pi^- p \rightarrow \omega \omega n$
$1550 \pm 10 \pm 20$	<sup>6</sup> AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
$1598 \pm 11 \pm 9$	BAKER	99B	SPEC 0 $\bar{p}p \rightarrow \omega \omega \pi^0$
$1598 \pm 72$	BALOSHIN	95	SPEC 40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
$1566^{+80}_{-50}$	<sup>7</sup> ANISOVICH	94	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0$
$1502 \pm 9$	ADAMO	93	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
$1488 \pm 10$	<sup>8</sup> ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
$1508 \pm 10$	<sup>8</sup> ARMSTRONG	93D	E760 $\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
$1525 \pm 10$	<sup>8</sup> ARMSTRONG	93D	E760 $\bar{p}p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
$\sim 1504$	<sup>9</sup> WEIDENAUER	93	ASTE 0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$
$1540 \pm 15$	<sup>8</sup> ADAMO	92	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
$1515 \pm 10$	<sup>10</sup> AKER	91	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0$
$1565 \pm 20$	MAY	90	ASTE 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$1477 \pm 5$	BRIDGES	86C	DBC 0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$

NODE=M123M

NEW

OCCUR=2

<sup>4</sup> Breit-Wigner mass.<sup>5</sup> Supersedes the  $\omega\omega$  state of BELADIDZE 92B earlier assigned to the  $f_2(1640)$ .<sup>6</sup> Breit-Wigner width.<sup>7</sup> From a simultaneous analysis of the annihilations  $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$  including AKER 91 data.<sup>8</sup>  $J^P$  not determined, could be partly  $f_0(1500)$ .<sup>9</sup>  $J^P$  not determined.<sup>10</sup> Superseded by AMSLER 95B.

NODE=M123M;LINKAGE=G

NODE=M123M;LINKAGE=AM

NODE=M123M;LINKAGE=D

NODE=M123M;LINKAGE=C

NODE=M123M;LINKAGE=E

NODE=M123M;LINKAGE=F

NODE=M123M;LINKAGE=BA

 **$f_2(1565)$  WIDTH**

NODE=M123W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>119 \pm 24</math> OUR AVERAGE</b>	[122 $\pm$ 13 MeV OUR 2022 AVERAGE]		
<b><math>119 \pm 24</math></b>	<sup>11</sup> BERTIN	98	OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$

NODE=M123W

NEW

- • • We do not use the following data for averages, fits, limits, etc. • • •

140± 11	<sup>12,13</sup> AMELIN	06	VES	36	$\pi^- p \rightarrow \omega \omega n$
130± 20±40	<sup>13</sup> AMELIN	00	VES	37	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$
263±101	BALOSHIN	95	SPEC	40	$\pi^- C \rightarrow K_S^0 K_S^0 X$
166 <sup>+</sup> <sub>-20</sub>	<sup>14</sup> ANISOVICH	94	CBAR	0.0	$\bar{p} p \rightarrow 3\pi^0, \eta \eta \pi^0$
130± 10	<sup>15</sup> ADAMO	93	OBLX		$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
148± 27	<sup>16</sup> ARMSTRONG	93C	E760		$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
103± 15	<sup>16</sup> ARMSTRONG	93D	E760		$\bar{p} p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111± 10	<sup>16</sup> ARMSTRONG	93D	E760		$\bar{p} p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 206	<sup>17</sup> WEIDENAUER	93	ASTE	0.0	$\bar{p} N \rightarrow 3\pi^- 2\pi^+$
132± 37	<sup>16</sup> ADAMO	92	OBLX		$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
120± 10	<sup>18</sup> AKER	91	CBAR	0.0	$\bar{p} p \rightarrow 3\pi^0$
170± 40	MAY	90	ASTE	0.0	$\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
116± 9	BRIDGES	86C	DBC	0.0	$\bar{p} N \rightarrow 3\pi^- 2\pi^+$

<sup>11</sup> Breit-Wigner width.

<sup>12</sup> Supersedes the  $\omega \omega$  state of BELADIDZE 92B earlier assigned to the  $f_2(1640)$ .

<sup>13</sup> Breit-Wigner width.

<sup>14</sup> From a simultaneous analysis of the annihilations  $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$  including AKER 91 data.

<sup>15</sup> Supersedes ADAMO 92.

<sup>16</sup>  $J^P$  not determined, could be partly  $f_0(1500)$ .

<sup>17</sup>  $J^P$  not determined.

<sup>18</sup> Superseded by AMSLER 95B.

OCCUR=2

NODE=M123W;LINKAGE=G  
 NODE=M123W;LINKAGE=AM  
 NODE=M123W;LINKAGE=H  
 NODE=M123W;LINKAGE=D

NODE=M123W;LINKAGE=C  
 NODE=M123W;LINKAGE=E  
 NODE=M123W;LINKAGE=F  
 NODE=M123W;LINKAGE=BA

## $f_2(1565)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi \pi$	seen
$\Gamma_2$ $\pi^+ \pi^-$	seen
$\Gamma_3$ $\pi^0 \pi^0$	seen
$\Gamma_4$ $\rho^0 \rho^0$	seen
$\Gamma_5$ $2\pi^+ 2\pi^-$	seen
$\Gamma_6$ $\eta \eta$	seen
$\Gamma_7$ $\omega \omega$	seen
$\Gamma_8$ $K \bar{K}$	seen
$\Gamma_9$ $\gamma \gamma$	seen

NODE=M123215;NODE=M123

DESIG=6;OUR EST;→ UNCHECKED ←  
 DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=3;OUR EST;→ UNCHECKED ←  
 DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=5;OUR EST;→ UNCHECKED ←  
 DESIG=4;OUR EST;→ UNCHECKED ←  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=9;OUR EST;→ UNCHECKED ←  
 DESIG=10;OUR EST;→ UNCHECKED ←

## $f_2(1565)$ PARTIAL WIDTHS

### $\Gamma(\eta \eta)$

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  $\Gamma_6$

- • • We do not use the following data for averages, fits, limits, etc. • • •

1.2±0.3 870 <sup>19</sup> SCHEGELSKY 06A RVUE  $\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M123225

NODE=M123W3  
 NODE=M123W3

### $\Gamma(K \bar{K})$

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  $\Gamma_8$

- • • We do not use the following data for averages, fits, limits, etc. • • •

2.0±1.0 870 <sup>19</sup> SCHEGELSKY 06A RVUE  $\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M123W1  
 NODE=M123W1

### $\Gamma(\gamma \gamma)$

VALUE (keV) EVTS DOCUMENT ID TECN COMMENT  $\Gamma_9$

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.14 870 <sup>19</sup> SCHEGELSKY 06A RVUE  $\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M123W2  
 NODE=M123W2

<sup>19</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $f_2(1565)$  mass of 1570 MeV, width of 160 MeV,  $\Gamma(\pi \pi) = 25$  MeV, and SU(3) relations.

NODE=M123W1;LINKAGE=SC

## $f_2(1565)$ BRANCHING RATIOS

### $\Gamma(\pi \pi)/\Gamma_{\text{total}}$

VALUE DOCUMENT ID TECN COMMENT  $\Gamma_1/\Gamma$

- • • We do not use the following data for averages, fits, limits, etc. • • •

seen BAKER 99B SPEC  $0 \bar{p} p \rightarrow \omega \omega \pi^0$

NODE=M123220

NODE=M123R5  
 NODE=M123R5



$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R1  
 NODE=M123R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BERTIN	98	OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
not seen	<sup>20</sup> ANISOVICH	94B	RVUE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
seen	MAY	89	ASTE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

<sup>20</sup> ANISOVICH 94B is from a reanalysis of MAY 90.

NODE=M123R1;LINKAGE=A

 $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R3  
 NODE=M123R3

seen	AMSLER	95B	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0$
------	--------	-----	--

 $\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$  $\Gamma_2/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R2  
 NODE=M123R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042±0.013	BRIDGES	86B	DBC $\bar{p}N \rightarrow 3\pi^-2\pi^+$
-------------	---------	-----	---

 $\Gamma(\eta\eta)/\Gamma(\pi^0\pi^0)$  $\Gamma_6/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R4  
 NODE=M123R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.024±0.005±0.012	<sup>21</sup> ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
-------------------	-------------------------	-----	---

<sup>21</sup>  $J^P$  not determined, could be partly  $f_0(1500)$ .

NODE=M123R4;LINKAGE=E

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R6  
 NODE=M123R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BAKER	99B	SPEC 0 $\bar{p}p \rightarrow \omega\omega\pi^0$
------	-------	-----	---

 $f_2(1565)$  REFERENCES

NODE=M123

ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=51574
		Translated from YAF 69 715.			
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
BAKER	99B	PL B467 147	C.A. Baker <i>et al.</i>		REFID=47398
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=44621
		Translated from YAF 58 50.			
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
ANISOVICH	94B	PR D50 1972	V.V. Anisovich <i>et al.</i>	(LOQM)	REFID=44071
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.)	REFID=43657
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43596
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)	REFID=42177
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42172
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	REFID=41587
MAY	90	ZPHY C46 203	B. May <i>et al.</i>	(ASTERIX Collab.)	REFID=41365
MAY	89	PL B225 450	B. May <i>et al.</i>	(ASTERIX Collab.) IJP	REFID=40921
BRIDGES	86B	PRL 56 215	D.L. Bridges <i>et al.</i>	(SYRA, CASE)	REFID=21376
BRIDGES	86C	PRL 57 1534	D.L. Bridges <i>et al.</i>	(SYRA)	REFID=21377

**$\rho(1570)$** 

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

May be an OZI-violating decay mode of  $\rho(1700)$ . See the review on "Spectroscopy of Light Meson Resonances."

NODE=M188

NODE=M188

 **$\rho(1570)$  MASS**

NODE=M188M

NODE=M188M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1570±36±62</b>	54	<sup>1</sup> AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••				
1585±15		<sup>2</sup> ACHASOV	20C SND	1.3–2.0 $e^+e^- \rightarrow K^+K^-\pi^0$
1480±40		<sup>3</sup> BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$

<sup>1</sup> From the fit with two resonances.<sup>2</sup> From a fit using a two resonance model in which the mass and width of the other resonance are fixed at the  $\rho(1700)$  values from PDG 20.<sup>3</sup> Systematic errors not estimated.

NODE=M188M;LINKAGE=AU

NODE=M188M;LINKAGE=A

NODE=M188M;LINKAGE=BI

 **$\rho(1570)$  WIDTH**

NODE=M188W

NODE=M188W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>144±75±43</b>	54	<sup>4</sup> AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••				
75±30		<sup>5</sup> ACHASOV	20C SND	1.3–2.0 $e^+e^- \rightarrow K^+K^-\pi^0$
130±60		<sup>6</sup> BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$

<sup>4</sup> From the fit with two resonances.<sup>5</sup> From a fit using a two resonance model in which the mass and width of the other resonance are fixed at the  $\rho(1700)$  values from PDG 20.<sup>6</sup> Systematic errors not estimated.

NODE=M188W;LINKAGE=AU

NODE=M188W;LINKAGE=A

NODE=M188W;LINKAGE=BI

 **$\rho(1570)$  DECAY MODES**

NODE=M188215;NODE=M188

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+e^-$	
$\Gamma_2$ $\phi\pi$	not seen
$\Gamma_3$ $\omega\pi$	

DESIG=1

DESIG=2

DESIG=3

 **$\rho(1570)$   $\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

NODE=M188225

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
<b>3.5±0.9±0.3</b>		54	<sup>7</sup> AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$	
••• We do not use the following data for averages, fits, limits, etc. •••						
<70	90		<sup>8</sup> AULCHENKO	87B ND	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$	

<sup>7</sup> From the fit with two resonances.<sup>8</sup> Using mass and width of BITYUKOV 87.

NODE=M188G01;LINKAGE=AU

NODE=M188G01;LINKAGE=AL

 **$\rho(1570)$  BRANCHING RATIOS**

NODE=M188220

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
<b>not seen</b>	ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$	
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.01	<sup>9</sup> DONNACHIE	91 RVUE		

<sup>9</sup> Using data from BISELLO 91B, DOLINSKY 86, and ALBRECHT 87L.

NODE=M188R01;LINKAGE=DO

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_3$
••• We do not use the following data for averages, fits, limits, etc. •••					
>0.5	95	BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$	

NODE=M188R02

NODE=M188R02

**$\rho(1570)$  REFERENCES**

ACHASOV	20C	EPJ C80 1139	M.N. Achasov <i>et al.</i>	(SND Collab.)	NODE=M188
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60935
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=60676
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=52242
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=45765
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41752
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41632
AULCHENKO	87B	JETPL 45 145	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=40418
		Translated from ZETFP 45 118.			REFID=41373
BITYUKOV	87	PL B188 383	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=40011
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246

 **$h_1(1595)$** 

$$I^G(J^{PC}) = 0^-(1^{+-})$$

OMITTED FROM SUMMARY TABLE

Seen in a partial-wave analysis of the  $\omega\eta$  system produced in the reaction  $\pi^- p \rightarrow \omega\eta n$  at 18 GeV/c.

 **$h_1(1595)$  MASS**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1594 \pm 15^{+10}_{-60}</math></b>	EUGENIO	01	SPEC 18 $\pi^- p \rightarrow \omega\eta n$

 **$h_1(1595)$  WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>384 \pm 60^{+70}_{-100}</math></b>	EUGENIO	01	SPEC 18 $\pi^- p \rightarrow \omega\eta n$

 **$h_1(1595)$  DECAY MODES**

<u>Mode</u>	<u>Fraction (<math>\Gamma_i/\Gamma</math>)</u>
$\Gamma_1 \quad \omega\eta$	seen

 **$h_1(1595)$  REFERENCES**

EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>	NODE=M166
				REFID=48010

NODE=M166

NODE=M166

NODE=M166M

NODE=M166M

NODE=M166W

NODE=M166W

NODE=M166215;NODE=M166

DESIG=1;OUR EST;→ UNCHECKED ←

$\pi_1(1600)$ 

$$I^G(J^{PC}) = 1^-(1^-+)$$

See the review on "Spectroscopy of Light Meson Resonances" and a note in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M164

NODE=M164

 **$\pi_1(1600)$  T-Matrix Pole  $\sqrt{s}$** 

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M164TMP

NODE=M164TMP

NODE=M164TMP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1480–1680) – i (150–300) OUR ESTIMATE</b>			
$(1623 \pm 47^{+24}_{-75}) - i(228 \pm 44^{+72}_{-88})$	<sup>1</sup> KOPF	21	RVUE 0.9 $p\bar{p} \rightarrow \pi^0\pi^0\eta$ , $\pi^0\eta\eta$ , $\pi^0K^+K^-$ and 191 $\pi^-p \rightarrow \pi^-\pi^-\pi^+p$
$(1564 \pm 24 \pm 86) - i(246 \pm 27 \pm 51)$	<sup>2</sup> RODAS	19	RVUE 191 $\pi^-p \rightarrow \eta(\prime)\pi^-p$
<sup>1</sup> From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi$ , $\eta'\pi$ and $K\bar{K}$ systems.			
<sup>2</sup> The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data.			

NODE=M164TMP;LINKAGE=B

NODE=M164TMP;LINKAGE=A

 **$\pi_1(1600)$  MASS**

NODE=M164M

NODE=M164M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1661^{+15}_{-11}</math> OUR AVERAGE</b> Error includes scale factor of 1.2.				
$1600^{+110}_{-60}$	46M	<sup>1</sup> AGHASYAN	18B	COMP 190 $\pi^-p \rightarrow \pi^-\pi^+\pi^-p$
$1664 \pm 8 \pm 10$	145k	<sup>2</sup> LU	05	B852 18 $\pi^-p \rightarrow \omega\pi^-\pi^0p$
$1709 \pm 24 \pm 41$	69k	<sup>3</sup> KUHN	04	B852 18 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$
$1597 \pm 10^{+45}_{-10}$		<sup>3</sup> IVANOV	01	B852 18 $\pi^-p \rightarrow \eta'\pi^-p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1660 \pm 10^{+0}_{-64}$	420k	<sup>4</sup> ALEKSEEV	10	COMP 190 $\pi^-Pb \rightarrow \pi^-\pi^-\pi^+Pb'$
$1593 \pm 8^{+29}_{-47}$		<sup>3,5</sup> ADAMS	98B	B852 18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$
<sup>1</sup> Statistical error negligible. See also the review ALEXEEV 22.				
<sup>2</sup> May be a different state: natural and unnatural parity exchanges.				
<sup>3</sup> Natural parity exchange.				
<sup>4</sup> Superseded by AGHASYAN 2018B.				
<sup>5</sup> Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with 2.6 M events of $\pi^-p \rightarrow \pi^-\pi^-\pi^+p$ and 3 M events of $\pi^-p \rightarrow \pi^-\pi^0\pi^0p$ of E852 data.				

NODE=M164M;LINKAGE=B

NODE=M164M;LINKAGE=LU

NODE=M164M;LINKAGE=A

NODE=M164M;LINKAGE=C

NODE=M164M;LINKAGE=DZ

 **$\pi_1(1600)$  WIDTH**


NODE=M164W

NODE=M164W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>240 \pm 50</math> OUR AVERAGE</b> Error includes scale factor of 1.7. See the ideogram below.				
$580^{+100}_{-230}$	46M	<sup>1</sup> AGHASYAN	18B	COMP 190 $\pi^-p \rightarrow \pi^-\pi^+\pi^-p$
$185 \pm 25 \pm 28$	145k	<sup>2</sup> LU	05	B852 18 $\pi^-p \rightarrow \omega\pi^-\pi^0p$
$403 \pm 80 \pm 115$	69k	<sup>3</sup> KUHN	04	B852 18 $\pi^-p \rightarrow \eta\pi^+\pi^-\pi^-p$
$340 \pm 40 \pm 50$		<sup>3</sup> IVANOV	01	B852 18 $\pi^-p \rightarrow \eta'\pi^-p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$269 \pm 21^{+42}_{-64}$	420k	<sup>4</sup> ALEKSEEV	10	COMP 190 $\pi^-Pb \rightarrow \pi^-\pi^-\pi^+Pb'$
$168 \pm 20^{+150}_{-12}$		<sup>3,5</sup> ADAMS	98B	B852 18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$

- 1 Statistical error negligible. See also the review ALEXEEV 22.
- 2 May be a different state: natural and unnatural parity exchanges.
- 3 Natural parity exchange.
- 4 Superseded by AGHASYAN 2018B.
- 5 Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with 2.6 M events of  $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$  and 3 M events of  $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$  of E852 data.

WEIGHTED AVERAGE  
240±50 (Error scaled by 1.7)



### $\pi_1(1600)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi \pi \pi$	seen
$\Gamma_2$ $\rho^0 \pi^-$	seen
$\Gamma_3$ $f_2(1270) \pi^-$	not seen
$\Gamma_4$ $b_1(1235) \pi$	seen
$\Gamma_5$ $\eta'(958) \pi^-$	seen
$\Gamma_6$ $\eta \pi$	seen
$\Gamma_7$ $f_1(1285) \pi$	seen

NODE=M164215;NODE=M164

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=2  
DESIG=4  
DESIG=5  
DESIG=3  
DESIG=7;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←

### $\pi_1(1600)$ BRANCHING RATIOS

$\Gamma(\rho^0 \pi^-)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ALEKSEEV	10	COMP 190 $\pi^- P b \rightarrow \pi^- \pi^- \pi^+ P b'$
• • •	We do not use the following data for averages, fits, limits, etc. • • •		
not seen	NOZAR	09	CLAS $\gamma p \rightarrow 2\pi^+ \pi^- n$
not seen	<sup>1</sup> DZIERBA	06	B852 18 $\pi^- p$

<sup>1</sup> From the PWA analysis of 2.6 M  $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$  and 3 M events of  $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$  of E852 data. Supersedes ADAMS 98B.

NODE=M164220

NODE=M164R1  
NODE=M164R1

NODE=M164R1;LINKAGE=DZ

$\Gamma(f_2(1270) \pi^-)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> DZIERBA	06	B852 18 $\pi^- p$

<sup>1</sup> From the PWA analysis of 2.6 M  $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$  and 3 M events of  $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$  of E852 data. Supersedes CHUNG 02.

NODE=M164R3  
NODE=M164R3

NODE=M164R3;LINKAGE=DZ

$\Gamma(b_1(1235) \pi)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	35280	<sup>1</sup> BAKER	03	SPEC $\bar{p} p \rightarrow \omega \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 145k LU 05 B852 18  $\pi^- p \rightarrow \omega \pi^- \pi^0 p$

<sup>1</sup>  $B((b_1 \pi)_{D\text{-wave}})/B((b_1 \pi)_{S\text{-wave}})=0.3 \pm 0.1$ .

NODE=M164R4  
NODE=M164R4

NODE=M164R;LINKAGE=RB

$\Gamma(\eta'(958)\pi^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	IVANOV	01	B852 18 $\pi^- p \rightarrow \eta' \pi^- p$

NODE=M164R2  
 NODE=M164R2

 $\Gamma(\eta'(958)\pi^-)/\Gamma(\eta\pi)$  $\Gamma_5/\Gamma_6$ 

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$5.54 \pm 1.1^{+1.8}_{-0.27}$	<sup>1</sup> KOPF	21	RVUE 0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$

NODE=M164R00  
 NODE=M164R00

<sup>1</sup>From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.

NODE=M164R00;LINKAGE=A

 $\Gamma(f_1(1285)\pi)/\Gamma(\eta'(958)\pi^-)$  $\Gamma_7/\Gamma_5$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.80 ± 0.78</b>	69k	<sup>1</sup> KUHN	04	B852 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

NODE=M164R5  
 NODE=M164R5

<sup>1</sup>Using  $\eta'(958)\pi$  data from IVANOV 01.

NODE=M164R;LINKAGE=KU

 $\pi_1(1600)$  REFERENCES

ALEXEEV	22	PR D105 012005	G.D. Alexeev <i>et al.</i>	(COMPASS Collab.)
KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
NOZAR	09	PRL 102 102002	M. Nozar <i>et al.</i>	(JLab CLAS Collab.)
DZIERBA	06	PR D73 072001	A.R. Dzierba <i>et al.</i>	(BNL E852 Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)
ADAMS	98B	PRL 81 5760	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)

NODE=M164

REFID=61491  
 REFID=61470  
 REFID=60439  
 REFID=59554  
 REFID=59471  
 REFID=56385  
 REFID=53356  
 REFID=52758  
 REFID=51077  
 REFID=51004  
 REFID=50459  
 REFID=49773  
 REFID=49414  
 REFID=48837  
 REFID=48317  
 REFID=46610

 $a_1(1640)$ 

$$I^G(J^{PC}) = 1^-(1^{++})$$

Possibly seen in the study of the hadronic structure in decay  $\tau \rightarrow 3\pi\nu_\tau$  (ABREU 98G and ASNER 00).

NODE=M161

NODE=M161

 $a_1(1640)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1655 ± 16 OUR AVERAGE</b>		Error includes scale factor of 1.2.		
$1700^{+35}_{-130}$	46M	<sup>1</sup> AGHASYAN	18B	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$1691 \pm 18 \pm 30$		DARGENT	17	RVUE $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
$1630 \pm 20$	35k	<sup>2</sup> BAKER	03	SPEC $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
$1714 \pm 9 \pm 36$		CHUNG	02	B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
$1640 \pm 12 \pm 30$		BAKER	99	SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$1670 \pm 90$		BELLINI	85	SPEC 40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$

<sup>1</sup>Statistical error negligible.

<sup>2</sup>Using the  $a_1(1260)$  mass and width results of BOWLER 88.

NODE=M161M

NODE=M161M

NODE=M161M;LINKAGE=A

NODE=M161M;LINKAGE=KB

 $a_1(1640)$  WIDTH


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>254 ± 40 OUR AVERAGE</b>		Error includes scale factor of 1.8. See the ideogram below.		
$510^{+170}_{-90}$	46M	<sup>1</sup> AGHASYAN	18B	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$171 \pm 33 \pm 40$		DARGENT	17	RVUE $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
$225 \pm 30$	35k	<sup>2</sup> BAKER	03	SPEC $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
$308 \pm 37 \pm 62$		CHUNG	02	B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
$300 \pm 22 \pm 40$		BAKER	99	SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$300 \pm 100$		BELLINI	85	SPEC 40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$

NODE=M161W

NODE=M161W

- <sup>1</sup> Statistical error negligible.
- <sup>2</sup> Using the  $a_1(1260)$  mass and width results of BOWLER 88.

NODE=M161W;LINKAGE=A  
 NODE=M161W;LINKAGE=KB



**$a_1(1640)$  DECAY MODES**

NODE=M161215;NODE=M161

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi\pi$	seen
$\Gamma_2$ $f_2(1270)\pi$	seen
$\Gamma_3$ $\sigma\pi$	seen
$\Gamma_4$ $\rho\pi S\text{-wave}$	seen
$\Gamma_5$ $\rho\pi D\text{-wave}$	seen
$\Gamma_6$ $\omega\pi\pi$	seen
$\Gamma_7$ $f_1(1285)\pi$	seen
$\Gamma_8$ $a_1(1260)\eta$	not seen

DESIG=3;OUR EST;→ UNCHECKED ←  
 DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=4;OUR EST;→ UNCHECKED ←  
 DESIG=5;OUR EST;→ UNCHECKED ←  
 DESIG=6;OUR EST;→ UNCHECKED ←  
 DESIG=8

**$a_1(1640)$  BRANCHING RATIOS**

NODE=M161220

$\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$	$\Gamma_2/\Gamma_3$
VALUE	DOCUMENT ID TECN COMMENT
<b>0.24±0.07</b>	BAKER 99 SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$

NODE=M161R1  
 NODE=M161R1

$\Gamma(\rho\pi D\text{-wave})/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma$
VALUE	DOCUMENT ID TECN COMMENT

NODE=M161R2  
 NODE=M161R2

- • • We do not use the following data for averages, fits, limits, etc. • • •
- seen CHUNG 02 B852 18.3  $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
- seen AMELIN 95B VES 36  $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$	$\Gamma_6/\Gamma$
VALUE	EVTS DOCUMENT ID TECN COMMENT

NODE=M161R3  
 NODE=M161R3

- • • We do not use the following data for averages, fits, limits, etc. • • •
- seen 35280 <sup>1</sup> BAKER 03 SPEC  $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

<sup>1</sup> Assuming the  $\omega\rho$  mechanism for the  $\omega\pi\pi$  state.

NODE=M161R;LINKAGE=KB

$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$	$\Gamma_7/\Gamma$
VALUE	DOCUMENT ID TECN COMMENT

NODE=M161R4  
 NODE=M161R4

- • • We do not use the following data for averages, fits, limits, etc. • • •
- not seen KUHN 04 B852 18  $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
- seen LEE 94 MPS2 18  $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- \pi^- p$

$\Gamma(a_1(1260)\eta)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	KUHN	04 B852	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

NODE=M161R5  
 NODE=M161R5

 **$a_1(1640)$  REFERENCES**

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
DARGENT	17	JHEP 1705 143	P. d'Argent <i>et al.</i>	(HEID, BRIS)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	(DELPHI Collab.)
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(SERP, TBIL)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(BNL, IND, KYUN, MASD+)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(OXF)
BOWLER	88	PL B209 99	M.G. Bowler	
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	

Translated from YAF 41 1223.

NODE=M161

REFID=59471  
 REFID=58121  
 REFID=49773  
 REFID=49414  
 REFID=48837  
 REFID=47339  
 REFID=46888  
 REFID=45909  
 REFID=44433  
 REFID=44092  
 REFID=40578  
 REFID=47490

NODE=M117

 **$f_2(1640)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

 **$f_2(1640)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1639 ± 6 OUR AVERAGE</b>	Error includes scale factor of 1.2.		
1620 ± 16	BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1647 ± 7	ADAMO	92 OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
1635 ± 7	ALDE	90 GAM2	$38 \pi^- p \rightarrow \omega \omega n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1640 ± 5	AMSLER	06 CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$
1659 ± 6	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1643 ± 7	<sup>1</sup> ALDE	89B GAM2	$38 \pi^- p \rightarrow \omega \omega n$

<sup>1</sup>Superseded by ALDE 90.

NODE=M117M

NODE=M117M

NODE=M117M;LINKAGE=BB

 **$f_2(1640)$  WIDTH**

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>99<sup>+60</sup><sub>-40</sub> OUR AVERAGE</b>		Error includes scale factor of 2.9.		
140 <sup>+60</sup> <sub>-20</sub>		BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
58 ± 20		ADAMO	92 OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
44 ± 9		AMSLER	06 CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$
152 ± 18		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
< 70	90	ALDE	90 GAM2	$38 \pi^- p \rightarrow \omega \omega n$

NODE=M117W

NODE=M117W

 **$f_2(1640)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\omega\omega$	seen
$\Gamma_2$ $4\pi$	seen
$\Gamma_3$ $K\bar{K}$	seen

NODE=M117215;NODE=M117

DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=3

 **$f_2(1640)$  BRANCHING RATIOS** $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AMSLER	06 CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$

NODE=M117220

NODE=M117R2  
 NODE=M117R2

 **$f_2(1640)$  REFERENCES**

AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
Translated from YAF 69 515.				
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH) JP
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	89B	PL B216 451	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+) IGJPC

NODE=M117

REFID=51136  
 REFID=51191  
 REFID=44438  
 REFID=42177  
 REFID=40935  
 REFID=40735



$\eta_2(1645)$ 

$$I^G(J^{PC}) = 0^+(2^-+)$$

NODE=M154

 $\eta_2(1645)$  MASS

NODE=M154M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>1617± 5 OUR AVERAGE</b>				
1613± 8	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1617± 8	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
1620±20	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1645±14±15	ADOMEIT	96	CBAR 0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1645± 6±20	ANISOVICH	00E	SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M154M

 $\eta_2(1645)$  WIDTH

NODE=M154W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>181±11 OUR AVERAGE</b>				
185±17	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
177±18	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
180±25	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
180 <sup>+40</sup> <sub>-21</sub> ±25	ADOMEIT	96	CBAR 0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200±25	ANISOVICH	00E	SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M154W

 $\eta_2(1645)$  DECAY MODES

NODE=M154215;NODE=M154

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $a_2(1320)\pi$	seen
$\Gamma_2$ $K\bar{K}\pi$	seen
$\Gamma_3$ $K^*\bar{K}$	seen
$\Gamma_4$ $\eta\pi^+\pi^-$	seen
$\Gamma_5$ $a_0(980)\pi$	seen
$\Gamma_6$ $f_2(1270)\eta$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=3;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←

 $\eta_2(1645)$  BRANCHING RATIOS

NODE=M154220

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_1$
<b>0.07±0.03</b>	<sup>1</sup> BARBERIS	97C	OMEG 450 $pp \rightarrow ppK\bar{K}\pi$	

NODE=M154R1  
NODE=M154R1

<sup>1</sup> Using  $2(\pi^+\pi^-)$  data from BARBERIS 97B.

NODE=M154R1;LINKAGE=A

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma_5$
<b>13.1±2.3 OUR AVERAGE</b>				
13.5±4.6	<sup>2</sup> ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$	
13.0±2.7	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M154R3  
NODE=M154R3

<sup>2</sup> Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M154R3;LINKAGE=AN

VALUE	DOCUMENT ID	COMMENT	$\Gamma_6/\Gamma$
not seen	BARBERIS	00B 450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M154R4  
NODE=M154R4

 $\eta_2(1645)$  REFERENCES

NODE=M154

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47958
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202

$\omega(1650)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

See also the  $\omega(1420)$  particle listing.

NODE=M126

NODE=M126

NODE=M126M

NODE=M126M

→ UNCHECKED ←

 $\omega(1650)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1670± 30 OUR ESTIMATE</b>				
1698± 10	267	<sup>1</sup> ACHASOV	20B SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
1651± $3^{+16}_-6$	183k	<sup>2</sup> ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+K^-\pi^0$
1673 $^{+6}_-7$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
1671± 6±10	824	<sup>3</sup> AKHMETSHIN	17A CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
1660± 10	898	<sup>4</sup> ACHASOV	16B SND	1.34–2.00 $e^+e^- \rightarrow \omega\eta$
1680± 10	13.1k	<sup>5</sup> AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1667± 13± 6		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1645± 8	13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
1660± 10± 2		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1770± 50±60	1.2M	<sup>6</sup> ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1619± 5		<sup>7</sup> HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi,$ $\omega\pi\pi$
1700± 20		EUGENIO	01 SPEC	18 $\pi^-p \rightarrow \omega\eta n$
1705± 26	612	<sup>8</sup> AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1820 $^{+190}_-150$		<sup>9</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1840 $^{+100}_-70$		<sup>10</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1780 $^{+170}_-300$		<sup>11</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow K^+K^-$
~ 2100		<sup>12</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606± 9		<sup>13</sup> CLEGG	94 RVUE	
1662± 13	750	<sup>14</sup> ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi,$ $\omega\pi\pi$
1670± 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
1657± 13		CORDIER	81 DM1	$e^+e^- \rightarrow \omega 2\pi$
1679± 34	21	ESPOSITO	80 FRAM	$e^+e^- \rightarrow 3\pi$
1652± 17		COSME	79 OSPK	$e^+e^- \rightarrow 3\pi$

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=4

NODE=M126M;LINKAGE=H

NODE=M126M;LINKAGE=G

NODE=M126M;LINKAGE=F

NODE=M126M;LINKAGE=E

NODE=M126M;LINKAGE=A

NODE=M126M;LINKAGE=VH

NODE=M126M;LINKAGE=AB

NODE=M126M;LINKAGE=KI

NODE=M126M;LINKAGE=L1

NODE=M126M;LINKAGE=L2

NODE=M126M;LINKAGE=L3

NODE=M126M;LINKAGE=L4

NODE=M126M;LINKAGE=AD

NODE=M126M;LINKAGE=AE

<sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass of  $\omega(1420)$  is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of  $\omega(1420)$  to the PDG 18 value of 220 MeV results in  $1694 \pm 9$  MeV measurement.

<sup>2</sup> Could also be  $\rho(1700)$ . Branching ratio  $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (5.3 \pm 0.3^{+0.6}_{-0.5}) \times 10^{-5}$ .

<sup>3</sup> From a fit of the interfering  $\omega(1420)$  and  $\omega(1650)$  with a relative phase of  $\pi$  and other parameters floating.

<sup>4</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ .

<sup>5</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ . See ACHASOV 20A for a further analysis of the  $\pi^+\pi^-\pi^0$  data.

<sup>6</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

<sup>7</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

<sup>8</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

<sup>9</sup> Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.

<sup>10</sup> Using the data from ANTONELLI 92.

<sup>11</sup> Using the data from IVANOV 81 and BISELLO 88B.

<sup>12</sup> Using the data from BISELLO 91C.

<sup>13</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

<sup>14</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.

 $\omega(1650)$  WIDTH

NODE=M126W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

**315± 35 OUR ESTIMATE**

• • • We do not use the following data for averages, fits, limits, etc. • • •

110± 16	267	1	ACHASOV	20B	SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
194± 8 <sup>+</sup> <sub>-7</sub>	183k	2	ABLIKIM	19AQ	BES	$J/\psi \rightarrow K^+K^-\pi^0$
95± 11			ACHASOV	19	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
113± 9± 10	824	3	AKHMETSHIN	17A	CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$
110± 20	898	4	ACHASOV	16B	SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$
310± 30	13.1k	5	AULCHENKO	15A	SND	$1.05-1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
222± 25± 20			AUBERT	07AU	BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
114± 14	13		AUBERT	06D	BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
230± 30± 20			AUBERT,B	04N	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
490 <sup>+</sup> <sub>-150</sub> ±130	1.2M	6	ACHASOV	03D	RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
250± 14		7	HENNER	02	RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
250± 50			EUGENIO	01	SPEC	$18 \pi^-p \rightarrow \omega\eta n$
370± 25	612	8	AKHMETSHIN	00D	CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
113± 20		9	CLEGG	94	RVUE	
280± 24	750	10	ANTONELLI	92	DM2	$1.34-2.4e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
160± 20			ATKINSON	83B	OMEG	$20-70 \gamma p \rightarrow 3\pi X$
136± 46			CORDIER	81	DM1	$e^+e^- \rightarrow \omega 2\pi$
99± 49	21		ESPOSITO	80	FRAM	$e^+e^- \rightarrow 3\pi$
42± 17			COSME	79	OSPK	$e^+e^- \rightarrow 3\pi$

NODE=M126W  
→ UNCHECKED ←

OCCUR=2  
OCCUR=5  
OCCUR=4

<sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass of  $\omega(1420)$  is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of  $\omega(1420)$  to the PDG 18 value of 220 MeV results in  $94 \pm 13$  MeV measurement.

NODE=M126W;LINKAGE=H

<sup>2</sup> Could also be  $\rho(1700)$ . Branching ratio  $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (5.3 \pm 0.3^{+0.6}_{-0.5}) \times 10^{-5}$ .

NODE=M126W;LINKAGE=G

<sup>3</sup> From a fit of the interfering  $\omega(1420)$  and  $\omega(1650)$  with a relative phase of  $\pi$  and other parameters floating.

NODE=M126W;LINKAGE=F

<sup>4</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ .

NODE=M126W;LINKAGE=E

<sup>5</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ . See ACHASOV 20A for a further analysis of the  $\pi^+\pi^-\pi^0$  data.

NODE=M126W;LINKAGE=A

<sup>6</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

NODE=M126W;LINKAGE=VH

<sup>7</sup> Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

NODE=M126W;LINKAGE=AB

<sup>8</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.

NODE=M126W;LINKAGE=KI

<sup>9</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

NODE=M126W;LINKAGE=AD

<sup>10</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.

NODE=M126W;LINKAGE=AE

**$\omega(1650)$  DECAY MODES**

NODE=M126215;NODE=M126

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\pi$	seen
$\Gamma_2$ $\rho(1450)\pi$	seen
$\Gamma_3$ $\omega\pi\pi$	seen
$\Gamma_4$ $\omega\eta$	seen
$\Gamma_5$ $e^+e^-$	seen
$\Gamma_6$ $\pi^0\gamma$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=6  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=3;OUR EST;→ UNCHECKED ←  
DESIG=5

**$\omega(1650) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$**

NODE=M126230

$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma \times \Gamma_5/\Gamma$

NODE=M126G3  
NODE=M126G3

VALUE (units  $10^{-6}$ ) EVTS DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.56 ±0.23	13.1k	1	AULCHENKO	15A	SND	$1.05-1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1.3 ±0.1 ±0.1			AUBERT,B	04N	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1.2 <sup>+0.4</sup> <sub>-0.1</sub> ±0.8	1.2M	2,3	ACHASOV	03D	RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.921±0.230		4,5	CLEGG	94	RVUE	
0.479±0.050	750	6,7	ANTONELLI	92	DM2	$1.34-2.4e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

- <sup>1</sup> From a fit with contributions from  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ , and  $\omega(1650)$ . See ACHASOV 20A for a further analysis of the  $\pi^+\pi^-\pi^0$  data.
- <sup>2</sup> Calculated by us from the cross section at the peak.
- <sup>3</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- <sup>4</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- <sup>5</sup> From the partial and leptonic width given by the authors.
- <sup>6</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.
- <sup>7</sup> From the product of the leptonic width and partial branching ratio given by the authors.

NODE=M126G3;LINKAGE=A

NODE=M126G;LINKAGE=AW  
NODE=M126G;LINKAGE=VH

NODE=M126G;LINKAGE=AD

NODE=M126G;LINKAGE=SE  
NODE=M126G;LINKAGE=AE  
NODE=M126G;LINKAGE=ES

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma \times \Gamma_5/\Gamma$

VALUE (units  $10^{-7}$ ) EVTS DOCUMENT ID TECN COMMENT

NODE=M126G4  
NODE=M126G4

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

7.0 ± 0.5		AUBERT	07AU	BABR	10.6	$e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
4.1 ± 0.9 ± 1.3	1.2M	<sup>1,2</sup> ACHASOV	03D	RVUE	0.44–2.00	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 ± 0.95		<sup>3</sup> AKHMETSHIN	00D	CMD2	1.2–1.38	$e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 ± 0.80		<sup>4,5</sup> CLEGG	94	RVUE		
6.07 ± 0.61	750	<sup>6,7</sup> ANTONELLI	92	DM2	1.34–2.4	$e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

- <sup>1</sup> Calculated by us from the cross section at the peak.
- <sup>2</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- <sup>3</sup> Using the data of AKHMETSHIN 00D and ANTONELLI 92. The  $\rho\pi$  dominance for the energy dependence of the  $\omega(1420)$  and  $\omega(1650)$  width assumed.
- <sup>4</sup> From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- <sup>5</sup> From the partial and leptonic width given by the authors.
- <sup>6</sup> From the combined fit of the  $\rho\pi$  and  $\omega\pi\pi$  final states.
- <sup>7</sup> From the product of the leptonic width and partial branching ratio given by the authors.

NODE=M126G4;LINKAGE=AW  
NODE=M126G4;LINKAGE=VH

NODE=M126G4;LINKAGE=KL

NODE=M126G4;LINKAGE=AD

NODE=M126G4;LINKAGE=SE  
NODE=M126G4;LINKAGE=AE  
NODE=M126G4;LINKAGE=ES

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$

VALUE (units  $10^{-7}$ ) EVTS DOCUMENT ID TECN COMMENT

NODE=M126G5  
NODE=M126G5

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

6.4 ± 0.9	267	<sup>1</sup> ACHASOV	20B	SND		$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
5.62 <sup>+0.45</sup> <sub>-0.42</sub>		ACHASOV	19	SND		$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
4.5 ± 0.3 ± 0.3	824	<sup>2</sup> AKHMETSHIN	17A	CMD3	1.4–2.0	$e^+e^- \rightarrow \omega\eta$
4.4 ± 0.5	898	<sup>3</sup> ACHASOV	16B	SND	1.34–2.00	$e^+e^- \rightarrow \omega\eta$
5.7 ± 0.6	13	AUBERT	06D	BABR	10.6	$e^+e^- \rightarrow \omega\eta\gamma$
< 60 at 90% CL		<sup>4</sup> AKHMETSHIN	03B	CMD2		$e^+e^- \rightarrow \eta\pi^0\gamma$

- <sup>1</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ . The mass of  $\omega(1420)$  is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of  $\omega(1420)$  to the PDG 18 value of 220 MeV results in  $(5.4 \pm 0.6) \times 10^{-7}$  measurement.
- <sup>2</sup> From a fit of the interfering  $\omega(1420)$  and  $\omega(1650)$  with a relative phase of  $\pi$  and other parameters floating. From an alternative fit  $\Gamma(\omega(1650) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1650) \rightarrow e^+e^-) = 51 \pm 3$  eV.
- <sup>3</sup> From a fit with contributions from  $\omega(1420)$ ,  $\omega(1650)$ , and  $\phi(1680)$ .
- <sup>4</sup>  $\omega(1650)$  mass and width fixed at 1700 MeV and 250 MeV, respectively.

NODE=M126G5;LINKAGE=C

NODE=M126G5;LINKAGE=B

NODE=M126G5;LINKAGE=A  
NODE=M126G5;LINKAGE=KH

### $\omega(1650)$ BRANCHING RATIOS

NODE=M126225

$\Gamma(\rho\pi)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

VALUE EVTS DOCUMENT ID TECN COMMENT

NODE=M126R3  
NODE=M126R3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 0.65	1.2M	<sup>1</sup> ACHASOV	03D	RVUE	0.44–2.00	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.380 ± 0.014		<sup>2</sup> HENNER	02	RVUE	1.2–2.0	$e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

- <sup>1</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- <sup>2</sup> Assuming that the  $\omega(1650)$  decays into  $\rho\pi$  and  $\omega\pi\pi$  only.

NODE=M126R;LINKAGE=VH

NODE=M126R;LINKAGE=AC

$\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

NODE=M126R02  
NODE=M126R02

seen ACHASOV 20A SND 1.15–2.00  $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 0.35$	1.2M	<sup>1</sup> ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$0.620 \pm 0.014$		<sup>2</sup> HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

<sup>1</sup> From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the  $\pi^+\pi^-\pi^0$  and ANTONELLI 92 on the  $\omega\pi^+\pi^-$  final states. Supersedes ACHASOV 99E and ACHASOV 02E.

<sup>2</sup> Assuming that the  $\omega(1650)$  decays into  $\rho\pi$  and  $\omega\pi\pi$  only.

NODE=M126R2  
NODE=M126R2

NODE=M126R2;LINKAGE=VH

NODE=M126R2;LINKAGE=AC

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 18$	1.2M	<sup>1,2</sup> ACHASOV	03D	RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$32 \pm 1$		<sup>2</sup> HENNER	02	RVUE 1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

<sup>1</sup> Calculated by us from the cross section at the peak.

<sup>2</sup> Assuming that the  $\omega(1650)$  decays into  $\rho\pi$  and  $\omega\pi\pi$  only.

NODE=M126R4  
NODE=M126R4

NODE=M126R;LINKAGE=AW  
NODE=M126R4;LINKAGE=AC

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen <sup>1</sup> ACHASOV 10D SND 1.075–2.0  $e^+e^- \rightarrow \pi^0\gamma$

<sup>1</sup> From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states  $\omega(1420)$ ,  $\rho(1450)$ ,  $\omega(1650)$ , and  $\rho(1700)$ . The width of the highest mass effective resonance is fixed at 315 MeV.

NODE=M126R00  
NODE=M126R00

NODE=M126R00;LINKAGE=A

 $\omega(1650)$  REFERENCES

NODE=M126

ACHASOV	20A	EPJ C80 993	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60923
ACHASOV	20B	EPJ C80 1008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60924
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59887
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=58239
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57537
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56843
		Translated from ZETF 148 34.			
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>		REFID=49177
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>		REFID=48010
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41867
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21502
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)	REFID=21586
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
ESPOSITO	80	LNC 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=21584
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)	REFID=21475

$\omega_3(1670)$ 

$$I^G(J^{PC}) = 0^-(3^{--})$$

NODE=M045

 $\omega_3(1670)$  MASS

NODE=M045M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1667 ± 4 OUR AVERAGE</b>				
1665.3 ± 5.2 ± 4.5	23400	AMELIN	96 VES	$36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1685 ± 20	60	BAUBILLIER	79 HBC	$8.2 K^- p$ backward
1673 ± 12	430	<sup>1,2</sup> BALTAY	78E HBC	$15 \pi^+ p \rightarrow \Delta 3\pi$
1650 ± 12		CORDEN	78B OMEG	$8-12 \pi^- p \rightarrow N 3\pi$
1669 ± 11	600	<sup>2</sup> WAGNER	75 HBC	$7 \pi^+ p \rightarrow \Delta^{++} 3\pi$
1678 ± 14	500	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 3\pi^0$
1660 ± 13	200	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p \omega \pi^0 \pi^0$
1679 ± 17	200	MATTHEWS	71D DBC	$7.0 \pi^+ n \rightarrow p 3\pi^0$
1670 ± 20		KENYON	69 DBC	$8 \pi^+ n \rightarrow p 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1700	110	<sup>1</sup> CERRADA	77B HBC	$4.2 K^- p \rightarrow \Lambda 3\pi$
1695 ± 20		BARNES	69B HBC	$4.6 K^- p \rightarrow \omega 2\pi X$
1636 ± 20		ARMENISE	68B DBC	$5.1 \pi^+ n \rightarrow p 3\pi^0$

NODE=M045M

OCCUR=2

<sup>1</sup> Phase rotation seen for  $J^P = 3^- \rho\pi$  wave.<sup>2</sup> From a fit to  $I(J^P) = 0(3^-) \rho\pi$  partial wave.NODE=M045M;LINKAGE=E  
NODE=M045M;LINKAGE=P $\omega_3(1670)$  WIDTH

NODE=M045W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>168 ± 10 OUR AVERAGE</b>				
149 ± 19 ± 7	23400	AMELIN	96 VES	$36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
160 ± 80	60	<sup>3</sup> BAUBILLIER	79 HBC	$8.2 K^- p$ backward
173 ± 16	430	<sup>4,5</sup> BALTAY	78E HBC	$15 \pi^+ p \rightarrow \Delta 3\pi$
253 ± 39		CORDEN	78B OMEG	$8-12 \pi^- p \rightarrow N 3\pi$
173 ± 28	600	<sup>3,5</sup> WAGNER	75 HBC	$7 \pi^+ p \rightarrow \Delta^{++} 3\pi$
167 ± 40	500	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 3\pi^0$
122 ± 39	200	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p \omega \pi^0 \pi^0$
155 ± 40	200	<sup>3</sup> MATTHEWS	71D DBC	$7.0 \pi^+ n \rightarrow p 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
90 ± 20		BARNES	69B HBC	$4.6 K^- p \rightarrow \omega 2\pi$
100 ± 40		KENYON	69 DBC	$8 \pi^+ n \rightarrow p 3\pi^0$
112 ± 60		ARMENISE	68B DBC	$5.1 \pi^+ n \rightarrow p 3\pi^0$

NODE=M045W

OCCUR=2

<sup>3</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.<sup>4</sup> Phase rotation seen for  $J^P = 3^- \rho\pi$  wave.<sup>5</sup> From a fit to  $I(J^P) = 0(3^-) \rho\pi$  partial wave.NODE=M045W;LINKAGE=S  
NODE=M045W;LINKAGE=E  
NODE=M045W;LINKAGE=P $\omega_3(1670)$  DECAY MODES

NODE=M045215;NODE=M045

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \rho\pi$	seen
$\Gamma_2 \quad \omega\pi\pi$	seen
$\Gamma_3 \quad b_1(1235)\pi$	possibly seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

 $\omega_3(1670)$  BRANCHING RATIOS

NODE=M045220

$\Gamma(\omega\pi\pi)/\Gamma(\rho\pi)$					$\Gamma_2/\Gamma_1$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.71 ± 0.27	100	DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 5\pi^0$	
$\Gamma(b_1(1235)\pi)/\Gamma(\rho\pi)$					$\Gamma_3/\Gamma_1$
VALUE		DOCUMENT ID	TECN	COMMENT	
possibly seen		DIAZ	74 DBC	$6 \pi^+ n \rightarrow p 5\pi^0$	

NODE=M045R3  
NODE=M045R3NODE=M045R4  
NODE=M045R4

$\Gamma(b_1(1235)\pi)/\Gamma(\omega\pi\pi)$  $\Gamma_3/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

>0.75	68	BAUBILLIER 79	HBC	8.2 $K^- p$ backward
-------	----	---------------	-----	----------------------

NODE=M045R5  
NODE=M045R5

 $\omega_3(1670)$  REFERENCES

AMELIN 96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44649
BAUBILLIER 79	PL 89B 131	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=21522
BALTAY 78E	PRL 40 87	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP	REFID=21520
CORDEN 78B	NP B138 235	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=21269
CERRADA 77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+) JP	REFID=20537
WAGNER 75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP	REFID=20843
DIAZ 74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)	REFID=21248
MATTHEWS 71D	PR D3 2561	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC)	REFID=21515
BARNES 69B	PRL 23 142	V.E. Barnes <i>et al.</i>	(BNL)	REFID=21512
KENYON 69	PRL 23 146	I.R. Kenyon <i>et al.</i>	(BNL, UCND, ORNL)	REFID=20800
ARMENISE 68B	PL 26B 336	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20783

NODE=M045

NODE=M034

 $\pi_2(1670)$ 

$$I^G(J^{PC}) = 1^-(2^-+)$$

 $\pi_2(1670)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**1670.6<sup>+2.9</sup><sub>-1.2</sub> OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

1642	$^{+12}_{-1}$	46M	1	AGHASYAN	18B	COMP	190	$\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$		
1749	$\pm 10$	$\pm 100$	145k	LU	05	B852	18	$\pi^- p \rightarrow \omega \pi^- \pi^0 p$		
1676	$\pm 3$	$\pm 8$		2	CHUNG	02	B852	18.3	$\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
1685	$\pm 10$	$\pm 30$		BARBERIS	01		450	$pp \rightarrow p_f 3\pi^0 p_s$		
1687	$\pm 9$	$\pm 15$		AMELIN	99	VES	37	$\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$		
1669	$\pm 4$			BARBERIS	98B		450	$pp \rightarrow p_f \rho \pi p_s$		
1670	$\pm 4$			BARBERIS	98B		450	$pp \rightarrow p_f f_2(1270) \pi p_s$		
1690	$\pm 14$			3	BERDNIKOV	94	VES	37	$\pi^- A \rightarrow K^+ K^- \pi^- A$	
1710	$\pm 20$	700		ANTIPOV	87	SIGM	-	50	$\pi^- \text{Cu} \rightarrow \mu^+ \mu^- \pi^- \text{Cu}$	
1676	$\pm 6$			3	EVANGELIS...	81	OMEG	-	12	$\pi^- p \rightarrow 3\pi p$
1657	$\pm 14$			3,4	DAUM	80D	SPEC	-	63-94	$\pi p \rightarrow 3\pi X$
1662	$\pm 10$	2000		3	BALTAY	77	HBC	+	15	$\pi^+ p \rightarrow p 3\pi$

NODE=M034M

NODE=M034M

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

1658	$\pm 3$	$^{+24}_{-8}$	420k	5	ALEKSEEV	10	COMP	190	$\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
1730	$\pm 20$			6	AMELIN	95B	VES	36	$\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
1742	$\pm 31$	$\pm 49$			ANTREASYAN	90	CBAL		$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$	
1624	$\pm 21$			2	BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1622	$\pm 35$			7	BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1693	$\pm 28$			8	BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1710	$\pm 20$			9	DAUM	81B	SPEC	-	63,94	$\pi^- p$
1660	$\pm 10$			3	ASCOLI	73	HBC	-	5-25	$\pi^- p \rightarrow p \pi_2$

OCCUR=2

OCCUR=3

<sup>1</sup> Statistical error negligible.

<sup>2</sup> From  $f_2(1270)\pi$  decay.

<sup>3</sup> From a fit to  $J^P = 2^- S$ -wave  $f_2(1270)\pi$  partial wave.

<sup>4</sup> Clear phase rotation seen in  $2^- S$ ,  $2^- P$ ,  $2^- D$  waves. We quote central value and spread of single-resonance fits to three channels.

<sup>5</sup> Superseded by AGHASYAN 2018B.

<sup>6</sup>  $J^{PC}$  ambiguous.

<sup>7</sup> From  $\rho\pi$  decay.

<sup>8</sup> From  $\sigma\pi$  decay.

NODE=M034M;LINKAGE=A

NODE=M034M;LINKAGE=F2

NODE=M034M;LINKAGE=P

NODE=M034M;LINKAGE=D

NODE=M034M;LINKAGE=B


NODE=M034M;LINKAGE=AX

NODE=M034M;LINKAGE=R2

NODE=M034M;LINKAGE=S2

<sup>9</sup> From a two-resonance fit to four  $2^-0^+$  waves. This should not be averaged with all the single resonance fits.

NODE=M034M;LINKAGE=L



**$\pi_2(1670)$  WIDTH**

NODE=M034W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>258<sup>+8</sup><sub>-9</sub></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.			
311 <sup>+12</sup> <sub>-23</sub>	46M	<sup>10</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
408 $\pm$ 60 $\pm$ 250	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
254 $\pm$ 3 $\pm$ 31		<sup>11</sup> CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
265 $\pm$ 30 $\pm$ 40		BARBERIS	01		450 $pp \rightarrow p_f 3\pi^0 p_s$
168 $\pm$ 43 $\pm$ 53		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
268 $\pm$ 15		BARBERIS	98B		450 $pp \rightarrow p_f p \pi p_s$
256 $\pm$ 15		BARBERIS	98B		450 $pp \rightarrow p_f f_2(1270) \pi p_s$
190 $\pm$ 50		<sup>12</sup> BERDNIKOV	94	VES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
170 $\pm$ 80	700	ANTIPOV	87	SIGM	- 50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
260 $\pm$ 20		<sup>12</sup> EVANGELIS...	81	OMEG	- 12 $\pi^- p \rightarrow 3\pi p$
219 $\pm$ 20		<sup>12,13</sup> DAUM	80D	SPEC	- 63-94 $\pi p \rightarrow 3\pi X$
285 $\pm$ 60	2000	<sup>12</sup> BALTAY	77	HBC	+ 15 $\pi^+ p \rightarrow p 3\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
271 $\pm$ 9 <sup>+22</sup> <sub>-24</sub>	420k	<sup>14</sup> ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
310 $\pm$ 20		<sup>15</sup> AMELIN	95B	VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
236 $\pm$ 49 $\pm$ 36		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
304 $\pm$ 22		<sup>11</sup> BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
404 $\pm$ 108		<sup>16</sup> BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
330 $\pm$ 90		<sup>17</sup> BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
312 $\pm$ 50		<sup>18</sup> DAUM	81B	SPEC	- 63,94 $\pi^- p$
270 $\pm$ 60		<sup>12</sup> ASCOLI	73	HBC	- 5-25 $\pi^- p \rightarrow p \pi_2$

NODE=M034W

NODE=M034W

OCCUR=2

OCCUR=2

OCCUR=3



- <sup>10</sup> Statistical error negligible.  
<sup>11</sup> From  $f_2(1270)\pi$  decay.  
<sup>12</sup> From a fit to  $J^P = 2^- f_2(1270)\pi$  partial wave.  
<sup>13</sup> Clear phase rotation seen in  $2^- S, 2^- P, 2^- D$  waves. We quote central value and spread of single-resonance fits to three channels.  
<sup>14</sup> Superseded by AGHASYAN 2018B.  
<sup>15</sup>  $J^{PC}$  ambiguous.  
<sup>16</sup> From  $\rho\pi$  decay.  
<sup>17</sup> From  $\sigma\pi$  decay.  
<sup>18</sup> From a two-resonance fit to four  $2^- 0^+$  waves. This should not be averaged with all the single resonance fits.

NODE=M034W;LINKAGE=A  
 NODE=M034W;LINKAGE=F2  
 NODE=M034W;LINKAGE=P  
 NODE=M034W;LINKAGE=D  
  
 NODE=M034W;LINKAGE=B  
 NODE=M034W;LINKAGE=AX  
 NODE=M034W;LINKAGE=R2  
 NODE=M034W;LINKAGE=S2  
 NODE=M034W;LINKAGE=L

### $\pi_2(1670)$ DECAY MODES

NODE=M034215;NODE=M034

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $3\pi$	(95.8±1.4) %	
$\Gamma_2$ $\pi^+\pi^-\pi^0$		
$\Gamma_3$ $\pi^0\pi^0\pi^0$		
$\Gamma_4$ $f_2(1270)\pi$	(56.3±3.2) %	
$\Gamma_5$ $\rho\pi$	(31 ±4 ) %	
$\Gamma_6$ $\sigma\pi$	(10 ±4 ) %	
$\Gamma_7$ $\pi(\pi\pi)_S$ -wave	( 8.7±3.4) %	
$\Gamma_8$ $\pi^\pm\pi^+\pi^-$	(53 ±4 ) %	
$\Gamma_9$ $K\bar{K}^*(892)+$ c.c.	( 4.2±1.4) %	
$\Gamma_{10}$ $\omega\rho$	( 2.7±1.1) %	
$\Gamma_{11}$ $\pi^\pm\gamma$	( 7.0±1.2) × 10 <sup>-4</sup>	
$\Gamma_{12}$ $\gamma\gamma$	< 2.8 × 10 <sup>-7</sup>	90%
$\Gamma_{13}$ $\eta\pi$	< 5 %	
$\Gamma_{14}$ $\pi^\pm 2\pi^+ 2\pi^-$	< 5 %	
$\Gamma_{15}$ $\rho(1450)\pi$	< 3.6 × 10 <sup>-3</sup>	97.7%
$\Gamma_{16}$ $b_1(1235)\pi$	< 1.9 × 10 <sup>-3</sup>	97.7%
$\Gamma_{17}$ $\eta 3\pi$		
$\Gamma_{18}$ $f_1(1285)\pi$	possibly seen	
$\Gamma_{19}$ $a_2(1320)\pi$	not seen	

DESIG=20  
 DESIG=22  
 DESIG=23  
 DESIG=8  
 DESIG=2  
 DESIG=13  
 DESIG=11  
 DESIG=10  
 DESIG=5  
 DESIG=14  
 DESIG=27  
 DESIG=12  
 DESIG=3  
 DESIG=4  
 DESIG=15  
 DESIG=16  
 DESIG=24  
 DESIG=25  
 DESIG=26

### CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 1.9$  for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_5$	-53		
$x_7$	-29	-59	
$x_9$	-8	-21	-9
	$x_4$	$x_5$	$x_7$

### $\pi_2(1670)$ PARTIAL WIDTHS

NODE=M034217

$\Gamma(\pi^\pm\gamma)$

$\Gamma_{11}$

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>181±11±27</b>	19 ADOLPH	14	COMP	- 190 $\pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$

NODE=M034W2  
 NODE=M034W2

- <sup>19</sup> Primakoff reaction. Assumes incoherent  $f_2(1270)\pi$  contribution to  $3\pi$  final state and uses  $B(\pi_2(1670) \rightarrow f_2\pi) = 56\%$ .

NODE=M034W2;LINKAGE=AD

$\Gamma(\gamma\gamma)$  $\Gamma_{12}$ 

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.072	90	20 ACCIARRI	97T L3		$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.19	90	20 ALBRECHT	97B ARG		$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.41 ±0.23±0.28		ANTREASYAN 90	CBAL 0		$e^+e^- \rightarrow e^+e^-\pi^0\pi^0\pi^0$
0.8 ±0.3 ±0.12		21 BEHREND	90C CELL 0		$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.3 ±0.3 ±0.2		22 BEHREND	90C CELL 0		$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$

NODE=M034W1  
NODE=M034W1

OCCUR=2

<sup>20</sup>Decaying into  $f_2(1270)\pi$  and  $\rho\pi$ .<sup>21</sup>Constructive interference between  $f_2(1270)\pi, \rho\pi$  and background.<sup>22</sup>Incoherent Ansatz.

NODE=M034W1;LINKAGE=QQ

NODE=M034W1;LINKAGE=C

NODE=M034W1;LINKAGE=G

 $\pi_2(1670) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M034230

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_2\Gamma_{12}/\Gamma$ 

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	95	23 SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

NODE=M034G01  
NODE=M034G01<sup>23</sup>From analysis of L3 data at 183–209 GeV.

NODE=M034G01;LINKAGE=SC

 $\pi_2(1670) \text{ BRANCHING RATIOS}$ 

NODE=M034220

 $\Gamma(3\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma = (\Gamma_4+\Gamma_5+\Gamma_7)/\Gamma$ 

VALUE	DOCUMENT ID
<b>0.958±0.014 OUR FIT</b>	

NODE=M034R20  
NODE=M034R20 $\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$  $\Gamma_3/\Gamma_2$ 

VALUE	DOCUMENT ID	COMMENT
<b>0.29±0.03±0.05</b>	BARBERIS 01	450 $\rho\rho \rightarrow \rho_f 3\pi^0 \rho_s$

NODE=M034R21  
NODE=M034R21 $\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi)$  $\Gamma_5/0.565\Gamma_4$ (With  $f_2(1270) \rightarrow \pi^+\pi^-$ .)

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.97±0.09 OUR AVERAGE</b>	Error includes scale factor of 1.9.		
0.76±0.07±0.10	CHUNG 02	B852	18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$
1.01±0.05	BARBERIS 98B		450 $\rho\rho \rightarrow \rho_f \pi^+\pi^-\pi^0 \rho_s$

NODE=M034R16  
NODE=M034R16  
NODE=M034R16 $\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi)$  $\Gamma_6/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.17±0.02±0.07</b>	CHUNG 02	B852	18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$

NODE=M034R15  
NODE=M034R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.24±0.10 <sup>24,25</sup>BAKER 99 SPEC 1.94  $\bar{p}p \rightarrow 4\pi^0$  $\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$  $\frac{1}{2}\Gamma_5/\Gamma_8 = \frac{1}{2}\Gamma_5/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.29±0.04 OUR FIT</b>				
<b>0.29±0.05</b>	<sup>26</sup> DAUM	81B	SPEC	63.94 $\pi^-p$

NODE=M034R2  
NODE=M034R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.3 BARTSCH 68 HBC + 8  $\pi^+p \rightarrow 3\pi p$  $0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$  $0.565\Gamma_4/\Gamma_8 = 0.565\Gamma_4/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ (With  $f_2(1270) \rightarrow \pi^+\pi^-$ .)

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.604±0.035 OUR FIT</b>				
<b>0.60 ±0.05 OUR AVERAGE</b>	Error includes scale factor of 1.3.			

NODE=M034R3

NODE=M034R3

NODE=M034R3

0.61 ±0.04 <sup>26</sup>DAUM 81B SPEC 63.94  $\pi^-p$ 0.76 <sup>+0.24</sup><sub>-0.34</sub> ARMENISE 69 DBC + 5.1  $\pi^+d \rightarrow d3\pi$ 0.35 ±0.20 BALTAY 68 HBC + 7–8.5  $\pi^+p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.59 BARTSCH 68 HBC + 8  $\pi^+p \rightarrow 3\pi p$

**0.624 $\Gamma(\pi(\pi\pi)_{S\text{-wave}})/\Gamma(\pi^\pm\pi^+\pi^-)$** 

**0.624 $\Gamma_7/\Gamma_8 = 0.624\Gamma_7/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$**

(With  $(\pi\pi)_{S\text{-wave}} \rightarrow \pi^+\pi^-$ .)

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.10<math>\pm</math>0.04 OUR FIT</b>			
<b>0.10<math>\pm</math>0.05</b>	26 DAUM	81B SPEC	63,94 $\pi^- p$

NODE=M034R11  
 NODE=M034R11  
 NODE=M034R11

 **$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(f_2(1270)\pi)$**  **$\Gamma_9/\Gamma_4$** 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.075<math>\pm</math>0.025 OUR FIT</b>				
<b>0.075<math>\pm</math>0.025</b>	27 ARMSTRONG	82B OMEG	-	16 $\pi^- p \rightarrow K^+ K^- \pi^- p$

NODE=M034R13  
 NODE=M034R13

 **$\Gamma(\omega\rho)/\Gamma_{\text{total}}$**  **$\Gamma_{10}/\Gamma$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.027<math>\pm</math>0.004<math>\pm</math>0.010</b>	28 AMELIN	99 VES	37 $\pi^- A \rightarrow \omega\pi^- \pi^0 A^*$

NODE=M034R17  
 NODE=M034R17

 **$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$** 

**$\Gamma_{13}/\Gamma_8 = \Gamma_{13}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$**

(All  $\eta$  decays.)

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.09</b>	BALTAY	68 HBC	+	7-8.5 $\pi^+ p$
<b>&lt;0.10</b>	CRENNELL	70 HBC	-	6 $\pi^- p \rightarrow f_2\pi^- N$

NODE=M034R5  
 NODE=M034R5  
 NODE=M034R5

 **$\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$** 

**$\Gamma_{14}/\Gamma_8 = \Gamma_{14}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.10</b>	CRENNELL	70 HBC	-	6 $\pi^- p \rightarrow f_2\pi^- N$
<b>&lt;0.1</b>	BALTAY	68 HBC	+	7,8.5 $\pi^+ p$

NODE=M034R6  
 NODE=M034R6

 **$\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$**  **$\Gamma_{15}/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0036</b>	97.7	AMELIN	99 VES	37 $\pi^- A \rightarrow \omega\pi^- \pi^0 A^*$

NODE=M034R18  
 NODE=M034R18

 **$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$**  **$\Gamma_{16}/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0019</b>	97.7	AMELIN	99 VES	37 $\pi^- A \rightarrow \omega\pi^- \pi^0 A^*$

NODE=M034R19  
 NODE=M034R19

 **$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$**  **$\Gamma_{18}/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	69k	KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M034R23  
 NODE=M034R23

 **$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$**  **$\Gamma_{19}/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	69k	KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M034R24  
 NODE=M034R24

**D-wave/S-wave RATIO FOR  $\pi_2(1670) \rightarrow f_2(1270)\pi$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.18<math>\pm</math>0.06</b>	24 BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$
<b>0.22<math>\pm</math>0.10</b>	26 DAUM	81B SPEC	63,94 $\pi^- p$

NODE=M034R14  
 NODE=M034R14

**F-wave/P-wave RATIO FOR  $\pi_2(1670) \rightarrow \rho\pi$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.72<math>\pm</math>0.07<math>\pm</math>0.14</b>	CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M034R22  
 NODE=M034R22

24 Using preliminary CBAR data.

25 With the  $\sigma\pi$  in  $L=2$  and the  $f_2(1270)\pi$  in  $L=0$ .26 From a two-resonance fit to four  $2^-0^+$  waves.27 From a partial-wave analysis of  $K^+K^-\pi^-$  system.28 Normalized to the  $B(\pi_2(1670) \rightarrow f_2\pi)$ .

NODE=M034R;LINKAGE=BK  
 NODE=M034R15;LINKAGE=A  
 NODE=M034R;LINKAGE=L  
 NODE=M034R13;LINKAGE=M  
 NODE=M034R;LINKAGE=DM

$\pi_2(1670)$  REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>	
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)
		Translated from YAF 62 487.		
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)
ANTREASYAN	90	ZPHY C48 561	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)
ANTIPOV	87	EPL 4 403	Y.M. Antipov <i>et al.</i>	(SERP, JINR, INRM+)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
		Translated from YAF 41 1223.		
ARMSTRONG	82B	NP B202 1	T.A. Armstrong, B. Baccari	(AACH3, BARI, BONN+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
		Also NP B186 594	C. Evangelista	
DAUM	80D	PL 89B 285	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
ASCOLI	73	PR D7 669	G. Ascoli	(ILL, TINTO, GENO, HAMB, MILA+) JP
CRENNELL	70	PRL 24 781	D.J. Crennell <i>et al.</i>	(BNL)
ARMENISE	69	LCN 2 501	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
BARTSCH	68	NP B7 345	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN) JP

NODE=M034

REFID=59471  
 REFID=55911  
 REFID=53356  
 REFID=51186  
 REFID=50459  
 REFID=49773  
 REFID=48837  
 REFID=48324  
 REFID=46910  
 REFID=46888  
 REFID=46345  
 REFID=45761  
 REFID=45418  
 REFID=44433  
 REFID=44073  
 REFID=41372  
 REFID=41356  
 REFID=40004  
 REFID=47490

 $\phi(1680)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M067

 $\phi(1680)$  MASS

NODE=M067205

 $e^+e^-$  PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1680±20 OUR ESTIMATE</b>				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1683± 7± 9		<sup>1</sup> ZHU	23 BELL	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
1673± 5		<sup>2</sup> ABLIKIM	22L BES3	2.0–3.08 $e^+e^- \rightarrow K^+K^-\pi^0$
$1680^{+12}_{-13} \pm 21$	1.8k	<sup>3</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+K^-\eta$
1662±20		<sup>4</sup> ACHASOV	20C SND	1.3–2.0 $e^+e^- \rightarrow K^+K^-\pi^0$
$1641^{+24}_{-18}$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
1667± 5±11	3k	<sup>5</sup> IVANOV	19A CMD3	1.59–2.007 $e^+e^- \rightarrow K^+K^-\eta$
1700±23	2k	<sup>6</sup> ACHASOV	18A SND	1.3–2.0 $e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$
1674±12± 6	6.2k	<sup>7</sup> LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1733±10±10		<sup>8</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
1689± 7±10	4.8k	<sup>9</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
1709±20±43		<sup>10</sup> AUBERT	08S BABR	10.6 $e^+e^- \rightarrow$ hadrons
1623±20	948	<sup>11</sup> AKHMETSHIN	03 CMD2	1.05–1.38 $e^+e^- \rightarrow K_L^0 K_S^0$
~ 1500		<sup>12</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0, \omega\pi^+\pi^-,$ $K^+K^-$
~ 1900		<sup>13</sup> ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1700±20		<sup>14</sup> CLEGG	94 RVUE	$e^+e^- \rightarrow K^+K^-, K_S^0 K\pi$
1657±27	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655±17		<sup>15</sup> BISELLO	88B DM2	$e^+e^- \rightarrow K^+K^-$
1680±10		<sup>16</sup> BUON	82 DM1	$e^+e^- \rightarrow$ hadrons
1677±12		<sup>17</sup> MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K\pi$

NODE=M067M1  
 NODE=M067M1

→ UNCHECKED ←

OCCUR=4

<sup>1</sup> From a fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution.

NODE=M067M1;LINKAGE=H

<sup>2</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

NODE=M067M1;LINKAGE=G

<sup>3</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+K^-\eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$ .

NODE=M067M1;LINKAGE=E

<sup>4</sup> From a fit using a vector meson dominance model with contribution from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ ,  $\rho(1450)$ .

NODE=M067M1;LINKAGE=F

- <sup>5</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.  
<sup>6</sup> Assuming the  $K\bar{K}^*(892) + c.c.$  dynamics. Systematic uncertainties not estimated.  
<sup>7</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .  
<sup>8</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.  
<sup>9</sup> From a fit with two incoherent Breit-Wigners.  
<sup>10</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + c.c.$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.  
<sup>11</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.  
<sup>12</sup> Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92.  
<sup>13</sup> Using the data from BISELLO 91C.  
<sup>14</sup> Using BISELLO 88B and MANE 82 data.  
<sup>15</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$  assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitation.  
<sup>16</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0 K_L^0$ ,  $K_S^0 K^\pm\pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.  
<sup>17</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

NODE=M067M1;LINKAGE=D  
 NODE=M067M1;LINKAGE=C  
 NODE=M067M1;LINKAGE=B  
  
 NODE=M067M1;LINKAGE=A  
 NODE=M067M1;LINKAGE=SH  
 NODE=M067M1;LINKAGE=AU  
  
 NODE=M067M;LINKAGE=HK  
  
 NODE=M067M1;LINKAGE=L1  
  
 NODE=M067M1;LINKAGE=L4  
 NODE=M067M;LINKAGE=A  
 NODE=M067M;LINKAGE=E  
  
 NODE=M067M;LINKAGE=C  
  
 NODE=M067M;LINKAGE=D  
  
 NODE=M067M3  
 NODE=M067M3

### $p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
$1700 \pm 8$	<sup>1</sup> AMSLER	06 CBAR	$0.9 p\bar{p} \rightarrow K^+ K^- \pi^0$
<sup>1</sup> Could also be $\rho(1700)$ .			

NODE=M067M3;LINKAGE=AM

### $\phi(1680)$ WIDTH

NODE=M067210

### $e^+e^-$ PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>150 \pm 50</math></b>	<b>OUR ESTIMATE</b>	This is only an educated guess; the error given is larger than the error on the average of the published values.		

NODE=M067W1  
 NODE=M067W1  
 → UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

$149 \pm 12 \pm 13$	<sup>1</sup> ZHU	23 BELL	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
$172 \pm 8$	<sup>2</sup> ABLIKIM	22L BES3	$2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$
$185^{+30+25}_{-26-47}$ 1.8k	<sup>3</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+K^-\eta$
$159 \pm 32$	<sup>4</sup> ACHASOV	20C SND	$1.3-2.0 e^+e^- \rightarrow K^+K^-\pi^0$
$103^{+26}_{-24}$	ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
$176 \pm 23 \pm 38$ 3k	<sup>5</sup> IVANOV	19A CMD3	$1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$
$300 \pm 50$ 2k	<sup>6</sup> ACHASOV	18A SND	$1.3-2.0 e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$
$165 \pm 38 \pm 70$ 6.2k	<sup>7</sup> LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
$300 \pm 15 \pm 37$	<sup>8</sup> LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
$211 \pm 14 \pm 19$ 4.8k	<sup>9</sup> SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
$322 \pm 77 \pm 160$	<sup>10</sup> AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \text{hadrons}$
$139 \pm 60$ 948	<sup>11</sup> AKHMETSHIN	03 CMD2	$1.05-1.38 e^+e^- \rightarrow K_L^0 K_S^0$
$300 \pm 60$	<sup>12</sup> CLEGG	94 RVUE	$e^+e^- \rightarrow K^+K^-, K_S^0 K\pi$
$146 \pm 55$ 367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm\pi^\mp$
$207 \pm 45$	<sup>13</sup> BISELLO	88B DM2	$e^+e^- \rightarrow K^+K^-$
$185 \pm 22$	<sup>14</sup> BUON	82 DM1	$e^+e^- \rightarrow \text{hadrons}$
$102 \pm 36$	<sup>15</sup> MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K\pi$

- <sup>1</sup> From a fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution.  
<sup>2</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.  
<sup>3</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+K^-\eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$ .  
<sup>4</sup> From a fit using a vector meson dominance model with contribution from  $\rho(770)$ ,  $\omega(782)$ ,  $\phi(1020)$ ,  $\omega(1420)$ ,  $\rho(1450)$ .  
<sup>5</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.  
<sup>6</sup> Assuming the  $K\bar{K}^*(892) + c.c.$  dynamics. Systematic uncertainties not estimated.  
<sup>7</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .  
<sup>8</sup> Using events with  $\pi\pi$  invariant mass less than 0.85 GeV.

NODE=M067W1;LINKAGE=H  
 NODE=M067W1;LINKAGE=G  
  
 NODE=M067W1;LINKAGE=E  
  
 NODE=M067W1;LINKAGE=F  
  
 NODE=M067W1;LINKAGE=D  
 NODE=M067W1;LINKAGE=C  
 NODE=M067W1;LINKAGE=B  
  
 NODE=M067W1;LINKAGE=A

<sup>9</sup> From a fit with two incoherent Breit-Wigners.

<sup>10</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + c.c.$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

<sup>11</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known.

<sup>12</sup> Using BISELLO 88B and MANE 82 data.

<sup>13</sup> From global fit including  $\rho$ ,  $\omega$ ,  $\phi$  and  $\rho(1700)$

<sup>14</sup> From global fit of  $\rho$ ,  $\omega$ ,  $\phi$  and their radial excitations to channels  $\omega\pi^+\pi^-$ ,  $K^+K^-$ ,  $K_S^0K_L^0$ ,  $K_S^0K^\pm\pi^\mp$ . Assume mass 1570 MeV and width 510 MeV for  $\rho$  radial excitations, mass 1570 and width 500 MeV for  $\omega$  radial excitation.

<sup>15</sup> Fit to one channel only, neglecting interference with  $\omega$ ,  $\rho(1700)$ .

NODE=M067W1;LINKAGE=SH

NODE=M067W1;LINKAGE=AU

NODE=M067W;LINKAGE=HK

NODE=M067W;LINKAGE=A

NODE=M067W;LINKAGE=E

NODE=M067W;LINKAGE=C

NODE=M067W;LINKAGE=D

NODE=M067W3

NODE=M067W3

## $p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

143±24	<sup>1</sup> AMSLER	06	CBAR 0.9 $p\bar{p} \rightarrow K^+K^-\pi^0$
--------	---------------------	----	---

<sup>1</sup> Could also be  $\rho(1700)$ .

NODE=M067W3;LINKAGE=AM

## $\phi(1680)$ DECAY MODES

NODE=M067215;NODE=M067

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\bar{K}^*(892) + c.c.$	seen
$\Gamma_2$ $K_S^0K\pi$	seen
$\Gamma_3$ $K\bar{K}$	seen
$\Gamma_4$ $K_L^0K_S^0$	
$\Gamma_5$ $e^+e^-$	seen
$\Gamma_6$ $\omega\pi\pi$	not seen
$\Gamma_7$ $\phi\pi\pi$	
$\Gamma_8$ $K^+K^-\pi^+\pi^-$	seen
$\Gamma_9$ $\eta\phi$	seen
$\Gamma_{10}$ $K^+K^-\eta$	
$\Gamma_{11}$ $\eta\gamma$	seen
$\Gamma_{12}$ $K^+K^-\pi^0$	
$\Gamma_{13}$ $f_2'(1525)\gamma$	not seen

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=11

DESIG=12;OUR EVAL;→ UNCHECKED ←

DESIG=10

DESIG=14

DESIG=13

DESIG=2

DESIG=15

## $\phi(1680)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M067220

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the integrated cross section into channel (I) in  $e^+e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(I)$  or the branching ratio  $\Gamma(I)/\text{total}$ .

NODE=M067220

$\Gamma(K_L^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_4\Gamma_5/\Gamma$
--	---------------------------

NODE=M067A00

NODE=M067A00

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.3±2.4±6.2	6.2k	<sup>1</sup> LEES	14H	BABR $e^+e^- \rightarrow K_S^0K_L^0\gamma$
--------------	------	-------------------	-----	--

<sup>1</sup> Using a vector meson dominance model with contribution from  $\phi(1020)$ ,  $\phi(1680)$ , and higher mass excitations of  $\rho(770)$  and  $\omega(782)$ .

NODE=M067A00;LINKAGE=A

$\Gamma(\phi\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_7\Gamma_5/\Gamma$
--	---------------------------

NODE=M067G02

NODE=M067G02

VALUE ( $10^{-2}$ keV)	DOCUMENT ID	TECN	COMMENT
------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2±0.2±0.3	LEES	12F	BABR 10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
-------------	------	-----	---

$\Gamma(\eta\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_9\Gamma_5/\Gamma$
--	---------------------------

NODE=M067R00

NODE=M067R00

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

122±6±13		<sup>1</sup> ZHU	23	BELL $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
----------	--	------------------	----	---

94±13±15	3k	<sup>2</sup> IVANOV	19A	CMD3 1.59-2.007 $e^+e^- \rightarrow K^+K^-\eta$
----------	----	---------------------	-----	---

<sup>1</sup> From a solution of the fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution. Other solutions with equal fit quality give  $(219 \pm 15 \pm 18)$  eV,  $(163 \pm 11 \pm 13)$  eV and  $(203 \pm 12 \pm 18)$  eV.

<sup>2</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

NODE=M067R00;LINKAGE=B

NODE=M067R00;LINKAGE=A

$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$ 

NODE=M067223

This combination of a branching ratio into channel ( $i$ ) and branching ratio into  $e^+e^-$  is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into ( $i$ ) or  $e^+e^-$ .

NODE=M067223

$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units  $10^{-6}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

NODE=M067G5  
NODE=M067G5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.131 ± 0.059    948    <sup>1</sup> AKHMETSHIN 03    CMD2    1.05–1.38  $e^+e^- \rightarrow K_L^0 K_S^0$

<sup>1</sup> From the combined fit of AKHMETSHIN 03 and MANE 81 also including  $\rho$ ,  $\omega$ , and  $\phi$ . Neither isospin nor flavor structure known. Recalculated by us.

NODE=M067G;LINKAGE=GK

$$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units  $10^{-6}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

NODE=M067G6  
NODE=M067G6

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.15 ± 0.16 ± 0.01    <sup>1</sup> AUBERT    08S    BABR    10.6  $e^+e^- \rightarrow K\bar{K}^*(892)\gamma +$

3.29 ± 1.57    367    <sup>2</sup> BISELLO    91C    DM2    1.35–2.40  $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

NODE=M067G6;LINKAGE=AU

<sup>2</sup> Recalculated by us with the published value of  $B(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)$ .

NODE=M067G;LINKAGE=GL

$$\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units  $10^{-7}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

NODE=M067G01  
NODE=M067G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.86 ± 0.14 ± 0.21    4.8k    <sup>1</sup> SHEN    09    BELL    10.6  $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

<sup>1</sup> Multiplied by 3/2 to take into account the  $\phi\pi^0\pi^0$  mode. Using  $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$ .

NODE=M067G01;LINKAGE=SH

$$\Gamma(\eta\phi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \Gamma_5/\Gamma$$

VALUE (units  $10^{-7}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

NODE=M067G7  
NODE=M067G7

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.64 <sup>+1.74</sup> <sub>-1.80</sub>    ACHASOV    19    SND     $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$

5.3 ± 0.6 ± 0.9    3k    <sup>1</sup> IVANOV    19A    CMD3    1.59–2.007  $e^+e^- \rightarrow$   
 $K^+K^-\eta$

4.3 ± 1.0 ± 0.9    <sup>2</sup> AUBERT    08S    BABR    10.6  $e^+e^- \rightarrow \phi\eta\gamma$

<sup>1</sup> From a fit with coherent interference of the  $\phi(1680)$  with a non-resonant contribution.

NODE=M067G7;LINKAGE=A

<sup>2</sup> From the simultaneous fit to the  $K\bar{K}^*(892) + \text{c.c.}$  and  $\phi\eta$  data from AUBERT 08S using the results of AUBERT 07AK.

NODE=M067G7;LINKAGE=AU

 $\phi(1680)$  BRANCHING RATIOS

NODE=M067225

$$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K\pi) \quad \Gamma_1/\Gamma_2$$

VALUE    DOCUMENT ID    TECN    COMMENT

NODE=M067R3  
NODE=M067R3

dominant    MANE    82    DM1     $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892) + \text{c.c.}) \quad \Gamma_3/\Gamma_1$$

VALUE    DOCUMENT ID    TECN    COMMENT

NODE=M067R2  
NODE=M067R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 ± 0.01    BUON    82    DM1     $e^+e^-$

$$\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892) + \text{c.c.}) \quad \Gamma_6/\Gamma_1$$

VALUE    DOCUMENT ID    TECN    COMMENT

NODE=M067R1  
NODE=M067R1

<0.10    BUON    82    DM1     $e^+e^-$

$$\Gamma(\eta\phi)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE    EVTS    DOCUMENT ID    TECN    COMMENT

NODE=M067R01  
NODE=M067R01

seen    35    <sup>1</sup> ACHASOV    14    SND    1.15–2.00  $e^+e^- \rightarrow \eta\gamma$

<sup>1</sup> From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

NODE=M067R01;LINKAGE=A

$\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892)+c.c.)$  $\Gamma_9/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\approx 0.37$  <sup>1</sup> AUBERT 08S BABR 10.6  $e^+e^- \rightarrow$  hadrons

<sup>1</sup> From the fit including data from AUBERT 07AK.

NODE=M067R5  
NODE=M067R5

NODE=M067R5;LINKAGE=AU

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**seen** 35 <sup>1</sup> ACHASOV 14 SND 1.15–2.00  $e^+e^- \rightarrow \eta\gamma$

<sup>1</sup> From a phenomenological model based on vector meson dominance with  $\rho(1450)$  and  $\phi(1680)$  masses and widths from the PDG 12.

NODE=M067R02  
NODE=M067R02

NODE=M067R02;LINKAGE=A

 $\Gamma(f_2'(1525)\gamma)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**not seen** <sup>1</sup> ACHASOV 22 SND 1.17–2.00  $e^+e^- \rightarrow \eta\eta\gamma$

<sup>1</sup> The 90% CL upper limit on the Born cross sections  $\sigma(e^+e^- \rightarrow \phi(1680) \rightarrow f_2'(1525)\gamma \rightarrow \eta\eta\gamma)$  and  $\sigma(e^+e^- \rightarrow \rho(1700) \rightarrow f_0(1500)\gamma \rightarrow \eta\eta\gamma)$  is 10.6 pb.

NODE=M067R03  
NODE=M067R03

NODE=M067R03;LINKAGE=A

 $\phi(1680)$  REFERENCES

NODE=M067

ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)	REFID=61911
ABLIKIM	22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61649
ACHASOV	22	EPJ C82 168	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61655
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
ACHASOV	20C	EPJ C80 1139	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60935
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59887
IVANOV	19A	PL B798 134946	V.L. Ivanov <i>et al.</i>	(CMD-3 Collab.)	REFID=60133
ACHASOV	18A	PR D97 032011	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58883
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55912
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49172
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin		REFID=48827
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41867
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)	REFID=21494
MANE	82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)	REFID=21590
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORSAY)	REFID=21588



$\rho_3(1690)$ 

$$I^G(J^{PC}) = 1^+(3^{--})$$

NODE=M015

 **$\rho_3(1690)$  MASS**

NODE=M015205

VALUE (MeV)

DOCUMENT ID

**1688.8±2.1 OUR AVERAGE** Includes data from the 5 datablocks that follow this one.

NODE=M015M

**2 $\pi$  MODE**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015M1  
NODE=M015M1**1686± 4 OUR AVERAGE**

1677±14		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow 2\pi p$
1679±11	476	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow \pi^+ \pi^- n$
1678±12	175	<sup>1</sup> ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
1690± 7	600	<sup>1</sup> ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
1693± 8		<sup>2</sup> GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1678±12		MATTHEWS	71C	DBC	0	7 $\pi^+ N$
1734±10		<sup>3</sup> CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow n2\pi$
1692±12		<sup>2,4</sup> ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1737±23		ARMENISE	70	DBC	0	9 $\pi^+ N$
1650±35	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
1687±21		STUNTEBECK	70	HDBC	0	8 $\pi^- p, 5.4 \pi^+ d$
1683±13		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
1670±30		GOLDBERG	65	HBC	0	6 $\pi^+ d, 8 \pi^- p$

<sup>1</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.<sup>2</sup> Uses same data as HYAMS 75.<sup>3</sup> From a phase shift solution containing a  $f'_2(1525)$  width two times larger than the  $K\bar{K}$  result.<sup>4</sup> From phase-shift analysis. Error takes account of spread of different phase-shift solutions.NODE=M015M1;LINKAGE=E  
NODE=M015M1;LINKAGE=G  
NODE=M015M1;LINKAGE=M

NODE=M015M1;LINKAGE=I

 **$K\bar{K}$  AND  $K\bar{K}\pi$  MODES**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015M2  
NODE=M015M2**1696± 4 OUR AVERAGE**

1699± 5		ALPER	80	CNTR	0	62 $\pi^- p \rightarrow K^+ K^- n$
1698±12	6k	<sup>5,6</sup> MARTIN	78D	SPEC		10 $\pi p \rightarrow K_S^0 K^- p$
1692± 6		BLUM	75	ASPK	0	18.4 $\pi^- p \rightarrow nK^+ K^-$
1690±16		ADERHOLZ	69	HBC	+	8 $\pi^+ p \rightarrow K\bar{K}\pi$

••• We do not use the following data for averages, fits, limits, etc. •••

1694± 8 <sup>7</sup> COSTA 80 OMEG 10  $\pi^- p \rightarrow K^+ K^- n$ <sup>5</sup> From a fit to  $J^P = 3^-$  partial wave.<sup>6</sup> Systematic error on mass scale subtracted.<sup>7</sup> They cannot distinguish between  $\rho_3(1690)$  and  $\omega_3(1670)$ .NODE=M015M2;LINKAGE=P  
NODE=M015M2;LINKAGE=S  
NODE=M015M2;LINKAGE=L**(4 $\pi$ ) $\pm$  MODE**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015M3  
NODE=M015M3**1686± 5 OUR AVERAGE** Error includes scale factor of 1.1.

1694± 6		<sup>8</sup> EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
1665±15	177	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
1670±10		THOMPSON	74	HBC	+	13 $\pi^+ p$
1687±20		CASON	73	HBC	-	8,18.5 $\pi^- p$
1685±14		<sup>9</sup> CASON	73	HBC	-	8,18.5 $\pi^- p$
1680±40	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N4\pi$
1689±20	102	<sup>9</sup> BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\rho$
1705±21		CASO	70	HBC	-	11.2 $\pi^- p \rightarrow n\rho2\pi$

••• We do not use the following data for averages, fits, limits, etc. •••

1718±10 <sup>10</sup> EVANGELIS... 81 OMEG - 12  $\pi^- p \rightarrow p4\pi$ 1673± 9 <sup>11</sup> EVANGELIS... 81 OMEG - 12  $\pi^- p \rightarrow p4\pi$ 1733± 9 66 <sup>9</sup> KLIGER 74 HBC - 4.5  $\pi^- p \rightarrow p4\pi$ 1630±15 HOLMES 72 HBC + 10-12  $K^+ p$ 1720±15 BALTAY 68 HBC + 7, 8.5  $\pi^+ p$ 

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

- <sup>8</sup> From  $\rho^- \rho^0$  mode, not independent of the other two EVANGELISTA 81 entries.
- <sup>9</sup> From  $\rho^\pm \rho^0$  mode.
- <sup>10</sup> From  $a_2(1320)^- \pi^0$  mode, not independent of the other two EVANGELISTA 81 entries.
- <sup>11</sup> From  $a_2(1320)^0 \pi^-$  mode, not independent of the other two EVANGELISTA 81 entries.

NODE=M015M3;LINKAGE=A  
 NODE=M015M3;LINKAGE=F  
 NODE=M015M3;LINKAGE=B  
 NODE=M015M3;LINKAGE=C

**$\omega\pi$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

**1681 ± 7 OUR AVERAGE**

1670 ± 25	<sup>12</sup> ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$
1690 ± 15	EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow \omega \pi p$
1666 ± 14	GESSAROLI	77	HBC	11 $\pi^- p \rightarrow \omega \pi p$
1686 ± 9	THOMPSON	74	HBC +	13 $\pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1654 ± 24	BARNHAM	70	HBC +	10 $K^+ p \rightarrow \omega \pi X$

<sup>12</sup> Supersedes ALDE 92C.

NODE=M015M5  
 NODE=M015M5

NODE=M015M5;LINKAGE=A

**$\eta\pi^+ \pi^-$  MODE**

(For difficulties with MMS experiments, see the  $a_2(1320)$  mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

**1682 ± 12 OUR AVERAGE**

1685 ± 10 ± 20	AMELIN	00	VES	37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1680 ± 15	FUKUI	88	SPEC 0	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1700 ± 47	<sup>13</sup> ANDERSON	69	MMS -	16 $\pi^- p$ backward
1632 ± 15	<sup>13,14</sup> FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow pMM$
1700 ± 15	<sup>13,14</sup> FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow pMM$
1748 ± 15	<sup>13,14</sup> FOCACCI	66	MMS -	7-12 $\pi^- p \rightarrow pMM$

<sup>13</sup> Seen in 2.5-3 GeV/c  $\bar{p}p$ .  $2\pi^+ 2\pi^-$ , with 0, 1, 2  $\pi^+ \pi^-$  pairs in  $\rho$  band not seen by OREN 74 (2.3 GeV/c  $\bar{p}p$ ) with more statistics. (Jan. 1976)

<sup>14</sup> Not seen by BOWEN 72.

NODE=M015M6  
 NODE=M015M6  
 NODE=M015M6

OCCUR=2  
 OCCUR=3

NODE=M015M6;LINKAGE=R

NODE=M015M6;LINKAGE=N

**$\rho_3(1690)$  WIDTH**


**2 $\pi$ ,  $K\bar{K}$ , AND  $K\bar{K}\pi$  MODES**

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**161 ± 10 OUR AVERAGE** Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.

NODE=M015W  
 NODE=M015W

NODE=M015210



**2π MODE**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W1  
NODE=M015W1

**186±14 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

220±29		DENNEY	83	LASS		10 π <sup>+</sup> N
246±37		EVANGELIS...	81	OMEG	-	12 π <sup>-</sup> p → 2π p
116±30	476	BALTAY	78B	HBC	0	15 π <sup>+</sup> p → π <sup>+</sup> π <sup>-</sup> n
162±50	175	<sup>15</sup> ANTIPOV	77	CIBS	0	25 π <sup>-</sup> p → p3π
167±40	600	ENGLER	74	DBC	0	6 π <sup>+</sup> n → π <sup>+</sup> π <sup>-</sup> p
200±18		<sup>16</sup> GRAYER	74	ASPK	0	17 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> n
156±36		MATTHEWS	71C	DBC	0	7 π <sup>+</sup> N
171±65		ARMENISE	70	DBC	0	9 π <sup>+</sup> d

••• We do not use the following data for averages, fits, limits, etc. •••

322±35		<sup>17</sup> CORDEN	79	OMEG		12-15 π <sup>-</sup> p → n2π
240±30		<sup>16,18</sup> ESTABROOKS	75	RVUE		17 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> n
180±30	122	BARTSCH	70B	HBC	+	8 π <sup>+</sup> p → N2π
267 <sup>+72</sup> <sub>-46</sub>		STUNTEBECK	70	HDBC	0	8 π <sup>-</sup> p, 5.4 π <sup>+</sup> d
188±49		ARMENISE	68	DBC	0	5.1 π <sup>+</sup> d
180±40		GOLDBERG	65	HBC	0	6 π <sup>+</sup> d, 8 π <sup>-</sup> p

<sup>15</sup> Width errors enlarged by us to 4Γ/√N; see the note with the K\*(892) mass.


<sup>16</sup> Uses same data as HYAMS 75 and BECKER 79.

<sup>17</sup> From a phase shift solution containing a f<sub>2</sub>'(1525) width two times larger than the K<sup>-</sup>K<sup>+</sup> result.

<sup>18</sup> From phase-shift analysis. Error takes account of spread of different phase-shift solutions.

NODE=M015W1;LINKAGE=T  
NODE=M015W1;LINKAGE=G  
NODE=M015W1;LINKAGE=M  
NODE=M015W1;LINKAGE=L

WEIGHTED AVERAGE  
186±14 (Error scaled by 1.3)



**K<sup>-</sup>K<sup>+</sup> AND K<sup>-</sup>K<sup>+</sup>π MODES**

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W2  
NODE=M015W2

**204±18 OUR AVERAGE**

199±40	6000	<sup>19</sup> MARTIN	78D	SPEC		10 π p → K <sub>S</sub> <sup>0</sup> K <sup>-</sup> p
205±20		BLUM	75	ASPK	0	18.4 π <sup>-</sup> p → nK <sup>+</sup> K <sup>-</sup>
219±4		ALPER	80	CNTR	0	62 π <sup>-</sup> p → K <sup>+</sup> K <sup>-</sup> n
186±11		<sup>20</sup> COSTA	80	OMEG		10 π <sup>-</sup> p → K <sup>+</sup> K <sup>-</sup> n
112±60		ADERHOLZ	69	HBC	+	8 π <sup>+</sup> p → K <sup>-</sup> K <sup>+</sup> π

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>19</sup> From a fit to J<sup>P</sup> = 3<sup>-</sup> partial wave.

<sup>20</sup> They cannot distinguish between ρ<sub>3</sub>(1690) and ω<sub>3</sub>(1670).

NODE=M015W2;LINKAGE=P  
NODE=M015W2;LINKAGE=L

**(4 $\pi$ ) $^\pm$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015W3  
 NODE=M015W3

**129 $\pm$ 10 OUR AVERAGE**

123 $\pm$ 13		21 EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
105 $\pm$ 30	177	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
169 $^{+70}_{-48}$		CASON	73	HBC	-	8,18.5 $\pi^- p$
135 $\pm$ 30	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N4\pi$
160 $\pm$ 30	102	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\rho$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
230 $\pm$ 28		22 EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
184 $\pm$ 33		23 EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow p4\pi$
150	66	24 KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$
106 $\pm$ 25		THOMPSON	74	HBC	+	13 $\pi^+ p$
125 $^{+83}_{-35}$		24 CASON	73	HBC	-	8,18.5 $\pi^- p$
130 $\pm$ 30		HOLMES	72	HBC	+	10-12 $K^+ p$
180 $\pm$ 30	90	24 BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow Na_2\pi$
100 $\pm$ 35		BALTAY	68	HBC	+	7, 8.5 $\pi^+ p$

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

21 From  $\rho^- \rho^0$  mode, not independent of the other two EVANGELISTA 81 entries.22 From  $a_2(1320)^- \pi^0$  mode, not independent of the other two EVANGELISTA 81 entries.23 From  $a_2(1320)^0 \pi^-$  mode, not independent of the other two EVANGELISTA 81 entries.24 From  $\rho^\pm \rho^0$  mode.

NODE=M015W3;LINKAGE=A

NODE=M015W3;LINKAGE=B

NODE=M015W3;LINKAGE=C

NODE=M015W3;LINKAGE=F

 **$\omega\pi$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015W5  
 NODE=M015W5

**190 $\pm$ 40 OUR AVERAGE**

230 $\pm$ 65		25 ALDE	95	GAM2		38 $\pi^- p \rightarrow \omega\pi^0 n$
190 $\pm$ 65		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow \omega\pi p$
160 $\pm$ 56		GESSAROLI	77	HBC		11 $\pi^- p \rightarrow \omega\pi p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
89 $\pm$ 25		THOMPSON	74	HBC	+	13 $\pi^+ p$
130 $^{+73}_{-43}$		BARNHAM	70	HBC	+	10 $K^+ p \rightarrow \omega\pi X$

25 Supersedes ALDE 92C.

NODE=M015W5;LINKAGE=A

 **$\eta\pi^+\pi^-$  MODE**(For difficulties with MMS experiments, see the  $a_2(1320)$  mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015W6

NODE=M015W6

NODE=M015W6

**126 $\pm$ 40 OUR AVERAGE** Error includes scale factor of 1.8.

220 $\pm$ 30 $\pm$ 50		AMELIN	00	VES		37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
106 $\pm$ 27		FUKUI	88	SPEC	0	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
195		26 ANDERSON	69	MMS	-	16 $\pi^- p$ backward
< 21		26,27 FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow pMM$
< 30		26,27 FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow pMM$
< 38		26,27 FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow pMM$

OCCUR=2

OCCUR=3

26 Seen in 2.5-3 GeV/c  $\bar{p}p$ .  $2\pi^+2\pi^-$ , with 0, 1, 2  $\pi^+\pi^-$  pairs in  $\rho^0$  band not seen by OREN 74 (2.3 GeV/c  $\bar{p}p$ ) with more statistics. (Jan. 1979)

NODE=M015W6;LINKAGE=R

27 Not seen by BOWEN 72.

NODE=M015W6;LINKAGE=N

$\rho_3(1690)$  DECAY MODES

NODE=M015215;NODE=M015

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $4\pi$	(71.1 $\pm$ 1.9 ) %	
$\Gamma_2$ $\pi^\pm \pi^+ \pi^- \pi^0$	(67 $\pm$ 22 ) %	
$\Gamma_3$ $\omega \pi$	(16 $\pm$ 6 ) %	
$\Gamma_4$ $\pi \pi$	(23.6 $\pm$ 1.3 ) %	
$\Gamma_5$ $K \bar{K} \pi$	( 3.8 $\pm$ 1.2 ) %	
$\Gamma_6$ $K \bar{K}$	( 1.58 $\pm$ 0.26 ) %	1.2
$\Gamma_7$ $\eta \pi^+ \pi^-$	seen	
$\Gamma_8$ $\rho(770) \eta$	seen	
$\Gamma_9$ $\pi \pi \rho$	seen	
$\Gamma_{10}$ $a_2(1320) \pi$	seen	
$\Gamma_{11}$ $\rho \rho$	seen	
$\Gamma_{12}$ $\phi \pi$		
$\Gamma_{13}$ $\eta \pi$		
$\Gamma_{14}$ $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

DESIG=2  
DESIG=11  
DESIG=7  
DESIG=1  
DESIG=3  
DESIG=4  
DESIG=13  
DESIG=14;OUR EST;→ UNCHECKED ←  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←  
DESIG=8;OUR EST;→ UNCHECKED ←  
DESIG=9  
DESIG=10  
DESIG=12

## CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 14.7$  for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_4$	-77		
$x_5$	-74	17	
$x_6$	-15	2	0
	$x_1$	$x_4$	$x_5$

 $\rho_3(1690)$  BRANCHING RATIOS

NODE=M015220

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.236<math>\pm</math>0.013 OUR FIT</b>					
<b>0.243<math>\pm</math>0.013 OUR AVERAGE</b>					
0.259 $^{+0.018}_{-0.019}$	BECKER	79	ASPK	0	17 $\pi^- p$ polarized
0.23 $\pm$ 0.02	CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow n 2\pi$
0.22 $\pm$ 0.04	<sup>28</sup> MATTHEWS	71c	HDBC	0	7 $\pi^+ n \rightarrow \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.245 $\pm$ 0.006	<sup>29</sup> ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
<sup>28</sup> One-pion-exchange model used in this estimation.					
<sup>29</sup> From phase-shift analysis of HYAMS 75 data.					

NODE=M015R1  
NODE=M015R1NODE=M015R1;LINKAGE=P  
NODE=M015R1;LINKAGE=G

$\Gamma(\pi\pi)/\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)$					$\Gamma_4/\Gamma_2$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.35<math>\pm</math>0.11</b>	CASON	73	HBC	-	8,18.5 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.2	HOLMES	72	HBC	+	10-12 $K^+ p$
<0.12	BALLAM	71B	HBC	-	16 $\pi^- p$

NODE=M015R2  
NODE=M015R2

$\Gamma(\pi\pi)/\Gamma(4\pi)$					$\Gamma_4/\Gamma_1$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.332<math>\pm</math>0.026 OUR FIT</b>	Error includes scale factor of 1.1.				
<b>0.30 <math>\pm</math> 0.10</b>	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow p 4\pi$

NODE=M015R3  
NODE=M015R3

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

$\Gamma_6/\Gamma_4$

VALUE DOCUMENT ID TECN CHG COMMENT


NODE=M015R4  
NODE=M015R4

**0.067±0.011 OUR FIT** Error includes scale factor of 1.2.

**0.118<sup>+0.040</sup><sub>-0.032</sub> OUR AVERAGE** Error includes scale factor of 1.7. See the ideogram below.

0.191 <sup>+0.040</sup> <sub>-0.037</sub>	GORLICH	80	ASPK	0	17,18 $\pi^- p$ polarized
0.08 ±0.03	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.08 <sup>+0.08</sup> <sub>-0.03</sub>	CRENNELL	68B	HBC		6.0 $\pi^- p$

WEIGHTED AVERAGE  
0.118+0.040-0.032 (Error scaled by 1.7)



$\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$

$\Gamma_5/\Gamma_4$

VALUE DOCUMENT ID TECN CHG COMMENT

NODE=M015R5  
NODE=M015R5

**0.16±0.05 OUR FIT**

**0.16±0.05** <sup>30</sup> BARTSCH 70B HBC + 8  $\pi^+ p$

<sup>30</sup> Increased by us to correspond to  $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$ .

NODE=M015R5;LINKAGE=A

$[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$   $(\Gamma_9+\Gamma_{10}+\Gamma_{11})/\Gamma_2$

VALUE DOCUMENT ID TECN CHG COMMENT

NODE=M015R6  
NODE=M015R6

**0.94±0.09 OUR AVERAGE**

0.96±0.21	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.88±0.15	BALLAM	71B	HBC	-	16 $\pi^- p$
1 ±0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
consistent with 1	CASO	68	HBC	-	11 $\pi^- p$

$\Gamma(\rho\rho)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$

$\Gamma_{11}/\Gamma_2$

VALUE EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M015R7  
NODE=M015R7

••• We do not use the following data for averages, fits, limits, etc. •••

0.12±0.11	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.56	66 KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$
0.13±0.09	<sup>31</sup> THOMPSON	74	HBC	+	13 $\pi^+ p$
0.7 ±0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$

<sup>31</sup>  $\rho\rho$  and  $a_2(1320)\pi$  modes are indistinguishable.

NODE=M015R7;LINKAGE=T

$\Gamma(\rho\rho)/[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]$   $\Gamma_{11}/(\Gamma_9+\Gamma_{10}+\Gamma_{11})$

VALUE DOCUMENT ID TECN CHG COMMENT

NODE=M015R8  
NODE=M015R8

••• We do not use the following data for averages, fits, limits, etc. •••

0.48±0.16	CASO	68	HBC	-	11 $\pi^- p$
-----------	------	----	-----	---	--------------

$\Gamma(a_2(1320)\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$  $\Gamma_{10}/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.66±0.08	BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
0.36±0.14	<sup>32</sup> THOMPSON	74	HBC	+ 13 $\pi^+ p$
not seen	CASON	73	HBC	- 8,18.5 $\pi^- p$
0.6 ±0.15	BARTSCH	70B	HBC	+ 8 $\pi^+ p$
0.6	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R9  
NODE=M015R9<sup>32</sup>  $\rho\rho$  and  $a_2(1320)\pi$  modes are indistinguishable.

NODE=M015R9;LINKAGE=T

 $\Gamma(\omega\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$  $\Gamma_3/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.23±0.05 OUR AVERAGE</b>		Error includes scale factor of 1.2.			
0.33±0.07		THOMPSON	74	HBC	+ 13 $\pi^+ p$
0.12±0.07		BALLAM	71B	HBC	- 16 $\pi^- p$
0.25±0.10		BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$
0.25±0.10		JOHNSTON	68	HBC	- 7.0 $\pi^- p$
<0.11	95	BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
<0.09		KLIGER	74	HBC	- 4.5 $\pi^- p \rightarrow p4\pi$

NODE=M015R10  
NODE=M015R10 $\Gamma(\phi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$  $\Gamma_{12}/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.11	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R11  
NODE=M015R11 $\Gamma(\pi^\pm 2\pi^+ 2\pi^- \pi^0)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$  $\Gamma_{14}/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.15	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R12  
NODE=M015R12 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$  $\Gamma_{13}/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.02	THOMPSON	74	HBC	+ 13 $\pi^+ p$

NODE=M015R13  
NODE=M015R13 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.0158±0.0026 OUR FIT</b>	Error includes scale factor of 1.2.			
<b>0.0130±0.0024 OUR AVERAGE</b>				

0.013 ±0.003	COSTA	80	OMEG 0	10 $\pi^- p \rightarrow K^+ K^- n$
0.013 ±0.004	<sup>33</sup> MARTIN	78B	SPEC	- 10 $\pi p \rightarrow K_S^0 K^- p$

<sup>33</sup> From  $(\Gamma_4\Gamma_6)^{1/2} = 0.056 \pm 0.034$  assuming  $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$ .NODE=M015R14  
NODE=M015R14

NODE=M015R14;LINKAGE=B

 $\Gamma(\omega\pi)/[\Gamma(\omega\pi) + \Gamma(\rho\rho)]$  $\Gamma_3/(\Gamma_3+\Gamma_{11})$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.22±0.08	CASON	73	HBC	- 8,18.5 $\pi^- p$

NODE=M015R16  
NODE=M015R16 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	FUKUI	88	SPEC 8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M015R17  
NODE=M015R17 $\Gamma(a_2(1320)\pi)/\Gamma(\rho(770)\eta)$  $\Gamma_{10}/\Gamma_8$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>5.5±2.0</b>	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M015R18  
NODE=M015R18

**$\rho_3(1690)$  REFERENCES**

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)
MARTIN	78B	NP B140 158	A.D. Martin <i>et al.</i>	(DURH, GEVA)
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA)
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
KLIGER	74	SJNP 19 428	G.K. Kliger <i>et al.</i>	(ITEP)

Translated from YAF 19 839.

OREN	74	NP B71 189	Y. Oren <i>et al.</i>	(ANL, OXF)
THOMPSON	74	NP B69 220	G. Thompson <i>et al.</i>	(PURD)
CASON	73	PR D7 1971	N.M. Cason <i>et al.</i>	(NDAM)
BOWEN	72	PRL 29 890	D.R. Bowen <i>et al.</i>	(NEAS, STON)
HOLMES	72	PR D6 3336	R. Holmes <i>et al.</i>	(ROCH)
BALLAM	71B	PR D3 2606	J. Ballam <i>et al.</i>	(SLAC)
MATTHEWS	71C	NP B33 1	J.A.J. Matthews <i>et al.</i>	(TNT0, WISC) JP
ARMENISE	70	LCN 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)
BARNHAM	70	PRL 24 1083	K.W.J. Barnham <i>et al.</i>	(BIRM)
BARTSCH	70B	NP B22 109	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN)
CASO	70	LCN 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)
ANDERSON	69	PRL 22 1390	E.W. Anderson <i>et al.</i>	(BNL, CMU)
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+) I
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I
CASO	68	NC 54A 983	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)
CRENNELL	68B	PL 28B 136	D.J. Crennell <i>et al.</i>	(BNL)
JOHNSTON	68	PRL 20 1414	T.F. Johnston <i>et al.</i>	(TNT0, WISC) IJP
FOCACCI	66	PRL 17 890	M.N. Focacci <i>et al.</i>	(CERN)
GOLDBERG	65	PL 17 354	M. Goldberg <i>et al.</i>	(CERN, EPOL, ORSAY+)

NODE=M015

REFID=47432  
 REFID=44371  
 REFID=41859  
 REFID=40273  
 REFID=20754  
 REFID=20462  
 REFID=21665  
 REFID=20737  
 REFID=20738  
 REFID=21084  
 REFID=20374  
 REFID=21265  
 REFID=21273  
 REFID=21272  
 REFID=20728  
 REFID=20230  
 REFID=21651  
 REFID=20642  
 REFID=20355  
 REFID=20110  
 REFID=20113  
 REFID=21648  
 REFID=20221  
 REFID=21650  
 REFID=20606  
 REFID=21711  
 REFID=21639  
 REFID=21630  
 REFID=21633  
 REFID=20693  
 REFID=21624  
 REFID=21625  
 REFID=20590  
 REFID=20696  
 REFID=20687  
 REFID=20795  
 REFID=20054  
 REFID=21531  
 REFID=20586  
 REFID=21616  
 REFID=21617  
 REFID=20402  
 REFID=21601

**$\rho(1700)$**

$$I^G(J^{PC}) = 1^+(1^{--})$$

NODE=M065

**$\rho(1700)$  MASS**

NODE=M065205

**$\eta\rho^0$  AND  $\pi^+\pi^-$  MODES**

VALUE (MeV)	DOCUMENT ID
<b>1720±20 OUR ESTIMATE</b>	

NODE=M065M0  
 NODE=M065M0  
 → UNCHECKED ←

**$\eta\rho^0$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M065M6  
 NODE=M065M6

The data in this block is included in the average printed for a previous datablock.

••• We do not use the following data for averages, fits, limits, etc. •••

1834±12	13.4k	<sup>1</sup> GRIBANOV	20	CMD3	1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1840±10	7.4k	<sup>2</sup> ACHASOV	18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1740±20		ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1701±15		<sup>3</sup> FUKUI	88	SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

<sup>1</sup> Mass and width of the  $\rho(770)$  fixed at 775 and 149 MeV, respectively; solution 2 of model 2,  $\eta \rightarrow \gamma\gamma$  decays used.

NODE=M065M6;LINKAGE=B

<sup>2</sup> From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering  $\rho(1450)$ ,  $\rho(1700)$  and  $\rho(2150)$  with the parameters of the  $\rho(1450)$  and  $\rho(1700)$  floating and the mass and width of the  $\rho(2150)$  fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are  $\pi$ , 0 and  $\pi$ , respectively.

NODE=M065M6;LINKAGE=A

<sup>3</sup> Assuming  $\rho^+ f_0(1370)$  decay mode interferes with  $a_1(1260)^+\pi$  background. From a two Breit-Wigner fit.

NODE=M065M;LINKAGE=B



**$\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1770.54 ± 5.49		1 BARTOS	17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1718.50 ± 65.44		2 BARTOS	17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1766.80 ± 52.36		3 BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
1644 ± 36	20k	4 LEES	17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
1780 ± 20	$^{+15}_{-20}$ 63.5k	5 ABRAMOWICZ12	ZEUS		$ep \rightarrow e\pi^+\pi^-p$
1861 ± 17		6 LEES	12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
1728 ± 17	± 89 5.4M	7,8 FUJIKAWA	08	BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
1780	$^{+37}_{-29}$	9 ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
1719 ± 15		9 BERTIN	97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1730 ± 30		CLEGG	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1768 ± 21		BISELLO	89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
1745.7 ± 91.9		DUBNICKA	89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1546 ± 26		GESHKEN...	89	RVUE	
1650		10 ERKAL	85	RVUE	20-70 $\gamma p \rightarrow \gamma\pi$
1550 ± 70		ABE	84B	HYBR	20 $\gamma p \rightarrow \pi^+\pi^-p$
1590 ± 20		11 ASTON	80	OMEG	20-70 $\gamma p \rightarrow p2\pi$
1600 ± 10		12 ATIYA	79B	SPEC	50 $\gamma C \rightarrow C2\pi$
1598	$^{+24}_{-22}$	BECKER	79	ASPK	17 $\pi^- p$ polarized
1659 ± 25		10 LANG	79	RVUE	
1575		10 MARTIN	78C	RVUE	17 $\pi^- p \rightarrow \pi^+\pi^-n$
1610 ± 30		10 FROGGATT	77	RVUE	17 $\pi^- p \rightarrow \pi^+\pi^-n$
1590 ± 20		13 HYAMS	73	ASPK	17 $\pi^- p \rightarrow \pi^+\pi^-n$

NODE=M065M1  
NODE=M065M1

OCCUR=2

1 Applies the Unitary &amp; Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

NODE=M065M1;LINKAGE=C

2 Applies the Unitary &amp; Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

NODE=M065M1;LINKAGE=D

3 Applies the Unitary &amp; Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.

NODE=M065M1;LINKAGE=E

4 From a Dalitz plot analysis in an isobar model with  $\rho(1450)$  and  $\rho(1700)$  masses and widths floating.

NODE=M065M1;LINKAGE=A

5 Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho-\omega$  interference.

NODE=M065M1;LINKAGE=AB

6 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

NODE=M065M1;LINKAGE=LE

7  $|F_\pi(0)|^2$  fixed to 1.

NODE=M065M1;LINKAGE=FU

8 From the GOUNARIS 68 parametrization of the pion form factor.

NODE=M065M1;LINKAGE=GO

9 T-matrix pole.

NODE=M065M;LINKAGE=QQ

10 From phase shift analysis of HYAMS 73 data.

NODE=M065M;LINKAGE=P

11 Simple relativistic Breit-Wigner fit with constant width.

NODE=M065M;LINKAGE=M

12 An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

NODE=M065M;LINKAGE=R

13 Included in BECKER 79 analysis.

NODE=M065M;LINKAGE=H

 **$\pi\omega$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1708 ± 41	7815	1 ACHASOV	13	SND	1.05-2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1550 to 1620		2 ACHASOV	00i	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1580 to 1710		3 ACHASOV	00i	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1710 ± 90		ACHASOV	97	RVUE	$e^+e^- \rightarrow \omega\pi^0$

NODE=M065M8  
NODE=M065M8

OCCUR=2

1 From a phenomenological model based on vector meson dominance with the interfering  $\rho(1450)$  and  $\rho(1700)$  and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M065M8;LINKAGE=AC

2 Taking into account both  $\rho(1450)$  and  $\rho(1700)$  contributions. Using the data of ACHASOV 00i on  $e^+e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^-\nu_\tau$ .  $\rho(1450)$  mass and width fixed at 1400 MeV and 500 MeV respectively.

NODE=M065M;LINKAGE=11

3 Taking into account the  $\rho(1700)$  contribution only. Using the data of ACHASOV 00i on  $e^+e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^-\nu_\tau$ .

NODE=M065M;LINKAGE=12

**$K\bar{K}$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1688.7 \pm 3.1^{+141.1}_{-1.3}$		<sup>1</sup> ALBRECHT	20	RVUE	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
$1541 \pm 12 \pm 33$	190k	<sup>2</sup> AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
$1740.8 \pm 22.2$	27k	<sup>3</sup> ABELE	99D	CBAR $\pm$	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$
$1582 \pm 36$	1600	CLELAND	82B	SPEC $\pm$	$50 \pi p \rightarrow K_S^0 K^\pm p$

<sup>1</sup> T-matrix pole, 2 poles, 3 channels, including  $\pi\pi$  scattering data from HYAMS 75.

<sup>2</sup> Using the GOUNARIS 68 parameterization with a fixed width. Value is average using different  $K\pi$  S-wave parameterizations in fit.

<sup>3</sup> K-matrix pole. Isospin not determined, could be  $\omega(1650)$  or  $\phi(1680)$ .

NODE=M065M2  
NODE=M065M2

NODE=M065M2;LINKAGE=D  
NODE=M065M2;LINKAGE=A

NODE=M065M2;LINKAGE=AN

 **$2(\pi^+\pi^-)$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1851^{+27}_{-24}$		ACHASOV	97	RVUE	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
$1570 \pm 20$		<sup>1</sup> CORDIER	82	DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
$1520 \pm 30$		<sup>2</sup> ASTON	81E	OMEG	$20-70 \gamma p \rightarrow p4\pi$
$1654 \pm 25$		<sup>3</sup> DIBIANCA	81	DBC	$\pi^+ d \rightarrow pp2(\pi^+\pi^-)$
$1666 \pm 39$		<sup>1</sup> BACCI	80	FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1780	34	KILLIAN	80	SPEC	$11 e^- p \rightarrow 2(\pi^+\pi^-)$
1500		<sup>4</sup> ATIYA	79B	SPEC	$50 \gamma C \rightarrow C4\pi^\pm$
$1570 \pm 60$	65	<sup>5</sup> ALEXANDER	75	HBC	$7.5 \gamma p \rightarrow p4\pi$
$1550 \pm 60$		<sup>2</sup> CONVERSI	74	OSPK	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
$1550 \pm 50$	160	SCHACHT	74	STRC	$5.5-9 \gamma p \rightarrow p4\pi$
$1450 \pm 100$	340	SCHACHT	74	STRC	$9-18 \gamma p \rightarrow p4\pi$
$1430 \pm 50$	400	BINGHAM	72B	HBC	$9.3 \gamma p \rightarrow p4\pi$

<sup>1</sup> Simple relativistic Breit-Wigner fit with model dependent width.

<sup>2</sup> Simple relativistic Breit-Wigner fit with constant width.

<sup>3</sup> One peak fit result.

<sup>4</sup> Parameters roughly estimated, not from a fit.

<sup>5</sup> Skew mass distribution compensated by Ross-Stodolsky factor.

NODE=M065M4  
NODE=M065M4

OCCUR=2

NODE=M065M;LINKAGE=A  
NODE=M065M4;LINKAGE=M  
NODE=M065M;LINKAGE=O  
NODE=M065M;LINKAGE=C  
NODE=M065M;LINKAGE=D

 **$\pi^+\pi^-\pi^0\pi^0$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1660 \pm 30$	ATKINSON	85B	OMEG 20-70 $\gamma p$
---------------	----------	-----	-----------------------

NODE=M065M5  
NODE=M065M5

 **$3(\pi^+\pi^-)$  AND  $2(\pi^+\pi^-\pi^0)$  MODES**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1730 \pm 34$	<sup>1</sup> FRABETTI	04	E687 $\gamma p \rightarrow 3\pi^+ 3\pi^- p$
$1783 \pm 15$	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

<sup>1</sup> From a fit with two resonances with the JACOB 72 continuum.

NODE=M065M7  
NODE=M065M7

NODE=M065M;LINKAGE=PI

 **$m_{\rho(1700)^0} - m_{\rho(1700)^\pm}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-48.30 \pm 83.81$	<sup>1</sup> BARTOS	17A	RVUE $e^+e^- \rightarrow \pi^+\pi^-$ , $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
--------------------	---------------------	-----	---

<sup>1</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M065DM

NODE=M065DM

NODE=M065DM;LINKAGE=A

 **$\rho(1700)$  WIDTH** **$\eta\rho^0$  AND  $\pi^+\pi^-$  MODES**

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**250 ± 100 OUR ESTIMATE**

NODE=M065210

NODE=M065W0  
NODE=M065W0

→ UNCHECKED ←

**$\eta\rho^0$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M065W6  
NODE=M065W6

• • • We do not use the following data for averages, fits, limits, etc. • • •

47 ± 19	13.4k	<sup>1</sup> GRIBANOV	20	CMD3	1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
132 ± 40	7.4k	<sup>2</sup> ACHASOV	18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
150 ± 30		ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
282 ± 44		<sup>3</sup> FUKUI	88	SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

<sup>1</sup> Mass and width of the  $\rho(770)$  fixed at 775 and 149 MeV, respectively; solution 2 of model 2,  $\eta \rightarrow \gamma\gamma$  decays used.

NODE=M065W6;LINKAGE=B

<sup>2</sup> From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering  $\rho(1450)$ ,  $\rho(1700)$  and  $\rho(2150)$  with the parameters of the  $\rho(1450)$  and  $\rho(1700)$  floating and the mass and width of the  $\rho(2150)$  fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are  $\pi$ , 0 and  $\pi$ , respectively.

NODE=M065W6;LINKAGE=A

<sup>3</sup> Assuming  $\rho^+ f_0(1370)$  decay mode interferes with  $a_1(1260)^+ \pi$  background. From a two Breit-Wigner fit.

NODE=M065W;LINKAGE=B

 **$\pi\pi$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M065W1  
NODE=M065W1

• • • We do not use the following data for averages, fits, limits, etc. • • •

268.98 ± 11.40		<sup>1</sup> BARTOS	17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
489.58 ± 16.95		<sup>2</sup> BARTOS	17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
414.71 ± 119.48		<sup>3</sup> BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
109 ± 19	20k	<sup>4</sup> LEES	17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
310 ± 30	$\begin{smallmatrix} +25 \\ -35 \end{smallmatrix}$ 63.5k	<sup>5</sup> ABRAMOWICZ12	ZEUS		$ep \rightarrow e\pi^+\pi^-p$
316 ± 26		<sup>6</sup> LEES	12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
164 ± 21	$\begin{smallmatrix} +89 \\ -26 \end{smallmatrix}$ 5.4M	<sup>7,8</sup> FUJIKAWA	08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
275 ± 45		<sup>9</sup> ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
310 ± 40		<sup>9</sup> BERTIN	97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
400 ± 100		CLEGG	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
224 ± 22		BISELLO	89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
242.5 ± 163.0		DUBNICKA	89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
620 ± 60		GESHKEN...	89	RVUE	
<315		<sup>10</sup> ERKAL	85	RVUE	20–70 $\gamma p \rightarrow \gamma\pi$
280 $\begin{smallmatrix} +30 \\ -80 \end{smallmatrix}$		ABE	84B	HYBR	20 $\gamma p \rightarrow \pi^+\pi^-p$
230 ± 80		<sup>11</sup> ASTON	80	OMEG	20–70 $\gamma p \rightarrow p2\pi$
283 ± 14		<sup>12</sup> ATIYA	79B	SPEC	50 $\gamma C \rightarrow C2\pi$
175 $\begin{smallmatrix} +98 \\ -53 \end{smallmatrix}$		BECKER	79	ASPK	17 $\pi^-p$ polarized
232 ± 34		<sup>10</sup> LANG	79	RVUE	
340		<sup>10</sup> MARTIN	78C	RVUE	17 $\pi^-p \rightarrow \pi^+\pi^-n$
300 ± 100		<sup>10</sup> FROGGATT	77	RVUE	17 $\pi^-p \rightarrow \pi^+\pi^-n$
180 ± 50		<sup>13</sup> HYAMS	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$

OCCUR=2

<sup>1</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

NODE=M065W1;LINKAGE=C

<sup>2</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

NODE=M065W1;LINKAGE=D

<sup>3</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.

NODE=M065W1;LINKAGE=E

<sup>4</sup> From a Dalitz plot analysis in an isobar model with  $\rho(1450)$  and  $\rho(1700)$  masses and widths floating.

NODE=M065W1;LINKAGE=A

<sup>5</sup> Using the KUHN 90 parametrization of the pion form factor, neglecting  $\rho-\omega$  interference.

NODE=M065W1;LINKAGE=AB

<sup>6</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

NODE=M065W1;LINKAGE=LE

<sup>7</sup>  $|F_\pi(0)|^2$  fixed to 1.

NODE=M065W1;LINKAGE=FU

<sup>8</sup> From the GOUNARIS 68 parametrization of the pion form factor.

NODE=M065W1;LINKAGE=GO

<sup>9</sup> T-matrix pole.

NODE=M065W;LINKAGE=QQ

<sup>10</sup> From phase shift analysis of HYAMS 73 data.

NODE=M065W;LINKAGE=P

<sup>11</sup> Simple relativistic Breit-Wigner fit with constant width.

NODE=M065W;LINKAGE=M

<sup>12</sup> An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

NODE=M065W;LINKAGE=R

<sup>13</sup> Included in BECKER 79 analysis.

NODE=M065W;LINKAGE=H

**$K\bar{K}$  MODE**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$150.9 \pm 2.5^{+60}_{-10.6}$		<sup>1</sup> ALBRECHT	20	RVUE	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
$187.2 \pm 26.7$	27k	<sup>2</sup> ABELE	99D	CBAR $\pm$	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$
$265 \pm 120$	1600	CLELAND	82B	SPEC $\pm$	$50 \pi p \rightarrow K_S^0 K^\pm p$

<sup>1</sup> T-matrix pole, 2 poles, 3 channels, including  $\pi\pi$  scattering data from HYAMS 75.

<sup>2</sup> K-matrix pole. Isospin not determined, could be  $\omega(1650)$  or  $\phi(1680)$ .

NODE=M065W2  
NODE=M065W2

NODE=M065W2;LINKAGE=A  
NODE=M065W2;LINKAGE=AN

 **$2(\pi^+\pi^-)$  MODE**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$510 \pm 40$		<sup>1</sup> CORDIER	82	DM1	$e^+ e^- \rightarrow 2(\pi^+\pi^-)$
$400 \pm 50$		<sup>2</sup> ASTON	81E	OMEG	$20-70 \gamma p \rightarrow p4\pi$
$400 \pm 146$		<sup>3</sup> DIBIANCA	81	DBC	$\pi^+ d \rightarrow pp2(\pi^+\pi^-)$
$700 \pm 160$		<sup>1</sup> BACCI	80	FRAG	$e^+ e^- \rightarrow 2(\pi^+\pi^-)$
100	34	KILLIAN	80	SPEC	$11 e^- p \rightarrow 2(\pi^+\pi^-)$
600		<sup>4</sup> ATIYA	79B	SPEC	$50 \gamma C \rightarrow C4\pi^\pm$
$340 \pm 160$	65	<sup>5</sup> ALEXANDER	75	HBC	$7.5 \gamma p \rightarrow p4\pi$
$360 \pm 100$		<sup>2</sup> CONVERSI	74	OSPK	$e^+ e^- \rightarrow 2(\pi^+\pi^-)$
$400 \pm 120$	160	<sup>6</sup> SCHACHT	74	STRC	$5.5-9 \gamma p \rightarrow p4\pi$
$850 \pm 200$	340	<sup>6</sup> SCHACHT	74	STRC	$9-18 \gamma p \rightarrow p4\pi$
$650 \pm 100$	400	BINGHAM	72B	HBC	$9.3 \gamma p \rightarrow p4\pi$

<sup>1</sup> Simple relativistic Breit-Wigner fit with model-dependent width.

<sup>2</sup> Simple relativistic Breit-Wigner fit with constant width.

<sup>3</sup> One peak fit result.

<sup>4</sup> Parameters roughly estimated, not from a fit.

<sup>5</sup> Skew mass distribution compensated by Ross-Stodolsky factor.

<sup>6</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

NODE=M065W4  
NODE=M065W4

OCCUR=2

NODE=M065W;LINKAGE=A  
NODE=M065W4;LINKAGE=M  
NODE=M065W;LINKAGE=O  
NODE=M065W;LINKAGE=C  
NODE=M065W;LINKAGE=D  
NODE=M065W;LINKAGE=E

 **$\pi^+\pi^-\pi^0\pi^0$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$300 \pm 50$	ATKINSON	85B	OMEG 20-70 $\gamma p$
--------------	----------	-----	-----------------------

NODE=M065W5  
NODE=M065W5

 **$\omega\pi^0$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

350 to 580	<sup>1</sup> ACHASOV	00I	SND $e^+ e^- \rightarrow \pi^0\pi^0\gamma$
490 to 1040	<sup>2</sup> ACHASOV	00I	SND $e^+ e^- \rightarrow \pi^0\pi^0\gamma$

<sup>1</sup> Taking into account both  $\rho(1450)$  and  $\rho(1700)$  contributions. Using the data of ACHASOV 00I on  $e^+ e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^-\nu_\tau$ .  $\rho(1450)$  mass and width fixed at 1400 MeV and 500 MeV respectively.

<sup>2</sup> Taking into account the  $\rho(1700)$  contribution only. Using the data of ACHASOV 00I on  $e^+ e^- \rightarrow \omega\pi^0$  and of EDWARDS 00A on  $\tau^- \rightarrow \omega\pi^-\nu_\tau$ .

NODE=M065W9  
NODE=M065W9

OCCUR=2

NODE=M065W;LINKAGE=I1

NODE=M065W;LINKAGE=I2

 **$3(\pi^+\pi^-)$  AND  $2(\pi^+\pi^-\pi^0)$  MODES**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$315 \pm 100$	<sup>1</sup> FRABETTI	04	E687 $\gamma p \rightarrow 3\pi^+ 3\pi^- p$
$285 \pm 20$	CLEGG	90	RVUE $e^+ e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

<sup>1</sup> From a fit with two resonances with the JACOB 72 continuum.

NODE=M065W7  
NODE=M065W7

NODE=M065W;LINKAGE=PI

$$\Gamma_{\rho(1700)^0} - \Gamma_{\rho(1700)^\pm}$$

NODE=M065DW

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$74.87 \pm 120.67$	<sup>1</sup> BARTOS	17A	RVUE $e^+ e^- \rightarrow \pi^+\pi^-, \tau^- \rightarrow \pi^-\pi^0\nu_\tau$
--------------------	---------------------	-----	--

<sup>1</sup> Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M065DW

NODE=M065DW;LINKAGE=A

**$\rho(1700)$  DECAY MODES**

NODE=M065215;NODE=M065

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $4\pi$	
$\Gamma_2$ $2(\pi^+\pi^-)$	seen
$\Gamma_3$ $\rho\pi\pi$	seen
$\Gamma_4$ $\rho^0\pi^+\pi^-$	seen
$\Gamma_5$ $\rho^0\pi^0\pi^0$	
$\Gamma_6$ $\rho^\pm\pi^\mp\pi^0$	seen
$\Gamma_7$ $a_1(1260)\pi$	seen
$\Gamma_8$ $h_1(1170)\pi$	seen
$\Gamma_9$ $\pi(1300)\pi$	seen
$\Gamma_{10}$ $\rho\rho$	seen
$\Gamma_{11}$ $\pi^+\pi^-$	seen
$\Gamma_{12}$ $\pi\pi$	seen
$\Gamma_{13}$ $K\bar{K}^*(892)+c.c.$	seen
$\Gamma_{14}$ $\eta\rho$	seen
$\Gamma_{15}$ $a_2(1320)\pi$	not seen
$\Gamma_{16}$ $K\bar{K}$	seen
$\Gamma_{17}$ $e^+e^-$	seen
$\Gamma_{18}$ $\pi^0\omega$	seen
$\Gamma_{19}$ $\pi^0\gamma$	not seen
$\Gamma_{20}$ $f_0(1500)\gamma$	not seen

DESIG=20  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=12;OUR EST;→ UNCHECKED ←  
DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=7  
DESIG=9;OUR EST;→ UNCHECKED ←  
DESIG=15;OUR EST;→ UNCHECKED ←  
DESIG=16;OUR EST;→ UNCHECKED ←  
DESIG=17;OUR EST;→ UNCHECKED ←  
DESIG=18;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=13;OUR EST;→ UNCHECKED ←  
DESIG=10;OUR EST;→ UNCHECKED ←  
DESIG=11;OUR EST;→ UNCHECKED ←  
DESIG=14;OUR EST;→ UNCHECKED ←  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=8;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←  
DESIG=194  
DESIG=195

 **$\rho(1700) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

NODE=M065225

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the cross-section into channel<sub>i</sub> in  $e^+e^-$  annihilation.

NODE=M065225

 **$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_2\Gamma_{17}/\Gamma$** NODE=M065G2  
NODE=M065G2

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
2.6 ± 0.2	DEL COURT	81B	DM1 $e^+e^- \rightarrow 2(\pi^+\pi^-)$
2.83 ± 0.42	BACCI	80	FRAG $e^+e^- \rightarrow 2(\pi^+\pi^-)$

 **$\Gamma(\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{11}\Gamma_{17}/\Gamma$** NODE=M065G4  
NODE=M065G4

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
0.13	<sup>1</sup> DIEKMAN	88	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
0.029 <sup>+0.016</sup> <sub>-0.012</sub>	KURDADZE	83	OLYA 0.64–1.4 $e^+e^- \rightarrow \pi^+\pi^-$

<sup>1</sup> Using total width = 220 MeV.

NODE=M065G4;LINKAGE=B

 **$\Gamma(K\bar{K}^*(892)+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{13}\Gamma_{17}/\Gamma$** NODE=M065G10  
NODE=M065G10

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
0.305 ± 0.071	<sup>1</sup> BIZOT	80	DM1 $e^+e^-$

<sup>1</sup> Model dependent.

NODE=M065G;LINKAGE=M

 **$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{14}\Gamma_{17}/\Gamma$** NODE=M065G11  
NODE=M065G11

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
1.35 ± 0.53 ± 0.08 13.4k		<sup>1</sup> GRIBANOV	20	CMD3 1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
84 ± 26 ± 4		<sup>2</sup> LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-$
7 ± 3		ANTONELLI	88	DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$

<sup>1</sup> Mass and width of the  $\rho(770)$  fixed at 775 and 149 MeV, respectively; solution 2 of model 2,  $\eta \rightarrow \gamma\gamma$  decays used.

NODE=M065G11;LINKAGE=B

<sup>2</sup> Includes non-resonant contribution. The selected fit model includes three  $\rho$  excited states. Model uncertainty is 80%.

NODE=M065G11;LINKAGE=A

$$\Gamma(K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{16}\Gamma_{17}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.035±0.029	<sup>1</sup> BIZOT	80	DM1 e <sup>+</sup> e <sup>-</sup>
-------------	--------------------	----	-----------------------------------

<sup>1</sup> Model dependent.

NODE=M065G5  
NODE=M065G5

NODE=M065G5;LINKAGE=M

$$\Gamma(\rho\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_3\Gamma_{17}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.510±0.090	<sup>1</sup> BIZOT	80	DM1 e <sup>+</sup> e <sup>-</sup>
-------------	--------------------	----	-----------------------------------

<sup>1</sup> Model dependent.

NODE=M065G12  
NODE=M065G12

NODE=M065G12;LINKAGE=M

$$\rho(1700) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$$

NODE=M065240

$$\Gamma(\pi^0\omega)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{18}/\Gamma \times \Gamma_{17}/\Gamma$$

VALUE (units 10 <sup>-6</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.09±0.05	10.2k	<sup>1</sup> ACHASOV	16D	SND 1.05–2.00 e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ
1.7 ±0.4	7815	<sup>2</sup> ACHASOV	13	SND 1.05–2.00 e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> γ

<sup>1</sup> From a phenomenological model based on vector meson dominance with interfering ρ(700), ρ(1450), and ρ(1700). The ρ(1700) mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainty not estimated. Supersedes ACHASOV 13.

<sup>2</sup> From a phenomenological model based on vector meson dominance with the interfering ρ(1450) and ρ(1700) and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M065R01  
NODE=M065R01

OCCUR=4

NODE=M065R01;LINKAGE=B

NODE=M065R01;LINKAGE=AC

$$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_{14}/\Gamma \times \Gamma_{17}/\Gamma$$

VALUE (units 10 <sup>-8</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.3 <sup>+3.8</sup> <sub>-3.1</sub>	7.4k	<sup>1</sup> ACHASOV	18	SND 1.22–2.00 e <sup>+</sup> e <sup>-</sup> → ηπ <sup>+</sup> π <sup>-</sup>
-------------------------------------	------	----------------------	----	--

<sup>1</sup> From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering ρ(1450), ρ(1700) and ρ(2150) with the parameters of the ρ(1450) and ρ(1700) floating and the mass and width of the ρ(2150) fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π, 0 and π, respectively.

NODE=M065R00  
NODE=M065R00

NODE=M065R00;LINKAGE=A

## ρ(1700) BRANCHING RATIOS

NODE=M065230

$$\Gamma(\rho\pi\pi)/\Gamma(4\pi)$$

$$\Gamma_3/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.28±0.06	<sup>1</sup> ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup> ωπ not included.

NODE=M065R19  
NODE=M065R19

NODE=M065R;LINKAGE=BL

$$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$$

$$\Gamma_4/\Gamma_2$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0		DEL COURT	81B	DM1 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )
0.7 ±0.1	500	SCHACHT	74	STRC 5.5–18 γp → p4π
0.80		<sup>1</sup> BINGHAM	72B	HBC 9.3 γp → p4π

<sup>1</sup> The ππ system is in S-wave.

NODE=M065R1  
NODE=M065R1

NODE=M065R1;LINKAGE=S

$$\Gamma(\rho^0\pi^0\pi^0)/\Gamma(\rho^\pm\pi^\mp\pi^0)$$

$$\Gamma_5/\Gamma_6$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10		ATKINSON	85B	OMEG 20–70 γp
<0.15		ATKINSON	82	OMEG 0 20–70 γp → p4π

NODE=M065R6  
NODE=M065R6

$$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$$

$$\Gamma_7/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.16±0.05	<sup>1</sup> ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

<sup>1</sup> ωπ not included.

NODE=M065R15  
NODE=M065R15

NODE=M065R15;LINKAGE=BL

$\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.06	<sup>1</sup> ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>  $\omega\pi$  not included.

 $\Gamma_8/\Gamma_1$ NODE=M065R16  
NODE=M065R16

NODE=M065R16;LINKAGE=BL

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.30±0.10	<sup>1</sup> ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>  $\omega\pi$  not included.

 $\Gamma_9/\Gamma_1$ NODE=M065R17  
NODE=M065R17

NODE=M065R17;LINKAGE=BL

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.09±0.03	<sup>1</sup> ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>  $\omega\pi$  not included.

 $\Gamma_{10}/\Gamma_1$ NODE=M065R18  
NODE=M065R18

NODE=M065R18;LINKAGE=BL

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.108±0.017 <sup>+0.162</sup> <sub>-0.004</sub>	<sup>1</sup> ALBRECHT	20	RVUE 0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
0.287 <sup>+0.043</sup> <sub>-0.042</sub>	BECKER	79	ASPK 17 $\pi^-p$ polarized
0.15 to 0.30	<sup>2</sup> MARTIN	78C	RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$
<0.20	<sup>3</sup> COSTA...	77B	RVUE $e^+e^- \rightarrow 2\pi, 4\pi$
0.30 ±0.05	<sup>2</sup> FROGGATT	77	RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$
<0.15	<sup>4</sup> EISENBERG	73	HBC 5 $\pi^+p \rightarrow \Delta^{++}2\pi$
0.25 ±0.05	<sup>5</sup> HYAMS	73	ASPK 17 $\pi^-p \rightarrow \pi^+\pi^-n$

 $\Gamma_{11}/\Gamma$ NODE=M065R5  
NODE=M065R5

<sup>1</sup> Residue from T-matrix pole, 2 poles, 3 channels, Chew-Mandelstam functions and simplified analytic continuation for the  $4\pi$  channel. Includes scattering data from HYAMS 75 and model-independent calculation of GARCIA-MARTIN 11A.

<sup>2</sup> From phase shift analysis of HYAMS 73 data.

<sup>3</sup> Estimate using unitarity, time reversal invariance, Breit-Wigner.

<sup>4</sup> Estimated using one-pion-exchange model.

<sup>5</sup> Included in BECKER 79 analysis.

NODE=M065R5;LINKAGE=A

NODE=M065R5;LINKAGE=P  
NODE=M065R5;LINKAGE=C  
NODE=M065R5;LINKAGE=E  
NODE=M065R5;LINKAGE=H $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.007±0.006 <sup>+0.041</sup> <sub>-0.002</sub>	<sup>1</sup> ALBRECHT	20	RVUE 0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$

 $\Gamma_{16}/\Gamma$ NODE=M065R03  
NODE=M065R03

<sup>1</sup> Residue from T-matrix pole, 2 poles, 3 channels, Chew-Mandelstam functions and simplified analytic continuation for the  $4\pi$  channel. Includes scattering data from HYAMS 75 and model-independent calculation of GARCIA-MARTIN 11A.

NODE=M065R03;LINKAGE=A

 $\Gamma(\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.13±0.05	ASTON	80	OMEG 20-70 $\gamma p \rightarrow p2\pi$
<0.14	<sup>1</sup> DAVIER	73	STRC 6-18 $\gamma p \rightarrow p4\pi$
<0.2	<sup>2</sup> BINGHAM	72B	HBC 9.3 $\gamma p \rightarrow p2\pi$

 $\Gamma_{11}/\Gamma_2$ NODE=M065R3  
NODE=M065R3

<sup>1</sup> Upper limit is estimate.

<sup>2</sup>  $2\sigma$  upper limit.

NODE=M065R3;LINKAGE=E  
NODE=M065R3;LINKAGE=S $\Gamma(\pi\pi)/\Gamma(4\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT
0.16±0.04	<sup>1,2</sup> ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Using ABELE 97.

<sup>2</sup>  $\omega\pi$  not included.

 $\Gamma_{12}/\Gamma_1$ NODE=M065R20  
NODE=M065R20NODE=M065R;LINKAGE=LK  
NODE=M065R20;LINKAGE=BL $\Gamma(K\bar{K}^*(892)+\text{c.c.})/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	COAN	04	CLEO $\tau^- \rightarrow K^-\pi^-K^+\nu_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma_{13}/\Gamma$ NODE=M065R21  
NODE=M065R21

$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(2(\pi^+\pi^-))$  $\Gamma_{13}/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.15±0.03	<sup>1</sup> DELCOURT	81B	DM1 $e^+e^- \rightarrow \bar{K}K\pi$
-----------	-----------------------	-----	--------------------------------------

<sup>1</sup> Assuming  $\rho(1700)$  and  $\omega$  radial excitations to be degenerate in mass.

NODE=M065R9  
NODE=M065R9

NODE=M065R9;LINKAGE=D

 $\Gamma(\eta\rho)/\Gamma_{total}$  $\Gamma_{14}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen		AKHMETSHIN	00D	CMD2 $e^+e^- \rightarrow \eta\pi^+\pi^-$
<0.04		DONNACHIE	87B	RVUE
<0.02	58	ATKINSON	86B	OMEG 20-70 $\gamma p$

NODE=M065R12  
NODE=M065R12

 $\Gamma(\eta\rho)/\Gamma(2(\pi^+\pi^-))$  $\Gamma_{14}/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123±0.027	DELCOURT	82	DM1 $e^+e^- \rightarrow \pi^+\pi^-MM$
~0.1	ASTON	80	OMEG 20-70 $\gamma p$

NODE=M065R8  
NODE=M065R8

 $\Gamma(\pi^+\pi^- \text{ neutrals})/\Gamma(2(\pi^+\pi^-))$  $(\Gamma_5+\Gamma_6+0.714\Gamma_{14})/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6±0.4	<sup>1</sup> BALLAM	74	HBC 9.3 $\gamma p$
---------	---------------------	----	--------------------

<sup>1</sup> Upper limit. Background not subtracted.

NODE=M065R7  
NODE=M065R7

NODE=M065R7;LINKAGE=U

 $\Gamma(a_2(1320)\pi)/\Gamma_{total}$  $\Gamma_{15}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMELIN	00	VES 37 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
----------	--------	----	---

NODE=M065R14  
NODE=M065R14

 $\Gamma(K\bar{K})/\Gamma(2(\pi^+\pi^-))$  $\Gamma_{16}/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.015±0.010		<sup>1</sup> DELCOURT	81B	DM1	$e^+e^- \rightarrow \bar{K}K$
<0.04	95	BINGHAM	72B	HBC	0 9.3 $\gamma p$

<sup>1</sup> Assuming  $\rho(1700)$  and  $\omega$  radial excitations to be degenerate in mass.

NODE=M065R4  
NODE=M065R4

NODE=M065R4;LINKAGE=D

 $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+c.c.)$  $\Gamma_{16}/\Gamma_{13}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.052±0.026	BUON	82	DM1 $e^+e^- \rightarrow \text{hadrons}$
-------------	------	----	---

NODE=M065R10  
NODE=M065R10

 $\Gamma(\pi^0\omega)/\Gamma_{total}$  $\Gamma_{18}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		MATVIENKO	15	BELL $\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
seen	1.6k	ACHASOV	12	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
not seen	2382	AKHMETSHIN	03B	CMD2 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
seen		ACHASOV	97	RVUE $e^+e^- \rightarrow \omega\pi^0$

NODE=M065R13  
NODE=M065R13

 $\Gamma(\pi^0\gamma)/\Gamma_{total}$  $\Gamma_{19}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen <sup>1</sup> ACHASOV 10D SND 1.075-2.0  $e^+e^- \rightarrow \pi^0\gamma$

<sup>1</sup> From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states  $\omega(1420)$ ,  $\rho(1450)$ ,  $\omega(1650)$ , and  $\rho(1700)$ . The width of the highest mass effective resonance is fixed at 315 MeV.

NODE=M065R02  
NODE=M065R02

NODE=M065R02;LINKAGE=A

 $\Gamma(f_0(1500)\gamma)/\Gamma_{total}$  $\Gamma_{20}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen <sup>1</sup> ACHASOV 22 SND 1.17-2.00  $e^+e^- \rightarrow \eta\eta\gamma$

<sup>1</sup> The 90% CL upper limit on the Born cross sections  $\sigma(e^+e^- \rightarrow \phi(1680) \rightarrow f'_2(1525)\gamma \rightarrow \eta\eta\gamma)$  and  $\sigma(e^+e^- \rightarrow \rho(1700) \rightarrow f_0(1500)\gamma \rightarrow \eta\eta\gamma)$  is 10.6 pb.

NODE=M065R04  
NODE=M065R04

NODE=M065R04;LINKAGE=A



$\rho(1700)$  REFERENCES

NODE=M065

ACHASOV	22	EPJ C82 168	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61655
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
GRIBANOV	20	JHEP 2001 112	S.S. Gribov <i>et al.</i>	(CMD-3 Collab.)	REFID=60620
ACHASOV	18	PR D97 012008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58785
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16D	PR D94 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57618
AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
		Translated from ZETFP 94 796.			
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
GARCIA-MAR...	11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=54121
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
FRABETTI	04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=49614
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACHASOV	00I	PL B486 29	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47931
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=41355
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
DIEKMAN	88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
ATKINSON	86B	ZPHY C30 531	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21508
ATKINSON	85B	ZPHY C26 499	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21506
ERKAL	85	ZPHY C29 485	C. Erkal, M.G. Olsson	(WISC)	REFID=20136
ABE	84B	PRL 53 751	K. Abe <i>et al.</i>	(SLAC HFP Collab.)	REFID=21503
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
		Translated from ZETFP 37 613.			
ATKINSON	82	PL 108B 55	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21493
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)	REFID=21494
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
CORDIER	82	PL 109B 129	A. Cordier <i>et al.</i>	(LALO)	REFID=21495
DELCOURT	82	PL 113B 93	B. Delcourt <i>et al.</i>	(LALO)	REFID=21496
ASTON	81E	NP B189 15	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21487
DELCOURT	81B	Bonn Conf. 205	B. Delcourt	(ORSAY)	REFID=21490
		Also			
		PL 109B 129	A. Cordier <i>et al.</i>	(LALO)	REFID=21495
DIBIANCA	81	PR D23 595	F.A. di Bianca <i>et al.</i>	(CASE, CMU)	REFID=21492
ASTON	80	PL 92B 215	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21478
BACCI	80	PL 95B 139	C. Bacci <i>et al.</i>	(ROMA, FRAS)	REFID=21481
BIZOT	80	Madison Conf. 546	J.C. Bizot <i>et al.</i>	(LALO, MONP)	REFID=21482
KILLIAN	80	PR D21 3005	T.J. Killian <i>et al.</i>	(CORN)	REFID=21484
ATIYA	79B	PRL 43 1691	M.S. Atiya <i>et al.</i>	(COLU, ILL, FNAL)	REFID=21470
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)	REFID=20126
MARTIN	78C	ANP 114 1	A.D. Martin, M.R. Pennington	(CERN)	REFID=21661
COSTA...	77B	PL 71B 345	B. Costa de Beauregard, B. Pire, T.N. Truong	(EPOL)	REFID=21465
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
ALEXANDER	75	PL 57B 487	G. Alexander <i>et al.</i>	(TELA)	REFID=21450
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
BALLAM	74	NP B76 375	J. Ballam <i>et al.</i>	(SLAC, LBL, MPIM)	REFID=20610
CONVERSI	74	PL 52B 493	M. Conversi <i>et al.</i>	(ROMA, FRAS)	REFID=20637
SCHACHT	74	NP B81 205	P. Schacht <i>et al.</i>	(MPIM)	REFID=21449
DAVIER	73	NP B58 31	M. Davier <i>et al.</i>	(SLAC)	REFID=21434
EISENBERG	73	PL 43B 149	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=21435
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
BINGHAM	72B	PL 41B 635	H.H. Bingham <i>et al.</i>	(LBL, UCB, SLAC) IGJP	REFID=21426
JACOB	72	PR D5 1847	M. Jacob, R. Slansky		REFID=49668
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

$a_2(1700)$ 

$$I^G(J^{PC}) = 1^-(2^{++})$$

NODE=M162

 $a_2(1700)$  MASS

NODE=M162M

NODE=M162M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1698 ± 44</b>		<sup>1</sup> AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1686 ± 22 $^{+19}_{-7}$		<sup>2</sup> KOPF 21	RVUE	0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
1638.9 ± 2.3 $^{+57.4}_{-0.1}$		<sup>3</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
1722 ± 15 ± 67		<sup>4</sup> RODAS 19	RVUE	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
1681 $^{+22}_{-35}$	46M	<sup>5,6</sup> AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow$ $\pi^- \pi^+ \pi^- p$
1720 ± 10 ± 60		<sup>7</sup> JACKURA 18	RVUE	$\pi^- p \rightarrow \eta \pi^- p$
1726 ± 12 ± 25		<sup>6</sup> ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
1675 ± 25		ANISOVICH 09	RVUE	0.0 $\bar{p}p$ , $\pi N$
1722 ± 9 ± 15	18k	<sup>8</sup> SCHEGELSKY 06	RVUE	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$
1702 ± 7	80k	<sup>9</sup> UMAN 06	E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
1721 ± 13 ± 44	145k	LU 05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1737 ± 5 ± 7		ABE 04	BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
1767 ± 14	221	<sup>10</sup> ACCIARRI 01H	L3	$\gamma \gamma \rightarrow K_S^0 K_S^0$ , $E_{cm}^{ee} =$ 91, 183–209 GeV
1660 ± 40		<sup>6</sup> ABELE 99B	CBAR	1.94 $\bar{p}p \rightarrow \pi^0 \eta \eta$
~ 1775		<sup>11</sup> GRYGOREV 99	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1752 ± 21 ± 4		ACCIARRI 97T	L3	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> T-matrix pole.<sup>2</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi$ ,  $\eta'\pi$  and  $K\bar{K}$  systems.<sup>3</sup> T-matrix pole, 2 poles, 2 channels ( $\pi\eta$ ,  $K\bar{K}$ ).<sup>4</sup> The coupled-channel analysis of both the  $\eta\pi$  and  $\eta'\pi$  systems using ADOLPH 15 data. The mass is extracted from the T-matrix pole.<sup>5</sup> Statistical error negligible.<sup>6</sup> Breit-Wigner mass.<sup>7</sup> Superseded by RODAS 19.<sup>8</sup> From analysis of L3 data at 183–209 GeV.<sup>9</sup> Statistical error only.<sup>10</sup> Spin 2 dominant, isospin not determined, could also be  $I=1$ .<sup>11</sup> Possibly two  $J^P = 2^+$  resonances with isospins 0 and 1.

NODE=M162M;LINKAGE=TT

NODE=M162M;LINKAGE=H

NODE=M162M;LINKAGE=G

NODE=M162M;LINKAGE=C

NODE=M162M;LINKAGE=B

NODE=M162M;LINKAGE=E

NODE=M162M;LINKAGE=D

NODE=M162M;LINKAGE=SC

NODE=M162M;LINKAGE=ST

NODE=M162M;LINKAGE=HA

NODE=M162M;LINKAGE=GR

 $a_2(1700)$  WIDTH

NODE=M162W

NODE=M162W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>265 ± 55</b>		<sup>1</sup> AMSLER 02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
421 ± 75 $^{+64}_{-57}$		<sup>2</sup> KOPF 21	RVUE	0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
224.0 ± 2.5 $^{+1.8}_{-48.3}$		<sup>3</sup> ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$ , $\pi^0 \eta \eta$ , $\pi^0 K^+ K^-$
247 ± 17 ± 63		<sup>4</sup> RODAS 19	RVUE	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
436 $^{+20}_{-16}$	46M	<sup>5,6</sup> AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
280 ± 10 ± 70		<sup>7</sup> JACKURA 18	RVUE	$\pi^- p \rightarrow \eta \pi^- p$
190 ± 18 ± 30		<sup>6</sup> ABLIKIM 17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
270 $^{+50}_{-20}$		ANISOVICH 09	RVUE	0.0 $\bar{p}p$ , $\pi N$

336 ± 20 ± 20	18k	<sup>8</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
417 ± 19	80k	<sup>9</sup> UMAN 06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
279 ± 49 ± 66	145k	LU 05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
151 ± 22 ± 24		ABE 04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
187 ± 60	221	<sup>10</sup> ACCIARRI 01H	L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
280 ± 70		<sup>6</sup> ABELE 99B	CBAR	$1.94 \bar{p}p \rightarrow \pi^0 \eta\eta$
150 ± 110 ± 34		ACCIARRI 97T	L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> T-matrix pole.

<sup>2</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi$ ,  $\eta'\pi$  and  $K\bar{K}$  systems.

<sup>3</sup> T-matrix pole, 2 poles, 2 channels ( $\pi\eta$ ,  $K\bar{K}$ ).

<sup>4</sup> The coupled-channel analysis of both the  $\eta\pi$  and  $\eta'\pi$  systems using ADOLPH 15 data. The width is extracted from the T-matrix pole.

<sup>5</sup> Statistical error negligible.

<sup>6</sup> Breit-Wigner width.

<sup>7</sup> Superseded by RODAS 19.

<sup>8</sup> From analysis of L3 data at 183–209 GeV.

<sup>9</sup> Statistical error only.

<sup>10</sup> Spin 2 dominant, isospin not determined, could also be  $I=1$ .

NODE=M162W;LINKAGE=TT  
NODE=M162W;LINKAGE=H

NODE=M162W;LINKAGE=G  
NODE=M162W;LINKAGE=C

NODE=M162W;LINKAGE=B  
NODE=M162W;LINKAGE=E  
NODE=M162W;LINKAGE=D  
NODE=M162W;LINKAGE=SC  
NODE=M162W;LINKAGE=ST  
NODE=M162W;LINKAGE=HA

### $a_2(1700)$ DECAY MODES

NODE=M162215;NODE=M162

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta\pi$	(3.6 ± 1.1) %
$\Gamma_2$ $\eta'\pi$	seen
$\Gamma_3$ $\gamma\gamma$	(1.13 ± 0.30) × 10 <sup>-6</sup>
$\Gamma_4$ $\rho\pi$	seen
$\Gamma_5$ $f_2(1270)\pi$	seen
$\Gamma_6$ $K\bar{K}$	(1.9 ± 1.2) %
$\Gamma_7$ $\omega\pi^-\pi^0$	seen
$\Gamma_8$ $\omega\rho$	seen

DESIG=4  
DESIG=8;OUR EVAL;→ UNCHECKED ←  
DESIG=1  
DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=3;OUR EVAL;→ UNCHECKED ←  
DESIG=5  
DESIG=6;OUR EVAL;→ UNCHECKED ←  
DESIG=7;OUR EVAL;→ UNCHECKED ←

### $a_2(1700)$ PARTIAL WIDTHS

NODE=M162220

$\Gamma(\eta\pi)$					$\Gamma_1$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>9.5 ± 2.0</b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M162W3  
NODE=M162W3

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W3;LINKAGE=SC

$\Gamma(\gamma\gamma)$					$\Gamma_3$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.30 ± 0.05</b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M162W2  
NODE=M162W2

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W2;LINKAGE=SC

$\Gamma(K\bar{K})$					$\Gamma_6$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>5.0 ± 3.0</b>	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M162W1  
NODE=M162W1

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV, using  $a_2(1700)$  mass of 1730 MeV and width of 340 MeV, and SU(3) relations.

NODE=M162W1;LINKAGE=SC

### $a_2(1700)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M162225

$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$(\Gamma_4 + \Gamma_5)\Gamma_3/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.29 ± 0.04 ± 0.02</b>		ACCIARRI 97T	L3	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	

NODE=M162G1  
NODE=M162G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.37 <sup>+0.12</sup> <sub>-0.08</sub> ± 0.10	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
---	-----	----------------------------	------	--

<sup>1</sup> From analysis of L3 data at 183–209 GeV.

NODE=M162G1;LINKAGE=SC

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_6\Gamma_3/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$20.6 \pm 4.2 \pm 4.6$	<sup>1</sup> ABE	04 BELL	$10.6 e^+e^- \rightarrow e^+e^- K^+K^-$
$49 \pm 11 \pm 13$	<sup>2</sup> ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183\text{--}209 \text{ GeV}$

<sup>1</sup> Assuming spin 2.<sup>2</sup> Spin 2 dominant, isospin not determined, could also be  $l=1$ .NODE=M162G2  
NODE=M162G2NODE=M162G2;LINKAGE=AB  
NODE=M162G;LINKAGE=HA $a_2(1700)$  BRANCHING RATIOS $\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$  $\Gamma_4/\Gamma_5$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$3.4 \pm 0.4 \pm 0.1$	18k	<sup>1</sup> SCHEGELSKY	06 RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> From analysis of L3 data at 183–209 GeV.

NODE=M162235

NODE=M162R01  
NODE=M162R01

NODE=M162R01;LINKAGE=SC

 $\Gamma(K\bar{K})/\Gamma(\eta\pi)$  $\Gamma_6/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.029 \pm 0.04 \begin{smallmatrix} +0.011 \\ -0.012 \end{smallmatrix}$	<sup>1</sup> KOPF	21 RVUE	$0.9 p\bar{p} \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0 K^+K^- \text{ and } 191 \pi^-p \rightarrow \pi^-\pi^-\pi^+p$
$4.134 \pm 0.106 \begin{smallmatrix} +4.909 \\ -2.988 \end{smallmatrix}$	<sup>2</sup> ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0 K^+K^-$

<sup>1</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.<sup>2</sup> Residues from T-matrix pole, 2 poles, 2 channels ( $\pi\eta, K\bar{K}$ ).NODE=M162R00  
NODE=M162R00

NODE=M162R00;LINKAGE=B

NODE=M162R00;LINKAGE=A

 $\Gamma(\eta'\pi)/\Gamma(\eta\pi)$  $\Gamma_2/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.035 \pm 0.044 \begin{smallmatrix} +0.069 \\ -0.012 \end{smallmatrix}$	<sup>1</sup> KOPF	21 RVUE	$0.9 p\bar{p} \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0 K^+K^- \text{ and } 191 \pi^-p \rightarrow \pi^-\pi^-\pi^+p$

<sup>1</sup> From T-matrix pole based on combined fit of Crystal Barrel and  $\pi\pi$  scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of  $\eta\pi, \eta'\pi$  and  $K\bar{K}$  systems.NODE=M162R02  
NODE=M162R02

NODE=M162R02;LINKAGE=A

 $a_2(1700)$  REFERENCES

KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)	REFID=61470
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=59554
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)	REFID=59003
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57953
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
AMSLER	02	EPJ C23 29	C. Amisler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46904
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>		REFID=46909
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761

NODE=M162

**$a_0(1710)$** 

$$I^G(J^{PC}) = 1^-(0^{++})$$

## OMITTED FROM SUMMARY TABLE

Evidence for this state is also inferred from the interference of the  $K^+K^-$  and  $K_S^0K_S^0$  decays of the  $f_0(1710)$  in  $D_s^+ \rightarrow f_0(1710)\pi^+$ , leading to a relative branching ratio an order of magnitude larger than expected from isospin symmetry (ABLIKIM 22F). See also the review on "Spectroscopy of Light Meson Resonances."

NODE=M263

NODE=M263

 **$a_0(1710)$  MASS**

NODE=M263M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1711 ± 27 OUR AVERAGE</b>	Error includes scale factor of 5.1.		
1817 ± 8 ± 20	<sup>1</sup> ABLIKIM	22AH BES3	$D_s^+ \rightarrow K_S^0 K^+ \pi^0$
1704 ± 5 ± 2	LEES	21A BABR	$\eta_c(1S) \rightarrow \pi^+ \pi^- \eta$
<sup>1</sup> Observed to decay into $K_S^0 K^+$ in a Breit-Wigner amplitude analysis involving $D_s^+$ decays into $\bar{K}^*(892)^0 K^+$ , $\bar{K}^*(892)^+ K_S^0$ , $\bar{K}^*(1410)^0 K^+$ , $a_0(980)^+ \pi^0$ , and $a_0(1817)^+ \pi^0$ .			

NODE=M263M

NODE=M263M;LINKAGE=A

 **$a_0(1710)$  WIDTH**

NODE=M263W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>106 ± 15 OUR AVERAGE</b>			
97 ± 22 ± 15	<sup>1</sup> ABLIKIM	22AH BES3	$D_s^+ \rightarrow K_S^0 K^+ \pi^0$
110 ± 15 ± 11	LEES	21A BABR	$\eta_c(1S) \rightarrow \pi^+ \pi^- \eta$
<sup>1</sup> Observed to decay into $K_S^0 K^+$ in a Breit-Wigner amplitude analysis involving $D_s^+$ decays into $\bar{K}^*(892)^0 K^+$ , $\bar{K}^*(892)^+ K_S^0$ , $\bar{K}^*(1410)^0 K^+$ , $a_0(980)^+ \pi^0$ , and $a_0(1817)^+ \pi^0$ .			

NODE=M263W

NODE=M263W;LINKAGE=A

 **$a_0(1710)$  DECAY MODES**

NODE=M263215;NODE=M263

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\eta$	seen
$\Gamma_2$ $K^+K^-$	
$\Gamma_3$ $K_S^0K_S^0$	
$\Gamma_4$ $K_S^0K^+$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

$\Gamma(\pi\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>seen</b>	LEES	21A	$\eta_c(1S) \rightarrow \pi^+ \pi^- \eta$	

NODE=M263R01

NODE=M263R01

$\Gamma(K^+K^-)/\Gamma(K_S^0K_S^0)$	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_3$
<b>0.32 ± 0.12</b>	<sup>1</sup> ABLIKIM	22F BES3	$D_s^+ \rightarrow K_S^0 K_S^0 \pi^+$	

NODE=M263R00

NODE=M263R00

OCCUR=2

<sup>1</sup> Using  $D_s^+ \rightarrow K^+K^- \pi^+$  from ABLIKIM 21AE. The apparent violation of isospin symmetry may be due to a destructive interference with the  $f_0(1710)$  in the  $K^+K^-$  channel, and a constructive interference in the  $K_S^0K_S^0$  channel.

NODE=M263R00;LINKAGE=B

$\Gamma(K_S^0K^+)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma$
<b>seen</b>	ABLIKIM	22AH BES3	$D_s^+ \rightarrow K_S^0 K^+ \pi^0$	

NODE=M263R02

NODE=M263R02

 **$a_0(1710)$  REFERENCES**

NODE=M263

ABLIKIM	22AH PRL 129 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61880
ABLIKIM	22F PR D105 L051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61641
ABLIKIM	21AE PR D104 012016	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61367
LEES	21A PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61442

$f_0(1710)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

 **$f_0(1710)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1680–1820) – <math>i</math> (50–180) OUR ESTIMATE</b>			
$(1769 \pm 8) - i(78 \pm 6)$	<sup>1</sup> RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
$(1700 \pm 18) - i(127 \pm 12)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1803 \pm 3.5^{+45.5}_{-10.4}) - i(145 \pm 2.5^{+16.3}_{-9.6})$	<sup>2</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1732 \pm 15) - i(160^{+25}_{-10})$	<sup>3</sup> ANISOVICH	03	RVUE $\pi\pi, K\bar{K}, \eta\eta, \eta\eta'$
$(1698 \pm 18) - i(60 \pm 13)$	BARBERIS	00E	OMEG 450 $p\bar{p} \rightarrow p_f \eta \eta p_s$
$(1770 \pm 12) - i(110 \pm 20)$	<sup>4</sup> ANISOVICH	99B	SPEC $0.6-1.2 p\bar{p} \rightarrow \eta\eta\pi^0$
$(1727 \pm 12 \pm 11) - i(63 \pm 8 \pm 9)$	BARBERIS	99D	OMEG 450 $p\bar{p} \rightarrow K^+ K^-$
$(1750 \pm 30) - i(125 \pm 70)$	ANISOVICH	98B	RVUE Compilation

<sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma\pi^0\pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).

<sup>2</sup> T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ), and BINON 84C ( $\eta\eta'$ ).

<sup>3</sup> Solution I.

<sup>4</sup> Not seen by AMSLER 02.

 **$f_0(1710)$  Breit-Wigner MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1733^{+8}_{-7}</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.5. See the ideogram below.		
$1757 \pm 24 \pm 9$		LEES	21A	BABR $\eta_c(1S) \rightarrow \eta' K^+ K^-$
$1759 \pm 6^{+14}_{-25}$	5.5k	<sup>1</sup> ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
$1750^{+6}_{-7}^{+29}_{-18}$		<sup>2</sup> UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
$1701 \pm 5^{+9}_{-2}$	4k	<sup>3</sup> CHEKANOV	08	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
$1765^{+4}_{-3} \pm 13$		ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+ \pi^-$
$1738 \pm 30$		ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
$1740 \pm 4^{+10}_{-25}$		BAI	03G	BES $J/\psi \rightarrow \gamma K\bar{K}$
$1740^{+30}_{-25}$		BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
$1710 \pm 25$		<sup>4</sup> FRENCH	99	300 $p\bar{p} \rightarrow p_f(K^+ K^-) p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1744 \pm 7 \pm 5$	381	<sup>5,6</sup> DOBBS	15	$J/\psi \rightarrow \gamma\pi^+ \pi^-$
$1705 \pm 11 \pm 5$	237	<sup>5,6</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+ \pi^-$
$1706 \pm 4 \pm 5$	1.0k	<sup>5,6</sup> DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
$1690 \pm 8 \pm 3$	349	<sup>5,6</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
$1750 \pm 13$		AMSLER	06	CBAR $1.64 \bar{p}p \rightarrow K^+ K^- \pi^0$
$1747 \pm 5$	80k	<sup>7</sup> UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
$1776 \pm 15$		VLADIMIRSK...	06	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1670 \pm 20$		BINON	05	GAMS $33 \pi^- p \rightarrow \eta\eta n$
$1682 \pm 16$		TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
$1670 \pm 26$	3.6k	<sup>8</sup> NICHITIU	02	OBLX $0 \bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1730 \pm 15$		BARBERIS	99	OMEG 450 $p\bar{p} \rightarrow p_s p_f K^+ K^-$
$1750 \pm 20$		BARBERIS	99B	OMEG 450 $p\bar{p} \rightarrow p_s p_f \pi^+ \pi^-$
$1720 \pm 39$		BAI	98H	BES $J/\psi \rightarrow \gamma\pi^0 \pi^0$
$1775 \pm 1.5$	57	<sup>9</sup> BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$
$1690 \pm 11$		<sup>10</sup> ABREU	96C	DLPH $Z^0 \rightarrow K^+ K^- + X$
$1696 \pm 5^{+9}_{-34}$		<sup>11</sup> BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$

NODE=M068

NODE=M068

NODE=M068PP

NODE=M068PP

NODE=M068PP

→ UNCHECKED ←

OCCUR=3

NODE=M068PP;LINKAGE=O

NODE=M068PP;LINKAGE=H

NODE=M068PP;LINKAGE=A

NODE=M068PP;LINKAGE=AV

NODE=M068M


NODE=M068M

OCCUR=2

OCCUR=3

OCCUR=4

1781± 8	$^{+10}_{-31}$	BAI	96C	BES	$J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=2
1768±14		BALOSHIN	95	SPEC	$40 \pi^- C \rightarrow K_S^0 K_S^0 X$	
1750±15		<sup>12</sup> BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$	
1620±16		<sup>11</sup> BUGG	95	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$	OCCUR=2
1748±10		<sup>13</sup> ARMSTRONG	93C	E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
~ 1750		BREAKSTONE	93	SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$	
1744±15		<sup>14</sup> ALDE	92D	GAM2	$38 \pi^- p \rightarrow \eta \eta n$	
1713±10		<sup>15</sup> ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K^+ K^-$	
1706±10		<sup>15</sup> ARMSTRONG	89D	OMEG	$300 p p \rightarrow p p K_S^0 K_S^0$	OCCUR=2
1707±10		<sup>13</sup> AUGUSTIN	88	DM2	$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$	
1700±15		<sup>11</sup> BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
1720±60		BOLONKIN	88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	OCCUR=2
1638±10		<sup>16</sup> FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	
1690± 4		<sup>17</sup> FALVARD	88	DM2	$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	OCCUR=2
1698±15		<sup>13</sup> AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$	
1720±10	$\pm 10$	<sup>11</sup> BALTRUSAIT..	87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$	
1755± 8		<sup>18</sup> ALDE	86C	GAM2	$38 \pi^- p \rightarrow n 2\eta$	
1730 $^{+2}_{-10}$		<sup>19</sup> LONGACRE	86	RVUE	$22 \pi^- p \rightarrow n 2K_S^0$	
1742±15		<sup>13</sup> WILLIAMS	84	MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$	
1670±50		BLOOM	83	CBAL	$J/\psi \rightarrow \gamma 2\eta$	
1650±50		BURKE	82	MRK2	$J/\psi \rightarrow \gamma 2\rho$	
1640±50		<sup>20,21</sup> EDWARDS	82D	CBAL	$J/\psi \rightarrow \gamma 2\eta$	
1730±10	$\pm 20$	<sup>22</sup> ETKIN	82C	MPS	$23 \pi^- p \rightarrow n 2K_S^0$	
		<sup>1</sup>			From partial wave analysis including all possible combinations of $0^{++}$ , $2^{++}$ , and $4^{++}$ resonances.	NODE=M068M;LINKAGE=D
		<sup>2</sup>			Spin 0 favored over spin 2.	NODE=M068M;LINKAGE=H
		<sup>3</sup>			In the SU(3) based model with a specific interference pattern of the $f_2(1270)$ , $a_2^0(1320)$ , and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.	NODE=M068M;LINKAGE=HE
		<sup>4</sup>			$J^P = 0^+$ , supersedes ARMSTRONG 89D.	NODE=M068M;LINKAGE=C3
		<sup>5</sup>			Using CLEO-c data but not authored by the CLEO Collaboration.	NODE=M068M;LINKAGE=F
		<sup>6</sup>			From a fit to a Breit-Wigner line shape with fixed $\Gamma = 135$ MeV.	NODE=M068M;LINKAGE=G
		<sup>7</sup>			Systematic errors not estimated.	NODE=M068M;LINKAGE=CH
		<sup>8</sup>			Decaying to $f_0(1370)\pi\pi$ .	NODE=M068M;LINKAGE=NC
		<sup>9</sup>			No $J^{PC}$ determination.	NODE=M068M;LINKAGE=4A
		<sup>10</sup>			No $J^{PC}$ determination, width not determined.	NODE=M068M;LINKAGE=A4
		<sup>11</sup>			$J^P = 2^+$ .	NODE=M068M;LINKAGE=A3
		<sup>12</sup>			From a fit to the $0^+$ partial wave.	NODE=M068M;LINKAGE=Q0
		<sup>13</sup>			No $J^{PC}$ determination.	NODE=M068M;LINKAGE=A1
		<sup>14</sup>			ALDE 92D combines all the GAMS-2000 data.	NODE=M068M;LINKAGE=AA
		<sup>15</sup>			$J^P = 2^+$ , superseded by FRENCH 99.	NODE=M068M;LINKAGE=C
		<sup>16</sup>			From an analysis ignoring interference with $f_2'(1525)$ .	NODE=M068M;LINKAGE=A
		<sup>17</sup>			From an analysis including interference with $f_2'(1525)$ .	NODE=M068M;LINKAGE=B
		<sup>18</sup>			Superseded by ALDE 92D.	NODE=M068M;LINKAGE=BB
		<sup>19</sup>			Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.	NODE=M068M;LINKAGE=A9
		<sup>20</sup>			$J^P = 2^+$ preferred.	NODE=M068M;LINKAGE=B2
		<sup>21</sup>			From fit neglecting nearby $f_2'(1525)$ . Replaced by BLOOM 83.	NODE=M068M;LINKAGE=E
		<sup>22</sup>			Superseded by LONGACRE 86.	NODE=M068M;LINKAGE=B1



**$f_0(1710)$  Breit-Wigner WIDTH**

NODE=M068W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>150 <math>\pm</math> 12 <math>\pm</math> 10</b>		<b>OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.
175 $\pm$ 23 $\pm$ 4		LEES	21A BABR	$\eta_c(1S) \rightarrow \eta' K^+ K^-$
172 $\pm$ 10 $\pm$ 32 $\pm$ 16	5.5k	1 ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
139 $\pm$ 11 $\pm$ 96 $\pm$ 12 $\pm$ 50		2 UEHARA	13 BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
100 $\pm$ 24 $\pm$ 7 $\pm$ 22	4k	3 CHEKANOV	08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
145 $\pm$ 8 $\pm$ 69		ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
125 $\pm$ 20		ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
166 $\pm$ 5 $\pm$ 7 $\pm$ 8 $\pm$ 15 $\pm$ 10		BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
120 $\pm$ 50 $\pm$ 40		BAI	00A BES	$J/\psi \rightarrow \gamma (\pi^+ \pi^- \pi^+ \pi^-)$
105 $\pm$ 34		4 FRENCH	99	300 $p p \rightarrow p_f (K^+ K^-) p_S$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
148 $\pm$ 40 $\pm$ 30		AMSLER	06 CBAR	1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$
188 $\pm$ 13	80k	5 UMAN	06 E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
250 $\pm$ 30		VLADIMIRSK..06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
260 $\pm$ 50		BINON	05 GAMS	33 $\pi^- p \rightarrow \eta \eta n$
102 $\pm$ 26		TIKHOMIROV	03 SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
267 $\pm$ 44	3651	6 NICHITIU	02 OBLX	0 $\bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
100 $\pm$ 25		BARBERIS	99 OMEG	450 $p p \rightarrow p_S p_f K^+ K^-$
160 $\pm$ 30		BARBERIS	99B OMEG	450 $p p \rightarrow p_S p_f \pi^+ \pi^-$
30 $\pm$ 7	57	7 BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$
103 $\pm$ 18 $\pm$ 30 $\pm$ 11		8 BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
85 $\pm$ 24 $\pm$ 22 $\pm$ 19		BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
56 $\pm$ 19		BALOSHIN	95 SPEC	40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
160 $\pm$ 40		9 BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
160 $\pm$ 60 $\pm$ 20		8 BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
264 $\pm$ 25		10 ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6 \gamma$
200 to 300		BREAKSTONE	93 SFM	$p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
< 80 90% CL		11 ALDE	92D GAM2	38 $\pi^- p \rightarrow \eta \eta N^*$
181 $\pm$ 30		12 ARMSTRONG	89D OMEG	300 $p p \rightarrow p p K^+ K^-$

NODE=M068W

OCCUR=2


OCCUR=2



104 ± 30	12 ARMSTRONG 89D OMEG	300 pp → ppK <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup>	OCCUR=2
166.4 ± 33.2	10 AUGUSTIN 88 DM2	J/ψ → γK <sup>+</sup> K <sup>-</sup> , K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup>	
30 ± 20	8 BOLONKIN 88 SPEC	40 π <sup>-</sup> p → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> n	
350 ± 150	BOLONKIN 88 SPEC	40 π <sup>-</sup> p → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> n	OCCUR=2
148 ± 17	13 FALVARD 88 DM2	J/ψ → φK <sup>+</sup> K <sup>-</sup> , K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup>	
184 ± 6	14 FALVARD 88 DM2	J/ψ → φK <sup>+</sup> K <sup>-</sup> , K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup>	OCCUR=2
136 ± 28	10 AUGUSTIN 87 DM2	J/ψ → γπ <sup>+</sup> π <sup>-</sup>	
130 ± 20	8 BALTRUSAIT..87 MRK3	J/ψ → γK <sup>+</sup> K <sup>-</sup>	
122 + 74 - 15	15 LONGACRE 86 RVUE	22 π <sup>-</sup> p → n2K <sub>S</sub> <sup>0</sup>	
57 ± 38	16 WILLIAMS 84 MPSF	200 π <sup>-</sup> N → 2K <sub>S</sub> <sup>0</sup> X	
160 ± 80	BLOOM 83 CBAL	J/ψ → γ2η	
200 ± 100	BURKE 82 MRK2	J/ψ → γ2ρ	
220 + 100 - 70	17,18 EDWARDS 82D CBAL	J/ψ → γ2η	
200 + 156 - 9	19 ETKIN 82B MPS	23 π <sup>-</sup> p → n2K <sub>S</sub> <sup>0</sup>	

- 1 From partial wave analysis including all possible combinations of 0<sup>++</sup>, 2<sup>++</sup>, and 4<sup>++</sup> resonances.
- 2 Spin 0 favored over spin 2.
- 3 In the SU(3) based model with a specific interference pattern of the f<sub>2</sub>(1270), a<sub>2</sub><sup>0</sup>(1320), and f<sub>2</sub>'(1525) mesons incoherently added to the f<sub>0</sub>(1710) and non-resonant background.
- 4 J<sup>P</sup> = 0<sup>+</sup>, supersedes ARMSTRONG 89D.
- 5 Systematic errors not estimated.
- 6 Decaying to f<sub>0</sub>(1370)ππ.
- 7 No J<sup>PC</sup> determination.
- 8 J<sup>P</sup> = 2<sup>+</sup>.
- 9 From a fit to the 0<sup>+</sup> partial wave.
- 10 No J<sup>PC</sup> determination.
- 11 ALDE 92D combines all the GAMS-2000 data.
- 12 J<sup>P</sup> = 2<sup>+</sup>, (0<sup>+</sup> excluded).
- 13 From an analysis ignoring interference with f<sub>2</sub>'(1525).
- 14 From an analysis including interference with f<sub>2</sub>'(1525).
- 15 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
- 16 No J<sup>PC</sup> determination.
- 17 J<sup>P</sup> = 2<sup>+</sup> preferred.
- 18 From fit neglecting nearby f<sub>2</sub>'(1525). Replaced by BLOOM 83.
- 19 From an amplitude analysis of the K<sub>S</sub><sup>0</sup>K<sub>S</sub><sup>0</sup> system, superseded by LONGACRE 86.

NODE=M068W;LINKAGE=F  
 NODE=M068W;LINKAGE=G  
 NODE=M068W;LINKAGE=HE  
 NODE=M068W;LINKAGE=C3  
 NODE=M068W;LINKAGE=CH  
 NODE=M068W;LINKAGE=NC  
 NODE=M068W;LINKAGE=4A  
 NODE=M068W;LINKAGE=A3  
 NODE=M068W;LINKAGE=Q0  
 NODE=M068W;LINKAGE=A1  
 NODE=M068W;LINKAGE=AA  
 NODE=M068W;LINKAGE=B  
 NODE=M068W;LINKAGE=C  
 NODE=M068W;LINKAGE=D  
 NODE=M068W;LINKAGE=A9  
 NODE=M068M;LINKAGE=WI  
 NODE=M068W;LINKAGE=B2  
 NODE=M068W;LINKAGE=E  
 NODE=M068W;LINKAGE=A



**$f_0(1710)$  DECAY MODES**

NODE=M068215;NODE=M068

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\bar{K}$	seen
$\Gamma_2$ $\eta\eta$	seen
$\Gamma_3$ $\eta\eta'$	
$\Gamma_4$ $\pi\pi$	seen
$\Gamma_5$ $\gamma\gamma$	seen
$\Gamma_6$ $\omega\omega$	seen

DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=7  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←  
DESIG=4

 **$f_0(1710)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$** 

NODE=M068220

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_5/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M068G2  
NODE=M068G2

$$12^{+3}_{-2} + 227^{+27}_{-8}$$

UEHARA 13 BELL  $\gamma\gamma \rightarrow K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<480	95	ALBRECHT	90G	ARG	$\gamma\gamma \rightarrow K^+ K^-$
<110	95	<sup>1</sup> BEHREND	89C	CELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
<280	95	<sup>1</sup> ALTHOFF	85B	TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

<sup>1</sup> Assuming helicity 2.

NODE=M068G2;LINKAGE=F

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_5/\Gamma$
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M068G3  
NODE=M068G3

<0.82 95 <sup>1</sup> BARATE 00E ALEP  $\gamma\gamma \rightarrow \pi^+ \pi^-$

<sup>1</sup> Assuming spin 0.

NODE=M068G;LINKAGE=Z

 **$f_0(1710)$  BRANCHING RATIOS**

NODE=M068225

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M068R2  
NODE=M068R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	1004	<sup>1</sup> DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$
seen	349	<sup>1</sup> DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$
0.36 ± 0.12		ALBALADEJO	08	RVUE	
0.38 <sup>+0.09</sup> <sub>-0.19</sub>		<sup>2</sup> LONGACRE	86	MPS	$22 \pi^- p \rightarrow n 2K_S^0$

OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M068R2;LINKAGE=A  
NODE=M068R;LINKAGE=L

<sup>2</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M068R1  
NODE=M068R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22 ± 0.12		ALBALADEJO	08	RVUE	
0.18 <sup>+0.03</sup> <sub>-0.13</sub>		<sup>1</sup> LONGACRE	86	RVUE	

<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

NODE=M068R1;LINKAGE=L

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M068R5  
NODE=M068R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	381	<sup>1</sup> DOBBS	15		$J/\psi \rightarrow \gamma \pi^+ \pi^-$
seen	237	<sup>1</sup> DOBBS	15		$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
not seen		AMSLER	02	CBAR	$0.9 \bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
0.039 <sup>+0.002</sup> <sub>-0.024</sub>		<sup>2</sup> LONGACRE	86	RVUE	

OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M068R5;LINKAGE=A  
NODE=M068R5;LINKAGE=L

<sup>2</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$  $\Gamma_4/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.23±0.05</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.		
0.64±0.27	±0.18	LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$ , $\gamma K^+K^-$
0.41 <sup>+0.11</sup> <sub>-0.17</sub>		ABLIKIM	06V BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
0.2 ±0.024±0.036		BARBERIS	99D OMEG 450	$p\bar{p} \rightarrow K^+K^-, \pi^+\pi^-$
0.39±0.14		ARMSTRONG	91 OMEG 300	$p\bar{p} \rightarrow p\bar{p}\pi\pi, p\bar{p}K\bar{K}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.32±0.14		ALBALADEJO	08 RVUE	
< 0.11	95	<sup>1</sup> ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+K^-$
5.8 <sup>+9.1</sup> <sub>-5.5</sub>		<sup>2</sup> ANISOVICH	02D SPEC	Combined fit

<sup>1</sup> Using data from ABLIKIM 04A.<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.NODE=M068R6  
NODE=M068R6NODE=M068R;LINKAGE=AB  
NODE=M068R;LINKAGE=CH $\Gamma(\eta\eta)/\Gamma(K\bar{K})$  $\Gamma_2/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>0.48±0.15</b>		BARBERIS	00E	450 $p\bar{p} \rightarrow p_f\eta\eta p_s$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.46 <sup>+0.70</sup> <sub>-0.38</sub>		<sup>1</sup> ANISOVICH	02D SPEC	Combined fit
<0.02	90	<sup>2</sup> PROKOSHKIN	91 GA24	300 $\pi^- p \rightarrow \pi^- p\eta\eta$

<sup>1</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.<sup>2</sup> Combining results of GAM4 with those of ARMSTRONG 89D.NODE=M068R7  
NODE=M068R7

NODE=M068R7;LINKAGE=CH

NODE=M068R;LINKAGE=A

 $\Gamma(\eta\eta')/\Gamma(\pi\pi)$  $\Gamma_3/\Gamma_4$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.61 × 10<sup>-3</sup></b>	90	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$  P-wave. The quoted measurement will be corrected in a forthcoming erratum.NODE=M068R00  
NODE=M068R00

NODE=M068R00;LINKAGE=A

 $\Gamma(\omega\omega)/\Gamma_{total}$  $\Gamma_6/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	180	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$

NODE=M068R3  
NODE=M068R3 $f_0(1710)$  REFERENCES

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61442
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58950
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ALBALADEJO	08	PRL 101 252002	M. Albaladejo, J.A. Oller		REFID=52656
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirov <i>et al.</i>	(ITEP, Moscow)	REFID=51191
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49740
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>		REFID=46886
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
Translated from UFN 168 481.					

NODE=M068

BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>		REFID=46616
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=44621
BUGG	95	Translated from YAF 58 50.	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ARMSTRONG	93C	PL B353 378	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BREAKSTONE	93	PL B307 394	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)	REFID=43312
ALDE	92D	ZPHY C58 251	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=41591
Also		PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=44696
		SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	
ARMSTRONG	91	Translated from YAF 54 745	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
PROKOSHKIN	91	ZPHY C51 351	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
		SPD 36 155			
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41010
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=21694
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=21418
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)	REFID=21693
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20391
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355

NODE=M255

## X(1750)

$$I^G(J^{PC}) = ?^-(1^{--})$$

### OMITTED FROM SUMMARY TABLE

The X(1750) was separated from the  $\phi(1680)$  in the 2022 listings due to its incompatible mass and incompatible pattern of  $\bar{K}K$  and  $\bar{K}^*(892)K$  branching fractions.

NODE=M255

### X(1750) MASS

NODE=M255M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1753.8 ± 2.7 OUR AVERAGE</b>			
1784 ± 12 $\begin{smallmatrix} +0 \\ -27 \end{smallmatrix}$	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
1753.5 ± 1.5 ± 2.3	LINK	02K FOCS	20-160 $\gamma p \rightarrow K^+ K^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1726 ± 22	BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
1760 ± 20	ATKINSON	85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$
1690 ± 10	ASTON	81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$

NODE=M255M

### X(1750) WIDTH

NODE=M255W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>120 ± 10 OUR AVERAGE</b>			
106 $\begin{smallmatrix} +22 +8 \\ -19 -36 \end{smallmatrix}$	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
122.2 ± 6.2 ± 8.0	LINK	02K FOCS	20-160 $\gamma p \rightarrow K^+ K^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
121 ± 47	BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
80 ± 40	ATKINSON	85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$
100 ± 40	ASTON	81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$

NODE=M255W

### X(1750) DECAY MODES

NODE=M255215;NODE=M255

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^+ K^-$	seen
$\Gamma_2$ $\bar{K}^*(892)^0 K_S^0$	not seen
$\Gamma_3$ $K^*(892)^\pm K^\mp$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←  
 DESIG=2;OUR EVAL;→ UNCHECKED ←  
 DESIG=3;OUR EVAL;→ UNCHECKED ←

### $\Gamma(\bar{K}^*(892)^0 K_S^0)/\Gamma(K^+ K^-)$

$\Gamma_2/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.065</b>	90	LINK	02K FOCS	$\gamma p \rightarrow K^+ K^- p$

NODE=M255R01  
 NODE=M255R01

$\Gamma(K^*(892)^\pm K^\mp)/\Gamma(K^+ K^-)$  $\Gamma_3/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.183	90	LINK	02K FOCS	$\gamma p \rightarrow K^+ K^- p$

NODE=M255R02  
 NODE=M255R02

## X(1750) REFERENCES

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
BUSENITZ	89	PR D40 1	J.K. Busewitz <i>et al.</i>	(ILL, FNAL)
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ASTON	81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)

NODE=M255

REFID=60256  
 REFID=48845  
 REFID=40927  
 REFID=21596  
 REFID=21585

NODE=M114

 $\eta(1760)$ 

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Seen by DM2 in the  $\rho\rho$  system (BISELLO 89B). Structure in this region has been reported before in the same system (BALTRUSAITIS 86B) and in the  $\omega\omega$  system (BALTRUSAITIS 85C, BISELLO 87).

NODE=M114

 $\eta(1760)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1751 ± 15 OUR AVERAGE</b>				
$1768^{+24}_{-25} \pm 10$	465	<sup>1</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
$1744 \pm 10 \pm 15$	1045	<sup>2</sup> ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
$1703^{+12}_{-11} \pm 2$		<sup>3</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
$1760 \pm 11$	320	<sup>4</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

NODE=M114M

NODE=M114M

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> From a single-resonance fit.
- <sup>2</sup> From a partial wave analysis including  $\eta(1760)$ ,  $f_0(1710)$ ,  $f_2(1640)$ , and  $f_2(1910)$ .
- <sup>3</sup> From a two-resonance fit.
- <sup>4</sup> Estimated by us from various fits. Systematic uncertainties not estimated.

OCCUR=2

NODE=M114M;LINKAGE=ZA  
 NODE=M114M;LINKAGE=MA  
 NODE=M114M;LINKAGE=ZH  
 NODE=M114M;LINKAGE=A

 $\eta(1760)$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>240 ± 30 OUR AVERAGE</b>				
$224^{+62}_{-56} \pm 25$	465	<sup>5</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
$244^{+24}_{-21} \pm 25$	1045	<sup>6</sup> ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
$42^{+36}_{-22} \pm 15$		<sup>7</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
$60 \pm 16$	320	<sup>8</sup> BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

NODE=M114W

NODE=M114W

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>5</sup> From a single-resonance fit.
- <sup>6</sup> From a partial wave analysis including  $\eta(1760)$ ,  $f_0(1710)$ ,  $f_2(1640)$ , and  $f_2(1910)$ .
- <sup>7</sup> From a two-resonance fit.
- <sup>8</sup> Estimated by us from various fits. Systematic uncertainties not estimated.

OCCUR=2

NODE=M114W;LINKAGE=ZA  
 NODE=M114W;LINKAGE=MA  
 NODE=M114W;LINKAGE=ZH  
 NODE=M114W;LINKAGE=B

 $\eta(1760)$  DECAY MODES

NODE=M114215;NODE=M114

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $4\pi$	
$\Gamma_2$ $2\pi^+ 2\pi^-$	seen
$\Gamma_3$ $\pi^+ \pi^- 2\pi^0$	seen
$\Gamma_4$ $\rho^0 \rho^0$	seen
$\Gamma_5$ $\rho^+ \rho^-$	seen
$\Gamma_6$ $\omega\omega$	seen
$\Gamma_7$ $\eta' \pi^+ \pi^-$	seen
$\Gamma_8$ $\gamma\gamma$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=7

DESIG=8;OUR EVAL;→ UNCHECKED ←

DESIG=9;OUR EVAL;→ UNCHECKED ←

$\eta(1760) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M114225

$$\Gamma(\eta' \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$$

 $\Gamma_7\Gamma_8/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$28.2^{+7.9}_{-7.5} \pm 3.7$	465	<sup>9</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$

NODE=M114G01  
NODE=M114G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.0^{+2.0}_{-1.2} \pm 0.8$	52	<sup>10</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
$18^{+13}_{-10} \pm 5$	315	<sup>11</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$

OCCUR=2

OCCUR=3

<sup>9</sup> From a single-resonance fit.<sup>10</sup> From a two-resonance fit. For constructive interference with the X(1835).<sup>11</sup> From a two-resonance fit. For destructive interference with the X(1835).

NODE=M114G01;LINKAGE=ZH

NODE=M114G01;LINKAGE=ZA

NODE=M114G01;LINKAGE=ZN

 $\eta(1760)$  BRANCHING RATIOS

NODE=M114210

$$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$$

 $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M114R01  
NODE=M114R01

$$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^- 2\pi^0$

NODE=M114R02  
NODE=M114R02

$$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$$

 $\Gamma_4/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma \rho^0 \rho^0$
seen	BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \gamma \rho^0 \rho^0$

NODE=M114R03  
NODE=M114R03

$$\Gamma(\rho^+ \rho^-)/\Gamma_{\text{total}}$$

 $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	89B DM2	$J/\psi \rightarrow \gamma \rho^+ \rho^-$
seen	BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \gamma \rho^+ \rho^-$

NODE=M114R04  
NODE=M114R04

$$\Gamma(\omega\omega)/\Gamma_{\text{total}}$$

 $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BISELLO	87 DM2	$J/\psi \rightarrow \omega\omega$
seen	BALTRUSAIT...85C	MRK3	$J/\psi \rightarrow \gamma\omega\omega$

NODE=M114R06  
NODE=M114R06

$$\Gamma(\gamma\gamma)/\Gamma(\omega\omega)$$

 $\Gamma_8/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.48 \times 10^{-3}$	90	<sup>12</sup> ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

NODE=M114R08  
NODE=M114R08<sup>12</sup> Using results from ABLIKIM 06H.

NODE=M114R08;LINKAGE=A

 $\eta(1760)$  REFERENCES

NODE=M114

ABLIKIM	180	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
BALTRUSAIT... 86		PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BALTRUSAIT... 86B		PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT... 85C		PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095

**$f_0(1770)$** 

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M264

NODE=M264

 **$f_0(1770)$  Breit-Wigner MASS**

NODE=M264M

NODE=M264M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**1784<sup>+16</sup><sub>-14</sub> OUR AVERAGE** Error includes scale factor of 1.1.1814±31 7.2k <sup>1</sup> KHOLODENK..21 VES 29  $\pi^- p \rightarrow n\omega\phi$ 1795±7<sup>+23</sup><sub>-20</sub> ABLIKIM 13J BES3  $J/\psi \rightarrow \gamma\omega\phi$ 1760±15<sup>+15</sup><sub>-10</sub> ABLIKIM 05Q BES2  $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$ 

●●● We do not use the following data for averages, fits, limits, etc. ●●●

1765±15 SARANTSEV 21 RVUE  $J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$ 1814±18 <sup>2,3</sup> AAIJ 14BR LHCB  $\bar{B}_s^0 \rightarrow J/\psi\pi^+\pi^-$ 1812<sup>+19</sup><sub>-26</sub>±18 <sup>4</sup> ABLIKIM 06J BES2  $J/\psi \rightarrow \gamma\omega\phi$ 1790<sup>+40</sup><sub>-30</sub> ABLIKIM 05 BES2  $J/\psi \rightarrow \phi\pi^+\pi^-$ <sup>1</sup> From partial wave analysis of  $\omega\phi$  invariant mass including  $0^{++}$ ,  $2^{++}$ , and  $0^{-+}$  resonances.<sup>2</sup> Second solution:  $1800 \pm 22$  MeV. The fit favors  $f_0(1770)$  to  $f_0(1710)$ .<sup>3</sup> Statistical error only.<sup>4</sup> Not seen by LIU 09 in  $B^\pm \rightarrow K^\pm\omega\phi$ .

NODE=M264M;LINKAGE=M

NODE=M264M;LINKAGE=A

NODE=M264M;LINKAGE=B

NODE=M264M;LINKAGE=L

 **$f_0(1770)$  Breit-Wigner WIDTH**

NODE=M264W

NODE=M264W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**161±21 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.182±19 7.2k <sup>1</sup> KHOLODENK..21 VES 29  $\pi^- p \rightarrow n\omega\phi$ 95±10<sup>+78</sup><sub>-82</sub> ABLIKIM 13J BES3  $J/\psi \rightarrow \gamma\omega\phi$ 125±25<sup>+10</sup><sub>-15</sub> ABLIKIM 05Q BES2  $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$ 

●●● We do not use the following data for averages, fits, limits, etc. ●●●

180±20 SARANTSEV 21 RVUE  $J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$ 328±34 <sup>2,3</sup> AAIJ 14BR LHCB  $\bar{B}_s^0 \rightarrow J/\psi\pi^+\pi^-$ 105±20±28 <sup>4</sup> ABLIKIM 06J BES2  $J/\psi \rightarrow \gamma\omega\phi$ 270<sup>+60</sup><sub>-30</sub> <sup>5</sup> ABLIKIM 05 BES2  $J/\psi \rightarrow \phi\pi^+\pi^-$ <sup>1</sup> From partial wave analysis of  $\omega\phi$  invariant mass including  $0^{++}$ ,  $2^{++}$ , and  $0^{-+}$  resonances.<sup>2</sup> Second solution:  $263 \pm 30$  MeV. The fit favors  $f_0(1770)$  to  $f_0(1710)$ .<sup>3</sup> Statistical error only.<sup>4</sup> Not seen by LIU 09 in  $B^\pm \rightarrow K^\pm\omega\phi$ .<sup>5</sup>  $f_0(1710)$  width fixed to PDG value.


NODE=M264W;LINKAGE=J

NODE=M264W;LINKAGE=A

NODE=M264W;LINKAGE=B

NODE=M264W;LINKAGE=I

NODE=M264W;LINKAGE=AB



**f<sub>0</sub>(1770) DECAY MODES**

NODE=M264215;NODE=M264

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $K\bar{K}$	seen
$\Gamma_3$ $\eta\eta$	seen
$\Gamma_4$ $\omega\phi$	seen

DESIG=1  
DESIG=2  
DESIG=3  
DESIG=4

**$\Gamma(\pi\pi)/\Gamma_{total}$**   **$\Gamma_1/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
seen	SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
seen	AAIJ 14BR	LHCB	$B_s^0 \rightarrow J/\psi\pi^+\pi^-$
<b>seen</b>	ABLIKIM 05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$

NODE=M264R01  
NODE=M264R01

**$\Gamma(K\bar{K})/\Gamma_{total}$**   **$\Gamma_2/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M264R02  
NODE=M264R02

**$\Gamma(\eta\eta)/\Gamma_{total}$**   **$\Gamma_3/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M264R03  
NODE=M264R03

**$\Gamma(\omega\phi)/\Gamma_{total}$**   **$\Gamma_4/\Gamma$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	7.2k	KHOLODENK..21	VES	$29 \pi^- p \rightarrow n\omega\phi$
<b>seen</b>		SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M264R04  
NODE=M264R04

**f<sub>0</sub>(1770) REFERENCES**

NODE=M264

KHOLODENK...21	PAN 83 1602	M.S. Kholodenko	(VES Collab.)	REFID=61410
SARANTSEV 21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
AAIJ 14BR	PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56035
ABLIKIM 13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54955
LIU 09	PR D79 071102	C. Liu <i>et al.</i>	(BELLE Collab.)	REFID=52752
ABLIKIM 06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM 05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM 05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958



# π(1800)

$$I^G(J^{PC}) = 1^-(0^{-+})$$

See the review on "Non- $q\bar{q}$  Mesons."

NODE=M075

NODE=M075

NODE=M075M

NODE=M075M

## π(1800) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**1810<sup>+9</sup><sub>-11</sub> OUR AVERAGE** Error includes scale factor of 2.2. See the ideogram below.

1804 <sup>+6</sup> <sub>-9</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 π <sup>-</sup> p → π <sup>-</sup> π <sup>+</sup> π <sup>-</sup> p
1876 ± 18 ± 16	4k	<sup>2</sup> EUGENIO	08	B852	- 18 π <sup>-</sup> p → ηηπ <sup>-</sup> p
1774 ± 18 ± 20		<sup>3</sup> CHUNG	02	B852	18.3 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> π <sup>-</sup> p
1863 ± 9 ± 10		<sup>4</sup> CHUNG	02	B852	18.3 π <sup>-</sup> p → π <sup>+</sup> π <sup>-</sup> π <sup>-</sup> p
1840 ± 10 ± 10	1.2k	AMELIN	96B	VES	- 37 π <sup>-</sup> A → ηηπ <sup>-</sup> A
1775 ± 7 ± 10		<sup>5</sup> AMELIN	95B	VES	- 36 π <sup>-</sup> A → π <sup>+</sup> π <sup>-</sup> π <sup>-</sup> A
1790 ± 14		<sup>6</sup> BERDNIKOV	94	VES	- 37 π <sup>-</sup> A → K <sup>+</sup> K <sup>-</sup> π <sup>-</sup> A
1873 ± 33 ± 20		BELADIDZE	92C	VES	- 36 π <sup>-</sup> Be → π <sup>-</sup> η' η Be
1814 ± 10 ± 23	426	BITYUKOV	91	VES	- 36 π <sup>-</sup> C → π <sup>-</sup> ηη C
1770 ± 30	1.1k	BELLINI	82	SPEC	- 40 π <sup>-</sup> A → 3π A

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1785 ± 9 <sup>+12</sup> <sub>-6</sub>	420k	<sup>7</sup> ALEKSEEV	10	COMP	190 π <sup>-</sup> Pb → π <sup>-</sup> π <sup>-</sup> π <sup>+</sup> Pb'
1737 ± 5 ± 15		AMELIN	99	VES	37 π <sup>-</sup> A → ω π <sup>-</sup> π <sup>0</sup> A*

<sup>1</sup> Statistical error negligible.

<sup>2</sup> From a single-pole fit.

<sup>3</sup> In the  $f_0(980)\pi$  wave.


<sup>4</sup> In the  $f_0(500)\pi$  wave.

<sup>5</sup> From a fit to  $J^{PC} = 0^{-+} f_0(980)\pi, f_0(1370)\pi$  waves.

<sup>6</sup> From a fit to  $J^{PC} = 0^{-+} K_0^*(1430)K^-, f_0(980)\pi^-$  waves.

<sup>7</sup> Superseded by AGHASYAN 2018B.

NODE=M075M;LINKAGE=B  
 NODE=M075M;LINKAGE=SP  
 NODE=M075M;LINKAGE=C1  
 NODE=M075M;LINKAGE=C2  
 NODE=M075M;LINKAGE=AX  
 NODE=M075M;LINKAGE=A  
 NODE=M075M;LINKAGE=C



## π(1800) WIDTH

NODE=M075W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>215<sup>+7</sup><sub>-8</sub> OUR AVERAGE</b>					
220 <sup>+8</sup> <sub>-11</sub>	46M	<sup>8</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
221 $\pm$ 26 $\pm$ 38	4k	<sup>9</sup> EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
223 $\pm$ 48 $\pm$ 50		<sup>10</sup> CHUNG	02	B852	18.3 $\pi^- p \rightarrow$ $\pi^+ \pi^- \pi^- p$
191 $\pm$ 21 $\pm$ 20		<sup>11</sup> CHUNG	02	B852	18.3 $\pi^- p \rightarrow$ $\pi^+ \pi^- \pi^- p$
210 $\pm$ 30 $\pm$ 30	1.2k	AMELIN	96B	VES	- 37 $\pi^- A \rightarrow \eta \eta \pi^- A$
190 $\pm$ 15 $\pm$ 15		<sup>12</sup> AMELIN	95B	VES	- 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
210 $\pm$ 70		<sup>13</sup> BERDNIKOV	94	VES	- 37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
225 $\pm$ 35 $\pm$ 20		BELADIDZE	92C	VES	- 36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$
205 $\pm$ 18 $\pm$ 32	426	BITYUKOV	91	VES	- 36 $\pi^- C \rightarrow \pi^- \eta \eta C$
310 $\pm$ 50	1.1k	BELLINI	82	SPEC	- 40 $\pi^- A \rightarrow 3\pi A$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
208 $\pm$ 22 <sup>+21</sup> <sub>-37</sub>	420k	<sup>14</sup> ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow$ $\pi^- \pi^- \pi^+ Pb'$
259 $\pm$ 19 $\pm$ 6		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$

NODE=M075W

OCCUR=2

<sup>8</sup> Statistical error negligible.<sup>9</sup> From a single-pole fit.<sup>10</sup> In the  $f_0(980)\pi$  wave.<sup>11</sup> In the  $f_0(500)\pi$  wave.<sup>12</sup> From a fit to  $J^{PC} = 0^- + f_0(980)\pi, f_0(1370)\pi$  waves.<sup>13</sup> From a fit to  $J^{PC} = 0^- + K_0^*(1430)K^-$  and  $f_0(980)\pi^-$  waves.<sup>14</sup> Superseded by AGHASYAN 2018B.

NODE=M075W;LINKAGE=B  
 NODE=M075W;LINKAGE=SP  
 NODE=M075W;LINKAGE=C1  
 NODE=M075W;LINKAGE=C2  
 NODE=M075W;LINKAGE=AX  
 NODE=M075W;LINKAGE=A  
 NODE=M075W;LINKAGE=C

 **$\pi(1800)$  DECAY MODES**

NODE=M075215;NODE=M075

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi^+ \pi^- \pi^-$	seen
$\Gamma_2$ $f_0(500)\pi^-$	seen
$\Gamma_3$ $f_0(980)\pi^-$	seen
$\Gamma_4$ $f_0(1370)\pi^-$	seen
$\Gamma_5$ $f_0(1500)\pi^-$	not seen
$\Gamma_6$ $\rho\pi^-$	not seen
$\Gamma_7$ $\eta\eta\pi^-$	seen
$\Gamma_8$ $a_0(980)\eta$	seen
$\Gamma_9$ $a_2(1320)\eta$	not seen
$\Gamma_{10}$ $f_2(1270)\pi$	not seen
$\Gamma_{11}$ $f_0(1370)\pi^-$	not seen
$\Gamma_{12}$ $f_0(1500)\pi^-$	seen
$\Gamma_{13}$ $\eta\eta'(958)\pi^-$	seen
$\Gamma_{14}$ $K_0^*(1430)K^-$	seen
$\Gamma_{15}$ $K^*(892)K^-$	not seen

DESIG=10;OUR EST;→ UNCHECKED ←  
 DESIG=11;OUR EST;→ UNCHECKED ←  
 DESIG=3;OUR EST;→ UNCHECKED ←  
 DESIG=1  
 DESIG=12  
 DESIG=2  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=5;OUR EST;→ UNCHECKED ←  
 DESIG=13  
 DESIG=14  
 DESIG=15  
 DESIG=6;OUR EST;→ UNCHECKED ←  
 DESIG=8;OUR EST;→ UNCHECKED ←  
 DESIG=4  
 DESIG=9

 **$\pi(1800)$  BRANCHING RATIOS**

NODE=M075220

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma_2$
<b>0.44<math>\pm</math>0.08<math>\pm</math>0.38</b>	<sup>15</sup> CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

NODE=M075R11  
 NODE=M075R11

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_3/\Gamma_4$
<b>1.7<math>\pm</math>1.3</b>	<sup>16</sup> AMELIN	95B	VES	- 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	

NODE=M075R5  
 NODE=M075R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_4/\Gamma$
<b>seen</b>	BELLINI	82	SPEC	- 40 $\pi^- A \rightarrow 3\pi A$	

NODE=M075R1  
 NODE=M075R1

$\Gamma(f_0(1500)\pi^-)/\Gamma_{\text{total}}$						$\Gamma_5/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT			NODE=M075R12 NODE=M075R12
not seen		CHUNG	02 B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$			
$\Gamma(\rho\pi^-)/\Gamma_{\text{total}}$						$\Gamma_6/\Gamma$	
VALUE		DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R2 NODE=M075R2
not seen		BELLINI	82 SPEC	-	$40 \pi^- A \rightarrow 3\pi A$		
$\Gamma(\rho\pi^-)/\Gamma(f_0(980)\pi^-)$						$\Gamma_6/\Gamma_3$	
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R6 NODE=M075R6
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<0.25		CHUNG	02 B852		$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$		
<0.14	90	AMELIN	95B VES	-	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$		
$\Gamma(\eta\eta\pi^-)/\Gamma(\pi^+\pi^-\pi^-)$						$\Gamma_7/\Gamma_1$	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R8 NODE=M075R8
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$0.5 \pm 0.1$	1200	<sup>16</sup> AMELIN	96B VES	-	$37 \pi^- A \rightarrow \eta\eta\pi^- A$		
$\Gamma(a_2(1320)\eta)/\Gamma_{\text{total}}$						$\Gamma_9/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT			NODE=M075R13 NODE=M075R13
not seen		EUGENIO	08 B852	$18 \pi^- p \rightarrow \eta\eta\pi^- p$			
$\Gamma(f_2(1270)\pi)/\Gamma_{\text{total}}$						$\Gamma_{10}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT			NODE=M075R14 NODE=M075R14
not seen		EUGENIO	08 B852	$18 \pi^- p \rightarrow \eta\eta\pi^- p$			
$\Gamma(f_0(1370)\pi^-)/\Gamma_{\text{total}}$						$\Gamma_{11}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT			NODE=M075R15 NODE=M075R15
not seen		EUGENIO	08 B852	$18 \pi^- p \rightarrow \eta\eta\pi^- p$			
$\Gamma(f_0(1500)\pi^-)/\Gamma(a_0(980)\eta)$						$\Gamma_{12}/\Gamma_8$	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R7 NODE=M075R7
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$0.48 \pm 0.17$	4k <sup>16,17</sup>	EUGENIO	08 B852	-	$18 \pi^- p \rightarrow \eta\eta\pi^- p$		
$0.030^{+0.014}_{-0.011}$	<sup>16</sup>	ANISOVICH	01B SPEC	0	$0.6-1.94 p\bar{p} \rightarrow \eta\eta\pi^0\pi^0$		
$0.08 \pm 0.03$	1200 <sup>16,18</sup>	AMELIN	96B VES	-	$37 \pi^- A \rightarrow \eta\eta\pi^- A$		
$\Gamma(\eta\eta'(958)\pi^-)/\Gamma(\eta\eta\pi^-)$						$\Gamma_{13}/\Gamma_7$	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R10 NODE=M075R10
• • • We do not use the following data for averages, fits, limits, etc. • • •							
$0.29 \pm 0.07$	<sup>16</sup>	BELADIDZE	92C VES	-	$36 \pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$		
$0.3 \pm 0.1$	$426 \pm 57$	<sup>16</sup> BITYUKOV	91 VES	-	$36 \pi^- \text{C} \rightarrow \pi^- \eta \eta \text{C}$		
$\Gamma(K_0^*(1430)K^-)/\Gamma_{\text{total}}$						$\Gamma_{14}/\Gamma$	
VALUE		DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R4 NODE=M075R4
seen		BERDNIKOV	94 VES	-	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$		
$\Gamma(K^*(892)K^-)/\Gamma_{\text{total}}$						$\Gamma_{15}/\Gamma$	
VALUE		DOCUMENT ID	TECN	CHG	COMMENT		NODE=M075R9 NODE=M075R9
not seen		BERDNIKOV	94 VES	-	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$		
<sup>15</sup> Assuming that $f_0(980)$ decays only to $\pi\pi$ .							
<sup>16</sup> Systematic errors not estimated.							
<sup>17</sup> From a single-pole fit.							
<sup>18</sup> Assuming that $f_0(1500)$ decays only to $\eta\eta$ and $a_0(980)$ decays only to $\eta\pi$ .							

 **$\pi(1800)$  REFERENCES**

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)	REFID=52160
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>		REFID=48318
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
AMELIN	96B	Translated from YAF 62 487			
		PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL) IGJPC	REFID=44725
AMELIN	95B	Translated from YAF 59 1021			
		PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44433
AMELIN	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)	REFID=44073
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)	REFID=44073
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)	REFID=43175
BELADIDZE	92C	Translated from YAF 55 2748			
BITYUKOV	91	PL B268 137	S.I. Bityukov <i>et al.</i>	(SERP, TBIL)	REFID=41749
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)	REFID=21134

NODE=M075

**$f_2(1810)$**

$I^G(J^{PC}) = 0^+(2^{++})$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M038

NODE=M038

NODE=M038M

NODE=M038M

**$f_2(1810)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1815 ±12</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.4. See the ideogram below.		
1822 $\begin{smallmatrix} +29 \\ -24 \end{smallmatrix}$	$\begin{smallmatrix} +66 \\ -57 \end{smallmatrix}$	5.5k	1 ABLIKIM	13N BES3 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
1737 ± 9	$\begin{smallmatrix} +198 \\ -65 \end{smallmatrix}$		2 UEHARA	10A BELL $10.6 e^+e^- \rightarrow e^+e^-\eta\eta$
1800 ±30	40		ALDE	88D GAM4 $300 \pi^-p \rightarrow \pi^-p4\pi^0$
1806 ±10	1600		ALDE	87 GAM4 $100 \pi^-p \rightarrow 4\pi^0n$
1870 ±40			3 ALDE	86D GAM4 $100 \pi^-p \rightarrow \eta\eta n$
1857 $\begin{smallmatrix} +35 \\ -24 \end{smallmatrix}$			4 COSTA	80 OMEG $10 \pi^-p \rightarrow K^+K^-n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1845.0 ± 2.2 $\begin{smallmatrix} + \\ - \end{smallmatrix}$	$\begin{smallmatrix} 1.6 \\ 7.2 \end{smallmatrix}$		5 ALBRECHT	20 RVUE $0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
1858 $\begin{smallmatrix} +18 \\ -71 \end{smallmatrix}$			6 LONGACRE	86 RVUE Compilation
1799 ±15			7 CASON	82 STRC $8 \pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>2</sup> Breit-Wigner mass. Could also be the  $f_2(1910)$ .

<sup>3</sup> Seen in only one solution.

<sup>4</sup> Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

<sup>5</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).

<sup>6</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

<sup>7</sup> From an amplitude analysis of the reaction  $\pi^+\pi^- \rightarrow 2\pi^0$ . The resonance in the  $2\pi^0$  final state is not confirmed by PROKOSHKIN 97.

NODE=M038M;LINKAGE=B

NODE=M038M;LINKAGE=UE


NODE=M038M;LINKAGE=F

NODE=M038M;LINKAGE=A

NODE=M038M;LINKAGE=C

NODE=M038M;LINKAGE=L

NODE=M038M;LINKAGE=P1



**$f_2(1810)$  WIDTH**

NODE=M038W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>197 ± 22</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.5. See the ideogram below.		
229 +52 -42	+88 -155	5.5k	1 ABLIKIM 13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
228 +21 -20	+234 -153		2 UEHARA 10A BELL	$10.6 e^+e^- \rightarrow e^+e^-\eta\eta$
160 ± 30	40	ALDE 88D	GAM4	$300 \pi^-p \rightarrow \pi^-p4\pi^0$
190 ± 20	1600	ALDE 87	GAM4	$100 \pi^-p \rightarrow 4\pi^0n$
250 ± 30		3 ALDE 86D	GAM4	$100 \pi^-p \rightarrow \eta\eta n$
185 +102 -139		4 COSTA 80	OMEG	$10 \pi^-p \rightarrow K^+K^-n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

260.9 ± 3.9	+199.9 -38.2	5 ALBRECHT 20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
388 +15 -21		6 LONGACRE 86	RVUE	Compilation
280 +42 -35		7 CASON 82	STRC	$8 \pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$

- 1 From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.
- 2 Breit-Wigner width. Could also be the  $f_2(1910)$ .
- 3 Seen in only one solution.
- 4 Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.
- 5 T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).
- 6 From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.
- 7 From an amplitude analysis of the reaction  $\pi^+\pi^- \rightarrow 2\pi^0$ . The resonance in the  $2\pi^0$  final state is not confirmed by PROKOSHKIN 97.


NODE=M038W

NODE=M038W;LINKAGE=B

NODE=M038W;LINKAGE=UE  
 NODE=M038W;LINKAGE=F  
 NODE=M038W;LINKAGE=A  
 NODE=M038W;LINKAGE=C

NODE=M038W;LINKAGE=L

NODE=M038W;LINKAGE=P1



### $f_2(1810)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	seen
$\Gamma_2$ $\eta\eta$	seen
$\Gamma_3$ $4\pi^0$	seen
$\Gamma_4$ $K^+K^-$	seen
$\Gamma_5$ $\gamma\gamma$	seen

NODE=M038215;NODE=M038

DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=3  
 DESIG=4;OUR EST;→ UNCHECKED ←  
 DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=5;OUR EST;→ UNCHECKED ←

$f_2(1810) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M038225

 $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_2\Gamma_5/\Gamma$ 

VALUE (eV)

DOCUMENT ID TECN COMMENT

 $5.2^{+0.9+37.3}_{-0.8-4.5}$ <sup>1</sup> UEHARA 10A BELL 10.6  $e^+e^- \rightarrow e^+e^-\eta\eta$ NODE=M038G01  
NODE=M038G01<sup>1</sup> Including interference with the  $f_2'(1525)$  (parameters fixed to the values from the 2008 edition of this review, PDG 08) and  $f_2(1270)$ . May also be the  $f_0(1500)$ .

NODE=M038G01;LINKAGE=UE

 $f_2(1810)$  BRANCHING RATIOS

NODE=M038220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

AMSLER 02 CBAR 0.9  $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$ 

not seen

PROKOSHKIN 97 GAM2 38  $\pi^-p \rightarrow \pi^0\pi^0n$  $0.21^{+0.02}_{-0.03}$ <sup>1</sup> LONGACRE 86 RVUE Compilation $0.44 \pm 0.03$ <sup>2</sup> CASON 82 STRC 8  $\pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$ NODE=M038R2  
NODE=M038R2<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R2;LINKAGE=L

<sup>2</sup> Included in LONGACRE 86 global analysis.

NODE=M038R;LINKAGE=C

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE

DOCUMENT ID TECN COMMENT

seen

ABLIKIM 13N BES3 PWA of  $J/\psi \rightarrow \gamma\eta\eta$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.008^{+0.028}_{-0.003}$ <sup>1</sup> LONGACRE 86 RVUE CompilationNODE=M038R3  
NODE=M038R3<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R3;LINKAGE=L

 $\Gamma(\pi\pi)/\Gamma(4\pi^0)$  $\Gamma_1/\Gamma_3$ 

VALUE

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt;0.75

ALDE 87 GAM4 100  $\pi^-p \rightarrow 4\pi^0n$ NODE=M038R4  
NODE=M038R4 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$  $\Gamma_3/\Gamma_2$ 

VALUE

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.8 \pm 0.3$ ALDE 87 GAM4 100  $\pi^-p \rightarrow 4\pi^0n$ NODE=M038R5  
NODE=M038R5 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $0.003^{+0.019}_{-0.002}$ <sup>1</sup> LONGACRE 86 RVUE Compilation

seen

COSTA 80 OMEG 10  $\pi^-p \rightarrow K^+K^-n$ NODE=M038R1  
NODE=M038R1<sup>1</sup> From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R1;LINKAGE=L

 $f_2(1810)$  REFERENCES

NODE=M038

ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
PROKOSHKIN	97	PD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)	REFID=45386
ALDE	88D	Translated from DANS 353 323. SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=44652
ALDE	87	Translated from YAF 47 1273. PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355

**X(1835)**

$$I^G(J^{PC}) = ?^?(0^{-+})$$

OMITTED FROM SUMMARY TABLE

Could be a superposition of two states, one with small width appearing as threshold enhancement in  $p\bar{p}$ , the other one with a larger width. For the former ABLIKIM 12D determine  $J^{PC} = 0^{-+}$ .

NODE=M085

NODE=M085

**X(1835) MASS**

NODE=M085M

NODE=M085M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1826.5<sup>+13.0</sup><sub>-3.4</sub> OUR AVERAGE</b>				
1825.3 ± 2.4 <sup>+17.3</sup> <sub>-2.4</sub>		<sup>1</sup> ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
1844 ± 9 <sup>+16</sup> <sub>-25</sub>		ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1839 ± 26 ± 26		<sup>2</sup> ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
1909.5 ± 15.9 <sup>+9.4</sup> <sub>-27.5</sub>		<sup>3</sup> ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
1842.2 ± 4.2 <sup>+7.1</sup> <sub>-2.6</sub>	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
1832 <sup>+19</sup> <sub>-5</sub> ± 26		<sup>4</sup> ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
1836.5 ± 3.0 <sup>+5.6</sup> <sub>-2.1</sub>	4265	<sup>5</sup> ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
1877.3 ± 6.3 <sup>+3.4</sup> <sub>-7.4</sub>		<sup>6</sup> ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
1837 <sup>+10</sup> <sub>-12</sub> ± 9 <sup>+9</sup> <sub>-7</sub>	231	<sup>7,8</sup> ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
1833.7 ± 6.1 ± 2.7	264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
1831 ± 7		<sup>8,9</sup> ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$
1859 <sup>+3</sup> <sub>-10</sub> ± 5 <sup>+5</sup> <sub>-25</sub>		<sup>8</sup> BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

OCCUR=2

OCCUR=2

<sup>1</sup> From a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors.

NODE=M085M;LINKAGE=B

<sup>2</sup> From a fit to  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded.

NODE=M085M;LINKAGE=C

<sup>3</sup> Pole mass from a fit of the measured  $\pi^+\pi^-\eta'$  lineshape to a Flatte formula that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold; the fit also includes known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit.

NODE=M085M;LINKAGE=A

<sup>4</sup> From the fit including final state interaction effects in isospin 0  $S$ -wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.

NODE=M085M;LINKAGE=AK

<sup>5</sup> From a fit of the  $\pi^+\pi^-\eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two states  $\gamma X(2120)$  and  $\gamma X(2370)$ , for  $M(\pi^+\pi^-\eta') < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ .

NODE=M085M;LINKAGE=AI

<sup>6</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$ . This state may be due also to  $\eta_2(1870)$  or to a combination of  $X(1835)$  and  $\eta_2(1870)$ .

NODE=M085M;LINKAGE=BL

<sup>7</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV.

NODE=M085M;LINKAGE=AE

<sup>8</sup> Evidence for a threshold enhancement in the  $p\bar{p}$  mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in  $B^+ \rightarrow p\bar{p}K^+$ , WANG 05A in  $B^0 \rightarrow p\bar{p}K_S^0$ , ABE 02W in  $\bar{B}^0 \rightarrow p\bar{p}D^0$ , DEL-AMO-SANCHEZ 12 in  $B \rightarrow D(D^*)p\bar{p}(\pi)$ , and WEI 08 in  $B^+ \rightarrow p\bar{p}\pi^+$  decays. Not seen by ATHAR 06 in  $\Upsilon(1S) \rightarrow p\bar{p}\gamma$ .

NODE=M085M;LINKAGE=HF

<sup>9</sup> From the fit including final state interaction effects in isospin 0  $S$ -wave according to SIBIRTSEV 05A. Systematic errors not estimated.

NODE=M085M;LINKAGE=AB

**X(1835) WIDTH**

NODE=M085W

NODE=M085W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>242<sup>+14</sup><sub>-15</sub> OUR AVERAGE</b>					
245.2 ± 13.1 <sup>+4.6</sup> <sub>-9.6</sub>			<sup>1</sup> ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
192 <sup>+20</sup> <sub>-17</sub> ± 62 <sup>+62</sup> <sub>-43</sub>			ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

175 ±57 ±25		<sup>2</sup> ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
273.5±21.4 <sup>+6.1</sup> <sub>-64.0</sub>		<sup>3</sup> ABLIKIM	16J	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
83 ±14 ±11	0.6k	ABLIKIM	13U	BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
< 76	90	<sup>4</sup> ABLIKIM	12D	BES3	$J/\psi \rightarrow \gamma\rho\bar{\rho}$
190 ± 9 <sup>+38</sup> <sub>-36</sub>	4265	<sup>5</sup> ABLIKIM	11C	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
57 ±12 <sup>+19</sup> <sub>-4</sub>		<sup>6</sup> ABLIKIM	11J	BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
0 <sup>+44</sup> <sub>-0</sub>	231	<sup>7,8</sup> ALEXANDER	10	CLEO	$J/\psi \rightarrow \gamma\rho\bar{\rho}$
67.7±20.3± 7.7	264	ABLIKIM	05R	BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
< 153	90	<sup>8,9</sup> ABLIKIM	05R	BES2	$J/\psi \rightarrow \gamma\rho\bar{\rho}$
< 30		<sup>8</sup> BAI	03F	BES2	$J/\psi \rightarrow \gamma\rho\bar{\rho}$

OCCUR=2

<sup>1</sup> From a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $\rho\bar{\rho}$  threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors.

NODE=M085W;LINKAGE=B

<sup>2</sup> From a fit to  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded.

NODE=M085W;LINKAGE=C

<sup>3</sup> Pole width from a fit of the measured  $\pi^+\pi^-\eta'$  lineshape to a Flatte formula that accounts for the abrupt distortion observed at the  $\rho\bar{\rho}$  threshold; the fit also includes known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit.

NODE=M085W;LINKAGE=A

<sup>4</sup> From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G.

NODE=M085W;LINKAGE=AK

<sup>5</sup> From a fit of the  $\pi^+\pi^-\eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two states  $\gamma X(2120)$  and  $\gamma X(2370)$ , for  $M(\pi^+\pi^-\eta') < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$ .

NODE=M085W;LINKAGE=AI

<sup>6</sup> The selected process is  $J/\psi \rightarrow \omega a_0(980)\pi$ . This state may be due also to  $\eta_2(1870)$  or to a combination of  $X(1835)$  and  $\eta_2(1870)$ .

NODE=M085W;LINKAGE=BL

<sup>7</sup> From a fit of the  $\rho\bar{\rho}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma\rho\bar{\rho}$  phase space, for  $M(\rho\bar{\rho}) < 2.85$  GeV.

NODE=M085W;LINKAGE=AE

<sup>8</sup> Evidence for a threshold enhancement in the  $\rho\bar{\rho}$  mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in  $B^+ \rightarrow \rho\bar{\rho}K^+$ , WANG 05A in  $B^0 \rightarrow \rho\bar{\rho}K_S^0$ , ABE 02W in  $\bar{B}^0 \rightarrow \rho\bar{\rho}D^0$ , DEL-AMO-SANCHEZ 12 in  $B \rightarrow D(D^*)\rho\bar{\rho}(\pi)$ , and WEI 08 in  $B^+ \rightarrow \rho\bar{\rho}\pi^+$  decays. Not seen by ATHAR 06 in  $\Upsilon(1S) \rightarrow \rho\bar{\rho}\gamma$ .

NODE=M085W;LINKAGE=HF

<sup>9</sup> From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Systematic errors not estimated.

NODE=M085W;LINKAGE=AB

### X(1835) DECAY MODES

NODE=M085215;NODE=M085

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\bar{\rho}$	seen
$\Gamma_2$ $\eta'\pi^+\pi^-$	seen
$\Gamma_3$ $\gamma\gamma$	
$\Gamma_4$ $K_S^0 K_S^0 \eta$	seen
$\Gamma_5$ $\gamma\phi(1020)$	possibly seen
$\Gamma_6$ $3(\pi^+\pi^-)$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2

DESIG=4

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6

DESIG=7

### X(1835) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M085225

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_3/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M085G01

NODE=M085G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

<35.6	90	<sup>1</sup> ZHANG	12A	BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
<83	90	<sup>2</sup> ZHANG	12A	BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

OCCUR=2

<sup>1</sup> From a two-resonance fit and constructive interference of the  $\eta(1760)$  and  $X(1835)$ , a significance of  $2.8\sigma$ .

NODE=M085G01;LINKAGE=ZH

<sup>2</sup> From a two-resonance fit and destructive interference of the  $\eta(1760)$  and  $X(1835)$ , a significance of  $2.8\sigma$ .

NODE=M085G01;LINKAGE=ZA



**X(1835) BRANCHING RATIOS**

$$\Gamma(p\bar{p})/\Gamma(\eta'\pi^+\pi^-)$$

$$\Gamma_1/\Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.333	ABLIKIM	05R	BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
-------	---------	-----	---

$$\Gamma(\eta'\pi^+\pi^-)/\Gamma(K_S^0 K_S^0 \eta)$$

$$\Gamma_2/\Gamma_4$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.7 \pm 1.8$	<sup>1</sup> ABLIKIM	15T	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
---------------	----------------------	-----	---

<sup>1</sup> Using results from ABLIKIM 05R.

$$\Gamma(\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>seen</b>	<sup>1</sup> ABLIKIM	16J	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
-------------	----------------------	-----	---

<sup>1</sup> ABLIKIM 16J quotes  $B(J/\psi \rightarrow \gamma X(1835)) \times B(X(1835) \rightarrow \pi^+\pi^-\eta') = (3.93 \pm 0.38^{+0.31}_{-0.84}) \times 10^{-4}$  from a fit of the measured  $\pi^+\pi^-\eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the  $X(1835)$ ; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

$$\Gamma(\gamma\phi(1020))/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

<b>possibly seen</b>	<sup>1</sup> ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
----------------------	----------------------	-----	--

<sup>1</sup> Seen as a peak in  $\gamma\phi$  invariant mass. Angular analysis consistent with  $J^{PC} = 0^{-+}$ . Other  $J^{PC}$  not excluded.

$$\Gamma(\gamma\gamma)/\Gamma(\eta'\pi^+\pi^-)$$

$$\Gamma_3/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 9.80 \times 10^{-3}$	90	<sup>1</sup> ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
-------------------------	----	----------------------	-----	--

<sup>1</sup> Using results from ABLIKIM 16J.

$$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

<b>seen</b>	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
-------------	------	---------	-----	--

**X(1835) REFERENCES**

ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57454
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56785
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.) JPC	REFID=54269
DEL-AMO-SA...	12	PR D85 092017	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=54286
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53931
ABLIKIM	10G	CP C34 421	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55685
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)	REFID=52086
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50993
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT,B	05L	PR D72 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50827
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer		REFID=51038
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)	REFID=50651
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48690
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48980

NODE=M085220

NODE=M085R01  
NODE=M085R01NODE=M085R00  
NODE=M085R00

NODE=M085R00;LINKAGE=A

NODE=M085R03  
NODE=M085R03

NODE=M085R03;LINKAGE=A

NODE=M085R04  
NODE=M085R04

NODE=M085R04;LINKAGE=A

NODE=M085R05  
NODE=M085R05

NODE=M085R05;LINKAGE=A

NODE=M085R06  
NODE=M085R06

NODE=M085

$\phi_3(1850)$ 

$$I^G(J^{PC}) = 0^-(3^{--})$$

NODE=M054

 $\phi_3(1850)$  MASS

NODE=M054M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1854 ± 7 OUR AVERAGE</b>				
1855 ± 10		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$ , $K_S^0 K^\pm \pi^\mp \Lambda$
1870 <sup>+30</sup> <sub>-20</sub>	430	ARMSTRONG	82 OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
1850 ± 10	123	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \Lambda$

NODE=M054M

 $\phi_3(1850)$  WIDTH

NODE=M054W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>87<sup>+28</sup><sub>-23</sub> OUR AVERAGE</b> Error includes scale factor of 1.2.				
64 ± 31		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$ , $K_S^0 K^\pm \pi^\mp \Lambda$
160 <sup>+90</sup> <sub>-50</sub>	430	ARMSTRONG	82 OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
80 <sup>+40</sup> <sub>-30</sub>	123	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \Lambda$

NODE=M054W

 $\phi_3(1850)$  DECAY MODES

NODE=M054215;NODE=M054

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \bar{K}$	seen
$\Gamma_2$ $K \bar{K}^*(892) + c.c.$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

 $\phi_3(1850)$  BRANCHING RATIOS

NODE=M054220

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_1$
<b><math>\Gamma(K \bar{K}^*(892) + c.c.)/\Gamma(K \bar{K})</math></b>				
<b>0.55<sup>+0.85</sup><sub>-0.45</sub></b>	ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$ , $K_S^0 K^\pm \pi^\mp \Lambda$	
0.8 ± 0.4	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \pi \Lambda$	

NODE=M054R1  
NODE=M054R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\phi_3(1850)$  REFERENCES

NODE=M054

ASTON	88E	PL B208 324	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) IGJPC
ARMSTRONG	82	PL 110B 77	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+) JP
ALHARRAN	81B	PL 101B 357	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)

REFID=40577  
REFID=21405  
REFID=21702

**$\eta_1(1855)$** 

$$I^G(J^{PC}) = 0^+(1^-+)$$

NODE=M267

## OMITTED FROM SUMMARY TABLE

Meson with exotic (non- $q\bar{q}$ ) quantum numbers. A state decaying into  $\eta\eta'$  with possible quantum numbers  $1^-+$  was reported earlier in this mass region BARBERIS 00A in high energy central  $pp$  production and by ALDE 91B in  $\pi^-p$  interactions, see the  $f_2(1910)$ , and the review on "Spectroscopy of Light Meson Resonances."

NODE=M267

 **$\eta_1(1855)$  MASS**

NODE=M267M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$1855 \pm 9_{-1}^{+6}$	<sup>1</sup> ABLIKIM	22A1	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$

NODE=M267M

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and the resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$   $P$ -wave. For analysis details see ABLIKIM 22AS.

NODE=M267M;LINKAGE=A

 **$\eta_1(1855)$  WIDTH**

NODE=M267W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$188 \pm 18_{-8}^{+3}$	<sup>1</sup> ABLIKIM	22A1	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$

NODE=M267W

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and the resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$   $P$ -wave. For analysis details see ABLIKIM 22AS.

NODE=M267W;LINKAGE=A

 **$\eta_1(1855)$  DECAY MODES**

NODE=M267215;NODE=M267

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \eta\eta'$	seen

DESIG=1

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
seen	ABLIKIM	22A1	BES3 $J/\psi \rightarrow \gamma\eta\eta'$	
seen	BARBERIS	00A	450 $pp \rightarrow p_f \eta\eta' p_s$	
seen	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$	

NODE=M267R00  
NODE=M267R00 **$\eta_1(1855)$  REFERENCES**

NODE=M267

ABLIKIM	22A1	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61881
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
ALDE	91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41844
Translated from YAF 54 751.					

$\eta_2(1870)$ 

$$I^G(J^{PC}) = 0^+(2^-+)$$

NODE=M101

 $\eta_2(1870)$  MASS

NODE=M101M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1842 ± 8 OUR AVERAGE</b>				
1835 ± 12		BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1844 ± 13		BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
1840 ± 25		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1875 ± 20 ± 35		ADOMEIT	96 CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
1881 ± 32 ± 40	26	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1860 ± 5 ± 15		ANISOVICH	00E SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
1840 ± 15		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M101M

 $\eta_2(1870)$  WIDTH

NODE=M101W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>225 ± 14 OUR AVERAGE</b>				
235 ± 22		BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
228 ± 23		BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
200 ± 40		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
200 ± 25 ± 45		ADOMEIT	96 CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
221 ± 92 ± 44	26	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
250 ± 25 $^{+50}_{-35}$		ANISOVICH	00E SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
170 ± 40		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M101W

 $\eta_2(1870)$  DECAY MODES

NODE=M101225;NODE=M101

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta \pi \pi$	seen
$\Gamma_2$ $a_2(1320) \pi$	seen
$\Gamma_3$ $f_2(1270) \eta$	seen
$\Gamma_4$ $a_0(980) \pi$	seen
$\Gamma_5$ $\gamma \gamma$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=4;OUR EVAL;→ UNCHECKED ←  
DESIG=8;OUR EVAL;→ UNCHECKED ←  
DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=9

 $\eta_2(1870)$  BRANCHING RATIOS

NODE=M101230

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_3$
<b>1.7 ± 0.4 OUR AVERAGE</b>				
1.60 ± 0.40	<sup>1</sup> ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$
20.4 ± 6.6	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	
4.1 ± 2.3	ADOMEIT	96	CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
<sup>1</sup> Reanalysis of ADOMEIT 96 and ANISOVICH 00E.				

NODE=M101R2  
NODE=M101R2

NODE=M101R2;LINKAGE=AN

VALUE	DOCUMENT ID	COMMENT	$\Gamma_2/\Gamma_4$
<b>32.6 ± 12.6</b>			
	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$

NODE=M101R4  
NODE=M101R4

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma_3$
<b>0.48 ± 0.45</b>				
	<sup>1</sup> ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$

NODE=M101R01  
NODE=M101R01

NODE=M101R01;LINKAGE=AN

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma$
<b>seen</b>				
	KARCH	92	CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$

NODE=M101R02  
NODE=M101R02

**$\eta_2(1870)$  REFERENCES**

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)

NODE=M101

REFID=53631  
REFID=47945  
REFID=47958  
REFID=47959  
REFID=46606  
REFID=45758  
REFID=45202  
REFID=42170

**$\pi_2(1880)$**

$$I^G(J^{PC}) = 1^-(2^-+)$$

NODE=M185

**$\pi_2(1880)$  MASS**

NODE=M185M

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M185M

**1874<sup>+26</sup><sub>-5</sub> OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.


1847 <sup>+20</sup> <sub>-3</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1929 $\pm$ 24 $\pm$ 18	4k	EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
1876 $\pm$ 11 $\pm$ 67	145k	LU	05	B852	- 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
2003 $\pm$ 88 $\pm$ 148	69k	KUHN	04	B852	- 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1880 $\pm$ 20		ANISOVICH	01B	SPEC	0 0.6-1.94 $\bar{p} p \rightarrow \eta \eta \pi^0 \pi^0$
---------------	--	-----------	-----	------	--

NODE=M185M;LINKAGE=A

<sup>1</sup>Statistical error negligible.



**$\pi_2(1880)$  WIDTH**

NODE=M185W

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M185W

**237<sup>+33</sup><sub>-30</sub> OUR AVERAGE** Error includes scale factor of 1.2.

246 <sup>+33</sup> <sub>-28</sub>	46M	<sup>2</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
323 $\pm$ 87 $\pm$ 43	4k	EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
146 $\pm$ 17 $\pm$ 62	145k	LU	05	B852	- 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
306 $\pm$ 132 $\pm$ 121	69k	KUHN	04	B852	- 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

255 $\pm$ 45		ANISOVICH	01B	SPEC	0 0.6-1.94 $\bar{p} p \rightarrow \eta \eta \pi^0 \pi^0$
--------------	--	-----------	-----	------	--

<sup>2</sup>Statistical error negligible.

NODE=M185W;LINKAGE=A

$\pi_2(1880)$  DECAY MODES

NODE=M185215;NODE=M185

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta\eta\pi^-$	seen
$\Gamma_2$ $a_0(980)\eta$	seen
$\Gamma_3$ $a_2(1320)\eta$	seen
$\Gamma_4$ $f_0(1500)\pi$	seen
$\Gamma_5$ $f_1(1285)\pi$	seen
$\Gamma_6$ $\omega\pi^-\pi^0$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=3;OUR EVAL;→ UNCHECKED ←  
DESIG=4;OUR EVAL;→ UNCHECKED ←  
DESIG=5;OUR EVAL;→ UNCHECKED ←  
DESIG=6;OUR EVAL;→ UNCHECKED ←

 $\Gamma(a_2(1320)\eta)/\Gamma(f_1(1285)\pi)$   $\Gamma_3/\Gamma_5$ 

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M185R01  
NODE=M185R01

••• We do not use the following data for averages, fits, limits, etc. •••

22.7±7.3	69k	KUHN	04	B852	-	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
----------	-----	------	----	------	---	--

 $\Gamma(f_0(1500)\pi)/\Gamma(a_0(980)\eta)$   $\Gamma_4/\Gamma_2$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

NODE=M185R02  
NODE=M185R02

••• We do not use the following data for averages, fits, limits, etc. •••

0.28 <sup>+0.20</sup> <sub>-0.15</sub>	<sup>3</sup> ANISOVICH	01B	SPEC	0	0.6–1.94 $\bar{p}p \rightarrow \eta\eta\pi^0\pi^0$
--	------------------------	-----	------	---	--

<sup>3</sup>Systematic errors not estimated.

NODE=M185R02;LINKAGE=NS

 $\pi_2(1880)$  REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>	

NODE=M185

REFID=59471  
REFID=52160  
REFID=50459  
REFID=49773  
REFID=48318

$\rho(1900)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M170

NODE=M170

 $\rho(1900)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M170M

NODE=M170M

••• We do not use the following data for averages, fits, limits, etc. •••

1880±10		<sup>1</sup> ABLIKIM	22L	BES3	2.0–3.08 $e^+e^- \rightarrow K^+K^-\pi^0$
1909±17±25	54	<sup>2</sup> AUBERT	08S	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
1880±30		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
1860±20		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
1910±10		<sup>3,4</sup> FRABETTI	04	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
1870±10		ANTONELLI	96	SPEC	$e^+e^- \rightarrow$ hadrons

OCCUR=2

<sup>1</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

<sup>2</sup> From the fit with two resonances.

<sup>3</sup> From a fit with two resonances with the JACOB 72 continuum.

<sup>4</sup> Supersedes FRABETTI 01.

NODE=M170M;LINKAGE=A

NODE=M170M;LINKAGE=AU  
NODE=M170M;LINKAGE=PI  
NODE=M170M;LINKAGE=RS

 $\rho(1900)$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M170W

NODE=M170W

••• We do not use the following data for averages, fits, limits, etc. •••

69±15		<sup>1</sup> ABLIKIM	22L	BES3	2.0–3.08 $e^+e^- \rightarrow K^+K^-\pi^0$
48±17±2	54	<sup>2</sup> AUBERT	08S	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
130±30		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
160±20		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
37±13		<sup>3,4</sup> FRABETTI	04	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
10±5		ANTONELLI	96	SPEC	$e^+e^- \rightarrow$ hadrons

OCCUR=2

<sup>1</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

<sup>2</sup> From the fit with two resonances.

<sup>3</sup> From a fit with two resonances with the JACOB 72 continuum.

<sup>4</sup> Supersedes FRABETTI 01.

NODE=M170W;LINKAGE=A

NODE=M170W;LINKAGE=AU

NODE=M170W;LINKAGE=PI

NODE=M170W;LINKAGE=RS

### $\rho(1900) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

$\Gamma(\phi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_6/\Gamma$

VALUE (units  $10^{-8}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.2 \pm 1.2 \pm 0.8$     54    <sup>1</sup> AUBERT    08S    BABR     $10.6 e^+e^- \rightarrow \phi\pi^0\gamma$

<sup>1</sup> From the fit with two resonances.

NODE=M170215

NODE=M170B01

NODE=M170B01

NODE=M170B01;LINKAGE=AU

### $\rho(1900)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $6\pi$	seen
$\Gamma_2$ $3\pi^+3\pi^-$	seen
$\Gamma_3$ $2\pi^+2\pi^-\pi^0$	
$\Gamma_4$ $\phi\pi$	seen
$\Gamma_5$ hadrons	seen
$\Gamma_6$ $e^+e^-$	seen
$\Gamma_7$ $\bar{N}N$	not seen

NODE=M170225;NODE=M170

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=6

DESIG=7;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

### $\rho(1900)$ BRANCHING RATIOS

$\Gamma(6\pi)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

VALUE    EVTS    DOCUMENT ID    TECN    COMMENT

seen	8k	AKHMETSHIN 13	CMD3	$e^+e^- \rightarrow 3\pi^+3\pi^-$
not seen		AGNELLO 02	OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-\pi^0$
seen		FRABETTI 01	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
seen		ANTONELLI 96	SPEC	$e^+e^- \rightarrow \text{hadrons}$

NODE=M170230

NODE=M170R1

NODE=M170R1

### $\rho(1900)$ REFERENCES

ABLIKIM 22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AKHMETSHIN 13	PL B723 82	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
AUBERT 08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
FRABETTI 04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AGNELLO 02	PL B527 39	M. Agnello <i>et al.</i>	(OBELIX Collab.)
FRABETTI 01	PL B514 240	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANTONELLI 96	PL B365 427	A. Antonelli <i>et al.</i>	(FENICE Collab.)
JACOB 72	PR D5 1847	M. Jacob, R. Slansky	

NODE=M170

REFID=61649

REFID=55370

REFID=52242

REFID=51047

REFID=49614

REFID=48576

REFID=48350

REFID=44633

REFID=49668

**$f_2(1910)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

We list here three different peaks with close masses and widths seen in the mass distributions of  $\omega\omega$ ,  $\eta\eta'$ , and  $K^+K^-$  final states. ALDE 91B argues that they are of different nature.

NODE=M142

NODE=M142

 **$f_2(1910)$  MASS**

NODE=M142205


NODE=M142MX

 **$f_2(1910)$   $\omega\omega$  MODE**NODE=M142M2  
NODE=M142M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1900 ± 9 OUR AVERAGE</b>	Error includes scale factor of 1.4. See the ideogram below.		
1890 ± 10	<sup>1</sup> AMELIN	06	VES 36 $\pi^- p \rightarrow \omega\omega n$
1897 ± 11	BARBERIS	00F	450 $p p \rightarrow p_f \omega \omega p_S$
1924 ± 14	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$

<sup>1</sup> Supersedes BELADIDZE 92B.

NODE=M142M2;LINKAGE=AM

 **$f_2(1910)$   $\eta\eta'$  MODE**NODE=M142M3  
NODE=M142M3

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1934 ± 16</b>	<sup>1</sup> BARBERIS	00A	450 $p p \rightarrow p_f \eta \eta' p_S$
1934 ± 20	<sup>2</sup> ANISOVICH	00J	SPEC
1911 ± 10	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<sup>1</sup> Also compatible with  $J^{PC} = 1^-+$ .<sup>2</sup> Combined fit with  $\eta\eta$ ,  $\pi\pi$ , and  $\eta\pi\pi$ .NODE=M142M3;LINKAGE=KS  
NODE=M142M3;LINKAGE=AN **$f_2(1910)$   $K^+K^-$  MODE**NODE=M142M4  
NODE=M142M4

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1941 ± 18	<sup>1</sup> AMSLER	06	CBAR 1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$

<sup>1</sup> Tentative, could be  $f_2(1950)$ .

NODE=M142M4;LINKAGE=A



**$f_2(1910)$  WIDTH**

NODE=M142210


NODE=M142WX

NODE=M142W2  
NODE=M142W2 **$f_2(1910)$   $\omega\omega$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>167±21 OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.		
165±19	<sup>1</sup> AMELIN	06 VES	36 $\pi^- p \rightarrow \omega\omega n$
202±32	BARBERIS	00F	450 $pp \rightarrow p_f\omega\omega p_S$
91±50	ALDE	90 GAM2	38 $\pi^- p \rightarrow \omega\omega n$

<sup>1</sup> Supersedes BELADIDZE 92B.

NODE=M142W2;LINKAGE=AM

 **$f_2(1910)$   $\eta\eta'$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>141±41</b>	<sup>1</sup> BARBERIS	00A	450 $pp \rightarrow p_f\eta\eta' p_S$
271±25	<sup>2</sup> ANISOVICH	00J SPEC	
90±35	ALDE	91B GAM2	38 $\pi^- p \rightarrow \eta\eta' n$

<sup>1</sup> Also compatible with  $J^{PC}=1^-+$ .<sup>2</sup> Combined fit with  $\eta\eta$ ,  $\pi\pi$ , and  $\eta\pi\pi$ .NODE=M142W3  
NODE=M142W3NODE=M142W3;LINKAGE=KS  
NODE=M142W3;LINKAGE=AN **$f_2(1910)$   $K^+K^-$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120±40	AMSLER	06 CBAR	1.64 $\bar{p}p \rightarrow K^+K^-\pi^0$

NODE=M142W4  
NODE=M142W4 **$f_2(1910)$  DECAY MODES**

NODE=M142215;NODE=M142

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi^0\pi^0$	
$\Gamma_2$ $K^+K^-$	seen
$\Gamma_3$ $K_S^0 K_S^0$	
$\Gamma_4$ $\eta\eta$	seen
$\Gamma_5$ $\omega\omega$	seen
$\Gamma_6$ $\eta\eta'$	seen
$\Gamma_7$ $\eta'\eta'$	
$\Gamma_8$ $\rho\rho$	seen
$\Gamma_9$ $a_2(1320)\pi$	seen
$\Gamma_{10}$ $f_2(1270)\eta$	seen

DESIG=6

DESIG=11

DESIG=8

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=12;OUR EST;→ UNCHECKED ←

DESIG=13;OUR EST;→ UNCHECKED ←

$f_2(1910)$  BRANCHING RATIOS $\Gamma(K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+K^-\pi^0$

<sup>1</sup> Tentative, could be  $f_2(1950)$ .

NODE=M142225

NODE=M142R11  
NODE=M142R11

NODE=M142R11;LINKAGE=A

 $\Gamma(\pi^0\pi^0)/\Gamma(\eta\eta')$  $\Gamma_1/\Gamma_6$ 

VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
<0.1	ALDE	89	GAM2 $38\pi^-p \rightarrow \eta\eta'n$

NODE=M142R4  
NODE=M142R4 $\Gamma(K_S^0K_S^0)/\Gamma(\eta\eta')$  $\Gamma_3/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.066	90	BALOSHIN	86	SPEC $40\pi p \rightarrow K_S^0K_S^0n$

NODE=M142R7  
NODE=M142R7 $\Gamma(\eta\eta)/\Gamma(\eta\eta')$  $\Gamma_4/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.05	90	ALDE	91B	GAM2 $38\pi^-p \rightarrow \eta\eta'n$

NODE=M142R6  
NODE=M142R6 $\Gamma(\omega\omega)/\Gamma(\eta\eta')$  $\Gamma_5/\Gamma_6$ 

VALUE	DOCUMENT ID	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••		
$2.6 \pm 0.6$	BARBERIS 00F	$450 pp \rightarrow p_f\omega p_S$

NODE=M142R10  
NODE=M142R10 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
probably not seen	BARBERIS 00A		$450 pp \rightarrow p_f\eta'\eta'p_S$
possibly seen	BELADIDZE 92D	VES	$37\pi^-p \rightarrow \eta'\eta'n$

NODE=M142R8  
NODE=M142R8 $\Gamma(\rho\rho)/\Gamma(\omega\omega)$  $\Gamma_8/\Gamma_5$ 

VALUE	DOCUMENT ID	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••		
$2.6 \pm 0.4$	BARBERIS 00F	$450 pp \rightarrow p_f\omega p_S$

NODE=M142R9  
NODE=M142R9 $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$  $\Gamma_{10}/\Gamma_9$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.09 \pm 0.05$	<sup>1</sup> ANISOVICH 11	SPEC	$0.9-1.94 p\bar{p}$

<sup>1</sup> Reanalysis of ADOMEIT 96 and ANISOVICH 00E.NODE=M142R12  
NODE=M142R12

NODE=M142R12;LINKAGE=AN

 $f_2(1910)$  REFERENCES

ANISOVICH 11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
AMELIN 06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=51574
AMSLER 06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
ANISOVICH 00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BARBERIS 00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
BARBERIS 00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47962
ADOMEIT 96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
BELADIDZE 92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42172
BELADIDZE 92D	ZPHY C57 13	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43309
ALDE 91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41844
Also	Translated from YAF 54 751.			
Also	PL B276 375	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41911
ALDE 90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=40935
ALDE 89	PL B216 447	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=40727
Also	SJNP 48 1035	D.M. Alde <i>et al.</i>	(BELG, SERP, LANL, LAPP)	REFID=44697
Also	Translated from YAF 48 1724.			
BALOSHIN 86	SJNP 43 959	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=40734
Also	Translated from YAF 43 1487.			

NODE=M142

**$a_0(1950)$** 

$$I^G(J^{PC}) = 1^-(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. Seen in  $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$  by LEES 16A  
with significance  $2.5\sigma$  in  $K_S^0 K^\pm \pi^\mp$  and  $4.2\sigma$  in  $K^+ K^- \pi^0$ .

NODE=M227

NODE=M227

 **$a_0(1950)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1931 \pm 14 \pm 22</math></b>	12k	<sup>1,2</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
$1949 \pm 32 \pm 76$	8k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
$1927 \pm 15 \pm 23$	4k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating width.

<sup>2</sup> Weighted average of the  $K_S^0 K^\pm$  and  $K^+ K^-$  decay modes.

NODE=M227M

NODE=M227M

OCCUR=3

OCCUR=2

NODE=M227M;LINKAGE=A

NODE=M227M;LINKAGE=B

 **$a_0(1950)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>271 \pm 22 \pm 29</math></b>	12k	<sup>1,2</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
$265 \pm 36 \pm 110$	8k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
$274 \pm 28 \pm 30$	4k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating mass.

<sup>2</sup> Weighted average of the  $K_S^0 K^\pm$  and  $K^+ K^-$  decay modes.

NODE=M227W

NODE=M227W

OCCUR=3

OCCUR=2

NODE=M227W;LINKAGE=A

NODE=M227W;LINKAGE=B

 **$a_0(1950)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\bar{K}$	seen

NODE=M227215;NODE=M227

DESIG=1

 **$a_0(1950)$  BRANCHING RATIOS**

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

<sup>1</sup> From a model-independent partial wave analysis.

NODE=M227225

NODE=M227R01

NODE=M227R01

NODE=M227R01;LINKAGE=A

 **$a_0(1950)$  REFERENCES**

LEES 16A PR D93 012005 J.P. Lees *et al.* (BABAR Collab.)

NODE=M227

REFID=57125

$f_2(1950)$ 

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M135

 $f_2(1950)$  T-MATRIX POLE  $\sqrt{s}$ Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

NODE=M135PP

NODE=M135PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1830-2020) - i (110-220) OUR ESTIMATE</b>			
$(1955 \pm 75) - i (175 \pm 57)$	<sup>1</sup> RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1978.2 \pm 1.8^{+28.4}_{-16.9}) - i$ $(118.8 \pm 0.8^{+20.8}_{-7.8})$	<sup>2</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
$(1867 \pm 46) - i (193 \pm 29)$	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
<sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).			
<sup>2</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).			

NODE=M135PP

→ UNCHECKED ←

NODE=M135PP;LINKAGE=A

NODE=M135PP;LINKAGE=B

 $f_2(1950)$  MASS

NODE=M135M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1936 ± 12 OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below.			
1940 ± 50	BAI	00A	BES $J/\psi \rightarrow \gamma (\pi^+ \pi^- \pi^+ \pi^-)$
1980 ± 22	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
1940 ± 22	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp2\pi2\pi^0$
1960 ± 30	BARBERIS	97B	OMEG $450 pp \rightarrow pp2(\pi^+ \pi^-)$
1918 ± 12	ANTINORI	95	OMEG $300,450 pp \rightarrow pp2(\pi^+ \pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$2038^{+13+12}_{-11-73}$	<sup>3</sup> UEHARA	09	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1930 ± 25	<sup>4</sup> BINON	05	GAMS $33 \pi^- p \rightarrow \eta \eta n$
1980 ± 2 ± 14	ABE	04	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
2010 ± 25	ANISOVICH	00J	SPEC
1980 ± 50	ANISOVICH	99B	SPEC $1.35-1.94 p\bar{p} \rightarrow \eta \eta \pi^0$
~ 1990	<sup>5</sup> OAKDEN	94	RVUE $0.36-1.55 \bar{p}p \rightarrow \pi\pi$
1950 ± 15	<sup>6</sup> ASTON	91	LASS $11 K^- p \rightarrow \Lambda K \bar{K} \pi \pi$
<sup>1</sup> Decaying into $\pi^+ \pi^- 2\pi^0$ .			
<sup>2</sup> Decaying into $2(\pi^+ \pi^-)$ .			
<sup>3</sup> Taking into account $f_4(2050)$ .			
<sup>4</sup> First solution, PWA is ambiguous.			
<sup>5</sup> From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.			
<sup>6</sup> Cannot determine spin to be 2.			

NODE=M135M

OCCUR=2

NODE=M135M;LINKAGE=A4


NODE=M135M;LINKAGE=B4

NODE=M135M;LINKAGE=UE

NODE=M135M;LINKAGE=BI

NODE=M135M;LINKAGE=BB

NODE=M135M;LINKAGE=A



**f<sub>2</sub>(1950) WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>464 ± 24 OUR AVERAGE</b>			
380 <sup>+120</sup> <sub>-90</sub>	BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
520 ± 50	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
485 ± 55	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
460 ± 40	BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
390 ± 60	ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

441 <sup>+27</sup> <sub>-25</sub> ± 28 <sub>-192</sub>	<sup>3</sup> UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
450 ± 50	<sup>4</sup> BINON	05 GAMS	33 $\pi^-p \rightarrow \eta\eta n$
297 ± 12 ± 6	ABE	04 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
495 ± 35	ANISOVICH	00J SPEC	
500 ± 100	ANISOVICH	99B SPEC	1.35-1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
~ 100	<sup>5</sup> OAKDEN	94 RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
250 ± 50	<sup>6</sup> ASTON	91 LASS	11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

<sup>1</sup> Decaying into  $\pi^+\pi^-2\pi^0$ .

<sup>2</sup> Decaying into  $2(\pi^+\pi^-)$ .

<sup>3</sup> Taking into account  $f_4(2050)$ .

<sup>4</sup> First solution, PWA is ambiguous.

<sup>5</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>6</sup> Cannot determine spin to be 2.

NODE=M135W

NODE=M135W

OCCUR=2

NODE=M135W;LINKAGE=A4

NODE=M135W;LINKAGE=B4

NODE=M135W;LINKAGE=UE

NODE=M135W;LINKAGE=BI

NODE=M135W;LINKAGE=BB

NODE=M135W;LINKAGE=A

**f<sub>2</sub>(1950) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\bar{K}^*(892)$	seen
$\Gamma_2$ $\pi\pi$	
$\Gamma_3$ $\pi^+\pi^-$	seen
$\Gamma_4$ $\pi^0\pi^0$	seen
$\Gamma_5$ $4\pi$	seen
$\Gamma_6$ $\pi^+\pi^-\pi^+\pi^-$	
$\Gamma_7$ $a_2(1320)\pi$	
$\Gamma_8$ $f_2(1270)\pi\pi$	
$\Gamma_9$ $\eta\eta$	seen
$\Gamma_{10}$ $K\bar{K}$	seen
$\Gamma_{11}$ $\gamma\gamma$	seen
$\Gamma_{12}$ $p\bar{p}$	seen

NODE=M135215;NODE=M135

DESIG=1

DESIG=11

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

DESIG=3

DESIG=4

DESIG=5

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=9;OUR EST;→ UNCHECKED ←

DESIG=12

$f_2(1950) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M135225

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{10}\Gamma_{11}/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M135G1  
NODE=M135G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$122 \pm 4 \pm 26$	<sup>1</sup> ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
--------------------	------------------	----	------	--

<sup>1</sup> Assuming spin 2.

NODE=M135G1;LINKAGE=AB

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_2\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M135G2  
NODE=M135G2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$162^{+69+1137}_{-42-204}$	<sup>1</sup> UEHARA	09	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
----------------------------	---------------------	----	------	--

<sup>1</sup> Taking into account  $f_4(2050)$ .

NODE=M135G2;LINKAGE=UE

 $f_2(1950)$  BRANCHING RATIOS

NODE=M135220

 $\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

NODE=M135R1  
NODE=M135R1

seen	ASTON	91	LASS	0	11 $K^- p \rightarrow \Lambda K \bar{K} \pi \pi$
------	-------	----	------	---	--

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M135R3  
NODE=M135R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
----------	----------	-----	--	---

not seen	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
----------	----------	-----	--	-----------------------------------

possibly seen	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
---------------	----------	-----	------	---------------------------------------

 $\Gamma(\eta\eta)/\Gamma(4\pi)$  $\Gamma_9/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

NODE=M135R5  
NODE=M135R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 5.0 \times 10^{-3}$	90	BARBERIS	00E	450 $pp \rightarrow p_f \eta \eta p_s$
------------------------	----	----------	-----	--

 $\Gamma(\eta\eta)/\Gamma(\pi^+ \pi^-)$  $\Gamma_9/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M135R6  
NODE=M135R6

<b>0.14 ± 0.05</b>	AMSLER	02	CBAR	0.9 $\bar{p} p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
--------------------	--------	----	------	--

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M135R07  
NODE=M135R07

seen	111	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p \bar{p}$
------	-----	-----------	----	--

 $f_2(1950)$  REFERENCES

NODE=M135

RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
		Translated from YAF 68 998.			
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47958
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>		REFID=46886
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=44437
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)	REFID=41746
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355

NODE=M017

**$a_4(1970)$**

$$I^G(J^{PC}) = 1^-(4^{++})$$

was  $a_4(2040)$

**$a_4(1970)$  MASS**

NODE=M017M

NODE=M017M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1967±16 OUR AVERAGE</b> Error includes scale factor of 2.1. See the ideogram below.					
1935 <sup>+11</sup> <sub>-13</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1900 <sup>+80</sup> <sub>-20</sub>		ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
1985±10±13	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1996±25±43		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi p$
2000±40 <sup>+60</sup> <sub>-20</sub>		IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
2010±20		<sup>2</sup> DONSKOV	96	GAM2 0	38 $\pi^- p \rightarrow \eta \pi^0 n$
2040±30		<sup>3</sup> CLELAND	82B	SPEC ±	50 $\pi p \rightarrow K_S^0 K^\pm p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1885±13 <sup>+50</sup> <sub>-2</sub>	420k	<sup>4</sup> ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
2004± 6	80k	<sup>5</sup> UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
2005 <sup>+25</sup> <sub>-45</sub>		<sup>6</sup> ANISOVICH	01F	SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
1944± 8±50		<sup>7</sup> AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1903±10		<sup>8</sup> BALDI	78	SPEC -	10 $\pi^- p \rightarrow p K_S^0 K^-$
2030±50		<sup>9</sup> CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi n$

- <sup>1</sup> Statistical error negligible.
- <sup>2</sup> From a simultaneous fit to the  $G_+$  and  $G_0$  wave intensities.
- <sup>3</sup> From an amplitude analysis.
- <sup>4</sup> Superseded by AGHASYAN 2018B.
- <sup>5</sup> Statistical error only.
- <sup>6</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.
- <sup>7</sup> May be a different state.
- <sup>8</sup> From a fit to the  $Y_8^0$  moment. Limited by phase space.
- <sup>9</sup>  $J^P = 4^+$  is favored, though  $J^P = 2^+$  cannot be excluded.

NODE=M017M;LINKAGE=B  
 NODE=M017M;LINKAGE=A  
 NODE=M017M;LINKAGE=C  
 NODE=M017M;LINKAGE=D  
 NODE=M017M;LINKAGE=ST  
 NODE=M017M;LINKAGE=AN  
 NODE=M017M;LINKAGE=DM  
 NODE=M017M;LINKAGE=Y  
 NODE=M017M;LINKAGE=M



**$a_4(1970)$  WIDTH**

NODE=M017W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>324<sup>+</sup><sub>-18</sub></b>	<b>OUR AVERAGE</b>				
333 <sup>+</sup> <sub>-21</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
300 <sup>+</sup> <sub>-100</sub>		ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
231 $\pm$ 30 $\pm$ 46	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
298 $\pm$ 81 $\pm$ 85		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi p$
350 $\pm$ 100 <sup>+</sup> <sub>-50</sub>		IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
370 $\pm$ 80		<sup>2</sup> DONSKOV	96	GAM2 0	38 $\pi^- p \rightarrow \eta \pi^0 n$
380 $\pm$ 150		<sup>3</sup> CLELAND	82B	SPEC $\pm$	50 $\pi p \rightarrow K_S^0 K^\pm p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

294 $\pm$ 25 <sup>+</sup> <sub>-19</sub>	420k	<sup>4</sup> ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ P_b'$
401 $\pm$ 16	80k	<sup>5</sup> UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
180 $\pm$ 30		<sup>6</sup> ANISOVICH	01F	SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
324 $\pm$ 26 $\pm$ 75		<sup>7</sup> AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
166 $\pm$ 43		<sup>8</sup> BALDI	78	SPEC $-$	10 $\pi^- p \rightarrow p K_S^0 K^-$
510 $\pm$ 200		<sup>9</sup> CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi n$

<sup>1</sup> Statistical error negligible.

<sup>2</sup> From a simultaneous fit to the  $G_+$  and  $G_0$  wave intensities.

<sup>3</sup> From an amplitude analysis.

<sup>4</sup> Superseded by AGHASYAN 2018B.

<sup>5</sup> Statistical error only.

<sup>6</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

<sup>7</sup> May be a different state.

<sup>8</sup> From a fit to the  $Y_8^0$  moment. Limited by phase space.

<sup>9</sup>  $J^P = 4^+$  is favored, though  $J^P = 2^+$  cannot be excluded.

NODE=M017W

NODE=M017W;LINKAGE=B  
 NODE=M017W;LINKAGE=A  
 NODE=M017W;LINKAGE=C  
 NODE=M017W;LINKAGE=D  
 NODE=M017W;LINKAGE=ST  
 NODE=M017W;LINKAGE=AN  
 NODE=M017W;LINKAGE=DM  
 NODE=M017W;LINKAGE=Y  
 NODE=M017W;LINKAGE=M

### $a_4(1970)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \bar{K}$	seen
$\Gamma_2$ $\pi^+ \pi^- \pi^0$	seen
$\Gamma_3$ $\rho \pi$	seen
$\Gamma_4$ $f_2(1270) \pi$	seen
$\Gamma_5$ $\omega \pi^- \pi^0$	seen
$\Gamma_6$ $\omega \rho$	seen
$\Gamma_7$ $\eta \pi$	seen
$\Gamma_8$ $\eta'(958) \pi$	seen

NODE=M017215;NODE=M017

DESIG=1  
 DESIG=2  
 DESIG=5;OUR EST;→ UNCHECKED ←  
 DESIG=6;OUR EST;→ UNCHECKED ←  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=8  
 DESIG=3  
 DESIG=4;OUR EST;→ UNCHECKED ←

### $a_4(1970)$ BRANCHING RATIOS

$\Gamma(K \bar{K})/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	BALDI	78	SPEC $\pm$	10 $\pi^- p \rightarrow K_S^0 K^- p$
$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi n$
$\Gamma(\rho \pi)/\Gamma(f_2(1270) \pi)$	$\Gamma_3/\Gamma_4$			
VALUE	DOCUMENT ID	TECN	COMMENT	

NODE=M017220

NODE=M017R1  
 NODE=M017R1

NODE=M017R2  
 NODE=M017R2

NODE=M017R4  
 NODE=M017R4

**1.7<sup>+</sup><sub>-0.8</sub>** OUR AVERAGE Error includes scale factor of 3.7.

2.9 <sup>+</sup> <sub>-0.4</sub>	46M	<sup>1</sup> AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1.1 $\pm$ 0.2 $\pm$ 0.2		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi p$

<sup>1</sup> Statistical error negligible.

NODE=M017R4;LINKAGE=A



$\Gamma(\eta\pi)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_7/\Gamma$
seen	DONSKOV	96	GAM2	0	$38 \pi^- p \rightarrow \eta \pi^0 n$

NODE=M017R3  
NODE=M017R3

 $\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma_7$
0.23±0.07	ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$

NODE=M017R01  
NODE=M017R01

 $\Gamma(\omega\rho)/\Gamma_{\text{total}}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma$
seen	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$

NODE=M017R5  
NODE=M017R5

 $a_4(1970)$  REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
		Translated from YAF 62 487.			
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>		REFID=46902
DONSKOV	96	PAN 59 982	S.V. Donskov <i>et al.</i>	(GAMS Collab.) IGJPC	REFID=45207
		Translated from YAF 59 1027.			
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
BALDI	78	PL 74B 413	R. Baldi <i>et al.</i>	(GEVA) JP	REFID=21783
CORDEN	78C	NP B136 77	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20859

NODE=M017

 $\rho_3(1990)$ 

$$I^G(J^{PC}) = 1^+(3^{--})$$

OMITTED FROM SUMMARY TABLE

NODE=M167

 $\rho_3(1990)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1982±14	<sup>1</sup> ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2007	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167M

NODE=M167M

NODE=M167M;LINKAGE=AY

 $\rho_3(1990)$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

188±24	<sup>2</sup> ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 287	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

<sup>2</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167W

NODE=M167W

NODE=M167W;LINKAGE=AY

 $\rho_3(1990)$  REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103

NODE=M167

$\pi_2(2005)$ 

$$I^G(J^{PC}) = 1^-(2^-+)$$

NODE=M239

OMITTED FROM SUMMARY TABLE

 $\pi_2(2005)$  MASS

NODE=M239M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1963<sup>+17</sup><sub>-27</sub> OUR AVERAGE</b>				
1962 <sup>+17</sup> <sub>-29</sub>	46M	<sup>1</sup> AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1974 ± 14 ± 83	145k	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2005 ± 15		ANISOVICH	01F SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
<sup>1</sup> Statistical uncertainty negligible.				

NODE=M239M

NODE=M239M;LINKAGE=A

 $\pi_2(2005)$  WIDTH

NODE=M239W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>370<sup>+16</sup><sub>-90</sub> OUR AVERAGE</b>				
371 <sup>+16</sup> <sub>-120</sub>	46M	<sup>1</sup> AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
341 ± 61 ± 139	145k	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200 ± 40		ANISOVICH	01F SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
<sup>1</sup> Statistical uncertainty negligible.				

NODE=M239W

NODE=M239W;LINKAGE=A

 $\pi_2(2005)$  DECAY MODES

NODE=M239215;NODE=M239

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi^- \pi^+ \pi^-$	seen
$\Gamma_2$ $\omega \pi^0 \pi^-$	seen

DESIG=1

DESIG=2

 $\pi_2(2005)$  BRANCHING RATIOS

NODE=M239220

$\Gamma(\pi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$\Gamma(\omega \pi^0 \pi^-)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$

NODE=M239R00  
NODE=M239R00NODE=M239R01  
NODE=M239R01 $\pi_2(2005)$  REFERENCES

NODE=M239

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352

**$f_2(2010)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M106

 **$f_2(2010)$  MASS**

NODE=M106M

NODE=M106M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2011^{+62}_{-76}$	<sup>1</sup> ETKIN	88 MPS	$22 \pi^- p \rightarrow \phi \phi n$
$2062 \pm 6^{+10}_{-7}$	<sup>2</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
$2005 \pm 12$	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1980 \pm 20$	<sup>3</sup> BOLONKIN	88 SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$2050^{+90}_{-50}$	ETKIN	85 MPS	$22 \pi^- p \rightarrow 2 \phi n$
$2120^{+20}_{-120}$	LINDENBAUM	84 RVUE	
$2160 \pm 50$	ETKIN	82 MPS	$22 \pi^- p \rightarrow 2 \phi n$

<sup>1</sup> Includes data of ETKIN 85. The percentage of the resonance going into  $\phi \phi 2^{++} S_2$ ,  $D_2$ , and  $D_0$  is  $98^{+1}_{-3}$ ,  $0^{+1}_{-0}$ , and  $2^{+2}_{-1}$ , respectively.

<sup>2</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

<sup>3</sup> Statistically very weak, only 1.4 s.d.

NODE=M106M;LINKAGE=C

NODE=M106M;LINKAGE=A

NODE=M106M;LINKAGE=E

 **$f_2(2010)$  WIDTH**

NODE=M106W

NODE=M106W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$202^{+67}_{-62}$	<sup>4</sup> ETKIN	88 MPS	$22 \pi^- p \rightarrow \phi \phi n$
$165 \pm 17^{+10}_{-5}$	<sup>5</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
$209 \pm 32$	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$145 \pm 50$	<sup>6</sup> BOLONKIN	88 SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$200^{+160}_{-50}$	ETKIN	85 MPS	$22 \pi^- p \rightarrow 2 \phi n$
$300^{+150}_{-50}$	LINDENBAUM	84 RVUE	
$310 \pm 70$	ETKIN	82 MPS	$22 \pi^- p \rightarrow 2 \phi n$

<sup>4</sup> Includes data of ETKIN 85.

<sup>5</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

<sup>6</sup> Statistically very weak, only 1.4 s.d.

NODE=M106W;LINKAGE=C

NODE=M106W;LINKAGE=A

NODE=M106W;LINKAGE=E

 **$f_2(2010)$  DECAY MODES**

NODE=M106215;NODE=M106

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \phi \phi$	seen
$\Gamma_2 \quad K \bar{K}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2

 **$f_2(2010)$  BRANCHING RATIOS**

NODE=M106230

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$

NODE=M106R01

NODE=M106R01

 **$f_2(2010)$  REFERENCES**

NODE=M106

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
VLADIMIRSK...06		PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)
ETKIN	82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)
Also		Brighton Conf. 351	S.J. Lindenbaum	(BNL, CUNY)

REFID=61891

REFID=51191

REFID=40580

REFID=40285

REFID=21871

REFID=21869

REFID=21866

REFID=21867

**$f_0(2020)$** 

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

 **$f_0(2020)$  T-MATRIX POLE  $\sqrt{s}$** Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1870–2080) – <math>i</math> (120–240) OUR ESTIMATE</b>			
$(2038 \pm 48) - i(156 \pm 41)$	<sup>1</sup> RODAS 22	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
$(1925 \pm 25) - i(160 \pm 18)$	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1910 \pm 50) - i(199 \pm 40)$	<sup>2</sup> ROPERTZ 18	RVUE	$\bar{B}_S^0 \rightarrow J/\psi(\pi^+\pi^-/K^+K^-)$
$(1992 \pm 16) - i(221 \pm 30)$	<sup>3</sup> BARBERIS 00C	450	$p\bar{p} \rightarrow p_f 4\pi p_s$
<sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma\pi^0\pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).			
<sup>2</sup> T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.			
<sup>3</sup> Average between $\pi^+\pi^-2\pi^0$ and $2(\pi^+\pi^-)$ .			

 **$f_0(2020)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$2010 \pm 6_{-4}^{+6}$		<sup>1</sup> ABLIKIM 22AS	BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
$1982 \pm 3_{-0}^{+54}$		<sup>2</sup> ABLIKIM 22C	BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
$2037 \pm 8$	80k	<sup>3</sup> UMAN 06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
$2040 \pm 38$		ANISOVICH 00J	SPEC	
$2010 \pm 60$		ALDE 98	GAM4	$100 \pi^- p \rightarrow \pi^0\pi^0 n$
$2020 \pm 35$		BARBERIS 97B	OMEG	$450 p\bar{p} \rightarrow p\bar{p} 2(\pi^+\pi^-)$
<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.				
<sup>2</sup> From a partial wave analysis of the systems $(\gamma X)$ , with $X \rightarrow \eta'\eta'$ , and $(\eta' X)$ , with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance $X$ is parametrized by a constant-width, relativistic Breit-Wigner.				
<sup>3</sup> Statistical error only.				

 **$f_0(2020)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$203 \pm 9_{-11}^{+13}$		<sup>1</sup> ABLIKIM 22AS	BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
$436 \pm 4_{-49}^{+46}$		<sup>2</sup> ABLIKIM 22C	BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
$296 \pm 17$	80k	<sup>3</sup> UMAN 06	E835	$5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
$405 \pm 40$		ANISOVICH 00J	SPEC	
$240 \pm 100$		ALDE 98	GAM4	$100 \pi^- p \rightarrow \pi^0\pi^0 n$
$410 \pm 50$		BARBERIS 97B	OMEG	$450 p\bar{p} \rightarrow p\bar{p} 2(\pi^+\pi^-)$
<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.				
<sup>2</sup> From a partial wave analysis of the systems $(\gamma X)$ , with $X \rightarrow \eta'\eta'$ , and $(\eta' X)$ , with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance $X$ is parametrized by a constant-width, relativistic Breit-Wigner.				
<sup>3</sup> Statistical error only.				

NODE=M156

NODE=M156

NODE=M156PP

NODE=M156PP

NODE=M156PP

→ UNCHECKED ←

NODE=M156PP;LINKAGE=A

NODE=M156PP;LINKAGE=C

NODE=M156PP;LINKAGE=B

NODE=M156M

NODE=M156M

NODE=M156M;LINKAGE=D

NODE=M156M;LINKAGE=C

NODE=M156M;LINKAGE=ST

NODE=M156W

NODE=M156W

NODE=M156W;LINKAGE=D

NODE=M156W;LINKAGE=C

NODE=M156W;LINKAGE=ST

$f_0(2020)$  DECAY MODES

NODE=M156215;NODE=M156

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\pi\pi$	seen
$\Gamma_2$ $\pi^0\pi^0$	seen
$\Gamma_3$ $\rho\rho$	seen
$\Gamma_4$ $\omega\omega$	seen
$\Gamma_5$ $\eta\eta$	seen
$\Gamma_6$ $\eta'\eta'$	seen

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=3;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=5  
DESIG=6

 $f_0(2020)$  BRANCHING RATIOS

NODE=M156220

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$	DOCUMENT ID	COMMENT	$\Gamma_3/\Gamma_4$
VALUE			

NODE=M156R1  
NODE=M156R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 3 BARBERIS 00F 450  $\rho\rho \rightarrow p_f\omega\omega p_s$

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma$
VALUE				

NODE=M156R01  
NODE=M156R01

seen UMAN 06 E835 5.2  $\bar{p}p \rightarrow \eta\eta\pi^0$

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma$
VALUE				

NODE=M156R00  
NODE=M156R00

seen <sup>1</sup> ABLIKIM 22C BES3  $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

<sup>1</sup> From a partial wave analysis of the systems ( $\gamma X$ ), with  $X \rightarrow \eta'\eta'$ , and ( $\eta' X$ ), with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M156R00;LINKAGE=A

 $f_0(2020)$  REFERENCES

NODE=M156

ABLIKIM 22AS PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 22C PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
RODAS 22 EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
SARANTSEV 21 PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM 18AA PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ROPERTZ 18 EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis	(BONN, JULI)
AAIJ 17V JHEP 1708 037	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM 15AE PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ 14BR PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)
UMAN 06 PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)
ANISOVICH 00J PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARBERIS 00C PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS 00F PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ALDE 98 EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
Translated from YAF 62 446.		
BARBERIS 97B PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)

REFID=61891  
REFID=61637  
REFID=61610  
REFID=61091  
REFID=59455  
REFID=59332  
REFID=57828  
REFID=56984  
REFID=56035  
REFID=51063  
REFID=47950  
REFID=47959  
REFID=47962  
REFID=46605  
REFID=46914  
REFID=45758

$f_4(2050)$ 

$$I^G(J^{PC}) = 0^+(4^{++})$$

NODE=M016

 $f_4(2050)$  MASS

NODE=M016M

NODE=M016M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>2018±11 OUR AVERAGE</b>		Error includes scale factor of 2.1. See the ideogram below.		
1960±15		AMELIN	06 VES	36 $\pi^- p \rightarrow \omega \omega n$
2005±10		<sup>1</sup> BINON	05 GAMS	33 $\pi^- p \rightarrow \eta \eta n$
1998±15		ALDE	98 GAM4	100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
2060±20		ALDE	90 GAM2	38 $\pi^- p \rightarrow \omega \omega n$
2038±30		AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2086±15		BALTRUSAIT..	87 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2000±60		ALDE	86D GAM4	100 $\pi^- p \rightarrow n 2 \eta$
2020±20	40k	<sup>2</sup> BINON	84B GAM2	38 $\pi^- p \rightarrow n 2 \pi^0$
2015±28		<sup>3</sup> CASON	82 STRC	8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
2031 <sup>+25</sup> <sub>-36</sub>		ETKIN	82B MPS	23 $\pi^- p \rightarrow n 2 K_S^0$
2020±30	700	APEL	75 NICE	40 $\pi^- p \rightarrow n 2 \pi^0$
2050±25		BLUM	75 ASPK	18.4 $\pi^- p \rightarrow n K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1966±25		<sup>4</sup> ANISOVICH	09 RVUE	0.0 $\bar{p} p, \pi N$
1885 <sup>+14+218</sup> <sub>-13-25</sub>		<sup>5</sup> UEHARA	09 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
2018± 6		ANISOVICH	00J SPEC	2.0 $\bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$ $\eta \eta, \eta \eta', \pi \pi$
~ 2000		<sup>6</sup> MARTIN	98 RVUE	$\bar{N} N \rightarrow \pi \pi$
~ 2010		<sup>7</sup> MARTIN	97 RVUE	$\bar{N} N \rightarrow \pi \pi$
~ 2040		<sup>8</sup> OAKDEN	94 RVUE	0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
~ 1990		<sup>9</sup> OAKDEN	94 RVUE	0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
1978± 5		<sup>10</sup> ALPER	80 CNTR	62 $\pi^- p \rightarrow K^+ K^- n$
2040±10		<sup>10</sup> ROZANSKA	80 SPRK	18 $\pi^- p \rightarrow p \bar{p} n$
1935±13		<sup>10</sup> CORDEN	79 OMEG	12-15 $\pi^- p \rightarrow n 2 \pi$
1988± 7		EVANGELIS...	79B OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
1922±14		<sup>11</sup> ANTIPOV	77 CIBS	25 $\pi^- p \rightarrow p 3 \pi$


OCCUR=2

<sup>1</sup> From the first PWA solution.<sup>2</sup> From a partial-wave analysis of the data.<sup>3</sup> From an amplitude analysis of the reaction  $\pi^+ \pi^- \rightarrow 2 \pi^0$ .<sup>4</sup> K matrix pole.<sup>5</sup> Taking into account the  $f_2(1950)$ . Helicity-2 production favored.<sup>6</sup> Energy-dependent analysis.<sup>7</sup> Single energy analysis.<sup>8</sup> From solution A of amplitude analysis of data on  $\bar{p} p \rightarrow \pi \pi$ . See however KLOET 96 who fit  $\pi^+ \pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.<sup>9</sup> From solution B of amplitude analysis of data on  $\bar{p} p \rightarrow \pi \pi$ . See however KLOET 96 who fit  $\pi^+ \pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.<sup>10</sup>  $I(J^P) = 0(4^+)$  from amplitude analysis assuming one-pion exchange.<sup>11</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

NODE=M016M;LINKAGE=BI  
 NODE=M016M;LINKAGE=N  
 NODE=M016M;LINKAGE=NN  
 NODE=M016M;LINKAGE=KM  
 NODE=M016M;LINKAGE=UE  
 NODE=M016M;LINKAGE=RB  
 NODE=M016M;LINKAGE=BR  
 NODE=M016M;LINKAGE=B

NODE=M016M;LINKAGE=BB

NODE=M016M;LINKAGE=M  
 NODE=M016M;LINKAGE=T



**$f_4(2050)$  WIDTH**

NODE=M016W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>237± 18 OUR AVERAGE</b>		Error includes scale factor of 1.9. See the ideogram below.		
290± 20		AMELIN	06	VES 36 $\pi^- p \rightarrow \omega \omega n$
340± 80		12 BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
395± 40		ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
170± 60		ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega \omega n$
304± 60		AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
210± 63		BALTRUSAIT..	87	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
400±100		ALDE	86D	GAM4 100 $\pi^- p \rightarrow n 2 \eta$
240± 40	40k	13 BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2 \pi^0$
190± 14		DENNEY	83	LASS 10 $\pi^+ n/\pi^+ p$
186 <sup>+103</sup> <sub>-58</sub>		14 CASON	82	STRC 8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
305 <sup>+36</sup> <sub>-119</sub>		ETKIN	82B	MPS 23 $\pi^- p \rightarrow n 2 K_S^0$
180± 60	700	APEL	75	NICE 40 $\pi^- p \rightarrow n 2 \pi^0$
225 <sup>+120</sup> <sub>-70</sub>		BLUM	75	ASPK 18.4 $\pi^- p \rightarrow n K^+ K^-$

NODE=M016W

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 260± 40 15 ANISOVICH 09 RVUE 0.0  $\bar{p} p, \pi N$
- 453± 20<sup>+31</sup><sub>-129</sub> 16 UEHARA 09 BELL 10.6  $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
- 182± 7 ANISOVICH 00J SPEC 2.0  $\bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$   
 $\eta \eta, \eta \eta', \pi \pi$
- ~ 170 17 MARTIN 98 RVUE  $N \bar{N} \rightarrow \pi \pi$
- ~ 200 18 MARTIN 97 RVUE  $\bar{N} N \rightarrow \pi \pi$
- ~ 60 19 OAKDEN 94 RVUE 0.36-1.55  $\bar{p} p \rightarrow \pi \pi$
- ~ 80 20 OAKDEN 94 RVUE 0.36-1.55  $\bar{p} p \rightarrow \pi \pi$
- 243± 16 21 ALPER 80 CNTR 62  $\pi^- p \rightarrow K^+ K^- n$
- 140± 15 21 ROZANSKA 80 SPRK 18  $\pi^- p \rightarrow p \bar{p} n$
- 263± 57 21 CORDEN 79 OMEG 12-15  $\pi^- p \rightarrow n 2 \pi$
- 100± 28 EVANGELIS... 79B OMEG 10  $\pi^- p \rightarrow K^+ K^- n$
- 107± 56 22 ANTIPOV 77 CIBS 25  $\pi^- p \rightarrow p 3 \pi$

OCCUR=2

12 From the first PWA solution.  
 13 From a partial-wave analysis of the data.  
 14 From an amplitude analysis of the reaction  $\pi^+ \pi^- \rightarrow 2 \pi^0$ .  
 15 K matrix pole.  
 16 Taking into account the  $f_2(1950)$ . Helicity-2 production favored.  
 17 Energy-dependent analysis.  
 18 Single energy analysis.


NODE=M016W;LINKAGE=BI  
 NODE=M016W;LINKAGE=N  
 NODE=M016W;LINKAGE=NN  
 NODE=M016W;LINKAGE=KM  
 NODE=M016W;LINKAGE=UE  
 NODE=M016W;LINKAGE=RB  
 NODE=M016W;LINKAGE=BR

- 19 From solution A of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.
- 20 From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.
- 21  $I(J^P) = 0(4^+)$  from amplitude analysis assuming one-pion exchange.
- 22 Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

NODE=M016W;LINKAGE=BW

NODE=M016W;LINKAGE=BB

NODE=M016W;LINKAGE=M  
NODE=M016W;LINKAGE=T



**$f_4(2050)$  DECAY MODES**

NODE=M016215;NODE=M016

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\omega\omega$	seen
$\Gamma_2$ $\pi\pi$	(17.0 ± 1.5) %
$\Gamma_3$ $K\bar{K}$	( 6.8 <sup>+3.4</sup> <sub>-1.8</sub> ) × 10 <sup>-3</sup>
$\Gamma_4$ $\eta\eta$	( 2.1 ± 0.8) × 10 <sup>-3</sup>
$\Gamma_5$ $4\pi^0$	< 1.2 %
$\Gamma_6$ $\gamma\gamma$	seen
$\Gamma_7$ $a_2(1320)\pi$	seen

DESIG=6  
DESIG=1  
DESIG=2  
DESIG=3  
DESIG=5  
DESIG=4;OUR EVAL;→ UNCHECKED ←  
DESIG=7

**$f_4(2050)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$**

NODE=M016220

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_6/\Gamma$
VALUE (keV) CL% DOCUMENT ID TECN COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••	
<0.29 95 ALTHOFF 85B TASS $\gamma\gamma \rightarrow K\bar{K}\pi$	

NODE=M016G2  
NODE=M016G2

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_6/\Gamma$
VALUE (eV) CL% EVTS DOCUMENT ID TECN COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••	
23.1 <sup>+3.6+70.5</sup> <sub>-3.3-15.6</sub> 23 UEHARA 09 BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	
<1100 95 13 ± 4 OEST 90 JADE $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	

NODE=M016G3  
NODE=M016G3

<sup>23</sup> Taking into account the  $f_2(1950)$ . Helicity-2 production favored.

NODE=M016G3;LINKAGE=UE

**$f_4(2050)$  BRANCHING RATIOS**

NODE=M016225

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
VALUE DOCUMENT ID TECN COMMENT	
seen AMELIN 06 VES 36 $\pi^-p \rightarrow \omega\omega n$	
••• We do not use the following data for averages, fits, limits, etc. •••	
not seen BARBERIS 00F 450 $pp \rightarrow pf\omega\omega p_S$	

NODE=M016R7  
NODE=M016R7



$\Gamma(\omega\omega)/\Gamma(\pi\pi)$				$\Gamma_1/\Gamma_2$	
VALUE	DOCUMENT ID	TECN	COMMENT		
<b>1.5±0.3</b>	ALDE	90	GAM2	38 $\pi^- p \rightarrow \omega\omega n$	NODE=M016R5 NODE=M016R5
$\Gamma(\pi\pi)/\Gamma_{total}$				$\Gamma_2/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT		
<b>0.170±0.015 OUR AVERAGE</b>					NODE=M016R1 NODE=M016R1
0.18 ±0.03	24 BINON	83C	GAM2	38 $\pi^- p \rightarrow n4\gamma$	
0.16 ±0.03	24 CASON	82	STRC	8 $\pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$	
0.17 ±0.02	24 CORDEN	79	OMEG	12-15 $\pi^- p \rightarrow n2\pi$	
<sup>24</sup> Assuming one pion exchange.					NODE=M016R1;LINKAGE=A
$\Gamma(K\bar{K})/\Gamma(\pi\pi)$				$\Gamma_3/\Gamma_2$	
VALUE	DOCUMENT ID	TECN	COMMENT		
<b>0.04<sup>+0.02</sup><sub>-0.01</sub></b>	ETKIN	82B	MPS	23 $\pi^- p \rightarrow n2K_S^0$	NODE=M016R2 NODE=M016R2
$\Gamma(\eta\eta)/\Gamma_{total}$				$\Gamma_4/\Gamma$	
VALUE (units 10 <sup>-3</sup> )	DOCUMENT ID	TECN	COMMENT		
<b>2.1±0.8</b>	ALDE	86D	GAM4	100 $\pi^- p \rightarrow n4\gamma$	NODE=M016R3 NODE=M016R3
$\Gamma(4\pi^0)/\Gamma_{total}$				$\Gamma_5/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT		
<b>&lt;0.012</b>	ALDE	87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$	NODE=M016R4 NODE=M016R4
$\Gamma(a_2(1320)\pi)/\Gamma_{total}$				$\Gamma_7/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT		
<b>seen</b>	AMELIN	00	VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$	NODE=M016R6 NODE=M016R6

**f<sub>4</sub>(2050) REFERENCES**

					NODE=M016
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=51574
BINON	05	Translated from YAF 69 715. PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
AMELIN	00	Translated from YAF 68 998. NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47962
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
MARTIN	98	Translated from YAF 62 446. PR C57 3492	B.R. Martin <i>et al.</i>		REFID=46373
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)	REFID=45685
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=40935
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP)	REFID=21780
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+)	REFID=40288
DENNEY	83	Translated from YAF 38 1199. PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21665
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)	REFID=21774
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21967
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP	REFID=21651

$\pi_2(2100)$ 

$$I^G(J^{PC}) = 1^-(2^-+)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M020

NODE=M020

NODE=M020M

NODE=M020M

NODE=M020M;LINKAGE=AX  
NODE=M020M;LINKAGE=L

NODE=M020W

NODE=M020W

NODE=M020W;LINKAGE=AX  
NODE=M020W;LINKAGE=L

NODE=M020215;NODE=M020

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=3;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←

NODE=M020220

NODE=M020R1  
NODE=M020R1NODE=M020R2  
NODE=M020R2NODE=M020R3  
NODE=M020R3NODE=M020R4  
NODE=M020R4

NODE=M020R;LINKAGE=L

NODE=M020

REFID=44433  
REFID=20872 $\pi_2(2100)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2090± 29 OUR AVERAGE</b>			
2090± 30	<sup>1</sup> AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
2100± 150	<sup>2</sup> DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$
<sup>1</sup> From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, (\pi\pi)_S\pi$ waves.			
<sup>2</sup> From a two-resonance fit to four $2^-0^+$ waves.			

 $\pi_2(2100)$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>625± 50 OUR AVERAGE</b>	Error includes scale factor of 1.2.		
520± 100	<sup>3</sup> AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
651± 50	<sup>4</sup> DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$
<sup>3</sup> From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, (\pi\pi)_S\pi$ waves.			
<sup>4</sup> From a two-resonance fit to four $2^-0^+$ waves.			

 $\pi_2(2100)$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $3\pi$	seen
$\Gamma_2$ $\rho\pi$	seen
$\Gamma_3$ $f_2(1270)\pi$	seen
$\Gamma_4$ $(\pi\pi)_S\pi$	seen

 $\pi_2(2100)$  BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(3\pi)$	$\Gamma_2/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.19±0.05</b>	<sup>5</sup> DAUM	81B CNTR	63,94 $\pi^- p$
$\Gamma(f_2(1270)\pi)/\Gamma(3\pi)$	$\Gamma_3/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.36±0.09</b>	<sup>5</sup> DAUM	81B CNTR	63,94 $\pi^- p$
$\Gamma((\pi\pi)_S\pi)/\Gamma(3\pi)$	$\Gamma_4/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.45±0.07</b>	<sup>5</sup> DAUM	81B CNTR	63,94 $\pi^- p$
<b>D-wave/S-wave RATIO FOR <math>\pi_2(2100) \rightarrow f_2(1270)\pi</math></b>			
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.39±0.23</b>	<sup>5</sup> DAUM	81B CNTR	63,94 $\pi^- p$
<sup>5</sup> From a two-resonance fit to four $2^-0^+$ waves.			

 $\pi_2(2100)$  REFERENCES

AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

$f_0(2100)$ 

$$J^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M168

NODE=M168

 $f_0(2100)$  MASS

NODE=M168M

NODE=M168M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2095<sup>+17</sup><sub>-19</sub> OUR AVERAGE</b>				
2116±27±17		LEES	21A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
2081±13 <sup>+24</sup> <sub>-36</sub>	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
2090±30		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2075±20		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
2090±10±6	529	<sup>2,3</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2099±17±8	283	<sup>2,3</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
2105±8	80k	<sup>4</sup> UMAN	06 E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
2102±13		<sup>5</sup> ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0, \eta \eta, \eta \eta', \pi^+ \pi^-$
2105±10		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p} p \rightarrow \eta \eta, \eta \eta'$
~ 2104		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
~ 2122		HASAN	94 RVUE	$\bar{p} p \rightarrow \pi \pi$

OCCUR=2

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>3</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 209$  MeV.<sup>4</sup> Statistical error only.<sup>5</sup> Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

NODE=M168M;LINKAGE=A

NODE=M168M;LINKAGE=B

NODE=M168M;LINKAGE=C

NODE=M168M;LINKAGE=ST

NODE=M168M;LINKAGE=AN

 $f_0(2100)$  WIDTH

NODE=M168W

NODE=M168W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>287<sup>+32</sup><sub>-24</sub> OUR AVERAGE</b>				
289±34±15		LEES	21A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
273 <sup>+27</sup> <sub>-24</sub> ±70 <sub>-23</sub>	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
330±100		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
260±25		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
236±14	80k	<sup>2</sup> UMAN	06 E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
211±29		<sup>3</sup> ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0, \eta \eta, \eta \eta', \pi^+ \pi^-$
200±25		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p} p \rightarrow \eta \eta, \eta \eta'$
~ 203		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
~ 273		HASAN	94 RVUE	$\bar{p} p \rightarrow \pi \pi$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.<sup>2</sup> Statistical error only.<sup>3</sup> Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

NODE=M168W;LINKAGE=A

NODE=M168W;LINKAGE=ST

NODE=M168W;LINKAGE=AN

 $f_0(2100)$  REFERENCES

NODE=M168

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61442
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103

**$f_2(2150)$** 

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called  $T_0$ .

NODE=M042

NODE=M042

NODE=M042205

NODE=M042M

NODE=M042M

 **$f_2(2150)$  MASS** **$f_2(2150)$  MASS, COMBINED MODES (MeV)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**2157±12 OUR AVERAGE** Includes data from the datablock that follows this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2170±6      80k      <sup>1</sup>UMAN      06      E835      5.2  $\bar{p}p \rightarrow \eta\eta\pi^0$ <sup>1</sup>Statistical error only.

NODE=M042M;LINKAGE=ST

 **$\eta\eta$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M042M3

NODE=M042M3

**2157±12 OUR AVERAGE**

2151±16	BARBERIS	00E	450 $pp \rightarrow pf\eta\eta p_S$
2175±20	PROKOSHKIN	95D	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$ , 450 $pp \rightarrow pp 2\eta$
2130±35	SINGOVSKI	94	GAM4 450 $pp \rightarrow pp 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2140±30      <sup>2</sup>ABELE      99B      CBAR      1.94  $\bar{p}p \rightarrow \pi^0\eta\eta$ 2104±20      <sup>3</sup>ARMSTRONG      93C      E760       $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$ <sup>2</sup>Spin not determined.<sup>3</sup>No  $J^{PC}$  determination.

NODE=M042M3;LINKAGE=K3

NODE=M042M3;LINKAGE=A

 **$\eta\pi\pi$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135±20±45      <sup>4</sup>ADOMEIT      96      CBAR      0      1.94  $\bar{p}p \rightarrow \eta 3\pi^0$ <sup>4</sup>ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single  $J^P = 2^+$  resonance.

NODE=M042M4;LINKAGE=AD

 **$\bar{p}p \rightarrow \pi\pi$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2090      <sup>5</sup>OAKDEN      94      RVUE      0.36–1.55  $\bar{p}p \rightarrow \pi\pi$ ~ 2120      <sup>6</sup>OAKDEN      94      RVUE      0.36–1.55  $\bar{p}p \rightarrow \pi\pi$ ~ 2170      <sup>7</sup>MARTIN      80B      RVUE~ 2150      <sup>7</sup>MARTIN      80C      RVUE~ 2150      <sup>8</sup>DULUDE      78B      OSPK      1–2  $\bar{p}p \rightarrow \pi^0\pi^0$ 

OCCUR=2

<sup>5</sup>OAKDEN 94 makes an amplitude analysis of LEAR data on  $\bar{p}p \rightarrow \pi\pi$  using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.<sup>6</sup>From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ .<sup>7</sup> $I(J^P) = 0(2^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^-\pi^+$  and  $\pi^0\pi^0$ .<sup>8</sup> $I^G(J^P) = 0^+(2^+)$  from partial-wave amplitude analysis.

NODE=M042M1;LINKAGE=B

NODE=M042M1;LINKAGE=BB

NODE=M042M1;LINKAGE=P

NODE=M042M1;LINKAGE=L

**S-CHANNEL  $\bar{p}p, \bar{N}N$  or  $\bar{K}K$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2139<sup>+8</sup><sub>-9</sub>      <sup>9</sup>EVANGELIS...      97      SPEC      0.6–2.4  $\bar{p}p \rightarrow K_S^0 K_S^0$ ~ 2190      <sup>9</sup>CUTTS      78B      CNTR      0.97–3  $\bar{p}p \rightarrow \bar{N}N$ 2155±15      <sup>9,10</sup>COUPLAND      77      CNTR      0      0.7–2.4  $\bar{p}p \rightarrow \bar{p}p$ 2193±2      <sup>9,11</sup>ALSPECTOR      73      CNTR       $\bar{p}p$  S channel<sup>9</sup>Isospins 0 and 1 not separated.<sup>10</sup>From a fit to the total elastic cross section.<sup>11</sup>Referred to as  $T$  or  $T$  region by ALSPECTOR 73.

NODE=M042M2

NODE=M042M2

NODE=M042M2;LINKAGE=I

NODE=M042M2;LINKAGE=E

NODE=M042M2;LINKAGE=M

**$K\bar{K}$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2200 \pm 13$	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$2150 \pm 20$	ABLIKIM	04E	$J/\psi \rightarrow \omega K^+ K^-$
$2130 \pm 35$	BARBERIS	99	$450 p p \rightarrow p_s p_f K^+ K^-$

NODE=M042M5  
 NODE=M042M5

• • • We do not use the following data for averages, fits, limits, etc. • • •

 **$f_2(2150)$  WIDTH**

NODE=M042210

 **$f_2(2150)$  WIDTH, COMBINED MODES (MeV)**

**$152 \pm 30$  OUR AVERAGE** Includes data from the datablock that follows this one. Error includes scale factor of 1.4. See the ideogram below.


NODE=M042W  
 NODE=M042W

• • • We do not use the following data for averages, fits, limits, etc. • • •

$182 \pm 11$       80k    <sup>12</sup> UMAN      06    E835     $5.2 \bar{p} p \rightarrow \eta \eta \pi^0$

<sup>12</sup>Statistical error only.

NODE=M042W;LINKAGE=ST

 **$\eta\eta$  MODE**

VALUE (MeV)      DOCUMENT ID      TECN      COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M042W3  
 NODE=M042W3

**$152 \pm 30$  OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

$280 \pm 70$       BARBERIS      00E       $450 p p \rightarrow p_f \eta \eta p_s$

$150 \pm 35$       PROKOSHKIN      95D       $300 \pi^- N \rightarrow \pi^- N 2\eta,$   
 $450 p p \rightarrow p p 2\eta$

$130 \pm 30$       SINGOVSKI      94       $450 p p \rightarrow p p 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •


$310 \pm 50$       <sup>13</sup> ABELE      99B      CBAR     $1.94 \bar{p} p \rightarrow \pi^0 \eta \eta$

$203 \pm 10$       <sup>14</sup> ARMSTRONG      93C      E760     $\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$

<sup>13</sup>Spin not determined.

<sup>14</sup>No  $J^{PC}$  determination.

NODE=M042W3;LINKAGE=K3  
 NODE=M042W3;LINKAGE=A

 **$\eta\pi\pi$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

250±25±45	<sup>15</sup> ADOMEIT	96	CBAR	0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
	<sup>15</sup> ANISOVICH	00E			recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

NODE=M042W4  
NODE=M042W4

NODE=M042W4;LINKAGE=AD

 **$\bar{p}p \rightarrow \pi\pi$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**250 OUR ESTIMATE**

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 70	<sup>16</sup> OAKDEN	94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	<sup>17</sup> MARTIN	80B	RVUE	
~ 250	<sup>17</sup> MARTIN	80C	RVUE	
~ 250	<sup>18</sup> DULUDE	78B	OSPK	1-2 $\bar{p}p \rightarrow \pi^0\pi^0$

NODE=M042W1  
NODE=M042W1  
→ UNCHECKED ←

NODE=M042W1;LINKAGE=CC

<sup>16</sup> See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>17</sup>  $I(J^P) = 0(2^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^-\pi^+$  and  $\pi^0\pi^0$ .

<sup>18</sup>  $I^G(J^P) = 0^+(2^+)$  from partial-wave amplitude analysis.

NODE=M042W1;LINKAGE=P

NODE=M042W1;LINKAGE=L

**S-CHANNEL  $\bar{p}p$ ,  $\bar{N}N$  or  $\bar{K}K$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

56 <sup>+31</sup> <sub>-16</sub>	<sup>19</sup> EVANGELIS...	97	SPEC	0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$	
135±75	<sup>20,21</sup> COUPLAND	77	CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
98± 8	<sup>21</sup> ALSPECTOR	73	CNTR		$\bar{p}p$ S channel

NODE=M042W2  
NODE=M042W2

<sup>19</sup> Isospin 0 and 2 not separated.

<sup>20</sup> From a fit to the total elastic cross section.

<sup>21</sup> Isospins 0 and 1 not separated.

NODE=M042W2;LINKAGE=F

NODE=M042W2;LINKAGE=E

NODE=M042W2;LINKAGE=I

 **$K\bar{K}$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

91±62	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
150±30	ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
270±50	BARBERIS	99	OMEG 450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

NODE=M042W5  
NODE=M042W5

**$f_2(2150)$  DECAY MODES**

NODE=M042215;NODE=M042

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	
$\Gamma_2$ $\eta\eta$	seen
$\Gamma_3$ $K\bar{K}$	seen
$\Gamma_4$ $f_2(1270)\eta$	seen
$\Gamma_5$ $a_2(1320)\pi$	seen
$\Gamma_6$ $\rho\bar{\rho}$	seen

DESIG=1

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=6

 **$f_2(2150)$  BRANCHING RATIOS**

NODE=M042220

 **$\Gamma(K\bar{K})/\Gamma(\eta\eta)$   $\Gamma_3/\Gamma_2$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>1.28±0.23</b>		BARBERIS	00E	450 $p\bar{p} \rightarrow p_f \eta \eta p_s$

NODE=M042R1

NODE=M042R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.1	95	<sup>22</sup> PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$ , 450 $p\bar{p} \rightarrow p\rho 2\eta$
------	----	------------------------------	------	---

<sup>22</sup> Using data from ARMSTRONG 89D.

NODE=M042R1;LINKAGE=A

 **$\Gamma(\pi\pi)/\Gamma(\eta\eta)$   $\Gamma_1/\Gamma_2$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.33	95	<sup>23</sup> PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$ , 450 $p\bar{p} \rightarrow p\rho 2\eta$

NODE=M042R2

NODE=M042R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>23</sup> Derived from a  $\pi^0\pi^0/\eta\eta$  limit.

NODE=M042R2;LINKAGE=A

 **$\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$   $\Gamma_4/\Gamma_5$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.79±0.11</b>	<sup>24</sup> ADOMEIT 96	CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M042R3

NODE=M042R3

<sup>24</sup> Using  $B(a_2(1320) \rightarrow \eta\pi) = 0.145$ 

NODE=M042R3;LINKAGE=A

 **$\Gamma(\rho\bar{\rho})/\Gamma_{total}$   $\Gamma_6/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	73	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma\rho\bar{\rho}$

NODE=M042R04

NODE=M042R04

 **$f_2(2150)$  REFERENCES**

NODE=M042

ALEXANDER 10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK... 06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
	Translated from YAF 69 515.			
ABLIKIM 04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
ANISOVICH 00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
BARBERIS 00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ABELE 99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46904
BARBERIS 99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
EVANGELIS... 97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)	REFID=45687
MARTIN 97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)	REFID=45685
ADOMEIT 96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
PROKOSHKIN 95D	PD 40 495	Y.D. Prokoshkin	(SERP) IGJPC	REFID=44647
	Translated from DANS 344 469.			
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
SINGOVSKI 94	NC A107 1911	A.V. Singovsky	(SERP)	REFID=44648
ARMSTRONG 93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
ARMSTRONG 89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41010
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP	REFID=21838
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP	REFID=21837
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)	REFID=21733
DULUDE 78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP	REFID=21850
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)	REFID=21830
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)	REFID=21813

REFID=53525

REFID=51063

REFID=51191

REFID=50174

REFID=47945

REFID=47961

REFID=46904

REFID=46921

REFID=45687

REFID=45685

REFID=45202

REFID=45212

REFID=44647

REFID=44103

REFID=45210

REFID=44648

REFID=43587

REFID=41010

REFID=21838

REFID=21837

REFID=21733

REFID=21850

REFID=21830

REFID=21813

**$\rho(2150)$** 

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called  $T_1(2190)$ . See the review on "Spectroscopy of Light Meson Resonances."

NODE=M032

NODE=M032

 **$\rho(2150)$  MASS**

NODE=M032205

 **$e^+e^-$  PRODUCED**

NODE=M032M3

NODE=M032M3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2034 $\pm 13 \pm 9$		<sup>1</sup> ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\pi^0$
2111 $\pm 43 \pm 25$		<sup>2</sup> ABLIKIM	21X BES3	$e^+e^- \rightarrow \eta'\pi^+\pi^-$
2255 $\begin{smallmatrix} +17 & +50 \\ -18 & -41 \end{smallmatrix}$	1.8k	<sup>3</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+K^-\eta$
2201 $\pm 19$		<sup>4</sup> LEES	20 BABR	$e^+e^- \rightarrow K^+K^-\gamma$
2227 $\pm 9 \pm 9$		<sup>5</sup> LEES	20 RVUE	$e^+e^- \rightarrow K^+K^-$
2039 $\pm 8 \begin{smallmatrix} +36 \\ -18 \end{smallmatrix}$		<sup>6</sup> ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+K^-\pi^0$
2239.2 $\pm 7.1 \pm 11.3$		<sup>7</sup> ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
2254 $\pm 22$		<sup>8</sup> LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
2150 $\pm 40 \pm 50$		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
1990 $\pm 80$		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$
2153 $\pm 37$		BIAGINI	91 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$ , $K^+K^-$
2110 $\pm 50$		<sup>9</sup> CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-)$ , $2(\pi^+\pi^-\pi^0)$

OCCUR=2

OCCUR=2

<sup>1</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from  $\rho(770)$ ,  $\rho(1450)$  and  $\rho(1700)$ . Could be another state.

NODE=M032M3;LINKAGE=G

<sup>2</sup> From a Breit-Wigner fit to the Born cross section, including an  $s$ -dependent continuum amplitude.

NODE=M032M3;LINKAGE=H

<sup>3</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+K^-\eta = (21.7 \pm 1.9 \begin{smallmatrix} +7.7 \\ -8.3 \end{smallmatrix}) \times 10^{-6}$ .

NODE=M032M3;LINKAGE=F

<sup>4</sup> From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is  $3.5\sigma$ .

NODE=M032M3;LINKAGE=C

<sup>5</sup> From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.

NODE=M032M3;LINKAGE=D

<sup>6</sup> Could also be another state. Seen in  $J/\psi$  decay with branching ratio  $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (6.7 \pm 1.1 \begin{smallmatrix} +2.2 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$ .

NODE=M032M3;LINKAGE=B

<sup>7</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .

NODE=M032M3;LINKAGE=E

<sup>8</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.

NODE=M032M3;LINKAGE=LE

<sup>9</sup> Includes ATKINSON 85.

NODE=M032M3;LINKAGE=A

 **$\bar{p}p \rightarrow \pi\pi$** 

NODE=M032M1

NODE=M032M1

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$\sim 2191$	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
$\sim 2070$	<sup>1</sup> OAKDEN	94 RVUE	$0.36-1.55 \bar{p}p \rightarrow \pi\pi$
$\sim 2170$	<sup>2</sup> MARTIN	80B RVUE	
$\sim 2100$	<sup>2</sup> MARTIN	80C RVUE	

<sup>1</sup> See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J=3$  to be important but not significantly resonant.

NODE=M032M1;LINKAGE=CC

<sup>2</sup>  $I(J^P) = 1(1^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^-\pi^+$  and  $\pi^0\pi^0$ .

NODE=M032M;LINKAGE=P

**S-CHANNEL  $\bar{N}N$** 

NODE=M032M2

NODE=M032M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2110 $\pm 35$	<sup>1</sup> ANISOVICH	02 SPEC	$0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
$\sim 2190$	<sup>2</sup> CUTTS	78B CNTR	$0.97-3 \bar{p}p \rightarrow \bar{N}N$
2155 $\pm 15$	<sup>2,3</sup> COUPLAND	77 CNTR	$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
2193 $\pm 2$	<sup>2,4</sup> ALSPECTOR	73 CNTR	$\bar{p}p$ S channel
2190 $\pm 10$	<sup>5</sup> ABRAMS	70 CNTR	S channel $\bar{p}N$



- <sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.  
<sup>2</sup> Isospins 0 and 1 not separated.  
<sup>3</sup> From a fit to the total elastic cross section.  
<sup>4</sup> Referred to as  $T$  or  $T$  region by ALSPECTOR 73.  
<sup>5</sup> Seen as bump in  $I = 1$  state. See also COOPER 68. PEASLEE 75 confirm  $\bar{p}p$  results of ABRAMS 70, no narrow structure.

NODE=M032M;LINKAGE=AY

NODE=M032M;LINKAGE=I  
 NODE=M032M;LINKAGE=E  
 NODE=M032M;LINKAGE=M  
 NODE=M032M;LINKAGE=B

 **$\pi^- p \rightarrow \omega \pi^0 n$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2140 ± 30	ALDE	95	GAM2 38 $\pi^- p \rightarrow \omega \pi^0 n$
2170 ± 30	ALDE	92C	GAM4 100 $\pi^- p \rightarrow \omega \pi^0 n$

NODE=M032M4  
 NODE=M032M4

 **$\rho(2150)$  WIDTH**

NODE=M032210

 **$e^+ e^-$  PRODUCED**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
234 ± 30 ± 25		<sup>1</sup> ABLIKIM	21A BES3	$e^+ e^- \rightarrow \omega \pi^0$
135 ± 34 ± 30		<sup>2</sup> ABLIKIM	21X BES3	$e^+ e^- \rightarrow \eta' \pi^+ \pi^-$
460 $\begin{smallmatrix} +54 \\ -48 \end{smallmatrix}$ $\begin{smallmatrix} +160 \\ -90 \end{smallmatrix}$	1.8k	<sup>3</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
70 ± 38		<sup>4</sup> LEES	20 BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
127 ± 14 ± 4		<sup>5</sup> LEES	20 RVUE	$e^+ e^- \rightarrow K^+ K^-$
196 ± 23 $\begin{smallmatrix} +25 \\ -27 \end{smallmatrix}$		<sup>6</sup> ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
139.8 ± 12.3 ± 20.6		<sup>7</sup> ABLIKIM	19L BES3	$e^+ e^- \rightarrow K^+ K^-$
109 ± 76		<sup>8</sup> LEES	12G BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
350 ± 40 ± 50		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
310 ± 140		AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow \eta' \pi^+ \pi^- \gamma$
389 ± 79		BIAGINI	91 RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- , K^+ K^-$
410 ± 100		<sup>9</sup> CLEGG	90 RVUE	$e^+ e^- \rightarrow 3(\pi^+ \pi^-), 2(\pi^+ \pi^- \pi^0)$

NODE=M032W3  
 NODE=M032W3

OCCUR=2

OCCUR=2

- <sup>1</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from  $\rho(770)$ ,  $\rho(1450)$  and  $\rho(1700)$ . Could be another state.  
<sup>2</sup> From a Breit-Wigner fit to the Born cross section, including an  $s$ -dependent continuum amplitude.  
<sup>3</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X \eta \rightarrow K^+ K^- \eta = (21.7 \pm 1.9 \begin{smallmatrix} +7.7 \\ -8.3 \end{smallmatrix}) \times 10^{-6}$ .  
<sup>4</sup> From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is  $3.5 \sigma$ .  
<sup>5</sup> From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.  
<sup>6</sup> Could also be another state. Seen in  $J/\psi$  decay with branching ratio  $J/\psi \rightarrow X \pi^0 \rightarrow K^+ K^- \pi^0 = (6.7 \pm 1.1 \begin{smallmatrix} +2.2 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$ .  
<sup>7</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .  
<sup>8</sup> Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the  $\rho(1450)$ ,  $\rho(1700)$ , and  $\rho(2150)$  resonances as free parameters of the fit.  
<sup>9</sup> Includes ATKINSON 85.

NODE=M032W3;LINKAGE=G

NODE=M032W3;LINKAGE=H

NODE=M032W3;LINKAGE=F

NODE=M032W3;LINKAGE=C

NODE=M032W3;LINKAGE=D

NODE=M032W3;LINKAGE=B

NODE=M032W3;LINKAGE=E

NODE=M032W3;LINKAGE=LE

NODE=M032W3;LINKAGE=A

 **$\bar{p}p \rightarrow \pi\pi$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 296	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 40	<sup>1</sup> OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	<sup>2</sup> MARTIN	80B	RVUE
~ 200	<sup>2</sup> MARTIN	80C	RVUE

NODE=M032W1  
 NODE=M032W1

- <sup>1</sup> See however KLOET 96 who fit  $\pi^+ \pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.  
<sup>2</sup>  $I(J^P) = 1(1^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .

NODE=M032W1;LINKAGE=CC

NODE=M032W;LINKAGE=P

**S-CHANNEL  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
230±50	<sup>1</sup> ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
135±75	<sup>2,3</sup> COUPLAND 77	CNTR	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98±8	<sup>3</sup> ALSPECTOR 73	CNTR	$\bar{p}p$ S channel
~ 85	<sup>4</sup> ABRAMS 70	CNTR	S channel $\bar{p}N$

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

<sup>2</sup> From a fit to the total elastic cross section.

<sup>3</sup> Isospins 0 and 1 not separated.

<sup>4</sup> Seen as bump in  $l = 1$  state. See also COOPER 68. PEASLEE 75 confirm  $\bar{p}p$  results of ABRAMS 70, no narrow structure.

NODE=M032W2  
NODE=M032W2

NODE=M032W;LINKAGE=AY

NODE=M032W;LINKAGE=E  
NODE=M032W;LINKAGE=I  
NODE=M032W;LINKAGE=B

 **$\pi^- p \rightarrow \omega\pi^0 n$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
320±70	ALDE 95	GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
~ 300	ALDE 92C	GAM4	100 $\pi^- p \rightarrow \omega\pi^0 n$

NODE=M032W4  
NODE=M032W4

 **$\rho(2150)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+e^-$	
$\Gamma_2$ $\pi^+\pi^-$	seen
$\Gamma_3$ $K^+K^-$	seen
$\Gamma_4$ $3(\pi^+\pi^-)$	seen
$\Gamma_5$ $2(\pi^+\pi^-\pi^0)$	seen
$\Gamma_6$ $\eta'\pi^+\pi^-$	seen
$\Gamma_7$ $f_1(1285)\pi^+\pi^-$	seen
$\Gamma_8$ $\omega\pi^0$	seen
$\Gamma_9$ $\omega\pi^0\eta$	seen
$\Gamma_{10}$ $p\bar{p}$	

NODE=M032215;NODE=M032

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=3;OUR EVAL;→ UNCHECKED ←  
DESIG=4;OUR EVAL;→ UNCHECKED ←  
DESIG=5;OUR EVAL;→ UNCHECKED ←  
DESIG=6;OUR EVAL;→ UNCHECKED ←  
DESIG=7;OUR EVAL;→ UNCHECKED ←  
DESIG=8;OUR EVAL;→ UNCHECKED ←  
DESIG=9;OUR EVAL;→ UNCHECKED ←  
DESIG=10

 **$\rho(2150) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_8\Gamma_1/\Gamma$
34±11±16	ABLIKIM 21A	BES3	$e^+e^- \rightarrow \omega\pi^0$	

NODE=M032220

NODE=M032R00  
NODE=M032R00

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_6\Gamma_1/\Gamma$
23.3±5.3±3.3	<sup>1</sup> ABLIKIM 21X	BES3	$e^+e^- \rightarrow \eta'\pi^+\pi^-$	

NODE=M032R02  
NODE=M032R02

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> From a Breit-Wigner fit to the Born cross section interfering constructively with the continuum. For destructive interference the value is  $0.64 \pm 0.49 \pm 0.42$  eV.

NODE=M032R02;LINKAGE=A

 **$\rho(2150) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$** 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma \times \Gamma_1/\Gamma$
3.1±0.6±0.5	<sup>1</sup> AUBERT 07AU BABR	10.6	$e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$	

NODE=M032230

NODE=M032G01  
NODE=M032G01

<sup>1</sup> Calculated by us from the reported value of cross section at the peak.

NODE=M032G01;LINKAGE=AU

VALUE (units $10^{-8}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma \times \Gamma_1/\Gamma$
4.9±1.9	<sup>1</sup> AUBERT 07AU BABR	10.6	$e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$	

NODE=M032G02  
NODE=M032G02

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Calculated by us from the reported value of cross section at the peak.

NODE=M032G02;LINKAGE=AU

$\rho(2150)$  REFERENCES

ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21X	PR D103 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	20	PR D101 012011	J.P. Lees <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
BIAGINI	91	NC 104A 363	M.E. Biagini <i>et al.</i>	(FRAS, PRAG)
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	

NODE=M032

REFID=61028  
REFID=61231  
REFID=60256  
REFID=60211  
REFID=59909  
REFID=59612  
REFID=55404  
REFID=54299  
REFID=52049  
REFID=48828  
REFID=48327  
REFID=48349  
REFID=47950  
REFID=45212  
REFID=44371  
REFID=44103  
REFID=45210  
REFID=41859  
REFID=41894  
REFID=41355  
REFID=22000  
REFID=21838  
REFID=21837  
REFID=21733  
REFID=21830  
REFID=21824  
REFID=21813  
REFID=21807  
REFID=21805  
REFID=48054

 $\phi(2170)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M103

NODE=M103

 $\phi(2170)$  MASS

NODE=M103M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2163 ± 7 OUR AVERAGE</b>		Error includes scale factor of 1.1. [2162 ± 7 MeV OUR 2022 AVERAGE Scale factor = 1.1]		
2190 ± 19 ± 37		<sup>1</sup> ABLIKIM	22L BES3	2.0–3.08 $e^+e^- \rightarrow K^+K^-\pi^0$
2176 ± 24 ± 3		<sup>2</sup> ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$
2163.5 ± 6.2 ± 3.0		<sup>3</sup> ABLIKIM	21T BES3	$e^+e^- \rightarrow \phi\eta$
2177.5 ± 4.8 ± 19.5		<sup>4</sup> ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$
2126.5 ± 16.8 ± 12.4		<sup>5</sup> ABLIKIM	20S BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2273.7 ± 5.7 ± 19.3		<sup>6</sup> ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$
2135 ± 8 ± 9	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
2239.2 ± 7.1 ± 11.3		<sup>7</sup> ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
2200 ± 6 ± 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
2180 ± 8 ± 8		<sup>8,9</sup> LEES	12F BABR	10.6 $e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
2079 ± 13 ± 79 -28	4.8k	<sup>10</sup> SHEN	09 BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2186 ± 10 ± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
2125 ± 22 ± 10	483	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
2192 ± 14	116	<sup>11</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2169 ± 20	149	<sup>11</sup> AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
2175 ± 10 ± 15	201	<sup>9,12</sup> AUBERT, BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

NODE=M103M

NEW

OCCUR=2

<sup>1</sup> By a simultaneous fit of the  $K_2^*(1430)^+K^-$  and  $K^*(892)^+K^-$  intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

<sup>2</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from  $\omega(1420)$  and  $\omega(1650)/\phi(1680)$ .

<sup>3</sup> From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ( $\phi(1680)$  and  $\phi(2170)$ ) and a nonresonant term.

<sup>4</sup> From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

NODE=M103M;LINKAGE=I

NODE=M103M;LINKAGE=F

NODE=M103M;LINKAGE=G

NODE=M103M;LINKAGE=D

- <sup>5</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.
- <sup>6</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to  $\rho(2150)$ .
- <sup>7</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .
- <sup>8</sup> Fit includes interference with the  $\phi(1680)$ .
- <sup>9</sup> From the  $\phi f_0(980)$  component.
- <sup>10</sup> From a fit with two incoherent Breit-Wigners.
- <sup>11</sup> From the  $K^+ K^- f_0(980)$  component.
- <sup>12</sup> Superseded by LEES 12F.

NODE=M103M;LINKAGE=E

NODE=M103M;LINKAGE=H

NODE=M103M;LINKAGE=C

NODE=M103M;LINKAGE=A

NODE=M103M;LINKAGE=AB

NODE=M103M;LINKAGE=SH

NODE=M103M;LINKAGE=AU

NODE=M103M;LINKAGE=B

 **$\phi(2170)$  WIDTH**

NODE=M103W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M103W

<b>103</b>	<b><math>^{+28}_{-21}</math></b>	<b>OUR AVERAGE</b>		Error includes scale factor of 2.2. See the ideogram below.
------------	----------------------------------	--------------------	--	---

NEW

[100	$^{+31}_{-23}$	MeV OUR 2022 AVERAGE		Scale factor = 2.5]
------	----------------	----------------------	--	---------------------

191	$\pm 28 \pm 60$	<sup>1</sup> ABLIKIM	22L BES3	$2.0-3.08 e^+ e^- \rightarrow K^+ K^- \pi^0$
89	$\pm 50 \pm 5$	<sup>2</sup> ABLIKIM	21A BES3	$e^+ e^- \rightarrow \omega \eta$
31.1	$^{+21.1}_{-11.6} \pm 1.1$	<sup>3</sup> ABLIKIM	21T BES3	$e^+ e^- \rightarrow \phi \eta$
149.0	$\pm 15.6 \pm 8.9$	<sup>4</sup> ABLIKIM	20M BES3	$e^+ e^- \rightarrow \eta' \phi$
106.9	$\pm 32.1 \pm 28.1$	<sup>5</sup> ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

86	$\pm 44 \pm 51$	<sup>6</sup> ABLIKIM	21AP BES3	$e^+ e^- \rightarrow K_S^0 K_L^0$
104	$\pm 24 \pm 12$	95 ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta \phi f_0(980)$
139.8	$\pm 12.3 \pm 20.6$	<sup>7</sup> ABLIKIM	19L BES3	$e^+ e^- \rightarrow K^+ K^-$
104	$\pm 15 \pm 15$	471 ABLIKIM	15H BES3	$J/\psi \rightarrow \eta \phi \pi^+ \pi^-$
77	$\pm 15 \pm 10$	<sup>8,9</sup> LEES	12F BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
192	$\pm 23 \pm 25$	<sup>10</sup> SHEN	09 BELL	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
65	$\pm 23 \pm 17$	52 ABLIKIM	08F BES	$J/\psi \rightarrow \eta \phi f_0(980)$
61	$\pm 50 \pm 13$	483 AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow \phi \eta \gamma$
71	$\pm 21$	116 <sup>11</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
102	$\pm 27$	149 <sup>11</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$
58	$\pm 16 \pm 20$	201 <sup>9,12</sup> AUBERT, BE	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi \pi \gamma$

OCCUR=2

- <sup>1</sup> By a simultaneous fit of the  $K_2^*(1430)^+ K^-$  and  $K^*(892)^+ K^-$  intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103W;LINKAGE=I

- <sup>2</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from  $\omega(1420)$  and  $\omega(1650)/\phi(1680)$ .

NODE=M103W;LINKAGE=F

- <sup>3</sup> From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ( $\phi(1680)$  and  $\phi(2170)$ ) and a nonresonant term.

NODE=M103W;LINKAGE=G

- <sup>4</sup> From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

NODE=M103W;LINKAGE=D

- <sup>5</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103W;LINKAGE=E

- <sup>6</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to  $\rho(2150)$ .

NODE=M103W;LINKAGE=H

- <sup>7</sup> The observed structure can be due to both the  $\phi(2170)$  and  $\rho(2150)$ .

NODE=M103W;LINKAGE=C

- <sup>8</sup> Fit includes interference with the  $\phi(1680)$ .

NODE=M103W;LINKAGE=A

- <sup>9</sup> From the  $\phi f_0(980)$  component.

NODE=M103W;LINKAGE=AB

- <sup>10</sup> From a fit with two incoherent Breit-Wigners.


NODE=M103W;LINKAGE=SH

- <sup>11</sup> From the  $K^+ K^- f_0(980)$  component.

NODE=M103W;LINKAGE=AU

- <sup>12</sup> Superseded by LEES 12F.

NODE=M103W;LINKAGE=B



**phi(2170) DECAY MODES**

NODE=M103215;NODE=M103

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	seen
$\Gamma_2$ $\phi\eta$	seen
$\Gamma_3$ $\omega\eta$	seen
$\Gamma_4$ $\phi\eta'$	seen
$\Gamma_5$ $\phi\pi\pi$	
$\Gamma_6$ $\phi f_0(980)$	seen
$\Gamma_7$ $K_S^0 K_L^0$	
$\Gamma_8$ $K^+ K^- \pi^+ \pi^-$	
$\Gamma_9$ $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
$\Gamma_{10}$ $K^+ K^- \pi^0 \pi^0$	
$\Gamma_{11}$ $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
$\Gamma_{12}$ $K^{*0} K^\pm \pi^\mp$	not seen
$\Gamma_{13}$ $K^*(892)^0 \bar{K}^*(892)^0$	not seen
$\Gamma_{14}$ $K^*(892)^+ K^*(892)^-$	
$\Gamma_{15}$ $K^*(892)^+ K^- + c.c.$	
$\Gamma_{16}$ $K(1460)^+ K^- + c.c.$	
$\Gamma_{17}$ $K_1(1270)^+ K^- + c.c.$	
$\Gamma_{18}$ $K_1(1400)^+ K^- + c.c.$	
$\Gamma_{19}$ $K_2^*(1430)^+ K^- + c.c.$	

DESIG=1;OUR EVAL;→ UNCHECKED ←  
 DESIG=5;OUR EVAL;→ UNCHECKED ←  
 DESIG=16;OUR EVAL;→ UNCHECKED ←  
 DESIG=11;OUR EVAL;→ UNCHECKED ←  
 DESIG=9  
 DESIG=2;OUR EVAL;→ UNCHECKED ←  
 DESIG=17  
 DESIG=3  
 DESIG=6  
 DESIG=4  
 DESIG=7  
 DESIG=8  
 DESIG=10  
 DESIG=15  
 DESIG=19  
 DESIG=12  
 DESIG=14  
 DESIG=13  
 DESIG=18

**phi(2170)  $\Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$**

NODE=M103230

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$	$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV) CL% EVTS DOCUMENT ID TECN COMMENT	
0.17 90 1 ZHU 23 BELL $e^+ e^- \rightarrow \gamma(nS) \rightarrow \phi\eta\gamma$	
$0.24^{+0.12}_{-0.07}$ 2 ABLIKIM 21T BES3 $e^+ e^- \rightarrow \phi\eta$	
$1.7 \pm 0.7 \pm 1.3$ 483 AUBERT 08S BABR $10.6 e^+ e^- \rightarrow \phi\eta\gamma$	

NODE=M103G2  
 NODE=M103G2

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> From a solution of the fit using a vector meson dominance model with contributions from  $\phi(1680)$ ,  $\phi(2170)$  and non resonant contribution with mass and width of  $\phi(2170)$  fixed at 2163.5 MeV and 31.1 MeV respectively. Four solutions are found with equal fit quality giving 0.17 eV (solution I and II) and 18.6 eV (III and IV) at 90% CL.

NODE=M103G2;LINKAGE=B

<sup>2</sup> From a solution of the fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ( $\phi(1680)$  and  $\phi(2170)$ ) and a nonresonant term. The other solution gives  $10.11^{+3.87}_{-3.13}$  eV.

NODE=M103G2;LINKAGE=A

$$\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>0.43±0.15±0.04</b>	<sup>1</sup> ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$

<sup>1</sup> For constructive interference with  $\omega(1420)$  and  $\omega(1650)/\phi(1680)$ . For destructive interference:  $1.25 \pm 0.48 \pm 0.18$  eV.

NODE=M103R09  
NODE=M103R09

NODE=M103R09;LINKAGE=A

$$\Gamma(\phi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>7.1±0.7±0.7</b>	<sup>1</sup> ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$

<sup>1</sup> From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

NODE=M103R00  
NODE=M103R00

NODE=M103R00;LINKAGE=A

$$\Gamma(\phi f_0(980)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_6\Gamma_1/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3±0.3±0.3</b>		<sup>1,2</sup> LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.5±0.8±0.4	201	<sup>2,3</sup> AUBERT,BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$
-------------	-----	--------------------------	----------	--

<sup>1</sup> From a fit with constructive interference with the  $\phi(1680)$ . In a fit with destructive interference, the value is larger by a factor of 12.

<sup>2</sup> From the  $\phi f_0(980)$  component.

<sup>3</sup> Superseded by LEES 12F.

NODE=M103G1  
NODE=M103G1

NODE=M103G1;LINKAGE=A

NODE=M103G1;LINKAGE=AB  
NODE=M103G1;LINKAGE=B

$$\Gamma(K_S^0 K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
0.9±0.6±0.7	<sup>1</sup> ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$

<sup>1</sup> From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to  $\rho(2150)$ .

NODE=M103R10  
NODE=M103R10

NODE=M103R10;LINKAGE=A

$$\Gamma(K^*(892)^+ K^*(892)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.9</b>	90	<sup>1</sup> ABLIKIM	20s BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103R08  
NODE=M103R08

NODE=M103R08;LINKAGE=A

$$\Gamma(K^*(892)^+ K^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.0±0.3	<sup>1</sup> ABLIKIM	22L BES3	$2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$

<sup>1</sup> From a solution of a simultaneous fit of the  $K_2^*(1430)^+ K^-$  and  $K^*(892)^+ K^-$  intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. The other solution gives  $7.1 \pm 0.9$  eV. Significance  $3.7 \sigma$ .

NODE=M103R11  
NODE=M103R11

NODE=M103R11;LINKAGE=A

$$\Gamma(K(1460)^+ K^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
3.0±3.8	<sup>1</sup> ABLIKIM	20s BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103R05  
NODE=M103R05

NODE=M103R05;LINKAGE=A

$$\Gamma(K_1(1270)^+ K^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;12.5</b>	90	<sup>1</sup> ABLIKIM	20s BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives an upper limit value of 297.6 eV.

NODE=M103R06  
NODE=M103R06

NODE=M103R06;LINKAGE=A

$$\Gamma(K_1(1400)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{18} \Gamma_1 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M103R07  
NODE=M103R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.7 ± 3.3      <sup>1</sup> ABLIKIM      20S    BES3    e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>π<sup>0</sup>π<sup>0</sup>

<sup>1</sup> By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives a value of 98.8 ± 7.8 eV.

NODE=M103R07;LINKAGE=A

$$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{19} \Gamma_1 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M103R12  
NODE=M103R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.6 ± 2.4      <sup>1</sup> ABLIKIM      22L    BES3    2.0-3.08 e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>π<sup>0</sup>

<sup>1</sup> From a solution of a simultaneous fit of the K<sub>2</sub><sup>\*</sup>(1430)<sup>+</sup>K<sup>-</sup> and K<sup>\*</sup>(892)<sup>+</sup>K<sup>-</sup> intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. The other solution gives 161.1 ± 20.6 eV.

NODE=M103R12;LINKAGE=A

### φ(2170) Γ(i)Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sup>2</sup>(total)

NODE=M103220

$$\Gamma(\phi\pi\pi) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_5 / \Gamma \times \Gamma_1 / \Gamma$$

VALUE (units 10 <sup>-7</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

NODE=M103G01  
NODE=M103G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.65 ± 0.15 ± 0.18    4.8k      <sup>1</sup> SHEN      09    BELL    10.6 e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>π<sup>+</sup>π<sup>-</sup>γ

<sup>1</sup> Multiplied by 3/2 to take into account the φπ<sup>0</sup>π<sup>0</sup> mode. Using B(φ → K<sup>+</sup>K<sup>-</sup>) = (49.2 ± 0.6)%.

NODE=M103G01;LINKAGE=SH

### φ(2170) BRANCHING RATIOS

NODE=M103225

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M103R01  
NODE=M103R01

seen      AUBERT      07AK    BABR    10.6 e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>π<sup>+</sup>π<sup>-</sup>γ

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{11} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M103R02  
NODE=M103R02

seen      AUBERT      07AK    BABR    10.6 e<sup>+</sup>e<sup>-</sup> → K<sup>+</sup>K<sup>-</sup>π<sup>0</sup>π<sup>0</sup>γ

$$\Gamma(K^{*0} K^\pm \pi^\mp) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M103R03  
NODE=M103R03

not seen      AUBERT      07AK    BABR    10.6 GeV e<sup>+</sup>e<sup>-</sup>

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M103R04  
NODE=M103R04

not seen      ABLIKIM      10C    BES2    J/ψ → ηK<sup>+</sup>π<sup>-</sup>K<sup>-</sup>π<sup>+</sup>

### φ(2170) REFERENCES

NODE=M103

ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)	REFID=61911
ABLIKIM	22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61649
ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61028
ABLIKIM	21AP	PR D104 092014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61448
ABLIKIM	21T	PR D104 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61154
ABLIKIM	20M	PR D102 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60337
ABLIKIM	20S	PRL 124 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60542
ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59605
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59612
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56773
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT_BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511

$f_0(2200)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in  $K_S^0 K_S^0$  (AUGUSTIN 88),  $K^+ K^-$  (ABLIKIM 05Q) and  $\eta\eta$  (BINON 05) system. Not seen in  $\Upsilon(1S)$  radiative decays (BARU 89).

NODE=M112

NODE=M112

 $f_0(2200)$  MASS

NODE=M112M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2187 ± 14 OUR AVERAGE</b>				
$2170 \pm 20^{+10}_{-15}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
$2197 \pm 17$		<sup>1</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2200 \pm 25$		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$2206 \pm 12 \pm 8$	381	<sup>2,3</sup> DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
$2188 \pm 17 \pm 16$	203	<sup>2,3</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
$2210 \pm 50$		<sup>4</sup> BINON	05 GAMS	$33 \pi^- p \rightarrow \eta\eta n$
~ 2122		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2321		HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$

NODE=M112M

OCCUR=2

OCCUR=2

<sup>1</sup> Cannot determine spin to be 0.<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>3</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 238$  MeV.<sup>4</sup> First solution, PWA is ambiguous.

NODE=M112M;LINKAGE=A  
 NODE=M112M;LINKAGE=B  
 NODE=M112M;LINKAGE=C  
 NODE=M112M;LINKAGE=BI

 $f_0(2200)$  WIDTH

NODE=M112W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>207 ± 40 OUR AVERAGE</b>			
$220 \pm 60^{+40}_{-45}$	ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
$201 \pm 51$	<sup>5</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$150 \pm 30$	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$380 \pm 90$	<sup>6</sup> BINON	05 GAMS	$33 \pi^- p \rightarrow \eta\eta n$
~ 273	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 223	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$

NODE=M112W

OCCUR=2

<sup>5</sup> Cannot determine spin to be 0.<sup>6</sup> First solution, PWA is ambiguous.

NODE=M112W;LINKAGE=A  
 NODE=M112W;LINKAGE=BI

 $f_0(2200)$  REFERENCES

NODE=M112

SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
		Translated from YAF 68 998.			
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574



$f_J(2220)$ 

$$I^G(J^{PC}) = 0^+(2^{++} \text{ or } 4^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. See our mini-review in the 2004 edition of this Review, PDG 04.

NODE=M082

NODE=M082

 $f_J(2220)$  MASS

NODE=M082M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2231.1 ± 3.5 OUR AVERAGE</b>				
2235 ± 4 ± 6	74	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
2230 $^{+6}_{-7}$ ±16	46	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
2232 $^{+8}_{-7}$ ±15	23	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
2235 ± 4 ± 5	32	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
2209 $^{+17}_{-15}$ ±10		ASTON	88F LASS	$11 K^-p \rightarrow K^+K^- \Lambda$
2230 ± 20		BOLONKIN	88 SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n$
2220 ± 10	41	<sup>1</sup> ALDE	86B GA24	$38-100 \pi p \rightarrow n\eta\eta'$
2230 ± 6 ± 14	93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
2232 ± 7 ± 7	23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0 K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2223.9 ± 2.5		<sup>2</sup> VLADIMIRSK...08	SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n + m\pi^0$
2246 ± 36		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
<sup>1</sup> ALDE 86B uses data from both the GAMS-2000 and GAMS-4000 detectors.				
<sup>2</sup> $J^{PC} = 2^{++}$ . Systematic uncertainties not evaluated				

NODE=M082M

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

NODE=M082M;LINKAGE=A  
NODE=M082M;LINKAGE=VL $f_J(2220)$  WIDTH

NODE=M082W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>23 <math>^{+8}_{-7}</math> OUR AVERAGE</b>					
19 $^{+13}_{-11}$ ±12		74	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
20 $^{+20}_{-15}$ ±17		46	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
20 $^{+25}_{-16}$ ±14		23	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
15 $^{+12}_{-9}$ ± 9		32	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
60 $^{+107}_{-57}$			ASTON	88F LASS	$11 K^-p \rightarrow K^+K^- \Lambda$
80 ± 30			BOLONKIN	88 SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n$
26 $^{+20}_{-16}$ ±17		93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
18 $^{+23}_{-15}$ ±10		23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0 K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
8.6 ± 2.5			<sup>1</sup> VLADIMIRSK...08	SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n + m\pi^0$
<80	90		ALDE	87C GAM2	$38 \pi^-p \rightarrow \eta'\eta n$
<sup>1</sup> $J^{PC} = 2^{++}$ . Systematic uncertainties not evaluated					

NODE=M082W

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

NODE=M082W;LINKAGE=VL

 $f_J(2220)$  DECAY MODES

NODE=M082215;NODE=M082

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi\pi$	not seen
$\Gamma_2$ $\pi^+\pi^-$	not seen
$\Gamma_3$ $K\bar{K}$	not seen
$\Gamma_4$ $p\bar{p}$	not seen
$\Gamma_5$ $\gamma\gamma$	not seen
$\Gamma_6$ $\eta\eta'(958)$	seen
$\Gamma_7$ $\phi\phi$	not seen
$\Gamma_8$ $\eta\eta$	not seen

DESIG=5

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=1

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

$f_J(2220) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M082220

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_3\Gamma_5/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	95	1 ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$

NODE=M082G1  
NODE=M082G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.6	95	1 GODANG	97 CLE2	$\gamma\gamma \rightarrow K_S^0 K_S^0$
< 86	95	1 ALBRECHT	90G ARG	$\gamma\gamma \rightarrow K^+ K^-$
<1000	95	2 ALTHOFF	85B TASS	$\gamma\gamma, K\bar{K}\pi$

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_5/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	95	ALAM	98C CLE2	$\gamma\gamma \rightarrow \pi^+ \pi^-$

NODE=M082G3  
NODE=M082G3<sup>1</sup> Assuming  $J^P = 2^+$ .<sup>2</sup> True for  $J^P = 0^+$  and  $J^P = 2^+$ .NODE=M082G1;LINKAGE=D  
NODE=M082G1;LINKAGE=C $f_J(2220) \Gamma(i)\Gamma(\rho\bar{\rho})/\Gamma^2(\text{total})$ 

NODE=M082223

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma \times \Gamma_1/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<18	95	1 AMSLER	01 CBAR	$1.4-1.5 \rho\bar{\rho} \rightarrow \pi^0 \pi^0$

NODE=M082GG1  
NODE=M082GG1

• • • We do not use the following data for averages, fits, limits, etc. • • •

<(11-42)	99	2 HASAN	96 SPEC	$1.35-1.55 \rho\bar{\rho} \rightarrow \pi^+ \pi^-$
----------	----	---------	---------	--

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma \times \Gamma_7/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6	95	3 EVANGELIS...	98 SPEC	$1.1-2.0 \rho\bar{\rho} \rightarrow \phi\phi$

NODE=M082GG2  
NODE=M082GG2 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma \times \Gamma_8/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4	95	1 AMSLER	01 CBAR	$1.4-1.5 \rho\bar{\rho} \rightarrow \eta\eta$

NODE=M082GG3  
NODE=M082GG3<sup>1</sup> For  $J^P = 2^+$  in the mass range 2222-2240 MeV and the total width between 10 and 20 MeV.<sup>2</sup> For  $J^P = 2^+$  and  $J^P = 4^+$  in the mass range 2220-2245 MeV and the total width of 15 MeV.<sup>3</sup> For  $J^P = 2^+$ , the mass of 2235 MeV and the total width of 15 MeV.

NODE=M082GG;LINKAGE=A

NODE=M082GG;LINKAGE=B

NODE=M082GG;LINKAGE=C

 $f_J(2220) \text{ BRANCHING RATIOS}$ 

NODE=M082225

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	COMMENT
not seen	1 DOBBS 15	$J/\psi \rightarrow \gamma\pi\pi$
not seen	1 DOBBS 15	$\psi(2S) \rightarrow \gamma\pi\pi$

NODE=M082R00  
NODE=M082R00

OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M082R00;LINKAGE=A

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	COMMENT
not seen	1 DOBBS 15	$J/\psi \rightarrow \gamma K\bar{K}$
not seen	1 DOBBS 15	$\psi(2S) \rightarrow \gamma K\bar{K}$

NODE=M082R01  
NODE=M082R01

OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M082R01;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$  $\Gamma_1/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
1.0±0.5	BAI 96B	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma 2\pi, K\bar{K}$

NODE=M082R2  
NODE=M082R2 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M082R1  
NODE=M082R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		1 AUBERT	07AV BABR	$B \rightarrow \rho\bar{\rho}K^*$
not seen		WANG	05A BELL	$B^+ \rightarrow \bar{p}\rho K^+$
<3.0	95	2 EVANGELIS...	97 SPEC	$1.96-2.40 \bar{p}\rho \rightarrow K_S^0 K_S^0$
<1.1	99.7	3 BARNES	93 SPEC	$1.3-1.57 \bar{p}\rho \rightarrow K_S^0 K_S^0$
<2.6	99.7	3 BARDIN	87 CNTR	$1.3-1.5 \bar{p}\rho \rightarrow K^+ K^-$
<3.6	99.7	3 SCULLI	87 CNTR	$1.29-1.55 \bar{p}\rho \rightarrow K^+ K^-$

- <sup>1</sup> Assuming  $\Gamma < 30$  MeV.  
<sup>2</sup> Assuming  $\Gamma \sim 20$  MeV,  $J^P = 2^+$  and  $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$ .  
<sup>3</sup> Assuming  $\Gamma = 30\text{-}35$  MeV,  $J^P = 2^+$  and  $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$ .

NODE=M082R1;LINKAGE=AU  
 NODE=M082R1;LINKAGE=C  
 NODE=M082R1;LINKAGE=B

 $\Gamma(p\bar{p})/\Gamma(K\bar{K})$  $\Gamma_4/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.17±0.09</b>	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}, K\bar{K}$

NODE=M082R3  
 NODE=M082R3

 $f_J(2220)$  REFERENCES

DOBBS 15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
VLADIMIRSK... 08	PAN 71 2129	V.V. Vladimirovsky <i>et al.</i>	(ITEP)
	Translated from YAF 71 2166.		
AUBERT 07AV	PR D76 092004	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG 05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
PDG 04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ACCIARRI 01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
AMSLER 01	PL B520 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ALAM 98C	PRL 81 3328	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BAI 98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
EVANGELIS... 98	PR D57 5370	C. Evangelista <i>et al.</i>	(JETSET Collab.)
EVANGELIS... 97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
GODANG 97	PRL 79 3829	R. Godang <i>et al.</i>	(CLEO Collab.)
BAI 96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
HASAN 96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BARNES 93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
ALBRECHT 90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASTON 88F	PL B215 199	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BOLONKIN 88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ALDE 87C	SJNP 45 255	D. Alde <i>et al.</i>	
	Translated from YAF 45 405.		
BARDIN 87	PL B195 292	G. Bardin <i>et al.</i>	(SACL, FERR, CERN, PADO+)
SCULLI 87	PRL 58 1715	J. Sculli <i>et al.</i>	(NYU, BNL)
ALDE 86B	PL B177 120	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BALTRUSAIT... 86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
ALTHOFF 85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)

NODE=M082

REFID=56805  
 REFID=52681

REFID=51990  
 REFID=50651  
 REFID=49653  
 REFID=48321  
 REFID=48558  
 REFID=46326  
 REFID=46342  
 REFID=46365  
 REFID=45687  
 REFID=45760  
 REFID=44736  
 REFID=45197  
 REFID=43601  
 REFID=41374  
 REFID=40585  
 REFID=40580  
 REFID=47474

## OTHER RELATED PAPERS

DEL-AMO-SA... 100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
-------------------	----------------	----------------------------------	-----------------

REFID=53533

NODE=M115

 $\eta(2225)$ 

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Seen in  $J/\psi \rightarrow \gamma\phi\phi$ . Possibly seen in  $B \rightarrow \phi\phi K$  by LEES 11A.

NODE=M115

 $\eta(2225)$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2221^{+13}_{-10}</math> OUR AVERAGE</b>				
$2216^{+4+21}_{-5-11}$		<sup>1</sup> ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$2240^{+30+30}_{-20-20}$	196 ± 19	ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$2230 \pm 25 \pm 15$		BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$2214 \pm 20 \pm 13$		BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

NODE=M115M

NODE=M115M

••• We do not use the following data for averages, fits, limits, etc. •••

~ 2220

BISELLO 86B DM2  $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>1</sup> From a partial wave analysis of  $J/\psi \rightarrow \gamma\phi\phi$  that also finds significant signals for  $\eta(2100)$ ,  $0^-+$  phase space,  $f_0(2100)$ ,  $f_2(2010)$ ,  $f_2(2300)$ ,  $f_2(2340)$ , and a previously unseen  $0^-+$  state  $X(2500)$  ( $M = 2470^{+15+101}_{-19-23}$  MeV,  $\Gamma = 230^{+64+56}_{-35-33}$  MeV).

NODE=M115M;LINKAGE=B

 $\eta(2225)$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>185^{+40}_{-20}</math> OUR AVERAGE</b>				
$185^{+12+43}_{-14-17}$		<sup>1</sup> ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
$190 \pm 30^{+60}_{-40}$	196 ± 19	ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
$150^{+300}_{-60} \pm 60$		BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M115W

NODE=M115W

••• We do not use the following data for averages, fits, limits, etc. •••

~ 80

BISELLO 86B DM2  $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>1</sup>From a partial wave analysis of  $J/\psi \rightarrow \gamma\phi\phi$  that also finds significant signals for for  $\eta(2100)$ ,  $0^{-+}$  phase space,  $f_0(2100)$ ,  $f_2(2010)$ ,  $f_2(2300)$ ,  $f_2(2340)$ , and a previously unseen  $0^{-+}$  state  $X(2500)$  ( $M = 2470_{-19}^{+15+101}$  MeV,  $\Gamma = 230_{-35}^{+64+56}$  MeV).

NODE=M115W;LINKAGE=A

## $\eta(2225)$ REFERENCES

ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)
LEES	11A	PR D84 012001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)

NODE=M115

REFID=57512  
REFID=16595  
REFID=52255  
REFID=41354  
REFID=22101

NODE=M044

$\rho_3(2250)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

## OMITTED FROM SUMMARY TABLE

Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also  $\rho(2150)$ ,  $f_2(2150)$ ,  $f_4(2300)$ ,  $\rho_5(2350)$ .

NODE=M044

## $\rho_3(2250)$ MASS

NODE=M044205

### $\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$2248 \pm 17_{-5}^{+59}$	1.8k	1 ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
~ 2232		HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2090		2 OAKDEN	94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2250		3 MARTIN	80B	RVUE	
~ 2300		3 MARTIN	80C	RVUE	
~ 2140		4 CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2150		5 CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

NODE=M044M1  
NODE=M044M1

<sup>1</sup>Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (1.9 \pm 0.4_{-1.3}^{+0.5}) \times 10^{-6}$ .

NODE=M044M1;LINKAGE=A

<sup>2</sup>See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

NODE=M044M1;LINKAGE=CC

<sup>3</sup> $I(J^P) = 1(3^-)$  from simultaneous analysis of  $\bar{p}p \rightarrow \pi^-\pi^+$  and  $\pi^0\pi^0$ .

NODE=M044M1;LINKAGE=P

<sup>4</sup> $I = 0, 1$ .  $J^P = 3^-$  from Barrelet-zero analysis.

NODE=M044M1;LINKAGE=K

<sup>5</sup> $I(J^P) = 1(3^-)$  from amplitude analysis.

NODE=M044M1;LINKAGE=J

### S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
$2260 \pm 20$	<sup>6</sup> ANISOVICH	02	SPEC	0.6-1.9 $\bar{p}p \rightarrow \omega\pi^0$ , $\omega\eta\pi^0, \pi^+\pi^-$
~ 2190	<sup>7</sup> CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
$2155 \pm 15$	<sup>7,8</sup> COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
$2193 \pm 2$	<sup>7,9</sup> ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
$2190 \pm 10$	<sup>10</sup> ABRAMS	70	CNTR	S channel $\bar{p}N$

NODE=M044M2  
NODE=M044M2

<sup>6</sup>From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M044M;LINKAGE=AY

<sup>7</sup>Isospins 0 and 1 not separated.

NODE=M044M2;LINKAGE=I

<sup>8</sup>From a fit to the total elastic cross section.

NODE=M044M2;LINKAGE=E

<sup>9</sup>Referred to as  $T$  or  $T$  region by ALSPECTOR 73.

NODE=M044M2;LINKAGE=M

<sup>10</sup>Seen as bump in  $I = 1$  state. See also COOPER 68. PEASLEE 75 confirm  $\bar{p}p$  results of ABRAMS 70, no narrow structure.

NODE=M044M2;LINKAGE=B

### Other processes

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2290 \pm 20 \pm 30$	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

NODE=M044M3  
NODE=M044M3

## $\rho_3(2250)$ WIDTH

NODE=M044210

**$\bar{p}p \rightarrow \pi\pi$  or  $K\bar{K}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$185^{+31}_{-26} + 17_{-103}$	1.8k	11 ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
$\sim 220$		HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
$\sim 60$		12 OAKDEN	94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
$\sim 250$		13 MARTIN	80B	RVUE	
$\sim 200$		13 MARTIN	80C	RVUE	
$\sim 150$		14 CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
$\sim 200$		15 CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

<sup>11</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (1.9 \pm 0.4^{+0.5}_{-1.3}) \times 10^{-6}$ .

<sup>12</sup> See however KLOET 96 who fit  $\pi^+ \pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>13</sup>  $I(J^P) = 1(3^-)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .

<sup>14</sup>  $I = 0, 1$ .  $J^P = 3^-$  from Barrelet-zero analysis.

<sup>15</sup>  $I(J^P) = 1(3^-)$  from amplitude analysis.

NODE=M044W1  
NODE=M044W1

NODE=M044W1;LINKAGE=A

NODE=M044W1;LINKAGE=CC

NODE=M044W1;LINKAGE=P

NODE=M044W1;LINKAGE=K

NODE=M044W1;LINKAGE=J

**S-CHANNEL  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
$160 \pm 25$	16 ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega \pi^0$ , $\omega \eta \pi^0, \pi^+ \pi^-$
$135 \pm 75$	17,18 COUPLAND	77	CNTR 0	0.7-2.4 $p\bar{p} \rightarrow \bar{p}p$
$98 \pm 8$	18 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
$\sim 85$	19 ABRAMS	70	CNTR	S channel $\bar{p}N$

<sup>16</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

<sup>17</sup> From a fit to the total elastic cross section.

<sup>18</sup> Isospins 0 and 1 not separated.

<sup>19</sup> Seen as bump in  $I = 1$  state. See also COOPER 68. PEASLEE 75 confirm  $\bar{p}p$  results of ABRAMS 70, no narrow structure.

NODE=M044W2  
NODE=M044W2

NODE=M044W;LINKAGE=AY

NODE=M044W2;LINKAGE=E

NODE=M044W2;LINKAGE=I

NODE=M044W2;LINKAGE=B

**Other processes**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$230 \pm 50 \pm 80$	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

NODE=M044W3  
NODE=M044W3

 **$\rho_3(2250)$  REFERENCES**

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP	REFID=21838
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP	REFID=21837
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)	REFID=21964
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)	REFID=21733
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP	REFID=21963
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)	REFID=21830
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)	REFID=21824
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)	REFID=21813
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)	REFID=21807
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)	REFID=21805

NODE=M044

$f_2(2300)$ 

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M107

 $f_2(2300)$  MASS

NODE=M107M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2297 ± 28</b>	<sup>1</sup> ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2262 ± 4 ± 28	<sup>2</sup> ABLIKIM	21A1 BES3	3.51–4.60 $e^+ e^- \rightarrow \phi \Lambda \bar{\Lambda}$
2243 <sup>+</sup> <sub>6</sub> <sup>7</sup> <sub>–29</sub> <sup>3</sup>	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
2270 ± 12	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2327 ± 9 ± 6	ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
2231 ± 10	BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
2220 <sup>+</sup> <sub>–20</sub> <sup>90</sup>	LINDENBAUM	84 RVUE	
2320 ± 40	ETKIN	82 MPS	22 $\pi^- p \rightarrow 2\phi n$

NODE=M107M

<sup>1</sup> Includes data of ETKIN 85. The percentage of the resonance going into  $\phi\phi 2^{++} S_2$ ,  $D_2$ , and  $D_0$  is  $6^{+15}_{-5}$ ,  $25^{+18}_{-14}$ , and  $69^{+16}_{-27}$ , respectively.

NODE=M107M;LINKAGE=C

<sup>2</sup> Threshold enhancement in  $\Lambda\bar{\Lambda}$ , preferred  $J^{PC}$  are  $2^{++}$ ,  $2^{-+}$ , or  $1^{++}$ . Could be another state.

NODE=M107M;LINKAGE=B

<sup>3</sup> Spin 2 preferred, tentatively assigned to  $f_2(2300)$ .

NODE=M107M;LINKAGE=A

 $f_2(2300)$  WIDTH

NODE=M107W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>149 ± 41</b>	<sup>1</sup> ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
72 ± 5 ± 43	<sup>2</sup> ABLIKIM	21A1 BES3	3.51–4.60 $e^+ e^- \rightarrow \phi \Lambda \bar{\Lambda}$
145 ± 12 <sup>+</sup> <sub>–34</sub> <sup>27</sup>	<sup>3</sup> UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
90 ± 29	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
275 ± 36 ± 20	ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
133 ± 50	BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
200 ± 50	LINDENBAUM	84 RVUE	
220 ± 70	ETKIN	82 MPS	22 $\pi^- p \rightarrow 2\phi n$

NODE=M107W

<sup>1</sup> Includes data of ETKIN 85.

NODE=M107W;LINKAGE=C

<sup>2</sup> Threshold enhancement in  $\Lambda\bar{\Lambda}$ , preferred  $J^{PC}$  are  $2^{++}$ ,  $2^{-+}$ , or  $1^{++}$ . Could be another state.

NODE=M107W;LINKAGE=B

<sup>3</sup> Spin 2 preferred, tentatively assigned to  $f_2(2300)$ .

NODE=M107W;LINKAGE=A

 $f_2(2300)$  DECAY MODES

NODE=M107215;NODE=M107

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\phi\phi$	seen
$\Gamma_2$ $K\bar{K}$	seen
$\Gamma_3$ $\gamma\gamma$	seen
$\Gamma_4$ $\Lambda\bar{\Lambda}$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

 $f_2(2300)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M107225

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_2\Gamma_3/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.2 <sup>+</sup> <sub>–0.4</sub> <sup>0.5</sup> <sub>–2.2</sub>	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
44 ± 6 ± 12	<sup>1</sup> ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$

NODE=M107G1

NODE=M107G1

<sup>1</sup> Assuming spin 2.

NODE=M107G1;LINKAGE=AB

$f_2(2340)$  BRANCHING RATIOS

NODE=M107220

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BOOTH	86	OMEG 85 $\pi^- \text{Be} \rightarrow 2\phi \text{Be}$
seen	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M107R01  
NODE=M107R01 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
seen	ABE	04	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$

NODE=M107R02  
NODE=M107R02 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M107R03  
NODE=M107R03 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> ABLIKIM	21AI	BES3 3.51-4.60 $e^+ e^- \rightarrow \phi \Lambda\bar{\Lambda}$

NODE=M107R00  
NODE=M107R00

<sup>1</sup> Threshold enhancement in  $\Lambda\bar{\Lambda}$ , preferred  $J^{PC}$  are  $2^{++}$ ,  $2^{-+}$ , or  $1^{++}$ . Could be another state.

NODE=M107R00;LINKAGE=A

 $f_2(2300)$  REFERENCES

NODE=M107

ABLIKIM	21AI	PR D104 052006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61440
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=40285
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIVP, GLAS, CERN)	REFID=21870
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21871
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)	REFID=21869
ETKIN	82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21866

NODE=M041

 $f_4(2300)$ 

$$I^G(J^{PC}) = 0^+(4^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called  $U_0(2350)$ . Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also  $\rho(2150)$ ,  $f_2(2150)$ ,  $\rho_3(2250)$ ,  $\rho_5(2350)$ .

NODE=M041

 $f_4(2300)$  MASS

NODE=M041205

NODE=M041M

 $\bar{p}p \rightarrow \pi\pi$  or  $\bar{K}K$ 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
•••	We do not use the following data for averages, fits, limits, etc. •••		
~ 2314	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2300	<sup>1</sup> MARTIN	80B	RVUE
~ 2300	<sup>1</sup> MARTIN	80C	RVUE
~ 2340	<sup>2</sup> CARTER	78B	CNTR 0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2330	DULUDE	78B	OSPK 1-2 $\bar{p}p \rightarrow \pi^0 \pi^0$
~ 2310	<sup>3</sup> CARTER	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

<sup>1</sup>  $I(J^P) = 0(4^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^- \pi^+$  and  $\pi^0 \pi^0$ .

<sup>2</sup>  $I(J^P) = 0(4^+)$  from Barrelet-zero analysis.

<sup>3</sup>  $I(J^P) = 0(4^+)$  from amplitude analysis.

NODE=M041M1  
NODE=M041M1NODE=M041M1;LINKAGE=P  
NODE=M041M1;LINKAGE=K  
NODE=M041M1;LINKAGE=J

**S-CHANNEL  $\bar{p}p$  or  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
2283±17	<sup>4</sup> ANISOVICH	00J	SPEC
~ 2380	<sup>5</sup> CUTTS	78B	CNTR 0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2345±15	<sup>5,6</sup> COUPLAND	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2359± 2	<sup>5,7</sup> ALSPECTOR	73	CNTR $\bar{p}p$ S channel
2375±10	ABRAMS	70	CNTR S channel $\bar{N}N$

NODE=M041M2  
 NODE=M041M2

<sup>4</sup> From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on  $\bar{p}p \rightarrow \eta\pi^0\pi^0$ ,  $\pi^0\pi^0$ ,  $\eta\eta$ ,  $\eta\eta'$ ,  $\pi^+\pi^-$ .

NODE=M041M2;LINKAGE=AN

<sup>5</sup> Isospins 0 and 1 not separated.

NODE=M041M2;LINKAGE=I

<sup>6</sup> From a fit to the total elastic cross section.

NODE=M041M2;LINKAGE=E

<sup>7</sup> Referred to as  $U$  or  $U$  region by ALSPECTOR 73.

NODE=M041M2;LINKAGE=M

 **$\pi^-p \rightarrow \eta\pi\pi n$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
2330±20±40	AMELIN	00	VES 37 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

NODE=M041M3  
 NODE=M041M3

 **$pp$  CENTRAL PRODUCTION**

VALUE (MeV)	DOCUMENT ID	COMMENT
<b>2320±60 OUR ESTIMATE</b>		

NODE=M041M4  
 NODE=M041M4

→ UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

2332±15	BARBERIS	00F	450 $pp \rightarrow p_f\omega\omega p_S$
---------	----------	-----	--

 **$f_4(2300)$  WIDTH**

NODE=M041210

 **$\bar{p}p \rightarrow \pi\pi$  or  $\bar{K}K$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 278	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 200	<sup>8</sup> MARTIN	80C	RVUE
~ 150	<sup>9</sup> CARTER	78B	CNTR 0.7-2.4 $\bar{p}p \rightarrow K^-K^+$
~ 210	<sup>10</sup> CARTER	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

NODE=M041W1  
 NODE=M041W1

<sup>8</sup>  $I(J^P) = 0(4^+)$  from simultaneous analysis of  $p\bar{p} \rightarrow \pi^-\pi^+$  and  $\pi^0\pi^0$ .

NODE=M041W1;LINKAGE=P

<sup>9</sup>  $I(J^P) = 0(4^+)$  from Barrelet-zero analysis.

NODE=M041W1;LINKAGE=K

<sup>10</sup>  $I(J^P) = 0(4^+)$  from amplitude analysis.

NODE=M041W1;LINKAGE=J

**S-CHANNEL  $\bar{p}p$  or  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
310± 25	<sup>11</sup> ANISOVICH	00J	SPEC
135 <sup>+150</sup> <sub>-65</sub>	<sup>12,13</sup> COUPLAND	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
165 <sup>+18</sup> <sub>-8</sub>	<sup>13</sup> ALSPECTOR	73	CNTR $\bar{p}p$ S channel
~ 190	ABRAMS	70	CNTR S channel $\bar{N}N$

NODE=M041W2  
 NODE=M041W2

<sup>11</sup> From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on  $\bar{p}p \rightarrow \eta\pi^0\pi^0$ ,  $\pi^0\pi^0$ ,  $\eta\eta$ ,  $\eta\eta'$ ,  $\pi^+\pi^-$ .

NODE=M041W2;LINKAGE=AN

<sup>12</sup> From a fit to the total elastic cross section.

NODE=M041W2;LINKAGE=E

<sup>13</sup> Isospins 0 and 1 not separated.

NODE=M041W2;LINKAGE=I

 **$\pi^-p \rightarrow \eta\pi\pi n$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
235±50±40	AMELIN	00	VES 37 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

NODE=M041W3  
 NODE=M041W3

 **$pp$  CENTRAL PRODUCTION**

VALUE (MeV)	DOCUMENT ID	COMMENT
<b>250±80 OUR ESTIMATE</b>		

NODE=M041W4  
 NODE=M041W4

→ UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

260±57	BARBERIS	00F	450 $pp \rightarrow p_f\omega\omega p_S$
--------	----------	-----	--



$f_4(2300)$  DECAY MODES

NODE=M041215;NODE=M041

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho\rho$	seen
$\Gamma_2$ $\omega\omega$	seen
$\Gamma_3$ $\eta\pi\pi$	seen
$\Gamma_4$ $\pi\pi$	seen
$\Gamma_5$ $K\bar{K}$	seen
$\Gamma_6$ $N\bar{N}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=2;OUR EST;→ UNCHECKED ←  
DESIG=3;OUR EST;→ UNCHECKED ←  
DESIG=4;OUR EST;→ UNCHECKED ←  
DESIG=5;OUR EST;→ UNCHECKED ←  
DESIG=6;OUR EST;→ UNCHECKED ←

 $f_4(2300)$  BRANCHING RATIOS

NODE=M041220

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$	$\Gamma_1/\Gamma_2$
VALUE	DOCUMENT ID COMMENT
2.8±0.5	BARBERIS 00F 450 $\rho\rho \rightarrow p_f\omega\omega p_s$

NODE=M041R1  
NODE=M041R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $f_4(2300)$  REFERENCES

NODE=M041

AMELIN 00 NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH 00J PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARBERIS 00F PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH 99C PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH 99F NP A651 253	A.V. Anisovich <i>et al.</i>	
HASAN 94 PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
MARTIN 80B NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN 80C NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER 78B NP B141 467	A.A. Carter	(LOQM)
CUTTS 78B PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE 78B PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
CARTER 77 PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND 77 PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR 73 PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS 70 PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)

REFID=47432  
REFID=47950  
REFID=47962  
REFID=46903  
REFID=46926  
REFID=44103  
REFID=21838  
REFID=21837  
REFID=21964  
REFID=21733  
REFID=21850  
REFID=21963  
REFID=21830  
REFID=21813  
REFID=21807

 $f_0(2330)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

NODE=M169

OMITTED FROM SUMMARY TABLE

 $f_0(2330)$  MASS

NODE=M169M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2312± 7 <sup>+</sup> <sub>3</sub>	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$
2312± 2 <sup>+</sup> <sub>0</sub> <sup>10</sup>	<sup>2</sup> ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
2419±64	<sup>3</sup> RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
2340±20	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
2314±25	<sup>4</sup> BUGG	04A RVUE	
2337±14	ANISOVICH	00J SPEC	2.0 $\bar{p}p \rightarrow \pi\pi, \eta\eta$
~ 2321	HASAN	94 RVUE	$\bar{p}p \rightarrow \pi\pi$

NODE=M169M

- <sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$  P-wave.  
<sup>2</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.  
<sup>3</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma\pi^0\pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).  
<sup>4</sup> Partial wave analysis of the data on  $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$  from BARNES 00.

NODE=M169M;LINKAGE=C

NODE=M169M;LINKAGE=B

NODE=M169M;LINKAGE=A

NODE=M169M;LINKAGE=BU

 $f_0(2330)$  WIDTH

NODE=M169W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
65±10 <sup>+</sup> <sub>12</sub> <sup>3</sup>	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

NODE=M169W

• • • We do not use the following data for averages, fits, limits, etc. • • •

$134 \pm 5_{-9}^{+30}$	<sup>2</sup> ABLIKIM	22C	BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
$274 \pm 94$	<sup>3</sup> RODAS	22	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$165 \pm 25$	SARANTSEV	21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$144 \pm 20$	<sup>4</sup> BUGG	04A	RVUE	
$217 \pm 33$	ANISOVICH	00J	SPEC	$2.0 \bar{p}p \rightarrow \pi\pi, \eta\eta$
$\sim 223$	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta\eta'$   $P$ -wave.

<sup>2</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>3</sup> T-matrix pole from coupled channel K-matrix fit to data on  $J/\psi \rightarrow \gamma\pi^0\pi^0$  (ABLIKIM 15AE) and  $J/\psi \rightarrow \gamma K_S^0 K_S^0$  (ABLIKIM 18AA).

<sup>4</sup> Partial wave analysis of the data on  $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$  from BARNES 00.

NODE=M169W;LINKAGE=C

NODE=M169W;LINKAGE=B

NODE=M169W;LINKAGE=A

NODE=M169W;LINKAGE=BU

### $f_0(2330)$ REFERENCES

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BUGG	04A	EPJ C36 161	D.V. Bugg	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARNES	00	PR C62 055203	P.D. Barnes <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)

NODE=M169

REFID=61891

REFID=61637

REFID=61610

REFID=61091

REFID=59455

REFID=56984

REFID=50158

REFID=47950

REFID=47965

REFID=44103

NODE=M108

$f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

### $f_2(2340)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2346_{-10}^{+21}</math> OUR AVERAGE</b>				
[ $2345_{-40}^{+50}$ MeV OUR 2022 AVERAGE]				
$2346 \pm 8_{-6}^{+22}$		<sup>1</sup> ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
$2362_{-30}^{+31} + 140_{-63}$	5.5k	<sup>2</sup> ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
$2339 \pm 55$		<sup>3</sup> ETKIN	88	MPS $22 \pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$2350 \pm 7$	80k	<sup>4</sup> UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
$2392 \pm 10$		BOOTH	86	OMEG $85 \pi^- Be \rightarrow 2\phi Be$
$2360 \pm 20$		LINDENBAUM	84	RVUE

NODE=M108M

NODE=M108M

NEW

<sup>1</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>2</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>3</sup> Includes data of ETKIN 85. The percentage of the resonance going into  $\phi\phi 2^{++} S_2$ ,  $D_2$ , and  $D_0$  is  $37 \pm 19$ ,  $4_{-4}^{+12}$ , and  $59_{-19}^{+21}$ , respectively.

<sup>4</sup> Statistical error only.

NODE=M108M;LINKAGE=B

NODE=M108M;LINKAGE=A

NODE=M108M;LINKAGE=C

NODE=M108M;LINKAGE=ST

### $f_2(2340)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>331_{-18}^{+27}</math> OUR AVERAGE</b>				
[ $322_{-60}^{+70}$ MeV OUR 2022 AVERAGE]				
$332 \pm 14_{-12}^{+26}$		<sup>1</sup> ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
$334_{-54}^{+62} + 165_{-100}$	5.5k	<sup>2</sup> ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
$319_{-69}^{+81}$		<sup>3</sup> ETKIN	88	MPS $22 \pi^- p \rightarrow \phi \phi n$

NODE=M108W

NODE=M108W

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

218 ± 16	80k	<sup>4</sup> UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
198 ± 50		BOOTH	86	OMEG	85 $\pi^- \text{Be} \rightarrow 2\phi\text{Be}$
150 <sup>+150</sup> <sub>-50</sub>		LINDENBAUM	84	RVUE	

<sup>1</sup> From a partial wave analysis of the systems ( $\gamma X$ ), with  $X \rightarrow \eta'\eta'$ , and ( $\eta' X$ ), with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

<sup>2</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>3</sup> Includes data of ETKIN 85.

<sup>4</sup> Statistical error only.

NODE=M108W;LINKAGE=B

NODE=M108W;LINKAGE=A

NODE=M108W;LINKAGE=C

NODE=M108W;LINKAGE=ST

NODE=M108215;NODE=M108

## $f_2(2340)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\phi\phi$	seen
$\Gamma_2$ $\eta\eta$	seen
$\Gamma_3$ $\eta'\eta'$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2

DESIG=3

## $f_2(2340)$ BRANCHING RATIOS

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
seen	UMAN	06	E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$

NODE=M108220

NODE=M108R01

NODE=M108R01

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma$
seen	<sup>1</sup> ABLIKIM	22C	BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M108R00

NODE=M108R00

<sup>1</sup> From a partial wave analysis of the systems ( $\gamma X$ ), with  $X \rightarrow \eta'\eta'$ , and ( $\eta' X$ ), with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M108R00;LINKAGE=A

## $f_2(2340)$ REFERENCES

ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61637
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=40285
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIVP, GLAS, CERN)	REFID=21870
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21871
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)	REFID=21869

NODE=M108

**$\rho_5(2350)$** 

$$I^G(J^{PC}) = 1^+(5^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called  $U_1(2400)$ . See also  $\rho(2150)$ ,  $f_2(2150)$ ,  $\rho_3(2250)$ ,  $f_4(2300)$ .

NODE=M033

NODE=M033

 **$\rho_5(2350)$  MASS**

NODE=M033205

NODE=M033M

 **$\pi^- p \rightarrow \omega \pi^0 n$** NODE=M033M3  
NODE=M033M3

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>2330±35</b>	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$

 **$\bar{p} p \rightarrow \pi \pi$  or  $\bar{K} K$** NODE=M033M1  
NODE=M033M1

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 2303	HASAN	94	RVUE	$\bar{p} p \rightarrow \pi \pi$
~ 2300	<sup>1</sup> MARTIN	80B	RVUE	
~ 2250	<sup>1</sup> MARTIN	80C	RVUE	
~ 2500	<sup>2</sup> CARTER	78B	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow K^- K^+$
~ 2480	<sup>3</sup> CARTER	77	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow \pi \pi$

**S-CHANNEL  $\bar{N} N$** NODE=M033M2  
NODE=M033M2

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
2300±45	<sup>4</sup> ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega \pi^0$ , $\omega \eta \pi^0$ , $\pi^+ \pi^-$
2295±30	ANISOVICH	00J	SPEC	
~ 2380	<sup>5</sup> CUTTS	78B	CNTR	0.97-3 $\bar{p} p \rightarrow \bar{N} N$
2345±15	<sup>5,6</sup> COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow \bar{p} p$
2359± 2	<sup>5,7</sup> ALSPECTOR	73	CNTR	$\bar{p} p$ S channel
2350±10	<sup>8</sup> ABRAMS	70	CNTR	S channel $\bar{N} N$
2360±25	<sup>9</sup> OH	70B	HDBC -0	$\bar{p}(p n)$ , $K^* K 2p$

 **$\pi^- p \rightarrow K^+ K^- n$** NODE=M033M4  
NODE=M033M4

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
2307±6	ALPER	80	CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$
<sup>1</sup> $I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$ .				
<sup>2</sup> $I = 0(1)$ ; $J^P = 5^-$ from Barrelet-zero analysis.				
<sup>3</sup> $I(J^P) = 1(5^-)$ from amplitude analysis.				
<sup>4</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.				
<sup>5</sup> Isospins 0 and 1 not separated.				
<sup>6</sup> From a fit to the total elastic cross section.				
<sup>7</sup> Referred to as $U$ or $U$ region by ALSPECTOR 73.				
<sup>8</sup> For $I = 1 \bar{N} N$ .				
<sup>9</sup> No evidence for this bump seen in the $\bar{p} p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.				

NODE=M033M1;LINKAGE=P  
NODE=M033M1;LINKAGE=K  
NODE=M033M1;LINKAGE=J  
NODE=M033M2;LINKAGE=AYNODE=M033M2;LINKAGE=I  
NODE=M033M2;LINKAGE=E  
NODE=M033M2;LINKAGE=M  
NODE=M033M2;LINKAGE=A  
NODE=M033M2;LINKAGE=N **$\rho_5(2350)$  WIDTH**

NODE=M033210

 **$\pi^- p \rightarrow \omega \pi^0 n$** NODE=M033W3  
NODE=M033W3

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>400±100</b>	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$

 **$\bar{p} p \rightarrow \pi \pi$  or  $\bar{K} K$** NODE=M033W1  
NODE=M033W1

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 169	HASAN	94	RVUE	$\bar{p} p \rightarrow \pi \pi$
~ 250	<sup>10</sup> MARTIN	80B	RVUE	
~ 300	<sup>10</sup> MARTIN	80C	RVUE	
~ 150	<sup>11</sup> CARTER	78B	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow K^- K^+$
~ 210	<sup>12</sup> CARTER	77	CNTR 0	0.7-2.4 $\bar{p} p \rightarrow \pi \pi$

**S-CHANNEL  $\bar{N}N$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$260 \pm 75$	13 ANISOVICH	02	SPEC	$0.6-1.9 \bar{p}\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
$235^{+65}_{-40}$	ANISOVICH	00J	SPEC	
$135^{+150}_{-65}$	14,15 COUPLAND	77	CNTR 0	$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
$165^{+18}_{-8}$	15 ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
< 60	16 OH	70B	HDBC -0	$\bar{p}(pn), K^*K2\pi$
~ 140	ABRAMS	67C	CNTR	S channel $\bar{p}N$

NODE=M033W2  
 NODE=M033W2

 **$\pi^-p \rightarrow K^+K^-n$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$245 \pm 20$	ALPER	80	CNTR 0	$62 \pi^-p \rightarrow K^+K^-n$
$10 I(J^P) = 1(5^-)$	from simultaneous analysis of $\bar{p}\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$ .			
$11 I = 0(1); J^P = 5^-$	from Barrelet-zero analysis.			
$12 I(J^P) = 1(5^-)$	from amplitude analysis.			
13	From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.			
14	From a fit to the total elastic cross section.			
15	Isospins 0 and 1 not separated.			
16	No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.			

NODE=M033W4  
 NODE=M033W4

OCCUR=2

NODE=M033W1;LINKAGE=P  
 NODE=M033W1;LINKAGE=K  
 NODE=M033W1;LINKAGE=J  
 NODE=M033W2;LINKAGE=AY

NODE=M033W2;LINKAGE=E  
 NODE=M033W2;LINKAGE=I  
 NODE=M033W2;LINKAGE=N

 **$\rho_5(2350)$  REFERENCES**

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington
CARTER	78B	NP B141 467	A.A. Carter
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>
OH	73	NP B51 57	B.Y. Oh <i>et al.</i>
CHAPMAN	71B	PR D4 1275	J.W. Chapman <i>et al.</i>
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>
OH	70B	PRL 24 1257	B.Y. Oh <i>et al.</i>
ABRAMS	67C	PRL 18 1209	R.J. Abrams <i>et al.</i>

NODE=M033

REFID=48828  
 REFID=48327  
 REFID=48349  
 REFID=47950  
 REFID=44371  
 REFID=44103  
 REFID=21665  
 REFID=21838  
 REFID=21837  
 REFID=21964  
 REFID=21733  
 REFID=21963  
 REFID=21830  
 REFID=21813  
 REFID=21931  
 REFID=21926  
 REFID=21807  
 REFID=21925  
 REFID=21804

**X(2370)**

$$I^G(J^{PC}) = ?^?(???)$$

NODE=M247

OMITTED FROM SUMMARY TABLE

**X(2370) MASS**

NODE=M247M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2357 ±17 OUR AVERAGE</b>		Error includes scale factor of 2.7. [2342 ± 9 MeV OUR 2022 AVERAGE]		
2341.6 ± 6.5 ± 5.7		<sup>1</sup> ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K \bar{K} \eta'$
2376.3 ± 8.7 <sup>+3.2</sup> <sub>-4.3</sub>	565	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M247M  
NEW

<sup>1</sup> The state observed by ABLIKIM 11C at 2120 MeV is not observed with 90% CL upper limit of  $1.49 \times 10^{-5}$  for  $J/\psi \rightarrow \gamma X(2120) \rightarrow \gamma K^+ K^- \eta'$  and  $6.38 \times 10^{-6}$  for  $K_S^0 K_S^0 \eta'$ .

NODE=M247M;LINKAGE=A

**X(2370) WIDTH**

NODE=M247W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>114<sup>+12</sup><sub>-10</sub> OUR AVERAGE</b>	[117 ± 13 MeV OUR 2022 AVERAGE]		
117 ± 10 ± 8	<sup>1</sup> ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K \bar{K} \eta'$
83 ± 17 <sup>+44</sup> <sub>-6</sub>	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M247W

NEW

<sup>1</sup> The state observed by ABLIKIM 11C at 2120 MeV is not observed with 90% CL upper limit of  $1.49 \times 10^{-5}$  for  $J/\psi \rightarrow \gamma X(2120) \rightarrow \gamma K^+ K^- \eta'$  and  $6.38 \times 10^{-6}$  for  $K_S^0 K_S^0 \eta'$ .

NODE=M247W;LINKAGE=A

**X(2370) DECAY MODES**

NODE=M247215;NODE=M247

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^+ K^- \eta'$	seen
$\Gamma_2$ $K_S^0 K_S^0 \eta'$	seen
$\Gamma_3$ $\pi^+ \pi^- \eta'$	seen

DESIG=1

DESIG=2

DESIG=3

**X(2370) BRANCHING RATIOS**

NODE=M247225

$\Gamma(K^+ K^- \eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
VALUE				
seen	ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K^+ K^- \eta'$	
$\Gamma(K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
VALUE				
seen	ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$	
$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma$
VALUE				
seen	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$	

NODE=M247R01  
NODE=M247R01NODE=M247R02  
NODE=M247R02NODE=M247R00  
NODE=M247R00**X(2370) REFERENCES**

NODE=M247

ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)

REFID=60457  
REFID=53684

**$f_0(2470)$** 

$$I^G(J^{PC}) = 0^+(0^{++})$$

NODE=M266

## OMITTED FROM SUMMARY TABLE

Seen by ABLIKIM 22C with a significance of  $5.2\sigma$  in a partial-wave analysis of the systems  $(\gamma X)$ ,  $X \rightarrow \eta'\eta'$  and  $(\eta' X)$ ,  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ .

NODE=M266

 **$f_0(2470)$  MASS**

NODE=M266M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2470 \pm 4_{-6}^{+4}$	<sup>1</sup> ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M266M

<sup>1</sup>From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M266M;LINKAGE=A

 **$f_0(2470)$  WIDTH**

NODE=M266W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$75 \pm 9_{-8}^{+11}$	<sup>1</sup> ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M266W

<sup>1</sup>From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M266W;LINKAGE=C

 **$f_0(2470)$  DECAY MODES**

NODE=M266215;NODE=M266

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \eta'\eta'$	seen

DESIG=1

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
seen	<sup>1</sup> ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$	

NODE=M266R00  
NODE=M266R00

<sup>1</sup>From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta'\eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma\eta'$  in the decay  $J/\psi \rightarrow \gamma\eta'\eta'$ . The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M266R00;LINKAGE=A

 **$f_0(2470)$  REFERENCES**

NODE=M266

ABLIKIM 22C PR D105 072002 M. Ablikim *et al.* (BESIII Collab.)

REFID=61637

$f_6(2510)$ 

$$I^G(J^{PC}) = 0^+(6^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M089

NODE=M089

 $f_6(2510)$  MASS

NODE=M089M

NODE=M089M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2465±50 OUR AVERAGE</b>	Error includes scale factor of 2.1.		
2420±30	ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
2510±30	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2485±40	<sup>1</sup> ANISOVICH	00J	SPEC 1.92–2.41 $p\bar{p}$
<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B.			

NODE=M089M;LINKAGE=AN

 $f_6(2510)$  WIDTH

NODE=M089W

NODE=M089W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>255±40 OUR AVERAGE</b>			
270±60	ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
240±60	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
410±90	<sup>2</sup> ANISOVICH	00J	SPEC 1.92–2.41 $p\bar{p}$
<sup>2</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B.			

NODE=M089W;LINKAGE=AN

 $f_6(2510)$  DECAY MODES

NODE=M089215;NODE=M089

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \pi\pi$	(6.0±1.0) %

DESIG=1

 $f_6(2510)$  BRANCHING RATIOS

NODE=M089220

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>0.06±0.01</b>	<sup>3</sup> BINON	83C	GAM2 38 $\pi^- p \rightarrow n 4\gamma$	
<sup>3</sup> Assuming one pion exchange and using data of BOLOTOV 74.				

NODE=M089R1  
NODE=M089R1

NODE=M089R1;LINKAGE=A

 $f_6(2510)$  REFERENCES

NODE=M089

ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>		REFID=46926
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>		REFID=47416
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP) JP	REFID=21780
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+)	REFID=40288
		Translated from YAF 38 1199.			
BOLOTOV	74	PL 52B 489	V.N. Bolotov <i>et al.</i>	(SERP)	REFID=44705



# STRANGE MESONS

## ( $S = \pm 1, C = B = 0$ )

$K^+ = u\bar{s}, K^0 = d\bar{s}, \bar{K}^0 = \bar{d}s, K^- = \bar{u}s$ , similarly for  $K^*$ 's

### $K_0^*(700)$

$$I(J^P) = \frac{1}{2}(0^+)$$

also known as  $\kappa$ ; was  $K_0^*(800)$

See the related review(s):

Scalar Mesons below 1 GeV

### $K_0^*(700)$ T-Matrix Pole $\sqrt{s}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(630–730) – <math>i</math> (260–340) OUR ESTIMATE</b> (see Fig. 64.1 in the review)			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(648 \pm 7) - i(280 \pm 16)$	<sup>1</sup> PELAEZ	20	RVUE $\pi K \rightarrow \pi K$
$(670 \pm 18) - i(295 \pm 28)$	<sup>2</sup> PELAEZ	17	RVUE $\pi K \rightarrow \pi K$
$(764 \pm 63_{-54}^{+71}) - i(306 \pm 149_{-85}^{+143})$	<sup>3</sup> ABLIKIM	11B	BES2 $1.3k J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
$(665 \pm 9) - i(268_{-6}^{+21})$	<sup>4</sup> GUO	11B	RVUE
$(849 \pm 77_{-14}^{+18}) - i(256 \pm 40_{-22}^{+46})$	<sup>3</sup> ABLIKIM	10E	BES2 $1.4k J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
$(663 \pm 8 \pm 34) - i(329 \pm 5 \pm 22)$	<sup>5</sup> BUGG	10	RVUE S-matrix pole
$(706.0 \pm 1.8 \pm 22.8) - i(319.4 \pm 2.2 \pm 20.2)$	<sup>6</sup> BONVICINI	08A	CLEO $141k D^+ \rightarrow K^- \pi^+ \pi^+$
$(841 \pm 30_{-73}^{+81}) - i(309 \pm 45_{-72}^{+48})$	<sup>3</sup> ABLIKIM	06C	BES2 $25k J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
$(750_{-55}^{+30}) - i(342 \pm 60)$	<sup>7</sup> BUGG	06	RVUE
$(658 \pm 13) - i(279 \pm 12)$	<sup>8</sup> DESCOTES-G.	06	RVUE $\pi K \rightarrow \pi K$
$(757 \pm 33) - i(279 \pm 41)$	<sup>9</sup> GUO	06	RVUE
$(694 \pm 53) - i(303 \pm 30)$	<sup>10</sup> ZHOU	06	RVUE $K p \rightarrow K^- \pi^+ n$
$(594 \pm 79) - i(362 \pm 166)$	<sup>10</sup> ZHENG	04	RVUE $K^- p \rightarrow K^- \pi^+ n$
$(722 \pm 60) - i(386 \pm 50)$	<sup>10</sup> BUGG	03	RVUE $11 K^- p \rightarrow K^- \pi^+ n$
$(875 \pm 75) - i(335 \pm 110)$	<sup>11</sup> ISHIDA	97B	RVUE $11 K^- p \rightarrow K^- \pi^+ n$
$727 - i 263$	<sup>12</sup> VANBEVEREN	86	RVUE

<sup>1</sup> Extracted employing  $\pi K$  partial wave analysis from ESTABROOKS 78 and ASTON 88, Roy-Steiner equations and once subtracted forward dispersion relations.

<sup>2</sup> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

<sup>3</sup> Extracted from Breit-Wigner parameters.

<sup>4</sup> Fit to scattering phase shifts using UChPT amplitudes with explicit resonances.

<sup>5</sup> Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an  $s$ -dependent width with couplings to  $K\pi$  and  $K\eta'$ , and the Adler zero near thresholds.

<sup>6</sup> From a complex pole included in the fit. Using parameters from the model that fits data best.

<sup>7</sup> Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C using for the  $\kappa$  an  $s$ -dependent width with an Adler zero near threshold.

<sup>8</sup> Using Roy-Steiner equations (ROY 71) consistent with unitarity, analyticity and crossing symmetry constraints.

<sup>9</sup> From UChPT fitted to MERCER 71, BINGHAM 72 and ESTABROOKS 78. Amplitude shown to be consistent with data of ABLIKIM 06C.

<sup>10</sup> Reanalysis of ASTON 88 data.

<sup>11</sup> Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes. Extracted from Breit-Wigner parameters.

<sup>12</sup> Unitarized Quark Model.

NODE=MXXX020

NODE=MXXX020

NODE=M174

NODE=M174TMP

NODE=M174TMP

→ UNCHECKED ←

NODE=M174TMP;LINKAGE=J

NODE=M174TMP;LINKAGE=N

NODE=M174TMP;LINKAGE=A

NODE=M174TMP;LINKAGE=D

NODE=M174TMP;LINKAGE=H

NODE=M174TMP;LINKAGE=B

NODE=M174TMP;LINKAGE=G

NODE=M174TMP;LINKAGE=I

NODE=M174TMP;LINKAGE=O

NODE=M174TMP;LINKAGE=F

NODE=M174TMP;LINKAGE=M

NODE=M174TMP;LINKAGE=E

**$K_0^*(700)$  Breit-Wigner Mass**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>845 ±17</b>	<b>OUR AVERAGE</b>			
826 ±49 $^{+49}_{-34}$	1.3k	<sup>1</sup> ABLIKIM	11B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
810 ±68 $^{+15}_{-24}$	1.4k	<sup>2</sup> ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
856 ±17 ±13	54k	LINK	07B FOCUS	$D^+ \rightarrow K^- \pi^+ \pi^+$
878 ±23 $^{+64}_{-55}$	25k	<sup>3</sup> ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
797 ±19 ±43	15k	<sup>4,5</sup> AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
888.0 ± 1.9	141k	<sup>6</sup> BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
855 ±15	0.6k	<sup>7</sup> CAWLFIELD	06A CLEO	$D^0 \rightarrow K^+ K^- \pi^0$
905 $^{+65}_{-30}$		<sup>8</sup> ISHIDA	97B RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M174M  
 NODE=M174M

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M174M;LINKAGE=LI

NODE=M174M;LINKAGE=BL

NODE=M174M;LINKAGE=EP

NODE=M174M;LINKAGE=A

NODE=M174M;LINKAGE=AU

NODE=M174M;LINKAGE=C

NODE=M174M;LINKAGE=CA

NODE=M174M;LINKAGE=IS

<sup>1</sup> The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is  $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$  MeV.

<sup>2</sup> From a fit including ten additional resonances and energy-independent Breit-Wigner width.

<sup>3</sup> A fit in the  $K_0^*(700) + K^*(892) + K^*(1410)$  model with mass and width of the  $K_0^*(700)$  from ABLIKIM 06C well describes the left slope of the  $K_S^0 \pi^-$  invariant mass spectrum in  $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$  decay studied by EPIFANOV 07. Averaged value from different parameterizations.

<sup>4</sup> Not seen by KOPP 01 using 7070 events of  $D^0 \rightarrow K^- \pi^+ \pi^0$ . LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than  $K_0^*(700)$  in their high statistics analysis of  $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$ .

<sup>5</sup> AUBERT 07T does not find evidence for the charged  $K_0^*(700)$  using 11k events of  $D^0 \rightarrow K^- K^+ \pi^0$ .

<sup>6</sup> Using parameters from the model that fits data best.

<sup>7</sup> Breit-Wigner parameters. A significant S-wave can be also modeled as a non-resonant contribution.

<sup>8</sup> Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.

 **$K_0^*(700)$  Breit-Wigner Width**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>468 ± 30</b>	<b>OUR AVERAGE</b>			
449 ±156 $^{+144}_{-81}$	1.3k	<sup>1</sup> ABLIKIM	11B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
536 ± 87 $^{+106}_{-47}$	1.4k	<sup>2</sup> ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
464 ± 28 ± 22	54k	LINK	07B FOCUS	$D^+ \rightarrow K^- \pi^+ \pi^+$
499 ± 52 $^{+55}_{-87}$	25k	<sup>3</sup> ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
410 ± 43 ± 87	15k	<sup>4,5</sup> AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
550.4 ± 11.8	141k	<sup>6</sup> BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
251 ± 48	0.6k	<sup>7</sup> CAWLFIELD	06A CLEO	$D^0 \rightarrow K^+ K^- \pi^0$
545 $^{+235}_{-110}$		<sup>8</sup> ISHIDA	97B RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M174W

NODE=M174W

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M174W;LINKAGE=LI

NODE=M174W;LINKAGE=BL

NODE=M174W;LINKAGE=EP

NODE=M174W;LINKAGE=A

NODE=M174W;LINKAGE=AU

NODE=M174W;LINKAGE=C

NODE=M174W;LINKAGE=CA

NODE=M174W;LINKAGE=IS

<sup>1</sup> The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is  $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$  MeV.

<sup>2</sup> From a fit including ten additional resonances and energy-independent Breit-Wigner width.

<sup>3</sup> A fit in the  $K_0^*(700) + K^*(892) + K^*(1410)$  model with mass and width of the  $K_0^*(700)$  from ABLIKIM 06C well describes the left slope of the  $K_S^0 \pi^-$  invariant mass spectrum in  $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$  decay studied by EPIFANOV 07. Averaged value from different parameterizations.

<sup>4</sup> Not seen by KOPP 01 using 7070 events of  $D^0 \rightarrow K^- \pi^+ \pi^0$ . LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than  $K_0^*(700)$  in their high statistics analysis of  $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$ .

<sup>5</sup> AUBERT 07T does not find evidence for the charged  $K_0^*(700)$  using 11k events of  $D^0 \rightarrow K^- K^+ \pi^0$ .

<sup>6</sup> Using parameters from the model that fits data best.

<sup>7</sup> Statistical error only. A fit to the Dalitz plot including the  $K_0^*(700)^\pm$ ,  $K^*(892)^\pm$ , and  $\phi$  resonances modeled as Breit-Wigners. A significant S-wave can be also modeled as a non-resonant contribution.

<sup>8</sup> Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.

**$K_0^*(700)$  DECAY MODES**

NODE=M174215;NODE=M174

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\pi$	100 %

DESIG=1;OUR EVAL;→ UNCHECKED ←

 **$K_0^*(700)$  REFERENCES**

NODE=M174

PELAEZ	20	PRL 124 172001	J.R. Pelaez <i>et al.</i>
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira
ABLIKIM	11B	PL B698 183	M. Ablikim <i>et al.</i> (BES II Collab.)
GUO	11B	PR D84 034005	Z.-H. Guo, J.A. Oller
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i> (BES II Collab.)
BUGG	10	PR D81 014002	D.V. Bugg (LOQM)
LINK	09	PL B681 14	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i> (CLEO Collab.)
AUBERT	07T	PR D76 011102	B. Aubert <i>et al.</i> (BABAR Collab.)
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i> (BELLE Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i> (BES Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
BUGG	06	PL B632 471	D.V. Bugg (LOQM)
CAWLFIELD	06A	PR D74 031108	C. Cawfield <i>et al.</i> (CLEO Collab.)
DESCOTES-G...	06	EPJ C48 553	S. Descotes-Genon, B. Moussallam
GUO	06	NP A773 78	F.K. Guo <i>et al.</i>
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng
LINK	05I	PL B621 72	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>
BUGG	03	PL B572 1	D.V. Bugg
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
LINK	02E	PL B535 43	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
KOPP	01	PR D63 092001	S. Kopp <i>et al.</i> (CLEO Collab.)
ISHIDA	97B	PTP 98 621	S. Ishida <i>et al.</i>
ASTON	88	NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i> (NIJM, BIEL)
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i> (MCGI, CARL, DURH+)
BINGHAM	72	NP B41 1	H.H. Bingham <i>et al.</i> (International $K^+$ Collab.)
MERCER	71	NP B32 381	R. Mercer <i>et al.</i> (JHU)
ROY	71	PL 36B 353	S.M. Roy

REFID=60559

REFID=57836

REFID=53683

REFID=58808

REFID=53361

REFID=53213

REFID=53056

REFID=52426

REFID=51726

REFID=51929

REFID=51875

REFID=51037

REFID=51051

REFID=51458

REFID=50996

REFID=51153

REFID=51518

REFID=51164

REFID=51198

REFID=50679

REFID=50165

REFID=49586

REFID=48807

REFID=48728

REFID=48134

REFID=48655

REFID=40262

REFID=45769

REFID=22443

REFID=22415

REFID=22412

REFID=51107

 **$K^*(892)$** 

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M018

 **$K^*(892)$  T-Matrix Pole  $\sqrt{s}$** 

NODE=M018TMP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**(890 ± 14) – i (26 ± 6) OUR ESTIMATE**

NODE=M018TMP

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

(890 ± 2) – i (25.6 ± 1.2)	<sup>1</sup> PELAEZ	20	RVUE	$\pi K \rightarrow \pi K$
(892 ± 1) – i (29 ± 1)	<sup>2</sup> PELAEZ	17	RVUE	$\pi K \rightarrow \pi K$
(889 ± 13) – i (24 ± 4)	<sup>3</sup> PELAEZ	04A	RVUE	$\pi K \rightarrow \pi K$

OCCUR=2

<sup>1</sup> Extracted employing  $\pi K$  partial wave analysis from ESTABROOKS 78 and ASTON 88, Roy-Steiner equations and once subtracted forward dispersion relations.

NODE=M018TMP;LINKAGE=A

<sup>2</sup> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M018TMP;LINKAGE=E

<sup>3</sup> Reanalysis of data from ESTABROOKS 78 and ASTON 88 in the unitarized ChPT model.

NODE=M018TMP;LINKAGE=B

 **$K^*(892)$  MASS**

NODE=M018205

**CHARGED ONLY, HADROPRODUCED**

NODE=M018M1

NODE=M018M1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**891.67 ± 0.26 OUR AVERAGE**

892.2 ± 0.5 ± 1.7		ALBRECHT	20	CBAR	0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
892.6 ± 0.5	5840	BAUBILLIER	84B	HBC	– 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
888 ± 3		NAPIER	84	SPEC	+ 200 $\pi^- p \rightarrow 2K_S^0 X$
891 ± 1		NAPIER	84	SPEC	– 200 $\pi^- p \rightarrow 2K_S^0 X$
891.7 ± 2.1	3700	BARTH	83	HBC	+ 70 $K^+ p \rightarrow K^0 \pi^+ X$
891 ± 1	4100	TOAFF	81	HBC	– 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892.8 ± 1.6		AJINENKO	80	HBC	+ 32 $K^+ p \rightarrow K^0 \pi^+ X$
890.7 ± 0.9	1800	AGUILAR-...	78B	HBC	± 0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
886.6 ± 2.4	1225	BALAND	78	HBC	± 12 $\bar{p}p \rightarrow (K\pi)^\pm X$

OCCUR=2

891.7 ± 0.6	6706	COOPER	78	HBC	±	0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$	
891.9 ± 0.7	9000	<sup>1</sup> PALER	75	HBC	-	14.3 $K^- p \rightarrow (K\pi)^- X$	
892.2 ± 1.5	4404	AGUILAR...	71B	HBC	-	3.9,4.6 $K^- p \rightarrow (K\pi)^- p$	
891 ± 2	1000	CRENNELL	69D	DBC	-	3.9 $K^- N \rightarrow K^0 \pi^- X$	
890 ± 3.0	720	BARLOW	67	HBC	±	1.2 $\bar{p}p \rightarrow (K^0 \pi)^\pm K^\mp$	
889 ± 3.0	600	BARLOW	67	HBC	±	1.2 $\bar{p}p \rightarrow (K^0 \pi)^\pm K \pi$	OCCUR=2
891 ± 2.3	620	<sup>2</sup> DEBAERE	67B	HBC	+	3.5 $K^+ p \rightarrow K^0 \pi^+ p$	
891.0 ± 1.2	1700	<sup>3</sup> WOJCICKI	64	HBC	-	1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

893.6 ± 0.1 <sup>+0.2</sup> / <sub>-0.3</sub>	183k	ABLIKIM	19AQ	BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$	
895.6 ± 0.8	4k	<sup>4</sup> LEES	17C	BABR		$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	
893.2 ± 0.1 ± 1.0	190k	<sup>5</sup> AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$	
893.5 ± 1.1	27k	<sup>6</sup> ABELE	99D	CBAR	±	0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$	
890.4 ± 0.2 ± 0.5	80k	<sup>7</sup> BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
890.0 ± 2.3	800	<sup>2,3</sup> CLELAND	82	SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$	
896.0 ± 1.1	3200	<sup>2,3</sup> CLELAND	82	SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$	OCCUR=2
893 ± 1	3600	<sup>2,3</sup> CLELAND	82	SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$	OCCUR=3
896.0 ± 1.9	380	DELFOSSÉ	81	SPEC	+	50 $K^\pm p \rightarrow K^\pm \pi^0 p$	
886.0 ± 2.3	187	DELFOSSÉ	81	SPEC	-	50 $K^\pm p \rightarrow K^\pm \pi^0 p$	OCCUR=2
894.2 ± 2.0	765	<sup>2</sup> CLARK	73	HBC	-	3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
894.3 ± 1.5	1150	<sup>2,3</sup> CLARK	73	HBC	-	3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$	OCCUR=2
892.0 ± 2.6	341	<sup>2</sup> SCHWEING...	68	HBC	-	5.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$	OCCUR=2

<sup>1</sup> Inclusive reaction. Complicated background and phase-space effects.

<sup>2</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ . See note.

<sup>3</sup> Number of events in peak reevaluated by us.

<sup>4</sup> From a Dalitz plot analysis in an isobar model with charged and neutral  $K^*$ (892) masses and widths floating.

<sup>5</sup> Average of fit results with different parametrizations for the  $K\pi$  S-wave.

<sup>6</sup> K-matrix pole.

<sup>7</sup> From a partial wave amplitude analysis.

NODE=M018M;LINKAGE=I  
 NODE=M018M;LINKAGE=D  
 NODE=M018M;LINKAGE=W  
 NODE=M018M1;LINKAGE=B

NODE=M018M1;LINKAGE=A  
 NODE=M018M1;LINKAGE=AN  
 NODE=M018M1;LINKAGE=F

NODE=M018MCT  
 NODE=M018MCT

## CHARGED ONLY, PRODUCED IN $\tau$ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>895.47 ± 0.20 ± 0.74</b>	53k	<sup>1</sup> EPIFANOV	07	BELL $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

892.0 ± 0.5		<sup>2</sup> BOITO	10	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
892.0 ± 0.9		<sup>3,4</sup> BOITO	09	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
895.3 ± 0.2		<sup>4,5</sup> JAMIN	08	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
896.4 ± 0.9	12k	<sup>6</sup> BONVICINI	02	CLEO $\tau^- \rightarrow K^- \pi^0 \nu_\tau$
895 ± 2		<sup>7</sup> BARATE	99R	ALEP $\tau^- \rightarrow K^- \pi^0 \nu_\tau$

<sup>1</sup> From a fit in the  $K_0^*(700) + K^*(892) + K^*(1410)$  model.

<sup>2</sup> From the pole position of the  $K\pi$  vector form factor using EPIFANOV 07 and constraints from  $K_{J3}$  decays in ANTONELLI 10.

<sup>3</sup> From the pole position of the  $K\pi$  vector form factor in the complex  $s$ -plane and using EPIFANOV 07 data.

<sup>4</sup> Systematic uncertainties not estimated.

<sup>5</sup> Reanalysis of EPIFANOV 07 using resonance chiral theory.

<sup>6</sup> Calculated by us from the shift by  $4.7 \pm 0.9$  MeV (statistical uncertainty only) reported in BONVICINI 02 with respect to the world average value from PDG 00.

<sup>7</sup> With mass and width of the  $K^*(1410)$  fixed at 1412 MeV and 227 MeV, respectively.

NODE=M018MCT;LINKAGE=EF  
 NODE=M018MCT;LINKAGE=BT

NODE=M018MCT;LINKAGE=BI

NODE=M018MCT;LINKAGE=NS  
 NODE=M018MCT;LINKAGE=JA  
 NODE=M018MCT;LINKAGE=BO

NODE=M018MCT;LINKAGE=BA

## NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>895.55 ± 0.20 OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.		
894.68 ± 0.25 ± 0.05		<sup>1</sup> ABLIKIM	16F	BES3 $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.4 ± 0.2 ± 0.2	243k	<sup>2</sup> DEL-AMO-SA..11I	BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.7 ± 0.2 ± 0.3	141k	<sup>3</sup> BONVICINI	08A	CLEO $D^+ \rightarrow K^- \pi^+ \pi^+$
895.41 ± 0.32 <sup>+0.35</sup> / <sub>-0.43</sub>	18k	<sup>4</sup> LINK	05I	FOCS $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
896 ± 2		BARBERIS	98E	OMEG 450 $pp \rightarrow p_f p_s K^* \bar{K}^*$
895.9 ± 0.5 ± 0.2		ASTON	88	LASS 11 $K^- p \rightarrow K^- \pi^+ n$
894.52 ± 0.63	25k	<sup>5</sup> ATKINSON	86	OMEG 20-70 $\gamma p$

NODE=M018M2  
 NODE=M018M2

894.63±0.76	20k	<sup>5</sup> ATKINSON	86	OMEG	20-70	$\gamma p$		OCCUR=2
897 ±1	28k	EVANGELIS...	80	OMEG	10	$\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$		
898.4 ±1.4	1180	AGUILAR-...	78B	HBC	0.76	$\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$		
894.9 ±1.6		WICKLUND	78	ASPK	3,4,6	$K^\pm N \rightarrow (K\pi)^0 N$		
897.6 ±0.9		BOWLER	77	DBC	5.4	$K^+ d \rightarrow K^+ \pi^- p p$		
895.5 ±1.0	3600	MCCUBBIN	75	HBC	3.6	$K^- p \rightarrow K^- \pi^+ n$		
897.1 ±0.7	22k	<sup>5</sup> PALER	75	HBC	14.3	$K^- p \rightarrow (K\pi)^0 X$		
896.0 ±0.6	10k	FOX	74	RVUE	2	$K^- p \rightarrow K^- \pi^+ n$		
896.0 ±0.6		FOX	74	RVUE	2	$K^+ n \rightarrow K^+ \pi^- p$		OCCUR=2
896 ±2		<sup>6</sup> MATISON	74	HBC	12	$K^+ p \rightarrow K^+ \pi^- \Delta$		
896 ±1	3186	LEWIS	73	HBC	2.1-2.7	$K^+ p \rightarrow K \pi \pi p$		
894.0 ±1.3		<sup>6</sup> LINGLIN	73	HBC	2-13	$K^+ p \rightarrow$ $K^+ \pi^- \pi^+ p$		
898.4 ±1.3	1700	<sup>7</sup> BUCHNER	72	DBC	4.6	$K^+ n \rightarrow K^+ \pi^- p$		
897.9 ±1.1	2934	<sup>7</sup> AGUILAR-...	71B	HBC	3.9,4.6	$K^- p \rightarrow K^- \pi^+ n$		
898.0 ±0.7	5362	<sup>7</sup> AGUILAR-...	71B	HBC	3.9,4.6	$K^- p \rightarrow$ $K^- \pi^+ \pi^- p$		OCCUR=2
895 ±1	4300	<sup>8</sup> HABER	70	DBC	3	$K^- N \rightarrow K^- \pi^+ X$		
893.7 ±2.0	10k	DAVIS	69	HBC	12	$K^+ p \rightarrow K^+ \pi^- \pi^+ p$		
894.7 ±1.4	1040	<sup>7</sup> DAUBER	67B	HBC	2.0	$K^- p \rightarrow K^- \pi^+ \pi^- p$		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●								
895.50±0.92±2.6		<sup>9</sup> ADUSZKIEW...	20A	NA61	158	$p p$		
898.1 ±1.0	4k	<sup>10</sup> LEES	17C	BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$			
895.53±0.17		LEES	13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$			
894.9 ±0.5 ±0.7	14.4k	<sup>11</sup> MITCHELL	09A	CLEO	$D_s^+ \rightarrow K^+ K^- \pi^+$			
896.2 ±0.3	20k	<sup>12</sup> AUBERT	07AK	BABR	$10.6 e^+ e^- \rightarrow$ $K^{*0} K^\pm \pi^\mp \gamma$			
900.7 ±1.1	5900	BARTH	83	HBC	70	$K^+ p \rightarrow K^+ \pi^- X$		

<sup>1</sup> Taking also into account the  $K_0^*(1430)^0$  and  $K_2^*(1430)^0$ .

<sup>2</sup> Taking into account the  $K^*(892)^0$ , S-wave and P-wave ( $K^*(1410)^0$ ).

<sup>3</sup> From the isobar model with a complex pole for the  $\kappa$ .

<sup>4</sup> Fit to  $K \pi$  mass spectrum includes a non-resonant scalar component.

<sup>5</sup> Inclusive reaction. Complicated background and phase-space effects.

<sup>6</sup> From pole extrapolation.

<sup>7</sup> Mass errors enlarged by us to  $\Gamma/\sqrt{N}$ . See note.

<sup>8</sup> Number of events in peak reevaluated by us.

<sup>9</sup> For transverse momenta between 0.6 and 0.8 GeV/c and rapidity  $0 < y < 0.5$ .

<sup>10</sup> From a Dalitz plot analysis in an isobar model with charged and neutral  $K^*(892)$  masses and widths floating.

<sup>11</sup> This value comes from a fit with  $\chi^2$  of 178/117.

<sup>12</sup> Systematic uncertainties not estimated.

NODE=M018M2;LINKAGE=B

NODE=M018M2;LINKAGE=DE

NODE=M018M2;LINKAGE=BO

NODE=M018M2;LINKAGE=LI

NODE=M018M2;LINKAGE=I

NODE=M018M;LINKAGE=C

NODE=M018M2;LINKAGE=D


NODE=M018M2;LINKAGE=W

NODE=M018M2;LINKAGE=C

NODE=M018M2;LINKAGE=A

NODE=M018M2;LINKAGE=MI

NODE=M018M2;LINKAGE=NS



### K\*(892) MASSES AND MASS DIFFERENCES

NODE=M018209

Unrealistically small errors have been reported by some experiments. We use simple “realistic” tests for the minimum errors on the determination of a mass and width from a sample of  $N$  events:

$$\delta_{\min}(m) = \frac{\Gamma}{\sqrt{N}}, \quad \delta_{\min}(\Gamma) = 4 \frac{\Gamma}{\sqrt{N}}. \quad (1)$$

We consistently increase unrealistic errors before averaging. For a detailed discussion, see the 1971 edition of this Note.

$m_{K^*(892)^0} - m_{K^*(892)^\pm}$					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>6.7±1.2 OUR AVERAGE</b>					
7.7±1.7	2980	AGUILAR-...	78B	HBC	±0 0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
5.7±1.7	7338	AGUILAR-...	71B	HBC	-0 3.9,4.6 $K^- p$
6.3±4.1	283	<sup>1</sup> BARASH	67B	HBC	0.0 $\bar{p}p$

<sup>1</sup>Number of events in peak reevaluated by us.

NODE=M018D

NODE=M018D

NODE=M018D;LINKAGE=W

### K\*(892) RANGE PARAMETER

All from partial wave amplitude analyses.

NODE=M018R

NODE=M018R

NODE=M018R

VALUE (GeV <sup>-1</sup> )	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.1 ±0.5 ±0.5	243k	<sup>1</sup> DEL-AMO-SA.11I	BABR	0	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
3.96±0.54 <sup>+1.31</sup> <sub>-0.90</sub>	18k	<sup>2</sup> LINK	05I	FOCS	0 $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
3.4 ±0.7		ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
••• We do not use the following data for averages, fits, limits, etc. •••					
12.1 ±3.2 ±3.0		BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

<sup>1</sup> Taking into account the  $K^*(892)^0$ ,  $S$ -wave and  $P$ -wave ( $K^*(1410)^0$ ).

<sup>2</sup> Fit to  $K\pi$  mass spectrum includes a non-resonant scalar component.

NODE=M018R;LINKAGE=DE  
NODE=M018R;LINKAGE=LI

## $K^*(892)$ WIDTH

NODE=M018215

### CHARGED ONLY, HADROPRODUCED

NODE=M018W1  
NODE=M018W1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

#### 51.4±0.8 OUR FIT

#### 51.4±0.8 OUR AVERAGE

54.4±0.9±1.7		ALBRECHT	20	CBAR	0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
49 ±2	5840	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
56 ±4		NAPIER	84	SPEC	- 200 $\pi^- p \rightarrow 2K_S^0 X$
51 ±2	4100	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
50.5±5.6		AJINENKO	80	HBC	+ 32 $K^+ p \rightarrow K^0 \pi^+ X$
45.8±3.6	1800	AGUILAR-...	78B	HBC	± 0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
52.0±2.5	6706	<sup>1</sup> COOPER	78	HBC	± 0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$
52.1±2.2	9000	<sup>2</sup> PALER	75	HBC	- 14.3 $K^- p \rightarrow (K\pi)^- X$
46.3±6.7	765	<sup>1</sup> CLARK	73	HBC	- 3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$
48.2±5.7	1150	<sup>1,3</sup> CLARK	73	HBC	- 3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$
54.3±3.3	4404	<sup>1</sup> AGUILAR-...	71B	HBC	- 3.9,4.6 $K^- p \rightarrow (K\pi)^- p$
46 ±5	1700	<sup>1,3</sup> WOJCICKI	64	HBC	- 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.7±0.2 <sup>+0.1</sup> <sub>-0.2</sub>	183k	ABLIKIM	19AQ	BES	± $J/\psi \rightarrow K^+ K^- \pi^0$
43.6±1.3	4k	<sup>4</sup> LEES	17C	BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
47.2±0.3±2.3	190k	<sup>5</sup> AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
54.8±1.7	27k	<sup>6</sup> ABELE	99D	CBAR	± 0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$
45.2±1 ±2	80k	<sup>7</sup> BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
42.8±7.1	3700	BARTH	83	HBC	+ 70 $K^+ p \rightarrow K^0 \pi^+ X$
64.0±9.2	800	<sup>1,3</sup> CLELAND	82	SPEC	+ 30 $K^+ p \rightarrow K_S^0 \pi^+ p$
62.0±4.4	3200	<sup>1,3</sup> CLELAND	82	SPEC	+ 50 $K^+ p \rightarrow K_S^0 \pi^+ p$
55 ±4	3600	<sup>1,3</sup> CLELAND	82	SPEC	- 50 $K^+ p \rightarrow K_S^0 \pi^- p$
62.6±3.8	380	DELFOSSÉ	81	SPEC	+ 50 $K^\pm p \rightarrow K^\pm \pi^0 p$
50.5±3.9	187	DELFOSSÉ	81	SPEC	- 50 $K^\pm p \rightarrow K^\pm \pi^0 p$

OCCUR=2

OCCUR=3

OCCUR=2

<sup>1</sup> Width errors enlarged by us to  $4 \times \Gamma/\sqrt{N}$ ; see note.

<sup>2</sup> Inclusive reaction. Complicated background and phase-space effects.

<sup>3</sup> Number of events in peak reevaluated by us.

<sup>4</sup> From a Dalitz plot analysis in an isobar model with charged and neutral  $K^*(892)$  masses and widths floating.

<sup>5</sup> Average of fit results with different parametrizations for the  $K\pi$   $S$ -wave.

<sup>6</sup>  $K$ -matrix pole.

<sup>7</sup> From a partial wave amplitude analysis.

NODE=M018W;LINKAGE=D  
NODE=M018W;LINKAGE=I  
NODE=M018W;LINKAGE=W  
NODE=M018W1;LINKAGE=B

NODE=M018W1;LINKAGE=A  
NODE=M018W1;LINKAGE=AN  
NODE=M018W1;LINKAGE=F

### CHARGED ONLY, PRODUCED IN $\tau$ LEPTON DECAYS

NODE=M018W5  
NODE=M018W5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**46.2±0.6±1.2**      53k      <sup>1</sup> EPIFANOV      07      BELL       $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.5±1.1		<sup>2</sup> BOITO	10	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
46.2±0.4		<sup>3,4</sup> BOITO	09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
47.5±0.4		<sup>4,5</sup> JAMIN	08	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
55 ±8		<sup>6</sup> BARATE	99R	ALEP	$\tau^- \rightarrow K^- \pi^0 \nu_\tau$

<sup>1</sup> From a fit in the  $K_0^*(700) + K^*(892) + K^*(1410)$  model.

<sup>2</sup> From the pole position of the  $K\pi$  vector form factor using EPIFANOV 07 and constraints from  $K_{J3}$  decays in ANTONELLI 10.

<sup>3</sup> From the pole position of the  $K\pi$  vector form factor in the complex  $s$ -plane and using EPIFANOV 07 data.

<sup>4</sup> Systematic uncertainties not estimated.

<sup>5</sup> Reanalysis of EPIFANOV 07 using resonance chiral theory.

<sup>6</sup> With mass and width of the  $K^*(1410)$  fixed at 1412 MeV and 227 MeV, respectively.

NODE=M018W5;LINKAGE=EF  
NODE=M018W5;LINKAGE=BT

NODE=M018W5;LINKAGE=BI

NODE=M018W5;LINKAGE=NS  
NODE=M018W5;LINKAGE=JA  
NODE=M018W5;LINKAGE=BA

**NEUTRAL ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M018W2  
 NODE=M018W2

**47.3 ±0.5 OUR FIT** Error includes scale factor of 1.9.

**47.3 ±0.5 OUR AVERAGE** Error includes scale factor of 2.0. See the ideogram below.

46.53 ±0.56 ±0.31		<sup>1</sup> ABLIKIM	16F BES3	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$	
46.5 ±0.3 ±0.2	243k	<sup>2</sup> DEL-AMO-SA..	11I BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$	
45.3 ±0.5 ±0.6	141k	<sup>3</sup> BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$	
47.79 ±0.86 <sup>+1.32</sup> <sub>-1.06</sub>	18k	<sup>4</sup> LINK	05I FOCS	$D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$	
54 ±3		BARBERIS	98E OMEG	450 $pp \rightarrow p_f p_s K^* \bar{K}^*$	
50.8 ±0.8 ±0.9		ASTON	88 LASS	11 $K^- p \rightarrow K^- \pi^+ n$	
46.5 ±4.3	5900	BARTH	83 HBC	70 $K^+ p \rightarrow K^+ \pi^- X$	
54 ±2	28k	EVANGELIS...	80 OMEG	10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$	
45.9 ±4.8	1180	AGUILAR-...	78B HBC	0.76 $\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$	
51.2 ±1.7		WICKLUND	78 ASPK	3,4,6 $K^\pm N \rightarrow (K\pi)^0 N$	
48.9 ±2.5		BOWLER	77 DBC	5.4 $K^+ d \rightarrow K^+ \pi^- pp$	
48 <sup>+3</sup> <sub>-2</sub>	3600	MCCUBBIN	75 HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$	
50.6 ±2.5	22k	<sup>5</sup> PALER	75 HBC	14.3 $K^- p \rightarrow (K\pi)^0 X$	
47 ±2	10k	FOX	74 RVUE	2 $K^- p \rightarrow K^- \pi^+ n$	
51 ±2		FOX	74 RVUE	2 $K^+ n \rightarrow K^+ \pi^- p$	OCCUR=2
46.0 ±3.3	3186	<sup>6</sup> LEWIS	73 HBC	2.1-2.7 $K^+ p \rightarrow K \pi \pi p$	
51.4 ±5.0	1700	<sup>6</sup> BUCHNER	72 DBC	4.6 $K^+ n \rightarrow K^+ \pi^- p$	
55.8 <sup>+4.2</sup> <sub>-3.4</sub>	2934	<sup>6</sup> AGUILAR-...	71B HBC	3.9,4.6 $K^- p \rightarrow K^- \pi^+ n$	
48.5 ±2.7	5362	AGUILAR-...	71B HBC	3.9,4.6 $K^- p \rightarrow$ $K^- \pi^+ \pi^- p$	OCCUR=2
54.0 ±3.3	4300	<sup>6,7</sup> HABER	70 DBC	3 $K^- N \rightarrow K^- \pi^+ X$	
53.2 ±2.1	10k	<sup>6</sup> DAVIS	69 HBC	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$	
44 ±5.5	1040	<sup>6</sup> DAUBER	67B HBC	2.0 $K^- p \rightarrow K^- \pi^+ \pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

48.8 ±1.8 ±2.0		<sup>8</sup> ADUSZKIEW...	20A NA61	158 $pp$	
52.6 ±1.7	4k	<sup>9</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	
44.90 ±0.30		LEES	13F BABR	$D^+ \rightarrow K^+ K^- \pi^+$	
45.7 ±1.1 ±0.5	14.4k	<sup>10</sup> MITCHELL	09A CLEO	$D_s^+ \rightarrow K^+ K^- \pi^+$	
50.6 ±0.9	20k	<sup>11</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow$ $K^{*0} K^\pm \pi^\mp \gamma$	

<sup>1</sup> Taking also into account the  $K_0^*(1430)^0$  and  $K_2^*(1430)^0$ .

<sup>2</sup> Taking into account the  $K^*(892)^0$ ,  $S$ -wave and  $P$ -wave ( $K^*(1410)^0$ ).

<sup>3</sup> From the isobar model with a complex pole for the  $\kappa$ .

<sup>4</sup> Fit to  $K\pi$  mass spectrum includes a non-resonant scalar component.

<sup>5</sup> Inclusive reaction. Complicated background and phase-space effects.

<sup>6</sup> Width errors enlarged by us to  $4 \times \Gamma/\sqrt{N}$ ; see note.

<sup>7</sup> Number of events in peak reevaluated by us.

<sup>8</sup> For transverse momenta between 0.6 and 0.8 GeV/c and rapidity  $0 < y < 0.5$ .

<sup>9</sup> From a Dalitz plot analysis in an isobar model with charged and neutral  $K^*(892)$  masses and widths floating.

<sup>10</sup> This value comes from a fit with  $\chi^2$  of 178/117.


<sup>11</sup> Systematic uncertainties not estimated.

NODE=M018W2;LINKAGE=B  
 NODE=M018W2;LINKAGE=DE  
 NODE=M018W2;LINKAGE=BO  
 NODE=M018W2;LINKAGE=LN  
 NODE=M018W2;LINKAGE=I  
 NODE=M018W2;LINKAGE=D  
 NODE=M018W2;LINKAGE=W  
 NODE=M018W2;LINKAGE=C  
 NODE=M018W2;LINKAGE=A

NODE=M018W2;LINKAGE=MI  
 NODE=M018W2;LINKAGE=NS



WEIGHTED AVERAGE  
47.3±0.5 (Error scaled by 2.0)



**K\*(892) DECAY MODES**

NODE=M018220;NODE=M018

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $K\pi$	$\sim 100$	%
$\Gamma_2$ $(K\pi)^\pm$	$(99.902 \pm 0.009) \%$	
$\Gamma_3$ $(K\pi)^0$	$(99.754 \pm 0.021) \%$	
$\Gamma_4$ $K^0\gamma$	$(2.46 \pm 0.21) \times 10^{-3}$	
$\Gamma_5$ $K^\pm\gamma$	$(9.8 \pm 0.9) \times 10^{-4}$	
$\Gamma_6$ $K\pi\pi$	$< 7 \times 10^{-4}$	95%

DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=11  
DESIG=12  
DESIG=4  
DESIG=3  
DESIG=2

**CONSTRAINED FIT INFORMATION**

An overall fit to the total width and a partial width uses 14 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 10.7$  for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$\begin{array}{c}
 x_5 \\
 \Gamma
 \end{array}
 \begin{array}{|c|}
 \hline
 \begin{array}{cc}
 -100 & \\
 17 & -17 \\
 \hline
 x_2 & x_5
 \end{array}
 \end{array}$$

Mode	Rate (MeV)
$\Gamma_2$ $(K\pi)^\pm$	51.4 ± 0.8
$\Gamma_5$ $K^\pm\gamma$	0.050 ± 0.005

DESIG=11  
DESIG=3

## CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 23 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 68.4$  for 21 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_4$	-100	
$\Gamma$	12	-12
	$x_3$	$x_4$

Mode	Rate (MeV)	Scale factor
$\Gamma_3$ ( $K\pi^0$ )	47.2 $\pm$ 0.5	1.9
$\Gamma_4$ ( $K^0\gamma$ )	0.117 $\pm$ 0.010	

DESIG=12

DESIG=4

### $K^*(892)$ PARTIAL WIDTHS

NODE=M018225

$\Gamma(K^0\gamma)$	$\Gamma_4$				
VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>116 <math>\pm</math> 10 OUR FIT</b>					
116.5 $\pm$ 9.9	584	CARLSMITH	86	SPEC	0 $K_L^0 A \rightarrow K_S^0 \pi^0 A$

NODE=M018W4  
NODE=M018W4

$\Gamma(K^\pm\gamma)$	$\Gamma_5$			
VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>50 <math>\pm</math> 5 OUR FIT</b>				
50 $\pm$ 5 OUR AVERAGE				
48 $\pm$ 11	BERG	83	SPEC	- 156 $K^- A \rightarrow \bar{K} \pi A$
51 $\pm$ 5	CHANDLEE	83	SPEC	+ 200 $K^+ A \rightarrow K \pi A$

NODE=M018W3  
NODE=M018W3

### $K^*(892)$ BRANCHING RATIOS

NODE=M018230

$\Gamma(K^0\gamma)/\Gamma_{\text{total}}$	$\Gamma_4/\Gamma$			
VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	CHG	COMMENT
<b>2.46 <math>\pm</math> 0.21 OUR FIT</b>				
2.46 $\pm$ 0.21	CARITHERS	75B	CNTR	0 8-16 $\bar{K}^0 A$

NODE=M018R3  
NODE=M018R3

$\Gamma(K^\pm\gamma)/\Gamma_{\text{total}}$	$\Gamma_5/\Gamma$				
VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.98 <math>\pm</math> 0.09 OUR FIT</b>					
<1.6	95	BEMPORAD	73	CNTR	+ 10-16 $K^+ A$

NODE=M018R2  
NODE=M018R2

$\Gamma(K\pi\pi)/\Gamma((K\pi)^\pm)$	$\Gamma_6/\Gamma_2$				
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt; 7 <math>\times</math> 10<sup>-4</sup></b>	95	JONGEJANS	78	HBC	4 $K^- p \rightarrow p \bar{K}^0 2\pi$
<20 $\times$ 10 <sup>-4</sup>		WOJCICKI	64	HBC	- 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NODE=M018R1  
NODE=M018R1

### $K^*(892)$ REFERENCES

NODE=M018

ADUSZKIEW... 20A	EPJ C80 460	A. Aduszkiewicz <i>et al.</i>	(CERN NA61 Collab.)	REFID=60631
ALBRECHT 20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
PELAEZ 20	PRL 124 172001	J.R. Pelaez <i>et al.</i>		REFID=60559
ABLIKIM 19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
LEES 17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
PELAEZ 17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira		REFID=57836
AAIJ 16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM 16F	PR D94 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57307
LEES 13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55127
DEL-AMO-SA... 11I	PR D83 072001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16493
ANTONELLI 10	EPJ C69 399	M. Antonelli <i>et al.</i>	(FlaviaNet Working Group)	REFID=53448
BOITO 10	JHEP 1009 031	D.R. Boito, R. Escribano, M. Jamin	(BARC)	REFID=53632
BOITO 09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin		REFID=52728

MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52756
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
JAMIN	08	PL B664 78	M. Jamin, A. Pich, J. Portoles		REFID=52285
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)	REFID=51929
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=50679
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
BONVICINI	02	PRL 88 111803	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=48701
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47366
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46348
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
CARLSMITH	86	PRL 56 18	D. Carlsmith <i>et al.</i>	(EFI, SACL)	REFID=22461
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
NAPIER	84	PL 149B 514	A. Napier <i>et al.</i>	(TUFTS, ARIZ, FNAL, FLOR+)	REFID=22460
BARTH	83	NP B223 296	M. Barth <i>et al.</i>	(BRUX, CERN, GENO, MONS+)	REFID=22456
BERG	83	Thesis UMI 83-21652	D.M. Berg	(ROCH)	REFID=22457
CHANDLEE	83	PRL 51 168	C. Handlee <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=22458
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
AJINENKO	80	ZPHY C5 177	I.V. Ajinenko <i>et al.</i>	(SERP, BRUX, MONS+)	REFID=22449
EVANGELIS...	80	NP B165 383	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=22450
AGUILAR-...	78B	NP B141 101	M. Aguilar-Benitez <i>et al.</i>	(MADR, TATA+)	REFID=22438
BALAND	78	NP B140 220	J.F. Baland <i>et al.</i>	(MONS, BELG, CERN+)	REFID=20369
COOPER	78	NP B136 365	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=22441
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
JONGEJANS	78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
BOWLER	77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
CARITHERS	75B	PRL 35 349	W.C.J. Carithers <i>et al.</i>	(ROCH, MCGI)	REFID=22433
MCCUBBIN	75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
PALER	75	NP B96 1	K. Paler <i>et al.</i>	(RHEL, SACL, EPOL)	REFID=22435
FOX	74	NP B80 403	G.C. Fox, M.L. Griss	(CIT)	REFID=22430
MATISON	74	PR D9 1872	M.J. Matison <i>et al.</i>	(LBL)	REFID=22431
BEMPORAD	73	NP B51 1	C. Bemporad <i>et al.</i>	(CERN, ETH, LOIC)	REFID=22416
CLARK	73	NP B54 432	A.G. Clark, L. Lyons, D. Radojicic	(OXF)	REFID=22426
LEWIS	73	NP B60 283	P.H. Lewis <i>et al.</i>	(LOWC, LOIC, CDEF)	REFID=22427
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
BUCHNER	72	NP B45 333	K. Buchner <i>et al.</i>	(MPIM, CERN, BRUX)	REFID=22418
AGUILAR-...	71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
HABER	70	NP B17 289	B. Haber <i>et al.</i>	(REHO, SACL, BGNA, EPOL)	REFID=22406
CRENNELL	69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS	69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
SCHWEING...	68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
DAUBER	67B	PR 153 1403	P.M. Dauber <i>et al.</i>	(UCLA)	REFID=22389
DEBAERE	67B	NC 51A 401	W. de Baere <i>et al.</i>	(BRUX, CERN)	REFID=22390
WOJCICKI	64	PR 135 B484	S.G. Wojcicki	(LRL)	REFID=22379

**$K_1(1270)$** 

$$I(J^P) = \frac{1}{2}(1^+)$$


NODE=M028

 **$K_1(1270)$  MASS**

NODE=M028MX

NODE=M028MX

VALUE (MeV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_

**1253±7 OUR AVERAGE** Includes data from the 4 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.**PRODUCED BY  $K^-$ , BACKWARD SCATTERING, HYPERON EXCHANGE**NODE=M028M2  
NODE=M028M2

VALUE (MeV) \_\_\_\_\_ EVTS \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ CHG \_\_\_\_\_ COMMENT \_\_\_\_\_

The data in this block is included in the average printed for a previous datablock.

**1275±10** 700 GAVILLET 78 HBC + 4.2  $K^- p \rightarrow \Xi^- (K\pi\pi)^+$ **PRODUCED BY K BEAMS**NODE=M028M3  
NODE=M028M3

VALUE (MeV) \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ CHG \_\_\_\_\_ COMMENT \_\_\_\_\_

The data in this block is included in the average printed for a previous datablock.

**1270±10** <sup>1</sup> DAUM 81C CNTR - 63  $K^- p \rightarrow K^- 2\pi p$

••• We do not use the following data for averages, fits, limits, etc. •••

~ 1276 <sup>2</sup> TORNQVIST 82B RVUE

~ 1300 VERGEEST 79 HBC - 4.2  $K^- p \rightarrow (\bar{K}\pi\pi)^- p$

1289±25 <sup>3</sup> CARNEGIE 77 ASPK ± 13  $K^\pm p \rightarrow (K\pi\pi)^\pm p$

~ 1300 BRANDENB... 76 ASPK ± 13  $K^\pm p \rightarrow (K\pi\pi)^\pm p$

~ 1270 OTTER 76 HBC - 10,14,16  $K^- p \rightarrow (\bar{K}\pi\pi)^- p$

1260 DAVIS 72 HBC + 12  $K^+ p$

1234±12 FIRESTONE 72B DBC + 12  $K^+ d$

<sup>1</sup> Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.<sup>2</sup> From a unitarized quark-model calculation.<sup>3</sup> From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

NODE=M028M3;LINKAGE=DA

NODE=M028M3;LINKAGE=T

NODE=M028M3;LINKAGE=E

**PRODUCED BY BEAMS OTHER THAN K MESONS**NODE=M028M1  
NODE=M028M1

VALUE (MeV) \_\_\_\_\_ EVTS \_\_\_\_\_ DOCUMENT ID \_\_\_\_\_ TECN \_\_\_\_\_ COMMENT \_\_\_\_\_

The data in this block is included in the average printed for a previous datablock.

**1248.1 ± 3.3 ± 1.4** GULER 11 BELL  $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1289.81 ± 0.56 ± 1.66	894k	AAIJ	18A1	LHCB	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
1279 ± 10	25k	<sup>1</sup> ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
1294 ± 10	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
1300	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
1242 +9 -10		<sup>2</sup> ASTIER	69	HBC	$\bar{p} p$
1300	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

<sup>1</sup> Systematic errors not estimated.  
<sup>2</sup> This was called the C meson.

NODE=M028M1;LINKAGE=AB  
 NODE=M028M1;LINKAGE=A

**PRODUCED IN  $\tau$  LEPTON DECAYS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M028MT  
 NODE=M028MT

<b>1254 ± 33 ± 34</b>	7k	ASNER	00B	CLEO	$\pm \tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
-----------------------	----	-------	-----	------	---

**$K_1(1270)$  WIDTH**


NODE=M028WX

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**90 ± 20 OUR ESTIMATE** This is only an educated guess; the error given is larger than the error on the average of the published values.

NODE=M028WX  
 → UNCHECKED ←

**101 ± 12 OUR AVERAGE** Includes data from the 4 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.



**PRODUCED BY  $K^-$ , BACKWARD SCATTERING, HYPERON EXCHANGE**

NODE=M028W2  
 NODE=M028W2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

<b>75 ± 15</b>	700	GAVILLET	78	HBC	+ 4.2 $K^- p \rightarrow \Xi^- K \pi \pi$
----------------	-----	----------	----	-----	---

**PRODUCED BY K BEAMS**

NODE=M028W3  
 NODE=M028W3

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

<b>90 ± 8</b>	<sup>1</sup> DAUM	81C	CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
---------------	-------------------	-----	------	---	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 150	VERGEEST	79	HBC	-	4.2 $K^- p \rightarrow (\bar{K} \pi \pi)^- p$
150 ± 71	<sup>2</sup> CARNEGIE	77	ASPK	±	13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$
~ 200	BRANDENB...	76	ASPK	±	13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$
120	DAVIS	72	HBC	+	12 $K^+ p$
188 ± 21	FIRESTONE	72B	DBC	+	12 $K^+ d$

<sup>1</sup> Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

NODE=M028W3;LINKAGE=DA

<sup>2</sup> From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

NODE=M028W3;LINKAGE=E

**PRODUCED BY BEAMS OTHER THAN K MESONS**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

<b>119.5 ± 5.2 ± 6.7</b>		GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
--------------------------	--	-------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

116.11 ± 1.65 ± 2.96	894k	AAIJ	18A1	LHCB	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
131 ± 21	25k	<sup>1</sup> ABLIIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
66 ± 15	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
127 <sup>+7</sup> / <sub>-25</sub>		ASTIER	69	HBC	$\bar{p} p$
60	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

<sup>1</sup> Systematic errors not estimated.NODE=M028W1  
NODE=M028W1

NODE=M028W1;LINKAGE=AB

**PRODUCED IN  $\tau$  LEPTON DECAYS**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

<b>260 <sup>+90</sup>/<sub>-70</sub> ± 80</b>	7k	ASNER	00B	CLEO	$\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
---	----	-------	-----	------	---

NODE=M028WT  
NODE=M028WT **$K_1(1270)$  DECAY MODES**

NODE=M028215;NODE=M028

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor
$\Gamma_1$ $K\rho$	(38 ± 13) %	2.2
$\Gamma_2$ $K_0^*(1430)\pi$	(28 ± 4) %	
$\Gamma_3$ $K^*(892)\pi$	(21 ± 10) %	2.2
$\Gamma_4$ $K\omega$	(11.0 ± 2.0) %	
$\Gamma_5$ $K f_0(1370)$	(3.0 ± 2.0) %	
$\Gamma_6$ $\gamma K^0$	seen	

DESIG=2  
DESIG=7  
DESIG=1  
DESIG=5  
DESIG=8  
DESIG=9;OUR EST;→ UNCHECKED ← **$K_1(1270)$  PARTIAL WIDTHS**

NODE=M028220

 **$\Gamma(K\rho)$**  **$\Gamma_1$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

57 ± 5	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
75 ± 6	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W5  
NODE=M028W5 **$\Gamma(K_0^*(1430)\pi)$**  **$\Gamma_2$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

26 ± 6	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
--------	----------	-----	------	---	--

NODE=M028W7  
NODE=M028W7 **$\Gamma(K^*(892)\pi)$**  **$\Gamma_3$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 11	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
2 ± 2	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W4  
NODE=M028W4 **$\Gamma(K\omega)$**  **$\Gamma_4$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 4	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
24 ± 3	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W6  
NODE=M028W6 **$\Gamma(K f_0(1370))$**  **$\Gamma_5$** 

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 5	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
--------	----------	-----	------	---	--

NODE=M028W8  
NODE=M028W8

$\Gamma(\gamma K^0)$  $\Gamma_6$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>73.2±6.1±28.3</b>	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$

NODE=M028W9  
 NODE=M028W9

 $K_1(1270)$  BRANCHING RATIOS

NODE=M028225

 $\Gamma(K\rho)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.38 ±0.13 OUR FIT** Error includes scale factor of 2.2.**0.42 ±0.06** <sup>1</sup> DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.584±0.043 <sup>2</sup> GULER 11 BELL  $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$   
dominant RODEBACK 81 HBC  $4 \pi^- p \rightarrow \Lambda K 2\pi$ 

NODE=M028R2  
 NODE=M028R2

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.28 ±0.04** <sup>1</sup> DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0201±0.0064 <sup>2</sup> GULER 11 BELL  $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$ 

NODE=M028R4  
 NODE=M028R4

 $\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.21 ±0.10 OUR FIT** Error includes scale factor of 2.2.**0.16 ±0.05** <sup>1</sup> DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.171±0.023 <sup>2</sup> GULER 11 BELL  $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$ 

NODE=M028R1  
 NODE=M028R1

 $\Gamma(K^*(892)\pi)/\Gamma(K\rho)$  $\Gamma_3/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.56±0.29 OUR FIT** Error includes scale factor of 2.2.**0.99±0.15±0.18** ABLIKIM 21U BES3  $D_s^+ \rightarrow \bar{K}_1(1270)^0 K^+$ 

NODE=M028R00  
 NODE=M028R00

 $\Gamma(K\omega)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.11 ±0.02** <sup>1</sup> DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.225±0.052 <sup>2</sup> GULER 11 BELL  $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$ 

NODE=M028R3  
 NODE=M028R3

 $\Gamma(K\omega)/\Gamma(K\rho)$  $\Gamma_4/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.30 95 RODEBACK 81 HBC  $4 \pi^- p \rightarrow \Lambda K 2\pi$ 

NODE=M028R6  
 NODE=M028R6

 $\Gamma(K f_0(1370))/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.03±0.02** <sup>1</sup> DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$ 

NODE=M028R5  
 NODE=M028R5

**D-wave/S-wave RATIO FOR  $K_1(1270) \rightarrow K^*(892)\pi$** 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**1.0±0.7** <sup>1</sup> DAUM 81C CNTR 63  $K^- p \rightarrow K^- 2\pi p$ 

NODE=M028R9  
 NODE=M028R9

<sup>1</sup> Average from low and high  $t$  data.<sup>2</sup> Assuming that decays are saturated by the  $K\rho$ ,  $K_0^*(1430)\pi$ ,  $K^*(892)\pi$ ,  $K\omega$  decay modes and neglecting interference between them. The values  $B(\omega \rightarrow \pi^+ \pi^-) = (1.53^{+0.11}_{-0.13})\%$  and  $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10)\%$  are used. Systematic uncertainties not estimated.

NODE=M028R;LINKAGE=F  
 NODE=M028R1;LINKAGE=GU

**$K_1(1270)$  REFERENCES**

ABLIKIM	21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18A1	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
GENG	07	PR D75 014017	L.S. Geng <i>et al.</i>	
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
RODEBACK	81	ZPHY C9 9	S. Rodeback <i>et al.</i>	(CERN, CDEF, MADR+)
MAZZUCATO	79	NP B156 532	M. Mazzucato <i>et al.</i>	(CERN, ZEEM, NIJM+)
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
GAVILLET	78	PL 76B 517	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+)
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
CARNEGIE	77B	PL 68B 287	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
OTTER	76	NP B106 77	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
CRENNELL	72	PR D6 1220	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)
ASTIER	69	NP B10 65	A. Astier <i>et al.</i>	(CDEF, CERN, IPNP, LIVP) JJP
CRENNELL	67	PRL 19 44	D.J. Crennell <i>et al.</i>	(BNL) I

NODE=M028

REFID=61155  
 REFID=59187  
 REFID=53668  
 REFID=51623  
 REFID=51037  
 REFID=48822  
 REFID=47766  
 REFID=20573  
 REFID=22548  
 REFID=22550  
 REFID=20867  
 REFID=22542  
 REFID=22538  
 REFID=22535  
 REFID=22536  
 REFID=22532  
 REFID=22533  
 REFID=22419  
 REFID=22505  
 REFID=22506  
 REFID=22482  
 REFID=22473

NODE=M064

 **$K_1(1400)$** 

$$I(J^P) = \frac{1}{2}(1^+)$$

 **$K_1(1400)$  MASS**

NODE=M064M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1403 ± 7 OUR AVERAGE</b>					
1463 ± 64 ± 68	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
1373 ± 14 ± 18		<sup>1</sup> ASTON	87	LASS	0 $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1392 ± 18		BAUBILLIER	82B	HBC	0 $8.25 K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
1410 ± 25		DAUM	81C	CNTR	- $63 K^- p \rightarrow K^- 2\pi p$
1415 ± 15		ETKIN	80	MPS	0 $6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1404 ± 10		<sup>2</sup> CARNEGIE	77	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1418 ± 8	25k	<sup>3</sup> ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 1350		<sup>4</sup> TORNQVIST	82B	RVUE	
~ 1400		VERGEEST	79	HBC	- $4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 1400		BRANDENB...	76	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
1420		DAVIS	72	HBC	+ $12 K^+ p$
1368 ± 18		FIRESTONE	72B	DBC	+ $12 K^+ d$

NODE=M064M

NODE=M064M

<sup>1</sup> From partial-wave analysis of  $K^0 \pi^+ \pi^-$  system.<sup>2</sup> From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.<sup>3</sup> Systematic errors not estimated.<sup>4</sup> From a unitarized quark-model calculation.

NODE=M064M;LINKAGE=P  
 NODE=M064M;LINKAGE=E  
 NODE=M064M;LINKAGE=AB  
 NODE=M064M;LINKAGE=T

 **$K_1(1400)$  WIDTH**

NODE=M064W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>174 ± 13 OUR AVERAGE</b>					
Error includes scale factor of 1.6. See the ideogram below.					
$300^{+370}_{-110} \pm 140$	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
188 ± 54 ± 60		<sup>1</sup> ASTON	87	LASS	0 $11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
276 ± 65		BAUBILLIER	82B	HBC	0 $8.25 K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
195 ± 25		DAUM	81C	CNTR	- $63 K^- p \rightarrow K^- 2\pi p$
180 ± 10		ETKIN	80	MPS	0 $6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
142 ± 16		<sup>2</sup> CARNEGIE	77	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
152 ± 16	25k	<sup>3</sup> ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 200		VERGEEST	79	HBC	- $4.2 K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 160		BRANDENB...	76	ASPK	± $13 K^\pm p \rightarrow (K\pi\pi)^\pm p$
80		DAVIS	72	HBC	+ $12 K^+ p$
241 ± 30		FIRESTONE	72B	DBC	+ $12 K^+ d$


NODE=M064W

NODE=M064W



- <sup>1</sup> From partial-wave analysis of  $K^0 \pi^+ \pi^-$  system.
- <sup>2</sup> From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.
- <sup>3</sup> Systematic errors not estimated.

NODE=M064W;LINKAGE=P  
 NODE=M064W;LINKAGE=E  
 NODE=M064W;LINKAGE=AB



### $K_1(1400)$ DECAY MODES

NODE=M064215;NODE=M064

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\pi$	(94 $\pm$ 6) %
$\Gamma_2$ $K\rho$	( 3.0 $\pm$ 3.0) %
$\Gamma_3$ $K f_0(1370)$	( 2.0 $\pm$ 2.0) %
$\Gamma_4$ $K\omega$	( 1.0 $\pm$ 1.0) %
$\Gamma_5$ $K_0^*(1430)\pi$	not seen
$\Gamma_6$ $\gamma K^0$	seen
$\Gamma_7$ $K\phi$	seen

DESIG=1  
 DESIG=2  
 DESIG=8  
 DESIG=5  
 DESIG=7;OUR EST;→ UNCHECKED ←  
 DESIG=9;OUR EST;→ UNCHECKED ←  
 DESIG=10

### $K_1(1400)$ PARTIAL WIDTHS

NODE=M064220

$\Gamma(K^*(892)\pi)$					$\Gamma_1$
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
<b>117<math>\pm</math>10</b>	CARNEGIE	77	ASPK	$\pm$ 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	
$\Gamma(K\rho)$					$\Gamma_2$
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
<b>2<math>\pm</math>1</b>	CARNEGIE	77	ASPK	$\pm$ 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	
$\Gamma(K\omega)$					$\Gamma_4$
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
<b>23<math>\pm</math>12</b>	CARNEGIE	77	ASPK	$\pm$ 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	
$\Gamma(\gamma K^0)$					$\Gamma_6$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
<b>280.8<math>\pm</math>23.2<math>\pm</math>40.4</b>	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$		

NODE=M064W1  
 NODE=M064W1  
 NODE=M064W2  
 NODE=M064W2  
 NODE=M064W5  
 NODE=M064W5  
 NODE=M064W6  
 NODE=M064W6

### $K_1(1400)$ BRANCHING RATIOS

NODE=M064225

$\Gamma(K^*(892)\pi)/\Gamma_{total}$				$\Gamma_1/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT	
<b>0.94<math>\pm</math>0.06</b>	<sup>1</sup> DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064R1  
 NODE=M064R1

<sup>1</sup> Average from low and high  $t$  data.

NODE=M064R1;LINKAGE=F

$\Gamma(K\rho)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.03 \pm 0.03$	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

<sup>1</sup> Average from low and high  $t$  data. $\Gamma_2/\Gamma$ NODE=M064R2  
NODE=M064R2

NODE=M064R2;LINKAGE=F

 $\Gamma(K f_0(1370))/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.02 \pm 0.02$	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

<sup>1</sup> Average from low and high  $t$  data. $\Gamma_3/\Gamma$ NODE=M064R5  
NODE=M064R5

NODE=M064R5;LINKAGE=F

 $\Gamma(K\omega)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.01 \pm 0.01$	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

<sup>1</sup> Average from low and high  $t$  data. $\Gamma_4/\Gamma$ NODE=M064R3  
NODE=M064R3

NODE=M064R3;LINKAGE=F

 $\Gamma(K\phi)/\Gamma_{\text{total}}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	<sup>1</sup> AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 9.2  $\sigma$ . $\Gamma_7/\Gamma$ NODE=M064R00  
NODE=M064R00

NODE=M064R00;LINKAGE=A

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

<sup>1</sup> Average from low and high  $t$  data. $\Gamma_5/\Gamma$ NODE=M064R4  
NODE=M064R4

NODE=M064R4;LINKAGE=F

**D-wave/S-wave RATIO FOR  $K_1(1400) \rightarrow K^*(892)\pi$** 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.04 \pm 0.01$	<sup>1</sup> DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

<sup>1</sup> Average from low and high  $t$  data.NODE=M064R9  
NODE=M064R9

NODE=M064R9;LINKAGE=F

 **$K_1(1400)$  REFERENCES**

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)	REFID=48822
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47766
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)	REFID=20573
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)	REFID=22542
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)	REFID=22535
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP	REFID=22532
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)	REFID=22505
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)	REFID=22506

NODE=M064

**$K^*(1410)$** 

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M094

 **$K^*(1410)$  T-MATRIX POLE  $\sqrt{s}$** 

NODE=M094PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M094PP

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(1368 \pm 38) - i(106^{+48}_{-59})$	<sup>1</sup> PELAEZ	17	RVUE $\pi K \rightarrow \pi K$
--------------------------------------	---------------------	----	--------------------------------

OCCUR=2

<sup>1</sup> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M094PP;LINKAGE=B

 **$K^*(1410)$  MASS**

NODE=M094M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M094M

**1414 ± 15 OUR AVERAGE** Error includes scale factor of 1.3.

$1380 \pm 21 \pm 19$		ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
$1420 \pm 7 \pm 10$		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1437 \pm 8 \pm 16$	190k	<sup>1</sup> AAIJ	16N	LHCB		$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
---------------------	------	-------------------	-----	------	--	---

OCCUR=2

$1426 \pm 8 \pm 24$	190k	<sup>2</sup> AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
---------------------	------	-------------------	-----	------	--	---

$1276^{+72}_{-77}$		<sup>3,4</sup> BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
--------------------	--	----------------------	----	------	--	---

$1367 \pm 54$		BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
---------------	--	------	----	------	---	--

$1474 \pm 25$		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
---------------	--	------------	-----	-----	---	---

$1500 \pm 30$		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
---------------	--	-------	----	-----	---	---

<sup>1</sup> Using a parametrization for the  $K\pi$   $S$ -wave similar to ASTON 88 with fixed resonance width.

NODE=M094M;LINKAGE=A

<sup>2</sup> Using a  $K\pi$   $S$ -wave parametrization with resonant and non-resonant contributions.

NODE=M094M;LINKAGE=C

<sup>3</sup> From the pole position of the  $K\pi$  vector form factor in the complex  $s$ -plane and using EPIFANOV 07 data.

NODE=M094M;LINKAGE=BI

<sup>4</sup> Systematic uncertainties not estimated.

NODE=M094M;LINKAGE=NS

 **$K^*(1410)$  WIDTH**

NODE=M094W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M094W

**232 ± 21 OUR AVERAGE** Error includes scale factor of 1.1.

$176 \pm 52 \pm 22$		ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
---------------------	--	-------	----	------	---	------------------------------------

$240 \pm 18 \pm 12$		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
---------------------	--	-------	----	------	---	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$210 \pm 20 \pm 60$	190k	<sup>1</sup> AAIJ	16N	LHCB		$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
---------------------	------	-------------------	-----	------	--	---

OCCUR=2

$270 \pm 20 \pm 40$	190k	<sup>1</sup> AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
---------------------	------	-------------------	-----	------	--	---

$198^{+61}_{-87}$		<sup>2,3</sup> BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
-------------------	--	----------------------	----	------	--	---

$114 \pm 101$		BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
---------------	--	------	----	------	---	--

$275 \pm 65$		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
--------------	--	------------	-----	-----	---	---

$500 \pm 100$		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
---------------	--	-------	----	-----	---	---

<sup>1</sup> Using a  $K\pi$   $S$ -wave parametrization with resonant and non-resonant contributions.

NODE=M094W;LINKAGE=A

<sup>2</sup> From the pole position of the  $K\pi$  vector form factor in the complex  $s$ -plane and using EPIFANOV 07 data.

NODE=M094W;LINKAGE=BI

<sup>3</sup> Systematic uncertainties not estimated.

NODE=M094W;LINKAGE=NS

 **$K^*(1410)$  DECAY MODES**

NODE=M094215;NODE=M094

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $K^*(892)\pi$	$> 40$ %	95%
$\Gamma_2$ $K\pi$	$(6.6 \pm 1.3)$ %	
$\Gamma_3$ $K\rho$	$< 7$ %	95%
$\Gamma_4$ $\gamma K^0$	$< 2.3$ $\times 10^{-4}$	90%
$\Gamma_5$ $K\phi$	seen	

DESIG=2

DESIG=1

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

DESIG=5

**$K^*(1410)$  PARTIAL WIDTHS** $\Gamma(\gamma K^0)$  $\Gamma_4$ 

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<52.9	90	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$

NODE=M094217

NODE=M094W1  
NODE=M094W1 **$K^*(1410)$  BRANCHING RATIOS** $\Gamma(K\rho)/\Gamma(K^*(892)\pi)$  $\Gamma_3/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.17	95	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M094220

NODE=M094R1  
NODE=M094R1 $\Gamma(K\pi)/\Gamma(K^*(892)\pi)$  $\Gamma_2/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.16	95	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M094R2  
NODE=M094R2 $\Gamma(K\pi)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.066 ± 0.010 ± 0.008</b>	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M094R3  
NODE=M094R3 $\Gamma(K\phi)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	24k	<sup>1</sup> AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$

NODE=M094R00  
NODE=M094R00<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 7.7  $\sigma$ .

NODE=M094R00;LINKAGE=A

 **$K^*(1410)$  REFERENCES**

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira	
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
BOITO	09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin	
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)

NODE=M094

REFID=61150  
REFID=57836  
REFID=57273  
REFID=52728  
REFID=51929  
REFID=48822  
REFID=41002  
REFID=40262  
REFID=40234  
REFID=22689  
REFID=22551  
REFID=22545  
REFID=22443 $K_0^*(1430)$ 

$$I(J^P) = \frac{1}{2}(0^+)$$

NODE=M019

 **$K_0^*(1430)$  T-MATRIX POLE  $\sqrt{s}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

(1431 ± 6) - i (110 ± 19) <sup>1</sup> PELAEZ 17 RVUE  $\pi K \rightarrow \pi K$ <sup>1</sup> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M019PP

NODE=M019PP

NODE=M019PP;LINKAGE=A

 **$K_0^*(1430)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1425 ± 50</b>	<b>OUR ESTIMATE</b>			

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1449 ± 17 ± 2	<sup>1</sup> LEES	21A	BABR	$\eta_c(1S) \rightarrow \eta' K^+ K^-$
1438 ± 8 ± 4 5.4k	<sup>2</sup> LEES	14E	BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
1427 ± 4 ± 13	<sup>3</sup> BUGG	10	RVUE	S-matrix pole
1466.6 ± 0.7 ± 3.4 141k	<sup>4</sup> BONVICINI	08A	CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1412	<sup>5</sup> LINK	07	FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
1461.0 ± 4.0 ± 2.1 54k	<sup>6</sup> LINK	07B	FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
1406 ± 29	<sup>7</sup> BUGG	06	RVUE	
1435 ± 6	<sup>8</sup> ZHOU	06	RVUE	$K p \rightarrow K^- \pi^+ n$
1455 ± 20 ± 15	ABLIKIM	05Q	BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
1456 ± 8	<sup>9</sup> ZHENG	04	RVUE	$K^- p \rightarrow K^- \pi^+ n$

NODE=M019M

NODE=M019M  
→ UNCHECKED ←

~ 1419		10	BUGG	03	RVUE	11	$K^- p \rightarrow K^- \pi^+ n$
~ 1440		11	LI	03	RVUE	11	$K^- p \rightarrow K^- \pi^+ n$
1459 ± 9	15k	12	AITALA	02	E791	$D^+ \rightarrow K^- \pi^+ \pi^+$	
~ 1440		13	JAMIN	00	RVUE	$K p \rightarrow K p$	
1436 ± 8		14	BARBERIS	98E	OMEG	$450 p p \rightarrow$ $p_f p_s K^+ K^- \pi^+ \pi^-$	
1415 ± 25		10	ANISOVICH	97C	RVUE	11	$K^- p \rightarrow K^- \pi^+ n$
~ 1450		15	TORNQVIST	96	RVUE	$\pi \pi \rightarrow \pi \pi, K \bar{K}, K \pi$	
1412 ± 6		16	ASTON	88	LASS	11	$K^- p \rightarrow K^- \pi^+ n$
~ 1430			BAUBILLIER	84B	HBC	8.25	$K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 1425		17	ESTABROOKS	78	ASPK	13	$K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$
~ 1450.0			MARTIN	78	SPEC	10	$K^\pm p \rightarrow K_S^0 \pi p$

<sup>1</sup> Using a  $K\pi-K\eta'$  coupled channel Breit-Wigner function.

<sup>2</sup> Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$ . From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed width  $\Gamma(K_0^*(1430)) = 210$  MeV.

<sup>3</sup> S-matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an  $s$ -dependent width with couplings to  $K\pi$  and  $K\eta'$ , and the Adler zero near thresholds.

<sup>4</sup> From the isobar model with a complex pole for the  $\kappa$ .

<sup>5</sup> From a non-parametric analysis.

<sup>6</sup> A Breit-Wigner mass and width.

<sup>7</sup> S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the  $\kappa$  with an  $s$ -dependent width and an Adler zero near threshold.

<sup>8</sup> S-matrix pole. Using ASTON 88 and assuming  $K_0^*(700)$ ,  $K_0^*(1950)$ .

<sup>9</sup> Using ASTON 88 and assuming  $K_0^*(700)$ .

<sup>10</sup> T-matrix pole. Reanalysis of ASTON 88 data.

<sup>11</sup> Breit-Wigner fit. Using ASTON 88.

<sup>12</sup> Assuming a low-mass scalar  $K\pi$  resonance,  $\kappa(700)$ .

<sup>13</sup> T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

<sup>14</sup>  $J^P$  not determined, could be  $K_2^*(1430)$ .

<sup>15</sup> T-matrix pole.

<sup>16</sup> Uses a model for the background, without this background they get a mass 1340 MeV, where the phase shift passes  $90^\circ$ .

<sup>17</sup> Mass defined by pole position. From elastic  $K\pi$  partial-wave analysis.

NODE=M019M;LINKAGE=C  
NODE=M019M;LINKAGE=LE

NODE=M019M;LINKAGE=BG

NODE=M019M;LINKAGE=BO

NODE=M019M;LINKAGE=LI

NODE=M019M;LINKAGE=BW

NODE=M019M;LINKAGE=BU

NODE=M019M;LINKAGE=ZU

NODE=M019M;LINKAGE=ZH

NODE=M019M;LINKAGE=A1

NODE=M019M;LINKAGE=E

NODE=M019M;LINKAGE=AO

NODE=M019M;LINKAGE=JM

NODE=M019M;LINKAGE=JP

NODE=M019M;LINKAGE=TT

NODE=M019M;LINKAGE=D

NODE=M019M;LINKAGE=A

### $K_0^*(1430)$ WIDTH

NODE=M019W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>270 ± 80</b>				<b>OUR ESTIMATE</b>
• • •				We do not use the following data for averages, fits, limits, etc. • • •
210 ± 20 ± 12	5.4k	1 LEES	14E	BABR $\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
270 ± 10 ± 40		2 BUGG	10	RVUE S-matrix pole
174.2 ± 1.9 ± 3.2	141k	3 BONVICINI	08A	CLEO $D^+ \rightarrow K^- \pi^+ \pi^+$
~ 500		4 LINK	07	FOCS $D^+ \rightarrow K^- K^+ \pi^+$
177.0 ± 8.0 ± 3.4	54k	5 LINK	07B	FOCS $D^+ \rightarrow K^- \pi^+ \pi^+$
350 ± 40		6 BUGG	06	RVUE
288 ± 22		7 ZHOU	06	RVUE $K p \rightarrow K^- \pi^+ n$
270 ± 45 <sup>+30</sup> <sub>-35</sub>		ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
217 ± 31		8 ZHENG	04	RVUE $K^- p \rightarrow K^- \pi^+ n$
~ 316		9 BUGG	03	RVUE 11 $K^- p \rightarrow K^- \pi^+ n$
~ 350		10 LI	03	RVUE 11 $K^- p \rightarrow K^- \pi^+ n$
175 ± 17	15k	11 AITALA	02	E791 $D^+ \rightarrow K^- \pi^+ \pi^+$
~ 300		12 JAMIN	00	RVUE $K p \rightarrow K p$
196 ± 45		13 BARBERIS	98E	OMEG 450 $p p \rightarrow$ $p_f p_s K^+ K^- \pi^+ \pi^-$
330 ± 50		9 ANISOVICH	97C	RVUE 11 $K^- p \rightarrow K^- \pi^+ n$
~ 320		14 TORNQVIST	96	RVUE $\pi \pi \rightarrow \pi \pi, K \bar{K}, K \pi$
294 ± 23		ASTON	88	LASS 11 $K^- p \rightarrow K^- \pi^+ n$
~ 200		BAUBILLIER	84B	HBC 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
200 to 300		15 ESTABROOKS	78	ASPK 13 $K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$

NODE=M019W

→ UNCHECKED ←

- <sup>1</sup> Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$ . From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed mass  $M(K_0^*(1430)) = 1435$  MeV.
- <sup>2</sup> S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to  $K\pi$  and  $K\eta'$ , and the Adler zero near thresholds.
- <sup>3</sup> From the isobar model with a complex pole for the  $\kappa$ .
- <sup>4</sup> From a non-parametric analysis.
- <sup>5</sup> A Breit-Wigner mass and width.
- <sup>6</sup> S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the  $\kappa$  with an s-dependent width and an Adler zero near threshold.
- <sup>7</sup> S-matrix pole. Using ASTON 88 and assuming  $K_0^*(700)$ ,  $K_0^*(1950)$ .
- <sup>8</sup> Using ASTON 88 and assuming  $K_0^*(700)$ .
- <sup>9</sup> T-matrix pole. Reanalysis of ASTON 88 data.
- <sup>10</sup> Breit-Wigner fit. Using ASTON 88.
- <sup>11</sup> Assuming a low-mass scalar  $K\pi$  resonance,  $\kappa(700)$ .
- <sup>12</sup> T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.
- <sup>13</sup>  $J^P$  not determined, could be  $K_2^*(1430)$ .
- <sup>14</sup> T-matrix pole.
- <sup>15</sup> From elastic  $K\pi$  partial-wave analysis.

NODE=M019W;LINKAGE=LE

NODE=M019W;LINKAGE=BG

NODE=M019W;LINKAGE=BO

NODE=M019W;LINKAGE=LI

NODE=M019W;LINKAGE=BW

NODE=M019W;LINKAGE=BU

NODE=M019W;LINKAGE=ZU

NODE=M019W;LINKAGE=ZH

NODE=M019W;LINKAGE=A1

NODE=M019W;LINKAGE=E

NODE=M019W;LINKAGE=A0

NODE=M019W;LINKAGE=JM

NODE=M019W;LINKAGE=JP

NODE=M019W;LINKAGE=TT

NODE=M019W;LINKAGE=C

 **$K_0^*(1430)$  DECAY MODES**

NODE=M019215;NODE=M019

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\pi$	(93 $\pm$ 10 ) %
$\Gamma_2$ $K\eta$	( 8.6 $^{+2.7}_{-3.4}$ ) %
$\Gamma_3$ $K\eta'(958)$	seen

DESIG=1

DESIG=2

DESIG=3

 **$K_0^*(1430)$  BRANCHING RATIOS**

NODE=M019220

$\Gamma(K\pi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.93<math>\pm</math>0.04<math>\pm</math>0.09</b>	ASTON	88	LASS	0 11 $K^-p \rightarrow K^- \pi^+ n$

NODE=M019R1

NODE=M019R1

$\Gamma(K\eta)/\Gamma(K\pi)$	$\Gamma_2/\Gamma_1$			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.2<math>\pm</math>2.5<math>^{+1.0}_{-2.5}</math></b>	5.4k	<sup>1</sup> LEES	14E	BABR $\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$

NODE=M019R01

NODE=M019R01

- <sup>1</sup> Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$ . From a Dalitz analysis in the presence of several interfering scalar-meson resonances.

NODE=M019R01;LINKAGE=LE

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M019R00

NODE=M019R00

$\Gamma(K\eta'(958))/\Gamma(K\pi)$	$\Gamma_3/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.397<math>\pm</math>0.064<math>\pm</math>0.054</b>	<sup>1</sup> LEES	21A	BABR $\eta_c(1S) \rightarrow \eta' K^+ K^-$

NODE=M019R02

NODE=M019R02

- <sup>1</sup> Using  $K\pi$  data from LEES 14E.

NODE=M019R02;LINKAGE=A

 **$K_0^*(1430)$  REFERENCES**

NODE=M019

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira	
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LINK	07	PL B648 156	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng	
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>	
BUGG	03	PL B572 1	D.V. Bugg	
LI	03	PR D67 034025	L. Li, B. Zou, G. Li	
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
JAMIN	00	NP B587 331	M. Jamin <i>et al.</i>	
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev	
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)

REFID=61442

REFID=57836

REFID=55901

REFID=55937

REFID=53213

REFID=53056

REFID=52426

REFID=51702

REFID=51875

REFID=51037

REFID=51051

REFID=51458

REFID=50996

REFID=51198

REFID=50958

REFID=50165

REFID=49586

REFID=49192

REFID=48807

REFID=47983

REFID=46348

REFID=45815

REFID=44507

REFID=40262

REFID=22459

REFID=22443

REFID=22443

REFID=22446

$K_2^*(1430)$ 

$$I(J^P) = \frac{1}{2}(2^+)$$

We consider that phase-shift analyses provide more reliable determinations of the mass and width.

NODE=M022

NODE=M022

 **$K_2^*(1430)$  T-MATRIX POLE  $\sqrt{s}$** 

NODE=M022PP

NODE=M022PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$(1424 \pm 4) - i(66 \pm 2)$	<sup>1</sup> PELAEZ	17 RVUE	$\pi K \rightarrow \pi K$
<sup>1</sup> Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.			

NODE=M022PP;LINKAGE=A

 **$K_2^*(1430)$  MASS**

NODE=M022205

**CHARGED ONLY, WITH FINAL STATE  $K\pi$** NODE=M022M1  
NODE=M022M1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1427.3 ± 1.5 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.			
$1432.7 \pm 0.7^{+2.2}_{-2.3}$	183k	ABLIKIM	19AQ BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
$1420 \pm 4$	1587	BAUBILLIER	84B HBC	-	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
$1436 \pm 5.5$	400	<sup>1,2</sup> CLELAND	82 SPEC	+	$30 K^+ p \rightarrow K_S^0 \pi^+ p$
$1430 \pm 3.2$	1500	<sup>1,2</sup> CLELAND	82 SPEC	+	$50 K^+ p \rightarrow K_S^0 \pi^+ p$
$1430 \pm 3.2$	1200	<sup>1,2</sup> CLELAND	82 SPEC	-	$50 K^+ p \rightarrow K_S^0 \pi^- p$
$1423 \pm 5$	935	TOAFF	81 HBC	-	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
$1428.0 \pm 4.6$		<sup>3</sup> MARTIN	78 SPEC	+	$10 K^\pm p \rightarrow K_S^0 \pi p$
$1423.8 \pm 4.6$		<sup>3</sup> MARTIN	78 SPEC	-	$10 K^\pm p \rightarrow K_S^0 \pi p$
$1420.0 \pm 3.1$	1400	AGUILAR-...	71B HBC	-	$3.9, 4.6 K^- p$
$1425 \pm 8.0$	225	<sup>1,2</sup> BARNHAM	71C HBC	+	$K^+ p \rightarrow K^0 \pi^+ p$
$1416 \pm 10$	220	CRENNELL	69D DBC	-	$3.9 K^- N \rightarrow \bar{K}^0 \pi^- N$
$1414 \pm 13.0$	60	<sup>1</sup> LIND	69 HBC	+	$9 K^+ p \rightarrow K^0 \pi^+ p$
$1427 \pm 12$	63	<sup>1</sup> SCHWEING...	68 HBC	-	$5.5 K^- p \rightarrow \bar{K} \pi N$
$1423 \pm 11.0$	39	<sup>1</sup> BASSANO	67 HBC	-	$4.6-5.0 K^- p \rightarrow \bar{K}^0 \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$1428 \pm 2$	4300	<sup>4</sup> ABLIKIM	22L BES3		$2.0-3.08 e^+ e^- \rightarrow K^+ K^- \pi^0$
$1423.4 \pm 2 \pm 3$	$24809 \pm 820$	<sup>5</sup> BIRD	89 LASS	-	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

<sup>1</sup> Errors enlarged by us to  $\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.<sup>2</sup> Number of events in peak re-evaluated by us.<sup>3</sup> Systematic error added by us.<sup>4</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.<sup>5</sup> From a partial wave amplitude analysis.


NODE=M022M;LINKAGE=D

NODE=M022M;LINKAGE=W

NODE=M022M;LINKAGE=B

NODE=M022M1;LINKAGE=A

NODE=M022M;LINKAGE=F



**NEUTRAL ONLY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1432.4± 1.3 OUR AVERAGE</b>				
1431.2± 1.8± 0.7		<sup>1</sup> ASTON 88 LASS		11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 4 ± 6		<sup>1</sup> ASTON 87 LASS		11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1433 ± 6 ± 10		<sup>1</sup> ASTON 84B LASS		11 $K^- p \rightarrow \bar{K}^0 2\pi n$
1471 ± 12		<sup>1</sup> BAUBILLIER 82B HBC		8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
1428 ± 3		<sup>1</sup> ASTON 81C LASS		11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 2		<sup>1</sup> ESTABROOKS 78 ASPK		13 $K^\pm p \rightarrow pK\pi$
1440 ± 10		<sup>1</sup> BOWLER 77 DBC		5.5 $K^+ d \rightarrow K\pi pp$
1428.5± 3.9	1786± 127	<sup>2</sup> AUBERT 07AK BABR		10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
1420 ± 7	300	HENDRICK 76 DBC		8.25 $K^+ N \rightarrow K^+ \pi N$
1421.6± 4.2	800	MCCUBBIN 75 HBC		3.6 $K^- p \rightarrow K^- \pi^+ n$
1420.1± 4.3		<sup>3</sup> LINGLIN 73 HBC		2-13 $K^+ p \rightarrow K^+ \pi^- X$
1419.1± 3.7	1800	AGUILAR-... 71B HBC		3.9,4.6 $K^- p$
1416 ± 6	600	CORDS 71 DBC		9 $K^+ n \rightarrow K^+ \pi^- p$
1421.1± 2.6	2200	DAVIS 69 HBC		12 $K^+ p \rightarrow K^+ \pi^- X$

••• We do not use the following data for averages, fits, limits, etc. •••

- <sup>1</sup> From phase shift or partial-wave analysis.
- <sup>2</sup> Systematic errors not estimated.
- <sup>3</sup> From pole extrapolation, using world  $K^+ p$  data summary tape.

NODE=M022M4  
NODE=M022M4

NODE=M022M;LINKAGE=P  
NODE=M022M4;LINKAGE=NS  
NODE=M022M;LINKAGE=C

**$K_2^*(1430)$  WIDTH**

NODE=M022210

**CHARGED ONLY, WITH FINAL STATE  $K\pi$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>100.0± 2.1 OUR FIT</b>					
<b>100.0± 2.2 OUR AVERAGE</b>					Error includes scale factor of 1.1.
102.5± 1.6 <sup>+3.1</sup> <sub>-2.8</sub>	183k	ABLIKIM 19AQ BES		±	$J/\psi \rightarrow K^+ K^- \pi^0$
109 ± 22	400	<sup>1,2</sup> CLELAND 82 SPEC		+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
124 ± 12.8	1500	<sup>1,2</sup> CLELAND 82 SPEC		+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
113 ± 12.8	1200	<sup>1,2</sup> CLELAND 82 SPEC		-	50 $K^+ p \rightarrow K_S^0 \pi^- p$
85 ± 16	935	TOAFF 81 HBC		-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
96.5± 3.8		MARTIN 78 SPEC		+	10 $K^\pm p \rightarrow K_S^0 \pi p$
97.7± 4.0		MARTIN 78 SPEC		-	10 $K^\pm p \rightarrow K_S^0 \pi p$
94.7 <sup>+15.1</sup> <sub>-12.5</sub>	1400	AGUILAR-... 71B HBC		-	3.9,4.6 $K^- p$

NODE=M022W1  
NODE=M022W1

OCCUR=2

OCCUR=3

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

107 ± 4	4300	<sup>3</sup> ABLIKIM 22L BES3			2.0-3.08 $e^+ e^- \rightarrow K^+ K^- \pi^0$
98 ± 4 ± 4	25k	<sup>4</sup> BIRD 89 LASS		-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$



- <sup>1</sup> Errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.
- <sup>2</sup> Number of events in peak re-evaluated by us.
- <sup>3</sup> From a partial wave amplitude analysis at  $\sqrt{s} = 2.125$  GeV which includes all the possible intermediate states that match  $J^{PC}$  conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.
- <sup>4</sup> From a partial wave amplitude analysis.

NODE=M022W;LINKAGE=D  
 NODE=M022W;LINKAGE=W  
 NODE=M022W1;LINKAGE=A

NODE=M022W;LINKAGE=F


NODE=M022W4  
 NODE=M022W4

**NEUTRAL ONLY**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>109 ± 5 OUR AVERAGE</b>				Error includes scale factor of 1.9. See the ideogram below.
116.5 ± 3.6 ± 1.7		<sup>1</sup> ASTON 88	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
129 ± 15 ± 15		<sup>1</sup> ASTON 87	LASS	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
131 ± 24 ± 20		<sup>1</sup> ASTON 84B	LASS	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
143 ± 34		<sup>1</sup> BAUBILLIER 82B	HBC	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
98 ± 8		<sup>1</sup> ASTON 81C	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
140 ± 30		<sup>1</sup> ETKIN 80	SPEC	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
98 ± 5		<sup>1</sup> ESTABROOKS 78	ASPK	13 $K^\pm p \rightarrow p K \pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
113.7 ± 9.2	1786 ± 127	<sup>2</sup> AUBERT 07AK	BABR	10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
125 ± 29	300	<sup>3</sup> HENDRICK 76	DBC	8.25 $K^+ N \rightarrow K^+ \pi N$
116 ± 18	800	MCCUBBIN 75	HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
61 ± 14		<sup>4</sup> LINGLIN 73	HBC	2-13 $K^+ p \rightarrow K^+ \pi^- X$
116.6 <sup>+10.3</sup> <sub>-15.5</sub>	1800	AGUILAR-...	71B	HBC 3.9,4.6 $K^- p$
144 ± 24.0	600	<sup>3</sup> CORDS 71	DBC	9 $K^+ n \rightarrow K^+ \pi^- p$
101 ± 10	2200	DAVIS 69	HBC	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$

NODE=M022W;LINKAGE=P  
 NODE=M022W4;LINKAGE=NS  
 NODE=M022W4;LINKAGE=D  
 NODE=M022W;LINKAGE=C

- <sup>1</sup> From phase shift or partial-wave analysis.
- <sup>2</sup> Systematic errors not estimated.
- <sup>3</sup> Errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.
- <sup>4</sup> From pole extrapolation, using world  $K^+ p$  data summary tape.



**$K_2^*(1430)$  DECAY MODES**

NODE=M022215;NODE=M022

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $K \pi$	(49.9 ± 1.2) %	
$\Gamma_2$ $K^*(892) \pi$	(24.7 ± 1.5) %	
$\Gamma_3$ $K^*(892) \pi \pi$	(13.4 ± 2.2) %	
$\Gamma_4$ $K \rho$	( 8.7 ± 0.8) %	S=1.2

DESIG=1  
 DESIG=2  
 DESIG=6  
 DESIG=3

$\Gamma_5$	$K\omega$	$(2.9 \pm 0.8) \%$		DESIG=4
$\Gamma_6$	$K^+\gamma$	$(2.4 \pm 0.5) \times 10^{-3}$	S=1.1	DESIG=8
$\Gamma_7$	$K\eta$	$(1.5^{+3.4}_{-1.0}) \times 10^{-3}$	S=1.3	DESIG=5
$\Gamma_8$	$K\omega\pi$	$< 7.2 \times 10^{-4}$	CL=95%	DESIG=7
$\Gamma_9$	$K^0\gamma$	$< 9 \times 10^{-4}$	CL=90%	DESIG=10;OUR EVAL;→ UNCHECKED ←

### CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 10 branching ratios uses 32 measurements and one constraint to determine 8 parameters. The overall fit has a  $\chi^2 = 21.1$  for 25 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-9						
$x_3$	-40	-73					
$x_4$	-8	36	-52				
$x_5$	-11	-3	-26	-7			
$x_6$	-1	-1	-1	-1	0		
$x_7$	-4	-7	-5	-5	-2	0	
$\Gamma$	0	0	0	0	0	-10	0
	$x_1$	$x_2$	$x_3$	$x_4$	$x_5$	$x_6$	$x_7$

	Mode	Rate (MeV)	Scale factor	
$\Gamma_1$	$K\pi$	$49.9 \pm 1.6$		DESIG=1
$\Gamma_2$	$K^*(892)\pi$	$24.7 \pm 1.6$		DESIG=2
$\Gamma_3$	$K^*(892)\pi\pi$	$13.5 \pm 2.3$		DESIG=6
$\Gamma_4$	$K\rho$	$8.7 \pm 0.8$	1.2	DESIG=3
$\Gamma_5$	$K\omega$	$2.9 \pm 0.8$		DESIG=4
$\Gamma_6$	$K^+\gamma$	$0.24 \pm 0.05$	1.1	DESIG=8
$\Gamma_7$	$K\eta$	$0.15^{+0.34}_{-0.10}$	1.3	DESIG=5

### $K_2^*(1430)$ PARTIAL WIDTHS

NODE=M022220

#### $\Gamma(K^+\gamma)$

 $\Gamma_6$ 

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>241 \pm 50</math> OUR FIT</b>				Error includes scale factor of 1.1.
<b><math>240 \pm 45</math></b>	CIHANGIR	82	SPEC +	$200 K^+ Z \rightarrow Z K^+ \pi^0,$ $Z K_S^0 \pi^+$

NODE=M022W8  
NODE=M022W8

#### $\Gamma(K^0\gamma)$

 $\Gamma_9$ 

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>&lt; 5.4</math></b>	90	ALAVI-HARATI02B	KTEV		$K + A \rightarrow K^* + A$
••• We do not use the following data for averages, fits, limits, etc. •••					
$< 84$	90	CARLSMITH	87	SPEC 0	$60-200 K_L^0 A \rightarrow$ $K_S^0 \pi^0 A$

NODE=M022W9  
NODE=M022W9

### $K_2^*(1430)$ BRANCHING RATIOS

NODE=M022225

#### $\Gamma(K\pi) / \Gamma_{\text{total}}$

 $\Gamma_1 / \Gamma$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>0.499 \pm 0.012</math> OUR FIT</b>				
<b><math>0.488 \pm 0.014</math> OUR AVERAGE</b>				
$0.485 \pm 0.006 \pm 0.020$	<sup>1</sup> ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
$0.49 \pm 0.02$	<sup>1</sup> ESTABROOKS	78	ASPK ±	$13 K^\pm p \rightarrow p K \pi$

NODE=M022R1  
NODE=M022R1<sup>1</sup> From phase shift analysis.

NODE=M022R;LINKAGE=P

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$

$\Gamma_2/\Gamma_1$

NODE=M022R4  
NODE=M022R4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.496±0.034 OUR FIT</b>				
<b>0.47 ±0.04 OUR AVERAGE</b>				
0.44 ±0.09	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
0.62 ±0.19	LAUSCHER	75	HBC	0 10,16 $K^- p \rightarrow K^- \pi^+ n$
0.54 ±0.16	DEHM	74	DBC	0 4.6 $K^+ N$
0.47 ±0.08	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
0.47 ±0.10	BASSANO	67	HBC	-0 4.6,5.0 $K^- p$
0.45 ±0.13	BADIER	65C	HBC	- 3 $K^- p$

$\Gamma(K\omega)/\Gamma(K\pi)$

$\Gamma_5/\Gamma_1$

NODE=M022R5  
NODE=M022R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.059±0.017 OUR FIT</b>				
<b>0.070±0.035 OUR AVERAGE</b>				
0.05 ±0.04	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
0.13 ±0.07	BASSOMPIE...	69	HBC	0 5 $K^+ p$

OCCUR=2

$\Gamma(K\rho)/\Gamma(K\pi)$

$\Gamma_4/\Gamma_1$

NODE=M022R6  
NODE=M022R6

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.174±0.017 OUR FIT</b> Error includes scale factor of 1.2.				
<b>0.150<sup>+0.029</sup><sub>-0.017</sub> OUR AVERAGE</b>				
0.18 ±0.05	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
0.02 <sup>+0.10</sup> <sub>-0.02</sub>	DEHM	74	DBC	0 4.6 $K^+ N$
0.16 ±0.05	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
0.14 ±0.10	BASSANO	67	HBC	-0 4.6,5.0 $K^- p$
0.14 ±0.07	BADIER	65C	HBC	- 3 $K^- p$


$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$

$\Gamma_4/\Gamma_2$

NODE=M022R7  
NODE=M022R7

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.350±0.031 OUR FIT</b> Error includes scale factor of 1.4.				
<b>0.354±0.033 OUR AVERAGE</b> Error includes scale factor of 1.4. See the ideogram below.				
0.293±0.032±0.020	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
0.38 ±0.09	BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
0.39 ±0.03	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$

WEIGHTED AVERAGE  
0.354±0.033 (Error scaled by 1.4)



$\Gamma(K\omega)/\Gamma(K^*(892)\pi)$

$\Gamma_5/\Gamma_2$

NODE=M022R8  
NODE=M022R8

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.118±0.034 OUR FIT</b>				
<b>0.10 ±0.04</b>	FIELD	67	HBC	- 3.8 $K^- p$

$\Gamma(K\eta)/\Gamma(K^*(892)\pi)$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_7/\Gamma_2$
-------	-------------	------	-----	---------	---------------------

**0.006<sup>+0.014</sup><sub>-0.004</sub> OUR FIT** Error includes scale factor of 1.2.

**0.07 ± 0.04** FIELD 67 HBC - 3.8  $K^- p$

NODE=M022R9  
NODE=M022R9

 $\Gamma(K\eta)/\Gamma(K\pi)$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_7/\Gamma_1$
-------	-----	-------------	------	-----	---------	---------------------

**0.0030<sup>+0.0070</sup><sub>-0.0020</sub> OUR FIT** Error includes scale factor of 1.3.

**0 ± 0.0056** <sup>1</sup> ASTON 88B LASS - 11  $K^- p \rightarrow K^- \eta p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04 95 AGUILAR-... 71B HBC 3.9,4.6  $K^- p$

<0.065 <sup>2</sup> BASSOMPIE... 69 HBC 5.0  $K^+ p$

<0.02 BISHOP 69 HBC 3.5  $K^+ p$

<sup>1</sup> ASTON 88B quote < 0.0092 at CL=95%. We convert this to a central value and 1 sigma error in order to be able to use it in our constrained fit.

<sup>2</sup> Restated by us.

NODE=M022R10  
NODE=M022R10

NODE=M022R;LINKAGE=PQ

NODE=M022R;LINKAGE=R

 $\Gamma(K^*(892)\pi\pi)/\Gamma_{total}$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_3/\Gamma$
-------	-------------	------	-----	---------	-------------------

**0.134 ± 0.022 OUR FIT**

**0.12 ± 0.04** <sup>1</sup> GOLDBERG 76 HBC - 3  $K^- p \rightarrow p \bar{K}^0 \pi \pi \pi$

<sup>1</sup> Assuming  $\pi\pi$  system has isospin 1, which is supported by the data.

NODE=M022R11  
NODE=M022R11

NODE=M022R;LINKAGE=T

 $\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_3/\Gamma_1$
-------	-------------	------	-----	---------	---------------------

**0.27 ± 0.05 OUR FIT**

**0.21 ± 0.08** <sup>1,2</sup> JONGEJANS 78 HBC - 4  $K^- p \rightarrow p \bar{K}^0 \pi \pi \pi$

<sup>1</sup> Restated by us.

<sup>2</sup> Assuming  $\pi\pi$  system has isospin 1, which is supported by the data.

NODE=M022R12  
NODE=M022R12

NODE=M022R12;LINKAGE=R  
NODE=M022R12;LINKAGE=T

 $\Gamma(K\omega\pi)/\Gamma_{total}$ 

VALUE (units 10 <sup>-3</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma$
---------------------------------	-----	------	-------------	------	---------	-------------------

**<0.72** 95 0 JONGEJANS 78 HBC 4  $K^- p \rightarrow p \bar{K}^0 4\pi$

NODE=M022R13  
NODE=M022R13

 $K_2^*(1430)$  REFERENCES

ABLIKIM 22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61649
ABLIKIM 19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
PELAEZ 17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira		REFID=57836
AUBERT 07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
ALAVI-HARATI 02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)	REFID=48822
BIRD 89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON 88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON 88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40281
ASTON 87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
CARLSMITH 87	PR D36 3502	D. Carlsmith <i>et al.</i>	(EFI, SACL)	REFID=40557
ASTON 84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)	REFID=22763
BAUBILLIER 84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
BAUBILLIER 82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
CIHANGIR 82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND 82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON 81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22821
DAUM 81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
TOAFF 81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
ETKIN 80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
ESTABROOKS 78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
Also	PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
JONGEJANS 78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
MARTIN 78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=22446
BOWLER 77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
GOLDBERG 76	LNC 17 253	J. Goldberg	(HAIF)	REFID=22742
HENDRICK 76	NP B112 189	K. Hendrickx <i>et al.</i>	(MONS, SACL, PARIS+)	REFID=22743
LAUSCHER 75	NP B86 189	P. Lauscher <i>et al.</i>	(ABCLV Collab.) JP	REFID=22582
MCCUBBIN 75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
DEHM 74	NP B75 47	G. Dehm <i>et al.</i>	(MPIM, BRUX, MONS, CERN)	REFID=22736
LINGLIN 73	NP B55 408	D. Linglin	(CERN)	REFID=22428
AGUILAR-... 71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
BARNHAM 71C	NP B28 171	K.W.J. Barnham <i>et al.</i>	(BIRM, GLAS)	REFID=22409
CORDS 71	PR D4 1974	D. Cords <i>et al.</i>	(PURD, UCD, IUPU)	REFID=22411
BASSOMPIE... 69	NP B13 189	G. Bassompierre <i>et al.</i>	(CERN, BRUX) JP	REFID=22710
BISHOP 69	NP B9 403	J.M. Bishop <i>et al.</i>	(WISC)	REFID=22485
CRENNELL 69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS 69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
LIND 69	NP B14 1	V.G. Lind <i>et al.</i>	(LRL) JP	REFID=22404
SCHWEING... 68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
Also	Thesis	F.L. Schweingruber	(NWES, NWES)	REFID=22709
BASSANO 67	PRL 19 968	D. Bassano <i>et al.</i>	(BNL, SYRA)	REFID=22695
FIELD 67	PL 24B 638	J.H. Field <i>et al.</i>	(UCSD)	REFID=22701
BADIER 65C	PL 19 612	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=22690

NODE=M022

**K(1460)**

$$I(J^P) = \frac{1}{2}(0^-)$$

Observed in  $K\pi\pi$  partial-wave analysis.

NODE=M021

NODE=M021

NODE=M021M

NODE=M021M

NODE=M021M;LINKAGE=A

NODE=M021W

NODE=M021W

NODE=M021W;LINKAGE=A

NODE=M021215;NODE=M021

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

NODE=M021220

NODE=M021W1  
NODE=M021W1NODE=M021W2  
NODE=M021W2NODE=M021W3  
NODE=M021W3NODE=M021R00  
NODE=M021R00

NODE=M021R00;LINKAGE=A

NODE=M021

REFID=61150  
REFID=59187  
REFID=22548  
REFID=22767**K(1460) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1482.40 ± 3.58 ± 15.22	894k	AAIJ	18AI	LHCB	$D^0 \rightarrow K^\mp 2\pi^\pm \pi^\mp$
~ 1460	63	DAUM	81C	CNTR	$K^- p \rightarrow K^- 2\pi p$
~ 1400	13	<sup>1</sup> BRANDENB...	76B	ASPK	$K^\pm p \rightarrow K^+ 2\pi p$

<sup>1</sup> Coupled mainly to  $K f_0(1370)$ . Decay into  $K^*(892)\pi$  seen.

**K(1460) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
335.60 ± 6.20 ± 8.65	894k	AAIJ	18AI	LHCB	$D^0 \rightarrow K^\mp 2\pi^\pm \pi^\mp$
~ 260	63	DAUM	81C	CNTR	$K^- p \rightarrow K^- 2\pi p$
~ 250	15	<sup>1</sup> BRANDENB...	76B	ASPK	$K^\pm p \rightarrow K^+ 2\pi p$

<sup>1</sup> Coupled mainly to  $K f_0(1370)$ . Decay into  $K^*(892)\pi$  seen.

**K(1460) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\pi$	seen
$\Gamma_2$ $K\rho$	seen
$\Gamma_3$ $K_0^*(1430)\pi$	seen
$\Gamma_4$ $K\phi$	seen

**K(1460) PARTIAL WIDTHS**

$\Gamma(K^*(892)\pi)$					$\Gamma_1$
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT		
~ 109	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$	
$\Gamma(K\rho)$					$\Gamma_2$
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT		
~ 34	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$	
$\Gamma(K_0^*(1430)\pi)$					$\Gamma_3$
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT		
~ 117	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$	
$\Gamma(K\phi)/\Gamma_{total}$					$\Gamma_4/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	24k	<sup>1</sup> AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi\phi K^+$

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 12  $\sigma$ .

**K(1460) REFERENCES**

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AI	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BRANDENB...	76B	PRL 36 1239	G.W. Brandenburg <i>et al.</i>	(SLAC) JP

**$K_2(1580)$** 

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the  $K^- \pi^+ \pi^-$  system. Needs confirmation.

NODE=M039

NODE=M039

 **$K_2(1580)$  MASS**

NODE=M039M

VALUE (MeV)	DOCUMENT ID	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1580	OTTER	79	- 10,14,16 $K^- p$

NODE=M039M

 **$K_2(1580)$  WIDTH**

NODE=M039W

VALUE (MeV)	DOCUMENT ID	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 110	OTTER	79	- 10,14,16 $K^- p$

NODE=M039W

 **$K_2(1580)$  DECAY MODES**

NODE=M039215;NODE=M039

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K^*(892)\pi$	seen
$\Gamma_2$ $K_2^*(1430)\pi$	possibly seen

DESIG=1

DESIG=2

 **$K_2(1580)$  BRANCHING RATIOS**

NODE=M039220

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
seen	OTTER	79	HBC	- 10,14,16 $K^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	

NODE=M039R1  
NODE=M039R1

$\Gamma(K_2^*(1430)\pi)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
possibly seen	OTTER	79	HBC	- 10,14,16 $K^- p$	

NODE=M039R2  
NODE=M039R2 **$K_2(1580)$  REFERENCES**

NODE=M039

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
OTTER	79	NP B147 1	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP

REFID=53668  
REFID=22772

**K(1630)**

$$I(J^P) = \frac{1}{2}(?^?)$$

OMITTED FROM SUMMARY TABLE

Seen as a narrow peak, compatible with the experimental resolution, in the invariant mass of the  $K_S^0 \pi^+ \pi^-$  system produced in  $\pi^- p$  interactions at high momentum transfers.

NODE=M160

NODE=M160

**K(1630) MASS**

NODE=M160M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1629±7</b>	~ 75	KARNAUKHOV98	BC	16.0 $\pi^- p \rightarrow (K_S^0 \pi^+ \pi^-)$ $X^+ \pi^- X^0$

NODE=M160M

**K(1630) WIDTH**

NODE=M160W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16<sup>+19</sup><sub>-16</sub></b>	~ 75	<sup>1</sup> KARNAUKHOV98	BC	16.0 $\pi^- p \rightarrow (K_S^0 \pi^+ \pi^-)$ $X^+ \pi^- X^0$

NODE=M160W

<sup>1</sup> Compatible with an experimental resolution of  $14 \pm 1$  MeV.

NODE=M160W;LINKAGE=A

**K(1630) DECAY MODES**

NODE=M160215;NODE=M160

Mode

 $\Gamma_1 \quad K_S^0 \pi^+ \pi^-$ 

DESIG=1

**K(1630) REFERENCES**

KARNAUKHOV 98 PAN 61 203 V.M. Karnaukhov, C. Coca, V.I. Moroz  
Translated from YAF 61 252.

NODE=M160

REFID=46371

NODE=M099

**K<sub>1</sub>(1650)**

$$I(J^P) = \frac{1}{2}(1^+)$$

This entry contains various peaks in strange meson systems ( $K^+ \phi$ ,  $K \pi \pi$ ) reported in partial-wave analysis in the 1600–1900 mass region.

NODE=M099

**K<sub>1</sub>(1650) MASS**

NODE=M099M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1650±50</b>		FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1861±10 <sup>+</sup> <sub>-46</sub>	24k	<sup>1</sup> AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1911±37 <sup>+</sup> <sub>-48</sub>	24k	<sup>1</sup> AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1793±59 <sup>+</sup> <sub>-101</sub>	4289	<sup>2,3</sup> AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
~ 1840		ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 1800		DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

NODE=M099M

OCCUR=2

<sup>1</sup> One of two  $K_1$  states reported by AAIJ 21E. From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $4.5 \sigma$ .

NODE=M099M;LINKAGE=C

<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $7.6 \sigma$ .

NODE=M099M;LINKAGE=A

<sup>3</sup> Superseded by AAIJ 21E.

NODE=M099M;LINKAGE=B

**K<sub>1</sub>(1650) WIDTH**

NODE=M099W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>150± 50</b>		FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$

NODE=M099W

• • • We do not use the following data for averages, fits, limits, etc. • • •

$149 \pm 41^{+231}_{-23}$	24k	<sup>1</sup> AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$276 \pm 50^{+319}_{-159}$	24k	<sup>1</sup> AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$365 \pm 157^{+138}_{-215}$	4289	<sup>2,3</sup> AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$\sim 250$		DAUM	81C	CNTR	$- 63 K^- p \rightarrow K^- 2\pi p$

OCCUR=2

<sup>1</sup>One of two  $K_1$  states reported by AAIJ 21E. From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $4.5 \sigma$ .

<sup>2</sup>From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $7.6 \sigma$ .

<sup>3</sup>Superseded by AAIJ 21E.

NODE=M099W;LINKAGE=C

NODE=M099W;LINKAGE=A  
NODE=M099W;LINKAGE=B

## $K_1(1650)$ DECAY MODES

NODE=M099215;NODE=M099

Mode	
$\Gamma_1$	$K \pi \pi$
$\Gamma_2$	$K \phi$

DESIG=1

DESIG=2

## $K_1(1650)$ REFERENCES

NODE=M099

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

REFID=61150  
REFID=57657  
REFID=57636  
REFID=20569  
REFID=22801  
REFID=22548

NODE=M095

$K^*(1680)$

$$I(J^P) = \frac{1}{2}(1^-)$$

## $K^*(1680)$ MASS

NODE=M095M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>1718 \pm 18</math> OUR AVERAGE</b>					
$1722 \pm 20^{+33}_{-109}$	4289	<sup>1</sup> AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$1677 \pm 10 \pm 32$		ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
$1735 \pm 10 \pm 20$		ASTON	87	LASS 0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M095M

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1678 \pm 64$		BIRD	89	LASS -	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
$1800 \pm 70$		ETKIN	80	MPS 0	$6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
$\sim 1650$		ESTABROOKS	78	ASPK 0	$13 K^\pm p \rightarrow K^\pm \pi^\pm n$

<sup>1</sup>From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $8.5 \sigma$ .

NODE=M095M;LINKAGE=A

## $K^*(1680)$ WIDTH

NODE=M095W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>322 \pm 110</math> OUR AVERAGE</b>					Error includes scale factor of 4.2.
$354 \pm 75^{+140}_{-181}$	4289	<sup>2</sup> AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$205 \pm 16 \pm 34$		ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
$423 \pm 18 \pm 30$		ASTON	87	LASS 0	$11 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M095W

• • • We do not use the following data for averages, fits, limits, etc. • • •

$454 \pm 270$		BIRD	89	LASS -	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$
$170 \pm 30$		ETKIN	80	MPS 0	$6 K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
250 to 300		ESTABROOKS	78	ASPK 0	$13 K^\pm p \rightarrow K^\pm \pi^\pm n$

<sup>2</sup>From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $8.5 \sigma$ .

NODE=M095W;LINKAGE=A



**K\*(1680) DECAY MODES**

NODE=M095215;NODE=M095

Mode	Fraction ( $\Gamma_i/\Gamma$ )	
$\Gamma_1$ $K\pi$	(38.7±2.5) %	DESIG=1
$\Gamma_2$ $K\rho$	(31.4 <sup>+5.0</sup> <sub>-2.1</sub> ) %	DESIG=3
$\Gamma_3$ $K^*(892)\pi$	(29.9 <sup>+2.2</sup> <sub>-5.0</sub> ) %	DESIG=2
$\Gamma_4$ $K\phi$	seen	DESIG=4
$\Gamma_5$ $K\eta$	( 1.4 <sup>+1.0</sup> <sub>-0.8</sub> ) %	DESIG=6

**CONSTRAINED FIT INFORMATION**

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 2.9$  for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-36	
$x_3$	-39	-72
	$x_1$	$x_2$

**K\*(1680) BRANCHING RATIOS**

NODE=M095220

$\Gamma(K\pi)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.387±0.026 OUR FIT</b>					NODE=M095R4
<b>0.388±0.014±0.022</b>	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$	NODE=M095R4

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$					$\Gamma_1/\Gamma_3$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>1.30<sup>+0.23</sup><sub>-0.14</sub> OUR FIT</b>					NODE=M095R2
<b>2.8 ±1.1</b>	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$	NODE=M095R2

$\Gamma(K\rho)/\Gamma(K\pi)$					$\Gamma_2/\Gamma_1$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.81<sup>+0.14</sup><sub>-0.09</sub> OUR FIT</b>					NODE=M095R3
<b>1.2 ±0.4</b>	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$	NODE=M095R3

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$					$\Gamma_2/\Gamma_3$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
<b>1.05<sup>+0.27</sup><sub>-0.11</sub> OUR FIT</b>					NODE=M095R1
<b>0.97±0.09<sup>+0.30</sup><sub>-0.10</sub></b>	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	NODE=M095R1

$\Gamma(K\phi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>seen</b>	24k	<sup>3</sup> AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$	NODE=M095R00
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	4289	<sup>4,5</sup> AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$	NODE=M095R00

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 4.7  $\sigma$ .

<sup>4</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 8.5  $\sigma$ .

<sup>5</sup> Superseded by AAIJ 21E.

NODE=M095R00;LINKAGE=B  
 NODE=M095R00;LINKAGE=A  
 NODE=M095R00;LINKAGE=C

$\Gamma(K\eta)/\Gamma(K\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma_1$
$0.037 \pm 0.007^{+0.024}_{-0.018}$	116k	<sup>6</sup> CHEN	20A BELL	$D^0 \rightarrow K^- \pi^+ \eta$	NODE=M095R02 NODE=M095R02

<sup>6</sup>CHEN 20A quotes the ratio  $\Gamma(K^*(1680)^- \rightarrow K^- \eta)/\Gamma(K^*(1680)^- \rightarrow K^- \pi^0) = 0.11 \pm 0.02^{+0.06}_{-0.04} \pm 0.04(\text{BPDG})$  where the last uncertainty comes from  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.20)\%$ . We divide it by 3 taking into account that  $\Gamma(K^*(1680)^- \rightarrow K^- \pi^0)/\Gamma(K^*(1680)^- \rightarrow (K\pi)^-) = 1/3$ .

NODE=M095R02;LINKAGE=A

$\Gamma(K\eta)/\Gamma_{\text{total}}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma$
$1.44 \pm 0.21^{+0.96}_{-0.73}$	116k	<sup>7</sup> CHEN	20A BELL	$D^0 \rightarrow K^- \pi^+ \eta$	NODE=M095R01 NODE=M095R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>7</sup>From an amplitude analysis of the decay  $D^0 \rightarrow K^- \pi^+ \eta$  with a significance of  $16 \sigma$ . Not independent of the CHEN 20A measurement of  $\Gamma(K^*(1680) \rightarrow K\eta)/\Gamma(K^*(1680) \rightarrow K\pi)$ .

NODE=M095R01;LINKAGE=A

**$K^*(1680)$  REFERENCES**

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)	REFID=60333
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22689
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP	REFID=22443

NODE=M095

**$K_2(1770)$**

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this Review, PDG 04.

NODE=M023

NODE=M023

**$K_2(1770)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>1773 \pm 8</math> OUR AVERAGE</b>					
$1777 \pm 35^{+122}_{-77}$	4289	<sup>1</sup> AAIJ	17C LHCb		$B^+ \rightarrow J/\psi \phi K^+$
$1773 \pm 8$		<sup>2</sup> ASTON	93 LASS		$11K^- p \rightarrow K^- \omega p$
$1743 \pm 15$		TIKHOMIROV	03 SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
$1810 \pm 20$		FRAME	86 OMEG +		$13 K^+ p \rightarrow \phi K^+ p$
$\sim 1730$		ARMSTRONG	83 OMEG -		$18.5 K^- p \rightarrow 3K p$
$\sim 1780$		<sup>3</sup> DAUM	81C CNTR -		$63 K^- p \rightarrow K^- 2\pi p$
$1710 \pm 15$	60	CHUNG	74 HBC -		$7.3 K^- p \rightarrow K^- \omega p$
$1767 \pm 6$		BLIEDEN	72 MMS -		$11-16 K^- p$
$1730 \pm 20$	306	<sup>4</sup> FIRESTONE	72B DBC +		$12 K^+ d$
$1765 \pm 40$		<sup>5</sup> COLLEY	71 HBC +		$10 K^+ p \rightarrow K 2\pi N$
1740		DENEGRI	71 DBC -		$12.6 K^- d \rightarrow K 2\pi d$
$1745 \pm 20$		AGUILAR-...	70C HBC -		$4.6 K^- p$
$1780 \pm 15$		BARTSCH	70C HBC -		$10.1 K^- p$
$1760 \pm 15$		LUDLAM	70 HBC -		$12.6 K^- p$

NODE=M023M

NODE=M023M

- <sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.0 \sigma$ .
- <sup>2</sup> From a partial wave analysis of the  $K^- \omega$  system.
- <sup>3</sup> From a partial wave analysis of the  $K^- 2\pi$  system.
- <sup>4</sup> Produced in conjunction with excited deuteron.
- <sup>5</sup> Systematic errors added correspond to spread of different fits.

NODE=M023M;LINKAGE=C  
NODE=M023M;LINKAGE=A  
NODE=M023M;LINKAGE=B  
NODE=M023M;LINKAGE=P  
NODE=M023M;LINKAGE=X

**$K_2(1770)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b><math>186 \pm 14</math> OUR AVERAGE</b>					
$217 \pm 116^{+221}_{-154}$	4289	<sup>6</sup> AAIJ	17C LHCb		$B^+ \rightarrow J/\psi \phi K^+$
$186 \pm 14$		<sup>7</sup> ASTON	93 LASS		$11K^- p \rightarrow K^- \omega p$

NODE=M023W

NODE=M023W

• • • We do not use the following data for averages, fits, limits, etc. • • •

147 ± 70		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow$ $K_S^0 K_S^0 K_L^0 X$
140 ± 40		FRAME 86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
~ 220		ARMSTRONG 83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 210		<sup>8</sup> DAUM 81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$
110 ± 50	60	CHUNG 74	HBC -	7.3 $K^- p \rightarrow K^- \omega p$
100 ± 26		BLIEDEN 72	MMS -	11-16 $K^- p$
210 ± 30	306	<sup>9</sup> FIRESTONE 72B	DBC +	12 $K^+ d$
90 ± 70		<sup>10</sup> COLLEY 71	HBC +	10 $K^+ p \rightarrow K 2\pi N$
130		DENEGRI 71	DBC -	12.6 $K^- d \rightarrow \bar{K} 2\pi d$
100 ± 50		AGUILAR-...	70C HBC -	4.6 $K^- p$
138 ± 40		BARTSCH 70C	HBC -	10.1 $K^- p$
50 <sup>+</sup> <sub>-</sub> 40 20		LUDLAM 70	HBC -	12.6 $K^- p$

<sup>6</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.0 \sigma$ .

<sup>7</sup> From a partial wave analysis of the  $K^- \omega$  system.

<sup>8</sup> From a partial wave analysis of the  $K^- 2\pi$  system.

<sup>9</sup> Produced in conjunction with excited deuteron.

<sup>10</sup> Systematic errors added correspond to spread of different fits.

NODE=M023W;LINKAGE=A  
 NODE=M023W;LINKAGE=B  
 NODE=M023W;LINKAGE=C  
 NODE=M023W;LINKAGE=P  
 NODE=M023W;LINKAGE=X

### $K_2(1770)$ DECAY MODES

NODE=M023215;NODE=M023

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \pi \pi$	
$\Gamma_2$ $K_2^*(1430)\pi$	seen
$\Gamma_3$ $K^*(892)\pi$	seen
$\Gamma_4$ $K f_2(1270)$	seen
$\Gamma_5$ $K f_0(980)$	
$\Gamma_6$ $K \phi$	seen
$\Gamma_7$ $K \omega$	seen

DESIG=1;OUR EST;→ UNCHECKED ←  
 DESIG=2;OUR EST;→ UNCHECKED ←  
 DESIG=4;OUR EST;→ UNCHECKED ←  
 DESIG=9;OUR EST;→ UNCHECKED ←  
 DESIG=11  
 DESIG=10  
 DESIG=8

### $K_2(1770)$ BRANCHING RATIOS

NODE=M023220

#### $\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$ $\Gamma_2/\Gamma_1$

$(K_2^*(1430) \rightarrow K\pi)$

NODE=M023R1  
 NODE=M023R1  
 NODE=M023R1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.03		DAUM 81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
~ 1.0		<sup>11</sup> FIRESTONE 72B	DBC +	12 $K^+ d$
< 1.0		COLLEY 71	HBC	10 $K^+ p$
0.2 ± 0.2		AGUILAR-...	70C HBC -	4.6 $K^- p$
< 1.0		BARTSCH 70C	HBC -	10.1 $K^- p$
1.0		BARBARO-...	69 HBC +	12.0 $K^+ p$

<sup>11</sup> Produced in conjunction with excited deuteron.

NODE=M023R1;LINKAGE=P

#### $\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$ $\Gamma_3/\Gamma_1$

NODE=M023R3  
 NODE=M023R3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.23		DAUM 81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
--------	--	----------	------	-----------------------------------

#### $\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$ $\Gamma_4/\Gamma_1$

$(f_2(1270) \rightarrow \pi\pi)$

NODE=M023R4  
 NODE=M023R4  
 NODE=M023R4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.74		DAUM 81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$
--------	--	----------	------	-----------------------------------

#### $\Gamma(K f_0(980))/\Gamma_{total}$ $\Gamma_5/\Gamma$

NODE=M023R6  
 NODE=M023R6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow$ $K_S^0 K_S^0 K_L^0 X$
---------------	--	---------------	------	---

$\Gamma(K\phi)/\Gamma_{\text{total}}$						$\Gamma_6/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	24k	12 AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi\phi K^+$	NODE=M023R5
seen		ARMSTRONG	83	OMEG	$18.5 K^- p \rightarrow K^- \phi N$	NODE=M023R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	4289	13,14 AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi\phi K^+$	NODE=M023R5;LINKAGE=C
					<sup>12</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 7.9 $\sigma$ .	NODE=M023R5;LINKAGE=A
					<sup>13</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.0 $\sigma$ .	NODE=M023R5;LINKAGE=B
					<sup>14</sup> Superseded by AAIJ 21E.	

$\Gamma(K\omega)/\Gamma_{\text{total}}$						$\Gamma_7/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		OTTER	81	HBC	$\pm$ 8.25,10,16 $K^\pm p$	NODE=M023R2
seen		CHUNG	74	HBC	$-$ 7.3 $K^- p \rightarrow K^- \omega p$	NODE=M023R2

## $K_2(1770)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=43597
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)	REFID=20569
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=22801
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
OTTER	81	NP B181 1	G. Otter	(AACH3, BERL, LOIC, VIEN, BIRM+)	REFID=22549
CHUNG	74	PL 51B 413	S.U. Chung <i>et al.</i>	(BNL)	REFID=22735
BLIEDEN	72	PL 39B 668	H.R. Blieden <i>et al.</i>	(STON, NEAS)	REFID=22788
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)	REFID=22506
COLLEY	71	NP B26 71	D.C. Colley <i>et al.</i>	(BIRM, GLAS)	REFID=22785
DENEGRI	71	NP B28 13	D. Denegri <i>et al.</i>	(JHU) JP	REFID=22497
AGUILAR...	70C	PRL 25 54	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=22782
BARTSCH	70C	PL 33B 186	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN+)	REFID=22783
LUDLAM	70	PR D2 1234	T. Ludlam, J. Sandweiss, A.J. Slaughter	(YALE)	REFID=22784
BARBARO...	69	PRL 22 1207	A. Barbaro-Galtieri <i>et al.</i>	(LRL)	REFID=22483

NODE=M023

$K_3^*(1780)$

$$I(J^P) = \frac{1}{2}(3^-)$$

NODE=M060

## $K_3^*(1780)$ T-MATRIX POLE $\sqrt{s}$

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT	
(1754 $\pm$ 13) - i (119 $\pm$ 14)		<sup>1</sup> PELAEZ	17	RVUE $\pi K \rightarrow \pi K$	NODE=M060PP

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M060PP

NODE=M060PP;LINKAGE=A

## $K_3^*(1780)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>1779 <math>\pm</math> 8 OUR AVERAGE</b>		Error includes scale factor of 1.2.				NODE=M060M
1813 $\pm$ 15 $^{+65}_{-16}$	18k	<sup>1</sup> ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$	NODE=M060M
1781 $\pm$ 8 $\pm$ 4		<sup>2</sup> ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$	
1740 $\pm$ 14 $\pm$ 15		<sup>2</sup> ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
1779 $\pm$ 11		<sup>3</sup> BALDI	76	SPEC	+ 10 $K^+ p \rightarrow K^0 \pi^+ p$	
1776 $\pm$ 26		<sup>4</sup> BRANDENB...	76D	ASPK	0 13 $K^\pm p \rightarrow K^\pm \pi^\mp N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1720 $\pm$ 10 $\pm$ 15	6111	<sup>5</sup> BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
1749 $\pm$ 10		ASTON	88B	LASS	- 11 $K^- p \rightarrow K^- \eta p$	
1780 $\pm$ 9	300	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
1790 $\pm$ 15		BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 2\pi N$	
1784 $\pm$ 9	2060	CLELAND	82	SPEC	$\pm$ 50 $K^+ p \rightarrow K_S^0 \pi^\pm p$	
1786 $\pm$ 15		<sup>6</sup> ASTON	81D	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$	
1762 $\pm$ 9	190	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
1850 $\pm$ 50		ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$	
1812 $\pm$ 28		BEUSCH	78	OMEG	0 10 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
1786 $\pm$ 8		CHUNG	78	MPS	0 6 $K^- p \rightarrow K^- \pi^+ n$	

NODE=M060M

NODE=M060M

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (2.0 \pm 0.4^{+1.9}_{-0.4}) \times 10^{-6}$ .

<sup>2</sup> From energy-independent partial-wave analysis.

<sup>3</sup> From a fit to  $Y_6^2$  moment.  $J^P = 3^-$  found.

<sup>4</sup> Confirmed by phase shift analysis of ESTABROOKS 78, yields  $J^P = 3^-$ .

<sup>5</sup> From a partial wave amplitude analysis.

<sup>6</sup> From a fit to the  $Y_6^0$  moment.

NODE=M060M;LINKAGE=D

NODE=M060M;LINKAGE=K

NODE=M060M;LINKAGE=M

NODE=M060M;LINKAGE=A

NODE=M060M;LINKAGE=F

NODE=M060M;LINKAGE=J

### $K_3^*(1780)$ WIDTH

NODE=M060W

NODE=M060W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>161±17 OUR AVERAGE</b>		Error includes scale factor of 1.1.			
191 <sup>+43+3</sup> <sub>-37-81</sub>	1.8k	<sup>1</sup> ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
203±30±8		<sup>2</sup> ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
171±42±20		<sup>2</sup> ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
135±22		<sup>3</sup> BALDI	76	SPEC	+ 10 $K^+ p \rightarrow K^0 \pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
187±31±20	6111	<sup>4</sup> BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
193 <sup>+51</sup> <sub>-37</sub>		ASTON	88B	LASS	- 11 $K^- p \rightarrow K^- \eta p$
99±30	300	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 130		BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 2\pi N$
191±24	2060	CLELAND	82	SPEC	± 50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
225±60		<sup>5</sup> ASTON	81D	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
~ 80	190	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
240±50		ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
181±44		<sup>6</sup> BEUSCH	78	OMEG	10 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
96±31		CHUNG	78	MPS	0 6 $K^- p \rightarrow K^- \pi^+ n$
270±70		<sup>7</sup> BRANDENB...	76D	ASPK	0 13 $K^\pm p \rightarrow K^\pm \pi^\mp N$

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (2.0 \pm 0.4^{+1.9}_{-0.4}) \times 10^{-6}$ .

<sup>2</sup> From energy-independent partial-wave analysis.

<sup>3</sup> From a fit to  $Y_6^2$  moment.  $J^P = 3^-$  found.

<sup>4</sup> From a partial wave amplitude analysis.

<sup>5</sup> From a fit to  $Y_6^0$  moment.

<sup>6</sup> Errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.

<sup>7</sup> ESTABROOKS 78 find that BRANDENBURG 76D data are consistent with 175 MeV width. Not averaged.

NODE=M060W;LINKAGE=A

NODE=M060W;LINKAGE=K

NODE=M060W;LINKAGE=M

NODE=M060W;LINKAGE=F

NODE=M060W;LINKAGE=J

NODE=M060W;LINKAGE=D

NODE=M060W;LINKAGE=E

### $K_3^*(1780)$ DECAY MODES

NODE=M060215;NODE=M060

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $K\rho$	(31 ± 9 ) %	DESIG=3
$\Gamma_2$ $K^*(892)\pi$	(20 ± 5 ) %	DESIG=2
$\Gamma_3$ $K\pi$	(18.8± 1.0) %	DESIG=1
$\Gamma_4$ $K\eta$	(30 ± 13 ) %	DESIG=6
$\Gamma_5$ $K_2^*(1430)\pi$	< 16 %	95% DESIG=4

## CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 4 measurements and one constraint to determine 4 parameters. The overall fit has a  $\chi^2 = 0.0$  for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	85		
$x_3$	18	21	
$x_4$	-98	-94	-27
	$x_1$	$x_2$	$x_3$

### K<sub>3</sub><sup>\*</sup>(1780) BRANCHING RATIOS

NODE=M060220

 $\Gamma(K\rho)/\Gamma(K^*(892)\pi)$ 
 $\Gamma_1/\Gamma_2$ NODE=M060R5  
NODE=M060R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>1.52±0.23 OUR FIT</b>				
<b>1.52±0.21±0.10</b>	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

 $\Gamma(K^*(892)\pi)/\Gamma(K\pi)$ 
 $\Gamma_2/\Gamma_3$ NODE=M060R7  
NODE=M060R7

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>1.09±0.26 OUR FIT</b>				
<b>1.09±0.26</b>	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

 $\Gamma(K\pi)/\Gamma_{\text{total}}$ 
 $\Gamma_3/\Gamma$ NODE=M060R4  
NODE=M060R4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.188±0.010 OUR FIT</b>				
<b>0.188±0.010 OUR AVERAGE</b>				
0.187±0.008±0.008	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
0.19 ±0.02	ESTABROOKS	78	ASPK	0 13 $K^\pm p \rightarrow K\pi N$

 $\Gamma(K\eta)/\Gamma(K\pi)$ 
 $\Gamma_4/\Gamma_3$ NODE=M060R8  
NODE=M060R8

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>1.6 ±0.7 OUR FIT</b>				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.41±0.050	<sup>1</sup> BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
0.50±0.18	ASTON	88B	LASS	- 11 $K^- p \rightarrow K^- \eta p$

<sup>1</sup> This result supersedes ASTON 88B.

NODE=M060R8;LINKAGE=H

 $\Gamma(K_2^*(1430)\pi)/\Gamma(K^*(892)\pi)$ 
 $\Gamma_5/\Gamma_2$ NODE=M060R6  
NODE=M060R6

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>&lt;0.78</b>	95	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

### K<sub>3</sub><sup>\*</sup>(1780) REFERENCES

NODE=M060

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira		REFID=57836
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP	REFID=40281
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
ASTON	84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)	REFID=22763
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81D	PL 99B 502	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22820
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
BEUSCH	78	PL 74B 282	W. Beusch <i>et al.</i>	(CERN, AACH3, ETH) JP	REFID=22537
CHUNG	78	PRL 40 355	S.U. Chung <i>et al.</i>	(BNL, BRAN, CUNY+) JP	REFID=22814
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
BALDI	76	PL 63B 344	R. Baldi <i>et al.</i>	(GEVA) JP	REFID=22807
BRANDENB...	76D	PL 60B 478	G.W. Brandenburg <i>et al.</i>	(SLAC) JP	REFID=22808

**$K_2(1820)$** 

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this *Review* (PDG 04) under  $K_2(1770)$ .

NODE=M146

NODE=M146

 **$K_2(1820)$  MASS**

NODE=M146M

NODE=M146M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1819±12 OUR AVERAGE</b>				
$1853 \pm 27^{+18}_{-35}$	4289	<sup>1</sup> AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$1816 \pm 13$		<sup>2</sup> ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 1840		<sup>3</sup> DAUM	81C CNTR	$63 K^- p \rightarrow K^- 2\pi p$
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 $\sigma$ .				
<sup>2</sup> From a partial wave analysis of the $K^- \omega$ system.				
<sup>3</sup> From a partial wave analysis of the $K^- 2\pi$ system.				

NODE=M146M;LINKAGE=B

NODE=M146M;LINKAGE=A

NODE=M146M;LINKAGE=C

 **$K_2(1820)$  WIDTH**

NODE=M146W

NODE=M146W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>264±34 OUR AVERAGE</b>				
$167 \pm 58^{+82}_{-72}$	4289	<sup>1</sup> AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
$276 \pm 35$		<sup>2</sup> ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 230		<sup>3</sup> DAUM	81C CNTR	$63 K^- p \rightarrow K^- 2\pi p$
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 $\sigma$ .				
<sup>2</sup> From a partial wave analysis of the $K^- \omega$ system.				
<sup>3</sup> From a partial wave analysis of the $K^- 2\pi$ system.				

NODE=M146W;LINKAGE=A

NODE=M146W;LINKAGE=B

NODE=M146W;LINKAGE=C

 **$K_2(1820)$  DECAY MODES**

NODE=M146215;NODE=M146

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \pi \pi$	seen
$\Gamma_2$ $K_2^*(1430)\pi$	seen
$\Gamma_3$ $K^*(892)\pi$	seen
$\Gamma_4$ $K f_2(1270)$	seen
$\Gamma_5$ $K \omega$	seen
$\Gamma_6$ $K \phi$	seen

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

DESIG=7

 **$K_2(1820)$  BRANCHING RATIOS**

NODE=M146220

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$	$\Gamma_2/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 0.77	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K} 2\pi p$
$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$	$\Gamma_3/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 0.05	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K} 2\pi p$
$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$	$\Gamma_4/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 0.18	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K} 2\pi p$

NODE=M146R1

NODE=M146R1

NODE=M146R2

NODE=M146R2

NODE=M146R3

NODE=M146R3

$\Gamma(K\phi)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
seen	4289	<sup>2,3</sup> AAIJ	17C LHCB	$B^+ \rightarrow J/\psi\phi K^+$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 5.8  $\sigma$ .  
<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 3.0  $\sigma$ .  
<sup>3</sup> Superseded by AAIJ 21E.

NODE=M146R00  
 NODE=M146R00

NODE=M146R00;LINKAGE=B  
 NODE=M146R00;LINKAGE=A  
 NODE=M146R00;LINKAGE=C

## $K_2(1820)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

NODE=M146

REFID=61150  
 REFID=57657  
 REFID=57636  
 REFID=49653  
 REFID=43597  
 REFID=22548

NODE=M088

$K(1830)$

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of  $K\phi$  system. Needs confirmation.

NODE=M088

## $K(1830)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$1874 \pm 43^{+59}_{-115}$	4289	<sup>1,2</sup> AAIJ	17C LHCB		$B^+ \rightarrow J/\psi\phi K^+$
$\sim 1830$		ARMSTRONG 83	OMEG	-	$18.5 K^- p \rightarrow 3K p$

- <sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 3.5  $\sigma$ .  
<sup>2</sup> A subsequent amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  by AAIJ 21E did not confirm this measurement.

NODE=M088M;LINKAGE=A  
 NODE=M088M;LINKAGE=B

NODE=M088M

NODE=M088M

## $K(1830)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$168 \pm 90^{+280}_{-104}$	4289	<sup>3,4</sup> AAIJ	17C LHCB		$B^+ \rightarrow J/\psi\phi K^+$
$\sim 250$		ARMSTRONG 83	OMEG	-	$18.5 K^- p \rightarrow 3K p$

- <sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 3.5  $\sigma$ .  
<sup>4</sup> A subsequent amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  by AAIJ 21E did not confirm this measurement.

NODE=M088W

NODE=M088W

NODE=M088W;LINKAGE=A  
 NODE=M088W;LINKAGE=B

## $K(1830)$ DECAY MODES

NODE=M088215;NODE=M088

Mode

$\Gamma_1$   $K\phi$

DESIG=1

## $K(1830)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)JP

NODE=M088

REFID=61150  
 REFID=57657  
 REFID=57636  
 REFID=22801



**$K_0^*(1950)$** 

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the  $K^- \pi^+$  system. Needs confirmation.

NODE=M134

NODE=M134

 **$K_0^*(1950)$  MASS**

NODE=M134M

NODE=M134M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>1944±18 OUR AVERAGE</b>				
1942±22±21	LEES	21A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' K^+ K^-$
1945±10±20	<sup>1</sup> ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1917±12	<sup>2</sup> ZHOU	06	RVUE	$K p \rightarrow K^- \pi^+ n$
1820±40	<sup>3</sup> ANISOVICH	97C	RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
<sup>1</sup> We take the central value of the two solutions and the larger error given.				
<sup>2</sup> S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$ , $K_0^*(1430)$ .				
<sup>3</sup> T-matrix pole. Reanalysis of ASTON 88 data.				

NODE=M134M;LINKAGE=A

NODE=M134M;LINKAGE=ZU

NODE=M134M;LINKAGE=A1

 **$K_0^*(1950)$  WIDTH**

NODE=M134W

NODE=M134W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>100± 40 OUR AVERAGE</b> Error includes scale factor of 1.3.				
80± 32±20	LEES	21A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' K^+ K^-$
201± 34±79	<sup>4</sup> ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
145± 38	<sup>5</sup> ZHOU	06	RVUE	$K p \rightarrow K^- \pi^+ n$
250±100	<sup>6</sup> ANISOVICH	97C	RVUE	$11 K^- p \rightarrow K^- \pi^+ n$
<sup>4</sup> We take the central value of the two solutions and the larger error given.				
<sup>5</sup> S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$ , $K_0^*(1430)$ .				
<sup>6</sup> T-matrix pole. Reanalysis of ASTON 88 data.				

NODE=M134W;LINKAGE=A

NODE=M134W;LINKAGE=ZU

NODE=M134W;LINKAGE=A1

 **$K_0^*(1950)$  DECAY MODES**

NODE=M134215;NODE=M134

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad K^- \pi^+$	(52±14) %

DESIG=1

 **$K_0^*(1950)$  BRANCHING RATIOS**

NODE=M134220

$\Gamma(K^- \pi^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.52±0.08±0.12</b>	<sup>7</sup> ASTON	88	LASS 0	$11 K^- p \rightarrow K^- \pi^+ n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 0.60	<sup>8</sup> ZHOU	06	RVUE	$K p \rightarrow K^- \pi^+ n$
<sup>7</sup> We take the central value of the two solutions and the larger error given.				
<sup>8</sup> S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$ , $K_0^*(1430)$ .				

NODE=M134R1

NODE=M134R1

NODE=M134R1;LINKAGE=A

NODE=M134R1;LINKAGE=ZU

 **$K_0^*(1950)$  REFERENCES**

NODE=M134

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng	
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev	
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

REFID=61442

REFID=51198

REFID=45815

REFID=40262

# K<sub>2</sub><sup>\*</sup>(1980)

$$I(J^P) = \frac{1}{2}(2^+)$$

Needs confirmation.

NODE=M104

NODE=M104

NODE=M104M

NODE=M104M

## K<sub>2</sub><sup>\*</sup>(1980) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>1994<sup>+60</sup><sub>-50</sub> OUR AVERAGE</b> Error includes scale factor of 2.8. See the ideogram below.					
2046 <sup>+17</sup> <sub>-16</sub>	67/15	1 ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
1868 ± 8	40/57	183k ABLIKIM	19AQ	BES ±	$J/\psi \rightarrow K^+ K^- \pi^0$
1973 ± 8	± 25	ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2073 ± 94	245/240	4289 2,3 AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
2020 ± 20		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1978 ± 40	241	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

<sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (7.0 \pm 0.5^{+3.7}_{-0.6}) \times 10^{-6}$ .


<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 5.4  $\sigma$ .

<sup>3</sup> A reanalysis by AAIJ 21E using a larger data sample did not confirm this measurement, the new result having a significance of only 1.6  $\sigma$ .

NODE=M104M;LINKAGE=C

NODE=M104M;LINKAGE=B

NODE=M104M;LINKAGE=E



## K<sub>2</sub><sup>\*</sup>(1980) WIDTH

NODE=M104W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>348<sup>+50</sup><sub>-30</sub> OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below.					
408 <sup>+38</sup> <sub>-34</sub>	72/44	1.8k 1 ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
272 ± 24	50/15	183k ABLIKIM	19AQ	BES ±	$J/\psi \rightarrow K^+ K^- \pi^0$
373 ± 33	± 60	ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
678 ± 311	1153/559	4289 2,3 AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
180 ± 70		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
398 ± 47	241	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$


NODE=M104W

- <sup>1</sup> Seen in  $\psi(2S)$  decay with branching ratio  $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (7.0 \pm 0.5^{+3.7}_{-0.6}) \times 10^{-6}$ .
- <sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.4 \sigma$ .
- <sup>3</sup> A reanalysis by AAIJ 21E using a larger data sample did not confirm this measurement, the new result having a significance of only  $1.6 \sigma$ .

NODE=M104W;LINKAGE=C

NODE=M104W;LINKAGE=B

NODE=M104W;LINKAGE=E

 **$K_2^*(1980)$  DECAY MODES**

NODE=M104215;NODE=M104

Mode	Fraction ( $\Gamma_i/\Gamma$ )	
$\Gamma_1$ $K^*(892)\pi$	possibly seen	DESIG=2
$\Gamma_2$ $K\rho$	possibly seen	DESIG=3
$\Gamma_3$ $K f_2(1270)$	possibly seen	DESIG=4
$\Gamma_4$ $K\phi$	seen	DESIG=5
$\Gamma_5$ $K\eta$	seen	DESIG=6

 **$K_2^*(1980)$  BRANCHING RATIOS**

NODE=M104220

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
possibly seen	GULER	11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
$\Gamma(K\rho)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
possibly seen	GULER	11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	$\Gamma_2/\Gamma_1$	
VALUE	DOCUMENT ID	TECN CHG COMMENT
<b>1.49 ± 0.24 ± 0.09</b>	ASTON	87 LASS 0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
$\Gamma(K f_2(1270))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
possibly seen	TIKHOMIROV	03 SPEC 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
$\Gamma(K\phi)/\Gamma_{\text{total}}$	$\Gamma_4/\Gamma$	
VALUE	EVTS	DOCUMENT ID TECN COMMENT
seen	4289	1,2 AAIJ 17C LHCB $B^+ \rightarrow J/\psi \phi K^+$

NODE=M104R01  
NODE=M104R01NODE=M104R02  
NODE=M104R02NODE=M104R1  
NODE=M104R1NODE=M104R3  
NODE=M104R3NODE=M104R00  
NODE=M104R00<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.4 \sigma$ .

NODE=M104R00;LINKAGE=A

<sup>2</sup> A reanalysis by AAIJ 21E using a larger data sample did not confirm this measurement, the new result having a significance of only  $1.6 \sigma$ .

NODE=M104R00;LINKAGE=C

$\Gamma(K\eta)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1.8k	<sup>1</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
seen	116k	<sup>2</sup> CHEN	20A BELLE	$D^0 \rightarrow K^- \pi^+ \eta$

<sup>1</sup> Seen decaying to  $K\eta$  in an amplitude analysis of  $\psi(2S) \rightarrow K^+ K^- \eta$ .

<sup>2</sup> From an amplitude analysis of the decay  $D^0 \rightarrow K^- \pi^+ \eta$  with a significance of 17  $\sigma$ .

NODE=M104R03  
NODE=M104R03

NODE=M104R03;LINKAGE=A  
NODE=M104R03;LINKAGE=B

### $K_2^*(1980)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
BIRD	89	SLAC-332	P.F. Bird	(SLAC)
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

NODE=M104

REFID=61150  
REFID=60256  
REFID=60333  
REFID=59909  
REFID=57657  
REFID=57636  
REFID=53668  
REFID=49423

REFID=41002  
REFID=40234

NODE=M035

$K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

### $K_4^*(2045)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**2048<sup>+</sup><sub>-9</sub> OUR AVERAGE** Error includes scale factor of 1.1.

2090 $\pm$ 9 <sup>+11</sup> <sub>-29</sub>	183k	ABLIKIM	19AQ BES	$\pm$	$J/\psi \rightarrow K^+ K^- \pi^0$
2062 $\pm$ 14 $\pm$ 13		<sup>1</sup> ASTON	86 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
2039 $\pm$ 10	400	<sup>2,3</sup> CLELAND	82 SPEC	$\pm$	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
2070 <sup>+100</sup> <sub>-40</sub>		<sup>4</sup> ASTON	81C LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
•••					We do not use the following data for averages, fits, limits, etc. •••
2079 $\pm$ 7	431	TORRES	86 MPSF		400 $pA \rightarrow 4KX$
2088 $\pm$ 20	650	BAUBILLIER	82 HBC	-	8.25 $K^- p \rightarrow K_S^0 \pi^- p$
2115 $\pm$ 46	488	CARMONY	77 HBC	0	9 $K^+ d \rightarrow K^+ \pi^+ s X$

<sup>1</sup> From a fit to all moments.

<sup>2</sup> From a fit to 8 moments.

<sup>3</sup> Number of events evaluated by us.

<sup>4</sup> From energy-independent partial-wave analysis.

NODE=M035M

NODE=M035M

NODE=M035M;LINKAGE=E  
NODE=M035M;LINKAGE=B  
NODE=M035M;LINKAGE=W  
NODE=M035M;LINKAGE=D

### $K_4^*(2045)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

**199<sup>+</sup><sub>-19</sub> OUR AVERAGE**

201 $\pm$ 19 <sup>+57</sup> <sub>-17</sub>	183k	ABLIKIM	19AQ BES	$\pm$	$J/\psi \rightarrow K^+ K^- \pi^0$
221 $\pm$ 48 $\pm$ 27		<sup>5</sup> ASTON	86 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
189 $\pm$ 35	400	<sup>6,7</sup> CLELAND	82 SPEC	$\pm$	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
•••					We do not use the following data for averages, fits, limits, etc. •••
61 $\pm$ 58	431	TORRES	86 MPSF		400 $pA \rightarrow 4KX$
170 <sup>+100</sup> <sub>-50</sub>	650	BAUBILLIER	82 HBC	-	8.25 $K^- p \rightarrow K_S^0 \pi^- p$
240 <sup>+500</sup> <sub>-100</sub>		<sup>8</sup> ASTON	81C LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
300 $\pm$ 200		CARMONY	77 HBC	0	9 $K^+ d \rightarrow K^+ \pi^+ s X$

<sup>5</sup> From a fit to all moments.

<sup>6</sup> From a fit to 8 moments.

<sup>7</sup> Number of events evaluated by us.

<sup>8</sup> From energy-independent partial-wave analysis.

NODE=M035W

NODE=M035W

NODE=M035W;LINKAGE=E  
NODE=M035W;LINKAGE=B  
NODE=M035W;LINKAGE=W  
NODE=M035W;LINKAGE=D

**$K_4^*(2045)$  DECAY MODES**

NODE=M035215;NODE=M035

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K\pi$	(9.9±1.2) %
$\Gamma_2$ $K^*(892)\pi\pi$	(9 ±5 ) %
$\Gamma_3$ $K^*(892)\pi\pi\pi$	(7 ±5 ) %
$\Gamma_4$ $\rho K\pi$	(5.7±3.2) %
$\Gamma_5$ $\omega K\pi$	(5.0±3.0) %
$\Gamma_6$ $\phi K\pi$	(2.8±1.4) %
$\Gamma_7$ $\phi K^*(892)$	(1.4±0.7) %

DESIG=1  
DESIG=2  
DESIG=5  
DESIG=3  
DESIG=4  
DESIG=6  
DESIG=7 **$K_4^*(2045)$  BRANCHING RATIOS**

NODE=M035220

 **$\Gamma(K\pi)/\Gamma_{\text{total}}$**   **$\Gamma_1/\Gamma$** NODE=M035R1  
NODE=M035R1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.099±0.012</b>	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

 **$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$**   **$\Gamma_2/\Gamma_1$** NODE=M035R2  
NODE=M035R2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.89±0.53</b>	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$

 **$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$**   **$\Gamma_3/\Gamma_1$** NODE=M035R5  
NODE=M035R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.75±0.49</b>	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$

 **$\Gamma(\rho K\pi)/\Gamma(K\pi)$**   **$\Gamma_4/\Gamma_1$** NODE=M035R3  
NODE=M035R3

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.58±0.32</b>	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$

 **$\Gamma(\omega K\pi)/\Gamma(K\pi)$**   **$\Gamma_5/\Gamma_1$** NODE=M035R4  
NODE=M035R4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.50±0.30</b>	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$

 **$\Gamma(\phi K\pi)/\Gamma_{\text{total}}$**   **$\Gamma_6/\Gamma$** NODE=M035R6  
NODE=M035R6

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.028±0.014</b>	<sup>9</sup> TORRES	86	MPSF 400 $pA \rightarrow 4KX$

 **$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$**   **$\Gamma_7/\Gamma$** NODE=M035R7  
NODE=M035R7

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.014±0.007</b>	<sup>9</sup> TORRES	86	MPSF 400 $pA \rightarrow 4KX$

<sup>9</sup> Error determination is model dependent.

NODE=M035R;LINKAGE=A

 **$K_4^*(2045)$  REFERENCES**

NODE=M035

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=22462
TORRES	86	PR D34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)	REFID=22845
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22842
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22821
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)	REFID=22811

**$K_2(2250)$** 

$$I(J^P) = \frac{1}{2}(2^-)$$

## OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems reported in the 2150–2260 MeV region, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the  $J^P = 2^-$  wave.

NODE=M040

NODE=M040

 **$K_2(2250)$  MASS**

NODE=M040M

NODE=M040M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>2247±17 OUR AVERAGE</b>						
2200±40		<sup>1</sup> ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$	
2235±50		<sup>1</sup> BAUBILLIER 81	HBC	-	8 $K^- p \rightarrow \Lambda \bar{p} X$	
2260±20		<sup>1</sup> CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2280±20		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
2147±4	37	CHLIAPNIK...	79	HBC	+	32 $K^+ p \rightarrow \bar{\Lambda} p X$
2240±20	20	LISSAUER 70	HBC		9 $K^+ p$	
<sup>1</sup> $J^P = 2^-$ from moments analysis.						

NODE=M040M;LINKAGE=Q

 **$K_2(2250)$  WIDTH**

NODE=M040W

NODE=M040W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>180±30 OUR AVERAGE</b>						
Error includes scale factor of 1.4.						
150±30		<sup>2</sup> ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$	
210±30		<sup>2</sup> CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
180±60		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
~ 200		<sup>2</sup> BAUBILLIER 81	HBC	-	8 $K^- p \rightarrow \Lambda \bar{p} X$	
~ 40	37	CHLIAPNIK...	79	HBC	+	32 $K^+ p \rightarrow \bar{\Lambda} p X$
80±20	20	LISSAUER 70	HBC		9 $K^+ p$	
<sup>2</sup> $J^P = 2^-$ from moments analysis.						

NODE=M040W;LINKAGE=Q

 **$K_2(2250)$  DECAY MODES**

NODE=M040215;NODE=M040

Mode	
$\Gamma_1$	$K \pi \pi$
$\Gamma_2$	$K f_2(1270)$
$\Gamma_3$	$K^*(892) f_0(980)$
$\Gamma_4$	$\rho \bar{\Lambda}$

DESIG=1

DESIG=3

DESIG=4

DESIG=2

 **$K_2(2250)$  REFERENCES**

NODE=M040

REFID=49423

TIKHOMIROV 03	PAN 66 828 Translated from YAF 66 860.	G.D. Tikhomirov <i>et al.</i>		
ARMSTRONG 83C	NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=22852
BAUBILLIER 81	NP B183 1	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+) JP	REFID=22850
CLELAND 81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+) JP	REFID=22851
CHLIAPNIK...	NP B158 253	P.V. Chliapnikov <i>et al.</i>	(CERN, BELG, MONS)	REFID=22849
LISSAUER 70	NP B18 491	D. Lissauer <i>et al.</i>	(LBL)	REFID=22847

**$K_3(2320)$** 

$$I(J^P) = \frac{1}{2}(3^+)$$

OMITTED FROM SUMMARY TABLE

Seen in the  $J^P = 3^+$  wave of the antihyperon-nucleon system.  
Needs confirmation.

 **$K_3(2320)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>2324 ± 24 OUR AVERAGE</b>				
2330 ± 40	<sup>1</sup> ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
2320 ± 30	<sup>1</sup> CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$

<sup>1</sup>  $J^P = 3^+$  from moments analysis.

 **$K_3(2320)$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>150 ± 30</b>	<sup>2</sup> ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 250	<sup>2</sup> CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$

<sup>2</sup>  $J^P = 3^+$  from moments analysis.

 **$K_3(2320)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\rho \bar{\Lambda}$	(6.1 ± 1.2) %

 **$K_3(2320)$  REFERENCES**

ARMSTRONG 83C NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
CLELAND 81 NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+)

 **$K_5^*(2380)$** 

$$I(J^P) = \frac{1}{2}(5^-)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

 **$K_5^*(2380)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>2382 ± 14 ± 19</b>	<sup>1</sup> ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

<sup>1</sup> From a fit to all the moments.

 **$K_5^*(2380)$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>178 ± 37 ± 32</b>	<sup>2</sup> ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

<sup>2</sup> From a fit to all the moments.

 **$K_5^*(2380)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $K \pi$	(6.1 ± 1.2) %

 **$K_5^*(2380)$  BRANCHING RATIOS**

$\Gamma(K\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_1/\Gamma$
<b>0.061 ± 0.012</b>	ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	

NODE=M090

NODE=M090

NODE=M090M

NODE=M090M

NODE=M090M;LINKAGE=P

NODE=M090W

NODE=M090W

NODE=M090W;LINKAGE=P

NODE=M090215;NODE=M090

DESIG=1

NODE=M090

REFID=22852  
REFID=22851

NODE=M098

NODE=M098

NODE=M098M

NODE=M098M

NODE=M098M;LINKAGE=E

NODE=M098W

NODE=M098W

NODE=M098W;LINKAGE=E

NODE=M098215;NODE=M098

DESIG=1

NODE=M098220

NODE=M098R1  
NODE=M098R1

**$K_5^*(2380)$  REFERENCES**

ASTON 88 NP B296 493 D. Aston *et al.* (SLAC, NAGO, CINC, INUS)  
 ASTON 86 PL B180 308 D. Aston *et al.* (SLAC, NAGO, CINC, INUS)

NODE=M098

REFID=40262  
REFID=22462

NODE=M091

 **$K_4(2500)$** 

$$I(J^P) = \frac{1}{2}(4^-)$$

OMITTED FROM SUMMARY TABLE  
 Needs confirmation.

NODE=M091

 **$K_4(2500)$  MASS**

NODE=M091M

NODE=M091M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
<b>2490±20</b>	<sup>1</sup> CLELAND	81	SPEC ±	50 $K^+ p \rightarrow \Lambda \bar{p}$

<sup>1</sup>  $J^P = 4^-$  from moments analysis.

NODE=M091M;LINKAGE=R

 **$K_4(2500)$  WIDTH**

NODE=M091W

NODE=M091W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
~ 250	<sup>2</sup> CLELAND	81	SPEC ±	50 $K^+ p \rightarrow \Lambda \bar{p}$

••• We do not use the following data for averages, fits, limits, etc. •••  
<sup>2</sup>  $J^P = 4^-$  from moments analysis.

NODE=M091W;LINKAGE=R

 **$K_4(2500)$  DECAY MODES**

NODE=M091215;NODE=M091

Mode

 $\Gamma_1 \quad p \bar{\Lambda}$ 

DESIG=1

 **$K_4(2500)$  REFERENCES**

CLELAND 81 NP B184 1 W.E. Cleland *et al.* (PITT, GEVA, LAUS+)

NODE=M091

REFID=22851

NODE=M129

 **$K(3100)$** 

$$I^G(J^{PC}) = ?^?(?^?)$$

OMITTED FROM SUMMARY TABLE

Narrow peak observed in several ( $\Lambda \bar{p} +$  pions) and ( $\bar{\Lambda} p +$  pions) states in  $\Sigma^-$  Be reactions by BOURQUIN 86 and in  $np$  and  $nA$  reactions by ALEEV 93. Not seen by BOEHNLEIN 91. If due to strong decays, this state has exotic quantum numbers ( $B=0, Q=+1, S=-1$  for  $\Lambda \bar{p} \pi^+ \pi^+$  and  $I \geq 3/2$  for  $\Lambda \bar{p} \pi^-$ ). Needs confirmation.

NODE=M129

 **$K(3100)$  MASS**

NODE=M129205

VALUE (MeV)	DOCUMENT ID
<b>≈ 3100 OUR ESTIMATE</b>	

NODE=M129M  
→ UNCHECKED ←**3-BODY DECAYS**NODE=M129M1  
NODE=M129M1

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>3054±11 OUR AVERAGE</b>			
3060± 7±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^+$
3056± 7±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^-$
3055± 8±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^-$
3045± 8±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^+$

OCCUR=2

OCCUR=3

OCCUR=4

**4-BODY DECAYS**NODE=M129M2  
NODE=M129M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>3059±11 OUR AVERAGE</b>			
3067± 6±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+$
3060± 8±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^-$
3055± 7±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^- \pi^-$
3052± 8±20	<sup>1</sup> ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^- \pi^+$

OCCUR=2

OCCUR=3

OCCUR=4

••• We do not use the following data for averages, fits, limits, etc. •••

3105±30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+$
3115±30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^-$

OCCUR=2



**5-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3095 ± 30	BOURQUIN	86	SPEC	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$
<sup>1</sup> Supersedes ALEEV 90.				

NODE=M129M3  
NODE=M129M3

NODE=M129M;LINKAGE=A

**K(3100) WIDTH**

NODE=M129210

**3-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

42 ± 16	<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^+$
36 ± 15	<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^-$
50 ± 18	<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^-$
30 ± 15	<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^+$

NODE=M129W1  
NODE=M129W1

OCCUR=2

OCCUR=3

OCCUR=4

**4-BODY DECAYS**

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 8		<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+$
28 ± 12		<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^-$
32 ± 15		<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^- \pi^-$
30 ± 15		<sup>2</sup> ALEEV	93	BIS2	$K(3100) \rightarrow \bar{\Lambda} p \pi^- \pi^+$
<30	90	BOURQUIN	86	SPEC	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+$
<80	90	BOURQUIN	86	SPEC	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^-$

NODE=M129W2  
NODE=M129W2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

**5-BODY DECAYS**

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	90	BOURQUIN	86	SPEC	$K(3100) \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$
<sup>2</sup> Supersedes ALEEV 90.					

NODE=M129W3  
NODE=M129W3

NODE=M129W;LINKAGE=A

**K(3100) DECAY MODES**

NODE=M129215;NODE=M129

Mode

$\Gamma_1$	$K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+$	DESIG=1
$\Gamma_2$	$K(3100)^{--} \rightarrow \Lambda \bar{p} \pi^-$	DESIG=2
$\Gamma_3$	$K(3100)^- \rightarrow \Lambda \bar{p} \pi^+ \pi^-$	DESIG=3
$\Gamma_4$	$K(3100)^+ \rightarrow \Lambda \bar{p} \pi^+ \pi^+$	DESIG=4
$\Gamma_5$	$K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$	DESIG=5
$\Gamma_6$	$K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$	DESIG=6

 **$\Gamma(\Sigma(1385)^+ \bar{p})/\Gamma(\Lambda \bar{p} \pi^+)$**  **$\Gamma_6/\Gamma_1$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.04	90	ALEEV	93	BIS2	$K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$

NODE=M129R1  
NODE=M129R1

**K(3100) REFERENCES**

NODE=M129

ALEEV	93	PAN 56 1358 Translated from YAF 56 100.	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)	REFID=43668
BOEHNLEIN	91	NPBPS B21 174	A. Boehnlein <i>et al.</i>	(FLOR, BNL, IND+)	REFID=41743
ALEEV	90	ZPHY C47 533	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)	REFID=42173
BOURQUIN	86	PL B172 113	M.H. Bourquin <i>et al.</i>	(GEVA, RAL, HEIDP+)	REFID=22928

# CHARMED MESONS

## (C = ±1)

$D^+ = c\bar{d}$ ,  $D^0 = c\bar{u}$ ,  $\bar{D}^0 = \bar{c}u$ ,  $D^- = \bar{c}d$ , similarly for  $D^{*}$ 's

### $D^*(2007)^0$

$I(J^P) = \frac{1}{2}(1^-)$   
I, J, P need confirmation.

J consistent with 1, value 0 ruled out (NGUYEN 77).

### $D^*(2007)^0$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ ,  
and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**2006.85 ± 0.05 OUR FIT** Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2006 ± 1.5	<sup>1</sup> GOLDHABER 77	MRK1	$e^+e^-$
------------	---------------------------	------	----------

<sup>1</sup>From simultaneous fit to  $D^*(2010)^+$ ,  $D^*(2007)^0$ ,  $D^+$ , and  $D^0$ .

### $m_{D^*(2007)^0} - m_{D^0}$

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ ,  
and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**142.014 ± 0.030 OUR FIT** Error includes scale factor of 1.5.

**142.016 ± 0.030 OUR AVERAGE** Error includes scale factor of 1.5.

142.007 ± 0.015 ± 0.014	10k	<sup>1</sup> TOMARADZE 15	CLEO	$e^+e^- \rightarrow$ hadrons
-------------------------	-----	---------------------------	------	------------------------------

142.2 ± 0.3 ± 0.2	145	ALBRECHT 95F	ARG	$e^+e^- \rightarrow$ hadrons
-------------------	-----	--------------	-----	------------------------------

142.12 ± 0.05 ± 0.05	1176	BORTOLETTO92B	CLE2	$e^+e^- \rightarrow$ hadrons
----------------------	------	---------------	------	------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

142.2 ± 2.0	SADROZINSKI 80	CBAL	$D^{*0} \rightarrow D^0\pi^0$
-------------	----------------	------	-------------------------------

142.7 ± 1.7	<sup>2</sup> GOLDHABER 77	MRK1	$e^+e^-$
-------------	---------------------------	------	----------

<sup>1</sup>Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. This value comes from the average of the results for two decay modes,  $D^0 \rightarrow K^-\pi^+$  and  $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$ .

<sup>2</sup>From simultaneous fit to  $D^*(2010)^+$ ,  $D^*(2007)^0$ ,  $D^+$ , and  $D^0$ .

### $D^*(2007)^0$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

<b>&lt;2.1</b>	90	<sup>1</sup> ABACHI	88B HRS	$D^{*0} \rightarrow D^+\pi^-$
----------------	----	---------------------	---------	-------------------------------

<sup>1</sup>Assuming  $m_{D^{*0}} = 2007.2 \pm 2.1$  MeV/ $c^2$ .

### $D^*(2007)^0$ DECAY MODES

$\bar{D}^*(2007)^0$  modes are charge conjugates of modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^0\pi^0$	(64.7 ± 0.9) %
$\Gamma_2$ $D^0\gamma$	(35.3 ± 0.9) %
$\Gamma_3$ $D^0e^+e^-$	(3.91 ± 0.33) × 10 <sup>-3</sup>

NODE=MXXX035

NODE=MXXX035

NODE=M061

NODE=M061

NODE=M061M

NODE=M061M

NODE=M061M

NODE=M061M;LINKAGE=G

NODE=M061DM

NODE=M061DM

NODE=M061DM

NODE=M061DM;LINKAGE=A

NODE=M061DM;LINKAGE=G

NODE=M061W

NODE=M061W

NODE=M061W;LINKAGE=A

NODE=M061220;NODE=M061

NODE=M061

DESIG=1

DESIG=2

DESIG=3

## CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 5 measurements and one constraint to determine 2 parameters. The overall fit has a  $\chi^2 = 2.5$  for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$x_2 \begin{vmatrix} & -100 \\ & x_1 \end{vmatrix}$$

### $D^*(2007)^0$ BRANCHING RATIOS

NODE=M061225

#### $\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$

 $\Gamma_1 / \Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**1.83 ± 0.07 OUR FIT** Error includes scale factor of 1.1.

**1.85 ± 0.07 OUR AVERAGE**

1.90 ± 0.07 ± 0.05	4.9k	ABLIKIM	15B BES3	10.6 $e^+ e^- \rightarrow$ hadrons
--------------------	------	---------	----------	------------------------------------

1.74 ± 0.02 ± 0.13		AUBERT,BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons
--------------------	--	-----------	----------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.789 ± 0.082		<sup>1</sup> AAIJ	22N LHCb	$B^0, B^0_s \rightarrow \bar{D}^{*0}(K\pi, \pi\pi)$
---------------	--	-------------------	----------	---

<sup>1</sup>Statistical error only.

NODE=M061R3  
NODE=M061R3

#### $\Gamma(D^0 e^+ e^-) / \Gamma(D^0 \gamma)$

 $\Gamma_3 / \Gamma_2$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**11.08 ± 0.76 ± 0.49** 421 ABLIKIM 21BD BES3 4.178 GeV  $e^+ e^-$

NODE=M061R00  
NODE=M061R00

#### $\Gamma(D^0 \pi^0) / \Gamma_{\text{total}}$

 $\Gamma_1 / \Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.647 ± 0.009 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.655 ± 0.008 ± 0.005	3.2k	<sup>1</sup> ABLIKIM	15B BES3	$e^+ e^- \rightarrow$ hadrons
-----------------------	------	----------------------	----------	-------------------------------

0.635 ± 0.003 ± 0.017	69k	<sup>1</sup> AUBERT,BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons
-----------------------	-----	------------------------	----------	------------------------------------

0.596 ± 0.035 ± 0.028	858	<sup>2</sup> ALBRECHT	95F ARG	$e^+ e^- \rightarrow$ hadrons
-----------------------	-----	-----------------------	---------	-------------------------------

0.636 ± 0.023 ± 0.033	1097	<sup>2</sup> BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons
-----------------------	------	---------------------	---------	-------------------------------

<sup>1</sup>Derived from the ratio  $\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$  assuming that the branching fractions of  $D^{*0} \rightarrow D^0 \pi^0$  and  $D^{*0} \rightarrow D^0 \gamma$  decays sum to 100%.

<sup>2</sup>The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

NODE=M061R2;LINKAGE=AU

NODE=M061R2;LINKAGE=A

#### $\Gamma(D^0 \gamma) / \Gamma_{\text{total}}$

 $\Gamma_2 / \Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.353 ± 0.009 OUR FIT**

**0.381 ± 0.029 OUR AVERAGE**

0.404 ± 0.035 ± 0.028	456	<sup>1</sup> ALBRECHT	95F ARG	$e^+ e^- \rightarrow$ hadrons
-----------------------	-----	-----------------------	---------	-------------------------------

0.364 ± 0.023 ± 0.033	621	<sup>1</sup> BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons
-----------------------	-----	---------------------	---------	-------------------------------

0.37 ± 0.08 ± 0.08		ADLER	88D MRK3	$e^+ e^-$
--------------------	--	-------	----------	-----------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.345 ± 0.008 ± 0.005	1.8k	<sup>2</sup> ABLIKIM	15B BES3	$e^+ e^- \rightarrow$ hadrons
-----------------------	------	----------------------	----------	-------------------------------

0.365 ± 0.003 ± 0.017	68k	<sup>2</sup> AUBERT,BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons
-----------------------	-----	------------------------	----------	------------------------------------

0.47 ± 0.23		LOW	87 HRS	29 GeV $e^+ e^-$
-------------	--	-----	--------	------------------

0.53 ± 0.13		BARTEL	85G JADE	$e^+ e^-$ , hadrons
-------------	--	--------	----------	---------------------

0.47 ± 0.12		COLES	82 MRK2	$e^+ e^-$
-------------	--	-------	---------	-----------

0.45 ± 0.15		GOLDHABER	77 MRK1	$e^+ e^-$
-------------	--	-----------	---------	-----------

<sup>1</sup>The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

<sup>2</sup>Derived from the ratio  $\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$  assuming that the branching fractions of  $D^{*0} \rightarrow D^0 \pi^0$  and  $D^{*0} \rightarrow D^0 \gamma$  decays sum to 100%.

NODE=M061R1  
NODE=M061R1

NODE=M061R;LINKAGE=A

NODE=M061R;LINKAGE=AU

**$D^*(2007)^0$  REFERENCES**

AAIJ	22N	PR D105 072005	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	21BD	PR D104 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15B	PR D91 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
TOMARADZE	15	PR D91 011102	A. Tomaradze <i>et al.</i>	(NWES)
AUBERT,BE	05G	PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BORTOLETTO	92B	PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)
ABACHI	88B	PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)
ADLER	88D	PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)
LOW	87	PL B183 232	E.H. Low <i>et al.</i>	(HRS Collab.)
BARTEL	85G	PL 161B 197	W. Bartel <i>et al.</i>	(JADE Collab.)
COLES	82	PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)
SADROZINSKI	80	Madison Conf. 681	H.F.W. Sadrozinski <i>et al.</i>	(PRIN, CIT+)
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
NGUYEN	77	PRL 39 262	H.K. Nguyen <i>et al.</i>	(LBL, SLAC) J

NODE=M061

REFID=61737  
 REFID=61535  
 REFID=56375  
 REFID=57142  
 REFID=50942  
 REFID=44374  
 REFID=43116  
 REFID=43170  
 REFID=40584  
 REFID=40579  
 REFID=40017  
 REFID=22880  
 REFID=22866  
 REFID=22877  
 REFID=11434  
 REFID=11543

 **$D^*(2010)^\pm$** 

$$I(J^P) = \frac{1}{2}(1^-)$$

$I, J, P$  need confirmation.

NODE=M062

 **$D^*(2010)^\pm$  MASS**

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ ,  
 and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M062M

NODE=M062M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

**2010.26 ± 0.05 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

2008 ± 3	<sup>1</sup> GOLDHABER 77	MRK1 ±	$e^+ e^-$
2008.6 ± 1.0	<sup>2</sup> PERUZZI 77	LGW ±	$e^+ e^-$

<sup>1</sup> From simultaneous fit to  $D^*(2010)^+, D^*(2007)^0, D^+,$  and  $D^0$ ; not independent of FELDMAN 77B mass difference below.

<sup>2</sup> PERUZZI 77 mass not independent of FELDMAN 77B mass difference below and PERUZZI 77  $D^0$  mass value.

NODE=M062M

NODE=M062M;LINKAGE=G

NODE=M062M;LINKAGE=P

 **$m_{D^*(2010)^+} - m_{D^+}$** 

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ ,  
 and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M062MD

NODE=M062MD

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**140.603 ± 0.015 OUR FIT****140.602 ± 0.014 OUR AVERAGE**

140.6010 ± 0.0068 ± 0.0129	151k	LEES	17F BABR	$e^+ e^- \rightarrow$ hadrons
140.64 ± 0.08 ± 0.06	620	BORTOLETTO92B	CLE2	$e^+ e^- \rightarrow$ hadrons

NODE=M062MD

 **$m_{D^*(2010)^+} - m_{D^0}$** 

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ ,  
 and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M062DM

NODE=M062DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**145.4258 ± 0.0017 OUR FIT****145.4258 ± 0.0020 OUR AVERAGE**

Error includes scale factor of 1.2.

145.4259 ± 0.0004 ± 0.0017	312.8k	LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K \pi, K 3 \pi) \pi^\pm$
145.412 ± 0.002 ± 0.012		ANASTASSOV 02	CLE2	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K \pi) \pi^\pm$
145.54 ± 0.08	611	<sup>3</sup> ADINOLFI 99	BEAT	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.45 ± 0.02		<sup>3</sup> BREITWEG 99	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K \pi) \pi^\pm$
145.42 ± 0.05		<sup>3</sup> BREITWEG 99	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- 3 \pi) \pi^\pm$
145.5 ± 0.15	103	<sup>4</sup> ADLOFF 97B	H1	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.44 ± 0.08	152	<sup>4</sup> BREITWEG 97	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm, D^0 \rightarrow K^- 3 \pi$

NODE=M062DM

OCCUR=3

OCCUR=2

145.42 ±0.11	199	<sup>4</sup> BREITWEG	97	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm,$ $D^0 \rightarrow K^- \pi^\pm$	OCCUR=2
145.4 ±0.2	48	<sup>4</sup> DERRICK	95	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.39 ±0.06 ±0.03		BARLAG	92B	ACCM	$\pi^-$ 230 GeV	
145.5 ±0.2	115	<sup>4</sup> ALEXANDER	91B	OPAL	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.30 ±0.06		<sup>4</sup> DECAMP	91J	ALEP	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.40 ±0.05 ±0.10		ABACHI	88B	HRS	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.46 ±0.07 ±0.03		ALBRECHT	85F	ARG	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.5 ±0.3	28	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	OCCUR=2
145.5 ±0.3	60	FITCH	81	SPEC	$\pi^- A$	
145.3 ±0.5	30	FELDMAN	77B	MRK1	$D^{*+} \rightarrow D^0 \pi^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

145.4256±0.0006±0.0017	138.5k	LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- \pi^+) \pi^\pm$	OCCUR=2
145.4266±0.0005±0.0019	174.3k	LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- 2\pi^+ \pi^-) \pi^\pm$	OCCUR=2
145.44 ±0.09	122	<sup>4</sup> BREITWEG	97B	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm,$ $D^0 \rightarrow K^- \pi^+$	
145.8 ±1.5	16	AHLEN	83	HRS	$D^{*+} \rightarrow D^0 \pi^+$	
145.1 ±1.8	12	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	OCCUR=3
145.1 ±0.5	14	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.5 ±0.5	14	YELTON	82	MRK2	$29 e^+ e^- \rightarrow$ $K^- \pi^+$	
~ 145.5		AVERY	80	SPEC	$\gamma A$	
145.2 ±0.6	2	BLIETSCHAU	79	BEBC	$\nu p$	

<sup>3</sup>Statistical errors only.

<sup>4</sup>Systematic error not evaluated.

NODE=M062DM;LINKAGE=AV  
NODE=M062DM;LINKAGE=A

### $m_{D^*(2010)^+} - m_{D^*(2007)^0}$

NODE=M062EM

VALUE (MeV) DOCUMENT ID TECN COMMENT

NODE=M062EM

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6±1.8		<sup>5</sup> PERUZZI	77	LGW	$e^+ e^-$	
---------	--	----------------------	----	-----	-----------	--

<sup>5</sup>Not independent of FELDMAN 77B mass difference above, PERUZZI 77  $D^0$  mass, and GOLDHABER 77  $D^*(2007)^0$  mass.

NODE=M062EM;LINKAGE=P

### $D^*(2010)^\pm$ WIDTH

NODE=M062W

VALUE (keV) CL% EVTS DOCUMENT ID TECN COMMENT

NODE=M062W

#### 83.4±1.8 OUR AVERAGE

83.3±1.2± 1.4	312.8k	<sup>6</sup> LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K \pi, K 3\pi) \pi^\pm$	OCCUR=3
96 ±4 ±22		<sup>6</sup> ANASTASSOV	02	CLE2	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K \pi) \pi^\pm$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
83.4±1.7± 1.5	138.5k	<sup>6</sup> LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- \pi^+) \pi^\pm$	
83.2±1.5± 2.6	174.3k	<sup>6</sup> LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow$ $(K^- 2\pi^+ \pi^-) \pi^\pm$	OCCUR=2
<131	90 110	BARLAG	92B	ACCM	$\pi^-$ 230 GeV	

<sup>6</sup>Ignoring the electromagnetic contribution from  $D^{*\pm} \rightarrow D^\pm \gamma$ .

NODE=M062W;LINKAGE=LE

### $D^*(2010)^\pm$ DECAY MODES

NODE=M062225;NODE=M062

$D^*(2010)^-$  modes are charge conjugates of the modes below.

NODE=M062

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^0 \pi^+$	(67.7±0.5) %
$\Gamma_2$ $D^+ \pi^0$	(30.7±0.5) %
$\Gamma_3$ $D^+ \gamma$	( 1.6±0.4) %

DESIG=1

DESIG=3

DESIG=2

## CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 6 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 0.3$  for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-62	
$x_3$	-43	-44
	$x_1$	$x_2$

### D\*(2010)<sup>+</sup> BRANCHING RATIOS

NODE=M062230

#### $\Gamma(D^0 \pi^+) / \Gamma_{\text{total}}$ $\Gamma_1 / \Gamma$

NODE=M062R1  
NODE=M062R1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.677 ± 0.005 OUR FIT****0.677 ± 0.006 OUR AVERAGE**

0.6759 ± 0.0029 ± 0.0064	7,8,9	BARTELT	98	CLE2	e <sup>+</sup> e <sup>-</sup>
0.688 ± 0.024 ± 0.013		ALBRECHT	95F	ARG	e <sup>+</sup> e <sup>-</sup> → hadrons
0.681 ± 0.010 ± 0.013	7	BUTLER	92	CLE2	e <sup>+</sup> e <sup>-</sup> → hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.57 ± 0.04 ± 0.04		ADLER	88D	MRK3	e <sup>+</sup> e <sup>-</sup>
0.44 ± 0.10		COLES	82	MRK2	e <sup>+</sup> e <sup>-</sup>
0.6 ± 0.15	9	GOLDHABER	77	MRK1	e <sup>+</sup> e <sup>-</sup>

#### $\Gamma(D^+ \pi^0) / \Gamma_{\text{total}}$ $\Gamma_2 / \Gamma$

NODE=M062R3  
NODE=M062R3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.307 ± 0.005 OUR FIT****0.3073 ± 0.0013 ± 0.0062**

0.312 ± 0.011 ± 0.008	1404	ALBRECHT	95F	ARG	e <sup>+</sup> e <sup>-</sup> → hadrons
0.308 ± 0.004 ± 0.008	410	7 BUTLER	92	CLE2	e <sup>+</sup> e <sup>-</sup> → hadrons
0.26 ± 0.02 ± 0.02		ADLER	88D	MRK3	e <sup>+</sup> e <sup>-</sup>
0.34 ± 0.07		COLES	82	MRK2	e <sup>+</sup> e <sup>-</sup>

#### $\Gamma(D^+ \gamma) / \Gamma_{\text{total}}$ $\Gamma_3 / \Gamma$

NODE=M062R2  
NODE=M062R2

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

**0.016 ± 0.004 OUR FIT****0.016 ± 0.005 OUR AVERAGE**

0.0168 ± 0.0042 ± 0.0029		7,8	BARTELT	98	CLE2 e <sup>+</sup> e <sup>-</sup>
0.011 ± 0.014 ± 0.016	12	7	BUTLER	92	CLE2 e <sup>+</sup> e <sup>-</sup> → hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 0.052	90		ALBRECHT	95F	ARG e <sup>+</sup> e <sup>-</sup> → hadrons
0.17 ± 0.05 ± 0.05			ADLER	88D	MRK3 e <sup>+</sup> e <sup>-</sup>
0.22 ± 0.12		10	COLES	82	MRK2 e <sup>+</sup> e <sup>-</sup>

<sup>7</sup> The branching ratios are not independent, they have been constrained by the authors to sum to 100%.

<sup>8</sup> Systematic error includes theoretical error on the prediction of the ratio of hadronic modes.

<sup>9</sup> Assuming that isospin is conserved in the decay.

<sup>10</sup> Not independent of  $\Gamma(D^0 \pi^+) / \Gamma_{\text{total}}$  and  $\Gamma(D^+ \pi^0) / \Gamma_{\text{total}}$  measurement.

NODE=M062R;LINKAGE=A

NODE=M062R;LINKAGE=B

NODE=M062R;LINKAGE=G

NODE=M062R;LINKAGE=C

**$D^*(2010)^\pm$  REFERENCES**

LEES	17F	PRL 119 202003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58277
LEES	13X	PRL 111 111801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55564
	Also	PR D88 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55547
	Also	PR D88 079902 (errata.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55695
ANASTASSOV	02	PR D65 032003	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=48550
ADINOLFI	99	NP B547 3	M. Adinolfi <i>et al.</i>	(Beatrice Collab.)	REFID=46925
BREITWEG	99	EPJ C6 67	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=46604
BARTELT	98	PRL 80 3919	J. Bartelt <i>et al.</i>	(CLEO Collab.)	REFID=46349
ADLOFF	97B	ZPHY C72 593	C. Adloff <i>et al.</i>	(H1 Collab.)	REFID=45421
BREITWEG	97	PL B401 192	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45520
BREITWEG	97B	PL B407 402	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45699
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44374
DERRICK	95	PL B349 225	M. Derrick <i>et al.</i>	(ZEUS Collab.)	REFID=44373
BARLAG	92B	PL B278 480	S. Barlag <i>et al.</i>	(ACCMOR Collab.)	REFID=42174
BORTOLETTO	92B	PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)	REFID=43116
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43170
ALEXANDER	91B	PL B262 341	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=41553
DECAMP	91J	PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)	REFID=41614
ABACHI	88B	PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)	REFID=40584
ADLER	88D	PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40579
ALBRECHT	85F	PL 150B 235	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=11527
AHLEN	83	PRL 51 1147	S.P. Ahlen <i>et al.</i>	(ANL, IND, LBL+)	REFID=22868
BAILEY	83	PL 132B 230	R. Bailey <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)	REFID=22870
COLES	82	PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)	REFID=22866
YELTON	82	PRL 49 430	J.M. Yelton <i>et al.</i>	(SLAC, LBL, UCB+)	REFID=22867
FITCH	81	PRL 46 761	V.L. Fitch <i>et al.</i>	(PRIN, SACL, TORI+)	REFID=22863
AVERY	80	PRL 44 1309	P. Avery <i>et al.</i>	(ILL, FNAL, COLU)	REFID=11498
BLIETSCHAU	79	PL 86B 108	J. Blietschau <i>et al.</i>	(AACH3, BONN, CERN+)	REFID=22861
FELDMAN	77B	PRL 38 1313	G.J. Feldman <i>et al.</i>	(Mark I Collab.)	REFID=22858
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)	REFID=11434
PERUZZI	77	PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)	REFID=11435

NODE=M062

NODE=M252

**$D_0^*(2300)$**

$$I(J^P) = \frac{1}{2}(0^+)$$

was  $D_0^*(2400)$

There is a strong evidence that recent data on  $B \rightarrow D\pi\pi$  (AAIJ 15Y, AAIJ 16AH) and  $B \rightarrow D\pi K$  (AAIJ 14BH, AAIJ 15V, AAIJ 15X) call for two poles in the scalar  $I = 1/2 \pi D$  amplitude in this mass range. The data are consistent with a lower pole at  $(2105^{+6}_{-8}) - i(102^{+10}_{-11})$  MeV and a higher pole at  $(2451^{+35}_{-26}) - i(134^{+7}_{-8})$  MeV (DU 18A, DU 19, DU 21). For details see review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M252

**$D_0^*(2300)$  MASS**

NODE=M252M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>2343±10 OUR AVERAGE</b>	Error	includes scale factor of 1.5. See the ideogram below.			
2360±15±30		1 AAIJ	15X LHCB	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2349± 6± 4		2 AAIJ	15Y LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2297± 8±20	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
2308±17±32		ABE	04D BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2354± 7±11		3 AAIJ	15Y LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2403±14±35	18.8k	4 LINK	04A FOCS	+	$\gamma A$
2407±21±35	9.8k	4 LINK	04A FOCS	0	$\gamma A$

NODE=M252M

OCCUR=2

OCCUR=2

<sup>1</sup> From the Dalitz plot analysis including various  $K^*$  and  $D^{**}$  mesons as well as broad structures in the  $K\pi$  S-wave and the  $D\pi$  S- and P-waves.

<sup>2</sup> Modeling the  $\pi^+ \pi^-$  S-wave with the Isobar formalism.

<sup>3</sup> Modeling the  $\pi^+ \pi^-$  S-wave with the K-matrix formalism.


<sup>4</sup> Possibly the feed-down from another state.

NODE=M252M;LINKAGE=A

NODE=M252M;LINKAGE=B

NODE=M252M;LINKAGE=C

NODE=M252M;LINKAGE=D



**$D_0^*(2300)$  WIDTH**

NODE=M252W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>229±16 OUR AVERAGE</b>					
255±26±51		<sup>1</sup> AAIJ	15X	LHCb	+ $B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
217±13±13		<sup>2</sup> AAIJ	15Y	LHCb	+ $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
273±12±48	3.4k	AUBERT	09AB	BABR	0 $B^- \rightarrow D^+ \pi^- \pi^-$
276±21±63		ABE	04D	BELL	0 $B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
230±15±21		<sup>3</sup> AAIJ	15Y	LHCb	+ $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
283±24±34	18.8k	<sup>4</sup> LINK	04A	FOCS	+ $\gamma A$
240±55±59	9.8k	<sup>4</sup> LINK	04A	FOCS	0 $\gamma A$

NODE=M252W

OCCUR=2

OCCUR=2

<sup>1</sup> From the Dalitz plot analysis including various  $K^*$  and  $D^{**}$  mesons as well as broad structures in the  $K\pi$  S-wave and the  $D\pi$  S- and P-waves.

<sup>2</sup> Modeling the  $\pi^+ \pi^-$  S-wave with the Isobar formalism.

<sup>3</sup> Modeling the  $\pi^+ \pi^-$  S-wave with the K-matrix formalism.

<sup>4</sup> Possibly the feed-down from another state.

NODE=M252W;LINKAGE=A

NODE=M252W;LINKAGE=B

NODE=M252W;LINKAGE=C

NODE=M252W;LINKAGE=D

**$D_0^*(2300)$  DECAY MODES**

NODE=M252215;NODE=M252

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D\pi^\pm$	seen

DESIG=1

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_1/\Gamma$
seen		AAIJ	15X	LHCb	+ $D^*(2300)^+ \rightarrow D^0 \pi^+$	
seen		AAIJ	15Y	LHCb	+ $D^*(2300)^+ \rightarrow D^0 \pi^+$	
seen	3.4k	AUBERT	09AB	BABR	0 $D^*(2300)^0 \rightarrow D^+ \pi^-$	
<b>seen</b>		ABE	04D	BELL	0 $D^*(2300)^0 \rightarrow D^+ \pi^-$	
<b>seen</b>	18.8k	LINK	04A	FOCS	+ $D^*(2300)^+ \rightarrow D^0 \pi^+$	

NODE=M252R01

NODE=M252R01

**$D_0^*(2300)$  REFERENCES**

NODE=M252

DU	21	PRL 126 192001	M.-L. Du <i>et al.</i>		REFID=61096
DU	19	PR D99 114002	M.-L. Du, F.-K. Guo, U.-G. Meissner		REFID=59893
DU	18A	PR D98 094018	M.-L. Du <i>et al.</i>		REFID=61424
AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57518
AAIJ	15V	PR D91 092002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56575
Also		PR D93 119901 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57289
AAIJ	15X	PR D92 012012	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56588
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56609
AAIJ	14BH	PR D90 072003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56208
AUBERT	09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52941
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50011
LINK	04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)	REFID=49775



$D_1(2420)$ 

$$I(J^P) = \frac{1}{2}(1^+)$$

NODE=M253

 **$D_1(2420)$  MASS**

NODE=M253M

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M253M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>2422.1±0.6 OUR FIT</b>		Error includes scale factor of 1.7.			
<b>2422.1±0.8 OUR AVERAGE</b>		Error includes scale factor of 2.1. See the ideogram below.			
2424.8±0.1±0.7	79k	<sup>1</sup> AAIJ	20D	LHCB	0 $B^- \rightarrow D^{*+} \pi^- \pi^-$
2427.2±1.0±1.2	4207	ABLIKIM	20P	BES3	+ $e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$
2419.6±0.1±0.7	210k	AAIJ	13CC	LHCB	0 $p p \rightarrow D^{*+} \pi^- X$
2423.1±1.5 <sup>+0.4</sup> <sub>-1.0</sub>	2.7k	<sup>2</sup> ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2421.9±4.7 <sup>+3.4</sup> <sub>-1.2</sub>	759	<sup>3</sup> ABRAMOWICZ13	ZEUS	+	$e^\pm p \rightarrow D^{(*)0} \pi^+ X$
2420.1±0.1±0.8	103k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2426 ±3 ±1	151	ABE	05A	BELL	0 $B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
2421 ±2 ±1	124	ABE	05A	BELL	+ $\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
2421.4±1.5±0.9		<sup>4</sup> ABE	04D	BELL	0 $B^- \rightarrow D^{*+} \pi^- \pi^-$
2421 <sup>+1</sup> <sub>-2</sub> ±2	286	AVERY	94C	CLE2	0 $e^+ e^- \rightarrow D^{*+} \pi^- X$
2425 ±2 ±2	146	BERGFELD	94B	CLE2	+ $e^+ e^- \rightarrow D^{*0} \pi^+ X$
2422 ±2 ±2	51	FRABETTI	94B	E687	0 $\gamma Be \rightarrow D^{*+} \pi^- X$
2428 ±3 ±2	279	AVERY	90	CLEO	0 $e^+ e^- \rightarrow D^{*+} \pi^- X$
2414 ±2 ±5	171	ALBRECHT	89H	ARG	0 $e^+ e^- \rightarrow D^{*+} \pi^- X$
2428 ±8 ±5	171	ANJOS	89C	TPS	0 $\gamma N \rightarrow D^{*+} \pi^- X$
2443 ±7 ±5	190	ANJOS	89C	TPS	+ $\gamma N \rightarrow D^0 \pi^+ X^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2420.5±2.1±0.9	3.1k	<sup>5</sup> CHEKANOV	09	ZEUS	0 $e^\pm p \rightarrow D^{*+} \pi^- X$
2421.7±0.7±0.6	7.5k	ABULENCIA	06A	CDF	0 $1900 p \bar{p} \rightarrow D^{*+} \pi^- X$
2425 ±3	235	<sup>6</sup> ABREU	98M	DLPH	0 $e^+ e^-$

NODE=M253M

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M253M;LINKAGE=B

NODE=M253M;LINKAGE=AR


NODE=M253M;LINKAGE=BA

NODE=M253M;LINKAGE=AB

NODE=M253M;LINKAGE=CH

NODE=M253M;LINKAGE=K

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.<sup>2</sup> From the combined fit of the  $M(D^+ \pi^-)$  and  $M(D^{*+} \pi^-)$  distributions. and  $A_{D_2}$  fixed to the theoretical prediction of  $-1$ .<sup>3</sup> From the fit of the  $M(D^0 \pi^+)$  distribution. The widths of the  $D_1^+$  and  $D_2^{*+}$  are fixed to 25 MeV and 37 MeV, and  $A_{D_1}$  and  $A_{D_2}$  are fixed to the theoretical predictions of 3 and  $-1$ , respectively.<sup>4</sup> Fit includes the contribution from  $D_1^*(2430)^0$ .<sup>5</sup> Calculated using the mass difference  $m(D_1^0) - m(D^{*+})_{PDG}$  reported below and  $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$  MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of 0.9 MeV.<sup>6</sup> No systematic error given.



$D_1(2420)$  mass (MeV)

$m_{D_1(2420)^0} - m_{D^{*+}}$

NODE=M253DM

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$  and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M253DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>411.8±0.6 OUR FIT</b>		Error includes scale factor of 1.7.		
<b>411.5±0.8 OUR AVERAGE</b>				
410.2±2.1±0.9	3.1k	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
411.7±0.7±0.4	7.5k	ABULENCIA 06A	CDF	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$

NODE=M253DM

$m_{D_1(2420)^\pm} - m_{D_1(2420)^0}$

NODE=M253DMC

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>4_{-3}^{+2} \pm 3</math></b>	BERGFELD 94B	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M253DMC

$D_1(2420)$  WIDTH

NODE=M253W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>31.3± 1.9 OUR AVERAGE</b>		Error includes scale factor of 2.8. See the ideogram below.			
33.6± 0.3± 2.7	79k	<sup>1</sup> AAIJ 20D	LHCB	0	$B^- \rightarrow D^{*+} \pi^- \pi^-$
23.2± 2.3± 2.3	4207	ABLIKIM 20P	BES3	+	$e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$
35.2± 0.4± 0.9	210k	AAIJ 13CC	LHCB	0	$p p \rightarrow D^{*+} \pi^- X$
38.8± 5.0 <sup>+1.9</sup> <sub>-5.4</sub>	2.7k	<sup>2</sup> ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
31.4± 0.5± 1.3	103k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
20.0± 1.7± 1.3	7.5k	ABULENCIA 06A	CDF	0	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$
24 ± 7 ± 8	151	ABE 05A	BELL	0	$B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
21 ± 5 ± 8	124	ABE 05A	BELL	+	$\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
23.7± 2.7± 4.0		<sup>3</sup> ABE 04D	BELL	0	$B^- \rightarrow D^{*+} \pi^- \pi^-$
20 <sup>+6</sup> <sub>-5</sub> ± 3	286	AVERY 94C	CLE2	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$

NODE=M253W


OCCUR=2

OCCUR=2

26	$\begin{matrix} +8 \\ -7 \end{matrix}$	$\pm 4$	146	BERGFELD	94B	CLE2	+	$e^+e^- \rightarrow D^{*0}\pi^+X$	OCCUR=2
15	$\pm 8$	$\pm 4$	51	FRABETTI	94B	E687	0	$\gamma Be \rightarrow D^{*+}\pi^-X$	
23	$\begin{matrix} +8 \\ -6 \end{matrix}$	$\begin{matrix} +10 \\ -3 \end{matrix}$	279	AVERY	90	CLEO	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	
13	$\pm 6$	$\begin{matrix} +10 \\ -5 \end{matrix}$	171	ALBRECHT	89H	ARG	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	
41	$\pm 19$	$\pm 8$	190	ANJOS	89C	TPS	+	$\gamma N \rightarrow D^0\pi^+X^0$	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●									
53.2	$\pm 7.2$	$\begin{matrix} +3.3 \\ -4.9 \end{matrix}$	3.1k	CHEKANOV	09	ZEUS	0	$e^\pm p \rightarrow D^{*+}\pi^-X$	
58	$\pm 14$	$\pm 10$	171	ANJOS	89C	TPS	0	$\gamma N \rightarrow D^{*+}\pi^-X$	

- <sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+}\pi^-\pi^-$  decay.
- <sup>2</sup> From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  $A_{D_2}$  fixed to the theoretical prediction of  $-1$ .
- <sup>3</sup> Fit includes the contribution from  $D_1^*(2430)^0$ .

NODE=M253W;LINKAGE=B  
 NODE=M253W;LINKAGE=AR  
 NODE=M253W;LINKAGE=AB



### $D_1(2420)$ DECAY MODES

NODE=M253215;NODE=M253  
 NODE=M253

$\bar{D}_1(2420)$  modes are charge conjugates of modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^*(2007)^0\pi$	seen
$\Gamma_2$ $D\pi^+\pi^-$	
$\Gamma_3$ $D\rho^0$	
$\Gamma_4$ $Df_0(500)$	
$\Gamma_5$ $D_0^*(2300)^0\pi$	
$\Gamma_6$ $D^0\pi$	
$\Gamma_7$ $D^*\pi^+\pi^-$	

DESIG=1  
 DESIG=3  
 DESIG=4  
 DESIG=5  
 DESIG=6  
 DESIG=2  
 DESIG=7

### $D_1(2420)$ BRANCHING RATIOS

NODE=M253220

$\Gamma(D^*(2007)^0\pi)/\Gamma_{total}$	$\Gamma_1/\Gamma$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ACKERSTAFF	97W	OPAL	0 $e^+e^- \rightarrow D^{*+}\pi^-X$
seen	AVERY	90	CLEO	0 $e^+e^- \rightarrow D^{*+}\pi^-X$
seen	ALBRECHT	89H	ARG	0 $e^+e^- \rightarrow D^*\pi^-X$
seen	ANJOS	89C	TPS	0 $\gamma N \rightarrow D^{*+}\pi^-X$
seen	ANJOS	89C	TPS	+ $\gamma N \rightarrow D^0\pi^+X^0$

NODE=M253R01  
 NODE=M253R01

OCCUR=2

$\Gamma(D^0\pi)/\Gamma(D^*(2007)^0\pi)$  $\Gamma_6/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.18	90	BERGFELD 94B	CLE2	+	$e^+e^- \rightarrow \text{hadrons}$
<0.24	90	AVERY 90	CLEO	0	$e^+e^- \rightarrow D^+\pi^-X$

NODE=M253R02  
 NODE=M253R02  
 OCCUR=2

 **$D_1(2420)$  POLARIZATION AMPLITUDE  $A_{D_1}$** 

NODE=M253PAH

A polarization amplitude  $A_{D_1}$  is a parameter that depends on the initial polarization of the  $D_1$  and is sensitive to a possible  $S$ -wave contribution to its decay. For  $D_1$  decays the helicity angle,  $\theta_h$ , distribution varies like  $1 + A_{D_1} \cos^2 \theta_h$ , where  $\theta_h$  is the angle in the  $D^*$  rest frame between the two pions emitted by the  $D_1 \rightarrow D^*\pi$  and the  $D^* \rightarrow D\pi$ .

NODE=M253PAH

Unpolarized  $D_1$  decaying purely via  $D$ -wave is predicted to give  $A_{D_1} = 3$ .

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>5.73±0.25 OUR AVERAGE</b>						
7.8	+6.7 -2.7	1 ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$	
5.72±0.25	103k	DEL-AMO-SA...10P	BABR	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	
5.9	+3.0 -1.7	CHEKANOV 09	ZEUS	0	$e^\pm p \rightarrow D^{*+}\pi^-X$	
3.30±0.48	210k	2 AAIJ	13CC	LHCB	0	$pp \rightarrow D^{*+}\pi^-X$
3.8 ±0.6 ±0.8		3 AUBERT	09Y	BABR	0	$B^+ \rightarrow D_1^0 \ell^+ \nu_\ell$
3.8 ±0.6 ±0.8		3 AUBERT	09Y	BABR	+	$B^0 \rightarrow D_1^- \ell^+ \nu_\ell$
2.74	+1.40 -0.93	4 AVERY	94C	CLE2	0	$e^+e^- \rightarrow D^{*+}\pi^-X$

NODE=M253PAH

••• We do not use the following data for averages, fits, limits, etc. •••

OCCUR=2

<sup>1</sup> From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  $A_{D_2}$  fixed to the theoretical prediction of  $-1$ . A pure  $D$ -wave not excluded although some  $\bar{S}$ -wave mixing possible.

NODE=M253PAH;LINKAGE=AR

<sup>2</sup> Systematic uncertainty not estimated. Resonance parameters fixed.

NODE=M253PAH;LINKAGE=A

<sup>3</sup> Assuming  $\Gamma(\Upsilon(4S) \rightarrow B^+B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.065 \pm 0.026$  and equal partial widths and helicity angle distributions for charged and neutral  $D_1$  mesons.

NODE=M253PAH;LINKAGE=AU

<sup>4</sup> Systematic uncertainties not estimated.

NODE=M253PAH;LINKAGE=AV

 **$D_1(2420)$  REFERENCES**

NODE=M253

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60253
ABLIKIM	20P	PL B804 135395	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60395
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
ABRAMOWICZ	13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54743
DEL-AMO-SA...	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52929
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)	REFID=51054
ABE	05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50755
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50011
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46315
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)	REFID=44096
BERGFELD	94B	PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)	REFID=44099
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=41001
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)	REFID=40737

$D_1(2430)^0$ 

$$I(J^P) = \frac{1}{2}(1^+)$$

 $J^P = 1^+$  determined by AAIJ 20D.

NODE=M180

NODE=M180

NODE=M180M

NODE=M180M

 $D_1(2430)^0$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2412 ± 9 OUR AVERAGE</b>				
2411 ± 3 ± 9	79k	<sup>1</sup> AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2427 ± 26 ± 25		ABE	04D BELLE	$B^- \rightarrow D^{*+} \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2477 ± 28		<sup>2</sup> AUBERT	06L BABR	$\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.<sup>2</sup> Systematic errors not estimated.NODE=M180M;LINKAGE=A  
NODE=M180M;LINKAGE=AU $D_1(2430)^0$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>314 ± 29 OUR AVERAGE</b>				
309 ± 9 ± 28	79k	<sup>1</sup> AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
384 <sup>+107</sup> <sub>-75</sub> ± 74		ABE	04D BELLE	$B^- \rightarrow D^{*+} \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
266 ± 97		<sup>2</sup> AUBERT	06L BABR	$\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.<sup>2</sup> Systematic errors not estimated.

NODE=M180W

NODE=M180W

NODE=M180W;LINKAGE=A  
NODE=M180W;LINKAGE=AU $D_1(2430)^0$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^*(2010)^+ \pi^-$	seen

NODE=M180215;NODE=M180

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D_1(2430)^0$  REFERENCES

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AUBERT	06L	PR D74 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M180

REFID=60253  
REFID=51140  
REFID=50011

$D_2^*(2460)$ 

$$I(J^P) = \frac{1}{2}(2^+)$$

NODE=M254

 **$D_2^*(2460)$  MASS**

NODE=M254M

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_S^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_S^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M254M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M254M

**2461.1<sup>+0.7</sup><sub>-0.8</sub> OUR FIT** Error includes scale factor of 6.2.

**2461.1±0.7 OUR AVERAGE** Error includes scale factor of 5.2. See the ideogram below.

2463.7±0.4±0.7	28k	<sup>1</sup> AAIJ	16AH	LHCB	0	$B^- \rightarrow D^+ \pi^- \pi^-$	OCCUR=2
2464.0±1.4±0.5	2k	<sup>2</sup> AAIJ	15V	LHCB	0	$B^- \rightarrow D^+ K^- \pi^-$	
2465.6±1.8±1.3		<sup>3</sup> AAIJ	15X	LHCB	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$	
2468.6±0.6±0.3		<sup>4</sup> AAIJ	15Y	LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$	
2460.4±0.4±1.2	82k	AAIJ	13CC	LHCB	0	$pp \rightarrow D^{*+} \pi^- X$	
2460.4±0.1±0.1	675k	AAIJ	13CC	LHCB	0	$pp \rightarrow D^+ \pi^- X$	OCCUR=2
2463.1±0.2±0.6	342k	AAIJ	13CC	LHCB	+	$pp \rightarrow D^0 \pi^+ X$	OCCUR=3
2462.5±2.4 <sup>+1.3</sup> <sub>-1.1</sub>	2.3k	<sup>5</sup> ABRAMOWICZ13	ZEUS		0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$	
2460.6±4.4 <sup>+3.6</sup> <sub>-0.8</sub>	1371	<sup>6</sup> ABRAMOWICZ13	ZEUS	+		$e^\pm p \rightarrow D^{(*)0} \pi^+ X$	OCCUR=2
2462.2±0.1±0.8	243k	DEL-AMO-SA..10P	BABR	0		$e^+ e^- \rightarrow D^+ \pi^- X$	
2465.4±0.2±1.1	111k	<sup>7</sup> DEL-AMO-SA..10P	BABR	+		$e^+ e^- \rightarrow D^0 \pi^+ X$	OCCUR=2
2460.4±1.2±2.2	3.4k	AUBERT	09AB	BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$	
2465.7±1.8 <sup>+1.4</sup> <sub>-4.8</sub>	2909	KUZMIN	07	BELL	+	$e^+ e^- \rightarrow \text{hadrons}$	
2461.6±2.1±3.3		<sup>8</sup> ABE	04D	BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$	
2464.5±1.1±1.9	5.8k	<sup>8</sup> LINK	04A	FOCS	0	$\gamma A$	
2465 ±3 ±3	486	AVERY	94C	CLE2	0	$e^+ e^- \rightarrow D^+ \pi^- X$	
2463 ±3 ±3	310	BERGFELD	94B	CLE2	+	$e^+ e^- \rightarrow D^0 \pi^+ X$	
2453 ±3 ±2	128	FRABETTI	94B	E687	0	$\gamma \text{Be} \rightarrow D^+ \pi^- X$	
2453 ±3 ±2	185	FRABETTI	94B	E687	+	$\gamma \text{Be} \rightarrow D^0 \pi^+ X$	OCCUR=2
2461 ±3 ±1	440	AVERY	90	CLEO	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
2455 ±3 ±5	337	ALBRECHT	89B	ARG	0	$e^+ e^- \rightarrow D^+ \pi^- X$	
2469 ±4 ±6		ALBRECHT	89F	ARG	+	$e^+ e^- \rightarrow D^0 \pi^+ X$	OCCUR=2
2459 ±3 ±2	153	ANJOS	89C	TPS	0	$\gamma N \rightarrow D^+ \pi^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
2468.1±0.6±0.5		<sup>9</sup> AAIJ	15Y	LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$	OCCUR=2
2469.1±3.7 <sup>+1.2</sup> <sub>-1.3</sub>	1.5k	<sup>10</sup> CHEKANOV	09	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$	
2463.3±0.6±0.8	20k	ABULENCIA	06A	CDF	0	$1900 p\bar{p} \rightarrow D^+ \pi^- X$	
2467.6±1.5±0.8	3.5k	<sup>11</sup> LINK	04A	FOCS	+	$\gamma A$	OCCUR=2
2461 ±6	126	<sup>12</sup> ABREU	98M	DLPH	0	$e^+ e^-$	
2466 ±7	1	ASRATYAN	95	BEBC	0	$53,40 \nu(\bar{\nu}) \rightarrow pX, dX$	

<sup>1</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

NODE=M254M;LINKAGE=B

<sup>2</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, nonresonant spin-0 and spin-1 components as well as the  $D_0^*(2400)^0$ ,  $D_2^*(2460)^0$  and  $D_1^*(2760)^0$  resonances.

NODE=M254M;LINKAGE=A

<sup>3</sup> From the Dalitz plot analysis including various  $K^*$  and  $D^{**}$  mesons as well as broad structures in the  $K\pi$   $S$ -wave and the  $D\pi$   $S$ - and  $P$ -waves.

NODE=M254M;LINKAGE=CA

<sup>4</sup> Modeling the  $\pi^+ \pi^-$   $S$ -wave with the Isobar formalism.

NODE=M254M;LINKAGE=BC

<sup>5</sup> From the combined fit of the  $M(D^+ \pi^-)$  and  $M(D^{*+} \pi^-)$  distributions. and  $A_{D_2}$  fixed to the theoretical prediction of  $-1$ .

NODE=M254M;LINKAGE=AR

<sup>6</sup> From the fit of the  $M(D^0 \pi^+)$  distribution. The widths of the  $D_1^+$  and  $D_2^{*+}$  are fixed to 25 MeV and 37 MeV, and  $A_{D_1}$  and  $A_{D_2}$  are fixed to the theoretical predictions of 3 and  $-1$ , respectively.

NODE=M254M;LINKAGE=AB

<sup>7</sup> At a fixed width of 50.5 MeV.

NODE=M254M;LINKAGE=DE

<sup>8</sup> Fit includes the contribution from  $D_0^*(2400)^0$ .

NODE=M254M;LINKAGE=LI

<sup>9</sup> Modeling the  $\pi^+ \pi^-$   $S$ -wave with the K-matrix formalism.

NODE=M254M;LINKAGE=CC

<sup>10</sup> Calculated using the mass difference  $m(D_2^{*0}) - m(D^{*+})_{PDG}$  reported below and  $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$  MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of  $^{+1.2}_{-1.3}$  MeV.


NODE=M254M;LINKAGE=CH

<sup>11</sup> Fit includes the contribution from  $D_0^*(2400)^\pm$ . Not independent of the corresponding mass difference measurement,  $(m_{D_2^*(2460)^\pm}) - (m_{D_2^*(2460)^0})$ .

NODE=M254M;LINKAGE=LC

<sup>12</sup> No systematic error given.

NODE=M254M;LINKAGE=K



$$m_{D_2^*(2460)^0} - m_{D^+}$$

NODE=M254DM

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M254DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>591.5<sup>+0.7</sup><sub>-0.8</sub> OUR FIT</b>				Error includes scale factor of 5.9.
<b>593.9±0.6±0.5</b>	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$

NODE=M254DM

$$m_{D_2^*(2460)^0} - m_{D^{*++}}$$

NODE=M254DM2

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M254DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>450.9<sup>+0.7</sup><sub>-0.8</sub> OUR FIT</b>				Error includes scale factor of 5.9.
<b>458.8±3.7<sup>+1.2</sup><sub>-1.3</sub></b>	1.5k	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$

NODE=M254DM2

$$m_{D_2^*(2460)^\pm} = m_{D_2^*(2460)^0}$$

NODE=M254DMC

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2.4±1.7 OUR AVERAGE</b>			
3.1±1.9±0.9	LINK	04A FOCS	$\gamma A$
- 2 ±4 ±4	BERGFELD	94B CLE2	$e^+e^- \rightarrow \text{hadrons}$
0 ±4	FRABETTI	94B E687	$\gamma Be \rightarrow D\pi X$
14 ±5 ±8	ALBRECHT	89F ARG	$e^+e^- \rightarrow D^0\pi^+X$

NODE=M254DMC

 **$D_2^*(2460)$  WIDTH**

NODE=M254W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>47.3± 0.8 OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.			
47.0± 0.8± 1.0	28k	<sup>1</sup> AAIJ	16AH LHCb	0	$B^- \rightarrow D^+\pi^-\pi^-$
43.8± 2.9± 1.8	2k	<sup>2</sup> AAIJ	15V LHCb	0	$B^- \rightarrow D^+K^-\pi^-$
46.0± 3.4± 3.2		<sup>3</sup> AAIJ	15X LHCb	+	$B^0 \rightarrow \bar{D}^0K^+\pi^-$
47.3± 1.5± 0.7		<sup>4</sup> AAIJ	15Y LHCb	+	$B^0 \rightarrow \bar{D}^0\pi^+\pi^-$
43.2± 1.2± 3.0	82k	AAIJ	13CC LHCb	0	$p\bar{p} \rightarrow D^{*+}\pi^-X$
45.6± 0.4± 1.1	675k	AAIJ	13CC LHCb	0	$p\bar{p} \rightarrow D^+\pi^-X$
48.6± 1.3± 1.9	342k	AAIJ	13CC LHCb	+	$p\bar{p} \rightarrow D^0\pi^+X$
46.6± 8.1 <sup>+</sup> <sub>-</sub> 5.9 <sub>3.8</sub>	2.3k	<sup>5</sup> ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$
50.5± 0.6± 0.7	243k	DEL-AMO-SA..10P	BABR	0	$e^+e^- \rightarrow D^+\pi^-X$
41.8± 2.5± 2.9	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+\pi^-\pi^-$
49.7± 3.8± 6.4	2909	KUZMIN	07 BELL	+	$e^+e^- \rightarrow \text{hadrons}$
49.2± 2.3± 1.3	20k	ABULENCIA	06A CDF	0	1900 $p\bar{p} \rightarrow D^+\pi^-X$
45.6± 4.4± 6.7		<sup>6</sup> ABE	04D BELL	0	$B^- \rightarrow D^+\pi^-\pi^-$
38.7± 5.3± 2.9	5.8k	<sup>6</sup> LINK	04A FOCS	0	$\gamma A$
34.1± 6.5± 4.2	3.5k	<sup>7</sup> LINK	04A FOCS	+	$\gamma A$
28 <sup>+</sup> <sub>-</sub> 8 <sub>7</sub> ± 6	486	AVERY	94C CLE2	0	$e^+e^- \rightarrow D^+\pi^-X$
27 <sup>+</sup> <sub>-</sub> 11 <sub>8</sub> ± 5	310	BERGFELD	94B CLE2	+	$e^+e^- \rightarrow D^0\pi^+X$
25 ±10 ± 5	128	FRABETTI	94B E687	0	$\gamma Be \rightarrow D^+\pi^-X$
23 ± 9 ± 5	185	FRABETTI	94B E687	+	$\gamma Be \rightarrow D^0\pi^+X$
20 <sup>+</sup> <sub>-</sub> 9 <sub>-12</sub> <sup>+</sup> <sub>-</sub> 9 <sub>-10</sub>	440	AVERY	90 CLEO	0	$e^+e^- \rightarrow D^{*+}\pi^-X$
15 <sup>+</sup> <sub>-</sub> 13 <sub>-10</sub> <sup>+</sup> <sub>-</sub> 5 <sub>-10</sub>	337	ALBRECHT	89B ARG	0	$e^+e^- \rightarrow D^+\pi^-X$
20 ±10 ± 5	153	ANJOS	89C TPS	0	$\gamma N \rightarrow D^+\pi^-X$

NODE=M254W

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

- <sup>8</sup> AAIJ 15Y LHCb +  $B^0 \rightarrow \bar{D}^0\pi^+\pi^-$
- <sup>1</sup> From the amplitude analysis in the model describing the  $D^+\pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.
  - <sup>2</sup> From the amplitude analysis in the model describing the  $D^+\pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, nonresonant spin-0 and spin-1 components as well as the  $D_0^*(2400)^0$ ,  $D_2^*(2460)^0$  and  $D_1^*(2760)^0$  resonances.
  - <sup>3</sup> From the Dalitz plot analysis including various  $K^*$  and  $D^{**}$  mesons as well as broad structures in the  $K\pi$  S-wave and the  $D\pi$  S- and P-waves.
  - <sup>4</sup> Modeling the  $\pi^+\pi^-$  S-wave with the Isobar formalism.
  - <sup>5</sup> From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  $A_{D_2}$  fixed to the theoretical prediction of -1.
  - <sup>6</sup> Fit includes the contribution from  $D_0^*(2400)^0$ .
  - <sup>7</sup> Fit includes the contribution from  $D_0^*(2400)^\pm$ .
  - <sup>8</sup> Modeling the  $\pi^+\pi^-$  S-wave with the K-matrix formalism.

NODE=M254W;LINKAGE=D

NODE=M254W;LINKAGE=A

NODE=M254W;LINKAGE=AC

NODE=M254W;LINKAGE=B

NODE=M254W;LINKAGE=AR


NODE=M254W;LINKAGE=LI

NODE=M254W;LINKAGE=LC

NODE=M254W;LINKAGE=C



WEIGHTED AVERAGE  
47.3±0.8 (Error scaled by 1.5)



**$D_2^*(2460)$  DECAY MODES**

NODE=M254215;NODE=M254

$\bar{D}_2^*(2460)$  modes are charge conjugates of modes below.

NODE=M254

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D\pi^-$	seen
$\Gamma_2$ $D^*(2010)\pi^-$	seen
$\Gamma_3$ $D\pi^+\pi^-$	
$\Gamma_4$ $D^*\pi^+\pi^-$	

DESIG=1  
DESIG=2  
DESIG=3  
DESIG=4

**$D_2^*(2460)$  BRANCHING RATIOS**

NODE=M254220

$\Gamma(D\pi^-)/\Gamma_{total}$						$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+\pi^-\pi^-$	
seen	337	ALBRECHT	89B ARG	0	$e^+e^- \rightarrow D^+\pi^-X$	
<b>seen</b>		ALBRECHT	89F ARG	+	$e^+e^- \rightarrow D^0\pi^+X$	
<b>seen</b>		ANJOS	89C TPS	0	$\gamma N \rightarrow D^+\pi^-X$	

NODE=M254R1  
NODE=M254R1

$\Gamma(D^*(2010)\pi^-)/\Gamma_{total}$						$\Gamma_2/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
seen	ACKERSTAFF 97W	OPAL	0	$e^+e^- \rightarrow D^{*+}\pi^-X$		
seen	AVERY 90	CLEO	0	$e^+e^- \rightarrow D^{*+}\pi^-X$		
<b>seen</b>	ALBRECHT 89H	ARG	0	$e^+e^- \rightarrow D^*\pi^-X$		

NODE=M254R2  
NODE=M254R2

$\Gamma(D\pi^-)/\Gamma(D^*(2010)\pi^-)$						$\Gamma_1/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>1.52±0.14 OUR AVERAGE</b>						
1.4 ±0.3 ±0.3	2.3k	1 ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$	
1.1 ±0.4 $\begin{smallmatrix} +0.3 \\ -0.2 \end{smallmatrix}$	1371	2 ABRAMOWICZ13	ZEUS	+	$e^\pm p \rightarrow D^{(*)0}\pi^+X$	OCCUR=2
1.47±0.03±0.16	379k	DEL-AMO-SA...10P	BABR	0	$e^+e^- \rightarrow D^{(*)+}\pi^-X$	
2.8 ±0.8 $\begin{smallmatrix} +0.5 \\ -0.6 \end{smallmatrix}$	1.5k	CHEKANOV 09	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$	
2.2 ±0.7 ±0.6		AVERY 94C	CLE2	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	

NODE=M254R3  
NODE=M254R3

1.9 ±1.1 ±0.3	BERGFELD	94B	CLE2	+	$e^+e^- \rightarrow \text{hadrons}$	OCCUR=2
2.3 ±0.8	AVERY	90	CLEO	0	$e^+e^-$	
3.0 ±1.1 ±1.5	ALBRECHT	89H	ARG	0	$e^+e^- \rightarrow D^*\pi^- X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.9 ±0.5	ABE	04D	BELL	0	$B^- \rightarrow D^{(*)+}\pi^-\pi^-$	
----------	-----	-----	------	---	--------------------------------------	--

<sup>1</sup> From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  $A_{D_2}$  fixed to the theoretical prediction of  $-1$ .

NODE=M254R3;LINKAGE=AR

<sup>2</sup> From the fit of the  $M(D^0\pi^+)$  distribution. The widths of the  $D_1^+$  and  $D_2^{*+}$  are fixed to 25 MeV and 37 MeV, and  $A_{D_1}$  and  $A_{D_2}$  are fixed to the theoretical predictions of 3 and  $-1$ , respectively.

NODE=M254R3;LINKAGE=AB

### $\Gamma(D\pi^-)/[\Gamma(D\pi^-) + \Gamma(D^*(2010)\pi^-)]$ $\Gamma_1/(\Gamma_1+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M254R01  
NODE=M254R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.62±0.03±0.02	8414	<sup>1</sup> AUBERT	09Y	BABR	0	$B^+ \rightarrow D_2^{*0}\ell^+\nu_\ell$	
0.62±0.03±0.02	3361	<sup>1</sup> AUBERT	09Y	BABR	+	$\bar{B}^0 \rightarrow D_2^{*+}\ell^-\nu_\ell$	OCCUR=2

OCCUR=2

<sup>1</sup> Assuming  $\Gamma(\Upsilon(4S) \rightarrow B^+B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.065 \pm 0.026$  and equal partial widths for charged and neutral  $D_2^*$  mesons.

NODE=M254R01;LINKAGE=AU

### $D_2^*(2460)$ POLARIZATION AMPLITUDE $A_{D_2}$

NODE=M254PAM

A polarization amplitude  $A_{D_2}$  is a parameter that depends on the initial polarization of the  $D_2$ . For  $D_2$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_{D_2} \cos^2(\theta_H)$ , where  $\theta_H$  is the angle in the  $D^*$  rest frame between the two pions emitted by the  $D_2 \rightarrow D^*\pi$  and  $D^* \rightarrow D\pi$ .

NODE=M254PAM

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M254PAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-1.16 \pm 0.35$	2.3k	<sup>1</sup> ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^- X$		
consistent with $-1$	243k	DEL-AMO-SA..10P	BABR	0	$e^+e^- \rightarrow D^+\pi^- X$		
$-0.74^{+0.49}_{-0.38}$		<sup>2</sup> AVERY	94C	CLE2	0	$e^+e^- \rightarrow D^{*+}\pi^- X$	

<sup>1</sup> From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions.

NODE=M254PAM;LINKAGE=AB

<sup>2</sup> Systematic uncertainties not estimated.

NODE=M254PAM;LINKAGE=AV

### $D_2^*(2460)$ REFERENCES

NODE=M254

AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57518
AAIJ	15V	PR D91 092002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56575
Also		PR D93 119901 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57289
AAIJ	15X	PR D92 012012	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56588
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56609
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
ABRAMOWICZ 13	NP	B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54743
DEL-AMO-SA...10P	PR	D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534
AUBERT	09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52941
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52929
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
KUZMIN	07	PR D76 012006	A. Kuzmin <i>et al.</i>	(BELLE Collab.)	REFID=51854
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)	REFID=51054
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50011
LINK	04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)	REFID=49775
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46315
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN	95	ZPHY C68 43	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)	REFID=44439
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)	REFID=44096
BERGFELD	94B	PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)	REFID=44099
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT	89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=40736
ALBRECHT	89F	PL B231 208	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40931
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=41001
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)	REFID=40737

$D_0(2550)^0$ 

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

 $J^P = 0^-$  determined by AAIJ 20D.

NODE=M198

NODE=M198

NODE=M198M


NODE=M198M

 $D_0(2550)^0$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2549 ±19 OUR AVERAGE</b>		Error includes scale factor of 4.6. See the ideogram below.		
2518 ± 2 ±7	79k	<sup>1</sup> AAIJ	20D LHCb	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2579.5 ± 3.4 ±5.5	60k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$
2539.4 ± 4.5 ±6.8	34k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.

NODE=M198M;LINKAGE=A

 $D_0(2550)^0$  WIDTH


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>165 ±24 OUR AVERAGE</b>		Error includes scale factor of 2.0. See the ideogram below.		
199 ± 5 ±17	79k	<sup>1</sup> AAIJ	20D LHCb	$B^- \rightarrow D^{*+} \pi^- \pi^-$
177.5 ±17.8 ±46.0	60k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$
130 ±12 ±13	34k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.

NODE=M198W

NODE=M198W

NODE=M198W;LINKAGE=A



**$D_0(2550)^0$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^{*+} \pi^-$	seen

NODE=M198215;NODE=M198

DESIG=1;OUR EVAL;→ UNCHECKED ←

**$D_0(2550)^0$  POLARIZATION AMPLITUDE  $A_{D_J}$**

A polarization amplitude  $A_{D_J}$  is a parameter that depends on the initial polarization of the  $D_J$ . For  $D_J$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_{D_J} \cos^2(\theta_H)$ , where  $\theta_H$  is the angle in the  $D_J$  rest frame between the two pions emitted in the  $D_J \rightarrow D^* \pi$  and  $D^* \rightarrow D \pi$  decays.

NODE=M198PAM

NODE=M198PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M198PAM

• • • We do not use the following data for averages, fits, limits, etc. • • •  
4.2±1.3                  60k                  <sup>1</sup> AAIJ                  13CC LHCb                   $pp \rightarrow D^{*+} \pi^- X$

<sup>1</sup>Systematic uncertainty not estimated.

NODE=M198PAM;LINKAGE=A

**$D_0(2550)^0$  REFERENCES**

NODE=M198

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA... 10P	PR D82	111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

REFID=60253  
REFID=55581  
REFID=53534

$D_1^*(2600)^0$ 

$$I(J^P) = \frac{1}{2}(1^-)$$

OMITTED FROM SUMMARY TABLE

was  $D_j^*(2600)$  $J^P = 1^-$  determined by AAIJ 20D.

NODE=M199

NODE=M199

NODE=M199M

NODE=M199M

OCCUR=2


NODE=M199M;LINKAGE=B

NODE=M199M;LINKAGE=A

NODE=M199M;LINKAGE=DE

 $D_1^*(2600)^0$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>2627 ±10</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 4.4. See the ideogram below.			
2641.9 ± 1.8 ± 4.5	79k	<sup>1</sup> AAIJ	20D	LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2681.1 ± 5.6 ± 14.0	28k	<sup>2</sup> AAIJ	16AH	LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
2649.2 ± 3.5 ± 3.5	51k	AAIJ	13CC	LHCB	$pp \rightarrow D^{*+} \pi^- X$
2608.7 ± 2.4 ± 2.5	26k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2621.3 ± 3.7 ± 4.2	13k	<sup>3</sup> DEL-AMO-SA..10P	BABR	+	$e^+ e^- \rightarrow D^0 \pi^+ X$

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.<sup>2</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.<sup>3</sup> At a fixed width of 93 MeV. $D_1^*(2600)^0$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>141 ±23</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 2.7. See the ideogram below.		
149 ± 4 ± 20	79k	<sup>1</sup> AAIJ	20D	LHCB
186.7 ± 8.5 ± 11.9	28k	<sup>2</sup> AAIJ	16AH	LHCB
140.2 ± 17.1 ± 18.6	51k	AAIJ	13CC	LHCB
93 ± 6 ± 13	26k	DEL-AMO-SA..10P	BABR	


<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.<sup>2</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

NODE=M199W

NODE=M199W

NODE=M199W;LINKAGE=B

NODE=M199W;LINKAGE=A



**$D_1^*(2600)^0$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D\pi$	seen
$\Gamma_2$ $D^+\pi^-$	seen
$\Gamma_3$ $D^0\pi^\pm$	seen
$\Gamma_4$ $D^*\pi$	seen
$\Gamma_5$ $D^{*+}\pi^-$	seen

NODE=M199215;NODE=M199

DESIG=1;OUR EVAL;→ UNCHECKED ←  
 DESIG=2;OUR EVAL;→ UNCHECKED ←  
 DESIG=3;OUR EVAL;→ UNCHECKED ←  
 DESIG=4;OUR EVAL;→ UNCHECKED ←  
 DESIG=5;OUR EVAL;→ UNCHECKED ←

**$D_1^*(2600)^0$  BRANCHING RATIOS**

$\Gamma(D^+\pi^-)/\Gamma(D^{*+}\pi^-)$					$\Gamma_2/\Gamma_5$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.32±0.02±0.09</b>	76k	DEL-AMO-SA...10P	BABR	$e^+e^- \rightarrow D^{(*)+}\pi^- X$	

NODE=M199220

NODE=M199R01  
 NODE=M199R01

**$D_1^*(2600)^0$  REFERENCES**

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA...10P	PR D82	111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

NODE=M199

REFID=60253  
 REFID=57518  
 REFID=55581  
 REFID=53534

**$D^*(2640)^\pm$** 

$$I(J^P) = \frac{1}{2}(??)$$

OMITTED FROM SUMMARY TABLE

Seen in Z decays by ABREU 98M. Not seen by ABBIENDI 01N and CHEKANOV 09. Needs confirmation.

NODE=M158

NODE=M158

 **$D^*(2640)^\pm$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2637 ± 2 ± 6</b>	66 ± 14	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

NODE=M158M

NODE=M158M

 **$D^*(2640)^\pm$  WIDTH**

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;15</b>	95	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

NODE=M158W

NODE=M158W

 **$D^*(2640)^+$  DECAY MODES** $D^*(2640)^-$  modes are charge conjugates of modes below.

NODE=M158215;NODE=M158

NODE=M158

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^*(2010)^+ \pi^+ \pi^-$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

 **$D^*(2640)^\pm$  REFERENCES**

CHEKANOV 09 EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABBIENDI 01N EPJ C20 445	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU 98M PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)

NODE=M158

REFID=52733

REFID=48296

REFID=46315

NODE=M228

 **$D_2(2740)^0$** 

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

was  $D(2740)^0$  $J^P = 2^-$  determined by (AAIJ 20D).

NODE=M228

 **$D_2(2740)^0$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2747 ± 6 OUR AVERAGE</b>				
2751 ± 3 ± 7	79k	<sup>1</sup> AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2737.0 ± 3.5 ± 11.2	7.7k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$

NODE=M228M

NODE=M228M

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.

NODE=M228M;LINKAGE=A

 **$D_2(2740)^0$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>88 ± 19 OUR AVERAGE</b>				
102 ± 6 ± 26	79k	<sup>1</sup> AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
73.2 ± 13.4 ± 25.0	7.7k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$

NODE=M228W

NODE=M228W

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.

NODE=M228W;LINKAGE=A

**$D_2(2740)^0$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^{*+}\pi^-$	seen

NODE=M228215;NODE=M228

DESIG=1;OUR EVAL;→ UNCHECKED ←

 **$D_2(2740)^0$  POLARIZATION AMPLITUDE  $A_{D_J}$** 

A polarization amplitude  $A_{D_J}$  is a parameter that depends on the initial polarization of the  $D_J$ . For  $D_J$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_{D_J} \cos^2(\theta_H)$ , where  $\theta_H$  is the angle in the  $D_J$  rest frame between the two pions emitted in the  $D_J \rightarrow D^*\pi$  and  $D^* \rightarrow D\pi$  decays.

NODE=M228PAM

NODE=M228PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M228PAM

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

3.1±2.2      7.7k      <sup>1</sup> AAIJ      13CC LHCB       $\rho\rho \rightarrow D^{*+}\pi^- X$ <sup>1</sup> Systematic uncertainty not estimated.

NODE=M228PAM;LINKAGE=A

 **$D_2(2740)^0$  REFERENCES**

AAIJ	20D PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M228

REFID=60253  
REFID=55581

NODE=M203

 **$D_3^*(2750)$** 

$$I(J^P) = \frac{1}{2}(3^-)$$

$J^P$  determined by AAIJ 15Y from the Dalitz plot analysis of  $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$  decays.

NODE=M203

 **$D_3^*(2750)$  MASS**

NODE=M203M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>2763.1± 3.2 OUR AVERAGE</b>		Error includes scale factor of 2.1. See the ideogram below.			
2753 ± 4 ± 6	79k	<sup>1</sup> AAIJ	20D	LHCB	$B^- \rightarrow D^{*+}\pi^-\pi^-$
2775.5± 4.5± 6.5	28k	<sup>2</sup> AAIJ	16AH	LHCB	$B^- \rightarrow D^+\pi^-\pi^-$
2798 ± 7 ± 7		<sup>3</sup> AAIJ	15Y	LHCB	$B^0 \rightarrow \bar{D}^0\pi^+\pi^-$
2761.1± 5.1± 6.5	14k	AAIJ	13CC	LHCB 0	$\rho\rho \rightarrow D^{*+}\pi^- X$
2760.1± 1.1± 3.7	56k	AAIJ	13CC	LHCB 0	$\rho\rho \rightarrow D^+\pi^- X$
2771.7± 1.7± 3.8	20k	AAIJ	13CC	LHCB +	$\rho\rho \rightarrow D^0\pi^+ X$
2752.4± 1.7± 2.7	23.5k	<sup>4</sup> DEL-AMO-SA..10P	BABR	0	$e^+e^- \rightarrow D^{*+}\pi^- X$
2763.3± 2.3± 2.3	11.3k	<sup>4</sup> DEL-AMO-SA..10P	BABR	0	$e^+e^- \rightarrow D^+\pi^- X$
2769.7± 3.8± 1.5	5.7k	<sup>4,5</sup> DEL-AMO-SA..10P	BABR	+	$e^+e^- \rightarrow D^0\pi^+ X$

NODE=M203M

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2802 ±11 ±10      <sup>6</sup> AAIJ      15Y LHCB       $B^0 \rightarrow \bar{D}^0\pi^+\pi^-$ 

OCCUR=2

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+}\pi^-\pi^-$  decay.

NODE=M203M;LINKAGE=D

<sup>2</sup> From the amplitude analysis in the model describing the  $D^+\pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

NODE=M203M;LINKAGE=C

<sup>3</sup> Modeling the  $\pi^+\pi^-$  S-wave with the Isobar formalism.<sup>4</sup> The states observed in the  $D^*\pi$  and  $D\pi$  final states are not necessarily the same.<sup>5</sup> At a fixed width of 60.9 MeV.<sup>6</sup> Modeling the  $\pi^+\pi^-$  S-wave with the K-matrix formalism.


NODE=M203M;LINKAGE=A

NODE=M203M;LINKAGE=DE

NODE=M203M;LINKAGE=DA

NODE=M203M;LINKAGE=B





**$D_3^*(2750)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>66 ± 5 OUR AVERAGE</b>					
66 ± 10 ± 14	79k	1 AAIJ 20D LHC B			$B^- \rightarrow D^{*+} \pi^- \pi^-$
95.3 ± 9.6 ± 34.0	28k	2 AAIJ 16AH LHC B			$B^- \rightarrow D^+ \pi^- \pi^-$
105 ± 18 ± 24		3 AAIJ 15Y LHC B			$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
74.4 ± 3.4 ± 37.0	14k	AAIJ 13CC LHC B 0			$pp \rightarrow D^{*+} \pi^- X$
74.4 ± 3.4 ± 19.1	56k	AAIJ 13CC LHC B 0			$pp \rightarrow D^+ \pi^- X$
66.7 ± 6.6 ± 10.5	20k	AAIJ 13CC LHC B +			$pp \rightarrow D^0 \pi^+ X$
71 ± 6 ± 11	23.5k	4 DEL-AMO-SA...10P BABR			$e^+ e^- \rightarrow D^{*+} \pi^- X$
60.9 ± 5.1 ± 3.6	11.3k	4 DEL-AMO-SA...10P BABR			$e^+ e^- \rightarrow D^+ \pi^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

154 ± 27 ± 16		5 AAIJ 15Y LHC B			$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
---------------	--	------------------	--	--	---

<sup>1</sup> From a full four-body amplitude analysis of the  $B^- \rightarrow D^{*+} \pi^- \pi^-$  decay.

<sup>2</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

<sup>3</sup> Modeling the  $\pi^+ \pi^-$  S-wave with the Isobar formalism.

<sup>4</sup> The states observed in the  $D^* \pi$  and  $D \pi$  final states are not necessarily the same.

<sup>5</sup> Modeling the  $\pi^+ \pi^-$  S-wave with the K-matrix formalism.

**$D_3^*(2750)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D \pi$	seen
$\Gamma_2$ $D^+ \pi^-$	seen
$\Gamma_3$ $D^0 \pi^\pm$	seen
$\Gamma_4$ $D^* \pi$	seen
$\Gamma_5$ $D^{*+} \pi^-$	seen

**$D_3^*(2750)$  BRANCHING RATIOS**

$\Gamma(D^+ \pi^-)/\Gamma(D^{*+} \pi^-)$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma_5$
	<b>0.42 ± 0.05 ± 0.11</b>	34.8k	1 DEL-AMO-SA...10P BABR		$e^+ e^- \rightarrow D^{(*)+} \pi^- X$	

NODE=M203W

NODE=M203W

OCCUR=2

OCCUR=4

OCCUR=2

OCCUR=2

NODE=M203W;LINKAGE=D

NODE=M203W;LINKAGE=C

NODE=M203W;LINKAGE=A

NODE=M203W;LINKAGE=DE

NODE=M203W;LINKAGE=B

NODE=M203215;NODE=M203

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=5;OUR EVAL;→ UNCHECKED ←

NODE=M203220

NODE=M203R01

NODE=M203R01

<sup>1</sup> The states observed in the  $D^* \pi$  and  $D \pi$  final states are not necessarily the same.

NODE=M203R01;LINKAGE=DE

### $D_3^*(2750)$ POLARIZATION AMPLITUDE $A_D$

NODE=M203PAM

NODE=M203PAM

A polarization amplitude  $A_D$  is a parameter that depends on the initial polarization of the  $D_3^*(2750)$ . For  $D_3^*(2750)$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_D \cos(\theta_H)$ , where  $\theta_H$  is the angle in the  $D^*$  rest frame between the two pions emitted by the  $D_3^*(2750) \rightarrow D^* \pi$  and  $D^* \rightarrow D \pi$ .

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M203PAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.33±0.28	23.5k	<sup>1</sup> DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$
------------	-------	-------------------------------	------	--------------------------------------

NODE=M203PAM;LINKAGE=DE

<sup>1</sup> Systematic uncertainties not estimated. The states observed in the  $D^* \pi$  and  $D \pi$  final states are not necessarily the same.

### $D_3^*(2750)$ REFERENCES

NODE=M203

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60253
AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57518
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=56609
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
DEL-AMO-SA... 10P	PR D82 111101		P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534

NODE=M249

$D_1^*(2760)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

OMITTED FROM SUMMARY TABLE

$J^P$  determined by AAIJ 15V from the Dalitz plot analysis of  $B^- \rightarrow D^+ K^- \pi^-$  decays.

NODE=M249

### $D_1^*(2760)^0$ MASS

NODE=M249M

NODE=M249M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2781±18±13</b>	2k	<sup>1</sup> AAIJ	15V	LHCB $B^- \rightarrow D^+ K^- \pi^-$

<sup>1</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, nonresonant spin-0 and spin-1 components as well as the  $D_0^*(2400)^0$ ,  $D_2^*(2460)^0$  and  $D_1^*(2760)^0$  resonances.

NODE=M249M;LINKAGE=A

### $D_1^*(2760)^0$ WIDTH

NODE=M249W

NODE=M249W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>177±32±21</b>	2k	<sup>1</sup> AAIJ	15V	LHCB $B^- \rightarrow D^+ K^- \pi^-$

<sup>1</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, nonresonant spin-0 and spin-1 components as well as the  $D_0^*(2400)^0$ ,  $D_2^*(2460)^0$  and  $D_1^*(2760)^0$  resonances.

NODE=M249W;LINKAGE=A

### $D_1^*(2760)^0$ DECAY MODES

NODE=M249215;NODE=M249

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^+ \pi^-$	seen

DESIG=1

### $D_1^*(2760)^0$ BRANCHING RATIOS

NODE=M249225

$\Gamma(D^+ \pi^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> AAIJ	15V	LHCB $B^- \rightarrow D^+ K^- \pi^-$

NODE=M249R01

NODE=M249R01

OCCUR=2

<sup>1</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, nonresonant spin-0 and spin-1 components as well as the  $D_0^*(2400)^0$ ,  $D_2^*(2460)^0$  and  $D_1^*(2760)^0$  resonances.

NODE=M249R01;LINKAGE=A

$D_1^*(2760)^0$  REFERENCES

AAIJ 15V PR D91 092002 R. Aaij *et al.* (LHCb Collab.) JP  
 Also PR D93 119901 (errat.) R. Aaij *et al.* (LHCb Collab.)

NODE=M249

REFID=56575  
REFID=57289 $D(3000)^0$ 

$$I(J^P) = \frac{1}{2}(?)^2$$

NODE=M229

## OMITTED FROM SUMMARY TABLE

Both natural- and unnatural-parity components observed depending on the decay mode (AAIJ 13CC).

NODE=M229

 $D(3000)^0$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3214 ± 29 ± 49</b>	28k	<sup>1</sup> AAIJ	16AH LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2971.8 ± 8.7	9.5k	<sup>2,3</sup> AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
3008.1 ± 4.0	17.6k	<sup>2,4</sup> AAIJ	13CC LHCB	$pp \rightarrow D^+ \pi^- X$

NODE=M229M

NODE=M229M

<sup>1</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

<sup>2</sup> Systematic uncertainty not estimated.

<sup>3</sup> Unnatural parity preferred.

<sup>4</sup> Natural parity state. A state  $D(3000)^+$  is possibly seen in  $D^0 \pi^+$  final state.

OCCUR=2

NODE=M229M;LINKAGE=D

NODE=M229M;LINKAGE=A

NODE=M229M;LINKAGE=B

NODE=M229M;LINKAGE=C

 $D(3000)^0$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>186 ± 38 ± 72</b>	28k	<sup>5</sup> AAIJ	16AH LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
188.1 ± 44.8	9.5k	<sup>6,7</sup> AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
110.5 ± 11.5	17.6k	<sup>6,8</sup> AAIJ	13CC LHCB	$pp \rightarrow D^+ \pi^- X$

NODE=M229W

NODE=M229W

<sup>5</sup> From the amplitude analysis in the model describing the  $D^+ \pi^-$  wave together with virtual contributions from the  $D^*(2007)^0$  and  $B^{*0}$  states, and components corresponding to the  $D_2^*(2460)^0$ ,  $D_1^*(2680)^0$ ,  $D_3^*(2760)^0$ , and  $D_2^*(3000)^0$  resonances.

<sup>6</sup> Systematic uncertainty not estimated.

<sup>7</sup> Unnatural parity preferred.

<sup>8</sup> Natural parity state. A state  $D(3000)^+$  is possibly seen in  $D^0 \pi^+$  final state.

OCCUR=2

NODE=M229W;LINKAGE=D

NODE=M229W;LINKAGE=A

NODE=M229W;LINKAGE=C

NODE=M229W;LINKAGE=B

 $D(3000)^0$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^{*+} \pi^-$	seen

NODE=M229215;NODE=M229

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D(3000)^0$  POLARIZATION AMPLITUDE  $A_{D_J}$ 

A polarization amplitude  $A_{D_J}$  is a parameter that depends on the initial polarization of the  $D_J$ . For  $D_J$  decays the helicity angle,  $\theta_H$ , distribution varies like  $1 + A_{D_J} \cos^2(\theta_H)$ , where  $\theta_H$  is the angle in the  $D_J$  rest frame between the two pions emitted in the  $D_J \rightarrow D^* \pi$  and  $D^* \rightarrow D \pi$  decays.

NODE=M229PAM

NODE=M229PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.5 ± 0.9</b>	9.5k	<sup>9</sup> AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$

NODE=M229PAM

<sup>9</sup> Systematic uncertainty not estimated.

NODE=M229PAM;LINKAGE=A

 $D(3000)^0$  REFERENCES

AAIJ 16AH PR D94 072001 R. Aaij *et al.* (LHCb Collab.)  
 AAIJ 13CC JHEP 1309 145 R. Aaij *et al.* (LHCb Collab.)

NODE=M229

REFID=57518  
REFID=55581

# CHARMED, STRANGE MESONS

## ( $C = \pm 1, S = \pm 1$ )

### (including possibly non- $q\bar{q}$ states)

$$D_s^+ = c\bar{s}, D_s^- = \bar{c}s, \quad \text{similarly for } D_s^{*\pm}$$

$D_s^{*\pm}$

$$I(J^P) = 0(?^?)$$

$J^P$  is natural, width and decay modes consistent with  $1^-$ .

### $D_s^{*\pm}$ MASS

The fit includes  $D_s^\pm, D_s^0, D_s^{\pm*}, D_s^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2112.2 ± 0.4 OUR FIT</b>			
<b>2106.6 ± 2.1 ± 2.7</b>	<sup>1</sup> BLAYLOCK	87	MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

<sup>1</sup> Assuming  $D_s^\pm$  mass =  $1968.7 \pm 0.9$  MeV.

### $m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes  $D_s^\pm, D_s^0, D_s^{\pm*}, D_s^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>143.8 ± 0.4 OUR FIT</b>				
<b>143.9 ± 0.4 OUR AVERAGE</b>				
143.76 ± 0.39 ± 0.40		GRONBERG	95	CLE2 $e^+e^-$
144.22 ± 0.47 ± 0.37		BROWN	94	CLE2 $e^+e^-$
142.5 ± 0.8 ± 1.5		<sup>2</sup> ALBRECHT	88	ARG $e^+e^- \rightarrow D_s^\pm \gamma X$
139.5 ± 8.3 ± 9.7	60	AIHARA	84D	TPC $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
143.0 ± 18.0	8	ASRATYAN	85	HLBC FNAL 15-ft, $\nu$ - <sup>2</sup> H
110 ± 46		BRANDELIK	79	DASP $e^+e^- \rightarrow D_s^\pm \gamma X$

<sup>2</sup> Result includes data of ALBRECHT 84B.

### $D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 1.9</b>	90	GRONBERG	95	CLE2 $e^+e^-$
<b>&lt; 4.5</b>	90	ALBRECHT	88	ARG $E_{cm}^{ee} = 10.2$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 4.9	90	BROWN	94	CLE2 $e^+e^-$
< 22	90	BLAYLOCK	87	MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

### $D_s^{*+}$ DECAY MODES

$D_s^{*-}$  modes are charge conjugates of the modes below.

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad D_s^+ \gamma$	(93.5 ± 0.7) %
$\Gamma_2 \quad D_s^+ \pi^0$	( 5.8 ± 0.7) %
$\Gamma_3 \quad D_s^+ e^+ e^-$	( 6.7 ± 1.6) × 10 <sup>-3</sup>

NODE=MXXX040

NODE=MXXX040

NODE=S074

NODE=S074

NODE=S074M

NODE=S074M

NODE=S074M

NODE=S074M;LINKAGE=E

NODE=S074DM

NODE=S074DM

NODE=S074DM

NODE=S074DM;LINKAGE=A

NODE=S074W

NODE=S074W

NODE=S074215;NODE=S074

NODE=S074

DESIG=1

DESIG=2

DESIG=3

**CONSTRAINED FIT INFORMATION**

An overall fit to 2 branching ratios uses 3 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 0.0$  for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	-97	
$x_3$	-19	-4
	$x_1$	$x_2$

 **$D_s^{*+}$  BRANCHING RATIOS**

NODE=S074220

 $\Gamma(D_s^+ \gamma) / \Gamma_{\text{total}}$  $\Gamma_1 / \Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=S074R1  
NODE=S074R1**0.935 ± 0.007 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ASRATYAN	91	HLBC	$\bar{\nu}_\mu \text{Ne}$
seen	ALBRECHT	88	ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$
seen	AIHARA	84D		
seen	ALBRECHT	84B		
seen	BRANDELIK	79		

 $\Gamma(D_s^+ \pi^0) / \Gamma(D_s^+ \gamma)$  $\Gamma_2 / \Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=S074R2  
NODE=S074R2**0.062 ± 0.008 OUR FIT****0.062 ± 0.008 OUR AVERAGE**

0.062 ± 0.005 ± 0.006	AUBERT, BE	05G	BABR	10.6 $e^+ e^- \rightarrow$ hadrons
0.062 $^{+0.020}_{-0.018} \pm 0.022$	GRONBERG	95	CLE2	$e^+ e^-$

 $\Gamma(D_s^+ e^+ e^-) / \Gamma(D_s^+ \gamma)$  $\Gamma_3 / \Gamma_1$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=S074R01  
NODE=S074R01**7.2 ± 1.7 OUR FIT**

7.2 $^{+1.5}_{-1.3} \pm 1.0$	38	CRONIN-HEN..12	CLEO	4.17 $e^+ e^- \rightarrow$ hadrons
------------------------------	----	----------------	------	------------------------------------

 **$D_s^{*\pm}$  REFERENCES**

NODE=S074

CRONIN-HEN...12	PR D86 072005	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=54627
AUBERT, BE	05G PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50942
GRONBERG	95 PRL 75 3232	J. Gronberg <i>et al.</i>	(CLEO Collab.)	REFID=44568
BROWN	94 PR D50 1884	D. Brown <i>et al.</i>	(CLEO Collab.)	REFID=43868
ASRATYAN	91 PL B257 525	A.E. Asratyan <i>et al.</i>	(ITEP, BELG, SACL+)	REFID=41582
ALBRECHT	88 PL B207 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40269
BLAYLOCK	87 PRL 58 2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)	REFID=40005
ASRATYAN	85 PL 156B 441	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=22887
AIHARA	84D PRL 53 2465	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=11561
ALBRECHT	84B PL 146B 111	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22886
BRANDELIK	79 PL 80B 412	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=11442

$$D_{s0}^*(2317)^\pm$$

$$I(J^P) = 0(0^+)$$

$J, P$  need confirmation.

AUBERT 06P and CHOI 15A do not observe neutral and doubly charged partners of the  $D_{s0}^*(2317)^\pm$ . See the review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M172

NODE=M172

### $D_{s0}^*(2317)^\pm$ MASS

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M172M

NODE=M172M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2317.8±0.5 OUR FIT</b>				
<b>2318.0±0.7 OUR AVERAGE</b>				
2318.3±1.2±1.2	115	<sup>1</sup> ABLIKIM	18J BES3	4.6 $e^+e^- \rightarrow D_s^{*\pm} D_{s0}^*(2317)^\mp$
2319.6±0.2±1.4	3.1k	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow D_s^+ \pi^0 X$
2317.3±0.4±0.8	1.0k	<sup>2</sup> AUBERT	04E BABR	10.6 $e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2317.2±1.3	88	<sup>3</sup> AUBERT,B	04S BABR	$B \rightarrow D_{s0}^{(*)}(2317)^+ \bar{D}^{(*)}$
2317.2±0.5±0.9	761	<sup>4</sup> MIKAMI	04 BELL	10.6 $e^+e^-$
2316.8±0.4±3.0	1.2k	<sup>4,5</sup> AUBERT	03G BABR	10.6 $e^+e^-$
2317.6±1.3	273	<sup>4,6</sup> AUBERT	03G BABR	10.6 $e^+e^-$
2319.8±2.1±2.0	24	<sup>4</sup> KROKOVNY	03B BELL	10.6 $e^+e^-$

NODE=M172M

<sup>1</sup> From a fit of the  $D_s^*$  recoil mass where the  $D_{s0}^*(2317)$  signal is described with a Crystal Ball function convolved with a Gaussian function.

<sup>2</sup> Supersedes AUBERT 03G.

<sup>3</sup> Systematic errors not evaluated.

<sup>4</sup> Not independent of the corresponding  $m_{D_{s0}^*(2317)} - m_{D_s}$ .

<sup>5</sup> From  $D_s^+ \rightarrow K^+ K^- \pi^+$  decay.

<sup>6</sup> From  $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$  decay.

NODE=M172M;LINKAGE=A

NODE=M172M;LINKAGE=AU

NODE=M172M;LINKAGE=AB

NODE=M172M;LINKAGE=B1

NODE=M172M;LINKAGE=A1

NODE=M172M;LINKAGE=A2

### $m_{D_{s0}^*(2317)^\pm} - m_{D_s^\pm}$

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M172DM

NODE=M172DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>349.4±0.5 OUR FIT</b>				
<b>349.2±0.7 OUR AVERAGE</b>				
348.7±0.5±0.7	761	MIKAMI	04 BELL	10.6 $e^+e^-$
350.0±1.2±1.0	135	BESSION	03 CLE2	10.6 $e^+e^-$
351.3±2.1±1.9	24	<sup>7</sup> KROKOVNY	03B BELL	10.6 $e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
349.6±0.4±3.0	1267	<sup>8,9</sup> AUBERT	03G BABR	10.6 $e^+e^-$
350.2±1.3	273	<sup>10,11</sup> AUBERT	03G BABR	10.6 $e^+e^-$
<sup>7</sup> Recalculated by us using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV.				
<sup>8</sup> From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.				
<sup>9</sup> Recalculated by us using $m_{D_s^+} = 1967.20 \pm 0.03$ MeV.				
<sup>10</sup> From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.				
<sup>11</sup> Recalculated by us using $m_{D_s^+} = 1967.4 \pm 0.2$ MeV. Systematic errors not estimated.				

NODE=M172DM

OCCUR=2

NODE=M172DM;LINKAGE=K3

NODE=M172DM;LINKAGE=A1

NODE=M172DM;LINKAGE=C1

NODE=M172DM;LINKAGE=A2

NODE=M172DM;LINKAGE=C2

### $D_{s0}^*(2317)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.8	95	3180	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow D_s^+ \pi^0 X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 4.6	90	761	MIKAMI	04 BELL	10.6 $e^+e^-$
< 10			AUBERT	03G BABR	10.6 $e^+e^-$
< 7	90	135	BESSION	03 CLE2	10.6 $e^+e^-$

NODE=M172W

NODE=M172W

$D_{s0}^*(2317)^\pm$  DECAY MODES

NODE=M172215;NODE=M172

 $D_{s0}^*(2317)^-$  modes are charge conjugates of modes below.

NODE=M172

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $D_s^+ \pi^0$	$(100^{+0}_{-20})\%$	
$\Gamma_2$ $D_s^+ \gamma$	$< 5\%$	90%
$\Gamma_3$ $D_s^*(2112)^+ \gamma$	$< 6\%$	90%
$\Gamma_4$ $D_s^+ \gamma \gamma$	$< 18\%$	95%
$\Gamma_5$ $D_s^*(2112)^+ \pi^0$	$< 11\%$	90%
$\Gamma_6$ $D_s^+ \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
$\Gamma_7$ $D_s^+ \pi^0 \pi^0$	not seen	

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7;OUR EVAL;→ UNCHECKED ←

 $D_{s0}^*(2317)^\pm$  BRANCHING RATIOS

NODE=M172220

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.00^{+0.00+0.00}_{-0.14-0.14}$	47	ABLIKIM	18J	BES3 4.6 $e^+ e^- \rightarrow D_s^{*\pm} D_{s0}^*(2317)^\mp$

NODE=M172R1  
NODE=M172R1

••• We do not use the following data for averages, fits, limits, etc. •••

seen 1.5k AUBERT 03G BABR 10.6  $e^+ e^-$ 

$\Gamma(D_s^+ \gamma)/\Gamma(D_s^+ \pi^0)$	$\Gamma_2/\Gamma_1$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.05$	90	MIKAMI	04	BELL 10.6 $e^+ e^-$
$<0.14$	95	AUBERT	06P	BABR 10.6 $e^+ e^-$
$<0.052$	90	BESSON	03	CLE2 10.6 $e^+ e^-$

NODE=M172R5  
NODE=M172R5

••• We do not use the following data for averages, fits, limits, etc. •••

$\Gamma(D_s^*(2112)^+ \gamma)/\Gamma(D_s^+ \pi^0)$	$\Gamma_3/\Gamma_1$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.059$	90	BESSON	03	CLE2 10.6 $e^+ e^-$
$<0.16$	95	AUBERT	06P	BABR 10.6 $e^+ e^-$
$<0.18$	90	MIKAMI	04	BELL 10.6 $e^+ e^-$

NODE=M172R6  
NODE=M172R6

••• We do not use the following data for averages, fits, limits, etc. •••

$\Gamma(D_s^+ \gamma \gamma)/\Gamma(D_s^+ \pi^0)$	$\Gamma_4/\Gamma_1$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.18$	95	AUBERT	06P	BABR 10.6 $e^+ e^-$
not seen		AUBERT	03G	BABR 10.6 $e^+ e^-$

NODE=M172R7  
NODE=M172R7

$\Gamma(D_s^*(2112)^+ \pi^0)/\Gamma(D_s^+ \pi^0)$	$\Gamma_5/\Gamma_1$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.11$	90	BESSON	03	CLE2 10.6 $e^+ e^-$

NODE=M172R8  
NODE=M172R8

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma(D_s^+ \pi^0)$	$\Gamma_6/\Gamma_1$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.004$	90	MIKAMI	04	BELL 10.6 $e^+ e^-$
$<0.005$	95	AUBERT	06P	BABR 10.6 $e^+ e^-$
$<0.019$	90	BESSON	03	CLE2 10.6 $e^+ e^-$

NODE=M172R9  
NODE=M172R9

$\Gamma(D_s^+ \pi^0 \pi^0)/\Gamma(D_s^+ \pi^0)$	$\Gamma_7/\Gamma_1$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.25$	95	AUBERT	06P	BABR 10.6 $e^+ e^-$

NODE=M172R10  
NODE=M172R10

$D_{s0}^*(2317)^\pm$  REFERENCES

ABLIKIM	18J	PR D97 051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58895
CHOI	15A	PR D91 092011	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=56577
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
AUBERT	03G	PRL 90 242001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49417
BESSION	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

NODE=M172

 $D_{s1}(2460)^\pm$ 

$$I(J^P) = 0(1^+)$$

See the review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M173

NODE=M173

 $D_{s1}(2460)^\pm$  MASS

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M173M

NODE=M173M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M173M

**2459.5±0.6 OUR FIT** Error includes scale factor of 1.1.**2459.6±0.9 OUR AVERAGE** Error includes scale factor of 1.3.2460.1±0.2±0.8 <sup>1</sup> AUBERT 06P BABR 10.6  $e^+e^-$ 2458.0±1.0±1.0 195 AUBERT 04E BABR 10.6  $e^+e^-$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

2459.5±1.2±3.7 920 AUBERT 06P BABR 10.6  $e^+e^- \rightarrow D_s^+ \gamma X$ 

OCCUR=2

2458.6±1.0±2.5 560 AUBERT 06P BABR 10.6  $e^+e^- \rightarrow D_s^+ \pi^0 \gamma X$ 

OCCUR=3

2460.2±0.2±0.8 123 AUBERT 06P BABR 10.6  $e^+e^- \rightarrow D_s^+ \pi^+ \pi^- X$ 

OCCUR=4

2458.9±1.5 112 <sup>2</sup> AUBERT,B 04S BABR  $B \rightarrow D_{s1}(2460)^+ \bar{D}^{(*)}$ 2461.1±1.6 139 <sup>3</sup> AUBERT,B 04S BABR  $B \rightarrow D_{s1}(2460)^+ \bar{D}^{(*)}$ 

OCCUR=2

2456.5±1.3±1.3 126 <sup>4,5</sup> MIKAMI 04 BELL 10.6  $e^+e^-$ 2459.5±1.3±2.0 152 <sup>6,7</sup> MIKAMI 04 BELL 10.6  $e^+e^-$ 

OCCUR=2

2459.9±0.9±1.6 60 <sup>6,7</sup> MIKAMI 04 BELL 10.6  $e^+e^-$ 

OCCUR=3

2459.2±1.6±2.0 57 KROKOVNY 03B BELL 10.6  $e^+e^-$ <sup>1</sup> The average of the values obtained from the  $D_s^+ \gamma, D_s^+ \pi^0 \gamma, D_s^+ \pi^+ \pi^-$  final state.

NODE=M173M;LINKAGE=UB

<sup>2</sup> Systematic errors not evaluated. From the decay to  $D_s^{*+} \pi^0$ .

NODE=M173M;LINKAGE=AU

<sup>3</sup> Systematic errors not evaluated. From the decay to  $D_s^+ \gamma$ .

NODE=M173M;LINKAGE=AB

<sup>4</sup> Not independent of the corresponding  $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$ .

NODE=M173M;LINKAGE=B1

<sup>5</sup> Using  $m_{D_s^{*+}} = 2112.4 \pm 0.7$  MeV.

NODE=M173M;LINKAGE=B2

<sup>6</sup> Not independent of the corresponding  $m_{D_{s1}(2460)^\pm} - m_{D_s^\pm}$ .

NODE=M173M;LINKAGE=B3

<sup>7</sup> Using  $m_{D_s^+} = 1968.5 \pm 0.6$  MeV.

NODE=M173M;LINKAGE=B4

 $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$ 

NODE=M173MD

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M173MD

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------


NODE=M173MD

**347.3±0.7 OUR FIT** Error includes scale factor of 1.2.**347.1±2.2 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.344.1±1.3±1.1 126 MIKAMI 04 BELL 10.6  $e^+e^-$ 351.2±1.7±1.0 41 BESSION 03 CLE2 10.6  $e^+e^-$ 346.8±1.6±1.9 57 <sup>8</sup> KROKOVNY 03B BELL 10.6  $e^+e^-$



<sup>8</sup> Recalculated by us using  $m_{D_s^{*+}} = 2112.4 \pm 0.7$  MeV.

NODE=M173MD;LINKAGE=K3



**$m_{D_{s1}(2460)^{\pm}} - m_{D_s^{\pm}}$**

NODE=M173DM

The fit includes  $D^{\pm}, D^0, D_s^{\pm}, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^{\pm}$  mass and mass difference measurements.

NODE=M173DM

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M173DM

**491.1±0.6 OUR FIT** Error includes scale factor of 1.1.

**491.3±1.4 OUR AVERAGE**

491.0±1.3±1.9	152	<sup>9</sup> MIKAMI	04	BELL	10.6 $e^+e^-$
491.4±0.9±1.5	60	<sup>10</sup> MIKAMI	04	BELL	10.6 $e^+e^-$

OCCUR=2

<sup>9</sup> From the decay to  $D_s^{\pm}\gamma$ .

NODE=M173DM;LINKAGE=M1

<sup>10</sup> From the decay to  $D_s^{\pm}\pi^+\pi^-$ .

NODE=M173DM;LINKAGE=M2

**$D_{s1}(2460)^{\pm}$  WIDTH**

NODE=M173W

VALUE (MeV) CL% EVTS DOCUMENT ID TECN COMMENT

NODE=M173W

< 3.5 95 123 AUBERT 06P BABR 10.6  $e^+e^- \rightarrow D_s^+\pi^+\pi^-X$

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.3 95 560 AUBERT 06P BABR 10.6  $e^+e^- \rightarrow D_s^+\pi^0\gamma X$

OCCUR=2

<10 195 AUBERT 04E BABR 10.6  $e^+e^-$

< 5.5 90 126 MIKAMI 04 BELL 10.6  $e^+e^-$

< 7 90 41 BESSON 03 CLE2 10.6  $e^+e^-$

**$D_{s1}(2460)^+$  DECAY MODES**

NODE=M173215;NODE=M173

$D_{s1}(2460)^-$  modes are charge conjugates of the modes below.

NODE=M173

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $D_s^{*+}\pi^0$	(48 ± 11 ) %	
$\Gamma_2$ $D_s^+\gamma$	(18 ± 4 ) %	
$\Gamma_3$ $D_s^+\pi^+\pi^-$	( 4.3± 1.3) %	S=1.1
$\Gamma_4$ $D_s^{*+}\gamma$	< 8 %	CL=90%

DESIG=1

DESIG=2

DESIG=3

DESIG=4

$\Gamma_5$	$D_{s0}^*(2317)^+\gamma$	$(3.7^{+5.0}_{-2.4})\%$	DESIG=5
$\Gamma_6$	$D_s^+\pi^0$		DESIG=7
$\Gamma_7$	$D_s^+\pi^0\pi^0$		DESIG=8
$\Gamma_8$	$D_s^+\gamma\gamma$		DESIG=9

### CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 8 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 3.4$  for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_2$	80		
$x_3$	68	62	
$x_5$	-3	25	26
	$x_1$	$x_2$	$x_3$

### $D_{s1}(2460)^\pm$ BRANCHING RATIOS

NODE=M173220

$\Gamma(D_s^{*+}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

VALUE EVTS DOCUMENT ID TECN COMMENT

**0.48±0.11 OUR FIT**

**0.56±0.13±0.09** <sup>11</sup> AUBERT 06N BABR  $B \rightarrow D_{s1}(2460)^-\bar{D}^*$

••• We do not use the following data for averages, fits, limits, etc. •••

seen 41 BESSON 03 CLE2  $10.6 e^+ e^-$

<sup>11</sup> Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.

NODE=M173R1  
NODE=M173R1

NODE=M173R1;LINKAGE=AU

$\Gamma(D_s^+\gamma)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

VALUE DOCUMENT ID TECN COMMENT

**0.18±0.04 OUR FIT**

**0.16±0.04±0.03** <sup>12</sup> AUBERT 06N BABR  $B \rightarrow D_{s1}(2460)^-\bar{D}^*$

<sup>12</sup> Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.

NODE=M173R6  
NODE=M173R6

NODE=M173R6;LINKAGE=AU

$\Gamma(D_s^+\gamma)/\Gamma(D_s^{*+}\pi^0)$   $\Gamma_2/\Gamma_1$

VALUE CL% EVTS DOCUMENT ID TECN COMMENT

**0.38 ±0.05 OUR FIT**

**0.44 ±0.09 OUR AVERAGE**

0.55 ±0.13 ±0.08 152 MIKAMI 04 BELL  $10.6 e^+ e^-$

0.38 ±0.11 ±0.04 38 KROKOVNY 03B BELL  $10.6 e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

0.274±0.045±0.020 251 <sup>13</sup> AUBERT,B 04S BABR  $B \rightarrow D_{s1}(2460)^+\bar{D}^*$

< 0.49 90 BESSON 03 CLE2  $10.6 e^+ e^-$

<sup>13</sup> Used by AUBERT 06N in their measurement of  $B(D_s^{*-}\pi^0)$  and  $B(D_s^-\gamma)$ .

NODE=M173R2  
NODE=M173R2

NODE=M173R2;LINKAGE=AU

$\Gamma(D_s^+\pi^+\pi^-)/\Gamma(D_s^{*+}\pi^0)$   $\Gamma_3/\Gamma_1$

VALUE CL% EVTS DOCUMENT ID TECN COMMENT

**0.090±0.020 OUR FIT** Error includes scale factor of 1.2.

**0.14 ±0.04 ±0.02** 60 MIKAMI 04 BELL  $10.6 e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.08 90 BESSON 03 CLE2  $10.6 e^+ e^-$

NODE=M173R3  
NODE=M173R3

$\Gamma(D_s^{*+}\gamma)/\Gamma(D_s^{*+}\pi^0)$   $\Gamma_4/\Gamma_1$

VALUE CL% DOCUMENT ID TECN COMMENT

**<0.16** 90 BESSON 03 CLE2  $10.6 e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.31 90 MIKAMI 04 BELL  $10.6 e^+ e^-$

NODE=M173R4  
NODE=M173R4

$\Gamma(D_{s0}^*(2317)^+\gamma)/\Gamma(D_s^{*+}\pi^0)$					$\Gamma_5/\Gamma_1$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.22	95	AUBERT	04E	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R5 NODE=M173R5
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.58	90	BESSON	03	CLE2	10.6 e <sup>+</sup> e <sup>-</sup>	
$\Gamma(D_s^{*+}\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_1/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>0.93±0.09 OUR FIT</b>						
<b>0.97±0.09±0.05</b>		AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R7 NODE=M173R7
$\Gamma(D_s^+\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_2/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>0.35 ±0.04 OUR FIT</b>						
<b>0.337±0.036±0.038</b>		AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R8 NODE=M173R8
$\Gamma(D_s^+\pi^+\pi^-)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_3/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>0.083±0.017 OUR FIT</b> Error includes scale factor of 1.2.						
<b>0.077±0.013±0.008</b>		AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R9 NODE=M173R9
$\Gamma(D_s^{*+}\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_4/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.24	95	AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R10 NODE=M173R10
$\Gamma(D_{s0}^*(2317)^+\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_5/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.25	95	AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R11 NODE=M173R11
$\Gamma(D_s^+\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_6/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.042	95	AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R12 NODE=M173R12
$\Gamma(D_s^+\pi^0\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_7/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.68	95	AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R13 NODE=M173R13
$\Gamma(D_s^+\gamma\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_8/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.33	95	AUBERT	06P	BABR	10.6 e <sup>+</sup> e <sup>-</sup>	NODE=M173R14 NODE=M173R14

 **$D_{s1}(2460)^\pm$  REFERENCES**

$D_{s1}(2460)^\pm$ REFERENCES					NODE=M173
AUBERT	06N	PR D74 031103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51142
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
BESSON	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

**$D_{s1}(2536)^\pm$**  $I(J^P) = 0(1^+)$   
 $J, P$  need confirmation.Seen in  $D^*(2010)^+ K^0$ ,  $D^*(2007)^0 K^+$ , and  $D_s^+ \pi^+ \pi^-$ . Not seen in  $D^+ K^0$  or  $D^0 K^+$ .  $J^P = 1^+$  assignment strongly favored.

NODE=M121

NODE=M121

 **$D_{s1}(2536)^\pm$  MASS**

NODE=M121M

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M121M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2535.11 ± 0.06 OUR FIT</b>				
<b>2535.21 ± 0.28 OUR AVERAGE</b>				
2537.7 ± 0.5 ± 3.1	24	<sup>1</sup> ABLIKIM	19P BES3	4.6 $e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
2535.7 ± 0.6 ± 0.5	46	<sup>2</sup> ABAZOV	09G D0	$B_s^0 \rightarrow D_{s1}^- \mu^+ \nu_\mu X$
2534.78 ± 0.31 ± 0.40	182	AUBERT	08B BABR	$B \rightarrow \bar{D}^{(*)} D^* K$
2534.6 ± 0.3 ± 0.7	193	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$
2535.3 ± 0.7	92	<sup>3</sup> HEISTER	02B ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2534.2 ± 1.2	9	ASRATYAN	94 BEBC	$\nu N \rightarrow D^* K^0 X, D^{*0} K^\pm X$
2535 ± 0.6 ± 1	75	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2535.2 ± 0.5 ± 1.5	28	ALBRECHT	92R ARG	10.4 $e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2536.6 ± 0.7 ± 0.4		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.9 ± 0.6 ± 2.0		ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2534.1 ± 0.6	116	<sup>4</sup> AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2535.08 ± 0.01 ± 0.15	8038	<sup>5</sup> LEES	11B BABR	10.6 $e^+ e^- \rightarrow D^{*+} K_S^0 X$
2535.57 <sup>+0.44</sup> <sub>-0.41</sub> ± 0.10	236	<sup>6</sup> CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
2535.3 ± 0.2 ± 0.5	134	<sup>7</sup> ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*0} K^+ X$
2534.8 ± 0.6 ± 0.6	44	<sup>8</sup> ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535 ± 28		<sup>9</sup> ASRATYAN	88 HLBC	$\nu N \rightarrow D_s \gamma \gamma X$

NODE=M121M

OCCUR=2

NODE=M121M;LINKAGE=D

NODE=M121M;LINKAGE=AB

NODE=M121M;LINKAGE=HI

NODE=M121M;LINKAGE=AU

NODE=M121M;LINKAGE=LE

NODE=M121M;LINKAGE=CH

NODE=M121M;LINKAGE=A

NODE=M121M;LINKAGE=C

NODE=M121M;LINKAGE=B

<sup>1</sup> From a fit of the  $D_s^+$  recoil mass distribution with an incoherent sum of the  $S$ -wave and  $D$ -wave Breit-Wigner line shapes.<sup>2</sup> Using the  $D^*(2010)^\pm$  mass of  $2010.0 \pm 0.4$  MeV from PDG 06.<sup>3</sup> Calculated using  $m(D^*(2010)^\pm) = 2010.0 \pm 0.5$  MeV,  $m(D^*(2007)^0) = 2006.7 \pm 0.5$  MeV, and the mass difference below.<sup>4</sup> Systematic uncertainties not evaluated.<sup>5</sup> Calculated using the mass difference  $m(D_{s1}^+) - m(D^{*+})_{PDG}$  below and  $m(D^{*+})_{PDG} = 2010.25 \pm 0.14$  MeV. Assuming  $S$ -wave decay of the  $D_{s1}(2536)$  to  $D^{*+} K_S^0$ , using a Breit-Wigner line shape corresponding to  $L=0$ .<sup>6</sup> Calculated using the mass difference  $m(D_{s1}^+) - m(D^{*+})_{PDG}$  reported below and  $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$  MeV.<sup>7</sup> Calculated using  $m(D^*(2007)^0) = 2006.6 \pm 0.5$  MeV and the mass difference below.<sup>8</sup> Calculated using  $m(D^*(2010)^\pm) = 2010.1 \pm 0.6$  MeV and the mass difference below.<sup>9</sup> Not seen in  $D^* K$ . **$m_{D_{s1}(2536)^\pm} - m_{D_s^*(2111)}$** 

NODE=M121DM

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M121DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>422.9 ± 0.4 OUR FIT</b>			
<b>424 ± 28</b>	ASRATYAN	88 HLBC	$D_s^{*\pm} \gamma$

NODE=M121DM

 **$m_{D_{s1}(2536)^\pm} - m_{D^*(2010)^\pm}$** 

NODE=M121DN

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ ,  
and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M121DN

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>524.85±0.04 OUR FIT</b>				
<b>524.84±0.04 OUR AVERAGE</b>				
524.83±0.01±0.04	8038	<sup>10</sup> LEES	11B BABR	10.6 $e^+e^- \rightarrow D^{*+} K_S^0 X$
525.30 <sup>+0.44</sup> <sub>-0.41</sub> ±0.10	236 ± 30	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
525.3 ±0.6 ±0.1	41	HEISTER 02B	ALEP	$e^+e^- \rightarrow D^{*+} K^0 X$
524.7 ±0.6 ±0.2	44	ALEXANDER93	CLE2	$e^+e^- \rightarrow D^{*+} K_S^0 X$
<sup>10</sup> Assuming S-wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$ , using a Breit-Wigner line shape corresponding to L=0.				

NODE=M121DN

OCCUR=2

NODE=M121DN;LINKAGE=LE

### $m_{D_{s1}(2536)^\pm} - m_{D^*(2007)^0}$

The fit includes  $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$ ,  
and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

NODE=M121DP

NODE=M121DP

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>528.26±0.05 OUR FIT</b> Error includes scale factor of 1.2.				
<b>528.68±0.28 OUR AVERAGE</b>				
528.7 ±1.9 ±0.5	51	HEISTER	02B ALEP	$e^+e^- \rightarrow D^{*0} K^+ X$
527.3 ±2.2	29	ACKERSTAFF 97W	OPAL	$e^+e^- \rightarrow D^{*0} K^+ X$
528.7 ±0.2 ±0.2	134	ALEXANDER 93	CLE2	$e^+e^- \rightarrow D^{*0} K^+ X$

NODE=M121DP

OCCUR=2

### $D_{s1}(2536)^\pm$ WIDTH

NODE=M121W

NODE=M121W

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.92±0.05 OUR AVERAGE</b>				
1.7 ±1.2 ±0.6	24	<sup>11</sup> ABLIKIM	19P BES3	4.6 $e^+e^- \rightarrow D_s^+ \bar{D}^0 K^-$
0.92±0.03±0.04	8038	<sup>12</sup> LEES	11B BABR	10.6 $e^+e^- \rightarrow D^{*+} K_S^0 X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.75±0.23	116	<sup>13</sup> AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
< 2.5	95 193	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow$ $D_s^+ \pi^+ \pi^- X$
< 3.2	90 75	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
< 2.3	90	ALEXANDER	93 CLEO	$e^+e^- \rightarrow D^{*0} K^+ X$
< 3.9	90	ALBRECHT	92R ARG	10.4 $e^+e^- \rightarrow D^{*0} K^+ X$
< 5.44	90	AVERY	90 CLEO	$e^+e^- \rightarrow D^{*+} K^0 X$
< 4.6	90	ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$

<sup>11</sup> From a fit of the  $D_s^+$  recoil mass distribution with an incoherent sum of the S-wave and S-wave Breit-Wigner line shapes.

<sup>12</sup> Assuming S-wave decay of the  $D_{s1}(2536)$  to  $D^{*+} K_S^0$ , using a Breit-Wigner line shape corresponding to L=0.

<sup>13</sup> Systematic uncertainties not evaluated.

NODE=M121W;LINKAGE=A

NODE=M121W;LINKAGE=LE

NODE=M121W;LINKAGE=AU

### $D_{s1}(2536)^+$ DECAY MODES

NODE=M121215;NODE=M121

Branching fractions are given relative to the one **DEFINED AS 1**.  
 $D_{s1}(2536)^-$  modes are charge conjugates of the modes below.

NODE=M121

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $D^*(2010)^+ K^0$	0.85 ±0.12	DESIG=1
$\Gamma_2$ $(D^*(2010)^+ K^0)_{S-wave}$	0.61 ±0.09	DESIG=7
$\Gamma_3$ $(D^*(2010)^+ K^0)_{D-wave}$		DESIG=9
$\Gamma_4$ $D^+ \pi^- K^+$	0.028±0.005	DESIG=8
$\Gamma_5$ $D^*(2007)^0 K^+$	<b>DEFINED AS 1</b>	DESIG=4
$\Gamma_6$ $D^+ K^0$	<0.34	90% DESIG=2
$\Gamma_7$ $D^0 K^+$	<0.12	90% DESIG=5
$\Gamma_8$ $D_s^{*+} \gamma$	possibly seen	DESIG=3
$\Gamma_9$ $D_s^+ \pi^+ \pi^-$	seen	DESIG=6

$D_{s1}(2536)^+$  BRANCHING RATIOS

NODE=M121220

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^*(2010)^+ K^0)$

 $\Gamma_5/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.18±0.16 OUR AVERAGE</b>				
0.88±0.24±0.08	116	AUSHEV 11	BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2.3 ±0.6 ±0.3	236 ± 30	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
1.32±0.47±0.23	92	<sup>14</sup> HEISTER 02B	ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
1.9 $\begin{smallmatrix} +1.1 \\ -0.9 \end{smallmatrix}$ ±0.4	35	<sup>14</sup> ACKERSTAFF 97W	OPAL	$e^+ e^- \rightarrow D^{*0} K^+ X,$ $D^{*+} K^0 X$
1.1 ±0.3		ALEXANDER 93	CLEO	$e^+ e^- \rightarrow$ $D^{*0} K^+ X, D^{*+} K^0 X$
1.4 ±0.3 ±0.2		<sup>15</sup> ALBRECHT 92R	ARG	10.4 $e^+ e^- \rightarrow$ $D^{*0} K^+ X, D^{*+} K^0 X$

NODE=M121R6  
NODE=M121R6<sup>14</sup> Ratio of the production rates measured in  $Z^0$  decays.<sup>15</sup> Evaluated by us from published inclusive cross-sections.NODE=M121R6;LINKAGE=6A  
NODE=M121R6;LINKAGE=A

$\Gamma((D^*(2010)^+ K^0)_{S-wave})/\Gamma(D^*(2010)^+ K^0)$

 $\Gamma_2/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.72±0.05±0.01</b>	5485	BALAGURA 08	BELL	10.6 $e^+ e^- \rightarrow D^{*+} K^0 X$

NODE=M121R8  
NODE=M121R8

$\Gamma(D^+ \pi^- K^+)/\Gamma(D^*(2010)^+ K^0)$

 $\Gamma_4/\Gamma_1$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.27±0.18±0.37</b>	1264	BALAGURA 08	BELL	10.6 $e^+ e^- \rightarrow D^+ \pi^- K^+ X$

NODE=M121R9  
NODE=M121R9

$\Gamma(D^+ K^0)/\Gamma(D^*(2010)^+ K^0)$

 $\Gamma_6/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.40</b>	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.43	90	ALBRECHT 89E	ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$

NODE=M121R1  
NODE=M121R1

$\Gamma(D^0 K^+)/\Gamma(D^*(2007)^0 K^+)$

 $\Gamma_7/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.12</b>	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121R4  
NODE=M121R4

$\Gamma(D_s^{*+} \gamma)/\Gamma_{total}$

 $\Gamma_8/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	ASRATYAN 88	HLBC	$\nu N \rightarrow D_s \gamma \gamma X$

NODE=M121R3  
NODE=M121R3

$\Gamma(D_s^{*+} \gamma)/\Gamma(D^*(2007)^0 K^+)$

 $\Gamma_8/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.42</b>	90	ALEXANDER 93	CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121R5  
NODE=M121R5

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma_{total}$

 $\Gamma_9/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AUBERT 06P	BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$

NODE=M121R7  
NODE=M121R7 $D_{s1}(2536)^\pm$  REFERENCES

NODE=M121

ABLIKIM 19P	CP C43 031001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59617
AUSHEV 11	PR D83 051102	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=16505
LEES 11B	PR D83 072003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16773
ABAZOV 09G	PRL 102 051801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52652
CHEKANOV 09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
AUBERT 08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BALAGURA 08	PR D77 032001	V. Balagura <i>et al.</i>	(BELLE Collab.)	REFID=52133
AUBERT 06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
PDG 06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
HEISTER 02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ACKERSTAFF 97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN 94	ZPHY C61 563	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)	REFID=43667
FRABETTI 94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
ALEXANDER 93	PL B303 377	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=43316
ALBRECHT 92R	PL B297 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=43179
AVERY 90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT 89E	PL B230 162	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40914
ASRATYAN 88	ZPHY C40 483	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=40916

NODE=M148

# $D_{s2}^*(2573)$

$$I(J^P) = 0(2^+)$$

## $D_{s2}^*(2573)$ MASS


NODE=M148M

NODE=M148M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2569.1 ±0.8</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 2.4. See the ideogram below.		
2570.7 ±2.0 ±1.7	62	<sup>1</sup> ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
2568.39±0.29±0.26		AAIJ	14AW LHCb	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
2569.4 ±1.6 ±0.5	82	AAIJ	11A LHCb	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$
2572.2 ±0.3 ±1.0		AUBERT,BE	06E BABR	$e^+ e^- \rightarrow D K X$
2574.5 ±3.3 ±1.6		ALBRECHT	96 ARG	$e^+ e^- \rightarrow D^0 K^+ X$
2573.2 $\begin{smallmatrix} +1.7 \\ -1.6 \end{smallmatrix}$ ±0.9	217	KUBOTA	94 CLE2	$e^+ e^- \sim 10.5 \text{ GeV}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2570.0 ±4.3	25	<sup>2</sup> EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$
2568.6 ±3.2	64	<sup>3</sup> HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$

- <sup>1</sup> From a fit of the  $D_s^+$  recoil mass distribution .
- <sup>2</sup> Not independent of the mass difference below.
- <sup>3</sup> Calculated using  $m_{D^0} = 1864.5 \pm 0.5 \text{ MeV}$  and the mass difference below.

NODE=M148M;LINKAGE=A  
 NODE=M148M;LINKAGE=EV  
 NODE=M148M;LINKAGE=HI



## $m_{D_{s2}^*(2573)} - m_{D^0}$

NODE=M148DM

NODE=M148DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>704 ±3 ±1</b>	64	HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
705.4±4.3	25	<sup>1</sup> EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$
<sup>1</sup> Systematic errors not estimated.				

NODE=M148DM;LINKAGE=EV

## $D_{s2}^*(2573)$ WIDTH

NODE=M148W

NODE=M148W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.9±0.7</b>	<b>OUR AVERAGE</b>			
17.2±3.6±1.1	62	<sup>1</sup> ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
16.9±0.5±0.6		AAIJ	14AW LHCb	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
12.1±4.5±1.6	82	AAIJ	11A LHCb	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$

27.1±0.6±5.6	AUBERT,BE	06E	BABR	$e^+e^- \rightarrow DKX$	
10.4±8.3±3.0	ALBRECHT	96	ARG	$e^+e^- \rightarrow D^0 K^+ X$	
16 $\begin{smallmatrix} +5 \\ -4 \end{smallmatrix}$ ±3	217	KUBOTA	94	CLE2	$e^+e^- \sim 10.5$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 $\begin{smallmatrix} +9 \\ -6 \end{smallmatrix}$	25	<sup>2</sup> EVDOKIMOV	04	SELX	600 $\Sigma^- A \rightarrow D^0 K^+ X$
---	----	------------------------	----	------	--

<sup>1</sup> From a fit of the  $D_s^+$  recoil mass distribution .

<sup>2</sup> Systematic errors not estimated.

NODE=M148W;LINKAGE=A  
NODE=M148W;LINKAGE=EV

### $D_{s2}^*(2573)^+$ DECAY MODES

$D_{s2}^*(2573)^-$  modes are charge conjugates of the modes below.

NODE=M148215;NODE=M148

NODE=M148

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^0 K^+$	seen
$\Gamma_2$ $D^*(2007)^0 K^+$	not seen
$\Gamma_3$ $D^+ K_S^0$	seen
$\Gamma_4$ $D^{*+} K_S^0$	seen

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=5;OUR EVAL;→ UNCHECKED ←

### $D_{s2}^*(2573)^+$ BRANCHING RATIOS

NODE=M148220

$\Gamma(D^0 K^+)/\Gamma_{\text{total}}$						$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>seen</b>	217	KUBOTA	94	CLE2	$\pm$ $e^+e^- \sim 10.5$ GeV	

NODE=M148R2  
NODE=M148R2

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^0 K^+)$						$\Gamma_2/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<b>&lt;0.33</b>	90	KUBOTA	94	CLE2	$+$ $e^+e^- \sim 10.5$ GeV	

NODE=M148R1  
NODE=M148R1

$\Gamma(D^{*+} K_S^0)/\Gamma(D^+ K_S^0)$						$\Gamma_4/\Gamma_3$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>0.044±0.005±0.011</b>	2000	<sup>1</sup> AAIJ	16AW	LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV	

NODE=M148R00  
NODE=M148R00

<sup>1</sup> First observation of the  $D_{s2}^*(2573)^+ \rightarrow D^{*+} K_S^0$  decay with a significance of 6.9  $\sigma$ .

NODE=M148R00;LINKAGE=A

### $D_{s2}^*(2573)$ REFERENCES

NODE=M148

ABLIKIM	19P	CP C43 031001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59617
AAIJ	16AW	JHEP 1602 133	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60464
AAIJ	14AW	PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=56105
AAIJ	11A	PL B698 14	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=16665
AUBERT,BE	06E	PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51512
EVDOKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)	REFID=50337
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ALBRECHT	96	ZPHY C69 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44631
KUBOTA	94	PRL 72 1972	Y. Kubota <i>et al.</i>	(CLEO Collab.)	REFID=43781



$D_{s0}(2590)^+$ 

$I(J^P) = 0(0^-)$

OMITTED FROM SUMMARY TABLE

 $D_{s0}(2590)^+$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2591±6±7</b>	444	<sup>1</sup> AAIJ	21A LHCb	$B^0 \rightarrow D^-(D^+K^+\pi^-)$

<sup>1</sup> The mass is calculated from the position of the T-matrix pole $D_{s0}(2590)^+$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>89±16±12</b>	444	<sup>1</sup> AAIJ	21A LHCb	$B^0 \rightarrow D^-(D^+K^+\pi^-)$

<sup>1</sup> The width is calculated from the position of the T-matrix pole $D_{s0}(2590)^+$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad D^+K^+\pi^-$	seen

$\Gamma(D^+K^+\pi^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	444	AAIJ	21A LHCb	$B^0 \rightarrow D^-(D^+K^+\pi^-)$

 $D_{s0}(2590)^+$  REFERENCESAAIJ 21A PRL 126 122002 R. Aaij *et al.* (LHCb Collab.) $D_{s1}^*(2700)^\pm$ 

$I(J^P) = 0(1^-)$

 $D_{s1}^*(2700)^+$  MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2714 ± 5 OUR AVERAGE</b>				Error includes scale factor of 1.5. See the ideogram below.
2732.3± 4.3± 5.8	15.7k	AAIJ	16AW LHCb	$pp \rightarrow D^{*+}K_S^0 X$ at 7, 8 TeV
2699 $^{+14}_{-7}$		<sup>1</sup> LEES	15C BABR	$B \rightarrow DD^0K^+$
2709.2± 1.9± 4.5	52k	<sup>2</sup> AAIJ	12AU LHCb	$pp \rightarrow (DK)^+X$ at 7 TeV
2710 ± 2 $^{+12}_{-7}$	10.4k	<sup>3</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$
2708 ± 9 $^{+11}_{-10}$	182	BRODZICKA	08 BELL	$B^+ \rightarrow D^0\bar{D}^0K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2694 ± 8 $^{+13}_{-3}$		LEES	15C BABR	$B^0 \rightarrow D^-D^0K^+$
2707 ± 8 ± 8		LEES	15C BABR	$B^+ \rightarrow \bar{D}^0D^0K^+$
2688 ± 4 ± 3		<sup>4</sup> AUBERT,BE	06E BABR	10.6 $e^+e^- \rightarrow DKX$

<sup>1</sup> From a combined analysis of  $B^0 \rightarrow D^-D^0K^+$  and  $B^+ \rightarrow \bar{D}^0D^0K^+$ .<sup>2</sup> From the combined fit of the  $D^+K_S^0$  and  $D^0K^+$  modes in the model including the  $D_{s2}^*(2573)^+$ ,  $D_{s1}^*(2700)^+$  and spin-0  $D_{sJ}^*(2860)^+$ .<sup>3</sup> From simultaneous fits to the two  $DK$  mass spectra and to the total  $D^*K$  mass spectrum.<sup>4</sup> Superseded by AUBERT 09AR.

NODE=M256

NODE=M256M

NODE=M256M

NODE=M256M;LINKAGE=B

NODE=M256W

NODE=M256W

NODE=M256W;LINKAGE=B

NODE=M256215;NODE=M256

DESIG=1

NODE=M256R01  
NODE=M256R01

NODE=M256

REFID=61092

NODE=M182

NODE=M182M

NODE=M182M

OCCUR=2


OCCUR=3

NODE=M182M;LINKAGE=B

NODE=M182M;LINKAGE=AA

NODE=M182M;LINKAGE=AB

NODE=M182M;LINKAGE=AU



**$D_{s1}^{*+}(2700)^+$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>122 ±10 OUR AVERAGE</b>				
136 ±19 ±24	15.7k	AAIJ	16AW LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
127 <sup>+24</sup> / <sub>-19</sub>		<sup>1</sup> LEES	15C BABR	$B \rightarrow D D^0 K^+$
115.8 ± 7.3 ±12.1	52k	<sup>2</sup> AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
149 ± 7 <sup>+39</sup> / <sub>-52</sub>	10.4k	<sup>3</sup> AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$
108 ±23 <sup>+36</sup> / <sub>-31</sub>	182	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

145 ±24 <sup>+22</sup> / <sub>-14</sub>		LEES	15C BABR	$B^0 \rightarrow D^- D^0 K^+$
113 ±21 <sup>+20</sup> / <sub>-16</sub>		LEES	15C BABR	$B^+ \rightarrow \bar{D}^0 D^0 K^+$
112 ± 7 ±36		<sup>4</sup> AUBERT,BE	06E BABR	10.6 $e^+ e^- \rightarrow D K X$

<sup>1</sup> From a combined analysis of  $B^0 \rightarrow D^- D^0 K^+$  and  $B^+ \rightarrow \bar{D}^0 D^0 K^+$ .

<sup>2</sup> From the combined fit of the  $D^+ K_S^0$  and  $D^0 K^+$  modes in the model including the  $D_{s2}^{*+}(2573)^+$ ,  $D_{s1}^{*+}(2700)^+$  and spin-0  $D_{sJ}^{*+}(2860)^+$ .

<sup>3</sup> From simultaneous fits to the two  $DK$  mass spectra and to the total  $D^* K$  mass spectrum.

<sup>4</sup> Superseded by AUBERT 09AR.

**$D_{s1}^{*+}(2700)^+$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $DK$	
$\Gamma_2$ $D^0 K^+$	seen
$\Gamma_3$ $D^+ K_S^0$	seen
$\Gamma_4$ $D^* K$	
$\Gamma_5$ $D^{*0} K^+$	seen
$\Gamma_6$ $D^{*+} K_S^0$	seen

**$D_{s1}^{*+}(2700)^+$  BRANCHING RATIOS**

$\Gamma(D^* K)/\Gamma(DK)$	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma_1$
<b>0.91±0.13±0.12</b>	10.4k	<sup>1</sup> AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$	

<sup>1</sup> From the average of the corresponding ratios with  $D^{(*)0} K^+$  and  $D^{(*)+} K_S^0$ .

NODE=M182W

NODE=M182W

OCCUR=2

OCCUR=3

NODE=M182W;LINKAGE=A

NODE=M182W;LINKAGE=AA

NODE=M182W;LINKAGE=AB

NODE=M182W;LINKAGE=AU

NODE=M182215;NODE=M182

DESIG=2

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

NODE=M182225

NODE=M182R01

NODE=M182R01

NODE=M182R01;LINKAGE=AU

$\Gamma(D^{*0}K^+)/\Gamma(D^0K^+)$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma_2$
<b>0.88±0.14±0.14</b>	7716	<sup>1</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$	

<sup>1</sup> From the  $D^{*0}K^+$  and  $D^0K^+$ , where  $D^{*0} \rightarrow D^0\pi^0$ .

NODE=M182R02  
NODE=M182R02

NODE=M182R02;LINKAGE=AU

 $\Gamma(D^{*+}K_S^0)/\Gamma(D^+K_S^0)$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma_3$
<b>1.14±0.39±0.23</b>	2700	<sup>1</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$	

<sup>1</sup> From the  $D^{*+}K_S^0$  and  $D^+K_S^0$ , where  $D^{*+} \rightarrow D^+\pi^0$ .

NODE=M182R03  
NODE=M182R03

NODE=M182R03;LINKAGE=AU

 $D_{s1}^*(2700)^\pm$  REFERENCES

AAIJ	16AW JHEP 1602 133	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60464
LEES	15C PR D91 052002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56412
AAIJ	12AU JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54735
AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53135
BRODZICKA	08 PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)	REFID=52144
AUBERT,BE	06E PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51512

NODE=M182

NODE=M196

$D_{s1}^*(2860)^\pm$

$$I(J^P) = 0(1^-)$$

OMITTED FROM SUMMARY TABLE

$J^P$  consistent with  $1^-$  from angular analysis of AAIJ 14AW.

NODE=M196

 $D_{s1}^*(2860)^+$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2859±12±24</b>	<sup>1</sup> AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$

<sup>1</sup> Separated from the spin-3 component  $D_{s3}^*(2860)^-$  by a fit of the helicity angle of the  $\bar{D}^0 K^-$  system, with a statistical significance of the spin-3 and spin-1 components in excess of  $10\sigma$ .

NODE=M196M

NODE=M196M

NODE=M196M;LINKAGE=A

 $D_{s1}^*(2860)^+$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>159±23±77</b>	<sup>1</sup> AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$

<sup>1</sup> Separated from the spin-3 component  $D_{s3}^*(2860)^-$  by a fit of the helicity angle of the  $\bar{D}^0 K^-$  system, with a statistical significance of the spin-3 and spin-1 components in excess of  $10\sigma$ .

NODE=M196W

NODE=M196W

NODE=M196W;LINKAGE=A

 $D_{s1}^*(2860)^\pm$  REFERENCES

AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=56105
------	---------------------	-----------------------	-------------------	-------------

NODE=M196

REFID=56105

NODE=M226

 $D_{s3}^*(2860)^\pm$ 

$$I(J^P) = 0(3^-)$$

$J^P$  consistent with  $3^-$  from angular analysis of AAIJ 14AW. Observed by AUBERT, BE 06E and AUBERT 09AR in inclusive production of  $DK$  and  $D^*K$  in  $e^+e^-$  annihilation.

NODE=M226

 **$D_{s3}^*(2860)^+$  MASS**

NODE=M226M

NODE=M226M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2860.5±2.6±6.5</b>		<sup>1</sup> AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2867.1±4.3±1.9	3.1k	AAIJ	16AW LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
2866.1±1.0±6.3	36k	<sup>2,3</sup> AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
2862 ±2 ± $\frac{+5}{-2}$	3122	<sup>2,4</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)} K X$
2856.6±1.5±5.0		<sup>5</sup> AUBERT, BE	06E BABR	$e^+e^- \rightarrow DK X$

NODE=M226M;LINKAGE=A

<sup>1</sup> Separated from the spin-1 component  $D_{s1}^*(2860)^-$  by a fit of the helicity angle of the  $\bar{D}^0 K^-$  system, with a statistical significance of the spin-3 and spin-1 components in excess of  $10\sigma$ .

<sup>2</sup> Possible contribution from the  $D_{s1}^*(2860)$  state.

<sup>3</sup> From the combined fit of the  $D^+ K_S^0$  and  $D^0 K^+$  modes in the model including the  $D_{s2}^*(2573)^+$ ,  $D_{s1}^*(2700)^+$  and spin-0  $D_{s,J}^*(2860)^+$ .

<sup>4</sup> From simultaneous fits to the two  $DK$  mass spectra and to the total  $D^*K$  mass spectrum.

<sup>5</sup> Superseded by AUBERT 09AR.

NODE=M226M;LINKAGE=D

NODE=M226M;LINKAGE=E

NODE=M226M;LINKAGE=C

NODE=M226M;LINKAGE=B

 **$D_{s3}^*(2860)^+$  WIDTH**

NODE=M226W

NODE=M226W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>53 ± 7 ± 7</b>		<sup>1</sup> AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
50 ±11 ±13	3.1k	AAIJ	16AW LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
69.9± 3.2± 6.6	36k	<sup>2,3</sup> AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
48 ± 3 ± 6	3122	<sup>2,4</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)} K X$
47 ± 7 ±10		<sup>5</sup> AUBERT, BE	06E BABR	$e^+e^- \rightarrow DK X$

NODE=M226W;LINKAGE=A

<sup>1</sup> Separated from the spin-1 component  $D_{s1}^*(2860)^-$  by a fit of the helicity angle of the  $\bar{D}^0 K^-$  system, with a statistical significance of the spin-3 and spin-1 components in excess of  $10\sigma$ .

<sup>2</sup> Possible contribution from the  $D_{s1}^*(2860)$  state.

<sup>3</sup> From the combined fit of the  $D^+ K_S^0$  and  $D^0 K^+$  modes in the model including the  $D_{s2}^*(2573)^+$ ,  $D_{s1}^*(2700)^+$  and spin-0  $D_{s,J}^*(2860)^+$ .

<sup>4</sup> From simultaneous fits to the two  $DK$  mass spectra and to the total  $D^*K$  mass spectrum.

<sup>5</sup> Superseded by AUBERT 09AR.

NODE=M226W;LINKAGE=D

NODE=M226W;LINKAGE=E

NODE=M226W;LINKAGE=C

NODE=M226W;LINKAGE=B

 **$D_{s3}^*(2860)^\pm$  DECAY MODES**

NODE=M226215;NODE=M226

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $DK$	
$\Gamma_2$ $D^0 K^+$	seen
$\Gamma_3$ $D^+ K_S^0$	seen
$\Gamma_4$ $D^* K$	
$\Gamma_5$ $D^{*0} K^+$	seen
$\Gamma_6$ $D^{*+} K_S^0$	seen

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

$D_{s3}^*(2860)^\pm$  BRANCHING RATIOS $\Gamma(D^*K)/\Gamma(DK)$  $\Gamma_4/\Gamma_1$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10 \pm 0.15 \pm 0.19$	3122	<sup>1</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$

<sup>1</sup> From the average of the corresponding ratios with  $D^{(*)0}K^+$  and  $D^{(*)+}K_S^0$ .

NODE=M226225

NODE=M226R01  
NODE=M226R01

NODE=M226R01;LINKAGE=AU

 $\Gamma(D^{*0}K^+)/\Gamma(D^0K^+)$  $\Gamma_5/\Gamma_2$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.04 \pm 0.17 \pm 0.20$	2241	<sup>1</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$

<sup>1</sup> From the  $D^{*0}K^+$  and  $D^0K^+$ , where  $D^{*0} \rightarrow D^0\pi^0$ .

NODE=M226R02  
NODE=M226R02

NODE=M226R02;LINKAGE=AU

 $\Gamma(D^{*+}K_S^0)/\Gamma(D^+K_S^0)$  $\Gamma_6/\Gamma_3$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.38 \pm 0.35 \pm 0.49$	881	<sup>1</sup> AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$

<sup>1</sup> From the  $D^{*+}K_S^0$  and  $D^+K_S^0$ , where  $D^{*+} \rightarrow D^+\pi^0$ .

NODE=M226R03  
NODE=M226R03

NODE=M226R03;LINKAGE=AU

 $D_{s3}^*(2860)^\pm$  REFERENCES

AAIJ	16AW JHEP 1602 133	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	12AU JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06E PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M226

REFID=60464  
REFID=56105  
REFID=54735  
REFID=53135  
REFID=51512

NODE=M197

 $D_{sJ}(3040)^\pm$  $I(J^P) = 0(?^?)$ 

OMITTED FROM SUMMARY TABLE

Observed by AUBERT 09AR in inclusive production of  $D^*K$  in  $e^+e^-$  annihilation.

NODE=M197

 $D_{sJ}(3040)^+$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$3044 \pm 8 \pm 30$ $-5$	AUBERT	09AR BABR	$e^+e^- \rightarrow D^*KX$

NODE=M197M

NODE=M197M

 $D_{sJ}(3040)^+$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$239 \pm 35 \pm 46$ $-42$	AUBERT	09AR BABR	$e^+e^- \rightarrow D^*KX$

NODE=M197W

NODE=M197W

 $D_{sJ}(3040)^\pm$  DECAY MODES

Mode		
$\Gamma_1$	$D^*K$	DESIG=1
$\Gamma_2$	$D^{*0}K^+$	DESIG=2
$\Gamma_3$	$D^{*+}K_S^0$	DESIG=3

NODE=M197215;NODE=M197

 $D_{sJ}(3040)^\pm$  REFERENCES

AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
--------	--------------------	-------------------------	-----------------

NODE=M197

REFID=53135

# BOTTOM MESONS

## ( $B = \pm 1$ )

$$B^+ = u\bar{b}, B^0 = d\bar{b}, \bar{B}^0 = \bar{d}b, B^- = \bar{u}b, \text{ similarly for } B^{*'}\text{'s}$$

### $B_1(5721)$

$$I(J^P) = \frac{1}{2}(1^+)$$

$I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

### $B_1(5721)$ MASS

#### $B_1(5721)^+$ mass

OUR FIT uses  $m_{B^{*0}}$  and  $m_{B_1^+} - m_{B^{*0}}$  to determine  $m_{B_1(5721)^+}$ .

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**5725.9<sup>+2.5</sup><sub>-2.7</sub> OUR FIT**

#### $m_{B_1^+} - m_{B^{*0}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**401.2<sup>+2.4</sup><sub>-2.7</sub> OUR FIT**

**401.2<sup>+2.4</sup><sub>-2.7</sub> OUR AVERAGE**

400.5 $\pm$ 1.8 $\pm$ 3.1	8k	<sup>1</sup> AAIJ	15AB	LHCB $p\bar{p}$ at 7, 8 TeV
402 $\pm$ 3 <sup>+1</sup> <sub>-3</sub>		<sup>2</sup> AALTONEN	14l	CDF $p\bar{p}$ at 1.96 TeV

<sup>1</sup> AAIJ 15AB reports  $[m_{B_1^+} - m_{B^0}] - (m_{B^{*0}} - m_{B^0}) - m_{\pi^+} = 260.9 \pm 1.8 \pm 3.1$

MeV which we adjust by the  $\pi^+$  mass and assume  $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV. The masses inside the square brackets were measured for each candidate event.

<sup>2</sup> AALTONEN 14l reports  $m_{B_1(5721)^+} - m_{B^{*0}} - m_{\pi^+} = 262 \pm 3^{+1}_{-3}$  MeV which we adjusted by the  $\pi^+$  mass.

#### $B_1(5721)^0$ mass

OUR FIT uses mass differences measurements listed below to determine the mass

$$m_{B_1(5721)^0}$$

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**5726.1 $\pm$ 1.3 OUR FIT** Error includes scale factor of 1.2.

#### $m_{B_1^0} - m_{B^+}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**446.7 $\pm$ 1.3 OUR FIT** Error includes scale factor of 1.2.

**441.5 $\pm$ 2.4 $\pm$ 1.3** <sup>1</sup> ABAZOV 07T D0  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

446.2 <sup>+1.9+1.0</sup> <sub>-2.1-1.2</sub>	<sup>1</sup> AALTONEN	09D	CDF	Repl. by AALTONEN 14l
---	-----------------------	-----	-----	-----------------------

<sup>1</sup> Observed in  $B_1^0 \rightarrow B^{*+} \pi^-$ .

#### $m_{B_1^0} - m_{B^{*+}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**401.4 $\pm$ 1.2 OUR FIT** Error includes scale factor of 1.2.

**402.8 $\pm$ 1.1 OUR AVERAGE**

403.4 $\pm$ 0.7 $\pm$ 1.5	35k	<sup>1</sup> AAIJ	15AB	LHCB $p\bar{p}$ at 7, 8 TeV
402.3 $\pm$ 0.9 <sup>+1.1</sup> <sub>-1.2</sub>		<sup>2</sup> AALTONEN	14l	CDF $p\bar{p}$ at 1.96 TeV

<sup>1</sup> AAIJ 15AB reports  $[m_{B_1^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 263.9 \pm 0.7 \pm 1.4$

MeV which we adjust by the  $\pi^-$  mass and  $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV. The masses inside the square brackets were measured for each candidate event.

<sup>2</sup> AALTONEN 14l reports  $m_{B_1(5721)^0} - m_{B^{*+}} - m_{\pi^-} = 262.7 \pm 0.9^{+1.1}_{-1.2}$  MeV which we adjusted by the  $\pi^-$  mass.

NODE=MXXX045

NODE=MXXX045

NODE=M244

NODE=M244

NODE=M244205

NODE=M244M+

NODE=M244M+  
NODE=M244M+NODE=M244DM+  
NODE=M244DM+

NODE=M244DM+;LINKAGE=A

NODE=M244DM+;LINKAGE=AA

NODE=M244M0  
NODE=M244M0

NODE=M244M0

NODE=M244DM0  
NODE=M244DM0

NODE=M244DM0;LINKAGE=AA

NODE=M244DM1  
NODE=M244DM1

NODE=M244DM1;LINKAGE=B

NODE=M244DM1;LINKAGE=AA

**$B_1(5721)$  WIDTH**

NODE=M244210

 **$B_1(5721)^+$  width**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>31 \pm 6</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.1.		
29.1 $\pm$ 3.6 $\pm$ 4.3	8k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
49 $^{+12}_{-10}$ $^{+2}_{-13}$		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M244W+  
NODE=M244W+ **$B_1(5721)^0$  width**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>27.5 \pm 3.4</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.1.		
30.1 $\pm$ 1.5 $\pm$ 3.5	35k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
23 $\pm$ 3 $\pm$ 4		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M244W0  
NODE=M244W0 **$B_1(5721)$  DECAY MODES**

NODE=M244215;NODE=M244

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^* \pi$	seen

DESIG=1

 **$\Gamma(B^* \pi)/\Gamma_{\text{total}}$**  **$\Gamma_1/\Gamma$** 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	AAIJ	15AB LHCB	$\pm 0$	$p\bar{p}$ at 7, 8 TeV
<b>seen</b>	AALTONEN	14I CDF	$\pm$	$p\bar{p}$ at 1.96 TeV
seen	AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV
<b>seen</b>	<sup>1</sup> ABZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV

NODE=M244R01  
NODE=M244R01

OCCUR=2

<sup>1</sup> Observed in  $B_1^0 \rightarrow B^{*+} \pi^-$  with  $B^{*+} \rightarrow B^+ \gamma$  and  $B^+ \rightarrow J/\psi \pi^+$ .

NODE=M244R01;LINKAGE=AB

 **$B_1(5721)$  REFERENCES**

NODE=M244

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=56628  
REFID=56029  
REFID=52700  
REFID=52014

NODE=M151

 **$B_J^*(5732)$** 

$$I(J^P) = ?(??)$$

OMITTED FROM SUMMARY TABLE  
also known as  $B^{**}$ 

Signal can be interpreted as stemming from several narrow and broad resonances.

NODE=M151

 **$B_J^*(5732)$  MASS**

NODE=M151M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5698 \pm 8</math></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.		
5710 $\pm$ 20		<sup>1</sup> AFFOLDER	01F CDF	$p\bar{p}$ at 1.8 TeV
5695 $^{+17}_{-19}$		<sup>2</sup> BARATE	98L ALEP	$e^+ e^- \rightarrow Z$
5704 $\pm$ 4 $\pm$ 10	1944	<sup>3</sup> BUSKULIC	96D ALEP	$E_{\text{cm}}^{ee} = 88-94$ GeV
5732 $\pm$ 5 $\pm$ 20	2157	ABREU	95B DLPH	$E_{\text{cm}}^{ee} = 88-94$ GeV
5681 $\pm$ 11	1738	AKERS	95E OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M151M

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

5713 $\pm$  2 <sup>4</sup> ACCIARRI 99N L3  $e^+ e^- \rightarrow Z$ <sup>1</sup> AFFOLDER 01F uses the reconstructed  $B$  meson through semileptonic decay channels. The fraction of light  $B$  mesons that are produced at  $L=1$   $B^{**}$  states is measured to be  $0.28 \pm 0.06 \pm 0.03$ .

NODE=M151M;LINKAGE=MF

<sup>2</sup> BARATE 98L uses fully reconstructed  $B$  mesons to search for  $B^{**}$  production in the  $B\pi^\pm$  system. In the framework of heavy quark symmetry (HQS), they also measured the mass of  $B_2^*$  to be  $5739^{+8+6}_{-11-4}$  MeV/ $c^2$  and the relative production rate of  $B(b \rightarrow B_2^* \rightarrow B^{(*)} \pi)/B(b \rightarrow B_{u,d}) = (31 \pm 9^{+6}_{-5})\%$ .

NODE=M151M;LINKAGE=B

<sup>3</sup> Using  $m_{B\pi} - m_B = 424 \pm 4 \pm 10$  MeV.

NODE=M151M;LINKAGE=A

<sup>4</sup> ACCIARRI 99N uses inclusive reconstructed  $B$  mesons to search for  $B^{**}$  production in the  $B^{(*)}\pi^{\pm}$  system. In the framework of HQET, they measured the mass of  $B_1^*$  and  $B_2^*$  to be  $5670 \pm 10 \pm 13$  MeV and  $5768 \pm 5 \pm 6$  with the  $B(b \rightarrow B^{**}) = (32 \pm 3 \pm 6) \times 10^{-2}$ . They also reported the evidence for the existence of an excited  $B$ -meson state or mixture of states in the region 5.9–6.0 GeV.

NODE=M151M;LINKAGE=N

 **$B_j^*(5732)$  WIDTH**

NODE=M151W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>128 ± 18 OUR AVERAGE</b>				
145 ± 28	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88-94$ GeV
116 ± 24	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88-94$ GeV

NODE=M151W

 **$B_j^*(5732)$  DECAY MODES**

NODE=M151215;NODE=M151

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^* \pi + B \pi$	seen
$\Gamma_2$ $B^* \pi (X)$	[a] (85 ± 29) %

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2

[a] X refers to decay modes with or without additional accompanying decay particles.

LINKAGE=151

 **$B_j^*(5732)$  BRANCHING RATIOS**

NODE=M151220

X refers to decay modes with or without additional accompanying decay particles.

NODE=M151220

$\Gamma(B^* \pi(X))/\Gamma_{total}$	$\Gamma_2/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>0.85^{+0.26}_{-0.27} \pm 0.12</math></b>	ABBIENDI	02E OPAL	$e^+ e^- \rightarrow Z$

NODE=M151R1  
NODE=M151R1 **$B_j^*(5732)$  REFERENCES**

NODE=M151

ABBIENDI	02E	EPJ C23 437	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48742
AFFOLDER	01F	PR D64 072002	T. Affolder <i>et al.</i>	(CDF Collab.)	REFID=48369
ACCIARRI	99N	PL B465 323	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47247
BARATE	98L	PL B425 215	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=46082
BUSKULIC	96D	ZPHY C69 393	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44677
ABREU	95B	PL B345 598	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44131
AKERS	95E	ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44182



**$B_2^*(5747)$** 

$$I(J^P) = \frac{1}{2}(2^+)$$

$I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M245

NODE=M245

 **$B_2^*(5747)$  MASS**

NODE=M245205

 **$B_2^*(5747)^+$  mass**

NODE=M245M+

OUR FIT uses  $m_{B^0}$  and  $m_{B_2^{*+}} - m_{B^0}$  to determine  $m_{B_2^*(5747)^+}$ .NODE=M245M+  
NODE=M245M+

VALUE (MeV)	DOCUMENT ID
<b>5737.2±0.7 OUR FIT</b>	

 **$m_{B_2^{*+}} - m_{B^0}$** NODE=M245DM+  
NODE=M245DM+

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>457.5 ±0.7 OUR FIT</b>				
<b>457.5 ±0.7 OUR AVERAGE</b>				
457.62±0.72±0.40	4k	<sup>1</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
457.3 ±1.3 $\begin{smallmatrix} +0.3 \\ -0.9 \end{smallmatrix}$		<sup>2</sup> AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

<sup>1</sup> AAIJ 15AB reports  $[m_{B_2^{*+}} - m_{B^0}] - m_{\pi^+} = 318.1 \pm 0.7 \pm 0.4$  MeV which we adjust by the  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event.

NODE=M245DM+;LINKAGE=B

<sup>2</sup> AALTONEN 14I reports  $m_{B_2^*(5747)^+} - m_{B^0} - m_{\pi^+} = 317.7 \pm 1.2 \begin{smallmatrix} +0.3 \\ -0.9 \end{smallmatrix}$  MeV which we adjusted by the  $\pi^+$  mass.

NODE=M245DM+;LINKAGE=A

 **$B_2^*(5747)^0$  mass**

NODE=M245M0

OUR FIT uses  $m_{B^+}$ ,  $m_{B_1^0} - m_{B^+}$ , and mass differences below to determine  $m_{B_2^*(5747)^0}$ . The  $-0.659$  correlation between statistical uncertainties of  $m_{B_1^0} - m_{B^+}$  and  $m_{B_2^{*0}} - m_{B^0}$  measurements reported by ABAZOV 07T is taken into account.

NODE=M245M0

VALUE (MeV)	DOCUMENT ID
<b>5739.5±0.7 OUR FIT</b>	Error includes scale factor of 1.4.

NODE=M245M0

 **$m_{B_2^{*0}} - m_{B_1^0}$** NODE=M245DM0  
NODE=M245DM0

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>13.4±1.4 OUR FIT</b>	Error includes scale factor of 1.3.		
<b>26.2±3.1±0.9</b>	<sup>1</sup> ABAZOV	07T D0	$p\bar{p}$ at 1.96 TeV
14.9 $\begin{smallmatrix} +2.2+1.2 \\ -2.5-1.4 \end{smallmatrix}$	<sup>1</sup> AALTONEN	09D CDF	Repl. by AALTONEN 14I

<sup>1</sup> Observed in  $B_2^{*0} \rightarrow B^{*+}\pi^-$  and  $B_2^{*0} \rightarrow B^+\pi^-$ .

NODE=M245DM0;LINKAGE=AB

 **$m_{B_2^{*0}} = m_{B^+}$** NODE=M245DM2  
NODE=M245DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>460.2 ±0.6 OUR FIT</b>	Error includes scale factor of 1.4.			
<b>459.9 ±0.8 OUR AVERAGE</b>	Error includes scale factor of 1.8.			
460.18±0.37±0.33	17k	<sup>1</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
457.5 ±1.2 $\begin{smallmatrix} +0.8 \\ -0.9 \end{smallmatrix}$		<sup>2</sup> AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

<sup>1</sup> AAIJ 15AB reports  $[m_{B_2^{*0}} - m_{B^+}] - m_{\pi^-} = 320.6 \pm 0.4 \pm 0.3$  MeV which we adjust by the  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event.

NODE=M245DM2;LINKAGE=A

<sup>2</sup> AALTONEN 14I reports  $m_{B_2^*(5747)^0} - m_{B^+} - m_{\pi^-} = 317.9 \pm 1.2 \begin{smallmatrix} +0.8 \\ -0.9 \end{smallmatrix}$  MeV which we adjusted by the  $\pi^-$  mass.

NODE=M245DM2;LINKAGE=AA

 **$B_2^*(5747)$  WIDTH**

NODE=M245210

 **$B_2^*(5747)^+$  width**NODE=M245W+  
NODE=M245W+

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20 ±5 OUR AVERAGE</b>	Error includes scale factor of 2.2.			
23.6±2.0±2.1	4k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
11 $\begin{smallmatrix} +4 +3 \\ -3 -4 \end{smallmatrix}$		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

**$B_2^*(5747)^0$  width**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>24.2±1.7 OUR AVERAGE</b>				
24.5±1.0± 1.5	17k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
22 $\begin{smallmatrix} +3 & +4 \\ -2 & -5 \end{smallmatrix}$		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
22.7 $\begin{smallmatrix} +3.8+ & 3.2 \\ -3.2- & 10.2 \end{smallmatrix}$		AALTONEN	09D CDF	Repl. by AALTONEN 14I

NODE=M245W0  
 NODE=M245W0

 **$B_2^*(5747)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B\pi$	seen
$\Gamma_2$ $B^*\pi$	seen

NODE=M245215;NODE=M245

DESIG=1  
 DESIG=2

$\Gamma(B\pi)/\Gamma_{\text{total}}$						$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	4k,17k	AAIJ	15AB LHCB	±0	$p\bar{p}$ at 7, 8 TeV	
<b>seen</b>		AALTONEN	14I CDF	±	$p\bar{p}$ at 1.96 TeV	
seen		AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV	
<b>seen</b>		ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV	

NODE=M245R01  
 NODE=M245R01

$\Gamma(B^*\pi)/\Gamma_{\text{total}}$						$\Gamma_2/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	4k,17k	AAIJ	15AB LHCB	±0	$p\bar{p}$ at 7, 8 TeV	
seen		AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV	
<b>seen</b>		ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV	

NODE=M245R02  
 NODE=M245R02

$\Gamma(B^*\pi)/\Gamma(B\pi)$						$\Gamma_2/\Gamma_1$
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>0.84±0.27 OUR AVERAGE</b>						
0.71±0.14±0.30	17k	AAIJ	15AB LHCB	0	$p\bar{p}$ at 7, 8 TeV	
1.0 ±0.5 ±0.8	4k	AAIJ	15AB LHCB	±	$p\bar{p}$ at 7, 8 TeV	
1.10±0.42±0.31		<sup>1</sup> ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV	

NODE=M245R03  
 NODE=M245R03

OCCUR=2

<sup>1</sup> Converted from measured ratio of  $R = B(B_2^{*0} \rightarrow B^{*+}\pi^-) / B(B_2^{*0} \rightarrow B^{(*)+}\pi^-)$   
 $= 0.475 \pm 0.095 \pm 0.069$ .

NODE=M245R03;LINKAGE=AB

 **$B_2^*(5747)$  REFERENCES**

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D	PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T	PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)

NODE=M245

REFID=56628  
 REFID=56029  
 REFID=52700  
 REFID=52014

**$B_J(5840)$** 

$$I(J^P) = \frac{1}{2}(?^?)$$

$I, J, P$  need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M246

NODE=M246

 **$B_J(5840)$  MASS**

NODE=M246205

 **$B_J(5840)^+$  MASS**

NODE=M246M+

OUR FIT uses  $m_{B^0}$  and  $m_{B_J(5840)^+} - m_{B^0}$  to determine  $m_{B_J(5840)^+}$ .NODE=M246M+  
NODE=M246M+

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**5851±19 OUR FIT** **$m_{B_J(5840)^+} - m_{B^0}$** NODE=M246DM+  
NODE=M246DM+

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**571±19 OUR FIT****571±13±14**      7k      <sup>1</sup> AAIJ      15AB LHCB       $pp$  at 7, 8 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

595±26±14      7k      <sup>2</sup> AAIJ      15AB LHCB       $pp$  at 7, 8 TeV

OCCUR=2

<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$  MeV which we adjust bythe  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses two relativistic Breit-Wigner functions in the fit for mass difference.<sup>2</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$  MeV which we adjust bythe  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM+;LINKAGE=A

NODE=M246DM+;LINKAGE=B

 **$m_{B_J(5840)^+} - m_{B^{*0}}$** NODE=M246DM1  
NODE=M246DM1

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

565±15±14      7k      <sup>1</sup> AAIJ      15AB LHCB       $pp$  at 7, 8 TeV<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$ MeV which we adjust by the  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = -(-1)^J$ ,  $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM1;LINKAGE=A

 **$B_J(5840)^0$  MASS**

NODE=M246M0

OUR FIT uses  $m_{B^+}$  and  $m_{B_J(5840)^0} - m_{B^+}$  to determine  $m_{B_J(5840)^0}$ .NODE=M246M0  
NODE=M246M0

VALUE (MeV)	DOCUMENT ID
-------------	-------------

**5863±9 OUR FIT** **$m_{B_J(5840)^0} - m_{B^+}$** NODE=M246DM0  
NODE=M246DM0

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**584± 9 OUR FIT****584± 5±7**      12k      <sup>1</sup> AAIJ      15AB LHCB       $pp$  at 7, 8 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

610±22±7      12k      <sup>2</sup> AAIJ      15AB LHCB       $pp$  at 7, 8 TeV

OCCUR=2

<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$  MeV which we adjust bythe  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses two relativistic Breit-Wigner functions in the fit for mass difference.<sup>2</sup> AAIJ 15AB reports  $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$  MeV which we adjust bythe  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM0;LINKAGE=A

NODE=M246DM0;LINKAGE=B

$m_{B_J(5840)^0} - m_{B^{*+}}$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$584 \pm 5 \pm 7$	12k	<sup>1</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
-------------------	-----	-------------------	-----------	------------------

<sup>1</sup> AAIJ 15AB reports  $[m_{B^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$  MeV

which we adjust by the  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = -(-1)^J$ ,  $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM2  
NODE=M246DM2

NODE=M246DM2;LINKAGE=A

 $B_J(5840)$  WIDTH

NODE=M246210

 $B_J(5840)^+$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

$224 \pm 24 \pm 80$	7k	<sup>1</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
---------------------	----	-------------------	-----------	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$215 \pm 27 \pm 80$	7k	<sup>2</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
---------------------	----	-------------------	-----------	------------------

$229 \pm 27 \pm 80$	7k	<sup>3</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
---------------------	----	-------------------	-----------	------------------

<sup>1</sup> Assuming  $P = (-1)^J$  and using two relativistic Breit-Wigner functions in the fit for mass difference.

<sup>2</sup> Assuming  $P = (-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

<sup>3</sup> Assuming  $P = -(-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246W+  
NODE=M246W+

OCCUR=2  
OCCUR=3

NODE=M246W+;LINKAGE=A

NODE=M246W+;LINKAGE=B

NODE=M246W+;LINKAGE=C

 $B_J(5840)^0$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

$127 \pm 17 \pm 34$	12k	<sup>1</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
---------------------	-----	-------------------	-----------	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$107 \pm 20 \pm 34$	12k	<sup>2</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
---------------------	-----	-------------------	-----------	------------------

$119 \pm 17 \pm 34$	12k	<sup>3</sup> AAIJ	15AB LHCB	$pp$ at 7, 8 TeV
---------------------	-----	-------------------	-----------	------------------

<sup>1</sup> Assuming  $P = (-1)^J$  and using two relativistic Breit-Wigner functions in the fit for mass difference.

<sup>2</sup> Assuming  $P = (-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

<sup>3</sup> Assuming  $P = -(-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246W0  
NODE=M246W0

OCCUR=2  
OCCUR=3

NODE=M246W0;LINKAGE=A

NODE=M246W0;LINKAGE=B

NODE=M246W0;LINKAGE=C

 $B_J(5840)$  DECAY MODES

NODE=M246215;NODE=M246

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^* \pi$	seen
$\Gamma_2$ $B \pi$	possibly seen

DESIG=1

DESIG=2

 $B_J(5840)$  BRANCHING RATIOS

NODE=M246220

$\Gamma(B^* \pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_1/\Gamma$
---	-------	------	-------------	------	-----	---------	-------------------

seen	7k	AAIJ	15AB LHCB	$\pm$		$pp$ at 7, 8 TeV	
------	----	------	-----------	-------	--	------------------	--

seen	12k	AAIJ	15AB LHCB	0		$pp$ at 7, 8 TeV	
------	-----	------	-----------	---	--	------------------	--

NODE=M246R01  
NODE=M246R01

OCCUR=2

$\Gamma(B \pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_2/\Gamma$
---------------------------------------	-------	------	-------------	------	-----	---------	-------------------

possibly seen	7k	<sup>1</sup> AAIJ	15AB LHCB	$\pm$		$pp$ at 7, 8 TeV	
---------------	----	-------------------	-----------	-------	--	------------------	--

possibly seen		<sup>1</sup> AAIJ	15AB LHCB	0		$pp$ at 7, 8 TeV	
---------------	--	-------------------	-----------	---	--	------------------	--

NODE=M246R02  
NODE=M246R02

OCCUR=2

<sup>1</sup> A  $B \pi$  decay is forbidden from a  $P = -(-1)^J$  parent, whereas  $B^* \pi$  is allowed.

NODE=M246R02;LINKAGE=A

 $B_J(5840)$  REFERENCES

NODE=M246

AAIJ	15AB JHEP 1504 024	R. Aaij et al.	(LHCb Collab.)
------	--------------------	----------------	----------------

REFID=56628

**$B_J(5970)$** 

$$I(J^P) = \frac{1}{2}(??)$$

$I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M248

NODE=M248

NODE=M248205

NODE=M248M+

NODE=M248M+

NODE=M248M+

NODE=M248DM+

NODE=M248DM+

OCCUR=2

NODE=M248DM+;LINKAGE=B

NODE=M248DM+;LINKAGE=A

NODE=M248DM+;LINKAGE=C

NODE=M248DM1

NODE=M248DM1

NODE=M248DM1;LINKAGE=A

NODE=M248M0

NODE=M248M0

NODE=M248M0

NODE=M248DM0

NODE=M248DM0

OCCUR=2

NODE=M248DM0;LINKAGE=B

NODE=M248DM0;LINKAGE=A

NODE=M248DM0;LINKAGE=C

 **$B_J(5970)$  MASS** **$B_J(5970)^+$  MASS**OUR FIT uses  $m_{B^0}$  and  $m_{B_J(5970)^+} - m_{B^0}$  to determine  $m_{B_J(5970)^+}$ .

VALUE (MeV)

DOCUMENT ID

**5964 ± 5 OUR FIT** **$m_{B_J(5970)^+} - m_{B^0}$** 

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

**685 ± 5 OUR FIT****685 ± 5 OUR AVERAGE**

685.3 ± 4.1 ± 2.5

2k

<sup>1</sup> AAIJ15AB LHCB  $pp$  at 7, 8 TeV

681 ± 5 ± 12

1.4k

<sup>2</sup> AALTONEN14l CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.8 ± 4.5 ± 2.5

2k

<sup>3</sup> AAIJ15AB LHCB  $pp$  at 7, 8 TeV<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 545.8 \pm 4.1 \pm 2.5$  MeV which we adjust bythe  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses two relativistic Breit-Wigner functions in the fit for mass difference.<sup>2</sup> AALTONEN 14l reports  $m_{B_J(5970)^+} - m_{B^0} - m_{\pi^+} = 541 \pm 5 \pm 12$  MeV which we adjusted by the  $\pi^+$  mass.<sup>3</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 547 \pm 5 \pm 3$  MeV which we adjust bythe  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$m_{B_J(5970)^+} - m_{B^{*0}}$** 

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.0 ± 4.0 ± 2.5

2k

<sup>1</sup> AAIJ15AB LHCB  $pp$  at 7, 8 TeV<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^+} - m_{B^0}] - (m_{B^{*0}} - m_{B^+}) - m_{\pi^+} = 547 \pm 4 \pm 3$  MeV whichwe adjust by the  $\pi^+$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = -(-1)^J$ ,  $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference. **$B_J(5970)^0$  MASS**OUR FIT uses  $m_{B^+}$  and  $m_{B_J(5970)^0} - m_{B^+}$  to determine  $m_{B_J(5970)^0}$ .

VALUE (MeV)

DOCUMENT ID

**5971 ± 5 OUR FIT** **$m_{B_J(5970)^0} - m_{B^+}$** 

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

**691 ± 5 OUR FIT****691 ± 5 OUR AVERAGE**

689.9 ± 2.9 ± 5.1

10k

<sup>1</sup> AAIJ15AB LHCB  $pp$  at 7, 8 TeV

698 ± 5 ± 12

2.6k

<sup>2</sup> AALTONEN14l CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

714.3 ± 6.4 ± 5.1

10k

<sup>3</sup> AAIJ15AB LHCB  $pp$  at 7, 8 TeV<sup>1</sup> AAIJ 15AB reports  $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 550.4 \pm 2.9 \pm 5.1$  MeV which we adjust bythe  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses two relativistic Breit-Wigner functions in the fit for mass difference.<sup>2</sup> AALTONEN 14l reports  $m_{B_J(5970)^0} - m_{B^+} - m_{\pi^-} = 558 \pm 5 \pm 12$  MeV which we adjusted by the  $\pi^-$  mass.<sup>3</sup> AAIJ 15AB reports  $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 575 \pm 6 \pm 5$  MeV which we adjust bythe  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = (-1)^J$  and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$m_{B_J(5970)^0} - m_{B^{*+}}$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

691.6 ± 3.7 ± 5.1	10k	<sup>1</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
		<sup>1</sup> AAIJ 15AB reports		$[m_{B_J^0} - m_{B^{*+}}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 552 \pm 4 \pm 5$ MeV

which we adjust by the  $\pi^-$  mass. The masses inside the square brackets were measured for each candidate event. The result assumes  $P = -(-1)^J$ ,  $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$  MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M248DM2  
NODE=M248DM2

NODE=M248DM2;LINKAGE=A

 $B_J(5970)$  WIDTH

NODE=M248210

 $B_J(5970)^+$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**62 ± 20 OUR AVERAGE**

63 ± 15 ± 17	2k	<sup>1</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
60 <sup>+30</sup> <sub>-20</sub> ± 40	1.4k	AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

61 ± 14 ± 17	2k	<sup>2</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
61 ± 15 ± 17	2k	<sup>3</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Assuming  $P = (-1)^J$  and using two relativistic Breit-Wigner functions in the fit for mass difference.

<sup>2</sup> Assuming  $P = (-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

<sup>3</sup> Assuming  $P = -(-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M248W+  
NODE=M248W+

OCCUR=2  
OCCUR=3

NODE=M248W+;LINKAGE=A

NODE=M248W+;LINKAGE=B

NODE=M248W+;LINKAGE=C

 $B_J(5970)^0$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**81 ± 12 OUR AVERAGE**

82 ± 8 ± 9	10k	<sup>1</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
70 <sup>+30</sup> <sub>-20</sub> ± 30	2.6k	AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

56 ± 7 ± 9	10k	<sup>2</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
82 ± 10 ± 9	10k	<sup>3</sup> AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Assuming  $P = (-1)^J$  and using two relativistic Breit-Wigner functions in the fit for mass difference.

<sup>2</sup> Assuming  $P = (-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

<sup>3</sup> Assuming  $P = -(-1)^J$  and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M248W0  
NODE=M248W0

OCCUR=2  
OCCUR=3

NODE=M248W0;LINKAGE=A

NODE=M248W0;LINKAGE=B

NODE=M248W0;LINKAGE=C

 $B_J(5970)$  DECAY MODES

NODE=M248215;NODE=M248

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B\pi$	possibly seen
$\Gamma_2$ $B^*\pi$	seen

DESIG=1  
DESIG=2

 $B_J(5970)$  BRANCHING RATIOS

NODE=M248220

 $\Gamma(B\pi)/\Gamma_{\text{total}}$ 

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_1/\Gamma$
-------	------	-------------	------	-----	---------	-------------------

possibly seen	2k	<sup>1</sup> AAIJ	15AB LHCB	±	$p\bar{p}$ at 7, 8 TeV	
possibly seen	10k	<sup>1</sup> AAIJ	15AB LHCB	0	$p\bar{p}$ at 7, 8 TeV	
<b>possibly seen</b>	2.6k	AALTONEN	14I CDF	0	$p\bar{p}$ at 1.96 TeV	
<b>possibly seen</b>	1.4k	AALTONEN	14I CDF	±	$p\bar{p}$ at 1.96 TeV	

<sup>1</sup> A  $B\pi$  decay is forbidden from a  $P = -(-1)^J$  parent, whereas  $B^*\pi$  is allowed.

NODE=M248R01  
NODE=M248R01

OCCUR=2

OCCUR=2

NODE=M248R01;LINKAGE=A

 $\Gamma(B^*\pi)/\Gamma_{\text{total}}$ 

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_2/\Gamma$
-------	------	-------------	------	-----	---------	-------------------

seen	10k	AAIJ	15AB LHCB	0	$p\bar{p}$ at 7, 8 TeV	
seen	2k	AAIJ	15AB LHCB	±	$p\bar{p}$ at 7, 8 TeV	
<b>seen</b>	2.6k	AALTONEN	14I CDF	0	$p\bar{p}$ at 1.96 TeV	
<b>seen</b>	1.4k	AALTONEN	14I CDF	±	$p\bar{p}$ at 1.96 TeV	

NODE=M248R02  
NODE=M248R02

OCCUR=2

OCCUR=2

 $B_J(5970)$  REFERENCES

NODE=M248

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)

REFID=56628  
REFID=56029

# BOTTOM, STRANGE MESONS

## ( $B = \pm 1, S = \mp 1$ )

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \text{ similarly for } B_s^{*0}$$

### $B_{s1}(5830)^0$

$$I(J^P) = 0(1^+)$$

$I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=MXXX046

NODE=MXXX046

NODE=M187

NODE=M187

NODE=M187M

NODE=M187M

### $B_{s1}(5830)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>5828.70<math>\pm</math>0.20 OUR FIT</b>			
<b>5828.65<math>\pm</math>0.24 OUR AVERAGE</b>			
5828.78 $\pm$ 0.09 $\pm$ 0.29	SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV
5828.40 $\pm$ 0.04 $\pm$ 0.41	<sup>1</sup> AAIJ	130 LHCb	$p\bar{p}$ at 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5829.4 $\pm$ 0.7	<sup>2</sup> AALTONEN	08K CDF	Repl. by AALTONEN 14I
<sup>1</sup> Uses $B_{s1}(5830)^0 \rightarrow B^{*+} K^-$ decay.			
<sup>2</sup> Uses two-body decays into $K^-$ and $B^+$ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$ , $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$ , $\bar{D}^0 \rightarrow K^+ \pi^-$ .			

NODE=M187M;LINKAGE=AI

NODE=M187M;LINKAGE=AA

$$m_{B_{s1}^0} - m_{B^{*+}}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>503.99<math>\pm</math>0.17 OUR FIT</b>			
<b>504.03<math>\pm</math>0.12<math>\pm</math>0.15</b>			
	<sup>1</sup> AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
504.41 $\pm$ 0.21 $\pm$ 0.14	<sup>2</sup> AALTONEN	08K CDF	Repl. by AALTONEN 14I
<sup>1</sup> AALTONEN 14I reports $m_{B_{s1}(5830)^0} - m_{B^{*+}} - m_{K^-} = 10.35 \pm 0.12 \pm 0.15$ MeV which we adjusted by the $K^-$ mass.			
<sup>2</sup> Uses two-body decays into $K^-$ and $B^+$ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$ , $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$ , $\bar{D}^0 \rightarrow K^+ \pi^-$ .			

NODE=M187DM

NODE=M187DM

NODE=M187DM;LINKAGE=AL

NODE=M187DM;LINKAGE=AA

### $B_{s1}(5830)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>0.5<math>\pm</math>0.3<math>\pm</math>0.3</b>	AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M187W

NODE=M187W

### $B_{s1}(5830)^0$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^{*+} K^-$	seen
$\Gamma_2$ $B^{*0} K_S^0$	

NODE=M187215;NODE=M187

DESIG=1

DESIG=2

### $B_{s1}(5830)^0$ BRANCHING RATIOS

$\Gamma(B^{*+} K^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AALTONEN	08K CDF	$p\bar{p}$ at 1.96 TeV
$\Gamma(B^{*0} K_S^0)/\Gamma(B^{*+} K^-)$	$\Gamma_2/\Gamma_1$		
VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.49<math>\pm</math>0.12<math>\pm</math>0.07</b>	<sup>1</sup> SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV

NODE=M187220

NODE=M187R01

NODE=M187R01

NODE=M187R00

NODE=M187R00

<sup>1</sup> With the branching fractions  $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$  and  $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$ .

NODE=M187R00;LINKAGE=A

**$B_{s1}(5830)^0$  REFERENCES**

SIRUNYAN	18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	130 PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=M187

REFID=59328  
REFID=56029  
REFID=54968  
REFID=52235 **$B_{s2}^*(5840)^0$**  $I(J^P) = 0(2^+)$   
 $I, J, P$  need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M186

NODE=M186

 **$B_{s2}^*(5840)^0$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**5839.86±0.12 OUR FIT****5839.92±0.14 OUR AVERAGE**

5839.86±0.09±0.17	SIRUNYAN	18DF CMS	$pp$ at 8 TeV
5839.99±0.05±0.20	AAIJ	130 LHCb	$pp$ at 7 TeV
5839.6 ±1.1 ±0.7	<sup>1</sup> ABAZOV	08E D0	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

5839.7 ±0.7 <sup>2</sup>AALTONEN 08K CDF Repl. by AALTONEN 14I<sup>1</sup> Observed in  $B_{s2}^{*0} \rightarrow B^+ K^-$ . Measured production rate of  $B_{s2}^{*0}$  relative to  $B^+$  to be  $(1.15 \pm 0.23 \pm 0.13)\%$ .<sup>2</sup> Uses two-body decays into  $K^-$  and  $B^+$  mesons reconstructed as  $B^+ \rightarrow J/\psi K^+$ ,  $J/\psi \rightarrow \mu^+ \mu^-$  or  $B^+ \rightarrow \bar{D}^0 \pi^+$ ,  $\bar{D}^0 \rightarrow K^+ \pi^-$ .

NODE=M186M

NODE=M186M

NODE=M186M;LINKAGE=AB

NODE=M186M;LINKAGE=AA

 **$m_{B_{s2}^{*0}} - m_{B_{s1}^0}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.5±0.6 <sup>1</sup>AALTONEN 08K CDF Repl. by AALTONEN 14I<sup>1</sup> Uses two-body decays into  $K^-$  and  $B^+$  mesons reconstructed as  $B^+ \rightarrow J/\psi K^+$ ,  $J/\psi \rightarrow \mu^+ \mu^-$  or  $B^+ \rightarrow \bar{D}^0 \pi^+$ ,  $\bar{D}^0 \rightarrow K^+ \pi^-$ . **$m_{B_{s2}^{*0}} - m_{B^+}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**560.52±0.14 OUR FIT****560.41±0.13±0.14**<sup>1</sup> AALTONEN 14I CDF  $p\bar{p}$  at 1.96 TeV<sup>1</sup> AALTONEN 14I reports  $m_{B_{s2}^*(5840)^0} - m_{B^+} - m_{K^-} = 66.73 \pm 0.13 \pm 0.14$  MeV which we adjusted by the  $K^-$  mass.

NODE=M186DM

NODE=M186DM

NODE=M186DM;LINKAGE=AA

NODE=M186DM2

NODE=M186DM2

NODE=M186DM2;LINKAGE=AL

 **$B_{s2}^*(5840)^0$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**1.49±0.27 OUR AVERAGE**

1.52±0.34±0.30	SIRUNYAN	18DF CMS	$pp$ at 8 TeV
1.4 ±0.4 ±0.2	AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
1.56±0.13±0.47	<sup>1</sup> AAIJ	130 LHCb	$pp$ at 7 TeV

<sup>1</sup> Uses  $B_{s2}^*(5840)^0 \rightarrow B^{*+} K^-$  decays.

NODE=M186W

NODE=M186W

NODE=M186W;LINKAGE=AI

 **$B_{s2}^*(5840)^0$  DECAY MODES**Branching fractions are given relative to the one **DEFINED AS 1**.

NODE=M186215;NODE=M186

NODE=M186

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^+ K^-$	<b>DEFINED AS 1</b>
$\Gamma_2$ $B^{*+} K^-$	$0.093 \pm 0.018$
$\Gamma_3$ $B^0 K_S^0$	$0.43 \pm 0.11$
$\Gamma_4$ $B^{*0} K_S^0$	$0.04 \pm 0.04$

DESIG=1

DESIG=2

DESIG=4

DESIG=3



**$B_{s2}^*(5840)^0$  BRANCHING RATIOS** $\Gamma(B^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AALTONEN	08K CDF	$p\bar{p}$ at 1.96 TeV
<b>seen</b>	<sup>1</sup> ABAZOV	08E D0	$p\bar{p}$ at 1.96 TeV

<sup>1</sup> Measured production rate of  $B_{s2}^{*0}$  relative to  $B^+$  to be  $(1.15 \pm 0.23 \pm 0.13)\%$ .

NODE=M186220

NODE=M186R01  
NODE=M186R01

NODE=M186R01;LINKAGE=AB

 $\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$  $\Gamma_2/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.093 ± 0.013 ± 0.012</b>	AAIJ	130 LHCB	$p\bar{p}$ at 7 TeV

NODE=M186R02  
NODE=M186R02 $\Gamma(B^{*0} K_S^0)/\Gamma(B^0 K_S^0)$  $\Gamma_4/\Gamma_3$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.093 ± 0.086 ± 0.014</b>	<sup>1</sup> SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV

NODE=M186R00  
NODE=M186R00<sup>1</sup> With the branching fraction  $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$ .

NODE=M186R00;LINKAGE=A

 $\Gamma(B^0 K_S^0)/\Gamma(B^+ K^-)$  $\Gamma_3/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.432 ± 0.077 ± 0.078</b>	<sup>1</sup> SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV

NODE=M186R03  
NODE=M186R03<sup>1</sup> With the branching fractions  $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$  and  $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$ .

NODE=M186R03;LINKAGE=A

 $\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$  $\Gamma_2/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.081 ± 0.021 ± 0.015</b>	<sup>1</sup> SIRUNYAN	18DF CMS	$p\bar{p}$ at 8 TeV

NODE=M186R04  
NODE=M186R04<sup>1</sup> With the branching fraction  $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$ .

NODE=M186R04;LINKAGE=A

 **$B_{s2}^*(5840)^0$  REFERENCES**

SIRUNYAN	18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59328
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=56029
AAIJ	130 PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54968
AALTONEN	08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52235
ABAZOV	08E PRL 100 082002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52232

NODE=M186

 **$B_{sJ}^*(5850)$**  $I(J^P) = ?(??)$ 

OMITTED FROM SUMMARY TABLE

Signal can be interpreted as coming from  $\bar{b}s$  states. Needs confirmation.

NODE=M153

NODE=M153

 **$B_{sJ}^*(5850)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5853 ± 15</b>	141	AKERS	95E OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M153M

NODE=M153M

 **$B_{sJ}^*(5850)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>47 ± 22</b>	141	AKERS	95E OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M153W

NODE=M153W

 **$B_{sJ}^*(5850)$  REFERENCES**

AKERS	95E ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)
-------	-----------------	------------------------	----------------

NODE=M153

REFID=44182

$B_{sJ}(6063)^0$ 

$I(J^P) = 0(?^?)$

OMITTED FROM SUMMARY TABLE

 $B_{sJ}(6063)^0$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$6063.5 \pm 1.2 \pm 0.8$	<sup>1</sup> AAIJ	21D LHCB	$pp$ at 7, 8, 13 TeV

<sup>1</sup> Seen in the decay channel  $B^\pm K^\mp$ . Integrated luminosity = 9 fb<sup>-1</sup>.

NODE=M257M

NODE=M257M

NODE=M257M;LINKAGE=A

 $B_{sJ}(6063)^0$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$26 \pm 4 \pm 4$	<sup>1</sup> AAIJ	21D LHCB	$pp$ at 7, 8, 13 TeV

<sup>1</sup> Seen in the decay channel  $B^\pm K^\mp$ . Integrated luminosity = 9 fb<sup>-1</sup>.

NODE=M257W

NODE=M257W

NODE=M257W;LINKAGE=A

 $B_{sJ}(6063)^0$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^+ K^-$	seen

NODE=M257215;NODE=M257

DESIG=1

$\Gamma(B^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
seen	AAIJ	21D LHCB	$B^\pm K^\mp$ mass spectrum	

NODE=M257R01  
NODE=M257R01 $B_{sJ}(6063)^0$  REFERENCESAAIJ 21D EPJ C81 601 R. Aaij *et al.* (LHCb Collab.)

NODE=M257

REFID=61130

NODE=M258

 $B_{sJ}(6114)^0$ 

$I(J^P) = 0(?^?)$

OMITTED FROM SUMMARY TABLE

 $B_{sJ}(6114)^0$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$6114 \pm 3 \pm 5$	<sup>1</sup> AAIJ	21D LHCB	$pp$ at 7, 8, 13 TeV

<sup>1</sup> Seen in the decay channel  $B^\pm K^\mp$ . Integrated luminosity = 9 fb<sup>-1</sup>.

NODE=M258M

NODE=M258M

NODE=M258M;LINKAGE=A

 $B_{sJ}(6114)^0$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$66 \pm 18 \pm 21$	<sup>1</sup> AAIJ	21D LHCB	$pp$ at 7, 8, 13 TeV

<sup>1</sup> Seen in the decay channel  $B^\pm K^\mp$ . Integrated luminosity = 9 fb<sup>-1</sup>.

NODE=M258W

NODE=M258W

NODE=M258W;LINKAGE=B

 $B_{sJ}(6114)^0$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B^+ K^-$	seen

NODE=M258215;NODE=M258

DESIG=1

$\Gamma(B^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
seen	AAIJ	21D LHCB	$B^\pm K^\mp$ mass spectrum	

NODE=M258R01  
NODE=M258R01 $B_{sJ}(6114)^0$  REFERENCESAAIJ 21D EPJ C81 601 R. Aaij *et al.* (LHCb Collab.)

NODE=M258

REFID=61130

# BOTTOM, CHARMED MESONS ( $B = C = \pm 1$ )

$$B_c^+ = c\bar{b}, B_c^- = \bar{c}b, \text{ similarly for } B_c^{*'}s$$

## $B_c(2S)^\pm$

$$I(J^P) = 0(0^-)$$

Quantum numbers neither measured nor confirmed.

### $B_c(2S)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6871.2±1.0 OUR AVERAGE</b>				
6871.7±1.3±0.3	24	1,2 AAIJ	19Y LHCB	$pp$ at 7, 8, 13 TeV
6870.6±1.4±0.3	51	3,4 SIRUNYAN	19M CMS	$pp$ at 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		5 AAIJ	18AL LHCB	$pp$ at 8 TeV
6842 ±4 ±5	57	6,7 AAD	14AQ ATLS	$pp$ at 7, 8 TeV

<sup>1</sup> AAIJ 19Y observed  $B_c(2S)^+$  in the decay mode  $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$  ( $B_c^+ \rightarrow J/\psi \pi^+$ ) with 2.2 (3.2) global (local) standard deviations significance.

<sup>2</sup> AAIJ 19Y reports mass difference measurement of  $M(B_c(2S)^+) - M(B_c^+) = 597.2 \pm 1.3 \pm 0.1$  MeV. We have adjusted this measurement with our best value of  $M(B_c^+) = 6274.47 \pm 0.32$  MeV. The first uncertainty of the  $M(B_c(2S)^+)$  value is a total of uncertainties reported by the experiment and the second one comes from our best value of  $M(B_c^+)$ .

<sup>3</sup> SIRUNYAN 19M observed  $B_c(2S)^+$  in the decay mode  $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$  ( $B_c^+ \rightarrow J/\psi \pi^+$ ) with 6.5 standard deviations significance.

<sup>4</sup> SIRUNYAN 19M reports mass difference measurement of  $M(B_c(2S)^+) - M(B_c^+) = 596.1 \pm 1.2 \pm 0.8$  MeV. We have adjusted this measurement with our best value of  $M(B_c^+) = 6274.47 \pm 0.32$  MeV. The first uncertainty of the  $M(B_c(2S)^+)$  value is a total of uncertainties reported by the experiment and the second one comes from our best value of  $M(B_c^+)$ .

<sup>5</sup> AAIJ 18AL reports an upper limit on the ratio of production cross sections for  $[\sigma(B_c(2S)^+)/\sigma(B_c^+)]: B(B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-) < 0.04-0.09$  at 95% CL for the mass value reported by AAD 14AQ.

<sup>6</sup> Observed in the decay mode  $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$  ( $B_c^+ \rightarrow J/\psi \pi^+$ ) with 5.2 standard deviations significance.

<sup>7</sup> Might be the  $B_c^{*'}(2S)$ .

### $B_c(2S)^\pm$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad B_c^+ \pi^+ \pi^-$	seen

### $B_c(2S)^\pm$ BRANCHING RATIOS

$\Gamma(B_c^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	57	1 AAD	14AQ ATLS	$pp$ at 7, 8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen <sup>2</sup> AAIJ 18AL LHCB  $pp$  at 8 TeV

<sup>1</sup> Observed with 5.2 standard deviations significance.

<sup>2</sup> AAIJ 18AL reports an upper limit on the ratio of production cross sections for  $[\sigma(B_c(2S)^+)/\sigma(B_c^+)]: B(B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-) < 0.04-0.09$  at 95% CL for the mass value reported by AAD 14AQ.

NODE=MXXX049

NODE=MXXX049

NODE=M217

NODE=M217

NODE=M217M

NODE=M217M

SYCLP=A

SYCLP=A

NODE=M217M;LINKAGE=E

NODE=M217M;LINKAGE=F

NODE=M217M;LINKAGE=B

NODE=M217M;LINKAGE=D

NODE=M217M;LINKAGE=A

NODE=M217M;LINKAGE=AA

NODE=M217M;LINKAGE=C

NODE=M217215;NODE=M217

DESIG=1

NODE=M217225

NODE=M217R01

NODE=M217R01

NODE=M217R01;LINKAGE=AA

NODE=M217R01;LINKAGE=A

**$B_c(2S)^\pm$  REFERENCES**

AAIJ	19Y	PRL 122 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	19M	PRL 122 132001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	18AL	JHEP 1801 138	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	14AQ	PRL 113 212004	G. Aad <i>et al.</i>	(ATLAS Collab.)

NODE=M217

REFID=59795  
REFID=59639  
REFID=59246  
REFID=56117


See the related review(s):  
[Spectroscopy of Mesons Containing Two Heavy Quarks](#)

**$c\bar{c}$  MESONS**  
**(including possibly non- $q\bar{q}$  states)**

NODE=MXXX025

NODE=M826

Updated March 2022.



The level scheme of meson states containing a minimal quark content of  $c\bar{c}$ . The name of a state is determined by its quantum numbers  $I^G J^{PC}$  (see the review “Naming Scheme for Hadrons”). States with unestablished quantum numbers are called  $X$  and are drawn according to our best estimate of their likely  $J^{PC}$ . States included in the Summary Tables are shown with solid lines; selected states not in the Summary Tables, but with assigned quantum numbers, are shown with dotted lines. The arrows indicate the most dominant hadronic transitions. Single photon transitions, including  $\psi(nS) \rightarrow \gamma\eta_c(mS)$ ,  $\psi(nS) \rightarrow \gamma\chi_{cJ}(1P)$ , and  $\chi_{cJ}(1P) \rightarrow \gamma J/\psi$ , are omitted for clarity. For orientation, the location of the thresholds related to a pair of ground state open charm mesons is indicated in the figure.

$\eta_c(1S)$ 

$$J^G(J^{PC}) = 0^+(0^{-+})$$

NODE=M026

 $\eta_c(1S)$  MASS

NODE=M026M

NODE=M026M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2983.9 ± 0.4</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.			
2983.9 ± 0.7 ± 0.1		<sup>1</sup> AAIJ	20H LHCB	$p\bar{p} \rightarrow bX \rightarrow p\bar{p}X$	
2985.9 ± 0.7 ± 2.1	1705	ABLIKIM	19AV BES3	$J/\psi \rightarrow \gamma\omega\omega$	
2984.6 ± 0.7 ± 2.2	2673	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	
2986.7 ± 0.5 ± 0.9	11k	<sup>2</sup> AAIJ	17AD LHCB	$p\bar{p} \rightarrow B^+X \rightarrow p\bar{p}K^+X$	
2982.8 ± 1.0 ± 0.5	6.4k	<sup>3</sup> AAIJ	17BB LHCB	$p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$	
2982.2 ± 1.5 ± 0.1	2.0k	<sup>4</sup> AAIJ	15BI LHCB	$p\bar{p} \rightarrow \eta_c(1S)X$	
2983.5 ± 1.4 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 1.6 \\ 3.6 \end{smallmatrix}$		<sup>5</sup> ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$	
2979.8 ± 0.8 ± 3.5	4.5k	<sup>6,7</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$	
2984.1 ± 1.1 ± 2.1	900	<sup>6,7,8</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$	OCCUR=2
2984.3 ± 0.6 ± 0.6		<sup>9,10</sup> ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$	
2984.49 ± 1.16 ± 0.52	832	<sup>6</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons	
2982.7 ± 1.8 ± 2.2	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$	
2984.5 ± 0.8 ± 3.1	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	
2985.4 ± 1.5 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 0.5 \\ 2.0 \end{smallmatrix}$	920	<sup>10</sup> VINOKUROVA	11 BELL	$B^\pm \rightarrow K_S^0 K^\pm \pi^\mp$	
2982.2 ± 0.4 ± 1.6	14k	<sup>11</sup> LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$	
2985.8 ± 1.5 ± 3.1	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K^(*) \rightarrow K\bar{K}\pi K^(*)$	
2986.1 ± 1.0 ± 2.5	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow$ hadrons	
2970 ± 5 ± 6	501	<sup>12</sup> ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$	
2971 ± 3 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2 \\ 1 \end{smallmatrix}$	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$	
2974 ± 7 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2 \\ 1 \end{smallmatrix}$	20	WU	06 BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$	OCCUR=2
2981.8 ± 1.3 ± 1.5	592	ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c' \rightarrow K_S^0 K^\pm \pi^\mp$	
2984.1 ± 2.1 ± 1.0	190	<sup>13</sup> AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2982.5 ± 0.4 ± 1.4	12k	<sup>14</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	OCCUR=2
2982.2 ± 0.6		<sup>15</sup> MITCHELL	09 CLEO	$e^+e^- \rightarrow \gamma X$	
2982 ± 5	270	<sup>16</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c\bar{c}$	
2982.5 ± 1.1 ± 0.9	2.5k	<sup>17</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow KK\pi$	
2977.5 ± 1.0 ± 1.2		<sup>15,18</sup> BAI	03 BES	$J/\psi \rightarrow \gamma\eta_c$	
2979.6 ± 2.3 ± 1.6	180	<sup>19</sup> FANG	03 BELL	$B \rightarrow \eta_c K$	
2976.3 ± 2.3 ± 1.2		<sup>15,20</sup> BAI	00F BES	$J/\psi, \psi(2S) \rightarrow \gamma\eta_c$	
2976.6 ± 2.9 ± 1.3	140	<sup>15,21</sup> BAI	00F BES	$J/\psi \rightarrow \gamma\eta_c$	OCCUR=2
2980.4 ± 2.3 ± 0.6		<sup>22</sup> BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$	
2975.8 ± 3.9 ± 1.2		<sup>21</sup> BAI	99B BES	Sup. by BAI 00F	
2999 ± 8	25	ABREU	98O DLPH	$e^+e^- \rightarrow e^+e^- +$ hadrons	
2988.3 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 3.3 \\ 3.1 \end{smallmatrix}$		ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$	
2974.4 ± 1.9		<sup>15,23</sup> BISELLO	91 DM2	$J/\psi \rightarrow \eta_c\gamma$	
2969 ± 4 ± 4	80	<sup>15</sup> BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^- K^+K^-$	
2956 ± 12 ± 12		<sup>15</sup> BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^- K_S^0 K_L^0$	OCCUR=3
2982.6 $\begin{smallmatrix} + \\ - \end{smallmatrix} \begin{smallmatrix} 2.7 \\ 2.3 \end{smallmatrix}$	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$	
2980.2 ± 1.6		<sup>15,23</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	

2984	$\pm 2.3 \pm 4.0$	15	GAISER	86	CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
2976	$\pm 8$	15,24	BALTRUSAIT..84	MRK3		$J/\psi \rightarrow 2\phi\gamma$
2982	$\pm 8$	18	25 HIMEL	80B	MRK2	$e^+e^-$
2980	$\pm 9$	25	PARTRIDGE	80B	CBAL	$e^+e^-$

<sup>1</sup>AAIJ 20H report  $m_{J/\psi} - m_{\eta_c(1S)} = 113.0 \pm 0.7 \pm 0.1$  MeV. We use the current value  $m_{J/\psi} = 3096.900 \pm 0.006$  MeV to obtain the quoted mass.

NODE=M026M;LINKAGE=H

<sup>2</sup>AAIJ 17AD report  $m_{J/\psi} - m_{\eta_c(1S)} = 110.2 \pm 0.5 \pm 0.9$  MeV. We use the current value  $m_{J/\psi} = 3096.900 \pm 0.006$  MeV to obtain the quoted mass.

NODE=M026M;LINKAGE=F

<sup>3</sup>From a fit of the  $\phi\phi$  invariant mass with the mass and width of  $\eta_c(1S)$  as free parameters.

NODE=M026M;LINKAGE=G

<sup>4</sup>AAIJ 15BI reports  $m_{J/\psi} - m_{\eta_c(1S)} = 114.7 \pm 1.5 \pm 0.1$  MeV from a sample of  $\eta_c(1S)$  and  $J/\psi$  produced in  $b$ -hadron decays. We have used current value of  $m_{J/\psi} = 3096.900 \pm 0.006$  MeV to arrive at the quoted  $m_{\eta_c(1S)}$  result.

NODE=M026M;LINKAGE=D

<sup>5</sup>Taking into account an asymmetric photon lineshape.

NODE=M026M;LINKAGE=E

<sup>6</sup>With floating width.

NODE=M026M;LINKAGE=AL

<sup>7</sup>Ignoring possible interference with the non-resonant  $0^-$  amplitude.

NODE=M026M;LINKAGE=LS

<sup>8</sup>Using both,  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+\pi^-\pi^0$  decays.

NODE=M026M;LINKAGE=EL

<sup>9</sup>From a simultaneous fit to six decay modes of the  $\eta_c$ .

NODE=M026M;LINKAGE=BL

<sup>10</sup>Accounts for interference with non-resonant continuum.

NODE=M026M;LINKAGE=VA

<sup>11</sup>Taking into account interference with the non-resonant  $J^P = 0^-$  amplitude.

NODE=M026M;LINKAGE=LE

<sup>12</sup>From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

NODE=M026M;LINKAGE=EB

<sup>13</sup>Using mass of  $\psi(2S) = 3686.00$  MeV.

NODE=M026M;LINKAGE=BG

<sup>14</sup>Not independent from the measurements reported by LEES 10.

NODE=M026M;LINKAGE=DE

<sup>15</sup>MITCHELL 09 observes a significant asymmetry in the lineshapes of  $\psi(2S) \rightarrow \gamma\eta_c$  and  $J/\psi \rightarrow \gamma\eta_c$  transitions. If ignored, this asymmetry could lead to significant bias whenever the mass and width are measured in  $\psi(2S)$  or  $J/\psi$  radiative decays.

NODE=M026M;LINKAGE=MI

<sup>16</sup>From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

NODE=M026M;LINKAGE=AU

<sup>17</sup>Superseded by LEES 10.

NODE=M026M;LINKAGE=UB

<sup>18</sup>From a simultaneous fit of five decay modes of the  $\eta_c$ .

NODE=M026M;LINKAGE=AK

<sup>19</sup>Superseded by VINOKUROVA 11.

NODE=M026M;LINKAGE=FA

<sup>20</sup>Weighted average of the  $\psi(2S)$  and  $J/\psi(1S)$  samples. Using an  $\eta_c$  width of 13.2 MeV.

NODE=M026M;LINKAGE=KZ

<sup>21</sup>Average of several decay modes. Using an  $\eta_c$  width of 13.2 MeV.

NODE=M026M;LINKAGE=C1

<sup>22</sup>Superseded by ASNER 04.

NODE=M026M;LINKAGE=NN

<sup>23</sup>Average of several decay modes.

NODE=M026M;LINKAGE=A

<sup>24</sup> $\eta_c \rightarrow \phi\phi$ .

NODE=M026M;LINKAGE=B

<sup>25</sup>Mass adjusted by us to correspond to  $J/\psi(1S)$  mass = 3097 MeV.

NODE=M026M;LINKAGE=M

### $\eta_c(1S)$ WIDTH

NODE=M026W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>32.0 ± 0.7 OUR FIT</b>				
<b>32.1 ± 0.8 OUR AVERAGE</b>				Error includes scale factor of 1.1.
33.8 ± 1.6 ± 4.1	1705	ABLIKIM	19AV BES3	$J/\psi \rightarrow \gamma\omega\omega$
30.8 <sup>+</sup> <sub>-2.2</sub> ± 2.3 ± 2.9	2673	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
34.0 ± 1.9 ± 1.3	11k	AAIJ	17AD LHCB	$pp \rightarrow B^+X \rightarrow p\bar{p}K^+X$
31.4 ± 3.5 ± 2.0	6.4k	<sup>1</sup> AAIJ	17BB LHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
27.2 ± 3.1 <sup>+</sup> <sub>-2.6</sub>		<sup>2</sup> ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
25.2 ± 2.6 ± 2.4	4.5k	<sup>3,4</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
34.8 ± 3.1 ± 4.0	900	<sup>3,4,5</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
32.0 ± 1.2 ± 1.0		<sup>6,7</sup> ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
36.4 ± 3.2 ± 1.7	832	<sup>3</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons
37.8 <sup>+</sup> <sub>-5.3</sub> ± 5.8 ± 3.1	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
36.2 ± 2.8 ± 3.0	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
35.1 ± 3.1 <sup>+</sup> <sub>-1.6</sub>	920	<sup>7</sup> VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm\pi^\mp)$
31.7 ± 1.2 ± 0.8	14k	<sup>8</sup> LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0 K^\pm\pi^\mp$
36.3 <sup>+</sup> <sub>-3.6</sub> ± 3.7 ± 4.4	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K^(*) \rightarrow K\bar{K}\pi K^(*)$
28.1 ± 3.2 ± 2.2	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow$ hadrons
48 <sup>+</sup> <sub>-7</sub> ± 8 ± 5	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$

NODE=M026W

OCCUR=2

40 ±19 ±5	20	WU	06	BELL	$B^+ \rightarrow \Lambda \bar{\Lambda} K^+$	OCCUR=2
24.8 ± 3.4 ± 3.5	592	ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$	
20.4 <sup>+</sup> <sub>-</sub> 7.7 ± 2.0	190	AMBROGIANI	03	E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$	
23.9 <sup>+</sup> <sub>-</sub> 12.6 ± 7.1		ARMSTRONG	95F	E760	$\bar{p}p \rightarrow \gamma\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

32.1 ± 1.1 ± 1.3	12k	<sup>9</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	OCCUR=2
34.3 ± 2.3 ± 0.9	2.5k	<sup>10</sup> AUBERT	04D	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K \bar{K} \pi$	
17.0 ± 3.7 ± 7.4		<sup>11</sup> BAI	03	BES $J/\psi \rightarrow \gamma \eta_c$	
29 ± 8 ± 6	180	<sup>12</sup> FANG	03	BELL $B \rightarrow \eta_c K$	
11.0 ± 8.1 ± 4.1		<sup>13</sup> BAI	00F	BES $J/\psi \rightarrow \gamma \eta_c$ and $\psi(2S) \rightarrow \gamma \eta_c$	
27.0 ± 5.8 ± 1.4		<sup>14</sup> BRANDENB...	00B	CLE2 $\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$	
7.0 <sup>+</sup> <sub>-</sub> 7.5 ± 7.0	12	BAGLIN	87B	SPEC $\bar{p}p \rightarrow \gamma\gamma$	
10.1 <sup>+</sup> <sub>-</sub> 33.0 ± 8.2	23	<sup>15</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma p \bar{p}$	
11.5 ± 4.5		GAISER	86	CBAL $J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$	
< 40 90% CL	18	HIMEL	80B	MRK2 $e^+ e^-$	
< 20 90% CL		PARTRIDGE	80B	CBAL $e^+ e^-$	

<sup>1</sup> From a fit of the  $\phi\phi$  invariant mass with the mass and width of  $\eta_c(1S)$  as free parameters.

<sup>2</sup> Taking into account an asymmetric photon lineshape.

<sup>3</sup> With floating mass.

<sup>4</sup> Ignoring possible interference with the non-resonant  $0^-$  amplitude.

<sup>5</sup> Using both,  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

<sup>6</sup> From a simultaneous fit to six decay modes of the  $\eta_c$ .

<sup>7</sup> Accounts for interference with non-resonant continuum.

<sup>8</sup> Taking into account interference with the non-resonant  $J^P = 0^-$  amplitude.

<sup>9</sup> Not independent from the measurements reported by LEES 10.

<sup>10</sup> Superseded by LEES 10.

<sup>11</sup> From a simultaneous fit of five decay modes of the  $\eta_c$ .

<sup>12</sup> Superseded by VINOKUROVA 11.

<sup>13</sup> From a fit to the 4-prong invariant mass in  $\psi(2S) \rightarrow \gamma \eta_c$  and  $J/\psi(1S) \rightarrow \gamma \eta_c$  decays.

<sup>14</sup> Superseded by ASNER 04.

<sup>15</sup> Positive and negative errors correspond to 90% confidence level.

NODE=M026W;LINKAGE=B  
 NODE=M026W;LINKAGE=A  
 NODE=M026W;LINKAGE=AL  
 NODE=M026W;LINKAGE=LS  
 NODE=M026W;LINKAGE=EL  
 NODE=M026W;LINKAGE=BL  
 NODE=M026W;LINKAGE=VA  
 NODE=M026W;LINKAGE=LE  
 NODE=M026W;LINKAGE=DE  
 NODE=M026W;LINKAGE=UB  
 NODE=M026W;LINKAGE=AK  
 NODE=M026W;LINKAGE=FA  
 NODE=M026W;LINKAGE=KZ  
 NODE=M026W;LINKAGE=NN  
 NODE=M026W;LINKAGE=L

## $\eta_c(1S)$ DECAY MODES

NODE=M026215;NODE=M026

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
<b>Decays involving hadronic resonances</b>		
$\Gamma_1$ $\eta'(958) \pi\pi$	( 1.87 ± 0.26 ) %	
$\Gamma_2$ $\eta'(958) K \bar{K}$	( 1.61 ± 0.25 ) %	
$\Gamma_3$ $\rho\rho$	( 1.5 ± 0.4 ) %	
$\Gamma_4$ $K^*(892)^0 K^- \pi^+ + c.c.$	( 1.5 ± 0.5 ) %	
$\Gamma_5$ $K^*(892) \bar{K}^*(892)$	( 6.3 ± 1.2 ) × 10 <sup>-3</sup>	
$\Gamma_6$ $K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-$	( 1.1 ± 0.5 ) %	
$\Gamma_7$ $\phi K^+ K^-$	( 2.9 ± 1.4 ) × 10 <sup>-3</sup>	
$\Gamma_8$ $\phi\phi$	( 1.58 ± 0.19 ) × 10 <sup>-3</sup>	
$\Gamma_9$ $\phi 2(\pi^+ \pi^-)$	< 4 × 10 <sup>-3</sup>	90%
$\Gamma_{10}$ $a_0(980) \pi$	seen	
$\Gamma_{11}$ $a_2(1320) \pi$		
$\Gamma_{12}$ $K^*(892) \bar{K} + c.c.$	< 1.28 %	90%
$\Gamma_{13}$ $f_2(1270) \eta$	seen	
$\Gamma_{14}$ $f_2(1270) \eta'$	seen	
$\Gamma_{15}$ $\omega\omega$	( 2.1 ± 0.5 ) × 10 <sup>-3</sup>	
$\Gamma_{16}$ $\omega\phi$	< 2.5 × 10 <sup>-4</sup>	90%
$\Gamma_{17}$ $f_2(1270) f_2(1270)$	( 9.7 ± 2.5 ) × 10 <sup>-3</sup>	
$\Gamma_{18}$ $f_2(1270) f'_2(1525)$	( 9.1 ± 3.0 ) × 10 <sup>-3</sup>	
$\Gamma_{19}$ $f_0(500) \eta$	seen	

NODE=M026;CLUMP=A  
 DESIG=24  
 DESIG=85  
 DESIG=19  
 DESIG=26  
 DESIG=18  
 DESIG=57  
 DESIG=28  
 DESIG=17  
 DESIG=58  
 DESIG=21  
 DESIG=22  
 DESIG=40  
 DESIG=23  
 DESIG=92  
 DESIG=20  
 DESIG=47  
 DESIG=46  
 DESIG=59  
 DESIG=86

Γ <sub>20</sub>	$f_0(500)\eta'$	seen	DESIG=87
Γ <sub>21</sub>	$f_0(980)\eta$	seen	DESIG=70
Γ <sub>22</sub>	$f_0(980)\eta'$	seen	DESIG=88
Γ <sub>23</sub>	$f_0(1500)\eta$	seen	DESIG=71
Γ <sub>24</sub>	$f_0(1710)\eta'$	seen	DESIG=90
Γ <sub>25</sub>	$f_0(2100)\eta'$	seen	DESIG=91
Γ <sub>26</sub>	$f_0(2200)\eta$	seen	DESIG=72
Γ <sub>27</sub>	$a_0(1320)\pi$	seen	DESIG=74
Γ <sub>28</sub>	$a_0(1450)\pi$	seen	DESIG=75
Γ <sub>29</sub>	$a_0(1700)\pi$	seen	DESIG=89
Γ <sub>30</sub>	$a_0(1950)\pi$	seen	DESIG=79
Γ <sub>31</sub>	$K_0^*(1430)\bar{K}$	seen	DESIG=76
Γ <sub>32</sub>	$K_2^*(1430)\bar{K}$	seen	DESIG=77
Γ <sub>33</sub>	$K_0^*(1950)\bar{K}$	seen	DESIG=78

**Decays into stable hadrons**

Γ <sub>34</sub>	$K\bar{K}\pi$	( 7.0 ± 0.4 ) %		NODE=M026;CLUMP=B
Γ <sub>35</sub>	$K\bar{K}\eta$	( 1.32±0.15) %		DESIG=14
Γ <sub>36</sub>	$\eta\pi^+\pi^-$	( 1.7 ± 0.5 ) %		DESIG=25
Γ <sub>37</sub>	$\eta 2(\pi^+\pi^-)$	( 4.6 ± 1.4 ) %		DESIG=16
Γ <sub>38</sub>	$K^+K^-\pi^+\pi^-$	( 6.5 ± 1.0 ) × 10 <sup>-3</sup>		DESIG=61
Γ <sub>39</sub>	$K^+K^-\pi^+\pi^-\pi^0$	( 3.4 ± 0.5 ) %		DESIG=15
Γ <sub>40</sub>	$K^0K^-\pi^+\pi^-\pi^+ + c.c.$	( 5.7 ± 1.6 ) %		DESIG=60
Γ <sub>41</sub>	$K^+K^- 2(\pi^+\pi^-)$	( 7.6 ± 2.4 ) × 10 <sup>-3</sup>		DESIG=62
Γ <sub>42</sub>	$2(K^+K^-)$	( 1.38±0.29) × 10 <sup>-3</sup>		DESIG=55
Γ <sub>43</sub>	$\pi^+\pi^-\pi^0$	< 5 × 10 <sup>-4</sup>	90%	DESIG=27
Γ <sub>44</sub>	$\pi^+\pi^-\pi^0\pi^0$	( 4.8 ± 1.1 ) %		DESIG=81
Γ <sub>45</sub>	$2(\pi^+\pi^-)$	( 8.7 ± 1.1 ) × 10 <sup>-3</sup>		DESIG=63
Γ <sub>46</sub>	$2(\pi^+\pi^-\pi^0)$	(16.2 ± 2.1) %		DESIG=11
Γ <sub>47</sub>	$3(\pi^+\pi^-)$	( 1.8 ± 0.4 ) %		DESIG=64
Γ <sub>48</sub>	$p\bar{p}$	( 1.35±0.13) × 10 <sup>-3</sup>		DESIG=56
Γ <sub>49</sub>	$p\bar{p}\pi^0$	( 3.6 ± 1.4 ) × 10 <sup>-3</sup>		DESIG=12
Γ <sub>50</sub>	$\Lambda\bar{\Lambda}$	( 1.02±0.23) × 10 <sup>-3</sup>		DESIG=65
Γ <sub>51</sub>	$K^+\bar{p}\Lambda + c.c.$	( 2.5 ± 0.4 ) × 10 <sup>-3</sup>		DESIG=45
Γ <sub>52</sub>	$\bar{\Lambda}(1520)\Lambda + c.c.$	( 3.1 ± 1.3 ) × 10 <sup>-3</sup>		DESIG=82
Γ <sub>53</sub>	$\Sigma^+\bar{\Sigma}^-$	( 2.1 ± 0.6 ) × 10 <sup>-3</sup>		DESIG=83
Γ <sub>54</sub>	$\Xi^-\bar{\Xi}^+$	( 9.0 ± 2.6 ) × 10 <sup>-4</sup>		DESIG=66
Γ <sub>55</sub>	$\pi^+\pi^-p\bar{p}$	( 5.5 ± 1.9 ) × 10 <sup>-3</sup>		DESIG=67

**Radiative decays**

Γ <sub>56</sub>	$\gamma\gamma$	( 1.68±0.12) × 10 <sup>-4</sup>		NODE=M026;CLUMP=C
				DESIG=31

**Charge conjugation (C), Parity (P),  
Lepton Family number (LF) violating modes**

Γ <sub>57</sub>	$\pi^+\pi^-$	$P, CP < 1.1$	× 10 <sup>-4</sup>	90%	NODE=M026;CLUMP=D
Γ <sub>58</sub>	$\pi^0\pi^0$	$P, CP < 4$	× 10 <sup>-5</sup>	90%	DESIG=51
Γ <sub>59</sub>	$K^+K^-$	$P, CP < 6$	× 10 <sup>-4</sup>	90%	DESIG=52
Γ <sub>60</sub>	$K_S^0K_S^0$	$P, CP < 3.1$	× 10 <sup>-4</sup>	90%	DESIG=53
					DESIG=54



## CONSTRAINED FIT INFORMATION

An overall fit to the total width, 10 combinations of partial widths obtained from integrated cross section, and 21 branching ratios uses 97 measurements and one constraint to determine 15 parameters. The overall fit has a  $\chi^2 = 120.8$  for 83 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_5$	13									
$x_8$	25	14								
$x_{15}$	10	5	10							
$x_{17}$	5	3	6	2						
$x_{34}$	30	17	35	13	7					
$x_{35}$	15	8	17	6	3	48				
$x_{38}$	16	9	17	7	3	21	10			
$x_{42}$	12	7	13	5	3	21	10	8		
$x_{45}$	18	10	20	8	4	24	12	13	10	
$x_{48}$	17	10	19	7	4	26	13	12	9	14
$x_{50}$	4	2	5	2	1	6	3	3	2	3
$x_{56}$	-45	-25	-50	-19	-10	-60	-29	-31	-24	-36
$\Gamma$	-2	-1	-2	-1	0	-3	-1	-2	-1	-2
	$x_1$	$x_5$	$x_8$	$x_{15}$	$x_{17}$	$x_{34}$	$x_{35}$	$x_{38}$	$x_{42}$	$x_{45}$
$x_{50}$	24									
$x_{56}$	-36	-9								
$\Gamma$	6	1	-28							
	$x_{48}$	$x_{50}$	$x_{56}$							

Mode	Rate (MeV)	
$\Gamma_1$ $\eta'(958)\pi\pi$	0.60 $\pm$ 0.08	DESIG=24
$\Gamma_5$ $K^*(892)\bar{K}^*(892)$	0.20 $\pm$ 0.04	DESIG=18
$\Gamma_8$ $\phi\phi$	0.051 $\pm$ 0.006	DESIG=17
$\Gamma_{15}$ $\omega\omega$	0.066 $\pm$ 0.015	DESIG=20
$\Gamma_{17}$ $f_2(1270)f_2(1270)$	0.31 $\pm$ 0.08	DESIG=46
$\Gamma_{34}$ $K\bar{K}\pi$	2.25 $\pm$ 0.14	DESIG=14
$\Gamma_{35}$ $K\bar{K}\eta$	0.42 $\pm$ 0.05	DESIG=25
$\Gamma_{38}$ $K^+K^-\pi^+\pi^-$	0.206 $\pm$ 0.032	DESIG=15
$\Gamma_{42}$ $2(K^+K^-)$	0.044 $\pm$ 0.009	DESIG=27
$\Gamma_{45}$ $2(\pi^+\pi^-)$	0.277 $\pm$ 0.035	DESIG=11
$\Gamma_{48}$ $p\bar{p}$	0.043 $\pm$ 0.004	DESIG=12
$\Gamma_{50}$ $\Lambda\bar{\Lambda}$	0.033 $\pm$ 0.007	DESIG=45
$\Gamma_{56}$ $\gamma\gamma$	0.0054 $\pm$ 0.0004	DESIG=31

## $\eta_c(1S)$ PARTIAL WIDTHS

NODE=M026217

 $\Gamma(\gamma\gamma)$  $\Gamma_{56}$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**5.4  $\pm$  0.4 OUR FIT**[5.15  $\pm$  0.35 keV OUR 2022 FIT]

••• We do not use the following data for averages, fits, limits, etc. •••

5.8 $\pm$ 1.1	486	<sup>1</sup> ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
5.2 $\pm$ 1.2	273 $\pm$ 43	<sup>2,3</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
5.5 $\pm$ 1.2 $\pm$ 1.8	157 $\pm$ 33	<sup>4</sup> KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$
7.4 $\pm$ 0.4 $\pm$ 2.3		<sup>5</sup> ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$
13.9 $\pm$ 2.0 $\pm$ 3.0	41	<sup>6</sup> ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \eta_c$
3.8 $^{+1.1}_{-1.0}$ $^{+1.9}_{-1.0}$	190	<sup>7</sup> AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$

NODE=M026W1

NODE=M026W1

NEW

7.6 ± 0.8 ± 2.3		5,8	BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
6.9 ± 1.7 ± 2.1	76	9	ACCIARRI	99T L3	$e^+ e^- \rightarrow e^+ e^- \eta_c$
27 ± 16 ± 10	5	5	SHIRAI	98 AMY	58 $e^+ e^-$
6.7 <sup>+</sup> <sub>-</sub> 2.4 <sup>+</sup> <sub>-</sub> 1.7 ± 2.3		4	ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
11.3 ± 4.2		10	ALBRECHT	94H ARG	$e^+ e^- \rightarrow e^+ e^- \eta_c$
8.0 ± 2.3 ± 2.4	17	11	ADRIANI	93N L3	$e^+ e^- \rightarrow e^+ e^- \eta_c$
5.9 <sup>+</sup> <sub>-</sub> 2.1 <sup>+</sup> <sub>-</sub> 1.8 ± 1.9		7	CHEN	90B CLEO	$e^+ e^- \rightarrow e^+ e^- \eta_c$
6.4 <sup>+</sup> <sub>-</sub> 5.0 <sup>+</sup> <sub>-</sub> 3.4		12	AIHARA	88D TPC	$e^+ e^- \rightarrow e^+ e^- X$
4.3 <sup>+</sup> <sub>-</sub> 3.4 <sup>+</sup> <sub>-</sub> 3.7 ± 2.4		4	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
28 ± 15		5,13	BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

<sup>1</sup> Assuming there is no interference with the non-resonant background.

<sup>2</sup> Calculated by us using  $\Gamma(\eta_c \rightarrow K\bar{K}\pi) \times \Gamma(\eta_c \rightarrow \gamma\gamma) / \Gamma = 0.44 \pm 0.05$  keV from PDG 06 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$  from AUBERT 06E.

<sup>3</sup> Systematic errors not evaluated.

<sup>4</sup> Normalized to  $B(\eta_c \rightarrow p\bar{p}) = (1.3 \pm 0.4) \times 10^{-3}$ .

<sup>5</sup> Normalized to  $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$ .

<sup>6</sup> Average of  $K_S^0 K^\pm \pi^\mp$ ,  $\pi^+ \pi^- K^+ K^-$ , and  $2(K^+ K^-)$  decay modes.

<sup>7</sup> Normalized to the sum of  $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$ ,  $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$ , and  $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$ .

<sup>8</sup> Superseded by ASNER 04.

<sup>9</sup> Normalized to the sum of 9 branching ratios.

<sup>10</sup> Normalized to the sum of  $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$ ,  $B(\eta_c \rightarrow \phi\phi)$ ,  $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$ , and  $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$ .

<sup>11</sup> Superseded by ACCIARRI 99T.

<sup>12</sup> Normalized to the sum of  $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$ ,  $B(\eta_c \rightarrow 2K^+ 2K^-)$ ,  $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$ , and  $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$ .

<sup>13</sup> Re-evaluated by AIHARA 88D.

NODE=M026W1;LINKAGE=ZH  
NODE=M026W1;LINKAGE=AU

NODE=M026W1;LINKAGE=NS  
NODE=M026W1;LINKAGE=N3  
NODE=M026W1;LINKAGE=N2  
NODE=M026W;LINKAGE=FF  
NODE=M026W1;LINKAGE=N6

NODE=M026W1;LINKAGE=NN  
NODE=M026W1;LINKAGE=N1  
NODE=M026W1;LINKAGE=N5

NODE=M026W1;LINKAGE=WD  
NODE=M026W1;LINKAGE=N4

NODE=M026W1;LINKAGE=A

### $\eta_c(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M026220

#### $\Gamma(\eta'(958)\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{56}/\Gamma$

NODE=M026G10  
NODE=M026G10

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

#### 101 ± 12 OUR FIT

98.1 ± 3.9 ± 11.7	2673	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
-------------------	------	----	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

75.8 <sup>+</sup> <sub>-</sub> 6.3 <sup>+</sup> <sub>-</sub> 6.2 ± 8.4	486	<sup>1</sup> ZHANG	12A BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
--	-----	--------------------	----------	---

<sup>1</sup> Superseded by XU 18.

NODE=M026G10;LINKAGE=A

#### $\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_{56}/\Gamma$

NODE=M026G09  
NODE=M026G09

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<39	90	< 1556	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+ \pi^-)$
-----	----	--------	--------	---------	---

#### $\Gamma(K^*(892)\bar{K}^*(892)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_{56}/\Gamma$

NODE=M026G08  
NODE=M026G08  
NEW

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

#### 34 ± 6 OUR FIT

[36 ± 6 eV OUR 2022 FIT]

32.4 ± 4.2 ± 5.8	882 ± 115	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+ \pi^- K^+ K^-$
------------------	-----------	--------	---------	--

#### $\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_8\Gamma_{56}/\Gamma$

NODE=M026G07  
NODE=M026G07  
NEW

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

#### 8.5 ± 0.9 OUR FIT

[9.0 ± 0.8 eV OUR 2022 FIT]

7.75 ± 0.66 ± 0.62	386 ± 31	<sup>1</sup> LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+ K^-)$
--------------------	----------	------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.8 ± 1.2 ± 1.3	132 ± 23	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(K^+ K^-)$
-----------------	----------	--------	---------	---------------------------------------

<sup>1</sup> Supersedes UEHARA 08. Using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ .

NODE=M026G07;LINKAGE=L1

$$\Gamma(\omega\gamma) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_{56}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**11.0 ± 2.5 OUR FIT**

**8.67 ± 2.86 ± 0.96** 85 ± 29 <sup>1</sup> LIU 12B BELL  $\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M026G03  
NODE=M026G03

NODE=M026G03;LINKAGE=LI

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{16}\Gamma_{56}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.49 90 <sup>1</sup> LIU 12B BELL  $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

<sup>1</sup> Using  $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$  and  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M026G04  
NODE=M026G04

NODE=M026G04;LINKAGE=LI

$$\Gamma(f_2(1270)f_2(1270)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_{56}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**52 ± 13 OUR FIT**

[50 ± 13 eV OUR 2022 FIT]

**69 ± 17 ± 12** 3182 ± 766 UEHARA 08 BELL  $\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

NODE=M026G19  
NODE=M026G19  
NEW

$$\Gamma(f_2(1270)f_2'(1525)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{18}\Gamma_{56}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**49 ± 9 ± 13** 1128 ± 206 UEHARA 08 BELL  $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

NODE=M026G20  
NODE=M026G20

$$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{34}\Gamma_{56}/\Gamma$$

VALUE (keV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
-------------	----------	-------------	------	---------

**0.378 ± 0.021 OUR FIT**

[0.374 ± 0.021 keV OUR 2022 FIT]

**0.407 ± 0.027 OUR AVERAGE** Error includes scale factor of 1.2.

0.374 ± 0.009 ± 0.031 14k <sup>1</sup> LEES 10 BABR  $10.6 e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$

0.407 ± 0.022 ± 0.028 <sup>2,3</sup> ASNER 04 CLEO  $\gamma\gamma \rightarrow \eta'_C \rightarrow K_S^0K^\pm\pi^\mp$

0.60 ± 0.12 ± 0.09 41 <sup>3,4</sup> ABDALLAH 03J DLPH  $\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$

1.47 ± 0.87 ± 0.27 <sup>3</sup> SHIRAI 98 AMY  $\gamma\gamma \rightarrow \eta_C \rightarrow K^\pm K_S^0\pi^\mp$

0.84 ± 0.21 <sup>3</sup> ALBRECHT 94H ARG  $\gamma\gamma \rightarrow K^\pm K_S^0\pi^\mp$

0.60 <sup>+0.23</sup> <sub>-0.20</sub> <sup>3</sup> CHEN 90B CLEO  $\gamma\gamma \rightarrow \eta_C K^\pm K_S^0\pi^\mp$

1.06 ± 0.41 ± 0.27 11 <sup>3</sup> BRAUNSCH... 89 TASS  $\gamma\gamma \rightarrow K\bar{K}\pi$

1.5 <sup>+0.60</sup> <sub>-0.45</sub> ± 0.3 7 <sup>3</sup> BERGER 86 PLUT  $\gamma\gamma \rightarrow K\bar{K}\pi$

NODE=M026G14  
NODE=M026G14  
NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.386 ± 0.008 ± 0.021 12k <sup>5</sup> DEL-AMO-SA...11M BABR  $\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$

0.418 ± 0.044 ± 0.022 <sup>3,6</sup> BRANDENB... 00B CLE2  $\gamma\gamma \rightarrow \eta_C \rightarrow K^\pm K_S^0\pi^\mp$

<0.63 95 <sup>3</sup> BEHREND 89 CELL  $\gamma\gamma \rightarrow K_S^0K^\pm\pi^\mp$

<4.4 95 ALTHOFF 85B TASS  $\gamma\gamma \rightarrow K\bar{K}\pi$

<sup>1</sup> From the corrected and unfolded mass spectrum.

<sup>2</sup> Calculated by us from the value reported in ASNER 04 that assumes  $B(\eta_C \rightarrow K\bar{K}\pi) = 5.5 \pm 1.7\%$

<sup>3</sup> We have multiplied  $K^\pm K_S^0\pi^\mp$  measurement by 3 to obtain  $K\bar{K}\pi$ .

<sup>4</sup> Calculated by us from the value reported in ABDALLAH 03J, which uses  $B(\eta_C \rightarrow K_S^0K^\pm\pi^\mp) = (1.5 \pm 0.4)\%$ .

<sup>5</sup> Not independent from the measurements reported by LEES 10.

<sup>6</sup> Superseded by ASNER 04.

NODE=M026G14;LINKAGE=LE  
NODE=M026G14;LINKAGE=AA

NODE=M026G14;LINKAGE=C  
NODE=M026G;LINKAGE=BB

NODE=M026G14;LINKAGE=DE  
NODE=M026G14;LINKAGE=NN

$$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{38}\Gamma_{56}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**35 ± 5 OUR FIT**

[34 ± 5 eV OUR 2022 FIT]

**27 ± 6 OUR AVERAGE**

25.7 ± 3.2 ± 4.9 2019 ± 248 UEHARA 08 BELL  $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

280 ± 100 ± 60 42 <sup>1</sup> ABDALLAH 03J DLPH  $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

170 ± 80 ± 20 13.9 ± 6.6 ALBRECHT 94H ARG  $\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

<sup>1</sup> Calculated by us from the value reported in ABDALLAH 03J, which uses  $B(\eta_C \rightarrow \pi^+\pi^-K^+K^-) = (2.0 \pm 0.7)\%$ .

NODE=M026G15  
NODE=M026G15  
NEW

NODE=M026G;LINKAGE=CC

$$\frac{\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\text{VALUE (keV)} \quad \text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}} \quad \Gamma_{39}\Gamma_{56}/\Gamma$$

NODE=M026G02  
NODE=M026G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.190 ± 0.006 ± 0.028 11k <sup>1</sup> DEL-AMO-SA..11M BABR  $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

<sup>1</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M026G02;LINKAGE=DE

$$\frac{\Gamma(2(K^+K^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\text{VALUE (eV)} \quad \text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}} \quad \Gamma_{42}\Gamma_{56}/\Gamma$$

NODE=M026G27  
NODE=M026G27

**7.4 ± 1.5 OUR FIT**  
[7.3 ± 1.5 eV OUR 2022 FIT]

**5.8 ± 1.9 OUR AVERAGE**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.6 ± 1.1 ± 1.6	216 ± 42	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$
350 ± 90 ± 60	46	<sup>1</sup> ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow 2(K^+K^-)$
231 ± 90 ± 23	9.1 ± 3.3	<sup>2</sup> ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(K^+K^-)$

<sup>1</sup> Calculated by us from the value reported in ABDALLAH 03J, which uses  $B(\eta_c \rightarrow 2(K^+K^-)) = (2.1 \pm 1.2)\%$ .

<sup>2</sup> Includes all topological modes except  $\eta_c \rightarrow \phi\phi$ .

NEW

NODE=M026G;LINKAGE=DD

NODE=M026G;LINKAGE=EE

$$\frac{\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\text{VALUE (eV)} \quad \text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}} \quad \Gamma_{45}\Gamma_{56}/\Gamma$$

NODE=M026G11  
NODE=M026G11

**47 ± 5 OUR FIT**  
[47 ± 6 eV OUR 2022 FIT]

**42 ± 6 OUR AVERAGE**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
40.7 ± 3.7 ± 5.3	5381 ± 492	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$
180 ± 70 ± 20	21.4 ± 8.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

NEW

$$\frac{\Gamma(p\bar{p}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\text{VALUE (eV)} \quad \text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}} \quad \Gamma_{48}\Gamma_{56}/\Gamma$$

NODE=M026G01  
NODE=M026G01

**7.3 ± 0.7 OUR FIT**  
[7.4 ± 0.7 eV OUR 2022 FIT]

**7.20 ± 1.53<sup>+0.67</sup><sub>-0.75</sub>** 157 ± 33 <sup>1</sup> KUO 05 BELL  $\gamma\gamma \rightarrow p\bar{p}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.6 <sup>+1.3</sup><sub>-1.1</sub> ± 0.4 190 <sup>1</sup> AMBROGIANI 03 E835  $\bar{p}p \rightarrow \gamma\gamma$

8.1 <sup>+2.9</sup><sub>-2.0</sub> <sup>1</sup> ARMSTRONG 95F E760  $\bar{p}p \rightarrow \gamma\gamma$

<sup>1</sup> Not independent from the  $\Gamma_{\gamma\gamma}$  reported by the same experiment.

NODE=M026G01;LINKAGE=GG

$$\frac{\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\text{VALUE (eV)} \quad \text{CL\%} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}} \quad \Gamma_{60}\Gamma_{56}/\Gamma$$

NODE=M026G05  
NODE=M026G05

**<1.6** 90 <sup>1</sup> UEHARA 13 BELL  $\gamma\gamma \rightarrow K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.29 90 <sup>2</sup> UEHARA 13 BELL  $\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> Taking into account interference with the non-resonant continuum.

<sup>2</sup> Neglecting interference with the non-resonant continuum.

OCCUR=2

NODE=M026G05;LINKAGE=U1

NODE=M026G05;LINKAGE=U2

## $\eta_c(1S)$ BRANCHING RATIOS

NODE=M026225

### HADRONIC DECAYS

NODE=M026305

$$\frac{\Gamma(\eta'(958)\pi\pi)/\Gamma_{\text{total}}}{\text{VALUE (units } 10^{-2}) \quad \text{EVTS} \quad \text{DOCUMENT ID} \quad \text{TECN} \quad \text{COMMENT}} \quad \Gamma_1/\Gamma$$

NODE=M026R14  
NODE=M026R14

**1.87 ± 0.26 OUR FIT**

**3.1 ± 1.2 OUR AVERAGE** [0.041 ± 0.017 OUR 2022 AVERAGE]

NEW

**3.1 ± 1.0 ± 0.7** 14 <sup>1,2</sup> BALTRUSAIT..86 MRK3  $J/\psi \rightarrow \eta_c \gamma$

<sup>1</sup> BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow \eta'(958)\pi\pi)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (5.25 \pm 1.65) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> The value reported by BALTRUSAITIS 86 has been multiplied by 3/2 to account for isospin symmetry.

NODE=M026R14;LINKAGE=A

NODE=M026R14;LINKAGE=E

$\Gamma(\rho\rho)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**1.5±0.4 OUR AVERAGE**[(18 ± 5) × 10<sup>-3</sup> OUR 2022 AVERAGE]

0.9±0.4±0.2		72	<sup>1</sup> ABLIKIM	05L BES2	$J/\psi \rightarrow \pi^+ \pi^- \pi^+ \pi^- \gamma$
2.0±0.4±0.4		113	<sup>2,3</sup> BISELLO	91 DM2	$J/\psi \rightarrow \gamma \rho^0 \rho^0$
1.8±0.8±0.4		32	<sup>4,5</sup> BISELLO	91 DM2	$J/\psi \rightarrow \gamma \rho^+ \rho^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14      90      <sup>6</sup> BALTRUSAIT..86    MRK3     $J/\psi \rightarrow \eta_c \gamma$ 

<sup>1</sup> ABLIKIM 05L reports  $[\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.6 \pm 0.6 \pm 0.4) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.30 \pm 0.30 \pm 0.60) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> The value reported by BISELLO 91 has been multiplied by 3 to account for isospin symmetry.

<sup>4</sup> BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.0 \pm 1.3 \pm 0.6) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup> The value reported by BISELLO 91 has been multiplied by 3/2 to account for isospin symmetry.

<sup>6</sup> Using  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R9

NODE=M026R9

NEW

OCCUR=2

NODE=M026R9;LINKAGE=F

NODE=M026R9;LINKAGE=A

NODE=M026R9;LINKAGE=B

NODE=M026R9;LINKAGE=C

NODE=M026R9;LINKAGE=D

NODE=M026R9;LINKAGE=G

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.5±0.5 OUR AVERAGE** [0.02 ± 0.007 OUR 2022 AVERAGE]**1.5±0.4±0.3**      63      <sup>1,2</sup> BALTRUSAIT..86    MRK3     $J/\psi \rightarrow \eta_c \gamma$ 

<sup>1</sup> BALTRUSAITIS 86 has an error according to Partridge.

<sup>2</sup> BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (2.6 \pm 0.6) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R16

NODE=M026R16

NEW

NODE=M026R;LINKAGE=03

NODE=M026R16;LINKAGE=A

 $\Gamma(K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**63±12 OUR FIT**[(69 ± 13) × 10<sup>-4</sup> OUR 2022 FIT]**69±23 OUR AVERAGE**[(91 ± 26) × 10<sup>-4</sup> OUR 2022 AVERAGE]

83±35±18		60	<sup>1</sup> ABLIKIM	05L BES2	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \gamma$
62±24±13		14	<sup>2,3</sup> BISELLO	91 DM2	$e^+ e^- \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
71±36±15		9	<sup>4,5</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

<sup>1</sup> ABLIKIM 05L reports  $[\Gamma(\eta_c(1S) \rightarrow K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.4 \pm 0.3 \pm 0.5) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.04 \pm 0.36 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> The value reported by BISELLO 91 has been multiplied by 2 to account for isospin symmetry.

<sup>4</sup> BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.2 \pm 0.6) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup> The value reported by BALTRUSAITIS 86 has been multiplied by 2 to account for isospin symmetry.

NODE=M026R8

NODE=M026R8

NEW

NEW

NODE=M026R8;LINKAGE=A

NODE=M026R8;LINKAGE=B

NODE=M026R8;LINKAGE=D

NODE=M026R8;LINKAGE=C

NODE=M026R8;LINKAGE=F

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_6 / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>113 ± 47 ± 24</b>	45	1 ABLIKIM	06A BES2	$J/\psi \rightarrow K^*0 \bar{K}^*0 \pi^+ \pi^- \gamma$

NODE=M026R25  
 NODE=M026R25

<sup>1</sup> ABLIKIM 06A reports  $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.91 \pm 0.64 \pm 0.48) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R25;LINKAGE=AB

 $\Gamma(\phi K^+ K^-) / \Gamma_{\text{total}}$  $\Gamma_7 / \Gamma$ 

VALUE (units $10^{-3}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>2.9<sup>+0.9</sup><sub>-0.8</sub> ± 1.1</b>	14.1 <sup>+4.4</sup> <sub>-3.7</sub>	1 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$

NODE=M026R21  
 NODE=M026R21

<sup>1</sup> Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12<sup>+0.10</sup><sub>-0.12</sub>) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R;LINKAGE=BB

 $\Gamma(\phi\phi) / \Gamma_{\text{total}}$  $\Gamma_8 / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>15.8 ± 1.9 OUR FIT</b>				

[(17.4 ± 1.9) × 10<sup>-4</sup> OUR 2022 FIT]

**25 ± 6 OUR AVERAGE**

[(28 ± 4) × 10<sup>-4</sup> OUR 2022 AVERAGE]

26 <sup>+4</sup> <sub>-8</sub> ± 5	1.2k	1 ABLIKIM	17P BES3	$J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$	
20 ± 5 ± 4	72	2 ABLIKIM	05L BES2	$J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$	
23 ± 7 ± 5	19	3 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
23 <sup>+14</sup> <sub>-10</sub> ± 5	5	4 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	OCCUR=2
55 ± 15 ± 12	80	5 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
50 ± 19 ± 11		6 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	OCCUR=3

NODE=M026R7  
 NODE=M026R7

NEW

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

20 ± 5 ± 4	357	7,8 BAI	04 BES	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
18 <sup>+8</sup> <sub>-6</sub> ± 7	7	9 HUANG	03 BELL	$B^+ \rightarrow (\phi\phi) K^+$	

<sup>1</sup> ABLIKIM 17P reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (4.3 \pm 0.5<sup>+0.5</sup><sub>-1.2</sub>) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=A

<sup>2</sup> ABLIKIM 05L reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (3.3 \pm 0.6 \pm 0.6) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=B

<sup>3</sup> BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.39 \pm 0.09 \pm 0.07) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=D

<sup>4</sup> BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.38<sup>+0.23</sup><sub>-0.15</sub> ± 0.07) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=F

<sup>5</sup> BAI 90B reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.93 \pm 0.20 \pm 0.16) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=G

<sup>6</sup> BAI 90B reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.85 \pm 0.27 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=H

<sup>7</sup> BAI 04 reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (3.3 \pm 0.6 \pm 0.6) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R7;LINKAGE=C

<sup>8</sup> Superseded by ABLIKIM 05L.

NODE=M026R7;LINKAGE=I

<sup>9</sup> Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12<sup>+0.10</sup><sub>-0.12</sub>) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R7;LINKAGE=BB

$\Gamma(\phi\phi)/\Gamma(K\bar{K}\pi)$  $\Gamma_8/\Gamma_{34}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.0225 ± 0.0025 OUR FIT**

[0.0240 ± 0.0025 OUR 2022 FIT]

**0.044  $\begin{smallmatrix} +0.012 \\ -0.010 \end{smallmatrix}$  OUR AVERAGE**

0.055 ± 0.014 ± 0.005

AUBERT,B 04B BABR  $B^\pm \rightarrow K^\pm \eta_c$ 0.032  $\begin{smallmatrix} +0.014 \\ -0.010 \end{smallmatrix}$  ± 0.0097 <sup>1</sup> HUANG 03 BELL  $B^\pm \rightarrow K^\pm \phi\phi$ 

<sup>1</sup> Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \begin{smallmatrix} +0.10 \\ -0.12 \end{smallmatrix}) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R39  
 NODE=M026R39  
 NEW

NODE=M026R39;LINKAGE=BB

 $\Gamma(\phi\phi)/\Gamma(p\bar{p})$  $\Gamma_8/\Gamma_{48}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**1.79 ± 0.14 ± 0.32** 6.4k <sup>1</sup> AAIJ 17BB LHCB  $p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$ 

<sup>1</sup> Using inputs from AAIJ 15AS and AAIJ 15BI and  $\Gamma(b \rightarrow J/\psi(1S)\text{anything})/\Gamma_{\text{total}} = (1.16 \pm 0.10)\%$  and  $\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} = (2.120 \pm 0.029) \times 10^{-3}$  from PDG 16.

NODE=M026R52  
 NODE=M026R52

NODE=M026R52;LINKAGE=A

 $\Gamma(\phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<40 90 <sup>1</sup> ABLIKIM 06A BES2  $J/\psi \rightarrow \phi 2(\pi^+\pi^-)\gamma$ 

<sup>1</sup> ABLIKIM 06A reports  $[\Gamma(\eta_c(1S) \rightarrow \phi 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.603 \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R26  
 NODE=M026R26

NODE=M026R26;LINKAGE=AB

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_c \rightarrow \pi^+\pi^-\eta$ seen LEES 14E BABR Dalitz anal. of  $\eta_c \rightarrow K^+K^-\pi^0$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02 90 <sup>1,2</sup> BALTRUSAIT..86 MRK3  $J/\psi \rightarrow \eta_c\gamma$ <sup>1</sup> The quoted branching ratio uses  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$ .<sup>2</sup> We are assuming  $B(a_0(980) \rightarrow \eta\pi) > 0.5$ .

NODE=M026R11  
 NODE=M026R11

NODE=M026R11;LINKAGE=E  
 NODE=M026R11;LINKAGE=F

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_c \rightarrow \pi^+\pi^-\eta$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02 90 <sup>1</sup> BALTRUSAIT..86 MRK3  $J/\psi \rightarrow \eta_c\gamma$ <sup>1</sup> The quoted branching ratio uses  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R12  
 NODE=M026R12

NODE=M026R12;LINKAGE=E

 $\Gamma(K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0128 90 BISELLO 91 DM2  $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ <0.0132 90 <sup>1</sup> BISELLO 91 DM2  $J/\psi \rightarrow \gamma K^+ K^- \pi^0$ <sup>1</sup> The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R17  
 NODE=M026R17

OCCUR=2

NODE=M026R17;LINKAGE=E

 $\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

seen [ $<0.011$  (CL = 90%) OUR 2022 BEST LIMIT]seen LEES 21A BABR Dalitz anal. of  $\eta_c \rightarrow \pi^+\pi^-\eta$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011 90 <sup>1</sup> BALTRUSAIT..86 MRK3  $J/\psi \rightarrow \eta_c\gamma$ <sup>1</sup> The quoted branching ratio uses  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R13  
 NODE=M026R13

NODE=M026R13;LINKAGE=E

 $\Gamma(f_2(1270)\eta')/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_c \rightarrow \pi^+\pi^-\eta'$ ;  $K^+K^-\eta'$

NODE=M026R60  
 NODE=M026R60

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**2.1±0.5 OUR FIT****2.9±0.5±0.6**1705 1 ABLIKIM 19AV BES3  $J/\psi \rightarrow \gamma\omega\omega$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.3 90 2 ABLIKIM 05L BES2  $J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$ <6.3 90 2 BISELLO 91 DM2  $J/\psi \rightarrow \gamma\omega\omega$ <3.1 90 2 BALTRUSAIT..86 MRK3  $J/\psi \rightarrow \eta_C\gamma$ 

<sup>1</sup> ABLIKIM 19AV reports  $[\Gamma(\eta_C(1S) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_C(1S))] = (4.90 \pm 0.17 \pm 0.77) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_C(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma\eta_C(1S)) = 0.0127 \pm 0.0036$ . Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

 $\Gamma_{15}/\Gamma$ NODE=M026R10  
NODE=M026R10 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< **2.5 × 10<sup>-4</sup>** 90 1 ABLIKIM 17P BES3  $J/\psi \rightarrow \pi^+\pi^-\pi^0 K^+ K^-\gamma$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17 × 10<sup>-4</sup> 90 2 ABLIKIM 05L BES2  $J/\psi \rightarrow \pi^+\pi^-\pi^0 K^+ K^-\gamma$ <sup>1</sup> Using  $B(J/\psi \rightarrow \gamma\eta_C) = 0.017 \pm 0.004$ .<sup>2</sup> The quoted branching ratio uses  $B(J/\psi(1S) \rightarrow \gamma\eta_C(1S)) = 0.0127 \pm 0.0036$ . $\Gamma_{16}/\Gamma$ NODE=M026R22  
NODE=M026R22 $\Gamma(f_2(1270)f_2(1270))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.97±0.25 OUR FIT**[(0.98 ± 0.25) × 10<sup>-2</sup> OUR 2022 FIT]**0.77<sup>+0.25</sup><sub>-0.30</sub>±0.17** 91.2 ± 19.8 1 ABLIKIM 04M BES  $J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$ 

<sup>1</sup> ABLIKIM 04M reports  $[\Gamma(\eta_C(1S) \rightarrow f_2(1270)f_2(1270))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_C(1S))] = (1.3 \pm 0.3<sup>+0.3</sup><sub>-0.4</sub>) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_C(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{17}/\Gamma$ NODE=M026R19  
NODE=M026R19

NEW

 $\Gamma(f_0(500)\eta)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C \rightarrow \pi^+\pi^-\eta$  $\Gamma_{19}/\Gamma$ NODE=M026R57  
NODE=M026R57 $\Gamma(f_0(500)\eta')/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C(1S) \rightarrow \pi^+\pi^-\eta'$  $\Gamma_{20}/\Gamma$ NODE=M026R58  
NODE=M026R58 $\Gamma(f_0(980)\eta)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C \rightarrow \pi^+\pi^-\eta$ seen LEES 14E BABR Dalitz anal. of  $\eta_C \rightarrow K^+K^-\eta$  $\Gamma_{21}/\Gamma$ NODE=M026R41  
NODE=M026R41 $\Gamma(f_0(980)\eta')/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C \rightarrow \pi^+\pi^-\eta'$ ,  
 $K^+K^-\eta'$  $\Gamma_{22}/\Gamma$ NODE=M026R56  
NODE=M026R56 $\Gamma(f_0(1500)\eta)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C \rightarrow \pi^+\pi^-\eta$ seen LEES 14E BABR Dalitz anal. of  $\eta_C \rightarrow K^+K^-\eta$  $\Gamma_{23}/\Gamma$ NODE=M026R42  
NODE=M026R42 $\Gamma(f_0(1710)\eta')/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C \rightarrow K^+K^-\eta'$  $\Gamma_{24}/\Gamma$ NODE=M026R59  
NODE=M026R59 $\Gamma(f_0(2100)\eta')/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen LEES 21A BABR Dalitz anal. of  $\eta_C \rightarrow \pi^+\pi^-\eta$  $\Gamma_{25}/\Gamma$ NODE=M026R61  
NODE=M026R61



$\Gamma(f_0(2200)\eta)/\Gamma_{\text{total}}$					$\Gamma_{26}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT		
seen		LEES	14E	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \eta$		NODE=M026R43 NODE=M026R43
$\Gamma(a_0(1320)\pi)/\Gamma_{\text{total}}$					$\Gamma_{27}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT		
seen		LEES	14E	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \pi^0$		NODE=M026R45 NODE=M026R45
$\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$					$\Gamma_{28}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT		
seen		LEES	21A	BABR Dalitz anal. of $\eta_C \rightarrow \pi^+ \pi^- \eta$		NODE=M026R46 NODE=M026R46
seen		LEES	14E	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \pi^0$		
$\Gamma(a_0(1700)\pi)/\Gamma_{\text{total}}$					$\Gamma_{29}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT		
seen		LEES	21A	BABR Dalitz anal. of $\eta_C \rightarrow \pi^+ \pi^- \eta'$		NODE=M026R62 NODE=M026R62
$\Gamma(a_0(1950)\pi)/\Gamma_{\text{total}}$					$\Gamma_{30}/\Gamma$	
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT		
seen		LEES	21A	BABR Dalitz anal. of $\eta_C(1S) \rightarrow \pi^+ \pi^- \eta'$		NODE=M026R00 NODE=M026R00
seen	12k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_C(1S) \rightarrow K\bar{K}\pi$		NODE=M026R00;LINKAGE=A
<sup>1</sup> From a model-independant partial wave analysis.						
$\Gamma(K_0^*(1430)\bar{K})/\Gamma_{\text{total}}$					$\Gamma_{31}/\Gamma$	
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT		
seen	12k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_C(1S) \rightarrow K\bar{K}\pi$		NODE=M026R47 NODE=M026R47
seen		LEES	14E	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \eta/\pi^0$		
<sup>1</sup> From a model-independant partial wave analysis.						
$\Gamma(K_2^*(1430)\bar{K})/\Gamma_{\text{total}}$					$\Gamma_{32}/\Gamma$	
VALUE		DOCUMENT ID	TECN	COMMENT		
seen		LEES	21A	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \eta'$		NODE=M026R48 NODE=M026R48
seen		LEES	14E	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \pi^0$		
$\Gamma(K_0^*(1950)\bar{K})/\Gamma_{\text{total}}$					$\Gamma_{33}/\Gamma$	
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT		
seen		LEES	21A	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \eta'$		NODE=M026R49 NODE=M026R49
seen	12k	<sup>1</sup> LEES	16A	BABR $\gamma\gamma \rightarrow \eta_C(1S) \rightarrow K\bar{K}\pi$		
seen		LEES	14E	BABR Dalitz anal. of $\eta_C \rightarrow K^+ K^- \eta/\pi^0$		
<sup>1</sup> From a Dalitz plot analysis using an isobar model.						
$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$					$\Gamma_{34}/\Gamma$	
VALUE (units $10^{-2}$ )	EVTs	DOCUMENT ID	TECN	COMMENT		
<b>7.0±0.4 OUR FIT</b>						NEW
[(7.3 ± 0.4) × 10 <sup>-2</sup> OUR 2022 FIT]						
<b>6.7±0.5 OUR AVERAGE</b>				Error includes scale factor of 1.1. [(6.9 ± 0.5) × 10 <sup>-2</sup> OUR 2022 AVERAGE]		NEW
6.9±0.7±0.6	146	<sup>1</sup> ABLIKIM	19AP	BES3 $h_C \rightarrow \gamma\eta_C$		
7.8±0.6±0.6	267	<sup>2</sup> ABLIKIM	19AP	BES3 $h_C \rightarrow \gamma\eta_C$		OCCUR=2
6.5±1.3±0.8	55	<sup>3,4</sup> ABLIKIM	12N	BES3 $\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^0$		
8.1±1.4 <sup>+0.9</sup> <sub>-1.0</sub>	107	<sup>5,6</sup> ABLIKIM	12N	BES3 $\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\pm$		OCCUR=2
8.5±1.8		<sup>7</sup> AUBERT	06E	BABR $B^\pm \rightarrow K^\pm X_{c\bar{c}}$		
3.9±1.0±0.8	0.6k	<sup>8,9</sup> BAI	04	BES $J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$		
5.2±1.5±1.1	33	<sup>10,11</sup> BISELLO	91	DM2 $J/\psi \rightarrow \gamma K^+ K^- \pi^0$		
4.1±1.0±0.9	68	<sup>12,13</sup> BISELLO	91	DM2 $J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$		OCCUR=2
4.6±1.8±1.0	32	<sup>14,15</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$		OCCUR=2
3.4±0.9±0.7	63	<sup>16,17</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$		OCCUR=3
13 <sup>+7</sup> <sub>-5</sub> ±2		<sup>18</sup> HIMEL	80B	MRK2 $\psi(2S) \rightarrow \eta_C \gamma$		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
4.8±1.7	95	<sup>19,20</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_C \gamma$		
< 10.7 90% CL		<sup>20</sup> PARTRIDGE	80B	CBAL $J/\psi \rightarrow \eta_C \gamma$		

- 1 ABLIKIM 19AP quotes  $B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.15 \pm 0.12 \pm 0.10) \times 10^{-2}$  which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=C
- 2 ABLIKIM 19AP quotes  $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (2.60 \pm 0.21 \pm 0.20) \times 10^{-2}$  which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=F
- 3 ABLIKIM 12N quotes  $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (4.54 \pm 0.76 \pm 0.48) \times 10^{-6}$  which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=BK
- 4 ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (27.24 \pm 4.56 \pm 2.88) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values. NODE=M026R4;LINKAGE=CK
- 5 ABLIKIM 12N quotes  $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (11.35 \pm 1.25 \pm 1.50) \times 10^{-6}$  which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=BL
- 6 ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (34.05 \pm 3.75 \pm 4.50) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values. NODE=M026R4;LINKAGE=CL
- 7 Determined from the ratio of  $B(B^\pm \rightarrow K^\pm \eta_c) B(\eta_c \rightarrow K \bar{K} \pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$  reported in AUBERT, B 04B and  $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$  reported in AUBERT 06E. NODE=M026R4;LINKAGE=AB
- 8 BAI 04 reports  $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (2.2 \pm 0.3 \pm 0.5) \times 10^{-4}$  which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=H
- 9 BAI 04 reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (6.6 \pm 0.9 \pm 1.5) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=I
- 10 BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (8.76 \pm 1.80 \pm 1.68) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=J
- 11 BISELLO 91 reports  $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.46 \pm 0.30 \pm 0.28) \times 10^{-4}$  which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=M
- 12 BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (6.9 \pm 1.2 \pm 1.2) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=L
- 13 BISELLO 91 reports  $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (2.3 \pm 0.4 \pm 0.4) \times 10^{-4}$  which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=N
- 14 BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (7.8 \pm 3.0) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=Q
- 15 BALTRUSAITIS 86 reports  $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.3 \pm 0.5) \times 10^{-4}$  which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=S
- 16 BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (5.7 \pm 1.5) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=R
- 17 BALTRUSAITIS 86 reports  $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (1.9 \pm 0.5) \times 10^{-4}$  which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=T
- 18 HIMEL 80B reports  $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \eta_c(1S))] = (4.5^{+2.4}_{-1.8}) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = (3.4 \pm 0.5) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=G
- 19 Average from  $K^+ K^- \pi^0$  and  $K^\pm K_S^0 \pi^\mp$  decay channels. NODE=M026R4;LINKAGE=D
- 20 The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$ . Where relevant, the error in this branching ratio is treated as a common systematic in computing averages. NODE=M026R4;LINKAGE=E

 $\Gamma(\phi K^+ K^-) / \Gamma(K \bar{K} \pi)$  $\Gamma_7 / \Gamma_{34}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.052^{+0.016}_{-0.014} \pm 0.014$	7	1 HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi \phi$

NODE=M026R02  
NODE=M026R02

1 Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R02;LINKAGE=BB

$\Gamma(K\bar{K}\eta)/\Gamma_{\text{total}}$  $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**1.32±0.15 OUR FIT**[(1.36 ± 0.15) × 10<sup>-2</sup> OUR 2022 FIT]

<b>1.0 ± 0.5 ± 0.1</b>		7	<sup>1,2</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta K^+ K^-$
------------------------	--	---	------------------------	----------	--

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<3.1		90	<sup>3</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
------	--	----	-----------------------------	------	------------------------------------

<sup>1</sup> ABLIKIM 12N quotes  $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \eta) = (2.11 \pm 1.01 \pm 0.32) \times 10^{-6}$  which we multiply by 2 to account for isospin symmetry.

<sup>2</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (4.22 \pm 2.02 \pm 0.64) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>3</sup> The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R15

NODE=M026R15

NEW

OCCUR=2

NODE=M026R15;LINKAGE=AK

NODE=M026R15;LINKAGE=AM

NODE=M026R15;LINKAGE=E

 $\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$  $\Gamma_{35}/\Gamma_{34}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.187±0.018 OUR FIT**

<b>0.190±0.008±0.017</b>	5.4k		<sup>1</sup> LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \eta/\pi^0$
--------------------------	------	--	-------------------	----------	---

<sup>1</sup> LEES 14E reports  $B(\eta_c(1S) \rightarrow K^+ K^- \eta)/B(\eta_c(1S) \rightarrow K^+ K^- \pi^0) = 0.571 \pm 0.025 \pm 0.051$ , which we divide by 3 to account for isospin symmetry. It uses both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

NODE=M026R40

NODE=M026R40

NODE=M026R40;LINKAGE=LE

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{36}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.7±0.5 OUR AVERAGE** [(1.7 ± 0.6) × 10<sup>-2</sup> OUR 2022 AVERAGE]

<b>1.7±0.4±0.2</b>	33		<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta \pi^+ \pi^-$
--------------------	----	--	----------------------	----------	--

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

5.4±2.0		75	<sup>2</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
---------	--	----	-----------------------------	------	------------------------------------

3.7±1.3±2.0		18	<sup>2</sup> PARTRIDGE	80B CBAL	$J/\psi \rightarrow \eta \pi^+ \pi^- \gamma$
-------------	--	----	------------------------	----------	--

<sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (7.22 \pm 1.47 \pm 1.11) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

<sup>2</sup> The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$ . Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R6

NODE=M026R6

NEW

NODE=M026R6;LINKAGE=AB

NODE=M026R6;LINKAGE=E

 $\Gamma(\eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{37}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**4.6±1.4 OUR AVERAGE** [(4.4 ± 1.6) × 10<sup>-2</sup> OUR 2022 AVERAGE]

<b>4.6±1.3±0.5</b>	39		<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta 2(\pi^+\pi^-)$
--------------------	----	--	----------------------	----------	--

<sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (19.17 \pm 3.77 \pm 3.72) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R05

NODE=M026R05

NEW

NODE=M026R05;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**6.5± 1.0 OUR FIT**[(6.6 ± 1.1) × 10<sup>-3</sup> OUR 2022 FIT]**10.6± 1.9 OUR AVERAGE**[(11.8 ± 2.3) × 10<sup>-3</sup> OUR 2022 AVERAGE]

9.9± 2.3±1.2	38		<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^+ \pi^-$
--------------	----	--	----------------------	----------	---

8.9± 1.7±1.9	0.4k		<sup>2</sup> BAI	04 BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
--------------	------	--	------------------	--------	---

16.0± 2.4±3.4	110		<sup>3</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
---------------	-----	--	-----------------------------	------	------------------------------------

12 $\frac{+17}{-7}$ ±2			<sup>4</sup> HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$
------------------------	--	--	--------------------	----------	--------------------------------------

NODE=M026R5

NODE=M026R5

NEW

NEW

- <sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (4.16 \pm 0.76 \pm 0.59) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.
- <sup>2</sup> BAI 04 reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.5 \pm 0.2 \pm 0.2) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>3</sup> BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (2.7 \pm 0.4) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>4</sup> HIMEL 80B reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\eta_c(1S))] = (4.0^{+6.0}_{-2.5}) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = (3.4 \pm 0.5) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R5;LINKAGE=AB

NODE=M026R5;LINKAGE=C

NODE=M026R5;LINKAGE=B

NODE=M026R5;LINKAGE=D

 **$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma(K\bar{K}\pi)$**  **$\Gamma_{39} / \Gamma_{34}$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.477 ± 0.017 ± 0.070</b>	11k	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M026R01  
NODE=M026R01

<sup>1</sup> We have multiplied the value of  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M026R01;LINKAGE=DE

 **$\Gamma(K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}) / \Gamma_{\text{total}}$**  **$\Gamma_{40} / \Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.7 ± 1.6 OUR AVERAGE</b>		[(5.6 ± 1.9) × 10 <sup>-2</sup> OUR 2022 AVERAGE]		
<b>5.7 ± 1.4 ± 0.7</b>	43	<sup>1,2</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\mp 2\pi^\pm$

NODE=M026R06  
NODE=M026R06

NEW

<sup>1</sup> ABLIKIM 12N quotes  $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^- \pi^- 2\pi^+)$  =  $(12.01 \pm 2.22 \pm 2.04) \times 10^{-6}$  which we multiply by 2 to take c.c. into account.

NODE=M026R06;LINKAGE=AA

<sup>2</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (24.02 \pm 4.44 \pm 4.08) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R06;LINKAGE=AB

 **$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$**  **$\Gamma_{41} / \Gamma$** 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.6 ± 2.4 OUR AVERAGE</b>		[(7.5 ± 2.4) × 10 <sup>-3</sup> OUR 2022 AVERAGE]		
9 ± 4 ± 1	10	<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- 2(\pi^+ \pi^-)$
7.2 ± 2.4 ± 1.5	100	<sup>2</sup> ABLIKIM	06A BES2	$J/\psi \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

NODE=M026R23  
NODE=M026R23

NEW

<sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (3.60 \pm 1.71 \pm 0.64) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R23;LINKAGE=AL

<sup>2</sup> ABLIKIM 06A reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.21 \pm 0.32 \pm 0.24) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R23;LINKAGE=AB

 **$\Gamma(2(K^+ K^-)) / \Gamma_{\text{total}}$**  **$\Gamma_{42} / \Gamma$** 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.38 ± 0.29 OUR FIT</b>		[(1.43 ± 0.30) × 10 <sup>-3</sup> OUR 2022 FIT]		
<b>2.2 ± 0.9 ± 0.3</b>	7	<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(K^+ K^-)$

NODE=M026R20  
NODE=M026R20

NEW

- • • We do not use the following data for averages, fits, limits, etc. • • •
- |   |  |                       |         |  |
|---|--|-----------------------|---------|--|
| 1.4 <sup>+0.5</sup> / <sub>-0.4</sub> ± 0.6 | 14.5 <sup>+4.6</sup> / <sub>-3.0</sub> | <sup>2</sup> HUANG    | 03 BELL | $B^+ \rightarrow 2(K^+ K^-) K^+$           |
| 21 ± 10 ± 6                                 |  | <sup>3</sup> ALBRECHT | 94H ARG | $\gamma\gamma \rightarrow K^+ K^- K^+ K^-$ |

- <sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (0.94 \pm 0.37 \pm 0.14) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.
- <sup>2</sup> Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$ .
- <sup>3</sup> Normalized to the sum of  $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$ ,  $B(\eta_c \rightarrow \phi\phi)$ ,  $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$ , and  $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$ .

NODE=M026R20;LINKAGE=AB

NODE=M026R20;LINKAGE=BB

NODE=M026R20;LINKAGE=AL

 **$\Gamma(2(K^+ K^-))/\Gamma(K\bar{K}\pi)$**  **$\Gamma_{42}/\Gamma_{34}$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.020±0.004 OUR FIT****0.024±0.007 OUR AVERAGE**

0.023±0.007±0.006		AUBERT,B	04B	BABR $B^\pm \rightarrow K^\pm \eta_c$
0.026 <sup>+0.009</sup> <sub>-0.007</sub> ±0.007	15	<sup>1</sup> HUANG	03	BELL $B^\pm \rightarrow K^\pm(2K^+ 2K^-)$

- <sup>1</sup> Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R38  
NODE=M026R38

NODE=M026R38;LINKAGE=BB

 **$\Gamma(\eta'(958) K\bar{K})/\Gamma(\eta'(958) \pi\pi)$**  **$\Gamma_2/\Gamma_1$** 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.859±0.052±0.043**

<sup>1</sup> LEES	21A	BABR	$\gamma\gamma \rightarrow \eta' K^+ K^-$ , $\eta' \pi^+ \pi^-$
-------------------	-----	------	---

- <sup>1</sup> Based on Dalitz-plot analysis of the  $\eta_c \rightarrow \eta' K^+ K^-$ ,  $\eta' \pi^+ \pi^-$  final states where the fit fractions and relative phases are determined for numerous two-body intermediate states.

NODE=M026R55  
NODE=M026R55

NODE=M026R55;LINKAGE=A

 **$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{43}/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<5 × 10<sup>-4</sup>** 90 <sup>1</sup> ABLIKIM 17AJ BES3  $\psi(2S) \rightarrow \gamma\pi^+ \pi^- \pi^0$ 

- <sup>1</sup> ABLIKIM 17AJ reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\eta_c(1S))] < 1.6 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = 3.4 \times 10^{-3}$ .

NODE=M026R51  
NODE=M026R51

NODE=M026R51;LINKAGE=A

 **$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{44}/\Gamma$** 

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

**4.8±1.1 OUR AVERAGE** [(4.7 ± 1.4) × 10<sup>-2</sup> OUR 2022 AVERAGE]**4.8±1.0±0.6** 118 <sup>1</sup> ABLIKIM 12N BES3  $\psi(2S) \rightarrow \pi^0 \gamma \pi^+ \pi^- 2\pi^0$ 

- <sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (20.31 \pm 2.20 \pm 3.33) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R07  
NODE=M026R07

NEW

NODE=M026R07;LINKAGE=AB

 **$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}}$**  **$\Gamma_{45}/\Gamma$** 

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

**0.87±0.11 OUR FIT**[(0.91 ± 0.12) × 10<sup>-2</sup> OUR 2022 FIT]**1.08±0.22 OUR AVERAGE** Error includes scale factor of 1.2. [(1.27 ± 0.23) × 10<sup>-2</sup> OUR 2022 AVERAGE]

1.79±0.33±0.21	100	<sup>1</sup> ABLIKIM	12N	BES3 $\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^-)$
0.77±0.27±0.17	0.5k	<sup>2</sup> BAI	04	BES $J/\psi \rightarrow \gamma 2(\pi^+ \pi^-)$
0.79±0.18±0.17	137	<sup>3</sup> BISELLO	91	DM2 $J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$
0.9 ± 0.4 ± 0.2	25	<sup>4</sup> BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
1.7 <sup>+1.1</sup> <sub>-0.7</sub> ± 0.2		<sup>5</sup> HIMEL	80B	MRK2 $\psi(2S) \rightarrow \eta_c \gamma$

- <sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (7.51 \pm 0.85 \pm 1.11) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R1  
NODE=M026R1

NEW

NEW

- <sup>2</sup> BAI 04 reports  $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.3 \pm 0.2 \pm 0.4) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R1;LINKAGE=B

<sup>3</sup>BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.33 \pm 0.22 \pm 0.20) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R1;LINKAGE=C

<sup>4</sup>BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.6 \pm 0.6) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R1;LINKAGE=D

<sup>5</sup>HIMEL 80B reports  $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\eta_c(1S))] = (5.7^{+3.9}_{-2.4}) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = (3.4 \pm 0.5) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R1;LINKAGE=F

 **$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$**  **$\Gamma_{46}/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**16.2±2.1 OUR AVERAGE**[(15.8 ± 2.3) × 10<sup>-2</sup> OUR 2022 AVERAGE]

15.3±1.8±1.8	333	ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$
17.9±3.0±2.1	175	<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma 2(\pi^+\pi^-\pi^0)$

NODE=M026R08

NODE=M026R08

NEW

<sup>1</sup>ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (75.13 \pm 7.42 \pm 9.99) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R08;LINKAGE=AB

 **$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$**  **$\Gamma_{47}/\Gamma$** 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**18 ±4 OUR AVERAGE**[(17 ± 4) × 10<sup>-3</sup> OUR 2022 AVERAGE]

21 ±5 ±2	51	<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma 3(\pi^+\pi^-)$
15.4±3.4±3.3	479	<sup>2</sup> ABLIKIM	06A BES2	$J/\psi \rightarrow 3(\pi^+\pi^-)\gamma$

NODE=M026R24

NODE=M026R24

NEW

<sup>1</sup>ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (8.82 \pm 1.57 \pm 1.59) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R24;LINKAGE=AL

<sup>2</sup>ABLIKIM 06A reports  $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (2.59 \pm 0.32 \pm 0.47) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R24;LINKAGE=AB

 **$\Gamma(p\bar{p})/\Gamma_{\text{total}}$**  **$\Gamma_{48}/\Gamma$** 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**13.5± 1.3 OUR FIT**[(14.4 ± 1.4) × 10<sup>-4</sup> OUR 2022 FIT]**11.5± 1.9 OUR AVERAGE**[(12.6 ± 2.1) × 10<sup>-4</sup> OUR 2022 AVERAGE]

12.0± 2.6±1.5	34	ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$
15 ± 5 ±2	15	<sup>1</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma p\bar{p}$
11.3± 2.5±2.4	213	<sup>2</sup> BAI	04 BES	$J/\psi \rightarrow \gamma p\bar{p}$
7.7± 3.0±1.7	18	<sup>3</sup> BISELLO	91 DM2	$J/\psi \rightarrow \gamma p\bar{p}$
8 ± 4 ±2	23	<sup>4</sup> BALTRUSAIT	86 MRK3	$J/\psi \rightarrow \eta_c\gamma$
23 <sup>+23</sup> <sub>-12</sub> ±3		<sup>5</sup> HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$

NODE=M026R2

NODE=M026R2

NEW

NEW

••• We do not use the following data for averages, fits, limits, etc. •••

13.1 <sup>+</sup> <sub>-2.1</sub> ±1.0	195	<sup>6</sup> WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
--	-----	-----------------	---------	-------------------------------

<sup>1</sup>ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (0.65 \pm 0.19 \pm 0.10) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R2;LINKAGE=AB

<sup>2</sup>BAI 04 reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.9 \pm 0.3 \pm 0.3) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=C

<sup>3</sup>BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (0.13 \pm 0.04 \pm 0.03) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup>BALTRUSAITIS 86 reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.4 \pm 0.7) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup>HIMEL 80B reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\eta_c(1S))] = (8_{-4}^{+8}) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = (3.4 \pm 0.5) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>6</sup>WU 06 reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (1.42 \pm 0.11_{-0.20}^{+0.16}) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c K^+) = (1.08 \pm 0.08) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=D

NODE=M026R2;LINKAGE=F

NODE=M026R2;LINKAGE=G

NODE=M026R2;LINKAGE=WU

 **$\Gamma(p\bar{p})/\Gamma(K\bar{K}\pi)$**  **$\Gamma_{48}/\Gamma_{34}$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0192±0.0019 OUR FIT</b>				
[0.0198 ± 0.0019 OUR 2022 FIT]				

NODE=M026R03  
NODE=M026R03  
NEW

<b>0.021 ± 0.002</b>	<b>+0.004</b> <b>-0.006</b>	195	<sup>1</sup> WU	06	BELL	$B^\pm \rightarrow K^\pm p\bar{p}$
----------------------	--------------------------------	-----	-----------------	----	------	------------------------------------

<sup>1</sup>Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12_{-0.12}^{+0.10}) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R03;LINKAGE=BB

 **$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$**  **$\Gamma_{48}/\Gamma \times \Gamma_8/\Gamma$** 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.214±0.035 OUR FIT</b>			
[(0.25 ± 0.04) × 10 <sup>-5</sup> OUR 2022 FIT]			

NODE=M026R33  
NODE=M026R33  
NEW

<b>4.0</b>	<b>+3.5</b> <b>-3.2</b>	BAGLIN	89	SPEC	$\bar{p}p \rightarrow K^+ K^- K^+ K^-$
------------	----------------------------	--------	----	------	--

 **$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{49}/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.36±0.14 OUR AVERAGE</b>				[(0.36 ± 0.15) × 10 <sup>-2</sup> OUR 2022 AVERAGE]

NODE=M026R09  
NODE=M026R09  
NEW

<b>0.36±0.13±0.04</b>	14	<sup>1</sup> ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}\pi^0$
-----------------------	----	----------------------	-----	------	---

<sup>1</sup>ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (1.53 \pm 0.49 \pm 0.23) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R09;LINKAGE=AB

 **$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$**  **$\Gamma_{50}/\Gamma$** 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.2±2.3 OUR FIT</b>					
[(10.6 ± 2.3) × 10 <sup>-4</sup> OUR 2022 FIT]					

NODE=M026R18  
NODE=M026R18  
NEW

<b>11.8±2.3±2.5</b>		<sup>1</sup> ABLIKIM	12B	BES3	
---------------------	--	----------------------	-----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$8.8_{-2.3}^{+2.4} \pm 0.7$	20	<sup>2</sup> WU	06	BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
<20	90	<sup>3</sup> BISELLO	91	DM2	$e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$

<sup>1</sup>ABLIKIM 12B reports  $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (0.198 \pm 0.021 \pm 0.032) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R18;LINKAGE=AB

<sup>2</sup>WU 06 reports  $[\Gamma(\eta_c(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (0.95_{-0.22-0.11}^{+0.25+0.08}) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c K^+) = (1.08 \pm 0.08) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R18;LINKAGE=WU

<sup>3</sup>The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R18;LINKAGE=E

$\Gamma(\Lambda\bar{\Lambda})/\Gamma(p\bar{p})$  $\Gamma_{50}/\Gamma_{48}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.76±0.17 OUR FIT</b> [0.74 ± 0.16 OUR 2022 FIT]			

NODE=M026R27  
 NODE=M026R27  
 NEW

**0.67<sup>+0.19</sup><sub>-0.16</sub>±0.12**

<sup>1</sup> WU 06 BELL  $B^+ \rightarrow p\bar{p}K^+, \Lambda\bar{\Lambda}K^+$

<sup>1</sup> Not independent from other  $\eta_c \rightarrow \Lambda\bar{\Lambda}, p\bar{p}$  branching ratios reported by WU 06.

NODE=M026R27;LINKAGE=WU

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{51}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.51<sup>+0.34</sup><sub>-0.32</sub>±0.19</b>	157	<sup>1</sup> LU	19	BELL $B^+ \rightarrow \bar{p}\Lambda K^+ K^+$

NODE=M026R53  
 NODE=M026R53

<sup>1</sup> LU 19 reports  $(2.83_{-0.34}^{+0.36} \pm 0.35) \times 10^{-3}$  from a measurement of  $[\Gamma(\eta_c(1S) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)]$  assuming  $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$ , which we rescale to our best value  $B(B^+ \rightarrow \eta_c K^+) = (1.08 \pm 0.08) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R53;LINKAGE=A

 $\Gamma(\bar{\Lambda}(1520)\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{52}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.1±1.3±0.2</b>	43	<sup>1</sup> LU	19	BELL $B^+ \rightarrow \bar{p}\Lambda K^+ K^+$

NODE=M026R54  
 NODE=M026R54

<sup>1</sup> LU 19 reports  $(3.48 \pm 1.48 \pm 0.46) \times 10^{-3}$  from a measurement of  $[\Gamma(\eta_c(1S) \rightarrow \bar{\Lambda}(1520)\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)]$  assuming  $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$ , which we rescale to our best value  $B(B^+ \rightarrow \eta_c K^+) = (1.08 \pm 0.08) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R54;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$  $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1±0.3±0.5</b>	112	<sup>1</sup> ABLIKIM	13C	BES3 $J/\psi \rightarrow \gamma p\bar{p}\pi^0\pi^0$

NODE=M026R28  
 NODE=M026R28

<sup>1</sup> ABLIKIM 13C reports  $[\Gamma(\eta_c(1S) \rightarrow \Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.60 \pm 0.48 \pm 0.31) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R28;LINKAGE=AB

 $\Gamma(\Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}$  $\Gamma_{54}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.90±0.18±0.19</b>	78	<sup>1</sup> ABLIKIM	13C	BES3 $J/\psi \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M026R29  
 NODE=M026R29

<sup>1</sup> ABLIKIM 13C reports  $[\Gamma(\eta_c(1S) \rightarrow \Xi^-\bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.51 \pm 0.27 \pm 0.14) \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R29;LINKAGE=AB

 $\Gamma(\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$  $\Gamma_{55}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.5±1.9 OUR AVERAGE</b>					[(5.3 ± 2.1) × 10 <sup>-3</sup> OUR 2022 AVERAGE]

NODE=M026R3  
 NODE=M026R3

<b>5.5±1.8±0.6</b>	19	<sup>1</sup> ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0\gamma p\bar{p}\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••					
<12	90	HIMEL	80B	MRK2	$\psi(2S) \rightarrow \eta_c\gamma$

NEW

<sup>1</sup> ABLIKIM 12N reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (2.30 \pm 0.65 \pm 0.36) \times 10^{-6}$  which we divide by our best values  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ ,  $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (57 \pm 5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R3;LINKAGE=AB

————— **RADIATIVE DECAYS** —————

NODE=M026310

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.68±0.12 OUR FIT</b>					[(1.61 ± 0.12) × 10 <sup>-4</sup> OUR 2022 FIT]

NODE=M026R31  
 NODE=M026R31

**2.2<sup>+0.9</sup><sub>-0.6</sub> OUR AVERAGE**

[(1.9<sup>+0.7</sup><sub>-0.6</sub>) × 10<sup>-4</sup> OUR 2022 AVERAGE]

NEW

2.7 ± 0.8 ± 0.6		<sup>1</sup> ABLIKIM	13i	BES3	
0.7 <sup>+1.6</sup> <sub>-0.7</sub> ± 0.2	1.2 <sup>+2.8</sup> <sub>-1.1</sub>	<sup>2</sup> ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+\pi^-J/\psi$

NEW



• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0	$\begin{smallmatrix} +0.9 \\ -0.7 \end{smallmatrix} \pm 0.2$	13	3	WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
2.80	$\begin{smallmatrix} +0.67 \\ -0.58 \end{smallmatrix} \pm 1.0$		4	ARMSTRONG	95F	E760	$\bar{p}p \rightarrow \gamma \gamma$
< 9	90		5	BISELLO	91	DM2	$J/\psi \rightarrow \gamma \gamma \gamma$
6	$\begin{smallmatrix} +4 \\ -3 \end{smallmatrix} \pm 4$		4	BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma \gamma$
< 18	90		6	BLOOM	83	CBAL	$J/\psi \rightarrow \eta_c \gamma$

<sup>1</sup> ABLIKIM 13I reports  $[\Gamma(\eta_c(1S) \rightarrow \gamma \gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (4.5 \pm 1.2 \pm 0.6) \times 10^{-6}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R31;LINKAGE=AL

<sup>2</sup> ADAMS 08 reports  $[\Gamma(\eta_c(1S) \rightarrow \gamma \gamma)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.2 \begin{smallmatrix} +2.7 \\ -1.1 \end{smallmatrix} \pm 0.3) \times 10^{-6}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R31;LINKAGE=AD

<sup>3</sup> WICHT 08 reports  $[\Gamma(\eta_c(1S) \rightarrow \gamma \gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (2.2 \begin{smallmatrix} +0.9+0.4 \\ -0.7-0.2 \end{smallmatrix}) \times 10^{-7}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c K^+) = (1.08 \pm 0.08) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R31;LINKAGE=WI

<sup>4</sup> Not independent from the values of the total and two-photon width quoted by the same experiment.

NODE=M026R31;LINKAGE=AB

<sup>5</sup> The quoted branching ratios use  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R31;LINKAGE=E

<sup>6</sup> Using  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$ .

NODE=M026R31;LINKAGE=C

### $\Gamma(\gamma \gamma)/\Gamma(K \bar{K} \pi)$

$\Gamma_{56}/\Gamma_{34}$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.40 ± 0.28 OUR FIT</b>				
$[(2.22 \pm 0.25) \times 10^{-3}]$				OUR 2022 FIT

NODE=M026R04

NODE=M026R04

NEW

3.2	$\begin{smallmatrix} +1.3 \\ -1.0 \end{smallmatrix} \begin{smallmatrix} +0.8 \\ -0.6 \end{smallmatrix}$	13	1	WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
-----	---	----	---	-------	----	------	---

<sup>1</sup> Using  $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12 \begin{smallmatrix} +0.10 \\ -0.12 \end{smallmatrix}) \times 10^{-3}$  from FANG 03 and  $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$ .

NODE=M026R04;LINKAGE=BB

### $\Gamma(p \bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma \gamma)/\Gamma_{\text{total}}$

$\Gamma_{48}/\Gamma \times \Gamma_{56}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.228 ± 0.022 OUR FIT</b>				
$[(0.232 \pm 0.022) \times 10^{-6}]$				OUR 2022 FIT

NODE=M026R32

NODE=M026R32

NEW

**0.26 ± 0.05 OUR AVERAGE** Error includes scale factor of 1.4.

0.224	$\begin{smallmatrix} +0.038 \\ -0.037 \end{smallmatrix} \pm 0.020$	190		AMBROGIANI	03	E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma \gamma$
0.336	$\begin{smallmatrix} +0.080 \\ -0.070 \end{smallmatrix}$			ARMSTRONG	95F	E760	$\bar{p}p \rightarrow \gamma \gamma$
0.68	$\begin{smallmatrix} +0.42 \\ -0.31 \end{smallmatrix}$	12		BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma \gamma$

————— Charge conjugation (C), Parity (P), —————  
 ————— Lepton family number (LF) violating modes —————

NODE=M026320

### $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$

$\Gamma_{57}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 11</b>	90	<sup>1</sup> ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^-$

NODE=M026R34

NODE=M026R34

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 70	90	<sup>2</sup> ABLIKIM	06B	BES2 $J/\psi \rightarrow \pi^+ \pi^- \gamma$
------	----	----------------------	-----	--

<sup>1</sup> ABLIKIM 11G reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 1.82 \times 10^{-6}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R34;LINKAGE=AL

<sup>2</sup> ABLIKIM 06B reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 1.1 \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R34;LINKAGE=AB

### $\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$

$\Gamma_{58}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 4</b>	90	<sup>1</sup> ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma \pi^0 \pi^0$

NODE=M026R35

NODE=M026R35

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 40	90	<sup>2</sup> ABLIKIM	06B	BES2 $J/\psi \rightarrow \pi^0 \pi^0 \gamma$
------	----	----------------------	-----	--

<sup>1</sup> ABLIKIM 11G reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 6.0 \times 10^{-7}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R35;LINKAGE=AL

<sup>2</sup> ABLIKIM 06B reports  $[\Gamma(\eta_c(1S) \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.71 \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R35;LINKAGE=AB

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<60	90	<sup>1</sup> ABLIKIM 06B	BES2	$J/\psi \rightarrow K^+ K^- \gamma$

NODE=M026R36  
NODE=M026R36

<sup>1</sup> ABLIKIM 06B reports  $[\Gamma(\eta_c(1S) \rightarrow K^+ K^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.96 \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R36;LINKAGE=AB

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{60}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<31	90	<sup>1</sup> ABLIKIM 06B	BES2	$J/\psi \rightarrow K_S^0 K_S^0 \gamma$

NODE=M026R37  
NODE=M026R37

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32 90 <sup>2</sup> UEHARA 13 BELL  $\gamma\gamma \rightarrow K_S^0 K_S^0$

< 5.6 90 <sup>3</sup> UEHARA 13 BELL  $\gamma\gamma \rightarrow K_S^0 K_S^0$

OCCUR=2

<sup>1</sup> ABLIKIM 06B reports  $[\Gamma(\eta_c(1S) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.53 \times 10^{-5}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.7 \times 10^{-2}$ .

NODE=M026R37;LINKAGE=AB

<sup>2</sup> Taking into account interference with the non-resonant continuum.

NODE=M026R37;LINKAGE=U1

<sup>3</sup> Neglecting interference with the non-resonant continuum.

NODE=M026R37;LINKAGE=U2

 $\eta_c(1S)$  CROSS-PARTICLE BRANCHING RATIOS

NODE=M026230

$$\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_8/\Gamma \times \Gamma_{222}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M026R80  
NODE=M026R80

**0.37<sup>+0.05</sup><sub>-0.07</sub> OUR AVERAGE**

0.43 ± 0.05<sup>+0.05</sup><sub>-0.12</sub> 1.2k ABLIKIM 17P BES3  $J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$

0.33 ± 0.06 ± 0.06 72 ABLIKIM 05L BES2  $J/\psi \rightarrow K^+ K^- K^+ K^- \gamma$

0.39 ± 0.09 ± 0.07 19 BISELLO 91 DM2  $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

23<sup>+14</sup><sub>-10</sub> ± 5 5 <sup>1</sup> BISELLO 91 DM2  $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

OCCUR=2

55 ± 15 ± 12 80 <sup>2</sup> BAI 90B MRK3  $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

50 ± 19 ± 11 3 BAI 90B MRK3  $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.33 ± 0.06 ± 0.06 357 <sup>4</sup> BAI 04 BES  $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>1</sup> BISELLO 91 reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.38<sup>+0.23</sup><sub>-0.15</sub> ± 0.07) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R80;LINKAGE=F

<sup>2</sup> BAI 90B reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.93 \pm 0.20 \pm 0.16) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R80;LINKAGE=G

<sup>3</sup> BAI 90B reports  $[\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (0.85 \pm 0.27 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.7 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R80;LINKAGE=H

<sup>4</sup> Superseded by ABLIKIM 05L.

NODE=M026R80;LINKAGE=E

 $\eta_c(1S)$  REFERENCES

NODE=M026

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61442
AAIJ	20H	EPJ C80 191	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60419
ABLIKIM	19AP	PR D100 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59901
ABLIKIM	19AV	PR D100 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59998
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)	REFID=59614
XU	18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)	REFID=59453
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58322
ABLIKIM	17P	PR D95 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57989
LEES	16A	PR D93 012005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57125
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
AAIJ	15AS	JHEP 1510 053	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56857

AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
ABLIKIM	13C	PR D87 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54878
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54954
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54267
ABLIKIM	12F	PRL 108 222002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54271
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54741
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53711
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
VINOKUROVA	11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=53927
LEES	10	PR D81 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53236
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52267
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ABLIKIM	06A	PL B633 19	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50987
ABLIKIM	06B	EPJ C45 337	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50988
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05L	PR D72 072005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50837
KUO	05	PL B621 41	C.C. Kuo <i>et al.</i>	(BELLE Collab.)	REFID=50801
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=49745
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49746
AUBERT,B	04B	PR D70 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50043
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49620
ABDALLAH	03J	EPJ C31 481	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49625
AMBROGIANI	03	PL B566 45	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=49465
BAI	03	PL B555 174	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49185
FANG	03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)	REFID=49206
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAI	00F	PR D62 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48546
BRANDENB...	00B	PRL 85 3095	G. Brandenburg <i>et al.</i>	(CLEO Collab.)	REFID=48553
ACCIARRI	99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47476
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46553
SHIRAI	98	PL B424 405	M. Shirai <i>et al.</i>	(AMY Collab.)	REFID=46381
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=44623
ALBRECHT	94H	PL B338 390	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44098
ADRIANI	93N	PL B318 575	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=43670
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
CHEN	90B	PL B243 169	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=41360
BAGLIN	89	PL B231 557	C. Baglin, S. Baird, G. Bassompierre	(R704 Collab.)	REFID=41015
BEHREND	89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40732
BRAUNSCH...	89	ZPHY C41 533	W. Braunschweig <i>et al.</i>	(TASSO Collab.)	REFID=40728
AIHARA	88D	PRL 60 2355	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=40588
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BERGER	86	PL 167B 120	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22010
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BALTRUSAIT...	84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+) JP	REFID=22006
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
HIMEL	80B	PRL 45 1146	T.M. Himel <i>et al.</i>	(SLAC, LBL, UCB)	REFID=22003
PARTRIDGE	80B	PRL 45 1150	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22004

**J/ψ(1S)**

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M070

**J/ψ(1S) MASS**

NODE=M070M

NODE=M070M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3096.900 ± 0.006 OUR AVERAGE</b>				
3096.900 ± 0.002 ± 0.006		<sup>1</sup> ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ± 0.09	502	<sup>2</sup> ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ± 0.03 ± 0.01		<sup>3</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ± 0.1 ± 0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.66 ± 0.19 ± 0.02	6.1k	<sup>4</sup> AAIJ 15Bl	LHCB	$pp \rightarrow J/\psi X$
3096.917 ± 0.010 ± 0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ± 0.3		GRIBUSHIN 96	FMPS	$515 \pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ± 2.0	38k	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3096.93 ± 0.09	502	<sup>5</sup> ZHOLENTZ 80	REDE	$e^+e^-$
3097.0 ± 1		<sup>6</sup> BRANDELIK 79c	DASP	$e^+e^-$

<sup>1</sup>Supersedes AULCHENKO 03.<sup>2</sup>Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).<sup>3</sup>Mass central value and systematic error recalculated by us according to Eq.(16) in ARMSTRONG 93B, using the value for the  $\psi(2S)$  mass from AULCHENKO 03.<sup>4</sup>From a sample of  $\eta_c(1S)$  and  $J/\psi$  produced in  $b$ -hadron decays. Systematic uncertainties not estimated.<sup>5</sup>Superseded by ARTAMONOV 00.<sup>6</sup>From a simultaneous fit to  $e^+e^-$ ,  $\mu^+\mu^-$  and hadronic channels assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ .NODE=M070M;LINKAGE=A  
NODE=M070M;LINKAGE=AR

NODE=M070M;LINKAGE=NW

NODE=M070M;LINKAGE=B

NODE=M070M;LINKAGE=RZ  
NODE=M070M;LINKAGE=FF**J/ψ(1S) WIDTH**

NODE=M070W

NODE=M070W

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>92.6 ± 1.7 OUR AVERAGE</b>				Error includes scale factor of 1.1.
92.45 ± 1.40 ± 1.48		<sup>1</sup> ANASHIN 20	KEDR	$e^+e^-$
96.1 ± 3.2	13k	<sup>2</sup> ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4 ± 8.9		BAI 95B	BES	$e^+e^-$
91 ± 11 ± 6		<sup>3</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 $\begin{matrix} + 6.1 \\ - 5.8 \end{matrix}$		<sup>4</sup> HSUEH 92	RVUE	See $\Upsilon$ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
92.94 ± 1.83		<sup>5,6</sup> ANASHIN 18A	KEDR	$e^+e^-$
94.1 ± 2.7		<sup>7</sup> ANASHIN 10	KEDR	$3.097 e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7 ± 3.5	7.8k	<sup>2</sup> AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

<sup>1</sup>Based on the same dataset as ANASHIN 18A and correlated to the values reported there<sup>2</sup>Calculated by us from the reported values of  $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$  using  $B(e^+e^-) = (5.94 \pm 0.06)\%$  and  $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$ .<sup>3</sup>The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].<sup>4</sup>Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.<sup>5</sup>Using  $\Gamma(e^+e^-)$  from ANASHIN 18A and  $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$  from PDG 16.<sup>6</sup>Superseded by ANASHIN 20 that is based on the same dataset.<sup>7</sup>Assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$  and using  $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$ .

NODE=M070W;LINKAGE=C

NODE=M070W;LINKAGE=AA

NODE=M070W;LINKAGE=AN  
NODE=M070W;LINKAGE=A

NODE=M070W;LINKAGE=B

NODE=M070W;LINKAGE=D  
NODE=M070W;LINKAGE=AS**J/ψ(1S) DECAY MODES**

NODE=M070215;NODE=M070

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons	(87.7 ± 0.5) %	DESIG=3
$\Gamma_2$ virtual $\gamma \rightarrow \text{hadrons}$	(13.50 ± 0.30) %	DESIG=4
$\Gamma_3$ $ggg$	(64.1 ± 1.0) %	DESIG=249
$\Gamma_4$ $\gamma gg$	( 8.8 ± 1.1) %	DESIG=250
$\Gamma_5$ $e^+e^-$	( 5.971 ± 0.032) %	DESIG=1
$\Gamma_6$ $e^+e^-\gamma$	[a] ( 8.8 ± 1.4) × 10 <sup>-3</sup>	DESIG=5
$\Gamma_7$ $\mu^+\mu^-$	( 5.961 ± 0.033) %	DESIG=2

## Decays involving hadronic resonances

Decays involving hadronic resonances				NODE=M070;CLUMP=A
Γ <sub>8</sub>	$\rho\pi$	( 1.69 ± 0.15 ) %	S=2.4	DESIG=20
Γ <sub>9</sub>	$\rho^0\pi^0$	( 5.6 ± 0.7 ) × 10 <sup>-3</sup>		DESIG=21
Γ <sub>10</sub>	$a_2(1320)^0\pi^+\pi^- \rightarrow$ $2(\pi^+\pi^-)\pi^0$	( 2.8 ± 0.6 ) × 10 <sup>-3</sup>		DESIG=442
Γ <sub>11</sub>	$a_2(1320)^+\pi^-\pi^0 + \text{c.c} \rightarrow$ $2(\pi^+\pi^-)\pi^0$	( 3.7 ± 0.7 ) × 10 <sup>-3</sup>		DESIG=443
Γ <sub>12</sub>	$a_2(1320)\rho$	( 1.09 ± 0.22 ) %		DESIG=43
Γ <sub>13</sub>	$\eta\pi^+\pi^-$	( 3.8 ± 0.7 ) × 10 <sup>-4</sup>		DESIG=239
Γ <sub>14</sub>	$\eta\pi^+\pi^-\pi^0$	( 1.17 ± 0.20 ) %		DESIG=420
Γ <sub>15</sub>	$\eta\pi^+\pi^-3\pi^0$	( 4.9 ± 1.0 ) × 10 <sup>-3</sup>		DESIG=422
Γ <sub>16</sub>	$\eta\rho$	( 1.93 ± 0.23 ) × 10 <sup>-4</sup>		DESIG=22
Γ <sub>17</sub>	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow$ $\eta\phi\pi^+\pi^-$	( 1.2 ± 0.4 ) × 10 <sup>-4</sup>		DESIG=287
Γ <sub>18</sub>	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0\bar{K}^*(892)^0$	< 2.52 × 10 <sup>-4</sup>	CL=90%	DESIG=253
Γ <sub>19</sub>	$\eta K^\pm K_S^0\pi^\mp$	[b] ( 2.2 ± 0.4 ) × 10 <sup>-3</sup>		DESIG=230
Γ <sub>20</sub>	$\eta K^*(892)^0\bar{K}^*(892)^0$	( 1.15 ± 0.26 ) × 10 <sup>-3</sup>		DESIG=252
Γ <sub>21</sub>	$\rho\eta'(958)$	( 8.1 ± 0.8 ) × 10 <sup>-5</sup>	S=1.6	DESIG=23
Γ <sub>22</sub>	$\rho^\pm\pi^\mp\pi^+\pi^-2\pi^0$	( 2.8 ± 0.8 ) %		DESIG=415
Γ <sub>23</sub>	$\rho^+\rho^-\pi^+\pi^-\pi^0$	( 6 ± 4 ) × 10 <sup>-3</sup>		DESIG=416
Γ <sub>24</sub>	$\rho^+K^+K^-\pi^- + \text{c.c} \rightarrow$ $K^+K^-\pi^+\pi^-\pi^0$	( 3.5 ± 0.8 ) × 10 <sup>-3</sup>		DESIG=444
Γ <sub>25</sub>	$\rho^\mp K^\pm K_S^0$	( 1.9 ± 0.4 ) × 10 <sup>-3</sup>		DESIG=342
Γ <sub>26</sub>	$h_1(1415)\eta' \rightarrow \gamma\eta\eta'$			DESIG=435
Γ <sub>27</sub>	$h_1(1595)\eta' \rightarrow \gamma\eta\eta'$			DESIG=437
Γ <sub>28</sub>	$\rho(1450)\pi$			DESIG=310
Γ <sub>29</sub>	$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	( 2.3 ± 0.7 ) × 10 <sup>-3</sup>		DESIG=328
Γ <sub>30</sub>	$\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp$	( 3.5 ± 0.6 ) × 10 <sup>-4</sup>		DESIG=329
Γ <sub>31</sub>	$\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0$	( 2.7 ± 0.6 ) × 10 <sup>-4</sup>		DESIG=312
Γ <sub>32</sub>	$\rho(1450)\eta'(958) \rightarrow$ $\pi^+\pi^-\eta'(958)$	( 3.3 ± 0.7 ) × 10 <sup>-6</sup>		DESIG=345
Γ <sub>33</sub>	$\rho(1700)\pi$			DESIG=325
Γ <sub>34</sub>	$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	( 1.7 ± 1.1 ) × 10 <sup>-4</sup>		DESIG=313
Γ <sub>35</sub>	$\rho(2150)\pi$			DESIG=326
Γ <sub>36</sub>	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	( 8 ± 40 ) × 10 <sup>-6</sup>		DESIG=314
Γ <sub>37</sub>	$\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$			DESIG=316
Γ <sub>38</sub>	$\omega\pi^0$	( 4.5 ± 0.5 ) × 10 <sup>-4</sup>	S=1.4	DESIG=32
Γ <sub>39</sub>	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$	( 1.7 ± 0.8 ) × 10 <sup>-5</sup>		DESIG=327
Γ <sub>40</sub>	$\omega\pi^+\pi^-$	( 8.5 ± 1.0 ) × 10 <sup>-3</sup>	S=1.3	DESIG=24
Γ <sub>41</sub>	$\omega\pi^0\pi^0$	( 3.4 ± 0.8 ) × 10 <sup>-3</sup>		DESIG=140
Γ <sub>42</sub>	$\omega 3\pi^0$	( 1.9 ± 0.6 ) × 10 <sup>-3</sup>		DESIG=421
Γ <sub>43</sub>	$\omega f_2(1270)$	( 4.3 ± 0.6 ) × 10 <sup>-3</sup>		DESIG=28
Γ <sub>44</sub>	$\omega\eta$	( 1.74 ± 0.20 ) × 10 <sup>-3</sup>	S=1.6	DESIG=30
Γ <sub>45</sub>	$\omega\pi^+\pi^-\pi^0$	( 4.0 ± 0.7 ) × 10 <sup>-3</sup>		DESIG=211
Γ <sub>46</sub>	$\omega\pi^0\eta$	( 3.4 ± 1.7 ) × 10 <sup>-4</sup>		DESIG=360
Γ <sub>47</sub>	$\omega\pi^+\pi^+\pi^-\pi^-$	( 8.5 ± 3.4 ) × 10 <sup>-3</sup>		DESIG=26
Γ <sub>48</sub>	$\omega\pi^+\pi^-2\pi^0$	( 3.3 ± 0.5 ) %		DESIG=412
Γ <sub>49</sub>	$\omega\eta'\pi^+\pi^-$	( 1.12 ± 0.13 ) × 10 <sup>-3</sup>		DESIG=385
Γ <sub>50</sub>	$\omega\eta'(958)$	( 1.89 ± 0.18 ) × 10 <sup>-4</sup>		DESIG=31
Γ <sub>51</sub>	$\omega f_0(980)$	( 1.4 ± 0.5 ) × 10 <sup>-4</sup>		DESIG=150
Γ <sub>52</sub>	$\omega f_0(1710) \rightarrow \omega K\bar{K}$	( 4.8 ± 1.1 ) × 10 <sup>-4</sup>		DESIG=130
Γ <sub>53</sub>	$\omega f_1(1420)$	( 6.8 ± 2.4 ) × 10 <sup>-4</sup>		DESIG=105
Γ <sub>54</sub>	$\omega f_2'(1525)$	< 2.2 × 10 <sup>-4</sup>	CL=90%	DESIG=29
Γ <sub>55</sub>	$\omega X(1835) \rightarrow \omega p\bar{p}$	< 3.9 × 10 <sup>-6</sup>	CL=95%	DESIG=263
Γ <sub>56</sub>	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	< 6.2 × 10 <sup>-5</sup>		DESIG=386
Γ <sub>57</sub>	$\omega K^+K^-$	( 1.52 ± 0.31 ) × 10 <sup>-3</sup>		DESIG=441
Γ <sub>58</sub>	$\omega K^\pm K_S^0\pi^\mp$	[b] ( 3.4 ± 0.5 ) × 10 <sup>-3</sup>		DESIG=101
Γ <sub>59</sub>	$\omega K\bar{K}$	( 1.9 ± 0.4 ) × 10 <sup>-3</sup>		DESIG=27
Γ <sub>60</sub>	$\omega K^*(892)\bar{K} + \text{c.c.}$	( 6.1 ± 0.9 ) × 10 <sup>-3</sup>		DESIG=102
Γ <sub>61</sub>	$\eta' K^{*\pm} K^\mp$	( 1.48 ± 0.13 ) × 10 <sup>-3</sup>		DESIG=355

Г62	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$		DESIG=357
Г63	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$		DESIG=353
Г64	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$		DESIG=354
Г65	$\eta' h_1(1415) \rightarrow \gamma \eta' \eta'$	$(4.7 \pm \frac{1.1}{2.0}) \times 10^{-7}$		DESIG=430
Г66	$\bar{K} K^*(892) + \text{c.c.}$			DESIG=331
Г67	$\bar{K} K^*(892) + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(5.0 \pm 0.5) \times 10^{-3}$		DESIG=332
Г68	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm \frac{0.8}{1.0}) \times 10^{-3}$	S=2.9	DESIG=121
Г69	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm \frac{0.13}{0.20}) \times 10^{-3}$		DESIG=231
Г70	$K^+ K^*(892)^- + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$		DESIG=232
Г71	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$		DESIG=122
Г72	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow$ $K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$		DESIG=233
Г73	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$		DESIG=214
Г74	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$		DESIG=343
Г75	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$		DESIG=299
Г76	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow$ $K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$		DESIG=300
Г77	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$		DESIG=445
Г78	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm \frac{0.6}{0.5}) \times 10^{-6}$		DESIG=376
Г79	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$		DESIG=344
Г80	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm \frac{1.0}{0.6}) \times 10^{-3}$		DESIG=257
Г81	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$		DESIG=46
Г82	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm \frac{0.22}{0.40}) \times 10^{-3}$		DESIG=256
Г83	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$		DESIG=132
Г84	$K^*(1410) \bar{K} + \text{c.c.}$			DESIG=317
Г85	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$		DESIG=330
Г86	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(8 \pm 6) \times 10^{-5}$		DESIG=318
Г87	$K_2^*(1430) \bar{K} + \text{c.c.}$			DESIG=319
Г88	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$		DESIG=321
Г89	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow$ $K_S^0 K^\pm \pi^\mp$	$(4.0 \pm 1.0) \times 10^{-4}$		DESIG=320
Г90	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%	DESIG=45
Г91	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(2.69 \pm \frac{0.25}{0.19}) \times 10^{-4}$		DESIG=381
Г92	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow$ $K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$		DESIG=446
Г93	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$		DESIG=301
Г94	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$		DESIG=48
Г95	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$		DESIG=303
Г96	$K_2^*(1430)^- K^*(892)^+ +$ $\text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- +$ $\text{c.c.}$	$(4 \pm 4) \times 10^{-4}$		DESIG=304
Г97	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3}$	CL=90%	DESIG=47
Г98	$\bar{K}_2^*(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow$ $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$		DESIG=235
Г99	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow$ $K^+ K^- \pi^0$	$(1.10 \pm \frac{0.60}{0.14}) \times 10^{-5}$		DESIG=382

$\Gamma_{100}$	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(6.2 \pm 2.9) \times 10^{-6}$		DESIG=383
$\Gamma_{101}$	$K_1(1270)^\pm K^\mp$	$< 3.0 \times 10^{-3}$	CL=90%	DESIG=131
$\Gamma_{102}$	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$		DESIG=377
$\Gamma_{103}$	$a_2(1320)^\pm \pi^\mp$	$[b] < 4.3 \times 10^{-3}$	CL=90%	DESIG=42
$\Gamma_{104}$	$\phi \pi^0$	$3 \times 10^{-6}$ or $1 \times 10^{-7}$		DESIG=33;OUR EVAL; $\rightarrow$ UNCHECKED $\leftarrow$
$\Gamma_{105}$	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7	DESIG=34
$\Gamma_{106}$	$\phi \pi^0 \pi^0$	$(5.0 \pm 1.0) \times 10^{-4}$		DESIG=76
$\Gamma_{107}$	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$		DESIG=35
$\Gamma_{108}$	$\phi \eta$	$(7.4 \pm 0.6) \times 10^{-4}$	S=1.2	DESIG=37
$\Gamma_{109}$	$\phi \eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2	DESIG=38
$\Gamma_{110}$	$\phi \eta \eta'$	$(2.32 \pm 0.17) \times 10^{-4}$		DESIG=387
$\Gamma_{111}$	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9	DESIG=41
$\Gamma_{112}$	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	$(2.60 \pm 0.34) \times 10^{-4}$		DESIG=236
$\Gamma_{113}$	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	$(1.8 \pm 0.5) \times 10^{-4}$		DESIG=237
$\Gamma_{114}$	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$		DESIG=278
$\Gamma_{115}$	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$		DESIG=279
$\Gamma_{116}$	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$		DESIG=229
$\Gamma_{117}$	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$	$(4.4 \pm 1.4) \times 10^{-6}$		DESIG=258
$\Gamma_{118}$	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$		DESIG=39
$\Gamma_{119}$	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$		DESIG=106
$\Gamma_{120}$	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$		DESIG=280
$\Gamma_{121}$	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi 3\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$		DESIG=281
$\Gamma_{122}$	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$		DESIG=128
$\Gamma_{123}$	$\phi f_2'(1525)$	$(8 \pm 4) \times 10^{-4}$	S=2.7	DESIG=40
$\Gamma_{124}$	$\phi X(1835) \rightarrow \phi p \bar{p}$	$< 2.1 \times 10^{-7}$	CL=90%	DESIG=291
$\Gamma_{125}$	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	$< 2.8 \times 10^{-4}$	CL=90%	DESIG=288
$\Gamma_{126}$	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	$< 6.13 \times 10^{-5}$	CL=90%	DESIG=289
$\Gamma_{127}$	$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3	DESIG=36
$\Gamma_{128}$	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$		DESIG=129
$\Gamma_{129}$	$\phi K^+ K^-$	$(8.3 \pm 1.1) \times 10^{-4}$		DESIG=295
$\Gamma_{130}$	$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$		DESIG=305
$\Gamma_{131}$	$\phi K^\pm K_S^0 \pi^\mp$	$[b] (7.2 \pm 0.8) \times 10^{-4}$		DESIG=103
$\Gamma_{132}$	$\phi K^*(892) \bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$		DESIG=104
$\Gamma_{133}$	$b_1(1235)^\pm \pi^\mp$	$[b] (3.0 \pm 0.5) \times 10^{-3}$		DESIG=49
$\Gamma_{134}$	$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$		DESIG=160
$\Gamma_{135}$	$f_2'(1525) K^+ K^-$	$(1.06 \pm 0.35) \times 10^{-3}$		DESIG=308
$\Gamma_{136}$	$\Delta(1232)^+ \bar{p}$	$< 1 \times 10^{-4}$	CL=90%	DESIG=112
$\Gamma_{137}$	$\Delta(1232)^{++} \bar{p} \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$		DESIG=70
$\Gamma_{138}$	$\Delta(1232)^{++} \bar{\Delta}(1232)^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$		DESIG=66
$\Gamma_{139}$	$\bar{\Sigma}(1385)^0 p K^-$	$(5.1 \pm 3.2) \times 10^{-4}$		DESIG=74
$\Gamma_{140}$	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	$< 8.2 \times 10^{-6}$	CL=90%	DESIG=111
$\Gamma_{141}$	$\Sigma(1385)^- \bar{\Sigma}^+ (\text{or c.c.})$	$[b] (3.1 \pm 0.5) \times 10^{-4}$		DESIG=68
$\Gamma_{142}$	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ (\text{or c.c.})$	$[b] (1.16 \pm 0.05) \times 10^{-3}$		DESIG=67
$\Gamma_{143}$	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(1.07 \pm 0.08) \times 10^{-3}$		DESIG=309
$\Gamma_{144}$	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	$< 4.1 \times 10^{-6}$	CL=90%	DESIG=260
$\Gamma_{145}$	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	$< 1.80 \times 10^{-3}$	CL=90%	DESIG=364
$\Gamma_{146}$	$\Xi^0 \Xi^0$	$(1.17 \pm 0.04) \times 10^{-3}$		DESIG=248
$\Gamma_{147}$	$\Xi(1530)^- \Xi^+ + \text{c.c.}$	$(3.18 \pm 0.08) \times 10^{-4}$		DESIG=107
$\Gamma_{148}$	$\Xi(1530)^0 \Xi^0$	$(3.2 \pm 1.4) \times 10^{-4}$		DESIG=108
$\Gamma_{149}$	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	$[c] < 1.1 \times 10^{-5}$	CL=90%	DESIG=205
$\Gamma_{150}$	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$[c] < 2.1 \times 10^{-5}$	CL=90%	DESIG=206
$\Gamma_{151}$	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$[c] < 1.6 \times 10^{-5}$	CL=90%	DESIG=207
$\Gamma_{152}$	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$[c] < 5.6 \times 10^{-5}$	CL=90%	DESIG=208
$\Gamma_{153}$	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$[c] < 1.1 \times 10^{-5}$	CL=90%	DESIG=209

## Decays into stable hadrons

NODE=M070;CLUMP=B

Γ <sub>154</sub>	$2(\pi^+\pi^-)\pi^0$	( 4.2 ± 0.4 ) %	S=2.1	DESIG=9
Γ <sub>155</sub>	$3(\pi^+\pi^-)\pi^0$	( 2.9 ± 0.6 ) %		DESIG=11
Γ <sub>156</sub>	$\pi^+\pi^-3\pi^0$	( 1.9 ± 0.9 ) %		DESIG=358
Γ <sub>157</sub>	$\pi^+\pi^-4\pi^0$	( 6.5 ± 1.3 ) × 10 <sup>-3</sup>		DESIG=419
Γ <sub>158</sub>	$\rho^\pm\pi^\mp\pi^0\pi^0$	( 1.41 ± 0.22 ) %		DESIG=362
Γ <sub>159</sub>	$\rho^+\rho^-\pi^0$	( 6.0 ± 1.1 ) × 10 <sup>-3</sup>		DESIG=363
Γ <sub>160</sub>	$\pi^+\pi^-\pi^0$	( 2.10 ± 0.08 ) %	S=1.6	DESIG=7
Γ <sub>161</sub>	$2(\pi^+\pi^-\pi^0)$	( 1.61 ± 0.20 ) %		DESIG=210
Γ <sub>162</sub>	$\pi^+\pi^-\pi^0K^+K^-$	( 1.52 ± 0.27 ) %	S=1.4	DESIG=18
Γ <sub>163</sub>	$\pi^+\pi^-$	( 1.47 ± 0.14 ) × 10 <sup>-4</sup>		DESIG=6
Γ <sub>164</sub>	$2(\pi^+\pi^-)$	( 3.20 ± 0.25 ) × 10 <sup>-3</sup>	S=1.2	DESIG=8
Γ <sub>165</sub>	$3(\pi^+\pi^-)$	( 4.3 ± 0.4 ) × 10 <sup>-3</sup>		DESIG=10
Γ <sub>166</sub>	$2(\pi^+\pi^-)3\pi^0$	( 6.2 ± 0.9 ) %		DESIG=411
Γ <sub>167</sub>	$4(\pi^+\pi^-)\pi^0$	( 9.0 ± 3.0 ) × 10 <sup>-3</sup>		DESIG=12
Γ <sub>168</sub>	$2(\pi^+\pi^-)\eta$	( 2.29 ± 0.28 ) × 10 <sup>-3</sup>		DESIG=201
Γ <sub>169</sub>	$3(\pi^+\pi^-)\eta$	( 7.2 ± 1.5 ) × 10 <sup>-4</sup>		DESIG=202
Γ <sub>170</sub>	$2(\pi^+\pi^-\pi^0)\eta$	( 1.6 ± 0.5 ) × 10 <sup>-3</sup>		DESIG=418
Γ <sub>171</sub>	$\pi^+\pi^-\pi^0\pi^0\eta$	( 2.4 ± 0.5 ) × 10 <sup>-3</sup>		DESIG=359
Γ <sub>172</sub>	$\rho^\pm\pi^\mp\pi^0\eta$	( 1.9 ± 0.8 ) × 10 <sup>-3</sup>		DESIG=361
Γ <sub>173</sub>	$K^+K^-$	( 2.86 ± 0.21 ) × 10 <sup>-4</sup>		DESIG=13
Γ <sub>174</sub>	$K_S^0K_L^0$	( 1.95 ± 0.11 ) × 10 <sup>-4</sup>	S=2.4	DESIG=75
Γ <sub>175</sub>	$K_S^0K_S^0$	< 1.4 × 10 <sup>-8</sup>	CL=95%	DESIG=14
Γ <sub>176</sub>	$K\bar{K}\pi$	( 6.1 ± 1.0 ) × 10 <sup>-3</sup>		DESIG=15
Γ <sub>177</sub>	$K^+K^-\pi^0$	( 2.88 ± 0.12 ) × 10 <sup>-3</sup>		DESIG=334
Γ <sub>178</sub>	$K_S^0K^\pm\pi^\mp$	( 5.6 ± 0.5 ) × 10 <sup>-3</sup>		DESIG=335
Γ <sub>179</sub>	$K_S^0K_L^0\pi^0$	( 2.06 ± 0.26 ) × 10 <sup>-3</sup>		DESIG=336
Γ <sub>180</sub>	$K^*(892)^0\bar{K}^0 + c.c. \rightarrow K_S^0K_L^0\pi^0$	( 1.21 ± 0.18 ) × 10 <sup>-3</sup>		DESIG=339
Γ <sub>181</sub>	$K_2^*(1430)^0\bar{K}^0 + c.c. \rightarrow K_S^0K_L^0\pi^0$	( 4.3 ± 1.3 ) × 10 <sup>-4</sup>		DESIG=338
Γ <sub>182</sub>	$K^+K^-\pi^+\pi^-$	( 7.0 ± 1.0 ) × 10 <sup>-3</sup>		DESIG=16
Γ <sub>183</sub>	$K^+K^-\pi^0\pi^0$	( 2.13 ± 0.22 ) × 10 <sup>-3</sup>		DESIG=234
Γ <sub>184</sub>	$K_S^0K_L^0\pi^+\pi^-$	( 3.8 ± 0.6 ) × 10 <sup>-3</sup>		DESIG=296
Γ <sub>185</sub>	$K_S^0K_L^0\pi^0\pi^0$	( 1.9 ± 0.4 ) × 10 <sup>-3</sup>		DESIG=337
Γ <sub>186</sub>	$K_S^0K_L^0\eta$	( 1.45 ± 0.33 ) × 10 <sup>-3</sup>		DESIG=340
Γ <sub>187</sub>	$K_S^0K_S^0\pi^+\pi^-$	( 1.68 ± 0.19 ) × 10 <sup>-3</sup>		DESIG=297
Γ <sub>188</sub>	$K^\mp K_S^0\pi^\pm\pi^0$	( 5.7 ± 0.5 ) × 10 <sup>-3</sup>		DESIG=341
Γ <sub>189</sub>	$K^+K^-2(\pi^+\pi^-)$	( 3.1 ± 1.3 ) × 10 <sup>-3</sup>		DESIG=17
Γ <sub>190</sub>	$K^+K^-\pi^+\pi^-\eta$	( 4.7 ± 0.7 ) × 10 <sup>-3</sup>		DESIG=238
Γ <sub>191</sub>	$2(K^+K^-)$	( 7.2 ± 0.8 ) × 10 <sup>-4</sup>		DESIG=19
Γ <sub>192</sub>	$K^+K^-K_S^0K_S^0$	( 4.2 ± 0.7 ) × 10 <sup>-4</sup>		DESIG=298
Γ <sub>193</sub>	$p\bar{p}$	( 2.120 ± 0.029 ) × 10 <sup>-3</sup>		DESIG=50
Γ <sub>194</sub>	$p\bar{p}\pi^0$	( 1.19 ± 0.08 ) × 10 <sup>-3</sup>	S=1.1	DESIG=52
Γ <sub>195</sub>	$p\bar{p}\pi^+\pi^-$	( 6.0 ± 0.5 ) × 10 <sup>-3</sup>	S=1.3	DESIG=54
Γ <sub>196</sub>	$p\bar{p}\pi^+\pi^-\pi^0$	[d] ( 2.3 ± 0.9 ) × 10 <sup>-3</sup>	S=1.9	DESIG=55
Γ <sub>197</sub>	$p\bar{p}\eta$	( 2.00 ± 0.12 ) × 10 <sup>-3</sup>		DESIG=56
Γ <sub>198</sub>	$p\bar{p}\rho$	< 3.1 × 10 <sup>-4</sup>	CL=90%	DESIG=57
Γ <sub>199</sub>	$p\bar{p}\omega$	( 9.8 ± 1.0 ) × 10 <sup>-4</sup>	S=1.3	DESIG=58
Γ <sub>200</sub>	$p\bar{p}\eta'(958)$	( 1.29 ± 0.14 ) × 10 <sup>-4</sup>	S=2.0	DESIG=59
Γ <sub>201</sub>	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	( 6.8 ± 1.8 ) × 10 <sup>-5</sup>		DESIG=276
Γ <sub>202</sub>	$p\bar{p}\phi$	( 5.19 ± 0.33 ) × 10 <sup>-5</sup>		DESIG=127
Γ <sub>203</sub>	$p\bar{n}\pi^-$	( 2.12 ± 0.09 ) × 10 <sup>-3</sup>		DESIG=53
Γ <sub>204</sub>	$n\bar{n}$	( 2.09 ± 0.16 ) × 10 <sup>-3</sup>		DESIG=64
Γ <sub>205</sub>	$n\bar{n}\pi^+\pi^-$	( 4 ± 4 ) × 10 <sup>-3</sup>		DESIG=65
Γ <sub>206</sub>	$nN(1440)$	seen		DESIG=215;OUR EST;→ UNCHECKED ←



Γ <sub>207</sub>	$nN(1520)$	seen		DESIG=216;OUR EST;→ UNCHECKED ←
Γ <sub>208</sub>	$nN(1535)$	seen		DESIG=217;OUR EST;→ UNCHECKED ←
Γ <sub>209</sub>	$\Lambda\bar{\Lambda}$		$(1.89 \pm 0.09) \times 10^{-3}$ S=2.8	DESIG=60
Γ <sub>210</sub>	$\Lambda\bar{\Lambda}\pi^0$		$(3.8 \pm 0.4) \times 10^{-5}$	DESIG=109
Γ <sub>211</sub>	$\Lambda\bar{\Lambda}\pi^+\pi^-$		$(4.3 \pm 1.0) \times 10^{-3}$	DESIG=261
Γ <sub>212</sub>	$\Lambda\bar{\Lambda}\eta$		$(1.62 \pm 0.17) \times 10^{-4}$	DESIG=228
Γ <sub>213</sub>	$\Lambda\bar{\Sigma}^-\pi^+$ (or c.c.)	[b]	$(8.3 \pm 0.7) \times 10^{-4}$ S=1.2	DESIG=71
Γ <sub>214</sub>	$pK^-\bar{\Lambda} + \text{c.c.}$		$(8.6 \pm 1.1) \times 10^{-4}$	DESIG=72
Γ <sub>215</sub>	$pK^-\bar{\Sigma}^0$		$(2.9 \pm 0.8) \times 10^{-4}$	DESIG=73
Γ <sub>216</sub>	$\bar{\Lambda}nK_S^0 + \text{c.c.}$		$(6.5 \pm 1.1) \times 10^{-4}$	DESIG=225
Γ <sub>217</sub>	$\Lambda\bar{\Sigma} + \text{c.c.}$		$(2.83 \pm 0.23) \times 10^{-5}$	DESIG=61
Γ <sub>218</sub>	$\Sigma^+\bar{\Sigma}^-$		$(1.07 \pm 0.04) \times 10^{-3}$	DESIG=247
Γ <sub>219</sub>	$\Sigma^0\bar{\Sigma}^0$		$(1.172 \pm 0.032) \times 10^{-3}$ S=1.4	DESIG=63
Γ <sub>220</sub>	$\Sigma^+\bar{\Sigma}^-\eta$		$(6.3 \pm 0.4) \times 10^{-5}$	DESIG=448
Γ <sub>221</sub>	$\Xi^-\bar{\Xi}^+$		$(9.7 \pm 0.8) \times 10^{-4}$ S=1.4	DESIG=62

## Radiative decays

				NODE=M070;CLUMP=C
Γ <sub>222</sub>	$\gamma\eta_c(1S)$	$(1.7 \pm 0.4) \%$	S=1.5	DESIG=85
Γ <sub>223</sub>	$\gamma\eta_c(1S) \rightarrow 3\gamma$	$(3.8 \pm 1.3) \times 10^{-6}$	S=1.1	DESIG=246
Γ <sub>224</sub>	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta'$	$(4.9 \pm 0.8) \times 10^{-5}$		DESIG=391
Γ <sub>225</sub>	$3\gamma$	$(1.16 \pm 0.22) \times 10^{-5}$		DESIG=81
Γ <sub>226</sub>	$4\gamma$	$< 9 \times 10^{-6}$	CL=90%	DESIG=244
Γ <sub>227</sub>	$5\gamma$	$< 1.5 \times 10^{-5}$	CL=90%	DESIG=245
Γ <sub>228</sub>	$\gamma\pi^0$	$(3.56 \pm 0.17) \times 10^{-5}$		DESIG=82
Γ <sub>229</sub>	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$		DESIG=283
Γ <sub>230</sub>	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9	DESIG=95
Γ <sub>231</sub>	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$		DESIG=203
Γ <sub>232</sub>	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$		DESIG=204
Γ <sub>233</sub>	$\gamma\pi^+\pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$		DESIG=99
Γ <sub>234</sub>	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$		DESIG=378
Γ <sub>235</sub>	$\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}]$	$(7 \pm 4) \times 10^{-4}$	S=2.1	DESIG=176
Γ <sub>236</sub>	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$		DESIG=143
Γ <sub>237</sub>	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$		DESIG=145
Γ <sub>238</sub>	$\gamma\eta$	$(1.085 \pm 0.018) \times 10^{-3}$		DESIG=83
Γ <sub>239</sub>	$\gamma\eta\pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$		DESIG=292
Γ <sub>240</sub>	$\gamma f_0(500) \rightarrow \gamma\pi\pi$			DESIG=398
Γ <sub>241</sub>	$\gamma f_0(500) \rightarrow \gamma K\bar{K}$			DESIG=399
Γ <sub>242</sub>	$\gamma f_0(500) \rightarrow \gamma\eta\eta$			DESIG=400
Γ <sub>243</sub>	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$	$< 2.5 \times 10^{-6}$	CL=95%	DESIG=293
Γ <sub>244</sub>	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$	$< 6.6 \times 10^{-6}$	CL=95%	DESIG=294
Γ <sub>245</sub>	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$		DESIG=96
Γ <sub>246</sub>	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$		DESIG=142
Γ <sub>247</sub>	$\gamma\eta'(958)$	$(5.25 \pm 0.07) \times 10^{-3}$	S=1.3	DESIG=84
Γ <sub>248</sub>	$\gamma f_0(980) \rightarrow \gamma\pi\pi$			DESIG=393
Γ <sub>249</sub>	$\gamma f_0(980) \rightarrow \gamma K\bar{K}$			DESIG=394
Γ <sub>250</sub>	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$		DESIG=94
Γ <sub>251</sub>	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	CL=90%	DESIG=226
Γ <sub>252</sub>	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90%	DESIG=227
Γ <sub>253</sub>	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$		DESIG=97
Γ <sub>254</sub>	$\gamma\phi\phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1	DESIG=98
Γ <sub>255</sub>	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6	DESIG=89
Γ <sub>256</sub>	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8	DESIG=171
Γ <sub>257</sub>	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=170
Γ <sub>258</sub>	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3	DESIG=124
Γ <sub>259</sub>	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95%	DESIG=212
Γ <sub>260</sub>	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	$< 2.63 \times 10^{-6}$	CL=90%	DESIG=348
Γ <sub>261</sub>	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	$< 1.86 \times 10^{-6}$	CL=90%	DESIG=349
Γ <sub>262</sub>	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$		DESIG=125
Γ <sub>263</sub>	$\gamma\eta(1760) \rightarrow \gamma\omega\omega$	$(1.98 \pm 0.33) \times 10^{-3}$		DESIG=224

Γ <sub>264</sub>	$\gamma\eta(1760) \rightarrow \gamma\gamma\gamma$	$< 4.80 \times 10^{-6}$	CL=90%	DESIG=347
Γ <sub>265</sub>	$\gamma\eta(2225)$	$(3.14 \pm_{-0.19}^{+0.50}) \times 10^{-4}$		DESIG=126
Γ <sub>266</sub>	$\gamma f_2(1270)$	$(1.63 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=86
Γ <sub>267</sub>	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm_{-0.22}^{+0.60}) \times 10^{-5}$		DESIG=373
Γ <sub>268</sub>	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$		DESIG=88
Γ <sub>269</sub>	$\gamma f_0(1370) \rightarrow \gamma\pi\pi$			DESIG=395
Γ <sub>270</sub>	$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$		DESIG=284
Γ <sub>271</sub>	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$		DESIG=368
Γ <sub>272</sub>	$\gamma f_0(1370) \rightarrow \gamma\eta\eta$			DESIG=396
Γ <sub>273</sub>	$\gamma f_0(1370) \rightarrow \gamma\eta\eta'$			DESIG=397
Γ <sub>274</sub>	$\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi$	$(7.9 \pm 1.3) \times 10^{-4}$		DESIG=175
Γ <sub>275</sub>	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	$(1.09 \pm 0.24) \times 10^{-4}$		DESIG=172
Γ <sub>276</sub>	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	$(1.7 \pm_{-1.4}^{+0.6}) \times 10^{-5}$		DESIG=265
Γ <sub>277</sub>	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm_{-0.60}^{+0.24}) \times 10^{-5}$		DESIG=369
Γ <sub>278</sub>	$\gamma f_0(1500) \rightarrow \gamma\eta\eta'$			DESIG=401
Γ <sub>279</sub>	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$		DESIG=141
Γ <sub>280</sub>	$\gamma f_2'(1525)$	$(5.7 \pm_{-0.5}^{+0.8}) \times 10^{-4}$	S=1.5	DESIG=87
Γ <sub>281</sub>	$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm_{-0.5}^{+0.7}) \times 10^{-5}$		DESIG=374
Γ <sub>282</sub>	$\gamma f_2'(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$		DESIG=268
Γ <sub>283</sub>	$\gamma f_2(1565) \rightarrow \gamma\eta\eta'$			DESIG=432
Γ <sub>284</sub>	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	$(2.8 \pm 1.8) \times 10^{-4}$		DESIG=222
Γ <sub>285</sub>	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=135
Γ <sub>286</sub>	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(9.5 \pm_{-0.5}^{+1.0}) \times 10^{-4}$	S=1.5	DESIG=91
Γ <sub>287</sub>	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$		DESIG=221
Γ <sub>288</sub>	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm_{-0.7}^{+1.2}) \times 10^{-4}$		DESIG=266
Γ <sub>289</sub>	$\gamma f_0(1710) \rightarrow \gamma\eta\eta'$			DESIG=402
Γ <sub>290</sub>	$\gamma f_0(1710) \rightarrow \gamma\omega\phi$	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=262
Γ <sub>291</sub>	$\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm_{-0.33}^{+0.20}) \times 10^{-5}$		DESIG=370
Γ <sub>292</sub>	$\gamma f_2(1810) \rightarrow \gamma\eta\eta$	$(5.4 \pm_{-2.4}^{+3.5}) \times 10^{-5}$		DESIG=269
Γ <sub>293</sub>	$\gamma\eta_1(1855) \rightarrow \gamma\eta\eta'$	$(2.7 \pm_{-0.5}^{+0.4}) \times 10^{-6}$		DESIG=447
Γ <sub>294</sub>	$\gamma f_0(1770) \rightarrow \gamma\eta\eta'$			DESIG=431
Γ <sub>295</sub>	$\gamma f_2(1910) \rightarrow \gamma\omega\omega$	$(2.0 \pm 1.4) \times 10^{-4}$		DESIG=223
Γ <sub>296</sub>	$\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$		DESIG=144
Γ <sub>297</sub>	$\gamma f_2(2010) \rightarrow \gamma\eta\eta'$			DESIG=440
Γ <sub>298</sub>	$\gamma f_0(2020) \rightarrow \gamma\pi\pi$			DESIG=403
Γ <sub>299</sub>	$\gamma f_0(2020) \rightarrow \gamma K \bar{K}$			DESIG=404
Γ <sub>300</sub>	$\gamma f_0(2020) \rightarrow \gamma\eta\eta$			DESIG=405
Γ <sub>301</sub>	$\gamma f_0(2020) \rightarrow \gamma\eta'\eta'$	$(2.63 \pm_{-0.50}^{+0.32}) \times 10^{-4}$		DESIG=426
Γ <sub>302</sub>	$\gamma f_0(2020) \rightarrow \gamma\eta\eta'$			DESIG=438
Γ <sub>303</sub>	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$		DESIG=100
Γ <sub>304</sub>	$\gamma f_4(2050) \rightarrow \gamma\eta\eta'$			DESIG=433
Γ <sub>305</sub>	$\gamma f_0(2100) \rightarrow \gamma\eta\eta$	$(1.13 \pm_{-0.30}^{+0.60}) \times 10^{-4}$		DESIG=267
Γ <sub>306</sub>	$\gamma f_0(2100) \rightarrow \gamma K \bar{K}$			DESIG=406
Γ <sub>307</sub>	$\gamma f_0(2100) \rightarrow \gamma\pi\pi$	$(6.2 \pm 1.0) \times 10^{-4}$		DESIG=286
Γ <sub>308</sub>	$\gamma f_0(2200)$			DESIG=123
Γ <sub>309</sub>	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$		DESIG=285
Γ <sub>310</sub>	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm_{-0.50}^{+0.19}) \times 10^{-4}$		DESIG=371
Γ <sub>311</sub>	$\gamma f_0(2200) \rightarrow \gamma\pi\pi$			DESIG=407

Γ <sub>312</sub>	$\gamma f_0(2200) \rightarrow \gamma \eta \eta$				DESIG=408
Γ <sub>313</sub>	$\gamma f_J(2220)$				DESIG=92
Γ <sub>314</sub>	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	< 3.9	$\times 10^{-5}$	CL=90%	DESIG=136
Γ <sub>315</sub>	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	< 4.1	$\times 10^{-5}$	CL=90%	DESIG=137
Γ <sub>316</sub>	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	( 1.5 ± 0.8 )	$\times 10^{-5}$		DESIG=138
Γ <sub>317</sub>	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	( 4.9 ± 0.7 )	$\times 10^{-5}$		DESIG=372
Γ <sub>318</sub>	$\gamma f_0(2330) \rightarrow \gamma \pi \pi$				DESIG=409
Γ <sub>319</sub>	$\gamma f_0(2330) \rightarrow \gamma \eta \eta$				DESIG=410
Γ <sub>320</sub>	$\gamma f_0(2330) \rightarrow \gamma \eta' \eta'$	( 6.1 + 4.0 / - 1.8 )	$\times 10^{-6}$		DESIG=427
Γ <sub>321</sub>	$\gamma f_0(2330) \rightarrow \gamma \eta \eta'$				DESIG=439
Γ <sub>322</sub>	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	( 5.6 + 2.4 / - 2.2 )	$\times 10^{-5}$		DESIG=270
Γ <sub>323</sub>	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	( 5.5 + 4.0 / - 1.5 )	$\times 10^{-5}$		DESIG=375
Γ <sub>324</sub>	$\gamma f_2(2340) \rightarrow \gamma \eta' \eta'$	( 8.7 + 0.9 / - 1.8 )	$\times 10^{-6}$		DESIG=428
Γ <sub>325</sub>	$\gamma f_0(2470) \rightarrow \gamma \eta' \eta'$	( 8.2 + 4.0 / - 2.8 )	$\times 10^{-7}$		DESIG=429
Γ <sub>326</sub>	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	( 2.7 + 0.6 / - 0.8 )	$\times 10^{-4}$	S=1.6	DESIG=213
Γ <sub>327</sub>	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	( 7.7 + 1.5 / - 0.9 )	$\times 10^{-5}$		DESIG=254
Γ <sub>328</sub>	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	( 3.3 + 2.0 / - 1.3 )	$\times 10^{-5}$		DESIG=282
Γ <sub>329</sub>	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$				DESIG=346
Γ <sub>330</sub>	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	< 3.56	$\times 10^{-6}$	CL=90%	DESIG=350
Γ <sub>331</sub>	$\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-)$	( 2.4 + 0.7 / - 0.8 )	$\times 10^{-5}$		DESIG=264
Γ <sub>332</sub>	$\gamma X(2370) \rightarrow \gamma K^+ K^- \eta'$	( 1.8 ± 0.7 )	$\times 10^{-5}$		DESIG=388
Γ <sub>333</sub>	$\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	( 1.2 ± 0.5 )	$\times 10^{-5}$		DESIG=389
Γ <sub>334</sub>	$\gamma X(2370) \rightarrow \gamma \eta \eta \eta'$	< 9.2	$\times 10^{-6}$	CL=90%	DESIG=390
Γ <sub>335</sub>	$\gamma p \bar{p}$	( 3.8 ± 1.0 )	$\times 10^{-4}$		DESIG=90
Γ <sub>336</sub>	$\gamma p \bar{p} \pi^+ \pi^-$	< 7.9	$\times 10^{-4}$	CL=90%	DESIG=93
Γ <sub>337</sub>	$\gamma \Lambda \bar{\Lambda}$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=200
Γ <sub>338</sub>	$\gamma A^0 \rightarrow \gamma$ invisible	[e] < 1.7	$\times 10^{-6}$	CL=90%	DESIG=251
Γ <sub>339</sub>	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[f] < 7.8	$\times 10^{-7}$	CL=90%	DESIG=259
<b>Dalitz decays</b>					
Γ <sub>340</sub>	$\pi^0 e^+ e^-$	( 7.6 ± 1.4 )	$\times 10^{-7}$		NODE=M070;CLUMP=G DESIG=271
Γ <sub>341</sub>	$\eta e^+ e^-$	( 1.42 ± 0.08 )	$\times 10^{-5}$		DESIG=272
Γ <sub>342</sub>	$\eta'(958) e^+ e^-$	( 6.59 ± 0.18 )	$\times 10^{-5}$		DESIG=273
Γ <sub>343</sub>	$X(1835) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta'$	( 3.58 ± 0.25 )	$\times 10^{-6}$		DESIG=423
Γ <sub>344</sub>	$X(2120) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta'$	( 8.2 ± 1.3 )	$\times 10^{-7}$		DESIG=425
Γ <sub>345</sub>	$X(2370) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta'$	( 1.08 ± 0.17 )	$\times 10^{-6}$		DESIG=424
Γ <sub>346</sub>	$\eta U \rightarrow \eta e^+ e^-$	[g] < 9.11	$\times 10^{-7}$	CL=90%	DESIG=352
Γ <sub>347</sub>	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	[g] < 2.0	$\times 10^{-7}$	CL=90%	DESIG=366
Γ <sub>348</sub>	$\phi e^+ e^-$	< 1.2	$\times 10^{-7}$	CL=90%	DESIG=384
<b>Weak decays</b>					
Γ <sub>349</sub>	$D^- e^+ \nu_e + c.c.$	< 7.1	$\times 10^{-8}$	CL=90%	NODE=M070;CLUMP=E DESIG=218
Γ <sub>350</sub>	$\bar{D}^0 e^+ e^- + c.c.$	< 8.5	$\times 10^{-8}$	CL=90%	DESIG=219
Γ <sub>351</sub>	$D_s^- e^+ \nu_e + c.c.$	< 1.3	$\times 10^{-6}$	CL=90%	DESIG=220
Γ <sub>352</sub>	$D_s^{*-} e^+ \nu_e + c.c.$	< 1.8	$\times 10^{-6}$	CL=90%	DESIG=290
Γ <sub>353</sub>	$D^- \pi^+ + c.c.$	< 7.5	$\times 10^{-5}$	CL=90%	DESIG=241
Γ <sub>354</sub>	$\bar{D}^0 \bar{K}^0 + c.c.$	< 1.7	$\times 10^{-4}$	CL=90%	DESIG=242
Γ <sub>355</sub>	$\bar{D}^0 \bar{K}^{*0} + c.c.$	< 2.5	$\times 10^{-6}$	CL=90%	DESIG=275
Γ <sub>356</sub>	$D_s^- \pi^+ + c.c.$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=243
Γ <sub>357</sub>	$D_s^- \rho^+ + c.c.$	< 1.3	$\times 10^{-5}$	CL=90%	DESIG=274

**Charge conjugation (C), Parity (P),  
Lepton Family number (LF) violating modes**

$\Gamma_{358}$	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%
$\Gamma_{359}$	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%
$\Gamma_{360}$	$e^{\pm}\mu^{\mp}$	LF	< 1.6	$\times 10^{-7}$	CL=90%
$\Gamma_{361}$	$e^{\pm}\tau^{\mp}$	LF	< 7.5	$\times 10^{-8}$	CL=90%
$\Gamma_{362}$	$\mu^{\pm}\tau^{\mp}$	LF	< 2.0	$\times 10^{-6}$	CL=90%
$\Gamma_{363}$	$\Lambda_c^+ e^- + c.c.$		< 6.9	$\times 10^{-8}$	CL=90%

NODE=M070;CLUMP=D

DESIG=80

DESIG=277

DESIG=177

DESIG=178

DESIG=179

DESIG=379

**Other decays**

$\Gamma_{364}$	invisible		< 7	$\times 10^{-4}$	CL=90%
----------------	-----------	--	-----	------------------	--------

NODE=M070;CLUMP=F

DESIG=240

[a] For  $E_\gamma > 100$  MeV.

LINKAGE=EGM

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

LINKAGE=SG

[c]  $\Theta(1540)$  is a hypothetical pentaquark state of  $1.54 \text{ GeV}/c^2$  mass and a width of less than  $25 \text{ MeV}/c^2$ .

LINKAGE=THT

[d] Includes  $p\bar{p}\pi^+\pi^-\gamma$  and excludes  $p\bar{p}\eta, p\bar{p}\omega, p\bar{p}\eta'$ .

LINKAGE=MF

[e] For a narrow state  $A$  with mass less than 960 MeV.

LINKAGE=NSA

[f] For a narrow scalar or pseudoscalar  $A^0$  with mass 0.21–3.0 GeV.

LINKAGE=NA0

[g] For a dark photon  $U$  with mass between 100 and 2100 MeV.

LINKAGE=DPH

**$J/\psi(1S)$  PARTIAL WIDTHS**

NODE=M070220

**$\Gamma(\text{hadrons})$**

 $\Gamma_1$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b><math>81.37 \pm 1.36 \pm 1.30</math></b>	<sup>1</sup> ANASHIN	20	KEDR $e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
74.1 $\pm$ 8.1	BAI	95B	BES $e^+e^-$
59 $\pm$ 24	BALDINI-...	75	FRAG $e^+e^-$
59 $\pm$ 14	BOYARSKI	75	MRK1 $e^+e^-$
50 $\pm$ 25	ESPOSITO	75B	FRAM $e^+e^-$

NODE=M070W3

NODE=M070W3

<sup>1</sup> Based on the same dataset as ANASHIN 18A and correlated to the values reported there

NODE=M070W3;LINKAGE=A

**$\Gamma(e^+e^-)$**

 $\Gamma_5$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.53 \pm 0.10</math></b>	<b>OUR AVERAGE</b>			
$5.550 \pm 0.056 \pm 0.089$		<sup>1,2</sup> ANASHIN	18A	KEDR $e^+e^-$
$5.36^{+0.29}_{-0.28}$		<sup>3</sup> HSUEH	92	RVUE See $\Upsilon$ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$5.58 \pm 0.05 \pm 0.08$		<sup>4</sup> ABLIKIM	16Q	BES3 $3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.71 \pm 0.16$	13k	<sup>5</sup> ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.57 \pm 0.19$	7.8k	<sup>5</sup> AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$
$5.14 \pm 0.39$		BAI	95B	BES $e^+e^-$
$4.72 \pm 0.35$		ALEXANDER	89	RVUE See $\Upsilon$ mini-review
$4.4 \pm 0.6$		<sup>3</sup> BRANDELIK	79C	DASP $e^+e^-$
$4.6 \pm 0.8$		<sup>6</sup> BALDINI-...	75	FRAG $e^+e^-$
$4.8 \pm 0.6$		BOYARSKI	75	MRK1 $e^+e^-$
$4.6 \pm 1.0$		ESPOSITO	75B	FRAM $e^+e^-$

NODE=M070W1

NODE=M070W1

OCCUR=4

<sup>1</sup> From the cross sections of  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow \text{hadrons}$  near the  $J/\psi(1S)$  peak.

NODE=M070W1;LINKAGE=D

<sup>2</sup> Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

NODE=M070W1;LINKAGE=E

<sup>3</sup> From a simultaneous fit to  $e^+e^-$ ,  $\mu^+\mu^-$ , and hadronic channels assuming  $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ .

NODE=M070W1;LINKAGE=F

<sup>4</sup> Using  $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$  from ABLIKIM 13R.

NODE=M070W1;LINKAGE=A

<sup>5</sup> Calculated by us from the reported values of  $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$  using  $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$ .

NODE=M070W1;LINKAGE=AA

<sup>6</sup> Assuming equal partial widths for  $e^+e^-$  and  $\mu^+\mu^-$ .

NODE=M070W1;LINKAGE=B

$\Gamma(\mu^+\mu^-)$  $\Gamma_7$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.13±0.52	BAI	95B	BES $e^+e^-$
4.8 ±0.6	BOYARSKI	75	MRK1 $e^+e^-$
5 ±1	ESPOSITO	75B	FRAM $e^+e^-$

NODE=M070W2  
 NODE=M070W2

 $\Gamma(\gamma\gamma)$  $\Gamma_{358}$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C	DASP $e^+e^-$

NODE=M070W70  
 NODE=M070W70

 $J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M070225

This combination of a partial width with the partial width into  $e^+e^-$  and with the total width is obtained from the integrated cross section into channel(l) in the  $e^+e^-$  annihilation.

NODE=M070225

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_5/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.884±0.048±0.078	<sup>1,2</sup> ANASHIN	18A	KEDR $e^+e^-$
4 ±0.8	<sup>3</sup> BALDINI-...	75	FRAG $e^+e^-$
3.9 ±0.8	<sup>3</sup> ESPOSITO	75B	FRAM $e^+e^-$

NODE=M070G3  
 NODE=M070G3

<sup>1</sup> From the cross sections of  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow \text{hadrons}$  near the  $J/\psi(1S)$  peak.

NODE=M070G3;LINKAGE=A

<sup>2</sup> Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

NODE=M070G3;LINKAGE=B

<sup>3</sup> Data redundant with branching ratios or partial widths above.

NODE=M070G3;LINKAGE=S

 $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_5\Gamma_5/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
333.1± 6.6±4.0	<sup>1,2</sup> ANASHIN	18A	KEDR $e^+e^-$
332.3± 6.4±4.8	ANASHIN	10	KEDR 3.097 $e^+e^- \rightarrow e^+e^-$
350 ± 20	BRANDELIK	79C	DASP $e^+e^-$
320 ± 70	<sup>3</sup> BALDINI-...	75	FRAG $e^+e^-$
340 ± 90	<sup>3</sup> ESPOSITO	75B	FRAM $e^+e^-$
360 ±100	<sup>3</sup> FORD	75	SPEC $e^+e^-$

NODE=M070G1  
 NODE=M070G1

OCCUR=2

<sup>1</sup> From the cross sections of  $e^+e^- \rightarrow e^+e^-$  and  $e^+e^- \rightarrow \text{hadrons}$  near the  $J/\psi(1S)$  peak.

NODE=M070G1;LINKAGE=A

<sup>2</sup> Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

NODE=M070G1;LINKAGE=B

<sup>3</sup> Data redundant with branching ratios or partial widths above.

NODE=M070G1;LINKAGE=S

 $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_7\Gamma_5/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>333 ± 4 OUR AVERAGE</b>				
333.4± 2.5±4.4		ABLIKIM	16Q	BES3 3.773 $e^+e^- \rightarrow \mu^+\mu^-\gamma$
331.8± 5.2±6.3		ANASHIN	10	KEDR 3.097 $e^+e^- \rightarrow \mu^+\mu^-$
338.4± 5.8±7.1	13k	ADAMS	06A	CLEO $e^+e^- \rightarrow \mu^+\mu^-\gamma$
330.1± 7.7±7.3	7.8k	AUBERT	04	BABR $e^+e^- \rightarrow \mu^+\mu^-\gamma$

NODE=M070G2  
 NODE=M070G2

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ±90		DASP	75	DASP $e^+e^-$
380 ±50	<sup>1</sup>	ESPOSITO	75B	FRAM $e^+e^-$

<sup>1</sup> Data redundant with branching ratios or partial widths above.

NODE=M070G2;LINKAGE=S

 $\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{13}\Gamma_5/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ±0.4 OUR AVERAGE</b>				
2.34±0.43±0.16	49	LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$
2.22±0.96±0.02	9	<sup>1</sup> AUBERT	07AU	BABR 10.6 $e^+e^- \rightarrow \eta\pi^+\pi^-\gamma$

NODE=M070G25  
 NODE=M070G25

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow \pi^+\pi^-\pi^0)] = 0.51 \pm 0.22 \pm 0.03$  eV which we divide by our best value  $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.02 \pm 0.25) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G25;LINKAGE=AU

$$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>64.8±11.1±0.4</b>	200	<sup>1</sup> LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

NODE=M070Q20  
NODE=M070Q20

<sup>1</sup> LEES 21C reports  $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] = 21.1 \pm 1.7 \pm 3.2$  eV which we divide by our best value  $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q20;LINKAGE=A

$$\Gamma(\eta\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>26.9±5.7±0.1</b>	101	<sup>1</sup> LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-3\pi^0\gamma\gamma)$

NODE=M070Q19  
NODE=M070Q19

<sup>1</sup> LEES 21C reports  $[\Gamma(J/\psi(1S) \rightarrow \eta\pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.6 \pm 1.6 \pm 1.6$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q19;LINKAGE=A

$$\Gamma(\eta K^\pm K_S^0 \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{19}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3±1.4±0.4</b>	44	LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

NODE=M070G38  
NODE=M070G38

$$\Gamma(\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{22}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>155±26±36</b>	14k	LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

NODE=M070Q13  
NODE=M070Q13

$$\Gamma(\rho^+ \rho^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{23}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>32±13±15</b>	14k	LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

NODE=M070Q14  
NODE=M070Q14

$$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.4±1.0±1.9</b>	130	LEES	17D BABR	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

NODE=M070G31  
NODE=M070G31

$$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{40}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>53.6±5.0±0.4</b>	788	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$

NODE=M070G24  
NODE=M070G24

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 47.8 \pm 3.1 \pm 3.2$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G24;LINKAGE=AU

$$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{41}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>27.8±3.5±0.2</b>	398	<sup>1</sup> LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

NODE=M070P54  
NODE=M070P54

<sup>1</sup> LEES 18E reports  $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 24.8 \pm 1.8 \pm 2.5$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P54;LINKAGE=A

$$\Gamma(\omega 3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.5±3.1±0.1</b>	89	<sup>1</sup> LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

NODE=M070Q18  
NODE=M070Q18

<sup>1</sup> LEES 21C reports  $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 9.4 \pm 2.3 \pm 1.5$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q18;LINKAGE=A

$$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{45}\Gamma_5/\Gamma$$

VALUE ( $10^{-2}$ keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2±0.3±0.2</b>	170	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\pi^0\gamma$

NODE=M070G8  
NODE=M070G8

$$\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{44}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.9±7.6±0.2</b>	1	LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

NODE=M070Q21  
NODE=M070Q21

<sup>1</sup> Different final state as in AUBERT 06. LEES 21C reports  $[\Gamma(J/\psi(1S) \rightarrow \omega\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 4.9 \pm 2.1 \pm 0.7$  eV which we divide by our best values  $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$ ,  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070Q21;LINKAGE=A

$$\Gamma(\omega\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{46}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.90±0.96±0.01</b>	27	1 LEES	18E	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\eta\gamma$

NODE=M070P58  
NODE=M070P58

<sup>1</sup> LEES 18E reports  $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 1.7 \pm 0.8 \pm 0.3$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P58;LINKAGE=A

$$\Gamma(\omega\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{48}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>185±30±1</b>	14k	1 LEES	21	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

NODE=M070Q11  
NODE=M070Q11

<sup>1</sup> LEES 21 reports  $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 165 \pm 9 \pm 25$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q11;LINKAGE=A

$$\Gamma(\omega K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{59}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.70±1.98±0.03</b>	24	1 AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow \omega K^+ K^- \gamma$

NODE=M070G29  
NODE=M070G29

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \omega K\bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 3.3 \pm 1.3 \pm 1.2$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G29;LINKAGE=AU

$$\Gamma(K^+K^*(892)^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{68}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>29.0±1.7±1.3</b>	AUBERT	08s	BABR $10.6 e^+e^- \rightarrow K^+ K^*(892)^- \gamma$

NODE=M070G18  
NODE=M070G18

$$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{69}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.96±0.85±0.70</b>	155	AUBERT	08s	BABR $10.6 e^+e^- \rightarrow K^+ K^- \pi^0 \gamma$

NODE=M070G20  
NODE=M070G20

$$\Gamma(K^+K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{70}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.76±1.70±1.00</b>	89	AUBERT	08s	BABR $10.6 e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

NODE=M070G21  
NODE=M070G21

$$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{71}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>26.6±2.5±1.5</b>	AUBERT	08s	BABR $10.6 e^+e^- \rightarrow K^0 \bar{K}^*(892)^0 \gamma$

NODE=M070G19  
NODE=M070G19

$$\Gamma(K^0\bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{72}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17.70±1.70±1.00</b>	94	AUBERT	08s	BABR $10.6 e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

NODE=M070G22  
NODE=M070G22

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{73}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>42.6±4.8±7.2</b>	99	1 LEES	17D	BABR $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

NODE=M070G39  
NODE=M070G39

<sup>1</sup> Dividing by 1/6 to account for  $B(K^*(892)^0 \rightarrow K_S^0 \pi^0) = 1/6$ .

NODE=M070G39;LINKAGE=A

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{74}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>22.8±2.8±6.8</b>	80	1 LEES	17D	BABR $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

NODE=M070G32  
NODE=M070G32

<sup>1</sup> Dividing by 1/4 to account for  $B(K^*(892)^\pm \rightarrow K_S^0 \pi^\pm) = 1/4$ .

NODE=M070G32;LINKAGE=A

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{75} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.0±2.8 OUR AVERAGE</b>				
9.2±1.2±3.2	64	<sup>1</sup> LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$
14.8±4.8±1.2	53	<sup>2</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by 1/2 to take into account  $B(K^*(892)^\pm \rightarrow K^\pm \pi^\mp) = 1/2$ .

<sup>2</sup> Dividing by 1/4 to take into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ .

NODE=M070GY4  
NODE=M070GY4

NODE=M070GY4;LINKAGE=B  
NODE=M070GY4;LINKAGE=A

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{76} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.7±1.2±0.3</b>	53	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GY5  
NODE=M070GY5

$$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{79} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.60±0.75±2.25</b>	34	<sup>1</sup> LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

<sup>1</sup> Dividing by 2/3 to account for  $B(K^*(892)^0 \rightarrow K^+ \pi^-) = 2/3$ .

NODE=M070G33  
NODE=M070G33

NODE=M070G33;LINKAGE=A

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{81} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.28±0.34±0.07</b>	47±12	<sup>1</sup> LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.28±0.40±0.11	25±8	<sup>1,2</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> Dividing by  $(2/3)^2$  to take twice into account that  $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K \pi)$ .

<sup>2</sup> Superseded by LEES 12F.

NODE=M070G01  
NODE=M070G01

NODE=M070G01;LINKAGE=AE

NODE=M070G01;LINKAGE=A

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{82} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.80±0.48±0.32</b>	1±5	<sup>1</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by  $(1/4)^2$  to take twice into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ .

NODE=M070GY8  
NODE=M070GY8

NODE=M070GY8;LINKAGE=A

$$\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{93} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.1±9.8±0.5</b>	35	<sup>1,2</sup> LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

<sup>1</sup> Dividing by 1/4 to take into account  $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$ .

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GY6  
NODE=M070GY6

NODE=M070GY6;LINKAGE=A

NODE=M070GY6;LINKAGE=B

$$\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{92} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.65±0.80±0.44</b>	1094	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M070Q43  
NODE=M070Q43

$$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{94} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.8±1.4±0.6</b>	710	<sup>1,2,3</sup> LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ±4 ±1 317 <sup>2,4</sup> AUBERT 07AK BABR 10.6  $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 12.89 \pm 0.54 \pm 0.41$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Dividing by 2/3 to take into account that  $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K \pi)$ .

<sup>3</sup> The  $K_2^*(1430)$  cannot be distinguished from the  $K_0^*(1430)$ .

<sup>4</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 16.4 \pm 1.1 \pm 1.4$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G02  
NODE=M070G02

NODE=M070G02;LINKAGE=A

NODE=M070G02;LINKAGE=AE

NODE=M070G02;LINKAGE=B

NODE=M070G02;LINKAGE=UB



$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{95} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18.6 ± 16.1 ± 0.4</b>	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GY9  
NODE=M070GY9

<sup>1</sup> Dividing by  $(1/4)^2$  to take into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$  and  $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4$   $B(K^*(1430) \rightarrow K \pi)$ .

NODE=M070GY9;LINKAGE=A

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 9.28 \pm 8.0 \pm 0.32$  eV which we divide by our best value  $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GY9;LINKAGE=B

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{96} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.32 ± 2.00 ± 0.08</b>	8 ± 8	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GZ0  
NODE=M070GZ0

<sup>1</sup> Dividing by 1/4 to take into account  $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ .

NODE=M070GZ0;LINKAGE=A

$$\Gamma(\bar{K}_2(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{98} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8 ± 0.4 ± 0.3</b>	110 ± 14	1 AUBERT 07AK	BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M070G03  
NODE=M070G03

<sup>1</sup> Dividing by 2/3 to take into account that  $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$ .

NODE=M070G03;LINKAGE=AE

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{105} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.48 ± 0.35 OUR AVERAGE</b>				

NODE=M070G14  
NODE=M070G14

4.46 ± 0.49 ± 0.05      181      <sup>1</sup> LEES      12F BABR       $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

4.51 ± 0.48 ± 0.05      254 ± 23      <sup>2</sup> SHEN      09 BELL       $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.3 ± 0.7 ± 0.1      103      <sup>3</sup> AUBERT, BE 06D BABR       $10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.19 \pm 0.23 \pm 0.07$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14;LINKAGE=B

<sup>2</sup> SHEN 09 reports  $4.50 \pm 0.41 \pm 0.26$  eV from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)]$  assuming  $B(\phi(1020) \rightarrow K^+ K^-) = (49.2 \pm 0.6) \times 10^{-2}$ , which we rescale to our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14;LINKAGE=SH

<sup>3</sup> Superseded by LEES 12F. AUBERT, BE 06D reports  $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.61 \pm 0.30 \pm 0.18$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14;LINKAGE=AU

$$\Gamma(\phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{106} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.77 ± 0.57 ± 0.03</b>	45	<sup>1</sup> LEES	12F BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

NODE=M070G15  
NODE=M070G15

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.13 ± 0.88 ± 0.03      23      <sup>2</sup> AUBERT, BE 06D BABR       $10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.36 \pm 0.27 \pm 0.07$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G15;LINKAGE=A

<sup>2</sup> Superseded by LEES 12F. AUBERT, BE 06D reports  $[\Gamma(J/\psi(1S) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.54 \pm 0.40 \pm 0.16$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G15;LINKAGE=AU

$$\Gamma(\phi 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{107} \Gamma_5 / \Gamma$$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.96 ± 0.19 ± 0.01</b>	35	<sup>1</sup> AUBERT	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → φ2(π <sup>+</sup> π <sup>-</sup> )γ
<sup>1</sup> AUBERT 06D reports [Γ(J/ψ(1S) → φ2(π <sup>+</sup> π <sup>-</sup> )) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(φ(1020) → K <sup>+</sup> K <sup>-</sup> )] = (0.47 ± 0.09 ± 0.03) × 10 <sup>-2</sup> keV which we divide by our best value B(φ(1020) → K <sup>+</sup> K <sup>-</sup> ) = (49.1 ± 0.5) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G10  
NODE=M070G10

NODE=M070G10;LINKAGE=AU

$$\Gamma(\phi \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{108} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1 ± 2.7 ± 0.4</b>	6	<sup>1</sup> AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → φηγ
<sup>1</sup> AUBERT 07AU quotes Γ <sub>ee</sub> <sup>J/ψ</sup> · B(J/ψ → φη) · B(φ → K <sup>+</sup> K <sup>-</sup> ) · B(η → 3π) = 0.84 ± 0.37 ± 0.05 eV.				

NODE=M070G28  
NODE=M070G28

NODE=M070G28;LINKAGE=AU

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{112} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.44 ± 0.19 OUR AVERAGE</b>				
1.40 ± 0.25 ± 0.02	57 ± 9	<sup>1</sup> LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
1.48 ± 0.27 ± 0.09	60 ± 11	<sup>2</sup> SHEN	09 BELL	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.02 ± 0.24 ± 0.01	20 ± 5	<sup>3</sup> AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
<sup>1</sup> LEES 12F reports [Γ(J/ψ(1S) → φf <sub>0</sub> (980) → φπ <sup>+</sup> π <sup>-</sup> ) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(φ(1020) → K <sup>+</sup> K <sup>-</sup> )] = 0.69 ± 0.11 ± 0.05 eV which we divide by our best value B(φ(1020) → K <sup>+</sup> K <sup>-</sup> ) = (49.1 ± 0.5) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>2</sup> Multiplied by 2/3 to take into account the φπ <sup>+</sup> π <sup>-</sup> mode only. Using B(φ → K <sup>+</sup> K <sup>-</sup> ) = (49.2 ± 0.6)%.				
<sup>3</sup> Superseded by LEES 12F. AUBERT 07AK reports [Γ(J/ψ(1S) → φf <sub>0</sub> (980) → φπ <sup>+</sup> π <sup>-</sup> ) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(φ(1020) → K <sup>+</sup> K <sup>-</sup> )] = 0.50 ± 0.11 ± 0.04 eV which we divide by our best value B(φ(1020) → K <sup>+</sup> K <sup>-</sup> ) = (49.1 ± 0.5) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G05  
NODE=M070G05

NODE=M070G05;LINKAGE=A

NODE=M070G05;LINKAGE=SH

NODE=M070G05;LINKAGE=UB

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{113} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.98 ± 0.26 ± 0.01</b>	16 ± 4	<sup>1</sup> LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.96 ± 0.40 ± 0.01	7.0 ± 2.8	<sup>2</sup> AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ
<sup>1</sup> LEES 12F reports [Γ(J/ψ(1S) → φf <sub>0</sub> (980) → φπ <sup>0</sup> π <sup>0</sup> ) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(φ(1020) → K <sup>+</sup> K <sup>-</sup> )] = 0.48 ± 0.12 ± 0.05 eV which we divide by our best value B(φ(1020) → K <sup>+</sup> K <sup>-</sup> ) = (49.1 ± 0.5) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports [Γ(J/ψ(1S) → φf <sub>0</sub> (980) → φπ <sup>0</sup> π <sup>0</sup> ) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(φ(1020) → K <sup>+</sup> K <sup>-</sup> )] = 0.47 ± 0.19 ± 0.05 eV which we divide by our best value B(φ(1020) → K <sup>+</sup> K <sup>-</sup> ) = (49.1 ± 0.5) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G06  
NODE=M070G06

NODE=M070G06;LINKAGE=A

NODE=M070G06;LINKAGE=UB

$$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{118} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.79 ± 0.32<sup>+0.02</sup><sub>-0.06</sub></b>	61	<sup>1,2,3</sup> LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.08 ± 0.73 <sup>+0.04</sup> <sub>-0.14</sub>	44	<sup>2,4</sup> AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
<sup>1</sup> LEES 12F reports [Γ(J/ψ(1S) → φf <sub>2</sub> (1270)) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(f <sub>2</sub> (1270) → ππ)] = 1.51 ± 0.25 ± 0.10 eV which we divide by our best value B(f <sub>2</sub> (1270) → ππ) = (84.3 <sup>+2.9</sup> <sub>-0.9</sub> ) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>2</sup> Using B(φ → K <sup>+</sup> K <sup>-</sup> ) = (48.9 ± 0.5)%.				
<sup>3</sup> Using π <sup>+</sup> π <sup>-</sup> invariant mass between 1.1 and 1.5 GeV. May include other sources such as f <sub>0</sub> (1370).				
<sup>4</sup> Superseded by LEES 12F. AUBERT 07AK reports [Γ(J/ψ(1S) → φf <sub>2</sub> (1270)) × Γ(J/ψ(1S) → e <sup>+</sup> e <sup>-</sup> )/Γ <sub>total</sub> ] × [B(f <sub>2</sub> (1270) → ππ)] = 3.44 ± 0.55 ± 0.28 eV which we divide by our best value B(f <sub>2</sub> (1270) → ππ) = (84.3 <sup>+2.9</sup> <sub>-0.9</sub> ) × 10 <sup>-2</sup> . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G07  
NODE=M070G07

NODE=M070G07;LINKAGE=A

NODE=M070G07;LINKAGE=AE  
NODE=M070G07;LINKAGE=B

NODE=M070G07;LINKAGE=UB

$$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{123} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.2 ± 3.2 ± 0.2</b>	11	1,2 LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

NODE=M070GZ2  
NODE=M070GZ2

<sup>1</sup> Dividing by 1/4 to take into account  $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K \bar{K})$  and using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ .

NODE=M070GZ2;LINKAGE=A

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = 7.2 \pm 2.8 \pm 0.3$  eV which we divide by our best value  $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GZ2;LINKAGE=B

$$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{129} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.60 ± 0.62 ± 0.05</b>	163	<sup>1</sup> LEES	12F	BABR $10.6 e^+ e^- \rightarrow K^+ K^- K^+ K^- \gamma$

NODE=M070G09  
NODE=M070G09

<sup>1</sup> LEES 12F reports  $[\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 2.26 \pm 0.26 \pm 0.16$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G09;LINKAGE=A

$$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{130} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.26 ± 0.84 ± 0.04</b>	29	<sup>1</sup> LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

NODE=M070GZ1  
NODE=M070GZ1

<sup>1</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+ K^-)] = 1.6 \pm 0.4 \pm 0.1$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+ K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GZ1;LINKAGE=A

$$\Gamma(f'_2(1525) K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{135} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.8 ± 1.9 ± 0.1</b>	16	1,2 LEES	14H	BABR $e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$

NODE=M070GZ4  
NODE=M070GZ4

<sup>1</sup> Dividing by 1/4 to take into account  $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K \bar{K})$ .

NODE=M070GZ4;LINKAGE=A

<sup>2</sup> LEES 14H reports  $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525) K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = 5.12 \pm 1.68 \pm 0.20$  eV which we divide by our best value  $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GZ4;LINKAGE=B

$$\Gamma(2(\pi^+ \pi^-) \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{154} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>303 ± 5 ± 18</b>	4990	AUBERT	07AU	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0 \gamma$

NODE=M070G23  
NODE=M070G23

$$\Gamma(\pi^+ \pi^- 3\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{156} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>104 ± 50 OUR AVERAGE</b>				Error includes scale factor of 4.3.

NODE=M070P53  
NODE=M070P53

55.4 ± 15.9 ± 0.5	14k	<sup>1</sup> LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$
150.0 ± 4.0 ± 15.0	2.3k	LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

<sup>1</sup> LEES 21 reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+ \pi^- 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}] = 19.2 \pm 4.5 \pm 3.2$  eV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) / \Gamma_{\text{total}} = 0.3468 \pm 0.0030$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P53;LINKAGE=A

$$\Gamma(\pi^+ \pi^- 4\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{157} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>35.8 ± 4.4 ± 5.4</b>	340	LEES	21C	BABR $e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

NODE=M070Q17  
NODE=M070Q17

$$\Gamma(\rho^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{158} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>78.0 ± 9.0 ± 8.0</b>	1.2k	LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

NODE=M070P55  
NODE=M070P55

$$\Gamma(\rho^+ \rho^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{159} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>33.0 ± 5.0 ± 3.3</b>	529	LEES	18E	BABR $10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

NODE=M070P56  
NODE=M070P56

$$\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{160}\Gamma_5/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.1248±0.0019±0.0026</b>	LEES	21B	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ
0.122 ±0.005 ±0.008	AUBERT,B	04N	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

NODE=M070G5  
NODE=M070G5

$$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{161}\Gamma_5/\Gamma$$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.9±0.5±1.0</b>	761	AUBERT	06D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> )γ

NODE=M070G7  
NODE=M070G7

$$\Gamma(\pi^+\pi^-\pi^0 K^+ K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{162}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>107.0±4.3±6.4</b>	768	AUBERT	07AU	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

NODE=M070G27  
NODE=M070G27

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{164}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.4±0.9±0.4</b>		LEES	12E	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2π <sup>+</sup> 2π <sup>-</sup> γ
19.5±1.4±1.3	270	<sup>1</sup> AUBERT	05D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )γ

NODE=M070G11  
NODE=M070G11

<sup>1</sup> Superseded by LEES 12E.

NODE=M070G11;LINKAGE=AU

$$\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{165}\Gamma_5/\Gamma$$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.37±0.16±0.14</b>	496	AUBERT	06D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 3(π <sup>+</sup> π <sup>-</sup> )γ

NODE=M070G6  
NODE=M070G6

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{166}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>345±10±50</b>	14k	LEES	21	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )3π <sup>0</sup> γ

NODE=M070Q12  
NODE=M070Q12

$$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{168}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.1±2.4±0.1</b>	85	<sup>1</sup> AUBERT	07AU	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )ηγ

NODE=M070G26  
NODE=M070G26

<sup>1</sup> AUBERT 07AU reports [Γ(J/ψ(1S) → 2(π<sup>+</sup>π<sup>-</sup>)η) × Γ(J/ψ(1S) → e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>] × [B(η → 2γ)] = 5.16 ± 0.85 ± 0.39 eV which we divide by our best value B(η → 2γ) = (39.36 ± 0.18) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G26;LINKAGE=AU

$$\Gamma(2(\pi^+\pi^-\pi^0)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{170}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.1±2.6±1.4</b>	14k	LEES	21	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )3π <sup>0</sup> γ

NODE=M070Q15  
NODE=M070Q15

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{171}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.1± 2.7 OUR AVERAGE</b>				
26.1±17.9±0.3	14k	<sup>1</sup> LEES	21	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )3π <sup>0</sup> γ
12.8± 1.8±2.0	203	LEES	18E	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> π <sup>0</sup> ηγ

NODE=M070P57  
NODE=M070P57

<sup>1</sup> LEES 21 reports [Γ(J/ψ(1S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>π<sup>0</sup>η) × Γ(J/ψ(1S) → e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>] × [B(η → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>)] = 6 ± 4 ± 1 eV which we divide by our best value B(η → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>) = (23.02 ± 0.25) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P57;LINKAGE=AU

$$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{172}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.5±4.1±1.6</b>	168	LEES	18E	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> π <sup>0</sup> ηγ

NODE=M070P59  
NODE=M070P59

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{173}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
1.78±0.11±0.05	462	<sup>1</sup> LEES	15J	BABR e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> γ
1.94±0.11±0.05	462	<sup>2</sup> LEES	15J	BABR e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> γ
1.42±0.23±0.08	51	<sup>3</sup> LEES	13Q	BABR e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> γ

NODE=M070G08  
NODE=M070G08

OCCUR=2

<sup>1</sup> sinφ > 0.

<sup>2</sup> sinφ < 0.

<sup>3</sup> Interference with non-resonant K<sup>+</sup>K<sup>-</sup> production not taken into account.

NODE=M070G08;LINKAGE=A  
NODE=M070G08;LINKAGE=B  
NODE=M070G08;LINKAGE=BA

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{179} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.4 ± 1.3 ± 0.6</b>	182	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

NODE=M070G41  
NODE=M070G41

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{180} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.7 ± 0.9 ± 0.4</b>	106	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

NODE=M070G42  
NODE=M070G42

$$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{181} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4 ± 0.7 ± 0.1</b>	37	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \gamma$

NODE=M070G43  
NODE=M070G43

$$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{182} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>37.94 ± 0.81 ± 1.10</b>	3.1k	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M070G12  
NODE=M070G12  
OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

36.3 ± 1.3 ± 2.1	1.5k	<sup>1</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
33.6 ± 2.7 ± 2.7	233	<sup>2</sup> AUBERT	05D BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

<sup>1</sup> Superseded by LEES 12F.

<sup>2</sup> Superseded by AUBERT 07AK.

NODE=M070G12;LINKAGE=B  
NODE=M070G12;LINKAGE=A

$$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{183} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.75 ± 0.81 ± 0.90</b>	388	LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$

NODE=M070G04  
NODE=M070G04

• • • We do not use the following data for averages, fits, limits, etc. • • •

13.6 ± 1.1 ± 1.3	203	<sup>1</sup> AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$
------------------	-----	---------------------	-----------	---

<sup>1</sup> Superseded by LEES 12F.

NODE=M070G04;LINKAGE=A

$$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{184} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.8 ± 2.3 ± 2.1</b>	248	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$

NODE=M070GY1  
NODE=M070GY1

$$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{185} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.3 ± 2.3 ± 0.5</b>	47	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$

NODE=M070G40  
NODE=M070G40

$$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{186} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.0 ± 1.8 ± 0.4</b>	45	LEES	17A BABR	$e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$

NODE=M070G35  
NODE=M070G35

$$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{187} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.3 ± 0.9 ± 0.5</b>	133	LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GY2  
NODE=M070GY2

$$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{188} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>31.7 ± 1.9 ± 1.8</b>	393	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

NODE=M070G34  
NODE=M070G34

$$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{189} \Gamma_5 / \Gamma$$

VALUE (10 <sup>-2</sup> keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.75 ± 0.23 ± 0.17</b>	205	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

NODE=M070G9  
NODE=M070G9

$$\Gamma(K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{190} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.9 ± 3.9 ± 0.1</b>	73	<sup>1</sup> AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$

NODE=M070G30  
NODE=M070G30

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow K^+ K^- \pi^+ \pi^- \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G30;LINKAGE=AU

$$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{191}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.00±0.33±0.29</b>	287 ± 24	LEES	12F	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(K <sup>+</sup> K <sup>-</sup> )γ
4.11±0.39±0.30	156 ± 15	<sup>1</sup> AUBERT	07AK	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(K <sup>+</sup> K <sup>-</sup> )γ
4.0 ±0.7 ±0.6	38	<sup>2</sup> AUBERT	05D	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → 2(K <sup>+</sup> K <sup>-</sup> )γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>Superseded by LEES 12F.

<sup>2</sup>Superseded by AUBERT 07AK.

NODE=M070G13  
NODE=M070G13

NODE=M070G13;LINKAGE=A  
NODE=M070G13;LINKAGE=A

$$\Gamma(K^+K^-K_S^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{192}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3±0.4±0.1</b>	29	LEES	14H	BABR e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>S</sub> <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ

NODE=M070GY3  
NODE=M070GY3

$$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{193}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.9±0.6 OUR AVERAGE</b>	Error includes scale factor of 1.8. See the ideogram below.			
11.3±0.4±0.3	821	<sup>1</sup> LEES	13O	BABR e <sup>+</sup> e <sup>-</sup> → p $\bar{p}$ γ
12.9±0.4±0.4	918	<sup>2</sup> LEES	13Y	BABR e <sup>+</sup> e <sup>-</sup> → p $\bar{p}$ γ
9.7±1.7		<sup>3</sup> ARMSTRONG	93B	E760 $\bar{p}p$ → e <sup>+</sup> e <sup>-</sup>
12.0±0.6±0.5	438	<sup>4</sup> AUBERT	06B	BABR e <sup>+</sup> e <sup>-</sup> → p $\bar{p}$ γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>ISR photon reconstructed in the detector

<sup>2</sup>ISR photon undetected

<sup>3</sup>Using  $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$  MeV.


<sup>4</sup>Superseded by LEES 13O

NODE=M070G4  
NODE=M070G4

NODE=M070G4;LINKAGE=B  
NODE=M070G4;LINKAGE=C

NODE=M070G;LINKAGE=A  
NODE=M070G4;LINKAGE=A

WEIGHTED AVERAGE  
11.9±0.6 (Error scaled by 1.8)



$$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{209}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>10.7±0.9±0.7</b>	AUBERT	07BD	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → Λ $\bar{\Lambda}$ γ

NODE=M070G16  
NODE=M070G16

$$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{219}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>6.4±1.2±0.6</b>	AUBERT	07BD	BABR 10.6 e <sup>+</sup> e <sup>-</sup> → Σ <sup>0</sup> Σ <sup>0</sup> γ

NODE=M070G17  
NODE=M070G17

**$J/\psi(1S)$  BRANCHING RATIOS**

NODE=M070230

For the first four branching ratios, see also the partial widths, and (partial widths)  $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  above.

NODE=M070300

 **$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$**  **$\Gamma_1/\Gamma$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.877±0.005 OUR AVERAGE</b>			
0.878±0.005	BAI	95B	BES $e^+e^-$
0.86 ±0.02	BOYARSKI	75	MRK1 $e^+e^-$

NODE=M070R3  
NODE=M070R3 **$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$**  **$\Gamma_2/\Gamma$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.135±0.003</b>	1,2 SETH	04	RVUE $e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.17 ±0.02	<sup>1</sup> BOYARSKI	75	MRK1 $e^+e^-$

NODE=M070R4  
NODE=M070R4<sup>1</sup> Included in  $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ .<sup>2</sup> Using  $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$  from RPP-2002 and  $R = 2.28 \pm 0.04$  determined by a fit to data from BAI 00 and BAI 02c.NODE=M070R4;LINKAGE=C  
NODE=M070R4;LINKAGE=SE **$\Gamma(gg)/\Gamma_{\text{total}}$**  **$\Gamma_3/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>64.1±1.0</b>	6 M	<sup>1</sup> BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- + \text{hadrons}$

NODE=M070S65  
NODE=M070S65

<sup>1</sup> Calculated using the value  $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$  from BESSON 08 and the PDG 08 values of  $B(\ell^+\ell^-)$ ,  $B(\text{virtual } \gamma \rightarrow \text{hadrons})$ , and  $B(\gamma\eta_c)$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma gg)/\Gamma_{\text{total}}$  measurement of BESSON 08.

NODE=M070S65;LINKAGE=BE

 **$\Gamma(\gamma gg)/\Gamma_{\text{total}}$**  **$\Gamma_4/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.79±1.05</b>	200 k	<sup>1</sup> BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^-\gamma + \text{hadrons}$

NODE=M070S66  
NODE=M070S66

<sup>1</sup> Calculated using the value  $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$  from BESSON 08 and the value of  $\Gamma(ggg)/\Gamma_{\text{total}}$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(ggg)/\Gamma_{\text{total}}$  measurement of BESSON 08.

NODE=M070S66;LINKAGE=BE

 **$\Gamma(\gamma gg)/\Gamma(ggg)$**  **$\Gamma_4/\Gamma_3$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.7±0.1±0.7</b>	6 M	BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

NODE=M070S67  
NODE=M070S67 **$\Gamma(e^+e^-)/\Gamma_{\text{total}}$**  **$\Gamma_5/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.971±0.032 OUR AVERAGE</b>				
5.983±0.007±0.037	720k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
5.945±0.067±0.042	15k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
5.90 ±0.05 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
6.09 ±0.33		BAI	95B	BES $e^+e^-$
5.92 ±0.15 ±0.20		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+e^-$

NODE=M070R1  
NODE=M070R1 **$\Gamma(e^+e^-\gamma)/\Gamma_{\text{total}}$**  **$\Gamma_6/\Gamma$** 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.8±1.3±0.4</b>	<sup>1</sup> ARMSTRONG	96	E760 $\bar{p}p \rightarrow e^+e^-\gamma$

NODE=M070S33  
NODE=M070S33<sup>1</sup> For  $E_\gamma > 100$  MeV.

NODE=M070S33;LINKAGE=A

 **$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$**  **$\Gamma_7/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.961±0.033 OUR AVERAGE</b>				
5.973±0.007±0.038	770k	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
5.960±0.065±0.050	17k	LI	05C	CLEO $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
5.84 ±0.06 ±0.10		BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
6.08 ±0.33		BAI	95B	BES $e^+e^-$
5.90 ±0.15 ±0.19		COFFMAN	92	MRK3 $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
6.9 ±0.9		BOYARSKI	75	MRK1 $e^+e^-$

NODE=M070R2  
NODE=M070R2

$\Gamma(e^+e^-)/\Gamma(\mu^+\mu^-)$

$\Gamma_5/\Gamma_7$

NODE=M070R5  
NODE=M070R5

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.0016 ± 0.0031 OUR AVERAGE</b>			
1.0022 ± 0.0044 ± 0.0048	<sup>1</sup> AULCHENKO 14	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
1.0017 ± 0.0017 ± 0.0033	<sup>2</sup> ABLIKIM 13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
1.002 ± 0.021 ± 0.013	<sup>3</sup> ANASHIN 10	KEDR	3.097 $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
0.997 ± 0.012 ± 0.006	LI 05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.011 ± 0.013 ± 0.016	BAI	98D	BES $\psi(2S) \rightarrow J/\psi \pi^+\pi^-$
1.00 ± 0.07	BAI	95B	BES $e^+e^-$
1.00 ± 0.05	BOYARSKI 75	MRK1	$e^+e^-$
0.91 ± 0.15	ESPOSITO 75B	FRAM	$e^+e^-$
0.93 ± 0.10	FORD 75	SPEC	$e^+e^-$

- <sup>1</sup> From 235.3k  $J/\psi \rightarrow e^+e^-$  and 156.6k  $J/\psi \rightarrow \mu^+\mu^-$  observed events.
- <sup>2</sup> Not independent of the corresponding measurements of  $\Gamma(e^+e^-)/\Gamma_{total}$  and  $\Gamma(\mu^+\mu^-)/\Gamma_{total}$ .
- <sup>3</sup> Not independent of the corresponding measurements of  $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{total}$  and  $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{total}$ .

NODE=M070R5;LINKAGE=A  
NODE=M070R5;LINKAGE=AB

NODE=M070R5;LINKAGE=AN

———— HADRONIC DECAYS ————

NODE=M070305

$\Gamma(\rho\pi)/\Gamma_{total}$

$\Gamma_8/\Gamma$

NODE=M070R20  
NODE=M070R20


VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.69 ± 0.15 OUR AVERAGE</b>				
Error includes scale factor of 2.4. See the ideogram below.				
2.18 ± 0.19		<sup>1,2</sup> AUBERT,B	04N	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
2.184 ± 0.005 ± 0.201	220k	<sup>2,3</sup> BAI	04H	BES $e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
2.091 ± 0.021 ± 0.116		<sup>2,4</sup> BAI	04H	BES $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
1.21 ± 0.20		BAI	96D	BES $e^+e^- \rightarrow \rho\pi$
1.42 ± 0.01 ± 0.19		COFFMAN	88	MRK3 $e^+e^-$
1.3 ± 0.3	150	FRANKLIN	83	MRK2 $e^+e^-$
1.6 ± 0.4	183	ALEXANDER	78	PLUT $e^+e^-$
1.33 ± 0.21		BRANDELIK	78B	DASP $e^+e^-$
1.0 ± 0.2	543	BARTEL	76	CNTR $e^+e^-$
1.3 ± 0.3	153	JEAN-MARIE	76	MRK1 $e^+e^-$

OCCUR=2

- <sup>1</sup> From the ratio of  $\Gamma(e^+e^-) B(\pi^+\pi^-\pi^0)$  and  $\Gamma(e^+e^-) B(\mu^+\mu^-)$  (AUBERT 04).
- <sup>2</sup> Not independent of their  $B(\pi^+\pi^-\pi^0)$ .
- <sup>3</sup> From  $J/\psi \rightarrow \pi^+\pi^-\pi^0$  events directly.
- <sup>4</sup> Obtained comparing the rates for  $\pi^+\pi^-\pi^0$  and  $\mu^+\mu^-$ , using  $J/\psi$  events produced via  $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$  and with  $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$ .

NODE=M070R20;LINKAGE=AU  
NODE=M070R20;LINKAGE=BU  
NODE=M070R20;LINKAGE=BA  
NODE=M070R20;LINKAGE=BI

WEIGHTED AVERAGE  
1.69±0.15 (Error scaled by 2.4)





$\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$  $\Gamma_8/\Gamma_{160}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.142±0.011±0.026</b>	20k	<sup>1</sup> LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
1.331±0.033	20k	<sup>2</sup> LEES	17C	BABR $J/\psi \rightarrow \pi^+\pi^-\pi^0$

NODE=M070P18  
NODE=M070P18

OCCUR=2

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.NODE=M070P18;LINKAGE=A  
NODE=M070P18;LINKAGE=B $\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$  $\Gamma_9/\Gamma_8$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.328±0.005±0.027</b>	COFFMAN	88	MRK3 $e^+e^-$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.35 ±0.08	ALEXANDER	78	PLUT $e^+e^-$
0.32 ±0.08	BRANDELIK	78B	DASP $e^+e^-$
0.39 ±0.11	BARTEL	76	CNTR $e^+e^-$
0.37 ±0.09	JEAN-MARIE	76	MRK1 $e^+e^-$

NODE=M070R21  
NODE=M070R21 $\Gamma(\pi^+K^+K^-\pi^- + c.c \rightarrow K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{total}$  $\Gamma_{24}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.53±0.16±0.81</b>	485	ANASHIN	22	KEDR $J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M070Q41  
NODE=M070Q41 $\Gamma(a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{total}$  $\Gamma_{10}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.84±0.08±0.60</b>	1317	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M070Q39  
NODE=M070Q39 $\Gamma(a_2(1320)^+\pi^-\pi^0 + c.c \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{total}$  $\Gamma_{11}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.67±0.09±0.73</b>	1628	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M070Q40  
NODE=M070Q40 $\Gamma(a_2(1320)\rho)/\Gamma_{total}$  $\Gamma_{12}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.9±2.2 OUR AVERAGE</b>				
11.7±0.7±2.5	7584	AUGUSTIN	89	DM2 $J/\psi \rightarrow \rho^0\rho^\pm\pi^\mp$
8.4±4.5	36	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M070R43  
NODE=M070R43 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{total}$  $\Gamma_{13}/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.78±0.68</b>	471	<sup>1</sup> ABLIKIM	19Q	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$

NODE=M070P81  
NODE=M070P81<sup>1</sup> From an energy scan of  $e^+e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$  assuming PDG 16 values for  $\Gamma(e^+e^-)$ ,  $\Gamma(\mu^+\mu^-)$ , and  $\Gamma(total)$ .

NODE=M070P81;LINKAGE=A

 $\Gamma(\eta\rho)/\Gamma_{total}$  $\Gamma_{16}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.193±0.023 OUR AVERAGE</b>				
0.194±0.017±0.029	299	JOUSSET	90	DM2 $J/\psi \rightarrow hadrons$
0.193±0.013±0.029		COFFMAN	88	MRK3 $e^+e^- \rightarrow \pi^+\pi^-\eta$

NODE=M070R22  
NODE=M070R22 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{total}$  $\Gamma_{17}/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.20±0.14±0.37</b>	471	ABLIKIM	15H	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$

NODE=M070B12  
NODE=M070B12 $\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{total}$  $\Gamma_{18}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.52 × 10<sup>-4</sup></b>	90	ABLIKIM	10C	BES2 $J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

NODE=M070S70  
NODE=M070S70 $\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{total}$  $\Gamma_{19}/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.8±2.2±3.4</b>	232 ± 23	ABLIKIM	08E	BES2 $e^+e^- \rightarrow J/\psi$

NODE=M070S57  
NODE=M070S57 $\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{total}$  $\Gamma_{20}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.15±0.13±0.22</b>	209	ABLIKIM	10C	BES2 $J/\psi \rightarrow \eta K^+\pi^- K^-\pi^+$

NODE=M070S69  
NODE=M070S69

$\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.1 ± 0.8 OUR AVERAGE</b>	Error	includes scale factor of 1.6.		
7.90 ± 0.19 ± 0.49	3476	<sup>1</sup> ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$
8.3 ± 3.0 ± 1.2	19	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
11.4 ± 1.4 ± 1.6		COFFMAN	88 MRK3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

<sup>1</sup> From a partial wave analysis of the decay  $J/\psi \rightarrow \pi^+ \pi^- \eta'$ .NODE=M070R23  
NODE=M070R23

NODE=M070R23;LINKAGE=A

 $\Gamma(\rho(1450)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$  $\Gamma_{29}/\Gamma_{160}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.9 ± 1.7 ± 2.7</b>	20k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.80 ± 0.27	20k	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.NODE=M070P25  
NODE=M070P25

OCCUR=2

NODE=M070P25;LINKAGE=A  
NODE=M070P25;LINKAGE=B $\Gamma(\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$  $\Gamma_{30}/\Gamma_{178}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.3 ± 0.8 ± 0.6</b>	4k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.NODE=M070P31  
NODE=M070P31

NODE=M070P31;LINKAGE=A

 $\Gamma(\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0)/\Gamma(K^+ K^- \pi^0)$  $\Gamma_{31}/\Gamma_{177}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.3 ± 2.0 ± 0.6</b>	2k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.NODE=M070P27  
NODE=M070P27

NODE=M070P27;LINKAGE=A

 $\Gamma(\rho(1450)\eta'(958) \rightarrow \pi^+ \pi^- \eta'(958))/\Gamma_{\text{total}}$  $\Gamma_{32}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.28 ± 0.55 ± 0.44</b>	119	<sup>1</sup> ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

<sup>1</sup> From a partial wave analysis of the decay  $J/\psi \rightarrow \pi^+ \pi^- \eta'$ .NODE=M070P36  
NODE=M070P36

NODE=M070P36;LINKAGE=A

 $\Gamma(\rho(1700)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$  $\Gamma_{34}/\Gamma_{160}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8 ± 2 ± 5</b>	20k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22 ± 6	20k	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.NODE=M070P21  
NODE=M070P21

OCCUR=2

NODE=M070P21;LINKAGE=A  
NODE=M070P21;LINKAGE=B $\Gamma(\rho(2150)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$  $\Gamma_{36}/\Gamma_{160}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4 ± 1 ± 20</b>	20k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
600 ± 250	20k	<sup>2</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in an isobar model.<sup>2</sup> From a Dalitz plot analysis in a Veneziano model.NODE=M070P22  
NODE=M070P22

OCCUR=2

NODE=M070P22;LINKAGE=A  
NODE=M070P22;LINKAGE=B $\Gamma(\rho_3(1690)\pi \rightarrow \pi^+ \pi^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$  $\Gamma_{37}/\Gamma_{160}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.0 ± 0.8	20k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> From a Dalitz plot analysis in a Veneziano model.NODE=M070P24  
NODE=M070P24

NODE=M070P24;LINKAGE=A


 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.45 ± 0.05 OUR AVERAGE</b>	Error	includes scale factor of 1.4. See the ideogram below.		
0.538 ± 0.012 ± 0.065	2090	<sup>1</sup> ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.482 ± 0.019 ± 0.064		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

NODE=M070R32  
NODE=M070R32

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$ .

NODE=M070R32;LINKAGE=BL



$\Gamma(\omega\pi^0 \rightarrow \pi^+ \pi^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$   $\Gamma_{39}/\Gamma_{160}$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8±3±2</b>	20k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

NODE=M070P23  
NODE=M070P23

<sup>1</sup> From a Dalitz plot analysis in an isobar model and significance 4.9  $\sigma$ .

NODE=M070P23;LINKAGE=A


$\Gamma(\omega\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.5±1.0 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.		
[(7.2 ± 1.0) × 10 <sup>-3</sup> OUR 2022 AVERAGE]				
10.6±1.2±0.1	3531	<sup>1</sup> ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$
7.8±1.6	215	BURMESTER	77D PLUT	$e^+ e^-$
6.8±1.9	348	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-)\pi^0$

NODE=M070R24  
NODE=M070R24  
NEW

<sup>1</sup> ANASHIN 22 reports [ $\Gamma(J/\psi(1S) \rightarrow \omega\pi^+ \pi^-)/\Gamma_{\text{total}}$ ] × [B( $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ )] = (0.946 ± 0.016 ± 0.108) × 10<sup>-2</sup> which we divide by our best value B( $\omega(782) \rightarrow \pi^+ \pi^- \pi^0$ ) = (89.2 ± 0.7) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R24;LINKAGE=A



$\Gamma(\omega\pi^0\pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{41}/\Gamma$
<b>3.4±0.3±0.7</b>	509	AUGUSTIN	89	DM2	$J/\psi \rightarrow \pi^+\pi^-3\pi^0$

NODE=M070S26  
NODE=M070S26

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{43}/\Gamma$
<b>4.3±0.6 OUR AVERAGE</b>					

NODE=M070R28  
NODE=M070R28

4.3±0.2±0.6 5860 AUGUSTIN 89 DM2  $e^+e^-$

4.0±1.6 70 BURMESTER 77D PLUT  $e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.9±0.8 81 VANNUCCI 77 MRK1  $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{44}/\Gamma$
<b>1.74 ±0.20 OUR AVERAGE</b>				Error includes scale factor of 1.6. See the ideogram below.	

NODE=M070R30  
NODE=M070R30

2.352±0.273 5k <sup>1</sup> ABLIKIM 06F BES2  $J/\psi \rightarrow \omega\eta$

1.44 ±0.40 ±0.14 13 <sup>2</sup> AUBERT 06D BABR 10.6  $e^+e^- \rightarrow \omega\eta\gamma$

1.43 ±0.10 ±0.21 378 JOUSSET 90 DM2  $J/\psi \rightarrow \text{hadrons}$


1.71 ±0.08 ±0.20 COFFMAN 88 MRK3  $e^+e^- \rightarrow 3\pi\eta$

<sup>1</sup> Using  $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$ ,  $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$ ,  $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$ , and  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$ .

NODE=M070R30;LINKAGE=BL

<sup>2</sup> Using  $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$  keV.

NODE=M070R30;LINKAGE=EE

 $\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{47}/\Gamma$
<b>85±34</b>	140	VANNUCCI	77	MRK1	$e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

NODE=M070R26  
NODE=M070R26

 $\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{49}/\Gamma$
<b>1.12±0.02±0.13</b>	14k	<sup>1</sup> ABLIKIM	19AC	BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

NODE=M070P83  
NODE=M070P83

<sup>1</sup> Using the decays  $\omega \rightarrow \pi^+\pi^-\pi^0$  and  $\eta' \rightarrow \eta\pi^+\pi^-$ .

NODE=M070P83;LINKAGE=A

 $\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{50}/\Gamma$
<b>1.89±0.18 OUR AVERAGE</b>					

NODE=M070R31  
NODE=M070R31

2.08±0.30±0.14 137 <sup>1</sup> ABLIKIM 17AK BES3  $J/\psi \rightarrow \pi^+\pi^-\eta'$

2.26±0.43 218 <sup>2</sup> ABLIKIM 06F BES2  $J/\psi \rightarrow \omega\eta'$

1.8  $\begin{matrix} +1.0 \\ -0.8 \end{matrix}$  ±0.3 6 JOUSSET 90 DM2  $J/\psi \rightarrow \text{hadrons}$

1.66±0.17±0.19 COFFMAN 88 MRK3  $e^+e^- \rightarrow 3\pi\eta'$

<sup>1</sup> From a partial wave analysis of the decay  $J/\psi \rightarrow \pi^+ \pi^- \eta'$ .

<sup>2</sup> Using  $B(\eta' \rightarrow \pi^+ \pi^- \eta) = (44.3 \pm 1.5)\%$ ,  $B(\eta' \rightarrow \pi^+ \pi^- \gamma) = 29.5 \pm 1.0\%$ ,  $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$ , and  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$ .

NODE=M070R31;LINKAGE=A  
NODE=M070R31;LINKAGE=BL

### $\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ $\Gamma_{51}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.41 ± 0.27 ± 0.47</b>	<sup>1</sup> AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M070S27  
NODE=M070S27

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 0.78$ .

NODE=M070S27;LINKAGE=K

### $\Gamma(\omega f_0(1710) \rightarrow \omega K \bar{K})/\Gamma_{\text{total}}$ $\Gamma_{52}/\Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.8 ± 1.1 ± 0.3</b>	<sup>1,2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S25  
NODE=M070S25

<sup>1</sup> Includes unknown branching fraction  $f_0(1710) \rightarrow K \bar{K}$ .

NODE=M070S25;LINKAGE=F  
NODE=M070S25;LINKAGE=G

<sup>2</sup> Addition of  $f_0(1710) \rightarrow K^+ K^-$  and  $f_0(1710) \rightarrow K^0 \bar{K}^0$  branching ratios.

### $\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ $\Gamma_{53}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.8<sup>+1.9</sup><sub>-1.6</sub> ± 1.7</b>	111 <sup>+31</sup> <sub>-26</sub>	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S5  
NODE=M070S5

### $\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ $\Gamma_{54}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.2 × 10<sup>-4</sup></b>	90	<sup>1</sup> VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 K^+ K^-$

NODE=M070R29  
NODE=M070R29

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.8 × 10 <sup>-4</sup>	90	<sup>1</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
--------------------------	----	----------------------	--------	-------------------------------------

<sup>1</sup> Re-evaluated assuming  $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$ .

NODE=M070R29;LINKAGE=C

### $\Gamma(\omega X(1835) \rightarrow \omega p \bar{p})/\Gamma_{\text{total}}$ $\Gamma_{55}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 3.9 × 10<sup>-6</sup></b>	95	ABLIKIM	13P BES3	$J/\psi \rightarrow \gamma \pi^0 p \bar{p}$

NODE=M070S81  
NODE=M070S81

### $\Gamma(\omega X(1835), X \rightarrow \eta' \pi^+ \pi^-)/\Gamma_{\text{total}}$ $\Gamma_{56}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6.2 × 10<sup>-5</sup></b>	<sup>1</sup> ABLIKIM	19AC BES3	$J/\psi \rightarrow \omega \eta' \pi^+ \pi^-$

NODE=M070P84  
NODE=M070P84

<sup>1</sup> Using the decays  $\omega \rightarrow \pi^+ \pi^- \pi^0$  and  $\eta' \rightarrow \eta \pi^+ \pi^-$ .

NODE=M070P84;LINKAGE=A

### $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ $\Gamma_{57}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.52 ± 0.30 ± 0.01</b>	276	<sup>1</sup> ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M070Q38  
NODE=M070Q38

<sup>1</sup> ANASHIN 22 reports  $[\Gamma(J/\psi(1S) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$  which we divide by our best value  $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q38;LINKAGE=A

### $\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ $\Gamma_{58}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>34 ± 5 OUR AVERAGE</b>				
37.7 ± 0.8 ± 5.8	1972 ± 41	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S1  
NODE=M070S1

### $\Gamma(\omega K \bar{K})/\Gamma_{\text{total}}$ $\Gamma_{59}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>19 ± 4 OUR AVERAGE</b>				
19.8 ± 2.1 ± 3.9		<sup>1</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN	77 MRK1	$e^+ e^-$

NODE=M070R27  
NODE=M070R27

<sup>1</sup> Addition of  $\omega K^+ K^-$  and  $\omega K^0 \bar{K}^0$  branching ratios.

NODE=M070R27;LINKAGE=B

### $\Gamma(\omega K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ $\Gamma_{60}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>61 ± 9 OUR AVERAGE</b>				
62.0 ± 6.8 ± 10.6	899 ± 98	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
65.3 ± 10.2 ± 13.5	176 ± 28	ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
53 ± 14 ± 14	530 ± 140	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S2  
NODE=M070S2

OCCUR=2

$\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**1.48±0.13 OUR AVERAGE**

1.50±0.02±0.19

1.47±0.03±0.17

<sup>1</sup> From  $\eta' K^+ K^- \pi^0$ .<sup>2</sup> From  $\eta' K_S^0 K^\pm \pi^\mp$ .

DOCUMENT ID TECN COMMENT

<sup>1</sup> ABLIKIM 18AB BES3  $J/\psi \rightarrow \eta' K^* \bar{K}$ <sup>2</sup> ABLIKIM 18AB BES3  $J/\psi \rightarrow \eta' K^* \bar{K}$  $\Gamma_{61}/\Gamma$ NODE=M070P48  
NODE=M070P48

OCCUR=2

NODE=M070P48;LINKAGE=A  
NODE=M070P48;LINKAGE=B $\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**1.66±0.03±0.21**<sup>1</sup> From  $\eta' K_S^0 K^\pm \pi^\mp$ .

DOCUMENT ID TECN COMMENT

<sup>1</sup> ABLIKIM 18AB BES3  $J/\psi \rightarrow \eta' K^* \bar{K}$  $\Gamma_{62}/\Gamma$ NODE=M070P49  
NODE=M070P49

NODE=M070P49;LINKAGE=A

 $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**2.16±0.12±0.29**<sup>1</sup> From  $\eta' K_S^0 K^\pm \pi^\mp$ .

EVTS

1.1k

DOCUMENT ID TECN COMMENT

<sup>1</sup> ABLIKIM 18AB BES3  $J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$  $\Gamma_{63}/\Gamma$ NODE=M070P52  
NODE=M070P52

NODE=M070P52;LINKAGE=A

 $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**1.51±0.09±0.21**<sup>1</sup> From  $\eta' K^+ K^- \pi^0$ .

EVTS

1.0k

DOCUMENT ID TECN COMMENT

<sup>1</sup> ABLIKIM 18AB BES3  $J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$  $\Gamma_{64}/\Gamma$ NODE=M070P44  
NODE=M070P44

NODE=M070P44;LINKAGE=A

 $\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$ VALUE (units  $10^{-7}$ )**4.69±0.80<sup>+0.74</sup><sub>-1.82</sub>**<sup>1</sup> From a partial wave analysis of the systems  $(\gamma X)$ , with  $X \rightarrow \eta' \eta'$ , and  $(\eta' X)$ , with  $X \rightarrow \gamma \eta'$  in the decay  $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

DOCUMENT ID TECN COMMENT

<sup>1</sup> ABLIKIM 22C BES3  $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$  $\Gamma_{65}/\Gamma$ NODE=M070Q28  
NODE=M070Q28

NODE=M070Q28;LINKAGE=A

 $\Gamma(h_1(1415) \eta' \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )**0.08±0.01<sup>+0.01</sup><sub>-0.02</sub>**<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 22AS BES3  $J/\psi(1S) \rightarrow \gamma \eta \eta'$  $\Gamma_{26}/\Gamma$ NODE=M070Q33  
NODE=M070Q33

NODE=M070Q33;LINKAGE=A

 $\Gamma(h_1(1595) \eta' \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )**0.16±0.02<sup>+0.03</sup><sub>-0.01</sub>**<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 22AS BES3  $J/\psi(1S) \rightarrow \gamma \eta \eta'$  $\Gamma_{27}/\Gamma$ NODE=M070Q34  
NODE=M070Q34

NODE=M070Q34;LINKAGE=A

 $\Gamma(\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ 

VALUE (%)

**90.5±0.9±3.8**<sup>1</sup> From a Dalitz plot analysis in an isobar model.

EVTS DOCUMENT ID TECN COMMENT

<sup>1</sup> LEES 17C BABR  $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$  $\Gamma_{67}/\Gamma_{178}$ NODE=M070P30  
NODE=M070P30

NODE=M070P30;LINKAGE=A

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**6.0<sup>+0.8</sup><sub>-1.0</sub> OUR AVERAGE**

Error includes scale factor of 2.9. See the ideogram below.

8.07±0.04<sup>+0.38</sup><sub>-0.61</sub>

4.57±0.17±0.70

5.26±0.13±0.53


EVTS DOCUMENT ID TECN COMMENT

183k ABLIKIM 19AQ BES  $J/\psi \rightarrow K^+ K^- \pi^0$ 2285 JOUSSET 90 DM2  $J/\psi \rightarrow \text{hadrons}$ COFFMAN 88 MRK3  $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp,$  $K^+ K^- \pi^0$  $\Gamma_{68}/\Gamma$ NODE=M070S15  
NODE=M070S15

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83	MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77	MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76	DASP	$J/\psi \rightarrow K^\pm X$

WEIGHTED AVERAGE  
6.0+0.8-1.0 (Error scaled by 2.9)



**$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma_{total}$**   **$\Gamma_{69} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.69 ± 0.01<sup>+0.13</sup><sub>-0.20</sub></b>	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P79  
NODE=M070P79

**$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$**   **$\Gamma_{69} / \Gamma_{177}$**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>92.4 ± 1.5 ± 3.4</b>	2k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P26  
NODE=M070P26

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

NODE=M070P26;LINKAGE=A

**$\Gamma(K^0 \bar{K}^*(892)^0 + c.c.) / \Gamma_{total}$**   **$\Gamma_{71} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.2 ± 0.4 OUR AVERAGE</b>				
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
4.33 ± 0.12 ± 0.45		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

NODE=M070S16  
NODE=M070S16

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.7 ± 0.6	45	VANNUCCI	77	MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
-----------	----	----------	----	------	--

**$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + c.c.) / \Gamma_{total}$**   **$\Gamma_{73} / \Gamma$**

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.73 ± 0.14 ± 0.82</b>	<sup>1</sup> ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M070S52  
NODE=M070S52

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>2</sup> ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
------	----------------------	----------	---

<sup>1</sup> Obtained from  $J/\psi \rightarrow K^*(892) K^- \pi^+ + c.c. \rightarrow K^+ K^- \pi^+ \pi^-$  taking the value 2/3 for the probability of the  $K^*(892)^0 \rightarrow K^+ \pi^-$  decay.

NODE=M070S52;LINKAGE=A

<sup>2</sup> A  $K_0^*(700)$  is observed by ABLIKIM 06C in the  $K^+ \pi^-$  mass spectrum of the  $\bar{K}^*(892)^0 K^+ \pi^-$  final state against the  $\bar{K}^*(892)$ . A corresponding branching fraction of the  $J/\psi(1S)$  is not presented.

NODE=M070S52;LINKAGE=AB

**$\Gamma(K^*(892)^0 K^- \pi^+ + c.c. \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{total}$**   **$\Gamma_{77} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.81 ± 0.10 ± 0.54</b>	1559	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M070Q42  
NODE=M070Q42

$$\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{78} / \Gamma$$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
$6.28^{+0.16+0.59}_{-0.17-0.52}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P71  
NODE=M070P71

$$\Gamma(K^*(892)^\pm K^*(700)^\mp) / \Gamma_{\text{total}} \quad \Gamma_{80} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.09 \pm 0.18^{+0.94}_{-0.54}$	655	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M070S74  
NODE=M070S74

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{81} / \Gamma$$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M070R46  
NODE=M070R46

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5                      90              VANNUCCI    77    MRK1     $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) / \Gamma_{\text{total}} \quad \Gamma_{82} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.00 \pm 0.19^{+0.11}_{-0.32}$	323	ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M070S73  
NODE=M070S73

$$\Gamma(K_1(1400)^\pm K^\mp) / \Gamma_{\text{total}} \quad \Gamma_{83} / \Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
$3.8 \pm 0.8 \pm 1.2$	<sup>1</sup> BAI	99C BES	$e^+ e^-$

NODE=M070S35  
NODE=M070S35

<sup>1</sup> Assuming  $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

NODE=M070S35;LINKAGE=M3

$$\Gamma(K^*(1410) \bar{K} + c.c. \rightarrow K^\pm K^\mp \pi^0) / \Gamma(K^+ K^- \pi^0) \quad \Gamma_{85} / \Gamma_{177}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.3 \pm 1.1 \pm 0.7$	2k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P28  
NODE=M070P28

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

NODE=M070P28;LINKAGE=A

$$\Gamma(K^*(1410) \bar{K} + c.c. \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp) \quad \Gamma_{86} / \Gamma_{178}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.9$	4k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

NODE=M070P32  
NODE=M070P32

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

NODE=M070P32;LINKAGE=A

$$\Gamma(K_2^*(1430) \bar{K} + c.c. \rightarrow K^\pm K^\mp \pi^0) / \Gamma(K^+ K^- \pi^0) \quad \Gamma_{88} / \Gamma_{177}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$3.5 \pm 1.3 \pm 0.9$	2k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P29  
NODE=M070P29

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

NODE=M070P29;LINKAGE=A

$$\Gamma(K_2^*(1430) \bar{K} + c.c. \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp) \quad \Gamma_{89} / \Gamma_{178}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$7.1 \pm 1.3 \pm 1.2$	4k	<sup>1</sup> LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

NODE=M070P33  
NODE=M070P33

<sup>1</sup> From a Dalitz plot analysis in an isobar model.

NODE=M070P33;LINKAGE=A

$$\Gamma(\bar{K}_2^*(1430) K + c.c.) / \Gamma_{\text{total}} \quad \Gamma_{90} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 40 \times 10^{-4}$	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$

NODE=M070R45  
NODE=M070R45

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 66 \times 10^{-4}$               90              BRAUNSCH...    76    DASP     $e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$

$$\Gamma(K_2^*(1430)^+ K^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{91} / \Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.04^{+0.25}_{-0.19}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P76  
NODE=M070P76

$$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + c.c.) / \Gamma_{\text{total}} \quad \Gamma_{94} / \Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R48  
NODE=M070R48

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.7 \pm 2.6$                       40              VANNUCCI    77    MRK1     $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$



$\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$   $\Gamma_{97}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-4}$	90	VANNUCCI 77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

NODE=M070R47  
NODE=M070R47

 $\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{99}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.1^{+0.6}_{-0.1}$	183k	ABLIKIM 19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P77  
NODE=M070P77

 $\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{100}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 0.7^{+2.8}_{-1.4}$	183k	ABLIKIM 19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P78  
NODE=M070P78

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{101}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	<sup>1</sup> BAI	99C BES	$e^+ e^-$

NODE=M070S34  
NODE=M070S34

<sup>1</sup> Assuming  $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

NODE=M070S34;LINKAGE=M2

 $\Gamma(K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{102}/\Gamma$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$8.54^{+1.07+2.35}_{-1.20-2.13}$	ABLIKIM 18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P72  
NODE=M070P72

 $\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{103}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^-$

NODE=M070R42  
NODE=M070R42

 $\Gamma(\phi \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{104}/\Gamma$ 

The two different fit values of ABLIKIM 15k below have the same statistical significance of 6.4  $\sigma$  and cannot be distinguished at this moment.

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	<sup>1</sup> ABLIKIM	15k BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$
$0.124 \pm 0.033 \pm 0.030$		35 $\pm$ 9	<sup>2</sup> ABLIKIM	15k BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$

NODE=M070R33  
NODE=M070R33

NODE=M070R33

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

$<6.4$	90	<sup>3</sup> ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \gamma \gamma$
$<6.8$	90	COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

<sup>1</sup> Corresponding to one of the two fit solutions with  $\delta = (-95.9 \pm 1.5)^\circ$  for the phase angle between the resonant  $J/\psi \rightarrow \phi \pi^0$  and non-phi  $J/\psi \rightarrow K^+ K^- \pi^0$  contributions.

NODE=M070R33;LINKAGE=A

<sup>2</sup> Corresponding to one of the two fit solutions with  $\delta = (-152.1 \pm 7.7)^\circ$  for the phase angle between the resonant  $J/\psi \rightarrow \phi \pi^0$  and non-phi  $J/\psi \rightarrow K^+ K^- \pi^0$  contributions.

NODE=M070R33;LINKAGE=C

<sup>3</sup> Superseded by ABLIKIM 15k.

NODE=M070R33;LINKAGE=B

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{105}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.94 <math>\pm</math> 0.15 OUR AVERAGE</b>		Error includes scale factor of 1.7.		
$1.09 \pm 0.02 \pm 0.13$		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$0.78 \pm 0.03 \pm 0.12$		FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
$2.1 \pm 0.9$	23	FELDMAN 77	MRK1	$e^+ e^-$

NODE=M070R34  
NODE=M070R34

 $\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{107}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>16.0 <math>\pm</math> 1.0 <math>\pm</math> 3.0</b>	FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070R35  
NODE=M070R35

 $\Gamma(\phi \eta)/\Gamma_{\text{total}}$   $\Gamma_{108}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.74 <math>\pm</math> 0.06 OUR AVERAGE</b>		Error includes scale factor of 1.2. $[(0.74 \pm 0.08) \times 10^{-3}]$		
		OUR 2022 AVERAGE Scale factor = 1.5]		
$0.71 \pm 0.10 \pm 0.05$	99 $\pm$ 14	<sup>1</sup> ZHU	23 BELL	$e^+ e^- \rightarrow \Upsilon(nS) \rightarrow \phi \eta \gamma$
$0.898 \pm 0.024 \pm 0.089$		ABLIKIM 05B	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET 90	DM2	$J/\psi \rightarrow \text{hadrons}$
$0.661 \pm 0.045 \pm 0.078$		COFFMAN 88	MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$


NODE=M070R37  
NODE=M070R37

NEW

<sup>1</sup> From a fit to the combined  $\phi\eta$  invariant mass spectrum with a Gaussian function for the  $J/\psi$  signals and a second-order polynomial function for the backgrounds.

NODE=M070R37;LINKAGE=A

$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$			$\Gamma_{109}/\Gamma$		
VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.6 ± 0.5</b>	<b>OUR AVERAGE</b>		Error includes scale factor of 2.2. See the ideogram below.		
5.10 ± 0.03 ± 0.32	31k		ABLIKIM	16M	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
5.46 ± 0.31 ± 0.56			ABLIKIM	05B	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
4.1 ± 0.3 ± 0.8	167		JOUSSET	90	DM2 $J/\psi \rightarrow \text{hadrons}$
3.08 ± 0.34 ± 0.36			COFFMAN	88	MRK3 $e^+e^- \rightarrow K^+K^-\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 13	90		VANNUCCI	77	MRK1 $e^+e^-$

NODE=M070R38  
NODE=M070R38

$\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$			$\Gamma_{110}/\Gamma$		
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>2.32 ± 0.06 ± 0.16</b>	2.2k	<sup>1</sup> ABLIKIM	19AN	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

NODE=M070P85  
NODE=M070P85

<sup>1</sup> Including contributions from intermediate resonances. Evidence for an intermediate resonance at  $M \approx 2$  GeV and  $\Gamma \approx 150$  MeV decaying to  $\phi\eta'$  with  $J^P = 1^+$  or  $J^P = 1^-$ , and  $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$ .

NODE=M070P85;LINKAGE=A

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$			$\Gamma_{111}/\Gamma$		
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.2 ± 0.9</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.9.			
4.6 ± 0.4 ± 0.8		<sup>1</sup> FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$
2.6 ± 0.6	50	<sup>1</sup> GIDAL	81	MRK2	$J/\psi \rightarrow K^+K^-K^+K^-$

NODE=M070R41  
NODE=M070R41

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 0.78$ .

NODE=M070R41;LINKAGE=A

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$			$\Gamma_{114}/\Gamma$		
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>4.50 ± 0.80 ± 0.61</b>	355	ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+K^-3\pi$

NODE=M070S97  
NODE=M070S97

$\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$			$\Gamma_{115}/\Gamma$		
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.67 ± 0.50 ± 0.24</b>	70	ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+K^-3\pi$

NODE=M070S98  
NODE=M070S98

$\Gamma(\phi f_0(980)\eta \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$			$\Gamma_{116}/\Gamma$		
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.23 ± 0.75 ± 0.73</b>	52	ABLIKIM	08F	BES	$J/\psi \rightarrow \eta\phi f_0(980)$

NODE=M070R08  
NODE=M070R08

$\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0) / \Gamma_{\text{total}}$  $\Gamma_{117} / \Gamma$ 

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.37 ± 1.35</b>	1 ABLIKIM	18D BES3	$J/\psi \rightarrow \phi \eta \pi^0$
5.0 ± 2.7 ± 2.5	2 ABLIKIM	11D BES3	$J/\psi \rightarrow \phi \eta \pi^0$

NODE=M070S75  
 NODE=M070S75

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Assuming constructive interference between  $a_0(980) - f_0(980)$  mixing and electromagnetic decay. Destructive interference gives a value of  $(4.93 \pm 1.77) \times 10^{-6}$  for this branching fraction.

NODE=M070S75;LINKAGE=A

<sup>2</sup> Assuming  $a_0(980) - f_0(980)$  mixing and isospin breaking via  $\gamma^*$  and  $K^* K$  loops.

NODE=M070S75;LINKAGE=AB

 $\Gamma(\phi f_2(1270)) / \Gamma_{\text{total}}$  $\Gamma_{118} / \Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 0.45	90	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

NODE=M070R39  
 NODE=M070R39

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\phi f_1(1285)) / \Gamma_{\text{total}}$  $\Gamma_{119} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6 ± 0.5 OUR AVERAGE</b>				
3.4 ± 1.8 ± 1.5	1.1k	1 ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
3.2 ± 0.6 ± 0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
2.1 ± 0.5 ± 0.4	25	2 JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070S6  
 NODE=M070S6

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2 ± 0.1 16 BECKER 87 MRK3  $J/\psi \rightarrow \phi K \bar{K} \pi$

OCCUR=2

<sup>1</sup> ABLIKIM 15H reports  $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$  which we divide by our best value  $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S6;LINKAGE=A

<sup>2</sup> We attribute to the  $f_1(1285)$  the signal observed in the  $\pi^+ \pi^- \eta$  invariant mass distribution at 1297 MeV.

NODE=M070S6;LINKAGE=Q

 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_{120} / \Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.36 ± 2.31 ± 1.54</b>	78	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S99  
 NODE=M070S99

 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi 3\pi^0) / \Gamma_{\text{total}}$  $\Gamma_{121} / \Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.08 ± 1.63 ± 1.47</b>	9	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S00  
 NODE=M070S00

 $\Gamma(\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_{122} / \Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.01 ± 0.58 ± 0.82</b>		172	1 ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070S23  
 NODE=M070S23

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 <sup>2</sup> FALVARD 88 DM2  $J/\psi \rightarrow \text{hadrons}$

<sup>1</sup> With 3.6  $\sigma$  significance.

NODE=M070S23;LINKAGE=B

<sup>2</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow \eta \pi \pi$ .

NODE=M070S23;LINKAGE=A

 $\Gamma(\phi f'_2(1525)) / \Gamma_{\text{total}}$  $\Gamma_{123} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8 ± 4 OUR AVERAGE</b>				Error includes scale factor of 2.7.
12.3 ± 0.6 ± 2.0		<sup>1,2</sup> FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
4.8 ± 1.8	46	<sup>1</sup> GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

NODE=M070R40  
 NODE=M070R40

<sup>1</sup> Re-evaluated using  $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$ .

NODE=M070R40;LINKAGE=B

<sup>2</sup> Including interference with  $f_0(1710)$ .

NODE=M070R40;LINKAGE=C

 $\Gamma(\phi X(1835) \rightarrow \phi \rho \bar{\rho}) / \Gamma_{\text{total}}$  $\Gamma_{124} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.1 × 10<sup>-7</sup></b>	90	1 ABLIKIM	16K BES3	$J/\psi \rightarrow \rho \bar{\rho} K_S^0 K_L^0, \rho \bar{\rho} K^+ K^-$

NODE=M070P00  
 NODE=M070P00

OCCUR=2

<sup>1</sup> Upper limit applies to any  $\rho \bar{\rho}$  mass enhancement near threshold.

NODE=M070P00;LINKAGE=A

**$\Gamma(\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{125} / \Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-4}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070B10  
 NODE=M070B10

 **$\Gamma(\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$**   **$\Gamma_{126} / \Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.13 \times 10^{-5}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070B11  
 NODE=M070B11

 **$\Gamma(\phi K \bar{K}) / \Gamma_{\text{total}}$**   **$\Gamma_{127} / \Gamma$** 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R36  
 NODE=M070R36

**17.7 ± 1.6 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.


16.6 ± 1.9 ± 1.2	163 ± 19	LEES	12F BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
21.4 ± 0.4 ± 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
48 $^{+20}_{-16} \pm 6$	9.0 $^{+3.7}_{-3.0}$	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 ± 0.8 ± 2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	$e^+ e^-$

<sup>1</sup> We have multiplied  $K^+ K^-$  measurement by 2 to obtain  $K \bar{K}$ .

<sup>2</sup> Using  $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$ .

<sup>3</sup> Addition of  $\phi K^+ K^-$  and  $\phi K^0 \bar{K}^0$  branching ratios.

NODE=M070R36;LINKAGE=AA  
 NODE=M070R36;LINKAGE=CC  
 NODE=M070R36;LINKAGE=A

 **$\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K}) / \Gamma_{\text{total}}$**   **$\Gamma_{128} / \Gamma$** 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.6 ± 0.2 ± 0.6</b>	1,2 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S24  
 NODE=M070S24

<sup>1</sup> Including interference with  $f_2'(1525)$ .

<sup>2</sup> Includes unknown branching fraction  $f_0(1710) \rightarrow K \bar{K}$ .

NODE=M070S24;LINKAGE=D  
 NODE=M070S24;LINKAGE=E

 **$\Gamma(\phi K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$**   **$\Gamma_{131} / \Gamma$** 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.2 ± 0.8 OUR AVERAGE</b>				
7.4 ± 0.6 ± 1.4	227 ± 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
7.4 ± 0.9 ± 1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 ± 0.6 ± 1.0	163 ± 15	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S3  
 NODE=M070S3

 **$\Gamma(\phi K^*(892) \bar{K} + \text{c.c.}) / \Gamma_{\text{total}}$**   **$\Gamma_{132} / \Gamma$** 


VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.8 ± 2.3 OUR AVERAGE</b>				
20.8 ± 2.7 ± 3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 ± 3.7 ± 4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7 ± 2.4 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S4  
 NODE=M070S4

OCCUR=2



$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$					$\Gamma_{146}/\Gamma$	
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>1.17 ± 0.04 OUR AVERAGE</b>						NODE=M070S64 NODE=M070S64
1.165 ± 0.004 ± 0.043	135k	ABLIKIM	17E BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$		
1.20 ± 0.12 ± 0.21	206	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$ hadrons		
$\Gamma(\Xi(1530)^- \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{147}/\Gamma$	
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>0.318 ± 0.008 OUR AVERAGE</b>						NODE=M070S9 NODE=M070S9
0.317 ± 0.002 ± 0.008	70k	ABLIKIM	20 BES3	$e^+ e^- \rightarrow J/\psi$		
0.59 ± 0.09 ± 0.12	75	HENRARD	87 DM2	$e^+ e^-$		
$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$					$\Gamma_{148}/\Gamma$	
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>0.32 ± 0.12 ± 0.07</b>	24 ± 9	HENRARD	87 DM2	$e^+ e^-$		NODE=M070S10 NODE=M070S10
$\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 \rho K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{149}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;1.1 × 10<sup>-5</sup></b>	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S47 NODE=M070S47
$\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$					$\Gamma_{150}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;2.1 × 10<sup>-5</sup></b>	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S48 NODE=M070S48
$\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					$\Gamma_{151}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;1.6 × 10<sup>-5</sup></b>	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S49 NODE=M070S49
$\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					$\Gamma_{152}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;5.6 × 10<sup>-5</sup></b>	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S50 NODE=M070S50
$\Gamma(\bar{\Theta}(1540) K_S^0 \rho \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$					$\Gamma_{153}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;1.1 × 10<sup>-5</sup></b>	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S51 NODE=M070S51
———— STABLE HADRONS ————						
$\Gamma(2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$					$\Gamma_{154}/\Gamma$	
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>4.2 ± 0.4 OUR AVERAGE</b>						NODE=M070R9 NODE=M070R9
Error includes scale factor of 2.1. See the ideogram below.						
[(3.71 ± 0.28) × 10 <sup>-2</sup> OUR 2022 AVERAGE Scale factor = 1.3]						
5.44 ± 0.07 ± 0.33	23K	ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$		
3.5 ± 0.8 ± 0.1	14k	<sup>1</sup> LEES	21 BABR	$10.6 e^+ e^- \rightarrow$ $2(\pi^+ \pi^-) 3\pi^0 \gamma$		
4.73 ± 0.44	228k	<sup>2</sup> ABLIKIM	19Q BES3	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$		
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$		
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons		
3.64 ± 0.52	1500	BURMESTER	77D PLUT	$e^+ e^-$		
4 ± 1	675	JEAN-MARIE	76 MRK1	$e^+ e^-$		
<sup>1</sup> LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times$ $[B(\psi(2S) \rightarrow J/\psi(1S) \pi^0 \pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3}$ keV which we divide by our best values $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV, $B(\psi(2S) \rightarrow J/\psi(1S) \pi^0 \pi^0) =$ $(18.24 \pm 0.31) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.						NODE=M070R9;LINKAGE=B
<sup>2</sup> From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$ , assuming PDG 16 values for $\Gamma(e^+ e^-)$ , $\Gamma(\mu^+ \mu^-)$ , and $\Gamma(\text{total})$ , and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$ . An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$ .						NODE=M070R9;LINKAGE=A



**$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{total}$**   **$\Gamma_{155}/\Gamma$**

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.029±0.006 OUR AVERAGE</b>				
0.028±0.009	11	FRANKLIN 83	MRK2	$e^+e^- \rightarrow$ hadrons
0.029±0.007	181	JEAN-MARIE 76	MRK1	$e^+e^-$

NODE=M070R11  
NODE=M070R11

**$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$**   **$\Gamma_{160}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>21.0 ±0.8 OUR AVERAGE</b>		Error includes scale factor of 1.6. See the ideogram below.		
21.37±0.04 <sup>+0.64</sup> <sub>-0.62</sub>	1.8M	1,2 ABLIKIM 12H	BES3	$e^+e^- \rightarrow J/\psi$
23.0 ±2.0 ±0.4	256	3 AUBERT 07AU	BABR 10.6	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
21.84±0.05±2.01	220k	1,4 BAI 04H	BES	$e^+e^-$
20.91±0.21±1.16		4,5 BAI 04H	BES	$e^+e^-$
15 ±2	168	FRANKLIN 83	MRK2	$e^+e^-$

NODE=M070R7  
NODE=M070R7

OCCUR=2

<sup>1</sup> From  $J/\psi \rightarrow \pi^+\pi^-\pi^0$  events directly.

<sup>2</sup> The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of  $J/\psi$  events.

<sup>3</sup> AUBERT 07AU reports  $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total} = 0.808 \pm 0.013$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.


<sup>4</sup> Mostly  $\rho\pi$ , see also  $\rho\pi$  subsection.

<sup>5</sup> Obtained comparing the rates for  $\pi^+\pi^-\pi^0$  and  $\mu^+\mu^-$ , using  $J/\psi$  events produced via  $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$  and with  $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$ .

NODE=M070R;LINKAGE=BA  
NODE=M070R7;LINKAGE=AB

NODE=M070R7;LINKAGE=AU

NODE=M070R;LINKAGE=BU  
NODE=M070R;LINKAGE=BI



**$\Gamma(\pi^+\pi^-\pi^0 K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{162}/\Gamma$**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.52±0.27 OUR AVERAGE</b>		Error includes scale factor of 1.4.		[(1.2 ± 0.3) × 10 <sup>-2</sup> OUR 2022 AVERAGE]	NEW
1.74±0.08±0.24	2616	ANASHIN	22	KEDR $J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$	
1.2 ±0.3	309	VANNUCCI	77	MRK1 $e^+e^-$	

**$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{163}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>1.47±0.14 OUR AVERAGE</b>					
1.47±0.13±0.13	140	<sup>1</sup> METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+\pi^-)$	
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	$e^+e^-$	
1.0 ±0.5	5	BRANDELIK	78B	DASP $e^+e^-$	
1.6 ±1.6	1	VANNUCCI	77	MRK1 $e^+e^-$	

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

**$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{164}/\Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>3.20±0.25 OUR AVERAGE</b>		Error includes scale factor of 1.2.		[(3.57 ± 0.30) × 10 <sup>-3</sup> OUR 2022 AVERAGE]	NEW
2.88±0.14±0.24	2654	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+\pi^-)$	
3.53±0.12±0.29	1107	<sup>1</sup> ABLIKIM	05H	BES2 $e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+\pi^-, J/\psi \rightarrow 2(\pi^+\pi^-)$	
4.0 ±1.0	76	JEAN-MARIE	76	MRK1 $e^+e^-$	

<sup>1</sup> Computed using  $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ .

**$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$   $\Gamma_{165}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
40±20	32	JEAN-MARIE	76	MRK1 $e^+e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

**$\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{167}/\Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>90±30</b>	13	JEAN-MARIE	76	MRK1 $e^+e^-$	



**$\Gamma(2(\pi^+\pi^-\eta))/\Gamma_{total}$**   **$\Gamma_{168}/\Gamma$**

VALUE (units  $10^{-3}$ ) EVTS DOCUMENT ID TECN COMMENT

**2.29±0.28 OUR AVERAGE**

3.1 ±1.5 ±0.1	14k	<sup>1</sup> LEES	21	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
2.26±0.08±0.27	4.8k	ABLIKIM	05c	BES2	$e^+e^- \rightarrow 2(\pi^+\pi^-)\eta$

<sup>1</sup> LEES 21 reports  $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta)/\Gamma_{total}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$  keV which we divide by our best values  $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$  keV,  $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070S42  
NODE=M070S42

NODE=M070S42;LINKAGE=A

**$\Gamma(3(\pi^+\pi^-)\eta)/\Gamma_{total}$**   **$\Gamma_{169}/\Gamma$**

VALUE (units  $10^{-4}$ ) EVTS DOCUMENT ID TECN COMMENT

<b>7.24±0.96±1.11</b>	616	ABLIKIM	05c	BES2	$e^+e^- \rightarrow 3(\pi^+\pi^-)\eta$
-----------------------	-----	---------	-----	------	--

NODE=M070S43  
NODE=M070S43

**$\Gamma(K^+K^-)/\Gamma_{total}$**   **$\Gamma_{173}/\Gamma$**

VALUE (units  $10^{-4}$ ) EVTS DOCUMENT ID TECN COMMENT

<b>2.86±0.09±0.19</b>	1k	<sup>1</sup> METREVELI	12		$\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$
-----------------------	----	------------------------	----	--	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.39±0.24±0.22	107	<sup>2</sup> BALTRUSAIT..85D	MRK3		$e^+e^-$
2.2 ±0.9	6	<sup>2</sup> BRANDELIK	79c	DASP	$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.  
<sup>2</sup> Interference with non-resonant  $K^+K^-$  production not taken into account.

NODE=M070R13  
NODE=M070R13

NODE=M070R13;LINKAGE=ME  
NODE=M070R13;LINKAGE=BA

**$\Gamma(K_S^0K_L^0)/\Gamma_{total}$**   **$\Gamma_{174}/\Gamma$**

VALUE (units  $10^{-4}$ ) EVTS DOCUMENT ID TECN COMMENT

**1.95±0.11 OUR AVERAGE** Error includes scale factor of 2.4. See the ideogram below.

1.93±0.01±0.05	110k	ABLIKIM	17AH	BES3	$J/\psi \rightarrow K_S^0K_L^0 \rightarrow \pi^+\pi^-X$
2.62±0.15±0.14	0.3k	<sup>1</sup> METREVELI	12		$\psi(2S) \rightarrow \pi^+\pi^-K_S^0K_L^0$
1.82±0.04±0.13	2.1k	<sup>2</sup> BAI	04A	BES2	$J/\psi \rightarrow K_S^0K_L^0 \rightarrow \pi^+\pi^-X$


• • • We do not use the following data for averages, fits, limits, etc. • • •

1.18±0.12±0.18		JOUSSET	90	DM2	$J/\psi \rightarrow$ hadrons
1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3		$e^+e^-$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.  
<sup>2</sup> Using  $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$ .

NODE=M070R75  
NODE=M070R75

NODE=M070R75;LINKAGE=ME  
NODE=M070R;LINKAGE=HZ



$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{175}/\Gamma$
$<1.4 \times 10^{-8}$	95	<sup>1</sup> ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$	

NODE=M070R14  
NODE=M070R14

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1 \times 10^{-6}$	95	<sup>1</sup> BAI	04D BES	$e^+ e^-$	
$<5.2 \times 10^{-6}$	90	<sup>1</sup> BALTRUSAIT..85C	MRK3	$e^+ e^-$	

<sup>1</sup> Forbidden by CP.

NODE=M070R14;LINKAGE=C

 $\Gamma(K \bar{K} \pi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{176}/\Gamma$
<b>61 ± 10 OUR AVERAGE</b>					
55.2 ± 12.0	25	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow K^+ K^- \pi^0$	
78.0 ± 21.0	126	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$	

NODE=M070R15  
NODE=M070R15

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{177}/\Gamma$
<b>2.88 ± 0.01 ± 0.12</b>	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$	

NODE=M070P80  
NODE=M070P80

 $\Gamma(K^+ K^- \pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{177}/\Gamma_{160}$
<b>12.0 ± 0.3 ± 0.9</b>	23k	LEES	17C BABR	$J/\psi \rightarrow h^+ h^- \pi^0$	

NODE=M070P34  
NODE=M070P34

 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+ \pi^- \pi^0)$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{178}/\Gamma_{160}$
<b>26.5 ± 0.5 ± 2.1</b>	24k	LEES	17C BABR	$J/\psi \rightarrow h^0 h^+ h^-$	

NODE=M070P35  
NODE=M070P35

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{182}/\Gamma$
<b>7.0 ± 1.0 OUR AVERAGE</b>				[(6.6 ± 0.5) × 10 <sup>-3</sup> OUR 2016 AVERAGE]	

NODE=M070R16  
NODE=M070R16

**7.04 ± 0.26 ± 0.92** 2671 ANASHIN 22 KEDR  $J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 2.3	205	VANNUCCI	77 MRK1	$e^+ e^-$	
-----------	-----	----------	---------	-----------	--

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{189}/\Gamma$
<b>31 ± 13</b>	30	VANNUCCI	77 MRK1	$e^+ e^-$	

NODE=M070R17  
NODE=M070R17

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{191}/\Gamma$
--------------------------	------	-------------	------	---------	-----------------------

NODE=M070R19  
NODE=M070R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.4^{+0.5}_{-0.4} \pm 0.2$	$11.0^{+4.3}_{-3.5}$	<sup>1</sup> HUANG	03 BELL	$B^+ \rightarrow 2(K^+ K^-) K^+$	
0.7 ± 0.3		VANNUCCI	77 MRK1	$e^+ e^-$	

<sup>1</sup> Using  $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$ .

NODE=M070R19;LINKAGE=CC

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{193}/\Gamma$
--------------------------	------	-------------	------	---------	-----------------------

NODE=M070R50  
NODE=M070R50

**2.120 ± 0.029 OUR AVERAGE**

2.112 ± 0.004 ± 0.031	314k	ABLIKIM	12C BES3	$e^+ e^-$	
2.17 ± 0.16 ± 0.04	317	<sup>1</sup> WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$	
2.26 ± 0.01 ± 0.14	63316	BAI	04E BES2	$e^+ e^- \rightarrow J/\psi$	
1.97 ± 0.22	99	BALDINI	98 FENI	$e^+ e^-$	
1.91 ± 0.04 ± 0.30		PALLIN	87 DM2	$e^+ e^-$	
2.16 ± 0.07 ± 0.15	1420	EATON	84 MRK2	$e^+ e^-$	
2.5 ± 0.4	133	BRANDELIK	79C DASP	$e^+ e^-$	
2.0 ± 0.5		BESCH	78 BONA	$e^+ e^-$	
2.2 ± 0.2	331	<sup>2</sup> PERUZZI	78 MRK1	$e^+ e^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3	48	ANTONELLI	93 SPEC	$e^+ e^-$	
-----------	----	-----------	---------	-----------	--

<sup>1</sup> WU 06 reports  $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R50;LINKAGE=WU

<sup>2</sup> Assuming angular distribution  $(1 + \cos^2\theta)$ .


NODE=M070R50;LINKAGE=A

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{194}/\Gamma$ NODE=M070R52  
NODE=M070R52

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.19±0.08 OUR AVERAGE</b>				Error includes scale factor of 1.1.
1.33±0.02±0.11	11k	ABLIKIM	09B BES2	$e^+e^-$
1.13±0.09±0.09	685	EATON	84 MRK2	$e^+e^-$
1.4 ±0.4		BRANDELIK	79C DASP	$e^+e^-$
1.00±0.15	109	PERUZZI	78 MRK1	$e^+e^-$

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{195}/\Gamma$ NODE=M070R54  
NODE=M070R54

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0 ±0.5 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
6.46±0.17±0.43	1435	EATON	84 MRK2	$e^+e^-$
3.8 ±1.6	48	BESCH	81 BONA	$e^+e^-$
5.5 ±0.6	533	PERUZZI	78 MRK1	$e^+e^-$

WEIGHTED AVERAGE  
6.0±0.5 (Error scaled by 1.3) $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{196}/\Gamma$ NODE=M070R55  
NODE=M070R55  
NODE=M070R55Including  $p\bar{p}\pi^+\pi^-\gamma$  and excluding  $\omega, \eta, \eta'$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ±0.9 OUR AVERAGE</b>				Error includes scale factor of 1.9.
3.36±0.65±0.28	364	EATON	84 MRK2	$e^+e^-$
1.6 ±0.6	39	PERUZZI	78 MRK1	$e^+e^-$

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$  $\Gamma_{197}/\Gamma$ NODE=M070R56  
NODE=M070R56

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.00±0.12 OUR AVERAGE</b>				
1.91±0.02±0.17	13k	<sup>1</sup> ABLIKIM	09 BES2	$e^+e^-$
2.03±0.13±0.15	826	EATON	84 MRK2	$e^+e^-$
2.5 ±1.2		BRANDELIK	79C DASP	$e^+e^-$
2.3 ±0.4	197	PERUZZI	78 MRK1	$e^+e^-$

<sup>1</sup> From the combination of  $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$  and  $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$  channels.


NODE=M070R56;LINKAGE=AB

 $\Gamma(p\bar{p}\rho)/\Gamma_{\text{total}}$  $\Gamma_{198}/\Gamma$ NODE=M070R57  
NODE=M070R57

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.31 × 10<sup>-3</sup></b>	90	EATON	84 MRK2	$e^+e^- \rightarrow \text{hadrons}\gamma$

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$  $\Gamma_{199}/\Gamma$ NODE=M070R58  
NODE=M070R58

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.98±0.10 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
0.90±0.02±0.09	2670	ABLIKIM	13P BES3	$e^+e^-$
0.98±0.03±0.14	2449	ABLIKIM	08 BES2	$e^+e^-$
1.10±0.17±0.18	486	EATON	84 MRK2	$e^+e^-$
1.6 ±0.3	77	PERUZZI	78 MRK1	$e^+e^-$



### $\Gamma(\rho\bar{\pi}^0(958))/\Gamma_{\text{total}}$

$\Gamma_{200}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.129 \pm 0.014</math> OUR AVERAGE</b>				Error includes scale factor of 2.0.
$0.126 \pm 0.002 \pm 0.007$	16k	<sup>1</sup> ABLIKIM	19N BES3	$e^+e^-$
$0.200 \pm 0.023 \pm 0.028$	$265 \pm 31$	<sup>2</sup> ABLIKIM	09 BES2	$e^+e^-$
$0.68 \pm 0.23 \pm 0.17$	19	EATON	84 MRK2	$e^+e^-$
$1.8 \pm 0.6$	19	PERUZZI	78 MRK1	$e^+e^-$

NODE=M070R59  
 NODE=M070R59

<sup>1</sup> From the combination of  $\rho\bar{\pi}^0 \rightarrow \rho\bar{\pi}^+\pi^-\eta$  and  $\rho\bar{\pi}^0 \rightarrow \rho\bar{\pi}^+\pi^-\gamma$  channels.

<sup>2</sup> From the combination of  $\rho\bar{\pi}^0 \rightarrow \rho\bar{\pi}^+\pi^-\eta$  and  $\rho\bar{\pi}^0 \rightarrow \rho\bar{\pi}^+\pi^-\rho^0$  channels.

NODE=M070R59;LINKAGE=A  
 NODE=M070R59;LINKAGE=AB

### $\Gamma(\rho\bar{\pi}^0(980) \rightarrow \rho\bar{\pi}^0\eta)/\Gamma_{\text{total}}$

$\Gamma_{201}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>6.8 \pm 1.2 \pm 1.3</math></b>	ABLIKIM	14N BES3	$e^+e^- \rightarrow J/\psi$

NODE=M070S94  
 NODE=M070S94

### $\Gamma(\rho\bar{\pi}^0\phi)/\Gamma_{\text{total}}$

$\Gamma_{202}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.519 \pm 0.033</math> OUR AVERAGE</b>				
$0.523 \pm 0.006 \pm 0.033$	14k	ABLIKIM	16K BES3	$J/\psi \rightarrow \rho\bar{\pi}^0 K_S^0 K_L^0,$ $\rho\bar{\pi}^0 K^+ K^-$
$0.45 \pm 0.13 \pm 0.07$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S22  
 NODE=M070S22

### $\Gamma(\rho\bar{\pi}^-\pi^-)/\Gamma_{\text{total}}$

$\Gamma_{203}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.12 \pm 0.09</math> OUR AVERAGE</b>				
$2.36 \pm 0.02 \pm 0.21$	59k	ABLIKIM	06K BES2	$J/\psi \rightarrow \rho\pi^-\bar{\pi}$
$2.47 \pm 0.02 \pm 0.24$	55k	ABLIKIM	06K BES2	$J/\psi \rightarrow \bar{\rho}\pi^+n$
$2.02 \pm 0.07 \pm 0.16$	1288	EATON	84 MRK2	$e^+e^- \rightarrow \rho\pi^-$
$1.93 \pm 0.07 \pm 0.16$	1191	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\rho}\pi^+$
$1.7 \pm 0.7$	32	BESCH	81 BONA	$e^+e^- \rightarrow \rho\pi^-$
$1.6 \pm 1.2$	5	BESCH	81 BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$
$2.16 \pm 0.29$	194	PERUZZI	78 MRK1	$e^+e^- \rightarrow \rho\pi^-$
$2.04 \pm 0.27$	204	PERUZZI	78 MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$

NODE=M070R53  
 NODE=M070R53

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

### $\Gamma(n\bar{n})/\Gamma_{\text{total}}$

$\Gamma_{204}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.09 \pm 0.16</math> OUR AVERAGE</b>				
$2.07 \pm 0.01 \pm 0.17$	36k	ABLIKIM	12C BES3	$e^+e^-$
$2.31 \pm 0.49$	79	BALDINI	98 FENI	$e^+e^-$
$1.8 \pm 0.9$		BESCH	78 BONA	$e^+e^-$
$1.90 \pm 0.55$	40	ANTONELLI	93 SPEC	$e^+e^-$

NODE=M070R64  
 NODE=M070R64

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{total}$   $\Gamma_{205}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8±3.6</b>	5	BESCH	81	BONA $e^+e^-$

NODE=M070R65  
NODE=M070R65


$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{total}$   $\Gamma_{209}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.89 ±0.09 OUR AVERAGE</b>		Error includes scale factor of 2.8. See the ideogram below.		
1.943±0.003±0.033	441k	ABLIKIM	17L	BES3 $e^+e^-$
2.03 ±0.03 ±0.15	8887	ABLIKIM	06	BES2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$
1.96 <sup>+0.47</sup> <sub>-0.44</sub> ±0.04	46	<sup>1</sup> WU	06	BELL $B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ±0.06 ±0.24	631	BAI	98G	BES $e^+e^-$
1.38 ±0.05 ±0.20	1847	PALLIN	87	DM2 $e^+e^-$
1.58 ±0.08 ±0.19	365	EATON	84	MRK2 $e^+e^-$
2.6 ±1.6	5	BESCH	81	BONA $e^+e^-$
1.1 ±0.2	196	PERUZZI	78	MRK1 $e^+e^-$

NODE=M070R60  
NODE=M070R60

<sup>1</sup>WU 06 reports  $[\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{total}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.00^{+0.34}_{-0.29} \pm 0.34) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R60;LINKAGE=WU



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{total}$   $\Gamma_{210}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.78±0.27±0.30</b>		323	<sup>1</sup> ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

NODE=M070S11  
NODE=M070S11

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.4	90	<sup>2</sup> ABLIKIM	07H	BES2	$e^+e^- \rightarrow \psi(2S)$
23 ±7 ±8	11	BAI	98G	BES	$e^+e^-$
22 ±5 ±5	19	HENRARD	87	DM2	$e^+e^-$

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ .

NODE=M070S11;LINKAGE=AL  
NODE=M070S11;LINKAGE=AB

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{total}$   $\Gamma_{211}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.30±0.13±0.99</b>	2.4k	ABLIKIM	12P	BES2 $J/\psi$

NODE=M070S78  
NODE=M070S78

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{total}$   $\Gamma_{212}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.2±1.7 OUR AVERAGE</b>				
15.7±0.80±1.54	454	<sup>1</sup> ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2±6.0 ±4.4	44	<sup>2</sup> ABLIKIM	07H	BES2 $e^+e^- \rightarrow \psi(2S)$

NODE=M070R07  
NODE=M070R07

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.31\%$ .

<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.4\%$ .

NODE=M070R07;LINKAGE=AL  
NODE=M070R07;LINKAGE=AB

$\Gamma(\Lambda\bar{\Sigma}^-\pi^+(\text{or c.c.}))/\Gamma_{\text{total}}$  $\Gamma_{213}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.83 ± 0.07 OUR AVERAGE</b>				Error includes scale factor of 1.2.
0.770 ± 0.051 ± 0.083	335	<sup>1</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
0.747 ± 0.056 ± 0.076	254	<sup>1</sup> ABLIKIM	07H BES2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
0.90 ± 0.06 ± 0.16	225 ± 15	HENRARD	87 DM2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
1.11 ± 0.06 ± 0.20	342 ± 18	HENRARD	87 DM2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$
1.53 ± 0.17 ± 0.38	135	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\Lambda}\Sigma^+\pi^-$
1.38 ± 0.21 ± 0.35	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\bar{\Sigma}^-\pi^+$

NODE=M070R71  
NODE=M070R71

OCCUR=2

OCCUR=2

OCCUR=2

<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$ .

NODE=M070R71;LINKAGE=AB

 $\Gamma(pK^-\bar{\Lambda}+\text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{214}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.86 ± 0.11 OUR AVERAGE</b>				
0.84 <sup>+0.17</sup> <sub>-0.15</sub> ± 0.02	45	<sup>1</sup> LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
0.89 ± 0.07 ± 0.14	307	EATON	84 MRK2	$e^+e^-$

NODE=M070R72  
NODE=M070R72

<sup>1</sup> LU 19 reports  $(8.32_{-1.45}^{+1.63} \pm 0.49) \times 10^{-4}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda}+\text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$  assuming  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$ , which we rescale to our best value  $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R72;LINKAGE=A

 $\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{215}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.29 ± 0.06 ± 0.05</b>	90	EATON	84 MRK2	$e^+e^-$

NODE=M070R73  
NODE=M070R73 $\Gamma(\bar{\Lambda}nK_S^0+\text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{216}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.46 ± 0.20 ± 1.07</b>	1058	<sup>1</sup> ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070S56  
NODE=M070S56<sup>1</sup> Using  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$ .

NODE=M070S56;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Sigma}^+\text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{217}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.83 ± 0.23 OUR AVERAGE</b>					
2.74 ± 0.24 ± 0.22		234 ± 21	<sup>1</sup> ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92 ± 0.22 ± 0.24		308 ± 24	<sup>2</sup> ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

NODE=M070R61  
NODE=M070R61

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18			<sup>2</sup> HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15	90		PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$

<sup>1</sup> ABLIKIM 12B quotes  $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$  which we multiply by 2.<sup>2</sup> ABLIKIM 12B and HENRARD 87 quote results for  $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$  which we multiply by 2.

NODE=M070R61;LINKAGE=AB

NODE=M070R61;LINKAGE=AC

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$  $\Gamma_{218}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.07 ± 0.04 OUR AVERAGE</b>				
1.061 ± 0.004 ± 0.036	87k	ABLIKIM	21AT BES3	$J/\psi \rightarrow p\pi^0\bar{p}\pi^0$
1.50 ± 0.10 ± 0.22	399	ABLIKIM	08O BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070S09  
NODE=M070S09 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{219}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.172 ± 0.032 OUR AVERAGE</b>				Error includes scale factor of 1.4.
1.164 ± 0.004 ± 0.023	111k	ABLIKIM	17L BES3	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.33 ± 0.04 ± 0.11	1.7k	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0\bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884	PALLIN	87 DM2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$

NODE=M070R63  
NODE=M070R63

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.4 ± 2.6	3	BESCH	81 BONA	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$
-----------	---	-------	---------	---

 $\Gamma(\Sigma^+\bar{\Sigma}^-\eta)/\Gamma_{\text{total}}$  $\Gamma_{220}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.34 ± 0.21 ± 0.37</b>	1821	ABLIKIM	22AY BES3	$J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-\eta$

NODE=M070Q45  
NODE=M070Q45


$\Gamma(\Xi^- \Xi^+)/\Gamma_{total}$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.97 ± 0.08 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
1.040 ± 0.006 ± 0.074	43k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Xi^- \Xi^+$
0.90 ± 0.03 ± 0.18	961	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^- \Xi^+$
0.70 ± 0.06 ± 0.12	132	HENRARD	87 DM2	$e^+ e^- \rightarrow \Xi^- \Xi^+$
1.14 ± 0.08 ± 0.20	194	EATON	84 MRK2	$e^+ e^- \rightarrow \Xi^- \Xi^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Xi^- \Xi^+$

$\Gamma_{221}/\Gamma$

NODE=M070R62  
NODE=M070R62

WEIGHTED AVERAGE  
0.97±0.08 (Error scaled by 1.4)



**RADIATIVE DECAYS**

$\Gamma(\gamma\eta_c(1S))/\Gamma_{total}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.7 ± 0.4 OUR AVERAGE</b>		Error includes scale factor of 1.5.		
2.00 ± 0.31 ± 0.02		<sup>1</sup> MITCHELL	09 CLEO	$e^+ e^- \rightarrow \gamma X$
1.27 ± 0.36		GAISER	86 CBAL	$J/\psi \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
seen		ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
0.79 ± 0.20	273 ± 43	<sup>2</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
seen	16	BALTRUSAITIS..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$

$\Gamma_{222}/\Gamma$

NODE=M070310

NODE=M070R85  
NODE=M070R85

<sup>1</sup> MITCHELL 09 reports  $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{total}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Calculated by the authors using an average of  $B(J/\psi \rightarrow \gamma\eta_c) \times B(\eta_c \rightarrow K\bar{K}\pi)$  from BALTRUSAITIS 86, BISELLO 91, BAI 04 and  $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$  from AUBERT 06E.

NODE=M070R85;LINKAGE=MI

NODE=M070R85;LINKAGE=AU

$\Gamma(\gamma\eta_c(1S) \rightarrow 3\gamma)/\Gamma_{total}$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8<sup>+1.3</sup><sub>-1.0</sub> OUR AVERAGE</b>		Error includes scale factor of 1.1.		
4.5 ± 1.2 ± 0.6	33 ± 9	ABLIKIM	13I BES3	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
1.2 <sup>+2.7</sup> <sub>-1.1</sub> ± 0.3	1.2 <sup>+2.8</sup> <sub>-1.1</sub>	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

$\Gamma_{223}/\Gamma$

NODE=M070S08  
NODE=M070S08

$\Gamma(\gamma\eta_c(1S) \rightarrow \gamma\eta\eta')/\Gamma_{total}$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.86 ± 0.62 ± 0.45</b>	137	ABLIKIM	21C BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta'$

$\Gamma_{224}/\Gamma$

NODE=M070P89  
NODE=M070P89

$\Gamma(3\gamma)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.6±2.2 OUR AVERAGE</b>					
11.3±1.8±2.0		113 ± 18	ABLIKIM	13I	BES3 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ±3 ±2		24.2 <sup>+7.2</sup> <sub>-6.0</sub>	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 $\Gamma_{225}/\Gamma$ 

NODE=M070R81  
NODE=M070R81

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55                      90                      PARTRIDGE    80    CBAL     $e^+e^-$

 $\Gamma(4\gamma)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;9 × 10<sup>-6</sup></b>	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 $\Gamma_{226}/\Gamma$ 

NODE=M070S06  
NODE=M070S06

 $\Gamma(5\gamma)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;15 × 10<sup>-6</sup></b>	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 $\Gamma_{227}/\Gamma$ 

NODE=M070S07  
NODE=M070S07

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.56±0.17 OUR AVERAGE</b>				
3.59±0.20±0.03	1.6k	<sup>1</sup> ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
3.63±0.36±0.13		PEDLAR	09	CLE3 $J/\psi \rightarrow \pi^0\gamma$
3.13 <sup>+0.65</sup> <sub>-0.47</sub>	586	ABLIKIM	06E	BES2 $J/\psi \rightarrow \pi^0\gamma$

 $\Gamma_{228}/\Gamma$ 

NODE=M070R82  
NODE=M070R82

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.6 ±1.1 ±0.7                      BLOOM                      83    CBAL     $e^+e^-$   
7.3 ±4.7                      10    BRANDELIK                      79C    DASP     $e^+e^-$

<sup>1</sup> ABLIKIM 180 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] = (3.57 \pm 0.12 \pm 0.16) \times 10^{-5}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}] \times [B(\pi^0 \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$ , which we rescale to our best values  $B(\pi^0 \rightarrow 2\gamma) = (98.823 \pm 0.034) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R82;LINKAGE=A

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.15±0.05</b>	<sup>1</sup> ABLIKIM	15AE	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$

 $\Gamma_{229}/\Gamma$ 

NODE=M070B00  
NODE=M070B00

<sup>1</sup> The uncertainty is systematic as statistical is negligible.

NODE=M070B00;LINKAGE=A

 $\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.8 ±0.5 OUR AVERAGE</b> Error includes scale factor of 1.9. See the ideogram below.			
4.32±0.14±0.73	<sup>1</sup> BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08±0.13±0.35	<sup>2</sup> BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
3.05±0.08±0.45	<sup>2</sup> BALTRUSAIT..	.86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$
4.85±0.45±1.20	<sup>3</sup> BURKE	82	MRK2 $e^+e^-$

 $\Gamma_{230}/\Gamma$ 


NODE=M070R95  
NODE=M070R95

<sup>1</sup>  $4\pi$  mass less than 3.0 GeV.  
<sup>2</sup>  $4\pi$  mass less than 2.0 GeV.  
<sup>3</sup>  $4\pi$  mass less than 2.5 GeV.

OCCUR=2

NODE=M070R95;LINKAGE=A  
NODE=M070R95;LINKAGE=B  
NODE=M070R95;LINKAGE=M





$\Gamma(\gamma f_2(1270) f_2(1270)) / \Gamma_{\text{total}}$   $\Gamma_{231} / \Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.5±0.7±1.6</b>	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070S45  
NODE=M070S45

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant})) / \Gamma_{\text{total}}$   $\Gamma_{232} / \Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.2±0.8±1.7</b>	<sup>1</sup> ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070S46  
NODE=M070S46

<sup>1</sup> Subtracting contribution from intermediate  $\eta_c(1S)$  decays.

NODE=M070S46;LINKAGE=AB

$\Gamma(\gamma \pi^+ \pi^- 2\pi^0) / \Gamma_{\text{total}}$   $\Gamma_{233} / \Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.3±0.2±3.1</b>	<sup>1</sup> BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

NODE=M070R99  
NODE=M070R99

<sup>1</sup>  $4\pi$  mass less than 2.0 GeV.

NODE=M070R99;LINKAGE=M

$\Gamma(\gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$   $\Gamma_{234} / \Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.1±0.4</b>	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P73  
NODE=M070P73

$\Gamma(\gamma (K \bar{K} \pi) [J^{PC} = 0^{-+}]) / \Gamma_{\text{total}}$   $\Gamma_{235} / \Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.7 ± 0.4 OUR AVERAGE</b>			Error includes scale factor of 2.1.
0.58 ± 0.03 ± 0.20	<sup>1</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ± 0.1 ± 0.7	<sup>2</sup> BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

NODE=M070S38  
NODE=M070S38

<sup>1</sup> For a broad structure around 1800 MeV.

<sup>2</sup> For a broad structure around 2040 MeV.

OCCUR=2

NODE=M070S38;LINKAGE=BD  
NODE=M070S38;LINKAGE=BE

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{236} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1±0.1±0.6</b>	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B05  
NODE=M070B05

$\Gamma(\gamma K^*(892) \bar{K}^*(892)) / \Gamma_{\text{total}}$   $\Gamma_{237} / \Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.0±0.3±1.3</b>	320	<sup>1</sup> BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B07  
NODE=M070B07

<sup>1</sup> Summed over all charges.

NODE=M070R;LINKAGE=B7

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ ) EVTS**1.085 ± 0.018 OUR AVERAGE**

1.067 ± 0.005 ± 0.023 87.9k

1.12 ± 0.05 ± 0.01 18.6k

1.101 ± 0.029 ± 0.022

1.123 ± 0.089 11k

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.88 ± 0.08 ± 0.11

0.82 ± 0.10

1.3 ± 0.4

DOCUMENT ID

TECN

COMMENT

ABLIKIM 21AMBES3  $e^+e^- \rightarrow J/\psi$ <sup>1</sup> ABLIKIM 18O BES3  $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$ PEDLAR 09 CLE3  $J/\psi \rightarrow \eta\gamma$ ABLIKIM 06E BES2  $J/\psi \rightarrow \eta\gamma$ BLOOM 83 CBAL  $e^+e^-$ BRANDELIK 79C DASP  $e^+e^-$ BARTEL 77 CNTR  $e^+e^-$ 

<sup>1</sup> ABLIKIM 18O reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$ , which we rescale to our best values  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma_{238}/\Gamma$ NODE=M070R83  
NODE=M070R83

NODE=M070R83;LINKAGE=A

 $\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$ VALUE (units  $10^{-6}$ ) EVTS**21.4 ± 1.8 ± 2.5**

596

DOCUMENT ID

TECN

COMMENT

ABLIKIM 16P BES3  $J/\psi \rightarrow 5\gamma$  $\Gamma_{239}/\Gamma$ NODE=M070P01  
NODE=M070P01 $\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ 

VALUE CL%

**< 2.5 × 10<sup>-6</sup>** 95

DOCUMENT ID

TECN

COMMENT

ABLIKIM 16P BES3  $J/\psi \rightarrow 5\gamma$  $\Gamma_{243}/\Gamma$ NODE=M070P02  
NODE=M070P02 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$ 

VALUE CL%

**< 6.6 × 10<sup>-6</sup>** 95

DOCUMENT ID

TECN

COMMENT

ABLIKIM 16P BES3  $J/\psi \rightarrow 5\gamma$  $\Gamma_{244}/\Gamma$ NODE=M070P03  
NODE=M070P03 $\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**6.1 ± 1.0 OUR AVERAGE**

5.85 ± 0.3 ± 1.05

7.8 ± 1.2 ± 2.4

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> EDWARDS 83B CBAL  $J/\psi \rightarrow \eta\pi^+\pi^-$ <sup>1</sup> EDWARDS 83B CBAL  $J/\psi \rightarrow \eta 2\pi^0$ <sup>1</sup> Broad enhancement at 1700 MeV. $\Gamma_{245}/\Gamma$ NODE=M070R96  
NODE=M070R96

OCCUR=2

NODE=M070R96;LINKAGE=M

 $\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**6.2 ± 2.2 ± 0.9**

DOCUMENT ID

TECN

COMMENT

BAI 99 BES  $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$  $\Gamma_{246}/\Gamma$ NODE=M070S37  
NODE=M070S37 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ ) EVTS**5.25 ± 0.07 OUR AVERAGE**

5.27 ± 0.03 ± 0.05 36k

5.43 ± 0.23 ± 0.09 5.0k

4.77 ± 0.22 ± 0.06

5.24 ± 0.12 ± 0.11

5.55 ± 0.44 35k

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.50 ± 0.14 ± 0.53

4.30 ± 0.31 ± 0.71

4.04 ± 0.16 ± 0.85

4.39 ± 0.09 ± 0.66

4.1 ± 0.3 ± 0.6

2.9 ± 1.1

2.4 ± 0.7

DOCUMENT ID

TECN

COMMENT

Error includes scale factor of 1.3. See the ideogram below.

ABLIKIM 19T BES  $J/\psi \rightarrow \gamma\eta'$ <sup>1</sup> ABLIKIM 18O BES3  $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$ <sup>2</sup> ABLIKIM 11 BES3  $J/\psi \rightarrow \eta'\gamma$ PEDLAR 09 CLE3  $J/\psi \rightarrow \eta'\gamma$ ABLIKIM 06E BES2  $J/\psi \rightarrow \eta'\gamma$ BOLTON 92B MRK3  $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow$  $\gamma\gamma$  $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow$  $\pi^+\pi^-\pi^0$  $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$  $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$ BLOOM 83 CBAL  $e^+e^- \rightarrow 3\gamma + \text{hadrons}$ BRANDELIK 79C DASP  $e^+e^- \rightarrow 3\gamma$ BARTEL 76 CNTR  $e^+e^- \rightarrow 2\gamma\rho$  $\Gamma_{247}/\Gamma$ NODE=M070R84  
NODE=M070R84

OCCUR=2


OCCUR=2

<sup>1</sup> ABLIKIM 180 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$ , which we rescale to our best values  $B(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R84;LINKAGE=A

<sup>2</sup> ABLIKIM 11 reports  $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$  from a measurement of  $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$  assuming  $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$ ,  $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$ , which we rescale to our best values  $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$ ,  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R84;LINKAGE=AB



$\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{240}/\Gamma$

VALUE (units  $10^{-4}$ )    DOCUMENT ID    TECN    COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

10.5±2.0    SARANTSEV    21    RVUE     $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P95  
NODE=M070P95

$\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{241}/\Gamma$

VALUE (units  $10^{-5}$ )    DOCUMENT ID    TECN    COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

5±5    SARANTSEV    21    RVUE     $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P97  
NODE=M070P97

$\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$   $\Gamma_{242}/\Gamma$

VALUE (units  $10^{-5}$ )    DOCUMENT ID    TECN    COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

4±3    SARANTSEV    21    RVUE     $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P98  
NODE=M070P98

$\Gamma(\gamma f_0(980) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{248}/\Gamma$

VALUE (units  $10^{-5}$ )    DOCUMENT ID    TECN    COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

1.3±0.2    SARANTSEV    21    RVUE     $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P90  
NODE=M070P90

$\Gamma(\gamma f_0(980) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$   $\Gamma_{249}/\Gamma$

VALUE (units  $10^{-5}$ )    DOCUMENT ID    TECN    COMMENT

••• We do not use the following data for averages, fits, limits, etc. •••

0.8±0.3    SARANTSEV    21    RVUE     $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P91  
NODE=M070P91

$\Gamma(\gamma\rho\rho)/\Gamma_{\text{total}}$ 

VALUE (units  $10^{-3}$ ) CL%  
**4.5 ± 0.8 OUR AVERAGE**

4.7 ± 0.3 ± 0.9  
 3.75 ± 1.05 ± 1.20

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.09 90 <sup>3</sup> BISELLO 89B  $J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup>  $4\pi$  mass less than 2.0 GeV.

<sup>2</sup>  $4\pi$  mass less than 2.0 GeV. We have multiplied  $2\rho^0$  measurement by 3 to obtain  $2\rho$ .

<sup>3</sup>  $4\pi$  mass in the range 2.0–25 GeV.

 $\Gamma_{250}/\Gamma$ 

NODE=M070R94  
 NODE=M070R94

 $\Gamma(\gamma\rho\omega)/\Gamma_{\text{total}}$ 

VALUE CL%  
**<5.4 × 10<sup>-4</sup>** 90

DOCUMENT ID TECN COMMENT  
 ABLIKIM 08A BES2  $e^+e^- \rightarrow J/\psi$

 $\Gamma_{251}/\Gamma$ 

NODE=M070R05  
 NODE=M070R05

 $\Gamma(\gamma\rho\phi)/\Gamma_{\text{total}}$ 

VALUE CL%  
**<8.8 × 10<sup>-5</sup>** 90

DOCUMENT ID TECN COMMENT  
 ABLIKIM 08A BES2  $e^+e^- \rightarrow J/\psi$

 $\Gamma_{252}/\Gamma$ 

NODE=M070R06  
 NODE=M070R06

 $\Gamma(\gamma\omega\omega)/\Gamma_{\text{total}}$ 

VALUE (units  $10^{-3}$ ) EVTS  
**1.61 ± 0.33 OUR AVERAGE**

6.0 ± 4.8 ± 1.8  
 1.41 ± 0.2 ± 0.42 120 ± 17  
 1.76 ± 0.09 ± 0.45

DOCUMENT ID TECN COMMENT  
 ABLIKIM 08A BES2  $J/\psi \rightarrow \gamma\omega\pi^+\pi^-$   
 BISELLO 87 SPEC  $e^+e^-$ , hadrons $\gamma$   
 BALTRUSAIT..85C MRK3  $e^+e^- \rightarrow$  hadrons $\gamma$

 $\Gamma_{253}/\Gamma$ 

NODE=M070R97  
 NODE=M070R97

 $\Gamma(\gamma\phi\phi)/\Gamma_{\text{total}}$ 

VALUE (units  $10^{-4}$ ) EVTS  
**4.0 ± 1.2 OUR AVERAGE**

7.5 ± 0.6 ± 1.2 168  
 3.4 ± 0.8 ± 0.6 33 ± 7  
 3.1 ± 0.7 ± 0.4

Error includes scale factor of 2.1. See the ideogram below.


DOCUMENT ID TECN COMMENT  
 BAI 90B MRK3  $J/\psi \rightarrow \gamma 4K$   
<sup>1</sup> BISELLO 90 DM2  $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$   
<sup>1</sup> BISELLO 86B DM2  $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

<sup>1</sup>  $\phi\phi$  mass less than 2.9 GeV,  $\eta_c$  excluded.

 $\Gamma_{254}/\Gamma$ 

NODE=M070R98  
 NODE=M070R98

NODE=M070R98;LINKAGE=C

 $\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ 

VALUE (units  $10^{-3}$ )  
**2.8 ± 0.6 OUR AVERAGE**

1.66 ± 0.1 ± 0.58  
 3.8 ± 0.3 ± 0.6  
 4.0 ± 0.7 ± 1.0  
 4.3 ± 1.7

DOCUMENT ID TECN COMMENT  
<sup>1,2</sup> BAI 00D BES  $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$   
<sup>3</sup> AUGUSTIN 90 DM2  $J/\psi \rightarrow \gamma K \bar{K} \pi$   
<sup>3</sup> EDWARDS 82E CBAL  $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$   
<sup>3,4</sup> SCHARRE 80 MRK2  $e^+e^-$

 $\Gamma_{255}/\Gamma$ 

NODE=M070R89  
 NODE=M070R89

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78 ± 0.21 ± 0.33	3,5,6	AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$	
0.83 ± 0.13 ± 0.18	3,7,8	AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$	OCCUR=2
0.66 <sup>+0.17+0.24</sup> <sub>-0.16-0.15</sub>	3,6,9	BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
1.03 <sup>+0.21+0.26</sup> <sub>-0.18-0.19</sub>	3,8,10	BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=2

<sup>1</sup> Interference with the  $J/\psi(1S)$  radiative transition to the broad  $K \bar{K} \pi$  pseudoscalar state around 1800 is  $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$ .

<sup>2</sup> Interference with  $J/\psi \rightarrow \gamma f_1(1420)$  is  $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$ .

<sup>3</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow K \bar{K} \pi$ .

<sup>4</sup> Corrected for spin-zero hypothesis for  $\eta(1405)$ .

<sup>5</sup> From fit to the  $a_0(980) \pi 0^-+$  partial wave.

<sup>6</sup>  $a_0(980) \pi$  mode.

<sup>7</sup> From fit to the  $K^*(892) K 0^-+$  partial wave.

<sup>8</sup>  $K^* K$  mode.

<sup>9</sup> From  $a_0(980) \pi$  final state.

<sup>10</sup> From  $K^*(890) K$  final state.

NODE=M070R89;LINKAGE=BD

NODE=M070R89;LINKAGE=BE

NODE=M070R89;LINKAGE=B

NODE=M070R89;LINKAGE=C

NODE=M070R89;LINKAGE=H


NODE=M070R89;LINKAGE=K9

NODE=M070R89;LINKAGE=J

NODE=M070R89;LINKAGE=K8

NODE=M070R89;LINKAGE=D

NODE=M070R89;LINKAGE=E



**$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$   $\Gamma_{256} / \Gamma$**

VALUE (units  $10^{-4}$ )      DOCUMENT ID      TECN      COMMENT

**0.78 ± 0.20 OUR AVERAGE** Error includes scale factor of 1.8.

1.07 ± 0.17 ± 0.11      <sup>1</sup> BAI      04J      BES2       $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

0.64 ± 0.12 ± 0.07      <sup>1</sup> COFFMAN      90      MRK3       $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

<sup>1</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow \gamma\rho^0$ .

NODE=M070S30  
NODE=M070S30

NODE=M070S30;LINKAGE=C

**$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$   $\Gamma_{257} / \Gamma$**

VALUE (units  $10^{-4}$ )      EVTS      DOCUMENT ID      TECN      COMMENT

**3.0 ± 0.5 OUR AVERAGE**

2.6 ± 0.7 ± 0.4      BAI      99      BES       $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

3.38 ± 0.33 ± 0.64      <sup>1</sup> BOLTON      92B      MRK3       $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ± 0.6 ± 1.1      261      <sup>2</sup> AUGUSTIN      90      DM2       $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

<sup>1</sup> Via  $a_0(980) \pi$ .

<sup>2</sup> Includes unknown branching fraction to  $\eta\pi^+\pi^-$ .

NODE=M070S29  
NODE=M070S29

NODE=M070S29;LINKAGE=RR  
NODE=M070S29;LINKAGE=R

**$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0) / \Gamma_{\text{total}}$   $\Gamma_{258} / \Gamma$**

VALUE (units  $10^{-3}$ )      DOCUMENT ID      TECN      COMMENT

**1.7 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.3.

2.1 ± 0.4      BUGG      95      MRK3       $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$

1.36 ± 0.38      <sup>1,2</sup> BISELLO      89B      DM2       $J/\psi \rightarrow 4\pi\gamma$

<sup>1</sup> Estimated by us from various fits.

<sup>2</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

NODE=M070S19  
NODE=M070S19

NODE=M070S19;LINKAGE=A  
NODE=M070S19;LINKAGE=B

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$  $\Gamma_{259}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<82	95		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+ K^-$
-----	----	--	-----	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.03 \pm 0.92 \pm 0.91$	1.3k	1	ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
--------------------------	------	---	---------	-----	--

$10.36 \pm 1.51 \pm 1.54$	1.9k	2	ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
---------------------------	------	---	---------	-----	--

<sup>1</sup> Constructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma\phi$  invariant mass.

<sup>2</sup> Destructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma\phi$  invariant mass.

NODE=M070R77  
NODE=M070R77

OCCUR=2

NODE=M070R77;LINKAGE=B

NODE=M070R77;LINKAGE=A

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{260}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.63 $\times 10^{-6}$	90	ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$
------------------------	----	---------	-----	--

NODE=M070P38  
NODE=M070P38

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{261}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.86 $\times 10^{-6}$	90	ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$
------------------------	----	---------	-----	--

NODE=M070P39  
NODE=M070P39

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$  $\Gamma_{262}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.13 $\pm 0.09$	1,2	BISELLO	89B DM2 $J/\psi \rightarrow 4\pi\gamma$
-----------------	-----	---------	---

NODE=M070S20  
NODE=M070S20

<sup>1</sup> Estimated by us from various fits.

<sup>2</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

NODE=M070S20;LINKAGE=A  
NODE=M070S20;LINKAGE=B

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$  $\Gamma_{263}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.98 $\pm 0.08 \pm 0.32$	1045	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$
--------------------------	------	---------	-----	---

NODE=M070R04  
NODE=M070R04

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{264}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<4.80 $\times 10^{-6}$	90	ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$
------------------------	----	---------	-----	--

NODE=M070P40  
NODE=M070P40

 $\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$  $\Gamma_{265}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.14 $^{+0.50}_{-0.19}$ OUR AVERAGE				
-------------------------------------	--	--	--	--

NODE=M070S21  
NODE=M070S21

2.40 $\pm 0.10$ $^{+2.47}_{-0.18}$	1,2	ABLIKIM	16N	BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
------------------------------------	-----	---------	-----	--

4.4 $\pm 0.4 \pm 0.8$	196	2	ABLIKIM	08I	BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
-----------------------	-----	---	---------	-----	---

3.3 $\pm 0.8 \pm 0.5$		2	BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
-----------------------	--	---	-----	-----	--

2.7 $\pm 0.6 \pm 0.6$		2	BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
-----------------------	--	---	-----	-----	--

OCCUR=2

2.4 $^{+1.5}_{-1.0}$	3,4	BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
----------------------	-----	---------	-----	-------------------------------------

<sup>1</sup> From a partial wave analysis of  $J/\psi \rightarrow \gamma\phi\phi$  that also finds significant signals for for  $\eta(2100)$ ,  $0^-+$  phase space,  $f_0(2100)$ ,  $f_2(2010)$ ,  $f_2(2300)$ ,  $f_2(2340)$ , and a previously unseen  $0^-+$  state  $X(2500)$  ( $M = 2470^{+15+101}_{-19-23}$  MeV,  $\Gamma = 230^{+64+56}_{-35-33}$  MeV).

NODE=M070S21;LINKAGE=C

<sup>2</sup> Includes unknown branching fraction to  $\phi\phi$ .

<sup>3</sup> Estimated by us from various fits.

<sup>4</sup> Includes unknown branching fraction to  $\rho^0\rho^0$ .

NODE=M070S21;LINKAGE=U  
NODE=M070S21;LINKAGE=A  
NODE=M070S21;LINKAGE=B

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$  $\Gamma_{266}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.63 $\pm 0.12$ OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
-----------------------------	--	--	--	---

[(1.64  $\pm 0.12$ )  $\times 10^{-3}$  OUR 2022 AVERAGE Scale factor = 1.3]

NODE=M070R86  
NODE=M070R86

NEW

2.07 $\pm 0.16$ $^{+0.02}_{-0.07}$	2.4k	1,2	DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
------------------------------------	------	-----	-------	----	-----------------------------------

1.63 $\pm 0.26$ $^{+0.02}_{-0.06}$		3	ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
------------------------------------	--	---	---------	-----	---

OCCUR=2

1.42 $\pm 0.21$ $^{+0.02}_{-0.05}$		4	ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^0\pi^0$
------------------------------------	--	---	---------	-----	---

1.33 $\pm 0.05 \pm 0.20$		5	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
--------------------------	--	---	----------	----	---

1.36 $\pm 0.09 \pm 0.23$		5	BALTRUSAIT	87	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-$
--------------------------	--	---	------------	----	--


1.48 $\pm 0.25 \pm 0.30$	178		EDWARDS	82B	CBAL $e^+e^- \rightarrow 2\pi^0\gamma$
--------------------------	-----	--	---------	-----	--

2.0 $\pm 0.7$	35		ALEXANDER	78	PLUT $e^+e^-$
---------------	----	--	-----------	----	---------------

1.2 $\pm 0.6$	30	6	BRANDELIK	78B	DASP $e^+e^- \rightarrow \pi^+\pi^-\gamma$
---------------	----	---	-----------	-----	--

- <sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.
- <sup>2</sup> DOBBS 15 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>3</sup> ABLIKIM 06V reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>4</sup> ABLIKIM 06V reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>5</sup> Estimated using  $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$ . The errors do not contain the uncertainty in the  $f_2(1270)$  decay.
- <sup>6</sup> Restated by us to take account of spread of E1, M2, E3 transitions.

NODE=M070R86;LINKAGE=A  
 NODE=M070R86;LINKAGE=DO  
 NODE=M070R86;LINKAGE=AI  
 NODE=M070R86;LINKAGE=AL  
 NODE=M070R86;LINKAGE=X  
 NODE=M070R86;LINKAGE=T



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	$\Gamma_{267}/\Gamma$		
VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.58<sup>+0.08+0.59</sup><sub>-0.09-0.20</sub></b>	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P68  
 NODE=M070P68

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$	$\Gamma_{268}/\Gamma$		
VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.61 ± 0.08 OUR AVERAGE</b>			
0.69 ± 0.16 ± 0.20	1 BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \rho^0$
0.61 ± 0.04 ± 0.21	2 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 ± 0.09 ± 0.17	3 BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.625 ± 0.063 ± 0.103	4 BOLTON	92 MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.70 ± 0.08 ± 0.16	5 BOLTON	92B MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M070R88  
 NODE=M070R88

- <sup>1</sup> Assuming  $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$ .
- <sup>2</sup> Assuming  $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$ .
- <sup>3</sup> Assuming  $\Gamma(f_1(1285) \rightarrow \eta \pi \pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$ .
- <sup>4</sup> Obtained summing the sequential decay channels  
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$ ;  
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$ ;  
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$ ;  
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$ .
- <sup>5</sup> Using  $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$ , and including unknown branching ratio for  $a_0(980) \rightarrow \eta \pi$ .

NODE=M070R88;LINKAGE=BI  
 NODE=M070R88;LINKAGE=BD  
 NODE=M070R88;LINKAGE=BA  
 NODE=M070R88;LINKAGE=B

NODE=M070R88;LINKAGE=A

$\Gamma(\gamma f_0(1370) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$  $\Gamma_{269} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$38 \pm 10$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	----	--

NODE=M070P92  
NODE=M070P92

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$  $\Gamma_{270} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$4.19 \pm 0.73 \pm 1.34$	478	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
--------------------------	-----	--------------------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.3 \pm 0.4$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

NODE=M070R00  
NODE=M070R00

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070R00;LINKAGE=A

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$  $\Gamma_{271} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.07^{+0.08+0.36}_{-0.07-0.34}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	---------	------	--

NODE=M070P63  
NODE=M070P63

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$  $\Gamma_{272} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.5 \pm 1.0$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

NODE=M070P93  
NODE=M070P93

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$  $\Gamma_{273} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.9 \pm 0.3$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

NODE=M070P94  
NODE=M070P94

 $\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$  $\Gamma_{274} / \Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**0.79 ± 0.13 OUR AVERAGE**

$0.68 \pm 0.04 \pm 0.24$	BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
--------------------------	-----	-----	---

$0.76 \pm 0.15 \pm 0.21$	<sup>1,2</sup> AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
--------------------------	-------------------------	----	---

$0.87 \pm 0.14^{+0.14}_{-0.11}$	<sup>1</sup> BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
---------------------------------	------------------	-----	--

NODE=M070S31  
NODE=M070S31

OCCUR=2

<sup>1</sup> Included unknown branching fraction  $f_1(1420) \rightarrow K \bar{K} \pi$ .

NODE=M070S31;LINKAGE=A

<sup>2</sup> From fit to the  $K^*(892)K 1^{++}$  partial wave.

NODE=M070S31;LINKAGE=D

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$  $\Gamma_{275} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.09 ± 0.24 OUR AVERAGE**

$1.21 \pm 0.29 \pm 0.24$	174	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
--------------------------	-----	--------------------	----	-------------------------------------

$1.00 \pm 0.03 \pm 0.45$		<sup>2</sup> ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
--------------------------	--	----------------------	-----	---

$1.02 \pm 0.09 \pm 0.45$		<sup>2</sup> ABLIKIM	06V	BES2 $e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
--------------------------	--	----------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.90 \pm 0.17$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------------	-----------	----	--

$5.7 \pm 0.8$	<sup>3,4</sup> BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
---------------	---------------------	----	--

NODE=M070S32  
NODE=M070S32

OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070S32;LINKAGE=C

<sup>2</sup> Including unknown branching fraction to  $\pi\pi$ .

NODE=M070S32;LINKAGE=AB

<sup>3</sup> Including unknown branching ratio for  $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ .

NODE=M070S32;LINKAGE=A

<sup>4</sup> Assuming that  $f_0(1500)$  decays only to two S-wave dipions.

NODE=M070S32;LINKAGE=B

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$  $\Gamma_{276} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	<sup>1</sup> ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.1 \pm 0.4$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	----	--

NODE=M070S83  
NODE=M070S83

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

NODE=M070S83;LINKAGE=A



$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{total}$   $\Gamma_{277} / \Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.16^{+0.18}_{-0.56}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$0.7 \pm 0.3$	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P64  
NODE=M070P64

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta') / \Gamma_{total}$   $\Gamma_{278} / \Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
$18.1 \pm 1.1^{+1.9}_{-1.3}$	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
$12 \pm 5$	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.			

NODE=M070P99  
NODE=M070P99

NODE=M070P99;LINKAGE=A

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{total}$   $\Gamma_{279} / \Gamma$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.7$	BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M070S36  
NODE=M070S36

$\Gamma(\gamma f_2'(1525)) / \Gamma_{total}$   $\Gamma_{280} / \Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7^{+0.8}_{-0.5}$			<b>OUR AVERAGE</b> Error includes scale factor of 1.5. See the ideogram below.		
$8.1 \pm 0.9 \pm 0.2$	750	1,2	DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		3	BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		3	BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		3	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		3	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		3	BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<3.4	90	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78 PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$


NODE=M070R87  
NODE=M070R87

OCCUR=3  
OCCUR=4  
OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.  
<sup>2</sup> DOBBS 15 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2'(1525)) / \Gamma_{total}] \times [B(f_2'(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$  which we divide by our best value  $B(f_2'(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.  
<sup>3</sup> Using  $B(f_2'(1525) \rightarrow K\bar{K}) = 0.888$ .  
<sup>4</sup> Assuming isotropic production and decay of the  $f_2'(1525)$  and isospin.

NODE=M070R87;LINKAGE=B  
NODE=M070R87;LINKAGE=D0

NODE=M070R87;LINKAGE=A1  
NODE=M070R87;LINKAGE=I



$\Gamma(\gamma f_2(1565) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$   $\Gamma_{283} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q31  
NODE=M070Q31

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.32 \pm 0.05^{+0.12}_{-0.02}$	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

NODE=M070Q31;LINKAGE=A

 $\Gamma(\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$   $\Gamma_{281} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070P69  
NODE=M070P69

$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	---------	-----------	---

 $\Gamma(\gamma f_2'(1525) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{282} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S86  
NODE=M070S86

$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	----------	---------------------------------------

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

NODE=M070S86;LINKAGE=A

 $\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$   $\Gamma_{284} / \Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R02  
NODE=M070R02

$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
--------------------------	-----	---------	---------	---

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$   $\Gamma_{285} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070B01  
NODE=M070B01

**3.8 ± 0.5 OUR AVERAGE**

$3.72 \pm 0.30 \pm 0.43$	483	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
--------------------------	-----	--------------------	----	-------------------------------------

$3.96 \pm 0.06 \pm 1.12$		<sup>2</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
--------------------------	--	----------------------	----------	---

$3.99 \pm 0.15 \pm 2.64$		<sup>2</sup> ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
--------------------------	--	----------------------	----------	---

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.6 \pm 0.2$		<sup>3</sup> SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	--	------------------------	---------	--

$2.5 \pm 1.6 \pm 0.8$		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
-----------------------	--	-----	---------	---

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Including unknown branching fraction to  $\pi \pi$ .

<sup>3</sup> There is a further  $(2.4 \pm 0.8) \times 10^{-4}$  scalar contribution at 1765 MeV.

NODE=M070B01;LINKAGE=A  
NODE=M070B01;LINKAGE=A  
NODE=M070B01;LINKAGE=B

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$   $\Gamma_{286} / \Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M070R91  
NODE=M070R91

**9.5 ± 1.0 OUR AVERAGE** Error includes scale factor of 1.5. See the ideogram below.

$8.00^{+0.12+1.24}_{-0.08-0.40}$		<sup>1</sup> ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	--	----------------------	-----------	---

$11.76 \pm 0.54 \pm 0.94$	1.2k	<sup>2</sup> DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
---------------------------	------	--------------------	----	---------------------------------------

$9.62 \pm 0.29^{+3.51}_{-1.86}$		<sup>3</sup> BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
---------------------------------	--	------------------	---------	---------------------------------------

$5.0 \pm 0.8^{+1.8}_{-0.4}$		<sup>1,4</sup> BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
-----------------------------	--	--------------------	---------	-------------------------------------

$9.2 \pm 1.4 \pm 1.4$		<sup>1</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
-----------------------	--	-----------------------	--------	-------------------------------------

$10.4 \pm 1.2 \pm 1.6$		<sup>1</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
------------------------	--	-----------------------	--------	---

OCCUR=2

$9.6 \pm 1.2 \pm 1.8$		<sup>1</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
-----------------------	--	-----------------------------	------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.3 \pm 0.8$		<sup>5</sup> SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
---------------	--	------------------------	---------	--

$1.6 \pm 0.2^{+0.6}_{-0.2}$		<sup>1,6</sup> BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
-----------------------------	--	--------------------	---------	-------------------------------------

OCCUR=2

$< 0.8$	90	<sup>7</sup> BISELLO	89B	$J/\psi \rightarrow 4\pi \gamma$
---------	----	----------------------	-----	----------------------------------


$1.6 \pm 0.4 \pm 0.3$		<sup>8</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
-----------------------	--	-----------------------------	------	---

OCCUR=2

$3.8 \pm 1.6$		<sup>9</sup> EDWARDS	82D CBAL	$e^+ e^- \rightarrow \eta \eta \gamma$
---------------	--	----------------------	----------	--

- <sup>1</sup> Includes unknown branching fraction to  $K^+ K^-$  or  $K_S^0 K_S^0$ . We have multiplied  $K^+ K^-$  measurement by 2, and  $K_S^0 K_S^0$  by 4 to obtain  $K\bar{K}$  result.
- <sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.
- <sup>3</sup> Includes unknown branching ratio to  $K^+ K^-$  or  $K_S^0 K_S^0$ .
- <sup>4</sup> Assuming  $J^P = 2^+$  for  $f_0(1710)$ .
- <sup>5</sup> There is a further  $(6 \pm 2) \times 10^{-4}$  scalar contribution at 1765 MeV.
- <sup>6</sup> Assuming  $J^P = 0^+$  for  $f_0(1710)$ .
- <sup>7</sup> Includes unknown branching fraction to  $\rho^0 \rho^0$ .
- <sup>8</sup> Includes unknown branching fraction to  $\pi^+ \pi^-$ .
- <sup>9</sup> Includes unknown branching fraction to  $\eta \eta$ .

NODE=M070R91;LINKAGE=B  
 NODE=M070R91;LINKAGE=D  
 NODE=M070R91;LINKAGE=K9  
 NODE=M070R91;LINKAGE=A1  
 NODE=M070R91;LINKAGE=E  
 NODE=M070R91;LINKAGE=A2  
 NODE=M070R91;LINKAGE=C  
 NODE=M070R91;LINKAGE=Z  
 NODE=M070R91;LINKAGE=A



**$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$**   **$\Gamma_{287} / \Gamma$**

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.31 \pm 0.06 \pm 0.08</math></b>	180	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma \omega \omega$

NODE=M070R01  
 NODE=M070R01

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$**   **$\Gamma_{288} / \Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2.35^{+0.13+1.24}_{-0.11-0.74}</math></b>	5.5k	<sup>1</sup> ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

NODE=M070S84  
 NODE=M070S84

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2  $\pm$  0.4 <sup>2</sup> SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

<sup>2</sup> There is a further  $(0.7 \pm 0.1) \times 10^{-4}$  scalar contribution at 1765 MeV.

NODE=M070S84;LINKAGE=A  
 NODE=M070S84;LINKAGE=B

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$**   **$\Gamma_{289} / \Gamma$**

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
6.5 $\pm$ 2.5	<sup>1</sup> SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

<sup>1</sup> There is a further  $(2.5 \pm 1.1) \times 10^{-5}$  scalar contribution at 1765 MeV.

NODE=M070Q00  
 NODE=M070Q00

NODE=M070Q00;LINKAGE=A

**$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \phi) / \Gamma_{\text{total}}$**   **$\Gamma_{290} / \Gamma$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.5 <math>\pm</math> 0.6 OUR AVERAGE</b>				
2.00 $\pm$ 0.08 $^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J	BES3 $J/\psi \rightarrow \gamma \omega \phi$
2.61 $\pm$ 0.27 $\pm$ 0.65	95	ABLIKIM	06J	BES2 $J/\psi \rightarrow \gamma \omega \phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1  $\pm$  0.1 <sup>1</sup> SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

<sup>1</sup> There is a further  $(2.2 \pm 0.4) \times 10^{-4}$  scalar contribution at 1765 MeV.

NODE=M070S79  
 NODE=M070S79

NODE=M070S79;LINKAGE=A

$\Gamma(\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$				$\Gamma_{291}/\Gamma$	NODE=M070P65 NODE=M070P65
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$		
$\Gamma(\gamma f_0(1770) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$				$\Gamma_{294}/\Gamma$	NODE=M070Q30 NODE=M070Q30
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.11 \pm 0.01^{+0.04}_{-0.03}$	<sup>1</sup> ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$		
<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.					NODE=M070Q30;LINKAGE=A
$\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$				$\Gamma_{292}/\Gamma$	NODE=M070S87 NODE=M070S87
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>COMMENT</u>		
$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	<sup>1</sup> ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$		
<sup>1</sup> From partial wave analysis including all possible combinations of $0^{++}$ , $2^{++}$ , and $4^{++}$ resonances.					NODE=M070S87;LINKAGE=A
$\Gamma(\gamma \eta_1(1855) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$				$\Gamma_{293}/\Gamma$	NODE=M070Q44 NODE=M070Q44
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.70 \pm 0.41^{+0.16}_{-0.35}$	<sup>1</sup> ABLIKIM	22AI	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$		
<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave. For analysis details see ABLIKIM 22AS.					NODE=M070Q44;LINKAGE=A
$\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega)/\Gamma_{\text{total}}$				$\Gamma_{295}/\Gamma$	NODE=M070R03 NODE=M070R03
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H BES $J/\psi \rightarrow \gamma \omega \omega$		
$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$				$\Gamma_{296}/\Gamma$	NODE=M070B06 NODE=M070B06
<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$0.7 \pm 0.1 \pm 0.2$	BAI	00B	BES $J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$		
$\Gamma(\gamma f_2(2010) \rightarrow \gamma \eta \eta')/\Gamma_{\text{total}}$				$\Gamma_{297}/\Gamma$	NODE=M070Q37 NODE=M070Q37
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$0.71 \pm 0.06^{+0.10}_{-0.06}$	<sup>1</sup> ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$		
<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.					NODE=M070Q37;LINKAGE=A
$\Gamma(\gamma f_0(2020) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$				$\Gamma_{298}/\Gamma$	NODE=M070Q01 NODE=M070Q01
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$42 \pm 10$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$		
$\Gamma(\gamma f_0(2020) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$				$\Gamma_{299}/\Gamma$	NODE=M070Q02 NODE=M070Q02
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$55 \pm 25$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$		
$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$				$\Gamma_{300}/\Gamma$	NODE=M070Q03 NODE=M070Q03
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$10 \pm 10$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$		
$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta' \eta')/\Gamma_{\text{total}}$				$\Gamma_{301}/\Gamma$	NODE=M070Q25 NODE=M070Q25
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.63 \pm 0.06^{+0.31}_{-0.46}$	<sup>1</sup> ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$		
<sup>1</sup> From a partial wave analysis of the systems $(\gamma X)$ , with $X \rightarrow \eta' \eta'$ , and $(\eta' X)$ , with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.					NODE=M070Q25;LINKAGE=A

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$  $\Gamma_{302} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$2.28 \pm 0.12^{+0.29}_{-0.20}$	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

NODE=M070Q35  
NODE=M070Q35

NODE=M070Q35;LINKAGE=A

 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$  $\Gamma_{303} / \Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
$2.7 \pm 0.5 \pm 0.5$	<sup>1</sup> BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Assuming branching fraction  $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$ .

NODE=M070S7  
NODE=M070S7

NODE=M070S7;LINKAGE=V

 $\Gamma(\gamma f_4(2050) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$  $\Gamma_{304} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$0.06 \pm 0.01^{+0.03}_{-0.01}$	<sup>1</sup> ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$  P-wave.

NODE=M070Q32  
NODE=M070Q32

NODE=M070Q32;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$  $\Gamma_{305} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	<sup>1</sup> ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 1.5

SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

NODE=M070S85  
NODE=M070S85

NODE=M070S85;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$  $\Gamma_{307} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.24 \pm 0.48 \pm 0.87$	744	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.8

SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070B08  
NODE=M070B08

NODE=M070B08;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$  $\Gamma_{306} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$32 \pm 20$	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M070Q04  
NODE=M070Q04

 $\Gamma(\gamma f_0(2200)) / \Gamma_{\text{total}}$  $\Gamma_{308} / \Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
1.5	<sup>1</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Includes unknown branching fraction to  $K_S^0 K_S^0$ .

NODE=M070S18  
NODE=M070S18

NODE=M070S18;LINKAGE=A

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$  $\Gamma_{311} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$5 \pm 2$	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M070Q05  
NODE=M070Q05

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$  $\Gamma_{309} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$5.86 \pm 0.49 \pm 1.20$	490	<sup>1</sup> DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.5 ± 0.5

SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070B09  
NODE=M070B09

NODE=M070B09;LINKAGE=A

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{310}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$2.72^{+0.08+0.17}_{-0.06-0.47}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P66  
NODE=M070P66

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$   $\Gamma_{312}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ± 0.4 SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070Q06  
NODE=M070Q06

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{313}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300			<sup>1</sup> BAI	96B BES	$e^+e^- \rightarrow \gamma p\bar{p}, K\bar{K}$
>250	99.9		<sup>2</sup> HASAN	96 SPEC	$\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		<sup>3</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95		<sup>3</sup> AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	<sup>3</sup> BALTRUSAIT...86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	<sup>3</sup> BALTRUSAIT...86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

NODE=M070R92  
NODE=M070R92

OCCUR=2

OCCUR=2

<sup>1</sup> Using BARNES 93.

<sup>2</sup> Using BAI 96B.

<sup>3</sup> Includes unknown branching fraction to  $K^+ K^-$  or  $K_S^0 K_S^0$ .

NODE=M070R92;LINKAGE=A  
NODE=M070R92;LINKAGE=M  
NODE=M070R92;LINKAGE=W

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$   $\Gamma_{314}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< **3.9** 90 <sup>1,2</sup>DOBBS 15  $J/\psi \rightarrow \gamma \pi \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$14 \pm 8 \pm 4$		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
$8.4 \pm 2.6 \pm 3.0$		BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $\pi^+ \pi^-$  and  $\pi^0 \pi^0$  are  $2.6/5.2 \times 10^{-5}$  and  $1.3/1.9 \times 10^{-5}$ , respectively.

NODE=M070B02  
NODE=M070B02

NODE=M070B02;LINKAGE=A  
NODE=M070B02;LINKAGE=D0

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$   $\Gamma_{315}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< **4.1** 90 <sup>1,2</sup>DOBBS 15  $J/\psi \rightarrow \gamma K \bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.6		<sup>3</sup> DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
< 2.9		<sup>3</sup> DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.6 \pm 2.9 \pm 2.4$		BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$		BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070B03  
NODE=M070B03

OCCUR=2

OCCUR=2

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $K^+ K^-$  and  $K_S^0 K_S^0$  are  $1.7/3.1 \times 10^{-5}$  and  $1.2/2.0 \times 10^{-5}$ , respectively.

<sup>3</sup> For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

NODE=M070B03;LINKAGE=A  
NODE=M070B03;LINKAGE=D0

NODE=M070B03;LINKAGE=DE

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$   $\Gamma_{316}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.5 \pm 0.6 \pm 0.5$  BAI 96B BES  $e^+e^- \rightarrow J/\psi \rightarrow \gamma p \bar{p}$

NODE=M070B04  
NODE=M070B04

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{317}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.95 \pm 0.21^{+0.66}_{-0.72}$  ABLIKIM 18AA BES3  $J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.1 SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P67  
NODE=M070P67

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$   $\Gamma_{318}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 2 SARANTSEV 21 RVUE  $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070Q07  
NODE=M070Q07

$\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{319} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ± 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	--------------	------	---

NODE=M070Q08  
NODE=M070Q08

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$   $\Gamma_{320} / \Gamma$ 

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>6.09 ± 0.64<sup>+4.00</sup><sub>-1.68</sub></b>	<sup>1</sup> ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
--	----------------------	-----	--

<sup>1</sup> From a partial wave analysis of the systems ( $\gamma X$ ), with  $X \rightarrow \eta' \eta'$ , and ( $\eta' X$ ), with  $X \rightarrow \gamma \eta'$  in the decay  $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q26  
NODE=M070Q26

NODE=M070Q26;LINKAGE=A

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$   $\Gamma_{321} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.10 ± 0.02 <sup>+0.01</sup> <sub>-0.02</sub>	<sup>1</sup> ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
---	----------------------	------	---

<sup>1</sup> From a Breit-Wigner fit involving 9 resonances and a resonating exotic  $\eta_1(1855) \rightarrow \eta \eta'$   $P$ -wave.

NODE=M070Q36  
NODE=M070Q36

NODE=M070Q36;LINKAGE=A

 $\Gamma(\gamma f_0(2470) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$   $\Gamma_{325} / \Gamma$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>8.18 ± 1.77<sup>+3.73</sup><sub>-2.23</sub></b>	<sup>1</sup> ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
--	----------------------	-----	--

<sup>1</sup> From a partial wave analysis of the systems ( $\gamma X$ ), with  $X \rightarrow \eta' \eta'$ , and ( $\eta' X$ ), with  $X \rightarrow \gamma \eta'$  in the decay  $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q27  
NODE=M070Q27

NODE=M070Q27;LINKAGE=A

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$   $\Gamma_{322} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>5.60<sup>+0.62+2.37</sup><sub>-0.65-2.07</sub></b>	5.5k	<sup>1</sup> ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$
---	------	----------------------	-----	--

<sup>1</sup> From partial wave analysis including all possible combinations of  $0^{++}$ ,  $2^{++}$ , and  $4^{++}$  resonances.

NODE=M070S88  
NODE=M070S88

NODE=M070S88;LINKAGE=A

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$   $\Gamma_{324} / \Gamma$ 

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>8.67 ± 0.70<sup>+0.61</sup><sub>-1.67</sub></b>	<sup>1</sup> ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
--	----------------------	-----	--

<sup>1</sup> From a partial wave analysis of the systems ( $\gamma X$ ), with  $X \rightarrow \eta' \eta'$ , and ( $\eta' X$ ), with  $X \rightarrow \gamma \eta'$  in the decay  $J/\psi \rightarrow \gamma \eta' \eta'$ . The intermediate resonance  $X$  is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q29  
NODE=M070Q29

NODE=M070Q29;LINKAGE=A

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$   $\Gamma_{323} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

<b>5.54<sup>+0.34+3.82</sup><sub>-0.40-1.49</sub></b>	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	------	--

NODE=M070P70  
NODE=M070P70

 $\Gamma(\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta') / \Gamma_{\text{total}}$   $\Gamma_{326} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>2.7<sup>+0.6</sup><sub>-0.8</sub></b>	OUR AVERAGE	Error includes scale factor of 1.6.		
--	-------------	-------------------------------------	--	--

3.93 ± 0.38 <sup>+0.31</sup> <sub>-0.84</sub>	<sup>1</sup> ABLIKIM	16J	BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
---	----------------------	-----	------	---

2.2 ± 0.4 ± 0.4	264	ABLIKIM	05R	BES2 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
-----------------	-----	---------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.87 ± 0.09 <sup>+0.49</sup> <sub>-0.52</sub>	4265	<sup>2</sup> ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
---	------	----------------------	-----	--

<sup>1</sup> From a fit of the measured  $\pi^+ \pi^- \eta'$  lineshape that accounts for the abrupt distortion observed at the  $p\bar{p}$  threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ( $M \approx 1919$  MeV;  $\Gamma \approx 51$  MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the  $X(1835)$ ; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

<sup>2</sup> From a fit of the  $\pi^+ \pi^- \eta'$  mass distribution to a combination of  $\gamma f_1(1510)$ ,  $\gamma X(1835)$ , and two states  $\gamma X(2120)$  and  $\gamma X(2370)$ , for  $M(\pi^+ \pi^- \eta') < 2.8$  GeV, and accounting for backgrounds from non- $\eta'$  events and  $J/\psi \rightarrow \pi^0 \pi^+ \pi^- \eta'$ .

NODE=M070R78  
NODE=M070R78

NODE=M070R78;LINKAGE=A

NODE=M070R78;LINKAGE=A

$\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$  $\Gamma_{327}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S71  
NODE=M070S71

**0.77<sup>+0.15</sup><sub>-0.09</sub> OUR AVERAGE**

0.90 <sup>+0.04+0.27</sup> <sub>-0.11-0.55</sub>		<sup>1</sup> ABLIKIM	12D	BES3	$J/\psi \rightarrow \gamma p \bar{p}$
1.14 <sup>+0.43+0.42</sup> <sub>-0.30-0.26</sub>	231	<sup>2</sup> ALEXANDER	10	CLEO	$J/\psi \rightarrow \gamma p \bar{p}$
0.70 $\pm$ 0.04 <sup>+0.19</sup> <sub>-0.08</sub>		BAI	03F	BES2	$J/\psi \rightarrow \gamma p \bar{p}$

<sup>1</sup> From the fit including final state interaction effects in isospin 0 S-wave according to SIBIRTSEV 05A.

<sup>2</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma X(1835)$ ,  $\gamma R$  with  $M(R) = 2100$  MeV and  $\Gamma(R) = 160$  MeV, and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV.

NODE=M070S71;LINKAGE=AK

NODE=M070S71;LINKAGE=AL

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$  $\Gamma_{328}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070S96  
NODE=M070S96

**3.31<sup>+0.33+1.96</sup><sub>-0.30-1.29</sub>** ABLIKIM 15T BES3  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \phi(1020))/\Gamma_{\text{total}}$  $\Gamma_{329}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070P37  
NODE=M070P37

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.77 $\pm$ 0.35 $\pm$ 0.25	305	<sup>1</sup> ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
8.09 $\pm$ 1.99 $\pm$ 1.36	1.3k	<sup>2</sup> ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$

OCCUR=2

<sup>1</sup> Constructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma\phi$  invariant mass.

<sup>2</sup> Destructive interference between the  $X(1835)$  and  $\eta(1405)/\eta(1475)$  is assumed in a fit to the  $\gamma\phi$  invariant mass.

NODE=M070P37;LINKAGE=A

NODE=M070P37;LINKAGE=B

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \gamma)/\Gamma_{\text{total}}$  $\Gamma_{330}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070P41  
NODE=M070P41

**<3.56  $\times$  10<sup>-6</sup>** 90 ABLIKIM 180 BES3  $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$

 $\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$  $\Gamma_{331}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S82  
NODE=M070S82

**2.44 $\pm$ 0.36<sup>+0.60</sup><sub>-0.74</sub>** 0.6k ABLIKIM 13U BES3  $J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$

 $\Gamma(\gamma X(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$  $\Gamma_{332}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070P86  
NODE=M070P86

**1.79 $\pm$ 0.23 $\pm$ 0.65** ABLIKIM 20Q BES3  $J/\psi \rightarrow \gamma K^+ K^- \eta'$

 $\Gamma(\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$  $\Gamma_{333}/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070P87  
NODE=M070P87

**1.18 $\pm$ 0.32 $\pm$ 0.39** ABLIKIM 20Q BES3  $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$

 $\Gamma(\gamma X(2370) \rightarrow \gamma \eta \eta \eta')/\Gamma_{\text{total}}$  $\Gamma_{334}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M070P88  
NODE=M070P88

**<9.2** 90 ABLIKIM 21C BES3  $J/\psi(1S) \rightarrow \gamma \eta \eta \eta'$

 $\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}}$  $\Gamma_{335}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M070R90  
NODE=M070R90

**0.38 $\pm$ 0.07 $\pm$ 0.07** 49 EATON 84 MRK2  $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11 90 PERUZZI 78 MRK1  $e^+ e^-$

 $\Gamma(\gamma p \bar{p} \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{336}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070R93  
NODE=M070R93

**<0.79  $\times$  10<sup>-3</sup>** 90 EATON 84 MRK2  $e^+ e^-$

 $\Gamma(\gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$  $\Gamma_{337}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070S8  
NODE=M070S8

**<0.13  $\times$  10<sup>-3</sup>** 90 HENRARD 87 DM2  $e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.16  $\times$  10<sup>-3</sup> 90 BAI 98G BES  $e^+ e^-$



$\Gamma(\gamma A^0 \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$  $\Gamma_{338}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-6}$	90	88M	<sup>1</sup> ABLIKIM	20K BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$<6.3 \times 10^{-6}$	90	3.7M	<sup>2</sup> INSLER	10 CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$

NODE=M070S68  
NODE=M070S68

<sup>1</sup> For a narrow state,  $A^0$ , with mass  $m_{A^0} < 1.2$  GeV. The limit varies with  $m_{A^0}$ , reaching its largest value of  $1.7 \times 10^{-6}$  at 1.2 GeV and being  $7.0 \times 10^{-7}$  for  $m_{A^0} = 0$ .

NODE=M070S68;LINKAGE=A

<sup>2</sup> The limit varies with mass  $m_{A^0}$  of a narrow state  $A^0$  and is  $4.3 \times 10^{-6}$  for  $m_{A^0} = 0$ , reaches its largest value of  $6.3 \times 10^{-6}$  at  $m_{A^0} = 500$  MeV, and is  $3.6 \times 10^{-6}$  at  $m_{A^0} = 960$  MeV.

NODE=M070S68;LINKAGE=IN

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_{339}/\Gamma$ (narrow state  $A^0$  with  $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$ )

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$ (CL = 90%)		$[<5 \times 10^{-6}$ (CL = 90%) OUR 2022 BEST LIMIT]		
$<7.8 \times 10^{-7}$	90	<sup>1</sup> ABLIKIM	22H BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$<0.5 \times 10^{-5}$	90	<sup>2</sup> ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
$<2.1 \times 10^{-5}$	90	<sup>3</sup> ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$

NODE=M070S76  
NODE=M070S76

<sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of  $m_{A^0}$  is in the range  $(1.2\text{--}778.0) \times 10^{-9}$ .

NODE=M070S76;LINKAGE=B

<sup>2</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of  $m_{A^0}$  is in the range  $(2.8\text{--}495.3) \times 10^{-8}$ .

NODE=M070S76;LINKAGE=A

<sup>3</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of  $m_{A^0}$  ranges from  $4 \times 10^{-7}$  to  $2.1 \times 10^{-5}$ .

NODE=M070S76;LINKAGE=AB

## DALITZ DECAYS

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{340}/\Gamma$ 

VALUE (units $10^{-7}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$

NODE=M070S89  
NODE=M070S89 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{341}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.42 \pm 0.04 \pm 0.07$	2.47k	<sup>1,2</sup> ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$
$1.16 \pm 0.07 \pm 0.06$	320	<sup>1</sup> ABLIKIM	14I BES3	$J/\psi \rightarrow \eta e^+ e^-$

NODE=M070S90  
NODE=M070S90

<sup>1</sup> Using both  $\eta \rightarrow \gamma \gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

NODE=M070S90;LINKAGE=A

<sup>2</sup> Approximation of the transition form factor squared as an incoherent sum of the  $\rho$ -meson and one-pole non-resonant amplitudes gives the pole mass  $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$  GeV. Supersedes ABLIKIM 14I.

NODE=M070S90;LINKAGE=C

 $\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{342}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.59 \pm 0.07 \pm 0.17$	8.9k	<sup>1</sup> ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$
$5.81 \pm 0.16 \pm 0.31$	1.4k	<sup>1,2</sup> ABLIKIM	14I BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

NODE=M070S91  
NODE=M070S91

<sup>1</sup> Using both  $\eta' \rightarrow \gamma \pi^+ \pi^-$  and  $\eta' \rightarrow \pi^+ \pi^- \eta$  decays.

NODE=M070S91;LINKAGE=A

<sup>2</sup> Superseded by ABLIKIM 19H.

NODE=M070S91;LINKAGE=B

 $\Gamma(X(1835) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$  $\Gamma_{343}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.58 \pm 0.19 \pm 0.16$	1364	<sup>1</sup> ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$

NODE=M070Q22  
NODE=M070Q22

<sup>1</sup> Assuming constructive interference. Destructive interference gives a value of  $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$  for this branching fraction.

NODE=M070Q22;LINKAGE=A

 $\Gamma(X(2120) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$  $\Gamma_{344}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.12 \pm 0.06$	310	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$

NODE=M070Q24  
NODE=M070Q24 $\Gamma(X(2370) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$  $\Gamma_{345}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.08 \pm 0.14 \pm 0.10$	397	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$

NODE=M070Q23  
NODE=M070Q23

$\Gamma(\eta U \rightarrow \eta e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{346}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.11 \times 10^{-7}$	90	<sup>1</sup> ABLIKIM 19A	BES3	$J/\psi \rightarrow \eta e^+ e^-$

NODE=M070P42  
 NODE=M070P42

<sup>1</sup> For a dark photon  $U$  with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of  $m_U$  range from  $1.9 \times 10^{-8}$  to  $91.1 \times 10^{-8}$ .

NODE=M070P42;LINKAGE=A

 $\Gamma(\eta'(958) U \rightarrow \eta'(958) e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{347}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-7}$	90	<sup>1</sup> ABLIKIM 19H	BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

NODE=M070P61  
 NODE=M070P61

<sup>1</sup> For a dark photon  $U$  with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of  $m_U$  range from  $1.8 \times 10^{-8}$  to  $2.0 \times 10^{-7}$ . The corresponding limits on the branching fraction  $J/\psi \rightarrow \eta' U$  range from  $5.7 \times 10^{-8}$  to  $7.4 \times 10^{-7}$ .

NODE=M070P61;LINKAGE=A

 $\Gamma(\phi e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{348}/\Gamma$ 

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2$	90	<sup>1</sup> ABLIKIM 19AB	BES3	$J/\psi \rightarrow \phi e^+ e^-$

NODE=M070P82  
 NODE=M070P82

<sup>1</sup> Using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$  and  $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) = (34.49 \pm 0.30)\%$ .

NODE=M070P82;LINKAGE=A

———— WEAK DECAYS ————

 $\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{349}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.1 \times 10^{-8}$	90	ABLIKIM 21Q	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S53  
 NODE=M070S53

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-5}$	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	-------------	------	------------------------------

 $\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{350}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-8}$	90	<sup>1</sup> ABLIKIM 17AF	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S54  
 NODE=M070S54

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-5}$	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	-------------	------	------------------------------

<sup>1</sup> Using  $D^0$  decays to  $K^- \pi^+$ ,  $K^- \pi^+ \pi^0$ , and  $K^- \pi^+ \pi^+ \pi^-$ .

NODE=M070S54;LINKAGE=A

 $\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{351}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S55  
 NODE=M070S55

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.6 \times 10^{-5}$	90	<sup>1</sup> ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	--------------------------	------	------------------------------

<sup>1</sup> Using  $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5 \%$ .

NODE=M070S55;LINKAGE=AB

 $\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{352}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070B13  
 NODE=M070B13

 $\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{353}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-5}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S61  
 NODE=M070S61

 $\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{354}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S62  
 NODE=M070S62

 $\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{355}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-6}$	90	ABLIKIM 14K	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S93  
 NODE=M070S93

 $\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{356}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S63  
 NODE=M070S63

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{357}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$	

NODE=M070S92  
NODE=M070S92

———— CHARGE CONJUGATION (C), PARITY (P), ————  
———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

NODE=M070315

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{358}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2.7 \times 10^{-7}$	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

NODE=M070R80  
NODE=M070R80

••• We do not use the following data for averages, fits, limits, etc. •••

$< 0.5 \times 10^{-5}$	90	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$< 1.6 \times 10^{-4}$	90	<sup>1</sup> WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
$< 2.2 \times 10^{-5}$	90	ABLIKIM	07J BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
$< 50 \times 10^{-5}$	90	BARTEL	77 CNTR	$e^+ e^-$

<sup>1</sup> WICHT 08 reports  $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.020 \times 10^{-3}$ .

NODE=M070R80;LINKAGE=WI

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$					$\Gamma_{359}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-6}$	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

NODE=M070S95  
NODE=M070S95

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					$\Gamma_{360}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-7}$	90	ABLIKIM	13L BES3	$e^+ e^- \rightarrow J/\psi$	

NODE=M070S39  
NODE=M070S39

••• We do not use the following data for averages, fits, limits, etc. •••

$<1.1 \times 10^{-6}$	90	BAI	03D BES	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	-----	---------	------------------------------

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$					$\Gamma_{361}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.5 \times 10^{-8}$	90	ABLIKIM	21M BES3	$e^+ e^- \rightarrow J/\psi$	

NODE=M070S40  
NODE=M070S40

••• We do not use the following data for averages, fits, limits, etc. •••

$<8.3 \times 10^{-6}$	90	<sup>1</sup> ABLIKIM	04 BES	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	----------------------	--------	------------------------------

<sup>1</sup> Superseded by ABLIKIM 21M.

NODE=M070S40;LINKAGE=A

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					$\Gamma_{362}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	ABLIKIM	04 BES	$e^+ e^- \rightarrow J/\psi$	

NODE=M070S41  
NODE=M070S41

$\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{363}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.9 \times 10^{-8}$	90	ABLIKIM	19AF BES3	$e^+ e^- \rightarrow J/\psi \rightarrow p K^- \pi^+ e^- (+ \text{c.c.})$	

NODE=M070P74  
NODE=M070P74

———— OTHER DECAYS ————

NODE=M070325

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$					$\Gamma_{364}/\Gamma_5$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-2}$	90	LEES	13I BABR	$B \rightarrow K^{(*)} J/\psi$	

NODE=M070S80  
NODE=M070S80

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$					$\Gamma_{364}/\Gamma_7$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	

NODE=M070S60  
NODE=M070S60

**J/ψ(1S) REFERENCES**

NODE=M070

ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)	REFID=61911
ABLIKIM	22AI	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61881
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61902
ABLIKIM	22B	PRL 129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61636
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61637
ABLIKIM	22H	PR D105 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61643
ANASHIN	22	EPJ C82 938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=61894
ABLIKIM	21AM	PR D104 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61445
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61463
ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61030
ABLIKIM	21M	PR D103 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61118

ABLIKIM	21Q	JHEP 2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61125
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61113
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61451
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60048
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60316
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60457
ANASHIN	20	JHEP 2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=60512
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59517
Also		PR D104 099901 (errata)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61449
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59846
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59850
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59857
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59890
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59604
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59615
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59747
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59773
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)	REFID=59614
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59456
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58849
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=59102
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58315
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58317
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58324
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57903
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57967
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57966
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57990
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57265
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57454
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57509
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57510
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57511
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)	REFID=57512
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57522
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57566
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56773
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56776
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56785
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55900
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55902
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55905
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56238
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56388
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=55655
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54920
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54954
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54955
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55300
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55392
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55402
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54265
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54267
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54268
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54269
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54273
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54863
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16715
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53220
DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53533
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)	REFID=53359
ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52718
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53099
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52047

ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52128
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52253
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52255
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52256
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52571
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
BESSION	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=52685
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52072
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (err.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50986
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51058
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51128
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51130
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51036
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
AUBERT_BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50496
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50507
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)	REFID=50802
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer		REFID=51038
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49739
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49611
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
BAI	04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49620
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49607
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49751
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49754
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49403
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47427
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46341
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)	REFID=46608
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45146
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44736
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45198
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)	REFID=45197
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44434
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)	REFID=43314
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)	REFID=43601
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41866
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)	REFID=41899
BISELLO	91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41359

COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)	REFID=41349
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=40346
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)	REFID=40002
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
HENRRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40261
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)	REFID=40243
BALTRUSAIT...	86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BALTRUSAIT... 86B	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT... 86D	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)	REFID=21865
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=22101
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
BALTRUSAIT... 85C	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095
BALTRUSAIT... 85D	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22097
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
BALTRUSAIT... 84	84	PRL 52 2126	Translated from YAF 41 733		
EATON	84	PR D29 804	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22006
BLOOM	83	ARNS 33 143	M.W. Eaton <i>et al.</i>	(LBL, SLAC)	REFID=22092
EDWARDS	83B	PRL 51 859	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
FRANKLIN	83	PRL 51 963	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
BURKE	82	PRL 49 632	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
EDWARDS	82B	PR D25 3065	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22080
Also		ARNS 33 143	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
EDWARDS	82E	PRL 49 259	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
LEMOIGNE	82	PL 113B 509	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
BESCH	81	ZPHY C8 1	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
GIDAL	81	PL 107B 153	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)	REFID=22077
PARTRIDGE	80	PRL 44 712	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
SCHARRE	80	PL 97B 329	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22073
ZHOLENTZ	80	PL 96B 214	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
BRANDELIK	79C	ZPHY C1 233	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
ALEXANDER	78	PL 72B 493	Translated from YAF 34 1471		
BESCH	78	PL 78B 347	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
BRANDELIK	78B	PL 74B 292	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22065
PERUZZI	78	PR D17 2901	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)	REFID=22066
BARTEL	77	PL 66B 489	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22067
BURMESTER	77D	PL 72B 135	I. Peruzzi <i>et al.</i>	(SLAC, LBL)	REFID=22068
FELDMAN	77	PRPL 33C 285	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22058
VANNUCCI	77	PR D15 1814	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22060
BARTEL	76	PL 64B 483	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
BRAUNSCH...	76	PL 63B 487	F. Vannucci <i>et al.</i>	(SLAC, LBL)	REFID=22063
JEAN-MARIE	76	PRL 36 291	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
BALDINI-...	75	PL 58B 471	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22054
BOYARSKI	75	PRL 34 1357	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG	REFID=22056
DASP	75	PL 56B 491	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)	REFID=22026
ESPOSITO	75B	LNC 14 73	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC	REFID=22030
FORD	75	PRL 34 604	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22036
			B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=22038
			R.L. Ford <i>et al.</i>	(SLAC, PENN)	REFID=22039

## BRANCHING RATIOS OF $\psi(2S)$ AND $\chi_{c0,1,2}$

Updated March 2022 by J.J. Hernández-Rey (IFIC, Valencia), S. Navas (U. of Granada), and C. Patrignani (Bologna Univ., INFN)

Since 2002, the treatment of the branching ratios of the  $\psi(2S)$  and  $\chi_{c0,1,2}$  has undergone an important restructuring.

When measuring a branching ratio experimentally, it is not always possible to normalize the number of events observed in the corresponding decay mode to the total number of particles produced. Therefore, the experimenters sometimes report the number of observed decays with respect to another decay mode of the same or another particle in the relevant decay chain. This is actually equivalent to measuring combinations of branching fractions of several decay modes.

To extract the branching ratio of a given decay mode, the collaborations use some previously reported measurements of the required branching ratios. However, the values are frequently taken from the *Review of Particle Physics* (RPP), which in turn uses the branching ratio reported by the experiment in the following edition, giving rise either to correlations or to plain vicious circles, as discussed in more detail in earlier editions of this review [1,2].

The way to avoid these dependencies and correlations is to extract the branching ratios through a fit that uses the truly measured combinations of branching fractions and partial widths. This fit, in fact, should involve decays from the four concerned particles,  $\psi(2S)$ ,  $\chi_{c0}$ ,  $\chi_{c1}$ , and  $\chi_{c2}$ , and occasionally some combinations of branching ratios of more than one of them. This is what is done since the 2002 edition [3].

The PDG policy is to quote the results of the collaborations in a manner as close as possible to what appears in their original publications. However, in order to avoid the problems mentioned above, we had in some cases to work out the values originally measured, using the number of events and detection efficiencies given by the collaborations, or rescaling back the published results. The information was sometimes spread over several articles, and some articles referred to papers still unpublished, which in turn contained the relevant numbers in footnotes.

Even though the experimental collaborations are entitled to extract whatever branching ratios they consider appropriate by using other published results, we would like to encourage them to also quote explicitly in their articles the actual quantities

measured, so that they can be used directly in averages and fits of different experimental determinations.

To inform the reader how we computed some of the values used in this edition of RPP, we use footnotes to indicate the branching ratios actually given by the experiments and the quantities they use to derive them from the true combination of branching ratios actually measured.

None of the branching ratios of the  $\chi_{c0,1,2}$  are measured independently of the  $\psi(2S)$  radiative decays. We tried to identify those branching ratios which can be correlated in a non-trivial way, and although we cannot preclude the existence of other cases, we are confident that the most relevant correlations have already been removed. Nevertheless, correlations in the errors of different quantities measured by the same experiment have not been taken into account.

### Fit information

This is an overall fit to 4 total widths, 1 partial width, 26 combinations of partial widths, 24 branching ratios, and 108 combinations of branching ratios. Of the latter 62 involve decays of more than one particle.

The overall fit uses 248 measurements to determine 49 parameters and has a  $\chi^2$  of 379.8 for 199 degrees of freedom.

The relatively high  $\chi^2$  of the fit, 1.9 per d.o.f., can be traced back to a few specific discrepancies in the data. No scaling factors to fit uncertainties have been applied.

In the listing we provide the inter-particle correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the corresponding parameter  $x_i$ .

### References

1. Y.F. Gu and X.H. Li, Phys. Lett. **B449**, 361 (1999).
  2. C. Patrignani, Phys. Rev. **D64**, 034017 (2001).
  3. K. Hagiwara *et al.* (Particle Data Group), Phys. Rev. **D68**, 010001 (2002).
-



$\chi_{c0}(1P)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

NODE=M056

 $\chi_{c0}(1P)$  MASS

NODE=M056M

NODE=M056M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3414.71 ± 0.30 OUR AVERAGE</b>				
3413.0 ± 1.9 ± 0.6	933	<sup>1</sup> AAIJ	17BB LHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3414.2 ± 0.5 ± 2.3	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
3406 ± 7 ± 6	230	<sup>2</sup> ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$
3414.21 ± 0.39 ± 0.27		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
3414.7 $\begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}$ ± 0.2		<sup>3</sup> ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
3415.5 ± 0.4 ± 0.4	392	<sup>4</sup> BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
3417.4 $\begin{smallmatrix} +1.8 \\ -1.9 \end{smallmatrix}$ ± 0.2		<sup>3</sup> AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
3414.1 ± 0.6 ± 0.8		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3417.8 ± 0.4 ± 4		<sup>3</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3416 ± 3 ± 4		<sup>5</sup> TANENBAUM	78 MRK1	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3414.6 ± 1.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
3416.5 ± 3.0		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c0}$
3422 ± 10		<sup>5</sup> BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3415 ± 9		<sup>5</sup> BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$

<sup>1</sup> From a fit of the  $\phi\phi$  invariant mass with the width of  $\chi_{c0}(1P)$  fixed to the PDG 16 value.

<sup>2</sup> From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>3</sup> Using mass of  $\psi(2S) = 3686.0$  MeV.

<sup>4</sup> Recalculated by ANDREOTTI 05A, using the value of  $\psi(2S)$  mass from AULCHENKO 03.

<sup>5</sup> Mass value shifted by us by amount appropriate for  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.

NODE=M056M;LINKAGE=A

NODE=M056M;LINKAGE=EB

NODE=M056M;LINKAGE=C

NODE=M056M;LINKAGE=NW

NODE=M056M;LINKAGE=D

 $\chi_{c0}(1P)$  WIDTH

NODE=M056W

NODE=M056W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.8 ± 0.6 OUR FIT</b>				
<b>10.5 ± 0.8 OUR AVERAGE</b> Error includes scale factor of 1.1.				
10.6 ± 1.9 ± 2.6	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$
12.6 $\begin{smallmatrix} +1.5+0.9 \\ -1.6-1.1 \end{smallmatrix}$		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
8.6 $\begin{smallmatrix} +1.7 \\ -1.3 \end{smallmatrix}$ ± 0.1		ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
9.7 ± 1.0	392	<sup>1</sup> BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
16.6 $\begin{smallmatrix} +5.2 \\ -3.7 \end{smallmatrix}$ ± 0.1		AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
14.3 ± 2.0 ± 3.0		BAI	98I BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
13.5 ± 3.3 ± 4.2		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X, \gamma\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
13.2 ± 2.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> Recalculated by ANDREOTTI 05A.

NODE=M056W;LINKAGE=AN

 $\chi_{c0}(1P)$  DECAY MODES

NODE=M056215;NODE=M056

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
<b>Hadronic decays</b>		
$\Gamma_1$ $2(\pi^+\pi^-)$	(2.34 ± 0.18) %	
$\Gamma_2$ $\rho^0\pi^+\pi^-$	(9.1 ± 2.9) × 10 <sup>-3</sup>	
$\Gamma_3$ $\rho^0\rho^0$		
$\Gamma_4$ $f_0(980)f_0(980)$	(6.6 ± 2.1) × 10 <sup>-4</sup>	
$\Gamma_5$ $\pi^+\pi^-\pi^0\pi^0$	(3.3 ± 0.4) %	
$\Gamma_6$ $\rho^+\pi^-\pi^0 + \text{c.c.}$	(2.9 ± 0.4) %	
$\Gamma_7$ $4\pi^0$	(3.3 ± 0.4) × 10 <sup>-3</sup>	
$\Gamma_8$ $\pi^+\pi^-K^+K^-$	(1.81 ± 0.14) %	

NODE=M056;CLUMP=A

DESIG=3

DESIG=9

DESIG=54

DESIG=20

DESIG=61

DESIG=62

DESIG=70

DESIG=5

$\Gamma_9$	$K_0^*(1430)^0 \bar{K}_0^*(1430)^0 \rightarrow \pi^+ \pi^- K^+ K^-$	$(9.8^{+4.0}_{-2.8}) \times 10^{-4}$		DESIG=31
$\Gamma_{10}$	$K_0^*(1430)^0 \bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(8.0^{+2.0}_{-2.4}) \times 10^{-4}$		DESIG=32
$\Gamma_{11}$	$K_1(1270)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$(6.3 \pm 1.9) \times 10^{-3}$		DESIG=33
$\Gamma_{12}$	$K_1(1400)^+ K^- + \text{c.c.} \rightarrow \pi^+ \pi^- K^+ K^-$	$< 2.7 \times 10^{-3}$	CL=90%	DESIG=34
$\Gamma_{13}$	$f_0(980) f_0(980)$	$(1.6^{+1.0}_{-0.9}) \times 10^{-4}$		DESIG=23
$\Gamma_{14}$	$f_0(980) f_0(2200)$	$(7.9^{+2.0}_{-2.5}) \times 10^{-4}$		DESIG=24
$\Gamma_{15}$	$f_0(1370) f_0(1370)$	$< 2.7 \times 10^{-4}$	CL=90%	DESIG=25
$\Gamma_{16}$	$f_0(1370) f_0(1500)$	$< 1.7 \times 10^{-4}$	CL=90%	DESIG=26
$\Gamma_{17}$	$f_0(1370) f_0(1710)$	$(6.7^{+3.5}_{-2.3}) \times 10^{-4}$		DESIG=27
$\Gamma_{18}$	$f_0(1500) f_0(1370)$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=28
$\Gamma_{19}$	$f_0(1500) f_0(1500)$	$< 5 \times 10^{-5}$	CL=90%	DESIG=29
$\Gamma_{20}$	$f_0(1500) f_0(1710)$	$< 7 \times 10^{-5}$	CL=90%	DESIG=30
$\Gamma_{21}$	$K^+ K^- \pi^+ \pi^- \pi^0$	$(8.6 \pm 0.9) \times 10^{-3}$		DESIG=75
$\Gamma_{22}$	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(4.2 \pm 0.4) \times 10^{-3}$		DESIG=87
$\Gamma_{23}$	$K^+ K^- \pi^0 \pi^0$	$(5.6 \pm 0.9) \times 10^{-3}$		DESIG=63
$\Gamma_{24}$	$K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(2.49 \pm 0.33) \%$		DESIG=65
$\Gamma_{25}$	$\rho^+ K^- K^0 + \text{c.c.}$	$(1.21 \pm 0.21) \%$		DESIG=66
$\Gamma_{26}$	$K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(4.6 \pm 1.2) \times 10^{-3}$		DESIG=67
$\Gamma_{27}$	$K_S^0 K_S^0 \pi^+ \pi^-$	$(5.7 \pm 1.1) \times 10^{-3}$		DESIG=41
$\Gamma_{28}$	$K^+ K^- \eta \pi^0$	$(3.0 \pm 0.7) \times 10^{-3}$		DESIG=68
$\Gamma_{29}$	$3(\pi^+ \pi^-)$	$(1.20 \pm 0.18) \%$		DESIG=4
$\Gamma_{30}$	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(7.5 \pm 1.6) \times 10^{-3}$		DESIG=10
$\Gamma_{31}$	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.7 \pm 0.6) \times 10^{-3}$		DESIG=21
$\Gamma_{32}$	$\pi \pi$	$(8.51 \pm 0.33) \times 10^{-3}$		DESIG=18
$\Gamma_{33}$	$\pi^0 \eta$	$< 1.8 \times 10^{-4}$		DESIG=35
$\Gamma_{34}$	$\pi^0 \eta'$	$< 1.1 \times 10^{-3}$		DESIG=36
$\Gamma_{35}$	$\pi^0 \eta_c$	$< 1.6 \times 10^{-3}$	CL=90%	DESIG=86
$\Gamma_{36}$	$\eta \eta$	$(3.01 \pm 0.19) \times 10^{-3}$		DESIG=13
$\Gamma_{37}$	$\eta \eta'$	$(9.1 \pm 1.1) \times 10^{-5}$		DESIG=37
$\Gamma_{38}$	$\eta' \eta'$	$(2.17 \pm 0.12) \times 10^{-3}$		DESIG=46
$\Gamma_{39}$	$\omega \omega$	$(9.7 \pm 1.1) \times 10^{-4}$		DESIG=22
$\Gamma_{40}$	$\omega \phi$	$(1.41 \pm 0.13) \times 10^{-4}$		DESIG=76
$\Gamma_{41}$	$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$		DESIG=88
$\Gamma_{42}$	$K^+ K^-$	$(6.05 \pm 0.31) \times 10^{-3}$		DESIG=2
$\Gamma_{43}$	$K_S^0 K_S^0$	$(3.16 \pm 0.17) \times 10^{-3}$		DESIG=15
$\Gamma_{44}$	$\pi^+ \pi^- \eta$	$< 2.0 \times 10^{-4}$	CL=90%	DESIG=50
$\Gamma_{45}$	$\pi^+ \pi^- \eta'$	$< 4 \times 10^{-4}$	CL=90%	DESIG=53
$\Gamma_{46}$	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$< 9 \times 10^{-5}$	CL=90%	DESIG=17
$\Gamma_{47}$	$K^+ K^- \pi^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=47
$\Gamma_{48}$	$K^+ K^- \eta$	$< 2.3 \times 10^{-4}$	CL=90%	DESIG=51
$\Gamma_{49}$	$K^+ K^- K_S^0 K_S^0$	$(1.4 \pm 0.5) \times 10^{-3}$		DESIG=42
$\Gamma_{50}$	$K_S^0 K_S^0 K_S^0 K_S^0$	$(5.8 \pm 0.5) \times 10^{-4}$		DESIG=94
$\Gamma_{51}$	$K^+ K^- K^+ K^-$	$(2.82 \pm 0.29) \times 10^{-3}$		DESIG=14
$\Gamma_{52}$	$K^+ K^- \phi$	$(9.7 \pm 2.5) \times 10^{-4}$		DESIG=44
$\Gamma_{53}$	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.7 \pm 0.6) \times 10^{-3}$		DESIG=91
$\Gamma_{54}$	$K^+ K^- \pi^0 \phi$	$(1.90 \pm 0.35) \times 10^{-3}$		DESIG=92
$\Gamma_{55}$	$\phi \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.15) \times 10^{-3}$		DESIG=89
$\Gamma_{56}$	$\phi \phi$	$(8.0 \pm 0.7) \times 10^{-4}$		DESIG=16
$\Gamma_{57}$	$\phi \phi \eta$	$(8.4 \pm 1.0) \times 10^{-4}$		DESIG=96
$\Gamma_{58}$	$\rho \bar{\rho}$	$(2.21 \pm 0.08) \times 10^{-4}$		DESIG=11
$\Gamma_{59}$	$\rho \bar{\rho} \pi^0$	$(7.0 \pm 0.7) \times 10^{-4}$	S=1.3	DESIG=48

Г60	$p\bar{p}\eta$	$(3.5 \pm 0.4) \times 10^{-4}$		DESIG=52
Г61	$p\bar{p}\omega$	$(5.2 \pm 0.6) \times 10^{-4}$		DESIG=69
Г62	$p\bar{p}\phi$	$(6.0 \pm 1.4) \times 10^{-5}$		DESIG=74
Г63	$p\bar{p}\pi^+\pi^-$	$(2.1 \pm 0.7) \times 10^{-3}$	S=1.4	DESIG=8
Г64	$p\bar{p}\pi^0\pi^0$	$(1.04 \pm 0.28) \times 10^{-3}$		DESIG=64
Г65	$p\bar{p}K^+K^-$ (non-resonant)	$(1.22 \pm 0.26) \times 10^{-4}$		DESIG=71
Г66	$p\bar{p}K_S^0 K_S^0$	$< 8.8 \times 10^{-4}$	CL=90%	DESIG=40
Г67	$p\bar{n}\pi^-$	$(1.27 \pm 0.11) \times 10^{-3}$		DESIG=43
Г68	$\bar{p}n\pi^+$	$(1.37 \pm 0.12) \times 10^{-3}$		DESIG=82
Г69	$p\bar{n}\pi^-\pi^0$	$(2.34 \pm 0.21) \times 10^{-3}$		DESIG=83
Г70	$\bar{p}n\pi^+\pi^0$	$(2.21 \pm 0.18) \times 10^{-3}$		DESIG=84
Г71	$\Lambda\bar{\Lambda}$	$(3.59 \pm 0.15) \times 10^{-4}$		DESIG=19
Г72	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(1.18 \pm 0.13) \times 10^{-3}$		DESIG=38
Г73	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$< 5 \times 10^{-4}$	CL=90%	DESIG=77
Г74	$\Lambda\bar{\Lambda}\eta$	$(2.3 \pm 0.4) \times 10^{-4}$		DESIG=102
Г75	$\Sigma(1385)^+\bar{\Lambda}\pi^- + c.c.$	$< 5 \times 10^{-4}$	CL=90%	DESIG=78
Г76	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + c.c.$	$< 5 \times 10^{-4}$	CL=90%	DESIG=79
Г77	$K^+\bar{p}\Lambda + c.c.$	$(1.25 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=49
Г78	$nK_S^0\bar{\Lambda} + c.c.$	$(6.6 \pm 0.5) \times 10^{-4}$		DESIG=101
Г79	$K^*(892)^+\bar{p}\Lambda + c.c.$	$(4.8 \pm 0.9) \times 10^{-4}$		DESIG=98
Г80	$K^+\bar{p}\Lambda(1520) + c.c.$	$(2.9 \pm 0.7) \times 10^{-4}$		DESIG=72
Г81	$\Lambda(1520)\bar{\Lambda}(1520)$	$(3.1 \pm 1.2) \times 10^{-4}$		DESIG=73
Г82	$\Sigma^0\bar{\Sigma}^0$	$(4.68 \pm 0.32) \times 10^{-4}$		DESIG=58
Г83	$\Sigma^+\bar{p}K_S^0 + c.c.$	$(3.52 \pm 0.27) \times 10^{-4}$		DESIG=97
Г84	$\Sigma^0\bar{p}K^+ + c.c.$	$(3.03 \pm 0.20) \times 10^{-4}$		DESIG=100
Г85	$\Sigma^+\bar{\Sigma}^-$	$(4.6 \pm 0.8) \times 10^{-4}$	S=2.6	DESIG=59
Г86	$\Sigma^-\bar{\Sigma}^+$	$(5.1 \pm 0.5) \times 10^{-4}$		DESIG=99
Г87	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(1.6 \pm 0.6) \times 10^{-4}$		DESIG=80
Г88	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$(2.3 \pm 0.7) \times 10^{-4}$		DESIG=81
Г89	$K^-\Lambda\bar{\Xi}^+ + c.c.$	$(1.94 \pm 0.35) \times 10^{-4}$		DESIG=85
Г90	$\Xi^0\bar{\Xi}^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.7	DESIG=60
Г91	$\Xi^-\bar{\Xi}^+$	$(4.45 \pm 0.19) \times 10^{-4}$		DESIG=39
Г92	$\eta_c\pi^+\pi^-$	$< 7 \times 10^{-4}$	CL=90%	DESIG=90
<b>Radiative decays</b>				
Г93	$\gamma J/\psi(1S)$	$(1.40 \pm 0.05) \%$		NODE=M056;CLUMP=B DESIG=6
Г94	$\gamma\rho^0$	$< 9 \times 10^{-6}$	CL=90%	DESIG=55
Г95	$\gamma\omega$	$< 8 \times 10^{-6}$	CL=90%	DESIG=56
Г96	$\gamma\phi$	$< 6 \times 10^{-6}$	CL=90%	DESIG=57
Г97	$\gamma\gamma$	$(2.04 \pm 0.09) \times 10^{-4}$		DESIG=7
Г98	$e^+e^- J/\psi(1S)$	$(1.33 \pm 0.29) \times 10^{-4}$		DESIG=93
Г99	$\mu^+\mu^- J/\psi(1S)$	$< 1.9 \times 10^{-5}$	CL=90%	DESIG=95

---

**CONSTRAINED FIT INFORMATION**

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_2$	24									
$x_8$	9	2								
$x_{30}$	5	1	28							
$x_{32}$	8	2	10	3						
$x_{36}$	4	1	5	1	14					
$x_{42}$	8	2	8	3	18	11				
$x_{43}$	7	2	8	2	18	10	14			
$x_{51}$	5	1	5	2	9	5	7	7		
$x_{56}$	7	2	6	2	9	5	7	7	4	
$x_{58}$	3	1	4	1	3	-1	7	7	3	3
$x_{71}$	7	2	9	2	23	13	18	18	8	9
$x_{93}$	5	1	6	2	17	11	13	12	6	6
$x_{97}$	-8	-2	-2	-3	14	9	10	10	3	1
$\Gamma$	-26	-6	-19	-10	-15	-7	-14	-12	-10	-13
	$x_1$	$x_2$	$x_8$	$x_{30}$	$x_{32}$	$x_{36}$	$x_{42}$	$x_{43}$	$x_{51}$	$x_{56}$
$x_{71}$	9									
$x_{93}$	-19	16								
$x_{97}$	6	15	13							
$\Gamma$	-4	-13	-9	-38						
	$x_{58}$	$x_{71}$	$x_{93}$	$x_{97}$						

### $\chi_{c0}(1P)$ PARTIAL WIDTHS

NODE=M056217

$$\chi_{c0}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$$

NODE=M056223

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{58} \Gamma_{93} / \Gamma$$

NODE=M056G1

NODE=M056G1

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**33.6 ± 2.3 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

26.6 ± 2.6 ± 1.4	392	1,2 BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi \gamma$
------------------	-----	--------------	---------	--

48.7 <sup>+11.3</sup> <sub>-8.9</sub> ± 2.4		1,2 AMBROGIANI	99B E835	$\bar{p}p \rightarrow \gamma J/\psi$
---	--	----------------	----------	--------------------------------------

<sup>1</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .

NODE=M056G;LINKAGE=7A

<sup>2</sup> Values in  $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$  and  $(\Gamma(p\bar{p}) / \Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}})$  are not independent. The latter is used in the fit since it is less correlated to the total width.

NODE=M056G;LINKAGE=KS

$$\chi_{c0}(1P) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$$

NODE=M056224

$$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}} \quad \Gamma_1 \Gamma_{97} / \Gamma$$

NODE=M056G2

NODE=M056G2

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**52 ± 4 OUR FIT****49 ± 10 OUR AVERAGE** Error includes scale factor of 1.8.

44.7 ± 3.6 ± 4.9	3.6k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$
------------------	------	--------	---------	---

75 ± 13 ± 8		EISENSTEIN	01 CLE2	$e^+ e^- \rightarrow e^+ e^- \chi_{c0}$
-------------	--	------------	---------	---

$$\Gamma(\rho^0 \rho^0) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}} \quad \Gamma_3 \Gamma_{97} / \Gamma$$

NODE=M056G07

NODE=M056G07

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	<252	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+ \pi^-)$
-----	----	------	--------	---------	---

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{8}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**40.0±3.5 OUR FIT**

38.8±3.7±4.7	1.7k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$
--------------	------	--------	----	--

NODE=M056G08  
NODE=M056G08

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{21}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**26±4±4**

1094		DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
------	--	-----------------	------	--

NODE=M056G01  
NODE=M056G01

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**16 ±4 OUR FIT**

16.7±6.1±3.0	495 ± 182	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$
--------------	-----------	--------	----	--

NODE=M056G09  
NODE=M056G09

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31}\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90	<148	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$
----	----	------	--------	----	--

NODE=M056G10  
NODE=M056G10

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**18.8± 1.3 OUR FIT**

**23 ± 5 OUR AVERAGE**

29.7 <sup>+17.4</sup> <sub>-12.0</sub> ± 4.8	103 <sup>+60</sup> <sub>-42</sub>	<sup>1</sup> UEHARA	09	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
--	-----------------------------------	---------------------	----	------	--

22.7 ± 3.2 ± 3.5	129 ± 18	<sup>2</sup> NAKAZAWA	05	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
------------------	----------	-----------------------	----	------	--

<sup>1</sup>We multiplied the measurement by 3 to convert from  $\pi^0\pi^0$  to  $\pi\pi$ . Interference with the continuum included.

<sup>2</sup>We have multiplied  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

NODE=M056G3  
NODE=M056G3

NODE=M056G3;LINKAGE=UE

NODE=M056G;LINKAGE=NA

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**9.4±2.3±1.2**

22		<sup>1</sup> UEHARA	10A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
----	--	---------------------	-----	------	--

<sup>1</sup>Interference with the continuum not included.

NODE=M056G06  
NODE=M056G06

NODE=M056G06;LINKAGE=UE

$$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{39}\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.9	90	<sup>1</sup> LIU	12B	BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$
------	----	------------------	-----	------	---

<sup>1</sup>Using  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M056G02  
NODE=M056G02

NODE=M056G02;LINKAGE=LI

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{40}\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.34	90	<sup>1</sup> LIU	12B	BELL	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
-------	----	------------------	-----	------	--

<sup>1</sup>Using  $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$  and  $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M056G03  
NODE=M056G03

NODE=M056G03;LINKAGE=LI

$$\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**13.4±1.0 OUR FIT**

14.3±1.6±2.3	153 ± 17	NAKAZAWA	05	BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
--------------	----------	----------	----	------	--

NODE=M056G4  
NODE=M056G4

$$\Gamma(K_S^0K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{43}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**7.0 ±0.5 OUR FIT**

8.7 ±1.7 ±0.9	266	<sup>1</sup> UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0K_S^0$
---------------	-----	---------------------	----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.00±0.65±0.71	134 ± 12	CHEN	07B	BELL	$e^+e^- \rightarrow e^+e^-\chi_{c0}$
----------------	----------	------	-----	------	--------------------------------------

<sup>1</sup>Supersedes CHEN 07B.

NODE=M056G5  
NODE=M056G5

NODE=M056G5;LINKAGE=UE

$$\Gamma(K^+K^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**6.2±0.7 OUR FIT**

7.9±1.3±1.1	215 ± 36	UEHARA	08	BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+K^-)$
-------------	----------	--------	----	------	--

NODE=M056G11  
NODE=M056G11

$$\frac{\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}}{\Gamma_{56}\Gamma_{97}/\Gamma}$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

**1.76±0.18 OUR FIT**

**1.72±0.33±0.14** 56 ± 11 <sup>1</sup> LIU 12B BELL  $\gamma\gamma \rightarrow 2(K^+K^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ± 0.9 ± 0.4 23.6 ± 9.6 UEHARA 08 BELL  $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+K^-)$

<sup>1</sup>Supersedes UEHARA 08. Using  $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ .

NODE=M056G12  
NODE=M056G12

NODE=M056G12;LINKAGE=LI

## $\chi_{c0}(1P)$ BRANCHING RATIOS

NODE=M056220

### HADRONIC DECAYS

NODE=M056305

$$\frac{\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}}{\Gamma_1/\Gamma}$$

VALUE	DOCUMENT ID
-------	-------------

**0.0234±0.0018 OUR FIT**

NODE=M056R2  
NODE=M056R2

$$\frac{\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))}{\Gamma_2/\Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.39±0.12 OUR FIT**

**0.39±0.12**

TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c0}$

NODE=M056R54  
NODE=M056R54

$$\frac{\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}}{\Gamma_2/\Gamma}$$

VALUE	DOCUMENT ID
-------	-------------

**0.0091±0.0029 OUR FIT**

NODE=M056R9  
NODE=M056R9

$$\frac{\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}}{\Gamma_4/\Gamma}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**6.6±2.1±0.1** 36 ± 9 <sup>1</sup> ABLIKIM 04G BES  $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

<sup>1</sup>ABLIKIM 04G reports  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (6.5 \pm 1.6 \pm 1.3) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R24  
NODE=M056R24

NODE=M056R24;LINKAGE=AB

$$\frac{\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}}{\Gamma_5/\Gamma}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

**3.3±0.4±0.1** 1751.4 <sup>1</sup> HE 08B CLEO  $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup>HE 08B reports  $3.54 \pm 0.10 \pm 0.43 \pm 0.18$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R62  
NODE=M056R62

NODE=M056R62;LINKAGE=HE

$$\frac{\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}}{\Gamma_6/\Gamma}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

**2.9±0.4±0.1** 1358.5 <sup>1,2</sup> HE 08B CLEO  $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup>HE 08B reports  $3.04 \pm 0.18 \pm 0.42 \pm 0.16$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup>Calculated by us. We have added the values from HE 08B for  $\rho^+\pi^-\pi^0$  and  $\rho^-\pi^+\pi^0$  decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M056R63  
NODE=M056R63

NODE=M056R63;LINKAGE=HE

NODE=M056R63;LINKAGE=OC

$$\frac{\Gamma(4\pi^0)/\Gamma_{\text{total}}}{\Gamma_7/\Gamma}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.3±0.4±0.1** 3296 <sup>1</sup> ABLIKIM 11A BES3  $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

<sup>1</sup>ABLIKIM 11A reports  $(3.34 \pm 0.06 \pm 0.44) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R71  
NODE=M056R71

NODE=M056R71;LINKAGE=AB

$$\frac{\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}}{\Gamma_8/\Gamma}$$

VALUE (units $10^{-3}$ )	DOCUMENT ID
--------------------------	-------------

**18.1±1.4 OUR FIT**

NODE=M056R3  
NODE=M056R3

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma(\pi^+\pi^-K^+K^-) \quad \Gamma_{30}/\Gamma_8$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.41±0.09 OUR FIT**

**0.41±0.10**

TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c0}$

NODE=M056R55  
NODE=M056R55

$$\Gamma(K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**9.8<sup>+3.6</sup><sub>-2.8</sub>±0.2**      83      <sup>1</sup> ABLIKIM      05Q      BES2       $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R36  
NODE=M056R36

<sup>1</sup> ABLIKIM 05Q reports  $(10.44 \pm 2.37^{+3.05}_{-1.90}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R36;LINKAGE=AB

$$\Gamma(K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**8.0<sup>+2.0</sup><sub>-2.4</sub>±0.2**      62      <sup>1</sup> ABLIKIM      05Q      BES2       $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R37  
NODE=M056R37

<sup>1</sup> ABLIKIM 05Q reports  $(8.49 \pm 1.66^{+1.32}_{-1.99}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R37;LINKAGE=AB

$$\Gamma(K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**6.3±1.9±0.1**      68      <sup>1</sup> ABLIKIM      05Q      BES2       $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R38  
NODE=M056R38

<sup>1</sup> ABLIKIM 05Q reports  $(6.66 \pm 1.31^{+1.60}_{-1.51}) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The measurement assumes  $B(K_1(1270) \rightarrow K\rho(770)) = 42 \pm 6\%$ .

NODE=M056R38;LINKAGE=AB

$$\Gamma(K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

**<2.7**      90      <sup>1</sup> ABLIKIM      05Q      BES2       $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R39  
NODE=M056R39

<sup>1</sup> ABLIKIM 05Q reports  $< 2.85 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The measurement assumes  $B(K_1(1400) \rightarrow K^*(892)\pi) = 94 \pm 6\%$ .

NODE=M056R39;LINKAGE=AB

$$\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**16.2<sup>+10.4</sup><sub>-9.0</sub>±0.3**      28      <sup>1</sup> ABLIKIM      05Q      BES2       $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R28  
NODE=M056R28

<sup>1</sup> ABLIKIM 05Q reports  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.59 \pm 0.50^{+0.89}_{-0.72}) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. One of the  $f_0(980)$  mesons is identified via decay to  $\pi^+\pi^-$  while the other via  $K^+K^-$  decay.

NODE=M056R28;LINKAGE=AB

$$\Gamma(f_0(980)f_0(2200))/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**7.9<sup>+2.0</sup><sub>-2.5</sub>±0.2**      77      <sup>1</sup> ABLIKIM      05Q      BES2       $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R29  
NODE=M056R29

<sup>1</sup> ABLIKIM 05Q reports  $(8.42 \pm 1.42^{+1.65}_{-2.29}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(2200))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The  $f_0$  mesons are identified via  $f_0(980) \rightarrow \pi^+\pi^-$  and  $f_0(2200) \rightarrow K^+K^-$  decays.

NODE=M056R29;LINKAGE=AB

### $\Gamma(f_0(1370)f_0(1370))/\Gamma_{\text{total}}$ $\Gamma_{15}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	<sup>1</sup> ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R30  
NODE=M056R30

<sup>1</sup> ABLIKIM 05Q reports  $< 2.9 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . One of the  $f_0(1370)$  mesons is identified via decay to  $\pi^+\pi^-$  while the other via  $K^+K^-$  decay. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

NODE=M056R30;LINKAGE=AB

### $\Gamma(f_0(1370)f_0(1500))/\Gamma_{\text{total}}$ $\Gamma_{16}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	<sup>1</sup> ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R31  
NODE=M056R31

<sup>1</sup> ABLIKIM 05Q reports  $< 1.8 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The  $f_0$  mesons are identified via  $f_0(1370) \rightarrow \pi^+\pi^-$  and  $f_0(1500) \rightarrow K^+K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

NODE=M056R31;LINKAGE=AB

### $\Gamma(f_0(1370)f_0(1710))/\Gamma_{\text{total}}$ $\Gamma_{17}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.7^{+3.5}_{-2.3} \pm 0.1$	61	<sup>1</sup> ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R32  
NODE=M056R32

<sup>1</sup> ABLIKIM 05Q reports  $(7.12 \pm 1.85^{+3.28}_{-1.68}) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The  $f_0$  mesons are identified via  $f_0(1370) \rightarrow \pi^+\pi^-$  and  $f_0(1710) \rightarrow K^+K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

NODE=M056R32;LINKAGE=AB

### $\Gamma(f_0(1500)f_0(1370))/\Gamma_{\text{total}}$ $\Gamma_{18}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	<sup>1</sup> ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R33  
NODE=M056R33

<sup>1</sup> ABLIKIM 05Q reports  $< 1.4 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The  $f_0$  mesons are identified via  $f_0(1500) \rightarrow \pi^+\pi^-$  and  $f_0(1370) \rightarrow K^+K^-$  decays. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

NODE=M056R33;LINKAGE=AB

### $\Gamma(f_0(1500)f_0(1500))/\Gamma_{\text{total}}$ $\Gamma_{19}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	<sup>1</sup> ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R34  
NODE=M056R34

<sup>1</sup> ABLIKIM 05Q reports  $< 0.55 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . One of the  $f_0(1500)$  is identified via decay to  $\pi^+\pi^-$  while the other via  $K^+K^-$  decay. Both branching fractions for these  $f_0$  decays are implicitly included in the quoted result.

NODE=M056R34;LINKAGE=AB



$\Gamma(f_0(1500) f_0(1710))/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.7</b>	90	<sup>1</sup> ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
<sup>1</sup> ABLIKIM 05Q reports $< 0.73 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500) f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ . The $f_0$ mesons are identified via $f_0(1500) \rightarrow \pi^+ \pi^-$ and $f_0(1710) \rightarrow K^+ K^-$ decays. Both branching fractions for these $f_0$ decays are implicitly included in the quoted result.				

NODE=M056R35  
NODE=M056R35

NODE=M056R35;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.61±0.13±0.94</b>	9.0k	<sup>1</sup> ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
<sup>1</sup> Using $1.06 \times 10^8$ $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$ .				

NODE=M056R85  
NODE=M056R85

NODE=M056R85;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.22±0.10±0.43</b>	2.7k	<sup>1</sup> ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
<sup>1</sup> Using $1.06 \times 10^8$ $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$ .				

NODE=M056R86  
NODE=M056R86

NODE=M056R86;LINKAGE=A

 $\Gamma(K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.56±0.09±0.01</b>	213.5	<sup>1</sup> HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
<sup>1</sup> HE 08B reports $0.59 \pm 0.05 \pm 0.08 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R64  
NODE=M056R64

NODE=M056R64;LINKAGE=HE

 $\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.49±0.33±0.05</b>	401.7	<sup>1</sup> HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
<sup>1</sup> HE 08B reports $2.64 \pm 0.15 \pm 0.31 \pm 0.14$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R66  
NODE=M056R66

NODE=M056R66;LINKAGE=HE

 $\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.21±0.21±0.02</b>	179.7	<sup>1</sup> HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
<sup>1</sup> HE 08B reports $1.28 \pm 0.16 \pm 0.15 \pm 0.07$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R67  
NODE=M056R67

NODE=M056R67;LINKAGE=HE

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.46±0.12±0.01</b>	64.1	<sup>1</sup> HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
<sup>1</sup> HE 08B reports $0.49 \pm 0.10 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R68  
NODE=M056R68

NODE=M056R68;LINKAGE=HE

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.7±1.0±0.1</b>	152 ± 14	<sup>1</sup> ABLIKIM	05o	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$
<sup>1</sup> ABLIKIM 05o reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(0.558 \pm 0.051 \pm 0.089) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R47  
NODE=M056R47

NODE=M056R47;LINKAGE=AB

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{28}/\Gamma$
<b>0.30±0.07±0.01</b>	56.4	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	NODE=M056R69 NODE=M056R69

<sup>1</sup> HE 08B reports  $0.32 \pm 0.05 \pm 0.05 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R69;LINKAGE=HE

 $\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{29}/\Gamma$
--------------------------	------	-------------	------	---------	----------------------

**12.0±1.8 OUR EVALUATION** Treating systematic error as correlated.  
**19.6±2.1 OUR AVERAGE** Error includes scale factor of 3.3. See the ideogram below.  
 [(12.0 ± 1.7) × 10<sup>-3</sup> OUR 2022 AVERAGE]

20.8±0.1±0.7	145K	<sup>1</sup> ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$	
11.7±1.0±1.9		<sup>2</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$	
12.5±2.9±0.5		<sup>2</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$	


<sup>1</sup> ABLIKIM 22Q reports  $(2.080 \pm 0.006 \pm 0.068) \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$ .

<sup>2</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$ .

NODE=M056R4  
NODE=M056R4→ UNCHECKED ←  
NEW

NODE=M056R4;LINKAGE=A

NODE=M056R;LINKAGE=X1

WEIGHTED AVERAGE  
19.6±2.1 (Error scaled by 3.3) $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + c.c.)/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	$\Gamma_{30}/\Gamma$
<b>0.0075±0.0016 OUR FIT</b>		NODE=M056R10 NODE=M056R10

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{31}/\Gamma$
--------------------------	------	-------------	------	---------	----------------------

**1.72<sup>+0.60</sup><sub>-0.54</sub>±0.04** 64 <sup>1</sup> ABLIKIM 05Q BES2  $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.56±0.40±0.03 30 ± 6 <sup>2,3</sup> ABLIKIM 04H BES Repl. by ABLIKIM 05Q

<sup>1</sup> ABLIKIM 05Q reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  =  $(0.168 \pm 0.035^{+0.047}_{-0.040}) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Assumes  $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$ .

<sup>3</sup> ABLIKIM 04H reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  =  $(1.53 \pm 0.29 \pm 0.26) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R26  
NODE=M056R26

NODE=M056R26;LINKAGE=A1

NODE=M056R;LINKAGE=AL  
NODE=M056R26;LINKAGE=AB

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**8.51±0.33 OUR FIT**

DOCUMENT ID

 $\Gamma_{32}/\Gamma$ NODE=M056R22  
NODE=M056R22 $\Gamma(\pi^0\eta_c)/\Gamma_{\text{total}}$ 

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

**<1.6 × 10<sup>-3</sup>**

90

<sup>1</sup> ABLIKIM 15N BES3  $\psi(2S)e^+e^- \rightarrow \gamma\pi^0\eta_c$ <sup>1</sup> Using  $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma\gamma) = (1.66 \pm 0.11) \times 10^{-2}$ . $\Gamma_{35}/\Gamma$ NODE=M056R00  
NODE=M056R00

NODE=M056R00;LINKAGE=A

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**3.01±0.19 OUR FIT**

DOCUMENT ID

 $\Gamma_{36}/\Gamma$ NODE=M056R13  
NODE=M056R13 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$ 

VALUE

**0.353±0.025 OUR FIT**

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{36}/\Gamma_{32}$ NODE=M056R20  
NODE=M056R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.09 <sup>+0.03</sup><sub>-0.02</sub><sup>1</sup> ANDREOTTI 05C E835  $\bar{p}p \rightarrow 2 \text{ mesons}$ 

0.24 ± 0.10 ± 0.08

<sup>1</sup> BAI 03C BES  $\psi(2S) \rightarrow 5\gamma$ <sup>1</sup> We have multiplied  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

NODE=M056R;LINKAGE=D1

 $\Gamma(\eta\eta')/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )

CL%

EVTS

DOCUMENT ID

TECN

COMMENT

**9.1±1.1±0.2**

85

<sup>1</sup> ABLIKIM 17AI BES3  $\psi(2S) \rightarrow \gamma\eta'\eta$  $\Gamma_{37}/\Gamma$ NODE=M056R03  
NODE=M056R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt;24

90

35 ± 13

<sup>2</sup> ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma\eta'\eta$ 

&lt;50

90

<sup>3</sup> ADAMS 07 CLEO  $\psi(2S) \rightarrow \gamma\chi_{c0}$ <sup>1</sup> ABLIKIM 17AI reports  $(8.92 \pm 0.84 \pm 0.65) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R03;LINKAGE=A

<sup>2</sup> ASNER 09 reports  $< 0.25 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R03;LINKAGE=AS

<sup>3</sup> Superseded by ASNER 09. ADAMS 07 reports  $< 0.5 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R03;LINKAGE=AD

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**2.17±0.12 OUR AVERAGE**

EVTS

DOCUMENT ID

TECN

COMMENT

2.23±0.13±0.05

2.5k

<sup>1</sup> ABLIKIM 17AI BES3  $\psi(2S) \rightarrow \gamma\eta'\eta'$ 

2.00±0.21±0.04

0.4k

<sup>2</sup> ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma\eta'\eta'$  $\Gamma_{38}/\Gamma$ NODE=M056R04  
NODE=M056R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.60±0.41±0.03

23

<sup>3</sup> ADAMS 07 CLEO  $\psi(2S) \rightarrow \gamma\chi_{c0}$ <sup>1</sup> ABLIKIM 17AI reports  $(2.19 \pm 0.03 \pm 0.14) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04;LINKAGE=A

<sup>2</sup> ASNER 09 reports  $(2.12 \pm 0.13 \pm 0.21) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04;LINKAGE=AS

<sup>3</sup> Superseded by ASNER 09. ADAMS 07 reports  $(1.7 \pm 0.4 \pm 0.2) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \eta'\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 0.0922 \pm 0.0011 \pm 0.0046$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04;LINKAGE=AD

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{39}/\Gamma$
<b>0.97±0.11 OUR AVERAGE</b>					
0.93±0.11±0.02	991	<sup>1</sup> ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons	
2.16±0.66±0.04	38.1±9.6	<sup>2</sup> ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma 6\pi$	

NODE=M056R27  
NODE=M056R27

<sup>1</sup> ABLIKIM 11K reports  $(0.95 \pm 0.03 \pm 0.11) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R27;LINKAGE=AL

<sup>2</sup> ABLIKIM 05N reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.212 \pm 0.053 \pm 0.037) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R27;LINKAGE=AB

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{40}/\Gamma$
<b>1.41±0.13±0.03</b>	486	<sup>1</sup> ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons	
1.18±0.22±0.02	76	<sup>2,3</sup> ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons	

NODE=M056R76  
NODE=M056R76

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 19J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (13.83 \pm 0.70 \pm 1.01) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R76;LINKAGE=A

<sup>2</sup> ABLIKIM 11K reports  $(1.2 \pm 0.1 \pm 0.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R76;LINKAGE=AL

<sup>3</sup> Superseded by ABLIKIM 19J.

NODE=M056R76;LINKAGE=B

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{41}/\Gamma$
<b>1.94±0.06±0.20</b>	1.4k	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$	

NODE=M056R87  
NODE=M056R87

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$ .

NODE=M056R87;LINKAGE=A

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	$\Gamma_{42}/\Gamma$
<b>6.05±0.31 OUR FIT</b>		

NODE=M056R6  
NODE=M056R6

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	$\Gamma_{43}/\Gamma$
<b>3.16±0.17 OUR FIT</b>		

NODE=M056R15  
NODE=M056R15

 $\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{43}/\Gamma_{32}$
<b>0.371±0.023 OUR FIT</b>				

NODE=M056R53  
NODE=M056R53

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.05 ± 0.05 <sup>1,2</sup> CHEN 07B BELL  $e^+e^- \rightarrow e^+e^-\chi_{c0}$

<sup>1</sup> Using  $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from the  $\pi^+\pi^-$  measurement of NAKAZAWA 05 rescaled by 3/2 to convert to  $\pi\pi$ .

NODE=M056R53;LINKAGE=CH

<sup>2</sup> Not independent from other measurements.

NODE=M056R53;LINKAGE=NI

 $\Gamma(K_S^0 K_S^0)/\Gamma(K^+ K^-)$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{43}/\Gamma_{42}$
<b>0.52±0.04 OUR FIT</b>				

NODE=M056R52  
NODE=M056R52

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.49±0.07±0.08 <sup>1,2</sup> CHEN 07B BELL  $e^+e^- \rightarrow e^+e^-\chi_{c0}$

<sup>1</sup> Using  $\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from NAKAZAWA 05.

NODE=M056R52;LINKAGE=CH

<sup>2</sup> Not independent from other measurements.

NODE=M056R52;LINKAGE=NI

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$  $\Gamma_{44}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt;0.20</b>	90	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------	----	--------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.0	90	<sup>2</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
------	----	----------------------	----------	---

<sup>1</sup> ATHAR 07 reports  $< 0.21 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R08  
NODE=M056R08

NODE=M056R08;LINKAGE=AT

NODE=M056R08;LINKAGE=AB

 $\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$  $\Gamma_{45}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt;0.4</b>	90	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
----------------	----	--------------------	---------	---

<sup>1</sup> ATHAR 07 reports  $< 0.38 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R51  
NODE=M056R51

NODE=M056R51;LINKAGE=AT

 $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{46}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt;0.09</b>	90	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------	----	--------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90	<sup>2,3</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
------	----	------------------------	----------	---

<0.7	90	<sup>3,4</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
------	----	--------------------	---------	---

<sup>1</sup> ATHAR 07 reports  $< 0.10 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R17  
NODE=M056R17

NODE=M056R17;LINKAGE=AT

<sup>2</sup> ABLIKIM 06R reports  $< 0.70 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R17;LINKAGE=AB

<sup>3</sup> We have multiplied the  $K_S^0 K^+ \pi^-$  measurement by a factor of 2 to convert to  $K^0 K^+ \pi^-$ .

NODE=M056R17;LINKAGE=BA

<sup>4</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$ .

NODE=M056R17;LINKAGE=X1

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{47}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt;0.06</b>	90	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------	----	--------------------	---------	---

<sup>1</sup> ATHAR 07 reports  $< 0.06 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R05  
NODE=M056R05

NODE=M056R05;LINKAGE=AT

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$  $\Gamma_{48}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt;0.23</b>	90	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------	----	--------------------	---------	---

<sup>1</sup> ATHAR 07 reports  $< 0.24 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R09  
NODE=M056R09

NODE=M056R09;LINKAGE=AT

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{49}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

<b>1.41 ± 0.47 ± 0.03</b>	16.8 ± 4.8	<sup>1</sup> ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$
---------------------------	------------	----------------------	----------	---

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.138 \pm 0.039 \pm 0.025) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R48  
NODE=M056R48

NODE=M056R48;LINKAGE=AB

$$\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-4}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**5.8±0.5±0.1**

319

<sup>1</sup> ABLIKIM 19AA BES3  $\psi(2S) \rightarrow \gamma 4K_S^0$

<sup>1</sup> Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ . ABLIKIM 19AA reports  $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (5.64 \pm 0.33 \pm 0.37) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value..

$\Gamma_{50}/\Gamma$

NODE=M056R95  
NODE=M056R95

NODE=M056R95;LINKAGE=A

$$\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

**2.82±0.29 OUR FIT**

$\Gamma_{51}/\Gamma$

NODE=M056R14  
NODE=M056R14

$$\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**0.97±0.25±0.02**

38

<sup>1</sup> ABLIKIM 06T BES2  $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> ABLIKIM 06T reports  $(1.03 \pm 0.22 \pm 0.15) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma_{52}/\Gamma$

NODE=M056R01  
NODE=M056R01

NODE=M056R01;LINKAGE=AB

$$\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**3.68±0.30±0.50**

ABLIKIM

15M

BES3

$\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma_{53}/\Gamma$

NODE=M056R90  
NODE=M056R90

$$\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**1.90±0.14±0.32**

ABLIKIM

15M

BES3

$\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma_{54}/\Gamma$

NODE=M056R91  
NODE=M056R91

$$\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**1.18±0.07±0.13**

538

<sup>1</sup> ABLIKIM 13B BES3  $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$ .

$\Gamma_{55}/\Gamma$

NODE=M056R88  
NODE=M056R88

NODE=M056R88;LINKAGE=A

$$\Gamma(\phi \phi)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

**0.80±0.07 OUR FIT**

$\Gamma_{56}/\Gamma$

NODE=M056R16  
NODE=M056R16

$$\Gamma(\phi \phi \eta)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-4}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**8.4±0.7±0.6**

186.6

<sup>1</sup> ABLIKIM 20B BES3  $\psi(2S) \rightarrow \gamma \phi \phi \eta$

<sup>1</sup> ABLIKIM 20B reports  $(8.41 \pm 0.74 \pm 0.62) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ .

$\Gamma_{57}/\Gamma$

NODE=M056R98  
NODE=M056R98

NODE=M056R98;LINKAGE=A

$$\Gamma(p \bar{p})/\Gamma_{\text{total}}$$

VALUE (units  $10^{-4}$ )

DOCUMENT ID

**2.21±0.08 OUR FIT**

$\Gamma_{58}/\Gamma$

NODE=M056R11  
NODE=M056R11

$$\Gamma(p \bar{p} \pi^0)/\Gamma_{\text{total}}$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**0.70±0.07 OUR AVERAGE** Error includes scale factor of 1.3.

0.73±0.06±0.01

<sup>1</sup> ONYISI 10 CLE3  $\psi(2S) \rightarrow \gamma p \bar{p} X$

0.56±0.12±0.01

<sup>2</sup> ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ONYISI 10 reports  $(7.76 \pm 0.37 \pm 0.51 \pm 0.39) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p \bar{p} \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma_{59}/\Gamma$

NODE=M056R06  
NODE=M056R06

NODE=M056R06;LINKAGE=ON

<sup>2</sup> ATHAR 07 reports  $(0.59 \pm 0.10 \pm 0.08) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p \bar{p} \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R06;LINKAGE=AT

$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.35±0.04 OUR AVERAGE**

0.35±0.04±0.01

0.37±0.11±0.01

DOCUMENT ID

TECN

COMMENT

1 ONYISI

10

CLE3

 $\psi(2S) \rightarrow \gamma p\bar{p}X$ 

2 ATHAR

07

CLEO

 $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ 

<sup>1</sup> ONYISI 10 reports  $(3.73 \pm 0.38 \pm 0.28 \pm 0.19) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ATHAR 07 reports  $(0.39 \pm 0.11 \pm 0.04) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{60}/\Gamma$ NODE=M056R50  
NODE=M056R50

NODE=M056R50;LINKAGE=ON

NODE=M056R50;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.52±0.06±0.01**

DOCUMENT ID

TECN

COMMENT

1 ONYISI

10

CLE3

 $\psi(2S) \rightarrow \gamma p\bar{p}X$ 

<sup>1</sup> ONYISI 10 reports  $(5.57 \pm 0.48 \pm 0.42 \pm 0.14) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{61}/\Gamma$ NODE=M056R70  
NODE=M056R70

NODE=M056R70;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )**6.0±1.4±0.1**

EVTS

42 ± 8

DOCUMENT ID

TECN

COMMENT

1 ABLIKIM

11F

BES3

 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$ 

<sup>1</sup> ABLIKIM 11F reports  $(6.12 \pm 1.18 \pm 0.86) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{62}/\Gamma$ NODE=M056R75  
NODE=M056R75

NODE=M056R75;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**2.1 ±0.7 OUR EVALUATION**

2.1 ±1.0 OUR AVERAGE

1.57±0.21±0.53

4.20±1.15±0.18

DOCUMENT ID

TECN

COMMENT

Error includes scale factor of 1.4. Treating systematic error as correlated.

Error includes scale factor of 2.0.

1 BAI

99B

BES

 $\psi(2S) \rightarrow \gamma\chi_{c0}$ 

1 TANENBAUM

78

MRK1

 $\psi(2S) \rightarrow \gamma\chi_{c0}$ 

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .

 $\Gamma_{63}/\Gamma$ NODE=M056R7  
NODE=M056R7

→ UNCHECKED ←

NODE=M056R7;LINKAGE=X1

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ 

VALUE (%)

**0.104±0.028±0.002**

EVTS

39.5

DOCUMENT ID

TECN

COMMENT

1 HE

08B

CLEO

 $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ 

<sup>1</sup> HE 08B reports  $0.11 \pm 0.02 \pm 0.02 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{64}/\Gamma$ NODE=M056R65  
NODE=M056R65

NODE=M056R65;LINKAGE=HE

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**1.22±0.26±0.02**

EVTS

48 ± 8

DOCUMENT ID

TECN

COMMENT

1 ABLIKIM

11F

BES3

 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$ 

<sup>1</sup> ABLIKIM 11F reports  $(1.24 \pm 0.20 \pm 0.18) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{65}/\Gamma$ NODE=M056R72  
NODE=M056R72

NODE=M056R72;LINKAGE=AB

$\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	<sup>1</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

NODE=M056R46  
 NODE=M056R46

<sup>1</sup> Using  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

NODE=M056R;LINKAGE=AB

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.7±1.1 OUR AVERAGE</b>				
12.9±1.1±0.3	5150	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
11.2±3.1±0.2		<sup>2</sup> ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

NODE=M056R49  
 NODE=M056R49

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.26 \pm 0.02 \pm 0.11) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R49;LINKAGE=AL

<sup>2</sup> ABLIKIM 06I reports  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.10 \pm 0.24 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R49;LINKAGE=AB

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.7±1.2±0.3</b>	5808	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

NODE=M056R82  
 NODE=M056R82

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.34 \pm 0.03 \pm 0.11) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R82;LINKAGE=AL

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>23.4±2.0±0.5</b>	2480	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

NODE=M056R83  
 NODE=M056R83

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.29 \pm 0.08 \pm 0.18) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R83;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>22.1±1.8±0.5</b>	2757	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

NODE=M056R84  
 NODE=M056R84

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.16 \pm 0.07 \pm 0.16) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R84;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$  $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>3.59±0.15 OUR FIT</b>	

NODE=M056R23  
 NODE=M056R23

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{72}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>118±12±2</b>		426	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M056R44  
 NODE=M056R44

• • • We do not use the following data for averages, fits, limits, etc. • • •

<400	90	<sup>2</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$
------	----	----------------------	----------	--

<sup>1</sup> ABLIKIM 12I reports  $(119.0 \pm 6.4 \pm 11.4) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R44;LINKAGE=AL

<sup>2</sup> Using  $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

NODE=M056R44;LINKAGE=AB



$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}$  $\Gamma_{73}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
<sup>1</sup> ABLIKIM 12I reports $< 54 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .				

NODE=M056R77  
NODE=M056R77

NODE=M056R77;LINKAGE=AL

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$
<sup>1</sup> ABLIKIM 12I reports $< 55 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .				

NODE=M056R78  
NODE=M056R78

NODE=M056R78;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{76}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$
<sup>1</sup> ABLIKIM 12I reports $< 50 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .				

NODE=M056R79  
NODE=M056R79

NODE=M056R79;LINKAGE=AL

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.25±0.12 OUR AVERAGE</b>		Error includes scale factor of 1.3.		
1.30±0.09±0.03	9k	<sup>1,2</sup> ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
1.01±0.19±0.02		<sup>3</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
<sup>1</sup> ABLIKIM 13D reports $(1.32 \pm 0.03 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>2</sup> Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$ .				
<sup>3</sup> ATHAR 07 reports $(1.07 \pm 0.17 \pm 0.12) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R07  
NODE=M056R07

NODE=M056R07;LINKAGE=AB

NODE=M056R07;LINKAGE=LB  
NODE=M056R07;LINKAGE=AT $\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{79}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.8±0.9±0.1</b>	254	<sup>1</sup> ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+}\bar{p}\Lambda$
<sup>1</sup> ABLIKIM 19AU reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ = $(4.7 \pm 0.7 \pm 0.5) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R99  
NODE=M056R99

NODE=M056R99;LINKAGE=A

 $\Gamma(K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.9±0.7±0.1</b>	62 ± 12	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$
<sup>1</sup> ABLIKIM 11F reports $(3.00 \pm 0.58 \pm 0.50) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+\bar{p}\Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R73  
NODE=M056R73

NODE=M056R73;LINKAGE=AB

$$\Gamma(nK_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{78}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.7±0.3±0.4</b>	1284	<sup>1</sup> ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$

NODE=M056P02  
NODE=M056P02

<sup>1</sup> ABLIKIM 21AV reports  $(6.65 \pm 0.26 \pm 0.41) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow n K_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.0979 \pm 0.0020$ . Also uses  $B(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = (63.9 \pm 0.5)\%$  and  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ .

NODE=M056P02;LINKAGE=A

$$\Gamma(\Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}} \quad \Gamma_{81}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.1±1.2±0.1</b>	28 ± 10	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M056R74  
NODE=M056R74

<sup>1</sup> ABLIKIM 11F reports  $(3.18 \pm 1.11 \pm 0.53) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R74;LINKAGE=AB

$$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}} \quad \Gamma_{82}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.68±0.32 OUR AVERAGE</b>				

NODE=M056R59  
NODE=M056R59

4.82±0.34±0.10      1046      <sup>1</sup> ABLIKIM      18v BES3       $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

4.2 ±0.7 ±0.1      78 ± 10      <sup>2</sup> NAIK      08 CLEO       $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.7 ±0.5 ±0.1      243      <sup>3,4</sup> ABLIKIM      13H BES3       $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<sup>1</sup> ABLIKIM 18v reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.72 \pm 0.18 \pm 0.28) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R59;LINKAGE=A

<sup>2</sup> NAIK 08 reports  $(4.41 \pm 0.56 \pm 0.47) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R59;LINKAGE=NA

<sup>3</sup> ABLIKIM 13H reports  $(4.78 \pm 0.34 \pm 0.39) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R59;LINKAGE=AB

<sup>4</sup> Superseded by ABLIKIM 18v

NODE=M056R59;LINKAGE=B

$$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}} \quad \Gamma_{85}/\Gamma$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.6 ±0.8 OUR AVERAGE</b>				Error includes scale factor of 2.6.

NODE=M056R60  
NODE=M056R60

5.10±0.35±0.10      747      <sup>1</sup> ABLIKIM      18v BES3       $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

3.1 ±0.7 ±0.1      39 ± 7      <sup>2</sup> NAIK      08 CLEO       $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.5 ±0.5 ±0.1      148      <sup>3,4</sup> ABLIKIM      13H BES3       $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

<sup>1</sup> ABLIKIM 18v reports  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (4.99 \pm 0.24 \pm 0.24) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R60;LINKAGE=A

<sup>2</sup> NAIK 08 reports  $(3.25 \pm 0.57 \pm 0.43) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R60;LINKAGE=NA

<sup>3</sup> ABLIKIM 13H reports  $(4.54 \pm 0.42 \pm 0.30) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R60;LINKAGE=AB

<sup>4</sup> Superseded by ABLIKIM 18v

NODE=M056R60;LINKAGE=B

$\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$  $\Gamma_{86}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.1 ± 0.2 ± 0.4</b>	2143	<sup>1</sup> ABLIKIM	20I	BES3 $\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$
<sup>1</sup> ABLIKIM 20I reports $(5.13 \pm 0.24 \pm 0.41) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ .				

NODE=M056P00  
NODE=M056P00

NODE=M056P00;LINKAGE=A

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$  $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.2 ± 5.8 ± 0.3</b>	27	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
<sup>1</sup> ABLIKIM 12I reports $(16.4 \pm 5.7 \pm 1.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R80  
NODE=M056R80

NODE=M056R80;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$  $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>23.2 ± 6.5 ± 0.5</b>	33	<sup>1</sup> ABLIKIM	12I	BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
<sup>1</sup> ABLIKIM 12I reports $(23.5 \pm 6.2 \pm 2.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R81  
NODE=M056R81

NODE=M056R81;LINKAGE=AL

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.94 ± 0.35 ± 0.04</b>	57	<sup>1</sup> ABLIKIM	15I	BES3 $\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$
<sup>1</sup> ABLIKIM 15I reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ $= (1.90 \pm 0.30 \pm 0.16) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R92  
NODE=M056R92

NODE=M056R92;LINKAGE=A

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$  $\Gamma_{90}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.5 ± 0.5 OUR AVERAGE</b>				Error includes scale factor of 1.7. $[(3.1 \pm 0.8) \times 10^{-4}$ OUR 2022 AVERAGE]
4.67 ± 0.19 ± 0.26	1741	<sup>1</sup> ABLIKIM	220	BES3 $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
3.1 ± 0.8 ± 0.1	23.3 ± 4.9	<sup>2</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
<sup>1</sup> ABLIKIM 220 reports $(4.67 \pm 0.19 \pm 0.26) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$ .				
<sup>2</sup> NAIK 08 reports $(3.34 \pm 0.70 \pm 0.48) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R61  
NODE=M056R61

NEW

NODE=M056R61;LINKAGE=A

NODE=M056R61;LINKAGE=NA

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$  $\Gamma_{91}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.45 ± 0.19 OUR AVERAGE</b>					$[(4.8 \pm 0.7) \times 10^{-4}$ OUR 2022 AVERAGE]
4.43 ± 0.08 ± 0.18		4932	<sup>1</sup> ABLIKIM	220	BES3 $\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
4.8 ± 0.7 ± 0.1		95	<sup>2</sup> NAIK	08	CLEO $\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<10.3		90	<sup>3</sup> ABLIKIM	06D	BES2 $\psi(2S) \rightarrow \chi_{c0} \gamma$
<sup>1</sup> ABLIKIM 220 reports $(4.43 \pm 0.08 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$ .					
<sup>2</sup> NAIK 08 reports $(5.14 \pm 0.60 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
<sup>3</sup> Using $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.2 \pm 0.5)\%$					

NODE=M056R45  
NODE=M056R45

NEW

NODE=M056R45;LINKAGE=A

NODE=M056R45;LINKAGE=NA

NODE=M056R45;LINKAGE=AB

$\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7 \times 10^{-4}$	90	1,2 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 41 \times 10^{-4}$	90	1,3 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
-----------------------	----	-------------	----------	---

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$ .

<sup>2</sup> From the  $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

<sup>3</sup> From the  $\eta_c \rightarrow K^+ K^- \pi^0$  decays.

NODE=M056R89  
NODE=M056R89

OCCUR=2

NODE=M056R89;LINKAGE=A  
NODE=M056R89;LINKAGE=B  
NODE=M056R89;LINKAGE=C

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma \times \Gamma_{32}/\Gamma$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>18.8 ± 1.0 OUR FIT</b>			
<b>15.3 ± 2.4 ± 0.8</b>	<sup>1</sup> ANDREOTTI 03	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0 \pi^0$

<sup>1</sup> We have multiplied  $B(p\bar{p}) \cdot B(\pi^0 \pi^0)$  measurement by 3 to obtain  $B(p\bar{p}) \cdot B(\pi\pi)$ .

NODE=M056R21  
NODE=M056R21

NODE=M056R;LINKAGE=AD

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0 \eta)/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma \times \Gamma_{33}/\Gamma$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.4</b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$

NODE=M056R41  
NODE=M056R41

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0 \eta')/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma \times \Gamma_{34}/\Gamma$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.5</b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$

NODE=M056R42  
NODE=M056R42

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma \times \Gamma_{36}/\Gamma$ 

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.7 ± 0.5 OUR FIT</b>			
<b>4.0 ± 1.2<sup>+0.5</sup><sub>-0.3</sub></b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \eta\eta$

NODE=M056R40  
NODE=M056R40

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta')/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma \times \Gamma_{37}/\Gamma$ 

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.1<sup>+2.3</sup><sub>-1.5</sub></b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<b>2.1<sup>+2.3</sup><sub>-1.5</sub></b>	ANDREOTTI 05C	E835	$\bar{p}p \rightarrow \pi^0 \eta$
--	---------------	------	-----------------------------------

NODE=M056R43  
NODE=M056R43

————— **RADIATIVE DECAYS** —————

NODE=M056310

 $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$   $\Gamma_{93}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.40 ± 0.05 OUR FIT</b>				

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.25 \pm 0.16 \pm 2.15$	12k	<sup>1</sup> ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
$2.0 \pm 0.2 \pm 0.2$		<sup>2</sup> ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> Not independent from  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$  and the product  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))$  also measured in ABLIKIM 17U.

<sup>2</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma \gamma J/\psi)$  from ADAM 05A and  $B(\psi(2S) \rightarrow \gamma \chi_{c0})$  from ATHAR 04.

NODE=M056R8  
NODE=M056R8

NODE=M056R8;LINKAGE=A

NODE=M056R8;LINKAGE=AD

 $\Gamma(\gamma \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt; 9</b>	90	$1.2 \pm 4.5$	<sup>1</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 10$	90	$6 \pm 12$	<sup>2</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
--------	----	------------	----------------------	----------	---

<sup>1</sup> BENNETT 08A reports  $< 9.6 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

<sup>2</sup> ABLIKIM 11E reports  $< 10.5 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .

NODE=M056R56  
NODE=M056R56

NODE=M056R56;LINKAGE=BE

NODE=M056R56;LINKAGE=AB

$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$  $\Gamma_{95}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8	90	$0.0 \pm 2.8$	<sup>1</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$

••• We do not use the following data for averages, fits, limits, etc. •••

<13	90	$5 \pm 11$	<sup>2</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
<sup>1</sup> BENNETT 08A reports $< 8.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .					
<sup>2</sup> ABLIKIM 11E reports $< 12.9 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .					

NODE=M056R57  
NODE=M056R57

NODE=M056R57;LINKAGE=BE

NODE=M056R57;LINKAGE=AB

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$  $\Gamma_{96}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 6	90	$0.1 \pm 1.6$	<sup>1</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$

••• We do not use the following data for averages, fits, limits, etc. •••

<16	90	$15 \pm 7$	<sup>2</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
<sup>1</sup> BENNETT 08A reports $< 6.4 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .					
<sup>2</sup> ABLIKIM 11E reports $< 16.2 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.79 \times 10^{-2}$ .					

NODE=M056R58  
NODE=M056R58

NODE=M056R58;LINKAGE=BE

NODE=M056R58;LINKAGE=AB

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{97}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>2.04±0.09 OUR FIT</b>				

••• We do not use the following data for averages, fits, limits, etc. •••

<7	90	<sup>1</sup> WICHT	08 BELL	$B^{\pm} \rightarrow K^{\pm}\gamma\gamma$
<sup>1</sup> WICHT 08 reports $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c0} K^+)] < 0.11 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c0} K^+) = 1.51 \times 10^{-4}$ .				

NODE=M056R1  
NODE=M056R1

NODE=M056R1;LINKAGE=WI

 $\Gamma(e^+e^-J/\psi(1S))/\Gamma_{\text{total}}$  $\Gamma_{98}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.54 \pm 0.33 \pm 0.03$	56	<sup>1,2</sup> ABLIKIM	17I BES3	$\psi(2S) \rightarrow \gamma e^+e^-J/\psi$

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> ABLIKIM 17I reports $(1.51 \pm 0.30 \pm 0.13) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow e^+e^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				
<sup>2</sup> Not independent from other measurements reported by ABLIKIM 17I				

NODE=M056R93  
NODE=M056R93

NODE=M056R93;LINKAGE=B

NODE=M056R93;LINKAGE=C

 $\Gamma(e^+e^-J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$  $\Gamma_{98}/\Gamma_{93}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.5±1.9±0.7</b>	56	<sup>1</sup> ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

NODE=M056R94  
NODE=M056R94

NODE=M056R94;LINKAGE=A

 $\Gamma(\mu^+\mu^-J/\psi(1S))/\Gamma(e^+e^-J/\psi(1S))$  $\Gamma_{99}/\Gamma_{98}$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.14	90	<9.5	ABLIKIM	19Z BES3	$\psi(2S) \rightarrow \gamma\chi_c \rightarrow \gamma(\mu^+\mu^-J/\psi)$

NODE=M056R97  
NODE=M056R97

 $\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$  $\Gamma_{97}/\Gamma_{93}$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.45±0.08 OUR FIT</b>			
<b>2.0 ±0.4 OUR AVERAGE</b>			

2.2 ±0.4 <sup>+0.1</sup><sub>-0.2</sub>

1.45±0.74	<sup>1</sup> ANDREOTTI	04 E835	$p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$
	<sup>2</sup> AMBROGIANI	00B E835	$\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$

<sup>1</sup> The values of  $B(p\bar{p})B(\gamma\gamma)$  and  $B(\gamma\gamma)B(\gamma J/\psi)$  measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

<sup>2</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .

NODE=M056R18  
NODE=M056R18

NODE=M056R;LINKAGE=AN

NODE=M056R;LINKAGE=7A

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{58}/\Gamma \times \Gamma_{93}/\Gamma$$

VALUE (units  $10^{-7}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**31.1±1.5 OUR FIT**

**28.2±2.1 OUR AVERAGE**

28.0±1.9±1.3    392    1,2,3 BAGNASCO    02    E835     $p\bar{p} \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$

29.3<sup>+5.7</sup><sub>-4.7</sub>±1.5    89    1,2 AMBROGIANI    99B     $p\bar{p} \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$

<sup>1</sup> Values in  $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$  and  $(\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$  are not independent. The latter is used in the fit since it is less correlated to the total width.

<sup>2</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .

<sup>3</sup> Recalculated by ANDREOTTI 05A.

NODE=M056R19  
NODE=M056R19

NODE=M056R;LINKAGE=KS

NODE=M056R19;LINKAGE=7A  
NODE=M056R19;LINKAGE=AN

$$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{58}/\Gamma \times \Gamma_{97}/\Gamma$$

VALUE (units  $10^{-8}$ )    DOCUMENT ID    TECN    COMMENT

**4.52±0.27 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.52±1.18<sup>+0.48</sup><sub>-0.72</sub>    <sup>1</sup> ANDREOTTI    04    E835     $p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$

<sup>1</sup> The values of  $B(p\bar{p})B(\gamma\gamma)$  and  $B(\gamma\gamma)B(\gamma J/\psi)$  measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

NODE=M056R25  
NODE=M056R25

NODE=M056R25;LINKAGE=AN

### $\chi_{c0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M056230

$$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{58}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units  $10^{-6}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**21.7±0.9 OUR FIT**

**23.7±1.0 OUR AVERAGE**

23.7±0.8±0.9    1222    ABLIKIM    13v    BES3     $\psi(2S) \rightarrow \gamma p\bar{p}$

23.7±1.4±1.4    383 ± 22    <sup>1</sup> NAIK    08    CLEO     $\psi(2S) \rightarrow \gamma p\bar{p}$

23.6<sup>+3.7</sup><sub>-3.4</sub>±3.4    89.5<sup>+14</sup><sub>-13</sub>    BAI    04F    BES     $\psi(2S) \rightarrow \gamma\chi_{c0}(1P) \rightarrow \gamma p\bar{p}$

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c0} \rightarrow p\bar{p}) = (25.7 \pm 1.5 \pm 1.5 \pm 1.3) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

NODE=M056B6  
NODE=M056B6

NODE=M056B6;LINKAGE=NA

$$\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{58}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units  $10^{-5}$ )    DOCUMENT ID    TECN    COMMENT

**6.25±0.26 OUR FIT**

**4.6 ±1.9**    <sup>1</sup> BAI    98i    BES     $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma p\bar{p}$

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow p\bar{p})$  reported in BAI 98i is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M056B1  
NODE=M056B1

NODE=M056B;LINKAGE=B1

$$\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{71}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units  $10^{-6}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**35.2±1.3 OUR FIT**

**35.1±1.4 OUR AVERAGE** Error includes scale factor of 1.1.

35.6±1.0±1.0    1486    ABLIKIM    21L    BES3     $\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+$

31.2±3.3±2.0    131    <sup>1</sup> NAIK    08    CLEO     $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

32.0±1.9±2.2    369    2,3 ABLIKIM    13H    BES3     $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.8 \pm 3.6 \pm 2.2 \pm 1.7) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

<sup>2</sup> Superseded by ABLIKIM 21L

<sup>3</sup> Calculated by us. ABLIKIM 13H reports  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.3 \pm 2.0 \pm 2.6) \times 10^{-5}$  from a measurement of  $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c0})$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.62 \pm 0.31)\%$ .

NODE=M056B20  
NODE=M056B20

NODE=M056B20;LINKAGE=NA

NODE=M056B20;LINKAGE=A  
NODE=M056B20;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{71}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**10.1 ± 0.4 OUR FIT**

13.0 <sup>+3.6</sup> <sub>-3.5</sub> ± 2.5	15.2 <sup>+4.2</sup> <sub>-4.0</sub>	1 BAI	03E BES	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda}$
--	--------------------------------------	-------	---------	---

<sup>1</sup> BAI 03E reports [  $B(\chi_{c0} \rightarrow \Lambda \bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c0}) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) ] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p \bar{p})] = (2.45^{+0.68}_{-0.65} \pm 0.46)\%$ . We calculate from this measurement the presented value using  $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$  and  $B(J/\psi \rightarrow p \bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$ .

NODE=M056B21  
NODE=M056B21

NODE=M056B21;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Lambda \bar{\Lambda} \eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{74}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.26 ± 0.30 ± 0.20	67	ABLIKIM	22A0 BES3	$\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+ \gamma \gamma$
--------------------	----	---------	-----------	---

NODE=M056P03  
NODE=M056P03

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.138 ± 0.005 OUR FIT****0.147 ± 0.029 OUR AVERAGE**

Error includes scale factor of 4.6.

0.158 ± 0.003 ± 0.006	4.8k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
0.024 ± 0.015 ± 0.205	12k	ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
0.069 ± 0.018		<sup>2</sup> OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.4 ± 0.3		<sup>3</sup> BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.16 ± 0.11		<sup>3</sup> BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma \chi_{c0}$
3.3 ± 1.7		<sup>4</sup> BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.151 ± 0.003 ± 0.010	4.3k	<sup>5</sup> ABLIKIM	120 BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.125 ± 0.007 ± 0.013	560	<sup>6</sup> MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.18 ± 0.01 ± 0.02	172	<sup>7</sup> ADAM	05A CLEO	Repl. by MENDEZ 08

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ .

<sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

<sup>3</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .

<sup>4</sup> Assumes isotropic gamma distribution.

<sup>5</sup> Superseded by ABLIKIM 17N.

<sup>6</sup> Not independent from other measurements of MENDEZ 08.

<sup>7</sup> Not independent from other values reported by ADAM 05A.

NODE=M056B2;LINKAGE=A

NODE=M056B;LINKAGE=3Q

NODE=M056B;LINKAGE=2Q

NODE=M056B;LINKAGE=EA

NODE=M056B2;LINKAGE=B

NODE=M056B2;LINKAGE=ME

NODE=M056B2;LINKAGE=AD

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})}{\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)}}$$

$$\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)} = \Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/(\Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \Gamma_{14}^{\psi(2S)} + 0.343 \Gamma_{167}^{\psi(2S)} + 0.190 \Gamma_{168}^{\psi(2S)})$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.224 ± 0.009 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.201 ± 0.011 ± 0.021	560	<sup>1</sup> MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0}$
0.31 ± 0.02 ± 0.03	172	ADAM	05A CLEO	Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

NODE=M056B7

NODE=M056B7

NODE=M056B7

NODE=M056B7;LINKAGE=ME

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{93}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.397 ± 0.015 OUR FIT**

0.358 ± 0.020 ± 0.037	560	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c0}$
-----------------------	-----	--------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.55 ± 0.04 ± 0.06	172	<sup>1</sup> ADAM	05A CLEO	Repl. by MENDEZ 08
--------------------	-----	-------------------	----------	--------------------

<sup>1</sup> Not independent from other values reported by ADAM 05A.

NODE=M056B8

NODE=M056B8

NODE=M056B;LINKAGE=AD

$$\Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{97}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.00±0.08 OUR FIT****1.95±0.09 OUR AVERAGE**

1.93±0.08±0.05	3.5k	ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
2.17±0.32±0.10	0.2k	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
3.7 ±1.8 ±1.0		LEE	85 CBAL	$\psi(2S) \rightarrow \gamma\chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.17±0.17±0.12	0.8k	<sup>1</sup> ABLIKIM	12A BES3	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow 3\gamma$
----------------	------	----------------------	----------	--

<sup>1</sup> Superseded by ABLIKIM 17AE.

NODE=M056B3  
NODE=M056B3

NODE=M056B3;LINKAGE=A

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{32}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**8.34±0.29 OUR FIT****8.80±0.34 OUR AVERAGE**

9.11±0.08±0.65	17k	<sup>1</sup> ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
8.81±0.11±0.43	8.9k	<sup>2</sup> ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
8.13±0.19±0.89	2.8k	<sup>3</sup> ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

<sup>1</sup> Calculated by us. ABLIKIM 10A reports  $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (3.23 \pm 0.03 \pm 0.23 \pm 0.14) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ . We have multiplied the  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

<sup>2</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow \pi^+\pi^-) = (6.37 \pm 0.08 \pm 0.31 \pm 0.32) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ . We have multiplied the  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

<sup>3</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow \pi^0\pi^0) = (2.94 \pm 0.07 \pm 0.32 \pm 0.15) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ . We have multiplied the  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

NODE=M056B22  
NODE=M056B22

OCCUR=2

NODE=M056B22;LINKAGE=AB

NODE=M056B22;LINKAGE=AS

NODE=M056B22;LINKAGE=AN

$$\Gamma(\chi_{c0}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{32}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**24.0±0.8 OUR FIT****20.7±1.7 OUR AVERAGE**

23.9±2.7±4.1	97 ± 11	<sup>1</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^0\pi^0$
20.2±1.1±1.5	720 ± 32	<sup>2</sup> BAI	98i BES	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\pi^+\pi^-$

<sup>1</sup> We have multiplied  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

<sup>2</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow \pi^+\pi^-)$  reported in BAI 98i is derived using  $B(\psi' \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi' \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D]. We have multiplied  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

NODE=M056B5  
NODE=M056B5

NODE=M056B;LINKAGE=D1

NODE=M056B;LINKAGE=D2

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{36}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.95±0.18 OUR FIT****3.12±0.19 OUR AVERAGE**

3.23±0.09±0.23	2132	<sup>1</sup> ABLIKIM	10A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$
2.93±0.12±0.29	0.9k	<sup>2</sup> ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.86±0.46±0.37	48	<sup>3</sup> ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma\chi_{c0}$
----------------	----	--------------------	---------	--

<sup>1</sup> Calculated by us. ABLIKIM 10A reports  $B(\chi_{c0} \rightarrow \eta\eta) = (3.44 \pm 0.10 \pm 0.24 \pm 0.13) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ .

<sup>2</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow \eta\eta) = (3.18 \pm 0.13 \pm 0.31 \pm 0.16) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

<sup>3</sup> Superseded by ASNER 09. Calculated by us. The value of  $B(\chi_{c0}(1P) \rightarrow \eta\eta)$  reported by ADAMS 07 was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46)\%$  (ATHAR 04).

NODE=M056B11  
NODE=M056B11

NODE=M056B11;LINKAGE=AB

NODE=M056B11;LINKAGE=AS

NODE=M056B11;LINKAGE=AD

$$\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{36}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**0.85 ±0.05 OUR FIT****0.578±0.241±0.158**

BAI	03C BES	$\psi(2S) \rightarrow \gamma\eta\eta$
-----	---------	---------------------------------------

NODE=M056B10  
NODE=M056B10



$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_{42} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.92±0.28 OUR FIT**

5.97±0.07±0.32	8.1k	<sup>1</sup> ASNER	09	CLEO $\psi(2S) \rightarrow \gamma K^+ K^-$
----------------	------	--------------------	----	--

<sup>1</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow K^+ K^-) = (6.47 \pm 0.08 \pm 0.35 \pm 0.32) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

NODE=M056B23  
NODE=M056B23

NODE=M056B23;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{42} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.71±0.08 OUR FIT**

1.63±0.10±0.15	774 ± 38	<sup>1</sup> BAI	98i	BES $\psi(2S) \rightarrow \gamma K^+ K^-$
----------------	----------	------------------	-----	---

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow K^+ K^-)$  reported by BAI 98i is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M056B9  
NODE=M056B9

NODE=M056B9;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_{43} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.10±0.16 OUR FIT****3.18±0.17 OUR AVERAGE**

3.22±0.07±0.17	2.1k	<sup>1</sup> ASNER	09	CLEO $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
----------------	------	--------------------	----	--

3.02±0.19±0.33	322	ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
----------------	-----	---------	-----	--

<sup>1</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c0} \rightarrow K_S^0 K_S^0) = (3.49 \pm 0.08 \pm 0.18 \pm 0.17) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$ .

NODE=M056B12  
NODE=M056B12

NODE=M056B12;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{43} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**8.9±0.5 OUR FIT**

5.6±0.8±1.3	<sup>1</sup> BAI	99B	BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
-------------	------------------	-----	---

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow K_S^0 K_S^0)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M056B13  
NODE=M056B13

NODE=M056B13;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_1 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**6.6±0.5 OUR FIT****6.9±2.4 OUR AVERAGE** Error includes scale factor of 3.8.

4.4±0.1±0.9	<sup>1</sup> BAI	99B	BES $\psi(2S) \rightarrow \gamma \chi_{c0}$
-------------	------------------	-----	---

9.3±0.9	<sup>2</sup> TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$
---------	------------------------	----	--

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$  reported in BAI 99B is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

<sup>2</sup> The value  $B(\psi(1S) \rightarrow \gamma \chi_{c0}) \times B(\chi_{c0} \rightarrow 2\pi^+ 2\pi^-)$  reported in TANENBAUM 78 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

NODE=M056B4  
NODE=M056B4

NODE=M056B;LINKAGE=B2

NODE=M056B;LINKAGE=J1

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_8 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**1.78±0.14 OUR FIT**

1.64±0.05±0.2	ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \chi_{c0}$
---------------	---------	-----	--

NODE=M056B18  
NODE=M056B18

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_8 / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID TECN COMMENT

NODE=M056B19  
NODE=M056B19

**5.1 ± 0.4 OUR FIT**

**5.8 ± 1.6 OUR AVERAGE** Error includes scale factor of 2.3.

4.22 ± 0.20 ± 0.97

BAI 99B BES  $\psi(2S) \rightarrow \gamma \chi_{c0}$

7.4 ± 1.0

<sup>1</sup> TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma \chi_{c0}$

<sup>1</sup> The reported value is derived using  $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using  $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

NODE=M056B19;LINKAGE=TA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_{51} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units  $10^{-4}$ ) EVTS

DOCUMENT ID TECN COMMENT

NODE=M056B14  
NODE=M056B14

**2.76 ± 0.28 OUR FIT**

**3.20 ± 0.11 ± 0.41**

278

<sup>1</sup> ABLIKIM 06T BES2  $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$ .

NODE=M056B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{51} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

NODE=M056B15  
NODE=M056B15

**8.0 ± 0.8 OUR FIT**

**6.1 ± 0.8 ± 0.9**

<sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow 2K^+ 2K^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M056B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_{56} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units  $10^{-4}$ ) EVTS

DOCUMENT ID TECN COMMENT

NODE=M056B16  
NODE=M056B16

**0.78 ± 0.07 OUR FIT**

**0.78 ± 0.08 OUR AVERAGE**

0.77 ± 0.03 ± 0.08

612

<sup>1</sup> ABLIKIM 11K BES3  $\psi(2S) \rightarrow \gamma$  hadrons

0.86 ± 0.19 ± 0.12

26

<sup>2</sup> ABLIKIM 06T BES2  $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow \phi \phi)$  reported by ABLIKIM 11K was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31)\%$ .

NODE=M056B16;LINKAGE=AL

<sup>2</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow \phi \phi)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4)\%$ .

NODE=M056B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{56} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

NODE=M056B17  
NODE=M056B17

**2.25 ± 0.21 OUR FIT**

**2.6 ± 1.0 ± 1.1**

<sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c0} \rightarrow \phi \phi)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.3 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M056B17;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) / \Gamma_{\text{total}}}{\Gamma_{83} / \Gamma \times \Gamma_{166}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units  $10^{-5}$ ) EVTS

DOCUMENT ID TECN COMMENT

NODE=M056B24  
NODE=M056B24

**3.45 ± 0.17 ± 0.19**

493

<sup>1</sup> ABLIKIM 19BB BES3  $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

<sup>1</sup> Calculated by us. ABLIKIM 19BB reports  $B(\chi_c^0 \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (3.52 \pm 0.19 \pm 0.21) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma \chi_c^0) = (9.79 \pm 0.20)\%$  and other branching fractions from PDG 18.

NODE=M056B24;LINKAGE=A

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{84}/\Gamma \times \Gamma_{166}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10 <sup>-5</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.97±0.12±0.14</b>	871	<sup>1</sup> ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+$ + c.c.

NODE=M056P01  
NODE=M056P01

<sup>1</sup> Calculated by us. ABLIKIM 20AE reports  $B(\chi_c^0 \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (3.03 \pm 0.12 \pm 0.15) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma \chi_c^0) = (9.79 \pm 0.20)\%$  and other branching fractions from PDG 20.

NODE=M056P01;LINKAGE=A

### χ<sub>c0</sub>(1P) REFERENCES

NODE=M056

ABLIKIM 22AO	PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61887
ABLIKIM 22O	JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61652
ABLIKIM 22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
ABLIKIM 21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61465
ABLIKIM 21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61117
ABLIKIM 20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60733
ABLIKIM 20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60212
ABLIKIM 20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60303
PDG 20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM 19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59844
ABLIKIM 19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM 19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60026
ABLIKIM 19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59606
ABLIKIM 19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59837
ABLIKIM 18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58990
PDG 18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AAJ 17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
ABLIKIM 17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58310
ABLIKIM 17AI	PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58322
ABLIKIM 17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM 17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM 17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
PDG 16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM 15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM 15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56778
ABLIKIM 15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56779
ABLIKIM 13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54877
ABLIKIM 13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM 13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM 13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
UEHARA 13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM 12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54266
ABLIKIM 12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM 12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM 12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
LIU 12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM 11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM 11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM 11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM 11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
DEL-AMO-SA... 11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM 10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53347
ONYISI 10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA 10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ASNER 09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA 09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT 08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND 08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE 08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ 08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK 08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA 08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT 08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE 07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ADAMS 07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR 07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN 07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM 06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM 06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM 06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM 06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
ABLIKIM 05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM 05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM 05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ABLIKIM 05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ADAM 05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI 05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
ANDREOTTI 05C	PR D72 112002	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50991
NAKAZAWA 05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABE 04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM 04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50187
ABLIKIM 04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ANDREOTTI 04	PL B584 16	M. Andreotti <i>et al.</i>	(E835 Collab.)	REFID=49744
ATHAR 04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI 04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
ANDREOTTI 03	PRL 91 091801	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=49578
AULCHENKO 03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI 03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI 03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE,K 02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAGNASCO 02	PL B533 237	S. Bagnasco <i>et al.</i>	(FNAL E835 Collab.)	REFID=48833
EISENSTEIN 01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI 00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
AMBROGIANI 99B	PRL 83 2902	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47389
BAI 99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
BAI 98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI 98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
GAISER 86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE 85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
OREGLIA 82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
BRANDELIK 79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL 78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM 78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also BIDDICK 77	Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059

$\chi_{c1}(1P)$ 

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the Review on " $\psi(2S)$  and  $\chi_c$  branching ratios" before the  $\chi_{c0}(1P)$  Listings.

NODE=M055

NODE=M055

 **$\chi_{c1}(1P)$  MASS**

NODE=M055M

NODE=M055M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3510.67 ± 0.05 OUR AVERAGE</b>		Error includes scale factor of 1.2.		
3508.4 ± 1.9 ± 0.7	460	<sup>1</sup> AAIJ	17BB LHCB	$p\bar{p} \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3510.71 ± 0.04 ± 0.09	4.8k	<sup>2</sup> AAIJ	17BI LHCB	$\chi_{c1} \rightarrow J/\psi\mu^+\mu^-$
3510.30 ± 0.14 ± 0.16		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
3510.719 ± 0.051 ± 0.019		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3509.4 ± 0.9		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3510.60 ± 0.087 ± 0.019	513	<sup>3</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3511.3 ± 0.4 ± 0.4	30	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
3512.3 ± 0.3 ± 4.0		<sup>4</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7	91	<sup>5</sup> LEMOIGNE	82 GOLI	$185\pi^-Be \rightarrow \gamma\mu^+\mu^-A$
3510.4 ± 0.6		OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1	254	<sup>6</sup> HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3509 ± 11	21	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3507 ± 3		<sup>6</sup> BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4		<sup>6,7</sup> TANENBAUM	78 MRK1	$e^+e^-$
3513 ± 7	367	<sup>6</sup> BIDDICK	77 CNTR	$\psi(2S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3500 ± 10	40	TANENBAUM	75 MRK1	Hadrons $\gamma$

OCCUR=2

NODE=M055M;LINKAGE=A

NODE=M055M;LINKAGE=B  
 NODE=M055M;LINKAGE=NW  
 NODE=M055M;LINKAGE=C  
 NODE=M055M;LINKAGE=P  
 NODE=M055M;LINKAGE=D

NODE=M055M;LINKAGE=M

- <sup>1</sup> From a fit of the  $\phi\phi$  invariant mass with the width of  $\chi_{c1}(1P)$  fixed to the PDG 16 value.  
<sup>2</sup> AAIJ 17BI reports also  $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$  MeV.  
<sup>3</sup> Recalculated by ANDREOTTI 05A, using the value of  $\psi(2S)$  mass from AULCHENKO 03.  
<sup>4</sup> Using mass of  $\psi(2S) = 3686.0$  MeV.  
<sup>5</sup>  $J/\psi(1S)$  mass constrained to 3097 MeV.  
<sup>6</sup> Mass value shifted by us by amount appropriate for  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.  
<sup>7</sup> From a simultaneous fit to radiative and hadronic decay channels.

 **$\chi_{c1}(1P)$  WIDTH**

NODE=M055W

NODE=M055W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.84 ± 0.04 OUR FIT</b>					
<b>0.88 ± 0.05 OUR AVERAGE</b>					
1.39 <sup>+0.40</sup> <sub>-0.38</sub> <sup>+0.26</sup> <sub>-0.77</sub>			ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
0.876 ± 0.045 ± 0.026			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
0.87 ± 0.11 ± 0.08		513	<sup>1</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.3		95	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-X$
<3.8		90	GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

- <sup>1</sup> Recalculated by ANDREOTTI 05A.

NODE=M055W;LINKAGE=AN

 **$\chi_{c1}(1P)$  DECAY MODES**

NODE=M055215;NODE=M055

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ hadrons		
$\Gamma_2$ $e^+e^-$	$(1.4^{+1.5}_{-1.0}) \times 10^{-7}$	

DESIG=112

DESIG=110

## Hadronic decays

				NODE=M055;CLUMP=A
Γ <sub>3</sub>	$3(\pi^+\pi^-)$	$(5.8 \pm 1.4) \times 10^{-3}$	S=1.2	DESIG=6
Γ <sub>4</sub>	$2(\pi^+\pi^-)$	$(7.6 \pm 2.6) \times 10^{-3}$		DESIG=5
Γ <sub>5</sub>	$\pi^+\pi^-\pi^0\pi^0$	$(1.19 \pm 0.15) \%$		DESIG=51
Γ <sub>6</sub>	$\rho^+\pi^-\pi^0 + \text{c.c.}$	$(1.45 \pm 0.24) \%$		DESIG=52
Γ <sub>7</sub>	$\rho^0\pi^+\pi^-$	$(3.9 \pm 3.5) \times 10^{-3}$		DESIG=9
Γ <sub>8</sub>	$4\pi^0$	$(5.4 \pm 0.8) \times 10^{-4}$		DESIG=60
Γ <sub>9</sub>	$\pi^+\pi^-K^+K^-$	$(4.5 \pm 1.0) \times 10^{-3}$		DESIG=7
Γ <sub>10</sub>	$K^+K^-\pi^0\pi^0$	$(1.12 \pm 0.27) \times 10^{-3}$		DESIG=53
Γ <sub>11</sub>	$K^+K^-\pi^+\pi^-\pi^0$	$(1.15 \pm 0.13) \%$		DESIG=79
Γ <sub>12</sub>	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(7.5 \pm 0.8) \times 10^{-3}$		DESIG=84
Γ <sub>13</sub>	$K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(8.6 \pm 1.4) \times 10^{-3}$		DESIG=55
Γ <sub>14</sub>	$\rho^-K^+\bar{K}^0 + \text{c.c.}$	$(5.0 \pm 1.2) \times 10^{-3}$		DESIG=56
Γ <sub>15</sub>	$K^*(892)^0\bar{K}^0\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	$(2.3 \pm 0.6) \times 10^{-3}$		DESIG=57
Γ <sub>16</sub>	$K^+K^-\eta\pi^0$	$(1.12 \pm 0.34) \times 10^{-3}$		DESIG=58
Γ <sub>17</sub>	$\pi^+\pi^-K_S^0K_S^0$	$(6.9 \pm 2.9) \times 10^{-4}$		DESIG=28
Γ <sub>18</sub>	$K^+K^-\eta$	$(3.2 \pm 1.0) \times 10^{-4}$		DESIG=42
Γ <sub>19</sub>	$\bar{K}^0K^+\pi^- + \text{c.c.}$	$(7.0 \pm 0.6) \times 10^{-3}$		DESIG=17
Γ <sub>20</sub>	$K^*(892)^0\bar{K}^0 + \text{c.c.}$	$(10 \pm 4) \times 10^{-4}$		DESIG=32
Γ <sub>21</sub>	$K^*(892)^+K^- + \text{c.c.}$	$(1.4 \pm 0.6) \times 10^{-3}$		DESIG=33
Γ <sub>22</sub>	$K_J^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K^+\pi^- + \text{c.c.}$	$< 8 \times 10^{-4}$	CL=90%	DESIG=34
Γ <sub>23</sub>	$K_J^*(1430)^+K^- + \text{c.c.} \rightarrow$ $K_S^0K^+\pi^- + \text{c.c.}$	$< 2.1 \times 10^{-3}$	CL=90%	DESIG=35
Γ <sub>24</sub>	$K^+K^-\pi^0$	$(1.81 \pm 0.24) \times 10^{-3}$		DESIG=38
Γ <sub>25</sub>	$\eta\pi^+\pi^-$	$(4.62 \pm 0.23) \times 10^{-3}$		DESIG=31
Γ <sub>26</sub>	$a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(3.2 \pm 0.4) \times 10^{-3}$	S=2.2	DESIG=36
Γ <sub>27</sub>	$a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(1.76 \pm 0.24) \times 10^{-4}$		DESIG=93
Γ <sub>28</sub>	$a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$(4.6 \pm 0.7) \times 10^{-5}$		DESIG=96
Γ <sub>29</sub>	$f_2(1270)\eta \rightarrow \eta\pi^+\pi^-$	$(3.5 \pm 0.6) \times 10^{-4}$		DESIG=94
Γ <sub>30</sub>	$f_4(2050)\eta \rightarrow \eta\pi^+\pi^-$	$(2.5 \pm 0.9) \times 10^{-5}$		DESIG=95
Γ <sub>31</sub>	$\pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 5 \times 10^{-5}$	CL=90%	DESIG=97
Γ <sub>32</sub>	$\pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 1.5 \times 10^{-5}$	CL=90%	DESIG=98
Γ <sub>33</sub>	$\pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	$< 8 \times 10^{-6}$	CL=90%	DESIG=99
Γ <sub>34</sub>	$f_2(1270)\eta$	$(6.7 \pm 1.1) \times 10^{-4}$		DESIG=37
Γ <sub>35</sub>	$\pi^+\pi^-\eta'$	$(2.2 \pm 0.4) \times 10^{-3}$		DESIG=44
Γ <sub>36</sub>	$K^+K^-\eta'(958)$	$(8.8 \pm 0.9) \times 10^{-4}$		DESIG=85
Γ <sub>37</sub>	$K_0^*(1430)^+K^- + \text{c.c.}$	$(6.4 \pm 2.2 \mp 2.8) \times 10^{-4}$		DESIG=86
Γ <sub>38</sub>	$f_0(980)\eta'(958)$	$(1.6 \pm 1.4 \mp 0.7) \times 10^{-4}$		DESIG=87
Γ <sub>39</sub>	$f_0(1710)\eta'(958)$	$(7 \pm 7 \mp 5) \times 10^{-5}$		DESIG=88
Γ <sub>40</sub>	$f_2'(1525)\eta'(958)$	$(9 \pm 6) \times 10^{-5}$		DESIG=89
Γ <sub>41</sub>	$\pi^0 f_0(980) \rightarrow \pi^0\pi^+\pi^-$	$(3.5 \pm 0.9) \times 10^{-7}$		DESIG=61
Γ <sub>42</sub>	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(3.2 \pm 2.1) \times 10^{-3}$		DESIG=10
Γ <sub>43</sub>	$K^*(892)^0\bar{K}^*(892)^0$	$(1.4 \pm 0.4) \times 10^{-3}$		DESIG=21
Γ <sub>44</sub>	$K^+K^-K_S^0K_S^0$	$< 4 \times 10^{-4}$	CL=90%	DESIG=29
Γ <sub>45</sub>	$K_S^0K_S^0K_S^0K_S^0$	$(3.5 \pm 1.0) \times 10^{-5}$		DESIG=102
Γ <sub>46</sub>	$K^+K^-K^+K^-$	$(5.4 \pm 1.1) \times 10^{-4}$		DESIG=14
Γ <sub>47</sub>	$K^+K^-\phi$	$(4.1 \pm 1.5) \times 10^{-4}$		DESIG=30
Γ <sub>48</sub>	$\bar{K}^0K^+\pi^-\phi + \text{c.c.}$	$(3.3 \pm 0.5) \times 10^{-3}$		DESIG=90
Γ <sub>49</sub>	$K^+K^-\pi^0\phi$	$(1.62 \pm 0.30) \times 10^{-3}$		DESIG=91
Γ <sub>50</sub>	$\phi\pi^+\pi^-\pi^0$	$(7.5 \pm 1.0) \times 10^{-4}$		DESIG=82
Γ <sub>51</sub>	$\omega\omega$	$(5.7 \pm 0.7) \times 10^{-4}$		DESIG=66
Γ <sub>52</sub>	$\omega K^+K^-$	$(7.8 \pm 0.9) \times 10^{-4}$		DESIG=81
Γ <sub>53</sub>	$\omega\phi$	$(2.7 \pm 0.4) \times 10^{-5}$		DESIG=67
Γ <sub>54</sub>	$\phi\phi$	$(4.2 \pm 0.5) \times 10^{-4}$		DESIG=68

Γ <sub>55</sub>	$\phi\phi\eta$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=104
Γ <sub>56</sub>	$\rho\bar{\rho}$	$(7.60 \pm 0.34) \times 10^{-5}$		DESIG=11
Γ <sub>57</sub>	$\rho\bar{\rho}\pi^0$	$(1.55 \pm 0.18) \times 10^{-4}$		DESIG=39
Γ <sub>58</sub>	$\rho\bar{\rho}\eta$	$(1.45 \pm 0.25) \times 10^{-4}$		DESIG=43
Γ <sub>59</sub>	$\rho\bar{\rho}\omega$	$(2.12 \pm 0.31) \times 10^{-4}$		DESIG=59
Γ <sub>60</sub>	$\rho\bar{\rho}\phi$	$< 1.7 \times 10^{-5}$	CL=90%	DESIG=65
Γ <sub>61</sub>	$\rho\bar{\rho}\pi^+\pi^-$	$(5.0 \pm 1.9) \times 10^{-4}$		DESIG=8
Γ <sub>62</sub>	$\rho\bar{\rho}\pi^0\pi^0$	$< 5 \times 10^{-4}$	CL=90%	DESIG=54
Γ <sub>63</sub>	$\rho\bar{\rho}K^+K^-$ (non-resonant)	$(1.27 \pm 0.22) \times 10^{-4}$		DESIG=62
Γ <sub>64</sub>	$\rho\bar{\rho}K_S^0 K_S^0$	$< 4.5 \times 10^{-4}$	CL=90%	DESIG=25
Γ <sub>65</sub>	$\rho\bar{n}\pi^-$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=74
Γ <sub>66</sub>	$\bar{\rho}n\pi^+$	$(3.9 \pm 0.5) \times 10^{-4}$		DESIG=75
Γ <sub>67</sub>	$\rho\bar{n}\pi^-\pi^0$	$(1.03 \pm 0.12) \times 10^{-3}$		DESIG=76
Γ <sub>68</sub>	$\bar{\rho}n\pi^+\pi^0$	$(1.01 \pm 0.12) \times 10^{-3}$		DESIG=77
Γ <sub>69</sub>	$\Lambda\bar{\Lambda}$	$(1.27 \pm 0.08) \times 10^{-4}$		DESIG=19
Γ <sub>70</sub>	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(2.9 \pm 0.5) \times 10^{-4}$		DESIG=24
Γ <sub>71</sub>	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=69
Γ <sub>72</sub>	$\Lambda\bar{\Lambda}\eta$	$(5.9 \pm 1.5) \times 10^{-5}$		DESIG=111
Γ <sub>73</sub>	$\Sigma(1385)^+\bar{\Lambda}\pi^- + c.c.$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=70
Γ <sub>74</sub>	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + c.c.$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=71
Γ <sub>75</sub>	$K^+\bar{p}\Lambda + c.c.$	$(4.2 \pm 0.4) \times 10^{-4}$	S=1.2	DESIG=40
Γ <sub>76</sub>	$nK_S^0\bar{\Lambda} + c.c.$	$(1.66 \pm 0.17) \times 10^{-4}$		DESIG=109
Γ <sub>77</sub>	$K^*(892)^+\bar{p}\Lambda + c.c.$	$(4.9 \pm 0.7) \times 10^{-4}$		DESIG=106
Γ <sub>78</sub>	$K^+\bar{p}\Lambda(1520) + c.c.$	$(1.7 \pm 0.4) \times 10^{-4}$		DESIG=63
Γ <sub>79</sub>	$\Lambda(1520)\bar{\Lambda}(1520)$	$< 9 \times 10^{-5}$	CL=90%	DESIG=64
Γ <sub>80</sub>	$\Sigma^0\bar{\Sigma}^0$	$(4.2 \pm 0.6) \times 10^{-5}$		DESIG=48
Γ <sub>81</sub>	$\Sigma^+\bar{p}K_S^0 + c.c.$	$(1.53 \pm 0.12) \times 10^{-4}$		DESIG=105
Γ <sub>82</sub>	$\Sigma^0\bar{p}K^+ + c.c.$	$(1.46 \pm 0.10) \times 10^{-4}$		DESIG=108
Γ <sub>83</sub>	$\Sigma^+\bar{\Sigma}^-$	$(3.6 \pm 0.7) \times 10^{-5}$		DESIG=49
Γ <sub>84</sub>	$\Sigma^-\bar{\Sigma}^+$	$(5.7 \pm 1.5) \times 10^{-5}$		DESIG=107
Γ <sub>85</sub>	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$< 9 \times 10^{-5}$	CL=90%	DESIG=72
Γ <sub>86</sub>	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$< 5 \times 10^{-5}$	CL=90%	DESIG=73
Γ <sub>87</sub>	$K^-\Lambda\bar{\Xi}^+ + c.c.$	$(1.35 \pm 0.24) \times 10^{-4}$		DESIG=92
Γ <sub>88</sub>	$\Xi^0\bar{\Xi}^0$	$(7.5 \pm 1.3) \times 10^{-5}$		DESIG=50
Γ <sub>89</sub>	$\Xi^-\bar{\Xi}^+$	$(6.0 \pm 0.6) \times 10^{-5}$		DESIG=26
Γ <sub>90</sub>	$\pi^+\pi^- + K^+K^-$	$< 2.1 \times 10^{-3}$		DESIG=23
Γ <sub>91</sub>	$K_S^0 K_S^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=27
Γ <sub>92</sub>	$\eta_c\pi^+\pi^-$	$< 3.2 \times 10^{-3}$	CL=90%	DESIG=83
<b>Radiative decays</b>				
Γ <sub>93</sub>	$\gamma J/\psi(1S)$	$(34.3 \pm 1.0) \%$		NODE=M055;CLUMP=B DESIG=1
Γ <sub>94</sub>	$\gamma\rho^0$	$(2.16 \pm 0.17) \times 10^{-4}$		DESIG=45
Γ <sub>95</sub>	$\gamma\omega$	$(6.8 \pm 0.8) \times 10^{-5}$		DESIG=46
Γ <sub>96</sub>	$\gamma\phi$	$(2.4 \pm 0.5) \times 10^{-5}$		DESIG=47
Γ <sub>97</sub>	$\gamma\gamma$	$< 6.3 \times 10^{-6}$	CL=90%	DESIG=4
Γ <sub>98</sub>	$e^+e^- J/\psi(1S)$	$(3.46 \pm 0.22) \times 10^{-3}$		DESIG=100
Γ <sub>99</sub>	$\mu^+\mu^- J/\psi(1S)$	$(2.33 \pm 0.29) \times 10^{-4}$		DESIG=103

## CONSTRAINED FIT INFORMATION

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_{46}$	3				
$x_{56}$	4	2			
$x_{69}$	11	4	5		
$x_{93}$	23	9	2	29	
$\Gamma$	-12	-5	-63	-15	-41
	$x_{19}$	$x_{46}$	$x_{56}$	$x_{69}$	$x_{93}$

### $\chi_{c1}(1P)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	$\Gamma_2$			
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.12^{+0.13}_{-0.08}$	250	<sup>1</sup> ABLIKIM	22AF BES3	$e^+e^- \rightarrow \chi_{c1} \rightarrow \gamma J/\psi$
<sup>1</sup> Assuming $\Gamma(\chi_{c1} \rightarrow \gamma J/\psi) = 0.28$ MeV.				

NODE=M055220

NODE=M055W1  
NODE=M055W1

NODE=M055W1;LINKAGE=A

### $\chi_{c1}(1P) \Gamma(i)\Gamma(\gamma J/\psi(1S))/\Gamma(\text{total})$

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$	$\Gamma_{56}\Gamma_{93}/\Gamma$		
VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>21.9±0.8 OUR FIT</b>			
<b>21.4±0.9 OUR AVERAGE</b>			
21.5±0.5±0.8	<sup>1</sup> ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
21.4±1.5±2.2	<sup>1,2</sup> ARMSTRONG	92 E760	$p\bar{p} \rightarrow e^+e^-\gamma$
19.9 <sup>+4.4</sup> <sub>-4.0</sub>	<sup>1</sup> BAGLIN	86B SPEC	$p\bar{p} \rightarrow e^+e^-X$
<sup>1</sup> Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .			
<sup>2</sup> Recalculated by ANDREOTTI 05A.			

NODE=M055223

NODE=M055G1  
NODE=M055G1NODE=M055G;LINKAGE=7A  
NODE=M055G;LINKAGE=AN

### $\chi_{c1}(1P)$ BRANCHING RATIOS

#### HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$	$\Gamma_3/\Gamma$			
VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.8 ± 1.4 OUR EVALUATION</b>				Error includes scale factor of 1.2. Treating systematic error as correlated.
<b>10.5 ± 1.5 OUR AVERAGE</b>				Error includes scale factor of 4.6. $[(5.8 \pm 1.1) \times 10^{-3}]$
				OUR 2022 AVERAGE]
10.92±0.04±0.35	84K	<sup>1</sup> ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+\pi^-)$
5.4 ± 0.7 ± 0.9		<sup>2</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
16.0 ± 5.9 ± 0.8		<sup>2</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$
<sup>1</sup> ABLIKIM 22Q reports $(1.092 \pm 0.004 \pm 0.035) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ .				
<sup>2</sup> Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .				

NODE=M055225

NODE=M055305

NODE=M055R6  
NODE=M055R6

→ UNCHECKED ←

NEW

NODE=M055R6;LINKAGE=A

NODE=M055R;LINKAGE=X2

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$	$\Gamma_4/\Gamma$		
VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>7.6±2.6 OUR EVALUATION</b>			Treating systematic error as correlated.
<b>8 ± 4 OUR AVERAGE</b>			Error includes scale factor of 1.5.
4.6±2.1±2.6	<sup>1</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$
12.5±4.2±0.6	<sup>1</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R4  
NODE=M055R4

→ UNCHECKED ←

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .

NODE=M055R4;LINKAGE=X2

### $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.19±0.15±0.03</b>	604.7	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R35  
NODE=M055R35

<sup>1</sup> HE 08B reports  $1.28 \pm 0.06 \pm 0.15 \pm 0.08\%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R35;LINKAGE=HE

### $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.45±0.24±0.04</b>	712.3	<sup>1,2</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R36  
NODE=M055R36

<sup>1</sup> HE 08B reports  $1.56 \pm 0.13 \pm 0.22 \pm 0.10\%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R36;LINKAGE=HE

<sup>2</sup> Calculated by us. We have added the values from HE 08B for  $\rho^+\pi^-\pi^0$  and  $\rho^-\pi^+\pi^0$  decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M055R36;LINKAGE=OC

### $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.9±3.5</b>	<sup>1</sup> TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R8  
NODE=M055R8

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

NODE=M055R;LINKAGE=T

### $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ $\Gamma_8/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.4±0.8±0.1</b>	608	<sup>1</sup> ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R44  
NODE=M055R44

<sup>1</sup> ABLIKIM 11A reports  $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R44;LINKAGE=AB

### $\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.5±1.0 OUR EVALUATION</b>	Treating systematic error as correlated.		
<b>4.5±0.9 OUR AVERAGE</b>			

NODE=M055R5  
NODE=M055R5

→ UNCHECKED ←

4.2±0.4±0.9

<sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 

7.3±3.0±0.4

<sup>1</sup> TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .

NODE=M055R5;LINKAGE=X2

### $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ $\Gamma_{10}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.12±0.27±0.03</b>	45.1	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M055R37  
NODE=M055R37

<sup>1</sup> HE 08B reports  $(0.12 \pm 0.02 \pm 0.02 \pm 0.01) \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R37;LINKAGE=HE

### $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{11}/\Gamma$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.46±0.12±1.29</b>	12k	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R00  
NODE=M055R00

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$ .

NODE=M055R00;LINKAGE=A



$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.52±0.11±0.79</b>	5.1k	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$ .

NODE=M055R60  
NODE=M055R60

NODE=M055R60;LINKAGE=A

 $\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.86±0.13±0.02</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.92 \pm 0.09 \pm 0.11 \pm 0.06\%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R39  
NODE=M055R39

NODE=M055R39;LINKAGE=HE

 $\Gamma(\rho^- K^+ \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.50±0.12±0.01</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.54 \pm 0.11 \pm 0.07 \pm 0.03\%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^- K^+ \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R40  
NODE=M055R40

NODE=M055R40;LINKAGE=HE

 $\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.23±0.06±0.01</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.25 \pm 0.06 \pm 0.03 \pm 0.02\%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R41  
NODE=M055R41

NODE=M055R41;LINKAGE=HE

 $\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.112±0.034±0.003</b>	141.3	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

<sup>1</sup> HE 08B reports  $0.12 \pm 0.03 \pm 0.02 \pm 0.01\%$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R42  
NODE=M055R42

NODE=M055R42;LINKAGE=HE

 $\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.9±2.9±0.2</b>	19.8±7.7	<sup>1</sup> ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1}\gamma$

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$   $= (0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R05  
NODE=M055R05

NODE=M055R05;LINKAGE=AB

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.2±1.0±0.1</b>	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ATHAR 07 reports  $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R25  
NODE=M055R25

NODE=M055R25;LINKAGE=AT

 $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID
<b>7.0±0.6 OUR FIT</b>	

NODE=M055R17  
NODE=M055R17

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.98 ± 0.37 ± 0.02</b>	22	<sup>1</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R09  
NODE=M055R09

<sup>1</sup> ABLIKIM 06R reports  $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R09;LINKAGE=AB

 $\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.43 ± 0.65 ± 0.03</b>	27	<sup>1</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R10  
NODE=M055R10

<sup>1</sup> ABLIKIM 06R reports  $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R10;LINKAGE=AB

 $\Gamma(K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 8 × 10<sup>-4</sup></b>	90	<sup>1</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R12  
NODE=M055R12

<sup>1</sup> ABLIKIM 06R reports  $< 0.9 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R12;LINKAGE=AB

 $\Gamma(K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.1 × 10<sup>-3</sup></b>	90	<sup>1</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R13  
NODE=M055R13

<sup>1</sup> ABLIKIM 06R reports  $< 2.4 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R13;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.81 ± 0.24 ± 0.04</b>	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M055R20  
NODE=M055R20

<sup>1</sup> ATHAR 07 reports  $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R20;LINKAGE=AT

 $\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.62 ± 0.23 OUR AVERAGE</b>				
4.58 ± 0.23 ± 0.11		<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
4.7 ± 0.5 ± 0.1		<sup>3</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
5.3 ± 0.9 ± 0.1	222	<sup>4</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R08  
NODE=M055R08

<sup>1</sup> From an amplitude analysis using an isobar model.  
<sup>2</sup> ABLIKIM 17K reports  $(4.67 \pm 0.03 \pm 0.23 \pm 0.16) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R08;LINKAGE=A  
NODE=M055R08;LINKAGE=B

<sup>3</sup> ATHAR 07 reports  $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R08;LINKAGE=AT

<sup>4</sup> ABLIKIM 06R reports  $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R08;LINKAGE=AB

$\Gamma(a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{26} / \Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.2 ± 0.4 OUR AVERAGE</b>				Error includes scale factor of 2.2.
3.33 ± 0.19 ± 0.08		<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
1.79 ± 0.63 ± 0.04	58	<sup>3</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R15  
NODE=M055R15

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(3.40 \pm 0.03 \pm 0.19 \pm 0.11) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R15;LINKAGE=A  
NODE=M055R15;LINKAGE=B

<sup>3</sup> ABLIKIM 06R reports  $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R15;LINKAGE=AB

 $\Gamma(a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{27} / \Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.176 ± 0.023 ± 0.004</b>	<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R72  
NODE=M055R72

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(0.18 \pm 0.01 \pm 0.02 \pm 0.01) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1320)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R72;LINKAGE=A  
NODE=M055R72;LINKAGE=B

 $\Gamma(a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{28} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.6 ± 0.7 ± 0.1</b>	<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R75  
NODE=M055R75

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(4.7 \pm 0.4 \pm 0.6 \pm 0.2) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1700)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R75;LINKAGE=A  
NODE=M055R75;LINKAGE=B

 $\Gamma(f_2(1270) \eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{29} / \Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.5 ± 0.6 ± 0.1</b>	<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R73  
NODE=M055R73

OCCUR=2

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(0.36 \pm 0.01 \pm 0.06 \pm 0.01) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270) \eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R73;LINKAGE=A  
NODE=M055R73;LINKAGE=D

 $\Gamma(f_4(2050) \eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{30} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.5 ± 0.9 ± 0.1</b>	<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R74  
NODE=M055R74

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $(2.6 \pm 0.4 \pm 0.8 \pm 0.1) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_4(2050) \eta \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R74;LINKAGE=A  
NODE=M055R74;LINKAGE=B

 $\Gamma(\pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{31} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 5 × 10<sup>-5</sup></b>	90	<sup>1,2</sup> ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R76  
NODE=M055R76

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $< 4.6 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1400)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R76;LINKAGE=A  
NODE=M055R76;LINKAGE=B

$\Gamma(\pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{32} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.5 \times 10^{-5}$	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R77  
NODE=M055R77

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $< 1.5 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1600)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R77;LINKAGE=A  
NODE=M055R77;LINKAGE=B

 $\Gamma(\pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$   $\Gamma_{33} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-6}$	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M055R78  
NODE=M055R78

<sup>1</sup> From an amplitude analysis using an isobar model.

<sup>2</sup> ABLIKIM 17K reports  $< 8 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(2015)^+ \pi^- + \text{c.c.} \rightarrow \eta \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R78;LINKAGE=A  
NODE=M055R78;LINKAGE=B

 $\Gamma(f_2(1270)\eta) / \Gamma_{\text{total}}$   $\Gamma_{34} / \Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.67 ± 0.11 OUR AVERAGE</b>				
0.63 ± 0.11 ± 0.02		1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
2.7 ± 0.8 ± 0.1	53	3 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R16  
NODE=M055R16

<sup>1</sup> ABLIKIM 17K reports  $(6.4 \pm 1.1) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R16;LINKAGE=B

<sup>2</sup> From an amplitude analysis using an isobar model.

<sup>3</sup> ABLIKIM 06R reports  $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R16;LINKAGE=D  
NODE=M055R16;LINKAGE=C

 $\Gamma(\pi^+ \pi^- \eta') / \Gamma_{\text{total}}$   $\Gamma_{35} / \Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.2 ± 0.4 ± 0.1</b>	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M055R28  
NODE=M055R28

<sup>1</sup> ATHAR 07 reports  $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R28;LINKAGE=AT

 $\Gamma(K^+ K^- \eta'(958)) / \Gamma_{\text{total}}$   $\Gamma_{36} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.75 ± 0.87</b>	310	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R64  
NODE=M055R64

<sup>1</sup> Derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.2 \pm 0.4)\%$ . Uncertainty includes both statistical and systematic contributions combined in quadrature.

NODE=M055R64;LINKAGE=A

 $\Gamma(K_0^*(1430)^+ K^- + \text{c.c.}) / \Gamma_{\text{total}}$   $\Gamma_{37} / \Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.41 ± 0.57<sup>+2.09</sup><sub>-2.71</sub></b>	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R65  
NODE=M055R65

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

NODE=M055R65;LINKAGE=A

 $\Gamma(f_0(980)\eta'(958)) / \Gamma_{\text{total}}$   $\Gamma_{38} / \Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.65 ± 0.47<sup>+1.32</sup><sub>-0.56</sub></b>	1 ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R66  
NODE=M055R66

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

NODE=M055R66;LINKAGE=A

$\Gamma(f_0(1710)\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$0.71 \pm 0.22^{+0.68}_{-0.48}$	<sup>1</sup> ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

NODE=M055R67  
NODE=M055R67

NODE=M055R67;LINKAGE=A

 $\Gamma(f_2'(1525)\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{40}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$0.92 \pm 0.23^{+0.55}_{-0.51}$	<sup>1</sup> ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

<sup>1</sup> Normalized to  $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$  branching fraction.

NODE=M055R68  
NODE=M055R68

NODE=M055R68;LINKAGE=A

 $\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{41}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$0.35 \pm 0.09$		ABLIKIM	18D	BES3 $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 <sup>1</sup> ABLIKIM 11D BES3  $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 11D reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 6.0 \times 10^{-7}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R18  
NODE=M055R18

NODE=M055R18;LINKAGE=BR

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{42}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
$32 \pm 21$	<sup>1</sup> TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

NODE=M055R9  
NODE=M055R9

NODE=M055R9;LINKAGE=T

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$   $\Gamma_{43}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.44 \pm 0.36 \pm 0.03$	$28.4 \pm 5.5$	<sup>1,2</sup> ABLIKIM	04H	BES $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

<sup>1</sup> ABLIKIM 04H reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Assumes  $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$ .

NODE=M055R26  
NODE=M055R26

NODE=M055R26;LINKAGE=AB

NODE=M055R26;LINKAGE=AL

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{44}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-4}$	90	$3.2 \pm 2.4$	<sup>1</sup> ABLIKIM	050	BES2 $\psi(2S) \rightarrow \chi_{c1} \gamma$

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R06  
NODE=M055R06

NODE=M055R06;LINKAGE=AB

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{45}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.35 \pm 0.10 \pm 0.01$	22	<sup>1</sup> ABLIKIM	19AA	BES3 $\psi(2S) \rightarrow \gamma 4K_S^0$

<sup>1</sup> Using  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ . ABLIKIM 19AA reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (3.4 \pm 0.9 \pm 0.3) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value..

NODE=M055R82  
NODE=M055R82

NODE=M055R82;LINKAGE=A

 $\Gamma(K^+ K^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{46}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID
$0.54 \pm 0.11$ OUR FIT	

NODE=M055R14  
NODE=M055R14

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.41 \pm 0.15 \pm 0.01$	17	<sup>1</sup> ABLIKIM	06T	BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> ABLIKIM 06T reports  $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R07  
NODE=M055R07

NODE=M055R07;LINKAGE=AB

$\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**3.27±0.28±0.46**

DOCUMENT ID

TECN

COMMENT

ABLIKIM 15M BES3  $\psi(2S) \rightarrow \gamma \chi_{c1}$  $\Gamma_{48}/\Gamma$ NODE=M055R69  
NODE=M055R69 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**1.62±0.12±0.28**

DOCUMENT ID

TECN

COMMENT

ABLIKIM 15M BES3  $\psi(2S) \rightarrow \gamma \chi_{c1}$  $\Gamma_{49}/\Gamma$ NODE=M055R70  
NODE=M055R70 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.75±0.06±0.08**

EVTS

373

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> ABLIKIM 13B BES3  $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$ <sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$ . $\Gamma_{50}/\Gamma$ NODE=M055R62  
NODE=M055R62

NODE=M055R62;LINKAGE=A

 $\Gamma(\omega \omega)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**5.7±0.7±0.1**

EVTS

597

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> ABLIKIM 11K BES3  $\psi(2S) \rightarrow \gamma$  hadrons<sup>1</sup> ABLIKIM 11K reports  $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma_{51}/\Gamma$ NODE=M055R49  
NODE=M055R49

NODE=M055R49;LINKAGE=AL

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.78±0.04±0.08**

EVTS

628

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> ABLIKIM 13B BES3  $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$ <sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$ . $\Gamma_{52}/\Gamma$ NODE=M055R61  
NODE=M055R61

NODE=M055R61;LINKAGE=A

 $\Gamma(\omega \phi)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**0.27±0.04±0.01**

EVTS

105

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> ABLIKIM 19J BES3  $\psi(2S) \rightarrow \gamma$  hadrons

••• We do not use the following data for averages, fits, limits, etc. •••

0.21±0.06±0.01

15

<sup>2,3</sup> ABLIKIM

11K

BES3

 $\psi(2S) \rightarrow \gamma$  hadrons<sup>1</sup> ABLIKIM 19J reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (2.67 \pm 0.31 \pm 0.27) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.<sup>2</sup> ABLIKIM 11K reports  $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.<sup>3</sup> Superseded by ABLIKIM 19J. $\Gamma_{53}/\Gamma$ NODE=M055R50  
NODE=M055R50

NODE=M055R50;LINKAGE=A

NODE=M055R50;LINKAGE=AL

NODE=M055R50;LINKAGE=B

 $\Gamma(\phi \phi)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**4.2±0.5±0.1**

EVTS

366

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> ABLIKIM 11K BES3  $\psi(2S) \rightarrow \gamma$  hadrons<sup>1</sup> ABLIKIM 11K reports  $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \phi \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. $\Gamma_{54}/\Gamma$ NODE=M055R51  
NODE=M055R51

NODE=M055R51;LINKAGE=AL

 $\Gamma(\phi \phi \eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**3.0±0.4±0.2**

EVTS

83.6

DOCUMENT ID

TECN

COMMENT

<sup>1</sup> ABLIKIM 20B BES3  $\psi(2S) \rightarrow \gamma \phi \phi \eta$ <sup>1</sup> ABLIKIM 20B reports  $(2.96 \pm 0.43 \pm 0.22) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \phi \phi \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . $\Gamma_{55}/\Gamma$ NODE=M055R85  
NODE=M055R85

NODE=M055R85;LINKAGE=A

 $\Gamma(\rho \bar{\rho})/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**0.760±0.034 OUR FIT**

DOCUMENT ID

 $\Gamma_{56}/\Gamma$ NODE=M055R11  
NODE=M055R11

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.155±0.018 OUR AVERAGE**

0.163±0.019±0.004

0.112±0.047±0.003

DOCUMENT ID TECN COMMENT

1 ONYISI 10 CLE3  $\psi(2S) \rightarrow \gamma p\bar{p}X$   
 2 ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ONYISI 10 reports  $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ATHAR 07 reports  $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{57}/\Gamma$ NODE=M055R21  
NODE=M055R21

NODE=M055R21;LINKAGE=ON

NODE=M055R21;LINKAGE=AT

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.145±0.024±0.004**

CL% DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.15 90 2 ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ 

<sup>1</sup> ONYISI 10 reports  $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ATHAR 07 reports  $< 0.16 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma_{58}/\Gamma$ NODE=M055R27  
NODE=M055R27

NODE=M055R27;LINKAGE=ON

NODE=M055R27;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.212±0.030±0.005**

DOCUMENT ID TECN COMMENT

1 ONYISI 10 reports  $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{59}/\Gamma$ NODE=M055R43  
NODE=M055R43

NODE=M055R43;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ 

VALUE CL%

**<1.7 × 10<sup>-5</sup>**

90

DOCUMENT ID TECN COMMENT

1 ABLIKIM 11F BES3  $\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$   
<sup>1</sup> ABLIKIM 11F reports  $< 1.82 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma_{60}/\Gamma$ NODE=M055R48  
NODE=M055R48

NODE=M055R48;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.50±0.19 OUR EVALUATION****0.50±0.19 OUR AVERAGE**

0.46±0.12±0.15

1.08±0.77±0.05

DOCUMENT ID TECN COMMENT

Treating systematic error as correlated.

1 BAI 99B BES  $\psi(2S) \rightarrow \gamma\chi_{c1}$   
 1 TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c1}$

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ .

 $\Gamma_{61}/\Gamma$ NODE=M055R7  
NODE=M055R7

→ UNCHECKED ←

NODE=M055R7;LINKAGE=X2

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ 

VALUE CL%

**<5 × 10<sup>-4</sup>**

90

DOCUMENT ID TECN COMMENT

1 HE 08B CLEO  $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ 

<sup>1</sup> HE 08B reports  $< 0.05 \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

 $\Gamma_{62}/\Gamma$ NODE=M055R38  
NODE=M055R38

NODE=M055R38;LINKAGE=HE

$\Gamma(\rho\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$  $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.27±0.22±0.03</b>	82 ± 9	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma\rho\bar{p}K^+K^-$
<sup>1</sup> ABLIKIM 11F reports $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R45  
NODE=M055R45

NODE=M055R45;LINKAGE=AB

 $\Gamma(\rho\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{64}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4.5</b>	90	<sup>1</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
<sup>1</sup> Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma)$ $(9.1 \pm 0.6)\%$ .				

NODE=M055R02  
NODE=M055R02

NODE=M055R;LINKAGE=AB

 $\Gamma(\rho\bar{n}\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{65}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8±0.5±0.1</b>	1412	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\rho\bar{n}\pi^-$
<sup>1</sup> ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.37 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R56  
NODE=M055R56

NODE=M055R56;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.9±0.5±0.1</b>	1625	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$
<sup>1</sup> ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.38 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R57  
NODE=M055R57

NODE=M055R57;LINKAGE=AL

 $\Gamma(\rho\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.3±1.1±0.2</b>	1082	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\rho\bar{n}\pi^-\pi^0$
<sup>1</sup> ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (1.00 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R58  
NODE=M055R58

NODE=M055R58;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.1±1.1±0.2</b>	1261	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$
<sup>1</sup> ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.98 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R59  
NODE=M055R59

NODE=M055R59;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$  $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>1.27±0.08 OUR FIT</b>	

NODE=M055R23  
NODE=M055R23 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>29±5±1</b>		105	<sup>1</sup> ABLIKIM	12i BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••					
<150		90	<sup>2</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
<sup>1</sup> ABLIKIM 12i reports $(31.1 \pm 3.4 \pm 3.9) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
<sup>2</sup> Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma)$ $(9.1 \pm 0.6)\%$ .					

NODE=M055R01  
NODE=M055R01

NODE=M055R01;LINKAGE=AL

NODE=M055R01;LINKAGE=AB



$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-(\text{non-resonant}))/\Gamma_{\text{total}}$  $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$25 \pm 6 \pm 1$	13	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M055R19  
 NODE=M055R19

<sup>1</sup> ABLIKIM 12I reports  $(26.2 \pm 5.5 \pm 3.3) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R19;LINKAGE=AL

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{73}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-4}$	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

NODE=M055R52  
 NODE=M055R52

<sup>1</sup> ABLIKIM 12I reports  $< 14 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R52;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{74}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 13$	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

NODE=M055R53  
 NODE=M055R53

<sup>1</sup> ABLIKIM 12I reports  $< 14 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R53;LINKAGE=AL

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 0.4$ OUR AVERAGE	Error includes scale factor of 1.2.			

NODE=M055R22  
 NODE=M055R22

$9.2^{+2.8}_{-2.4} \pm 0.4$	24	<sup>1</sup> LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
$4.2 \pm 0.4 \pm 0.1$	3k	<sup>2,3</sup> ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
$3.1 \pm 0.9 \pm 0.1$		<sup>4</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> LU 19 reports  $(9.15^{+2.63}_{-2.25} \pm 0.86) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(1P)K^+)]$  assuming  $B(B^+ \rightarrow \chi_{c1}(1P)K^+) = (4.79 \pm 0.23) \times 10^{-4}$ , which we rescale to our best value  $B(B^+ \rightarrow \chi_{c1}(1P)K^+) = (4.74 \pm 0.22) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R22;LINKAGE=A

<sup>2</sup> ABLIKIM 13D reports  $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R22;LINKAGE=AB

<sup>3</sup> Using  $B(\Lambda \rightarrow p\pi^-) = 63.9\%$ .

NODE=M055R22;LINKAGE=LB  
 NODE=M055R22;LINKAGE=AT

<sup>4</sup> ATHAR 07 reports  $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{76}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.66 \pm 0.12 \pm 0.12$	399	<sup>1</sup> ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma nK_S^0\bar{\Lambda} + \text{c.c.}$

NODE=M055R89  
 NODE=M055R89

<sup>1</sup> ABLIKIM 21AV reports  $(1.66 \pm 0.12 \pm 0.12) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow nK_S^0\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0975 \pm 0.0024$ . Also uses  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = (63.9 \pm 0.5)\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$ .

NODE=M055R89;LINKAGE=A

 $\Gamma(K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.9 \pm 0.7 \pm 0.1$	328	<sup>1</sup> ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+}\bar{p}\Lambda$

NODE=M055R86  
 NODE=M055R86

<sup>1</sup> ABLIKIM 19AU reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (4.8 \pm 0.5 \pm 0.4) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R86;LINKAGE=F

$\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{78}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.71 \pm 0.44 \pm 0.04</math></b>	$48 \pm 10$	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M055R46  
 NODE=M055R46

<sup>1</sup> ABLIKIM 11F reports  $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R46;LINKAGE=AB

 $\Gamma(\Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}$  $\Gamma_{79}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>&lt;9 \times 10^{-5}</math></b>	90	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M055R47  
 NODE=M055R47

<sup>1</sup> ABLIKIM 11F reports  $< 1.00 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R47;LINKAGE=AB

 $\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4.2 \pm 0.6 \pm 0.1</math></b>		103	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

NODE=M055R32  
 NODE=M055R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6$	90		<sup>2</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
$<4$	90	$3.8 \pm 2.5$	<sup>3</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.41 \pm 0.05 \pm 0.03) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R32;LINKAGE=B

<sup>2</sup> ABLIKIM 13H reports  $< 0.62 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R32;LINKAGE=AB

<sup>3</sup> NAIK 08 reports  $< 0.44 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R32;LINKAGE=NA

 $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$  $\Gamma_{83}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.6 \pm 0.6 \pm 0.1</math></b>		59	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

NODE=M055R33  
 NODE=M055R33

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<8$	90		<sup>2</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
$<6$	90	$4.3 \pm 2.3$	<sup>3</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.35 \pm 0.06 \pm 0.02) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R33;LINKAGE=B

<sup>2</sup> ABLIKIM 13H reports  $< 0.87 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R33;LINKAGE=AB

<sup>3</sup> NAIK 08 reports  $< 0.65 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R33;LINKAGE=NA

 $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$  $\Gamma_{84}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5.7 \pm 1.4 \pm 0.6</math></b>	214	<sup>1</sup> ABLIKIM	20I BES3	$\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

NODE=M055R87  
 NODE=M055R87

<sup>1</sup> ABLIKIM 20I reports  $(5.7 \pm 1.4 \pm 0.6) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ .

NODE=M055R87;LINKAGE=A

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9 \times 10^{-5}$	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
<sup>1</sup> ABLIKIM 12I reports $< 10 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .				

NODE=M055R54  
NODE=M055R54

NODE=M055R54;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$   $\Gamma_{86}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5 \times 10^{-5}$	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
<sup>1</sup> ABLIKIM 12I reports $< 5.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .				

NODE=M055R55  
NODE=M055R55

NODE=M055R55;LINKAGE=AL

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.35 \pm 0.24 \pm 0.03$	49	<sup>1</sup> ABLIKIM	15I BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$
<sup>1</sup> ABLIKIM 15I reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $= (1.32 \pm 0.20 \pm 0.12) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M055R71  
NODE=M055R71

NODE=M055R71;LINKAGE=A

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.5 \pm 1.1 \pm 0.6$		325	<sup>1</sup> ABLIKIM	220 BES3	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<6$	90	$1.7 \pm 2.4$	<sup>2</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
<sup>1</sup> ABLIKIM 220 reports $(0.75 \pm 0.11 \pm 0.06) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ .					
<sup>2</sup> NAIK 08 reports $< 0.60 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .					

NODE=M055R34  
NODE=M055R34

NODE=M055R34;LINKAGE=A

NODE=M055R34;LINKAGE=NA

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.60 \pm 0.06$ OUR AVERAGE					
[( $0.80 \pm 0.21$ ) $\times 10^{-4}$ OUR 2022 AVERAGE]					
$0.58 \pm 0.04 \pm 0.05$		692	<sup>1</sup> ABLIKIM	220 BES3	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
$0.80 \pm 0.21 \pm 0.02$		$16.4 \pm 4.3$	<sup>2</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 3.4$	90		<sup>3</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
<sup>1</sup> ABLIKIM 220 reports $(0.58 \pm 0.04 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ .					
<sup>2</sup> NAIK 08 reports $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
<sup>3</sup> Using $B(\psi(2S) \rightarrow \chi_{c1} \gamma)$ $(9.1 \pm 0.6)\%$ .					

NODE=M055R03  
NODE=M055R03

NEW

NODE=M055R03;LINKAGE=A

NODE=M055R03;LINKAGE=NA

NODE=M055R03;LINKAGE=AB

 $[\Gamma(\pi^+ \pi^-) + \Gamma(K^+ K^-)]/\Gamma_{\text{total}}$   $\Gamma_{90}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<21 \times 10^{-4}$		<sup>1</sup> FELDMAN	77 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<38 \times 10^{-4}$	90	<sup>1</sup> BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$
<sup>1</sup> Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the $\psi(2S)$ decay.				

NODE=M055R2  
NODE=M055R2

NODE=M055R2;LINKAGE=T

$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{91}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-5}$	90	<sup>1</sup> ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

NODE=M055R04  
 NODE=M055R04

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$   
 $< 0.6 \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R04;LINKAGE=AB

 $\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{92}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-3}$	90	<sup>1,2</sup> ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R63  
 NODE=M055R63

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<4.4 \times 10^{-3}$	90	<sup>1,3</sup> ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$
-----------------------	----	------------------------	----------	---

OCCUR=2

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$ .

NODE=M055R63;LINKAGE=A

<sup>2</sup> Using the  $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

NODE=M055R63;LINKAGE=B

<sup>3</sup> Using the  $\eta_c \rightarrow K^+ K^- \pi^0$  decays.

NODE=M055R63;LINKAGE=C

## RADIATIVE DECAYS

 $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$  $\Gamma_{93}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R1  
 NODE=M055R1

**34.3 ± 1.0 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

$34.75 \pm 0.11 \pm 1.70$	1.9M	<sup>1</sup> ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
$37.9 \pm 0.8 \pm 2.1$		<sup>2</sup> ADAM	05A CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Not independent from  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))$  and the product  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))$  also measured in ABLIKIM 17U.

NODE=M055R1;LINKAGE=A

<sup>2</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c1} \rightarrow \gamma \gamma J/\psi)$  from ADAM 05A and  $B(\psi(2S) \rightarrow \gamma \chi_{c1})$  from ATHAR 04.

NODE=M055R1;LINKAGE=AD

 $\Gamma(\gamma \rho^0)/\Gamma_{\text{total}}$  $\Gamma_{94}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R29  
 NODE=M055R29

**216 ± 17 OUR AVERAGE**

$215 \pm 22 \pm 5$	$432 \pm 25$	<sup>1</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
$217 \pm 24 \pm 5$	$186 \pm 15$	<sup>2</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$

NODE=M055R29;LINKAGE=AB

<sup>1</sup> ABLIKIM 11E reports  $(228 \pm 13 \pm 22) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R29;LINKAGE=BE

<sup>2</sup> BENNETT 08A reports  $(243 \pm 19 \pm 22) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\gamma \omega)/\Gamma_{\text{total}}$  $\Gamma_{95}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R30  
 NODE=M055R30

**68 ± 8 OUR AVERAGE**

$66 \pm 9 \pm 2$	$136 \pm 14$	<sup>1</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \omega$
$74 \pm 17 \pm 2$	$39 \pm 7$	<sup>2</sup> BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \omega$

NODE=M055R30;LINKAGE=AB

<sup>1</sup> ABLIKIM 11E reports  $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R30;LINKAGE=BE

<sup>2</sup> BENNETT 08A reports  $(83 \pm 15 \pm 12) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>24±5±1</b>		43 ± 9	<sup>1</sup> ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$

 $\Gamma_{96}/\Gamma$ NODE=M055R31  
NODE=M055R31

• • • We do not use the following data for averages, fits, limits, etc. • • •

<23 90 5.2 ± 3.1 <sup>2</sup> BENNETT 08A CLEO  $\psi(2S) \rightarrow \gamma\gamma\phi$ 

<sup>1</sup> ABLIKIM 11E reports  $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R31;LINKAGE=AB

<sup>2</sup> BENNETT 08A reports  $< 26 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$ .

NODE=M055R31;LINKAGE=BE

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6.3 × 10<sup>-6</sup></b>	90	ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$

 $\Gamma_{97}/\Gamma$ NODE=M055R3  
NODE=M055R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.5 × 10<sup>-5</sup> 90 ECKLUND 08A CLEO  $\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow 3\gamma$ <150 × 10<sup>-5</sup> 90 <sup>1</sup> YAMADA 77 DASP  $e^+e^- \rightarrow 3\gamma$ 

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$ . The errors do not contain the uncertainty in the  $\psi(2S)$  decay.

NODE=M055R;LINKAGE=T1

 $\Gamma(e^+e^- J/\psi(1S))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

 $\Gamma_{98}/\Gamma$ NODE=M055R79  
NODE=M055R79

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.65±0.23±0.09 1.9k <sup>1,2</sup> ABLIKIM 17I BES3  $\psi(2S) \rightarrow \gamma e^+e^- J/\psi$ 

<sup>1</sup> ABLIKIM 17I reports  $(3.73 \pm 0.09 \pm 0.25) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c1}(1P) \rightarrow e^+e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R79;LINKAGE=B

<sup>2</sup> Not independent from other measurements reported by ABLIKIM 17I

NODE=M055R79;LINKAGE=C

 $\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.1±0.3±0.5</b>	1.9k	<sup>1</sup> ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^- \gamma J/\psi$

 $\Gamma_{98}/\Gamma_{93}$ NODE=M055R80  
NODE=M055R80

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

NODE=M055R80;LINKAGE=A

 $\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S))$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.73±0.51±0.50</b>	222	ABLIKIM	19Z BES3	$\psi(2S) \rightarrow \gamma\chi_c \rightarrow \gamma(\mu^+\mu^- J/\psi)$

 $\Gamma_{99}/\Gamma_{98}$ NODE=M055R84  
NODE=M055R84 $\chi_{c1}(1P)$  CROSS-PARTICLE BRANCHING RATIOS

NODE=M055230

 $\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow$  $J/\psi(1S)\pi^+\pi^-)$  $\Gamma_{56}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M055B1  
NODE=M055B1**2.14±0.10 OUR FIT****1.1 ± 1.0** <sup>1</sup> BAI 98I BES  $\psi(2S) \rightarrow \gamma\chi_{c1} \rightarrow \gamma p\bar{p}$ 

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c1} \rightarrow p\bar{p})$  reported in BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.7 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M055B;LINKAGE=J2

$$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{69}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**12.4±0.8 OUR FIT****12.3±0.9 OUR AVERAGE** Error includes scale factor of 1.2.12.8±0.6±0.6 528 ABLIKIM 21L BES3  $\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+$ 10.5±1.6±0.6 46 <sup>1</sup> NAIK 08 CLEO  $\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.2±1.0±0.9 136 <sup>2,3</sup> ABLIKIM 13H BES3  $\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$ <sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$ .<sup>2</sup> Superseded by ABLIKIM 21L<sup>3</sup> Calculated by us. ABLIKIM 13H reports  $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (12.2 \pm 1.1 \pm 1.1) \times 10^{-5}$  from a measurement of  $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c1})$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.2 \pm 0.4)\%$ .NODE=M055B10  
NODE=M055B10

NODE=M055B10;LINKAGE=NA

NODE=M055B10;LINKAGE=A  
NODE=M055B10;LINKAGE=AB

$$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{69}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.58±0.22 OUR FIT**7.1  $\begin{matrix} +2.8 \\ -2.4 \end{matrix} \pm 1.3$  9.0  $\begin{matrix} +3.5 \\ -3.1 \end{matrix}$  <sup>1</sup> BAI 03E BES  $\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$ <sup>1</sup> BAI 03E reports  $[B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c1}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.52}_{-0.46} \pm 0.25)\%$ . We calculate from this measurement the presented value using  $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$  and  $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$ .NODE=M055B11  
NODE=M055B11

NODE=M055B11;LINKAGE=BA

$$\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{72}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.72±1.34±0.65** 21 ABLIKIM 22A0 BES3  $\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+ \gamma\gamma$ NODE=M055R90  
NODE=M055R90

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{93}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma^{\psi(2S)}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.34 ±0.06 OUR FIT****3.24 ±0.16 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.3.518±0.010±0.120 143k <sup>1</sup> ABLIKIM 17N BES3  $\psi(2S) \rightarrow \gamma\gamma J/\psi$ 3.442±0.010±0.132 1.9M ABLIKIM 17U BES3  $e^+e^- \rightarrow \gamma X$ 2.81 ±0.05 ±0.23 13k BAI 04I BES2  $\psi(2S) \rightarrow J/\psi\gamma\gamma$ 2.56 ±0.12 ±0.20 GAISER 86 CBAL  $\psi(2S) \rightarrow \gamma X$ 2.78 ±0.30 <sup>2</sup> OREGLIA 82 CBAL  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 2.2 ±0.5 <sup>3</sup> BRANDELIK 79B DASP  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 2.9 ±0.5 <sup>3</sup> BARTEL 78B CNTR  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 5.0 ±1.5 <sup>4</sup> BIDDICK 77 CNTR  $e^+e^- \rightarrow \gamma X$ 2.8 ±0.9 <sup>2</sup> WHITAKER 76 MRK1  $e^+e^-$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.377±0.009±0.183 142k <sup>5</sup> ABLIKIM 120 BES3  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 3.56 ±0.03 ±0.12 24.9k <sup>6</sup> MENDEZ 08 CLEO  $\psi(2S) \rightarrow \gamma\chi_{c1}$ 3.44 ±0.06 ±0.13 3.7k <sup>7</sup> ADAM 05A CLEO Repl. by MENDEZ 08<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$ .<sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$ .<sup>3</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ .<sup>4</sup> Assumes isotropic gamma distribution.<sup>5</sup> Superseded by ABLIKIM 17N.<sup>6</sup> Not independent from other measurements of MENDEZ 08.<sup>7</sup> Not independent from other values reported by ADAM 05A.NODE=M055B2  
NODE=M055B2

NODE=M055B2;LINKAGE=A

NODE=M055B;LINKAGE=3Q

NODE=M055B;LINKAGE=2Q


NODE=M055B;LINKAGE=EA

NODE=M055B2;LINKAGE=B

NODE=M055B2;LINKAGE=ME

NODE=M055B;LINKAGE=AD

WEIGHTED AVERAGE  
3.24±0.16 (Error scaled by 2.1)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{total} \text{ (units } 10^{-2}\text{)}$$

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{93}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)} = \Gamma_{93}/\Gamma \times \Gamma_{167}^{\psi(2S)}/(\Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \Gamma_{14}^{\psi(2S)} + 0.343\Gamma_{167}^{\psi(2S)} + 0.190\Gamma_{168}^{\psi(2S)})$$

NODE=M055B7

NODE=M055B7

NODE=M055B7

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.43±0.10 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.70±0.04±0.15	24.9k	<sup>1</sup> MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
5.77±0.10±0.12	3.7k	ADAM	05A CLEO	Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

NODE=M055B7;LINKAGE=ME

$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{93}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

NODE=M055B3  
NODE=M055B3

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**9.63±0.17 OUR FIT**

**10.15±0.28 OUR AVERAGE**

10.17±0.07±0.27	24.9k	MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
12.6 ±0.3 ±3.8	3k	<sup>1</sup> ABLIKIM	04B BES	$\psi(2S) \rightarrow J/\psi X$
8.5 ±2.1		<sup>2</sup> HIMEL	80 MRK2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10.24±0.17±0.23	3.7k	<sup>3</sup> ADAM	05A CLEO	Repl. by MENDEZ 08

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> The value for  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$  quoted in HIMEL 80 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$  and  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

<sup>3</sup> Not independent from other values reported by ADAM 05A.

NODE=M055B;LINKAGE=AB  
NODE=M055B;LINKAGE=J3

NODE=M055B3;LINKAGE=AD

$$\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + c.c.)/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{total}$$

$$\Gamma_{19}/\Gamma \times \Gamma_{167}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

NODE=M055B16  
NODE=M055B16

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**6.8±0.5 OUR FIT**

**7.2±0.6 OUR AVERAGE**

7.3±0.5±0.5	<sup>1</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
7.0±0.5±0.9	<sup>2</sup> ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow \bar{K}^0 K^+ \pi^- + c.c.)$  reported by ATHAR 07 was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$ .

<sup>2</sup> Calculated by us. ABLIKIM 06R reports  $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ . We use  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$ .

NODE=M055B16;LINKAGE=AT

NODE=M055B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + c.c.) / \Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{total}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{19} / \Gamma \times \Gamma_{167}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

NODE=M055B17  
NODE=M055B17

VALUE (units 10<sup>-4</sup>) DOCUMENT ID TECN COMMENT  
**19.6±1.6 OUR FIT**  
**13.2±2.4±3.2** <sup>1</sup>BAI 99B BES  $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M055B17;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{total}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{46} / \Gamma \times \Gamma_{167}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

NODE=M055B14  
NODE=M055B14

VALUE (units 10<sup>-4</sup>) EVTS DOCUMENT ID TECN COMMENT  
**0.53±0.11 OUR FIT**  
**0.61±0.11±0.08** 54 <sup>1</sup>ABLIKIM 06T BES2  $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ .

NODE=M055B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{total}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{46} / \Gamma \times \Gamma_{167}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

NODE=M055B15  
NODE=M055B15

VALUE (units 10<sup>-4</sup>) DOCUMENT ID TECN COMMENT  
**1.52±0.31 OUR FIT**  
**1.13±0.40±0.29** <sup>1</sup>BAI 99B BES  $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M055B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p \bar{p}) / \Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{total}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \times \frac{\Gamma_{56} / \Gamma \times \Gamma_{167}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$


NODE=M055B6  
NODE=M055B6

VALUE (units 10<sup>-6</sup>) EVTS DOCUMENT ID TECN COMMENT  
**7.41±0.35 OUR FIT**  
**7.8 ±0.6 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

7.9 ±0.4 ±0.3	453	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p \bar{p}$
8.2 ±0.7 ±0.4	141 ± 13	<sup>1</sup> NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
4.8 <sup>+1.4</sup> <sub>-1.3</sub> ±0.6	18.2 <sup>+5.5</sup> <sub>-4.9</sub>	BAI	04F	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}(1P) \rightarrow \gamma p \bar{p}$

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c1} \rightarrow p \bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$ .

NODE=M055B6;LINKAGE=NA





$$\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{81} / \Gamma \times \Gamma_{167}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.49 ± 0.09 ± 0.07</b>	258	<sup>1</sup> ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

NODE=M055B01  
NODE=M055B01

<sup>1</sup> Calculated by us. ABLIKIM 19BB reports  $B(\chi_{c1} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (1.53 \pm 0.10 \pm 0.08) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.75 \pm 0.24)\%$  and other branching fractions from PDG 18.

NODE=M055B01;LINKAGE=A

$$\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{82} / \Gamma \times \Gamma_{167}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.42 ± 0.07 ± 0.06</b>	493	<sup>1</sup> ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$

NODE=M055R88  
NODE=M055R88

<sup>1</sup> Calculated by us. ABLIKIM 20AE reports  $B(\chi_{c1} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (1.46 \pm 0.07 \pm 0.07) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c1}^0) = (9.75 \pm 0.24)\%$  and other branching fractions from PDG 20.

NODE=M055R88;LINKAGE=A

### MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

NODE=M055240

$a_2 = M2 / \sqrt{E1^2 + M2^2}$  Magnetic quadrupole fractional transition amplitude


NODE=M055A1  
NODE=M055A1

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-6.7 ± 0.9 OUR AVERAGE</b>				Error includes scale factor of 2.6. See the ideogram below.
-7.40 ± 0.33 ± 0.34	164k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
-6.26 ± 0.63 ± 0.24	39k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.2 ± 3.2 ± 0.4	2090	AMBROGIANI	02 E835	$p \bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi \gamma$
-0.2 <sup>+0.8</sup> <sub>-2.0</sub>	921	OREGLIA	82 CBAL	$\psi(2S) \rightarrow \chi_{c1} \gamma \rightarrow J/\psi \gamma \gamma$

<sup>1</sup> Correlated with  $b_2$  with correlation coefficient  $\rho_{a_2 b_2} = 0.133$ .

NODE=M055A1;LINKAGE=A

WEIGHTED AVERAGE  
-6.7 ± 0.9 (Error scaled by 2.6)



### MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma \chi_{c1}(1S)$ RADIATIVE DECAY

NODE=M055250

$b_2 = M2 / \sqrt{E1^2 + M2^2}$  Magnetic quadrupole fractional transition amplitude

NODE=M055QB2  
NODE=M055QB2

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.5 ± 0.4 OUR AVERAGE</b>				
2.29 ± 0.39 ± 0.27	164k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
2.76 ± 0.73 ± 0.23	39k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
7.7 <sup>+5.0</sup> <sub>-4.5</sub>	921	OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>1</sup> Correlated with  $a_2$  with correlation coefficient  $\rho_{a_2 b_2} = 0.133$ .

NODE=M055QB2;LINKAGE=A

**MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS**

$$\psi(2S) \rightarrow \gamma \chi_{c1}(1S) \text{ and } \chi_{c1} \rightarrow \gamma J/\psi(1S)$$

 **$a_2/b_2$  Magnetic quadrupole transition amplitude ratio**

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$-2.27^{+0.57}_{-0.99}$	39k	<sup>1</sup> ARTUSO	09	CLEO $\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>1</sup> Statistical and systematic errors combined. Not independent of  $a_2(\chi_{c1})$  and  $b_2(\chi_{c1})$  values from ARTUSO 09.

NODE=M055260

NODE=M055QAR  
NODE=M055QAR

NODE=M055QAR;LINKAGE=AR

 **$\chi_{c1}(1P)$  REFERENCES**

Author	Year	Journal	Volume	Page	RefID
ABLIKIM	22AF	PRL	129	122001	REFID=61878
ABLIKIM	22AO	PR	D106	072004	REFID=61887
ABLIKIM	22O	JHEP	2206	074	REFID=61652
ABLIKIM	22Q	PR	D106	032014	REFID=61663
ABLIKIM	21AV	JHEP	2111	217	REFID=61465
ABLIKIM	21L	PR	D103	112004	REFID=61117
ABLIKIM	20AE	PR	D102	092006	REFID=60733
ABLIKIM	20B	PR	D101	012012	REFID=60212
ABLIKIM	20I	PR	D101	092002	REFID=60303
PDG	20	PTEP	2020	083C01	REFID=60676
ABLIKIM	19AA	PR	D99	052008	REFID=59844
ABLIKIM	19AU	PR	D100	052010	REFID=59996
ABLIKIM	19BB	PR	D100	092006	REFID=60026
ABLIKIM	19J	PR	D99	012015	REFID=59606
ABLIKIM	19Z	PR	D99	051101	REFID=59837
LU	19	PR	D99	032003	REFID=59614
ABLIKIM	18D	PRL	121	022001	REFID=58849
ABLIKIM	18V	PR	D97	052011	REFID=58990
PDG	18	PR	D98	030001	REFID=59304
AAIJ	17BB	EPJ	C77	609	REFID=58191
AAIJ	17BI	PRL	119	221801	REFID=58278
ABLIKIM	17AE	PR	D96	092007	REFID=58310
ABLIKIM	17I	PRL	118	221802	REFID=57931
ABLIKIM	17K	PR	D95	032002	REFID=57953
ABLIKIM	17N	PR	D95	072004	REFID=57978
ABLIKIM	17U	PR	D96	032001	REFID=58026
PDG	16	CP	C40	100001	REFID=57140
ABLIKIM	15I	PR	D91	092006	REFID=56774
ABLIKIM	15M	PR	D91	112008	REFID=56778
ABLIKIM	14J	PR	D89	074030	REFID=55901
ABLIKIM	13B	PR	D87	012002	REFID=54877
ABLIKIM	13D	PR	D87	012007	REFID=54879
ABLIKIM	13H	PR	D87	032007	REFID=54953
ABLIKIM	13V	PR	D88	112001	REFID=55583
ABLIKIM	12I	PR	D86	052004	REFID=54736
ABLIKIM	12J	PR	D86	052011	REFID=54737
ABLIKIM	12O	PRL	109	172002	REFID=54742
ABLIKIM	11A	PR	D83	012006	REFID=53647
ABLIKIM	11D	PR	D83	032003	REFID=16715
ABLIKIM	11E	PR	D83	112005	REFID=16717
ABLIKIM	11F	PR	D83	112009	REFID=16719
ABLIKIM	11K	PRL	107	092001	REFID=53940
ONYISI	10	PR	D82	011103	REFID=53360
ARTUSO	09	PR	D80	112003	REFID=53206
BENNETT	08A	PRL	101	151801	REFID=52575
ECKLUND	08A	PR	D78	091501	REFID=52583
HE	08B	PR	D78	092004	REFID=52588
MENDEZ	08	PR	D78	011102	REFID=52684
NAIK	08	PR	D78	031101	REFID=52301
ATHAR	07	PR	D75	032002	REFID=51618
ABLIKIM	06D	PR	D73	052006	REFID=51049
ABLIKIM	06R	PR	D74	072001	REFID=51447
ABLIKIM	06T	PL	B642	197	REFID=51453
ABLIKIM	05G	PR	D71	092002	REFID=50756
ABLIKIM	05O	PL	B630	21	REFID=50846
ADAM	05A	PRL	94	232002	REFID=50763
ANDREOTTI	05A	NP	B717	34	REFID=50769
ABLIKIM	04B	PR	D70	012003	REFID=49741
ABLIKIM	04H	PR	D70	092003	REFID=50188
ATHAR	04	PR	D70	112002	REFID=50331
BAI	04F	PR	D69	092001	REFID=49752
BAI	04I	PR	D70	012006	REFID=49755
AULCHENKO	03	PL	B573	63	REFID=49579
BAI	03E	PR	D67	112001	REFID=49416
AMBROGIANI	02	PR	D65	052002	REFID=48552
BAI	99B	PR	D60	072001	REFID=47385
BAI	98D	PR	D58	092006	REFID=46338
BAI	98I	PRL	81	3091	REFID=46343
ARMSTRONG	92	NP	B373	35	REFID=41865
Also		PRL	68	1468	REFID=41907
BAGLIN	86B	PL	B172	455	REFID=22145
GAISER	86	PR	D34	711	REFID=22012
LEMOIGNE	82	PL	113B	509	REFID=22084
OREGLIA	82	PR	D25	2259	REFID=22120
Also		Private Comm.			REFID=22143
HIMEL	80	PRL	44	920	REFID=22119
Also		Private Comm.			REFID=22113
BRANDELIK	79B	NP	B160	426	REFID=22115
BARTEL	78B	PL	79B	492	REFID=22111
TANENBAUM	78	PR	D17	1731	REFID=22112
Also		Private Comm.			REFID=22113
BIDDICK	77	PRL	38	1324	REFID=22059
FELDMAN	77	PRPL	33C	285	REFID=22062
YAMADA	77	Hamburg Conf.		69	REFID=22064
WHITAKER	76	PRL	37	1596	REFID=22151
TANENBAUM	75	PRL	35	1323	REFID=22106

NODE=M055

**$h_c(1P)$** 

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Quantum numbers are quark model prediction,  $C = -$  established by  $\eta_c \gamma$  decay.

NODE=M144

NODE=M144

 **$h_c(1P)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3525.37±0.14 OUR AVERAGE</b>		Error includes scale factor of 1.2. [3525.38 ± 0.11 MeV OUR 2022 AVERAGE]		
3525.32±0.06±0.15	23k	ABLIKIM	22AQ BES3	$\psi(2S) \rightarrow \pi^0$ hadrons; $\pi^0 \gamma(\eta_c)$
3525.20±0.18±0.12	1282	<sup>1</sup> DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3525.8 ±0.2 ±0.2	13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3525.31±0.11±0.14	832	<sup>2,3</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
3525.40±0.13±0.18	3679	<sup>2</sup> ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
3525.6 ±0.5	92	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2(\pi^+ \pi^- \pi^0)$
3524.4 ±0.6 ±0.4	168	<sup>4</sup> ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3527 ±8	42	ANTONIAZZI	94 E705	$300 \pi^\pm, pLi \rightarrow J/\psi \pi^0 X$
3526.28±0.18±0.19	59	<sup>5</sup> ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$
3525.4 ±0.8 ±0.4	5	BAGLIN	86 SPEC	$\bar{p}p \rightarrow J/\psi X$

NODE=M144M

NODE=M144M

NEW

<sup>1</sup> Combination of exclusive and inclusive analyses for the reaction  $\psi(2S) \rightarrow \pi^0 h_c \rightarrow \pi^0 \eta_c \gamma$ . This result is the average of DOBBS 08A and ROSNER 05.

<sup>2</sup> Superseded by ABLIKIM 22AQ

<sup>3</sup> With floating width.

<sup>4</sup> Superseded by DOBBS 08A.

<sup>5</sup> Mass central value and systematic error recalculated by us according to Eq.(16) in ARMSTRONG 93B, using the value for the  $\psi(2S)$  mass from AULCHENKO 03.

NODE=M144M;LINKAGE=DO

NODE=M144M;LINKAGE=A  
NODE=M144M;LINKAGE=AB  
NODE=M144M;LINKAGE=RO  
NODE=M144M;LINKAGE=NW

 **$h_c(1P)$  WIDTH**

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.78±0.28 OUR AVERAGE</b>		[0.7 ± 0.4 MeV OUR 2022 AVERAGE]			
<b>0.78<sup>+0.27</sup><sub>-0.24</sub>±0.12</b>		23k	ABLIKIM	22AQ BES3	$\psi(2S) \rightarrow \pi^0$ hadrons; $\pi^0 \gamma(\eta_c)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.70±0.28±0.22		832	<sup>1,2</sup> ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
< 1.44	90	3679	<sup>3</sup> ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
< 1		13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
< 1.1	90	59	ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$

<sup>1</sup> Superseded by ABLIKIM 22AQ

<sup>2</sup> With floating mass.

<sup>3</sup> The central value is  $\Gamma = 0.73 \pm 0.45 \pm 0.28$  MeV.

NODE=M144W

NODE=M144W

NEW

NODE=M144W;LINKAGE=A  
NODE=M144W;LINKAGE=AL  
NODE=M144W;LINKAGE=AB

 **$h_c(1P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $J/\psi(1S) \pi^0$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_2$ $J/\psi(1S) \pi \pi$	not seen	
$\Gamma_3$ $J/\psi(1S) \pi^+ \pi^-$	< 2.7 × 10 <sup>-3</sup>	90%
$\Gamma_4$ $p \bar{p}$	< 1.7 × 10 <sup>-4</sup>	90%
$\Gamma_5$ $p \bar{p} \pi^0$	< 8 × 10 <sup>-4</sup>	90%
$\Gamma_6$ $p \bar{p} \pi^+ \pi^-$	( 3.3±0.6 ) × 10 <sup>-3</sup>	
$\Gamma_7$ $p \bar{p} \pi^0 \pi^0$	< 6 × 10 <sup>-4</sup>	90%
$\Gamma_8$ $p \bar{p} \pi^+ \pi^- \pi^0$	( 4.4±1.3 ) × 10 <sup>-3</sup>	
$\Gamma_9$ $p \bar{p} \eta$	( 7.4±2.2 ) × 10 <sup>-4</sup>	
$\Gamma_{10}$ $\pi^+ \pi^- \pi^0$	( 1.9±0.5 ) × 10 <sup>-3</sup>	
$\Gamma_{11}$ $\pi^+ \pi^- \pi^0 \eta$	( 8.3±2.4 ) × 10 <sup>-3</sup>	

NODE=M144215;NODE=M144

DESIG=1

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=10

DESIG=3

DESIG=24

DESIG=11

DESIG=13

DESIG=25

DESIG=23

DESIG=5

DESIG=14

$\Gamma_{12}$	$2\pi^+2\pi^-\pi^0$	$(9.4\pm 1.7)\times 10^{-3}$			DESIG=6
$\Gamma_{13}$	$3\pi^+3\pi^-\pi^0$	$< 1.0$	%	90%	DESIG=7
$\Gamma_{14}$	$K^+K^-\pi^+\pi^-$	$< 7$	$\times 10^{-4}$	90%	DESIG=12
$\Gamma_{15}$	$K^+K^-\pi^+\pi^-\pi^0$	$(3.8\pm 0.8)\times 10^{-3}$			DESIG=15
$\Gamma_{16}$	$K^+K^-\pi^+\pi^-\eta$	$< 2.7$	$\times 10^{-3}$	90%	DESIG=16
$\Gamma_{17}$	$K^+K^-\pi^0$	$< 6$	$\times 10^{-4}$	90%	DESIG=17
$\Gamma_{18}$	$K^+K^-\pi^0\eta$	$< 2.4$	$\times 10^{-3}$	90%	DESIG=18
$\Gamma_{19}$	$K^+K^-\eta$	$< 1.0$	$\times 10^{-3}$	90%	DESIG=19
$\Gamma_{20}$	$2K^+2K^-\pi^0$	$< 2.8$	$\times 10^{-4}$	90%	DESIG=20
$\Gamma_{21}$	$K_S^0 K^\pm \pi^\mp$	$< 6$	$\times 10^{-4}$	90%	DESIG=21
$\Gamma_{22}$	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(3.2\pm 1.0)\times 10^{-3}$			DESIG=22

### Radiative decays

$\Gamma_{23}$	$\gamma\eta$	$(4.7\pm 2.1)\times 10^{-4}$			NODE=M144;CLUMP=R
$\Gamma_{24}$	$\gamma\eta'(958)$	$(1.5\pm 0.4)\times 10^{-3}$			DESIG=9
$\Gamma_{25}$	$\gamma\eta_c(1S)$	$(57 \pm 5) \%$			DESIG=8

### $h_c(1P)$ PARTIAL WIDTHS

NODE=M144220

### $h_c(1P) \Gamma(i)\Gamma(\bar{p}p)/\Gamma(\text{total})$

NODE=M144223

### $\Gamma(\gamma\eta_c(1S)) \times \Gamma(\bar{p}p)/\Gamma_{\text{total}}$

 $\Gamma_{25}\Gamma_4/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M144G1  
NODE=M144G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.0 $\pm$ 4.5	13	<sup>1</sup> ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c\gamma$
----------------	----	------------------------	----------	-------------------------------------

<sup>1</sup> Assuming  $\Gamma = 1$  MeV.

NODE=M144G1;LINKAGE=AN

### $h_c(1P)$ BRANCHING RATIOS

NODE=M144225

### $\Gamma(J/\psi(1S)\pi^0)/\Gamma(\gamma\eta_c(1S))$

 $\Gamma_1/\Gamma_{25}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M144R24  
NODE=M144R24 $< 9 \times 10^{-4}$  90 <sup>1</sup> ABLIKIM 22N BES3  $e^+e^- \rightarrow \pi^+\pi^-h_c$ 

<sup>1</sup> ABLIKIM 22N reports  $[\Gamma(h_c(1P) \rightarrow J/\psi(1S)\pi^0)/\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))]/[B(\eta_c \rightarrow K^+K^-\pi^0)] < 7.5 \times 10^{-2}$  which we multiply by our best value  $B(\eta_c \rightarrow K^+K^-\pi^0) = 1/6 B(\eta_c(1S) \rightarrow K\bar{K}\pi) = 1/6 (7.0 \times 10^{-2})$ .

NODE=M144R24;LINKAGE=B

### $\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$

 $\Gamma_2/\Gamma_1$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M144R1  
NODE=M144R1 $< 0.18$  90 ARMSTRONG 92D E760  $\bar{p}p \rightarrow J/\psi\pi^0$ 

### $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$

 $\Gamma_3/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M144R07  
NODE=M144R07 $< 2.7 \times 10^{-3}$  (CL = 90%) [ $< 2.3 \times 10^{-3}$  (CL = 90%) OUR 2022 BEST LIMIT] $< 2.7 \times 10^{-3}$  90 <sup>1</sup> ABLIKIM 18M BES3  $\psi(2S) \rightarrow \pi^0\pi^+\pi^-J/\psi$ 

OCCUR=2

<sup>1</sup> ABLIKIM 18M reports  $[\Gamma(h_c(1P) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.0 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R07;LINKAGE=B

### $\Gamma(\bar{p}p)/\Gamma_{\text{total}}$

 $\Gamma_4/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M144R20  
NODE=M144R20 $< 1.7 \times 10^{-4}$  (CL = 90%) [ $< 1.5 \times 10^{-4}$  (CL = 90%) OUR 2022 BEST LIMIT] $< 1.7 \times 10^{-4}$  90 <sup>1</sup> ABLIKIM 13V BES3  $\psi(2S) \rightarrow \gamma\bar{p}p$ 

<sup>1</sup> ABLIKIM 13V reports  $[\Gamma(h_c(1P) \rightarrow \bar{p}p)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 1.3 \times 10^{-7}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R20;LINKAGE=A

### $\Gamma(\bar{p}p\pi^0)/\Gamma_{\text{total}}$

 $\Gamma_5/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M144R22  
NODE=M144R22 $< 8 \times 10^{-4}$  90 <sup>1</sup> ABLIKIM 22M BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

<sup>1</sup> ABLIKIM 22M reports  $[\Gamma(h_c(1P) \rightarrow \bar{p}p\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 5.67 \times 10^{-7}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R22;LINKAGE=A

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.3±0.6 OUR AVERAGE** [(2.9 ± 0.6) × 10<sup>-3</sup> OUR 2022 AVERAGE]NODE=M144R08  
NODE=M144R08

NEW

**3.3±0.5±0.2** 230 <sup>1</sup> ABLIKIM 19AG BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

<sup>1</sup> ABLIKIM 19AG reports [ $\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] = (2.49 ± 0.27 ± 0.28) × 10<sup>-6</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = (7.4 ± 0.5) × 10<sup>-4</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R08;LINKAGE=A

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

**<6 × 10<sup>-4</sup> (CL = 90%)** [ $<5 \times 10^{-4}$  (CL = 90%) OUR 2022 BEST LIMIT]NODE=M144R10  
NODE=M144R10**<6 × 10<sup>-4</sup>** 90 12 <sup>1</sup> ABLIKIM 20AH BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

<sup>1</sup> ABLIKIM 20AH reports [ $\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] < 4.4 × 10<sup>-7</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = 7.4 × 10<sup>-4</sup>.

NODE=M144R10;LINKAGE=A

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**4.4±1.2±0.3** 86 <sup>1</sup> ABLIKIM 22M BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

<sup>1</sup> ABLIKIM 22M reports [ $\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] = (3.30 ± 0.71 ± 0.59) × 10<sup>-6</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = (7.4 ± 0.5) × 10<sup>-4</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R23  
NODE=M144R23

NODE=M144R23;LINKAGE=A

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**7.4±2.1±0.5** 20 <sup>1</sup> ABLIKIM 22M BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

<sup>1</sup> ABLIKIM 22M reports [ $\Gamma(h_c(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] = (5.51 ± 1.50 ± 0.46) × 10<sup>-7</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = (7.4 ± 0.5) × 10<sup>-4</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R21  
NODE=M144R21

NODE=M144R21;LINKAGE=A

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**1.9±0.5 OUR AVERAGE** [(1.6 ± 0.5) × 10<sup>-3</sup> OUR 2022 AVERAGE]NODE=M144R01  
NODE=M144R01

NEW

**1.9±0.5±0.1** 101 <sup>1</sup> ABLIKIM 19AG BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.6 90 <sup>2</sup> ADAMS 09 CLEO  $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$ 

<sup>1</sup> ABLIKIM 19AG reports [ $\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] = (1.38 ± 0.35 ± 0.17) × 10<sup>-6</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = (7.4 ± 0.5) × 10<sup>-4</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R01;LINKAGE=A

<sup>2</sup> ADAMS 09 reports [ $\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] < 0.19 × 10<sup>-5</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = 7.4 × 10<sup>-4</sup>.

NODE=M144R01;LINKAGE=AD

 $\Gamma(\pi^+\pi^-\pi^0\eta)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**8.3±2.4 OUR AVERAGE** [(7.2 ± 2.3) × 10<sup>-3</sup> OUR 2022 AVERAGE]NODE=M144R11  
NODE=M144R11

NEW

**8.3±2.3±0.6** 35 <sup>1</sup> ABLIKIM 20AH BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 

<sup>1</sup> ABLIKIM 20AH reports [ $\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0\eta)/\Gamma_{\text{total}}$ ] × [B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ )] = (6.2 ± 1.6 ± 0.7) × 10<sup>-6</sup> which we divide by our best value B( $\psi(2S) \rightarrow h_c(1P)\pi^0$ ) = (7.4 ± 0.5) × 10<sup>-4</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R11;LINKAGE=A

 $\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.94±0.17 OUR AVERAGE** [(0.81 ± 0.18) × 10<sup>-2</sup> OUR 2022 AVERAGE]NODE=M144R02  
NODE=M144R02

NEW

0.86±0.16±0.06 254 <sup>1</sup> ABLIKIM 19AG BES3  $\psi(2S) \rightarrow \pi^0 h_c(1P)$ 2.5 <sup>+0.9</sup>/<sub>-0.7</sub> ±0.2 92 <sup>2</sup> ADAMS 09 CLEO  $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow 2\pi^+ 2\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (6.40 \pm 0.81 \pm 0.87) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R02;LINKAGE=A

<sup>2</sup> ADAMS 09 reports  $[\Gamma(h_c(1P) \rightarrow 2\pi^+ 2\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (1.88_{-0.45-0.30}^{+0.48+0.47}) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R02;LINKAGE=AD

 **$\Gamma(3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{13}/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.010 (CL = 90%)</b>	<b>[&lt;9 × 10<sup>-3</sup> (CL = 90%) OUR 2022 BEST LIMIT]</b>			
<b>&lt;0.010</b>	90	<sup>1</sup> ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R03  
NODE=M144R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.034 90 <sup>2</sup> ADAMS 09 CLEO  $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 7.5 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R03;LINKAGE=A

<sup>2</sup> ADAMS 09 reports  $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.5 \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R03;LINKAGE=AD

 **$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$**  **$\Gamma_{14}/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;7 × 10<sup>-4</sup> (CL = 90%)</b>	<b>[&lt;6 × 10<sup>-4</sup> (CL = 90%) OUR 2022 BEST LIMIT]</b>			
<b>&lt;7 × 10<sup>-4</sup></b>	90	<sup>1</sup> ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R09  
NODE=M144R09

<sup>1</sup> ABLIKIM 19AG reports  $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 0.5 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R09;LINKAGE=A

 **$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{15}/\Gamma$** 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8±0.8 OUR AVERAGE</b>	<b>[(3.2 ± 0.8) × 10<sup>-3</sup> OUR 2022 AVERAGE]</b>			
<b>3.8±0.8±0.3</b>	80	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R12  
NODE=M144R12

NEW

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (2.8 \pm 0.5 \pm 0.3) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R12;LINKAGE=A

 **$\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$**  **$\Gamma_{16}/\Gamma$** 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.7 × 10<sup>-3</sup> (CL = 90%)</b>	<b>[&lt;2.3 × 10<sup>-3</sup> (CL = 90%) OUR 2022 BEST LIMIT]</b>				
<b>&lt;2.7 × 10<sup>-3</sup></b>	90	24	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R13  
NODE=M144R13

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.0 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R13;LINKAGE=A

 **$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{17}/\Gamma$** 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;6 × 10<sup>-4</sup></b>	90	20	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R14  
NODE=M144R14

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 4.8 \times 10^{-7}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R14;LINKAGE=A

 **$\Gamma(K^+ K^- \pi^0 \eta)/\Gamma_{\text{total}}$**  **$\Gamma_{18}/\Gamma$** 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.4 × 10<sup>-3</sup> (CL = 90%)</b>	<b>[&lt;2.1 × 10<sup>-3</sup> (CL = 90%) OUR 2022 BEST LIMIT]</b>				
<b>&lt;2.4 × 10<sup>-3</sup></b>	90	20	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R15  
NODE=M144R15

<sup>1</sup> ABLIKIM 20AH reports  $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^0 \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 1.8 \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .

NODE=M144R15;LINKAGE=A

$\Gamma(K^+K^-\eta)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-3}$ (CL = 90%)					[ $<9 \times 10^{-4}$ (CL = 90%) OUR 2022 BEST LIMIT]
$<1.0 \times 10^{-3}$	90	18	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports [ $\Gamma(h_c(1P) \rightarrow K^+K^-\eta)/\Gamma_{\text{total}}$ ] $\times$ [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] $< 7.5 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .					

NODE=M144R16  
NODE=M144R16

NODE=M144R16;LINKAGE=A

 $\Gamma(2K^+2K^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-4}$ (CL = 90%)					[ $<2.4 \times 10^{-4}$ (CL = 90%) OUR 2022 BEST LIMIT]
$<2.8 \times 10^{-4}$	90	11	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports [ $\Gamma(h_c(1P) \rightarrow 2K^+2K^-\pi^0)/\Gamma_{\text{total}}$ ] $\times$ [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] $< 2.1 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .					

NODE=M144R17  
NODE=M144R17

NODE=M144R17;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-4}$	90	17	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports [ $\Gamma(h_c(1P) \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$ ] $\times$ [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] $< 4.8 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$ .					

NODE=M144R18  
NODE=M144R18

NODE=M144R18;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.2<math>\pm</math>1.0 OUR AVERAGE</b>				[(2.8 $\pm$ 1.0) $\times 10^{-3}$ OUR 2022 AVERAGE]
<b>3.2<math>\pm</math>1.0<math>\pm</math>0.2</b>	41	<sup>1</sup> ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
<sup>1</sup> ABLIKIM 20AH reports [ $\Gamma(h_c(1P) \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ ] $\times$ [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] $= (2.4 \pm 0.7 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M144R19  
NODE=M144R19

NEW

NODE=M144R19;LINKAGE=A

## ————— RADIATIVE DECAYS —————

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.7<math>\pm</math>1.5<math>\pm</math>1.4</b>	18	ABLIKIM	16I BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta$

NODE=M144R06  
NODE=M144R06 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.52<math>\pm</math>0.27<math>\pm</math>0.29</b>	44	ABLIKIM	16I BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta'(958)$

NODE=M144R00  
NODE=M144R00 $\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>57<math>\pm</math> 5 OUR AVERAGE</b>				[(50 $\pm$ 9) $\times 10^{-2}$ OUR 2022 AVERAGE]
57 $\pm$ 4 $\pm$ 4	23k	<sup>1</sup> ABLIKIM	22AQ BES3	$\psi(2S) \rightarrow \pi^0$ hadrons; $\pi^0 \gamma(\eta_c)$
56 $\pm$ 6 $\pm$ 4		<sup>2</sup> DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
62 $\pm$ 9 $\pm$ 4	3679	<sup>3,4</sup> ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
56 $\pm$ 7 $\pm$ 4	1282	<sup>5</sup> DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
54 $\pm$ 14 $\pm$ 4	168	<sup>6</sup> ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

NODE=M144R2  
NODE=M144R2

NEW

OCCUR=2

<sup>1</sup> ABLIKIM 22AQ reports [ $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] =  
(4.22 $^{+0.27}_{-0.26}$   $\pm$  0.19)  $\times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$   
= (7.4  $\pm$  0.5)  $\times 10^{-4}$ . Our first error is their experiment's error and our second error is  
the systematic error from using our best value.

NODE=M144R2;LINKAGE=C

<sup>2</sup> Average of DOBBS 08A and ROSNER 05. DOBBS 08A reports [ $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$ ]  
 $\times$  [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] = (4.16  $\pm$  0.30  $\pm$  0.37)  $\times 10^{-4}$  which we divide by  
our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their  
experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=DB

<sup>3</sup> ABLIKIM 10B reports [ $\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$ ] =  
(4.58  $\pm$  0.40  $\pm$  0.50)  $\times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0)$   
= (7.4  $\pm$  0.5)  $\times 10^{-4}$ . Our first error is their experiment's error and our second error is  
the systematic error from using our best value.

NODE=M144R2;LINKAGE=A

<sup>4</sup>Superseded by ABLIKIM 22AQ

<sup>5</sup>DOBBS 08A reports  $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.19 \pm 0.32 \pm 0.45) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>6</sup>ROSNER 05 reports  $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.0 \pm 0.8 \pm 0.7) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=B  
NODE=M144R2;LINKAGE=DO

NODE=M144R2;LINKAGE=RO

## $h_c(1P)$ REFERENCES

ABLIKIM 22AQ	PR D106 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61889
ABLIKIM 22M	JHEP 2205 108	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61650
ABLIKIM 22N	JHEP 2205 003	M. Ablikim	(BESIII Collab.)	REFID=61651
ABLIKIM 20AH	PR D102 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60750
ABLIKIM 19AG	PR D99 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59858
ABLIKIM 18M	PR D97 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58901
ABLIKIM 16I	PRL 116 251802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57450
ABLIKIM 13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
ABLIKIM 12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54741
ABLIKIM 10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53348
ADAMS 09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
DOBBS 08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
ANDREOTTI 05B	PR D72 032001	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50768
ROSNER 05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
AULCHENKO 03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
ANTONIAZZI 94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074
ARMSTRONG 93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ARMSTRONG 92D	PRL 69 2337	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43174
BAGLIN 86	PL B171 135	C. Baglin <i>et al.</i>	(LAPP, CERN, TORI, STRB+)	REFID=43180

NODE=M144

NODE=M057

$\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on " $\psi(2S)$  and  $\chi_c$  branching ratios" before the  $\chi_{c0}(1P)$  Listings.

NODE=M057

## $\chi_{c2}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3556.17 ± 0.07</b>	<b>OUR AVERAGE</b>			
3557.3 ± 1.7 ± 0.7	611	<sup>1</sup> AAIJ	17BB LHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
3556.10 ± 0.06 ± 0.11	4.0k	<sup>2</sup> AAIJ	17BI LHCB	$\chi_{c2} \rightarrow J/\psi\mu^+\mu^-$
3555.3 ± 0.6 ± 2.2	2.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \text{hadrons}$
3555.70 ± 0.59 ± 0.39		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3556.173 ± 0.123 ± 0.020		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+e^-\gamma$
3559.9 ± 2.9		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c2}$
3556.4 ± 0.7		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3556.22 ± 0.131 ± 0.020	585	<sup>3</sup> ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+e^-\gamma$
3556.9 ± 0.4 ± 0.5	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
3557.8 ± 0.2 ± 4		<sup>4</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3553.4 ± 2.2	66	<sup>5</sup> LEMOIGNE	82 GOLI	$185\pi^-\text{Be} \rightarrow \gamma\mu^+\mu^-A$
3555.9 ± 0.7		<sup>6</sup> OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3557 ± 1.5	69	<sup>7</sup> HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3551 ± 11	15	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4		<sup>7</sup> BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3553 ± 4 ± 4		<sup>7,8</sup> TANENBAUM	78 MRK1	$e^+e^-$
3563 ± 7	360	<sup>7</sup> BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3555.4 ± 1.3	53	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
3543 ± 10	4	WHITAKER	76 MRK1	$e^+e^- \rightarrow J/\psi 2\gamma$

NODE=M057M

NODE=M057M



- <sup>1</sup> From a fit of the  $\phi\phi$  invariant mass with the width of  $\chi_{c2}(1P)$  fixed to the PDG 16 value.  
<sup>2</sup> AAIJ 17Bl reports also  $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$  MeV.  
<sup>3</sup> Recalculated by ANDREOTTI 05A, using the value of  $\psi(2S)$  mass from AULCHENKO 03.  
<sup>4</sup> Using mass of  $\psi(2S) = 3686.0$  MeV.  
<sup>5</sup>  $J/\psi(1S)$  mass constrained to 3097 MeV.  
<sup>6</sup> Assuming  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.  
<sup>7</sup> Mass value shifted by us by amount appropriate for  $\psi(2S)$  mass = 3686 MeV and  $J/\psi(1S)$  mass = 3097 MeV.  
<sup>8</sup> From a simultaneous fit to radiative and hadronic decay channels.

NODE=M057M;LINKAGE=A

NODE=M057M;LINKAGE=B

NODE=M057M;LINKAGE=NW

NODE=M057M;LINKAGE=C

NODE=M057M;LINKAGE=P

NODE=M057M;LINKAGE=E

NODE=M057M;LINKAGE=D

NODE=M057M;LINKAGE=M

 **$\chi_{c2}(1P)$  WIDTH**

NODE=M057W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.97 ± 0.09</b>	<b>OUR FIT</b>			
<b>2.00 ± 0.11</b>	<b>OUR AVERAGE</b>			
2.10 ± 0.20 ± 0.02	4.0k	AAIJ	17Bl LHCb	$\chi_{c2} \rightarrow J/\psi \mu^+ \mu^-$
1.915 ± 0.188 ± 0.013		ANDREOTTI	05A E835	$\rho \bar{p} \rightarrow e^+ e^- \gamma$
1.96 ± 0.17 ± 0.07	585	<sup>1</sup> ARMSTRONG	92 E760	$\bar{p} p \rightarrow e^+ e^- \gamma$
2.6 $\begin{smallmatrix} +1.4 \\ -1.0 \end{smallmatrix}$	50	BAGLIN	86B SPEC	$\bar{p} p \rightarrow e^+ e^- X$
2.8 $\begin{smallmatrix} +2.1 \\ -2.0 \end{smallmatrix}$		<sup>2</sup> GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

NODE=M057W

<sup>1</sup> Recalculated by ANDREOTTI 05A.<sup>2</sup> Errors correspond to 90% confidence level; authors give only width range.

NODE=M057W;LINKAGE=AN

NODE=M057W;LINKAGE=E

 **$\chi_{c2}(1P)$  DECAY MODES**

NODE=M057215;NODE=M057

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
<b>Hadronic decays</b>		
$\Gamma_1$ $2(\pi^+ \pi^-)$	( 1.02 ± 0.09 ) %	
$\Gamma_2$ $\rho \rho$		
$\Gamma_3$ $\pi^+ \pi^- \pi^0 \pi^0$	( 1.83 ± 0.23 ) %	
$\Gamma_4$ $\rho^+ \pi^- \pi^0 + \text{c.c.}$	( 2.19 ± 0.34 ) %	
$\Gamma_5$ $4\pi^0$	( 1.11 ± 0.15 ) × 10 <sup>-3</sup>	
$\Gamma_6$ $K^+ K^- \pi^0 \pi^0$	( 2.1 ± 0.4 ) × 10 <sup>-3</sup>	
$\Gamma_7$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	( 1.38 ± 0.20 ) %	
$\Gamma_8$ $\rho^- K^+ \bar{K}^0 + \text{c.c.}$	( 4.1 ± 1.2 ) × 10 <sup>-3</sup>	
$\Gamma_9$ $K^*(892)^0 K^- \pi^+ \rightarrow$ $K^- \pi^+ K^0 \pi^0 + \text{c.c.}$	( 2.9 ± 0.8 ) × 10 <sup>-3</sup>	
$\Gamma_{10}$ $K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	( 3.8 ± 0.9 ) × 10 <sup>-3</sup>	
$\Gamma_{11}$ $K^*(892)^- K^+ \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	( 3.7 ± 0.8 ) × 10 <sup>-3</sup>	
$\Gamma_{12}$ $K^*(892)^+ \bar{K}^0 \pi^- \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	( 2.9 ± 0.8 ) × 10 <sup>-3</sup>	
$\Gamma_{13}$ $K^+ K^- \eta \pi^0$	( 1.3 ± 0.4 ) × 10 <sup>-3</sup>	
$\Gamma_{14}$ $K^+ K^- \pi^+ \pi^-$	( 8.4 ± 0.9 ) × 10 <sup>-3</sup>	
$\Gamma_{15}$ $K^+ K^- \pi^+ \pi^- \pi^0$	( 1.17 ± 0.13 ) %	
$\Gamma_{16}$ $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	( 7.3 ± 0.8 ) × 10 <sup>-3</sup>	
$\Gamma_{17}$ $K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	( 2.1 ± 1.1 ) × 10 <sup>-3</sup>	
$\Gamma_{18}$ $K^*(892)^0 \bar{K}^*(892)^0$	( 2.3 ± 0.4 ) × 10 <sup>-3</sup>	
$\Gamma_{19}$ $3(\pi^+ \pi^-)$	( 8.6 ± 1.8 ) × 10 <sup>-3</sup>	
$\Gamma_{20}$ $\phi\phi$	( 1.06 ± 0.09 ) × 10 <sup>-3</sup>	
$\Gamma_{21}$ $\phi\phi\eta$	( 5.3 ± 0.6 ) × 10 <sup>-4</sup>	
$\Gamma_{22}$ $\omega\omega$	( 8.4 ± 1.0 ) × 10 <sup>-4</sup>	
$\Gamma_{23}$ $\omega K^+ K^-$	( 7.3 ± 0.9 ) × 10 <sup>-4</sup>	
$\Gamma_{24}$ $\omega\phi$	( 9.6 ± 2.7 ) × 10 <sup>-6</sup>	
$\Gamma_{25}$ $\pi\pi$	( 2.23 ± 0.09 ) × 10 <sup>-3</sup>	
$\Gamma_{26}$ $\rho^0 \pi^+ \pi^-$	( 3.7 ± 1.6 ) × 10 <sup>-3</sup>	
$\Gamma_{27}$ $\pi^+ \pi^- \pi^0$ (non-resonant)	( 2.0 ± 0.4 ) × 10 <sup>-5</sup>	
$\Gamma_{28}$ $\rho(770)^\pm \pi^\mp$	( 6 ± 4 ) × 10 <sup>-6</sup>	
$\Gamma_{29}$ $\pi^+ \pi^- \eta$	( 4.8 ± 1.3 ) × 10 <sup>-4</sup>	
$\Gamma_{30}$ $\pi^+ \pi^- \eta'$	( 5.0 ± 1.8 ) × 10 <sup>-4</sup>	

NODE=M057;CLUMP=A

DESIG=3

DESIG=43

DESIG=50

DESIG=51

DESIG=62

DESIG=52

DESIG=54

DESIG=55

DESIG=60

DESIG=56

DESIG=57

DESIG=58

DESIG=59

DESIG=5

DESIG=67

DESIG=78

DESIG=10

DESIG=21

DESIG=4

DESIG=16

DESIG=99

DESIG=25

DESIG=79

DESIG=68

DESIG=22

DESIG=9

DESIG=95

DESIG=96

DESIG=39

DESIG=42

Γ <sub>31</sub>	$\eta\eta$	$(5.4 \pm 0.4) \times 10^{-4}$		DESIG=14
Γ <sub>32</sub>	$K^+K^-$	$(1.01 \pm 0.06) \times 10^{-3}$		DESIG=2
Γ <sub>33</sub>	$K_S^0 K_S^0$	$(5.2 \pm 0.4) \times 10^{-4}$		DESIG=15
Γ <sub>34</sub>	$K^{*(892)\pm} K^\mp$	$(1.44 \pm 0.21) \times 10^{-4}$		DESIG=87
Γ <sub>35</sub>	$K^{*(892)^0} \bar{K}^0 + \text{c.c.}$	$(1.24 \pm 0.27) \times 10^{-4}$		DESIG=88
Γ <sub>36</sub>	$K_2^{*(1430)\pm} K^\mp$	$(1.48 \pm 0.12) \times 10^{-3}$		DESIG=89
Γ <sub>37</sub>	$K_2^{*(1430)^0} \bar{K}^0 + \text{c.c.}$	$(1.24 \pm 0.17) \times 10^{-3}$		DESIG=90
Γ <sub>38</sub>	$K_3^{*(1780)\pm} K^\mp$	$(5.2 \pm 0.8) \times 10^{-4}$		DESIG=91
Γ <sub>39</sub>	$K_3^{*(1780)^0} \bar{K}^0 + \text{c.c.}$	$(5.6 \pm 2.1) \times 10^{-4}$		DESIG=92
Γ <sub>40</sub>	$a_2(1320)^0 \pi^0$	$(1.29 \pm 0.34) \times 10^{-3}$		DESIG=93
Γ <sub>41</sub>	$a_2(1320)^\pm \pi^\mp$	$(1.8 \pm 0.6) \times 10^{-3}$		DESIG=94
Γ <sub>42</sub>	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$(1.28 \pm 0.18) \times 10^{-3}$		DESIG=17
Γ <sub>43</sub>	$K^+ K^- \pi^0$	$(3.0 \pm 0.8) \times 10^{-4}$		DESIG=36
Γ <sub>44</sub>	$K^+ K^- \eta$	$< 3.2 \times 10^{-4}$	90%	DESIG=40
Γ <sub>45</sub>	$K^+ K^- \eta'(958)$	$(1.94 \pm 0.34) \times 10^{-4}$		DESIG=82
Γ <sub>46</sub>	$\eta\eta'$	$(2.2 \pm 0.5) \times 10^{-5}$		DESIG=34
Γ <sub>47</sub>	$\eta'\eta'$	$(4.6 \pm 0.6) \times 10^{-5}$		DESIG=35
Γ <sub>48</sub>	$\pi^+ \pi^- K_S^0 K_S^0$	$(2.2 \pm 0.5) \times 10^{-3}$		DESIG=29
Γ <sub>49</sub>	$K^+ K^- K_S^0 K_S^0$	$< 4 \times 10^{-4}$	90%	DESIG=30
Γ <sub>50</sub>	$K_S^0 K_S^0 K_S^0 K_S^0$	$(1.13 \pm 0.18) \times 10^{-4}$		DESIG=97
Γ <sub>51</sub>	$K^+ K^- K^+ K^-$	$(1.65 \pm 0.20) \times 10^{-3}$		DESIG=24
Γ <sub>52</sub>	$K^+ K^- \phi$	$(1.42 \pm 0.29) \times 10^{-3}$		DESIG=32
Γ <sub>53</sub>	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(4.8 \pm 0.7) \times 10^{-3}$		DESIG=83
Γ <sub>54</sub>	$K^+ K^- \pi^0 \phi$	$(2.7 \pm 0.5) \times 10^{-3}$		DESIG=84
Γ <sub>55</sub>	$\phi \pi^+ \pi^- \pi^0$	$(9.3 \pm 1.2) \times 10^{-4}$		DESIG=80
Γ <sub>56</sub>	$\rho\bar{\rho}$	$(7.33 \pm 0.33) \times 10^{-5}$		DESIG=11
Γ <sub>57</sub>	$\rho\bar{\rho}\pi^0$	$(4.7 \pm 0.4) \times 10^{-4}$		DESIG=37
Γ <sub>58</sub>	$\rho\bar{\rho}\eta$	$(1.74 \pm 0.25) \times 10^{-4}$		DESIG=41
Γ <sub>59</sub>	$\rho\bar{\rho}\omega$	$(3.6 \pm 0.4) \times 10^{-4}$		DESIG=61
Γ <sub>60</sub>	$\rho\bar{\rho}\phi$	$(2.8 \pm 0.9) \times 10^{-5}$		DESIG=66
Γ <sub>61</sub>	$\rho\bar{\rho}\pi^+ \pi^-$	$(1.32 \pm 0.34) \times 10^{-3}$		DESIG=8
Γ <sub>62</sub>	$\rho\bar{\rho}\pi^0 \pi^0$	$(7.8 \pm 2.3) \times 10^{-4}$		DESIG=53
Γ <sub>63</sub>	$\rho\bar{\rho}K^+ K^- (\text{non-resonant})$	$(1.91 \pm 0.32) \times 10^{-4}$		DESIG=63
Γ <sub>64</sub>	$\rho\bar{\rho}K_S^0 K_S^0$	$< 7.9 \times 10^{-4}$	90%	DESIG=28
Γ <sub>65</sub>	$\rho\bar{n}\pi^-$	$(8.5 \pm 0.9) \times 10^{-4}$		DESIG=31
Γ <sub>66</sub>	$\bar{\rho}n\pi^+$	$(8.9 \pm 0.8) \times 10^{-4}$		DESIG=75
Γ <sub>67</sub>	$\rho\bar{n}\pi^- \pi^0$	$(2.17 \pm 0.18) \times 10^{-3}$		DESIG=76
Γ <sub>68</sub>	$\bar{\rho}n\pi^+ \pi^0$	$(2.11 \pm 0.18) \times 10^{-3}$		DESIG=77
Γ <sub>69</sub>	$\Lambda\bar{\Lambda}$	$(1.83 \pm 0.16) \times 10^{-4}$		DESIG=19
Γ <sub>70</sub>	$\Lambda\bar{\Lambda}\pi^+ \pi^-$	$(1.25 \pm 0.15) \times 10^{-3}$		DESIG=27
Γ <sub>71</sub>	$\Lambda\bar{\Lambda}\pi^+ \pi^- (\text{non-resonant})$	$(6.6 \pm 1.5) \times 10^{-4}$		DESIG=70
Γ <sub>72</sub>	$\Lambda\bar{\Lambda}\eta$	$(1.05 \pm 0.26) \times 10^{-4}$		DESIG=105
Γ <sub>73</sub>	$\Sigma(1385)^+ \bar{\Lambda}\pi^- + \text{c.c.}$	$< 4 \times 10^{-4}$	90%	DESIG=71
Γ <sub>74</sub>	$\Sigma(1385)^- \bar{\Lambda}\pi^+ + \text{c.c.}$	$< 6 \times 10^{-4}$	90%	DESIG=72
Γ <sub>75</sub>	$K^+ \bar{p}\Lambda + \text{c.c.}$	$(7.8 \pm 0.5) \times 10^{-4}$		DESIG=38
Γ <sub>76</sub>	$nK_S^0 \bar{\Lambda} + \text{c.c.}$	$(3.58 \pm 0.28) \times 10^{-4}$		DESIG=104
Γ <sub>77</sub>	$K^{*(892)^+} \bar{p}\Lambda + \text{c.c.}$	$(8.2 \pm 1.1) \times 10^{-4}$		DESIG=101
Γ <sub>78</sub>	$K^+ \bar{p}\Lambda(1520) + \text{c.c.}$	$(2.8 \pm 0.7) \times 10^{-4}$		DESIG=64
Γ <sub>79</sub>	$\Lambda(1520)\bar{\Lambda}(1520)$	$(4.6 \pm 1.5) \times 10^{-4}$		DESIG=65
Γ <sub>80</sub>	$\Sigma^0 \bar{\Sigma}^0$	$(3.7 \pm 0.6) \times 10^{-5}$		DESIG=47
Γ <sub>81</sub>	$\Sigma^+ \bar{p}K_S^0 + \text{c.c.}$	$(8.2 \pm 0.9) \times 10^{-5}$		DESIG=100
Γ <sub>82</sub>	$\Sigma^0 \bar{p}K^+ + \text{c.c.}$	$(9.1 \pm 0.8) \times 10^{-5}$		DESIG=103
Γ <sub>83</sub>	$\Sigma^+ \bar{\Sigma}^-$	$(3.4 \pm 0.7) \times 10^{-5}$		DESIG=48
Γ <sub>84</sub>	$\Sigma^- \bar{\Sigma}^+$	$(4.4 \pm 1.8) \times 10^{-5}$		DESIG=102
Γ <sub>85</sub>	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$< 1.6 \times 10^{-4}$	90%	DESIG=73
Γ <sub>86</sub>	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$< 8 \times 10^{-5}$	90%	DESIG=74
Γ <sub>87</sub>	$K^- \Lambda\bar{\Xi}^+ + \text{c.c.}$	$(1.76 \pm 0.32) \times 10^{-4}$		DESIG=85
Γ <sub>88</sub>	$\Xi^0 \bar{\Xi}^0$	$(1.83 \pm 0.22) \times 10^{-4}$		DESIG=49
Γ <sub>89</sub>	$\Xi^- \bar{\Xi}^+$	$(1.44 \pm 0.12) \times 10^{-4}$		DESIG=26
Γ <sub>90</sub>	$J/\psi(1S)\pi^+ \pi^- \pi^0$	$< 1.5 \%$	90%	DESIG=12
Γ <sub>91</sub>	$\pi^0 \eta_c$	$< 3.2 \times 10^{-3}$	90%	DESIG=81
Γ <sub>92</sub>	$\eta_c(1S)\pi^+ \pi^-$	$< 5.4 \times 10^{-3}$	90%	DESIG=69

**Radiative decays**

$\Gamma_{93}$	$\gamma J/\psi(1S)$	(19.0 $\pm$ 0.5) %		NODE=M057;CLUMP=B
$\Gamma_{94}$	$\gamma \rho^0$	< 1.9 $\times 10^{-5}$	90%	DESIG=6
$\Gamma_{95}$	$\gamma \omega$	< 6 $\times 10^{-6}$	90%	DESIG=44
$\Gamma_{96}$	$\gamma \phi$	< 7 $\times 10^{-6}$	90%	DESIG=45
$\Gamma_{97}$	$\gamma \gamma$	(2.85 $\pm$ 0.10) $\times 10^{-4}$		DESIG=46
$\Gamma_{98}$	$e^+ e^- J/\psi(1S)$	(2.15 $\pm$ 0.14) $\times 10^{-3}$		DESIG=7
$\Gamma_{99}$	$\mu^+ \mu^- J/\psi(1S)$	(2.02 $\pm$ 0.33) $\times 10^{-4}$		DESIG=86
				DESIG=98

**CONSTRAINED FIT INFORMATION**

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_{14}$	7									
$x_{17}$	2	21								
$x_{18}$	4	3	1							
$x_{20}$	7	5	1	3						
$x_{25}$	7	6	1	4	10					
$x_{26}$	18	2	0	1	1	1				
$x_{31}$	3	3	1	2	5	12	1			
$x_{32}$	5	4	1	3	7	15	1	8		
$x_{33}$	5	4	1	2	6	13	1	7	8	
$x_{42}$	2	2	0	1	3	7	0	3	4	4
$x_{51}$	4	3	1	2	4	7	1	4	5	4
$x_{56}$	10	9	2	5	9	11	2	5	8	7
$x_{69}$	3	3	1	2	5	12	1	6	8	6
$x_{93}$	12	10	2	6	15	34	2	18	22	18
$x_{97}$	-6	-4	-1	-2	2	20	-2	12	12	10
$\Gamma$	-23	-19	-4	-11	-19	-25	-5	-12	-18	-15
	$x_1$	$x_{14}$	$x_{17}$	$x_{18}$	$x_{20}$	$x_{25}$	$x_{26}$	$x_{31}$	$x_{32}$	$x_{33}$
$x_{51}$	2									
$x_{56}$	4	5								
$x_{69}$	3	4	5							
$x_{93}$	10	11	4	17						
$x_{97}$	5	4	18	11	34					
$\Gamma$	-8	-11	-45	-12	-46	-43				
	$x_{42}$	$x_{51}$	$x_{56}$	$x_{69}$	$x_{93}$	$x_{97}$				

**$\chi_{c2}(1P)$  PARTIAL WIDTHS**

$\chi_{c2}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$

NODE=M057220

NODE=M057223

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{56} \Gamma_{93} / \Gamma$

NODE=M057G1  
NODE=M057G1

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

**27.5 $\pm$ 1.2 OUR FIT**

**27.5 $\pm$ 1.5 OUR AVERAGE**

27.0 $\pm$ 1.5 $\pm$ 1.1	<sup>1</sup> ANDREOTTI 05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
27.7 $\pm$ 1.5 $\pm$ 2.0	<sup>1,2</sup> ARMSTRONG 92 E760	$p\bar{p} \rightarrow e^+ e^- \gamma$
36 $\pm$ 8	<sup>1</sup> BAGLIN 86B SPEC	$p\bar{p} \rightarrow e^+ e^- X$

<sup>1</sup> Calculated by us using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .

<sup>2</sup> Recalculated by ANDREOTTI 05A.

NODE=M057G;LINKAGE=7A  
NODE=M057G;LINKAGE=AN

$$\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{97}\Gamma_{93}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>107 ± 5 OUR FIT</b>				
<b>117 ± 10 OUR AVERAGE</b>				
111 ± 12 ± 9	147 ± 15	<sup>1</sup> DOBBS	06 CLE3	10.4 e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>
114 ± 11 ± 9	136 ± 13.3	<sup>1,2</sup> ABE	02T BELL	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>
139 ± 55 ± 21		<sup>1,3</sup> ACCIARRI	99E L3	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>
242 ± 65 ± 51		<sup>1,4</sup> ACKER...,K...	98 OPAL	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>
150 ± 42 ± 36		<sup>1,5</sup> DOMINICK	94 CLE2	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>
470 ± 240 ± 120		<sup>1,6</sup> BAUER	93 TPC	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>

NODE=M057G2  
NODE=M057G2

<sup>1</sup> Calculated by us using  $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1187 \pm 0.0008$ .

<sup>2</sup> All systematic errors added in quadrature.

<sup>3</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in ACCIARRI 99E is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.0162 \pm 0.0014$ .

<sup>4</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in ACKERSTAFF,K 98 is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$  and  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1203 \pm 0.0038$ .

<sup>5</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in DOMINICK 94 is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ ,  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0627 \pm 0.0020$ , and  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0597 \pm 0.0025$ .

<sup>6</sup> The value for  $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$  reported in BAUER 93 is derived using  $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ ,  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0627 \pm 0.0020$ , and  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0597 \pm 0.0025$ .

NODE=M057G;LINKAGE=LL  
NODE=M057G;LINKAGE=GT  
NODE=M057G;LINKAGE=J4

NODE=M057G;LINKAGE=J5

NODE=M057G;LINKAGE=J6

NODE=M057G;LINKAGE=J7

$$\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$$

NODE=M057224

$$\Gamma(2(\pi^+ \pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.7 ± 0.5 OUR FIT</b>				
<b>5.2 ± 0.7 OUR AVERAGE</b>				
5.01 ± 0.44 ± 0.55	1597 ± 138	UEHARA	08 BELL	γγ → χ <sub>c2</sub> → 2(π <sup>+</sup> π <sup>-</sup> )
6.4 ± 1.8 ± 0.8		EISENSTEIN	01 CLE2	e <sup>+</sup> e <sup>-</sup> → e <sup>+</sup> e <sup>-</sup> χ <sub>c2</sub>

NODE=M057G3  
NODE=M057G3

$$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<7.8	90	<598	UEHARA	08 BELL	γγ → χ <sub>c2</sub> → 2(π <sup>+</sup> π <sup>-</sup> )

NODE=M057G08  
NODE=M057G08

$$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.7 ± 0.5 OUR FIT</b>				
<b>4.42 ± 0.42 ± 0.53</b>	780 ± 74	UEHARA	08 BELL	γγ → χ <sub>c2</sub> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup>

NODE=M057G09  
NODE=M057G09

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.5 ± 0.9 ± 1.5</b>	1250	DEL-AMO-SA..11M	BABR	γγ → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup>

NODE=M057G02  
NODE=M057G02

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{18}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.26 ± 0.24 OUR FIT</b>				
<b>0.8 ± 0.17 ± 0.27</b>	151 ± 30	UEHARA	08 BELL	γγ → χ <sub>c2</sub> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup>

NODE=M057G10  
NODE=M057G10

$$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{20}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.60 ± 0.05 OUR FIT</b>				
<b>0.62 ± 0.07 ± 0.05</b>	89 ± 11	<sup>1</sup> LIU	12B BELL	γγ → 2(K <sup>+</sup> K <sup>-</sup> )

NODE=M057G12  
NODE=M057G12

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.58 ± 0.18 ± 0.16 26.5 ± 8.1 UEHARA 08 BELL γγ → χ<sub>c2</sub> → 2(K<sup>+</sup>K<sup>-</sup>)

<sup>1</sup> Supersedes UEHARA 08. Using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ .

NODE=M057G12;LINKAGE=LI

$$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{22}\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.64	90	<sup>1</sup> LIU	12B BELL	γγ → 2(π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> )

NODE=M057G03  
NODE=M057G03

<sup>1</sup> Using  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M057G03;LINKAGE=LI

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M057G04  
NODE=M057G04

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	90	<sup>1</sup> LIU	12B	BELL $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-------	----	------------------	-----	---

<sup>1</sup> Using  $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$  and  $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$ .

NODE=M057G04;LINKAGE=LI

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G4  
NODE=M057G4

**1.25±0.07 OUR FIT**

**1.18±0.25 OUR AVERAGE**

1.44±0.54±0.47	34 ± 13	<sup>1</sup> UEHARA	09	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
----------------	---------	---------------------	----	---

1.14±0.21±0.17	54 ± 10	<sup>2</sup> NAKAZAWA	05	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
----------------	---------	-----------------------	----	---

<sup>1</sup> We multiplied the measurement by 3 to convert from  $\pi^0 \pi^0$  to  $\pi\pi$ . Interference with the continuum included.

<sup>2</sup> We have multiplied  $\pi^+ \pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

NODE=M057G4;LINKAGE=UE

NODE=M057G;LINKAGE=NA

$$\Gamma(\rho^0 \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{26}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G07  
NODE=M057G07

**2.1±0.9 OUR FIT**

<b>3.2±1.9±0.5</b>	986 ± 578	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+ \pi^-)$
--------------------	-----------	--------	----	--

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G13  
NODE=M057G13

<b>0.53±0.22±0.09</b>	8	<sup>1</sup> UEHARA	10A	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$
-----------------------	---	---------------------	-----	--

<sup>1</sup> Interference with the continuum not included.

NODE=M057G13;LINKAGE=UE

$$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G5  
NODE=M057G5

**0.56±0.04 OUR FIT**

<b>0.44±0.11±0.07</b>	33 ± 8	NAKAZAWA	05	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
-----------------------	--------	----------	----	---

$$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G6  
NODE=M057G6

**0.294±0.025 OUR FIT**

<b>0.27 <math>\begin{smallmatrix} +0.07 \\ -0.06 \end{smallmatrix} \pm 0.03</math></b>	53	<sup>1</sup> UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$
--	----	---------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.05 ± 0.03	38 ± 7	CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
--------------------	--------	------	-----	--

<sup>1</sup> Supersedes CHEN 07B.

NODE=M057G6;LINKAGE=UE

$$\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G01  
NODE=M057G01

**0.72±0.11 OUR FIT**

<b>1.20±0.33±0.13</b>	126	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
-----------------------	-----	------------------------------	------	--

<sup>1</sup> We have multiplied  $\bar{K} K \pi$  by 2/3 to obtain  $\bar{K}^0 K^+ \pi^- + \text{c.c.}$

NODE=M057G01;LINKAGE=DE

$$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_{97}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G11  
NODE=M057G11

**0.93±0.11 OUR FIT**

<b>1.10±0.21±0.15</b>	126 ± 24	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$
-----------------------	----------	--------	----	--

$$\Gamma(\eta_c(1S)\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{92}\Gamma_{97}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M057G05  
NODE=M057G05

< <b>15.7</b>	90	LEES	12AE	BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$
---------------	----	------	------	---

## $\chi_{c2}(1P)$ BRANCHING RATIOS

NODE=M057225

### HADRONIC DECAYS

NODE=M057305

$$\Gamma(2(\pi^+ \pi^-))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	DOCUMENT ID
-------	-------------

NODE=M057R2  
NODE=M057R2

**0.0102±0.0009 OUR FIT**

$$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-)) \quad \Gamma_{26}/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M057R38  
NODE=M057R38

**0.36±0.15 OUR FIT**

<b>0.31±0.17</b>	TANENBAUM	78	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c2}$
------------------	-----------	----	--

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.83±0.23±0.04</b>	903.5	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R46  
NODE=M057R46

<sup>1</sup> HE 08B reports  $1.87 \pm 0.07 \pm 0.22 \pm 0.13$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R46;LINKAGE=HE

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.19±0.34±0.05</b>	1031.9	<sup>1,2</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R47  
NODE=M057R47

<sup>1</sup> HE 08B reports  $2.23 \pm 0.11 \pm 0.32 \pm 0.16$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R47;LINKAGE=HE

<sup>2</sup> Calculated by us. We have added the values from HE 08B for  $\rho^+\pi^-\pi^0$  and  $\rho^-\pi^+\pi^0$  decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M057R47;LINKAGE=OC

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.11±0.15±0.02</b>	1164	<sup>1</sup> ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R58  
NODE=M057R58

<sup>1</sup> ABLIKIM 11A reports  $(1.21 \pm 0.05 \pm 0.16) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R58;LINKAGE=AB

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.206±0.040±0.004</b>	76.9	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R48  
NODE=M057R48

<sup>1</sup> HE 08B reports  $0.21 \pm 0.03 \pm 0.03 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R48;LINKAGE=HE

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.38±0.19±0.03</b>	211.6	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R50  
NODE=M057R50

<sup>1</sup> HE 08B reports  $1.41 \pm 0.11 \pm 0.16 \pm 0.10$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R50;LINKAGE=HE

 $\Gamma(\rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_8/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.41±0.12±0.01</b>	62.9	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R51  
NODE=M057R51

<sup>1</sup> HE 08B reports  $0.42 \pm 0.11 \pm 0.06 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R51;LINKAGE=HE

 $\Gamma(K^*(892)^0 K^- \pi^+ \rightarrow K^- \pi^+ K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_9/\Gamma$ 

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.29±0.08±0.01</b>	38.7	<sup>1</sup> HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R57  
NODE=M057R57

<sup>1</sup> HE 08B reports  $0.30 \pm 0.07 \pm 0.04 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^- \pi^+ \rightarrow K^- \pi^+ K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R57;LINKAGE=HE

$$\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{10}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.38 ± 0.09 ± 0.01</b>	63.0	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R52  
NODE=M057R52

<sup>1</sup> HE 08B reports  $0.39 \pm 0.07 \pm 0.05 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R52;LINKAGE=HE

$$\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.37 ± 0.08 ± 0.01</b>	51.1	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R53  
NODE=M057R53

<sup>1</sup> HE 08B reports  $0.38 \pm 0.07 \pm 0.04 \pm 0.03$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R53;LINKAGE=HE

$$\Gamma(K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{12}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.29 ± 0.08 ± 0.01</b>	39.3	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R54  
NODE=M057R54

<sup>1</sup> HE 08B reports  $0.30 \pm 0.07 \pm 0.04 \pm 0.02$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R54;LINKAGE=HE

$$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{13}/\Gamma$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.127 ± 0.044 ± 0.003</b>	22.9	<sup>1</sup> HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R55  
NODE=M057R55

<sup>1</sup> HE 08B reports  $0.13 \pm 0.04 \pm 0.02 \pm 0.01$  % from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R55;LINKAGE=HE

$$\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID
<b>8.4 ± 0.9 OUR FIT</b>	

NODE=M057R3  
NODE=M057R3

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.69 ± 0.13 ± 1.31</b>	11k	<sup>1</sup> ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R00  
NODE=M057R00

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$ .

NODE=M057R00;LINKAGE=A

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.30 ± 0.11 ± 0.75</b>	4.5k	<sup>1</sup> ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R73  
NODE=M057R73

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$ .

NODE=M057R73;LINKAGE=A

$$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(K^+ K^- \pi^+ \pi^-) \quad \Gamma_{17}/\Gamma_{14}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.25 ± 0.13 OUR FIT</b>			
<b>0.25 ± 0.13</b>	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R39  
NODE=M057R39

$$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>21 ± 11 OUR FIT</b>	

NODE=M057R9  
NODE=M057R9

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units $10^{-3}$ )	DOCUMENT ID
<b>2.3 ± 0.4 OUR FIT</b>	

NODE=M057R26  
NODE=M057R26

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**8.6±1.8 OUR EVALUATION** Treating systematic error as correlated.NODE=M057R4  
NODE=M057R4

→ UNCHECKED ←

**15.2±1.7 OUR AVERAGE** Error includes scale factor of 3.7.  $[(8.6 \pm 1.8) \times 10^{-3}$  OUR 2022 AVERAGE]

NEW

15.7±0.1±0.5	112K	<sup>1</sup> ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+\pi^-)$
8.6±0.9±1.6		<sup>2</sup> BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
8.7±5.9±0.4		<sup>2</sup> TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

<sup>1</sup> ABLIKIM 22Q reports  $(1.565 \pm 0.005 \pm 0.048) \times 10^{-2}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ .

NODE=M057R4;LINKAGE=A

<sup>2</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ . Multiplied by a factor of 2 to convert from  $K_S^0 K^+\pi^-$  to  $K^0 K^+\pi^-$  decay.

NODE=M057R;LINKAGE=X3

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID
--------------------------	-------------

**1.06±0.09 OUR FIT**NODE=M057R20  
NODE=M057R20 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.3±0.5±0.4** 143.6 <sup>1</sup> ABLIKIM 20B BES3  $\psi(2S) \rightarrow \gamma \phi\phi\eta$ NODE=M057R97  
NODE=M057R97<sup>1</sup> ABLIKIM 20B reports  $(5.33 \pm 0.52 \pm 0.39) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ .

NODE=M057R97;LINKAGE=A

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.84±0.10 OUR AVERAGE**

0.82±0.10±0.02	762	<sup>1</sup> ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
1.73±0.57±0.04	27.7±7.4	<sup>2</sup> ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma 6\pi$

NODE=M057R28  
NODE=M057R28<sup>1</sup> ABLIKIM 11K reports  $(8.9 \pm 0.3 \pm 1.1) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R28;LINKAGE=AL

<sup>2</sup> ABLIKIM 05N reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R28;LINKAGE=AB

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.73±0.04±0.08** 512 <sup>1</sup> ABLIKIM 13B BES3  $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$ NODE=M057R74  
NODE=M057R74<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$ .

NODE=M057R74;LINKAGE=A

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**9.6±2.7±0.2** 33 <sup>1</sup> ABLIKIM 19J BES3  $\psi(2S) \rightarrow \gamma$  hadronsNODE=M057R63  
NODE=M057R63

●●● We do not use the following data for averages, fits, limits, etc. ●●●

<18 90 <sup>2,3</sup> ABLIKIM 11K BES3  $\psi(2S) \rightarrow \gamma$  hadrons<sup>1</sup> ABLIKIM 19J reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.91 \pm 0.23 \pm 0.12) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R63;LINKAGE=A

<sup>2</sup> ABLIKIM 11K reports  $< 2 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R63;LINKAGE=AL

<sup>3</sup> Superseded by ABLIKIM 19J.

NODE=M057R63;LINKAGE=B

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID
--------------------------	-------------

**2.23±0.09 OUR FIT**NODE=M057R27  
NODE=M057R27



$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**37±16 OUR FIT**

DOCUMENT ID

 $\Gamma_{26}/\Gamma$ NODE=M057R8  
NODE=M057R8 $\Gamma(\pi^+ \pi^- \pi^0 (\text{non-resonant}))/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**2.01±0.42±0.04**

64

<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$ 

<sup>1</sup> ABLIKIM 17AG reports  $(2.1 \pm 0.4 \pm 0.2) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- \pi^0 (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{27}/\Gamma$ NODE=M057R84  
NODE=M057R84

OCCUR=2

NODE=M057R84;LINKAGE=B

 $\Gamma(\rho(770)^\pm \pi^\mp)/\Gamma_{\text{total}}$ VALUE (units  $10^{-5}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**0.61±0.38±0.01**

15

<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$ 

<sup>1</sup> ABLIKIM 17AG reports  $(0.64 \pm 0.39 \pm 0.07) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \rho(770)^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{28}/\Gamma$ NODE=M057R85  
NODE=M057R85

NODE=M057R85;LINKAGE=A

 $\Gamma(\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )

CL%

DOCUMENT ID

TECN

COMMENT

**0.48±0.13±0.01**<sup>1</sup> ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt;1.4

90

<sup>2</sup> ABLIKIM 06R BES2  $\psi(2S) \rightarrow \gamma \chi_{c2}$ 

<sup>1</sup> ATHAR 07 reports  $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R08  
NODE=M057R08

NODE=M057R08;LINKAGE=AT

<sup>2</sup> ABLIKIM 06R reports  $< 1.7 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R08;LINKAGE=AB

 $\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**0.50±0.18±0.01**<sup>1</sup> ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ 

<sup>1</sup> ATHAR 07 reports  $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{30}/\Gamma$ NODE=M057R35  
NODE=M057R35

NODE=M057R35;LINKAGE=AT

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ VALUE (units  $10^{-4}$ )**5.4±0.4 OUR FIT**

DOCUMENT ID

 $\Gamma_{31}/\Gamma$ NODE=M057R16  
NODE=M057R16 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**1.01±0.06 OUR FIT**

DOCUMENT ID

 $\Gamma_{32}/\Gamma$ NODE=M057R11  
NODE=M057R11 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ VALUE (units  $10^{-3}$ )**0.52±0.04 OUR FIT**

DOCUMENT ID

 $\Gamma_{33}/\Gamma$ NODE=M057R19  
NODE=M057R19 $\Gamma(K_S^0 K_S^0)/\Gamma(\pi\pi)$ 

VALUE

**0.235±0.019 OUR FIT**

DOCUMENT ID

TECN

COMMENT

 $\Gamma_{33}/\Gamma_{25}$ NODE=M057R36  
NODE=M057R36

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.27 ± 0.07 ± 0.04

<sup>1,2</sup> CHEN07B BELL  $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$ 

<sup>1</sup> Using  $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from the  $\pi^+ \pi^-$  measurement of NAKAZAWA 05 rescaled by 3/2 to convert to  $\pi\pi$ .

NODE=M057R36;LINKAGE=CH

<sup>2</sup> Not independent from other measurements.

NODE=M057R36;LINKAGE=NI

$\Gamma(K_S^0 \bar{K}_S^0)/\Gamma(K^+ K^-)$  $\Gamma_{33}/\Gamma_{32}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M057R37  
 NODE=M057R37

**0.52±0.05 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.21±0.12	1,2 CHEN	07B BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
----------------	----------	----------	---

<sup>1</sup> Using  $\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  from NAKAZAWA 05.

<sup>2</sup> Not independent from other measurements.

NODE=M057R37;LINKAGE=CH  
 NODE=M057R37;LINKAGE=NI

 $\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$  $\Gamma_{34}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057R86  
 NODE=M057R86

**1.44±0.21±0.03**

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.72±0.26±0.04	<sup>2</sup> ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
1.34±0.27±0.03	<sup>3</sup> ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

OCCUR=2  
 OCCUR=3

<sup>1</sup> ABLIKIM 17AG reports  $(1.5 \pm 0.1 \pm 0.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R86;LINKAGE=A

<sup>2</sup> ABLIKIM 17AG reports  $(1.8 \pm 0.2 \pm 0.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R86;LINKAGE=B

<sup>3</sup> ABLIKIM 17AG reports  $(1.4 \pm 0.2 \pm 0.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R86;LINKAGE=C

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057R87  
 NODE=M057R87

**1.24±0.27±0.03**

	<sup>1</sup> ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
--	----------------------	-----------	---

<sup>1</sup> ABLIKIM 17AG reports  $(1.3 \pm 0.2 \pm 0.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R87;LINKAGE=A

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$  $\Gamma_{36}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057R88  
 NODE=M057R88

**14.8±1.2±0.3**

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.4±1.6±0.4	<sup>2</sup> ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
13.0±1.5±0.3	<sup>3</sup> ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

OCCUR=2  
 OCCUR=3

<sup>1</sup> ABLIKIM 17AG reports  $(15.5 \pm 0.6 \pm 1.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R88;LINKAGE=A

<sup>2</sup> ABLIKIM 17AG reports  $(18.2 \pm 0.8 \pm 1.6) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R88;LINKAGE=B

<sup>3</sup> ABLIKIM 17AG reports  $(13.6 \pm 0.8 \pm 1.4) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R88;LINKAGE=C

$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{37}/\Gamma$ VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

**12.4 ± 1.7 ± 0.3**<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ 

<sup>1</sup> ABLIKIM 17AG reports  $(13.0 \pm 1.0 \pm 1.5) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R89  
NODE=M057R89

NODE=M057R89;LINKAGE=A

 $\Gamma(K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

**5.2 ± 0.8 ± 0.1**<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K \bar{K} \pi$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.1 ± 1.0 ± 0.1

<sup>2</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ 

5.6 ± 1.8 ± 0.1

<sup>3</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ 

<sup>1</sup> ABLIKIM 17AG reports  $(5.4 \pm 0.5 \pm 0.7) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> ABLIKIM 17AG reports  $(5.3 \pm 0.5 \pm 0.9) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> ABLIKIM 17AG reports  $(5.9 \pm 1.1 \pm 1.5) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R90  
NODE=M057R90

OCCUR=2

OCCUR=3

NODE=M057R90;LINKAGE=A

NODE=M057R90;LINKAGE=B

NODE=M057R90;LINKAGE=C

 $\Gamma(K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{39}/\Gamma$ VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

**5.6 ± 2.1 ± 0.1**<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ 

<sup>1</sup> ABLIKIM 17AG reports  $(5.9 \pm 1.6 \pm 1.5) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R91  
NODE=M057R91

NODE=M057R91;LINKAGE=A

 $\Gamma(a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{40}/\Gamma$ VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

**12.9 ± 3.4 ± 0.3**<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$ 

<sup>1</sup> ABLIKIM 17AG reports  $(13.5 \pm 1.6 \pm 3.2) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R92  
NODE=M057R92

NODE=M057R92;LINKAGE=A

 $\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$  $\Gamma_{41}/\Gamma$ VALUE (units  $10^{-4}$ )

DOCUMENT ID TECN COMMENT

**17.6 ± 6.1 ± 0.4**<sup>1</sup> ABLIKIM 17AG BES3  $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$ 

<sup>1</sup> ABLIKIM 17AG reports  $(18.4 \pm 3.3 \pm 5.5) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R93  
NODE=M057R93

NODE=M057R93;LINKAGE=A

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{43}/\Gamma$ VALUE (units  $10^{-3}$ )

DOCUMENT ID TECN COMMENT

**0.30±0.08±0.01**<sup>1</sup> ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ 

<sup>1</sup> ATHAR 07 reports  $(0.31 \pm 0.07 \pm 0.04) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R05  
NODE=M057R05

NODE=M057R05;LINKAGE=AT

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$  $\Gamma_{44}/\Gamma$ VALUE (units  $10^{-3}$ )

CL%

DOCUMENT ID TECN COMMENT

**<0.32**

90

<sup>1</sup> ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ 

<sup>1</sup> ATHAR 07 reports  $< 0.33 \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R09  
NODE=M057R09

NODE=M057R09;LINKAGE=AT

 $\Gamma(K^+ K^- \eta'(958))/\Gamma_{\text{total}}$  $\Gamma_{45}/\Gamma$ VALUE (units  $10^{-4}$ )

EVTS

DOCUMENT ID TECN COMMENT

**1.94±0.34**

107

<sup>1</sup> ABLIKIM 14J BES3  $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$ 

<sup>1</sup> Derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.72 \pm 0.34)\%$ . Uncertainty includes both statistical and systematic contributions combined in quadrature.

NODE=M057R78  
NODE=M057R78

NODE=M057R78;LINKAGE=A

 $\Gamma(\eta\eta')/\Gamma_{\text{total}}$  $\Gamma_{46}/\Gamma$ VALUE (units  $10^{-5}$ )

CL%

EVTS

DOCUMENT ID TECN COMMENT

**2.17±0.47±0.05**

20

<sup>1</sup> ABLIKIM 17AI BES3  $\psi(2S) \rightarrow \gamma \eta' \eta$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt; 6

90

3.3 ± 8.0

<sup>2</sup> ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma \eta \eta'$ 

&lt;23

90

<sup>3</sup> ADAMS 07 CLEO  $\psi(2S) \rightarrow \gamma \chi_{c2}$ 

<sup>1</sup> ABLIKIM 17AI reports  $(2.27 \pm 0.43 \pm 0.25) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R03  
NODE=M057R03

NODE=M057R03;LINKAGE=A

<sup>2</sup> ASNER 09 reports  $< 0.6 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R03;LINKAGE=AS

<sup>3</sup> Superseded by ASNER 09. ADAMS 07 reports  $< 2.3 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R03;LINKAGE=AD

 $\Gamma(\eta' \eta')/\Gamma_{\text{total}}$  $\Gamma_{47}/\Gamma$ VALUE (units  $10^{-5}$ )

CL%

EVTS

DOCUMENT ID TECN COMMENT

**4.6±0.6±0.1**

60

<sup>1</sup> ABLIKIM 17AI BES3  $\psi(2S) \rightarrow \gamma \eta' \eta'$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

&lt;10

90

12 ± 7

<sup>2</sup> ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma \eta' \eta'$ 

&lt;30

90

<sup>3</sup> ADAMS 07 CLEO  $\psi(2S) \rightarrow \gamma \chi_{c2}$ 

<sup>1</sup> ABLIKIM 17AI reports  $(4.76 \pm 0.56 \pm 0.38) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R04  
NODE=M057R04

NODE=M057R04;LINKAGE=A

<sup>2</sup> ASNER 09 reports  $< 1.0 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R04;LINKAGE=AS

<sup>3</sup> Superseded by ASNER 09. ADAMS 07 reports  $< 3.1 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R04;LINKAGE=AD

$\Gamma(\pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.17±0.54±0.05</b>	57 ± 11	<sup>1</sup> ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R31  
NODE=M057R31

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R31;LINKAGE=AB

 $\Gamma(K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;4</b>	90	2.3 ± 2.2	<sup>1</sup> ABLIKIM	050 BES2	$e^+e^- \rightarrow \chi_{c2}\gamma$

NODE=M057R32  
NODE=M057R32

<sup>1</sup> ABLIKIM 050 reports  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] < 3.5 \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R32;LINKAGE=AB

 $\Gamma(K_S^0K_S^0K_S^0K_S^0)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.13±0.18±0.02</b>	68	<sup>1</sup> ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$

NODE=M057R94  
NODE=M057R94

<sup>1</sup> Using  $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$ . ABLIKIM 19AA reports  $[\Gamma(\chi_{c2}(1P) \rightarrow K_S^0K_S^0K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (10.8 \pm 1.5 \pm 0.8) \times 10^{-6}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value..

NODE=M057R94;LINKAGE=A

 $\Gamma(K^+K^-K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID
<b>1.65±0.20 OUR FIT</b>	

NODE=M057R18  
NODE=M057R18

 $\Gamma(K^+K^-\phi)/\Gamma_{\text{total}}$   $\Gamma_{52}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.42±0.29±0.03</b>	52	<sup>1</sup> ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M057R01  
NODE=M057R01

<sup>1</sup> ABLIKIM 06T reports  $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R01;LINKAGE=AB

 $\Gamma(\bar{K}^0K^+\pi^-\phi + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.83±0.32±0.66</b>	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R79  
NODE=M057R79

 $\Gamma(K^+K^-\pi^0\phi)/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.74±0.16±0.44</b>	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R80  
NODE=M057R80

 $\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.93±0.06±0.10</b>	408	<sup>1</sup> ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R75  
NODE=M057R75

<sup>1</sup> Using  $1.06 \times 10^8 \psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$ .

NODE=M057R75;LINKAGE=A

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>0.733±0.033 OUR FIT</b>	

NODE=M057R12  
NODE=M057R12

 $\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.47±0.04 OUR AVERAGE</b>			

NODE=M057R06  
NODE=M057R06

0.47±0.04±0.01	<sup>1</sup> ONYISI	10	CLE3	$\psi(2S) \rightarrow \gamma\rho\bar{\rho}X$
0.43±0.09±0.01	<sup>2</sup> ATHAR	07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ONYISI 10 reports  $(4.83 \pm 0.25 \pm 0.35 \pm 0.31) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=ON

<sup>2</sup> ATHAR 07 reports  $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=AT

 **$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$**  **$\Gamma_{58}/\Gamma$** VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**0.174 ± 0.025 OUR AVERAGE**

0.172 ± 0.026 ± 0.004

<sup>1</sup> ONYISI 10 CLE3  $\psi(2S) \rightarrow \gamma p\bar{p}X$ 

0.186 ± 0.070 ± 0.004

<sup>2</sup> ATHAR 07 CLEO  $\psi(2S) \rightarrow \gamma h^+ h^- h^0$ NODE=M057R34  
NODE=M057R34

<sup>1</sup> ONYISI 10 reports  $(1.76 \pm 0.23 \pm 0.14 \pm 0.11) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R34;LINKAGE=ON

<sup>2</sup> ATHAR 07 reports  $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R34;LINKAGE=AT

 **$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$**  **$\Gamma_{59}/\Gamma$** VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**0.36 ± 0.04 ± 0.01**<sup>1</sup> ONYISI 10 CLE3  $\psi(2S) \rightarrow \gamma p\bar{p}X$ 

<sup>1</sup> ONYISI 10 reports  $(3.68 \pm 0.35 \pm 0.26 \pm 0.24) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R56  
NODE=M057R56

NODE=M057R56;LINKAGE=ON

 **$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$**  **$\Gamma_{60}/\Gamma$** VALUE (units  $10^{-5}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**2.8 ± 0.9 ± 0.1**

24 ± 7

<sup>1</sup> ABLIKIM 11F BES3  $\psi(2S) \rightarrow \gamma p\bar{p}K^+ K^-$ 

<sup>1</sup> ABLIKIM 11F reports  $(3.04 \pm 0.85 \pm 0.43) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R62  
NODE=M057R62

NODE=M057R62;LINKAGE=AB

 **$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$**  **$\Gamma_{61}/\Gamma$** VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**1.32 ± 0.34 OUR EVALUATION**

Treating systematic error as correlated.

**1.3 ± 0.4 OUR AVERAGE**

Error includes scale factor of 1.3.

1.17 ± 0.19 ± 0.30

<sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma\chi_{c2}$ 

2.64 ± 1.03 ± 0.14

<sup>1</sup> TANENBAUM 78 MRK1  $\psi(2S) \rightarrow \gamma\chi_{c2}$ 

<sup>1</sup> Rescaled by us using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$ . Multiplied by a factor of 2 to convert from  $K_S^0 K^+\pi^-$  to  $K^0 K^+\pi^-$  decay.

NODE=M057R6  
NODE=M057R6

→ UNCHECKED ←

NODE=M057R6;LINKAGE=X3

 **$\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$**  **$\Gamma_{62}/\Gamma$** 

VALUE (%)

EVTS

DOCUMENT ID

TECN

COMMENT

**0.078 ± 0.023 ± 0.002**

29.2

<sup>1</sup> HE 08B CLEO  $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ 

<sup>1</sup> HE 08B reports  $0.08 \pm 0.02 \pm 0.01 \pm 0.01\%$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R49  
NODE=M057R49

NODE=M057R49;LINKAGE=HE

$\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$  $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.91±0.32±0.04</b>	131 ± 12	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M057R59  
 NODE=M057R59

<sup>1</sup> ABLIKIM 11F reports  $(2.08 \pm 0.19 \pm 0.30) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] \text{ assuming } B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R59;LINKAGE=AB

 $\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{64}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;7.9</b>	90	<sup>1</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$

NODE=M057R30  
 NODE=M057R30

<sup>1</sup> Using  $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$ .

NODE=M057R;LINKAGE=AB

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{65}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.5±0.9 OUR AVERAGE</b>				
8.4±1.0±0.2	3309	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
10.2±3.4±0.2		<sup>2</sup> ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

NODE=M057R33  
 NODE=M057R33

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.80 \pm 0.02 \pm 0.09) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R33;LINKAGE=AL

<sup>2</sup> ABLIKIM 06I reports  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R33;LINKAGE=AB

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$  $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.9±0.8±0.2</b>	3732	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

NODE=M057R70  
 NODE=M057R70

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.85 \pm 0.02 \pm 0.07) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R70;LINKAGE=AL

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.7±1.7±0.5</b>	2128	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

NODE=M057R71  
 NODE=M057R71

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.07 \pm 0.06 \pm 0.15) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R71;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>21.1±1.8±0.4</b>	2352	<sup>1</sup> ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

NODE=M057R72  
 NODE=M057R72

<sup>1</sup> ABLIKIM 12J reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.01 \pm 0.06 \pm 0.16) \times 10^{-4}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R72;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$  $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>1.83±0.16 OUR FIT</b>	

NODE=M057R25  
 NODE=M057R25

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>125±15±3</b>		371	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
<350	90		<sup>2</sup> ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$

NODE=M057R29  
 NODE=M057R29

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> ABLIKIM 12I reports  $(137.0 \pm 7.6 \pm 15.7) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda} \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R29;LINKAGE=AL

<sup>2</sup> Using  $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$ .

NODE=M057R29;LINKAGE=AB

### $\Gamma(\Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant})) / \Gamma_{\text{total}}$

 $\Gamma_{71} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>66 ± 15 ± 1</b>	36	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M057R65  
NODE=M057R65

<sup>1</sup> ABLIKIM 12I reports  $(71.8 \pm 14.5 \pm 8.2) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant})) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R65;LINKAGE=AL

### $\Gamma(\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}) / \Gamma_{\text{total}}$

 $\Gamma_{73} / \Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 40</b>	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Lambda} \pi^-$

NODE=M057R66  
NODE=M057R66

<sup>1</sup> ABLIKIM 12I reports  $< 42 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R66;LINKAGE=AL

### $\Gamma(\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}) / \Gamma_{\text{total}}$

 $\Gamma_{74} / \Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 60</b>	90	<sup>1</sup> ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Lambda} \pi^+$

NODE=M057R67  
NODE=M057R67

<sup>1</sup> ABLIKIM 12I reports  $< 61 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R67;LINKAGE=AL

### $\Gamma(K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}$

 $\Gamma_{75} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.8 ± 0.5 OUR AVERAGE</b>				

NODE=M057R07  
NODE=M057R07

7.7 ± 0.5 ± 0.2	5k	<sup>1,2</sup> ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$
8.3 ± 1.6 ± 0.2		<sup>3</sup> ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

<sup>1</sup> ABLIKIM 13D reports  $(8.4 \pm 0.3 \pm 0.6) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R07;LINKAGE=AB

<sup>2</sup> Using  $B(\Lambda \rightarrow p \pi^-) = 63.9\%$ .

NODE=M057R07;LINKAGE=LB

<sup>3</sup> ATHAR 07 reports  $(8.5 \pm 1.4 \pm 1.0) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R07;LINKAGE=AT

### $\Gamma(n K_S^0 \bar{\Lambda} + \text{c.c.}) / \Gamma_{\text{total}}$

 $\Gamma_{76} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.58 ± 0.16 ± 0.23</b>	879	<sup>1</sup> ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$

NODE=M057P01  
NODE=M057P01

<sup>1</sup> ABLIKIM 21AV reports  $(3.58 \pm 0.16 \pm 0.23) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow n K_S^0 \bar{\Lambda} + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0952 \pm 0.0020$ . Also uses  $B(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = (63.9 \pm 0.5)\%$  and  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ .

NODE=M057P01;LINKAGE=B

### $\Gamma(K^*(892)^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}$

 $\Gamma_{77} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.2 ± 1.1 ± 0.2</b>	476	<sup>1</sup> ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+} \bar{p} \Lambda$

NODE=M057R98  
NODE=M057R98

<sup>1</sup> ABLIKIM 19AU reports  $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (7.8 \pm 0.9 \pm 0.6) \times 10^{-5}$  which we divide by our best value  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R98;LINKAGE=F



$\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{78}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.8±0.7±0.1</b>	79 ± 13	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M057R60  
NODE=M057R60

<sup>1</sup> ABLIKIM 11F reports  $(3.06 \pm 0.50 \pm 0.54) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R60;LINKAGE=AB

 $\Gamma(\Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}$  $\Gamma_{79}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.6±1.4±0.1</b>	29 ± 7	<sup>1</sup> ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M057R61  
NODE=M057R61

<sup>1</sup> ABLIKIM 11F reports  $(5.05 \pm 1.29 \pm 0.93) \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R61;LINKAGE=AB

 $\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.7±0.6±0.1</b>		91	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

NODE=M057R43  
NODE=M057R43

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90		<sup>2</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
<7	90	7.5 ± 3.4	<sup>3</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  =  $(0.35 \pm 0.05 \pm 0.02) \times 10^{-5}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R43;LINKAGE=A

<sup>2</sup> ABLIKIM 13H reports  $< 0.65 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R43;LINKAGE=AB

<sup>3</sup> NAIK 08 reports  $< 0.75 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R43;LINKAGE=NA

 $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$  $\Gamma_{83}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.4±0.7±0.1</b>		55	<sup>1</sup> ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

NODE=M057R44  
NODE=M057R44

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8	90		<sup>2</sup> ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
<7	90	4.0 ± 3.5	<sup>3</sup> NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

<sup>1</sup> ABLIKIM 18V reports  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  =  $(0.32 \pm 0.06 \pm 0.03) \times 10^{-5}$  which we divide by our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R44;LINKAGE=A

<sup>2</sup> ABLIKIM 13H reports  $< 0.88 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R44;LINKAGE=AB

<sup>3</sup> NAIK 08 reports  $< 0.67 \times 10^{-4}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R44;LINKAGE=NA

 $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$  $\Gamma_{84}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.4±1.7±0.5</b>	131	<sup>1</sup> ABLIKIM	20I BES3	$\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

NODE=M057R99  
NODE=M057R99

<sup>1</sup> ABLIKIM 20I reports  $(4.4 \pm 1.7 \pm 0.5) \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$  assuming  $\text{B}(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ .

NODE=M057R99;LINKAGE=A

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$  $\Gamma_{85}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	<sup>1</sup> ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
<sup>1</sup> ABLIKIM 12I reports $< 17 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .				

NODE=M057R68  
NODE=M057R68

NODE=M057R68;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$  $\Gamma_{86}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	<sup>1</sup> ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Sigma}(1385)^+$
<sup>1</sup> ABLIKIM 12I reports $< 8.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .				

NODE=M057R69  
NODE=M057R69

NODE=M057R69;LINKAGE=AL

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.76 ± 0.32 ± 0.04</b>	51	<sup>1</sup> ABLIKIM 15I	BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$
<sup>1</sup> ABLIKIM 15I reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ $= (1.68 \pm 0.26 \pm 0.15) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M057R81  
NODE=M057R81

NODE=M057R81;LINKAGE=A

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$  $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.83 ± 0.15 ± 0.16</b>		804	<sup>1</sup> ABLIKIM 220	BES3	$\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.0	90	3	<sup>2</sup> NAIK 08	CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
<sup>1</sup> ABLIKIM 220 reports $(1.83 \pm 0.15 \pm 0.16) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ .					
<sup>2</sup> NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .					

NODE=M057R45  
NODE=M057R45

NODE=M057R45;LINKAGE=A

NODE=M057R45;LINKAGE=NA

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$  $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.44 ± 0.12 OUR AVERAGE</b>					
[(1.42 ± 0.32) × 10 <sup>-4</sup> OUR 2022 AVERAGE]					
1.44 ± 0.06 ± 0.11		1691	<sup>1</sup> ABLIKIM 220	BES3	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
1.42 ± 0.31 ± 0.03		29 ± 5	<sup>2</sup> NAIK 08	CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.7	90		<sup>3</sup> ABLIKIM 06D	BES2	$\psi(2S) \rightarrow \chi_{c2} \gamma$
<sup>1</sup> ABLIKIM 220 reports $(1.44 \pm 0.06 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ .					
<sup>2</sup> NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
<sup>3</sup> Using $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$ .					

NODE=M057R17  
NODE=M057R17

NEW

NODE=M057R17;LINKAGE=A

NODE=M057R17;LINKAGE=NA

NODE=M057R17;LINKAGE=AB

 $\Gamma(J/\psi(1S) \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{90}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	90	BARATE 81	SPEC	190 GeV $\pi^- \text{Be} \rightarrow 2\pi 2\mu$

NODE=M057R13  
NODE=M057R13 $\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$  $\Gamma_{91}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10 <sup>-3</sup>	90	<sup>1</sup> ABLIKIM 15N	BES3	$\psi(2S) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$
<sup>1</sup> Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$ .				

NODE=M057R77  
NODE=M057R77

NODE=M057R77;LINKAGE=A

$\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.54 \times 10^{-2}$	90	1,2 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-2}$	90	1,3 ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
-----------------------	----	-------------	----------	---

<sup>1</sup> Using  $1.06 \times 10^8$   $\psi(2S)$  mesons and  $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$ .

<sup>2</sup> From the  $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$  decays.

<sup>3</sup> From the  $\eta_c \rightarrow K^+ K^- \pi^0$  decays.

NODE=M057R76  
NODE=M057R76

OCCUR=2

NODE=M057R76;LINKAGE=A

NODE=M057R76;LINKAGE=B

NODE=M057R76;LINKAGE=C

 $\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})$   $\Gamma_{92}/\Gamma_{42}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<16.4$	90	1 LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

<sup>1</sup> We divided the reported limit by 2 to take into account the  $K_L^0 K^+ \pi^-$  mode.

NODE=M057R64  
NODE=M057R64

NODE=M057R64;LINKAGE=LE

NODE=M057310

————— RADIATIVE DECAYS —————

 $\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$   $\Gamma_{93}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**19.0 ± 0.5 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

$18.64 \pm 0.08 \pm 1.69$	1.0M	1 ABLIKIM	17U BES3	$e^+e^- \rightarrow \gamma X$
---------------------------	------	-----------	----------	-------------------------------

$19.9 \pm 0.5 \pm 1.2$		2 ADAM	05A CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
------------------------	--	--------	----------	---

<sup>1</sup> Not independent from  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))$  and the product  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))$  also measured in ABLIKIM 17U.

<sup>2</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi)$  from ADAM 05A and  $B(\psi(2S) \rightarrow \gamma\chi_{c2})$  from ATHAR 04.

NODE=M057R7  
NODE=M057R7

NODE=M057R7;LINKAGE=A

NODE=M057R7;LINKAGE=AD

 $\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$<19$	90	$13 \pm 11$	1 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
-------	----	-------------	-----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<40$	90	$17.2 \pm 6.8$	2 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
-------	----	----------------	-----------	----------	---

<sup>1</sup> ABLIKIM 11E reports  $< 20.8 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

<sup>2</sup> BENNETT 08A reports  $< 50 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R40  
NODE=M057R40

NODE=M057R40;LINKAGE=AB

NODE=M057R40;LINKAGE=BE

 $\Gamma(\gamma\omega)/\Gamma_{\text{total}}$   $\Gamma_{95}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$<6$	90	$1 \pm 6$	1 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
------	----	-----------	-----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6$	90	$0.0 \pm 1.8$	2 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$
------	----	---------------	-----------	----------	---

<sup>1</sup> ABLIKIM 11E reports  $< 6.1 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

<sup>2</sup> BENNETT 08A reports  $< 7.0 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R41  
NODE=M057R41

NODE=M057R41;LINKAGE=AB

NODE=M057R41;LINKAGE=BE

 $\Gamma(\gamma\phi)/\Gamma_{\text{total}}$   $\Gamma_{96}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$<7$	90	$5 \pm 5$	1 ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
------	----	-----------	-----------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<11$	90	$1.3 \pm 2.5$	2 BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
-------	----	---------------	-----------	----------	---

<sup>1</sup> ABLIKIM 11E reports  $< 8.1 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

<sup>2</sup> BENNETT 08A reports  $< 13 \times 10^{-6}$  from a measurement of  $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.52 \times 10^{-2}$ .

NODE=M057R42  
NODE=M057R42

NODE=M057R42;LINKAGE=AB

NODE=M057R42;LINKAGE=BE

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ VALUE (units $10^{-4}$ ) <b>2.85±0.10 OUR FIT</b>	DOCUMENT ID				$\Gamma_{97}/\Gamma$	NODE=M057R1 NODE=M057R1
$\Gamma(e^+e^-J/\psi(1S))/\Gamma_{\text{total}}$ VALUE (units $10^{-3}$ ) EVTS DOCUMENT ID TECN COMMENT					$\Gamma_{98}/\Gamma$	NODE=M057R82 NODE=M057R82
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2.37±0.15±0.05	1.3k	<sup>1,2</sup> ABLIKIM	17I	BES3	$\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$	
<sup>1</sup> ABLIKIM 17I reports $(2.48 \pm 0.08 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.						NODE=M057R82;LINKAGE=B
<sup>2</sup> Not independent from other measurements reported by ABLIKIM 17I						NODE=M057R82;LINKAGE=C
$\Gamma(e^+e^-J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ VALUE (units $10^{-3}$ ) EVTS DOCUMENT ID TECN COMMENT					$\Gamma_{98}/\Gamma_{93}$	NODE=M057R83 NODE=M057R83
<b>11.3±0.4±0.5</b>	1.3k	<sup>1</sup> ABLIKIM	17I	BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$	
<sup>1</sup> Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.						NODE=M057R83;LINKAGE=A
$\Gamma(\mu^+\mu^-J/\psi(1S))/\Gamma(e^+e^-J/\psi(1S))$ VALUE (units $10^{-2}$ ) EVTS DOCUMENT ID TECN COMMENT					$\Gamma_{99}/\Gamma_{98}$	NODE=M057R96 NODE=M057R96
<b>9.40±0.79±1.15</b>	219	ABLIKIM	19Z	BES3	$\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma(\mu^+\mu^-J/\psi)$	
$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$ VALUE (units $10^{-3}$ ) DOCUMENT ID TECN COMMENT					$\Gamma_{97}/\Gamma_{93}$	NODE=M057R23 NODE=M057R23
<b>1.50±0.05 OUR FIT</b> <b>0.99±0.18</b>		<sup>1</sup> AMBROGIANI	00B	E835	$\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$	
<sup>1</sup> Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .						NODE=M057R;LINKAGE=7A
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$ VALUE (units $10^{-8}$ ) DOCUMENT ID TECN COMMENT					$\Gamma_{97}/\Gamma \times \Gamma_{56}/\Gamma$	NODE=M057R24 NODE=M057R24
<b>2.09±0.13 OUR FIT</b> <b>1.7 ±0.4 OUR AVERAGE</b>						
1.60±0.42		ARMSTRONG	93	E760	$\bar{p}p \rightarrow \gamma\gamma X$	
9.9 ±4.5		BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma X$	
<b><math>\chi_{c2}(1P)</math> CROSS-PARTICLE BRANCHING RATIOS</b>						
$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$ VALUE (units $10^{-3}$ ) DOCUMENT ID TECN COMMENT					$\Gamma_{14}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$	NODE=M057230 NODE=M057B18 NODE=M057B18
<b>2.31±0.26 OUR FIT</b> <b>2.5 ±0.9 OUR AVERAGE</b> Error includes scale factor of 2.3.						
1.90±0.14±0.44		BAI	99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$	
3.8 ±0.67		<sup>1</sup> TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$	
<sup>1</sup> The reported value is derived using $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .						NODE=M057B18;LINKAGE=TA
$\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$ VALUE (units $10^{-4}$ ) DOCUMENT ID TECN COMMENT					$\Gamma_{18}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{\psi(2S)}$	NODE=M057B19 NODE=M057B19
<b>2.1 ±0.4 OUR FIT</b> <b>3.11±0.36±0.48</b>		ABLIKIM	04H	BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$	
$\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$ VALUE (units $10^{-5}$ ) DOCUMENT ID TECN COMMENT					$\Gamma_{56}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$	NODE=M057B1 NODE=M057B1
<b>2.01±0.09 OUR FIT</b> <b>1.4 ±1.1</b>		<sup>1</sup> BAI	98I	BES	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma \bar{p}p$	
<sup>1</sup> Calculated by us. The value for $B(\chi_{c2} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].						NODE=M057B;LINKAGE=J8

$$\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{56}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units  $10^{-6}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**6.98±0.32 OUR FIT**

**7.1 ±0.5 OUR AVERAGE** Error includes scale factor of 1.2.

7.3 ±0.4 ±0.3	405	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
7.2 ±0.7 ±0.4	121 ± 12	<sup>1</sup> NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
4.4 $\begin{smallmatrix} +1.6 \\ -1.4 \end{smallmatrix}$ ±0.6	14.3 $\begin{smallmatrix} +5.2 \\ -4.7 \end{smallmatrix}$	BAI	04F	BES	$\psi(2S) \rightarrow \gamma\chi_{c2}(1P) \rightarrow \gamma p\bar{p}$

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c2} \rightarrow p\bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .

NODE=M057B6  
NODE=M057B6

NODE=M057B6;LINKAGE=NA

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{69}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units  $10^{-6}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**17.4±1.4 OUR FIT**

**17.3±1.5 OUR AVERAGE**

18.2±0.8±1.7	670	ABLIKIM	21L	BES3	$\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+$
15.9±2.1±1.0	71	<sup>1</sup> NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.2±1.4±0.9	207	<sup>2,3</sup> ABLIKIM	13H	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
--------------	-----	------------------------	-----	------	---

<sup>1</sup> Calculated by us. NAIK 08 reports  $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .

<sup>2</sup> Superseded by ABLIKIM 21L

<sup>3</sup> Calculated by us. ABLIKIM 13H reports  $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (20.8 \pm 1.6 \pm 2.3) \times 10^{-5}$  from a measurement of  $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c2})$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.74 \pm 0.35)\%$ .

NODE=M057B10  
NODE=M057B10

NODE=M057B10;LINKAGE=NA

NODE=M057B10;LINKAGE=A  
NODE=M057B10;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{69}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units  $10^{-5}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**5.0±0.4 OUR FIT**

7.1 $\begin{smallmatrix} +3.1 \\ -2.9 \end{smallmatrix}$ ±1.3	8.3 $\begin{smallmatrix} +3.7 \\ -3.4 \end{smallmatrix}$	<sup>1</sup> BAI	03E	BES	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
---	--	------------------	-----	-----	---

<sup>1</sup> BAI 03E reports  $[B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c2}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33  $\begin{smallmatrix} +0.59 \\ -0.55 \end{smallmatrix}$  \pm 0.25)\%$ . We calculate from this measurement the presented value using  $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$  and  $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$ .

NODE=M057B11  
NODE=M057B11

NODE=M057B11;LINKAGE=BA

$$\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{72}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units  $10^{-5}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

<b>1.00±0.20±0.14</b>	32	ABLIKIM	22A0	BES3	$\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+\gamma\gamma$
-----------------------	----	---------	------	------	--

NODE=M057P02  
NODE=M057P02

$$\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{25}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units  $10^{-4}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**2.12±0.08 OUR FIT**

**2.17±0.09 OUR AVERAGE**

2.19±0.05±0.15	4.5k	<sup>1</sup> ABLIKIM	10A	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
2.23±0.06±0.10	2.5k	<sup>2</sup> ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1.90±0.08±0.20	0.8k	<sup>3</sup> ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

<sup>1</sup> Calculated by us. ABLIKIM 10A reports  $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.88 \pm 0.02 \pm 0.06 \pm 0.04) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ . We have multiplied the  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

<sup>2</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c2} \rightarrow \pi^+\pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ . We have multiplied the  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .

<sup>3</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c2} \rightarrow \pi^0\pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ . We have multiplied the  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .

NODE=M057B02  
NODE=M057B02

OCCUR=2

NODE=M057B02;LINKAGE=AB

NODE=M057B02;LINKAGE=AS

NODE=M057B02;LINKAGE=AN

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{25}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.612±0.023 OUR FIT****0.54 ±0.06 OUR AVERAGE**0.66 ±0.18 ±0.37 21 ± 6 <sup>1</sup> BAI 03C BES  $\psi(2S) \rightarrow \gamma\pi^0\pi^0$ 0.54 ±0.05 ±0.04 185 ± 16 <sup>2</sup> BAI 98I BES  $\psi(2S) \rightarrow \gamma\pi^+\pi^-$ <sup>1</sup> We have multiplied  $\pi^0\pi^0$  measurement by 3 to obtain  $\pi\pi$ .<sup>2</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow \pi^+\pi^-)$  reported by BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D]. We have multiplied  $\pi^+\pi^-$  measurement by 3/2 to obtain  $\pi\pi$ .NODE=M057B9  
NODE=M057B9NODE=M057B;LINKAGE=BM  
NODE=M057B;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{31}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**0.52±0.04 OUR FIT****0.52±0.04 OUR AVERAGE**0.54±0.03±0.04 386 <sup>1</sup> ABLIKIM 10A BES3  $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$ 0.47±0.05±0.05 156 ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma\eta\eta$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.44 90 <sup>2</sup> ADAMS 07 CLEO  $\psi(2S) \rightarrow \gamma\chi_{c2}$ < 3 90 BAI 03C BES  $\psi(2S) \rightarrow \gamma\eta\eta \rightarrow 5\gamma$ 0.62±0.31±0.19 LEE 85 CBAL  $\psi(2S) \rightarrow \text{photons}$ <sup>1</sup> Calculated by us. ABLIKIM 10A reports  $B(\chi_{c2} \rightarrow \eta\eta) = (0.65 \pm 0.04 \pm 0.05 \pm 0.03) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ .<sup>2</sup> Superseded by ASNER 09.NODE=M057B04  
NODE=M057B04

NODE=M057B04;LINKAGE=AB

NODE=M057B04;LINKAGE=AD

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{32}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**9.6±0.6 OUR FIT****10.5±0.3±0.6**1.6k <sup>1</sup> ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma K^+K^-$ <sup>1</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c2} \rightarrow K^+K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .NODE=M057B03  
NODE=M057B03

NODE=M057B03;LINKAGE=AS

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{32}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.276±0.017 OUR FIT****0.190±0.034±0.019**115 ± 13 <sup>1</sup> BAI 98I BES  $\psi(2S) \rightarrow \gamma K^+K^-$ <sup>1</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow K^+K^-)$  reported by BAI 98I is derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].NODE=M057B8  
NODE=M057B8

NODE=M057B;LINKAGE=BI

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{33}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.0 ±0.4 OUR FIT****5.0 ±0.4 OUR AVERAGE**4.9 ±0.3 ±0.3 373 ± 20 <sup>1</sup> ASNER 09 CLEO  $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ 5.72±0.76±0.63 65 ABLIKIM 050 BES2  $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ <sup>1</sup> Calculated by us. ASNER 09 reports  $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .NODE=M057B12  
NODE=M057B12

NODE=M057B12;LINKAGE=AS

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{33}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**14.4±1.1 OUR FIT****14.7±4.1±3.3**<sup>1</sup> BAI 99B BES  $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$ <sup>1</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].NODE=M057B13  
NODE=M057B13

NODE=M057B13;LINKAGE=BA

$$\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{42}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.22±0.17 OUR FIT</b>				
<b>1.15±0.18 OUR AVERAGE</b>				

1.21±0.19±0.09	37	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
0.97±0.32±0.13	28	2 ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Calculated by us. ATHAR 07 reports  $B(\chi_{c2} \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (1.3 \pm 0.2 \pm 0.1 \pm 0.1) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$ .

<sup>2</sup> Calculated by us. ABLIKIM 06R reports  $B(\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp) = (0.6 \pm 0.2 \pm 0.1) \times 10^{-3}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.1 \pm 0.6)\%$ . We have multiplied by 2 to obtain  $\bar{K}^0 K^+ \pi^- + \text{c.c.}$  from  $K_S^0 K^\pm \pi^\mp$ .

NODE=M057B05  
NODE=M057B05

NODE=M057B05;LINKAGE=AT

NODE=M057B05;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_1/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10 <sup>-3</sup> )	DOCUMENT ID	TECN	COMMENT
<b>2.79±0.26 OUR FIT</b>			
<b>3.1 ±1.0 OUR AVERAGE</b>			Error includes scale factor of 2.5.

2.3 ±0.1 ±0.5	1 BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
4.3 ±0.6	2 TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

<sup>1</sup> Calculated by us. The value for  $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$  reported in BAI 99B is derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

<sup>2</sup> The value for  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$  reported in TANENBAUM 78 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \times B(J/\psi(1S) \ell^+ \ell^-) = (4.6 \pm 0.7)\%$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

NODE=M057B5  
NODE=M057B5

NODE=M057B;LINKAGE=K1

NODE=M057B;LINKAGE=K2

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{51}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.57±0.19 OUR FIT</b>				
<b>1.76±0.16±0.24</b>	160	1 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$ .

NODE=M057B14  
NODE=M057B14

NODE=M057B14;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{51}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10 <sup>-4</sup> )	DOCUMENT ID	TECN	COMMENT
<b>4.5±0.5 OUR FIT</b>			
<b>3.6±0.6±0.6</b>	1 BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M057B15  
NODE=M057B15

NODE=M057B15;LINKAGE=BA

$$\Gamma(\chi_{c2}(1P) \rightarrow \phi \phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_{20}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.01±0.08 OUR FIT</b>				
<b>0.98±0.13 OUR AVERAGE</b>				Error includes scale factor of 1.3.

0.94±0.03±0.10	849	1 ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma \text{hadrons}$
1.38±0.24±0.23	41	2 ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow \phi \phi)$  reported by ABLIKIM 11K was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35)\%$ .

<sup>2</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow \phi \phi)$  reported by ABLIKIM 06T was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$ .

NODE=M057B16  
NODE=M057B16

NODE=M057B16;LINKAGE=AL

NODE=M057B16;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow \phi \phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{20}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10 <sup>-4</sup> )	DOCUMENT ID	TECN	COMMENT
<b>2.92±0.24 OUR FIT</b>			
<b>4.8 ±1.3 ±1.3</b>	1 BAI	99B BES	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

<sup>1</sup> Calculated by us. The value of  $B(\chi_{c2} \rightarrow \phi \phi)$  reported by BAI 99B was derived using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$  and  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$  [BAI 98D].

NODE=M057B17  
NODE=M057B17

NODE=M057B17;LINKAGE=BA

$$\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{81} / \Gamma \times \Gamma_{168}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.85 ± 0.77 ± 0.44</b>	129	<sup>1</sup> ABLIKIM	19BB BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

NODE=M057B07  
NODE=M057B07

<sup>1</sup> Calculated by us. ABLIKIM 19BB reports  $B(\chi_{c2} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (8.25 \pm 0.83 \pm 0.49) \times 10^{-5}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$  and other branching fractions from PDG 18.

NODE=M057B07;LINKAGE=A

$$\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{82} / \Gamma \times \Gamma_{168}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.87 ± 0.06 ± 0.04</b>	271	<sup>1</sup> ABLIKIM	20AE BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$

NODE=M057P00  
NODE=M057P00

<sup>1</sup> Calculated by us. ABLIKIM 20AE reports  $B(\chi_{c2} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (0.91 \pm 0.06 \pm 0.05) \times 10^{-4}$  using  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$  and other branching fractions from PDG 20.

NODE=M057P00;LINKAGE=A

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{93} / \Gamma \times \Gamma_{168}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.81 ± 0.04 OUR FIT</b>				
<b>1.69 ± 0.16 OUR AVERAGE</b>				Error includes scale factor of 3.4. See the ideogram below.

1.996 ± 0.008 ± 0.070	81k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
1.793 ± 0.008 ± 0.163	1.0M	ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
1.62 ± 0.04 ± 0.12	5.8k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
0.99 ± 0.10 ± 0.08		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
1.47 ± 0.17		<sup>2</sup> OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.8 ± 0.5		<sup>3</sup> BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.2 ± 0.2		<sup>3</sup> BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma \chi_{c2}$
2.2 ± 1.2		<sup>4</sup> BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$
1.2 ± 0.7		<sup>2</sup> WHITAKER	76 MRK1	$e^+ e^-$

NODE=M057B2  
NODE=M057B2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.874 ± 0.007 ± 0.102	76k	<sup>5</sup> ABLIKIM	12O BES3	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.95 ± 0.02 ± 0.07	12.4k	<sup>6</sup> MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
1.85 ± 0.04 ± 0.07	1.9k	<sup>7</sup> ADAM	05A CLEO	Repl. by MENDEZ 08

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ .

NODE=M057B2;LINKAGE=A

<sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

NODE=M057B;LINKAGE=3Q

<sup>3</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .

NODE=M057B;LINKAGE=2Q

<sup>4</sup> Assumes isotropic gamma distribution.

NODE=M057B;LINKAGE=EA

<sup>5</sup> Superseded by ABLIKIM 17N.

NODE=M057B2;LINKAGE=B

<sup>6</sup> Not independent from other measurements of MENDEZ 08.


NODE=M057B2;LINKAGE=ME

<sup>7</sup> Not independent from other values reported by ADAM 05A.

NODE=M057B;LINKAGE=AD



WEIGHTED AVERAGE  
1.69±0.16 (Error scaled by 3.4)



$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{total} \text{ (units } 10^{-2}\text{)}$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \text{ anything})$$

$$\Gamma_{93}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{10}^{\psi(2S)} = \Gamma_{93}/\Gamma \times \Gamma_{168}^{\psi(2S)}/(\Gamma_{12}^{\psi(2S)} + \Gamma_{13}^{\psi(2S)} + \Gamma_{14}^{\psi(2S)} + 0.343\Gamma_{167}^{\psi(2S)} + 0.190\Gamma_{168}^{\psi(2S)})$$

NODE=M057B7

NODE=M057B7

NODE=M057B7

VALUE (units  $10^{-2}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**2.95±0.06 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.12±0.03±0.09	12.4k	<sup>1</sup> MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
3.11±0.07±0.07	1.9k	ADAM	05A	CLEO	Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

NODE=M057B7;LINKAGE=ME

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)$$

$$\Gamma_{93}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

NODE=M057B3

NODE=M057B3

VALUE (units  $10^{-2}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**5.22±0.11 OUR FIT**  
**5.53±0.17 OUR AVERAGE**

5.56±0.05±0.16	12.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
6.0 ±2.8	1.3k	<sup>1</sup> ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
3.9 ±1.2		<sup>2</sup> HIMEL	80	MRK2	$\psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.52±0.13±0.13	1.9k	<sup>3</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
----------------	------	-------------------	-----	------	--------------------

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> The value for  $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$  reported in HIMEL 80 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$  and  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$ .

<sup>3</sup> Not independent from other values reported by ADAM 05A.

NODE=M057B;LINKAGE=AB

NODE=M057B;LINKAGE=H8

NODE=M057B3;LINKAGE=AD

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma \gamma)/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{total}$$

$$\Gamma_{97}/\Gamma \times \Gamma_{168}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

NODE=M057B4

NODE=M057B4

VALUE (units  $10^{-5}$ )    EVTS    DOCUMENT ID    TECN    COMMENT

**2.71±0.08 OUR FIT**  
**2.82±0.10 OUR AVERAGE**

2.83±0.08±0.06	5k	<sup>1</sup> ABLIKIM	17AE	BES3	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$
2.68±0.28±0.15	0.3k	ECKLUND	08A	CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$
7.0 ±2.1 ±2.0		LEE	85	CBAL	$\psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.81±0.17±0.15	1.1k	<sup>2</sup> ABLIKIM	12A	BES3	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow 3\gamma$
----------------	------	----------------------	-----	------	---

- <sup>1</sup> ABLIKIM 17AE measures the ratio of two-photon partial widths for the helicity  $\lambda = 0$  and helicity  $\lambda = 2$  components to be  $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0} / \Gamma_{\gamma\gamma}^{\lambda=2} = 0.000 \pm 0.006 \pm 0.012$ .
- <sup>2</sup> ABLIKIM 12A measures the ratio of two-photon partial widths for the helicity  $\lambda = 0$  and helicity  $\lambda = 2$  components to be  $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0} / \Gamma_{\gamma\gamma}^{\lambda=2} = 0.00 \pm 0.02 \pm 0.02$ . Superseded by ABLIKIM 17AE.

NODE=M057B4;LINKAGE=A

NODE=M057B4;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma) / \Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma) \quad \Gamma_{97} / \Gamma_{97}^{\chi_{c0}(1P)}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.292 ± 0.028 OUR AVERAGE</b>				
0.295 ± 0.014 ± 0.028	8k	<sup>1</sup> ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
0.278 ± 0.050 ± 0.036	0.5k	<sup>1</sup> ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.271 ± 0.029 ± 0.030	1.9k	<sup>1,2</sup> ABLIKIM	12A BES3	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$

NODE=M057B06  
NODE=M057B06<sup>1</sup> Not independent from the values of  $\Gamma(\chi_{c0}, \chi_{c2})$  and  $B(\psi(2S) \rightarrow \chi_{c0}, \chi_{c2})$ .<sup>2</sup> Superseded by ABLIKIM 17AE.NODE=M057B06;LINKAGE=AB  
NODE=M057B06;LINKAGE=A

### MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

NODE=M057240

#### $a_2 = M2 / \sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-11.0 ± 1.0 OUR AVERAGE</b>				
-12.0 ± 1.3 ± 0.4	89k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-9.3 ± 1.6 ± 0.3	19.8k	<sup>2</sup> ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-9.3 <sup>+</sup> <sub>-</sub> 3.9 <sup>+</sup> <sub>-</sub> 4.1 ± 0.6	5.9k	<sup>3</sup> AMBROGIANI	02 E835	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
-14 ± 6	1.9k	<sup>3</sup> ARMSTRONG	93E E760	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
-33.3 <sup>+</sup> <sub>-</sub> 11.6 <sub>-</sub> 29.2	441	<sup>3</sup> OREGLIA	82 CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$

NODE=M057A1  
NODE=M057A1

• • • We do not use the following data for averages, fits, limits, etc. • • •

- 7.9 ± 1.9 ± 0.3 19.8k <sup>4</sup> ARTUSO 09 CLEO  $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$ 

OCCUR=2

<sup>1</sup> Correlated with  $a_3$ ,  $b_2$ , and  $b_3$  with correlation coefficients  $\rho_{a_2 a_3} = 0.733$ ,  $\rho_{a_2 b_2} = -0.605$ , and  $\rho_{a_2 b_3} = -0.095$ .

NODE=M057A1;LINKAGE=B

<sup>2</sup> From a fit with floating  $M2$  amplitudes  $a_2$  and  $b_2$ , and fixed  $E3$  amplitudes  $a_3 = b_3 = 0$ .

NODE=M057A1;LINKAGE=AR

<sup>3</sup> Assuming  $a_3 = 0$ .

NODE=M057A1;LINKAGE=A

<sup>4</sup> From a fit with floating  $M2$  and  $E3$  amplitudes  $a_2$ ,  $b_2$ , and  $a_3$ , and  $b_3$ .

NODE=M057A1;LINKAGE=AT

#### $a_3 = E3 / \sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude


VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-0.3 ± 1.0 OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below.				
-1.3 ± 0.9 ± 0.4	89k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.7 ± 1.4 ± 0.3	19.8k	<sup>2</sup> ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.0 <sup>+</sup> <sub>-</sub> 5.5 <sub>-</sub> 4.4 ± 0.9	5908	AMBROGIANI	02 E835	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
0 <sup>+</sup> <sub>-</sub> 6 <sub>-</sub> 5	1904	ARMSTRONG	93E E760	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$

NODE=M057A2  
NODE=M057A2<sup>1</sup> Correlated with  $a_2$ ,  $b_2$ , and  $b_3$  with correlation coefficients  $\rho_{a_2 a_3} = 0.733$ ,  $\rho_{a_3 b_2} = -0.422$ , and  $\rho_{a_3 b_3} = -0.024$ .

NODE=M057A2;LINKAGE=A

<sup>2</sup> From a fit with floating  $M2$  and  $E3$  amplitudes  $a_2$ ,  $b_2$ , and  $a_3$ , and  $b_3$ .

NODE=M057A2;LINKAGE=AR



$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$  Electric octupole fractional transition amplitude (units  $10^{-2}$ )

**MULTIPOLE AMPLITUDES IN  $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$  RADIATIVE DECAY**

NODE=M057250

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$  Magnetic quadrupole fractional transition amplitude

NODE=M057QB2  
NODE=M057QB2

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.9±0.9 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
1.7±0.8±0.2	89k	1 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
4.6±1.0±1.3	13.8k	2 ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.2±1.5±0.4	19.8k	3 ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 5.1 <sup>+5.4</sup> <sub>-3.6</sub>	721	2 ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
13.2 <sup>+9.8</sup> <sub>-7.5</sub>	441	4 OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.0±1.3±0.3 19.8k 4 ARTUSO 09 CLEO  $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M057QB2;LINKAGE=A

<sup>1</sup> Correlated with  $a_2$ ,  $a_3$ , and  $b_3$  with correlation coefficients  $\rho_{a_2 b_2} = -0.605$ ,  $\rho_{a_3 b_2} = -0.422$ , and  $\rho_{b_2 b_3} = 0.384$ .

<sup>2</sup> From a fit with floating  $M2$  and  $E3$  amplitudes  $b_2$  and  $b_3$ .


NODE=M057QB2;LINKAGE=AB

<sup>3</sup> From a fit with floating  $M2$  and  $E3$  amplitudes  $a_2$ ,  $b_2$ , and  $a_3$ , and  $b_3$ .

NODE=M057QB2;LINKAGE=AT

<sup>4</sup> From a fit with floating  $M2$  amplitudes  $a_2$  and  $b_2$ , and fixed  $E3$  amplitudes  $a_3=b_3=0$ .

NODE=M057QB2;LINKAGE=AR



$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$  Magnetic quadrupole fractional transition amplitude (units  $10^{-2}$ )

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$  Electric octupole fractional transition amplitude

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-1.0±0.6 OUR AVERAGE</b>				
-1.4±0.7±0.4	89k	<sup>1</sup> ABLIKIM 17N BES3		$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.5±0.8±1.8	13.8k	<sup>2</sup> ABLIKIM 11I BES3		$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
-0.8±1.2±0.2	19.8k	ARTUSO 09 CLEO		$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-2.7 <sup>+4.3</sup> <sub>-2.9</sub>	721	<sup>2</sup> ABLIKIM 04I BES2		$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$

<sup>1</sup> Correlated with  $a_2$ ,  $a_3$ , and  $b_2$  with correlation coefficients  $\rho_{a_2 b_3} = -0.095$ ,  $\rho_{a_3 b_3} = -0.024$ , and  $\rho_{b_2 b_3} = 0.384$ .

<sup>2</sup> From a fit with floating  $M2$  and  $E3$  amplitudes  $b_2$  and  $b_3$ .

NODE=M057QB3  
NODE=M057QB3

NODE=M057QB3;LINKAGE=A

NODE=M057QB3;LINKAGE=AB

**MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS**

$\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$  and  $\chi_{c2} \rightarrow \gamma J/\psi(1S)$

NODE=M057260

$b_2/a_2$  Magnetic quadrupole transition amplitude ratio

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>-11<sup>+14</sup></b> <b>-15</b>	19.8k	<sup>1</sup> ARTUSO 09 CLEO		$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M057QAR  
NODE=M057QAR

<sup>1</sup> Statistical and systematic errors combined. From a fit with floating  $M2$  amplitudes  $a_2$  and  $b_2$ , and fixed  $E3$  amplitudes  $a_3=b_3=0$ . Not independent of values for  $a_2(\chi_{c2}(1P))$  and  $b_2(\chi_{c2}(1P))$  from ARTUSO 09.

NODE=M057QAR;LINKAGE=AR

**$\chi_{c2}(1P)$  REFERENCES**

NODE=M057

ABLIKIM 22AO PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61887
ABLIKIM 22O JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61652
ABLIKIM 22Q PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
ABLIKIM 21AV JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61465
ABLIKIM 21L PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61117
ABLIKIM 20AE PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60733
ABLIKIM 20B PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60212
ABLIKIM 20I PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60303
PDG 20 PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM 19AA PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59844
ABLIKIM 19AU PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM 19BB PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60026
ABLIKIM 19J PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59606
ABLIKIM 19Z PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59837
ABLIKIM 18V PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58990
PDG 18 PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AAIJ 17BB EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
AAIJ 17BI PRL 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58278
ABLIKIM 17AE PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58310
ABLIKIM 17AG PR D96 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58316
ABLIKIM 17AI PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58321
ABLIKIM 17I PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM 17N PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978

ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56778
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56779
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55901
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11I	PR D84 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53930
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
DOBBS	06	PR D73 071101	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51062
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50189
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE	02T	PL B540 33	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48813
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48552
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
ACCIARRI	99E	PL B453 73	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46943
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ACKER.,K...	98	PL B439 197	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46324
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
DOMINICK	94	PR D50 4265	J. Dominick <i>et al.</i>	(CLEO Collab.)	REFID=44077
ARMSTRONG	93	PRL 70 2988	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43306
ARMSTRONG	93E	PR D48 3037	T.A. Armstrong <i>et al.</i>	(FNAL-E760 Collab.)	REFID=48616
BAUER	93	PL B302 345	D.A. Bauer <i>et al.</i>	(TPC Collab.)	REFID=43315
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41865
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41907
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BAGLIN	86B	PL B172 455	C. Baglin	(LAPP, CERN, GENO, LYON, OSLO+)	REFID=22145
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
Also		Private Comm.	M.J. Oreglia	(EFI)	REFID=22143
BARATE	81	PR D24 2994	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, CERN+)	REFID=22164
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Quantum numbers are quark model predictions.

NODE=M059

NODE=M059

NODE=M059M

NODE=M059M

NEW

### $\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3637.7±1.1 OUR AVERAGE</b>				Error includes scale factor of 1.2. [3637.5±1.1 MeV OUR 2022 AVERAGE Scale factor = 1.2]
3643.4±2.3±4.4	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
3635.1±3.7±2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6±1.7±0.6	106	<sup>1</sup> AAIJ	17ADLHCB	$pp \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$
3636.4±4.1±0.7	365	<sup>2</sup> AAIJ	17BBLHCB	$pp \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-) X$
3637.0±5.7±3.4	178	<sup>3,4</sup> LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$
3635.1±5.8±2.1	47	<sup>3,5</sup> LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$
3646.9±1.6±3.6	57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6±2.9±1.6	127 ± 18	<sup>6</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
3638.5±1.5±0.8	624	<sup>3</sup> DEL-AMO-SA...11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5±3.2±2.5	1201	<sup>3</sup> DEL-AMO-SA...11M	BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 <sup>+3.9+0.7</sup> <sub>-4.2-2.0</sub>	128	<sup>7</sup> VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	<sup>8</sup> ABE	07 BELL	$e^+ e^- \rightarrow J/\psi (c \bar{c})$
3645.0±5.5 <sup>+4.9</sup> <sub>-7.8</sub>	121 ± 27	AUBERT	05c BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
3642.9±3.1±1.5	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3639 ± 7	98 ± 52	<sup>9</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X c \bar{c}$
3630.8±3.4±1.0	112 ± 24	<sup>10</sup> AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39 ± 11	<sup>11</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		<sup>12</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup>AAIJ 17AD report  $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$  MeV. We use the current value  $m_{\psi(2S)} = 3686.097 \pm 0.025$  MeV to obtain the quoted mass.

<sup>2</sup>From a fit of the  $\phi\phi$  invariant mass with the width of  $\eta_c(2S)$  fixed to the PDG 16 value.

<sup>3</sup>Ignoring possible interference with continuum.

<sup>4</sup>With a width fixed to 11.3 MeV.

<sup>5</sup>With a width fixed to 11.3 MeV. Using both  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow \pi^+ \pi^- \pi^0$  decays.

<sup>6</sup>From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>7</sup>Accounts for interference with non-resonant continuum.

<sup>8</sup>From a fit of the  $J/\psi$  recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

<sup>9</sup>From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>10</sup>Superseded by DEL-AMO-SANCHEZ 11M.

<sup>11</sup>Superseded by VINOKUROVA 11.

<sup>12</sup>Assuming mass of  $\psi(2S) = 3686$  MeV.

NODE=M059M;LINKAGE=B

NODE=M059M;LINKAGE=C

NODE=M059M;LINKAGE=DE

NODE=M059M;LINKAGE=LE

NODE=M059M;LINKAGE=LS

NODE=M059M;LINKAGE=AB

NODE=M059M;LINKAGE=VA

NODE=M059M;LINKAGE=EB

NODE=M059M;LINKAGE=AU

NODE=M059M;LINKAGE=AR

NODE=M059M;LINKAGE=CH

NODE=M059M;LINKAGE=A

### $\eta_c(2S)$ WIDTH

NODE=M059W

NODE=M059W

NEW

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.9± 2.6 OUR AVERAGE</b>					
[11.3 <sup>+3.2</sup> <sub>-2.9</sub> MeV OUR 2022 AVERAGE]					
19.8± 3.9±3.1		569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
9.9± 4.8±2.9		57 ± 17	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9± 6.4±4.8		127 ± 18	<sup>1</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
13.4± 4.6±3.2		624	<sup>2</sup> DEL-AMO-SA...11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 <sup>+ 8.4+2.6</sup> <sub>- 5.1-0.9</sub>		128	<sup>3</sup> VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3±12.4±4.0		61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 23	90	98 ± 52	<sup>4</sup> AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
22 ± 14		121 ± 27	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c\bar{c}$
17.0 ± 8.3 ± 2.5		112 ± 24	<sup>5</sup> AUBERT	04D BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K\bar{K}\pi$
< 55	90	39 ± 11	<sup>6</sup> CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
< 8.0	95		<sup>7</sup> EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> From a simultaneous fit to  $K_S^0 K^\pm \pi^\mp$  and  $K^+ K^- \pi^0$  decay modes.

<sup>2</sup> Ignoring possible interference with continuum.

<sup>3</sup> Accounts for interference with non-resonant continuum.

<sup>4</sup> From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

<sup>5</sup> Superseded by DEL-AMO-SANCHEZ 11M.

<sup>6</sup> For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.

<sup>7</sup> For a mass value of 3594 ± 5 MeV

NODE=M059W;LINKAGE=AB  
 NODE=M059W;LINKAGE=DE  
 NODE=M059W;LINKAGE=VA  
 NODE=M059W;LINKAGE=AU  
 NODE=M059W;LINKAGE=AR  
 NODE=M059W;LINKAGE=W2  
 NODE=M059W;LINKAGE=W

## $\eta_c(2S)$ DECAY MODES

NODE=M059215;NODE=M059

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	not seen	
$\Gamma_2$ $K\bar{K}\pi$	( 1.9±1.2 ) %	
$\Gamma_3$ $K\bar{K}\eta$	( 5 ± 4 ) × 10 <sup>-3</sup>	
$\Gamma_4$ $2\pi^+ 2\pi^-$	< 2.1 %	90%
$\Gamma_5$ $\rho^0 \rho^0$	< 1.9 × 10 <sup>-3</sup>	90%
$\Gamma_6$ $3\pi^+ 3\pi^-$	( 1.3±0.9 ) %	
$\Gamma_7$ $K^+ K^- \pi^+ \pi^-$	< 1.4 %	90%
$\Gamma_8$ $K^{*0} \bar{K}^{*0}$	< 2.9 × 10 <sup>-3</sup>	90%
$\Gamma_9$ $K^+ K^- \pi^+ \pi^- \pi^0$	( 1.4±1.0 ) %	
$\Gamma_{10}$ $K^+ K^- 2\pi^+ 2\pi^-$	< 1.4 %	90%
$\Gamma_{11}$ $K_S^0 K^- 2\pi^+ \pi^- + c.c.$	( 1.0±0.8 ) %	
$\Gamma_{12}$ $2K^+ 2K^-$	< 1.3 × 10 <sup>-3</sup>	90%
$\Gamma_{13}$ $\phi\phi$	< 1.1 × 10 <sup>-3</sup>	90%
$\Gamma_{14}$ $p\bar{p}$	< 2.0 × 10 <sup>-3</sup>	90%
$\Gamma_{15}$ $p\bar{p}\pi^+ \pi^-$	seen	
$\Gamma_{16}$ $\gamma\gamma$	( 1.6±1.0 ) × 10 <sup>-4</sup>	
$\Gamma_{17}$ $\gamma J/\psi(1S)$	< 1.4 %	90%
$\Gamma_{18}$ $\pi^+ \pi^- \eta$	< 6 × 10 <sup>-3</sup>	90%
$\Gamma_{19}$ $\pi^+ \pi^- \eta'$	( 2.6±1.9 ) × 10 <sup>-3</sup>	
$\Gamma_{20}$ $\pi^+ \pi^- \eta_c(1S)$	< 25 %	90%

DESIG=1  
 DESIG=4  
 DESIG=20  
 DESIG=5  
 DESIG=16  
 DESIG=8  
 DESIG=6  
 DESIG=17  
 DESIG=9  
 DESIG=10  
 DESIG=11  
 DESIG=7  
 DESIG=18  
 DESIG=3  
 DESIG=22  
 DESIG=2  
 DESIG=21  
 DESIG=12  
 DESIG=13  
 DESIG=15

## $\eta_c(2S)$ PARTIAL WIDTHS

NODE=M059216

$\Gamma(\gamma\gamma)$

$\Gamma_{16}$

VALUE (keV) EVTS DOCUMENT ID TECN COMMENT

NODE=M059W1  
 NODE=M059W1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.44±0.14	106	<sup>1</sup> XU	18	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
1.3 ± 0.6		<sup>2</sup> ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

<sup>1</sup> Assuming that the branching fraction into  $\eta' \pi^+ \pi^-$  is the same as for  $\eta_c(1S)$ .

<sup>2</sup> They measure  $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K\bar{K}\pi)$ . The value for  $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$  is derived assuming that the branching fractions for  $\eta_c(2S)$  and  $\eta_c(1S)$  decays to  $K_S K\pi$  are equal and using  $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$  keV.

NODE=M059W1;LINKAGE=A  
 NODE=M059W1;LINKAGE=AS

$\Gamma(\gamma\gamma) \times \Gamma(\pi^+ \pi^- \eta')/\Gamma_{total}$

$\Gamma_{16}\Gamma_{19}/\Gamma$

VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M059R29  
 NODE=M059R29

<b>5.6<sup>+1.2</sup><sub>-1.1</sub> ± 1.1</b>	106	XU	18	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
--	-----	----	----	------	---

## $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(total)$

NODE=M059218

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{total}$

$\Gamma_2\Gamma_{16}/\Gamma$

VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

NODE=M059G04  
 NODE=M059G04

<b>41 ± 4 ± 6</b>	624	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
-------------------	-----	------------------------------	------	--

<sup>1</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059G04;LINKAGE=DE

$$\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_{16}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

NODE=M059G01  
NODE=M059G01

$$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_7\Gamma_{16}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

NODE=M059G02  
NODE=M059G02

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_9\Gamma_{16}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$30 \pm 6 \pm 5$	1201	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M059G05  
NODE=M059G05

<sup>1</sup> Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059G05;LINKAGE=DE

$$\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{12}\Gamma_{16}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

NODE=M059G03  
NODE=M059G03

$$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{20}\Gamma_{16}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<133	90	LEES	12AE	BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

NODE=M059G06  
NODE=M059G06

$$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$$

NODE=M059217

$$\Gamma(\bar{p}p)/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$$

VALUE (units $10^{-8}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	<sup>1,2,3</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

NODE=M059G1  
NODE=M059G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.0	90	<sup>1,2,4</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
-------	----	--------------------------------	------	-------------------------------------

OCCUR=2

<12.0	90	<sup>2,4</sup> AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$
-------	----	------------------------------	------	-------------------------------------

OCCUR=3

<sup>1</sup> Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

NODE=M059G1;LINKAGE=A

<sup>2</sup> For a total width  $\Gamma=5$  MeV.

NODE=M059G1;LINKAGE=B

<sup>3</sup> For the resonance mass region 3589–3599 MeV/ $c^2$ .

NODE=M059G1;LINKAGE=C1

<sup>4</sup> For the resonance mass region 3575–3660 MeV/ $c^2$ .

NODE=M059G1;LINKAGE=C2

## $\eta_c(2S)$ BRANCHING RATIOS

NODE=M059220

$$\Gamma(\text{hadrons})/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABREU	98O	DLPH $e^+e^- \rightarrow e^+e^- + \text{hadrons}$

NODE=M059R1  
NODE=M059R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>1</sup> EDWARDS	82C	CBAL $e^+e^- \rightarrow \gamma X$
------	----------------------	-----	------------------------------------

<sup>1</sup> For a mass value of  $3594 \pm 5$  MeV

NODE=M059R;LINKAGE=W

$$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.4 \pm 1.1$	$59 \pm 12$	<sup>1</sup> AUBERT	08AB	BABR $B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$

NODE=M059R3  
NODE=M059R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	$127 \pm 18$	ABLIKIM	12G	BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$
------	--------------	---------	-----	--

seen	$39 \pm 11$	<sup>2</sup> CHOI	02	BELL $B \rightarrow K K_S K^- \pi^+$
------	-------------	-------------------	----	--------------------------------------

<sup>1</sup> Derived from a measurement of  $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6_{-1.9}^{+2.0} \pm 2.5)\%$  and using  $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$ , and  $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77_{-0.66}^{+0.55}) \times 10^{-5}$ .

NODE=M059R3;LINKAGE=AU

<sup>2</sup> For a mass value of  $3654 \pm 6$  MeV

NODE=M059R;LINKAGE=W2

$$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi) \quad \Gamma_3/\Gamma_2$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$27.3 \pm 7.0 \pm 9.0$	225	<sup>1</sup> LEES	14E	BABR $\gamma\gamma \rightarrow K^+K^-\gamma\gamma$

NODE=M059R26  
NODE=M059R26

<sup>1</sup> LEES 14E reports  $B(\eta_c(2S) \rightarrow K^+K^-\eta)/B(\eta_c(2S) \rightarrow K^+K^-\pi^0) = 0.82 \pm 0.21 \pm 0.27$ , which we divide by 3 to account for isospin symmetry.

NODE=M059R26;LINKAGE=LE



$\Gamma(2\pi^+2\pi^-)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M059R01  
NODE=M059R01

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	--

 $\Gamma(\rho^0\rho^0)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M059R15  
NODE=M059R15

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
----------	---------	-----	--

 $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M059R02  
NODE=M059R02

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	--

 $\Gamma(K^{*0}\bar{K}^{*0})/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M059R16  
NODE=M059R16

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
----------	---------	-----	--

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K\bar{K}\pi)$  $\Gamma_9/\Gamma_2$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M059R21  
NODE=M059R21

<b>0.73±0.17±0.17</b>	1201	<sup>1</sup> DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-----------------------	------	------------------------------	------	--

<sup>1</sup>We have multiplied the value of  $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$  reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain  $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma(K\bar{K}\pi)$ . Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059R21;LINKAGE=DE

 $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M059R22  
NODE=M059R22

••• We do not use the following data for averages, fits, limits, etc. •••

seen	57±17	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
------	-------	---------	-----	--

 $\Gamma(2K^+2K^-)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M059R03  
NODE=M059R03

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$
----------	--------	----	--

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M059R17  
NODE=M059R17

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$
----------	---------	-----	--

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M059R04  
NODE=M059R04

••• We do not use the following data for averages, fits, limits, etc. •••

seen	106	<sup>1</sup> AAIJ	17AD	LHCB $pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$
------	-----	-------------------	------	---

OCCUR=2

<sup>1</sup>AAIJ 17AD report a 6.4 standard deviation signal, with  $B(B^+ \rightarrow \eta_c(2S)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$ .

NODE=M059R04;LINKAGE=A

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M059R30  
NODE=M059R30

seen	110	<sup>1</sup> CHILIKIN	19	BELL $e^+e^- \rightarrow \mathcal{T}(4S)$
------	-----	-----------------------	----	---

<sup>1</sup>CHILIKIN 19 reports signals in  $B^+ \rightarrow \eta_c(2S)K^+$  and  $B^0 \rightarrow \eta_c(2S)K_S^0$  with 12.3 and 5.9 standard deviations, respectively.

NODE=M059R30;LINKAGE=A

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R2  
NODE=M059R2

••• We do not use the following data for averages, fits, limits, etc. •••

<4 × 10 <sup>-4</sup>	90	<sup>1</sup> WICHT	08	BELL $B^\pm \rightarrow K^\pm \gamma\gamma$
not seen		AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$

<0.01	90	LEE	85	CBAL $\psi' \rightarrow \text{photons}$
-------	----	-----	----	---

<sup>1</sup>WICHT 08 reports  $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$ .

NODE=M059R2;LINKAGE=WI

$$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma(K \bar{K} \pi)$$

$$\Gamma_{20}/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.33	90	<sup>1</sup> LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

<sup>1</sup>We divided the reported limit by 3 to take into account isospin relations.

NODE=M059R23  
NODE=M059R23

NODE=M059R23;LINKAGE=LE

### $\eta_c(2S)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M059230

$$\Gamma(\eta_c(2S) \rightarrow K \bar{K} \eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<11.8 × 10 <sup>-6</sup>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \eta$
<sup>1</sup> CRONIN-HENNESSY 10 reports a limit of < 5.9 × 10 <sup>-6</sup> for the decay $\eta_c(2S) \rightarrow K^+ K^- \eta$ which we multiply by 2 account for isospin symmetry. It assumes $\Gamma(\eta_c(2S)) = 14$ MeV. It also gives the analytic dependence of limits on width.				

NODE=M059R25  
NODE=M059R25

NODE=M059R25;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_4/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<14.6 × 10 <sup>-6</sup>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
<sup>1</sup> Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.				

NODE=M059R05  
NODE=M059R05

NODE=M059R05;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \rho^0 \rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<12.7 × 10 <sup>-7</sup>	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$
--------------------------	----	---------	----------	---

NODE=M059R18  
NODE=M059R18

$$\Gamma(\eta_c(2S) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE (units 10 <sup>-6</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

9.2 ± 1.0 ± 1.2	569		ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
-----------------	-----	--	---------	----------	--

••• We do not use the following data for averages, fits, limits, etc. •••

<13.2	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<sup>1</sup> Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.				

NODE=M059R06  
NODE=M059R06

NODE=M059R06;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_7/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<9.6 × 10 <sup>-6</sup>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
-------------------------	----	-----------------------------	------	---

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R07  
NODE=M059R07

NODE=M059R07;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^{*0} \bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_8/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<19.6 × 10 <sup>-7</sup>	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
--------------------------	----	---------	----------	---

NODE=M059R19  
NODE=M059R19

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_9/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<43.0 × 10 <sup>-6</sup>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$
--------------------------	----	-----------------------------	------	---

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R08  
NODE=M059R08

NODE=M059R08;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

$$\Gamma_{10}/\Gamma \times \Gamma_{170}^{\psi(2S)}/\Gamma\psi(2S)$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<9.7 × 10 <sup>-6</sup>	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$
-------------------------	----	-----------------------------	------	---

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R09  
NODE=M059R09

NODE=M059R09;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE (units $10^{-6}$ )	CL% EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60	ABLIKIM 13K	BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

NODE=M059R10  
NODE=M059R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.2	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
--------	----	-----------------------------	------	--

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R10;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \phi \phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< $7.8 \times 10^{-7}$	90	ABLIKIM 11H	BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M059R20  
NODE=M059R20

$$\Gamma(\eta_c(2S) \rightarrow \rho \bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< $1.4 \times 10^{-6}$	90	ABLIKIM 13V	BES3	$\psi(2S) \rightarrow \gamma \rho \bar{\rho}$

NODE=M059R24  
NODE=M059R24

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< $9.7 \times 10^{-6}$	90	33	<sup>1</sup> ABLIKIM 17N	BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$

NODE=M059R27  
NODE=M059R27

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ .

NODE=M059R27;LINKAGE=A

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< $4.3 \times 10^{-6}$	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$

NODE=M059R11  
NODE=M059R11

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R11;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< $14.2 \times 10^{-6}$	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M059R12  
NODE=M059R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R12;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{\text{total}}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< $1.7 \times 10^{-4}$	90	<sup>1</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

NODE=M059R14  
NODE=M059R14

<sup>1</sup> Assuming  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R14;LINKAGE=CR

## $\eta_c(2S)$ REFERENCES


ABLIKIM 22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
CHILIKIN 19	PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)	REFID=59899
XU 18	PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)	REFID=59453
AAIJ 17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
AAIJ 17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
ABLIKIM 17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
PDG 16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
LEES 14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
ABLIKIM 13K	PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54956
ABLIKIM 13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
ABLIKIM 12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54272
LEES 12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
ABLIKIM 11H	PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53929
DEL-AMO-SA... 11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
VINOKUROVA 11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=53927
CRONIN-HEN... 10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=53233

NODE=M059

AUBERT	08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)
AUBERT	06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	05C	PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)
ASNER	04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)
AUBERT	04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)
CHOI	02	PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)
AMBROGIANI	01	PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)
ABREU	98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ARMSTRONG	95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
LEE	85	SLAC 282	R.A. Lee	(SLAC)
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)

REFID=52267  
REFID=52064  
REFID=52204  
REFID=51627  
REFID=51059  
REFID=50773  
REFID=50182  
REFID=49745  
REFID=49746  
REFID=49188  
REFID=48760  
REFID=48340  
REFID=46553  
REFID=44623  
REFID=40589  
REFID=22173

NODE=M071



$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on " $\psi(2S)$  and  $\chi_c$  branching ratios" before the  $\chi_{c0}(1P)$  Listings.

NODE=M071

### $\psi(2S)$ MASS

NODE=M071M

OUR FIT includes measurements of  $m_{\psi(2S)}$ ,  $m_{\psi(3770)}$ , and  $m_{\psi(3770)} - m_{\psi(2S)}$ .

NODE=M071M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3686.10 ± 0.06</b>	<b>OUR FIT</b>	Error includes scale factor of 5.9.		
<b>3686.097 ± 0.010</b>	<b>OUR AVERAGE</b>			
3686.099 ± 0.004 ± 0.009		<sup>1</sup> ANASHIN 15	KEDR	$e^+e^- \rightarrow$ hadrons
3686.12 ± 0.06 ± 0.10	4k	AAIJ 12H	LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
3685.95 ± 0.10	413	<sup>2</sup> ARTAMONOV 00	OLYA	$e^+e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		<sup>3</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3686.114 ± 0.007 <sup>+0.011</sup> <sub>-0.016</sub>		<sup>4</sup> ANASHIN 12	KEDR	$e^+e^- \rightarrow$ hadrons
3686.111 ± 0.025 ± 0.009		AULCHENKO 03	KEDR	$e^+e^- \rightarrow$ hadrons
3686.00 ± 0.10	413	<sup>5</sup> ZHOLENTZ 80	OLYA	$e^+e^-$

NODE=M071M

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup>Supersedes AULCHENKO 03 and ANASHIN 12.
- <sup>2</sup>Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).
- <sup>3</sup>Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the  $J/\psi(1S)$  mass from AULCHENKO 03.
- <sup>4</sup>From the scans in 2004 and 2006. ANASHIN 12 reports the value  $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$  MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.
- <sup>5</sup>Superseded by ARTAMONOV 00.

NODE=M071M;LINKAGE=A  
NODE=M071M;LINKAGE=AR

NODE=M071M;LINKAGE=NW

NODE=M071M;LINKAGE=AN

NODE=M071M;LINKAGE=RZ

### $m_{\psi(2S)} - m_{J/\psi(1S)}$

NODE=M071DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>589.188 ± 0.028</b>	<b>OUR AVERAGE</b>		
589.194 ± 0.027 ± 0.011	<sup>1</sup> AULCHENKO 03	KEDR	$e^+e^- \rightarrow$ hadrons
589.7 ± 1.2	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
589.07 ± 0.13	<sup>1</sup> ZHOLENTZ 80	OLYA	$e^+e^-$
588.7 ± 0.8	LUTH 75	MRK1	
588 ± 1	<sup>2</sup> BAI 98E	BES	$e^+e^-$

NODE=M071DM

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup>Redundant with data in mass above.
- <sup>2</sup>Systematic errors not evaluated.

NODE=M071DM;LINKAGE=R  
NODE=M071DM;LINKAGE=BD

### $\psi(2S)$ WIDTH

NODE=M071W

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>294 ± 8</b>	<b>OUR FIT</b>			
<b>286 ± 16</b>	<b>OUR AVERAGE</b>			
358 ± 88 ± 4		ABLIKIM 08B	BES2	$e^+e^- \rightarrow$ hadrons
290 ± 25 ± 4	2.7k	ANDREOTTI 07	E835	$\bar{p}p \rightarrow e^+e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM 06L	BES2	$e^+e^- \rightarrow$ hadrons
264 ± 27		<sup>1</sup> BAI 02B	BES2	$e^+e^-$
287 ± 37 ± 16		<sup>2</sup> ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$

NODE=M071W

<sup>1</sup>From a simultaneous fit to the hadronic and  $\mu^+\mu^-$  cross section, assuming  $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$  and lepton universality. Does not include vacuum polarization correction.

<sup>2</sup>The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

NODE=M071W;LINKAGE=BC

NODE=M071W;LINKAGE=AN

NODE=M071220;NODE=M071

 **$\psi(2S)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level	
$\Gamma_1$ hadrons	(97.85 $\pm$ 0.13 ) %		DESIG=3
$\Gamma_2$ virtual $\gamma \rightarrow$ hadrons	( 1.73 $\pm$ 0.14 ) %	S=1.5	DESIG=4
$\Gamma_3$ $ggg$	(10.6 $\pm$ 1.6 ) %		DESIG=255
$\Gamma_4$ $\gamma gg$	( 1.03 $\pm$ 0.29 ) %		DESIG=256
$\Gamma_5$ light hadrons	(15.4 $\pm$ 1.5 ) %		DESIG=226
$\Gamma_6$ $K_S^0$ anything	(16.0 $\pm$ 1.1 ) %		DESIG=325
$\Gamma_7$ $e^+e^-$	( 7.93 $\pm$ 0.17 ) $\times 10^{-3}$		DESIG=1
$\Gamma_8$ $\mu^+\mu^-$	( 8.0 $\pm$ 0.6 ) $\times 10^{-3}$		DESIG=2
$\Gamma_9$ $\tau^+\tau^-$	( 3.1 $\pm$ 0.4 ) $\times 10^{-3}$		DESIG=68
<b>Decays into <math>J/\psi(1S)</math> and anything</b>			
$\Gamma_{10}$ $J/\psi(1S)$ anything	(61.4 $\pm$ 0.6 ) %		NODE=M071;CLUMP=A DESIG=11
$\Gamma_{11}$ $J/\psi(1S)$ neutrals	(25.38 $\pm$ 0.32 ) %		DESIG=12
$\Gamma_{12}$ $J/\psi(1S)\pi^+\pi^-$	(34.68 $\pm$ 0.30 ) %		DESIG=13
$\Gamma_{13}$ $J/\psi(1S)\pi^0\pi^0$	(18.24 $\pm$ 0.31 ) %		DESIG=14
$\Gamma_{14}$ $J/\psi(1S)\eta$	( 3.37 $\pm$ 0.05 ) %		DESIG=15
$\Gamma_{15}$ $J/\psi(1S)\pi^0$	( 1.268 $\pm$ 0.032) $\times 10^{-3}$		DESIG=18
<b>Hadronic decays</b>			
$\Gamma_{16}$ $\pi^+\pi^-$	( 7.8 $\pm$ 2.6 ) $\times 10^{-6}$		NODE=M071;CLUMP=B DESIG=21
$\Gamma_{17}$ $\pi^+\pi^-\pi^0$	( 2.01 $\pm$ 0.17 ) $\times 10^{-4}$	S=1.7	DESIG=36
$\Gamma_{18}$ $\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$	( 3.2 $\pm$ 1.2 ) $\times 10^{-5}$	S=1.8	DESIG=22
$\Gamma_{19}$ $\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	( 1.9 $\pm$ 1.2 -0.4 ) $\times 10^{-4}$		DESIG=201
$\Gamma_{20}$ $2(\pi^+\pi^-)$	( 2.4 $\pm$ 0.6 ) $\times 10^{-4}$	S=2.2	DESIG=24
$\Gamma_{21}$ $\rho^0\pi^+\pi^-$	( 2.2 $\pm$ 0.6 ) $\times 10^{-4}$	S=1.4	DESIG=33
$\Gamma_{22}$ $2(\pi^+\pi^-)\pi^0$	( 2.9 $\pm$ 1.0 ) $\times 10^{-3}$	S=4.7	DESIG=25
$\Gamma_{23}$ $\rho a_2(1320)$	( 2.6 $\pm$ 0.9 ) $\times 10^{-4}$		DESIG=65
$\Gamma_{24}$ $\pi^+\pi^-\pi^0\pi^0\pi^0$	( 5.3 $\pm$ 0.9 ) $\times 10^{-3}$		DESIG=312
$\Gamma_{25}$ $\pi^+\pi^-4\pi^0$	( 1.4 $\pm$ 1.0 ) $\times 10^{-3}$		DESIG=332
$\Gamma_{26}$ $\rho^\pm\pi^\mp\pi^0\pi^0$	< 2.7 $\times 10^{-3}$	CL=90%	DESIG=315
$\Gamma_{27}$ $3(\pi^+\pi^-)$	( 3.5 $\pm$ 2.0 ) $\times 10^{-4}$	S=2.8	DESIG=32
$\Gamma_{28}$ $2(\pi^+\pi^-\pi^0)$	( 4.8 $\pm$ 1.5 ) $\times 10^{-3}$		DESIG=221
$\Gamma_{29}$ $3(\pi^+\pi^-)\pi^0$	( 3.5 $\pm$ 1.6 ) $\times 10^{-3}$		DESIG=37
$\Gamma_{30}$ $2(\pi^+\pi^-)3\pi^0$	( 1.42 $\pm$ 0.31 ) %		DESIG=329
$\Gamma_{31}$ $\eta\pi^+\pi^-$	< 1.6 $\times 10^{-4}$	CL=90%	DESIG=202
$\Gamma_{32}$ $\eta\pi^+\pi^-\pi^0$	( 9.5 $\pm$ 1.7 ) $\times 10^{-4}$		DESIG=203
$\Gamma_{33}$ $\eta 2(\pi^+\pi^-)$	( 1.2 $\pm$ 0.6 ) $\times 10^{-3}$		DESIG=251
$\Gamma_{34}$ $\eta\pi^+\pi^-\pi^0\pi^0$	< 4 $\times 10^{-4}$	CL=90%	DESIG=313
$\Gamma_{35}$ $\eta\pi^+\pi^-3\pi^0$	< 2.1 $\times 10^{-3}$	CL=90%	DESIG=334
$\Gamma_{36}$ $\eta 2(\pi^+\pi^-\pi^0)$	< 2.1 $\times 10^{-3}$	CL=90%	DESIG=328
$\Gamma_{37}$ $\rho\eta$	( 2.2 $\pm$ 0.6 ) $\times 10^{-5}$	S=1.1	DESIG=94
$\Gamma_{38}$ $\eta'\pi^+\pi^-\pi^0$	( 4.5 $\pm$ 2.1 ) $\times 10^{-4}$		DESIG=204
$\Gamma_{39}$ $\eta'\rho$	( 1.9 $\pm$ 1.7 -1.2 ) $\times 10^{-5}$		DESIG=93
$\Gamma_{40}$ $\omega\pi^0$	( 2.1 $\pm$ 0.6 ) $\times 10^{-5}$		DESIG=92
$\Gamma_{41}$ $\omega\pi^+\pi^-$	( 7.3 $\pm$ 1.2 ) $\times 10^{-4}$	S=2.1	DESIG=75
$\Gamma_{42}$ $\omega\pi^+\pi^-2\pi^0$	( 8.7 $\pm$ 2.4 ) $\times 10^{-3}$		DESIG=327
$\Gamma_{43}$ $b_1^\pm\pi^\mp$	( 4.0 $\pm$ 0.6 ) $\times 10^{-4}$	S=1.1	DESIG=40
$\Gamma_{44}$ $\omega f_2(1270)$	( 2.2 $\pm$ 0.4 ) $\times 10^{-4}$		DESIG=64
$\Gamma_{45}$ $\omega\pi^0\pi^0$	( 1.11 $\pm$ 0.35 ) $\times 10^{-3}$		DESIG=314
$\Gamma_{46}$ $\omega 3\pi^0$	< 8 $\times 10^{-4}$	CL=90%	DESIG=333
$\Gamma_{47}$ $b_1^0\pi^0$	( 2.4 $\pm$ 0.6 ) $\times 10^{-4}$		DESIG=193
$\Gamma_{48}$ $\omega\eta$	< 1.1 $\times 10^{-5}$	CL=90%	DESIG=95

Г49	$\omega\eta'$	$(3.2^{+2.5}_{-2.1}) \times 10^{-5}$		DESIG=91
Г50	$\phi\pi^0$	$< 4 \times 10^{-7}$	CL=90%	DESIG=96
Г51	$\phi\pi^+\pi^-$	$(1.18 \pm 0.26) \times 10^{-4}$	S=1.5	DESIG=78
Г52	$\phi f_0(980) \rightarrow \pi^+\pi^-$	$(7.5 \pm 3.3) \times 10^{-5}$	S=1.6	DESIG=81
Г53	$\phi\eta$	$(3.10 \pm 0.31) \times 10^{-5}$		DESIG=89
Г54	$\eta\phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+\pi^-$	$< 2.2 \times 10^{-6}$	CL=90%	DESIG=316
Г55	$\phi\eta'$	$(1.54 \pm 0.20) \times 10^{-5}$		DESIG=90
Г56	$\phi f_1(1285)$	$(3.0 \pm 1.3) \times 10^{-5}$		DESIG=319
Г57	$\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta$	$(8.5 \pm 1.7) \times 10^{-6}$		DESIG=320
Г58	$\phi f_2'(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$		DESIG=67
Г59	$K^+K^-$	$(7.5 \pm 0.5) \times 10^{-5}$		DESIG=23
Г60	$K^+K^-\pi^+$	$(7.3 \pm 0.5) \times 10^{-4}$		DESIG=26
Г61	$K^+K^-\pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$		DESIG=38
Г62	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$		DESIG=86
Г63	$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$		DESIG=85
Г64	$K_S^0 K_L^0 \pi^0$	$< 3.0 \times 10^{-4}$	CL=90%	DESIG=303
Г65	$K^+K^-\pi^0\pi^0$	$(2.6 \pm 1.3) \times 10^{-4}$		DESIG=298
Г66	$K^+K^-\pi^+\pi^-\pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$		DESIG=206
Г67	$\omega f_0(1710) \rightarrow \omega K^+K^-$	$(5.9 \pm 2.2) \times 10^{-5}$		DESIG=216
Г68	$K^*(892)^0 K^-\pi^+\pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$		DESIG=217
Г69	$K^*(892)^+ K^-\pi^+\pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$		DESIG=218
Г70	$K^*(892)^+ K^-\rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$		DESIG=219
Г71	$K^*(892)^0 K^-\rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$		DESIG=220
Г72	$K_S^0 K_S^0 \pi^+\pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=225
Г73	$K_S^0 K_L^0 \pi^0\pi^0$	$(1.3 \pm 0.6) \times 10^{-3}$		DESIG=304
Г74	$K_S^0 K_L^0 \eta$	$(1.3 \pm 0.5) \times 10^{-3}$		DESIG=305
Г75	$K^+K^-\rho^0$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=205
Г76	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$		DESIG=66
Г77	$K^+K^-\pi^+\pi^-\eta$	$(1.3 \pm 0.7) \times 10^{-3}$		DESIG=252
Г78	$K^+K^-2(\pi^+\pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$		DESIG=222
Г79	$K^+K^-2(\pi^+\pi^-)\pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$		DESIG=240
Г80	$K^+K^*(892)^- + \text{c.c.}$	$(2.9 \pm 0.4) \times 10^{-5}$	S=1.2	DESIG=39
Г81	$2(K^+K^-)$	$(6.3 \pm 1.3) \times 10^{-5}$		DESIG=208
Г82	$2(K^+K^-)\pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$		DESIG=209
Г83	$K^+K^-\phi$	$(7.0 \pm 1.6) \times 10^{-5}$		DESIG=79
Г84	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$		DESIG=41
Г85	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$		DESIG=34
Г86	$\eta K^+K^-$ , no $\eta\phi$	$(3.49 \pm 0.17) \times 10^{-5}$		DESIG=207
Г87	$X(1750)\eta \rightarrow K^+K^-\eta$	$(4.8 \pm 2.8) \times 10^{-6}$		DESIG=324
Г88	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%	DESIG=42
Г89	$K_2^*(1430)^\pm K^\mp$	$(7.1^{+1.3}_{-0.9}) \times 10^{-5}$		DESIG=265
Г90	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$		DESIG=194
Г91	$\omega K^+K^-$	$(1.62 \pm 0.11) \times 10^{-4}$	S=1.1	DESIG=76
Г92	$\omega K_S^0 K_S^0$	$(7.0 \pm 0.5) \times 10^{-5}$		DESIG=330
Г93	$\omega K^*(892)^+ K^- + \text{c.c.}$	$(2.07 \pm 0.26) \times 10^{-4}$		DESIG=276
Г94	$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$(6.1 \pm 1.2) \times 10^{-5}$		DESIG=277
Г95	$\omega \bar{K}^*(892)^0 K^0$	$(1.68 \pm 0.30) \times 10^{-4}$		DESIG=278
Г96	$\omega \bar{K}_2^*(1430)^0 K^0$	$(5.8 \pm 2.2) \times 10^{-5}$		DESIG=279
Г97	$\omega X(1440) \rightarrow \omega K_S^0 K^-\pi^+ + \text{c.c.}$	$(1.6 \pm 0.4) \times 10^{-5}$		DESIG=282
Г98	$\omega X(1440) \rightarrow \omega K^+K^-\pi^0$	$(1.09 \pm 0.26) \times 10^{-5}$		DESIG=283
Г99	$\omega f_1(1285) \rightarrow \omega K_S^0 K^-\pi^+ + \text{c.c.}$	$(3.0 \pm 1.0) \times 10^{-6}$		DESIG=284
Г100	$\omega f_1(1285) \rightarrow \omega K^+K^-\pi^0$	$(1.2 \pm 0.7) \times 10^{-6}$		DESIG=285
Г101	$p\bar{p}$	$(2.94 \pm 0.08) \times 10^{-4}$		DESIG=27
Г102	$n\bar{n}$	$(3.06 \pm 0.15) \times 10^{-4}$		DESIG=309

$\Gamma_{103}$	$\rho\bar{\rho}\pi^0$	$(1.53 \pm 0.07) \times 10^{-4}$		DESIG=35
$\Gamma_{104}$	$N(940)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(6.4 \begin{smallmatrix} +1.8 \\ -1.3 \end{smallmatrix}) \times 10^{-5}$		DESIG=267
$\Gamma_{105}$	$N(1440)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(7.3 \begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}) \times 10^{-5}$	S=2.5	DESIG=261
$\Gamma_{106}$	$N(1520)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(6.4 \begin{smallmatrix} +2.3 \\ -1.8 \end{smallmatrix}) \times 10^{-6}$		DESIG=268
$\Gamma_{107}$	$N(1535)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(2.5 \pm 1.0) \times 10^{-5}$		DESIG=269
$\Gamma_{108}$	$N(1650)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(3.8 \begin{smallmatrix} +1.4 \\ -1.7 \end{smallmatrix}) \times 10^{-5}$		DESIG=270
$\Gamma_{109}$	$N(1720)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(1.79 \begin{smallmatrix} +0.26 \\ -0.70 \end{smallmatrix}) \times 10^{-5}$		DESIG=271
$\Gamma_{110}$	$N(2300)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(2.6 \begin{smallmatrix} +1.2 \\ -0.7 \end{smallmatrix}) \times 10^{-5}$		DESIG=272
$\Gamma_{111}$	$N(2570)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\pi^0$	$(2.13 \begin{smallmatrix} +0.40 \\ -0.31 \end{smallmatrix}) \times 10^{-5}$		DESIG=273
$\Gamma_{112}$	$\rho\bar{\rho}\pi^+\pi^-$	$(6.0 \pm 0.4) \times 10^{-4}$		DESIG=31
$\Gamma_{113}$	$\rho\bar{\rho}K^+K^-$	$(2.7 \pm 0.7) \times 10^{-5}$		DESIG=212
$\Gamma_{114}$	$\rho\bar{\rho}\eta$	$(6.0 \pm 0.4) \times 10^{-5}$		DESIG=200
$\Gamma_{115}$	$N(1535)\bar{\rho} + \text{c.c.} \rightarrow \rho\bar{\rho}\eta$	$(4.5 \begin{smallmatrix} +0.7 \\ -0.6 \end{smallmatrix}) \times 10^{-5}$		DESIG=264
$\Gamma_{116}$	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$		DESIG=211
$\Gamma_{117}$	$\rho\bar{\rho}\rho^0$	$(5.0 \pm 2.2) \times 10^{-5}$		DESIG=210
$\Gamma_{118}$	$\rho\bar{\rho}\omega$	$(6.9 \pm 2.1) \times 10^{-5}$		DESIG=77
$\Gamma_{119}$	$\rho\bar{\rho}\eta'$	$(1.10 \pm 0.13) \times 10^{-5}$		DESIG=317
$\Gamma_{120}$	$\rho\bar{\rho}\phi$	$(6.1 \pm 0.6) \times 10^{-6}$		DESIG=80
$\Gamma_{121}$	$\phi X(1835) \rightarrow \rho\bar{\rho}\phi$	$< 1.82 \times 10^{-7}$	CL=90%	DESIG=318
$\Gamma_{122}$	$\rho\bar{n}\pi^- \text{ or c.c.}$	$(2.48 \pm 0.17) \times 10^{-4}$		DESIG=227
$\Gamma_{123}$	$\rho\bar{n}\pi^-\pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$		DESIG=228
$\Gamma_{124}$	$\Lambda\bar{\Lambda}$	$(3.81 \pm 0.13) \times 10^{-4}$	S=1.4	DESIG=28
$\Gamma_{125}$	$\Lambda\bar{\Lambda}\pi^0$	$(1.4 \pm 0.7) \times 10^{-6}$		DESIG=238
$\Gamma_{126}$	$\Lambda\bar{\Lambda}\eta$	$(2.43 \pm 0.32) \times 10^{-5}$		DESIG=239
$\Gamma_{127}$	$\Lambda\bar{\Lambda}\omega(782)$	$(3.3 \pm 0.4) \times 10^{-5}$		DESIG=340
$\Gamma_{128}$	$\Lambda(1670)\bar{\Lambda} \rightarrow \Lambda\bar{\Lambda}\eta$	$(1.3 \pm 0.7) \times 10^{-5}$		DESIG=336
$\Gamma_{129}$	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(2.8 \pm 0.6) \times 10^{-4}$		DESIG=213
$\Gamma_{130}$	$\Lambda\bar{\rho}K^+$	$(1.00 \pm 0.14) \times 10^{-4}$		DESIG=214
$\Gamma_{131}$	$\Lambda\bar{\rho}K^*(892)^+ + \text{c.c.}$	$(6.3 \pm 0.7) \times 10^{-5}$		DESIG=321
$\Gamma_{132}$	$\Lambda\bar{\rho}K^+\pi^+\pi^-$	$(1.8 \pm 0.4) \times 10^{-4}$		DESIG=215
$\Gamma_{133}$	$\Lambda\bar{n}K_S^0 + \text{c.c.}$	$(8.1 \pm 1.8) \times 10^{-5}$		DESIG=237
$\Gamma_{134}$	$\Delta^{++}\bar{\Delta}^{--}$	$(1.28 \pm 0.35) \times 10^{-4}$		DESIG=70
$\Gamma_{135}$	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.40 \pm 0.13) \times 10^{-4}$		DESIG=280
$\Gamma_{136}$	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	$(1.54 \pm 0.14) \times 10^{-4}$		DESIG=281
$\Gamma_{137}$	$\Lambda\bar{\Sigma}^0 + \text{c.c.}$	$(1.6 \pm 0.7) \times 10^{-6}$		DESIG=326
$\Gamma_{138}$	$\Lambda\bar{\Sigma}^0$			DESIG=307
$\Gamma_{139}$	$\Sigma^0\bar{\rho}K^+ + \text{c.c.}$	$(1.67 \pm 0.18) \times 10^{-5}$		DESIG=274
$\Gamma_{140}$	$\Sigma^+\bar{\Sigma}^-$	$(2.43 \pm 0.10) \times 10^{-4}$	S=1.4	DESIG=223
$\Gamma_{141}$	$\Sigma^0\bar{\Sigma}^0$	$(2.35 \pm 0.09) \times 10^{-4}$	S=1.1	DESIG=71
$\Gamma_{142}$	$\Sigma^-\bar{\Sigma}^+$	$(2.82 \pm 0.09) \times 10^{-4}$		DESIG=335
$\Gamma_{143}$	$\Sigma^+\bar{\Sigma}^-\eta$	$(9.6 \pm 2.4) \times 10^{-6}$		DESIG=339
$\Gamma_{144}$	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$(8.5 \pm 0.7) \times 10^{-5}$		DESIG=72
$\Gamma_{145}$	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$(8.5 \pm 0.8) \times 10^{-5}$		DESIG=297
$\Gamma_{146}$	$\Sigma(1385)^0\bar{\Sigma}(1385)^0$	$(6.9 \pm 0.7) \times 10^{-5}$		DESIG=299
$\Gamma_{147}$	$\Xi^-\bar{\Xi}^+$	$(2.87 \pm 0.11) \times 10^{-4}$	S=1.1	DESIG=29
$\Gamma_{148}$	$\Xi^0\bar{\Xi}^0$	$(2.3 \pm 0.4) \times 10^{-4}$	S=4.2	DESIG=224
$\Gamma_{149}$	$\Xi(1530)^0\bar{\Xi}(1530)^0$	$(6.8 \pm 0.4) \times 10^{-5}$		DESIG=73
$\Gamma_{150}$	$\Lambda\bar{\Xi}^+K^- + \text{c.c.}$	$(3.9 \pm 0.4) \times 10^{-5}$		DESIG=293
$\Gamma_{151}$	$\Xi(1530)^-\bar{\Xi}(1530)^+$	$(1.15 \pm 0.07) \times 10^{-4}$		DESIG=322
$\Gamma_{152}$	$\Xi(1530)^-\bar{\Xi}^+$	$(7.0 \pm 1.2) \times 10^{-6}$		DESIG=323
$\Gamma_{153}$	$\Xi(1530)^0\bar{\Xi}^0$	$(5.3 \pm 0.5) \times 10^{-6}$		DESIG=331
$\Gamma_{154}$	$\Xi(1690)^-\bar{\Xi}^+ \rightarrow K^-\Lambda\bar{\Xi}^+ + \text{c.c.}$	$(5.2 \pm 1.6) \times 10^{-6}$		DESIG=294

Γ <sub>155</sub>	$\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ +$	$(1.20 \pm 0.32) \times 10^{-5}$		DESIG=295
Γ <sub>156</sub>	$\Sigma^0 \Xi^+ \bar{K}^- + c.c.$	$(3.7 \pm 0.4) \times 10^{-5}$		DESIG=296
Γ <sub>157</sub>	$\Omega^- \bar{\Omega}^+$	$(5.66 \pm 0.30) \times 10^{-5}$	S=1.3	DESIG=74
Γ <sub>158</sub>	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%	DESIG=229
Γ <sub>159</sub>	$h_c(1P) \pi^0$	$(7.4 \pm 0.5) \times 10^{-4}$		DESIG=254
Γ <sub>160</sub>	$\Lambda_c^+ \bar{p} e^+ e^- + c.c.$	$< 1.7 \times 10^{-6}$	CL=90%	DESIG=310
Γ <sub>161</sub>	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + c.c.$	[a] $< 8.8 \times 10^{-6}$	CL=90%	DESIG=195
Γ <sub>162</sub>	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[a] $< 1.0 \times 10^{-5}$	CL=90%	DESIG=196
Γ <sub>163</sub>	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[a] $< 7.0 \times 10^{-6}$	CL=90%	DESIG=197
Γ <sub>164</sub>	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[a] $< 2.6 \times 10^{-5}$	CL=90%	DESIG=198
Γ <sub>165</sub>	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[a] $< 6.0 \times 10^{-6}$	CL=90%	DESIG=199

## Radiative decays

Γ <sub>166</sub>	$\gamma \chi_{c0}(1P)$	$(9.79 \pm 0.20) \%$		NODE=M071;CLUMP=C DESIG=56
Γ <sub>167</sub>	$\gamma \chi_{c1}(1P)$	$(9.75 \pm 0.24) \%$		DESIG=58
Γ <sub>168</sub>	$\gamma \chi_{c2}(1P)$	$(9.52 \pm 0.20) \%$		DESIG=59
Γ <sub>169</sub>	$\gamma \eta_c(1S)$	$(3.4 \pm 0.5) \times 10^{-3}$	S=1.3	DESIG=61
Γ <sub>170</sub>	$\gamma \eta_c(2S)$	$(7 \pm 5) \times 10^{-4}$		DESIG=63
Γ <sub>171</sub>	$\gamma \pi_0$	$(1.04 \pm 0.22) \times 10^{-6}$	S=1.4	DESIG=52
Γ <sub>172</sub>	$\gamma 2(\pi^+ \pi^-)$	$(4.0 \pm 0.6) \times 10^{-4}$		DESIG=241
Γ <sub>173</sub>	$\gamma 3(\pi^+ \pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%	DESIG=249
Γ <sub>174</sub>	$\gamma \eta'(958)$	$(1.24 \pm 0.04) \times 10^{-4}$		DESIG=54
Γ <sub>175</sub>	$\gamma f_2(1270)$	$(2.73 \pm 0.29 \mp 0.25) \times 10^{-4}$	S=1.8	DESIG=82
Γ <sub>176</sub>	$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(3.1 \pm 1.7) \times 10^{-5}$		DESIG=286
Γ <sub>177</sub>	$\gamma f_0(1500)$	$(9.3 \pm 1.9) \times 10^{-5}$		DESIG=287
Γ <sub>178</sub>	$\gamma f_2'(1525)$	$(3.3 \pm 0.8) \times 10^{-5}$		DESIG=288
Γ <sub>179</sub>	$\gamma f_0(1710)$			DESIG=236
Γ <sub>180</sub>	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.5 \pm 0.6) \times 10^{-5}$		DESIG=83
Γ <sub>181</sub>	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(6.6 \pm 0.7) \times 10^{-5}$		DESIG=84
Γ <sub>182</sub>	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(4.8 \pm 1.0) \times 10^{-6}$		DESIG=289
Γ <sub>183</sub>	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(3.2 \pm 1.0) \times 10^{-6}$		DESIG=290
Γ <sub>184</sub>	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 5.8 \times 10^{-6}$	CL=90%	DESIG=291
Γ <sub>185</sub>	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 9.5 \times 10^{-6}$	CL=90%	DESIG=292
Γ <sub>186</sub>	$\gamma \eta$	$(9.2 \pm 1.8) \times 10^{-7}$		DESIG=53
Γ <sub>187</sub>	$\gamma \eta \pi^+ \pi^-$	$(8.7 \pm 2.1) \times 10^{-4}$		DESIG=230
Γ <sub>188</sub>	$\gamma \eta(1405)$			DESIG=231
Γ <sub>189</sub>	$\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$	$< 9 \times 10^{-5}$	CL=90%	DESIG=62
Γ <sub>190</sub>	$\gamma \eta(1405) \rightarrow \gamma \eta \pi^+ \pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$		DESIG=232
Γ <sub>191</sub>	$\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow$ $\gamma \pi^+ \pi^- \pi^0$	$< 5.0 \times 10^{-7}$	CL=90%	DESIG=308
Γ <sub>192</sub>	$\gamma \eta(1475)$			DESIG=233
Γ <sub>193</sub>	$\gamma \eta(1475) \rightarrow \gamma K \bar{K} \pi$	$< 1.4 \times 10^{-4}$	CL=90%	DESIG=234
Γ <sub>194</sub>	$\gamma \eta(1475) \rightarrow \gamma \eta \pi^+ \pi^-$	$< 8.8 \times 10^{-5}$	CL=90%	DESIG=235
Γ <sub>195</sub>	$\gamma K^{*0} K^+ \pi^- + c.c.$	$(3.7 \pm 0.9) \times 10^{-4}$		DESIG=242
Γ <sub>196</sub>	$\gamma K^{*0} \bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$		DESIG=243
Γ <sub>197</sub>	$\gamma K_S^0 K^+ \pi^- + c.c.$	$(2.6 \pm 0.5) \times 10^{-4}$		DESIG=244
Γ <sub>198</sub>	$\gamma K^+ K^- \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$		DESIG=245
Γ <sub>199</sub>	$\gamma K^+ K^- 2(\pi^+ \pi^-)$	$< 2.2 \times 10^{-4}$	CL=90%	DESIG=248
Γ <sub>200</sub>	$\gamma 2(K^+ K^-)$	$< 4 \times 10^{-5}$	CL=90%	DESIG=250
Γ <sub>201</sub>	$\gamma p \bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0	DESIG=246
Γ <sub>202</sub>	$\gamma f_2(1950) \rightarrow \gamma p \bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$		DESIG=257
Γ <sub>203</sub>	$\gamma f_2(2150) \rightarrow \gamma p \bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$		DESIG=258
Γ <sub>204</sub>	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(4.6 \pm 1.8 \mp 4.0) \times 10^{-6}$		DESIG=259
Γ <sub>205</sub>	$\gamma X \rightarrow \gamma p \bar{p}$	[b] $< 2 \times 10^{-6}$	CL=90%	DESIG=260
Γ <sub>206</sub>	$\gamma p \bar{p} \pi^+ \pi^-$	$(2.8 \pm 1.4) \times 10^{-5}$		DESIG=247
Γ <sub>207</sub>	$\gamma \gamma$	$< 1.5 \times 10^{-4}$	CL=90%	DESIG=51
Γ <sub>208</sub>	$\gamma \gamma J/\psi$	$(3.1 \pm 1.0 \mp 1.2) \times 10^{-4}$		DESIG=266
Γ <sub>209</sub>	$e^+ e^- \eta'$	$(1.90 \pm 0.26) \times 10^{-6}$		DESIG=311
Γ <sub>210</sub>	$e^+ e^- \eta_c(1S)$	$(3.8 \pm 0.4) \times 10^{-5}$		DESIG=338
Γ <sub>211</sub>	$e^+ e^- \chi_{c0}(1P)$	$(1.06 \pm 0.24) \times 10^{-3}$		DESIG=300
Γ <sub>212</sub>	$e^+ e^- \chi_{c1}(1P)$	$(8.5 \pm 0.6) \times 10^{-4}$		DESIG=301
Γ <sub>213</sub>	$e^+ e^- \chi_{c2}(1P)$	$(7.0 \pm 0.8) \times 10^{-4}$		DESIG=302



Weak decays			
$\Gamma_{214}$	$D^0 e^+ e^- + \text{c.c.}$	< 1.4	$\times 10^{-7}$ CL=90%
$\Gamma_{215}$	$\Lambda_c^+ \bar{\Sigma}^- + \text{c.c.}$	< 1.4	$\times 10^{-5}$ CL=90%

NODE=M071;CLUMP=E  
DESIG=306  
DESIG=337

Other decays			
$\Gamma_{216}$	invisible	< 1.6	% CL=90%

NODE=M071;CLUMP=D  
DESIG=275

[a]  $\Theta(1540)$  is a hypothetical pentaquark state of 1.54 GeV/c<sup>2</sup> mass and a width of less than 25 MeV/c<sup>2</sup>.

LINKAGE=THT

[b] For a narrow resonance in the range  $2.2 < M(X) < 2.8$  GeV.

LINKAGE=NMR

### CONSTRAINED FIT INFORMATION

A multiparticle fit to  $\chi_{c1}(1P)$ ,  $\chi_{c0}(1P)$ ,  $\chi_{c2}(1P)$ , and  $\psi(2S)$  with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 84 branching ratios uses 248 measurements to determine 49 parameters. The overall fit has a  $\chi^2 = 379.8$  for 199 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ .

$x_8$	3									
$x_9$	1	0								
$x_{12}$	29	11	2							
$x_{13}$	28	6	1	48						
$x_{14}$	13	4	1	36	15					
$x_{101}$	0	0	0	4	3	2				
$x_{166}$	1	0	0	2	1	1	0			
$x_{167}$	1	0	0	2	1	1	0	0		
$x_{168}$	1	0	0	3	1	1	0	0	0	
$\Gamma$	-81	-4	-1	-38	-34	-16	-7	-1	-1	-1
	$x_7$	$x_8$	$x_9$	$x_{12}$	$x_{13}$	$x_{14}$	$x_{101}$	$x_{166}$	$x_{167}$	$x_{168}$

### $\psi(2S)$ PARTIAL WIDTHS

NODE=M071225

#### $\Gamma(\text{hadrons})$

$\Gamma_1$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
258 ± 26	BAI	02B	BES2 $e^+ e^-$
224 ± 56	LUTH	75	MRK1 $e^+ e^-$

NODE=M071W3  
NODE=M071W3

#### $\Gamma(e^+ e^-)$

$\Gamma_7$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>2.33 ± 0.04 OUR FIT</b>			
<b>2.29 ± 0.06 OUR AVERAGE</b>			
2.23 ± 0.10 ± 0.02	<sup>1</sup> ABLIKIM	15V	BES3 $4.0\text{--}4.4 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
2.338 ± 0.037 ± 0.096	ABLIKIM	08B	BES2 $e^+ e^- \rightarrow \text{hadrons}$
2.330 ± 0.036 ± 0.110	ABLIKIM	06L	BES2 $e^+ e^- \rightarrow \text{hadrons}$
2.44 ± 0.21	<sup>2</sup> BAI	02B	BES2 $e^+ e^-$
2.14 ± 0.21	ALEXANDER	89	RVUE See $\Upsilon$ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.279 ± 0.015 ± 0.042	<sup>3</sup> ANASHIN	18	KEDR $e^+ e^-$
2.282 ± 0.015 ± 0.042	<sup>4</sup> ANASHIN	18	KEDR $e^+ e^-$
2.0 ± 0.3	BRANDELIK	79C	DASP $e^+ e^-$
2.1 ± 0.3	<sup>5</sup> LUTH	75	MRK1 $e^+ e^-$

NODE=M071W1  
NODE=M071W1

OCCUR=2

- <sup>1</sup> ABLIKIM 15V reports  $2.213 \pm 0.018 \pm 0.099$  keV from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+ e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$  assuming  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>2</sup> From a simultaneous fit to  $e^+ e^-$ ,  $\mu^+ \mu^-$ , and hadronic channel, assuming  $\Gamma_e = \Gamma_\mu = \Gamma_\tau/0.38847$ .
- <sup>3</sup> Combining  $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$  from ANASHIN 18 with  $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$  from ANASHIN 12 and assuming lepton universality.
- <sup>4</sup> From the sum of  $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$  from ANASHIN 12,  $\Gamma_{e^+e^-} \cdot B(e^+e^-)$  and  $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$  from ANASHIN 18, and  $\Gamma_{e^+e^-} \cdot B(\tau^+\tau^-)$  from ANASHIN 07.
- <sup>5</sup> From a simultaneous fit to  $e^+ e^-$ ,  $\mu^+ \mu^-$ , and hadronic channels assuming  $\Gamma(e^+ e^-) = \Gamma(\mu^+ \mu^-)$ .

NODE=M071W1;LINKAGE=A

NODE=M071W;LINKAGE=BB

NODE=M071W1;LINKAGE=B

NODE=M071W1;LINKAGE=C

NODE=M071W1;LINKAGE=F

 $\Gamma(\gamma\gamma)$  $\Gamma_{207}$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<43	90	BRANDELIK	79C DASP	$e^+ e^-$

NODE=M071W51  
NODE=M071W51 $\psi(2S) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$ 

NODE=M071230

This combination of a partial width with the partial width into  $e^+ e^-$  and with the total width is obtained from the integrated cross section into channel(i) in the  $e^+ e^-$  annihilation. We list only data that have not been used to determine the partial width  $\Gamma(i)$  or the branching ratio  $\Gamma(i)/\text{total}$ .

NODE=M071230

 $\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_1\Gamma_7/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>2.233±0.015±0.042</b>	<sup>1</sup> ANASHIN 12	KEDR	$e^+ e^- \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.2 ±0.4	ABRAMS 75	MRK1	$e^+ e^-$

NODE=M071G3  
NODE=M071G3

- <sup>1</sup> ANASHIN 12 reports the value  $2.233 \pm 0.015 \pm 0.037 \pm 0.020$  keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

NODE=M071G3;LINKAGE=AN

 $\Gamma(K_S^0 \text{ anything}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_6\Gamma_7/\Gamma$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.3738±0.0067±0.0200</b>	ABLIKIM 21s	BES3	$e^+ e^- \rightarrow K_S^0 \text{ anything}$

NODE=M071P30  
NODE=M071P30 $\Gamma(e^+ e^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_7\Gamma_7/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>21.2±0.7±1.2</b>	<sup>1</sup> ANASHIN 18	KEDR	$e^+ e^-$

NODE=M071P14  
NODE=M071P14

- <sup>1</sup> From the average of nine scans of the  $\psi(2S)$ .

NODE=M071P14;LINKAGE=A

 $\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_8\Gamma_7/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>19.3±0.3±0.5</b>	<sup>1</sup> ANASHIN 18	KEDR	$\psi(2S) \rightarrow \mu^+ \mu^-$

NODE=M071P13  
NODE=M071P13

- <sup>1</sup> From the average of nine scans of the  $\psi(2S)$ .

NODE=M071P13;LINKAGE=A

 $\Gamma(\tau^+ \tau^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_9\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9.0±2.6	79	<sup>1</sup> ANASHIN 07	KEDR	$e^+ e^- \rightarrow \psi(2S) \rightarrow \tau^+ \tau^-$

NODE=M071G9  
NODE=M071G9

- <sup>1</sup> Using  $\psi(2S)$  total width of  $337 \pm 13$  keV. Systematic errors not evaluated.

NODE=M071G9;LINKAGE=AN

 $\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_{12}\Gamma_7/\Gamma$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.808±0.013 OUR FIT</b>				
<b>0.837±0.025 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
0.837±0.028±0.005		<sup>1</sup> LEES 12E	BABR	10.6 $e^+ e^- \rightarrow 2\pi^+ 2\pi^- \gamma$
0.852±0.010±0.026	19.5k	ADAM 06	CLEO	3.773 $e^+ e^- \rightarrow \gamma\psi(2S)$
0.68 ±0.09		<sup>2</sup> BAI 98E	BES	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.88 ±0.08 ±0.03	256	<sup>3</sup> AUBERT 07AU	BABR	10.6 $e^+ e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
0.755±0.048±0.004	544	<sup>4</sup> AUBERT 05D	BABR	10.6 $e^+ e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$

NODE=M071G1  
NODE=M071G1


- <sup>1</sup> LEES 12E reports  $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$  keV which we divide by our best value  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>2</sup> The value of  $\Gamma(e^+e^-)$  quoted in BAI 98E is derived using  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$  and  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$ . Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$ .
- <sup>3</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$  keV which we divide by our best value  $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>4</sup> AUBERT 05D reports  $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$  keV which we divide by our best value  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.

NODE=M071G1;LINKAGE=LE

NODE=M071G1;LINKAGE=A

NODE=M071G1;LINKAGE=UB

NODE=M071G1;LINKAGE=AU



**$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{13}\Gamma_7/\Gamma$**

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.425±0.009 OUR FIT</b>				
<b>0.411±0.008±0.018</b>	3.6k	ADAM	06 CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.48 ± 0.09 ± 0.02	142	<sup>1</sup> LEES	18E BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^0\pi^0\gamma$
--------------------	-----	-------------------	----------	--

NODE=M071G6  
NODE=M071G6

NODE=M071G6;LINKAGE=A

- <sup>1</sup> LEES 18E reports  $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0101 \pm 0.0015 \pm 0.0011$  keV which we divide by our best value  $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.10 \pm 0.08) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

**$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{14}\Gamma_7/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>78.6± 1.6 OUR FIT</b>				
<b>87 ± 9 OUR AVERAGE</b>				
83 ± 25 ± 5	14	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ± 7	291 ± 24	ADAM	06 CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi\eta) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07$  eV.

NODE=M071G7  
NODE=M071G7

NODE=M071G7;LINKAGE=UB

**$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$   $\Gamma_{15}\Gamma_7/\Gamma$**

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;8</b>	90	<37	ADAM	06 CLEO	3.773 $e^+e^- \rightarrow \gamma\psi(2S)$

NODE=M071G8  
NODE=M071G8

$$\Gamma(2(\pi^+\pi^-\pi^0)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{22}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>29.7±2.2±1.8</b>	410	AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )π <sup>0</sup> γ

NODE=M071G01  
NODE=M071G01

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.4±1.8±1.2</b>	177	LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γ

NODE=M071P16  
NODE=M071P16

$$\Gamma(\pi^+\pi^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.3±2.3±0.5</b>	18	LEES	21C BABR	e <sup>+</sup> e <sup>-</sup> → γ <sub>ISR</sub> (π <sup>+</sup> π <sup>-</sup> 4π <sup>0</sup> )

NODE=M071P36  
NODE=M071P36

$$\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{26}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;6.2</b>	90	LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γ

NODE=M071P18  
NODE=M071P18

$$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{28}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.2±3.3±1.3</b>	43	AUBERT	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> )γ

NODE=M071G4  
NODE=M071G4

$$\Gamma(2(\pi^+\pi^-\pi^0)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>33±5±5</b>	14k	LEES	21 BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )3π <sup>0</sup> γ

NODE=M071P31  
NODE=M071P31

$$\Gamma(\eta 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.87±1.41±0.01</b>		16	<sup>1</sup> AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )ηγ

NODE=M071G03  
NODE=M071G03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<b>&lt;7</b>	90	14k	<sup>2</sup> LEES	21 BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )3π <sup>0</sup> γ
--------------	----	-----	-------------------	---------	---

<sup>1</sup>AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+\pi^-)) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.13 \pm 0.55 \pm 0.08$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G03;LINKAGE=UB

<sup>2</sup>LEES 21 reports  $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+\pi^-)) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] < 2.3$  eV which we divide by our best value  $B(\eta \rightarrow 3\pi^0) = 32.57 \times 10^{-2}$ .

NODE=M071G03;LINKAGE=A

$$\Gamma(\eta\pi^+\pi^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{34}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.85</b>	90	LEES	18E BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> π <sup>0</sup> ηγ

NODE=M071P15  
NODE=M071P15

$$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{35}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	<sup>1</sup> LEES	21C BABR	e <sup>+</sup> e <sup>-</sup> → γ <sub>ISR</sub> (π <sup>+</sup> π <sup>-</sup> 3π <sup>0</sup> γγ)

NODE=M071P40  
NODE=M071P40

<sup>1</sup>LEES 21C reports  $[\Gamma(\psi(2S) \rightarrow \eta\pi^+\pi^-\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 1.9$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M071P40;LINKAGE=A

$$\Gamma(\eta 2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	14k	<sup>1</sup> LEES	21 BABR	10.6 e <sup>+</sup> e <sup>-</sup> → 2(π <sup>+</sup> π <sup>-</sup> )3π <sup>0</sup> γ

NODE=M071P33  
NODE=M071P33

<sup>1</sup>LEES 21 reports  $[\Gamma(\psi(2S) \rightarrow \eta 2(\pi^+\pi^-\pi^0)) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 1.9$  eV which we divide by our best value  $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$ .

NODE=M071P33;LINKAGE=A

$$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{41}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.01±0.84±0.02</b>	37	<sup>1</sup> AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → ωπ <sup>+</sup> π <sup>-</sup> γ

NODE=M071G02  
NODE=M071G02

<sup>1</sup>AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = 2.69 \pm 0.73 \pm 0.16$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G02;LINKAGE=UB

$$\Gamma(\omega\pi^+\pi^-2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.2±5.6±0.1</b>	14k	<sup>1</sup> LEES	21	BABR 10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$

NODE=M071P32  
NODE=M071P32

<sup>1</sup> LEES 21 reports [ $\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-2\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$ ] =  $18 \pm 4 \pm 3$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P32;LINKAGE=A

$$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{45}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.58±0.82±0.02</b>	33	<sup>1</sup> LEES	18E	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

NODE=M071P17  
NODE=M071P17

<sup>1</sup> LEES 18E reports [ $\Gamma(\psi(2S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$ ] =  $2.3 \pm 0.7 \pm 0.2$  eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P17;LINKAGE=A

$$\Gamma(\omega3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{46}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.8</b>	90	<sup>1</sup> LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

NODE=M071P39  
NODE=M071P39

<sup>1</sup> LEES 21C reports [ $\Gamma(\psi(2S) \rightarrow \omega3\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$ ] < 1.6 eV which we divide by our best value  $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = 89.2 \times 10^{-2}$ .

NODE=M071P39;LINKAGE=A

$$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.55±0.19±0.01</b>	19	<sup>1</sup> LEES	12F	BABR 10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

NODE=M071G10  
NODE=M071G10

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.57±0.23±0.01	10	<sup>2</sup> AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
----------------	----	-----------------------------	------	--

<sup>1</sup> LEES 12F reports [ $\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\phi(1020) \rightarrow K^+K^-)$ ] =  $0.27 \pm 0.09 \pm 0.02$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G10;LINKAGE=A

<sup>2</sup> Superseded by LEES 12F. AUBERT, BE 06D reports [ $\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\phi(1020) \rightarrow K^+K^-)$ ] =  $0.28 \pm 0.11 \pm 0.02$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G10;LINKAGE=AU

$$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{52}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.346±0.129±0.004</b>	12	<sup>1</sup> LEES	12F	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

NODE=M071G13  
NODE=M071G13

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.346±0.168±0.004	6±3	<sup>2</sup> AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
-------------------	-----	---------------------	------	---

<sup>1</sup> LEES 12F reports [ $\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\phi(1020) \rightarrow K^+K^-)$ ] =  $0.17 \pm 0.06 \pm 0.02$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G13;LINKAGE=A

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports [ $\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\phi(1020) \rightarrow K^+K^-)$ ] =  $0.17 \pm 0.08 \pm 0.02$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G13;LINKAGE=AU

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{59}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.147±0.035±0.005	66	<sup>1</sup> LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.197±0.035±0.005	66	<sup>2</sup> LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.35±0.14±0.03	11	<sup>3</sup> LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

NODE=M071G06  
NODE=M071G06

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.147±0.035±0.005	66	<sup>1</sup> LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.197±0.035±0.005	66	<sup>2</sup> LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.35±0.14±0.03	11	<sup>3</sup> LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

OCCUR=2

<sup>1</sup>  $\sin\phi > 0$ .

<sup>2</sup>  $\sin\phi < 0$ .

<sup>3</sup> Interference with non-resonant  $K^+K^-$  production not taken into account.

NODE=M071G06;LINKAGE=A  
NODE=M071G06;LINKAGE=B  
NODE=M071G06;LINKAGE=BA

$\Gamma(K^+K^-\pi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{60}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.92±0.30±0.06</b>	133	LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.56±0.42±0.16	85	<sup>1</sup> AUBERT	07AK BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ
----------------	----	---------------------	-----------	--

<sup>1</sup> Superseded by LEES 12F.

NODE=M071G12  
NODE=M071G12

NODE=M071G12;LINKAGE=A

 $\Gamma(K_S^0K_L^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{64}\Gamma_7/\Gamma$ 

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.7</b>	90	8	LEES	17A BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>L</sub> <sup>0</sup> π <sup>0</sup> γ

NODE=M071G15  
NODE=M071G15

 $\Gamma(K^+K^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{65}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.60±0.31±0.03</b>	17	LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → π <sup>0</sup> π <sup>0</sup> K <sup>+</sup> K <sup>-</sup> γ

NODE=M071G08  
NODE=M071G08

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{66}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.4±1.3±0.3</b>	32	AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> π <sup>0</sup> γ

NODE=M071G04  
NODE=M071G04

 $\Gamma(K_S^0K_L^0\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{73}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.92±1.27±0.15</b>	14	LEES	17A BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>L</sub> <sup>0</sup> π <sup>0</sup> π <sup>0</sup> γ

NODE=M071G14  
NODE=M071G14

 $\Gamma(K_S^0K_L^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{74}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.14±1.08±0.16</b>	16	LEES	17A BABR	e <sup>+</sup> e <sup>-</sup> → K <sub>S</sub> <sup>0</sup> K <sub>L</sub> <sup>0</sup> ηγ

NODE=M071G16  
NODE=M071G16

 $\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{77}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.05±1.80±0.01</b>	7	<sup>1</sup> AUBERT	07AU BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> π <sup>+</sup> π <sup>-</sup> ηγ

<sup>1</sup> AUBERT 07AU reports [Γ(ψ(2S) → K<sup>+</sup>K<sup>-</sup>π<sup>+</sup>π<sup>-</sup>η) × Γ(ψ(2S) → e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>] × [B(η → 2γ)] = 1.2 ± 0.7 ± 0.1 eV which we divide by our best value B(η → 2γ) = (39.36 ± 0.18) × 10<sup>-2</sup>. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G05  
NODE=M071G05

NODE=M071G05;LINKAGE=UB

 $\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{78}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.4±2.1±0.3</b>	26	AUBERT	06D BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> 2(π <sup>+</sup> π <sup>-</sup> )γ

NODE=M071G5  
NODE=M071G5

 $\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{81}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.22±0.10±0.02</b>	13	LEES	12F BABR	10.6 e <sup>+</sup> e <sup>-</sup> → K <sup>+</sup> K <sup>-</sup> K <sup>+</sup> K <sup>-</sup> γ

NODE=M071G07  
NODE=M071G07

 $\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{101}\Gamma_7/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.686±0.019 OUR FIT</b>				
<b>0.63 ±0.05 OUR AVERAGE</b>				Error includes scale factor of 1.2.

0.67 ±0.12 ±0.02	43	<sup>1</sup> LEES	130 BABR	e <sup>+</sup> e <sup>-</sup> → p $\bar{p}$ γ
------------------	----	-------------------	----------	---

0.74 ±0.07 ±0.04	142	<sup>2</sup> LEES	13Y BABR	e <sup>+</sup> e <sup>-</sup> → p $\bar{p}$ γ
------------------	-----	-------------------	----------	---

0.579±0.038±0.036	2.7k	ANDREOTTI	07 E835	p $\bar{p}$ → e <sup>+</sup> e <sup>-</sup> , J/ψX
-------------------	------	-----------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.70 ±0.17 ±0.03	22	<sup>3</sup> AUBERT	06B BABR	e <sup>+</sup> e <sup>-</sup> → p $\bar{p}$ γ
------------------	----	---------------------	----------	---

<sup>1</sup> ISR photon reconstructed in the detector

<sup>2</sup> ISR photon undetected

<sup>3</sup> Superseded by LEES 130

NODE=M071G2  
NODE=M071G2

NODE=M071G2;LINKAGE=C  
NODE=M071G2;LINKAGE=B  
NODE=M071G2;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_{124}\Gamma_7/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>1.5±0.4±0.1</b>	AUBERT	07BD BABR	10.6 e <sup>+</sup> e <sup>-</sup> → Λ $\bar{\Lambda}$ γ

NODE=M071G11  
NODE=M071G11

$\psi(2S)$  BRANCHING RATIOS

NODE=M071235

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.9785 ± 0.0013 OUR AVERAGE</b>			
0.9779 ± 0.0015	<sup>1</sup> BAI	02B	BES2 $e^+ e^-$
0.981 ± 0.003	<sup>1</sup> LUTH	75	MRK1 $e^+ e^-$

NODE=M071R3  
NODE=M071R3<sup>1</sup> Includes cascade decay into  $J/\psi(1S)$ .

NODE=M071R;LINKAGE=P

 $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0173 ± 0.0014 OUR AVERAGE</b>			Error includes scale factor of 1.5.
0.0166 ± 0.0010	<sup>1,2</sup> SETH	04	RVUE $e^+ e^-$
0.0199 ± 0.0019	<sup>1</sup> BAI	02B	BES2 $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.029 ± 0.004	<sup>1</sup> LUTH	75	MRK1 $e^+ e^-$

NODE=M071R5  
NODE=M071R5<sup>1</sup> Included in  $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ .NODE=M071R;LINKAGE=Z  
NODE=M071R5;LINKAGE=SE<sup>2</sup> Using  $B(\psi(2S) \rightarrow \ell^+ \ell^-) = (0.73 \pm 0.04)\%$  from RPP-2002 and  $R = 2.28 \pm 0.04$  determined by a fit to data from BAI 00 and BAI 02c. $\Gamma(g g g)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.58 ± 1.62</b>	2.9 M	<sup>1</sup> LIBBY	09	CLEO $\psi(2S) \rightarrow \text{hadrons}$

NODE=M071S43  
NODE=M071S43<sup>1</sup> Calculated using  $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$  from LIBBY 09,  $B(\psi(2S) \rightarrow X J/\psi)$  relative and absolute branching fractions from MENDEZ 08,  $B(\psi(2S) \rightarrow \gamma \eta_c)$  from MITCHELL 09, and  $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$ ,  $B(\psi(2S) \rightarrow \gamma \chi_{cJ})$ , and  $B(\psi(2S) \rightarrow \ell^+ \ell^-)$  from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of  $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  LIBBY 09 measurement.

NODE=M071S43;LINKAGE=LI

 $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.025 ± 0.288</b>	200 k	<sup>1</sup> LIBBY	09	CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$

NODE=M071S44  
NODE=M071S44<sup>1</sup> Calculated using  $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$  from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of  $\Gamma(g g g)/\Gamma_{\text{total}}$  LIBBY 09 measurement.

NODE=M071S44;LINKAGE=LI

 $\Gamma(\gamma g g)/\Gamma(g g g)$  $\Gamma_4/\Gamma_3$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.7 ± 2.6 ± 1.6</b>	2.9 M	LIBBY	09	CLEO $\psi(2S) \rightarrow (\gamma +) \text{hadrons}$

NODE=M071S45  
NODE=M071S45 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.154 ± 0.015</b>	<sup>1</sup> MENDEZ	08	CLEO $e^+ e^- \rightarrow \psi(2S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.169 ± 0.026	<sup>2</sup> ADAM	05A	CLEO $e^+ e^- \rightarrow \psi(2S)$

NODE=M071S27  
NODE=M071S27<sup>1</sup> Uses  $B(\psi(2S) \rightarrow J/\psi X)$  from MENDEZ 08 and other branching fractions from PDG 07.

NODE=M071S27;LINKAGE=ME

<sup>2</sup> Uses  $B(J/\psi X)$  from ADAM 05A,  $B(\chi_{cJ} \gamma)$ ,  $B(\eta_c \gamma)$  from ATHAR 04 and  $B(\ell^+ \ell^-)$  from PDG 04. Superseded by MENDEZ 08.

NODE=M071S27;LINKAGE=AD

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>79.3 ± 1.7 OUR FIT</b>			

NODE=M071R1  
NODE=M071R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

88 ± 13 <sup>1</sup> FELDMAN 77 RVUE  $e^+ e^-$ <sup>1</sup> From an overall fit assuming equal partial widths for  $e^+ e^-$  and  $\mu^+ \mu^-$ . For a measurement of the ratio see the entry  $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$  below. Includes LUTH 75, HILGER 75, BURMESTER 77.

NODE=M071R;LINKAGE=L

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID
<b>80 ± 6 OUR FIT</b>	

NODE=M071R2  
NODE=M071R2

$\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$

$\Gamma_8/\Gamma_7$

NODE=M071R4  
NODE=M071R4

VALUE DOCUMENT ID TECN COMMENT

**1.00±0.08 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89±0.16 BOYARSKI 75C MRK1  $e^+ e^-$

$\Gamma(\tau^+ \tau^-)/\Gamma_{total}$

$\Gamma_9/\Gamma$

NODE=M071R75  
NODE=M071R75

VALUE (units  $10^{-4}$ ) DOCUMENT ID TECN COMMENT

**31 ±4 OUR FIT**

**30.8±2.1±3.8** <sup>1</sup> ABLIKIM 06W BES  $e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> Computed using PDG 02 value of  $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$  to estimate the total number of  $\psi(2S)$  events.

NODE=M071R75;LINKAGE=AB

———— DECAYS INTO  $J/\psi(1S)$  AND ANYTHING ————

NODE=M071305

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{total}$

$\Gamma_{10}/\Gamma = (\Gamma_{12} + \Gamma_{13} + \Gamma_{14} + 0.343\Gamma_{167} + 0.190\Gamma_{168})/\Gamma$

NODE=M071R10  
NODE=M071R10

VALUE EVTS DOCUMENT ID TECN COMMENT

**0.614 ±0.006 OUR FIT**

**0.55 ±0.07 OUR AVERAGE**

0.51 ±0.12 BRANDELIK 79C DASP  $e^+ e^- \rightarrow \mu^+ \mu^- X$

0.57 ±0.08 ABRAMS 75B MRK1  $e^+ e^- \rightarrow \mu^+ \mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.644 ±0.006 ±0.016 <sup>1</sup> ABLIKIM 21Z BES3  $e^+ e^- \rightarrow \ell^+ \ell^- X$

0.6254±0.0016±0.0155 1.1M <sup>2</sup> MENDEZ 08 CLEO  $\psi(2S) \rightarrow \ell^+ \ell^- X$

0.5950±0.0015±0.0190 151k ADAM 05A CLEO Repl. by MENDEZ 08

<sup>1</sup> From a fit to the  $e^+ e^- \rightarrow J/\psi X$  cross section between 3.645 and 3.891 GeV, with  $\Gamma(ee)$  and  $\Gamma$  fixed to the PDG 20 values of the cross particle fit which are correlated to "OUR FIT" value for  $B(\psi(2S) \rightarrow J/\psi X)$ .

NODE=M071R10;LINKAGE=A

<sup>2</sup> Not independent from other measurements of MENDEZ 08.

NODE=M071R10;LINKAGE=ME

$\Gamma(e^+ e^-)/\Gamma(J/\psi(1S)\text{anything})$

$\Gamma_7/\Gamma_{10}$

NODE=M071R72  
NODE=M071R72

VALUE (units  $10^{-2}$ ) EVTS DOCUMENT ID TECN COMMENT

**1.291±0.026 OUR FIT**

**1.28 ±0.04 OUR AVERAGE** Error includes scale factor of 1.6. See the ideogram below.

1.22 ±0.02 ±0.05 5097 ± 73 <sup>1</sup> ANDREOTTI 05 E835  $p\bar{p} \rightarrow \psi(2S) \rightarrow e^+ e^-$


1.28 ±0.03 ±0.02 <sup>1</sup> AMBROGIANI 00A E835  $p\bar{p} \rightarrow \psi(2S)$

1.44 ±0.08 ±0.02 <sup>1</sup> ARMSTRONG 97 E760  $p\bar{p} \rightarrow \psi(2S)$

<sup>1</sup> Using  $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ .

NODE=M071R;LINKAGE=7A

WEIGHTED AVERAGE  
1.28±0.04 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(e^+ e^-)/\Gamma(J/\psi(1S)\text{anything})$   
(units  $10^{-2}$ )

$\Gamma_7/\Gamma_{10}$



$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\text{anything})$  $\Gamma_8/\Gamma_{10}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0130±0.0010 OUR FIT</b>			
<b>0.014 ±0.003</b>	HILGER	75	SPEC $e^+e^-$

NODE=M071R74  
 NODE=M071R74

 $\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID
<b>0.2538±0.0032 OUR FIT</b>	

NODE=M071R18  
 NODE=M071R18

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.3468±0.0030 OUR FIT</b>				

NODE=M071R12  
 NODE=M071R12

**0.348 ±0.005 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.


0.3498±0.0002±0.0045	20M	ABLIKIM	13R	BES3	$\psi(2S) \rightarrow J/\psi\pi^+\pi^-$
0.3504±0.0007±0.0077	565k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.323 ±0.014		BAI	02B	BES2	$e^+e^-$
0.32 ±0.04		ABRAMS	75B	MRK1	$e^+e^- \rightarrow J/\psi\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3354±0.0014±0.0110 60k <sup>1</sup>ADAM 05A CLEO Repl. by MENDEZ 08

<sup>1</sup>Not independent from other values reported by ADAM 05A.

NODE=M071R;LINKAGE=AD

 $\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$  $\Gamma_7/\Gamma_{12}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0229±0.0005 OUR FIT</b>			
<b>0.0252±0.0028±0.0011</b>	<sup>1</sup> AUBERT	02B	BABR $e^+e^-$

NODE=M071R73  
 NODE=M071R73

<sup>1</sup>Using  $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$ .

NODE=M071R73;LINKAGE=7A

 $\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$  $\Gamma_8/\Gamma_{12}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0230±0.0017 OUR FIT</b>			
<b>0.0228±0.0018 OUR AVERAGE</b>			

NODE=M071R63  
 NODE=M071R63

0.0230±0.0020±0.0012	<sup>1</sup> AAIJ	16Y	LHCB $\Lambda_b^0 \rightarrow \psi(2S)X$
0.0216±0.0026±0.0014	<sup>2</sup> AUBERT	02B	BABR $e^+e^-$
0.0327±0.0077±0.0072	<sup>2</sup> GRIBUSHIN	96	FMPS $515\pi^-Be \rightarrow 2\mu X$

<sup>1</sup>Using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$ .

<sup>2</sup>Using  $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$ .

NODE=M071R63;LINKAGE=A  
 NODE=M071R;LINKAGE=Q2

 $\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$  $\Gamma_9/\Gamma_{12}$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.8 ±1.1 OUR FIT</b>			
<b>8.73±1.39±1.57</b>	BAI	02	BES $e^+e^-$

NODE=M071R76  
 NODE=M071R76

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$   $\Gamma_{12}/\Gamma_{10}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M071R70  
NODE=M071R70

**0.5645 ± 0.0026 OUR FIT**

**0.554 ± 0.008 OUR AVERAGE** Error includes scale factor of 1.3. See the ideogram below.

0.5604 ± 0.0009 ± 0.0062	565k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \pi^+ \pi^-$
0.525 ± 0.009 ± 0.022	4k	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.536 ± 0.007 ± 0.016	20k	<sup>1,2</sup> ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.496 ± 0.037		ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •


0.5637 ± 0.0027 ± 0.0046	60k	ADAM	05A	CLEO	Repl. by MENDEZ 08
--------------------------	-----	------	-----	------	--------------------

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.

<sup>2</sup> ABLIKIM 04B quotes  $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-)$ .

NODE=M071R;LINKAGE=AB  
NODE=M071R;LINKAGE=AL

WEIGHTED AVERAGE  
0.554 ± 0.008 (Error scaled by 1.3)


 $\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$   
 $\Gamma_{11}/\Gamma_{12} = (0.9761\Gamma_{13} + 0.719\Gamma_{14} + 0.343\Gamma_{167} + 0.190\Gamma_{168})/\Gamma_{12}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M071R11  
NODE=M071R11

**0.732 ± 0.008 OUR FIT**

**0.73 ± 0.09** TANENBAUM 76 MRK1  $e^+ e^-$

 $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M071R17  
NODE=M071R17

**0.1824 ± 0.0031 OUR FIT**

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.1769 ± 0.0008 ± 0.0053	61k	<sup>1</sup> MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.1652 ± 0.0014 ± 0.0058	13.4k	<sup>2</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

<sup>2</sup> Not independent from other values reported by ADAM 05A.

NODE=M071R17;LINKAGE=ME  
NODE=M071R17;LINKAGE=AD

 $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$   $\Gamma_{13}/\Gamma_{10}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M071R69  
NODE=M071R69

**0.2968 ± 0.0031 OUR FIT**

**0.320 ± 0.012 OUR AVERAGE**

0.300 ± 0.008 ± 0.022	1655 ± 44	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.328 ± 0.013 ± 0.008		AMBROGIANI	00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2829 ± 0.0012 ± 0.0056	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.2776 ± 0.0025 ± 0.0043	13.4k	ADAM	05A	CLEO	Repl. by MENDEZ 08

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{13}/\Gamma_{12}$

VALUE EVTS DOCUMENT ID TECN COMMENT

**0.526 ± 0.008 OUR FIT**

**0.513 ± 0.022 OUR AVERAGE** Error includes scale factor of 2.2.

0.5047 ± 0.0022 ± 0.0102	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.570 ± 0.009 ± 0.026	14k	<sup>1</sup> ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.4924 ± 0.0047 ± 0.0086	73k	<sup>2,3</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
0.571 ± 0.018 ± 0.044		<sup>4</sup> ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.53 ± 0.06		TANENBAUM	76	MRK1	$e^+ e^-$
0.64 ± 0.15		<sup>5</sup> HILGER	75	SPEC	$e^+ e^-$

- <sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.
- <sup>2</sup> Not independent from other values reported by ADAM 05A.
- <sup>3</sup> Using 13,217  $J/\psi\pi^0\pi^0$  and 60,010  $J/\psi\pi^+\pi^-$  events.
- <sup>4</sup> Not independent from other values reported by ANDREOTTI 05.
- <sup>5</sup> Ignoring the  $J/\psi(1S)\eta$  and  $J/\psi(1S)\gamma\gamma$  decays.

NODE=M071R14  
NODE=M071R14

NODE=M071R14;LINKAGE=AB  
NODE=M071R14;LINKAGE=AD  
NODE=M071R14;LINKAGE=AM  
NODE=M071R;LINKAGE=AN  
NODE=M071R;LINKAGE=I

$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$

$\Gamma_{14}/\Gamma$

VALUE (units  $10^{-3}$ ) EVTS DOCUMENT ID TECN COMMENT

**33.7 ± 0.5 OUR FIT**


**32.9 ± 1.7 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

33.75 ± 0.17 ± 0.86	68.2k	ABLIKIM	12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
29.8 ± 0.9 ± 2.3	5.7k	BAI	04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
25.5 ± 2.9	386	<sup>1</sup> OREGLIA	80	CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
45 ± 12	17	<sup>2</sup> BRANDELIK	79B	DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
42 ± 6	164	<sup>2</sup> BARTEL	78B	CNTR	$e^+ e^-$
34.3 ± 0.4 ± 0.9	18.4k	<sup>3</sup> MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
32.5 ± 0.6 ± 1.1	2.8k	<sup>4</sup> ADAM	05A	CLEO	Repl. by MENDEZ 08
43 ± 8	44	TANENBAUM	76	MRK1	$e^+ e^-$

- <sup>1</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .
- <sup>2</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .
- <sup>3</sup> Not independent from other measurements of MENDEZ 08.
- <sup>4</sup> Not independent from other values reported by ADAM 05A.

NODE=M071R15  
NODE=M071R15

NODE=M071R;LINKAGE=3Q  
NODE=M071R;LINKAGE=2Q  
NODE=M071R15;LINKAGE=ME  
NODE=M071R15;LINKAGE=AD



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$  $\Gamma_{14}/\Gamma_{10}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M071R68  
 NODE=M071R68

**0.0549 ± 0.0008 OUR FIT****0.058 ± 0.007 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

0.050 ± 0.006 ± 0.003	298 ± 20	ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$
0.072 ± 0.009		AMBROGIANI 00A	E835	$\rho\bar{\rho} \rightarrow \psi(2S)$
0.061 ± 0.015		ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$


OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.0549 ± 0.0006 ± 0.0009	18.4k	<sup>1</sup> MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.0546 ± 0.0010 ± 0.0007	2.8k	ADAM 05A	CLEO	Repl. by MENDEZ 08

<sup>1</sup> Not independent from other measurements of MENDEZ 08.

NODE=M071R68;LINKAGE=ME

 $\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$  $\Gamma_{14}/\Gamma_{12}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M071R71  
 NODE=M071R71

**0.0972 ± 0.0014 OUR FIT****0.0979 ± 0.0018 OUR AVERAGE**

0.0979 ± 0.0010 ± 0.0015	18.4k	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.098 ± 0.005 ± 0.010	2k	<sup>1</sup> ABLIKIM 04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ± 0.021		<sup>2</sup> HIMEL 80	MRK2	$e^+ e^- \rightarrow \psi(2S) X$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.0968 ± 0.0019 ± 0.0013	2.8k	<sup>3</sup> ADAM 05A	CLEO	Repl. by MENDEZ 08
0.095 ± 0.007 ± 0.007		<sup>4</sup> ANDREOTTI 05	E835	$\psi(2S) \rightarrow J/\psi X$

<sup>1</sup> From a fit to the  $J/\psi$  recoil mass spectra.<sup>2</sup> The value for  $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$  reported in HIMEL 80 is derived using  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$  and  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$ . Calculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$ .<sup>3</sup> Not independent from other values reported by ADAM 05A.<sup>4</sup> Not independent from other values reported by ANDREOTTI 05.

NODE=M071R71;LINKAGE=AB  
 NODE=M071R;LINKAGE=8H

NODE=M071R71;LINKAGE=AD  
 NODE=M071R71;LINKAGE=AN

 $\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M071R16  
 NODE=M071R16

**12.68 ± 0.32 OUR AVERAGE**

12.6 ± 0.2 ± 0.3	4.1k	ABLIKIM 12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
13.3 ± 0.8 ± 0.3	530	MENDEZ 08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
14.3 ± 1.4 ± 1.2	280	BAI 04I	BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
14 ± 6	7	HIMEL 80	MRK2	$e^+ e^-$
9 ± 2 ± 1	23	<sup>1</sup> OREGLIA 80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

13 ± 1 ± 1	88	ADAM 05A	CLEO	Repl. by MENDEZ 08
------------	----	----------	------	--------------------

<sup>1</sup> Recalculated by us using  $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$ .

NODE=M071R16;LINKAGE=3Q

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\text{anything})$   
 $\Gamma_{15}/\Gamma_{10} = \Gamma_{15}/(\Gamma_{12}+\Gamma_{13}+\Gamma_{14}+0.343\Gamma_{167}+0.190\Gamma_{168})$

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
0.213±0.012±0.003	527	<sup>1</sup> MENDEZ	08 CLEO	e <sup>+</sup> e <sup>-</sup> → J/ψγγ
0.22 ±0.02 ±0.01		<sup>2</sup> ADAM	05A CLEO	e <sup>+</sup> e <sup>-</sup> → ψ(2S) → J/ψγγ

<sup>1</sup> Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.  
<sup>2</sup> Not independent from other values reported by ADAM 05A.

NODE=M071S25  
 NODE=M071S25

NODE=M071S25;LINKAGE=ME  
 NODE=M071S25;LINKAGE=AD

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_{15}/\Gamma_{12}$

VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
0.380±0.022±0.005	527	<sup>1</sup> MENDEZ	08 CLEO	e <sup>+</sup> e <sup>-</sup> → J/ψγγ
0.39 ±0.04 ±0.01		<sup>2</sup> ADAM	05A CLEO	e <sup>+</sup> e <sup>-</sup> → ψ(2S) → J/ψγγ

<sup>1</sup> Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.  
<sup>2</sup> Not independent from other values reported by ADAM 05A.

NODE=M071S26  
 NODE=M071S26

NODE=M071S26;LINKAGE=ME  
 NODE=M071S26;LINKAGE=AD

———— HADRONIC DECAYS ————

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$

VALUE (units 10 <sup>-5</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.78±0.26 OUR AVERAGE</b>					
0.76±0.25±0.06		30	<sup>1</sup> METREVELI	12	ψ(2S) → π <sup>+</sup> π <sup>-</sup>
8 ±5			BRANDELIK	79C DASP	e <sup>+</sup> e <sup>-</sup>
<2.1	90		DOBBS	06A CLEO	e <sup>+</sup> e <sup>-</sup> → ψ(2S)
<5	90		FELDMAN	77 MRK1	e <sup>+</sup> e <sup>-</sup>

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using ψ(3770) → π<sup>+</sup>π<sup>-</sup> for continuum subtraction.

NODE=M071310

NODE=M071R20  
 NODE=M071R20

NODE=M071R20;LINKAGE=ME

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.01±0.17 OUR AVERAGE</b> Error includes scale factor of 1.7. See the ideogram below.				
2.14±0.03 <sup>+0.12</sup> <sub>-0.11</sub>	7k	<sup>1</sup> ABLIKIM	12H BES3	e <sup>+</sup> e <sup>-</sup> → ψ(2S)
1.81±0.18±0.19	260 ± 19	<sup>2</sup> ABLIKIM	05J BES2	e <sup>+</sup> e <sup>-</sup> → ψ(2S)
1.88 <sup>+0.16</sup> <sub>-0.15</sub> ±0.28	194	ADAM	05 CLEO	e <sup>+</sup> e <sup>-</sup> → ψ(2S)
0.85±0.46	4	FRANKLIN	83 MRK2	e <sup>+</sup> e <sup>-</sup> → hadrons


<sup>1</sup> From ψ(2S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup> events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of ψ(2S) events.

<sup>2</sup> From a PW analysis of ψ(2S) → π<sup>+</sup>π<sup>-</sup>π<sup>0</sup>.

NODE=M071R36  
 NODE=M071R36

NODE=M071R36;LINKAGE=AB

NODE=M071R;LINKAGE=AK



$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.32 ± 0.12 OUR AVERAGE</b>					Error includes scale factor of 1.8.
0.51 ± 0.07 ± 0.11			<sup>1</sup> ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
0.24 <sup>+0.08</sup> <sub>-0.07</sub> ± 0.02		22	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83 MRK2	$e^+e^-$
<10	90		BARTEL	76 CNTR	$e^+e^-$
<10	90		<sup>2</sup> ABRAMS	75 MRK1	$e^+e^-$

<sup>1</sup> From a PW analysis of  $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ .

<sup>2</sup> Final state  $\rho^0\pi^0$ .

NODE=M071R26  
NODE=M071R26

NODE=M071R26;LINKAGE=AK  
NODE=M071R;LINKAGE=N

 $\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.94 ± 0.25<sup>+1.15</sup><sub>-0.34</sub></b>	<sup>1</sup> ABLIKIM	05J BES2	$\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

<sup>1</sup> From a PW analysis of  $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ .

NODE=M071R57  
NODE=M071R57

NODE=M071R57;LINKAGE=AK

 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4 ± 0.6 OUR AVERAGE</b>				Error includes scale factor of 2.2.
2.2 ± 0.2 ± 0.2	308	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$
4.5 ± 1.0		TANENBAUM	78 MRK1	$e^+e^-$

NODE=M071R27  
NODE=M071R27

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2 ± 0.6 OUR AVERAGE</b>				Error includes scale factor of 1.4.
2.0 ± 0.2 ± 0.4	285.5	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$
4.2 ± 1.5		TANENBAUM	78 MRK1	$e^+e^-$

NODE=M071R33  
NODE=M071R33


 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>29 ± 10 OUR AVERAGE</b>				Error includes scale factor of 4.7. See the ideogram below.
24.9 ± 0.7 ± 3.6	2173	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
127 ± 12 ± 2	410	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$
26.1 ± 0.7 ± 3.0	1703	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
30 ± 8	42	FRANKLIN	83 MRK2	$e^+e^-$

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R22  
NODE=M071R22

NODE=M071R22;LINKAGE=UB



**$\Gamma(\rho a_2(1320))/\Gamma_{\text{total}}$**

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{23}/\Gamma$
<b>2.55±0.73±0.47</b>		112 ± 31	BAI	04c	BES2 $\psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
<2.3		90	BAI	98J	BES $e^+e^-$	

NODE=M071R65  
NODE=M071R65

**$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{27}/\Gamma$
<b>3.5 ±2.0 OUR AVERAGE</b>		Error includes scale factor of 2.8.			
5.45±0.42±0.87	671	ABLIKIM	05H	BES2 $e^+e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+\pi^-)$	
1.5 ±1.0		<sup>1</sup> TANENBAUM	78	MRK1 $e^+e^-$	
<sup>1</sup> Assuming entirely strong decay.					

NODE=M071R32  
NODE=M071R32

NODE=M071R32;LINKAGE=K

**$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{29}/\Gamma$
<b>35±16</b>	6	FRANKLIN	83	MRK2 $e^+e^- \rightarrow \text{hadrons}$	

NODE=M071R37  
NODE=M071R37

**$\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$**

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{31}/\Gamma$
<b>&lt;1.6</b>	90	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-\pi^0)$	

NODE=M071S06  
NODE=M071S06

**$\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$**

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{32}/\Gamma$
<b>9.5±0.7±1.5</b>		<sup>1</sup> BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
10.3±0.8±1.4	201.7	<sup>2</sup> BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$	OCCUR=2
8.1±1.4±1.6	50.0	<sup>2</sup> BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$	OCCUR=3

NODE=M071S07  
NODE=M071S07

<sup>1</sup> Average of  $\eta \rightarrow \gamma\gamma$  and  $\eta \rightarrow 3\pi$ .

<sup>2</sup> Not independent from other values reported by BRIERE 05.

NODE=M071S07;LINKAGE=BR  
NODE=M071S07;LINKAGE=BI

**$\Gamma(\rho\eta)/\Gamma_{\text{total}}$**

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{37}/\Gamma$
<b>2.2 ±0.6 OUR AVERAGE</b>		Error includes scale factor of 1.1.			
3.0 $\begin{smallmatrix} +1.1 \\ -0.9 \end{smallmatrix} \pm 0.2$	18	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$	
1.78 $\begin{smallmatrix} +0.67 \\ -0.62 \end{smallmatrix} \pm 0.17$	13	ABLIKIM	04L	BES $e^+e^- \rightarrow \psi(2S)$	

NODE=M071R94  
NODE=M071R94

$\Gamma(\eta' \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$					$\Gamma_{38}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>4.5 \pm 1.6 \pm 1.3</math></b>	12.8	BRIERE	05	CLEO $e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$	

NODE=M071S08  
NODE=M071S08

$\Gamma(\eta' \rho)/\Gamma_{\text{total}}$					$\Gamma_{39}/\Gamma$
VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>1.87 \begin{smallmatrix} +1.64 \\ -1.11 \end{smallmatrix} \pm 0.33</math></b>	2	ABLIKIM	04L	BES $e^+ e^- \rightarrow \psi(2S)$	

NODE=M071R93  
NODE=M071R93

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.02 \pm 0.11 \pm 0.24$	143	<sup>1</sup> ABLIKIM	17AK	BES3	$e^+ e^- \rightarrow \psi(2S)$
$0.569 \pm 0.128 \pm 0.236$	80	<sup>2</sup> ABLIKIM	17AK	BES3	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> Destructive-interference solution of a partial wave analysis of the decay  $\psi(2S) \rightarrow \pi^+ \pi^- \eta'$ .  
<sup>2</sup> Constructive-interference solution of a partial wave analysis of the decay  $\psi(2S) \rightarrow \pi^+ \pi^- \eta'$ .

OCCUR=2  
NODE=M071R93;LINKAGE=A  
NODE=M071R93;LINKAGE=B

$\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$					$\Gamma_{40}/\Gamma$
VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
<b><math>2.1 \pm 0.6</math> OUR AVERAGE</b>					
$2.5 \begin{smallmatrix} +1.2 \\ -1.0 \end{smallmatrix} \pm 0.2$	14	ADAM	05	CLEO $e^+ e^- \rightarrow \psi(2S)$	
$1.87 \begin{smallmatrix} +0.68 \\ -0.62 \end{smallmatrix} \pm 0.28$	14	ABLIKIM	04L	BES $e^+ e^- \rightarrow \psi(2S)$	

NODE=M071R92  
NODE=M071R92


$\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$					$\Gamma_{41}/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M071R77  
NODE=M071R77

<b><math>7.3 \pm 1.2</math> OUR AVERAGE</b>		Error includes scale factor of 2.1. See the ideogram below.			
$8.4 \pm 0.5 \pm 1.2$	386	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$
$12.2 \pm 2.2 \pm 0.7$	37	<sup>1</sup> AUBERT	07AU	BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$
$8.2 \pm 0.5 \pm 0.7$	391	BRIERE	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
$4.8 \pm 0.6 \pm 0.7$	$100 \pm 22$	<sup>2</sup> BAI	03B	BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

<sup>1</sup> AUBERT 07AU quotes  $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega \pi^+ \pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$  eV.  
<sup>2</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

NODE=M071R77;LINKAGE=UB  
NODE=M071R77;LINKAGE=B3





$\Gamma(b_1^\pm \pi^\mp)/\Gamma_{\text{total}}$  $\Gamma_{43}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.0 ± 0.6 OUR AVERAGE</b>				Error includes scale factor of 1.1.
5.1 ± 0.6 ± 0.8	202	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$
4.18 <sup>+0.43</sup> <sub>-0.42</sub> ± 0.92	170	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.2 ± 0.6 ± 0.5	61 ± 11	1,2 BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
5.2 ± 0.8 ± 1.0		<sup>1</sup> BAI	99c BES	Repl. by BAI 03B

NODE=M071R40  
 NODE=M071R40

<sup>1</sup> Assuming  $B(b_1 \rightarrow \omega \pi) = 1$ .

<sup>2</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

NODE=M071R;LINKAGE=M1  
 NODE=M071R40;LINKAGE=B3

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$  $\Gamma_{44}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2 ± 0.4 OUR AVERAGE</b>					
2.3 ± 0.5 ± 0.4		57	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$
2.05 ± 0.41 ± 0.38		62 ± 12	BAI	04c BES2	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
<1.5	90		<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+ \pi^-) \pi^0$
<1.7	90		BAI	98J BES	Repl. by BAI 03B

NODE=M071R64  
 NODE=M071R64

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

NODE=M071R64;LINKAGE=B3

 $\Gamma(b_1^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{47}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.35<sup>+0.47</sup><sub>-0.42</sub> ± 0.40</b>	45	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R21  
 NODE=M071R21

 $\Gamma(\omega \eta)/\Gamma_{\text{total}}$  $\Gamma_{48}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.1</b>	90	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
<3.1	90	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R95  
 NODE=M071R95

 $\Gamma(\omega \eta')/\Gamma_{\text{total}}$  $\Gamma_{49}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.2<sup>+2.4</sup><sub>-2.0</sub> ± 0.7</b>	4	<sup>1</sup> ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R91  
 NODE=M071R91

<sup>1</sup> Calculated combining  $\eta' \rightarrow \gamma \rho$  and  $\eta \pi^+ \pi^-$  channels.

NODE=M071R91;LINKAGE=AI

 $\Gamma(\phi \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{50}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.04</b>	90	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
<0.7	90	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
<0.4	90	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R96  
 NODE=M071R96

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{51}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.18 ± 0.26 OUR AVERAGE</b>				Error includes scale factor of 1.5. See the ideogram below.
2.3 ± 0.8 ± 0.1	19 ± 6	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.9 ± 0.2 ± 0.1	47.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
1.5 ± 0.2 ± 0.2	51.5 ± 8.3	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
2.45 ± 0.96 ± 0.04	10 ± 4	<sup>2,3</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M071R80  
 NODE=M071R80


<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(\psi(2S) \rightarrow \phi \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Using  $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$ .

NODE=M071R80;LINKAGE=B3  
 NODE=M071R80;LINKAGE=BE

NODE=M071R80;LINKAGE=UB



### $\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$

$\Gamma_{52}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.75±0.33 OUR AVERAGE</b>				Error includes scale factor of 1.6.
1.5 ±0.5 ±0.1	12 ± 4	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.6 ±0.2 ±0.1	18.4 ± 6.4	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.46±0.71±0.02	6 ± 3	<sup>2,3</sup> AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M071R83  
NODE=M071R83

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Using  $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$ .

NODE=M071R83;LINKAGE=B3  
NODE=M071R83;LINKAGE=BE

NODE=M071R83;LINKAGE=UB

### $\Gamma(\phi \eta) / \Gamma_{\text{total}}$

$\Gamma_{53}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.10±0.31 OUR AVERAGE</b>				
3.14±0.23±0.23	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.0 $^{+1.5}_{-1.1}$ ±0.4	6	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.3 ±1.1 ±0.5	17	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R89  
NODE=M071R89

### $\Gamma(\eta \phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$

$\Gamma_{54}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.2 × 10<sup>-6</sup></b>	90	ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta \phi f_0(980)$

NODE=M071P19  
NODE=M071P19

### $\Gamma(\phi \eta') / \Gamma_{\text{total}}$

$\Gamma_{55}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.54±0.20 OUR AVERAGE</b>				
1.51±0.16±0.12	201	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
3.1 ±1.4 ±0.7	8	<sup>1</sup> ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R90  
NODE=M071R90

<sup>1</sup> Calculated combining  $\eta' \rightarrow \gamma \rho$  and  $\eta \pi^+ \pi^-$  channels.

NODE=M071R;LINKAGE=AI

### $\Gamma(\phi f_1(1285)) / \Gamma_{\text{total}}$

$\Gamma_{56}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.0±0.4±1.3</b>	234	<sup>1</sup> ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071P22  
NODE=M071P22

<sup>1</sup> ABLIKIM 19BA reports  $[\Gamma(\psi(2S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.03 \pm 0.10 \pm 0.09) \times 10^{-5}$  which we divide by our best value  $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P22;LINKAGE=B

$\Gamma(\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.46±1.37±0.92</b>	195	ABLIKIM	19BA BES3	$e^+e^- \rightarrow \psi(2S)$

NODE=M071P24  
 NODE=M071P24

 $\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.44±0.12±0.11</b>	20 ± 6	BAI	04C		$\psi(2S) \rightarrow 2(K^+K^-)$

NODE=M071R67  
 NODE=M071R67

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.45	90	BAI	98J BES		$e^+e^- \rightarrow 2(K^+K^-)$
-------	----	-----	---------	--	--------------------------------

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.48±0.23±0.39</b>	1.3k	<sup>1</sup> METREVELI	12		$\psi(2S) \rightarrow K^+K^-$

NODE=M071R23  
 NODE=M071R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.2 ±1.5 ±0.2	66	<sup>2,3</sup> LEES	15J BABR		$e^+e^- \rightarrow K^+K^-\gamma$
8.3 ±1.5 ±0.2	66	<sup>3,4</sup> LEES	15J BABR		$e^+e^- \rightarrow K^+K^-\gamma$
6.3 ±0.6 ±0.3		<sup>5</sup> DOBBS	06A CLEO		$e^+e^-$
10 ±7		<sup>5</sup> BRANDELIK	79C DASP		$e^+e^-$
< 5	90	FELDMAN	77 MRK1		$e^+e^-$

OCCUR=2

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup>  $\sin\phi > 0$ .

<sup>3</sup> Using  $\Gamma(\psi(2S) \rightarrow e^+e^-) = (2.37 \pm 0.04)$  keV.

<sup>4</sup>  $\sin\phi < 0$ .

<sup>5</sup> Interference with non-resonant  $K^+K^-$  production not taken into account.

NODE=M071R23;LINKAGE=ME  
 NODE=M071R23;LINKAGE=A  
 NODE=M071R23;LINKAGE=B  
 NODE=M071R23;LINKAGE=C  
 NODE=M071R23;LINKAGE=BA

 $\Gamma(K^+K^-\pi^+)/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3±0.5 OUR AVERAGE</b>				

NODE=M071R24  
 NODE=M071R24

8.1±1.3±0.3	133	LEES	12F BABR		$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
7.1±0.3±0.4	817.2	BRIERE	05 CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
16 ±4		<sup>1</sup> TANENBAUM	78 MRK1		$e^+e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.0±1.9±0.2	85	<sup>2</sup> AUBERT	07AK BABR		$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
--------------	----	---------------------	-----------	--	--

<sup>1</sup> Assuming entirely strong decay.

<sup>2</sup> Superseded by LEES 12F. AUBERT 07AK reports  $[\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R24;LINKAGE=K  
 NODE=M071R24;LINKAGE=BE

 $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.07±0.16±0.26</b>	0.9k	ABLIKIM	12L BES3		$e^+e^- \rightarrow \psi(2S)$

NODE=M071R38  
 NODE=M071R38

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8.9	90	1	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$
------	----	---	----------	---------	-------------------------------------

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.046</b>	<sup>1</sup> BAI	04D BES	$e^+e^-$

NODE=M071R88  
 NODE=M071R88

<sup>1</sup> Forbidden by CP.

NODE=M071R;LINKAGE=BA

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.34±0.33 OUR AVERAGE</b>				

NODE=M071R87  
 NODE=M071R87

5.28±0.25±0.34	478 ± 23	<sup>1</sup> METREVELI	12		$\psi(2S) \rightarrow K_S^0 K_L^0$
5.8 ±0.8 ±0.4		DOBBS	06A CLEO		$e^+e^-$
5.24±0.47±0.48	156 ± 14	<sup>2</sup> BAI	04B BES2		$\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$

<sup>1</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Using  $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$ .

NODE=M071R87;LINKAGE=ME  
 NODE=M071R;LINKAGE=KZ

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.6±0.9 OUR AVERAGE</b>				
18.9±5.7±0.3	32	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$
11.7±1.0±1.5	597	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
12.7±0.5±1.0	711.6	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S10  
NODE=M071S10

<sup>1</sup> AUBERT 07AU reports  $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071S10;LINKAGE=UB

 $\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.9±2.0±0.9</b>	19	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S20  
NODE=M071S20 $\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6±1.3±1.8</b>	238	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S21  
NODE=M071S21 $\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.6±2.2±1.7</b>	133	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S22  
NODE=M071S22 $\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3±2.2±1.4</b>	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S23  
NODE=M071S23 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1±1.3±1.2</b>	125	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S24  
NODE=M071S24 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{72}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.20±0.25±0.37</b>	83 ± 9	ABLIKIM	05o BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R49  
NODE=M071R49 $\Gamma(K^+ K^- \rho^0)/\Gamma_{\text{total}}$  $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.2±0.2±0.4</b>	223.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M071S09  
NODE=M071S09 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$  $\Gamma_{76}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.86±0.32±0.43</b>	93 ± 16	BAI	04C		$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
<1.2	90	BAI	98J BES		$e^+e^-$

NODE=M071R66  
NODE=M071R66 $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$  $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.3±0.7±0.1</b>	7	<sup>1</sup> AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$
		<sup>1</sup> AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^- \eta)) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1$ eV.		

NODE=M071S39  
NODE=M071S39

NODE=M071S39;LINKAGE=UB

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{79}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.0±2.5±1.8</b>	65	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R09  
NODE=M071R09

$\Gamma(K^+ K^*(892)^- + c.c.)/\Gamma_{\text{total}}$   $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.9 ± 0.4</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.			
3.18 ± 0.30 <sup>+0.26</sup> <sub>-0.31</sub>		0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.9 <sup>+1.3</sup> <sub>-1.7</sub> ± 0.4		9.6 ± 4.2	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
1.3 <sup>+1.0</sup> <sub>-0.7</sub> ± 0.3		7	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.4	90		FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$
------	----	--	----------	---------	--------------------------------------

NODE=M071R39  
NODE=M071R39

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$   $\Gamma_{81}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.63 ± 0.13</b>	<b>OUR AVERAGE</b>			
0.9 ± 0.4 ± 0.1	13	LEES	12F BABR	$10.6 e^+ e^- \rightarrow 2(K^+ K^-)\gamma$
0.6 ± 0.1 ± 0.1	59.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

NODE=M071S12  
NODE=M071S12

 $\Gamma(2(K^+ K^-)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{82}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.1 ± 0.2 ± 0.2</b>	44.7	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)\pi^0$

NODE=M071S13  
NODE=M071S13

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$   $\Gamma_{83}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.70 ± 0.16</b>	<b>OUR AVERAGE</b>			
0.8 ± 0.2 ± 0.1	36.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ± 0.2 ± 0.1	16.1 ± 5.0	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow 2(K^+ K^-)$

<sup>1</sup>Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

NODE=M071R81  
NODE=M071R81

NODE=M071R81;LINKAGE=B3

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{84}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>10.0 ± 1.8 ± 2.1</b>	<sup>1</sup> BAI	99C BES	$e^+ e^-$

<sup>1</sup>Assuming  $B(K_1(1270) \rightarrow K\rho) = 0.42 \pm 0.06$

NODE=M071R41  
NODE=M071R41

NODE=M071R;LINKAGE=M2

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + c.c.)/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.7 ± 2.5</b>	TANENBAUM 78	MRK1	$e^+ e^-$

NODE=M071R34  
NODE=M071R34

 $\Gamma(\eta K^+ K^-, \text{no } \eta\phi)/\Gamma_{\text{total}}$   $\Gamma_{86}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.49 ± 0.09 ± 0.15</b>		1.8k	<sup>1</sup> ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$
3.08 ± 0.29 ± 0.25		0.3k	<sup>1,2</sup> ABLIKIM	12L BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$
<13	90		BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>Excluding  $\eta\phi$ .

<sup>2</sup>Superseded by ABLIKIM 20F.

NODE=M071S11  
NODE=M071S11

NODE=M071S11;LINKAGE=AB  
NODE=M071S11;LINKAGE=A

 $\Gamma(X(1750)\eta \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.8 ± 1.0 ± 2.6</b>	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$

NODE=M071P28  
NODE=M071P28

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;3.1</b>	90	<sup>1</sup> BAI	99C BES	$e^+ e^-$

<sup>1</sup>Assuming  $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

NODE=M071R45  
NODE=M071R45

NODE=M071R;LINKAGE=M3

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.12 ± 0.62<sup>+1.13</sup><sub>-0.61</sub></b>	251 ± 22	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S54  
NODE=M071S54

$\Gamma(K^*(892)^0 \bar{K}^0 + c.c.)/\Gamma_{\text{total}}$  $\Gamma_{90}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.9±2.0 OUR AVERAGE</b>				
$13.3^{+2.4}_{-2.8} \pm 1.7$	$65.6 \pm 9.0$	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
$9.2^{+2.7}_{-2.2} \pm 0.9$	25	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R30  
NODE=M071R30 $\Gamma(K^+ K^*(892)^- + c.c.)/\Gamma(K^*(892)^0 \bar{K}^0 + c.c.)$  $\Gamma_{80}/\Gamma_{90}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.16±0.06 OUR AVERAGE</b>			
$0.22^{+0.10}_{-0.14}$	ABLIKIM	05I BES2	$e^+ e^- \rightarrow \psi(2S)$
$0.14^{+0.08}_{-0.06}$	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R46  
NODE=M071R46 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{91}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.62±0.11 OUR AVERAGE</b>				
Error includes scale factor of 1.1.				
$1.56 \pm 0.04 \pm 0.11$	2.8k	ABLIKIM	14G BES3	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$2.38 \pm 0.37 \pm 0.29$	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1.9 \pm 0.3 \pm 0.3$	76.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
$1.5 \pm 0.3 \pm 0.2$	23	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071R78  
NODE=M071R78<sup>1</sup>Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

NODE=M071R78;LINKAGE=B3

 $\Gamma(\omega K_S^0 K_S^0)/\Gamma_{\text{total}}$  $\Gamma_{92}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.04±0.39±0.36</b>				
1.5k	ABLIKIM	21AL BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \pi^0 K_S^0 K_S^0$	

NODE=M071P34  
NODE=M071P34 $\Gamma(\omega K^*(892)^+ K^- + c.c.)/\Gamma_{\text{total}}$  $\Gamma_{93}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>20.7±2.6 OUR AVERAGE</b>				
$18.9 \pm 2.9 \pm 2.2$	396	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
$22.6 \pm 3.0 \pm 2.4$	535	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S67  
NODE=M071S67

OCCUR=2

 $\Gamma(\omega K_2^*(1430)^+ K^- + c.c.)/\Gamma_{\text{total}}$  $\Gamma_{94}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1 ±1.2 OUR AVERAGE</b>				
$6.39 \pm 1.50 \pm 0.78$	128	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
$5.86 \pm 1.61 \pm 0.83$	143	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S68  
NODE=M071S68

OCCUR=2

 $\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$  $\Gamma_{95}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>16.8±2.5±1.6</b>				
356	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	

NODE=M071S69  
NODE=M071S69 $\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$  $\Gamma_{96}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.82±2.08±0.72</b>				
116	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	

NODE=M071S70  
NODE=M071S70 $\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + c.c.)/\Gamma_{\text{total}}$  $\Gamma_{97}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.60±0.27±0.24</b>				
109	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	

NODE=M071S71  
NODE=M071S71<sup>1</sup>X(1440) compatible with  $\eta(1405)$  and  $\eta(1475)$ . A  $f_1(1420)$  is also possible.

NODE=M071S71;LINKAGE=AB

 $\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{98}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.09±0.20±0.16</b>				
82	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$	

NODE=M071S72  
NODE=M071S72<sup>1</sup>X(1440) compatible with  $\eta(1405)$  and  $\eta(1475)$ . A  $f_1(1420)$  is also possible.

NODE=M071S72;LINKAGE=AB

 $\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + c.c.)/\Gamma_{\text{total}}$  $\Gamma_{99}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.302±0.098±0.027</b>				
22	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$	

NODE=M071S73  
NODE=M071S73<sup>1</sup>Statistical significance  $4.5 \sigma$ . This measurement is equivalent to a limit of  $< 0.478 \times 10^{-5}$  at 90% C.L.

NODE=M071S73;LINKAGE=AB

$\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0) / \Gamma_{\text{total}}$  $\Gamma_{100} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.125 ± 0.070 ± 0.013</b>	10	<sup>1</sup> ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S74  
NODE=M071S74

<sup>1</sup> Statistical significance 3.2  $\sigma$ . This measurement is equivalent to a limit of  $< 0.221 \times 10^{-5}$  at 90% C.L.

NODE=M071S74;LINKAGE=AB

 $\Gamma(p\bar{p}) / \Gamma_{\text{total}}$  $\Gamma_{101} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.94 ± 0.08 OUR FIT****3.02 ± 0.08 OUR AVERAGE**

3.05 ± 0.02 ± 0.12	19k	ABLIKIM	18T BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.08 ± 0.05 ± 0.18	4.5k	<sup>1</sup> DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.36 ± 0.09 ± 0.25	1.6k	ABLIKIM	07C BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.87 ± 0.12 ± 0.15	557	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
1.4 ± 0.8	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.3 ± 0.7		FELDMAN	77 MRK1	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

NODE=M071R25  
NODE=M071R25

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071R25;LINKAGE=A

 $\Gamma(p\bar{p}) / \Gamma(J/\psi(1S) \pi^+ \pi^-)$  $\Gamma_{101} / \Gamma_{12}$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**8.49 ± 0.23 OUR FIT****6.98 ± 0.49 ± 0.97**

BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
-----	----	-----	---

NODE=M071S40  
NODE=M071S40 $\Gamma(n\bar{n}) / \Gamma_{\text{total}}$  $\Gamma_{102} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**3.06 ± 0.06 ± 0.14**

6k	ABLIKIM	18T BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$
----	---------	----------	---

NODE=M071P10  
NODE=M071P10 $\Gamma(p\bar{p}\pi^0) / \Gamma_{\text{total}}$  $\Gamma_{103} / \Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**1.53 ± 0.07 OUR AVERAGE**

1.65 ± 0.03 ± 0.15	4.5k	ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
1.54 ± 0.06 ± 0.06	948	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$
1.32 ± 0.10 ± 0.15	256	<sup>1</sup> ABLIKIM	05E BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
1.4 ± 0.5	9	FRANKLIN	83 MRK2	$e^+ e^-$

NODE=M071R35  
NODE=M071R35

<sup>1</sup> Computed using  $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$ .

NODE=M071R35;LINKAGE=AB

 $\Gamma(N(940)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0) / \Gamma_{\text{total}}$  $\Gamma_{104} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**6.42 ± 0.20<sup>+1.78</sup><sub>-1.28</sub>**

<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
----------------------	----------	--------------------------------------

NODE=M071S56  
NODE=M071S56

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S56;LINKAGE=AB

 $\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0) / \Gamma_{\text{total}}$  $\Gamma_{105} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**7.3<sup>+1.7</sup><sub>-1.5</sub> OUR AVERAGE** Error includes scale factor of 2.5.

3.58 ± 0.25 <sup>+1.59</sup> <sub>-0.84</sub>	1.1k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
8.1 ± 0.7 ± 0.3	474	<sup>2</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$

NODE=M071S50;LINKAGE=AB  
NODE=M071S50;LINKAGE=AL

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

<sup>2</sup> From a fit of the  $p\bar{p}$  and  $p\pi^0$  mass distributions to a combination of  $N(1440)\bar{p}$ , a broad  $p\bar{p}$  enhancement around 2100 MeV, and two other broad, unestablished resonances.

 $\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0) / \Gamma_{\text{total}}$  $\Gamma_{106} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**0.64 ± 0.05<sup>+0.22</sup><sub>-0.17</sub>**

<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
----------------------	----------	--------------------------------------

NODE=M071S57  
NODE=M071S57

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S57;LINKAGE=AB

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0) / \Gamma_{\text{total}}$  $\Gamma_{107} / \Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**2.47 ± 0.28<sup>+0.99</sup><sub>-0.97</sub>**

<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
----------------------	----------	--------------------------------------

NODE=M071S58  
NODE=M071S58

<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S58;LINKAGE=AB

$\Gamma(N(1650)\bar{p} + c.c. \rightarrow p\bar{p}\pi^0)/\Gamma_{total}$  $\Gamma_{108}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.76 \pm 0.28^{+1.37}_{-1.66}$	1.1k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

NODE=M071S59  
NODE=M071S59<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S59;LINKAGE=AB

 $\Gamma(N(1720)\bar{p} + c.c. \rightarrow p\bar{p}\pi^0)/\Gamma_{total}$  $\Gamma_{109}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.79 \pm 0.10^{+0.24}_{-0.71}$	0.5k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

NODE=M071S60  
NODE=M071S60<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S60;LINKAGE=AB

 $\Gamma(N(2300)\bar{p} + c.c. \rightarrow p\bar{p}\pi^0)/\Gamma_{total}$  $\Gamma_{110}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.62 \pm 0.28^{+1.12}_{-0.64}$	0.9k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

NODE=M071S61  
NODE=M071S61<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S61;LINKAGE=AB

 $\Gamma(N(2570)\bar{p} + c.c. \rightarrow p\bar{p}\pi^0)/\Gamma_{total}$  $\Gamma_{111}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.13 \pm 0.08^{+0.40}_{-0.30}$	0.8k	<sup>1</sup> ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$

NODE=M071S62  
NODE=M071S62<sup>1</sup> From a fit of  $\pi^0 p\bar{p}$  data to eight distinct intermediate  $N\bar{p}$  resonant states.

NODE=M071S62;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{total}$  $\Gamma_{112}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0 ± 0.4 OUR AVERAGE</b>				
$5.9 \pm 0.2 \pm 0.4$	904.5	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-$
$8 \pm 2$		<sup>1</sup> TANENBAUM	78 MRK1	$e^+e^-$

NODE=M071R31  
NODE=M071R31<sup>1</sup> Assuming entirely strong decay.

NODE=M071R;LINKAGE=K

 $\Gamma(p\bar{p}K^+K^-)/\Gamma_{total}$  $\Gamma_{113}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

NODE=M071S16  
NODE=M071S16 $\Gamma(p\bar{p}\eta)/\Gamma_{total}$  $\Gamma_{114}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.0 ± 0.4 OUR AVERAGE</b>				
$6.4 \pm 0.2 \pm 0.6$	679	<sup>1</sup> ABLIKIM	13S BES3	$\psi(2S) \rightarrow \eta p\bar{p}$
$5.6 \pm 0.6 \pm 0.3$	154	<sup>1</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta p\bar{p}$
$5.8 \pm 1.1 \pm 0.7$	$44.8 \pm 8.5$	<sup>2</sup> ABLIKIM	05E BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
$8 \pm 3 \pm 3$	9.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

NODE=M071R56  
NODE=M071R56<sup>1</sup> With  $N(1535)$  decaying to  $p\eta$ .<sup>2</sup> Computed using  $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$ .NODE=M071R56;LINKAGE=A  
NODE=M071R56;LINKAGE=AB $\Gamma(N(1535)\bar{p} + c.c. \rightarrow p\bar{p}\eta)/\Gamma_{total}$  $\Gamma_{115}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.5<sup>+0.7</sup><sub>-0.6</sub> OUR AVERAGE</b>				
$5.2 \pm 0.3^{+3.2}_{-1.2}$	527	<sup>1</sup> ABLIKIM	13S BES3	$\psi(2S) \rightarrow \eta p\bar{p}$
$4.4 \pm 0.6 \pm 0.3$	123	<sup>2</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta p\bar{p}$

NODE=M071S53  
NODE=M071S53<sup>1</sup> With  $N(1535)$  decaying to  $p\eta$ .<sup>2</sup> From a fit of the  $p\bar{p}$  and  $p\eta$  distributions to a combination of  $N^*(1535)\bar{p}$  and a broad  $p\bar{p}$  enhancement around 2100 MeV.NODE=M071S53;LINKAGE=A  
NODE=M071S53;LINKAGE=AL $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{total}$  $\Gamma_{116}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.3 ± 0.4 ± 0.6</b>	434.9	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

NODE=M071S15  
NODE=M071S15



$\Gamma(\rho\bar{\rho}\rho^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.5±0.1±0.2</b>	61.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-$

 $\Gamma_{117}/\Gamma$ NODE=M071S14  
NODE=M071S14 $\Gamma(\rho\bar{\rho}\omega)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.69±0.21 OUR AVERAGE</b>				
0.6 ±0.2 ±0.2	21.2	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-\pi^0$
0.8 ±0.3 ±0.1	14.9 ± 0.1	<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow \rho\bar{\rho}\pi^+\pi^-\pi^0$

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

 $\Gamma_{118}/\Gamma$ NODE=M071R79  
NODE=M071R79

NODE=M071R;LINKAGE=B3

 $\Gamma(\rho\bar{\rho}\eta')/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.10±0.10±0.08</b>	491	<sup>1</sup> ABLIKIM	19N BES3	$\psi(2S) \rightarrow \eta'\rho\bar{\rho}$

<sup>1</sup> From the combination of  $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\pi^+\pi^-\eta$  and  $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\pi^+\pi^-\gamma$  channels.

 $\Gamma_{119}/\Gamma$ NODE=M071P20  
NODE=M071P20

NODE=M071P20;LINKAGE=A

 $\Gamma(\rho\bar{\rho}\phi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.06±0.38±0.48</b>		753	ABLIKIM	19AO BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}K^+K^-$
<24	90		BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}K^+K^-$
<26	90		<sup>1</sup> BAI	03B BES	$\psi(2S) \rightarrow K^+K^-\rho\bar{\rho}$

<sup>1</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

 $\Gamma_{120}/\Gamma$ NODE=M071R82  
NODE=M071R82

NODE=M071R82;LINKAGE=B3

 $\Gamma(\phi X(1835) \rightarrow \rho\bar{\rho}\phi)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.82 × 10<sup>-7</sup></b>	90	ABLIKIM	19AO BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{\rho}K^+K^-$

 $\Gamma_{121}/\Gamma$ NODE=M071P21  
NODE=M071P21 $\Gamma(\rho\bar{\eta}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.48±0.17 OUR AVERAGE</b>				
2.45±0.11±0.21	851	ABLIKIM	06I BES2	$e^+e^- \rightarrow \rho\pi^-X$
2.52±0.12±0.22	849	ABLIKIM	06I BES2	$e^+e^- \rightarrow \bar{\rho}\pi^+X$

 $\Gamma_{122}/\Gamma$ NODE=M071R01  
NODE=M071R01

OCCUR=2


 $\Gamma(\rho\bar{\eta}\pi^-\pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.18±0.50±0.50</b>	135 ± 21	ABLIKIM	06I BES2	$e^+e^- \rightarrow \rho\pi^-\pi^0X$

 $\Gamma_{123}/\Gamma$ NODE=M071R02  
NODE=M071R02 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.81±0.13 OUR AVERAGE</b>					Error includes scale factor of 1.4. See the ideogram below.
3.97±0.02±0.12	31k	ABLIKIM	17L BES3		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
3.71±0.05±0.15	6.5k	<sup>1</sup> DOBBS	17		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
3.39±0.20±0.32	337	ABLIKIM	07C BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
6.4 ±1.8 ±0.1		<sup>2</sup> AUBERT	07BD BABR	10.6	$e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
3.28±0.23±0.25	208	PEDLAR	05 CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
3.75±0.09±0.23	1.9k	<sup>1,3</sup> DOBBS	14		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
1.81±0.20±0.27	80	<sup>4</sup> BAI	01 BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 4	90	FELDMAN	77 MRK1		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 $\Gamma_{124}/\Gamma$ NODE=M071R28  
NODE=M071R28NODE=M071R28;LINKAGE=A  
NODE=M071R28;LINKAGE=AU<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> AUBERT 07BD reports  $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$  keV which we divide by our best value  $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$  keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.<sup>3</sup> Superseded by DOBBS 17.<sup>4</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .NODE=M071R28;LINKAGE=B  
NODE=M071R28;LINKAGE=PP

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{125}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.42±0.39±0.59</b>		23	<sup>1</sup> ABLIKIM 22AP	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
< 2.9	90		<sup>2</sup> ABLIKIM 13F	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
< 120	90		<sup>3</sup> ABLIKIM 07H	BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R6  
NODE=M071R6<sup>1</sup> With a significance of 3.7  $\sigma$ . The corresponding 90% CL upper limit is  $2.47 \times 10^{-6}$ .<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$ .<sup>3</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.4\%$ .NODE=M071R6;LINKAGE=A  
NODE=M071R6;LINKAGE=AL  
NODE=M071R6;LINKAGE=AB $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$  $\Gamma_{126}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.43±0.32 OUR AVERAGE</b>					
[(2.5 ± 0.4) × 10 <sup>-5</sup> OUR 2022 AVERAGE]					
2.34±0.18±0.52		218	ABLIKIM 22AP	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
2.48±0.34±0.19		60	<sup>1</sup> ABLIKIM 13F	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
●●● We do not use the following data for averages, fits, limits, etc. ●●●					
< 4.9	90		<sup>2</sup> ABLIKIM 07H	BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R7  
NODE=M071R7  
NEW<sup>1</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$  and  $B(\eta \rightarrow \gamma\gamma) = 39.31\%$ .<sup>2</sup> Using  $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ .NODE=M071R7;LINKAGE=AL  
NODE=M071R7;LINKAGE=AB $\Gamma(\Lambda\bar{\Lambda}\omega(782))/\Gamma_{\text{total}}$  $\Gamma_{127}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.30±0.34±0.29</b>	207	<sup>1</sup> ABLIKIM 22AZ	BES3	$e^+e^- \rightarrow \psi(2S)$
<sup>1</sup> Using $B(\Lambda \rightarrow \pi^- p) = 0.639$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.893$ .				

NODE=M071P46  
NODE=M071P46

OCCUR=2

NODE=M071P46;LINKAGE=B

 $\Gamma(\Lambda(1670)\bar{\Lambda} \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$  $\Gamma_{128}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.29±0.31±0.62</b>	116	<sup>1</sup> ABLIKIM 22AP	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
<sup>1</sup> From a partial wave analysis of the $\Lambda\eta$ system.				

NODE=M071P42  
NODE=M071P42

NODE=M071P42;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{129}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.8±0.4±0.5</b>	73.4	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$

NODE=M071S17  
NODE=M071S17 $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$  $\Gamma_{130}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.0±0.1±0.1</b>	74.0	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^-$

NODE=M071S18  
NODE=M071S18

$\Gamma(\Lambda\bar{p}K^*(892)^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{131}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.3±0.5±0.5</b>	1011	ABLIKIM	19AU BES3	$e^+e^- \rightarrow \psi(2S)$

NODE=M071P25  
NODE=M071P25 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{132}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.8±0.3±0.3</b>	45.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+\pi^+\pi^-\pi^-$

NODE=M071S19  
NODE=M071S19 $\Gamma(\Lambda n K_S^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{133}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.81±0.11±0.14</b>	50	<sup>1</sup> ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

NODE=M071R08  
NODE=M071R08<sup>1</sup> Using  $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$  and  $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$ .

NODE=M071R08;LINKAGE=AB

 $\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$  $\Gamma_{134}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.8±1.0±3.4</b>	157	<sup>1</sup> BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

NODE=M071R50  
NODE=M071R50<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$ .

NODE=M071R50;LINKAGE=PP

 $\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{135}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.40±0.03±0.13</b>	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow$ hadrons

NODE=M071S65  
NODE=M071S65 $\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{136}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.54±0.04±0.13</b>	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow$ hadrons

NODE=M071S66  
NODE=M071S66 $\Gamma(\Lambda\bar{\Sigma}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{137}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.60±0.31±0.59</b>	60	ABLIKIM	21L BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

NODE=M071P29  
NODE=M071P29 $\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{138}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	COMMENT
<b>1.23±0.23±0.08</b>	30	<sup>1</sup> DOBBS	17 $e^+e^- \rightarrow \psi(2S) \rightarrow$ hadrons

NODE=M071P08  
NODE=M071P08

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071P08;LINKAGE=A

 $\Gamma(\Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{139}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.67±0.13±0.12</b>	276	<sup>1</sup> ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$

NODE=M071S63  
NODE=M071S63<sup>1</sup> Using  $B(\Lambda \rightarrow p\pi^-) = 63.9\%$ , and  $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$ .

NODE=M071S63;LINKAGE=AB

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$  $\Gamma_{140}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.43±0.10 OUR AVERAGE</b>				Error includes scale factor of 1.4.

NODE=M071R47  
NODE=M071R472.52±0.04±0.09 5.4k ABLIKIM 21AT BES3  $\psi(2S) \rightarrow p\pi^0\bar{p}\pi^0$ 2.31±0.06±0.10 1.9k <sup>1</sup> DOBBS 17  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons2.57±0.44±0.68 35 PEDLAR 05 CLEO  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.51±0.15±0.16 281 <sup>1,2</sup> DOBBS 14  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> Superseded by DOBBS 17.NODE=M071R47;LINKAGE=A  
NODE=M071R47;LINKAGE=B $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{141}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.35±0.09 OUR AVERAGE</b>				Error includes scale factor of 1.1.

NODE=M071R51  
NODE=M071R512.44±0.03±0.11 7k ABLIKIM 17L BES3  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons2.22±0.05±0.11 2.6k <sup>1</sup> DOBBS 17  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons2.35±0.36±0.32 59 ABLIKIM 07C BES  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons2.63±0.35±0.21 58 PEDLAR 05 CLEO  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25±0.11±0.16 439 <sup>1,2</sup> DOBBS 14  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons1.2 ±0.4 ±0.4 8 <sup>3</sup> BAI 01 BES  $e^+e^- \rightarrow \psi(2S) \rightarrow$  hadrons

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Superseded by DOBBS 17.

<sup>3</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .

NODE=M071R51;LINKAGE=A  
NODE=M071R51;LINKAGE=B  
NODE=M071R51;LINKAGE=PP

### $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$ $\Gamma_{142}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.82 ± 0.04 ± 0.08</b>	6.6k	ABLIKIM	22AV BES3	$\psi(2S) \rightarrow n \pi^- \bar{n} \pi^+$

NODE=M071P41  
NODE=M071P41

### $\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$ $\Gamma_{143}/\Gamma$

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.59 ± 2.37 ± 0.61</b>	21	ABLIKIM	22AY BES3	$\psi(2S) \rightarrow \Sigma^+ \bar{\Sigma}^- \eta$

NODE=M071P45  
NODE=M071P45

### $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ $\Gamma_{144}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.5 ± 0.7 OUR AVERAGE</b>				

NODE=M071R52  
NODE=M071R52

8.4 ± 0.5 ± 0.5	1.5k	ABLIKIM	16L BES3	$\psi(2S) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
11 ± 3 ± 3	14	<sup>1</sup> BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

OCCUR=2

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .

NODE=M071R52;LINKAGE=PP

### $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ $\Gamma_{145}/\Gamma$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.5 ± 0.6 ± 0.6</b>	1.4k	ABLIKIM	16L BES3	$\psi(2S) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$

NODE=M071R00  
NODE=M071R00

### $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ $\Gamma_{146}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.69 ± 0.05 ± 0.05</b>	2.2k	ABLIKIM	17E BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071P00  
NODE=M071P00

### $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ $\Gamma_{147}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.87 ± 0.11 OUR AVERAGE</b>					Error includes scale factor of 1.1.

NODE=M071R29  
NODE=M071R29

3.03 ± 0.05 ± 0.14	3.6k	<sup>1</sup> DOBBS	17		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.78 ± 0.05 ± 0.14	5k	ABLIKIM	16L BES3		$\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$
3.03 ± 0.40 ± 0.32	67	ABLIKIM	07C BES		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.38 ± 0.30 ± 0.21	63	PEDLAR	05 CLEO		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.66 ± 0.12 ± 0.20	548	<sup>1,2</sup> DOBBS	14		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
0.94 ± 0.27 ± 0.15	12	<sup>3</sup> BAI	01 BES		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<2	90	FELDMAN	77 MRK1		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Superseded by DOBBS 17.

<sup>3</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .

NODE=M071R29;LINKAGE=A  
NODE=M071R29;LINKAGE=B  
NODE=M071R29;LINKAGE=PP

### $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$ $\Gamma_{148}/\Gamma$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.3 ± 0.4 OUR AVERAGE</b>				Error includes scale factor of 4.2.

NODE=M071R48  
NODE=M071R48

2.73 ± 0.03 ± 0.13	11k	ABLIKIM	17E BES3		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
1.97 ± 0.06 ± 0.11	1.2k	<sup>1</sup> DOBBS	17		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.75 ± 0.64 ± 0.61	19	PEDLAR	05 CLEO		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.02 ± 0.19 ± 0.15	112	<sup>1,2</sup> DOBBS	14		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
--------------------	-----	----------------------	----	--	---

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Superseded by DOBBS 17.

NODE=M071R48;LINKAGE=A  
NODE=M071R48;LINKAGE=B

### $\Gamma(\Xi(1530)^0 \bar{\Xi}(1530)^0)/\Gamma_{\text{total}}$ $\Gamma_{149}/\Gamma$

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.77 ± 0.14 ± 0.39</b>		2951	ABLIKIM	21AO BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071R53  
NODE=M071R53

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	PEDLAR	05 CLEO		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 8.1	90	<sup>1</sup> BAI	01 BES		$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .

NODE=M071R53;LINKAGE=PP

$\Gamma(\Lambda\bar{\Xi}^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{150}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.86 ± 0.27 ± 0.32</b>	236	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda\bar{\Xi}^+ + \text{c.c.}$

NODE=M071S82  
NODE=M071S82 $\Gamma(\Xi(1530)^- \bar{\Xi}(1530)^+)/\Gamma_{\text{total}}$  $\Gamma_{151}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>11.45 ± 0.40 ± 0.59</b>	5k	ABLIKIM	19AT BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071P26  
NODE=M071P26 $\Gamma(\Xi(1530)^- \bar{\Xi}^+)/\Gamma_{\text{total}}$  $\Gamma_{152}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.0 ± 1.1 ± 0.4</b>	399	ABLIKIM	19AT BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071P27  
NODE=M071P27 $\Gamma(\Xi(1530)^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$  $\Gamma_{153}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.53 ± 0.04 ± 0.03</b>	278	ABLIKIM	21AO BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071P35  
NODE=M071P35 $\Gamma(\Xi(1690)^- \bar{\Xi}^+ \rightarrow K^- \Lambda\bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{154}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.21 ± 1.48 ± 0.57</b>	74	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda\bar{\Xi}^+ + \text{c.c.}$

NODE=M071S83  
NODE=M071S83 $\Gamma(\Xi(1820)^- \bar{\Xi}^+ \rightarrow K^- \Lambda\bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{155}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.03 ± 2.94 ± 1.22</b>	136	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda\bar{\Xi}^+ + \text{c.c.}$

NODE=M071S84  
NODE=M071S84 $\Gamma(\Sigma^0 \bar{\Xi}^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{156}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.67 ± 0.33 ± 0.28</b>	142	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \bar{\Xi}^+ + \text{c.c.}$

NODE=M071S85  
NODE=M071S85 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$  $\Gamma_{157}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.66 ± 0.30 OUR AVERAGE</b>			Error includes scale factor of 1.3.		
5.85 ± 0.12 ± 0.25		4k	<sup>1</sup> ABLIKIM	21E BES3	$\psi(2S) \rightarrow \Omega^- \bar{\Omega}^+ \rightarrow \Lambda K^- \bar{\Lambda} K^+$
5.2 ± 0.3 ± 0.3		326	<sup>1,2</sup> DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
4.7 ± 0.9 ± 0.5		27	<sup>1,2,3</sup> DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<15	90		ABLIKIM	12Q BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<16	90		PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 7.3	90		<sup>4</sup> BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071R54  
NODE=M071R54<sup>1</sup> Using  $B(\Omega^- \rightarrow \Lambda K^-) = (67.8 \pm 0.7)\%$  and  $B(\Lambda \rightarrow p \pi^-) = (63.9 \pm 0.5)\%$ .<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>3</sup> Superseded by DOBBS 17.<sup>4</sup> Estimated using  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$ .NODE=M071R54;LINKAGE=D  
NODE=M071R54;LINKAGE=A  
NODE=M071R54;LINKAGE=B  
NODE=M071R54;LINKAGE=PP $\Gamma(\eta_c \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{158}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.0</b>	90	PEDLAR	07 CLEO	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R03  
NODE=M071R03 $\Gamma(h_c(1P) \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{159}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.4 ± 0.5 OUR AVERAGE</b>				
[(8.6 ± 1.3) × 10 <sup>-4</sup> OUR 2022 AVERAGE]				
7.32 ± 0.34 ± 0.41	46k	ABLIKIM	22AQ BES3	$\psi(2S) \rightarrow \pi^0 \text{ hadrons}$
9.0 ± 1.5 ± 1.3	3k	<sup>1</sup> GE	11 CLEO	$\psi(2S) \rightarrow \pi^0 \text{ anything}$

NODE=M071S42  
NODE=M071S42

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.4 ± 1.3 ± 1.0	11k	<sup>2</sup> ABLIKIM	10B	BES3	$\psi(2S) \rightarrow \pi^0 h_c$
seen	92 <sup>+23</sup> <sub>-22</sub>	ADAMS	09	CLEO	$\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$
seen	1282	DOBBS	08A	CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
seen	168 ± 40	ROSNER	05	CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

<sup>1</sup> Assuming a width  $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$ , a measured dependence of the central value of  $B = (7.6 + 1.4 \times \Gamma(h_c(1P))/\Gamma_0) \times 10^{-4}$ , and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

<sup>2</sup> Superseded by ABLIKIM 22AQ

NODE=M071S42;LINKAGE=GE

NODE=M071S42;LINKAGE=A

$\Gamma(\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$						$\Gamma_{160}/\Gamma$	
VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT		
<b>&lt;1.7 × 10<sup>-6</sup></b>	90	450M	ABLIKIM	18Q	BES3	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071P11 NODE=M071P11

$\Gamma(\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 \rho K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$						$\Gamma_{161}/\Gamma$	
VALUE (units 10 <sup>-5</sup> )	CL%		DOCUMENT ID	TECN	COMMENT		
<b>&lt;0.88</b>	90		BAI	04G	BES2	$e^+ e^-$	NODE=M071S01 NODE=M071S01

$\Gamma(\Theta(1540) K^- \bar{n} \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$						$\Gamma_{162}/\Gamma$	
VALUE (units 10 <sup>-5</sup> )	CL%		DOCUMENT ID	TECN	COMMENT		
<b>&lt;1.0</b>	90		BAI	04G	BES2	$e^+ e^-$	NODE=M071S02 NODE=M071S02

$\Gamma(\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$						$\Gamma_{163}/\Gamma$	
VALUE (units 10 <sup>-5</sup> )	CL%		DOCUMENT ID	TECN	COMMENT		
<b>&lt;0.70</b>	90		BAI	04G	BES2	$e^+ e^-$	NODE=M071S03 NODE=M071S03

$\Gamma(\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$						$\Gamma_{164}/\Gamma$	
VALUE (units 10 <sup>-5</sup> )	CL%		DOCUMENT ID	TECN	COMMENT		
<b>&lt;2.6</b>	90		BAI	04G	BES2	$e^+ e^-$	NODE=M071S04 NODE=M071S04

$\Gamma(\bar{\Theta}(1540) K_S^0 \rho \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$						$\Gamma_{165}/\Gamma$	
VALUE (units 10 <sup>-5</sup> )	CL%		DOCUMENT ID	TECN	COMMENT		
<b>&lt;0.60</b>	90		BAI	04G	BES2	$e^+ e^-$	NODE=M071S05 NODE=M071S05

### RADIATIVE DECAYS

$\Gamma(\gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$						$\Gamma_{166}/\Gamma$
VALUE (units 10 <sup>-2</sup> )		EVTs	DOCUMENT ID	TECN	COMMENT	
<b>9.79 ± 0.20</b>	<b>OUR FIT</b>					
<b>9.33 ± 0.26</b>	<b>OUR AVERAGE</b>					
9.389 ± 0.014 ± 0.332	4.7M		ABLIKIM	17U	BES3	$e^+ e^- \rightarrow \gamma X$
9.22 ± 0.11 ± 0.46	72k		ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
9.9 ± 0.5 ± 0.8			<sup>1</sup> GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.2 ± 2.3			<sup>1</sup> BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$
7.5 ± 2.6			<sup>1</sup> WHITAKER	76	MRK1	$e^+ e^-$

<sup>1</sup> Angular distribution  $(1 + \cos^2\theta)$  assumed.

NODE=M071315

NODE=M071R55  
NODE=M071R55

NODE=M071R;LINKAGE=A

$\Gamma(\gamma \chi_{c1}(1P))/\Gamma_{\text{total}}$						$\Gamma_{167}/\Gamma$
VALUE (units 10 <sup>-2</sup> )		EVTs	DOCUMENT ID	TECN	COMMENT	
<b>9.75 ± 0.24</b>	<b>OUR FIT</b>					
<b>9.54 ± 0.29</b>	<b>OUR AVERAGE</b>					
9.905 ± 0.011 ± 0.353	5.0M		ABLIKIM	17U	BES3	$e^+ e^- \rightarrow \gamma X$
9.07 ± 0.11 ± 0.54	76k		ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
9.0 ± 0.5 ± 0.7			<sup>1</sup> GAISER	86	CBAL	$e^+ e^- \rightarrow \gamma X$
7.1 ± 1.9			<sup>2</sup> BIDDICK	77	CNTR	$e^+ e^- \rightarrow \gamma X$

<sup>1</sup> Angular distribution  $(1 - 0.189 \cos^2\theta)$  assumed.

<sup>2</sup> Valid for isotropic distribution of the photon.

NODE=M071R58  
NODE=M071R58

NODE=M071R;LINKAGE=G  
NODE=M071R;LINKAGE=B

$\Gamma(\gamma \chi_{c0}(1P))/\Gamma(\gamma \chi_{c1}(1P))$				$\Gamma_{166}/\Gamma_{167}$
VALUE		DOCUMENT ID	TECN	COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.02 ± 0.01 ± 0.07	<sup>1</sup> ATHAR	04	CLEO	$e^+ e^- \rightarrow \gamma X$
--------------------	--------------------	----	------	--------------------------------

<sup>1</sup> Not independent from ATHAR 04 measurements of  $B(\gamma \chi_{cJ})$ .

NODE=M071R97  
NODE=M071R97

NODE=M071R97;LINKAGE=AH


$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{total}$   $\Gamma_{168}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.52 ±0.20 OUR FIT</b>				
<b>9.42 ±0.31 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
9.621±0.013±0.272	4.2M	ABLIKIM	17U	BES3 $e^+e^- \rightarrow \gamma X$
9.33 ±0.14 ±0.61	79k	ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
8.0 ±0.5 ±0.7		<sup>1</sup> GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$
7.0 ±2.0		<sup>2</sup> BIDDICK	77	CNTR $e^+e^- \rightarrow \gamma X$

NODE=M071R59  
NODE=M071R59

<sup>1</sup> Angular distribution  $(1-0.052 \cos^2\theta)$  assumed.  
<sup>2</sup> Valid for isotropic distribution of the photon.

NODE=M071R;LINKAGE=F  
NODE=M071R59;LINKAGE=B



$[\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))]/\Gamma_{total}$  ( $\Gamma_{166}+\Gamma_{167}+\Gamma_{168}$ )/ $\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
27.6±0.3±2.0	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$

NODE=M071R19  
NODE=M071R19

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Not independent from ATHAR 04 measurements of  $B(\gamma\chi_{cJ})$ .

NODE=M071R;LINKAGE=AH

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c2}(1P))$   $\Gamma_{166}/\Gamma_{168}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.99±0.02±0.08	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$

NODE=M071R99  
NODE=M071R99

<sup>1</sup> Not independent from ATHAR 04 measurements of  $B(\gamma\chi_{cJ})$ .

NODE=M071R99;LINKAGE=AH

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma(\gamma\chi_{c1}(1P))$   $\Gamma_{168}/\Gamma_{167}$

VALUE	DOCUMENT ID	TECN	COMMENT
1.03±0.02±0.03	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$

NODE=M071R98  
NODE=M071R98

<sup>1</sup> Not independent from ATHAR 04 measurements of  $B(\gamma\chi_{cJ})$ .

NODE=M071R98;LINKAGE=AH

$\Gamma(\gamma\eta_c(1S))/\Gamma_{total}$   $\Gamma_{169}/\Gamma$


VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.34 ±0.05 OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.
0.432±0.016±0.060		MITCHELL	09	CLEO $e^+e^- \rightarrow \gamma X$
0.32 ±0.04 ±0.06	2.5k	<sup>1</sup> ATHAR	04	CLEO $e^+e^- \rightarrow \gamma X$
0.28 ±0.06		<sup>2</sup> GAISER	86	CBAL $e^+e^- \rightarrow \gamma X$

NODE=M071R60  
NODE=M071R60

<sup>1</sup> ATHAR 04 used  $\Gamma_{\eta_c(1S)} = 24.8 \pm 4.9$  MeV to obtain this result.

<sup>2</sup> GAISER 86 used  $\Gamma_{\eta_c(1S)} = 11.5 \pm 4.5$  MeV to obtain this result.

NODE=M071R60;LINKAGE=AT  
 NODE=M071R60;LINKAGE=GA



**$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$**   **$\Gamma_{170}/\Gamma$**

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b><math>7 \pm 2 \pm 4</math></b>		<sup>1</sup> ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K\pi,$ $K K \pi^0$

NODE=M071R62  
 NODE=M071R62

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8	90	<sup>2</sup> CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K \bar{K} \pi$
< 20	90	ATHAR	04 CLEO	$e^+ e^- \rightarrow \gamma X$
20-130	95	EDWARDS	82c CBAL	$e^+ e^- \rightarrow \gamma X$

NODE=M071R62;LINKAGE=AB

<sup>1</sup> ABLIKIM 12G reports  $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$  which we divide by our best value  $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> CRONIN-HENNESSY 10 reports  $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$  which we divide by our best value  $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$ . This measurement assumes  $\Gamma(\eta_c(2S)) = 14$  MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M071R62;LINKAGE=CR

**$\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$**   **$\Gamma_{171}/\Gamma$**

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.04 \pm 0.22</math> OUR AVERAGE</b>			Error includes scale factor of 1.4.		
$0.95 \pm 0.16 \pm 0.05$		423	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^0$
$1.58 \pm 0.40 \pm 0.13$		37	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^0$

NODE=M071R42  
 NODE=M071R42

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5	90	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
< 5400	95	<sup>1</sup> LIBERMAN	75 SPEC	$e^+ e^-$
< $1 \times 10^4$	90	WIJK	75 DASP	$e^+ e^-$

<sup>1</sup> Restated by us using  $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$ .

NODE=M071R;LINKAGE=U

**$\Gamma(\gamma 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$**   **$\Gamma_{172}/\Gamma$**

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>39.6 \pm 2.8 \pm 5.0</math></b>	583	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S28  
 NODE=M071S28

**$\Gamma(\gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$**   **$\Gamma_{173}/\Gamma$**

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 17</b>	90	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S36  
 NODE=M071S36



$\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.24 ± 0.04</b>		<b>OUR AVERAGE</b>			
1.251 ± 0.022 ± 0.062		56k	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta,$ $\gamma\pi^0\pi^0\eta$
1.26 ± 0.03 ± 0.08		2226	<sup>1</sup> ABLIKIM	10F BES3	$\psi(2S) \rightarrow 3\gamma\pi^+\pi^-,$ $2\gamma\pi^+\pi^-$
1.19 ± 0.08 ± 0.03			PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
1.24 ± 0.27 ± 0.15		23	ABLIKIM	06R BES2	$e^+e^- \rightarrow \psi(2S)$
1.54 ± 0.31 ± 0.20		~ 43	BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^-2\gamma,$ $\pi^+\pi^-3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60	90	<sup>2</sup> BRAUNSCH...	77	DASP	$e^+e^-$
< 11	90	<sup>3</sup> BARTEL	76	CNTR	$e^+e^-$

<sup>1</sup> Combining the results from  $\eta' \rightarrow \pi^+\pi^-\eta$  and  $\eta' \rightarrow \pi^+\pi^-\gamma$  decay modes.

<sup>2</sup> Restated by us using total decay width 228 keV.

<sup>3</sup> The value is normalized to the branching ratio for  $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ .

NODE=M071R44  
NODE=M071R44

NODE=M071R44;LINKAGE=AB  
NODE=M071R;LINKAGE=R  
NODE=M071R;LINKAGE=C

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.73<sup>+0.29</sup><sub>-0.25</sub></b>		<b>OUR AVERAGE</b>		Error includes scale factor of 1.8.
2.84 ± 0.15 <sup>+0.03</sup> <sub>-0.10</sub>	1.9k	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
2.12 ± 0.19 ± 0.32		<sup>3,4</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.08 ± 0.19 ± 0.33	200.6 ± 18.8	<sup>3</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
2.90 ± 1.08 ± 1.07	29.9 ± 11.1	<sup>3</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> DOBBS 15 reports  $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$  which we divide by our best value  $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.9}_{-0.9}) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

<sup>4</sup> Combining the results from  $\pi^+\pi^-$  and  $\pi^0\pi^0$  decay modes.

NODE=M071R84  
NODE=M071R84

OCCUR=2  
OCCUR=3

NODE=M071R84;LINKAGE=A  
NODE=M071R84;LINKAGE=B

NODE=M071R;LINKAGE=3B  
NODE=M071R;LINKAGE=B9

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.1 ± 1.0 ± 1.4</b>	175	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S75  
NODE=M071S75

NODE=M071S75;LINKAGE=A

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.3 ± 1.8 ± 0.6</b>	274	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$

<sup>1</sup> DOBBS 15 reports  $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi\pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$  which we divide by our best value  $B(f_0(1500) \rightarrow \pi\pi) = (34.5 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S76  
NODE=M071S76

NODE=M071S76;LINKAGE=A

NODE=M071S76;LINKAGE=B

 $\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.3 ± 0.8 ± 0.1</b>	136	<sup>1,2</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K\bar{K}$

<sup>1</sup> DOBBS 15 reports  $[\Gamma(\psi(2S) \rightarrow \gamma f_2'(1525))/\Gamma_{\text{total}}] \times [B(f_2'(1525) \rightarrow K\bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$  which we divide by our best value  $B(f_2'(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S77  
NODE=M071S77

NODE=M071S77;LINKAGE=A

NODE=M071S77;LINKAGE=B

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.5 ± 0.6</b>		<b>OUR AVERAGE</b>		
3.6 ± 0.4 ± 0.5	290	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
3.01 ± 0.41 ± 1.24	35.6 ± 4.8	<sup>2</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$

<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>2</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$ .

NODE=M071R85  
NODE=M071R85

NODE=M071R85;LINKAGE=A  
NODE=M071R85;LINKAGE=3B

$\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$  $\Gamma_{181}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.6 ± 0.7 OUR AVERAGE</b>					
6.7 ± 0.6 ± 0.6		375	<sup>1</sup> DOBBS	15	$\psi(2S) \rightarrow \gamma K \bar{K}$
6.04 ± 0.90 ± 1.32	39.6 ± 5.9		<sup>2,3</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 15.6	90	6.8 ± 3.1	<sup>2,3</sup> BAI	03C BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

NODE=M071R86  
NODE=M071R86<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> Includes unknown branching fractions to  $K^+ K^-$  or  $K_S^0 K_S^0$ . We have multiplied the  $K^+ K^-$  result by a factor of 2 and the  $K_S^0 K_S^0$  result by a factor of 4 to obtain the  $K \bar{K}$  result.<sup>3</sup> Normalized to  $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$ .

OCCUR=2

NODE=M071R86;LINKAGE=A

NODE=M071R;LINKAGE=CK

NODE=M071R86;LINKAGE=3B

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$  $\Gamma_{182}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	COMMENT
<b>4.8 ± 0.5 ± 0.9</b>			
	373	<sup>1</sup> DOBBS	15 $\psi(2S) \rightarrow \gamma \pi \pi$

NODE=M071S78  
NODE=M071S78<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S78;LINKAGE=A

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$  $\Gamma_{183}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	COMMENT
<b>3.2 ± 0.6 ± 0.8</b>			
	207	<sup>1</sup> DOBBS	15 $\psi(2S) \rightarrow \gamma K \bar{K}$

NODE=M071S79  
NODE=M071S79<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S79;LINKAGE=A

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$  $\Gamma_{184}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	COMMENT
<b>&lt; 5.8 × 10<sup>-6</sup></b>			
	90	<sup>1,2</sup> DOBBS	15 $\psi(2S) \rightarrow \gamma \pi \pi$

NODE=M071S80  
NODE=M071S80<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $\pi^+ \pi^-$  and  $\pi^0 \pi^0$  are  $3.2/4.3 \times 10^{-6}$  and  $2.6/4.0 \times 10^{-6}$ , respectively.

NODE=M071S80;LINKAGE=A

NODE=M071S80;LINKAGE=DO

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$  $\Gamma_{185}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	COMMENT
<b>&lt; 9.5 × 10<sup>-6</sup></b>			
	90	<sup>1,2</sup> DOBBS	15 $\psi(2S) \rightarrow \gamma K \bar{K}$

NODE=M071S81  
NODE=M071S81<sup>1</sup> Using CLEO-c data but not authored by the CLEO Collaboration.<sup>2</sup> For  $\Gamma = 20/50$  MeV, the 90% CL upper limits for  $K^+ K^-$  and  $K_S^0 K_S^0$  are  $2.1/4.3 \times 10^{-6}$  and  $3.7/5.5 \times 10^{-6}$ , respectively.

NODE=M071S81;LINKAGE=A

NODE=M071S81;LINKAGE=DO

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$  $\Gamma_{186}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.92 ± 0.18 OUR AVERAGE</b>					
0.85 ± 0.18 ± 0.04		382	<sup>1</sup> ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0,$ $\gamma 3\pi^0$
1.38 ± 0.48 ± 0.09		13	<sup>1</sup> ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0,$ $\gamma 3\pi^0$

NODE=M071R43  
NODE=M071R43

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2	90	PEDLAR	09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 90	90	BAI	98F	BES	$\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$
< 200	90	YAMADA	77	DASP	$e^+ e^- \rightarrow 3\gamma$

<sup>1</sup> Combining the results from  $\eta \rightarrow \pi^+ \pi^- \pi^0$  and  $\eta \rightarrow 3\pi^0$  decay modes.

NODE=M071R43;LINKAGE=AB

 $\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{187}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.71 ± 1.25 ± 1.64</b>				
	418	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M071R04  
NODE=M071R04 $\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$  $\Gamma_{189}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.9</b>				
	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

NODE=M071R61  
NODE=M071R61

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$	OCCUR=2
< 1.2	90	<sup>1</sup> SCHARRE	80 MRK1	$e^+ e^-$	

<sup>1</sup> Includes unknown branching fraction  $\eta(1405) \rightarrow K \bar{K} \pi$ .

NODE=M071R;LINKAGE=E

$\Gamma(\gamma\eta(1405) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{190}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.36±0.25±0.05</b>	10	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M071R05  
NODE=M071R05 $\Gamma(\gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{191}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.0 × 10<sup>-7</sup></b>	90	ABLIKIM	17AJ BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

NODE=M071P09  
NODE=M071P09 $\Gamma(\gamma\eta(1475) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{\text{total}}$   $\Gamma_{193}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.4</b>	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.5	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

NODE=M071R06  
NODE=M071R06

OCCUR=2

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{194}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.88</b>	90	ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M071R07  
NODE=M071R07 $\Gamma(\gamma K^*0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{195}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>37.0±6.1±7.2</b>	237	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S29  
NODE=M071S29 $\Gamma(\gamma K^*0 \bar{K}^*0)/\Gamma_{\text{total}}$   $\Gamma_{196}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>24.0±4.5±5.0</b>	41	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S30  
NODE=M071S30 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{197}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>25.6±3.6±3.6</b>	115	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S31  
NODE=M071S31 $\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{198}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>19.1±2.7±4.3</b>	132	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S32  
NODE=M071S32 $\Gamma(\gamma K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{199}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;22</b>	90	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S35  
NODE=M071S35 $\Gamma(\gamma 2(K^+ K^-))/\Gamma_{\text{total}}$   $\Gamma_{200}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4</b>	90	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S37  
NODE=M071S37 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{201}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.9 ±0.5 OUR AVERAGE</b>				Error includes scale factor of 2.0.
4.18±0.26±0.18	348	<sup>1</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
2.9 ±0.4 ±0.4	142	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S33  
NODE=M071S33

<sup>1</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma f_2(1950)$ ,  $\gamma f_2(2150)$ , and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV, and accounting for backgrounds from  $\psi(2S) \rightarrow \pi^0 p\bar{p}$  and continuum.

NODE=M071S33;LINKAGE=AL

 $\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{202}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.2±0.2±0.1</b>	111	<sup>1</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M071S46  
NODE=M071S46

<sup>1</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma f_2(1950)$ ,  $\gamma f_2(2150)$ , and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV, and accounting for backgrounds from  $\psi(2S) \rightarrow \pi^0 p\bar{p}$  and continuum.

NODE=M071S46;LINKAGE=AL

 $\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{203}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.72±0.18±0.03</b>	73	<sup>1</sup> ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$

NODE=M071S47  
NODE=M071S47

<sup>1</sup> From a fit of the  $p\bar{p}$  mass distribution to a combination of  $\gamma f_2(1950)$ ,  $\gamma f_2(2150)$ , and  $\gamma p\bar{p}$  phase space, for  $M(p\bar{p}) < 2.85$  GeV, and accounting for backgrounds from  $\psi(2S) \rightarrow \pi^0 p\bar{p}$  and continuum.

NODE=M071S47;LINKAGE=AL

$\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$  $\Gamma_{204}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$4.57 \pm 0.36^{+1.77}_{-4.26}$		ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p \bar{p}$

NODE=M071S48  
 NODE=M071S48

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6	90	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$
<5.4	90	ABLIKIM	07D BES	$\psi(2S) \rightarrow \gamma p \bar{p}$

 $\Gamma(\gamma X \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$  $\Gamma_{205}/\Gamma$ 

For a narrow resonance in the range  $2.2 < M(X) < 2.8$  GeV.

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \gamma p \bar{p}$

NODE=M071S49  
 NODE=M071S49  
 NODE=M071S49

 $\Gamma(\gamma p \bar{p} \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_{206}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 1.2 \pm 0.7$	17	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S34  
 NODE=M071S34

 $\Gamma(\gamma \gamma J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{208}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.1 \pm 0.6^{+0.8}_{-1.0}$	1.1k	ABLIKIM	120 BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S55  
 NODE=M071S55

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.2 \pm 0.6$	1.1k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
---------------	------	----------------------	----------	---

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$ . No systematic error estimation.

NODE=M071S55;LINKAGE=A

 $\Gamma(e^+ e^- \eta')/\Gamma_{\text{total}}$  $\Gamma_{209}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.90 ± 0.26 OUR AVERAGE</b>				
$1.99 \pm 0.33 \pm 0.12$	57	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$ , $\eta' \rightarrow \gamma \pi^+ \pi^-$
$1.79 \pm 0.38 \pm 0.11$	20	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$ , $\eta' \rightarrow \eta \pi^+ \pi^-$

NODE=M071P12  
 NODE=M071P12

OCCUR=2

 $\Gamma(e^+ e^- \eta_c(1S))/\Gamma_{\text{total}}$  $\Gamma_{210}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.77 ± 0.40 ± 0.18</b>	3k	<sup>1</sup> ABLIKIM	22AX BES3	$e^+ e^- \rightarrow \psi(2S)$

<sup>1</sup> From a fit to the recoil mass distribution of  $e^+ e^-$  with inclusive  $\eta_c(1S)$  decays.

NODE=M071P44  
 NODE=M071P44

NODE=M071P44;LINKAGE=A

 $\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}$  $\Gamma_{211}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.6 ± 2.4 ± 0.4</b>	48	<sup>1</sup> ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> ABLIKIM 17I reports  $(11.7 \pm 2.5 \pm 1.0) \times 10^{-4}$  from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c0}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.27 \pm 0.06) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (1.40 \pm 0.05) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P01  
 NODE=M071P01

NODE=M071P01;LINKAGE=B

 $\Gamma(e^+ e^- \chi_{c0}(1P))/\Gamma(\gamma \chi_{c0}(1P))$  $\Gamma_{211}/\Gamma_{166}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.4 ± 1.9 ± 0.6</b>	48	<sup>1</sup> ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

NODE=M071P04  
 NODE=M071P04

NODE=M071P04;LINKAGE=A

 $\Gamma(e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}$  $\Gamma_{212}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.5 ± 0.6 ± 0.2</b>	873	<sup>1</sup> ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$

<sup>1</sup> ABLIKIM 17I reports  $(8.6 \pm 0.3 \pm 0.6) \times 10^{-4}$  from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+ e^- \chi_{c1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P02  
 NODE=M071P02

NODE=M071P02;LINKAGE=B

$\Gamma(e^+e^-\chi_{c1}(1P))/\Gamma(\gamma\chi_{c1}(1P))$  $\Gamma_{212}/\Gamma_{167}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.3±0.3±0.4</b>	873	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

NODE=M071P05  
NODE=M071P05

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

NODE=M071P05;LINKAGE=A

 $\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma_{\text{total}}$  $\Gamma_{213}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.0±0.7±0.2</b>	227	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

NODE=M071P03  
NODE=M071P03

<sup>1</sup> ABLIKIM 17I reports  $(6.9 \pm 0.5 \pm 0.6) \times 10^{-4}$  from a measurement of  $[\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P03;LINKAGE=B

 $\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma(\gamma\chi_{c2}(1P))$  $\Gamma_{213}/\Gamma_{168}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.6±0.5±0.4</b>	227	1 ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

NODE=M071P06  
NODE=M071P06

<sup>1</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$  from ABLIKIM 17N and accounts for common systematic errors.

NODE=M071P06;LINKAGE=A

## WEAK DECAYS

NODE=M071330

 $\Gamma(D^0e^+e^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{214}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.4 × 10<sup>-7</sup></b>	90	1 ABLIKIM	17AF BES3	$e^+e^- \rightarrow \psi(2S)$

NODE=M071P07  
NODE=M071P07

<sup>1</sup> Using  $D^0$  decays to  $K^-\pi^+$ ,  $K^-\pi^+\pi^0$ , and  $K^-\pi^+\pi^+\pi^-$ .

NODE=M071P07;LINKAGE=A

 $\Gamma(\Lambda_c^+\bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{215}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.4 × 10<sup>-5</sup></b>	90	1 ABLIKIM	23 BES3	$e^+e^- \rightarrow \psi(2S)$

NODE=M071P43  
NODE=M071P43

<sup>1</sup> Using  $\Lambda_c^+ \rightarrow pK^-\pi^+$  and  $\bar{\Sigma}^- \rightarrow \bar{p}\pi^0$ .

NODE=M071P43;LINKAGE=A

## OTHER DECAYS

NODE=M071320

 $\Gamma(\text{invisible})/\Gamma(e^+e^-)$  $\Gamma_{216}/\Gamma_7$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.0</b>	90	LEES	13I BABR	$B \rightarrow K^{(*)}\psi(2S)$

NODE=M071S64  
NODE=M071S64 $\psi(2S)$  CROSS-PARTICLE BRANCHING RATIOS

NODE=M071240

For measurements involving  $B(\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$  see the corresponding entries in the  $\chi_{cJ}(1P)$  sections.

NODE=M071240

## MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

NODE=M071250

 $\psi(2S) \rightarrow \gamma\chi_{cJ}(1P)$  and  $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$  $a_2(\chi_{c1})/a_2(\chi_{c2})$  Magnetic quadrupole transition amplitude ratioNODE=M071QAR  
NODE=M071QAR

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>63 ± 7 OUR AVERAGE</b>				

61.7 ± 8.3	253k	1 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
------------	------	-----------	----------	---

67 <sup>+19</sup> <sub>-13</sub>	59k	2 ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
----------------------------------	-----	----------	---------	---

<sup>1</sup> Statistical and systematic errors combined.

NODE=M071QAR;LINKAGE=A

<sup>2</sup> Statistical and systematic errors combined. Using values from fits with floating  $M2$  amplitudes  $a_2(\chi_{c1})$ ,  $a_2(\chi_{c2})$ ,  $b_2(\chi_{c1})$ ,  $b_2(\chi_{c2})$  and fixed  $E3$  amplitudes of  $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$ . Not independent of values for  $a_2(\chi_{c1}(1P))$  and  $a_2(\chi_{c2}(1P))$  from ARTUSO 09.

NODE=M071QAR;LINKAGE=AR

**$b_2(\chi_{c2})/b_2(\chi_{c1})$  Magnetic quadrupole transition amplitude ratio**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>60±31 OUR AVERAGE</b>				
74±40	253k	<sup>1</sup> ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
37 <sup>+53</sup> <sub>-47</sub>	59k	<sup>2</sup> ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>1</sup> Statistical and systematic errors combined. Derived from the reported measurement of  $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$ .

<sup>2</sup> Statistical and systematic errors combined. Using values from fits with floating  $M2$  amplitudes  $a_2(\chi_{c1})$ ,  $a_2(\chi_{c2})$ ,  $b_2(\chi_{c1})$ ,  $b_2(\chi_{c2})$  and fixed  $E3$  amplitudes of  $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$ . Not independent of values for  $b_2(\chi_{c1}(1P))$  and  $b_2(\chi_{c2}(1P))$  from ARTUSO 09.

NODE=M071QBR  
NODE=M071QBR

NODE=M071QBR;LINKAGE=A

NODE=M071QBR;LINKAGE=AR

 **$\psi(2S)$  REFERENCES**

ABLIKIM	23	CP C47 013002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61897
ABLIKIM	22AP	PR D106 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61888
ABLIKIM	22AQ	PR D106 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61889
ABLIKIM	22AV	JHEP 2212 016	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61899
ABLIKIM	22AX	PR D106 112002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61901
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61902
ABLIKIM	22AZ	PR D106 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61903
ABLIKIM	21AL	PR D104 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61444
ABLIKIM	21AO	PR D104 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61447
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61463
ABLIKIM	21E	PRL 126 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61033
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61117
ABLIKIM	21S	PL B820 136576	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61152
ABLIKIM	21Z	PRL 127 082002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61265
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61113
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61451
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19AO	PR D99 112010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59892
ABLIKIM	19AT	PR D100 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59989
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60024
ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59605
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59615
ABLIKIM	18Q	PR D97 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58933
ABLIKIM	18T	PR D98 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58975
ABLIKIM	18Z	PL B783 452	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59038
ANASHIN	18	PL B781 174	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=59013
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58315
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58322
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58324
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57903
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57967
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
ABLIKIM	17X	PR D96 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58216
DOBBS	17	PR D96 092004	S. Dobbs <i>et al.</i>	(NWES, WAYN)	REFID=58670
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57966
AAIJ	16Y	JHEP 1605 132	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57333
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57510
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56787
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14G	PR D89 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55898
DOBBS	14	PL B739 90	S. Dobbs <i>et al.</i>	(NWES, WAYN)	REFID=56333
ABLIKIM	13A	PRL 110 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54834
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54920
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55386
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55402
ABLIKIM	13S	PR D88 032010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55403
ABLIKIM	13W	PR D88 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55634
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54269
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54272
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54273
ABLIKIM	12L	PR D86 072011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54739
ABLIKIM	12M	PR D86 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54740
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
ABLIKIM	12Q	CP C36 1040	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54864
ANASHIN	12	PL B711 280	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54038
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)	REFID=53960
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53348
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53630
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
CRONIN-HEN...	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=53233
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=53124

NODE=M071

MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51636
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51725
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=51655
		Translated from ZETFP 85 429.			
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (errat.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)	REFID=52717; ERROR=8
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=51630
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51048
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51560
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT_BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51158
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50757
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50758
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50760
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50451
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50497
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50785
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=50808
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50327
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50328
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49608
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49749
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49186
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48548
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48578
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49171
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)	REFID=48632
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48003
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47939
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46340
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46554
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45416
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41 733.			
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22173
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22207
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
		Translated from YAF 34 1471.			
BRANDELIC	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BRANDELIC	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22197
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22198
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)	REFID=22064
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG	REFID=22194
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)	REFID=22176
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)	REFID=22177
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)	REFID=22179
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)	REFID=22186
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)	REFID=22046
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC	REFID=22188
WIJK	75	Stanford Symp. 69	B.H. Wiik	(DESY)	REFID=22050

$\psi(3770)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M053

 **$\psi(3770)$  MASS (MeV)**

NODE=M053M

OUR FIT includes measurements of  $m_{\psi(2S)}$ ,  $m_{\psi(3770)}$ , and  $m_{\psi(3770)} - m_{\psi(2S)}$ .

NODE=M053M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>3773.7±0.4 OUR FIT</b>				Error includes scale factor of 1.4.
<b>3778.1±0.7 OUR AVERAGE</b>				
3778.1±0.7±0.6		<sup>1</sup> AAIJ	19M LHCB	$pp \rightarrow D\bar{D} + \text{anything}$
3779.2 <sup>+1.8+0.6</sup> <sub>-1.7-0.8</sub>		<sup>2</sup> ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
3775.5±2.4±0.5	57	AUBERT	08B BABR	$B \rightarrow D\bar{D}K$
3776 ±5 ±4	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0\bar{D}^0K^+$
3778.8±1.9±0.9		AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3779.8±0.6		<sup>3</sup> SHAMOV	17 RVUE	$e^+e^- \rightarrow D\bar{D}$ , hadrons
3772.0±1.9		<sup>4,5</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$
3778.4±3.0±1.3	34	CHISTOV	04 BELL	Sup. by BRODZICKA 08

NODE=M053M

<sup>1</sup> Measured in prompt hadroproduction.

<sup>2</sup> Taking into account interference between the resonant and non-resonant  $D\bar{D}$  production.

<sup>3</sup> From the joint analysis of the data on the  $D\bar{D}$  and inclusive hadronic cross sections in the  $\psi(3770)$  region from BaBar, Belle, BES-II, CLEO and KEDR.

<sup>4</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = 0^\circ$ .

<sup>5</sup> Interference between the resonant and non-resonant  $D\bar{D}$  production not taken into account.

NODE=M053M;LINKAGE=B  
 NODE=M053M;LINKAGE=AN  
 NODE=M053M;LINKAGE=A

NODE=M053M;LINKAGE=AB

NODE=M053M;LINKAGE=NI

 **$m_{\psi(3770)} - m_{\psi(2S)}$** 

NODE=M053DM

OUR FIT includes measurements of  $m_{\psi(2S)}$ ,  $m_{\psi(3770)}$ , and  $m_{\psi(3770)} - m_{\psi(2S)}$ .

NODE=M053DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>87.6±0.4 OUR FIT</b>			Error includes scale factor of 1.4.
<b>86.6±0.7 OUR AVERAGE</b>			Error includes scale factor of 2.0. See the ideogram below.
86.9±0.4	<sup>1</sup> ABLIKIM	07E BES2	$e^+e^- \rightarrow \text{hadrons}$
86.7±0.7	ABLIKIM	06L BES2	$e^+e^- \rightarrow \text{hadrons}$
80 ±2	SCHINDLER	80 MRK2	$e^+e^-$
86 ±2	<sup>2</sup> BACINO	78 DLCO	$e^+e^-$
88 ±3	RAPIDIS	77 LGW	$e^+e^-$


NODE=M053DM

<sup>1</sup> BES-II  $\psi(2S)$  mass subtracted (see ABLIKIM 06L).

<sup>2</sup> SPEAR  $\psi(2S)$  mass subtracted (see SCHINDLER 80).

NODE=M053DM;LINKAGE=AK  
 NODE=M053DM;LINKAGE=S





**ψ(3770) WIDTH**

NODE=M053W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
<b>27.2± 1.0 OUR FIT</b>				
<b>27.5± 0.9 OUR AVERAGE</b>				
24.9 <sup>+</sup> 4.6 <sup>+</sup> 0.5 - 4.0 <sup>-</sup> 1.1		1 ANASHIN 12A	KEDR	$e^+e^- \rightarrow D\bar{D}$
30.4± 8.5		2,3 ABLIKIM 08D	BES2	$e^+e^- \rightarrow$ hadrons
27 ±10 ±5	68	BRODZICKA 08	BELL	$B^+ \rightarrow D^0\bar{D}^0 K^+$
28.5± 1.2±0.2		3 ABLIKIM 07E	BES2	$e^+e^- \rightarrow$ hadrons
23.5± 3.7±0.9		AUBERT 07BE	BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
26.9± 2.4±0.3		3 ABLIKIM 06L	BES2	$e^+e^- \rightarrow$ hadrons
24 ± 5		3 SCHINDLER 80	MRK2	$e^+e^-$
24 ± 5		3 BACINO 78	DLCO	$e^+e^-$
28 ± 5		3 RAPIDIS 77	LGW	$e^+e^-$

NODE=M053W

• • • We do not use the following data for averages, fits, limits, etc. • • •

25.8± 1.3		4 SHAMOV 17	RVUE	$e^+e^- \rightarrow D\bar{D},$ hadrons
-----------	--	-------------	------	--

<sup>1</sup> Taking into account interference between the resonant and non-resonant  $D\bar{D}$  production.

<sup>2</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = 0^\circ$ .

<sup>3</sup> Interference between the resonant and non-resonant  $D\bar{D}$  production not taken into account.

<sup>4</sup> From the joint analysis of the data on the  $D\bar{D}$  and inclusive hadronic cross sections in the  $\psi(3770)$  region from BaBar, Belle, BES-II, CLEO and KEDR.

NODE=M053W;LINKAGE=AN

NODE=M053W;LINKAGE=AB

NODE=M053W;LINKAGE=NI

NODE=M053W;LINKAGE=A

**ψ(3770) DECAY MODES**

NODE=M053220;NODE=M053

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $D\bar{D}$	(93 <sup>+8</sup> / <sub>-9</sub> ) %	S=2.0
$\Gamma_2$ $D^0\bar{D}^0$	(52 <sup>+4</sup> / <sub>-5</sub> ) %	S=2.0
$\Gamma_3$ $D^+D^-$	(41 ± 4) %	S=2.0
$\Gamma_4$ $J/\psi X$	( 5.0 ± 2.2 ) × 10 <sup>-3</sup>	
$\Gamma_5$ $J/\psi \pi^+\pi^-$	( 1.93 ± 0.28 ) × 10 <sup>-3</sup>	
$\Gamma_6$ $J/\psi \pi^0\pi^0$	( 8.0 ± 3.0 ) × 10 <sup>-4</sup>	
$\Gamma_7$ $J/\psi \eta$	( 9 ± 4 ) × 10 <sup>-4</sup>	
$\Gamma_8$ $J/\psi \pi^0$	< 2.8 × 10 <sup>-4</sup>	CL=90%
$\Gamma_9$ $e^+e^-$	( 9.6 ± 0.7 ) × 10 <sup>-6</sup>	S=1.3

DESIG=2

DESIG=5

DESIG=6

DESIG=235

DESIG=4

DESIG=46

DESIG=47

DESIG=48

DESIG=1

## Decays to light hadrons

NODE=M053;CLUMP=H

Γ <sub>10</sub>	$b_1(1235)\pi$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=20
Γ <sub>11</sub>	$\phi\eta'$	< 7	$\times 10^{-4}$	CL=90%	DESIG=17
Γ <sub>12</sub>	$\omega\eta'$	< 4	$\times 10^{-4}$	CL=90%	DESIG=16
Γ <sub>13</sub>	$\rho^0\eta'$	< 6	$\times 10^{-4}$	CL=90%	DESIG=15
Γ <sub>14</sub>	$\phi\eta$	( 3.1 ± 0.7 )	$\times 10^{-4}$		DESIG=8
Γ <sub>15</sub>	$\omega\eta$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=14
Γ <sub>16</sub>	$\rho^0\eta$	< 5	$\times 10^{-4}$	CL=90%	DESIG=13
Γ <sub>17</sub>	$\phi\pi^0$	< 3	$\times 10^{-5}$	CL=90%	DESIG=12
Γ <sub>18</sub>	$\omega\pi^0$	< 6	$\times 10^{-4}$	CL=90%	DESIG=11
Γ <sub>19</sub>	$\pi^+\pi^-\pi^0$	< 5	$\times 10^{-6}$	CL=90%	DESIG=9
Γ <sub>20</sub>	$\rho\pi$	< 5	$\times 10^{-6}$	CL=90%	DESIG=10
Γ <sub>21</sub>	$K^+K^-$				DESIG=234
Γ <sub>22</sub>	$K^*(892)^+K^- + \text{c.c.}$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=19
Γ <sub>23</sub>	$K^*(892)^0K^0 + \text{c.c.}$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=18
Γ <sub>24</sub>	$K_S^0K_L^0$	< 1.2	$\times 10^{-5}$	CL=90%	DESIG=3
Γ <sub>25</sub>	$2(\pi^+\pi^-)$	< 1.12	$\times 10^{-3}$	CL=90%	DESIG=21
Γ <sub>26</sub>	$2(\pi^+\pi^-)\pi^0$	< 1.06	$\times 10^{-3}$	CL=90%	DESIG=22
Γ <sub>27</sub>	$2(\pi^+\pi^-\pi^0)$	< 5.85	%	CL=90%	DESIG=208
Γ <sub>28</sub>	$\omega\pi^+\pi^-$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=24
Γ <sub>29</sub>	$3(\pi^+\pi^-)$	< 9.1	$\times 10^{-3}$	CL=90%	DESIG=52
Γ <sub>30</sub>	$3(\pi^+\pi^-)\pi^0$	< 1.37	%	CL=90%	DESIG=55
Γ <sub>31</sub>	$3(\pi^+\pi^-)2\pi^0$	< 11.74	%	CL=90%	DESIG=210
Γ <sub>32</sub>	$\eta\pi^+\pi^-$	< 1.24	$\times 10^{-3}$	CL=90%	DESIG=23
Γ <sub>33</sub>	$\pi^+\pi^-2\pi^0$	< 8.9	$\times 10^{-3}$	CL=90%	DESIG=206
Γ <sub>34</sub>	$\rho^0\pi^+\pi^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=64
Γ <sub>35</sub>	$\eta3\pi$	< 1.34	$\times 10^{-3}$	CL=90%	DESIG=25
Γ <sub>36</sub>	$\eta2(\pi^+\pi^-)$	< 2.43	%	CL=90%	DESIG=53
Γ <sub>37</sub>	$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%	DESIG=221
Γ <sub>38</sub>	$\eta'3\pi$	< 2.44	$\times 10^{-3}$	CL=90%	DESIG=26
Γ <sub>39</sub>	$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%	DESIG=27
Γ <sub>40</sub>	$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=28
Γ <sub>41</sub>	$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%	DESIG=207
Γ <sub>42</sub>	$4(\pi^+\pi^-)$	< 1.67	%	CL=90%	DESIG=62
Γ <sub>43</sub>	$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%	DESIG=63
Γ <sub>44</sub>	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%	DESIG=29
Γ <sub>45</sub>	$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%	DESIG=30
Γ <sub>46</sub>	$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%	DESIG=67
Γ <sub>47</sub>	$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%	DESIG=68
Γ <sub>48</sub>	$\omega K^+K^-$	< 3.4	$\times 10^{-4}$	CL=90%	DESIG=32
Γ <sub>49</sub>	$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%	DESIG=69
Γ <sub>50</sub>	$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%	DESIG=70
Γ <sub>51</sub>	$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%	DESIG=71
Γ <sub>52</sub>	$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%	DESIG=209
Γ <sub>53</sub>	$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%	DESIG=57
Γ <sub>54</sub>	$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%	DESIG=58
Γ <sub>55</sub>	$\eta K^+K^-$	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=31
Γ <sub>56</sub>	$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%	DESIG=222
Γ <sub>57</sub>	$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%	DESIG=65
Γ <sub>58</sub>	$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=33
Γ <sub>59</sub>	$\phi K^+K^-$	< 7.5	$\times 10^{-4}$	CL=90%	DESIG=34
Γ <sub>60</sub>	$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=35
Γ <sub>61</sub>	$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=59
Γ <sub>62</sub>	$K_S^0K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=200
Γ <sub>63</sub>	$K_S^0K^-\pi^+\pi^0$	< 1.33	%	CL=90%	DESIG=201
Γ <sub>64</sub>	$K_S^0K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%	DESIG=214
Γ <sub>65</sub>	$K_S^0K^-2\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%	DESIG=202
Γ <sub>66</sub>	$K_S^0K^-\pi^+\rho^0$	< 1.6	%	CL=90%	DESIG=215

Γ <sub>67</sub>	$K_S^0 K^- \pi^+ \eta$	< 1.3	%	CL=90%	DESIG=216
Γ <sub>68</sub>	$K_S^0 K^- 2\pi^+ \pi^- \pi^0$	< 4.18	%	CL=90%	DESIG=203
Γ <sub>69</sub>	$K_S^0 K^- 2\pi^+ \pi^- \eta$	< 4.8	%	CL=90%	DESIG=217
Γ <sub>70</sub>	$K_S^0 K^- \pi^+ 2(\pi^+ \pi^-)$	< 1.22	%	CL=90%	DESIG=204
Γ <sub>71</sub>	$K_S^0 K^- \pi^+ 2\pi^0$	< 2.65	%	CL=90%	DESIG=205
Γ <sub>72</sub>	$K_S^0 K^- K^+ K^- \pi^+$	< 4.9	$\times 10^{-3}$	CL=90%	DESIG=218
Γ <sub>73</sub>	$K_S^0 K^- K^+ K^- \pi^+ \pi^0$	< 3.0	%	CL=90%	DESIG=219
Γ <sub>74</sub>	$K_S^0 K^- K^+ K^- \pi^+ \eta$	< 2.2	%	CL=90%	DESIG=220
Γ <sub>75</sub>	$K_S^{*0} K^- \pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%	DESIG=60
Γ <sub>76</sub>	$p\bar{p}$				DESIG=233
Γ <sub>77</sub>	$p\bar{p}\pi^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=54
Γ <sub>78</sub>	$p\bar{p}\pi^+\pi^-$	< 5.8	$\times 10^{-4}$	CL=90%	DESIG=36
Γ <sub>79</sub>	$\Lambda\bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%	DESIG=42
Γ <sub>80</sub>	$p\bar{p}\pi^+\pi^-\pi^0$	< 1.85	$\times 10^{-3}$	CL=90%	DESIG=37
Γ <sub>81</sub>	$\omega p\bar{p}$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=39
Γ <sub>82</sub>	$\Lambda\bar{\Lambda}\pi^0$	< 7	$\times 10^{-5}$	CL=90%	DESIG=72
Γ <sub>83</sub>	$p\bar{p}2(\pi^+\pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%	DESIG=61
Γ <sub>84</sub>	$\eta p\bar{p}$	< 5.4	$\times 10^{-4}$	CL=90%	DESIG=38
Γ <sub>85</sub>	$\eta p\bar{p}\pi^+\pi^-$	< 3.3	$\times 10^{-3}$	CL=90%	DESIG=223
Γ <sub>86</sub>	$\rho^0 p\bar{p}$	< 1.7	$\times 10^{-3}$	CL=90%	DESIG=66
Γ <sub>87</sub>	$p\bar{p}K^+K^-$	< 3.2	$\times 10^{-4}$	CL=90%	DESIG=40
Γ <sub>88</sub>	$\eta p\bar{p}K^+K^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=224
Γ <sub>89</sub>	$\pi^0 p\bar{p}K^+K^-$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=225
Γ <sub>90</sub>	$\phi p\bar{p}$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=41
Γ <sub>91</sub>	$\Lambda\bar{\Lambda}\pi^+\pi^-$	< 2.5	$\times 10^{-4}$	CL=90%	DESIG=43
Γ <sub>92</sub>	$\Lambda\bar{p}K^+$	< 2.8	$\times 10^{-4}$	CL=90%	DESIG=44
Γ <sub>93</sub>	$\Lambda\bar{p}K^+\pi^+\pi^-$	< 6.3	$\times 10^{-4}$	CL=90%	DESIG=45
Γ <sub>94</sub>	$\Lambda\bar{\Lambda}\eta$	< 1.9	$\times 10^{-4}$	CL=90%	DESIG=226
Γ <sub>95</sub>	$\Sigma^+\Sigma^-$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=227
Γ <sub>96</sub>	$\Sigma^0\Sigma^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=228
Γ <sub>97</sub>	$\Xi^+\Xi^-$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=229
Γ <sub>98</sub>	$\Xi^0\Xi^0$	< 1.4	$\times 10^{-4}$	CL=90%	DESIG=230
<b>Radiative decays</b>					
Γ <sub>99</sub>	$\gamma\chi_{c2}$	< 6.4	$\times 10^{-4}$	CL=90%	NODE=M053;CLUMP=R DESIG=51
Γ <sub>100</sub>	$\gamma\chi_{c1}$	( 2.49±0.23 )	$\times 10^{-3}$		DESIG=50
Γ <sub>101</sub>	$\gamma\chi_{c0}$	( 6.9 ±0.6 )	$\times 10^{-3}$		DESIG=49
Γ <sub>102</sub>	$\gamma\eta_c$	< 7	$\times 10^{-4}$	CL=90%	DESIG=231
Γ <sub>103</sub>	$\gamma\eta_c(2S)$	< 9	$\times 10^{-4}$	CL=90%	DESIG=232
Γ <sub>104</sub>	$\gamma\eta'$	< 1.8	$\times 10^{-4}$	CL=90%	DESIG=213
Γ <sub>105</sub>	$\gamma\eta$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=212
Γ <sub>106</sub>	$\gamma\pi^0$	< 2	$\times 10^{-4}$	CL=90%	DESIG=211

### CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 23 measurements and one constraint to determine 5 parameters. The overall fit has a  $\chi^2 = 20.1$  for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$ , in percent, from the fit to parameters  $p_i$ , including the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$x_3$	99		
$x_9$	0	0	
$\Gamma$	0	0	-44
	$x_2$	$x_3$	$x_9$

	Mode	Rate (MeV)	Scale factor	
Γ <sub>2</sub>	$D^0\bar{D}^0$	14.0 ±1.4	1.8	DESIG=5
Γ <sub>3</sub>	$D^+D^-$	11.2 ±1.1	1.7	DESIG=6
Γ <sub>9</sub>	$e^+e^-$	( 2.62±0.18 ) $\times 10^{-4}$	1.4	DESIG=1

$\psi(3770)$  PARTIAL WIDTHS

NODE=M053225

 $\Gamma(e^+e^-)$  $\Gamma_9$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.262 ± 0.018</b>	<b>OUR FIT</b>	Error includes scale factor of 1.4.		
<b>0.256 ± 0.016</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.2.		
0.154 <sup>+0.079+0.021</sup> <sub>-0.058-0.027</sub>		1,2 ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
0.22 ± 0.05		3,4 ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
0.277 ± 0.011 ± 0.013		4 ABLIKIM	07E BES2	$e^+e^- \rightarrow$ hadrons
0.203 ± 0.003 <sup>+0.041</sup> <sub>-0.027</sub>	1.4M	4,5 BESSON	06 CLEO	$e^+e^- \rightarrow$ hadrons
0.276 ± 0.050		4 SCHINDLER	80 MRK2	$e^+e^-$
0.18 ± 0.06		4 BACINO	78 DLCO	$e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.196 ± 0.018		6 SHAMOV	17 RVUE	$e^+e^- \rightarrow D\bar{D}$ , hadrons
0.414 <sup>+0.072+0.093</sup> <sub>-0.080-0.028</sub>		2,7 ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
0.37 ± 0.09		8 RAPIDIS	77 LGW	$e^+e^-$

NODE=M053W1  
NODE=M053W1OCCUR=2  
OCCUR=2NODE=M053W1;LINKAGE=A1  
NODE=M053W1;LINKAGE=AN  
NODE=M053W1;LINKAGE=AB

NODE=M053W1;LINKAGE=NI

NODE=M053W1;LINKAGE=BE

NODE=M053W1;LINKAGE=B

NODE=M053W1;LINKAGE=A2  
NODE=M053W1;LINKAGE=R<sup>1</sup> Solution I of the two solutions.<sup>2</sup> Taking into account interference between the resonant and non-resonant  $D\bar{D}$  production.<sup>3</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = 0^\circ$ .<sup>4</sup> Interference between the resonant and non-resonant  $D\bar{D}$  production not taken into account.<sup>5</sup> BESSON 06 (as corrected in BESSON 10) measure  $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow$  hadrons) =  $6.36 \pm 0.08^{+0.41}_{-0.30}$  nb at  $\sqrt{s} = 3773 \pm 1$  MeV, and obtain  $\Gamma_{e^e}$  from the Born-level cross section calculated using  $\psi(3770)$  mass and width from our 2004 edition, PDG 04.<sup>6</sup> From the joint analysis of the data on the  $D\bar{D}$  and inclusive hadronic cross sections in the  $\psi(3770)$  region from BaBar, Belle, BES-II, CLEO and KEDR.<sup>7</sup> Solution II of the two solutions.<sup>8</sup> See also  $\Gamma(e^+e^-)/\Gamma_{\text{total}}$  below. $\psi(3770)$  BRANCHING RATIOS

NODE=M053230

 $\Gamma(D\bar{D})/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma = (\Gamma_2 + \Gamma_3)/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.93<sup>+0.08</sup><sub>-0.09</sub></b>	<b>OUR FIT</b>	Error includes scale factor of 2.0.		
<b>0.93<sup>+0.08</sup><sub>-0.09</sub></b>	<b>OUR AVERAGE</b>	Error includes scale factor of 2.1.		
0.849 ± 0.056 ± 0.018		1 ABLIKIM	08B BES2	$e^+e^- \rightarrow$ non- $D\bar{D}$
1.033 ± 0.014 <sup>+0.048</sup> <sub>-0.066</sub>	1.427M	2 BESSON	06 CLEO	$e^+e^- \rightarrow$ hadrons

NODE=M053R1  
NODE=M053R1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.836 ± 0.049		3 SHAMOV	17 RVUE	$e^+e^- \rightarrow D\bar{D}$ , hadrons
0.866 ± 0.050 ± 0.036		4,5 ABLIKIM	07K BES2	$e^+e^- \rightarrow$ non- $D\bar{D}$
0.836 ± 0.073 ± 0.042		5 ABLIKIM	06L BES2	$e^+e^- \rightarrow D\bar{D}$
0.855 ± 0.017 ± 0.058		5,6 ABLIKIM	06N BES2	$e^+e^- \rightarrow D\bar{D}$

NODE=M053R1;LINKAGE=AI  
NODE=M053R1;LINKAGE=BE

NODE=M053R1;LINKAGE=A

NODE=M053R1;LINKAGE=AL  
NODE=M053R1;LINKAGE=SU

NODE=M053R1;LINKAGE=AB

<sup>1</sup> Neglecting interference.<sup>2</sup> Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of  $D\bar{D}$  reported by CLEO in DOBBS 07.<sup>3</sup> From the joint analysis of the data on the  $D\bar{D}$  and inclusive hadronic cross sections in the  $\psi(3770)$  region from BaBar, Belle, BES-II, CLEO and KEDR.<sup>4</sup> Using  $\sigma^{obs} = 7.07 \pm 0.58$  nb and neglecting interference.<sup>5</sup> Not independent of ABLIKIM 08B.<sup>6</sup> From a measurement of  $\sigma(e^+e^- \rightarrow D\bar{D})$  at  $\sqrt{s} = 3773$  MeV, using the  $\psi(3770)$  resonance parameters measured by ABLIKIM 06L. $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.52<sup>+0.04</sup><sub>-0.05</sub></b>	<b>OUR FIT</b>	Error includes scale factor of 2.0.	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.467 ± 0.047 ± 0.023	ABLIKIM	06L BES2	$e^+e^- \rightarrow D^0\bar{D}^0$
0.499 ± 0.013 ± 0.038	1 ABLIKIM	06N BES2	$e^+e^- \rightarrow D^0\bar{D}^0$

NODE=M053R46  
NODE=M053R46<sup>1</sup> From a measurement of  $\sigma(e^+e^- \rightarrow D\bar{D})$  at  $\sqrt{s} = 3773$  MeV, using the  $\psi(3770)$  resonance parameters measured by ABLIKIM 06L.

NODE=M053R46;LINKAGE=AB

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M053R47  
 NODE=M053R47

**0.41 ± 0.04 OUR FIT** Error includes scale factor of 2.0.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.369 ± 0.037 ± 0.028	ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D^+ D^-$
0.357 ± 0.011 ± 0.034	<sup>1</sup> ABLIKIM	06N	BES2	$e^+ e^- \rightarrow D^+ D^-$

<sup>1</sup> From a measurement of  $\sigma(e^+ e^- \rightarrow D\bar{D})$  at  $\sqrt{s} = 3773$  MeV, using the  $\psi(3770)$  resonance parameters measured by ABLIKIM 06L.

NODE=M053R47;LINKAGE=AB

 $\Gamma(D^0 \bar{D}^0)/\Gamma(D^+ D^-)$  $\Gamma_2/\Gamma_3$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M053R5  
 NODE=M053R5

**1.253 ± 0.016 OUR FIT****1.253 ± 0.016 OUR AVERAGE**

1.252 ± 0.009 ± 0.013	5.3M	BONVICINI	14	CLEO	$e^+ e^- \rightarrow D\bar{D}$
1.39 ± 0.31 ± 0.12		PAKHLOVA	08	BELL	10.6 $e^+ e^- \rightarrow D\bar{D}\gamma$
1.78 ± 0.33 ± 0.24		AUBERT	07BE	BABR	$e^+ e^- \rightarrow D\bar{D}\gamma$
1.27 ± 0.12 ± 0.08		ABLIKIM	06L	BES2	$e^+ e^- \rightarrow D\bar{D}$
2.43 ± 1.50 ± 0.43	34	<sup>1</sup> CHISTOV	04	BELL	$B^+ \rightarrow \psi(3770) K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.258 ± 0.016 ± 0.014		<sup>2</sup> DOBBS	07	CLEO	$e^+ e^- \rightarrow D\bar{D}$
-----------------------	--	--------------------	----	------	--------------------------------

<sup>1</sup> See ADLER 88C for older measurements of this quantity.

<sup>2</sup> Superseded by BONVICINI 14.

NODE=M053R5;LINKAGE=CH  
 NODE=M053R5;LINKAGE=DO

 $\Gamma(J/\psi X)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

NODE=M053P00  
 NODE=M053P00

**0.5 ± 0.2 ± 0.1**

<sup>1</sup> ABLIKIM	21Z	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- X$
----------------------	-----	------	---------------------------------------

<sup>1</sup> From a fit to the  $e^+ e^- \rightarrow J/\psi X$  cross section between 3.645 and 3.891 GeV, with  $\psi(2S)$  and  $\psi(3770)$  masses, total widths and leptonic widths fixed to the values from the PDG 20. An alternative fit with an improved  $\chi^2$ , corresponding to a significance of  $5.3 \sigma$ , uses an additional resonance with a mass of  $3766.2 \pm 3.8 \pm 0.4$  MeV/ $c^2$ , a total width of  $22.2 \pm 5.9 \pm 1.4$  MeV, and  $\Gamma(e \cdot e) \cdot B(J/\psi X) = 79.4 \pm 85.5 \pm 11.7$  eV, possibly compatible with the results of ABLIKIM 08H.

NODE=M053P00;LINKAGE=A

 $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M053R4  
 NODE=M053R4

**1.93 ± 0.28 OUR AVERAGE**

1.89 ± 0.20 ± 0.20	231 ± 33	ADAM	06	CLEO	$e^+ e^- \rightarrow \psi(3770)$
3.4 ± 1.4 ± 0.9	17.8 ± 4.8	BAI	05	BES2	$e^+ e^- \rightarrow \psi(3770)$

 $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M053R7  
 NODE=M053R7

**0.080 ± 0.025 ± 0.016** 39 ± 14 ADAM 06 CLEO  $e^+ e^- \rightarrow \psi(3770)$  $\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M053R8  
 NODE=M053R8

**87 ± 33 ± 22** 22 ± 10 ADAM 06 CLEO  $e^+ e^- \rightarrow \psi(3770)$  $\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M053R9  
 NODE=M053R9

**<28** 90 <10 ADAM 06 CLEO  $e^+ e^- \rightarrow \psi(3770)$  $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M053R2  
 NODE=M053R2

**0.96 ± 0.07 OUR FIT** Error includes scale factor of 1.3.**1.3 ± 0.2** RAPIDIS 77 LGW  $e^+ e^-$ 

## ————— DECAYS TO LIGHT HADRONS —————

NODE=M053250

 $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M053R82  
 NODE=M053R82

**<1.4** 90 <sup>1</sup> ADAMS 06 CLEO  $e^+ e^- \rightarrow \psi(3770)$ 

<sup>1</sup> Comparing cross sections at  $\sqrt{s} = 3.773$  GeV and  $\sqrt{s} = 3.671$  GeV, neglecting interference, and using  $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$  nb.

NODE=M053R82;LINKAGE=AD

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$					$\Gamma_{11}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<7	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R83 NODE=M053R83
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R83;LINKAGE=AD
$\Gamma(\omega\eta')/\Gamma_{\text{total}}$					$\Gamma_{12}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<4	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R84 NODE=M053R84
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R84;LINKAGE=AD
$\Gamma(\rho^0\eta')/\Gamma_{\text{total}}$					$\Gamma_{13}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<6	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R85 NODE=M053R85
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R85;LINKAGE=AD
$\Gamma(\phi\eta)/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<b>3.1±0.6±0.3</b>		<sup>1</sup> ADAMS	06	CLEO $3.773 e^+e^- \rightarrow \phi\eta$	NODE=M053R6 NODE=M053R6
••• We do not use the following data for averages, fits, limits, etc. •••					
<19	90	<sup>2</sup> ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$	
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R6;LINKAGE=AD
<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.					NODE=M053R6;LINKAGE=AK
$\Gamma(\omega\eta)/\Gamma_{\text{total}}$					$\Gamma_{15}/\Gamma$
VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<1.4	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R86 NODE=M053R86
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R86;LINKAGE=AD
$\Gamma(\rho^0\eta)/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<5	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R87 NODE=M053R87
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R87;LINKAGE=AD
$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{17}/\Gamma$
VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
< 3	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R11 NODE=M053R11
••• We do not use the following data for averages, fits, limits, etc. •••					
<50	90	<sup>2</sup> ABLIKIM	07B	BES2 $e^+e^- \rightarrow \psi(3770)$	
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R11;LINKAGE=AD
<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.					NODE=M053R11;LINKAGE=AK
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{18}/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<6	90	<sup>1</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R88 NODE=M053R88
<sup>1</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R88;LINKAGE=AD
$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{19}/\Gamma$
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	
<5	90	<sup>1,2</sup> ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R89 NODE=M053R89
<sup>1</sup> Data suggest possible destructive interference with continuum.					NODE=M053R89;LINKAGE=AD
<sup>2</sup> Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.					NODE=M053R89;LINKAGE=AS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	1,2 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

NODE=M053R90  
 NODE=M053R90

<sup>1</sup> Comparing cross sections at  $\sqrt{s} = 3.773$  GeV and  $\sqrt{s} = 3.671$  GeV, neglecting interference, and using  $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$  nb.

NODE=M053R90;LINKAGE=AD

<sup>2</sup> Data suggest possible destructive interference with continuum.

NODE=M053R90;LINKAGE=AS

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$\sim 10^{-5}$	1 DRUZHININ	15	RVUE $e^+e^- \rightarrow \psi(3770)$

NODE=M053R00  
 NODE=M053R00

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes  $e^+e^- \rightarrow K^+K^-$  and  $e^+e^- \rightarrow K_S^0 K_L^0$ .

NODE=M053R00;LINKAGE=A

 $\Gamma(K^*(892)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	1 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

NODE=M053R91  
 NODE=M053R91

<sup>1</sup> Comparing cross sections at  $\sqrt{s} = 3.773$  GeV and  $\sqrt{s} = 3.671$  GeV, neglecting interference, and using  $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$  nb.

NODE=M053R91;LINKAGE=AS

 $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	1 ADAMS	06	CLEO $e^+e^- \rightarrow \psi(3770)$

NODE=M053R92  
 NODE=M053R92

<sup>1</sup> Comparing cross sections at  $\sqrt{s} = 3.773$  GeV and  $\sqrt{s} = 3.671$  GeV, neglecting interference, and using  $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$  nb.

NODE=M053R92;LINKAGE=AD

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	1 CRONIN-HEN..06	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R3  
 NODE=M053R3

••• We do not use the following data for averages, fits, limits, etc. •••

<21 90 <sup>2</sup> ABLIKIM 04F BES  $e^+e^- \rightarrow \psi(3770)$

<sup>1</sup> Using  $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08_{-0.30}^{+0.41})$  nb from BESSON 06 and  $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895 \pm 0.0014$ .

NODE=M053R3;LINKAGE=CR

<sup>2</sup> Using  $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$ .

NODE=M053R3;LINKAGE=AB

 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<11.2	90	1 HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

NODE=M053R21  
 NODE=M053R21

••• We do not use the following data for averages, fits, limits, etc. •••

<48 90 <sup>2</sup> ABLIKIM 07B BES2  $e^+e^- \rightarrow \psi(3770)$

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R21;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R21;LINKAGE=AK

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<10.6	90	1 HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$

NODE=M053R22  
 NODE=M053R22

••• We do not use the following data for averages, fits, limits, etc. •••

<62 90 <sup>2</sup> ABLIKIM 07B BES2  $e^+e^- \rightarrow \psi(3770)$

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R22;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R22;LINKAGE=AK

 $\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<58.5	90	305	ABLIKIM	08N	BES2 $e^+e^- \rightarrow \psi(3770)$

NODE=M053R72  
 NODE=M053R72

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{28}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 6.0	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R24  
 NODE=M053R24

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	<sup>2</sup> ABLIKIM 07I	BES2	3.77 $e^+e^-$
-----	----	--------------------------	------	---------------

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R24;LINKAGE=HU  
 NODE=M053R24;LINKAGE=AK

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{29}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<91	90	<sup>1</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R07  
 NODE=M053R07

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R07;LINKAGE=AK

 $\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$  $\Gamma_{30}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<137	90	<sup>1</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R10  
 NODE=M053R10

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R10;LINKAGE=AK

 $\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<117.4	90	59	ABLIKIM 08N	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R74  
 NODE=M053R74

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{32}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.24	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R23  
 NODE=M053R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.3	90	<sup>2</sup> ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$
------	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R23;LINKAGE=HU  
 NODE=M053R23;LINKAGE=AK

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{33}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8.9	90	218	ABLIKIM 08N	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R70  
 NODE=M053R70

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{34}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	<sup>1</sup> ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R53  
 NODE=M053R53

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R53;LINKAGE=AK

 $\Gamma(\eta3\pi)/\Gamma_{\text{total}}$  $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<13.4	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R25  
 NODE=M053R25

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R25;LINKAGE=HU

 $\Gamma(\eta2(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{36}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<243	90	<sup>1</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R08  
 NODE=M053R08

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R08;LINKAGE=AK

 $\Gamma(\eta\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{37}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.45	90	<sup>1</sup> ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R77  
 NODE=M053R77

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R77;LINKAGE=AK



$\Gamma(\eta'/3\pi)/\Gamma_{\text{total}}$  $\Gamma_{38}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<24.4	90	<sup>1</sup> HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R26  
 NODE=M053R26

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R26;LINKAGE=HU

 $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{39}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 9.0	90	<sup>1</sup> HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R27  
 NODE=M053R27

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48	90	<sup>2</sup> ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R27;LINKAGE=HU  
 NODE=M053R27;LINKAGE=AK

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{40}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	<sup>1</sup> HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R28  
 NODE=M053R28

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	<sup>2</sup> ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R28;LINKAGE=HU  
 NODE=M053R28;LINKAGE=AK

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

 $\Gamma(K^+K^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{41}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4.2	90	14	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R71  
 NODE=M053R71

 $\Gamma(4(\pi^+\pi^-))/\Gamma_{\text{total}}$  $\Gamma_{42}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<16.7	90	<sup>1</sup> ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R50  
 NODE=M053R50

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R50;LINKAGE=AK

 $\Gamma(4(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{43}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<30.6	90	<sup>1</sup> ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R52  
 NODE=M053R52

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R52;LINKAGE=AK

 $\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$  $\Gamma_{44}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	<sup>1</sup> HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R29  
 NODE=M053R29

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R29;LINKAGE=HU

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{45}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 23.6	90	<sup>1</sup> HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R30  
 NODE=M053R30

• • • We do not use the following data for averages, fits, limits, etc. • • •

<111	90	<sup>2</sup> ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R30;LINKAGE=HU  
 NODE=M053R30;LINKAGE=AK

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

 $\Gamma(K^+K^-\rho^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{46}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	<sup>1</sup> ABLIKIM	07I BES2	$3.77 e^+e^-$

NODE=M053R58  
 NODE=M053R58

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R58;LINKAGE=AK

$\Gamma(K^+ K^- \rho^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<146	90	<sup>1</sup> ABLIKIM	07I	BES2 3.77 $e^+ e^-$

NODE=M053R59  
NODE=M053R59

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R59;LINKAGE=AK

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 3.4	90	<sup>1</sup> HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

NODE=M053R32  
NODE=M053R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	<sup>2</sup> ABLIKIM	07I	BES2 3.77 $e^+ e^-$
-----	----	----------------------	-----	---------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R32;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R32;LINKAGE=AK

 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<38	90	<sup>1</sup> ABLIKIM	07I	BES2 3.77 $e^+ e^-$

NODE=M053R60  
NODE=M053R60

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R60;LINKAGE=AK

 $\Gamma(K^{*0} K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<162	90	<sup>1</sup> ABLIKIM	07I	BES2 3.77 $e^+ e^-$

NODE=M053R61  
NODE=M053R61

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R61;LINKAGE=AK

 $\Gamma(K^{*+} K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<323	90	<sup>1</sup> ABLIKIM	07I	BES2 3.77 $e^+ e^-$

NODE=M053R62  
NODE=M053R62

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R62;LINKAGE=AK

 $\Gamma(K^+ K^- \pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{52}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.7	90	24	ABLIKIM	08N	BES2 $e^+ e^- \rightarrow \psi(3770)$

NODE=M053R73  
NODE=M053R73

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<10.3	90	<sup>1</sup> ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

NODE=M053R57  
NODE=M053R57

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R57;LINKAGE=AK

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<36.0	90	<sup>1</sup> ABLIKIM	07F	BES2 $e^+ e^- \rightarrow \psi(3770)$

NODE=M053R51  
NODE=M053R51

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R51;LINKAGE=AK

 $\Gamma(\eta K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	<sup>1</sup> HUANG	06A	CLEO $e^+ e^- \rightarrow \psi(3770)$

NODE=M053R31  
NODE=M053R31

• • • We do not use the following data for averages, fits, limits, etc. • • •

<31	90	<sup>2</sup> ABLIKIM	10D	BES2 $e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------------	-----	---------------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R31;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R31;LINKAGE=AK

$\Gamma(\eta K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.24	90	<sup>1</sup> ABLIKIM	10D BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R78  
NODE=M053R78

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R78;LINKAGE=AK

 $\Gamma(\rho^0 K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	<sup>1</sup> ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R54  
NODE=M053R54

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R54;LINKAGE=AK

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 6.0	90	<sup>1</sup> HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R33  
NODE=M053R33

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17	90	<sup>2</sup> ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	----------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R33;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R33;LINKAGE=AK

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 7.5	90	<sup>1</sup> HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R34  
NODE=M053R34

• • • We do not use the following data for averages, fits, limits, etc. • • •

<24	90	<sup>2</sup> ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	----------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R34;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R34;LINKAGE=AK

 $\Gamma(2(K^+ K^- \pi^0))/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	<sup>1</sup> HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R35  
NODE=M053R35

• • • We do not use the following data for averages, fits, limits, etc. • • •

<46	90	<sup>2</sup> ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	----------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R35;LINKAGE=HU

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R35;LINKAGE=AK

 $\Gamma(2(K^+ K^- \pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	<sup>1</sup> ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R48  
NODE=M053R48

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R48;LINKAGE=AK

 $\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.2	90	18	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R64  
NODE=M053R64

 $\Gamma(K_S^0 K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<13.3	90	40	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R65  
NODE=M053R65

 $\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R15  
NODE=M053R15

 $\Gamma(K_S^0 K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8.7	90	39	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R66  
NODE=M053R66

$\Gamma(K_S^0 K^- \pi^+ \rho^0)/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R16  
NODE=M053R16

 $\Gamma(K_S^0 K^- \pi^+ \eta)/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R17  
NODE=M053R17

 $\Gamma(K_S^0 K^- 2\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<41.8	90	23	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R67  
NODE=M053R67

 $\Gamma(K_S^0 K^- 2\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.8	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R18  
NODE=M053R18

 $\Gamma(K_S^0 K^- \pi^+ 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<12.2	90	4	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R68  
NODE=M053R68

 $\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.5	90	17	ABLIKIM	08M BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R69  
NODE=M053R69

 $\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.9	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R19  
NODE=M053R19

 $\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R75  
NODE=M053R75

 $\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	ABLIKIM	09C BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R76  
NODE=M053R76

 $\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<9.7	90	<sup>1</sup> ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R55  
NODE=M053R55

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$ .

NODE=M053R55;LINKAGE=AK

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		<sup>1</sup> AAIJ	17AD LHCb	$pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$
$7.1^+_{-2.9}$	684	<sup>2</sup> ABLIKIM	14L BES3	$e^+ e^- \rightarrow \psi(3770)$
310 ± 30	684	<sup>3</sup> ABLIKIM	14L BES3	$e^+ e^- \rightarrow \psi(3770)$

OCCUR=2

<sup>1</sup> AAIJ 17AD reports  $B(B^+ \rightarrow \psi(3770)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) < 0.09$  (0.10) at 90% (95%) CL.

NODE=M053R98;LINKAGE=C

<sup>2</sup> Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

NODE=M053R98;LINKAGE=A

<sup>3</sup> Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

NODE=M053R98;LINKAGE=B

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 0.4	90	<sup>1,2</sup> ABLIKIM	14O BES3	$e^+ e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$59^{+3}_{-2} \pm 5$		<sup>1,3</sup> ABLIKIM	14O BES3	$e^+ e^- \rightarrow \psi(3770)$
<12	90	<sup>4</sup> ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

OCCUR=2

NODE=M053R09  
NODE=M053R09

- <sup>1</sup> Calculated by the authors using  $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30}$  nb from BESSON 10.
- <sup>2</sup> Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.
- <sup>3</sup> Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.
- <sup>4</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R09;LINKAGE=A

NODE=M053R09;LINKAGE=B

NODE=M053R09;LINKAGE=C

NODE=M053R09;LINKAGE=AK

 $\Gamma(\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{total}$  $\Gamma_{78}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 5.8	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	<sup>2</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.NODE=M053R36  
NODE=M053R36

NODE=M053R36;LINKAGE=HU

NODE=M053R36;LINKAGE=AK

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{total}$  $\Gamma_{79}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 $\times 10^{-4}$	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.8 $\times 10^{-4}$	90	<sup>2</sup> ABLIKIM 21AS	BES3	$e^+e^- \rightarrow \psi(3770)$
-----------------------	----	---------------------------	------	---------------------------------

<4 $\times 10^{-4}$	90	<sup>3</sup> ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$
---------------------	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.<sup>2</sup> From a measurement of the  $e^+e^- \rightarrow \Lambda\bar{\Lambda}$  cross section between 3.5 and 4.6 GeV. At a 90% CL the lower bound is  $> 2.4 \times 10^{-6}$ .<sup>3</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R42

NODE=M053R42

NODE=M053R42;LINKAGE=HU

NODE=M053R42;LINKAGE=A

NODE=M053R42;LINKAGE=AK

 $\Gamma(\rho\bar{\rho}\pi^+\pi^-\pi^0)/\Gamma_{total}$  $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<18.5	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<73	90	<sup>2</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R37

NODE=M053R37

NODE=M053R37;LINKAGE=HU

NODE=M053R37;LINKAGE=AK

 $\Gamma(\omega\rho\bar{\rho})/\Gamma_{total}$  $\Gamma_{81}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	90	<sup>2</sup> ABLIKIM 07I	BES2	3.77 $e^+e^-$
-----	----	--------------------------	------	---------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.<sup>2</sup> Using  $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$  nb and neglecting interference.

NODE=M053R39

NODE=M053R39

NODE=M053R39;LINKAGE=HU

NODE=M053R39;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{total}$  $\Gamma_{82}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 0.7	90	<sup>1</sup> ABLIKIM 13Q	BES3	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	<sup>2</sup> ABLIKIM 07I	BES2	3.77 $e^+e^-$
-----	----	--------------------------	------	---------------

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected.<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R63

NODE=M053R63

NODE=M053R63;LINKAGE=A

NODE=M053R63;LINKAGE=AK

 $\Gamma(\rho\bar{\rho}2(\pi^+\pi^-))/\Gamma_{total}$  $\Gamma_{83}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	<sup>1</sup> ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R49

NODE=M053R49

NODE=M053R49;LINKAGE=AK

$\Gamma(\eta\rho\bar{p})/\Gamma_{\text{total}}$  $\Gamma_{84}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 5.4	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R38  
 NODE=M053R38

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11	90	<sup>2</sup> ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R38;LINKAGE=HU  
 NODE=M053R38;LINKAGE=AK

 $\Gamma(\eta\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{85}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	<sup>1</sup> ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R79  
 NODE=M053R79

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R79;LINKAGE=AK

 $\Gamma(\rho^0\rho\bar{p})/\Gamma_{\text{total}}$  $\Gamma_{86}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	<sup>1</sup> ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R56  
 NODE=M053R56

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R56;LINKAGE=AK

 $\Gamma(\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R40  
 NODE=M053R40

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11	90	<sup>2</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R40;LINKAGE=HU  
 NODE=M053R40;LINKAGE=AK

 $\Gamma(\eta\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	<sup>1</sup> ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R80  
 NODE=M053R80

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R80;LINKAGE=AK

 $\Gamma(\pi^0\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	<sup>1</sup> ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R81  
 NODE=M053R81

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R81;LINKAGE=AK

 $\Gamma(\phi\rho\bar{p})/\Gamma_{\text{total}}$  $\Gamma_{90}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R41  
 NODE=M053R41

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9	90	<sup>2</sup> ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
----	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R41;LINKAGE=HU  
 NODE=M053R41;LINKAGE=AK

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{91}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	<sup>1</sup> HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R43  
 NODE=M053R43

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.7	90	<sup>2</sup> ABLIKIM 13Q	BES3	$e^+e^- \rightarrow \psi(3770)$
-------	----	--------------------------	------	---------------------------------

<39	90	<sup>3</sup> ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

<sup>1</sup> Using  $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

<sup>2</sup> Assuming that interference effects between resonance and continuum can be neglected.

<sup>3</sup> Assuming that interference effects between resonance and continuum can be neglected and using  $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$  nb.

NODE=M053R43;LINKAGE=HU  
 NODE=M053R43;LINKAGE=A  
 NODE=M053R43;LINKAGE=AK

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{92}/\Gamma$
<2.8	90	<sup>1</sup> HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R44 NODE=M053R44

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R44;LINKAGE=HU

 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{93}/\Gamma$
<6.3	90	<sup>1</sup> HUANG	06A	CLEO $e^+e^- \rightarrow \psi(3770)$	NODE=M053R45 NODE=M053R45

<sup>1</sup> Using  $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$  nb at the resonance.

NODE=M053R45;LINKAGE=HU

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{94}/\Gamma$
<1.9	90	<sup>1</sup> ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R93 NODE=M053R93

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R93;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{95}/\Gamma$
<1.0	90	<sup>1</sup> ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R94 NODE=M053R94

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R94;LINKAGE=A

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{96}/\Gamma$
<0.4	90	<sup>1</sup> ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R95 NODE=M053R95

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R95;LINKAGE=A

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{97}/\Gamma$
<1.5	90	<sup>1</sup> ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R96 NODE=M053R96

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R96;LINKAGE=A

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{98}/\Gamma$
<1.4	90	<sup>1</sup> ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(3770)$	NODE=M053R97 NODE=M053R97

<sup>1</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R97;LINKAGE=A

————— RADIATIVE DECAYS —————

 $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{99}/\Gamma$
<0.64	90	<sup>1</sup> ABLIKIM	15J	BES3 $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	NODE=M053R03 NODE=M053R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0                      90                      <sup>2</sup> BRIERE                      06                      CLEO                       $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$

<0.9                      90                      <sup>3</sup> COAN                      06A                      CLEO                       $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> This limit is equivalent to  $(0.25 \pm 0.21 \pm 0.18) \times 10^{-3}$  branching fraction value.

<sup>2</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$  from ATHAR 04,  $\psi(2S)$  mass and width from PDG 04, and  $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$  keV from ADAM 06.

<sup>3</sup> Using  $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$  keV from ADAM 06 and taking  $\sigma(e^+e^- \rightarrow D\bar{D})$  from HE 05 for  $\sigma(e^+e^- \rightarrow \psi(3770))$ .

NODE=M053R03;LINKAGE=A

NODE=M053R03;LINKAGE=BR

NODE=M053R03;LINKAGE=CO

NODE=M053240

 $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{100}/\Gamma$
<b>2.49±0.23 OUR AVERAGE</b>					

1.98±0.78±0.05                      202                      <sup>1</sup> ABLIKIM                      16B                      BES3                       $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$

2.48±0.15±0.23                      0.6k                      ABLIKIM                      15J                      BES3                       $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

2.4 ± 0.8 ± 0.2                      <sup>2</sup> ABLIKIM                      14H                      BES3                       $e^+e^- \rightarrow \psi(3770) \rightarrow K_S^0 K^\pm \pi^\mp$

2.9 ± 0.5 ± 0.4                      <sup>3</sup> BRIERE                      06                      CLEO                       $e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma\gamma J/\psi$

NODE=M053R02

NODE=M053R02

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.9 ±1.4 ±0.6	54	<sup>4</sup> BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$
2.8 ±0.5 ±0.4	53	<sup>5</sup> COAN	06A	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$

<sup>1</sup> ABLIKIM 16B reports  $(1.94 \pm 0.42 \pm 0.64) \times 10^{-3}$  from a measurement of  $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M053R02;LINKAGE=A

<sup>2</sup> ABLIKIM 14H reports  $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)] = (8.51 \pm 2.39 \pm 1.42) \times 10^{-6}$  which we divide by our best value  $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp) = 0.00349 \pm 0.00029$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. We have calculated the best value of  $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)$  as 1/2 of  $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$ .

NODE=M053R02;LINKAGE=AB

<sup>3</sup> Averages the two measurements from COAN 06A and BRIERE 06.

NODE=M053R02;LINKAGE=BI  
NODE=M053R02;LINKAGE=BR

<sup>4</sup> Uses  $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$  from ATHAR 04,  $\psi(2S)$  mass and width from PDG 04, and  $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$  keV from ADAM 06.

<sup>5</sup> Using  $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$  keV from ADAM 06 and taking  $\sigma(e^+e^- \rightarrow D\bar{D})$  from HE 05 for  $\sigma(e^+e^- \rightarrow \psi(3770))$ .

NODE=M053R02;LINKAGE=CO

$\Gamma(\gamma\chi_{c1})/\Gamma(J/\psi\pi^+\pi^-)$		$\Gamma_{100}/\Gamma_5$				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>1.49±0.31±0.26</b>	53 ± 10	<sup>1</sup> COAN	06A	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	NODE=M053R04 NODE=M053R04

<sup>1</sup> Using  $B(\psi(3770) \rightarrow J/\psi\pi^+\pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$  from ADAM 06.

NODE=M053R04;LINKAGE=CO

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$		$\Gamma_{101}/\Gamma$				
VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>6.9±0.6 OUR AVERAGE</b>						NODE=M053R01 NODE=M053R01
6.7±0.7±0.1	2.2k	<sup>1</sup> ABLIKIM	16B	BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
7.3±0.7±0.6	274	BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
•••		We do not use the following data for averages, fits, limits, etc. •••				
< 44	90	<sup>2</sup> COAN	06A	CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

<sup>1</sup> ABLIKIM 16B reports  $(6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$  from a measurement of  $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c0})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$  assuming  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$ , which we rescale to our best value  $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M053R01;LINKAGE=B

<sup>2</sup> Using  $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$  keV from ADAM 06 and taking  $\sigma(e^+e^- \rightarrow D\bar{D})$  from HE 05 for  $\sigma(e^+e^- \rightarrow \psi(3770))$ .

NODE=M053R01;LINKAGE=CO

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$		$\Gamma_{101}/\Gamma_{99}$				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
•••		We do not use the following data for averages, fits, limits, etc. •••				NODE=M053R06 NODE=M053R06
>8	90	<sup>1</sup> BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	
		<sup>1</sup> Not independent of other results in BRIERE 06.				NODE=M053R06;LINKAGE=BR

NODE=M053R06;LINKAGE=BR

$\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$		$\Gamma_{101}/\Gamma_{100}$				
VALUE		DOCUMENT ID	TECN	COMMENT		
•••		We do not use the following data for averages, fits, limits, etc. •••				NODE=M053R05 NODE=M053R05
2.5±0.6		<sup>1</sup> BRIERE	06	CLEO	$e^+e^- \rightarrow \psi(3770)$	
		<sup>1</sup> Not independent of other results in BRIERE 06.				NODE=M053R05;LINKAGE=BR

NODE=M053R05;LINKAGE=BR

$\Gamma(\gamma\eta_c)/\Gamma_{\text{total}}$		$\Gamma_{102}/\Gamma$				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;7 × 10<sup>-4</sup></b>	90	<sup>1</sup> ABLIKIM	14H	BES3		NODE=M053R99 NODE=M053R99
		<sup>1</sup> ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 16 \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = 2.34 \times 10^{-2}$ . We have calculated the best value of $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = 7.0 \times 10^{-2}$ .				NODE=M053R99;LINKAGE=AB

NODE=M053R99;LINKAGE=AB



$\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN
$<9 \times 10^{-4}$	90	<sup>1</sup> ABLIKIM	14H BES3

<sup>1</sup> ABLIKIM 14H reports  $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 5.6 \times 10^{-6}$  which we divide by our best value  $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) = 6 \times 10^{-3}$ . We have calculated the best value of  $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)$  as 1/3 of  $B(\eta_c(2S) \rightarrow K \bar{K} \pi) = 1.9 \times 10^{-2}$ .

 $\Gamma_{103}/\Gamma$ NODE=M053R20  
NODE=M053R20

NODE=M053R20;LINKAGE=AB

 $\Gamma(\gamma\eta')/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8$	90	<sup>1</sup> PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$

<sup>1</sup> Assuming maximal destructive interference between  $\psi(3770)$  and continuum sources.

NODE=M053R14  
NODE=M053R14

NODE=M053R14;LINKAGE=PE

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5$	90	<sup>1</sup> PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$

<sup>1</sup> Assuming maximal destructive interference between  $\psi(3770)$  and continuum sources.

NODE=M053R13  
NODE=M053R13

NODE=M053R13;LINKAGE=PE

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<2$	90	PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$

NODE=M053R12  
NODE=M053R12 $\psi(3770)$  REFERENCES

NODE=M053

ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21Z	PRL 127 082002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61265
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59697
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
SHAMOV	17	PL B769 187	A.G. Shamov, K.Yu. Todyshev		REFID=57900
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57126
ABLIKIM	15J	PR D91 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56775
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)	REFID=56962
ABLIKIM	14H	PR D89 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55899
ABLIKIM	14L	PL B735 101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55903
ABLIKIM	14O	PR D90 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55906
BONVICINI	14	PR D89 072002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=55798
ABLIKIM	13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55393
ANASHIN	12A	PL B711 292	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54055
ABLIKIM	10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53350
BESSION	10	PRL 104 159901 (errata.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
ABLIKIM	09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=53134
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
ABLIKIM	08H	PRL 101 102004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52254
ABLIKIM	08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52569
ABLIKIM	08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52570
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)	REFID=52144
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
ABLIKIM	07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51704
ABLIKIM	07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51882
ABLIKIM	07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51940
ABLIKIM	07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52045
ABLIKIM	07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52073
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52075
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51131
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
ADAMS	06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50990
BESSION	06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51041
Also		PRL 104 159901 (errata.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
BRIERE	06	PR D74 031106	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=51149
COAN	06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51155
CRONIN-HEN...	06	PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=51156
HUANG	06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=50999
BAI	05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50332
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=50924
Also		PRL 96 199903 (errata.)	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51211
ABLIKIM	04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50185
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40361
SCHINDLER	80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)	REFID=22222
BACINO	78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)	REFID=11437
RAPIDIS	77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)	REFID=22220

$\psi_2(3823)$ 
 $I^G(J^{PC}) = 0^-(2^{--})$   
 $I, J, P$  need confirmation.
was  $\psi(3823)$ ,  $X(3823)$ 

Seen by BHARDWAJ 13 in  $B \rightarrow \chi_{c1} \gamma K$  and ABLIKIM 15S in  $e^+ e^- \rightarrow \pi^+ \pi^- \gamma \chi_{c1}$  decays as a narrow peak in the invariant mass distribution of the  $\chi_{c1} \gamma$  system. Properties consistent with the  $\psi_2(1^3D_2) c\bar{c}$  state.

NODE=M212

NODE=M212

 **$\psi_2(3823)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3823.5 ± 0.5 OUR AVERAGE</b>				Error includes scale factor of 1.4. [3823.7 ± 0.5 MeV OUR 2022 AVERAGE Scale factor = 1.1]

3823.12 ± 0.43 ± 0.13	120	ABLIKIM	22R	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
3824.08 ± 0.53 ± 0.14	137	<sup>1</sup> AAIJ	20S	LHCB $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
3823.1 ± 1.8 ± 0.7	33 ± 10	<sup>2</sup> BHARDWAJ	13	BELL $B^\pm \rightarrow \chi_{c1} \gamma K^\pm$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3821.7 ± 1.3 ± 0.7	19 ± 5	<sup>3</sup> ABLIKIM	15S	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
--------------------	--------	----------------------	-----	---

<sup>1</sup> Using the measured  $m_{\psi_2(3823)} - m_{\psi(2S)} = 137.98 \pm 0.53 \pm 0.14$  MeV.<sup>2</sup> From a simultaneous fit to  $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$  and  $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$  with significance  $4.0\sigma$  including systematics. Corrected for the measured  $\psi(2S)$  mass using  $B \rightarrow \psi(2S) K \rightarrow (\gamma \chi_{c1}) K$  decays.<sup>3</sup> From a simultaneous unbinned maximum likelihood fit of  $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$  data (the  $\pi^+ \pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both  $\psi(2S) \rightarrow \chi_{c1} \gamma$  and  $\psi_2(3823) \rightarrow \chi_{c1} \gamma$  together, with floating mass scale offset for  $\psi(2S)$ , floating  $\psi_2(3823)$  mass, and zero  $\psi_2(3823)$  width, resulting in a significance of  $5.9\sigma$  when including systematic uncertainties. Superseded by ABLIKIM 22R.

NODE=M212M

NODE=M212M

NEW

NODE=M212M;LINKAGE=C

NODE=M212M;LINKAGE=A

NODE=M212M;LINKAGE=B

 **$m_{\psi_2(3823)} - m_{\psi(2S)}$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
137.98 ± 0.53 ± 0.14	137	<sup>1</sup> AAIJ	20S	LHCB $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$

<sup>1</sup> AAIJ 20S also reports  $m_{\chi_{c1}(3872)} - m_{\psi_2(3823)} = 47.50 \pm 0.53 \pm 0.13$  MeV.

NODE=M212A00

NODE=M212A00

NODE=M212A00;LINKAGE=A

 **$\psi_2(3823)$  WIDTH**

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>&lt; 2.9 (CL = 90%)</b>					[<5.2 MeV (CL = 90%) OUR 2022 BEST LIMIT]

< 2.9	90	120	ABLIKIM	22R	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
-------	----	-----	---------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.2	90		<sup>1</sup> AAIJ	20S	LHCB $B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
< 16	90		<sup>2</sup> ABLIKIM	15S	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
< 24	90		<sup>3</sup> BHARDWAJ	13	BELL $B^\pm \rightarrow \chi_{c1} \gamma K^\pm$

<sup>1</sup> AAIJ 20S also provides a limit of < 6.6 MeV with 95% CL.<sup>2</sup> From a fit of  $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$  data (the  $\pi^+ \pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.<sup>3</sup> From a simultaneous fit to  $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$  and  $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$  with significance  $4.0\sigma$  including systematics.

NODE=M212W

NODE=M212W

NODE=M212W;LINKAGE=C

NODE=M212W;LINKAGE=B

NODE=M212W;LINKAGE=A

 **$\psi_2(3823)$  DECAY MODES**Branching fractions are given relative to the one **DEFINED AS 1**.

NODE=M212215;NODE=M212

NODE=M212

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $J/\psi(1S) \pi^+ \pi^-$	<0.06	90%
$\Gamma_2$ $J/\psi(1S) \pi^0 \pi^0$	<0.11	90%
$\Gamma_3$ $J/\psi(1S) \pi^0$	<0.030	90%
$\Gamma_4$ $J/\psi(1S) \eta$	<0.14	90%
$\Gamma_5$ $\chi_{c0} \gamma$	<0.24	90%
$\Gamma_6$ $\chi_{c1} \gamma$	<b>DEFINED AS 1</b>	DESIG=1
$\Gamma_7$ $\chi_{c2} \gamma$	0.28 $\begin{smallmatrix} +0.14 \\ -0.11 \end{smallmatrix}$	DESIG=2

$\psi_2(3823)$  BRANCHING RATIOS $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		<sup>1</sup> ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	137 ± 26	AAIJ	20S	LHCB	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212225

NODE=M212R00  
NODE=M212R00

NODE=M212R00;LINKAGE=A

 $\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(\chi_{c1}\gamma)$   $\Gamma_1/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<0.06** 90 <sup>1</sup> ABLIKIM 210 BES3  $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R06  
NODE=M212R06

NODE=M212R06;LINKAGE=A

 $\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(\chi_{c1}\gamma)$   $\Gamma_2/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<0.11** 90 <sup>1</sup> ABLIKIM 210 BES3  $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R07  
NODE=M212R07

NODE=M212R07;LINKAGE=A

 $\Gamma(J/\psi(1S)\pi^0)/\Gamma(\chi_{c1}\gamma)$   $\Gamma_3/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<0.03** 90 <sup>1</sup> ABLIKIM 210 BES3  $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R08  
NODE=M212R08

NODE=M212R08;LINKAGE=A

 $\Gamma(J/\psi(1S)\eta)/\Gamma(\chi_{c1}\gamma)$   $\Gamma_4/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**<0.14** 90 <sup>1</sup> ABLIKIM 210 BES3  $e^+e^- \rightarrow \pi^+\pi^-X$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R09  
NODE=M212R09

NODE=M212R09;LINKAGE=A

 $\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$   $\Gamma_4/\Gamma_1$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**$4.4^{+2.5}_{-1.9} \pm 0.9$**  <sup>1</sup> AAIJ 22D LHCB  $B^+ \rightarrow J/\psi(1S)\eta K^+$

<sup>1</sup> Using the branching ratio for  $B^+ \rightarrow \psi_2(3823)K^+$  with  $\psi_2(3823) \rightarrow J/\psi(1S)\pi^+\pi^-$  from AAIJ 20S.

NODE=M212R10  
NODE=M212R10

NODE=M212R10;LINKAGE=A

 $\Gamma(\chi_{c0}\gamma)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		<sup>1</sup> ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
----------	--	----------------------	-----	------	----------------------------------

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R04  
NODE=M212R04

NODE=M212R04;LINKAGE=A

 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	120	<sup>1</sup> ABLIKIM	22R	BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
seen	63 ± 9	<sup>2</sup> ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	16 ± 5	<sup>3</sup> ABLIKIM	210	BES3	$e^+e^- \rightarrow \pi^0\pi^0X$
seen	33 ± 10	<sup>4</sup> BHARDWAJ	13	BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak  $R$  with mass  $4417.5 \pm 26.2 \pm 3.5$  MeV and width  $245 \pm 48 \pm 13$  MeV, which leads to  $\Gamma(e^+e^-)B(R \rightarrow \pi^+\pi^-\psi_2(3823))B(\psi_2(3823) \rightarrow \chi_{c1}\gamma) = 0.57 \pm 0.08$  eV.

<sup>2</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ . Signal has a 11.8  $\sigma$  significance.

<sup>3</sup> From a fit of the invariant  $\pi^0\pi^0$  recoil-mass distribution. Signal has a 4.3  $\sigma$  significance.

<sup>4</sup> BHARDWAJ 13 reports  $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$  with statistical significance 3.8  $\sigma$ .

NODE=M212R01  
NODE=M212R01

OCCUR=2

NODE=M212R01;LINKAGE=D

NODE=M212R01;LINKAGE=B

NODE=M212R01;LINKAGE=C

NODE=M212R01;LINKAGE=A

$\Gamma(\chi_{c0}\gamma)/\Gamma(\chi_{c1}\gamma)$  $\Gamma_5/\Gamma_6$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.24</b>	90	<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M212R05  
 NODE=M212R05

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R05;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M212R02  
 NODE=M212R02

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
not seen	<sup>2</sup> ABLIKIM	15s BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
not seen	<sup>3</sup> BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2}\gamma K^\pm$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ . Signal has a 3.2  $\sigma$  significance.

NODE=M212R02;LINKAGE=C

<sup>2</sup> From a simultaneous unbinned maximum likelihood fit of  $e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$  data (the  $\pi^+\pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both  $\psi(2S) \rightarrow \chi_{c2}\gamma$  and  $\psi_2(3823) \rightarrow \chi_{c2}\gamma$  together, with floating mass scale offset for  $\psi(2S)$ ,  $\psi_2(3823)$  mass floating (fixed to that above), and zero  $\psi_2(3823)$  width.

NODE=M212R02;LINKAGE=B

<sup>3</sup> BHARDWAJ 13 reports  $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c2}) < 3.6 \times 10^{-6}$  at 90% CL.

NODE=M212R02;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$  $\Gamma_7/\Gamma_6$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>0.28^{+0.14}_{-0.11} \pm 0.02</math></b>	9 $\pm$ 4		<sup>1</sup> ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$

NODE=M212R03  
 NODE=M212R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.42	90	<sup>2</sup> ABLIKIM	15s BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
<0.41	90	BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1/c2}\gamma K^\pm$

<sup>1</sup> From a simultaneous unbinned maximum likelihood fit of the  $\pi^+\pi^-$  recoil mass distributions of seven decay channels in the process  $e^+e^- \rightarrow \pi^+\pi^-X$ .

NODE=M212R03;LINKAGE=B

<sup>2</sup> From a simultaneous unbinned maximum likelihood fit of  $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1(2)}\gamma$  data (the  $\pi^+\pi^-$  recoil mass) taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both  $\psi(2S) \rightarrow \chi_{c1(2)}\gamma$  and  $\psi_2(3823) \rightarrow \chi_{c1(2)}\gamma$  together, with floating mass scale offset for  $\psi(2S)$ ,  $\psi_2(3823)$  mass floating (fixed to that above), and zero  $\psi_2(3823)$  width.

NODE=M212R03;LINKAGE=A

 $\psi_2(3823)$  REFERENCES

NODE=M212

AAIJ	22D	JHEP 2204 046	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61647
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61664
ABLIKIM	210	PR D103 L091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61121
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60526
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56784
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=55412

$\psi_3(3842)$ 
 $I^G(J^{PC}) = 0^-(3^{--})$   
 $J, P$  need confirmation.

 $J^P$  has not been measured,  $3^-$  is the quark model prediction.

NODE=M241

NODE=M241

NODE=M241M

NODE=M241M

 $\psi_3(3842)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>3842.71 ± 0.16 ± 0.12</b>	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$

NODE=M241W

NODE=M241W

 $\psi_3(3842)$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>2.79 ± 0.51 ± 0.35</b>	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$

NODE=M241215;NODE=M241

 $\psi_3(3842)$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^+ D^-$	seen
$\Gamma_2$ $D^0 \bar{D}^0$	seen

DESIG=1

DESIG=2

 $\psi_3(3842)$  BRANCHING RATIOS

NODE=M241225

 $\Gamma(D^+ D^-)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$

NODE=M241R01  
NODE=M241R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen <sup>1</sup> ABLIKIM 22AL BES3  $e^+ e^- \rightarrow \pi^+ \pi^- D^+ D^-$

<sup>1</sup> From a fit to the  $\pi^+ \pi^-$  recoil mass for  $e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$ .

NODE=M241R01;LINKAGE=A

 $\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$

NODE=M241R02  
NODE=M241R02 $\psi_3(3842)$  REFERENCES

NODE=M241

ABLIKIM	22AL PR D106 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	19M JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=61884  
REFID=59697

$\chi_{c0}(3860)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

NODE=M237

## OMITTED FROM SUMMARY TABLE

The assignment  $J^P = 0^+$  is preferred over  $2^+$  by 2.5 sigma.

NODE=M237

Observed by CHILIKIN 17 using full amplitude analysis of the process  $e^+ e^- \rightarrow J/\psi D \bar{D}$ , where  $D = D^0, D^+$ . Not seen by AAIJ 20AI in the decay  $B^+ \rightarrow D^+ D^- K^+$ .

 $\chi_{c0}(3860)$  MASS

NODE=M237M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$3862^{+26+40}_{-32-13}$	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D \bar{D}$

NODE=M237M

 $\chi_{c0}(3860)$  WIDTH

NODE=M237W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$201^{+154+88}_{-67-82}$	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D \bar{D}$

NODE=M237W

 $\chi_{c0}(3860)$  DECAY MODES

NODE=M237215;NODE=M237

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad D^0 \bar{D}^0$	seen
$\Gamma_2 \quad D^+ D^-$	seen

DESIG=1

DESIG=2

 $\chi_{c0}(3860)$  BRANCHING RATIOS

NODE=M237220

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
seen	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D^0 \bar{D}^0$	

NODE=M237R00  
NODE=M237R00

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
seen	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D^+ D^-$	

NODE=M237R01  
NODE=M237R01 $\chi_{c0}(3860)$  REFERENCES

NODE=M237

AAIJ	20AI	PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)
CHILIKIN	17	PR D95 112003	K. Chilikin <i>et al.</i>	(BELLE Collab.) JPC

REFID=60739  
REFID=57995

$\chi_{c1}(3872)$ 

$$I^G(J^{PC}) = 0^+(1^{++})$$

also known as  $X(3872)$ 

This state shows properties different from a conventional  $q\bar{q}$  state. A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

First observed by CHOI 03 in  $B \rightarrow K\pi^+\pi^- J/\psi(1S)$  decays as a narrow peak in the invariant mass distribution of the  $\pi^+\pi^- J/\psi(1S)$  final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in  $B^+ \rightarrow \chi_{c1}(3872)K^+$  decays, where  $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$  and  $J/\psi \rightarrow \mu^+\mu^-$ , which unambiguously gives the  $J^{PC} = 1^{++}$  assignment under the assumption that the  $\pi^+\pi^-$  and  $J/\psi$  are in an  $S$ -wave. AAIJ 15AO extend this analysis with more data to limit  $D$ -wave contributions to  $< 4\%$  at 95% CL.

See the review on "Spectroscopy of Mesons Containing Two Heavy Quarks."

 **$\chi_{c1}(3872)$  MASS FROM  $J/\psi X$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3871.65 ± 0.06 OUR AVERAGE</b>				
3871.64 ± 0.06 ± 0.01	19.8k	<sup>1</sup> AAIJ	20S LHC	$B^+ \rightarrow J/\psi\pi^+\pi^- K^+$
3871.9 ± 0.7 ± 0.2	20	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
3871.95 ± 0.48 ± 0.12	0.6k	AAIJ	12H LHC	$pp \rightarrow J/\psi\pi^+\pi^- X$
3871.85 ± 0.27 ± 0.19	170	<sup>2</sup> CHOI	11 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3873 ± 1.8 ± 1.3	27	<sup>3</sup> DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61 ± 0.16 ± 0.19	6k	<sup>3,4</sup> AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3871.4 ± 0.6 ± 0.1	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3868.7 ± 1.5 ± 0.4	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.8 ± 3.1 ± 3.0	522	<sup>3,5</sup> ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3871.695 ± 0.067 ± 0.068	15.6k	<sup>6</sup> AAIJ	20AD LHC	$pp \rightarrow J/\psi\pi^+\pi^- X$
3871.59 ± 0.06 ± 0.03	4.2k	<sup>7</sup> AAIJ	20S LHC	$B^+ \rightarrow J/\psi\pi^+\pi^- K^+$
3873.3 ± 1.1 ± 1.0	45	<sup>8</sup> ABLIKIM	19V BES	$e^+e^- \rightarrow \gamma\omega J/\psi$
3860.0 ± 10.4	13.6	<sup>3,9</sup> AGHASYAN	18A COMP	$\gamma^* N \rightarrow X\pi^\pm N'$
3868.6 ± 1.2 ± 0.2	8	<sup>10</sup> AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.3 ± 0.6 ± 0.1	61	<sup>10</sup> AUBERT	06 BABR	$B^- \rightarrow K^- J/\psi\pi^+\pi^-$
3873.4 ± 1.4	25	<sup>11</sup> AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3871.3 ± 0.7 ± 0.4	730	<sup>3,12</sup> ACOSTA	04 CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3872.0 ± 0.6 ± 0.5	36	<sup>13</sup> CHOI	03 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3836 ± 13	58	<sup>3,14</sup> ANTONIAZZI	94 E705	$300 \pi^\pm L_i \rightarrow J/\psi\pi^+\pi^- X$

<sup>1</sup> Calculated from  $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.54 \pm 0.06$  MeV obtained by combining the data with  $\chi_{c1}(3872)$  produced in  $B^+$  decays from AAIJ 20S and inclusive  $b$ -hadron decays from AAIJ 20AD and using  $m_{\psi(2S)} = 3686.097$  MeV. Breit-Wigner parametrization.

<sup>2</sup> The mass difference for the  $\chi_{c1}(3872)$  produced in  $B^+$  and  $B^0$  decays is  $(-0.71 \pm 0.96 \pm 0.19)$  MeV.

<sup>3</sup> Width consistent with detector resolution.

<sup>4</sup> A possible equal mixture of two states with a mass difference greater than 3.6 MeV/ $c^2$  is excluded at 95% CL.

<sup>5</sup> Calculated from the corresponding  $m_{\chi_{c1}(3872)} - m_{J/\psi}$  using  $m_{J/\psi} = 3096.916$  MeV.

NODE=M176

NODE=M176

NODE=M176M

NODE=M176M

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M176M;LINKAGE=F

NODE=M176M;LINKAGE=CO

NODE=M176M;LINKAGE=AC

NODE=M176M;LINKAGE=AA

NODE=M176M;LINKAGE=AB

- <sup>6</sup> Using  $\chi_{c1}(3872)$  produced in inclusive  $b$ -hadron decays and  $m_{\psi(2S)} = 3686.097 \pm 0.010$  MeV. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.
- <sup>7</sup> Using Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.
- <sup>8</sup> Fit with fixed width and including two resonances,  $\chi_{c0}(3915)$  and  $X(3960)$ .
- <sup>9</sup> Could be a different state.
- <sup>10</sup> Calculated from the corresponding  $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$  using  $m_{\psi(2S)} = 3686.093$  MeV. Superseded by AUBERT 08Y.
- <sup>11</sup> Calculated from the corresponding  $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$  using  $m_{\psi(2S)} = 3685.96$  MeV. Superseded by AUBERT 06.
- <sup>12</sup> Superseded by AALTONEN 09AU.
- <sup>13</sup> Superseded by CHOI 11.
- <sup>14</sup> A lower mass value can be due to an incorrect momentum scale for soft pions.

NODE=M176M;LINKAGE=D

NODE=M176M;LINKAGE=E

NODE=M176M;LINKAGE=B

NODE=M176M;LINKAGE=A

NODE=M176M;LINKAGE=AE

NODE=M176M;LINKAGE=AU

NODE=M176M;LINKAGE=AT

NODE=M176M;LINKAGE=CH

NODE=M176M;LINKAGE=AN

### $\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3872.9^{+0.6+0.4}_{-0.4-0.5}$	50	<sup>1,2</sup> AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
$3875.1^{+0.7}_{-0.5} \pm 0.5$	$33 \pm 6$	<sup>2</sup> AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$
$3875.2 \pm 0.7^{+0.9}_{-1.8}$	$24 \pm 6$	<sup>2,3</sup> GOKHROO	06	BELL $B \rightarrow D^0 \bar{D}^0 \pi^0 K$

- <sup>1</sup> Calculated from the measured  $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{\bar{D}^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$  MeV.
- <sup>2</sup> Experiments report  $D^{*0} \bar{D}^0$  invariant mass above  $D^{*0} \bar{D}^0$  threshold because  $D^{*0}$  decay products are kinematically constrained to the  $D^{*0}$  mass, even though the  $D^{*0}$  may decay off-shell.
- <sup>3</sup> Superseded by AUSHEV 10.

NODE=M176MD0

NODE=M176MD0

NODE=M176MD0;LINKAGE=AS

NODE=M176MD0;LINKAGE=AU

NODE=M176MD0;LINKAGE=GO

### $m_{\chi_{c1}(3872)} - m_{J/\psi}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>774.9 \pm 3.1 \pm 3.0</math></b>	522	ABAZOV	04F D0	$\rho \bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

### $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$185.598 \pm 0.067 \pm 0.068$	15.6k	<sup>1</sup> AAIJ	20AD LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
$185.54 \pm 0.06$	19.8k	<sup>2</sup> AAIJ	20S LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
$187.4 \pm 1.4$	25	<sup>3</sup> AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

- <sup>1</sup> Using  $\chi_{c1}(3872)$  produced in inclusive  $b$ -hadron decays. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.
- <sup>2</sup> Combining  $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.49 \pm 0.06 \pm 0.03$  MeV from AAIJ 20S and the measured mass difference from AAIJ 20AD. Breit-Wigner parametrization.
- <sup>3</sup> Superseded by AUBERT 06.

NODE=M176DM

NODE=M176DM

NODE=M176DM2

NODE=M176DM2

NODE=M176DM2;LINKAGE=A

NODE=M176DM2;LINKAGE=E

NODE=M176DM2;LINKAGE=AU

### $\chi_{c1}(3872)$ WIDTH

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>1.19 \pm 0.21</math> OUR AVERAGE</b>		Error includes scale factor of 1.1.		
$1.39 \pm 0.24 \pm 0.10$	15.6k	<sup>1</sup> AAIJ	20AD LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
$0.96^{+0.19}_{-0.18} \pm 0.21$	4.2k	<sup>2</sup> AAIJ	20S LHCB	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$

- • • We do not use the following data for averages, fits, limits, etc. • • •**
- <2.4 90 ABLIKIM 14 BES3  $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
- <1.2 90 CHOI 11 BELL  $B \rightarrow K \pi^+ \pi^- J/\psi$
- <3.3 90 AUBERT 08Y BABR  $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
- <4.1 90 69 AUBERT 06 BABR  $B \rightarrow K \pi^+ \pi^- J/\psi$
- <2.3 90 36 <sup>3</sup> CHOI 03 BELL  $B \rightarrow K \pi^+ \pi^- J/\psi$

- <sup>1</sup> Using  $\chi_{c1}(3872)$  produced in inclusive  $b$ -hadron decays. Breit-Wigner parametrization.
- <sup>2</sup> Using Breit-Wigner parametrization. Partially overlapping dataset with that of AAIJ 20AD.
- <sup>3</sup> Superseded by CHOI 11.

NODE=M176W

NODE=M176W

OCCUR=3

OCCUR=2

NODE=M176W;LINKAGE=E

NODE=M176W;LINKAGE=F

NODE=M176W;LINKAGE=CH



**$\chi_{c1}(3872)$  WIDTH FROM  $\bar{D}^{*0} D^0$  MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.9^{+2.8+0.2}_{-1.4-1.1}$	50	<sup>1</sup> AUSHEV	10	BELL $B \rightarrow \bar{D}^{*0} D^0 K$
$3.0^{+1.9}_{-1.4} \pm 0.9$	$33 \pm 6$	AUBERT	08B	BABR $B \rightarrow \bar{D}^{*0} D^0 K$

<sup>1</sup>With a measured value of  $B(B \rightarrow \chi_{c1}(3872) K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} D^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$ , assumed to be equal for both charged and neutral modes.

NODE=M176WD0  
NODE=M176WD0

NODE=M176WD0;LINKAGE=AU

 **$\chi_{c1}(3872)$  DECAY MODES**

NODE=M176215;NODE=M176

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $e^+ e^-$	$< 2.8 \times 10^{-6}$	90%
$\Gamma_2$ $\pi^+ \pi^- \pi^0$	$< 9 \times 10^{-3}$	90%
$\Gamma_3$ $\pi^+ \pi^- J/\psi(1S)$	$(3.8 \pm 1.2) \%$	
$\Gamma_4$ $\pi^+ \pi^- \pi^0 J/\psi(1S)$	not seen	
$\Gamma_5$ $\omega \eta_c(1S)$	$< 33 \%$	90%
$\Gamma_6$ $\omega J/\psi(1S)$	$(4.3 \pm 2.1) \%$	
$\Gamma_7$ $\phi \phi$	not seen	
$\Gamma_8$ $D^0 \bar{D}^0 \pi^0$	$(49^{+18}_{-20}) \%$	
$\Gamma_9$ $\bar{D}^{*0} D^0$	$(37 \pm 9) \%$	
$\Gamma_{10}$ $\gamma \gamma$	$< 11 \%$	90%
$\Gamma_{11}$ $D^0 \bar{D}^0$	$< 29 \%$	90%
$\Gamma_{12}$ $D^+ D^-$	$< 19 \%$	90%
$\Gamma_{13}$ $\pi^0 \chi_{c2}$	$< 4 \%$	90%
$\Gamma_{14}$ $\pi^0 \chi_{c1}$	$(3.4 \pm 1.6) \%$	
$\Gamma_{15}$ $\pi^0 \chi_{c0}$	$< 14 \%$	90%
$\Gamma_{16}$ $\pi^+ \pi^- \eta_c(1S)$	$< 14 \%$	90%
$\Gamma_{17}$ $\pi^0 \pi^0 \chi_{c0}$	$< 7 \%$	90%
$\Gamma_{18}$ $\pi^+ \pi^- \chi_{c0}$	$< 2.1 \%$	90%
$\Gamma_{19}$ $\pi^+ \pi^- \chi_{c1}$	$< 7 \times 10^{-3}$	90%
$\Gamma_{20}$ $p \bar{p}$	$< 2.4 \times 10^{-5}$	95%

DESIG=1  
DESIG=29  
DESIG=2  
DESIG=25  
DESIG=24  
DESIG=13  
DESIG=26  
DESIG=8  
DESIG=12  
DESIG=5  
DESIG=6  
DESIG=7  
DESIG=20  
DESIG=18  
DESIG=19  
DESIG=14  
DESIG=28  
DESIG=27  
DESIG=17  
DESIG=16

**Radiative decays**

$\Gamma_{21}$ $\gamma D^+ D^-$	$< 4 \%$	90%
$\Gamma_{22}$ $\gamma \bar{D}^0 D^0$	$< 6 \%$	90%
$\Gamma_{23}$ $\gamma J/\psi$	$(8 \pm 4) \times 10^{-3}$	
$\Gamma_{24}$ $\gamma \chi_{c1}$	$< 9 \times 10^{-3}$	90%
$\Gamma_{25}$ $\gamma \chi_{c2}$	$< 3.2 \%$	90%
$\Gamma_{26}$ $\gamma \psi(2S)$	$(4.5 \pm 2.0) \%$	

NODE=M176;CLUMP=B  
DESIG=21  
DESIG=23  
DESIG=9  
DESIG=3  
DESIG=15  
DESIG=11

**C-violating decays**

$\Gamma_{27}$ $\eta J/\psi$	$< 1.8 \%$	90%
-----------------------------	------------	-----

NODE=M176;CLUMP=A  
DESIG=4

 **$\chi_{c1}(3872)$  PARTIAL WIDTHS**

NODE=M176220

 **$\Gamma(e^+ e^-)$**  **$\Gamma_1$** 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 4.3$	90	<sup>1</sup> ABLIKIM	15V	BES3 $4.0-4.4 e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$
$< 280$	90	<sup>2</sup> YUAN	04	RVUE $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$

<sup>1</sup>ABLIKIM 15V reports this limit from the measurement of  $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+ e^-) / \Gamma < 0.13$  eV using  $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) / \Gamma = 3\%$ .

<sup>2</sup>Using BAI 98E data on  $e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$ . Assuming that  $\Gamma(\pi^+ \pi^- J/\psi)$  of  $\chi_{c1}(3872)$  is the same as that of  $\psi(2S)$  (85.4 keV).

NODE=M176W1  
NODE=M176W1

NODE=M176W1;LINKAGE=B

NODE=M176W1;LINKAGE=A

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.9</b>	90	1,2 ADACHI	23 BELL	$B^+ \rightarrow \chi_{c1}(3872)K^+$
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.4	90	2,3 ADACHI	23 BELL	$B^0 \rightarrow \chi_{c1}(3872)K^0$
1 ADACHI 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .				
2 Assuming the decay products, $\pi^+\pi^-\pi^0$ , are uniformly distributed in phase space. The limit is the 90% "credible" upper limit (i.e. Bayesian).				
3 ADACHI 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \chi_{c1}(3872)K^0)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^0 \rightarrow \chi_{c1}(3872)K^0) = 1.1 \times 10^{-4}$ .				

NODE=M176R30  
NODE=M176R30

OCCUR=2

NODE=M176R30;LINKAGE=A

NODE=M176R30;LINKAGE=E

NODE=M176R30;LINKAGE=D

 $\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M176230

 $\Gamma(\pi^+\pi^-J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$   $\Gamma_3\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 0.13</b>	90	ABLIKIM	15V BES3	$4.0-4.4 e^+e^- \rightarrow \pi^+\pi^-J/\psi$
••• We do not use the following data for averages, fits, limits, etc. •••				
< 6.2	90	1,2 AUBERT	05D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
< 8.3	90	2 DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
<10	90	3 YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
1 Using $B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.				
2 Assuming $\chi_{c1}(3872)$ has $J^{PC} = 1^{--}$ .				
3 Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$ . From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$ .				

NODE=M176G1  
NODE=M176G1

NODE=M176G1;LINKAGE=AU

NODE=M176G1;LINKAGE=DO

NODE=M176G1;LINKAGE=A

 $\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M176232

 $\Gamma(\pi^+\pi^-J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_3\Gamma_{10}/\Gamma$ 

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
$5.5^{+4.1}_{-3.8} \pm 0.7$		3	1 TERAMOTO	21 BELL	$e^+e^- \rightarrow \gamma^*\gamma$ at $\Upsilon(nS)$
<12.9	90		2 DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi\gamma$
1 Measured in single-tag two-photon production assuming $Q^2$ dependence of a $c\bar{c}$ meson model. Here, $\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)$ is the reduced two-photon decay width, $\bar{\Gamma}_{\gamma\gamma}$ .					
2 Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.					

NODE=M176H1  
NODE=M176H1

NODE=M176H1;LINKAGE=A

NODE=M176H1;LINKAGE=DO

 $\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_6\Gamma_{10}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.7	90	1 LEES	12AD BABR	$e^+e^- \rightarrow e^+e^-\omega J/\psi$
1 Assuming $\chi_{c1}(3872)$ has spin 2.				

NODE=M176G01  
NODE=M176G01

NODE=M176G01;LINKAGE=LE

 $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_{16}\Gamma_{10}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;11.1</b>	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

NODE=M176G02  
NODE=M176G02 $\chi_{c1}(3872)$  BRANCHING RATIOS

NODE=M176235

 $\Gamma(\pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.038±0.012 OUR AVERAGE</b>				
0.038±0.002±0.012		1 AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
0.041±0.005±0.013		2 CHOI	11 BELL	$B^+ \rightarrow \pi^+\pi^-J/\psi K^+$
0.040±0.008±0.013	93	3,4 AUBERT	08Y BABR	$B \rightarrow \chi_{c1}(3872)K$
••• We do not use the following data for averages, fits, limits, etc. •••				
seen	151	5 BALA	15 BELL	$B \rightarrow \chi_{c1}(3872)K\pi$
0.061±0.020±0.020	30	6 AUBERT	05R BABR	$B^+ \rightarrow K^+\pi^+\pi^-J/\psi$
0.065±0.014±0.021	36	7 CHOI	03 BELL	$B^+ \rightarrow K^+\pi^+\pi^-J/\psi$

NODE=M176R6  
NODE=M176R6

SYCLP=A

SYCLP=A

SYCLP=A

SYCLP=A

- <sup>1</sup>AAIJ 20s reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (7.95 \pm 0.15 \pm 0.33) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M176R6;LINKAGE=E
- <sup>2</sup>CHOI 11 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.63 \pm 0.82 \pm 0.52) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M176R6;LINKAGE=F
- <sup>3</sup>AUBERT 08Y reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M176R6;LINKAGE=AB
- <sup>4</sup>superseded by LEES 20C NODE=M176R6;LINKAGE=C
- <sup>5</sup>BALA 15 reports  $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872)K^+\pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$  and  $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872)K^0\pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$ . NODE=M176R6;LINKAGE=A
- <sup>6</sup>Superseded by AUBERT 08Y. AUBERT 05R reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.28 \pm 0.41) \times 10^{-5}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M176R6;LINKAGE=AE
- <sup>7</sup>CHOI 03 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] / [B(B^+ \rightarrow \psi(2S)K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = 0.063 \pm 0.012 \pm 0.007$  which we multiply or divide by our best values  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ ,  $B(B^+ \rightarrow \psi(2S)K^+) = (6.24 \pm 0.20) \times 10^{-4}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values. NODE=M176R6;LINKAGE=CH

 **$\Gamma(\pi^+\pi^-\pi^0J/\psi(1S))/\Gamma_{\text{total}}$**  **$\Gamma_4/\Gamma$** 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> WANG	11B	BELL $\Upsilon(2S) \rightarrow \gamma X$
not seen	<sup>2</sup> SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$

NODE=M176R25  
NODE=M176R25

- <sup>1</sup>WANG 11B reports  $B(\Upsilon(2S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0J/\psi) < 2.4 \times 10^{-6}$  at 95% CL. NODE=M176R25;LINKAGE=B
- <sup>2</sup>SHEN 10A reports  $B(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0J/\psi) < 2.8 \times 10^{-6}$  at 95% CL. NODE=M176R25;LINKAGE=A

 **$\Gamma(\omega\eta_c(1S))/\Gamma_{\text{total}}$**  **$\Gamma_5/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.33</b>	90	<sup>1</sup> VINOKUROVA 15	BELL	$B^+ \rightarrow \omega\eta_c K^+$

NODE=M176R24  
NODE=M176R24

- <sup>1</sup>VINOKUROVA 15 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \omega\eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.9 \times 10^{-5}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ . NODE=M176R24;LINKAGE=A

 **$\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$**  **$\Gamma_6/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M176R14  
NODE=M176R14

●●● We do not use the following data for averages, fits, limits, etc. ●●●

0.029±0.011±0.009 21±7 <sup>1</sup>DEL-AMO-SA..10B BABR  $B^+ \rightarrow \omega J/\psi K^+$ 

- <sup>1</sup>DEL-AMO-SANCHEZ 10B reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. DEL-AMO-SANCHEZ 10B also reports  $B(B^0 \rightarrow \chi_{c1}(3872)K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi\omega) = (6 \pm 3 \pm 1) \times 10^{-6}$ . NODE=M176R14;LINKAGE=DE

 **$\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+\pi^-J/\psi(1S))$**  **$\Gamma_6/\Gamma_3$** 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.1±0.4 OUR AVERAGE</b>	Error includes scale factor of 1.7.		

NODE=M176R15  
NODE=M176R15

- 1.6<sup>+0.4</sup><sub>-0.3</sub>±0.2 <sup>1</sup>ABLIKIM 19V BES  $e^+e^- \rightarrow \gamma\omega J/\psi$
- 0.8±0.3 <sup>2</sup>DEL-AMO-SA..10B BABR  $B \rightarrow \omega J/\psi K$

<sup>1</sup>Fit with fixed width and including two resonances,  $\chi_{c0}(3915)$  and  $X(3960)$ .<sup>2</sup>Statistical and systematic errors added in quadrature. Uses the values of  $B(B \rightarrow \chi_{c1}(3872)K) \times B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-)$  reported in AUBERT 08Y, taking into account the common systematics.NODE=M176R15;LINKAGE=A  
NODE=M176R15;LINKAGE=DE

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> AAIJ	17BB LHCB	$pp$ at 7, 8 TeV

NODE=M176R26  
 NODE=M176R26

<sup>1</sup> AAIJ 17BB reports  $B(b \rightarrow \chi_{c1}(3872)\text{anything}) \times B(\chi_{c1}(3872) \rightarrow \phi\phi) < 4.5 \times 10^{-7}$  at 95% CL.

NODE=M176R26;LINKAGE=A

 $\Gamma(D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$0.49^{+0.18}_{-0.20} \pm 0.16$		17	<sup>1</sup> GOKHROO	06	BELL $B^+ \rightarrow D^0\bar{D}^0\pi^0 K^+$

NODE=M176R12  
 NODE=M176R12

••• We do not use the following data for averages, fits, limits, etc. •••

<0.29 90 <sup>2</sup> CHISTOV 04 BELL Sup. by GOKHROO 06

<sup>1</sup> GOKHROO 06 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R12;LINKAGE=GO

<sup>2</sup> CHISTOV 04 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 0.6 \times 10^{-4}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .

NODE=M176R12;LINKAGE=A

 $\Gamma(D^0\bar{D}^0\pi^0)/\Gamma(\pi^+\pi^-J/\psi(1S))$  $\Gamma_8/\Gamma_3$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.16	90	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R17  
 NODE=M176R17

••• We do not use the following data for averages, fits, limits, etc. •••

 $\Gamma(\bar{D}^{*0}D^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.37 \pm 0.09 \pm 0.12$	$41^{+9}_{-8}$	<sup>1</sup> AUSHEV	10	BELL $B^+ \rightarrow D^{*0}\bar{D}^0 K^+$

NODE=M176R13  
 NODE=M176R13  
 SYCLP=A

••• We do not use the following data for averages, fits, limits, etc. •••

0.80 ± 0.28 ± 0.26 27 ± 6 <sup>2</sup> AUBERT 08B BABR  $B^+ \rightarrow \bar{D}^{*0}D^0 K^+$

SYCLP=A

<sup>1</sup> AUSHEV 10 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R13;LINKAGE=AS

<sup>2</sup> AUBERT 08B reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0}D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R13;LINKAGE=AU

 $\Gamma(\bar{D}^{*0}D^0)/\Gamma(\pi^+\pi^-J/\psi(1S))$  $\Gamma_9/\Gamma_3$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$11.77 \pm 3.09$	50	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R16  
 NODE=M176R16

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.11	90	<sup>1</sup> WICHT	08	BELL $e^+e^- \rightarrow \Upsilon(4S)$

NODE=M176R09  
 NODE=M176R09

<sup>1</sup> WICHT 08 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 2.4 \times 10^{-5}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .

NODE=M176R09;LINKAGE=A

 $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.29	90	<sup>1</sup> CHISTOV	04	BELL $B \rightarrow K D^0\bar{D}^0$

NODE=M176R3  
 NODE=M176R3

<sup>1</sup> CHISTOV 04 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow D^0\bar{D}^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6 \times 10^{-5}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .

NODE=M176R3;LINKAGE=A

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.19	90	<sup>1</sup> CHISTOV	04	BELL $B \rightarrow K D^+ D^-$

NODE=M176R4  
 NODE=M176R4

<sup>1</sup> CHISTOV 04 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow D^+D^-)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 4 \times 10^{-5}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .

NODE=M176R4;LINKAGE=A

 $\Gamma(\pi^0\chi_{c2})/\Gamma(\pi^+\pi^-J/\psi(1S))$  $\Gamma_{13}/\Gamma_3$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ABLIKIM	19U BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R06  
 NODE=M176R06

$\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
••• We do not use the following data for averages, fits, limits, etc. •••						
<0.04	90	<sup>1</sup> BHARDWAJ 19	BELL	$B^\pm \rightarrow \pi^0 \chi_{c1} K^\pm$		NODE=M176R23 NODE=M176R23
<sup>1</sup> BHARDWAJ 19 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^0 \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 8.1 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .						
$\Gamma(\pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$					$\Gamma_{14}/\Gamma_3$	
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$88^{+33}_{-27} \pm 10$	10.8	ABLIKIM 19U	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		NODE=M176R05 NODE=M176R05
$\Gamma(\pi^0 \pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$					$\Gamma_{17}/\Gamma_3$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.7	90	ABLIKIM 22D	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		NODE=M176R28 NODE=M176R28
$\Gamma(\pi^+ \pi^- \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$					$\Gamma_{18}/\Gamma_3$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.56	90	ABLIKIM 22D	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		NODE=M176R29 NODE=M176R29
$\Gamma(\pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$					$\Gamma_{15}/\Gamma_3$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< 3.6 (CL = 90%)	[<19 (CL = 90%) OUR 2022 BEST LIMIT]					
< 3.6	90	ABLIKIM 22D	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		NODE=M176R04 NODE=M176R04
••• We do not use the following data for averages, fits, limits, etc. •••						
<19	90	ABLIKIM 19U	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		
$\Gamma(\pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.14	90	<sup>1</sup> VINOKUROVA 15	BELL	$B^+ \rightarrow \pi^+ \pi^- \eta_c K^+$		NODE=M176R22 NODE=M176R22
<sup>1</sup> VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 3.0 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .						
$\Gamma(\pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}$					$\Gamma_{19}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< $7 \times 10^{-3}$	90	<sup>1</sup> BHARDWAJ 16	BELL	$B^+ \rightarrow \pi^+ \pi^- \chi_{c1} K^+$		NODE=M176R00 NODE=M176R00
<sup>1</sup> BHARDWAJ 16 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .						
$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$					$\Gamma_{20}/\Gamma$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
< $2.4 \times 10^{-5}$	95	<sup>1</sup> AAIJ 17AD	LHCB	$B^+ \rightarrow \rho\bar{\rho} K^+$		NODE=M176R03 NODE=M176R03 SYCLP=A
••• We do not use the following data for averages, fits, limits, etc. •••						
< $8 \times 10^{-5}$	95	<sup>2</sup> AAIJ 13S	LHCB	$B^+ \rightarrow \rho\bar{\rho} K^+$		SYCLP=A
<sup>1</sup> AAIJ 17AD reports $[\Gamma(\chi_{c1}(3872) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 0.5 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .						
<sup>2</sup> AAIJ 13S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .						
———— Radiative decays ————						
$\Gamma(\gamma D^+ D^-)/\Gamma(\pi^+ \pi^- J/\psi(1S))$					$\Gamma_{21}/\Gamma_3$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.99	90	ABLIKIM 20W	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		NODE=M176R20 NODE=M176R20
$\Gamma(\gamma \bar{D}^0 D^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$					$\Gamma_{22}/\Gamma_3$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.58	90	ABLIKIM 20W	BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$		NODE=M176R21 NODE=M176R21

$\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.0085<sup>+0.0024</sup><sub>-0.0022</sub> ± 0.0027**<sup>1</sup> BHARDWAJ 11 BELL  $B^\pm \rightarrow \gamma J/\psi K^\pm$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.013 ± 0.004 ± 0.004 20 <sup>2</sup> AUBERT 09B BABR  $B^+ \rightarrow \gamma J/\psi K^+$ 0.016 ± 0.005 ± 0.005 19 <sup>3</sup> AUBERT, BE 06M BABR  $B^+ \rightarrow \gamma J/\psi K^+$ 

<sup>1</sup> BHARDWAJ 11 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$   
 $= (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> AUBERT 09B reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$   
 $= (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> Superseded by AUBERT 09B. AUBERT, BE 06M reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma_{23}/\Gamma$ 

NODE=M176R7

NODE=M176R7

SYCLP=A

SYCLP=A

SYCLP=A

NODE=M176R7;LINKAGE=BA

NODE=M176R7;LINKAGE=AB

NODE=M176R7;LINKAGE=AU

 $\Gamma(\gamma J/\psi)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.79 ± 0.28**ABLIKIM 20W BES3  $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$  $\Gamma_{23}/\Gamma_3$ 

NODE=M176R18

NODE=M176R18

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 9 × 10<sup>-3</sup>**

90

<sup>1</sup> BHARDWAJ 13 BELL  $B^\pm \rightarrow \chi_{c1} \gamma K^\pm$ 

<sup>1</sup> BHARDWAJ 13 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$   
 $< 1.9 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .

 $\Gamma_{24}/\Gamma$ 

NODE=M176R08

NODE=M176R08

NODE=M176R08;LINKAGE=B

 $\Gamma(\gamma \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 0.89**

90

CHOI 03 BELL  $B \rightarrow K \pi^+ \pi^- J/\psi$  $\Gamma_{24}/\Gamma_3$ 

NODE=M176R1

NODE=M176R1

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 0.032**

90

<sup>1</sup> BHARDWAJ 13 BELL  $B^\pm \rightarrow \chi_{c2} \gamma K^\pm$ 

<sup>1</sup> BHARDWAJ 13 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \chi_{c2})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$   
 $< 6.7 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.1 \times 10^{-4}$ .

 $\Gamma_{25}/\Gamma$ 

NODE=M176R01

NODE=M176R01

NODE=M176R01;LINKAGE=B

 $\Gamma(\gamma \psi(2S))/\Gamma_{\text{total}}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

**0.045 ± 0.013 ± 0.015**

25 ± 7

<sup>1</sup> AUBERT 09B BABR  $B^+ \rightarrow \gamma \psi(2S) K^+$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 36 ± 9 <sup>2</sup> AAIJ 14AH LHCB  $B^+ \rightarrow \gamma \psi(2S) K^+$ not seen <sup>3</sup> BHARDWAJ 11 BELL  $B^+ \rightarrow \gamma \psi(2S) K^+$ 

<sup>1</sup> AUBERT 09B reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma \psi(2S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)]$   
 $= (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.1 \pm 0.7) \times 10^{-4}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> From 36.4 ± 9.0 events of  $\chi_{c1}(3872) \rightarrow J/\psi \gamma$  decays with a statistical significance of 4.4σ.

<sup>3</sup> BHARDWAJ 11 reports  $B(B^+ \rightarrow K^+ \chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma \psi(2S)) < 3.45 \times 10^{-6}$  at 90% CL.

 $\Gamma_{26}/\Gamma$ 

NODE=M176R10

NODE=M176R10

NODE=M176R10;LINKAGE=AU

NODE=M176R10;LINKAGE=A

NODE=M176R10;LINKAGE=BH

 $\Gamma(\gamma \psi(2S))/\Gamma(\pi^+ \pi^- J/\psi(1S))$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

**< 0.42**

90

ABLIKIM 20W BES3  $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$  $\Gamma_{26}/\Gamma_3$ 

NODE=M176R19

NODE=M176R19

$\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$  $\Gamma_{26}/\Gamma_{23}$ 

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6 ± 0.6</b>					<b>OUR AVERAGE</b>
2.46 ± 0.64 ± 0.29	36 ± 9		<sup>1</sup> AAIJ	14AH LHCB	$B^+ \rightarrow \gamma\psi(2S)K^+$
3.4 ± 1.4			AUBERT	09B BABR	$B^+ \rightarrow \gamma c\bar{c}K'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.1	90		BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$

<sup>1</sup> From  $36.4 \pm 9.0$  events of  $\chi_{c1}(3872) \rightarrow J/\psi\gamma$  decays with a statistical significance of  $4.4\sigma$ .

NODE=M176R11  
 NODE=M176R11

NODE=M176R11;LINKAGE=A

NODE=M176405

NODE=M176R2  
 NODE=M176R2

NODE=M176R2;LINKAGE=A

NODE=M176R2;LINKAGE=C

NODE=M176R2;LINKAGE=D

## C-violating decays

 $\Gamma(\eta J/\psi)/\Gamma_{total}$  $\Gamma_{27}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.018</b>	90	<sup>1,2</sup> IWASHITA	14 BELL	$B \rightarrow K\eta J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.04	90	<sup>3</sup> AUBERT	04Y BABR	$B \rightarrow K\eta J/\psi$

<sup>1</sup> IWASHITA 14 reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 3.8 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .

<sup>2</sup> IWASHITA 14 also scans the  $\eta J/\psi$  mass range 3.8–4.75 GeV and sets upper limits for  $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$  in 5 MeV intervals.

<sup>3</sup> AUBERT 04Y reports  $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 7.7 \times 10^{-6}$  which we divide by our best value  $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.1 \times 10^{-4}$ .

 $\chi_{c1}(3872)$  REFERENCES

NODE=M176

ADACHI	23	PR D107 052004	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=61909
ABLIKIM	22D	PR D105 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61638
TERAMOTO	21	PRL 126 122001	Y. Teramoto <i>et al.</i>	(BELLE Collab.)	REFID=61098
AAIJ	20AD	PR D102 092005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60673
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60526
ABLIKIM	20W	PRL 124 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60570
LEES	20C	PRL 124 152001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=60554
ABLIKIM	19U	PRL 122 202001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59786
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=59884
AGHASYAN	18A	PL B783 334	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59036
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=57272
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56771
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56787
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)	REFID=56408
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (err.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)	REFID=57795
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55897
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55647
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)	REFID=55925
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54985
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55008
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=55412
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54751
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=53779
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=53934
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=53939
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=53225
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53362
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53098
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52722
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52265
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT_BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51454
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)	REFID=51432
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50498
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50627
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=50458
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=50200
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=49742
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49997
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>		REFID=49677
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=49628
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074

NODE=M159

# $\chi_{c0}(3915)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

was  $X(3915)$

The  $\chi_{c0}(3915)$  was originally seen by BELLE in its  $\omega J/\psi$  decay mode and was produced in both  $B$  decays in CHOI 05 and  $\gamma\gamma$  collisions in UEHARA 10. The  $J^{PC}$  was determined to be  $0^{++}$  by BABAR in LEES 12AD but this assignment was questioned by ZHOU 15C. In AAIJ 20AI LHCb found the  $D^+ D^-$  decay mode of the  $\chi_{c0}(3915)$  using  $B$  decays and determined its  $J^{PC}$  to be  $0^{++}$ . Based on their compatible mass, width, and  $J^{PC}$ , we assume the state decaying to  $\omega J/\psi$  and the state decaying to  $D^+ D^-$  are both the  $\chi_{c0}(3915)$ . See also the  $\chi_{c2}(3930)$ .

NODE=M159

## $\chi_{c0}(3915)$ MASS

NODE=M159M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3921.7 ± 1.8 OUR AVERAGE</b>		Error includes scale factor of 1.5. See the ideogram below.		
3923.8 ± 1.5 ± 0.4	1.2k	<sup>1</sup> AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$
3919.4 ± 2.2 ± 1.6	59 ± 10	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
3919.1 <sup>+</sup> <sub>-</sub> ± 3.8 <sup>+</sup> <sub>3.4</sub> ± 2.0		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3915 ± 3 ± 2	49 ± 15	UEHARA	10 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
3943 ± 11 ± 13	58 ± 11	<sup>2</sup> CHOI	05 BELL	$B \rightarrow \omega J/\psi K$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3922.4 ± 6.5 ± 2.0		<sup>3</sup> WANG	22A BELL	$\gamma\gamma \rightarrow \gamma\psi(2S)$
3926.4 ± 2.2 ± 1.2		<sup>4</sup> ABLIKIM	19V BES	$e^+ e^- \rightarrow \gamma\omega J/\psi$
3914.6 <sup>+</sup> <sub>-</sub> ± 3.8 <sup>+</sup> <sub>3.4</sub> ± 2.0		<sup>2</sup> AUBERT	08W BABR	Superseded by DEL-AMO-SANCHEZ 10B

NODE=M159M

- <sup>1</sup> Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape.
- <sup>2</sup>  $\omega J/\psi$  threshold enhancement fitted as an S-wave Breit-Wigner resonance.
- <sup>3</sup> Not distinguished from the  $\chi_{c2}(3930)$ .
- <sup>4</sup> Could also be  $X(3940)$ . Significance  $3.1\sigma$ . Fit with additional resonance at  $3963.7 \pm 5.7$  MeV, significance  $3.4\sigma$ .

NODE=M159M;LINKAGE=B

NODE=M159M;LINKAGE=CH  
 NODE=M159M;LINKAGE=C  
 NODE=M159M;LINKAGE=A



## $\chi_{c0}(3915)$ WIDTH

NODE=M159W



VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18.8 ± 3.5 OUR AVERAGE</b>				
17.4 ± 5.1 ± 0.8	1.2k	<sup>1</sup> AAIJ	20A1	LHCB $B^+ \rightarrow D^+ D^- K^+$
13 ± 6 ± 3	59	LEES	12AD	BABR $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
31 $\begin{smallmatrix} +10 \\ -8 \end{smallmatrix}$ ± 5		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
17 ± 10 ± 3	49	UEHARA	10	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
87 ± 22 ± 26	58	<sup>2</sup> CHOI	05	BELL $B \rightarrow \omega J/\psi K$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
22 ± 17 ± 4		<sup>3</sup> WANG	22A	BELL $\gamma\gamma \rightarrow \gamma\psi(2S)$
3.8 ± 7.5 ± 2.6		<sup>4</sup> ABLIKIM	19V	BES $e^+ e^- \rightarrow \gamma\omega J/\psi$
34 $\begin{smallmatrix} +12 \\ -8 \end{smallmatrix}$ ± 5		<sup>2</sup> AUBERT	08W	BABR Superseded by DEL-AMO-SANCHEZ 10B
<sup>1</sup> Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape.				
<sup>2</sup> $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.				
<sup>3</sup> Not distinguished from the $\chi_{c2}(3930)$ .				
<sup>4</sup> Could also be $X(3940)$ . Significance $3.1\sigma$ . Fit with additional resonance at $3963.7 \pm 5.7$ MeV, significance $3.4\sigma$ .				

NODE=M159W

NODE=M159W;LINKAGE=B

NODE=M159W;LINKAGE=CH

NODE=M159W;LINKAGE=C

NODE=M159W;LINKAGE=A

 **$\chi_{c0}(3915)$  DECAY MODES**

NODE=M159215;NODE=M159

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\omega J/\psi$	seen
$\Gamma_2$ $\bar{D}^{*0} D^0$	not seen
$\Gamma_3$ $D^+ D^-$	seen
$\Gamma_4$ $\pi^+ \pi^- \eta_c(1S)$	not seen
$\Gamma_5$ $\eta_c \eta$	not seen
$\Gamma_6$ $\eta_c \pi^0$	not seen
$\Gamma_7$ $K \bar{K}$	not seen
$\Gamma_8$ $\gamma\gamma$	seen
$\Gamma_9$ $\gamma\psi(2S)$	
$\Gamma_{10}$ $\pi^0 \chi_{c1}$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←  
DESIG=3;OUR EVAL;→ UNCHECKED ←  
DESIG=9  
DESIG=4;OUR EVAL;→ UNCHECKED ←  
DESIG=6  
DESIG=7  
DESIG=5;OUR EVAL;→ UNCHECKED ←  
DESIG=2  
DESIG=10  
DESIG=8

 **$\chi_{c0}(3915)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$** 

NODE=M159220

$\Gamma(\omega J/\psi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_1 \Gamma_8/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>54 ± 9 OUR AVERAGE</b>				
52 ± 10 ± 3	59 ± 10	<sup>1</sup> LEES	12AD	BABR $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
61 ± 17 ± 8	49 ± 15	<sup>1</sup> UEHARA	10	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
18 ± 5 ± 2	49 ± 15	<sup>2</sup> UEHARA	10	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
<sup>1</sup> For $J^P = 0^+$ .				
<sup>2</sup> For $J^P = 2^+$ , helicity-2.				

NODE=M159G01  
NODE=M159G01

OCCUR=2

NODE=M159G01;LINKAGE=UH  
NODE=M159G01;LINKAGE=UR

$\Gamma(\gamma\psi(2S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_9 \Gamma_8/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT

NODE=M159R07  
NODE=M159R07

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
9.8 ± 3.6 ± 1.3		<sup>1</sup> WANG	22A	BELL $\gamma\gamma \rightarrow \gamma\psi(2S)$
<sup>1</sup> Not distinguished from the $\chi_{c2}(3930)$ .				

NODE=M159R07;LINKAGE=A

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_4 \Gamma_8/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	LEES	12AE	BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

NODE=M159G02  
NODE=M159G02

$\Gamma(K \bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_7 \Gamma_8/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.96	90	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M159G03  
NODE=M159G03

$\chi_{c0}(3915)$  BRANCHING RATIOS $\Gamma(\omega J/\psi)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> DEL-AMO-SA...10B	BABR	$B \rightarrow \omega J/\psi K$
seen	<sup>2</sup> CHOI	05 BELL	$B \rightarrow \omega J/\psi K$

<sup>1</sup> DEL-AMO-SANCHEZ 10B reports  $B(B^\pm \rightarrow \chi_{c0}(3915) K^\pm) \times B(\chi_{c0}(3915) \rightarrow J/\psi \omega)$   
 $= (3.0^{+0.7+0.5}_{-0.6-0.3}) \times 10^{-5}$  and  $B(B^0 \rightarrow \chi_{c0}(3915) K^0) \times B(\chi_{c0}(3915) \rightarrow J/\psi \omega)$   
 $= (2.1 \pm 0.9 \pm 0.3) \times 10^{-5}$ .

<sup>2</sup> CHOI 05 reports  $B(B \rightarrow \chi_{c0}(3915) K) \times B(\chi_{c0}(3915) \rightarrow J/\psi \omega) = (7.1 \pm 1.3 \pm 3.1) \times 10^{-5}$ .

NODE=M159225

NODE=M159R03  
NODE=M159R03

NODE=M159R03;LINKAGE=DE

NODE=M159R03;LINKAGE=CH

 $\Gamma(\omega J/\psi)/\Gamma(\bar{D}^{*0} D^0)$  $\Gamma_1/\Gamma_2$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.71	90	<sup>1</sup> AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$

<sup>1</sup> By combining the upper limit  $B(B \rightarrow \chi_{c0}(3915) K) \times B(\chi_{c0}(3915) \rightarrow D^{*0} \bar{D}^0)$   
 $< 0.67 \times 10^{-4}$  from AUSHEV 10 with the average of CHOI 05 and AUBERT 08W  
measurements  $B(B \rightarrow \chi_{c0}(3915) K) \times B(\chi_{c0}(3915) \rightarrow \omega J/\psi) = (0.51 \pm 0.11) \times 10^{-4}$ .

NODE=M159R02  
NODE=M159R02

NODE=M159R02;LINKAGE=AU

 $\Gamma(D^+ D^-)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M159R06  
NODE=M159R06 $\Gamma(\eta_c \eta)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
not seen	90	<sup>1</sup> VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \eta$

<sup>1</sup> VINOKUROVA 15 reports  $B(B^+ \rightarrow K^+ \chi_{c0}(3915)) \times B(\chi_{c0}(3915) \rightarrow \eta_c \eta) < 4.7 \times 10^{-5}$  at 90% CL.

NODE=M159R00  
NODE=M159R00

OCCUR=2

NODE=M159R00;LINKAGE=A

 $\Gamma(\eta_c \pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
not seen	90	<sup>1</sup> VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \pi^0$

<sup>1</sup> VINOKUROVA 15 reports  $B(B^+ \rightarrow K^+ \chi_{c0}(3915)^0) \times B(\chi_{c0}(3915) \rightarrow \eta_c \pi^0) < 1.7 \times 10^{-5}$  at 90% CL.

NODE=M159R04  
NODE=M159R04

OCCUR=2

NODE=M159R04;LINKAGE=A

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	59 ± 10	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
seen		UEHARA	10 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$

NODE=M159R01  
NODE=M159R01 $\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
not seen	42 ± 14	<sup>1</sup> BHARDWAJ 19	BELL	$B^\pm \rightarrow \chi_{c1} \pi^0 K^\pm$

<sup>1</sup> BHARDWAJ 19 reports  $B(B^+ \rightarrow K^+ \chi_{c0}(3915)) \times B(\chi_{c0}(3915) \rightarrow \chi_{c1} \pi^0) < 3.8 \times 10^{-5}$  at 90% CL. A signal significance 2.3 standard deviations.

NODE=M159R05  
NODE=M159R05

NODE=M159R05;LINKAGE=A

 $\chi_{c0}(3915)$  REFERENCES

NODE=M159

WANG	22A	PR D105 112011	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=61640
AAIJ	20AI	PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.) JPC	REFID=60739
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=59884
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (err.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)	REFID=57795
ZHOU	15C	PRL 115 022001	Z.-Y. Zhou, Z. Xiao, H.-Q. Zhou	(BEIJT, NANJ)	REFID=56842
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54751
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=53225
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53362
UEHARA	10	PRL 104 092001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53232
AUBERT	08W	PRL 101 082001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52263
CHOI	05	PRL 94 182002	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=50737

$\chi_{c2}(3930)$ 

$$J^G(J^{PC}) = 0^+(2^{++})$$

NODE=M050

 **$\chi_{c2}(3930)$  MASS**

NODE=M050M

NODE=M050M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3922.5±1.0 OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.		
3926.8±2.4±0.8	1.2k	<sup>1</sup> AAIJ	20AI LHCb	$B^+ \rightarrow D^+ D^- K^+$
3921.9±0.6±0.2		<sup>2</sup> AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$
3926.7±2.7±1.1	76 ± 17	AUBERT	10G BABR	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$
3929 ± 5 ± 2	64	UEHARA	06 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$

<sup>1</sup> Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Previous measurements assumed a single state in this region. This analysis revealed the presence of  $\chi_{c0}(3930)$  with the same mass.

<sup>2</sup> Measured in prompt hadroproduction.

NODE=M050M;LINKAGE=B

NODE=M050M;LINKAGE=A

 **$\chi_{c2}(3930)$  WIDTH**

NODE=M050W

NODE=M050W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>35.2± 2.2 OUR AVERAGE</b>		Error includes scale factor of 1.2.		
34.2± 6.6±1.1	1.2k	<sup>1</sup> AAIJ	20AI LHCb	$B^+ \rightarrow D^+ D^- K^+$
36.6± 1.9±0.9		<sup>2</sup> AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$
21.3± 6.8±3.6	76 ± 17	AUBERT	10G BABR	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$
29 ± 10 ± 2	64	UEHARA	06 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$

<sup>1</sup> Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Previous measurements assumed a single state in this region. This analysis revealed the presence of  $\chi_{c0}(3930)$  with the same mass.

<sup>2</sup> Measured in prompt hadroproduction.

NODE=M050W;LINKAGE=B

NODE=M050W;LINKAGE=A

 **$\chi_{c2}(3930)$  DECAY MODES**

NODE=M050215;NODE=M050

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\gamma\gamma$	seen
$\Gamma_2$ $K\bar{K}\pi$	
$\Gamma_3$ $K^+ K^- \pi^+ \pi^- \pi^0$	
$\Gamma_4$ $D\bar{D}$	seen
$\Gamma_5$ $D^+ D^-$	seen
$\Gamma_6$ $D^0 \bar{D}^0$	seen
$\Gamma_7$ $\pi^+ \pi^- \eta_c(1S)$	not seen
$\Gamma_8$ $K\bar{K}$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=5

DESIG=6

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=7;OUR EVAL;→ UNCHECKED ←

DESIG=8;OUR EVAL;→ UNCHECKED ←

$\chi_{c2}(3930)$  PARTIAL WIDTHS

NODE=M050220

 $\chi_{c2}(3930) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M050222

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.1	90	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	

NODE=M050G01  
NODE=M050G01

$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<3.4	90	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

NODE=M050G02  
NODE=M050G02

$\Gamma(D\bar{D}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_1/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.21±0.04 OUR AVERAGE</b>					

NODE=M050G1  
NODE=M050G1

0.24±0.05±0.04	76 ± 17	AUBERT	10G	BABR	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$
0.18±0.05±0.03	64	<sup>1</sup> UEHARA	06	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$

<sup>1</sup> Assuming  $B(D^+ D^-) = 0.89 B(D^0 \bar{D}^0)$ .

NODE=M050G1;LINKAGE=UE

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<18	90	LEES	12AE	BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

NODE=M050G03  
NODE=M050G03

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.256	90	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M050G04  
NODE=M050G04 $\chi_{c2}(3930)$  BRANCHING RATIOS

NODE=M050225

$\Gamma(D^+ D^-)/\Gamma(D^0 \bar{D}^0)$					$\Gamma_5/\Gamma_6$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>0.74±0.43±0.16</b>	64	UEHARA	06	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- D\bar{D}$

NODE=M050R01  
NODE=M050R01 $\chi_{c2}(3930)$  REFERENCES

NODE=M050

AAIJ	20AI	PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60739
AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59697
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
DEL-AMO-SA...11M	PR D84	012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
AUBERT	10G	PR D81 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53357
UEHARA	06	PRL 96 082003	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=51039

**X(3940)**

$$I^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Reported by ABE 07, observed in  $e^+e^- \rightarrow J/\psi X$ .

NODE=M029

NODE=M029

NODE=M029M

NODE=M029M

**X(3940) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3942^{+7}_{-6} \pm 6$	52	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$3943 \pm 6 \pm 6$	25	<sup>1</sup> ABE	07	BELL $e^+e^- \rightarrow J/\psi X$
$3936 \pm 14$	266	<sup>2</sup> ABE	07	BELL $e^+e^- \rightarrow J/\psi(c\bar{c})$
<sup>1</sup> From a fit to $D^{*+}D^-$ and $D^{*0}\bar{D}^0$ events.				
<sup>2</sup> From the inclusive fit. Not independent of the exclusive measurement by ABE 07.				

OCCUR=2

NODE=M029M;LINKAGE=EB

NODE=M029M;LINKAGE=EM

**X(3940) WIDTH**

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$37^{+26}_{-15} \pm 8$		52	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<52	90	25	ABE	07	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M029W

NODE=M029W

**X(3940) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D\bar{D}^* + c.c.$	seen
$\Gamma_2$ $D\bar{D}$	not seen
$\Gamma_3$ $J/\psi\omega$	not seen

NODE=M029215;NODE=M029

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

**X(3940) BRANCHING RATIOS**

$\Gamma(D\bar{D}^* + c.c.)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
>0.45	90	25	<sup>1,2</sup> ABE	07	BELL $e^+e^- \rightarrow J/\psi X$
<sup>1</sup> For X(3940) decaying to final states with more than two tracks.					
<sup>2</sup> PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c <sup>2</sup> may consist of several states.					

NODE=M029225

NODE=M029R01

NODE=M029R01

NODE=M029R01;LINKAGE=AB

NODE=M029R01;LINKAGE=AE

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.41	90	<sup>1,2</sup> ABE	07	BELL $e^+e^- \rightarrow J/\psi X$	
<sup>1</sup> For X(3940) decaying to final states with more than two tracks.					
<sup>2</sup> PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c <sup>2</sup> may consist of several states.					

NODE=M029R02

NODE=M029R02

NODE=M029R02;LINKAGE=AB

NODE=M029R02;LINKAGE=AE

$\Gamma(J/\psi\omega)/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.26	90	<sup>1,2</sup> ABE	07	BELL $e^+e^- \rightarrow J/\psi X$	
<sup>1</sup> For X(3940) decaying to final states with more than two tracks.					
<sup>2</sup> PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c <sup>2</sup> may consist of several states.					

NODE=M029R03

NODE=M029R03

NODE=M029R03;LINKAGE=AB

NODE=M029R03;LINKAGE=AE

**X(3940) REFERENCES**

PAKHLOV 08	PRL 100 202001	P. Pakhlov <i>et al.</i>	(BELLE Collab.)
ABE 07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M029

REFID=52302

REFID=51627

$\psi(4040)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M072

 $\psi(4040)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>4039 ± 1 OUR ESTIMATE</b>			
<b>4039.6 ± 4.3</b>	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4034 ± 6	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4037 ± 2	<sup>3</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4040 ± 1	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4040 ± 10	BRANDELIK	78C DASP	$e^+e^-$
<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$ , $\psi(4040)$ , $\psi(4160)$ , and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$ .			
<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$ , $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.			
<sup>4</sup> From a fit to BES (BAI 02C) data.			

NODE=M072M

NODE=M072M  
→ UNCHECKED ←

OCCUR=2

NODE=M072M;LINKAGE=AB

NODE=M072M;LINKAGE=MO

NODE=M072M;LINKAGE=ST  
NODE=M072M;LINKAGE=SE $\psi(4040)$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>80 ± 10 OUR ESTIMATE</b>			
<b>84.5 ± 12.3</b>	<sup>5</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87 ± 11	<sup>6</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
85 ± 10	<sup>7</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
89 ± 6	<sup>8</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
52 ± 10	BRANDELIK	78C DASP	$e^+e^-$
<sup>5</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$ , $\psi(4040)$ , $\psi(4160)$ , and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$ .			
<sup>6</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$ , $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
<sup>7</sup> From a fit to Crystal Ball (OSTERHELD 86) data.			
<sup>8</sup> From a fit to BES (BAI 02C) data.			

NODE=M072W

NODE=M072W  
→ UNCHECKED ←

OCCUR=2

NODE=M072W;LINKAGE=AB

NODE=M072W;LINKAGE=MO

NODE=M072W;LINKAGE=ST  
NODE=M072W;LINKAGE=SE $\psi(4040)$  DECAY MODES

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle’s central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

NODE=M072215;NODE=M072

NODE=M072

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $e^+e^-$	$(1.07 \pm 0.16) \times 10^{-5}$	DESIG=5
$\Gamma_2$ $D\bar{D}$	seen	DESIG=17;OUR EST;→ UNCHECKED ←
$\Gamma_3$ $D^0\bar{D}^0$	seen	DESIG=1
$\Gamma_4$ $D^+D^-$	seen	DESIG=18
$\Gamma_5$ $D^*\bar{D} + c.c.$	seen	DESIG=19;OUR EST;→ UNCHECKED ←
$\Gamma_6$ $D^*(2007)^0\bar{D}^0 + c.c.$	seen	DESIG=2
$\Gamma_7$ $D^*(2010)^+D^- + c.c.$	seen	DESIG=20
$\Gamma_8$ $D^*\bar{D}^*$	seen	DESIG=21;OUR EST;→ UNCHECKED ←
$\Gamma_9$ $D^*(2007)^0\bar{D}^*(2007)^0$	seen	DESIG=3
$\Gamma_{10}$ $D^*(2010)^+D^*(2010)^-$	seen	DESIG=22
$\Gamma_{11}$ $D\bar{D}\pi$ (excl. $D^*\bar{D}$ )		DESIG=23
$\Gamma_{12}$ $D^0D^-\pi^+ + c.c.$ (excl. $D^*(2007)^0\bar{D}^0 + c.c.$ , $D^*(2010)^+D^- + c.c.$ )	not seen	DESIG=24
$\Gamma_{13}$ $D\bar{D}^*\pi$ (excl. $D^*\bar{D}^*$ )	not seen	DESIG=25

Γ <sub>14</sub>	$D^0 \bar{D}^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)$	seen			DESIG=26
Γ <sub>15</sub>	$D_s^+ D_s^-$	seen			DESIG=27
Γ <sub>16</sub>	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$				DESIG=37
Γ <sub>17</sub>	$J/\psi(1S)$ hadrons				DESIG=4
Γ <sub>18</sub>	$J/\psi \pi^+ \pi^-$	< 4	$\times 10^{-3}$	90%	DESIG=7
Γ <sub>19</sub>	$J/\psi \pi^0 \pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=8
Γ <sub>20</sub>	$J/\psi \eta$	(5.2 ± 0.7)	$\times 10^{-3}$		DESIG=9
Γ <sub>21</sub>	$J/\psi \pi^0$	< 2.8	$\times 10^{-4}$	90%	DESIG=10
Γ <sub>22</sub>	$J/\psi \pi^+ \pi^- \pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=11
Γ <sub>23</sub>	$\chi_{c1} \gamma$	< 3.4	$\times 10^{-3}$	90%	DESIG=12
Γ <sub>24</sub>	$\chi_{c2} \gamma$	< 5	$\times 10^{-3}$	90%	DESIG=13
Γ <sub>25</sub>	$\chi_{c1} \pi^+ \pi^- \pi^0$	< 1.1	%	90%	DESIG=14
Γ <sub>26</sub>	$\chi_{c2} \pi^+ \pi^- \pi^0$	< 3.2	%	90%	DESIG=15
Γ <sub>27</sub>	$h_c(1P) \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=28
Γ <sub>28</sub>	$\phi \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=16
Γ <sub>29</sub>	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.9	$\times 10^{-4}$	90%	DESIG=29
Γ <sub>30</sub>	$\Lambda \bar{\Lambda} \pi^0$	< 9	$\times 10^{-5}$	90%	DESIG=30
Γ <sub>31</sub>	$\Lambda \bar{\Lambda} \eta$	< 3.0	$\times 10^{-4}$	90%	DESIG=31
Γ <sub>32</sub>	$\Lambda \bar{\Lambda}$	< 6	$\times 10^{-6}$	90%	DESIG=36
Γ <sub>33</sub>	$\Sigma^+ \bar{\Sigma}^-$	< 1.3	$\times 10^{-4}$	90%	DESIG=32
Γ <sub>34</sub>	$\Sigma^0 \bar{\Sigma}^0$	< 7	$\times 10^{-5}$	90%	DESIG=33
Γ <sub>35</sub>	$\Xi^+ \bar{\Xi}^-$	< 1.6	$\times 10^{-4}$	90%	DESIG=34
Γ <sub>36</sub>	$\Xi^0 \bar{\Xi}^0$	< 1.8	$\times 10^{-4}$	90%	DESIG=35
Γ <sub>37</sub>	$\mu^+ \mu^-$	(9 ± 6)	$\times 10^{-6}$		DESIG=6

### ψ(4040) PARTIAL WIDTHS

NODE=M072220

#### Γ(e<sup>+</sup>e<sup>-</sup>)

Γ<sub>1</sub>

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.86±0.07 OUR ESTIMATE</b>			
<b>0.83±0.20</b>	<sup>9</sup> ABLIKIM	08D BES2	e <sup>+</sup> e <sup>-</sup> → hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.6 to 1.4	<sup>10</sup> MO	10 RVUE	e <sup>+</sup> e <sup>-</sup> → hadrons
0.88±0.11	<sup>11</sup> SETH	05A RVUE	e <sup>+</sup> e <sup>-</sup> → hadrons
0.91±0.13	<sup>12</sup> SETH	05A RVUE	e <sup>+</sup> e <sup>-</sup> → hadrons
0.75±0.15	BRANDELIK	78C DASP	e <sup>+</sup> e <sup>-</sup>

NODE=M072W5  
NODE=M072W5

→ UNCHECKED ←

OCCUR=2

<sup>9</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the ψ(3770), ψ(4040), ψ(4160), and ψ(4415) resonances. Phase angle fixed in the fit to δ = (130 ± 46)°.

NODE=M072W5;LINKAGE=AB

<sup>10</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the ψ(4040), ψ(4160) and ψ(4415) resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e<sup>+</sup>e<sup>-</sup> partial widths. We quote only the range of values.

NODE=M072W5;LINKAGE=MO

<sup>11</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M072W5;LINKAGE=ST

<sup>12</sup> From a fit to BES (BAI 02C) data.

NODE=M072W5;LINKAGE=SE

#### Γ(μ<sup>+</sup>μ<sup>-</sup>)

Γ<sub>37</sub>

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.73±0.48±0.12</b>	<sup>13,14</sup> ABLIKIM	20AG BES3	e <sup>+</sup> e <sup>-</sup> → μ <sup>+</sup> μ <sup>-</sup>
<sup>13</sup> From a fit to the e <sup>+</sup> e <sup>-</sup> → μ <sup>+</sup> μ <sup>-</sup> cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming Γ(μ <sup>+</sup> μ <sup>-</sup> ) = Γ(e <sup>+</sup> e <sup>-</sup> ).			
<sup>14</sup> From solution 1 of 8 with equal fit quality. Other solutions range from 0.58 ± 0.52 ± 0.10 to 0.80 ± 0.48 ± 0.13 keV.			

NODE=M072W2  
NODE=M072W2

NODE=M072W2;LINKAGE=A

NODE=M072W2;LINKAGE=B

### ψ(4040) Γ(i) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ(total)

NODE=M072235

#### Γ(χ<sub>c1</sub>γ) × Γ(e<sup>+</sup>e<sup>-</sup>)/Γ<sub>total</sub>

Γ<sub>23</sub>Γ<sub>1</sub>/Γ

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.9</b>	90	<sup>15</sup> HAN	15 BELL	10.58 e <sup>+</sup> e <sup>-</sup> → χ <sub>c1</sub> γ

NODE=M072G01  
NODE=M072G01

<sup>15</sup> Using B(η → γγ) = (39.41 ± 0.21)%.

NODE=M072G01;LINKAGE=A

$$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<4.6	90	<sup>16</sup> HAN	15	BELL 10.58 e <sup>+</sup> e <sup>-</sup> → χ <sub>c2</sub> γ

<sup>16</sup> Using B(η → γγ) = (39.41 ± 0.21)%.

NODE=M072G02  
NODE=M072G02

NODE=M072G02;LINKAGE=A

$$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{20}\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ± 0.3	<sup>17</sup> ABLIKIM	200	BES3 e <sup>+</sup> e <sup>-</sup> → ηJ/ψ
1.4 ± 0.3	<sup>18</sup> ABLIKIM	200	BES3 e <sup>+</sup> e <sup>-</sup> → ηJ/ψ
7.0 ± 0.6	<sup>19</sup> ABLIKIM	200	BES3 e <sup>+</sup> e <sup>-</sup> → ηJ/ψ

NODE=M072R00  
NODE=M072R00

OCCUR=2

OCCUR=3

NODE=M072R00;LINKAGE=A  
NODE=M072R00;LINKAGE=B  
NODE=M072R00;LINKAGE=C

<sup>17</sup> Solution 1 of three equivalent fit solutions using three resonant structures.

<sup>18</sup> Solution 2 of three equivalent fit solutions using three resonant structures.

<sup>19</sup> Solution 3 of three equivalent fit solutions using three resonant structures.

$$\psi(4040) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$$

NODE=M072230

$$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$$

VALUE (units 10 <sup>-8</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.1 ± 1.4 ± 1.5	<sup>20</sup> WANG	13B	BELL e <sup>+</sup> e <sup>-</sup> → J/ψηγ
12.8 ± 2.1 ± 1.9	<sup>21</sup> WANG	13B	BELL e <sup>+</sup> e <sup>-</sup> → J/ψηγ

NODE=M072R25  
NODE=M072R25

OCCUR=2

<sup>20</sup> Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

NODE=M072R25;LINKAGE=A

<sup>21</sup> Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

NODE=M072R25;LINKAGE=B

$$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{32}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<5.5 × 10<sup>-3</sup> 90 <sup>22</sup> ABLIKIM 21AS BES3 e<sup>+</sup>e<sup>-</sup> → ψ(4040)

<sup>22</sup> From a measurement of the e<sup>+</sup>e<sup>-</sup> → ΛΛ̄ cross section between 3.5 and 4.6 GeV.

NODE=M072R33  
NODE=M072R33

NODE=M072R33;LINKAGE=A

### ψ(4040) BRANCHING RATIOS

NODE=M072225

$$\Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE (units 10 <sup>-5</sup> )	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0 FELDMAN 77 MRK1 e<sup>+</sup>e<sup>-</sup>

NODE=M072R4  
NODE=M072R4

$$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen AUBERT 09M BABR e<sup>+</sup>e<sup>-</sup> → D<sup>0</sup>̄D<sup>0</sup>γ

seen CRONIN-HEN..09 CLEO e<sup>+</sup>e<sup>-</sup> → D<sup>0</sup>̄D<sup>0</sup>

seen PAKHLOVA 08 BELL e<sup>+</sup>e<sup>-</sup> → D<sup>0</sup>̄D<sup>0</sup>γ

NODE=M072R14  
NODE=M072R14

$$\Gamma(D^+D^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen AUBERT 09M BABR e<sup>+</sup>e<sup>-</sup> → D<sup>+</sup>D<sup>-</sup>γ

seen CRONIN-HEN..09 CLEO e<sup>+</sup>e<sup>-</sup> → D<sup>+</sup>D<sup>-</sup>

seen PAKHLOVA 08 BELL e<sup>+</sup>e<sup>-</sup> → D<sup>+</sup>D<sup>-</sup>γ

NODE=M072R15  
NODE=M072R15

$$\Gamma(D\bar{D})/\Gamma(D^*\bar{D} + \text{c.c.}) \quad \Gamma_2/\Gamma_5$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.24 ± 0.05 ± 0.12 AUBERT 09M BABR e<sup>+</sup>e<sup>-</sup> → γD(\*)̄D

NODE=M072R12  
NODE=M072R12

$$\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.}) \quad \Gamma_3/\Gamma_6$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.05 ± 0.03 <sup>23</sup> GOLDHABER 77 MRK1 e<sup>+</sup>e<sup>-</sup>

NODE=M072R1  
NODE=M072R1

<sup>23</sup> Phase-space factor (p<sup>3</sup>) explicitly removed.

NODE=M072R;LINKAGE=P

$$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen AUBERT 09M BABR e<sup>+</sup>e<sup>-</sup> → D<sup>\*0</sup>̄D<sup>0</sup>γ

seen CRONIN-HEN..09 CLEO e<sup>+</sup>e<sup>-</sup> → D<sup>\*0</sup>̄D<sup>0</sup>

NODE=M072R16  
NODE=M072R16



$$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$$

 $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>24</sup> ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
<b>seen</b>	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
<sup>24</sup> Supersedes PAKHLOVA 07.			

NODE=M072R17  
 NODE=M072R17  
 OCCUR=3

NODE=M072R17;LINKAGE=C

$$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})$$

 $\Gamma_7/\Gamma_6$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.95 ± 0.09 ± 0.10</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^* \bar{D}$

NODE=M072R11  
 NODE=M072R11

$$\Gamma(D^* \bar{D}^*)/\Gamma(D^* \bar{D} + \text{c.c.})$$

 $\Gamma_8/\Gamma_5$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.18 ± 0.14 ± 0.03</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

NODE=M072R13  
 NODE=M072R13

$$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$$

 $\Gamma_9/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$

NODE=M072R18  
 NODE=M072R18

$$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.})$$

 $\Gamma_9/\Gamma_6$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>32.0 ± 12.0</b>	<sup>25</sup> GOLDHABER 77	MRK1	$e^+ e^-$

NODE=M072R2  
 NODE=M072R2

<sup>25</sup>Phase-space factor ( $p^3$ ) explicitly removed.

NODE=M072R2;LINKAGE=P

$$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$$

 $\Gamma_{10}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>26</sup> ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<b>seen</b>	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
<b>seen</b>	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
<sup>26</sup> Supersedes PAKHLOVA 07.			

NODE=M072R19  
 NODE=M072R19  
 OCCUR=2

NODE=M072R19;LINKAGE=B

$$\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.})/$$

 $\Gamma_{\text{total}}$ 
 $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

NODE=M072R20  
 NODE=M072R20

$$\Gamma(D \bar{D}^* \pi (\text{excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$$

 $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

NODE=M072R21  
 NODE=M072R21

$$\Gamma(D^0 \bar{D}^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$$

 $\Gamma_{14}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

NODE=M072R22  
 NODE=M072R22

$$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$$

 $\Gamma_{15}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>seen</b>	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
<b>seen</b>	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

NODE=M072R23  
 NODE=M072R23

$$\Gamma(\pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_{16}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$(3.51 \pm 1.89 \pm 1.24) \times 10^{-5}$	<sup>27</sup> ABLIKIM 21AW	BES3	$e^+ e^- \rightarrow 2\pi^+ 2\pi^- \pi^0$
$(2.41 \pm 0.05 \pm 0.79) \times 10^{-2}$	<sup>28</sup> ABLIKIM 21AW	BES3	$e^+ e^- \rightarrow 2\pi^+ 2\pi^- \pi^0$

NODE=M072R34  
 NODE=M072R34

OCCUR=2

<sup>27</sup>Solution 1 of two solutions with equal fit quality. The significance of the  $\psi(4040)$  signal is 3.6  $\sigma$ .

NODE=M072R34;LINKAGE=A

<sup>28</sup>Solution 2 of two solutions with equal fit quality. The significance of the  $\psi(4040)$  signal is 3.6  $\sigma$ .

NODE=M072R34;LINKAGE=B

$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R01  
NODE=M072R01 $\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R02  
NODE=M072R02 $\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>5.2±0.5±0.5</b>		<sup>29</sup> ABLIKIM	12K	BES3 $e^+e^- \rightarrow \ell^+\ell^-2\gamma$

NODE=M072R03  
NODE=M072R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons
----	----	------	----	---

<sup>29</sup> ABLIKIM 12K measure  $\sigma(e^+e^- \rightarrow J/\psi\eta) = 32.1 \pm 2.8 \pm 1.3$  pb. They assume the  $\eta J/\psi$  fully originates from  $\psi(4040)$  decays.

NODE=M072R03;LINKAGE=AB

 $\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.28	90	<sup>30</sup> ABLIKIM	12K	BES3 $e^+e^- \rightarrow \ell^+\ell^-2\gamma$

NODE=M072R04  
NODE=M072R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons
----	----	------	----	---

<sup>30</sup> ABLIKIM 12K measure  $\sigma(e^+e^- \rightarrow J/\psi\pi^0) < 1.6$  pb. They assume the  $\eta J/\psi$  fully originates from  $\psi(4040)$  decays.

NODE=M072R04;LINKAGE=AB

 $\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R05  
NODE=M072R05 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R06  
NODE=M072R06

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<17	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R07  
NODE=M072R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R08  
NODE=M072R08 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<32	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R09  
NODE=M072R09 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	<sup>31</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

NODE=M072R24  
NODE=M072R24<sup>31</sup> From several values of  $\sqrt{s}$  near the peak of the  $\psi(4040)$ , PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 1.0 \pm 8.0 \pm 5.4 \pm 0.2$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

NODE=M072R24;LINKAGE=PE

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN	06	CLEO 3.97–4.06 $e^+e^- \rightarrow$ hadrons

NODE=M072R10  
NODE=M072R10 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	<sup>32</sup> ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

NODE=M072R26  
NODE=M072R26<sup>32</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R26;LINKAGE=A

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{30}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	33 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

<sup>33</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R27  
NODE=M072R27

NODE=M072R27;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$  $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	34 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

<sup>34</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R28  
NODE=M072R28

NODE=M072R28;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$  $\Gamma_{33}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	35 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

<sup>35</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R29  
NODE=M072R29

NODE=M072R29;LINKAGE=A

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$  $\Gamma_{34}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	36 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

<sup>36</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R30  
NODE=M072R30

NODE=M072R30;LINKAGE=A

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$  $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	37 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

<sup>37</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R31  
NODE=M072R31

NODE=M072R31;LINKAGE=A

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$  $\Gamma_{36}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	38 ABLIKIM	13Q	BES3 $e^+e^- \rightarrow \psi(4040)$

<sup>38</sup> Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R32  
NODE=M072R32

NODE=M072R32;LINKAGE=A

 $\psi(4040)$  REFERENCES

NODE=M072

ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21AW	PR D104 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61479
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	200	PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60344
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
ABLIKIM	13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55393
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
ABLIKIM	12K	PR D86 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54738
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51075
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232
Also		ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)	REFID=11434

NODE=M193

# $\chi_{c1}(4140)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

was  $X(4140)$

This state shows properties different from a conventional  $q\bar{q}$  state. A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

NODE=M193

Seen by AALTONEN 09AH, ABAZOV 14A, CHATRCHYAN 14M, AAIJ 17C in  $B^+ \rightarrow \chi_{c1} K^+$ ,  $\chi_{c1} \rightarrow J/\psi \phi$ , and by ABAZOV 15M separately in both prompt ( $4.7 \sigma$ ) and non-prompt ( $5.6 \sigma$ ) production in  $p\bar{p} \rightarrow J/\psi \phi + \text{anything}$ . Not seen by SHEN 10 in  $\gamma\gamma \rightarrow J/\psi \phi$  and ABLIKIM 15 in  $e^+e^- \rightarrow \gamma J/\psi \phi$  at  $\sqrt{s} = 4.23, 4.26, 4.36$  GeV.

## $\chi_{c1}(4140)$ MASS

NODE=M193M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4146.5 ± 3.0 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.		
4118 ± 11 <sup>+19</sup> <sub>-36</sub>	24k	1 AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$
4143.4 <sup>+2.9</sup> <sub>-3.0</sub> ± 0.6	19	2 AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$
4152.5 ± 1.7 <sup>+6.2</sup> <sub>-5.4</sub>	616	3 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi \phi + \text{anything}$
4159.0 ± 4.3 ± 6.6	52	4 ABAZOV	14A D0	$B^+ \rightarrow J/\psi \phi K^+$
4148.0 ± 2.4 ± 6.3	0.3k	5 CHATRCHYAN 14M	CMS	$B^+ \rightarrow J/\psi \phi K^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4146.5 ± 4.5 <sup>+4.6</sup> <sub>-2.8</sub>	4289	6,7 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
4143.0 ± 2.9 ± 1.2	14	8,9 AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M193M

- <sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $13 \sigma$ .
- <sup>2</sup> Statistical significance of more than  $5 \sigma$ .
- <sup>3</sup> Statistical significance of more than  $6 \sigma$ .
- <sup>4</sup> Statistical significance of  $3.1 \sigma$ .
- <sup>5</sup> From a fit assuming an  $S$ -wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than  $5 \sigma$ .
- <sup>6</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $8.4 \sigma$ .
- <sup>7</sup> Superseded by AAIJ 21E.
- <sup>8</sup> Statistical significance of  $3.8 \sigma$ .
- <sup>9</sup> Superseded by AALTONEN 17.

NODE=M193M;LINKAGE=G  
 NODE=M193M;LINKAGE=E  
 NODE=M193M;LINKAGE=C  
 NODE=M193M;LINKAGE=A  
 NODE=M193M;LINKAGE=B

NODE=M193M;LINKAGE=D  
 NODE=M193M;LINKAGE=H  
 NODE=M193M;LINKAGE=AA  
 NODE=M193M;LINKAGE=F



## $\chi_{c1}(4140)$ WIDTH

NODE=M193W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>19 <math>\pm \frac{7}{5}</math> OUR AVERAGE</b>				
162 $\pm 21$ $^{+24}_{-49}$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
15.3 $^{+10.4}_{-6.1} \pm 2.5$	19	<sup>2</sup> AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$
16.3 $\pm 5.6 \pm 11.4$	616	<sup>3</sup> ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi \phi + \text{anything}$
20 $\pm 13$ $^{+3}_{-8}$	52	<sup>4</sup> ABAZOV	14A D0	$B^+ \rightarrow J/\psi \phi K^+$
28 $^{+15}_{-11} \pm 19$	0.3k	<sup>5</sup> CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
83 $\pm 21$ $^{+21}_{-14}$	4289	<sup>6,7</sup> AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
11.7 $^{+8.3}_{-5.0} \pm 3.7$	14	<sup>8,9</sup> AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi \phi K^+$
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 13 $\sigma$ .				
<sup>2</sup> Statistical significance of more than 5 $\sigma$ .				
<sup>3</sup> Statistical significance of more than 6 $\sigma$ .				
<sup>4</sup> Statistical significance of 3.1 $\sigma$ .				
<sup>5</sup> From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5 $\sigma$ .				
<sup>6</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.4 $\sigma$ .				
<sup>7</sup> Superseded by AAIJ 21E.				
<sup>8</sup> Statistical significance of 3.8 $\sigma$ .				
<sup>9</sup> Superseded by AALTONEN 17.				

NODE=M193W

NODE=M193W;LINKAGE=G  
 NODE=M193W;LINKAGE=E  
 NODE=M193W;LINKAGE=C  
 NODE=M193W;LINKAGE=A  
 NODE=M193W;LINKAGE=B

NODE=M193W;LINKAGE=D  
 NODE=M193W;LINKAGE=H  
 NODE=M193W;LINKAGE=AA  
 NODE=M193W;LINKAGE=F

 **$\chi_{c1}(4140)$  DECAY MODES**

NODE=M193215;NODE=M193

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi \phi$	seen
$\Gamma_2$ $\gamma\gamma$	not seen

DESIG=1

DESIG=2

 **$\chi_{c1}(4140)$   $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$** 

NODE=M193220

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
<b>&lt;41</b>	90	<sup>1</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi \phi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6	90	<sup>2</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi \phi$	
<sup>1</sup> For $J^P = 0^+$ .					
<sup>2</sup> For $J^P = 2^+$ .					

NODE=M193G01  
 NODE=M193G01

OCCUR=2

NODE=M193G01;LINKAGE=S0  
 NODE=M193G01;LINKAGE=S2

 **$\chi_{c1}(4140)$  BRANCHING RATIOS**

NODE=M193225

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>seen</b>	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$	
<b>seen</b>	616	<sup>2</sup> ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi \phi + \text{anything}$	
<b>seen</b>	52	<sup>3</sup> ABAZOV	14A D0	$B^+ \rightarrow J/\psi \phi K^+$	
<b>seen</b>	0.3k	<sup>4</sup> CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi \phi K^+$	
<b>seen</b>	14	<sup>5</sup> AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi \phi K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<b>seen</b>	4289	<sup>6,7</sup> AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$	
<b>not seen</b>		<sup>8</sup> ABLIKIM	15 BES3	$e^+ e^- \rightarrow \gamma \phi J/\psi$	
<b>not seen</b>		<sup>9</sup> AAIJ	12AA LHCB	$p\bar{p} \rightarrow B^+ X$ at 7 TeV	

NODE=M193R01  
 NODE=M193R01

NODE=M193R01;LINKAGE=F  
 NODE=M193R01;LINKAGE=D  
 NODE=M193R01;LINKAGE=A

NODE=M193R01;LINKAGE=C

NODE=M193R01;LINKAGE=AA  
 NODE=M193R01;LINKAGE=E  
 NODE=M193R01;LINKAGE=G  
 NODE=M193R01;LINKAGE=B

NODE=M193R01;LINKAGE=AI

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 13  $\sigma$ .<sup>2</sup> Statistical significance of more than 6  $\sigma$ .<sup>3</sup> ABAZOV 14A reports  $B(B^+ \rightarrow \chi_{c1}(4140) K^+ \rightarrow J/\psi \phi K^+)/B(B^+ \rightarrow J/\psi \phi K^+) = (19 \pm 7 \pm 4)\%$  with 3.1  $\sigma$  significance.<sup>4</sup> From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5  $\sigma$ .<sup>5</sup> Statistical significance of 3.8  $\sigma$ .<sup>6</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 8.4  $\sigma$ .<sup>7</sup> Superseded by AAIJ 21E.<sup>8</sup> Reported  $\sigma(e^+ e^- \rightarrow \gamma \chi_{c1}(4140)) \cdot B(\chi_{c1}(4140) \rightarrow J/\psi \phi) < 0.35, 0.28, \text{ and } 0.33$  pb at 4.23, 4.26, and 4.36 GeV, respectively, at 90% CL.<sup>9</sup> Reported  $B(B^+ \rightarrow \chi_{c1}(4140) K^+) \cdot B(\chi_{c1}(4140) \rightarrow J/\psi \phi)/B(B^+ \rightarrow J/\psi \phi K^+) < 0.07$  at 90% CL.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

NODE=M193R02  
 NODE=M193R02

 $\chi_{c1}(4140)$  REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
	Also	PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	17	MPL A32 1750139	T. Altonen <i>et al.</i>	(CDF Collab.)
ABAZOV	15M	PRL 115 232001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABLIKIM	15	PR D91 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABAZOV	14A	PR D89 012004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
CHATRCHYAN	14M	PL B734 261	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAIJ	12AA	PR D85 091103	R. Aaij <i>et al.</i>	(LHCb Collab.)
SHEN	10	PRL 104 112004	C.P. Shen <i>et al.</i>	(BELLE Collab.)
AALTONEN	09AH	PRL 102 242002	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=M193

REFID=61150  
 REFID=57657  
 REFID=57636  
 REFID=58161  
 REFID=56957  
 REFID=56368  
 REFID=55650  
 REFID=55753  
 REFID=54263  
 REFID=53235  
 REFID=52968

 $\psi(4160)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M025

 $\psi(4160)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>4191 ± 5 OUR AVERAGE</b>			
4191 $^{+9}_{-8}$	AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
4191.7 ± 6.5	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
4193 ± 7	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4151 ± 4	<sup>3</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4155 ± 5	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C DASP	$e^+e^-$

NODE=M025M

NODE=M025M

OCCUR=2

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

NODE=M025M;LINKAGE=AB

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

NODE=M025M;LINKAGE=MO

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M025M;LINKAGE=ST

<sup>4</sup> From a fit to BES (BAI 02C) data.

NODE=M025M;LINKAGE=SE

 $\psi(4160)$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>70 ± 10 OUR AVERAGE</b>			
65 $^{+22}_{-16}$	AAIJ	13BC LHCB	$B^+ \rightarrow K^+ \mu^+ \mu^-$
71.8 ± 12.3	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
79 ± 14	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
107 ± 10	<sup>3</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
107 ± 16	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
78 ± 20	BRANDELIK	78C DASP	$e^+e^-$

NODE=M025W

NODE=M025W

OCCUR=2

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

NODE=M025W;LINKAGE=AB

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

NODE=M025W;LINKAGE=MO

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M025W;LINKAGE=ST

<sup>4</sup> From a fit to BES (BAI 02C) data.

NODE=M025W;LINKAGE=SE

**$\psi(4160)$  DECAY MODES**

NODE=M025215;NODE=M025

NODE=M025

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, "seen" ("not seen") means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle's central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	
$\Gamma_1$ $e^+e^-$	$(6.9 \pm 3.3) \times 10^{-6}$		DESIG=1
$\Gamma_2$ $\mu^+\mu^-$	seen		DESIG=33
$\Gamma_3$ $D\bar{D}$	seen		DESIG=15;OUR EVAL;→ UNCHECKED ←
$\Gamma_4$ $D^0\bar{D}^0$	seen		DESIG=16
$\Gamma_5$ $D^+D^-$	seen		DESIG=17
$\Gamma_6$ $D^*\bar{D} + \text{c.c.}$	seen		DESIG=18;OUR EVAL;→ UNCHECKED ←
$\Gamma_7$ $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen		DESIG=19
$\Gamma_8$ $D^*(2010)^+D^- + \text{c.c.}$	seen		DESIG=20
$\Gamma_9$ $D^*\bar{D}^*$	seen		DESIG=21;OUR EVAL;→ UNCHECKED ←
$\Gamma_{10}$ $D^*(2007)^0\bar{D}^*(2007)^0$	seen		DESIG=22
$\Gamma_{11}$ $D^*(2010)^+D^*(2010)^-$	seen		DESIG=23
$\Gamma_{12}$ $D^0D^-\pi^+ + \text{c.c. (excl. } D^*(2007)^0\bar{D}^0 + \text{c.c., } D^*(2010)^+D^- + \text{c.c.)}$	not seen		DESIG=24
$\Gamma_{13}$ $D\bar{D}^*\pi + \text{c.c. (excl. } D^*\bar{D}^*)$	seen		DESIG=25
$\Gamma_{14}$ $D^0D^{*-}\pi^+ + \text{c.c. (excl. } D^*(2010)^+D^*(2010)^-)$	not seen		DESIG=26
$\Gamma_{15}$ $D_s^+D_s^-$	not seen		DESIG=27
$\Gamma_{16}$ $D_s^{*+}D_s^- + \text{c.c.}$	seen		DESIG=28
$\Gamma_{17}$ $J/\psi\pi^+\pi^-$	$< 3 \times 10^{-3}$	90%	DESIG=2
$\Gamma_{18}$ $J/\psi\pi^0\pi^0$	$< 3 \times 10^{-3}$	90%	DESIG=3
$\Gamma_{19}$ $J/\psi K^+K^-$	$< 2 \times 10^{-3}$	90%	DESIG=4
$\Gamma_{20}$ $J/\psi\eta$	$< 8 \times 10^{-3}$	90%	DESIG=5
$\Gamma_{21}$ $J/\psi\pi^0$	$< 1 \times 10^{-3}$	90%	DESIG=6
$\Gamma_{22}$ $J/\psi\eta'$	$< 5 \times 10^{-3}$	90%	DESIG=7
$\Gamma_{23}$ $J/\psi\pi^+\pi^-\pi^0$	$< 1 \times 10^{-3}$	90%	DESIG=8
$\Gamma_{24}$ $\psi(2S)\pi^+\pi^-$	$< 4 \times 10^{-3}$	90%	DESIG=9
$\Gamma_{25}$ $\chi_{c1}\gamma$	$< 5 \times 10^{-3}$	90%	DESIG=10
$\Gamma_{26}$ $\chi_{c2}\gamma$	$< 1.3 \%$	90%	DESIG=11
$\Gamma_{27}$ $\chi_{c1}\pi^+\pi^-\pi^0$	$< 2 \times 10^{-3}$	90%	DESIG=12
$\Gamma_{28}$ $\chi_{c2}\pi^+\pi^-\pi^0$	$< 8 \times 10^{-3}$	90%	DESIG=13
$\Gamma_{29}$ $h_c(1P)\pi^+\pi^-$	$< 5 \times 10^{-3}$	90%	DESIG=29
$\Gamma_{30}$ $h_c(1P)\pi^0\pi^0$	$< 2 \times 10^{-3}$	90%	DESIG=30
$\Gamma_{31}$ $h_c(1P)\eta$	$< 2 \times 10^{-3}$	90%	DESIG=31
$\Gamma_{32}$ $h_c(1P)\pi^0$	$< 4 \times 10^{-4}$	90%	DESIG=32
$\Gamma_{33}$ $\phi\pi^+\pi^-$	$< 2 \times 10^{-3}$	90%	DESIG=14
$\Gamma_{34}$ $\gamma\chi_{c1}(3872)$	$< 1.8 \times 10^{-3}$	90%	DESIG=44
$\Gamma_{35}$ $\gamma\chi_{c0}(3915) \rightarrow \gamma J/\psi\pi^+\pi^-$	$< 1.36 \times 10^{-4}$	90%	DESIG=35
$\Gamma_{36}$ $\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-$	$< 1.18 \times 10^{-4}$	90%	DESIG=36
$\Gamma_{37}$ $\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-$	$< 1.47 \times 10^{-4}$	90%	DESIG=37
$\Gamma_{38}$ $\gamma\chi_{c0}(3915) \rightarrow \gamma\gamma J/\psi$	$< 1.26 \times 10^{-4}$	90%	DESIG=39
$\Gamma_{39}$ $\gamma X(3930) \rightarrow \gamma\gamma J/\psi$	$< 8.8 \times 10^{-5}$	90%	DESIG=40
$\Gamma_{40}$ $\gamma X(3940) \rightarrow \gamma\gamma J/\psi$	$< 1.79 \times 10^{-4}$	90%	DESIG=41
$\Gamma_{41}$ $\omega\pi^0$	not seen		DESIG=47
$\Gamma_{42}$ $\omega\eta$	not seen		DESIG=48
$\Gamma_{43}$ $K^+K^-$			DESIG=42
$\Gamma_{44}$ $K_S^0 K^\pm \pi^\mp$			DESIG=43
$\Gamma_{45}$ $p\bar{p}p\bar{p}$	not seen		DESIG=45
$\Gamma_{46}$ $\Lambda\bar{\Lambda}$	$< 1.5 \times 10^{-6}$	90%	DESIG=46

$\psi(4160)$  PARTIAL WIDTHS

NODE=M025220

 $\Gamma(e^+e^-)$  $\Gamma_1$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.48±0.22</b>	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
••• We do not use the following data for averages, fits, limits, etc. •••			
0.4 to 1.1	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
0.83±0.08	<sup>3</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.84±0.13	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.77±0.23	BRANDELIK	78C DASP	$e^+e^-$

NODE=M025W1  
NODE=M025W1

OCCUR=2

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (293 \pm 57)^\circ$ .

NODE=M025W1;LINKAGE=AB

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different  $e^+e^-$  partial widths. We quote only the range of values.

NODE=M025W1;LINKAGE=MO

<sup>3</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M025W1;LINKAGE=ST

<sup>4</sup> From a fit to BES (BAI 02C) data.

NODE=M025W1;LINKAGE=SE

 $\Gamma(\mu^+\mu^-)$  $\Gamma_2$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>2.45±1.24±0.94</b>	<sup>1,2</sup> ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M025W2  
NODE=M025W2

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \mu^+\mu^-$  cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming  $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$ .

NODE=M025W2;LINKAGE=A

<sup>2</sup> From solution 1 of 8 with equal fit quality. Other solutions range from  $2.08 \pm 0.99 \pm 0.80$  to  $2.45 \pm 1.24 \pm 0.94$  keV.

NODE=M025W2;LINKAGE=B

 $\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M025235

 $\Gamma(J/\psi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{22}\Gamma_1/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.17±0.04	86	<sup>1,2</sup> ABLIKIM	20A BES3	$e^+e^- \rightarrow \eta' J/\psi$
1.07±0.09	86	<sup>1,3</sup> ABLIKIM	20A BES3	$e^+e^- \rightarrow \eta' J/\psi$

NODE=M025R42  
NODE=M025R42

••• We do not use the following data for averages, fits, limits, etc. •••

<sup>1</sup> Based on a fit to  $\sigma(e^+e^- \rightarrow \eta' J/\psi)$  from  $\sqrt{s} = 4.18$  to 4.60 GeV assuming interfering  $\psi(4160)$  and  $\psi(4260)$  contributions. At  $\sqrt{s} = 4.18$  GeV,  $\sigma(e^+e^- \rightarrow \eta' J/\psi) = 2.4 \pm 0.3 \pm 0.2$  pb.

OCCUR=2

NODE=M025R42;LINKAGE=A

<sup>2</sup> Solution I of the fit, corresponding to a phase of  $-0.03 \pm 0.44$  rad.

NODE=M025R42;LINKAGE=B

<sup>3</sup> Solution II of the fit, corresponding to a phase of  $2.54 \pm 0.04$  rad.

NODE=M025R42;LINKAGE=C

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{25}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.2</b>	90	<sup>1</sup> HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

NODE=M025G01  
NODE=M025G01

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

NODE=M025G01;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{26}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.1	90	<sup>1</sup> HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

NODE=M025G02  
NODE=M025G02

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

NODE=M025G02;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{44}\Gamma_1/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
2.71 ±0.13 ±0.12	<sup>1</sup> ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0095±0.0088±0.0004	<sup>2</sup> ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

NODE=M025R00  
NODE=M025R00

••• We do not use the following data for averages, fits, limits, etc. •••

OCCUR=5

<sup>1</sup> Solution I of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

NODE=M025R00;LINKAGE=A

<sup>2</sup> Solution II of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

NODE=M025R00;LINKAGE=D



$\psi(4160) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$ 

NODE=M025230

 $\Gamma(J/\psi\eta) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$  $\Gamma_{20} / \Gamma \times \Gamma_1 / \Gamma$ NODE=M025R32  
NODE=M025R32VALUE (units  $10^{-8}$ )

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $2.8 \pm 0.9 \pm 0.9$ <sup>1</sup> WANG

13B

BELL

 $e^+ e^- \rightarrow J/\psi \eta \gamma$  $12.8 \pm 1.7 \pm 2.0$ <sup>2</sup> WANG

13B

BELL

 $e^+ e^- \rightarrow J/\psi \eta \gamma$ 

OCCUR=2

<sup>1</sup> Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

NODE=M025R32;LINKAGE=A

<sup>2</sup> Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

NODE=M025R32;LINKAGE=B

 $\psi(4160)$  BRANCHING RATIOS

NODE=M025225

 $\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$  $\Gamma_2 / \Gamma$ 

VALUE

DOCUMENT ID

TECN

COMMENT

seen

<sup>1</sup> AAIJ

13BC

LHCB

 $B^+ \rightarrow K^+ \mu^+ \mu^-$ NODE=M025R31  
NODE=M025R31<sup>1</sup> AAIJ 13BC report  $B(B^+ \rightarrow K^+ \psi(4160)) B(\psi(4160) \rightarrow \mu^+ \mu^-) = (3.5_{-0.8}^{+0.9}) \times 10^{-9}$ .

NODE=M025R31;LINKAGE=A

 $\Gamma(D\bar{D}) / \Gamma(D^* \bar{D}^*)$  $\Gamma_3 / \Gamma_9$ 

VALUE

DOCUMENT ID

TECN

COMMENT

 $0.02 \pm 0.03 \pm 0.02$ 

AUBERT

09M

BABR

 $e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$ NODE=M025R14  
NODE=M025R14 $\Gamma(D^0 \bar{D}^0) / \Gamma_{\text{total}}$  $\Gamma_4 / \Gamma$ 

VALUE

DOCUMENT ID

TECN

COMMENT

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^0 \bar{D}^0$ 

seen

PAKHLOVA 08

BELL

 $e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$ NODE=M025R16  
NODE=M025R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^0 \bar{D}^0 \gamma$  $\Gamma(D^+ D^-) / \Gamma_{\text{total}}$  $\Gamma_5 / \Gamma$ 

VALUE

DOCUMENT ID

TECN

COMMENT

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^+ D^-$ 

seen

PAKHLOVA 08

BELL

 $e^+ e^- \rightarrow D^+ D^- \gamma$ NODE=M025R17  
NODE=M025R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^+ D^- \gamma$  $\Gamma(D^*(2007)^0 \bar{D}^0 + \text{c.c.}) / \Gamma_{\text{total}}$  $\Gamma_7 / \Gamma$ 

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^{*0} \bar{D}^0 \gamma$ 

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^{*0} \bar{D}^0$ NODE=M025R18  
NODE=M025R18 $\Gamma(D^*(2010)^+ D^- + \text{c.c.}) / \Gamma_{\text{total}}$  $\Gamma_8 / \Gamma$ 

VALUE

DOCUMENT ID

TECN

COMMENT

seen

<sup>1</sup> ZHUKOVA

18

BELL

 $e^+ e^- \rightarrow D^{*+} D^- \gamma$ 

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^{*+} D^- \gamma$ 

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^{*+} D^-$ NODE=M025R19  
NODE=M025R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

PAKHLOVA

07

BELL

 $e^+ e^- \rightarrow D^{*+} D^- \gamma$ <sup>1</sup> Supersedes PAKHLOVA 07.

NODE=M025R19;LINKAGE=A

 $\Gamma(D^* \bar{D} + \text{c.c.}) / \Gamma(D^* \bar{D}^*)$  $\Gamma_6 / \Gamma_9$ 

VALUE

DOCUMENT ID

TECN

COMMENT

 $0.34 \pm 0.14 \pm 0.05$ 

AUBERT

09M

BABR

 $e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$ NODE=M025R15  
NODE=M025R15 $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0) / \Gamma_{\text{total}}$  $\Gamma_{10} / \Gamma$ 

VALUE

DOCUMENT ID

TECN

COMMENT

seen

AUBERT

09M

BABR

 $e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$ 

seen

CRONIN-HEN..09

CLEO

 $e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$ NODE=M025R20  
NODE=M025R20

$\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$   $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

NODE=M025R21  
 NODE=M025R21

<sup>1</sup>Supersedes PAKHLOVA 07.

NODE=M025R21;LINKAGE=A

 $\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.}))/\Gamma_{\text{total}}$   $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 08A	BELL	$e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

NODE=M025R22  
 NODE=M025R22

 $\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$   $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D \bar{D}^* \pi$

NODE=M025R23  
 NODE=M025R23

 $\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

NODE=M025R24  
 NODE=M025R24

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$

NODE=M025R25  
 NODE=M025R25

 $\Gamma(D_s^{*+} D_s^- + \text{c.c.}))/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$

NODE=M025R26  
 NODE=M025R26

 $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

NODE=M025R01  
 NODE=M025R01

 $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

NODE=M025R02  
 NODE=M025R02

 $\Gamma(J/\psi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

NODE=M025R03  
 NODE=M025R03

 $\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

NODE=M025R04  
 NODE=M025R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen <sup>1</sup> ABLIKIM 15L BES3  $e^+ e^- \rightarrow J/\psi \eta$   
 seen WANG 13B BELL  $e^+ e^- \rightarrow J/\psi \eta \gamma$

<sup>1</sup> An enhancement around 4.2 GeV is observed.

NODE=M025R04;LINKAGE=A

 $\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

NODE=M025R05  
 NODE=M025R05

 $\Gamma(J/\psi \eta')/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	COAN 06	CLEO	4.12–4.2 $e^+ e^- \rightarrow$ hadrons

NODE=M025R06  
 NODE=M025R06

$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R07  
NODE=M025R07

 $\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{24}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R08  
NODE=M025R08

 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$   $\Gamma_{25}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<7	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons
----	----	------	----	--

NODE=M025R09  
NODE=M025R09

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$   $\Gamma_{26}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<13	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R10  
NODE=M025R10

 $\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R11  
NODE=M025R11

 $\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R12  
NODE=M025R12

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

NODE=M025R27  
NODE=M025R27

NODE=M025R27;LINKAGE=PE

 $\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

NODE=M025R28  
NODE=M025R28

NODE=M025R28;LINKAGE=PE

 $\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2	90		<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen 41 <sup>2</sup> ABLIKIM 17R BES3  $e^+e^- \rightarrow h_c(1P)\eta$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

<sup>2</sup> An enhancement around 4.2 GeV is observed.

NODE=M025R29  
NODE=M025R29

NODE=M025R29;LINKAGE=PE

NODE=M025R29;LINKAGE=A

 $\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	<sup>1</sup> PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^0$

<sup>1</sup> At  $\sqrt{s} = 4170$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

NODE=M025R30  
NODE=M025R30

NODE=M025R30;LINKAGE=PE

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 4.12–4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R13  
NODE=M025R13

$\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{total}$					$\Gamma_{34}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.8 \times 10^{-3}$	90	1,2 XIAO	13	$\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$	NODE=M025R43 NODE=M025R43
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<0.012$	90	1,3 XIAO	13	$\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$	OCCUR=2
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
<sup>2</sup> XIAO 13 reports $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{total}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 0.68 \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 3.8 \times 10^{-2}$ .					
<sup>3</sup> XIAO 13 reports $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{total}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 1.05 \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 8 \times 10^{-3}$ .					
$\Gamma(\gamma\chi_{c0}(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-)/\Gamma_{total}$					$\Gamma_{35}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.36 \times 10^{-4}$	90	1 XIAO	13	$\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$	NODE=M025R35 NODE=M025R35
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-)/\Gamma_{total}$					$\Gamma_{36}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.18 \times 10^{-4}$	90	1 XIAO	13	$\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$	NODE=M025R36 NODE=M025R36
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-)/\Gamma_{total}$					$\Gamma_{37}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.47 \times 10^{-4}$	90	1 XIAO	13	$\psi(4160) \rightarrow \gamma J/\psi \pi^+ \pi^-$	NODE=M025R37 NODE=M025R37
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma\chi_{c0}(3915) \rightarrow \gamma\gamma J/\psi)/\Gamma_{total}$					$\Gamma_{38}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.26 \times 10^{-4}$	90	1 XIAO	13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$	NODE=M025R39 NODE=M025R39
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3930) \rightarrow \gamma\gamma J/\psi)/\Gamma_{total}$					$\Gamma_{39}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<0.88 \times 10^{-4}$	90	1 XIAO	13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$	NODE=M025R40 NODE=M025R40
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\gamma X(3940) \rightarrow \gamma\gamma J/\psi)/\Gamma_{total}$					$\Gamma_{40}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.79 \times 10^{-4}$	90	1 XIAO	13	$\psi(4160) \rightarrow \gamma\gamma J/\psi$	NODE=M025R41 NODE=M025R41
<sup>1</sup> Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.					
$\Gamma(\omega\pi^0)/\Gamma_{total}$					$\Gamma_{41}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
not seen		ABLIKIM	22K	BES3 $e^+ e^- \rightarrow \omega\pi^0$	NODE=M025R46 NODE=M025R46
$\Gamma(\omega\eta)/\Gamma_{total}$					$\Gamma_{42}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
not seen		ABLIKIM	22K	BES3 $e^+ e^- \rightarrow \omega\eta$	NODE=M025R47 NODE=M025R47
$\Gamma(K^+ K^-)/\Gamma_{total}$					$\Gamma_{43}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2 \times 10^{-5}$	90	1 DRUZHININ	15	RVUE $e^+ e^- \rightarrow \psi(3770)$	NODE=M025R33 NODE=M025R33
<sup>1</sup> DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+ e^- \rightarrow K^+ K^-$ and $e^+ e^- \rightarrow K_S^0 K_L^0$ .					
$\Gamma(p\bar{p}p\bar{p})/\Gamma_{total}$					$\Gamma_{45}/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
not seen		ABLIKIM	21D	BES3 $4.0-4.6 e^+ e^- \rightarrow p\bar{p}p\bar{p}$	NODE=M025R44 NODE=M025R44

$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$		$\Gamma_{46}\Gamma_1/\Gamma$		
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<0.7 \times 10^{-3}$	90	<sup>1</sup> ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4160)$

<sup>1</sup> From a measurement of the  $e^+e^- \rightarrow \Lambda\bar{\Lambda}$  cross section between 3.5 and 4.6 GeV.

NODE=M025R45  
NODE=M025R45

NODE=M025R45;LINKAGE=A

NODE=M025

### $\psi(4160)$ REFERENCES

ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21D	PR D103 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61031
ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60210
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59856
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
ABLIKIM	17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58009
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan		REFID=57991
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56777
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)	REFID=56962
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55229
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
XIAO	13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)	REFID=55381
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51075
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232

NODE=M190

## X(4160)

$$I^G(J^{PC}) = ?^?(?^{??})$$

### OMITTED FROM SUMMARY TABLE

Seen by PAKHLOV 08 in  $e^+e^- \rightarrow J/\psi X$ ,  $X \rightarrow D^*\bar{D}^*$

NODE=M190

A state with consistent mass and width is seen by AAIJ 21E in  $B^+ \rightarrow X(4160)K^+$  with  $X(4160) \rightarrow J/\psi\phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  with a significance (accounting for systematic uncertainties) of  $4.8\sigma$ . The  $J^{PC} = 2^{-+}$  assignment is favored over other assignments with a significance of more than  $4\sigma$ .

### X(4160) MASS

NODE=M190M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4153^{+23}_{-21}</math> OUR AVERAGE</b>				
$4146 \pm 18 \pm 33$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
$4156^{+25}_{-20} \pm 15$	24	PAKHLOV	08 BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M190M

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $4.8\sigma$ .

NODE=M190M;LINKAGE=A

### X(4160) WIDTH

NODE=M190W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>136^{+60}_{-35}</math> OUR AVERAGE</b>				
$135 \pm 28^{+59}_{-30}$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
$139^{+111}_{-61} \pm 21$	24	PAKHLOV	08 BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M190W

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $4.8\sigma$ .

NODE=M190W;LINKAGE=A

**X(4160) DECAY MODES**

NODE=M190215;NODE=M190

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D\bar{D}$	not seen
$\Gamma_2$ $D^*\bar{D} + \text{c.c.}$	not seen
$\Gamma_3$ $D^*\bar{D}^*$	seen
$\Gamma_4$ $J/\psi\phi$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=3;OUR EVAL;→ UNCHECKED ←  
DESIG=4

**X(4160) BRANCHING RATIOS**

NODE=M190225

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$					$\Gamma_1/\Gamma_3$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.09</b>	90	PAKHLOV	08	BELL $e^+e^- \rightarrow J/\psi X$	

NODE=M190R01  
NODE=M190R01

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$					$\Gamma_2/\Gamma_3$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.22</b>	90	PAKHLOV	08	BELL $e^+e^- \rightarrow J/\psi X$	

NODE=M190R02  
NODE=M190R02

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
<b>seen</b>	24k	<sup>1</sup> AAIJ	21E	LHCB $B^+ \rightarrow J/\psi\phi K^+$	

NODE=M190R00  
NODE=M190R00

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 4.8  $\sigma$ .

NODE=M190R00;LINKAGE=A

**X(4160) REFERENCES**

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
PAKHLOV	08	PRL 100 202001	P. Pakhlov <i>et al.</i>	(BELLE Collab.)

NODE=M190

REFID=61150  
REFID=52302

NODE=M074

 **$\psi(4230)$** 

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as  $Y(4230)$ ; was  $\psi(4260)$

NODE=M074

The original  $\psi(4260)$  (also known as  $Y(4260)$ ) was observed by AUBERT,B 05I as a peak in the energy dependence of the  $e^+e^- \rightarrow \pi^+\pi^- J/\psi$  cross section and was confirmed by HE 06B, YUAN 07, LEES 12AC, and LIU 13B in the same process. A higher-statistics analysis by ABLIKIM 17B revealed an asymmetry in the cross section and resulted in a shift of the peak position to a lower mass. The  $\psi(4260)$  was therefore renamed  $\psi(4230)$ . The energy-dependent cross sections for  $e^+e^-$  to other channels also exhibit peaks in the same mass region. The parameters corresponding to those peaks are also listed here, but the number of states in this region remains to be determined.

For details see the review on "Spectroscopy of mesons containing two heavy quarks."

 **$\psi(4230)$  MASS**

NODE=M074M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>4222.5 \pm 2.4</math> OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below. [4222.7 $\pm$ 2.6 MeV OUR 2022 AVERAGE Scale factor = 1.7]		
$4221.4 \pm 1.5 \pm 2.0$	1	ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
$4225.3 \pm 2.3 \pm 21.5$	2	ABLIKIM	22AU BES3	$e^+e^- \rightarrow K^+K^- J/\psi$
$4234.4 \pm 3.2 \pm 0.2$	3	ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
$4216.7 \pm 8.9 \pm 4.1$	4	ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$
$4220.4 \pm 2.4 \pm 2.3$	5	ABLIKIM	20N BES3	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
$4218.6 \pm 3.8 \pm 2.5$	5	ABLIKIM	20O BES3	$e^+e^- \rightarrow \eta J/\psi$
$4218.5 \pm 1.6 \pm 4.0$	6	ABLIKIM	19AI BES3	$e^+e^- \rightarrow \omega\chi_{c0}$
$4228.6 \pm 4.1 \pm 6.3$		ABLIKIM	19R BES3	$e^+e^- \rightarrow \pi^+D^0 D^{*-} + \text{c.c.}$
$4200.6^{+7.9}_{-13.3} \pm 3.0$	7	ABLIKIM	19V BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$
$4218^{+5.5}_{-4.5} \pm 0.9$		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^- h_c$

NODE=M074M


NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

4231.9 ± 5.3 ± 4.9	ABLIKIM	20N	BES3	$e^+e^- \rightarrow \pi^0 Z_c(3900)^0, Z_c^0 \rightarrow \pi^0 J/\psi$	OCCUR=2
4222.0 ± 3.1 ± 1.4	<sup>8</sup> ABLIKIM	17B	BES3	$e^+e^- \rightarrow \pi^+ \pi^- J/\psi$	
4209.5 ± 7.4 ± 1.4	<sup>9</sup> ABLIKIM	17V	BES3	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
4209.1 ± 6.8 ± 7.0	<sup>10</sup> ZHANG	17B	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
4223.3 ± 1.6 ± 2.5	<sup>11</sup> ZHANG	17C	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$	OCCUR=2
4230 ± 8 ± 6 180	<sup>12</sup> ABLIKIM	15C	BES3	$e^+e^- \rightarrow \omega \chi_{c0}$	
4258.6 ± 8.3 ± 12.1	<sup>13</sup> LIU	13B	BELL	$e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
4245 ± 5 ± 4	<sup>14</sup> LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
4247 ± 12 $\begin{smallmatrix} +17 \\ -32 \end{smallmatrix}$	<sup>13,15</sup> YUAN	07	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
4284 $\begin{smallmatrix} +17 \\ -16 \end{smallmatrix}$ ± 413.6	HE	06B	CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
4259 ± 8 $\begin{smallmatrix} + \\ - \end{smallmatrix} \frac{2}{6}$ 125	<sup>16</sup> AUBERT,B	05I	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	

- <sup>1</sup> From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 3.7730\text{--}4.7008$  GeV.
- <sup>2</sup> From a two-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.127\text{--}4.600$  GeV. The second resonance has a mass of  $4484.7 \pm 13.3 \pm 24.1$  MeV and a total width of  $111.1 \pm 30.1 \pm 15.2$  MeV.
- <sup>3</sup> From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.008\text{--}4.698$  GeV.
- <sup>4</sup> Solution 1 of 8 with equal fit quality to the  $e^+e^- \rightarrow \mu^+\mu^-$  cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes. Other solutions range from  $4212.8 \pm 7.2 \pm 4.0$  to  $4219.4 \pm 11.2 \pm 4.1$  MeV.
- <sup>5</sup> From a fit of the measured cross section in the range  $\sqrt{s} = 3.808\text{--}4.600$  GeV.
- <sup>6</sup> From a fit of the measured cross section from  $\sqrt{s} = 4.178\text{--}4.278$  GeV. Supersedes ABLIKIM 15C.
- <sup>7</sup> Simultaneous fit to  $\chi_{c1} \rightarrow \omega J/\psi$  and  $\chi_{c1} \rightarrow \pi^+ \pi^- J/\psi$ .
- <sup>8</sup> From a three-resonance fit. Superseded by ABLIKIM 22AM.
- <sup>9</sup> From a fit to the cross section for  $e^+e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-) \ell^+ \ell^-$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1 \text{ fb}^{-1}$ . Superseded by ABLIKIM 21AJ.
- <sup>10</sup> From a three-resonance fit.
- <sup>11</sup> From a combined fit of BELLE, BABAR and BES3  $e^+e^- \rightarrow \pi^+ \pi^- J/\psi$  and  $e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$  data.
- <sup>12</sup> From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+ \pi^-$ ,  $\chi_{c0} \rightarrow K^+ K^-$ , and  $\omega \rightarrow \pi^+ \pi^- \pi^0$ .
- <sup>13</sup> From a two-resonance fit.
- <sup>14</sup> From a single-resonance fit. Supersedes AUBERT,B 05I.
- <sup>15</sup> Superseded by LIU 13B.
- <sup>16</sup> From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074M;LINKAGE=F  
 NODE=M074M;LINKAGE=H  
 NODE=M074M;LINKAGE=C  
 NODE=M074M;LINKAGE=HP  
 NODE=M074M;LINKAGE=GP  
 NODE=M074M;LINKAGE=CP  
 NODE=M074M;LINKAGE=FP  
 NODE=M074M;LINKAGE=G  
 NODE=M074M;LINKAGE=BP  
 NODE=M074M;LINKAGE=A  
 NODE=M074M;LINKAGE=CA  
 NODE=M074M;LINKAGE=AP  
 NODE=M074M;LINKAGE=YU  
 NODE=M074M;LINKAGE=LE  
 NODE=M074M;LINKAGE=YN  
 NODE=M074M;LINKAGE=AU



$\psi(4230)$  WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>48 ± 8 OUR AVERAGE</b>				Error includes scale factor of 3.6. See the ideogram below.
[49 ± 8 MeV OUR 2022 AVERAGE				Scale factor = 3.5]
41.8 ± 2.9 ± 2.7	1	ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
72.9 ± 6.1 ± 30.8	2	ABLIKIM	22AU BES3	$e^+e^- \rightarrow K^+K^- J/\psi$
17.6 ± 18.1 ± 0.9	3	ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
47.2 ± 22.8 ± 10.5	4	ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$
46.2 ± 4.7 ± 2.1	5	ABLIKIM	20N BES3	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
82.0 ± 5.7 ± 0.4	5	ABLIKIM	20O BES3	$e^+e^- \rightarrow \eta J/\psi$
28.2 ± 3.9 ± 1.6	6	ABLIKIM	19AI BES3	$e^+e^- \rightarrow \omega\chi_{c0}$
77.0 ± 6.8 ± 6.3		ABLIKIM	19R BES3	$e^+e^- \rightarrow \pi^+D^0 D^{*-} + \text{c.c.}$
115 $\begin{smallmatrix} +38 \\ -26 \end{smallmatrix}$ ± 12	7	ABLIKIM	19V BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$
66.0 $\begin{smallmatrix} +12.3 \\ -8.3 \end{smallmatrix}$ ± 0.4		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^- h_c$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
41.2 ± 16.0 ± 16.4		ABLIKIM	20N BES3	$e^+e^- \rightarrow \pi^0 Z_c(3900)^0, Z_c^0 \rightarrow \pi^0 J/\psi$
44.1 ± 4.3 ± 2.0	8	ABLIKIM	17B BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
80.1 ± 24.6 ± 2.9	9	ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
76.6 ± 14.2 ± 2.4	10	ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
54.2 ± 2.6 ± 1.0	11	ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
38 ± 12 ± 2 180	12	ABLIKIM	15C BES3	$e^+e^- \rightarrow \omega\chi_{c0}$
134.1 ± 16.4 ± 5.5	13	LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
114 $\begin{smallmatrix} +16 \\ -15 \end{smallmatrix}$ ± 7	14	LEES	12AC BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
108 ± 19 ± 10	13,15	YUAN	07 BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
73 $\begin{smallmatrix} +39 \\ -25 \end{smallmatrix}$ ± 5 13.6		HE	06B CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
88 ± 23 $\begin{smallmatrix} +6 \\ -4 \end{smallmatrix}$ 125	16	AUBERT,B	05I BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
<p><sup>1</sup> From a three-resonance fit to the Born cross section in the range <math>\sqrt{s} = 3.7730\text{--}4.7008</math> GeV.</p> <p><sup>2</sup> From a two-resonance fit to the Born cross section in the range <math>\sqrt{s} = 4.127\text{--}4.600</math> GeV. The second resonance has a mass of <math>4484.7 \pm 13.3 \pm 24.1</math> MeV and a total width of <math>111.1 \pm 30.1 \pm 15.2</math> MeV.</p> <p><sup>3</sup> From a three-resonance fit to the Born cross section in the range <math>\sqrt{s} = 4.008\text{--}4.698</math> GeV.</p> <p><sup>4</sup> Solution 1 of 8 with equal fit quality to the <math>e^+e^- \rightarrow \mu^+\mu^-</math> cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes. Other solutions range from <math>36.4 \pm 16.8 \pm 8.1</math> to <math>49.6 \pm 22.6 \pm 11.0</math> MeV.</p> <p><sup>5</sup> From a fit of the measured cross section in the range <math>\sqrt{s} = 3.808\text{--}4.600</math> GeV.</p> <p><sup>6</sup> From a fit of the measured cross section from <math>\sqrt{s} = 4.178\text{--}4.278</math> GeV. Supersedes ABLIKIM 15C.</p> <p><sup>7</sup> Simultaneous fit to <math>\chi_{c1} \rightarrow \omega J/\psi</math> and <math>\chi_{c1} \rightarrow \pi^+\pi^- J/\psi</math>.</p> <p><sup>8</sup> From a three-resonance fit. Superseded by ABLIKIM 22AM.</p> <p><sup>9</sup> From a fit to the cross section for <math>e^+e^- \rightarrow \pi^+\pi^- \psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-</math> obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising <math>5.1 \text{ fb}^{-1}</math>. Superseded by ABLIKIM 21AJ.</p> <p><sup>10</sup> From a three-resonance fit.</p> <p><sup>11</sup> From a combined fit of BELLE, BABAR and BES3 <math>e^+e^- \rightarrow \pi^+\pi^- J/\psi</math> and <math>e^+e^- \rightarrow \pi^+\pi^- \psi(2S)</math> data.</p> <p><sup>12</sup> From a 3-parameter fit of measured cross sections from <math>\sqrt{s} = 4.21\text{--}4.42</math> GeV to a phase-space modified Breit-Wigner function, using the decays <math>\chi_{c0} \rightarrow \pi^+\pi^-</math>, <math>\chi_{c0} \rightarrow K^+K^-</math>, and <math>\omega \rightarrow \pi^+\pi^-\pi^0</math>.</p> <p><sup>13</sup> From a two-resonance fit.</p> <p><sup>14</sup> From a single-resonance fit. Supersedes AUBERT,B 05I.</p> <p><sup>15</sup> Superseded by LIU 13B.</p> <p><sup>16</sup> From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.</p>				

NODE=M074W

NODE=M074W

NEW

OCCUR=2

NODE=M074W;LINKAGE=F

NODE=M074W;LINKAGE=G

NODE=M074W;LINKAGE=D

NODE=M074W;LINKAGE=GP

NODE=M074W;LINKAGE=FP

NODE=M074W;LINKAGE=CP

NODE=M074W;LINKAGE=EP

NODE=M074W;LINKAGE=E

NODE=M074W;LINKAGE=BP

NODE=M074W;LINKAGE=C

NODE=M074W;LINKAGE=B

NODE=M074W;LINKAGE=AP


NODE=M074W;LINKAGE=YU

NODE=M074W;LINKAGE=LE

NODE=M074W;LINKAGE=YN

NODE=M074W;LINKAGE=AU





### $\psi(4230)$ DECAY MODES

NODE=M074215;NODE=M074

Mode	Fraction ( $\Gamma_i/\Gamma$ )	
$\Gamma_1$ $e^+ e^-$		DESIG=1
$\Gamma_2$ $\mu^+ \mu^-$	$(3.2 \pm 2.9) \times 10^{-5}$	DESIG=63
$\Gamma_3$ $\eta_c(1S) \pi^+ \pi^-$	not seen	DESIG=65
$\Gamma_4$ $\eta_c(1S) \pi^+ \pi^- \pi^0$	seen	DESIG=64
$\Gamma_5$ $J/\psi \pi^+ \pi^-$	seen	DESIG=2
$\Gamma_6$ $J/\psi f_0(980), f_0(980) \rightarrow \pi^+ \pi^-$	seen	DESIG=41;OUR EVAL;→ UNCHECKED ←
$\Gamma_7$ $Z_c(3900)^\pm \pi^\mp, Z_c^\pm \rightarrow J/\psi \pi^\pm$	seen	DESIG=43;OUR EVAL;→ UNCHECKED ←
$\Gamma_8$ $J/\psi \pi^0 \pi^0$	seen	DESIG=4
$\Gamma_9$ $J/\psi K^+ K^-$	seen	DESIG=5;OUR EVAL;→ UNCHECKED ←
$\Gamma_{10}$ $J/\psi K_S^0 K_S^0$	not seen	DESIG=44
$\Gamma_{11}$ $J/\psi \eta$	seen	DESIG=6
$\Gamma_{12}$ $J/\psi \pi^0$	not seen	DESIG=7;OUR EVAL;→ UNCHECKED ←
$\Gamma_{13}$ $J/\psi \eta'$	seen	DESIG=8;OUR EVAL;→ UNCHECKED ←
$\Gamma_{14}$ $J/\psi \pi^+ \pi^- \pi^0$	not seen	DESIG=9;OUR EVAL;→ UNCHECKED ←
$\Gamma_{15}$ $J/\psi \eta \pi^0$	not seen	DESIG=45
$\Gamma_{16}$ $J/\psi \eta \eta$	not seen	DESIG=10;OUR EVAL;→ UNCHECKED ←
$\Gamma_{17}$ $\psi(2S) \pi^+ \pi^-$	seen	DESIG=11
$\Gamma_{18}$ $\psi(2S) \eta$	not seen	DESIG=12;OUR EVAL;→ UNCHECKED ←
$\Gamma_{19}$ $\chi_{c0} \omega$	seen	DESIG=13
$\Gamma_{20}$ $\chi_{c1} \pi^+ \pi^- \pi^0$	not seen	DESIG=16;OUR EVAL;→ UNCHECKED ←
$\Gamma_{21}$ $\chi_{c2} \pi^+ \pi^- \pi^0$	not seen	DESIG=17;OUR EVAL;→ UNCHECKED ←
$\Gamma_{22}$ $h_c(1P) \pi^+ \pi^-$	seen	DESIG=40
$\Gamma_{23}$ $\phi \pi^+ \pi^-$	not seen	DESIG=18;OUR EVAL;→ UNCHECKED ←
$\Gamma_{24}$ $\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	not seen	DESIG=22;OUR EVAL;→ UNCHECKED ←
$\Gamma_{25}$ $D \bar{D}$	not seen	DESIG=19;OUR EVAL;→ UNCHECKED ←
$\Gamma_{26}$ $D^0 \bar{D}^0$	not seen	DESIG=31
$\Gamma_{27}$ $D^+ D^-$	not seen	DESIG=32
$\Gamma_{28}$ $D^* \bar{D} + c.c.$	not seen	DESIG=23;OUR EVAL;→ UNCHECKED ←
$\Gamma_{29}$ $D^*(2007)^0 \bar{D}^0 + c.c.$	not seen	DESIG=33
$\Gamma_{30}$ $D^*(2010)^+ D^- + c.c.$	not seen	DESIG=34
$\Gamma_{31}$ $D^* \bar{D}^*$		DESIG=24
$\Gamma_{32}$ $D^*(2007)^0 \bar{D}^*(2007)^0$	not seen	DESIG=35
$\Gamma_{33}$ $D^*(2010)^+ D^*(2010)^-$	not seen	DESIG=36
$\Gamma_{34}$ $D \bar{D} \pi + c.c.$		DESIG=37

Γ <sub>35</sub>	$D^0 D^- \pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0 \bar{D}^{*0} + \text{c.c.}$ , $D^*(2010)^+ D^- + \text{c.c.}$ )	not seen	DESIG=38
Γ <sub>36</sub>	$D \bar{D}^* \pi + \text{c.c.}$ (excl. $D^* \bar{D}^*$ )	not seen	DESIG=25
Γ <sub>37</sub>	$D^0 D^{*-} \pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+ D^*(2010)^-$ )	not seen	DESIG=39
Γ <sub>38</sub>	$D^0 D^*(2010)^- \pi^+ + \text{c.c.}$	seen	DESIG=30
Γ <sub>39</sub>	$D_1(2420) \bar{D} + \text{c.c.}$	not seen	DESIG=50
Γ <sub>40</sub>	$D^* \bar{D}^* \pi$	not seen	DESIG=26
Γ <sub>41</sub>	$D_s^+ D_s^-$	not seen	DESIG=27
Γ <sub>42</sub>	$D_s^{*+} D_s^- + \text{c.c.}$	not seen	DESIG=28
Γ <sub>43</sub>	$D_s^{*+} D_s^{*-}$	not seen	DESIG=29
Γ <sub>44</sub>	$p \bar{p}$	not seen	DESIG=3;OUR EVAL;→ UNCHECKED ←
Γ <sub>45</sub>	$p \bar{p} \pi^0$	not seen	DESIG=46;OUR EVAL;→ UNCHECKED ←
Γ <sub>46</sub>	$p \bar{p} \eta$	not seen	DESIG=61
Γ <sub>47</sub>	$p \bar{p} \omega$	not seen	DESIG=62
Γ <sub>48</sub>	$\Xi^- \bar{\Xi}^+$	not seen	DESIG=51;OUR EVAL;→ UNCHECKED ←
Γ <sub>49</sub>	$\pi^+ \pi^+ \pi^- \pi^-$	not seen	DESIG=53;OUR EVAL;→ UNCHECKED ←
Γ <sub>50</sub>	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	not seen	DESIG=54;OUR EVAL;→ UNCHECKED ←
Γ <sub>51</sub>	$\omega \pi^0$	not seen	DESIG=68
Γ <sub>52</sub>	$\omega \eta$	not seen	DESIG=69
Γ <sub>53</sub>	$K_S^0 K^\pm \pi^\mp$	not seen	DESIG=20;OUR EVAL;→ UNCHECKED ←
Γ <sub>54</sub>	$K_S^0 K^\pm \pi^\mp \pi^0$	not seen	DESIG=48;OUR EVAL;→ UNCHECKED ←
Γ <sub>55</sub>	$K_S^0 K^\pm \pi^\mp \eta$	not seen	DESIG=49;OUR EVAL;→ UNCHECKED ←
Γ <sub>56</sub>	$K^+ K^- \pi^0$	not seen	DESIG=21;OUR EVAL;→ UNCHECKED ←
Γ <sub>57</sub>	$K^+ K^- \pi^+ \pi^-$	not seen	DESIG=55;OUR EVAL;→ UNCHECKED ←
Γ <sub>58</sub>	$K^+ K^- \pi^+ \pi^- \pi^0$	not seen	DESIG=56;OUR EVAL;→ UNCHECKED ←
Γ <sub>59</sub>	$K^+ K^+ K^- K^-$	not seen	DESIG=57;OUR EVAL;→ UNCHECKED ←
Γ <sub>60</sub>	$K^+ K^+ K^- K^- \pi^0$	not seen	DESIG=58;OUR EVAL;→ UNCHECKED ←
Γ <sub>61</sub>	$p \bar{p} \pi^+ \pi^-$	not seen	DESIG=59;OUR EVAL;→ UNCHECKED ←
Γ <sub>62</sub>	$p \bar{p} \pi^+ \pi^- \pi^0$	not seen	DESIG=60;OUR EVAL;→ UNCHECKED ←
Γ <sub>63</sub>	$p \bar{p} p \bar{p}$	not seen	DESIG=67
Γ <sub>64</sub>	$\Lambda \bar{\Lambda}$	not seen	DESIG=52;OUR EVAL;→ UNCHECKED ←

## Radiative decays

Γ <sub>65</sub>	$\eta_c(1S) \gamma$	possibly seen	NODE=M074;CLUMP=C DESIG=47
Γ <sub>66</sub>	$\eta_c(1S) \pi^0 \gamma$	not seen	DESIG=66
Γ <sub>67</sub>	$\chi_{c1} \gamma$	not seen	DESIG=14;OUR EVAL;→ UNCHECKED ←
Γ <sub>68</sub>	$\chi_{c2} \gamma$	not seen	DESIG=15;OUR EVAL;→ UNCHECKED ←
Γ <sub>69</sub>	$\chi_{c1}(3872) \gamma$	seen	DESIG=42

 $\psi(4230)$  PARTIAL WIDTHS $\Gamma(\mu^+ \mu^-)$ Γ<sub>2</sub>

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>1.53 ± 1.26 ± 0.54</b>	<sup>1,2</sup> ABLIKIM	20AG BES3	$e^+ e^- \rightarrow \mu^+ \mu^-$

<sup>1</sup> From a fit to the  $e^+ e^- \rightarrow \mu^+ \mu^-$  cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming  $\Gamma(\mu^+ \mu^-) = \Gamma(e^+ e^-)$ .

<sup>2</sup> From solution 1 of 8 with equal fit quality. Other solutions range from  $1.09 \pm 0.84 \pm 0.39$  to  $1.53 \pm 1.26 \pm 0.54$  keV.

NODE=M074235

NODE=M074W01  
NODE=M074W01

NODE=M074W01;LINKAGE=A

NODE=M074W01;LINKAGE=B

 $\psi(4230) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$ 

NODE=M074230

 $\Gamma(J/\psi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ Γ<sub>5</sub>Γ<sub>1</sub>/Γ

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.2 ± 1.0 OUR AVERAGE</b>				
$9.2 \pm 0.8 \pm 0.7$	<sup>1</sup> LEES	12AC BABR	10.58	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
$8.9_{-3.1}^{+3.9} \pm 1.8$	8.1	HE	06B CLEO	$9.4\text{--}10.6 e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

NODE=M074G1  
NODE=M074G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$6.4 \pm 0.8 \pm 0.6$	2	LIU	13B	BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
$20.5 \pm 1.4 \pm 2.0$	3	LIU	13B	BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	OCCUR=2
$6.0 \pm 1.2^{+4.7}_{-0.5}$	2,4	YUAN	07	BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	
$20.6 \pm 2.3^{+9.1}_{-1.7}$	3,4	YUAN	07	BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	OCCUR=2
$5.5 \pm 1.0^{+0.8}_{-0.7}$	125	5 AUBERT,B	05I	BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$	

<sup>1</sup> From a single-resonance fit. Supersedes AUBERT,B 05I.

<sup>2</sup> Solution I of two equivalent solutions in a fit using two interfering resonances.

<sup>3</sup> Solution II of two equivalent solutions in a fit using two interfering resonances.

<sup>4</sup> Superseded by LIU 13B.

<sup>5</sup> From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074G1;LINKAGE=LE

NODE=M074G1;LINKAGE=YOU

NODE=M074G1;LINKAGE=YA

NODE=M074G1;LINKAGE=YN

NODE=M074G1;LINKAGE=AU

$\Gamma(J/\psi K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_9 \Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M074G3

NODE=M074G3

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.42 \pm 0.04 \pm 0.15$	1	ABLIKIM	22AU	BES3	$e^+ e^- \rightarrow K^+ K^- J/\psi$	
$0.29 \pm 0.02 \pm 0.10$	2	ABLIKIM	22AU	BES3	$e^+ e^- \rightarrow K^+ K^- J/\psi$	OCCUR=2
<1.7	90	3 SHEN	14	BELL	$9.4-10.9 e^+ e^- \rightarrow \gamma K^+ K^- J/\psi$	
<1.2	90	4 YUAN	08	BELL	$e^+ e^- \rightarrow \gamma K^+ K^- J/\psi$	

<sup>1</sup> Solution I from a two-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.127-4.600$  GeV. The second resonance has a mass of  $4484.7 \pm 13.3 \pm 24.1$  MeV, a total width of  $111.1 \pm 30.1 \pm 15.2$  MeV, and  $\Gamma_{ee} \cdot B = 1.35 \pm 0.14 \pm 0.07$  eV. The phase difference is  $1.72 \pm 0.09 \pm 0.52$  rad.

<sup>2</sup> Solution II from a two-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.127-4.600$  GeV. The second resonance has a mass of  $4484.7 \pm 13.3 \pm 24.1$  MeV, a total width of  $111.1 \pm 30.1 \pm 15.2$  MeV, and  $\Gamma_{ee} \cdot B = 0.41 \pm 0.08 \pm 0.13$  eV. The phase difference is  $5.49 \pm 0.35 \pm 0.58$  rad.

<sup>3</sup> From a fit of the broad  $K^+ K^- J/\psi$  enhancement including a coherent  $\psi(4260)$  amplitude with mass and width from LIU 13B. Supersedes YUAN 08. The shape of the cross section observed by ABLIKIM 18N between 4.2 and 4.3 GeV is incompatible with that of  $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$  in ABLIKIM 13T and ABLIKIM 17B. They also observe a broad enhancement around 4.5 GeV.

<sup>4</sup> From a fit of the broad  $K^+ K^- J/\psi$  enhancement including a coherent  $\psi(4260)$  amplitude with mass and width from YUAN 07.

NODE=M074G3;LINKAGE=B

NODE=M074G3;LINKAGE=C

NODE=M074G3;LINKAGE=A

NODE=M074G3;LINKAGE=YOU

$\Gamma(J/\psi K_S^0 K_S^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{10} \Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M074G02

NODE=M074G02

<0.85	90	1 SHEN	14	BELL	$9.4-10.9 e^+ e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$	
-------	----	--------	----	------	--	--

<sup>1</sup> From a fit of the  $K_S^0 K_S^0 J/\psi$  mass range from 4.4 to 5.5 GeV including a coherent  $\psi(4260)$  amplitude with mass and width from LIU 13B.

NODE=M074G02;LINKAGE=A

$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{11} \Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	

NODE=M074G01

NODE=M074G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

$8.0 \pm 1.7$	1	ABLIKIM	200	BES3	$e^+ e^- \rightarrow \eta J/\psi$	
$4.8 \pm 1.0$	2	ABLIKIM	200	BES3	$e^+ e^- \rightarrow \eta J/\psi$	OCCUR=2
$7.0 \pm 1.5$	3	ABLIKIM	200	BES3	$e^+ e^- \rightarrow \eta J/\psi$	OCCUR=3
<14.2	90	WANG	13B	BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$	

<sup>1</sup> Solution 1 of three equivalent fit solutions using three resonant structures.

<sup>2</sup> Solution 2 of three equivalent fit solutions using three resonant structures.

<sup>3</sup> Solution 3 of three equivalent fit solutions using three resonant structures.

NODE=M074G01;LINKAGE=A

NODE=M074G01;LINKAGE=B

NODE=M074G01;LINKAGE=C

$\Gamma(J/\psi \eta') \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{13} \Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	

NODE=M074R34

NODE=M074R34

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06 \pm 0.03$	46	1,2 ABLIKIM	20A	BES3	$e^+ e^- \rightarrow \eta' J/\psi$	
$1.38 \pm 0.11$	46	1,3 ABLIKIM	20A	BES3	$e^+ e^- \rightarrow \eta' J/\psi$	OCCUR=2

<sup>1</sup> Based on a fit to  $\sigma(e^+ e^- \rightarrow \eta' J/\psi)$  from  $\sqrt{s} = 4.18$  to 4.60 GeV assuming interfering  $\psi(4160)$  and  $\psi(4260)$  contributions. At  $\sqrt{s} = 4.23$  GeV,  $\sigma(e^+ e^- \rightarrow \eta' J/\psi) = 3.6 \pm 0.6 \pm 0.3$  pb.

<sup>2</sup> Solution I of the fit, corresponding to a phase of  $-0.03 \pm 0.44$  rad.

<sup>3</sup> Solution II of the fit, corresponding to a phase of  $2.54 \pm 0.04$  rad.

NODE=M074R34;LINKAGE=A

NODE=M074R34;LINKAGE=B

NODE=M074R34;LINKAGE=C

$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{17}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.75$		1 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$1.63 \pm 0.78$		2 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$0.02 \pm 0.01$		3 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$1.6 \pm 1.3$		4 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$1.8 \pm 1.4$		5 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
$< 4.3$	90	6 LIU	08H RVUE	$10.58 e^+e^- \rightarrow \psi(2S)\pi^+\pi^-\gamma$
$7.4^{+2.1}_{-1.7}$		7 LIU	08H RVUE	$10.58 e^+e^- \rightarrow \psi(2S)\pi^+\pi^-\gamma$

NODE=M074G7  
NODE=M074G7

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.59 \pm 0.75$		1 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
$1.63 \pm 0.78$		2 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=2
$0.02 \pm 0.01$		3 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=3
$1.6 \pm 1.3$		4 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
$1.8 \pm 1.4$		5 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=2
$< 4.3$	90	6 LIU	08H RVUE	$10.58 e^+e^- \rightarrow \psi(2S)\pi^+\pi^-\gamma$	
$7.4^{+2.1}_{-1.7}$		7 LIU	08H RVUE	$10.58 e^+e^- \rightarrow \psi(2S)\pi^+\pi^-\gamma$	OCCUR=2

1 Solution I of four equivalent solutions in a fit using three interfering resonances.

2 Solution II of four equivalent solutions in a fit using three interfering resonances

3 Solutions III and IV of four equivalent solutions in a fit using three interfering resonances.

4 Solution I of two equivalent solutions in a fit using two interfering resonances.

5 Solution II of two equivalent solutions in a fit using two interfering resonances.

6 For constructive interference with the  $\psi(4360)$  in a combined fit of AUBERT 07S and WANG 07D data with three resonances.7 For destructive interference with the  $\psi(4360)$  in a combined fit of AUBERT 07S and WANG 07D data with three resonances.NODE=M074G7;LINKAGE=A  
NODE=M074G7;LINKAGE=B  
NODE=M074G7;LINKAGE=C  
NODE=M074G7;LINKAGE=AA  
NODE=M074G7;LINKAGE=BB  
NODE=M074G7;LINKAGE=LI  
NODE=M074G7;LINKAGE=LU $\Gamma(\chi_{c0}\omega) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{19}\Gamma_1/\Gamma$ 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.2 \pm 0.3$		1 ABLIKIM	19AI BES3	$e^+e^- \rightarrow \omega\chi_{c0}$
$2.7 \pm 0.5 \pm 0.4$	180	2 ABLIKIM	15C BES3	$e^+e^- \rightarrow \omega\chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1 From a fit of the measured cross section from  $\sqrt{s} = 4.178\text{--}4.278$  GeV. Supersedes ABLIKIM 15C.2 From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+\pi^-$ ,  $\chi_{c0} \rightarrow K^+K^-$ , and  $\omega \rightarrow \pi^+\pi^-\pi^0$ .NODE=M074G05  
NODE=M074G05

NODE=M074G05;LINKAGE=B

NODE=M074G05;LINKAGE=A

 $\Gamma(h_c(1P)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{22}\Gamma_1/\Gamma$ 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$4.6^{+2.9}_{-1.4} \pm 0.8$	ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$

NODE=M074R47  
NODE=M074R47 $\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{23}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.4$	90	AUBERT, BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

NODE=M074G2  
NODE=M074G2 $\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{24}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.28$	90	1 AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

NODE=M074G6  
NODE=M074G61 AUBERT 07AK reports  $[\Gamma(\psi(4230) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(4230) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] < 0.14$  eV which we divide by our best value  $B(\phi(1020) \rightarrow K^+K^-) = 49.1 \times 10^{-2}$ .

NODE=M074G6;LINKAGE=AU

 $\Gamma(\Xi^-\Xi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{48}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.7 \times 10^{-4}$	90	ABLIKIM	20C BES3	$e^+e^- \rightarrow \Xi^-\Xi^+$

NODE=M074R35  
NODE=M074R35 $\Gamma(\pi^+\pi^+\pi^-\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{49}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 32$	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow 2\pi^+2\pi^-$

NODE=M074R37  
NODE=M074R37 $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{50}\Gamma_1/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 16$	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow 2\pi^+2\pi^-\pi^0$

NODE=M074R38  
NODE=M074R38

$$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{53} \Gamma_1 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.04 ± 0.19 ± 0.09	<sup>1</sup> ABLIKIM	19AE BES3	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0027 ± 0.0023 ± 0.0001	<sup>2</sup> ABLIKIM	19AE BES3	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp$
< 0.5 at 90% CL	AUBERT	08S BABR	$10.6 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

NODE=M074G4  
NODE=M074G4

OCCUR=2

<sup>1</sup> Solution I of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

NODE=M074G4;LINKAGE=A

<sup>2</sup> Solution II of the fit including the  $\psi(4160)$  with mass  $4191 \pm 5$  MeV and width  $70 \pm 10$  MeV from PDG 16 and the  $\psi(4230)$  with mass  $4219.6 \pm 3.3 \pm 5.1$  MeV and width  $56.0 \pm 3.6 \pm 6.9$  MeV from GAO 17.

NODE=M074G4;LINKAGE=B

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{54} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.05      90      ABLIKIM      19      BES3       $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$

NODE=M074R31  
NODE=M074R31

$$\Gamma(K_S^0 K^\pm \pi^\mp \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{55} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.19      90      ABLIKIM      19      BES3       $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \eta$

NODE=M074R32  
NODE=M074R32

$$\Gamma(K^+ K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{56} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.6      90      AUBERT      08S      BABR       $10.6 e^+ e^- \rightarrow K^+ K^- \pi^0 \gamma$

NODE=M074G5  
NODE=M074G5

$$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{57} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<20      90      ABLIKIM      21AW BES3       $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M074R39  
NODE=M074R39

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{58} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<43      90      ABLIKIM      21AW BES3       $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M074R40  
NODE=M074R40

$$\Gamma(K^+ K^+ K^- K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{59} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<3.8      90      ABLIKIM      21AW BES3       $e^+ e^- \rightarrow 2K^+ 2K^-$

NODE=M074R41  
NODE=M074R41

$$\Gamma(K^+ K^+ K^- K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{60} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<2.1      90      ABLIKIM      21AW BES3       $e^+ e^- \rightarrow 2K^+ 2K^- \pi^0$

NODE=M074R42  
NODE=M074R42

$$\Gamma(p\bar{p}\pi^+\pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{61} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<7.2      90      ABLIKIM      21AW BES3       $e^+ e^- \rightarrow p\bar{p}\pi^+\pi^-$

NODE=M074R43  
NODE=M074R43

$$\Gamma(p\bar{p}\pi^+\pi^-\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{62} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<15      90      ABLIKIM      21AW BES3       $e^+ e^- \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

NODE=M074R44  
NODE=M074R44

$$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{64} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.8 × 10<sup>-3</sup>      90      <sup>1</sup> ABLIKIM      21AS BES3       $e^+ e^- \rightarrow \psi(4260)$

NODE=M074R36  
NODE=M074R36

<sup>1</sup> From a measurement of the  $e^+ e^- \rightarrow \Lambda\bar{\Lambda}$  cross section between 3.5 and 4.6 GeV.

NODE=M074R36;LINKAGE=A

$$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{67} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.4      90      <sup>1</sup> HAN      15      BELL       $10.58 e^+ e^- \rightarrow \chi_{c1}\gamma$

NODE=M074G03  
NODE=M074G03

<sup>1</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

NODE=M074G03;LINKAGE=A

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{68}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;4.0</b>	90	<sup>1</sup> HAN	15	BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$
<sup>1</sup> Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .					

NODE=M074G04  
NODE=M074G04

NODE=M074G04;LINKAGE=A

### $\psi(4230)$ BRANCHING RATIOS

NODE=M074225

$\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_3/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>not seen</b>		<sup>1</sup> ABLIKIM	21B	BES3	$e^+e^- \rightarrow \pi^+\pi^-\eta_c$
<sup>1</sup> Not seen in $e^+e^- \rightarrow \pi^+\pi^-\eta_c$ at $\sqrt{s} = 4.226$ GeV with a 90% C.L. upper limit on the cross section of 16.8 pb.					

NODE=M074R56  
NODE=M074R56

NODE=M074R56;LINKAGE=A

$\Gamma(\eta_c(1S)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>seen</b>		<sup>1</sup> ABLIKIM	21B	BES3	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c$
<sup>1</sup> Seen as a peak in the $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c$ cross section with a peak value of $46.1^{+9.5}_{-9.4} \pm 6.6$ pb at $\sqrt{s} = 4.226$ GeV.					

NODE=M074R55  
NODE=M074R55

NODE=M074R55;LINKAGE=A

$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_5/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>seen</b>		<sup>1</sup> ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$	
••• We do not use the following data for averages, fits, limits, etc. •••					
<b>seen</b>		<sup>2</sup> ABLIKIM	17B	BES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
<sup>1</sup> From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730\text{--}4.7008$ GeV.					
<sup>2</sup> From a three-resonance fit. Superseded by ABLIKIM 22AM.					

NODE=M074R51  
NODE=M074R51

NODE=M074R51;LINKAGE=A

NODE=M074R51;LINKAGE=B

$\Gamma(J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$					$\Gamma_6/\Gamma_5$
VALUE		DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••					
0.17±0.13		<sup>1</sup> LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
<sup>1</sup> Systematic uncertainties not estimated.					

NODE=M074R02  
NODE=M074R02

NODE=M074R02;LINKAGE=LE

$\Gamma(Z_c(3900)^\pm\pi^\mp, Z_c^\pm \rightarrow J/\psi\pi^\pm)/\Gamma(J/\psi\pi^+\pi^-)$					$\Gamma_7/\Gamma_5$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>0.215±0.033±0.075</b>		<sup>1</sup> ABLIKIM	13T	BES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$
••• We do not use the following data for averages, fits, limits, etc. •••					
0.29 ±0.08		<sup>2</sup> LIU	13B	BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
<sup>1</sup> Assuming that the cross section of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ is fully due to the $\psi(4260)$ .					
<sup>2</sup> Systematic error not evaluated.					

NODE=M074R01  
NODE=M074R01

NODE=M074R01;LINKAGE=AB  
NODE=M074R01;LINKAGE=A

$\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$					$\Gamma_8/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>seen</b>		<sup>1</sup> ABLIKIM	20N	BES3	$e^+e^- \rightarrow \pi^0\pi^0J/\psi$
<sup>1</sup> From a fit to the cross section $e^+e^- \rightarrow \pi^0\pi^0J/\psi$ at center-of-mass energies between 3.808 and 4.600 GeV.					

NODE=M074R50  
NODE=M074R50

NODE=M074R50;LINKAGE=A

$\Gamma(J/\psi K_S^0 K_S^0)/\Gamma_{\text{total}}$					$\Gamma_{10}/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>not seen</b>		SHEN	14	BELL	9.4–10.9 $e^+e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$

NODE=M074R27  
NODE=M074R27

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$					$\Gamma_{11}/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>seen</b>		ABLIKIM	200	BES3	$e^+e^- \rightarrow \eta J/\psi$

NODE=M074R52  
NODE=M074R52

$\Gamma(J/\psi\eta\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{15}/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
<b>not seen</b>		ABLIKIM	15Q	BES3	4.0–4.6 $e^+e^- \rightarrow J/\psi\eta\pi^0$

NODE=M074R28  
NODE=M074R28

$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$

NODE=M074R53  
NODE=M074R53

<sup>1</sup> From a fit to the cross section for  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-\ell^+\ell^-)$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1 \text{ fb}^{-1}$ .

NODE=M074R53;LINKAGE=A

 $\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$  $\Gamma_{17}/\Gamma_5$ 

VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
(0.11 ± 0.03 ± 0.03) to (0.55 ± 0.18 ± 0.19)	<sup>1</sup> ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$

NODE=M074R30  
NODE=M074R30

<sup>1</sup> From a combined fit of BELLE, BABAR and BES3  $e^+e^- \rightarrow \pi^+\pi^-J/\psi$  and  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$  data.

NODE=M074R30;LINKAGE=A

 $\Gamma(\chi_{c0}\omega)/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	180	<sup>1</sup> ABLIKIM	15C BES3	$e^+e^- \rightarrow \omega\chi_{c0}$

NODE=M074R48  
NODE=M074R48

<sup>1</sup> From a 3-parameter fit of measured cross sections from  $\sqrt{s} = 4.21\text{--}4.42$  GeV to a phase-space modified Breit-Wigner function, using the decays  $\chi_{c0} \rightarrow \pi^+\pi^-$ ,  $\chi_{c0} \rightarrow K^+K^-$ , and  $\omega \rightarrow \pi^+\pi^-\pi^0$ .

NODE=M074R48;LINKAGE=A

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$

NODE=M074R49  
NODE=M074R49 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$  $\Gamma_{22}/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	<sup>1</sup> PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

NODE=M074R25  
NODE=M074R25

<sup>1</sup> At  $\sqrt{s} = 4260$  MeV, PEDLAR 11 measures  $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 32 \pm 17 \pm 6 \pm 6$  pb, where the errors are statistical, systematic, and due to uncertainty in  $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$ , respectively.

NODE=M074R25;LINKAGE=PE

 $\Gamma(D\bar{D})/\Gamma(J/\psi\pi^+\pi^-)$  $\Gamma_{25}/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	<sup>1</sup> AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$

NODE=M074R2  
NODE=M074R2

••• We do not use the following data for averages, fits, limits, etc. •••

<4.0	90	CRONIN-HEN..09	CLEO	$e^+e^-$
------	----	----------------	------	----------

<sup>1</sup> Using  $4259 \pm 10$  MeV for the mass and  $88 \pm 24$  MeV for the width of  $\psi(4260)$ .

NODE=M074R2;LINKAGE=AU

 $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$

NODE=M074R12  
NODE=M074R12

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$
not seen	PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$  $\Gamma_{27}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$

NODE=M074R13  
NODE=M074R13

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^+D^-\gamma$
not seen	PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^+D^-\gamma$

 $\Gamma(D^*\bar{D}^0+c.c.)/\Gamma(J/\psi\pi^+\pi^-)$  $\Gamma_{28}/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<34	90	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}^0$

NODE=M074R03  
NODE=M074R03

••• We do not use the following data for averages, fits, limits, etc. •••

<45	90	CRONIN-HEN..09	CLEO	$e^+e^-$
-----	----	----------------	------	----------

 $\Gamma(D^*(2007)^0\bar{D}^0+c.c.)/\Gamma_{\text{total}}$  $\Gamma_{29}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$

NODE=M074R14  
NODE=M074R14

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$
----------	--------	----------	--

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^-$
not seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$

NODE=M074R15  
NODE=M074R15 $\Gamma(D^* \bar{D}^*)/\Gamma(J/\psi \pi^+ \pi^-)$   $\Gamma_{31}/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	CRONIN-HEN..09	CLEO	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<40	90	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^* \bar{D}^*$

NODE=M074R04  
NODE=M074R04 $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$

NODE=M074R17  
NODE=M074R17 $\Gamma(D^*(2010)^+ D^*(2010)^-)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^{*+} D^{*-}$
not seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

NODE=M074R18  
NODE=M074R18 $\Gamma(D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^{*0} + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.}))/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 08A	BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

NODE=M074R16  
NODE=M074R16 $\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma_{\text{total}}$   $\Gamma_{36}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D} \pi$

NODE=M074R22  
NODE=M074R22 $\Gamma(D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*))/\Gamma(J/\psi \pi^+ \pi^-)$   $\Gamma_{36}/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R05  
NODE=M074R05 $\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

NODE=M074R23  
NODE=M074R23 $\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM 19R	BES3	$e^+ e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$

NODE=M074R54  
NODE=M074R54 $\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-)$   $\Gamma_{38}/\Gamma_5$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R10  
NODE=M074R10 $\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$   $\Gamma_{38}/\Gamma \times \Gamma_1/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42 × 10 <sup>-6</sup>	90	<sup>1</sup> PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R11  
NODE=M074R11<sup>1</sup> Using  $4263^{+8}_{-9}$  MeV for the mass of  $\psi(4260)$ .

NODE=M074R11;LINKAGE=PA

 $\Gamma(D_1(2420) \bar{D} + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> ABLIKIM 19AR	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$

NODE=M074R33  
NODE=M074R33<sup>1</sup> Results from a measurement of  $\sigma(e^+ e^- \rightarrow D_1(2420) \bar{D} + \text{c.c.})$  between  $\sqrt{s} = 4.3$  and 4.6 GeV.

NODE=M074R33;LINKAGE=A



$\Gamma(D^* \bar{D}^* \pi) / \Gamma_{\text{total}}$					$\Gamma_{40} / \Gamma$	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M074R24 NODE=M074R24
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D}^* \pi$			
$\Gamma(D^* \bar{D}^* \pi) / \Gamma(J/\psi \pi^+ \pi^-)$					$\Gamma_{40} / \Gamma_5$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M074R06 NODE=M074R06
<8.2	90	CRONIN-HEN..09	CLEO	$e^+ e^-$		
$\Gamma(D_s^+ D_s^-) / \Gamma_{\text{total}}$					$\Gamma_{41} / \Gamma$	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M074R19 NODE=M074R19
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$			
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$			
$\Gamma(D_s^+ D_s^-) / \Gamma(J/\psi \pi^+ \pi^-)$					$\Gamma_{41} / \Gamma_5$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M074R07 NODE=M074R07
<0.7	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.3	90	CRONIN-HEN..09	CLEO	$e^+ e^-$		
$\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma_{\text{total}}$					$\Gamma_{42} / \Gamma$	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M074R20 NODE=M074R20
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$			
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$			
$\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma(J/\psi \pi^+ \pi^-)$					$\Gamma_{42} / \Gamma_5$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M074R08 NODE=M074R08
< 0.8	90	CRONIN-HEN..09	CLEO	$e^+ e^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<44	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$		
$\Gamma(D_s^{*+} D_s^{*-}) / \Gamma_{\text{total}}$					$\Gamma_{43} / \Gamma$	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M074R21 NODE=M074R21
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$			
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$			
$\Gamma(D_s^{*+} D_s^{*-}) / \Gamma(J/\psi \pi^+ \pi^-)$					$\Gamma_{43} / \Gamma_5$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M074R09 NODE=M074R09
< 9.5	90	CRONIN-HEN..09	CLEO	$e^+ e^-$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<30	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$		
$\Gamma(p \bar{p}) / \Gamma(J/\psi \pi^+ \pi^-)$					$\Gamma_{44} / \Gamma_5$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M074R1 NODE=M074R1
<0.13	90	<sup>1</sup> AUBERT 06B	BABR	$e^+ e^- \rightarrow p \bar{p} \gamma$		
<sup>1</sup> Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$ .						NODE=M074R1;LINKAGE=AU
$\Gamma(p \bar{p} \pi^0) / \Gamma(J/\psi \pi^+ \pi^-)$					$\Gamma_{45} / \Gamma_5$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M074R00 NODE=M074R00
<2 × 10 <sup>-4</sup>	90	ABLIKIM 17F	BES3	$e^+ e^- \rightarrow \psi(4260) \rightarrow$ hadrons		OCCUR=2
$\Gamma(\omega \pi^0) / \Gamma_{\text{total}}$					$\Gamma_{51} / \Gamma$	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M074R59 NODE=M074R59
not seen	ABLIKIM 22K	BES3	$e^+ e^- \rightarrow \omega \pi^0$			

$\Gamma(\omega\eta)/\Gamma_{\text{total}}$	$\Gamma_{52}/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
not seen	ABLIKIM 22K BES3	$e^+e^- \rightarrow \omega\eta$
		NODE=M074R60 NODE=M074R60
$\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$	$\Gamma_{46}/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
not seen	ABLIKIM 21AN BES3	$e^+e^- \rightarrow p\bar{p}\eta$
		NODE=M074R45 NODE=M074R45
$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$	$\Gamma_{47}/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
not seen	ABLIKIM 21AN BES3	$e^+e^- \rightarrow p\bar{p}\omega$
		NODE=M074R46 NODE=M074R46
$\Gamma(p\bar{p}p\bar{p})/\Gamma_{\text{total}}$	$\Gamma_{63}/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
not seen	ABLIKIM 21D BES3	$4.0\text{--}4.6 e^+e^- \rightarrow p\bar{p}p\bar{p}$
		NODE=M074R58 NODE=M074R58

## Radiative decays

$\Gamma(\eta_c(1S)\gamma)/\Gamma_{\text{total}}$	$\Gamma_{65}/\Gamma$	
VALUE	DOCUMENT ID	COMMENT
possibly seen	<sup>1</sup> ABLIKIM 17W	$e^+e^- \rightarrow \gamma\eta_c(1S)$
		NODE=M074310
		NODE=M074R29 NODE=M074R29
		NODE=M074R29;LINKAGE=A

<sup>1</sup>Significance ranges from 4.2  $\sigma$  to as low as 1.5  $\sigma$  for a flat component plus  $\psi(4260)$  spectrum. Needs confirmation.

$\Gamma(\eta_c(1S)\pi^0\gamma)/\Gamma_{\text{total}}$	$\Gamma_{66}/\Gamma$	
VALUE	DOCUMENT ID	TECN COMMENT
not seen	<sup>1</sup> ABLIKIM 21B BES3	$e^+e^- \rightarrow \gamma\pi^0\eta_c$
		NODE=M074R57 NODE=M074R57
		NODE=M074R57;LINKAGE=A

<sup>1</sup>Not seen in  $e^+e^- \rightarrow \gamma\pi^0\eta_c$  at  $\sqrt{s} = 4.226$  GeV with a 90% C.L. upper limit on the cross section of 11.2 pb.

$\Gamma(\chi_{c1}(3872)\gamma)/\Gamma_{\text{total}}$	$\Gamma_{69}/\Gamma$	
VALUE	EVTS	DOCUMENT ID TECN COMMENT
seen		ABLIKIM 19v BES3 $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$
seen	20 $\pm$ 5	ABLIKIM 14 BES3 $e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
		NODE=M074R26 NODE=M074R26

 $\psi(4230)$  REFERENCES

ABLIKIM	22AM PR D106 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61885
ABLIKIM	22AU CP C46 111002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61896
ABLIKIM	22K JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	21AJ PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61441
ABLIKIM	21AN PR D104 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61446
ABLIKIM	21AS PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21AW PR D104 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61479
ABLIKIM	21B PR D103 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61029
ABLIKIM	21D PR D103 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61031
ABLIKIM	20A PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60210
ABLIKIM	20AG PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	20C PRL 124 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60215
ABLIKIM	20N PR D102 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60338
ABLIKIM	20O PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60344
ABLIKIM	19 PR D99 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59515
ABLIKIM	19AE PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59856
ABLIKIM	19AI PR D99 091103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59871
ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ABLIKIM	19K PR D99 019903 (err.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	19R PRL 122 102002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59765
ABLIKIM	19V PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796
ABLIKIM	18N PR D97 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58914
ABLIKIM	17B PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57755
ABLIKIM	17F PL B771 45	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57909
ABLIKIM	17G PRL 118 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57915
ABLIKIM	17V PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58029
Also	PR D99 019903 (err.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	17W PR D96 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58212
GAO	17 PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan		REFID=57991
ZHANG	17B PR D96 054008	J. Zhang, J. Zhang		REFID=58219
ZHANG	17C EPJ C77 727	J. Zhang, L. Yuan		REFID=58463
PDG	16 CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15C PRL 114 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56401
ABLIKIM	15Q PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56782
HAN	15 PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
ABLIKIM	14 PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55647
SHEN	14 PR D89 072015	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55944

NODE=M074

ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55409
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
LEES	12AC	PR D86 051102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54750
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=52296
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
YUAN	08	PR D77 011105	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=52135
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51724
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51959
YUAN	07	PRL 99 182004	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=51960
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
HE	06B	PR D74 091104	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51523
AUBERT,B	05I	PRL 95 142001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50776

NODE=M233

 $\chi_{c1}(4274)$ 

$$J^{PC} = 0^+(1^+)$$

was  $X(4274)$ 

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

NODE=M233

Seen by AAIJ 17C in  $B^+ \rightarrow \chi_{c1} K^+$ ,  $\chi_{c1} \rightarrow J/\psi \phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi \phi K^+$  with a significance (accounting for systematic uncertainties) of  $6.0 \sigma$ .

 **$\chi_{c1}(4274)$  MASS**

NODE=M233M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4286 <math>\pm 8</math><sub>-9</sub></b>		<b>OUR AVERAGE</b>		Error includes scale factor of 1.7.
4294 $\pm 4$ $\pm 3$ <sub>-6</sub>	24k	<sup>1</sup> AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$
4274.4 $\pm 8.4$ <sub>-6.7</sub> $\pm 1.9$	22	<sup>2</sup> AALTONEN	17	CDF $B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4273.3 $\pm 8.3$ $\pm 17.2$ <sub>-3.6</sub>	4289	<sup>3,4</sup> AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$

NODE=M233M

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $18 \sigma$ .

NODE=M233M;LINKAGE=C

<sup>2</sup> From a fit to the invariant mass spectrum with a significance of  $3.1 \sigma$ .

NODE=M233M;LINKAGE=B

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $6.0 \sigma$ .

NODE=M233M;LINKAGE=A

<sup>4</sup> Superseded by AAIJ 21E.

NODE=M233M;LINKAGE=D

 **$\chi_{c1}(4274)$  WIDTH**

NODE=M233W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>51 <math>\pm 7</math></b>		<b>OUR AVERAGE</b>		
53 $\pm 5$ $\pm 5$	24k	<sup>1</sup> AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$
32.3 $\pm 21.9$ <sub>-15.3</sub> $\pm 7.6$	22	<sup>2</sup> AALTONEN	17	CDF $B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
56 $\pm 11$ $\pm 8$ <sub>-11</sub>	4289	<sup>3,4</sup> AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$

NODE=M233W

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $18 \sigma$ .

NODE=M233W;LINKAGE=C

<sup>2</sup> From a fit to the invariant mass spectrum with a significance of  $3.1 \sigma$ .

NODE=M233W;LINKAGE=B

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $6.0 \sigma$ .

NODE=M233W;LINKAGE=A

<sup>4</sup> Superseded by AAIJ 21E.

NODE=M233W;LINKAGE=D

 **$\chi_{c1}(4274)$  DECAY MODES**

NODE=M233215;NODE=M233

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi \phi$	seen

DESIG=1

$\chi_{c1}(4274)$  BRANCHING RATIOS

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
••• We do not use the following data for averages, fits, limits, etc. •••					
seen	4289	<sup>2,3</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 18 $\sigma$ .					
<sup>2</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 6.0 $\sigma$ .					
<sup>3</sup> Superseded by AAIJ 21E.					

NODE=M233220

NODE=M233R01  
NODE=M233R01NODE=M233R01;LINKAGE=B  
NODE=M233R01;LINKAGE=A  
NODE=M233R01;LINKAGE=C $\chi_{c1}(4274)$  REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	17	MPL A32 1750139	T. Altonen <i>et al.</i>	(CDF Collab.)

NODE=M233

REFID=61150  
REFID=57657  
REFID=57636  
REFID=58161

NODE=M194

**X(4350)**

$$I^G(J^{PC}) = 0^+(?^?+)$$

OMITTED FROM SUMMARY TABLE

Seen by SHEN 10 in the  $\gamma\gamma \rightarrow J/\psi\phi$ . Needs confirmation.

NODE=M194

**X(4350) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4350.6^{+4.6}_{-5.1} \pm 0.7$	$8.8^{+4.2}_{-3.2}$	<sup>1</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi\phi$
<sup>1</sup> Statistical significance of 3.2 $\sigma$ .				

NODE=M194M

NODE=M194M

NODE=M194M;LINKAGE=SH

**X(4350) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$13^{+18}_{-9} \pm 4$	$8.8^{+4.2}_{-3.2}$	<sup>1</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi\phi$
<sup>1</sup> Statistical significance of 3.2 $\sigma$ .				

NODE=M194W

NODE=M194W

NODE=M194W;LINKAGE=SH

**X(4350) DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi\phi$	seen
$\Gamma_2$ $\gamma\gamma$	seen

NODE=M194215;NODE=M194

DESIG=1

DESIG=2

**X(4350)  $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$** 

$\Gamma(\gamma\gamma) \times \Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$6.7^{+3.2}_{-2.4} \pm 1.1$	$8.8^{+4.2}_{-3.2}$	<sup>1</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi\phi$	
••• We do not use the following data for averages, fits, limits, etc. •••					
$1.5^{+0.7}_{-0.6} \pm 0.3$	$8.8^{+4.2}_{-3.2}$	<sup>2</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi\phi$	
<sup>1</sup> For $J^P = 0^+$ . Statistical significance of 3.2 $\sigma$ .					
<sup>2</sup> For $J^P = 2^+$ . Statistical significance of 3.2 $\sigma$ .					

NODE=M194220

NODE=M194G01  
NODE=M194G01

OCCUR=2

NODE=M194G01;LINKAGE=S0  
NODE=M194G01;LINKAGE=S2**X(4350) BRANCHING RATIOS**

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		<sup>1</sup> SHEN	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- J/\psi\phi$	
<sup>1</sup> Statistical significance of 3.2 $\sigma$ .					

NODE=M194225

NODE=M194R01  
NODE=M194R01

NODE=M194R01;LINKAGE=SH

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	1 SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

<sup>1</sup> Statistical significance of 3.2  $\sigma$ .NODE=M194R02  
NODE=M194R02

NODE=M194R02;LINKAGE=SH

**X(4350) REFERENCES**SHEN 10 PRL 104 112004 C.P. Shen *et al.* (BELLE Collab.)

NODE=M194

REFID=53235

NODE=M181

 **$\psi(4360)$** 

$$J^{PC} = 0^-(1^-)$$

also known as  $Y(4360)$ ; was  $X(4360)$ 

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

NODE=M181

Seen in radiative return from  $e^+e^-$  collisions at  $\sqrt{s} = 9.54\text{--}10.58$  GeV by AUBERT 07S, WANG 07D, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

 **$\psi(4360)$  MASS**

NODE=M181M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4374 <math>\pm</math> 7 OUR AVERAGE</b>		Error includes scale factor of 2.4. See the ideogram below. [4372 $\pm$ 9 MeV OUR 2022 AVERAGE Scale factor = 2.9]		
4371.6 $\pm$ 2.5 $\pm$ 9.2		1 ABLIKIM	22AL BES3	$e^+e^- \rightarrow \pi^+\pi^- D^+ D^-$
4298 $\pm$ 12 $\pm$ 26		2 ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
4390.3 $\pm$ 6.0 $\pm$ 0.7		3 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
4371.7 $\pm$ 7.5 $\pm$ 1.8		4 ABLIKIM	21AK BES3	$e^+e^- \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi$
4382.0 $\pm$ 13.3 $\pm$ 1.7		5 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$
4391.5 $^{+6.3}_{-6.8} \pm$ 1.0		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^- h_c$
4347 $\pm$ 6 $\pm$ 3	279	6 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
4340 $\pm$ 16 $\pm$ 9	37	7 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4406.9 $\pm$ 17.2 $\pm$ 4.5		8 ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^- \chi_{c1}\gamma$
4320.0 $\pm$ 10.4 $\pm$ 7.0		9 ABLIKIM	17B BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
4383.8 $\pm$ 4.2 $\pm$ 0.8		10 ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
4383.7 $\pm$ 2.9 $\pm$ 6.2		11 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
4386.4 $\pm$ 2.1 $\pm$ 6.4		12 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
4355 $^{+9}_{-10} \pm$ 9	74	13 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
4324 $\pm$ 24		14 AUBERT	07S BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
4361 $\pm$ 9 $\pm$ 9	47	7 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$

NODE=M181M

NEW

<sup>1</sup> From a fit to the cross section for  $e^+e^- \rightarrow D^+D^-\pi^+\pi^-$  in the range  $\sqrt{s} = 4.190\text{--}4.946$  GeV.

NODE=M181M;LINKAGE=M

<sup>2</sup> From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 3.7730\text{--}4.7008$  GeV. Parameters depend on the existence or non-existence of a state near 4.5 GeV.

NODE=M181M;LINKAGE=L

<sup>3</sup> From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.008\text{--}4.698$  GeV.

NODE=M181M;LINKAGE=H

<sup>4</sup> From a five-resonance fit to the cross section for  $e^+e^- \rightarrow \gamma\gamma J/\psi \rightarrow \gamma\gamma\ell^+\ell^-$ .

NODE=M181M;LINKAGE=G

<sup>5</sup> From a fit of the measured cross section in the range  $\sqrt{s} = 3.808\text{--}4.600$  GeV.

NODE=M181M;LINKAGE=BA

<sup>6</sup> From a two-resonance fit. Supersedes WANG 07D.

NODE=M181M;LINKAGE=A

<sup>7</sup> From a two-resonance fit.

NODE=M181M;LINKAGE=WA

<sup>8</sup> From a fit to the  $e^+e^- \rightarrow \pi^+\pi^- \psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass  $4417.5 \pm 26.2 \pm 3.5$  MeV and width  $245 \pm 48 \pm 13$  MeV.

NODE=M181M;LINKAGE=J

<sup>9</sup> From a three-resonance fit. Superseded by ABLIKIM 22AM.

NODE=M181M;LINKAGE=K

<sup>10</sup> From a fit to the cross section for  $e^+e^- \rightarrow \pi^+\pi^- \psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1 \text{ fb}^{-1}$ . Superseded by ABLIKIM 21AJ.

NODE=M181M;LINKAGE=C

<sup>11</sup> From a three-resonance fit.

NODE=M181M;LINKAGE=E

<sup>12</sup> From a combined fit of BELLE, BABAR and BES3  $e^+e^- \rightarrow \pi^+\pi^- J/\psi$  and  $e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$  data.


NODE=M181M;LINKAGE=D

<sup>13</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M181M;LINKAGE=LI

<sup>14</sup> From a single-resonance fit. Systematic errors not estimated.

NODE=M181M;LINKAGE=AU



**psi(4360) WIDTH**

NODE=M181W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  
**118 ±12 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.  
 [115 ± 13 MeV OUR 2022 AVERAGE Scale factor = 2.2]

NODE=M181W  
NEW

167 ± 4 ±29	1	ABLIKIM	22AL BES3	$e^+e^- \rightarrow \pi^+\pi^- D^+ D^-$
127 ±17 ±10	2	ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
143.3±10.0± 0.5	3	ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
51.1±17.6± 1.9	4	ABLIKIM	21AK BES3	$e^+e^- \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi$
135.8±60.8±22.5	5	ABLIKIM	20O BES3	$e^+e^- \rightarrow \eta J/\psi$
139.5 <sup>+16.2</sup> <sub>-20.6</sub> ± 0.6		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^- h_c$
103 ± 9 ± 5	279	6 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
94 ±32 ±13	37	7 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
128.1±37.2± 2.3		8 ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^- \chi_{c1}\gamma$
101.4 <sup>+25.3</sup> <sub>-19.7</sub> ±10.2		9 ABLIKIM	17B BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
84.2±12.5± 2.1		10 ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
94.2± 7.3± 2.0		11 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
96.0± 6.7± 2.7		12 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
103 <sup>+17</sup> <sub>-15</sub> ±11	74	13 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
172 ±33		14 AUBERT	07S BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
74 ±15 ±10	47	7 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$

- 1 From a fit to the cross section for  $e^+e^- \rightarrow D^+ D^- \pi^+\pi^-$  in the range  $\sqrt{s} = 4.190-4.946$  GeV.
- 2 From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 3.7730-4.7008$  GeV. Parameters depend on the existence or non-existence of a state near 4.5 GeV.
- 3 From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.008-4.698$  GeV.
- 4 From a five-resonance fit to the cross section for  $e^+e^- \rightarrow \gamma\gamma J/\psi \rightarrow \gamma\gamma\ell^+\ell^-$ .
- 5 From a fit of the measured cross section in the range  $\sqrt{s} = 3.808-4.600$  GeV.
- 6 From a two-resonance fit. Supersedes WANG 07D.
- 7 From a two-resonance fit.
- 8 From a fit to the  $e^+e^- \rightarrow \pi^+\pi^- \psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass  $4417.5 \pm 26.2 \pm 3.5$  MeV and width  $245 \pm 48 \pm 13$  MeV.
- 9 From a three-resonance fit. Superseded by ABLIKIM 22AM.
- 10 From a fit to the cross section for  $e^+e^- \rightarrow \pi^+\pi^- \psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$  obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising  $5.1 \text{ fb}^{-1}$ . Superseded by ABLIKIM 21AJ.
- 11 From a three-resonance fit.
- 12 From a combined fit of BELLE, BABAR and BES3  $e^+e^- \rightarrow \pi^+\pi^- J/\psi$  and  $e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$  data.

NODE=M181W;LINKAGE=L

NODE=M181W;LINKAGE=K

NODE=M181W;LINKAGE=G


NODE=M181W;LINKAGE=F  
 NODE=M181W;LINKAGE=BA  
 NODE=M181W;LINKAGE=A  
 NODE=M181W;LINKAGE=WA  
 NODE=M181W;LINKAGE=I

NODE=M181W;LINKAGE=J  
 NODE=M181W;LINKAGE=C

NODE=M181W;LINKAGE=E  
 NODE=M181W;LINKAGE=D

<sup>13</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.  
<sup>14</sup> From a single-resonance fit. Systematic errors not estimated.

NODE=M181W;LINKAGE=LI  
 NODE=M181W;LINKAGE=AU



**psi(4360) DECAY MODES**

NODE=M181215;NODE=M181

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	
$\Gamma_2$ $h_c \pi^+ \pi^-$	seen
$\Gamma_3$ $J/\psi \pi^+ \pi^-$	
$\Gamma_4$ $\psi(2S) \pi^+ \pi^-$	seen
$\Gamma_5$ $\psi(3770) \pi^+ \pi^-$	possibly seen
$\Gamma_6$ $\psi_2(3823) \pi^+ \pi^-$	seen
$\Gamma_7$ $J/\psi \eta$	seen
$\Gamma_8$ $D^0 D^{*-} \pi^+$	
$\Gamma_9$ $D^+ D^- \pi^+ \pi^-$	seen
$\Gamma_{10}$ $D_1(2420) \bar{D} + c.c.$	possibly seen
$\Gamma_{11}$ $\omega \pi^0$	not seen
$\Gamma_{12}$ $\omega \eta$	not seen
$\Gamma_{13}$ $\rho \bar{\rho} \eta$	not seen
$\Gamma_{14}$ $\rho \bar{\rho} \omega$	not seen
$\Gamma_{15}$ $\chi_{c1} \gamma$	
$\Gamma_{16}$ $\chi_{c2} \gamma$	

DESIG=1  
 DESIG=12  
 DESIG=8  
 DESIG=2  
 DESIG=11  
 DESIG=5  
 DESIG=4  
 DESIG=3  
 DESIG=17  
 DESIG=10  
 DESIG=15  
 DESIG=16  
 DESIG=13  
 DESIG=14  
 DESIG=6  
 DESIG=7

**psi(4360)  $\Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$**

NODE=M181230

**$\Gamma(h_c \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_2 \Gamma_1 / \Gamma$**

NODE=M181R11  
 NODE=M181R11

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$11.6^{+5.0}_{-4.4} \pm 1.9$	ABLIKIM	17G	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- h_c$

**$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$   $\Gamma_4 \Gamma_1 / \Gamma$**

NODE=M181G1  
 NODE=M181G1

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.7±4.1	1	ABLIKIM	21AJ	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$		
20.7±2.5	2	ABLIKIM	21AJ	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	OCCUR=2	
9.9±4.1	3	ABLIKIM	21AJ	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	OCCUR=3	
19.4±2.0	4	ABLIKIM	21AJ	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	OCCUR=4	
7.3±2.8	5	ABLIKIM	19K	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$		
11.0±3.8	6	ABLIKIM	19K	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	OCCUR=2	
9.2±0.6±0.6	279	7	WANG	15A	BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	

10.9±0.6±0.7	279	<sup>8</sup> WANG	15A BELL	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	OCCUR=2
6.0±1.0±0.5	37	<sup>5</sup> LEES	14F BABR	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	
7.2±1.0±0.6	37	<sup>6</sup> LEES	14F BABR	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	OCCUR=2
11.1 <sup>+1.3</sup> <sub>-1.2</sub>	74	<sup>9</sup> LIU	08H RVUE	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	
12.3±1.2	74	<sup>10</sup> LIU	08H RVUE	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	OCCUR=2
10.4±1.7±1.5	47	<sup>5</sup> WANG	07D BELL	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	
11.8±1.8±1.4	47	<sup>6</sup> WANG	07D BELL	10.58 e <sup>+</sup> e <sup>-</sup> → γπ <sup>+</sup> π <sup>-</sup> ψ(2S)	OCCUR=2
<sup>1</sup> Solution I of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.					NODE=M181G1;LINKAGE=E
<sup>2</sup> Solution II of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.					NODE=M181G1;LINKAGE=F
<sup>3</sup> Solution III of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.					NODE=M181G1;LINKAGE=G
<sup>4</sup> Solution IV of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.					NODE=M181G1;LINKAGE=H
<sup>5</sup> Solution I of two equivalent solutions in a fit using two interfering resonances.					NODE=M181G1;LINKAGE=WA
<sup>6</sup> Solution II of two equivalent solutions in a fit using two interfering resonances.					NODE=M181G1;LINKAGE=WN
<sup>7</sup> Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.					NODE=M181G1;LINKAGE=A
<sup>8</sup> Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.					NODE=M181G1;LINKAGE=B
<sup>9</sup> Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.					NODE=M181G1;LINKAGE=LI
<sup>10</sup> Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.					NODE=M181G1;LINKAGE=LU

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3.4±2.2		<sup>1</sup> ABLIKIM	200 BES3	e <sup>+</sup> e <sup>-</sup> → ηJ/ψ	
1.5±1.0		<sup>2</sup> ABLIKIM	200 BES3	e <sup>+</sup> e <sup>-</sup> → ηJ/ψ	OCCUR=2
1.7±1.1		<sup>3</sup> ABLIKIM	200 BES3	e <sup>+</sup> e <sup>-</sup> → ηJ/ψ	OCCUR=3
<6.8	90	WANG	13B BELL	e <sup>+</sup> e <sup>-</sup> → J/ψηη	
<sup>1</sup> Solution 1 of three equivalent fit solutions using three resonant structures.					NODE=M181G01;LINKAGE=A
<sup>2</sup> Solution 2 of three equivalent fit solutions using three resonant structures.					NODE=M181G01;LINKAGE=B
<sup>3</sup> Solution 3 of three equivalent fit solutions using three resonant structures.					NODE=M181G01;LINKAGE=C

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{15}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.57	90	<sup>1</sup> HAN	15 BELL	10.58 e <sup>+</sup> e <sup>-</sup> → χ <sub>c1</sub> γ	NODE=M181G02 NODE=M181G02
<sup>1</sup> Using B(η → γγ) = (39.41 ± 0.21)%.					NODE=M181G02;LINKAGE=A
$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{16}\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.9	90	<sup>1</sup> HAN	15 BELL	10.58 e <sup>+</sup> e <sup>-</sup> → χ <sub>c2</sub> γ	NODE=M181G03 NODE=M181G03
<sup>1</sup> Using B(η → γγ) = (39.41 ± 0.21)%.					NODE=M181G03;LINKAGE=A

### ψ(4360) BRANCHING RATIOS

$\Gamma(h_c\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_2/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		ABLIKIM	17G BES3	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> h <sub>c</sub>	NODE=M181R08 NODE=M181R08
$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		<sup>1</sup> ABLIKIM	17V BES3	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> ψ(2S)	NODE=M181R00 NODE=M181R00
<sup>1</sup> From a fit to the cross section for e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> ψ(2S) → 2(π <sup>+</sup> π <sup>-</sup> )ℓ <sup>+</sup> ℓ <sup>-</sup> obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb <sup>-1</sup> .					NODE=M181R00;LINKAGE=A
$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$					$\Gamma_4/\Gamma_3$
VALUE		DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
(0.81 ± 0.12 ± 0.13) to (42 ± 15 ± 15)		<sup>1</sup> ZHANG	17C RVUE	e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> J/ψ or ψ(2S)	NODE=M181R04 NODE=M181R04
<sup>1</sup> From a combined fit of BELLE, BABAR and BES3 e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> J/ψ and e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> ψ(2S) data.					NODE=M181R04;LINKAGE=A



$\Gamma(\psi(3770)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	<sup>1</sup> ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^- D\bar{D}$

NODE=M181R06  
NODE=M181R06

<sup>1</sup> Observe  $e^+e^- \rightarrow \pi^+\pi^-\psi(3770)$  at  $\sqrt{s} = 4.26, 4.36,$  and  $4.42$  GeV but cannot establish if continuum or resonant.

NODE=M181R06;LINKAGE=A

 $\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	[possibly seen OUR 2022 BEST LIMIT]			

NODE=M181R03  
NODE=M181R03

<b>seen</b>	<sup>1</sup> ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
-------------	----------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<b>possibly seen</b>	19	<sup>2</sup> ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
----------------------	----	----------------------	----------	--

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances.

NODE=M181R03;LINKAGE=C

<sup>2</sup> From a fit of  $e^+e^- \rightarrow \pi^+\pi^-\psi_2(3823), \psi_2(3823) \rightarrow \chi_{c1}\gamma$  cross sections taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the  $\psi(4360)$  line shape.

NODE=M181R03;LINKAGE=A

 $\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$

NODE=M181R07  
NODE=M181R07

<sup>1</sup> With a significance of  $6.0\sigma$ .

NODE=M181R07;LINKAGE=A

 $\Gamma(D^0 D^{*-}\pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma \times \Gamma_1/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.72 \times 10^{-6}$	90	<sup>1</sup> PAKHLOVA	09 BELL	$e^+e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-}\pi^+$

NODE=M181R02  
NODE=M181R02

<sup>1</sup> Using  $4355_{-10}^{+9} \pm 9$  MeV for the mass of  $\psi(4360)$ .

NODE=M181R02;LINKAGE=PA

 $\Gamma(D^0 D^{*-}\pi^+)/\Gamma(\psi(2S)\pi^+\pi^-)$  $\Gamma_8/\Gamma_4$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8$	90	PAKHLOVA	09 BELL	$e^+e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-}\pi^+$

NODE=M181R01  
NODE=M181R01 $\Gamma(D^+ D^- \pi^+ \pi^-)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> ABLIKIM	22AL BES3	$e^+e^- \rightarrow \pi^+\pi^- D^+ D^-$

NODE=M181R14  
NODE=M181R14

<sup>1</sup> From a fit to the cross section for  $e^+e^- \rightarrow D^+ D^- \pi^+ \pi^-$  in the range  $\sqrt{s} = 4.190-4.946$  GeV.

NODE=M181R14;LINKAGE=A

 $\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>possibly seen</b>	<sup>1</sup> ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^- D\bar{D}$

NODE=M181R05  
NODE=M181R05

<sup>1</sup> Evidence for  $e^+e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$  between  $\sqrt{s} = 4.3$  and  $4.6$  GeV, not necessarily resonant.

NODE=M181R05;LINKAGE=A

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	22K BES3	$e^+e^- \rightarrow \omega\pi^0$

NODE=M181R12  
NODE=M181R12 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	22K BES3	$e^+e^- \rightarrow \omega\eta$

NODE=M181R13  
NODE=M181R13 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	21AN BES3	$e^+e^- \rightarrow p\bar{p}\eta$

NODE=M181R09  
NODE=M181R09 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	ABLIKIM	21AN BES3	$e^+e^- \rightarrow p\bar{p}\omega$

NODE=M181R10  
NODE=M181R10

$\psi(4360)$  REFERENCES

ABLIKIM	22AL	PR D106 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61884
ABLIKIM	22AM	PR D106 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61885
ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61664
ABLIKIM	21AJ	PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61441
ABLIKIM	21AK	PR D104 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61443
ABLIKIM	21AN	PR D104 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61446
ABLIKIM	20O	PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60344
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ABLIKIM	19K	PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57755
ABLIKIM	17G	PRL 118 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57915
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58029
Also		PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang		REFID=58219
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan		REFID=58463
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=56839
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55938
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=52296
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51724
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51959

NODE=M181

 $\psi(4415)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M073

 $\psi(4415)$  MASS

NODE=M073M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>4421 ± 4 OUR ESTIMATE</b>			
<b>4415.1 ± 7.9</b>	<sup>1</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4412 ± 15	<sup>2</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4411 ± 7	<sup>3</sup> PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	<sup>4</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4429 ± 9	<sup>5</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C DASP	$e^+e^-$
4414 ± 7	SIEGRIST	76 MRK1	$e^+e^-$

NODE=M073M

→ UNCHECKED ←

<sup>1</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (234 \pm 88)^\circ$ .

NODE=M073M;LINKAGE=AB

<sup>2</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

NODE=M073M;LINKAGE=MO

<sup>3</sup> Systematic uncertainties not estimated.

NODE=M073M;LINKAGE=NS

<sup>4</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M073M;LINKAGE=ST

<sup>5</sup> From a fit to BES (BAI 02C) data.

NODE=M073M;LINKAGE=SE

 $\psi(4415)$  WIDTH

NODE=M073W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>62 ± 20 OUR ESTIMATE</b>			
<b>71.5 ± 19.0</b>	<sup>6</sup> ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
118 ± 32	<sup>7</sup> MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
77 ± 20	<sup>8</sup> PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ± 16	<sup>9</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
118 ± 35	<sup>10</sup> SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
66 ± 15	BRANDELIK	78C DASP	$e^+e^-$
33 ± 10	SIEGRIST	76 MRK1	$e^+e^-$

NODE=M073W

→ UNCHECKED ←

<sup>6</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (234 \pm 88)^\circ$ .

NODE=M073W;LINKAGE=AB

<sup>7</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects.

NODE=M073W;LINKAGE=MO

<sup>8</sup> Systematic uncertainties not estimated.

NODE=M073W;LINKAGE=NS

<sup>9</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M073W;LINKAGE=ST

<sup>10</sup> From a fit to BES (BAI 02C) data.

NODE=M073W;LINKAGE=SE

**$\psi(4415)$  DECAY MODES**

Due to the complexity of the  $c\bar{c}$  threshold region, in this listing, "seen" ("not seen") means that a cross section for the mode in question has been measured at effective  $\sqrt{s}$  near this particle's central mass value, more (less) than  $2\sigma$  above zero, without regard to any peaking behavior in  $\sqrt{s}$  or absence thereof. See mode listing(s) for details and references.

NODE=M073215;NODE=M073

NODE=M073

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $D\bar{D}$	seen	
$\Gamma_2$ $D^0\bar{D}^0$	seen	
$\Gamma_3$ $D^+D^-$	seen	
$\Gamma_4$ $D^*\bar{D} + \text{c.c.}$	seen	
$\Gamma_5$ $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	
$\Gamma_6$ $D^*(2010)^+D^- + \text{c.c.}$	seen	
$\Gamma_7$ $D^*\bar{D}^*$	seen	
$\Gamma_8$ $D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen	
$\Gamma_9$ $D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen	
$\Gamma_{10}$ $D^0D^-\pi^+$ (excl. $D^*(2007)^0\bar{D}^0$ +c.c., $D^*(2010)^+D^-$ +c.c.)	< 2.3 %	90%
$\Gamma_{11}$ $D\bar{D}_2^*(2460) \rightarrow D^0D^-\pi^+$ +c.c.	(10 $\pm$ 4) %	
$\Gamma_{12}$ $D^0D^{*-}\pi^+$ +c.c.	< 11 %	90%
$\Gamma_{13}$ $D_1(2420)\bar{D} + \text{c.c.}$	possibly seen	
$\Gamma_{14}$ $D_s^+D_s^-$	not seen	
$\Gamma_{15}$ $\omega\chi_{c2}$	possibly seen	
$\Gamma_{16}$ $D_s^{*+}D_s^- + \text{c.c.}$	seen	
$\Gamma_{17}$ $D_s^{*+}D_s^{*-}$	not seen	
$\Gamma_{18}$ $\psi_2(3823)\pi^+\pi^-$	possibly seen	
$\Gamma_{19}$ $\psi(3770)\pi^+\pi^-$	possibly seen	
$\Gamma_{20}$ $J/\psi\eta$	< 6 $\times 10^{-3}$	90%
$\Gamma_{21}$ $\chi_{c1}\gamma$	< 8 $\times 10^{-4}$	90%
$\Gamma_{22}$ $\chi_{c2}\gamma$	< 4 $\times 10^{-3}$	90%
$\Gamma_{23}$ $\Lambda\bar{\Lambda}$	< 3.1 $\times 10^{-6}$	90%
$\Gamma_{24}$ $\omega\pi^0$	not seen	
$\Gamma_{25}$ $\omega\eta$	not seen	
$\Gamma_{26}$ $e^+e^-$	(9.4 $\pm$ 3.2) $\times 10^{-6}$	
$\Gamma_{27}$ $\mu^+\mu^-$	(2.0 $\pm$ 1.0) $\times 10^{-5}$	

DESIG=7;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=8  
DESIG=9  
DESIG=10;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=11  
DESIG=12  
DESIG=13;OUR EVAL; $\rightarrow$  UNCHECKED  $\leftarrow$   
DESIG=14  
DESIG=15  
DESIG=4  
DESIG=5  
DESIG=6  
DESIG=25  
DESIG=16  
DESIG=20  
DESIG=17  
DESIG=18  
DESIG=21  
DESIG=24  
DESIG=19  
DESIG=22  
DESIG=23  
DESIG=27  
DESIG=28  
DESIG=29  
DESIG=1  
DESIG=26

 **$\psi(4415)$  PARTIAL WIDTHS**

NODE=M073220

 **$\Gamma(e^+e^-)$**  **$\Gamma_{26}$** 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.58<math>\pm</math>0.07 OUR ESTIMATE</b>			
<b>0.35<math>\pm</math>0.12</b>	11 ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.4 to 0.8	12 MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
0.72 $\pm$ 0.11	13 SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.64 $\pm$ 0.23	14 SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
0.49 $\pm$ 0.13	BRANDELIK	78C DASP	$e^+e^-$
0.44 $\pm$ 0.14	SIEGRIST	76 MRK1	$e^+e^-$

NODE=M073W1  
NODE=M073W1  
 $\rightarrow$  UNCHECKED  $\leftarrow$

OCCUR=2

<sup>11</sup> Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the  $\psi(3770)$ ,  $\psi(4040)$ ,  $\psi(4160)$ , and  $\psi(4415)$  resonances. Phase angle fixed in the fit to  $\delta = (234 \pm 88)^\circ$ .

NODE=M073W1;LINKAGE=AB

<sup>12</sup> Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the  $\psi(4040)$ ,  $\psi(4160)$  and  $\psi(4415)$  resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different  $e^+e^-$  partial widths. We quote only the range of values.

NODE=M073W1;LINKAGE=MO

<sup>13</sup> From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M073W1;LINKAGE=ST

<sup>14</sup> From a fit to BES (BAI 02C) data.

NODE=M073W1;LINKAGE=SE

$\Gamma(\mu^+\mu^-)$  $\Gamma_{27}$ 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$1.25 \pm 0.28 \pm 0.35$	<sup>15,16</sup> ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M073W2  
 NODE=M073W2

<sup>15</sup> From a fit to the  $e^+e^- \rightarrow \mu^+\mu^-$  cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming  $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$ .

NODE=M073W2;LINKAGE=A

<sup>16</sup> From solution 1 of 8 with equal fit quality. Other solutions range from  $1.24 \pm 0.28 \pm 0.35$  to  $1.27 \pm 0.41 \pm 0.36$  keV.

NODE=M073W2;LINKAGE=B

 $\psi(4415) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M073230

 $\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{20}\Gamma_{26}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6$	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

NODE=M073G01  
 NODE=M073G01

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{21}\Gamma_{26}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<0.47$	90	<sup>17</sup> HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

NODE=M073G02  
 NODE=M073G02

<sup>17</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

NODE=M073G02;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{22}\Gamma_{26}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<2.3$	90	<sup>18</sup> HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

NODE=M073G03  
 NODE=M073G03

<sup>18</sup> Using  $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .

NODE=M073G03;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_{23}\Gamma_{26}/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-3}$	90	<sup>19</sup> ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4415)$

NODE=M073R16  
 NODE=M073R16

<sup>19</sup> From a measurement of the  $e^+e^- \rightarrow \Lambda\bar{\Lambda}$  cross section between 3.5 and 4.6 GeV.

NODE=M073R16;LINKAGE=A

 $\psi(4415)$  BRANCHING RATIOS

NODE=M073225

 $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 08 BELL		$e^+e^- \rightarrow D^0\bar{D}^0\gamma$
not seen	AUBERT 09M BABR		$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

NODE=M073R04  
 NODE=M073R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen AUBERT 09M BABR  $e^+e^- \rightarrow D^0\bar{D}^0\gamma$

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 08 BELL		$e^+e^- \rightarrow D^+D^-\gamma$
not seen	AUBERT 09M BABR		$e^+e^- \rightarrow D^+D^-\gamma$

NODE=M073R05  
 NODE=M073R05

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen AUBERT 09M BABR  $e^+e^- \rightarrow D^+D^-\gamma$

 $\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$  $\Gamma_1/\Gamma_7$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.14 \pm 0.12 \pm 0.03$	AUBERT 09M BABR		$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

NODE=M073R02  
 NODE=M073R02

 $\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M BABR		$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$

NODE=M073R06  
 NODE=M073R06

 $\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>20</sup> ZHUKOVA 18 BELL		$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	AUBERT 09M BABR		$e^+e^- \rightarrow D^{*+}D^-\gamma$

NODE=M073R07  
 NODE=M073R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen PAKHLOVA 07 BELL  $e^+e^- \rightarrow D^{*+}D^-\gamma$

<sup>20</sup> Supersedes PAKHLOVA 07.

NODE=M073R07;LINKAGE=A

 $\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$  $\Gamma_4/\Gamma_7$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.17 \pm 0.25 \pm 0.03$	AUBERT 09M BABR		$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

NODE=M073R03  
 NODE=M073R03

$$\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0 + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_8 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$

NODE=M073R08  
NODE=M073R08

$$\Gamma(D^*(2010)^+ D^*(2010)^- + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>21</sup> ZHUKOVA	18 BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$

NODE=M073R09  
NODE=M073R09

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	PAKHLOVA	07 BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
------	----------	---------	--

<sup>21</sup> Supersedes PAKHLOVA 07.

NODE=M073R09;LINKAGE=A

$$\Gamma(D \bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{11} / \Gamma$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
$10.5 \pm 2.4 \pm 3.8$	<sup>22</sup> PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

<sup>22</sup> Using  $4421 \pm 4$  MeV for the mass and  $62 \pm 20$  MeV for the width of  $\psi(4415)$ .

NODE=M073R3  
NODE=M073R3

NODE=M073R3;LINKAGE=PA

$$\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c.}, D^*(2010)^+ D^- + \text{c.c.}) / \Gamma(D \bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.}) \quad \Gamma_{10} / \Gamma_{11}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.22$	90	<sup>23</sup> PAKHLOVA	08A BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$

<sup>23</sup> Using  $4421 \pm 4$  MeV for the mass and  $62 \pm 20$  MeV for the width of  $\psi(4415)$ .

NODE=M073R4  
NODE=M073R4

NODE=M073R4;LINKAGE=PA

$$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma \times \Gamma_{26} / \Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.99 \times 10^{-6}$	90	<sup>24</sup> PAKHLOVA	09 BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

<sup>24</sup> Using  $4421 \pm 4$  MeV for the mass of  $\psi(4415)$ .

NODE=M073R01  
NODE=M073R01

NODE=M073R01;LINKAGE=PA

$$\Gamma(D_1(2420) \bar{D} + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	<sup>25</sup> ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$

<sup>25</sup> Evidence for  $e^+ e^- \rightarrow D_1(2420) \bar{D} + \text{c.c.}$  between  $\sqrt{s} = 4.3$  and 4.6 GeV, not necessarily resonant.

NODE=M073R15  
NODE=M073R15

NODE=M073R15;LINKAGE=A

$$\Gamma(D_s^+ D_s^-) / \Gamma_{\text{total}} \quad \Gamma_{14} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

NODE=M073R10  
NODE=M073R10

$$\Gamma(\omega \chi_{c2}) / \Gamma_{\text{total}} \quad \Gamma_{15} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	ABLIKIM	16A BES3	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$

NODE=M073R00  
NODE=M073R00

$$\Gamma(D_s^{*+} D_s^- + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{16} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

NODE=M073R11  
NODE=M073R11

$$\Gamma(D_s^{*+} D_s^{*-}) / \Gamma_{\text{total}} \quad \Gamma_{17} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$

NODE=M073R12  
NODE=M073R12

$$\Gamma(\psi(3770) \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{19} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	<sup>26</sup> ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$

<sup>26</sup> Observe  $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$  at  $\sqrt{s} = 4.26, 4.36, \text{ and } 4.42$  GeV but cannot establish if continuum or resonant.

NODE=M073R14  
NODE=M073R14

NODE=M073R14;LINKAGE=A

$$\Gamma(\psi_2(3823) \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{18} / \Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
possibly seen	19	<sup>27</sup> ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$

<sup>27</sup> From a fit of  $e^+ e^- \rightarrow \pi^+ \pi^- \psi_2(3823)$ ,  $\psi_2(3823) \rightarrow \chi_{c1} \gamma$  cross sections taken at  $\sqrt{s}$  values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the  $\psi(4415)$  line shape.

NODE=M073R13  
NODE=M073R13

NODE=M073R13;LINKAGE=A

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ 

VALUE

not seen

DOCUMENT ID

TECN

COMMENT

ABLIKIM

22K

BES3

 $e^+e^- \rightarrow \omega\pi^0$  $\Gamma_{24}/\Gamma$ NODE=M073R17  
NODE=M073R17 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ 

VALUE

not seen

DOCUMENT ID

TECN

COMMENT

ABLIKIM

22K

BES3

 $e^+e^- \rightarrow \omega\eta$  $\Gamma_{25}/\Gamma$ NODE=M073R18  
NODE=M073R18 $\psi(4415)$  REFERENCES

NODE=M073

ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
ABLIKIM	16A	PR D93 011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57122
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)	REFID=22243

NODE=M234

 $\chi_{c0}(4500)$ 

$$I^{G(J^{PC})} = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE  
was  $X(4500)$ This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

NODE=M234

Seen by AAIJ 17C in  $B^+ \rightarrow \chi_{c0} K^+$ ,  $\chi_{c0} \rightarrow J/\psi\phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  with a significance (accounting for systematic uncertainties) of 6.1  $\sigma$ . $\chi_{c0}(4500)$  MASS

NODE=M234M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

**4474 ± 3 ± 3** 24k <sup>1</sup> AAIJ 21E LHCB  $B^+ \rightarrow J/\psi\phi K^+$ 

••• We do not use the following data for averages, fits, limits, etc. •••

4506 ± 11<sup>+12</sup><sub>-15</sub> 4289 <sup>2,3</sup> AAIJ 17C LHCB  $B^+ \rightarrow J/\psi\phi K^+$ <sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 20  $\sigma$ .<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 6.1  $\sigma$ .<sup>3</sup> Superseded by AAIJ 21E.NODE=M234M;LINKAGE=B  
NODE=M234M;LINKAGE=A  
NODE=M234M;LINKAGE=C $\chi_{c0}(4500)$  WIDTH

NODE=M234W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

**77 ± 6<sup>+10</sup><sub>-8</sub>** 24k <sup>1</sup> AAIJ 21E LHCB  $B^+ \rightarrow J/\psi\phi K^+$ 

••• We do not use the following data for averages, fits, limits, etc. •••

92 ± 21<sup>+21</sup><sub>-20</sub> 4289 <sup>2,3</sup> AAIJ 17C LHCB  $B^+ \rightarrow J/\psi\phi K^+$ <sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 20  $\sigma$ .<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of 6.1  $\sigma$ .<sup>3</sup> Superseded by AAIJ 21E.NODE=M234W;LINKAGE=B  
NODE=M234W;LINKAGE=A  
NODE=M234W;LINKAGE=C

$\chi_{c0}(4500)$  DECAY MODES

NODE=M234215;NODE=M234

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi\phi$	seen

DESIG=1

 $\chi_{c0}(4500)$  BRANCHING RATIOS

NODE=M234220

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
seen	4289	<sup>2,3</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of $20\sigma$ .					
<sup>2</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of $6.1\sigma$ .					
<sup>3</sup> Superseded by AAIJ 21E.					

NODE=M234R01  
NODE=M234R01NODE=M234R01;LINKAGE=B  
NODE=M234R01;LINKAGE=A  
NODE=M234R01;LINKAGE=C $\chi_{c0}(4500)$  REFERENCES

NODE=M234

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636

REFID=61150  
REFID=57657  
REFID=57636

NODE=M262

**X(4630)**

$$J^G(J^{PC}) = 0^+(?^?+)$$

## OMITTED FROM SUMMARY TABLE

NODE=M262

This state shows properties different from a conventional  $q\bar{q}$  state. A candidate for an exotic structure. See the review on "Heavy Non- $q\bar{q}$  Mesons."

Seen by AAIJ 21E in  $B^+ \rightarrow X(4630)K^+$  with  $X(4630) \rightarrow J/\psi\phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  with a significance (accounting for systematic uncertainties) of  $5.5\sigma$ . The  $J^P = 1^-$  assignment is favored over  $2^-$  with a significance of  $3\sigma$  and other assignments are disfavored by more than  $5\sigma$ .

**X(4630) MASS**

NODE=M262M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4626 \pm 16^{+18}_{-110}$	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi\phi K^+$
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of $5.5\sigma$ .				

NODE=M262M

NODE=M262M;LINKAGE=A

**X(4630) WIDTH**

NODE=M262W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$174 \pm 27^{+134}_{-73}$	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi\phi K^+$
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of $5.5\sigma$ .				

NODE=M262W

NODE=M262W;LINKAGE=A

**X(4630) DECAY MODES**

NODE=M262215;NODE=M262

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi\phi$	seen

DESIG=1

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
<sup>1</sup> From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of $5.5\sigma$ .					

NODE=M262R01  
NODE=M262R01

NODE=M262R01;LINKAGE=A

**X(4630) REFERENCES**

NODE=M262

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
------	-----	----------------	-----------------------	----------------	-------------

REFID=61150

**$\psi(4660)$** 

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as  $Y(4660)$ ; was  $X(4660)$ 

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

Seen in radiative return from  $e^+e^-$  collisions at  $\sqrt{s} = 9.54\text{--}10.58$  GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

NODE=M189

NODE=M189

 **$\psi(4660)$  MASS**

NODE=M189M

NODE=M189M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4630 ± 6</b>	<b>OUR AVERAGE</b>	Error includes scale factor of 1.4. See the ideogram below.		
4651.0 ± 37.8 ± 2.1		<sup>1</sup> ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4619.8 <sup>+8.9</sup> <sub>-8.0</sub> ± 2.3	66	<sup>2</sup> JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$
4625.9 <sup>+6.2</sup> <sub>-6.0</sub> ± 0.4	89	<sup>3</sup> JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
4652 ± 10 ± 11	279	<sup>4</sup> WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4669 ± 21 ± 3	37	<sup>5</sup> LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4634 <sup>+8</sup> <sub>-7</sub> <sup>+5</sup> <sub>-8</sub>	142	<sup>6</sup> PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4647.9 ± 8.6 ± 0.8		<sup>7</sup> ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
4652.5 ± 3.4 ± 1.1		<sup>8</sup> DAI	17 RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
4645.2 ± 9.5 ± 6.0		<sup>9</sup> ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4646.4 ± 9.7 ± 4.8		<sup>10</sup> ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
4661 <sup>+9</sup> <sub>-8</sub> ± 6	44	<sup>11</sup> LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4664 ± 11 ± 5	44	WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

<sup>1</sup> From a three-resonance fit to the Born cross section in the range  $\sqrt{s} = 4.008\text{--}4.698$  GeV.

<sup>2</sup> Using  $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$  decays.

<sup>3</sup> From a fit of a Breit-Wigner convolved with a Gaussian.

<sup>4</sup> From a two-resonance fit. Supersedes WANG 07D.

<sup>5</sup> From a two-resonance fit.

<sup>6</sup> The  $\pi^+\pi^-\psi(2S)$  and  $\Lambda_c^+ \Lambda_c^-$  states are not necessarily the same.

<sup>7</sup> From a fit to the  $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass  $4417.5 \pm 26.2 \pm 3.5$  MeV and width  $245 \pm 48 \pm 13$  MeV.

<sup>8</sup> The pole parameters are extracted from the speed plot.

<sup>9</sup> From a three-resonance fit.

<sup>10</sup> From a combined fit of BELLE, BABAR and BES3  $e^+e^- \rightarrow \pi^+\pi^- J/\psi$  and  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$  data.

<sup>11</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189M;LINKAGE=G

NODE=M189M;LINKAGE=F

NODE=M189M;LINKAGE=E

NODE=M189M;LINKAGE=A

NODE=M189M;LINKAGE=LE

NODE=M189M;LINKAGE=PA

NODE=M189M;LINKAGE=I


NODE=M189M;LINKAGE=C

NODE=M189M;LINKAGE=D

NODE=M189M;LINKAGE=B

NODE=M189M;LINKAGE=LI





**psi(4660) WIDTH**

NODE=M189W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M189W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>72 <math>^{+14}_{-12}</math> OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.		
155.4 ± 24.8 ± 0.8		1 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
47.0 $^{+31.3}_{-14.8}$ ± 4.6	66	2 JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$
49.8 $^{+13.9}_{-11.5}$ ± 4.0	89	3 JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
68 ± 11 ± 5	279	4 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
104 ± 48 ± 10	37	5 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
92 $^{+40}_{-24}$ $^{+10}_{-21}$	142	6 PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
33.1 ± 18.6 ± 4.1		7 ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
62.6 ± 5.6 ± 4.3		8 DAI	17 RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
113.8 ± 18.1 ± 3.4		9 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
103.5 ± 15.6 ± 4.0		10 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
42 $^{+17}_{-12}$ ± 6	44	11 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
48 ± 15 ± 3	44	WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+\pi^-\psi(2S)$
<sup>1</sup> From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008-4.698$ GeV. <sup>2</sup> Using $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$ decays. <sup>3</sup> From a fit of a Breit-Wigner convolved with a Gaussian. <sup>4</sup> From a two-resonance fit. Supersedes WANG 07D. <sup>5</sup> From a two-resonance fit. <sup>6</sup> The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same. <sup>7</sup> From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV. <sup>8</sup> The pole parameters are extracted from the speed plot. <sup>9</sup> From a three-resonance fit. <sup>10</sup> From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data. <sup>11</sup> From a combined fit of AUBERT 07S and WANG 07D data with two resonances.				


NODE=M189W;LINKAGE=H

NODE=M189W;LINKAGE=G  
 NODE=M189W;LINKAGE=F  
 NODE=M189W;LINKAGE=A  
 NODE=M189W;LINKAGE=LE

NODE=M189W;LINKAGE=B  
 NODE=M189W;LINKAGE=I

NODE=M189W;LINKAGE=D  
 NODE=M189W;LINKAGE=E  
 NODE=M189W;LINKAGE=C

NODE=M189W;LINKAGE=LI



**psi(4660) DECAY MODES**

NODE=M189215;NODE=M189

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $e^+ e^-$	not seen
$\Gamma_2$ $\psi(2S)\pi^+\pi^-$	seen
$\Gamma_3$ $J/\psi\eta$	not seen
$\Gamma_4$ $D^0 D^{*-}\pi^+$	not seen
$\Gamma_5$ $\psi_2(3823)\pi^+\pi^-$	seen
$\Gamma_6$ $\chi_{c1}\gamma$	not seen
$\Gamma_7$ $\chi_{c2}\gamma$	not seen
$\Gamma_8$ $\Lambda_c^+ \Lambda_c^-$	seen
$\Gamma_9$ $D_s^+ D_{s1}(2536)^-$	seen
$\Gamma_{10}$ $D_s^+ D_{s2}^*(2573)^-$	seen
$\Gamma_{11}$ $\omega\pi^0$	not seen
$\Gamma_{12}$ $\omega\eta$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←  
 DESIG=2;OUR EVAL;→ UNCHECKED ←  
 DESIG=4;OUR EVAL;→ UNCHECKED ←  
 DESIG=3;OUR EVAL;→ UNCHECKED ←  
 DESIG=10  
 DESIG=6;OUR EVAL;→ UNCHECKED ←  
 DESIG=7;OUR EVAL;→ UNCHECKED ←  
 DESIG=5;OUR EVAL;→ UNCHECKED ←  
 DESIG=8;OUR EVAL;→ UNCHECKED ←  
 DESIG=9  
 DESIG=11  
 DESIG=12

**psi(4660)  $\Gamma(i) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$**

NODE=M189230

$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
4.7±3.8		1 ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+\pi^-\psi(2S)$	
11.2±3.2		2 ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=2
4.7±4.2		3 ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=3
11.3±3.3		4 ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=4
2.0±0.3±0.2	279	5 WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
8.1±1.1±1.0	279	6 WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2
2.7±1.3±0.5	37	7 LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
7.5±1.7±0.7	37	8 LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2
2.2 <sup>+0.7</sup> <sub>-0.6</sub>	44	9 LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
5.9±1.6	44	10 LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2
3.0±0.9±0.3	44	7 WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
7.6±1.8±0.8	44	8 WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2

NODE=M189G1  
 NODE=M189G1

- <sup>1</sup> Solution I of four equivalent solutions in a fit using three interfering resonances.
- <sup>2</sup> Solution II of four equivalent solutions in a fit using three interfering resonances.
- <sup>3</sup> Solution III of four equivalent solutions in a fit using three interfering resonances.
- <sup>4</sup> Solution IV of four equivalent solutions in a fit using three interfering resonances.
- <sup>5</sup> Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.
- <sup>6</sup> Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.
- <sup>7</sup> Solution I of two equivalent solutions in a fit using two interfering resonances.
- <sup>8</sup> Solution II of two equivalent solutions in a fit using two interfering resonances.
- <sup>9</sup> Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.
- <sup>10</sup> Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189G1;LINKAGE=C  
 NODE=M189G1;LINKAGE=D  
 NODE=M189G1;LINKAGE=E  
 NODE=M189G1;LINKAGE=F  
 NODE=M189G1;LINKAGE=A

NODE=M189G1;LINKAGE=B

NODE=M189G1;LINKAGE=WA  
 NODE=M189G1;LINKAGE=WN  
 NODE=M189G1;LINKAGE=LI  
 NODE=M189G1;LINKAGE=LU

$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_5/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	<sup>1</sup> ABLIKIM	22R	BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$	NODE=M189R05 NODE=M189R05

<sup>1</sup> From a fit to the  $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$  cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances.

NODE=M189R05;LINKAGE=A

$\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.94	90	WANG	13B	BELL $e^+e^- \rightarrow J/\psi\eta\gamma$	NODE=M189G01 NODE=M189G01

$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_6\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.45	90	<sup>1</sup> HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c1}\gamma$	NODE=M189G02 NODE=M189G02
				<sup>1</sup> Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .	NODE=M189G02;LINKAGE=A

$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.1	90	<sup>1</sup> HAN	15	BELL $10.58 e^+e^- \rightarrow \chi_{c2}\gamma$	NODE=M189G03 NODE=M189G03
				<sup>1</sup> Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$ .	NODE=M189G03;LINKAGE=A

$\Gamma(D_s^+ D_{s1}^-(2536)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_9\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$14.3_{-2.6}^{+2.8} \pm 1.5$	89	<sup>1</sup> JIA	19A	BELL $e^+e^- \rightarrow \gamma D_s^+ D_{s1}^-(2536)^-$	NODE=M189R00 NODE=M189R00
				<sup>1</sup> Assuming $B(D_{s1}^-(2536)^- \rightarrow \bar{D}^{*0} K^-) = 1$ .	NODE=M189R00;LINKAGE=A

$\Gamma(D_s^+ D_{s2}^*(2573)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_{10}\Gamma_1/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$14.7_{-4.5}^{+5.9} \pm 3.6$	66	<sup>1</sup> JIA	20	BELL $e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$	NODE=M189R04 NODE=M189R04
				<sup>1</sup> Assuming $B(D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-) = 1$ .	NODE=M189R04;LINKAGE=A

### $\psi(4660)$ BRANCHING RATIOS

$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S)\pi^+\pi^-)$					$\Gamma_4/\Gamma_2$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<10	90	PAKHOVA	09	BELL $e^+e^- \rightarrow D^0 D^{*-} \pi^+$	NODE=M189R01 NODE=M189R01

$\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma \times \Gamma_1/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
< $0.37 \times 10^{-6}$	90	<sup>1</sup> PAKHOVA	09	BELL $e^+e^- \rightarrow D^0 D^{*-} \pi^+$	NODE=M189R02 NODE=M189R02
				<sup>1</sup> Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$ .	NODE=M189R02;LINKAGE=PA

$\Gamma(\Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_8/\Gamma \times \Gamma_1/\Gamma$
VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.68_{-0.15}^{+0.16+0.29} - 0.30$	142	<sup>1</sup> PAKHOVA	08B	BELL $e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$	NODE=M189R03 NODE=M189R03

<sup>1</sup> The  $\pi^+\pi^-\psi(2S)$  and  $\Lambda_c^+\Lambda_c^-$  states are not necessarily the same.

NODE=M189225

NODE=M189R03;LINKAGE=A

$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ 

VALUE

not seen

DOCUMENT ID

TECN

COMMENT

ABLIKIM

22K

BES3

 $e^+e^- \rightarrow \omega\pi^0$  $\Gamma_{11}/\Gamma$ NODE=M189R06  
NODE=M189R06 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ 

VALUE

not seen

DOCUMENT ID

TECN

COMMENT

ABLIKIM

22K

BES3

 $e^+e^- \rightarrow \omega\eta$  $\Gamma_{12}/\Gamma$ NODE=M189R07  
NODE=M189R07 $\psi(4660)$  REFERENCES

NODE=M189

ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AJ	PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
JIA	20	PR D101 091101	S. Jia <i>et al.</i>	(BELLE Collab.)
JIA	19A	PR D100 111103	S. Jia <i>et al.</i>	(BELLE Collab.)
DAI	17	PR D96 116001	L.-Y. Dai, J. Haidenbauer, U.-G. Meissner	(JULI+)
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang	
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan	
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	
PAKHLOVA	08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)

REFID=61648  
REFID=61664  
REFID=61441  
REFID=60301  
REFID=60037  
REFID=58704  
REFID=58219  
REFID=58463  
REFID=56816  
REFID=56839  
REFID=55938  
REFID=55377  
REFID=53143  
REFID=52296  
REFID=52596  
REFID=51724  
REFID=51959 $\chi_{c1}(4685)$  $I^G(J^{PC}) = 0^+(1^{++})$ 

NODE=M261

## OMITTED FROM SUMMARY TABLE

This state shows properties different from a conventional  $q\bar{q}$  state. A candidate for an exotic structure. See the review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M261

Seen by AAIJ 21E in  $B^+ \rightarrow \chi_{c1}(4685)K^+$  with  $\chi_{c1}(4685) \rightarrow J/\psi\phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi\phi K^+$  with a significance (accounting for systematic uncertainties) of  $15\sigma$ . The  $J^P = 1^+$  assignment is favored with high significance.

 $\chi_{c1}(4685)$  MASS

NODE=M261M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4684 \pm 7^{+13}_{-16}$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M261M

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $15\sigma$ .

NODE=M261M;LINKAGE=A

 $\chi_{c1}(4685)$  WIDTH

NODE=M261W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$126 \pm 15^{+37}_{-41}$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M261W

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $15\sigma$ .

NODE=M261W;LINKAGE=A

 $\chi_{c1}(4685)$  DECAY MODES

NODE=M261215;NODE=M261

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi\phi$	seen

DESIG=1

 $\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$ 

VALUE

seen

EVTS

DOCUMENT ID

TECN

COMMENT

24k

<sup>1</sup> AAIJ

21E

LHCB

 $B^+ \rightarrow J/\psi\phi K^+$  $\Gamma_1/\Gamma$ NODE=M261R01  
NODE=M261R01

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi\phi K^+$  with a significance of  $15\sigma$ .

NODE=M261R01;LINKAGE=A

 $\chi_{c1}(4685)$  REFERENCES

NODE=M261

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
------	-----	----------------	-----------------------	-------------------

REFID=61150

$\chi_{c0}(4700)$ 

$$J^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE  
was  $X(4700)$

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

Seen by AAIJ 17C in  $B^+ \rightarrow \chi_{c0} K^+$ ,  $\chi_{c0} \rightarrow J/\psi \phi$  using an amplitude analysis of  $B^+ \rightarrow J/\psi \phi K^+$  with a significance (accounting for systematic uncertainties) of  $5.6 \sigma$ .

NODE=M235

NODE=M235

 **$\chi_{c0}(4700)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4694 \pm 4^{+16}_{-3}$	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$
$4741 \pm 6 \pm 6$	175	<sup>2</sup> AAIJ	21C LHCb	$B_s^0 \rightarrow J/\psi \phi \pi^+ \pi^-$
$4704 \pm 10^{+14}_{-24}$	4289	<sup>3,4</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $17 \sigma$ .

<sup>2</sup> From a 1D fit to the  $J/\psi \phi$  mass distribution with a significance of  $5.3 \sigma$ . The identification of this structure as the  $\chi_{c0}(4700)$  needs confirmation.

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.6 \sigma$ .

<sup>4</sup> Superseded by AAIJ 21E.

NODE=M235M

NODE=M235M

NODE=M235M;LINKAGE=C  
NODE=M235M;LINKAGE=BNODE=M235M;LINKAGE=A  
NODE=M235M;LINKAGE=D **$\chi_{c0}(4700)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$87 \pm 8^{+16}_{-6}$	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$
$53 \pm 15 \pm 11$	175	<sup>2</sup> AAIJ	21C LHCb	$B_s^0 \rightarrow J/\psi \phi \pi^+ \pi^-$
$120 \pm 31^{+42}_{-33}$	4289	<sup>3,4</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $17 \sigma$ .

<sup>2</sup> From a 1D fit to the  $J/\psi \phi$  mass distribution with a significance of  $5.3 \sigma$ . The identification of this structure as the  $\chi_{c0}(4700)$  needs confirmation.

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.6 \sigma$ .

<sup>4</sup> Superseded by AAIJ 21E.

NODE=M235W

NODE=M235W

NODE=M235W;LINKAGE=C  
NODE=M235W;LINKAGE=BNODE=M235W;LINKAGE=A  
NODE=M235W;LINKAGE=D **$\chi_{c0}(4700)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi \phi$	seen

NODE=M235215;NODE=M235

DESIG=1

 **$\chi_{c0}(4700)$  BRANCHING RATIOS**

$\Gamma(J/\psi \phi)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	<sup>1</sup> AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$
seen	175	<sup>2</sup> AAIJ	21C LHCb	$B_s^0 \rightarrow J/\psi \phi \pi^+ \pi^-$
seen	4289	<sup>3,4</sup> AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $17 \sigma$ .

<sup>2</sup> From a 1D fit to the  $J/\psi \phi$  mass distribution with a significance of  $5.3 \sigma$ . The identification of this structure as the  $\chi_{c0}(4700)$  needs confirmation.

<sup>3</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.6 \sigma$ .

<sup>4</sup> Superseded by AAIJ 21E.

NODE=M235220

NODE=M235R01  
NODE=M235R01NODE=M235R01;LINKAGE=C  
NODE=M235R01;LINKAGE=BNODE=M235R01;LINKAGE=A  
NODE=M235R01;LINKAGE=D **$\chi_{c0}(4700)$  REFERENCES**

AAIJ	21C	JHEP 2102 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M235


REFID=61124  
REFID=61150  
REFID=57657  
REFID=57636

# $b\bar{b}$ MESONS (including possibly non- $q\bar{q}$ states)

NODE=MXXX030

NODE=M849

Updated July 2021.



The level scheme of meson states containing a minimal quark content of  $b\bar{b}$ . The name of a state is determined by its quantum numbers  $I^G J^{PC}$  (see the review “Naming Scheme for Hadrons”). States included in the Summary Tables are shown with solid lines; those requiring confirmation are shown with dotted lines. The arrows indicate the most dominant hadronic transitions. Single photon transitions, including  $Y(nS) \rightarrow \gamma\eta_b(mS)$ ,  $Y(nS) \rightarrow \gamma\chi_{bJ}(mP)$ , and  $\chi_{bJ}(nP) \rightarrow \gamma Y(mS)$ , are omitted for clarity. For orientation, the location of the thresholds related to a pair of ground state open bottom mesons is indicated in the figure.

NODE=M849

## WIDTH DETERMINATIONS OF THE $Y$ STATES

As is the case for the  $J/\psi(1S)$  and  $\psi(2S)$ , the full widths of the  $b\bar{b}$  states  $Y(1S)$ ,  $Y(2S)$ , and  $Y(3S)$  are not directly measurable, since they are much narrower than the energy resolution of the  $e^+e^-$  storage rings where these states are produced. The common indirect method to determine  $\Gamma$  starts from

$$\Gamma = \Gamma_{\ell\ell}/B_{\ell\ell} , \quad (1)$$

where  $\Gamma_{\ell\ell}$  is one leptonic partial width and  $B_{\ell\ell}$  is the corresponding branching fraction ( $\ell = e, \mu, \text{ or } \tau$ ). One then assumes  $e\text{-}\mu\text{-}\tau$  universality and uses

$$\begin{aligned}\Gamma_{\ell\ell} &= \Gamma_{ee} \\ B_{\ell\ell} &= \text{average of } B_{ee}, B_{\mu\mu}, \text{ and } B_{\tau\tau} .\end{aligned}\quad (2)$$

The electronic partial width  $\Gamma_{ee}$  is also not directly measurable at  $e^+e^-$  storage rings, only in the combination  $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$ , where  $\Gamma_{\text{had}}$  is the hadronic partial width and

$$\Gamma_{\text{had}} + 3\Gamma_{ee} = \Gamma .\quad (3)$$

This combination is obtained experimentally from the energy-integrated hadronic cross section

$$\begin{aligned}\int_{\text{resonance}} \sigma(e^+e^- \rightarrow \Upsilon \rightarrow \text{hadrons})dE \\ = \frac{6\pi^2}{M^2} \frac{\Gamma_{ee}\Gamma_{\text{had}}}{\Gamma} C_r = \frac{6\pi^2}{M^2} \frac{\Gamma_{ee}^{(0)}\Gamma_{\text{had}}}{\Gamma} C_r^{(0)} ,\end{aligned}\quad (4)$$

where  $M$  is the  $\Upsilon$  mass, and  $C_r$  and  $C_r^{(0)}$  are radiative correction factors.  $C_r$  is used for obtaining  $\Gamma_{ee}$  as defined in Eq. (1), and contains corrections from all orders of QED for describing  $(b\bar{b}) \rightarrow e^+e^-$ . The lowest order QED value  $\Gamma_{ee}^{(0)}$ , relevant for comparison with potential-model calculations, is defined by the lowest order QED graph (Born term) alone, and is about 7% lower than  $\Gamma_{ee}$ .

The Listings give experimental results on  $B_{ee}$ ,  $B_{\mu\mu}$ ,  $B_{\tau\tau}$ , and  $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$ . The entries of the last quantity have been re-evaluated consistently using the correction procedure of KURA EV 85 [1]. The partial width  $\Gamma_{ee}$  is obtained from the average values for  $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$  and  $B_{\ell\ell}$  using

$$\Gamma_{ee} = \frac{\Gamma_{ee}\Gamma_{\text{had}}}{\Gamma(1 - 3B_{\ell\ell})} .\quad (5)$$

The total width  $\Gamma$  is then obtained from Eq. (1). We do not list  $\Gamma_{ee}$  and  $\Gamma$  values of individual experiments. The  $\Gamma_{ee}$  values in the Meson Summary Table are also those defined in Eq. (1).

## References

1. E.A. Kuraev, V.S. Fadin, Sov. J. Nucl. Phys. **41**, 466 (1985).

$\eta_b(1S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ .

NODE=M171

NODE=M171

NODE=M171M

NODE=M171M

$\eta_b(1S)$  MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9398.7 ± 2.0 OUR AVERAGE</b> Error includes scale factor of 1.5. See the ideogram below.				
9394.8 <sup>+2.7</sup> <sub>-3.1</sub> ± 4.5 <sub>-2.7</sub>	29k	FULSOM	18 BELL	$\Upsilon(2S) \rightarrow \gamma X$
9400.7 ± 1.7 ± 1.6	33.1k	TAMPONI	15 BELL	$e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
9402.4 ± 1.5 ± 1.8	34k	<sup>1</sup> MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
9391.8 ± 6.6 ± 2.0	2.3k	<sup>2</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
9394.2 <sup>+4.8</sup> <sub>-4.9</sub> ± 2.0	13k	<sup>2</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
9388.9 <sup>+3.1</sup> <sub>-2.3</sub> ± 2.7	19k	<sup>2</sup> AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9393.2 ± 3.4 ± 2.3	10	<sup>2,3</sup> DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$
9300 ± 20 ± 20		HEISTER	02D ALEP	181–209 $e^+e^-$

- <sup>1</sup> With floating width. Not independent of the corresponding mass difference measurement.
- <sup>2</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV. Not independent of the corresponding  $\gamma$  energy or mass difference measurements.
- <sup>3</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M;LINKAGE=MI

NODE=M171M;LINKAGE=AU

NODE=M171M;LINKAGE=DO



$m_{\Upsilon(1S)} - m_{\eta_b}$

NODE=M171M2


NODE=M171M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>62.3 ± 3.2 OUR AVERAGE</b> Error includes scale factor of 1.8. See the ideogram below.				
57.9 ± 1.5 ± 1.8	34k	<sup>1</sup> MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
68.5 ± 6.6 ± 2.0	2.3 ± 0.5k	<sup>2</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
66.1 <sup>+4.8</sup> <sub>-4.9</sub> ± 2.0	13 ± 5k	<sup>2</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
71.4 <sup>+2.3</sup> <sub>-3.1</sub> ± 2.7	19 ± 3k	<sup>2</sup> AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1 ± 3.4 ± 2.3	10 <sup>+5</sup> <sub>-4</sub>	<sup>2,3</sup> DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$



- <sup>1</sup>With floating width. Not independent of the corresponding mass measurement.
- <sup>2</sup>Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV. Not independent of the corresponding  $\gamma$  energy or mass measurements.
- <sup>3</sup>Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M2;LINKAGE=MI  
 NODE=M171M2;LINKAGE=AU  
 NODE=M171M2;LINKAGE=DO



**$\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY**

NODE=M171DM  
 NODE=M171DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>920.6<sup>+2.8</sup><sub>-3.2</sub> OUR AVERAGE</b>				
918.6 ± 6.0 ± 1.9	2.3 ± 0.5k	<sup>1</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
921.2 <sup>+2.1</sup> <sub>-2.8</sub> ± 2.4	19 ± 3k	<sup>1</sup> AUBERT	08v BABR	$\Upsilon(3S) \rightarrow \gamma X$

<sup>1</sup>Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171DM;LINKAGE=BO

**$\gamma$  ENERGY IN  $\Upsilon(2S)$  DECAY**

NODE=M171U2S  
 NODE=M171U2S

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>609.3<sup>+4.6</sup><sub>-4.5</sub> ± 1.9</b>				
	13 ± 5k	<sup>1</sup> AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$

<sup>1</sup>Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171U2S;LINKAGE=AU

**$\eta_b(1S)$  WIDTH**

NODE=M171W  
 NODE=M171W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10<sup>+5</sup><sub>-4</sub> OUR AVERAGE</b>				
8 <sup>+6</sup> <sub>-5</sub> ± 5	33.1k	<sup>1</sup> TAMPONI	15 BELL	$e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
10.8 <sup>+4.0</sup> <sub>-3.7</sub> <sup>+4.5</sup> <sub>-2.0</sub>	34k	<sup>1</sup> MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$

<sup>1</sup>With floating mass.

NODE=M171W;LINKAGE=MI

$\eta_b(1S)$  DECAY MODES

NODE=M171225;NODE=M171

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ hadrons	seen	
$\Gamma_2$ $3h^+3h^-$	not seen	
$\Gamma_3$ $2h^+2h^-$	not seen	
$\Gamma_4$ $4h^+4h^-$	not seen	
$\Gamma_5$ $\gamma\gamma$	not seen	
$\Gamma_6$ $\mu^+\mu^-$	$<9 \times 10^{-3}$	90%
$\Gamma_7$ $\tau^+\tau^-$	$<8\%$	90%

DESIG=7

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=5

DESIG=6

 $\eta_b(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$ 

NODE=M171230

 $\Gamma(3h^+3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_2\Gamma_5/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<470	95	ABDALLAH	06	DLPH	161-209 $e^+e^-$
------	----	----------	----	------	------------------

<132	95	HEISTER	02D	ALEP	181-209 $e^+e^-$
------	----	---------	-----	------	------------------

NODE=M171G1

NODE=M171G1

 $\Gamma(2h^+2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_3\Gamma_5/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<190	95	ABDALLAH	06	DLPH	161-209 $e^+e^-$
------	----	----------	----	------	------------------

< 48	95	HEISTER	02D	ALEP	181-209 $e^+e^-$
------	----	---------	-----	------	------------------

NODE=M171G2

NODE=M171G2

 $\Gamma(4h^+4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$   $\Gamma_4\Gamma_5/\Gamma$ 

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<660	95	ABDALLAH	06	DLPH	161-209 $e^+e^-$
------	----	----------	----	------	------------------

NODE=M171G3

NODE=M171G3

 $\eta_b(1S)$  BRANCHING RATIOS

NODE=M171235

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	34k	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
------	-----	-------	----	---

NODE=M171R03

NODE=M171R03

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<9 \times 10^{-3}$	90	<sup>1</sup> AUBERT	09Z	BABR $e^+e^- \rightarrow \Upsilon(2S, 3S) \rightarrow \gamma\eta_b$
---------------------	----	---------------------	-----	---

<sup>1</sup> Obtained using  $B(\Upsilon(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$  and  $B(\Upsilon(3S) \rightarrow \gamma\eta_b)$

$= (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$ . This limit is equivalent to  $B(\eta_b \rightarrow \mu^+\mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$  measurement.

NODE=M171R01

NODE=M171R01

NODE=M171R01;LINKAGE=AU

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<8 \times 10^{-2}$	90	AUBERT	09P	BABR $e^+e^- \rightarrow \gamma\tau^+\tau^-$
---------------------	----	--------	-----	--

NODE=M171R02

NODE=M171R02

 $\eta_b(1S)$  REFERENCES

NODE=M171

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)

REFID=59535

REFID=56996

REFID=54288

REFID=54718

REFID=53231

REFID=53106

REFID=53062

REFID=52930

REFID=52262

REFID=51042

REFID=48577

$\Upsilon(1S)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M049

 **$\Upsilon(1S)$  MASS**

NODE=M049M

VALUE (MeV) DOCUMENT ID TECN COMMENT  
**9460.40 ± 0.10 OUR AVERAGE** [9460.30 ± 0.26 MeV OUR 2022 AVERAGE Scale factor = 3.3]

NODE=M049M  
NEW

**9460.40 ± 0.09 ± 0.04** <sup>1</sup> SHAMOV 23 RVUE  $e^+e^- \rightarrow$  hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

9460.11 ± 0.11 ± 0.07 <sup>2</sup> SHAMOV 23 RVUE  $e^+e^- \rightarrow$  hadrons

OCCUR=2

9460.51 ± 0.09 ± 0.05 <sup>3,4</sup> ARTAMONOV 00 MD1  $e^+e^- \rightarrow$  hadrons

9460.60 ± 0.09 ± 0.05 <sup>5,6</sup> BARU 92B MD1  $e^+e^- \rightarrow$  hadrons

9460.59 ± 0.12 BARU 86 MD1  $e^+e^- \rightarrow$  hadrons

9460.6 ± 0.4 <sup>6,7</sup> ARTAMONOV 84 MD1  $e^+e^- \rightarrow$  hadrons

9459.97 ± 0.11 ± 0.07 <sup>8</sup> MACKAY 84 CUSB  $e^+e^- \rightarrow$  hadrons

<sup>1</sup> Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.

NODE=M049M;LINKAGE=D

<sup>2</sup> Obtained by reanalysing CUSB data (MACKAY 84), but not authored by the CUSB collaboration.

NODE=M049M;LINKAGE=E

<sup>3</sup> Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).

NODE=M049M;LINKAGE=AR

<sup>4</sup> Superseded by SHAMOV 23.

NODE=M049M;LINKAGE=B

<sup>5</sup> Supersedes BARU 86.

NODE=M049M;LINKAGE=A

<sup>6</sup> Superseded by ARTAMONOV 00.

NODE=M049M;LINKAGE=RZ

<sup>7</sup> Value includes data of ARTAMONOV 82.

NODE=M049M;LINKAGE=G

<sup>8</sup> Reanalysed by SHAMOV 23.

NODE=M049M;LINKAGE=C

 **$\Upsilon(1S)$  WIDTH**

NODE=M049W

VALUE (keV) DOCUMENT ID  
**54.02 ± 1.25 OUR EVALUATION** See the Note on "Width Determinations of the  $\Upsilon$  States"

NODE=M049W  
→ UNCHECKED ← **$\Upsilon(1S)$  DECAY MODES**

NODE=M049215;NODE=M049

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\tau^+ \tau^-$	( 2.60 ± 0.10 ) %	
$\Gamma_2$ $e^+ e^-$	( 2.39 ± 0.08 ) %	
$\Gamma_3$ $\mu^+ \mu^-$	( 2.48 ± 0.04 ) %	

DESIG=3

DESIG=2

DESIG=1

**Hadronic decays**

NODE=M049;CLUMP=A

$\Gamma_4$ $g g g$	(81.7 ± 0.7 ) %	
$\Gamma_5$ $\gamma g g$	( 2.2 ± 0.6 ) %	
$\Gamma_6$ $\eta'(958)$ anything	( 2.94 ± 0.24 ) %	
$\Gamma_7$ $J/\psi(1S)$ anything	( 5.4 ± 0.4 ) × 10 <sup>-4</sup>	S=1.4
$\Gamma_8$ $J/\psi(1S)\eta_c$	< 2.2	× 10 <sup>-6</sup> CL=90%
$\Gamma_9$ $J/\psi(1S)\chi_{c0}$	< 3.4	× 10 <sup>-6</sup> CL=90%
$\Gamma_{10}$ $J/\psi(1S)\chi_{c1}$	( 3.9 ± 1.2 ) × 10 <sup>-6</sup>	
$\Gamma_{11}$ $J/\psi(1S)\chi_{c2}$	< 1.4	× 10 <sup>-6</sup> CL=90%
$\Gamma_{12}$ $J/\psi(1S)\eta_c(2S)$	< 2.2	× 10 <sup>-6</sup> CL=90%
$\Gamma_{13}$ $J/\psi(1S)X(3940)$	< 5.4	× 10 <sup>-6</sup> CL=90%
$\Gamma_{14}$ $J/\psi(1S)X(4160)$	< 5.4	× 10 <sup>-6</sup> CL=90%
$\Gamma_{15}$ $X(4350)$ anything, $X \rightarrow J/\psi(1S)\phi$	< 8.1	× 10 <sup>-6</sup> CL=90%
$\Gamma_{16}$ $Z_c(3900)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 1.3	× 10 <sup>-5</sup> CL=90%
$\Gamma_{17}$ $Z_c(4200)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 6.0	× 10 <sup>-5</sup> CL=90%
$\Gamma_{18}$ $Z_c(4430)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 4.9	× 10 <sup>-5</sup> CL=90%
$\Gamma_{19}$ $X_{cs}^\pm$ anything, $X \rightarrow J/\psi K^\pm$	< 5.7	× 10 <sup>-6</sup> CL=90%
$\Gamma_{20}$ $\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)\pi^+\pi^-$	< 3.8	× 10 <sup>-5</sup> CL=90%

DESIG=117

DESIG=118

DESIG=73

DESIG=12

DESIG=146

DESIG=147

DESIG=148

DESIG=149

DESIG=150

DESIG=151

DESIG=152

DESIG=167

DESIG=168

DESIG=169

DESIG=170

DESIG=173

DESIG=161

Γ <sub>21</sub>	$\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)K^+K^-$	< 7.5	$\times 10^{-6}$	CL=90%	DESIG=165
Γ <sub>22</sub>	$\chi_{c1}(4140)$ anything, $\chi_{c1} \rightarrow J/\psi(1S)\phi$	< 5.2	$\times 10^{-6}$	CL=90%	DESIG=166
Γ <sub>23</sub>	$\chi_{c0}$ anything	< 4	$\times 10^{-3}$	CL=90%	DESIG=5
Γ <sub>24</sub>	$\chi_{c1}$ anything	( 1.90 ± 0.35 )	$\times 10^{-4}$		DESIG=6
Γ <sub>25</sub>	$\chi_{c1}(1P)X_{tetra}$	< 3.78	$\times 10^{-5}$	CL=90%	DESIG=175
Γ <sub>26</sub>	$\chi_{c2}$ anything	( 2.8 ± 0.8 )	$\times 10^{-4}$		DESIG=7
Γ <sub>27</sub>	$\psi(2S)$ anything	( 1.23 ± 0.20 )	$\times 10^{-4}$		DESIG=8
Γ <sub>28</sub>	$\psi(2S)\eta_c$	< 3.6	$\times 10^{-6}$	CL=90%	DESIG=153
Γ <sub>29</sub>	$\psi(2S)\chi_{c0}$	< 6.5	$\times 10^{-6}$	CL=90%	DESIG=154
Γ <sub>30</sub>	$\psi(2S)\chi_{c1}$	< 4.5	$\times 10^{-6}$	CL=90%	DESIG=155
Γ <sub>31</sub>	$\psi(2S)\chi_{c2}$	< 2.1	$\times 10^{-6}$	CL=90%	DESIG=156
Γ <sub>32</sub>	$\psi(2S)\eta_c(2S)$	< 3.2	$\times 10^{-6}$	CL=90%	DESIG=157
Γ <sub>33</sub>	$\psi(2S)X(3940)$	< 2.9	$\times 10^{-6}$	CL=90%	DESIG=158
Γ <sub>34</sub>	$\psi(2S)X(4160)$	< 2.9	$\times 10^{-6}$	CL=90%	DESIG=159
Γ <sub>35</sub>	$\psi(4230)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 7.9	$\times 10^{-5}$	CL=90%	DESIG=162
Γ <sub>36</sub>	$\psi(4360)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 5.2	$\times 10^{-5}$	CL=90%	DESIG=163
Γ <sub>37</sub>	$\psi(4660)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 2.2	$\times 10^{-5}$	CL=90%	DESIG=164
Γ <sub>38</sub>	$X(4050)^\pm$ anything, $X \rightarrow \psi(2S)\pi^\pm$	< 8.8	$\times 10^{-5}$	CL=90%	DESIG=171
Γ <sub>39</sub>	$Z_c(4430)^\pm$ anything, $Z_c \rightarrow \psi(2S)\pi^\pm$	< 6.7	$\times 10^{-5}$	CL=90%	DESIG=172
Γ <sub>40</sub>	$\chi_{c1}(3872)$ anything	< 2.5	$\times 10^{-4}$	CL=90%	DESIG=194
Γ <sub>41</sub>	$Z_c(4200)^+Z_c(4200)^-$	< 2.23	$\times 10^{-5}$	CL=90%	DESIG=178
Γ <sub>42</sub>	$Z_c(3900)^\pm Z_c(4200)^\mp$	< 8.1	$\times 10^{-6}$	CL=90%	DESIG=179
Γ <sub>43</sub>	$Z_c(3900)^+Z_c(3900)^-$	< 1.8	$\times 10^{-6}$	CL=90%	DESIG=180
Γ <sub>44</sub>	$X(4050)^+X(4050)^-$	< 1.58	$\times 10^{-5}$	CL=90%	DESIG=181
Γ <sub>45</sub>	$X(4250)^+X(4250)^-$	< 2.66	$\times 10^{-5}$	CL=90%	DESIG=182
Γ <sub>46</sub>	$X(4050)^\pm X(4250)^\mp$	< 4.42	$\times 10^{-5}$	CL=90%	DESIG=183
Γ <sub>47</sub>	$Z_c(4430)^+Z_c(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%	DESIG=184
Γ <sub>48</sub>	$X(4055)^\pm X(4055)^\mp$	< 2.33	$\times 10^{-5}$	CL=90%	DESIG=186
Γ <sub>49</sub>	$X(4055)^\pm Z_c(4430)^\mp$	< 4.55	$\times 10^{-5}$	CL=90%	DESIG=189
Γ <sub>50</sub>	$\rho\pi$	< 3.68	$\times 10^{-6}$	CL=90%	DESIG=11
Γ <sub>51</sub>	$\omega\pi^0$	< 3.90	$\times 10^{-6}$	CL=90%	DESIG=131
Γ <sub>52</sub>	$\pi^+\pi^-$	< 5	$\times 10^{-4}$	CL=90%	DESIG=23
Γ <sub>53</sub>	$K^+K^-$	< 5	$\times 10^{-4}$	CL=90%	DESIG=24
Γ <sub>54</sub>	$\rho\bar{\rho}$	< 5	$\times 10^{-4}$	CL=90%	DESIG=25
Γ <sub>55</sub>	$\pi^+\pi^-\pi^0$	( 2.1 ± 0.8 )	$\times 10^{-6}$		DESIG=72
Γ <sub>56</sub>	$\phi K^+K^-$	( 2.4 ± 0.5 )	$\times 10^{-6}$		DESIG=136
Γ <sub>57</sub>	$\omega\pi^+\pi^-$	( 4.5 ± 1.0 )	$\times 10^{-6}$		DESIG=137
Γ <sub>58</sub>	$K^*(892)^0 K^- \pi^+ + c.c.$	( 4.4 ± 0.8 )	$\times 10^{-6}$		DESIG=138
Γ <sub>59</sub>	$\phi f'_2(1525)$	< 1.63	$\times 10^{-6}$	CL=90%	DESIG=139
Γ <sub>60</sub>	$\omega f_2(1270)$	< 1.79	$\times 10^{-6}$	CL=90%	DESIG=140
Γ <sub>61</sub>	$\rho(770)a_2(1320)$	< 2.24	$\times 10^{-6}$	CL=90%	DESIG=141
Γ <sub>62</sub>	$K^*(892)^0 \bar{K}_2^*(1430)^0 + c.c.$	( 3.0 ± 0.8 )	$\times 10^{-6}$		DESIG=142
Γ <sub>63</sub>	$K_1(1270)^\pm K^\mp$	< 2.41	$\times 10^{-6}$	CL=90%	DESIG=143
Γ <sub>64</sub>	$K_1(1400)^\pm K^\mp$	( 1.0 ± 0.4 )	$\times 10^{-6}$		DESIG=144
Γ <sub>65</sub>	$b_1(1235)^\pm \pi^\mp$	< 1.25	$\times 10^{-6}$	CL=90%	DESIG=145
Γ <sub>66</sub>	$\pi^+\pi^-\pi^0\pi^0$	( 1.28 ± 0.30 )	$\times 10^{-5}$		DESIG=132
Γ <sub>67</sub>	$K_S^0 K^+ \pi^- + c.c.$	( 1.6 ± 0.4 )	$\times 10^{-6}$		DESIG=133
Γ <sub>68</sub>	$K^*(892)^0 \bar{K}^0 + c.c.$	( 2.9 ± 0.9 )	$\times 10^{-6}$		DESIG=134
Γ <sub>69</sub>	$K^*(892)^- K^+ + c.c.$	< 1.11	$\times 10^{-6}$	CL=90%	DESIG=135
Γ <sub>70</sub>	$f_1(1285)$ anything	( 4.6 ± 3.1 )	$\times 10^{-3}$		DESIG=174
Γ <sub>71</sub>	$D^*(2010)^\pm$ anything	( 2.52 ± 0.20 )	%		DESIG=30
Γ <sub>72</sub>	$f_1(1285)X_{tetra}$	< 6.24	$\times 10^{-5}$	CL=90%	DESIG=176
Γ <sub>73</sub>	${}^2H$ anything	( 2.85 ± 0.25 )	$\times 10^{-5}$		DESIG=107
Γ <sub>74</sub>	Sum of 100 exclusive modes	( 1.200 ± 0.017 )	%		DESIG=128

## Radiative decays

				NODE=M049;CLUMP=B
Γ <sub>75</sub>	$\gamma\pi^+\pi^-$	( 6.3 ±1.8 )	$\times 10^{-5}$	DESIG=70
Γ <sub>76</sub>	$\gamma\pi^0\pi^0$	( 1.7 ±0.7 )	$\times 10^{-5}$	DESIG=71
Γ <sub>77</sub>	$\gamma\pi\pi$ (S-wave)	( 4.6 ±0.7 )	$\times 10^{-5}$	DESIG=190
Γ <sub>78</sub>	$\gamma\pi^0\eta$	< 2.4	$\times 10^{-6}$	CL=90% DESIG=111
Γ <sub>79</sub>	$\gamma K^+ K^-$	[a] ( 1.14 ±0.13 )	$\times 10^{-5}$	DESIG=102
Γ <sub>80</sub>	$\gamma p\bar{p}$	[b] < 6	$\times 10^{-6}$	CL=90% DESIG=103
Γ <sub>81</sub>	$\gamma 2h^+ 2h^-$	( 7.0 ±1.5 )	$\times 10^{-4}$	DESIG=20
Γ <sub>82</sub>	$\gamma 3h^+ 3h^-$	( 5.4 ±2.0 )	$\times 10^{-4}$	DESIG=21
Γ <sub>83</sub>	$\gamma 4h^+ 4h^-$	( 7.4 ±3.5 )	$\times 10^{-4}$	DESIG=22
Γ <sub>84</sub>	$\gamma\pi^+\pi^- K^+ K^-$	( 2.9 ±0.9 )	$\times 10^{-4}$	DESIG=14
Γ <sub>85</sub>	$\gamma 2\pi^+ 2\pi^-$	( 2.5 ±0.9 )	$\times 10^{-4}$	DESIG=13
Γ <sub>86</sub>	$\gamma 3\pi^+ 3\pi^-$	( 2.5 ±1.2 )	$\times 10^{-4}$	DESIG=17
Γ <sub>87</sub>	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	( 2.4 ±1.2 )	$\times 10^{-4}$	DESIG=18
Γ <sub>88</sub>	$\gamma\pi^+\pi^- p\bar{p}$	( 1.5 ±0.6 )	$\times 10^{-4}$	DESIG=15
Γ <sub>89</sub>	$\gamma 2\pi^+ 2\pi^- p\bar{p}$	( 4 ±6 )	$\times 10^{-5}$	DESIG=19
Γ <sub>90</sub>	$\gamma 2K^+ 2K^-$	( 2.0 ±2.0 )	$\times 10^{-5}$	DESIG=16
Γ <sub>91</sub>	$\gamma\eta'(958)$	< 1.9	$\times 10^{-6}$	CL=90% DESIG=55
Γ <sub>92</sub>	$\gamma\eta$	< 1.0	$\times 10^{-6}$	CL=90% DESIG=54
Γ <sub>93</sub>	$\gamma f_0(980)$	< 3	$\times 10^{-5}$	CL=90% DESIG=105
Γ <sub>94</sub>	$\gamma f_2'(1525)$	( 2.9 ±0.6 )	$\times 10^{-5}$	DESIG=52
Γ <sub>95</sub>	$\gamma f_2(1270)$	( 1.01 ±0.06 )	$\times 10^{-4}$	DESIG=51
Γ <sub>96</sub>	$\gamma\eta(1405)$	< 8.2	$\times 10^{-5}$	CL=90% DESIG=65
Γ <sub>97</sub>	$\gamma f_0(1500)$	< 1.5	$\times 10^{-5}$	CL=90% DESIG=108
Γ <sub>98</sub>	$\gamma f_0(1500) \rightarrow \gamma K^+ K^-$	( 1.0 ±0.4 )	$\times 10^{-5}$	DESIG=192
Γ <sub>99</sub>	$\gamma f_0(1710)$	< 2.6	$\times 10^{-4}$	CL=90% DESIG=53
Γ <sub>100</sub>	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	( 1.01 ±0.32 )	$\times 10^{-5}$	DESIG=112
Γ <sub>101</sub>	$\gamma f_0(1710) \rightarrow \gamma\pi^+\pi^-$	( 5.3 ±2.0 )	$\times 10^{-6}$	DESIG=191
Γ <sub>102</sub>	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$	< 1.4	$\times 10^{-6}$	CL=90% DESIG=109
Γ <sub>103</sub>	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	< 1.8	$\times 10^{-6}$	CL=90% DESIG=110
Γ <sub>104</sub>	$\gamma f_4(2050)$	< 5.3	$\times 10^{-5}$	CL=90% DESIG=104
Γ <sub>105</sub>	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	< 2	$\times 10^{-4}$	CL=90% DESIG=69
Γ <sub>106</sub>	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	< 8	$\times 10^{-7}$	CL=90% DESIG=60
Γ <sub>107</sub>	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	< 6	$\times 10^{-7}$	CL=90% DESIG=61
Γ <sub>108</sub>	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	< 1.1	$\times 10^{-6}$	CL=90% DESIG=62
Γ <sub>109</sub>	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	< 3	$\times 10^{-3}$	CL=90% DESIG=68
Γ <sub>110</sub>	$\gamma\eta_c(1S)$	< 2.9	$\times 10^{-5}$	CL=90% DESIG=119
Γ <sub>111</sub>	$\gamma\eta_c(2S)$	< 4	$\times 10^{-4}$	CL=90% DESIG=193
Γ <sub>112</sub>	$\gamma\chi_{c0}$	< 6.6	$\times 10^{-5}$	CL=90% DESIG=120
Γ <sub>113</sub>	$\gamma\chi_{c1}$	( 4.7 <sup>+2.4</sup> <sub>-1.9</sub> )	$\times 10^{-5}$	DESIG=121
Γ <sub>114</sub>	$\gamma\chi_{c2}$	< 7.6	$\times 10^{-6}$	CL=90% DESIG=122
Γ <sub>115</sub>	$\gamma\chi_{c1}(3872)$	< 4	$\times 10^{-5}$	CL=90% DESIG=195
Γ <sub>116</sub>	$\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi$	< 2.8	$\times 10^{-6}$	CL=90% DESIG=124
Γ <sub>117</sub>	$\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi$	< 3.0	$\times 10^{-6}$	CL=90% DESIG=125
Γ <sub>118</sub>	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	< 2.2	$\times 10^{-6}$	CL=90% DESIG=126
Γ <sub>119</sub>	$\gamma X\bar{X} (m_X < 3.1 \text{ GeV})$	[c] < 1	$\times 10^{-3}$	CL=90% DESIG=67
Γ <sub>120</sub>	$\gamma X\bar{X} (m_X < 4.5 \text{ GeV})$	[d] < 2.4	$\times 10^{-4}$	CL=90% DESIG=127
Γ <sub>121</sub>	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[e] < 1.78	$\times 10^{-4}$	CL=95% DESIG=113
Γ <sub>122</sub>	$\gamma A^0$	[f]		DESIG=66
Γ <sub>123</sub>	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	[g] < 9	$\times 10^{-6}$	CL=90% DESIG=114
Γ <sub>124</sub>	$\gamma A^0 \rightarrow \gamma\tau^+\tau^-$	[a] < 1.30	$\times 10^{-4}$	CL=90% DESIG=115
Γ <sub>125</sub>	$\gamma A^0 \rightarrow \gamma g g$	[h] < 1	%	CL=90% DESIG=129
Γ <sub>126</sub>	$\gamma A^0 \rightarrow \gamma s\bar{s}$	[h] < 1	$\times 10^{-3}$	CL=90% DESIG=130

## Lepton Family number (LF) violating modes

				NODE=M049;CLUMP=C
Γ <sub>127</sub>	$e^\pm \mu^\mp$	LF	< 3.9	$\times 10^{-7}$ CL=90% DESIG=196
Γ <sub>128</sub>	$\mu^\pm \tau^\mp$	LF	< 2.7	$\times 10^{-6}$ CL=90% DESIG=116
Γ <sub>129</sub>	$e^\pm \tau^\mp$	LF	< 2.7	$\times 10^{-6}$ CL=90% DESIG=197
Γ <sub>130</sub>	$\gamma e^\pm \mu^\mp$	LF	< 4.2	$\times 10^{-7}$ CL=90% DESIG=198
Γ <sub>131</sub>	$\gamma \mu^\pm \tau^\mp$	LF	< 6.1	$\times 10^{-6}$ CL=90% DESIG=199
Γ <sub>132</sub>	$\gamma e^\pm \tau^\mp$	LF	< 6.5	$\times 10^{-6}$ CL=90% DESIG=200

**Other decays**

$\Gamma_{133}$  invisible  $< 3.0 \times 10^{-4}$  CL=90%  
 $\Gamma_{134}$  hadrons  $(96 \pm 4) \%$

NODE=M049;CLUMP=D

DESIG=106

DESIG=101

[a]  $2m_\tau < M(\tau^+\tau^-) < 9.2$  GeV

LINKAGE=E49

[b]  $2$  GeV  $< m_{K^+K^-} < 3$  GeV

LINKAGE=G49

[c]  $X\bar{X}$  = vectors with  $m < 3.1$  GeV

LINKAGE=B49

[d]  $X$  and  $\bar{X}$  = zero spin with  $m < 4.5$  GeV

LINKAGE=F49

[e]  $1.5$  GeV  $< m_X < 5.0$  GeV

LINKAGE=C49

[f]  $A^0$  = scalar with  $m < 8.0$  GeV

LINKAGE=A49

[g]  $201$  MeV  $< M(\mu^+\mu^-) < 3565$  MeV

LINKAGE=D49

[h]  $0.5$  GeV  $< m_X < 9.0$  GeV, where  $m_X$  is the invariant mass of the hadronic final state.

LINKAGE=I49

 **$\Upsilon(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

NODE=M049218

 **$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**  **$\Gamma_3\Gamma_2/\Gamma$** 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>31.2±1.6±1.7</b>	KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$

NODE=M049G1

NODE=M049G1

 **$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**  **$\Gamma_{134}\Gamma_2/\Gamma$** 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>1.240±0.016 OUR AVERAGE</b>			
1.252±0.004±0.019	<sup>1</sup> ROSNER	06	CLEO $9.5 e^+e^- \rightarrow \text{hadrons}$
1.187±0.023±0.031	<sup>1</sup> BARU	92B	MD1 $e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	<sup>1</sup> JAKUBOWSKI	88	CBAL $e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	<sup>2</sup> GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	<sup>2</sup> ALBRECHT	82	DASP $e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	<sup>2</sup> NICZYPORUK	82	LENA $e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	<sup>2</sup> BOCK	80	CNTR $e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	<sup>3</sup> BERGER	79	PLUT $e^+e^- \rightarrow \text{hadrons}$

NODE=M049G2

NODE=M049G2

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.<sup>3</sup> Radiative corrections reevaluated by ALEXANDER 89 using  $B(\mu\mu) = 0.026$ .

NODE=M049G2;LINKAGE=B

NODE=M049G2;LINKAGE=R

NODE=M049G2;LINKAGE=P

 **$\Upsilon(1S)$  PARTIAL WIDTHS**

NODE=M049220

 **$\Gamma(e^+e^-)$**  **$\Gamma_2$** 

VALUE (keV)	DOCUMENT ID
<b>1.340±0.018 OUR EVALUATION</b>	

NODE=M049W2

NODE=M049W2

→ UNCHECKED ←

 **$\Upsilon(1S)$  BRANCHING RATIOS**

NODE=M049225

 **$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$**  **$\Gamma_1/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.60±0.10 OUR AVERAGE</b>				
2.53±0.13±0.04	60k	<sup>1</sup> BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+\tau^-$
2.61±0.12 <sup>+0.09</sup> <sub>-0.13</sub>	25k	CINABRO	94B	CLE2 $e^+e^- \rightarrow \tau^+\tau^-$
2.7 ±0.4 ±0.2		<sup>2</sup> ALBRECHT	85C	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$
3.4 ±0.4 ±0.4		GILES	83	CLEO $e^+e^- \rightarrow \tau^+\tau^-$

NODE=M049R3

NODE=M049R3

<sup>1</sup> BESSON 07 reports  $[\Gamma(\Upsilon(1S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = 1.02 \pm 0.02 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$ ; not used for width evaluations.

NODE=M049R3;LINKAGE=BE

NODE=M049R3;LINKAGE=A

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.39±0.08 OUR AVERAGE</b>				
[(2.38 ± 0.11) × 10 <sup>-2</sup> OUR 2022 AVERAGE]				
2.40±0.01±0.12	191k	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.29±0.08±0.11		ALEXANDER	98	CLE2 $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.42±0.14±0.14	307	ALBRECHT	87	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.8 ±0.3 ±0.2	826	BESSON	84	CLEO $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
5.1 ±3.0		BERGER	80c	PLUT $e^+e^- \rightarrow e^+e^-$

NODE=M049R2  
 NODE=M049R2  
 NEW

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.48±0.04 OUR AVERAGE</b>				
[0.0248 ± 0.0005 OUR 2022 AVERAGE]				
2.46±0.01±0.11	246k	PATRA	22	BELL $\Upsilon(2S) \rightarrow$
				$\pi^+\pi^-\mu^+\mu^-$
2.49±0.02±0.07	345k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
2.49±0.08±0.13		ALEXANDER	98	CLE2 $\Upsilon(2S) \rightarrow$
				$\pi^+\pi^-\mu^+\mu^-$
2.12±0.20±0.10		<sup>1</sup> BARU	92	MD1 $e^+e^- \rightarrow \mu^+\mu^-$
2.31±0.12±0.10		<sup>1</sup> KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$
2.52±0.07±0.07		CHEN	89B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
2.61±0.09±0.11		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
2.30±0.25±0.13	86	ALBRECHT	87	ARG $\Upsilon(2S) \rightarrow$
				$\pi^+\pi^-\mu^+\mu^-$
2.9 ±0.3 ±0.2	864	BESSON	84	CLEO $\Upsilon(2S) \rightarrow$
				$\pi^+\pi^-\mu^+\mu^-$
2.7 ±0.3 ±0.3		ANDREWS	83	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
3.2 ±1.3 ±0.3		ALBRECHT	82	DASP $e^+e^- \rightarrow \mu^+\mu^-$
3.8 ±1.5 ±0.2		NICZYPORUK	82	LENA $e^+e^- \rightarrow \mu^+\mu^-$
1.4 +3.4 -1.4		BOCK	80	CNTR $e^+e^- \rightarrow \mu^+\mu^-$
2.2 ±2.0		BERGER	79	PLUT $e^+e^- \rightarrow \mu^+\mu^-$

NODE=M049R1  
 NODE=M049R1  
 NEW

<sup>1</sup> Taking into account interference between the resonance and continuum.

NODE=M049R1;LINKAGE=G

 $\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$  $\Gamma_1/\Gamma_3$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.008±0.023 OUR AVERAGE</b>				
1.005±0.013±0.022	0.7M	<sup>1</sup> DEL-AMO-SA..10c	BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\Upsilon(1S)$
1.02 ±0.02 ±0.05	60k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S)$

NODE=M049R43  
 NODE=M049R43

<sup>1</sup> Allows any number of extra photons with total energy < 500 MeV.

NODE=M049R43;LINKAGE=DE

 $\Gamma(ggg)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>81.7±0.7</b>	20M	<sup>1</sup> BESSON	06A	CLEO $\Upsilon(1S) \rightarrow$ hadrons

NODE=M049R35  
 NODE=M049R35

<sup>1</sup> Calculated using the value  $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$  from BESSON 06A and PDG 08 values of  $B(\mu^+\mu^-) = (2.48 \pm 0.05)\%$  and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma gg)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

NODE=M049R35;LINKAGE=BE

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.20±0.60</b>	400k	<sup>1</sup> BESSON	06A	CLEO $\Upsilon(1S) \rightarrow \gamma +$ hadrons

NODE=M049R36  
 NODE=M049R36

<sup>1</sup> Calculated using BESSON 06A values of  $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$  and  $\Gamma(ggg)/\Gamma_{\text{total}}$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(ggg)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

NODE=M049R36;LINKAGE=BE

 $\Gamma(\gamma gg)/\Gamma(ggg)$  $\Gamma_5/\Gamma_4$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.70±0.01±0.27</b>	20M	BESSON	06A	CLEO $\Upsilon(1S) \rightarrow (\gamma +)$ hadrons

NODE=M049R37  
 NODE=M049R37

 $\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0294±0.0024 OUR AVERAGE</b>			
0.030 ±0.002 ±0.002	AQUINES	06A	CLE3 $\Upsilon(1S) \rightarrow \eta'$ anything
0.028 ±0.004 ±0.002	ARTUSO	03	CLE2 $\Upsilon(1S) \rightarrow \eta'$ anything

NODE=M049R73  
 NODE=M049R73

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>5.4 ± 0.4 OUR FIT</b>					Error includes scale factor of 1.4.
<b>5.4 ± 0.4 OUR AVERAGE</b>					Error includes scale factor of 1.5.
5.25 ± 0.13 ± 0.25		3k	SHEN	16	BELL $e^+e^- \rightarrow J/\psi X$
6.4 ± 0.4 ± 0.6		730	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$
11 ± 4 ± 2			<sup>1</sup> FULTON	89	CLEO $e^+e^- \rightarrow \mu^+\mu^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.8	90		ALBRECHT	92J	ARG $e^+e^- \rightarrow e^+e^- X$ , $\mu^+\mu^- X$
<17	90		MASCHMANN	90	CBAL $e^+e^- \rightarrow \text{hadrons}$
<200	90		NICZYPORUK	83	LENA
<sup>1</sup> Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$ .					

NODE=M049R12  
NODE=M049R12

NODE=M049R12;LINKAGE=K

 $\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10 <sup>-6</sup>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R85  
NODE=M049R85 $\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 × 10 <sup>-6</sup>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R86  
NODE=M049R86 $\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>3.90 ± 1.21 ± 0.23</b>	20	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R87  
NODE=M049R87 $\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10 <sup>-6</sup>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R88  
NODE=M049R88 $\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10 <sup>-6</sup>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R89  
NODE=M049R89 $\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10 <sup>-6</sup>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R90  
NODE=M049R90 $\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10 <sup>-6</sup>	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R91  
NODE=M049R91 $\Gamma(X(4350) \text{ anything}, X \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.1 × 10 <sup>-6</sup>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

NODE=M049P05  
NODE=M049P05 $\Gamma(Z_c(3900)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10 <sup>-5</sup>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

NODE=M049P06  
NODE=M049P06 $\Gamma(Z_c(4200)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.0 × 10 <sup>-5</sup>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

NODE=M049P07  
NODE=M049P07 $\Gamma(Z_c(4430)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.9 × 10 <sup>-5</sup>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

NODE=M049P08  
NODE=M049P08 $\Gamma(X_{cs}^\pm \text{ anything}, X \rightarrow J/\psi K^\pm)/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.7 × 10 <sup>-6</sup>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^- X$

NODE=M049P11  
NODE=M049P11 $\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.8 × 10 <sup>-5</sup>	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^+ \pi^- X$

NODE=M049R99  
NODE=M049R99



<b><math>\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S) K^+ K^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{21}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P03 NODE=M049P03
$<7.5 \times 10^{-6}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$		
<b><math>\Gamma(\chi_{c1}(4140) \text{ anything}, \chi_{c1} \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{22}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P04 NODE=M049P04
$<5.2 \times 10^{-6}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$		
<b><math>\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})</math></b>					<b><math>\Gamma_{23}/\Gamma_7</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R25 NODE=M049R25
$<7.4$	90	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$		
<b><math>\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{24}/\Gamma</math></b>	
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P13 NODE=M049P13
<b>1.90±0.35 OUR FIT</b>						
<b>1.90±0.43±0.14</b>	215	JIA	17	BELL $\Upsilon(1S) \rightarrow \gamma J/\psi(1S)$		
<b><math>\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})</math></b>					<b><math>\Gamma_{24}/\Gamma_7</math></b>	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R26 NODE=M049R26
<b>0.35±0.07 OUR FIT</b>						
<b>0.35±0.08±0.06</b>	52 ± 12	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$		
<b><math>\Gamma(\chi_{c1}(1P)\chi_{tetra})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{25}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P15 NODE=M049P15
$<37.8 \times 10^{-6}$	90	<sup>1</sup> JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$		
<sup>1</sup> For a tetraquark state $\chi_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of $\chi_{tetra}$ mass and width range from $4.4 \times 10^{-6}$ to $37.8 \times 10^{-6}$ .						NODE=M049P15;LINKAGE=A
<b><math>\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})</math></b>					<b><math>\Gamma_{26}/\Gamma_7</math></b>	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R27 NODE=M049R27
<b>0.52±0.12±0.09</b>	47 ± 11	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$		
<b><math>\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{27}/\Gamma</math></b>	
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049P12 NODE=M049P12
<b>1.23±0.17±0.11</b>	215	SHEN	16	BELL $e^+ e^- \rightarrow \psi(2S) X$		
<b><math>\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})</math></b>					<b><math>\Gamma_{27}/\Gamma_7</math></b>	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R28 NODE=M049R28
<b>0.41±0.11±0.08</b>	42 ± 11	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- X$		
<b><math>\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{28}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R92 NODE=M049R92
$<3.6 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
<b><math>\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{29}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R93 NODE=M049R93
$<6.5 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
<b><math>\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{30}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R94 NODE=M049R94
$<4.5 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
<b><math>\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{31}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R95 NODE=M049R95
$<2.1 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
<b><math>\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{32}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R96 NODE=M049R96
$<3.2 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		
<b><math>\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{33}/\Gamma</math></b>	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M049R97 NODE=M049R97
$<2.9 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$		

<b><math>\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{34}/\Gamma</math></b>	NODE=M049R98 NODE=M049R98
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.9 \times 10^{-6}$	90	YANG	14	BELL $e^+e^- \rightarrow \psi(2S)X$		
<b><math>\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{35}/\Gamma</math></b>	NODE=M049P00 NODE=M049P00
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<7.9 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$		
<b><math>\Gamma(\psi(4360) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{36}/\Gamma</math></b>	NODE=M049P01 NODE=M049P01
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<5.2 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$		
<b><math>\Gamma(\psi(4660) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{37}/\Gamma</math></b>	NODE=M049P02 NODE=M049P02
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.2 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^-X$		
<b><math>\Gamma(X(4050)^\pm \text{ anything}, X \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{38}/\Gamma</math></b>	NODE=M049P09 NODE=M049P09
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<8.8 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$		
<b><math>\Gamma(Z_c(4430)^\pm \text{ anything}, Z_c \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{39}/\Gamma</math></b>	NODE=M049P10 NODE=M049P10
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<6.7 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$		
<b><math>\Gamma(\chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{40}/\Gamma</math></b>	NODE=M049P31 NODE=M049P31
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.5 \times 10^{-4}$	90	<sup>1</sup> SHEN	16	BELL $\Upsilon(1S) \rightarrow$ $J/\psi\pi^+\pi^-X$		
				<sup>1</sup> SHEN 16 reports $[\Gamma(\Upsilon(1S) \rightarrow \chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))] < 9.5 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S)) = 3.8 \times 10^{-2}$ .		NODE=M049P31;LINKAGE=A
<b><math>\Gamma(Z_c(4200)^+Z_c(4200)^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{41}/\Gamma</math></b>	NODE=M049P17 NODE=M049P17
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<22.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$		
				<sup>1</sup> Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1$ .		NODE=M049P17;LINKAGE=A
<b><math>\Gamma(Z_c(3900)^\pm Z_c(4200)^\mp)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{42}/\Gamma</math></b>	NODE=M049P18 NODE=M049P18
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<8.1 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$		
				<sup>1</sup> Assuming $B(Z_c(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1 = B(Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm)$ .		NODE=M049P18;LINKAGE=A
<b><math>\Gamma(Z_c(3900)^+Z_c(3900)^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{43}/\Gamma</math></b>	NODE=M049P19 NODE=M049P19
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<1.8 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$		
				<sup>1</sup> Assuming $B(Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm) = 1$		NODE=M049P19;LINKAGE=A
<b><math>\Gamma(X(4050)^+X(4050)^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{44}/\Gamma</math></b>	NODE=M049P20 NODE=M049P20
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<15.8 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P)\pi^\pm X$		
				<sup>1</sup> Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1$		NODE=M049P20;LINKAGE=A
<b><math>\Gamma(X(4250)^+X(4250)^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{45}/\Gamma</math></b>	NODE=M049P21 NODE=M049P21
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<26.6 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P)\pi^\pm X$		
				<sup>1</sup> Assuming $B(X(4250)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1$		NODE=M049P21;LINKAGE=A
<b><math>\Gamma(X(4050)^\pm X(4250)^\mp)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_{46}/\Gamma</math></b>	NODE=M049P22 NODE=M049P22
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<44.2 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P)\pi^\pm X$		
				<sup>1</sup> Assuming $B(X(4050)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1 = B(X(4250)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm)$		NODE=M049P22;LINKAGE=A

$\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$   $\Gamma_{47}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S)\pi^\pm X$

<sup>1</sup> Assuming  $B(Z_c(4430)^\pm \rightarrow \psi(2S)\pi^\pm) = 1$

NODE=M049P23  
NODE=M049P23

NODE=M049P23;LINKAGE=A

 $\Gamma(X(4055)^\pm X(4055)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{48}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<23.3 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

<sup>1</sup> Assuming  $B(X(4055)^\pm \rightarrow \psi(2S)\pi^\pm) = 1$

NODE=M049P25  
NODE=M049P25

NODE=M049P25;LINKAGE=A

 $\Gamma(X(4055)^\pm Z_c(4430)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{49}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<45.5 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

<sup>1</sup> Assuming  $B(X(4055)^\pm \rightarrow \psi(2S)\pi^\pm) = 1 = B(Z_c(4430)^\pm \rightarrow \psi(2S)\pi^\pm)$

NODE=M049P26  
NODE=M049P26

NODE=M049P26;LINKAGE=A

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$   $\Gamma_{50}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.68$	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1 \times 10^3$	90	BLINOV	90	MD1 $\Upsilon(1S) \rightarrow \rho^0 \pi^0$
$<2 \times 10^2$	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0 \pi^0$
$<2.1 \times 10^3$	90	NICZYPORUK	83	LENA $\Upsilon(1S) \rightarrow \rho^0 \pi^0$

NODE=M049R11  
NODE=M049R11

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{51}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.90$	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

NODE=M049R05  
NODE=M049R05

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{52}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<5$	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow \pi^+ \pi^-$

NODE=M049R57  
NODE=M049R57

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{53}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<5$	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow K^+ K^-$

NODE=M049R58  
NODE=M049R58

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{54}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<5$	90	<sup>1</sup> BARU	96	MD1 $\Upsilon(1S) \rightarrow \rho\bar{\rho}$

<sup>1</sup> Supersedes BARU 92 in this node.

NODE=M049R59  
NODE=M049R59

NODE=M049R59;LINKAGE=A

 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{55}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.14 \pm 0.72 \pm 0.34$		$26 \pm 9$	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<18.4$	90		ANASTASSOV	99	CLE2 $e^+ e^- \rightarrow \text{hadrons}$
---------	----	--	------------	----	---

NODE=M049R72  
NODE=M049R72

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.36 \pm 0.37 \pm 0.29$	56	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+ K^-)$

NODE=M049R75  
NODE=M049R75

 $\Gamma(\omega\pi^+ \pi^-)/\Gamma_{\text{total}}$   $\Gamma_{57}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.46 \pm 0.67 \pm 0.72$	64	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-)\pi^0$

NODE=M049R76  
NODE=M049R76

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{58}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.42 \pm 0.50 \pm 0.58$	173	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R77  
NODE=M049R77

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$   $\Gamma_{59}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.63$	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+ K^-)$

NODE=M049R78  
NODE=M049R78

$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.79	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M049R79  
NODE=M049R79 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.24	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M049R80  
NODE=M049R80 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.02 \pm 0.68 \pm 0.34$	42	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R81  
NODE=M049R81 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.41	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R82  
NODE=M049R82 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.02 \pm 0.35 \pm 0.22$	24	SHEN	12A BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R83  
NODE=M049R83 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.25	90	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M049R84  
NODE=M049R84 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$12.8 \pm 2.0 \pm 2.3$	143 $\pm$ 22	SHEN	13 BELL	$\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

NODE=M049R06  
NODE=M049R06 $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.33 \pm 0.18$		37 $\pm$ 8	SHEN	13 BELL	$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R07  
NODE=M049R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.4	90		<sup>1</sup> DOBBS	12A	$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M049R07;LINKAGE=DO

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.92 \pm 0.85 \pm 0.37$	16 $\pm$ 5	SHEN	13 BELL	$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R08  
NODE=M049R08 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.11	90	SHEN	13 BELL	$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R09  
NODE=M049R09 $\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 2.8 \pm 1.3$	3.1k	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049P14  
NODE=M049P14 $\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$25.2 \pm 1.3 \pm 1.5$		$\approx$ 2k	<sup>1</sup> AUBERT	10C BABR	$\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

NODE=M049R32  
NODE=M049R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (units $10^{-3}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<19	90		<sup>2</sup> ALBRECHT	92J ARG	$e^+ e^- \rightarrow D^0 \pi^\pm X$

<sup>1</sup> For  $x_p > 0.1$ .<sup>2</sup> For  $x_p > 0.2$ .

NODE=M049R32;LINKAGE=AU

NODE=M049R32;LINKAGE=B

 $\Gamma(f_1(1285) X_{tetra})/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<62.4 $\times$ 10 <sup>-6</sup>	90	<sup>1</sup> JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049P16  
NODE=M049P16<sup>1</sup> For a tetraquark state  $X_{tetra}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{tetra}$  mass and width range from  $4.6 \times 10^{-6}$  to  $62.4 \times 10^{-6}$ .

NODE=M049P16;LINKAGE=A

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{73}/\Gamma$
<b>2.85±0.25 OUR AVERAGE</b>					
$2.81 \pm 0.49^{+0.20}_{-0.24}$		LEES	14G	BABR $e^+ e^- \rightarrow \overline{2H} X$	OCCUR=2
$2.86 \pm 0.19 \pm 0.21$	455	ASNER	07	CLEO $e^+ e^- \rightarrow \overline{2H} X$	

NODE=M049R33  
NODE=M049R33

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	COMMENT	$\Gamma_{74}/\Gamma$
<b>1.200±0.017</b>	1,2 DOBBS	12A $\Upsilon(1S) \rightarrow \text{hadrons}$	

NODE=M049R02  
NODE=M049R02

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

NODE=M049R02;LINKAGE=DO

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M049R02;LINKAGE=NC

 $\Gamma(ggg, \gamma g g \rightarrow \bar{d} \text{ anything})/\Gamma(ggg, \gamma g g \rightarrow \text{anything})$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{75}/\Gamma$
<b>3.36±0.23±0.25</b>	455	ASNER	07	CLEO $e^+ e^- \rightarrow \bar{d} X$	

NODE=M049R34  
NODE=M049R34

 $\Gamma(\gamma \pi^+ \pi^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_{75}/\Gamma$
<b>6.3±1.2±1.3</b>	<sup>1</sup> ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R70  
NODE=M049R70

<sup>1</sup> For  $m_{\pi\pi} > 1$  GeV.

NODE=M049R70;LINKAGE=A

 $\Gamma(\gamma \pi^0 \pi^0)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_{76}/\Gamma$
<b>1.7±0.6±0.3</b>	<sup>1</sup> ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R71  
NODE=M049R71

<sup>1</sup> For  $m_{\pi\pi} > 1$  GeV.

NODE=M049R71;LINKAGE=A

 $\Gamma(\gamma \pi \pi (\text{S-wave}))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_{77}/\Gamma$
<b>4.63±0.56±0.48</b>	LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	

NODE=M049P27  
NODE=M049P27

 $\Gamma(\gamma \pi^0 \eta)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{78}/\Gamma$
<b>&lt;2.4</b>	90	<sup>1</sup> BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S)$	

NODE=M049R47  
NODE=M049R47

<sup>1</sup> BESSON 07A obtained this limit for  $0.7 < m_{\pi^0 \eta} < 3$  GeV.

NODE=M049R47;LINKAGE=BE

 $\Gamma(\gamma K^+ K^-)/\Gamma_{\text{total}}$ 

( $2 < m_{K^+ K^-} < 3$  GeV)

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{79}/\Gamma$
<b>1.14±0.08±0.10</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

NODE=M049R24  
NODE=M049R24

NODE=M049R24

 $\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}}$ 

( $2 < m_{p \bar{p}} < 3$  GeV)

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{80}/\Gamma$
<b>&lt;0.6</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma p \bar{p}$	

NODE=M049R29

NODE=M049R29  
NODE=M049R29

 $\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{81}/\Gamma$
<b>7.0±1.1±1.0</b>	80 ± 12	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R20  
NODE=M049R20

 $\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{82}/\Gamma$
<b>5.4±1.5±1.3</b>	39 ± 11	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R21  
NODE=M049R21

 $\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{83}/\Gamma$
<b>7.4±2.5±2.5</b>	36 ± 12	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R22  
NODE=M049R22

 $\Gamma(\gamma \pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{84}/\Gamma$
<b>2.9±0.7±0.6</b>	29 ± 8	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R14  
NODE=M049R14

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{85}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.7 \pm 0.5$	$26 \pm 7$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R13  
 NODE=M049R13

 $\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{86}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.9 \pm 0.8$	$17 \pm 5$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R17  
 NODE=M049R17

 $\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{87}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 0.9 \pm 0.8$	$18 \pm 7$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R18  
 NODE=M049R18

 $\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{88}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.5 \pm 0.5 \pm 0.3$	$22 \pm 6$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R15  
 NODE=M049R15

 $\Gamma(\gamma 2\pi^+ 2\pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$   $\Gamma_{89}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.4 \pm 0.4 \pm 0.4$	$7 \pm 6$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R19  
 NODE=M049R19

 $\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$   $\Gamma_{90}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.2 \pm 0.2$	$2 \pm 2$	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M049R16  
 NODE=M049R16

 $\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{91}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.9$	90	ATHAR 07A	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \pi^+ \pi^- \eta, \gamma \rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 16$	90	RICHICHI 01B	CLE2	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M049R55  
 NODE=M049R55

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$   $\Gamma_{92}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.0$	90	ATHAR 07A	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta \rightarrow \gamma \gamma \gamma, \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 21$	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta$

NODE=M049R54  
 NODE=M049R54

 $\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$   $\Gamma_{93}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$< 3$	90	<sup>1</sup> ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M049R31  
 NODE=M049R31

<sup>1</sup> Assuming  $B(f_0(980) \rightarrow \pi\pi) = 1$ .

NODE=M049R31;LINKAGE=AT

 $\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}}$   $\Gamma_{94}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.9 \pm 0.6$			<b>OUR AVERAGE</b>		
$2.13 \pm 0.28 \pm 0.72$			<sup>1</sup> LEES 18A	BABR	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
$4.1 \pm 1.4 \pm 0.1$		17	<sup>2</sup> BESSON 11	CLEO	$\Upsilon(1S) \rightarrow K_S^0 K_S^0$
$3.7 \pm 0.9 \pm 0.8$			ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R52  
 NODE=M049R52

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 14$	90	<sup>3</sup> FULTON 90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
$< 19.4$	90	<sup>3</sup> ALBRECHT 89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$

<sup>1</sup> Using  $B(f_2'(1525) \rightarrow K\bar{K}) = 0.887 \pm 0.022$  and  $B(K^0 \bar{K}^0) = 1/2 B(K\bar{K})$ .

<sup>2</sup> BESSON 11 reports  $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$  from a measurement of  $[\Gamma(\Upsilon(1S) \rightarrow \gamma f_2'(1525))/\Gamma_{\text{total}}] \times [B(f_2'(1525) \rightarrow K\bar{K})]$  assuming  $B(f_2'(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$ , which we rescale to our best value  $B(f_2'(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The result also assumes  $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$  and  $B(f_2'(1525) \rightarrow K\bar{K}) = 4 B(f_2'(1525) \rightarrow K_S^0 K_S^0)$ .

<sup>3</sup> Assuming  $B(f_2'(1525) \rightarrow K\bar{K}) = 0.71$ .

NODE=M049R52;LINKAGE=A  
 NODE=M049R52;LINKAGE=BE

NODE=M049R52;LINKAGE=D

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{95}/\Gamma$
<b>10.1 ± 0.6</b>				<b>OUR AVERAGE</b>	
10.15 ± 0.59 <sup>+0.54</sup> <sub>-0.43</sub>		1 LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
10.5 ± 1.6 <sup>+1.9</sup> <sub>-1.8</sub>		2 BESSON	07A CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$	
10.2 ± 0.8 ± 0.7		ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
8.1 ± 2.3 <sup>+2.9</sup> <sub>-2.7</sub>		3 ANASTASSOV	99 CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	3 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<13	90	3 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<81	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	
<sup>1</sup> Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = 1/3 B(f_2(1270) \rightarrow \pi \pi)$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9})\%$ .					
<sup>2</sup> Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi \pi)/3$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.7^{+2.5}_{-1.2})\%$ .					
<sup>3</sup> Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$ .					

NODE=M049R51  
NODE=M049R51

NODE=M049R51;LINKAGE=A

NODE=M049R51;LINKAGE=BE

NODE=M049R51;LINKAGE=C

 $\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{96}/\Gamma$
<b>&lt;8.2</b>	90	1 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$	
<sup>1</sup> Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$ .					

NODE=M049R23  
NODE=M049R23

NODE=M049R23;LINKAGE=J

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{97}/\Gamma$
<b>&lt;1.5</b>	90	1 BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<6.1	90	2 BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$	OCCUR=2
<sup>1</sup> Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi \pi)/3$ and $B(f_0(1500) \rightarrow \pi \pi) = (0.349 \pm 0.023)\%$ .					
<sup>2</sup> Calculated by us using $B(f_0(1500) \rightarrow \eta \eta) = (5.1 \pm 0.9)\%$ .					

NODE=M049R44  
NODE=M049R44

NODE=M049R44;LINKAGE=BE

NODE=M049R44;LINKAGE=BS

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT	$\Gamma_{98}/\Gamma$
<b>1.04 ± 0.14 ± 0.33</b>	1 LEES	18A BABR	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	
<sup>1</sup> LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1500) \rightarrow \gamma K \bar{K}) = (2.08 \pm 0.27 \pm 0.65) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$ .				

NODE=M049P29  
NODE=M049P29

NODE=M049P29;LINKAGE=A

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{99}/\Gamma$
<b>&lt; 2.6</b>	90	1 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6.3	90	1 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
< 19	90	1 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$	OCCUR=2
< 8	90	2 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	OCCUR=2
< 24	90	3 SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	
<sup>1</sup> Assuming $B(f_0(1710) \rightarrow K \bar{K}) = 0.38$ .					
<sup>2</sup> Assuming $B(f_0(1710) \rightarrow \pi \pi) = 0.04$ .					
<sup>3</sup> Assuming $B(f_0(1710) \rightarrow \eta \eta) = 0.18$ .					

NODE=M049R53  
NODE=M049R53

NODE=M049R53;LINKAGE=E

NODE=M049R53;LINKAGE=F

NODE=M049R53;LINKAGE=A

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{100}/\Gamma$
<b>1.01 ± 0.26 ± 0.18</b>		1 LEES	18A BABR	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.7	90	ATHAR	06 CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	
<sup>1</sup> LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma K \bar{K}) = (2.02 \pm 0.51 \pm 0.35) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$ .					

NODE=M049R50  
NODE=M049R50

NODE=M049R50;LINKAGE=A

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_{101} / \Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>0.53 \pm 0.17 \pm 0.11</math></b>	<sup>1</sup> LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<sup>1</sup> LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi \pi) = (0.79 \pm 0.26 \pm 0.17) \times 10^{-5}$ assuming $B(\pi^0 \pi^0) = 1/3 B(\pi \pi)$ .			

NODE=M049P28  
NODE=M049P28

NODE=M049P28;LINKAGE=A

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0) / \Gamma_{\text{total}}$  $\Gamma_{102} / \Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.4</b>	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

NODE=M049R45  
NODE=M049R45 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$  $\Gamma_{103} / \Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.8</b>	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$

NODE=M049R46  
NODE=M049R46 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$  $\Gamma_{104} / \Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.3</b>	90	<sup>1</sup> ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
<sup>1</sup> Assuming $B(f_4(2050) \rightarrow \pi \pi) = 0.17$ .				

NODE=M049R30  
NODE=M049R30

NODE=M049R30;LINKAGE=AT

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$  $\Gamma_{105} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.0002</b>	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R63  
NODE=M049R63 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$  $\Gamma_{106} / \Gamma$ 

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 8</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
<2000	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R56  
NODE=M049R56 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$  $\Gamma_{107} / \Gamma$ 

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<120	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M049R41  
NODE=M049R41 $\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p}) / \Gamma_{\text{total}}$  $\Gamma_{108} / \Gamma$ 

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt; 11</b>	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma p \bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma p \bar{p}$

NODE=M049R42  
NODE=M049R42 $\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi) / \Gamma_{\text{total}}$  $\Gamma_{109} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.003</b>	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M049R62  
NODE=M049R62 $\Gamma(\gamma \eta_c(1S)) / \Gamma_{\text{total}}$  $\Gamma_{110} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.9 <math>\times 10^{-5}</math></b>	90	<sup>1</sup> KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<5.7 $\times 10^{-5}$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$
<sup>1</sup> Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

NODE=M049R38  
NODE=M049R38

NODE=M049R38;LINKAGE=A

 $\Gamma(\gamma \eta_c(2S)) / \Gamma_{\text{total}}$  $\Gamma_{111} / \Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4 <math>\times 10^{-4}</math></b>	90	<sup>1</sup> KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$
<sup>1</sup> Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

NODE=M049P30  
NODE=M049P30

NODE=M049P30;LINKAGE=A



$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.6 \times 10^{-5}$	90	<sup>1</sup> KATRENKO 20	BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.5 \times 10^{-4}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
<sup>1</sup> Using $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ decays.				

 $\Gamma_{112}/\Gamma$ NODE=M049R39  
NODE=M049R39

NODE=M049R39;LINKAGE=A

 $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7^{+2.4+0.4}_{-1.8-0.5}$		5	<sup>1</sup> KATRENKO 20	BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<2.3$	90		SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
<sup>1</sup> Using $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ decays.					

 $\Gamma_{113}/\Gamma$ NODE=M049R40  
NODE=M049R40

NODE=M049R40;LINKAGE=A

 $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.6 \times 10^{-6}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<3.3 \times 10^{-5}$	90	<sup>1</sup> KATRENKO 20	BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
<sup>1</sup> Using $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ decays.				

 $\Gamma_{114}/\Gamma$ NODE=M049R48  
NODE=M049R48

NODE=M049R48;LINKAGE=A

 $\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4 \times 10^{-5}$	90	<sup>1</sup> SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
<sup>1</sup> SHEN 10A reports $[\Gamma(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 1.6 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.8 \times 10^{-2}$ .				

 $\Gamma_{115}/\Gamma$ NODE=M049P32  
NODE=M049P32

NODE=M049P32;LINKAGE=A

 $\Gamma(\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-6}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 $\Gamma_{116}/\Gamma$ NODE=M049R68  
NODE=M049R68 $\Gamma(\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 $\Gamma_{117}/\Gamma$ NODE=M049R69  
NODE=M049R69 $\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 $\Gamma_{118}/\Gamma$ NODE=M049R74  
NODE=M049R74 $\Gamma(\gamma X \bar{X} (m_X < 3.1 \text{ GeV}))/\Gamma_{\text{total}}$  $(X \bar{X} = \text{vectors with } m < 3.1 \text{ GeV})$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1$	90	<sup>1</sup> BALEST 95	CLEO	$e^+e^- \rightarrow \gamma + X \bar{X}$

 $\Gamma_{119}/\Gamma$ NODE=M049R61  
NODE=M049R61  
NODE=M049R61

NODE=M049R61;LINKAGE=A

 $\Gamma(\gamma X \bar{X} (m_X < 4.5 \text{ GeV}))/\Gamma_{\text{total}}$  $X \text{ and } \bar{X} = \text{zero spin with } m < 4.5 \text{ GeV}$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<24$	90	<sup>1</sup> DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X \bar{X}$

 $\Gamma_{120}/\Gamma$ NODE=M049R01  
NODE=M049R01  
NODE=M049R01

NODE=M049R01;LINKAGE=DA

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$  $(1.5 \text{ GeV} < m_X < 5.0 \text{ GeV})$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.78$	95	ROSNER 07A	CLEO	$e^+e^- \rightarrow \gamma X$

 $\Gamma_{121}/\Gamma$ NODE=M049R64  
NODE=M049R64  
NODE=M049R64

$\Gamma(\gamma A^0)/\Gamma_{\text{total}}$ (A<sup>0</sup> = scalar with  $m < 8.0$  GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 4.5 \times 10^{-6}$	90	<sup>1</sup> DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X$
$< 3 \times 10^{-5}$	90	<sup>2</sup> BALEST	95 CLEO	$e^+e^- \rightarrow \gamma + X$
$< 5.6 \times 10^{-5}$	90	<sup>2</sup> ANTREASYAN 90C	CBAL	$e^+e^- \rightarrow \gamma + X$

<sup>1</sup> For a non-interacting scalar or pseudoscalar, A<sup>0</sup>, with mass  $m_{A^0} < 8.0$  GeV. 90% CL upper limits range from  $1.9 \times 10^{-6}$  to  $4.5 \times 10^{-6}$ .

<sup>2</sup> For any non-interacting long-lived particle with mass  $< 7.2$  GeV.

 $\Gamma_{122}/\Gamma$ 

NODE=M049R60

NODE=M049R60

NODE=M049R60

NODE=M049R60;LINKAGE=DA

NODE=M049R60;LINKAGE=A

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ (201 < M( $\mu^+ \mu^-$ ) < 3565 MeV)

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt; 9</b>	90	<sup>1</sup> LOVE	08 CLEO	$e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$
---------------	----	-------------------	---------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 16$	90	<sup>2</sup> JIA	22 BELL	$\Upsilon(2S) \rightarrow \gamma \mu^+ \mu^- \pi^+ \pi^-$
$< 9.7$	90	<sup>3</sup> LEES	13C BABR	$e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

<sup>1</sup> For a narrow scalar or pseudoscalar, A<sup>0</sup>, with  $201 < M(\mu^+ \mu^-) < 3565$  MeV, excluding  $J/\psi$ . Measured 90% CL limits as a function of  $M(\mu^+ \mu^-)$  range from  $1-9 \times 10^{-6}$ .

<sup>2</sup> For a narrow scalar or pseudoscalar, A<sup>0</sup>, with  $0.22 < M(A^0) < 9.2$  GeV, resulting in 90% CL upper limits ranging from  $3.1 \times 10^{-7}$  at  $M(A^0) = 0.22$  GeV to  $1.6 \times 10^{-5}$  at  $M(A^0) = 9.2$  GeV.

<sup>3</sup> For a narrow scalar or pseudoscalar, A<sup>0</sup>, with mass in the range 0.212–9.2 GeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{A^0}$  are in the range  $0.28-9.7 \times 10^{-6}$ .

 $\Gamma_{123}/\Gamma$ 

NODE=M049R65

NODE=M049R65

NODE=M049R65

NODE=M049R65;LINKAGE=LO

NODE=M049R65;LINKAGE=A

NODE=M049R65;LINKAGE=LE

 $\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ (2m<sub>τ</sub> < M( $\tau^+ \tau^-$ ) < 9.2 GeV)

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<b>&lt; 130</b>	90	<sup>1</sup> LEES	13R BABR	$\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
-----------------	----	-------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 150$	90	<sup>2</sup> JIA	22 BELL	$\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
$< 50$	90	<sup>3</sup> LOVE	08 CLEO	$e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

<sup>1</sup> For a narrow scalar or pseudoscalar, A<sup>0</sup>, with  $2m_\tau < M(A^0) < 9.2$  GeV, resulting in 90% CL upper limits of  $0.9 \times 10^{-5}$  at  $M(A^0) = 2m_\tau$ ,  $\approx 1.5 \times 10^{-5}$  at  $M(A^0) = 7.5$  GeV, and  $13 \times 10^{-5}$  at  $M(A^0) = 9.2$  GeV.

<sup>2</sup> For a narrow scalar or pseudoscalar, A<sup>0</sup>, with  $2m_\tau < M(A^0) < 9.2$  GeV, resulting in 90% CL upper limits ranging from  $3.8 \times 10^{-6}$  at  $M(A^0) = 2m_\tau$  to  $1.5 \times 10^{-4}$  at  $M(A^0) = 9.2$  GeV.

<sup>3</sup> For a narrow scalar or pseudoscalar, A<sup>0</sup>, with  $2m_\tau < M(A^0) < 7.5$  GeV, resulting in 90% CL limits ranging from  $1 \times 10^{-5}$  at  $M(A^0) = 2m_\tau$  to  $5 \times 10^{-5}$  at  $M(A^0) = 7.5$  GeV.

 $\Gamma_{124}/\Gamma$ 

NODE=M049R66

NODE=M049R66

NODE=M049R66

NODE=M049R66;LINKAGE=A

NODE=M049R66;LINKAGE=B

NODE=M049R66;LINKAGE=LO

 $\Gamma(\gamma A^0 \rightarrow \gamma g g)/\Gamma_{\text{total}}$ 

(0.5 GeV &lt; m &lt; 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b>&lt; 1 × 10<sup>-2</sup></b>	90	<sup>1</sup> LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$
---------------------------------	----	-------------------	----------	-------------------------------------

<sup>1</sup> For a narrow, CP-odd pseudoscalar, A<sup>0</sup>, searched for in 26 hadronic decay modes with invariant mass  $0.5 \text{ GeV} < m_{A^0} < 9.0$  GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $10^{-6}$  to  $10^{-2}$ .

 $\Gamma_{125}/\Gamma$ 

NODE=M049R03

NODE=M049R03

NODE=M049R03

NODE=M049R03;LINKAGE=A

 $\Gamma(\gamma A^0 \rightarrow \gamma s \bar{s})/\Gamma_{\text{total}}$ 

(0.5 GeV &lt; m &lt; 9.0 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<b>&lt; 1 × 10<sup>-3</sup></b>	90	<sup>1</sup> LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$
---------------------------------	----	-------------------	----------	-------------------------------------

<sup>1</sup> For a narrow, CP-odd pseudoscalar, A<sup>0</sup>, searched for in 14 hadronic decay modes with invariant mass  $1.5 \text{ GeV} < m_{A^0} < 9.0$  GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $10^{-5}$  to  $10^{-3}$ .

 $\Gamma_{126}/\Gamma$ 

NODE=M049R04

NODE=M049R04

NODE=M049R04

NODE=M049R04;LINKAGE=A

LEPTON FAMILY NUMBER ( $LF$ ) VIOLATING MODES

LEPTON FAMILY NUMBER ( $LF$ ) VIOLATING MODES						NODE=M049230
$\Gamma(e^\pm \mu^\mp)/\Gamma_{total}$					$\Gamma_{127}/\Gamma$	NODE=M049P33 NODE=M049P33
VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;3.9</b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \mu^\mp$		
$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{total}$					$\Gamma_{128}/\Gamma$	NODE=M049R67 NODE=M049R67
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;2.7 <math>\times 10^{-6}</math> (CL = 90%)</b>				[<6.0 $\times 10^{-6}$ (CL = 95%) OUR 2022 BEST LIMIT]		
<b>&lt;2.7 <math>\times 10^{-6}</math></b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^\pm \tau^\mp$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<6.0 $\times 10^{-6}$	95	LOVE	08A	CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$		
$\Gamma(e^\pm \tau^\mp)/\Gamma_{total}$					$\Gamma_{129}/\Gamma$	NODE=M049P34 NODE=M049P34
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;2.7</b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \tau^\mp$		
$\Gamma(\gamma e^\pm \mu^\mp)/\Gamma_{total}$					$\Gamma_{130}/\Gamma$	NODE=M049P35 NODE=M049P35
VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;4.2</b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \mu^\mp$		
$\Gamma(\gamma \mu^\pm \tau^\mp)/\Gamma_{total}$					$\Gamma_{131}/\Gamma$	NODE=M049P36 NODE=M049P36
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;6.1</b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma \mu^\pm \tau^\mp$		
$\Gamma(\gamma e^\pm \tau^\mp)/\Gamma_{total}$					$\Gamma_{132}/\Gamma$	NODE=M049P37 NODE=M049P37
VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt;6.5</b>	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \tau^\mp$		

## OTHER DECAYS

OTHER DECAYS						NODE=M049235
$\Gamma(\text{invisible})/\Gamma_{total}$					$\Gamma_{133}/\Gamma$	NODE=M049R10 NODE=M049R10
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt; 3.0</b>	90	AUBERT	09AX	BABR $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<39	90	RUBIN	07	CLEO $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$		
<25	90	TAJIMA	07	BELL $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$		

 $\Upsilon(1S)$  REFERENCES

$\Upsilon(1S)$ REFERENCES						NODE=M049
SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)	REFID=62012	
JIA	22	PRL 128 081804	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=61657	
PATRA	22	JHEP 2205 095	S. Patra <i>et al.</i>	(BELLE Collab.)	REFID=61653	
KATRENKO	20	PRL 124 122001	P. Katrenko <i>et al.</i>	(BELLE Collab.)	REFID=60544	
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58949	
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58950	
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=57635	
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318	
SHEN	16	PR D93 112013	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=57515	
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939	
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345	
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54949	
LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55167	
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55451	
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395	
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>		REFID=54746	
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314	
BESSION	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=16737	
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16495	
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53211	
DEL-AMO-SA...	10C	PRL 104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53354	
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545	
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53201	
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52565	
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592	
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166	
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617	
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51945	
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620	
BESSION	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51638	
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079	
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)	REFID=51629	
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=51645	
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51510	
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50993	
BESSION	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147	
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035	
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452	

BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50183
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=49395
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)	REFID=48846
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)	REFID=48345
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
ANASTASSOV	99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=46609
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)	REFID=44146
CINABRO	94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)	REFID=44102
ALBRECHT	92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=42167
BARU	92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)	REFID=41860
BARU	92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)	REFID=42168
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
ANTREASYAN	90C	PL B251 204	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=41455
BLINOV	90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41361
FULTON	90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=41012
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
FULTON	89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=40918
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding, World Scientific, Singapore					
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC	REFID=40742
SCHMITT	88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)	REFID=40582
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU	86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22284
ALBRECHT	85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22282
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
MACKAY	84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)	REFID=22281
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)	REFID=22273
GILES	83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)	REFID=22274
NICZYPORUK	83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=12488
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
ARTAMONOV	82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22271
NICZYPORUK	82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22272
BERGER	80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22263
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264
BERGER	79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22259

$\chi_{b0}(1P)$ 

$$I^G(J^{PC}) = 0^+(0^{++})$$

$J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(2S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

NODE=M076

NODE=M076

 $\chi_{b0}(1P)$  MASS

NODE=M076M

VALUE (MeV)

DOCUMENT ID

**9859.44 ± 0.42 ± 0.31 OUR EVALUATION** From average  $\gamma$  energy below, using  $\Upsilon(2S)$   
mass = 10023.26 ± 0.31 MeV

NODE=M076M

→ UNCHECKED ←

 $m_{\chi_{b1}(1P)} - m_{\chi_{b0}(1P)}$ 

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

**32.49 ± 0.93**

LEES

14M

BABR

 $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ 

NODE=M076M2

NODE=M076M2

 $\gamma$  ENERGY IN  $\Upsilon(2S)$  DECAY

NODE=M076DM

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

**162.5 ± 0.4 OUR AVERAGE**

162.56 ± 0.19 ± 0.42

ARTUSO

05

CLEO

 $\Upsilon(2S) \rightarrow \gamma X$ 

162.0 ± 0.8 ± 1.2

EDWARDS

99

CLE2

 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$ 

162.1 ± 0.5 ± 1.4

ALBRECHT

85E

ARG

 $\Upsilon(2S) \rightarrow \text{conv.}\gamma X$ 

163.8 ± 1.6 ± 2.7

NERNST

85

CBAL

 $\Upsilon(2S) \rightarrow \gamma X$ 

158.0 ± 7 ± 1

HAAS

84

CLEO

 $\Upsilon(2S) \rightarrow \text{conv.}\gamma X$ 

• • • We do not use the following data for averages, fits, limits, etc. • • •

149.4 ± 0.7 ± 5.0

KLOPFEN...

83

CUSB

 $\Upsilon(2S) \rightarrow \gamma X$ 

NODE=M076DM

 $\chi_{b0}(1P)$  DECAY MODES

NODE=M076215;NODE=M076

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\gamma \Upsilon(1S)$	( 1.94 ± 0.27 ) %	
$\Gamma_2$ $D^0 X$	< 10.4	% 90%
$\Gamma_3$ $\pi^+ \pi^- K^+ K^- \pi^0$	< 1.6	$\times 10^{-4}$ 90%
$\Gamma_4$ $2\pi^+ \pi^- K^- K_S^0$	< 5	$\times 10^{-5}$ 90%
$\Gamma_5$ $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 5	$\times 10^{-4}$ 90%
$\Gamma_6$ $2\pi^+ 2\pi^- 2\pi^0$	< 2.1	$\times 10^{-4}$ 90%
$\Gamma_7$ $2\pi^+ 2\pi^- K^+ K^-$	( 1.1 ± 0.6 ) $\times 10^{-4}$	
$\Gamma_8$ $2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.7	$\times 10^{-4}$ 90%
$\Gamma_9$ $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 5	$\times 10^{-4}$ 90%
$\Gamma_{10}$ $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 1.6	$\times 10^{-4}$ 90%
$\Gamma_{11}$ $3\pi^+ 3\pi^-$	< 8	$\times 10^{-5}$ 90%
$\Gamma_{12}$ $3\pi^+ 3\pi^- 2\pi^0$	< 6	$\times 10^{-4}$ 90%
$\Gamma_{13}$ $3\pi^+ 3\pi^- K^+ K^-$	( 2.4 ± 1.2 ) $\times 10^{-4}$	
$\Gamma_{14}$ $3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 1.0	$\times 10^{-3}$ 90%
$\Gamma_{15}$ $4\pi^+ 4\pi^-$	< 8	$\times 10^{-5}$ 90%
$\Gamma_{16}$ $4\pi^+ 4\pi^- 2\pi^0$	< 2.1	$\times 10^{-3}$ 90%
$\Gamma_{17}$ $J/\psi J/\psi$	< 7	$\times 10^{-5}$ 90%
$\Gamma_{18}$ $J/\psi \psi(2S)$	< 1.2	$\times 10^{-4}$ 90%
$\Gamma_{19}$ $\psi(2S) \psi(2S)$	< 3.1	$\times 10^{-5}$ 90%
$\Gamma_{20}$ $J/\psi(1S)$ anything	< 2.3	$\times 10^{-3}$ 90%

 $\chi_{b0}(1P)$  BRANCHING RATIOS

NODE=M076220

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE (%)

CL% EVTS

DOCUMENT ID

TECN

COMMENT

**1.94 ± 0.27 OUR AVERAGE**

2.07 ± 0.24 ± 0.21

1,2

LEES

14M

BABR

 $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ 

1.76 ± 0.30 ± 0.18

87

3,4

KORNICER

11

CLEO

 $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$ 

NODE=M076R1

NODE=M076R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.6	90	<sup>5</sup> LEES	11J	BABR	$\Upsilon(2S) \rightarrow X\gamma$
< 6	90	WALK	86	CBAL	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
< 11	90	PAUSS	83	CUSB	$\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>1</sup> LEES 14M quotes  $\Gamma(\chi_{b0}(1P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$   
 $= (7.75 \pm 0.91) \times 10^{-4}$  combining the results from samples of  $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$   
with and without converted photons. Assumes  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M076R1;LINKAGE=A

<sup>2</sup> LEES 14M reports  $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$   
 $= (7.75 \pm 0.91) \times 10^{-4}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) =$   
 $(3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is  
the systematic error from using our best value.

NODE=M076R1;LINKAGE=B

<sup>3</sup> Assuming  $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (2.48 \pm 0.05)\%$ .

NODE=M076R1;LINKAGE=KA

<sup>4</sup> KORNICER 11 reports  $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$   
 $= (6.59 \pm 0.96 \pm 0.60) \times 10^{-4}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) =$   
 $(3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is  
the systematic error from using our best value.

NODE=M076R1;LINKAGE=KR

<sup>5</sup> LEES 11J quotes a central value of  $\Gamma(\chi_{b0}(1P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow$   
 $\gamma\chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$ .

NODE=M076R1;LINKAGE=LE

$\Gamma(D^0 X)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<10.4 × 10 <sup>-2</sup>	90	<sup>6,7</sup> BRIERE	08	CLEO $\Upsilon(2S) \rightarrow \gamma D^0 X$

NODE=M076R01  
NODE=M076R01

<sup>6</sup> For  $p_{D^0} > 2.5$  GeV/c.

NODE=M076R01;LINKAGE=BR

<sup>7</sup> The authors also present their result as  $(5.6 \pm 3.6 \pm 0.5) \times 10^{-2}$ .

NODE=M076R01;LINKAGE=RI

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	<sup>8</sup> ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$

NODE=M076R02  
NODE=M076R02

<sup>8</sup> ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow$   
 $\gamma\chi_{b0}(1P))]$  <  $6 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) =$   
 $3.8 \times 10^{-2}$ .

NODE=M076R02;LINKAGE=AS

$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	<sup>9</sup> ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

NODE=M076R03  
NODE=M076R03

<sup>9</sup> ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow$   
 $\gamma\chi_{b0}(1P))]$  <  $2 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) =$   
 $3.8 \times 10^{-2}$ .

NODE=M076R03;LINKAGE=AS

$\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	<sup>10</sup> ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M076R04  
NODE=M076R04

<sup>10</sup> ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow$   
 $\gamma\chi_{b0}(1P))]$  <  $18 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) =$   
 $3.8 \times 10^{-2}$ .

NODE=M076R04;LINKAGE=AS

$\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	<sup>11</sup> ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

NODE=M076R05  
NODE=M076R05

<sup>11</sup> ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$   
<  $8 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R05;LINKAGE=AS

$\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$

VALUE (units 10 <sup>-4</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.6±0.1	7	<sup>12</sup> ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$

NODE=M076R06  
NODE=M076R06

<sup>12</sup> ASNER 08A reports  $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow$   
 $\gamma\chi_{b0}(1P))]$  =  $(4 \pm 2 \pm 1) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow$   
 $\gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our  
second error is the systematic error from using our best value.

NODE=M076R06;LINKAGE=AS

$\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	13 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$ 13 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 10 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R07  
NODE=M076R07

NODE=M076R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	14 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$ 14 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 20 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R08  
NODE=M076R08

NODE=M076R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	15 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$ 15 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R09  
NODE=M076R09

NODE=M076R09;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	16 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-$ 16 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R10  
NODE=M076R10

NODE=M076R10;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	17 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-2\pi^0$ 17 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 22 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R11  
NODE=M076R11

NODE=M076R11;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 1.2 \pm 0.2$	9	18 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-K^+K^-$ 18 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+3\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] = (9 \pm 4 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M076R12  
NODE=M076R12

NODE=M076R12;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<10	90	19 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-K^+K^-\pi^0$ 19 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+3\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 37 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R13  
NODE=M076R13

NODE=M076R13;LINKAGE=AS

 $\Gamma(4\pi^+4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	20 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+4\pi^-$ 20 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R14  
NODE=M076R14

NODE=M076R14;LINKAGE=AS

 $\Gamma(4\pi^+4\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<21	90	21 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+4\pi^-2\pi^0$ 21 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+4\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] < 77 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M076R15  
NODE=M076R15

NODE=M076R15;LINKAGE=AS

$\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	22 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma\psi X$
22 SHEN 12 reports $< 7.1 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ .				

NODE=M076R16  
NODE=M076R16

NODE=M076R16;LINKAGE=SH

 $\Gamma(J/\psi\psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	23 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma\psi X$
23 SHEN 12 reports $< 12 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi\psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ .				

NODE=M076R17  
NODE=M076R17

NODE=M076R17;LINKAGE=SH

 $\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	24 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma\psi X$
24 SHEN 12 reports $< 3.1 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$ .				

NODE=M076R18  
NODE=M076R18

NODE=M076R18;LINKAGE=SH

 $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.3 $\times 10^{-3}$	90	JIA	17A BELL	$e^+e^- \rightarrow \text{hadrons}$

NODE=M076R00  
NODE=M076R00 $\chi_{b0}(1P)$  CROSS-PARTICLE BRANCHING RATIOS

NODE=M076230

 $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$   
 $\Gamma_1/\Gamma \times \Gamma_{61}^{\Upsilon(2S)}/\Gamma^{\Upsilon(2S)}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 $\times 10^{-3}$	90	25 LEES	11J BABR	$\Upsilon(2S) \rightarrow X\gamma$
25 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}} < 4.6\%$ using $B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) = (3.8 \pm 0.4)\%$ .				

NODE=M076B02  
NODE=M076B02

NODE=M076B02;LINKAGE=LE

 $B(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$ NODE=M076B01  
NODE=M076B01

VALUE (units $10^{-5}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1.67 ± 0.28 OUR AVERAGE</b>				
2.9 $^{+1.7}_{-1.4}$ $^{+0.1}_{-0.8}$		26 LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$
1.63 ± 0.24 ± 0.15	87	KORNICER	11 CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
26 From a sample of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ with one converted photon.				

NODE=M076B01;LINKAGE=A

 $[B(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma\chi_{b0}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))]$ NODE=M076A01  
NODE=M076A01

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>3.28 ± 0.37</b>	27 LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$
27 From a sample of $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$ without converted photons.			

NODE=M076A01;LINKAGE=A

 $\chi_{b0}(1P)$  REFERENCES

NODE=M076

JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54313
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)	REFID=22290
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)	REFID=22286



**$\chi_{b1}(1P)$**

$I^G(J^{PC}) = 0^+(1^{++})$   
 $J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(2S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .  $J = 1$  from SKWARNICKI 87.

NODE=M077

NODE=M077

**$\chi_{b1}(1P)$  MASS**

NODE=M077M

VALUE (MeV)                      DOCUMENT ID  
**9892.78 ± 0.26 ± 0.31 OUR EVALUATION** From average  $\gamma$  energy below, using  $\Upsilon(2S)$   
 mass = 10023.26 ± 0.31 MeV

NODE=M077M  
 → UNCHECKED ←


**$\gamma$  ENERGY IN  $\Upsilon(2S)$  DECAY**

NODE=M077DM

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>129.63 ± 0.33 OUR AVERAGE</b>	Error includes scale factor of 1.3. See the ideogram below.		
129.58 ± 0.09 ± 0.29	ARTUSO	05	CLEO $\Upsilon(2S) \rightarrow \gamma X$
128.8 ± 0.4 ± 0.6	EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
131.7 ± 0.9 ± 1.3	WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
131.7 ± 0.3 ± 1.1	ALBRECHT	85E	ARG $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
130.6 ± 0.8 ± 2.4	NERNST	85	CBAL $\Upsilon(2S) \rightarrow \gamma X$
129 ± 0.8 ± 1	HAAS	84	CLEO $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
128.1 ± 0.4 ± 3.0	KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma X$
130.6 ± 3.0	PAUSS	83	CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M077DM

WEIGHTED AVERAGE  
 129.63 ± 0.33 (Error scaled by 1.3)



**$\chi_{b1}(1P)$  DECAY MODES**

NODE=M077215;NODE=M077

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\gamma \Upsilon(1S)$	(35.2 ± 2.0) %	DESIG=1
$\Gamma_2$ $D^0 X$	(12.6 ± 2.2) %	DESIG=2
$\Gamma_3$ $\pi^+ \pi^- K^+ K^- \pi^0$	( 2.0 ± 0.6) × 10 <sup>-4</sup>	DESIG=3
$\Gamma_4$ $2\pi^+ \pi^- K^- K_S^0$	( 1.3 ± 0.5) × 10 <sup>-4</sup>	DESIG=4
$\Gamma_5$ $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 6 × 10 <sup>-4</sup>	90% DESIG=5
$\Gamma_6$ $2\pi^+ 2\pi^- 2\pi^0$	( 8.0 ± 2.5) × 10 <sup>-4</sup>	DESIG=6
$\Gamma_7$ $2\pi^+ 2\pi^- K^+ K^-$	( 1.5 ± 0.5) × 10 <sup>-4</sup>	DESIG=7
$\Gamma_8$ $2\pi^+ 2\pi^- K^+ K^- \pi^0$	( 3.5 ± 1.2) × 10 <sup>-4</sup>	DESIG=8
$\Gamma_9$ $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	( 8.6 ± 3.2) × 10 <sup>-4</sup>	DESIG=9
$\Gamma_{10}$ $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	( 9.3 ± 3.3) × 10 <sup>-4</sup>	DESIG=10
$\Gamma_{11}$ $3\pi^+ 3\pi^-$	( 1.9 ± 0.6) × 10 <sup>-4</sup>	DESIG=11

$\Gamma_{12}$	$3\pi^+ 3\pi^- 2\pi^0$	$(1.7 \pm 0.5) \times 10^{-3}$	DESIG=12
$\Gamma_{13}$	$3\pi^+ 3\pi^- K^+ K^-$	$(2.6 \pm 0.8) \times 10^{-4}$	DESIG=13
$\Gamma_{14}$	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(7.5 \pm 2.6) \times 10^{-4}$	DESIG=14
$\Gamma_{15}$	$4\pi^+ 4\pi^-$	$(2.6 \pm 0.9) \times 10^{-4}$	DESIG=15
$\Gamma_{16}$	$4\pi^+ 4\pi^- 2\pi^0$	$(1.4 \pm 0.6) \times 10^{-3}$	DESIG=16
$\Gamma_{17}$	$\omega$ anything	$(4.9 \pm 1.4) \%$	DESIG=21
$\Gamma_{18}$	$\omega X_{tetra}$	$< 4.44 \times 10^{-4}$	90% DESIG=22
$\Gamma_{19}$	$J/\psi J/\psi$	$< 2.7 \times 10^{-5}$	90% DESIG=17
$\Gamma_{20}$	$J/\psi \psi(2S)$	$< 1.7 \times 10^{-5}$	90% DESIG=18
$\Gamma_{21}$	$\psi(2S)\psi(2S)$	$< 6 \times 10^{-5}$	90% DESIG=19
$\Gamma_{22}$	$J/\psi(1S)$ anything	$< 1.1 \times 10^{-3}$	90% DESIG=20
$\Gamma_{23}$	$J/\psi(1S) X_{tetra}$	$< 2.27 \times 10^{-4}$	90% DESIG=23

 **$\chi_{b1}(1P)$  BRANCHING RATIOS**

NODE=M077220

 **$\Gamma(\gamma \Upsilon(1S))/\Gamma_{total}$**  **$\Gamma_1/\Gamma$** 

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.352 ± 0.020 OUR AVERAGE</b>				
0.356 <sup>+0.016</sup> <sub>-0.022</sub> ± 0.019	964k	<sup>1</sup> FULSOM	18 BELL	$\Upsilon(2S) \rightarrow \gamma X$
0.364 ± 0.017 ± 0.019		<sup>2,3,4</sup> LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.331 ± 0.018 ± 0.017	3222	<sup>4,5</sup> KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
0.350 ± 0.023 ± 0.018	13k	<sup>6</sup> LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$
0.34 ± 0.07 ± 0.02	53	<sup>4,7,8</sup> WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.47 ± 0.18		KLOPFEN...	83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M077R1  
NODE=M077R1

OCCUR=4

<sup>1</sup> FULSOM 18 reports  $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (2.45 \pm 0.02 \pm 0.11 \pm 0.15) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=J

<sup>2</sup> LEES 14M quotes  $\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{total} = (2.51 \pm 0.12) \%$  combining the results from samples of  $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with and without converted photons.

NODE=M077R1;LINKAGE=B

<sup>3</sup> LEES 14M reports  $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (2.51 \pm 0.12) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=D

<sup>4</sup> Assuming  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05) \%$ .

NODE=M077R1;LINKAGE=KA

<sup>5</sup> KORNICER 11 reports  $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (22.8 \pm 0.4 \pm 1.2) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=KR

<sup>6</sup> LEES 11J reports  $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (24.1 \pm 0.6 \pm 1.5) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=LE

<sup>7</sup> WALK 86 quotes  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (5.8 \pm 0.9 \pm 0.7) \%$ .

NODE=M077R1;LINKAGE=A

<sup>8</sup> WALK 86 reports  $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (23.4 \pm 3.63 \pm 2.82) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=C

 **$\Gamma(D^0 X)/\Gamma_{total}$**  **$\Gamma_2/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>12.6 ± 1.9 ± 1.1</b>	2310	<sup>1</sup> BRIERE	08 CLEO	$\Upsilon(2S) \rightarrow \gamma D^0 X$

NODE=M077R01  
NODE=M077R01

<sup>1</sup> For  $p_{D^0} > 2.5$  GeV/c.

NODE=M077R01;LINKAGE=BR

 **$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{total}$**  **$\Gamma_3/\Gamma$** 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>2.0 ± 0.6 ± 0.1</b>	18	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

NODE=M077R02  
NODE=M077R02

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{total}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (14 \pm 3 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R02;LINKAGE=AS

$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.3±0.5±0.1</b>	11	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

NODE=M077R03  
NODE=M077R03

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (9 \pm 3 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R03;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;6</b>	90	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M077R04  
NODE=M077R04

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] < 42 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = 6.9 \times 10^{-2}$ .

NODE=M077R04;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.0±2.4±0.4</b>	46	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

NODE=M077R05  
NODE=M077R05

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (55 \pm 9 \pm 14) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R05;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.5±0.5±0.1</b>	18	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$

NODE=M077R06  
NODE=M077R06

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (10 \pm 3 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.5±1.2±0.2</b>	22	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$

NODE=M077R07  
NODE=M077R07

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (24 \pm 6 \pm 6) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6±3.2±0.4</b>	26	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$

NODE=M077R08  
NODE=M077R08

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (59 \pm 14 \pm 17) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.3±3.3±0.5</b>	21	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$

NODE=M077R09  
NODE=M077R09

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (64 \pm 16 \pm 16) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R09;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.9±0.6±0.1</b>	25	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-$

NODE=M077R10  
NODE=M077R10

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))] = (13 \pm 3 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R10;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17±5±1</b>	56	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

NODE=M077R11  
NODE=M077R11

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$   
 $= (119 \pm 18 \pm 32) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$   
 $= (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6±0.8±0.1</b>	21	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$

NODE=M077R12  
NODE=M077R12

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$   
 $= (18 \pm 4 \pm 4) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$   
 $= (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.5±2.6±0.4</b>	28	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$

NODE=M077R13  
NODE=M077R13

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$   
 $= (52 \pm 11 \pm 14) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$   
 $= (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.6±0.9±0.1</b>	24	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$

NODE=M077R14  
NODE=M077R14

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$   
 $= (18 \pm 4 \pm 5) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$   
 $= (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>14±5±1</b>	26	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$

NODE=M077R15  
NODE=M077R15

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$   
 $= (96 \pm 24 \pm 29) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))$   
 $= (6.9 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R15;LINKAGE=AS

 $\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.9±1.3±0.6</b>	51k	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M077R19  
NODE=M077R19 $\Gamma(\omega X_{\text{tetra}})/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;44.4 × 10<sup>-5</sup></b>	90	<sup>1</sup> JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M077R23  
NODE=M077R23

<sup>1</sup> For a tetraquark state  $X_{\text{tetra}}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{\text{tetra}}$  mass and width range from  $3.3 \times 10^{-5}$  to  $44.4 \times 10^{-5}$ .

NODE=M077R23;LINKAGE=A

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.7</b>	90	<sup>1</sup> SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$

NODE=M077R16  
NODE=M077R16

<sup>1</sup> SHEN 12 reports  $< 2.7 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$  assuming  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ .

NODE=M077R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.7</b>	90	<sup>1</sup> SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$

NODE=M077R17  
NODE=M077R17

<sup>1</sup> SHEN 12 reports  $< 1.7 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$  assuming  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ .

NODE=M077R17;LINKAGE=SH

$\Gamma(\psi(2S)\psi(2S))/\Gamma_{\text{total}}$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	<sup>1</sup> SHEN	12	BELL $\Upsilon(2S) \rightarrow \gamma\psi X$

<sup>1</sup> SHEN 12 reports  $< 6.2 \times 10^{-5}$  from a measurement of  $[\Gamma(\chi_{b1}(1P) \rightarrow \psi(2S)\psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P))]$  assuming  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$ .

 $\Gamma_{21}/\Gamma$ NODE=M077R18  
NODE=M077R18

NODE=M077R18;LINKAGE=SH

 $\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 $\times 10^{-3}$	90	JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma_{22}/\Gamma$ NODE=M077R00  
NODE=M077R00 $\Gamma(J/\psi(1S)X_{\text{tetra}})/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<22.7 $\times 10^{-5}$	90	<sup>1</sup> JIA	17A	BELL $e^+e^- \rightarrow \text{hadrons}$

 $\Gamma_{23}/\Gamma$ NODE=M077R22  
NODE=M077R22

<sup>1</sup> For a tetraquark state  $X_{\text{tetra}}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{\text{tetra}}$  mass and width range from  $1.8 \times 10^{-5}$  to  $22.7 \times 10^{-5}$ .

NODE=M077R22;LINKAGE=A

 $\chi_{b1}(1P)$  Cross-Particle Branching Ratios

NODE=M077230

$$\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}} \times \Gamma_1/\Gamma \times \Gamma_{59}^{\Upsilon(2S)}/\Gamma \Upsilon(2S)$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$24.1 \pm 0.6 \pm 1.5$	13k	LEES	11J	BABR $\Upsilon(2S) \rightarrow X\gamma$

NODE=M077B03  
NODE=M077B03

$$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.90 <math>\pm 0.34</math> OUR AVERAGE</b>				Error includes scale factor of 1.3. See the ideogram below.


NODE=M077B01  
NODE=M077B01

$6.86^{+0.47+0.44}_{-0.45-0.35}$		<sup>1</sup> LEES	14M	BABR $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$
$5.65 \pm 0.11 \pm 0.27$	3222	KORNICER	11	CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
$5.8 \pm 0.9 \pm 0.7$	53	WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>1</sup> From a sample of  $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$  with one converted photon.

NODE=M077B01;LINKAGE=A

WEIGHTED AVERAGE  
5.90  $\pm 0.34$  (Error scaled by 1.3)



$$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

(units  $10^{-4}$ )

$$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.30 <math>\pm 0.34</math> OUR AVERAGE</b>				

NODE=M077B02  
NODE=M077B02

$1.16^{+0.78+0.14}_{-0.67-0.16}$		<sup>1</sup> LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$1.33 \pm 0.30 \pm 0.23$	50	KORNICER	11	CLEO $e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>1</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with converted photons.

NODE=M077B02;LINKAGE=A

**B( $\chi_{b2}(1P) \rightarrow pX + \bar{p}X$ )/B( $\chi_{b1}(1P) \rightarrow pX + \bar{p}X$ )**

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.068±0.010±0.040</b>	BRIERE	07	CLEO $\Upsilon(2S) \rightarrow \gamma\chi_{bJ}(1P)$

NODE=M077R20  
NODE=M077R20**B( $\chi_{b0}(1P) \rightarrow pX + \bar{p}X$ )/B( $\chi_{b1}(1P) \rightarrow pX + \bar{p}X$ )**

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.11±0.15±0.20</b>	BRIERE	07	CLEO $\Upsilon(2S) \rightarrow \gamma\chi_{bJ}(1P)$

NODE=M077R21  
NODE=M077R21 **$\chi_{b1}(1P)$  REFERENCES**

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.) J
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)

NODE=M077

REFID=59535  
REFID=58318  
REFID=56343  
REFID=54313  
REFID=16769  
REFID=53936  
REFID=52574  
REFID=52577  
REFID=51887  
REFID=50454  
REFID=46612  
REFID=40019  
REFID=22290  
REFID=22288  
REFID=22289  
REFID=22287  
REFID=22285  
REFID=22286 **$h_b(1P)$** 

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Quantum numbers are quark model predictions,  $C = -$  established by  $\eta_b\gamma$  decay.

NODE=M204

NODE=M204

 **$h_b(1P)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9899.3±0.8 OUR AVERAGE</b>				
9899.3±0.4±1.0	112k	TAMPONI	15	BELL $e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
9899.1±0.4±1.0	70k	MIZUK	12	BELL $e^+e^- \rightarrow \pi^+\pi^- \text{ hadrons}$
9902 ±4 ±2	10.8k	LEES	11K	BABR $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$
9898.2 <sup>+1.1+1.0</sup> <sub>-1.0-1.1</sub>	50.0k	<sup>1</sup> ADACHI	12	BELL 10.86 $e^+e^- \rightarrow \pi^+\pi^- \text{ MM}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>Superseded by MIZUK 12.

NODE=M204M

NODE=M204M

NODE=M204M;LINKAGE=AD

 **$h_b(1P)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \eta_b(1S)\gamma$	(52 <sup>+6</sup> <sub>-5</sub> ) %

NODE=M204215;NODE=M204

DESIG=1

 **$h_b(1P)$  BRANCHING RATIOS**

$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE (units 10 <sup>-2</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>52<sup>+6</sup><sub>-5</sub> OUR AVERAGE</b>				
56 ±8 ±4	33.1k	<sup>1</sup> TAMPONI	15	BELL $e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
49.2±5.7 <sup>+5.6</sup> <sub>-3.3</sub>	24k	MIZUK	12	BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^- \text{ hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 10.8k LEES 11K BABR  $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$ <sup>1</sup>Using B( $\eta \rightarrow 2\gamma$ ) = (39.41 ± 0.20)%.

NODE=M204225

NODE=M204R01  
NODE=M204R01

NODE=M204R01;LINKAGE=A

 **$h_b(1P)$  REFERENCES**

TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)

NODE=M204

REFID=56996  
REFID=53962  
REFID=54718  
REFID=53937

$\chi_{b2}(1P)$ 

$$I^G(J^{PC}) = 0^+(2^{++})$$

$J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(2S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .  $J = 2$  from SKWARNICKI 87.

NODE=M078

NODE=M078

 $\chi_{b2}(1P)$  MASS

NODE=M078M

VALUE (MeV)                      DOCUMENT ID  
**9912.21±0.26±0.31 OUR EVALUATION** From average  $\gamma$  energy below, using  $\Upsilon(2S)$   
 mass = 10023.26 ± 0.31 MeV

NODE=M078M  
 → UNCHECKED ←

 $m_{\chi_{b2}(1P)} - m_{\chi_{b1}(1P)}$ 

NODE=M078DM2

VALUE (MeV)                      DOCUMENT ID      TECN      COMMENT  
**19.10±0.25 OUR AVERAGE** Error includes scale factor of 1.1.  
 19.81±0.65±0.20                      <sup>1</sup> AAIJ                      14BG LHCB       $pp \rightarrow \gamma\mu^+\mu^-X$   
 19.01±0.24                              LEES                      14M BABR       $\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M078DM2

<sup>1</sup> From the  $\chi_{bj}(1P) \rightarrow \Upsilon(1S)\gamma$  transition.

NODE=M078DM2;LINKAGE=A

 $\gamma$  ENERGY IN  $\Upsilon(2S)$  DECAY

NODE=M078DM

VALUE (MeV)                      DOCUMENT ID      TECN      COMMENT  
**110.44±0.29 OUR AVERAGE** Error includes scale factor of 1.1.  
 110.58±0.08±0.30                      ARTUSO      05      CLEO       $\Upsilon(2S) \rightarrow \gamma X$   
 110.8 ±0.3 ±0.6                      EDWARDS      99      CLE2       $\Upsilon(2S) \rightarrow \gamma\chi(1P)$   
 107.0 ±1.1 ±1.3                      WALK              86      CBAL       $\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$   
 110.6 ±0.3 ±0.9                      ALBRECHT      85E      ARG       $\Upsilon(2S) \rightarrow \text{conv.}\gamma X$   
 110.4 ±0.8 ±2.2                      NERNST              85      CBAL       $\Upsilon(2S) \rightarrow \gamma X$   
 109.5 ±0.7 ±1.0                      HAAS              84      CLEO       $\Upsilon(2S) \rightarrow \text{conv.}\gamma X$   
 108.2 ±0.3 ±2.0                      KLOPFEN...      83      CUSB       $\Upsilon(2S) \rightarrow \gamma X$   
 108.8 ±4.0                              PAUSS              83      CUSB       $\Upsilon(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M078DM

 $\chi_{b2}(1P)$  DECAY MODES

NODE=M078215;NODE=M078

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\gamma \Upsilon(1S)$	(18.0±1.0) %	
$\Gamma_2$ $D^0 X$	< 7.9 %	90%
$\Gamma_3$ $\pi^+\pi^-\pi^+K^-\pi^0$	( 8 ±5 ) × 10 <sup>-5</sup>	
$\Gamma_4$ $2\pi^+\pi^-\pi^+K_S^0$	< 1.0 × 10 <sup>-4</sup>	90%
$\Gamma_5$ $2\pi^+\pi^-\pi^+K_S^0 2\pi^0$	( 5.3±2.4 ) × 10 <sup>-4</sup>	
$\Gamma_6$ $2\pi^+2\pi^-\pi^0$	( 3.5±1.4 ) × 10 <sup>-4</sup>	
$\Gamma_7$ $2\pi^+2\pi^-\pi^+K^+K^-$	( 1.1±0.4 ) × 10 <sup>-4</sup>	
$\Gamma_8$ $2\pi^+2\pi^-\pi^+K^+K^-\pi^0$	( 2.1±0.9 ) × 10 <sup>-4</sup>	
$\Gamma_9$ $2\pi^+2\pi^-\pi^+K^+K^-2\pi^0$	( 3.9±1.8 ) × 10 <sup>-4</sup>	
$\Gamma_{10}$ $3\pi^+2\pi^-\pi^+K^+K_S^0\pi^0$	< 5 × 10 <sup>-4</sup>	90%
$\Gamma_{11}$ $3\pi^+3\pi^-$	( 7.0±3.1 ) × 10 <sup>-5</sup>	
$\Gamma_{12}$ $3\pi^+3\pi^-2\pi^0$	( 1.0±0.4 ) × 10 <sup>-3</sup>	
$\Gamma_{13}$ $3\pi^+3\pi^-\pi^+K^+K^-$	< 8 × 10 <sup>-5</sup>	90%
$\Gamma_{14}$ $3\pi^+3\pi^-\pi^+K^+K^-\pi^0$	( 3.6±1.5 ) × 10 <sup>-4</sup>	
$\Gamma_{15}$ $4\pi^+4\pi^-$	( 8 ±4 ) × 10 <sup>-5</sup>	
$\Gamma_{16}$ $4\pi^+4\pi^-2\pi^0$	( 1.8±0.7 ) × 10 <sup>-3</sup>	
$\Gamma_{17}$ $J/\psi J/\psi$	< 4 × 10 <sup>-5</sup>	90%
$\Gamma_{18}$ $J/\psi\psi(2S)$	< 5 × 10 <sup>-5</sup>	90%
$\Gamma_{19}$ $\psi(2S)\psi(2S)$	< 1.6 × 10 <sup>-5</sup>	90%
$\Gamma_{20}$ $J/\psi(1S)$ anything	( 1.5±0.4 ) × 10 <sup>-3</sup>	

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

DESIG=9

DESIG=10

DESIG=11

DESIG=12

DESIG=13

DESIG=14

DESIG=15

DESIG=16

DESIG=17

DESIG=18

DESIG=19

DESIG=20

$\chi_{b2}(1P)$  BRANCHING RATIOS

NODE=M078220

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$						$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>0.180 ± 0.010 OUR AVERAGE</b>						
$0.164^{+0.009}_{-0.010} \pm 0.008$	503k	<sup>1</sup> FULSOM	18	BELL	$\Upsilon(2S) \rightarrow \gamma X$	
$0.185 \pm 0.008 \pm 0.009$		<sup>2,3,4</sup> LEES	14M	BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
$0.186 \pm 0.011 \pm 0.009$	1770	<sup>4,5</sup> KORNICER	11	CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$	
$0.194^{+0.014}_{-0.017} \pm 0.009$	8k	<sup>6</sup> LEES	11J	BABR	$\Upsilon(2S) \rightarrow X \gamma$	
$0.25 \pm 0.06 \pm 0.01$	35	<sup>4,7,8</sup> WALK	86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
$0.20 \pm 0.05$		KLOPFEN...	83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

NODE=M078R1  
NODE=M078R1

OCCUR=2

<sup>1</sup> FULSOM 18 reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.17 \pm 0.01^{+0.06}_{-0.07}) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=H

<sup>2</sup> LEES 14M quotes  $\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}} = (1.32 \pm 0.06)\%$  combining the results from samples of  $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with and without converted photons.

NODE=M078R1;LINKAGE=B

<sup>3</sup> LEES 14M reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.32 \pm 0.06) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=F

<sup>4</sup> Assuming  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M078R1;LINKAGE=KA

<sup>5</sup> KORNICER 11 reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.33 \pm 0.04 \pm 0.07) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=KR

<sup>6</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (13.9 \pm 0.5^{+0.9}_{-1.1}) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=LE

<sup>7</sup> WALK 86 quotes  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (4.4 \pm 0.9 \pm 0.5) \%$ .

NODE=M078R1;LINKAGE=A

<sup>8</sup> WALK 86 reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (17.7 \pm 3.6 \pm 2.0) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=E

$\Gamma(D^0 X)/\Gamma_{\text{total}}$						$\Gamma_2/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$< 7.9 \times 10^{-2}$	90	<sup>1,2</sup> BRIERE	08	CLEO	$\Upsilon(2S) \rightarrow \gamma D^0 X$	

NODE=M078R01  
NODE=M078R01

<sup>1</sup> For  $p_{D^0} > 2.5$  GeV/c.

NODE=M078R01;LINKAGE=BR

<sup>2</sup> The authors also present their result as  $(5.4 \pm 1.9 \pm 0.5) \times 10^{-2}$ .

NODE=M078R01;LINKAGE=RI

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$						$\Gamma_3/\Gamma$
VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
<b>0.84 ± 0.50 ± 0.04</b>	8	<sup>1</sup> ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$	

NODE=M078R02  
NODE=M078R02

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 3 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$						$\Gamma_4/\Gamma$
VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT		
<b>&lt; 1.0</b>	90	<sup>1</sup> ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$	

NODE=M078R03  
NODE=M078R03

<sup>1</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 7 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .

NODE=M078R03;LINKAGE=AS



$\Gamma(2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.3±2.4±0.3</b>	11	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M078R04  
 NODE=M078R04

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(38 \pm 14 \pm 10) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R04;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.5±1.4±0.2</b>	19	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

NODE=M078R05  
 NODE=M078R05

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(25 \pm 8 \pm 6) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R05;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.1±0.4±0.1</b>	14	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$

NODE=M078R06  
 NODE=M078R06

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(8 \pm 2 \pm 2) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.1±0.9±0.1</b>	13	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$

NODE=M078R07  
 NODE=M078R07

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(15 \pm 5 \pm 4) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.9±1.8±0.2</b>	11	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$

NODE=M078R08  
 NODE=M078R08

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(28 \pm 11 \pm 7) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$

NODE=M078R09  
 NODE=M078R09

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] <  $36 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .

NODE=M078R09;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.70±0.31±0.03</b>	9	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-$

NODE=M078R10  
 NODE=M078R10

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(5 \pm 2 \pm 1) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R10;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.2±3.6±0.5</b>	34	<sup>1</sup> ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-2\pi^0$

NODE=M078R11  
 NODE=M078R11

<sup>1</sup> ASNER 08A reports [ $\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$ ]  $\times$  [ $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))$ ] =  $(73 \pm 16 \pm 20) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R11;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.8</b>	90	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .				

NODE=M078R12  
NODE=M078R12

NODE=M078R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.6±1.5±0.2</b>	14	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (26 \pm 8 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R13  
NODE=M078R13

NODE=M078R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.84±0.40±0.04</b>	7	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R14  
NODE=M078R14

NODE=M078R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18±7±1</b>	29	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (132 \pm 31 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R15  
NODE=M078R15

NODE=M078R15;LINKAGE=AS

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	1 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 4.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ .				

NODE=M078R16  
NODE=M078R16

NODE=M078R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5</b>	90	1 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 4.9 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ .				

NODE=M078R17  
NODE=M078R17

NODE=M078R17;LINKAGE=SH

 $\Gamma(\psi(2S) \psi(2S))/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.6</b>	90	1 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 1.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \psi(2S) \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$ .				

NODE=M078R18  
NODE=M078R18

NODE=M078R18;LINKAGE=SH

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.50±0.34±0.22</b>	462	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M078R00  
NODE=M078R00 $\chi_{b2}(1P)$  Cross-Particle Branching Ratios

NODE=M078230

$$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_{60}^{\Upsilon(2S)}/\Gamma \Upsilon(2S)$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>13.9±0.5<sup>+0.9</sup><sub>-1.1</sub></b>	8k	LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$

NODE=M078B03  
NODE=M078B03

**$B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$** 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.38 \pm 0.16</math></b>	<b>OUR AVERAGE</b>			
$3.63^{+0.36+0.18}_{-0.34-0.19}$		<sup>1</sup> LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$3.29 \pm 0.09 \pm 0.16$	1770	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
$4.4 \pm 0.9 \pm 0.5$	35	WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>1</sup> From a sample of  $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with converted photons.

NODE=M078B01  
NODE=M078B01

NODE=M078B01;LINKAGE=A

 **$[B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$** 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b><math>55.6 \pm 1.6</math></b>	<sup>1</sup> LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$

<sup>1</sup> From a sample of  $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$  events without converted photons.

NODE=M078A00  
NODE=M078A00

NODE=M078A00;LINKAGE=A

 **$B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$** 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>3.8 \pm 0.5</math></b>	<b>OUR AVERAGE</b>			
$4.68^{+0.99}_{-0.92} \pm 0.37$		<sup>1</sup> LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$3.56 \pm 0.40 \pm 0.41$	126	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>1</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with converted photons.

NODE=M078B02  
NODE=M078B02

NODE=M078B02;LINKAGE=A

 **$\chi_{b2}(1P)$  REFERENCES**

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)	REFID=59535
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56199
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54313
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)	REFID=40019
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)	REFID=22290
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)	REFID=22286

NODE=M078

REFID=59535  
REFID=58318  
REFID=56199  
REFID=56343  
REFID=54313  
REFID=16769  
REFID=53936  
REFID=52574  
REFID=52577  
REFID=50454  
REFID=46612  
REFID=40019  
REFID=22290  
REFID=22288  
REFID=22289  
REFID=22287  
REFID=22285  
REFID=22286

NODE=M200

**$\eta_b(2S)$**

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M200

 **$\eta_b(2S)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>9999.0 \pm 3.5^{+2.8}_{-1.9}</math></b>	26k	<sup>1</sup> MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- + \text{hadrons}$

••• We do not use the following data for averages, fits, limits, etc. •••

9974.6  $\pm$  2.3  $\pm$  2.1    11  $\pm$  4    <sup>2,3,4</sup> DOBBS    12     $\Upsilon(2S) \rightarrow \gamma \text{ hadrons}$

<sup>1</sup> Assuming  $\Gamma_{\eta_b(2S)} = 4.9$  MeV. Not independent of the corresponding mass difference measurement.

<sup>2</sup> SANDILYA 13 (Belle Collab.) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of  $(157.8 \pm 3.6) \times 10^6 \Upsilon(2S)$  decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction  $B(\Upsilon(2S) \rightarrow \eta_b(2S) \gamma) \times \sum_i B(\eta_b(2S) \rightarrow X_i) < 4.9 \times 10^{-6}$ , summed over the exclusive hadronic final states  $X_i$ , is an order of magnitude smaller than that reported by DOBBS 12.

<sup>3</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

<sup>4</sup> Assuming  $\Gamma_{\eta_b(2S)} = 5$  MeV. Not independent of the corresponding mass difference measurement.

NODE=M200M

NODE=M200M

NODE=M200M;LINKAGE=MI

NODE=M200M;LINKAGE=A

NODE=M200M;LINKAGE=DO

NODE=M200M;LINKAGE=NI

 **$m_{\Upsilon(2S)} - m_{\eta_b(2S)}$** 

NODE=M200DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$24.3 \pm 3.5^{+2.8}_{-1.9}$	26k	<sup>5</sup> MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons

NODE=M200DM

• • • We do not use the following data for averages, fits, limits, etc. • • •

48.7±2.3±2.1      11 ± 4    <sup>6,7,8</sup> DOBBS      12       $\Upsilon(2S) \rightarrow \gamma$  hadrons

<sup>5</sup> Assuming  $\Gamma_{\eta_b(2S)} = 4.9$  MeV. Not independent of the corresponding mass measurement.

NODE=M200DM;LINKAGE=MI

<sup>6</sup> SANDILYA 13 (Belle Collab.) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of  $(157.8 \pm 3.6) \times 10^6$   $\Upsilon(2S)$  decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction  $B(\Upsilon(2S) \rightarrow \eta_b(2S)\gamma) \times \sum_i B(\eta_b(2S) \rightarrow X_i) < 4.9 \times 10^{-6}$ , summed over the exclusive hadronic final states  $X_i$ , is an order of magnitude smaller than that reported by DOBBS 12.

NODE=M200DM;LINKAGE=A

<sup>7</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M200DM;LINKAGE=DO

<sup>8</sup> Assuming  $\Gamma_{\eta_b(2S)} = 5$  MeV. Not independent of the corresponding mass measurement.

NODE=M200DM;LINKAGE=NI

### $\eta_b(2S)$ WIDTH

NODE=M200W

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^-$ hadrons

NODE=M200W

### $\eta_b(2S)$ DECAY MODES

NODE=M200215;NODE=M200

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ hadrons	seen

DESIG=1

### $\eta_b(2S)$ BRANCHING RATIOS

NODE=M200225

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	26k	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^-$ hadrons

NODE=M200R01  
NODE=M200R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen      <sup>9,10</sup> DOBBS      12       $\Upsilon(2S) \rightarrow \gamma$  hadrons

<sup>9</sup> SANDILYA 13 (Belle Collab.) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of  $(157.8 \pm 3.6) \times 10^6$   $\Upsilon(2S)$  decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction  $B(\Upsilon(2S) \rightarrow \eta_b(2S)\gamma) \times \sum_i B(\eta_b(2S) \rightarrow X_i) < 4.9 \times 10^{-6}$ , summed over the exclusive hadronic final states  $X_i$ , is an order of magnitude smaller than that reported by DOBBS 12.

NODE=M200R01;LINKAGE=A

<sup>10</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M200R01;LINKAGE=DO

### $\eta_b(2S)$ REFERENCES

NODE=M200

SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

REFID=55590  
REFID=54288  
REFID=54718

$\Upsilon(2S)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M052

 **$\Upsilon(2S)$  MASS**

NODE=M052M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10023.4±0.5 OUR AVERAGE</b>	[10023.26 ± 0.31 MeV OUR 2022 AVERAGE]		
<b>10023.4±0.5</b>	<sup>1</sup> SHAMOV	23	RVUE e <sup>+</sup> e <sup>-</sup> → hadrons
10022.7±0.4	<sup>2</sup> SHAMOV	23	RVUE e <sup>+</sup> e <sup>-</sup> → hadrons
10023.5±0.5	<sup>3,4</sup> ARTAMONOV	00	MD1 e <sup>+</sup> e <sup>-</sup> → hadrons
10023.6±0.5	<sup>5,6</sup> BARU	86B	MD1 e <sup>+</sup> e <sup>-</sup> → hadrons
10023.1±0.4	<sup>7</sup> BARBER	84	ARG e <sup>+</sup> e <sup>-</sup> → hadrons

NODE=M052M  
NEW

••• We do not use the following data for averages, fits, limits, etc. •••

OCCUR=2

<sup>1</sup> Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.

NODE=M052M;LINKAGE=A

<sup>2</sup> Obtained by reanalysing ARGUS and Crystal Ball data (BARBER 84), but not authored by the ARGUS and Crystal Ball collaboration.

NODE=M052M;LINKAGE=B

<sup>3</sup> Reanalysis of BARU 86B using new electron mass (COHEN 87).

NODE=M052M;LINKAGE=AR

<sup>4</sup> Superseded by SHAMOV 23.

NODE=M052M;LINKAGE=E

<sup>5</sup> Reanalysis of ARTAMONOV 84.

NODE=M052M;LINKAGE=C

<sup>6</sup> Superseded by ARTAMONOV 00.

NODE=M052M;LINKAGE=RZ

<sup>7</sup> Reanalysed by SHAMOV 23.

NODE=M052M;LINKAGE=D

 **$m\Upsilon(3S) - m\Upsilon(2S)$** 

NODE=M052DM3

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>331.50±0.02±0.13</b>	LEES	11C	BABR e <sup>+</sup> e <sup>-</sup> → π <sup>+</sup> π <sup>-</sup> X

NODE=M052DM3

 **$\Upsilon(2S)$  WIDTH**

NODE=M052W

VALUE (keV)	DOCUMENT ID	COMMENT
<b>31.98±2.63 OUR EVALUATION</b>	See the Note on "Width Determinations of the $\Upsilon$ States"	

NODE=M052W  
→ UNCHECKED ← **$\Upsilon(2S)$  DECAY MODES**

NODE=M052215;NODE=M052

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\Upsilon(1S)\pi^+\pi^-$	(17.85 ± 0.26) %	
$\Gamma_2$ $\Upsilon(1S)\pi^0\pi^0$	( 8.6 ± 0.4 ) %	
$\Gamma_3$ $\tau^+\tau^-$	( 2.00 ± 0.21 ) %	
$\Gamma_4$ $\mu^+\mu^-$	( 1.93 ± 0.17 ) %	S=2.2
$\Gamma_5$ $e^+e^-$	( 1.91 ± 0.16 ) %	
$\Gamma_6$ $\Upsilon(1S)\pi^0$	< 4 × 10 <sup>-5</sup>	CL=90%
$\Gamma_7$ $\Upsilon(1S)\eta$	( 2.9 ± 0.4 ) × 10 <sup>-4</sup>	S=2.0
$\Gamma_8$ $J/\psi(1S)$ anything	< 6 × 10 <sup>-3</sup>	CL=90%
$\Gamma_9$ $J/\psi(1S)\eta_c$	< 5.4 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{10}$ $J/\psi(1S)\chi_{c0}$	< 3.4 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{11}$ $J/\psi(1S)\chi_{c1}$	< 1.2 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{12}$ $J/\psi(1S)\chi_{c2}$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{13}$ $J/\psi(1S)\eta_c(2S)$	< 2.5 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{14}$ $J/\psi(1S)X(3940)$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{15}$ $J/\psi(1S)X(4160)$	< 2.0 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{16}$ $\chi_{c1}$ anything	( 2.2 ± 0.5 ) × 10 <sup>-4</sup>	
$\Gamma_{17}$ $\chi_{c1}(1P)^0 X_{tetra}$	< 3.67 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{18}$ $\chi_{c2}$ anything	( 2.3 ± 0.8 ) × 10 <sup>-4</sup>	
$\Gamma_{19}$ $\psi(2S)\eta_c$	< 5.1 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{20}$ $\psi(2S)\chi_{c0}$	< 4.7 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{21}$ $\psi(2S)\chi_{c1}$	< 2.5 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{22}$ $\psi(2S)\chi_{c2}$	< 1.9 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{23}$ $\psi(2S)\eta_c(2S)$	< 3.3 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{24}$ $\psi(2S)X(3940)$	< 3.9 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{25}$ $\psi(2S)X(4160)$	< 3.9 × 10 <sup>-6</sup>	CL=90%

DESIG=4

DESIG=5

DESIG=3

DESIG=1

DESIG=2

DESIG=10

DESIG=6

DESIG=20

DESIG=143

DESIG=144

DESIG=145

DESIG=146

DESIG=147

DESIG=148

DESIG=149

DESIG=157

DESIG=160

DESIG=158

DESIG=150

DESIG=151

DESIG=152

DESIG=153

DESIG=154

DESIG=155

DESIG=156

Γ <sub>26</sub>	$Z_c(3900)^+ Z_c(3900)^-$	< 1.0	$\times 10^{-6}$	CL=90%	DESIG=162
Γ <sub>27</sub>	$Z_c(4200)^+ Z_c(4200)^-$	< 1.67	$\times 10^{-5}$	CL=90%	DESIG=163
Γ <sub>28</sub>	$Z_c(3900)^\pm Z_c(4200)^\mp$	< 7.3	$\times 10^{-6}$	CL=90%	DESIG=164
Γ <sub>29</sub>	$X(4050)^+ X(4050)^-$	< 1.35	$\times 10^{-5}$	CL=90%	DESIG=165
Γ <sub>30</sub>	$X(4250)^+ X(4250)^-$	< 2.67	$\times 10^{-5}$	CL=90%	DESIG=166
Γ <sub>31</sub>	$X(4050)^\pm X(4250)^\mp$	< 2.72	$\times 10^{-5}$	CL=90%	DESIG=167
Γ <sub>32</sub>	$Z_c(4430)^+ Z_c(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%	DESIG=168
Γ <sub>33</sub>	$X(4055)^\pm X(4055)^\mp$	< 1.11	$\times 10^{-5}$	CL=90%	DESIG=170
Γ <sub>34</sub>	$X(4055)^\pm Z_c(4430)^\mp$	< 2.11	$\times 10^{-5}$	CL=90%	DESIG=171
Γ <sub>35</sub>	${}^2H$ anything	$(2.78^{+0.30}_{-0.26}) \times 10^{-5}$		S=1.2	DESIG=16
Γ <sub>36</sub>	hadrons	$(94 \pm 11) \%$			DESIG=101
Γ <sub>37</sub>	$ggg$	$(58.8 \pm 1.2) \%$			DESIG=105
Γ <sub>38</sub>	$\gamma g g$	$(1.87 \pm 0.28) \%$			DESIG=106
Γ <sub>39</sub>	$\phi K^+ K^-$	$(1.6 \pm 0.4) \times 10^{-6}$			DESIG=133
Γ <sub>40</sub>	$\omega \pi^+ \pi^-$	< 2.58	$\times 10^{-6}$	CL=90%	DESIG=134
Γ <sub>41</sub>	$K^*(892)^0 K^- \pi^+ + c.c.$	$(2.3 \pm 0.7) \times 10^{-6}$			DESIG=135
Γ <sub>42</sub>	$\phi f'_2(1525)$	< 1.33	$\times 10^{-6}$	CL=90%	DESIG=136
Γ <sub>43</sub>	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%	DESIG=137
Γ <sub>44</sub>	$\rho(770) a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%	DESIG=138
Γ <sub>45</sub>	$K^*(892)^0 K_2^*(1430)^0 + c.c.$	$(1.5 \pm 0.6) \times 10^{-6}$			DESIG=139
Γ <sub>46</sub>	$K_1(1270)^\pm K^\mp$	< 3.22	$\times 10^{-6}$	CL=90%	DESIG=140
Γ <sub>47</sub>	$K_1(1400)^\pm K^\mp$	< 8.3	$\times 10^{-7}$	CL=90%	DESIG=141
Γ <sub>48</sub>	$b_1(1235)^\pm \pi^\mp$	< 4.0	$\times 10^{-7}$	CL=90%	DESIG=142
Γ <sub>49</sub>	$\rho \pi$	< 1.16	$\times 10^{-6}$	CL=90%	DESIG=126
Γ <sub>50</sub>	$\pi^+ \pi^- \pi^0$	< 8.0	$\times 10^{-7}$	CL=90%	DESIG=127
Γ <sub>51</sub>	$\omega \pi^0$	< 1.63	$\times 10^{-6}$	CL=90%	DESIG=128
Γ <sub>52</sub>	$\pi^+ \pi^- \pi^0 \pi^0$	$(1.30 \pm 0.28) \times 10^{-5}$			DESIG=129
Γ <sub>53</sub>	$K_S^0 K^+ \pi^- + c.c.$	$(1.14 \pm 0.33) \times 10^{-6}$			DESIG=130
Γ <sub>54</sub>	$K^*(892)^0 \bar{K}^0 + c.c.$	< 4.22	$\times 10^{-6}$	CL=90%	DESIG=131
Γ <sub>55</sub>	$K^*(892)^- K^+ + c.c.$	< 1.45	$\times 10^{-6}$	CL=90%	DESIG=132
Γ <sub>56</sub>	$f_1(1285)$ anything	$(2.2 \pm 1.6) \times 10^{-3}$			DESIG=159
Γ <sub>57</sub>	$f_1(1285) X_{tetra}$	< 6.47	$\times 10^{-5}$	CL=90%	DESIG=161
Γ <sub>58</sub>	Sum of 100 exclusive modes	$(2.90 \pm 0.30) \times 10^{-3}$			DESIG=121

## Radiative decays

Γ <sub>59</sub>	$\gamma \chi_{b1}(1P)$	$(6.9 \pm 0.4) \%$			NODE=M052;CLUMP=A DESIG=8
Γ <sub>60</sub>	$\gamma \chi_{b2}(1P)$	$(7.15 \pm 0.35) \%$			DESIG=7
Γ <sub>61</sub>	$\gamma \chi_{b0}(1P)$	$(3.8 \pm 0.4) \%$			DESIG=9
Γ <sub>62</sub>	$\gamma f_0(1710)$	< 5.9	$\times 10^{-4}$	CL=90%	DESIG=13
Γ <sub>63</sub>	$\gamma f'_2(1525)$	< 5.3	$\times 10^{-4}$	CL=90%	DESIG=12
Γ <sub>64</sub>	$\gamma f_2(1270)$	< 2.41	$\times 10^{-4}$	CL=90%	DESIG=11
Γ <sub>65</sub>	$\gamma f_J(2220)$				DESIG=14
Γ <sub>66</sub>	$\gamma \eta_c(1S)$	< 2.7	$\times 10^{-5}$	CL=90%	DESIG=111
Γ <sub>67</sub>	$\gamma \chi_{c0}$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=112
Γ <sub>68</sub>	$\gamma \chi_{c1}$	< 3.6	$\times 10^{-6}$	CL=90%	DESIG=113
Γ <sub>69</sub>	$\gamma \chi_{c2}$	< 1.5	$\times 10^{-5}$	CL=90%	DESIG=114
Γ <sub>70</sub>	$\gamma \chi_{c1}(3872)$	< 2.1	$\times 10^{-5}$	CL=90%	DESIG=172
Γ <sub>71</sub>	$\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi$	< 2.4	$\times 10^{-6}$	CL=90%	DESIG=116
Γ <sub>72</sub>	$\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi$	< 2.8	$\times 10^{-6}$	CL=90%	DESIG=117
Γ <sub>73</sub>	$\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi$	< 1.2	$\times 10^{-6}$	CL=90%	DESIG=118
Γ <sub>74</sub>	$\gamma X(4350) \rightarrow \phi J/\psi$	< 1.3	$\times 10^{-6}$	CL=90%	DESIG=119
Γ <sub>75</sub>	$\gamma \eta_b(1S)$	$(5.5^{+1.1}_{-0.9}) \times 10^{-4}$		S=1.2	DESIG=102
Γ <sub>76</sub>	$\gamma \eta_b(1S) \rightarrow \gamma$ Sum of 26 exclusive modes	< 3.7	$\times 10^{-6}$	CL=90%	DESIG=124
Γ <sub>77</sub>	$\gamma X_{b\bar{b}} \rightarrow \gamma$ Sum of 26 exclusive modes	< 4.9	$\times 10^{-6}$	CL=90%	DESIG=125
Γ <sub>78</sub>	$\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 1.95	$\times 10^{-4}$	CL=95%	DESIG=103
Γ <sub>79</sub>	$\gamma A^0 \rightarrow \gamma$ hadrons	< 8	$\times 10^{-5}$	CL=90%	DESIG=108
Γ <sub>80</sub>	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	< 8.3	$\times 10^{-6}$	CL=90%	DESIG=123

Lepton Family number (*LF*) violating modes

$\Gamma_{81}$	$e^\pm \tau^\mp$	<i>LF</i>	< 3.2	$\times 10^{-6}$	CL=90%
$\Gamma_{82}$	$\mu^\pm \tau^\mp$	<i>LF</i>	< 3.3	$\times 10^{-6}$	CL=90%

NODE=M052;CLUMP=B

DESIG=107

DESIG=104

[a] 1.5 GeV <  $m_\chi$  < 5.0 GeV

LINKAGE=C52

## CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a  $\chi^2 = 11.8$  for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

$$x_7 \left| \begin{array}{c} 2 \\ \hline x_1 \end{array} \right.$$

 $\Upsilon(2S)$   $\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$ 

NODE=M052218

$$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_5/\Gamma$$

NODE=M052G1  
NODE=M052G1

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<b>6.5±1.5±1.0</b>	KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$

$$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_5/\Gamma$$

NODE=M052G03  
NODE=M052G03

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>105.4±1.0±4.2</b>	11.8k	<sup>1</sup> AUBERT	08BP BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
<sup>1</sup> Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .				

NODE=M052G03;LINKAGE=AU

$$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_5/\Gamma$$

NODE=M052G2  
NODE=M052G2

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.577±0.009 OUR AVERAGE</b>			
0.581±0.004±0.009	<sup>1</sup> ROSNER	06	CLEO $e^+e^- \rightarrow \text{hadrons}$
0.552±0.031±0.017	<sup>1</sup> BARU	96	MD1 $e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.04 ±0.02	<sup>1</sup> JAKUBOWSKI	88	CBAL $e^+e^- \rightarrow \text{hadrons}$
0.58 ±0.03 ±0.04	<sup>2</sup> GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$
0.60 ±0.12 ±0.07	<sup>2</sup> ALBRECHT	82	DASP $e^+e^- \rightarrow \text{hadrons}$
0.54 ±0.07 $^{+0.09}_{-0.05}$	<sup>2</sup> NICZYPORUK	81C	LENA $e^+e^- \rightarrow \text{hadrons}$
0.41 ±0.18	<sup>2</sup> BOCK	80	CNTR $e^+e^- \rightarrow \text{hadrons}$

<sup>1</sup> Radiative corrections evaluated following KURAEV 85.<sup>2</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.NODE=M052G2;LINKAGE=P  
NODE=M052G2;LINKAGE=R $\Upsilon(2S)$  PARTIAL WIDTHS

NODE=M052220

$$\Gamma(e^+e^-) \quad \Gamma_5$$

NODE=M052W2  
NODE=M052W2

VALUE (keV)	DOCUMENT ID
<b>0.612±0.011 OUR EVALUATION</b>	

→ UNCHECKED ←

 $\Upsilon(2S)$  BRANCHING RATIOS

NODE=M052225

$$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

NODE=M052R4

NODE=M052R4  
NODE=M052R4

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>17.85±0.26 OUR FIT</b>				
<b>17.92±0.26 OUR AVERAGE</b>				
16.8 ±1.1 ±1.3	906k	<sup>1</sup> LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
17.80±0.05±0.37	170k	<sup>2</sup> LEES	11L BABR	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02±0.02±0.61	851k	<sup>3</sup> BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^- \text{MM}$
17.22±0.17±0.75	11.8k	<sup>4</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ±0.2 ±1.0	52.6k	<sup>5</sup> ALEXANDER	98 CLE2	$\pi^+\pi^-\ell^+\ell^-, \pi^+\pi^- \text{MM}$

18.1 ±0.5 ±1.0	11.6k	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-$	MM
16.9 ±4.0		GELPHMAN	85	CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$	
19.1 ±1.2 ±0.6		BESSION	84	CLEO	$\pi^+\pi^-$	MM
18.9 ±2.6		FONSECA	84	CUSB	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$	
21 ±7	7	NICZYPORUK	81B	LENA	$e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$	

<sup>1</sup>LEES 11c reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M052R4;LINKAGE=ES

<sup>2</sup>Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M052R4;LINKAGE=LE

<sup>3</sup>A weighted average of the inclusive and exclusive results.

NODE=M052R4;LINKAGE=BH

<sup>4</sup>Using  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ ,  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$  and,  $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

NODE=M052R4;LINKAGE=AU

<sup>5</sup>Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

NODE=M052R4;LINKAGE=T

### $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.6 ±0.4 OUR AVERAGE</b>				
8.43±0.16±0.42	38k	<sup>1</sup> BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 ±0.6 ±0.8	275	<sup>2</sup> ALEXANDER	98	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 ±1.9 ±1.9	25	ALBRECHT	87	ARG $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ±1.5		GELPHMAN	85	CBAL $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ±2.3		FONSECA	84	CUSB $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

NODE=M052R5  
NODE=M052R5

<sup>1</sup>Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

NODE=M052R5;LINKAGE=BH

<sup>2</sup>Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$ .

NODE=M052R5;LINKAGE=T

### $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ $\Gamma_2/\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
0.462±0.037	<sup>1</sup> BHARI	09	CLEO $e^+e^- \rightarrow \Upsilon(2S)$

NODE=M052R21  
NODE=M052R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup>Not independent of other values reported by BHARI 09.

NODE=M052R21;LINKAGE=BH

### $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.00±0.21 OUR AVERAGE</b>				
2.00±0.12±0.18	22k	<sup>1</sup> BESSION	07	CLEO $e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 ±1.5 ±0.6		HAAS	84B	CLEO $e^+e^- \rightarrow \tau^+\tau^-$

NODE=M052R3  
NODE=M052R3

<sup>1</sup>BESSION 07 reports  $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M052R3;LINKAGE=BE

### $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.0193±0.0017 OUR AVERAGE</b>					Error includes scale factor of 2.2. See the ideogram below.
0.0203±0.0003±0.0008		120k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
0.0122±0.0028±0.0019			<sup>1</sup> KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$
0.0138±0.0025±0.0015			KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
0.009 ±0.006 ±0.006			<sup>2</sup> ALBRECHT	85	ARG $e^+e^- \rightarrow \mu^+\mu^-$
0.018 ±0.008 ±0.005			HAAS	84B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$

NODE=M052R1  
NODE=M052R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.038 90 NICZYPORUK 81C LENA  $e^+e^- \rightarrow \mu^+\mu^-$


<sup>1</sup>Taking into account interference between the resonance and continuum.

NODE=M052R1;LINKAGE=A

<sup>2</sup>Re-evaluated using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$ .

NODE=M052R1;LINKAGE=R





$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$   $\Gamma_3/\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.04±0.04±0.05</b>	22k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(2S)$

NODE=M052R17  
NODE=M052R17

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{total}$   $\Gamma_6/\Gamma$

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

< 4	90	<sup>1</sup> TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	<sup>2</sup> HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<110	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
<800	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

<sup>1</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0)/\Gamma_{total}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$ .

<sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

NODE=M052R10;LINKAGE=TA

NODE=M052R10;LINKAGE=HE

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$   $\Gamma_6/\Gamma_1$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.3</b>	90	TAMPONI	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

NODE=M052R09  
NODE=M052R09

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{total}$   $\Gamma_7/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

**2.9 ±0.4 OUR FIT** Error includes scale factor of 2.0.

**2.9 ±0.4 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.

2.39±0.31±0.14	112	<sup>1</sup> LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+\ell^-\eta$
2.1 <sup>+0.7</sup> / <sub>-0.6</sub> ±0.3	14	<sup>2</sup> HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$

••• We use the following data for averages but not for fits. •••

3.55±0.32±0.05	241	<sup>3</sup> TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\eta$
----------------	-----	----------------------	----	------	---------------------------------------

NOTFITTED

••• We do not use the following data for averages, fits, limits, etc. •••

< 9	90	<sup>1,4</sup> AUBERT	08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
< 28	90	ALEXANDER	98	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\eta$
< 50	90	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-MM$
< 70	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-MM$
< 20	90	FONSECA	84	CUSB	$e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, \pi^+\pi^-\pi^0)$


- <sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .
- <sup>2</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+ e^-) + B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 4.96\%$ .
- <sup>3</sup> TAMPONI 13 reports  $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (17.85 \pm 0.26) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.
- <sup>4</sup> Using  $\Gamma_{ec}(\Upsilon(2S)) = 0.612 \pm 0.011$  keV.

NODE=M052R6;LINKAGE=AU

NODE=M052R6;LINKAGE=HE

NODE=M052R6;LINKAGE=TA

NODE=M052R6;LINKAGE=UB



**$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$**

**$\Gamma_7/\Gamma_1$**

NODE=M052R22  
NODE=M052R22

VALUE (units $10^{-3}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1.64±0.25 OUR FIT</b>					Error includes scale factor of 2.0.

**1.99±0.14±0.11**    241    TAMPONI 13    BELL     $e^+ e^- \rightarrow \Upsilon(1S)\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.17±0.08    <sup>1</sup> LEES    11L BABR     $\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$   
 < 5.2    90    <sup>2</sup> AUBERT    08BP BABR     $e^+ e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

<sup>1</sup> Not independent of other values reported by LEES 11L.

<sup>2</sup> Not independent of other values reported by AUBERT 08BP.

NODE=M052R22;LINKAGE=LE  
NODE=M052R22;LINKAGE=AU

**$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$**

**$\Gamma_6/\Gamma_7$**

NODE=M052R23  
NODE=M052R23

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.13	90	TAMPONI 13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

**$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$**

**$\Gamma_8/\Gamma$**

NODE=M052R16  
NODE=M052R16

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.006	90	MASCHMANN 90	CBAL	$e^+ e^- \rightarrow \text{hadrons}$

**$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$**

**$\Gamma_9/\Gamma$**

NODE=M052R53  
NODE=M052R53

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.4 × 10 <sup>-6</sup>	90	YANG 14	BELL	$e^+ e^- \rightarrow J/\psi X$

**$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$**

**$\Gamma_{10}/\Gamma$**

NODE=M052R54  
NODE=M052R54

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.4 × 10 <sup>-6</sup>	90	YANG 14	BELL	$e^+ e^- \rightarrow J/\psi X$

**$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$**

**$\Gamma_{11}/\Gamma$**

NODE=M052R55  
NODE=M052R55

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2 × 10 <sup>-6</sup>	90	YANG 14	BELL	$e^+ e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$					$\Gamma_{12}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R56 NODE=M052R56
$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_{13}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R57 NODE=M052R57
$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$					$\Gamma_{14}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R58 NODE=M052R58
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					$\Gamma_{15}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.0 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R59 NODE=M052R59
$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{16}/\Gamma$	
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.24 \pm 0.44 \pm 0.20$	376	JIA	17	BELL	$\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$	NODE=M052R00 NODE=M052R00
$\Gamma(\chi_{c1}(1P)^0 X_{\text{tetra}})/\Gamma_{\text{total}}$					$\Gamma_{17}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<36.7 \times 10^{-6}$	90	<sup>1</sup> JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M052R69 NODE=M052R69
<sup>1</sup> For a tetraquark state $X_{\text{tetra}}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of $X_{\text{tetra}}$ mass and width range from $4.4 \times 10^{-6}$ to $36.7 \times 10^{-6}$ .						
$\Gamma(\chi_{c2} \text{ anything})/\Gamma_{\text{total}}$					$\Gamma_{18}/\Gamma$	
<u>VALUE (units <math>10^{-4}</math>)</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$2.28 \pm 0.73 \pm 0.34$		JIA	17	BELL	$\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$	NODE=M052R67 NODE=M052R67
$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$					$\Gamma_{19}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<5.1 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R60 NODE=M052R60
$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$					$\Gamma_{20}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<4.7 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R61 NODE=M052R61
$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$					$\Gamma_{21}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<2.5 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R62 NODE=M052R62
$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$					$\Gamma_{22}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<1.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R63 NODE=M052R63
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$					$\Gamma_{23}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<3.3 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R64 NODE=M052R64
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$					$\Gamma_{24}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R65 NODE=M052R65
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$					$\Gamma_{25}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R66 NODE=M052R66
$\Gamma(Z_c(3900)^+ Z_c(3900)^-)/\Gamma_{\text{total}}$					$\Gamma_{26}/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
$<1.0 \times 10^{-6}$	90	<sup>1</sup> JIA	18	BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$	NODE=M052R71 NODE=M052R71
<sup>1</sup> Assuming $B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$ .						
						NODE=M052R71;LINKAGE=A

$\Gamma(Z_c(4200)^+ Z_c(4200)^-)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<16.7 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

<sup>1</sup> Assuming  $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$

NODE=M052R72  
NODE=M052R72

NODE=M052R72;LINKAGE=A

 $\Gamma(Z_c(3900)^\pm Z_c(4200)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$

<sup>1</sup> Assuming  $B(Z_c(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(Z_c(3900)^\pm \rightarrow J/\psi \pi^\pm)$ .

NODE=M052R73  
NODE=M052R73

NODE=M052R73;LINKAGE=A

 $\Gamma(X(4050)^+ X(4050)^-)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<13.5 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

<sup>1</sup> Assuming  $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

NODE=M052R74  
NODE=M052R74

NODE=M052R74;LINKAGE=A

 $\Gamma(X(4250)^+ X(4250)^-)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<26.7 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

<sup>1</sup> Assuming  $B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$

NODE=M052R75  
NODE=M052R75

NODE=M052R75;LINKAGE=A

 $\Gamma(X(4050)^\pm X(4250)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<27.2 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$

<sup>1</sup> Assuming  $B(X(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(X(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$

NODE=M052R76  
NODE=M052R76

NODE=M052R76;LINKAGE=A

 $\Gamma(Z_c(4430)^+ Z_c(4430)^-)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$

<sup>1</sup> Assuming  $B(Z_c(4430)^\pm \rightarrow \psi(2P) \pi^\pm) = 1$

NODE=M052R77  
NODE=M052R77

NODE=M052R77;LINKAGE=A

 $\Gamma(X(4055)^\pm X(4055)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<11.1 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$

<sup>1</sup> Assuming  $B(X(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$

NODE=M052R79  
NODE=M052R79

NODE=M052R79;LINKAGE=A

 $\Gamma(X(4055)^\pm Z_c(4430)^\mp)/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<21.1 \times 10^{-6}$	90	1 JIA	18 BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$

<sup>1</sup> Assuming  $B(X(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(Z_c(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$

NODE=M052R80  
NODE=M052R80

NODE=M052R80;LINKAGE=A

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{35}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.78^{+0.30}_{-0.26}$				<b>OUR AVERAGE</b> Error includes scale factor of 1.2.

$2.64 \pm 0.11^{+0.26}_{-0.21}$  LEES 14G BABR  $e^+ e^- \rightarrow \overline{2H} X$

$3.37 \pm 0.50 \pm 0.25$  58 ASNER 07 CLEO  $e^+ e^- \rightarrow \overline{2H} X$

NODE=M052R18  
NODE=M052R18

 $\Gamma(g g g)/\Gamma_{\text{total}}$   $\Gamma_{37}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$58.8 \pm 1.2$	6M	1 BESSON	06A CLEO	$\Upsilon(2S) \rightarrow \text{hadrons}$

<sup>1</sup> Calculated using the value  $\Gamma(\gamma g g)/\Gamma(g g g) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$  from BESSON 06A and PDG 08 values of  $B(\pi^+ \pi^- \Upsilon(1S)) = (18.1 \pm 0.4)\%$ ,  $B(\pi^0 \pi^0 \Upsilon(1S)) = (8.6 \pm 0.4)\%$ ,  $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$ , and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with that of  $\Gamma(\gamma g g)/\Gamma_{\text{total}}$  measurement of BESSON 06A.

NODE=M052R01  
NODE=M052R01

NODE=M052R01;LINKAGE=BE

 $\Gamma(\gamma g g)/\Gamma(g g g)$   $\Gamma_{38}/\Gamma_{37}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.18 \pm 0.04 \pm 0.47$	6M	BESSON	06A CLEO	$\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$

NODE=M052R03  
NODE=M052R03

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$   $\Gamma_{39}/\Gamma$ 

VALUE (units $10^{-6}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.58 \pm 0.33 \pm 0.18$	58	SHEN	12A BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$

NODE=M052R43  
NODE=M052R43

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_{40}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<2.58	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$	NODE=M052R44 NODE=M052R44
$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{41}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
2.32±0.40±0.54	135	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	NODE=M052R45 NODE=M052R45
$\Gamma(\phi f_2'(1525))/\Gamma_{\text{total}}$					$\Gamma_{42}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.33	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(K^+ K^-)$	NODE=M052R46 NODE=M052R46
$\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$					$\Gamma_{43}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.57	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$	NODE=M052R47 NODE=M052R47
$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$					$\Gamma_{44}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.88	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$	NODE=M052R48 NODE=M052R48
$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{45}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.53±0.52±0.19	32	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	NODE=M052R49 NODE=M052R49
$\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$					$\Gamma_{46}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<3.22	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	NODE=M052R50 NODE=M052R50
$\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$					$\Gamma_{47}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.83	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$	NODE=M052R51 NODE=M052R51
$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$					$\Gamma_{48}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.40	90	SHEN	12A	BELL	$\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$	NODE=M052R52 NODE=M052R52
$\Gamma(\rho\pi)/\Gamma_{\text{total}}$					$\Gamma_{49}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.16	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0$	NODE=M052R27 NODE=M052R27
$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{50}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<0.80	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0$	NODE=M052R28 NODE=M052R28
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{51}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.63	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$	NODE=M052R29 NODE=M052R29
$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$					$\Gamma_{52}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
13.0±1.9±2.1	261 ± 37	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$	NODE=M052R30 NODE=M052R30
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{53}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	
1.14±0.30±0.13	40 ± 10	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	NODE=M052R40 NODE=M052R40
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<3.2	90	<sup>1</sup> DOBBS	12A		$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	
<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.						
$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$					$\Gamma_{54}/\Gamma$	
<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<4.22	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$	NODE=M052R41 NODE=M052R41

$\Gamma(K^*(892)^- K^+ + c.c.)/\Gamma_{\text{total}}$  $\Gamma_{55}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.45	90	SHEN	13	BELL $\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$

NODE=M052R42  
NODE=M052R42 $\Gamma(f_1(1285)\text{anything})/\Gamma_{\text{total}}$  $\Gamma_{56}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.20 \pm 1.50 \pm 0.63$	2.9k	JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

NODE=M052R68  
NODE=M052R68 $\Gamma(f_1(1285)X_{\text{tetra}})/\Gamma_{\text{total}}$  $\Gamma_{57}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< $64.7 \times 10^{-6}$	90	<sup>1</sup> JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

NODE=M052R70  
NODE=M052R70

<sup>1</sup> For a tetraquark state  $X_{\text{tetra}}$ , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of  $X_{\text{tetra}}$  mass and width range from  $7.8 \times 10^{-6}$  to  $64.7 \times 10^{-6}$ .

NODE=M052R70;LINKAGE=A

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$  $\Gamma_{58}/\Gamma$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	COMMENT
$0.29 \pm 0.03$	<sup>1,2</sup> DOBBS 12A	$\Upsilon(2S) \rightarrow \text{hadrons}$

NODE=M052R08  
NODE=M052R08

<sup>1</sup> DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

NODE=M052R08;LINKAGE=DO

<sup>2</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M052R08;LINKAGE=NC

 $\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{\text{total}}$  $\Gamma_{59}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.069 \pm 0.004$ OUR AVERAGE				
$0.0693 \pm 0.0012 \pm 0.0041$	407k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
$0.069 \pm 0.005 \pm 0.009$		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
$0.091 \pm 0.018 \pm 0.022$		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
$0.065 \pm 0.007 \pm 0.012$		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
$0.080 \pm 0.017 \pm 0.016$		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
$0.059 \pm 0.014$		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

NODE=M052R8  
NODE=M052R8 $\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{\text{total}}$  $\Gamma_{60}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.0715 \pm 0.0035$ OUR AVERAGE				
$0.0724 \pm 0.0011 \pm 0.0040$	410k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
$0.074 \pm 0.005 \pm 0.008$		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
$0.098 \pm 0.021 \pm 0.024$		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
$0.058 \pm 0.007 \pm 0.010$		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
$0.102 \pm 0.018 \pm 0.021$		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
$0.061 \pm 0.014$		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

NODE=M052R7  
NODE=M052R7 $\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{\text{total}}$  $\Gamma_{61}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.038 \pm 0.004$ OUR AVERAGE				
$0.0375 \pm 0.0012 \pm 0.0047$	198k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
$0.034 \pm 0.005 \pm 0.006$		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma\chi(1P)$
$0.064 \pm 0.014 \pm 0.016$		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
$0.036 \pm 0.008 \pm 0.009$		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
$0.044 \pm 0.023 \pm 0.009$		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.035 \pm 0.014$		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

NODE=M052R9  
NODE=M052R9 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$  $\Gamma_{62}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<59	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 5.9	90	<sup>2</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M052R13  
NODE=M052R13

<sup>1</sup> Re-evaluated assuming  $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$ .

NODE=M052R13;LINKAGE=M

<sup>2</sup> Includes unknown branching ratio of  $f_0(1710) \rightarrow \pi^+ \pi^-$ .

NODE=M052R13;LINKAGE=N

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$  $\Gamma_{63}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<53	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$

NODE=M052R12  
NODE=M052R12

<sup>1</sup> Re-evaluated assuming  $B(f'_2(1525) \rightarrow K\bar{K}) = 0.71$ .

NODE=M052R12;LINKAGE=L

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<24.1	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$

<sup>1</sup> Using  $B(f_2(1270) \rightarrow \pi \pi) = 0.84$ .

NODE=M052R11  
NODE=M052R11

NODE=M052R11;LINKAGE=K

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6.8	90	<sup>1</sup> ALBRECHT 89	ARG	$\Upsilon(2S) \rightarrow \gamma K^+ K^-$

<sup>1</sup> Includes unknown branching ratio of  $f_J(2220) \rightarrow K^+ K^-$ .

NODE=M052R14  
NODE=M052R14

NODE=M052R14;LINKAGE=S

 $\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.7 $\times 10^{-5}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R31  
NODE=M052R31

 $\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$   $\Gamma_{67}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.0 $\times 10^{-4}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R32  
NODE=M052R32

 $\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$   $\Gamma_{68}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.6 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R33  
NODE=M052R33

 $\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$   $\Gamma_{69}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.5 $\times 10^{-5}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R34  
NODE=M052R34

 $\Gamma(\gamma \chi_{c1}(3872))/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.1 $\times 10^{-5}$	90	<sup>1</sup> WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

<sup>1</sup> WANG 11B reports  $[\Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 0.8 \times 10^{-6}$  which we divide by our best value  $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 3.8 \times 10^{-2}$ .

NODE=M052R81  
NODE=M052R81

NODE=M052R81;LINKAGE=A

 $\Gamma(\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{71}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.4 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R36  
NODE=M052R36

 $\Gamma(\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{72}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.8 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R37  
NODE=M052R37

 $\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{73}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R38  
NODE=M052R38

 $\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$   $\Gamma_{74}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 $\times 10^{-6}$	90	WANG 11B	BELL	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M052R39  
NODE=M052R39

 $\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$   $\Gamma_{75}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.5<sup>+1.1</sup><sub>-0.9</sub></b>					<b>OUR AVERAGE</b> Error includes scale factor of 1.2.
6.1 <sup>+0.6+0.9</sup> <sub>-0.7-0.6</sub>		29k	FULSOM 18	BELL	$\Upsilon(2S) \rightarrow \gamma X$
3.9 $\pm$ 1.1 <sup>+1.1</sup> <sub>-0.9</sub>		13 $\pm$ 5k	<sup>1</sup> AUBERT 09AQ	BABR	$\Upsilon(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21 90 LEES 11J BABR  $\Upsilon(2S) \rightarrow X \gamma$

< 8.4 90 <sup>1</sup> BONVICINI 10 CLEO  $\Upsilon(2S) \rightarrow \gamma X$

< 5.1 90 <sup>2</sup> ARTUSO 05 CLEO  $e^+ e^- \rightarrow \gamma X$

<sup>1</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10$  MeV.

<sup>2</sup> Superseded by BONVICINI 10.

NODE=M052R15  
NODE=M052R15

NODE=M052R15;LINKAGE=BO  
NODE=M052R15;LINKAGE=SU

$\Gamma(\gamma\eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{76}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.7 \times 10^{-6}$	90	SANDILYA 13	BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons

NODE=M052R25  
 NODE=M052R25

 $\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$   $\Gamma_{77}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<4.9$	90		SANDILYA 13	BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons

NODE=M052R26  
 NODE=M052R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

$46.2^{+29.7}_{-14.2} \pm 10.6$	10	<sup>1</sup> DOBBS 12			$\Upsilon(2S) \rightarrow \gamma$ hadrons
---------------------------------	----	-----------------------	--	--	---

<sup>1</sup> Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M052R26;LINKAGE=DO

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{78}/\Gamma$ 

(1.5 GeV  $< m_X < 5.0$  GeV)

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.95$	95	ROSNER 07A	CLEO	$e^+e^- \rightarrow \gamma X$

NODE=M052R19

NODE=M052R19  
 NODE=M052R19

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_{79}/\Gamma$ 

(0.3 GeV  $< m_{A^0} < 7$  GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-5}$	90	<sup>1</sup> LEES 11H	BABR	$\Upsilon(2S) \rightarrow \gamma$ hadrons

NODE=M052R06

NODE=M052R06  
 NODE=M052R06

<sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $1 \times 10^{-6}$  to  $8 \times 10^{-5}$ .

NODE=M052R06;LINKAGE=LE

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{80}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3$	90	<sup>1</sup> AUBERT 09Z	BABR	$e^+e^- \rightarrow A^0 \rightarrow \gamma \mu^+ \mu^-$

NODE=M052R24  
 NODE=M052R24

<sup>1</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with mass in the range 212–9300 MeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{A^0}$  range from 0.26–8.3  $\times 10^{-6}$ .

NODE=M052R24;LINKAGE=AU

## LEPTON FAMILY NUMBER (LF) VIOLATING MODES

NODE=M052230

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{81}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2$	90	LEES 10B	BABR	$e^+e^- \rightarrow e^\pm \tau^\mp$

NODE=M052R04  
 NODE=M052R04

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{82}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3$	90	LEES 10B	BABR	$e^+e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M052R20  
 NODE=M052R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<14.4$	95	LOVE 08A	CLEO	$e^+e^- \rightarrow \mu^\pm \tau^\mp$
---------	----	----------	------	---------------------------------------

 $\Upsilon(2S)$  Cross-Particle Branching Ratios

NODE=M052240

 $B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.78 \pm 0.02 \pm 0.11$	906k	LEES 11C	BABR	$e^+e^- \rightarrow \pi^+ \pi^- X$

NODE=M052R05  
 NODE=M052R05

 $\Upsilon(2S)$  REFERENCES

NODE=M052

SHAMOV 23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)	REFID=62012
FULSOM 18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)	REFID=59535
JIA 18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58949
JIA 17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=57635
JIA 17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
LEES 14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
YANG 14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345
SANDILYA 13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)	REFID=55590
SHEN 13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395
TAMPONI 13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=54919
DOBBS 12	PRL 109 082001	S. Dobbs <i>et al.</i>		REFID=54288
DOBBS 12A	PR D86 052003	S. Dobbs <i>et al.</i>		REFID=54746
SHEN 12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314
LEES 11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES 11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
LEES 11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
LEES 11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53938
WANG 11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=53939



BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding, World Scientific, Singapore					
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC	REFID=40742
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)	REFID=40021
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22334
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)	REFID=22336
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>		REFID=22327; ERROR=9
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)	REFID=22329
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22332
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22319
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22318
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264

$\Upsilon_2(1D)$ 

$$I^G(J^{PC}) = 0^-(2^{--})$$

was  $\Upsilon(1D)$ 

First observed by BONVICINI 04 in the decay to  $\gamma\gamma \Upsilon(1S)$  and confirmed by DEL-AMO-SANCHEZ 10R in the decay to  $\pi^+\pi^- \Upsilon(1S)$ .

Data consistent with  $J^P = 2^-$ . The states with  $J = 1$  and 3 also possibly seen, but need confirmation.

NODE=M177

NODE=M177

 **$\Upsilon_2(1D)$  MASS**

NODE=M177M

NODE=M177M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10163.7 ± 1.4 OUR AVERAGE</b>				Error includes scale factor of 1.7.
10164.5 ± 0.8 ± 0.5		DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
10161.1 ± 0.6 ± 1.6	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

 **$\Upsilon_2(1D)$  DECAY MODES**

NODE=M177215;NODE=M177

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \gamma\gamma \Upsilon(1S)$	seen
$\Gamma_2 \quad \gamma\chi_{bJ}(1P)$	seen
$\Gamma_3 \quad \eta \Upsilon(1S)$	not seen
$\Gamma_4 \quad \pi^+\pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$

DESIG=1;OUR EVAL;→ UNCHECKED ←  
DESIG=2;OUR EVAL;→ UNCHECKED ←  
DESIG=3;OUR EVAL;→ UNCHECKED ←  
DESIG=4

 **$\Upsilon_2(1D)$  BRANCHING RATIOS**

NODE=M177225

$\Gamma(\eta \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$					$\Gamma_3/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;0.25</b>	90	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	

NODE=M177R01  
NODE=M177R01

$\Gamma(\pi^+\pi^- \Upsilon(1S))/\Gamma_{\text{total}}$					$\Gamma_4/\Gamma$
VALUE (units $10^{-2}$ )		DOCUMENT ID	TECN	COMMENT	
<b>0.66<sup>+0.15</sup><sub>-0.14</sub> ± 0.06</b>		<sup>1</sup> DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$	

NODE=M177R03  
NODE=M177R03

<sup>1</sup> Using theoretical predictions for  $B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon_2(1D))$ .

NODE=M177R03;LINKAGE=DE

$\Gamma(\pi^+\pi^- \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$					$\Gamma_4/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<b>&lt;1.2</b>	90	<sup>2</sup> BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$	

NODE=M177R02  
NODE=M177R02

<sup>2</sup> Assuming  $J = 2$ .

NODE=M177R02;LINKAGE=BO

 **$\Upsilon_2(1D)$  REFERENCES**

NODE=M177

DEL-AMO-SA... 10R PR D82 111102 P. del Amo Sanchez *et al.* (BABAR Collab.)  
BONVICINI 04 PR D70 032001 G. Bonvicini *et al.* (CLEO Collab.)

REFID=53634  
REFID=49759

$\chi_{b0}(2P)$

$I^G(J^{PC}) = 0^+(0^{++})$   
 J needs confirmation.

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

NODE=M079

NODE=M079

$\chi_{b0}(2P)$  MASS

VALUE (MeV) DOCUMENT ID  
**10232.5 ± 0.4 ± 0.5 OUR EVALUATION** From  $\gamma$  energy below, using  $\Upsilon(3S)$  mass = 10355.2 ± 0.5 MeV

NODE=M079M

NODE=M079M

→ UNCHECKED ←

$m_{\chi_{b1}(2P)} = m_{\chi_{b0}(2P)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>23.8 ± 1.7</b>	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M079M2  
 NODE=M079M2

$\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>121.9 ± 0.4 OUR EVALUATION</b>		Treating systematic errors as correlated		
<b>122.2 ± 0.5 OUR AVERAGE</b>		Error includes scale factor of 1.4. See the ideogram below.		
121.55 ± 0.16 ± 0.46		ARTUSO	05	CLEO $\Upsilon(3S) \rightarrow \gamma X$
123.0 ± 0.8	4959	<sup>1</sup> HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma X$
124.6 ± 1.4	17	<sup>2</sup> HEINTZ	92	CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
122.3 ± 0.3 ± 0.6	9903	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma X$

NODE=M079DM

NODE=M079DM


→ UNCHECKED ←

OCCUR=2

NODE=M079DM;LINKAGE=A

NODE=M079DM;LINKAGE=B

<sup>1</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.  
<sup>2</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.



$\chi_{b0}(2P)$  DECAY MODES

NODE=M079215;NODE=M079

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1 \quad \gamma \Upsilon(2S)$	(1.38 ± 0.30) %	
$\Gamma_2 \quad \gamma \Upsilon(1S)$	(3.8 ± 1.7) × 10 <sup>-3</sup>	
$\Gamma_3 \quad D^0 X$	< 8.2 %	90%
$\Gamma_4 \quad \pi^+\pi^-K^+K^-\pi^0$	< 3.4 × 10 <sup>-5</sup>	90%
$\Gamma_5 \quad 2\pi^+\pi^-K^-K_S^0$	< 5 × 10 <sup>-5</sup>	90%
$\Gamma_6 \quad 2\pi^+\pi^-K^-K_S^0 2\pi^0$	< 2.2 × 10 <sup>-4</sup>	90%
$\Gamma_7 \quad 2\pi^+ 2\pi^- 2\pi^0$	< 2.4 × 10 <sup>-4</sup>	90%
$\Gamma_8 \quad 2\pi^+ 2\pi^- K^+ K^-$	< 1.5 × 10 <sup>-4</sup>	90%

DESIG=2

DESIG=1

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

$\Gamma_9$	$2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.2	$\times 10^{-4}$	90%	DESIG=9
$\Gamma_{10}$	$2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 1.1	$\times 10^{-3}$	90%	DESIG=10
$\Gamma_{11}$	$3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=11
$\Gamma_{12}$	$3\pi^+ 3\pi^-$	< 7	$\times 10^{-5}$	90%	DESIG=12
$\Gamma_{13}$	$3\pi^+ 3\pi^- 2\pi^0$	< 1.2	$\times 10^{-3}$	90%	DESIG=13
$\Gamma_{14}$	$3\pi^+ 3\pi^- K^+ K^-$	< 1.5	$\times 10^{-4}$	90%	DESIG=14
$\Gamma_{15}$	$3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 7	$\times 10^{-4}$	90%	DESIG=15
$\Gamma_{16}$	$4\pi^+ 4\pi^-$	< 1.7	$\times 10^{-4}$	90%	DESIG=16
$\Gamma_{17}$	$4\pi^+ 4\pi^- 2\pi^0$	< 6	$\times 10^{-4}$	90%	DESIG=17

 **$\chi_{b0}(2P)$  BRANCHING RATIOS**

NODE=M079220

 **$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$**  **$\Gamma_1/\Gamma$** 

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

NODE=M079R2  
NODE=M079R2**1.38 ± 0.30 OUR AVERAGE**

1.31 ± 0.27 <sup>+0.13</sup> <sub>-0.12</sub>	3,4	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
---	-----	------	-----	------	--

3.6 ± 1.6 ± 0.3	3,5	HEINTZ	92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
-----------------	-----	--------	----	------	---

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<2.8	90	6	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X \gamma$
------	----	---	------	-----	------	-------------------------------------

<8.9	90	7	CRAWFORD	92B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
------	----	---	----------	-----	------	---

<sup>3</sup> Assuming  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$ .

NODE=M079R2;LINKAGE=D  
NODE=M079R2;LINKAGE=E

<sup>4</sup> LEES 14M reports  $[\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] = (7.7 \pm 1.6) \times 10^{-4}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>5</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$  using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ . Supersedes HEINTZ 91.

NODE=M079R2;LINKAGE=C

<sup>6</sup> LEES 11J quotes a central value of  $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$ .

NODE=M079R2;LINKAGE=LE

<sup>7</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) < 1.19 \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P) \gamma) = 0.049$ .

NODE=M079R2;LINKAGE=B

 **$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$**  **$\Gamma_2/\Gamma$** 

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

NODE=M079R1  
NODE=M079R1**0.38 ± 0.17 OUR AVERAGE**

0.36 ± 0.17 ± 0.03	8,9,10	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
--------------------	--------	------	-----	------	--

0.9 ± 0.7 ± 0.1	9,11	HEINTZ	92	CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
-----------------	------	--------	----	------	---

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<1.2	90	12	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X \gamma$
------	----	----	------	-----	------	-------------------------------------

<2.5	90	13	CRAWFORD	92B	CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
------	----	----	----------	-----	------	---

<sup>8</sup> LEES 14M quotes  $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (2.1 \pm 1.0) \times 10^{-4}$  combining the results from  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$  samples with and without photon conversions.

NODE=M079R1;LINKAGE=D

<sup>9</sup> Assuming  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M079R1;LINKAGE=E

<sup>10</sup> LEES 14M reports  $[\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] = (2.1 \pm 1.0) \times 10^{-4}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M079R1;LINKAGE=F

<sup>11</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$  using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$ . Supersedes HEINTZ 91.

NODE=M079R1;LINKAGE=C

<sup>12</sup> LEES 11J quotes a central value of  $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$ .

NODE=M079R1;LINKAGE=LE

<sup>13</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$ ,  $B(\Upsilon(3S) \rightarrow \gamma \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) < 0.63 \times 10^{-4}$ , and  $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P) \gamma) = 0.049$ .

NODE=M079R1;LINKAGE=B

 **$\Gamma(D^0 X)/\Gamma_{\text{total}}$**  **$\Gamma_3/\Gamma$** 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M079R01  
NODE=M079R01

<b>&lt;8.2 × 10<sup>-2</sup></b>	90	14,15	BRIERE	08	CLEO	$\Upsilon(3S) \rightarrow \gamma D^0 X$
----------------------------------	----	-------	--------	----	------	---

<sup>14</sup> For  $p_{D^0} > 2.5$  GeV/c.

NODE=M079R01;LINKAGE=BR

<sup>15</sup> The authors also present their result as  $(4.1 \pm 3.0 \pm 0.4) \times 10^{-2}$ .

NODE=M079R01;LINKAGE=RI

$$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_4/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<0.34	90	16 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$ 16 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R02  
NODE=M079R02

NODE=M079R02;LINKAGE=AS

$$\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$$

 $\Gamma_5/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	17 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$ 17 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R03  
NODE=M079R03

NODE=M079R03;LINKAGE=AS

$$\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_6/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	18 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$ 18 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R04  
NODE=M079R04

NODE=M079R04;LINKAGE=AS

$$\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_7/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<2.4	90	19 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$ 19 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R05  
NODE=M079R05

NODE=M079R05;LINKAGE=AS

$$\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$$

 $\Gamma_8/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	20 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$ 20 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 9 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R06  
NODE=M079R06

NODE=M079R06;LINKAGE=AS

$$\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_9/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	21 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$ 21 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R07  
NODE=M079R07

NODE=M079R07;LINKAGE=AS

$$\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_{10}/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	22 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$ 22 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 63 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R08  
NODE=M079R08

NODE=M079R08;LINKAGE=AS

$$\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$$

 $\Gamma_{11}/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$ 23 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 39 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R09  
NODE=M079R09

NODE=M079R09;LINKAGE=AS

$$\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$$

 $\Gamma_{12}/\Gamma$ 

VALUE (units 10 <sup>-4</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+3\pi^-$ 24 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P))] < 4 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R10  
NODE=M079R10

NODE=M079R10;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	25 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$ 25 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < $72 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R11  
NODE=M079R11

NODE=M079R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	26 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$ 26 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < $9 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R12  
NODE=M079R12

NODE=M079R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	27 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$ 27 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < $43 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R13  
NODE=M079R13

NODE=M079R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	28 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$ 28 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < $10 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R14  
NODE=M079R14

NODE=M079R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$ 29 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < $38 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$ .

NODE=M079R15  
NODE=M079R15

NODE=M079R15;LINKAGE=AS

 $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}}$   
 $\Gamma_2/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	30 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$ 30 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) < 1.2\%$ using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$ .

NODE=M079B01  
NODE=M079B01

NODE=M079B01;LINKAGE=LE

 $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.4 ± 0.9 OUR AVERAGE</b>			

NODE=M079A02  
NODE=M079A02

$1.7^{+1.5+0.1}_{-1.4-1.2}$	31 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$1.3 \pm 1.0 \pm 0.3$	32 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
31 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.			
32 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$ .			

NODE=M079A02;LINKAGE=A  
NODE=M079A02;LINKAGE=K $[B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ 

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>1.71 ± 0.80</b>	33 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
33 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ without converted photons.			

NODE=M079A00  
NODE=M079A00

NODE=M079A00;LINKAGE=A

 $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}}$   
 $\Gamma_1/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	34 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$ 34 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) < 2.8\%$ using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$ .

NODE=M079B02  
NODE=M079B02

NODE=M079B02;LINKAGE=LE

**$B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$**

VALUE (units  $10^{-5}$ ) DOCUMENT ID TECN COMMENT

**4.4 ± 1.6 OUR AVERAGE**

$6.6^{+4.9+2.0}_{-4.0-0.3}$	35	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$4.0 \pm 1.7 \pm 0.3$	36	HEINTZ	92	CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>35</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with one converted photon.

<sup>36</sup> Calculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$  using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ .

NODE=M079A03  
NODE=M079A03

NODE=M079A03;LINKAGE=A  
NODE=M079A03;LINKAGE=B

**$[B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$**

VALUE (%) DOCUMENT ID TECN COMMENT

**3.31 ± 0.56**

37	LEES	14M	BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
----	------	-----	------	--

<sup>37</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$  without converted photons.

NODE=M079A01  
NODE=M079A01

NODE=M079A01;LINKAGE=A

**$\chi_{b0}(2P)$  REFERENCES**

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

NODE=M079

REFID=56343  
REFID=53936  
REFID=52574  
REFID=52577  
REFID=50454  
REFID=43177  
REFID=43604  
REFID=41580  
REFID=41634  
REFID=41586

**$\chi_{b1}(2P)$**

$J^G(J^{PC}) = 0^+(1^{++})$   
J needs confirmation.

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

NODE=M080

NODE=M080

**$\chi_{b1}(2P)$  MASS**

VALUE (MeV)	DOCUMENT ID
<b>10255.46 ± 0.22 ± 0.50 OUR EVALUATION</b>	From $\gamma$ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

NODE=M080M  
NODE=M080M  
→ UNCHECKED ←

**$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>3.5 ± 0.7 ± 0.7</b>	<sup>1</sup> HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$

<sup>1</sup> From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

NODE=M080M2  
NODE=M080M2  
NODE=M080M2;LINKAGE=A

**$\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>99.26 ± 0.22 OUR EVALUATION</b>		Treating systematic errors as correlated		
<b>99.53 ± 0.23 OUR AVERAGE</b>		Error includes scale factor of 1.3. See the ideogram below.		
$99.15 \pm 0.07 \pm 0.25$		ARTUSO	05	CLEO $\Upsilon(3S) \rightarrow \gamma X$
$99 \pm 1$	169	CRAWFORD	92B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
$100.1 \pm 0.4$	11147	<sup>2</sup> HEINTZ	92	CSB2 $e^+ e^- \rightarrow \gamma X$
$100.2 \pm 0.5$	223	<sup>3</sup> HEINTZ	92	CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
$99.5 \pm 0.1 \pm 0.5$	25759	MORRISON	91	CLE2 $e^+ e^- \rightarrow \gamma X$


NODE=M080DM  
NODE=M080DM  
→ UNCHECKED ←

OCCUR=2

- 2 A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.
- 3 A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M080DM;LINKAGE=A

NODE=M080DM;LINKAGE=B



### $\chi_{b1}(2P)$ DECAY MODES

NODE=M080215;NODE=M080

Mode	Fraction ( $\Gamma_i/\Gamma$ )	
$\Gamma_1$ $\omega \Upsilon(1S)$	$(1.63^{+0.40}_{-0.34})\%$	DESIG=3
$\Gamma_2$ $\gamma \Upsilon(2S)$	$(18.1 \pm 1.9)\%$	DESIG=2
$\Gamma_3$ $\gamma \Upsilon(1S)$	$(9.9 \pm 1.0)\%$	DESIG=1
$\Gamma_4$ $\pi\pi \chi_{b1}(1P)$	$(9.1 \pm 1.3) \times 10^{-3}$	DESIG=4
$\Gamma_5$ $D^0 X$	$(8.8 \pm 1.7)\%$	DESIG=5
$\Gamma_6$ $\pi^+ \pi^- K^+ K^- \pi^0$	$(3.1 \pm 1.0) \times 10^{-4}$	DESIG=6
$\Gamma_7$ $2\pi^+ \pi^- K^- K_S^0$	$(1.1 \pm 0.5) \times 10^{-4}$	DESIG=7
$\Gamma_8$ $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$(7.7 \pm 3.2) \times 10^{-4}$	DESIG=8
$\Gamma_9$ $2\pi^+ 2\pi^- 2\pi^0$	$(5.9 \pm 2.0) \times 10^{-4}$	DESIG=9
$\Gamma_{10}$ $2\pi^+ 2\pi^- K^+ K^-$	$(10 \pm 4) \times 10^{-5}$	DESIG=10
$\Gamma_{11}$ $2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(5.5 \pm 1.8) \times 10^{-4}$	DESIG=11
$\Gamma_{12}$ $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(10 \pm 4) \times 10^{-4}$	DESIG=12
$\Gamma_{13}$ $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$(6.7 \pm 2.6) \times 10^{-4}$	DESIG=13
$\Gamma_{14}$ $3\pi^+ 3\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	DESIG=14
$\Gamma_{15}$ $3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$	DESIG=15
$\Gamma_{16}$ $3\pi^+ 3\pi^- K^+ K^-$	$(2.0 \pm 0.8) \times 10^{-4}$	DESIG=16
$\Gamma_{17}$ $3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(6.1 \pm 2.2) \times 10^{-4}$	DESIG=17
$\Gamma_{18}$ $4\pi^+ 4\pi^-$	$(1.7 \pm 0.6) \times 10^{-4}$	DESIG=18
$\Gamma_{19}$ $4\pi^+ 4\pi^- 2\pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$	DESIG=19

### $\chi_{b1}(2P)$ BRANCHING RATIOS

NODE=M080220

$\Gamma(\omega \Upsilon(1S))/\Gamma_{total}$	$\Gamma_1/\Gamma$
VALUE (units $10^{-2}$ )	
EVTS	DOCUMENT ID
TECN	COMMENT
$1.63^{+0.35+0.16}_{-0.31-0.15}$	
32.6 <sup>+6.9</sup> <sub>-6.1</sub>	4 CRONIN-HEN..04
	CLE3 $\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

NODE=M080R3  
NODE=M080R3

<sup>4</sup> Using  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.3 \pm 0.6)\%$  and  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$ .

NODE=M080R3;LINKAGE=CR



$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.181±0.019 OUR AVERAGE</b>				
0.211±0.017±0.019	5,6,7	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
0.190±0.018±0.017	4.3k	8 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
0.206±0.035±0.019	5,9	CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
0.132±0.018±0.012	5,10	HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M080R2  
 NODE=M080R2

<sup>5</sup> Assuming  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ .

<sup>6</sup> LEES 14M quotes  $\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))/\Gamma_{\text{total}} = (2.66 \pm 0.22)\%$  combining the results from  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  samples with and without photon conversions.

<sup>7</sup> LEES 14M reports  $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (2.66 \pm 0.22) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>8</sup> LEES 11J reports  $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>9</sup> CRAWFORD 92B quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \ell^+\ell^-) = (10.23 \pm 1.20 \pm 1.26) 10^{-4}$ .

<sup>10</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21) \%$  using  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ . Supersedes HEINTZ 91.

NODE=M080R2;LINKAGE=D  
 NODE=M080R2;LINKAGE=E

NODE=M080R2;LINKAGE=F

NODE=M080R2;LINKAGE=LE

NODE=M080R2;LINKAGE=B

NODE=M080R2;LINKAGE=C

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$  $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.099±0.010 OUR AVERAGE</b>				
0.107±0.006±0.010	11,12,13	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
0.098±0.005±0.009	15k	14 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
0.103±0.023±0.009	11,15	CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
0.075±0.010±0.007	11,16	HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M080R1  
 NODE=M080R1

<sup>11</sup> Assuming  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

<sup>12</sup> LEES 14M quotes  $\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))/\Gamma_{\text{total}} = (13.48 \pm 0.72) \times 10^{-3}$  combining the results from samples of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with and without converted photons.

<sup>13</sup> LEES 14M reports  $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (13.48 \pm 0.72) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>14</sup> LEES 11J reports  $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>15</sup> CRAWFORD 92B quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (6.47 \pm 1.12 \pm 0.82) 10^{-4}$ .

<sup>16</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$  using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$ . Supersedes HEINTZ 91.

NODE=M080R1;LINKAGE=D  
 NODE=M080R1;LINKAGE=E

NODE=M080R1;LINKAGE=G

NODE=M080R1;LINKAGE=LE

NODE=M080R1;LINKAGE=B

NODE=M080R1;LINKAGE=F

 $\Gamma(\pi\pi\chi_{b1}(1P))/\Gamma_{\text{total}}$  $\Gamma_4/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.1±1.3 OUR AVERAGE</b>				
9.2±1.1±0.8	31k	17 LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
8.6±2.3±2.1	18	CAWLFIELD	06 CLE3	$\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$

NODE=M080R4  
 NODE=M080R4

<sup>17</sup> LEES 11c measures  $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) \times B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$ . We derive the value assuming  $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) = B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$ .

<sup>18</sup> CAWLFIELD 06 quote  $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$  keV assuming l-spin conservation, no D-wave contribution,  $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$  keV, and  $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$  keV.

NODE=M080R4;LINKAGE=LE

NODE=M080R4;LINKAGE=CA

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.8±1.5±0.8</b>	2243	19 BRIERE	08 CLEO	$\Upsilon(3S) \rightarrow \gamma D^0 X$

NODE=M080R01  
 NODE=M080R01

<sup>19</sup> For  $p_{D^0} > 2.5$  GeV/c.

NODE=M080R01;LINKAGE=BR

$\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.1±1.0±0.3</b>	30	20 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$
<p><sup>20</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(39 \pm 8 \pm 9) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R02  
NODE=M080R02

NODE=M080R02;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.1±0.5±0.1</b>	10	21 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$
<p><sup>21</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(14 \pm 5 \pm 3) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R03  
NODE=M080R03

NODE=M080R03;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>7.7±3.1±0.7</b>	15	22 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$
<p><sup>22</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(97 \pm 30 \pm 26) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R04  
NODE=M080R04

NODE=M080R04;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.9±2.0±0.5</b>	36	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$
<p><sup>23</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(74 \pm 16 \pm 19) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R05  
NODE=M080R05

NODE=M080R05;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.0±0.4±0.1</b>	12	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$
<p><sup>24</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(12 \pm 4 \pm 3) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R06  
NODE=M080R06

NODE=M080R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.5±1.7±0.5</b>	38	25 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$
<p><sup>25</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(69 \pm 13 \pm 17) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R07  
NODE=M080R07

NODE=M080R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>9.6±3.5±0.9</b>	27	26 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$
<p><sup>26</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(121 \pm 29 \pm 33) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R08  
NODE=M080R08

NODE=M080R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.7±2.5±0.6</b>	17	27 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$
<p><sup>27</sup> ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))]</math> = <math>(85 \pm 23 \pm 22) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M080R09  
NODE=M080R09

NODE=M080R09;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.2±0.4±0.1</b>	18	28 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<p>28 ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]</math>  <math>= (15 \pm 4 \pm 3) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) =</math>  <math>(12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is  the systematic error from using our best value.</p>				

NODE=M080R10  
NODE=M080R10

NODE=M080R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12±4±1</b>	44	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
<p>29 ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]</math>  <math>= (150 \pm 30 \pm 40) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) =</math>  <math>(12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error  is the systematic error from using our best value.</p>				

NODE=M080R11  
NODE=M080R11

NODE=M080R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.0±0.7±0.2</b>	16	30 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<p>30 ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]</math>  <math>= (25 \pm 7 \pm 6) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) =</math>  <math>(12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our  second error is the systematic error from using our best value.</p>				

NODE=M080R12  
NODE=M080R12

NODE=M080R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>6.1±2.1±0.6</b>	25	31 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<p>31 ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]</math>  <math>= (77 \pm 17 \pm 21) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) =</math>  <math>(12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our  second error is the systematic error from using our best value.</p>				

NODE=M080R13  
NODE=M080R13

NODE=M080R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.7±0.6±0.2</b>	16	32 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>32 ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]</math>  <math>= (22 \pm 6 \pm 5) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) =</math>  <math>(12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error is  the systematic error from using our best value.</p>				

NODE=M080R14  
NODE=M080R14

NODE=M080R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>19±7±2</b>	41	33 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
<p>33 ASNER 08A reports <math>[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]</math>  <math>= (241 \pm 47 \pm 72) \times 10^{-6}</math> which we divide by our best value <math>B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) =</math>  <math>(12.6 \pm 1.2) \times 10^{-2}</math>. Our first error is their experiment's error and our second error  is the systematic error from using our best value.</p>				

NODE=M080R15  
NODE=M080R15

NODE=M080R15;LINKAGE=AS

 $\chi_{b1}(2P)$  Cross-Particle Branching Ratios

NODE=M080230

$$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \Gamma_{21}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>12.4±0.3±0.6</b>	15k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$


NODE=M080B01  
NODE=M080B01 $B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ NODE=M080A00  
NODE=M080A00

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.9 ±0.4 OUR AVERAGE</b>		Error includes scale factor of 1.9. See the ideogram below.		
3.52 <sup>+0.28</sup> +0.17 -0.27-0.18		34 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
3.24±0.56±0.41	58	35 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
2.34±0.28±0.15		36 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$

- 34 From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with one converted photon.
- 35 CRAWFORD 92B quotes  $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS))$   
 $B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$ .
- 36 Calculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$  using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$ .

NODE=M080A00;LINKAGE=A  
 NODE=M080A00;LINKAGE=C

NODE=M080A00;LINKAGE=B



$$B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$$

(units  $10^{-4}$ )

$$\frac{\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))}{\Gamma_{\text{total}}} \times \frac{\Gamma_{21}^{\Upsilon(3S)}}{\Gamma_{21}^{\Upsilon(3S)}} \times \frac{\Gamma_{21}^{\Upsilon(3S)}}{\Gamma_{\text{total}}^{\Upsilon(3S)}}$$

NODE=M080B02  
 NODE=M080B02

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.4 ± 0.1 ± 0.2</b>	4.3k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

$$B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+\ell^-)$$


NODE=M080A01  
 NODE=M080A01

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.8 ± 0.6 OUR AVERAGE</b>		Error includes scale factor of 1.8. See the ideogram below.		
$4.95^{+0.75}_{-0.70} \pm 1.01 \pm 0.24$		37 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$5.12 \pm 0.60 \pm 0.63$	111	38 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
$3.30 \pm 0.33 \pm 0.20$		39 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

- 37 From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with one converted photon.
- 38 CRAWFORD 92B quotes  $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon(nS))$   
 $B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$ .
- 39 Calculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21)\%$  using  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ .

NODE=M080A01;LINKAGE=A  
 NODE=M080A01;LINKAGE=C

NODE=M080A01;LINKAGE=B



$$B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$$

(units  $10^{-4}$ )

$$B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X)$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.16±0.07±0.12</b>	31k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M080R16  
NODE=M080R16

$$B(\chi_{b2}(2P) \rightarrow \rho X + \bar{\rho} X) / B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.109±0.007±0.040</b>	BRIERE	07 CLEO	$\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

NODE=M080R20  
NODE=M080R20

$$B(\chi_{b0}(2P) \rightarrow \rho X + \bar{\rho} X) / B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.082±0.025±0.060</b>	BRIERE	07 CLEO	$\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)$

NODE=M080R21  
NODE=M080R21

### $\chi_{b1}(2P)$ REFERENCES

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=51887
CRAWFIELD	06	PR D73 012003	C. Cawfield <i>et al.</i>	(CLEO Collab.)	REFID=50997
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
CRONIN-HEN...	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=49766
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)	REFID=43177
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586

NODE=M080

**$h_b(2P)$** 

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Quantum numbers are quark model predictions.  $C = -$  established by  $\eta_b\gamma$  decay.

NODE=M205

NODE=M205

 **$h_b(2P)$  MASS**

NODE=M205M

NODE=M205M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10259.8±0.5±1.1</b>	90k	<sup>1</sup> MIZUK	12 BELL	$e^+e^- \rightarrow \pi^+\pi^-$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
10259.8±0.6 <sup>+1.4</sup> <sub>-1.0</sub>	83.9k	<sup>2</sup> ADACHI	12 BELL	10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM

<sup>1</sup> Observed with 9 standard deviations significance.

<sup>2</sup> Superseded by MIZUK 12.

NODE=M205M;LINKAGE=A  
NODE=M205M;LINKAGE=AD

 **$h_b(2P)$  DECAY MODES**

NODE=M205215;NODE=M205

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ hadrons	not seen
$\Gamma_2$ $\eta_b(1S)\gamma$	(22±5) %
$\Gamma_3$ $\eta_b(2S)\gamma$	(48±13) %

DESIG=1  
DESIG=2  
DESIG=3

 **$h_b(2P)$  BRANCHING RATIOS**

NODE=M205225

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	83.9k	ADACHI	12 BELL	10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM

NODE=M205R01  
NODE=M205R01

$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$				$\Gamma_2/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>22.3±3.8<sup>+3.1</sup><sub>-3.3</sub></b>	10k	MIZUK	12 BELL	$e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons

NODE=M205R02  
NODE=M205R02

$\Gamma(\eta_b(2S)\gamma)/\Gamma_{\text{total}}$				$\Gamma_3/\Gamma$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>47.5±10.5<sup>+6.8</sup><sub>-7.7</sub></b>	26k	MIZUK	12 BELL	$e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons

NODE=M205R03  
NODE=M205R03

 **$h_b(2P)$  REFERENCES**

NODE=M205

ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

REFID=53962  
REFID=54718

$\chi_{b2}(2P)$ 

$$J^G(J^{PC}) = 0^+(2^{++})$$

$J$  needs confirmation.

Observed in radiative decay of the  $\Upsilon(3S)$ , therefore  $C = +$ . Branching ratio requires E1 transition, M1 is strongly disfavored, therefore  $P = +$ .

NODE=M081

NODE=M081

 $\chi_{b2}(2P)$  MASS

NODE=M081M

VALUE (MeV)

DOCUMENT ID

**10268.65 ± 0.22 ± 0.50 OUR EVALUATION** From  $\Upsilon(3S)$  mass = 10355.2 ± 0.5 MeV

NODE=M081M

→ UNCHECKED ←

 $m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$ 

NODE=M081M2

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

**13.10 ± 0.24 OUR AVERAGE**

12.3 ± 2.6 ± 0.6

<sup>1</sup> AAIJ

14BG LHCB

 $pp \rightarrow \gamma \mu^+ \mu^- X$ 

13.04 ± 0.26

LEES

14M BABR

 $\Upsilon(3S) \rightarrow \gamma \mu^+ \mu^-$ 

13.5 ± 0.4 ± 0.5

<sup>2</sup> HEINTZ

92 CSB2

 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$ <sup>1</sup> From the  $\chi_{bj}(2P) \rightarrow \Upsilon(1S) \gamma$  transition.<sup>2</sup> From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

NODE=M081M2

NODE=M081M2;LINKAGE=B

NODE=M081M2;LINKAGE=A

 $\gamma$  ENERGY IN  $\Upsilon(3S)$  DECAY

NODE=M081DM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

**86.19 ± 0.22 OUR EVALUATION**

Treating systematic errors as correlated

**86.40 ± 0.18 OUR AVERAGE**

86.04 ± 0.06 ± 0.27

ARTUSO

05

CLEO

 $\Upsilon(3S) \rightarrow \gamma X$ 

86 ± 1

101

CRAWFORD

92B

CLE2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ 

86.7 ± 0.4

10319

<sup>3</sup> HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \gamma X$ 

86.9 ± 0.4

157

<sup>4</sup> HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$ 

86.4 ± 0.1 ± 0.4

30741

MORRISON

91

CLE2

 $e^+ e^- \rightarrow \gamma X$ <sup>3</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.<sup>4</sup> A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M081DM

NODE=M081DM

→ UNCHECKED ←

OCCUR=2

NODE=M081DM;LINKAGE=A

NODE=M081DM;LINKAGE=B

 $\chi_{b2}(2P)$  DECAY MODES

NODE=M081215;NODE=M081

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $\omega \Upsilon(1S)$	$(1.10^{+0.34}_{-0.30}) \%$	DESIG=3
$\Gamma_2$ $\gamma \Upsilon(2S)$	$(8.9 \pm 1.2) \%$	DESIG=2
$\Gamma_3$ $\gamma \Upsilon(1S)$	$(6.6 \pm 0.8) \%$	DESIG=1
$\Gamma_4$ $\pi \pi \chi_{b2}(1P)$	$(5.1 \pm 0.9) \times 10^{-3}$	DESIG=4
$\Gamma_5$ $D^0 X$	$< 2.4 \%$	90% DESIG=5
$\Gamma_6$ $\pi^+ \pi^- K^+ K^- \pi^0$	$< 1.1 \times 10^{-4}$	90% DESIG=6
$\Gamma_7$ $2\pi^+ \pi^- K^- K_S^0$	$< 9 \times 10^{-5}$	90% DESIG=7
$\Gamma_8$ $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	$< 7 \times 10^{-4}$	90% DESIG=8
$\Gamma_9$ $2\pi^+ 2\pi^- 2\pi^0$	$(3.9 \pm 1.6) \times 10^{-4}$	DESIG=9
$\Gamma_{10}$ $2\pi^+ 2\pi^- K^+ K^-$	$(9 \pm 4) \times 10^{-5}$	DESIG=10
$\Gamma_{11}$ $2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(2.4 \pm 1.1) \times 10^{-4}$	DESIG=11
$\Gamma_{12}$ $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(4.7 \pm 2.3) \times 10^{-4}$	DESIG=12
$\Gamma_{13}$ $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$< 4 \times 10^{-4}$	90% DESIG=13
$\Gamma_{14}$ $3\pi^+ 3\pi^-$	$(9 \pm 4) \times 10^{-5}$	DESIG=14
$\Gamma_{15}$ $3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$	DESIG=15
$\Gamma_{16}$ $3\pi^+ 3\pi^- K^+ K^-$	$(1.4 \pm 0.7) \times 10^{-4}$	DESIG=16
$\Gamma_{17}$ $3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(4.2 \pm 1.7) \times 10^{-4}$	DESIG=17
$\Gamma_{18}$ $4\pi^+ 4\pi^-$	$(9 \pm 5) \times 10^{-5}$	DESIG=18
$\Gamma_{19}$ $4\pi^+ 4\pi^- 2\pi^0$	$(1.3 \pm 0.5) \times 10^{-3}$	DESIG=19

$\chi_{b2}(2P)$  BRANCHING RATIOS

NODE=M081220

 $\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.10<sup>+0.32+0.11</sup><sub>-0.28-0.10</sub></b>	20.1 <sup>+5.8</sup> <sub>-5.1</sub>	5 CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

NODE=M081R3  
NODE=M081R3

<sup>5</sup> Using  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$  and  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2$   
 $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$ .

NODE=M081R3;LINKAGE=CR

 $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.089<math>\pm</math>0.012 OUR AVERAGE</b>				
0.085 $\pm$ 0.010 $\pm$ 0.010	6,7,8	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.084 $\pm$ 0.011 $\pm$ 0.010	2.5k	<sup>9</sup> LEES	11J	BABR $\Upsilon(3S) \rightarrow X \gamma$
0.096 $\pm$ 0.022 $\pm$ 0.012	7,10	CRAWFORD	92B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
0.106 $\pm$ 0.016 $\pm$ 0.013	7,11	HEINTZ	92	CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

NODE=M081R2  
NODE=M081R2

<sup>6</sup> LEES 14M quotes  $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$   
 $= (1.12 \pm 0.13)\%$  combining the results from samples of  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$  with and  
without converted photons.

NODE=M081R2;LINKAGE=D

<sup>7</sup> Assuming  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$ .

NODE=M081R2;LINKAGE=E

<sup>8</sup> LEES 14M reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$   
 $(1.12 \pm 0.13) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$   
 $(13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is  
the systematic error from using our best value.

NODE=M081R2;LINKAGE=F

<sup>9</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$   
 $(1.1 \pm 0.1 \pm 0.1) \times 10^{-2}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$   
 $(13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error  
is the systematic error from using our best value.

NODE=M081R2;LINKAGE=LE

<sup>10</sup> CRAWFORD 92B quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2$   
 $B(\Upsilon(2S) \rightarrow \ell^+ \ell^-) = (4.98 \pm 0.94 \pm 0.62) 10^{-4}$ .

NODE=M081R2;LINKAGE=B

<sup>11</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow$   
 $\gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18) \%$  using  $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$ .  
Supersedes HEINTZ 91.

NODE=M081R2;LINKAGE=C

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.066<math>\pm</math>0.008 OUR AVERAGE</b>				
0.061 $\pm$ 0.004 $\pm$ 0.007	12,13,14	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.070 $\pm$ 0.004 $\pm$ 0.008	11k	<sup>15</sup> LEES	11J	BABR $\Upsilon(3S) \rightarrow X \gamma$
0.077 $\pm$ 0.018 $\pm$ 0.009	13,16	CRAWFORD	92B	CLE2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
0.061 $\pm$ 0.009 $\pm$ 0.007	13,17	HEINTZ	92	CSB2 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

NODE=M081R1  
NODE=M081R1

<sup>12</sup> LEES 14M quotes  $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$   
 $= (8.03 \pm 0.50) \times 10^{-3}$  combining the results from samples of  $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$   
with and without converted photons.

NODE=M081R1;LINKAGE=D

<sup>13</sup> Assuming  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M081R1;LINKAGE=E

<sup>14</sup> LEES 14M reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$   
 $(8.03 \pm 0.50) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$   
 $(13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is  
the systematic error from using our best value.

NODE=M081R1;LINKAGE=F

<sup>15</sup> LEES 11J reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$   
 $(9.2 \pm 0.3 \pm 0.4) \times 10^{-3}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$   
 $(13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error  
is the systematic error from using our best value.

NODE=M081R1;LINKAGE=LE

<sup>16</sup> CRAWFORD 92B quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times 2$   
 $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (5.03 \pm 0.94 \pm 0.63) 10^{-4}$ .

NODE=M081R1;LINKAGE=B

<sup>17</sup> Recalculated by us. HEINTZ 92 quotes  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow$   
 $\gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$  using  $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$ .  
Supersedes HEINTZ 91.

NODE=M081R1;LINKAGE=C

 $\Gamma(\pi \pi \chi_{b2}(1P))/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.1<math>\pm</math>0.9 OUR AVERAGE</b>				
4.9 $\pm$ 0.7 $\pm$ 0.6	17k	<sup>18</sup> LEES	11C	BABR $e^+ e^- \rightarrow \pi^+ \pi^- X$
6.0 $\pm$ 1.6 $\pm$ 1.4		<sup>19</sup> CAWLFIELD	06	CLE3 $\Upsilon(3S) \rightarrow 2(\gamma \pi \ell)$

NODE=M081R4  
NODE=M081R4

<sup>18</sup>  $(0.64 \pm 0.05 \pm 0.08) \times 10^{-3}$ . We derive the value assuming  $B(\Upsilon(3S) \rightarrow \chi_{b2}(2P) X)$   
 $= B(\Upsilon(3S) \rightarrow \chi_{b2}(2P) \gamma) = (13.1 \pm 1.6) \times 10^{-2}$ .

NODE=M081R4;LINKAGE=LE

<sup>19</sup> CAWLFIELD 06 quote  $\Gamma(\chi_{b2}(2P) \rightarrow \pi \pi \chi_{b1}(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$  keV  
assuming l-spin conservation, no D-wave contribution,  $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$  keV, and  
 $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$  keV.

NODE=M081R4;LINKAGE=CA



$\Gamma(D^0 X)/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-2}$	90	20,21 BRIERE	08	CLEO $\Upsilon(3S) \rightarrow \gamma D^0 X$

NODE=M081R01  
 NODE=M081R01

<sup>20</sup> For  $p_{D^0} > 2.5$  GeV/c.

<sup>21</sup> The authors also present their result as  $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$ .

NODE=M081R01;LINKAGE=BR  
 NODE=M081R01;LINKAGE=RI

 $\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1$	90	<sup>22</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

NODE=M081R02  
 NODE=M081R02

<sup>22</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 14 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

NODE=M081R02;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<0.9$	90	<sup>23</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

NODE=M081R03  
 NODE=M081R03

<sup>23</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 12 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

NODE=M081R03;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
$<7$	90	<sup>24</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

NODE=M081R04  
 NODE=M081R04

<sup>24</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 87 \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .

NODE=M081R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9 \pm 1.6 \pm 0.5$	23	<sup>25</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

NODE=M081R05  
 NODE=M081R05

<sup>25</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.9 \pm 0.4 \pm 0.1$	11	<sup>26</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

NODE=M081R06  
 NODE=M081R06

<sup>26</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 1.0 \pm 0.3$	16	<sup>27</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

NODE=M081R07  
 NODE=M081R07

<sup>27</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (32 \pm 11 \pm 8) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 2.2 \pm 0.6$	14	<sup>28</sup> ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

NODE=M081R08  
 NODE=M081R08

<sup>28</sup> ASNER 08A reports  $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (62 \pm 23 \pm 17) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R08;LINKAGE=AS

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4</b>	90	<sup>29</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
<sup>29</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $< 58 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$ .				

NODE=M081R09  
NODE=M081R09

NODE=M081R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.9±0.4±0.1</b>	14	<sup>30</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
<sup>30</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R10  
NODE=M081R10

NODE=M081R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>12±4±1</b>	45	<sup>31</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
<sup>31</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (159 \pm 33 \pm 43) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R11  
NODE=M081R11

NODE=M081R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$  $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>1.4±0.7±0.2</b>	12	<sup>32</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<sup>32</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (19 \pm 7 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R12  
NODE=M081R12

NODE=M081R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$  $\Gamma_{17}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>4.2±1.7±0.5</b>	16	<sup>33</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<sup>33</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (55 \pm 16 \pm 15) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R13  
NODE=M081R13

NODE=M081R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.9±0.4±0.1</b>	9	<sup>34</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<sup>34</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (12 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R14  
NODE=M081R14

NODE=M081R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>13±5±2</b>	27	<sup>35</sup> ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
<sup>35</sup> ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ $= (165 \pm 46 \pm 50) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R15  
NODE=M081R15

NODE=M081R15;LINKAGE=AS

 $\chi_{b2}(2P)$  Cross-Particle Branching Ratios

NODE=M081230

 $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$   
 $\Gamma_3/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$ 

VALUE (units $10^{-3}$ )	EVTs	DOCUMENT ID	TECN	COMMENT
<b>9.2±0.3±0.4</b>	11k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M081B01  
NODE=M081B01

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) / \Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) / \Gamma_{\text{total}}$$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.1±0.1±0.1</b>	2.5k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M081B02  
NODE=M081B02

$$\mathbf{B}(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times \mathbf{B}(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X)$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.64±0.05±0.08</b>	17k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M081R16  
NODE=M081R16

$$\mathbf{B}(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times \mathbf{B}(\Upsilon(1S) \rightarrow \ell^+\ell^-)$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.02±0.18 OUR AVERAGE</b>				

NODE=M081A01  
NODE=M081A01

1.95 <sup>+0.22+0.10</sup> <sub>-0.21-0.16</sub>		36 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.52±0.47±0.32	48	37 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.98±0.28±0.12		38 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>36</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with converted photons.

<sup>37</sup> CRAWFORD 92B quotes  $2 \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)) \mathbf{B}(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) \mathbf{B}(\Upsilon(nS) \rightarrow \ell^+\ell^-)$ .

<sup>38</sup> Calculated by us. HEINTZ 92 quotes  $\mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times \mathbf{B}(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$  using  $\mathbf{B}(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$ .

NODE=M081A01;LINKAGE=A  
NODE=M081A01;LINKAGE=C

NODE=M081A01;LINKAGE=B

$$\mathbf{[B}(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / \mathbf{[B}(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>66.6±3.0</b>	39 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M081A00  
NODE=M081A00

<sup>39</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  events without converted photons.

NODE=M081A00;LINKAGE=A

$$\mathbf{B}(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times \mathbf{B}(\Upsilon(2S) \rightarrow \ell^+\ell^-)$$

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.74±0.29 OUR AVERAGE</b>				

NODE=M081A02  
NODE=M081A02

3.22 <sup>+0.58+0.16</sup> <sub>-0.53-0.71</sub>		40 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.49±0.47±0.31	53	41 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.74±0.33±0.18		42 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>40</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  with converted photons.

<sup>41</sup> CRAWFORD 92B quotes  $2 \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)) \mathbf{B}(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) \mathbf{B}(\Upsilon(nS) \rightarrow \ell^+\ell^-)$ .

<sup>42</sup> Calculated by us. HEINTZ 92 quotes  $\mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times \mathbf{B}(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18)\%$  using  $\mathbf{B}(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ .

NODE=M081A02;LINKAGE=A  
NODE=M081A02;LINKAGE=C

NODE=M081A02;LINKAGE=B

$$\mathbf{[B}(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / \mathbf{[B}(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times \mathbf{B}(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>46.9±2.0</b>	43 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M081A03  
NODE=M081A03

<sup>43</sup> From a sample of  $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$  without converted photons.

NODE=M081A03;LINKAGE=A

## $\chi_{b2}(2P)$ REFERENCES

AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56199
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
CAWLFIELD	06	PR D73 012003	C. Cawlfeld <i>et al.</i>	(CLEO Collab.)	REFID=50997
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
CRONIN-HEN...04	PRL	92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=49766
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)	REFID=43177
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586

NODE=M081

$\Upsilon(3S)$ 

$$J^{PC} = 0^{-}(1^{-}-)$$

NODE=M048

 **$\Upsilon(3S)$  MASS**

NODE=M048M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10355.1±0.5 OUR AVERAGE</b>	[10355.2 ± 0.5 MeV OUR 2022 AVERAGE]		
<b>10355.1±0.5</b>	<sup>1</sup> SHAMOV	23	RVUE $e^+e^- \rightarrow$ hadrons
10355.2±0.5	<sup>2,3</sup> ARTAMONOV	00	MD1 $e^+e^- \rightarrow$ hadrons
10355.3±0.5	<sup>4,5</sup> BARU	86B	MD1 $e^+e^- \rightarrow$ hadrons

NODE=M048M

NEW

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <sup>1</sup> Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.  
<sup>2</sup> Reanalysis of BARU 86B using new electron mass (COHEN 87).  
<sup>3</sup> Superseded by SHAMOV 23.  
<sup>4</sup> Reanalysis of ARTAMONOV 84.  
<sup>5</sup> Superseded by ARTAMONOV 00.

NODE=M048M;LINKAGE=A

NODE=M048M;LINKAGE=AR

NODE=M048M;LINKAGE=B

NODE=M048M;LINKAGE=C

NODE=M048M;LINKAGE=RZ

 **$m\Upsilon(3S) - m\Upsilon(2S)$** 

NODE=M048DM2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>331.50±0.02±0.13</b>	LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M048DM2

 **$\Upsilon(3S)$  WIDTH**

NODE=M048W

VALUE (keV)	DOCUMENT ID	COMMENT
<b>20.32±1.85 OUR EVALUATION</b>	See the Note on "Width Determinations of the $\Upsilon$ States"	

NODE=M048W

→ UNCHECKED ←

 **$\Upsilon(3S)$  DECAY MODES**

NODE=M048215;NODE=M048

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $\Upsilon(2S)$ anything	(10.6 ± 0.8) %	
$\Gamma_2$ $\Upsilon(2S)\pi^+\pi^-$	(2.82 ± 0.18) %	S=1.6
$\Gamma_3$ $\Upsilon(2S)\pi^0\pi^0$	(1.85 ± 0.14) %	
$\Gamma_4$ $\Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
$\Gamma_5$ $\Upsilon(2S)\pi^0$	< 5.1 × 10 <sup>-4</sup>	CL=90%
$\Gamma_6$ $\Upsilon(1S)\pi^+\pi^-$	(4.37 ± 0.08) %	
$\Gamma_7$ $\Upsilon(1S)\pi^0\pi^0$	(2.20 ± 0.13) %	
$\Gamma_8$ $\Upsilon(1S)\eta$	< 1 × 10 <sup>-4</sup>	CL=90%
$\Gamma_9$ $\Upsilon(1S)\pi^0$	< 7 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{10}$ $h_b(1P)\pi^0$	< 1.2 × 10 <sup>-3</sup>	CL=90%
$\Gamma_{11}$ $h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 ± 1.4) × 10 <sup>-4</sup>	
$\Gamma_{12}$ $h_b(1P)\pi^+\pi^-$	< 1.2 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{13}$ $\tau^+\tau^-$	(2.29 ± 0.30) %	
$\Gamma_{14}$ $\mu^+\mu^-$	(2.18 ± 0.21) %	S=2.1
$\Gamma_{15}$ $e^+e^-$	(2.18 ± 0.20) %	
$\Gamma_{16}$ hadrons	(93 ± 12) %	
$\Gamma_{17}$ $ggg$	(35.7 ± 2.6) %	
$\Gamma_{18}$ $\gamma gg$	(9.7 ± 1.8) × 10 <sup>-3</sup>	
$\Gamma_{19}$ $^2H$ anything	(2.33 ± 0.33) × 10 <sup>-5</sup>	

**Radiative decays**

NODE=M048;CLUMP=B

$\Gamma_{20}$ $\gamma\chi_{b2}(2P)$	(13.1 ± 1.6) %	S=3.4
$\Gamma_{21}$ $\gamma\chi_{b1}(2P)$	(12.6 ± 1.2) %	S=2.4
$\Gamma_{22}$ $\gamma\chi_{b0}(2P)$	(5.9 ± 0.6) %	S=1.4
$\Gamma_{23}$ $\gamma\chi_{b2}(1P)$	(10.0 ± 1.0) × 10 <sup>-3</sup>	S=1.7
$\Gamma_{24}$ $\gamma\chi_{b1}(1P)$	(9 ± 5) × 10 <sup>-4</sup>	S=1.8
$\Gamma_{25}$ $\gamma\chi_{b0}(1P)$	(2.7 ± 0.4) × 10 <sup>-3</sup>	
$\Gamma_{26}$ $\gamma\eta_b(2S)$	< 6.2 × 10 <sup>-4</sup>	CL=90%
$\Gamma_{27}$ $\gamma\eta_b(1S)$	(5.1 ± 0.7) × 10 <sup>-4</sup>	
$\Gamma_{28}$ $\gamma A^0 \rightarrow \gamma$ hadrons	< 8 × 10 <sup>-5</sup>	CL=90%
$\Gamma_{29}$ $\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 × 10 <sup>-4</sup>	CL=95%
$\Gamma_{30}$ $\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	< 5.5 × 10 <sup>-6</sup>	CL=90%
$\Gamma_{31}$ $\gamma A^0 \rightarrow \gamma\tau^+\tau^-$	[b] < 1.6 × 10 <sup>-4</sup>	CL=90%

DESIG=8

DESIG=4

DESIG=10

DESIG=12

DESIG=107

DESIG=3

DESIG=11

DESIG=9

DESIG=106

DESIG=112

DESIG=113

DESIG=114

DESIG=16

DESIG=1

DESIG=2

DESIG=101

DESIG=109

DESIG=110

DESIG=117

DESIG=117

**Lepton Family number (LF) violating modes**

$\Gamma_{32}$	$e^\pm \tau^\mp$	LF	< 4.2	$\times 10^{-6}$	CL=90%
$\Gamma_{33}$	$e^\pm \mu^\mp$	LF	< 3.6	$\times 10^{-7}$	CL=90%
$\Gamma_{34}$	$\mu^\pm \tau^\mp$	LF	< 3.1	$\times 10^{-6}$	CL=90%

NODE=M048;CLUMP=C

DESIG=111

DESIG=119

DESIG=105

[a]  $1.5 \text{ GeV} < m_X < 5.0 \text{ GeV}$ 

LINKAGE=C48

[b] For  $m_{\tau^+\tau^-}$  in the ranges 4.03–9.52 and 9.61–10.10 GeV.

LINKAGE=MRG

 **$\Upsilon(3S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$** 

NODE=M048218

 **$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_{16}\Gamma_{15}/\Gamma$** 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.414±0.007 OUR AVERAGE</b>			
0.413±0.004±0.006	ROSNER	06	CLEO $10.4 e^+e^- \rightarrow \text{hadrons}$
0.45 ±0.03 ±0.03	<sup>6</sup> GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$

NODE=M048G2

NODE=M048G2

<sup>6</sup> Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

NODE=M048G2;LINKAGE=R

 **$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$**   **$\Gamma_6\Gamma_{15}/\Gamma$** 

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>18.46±0.27±0.77</b>	6.4k	<sup>7</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

NODE=M048G01

NODE=M048G01

<sup>7</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M048G01;LINKAGE=AU

 **$\Upsilon(3S)$  PARTIAL WIDTHS**

NODE=M048220

 **$\Gamma(e^+e^-)$**   **$\Gamma_{15}$** 

VALUE (keV)	DOCUMENT ID
<b>0.443±0.008 OUR EVALUATION</b>	

NODE=M048W2

NODE=M048W2

→ UNCHECKED ←

 **$\Upsilon(3S)$  BRANCHING RATIOS**

NODE=M048225

 **$\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}}$**   **$\Gamma_1/\Gamma$** 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.106 ±0.008 OUR AVERAGE</b>				
0.1023±0.0105	4625	<sup>8,9,10</sup> BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-X$
0.111 ±0.012	4891	<sup>9,10,11</sup> BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

NODE=M048R8

NODE=M048R8

<sup>8</sup> Using  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$ , and  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$ .

NODE=M048R;LINKAGE=A

<sup>9</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ . With the assumption of  $e\mu$  universality.

NODE=M048R;LINKAGE=B

<sup>10</sup> Using  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$ .

NODE=M048R;LINKAGE=D

<sup>11</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ ,  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$ , and  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$ . With the assumption of  $e\mu$  universality.

NODE=M048R;LINKAGE=C

 **$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_2/\Gamma$** 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.82±0.18 OUR AVERAGE</b>				Error includes scale factor of 1.6. See the ideogram below.

NODE=M048R4

NODE=M048R4

3.00±0.02±0.14	543k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
2.40±0.10±0.26	800	<sup>12</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-e^+e^-$
3.12±0.49	980	<sup>13,14</sup> BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
2.13±0.38	974	<sup>15</sup> BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.82±0.65±0.53	138	<sup>15</sup> WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.1 ±2.0	5	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

<sup>12</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ ,  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ , and  $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008 \text{ keV}$ .

NODE=M048R4;LINKAGE=AU

<sup>13</sup> From the exclusive mode.


NODE=M048R;LINKAGE=M

<sup>14</sup> Using  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$ , and  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$ .

NODE=M048R4;LINKAGE=A

<sup>15</sup> Using  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ ,  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$ , and  $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$ . With the assumption of  $e\mu$  universality.

NODE=M048R4;LINKAGE=C



**$\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{total}$   $\Gamma_3/\Gamma$**

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.85±0.14 OUR AVERAGE</b>				
1.82±0.09±0.12	4391	<sup>16</sup> BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16±0.39		<sup>17,18</sup> BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.7 ±0.5 ±0.2	10	<sup>19</sup> HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

NODE=M048R10  
NODE=M048R10

<sup>16</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$ .  
<sup>17</sup>  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$  and assuming  $e\mu$  universality.  
<sup>18</sup> From the exclusive mode.  
<sup>19</sup>  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

NODE=M048R10;LINKAGE=BH  
 NODE=M048R;LINKAGE=K  
 NODE=M048R10;LINKAGE=M  
 NODE=M048R;LINKAGE=G

**$\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{total}$   $\Gamma_4/\Gamma$**

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.0502±0.0069</b>	<sup>20</sup> BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

NODE=M048R12  
NODE=M048R12

<sup>20</sup> From the exclusive mode.

NODE=M048R12;LINKAGE=M

**$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{total}$   $\Gamma_5/\Gamma$**

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.51</b>	90	<sup>21</sup> HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M048R25  
NODE=M048R25

<sup>21</sup> Authors assume  $B(\Upsilon(2S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$ .

NODE=M048R25;LINKAGE=HE

**$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{total}$   $\Gamma_6/\Gamma$**

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>4.37±0.08 OUR AVERAGE</b>				
4.32±0.07±0.13	90k	<sup>22</sup> LEES	11L BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.01±0.13	190k	<sup>23</sup> BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^-$ MM
4.17±0.06±0.19	6.4k	<sup>24</sup> AUBERT	08BP BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52±0.35	11830	<sup>25</sup> BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$
4.46±0.34±0.50	451	<sup>25</sup> WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.30	11221	<sup>25</sup> BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

NODE=M048R3  
 NODE=M048R3  
 NODE=M048R3

••• We do not use the following data for averages, fits, limits, etc. •••

4.9 ±1.0	22	GREEN	82 CLEO	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.9 ±1.3	26	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

<sup>22</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M048R3;LINKAGE=LE

<sup>23</sup> A weighted average of the inclusive and exclusive results.

NODE=M048R3;LINKAGE=BH  
 NODE=M048R3;LINKAGE=AU

<sup>24</sup> Using  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ ,  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ , and  $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$  keV.

<sup>25</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ . With the assumption of  $e\mu$  universality.

NODE=M048R3;LINKAGE=B

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$  $\Gamma_2/\Gamma_6$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M048R28  
 NODE=M048R28

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.577±0.026±0.060	800	<sup>26</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
-------------------	-----	----------------------	-----------	---

<sup>26</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ ,  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ ,  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ , and  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ . Not independent of other values reported by AUBERT 08BP.

NODE=M048R28;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M048R11  
 NODE=M048R11

**2.20±0.13 OUR AVERAGE**

2.24±0.09±0.11	6584	<sup>27</sup> BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99±0.34	56	<sup>28</sup> BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 ±0.4 ±0.3	33	<sup>29</sup> HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

<sup>27</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

<sup>28</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$  and assuming  $e\mu$  universality.

<sup>29</sup> Using  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$  and assuming  $e\mu$  universality. Supersedes HEINTZ 91.

NODE=M048R11;LINKAGE=BH  
 NODE=M048R11;LINKAGE=B  
 NODE=M048R;LINKAGE=I

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$  $\Gamma_7/\Gamma_6$ 

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M048R26  
 NODE=M048R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.501±0.043	<sup>30</sup> BHARI	09 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$
-------------	---------------------	---------	-----------------------------------

<sup>30</sup> Not independent of other values reported by BHARI 09.

NODE=M048R26;LINKAGE=BH

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R9  
 NODE=M048R9

<0.1	90	<sup>31</sup> LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
------	----	--------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90	<sup>31,32</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
------	----	-------------------------	-----------	--

<0.18	90	<sup>33</sup> HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
-------	----	------------------	----------	---------------------------------------

<2.2	90	BROCK	91 CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
------	----	-------	---------	---------------------------------------

<sup>31</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ ,  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

<sup>32</sup> Using  $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$  keV.

<sup>33</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

NODE=M048R9;LINKAGE=LE  
 NODE=M048R9;LINKAGE=AU  
 NODE=M048R9;LINKAGE=HE

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$  $\Gamma_8/\Gamma_6$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R27  
 NODE=M048R27

<0.23	90	<sup>34</sup> LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
-------	----	--------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9	90	<sup>35</sup> AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$
------	----	----------------------	-----------	--

<sup>34</sup> Not independent of other values reported by LEES 11L.

<sup>35</sup> Not independent of other values reported by AUBERT 08BP.

NODE=M048R27;LINKAGE=LE  
 NODE=M048R27;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE (units $10^{-3}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R24  
 NODE=M048R24

<0.07	90	<sup>36</sup> HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
-------	----	------------------	----------	---

<sup>36</sup> Authors assume  $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$ .

NODE=M048R24;LINKAGE=HE

 $\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M048R03  
 NODE=M048R03

<1.2 × 10 <sup>-3</sup>	90	<sup>37</sup> GE	11 CLEO	$\Upsilon(3S) \rightarrow \pi^0$ anything
-------------------------	----	------------------	---------	---

<sup>37</sup> Assuming  $M(h_b(1P)) = 9900$  MeV and  $\Gamma(h_b(1P)) = 0$  MeV, and allowing  $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$  to vary from 0–100%.

NODE=M048R03;LINKAGE=GE

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M048R33  
 NODE=M048R33

<b>4.3±1.1±0.9</b>	LEES	11K BABR	$\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$
--------------------	------	----------	--

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{total}$   $\Gamma_{12}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	38 LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^- X$
<18		38 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^- X$
<15		38 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^- X$

<sup>38</sup> For  $M(h_b(1P)) = 9900$  MeV.

NODE=M048R34  
NODE=M048R34

NODE=M048R34;LINKAGE=MH

$\Gamma(\tau^+\tau^-)/\Gamma_{total}$   $\Gamma_{13}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.29±0.21±0.22</b>	15k	<sup>39</sup> BESSON	07 CLEO	$e^+e^- \rightarrow \Upsilon(3S) \rightarrow \tau^+\tau^-$

<sup>39</sup> BESSON 07 reports  $[\Gamma(\Upsilon(3S) \rightarrow \tau^+\tau^-)/\Gamma_{total}] / [B(\Upsilon(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$  which we multiply by our best value  $B(\Upsilon(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M048R18  
NODE=M048R18

NODE=M048R18;LINKAGE=BE

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$   $\Gamma_{13}/\Gamma_{14}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.968±0.016 OUR AVERAGE</b>				
0.966±0.008±0.014	2.2M	LEES	20E BABR	$e^+e^- \rightarrow \Upsilon(3S)$
1.05 ±0.08 ±0.05	15k	BESSON	07 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$


NODE=M048R19  
NODE=M048R19

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$   $\Gamma_{14}/\Gamma$

**0.0218±0.0021 OUR AVERAGE** Error includes scale factor of 2.1. See the ideogram below.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0239±0.0007±0.0010	81k	ADAMS	05 CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0202±0.0019±0.0033		CHEN	89B CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0173±0.0015±0.0011		KAARSBERG	89 CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.033 ±0.013 ±0.007	1096	ANDREWS	83 CLEO	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M048R1  
NODE=M048R1



$\Gamma(ggg)/\Gamma_{total}$   $\Gamma_{17}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>35.7±2.6</b>	3M	<sup>40</sup> BESSON	06A CLEO	$\Upsilon(3S) \rightarrow \text{hadrons}$

<sup>40</sup> Calculated using BESSON 06A value of  $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$  and the PDG 08 values of  $B(\Upsilon(2S) + \text{anything}) = (10.6 \pm 0.8)\%$ ,  $B(\pi^+\pi^-\Upsilon(1S)) = (4.40 \pm 0.10)\%$ ,  $B(\pi^0\pi^0\Upsilon(1S)) = (2.20 \pm 0.13)\%$ ,  $B(\gamma\chi_{b2}(2P)) = (13.1 \pm 1.6)\%$ ,  $B(\gamma\chi_{b1}(2P)) = (12.6 \pm 1.2)\%$ ,  $B(\gamma\chi_{b0}(2P)) = (5.9 \pm 0.6)\%$ ,  $B(\gamma\chi_{b0}(1P)) = (0.30 \pm 0.11)\%$ ,  $B(\mu^+\mu^-) = (2.18 \pm 0.21)\%$ , and  $R_{\text{hadrons}} = 3.51$ . The statistical error is negligible and the systematic error is partially correlated with  $\Gamma(\gamma gg)/\Gamma_{total}$  BESSON 06A value.

NODE=M048R30  
NODE=M048R30

NODE=M048R30;LINKAGE=BE



$\Gamma(\gamma g g)/\Gamma_{total}$   $\Gamma_{18}/\Gamma$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.97±0.18</b>	60k	<sup>41</sup> BESSON	06A CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

NODE=M048R31  
NODE=M048R31

<sup>41</sup> Calculated using BESSON 06A values of  $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$  and  $\Gamma(g g g)/\Gamma_{total}$ . The statistical error is negligible and the systematic error is partially correlated with  $\Gamma(g g g)/\Gamma_{total}$  BESSON 06A value.

NODE=M048R31;LINKAGE=BE

$\Gamma(\gamma g g)/\Gamma(g g g)$   $\Gamma_{18}/\Gamma_{17}$

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>2.72±0.06±0.49</b>	3M	BESSON	06A CLEO	$\Upsilon(3S) \rightarrow (\gamma +) \text{hadrons}$

NODE=M048R32  
NODE=M048R32

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{total}$   $\Gamma_{19}/\Gamma$

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.33±0.15<sup>+0.31</sup><sub>-0.28</sub></b>	LEES	14G BABR	$e^+ e^- \rightarrow \overline{2H} X$

NODE=M048R00  
NODE=M048R00


$\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{total}$   $\Gamma_{20}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.131 ±0.016 OUR AVERAGE</b>				Error includes scale factor of 3.4. See the ideogram below.
0.1579±0.0017±0.0073	568k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.111 ±0.005 ±0.004	10319	<sup>42</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.135 ±0.003 ±0.017	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

NODE=M048R5  
NODE=M048R5

<sup>42</sup>Supersedes NARAIN 91.

NODE=M048R;LINKAGE=H




$\Gamma(\gamma \chi_{b1}(2P))/\Gamma_{total}$   $\Gamma_{21}/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.126 ±0.012 OUR AVERAGE</b>				Error includes scale factor of 2.4. See the ideogram below.
0.1454±0.0018±0.0073	537k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.115 ±0.005 ±0.005	11147	<sup>43</sup> HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.105 <sup>+0.003</sup> <sub>-0.002</sub> ±0.013	25759	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

NODE=M048R6  
NODE=M048R6

<sup>43</sup>Supersedes NARAIN 91.

NODE=M048R6;LINKAGE=H




**$\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{total}$**   **$\Gamma_{22}/\Gamma$**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.059 ±0.006 OUR AVERAGE</b>				Error includes scale factor of 1.4. See the ideogram below.
0.0677±0.0020±0.0065	225k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
0.060 ±0.004 ±0.006	4959	<sup>44</sup> HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma X$
0.049 <sup>+0.003</sup> <sub>-0.004</sub> ±0.006	9903	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma X$

NODE=M048R7  
NODE=M048R7

<sup>44</sup>Supersedes NARAIN 91.

NODE=M048R7;LINKAGE=H



**$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{total}$**   **$\Gamma_{23}/\Gamma$**

VALUE (units 10 <sup>-3</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10.0±1.0 OUR AVERAGE</b>					Error includes scale factor of 1.7.
8.0±1.3±0.4	126	<sup>45,46</sup> KORNICER	11	CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
10.5±0.3 <sup>+0.7</sup> <sub>-0.6</sub>	9.7k	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M048R21  
NODE=M048R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	<sup>47</sup> ASNER	08A	CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
seen		<sup>48</sup> HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

<sup>45</sup> Assuming  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$ .

<sup>46</sup> KORNICER 11 reports  $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))]$   
 $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$  which we divide by our best value  $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) = (18.0 \pm 1.0) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>47</sup> ASNER 08A reports  $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$   
 $< 27.1 \times 10^{-2}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$ .

<sup>48</sup> HEINTZ 92, while unable to distinguish between different  $J$  states, measures  $\sum_J B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma \Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$  for  $J = 0,1,2$  using inclusive  $\Upsilon(1S)$  decays and  $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$  for  $J = 1,2$  using  $\Upsilon(1S) \rightarrow \ell^+ \ell^-$ .

NODE=M048R21;LINKAGE=KA  
 NODE=M048R21;LINKAGE=KR

NODE=M048R21;LINKAGE=AS

NODE=M048R21;LINKAGE=HE

### $\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$

$\Gamma_{24}/\Gamma$

VALUE (units $10^{-3}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.9±0.5 OUR AVERAGE</b> Error includes scale factor of 1.8.					
$1.5 \pm 0.4 \pm 0.1$		50	<sup>49,50</sup> KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
$0.5 \pm 0.3^{+0.2}_{-0.1}$			LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$

NODE=M048R22  
 NODE=M048R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 1.7$  90 <sup>51</sup> ASNER 08A CLEO  $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$   
 seen <sup>52</sup> HEINTZ 92 CSB2  $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

<sup>49</sup> Assuming  $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$ .

<sup>50</sup> KORNICER 11 reports  $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))]$   
 $= (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$  which we divide by our best value  $B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) = (35.2 \pm 2.0) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>51</sup> ASNER 08A reports  $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] < 2.5 \times 10^{-2}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = 6.9 \times 10^{-2}$ .

<sup>52</sup> HEINTZ 92, while unable to distinguish between different  $J$  states, measures  $\sum_J B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma \Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$  for  $J = 0,1,2$  using inclusive  $\Upsilon(1S)$  decays and  $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$  for  $J = 1,2$  using  $\Upsilon(1S) \rightarrow \ell^+ \ell^-$ .

NODE=M048R22;LINKAGE=KA  
 NODE=M048R22;LINKAGE=KR

NODE=M048R22;LINKAGE=AS

NODE=M048R22;LINKAGE=HE

### $\Gamma(\gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$

$\Gamma_{25}/\Gamma$

VALUE (units $10^{-2}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>0.27±0.04 OUR AVERAGE</b>					
$0.27 \pm 0.04 \pm 0.02$		2.3k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
$0.30 \pm 0.04 \pm 0.10$		8.7k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$

NODE=M048R15  
 NODE=M048R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 0.8$  90 <sup>53</sup> ASNER 08A CLEO  $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

<sup>53</sup> ASNER 08A reports  $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 21.9 \times 10^{-2}$  which we multiply by our best value  $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$ .

NODE=M048R15;LINKAGE=AS

### $\Gamma(\gamma \eta_b(2S))/\Gamma_{\text{total}}$

$\Gamma_{26}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>&lt; 6.2</b>	90		ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
$< 19$	90		LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$

NODE=M048R16  
 NODE=M048R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

### $\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$

$\Gamma_{27}/\Gamma$

VALUE (units $10^{-4}$ )	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<b>5.1±0.7 OUR AVERAGE</b>					
$7.1 \pm 1.8 \pm 1.3$		$2.3 \pm 0.5k$	<sup>54</sup> BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
$4.8 \pm 0.5 \pm 0.6$		$19 \pm 3k$	<sup>54</sup> AUBERT	09AQ BABR	$\Upsilon(3S) \rightarrow \gamma X$
$< 8.5$	90		LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
$4.8 \pm 0.5 \pm 1.2$		$19 \pm 3k$	<sup>54,55</sup> AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
$< 4.3$	90		<sup>56</sup> ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$

NODE=M048R17  
 NODE=M048R17

<sup>54</sup> Assuming  $\Gamma_{\eta_b(1S)} = 10 \text{ MeV}$ .

<sup>55</sup> Systematic error re-evaluated by AUBERT 09AQ.

<sup>56</sup> Superseded by BONVICINI 10.

NODE=M048R17;LINKAGE=BO  
 NODE=M048R17;LINKAGE=AU  
 NODE=M048R17;LINKAGE=SU

$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$   
(0.3 GeV <  $m_{A^0}$  < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<8 \times 10^{-5}$  90 57 LEES 11H BABR  $\Upsilon(3S) \rightarrow \gamma \text{hadrons}$

<sup>57</sup> For a narrow scalar or pseudoscalar,  $A^0$ , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of  $m_{A^0}$  range from  $1 \times 10^{-6}$  to  $8 \times 10^{-5}$ .

NODE=M048R02

NODE=M048R02  
NODE=M048R02

NODE=M048R02;LINKAGE=LE

$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$   
(1.5 GeV <  $m_X$  < 5.0 GeV)

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<2.2$  95 ROSNER 07A CLEO  $e^+e^- \rightarrow \gamma X$

NODE=M048R20

NODE=M048R20  
NODE=M048R20

$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<5.5$  90 58 AUBERT 09Z BABR  $e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

<sup>58</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with mass in the range 212–9300 MeV, excluding  $J/\psi$  and  $\psi(2S)$ . Measured 90% CL limits as a function of  $m_{A^0}$  range from 0.27–5.5  $\times 10^{-6}$ .

NODE=M048R04  
NODE=M048R04

NODE=M048R04;LINKAGE=AU

$\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<1.6 \times 10^{-4}$  90 59 AUBERT 09P BABR  $e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

<sup>59</sup> For a narrow scalar or pseudoscalar,  $A^0$ , with  $M(\tau^+ \tau^-)$  in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of  $M(\tau^+ \tau^-)$  range from  $1.5-16 \times 10^{-5}$ .

NODE=M048R29  
NODE=M048R29

NODE=M048R29;LINKAGE=AU

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<4.2$  90 LEES 10B BABR  $e^+e^- \rightarrow e^\pm \tau^\mp$

NODE=M048230

NODE=M048R01  
NODE=M048R01

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$   $\Gamma_{33}/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<3.6 \times 10^{-7}$  90 LEES 22A BABR  $e^+e^- \rightarrow e^\pm \mu^\mp$

NODE=M048R07  
NODE=M048R07

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$   $\Gamma_{34}/\Gamma$

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

$<3.1$  90 LEES 10B BABR  $e^+e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M048R23  
NODE=M048R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<20.3$  95 LOVE 08A CLEO  $e^+e^- \rightarrow \mu^\pm \tau^\mp$

$\Upsilon(3S)$  REFERENCES

SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVO)	REFID=62012
LEES	22A	PRL 128 091804	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61659
LEES	20E	PRL 125 241801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=60700
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)	REFID=53960
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53937
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53938
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53062
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52262
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079

NODE=M048

BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43799
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)	REFID=43313
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)	REFID=41579
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUELLER...	88	HE $e^+e^-$ Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding,		World Scientific, Singapore			
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU	86B	ZPHY C32 622 (erratum)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41 733.			
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)	REFID=22273
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)	REFID=22321
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)	REFID=22359

NODE=M206

 $\chi_{b1}(3P)$ 

$$I^G(J^{PC}) = 0^+(1^{++})$$

Observed in the radiative decay to  $\Upsilon(1S, 2S, 3S)$ , therefore  $C = +$ .  
 $J$  needs confirmation.

NODE=M206

### $\chi_{b1}(3P)$ MASS

NODE=M206M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>10513.42 ± 0.41 ± 0.53</b>		<sup>1</sup> SIRUNYAN	18N CMS	$pp \rightarrow \gamma \mu^+ \mu^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
10515.7 <sup>+2.2</sup> <sub>-3.9</sub> <sup>+1.5</sup> <sub>-2.1</sub>	169	<sup>2</sup> AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10512.1 ± 2.1 ± 0.9	351	<sup>3</sup> AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10511.3 ± 1.7 ± 2.5	182	<sup>4</sup> AAIJ	14BI LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10530 ± 5 ± 9		<sup>5</sup> AAD	12A ATLS	$pp \rightarrow \gamma \mu^+ \mu^- X$
10551 ± 14 ± 17		<sup>5</sup> ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma \mu^+ \mu^- X$

NODE=M206M

<sup>1</sup> Systematic error includes an additional 0.5 MeV for the uncertainty on the  $\Upsilon(3S)$  mass. Also measures  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$  MeV. A total of 372  $\chi_{b1}(3P)$  and  $\chi_{b2}(3P)$  events was observed.

NODE=M206M;LINKAGE=D

<sup>2</sup> From  $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$  transitions assuming  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$  MeV and allowing for  $\pm 30\%$  variation in the  $\chi_{b2}(3P)$  production rate relative to that of  $\chi_{b1}(3P)$ .

NODE=M206M;LINKAGE=A

<sup>3</sup> The mass of the  $\chi_{b1}(3P)$  state obtained by combining the results of AAIJ 14BG with that of AAIJ 14BI. The first uncertainty is experimental and the second attributable to the unknown mass splitting, assumed to be  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$  MeV.

NODE=M206M;LINKAGE=B

<sup>4</sup> From  $\chi_{b1}(3P) \rightarrow \Upsilon(3S)\gamma$  transition assuming  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$  MeV.

NODE=M206M;LINKAGE=C

<sup>5</sup> The mass barycenter of the merged lineshapes from the  $J = 1$  and 2 states.

NODE=M206M;LINKAGE=AA

### $\chi_{b1}(3P)$ DECAY MODES

NODE=M206215;NODE=M206

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\Upsilon(1S)\gamma$	seen
$\Gamma_2$ $\Upsilon(2S)\gamma$	seen
$\Gamma_3$ $\Upsilon(3S)\gamma$	seen

DESIG=1

DESIG=2

DESIG=3

### $\chi_{b1}(3P)$ BRANCHING RATIOS

NODE=M206225

$\Gamma(\Upsilon(1S)\gamma)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>seen</b>		169	<sup>1</sup> AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
seen			AAD	12A ATLS	$pp \rightarrow \gamma \mu^+ \mu^- X$	
seen			ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma \mu^+ \mu^- X$	

NODE=M206R01

NODE=M206R01

<sup>1</sup> From  $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$  transitions assuming  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$  MeV and allowing for  $\pm 30\%$  variation in the  $\chi_{b2}(3P)$  production rate relative to that of  $\chi_{b1}(3P)$ .

NODE=M206R01;LINKAGE=A

$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	169	<sup>1</sup> AAIJ	14BG LHCb	$pp \rightarrow \gamma\mu^+\mu^-X$
seen		AAD	12A ATLAS	$pp \rightarrow \gamma\mu^+\mu^-X$

<sup>1</sup> From  $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$  transitions assuming  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$  MeV and allowing for  $\pm 30\%$  variation in the  $\chi_{b2}(3P)$  production rate relative to that of  $\chi_{b1}(3P)$ .

NODE=M206R02  
NODE=M206R02

NODE=M206R02;LINKAGE=A

 $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen		SIRUNYAN	18N CMS	$pp \rightarrow \gamma\mu^+\mu^-X$
seen	182	AAIJ	14BI LHCb	$pp \rightarrow \gamma\mu^+\mu^-X$

NODE=M206R03  
NODE=M206R03

 $\chi_{b1}(3P)$  REFERENCES

SIRUNYAN	18N	PRL 121 092002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BI	EPJ C74 3092	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	12A	PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV	12Q	PR D86 031103	V.M. Abazov <i>et al.</i>	(D0 Collab.)

NODE=M206

REFID=58873  
REFID=56199  
REFID=56235  
REFID=54037  
REFID=54264

$\chi_{b2}(3P)$

$$J^{PC} = 0^+(2^{++})$$

Observed in the radiative decay to  $\Upsilon(3S)$ , therefore  $C = +$ .  $J$  needs confirmation.

NODE=M238

NODE=M238

 $\chi_{b2}(3P)$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10524.02 ± 0.57 ± 0.53</b>	<sup>1</sup> SIRUNYAN	18N CMS	$pp \rightarrow \gamma\mu^+\mu^-X$
10530 ± 5 ± 9	<sup>2</sup> AAD	12A ATLAS	$pp \rightarrow \gamma\mu^+\mu^-X$

<sup>1</sup> Systematic error includes an additional 0.5 MeV for the uncertainty on the  $\Upsilon(3S)$  mass. Also measures  $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$  MeV. A total of 372  $\chi_{b1}(3P)$  and  $\chi_{b2}(3P)$  events was observed.

<sup>2</sup> The mass barycenter of the merged lineshapes from the  $J = 1$  and 2 states.

NODE=M238M

NODE=M238M

NODE=M238M;LINKAGE=A

NODE=M238M;LINKAGE=AA

 $\chi_{b2}(3P)$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\Upsilon(3S)\gamma$	seen

NODE=M238215;NODE=M238

DESIG=1

 $\chi_{b2}(3P)$  BRANCHING RATIOS $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	SIRUNYAN	18N CMS	$pp \rightarrow \gamma\mu^+\mu^-X$

NODE=M238225

NODE=M238R01  
NODE=M238R01

 $\chi_{b2}(3P)$  REFERENCES

SIRUNYAN	18N	PRL 121 092002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAD	12A	PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)

NODE=M238

REFID=58873  
REFID=54037

$\Upsilon(4S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as  $\Upsilon(10580)$ 

NODE=M047

 **$\Upsilon(4S)$  MASS**

NODE=M047M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10579.4±1.2 OUR AVERAGE</b>			
10579.3±0.4±1.2	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
10580.0±3.5	<sup>1</sup> BEBEK	87 CLEO	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10577.4±1.0	<sup>2</sup> LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons
<sup>1</sup> Reanalysis of BESSON 85.			
<sup>2</sup> No systematic error given.			

NODE=M047M

NODE=M047M;LINKAGE=C  
NODE=M047M;LINKAGE=B **$\Upsilon(4S)$  WIDTH**

NODE=M047W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>20.5±2.5 OUR AVERAGE</b>			
20.7±1.6±2.5	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
20 ±2 ±4	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
25 ±2.5	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

NODE=M047W

 **$\Upsilon(4S)$  DECAY MODES**

NODE=M047215;NODE=M047

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level
$\Gamma_1$ $B\bar{B}$	> 96 %	95%
$\Gamma_2$ $B^+B^-$	(51.4 ±0.6) %	
$\Gamma_3$ $D^+$ anything + c.c.	(17.8 ±2.6) %	
$\Gamma_4$ $B^0\bar{B}^0$	(48.6 ±0.6) %	
$\Gamma_5$ $J/\psi K_S^0 + (J/\psi, \eta_c) K_S^0$	< 4 × 10 <sup>-7</sup>	90%
$\Gamma_6$ non- $B\bar{B}$	< 4 %	95%
$\Gamma_7$ $e^+e^-$	(1.57±0.08) × 10 <sup>-5</sup>	
$\Gamma_8$ $\rho^+\rho^-$	< 5.7 × 10 <sup>-6</sup>	90%
$\Gamma_9$ $K^*(892)^0\bar{K}^0$	< 2.0 × 10 <sup>-6</sup>	90%
$\Gamma_{10}$ $J/\psi(1S)$ anything	< 1.9 × 10 <sup>-4</sup>	95%
$\Gamma_{11}$ $D^{*+}$ anything + c.c.	< 7.4 %	90%
$\Gamma_{12}$ $\phi$ anything	(7.1 ±0.6) %	
$\Gamma_{13}$ $\phi\eta$	< 1.8 × 10 <sup>-6</sup>	90%
$\Gamma_{14}$ $\phi\eta'$	< 4.3 × 10 <sup>-6</sup>	90%
$\Gamma_{15}$ $\rho\eta$	< 1.3 × 10 <sup>-6</sup>	90%
$\Gamma_{16}$ $\rho\eta'$	< 2.5 × 10 <sup>-6</sup>	90%
$\Gamma_{17}$ $\Upsilon(1S)$ anything	< 4 × 10 <sup>-3</sup>	90%
$\Gamma_{18}$ $\Upsilon(1S)\pi^+\pi^-$	(8.2 ±0.4) × 10 <sup>-5</sup>	
$\Gamma_{19}$ $\Upsilon(1S)\eta$	(1.81±0.18) × 10 <sup>-4</sup>	
$\Gamma_{20}$ $\Upsilon(1S)\eta'$	(3.4 ±0.9) × 10 <sup>-5</sup>	
$\Gamma_{21}$ $\Upsilon(2S)\pi^+\pi^-$	(8.2 ±0.8) × 10 <sup>-5</sup>	
$\Gamma_{22}$ $h_b(1P)\pi^+\pi^-$	not seen	
$\Gamma_{23}$ $h_b(1P)\eta$	(2.18±0.21) × 10 <sup>-3</sup>	
$\Gamma_{24}$ $\eta_b(1S)\omega$	< 1.8 × 10 <sup>-4</sup>	90%
$\Gamma_{25}$ ${}^2H$ anything	< 1.3 × 10 <sup>-5</sup>	90%

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=10

DESIG=12

DESIG=11

DESIG=15

DESIG=6

DESIG=1

DESIG=16

DESIG=22

DESIG=2

DESIG=3

DESIG=4

DESIG=13

DESIG=18

DESIG=19

DESIG=20

DESIG=5

DESIG=7

DESIG=17

DESIG=26

DESIG=9

DESIG=21

DESIG=23

DESIG=27

DESIG=14

**Double Radiative Decays**

$\Gamma_{26}$ $\gamma\gamma\Upsilon(D) \rightarrow \gamma\gamma\eta\Upsilon(1S)$	< 2.3 × 10 <sup>-5</sup>	90%
--	--------------------------	-----

NODE=M047;CLUMP=B

DESIG=24

### $\Upsilon(4S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

$\Gamma_7$


VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.272 ± 0.029 OUR AVERAGE</b>	Error includes scale factor of 1.5. See the ideogram below.		
0.321 ± 0.017 ± 0.029	AUBERT	05Q	BABR $e^+e^- \rightarrow$ hadrons
0.28 ± 0.05 ± 0.01	<sup>1</sup> ALBRECHT	95E	ARG $e^+e^- \rightarrow$ hadrons
0.192 ± 0.007 ± 0.038	BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
0.283 ± 0.037	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

<sup>1</sup> Using LEYAOUANC 77 parametrization of  $\Gamma(s)$ .

NODE=M047220

NODE=M047W1  
NODE=M047W1

NODE=M047W1;LINKAGE=A



### $\Upsilon(4S)$ BRANCHING RATIOS

#### $B\bar{B}$ DECAYS

The ratio of branching fraction to charged and neutral B mesons is often derived assuming isospin invariance in the decays, and relies on the knowledge of the  $B^+/B^0$  lifetime ratio. "OUR EVALUATION" is obtained based on averages of rescaled data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <https://hflav.web.cern.ch/>. The averaging/rescaling procedure takes into account the common dependence of the measurement on the value of the lifetime ratio.

NODE=M047230

NODE=M047BBD

NODE=M047BBD

$\Gamma(B^+B^-)/\Gamma_{total}$

$\Gamma_2/\Gamma$

VALUE	DOCUMENT ID	COMMENT
<b>0.514 ± 0.006 OUR EVALUATION</b>	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$	

NODE=M047R11  
NODE=M047R11  
→ UNCHECKED ←

$\Gamma(D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{total}$

$\Gamma_3/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.178 ± 0.021 ± 0.016</b>	<sup>1</sup> ARTUSO	05B	CLE3 $e^+e^- \rightarrow D_s X$
<sup>1</sup> ARTUSO 05B reports $[\Gamma(\Upsilon(4S) \rightarrow D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{total}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (8.0 \pm 0.2 \pm 0.9) \times 10^{-3}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.			

NODE=M047R13  
NODE=M047R13  
NODE=M047R13;LINKAGE=AR

$\Gamma(B^0\bar{B}^0)/\Gamma_{total}$

$\Gamma_4/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.486 ± 0.006 OUR EVALUATION</b>	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$		

NODE=M047R12  
NODE=M047R12  
→ UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

0.487 ± 0.010 ± 0.008	<sup>1</sup> AUBERT,B	05H	BABR $\Upsilon(4S) \rightarrow \bar{B}B \rightarrow D^* \ell \nu_\ell$
<sup>1</sup> Direct measurement. This value is averaged with the value extracted from the $\Gamma(B^+B^-)/\Gamma(B^0\bar{B}^0)$ measurements.			

NODE=M047R12;LINKAGE=AU



$\Gamma(B^+ B^-)/\Gamma(B^0 \bar{B}^0)$  $\Gamma_2/\Gamma_4$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.058 ± 0.024 OUR EVALUATION</b>			
1.006 ± 0.036 ± 0.031	<sup>1</sup> AUBERT 04F	BABR	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K$
1.01 ± 0.03 ± 0.09	<sup>1</sup> HASTINGS 03	BELL	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow \text{dileptons}$
1.058 ± 0.084 ± 0.136	<sup>2</sup> ATHAR 02	CLEO	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow D^* \ell \nu$
1.10 ± 0.06 ± 0.05	<sup>3</sup> AUBERT 02	BABR	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow (c\bar{c})K^*$
1.04 ± 0.07 ± 0.04	<sup>4</sup> ALEXANDER 01	CLEO	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K^*$

NODE=M047R10  
 NODE=M047R10  
 → UNCHECKED ←

<sup>1</sup> HASTINGS 03 and AUBERT 04F assume  $\tau(B^+)/\tau(B^0) = 1.083 \pm 0.017$ .

<sup>2</sup> ATHAR 02 assumes  $\tau(B^+)/\tau(B^0) = 1.074 \pm 0.028$ . Supersedes BARISH 95.

<sup>3</sup> AUBERT 02 assumes  $\tau(B^+)/\tau(B^0) = 1.062 \pm 0.029$ .

<sup>4</sup> ALEXANDER 01 assumes  $\tau(B^+)/\tau(B^0) = 1.066 \pm 0.024$ .

NODE=M047R10;LINKAGE=F  
 NODE=M047R10;LINKAGE=D  
 NODE=M047R10;LINKAGE=E  
 NODE=M047R10;LINKAGE=C

 $[\Gamma(J/\psi K_S^0) + \Gamma((J/\psi, \eta_c) K_S^0)]/\Gamma_{\text{total}}$  $\Gamma_5/\Gamma$ 

Forbidden by  $CP$  invariance.

VALUE (units $10^{-7}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;4</b>	90	<sup>1</sup> TAJIMA 07A	BELL	$\Upsilon(4S) \rightarrow B^0 \bar{B}^0$

<sup>1</sup>  $\Upsilon(4S)$  with  $CP = +1$  decays to the final state with  $CP = -1$ .

NODE=M047R16  
 NODE=M047R16  
 NODE=M047R16

NODE=M047R16;LINKAGE=TA

NODE=M047NBB

**non- $B\bar{B}$  DECAYS** $\Gamma(\text{non-}B\bar{B})/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.04</b>	95	BARISH 96B	CLEO	$e^+ e^-$

NODE=M047R6  
 NODE=M047R6

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**1.57 ± 0.08 OUR AVERAGE**

1.55 ± 0.04 ± 0.07	AUBERT 05Q	BABR	$e^+ e^- \rightarrow \text{hadrons}$
2.77 ± 0.50 ± 0.49	<sup>1</sup> ALBRECHT 95E	ARG	$e^+ e^- \rightarrow \text{hadrons}$

<sup>1</sup> Using LEYAQUANC 77 parametrization of  $\Gamma(s)$ .

NODE=M047R5  
 NODE=M047R5

NODE=M047R5;LINKAGE=A

 $\Gamma(\rho^+ \rho^-)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;5.7 × 10<sup>-6</sup></b>	90	AUBERT 08B0	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0$

NODE=M047R17  
 NODE=M047R17

 $\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;2.0 × 10<sup>-6</sup></b>	90	SHEN 13A	BELL	$e^+ e^- \rightarrow K^*(892)^0 \bar{K}^0$

NODE=M047R02  
 NODE=M047R02

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.9</b>	95	<sup>1</sup> ABE 02D	BELL	$e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	90	<sup>1</sup> AUBERT 01c	BABR	$e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$
------	----	-------------------------	------	--

<sup>1</sup> Uses  $B(J/\psi \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$  and  $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$ .

NODE=M047R1  
 NODE=M047R1

NODE=M047R;LINKAGE=AC

 $\Gamma(D^{*+} \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;0.074</b>	90	<sup>1</sup> ALEXANDER 90c	CLEO	$e^+ e^-$

<sup>1</sup> For  $x > 0.473$ .

NODE=M047R2  
 NODE=M047R2

NODE=M047R2;LINKAGE=A

 $\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_{12}/\Gamma$ 

VALUE (units $10^{-2}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>7.1 ± 0.1 ± 0.6</b>		HUANG 07	CLEO	$\Upsilon(4S) \rightarrow \phi X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23	90	<sup>1</sup> ALEXANDER 90c	CLEO	$e^+ e^-$
-------	----	----------------------------	------	-----------

<sup>1</sup> For  $x > 0.52$ .

NODE=M047R3  
 NODE=M047R3

NODE=M047R3;LINKAGE=A

 $\Gamma(\phi \eta)/\Gamma_{\text{total}}$  $\Gamma_{13}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<b>&lt;1.8</b>	90	<sup>1</sup> BELOUS 09	BELL	$e^+ e^- \rightarrow \phi \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.5	90	AUBERT, BE 06F	BABR	$e^+ e^- \rightarrow \phi \eta$
------	----	----------------	------	---------------------------------

<sup>1</sup> Using all intermediate branching fraction values from PDG 08.

NODE=M047R14  
 NODE=M047R14

NODE=M047R14;LINKAGE=BE

$\Gamma(\phi\eta')/\Gamma_{\text{total}}$   $\Gamma_{14}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<4.3	90	<sup>1</sup> BELOUS	09 BELL	$e^+e^- \rightarrow \phi\eta'$

<sup>1</sup> Using all intermediate branching fraction values from PDG 08.

NODE=M047R21  
NODE=M047R21

NODE=M047R21;LINKAGE=BE

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$   $\Gamma_{15}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	<sup>1</sup> BELOUS	09 BELL	$e^+e^- \rightarrow \rho\eta$

<sup>1</sup> Using all intermediate branching fraction values from PDG 08.

NODE=M047R22  
NODE=M047R22

NODE=M047R22;LINKAGE=BE

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$   $\Gamma_{16}/\Gamma$ 

VALUE (units $10^{-6}$ )	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	<sup>1</sup> BELOUS	09 BELL	$e^+e^- \rightarrow \rho\eta'$

<sup>1</sup> Using all intermediate branching fraction values from PDG 08.

NODE=M047R23  
NODE=M047R23

NODE=M047R23;LINKAGE=BE

 $\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{\text{total}}$   $\Gamma_{17}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	90	ALEXANDER	90C CLEO	$e^+e^-$

NODE=M047R4  
NODE=M047R4

 $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{18}/\Gamma$ 

VALUE (units $10^{-5}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>8.2 ± 0.4 OUR AVERAGE</b>					
8.2 ± 0.5 ± 0.4		515	GUIDO	17 BELL	$\Upsilon(4S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
8.5 ± 1.3 ± 0.1	113 ± 16		<sup>1</sup> SOKOLOV	09 BELL	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
8.00 ± 0.64 ± 0.27	430		<sup>2</sup> AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.8 ± 4.0 ± 0.3			<sup>3,4</sup> SOKOLOV	07 BELL	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
9.0 ± 1.5 ± 0.2	167 ± 19		<sup>5</sup> AUBERT	06R BABR	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
<12	90		GLENN	99 CLE2	$e^+e^-$

NODE=M047R7  
NODE=M047R7

NODE=M047R7;LINKAGE=SK

<sup>1</sup> SOKOLOV 09 reports  $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (0.211 \pm 0.030 \pm 0.014) \times 10^{-5}$  which we divide by our best value  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>2</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M047R7;LINKAGE=UB

<sup>3</sup> SOKOLOV 07 reports  $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (4.42 \pm 0.81 \pm 0.56) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R7;LINKAGE=SO

<sup>4</sup> According to the authors, systematic errors were underestimated.

<sup>5</sup> Superseded by AUBERT 08BP. AUBERT 06R reports  $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (2.23 \pm 0.25 \pm 0.27) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R7;LINKAGE=US  
NODE=M047R7;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$   $\Gamma_{19}/\Gamma$ 

VALUE (units $10^{-4}$ )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.81 ± 0.18 OUR AVERAGE</b>					
1.70 ± 0.23 ± 0.08	49		GUIDO	17 BELL	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\mu^+\mu^-$
1.96 ± 0.26 ± 0.09	56		<sup>1</sup> AUBERT	08BP BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.7	90		<sup>2</sup> TAMPONI	15 BELL	$e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
------	----	--	----------------------	---------	--

<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$  and  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ .

NODE=M047R18  
NODE=M047R18

NODE=M047R18;LINKAGE=UB

<sup>2</sup> Using  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$ .

NODE=M047R18;LINKAGE=A

 $\Gamma(\Upsilon(1S)\eta')/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>3.43 ± 0.88 ± 0.21</b>	27	GUIDO	18 BELL	$\Upsilon(4S) \rightarrow (\rho^0\gamma, \pi^+\pi^-\eta)\mu^+\mu^-$

NODE=M047R03  
NODE=M047R03

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$  $\Gamma_{19}/\Gamma_{18}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M047R19  
 NODE=M047R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.41±0.40±0.12 56 <sup>1</sup> AUBERT 08BP BABR  $\Upsilon(4S) \rightarrow \pi^+\pi^-(\pi^0)\ell^+\ell^-$

<sup>1</sup> Not independent of other values reported by AUBERT 08BP.

NODE=M047R19;LINKAGE=UB

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{21}/\Gamma$ 

VALUE (units 10 <sup>-5</sup> )	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

NODE=M047R9  
 NODE=M047R9

**8.2±0.8 OUR AVERAGE**

7.9±1.0±0.4 181 GUIDO 17 BELL  $\Upsilon(4S) \rightarrow \pi^+\pi^-\mu^+\mu^-$

8.6±1.1±0.7 220 <sup>1</sup> AUBERT 08BP BABR  $\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.8±1.7±0.8 97 ± 15 <sup>2</sup> AUBERT 06R BABR  $e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$

<3.9 90 GLENN 99 CLE2  $e^+e^-$

<sup>1</sup> Using  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$  and  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ .

NODE=M047R9;LINKAGE=UB

<sup>2</sup> Superseded by AUBERT 08BP. AUBERT 06R reports  $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = (1.69 \pm 0.26 \pm 0.20) \times 10^{-6}$  which we divide by our best value  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R9;LINKAGE=AU

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$  $\Gamma_{21}/\Gamma_{18}$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M047R20  
 NODE=M047R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.16±0.16±0.14 220 <sup>1</sup> AUBERT 08BP BABR  $\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

<sup>1</sup> Using  $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ ,  $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$ ,  $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$ , and  $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ . Not independent of other values reported by AUBERT 08BP.

NODE=M047R20;LINKAGE=UB

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{22}/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M047R01  
 NODE=M047R01

not seen (35<sup>+32</sup>/<sub>-26</sub>)k <sup>1</sup> ADACHI 12 BELL 10.58  $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

<sup>1</sup> From the upper limit on the ratio of  $\sigma(e^+e^- \rightarrow h_b(1P)\pi^+\pi^-)$  at the  $\Upsilon(4S)$  to that at the  $\Upsilon(5S)$  of 0.27.

NODE=M047R01;LINKAGE=AD

 $\Gamma(h_b(1P)\eta)/\Gamma_{\text{total}}$  $\Gamma_{23}/\Gamma$ 

VALUE (units 10 <sup>-3</sup> )	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

NODE=M047R00  
 NODE=M047R00

2.18±0.11±0.18 112k <sup>1</sup> TAMPONI 15 BELL  $e^+e^- \rightarrow h_b(1P)\eta$

<sup>1</sup> Using  $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$ .

NODE=M047R00;LINKAGE=A

 $\Gamma(\eta_b(1S)\omega)/\Gamma_{\text{total}}$  $\Gamma_{24}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M047R04  
 NODE=M047R04

<1.8 × 10<sup>-4</sup> 90 OSKIN 20 BELL  $e^+e^- \rightarrow \omega X$

 $\Gamma(\eta_b(1S)\omega)/\Gamma(h_b(1P)\eta)$  $\Gamma_{24}/\Gamma_{23}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M047R05  
 NODE=M047R05

<8.4 × 10<sup>-2</sup> 90 <sup>1</sup> OSKIN 20 BELL  $e^+e^- \rightarrow \omega X$

<sup>1</sup> Using  $B(\Upsilon(4S) \rightarrow h_b(1P)\eta) = (2.18 \pm 0.11 \pm 0.18) \times 10^{-3}$  from TAMPONI 15.

NODE=M047R05;LINKAGE=A

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$  $\Gamma_{25}/\Gamma$ 

VALUE (units 10 <sup>-5</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M047R15  
 NODE=M047R15

<1.3 90 ASNER 07 CLEO  $e^+e^- \rightarrow \overline{d}X$

————— Double Radiative Decays —————

NODE=M047240

 $\Gamma(\gamma\gamma \Upsilon(D) \rightarrow \gamma\gamma\eta \Upsilon(1S))/\Gamma_{\text{total}}$  $\Gamma_{26}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M047R24  
 NODE=M047R24

<2.3 × 10<sup>-5</sup> 90 GUIDO 17 BELL  $\Upsilon(4S) \rightarrow \gamma\gamma\pi^+\pi^-\pi^0\mu^+\mu^-$

$\Upsilon(4S)$  REFERENCES

OSKIN	20	PR D102 092011	P. Oskin <i>et al.</i>	(BELLE Collab.)	REFID=60735
GUIDO	18	PRL 121 062001	E. Guido <i>et al.</i>	(BELLE Collab.)	REFID=58860
GUIDO	17	PR D96 052005	E. Guido <i>et al.</i>	(BELLE Collab.)	REFID=58218
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=56996
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
BELOUS	09	PL B681 400	K. Belous <i>et al.</i>	(BELLE Collab.)	REFID=53107
SOKOLOV	09	PR D79 051103	A. Sokolov <i>et al.</i>	(BELLE Collab.)	REFID=52760
AUBERT	08BO	PR D78 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52659
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
SOKOLOV	07	PR D75 071103	A. Sokolov <i>et al.</i>	(BELLE Collab.)	REFID=51715
TAJIMA	07A	PRL 99 211601	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=52066
AUBERT	06R	PRL 96 232001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51143
AUBERT,BE	06F	PR D74 111103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51563
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
AUBERT	05Q	PR D72 032005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50774
AUBERT,B	05H	PRL 95 042001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50777
AUBERT	04F	PR D69 071101	B. Aubert <i>et al.</i>		REFID=49748
HASTINGS	03	PR D67 052004	N.C. Hastings <i>et al.</i>	(BELLE Collab.)	REFID=49209
ABE	02D	PRL 88 052001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48557
ATHAR	02	PR D66 052003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=48832
AUBERT	02	PR D65 032001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48514
ALEXANDER	01	PRL 86 2737	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48316
AUBERT	01C	PRL 87 162002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48346
GLENN	99	PR D59 052003	S. Glenn <i>et al.</i>		REFID=46890
BARISH	96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44693
ALBRECHT	95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44372
BARISH	95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44139
ALEXANDER	90C	PRL 64 2226	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=41346
BEBEK	87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=40270
BESSION	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSP Collab.)	REFID=22369
LEYAOUANC	77	PL B71 397	A. Le Yaouanc <i>et al.</i>	(ORSAY)	REFID=44695

NODE=M243

 $\Upsilon(10753)$ 

$$I^G(J^{PC}) = ?(1^{--})$$

OMITTED FROM SUMMARY TABLE

A candidate for  $\Upsilon(3D)$  state or an exotic structure.

NODE=M243

Seen by MIZUK 19 in  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$  ( $n=1,2,3$ ) with a significance of  $5.2\sigma$ . $\Upsilon(10753)$  MASS

NODE=M243M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$10752.7 \pm 5.9^{+0.7}_{-1.1}$	<sup>1</sup> MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

NODE=M243M

••• We do not use the following data for averages, fits, limits, etc. •••

10761 $\pm 2$	<sup>2</sup> DONG	20A	$e^+e^- \rightarrow b\bar{b}$
---------------	-------------------	-----	-------------------------------

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.63\text{--}11.02$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M243M;LINKAGE=A

<sup>2</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M243M;LINKAGE=B

 $\Upsilon(10753)$  WIDTH

NODE=M243W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$35.5^{+17.6+3.9}_{-11.3-3.3}$	<sup>1</sup> MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

NODE=M243W

••• We do not use the following data for averages, fits, limits, etc. •••

48.5 $\pm 3.0$	<sup>2</sup> DONG	20A	$e^+e^- \rightarrow b\bar{b}$
----------------	-------------------	-----	-------------------------------

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.63\text{--}11.02$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M243W;LINKAGE=A

<sup>2</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M243W;LINKAGE=B

**$\Upsilon(10753)$  DECAY MODES**

NODE=M243215;NODE=M243

Mode					
$\Gamma_1$	$\Upsilon(1S)\pi^+\pi^-$				DESIG=3
$\Gamma_2$	$\Upsilon(2S)\pi^+\pi^-$				DESIG=4
$\Gamma_3$	$\Upsilon(3S)\pi^+\pi^-$				DESIG=5
$\Gamma_4$	$e^+e^-$				DESIG=2
<b><math>\Upsilon(10753) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})</math></b>					
<b><math>\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_1\Gamma_4/\Gamma</math></b>
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M243225
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.295±0.175	1,2 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$		NODE=M243R00
			<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$ .		NODE=M243R00;LINKAGE=A
			<sup>2</sup> Reported as the range 0.12–0.47 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.		NODE=M243R00;LINKAGE=B
<b><math>\Gamma(\Upsilon(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_2\Gamma_4/\Gamma</math></b>
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M243R02
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.875±0.345	1,2 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$		NODE=M243R02
			<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$ .		NODE=M243R02;LINKAGE=A
			<sup>2</sup> Reported as the range 0.53–1.22 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.		NODE=M243R02;LINKAGE=B
<b><math>\Gamma(\Upsilon(3S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}</math></b>					<b><math>\Gamma_3\Gamma_4/\Gamma</math></b>
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M243R03
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.235±0.025	1,2 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$		NODE=M243R03
			<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$ .		NODE=M243R03;LINKAGE=A
			<sup>2</sup> Reported as the range 0.21–0.26 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.		NODE=M243R03;LINKAGE=B

 **$\Upsilon(10753)$  REFERENCES**

NODE=M243

DONG	20A	CP C44 083001	X.-K. Dong <i>et al.</i>		REFID=60595
MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60090
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661

$\Upsilon(10860)$ 

$$J^{PC} = 0^{-}(1^{-}-)$$

NODE=M092

 $\Upsilon(10860)$  MASS

NODE=M092M

NODE=M092M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>10885.2^{+2.6}_{-1.6}</math> OUR AVERAGE</b>			
$10885.3 \pm 1.5^{+2.2}_{-0.9}$	1 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
$10884.7^{+3.6+8.9}_{-3.4-1.0}$	2 MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P,2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$10882 \pm 1$	<sup>3</sup> DONG	20A	$e^+e^- \rightarrow b\bar{b}$
$10881.8^{+1.0}_{-1.1} \pm 1.2$	4,5 SANTEL	16	BELL $e^+e^- \rightarrow \text{hadrons}$
$10891.1 \pm 3.2^{+1.2}_{-2.0}$	6,7 SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S,2S,3S)\pi^+\pi^-$
$10879 \pm 3$	8,9 CHEN	10	BELL $e^+e^- \rightarrow \text{hadrons}$
$10888.4^{+2.7}_{-2.6} \pm 1.2$	<sup>10</sup> CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S,2S,3S)\pi^+\pi^-$
$10876 \pm 2$	<sup>8</sup> AUBERT	09E	BABR $e^+e^- \rightarrow \text{hadrons}$
$10869 \pm 2$	<sup>11</sup> AUBERT	09E	BABR $e^+e^- \rightarrow \text{hadrons}$
$10868 \pm 6 \pm 5$	<sup>12</sup> BESSON	85	CLEO $e^+e^- \rightarrow \text{hadrons}$
$10845 \pm 20$	<sup>13</sup> LOVELOCK	85	CUSB $e^+e^- \rightarrow \text{hadrons}$

OCCUR=2

OCCUR=2

OCCUR=2

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M092M;LINKAGE=E

<sup>2</sup> From a simultaneous fit to the  $h_b(nP)\pi^+\pi^-$ ,  $n = 1, 2$  cross sections at 22 energy points within  $\sqrt{s} = 10.77\text{--}11.02$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single relative phase, a single relative amplitude, and two overall normalization factors, one for each  $n$ ). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

NODE=M092M;LINKAGE=D

<sup>3</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M092M;LINKAGE=H

<sup>4</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).

NODE=M092M;LINKAGE=A

<sup>5</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions.

NODE=M092M;LINKAGE=B

<sup>6</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M092M;LINKAGE=C

<sup>7</sup> Superseded by MIZUK 19.

NODE=M092M;LINKAGE=F

<sup>8</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M092M;LINKAGE=AU

<sup>9</sup> The parameters of the  $\Upsilon(11020)$  are fixed to those in AUBERT 09E.

NODE=M092M;LINKAGE=CH

<sup>10</sup> In a model where a flat nonresonant  $\Upsilon(1S,2S,3S)\pi^+\pi^-$  continuum interferes with a single Breit-Wigner resonance.

NODE=M092M;LINKAGE=CE

<sup>11</sup> In a model where a non-resonant  $b\bar{b}$ -continuum represented by a threshold function at  $\sqrt{s}=2m_B$  is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.

NODE=M092M;LINKAGE=UB

<sup>12</sup> Assuming four Gaussians with radiative tails and a single step in  $R$ .

NODE=M092M;LINKAGE=BE

<sup>13</sup> In a coupled-channel model with three resonances and a smooth step in  $R$ .

NODE=M092M;LINKAGE=LO

 $\Upsilon(10860)$  WIDTH

NODE=M092W

NODE=M092W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>37 \pm 4</math> OUR AVERAGE</b>			
$36.6^{+4.5+0.5}_{-3.9-1.1}$	1 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
$40.6^{+12.7+1.1}_{-8.0-19.1}$	2 MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P,2P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

49.5 ± 1.5	<sup>3</sup> DONG	20A	$e^+e^- \rightarrow b\bar{b}$	
48.5 <sup>+1.9+2.0</sup> <sub>-1.8-2.8</sub>	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow \text{hadrons}$	
53.7 <sup>+7.1+1.3</sup> <sub>-5.6-5.4</sub>	6,7 SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$	OCCUR=2
46 <sup>+9</sup> <sub>-7</sub>	8,9 CHEN	10 BELL	$e^+e^- \rightarrow \text{hadrons}$	
30.7 <sup>+8.3</sup> <sub>-7.0</sub> ± 3.1	<sup>10</sup> CHEN	10 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$	OCCUR=2
43 ± 4	<sup>8</sup> AUBERT	09E BABR	$e^+e^- \rightarrow \text{hadrons}$	
74 ± 4	<sup>11</sup> AUBERT	09E BABR	$e^+e^- \rightarrow \text{hadrons}$	OCCUR=2
112 ± 17 ± 23	<sup>12</sup> BESSON	85 CLEO	$e^+e^- \rightarrow \text{hadrons}$	
110 ± 15	<sup>13</sup> LOVELOCK	85 CUSB	$e^+e^- \rightarrow \text{hadrons}$	
<sup>1</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$ .				NODE=M092W;LINKAGE=F
<sup>2</sup> From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$ , $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$ , a single relative phase, a single relative amplitude, and two overall normalization factors, one for each $n$ ). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.				NODE=M092W;LINKAGE=D
<sup>3</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.				NODE=M092W;LINKAGE=G
<sup>4</sup> From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).				NODE=M092W;LINKAGE=A
<sup>5</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.				NODE=M092W;LINKAGE=B
<sup>6</sup> From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$ , $n = 1, 2, 3$ , cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.				NODE=M092W;LINKAGE=C
<sup>7</sup> Superseded by MIZUK 19.				NODE=M092W;LINKAGE=E
<sup>8</sup> In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.				NODE=M092W;LINKAGE=AU
<sup>9</sup> The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.				NODE=M092W;LINKAGE=CH
<sup>10</sup> In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.				NODE=M092W;LINKAGE=CE
<sup>11</sup> In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.				NODE=M092W;LINKAGE=UB
<sup>12</sup> Assuming four Gaussians with radiative tails and a single step in $R$ .				NODE=M092W;LINKAGE=BE
<sup>13</sup> In a coupled-channel model with three resonances and a smooth step in $R$ .				NODE=M092W;LINKAGE=LO

## $\Upsilon(10860)$ DECAY MODES

NODE=M092215;NODE=M092

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Confidence level	
$\Gamma_1$ $B\bar{B}X$	( 76.2 <sup>+2.7</sup> <sub>-4.0</sub> ) %		DESIG=9
$\Gamma_2$ $B\bar{B}$	( 5.5 ± 1.0 ) %		DESIG=2
$\Gamma_3$ $B\bar{B}^* + \text{c.c.}$	( 13.7 ± 1.6 ) %		DESIG=3
$\Gamma_4$ $B^*\bar{B}^*$	( 38.1 ± 3.4 ) %		DESIG=4
$\Gamma_5$ $B\bar{B}^{(*)}\pi$	< 19.7 %	90%	DESIG=10
$\Gamma_6$ $B\bar{B}\pi$	( 0.0 ± 1.2 ) %		DESIG=23
$\Gamma_7$ $B^*\bar{B}\pi + B\bar{B}^*\pi$	( 7.3 ± 2.3 ) %		DESIG=24
$\Gamma_8$ $B^*\bar{B}^*\pi$	( 1.0 ± 1.4 ) %		DESIG=25
$\Gamma_9$ $B\bar{B}\pi\pi$	< 8.9 %	90%	DESIG=11
$\Gamma_{10}$ $B_s^{(*)}\bar{B}_s^{(*)}$	( 20.1 ± 3.1 ) %		DESIG=16
$\Gamma_{11}$ $B_s\bar{B}_s$	( 5 ± 5 ) × 10 <sup>-3</sup>		DESIG=5
$\Gamma_{12}$ $B_s\bar{B}_s^* + \text{c.c.}$	( 1.35 ± 0.32 ) %		DESIG=7
$\Gamma_{13}$ $B_s^*\bar{B}_s^*$	( 17.6 ± 2.7 ) %		DESIG=8
$\Gamma_{14}$ no open-bottom	( 3.8 <sup>+5.0</sup> <sub>-0.5</sub> ) %		DESIG=28

$\Gamma_{15}$	$e^+ e^-$	$( 8.3 \pm 2.1 ) \times 10^{-6}$		DESIG=1
$\Gamma_{16}$	$K^*(892)^0 \bar{K}^0$	$< 1.0 \times 10^{-5}$	90%	DESIG=29
$\Gamma_{17}$	$\Upsilon(1S) \pi^+ \pi^-$	$( 5.3 \pm 0.6 ) \times 10^{-3}$		DESIG=17
$\Gamma_{18}$	$\Upsilon(1S) \eta$	$( 8.5 \pm 1.7 ) \times 10^{-4}$		DESIG=44
$\Gamma_{19}$	$\Upsilon(1S) \eta'$	$< 6.9 \times 10^{-5}$	90%	DESIG=45
$\Gamma_{20}$	$\Upsilon(2S) \pi^+ \pi^-$	$( 7.8 \pm 1.3 ) \times 10^{-3}$		DESIG=18
$\Gamma_{21}$	$\Upsilon(2S) \eta$	$( 4.1 \pm 0.6 ) \times 10^{-3}$		DESIG=46
$\Gamma_{22}$	$\Upsilon(3S) \pi^+ \pi^-$	$( 4.8 \begin{smallmatrix} +1.9 \\ -1.7 \end{smallmatrix} ) \times 10^{-3}$		DESIG=19
$\Gamma_{23}$	$\Upsilon(1S) K^+ K^-$	$( 6.1 \pm 1.8 ) \times 10^{-4}$		DESIG=20
$\Gamma_{24}$	$\eta \Upsilon_J(1D)$	$( 4.8 \pm 1.1 ) \times 10^{-3}$		DESIG=40
$\Gamma_{25}$	$h_b(1P) \pi^+ \pi^-$	$( 3.5 \begin{smallmatrix} +1.0 \\ -1.3 \end{smallmatrix} ) \times 10^{-3}$		DESIG=26
$\Gamma_{26}$	$h_b(2P) \pi^+ \pi^-$	$( 5.7 \begin{smallmatrix} +1.7 \\ -2.1 \end{smallmatrix} ) \times 10^{-3}$		DESIG=27
$\Gamma_{27}$	$\chi_{bJ}(1P) \pi^+ \pi^- \pi^0$	$( 2.5 \pm 2.3 ) \times 10^{-3}$		DESIG=41
$\Gamma_{28}$	$\chi_{b0}(1P) \pi^+ \pi^- \pi^0$	$< 6.3 \times 10^{-3}$	90%	DESIG=30
$\Gamma_{29}$	$\chi_{b0}(1P) \omega$	$< 3.9 \times 10^{-3}$	90%	DESIG=31
$\Gamma_{30}$	$\chi_{b0}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	$< 4.8 \times 10^{-3}$	90%	DESIG=32
$\Gamma_{31}$	$\chi_{b1}(1P) \pi^+ \pi^- \pi^0$	$( 1.85 \pm 0.33 ) \times 10^{-3}$		DESIG=33
$\Gamma_{32}$	$\chi_{b1}(1P) \omega$	$( 1.57 \pm 0.30 ) \times 10^{-3}$		DESIG=34
$\Gamma_{33}$	$\chi_{b1}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	$( 5.2 \pm 1.9 ) \times 10^{-4}$		DESIG=35
$\Gamma_{34}$	$\chi_{b2}(1P) \pi^+ \pi^- \pi^0$	$( 1.17 \pm 0.30 ) \times 10^{-3}$		DESIG=36
$\Gamma_{35}$	$\chi_{b2}(1P) \omega$	$( 6.0 \pm 2.7 ) \times 10^{-4}$		DESIG=37
$\Gamma_{36}$	$\chi_{b2}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	$( 6 \pm 4 ) \times 10^{-4}$		DESIG=38
$\Gamma_{37}$	$\gamma X_b \rightarrow \gamma \Upsilon(1S) \omega$	$< 3.8 \times 10^{-5}$	90%	DESIG=39
$\Gamma_{38}$	$\eta_b(1S) \omega$	$< 1.3 \times 10^{-3}$	90%	DESIG=42
$\Gamma_{39}$	$\eta_b(2S) \omega$	$< 5.6 \times 10^{-3}$	90%	DESIG=43

**Inclusive Decays.**

NODE=M092;CLUMP=I

These decay modes are submodes of one or more of the decay modes above.

NODE=M092

$\Gamma_{40}$	$\phi$ anything	$( 13.8 \begin{smallmatrix} +2.4 \\ -1.7 \end{smallmatrix} ) \%$		DESIG=12
$\Gamma_{41}$	$D^0$ anything + c.c.	$( 108 \pm 8 ) \%$		DESIG=13
$\Gamma_{42}$	$D_s$ anything + c.c.	$( 46 \pm 6 ) \%$		DESIG=6
$\Gamma_{43}$	$J/\psi$ anything	$( 2.06 \pm 0.21 ) \%$		DESIG=14
$\Gamma_{44}$	$B^0$ anything + c.c.	$( 77 \pm 8 ) \%$		DESIG=21
$\Gamma_{45}$	$B^+$ anything + c.c.	$( 72 \pm 6 ) \%$		DESIG=22

 **$\Upsilon(10860)$  PARTIAL WIDTHS**

NODE=M092220

 **$\Gamma(e^+ e^-)$**  **$\Gamma_{15}$** 

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
<b>0.31 <math>\pm</math> 0.07 OUR AVERAGE</b>	Error includes scale factor of 1.3.		
0.22 $\pm$ 0.05 $\pm$ 0.07	BESSION	85	CLEO $e^+ e^- \rightarrow$ hadrons
0.365 $\pm$ 0.070	LOVELOCK	85	CUSB $e^+ e^- \rightarrow$ hadrons

NODE=M092W1  
NODE=M092W1 **$\Gamma(e^+ e^-) \times \Gamma(\Upsilon(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}$**  **$\Gamma_{15} \Gamma_{17} / \Gamma$** 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.09 $\pm$ 0.34	<sup>1,2</sup> MIZUK	19	BELL $e^+ e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-$

NODE=M092R50  
NODE=M092R50

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS) \pi^+ \pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6$ –11.05 GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M092R50;LINKAGE=A

<sup>2</sup> Reported as the range 0.75–1.43 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M092R50;LINKAGE=B

 **$\Gamma(e^+ e^-) \times \Gamma(\Upsilon(2S) \pi^+ \pi^-) / \Gamma_{\text{total}}$**  **$\Gamma_{15} \Gamma_{20} / \Gamma$** 

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.58 $\pm$ 1.22	<sup>1,2</sup> MIZUK	19	BELL $e^+ e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-$

NODE=M092R51  
NODE=M092R51

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS) \pi^+ \pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6$ –11.05 GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M092R51;LINKAGE=A

<sup>2</sup> Reported as the range 1.35–3.80 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M092R51;LINKAGE=B



$$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_{22}/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M092R52  
NODE=M092R52

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.73±0.30	1,2 MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
-----------	-----------	----	------	---

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M092R52;LINKAGE=A

<sup>2</sup> Reported as the range 0.43–1.03 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M092R52;LINKAGE=B

## $\Upsilon(10860)$ BRANCHING RATIOS

NODE=M092230

"OUR EVALUATION" is obtained based on averages of rescaled data listed below. The averages and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <https://hflav.web.cern.ch/>.

NODE=M092230

$$\Gamma(B\bar{B}X)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M092R13  
NODE=M092R13

**0.762<sup>+0.027</sup><sub>-0.043</sub> OUR EVALUATION**

→ UNCHECKED ←

**0.71 ± 0.06 OUR AVERAGE**

0.737±0.032±0.051	1063	<sup>1</sup> DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
-------------------	------	-----------------------	----	------	---------------------------------------

0.589±0.100±0.092		<sup>2</sup> HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------------------	--	--------------------	----	------	---

<sup>1</sup> Not independent of DRUTSKOY 10 values for  $\Upsilon(5S) \rightarrow B^{\pm,0}$  anything.

NODE=M092R13;LINKAGE=DR

<sup>2</sup> Using measurements or limits from AQUINES 06.

NODE=M092R13;LINKAGE=HU

$$\Gamma(B\bar{B})/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE (units 10 <sup>-2</sup> )	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M092R16  
NODE=M092R16

<b>5.5<sup>+1.0</sup><sub>-0.9</sub> ± 0.4</b>		<sup>1</sup> DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
--	--	-----------------------	----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.8	90	<sup>2</sup> HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------	----	--------------------	----	------	---

<sup>1</sup> Assuming isospin conservation.

NODE=M092R16;LINKAGE=DR

<sup>2</sup> Using measurements or limits from AQUINES 06.

NODE=M092R16;LINKAGE=HU

$$\Gamma(B\bar{B})/\Gamma(B\bar{B}X) \quad \Gamma_2/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M092R05  
NODE=M092R05

<b>&lt;0.22</b>	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$
-----------------	----	---------	----	------	---

$$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M092R15  
NODE=M092R15

**0.137±0.016 OUR AVERAGE**

0.137±0.013±0.011		<sup>1</sup> DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
-------------------	--	-----------------------	----	------	---------------------------------------

0.143±0.053±0.027		<sup>2</sup> HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------------------	--	--------------------	----	------	---

<sup>1</sup> Assuming isospin conservation.

NODE=M092R15;LINKAGE=DR

<sup>2</sup> Using measurements or limits from AQUINES 06.

NODE=M092R15;LINKAGE=HU

$$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma(B\bar{B}X) \quad \Gamma_3/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M092R06  
NODE=M092R06

<b>0.24±0.09±0.03</b>	10	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$
-----------------------	----	---------	----	------	---

$$\Gamma(B^*\bar{B}^*)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M092R14  
NODE=M092R14

**0.381±0.034 OUR AVERAGE**

0.375 <sup>+0.021</sup> <sub>-0.019</sub> ± 0.030		<sup>1</sup> DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
---	--	-----------------------	----	------	---------------------------------------

0.436±0.083±0.072		<sup>2</sup> HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------------------	--	--------------------	----	------	---

<sup>1</sup> Assuming isospin conservation.

NODE=M092R14;LINKAGE=DR

<sup>2</sup> Using measurements or limits from AQUINES 06.

NODE=M092R14;LINKAGE=HU

$$\Gamma(B^*\bar{B}^*)/\Gamma(B\bar{B}X) \quad \Gamma_4/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M092R07  
NODE=M092R07

<b>0.74±0.15±0.08</b>	31	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$
-----------------------	----	---------	----	------	---

$$\Gamma(B\bar{B}^{(*)}\pi)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M092R17  
NODE=M092R17

<b>&lt;0.197</b>	90	<sup>1</sup> HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
------------------	----	--------------------	----	------	---

<sup>1</sup> Using measurements or limits from AQUINES 06.

NODE=M092R17;LINKAGE=HU

$\Gamma(B\bar{B}^*\pi)/\Gamma(B\bar{B}X)$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma_1$
<b>&lt;0.32</b>	90	AQUINES	06	CLE3 $\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092R08  
NODE=M092R08

 $\Gamma(B\bar{B}\pi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma$
<b><math>0.0 \pm 1.2 \pm 0.3</math></b>	0	<sup>1</sup> DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$	

NODE=M092R28  
NODE=M092R28

<sup>1</sup> Assuming isospin conservation.

NODE=M092R28;LINKAGE=DR

 $[\Gamma(B^*\bar{B}\pi) + \Gamma(B\bar{B}^*\pi)]/\Gamma_{\text{total}}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma$
<b><math>7.3^{+2.3}_{-2.1} \pm 0.8</math></b>	38	<sup>1</sup> DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$	

NODE=M092R29  
NODE=M092R29

<sup>1</sup> Assuming isospin conservation.

NODE=M092R29;LINKAGE=DR

 $\Gamma(B^*\bar{B}^*\pi)/\Gamma_{\text{total}}$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_8/\Gamma$
<b><math>1.0^{+1.4}_{-1.3} \pm 0.4</math></b>	5	<sup>1</sup> DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+,0}\pi^- X$	

NODE=M092R30  
NODE=M092R30

<sup>1</sup> Assuming isospin conservation.

NODE=M092R30;LINKAGE=DR

 $\Gamma(B\bar{B}\pi\pi)/\Gamma_{\text{total}}$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma$
<b>&lt;0.089</b>	90	<sup>1</sup> HUANG	07	CLEO $\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092R18  
NODE=M092R18

<sup>1</sup> Using measurements or limits from AQUINES 06.

NODE=M092R18;LINKAGE=HU

 $\Gamma(B\bar{B}\pi\pi)/\Gamma(B\bar{B}X)$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma_1$
<b>&lt;0.14</b>	90	AQUINES	06	CLE3 $\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092R09  
NODE=M092R09

 $\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma_{\text{total}}$ 

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{10}/\Gamma = (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})/\Gamma$
-------	-------------	------	---------	---

NODE=M092R01  
NODE=M092R01

**$0.201^{+0.030}_{-0.031}$  OUR EVALUATION**

→ UNCHECKED ←

**$0.189^{+0.027}_{-0.021}$  OUR AVERAGE**

0.172 ± 0.030 <sup>1</sup> ESEN 13 BELL  $\Upsilon(5S) \rightarrow D^0 X, D_s X$

0.21  $^{+0.06}_{-0.03}$  <sup>2</sup> HUANG 07 CLEO  $\Upsilon(5S) \rightarrow D_s X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.180 ± 0.013 ± 0.032 <sup>3</sup> DRUTSKOY 07 BELL  $\Upsilon(5S) \rightarrow D^0 X, D_s X$

0.160 ± 0.026 ± 0.058 <sup>4</sup> ARTUSO 05B CLEO  $e^+ e^- \rightarrow D_s X$

<sup>1</sup> Supersedes DRUTSKOY 07.

<sup>2</sup> Supersedes ARTUSO 05B. Combining inclusive  $\phi$ ,  $D_s$ , and  $B$  measurements. Using  $B(D_s^+ \rightarrow \phi\pi^+) = 4.4 \pm 0.6\%$  from PDG 06.

<sup>3</sup> Using  $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6)\%$  from PDG 06.

<sup>4</sup> Uses a model-dependent estimate  $B(B_s \rightarrow D_s X) = (92 \pm 11)\%$ .

NODE=M092R01;LINKAGE=ES  
NODE=M092R01;LINKAGE=HU

NODE=M092R01;LINKAGE=DR  
NODE=M092R01;LINKAGE=AR

 $\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma(B\bar{B}X)$ 

VALUE	DOCUMENT ID	$\Gamma_{10}/\Gamma_1$
-------	-------------	------------------------

NODE=M092R34  
NODE=M092R34

**$0.264^{+0.052}_{-0.045}$  OUR EVALUATION**

→ UNCHECKED ←

 $\Gamma(B_s^*\bar{B}_s^*)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{13}/\Gamma_{10} = \Gamma_{13}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$
--------------------------	------	-------------	------	---------	---

NODE=M092R19  
NODE=M092R19

**$87.8 \pm 1.5$  OUR AVERAGE**

87.0 ± 1.7 <sup>1,2</sup> ESEN 13 BELL  $B_s^0 \rightarrow D_s^- \pi^+$

90.5 ± 3.2 ± 0.1 <sup>2,3</sup> LI 12 BELL  $B_s^0 \rightarrow J/\psi\eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

90.1  $^{+3.8}_{-4.0} \pm 0.2$  <sup>4</sup> LOUVOT 09 BELL  $10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

93  $^{+7}_{-9} \pm 1$  <sup>4</sup> DRUTSKOY 07A BELL Superseded by LOUVOT 09

<sup>1</sup>Supersedes LOUVOT 09.

<sup>2</sup>With  $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$ .

<sup>3</sup>The ratios  $N(B_s^{(*)}\bar{B}_s^{(*)}) / N(B_s^{(*)}\bar{B}_s^{(*)})$  and  $N(B_s^{(*)}\bar{B}_s^0) / N(B_s^{(*)}\bar{B}_s^{(*)})$  are measured with a correlation coefficient of  $-0.72$ .

<sup>4</sup>From a measurement of  $\sigma(e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}) / \sigma(e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)})$  at  $\sqrt{s} = 10.86$  GeV.

$$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

$$\Gamma_{11}/\Gamma_{10} = \Gamma_{11}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$$

VALUE (units  $10^{-2}$ )

DOCUMENT ID

TECN

COMMENT

$2.6^{+2.6}_{-2.5}$

LOUVOT

09

BELL

10.86  $e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

NODE=M092R19;LINKAGE=ES

NODE=M092R19;LINKAGE=IL

NODE=M092R19;LINKAGE=L1

NODE=M092R19;LINKAGE=DR

NODE=M092R24

NODE=M092R24

$$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

$$\Gamma_{11}/\Gamma_{13}$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

$<0.16$

90

BONVICINI

06

CLE3

$e^+e^-$

NODE=M092R03

NODE=M092R03

$$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

$$\Gamma_{12}/\Gamma_{10} = \Gamma_{12}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$$

VALUE (units  $10^{-2}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**6.7 $\pm$ 1.2 OUR AVERAGE**

7.3 $\pm$ 1.4

<sup>1,2</sup>ESEN

13

BELL

$B_s^0 \rightarrow D_s^- \pi^+$

4.9 $\pm$ 2.5 $\pm$ 0.0

227

<sup>2,3</sup>LI

12

BELL

$B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.3 $^{+3.3}_{-3.0}$  $\pm$ 0.1

LOUVOT

09

BELL

10.86  $e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

NODE=M092R25

NODE=M092R25

<sup>1</sup>Supersedes LOUVOT 09.

<sup>2</sup>With  $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$ .

<sup>3</sup>The ratios  $N(B_s^{(*)}\bar{B}_s^{(*)}) / N(B_s^{(*)}\bar{B}_s^{(*)})$  and  $N(B_s^{(*)}\bar{B}_s^0) / N(B_s^{(*)}\bar{B}_s^{(*)})$  are measured with a correlation coefficient of  $-0.72$ .

$$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$$

$$\Gamma_{12}/\Gamma_{13}$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

$<0.16$

90

BONVICINI

06

CLE3

$e^+e^-$

NODE=M092R04

NODE=M092R04

$$\Gamma(\text{no open-bottom})/\Gamma_{\text{total}}$$

$$\Gamma_{14}/\Gamma$$

VALUE

DOCUMENT ID

**0.038 $^{+0.051}_{-0.005}$  OUR EVALUATION**

NODE=M092R33

NODE=M092R33

→ UNCHECKED ←

$$\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma_{\text{total}}$$

$$\Gamma_{16}/\Gamma$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

$<1.0 \times 10^{-5}$

90

SHEN

13A

BELL

$e^+e^- \rightarrow K^*(892)^0 \bar{K}^0$

NODE=M092R35

NODE=M092R35

$$\Gamma(\eta \mathcal{T}_J(1D))/\Gamma_{\text{total}}$$

$$\Gamma_{24}/\Gamma$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**4.82 $\pm$ 0.92 $\pm$ 0.67**

<sup>1</sup>TAMPONI

18

BELL

$e^+e^- \rightarrow \mathcal{T}(5S) \rightarrow \eta X$

NODE=M092R48

NODE=M092R48

<sup>1</sup>Mainly  $J = 2$ , assumes no continuum contribution under  $\mathcal{T}(5S)$ .

NODE=M092R48;LINKAGE=A

$$\Gamma(\mathcal{T}(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma$$

VALUE (units  $10^{-3}$ )

EVTS

DOCUMENT ID

TECN

COMMENT

**5.3 $\pm$ 0.3 $\pm$ 0.5**

325

<sup>1</sup>CHEN

08

BELL

10.87  $e^+e^- \rightarrow \mathcal{T}(1S)\pi^+\pi^-$

NODE=M092R20

NODE=M092R20

<sup>1</sup>Assuming that the observed events are solely due to the  $\mathcal{T}(5S)$  resonance.

NODE=M092R20;LINKAGE=CH

$$\Gamma(\mathcal{T}(1S)\eta)/\Gamma_{\text{total}}$$

$$\Gamma_{18}/\Gamma$$

VALUE (units  $10^{-3}$ )

DOCUMENT ID

TECN

COMMENT

**0.85 $\pm$ 0.15 $\pm$ 0.08**

<sup>1,2</sup>KOVALENKO

21

BELL

$e^+e^- \rightarrow \mathcal{T}(5S)$

NODE=M092R55

NODE=M092R55

<sup>1</sup>Assuming that the observed events are solely due to the  $\mathcal{T}(5S)$  resonance.

NODE=M092R55;LINKAGE=A

<sup>2</sup>Using a data sample of 118.3 fb $^{-1}$  of  $e^+e^-$  collisions at  $\sqrt{s} = 10.866$  GeV.

NODE=M092R55;LINKAGE=B

$$\Gamma(\mathcal{T}(1S)\eta')/\Gamma_{\text{total}}$$

$$\Gamma_{19}/\Gamma$$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

$<6.9 \times 10^{-5}$

90

<sup>1,2</sup>KOVALENKO

21

BELL

$e^+e^- \rightarrow \mathcal{T}(5S)$

NODE=M092R56

NODE=M092R56

<sup>1</sup>Assuming that the observed events are solely due to the  $\mathcal{T}(5S)$  resonance.

NODE=M092R56;LINKAGE=A

<sup>2</sup>Using a data sample of 118.3 fb $^{-1}$  of  $e^+e^-$  collisions at  $\sqrt{s} = 10.866$  GeV.

NODE=M092R56;LINKAGE=B

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{20}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$7.8 \pm 0.6 \pm 1.1$	186	<sup>1</sup> CHEN 08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

NODE=M092R21  
 NODE=M092R21

<sup>1</sup> Assuming that the observed events are solely due to the  $\Upsilon(5S)$  resonance.

NODE=M092R21;LINKAGE=CH

 $\Gamma(\Upsilon(2S)\eta)/\Gamma_{\text{total}}$   $\Gamma_{21}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
$4.13 \pm 0.41 \pm 0.37$	<sup>1,2</sup> KOVALENKO 21	BELL	$e^+ e^- \rightarrow \Upsilon(5S)$

NODE=M092R57  
 NODE=M092R57

<sup>1</sup> Assuming that the observed events are solely due to the  $\Upsilon(5S)$  resonance.

NODE=M092R57;LINKAGE=A

<sup>2</sup> Using a data sample of  $118.3 \text{ fb}^{-1}$  of  $e^+ e^-$  collisions at  $\sqrt{s} = 10.866 \text{ GeV}$ .

NODE=M092R57;LINKAGE=B

 $\Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{22}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$4.8 \pm 1.8 \pm 0.7$	10	<sup>1</sup> CHEN 08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

NODE=M092R22  
 NODE=M092R22

<sup>1</sup> Assuming that the observed events are solely due to the  $\Upsilon(5S)$  resonance.

NODE=M092R22;LINKAGE=CH

 $\Gamma(\Upsilon(1S)K^+K^-)/\Gamma_{\text{total}}$   $\Gamma_{23}/\Gamma$ 

VALUE (units $10^{-4}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$6.1 \pm 1.6 \pm 1.0$	20	<sup>1</sup> CHEN 08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(1S)K^+K^-$

NODE=M092R23  
 NODE=M092R23

<sup>1</sup> Assuming that the observed events are solely due to the  $\Upsilon(5S)$  resonance.

NODE=M092R23;LINKAGE=CH

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$   $\Gamma_{25}/\Gamma_{20}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.45 \pm 0.08 \pm 0.07$ $-0.12$	ADACHI 12	BELL	$10.86 e^+ e^- \rightarrow \text{hadrons}$

NODE=M092R31  
 NODE=M092R31

 $\Gamma(h_b(2P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$   $\Gamma_{26}/\Gamma_{20}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.77 \pm 0.08 \pm 0.22$ $-0.17$	ADACHI 12	BELL	$10.86 e^+ e^- \rightarrow \text{hadrons}$

NODE=M092R32  
 NODE=M092R32

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(h_b(2P)\pi^+\pi^-)$   $\Gamma_{25}/\Gamma_{26}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.616 \pm 0.052 \pm 0.017$	MIZUK 16	BELL	$e^+ e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$

NODE=M092R00  
 NODE=M092R00

 $\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{27}/\Gamma$ 

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.6 \pm 2.2$	YIN 18	BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M092R49  
 NODE=M092R49

 $\Gamma(\chi_{b0}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{28}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-3}$	90	<sup>1</sup> HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R36  
 NODE=M092R36

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$  from ESEN 13. Correlated with other results from HE 14.

NODE=M092R36;LINKAGE=A

 $\Gamma(\chi_{b0}(1P)\omega)/\Gamma_{\text{total}}$   $\Gamma_{29}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-3}$	90	<sup>1</sup> HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R37  
 NODE=M092R37

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$  from ESEN 13. Correlated with other results from HE 14.

NODE=M092R37;LINKAGE=A

 $\Gamma(\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.8 \times 10^{-3}$	90	<sup>1</sup> HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R38  
 NODE=M092R38

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$  from ESEN 13. Correlated with other results from HE 14.

NODE=M092R38;LINKAGE=A

 $\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.85 \pm 0.23 \pm 0.23$	80	<sup>1</sup> HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R39  
 NODE=M092R39

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$  from ESEN 13. Correlated with other results from HE 14.

NODE=M092R39;LINKAGE=A

$$\Gamma(\chi_{b1}(1P)\omega)/\Gamma_{\text{total}} \quad \Gamma_{32}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.57 \pm 0.22 \pm 0.21$	60	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R40  
NODE=M092R40

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$  nb from ESEN 13. Correlated with other results from HE 14.

NODE=M092R40;LINKAGE=A

$$\Gamma(\chi_{b1}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}} \quad \Gamma_{33}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.52 \pm 0.15 \pm 0.11$	24	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R41  
NODE=M092R41

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$  nb from ESEN 13. Correlated with other results from HE 14.

NODE=M092R41;LINKAGE=A

$$\Gamma(\chi_{b2}(1P)\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}} \quad \Gamma_{34}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$1.17 \pm 0.27 \pm 0.14$	29	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R42  
NODE=M092R42

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$  nb from ESEN 13. Correlated with other results from HE 14.

NODE=M092R42;LINKAGE=A

$$\Gamma(\chi_{b2}(1P)\omega)/\Gamma_{\text{total}} \quad \Gamma_{35}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.60 \pm 0.23 \pm 0.15$	13	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R43  
NODE=M092R43

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$  nb from ESEN 13. Correlated with other results from HE 14.

NODE=M092R43;LINKAGE=A

$$\Gamma(\chi_{b2}(1P)\omega)/\Gamma(\chi_{b1}(1P)\omega) \quad \Gamma_{35}/\Gamma_{32}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.38 \pm 0.16 \pm 0.09$	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R44  
NODE=M092R44

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Accounting for correlated systematics.

NODE=M092R44;LINKAGE=A

$$\Gamma(\chi_{b2}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}} \quad \Gamma_{36}/\Gamma$$

VALUE (units $10^{-3}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$0.61 \pm 0.22 \pm 0.28$	16	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R45  
NODE=M092R45

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$  nb from ESEN 13. Correlated with other results from HE 14.

NODE=M092R45;LINKAGE=A

$$\Gamma(\chi_{b2}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega})/\Gamma(\chi_{b1}(1P)(\pi^+ \pi^- \pi^0)_{\text{non-}\omega}) \quad \Gamma_{36}/\Gamma_{33}$$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.20 \pm 0.55 \pm 0.65$	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R46  
NODE=M092R46

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> Accounting for correlated systematics.

NODE=M092R46;LINKAGE=A

$$\Gamma(\eta_b(1S)\omega)/\Gamma_{\text{total}} \quad \Gamma_{38}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-3}$	90	<sup>1</sup> OSKIN	20	BELL $e^+e^- \rightarrow \omega X$

NODE=M092R53  
NODE=M092R53

<sup>1</sup> Using  $\sigma_{b\bar{b}} = 0.340 \pm 0.016$  nb from TAMPONI 15.

NODE=M092R53;LINKAGE=A

$$\Gamma(\eta_b(2S)\omega)/\Gamma_{\text{total}} \quad \Gamma_{39}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-3}$	90	<sup>1</sup> OSKIN	20	BELL $e^+e^- \rightarrow \omega X$

NODE=M092R54  
NODE=M092R54

<sup>1</sup> Using  $\sigma_{b\bar{b}} = 0.340 \pm 0.016$  nb from TAMPONI 15.

NODE=M092R54;LINKAGE=A

$$\Gamma(\gamma X_b \rightarrow \gamma \Upsilon(1S)\omega)/\Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.8 \times 10^{-5}$	90	<sup>1</sup> HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+ \pi^- \pi^0 \gamma \Upsilon(1S)$

NODE=M092R47  
NODE=M092R47

<sup>1</sup> Assuming that all the  $b\bar{b}$  events are from  $\Upsilon(5S)$  resonance decays and using  $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$  nb from ESEN 13. Correlated with other results from HE 14. For a state  $X_b$  with mass between  $10.55 \text{ GeV}/c^2$  and  $10.65 \text{ GeV}/c^2$ , the obtained 90% upper limit as a function of  $m_{X_b}$  varies from  $2.6 \times 10^{-5}$  to  $3.8 \times 10^{-5}$ .

NODE=M092R47;LINKAGE=A

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$				$\Gamma_{40}/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT		
$0.138 \pm 0.007^{+0.023}_{-0.015}$	HUANG	07	CLEO $\Upsilon(5S) \rightarrow \phi X$		NODE=M092R12 NODE=M092R12
$\Gamma(D^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$				$\Gamma_{41}/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT		
$1.076 \pm 0.040 \pm 0.068$	DRUTSKOY	07	BELL $\Upsilon(5S) \rightarrow D^0 X$		NODE=M092R10 NODE=M092R10
$\Gamma(D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$				$\Gamma_{42}/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT		
<b>0.46 ± 0.06 OUR AVERAGE</b>					NODE=M092R02 NODE=M092R02
0.472 ± 0.024 ± 0.072	<sup>1</sup> DRUTSKOY	07	BELL $\Upsilon(5S) \rightarrow D_s X$		
0.44 ± 0.09 ± 0.04	<sup>2</sup> ARTUSO	05B	CLE3 $e^+ e^- \rightarrow D_x X$		
<sup>1</sup> Using $B(D_s^+ \rightarrow \phi \pi^+) = (4.4 \pm 0.6)\%$ from PDG 06. <sup>2</sup> ARTUSO 05B reports $[\Gamma(\Upsilon(10860) \rightarrow D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(D_s^+ \rightarrow \phi \pi^+)] = 0.0198 \pm 0.0019 \pm 0.0038$ which we divide by our best value $B(D_s^+ \rightarrow \phi \pi^+) = (4.5 \pm 0.4) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.					
$\Gamma(J/\psi \text{ anything})/\Gamma_{\text{total}}$				$\Gamma_{43}/\Gamma$	
VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT		
$2.060 \pm 0.160 \pm 0.134$	DRUTSKOY	07	BELL $\Upsilon(5S) \rightarrow J/\psi X$		NODE=M092R11 NODE=M092R11
$\Gamma(B^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$				$\Gamma_{44}/\Gamma$	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.770^{+0.058}_{-0.056} \pm 0.061$	352	DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^0 X$	NODE=M092R26 NODE=M092R26
$\Gamma(B^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$				$\Gamma_{45}/\Gamma$	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.721^{+0.039}_{-0.038} \pm 0.050$	711	DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^+ X$	NODE=M092R27 NODE=M092R27

### $\Upsilon(10860)$ REFERENCES

KOVALENKO	21	PR D104 112006	E. Kovalenko <i>et al.</i>	(BELLE Collab.)	REFID=61452
DONG	20A	CP C44 083001	X.-K. Dong <i>et al.</i>		REFID=60595
OSKIN	20	PR D102 092011	P. Oskin <i>et al.</i>	(BELLE Collab.)	REFID=60735
MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60090
TAMPONI	18	EPJ C78 633	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=59195
YIN	18	PR D98 091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)	REFID=59468
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=56996
HE	14	PRL 113 142001	X.H. He <i>et al.</i>	(BELLE Collab.)	REFID=55927
ESEN	13	PR D87 031101	S. Esen <i>et al.</i>	(BELLE Collab.)	REFID=54894
SHEN	13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
LI	12	PRL 108 181808	J. Li <i>et al.</i>	(BELLE Collab.)	REFID=54116
CHEN	10	PR D82 091106	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53531
DRUTSKOY	10	PR D81 112003	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=53358
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661
LOUVOT	09	PRL 102 021801	R. Louvot <i>et al.</i>	(BELLE Collab.)	REFID=52646
CHEN	08	PRL 100 112001	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=52153
DRUTSKOY	07	PRL 98 052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51621
DRUTSKOY	07A	PR D76 012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51852
HUANG	07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
AQUINES	06	PRL 96 152001	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51106
BONVICINI	06	PRL 96 022002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=50995
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
ARTUSO	05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
BESSON	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369

NODE=M092

$\Upsilon(11020)$ 

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M093

 **$\Upsilon(11020)$  MASS**

NODE=M093M

NODE=M093M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>11000 ± 4 OUR AVERAGE</b>			
11000.0 <sup>+4.0</sup> <sub>-4.5</sub> <sup>+1.0</sup> <sub>-1.3</sub>	1 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10999.0 <sup>+7.3</sup> <sub>-7.8</sub> <sup>+16.9</sup> <sub>-1.0</sub>	2 MIZUK	16 BELL	$e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
11001 ± 1	3 DONG	20A	$e^+e^- \rightarrow b\bar{b}$
11003.0 ± 1.1 <sup>+</sup> <sub>-1.0</sub> <sup>+0.9</sup> <sub>-1.0</sub>	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow$ hadrons
10987.5 <sup>+6.4</sup> <sub>-2.5</sub> <sup>+9.1</sup> <sub>-2.3</sub>	6,7 SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10996 ± 2	8 AUBERT	09E BABR	$e^+e^- \rightarrow$ hadrons
11019 ± 5 ± 7	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
11020 ± 30	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

OCCUR=2

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M093M;LINKAGE=E

<sup>2</sup> From a simultaneous fit to the  $h_b(nP)\pi^+\pi^-$ ,  $n = 1, 2$  cross sections at 22 energy points within  $\sqrt{s} = 10.77\text{--}11.02$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single relative phase, a single relative amplitude, and two overall normalization factors, one for each  $n$ ). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

NODE=M093M;LINKAGE=D

<sup>3</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M093M;LINKAGE=G

<sup>4</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient).

NODE=M093M;LINKAGE=A

<sup>5</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions.

NODE=M093M;LINKAGE=B

<sup>6</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M093M;LINKAGE=C

<sup>7</sup> Superseded by MIZUK 19.

NODE=M093M;LINKAGE=F

<sup>8</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M093M;LINKAGE=AU

 **$\Upsilon(11020)$  WIDTH**

NODE=M093W

NODE=M093W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>24 <sup>+8</sup><sub>-6</sub> OUR AVERAGE</b>			
23.8 <sup>+8.0</sup> <sub>-6.8</sub> <sup>+0.7</sup> <sub>-1.8</sub>	1 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
27 <sup>+27</sup> <sub>-11</sub> <sup>+5</sup> <sub>-12</sub>	2 MIZUK	16 BELL	$e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
35.1 ± 1.2	3 DONG	20A	$e^+e^- \rightarrow b\bar{b}$
39.3 <sup>+1.7</sup> <sub>-1.6</sub> <sup>+1.3</sup> <sub>-2.4</sub>	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow$ hadrons
61 <sup>+9</sup> <sub>-19</sub> <sup>+2</sup> <sub>-20</sub>	6,7 SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
37 ± 3	8 AUBERT	09E BABR	$e^+e^- \rightarrow$ hadrons
61 ± 13 ± 22	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
90 ± 20	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

OCCUR=2

- <sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ . NODE=M093W;LINKAGE=E
- <sup>2</sup> From a simultaneous fit to the  $h_b(nP)\pi^+\pi^-$ ,  $n = 1, 2$  cross sections at 22 energy points within  $\sqrt{s} = 10.77\text{--}11.02$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single relative phase, a single relative amplitude, and two overall normalization factors, one for each  $n$ ). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude. NODE=M093W;LINKAGE=D
- <sup>3</sup> From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths. NODE=M093W;LINKAGE=G
- <sup>4</sup> From a fit to the total hadronic cross sections measured at 60 energy points within  $\sqrt{s} = 10.82\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with  $1/\sqrt{s}$  dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , one relative phase, and one decoherence coefficient). NODE=M093W;LINKAGE=A
- <sup>5</sup> Not including uncertain and potentially large systematic errors due to assumed continuum amplitude  $1/\sqrt{s}$  dependence and related interference contributions. NODE=M093W;LINKAGE=B
- <sup>6</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n=1, 2, 3$ , cross sections at 25 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of  $\Upsilon(10860)$  and  $\Upsilon(11020)$ , a single universal relative phase, and three decoherence coefficients, one for each  $n$ ). Continuum contributions were measured (and therefore fixed) to be zero. NODE=M093W;LINKAGE=C
- <sup>7</sup> Superseded by MIZUK 19. NODE=M093W;LINKAGE=F
- <sup>8</sup> In a model where a flat non-resonant  $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated. NODE=M093W;LINKAGE=AU

### $\Upsilon(11020)$ DECAY MODES

NODE=M093215;NODE=M093

Mode	Fraction ( $\Gamma_i/\Gamma$ )	
$\Gamma_1$ $e^+e^-$	$(5.4^{+1.9}_{-2.1}) \times 10^{-6}$	DESIG=1
$\Gamma_2$ $\Upsilon(1S)\pi^+\pi^-$		DESIG=5
$\Gamma_3$ $\Upsilon(2S)\pi^+\pi^-$		DESIG=6
$\Gamma_4$ $\Upsilon(3S)\pi^+\pi^-$		DESIG=7
$\Gamma_5$ $\chi_{bJ}(1P)\pi^+\pi^-\pi^0$	$(9^{+9}_{-8}) \times 10^{-3}$	DESIG=2
$\Gamma_6$ $\chi_{b1}(1P)\pi^+\pi^-\pi^0$	seen	DESIG=3
$\Gamma_7$ $\chi_{b2}(1P)\pi^+\pi^-\pi^0$	seen	DESIG=4

### $\Upsilon(11020)$ PARTIAL WIDTHS

NODE=M093220

$\Gamma(e^+e^-)$					$\Gamma_1$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
<b>0.130 ± 0.030 OUR AVERAGE</b>					NODE=M093W1 NODE=M093W1
0.095 ± 0.03 ± 0.035	BESSION	85	CLEO $e^+e^- \rightarrow$ hadrons		
0.156 ± 0.040	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons		

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_2/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.46 ± 0.08 <sup>1,2</sup> MIZUK 19 BELL  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ . NODE=M093R04;LINKAGE=A

<sup>2</sup> Reported as the range 0.38–0.54 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions. NODE=M093R04;LINKAGE=B

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_3/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.65 ± 0.52 <sup>1,2</sup> MIZUK 19 BELL  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ . NODE=M093R05;LINKAGE=A

<sup>2</sup> Reported as the range 0.13–1.16 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions. NODE=M093R05;LINKAGE=B



$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_1\Gamma_4/\Gamma$   
 VALUE (eV) DOCUMENT ID TECN COMMENT

NODE=M093R06  
 NODE=M093R06

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.33±0.16 <sup>1,2</sup> MIZUK 19 BELL  $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

<sup>1</sup> From a simultaneous fit to the  $\Upsilon(nS)\pi^+\pi^-$ ,  $n = 1, 2, 3$ , cross sections at 28 energy points within  $\sqrt{s} = 10.6\text{--}11.05$  GeV, including the initial-state radiation at  $\Upsilon(10860)$ .

NODE=M093R06;LINKAGE=A

<sup>2</sup> Reported as the range 0.17–0.49 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M093R06;LINKAGE=B

$\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$   
 VALUE (units  $10^{-3}$ ) DOCUMENT ID TECN COMMENT

NODE=M093R00  
 NODE=M093R00

$8.7 \pm 4.3^{+7.6}_{-6.6}$  YIN 18 BELL  $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_6/\Gamma$   
 VALUE DOCUMENT ID TECN COMMENT

NODE=M093R01  
 NODE=M093R01

seen YIN 18 BELL  $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$   $\Gamma_7/\Gamma$   
 VALUE DOCUMENT ID TECN COMMENT

NODE=M093R02  
 NODE=M093R02

seen YIN 18 BELL  $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)$   $\Gamma_7/\Gamma_6$   
 VALUE DOCUMENT ID TECN COMMENT

NODE=M093R03  
 NODE=M093R03

$0.4 \pm 0.2$  YIN 18 BELL  $e^+e^- \rightarrow \text{hadrons}$

## $\Upsilon(11020)$ REFERENCES

NODE=M093

DONG	20A	CP C44 083001	X.-K. Dong <i>et al.</i>		
MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60595
YIN	18	PR D98 091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)	REFID=60090
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=59468
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57465
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=57121
BESSION	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=52661
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSP Collab.)	REFID=22368
					REFID=22369

## OTHER MESONS

NODE=MXXX050

$X_0(2900)$

$I(J^P) = ?(0^+)$

NODE=M250

OMITTED FROM SUMMARY TABLE

An exotic state with minimal quark content  $\bar{c}d\bar{s}u$ . Observed by AAIJ 20AI using full amplitude analysis of  $B^+ \rightarrow D^+ D^- K^+$  decays.

NODE=M250

## $X_0(2900)$ MASS

NODE=M250M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  
 $2866 \pm 7 \pm 2$  1.2k <sup>1</sup> AAIJ 20AI LHCB  $B^+ \rightarrow D^+ D^- K^+$

NODE=M250M

<sup>1</sup> Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M250M;LINKAGE=A

## $X_0(2900)$ WIDTH

NODE=M250W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT  
 $57 \pm 12 \pm 4$  1.2k <sup>1</sup> AAIJ 20AI LHCB  $B^+ \rightarrow D^+ D^- K^+$

NODE=M250W

<sup>1</sup> Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M250W;LINKAGE=A

**$X_0(2900)$  DECAY MODES**

NODE=M250215;NODE=M250

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^- K^+$	seen

DESIG=1

 **$X_0(2900)$  BRANCHING RATIOS**

NODE=M250225

$\Gamma(D^- K^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M250R01  
NODE=M250R01 **$X_0(2900)$  REFERENCES**

NODE=M250

AAIJ	20AF PRL 125 242001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20AI PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=60702  
REFID=60739 **$X_1(2900)$** 

$$I(J^P) = ?(1^-)$$

NODE=M251

OMITTED FROM SUMMARY TABLE

An exotic state with minimal quark content  $\bar{c}d\bar{s}u$ . Observed by AAIJ 20AI using full amplitude analysis of  $B^+ \rightarrow D^+ D^- K^+$  decays.

NODE=M251

 **$X_1(2900)$  MASS**

NODE=M251M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>2904 \pm 5 \pm 1</math></b>	1.2k	<sup>1</sup> AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M251M

<sup>1</sup>Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M251M;LINKAGE=B

 **$X_1(2900)$  WIDTH**

NODE=M251W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>110 \pm 11 \pm 4</math></b>	1.2k	<sup>1</sup> AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M251W

<sup>1</sup>Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M251W;LINKAGE=A

 **$X_1(2900)$  DECAY MODES**

NODE=M251215;NODE=M251

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^- K^+$	seen

DESIG=1

 **$X_1(2900)$  BRANCHING RATIOS**

NODE=M251225

$\Gamma(D^- K^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M251R01  
NODE=M251R01 **$X_1(2900)$  REFERENCES**

NODE=M251

AAIJ	20AF PRL 125 242001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20AI PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=60702  
REFID=60739

$T_{cc}(3875)^+$ 

$I(J^P) = ?(??)$

NODE=M265

## OMITTED FROM SUMMARY TABLE

Observed with large significance by AAIJ 22E in the doubly-charmed (C = 2) decay mode  $D^0 D^0 \pi^+$  using inclusive  $pp$  collisions at 7, 8, and 13 TeV.

NODE=M265

 $T_{cc}(3875)^+$  MASS

NODE=M265M

OUR FIT value comes from the measurement of  $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$  below and  $m_{D^{*+}} + m_{D^0}$  values.

NODE=M265M

VALUE (MeV) DOCUMENT ID  
**3874.83 ± 0.11 OUR FIT**

NODE=M265M

 $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$ 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>= 0.27 ± 0.06 OUR FIT</b>				
<b>= 0.273 ± 0.061<sup>+0.012</sup><sub>-0.015</sub></b>	117	<sup>1</sup> AAIJ	22E LHCB	$pp \rightarrow D^0 D^0 \pi^+ X$

NODE=M265DM  
 NODE=M265DM

<sup>1</sup> The fit assumes a relativistic  $P$ -wave Breit Wigner function modified by Blatt-Weisskopf form factor with radius 3.5 GeV<sup>-1</sup>. The significance for  $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0}) < 0$  is 4.3  $\sigma$ .

NODE=M265DM;LINKAGE=A

 $T_{cc}(3875)^+$  WIDTH

NODE=M265W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>0.410 ± 0.165<sup>+0.047</sup><sub>-0.057</sub></b>	117	<sup>1</sup> AAIJ	22E LHCB	$pp \rightarrow D^0 D^0 \pi^+ X$

NODE=M265W

<sup>1</sup> The fit assumes a relativistic  $P$ -wave Breit Wigner function modified by Blatt-Weisskopf form factor with radius 3.5 GeV<sup>-1</sup>.

NODE=M265W;LINKAGE=A

 $T_{cc}(3875)^+$  DECAY MODES

NODE=M265215;NODE=M265

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $D^0 D^0 \pi^+$	seen

DESIG=1

 $T_{cc}(3875)^+$  BRANCHING RATIOS

NODE=M265225

$\Gamma(D^0 D^0 \pi^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	117	AAIJ	22E LHCB	$pp \rightarrow D^0 D^0 \pi^+ X$

NODE=M265R01  
 NODE=M265R01

 $T_{cc}(3875)^+$  REFERENCES

NODE=M265

AAIJ 22E NATP 18 751 R. Aaij *et al.* (LHCb Collab.)

REFID=61658

**Z<sub>c</sub>(3900)**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

NODE=M210

was X(3900)

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

NODE=M210

Charged  $Z_c(3900)$  seen as a peak in the invariant mass distribution of the  $J/\psi\pi^\pm$  system by BES III (ABLIKIM 13T) in  $e^+e^- \rightarrow \pi^+\pi^-J/\psi$  at c.m. energy of 4.26 GeV and by radiative return from  $e^+e^-$  collisions at  $\sqrt{s}$  from 9.46 to 10.86 GeV at Belle (LIU 13B). Partial wave analysis of ABLIKIM 17J determines  $J^P = 1^+$  with more than  $7\sigma$  significance. Neutral  $Z_c(3900)$  seen in the  $J/\psi\pi^0$  invariant mass distribution in  $e^+e^- \rightarrow \pi^0\pi^0J/\psi$  at c.m. energies of 4.23, 4.26, and 4.36 GeV by BES III (ABLIKIM 15U) and at 4.17 GeV by XIAO 13A. Peaks in  $(D\bar{D}^*)^{0,\pm}$  reported by BES III (ABLIKIM 14A, ABLIKIM 15AB) are assumed to be related.

**Z<sub>c</sub>(3900) MASS**

NODE=M210M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>3887.1±2.6 OUR AVERAGE</b>		Error includes scale factor of 1.7. See the ideogram below.			
3893.1±2.2± 3.0		<sup>1</sup> ABLIKIM	20N	BES3	0 $e^+e^- \rightarrow \pi^0\pi^0J/\psi$
3902.6 <sup>+5.2+ 3.3</sup> <sub>-5.0- 1.4</sub>		<sup>2,3</sup> ABAZOV	19	D0	± 1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3881.2±4.2±52.7	6k	<sup>4</sup> ABLIKIM	17J	BES3	± $e^+e^- \rightarrow \pi^+\pi^-J/\psi$
3885.7 <sup>+4.3</sup> <sub>-5.7</sub> ± 8.4		<sup>2,4</sup> ABLIKIM	15AB	BES3	0 $e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
3881.7±1.6± 1.6	1.2k	<sup>2,4</sup> ABLIKIM	15AC	BES3	± $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3883.9±1.5± 4.2	1.2k	<sup>2,4</sup> ABLIKIM	14A	BES3	± $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3894.5±6.6± 4.5	159	<sup>2</sup> LIU	13B	BELL	± $e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$
3886 ± 4 ± 2	81	<sup>2,5</sup> XIAO	13A		± 4.17 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$
3904 ± 9 ± 5	25	<sup>2,5</sup> XIAO	13A	0	4.17 $e^+e^- \rightarrow \pi^0\pi^0J/\psi$

NODE=M210M

• • • We do not use the following data for averages, fits, limits, etc. • • •

3895.0±5.2 <sup>+ 4.0</sup> <sub>- 2.7</sub>	502	<sup>2,6</sup> ABAZOV	18B	D0	± 1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3894.8±2.3± 3.2	356	<sup>2,7</sup> ABLIKIM	15U	BES3	0 $e^+e^- \rightarrow \pi^0\pi^0J/\psi$
3899.0±3.6± 4.9	307	<sup>2,8</sup> ABLIKIM	13T	BES3	± $e^+e^- \rightarrow \pi^+\pi^-J/\psi$

OCCUR=2

<sup>1</sup> Pole mass obtained from a fit to a relativistic Breit-Wigner.

<sup>2</sup> Neglecting interference between the  $Z_c(3900)$  and other processes.

<sup>3</sup> Measured in weak decays of  $b$ -flavored hadrons (nonprompt).

<sup>4</sup> Pole mass obtained from a fit to a Flatte-like formula.

<sup>5</sup> For  $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$ . Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

<sup>6</sup> The signal of the  $Z_c(3900)$  is correlated with a parent  $J/\psi\pi^+\pi^-$  system in the invariant mass range 4.2–4.7 GeV. Superseded by ABAZOV 19.

<sup>7</sup> Superseded by ABLIKIM 20N.

<sup>8</sup> Superseded by ABLIKIM 17J.

NODE=M210M;LINKAGE=L

NODE=M210M;LINKAGE=A

NODE=M210M;LINKAGE=G


NODE=M210M;LINKAGE=E

NODE=M210M;LINKAGE=B

NODE=M210M;LINKAGE=F

NODE=M210M;LINKAGE=J

NODE=M210M;LINKAGE=K



### Z<sub>c</sub>(3900) WIDTH

NODE=M210W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
<b>28.4 ± 2.6 OUR AVERAGE</b>						
44.4 ± 5.2 ± 14.0		<sup>1</sup> ABLIKIM	20N	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
32 <sup>+28</sup> <sub>-21</sub> <sup>+26</sup> <sub>-7</sub>		<sup>2,3</sup> ABAZOV	19	D0	±	1.96 TeV $p\bar{p} \rightarrow \pi^+\pi^- J/\psi X$ (non-prompt)
51.8 ± 4.6 ± 36.0	6 k	<sup>4</sup> ABLIKIM	17J	BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
35 <sup>+11</sup> <sub>-12</sub> ± 15		<sup>2,4</sup> ABLIKIM	15AB	BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
26.6 ± 2.0 ± 2.1	1248	<sup>2,4</sup> ABLIKIM	15AC	BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
24.8 ± 3.3 ± 11.0	1212	<sup>2,4</sup> ABLIKIM	14A	BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
63 ± 24 ± 26	159	<sup>2</sup> LIU	13B	BELL	±	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
37 ± 4 ± 8	81	<sup>2,5</sup> XIAO	13A		±	4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
29.6 ± 8.2 ± 8.2	356	<sup>2,6</sup> ABLIKIM	15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
46 ± 10 ± 20	307	<sup>2,7</sup> ABLIKIM	13T	BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

NODE=M210W

- <sup>1</sup> Pole width obtained from a fit to a relativistic Breit-Wigner.
- <sup>2</sup> Neglecting interference between the Z<sub>c</sub>(3900) and other processes.
- <sup>3</sup> Measured in weak decays of b-flavored hadrons (nonprompt).
- <sup>4</sup> Pole width obtained from a fit to a Flatte-like formula.
- <sup>5</sup> For  $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$ . Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
- <sup>6</sup> Superseded by ABLIKIM 20N.
- <sup>7</sup> Superseded by ABLIKIM 17J.

NODE=M210W;LINKAGE=G  
 NODE=M210W;LINKAGE=A  
 NODE=M210W;LINKAGE=F  
 NODE=M210W;LINKAGE=E  
 NODE=M210W;LINKAGE=B

NODE=M210W;LINKAGE=H  
 NODE=M210W;LINKAGE=I

### Z<sub>c</sub>(3900) DECAY MODES

NODE=M210215;NODE=M210

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi\pi$	seen
$\Gamma_2$ $h_c\pi^\pm$	not seen
$\Gamma_3$ $\eta_c\pi^+\pi^-$	not seen
$\Gamma_4$ $\eta_c(1S)\rho(770)^\pm$	
$\Gamma_5$ $(D\bar{D}^*)^\pm$	seen
$\Gamma_6$ $D^0D^{*-} + \text{c.c.}$	seen
$\Gamma_7$ $D^-D^{*0} + \text{c.c.}$	seen
$\Gamma_8$ $\omega\pi^\pm$	not seen
$\Gamma_9$ $J/\psi\eta$	not seen
$\Gamma_{10}$ $D^+D^{*-} + \text{c.c.}$	seen
$\Gamma_{11}$ $D^0\bar{D}^{*0} + \text{c.c.}$	seen

DESIG=1  
 DESIG=2  
 DESIG=10  
 DESIG=11  
 DESIG=3;OUR EVAL;→ UNCHECKED ←  
 DESIG=8  
 DESIG=9  
 DESIG=4  
 DESIG=5  
 DESIG=6  
 DESIG=7

**Z<sub>c</sub>(3900) BRANCHING RATIOS**

NODE=M210225

$\Gamma(J/\psi\pi)/\Gamma_{\text{total}}$						$\Gamma_1/\Gamma$
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	20N	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
seen		<sup>1</sup> ABAZOV	19	D0	$\pm$	1.96 TeV $p\bar{p} \rightarrow \pi^+\pi^- J/\psi X$ (prompt)
seen		ABLIKIM	17J	BES3	$\pm$	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
seen	356	ABLIKIM	15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
not seen		<sup>2</sup> ADOLPH	15D	COMP	$\pm$	$\gamma N \rightarrow J/\psi\pi^\pm N$
seen	307	ABLIKIM	13T	BES3	$\pm$	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
seen	25	<sup>3</sup> XIAO	13A		0	4.17 $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$

NODE=M210R01  
NODE=M210R01<sup>1</sup> But not seen in the "prompt" sample (no b-hadron enhancement).<sup>2</sup> ADOLPH 15D measure  $B(Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm) \sigma(\gamma N \rightarrow Z_c(3900)^\pm N) / \sigma(\gamma N \rightarrow J/\psi N) < 3.7 \times 10^{-3}$  at 90% CL.<sup>3</sup> Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.NODE=M210R01;LINKAGE=C  
NODE=M210R01;LINKAGE=A

NODE=M210R01;LINKAGE=XI

$\Gamma(h_c\pi^\pm)/\Gamma_{\text{total}}$						$\Gamma_2/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	13X	BES3	$\pm$	$e^+e^- \rightarrow h_c\pi^+\pi^-$

NODE=M210R02  
NODE=M210R02

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$						$\Gamma_3/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		<sup>1</sup> VINOKUROVA 15	BELL	0	$B^+ \rightarrow K^+\eta_c\pi^+\pi^-$	

NODE=M210R11  
NODE=M210R11<sup>1</sup> VINOKUROVA 15 reports  $B(B^+ \rightarrow K^+ Z_c(3900)^0) \times B(X \rightarrow \eta_c\pi^+\pi^-) < 4.7 \times 10^{-5}$  at 90% CL.

NODE=M210R11;LINKAGE=VI

$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi\pi)$						$\Gamma_5/\Gamma_1$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
$6.2 \pm 1.1 \pm 2.7$		<sup>1</sup> ABLIKIM	14A	BES3	$\pm$	$e^+e^- \rightarrow \pi^\pm (D\bar{D}^*)^\mp$

NODE=M210R03  
NODE=M210R03<sup>1</sup> Assuming the same origin of the  $(D\bar{D}^*)^\pm$  and  $\pi^\pm J/\psi$  decay modes.

NODE=M210R03;LINKAGE=A

$\Gamma(D^0 D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$						$\Gamma_6/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	15AC	BES3	$\pm$	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$
seen		ABLIKIM	14A	BES3	$\pm$	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$

NODE=M210R09  
NODE=M210R09

$\Gamma(D^- D^{*0} + \text{c.c.})/\Gamma_{\text{total}}$						$\Gamma_7/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	15AC	BES3	$\pm$	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$
seen		ABLIKIM	14A	BES3	$\pm$	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$

NODE=M210R10  
NODE=M210R10

$\Gamma(\omega\pi^\pm)/\Gamma_{\text{total}}$						$\Gamma_8/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	15R	BES3	$\pm$	$e^+e^- \rightarrow \omega\pi^+\pi^-$

NODE=M210R00  
NODE=M210R00

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$						$\Gamma_9/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	15Q	BES3	0	4.0–4.6 $e^+e^- \rightarrow J/\psi\eta\pi^0$

NODE=M210R04  
NODE=M210R04

$\Gamma(J/\psi\eta)/\Gamma(J/\psi\pi)$						$\Gamma_9/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.15	90	ABLIKIM	15Q	BES3	0	4.226 $e^+e^- \rightarrow J/\psi\eta\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.65	90	ABLIKIM	15Q	BES3	0	4.257 $e^+e^- \rightarrow J/\psi\eta\pi^0$

NODE=M210R05  
NODE=M210R05

OCCUR=2

$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(J/\psi\pi)$						$\Gamma_4/\Gamma_1$
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
$2.3 \pm 0.8$	332	<sup>1</sup> ABLIKIM	19BC	BES3	$\pm$	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$

NODE=M210R12  
NODE=M210R12<sup>1</sup> Using  $e^+e^- \rightarrow \pi^\mp (Z_c(3900)^\pm \rightarrow J/\psi\pi^\pm)$  cross section at 4.23 and 4.26 GeV from ABLIKIM 17J.

NODE=M210R12;LINKAGE=A

$\Gamma(D^+ D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$						$\Gamma_{10}/\Gamma$
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	15AB	BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

NODE=M210R06  
NODE=M210R06

$\Gamma(D^0\bar{D}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$  $\Gamma_{11}/\Gamma$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

NODE=M210R07  
NODE=M210R07 $\Gamma(D^+D^{*-} + \text{c.c.})/\Gamma(D^0\bar{D}^{*0} + \text{c.c.})$  $\Gamma_{10}/\Gamma_{11}$ 

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<b>0.96±0.18±0.12</b>	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

NODE=M210R08  
NODE=M210R08**Z<sub>c</sub>(3900) REFERENCES**

NODE=M210

ABLIKIM	20N	PR D102 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=60338
ABAZOV	19	PR D100 012005	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=59903
ABLIKIM	19BC	PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60036
ABAZOV	18B	PR D98 052010	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=59442
ABLIKIM	17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=57950
ABLIKIM	15AB	PRL 115 222002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56954
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=56967
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56782
ABLIKIM	15R	PR D92 032009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56783
ABLIKIM	15U	PRL 115 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56786
ADOLPH	15D	PL B742 330	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56791
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (errat.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)	REFID=57795
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=55648
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55409
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55635
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)	REFID=55593

NODE=M259

**Z<sub>cs</sub>(4000)**

$$I(J^P) = \frac{1}{2}(1^+)$$

## OMITTED FROM SUMMARY TABLE

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M259

Seen by AAIJ 21E in  $B^+ \rightarrow Z_{cs}(4000)^+ \phi$  with  $Z_{cs}(4000)^+ \rightarrow J/\psi K^+$  using an amplitude analysis of  $B^+ \rightarrow J/\psi \phi K^+$  with a significance (accounting for systematic uncertainties) of  $15\sigma$ . The  $J^P = 1^+$  assignment is favored with high significance. ABLIKIM 21G also reports a  $J^P = 1^+$   $Z_{cs}$  state in this mass region using  $e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$  with a significance of  $5.3\sigma$ . The incompatible values for the widths reported by AAIJ 21E and ABLIKIM 21G could either indicate the existence of two separate states or possibly be explained in a coupled channel model (see ORTEGA 21).

**Z<sub>cs</sub>(4000) MASS**

NODE=M259M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**3980-4010 OUR EVALUATION**NODE=M259M  
→ UNCHECKED ←

**3988 ±5 OUR AVERAGE** Error includes scale factor of 2.7.  $[3983.2^{+2.6}_{-3.3}$  MeV OUR 2022 AVERAGE]

NEW

$3992.2 \pm 1.7 \pm 1.6$	1	ABLIKIM	22AE BES3	$e^+e^- \rightarrow K_S^0(D_s^- D^{*+} + D_s^{*-} D^+)$
$4003 \pm 6 \begin{smallmatrix} +4 \\ -14 \end{smallmatrix}$	24k	2	AAIJ	21E LHCB $B^+ \rightarrow J/\psi \phi K^+$
$3982.5 \begin{smallmatrix} +1.8 \\ -2.6 \end{smallmatrix} \pm 2.1$	3	ABLIKIM	21G BES3	$e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$

<sup>1</sup> Pole mass for a mass-, width-dependent Breit-Wigner fit to the mass spectrum recoiling against  $K_S^0$  at center of mass energies between 4.628 and 4.699 GeV, with a significance of  $4.6\sigma$ .

NODE=M259M;LINKAGE=D

<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $15\sigma$ .

NODE=M259M;LINKAGE=A

<sup>3</sup> Pole mass for a mass-dependent Breit-Wigner fit to the mass spectrum recoiling against  $K^+$  at center of mass energies between 4.628 and 4.698 GeV, with a significance of  $5.3\sigma$ .

NODE=M259M;LINKAGE=B

**Z<sub>cs</sub>(4000) WIDTH**

NODE=M259W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

**5-150 OUR EVALUATION****13 ± 4 OUR AVERAGE**[17<sup>+6</sup><sub>-5</sub> MeV OUR 2022 AVERAGE]

7.7 <sup>+</sup> <sub>-</sub> 4.1 <sup>±</sup> 4.3		<sup>1</sup> ABLIKIM	22AE BES3	$e^+e^- \rightarrow K_S^0(D_s^- D^{*+} + D_s^{*-} D^+)$
131 ±15 ±26	24k	<sup>2</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
12.8 <sup>+</sup> <sub>-</sub> 5.3 <sup>±</sup> 3.0		<sup>3</sup> ABLIKIM	21G BES3	$e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$

<sup>1</sup> Pole width for a mass-, width-dependent Breit-Wigner fit to the mass spectrum recoiling against  $K_S^0$  at center of mass energies between 4.628 and 4.699 GeV, with a significance of 4.6  $\sigma$ .

<sup>2</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 15  $\sigma$ .

<sup>3</sup> Pole width for a mass-dependent Breit-Wigner fit to the mass spectrum recoiling against  $K^+$  at center of mass energies between 4.628 and 4.698 GeV, with a significance of 5.3  $\sigma$ .

NODE=M259W  
→ UNCHECKED ←  
NEW

NODE=M259W;LINKAGE=C

NODE=M259W;LINKAGE=A  
NODE=M259W;LINKAGE=B

**Z<sub>cs</sub>(4000) DECAY MODES**

NODE=M259215;NODE=M259

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi K^+$	seen
$\Gamma_2$ $D_s^+ \bar{D}^{*0}$ or $D_s^{*+} \bar{D}^0$	seen

DESIG=1

DESIG=2

$\Gamma(J/\psi K^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
--	-------------------

NODE=M259R01  
NODE=M259R01

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
------	-----	-------------------	----------	-----------------------------------

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of 15  $\sigma$ .

NODE=M259R01;LINKAGE=A

$\Gamma(D_s^+ \bar{D}^{*0}$ or $D_s^{*+} \bar{D}^0)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma$
--	-------------------

NODE=M259R00  
NODE=M259R00

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	<sup>1</sup> ABLIKIM	22AE BES3	$e^+e^- \rightarrow K_S^0(D_s^- D^{*+} + D_s^{*-} D^+)$
------	----------------------	-----------	---

seen	<sup>2</sup> ABLIKIM	21G BES3	$e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$
------	----------------------	----------	---

<sup>1</sup> Seen in the mass spectrum recoiling against  $K_S^0$  at center of mass energies between 4.628 and 4.699 GeV, with a significance of 4.6  $\sigma$ .

NODE=M259R00;LINKAGE=B

<sup>2</sup> Seen in the spectrum recoiling against  $K^+$  in  $e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$  collisions at center of mass energies between 4.628 and 4.698 GeV, with a significance of 5.3  $\sigma$ .

NODE=M259R00;LINKAGE=A

**Z<sub>cs</sub>(4000) REFERENCES**

NODE=M259

ABLIKIM	22AE PRL 129 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	21E PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
ABLIKIM	21G PRL 126 102001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ORTEGA	21 PL B818 136382	P.G. Ortega, D.R. Entem, F. Fernandez	

REFID=61877

REFID=61150

REFID=61065

REFID=61108



**$X(4020)^\pm$** 

$$I^G(J^{PC}) = 1^+(\text{?}^-)$$

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

Charged  $X(4020)$  seen by ABLIKIM 13X from  $e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$  at c.m. energy from 3.90 to 4.42 GeV as a peak in the invariant mass distribution of the  $\pi^\pm h_c(1P)$  system, and by ABLIKIM 14B from  $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$  events in  $(D^*\bar{D}^*)^\pm$  mass. A neutral  $X(4020)$  seen by ABLIKIM 14P at three c.m. energies in the same range in  $e^+e^- \rightarrow \pi^0\pi^0h_c(1P)$  as a peak in the larger of the two masses recoiling against a  $\pi^0$ . ABLIKIM 15AA observes a  $5.9\sigma$  signal in  $(D^*\bar{D}^*)^0$  in  $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$  events using collisions at two c.m. energies. Production rates and mass values support grouping neutral and charged  $X(4020)$  together as manifestations of a single  $I = 1$  particle.

NODE=M213

NODE=M213

 **$X(4020)^\pm$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>4024.1 ± 1.9 OUR AVERAGE</b>					
$4025.5^{+2.0}_{-4.7} \pm 3.1$	116	<sup>1</sup> ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
$4026.3 \pm 2.6 \pm 3.7$	401	<sup>1</sup> ABLIKIM 14B	BES3	$\pm$	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
$4023.9 \pm 2.2 \pm 3.8$	61	<sup>1,2</sup> ABLIKIM 14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0h_c$
$4022.9 \pm 0.8 \pm 2.7$	253	<sup>1</sup> ABLIKIM 13X	BES3	$\pm$	$e^+e^- \rightarrow \pi^+\pi^-h_c$

NODE=M213M

NODE=M213M

<sup>1</sup> Neglecting interference between the  $X(4020)$  and non-resonant continuum.

<sup>2</sup> Assuming  $J^P = 1^+$  and width of  $7.9 \pm 2.6$  MeV.

NODE=M213M;LINKAGE=AB

NODE=M213M;LINKAGE=B

 **$X(4020)^\pm$  WIDTH**


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>13 ± 5 OUR AVERAGE</b> Error includes scale factor of 1.7. See the ideogram below.					
$23.0 \pm 6.0 \pm 1.0$	116	<sup>1</sup> ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
$24.8 \pm 5.6 \pm 7.7$	401	<sup>1</sup> ABLIKIM 14B	BES3	$\pm$	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
$7.9 \pm 2.7 \pm 2.6$	253	<sup>1</sup> ABLIKIM 13X	BES3	$\pm$	$e^+e^- \rightarrow \pi^+\pi^-h_c$

NODE=M213W

NODE=M213W

<sup>1</sup> Neglecting interference between the  $X(4020)$  and non-resonant continuum.

NODE=M213W;LINKAGE=AB



**$X(4020)^\pm$  DECAY MODES**

NODE=M213215;NODE=M213

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $h_c(1P)\pi$	seen
$\Gamma_2$ $D^*\bar{D}^*$	seen
$\Gamma_3$ $D\bar{D}^* + \text{c.c.}$	not seen
$\Gamma_4$ $\eta_c\pi^+\pi^-$	not seen
$\Gamma_5$ $\eta_c(1S)\rho(770)^\pm$	
$\Gamma_6$ $J/\psi(1S)\pi^\pm$	not seen

DESIG=1

DESIG=2

DESIG=4

DESIG=3

DESIG=6

DESIG=5

 **$X(4020)^\pm$  BRANCHING RATIOS**

NODE=M213225

$\Gamma(h_c(1P)\pi)/\Gamma_{\text{total}}$						$\Gamma_1/\Gamma$	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
seen	61	ABLIKIM	14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 h_c$	NODE=M213R01
seen	253	ABLIKIM	13X	BES3	$\pm$	$e^+e^- \rightarrow \pi^+\pi^- h_c$	NODE=M213R01

$\Gamma(D^*\bar{D}^*)/\Gamma_{\text{total}}$						$\Gamma_2/\Gamma$	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
seen	116	<sup>1</sup> ABLIKIM	15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$	NODE=M213R02
seen	401	<sup>1</sup> ABLIKIM	14B	BES3	$\pm$	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$	NODE=M213R02

NODE=M213R02  
NODE=M213R02<sup>1</sup> Neglecting interference between the  $X(4020)$  and non-resonant continuum.

NODE=M213R02;LINKAGE=A

$\Gamma(D\bar{D}^* + \text{c.c.})/\Gamma_{\text{total}}$						$\Gamma_3/\Gamma$
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
not seen	ABLIKIM	15AC	BES3	$\pm$	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$	NODE=M213R03

NODE=M213R03  
NODE=M213R03

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$						$\Gamma_4/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT			
not seen	<sup>1</sup> VINOKUROVA 15	BELL	$B^+ \rightarrow K^+\eta_c\pi^+\pi^-$			

NODE=M213R00  
NODE=M213R00<sup>1</sup> VINOKUROVA 15 reports  $B(B^+ \rightarrow K^+X(4020)^0) \times B(X \rightarrow \eta_c\pi^+\pi^-) < 1.6 \times 10^{-5}$  at 90% CL.

NODE=M213R00;LINKAGE=VI

$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(h_c(1P)\pi)$						$\Gamma_5/\Gamma_1$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.2	90	<sup>1</sup> ABLIKIM	19BC	BES3	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$	NODE=M213R05

NODE=M213R05  
NODE=M213R05<sup>1</sup> Using  $e^+e^- \rightarrow \pi^\mp(Z_c(4020)^\pm \rightarrow h_c(1P)\pi^\pm)$  cross section at 4.23, 4.26 and 4.36 GeV from ABLIKIM 13X.

NODE=M213R05;LINKAGE=A

$\Gamma(J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$						$\Gamma_6/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT			
not seen	<sup>1</sup> ABLIKIM	17J	BES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$		

NODE=M213R04  
NODE=M213R04<sup>1</sup> From Partial Wave Analysis assuming  $J^P = 1^+$ .

NODE=M213R04;LINKAGE=A

 **$X(4020)^\pm$  REFERENCES**

NODE=M213

ABLIKIM	19BC	PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60036
ABLIKIM	17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57950
ABLIKIM	15AA	PRL 115 182002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56951
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56967
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (errata.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)	REFID=57795
ABLIKIM	14B	PRL 112 132001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55654
ABLIKIM	14P	PRL 113 212002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56118
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55635

**X(4050)<sup>±</sup>**
 $I^G(J^{PC}) = 1^-(?^{?+})$   
 I, G, C need confirmation.

## OMITTED FROM SUMMARY TABLE

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

Observed by MIZUK 08 in the  $\pi^+\chi_{c1}(1P)$  invariant mass distribution in  $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$  decays. Not seen by LEES 12B in this same mode after accounting for  $K\pi$  resonant mass and angular structure.

NODE=M191

NODE=M191

**X(4050)<sup>±</sup> MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4051 \pm 14^{+20}_{-41}$	<sup>1</sup> MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

<sup>1</sup> From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M191M

NODE=M191M

NODE=M191M;LINKAGE=MI

**X(4050)<sup>±</sup> WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$82^{+21+47}_{-17-22}$	<sup>1</sup> MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

<sup>1</sup> From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M191W

NODE=M191W

NODE=M191W;LINKAGE=MI

**X(4050)<sup>±</sup> DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi^+\chi_{c1}(1P)$	seen
$\Gamma_2$ $\pi^\pm\psi(3770)$	not seen
$\Gamma_3$ $\pi^\pm\chi_{c0}(1P)$	not seen
$\Gamma_4$ $\pi^\pm\chi_{c2}(1P)$	not seen

NODE=M191215;NODE=M191

DESIG=1

DESIG=2

DESIG=3

DESIG=4

**X(4050)<sup>±</sup> BRANCHING RATIOS**

$\Gamma(\pi^+\chi_{c1}(1P))/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
seen			<sup>1</sup> MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$	

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	16	<sup>2</sup> ABLIKIM	21W	BES3	$e^+e^- \rightarrow \chi_{cJ}\pi^+\pi^+$
not seen		<sup>3</sup> LEES	12B	BABR	$B \rightarrow K\pi\chi_{c1}(1P)$

<sup>1</sup> With a product branching fraction measurement of  $B(\bar{B}^0 \rightarrow K^-\chi(4050)^+) \times B(\chi(4050)^+ \rightarrow \pi^+\chi_{c1}(1P)) = (3.0^{+1.5+3.7}_{-0.8-1.6}) \times 10^{-5}$ .

<sup>2</sup> ABLIKIM 21W measurement is limited by statistics.

<sup>3</sup> With a product branching fraction limit of  $B(\bar{B}^0 \rightarrow \chi(4050)^+K^-) \times B(\chi(4050)^+ \rightarrow \chi_{c1}\pi^+) < 1.8 \times 10^{-5}$  at 90% CL.

NODE=M191225

NODE=M191R01

NODE=M191R01

NODE=M191R01;LINKAGE=MI

NODE=M191R01;LINKAGE=A

NODE=M191R01;LINKAGE=LE

$\Gamma(\pi^\pm\chi_{c0}(1P))/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_3/\Gamma$
not seen		18	<sup>1</sup> ABLIKIM	21W	BES3 $e^+e^- \rightarrow \chi_{cJ}\pi^+\pi^+$	

<sup>1</sup> ABLIKIM 21W measurement is limited by statistics.

NODE=M191R02

NODE=M191R02

NODE=M191R02;LINKAGE=A

$\Gamma(\pi^\pm\chi_{c2}(1P))/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma$
not seen		14	<sup>1</sup> ABLIKIM	21W	BES3 $e^+e^- \rightarrow \chi_{cJ}\pi^+\pi^+$	

<sup>1</sup> ABLIKIM 21W measurement is limited by statistics.

NODE=M191R03

NODE=M191R03

NODE=M191R03;LINKAGE=A

$\Gamma(\pi^\pm\psi(3770))/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma$
not seen		<sup>1</sup> ABLIKIM	19AR	BES3 $e^+e^- \rightarrow \pi^+\pi^-D\bar{D}$	

<sup>1</sup> From a measurement of  $\sigma(e^+e^- \rightarrow \pi^+\pi^-D\bar{D})$  between  $\sqrt{s} = 4.08$  and 4.6 GeV.

NODE=M191R00

NODE=M191R00

NODE=M191R00;LINKAGE=A

**X(4050)<sup>±</sup> REFERENCES**

ABLIKIM	21W	PR D103 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12B	PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08	PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M191

REFID=61221  
REFID=59910  
REFID=54042  
REFID=52535

NODE=M223

**X(4055)<sup>±</sup>**

$$I^G(J^{PC}) = 1^+(?^{?^-})$$

*I, G, C* need confirmation.

**OMITTED FROM SUMMARY TABLE**

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

NODE=M223

Needs confirmation. Seen by WANG 15A in the  $\psi(2S)\pi^+$  invariant mass distribution in  $\psi(4360) \rightarrow \psi(2S)\pi^+\pi^-$  decay.

**X(4055)<sup>±</sup> MASS**

NODE=M223M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>4054 ±3 ±1</b>	<sup>1</sup> WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
4039.3±6.0	<sup>2</sup> ABLIKIM	18K BES3	$e^+e^- \rightarrow \pi^0\pi^0\psi(2S)$
4032.1±2.4	<sup>3</sup> ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$

NODE=M223M

OCCUR=2

<sup>1</sup> Statistical significance of 3.5  $\sigma$ .<sup>2</sup> Statistical error only, with significance of 5.9  $\sigma$  (from a fit with a 19% CL). Identified as the same structure observed in ABLIKIM 17V in  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$  decays.<sup>3</sup> Statistical error only, with significance of 9.2  $\sigma$ . From an unbinned maximum likelihood fit of the  $\pi^+\pi^-\psi(2S)$  Dalitz plot from data collected at  $\sqrt{s} = 4.416$  GeV for a  $J^C = 1^+$  state. The fit does not match the detailed structure of the data, having a C.L. of only 8%.NODE=M223M;LINKAGE=A  
NODE=M223M;LINKAGE=C

NODE=M223M;LINKAGE=B

**X(4055)<sup>±</sup> WIDTH**

NODE=M223W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>45 ±11 ±6</b>	<sup>1</sup> WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
31.9±14.8	<sup>2</sup> ABLIKIM	18K BES3	$e^+e^- \rightarrow \pi^0\pi^0\psi(2S)$
26.1± 5.3	<sup>3</sup> ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$

NODE=M223W

<sup>1</sup> Statistical significance of 3.5  $\sigma$ .<sup>2</sup> Statistical error only, with significance of 5.9  $\sigma$  (from a fit with a 19% CL). Identified as the same structure observed in ABLIKIM 17V in  $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$  decays.<sup>3</sup> Statistical error only, with significance of 9.2  $\sigma$ . From an unbinned maximum likelihood fit of the  $\pi^+\pi^-\psi(2S)$  Dalitz plot from data collected at  $\sqrt{s} = 4.416$  GeV for a  $J^C = 1^+$  state. The fit does not match the detailed structure of the data, having a C.L. of only 8%.NODE=M223W;LINKAGE=A  
NODE=M223W;LINKAGE=C

NODE=M223W;LINKAGE=B

**X(4055)<sup>±</sup> DECAY MODES**

NODE=M223215;NODE=M223

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi^+\psi(2S)$	seen
$\Gamma_2$ $\pi^\pm\psi(3770)$	not seen

DESIG=1

DESIG=2

**X(4055)<sup>±</sup> BRANCHING RATIOS**

NODE=M223225

$\Gamma(\pi^+\psi(2S))/\Gamma_{\text{total}}$		$\Gamma_1/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

NODE=M223R01  
NODE=M223R01<sup>1</sup> Statistical significance of 3.5  $\sigma$ .

NODE=M223R01;LINKAGE=A

$\Gamma(\pi^\pm\psi(3770))/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^- D\bar{D}$

<sup>1</sup> From a measurement of  $\sigma(e^+e^- \rightarrow \pi^+\pi^- D\bar{D})$  between  $\sqrt{s} = 4.08$  and 4.6 GeV.

NODE=M223R00  
NODE=M223R00

NODE=M223R00;LINKAGE=A

 $X(4055)^\pm$  REFERENCES

ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18K	PR D97 052001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)

NODE=M223

REFID=59910  
REFID=58896  
REFID=58029  
REFID=59611  
REFID=56839

NODE=M240

 $X(4100)^\pm$ 

$$I^G(J^{PC}) = 1^-(???)$$

## OMITTED FROM SUMMARY TABLE

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

NODE=M240

Reported by AAIJ 18AN in the  $\eta_c(1S)\pi^-$  invariant mass distribution in  $B^0 \rightarrow \eta_c(1S)K^+\pi^-$  decays with a significance of  $3.4\sigma$ .  $J^P = 0^+$  or  $1^-$  assignment consistent with data.

 $X(4100)^\pm$  MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4096 \pm 20^{+18}_{-22}$	AAIJ	18AN LHCB	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

NODE=M240M

NODE=M240M

 $X(4100)^\pm$  WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$152 \pm 58^{+60}_{-35}$	AAIJ	18AN LHCB	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

NODE=M240W

NODE=M240W

 $X(4100)^\pm$  DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\eta_c(1S)\pi^-$	seen
$\Gamma_2$ $\pi^\pm\psi(3770)$	not seen

NODE=M240215;NODE=M240

DESIG=1

DESIG=2

 $X(4100)^\pm$  BRANCHING RATIOS $\Gamma(\eta_c(1S)\pi^-)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> AAIJ	18AN LHCB	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$

<sup>1</sup> AAIJ 18AN quotes a fit fraction for  $B^0 \rightarrow X(4100)^- K^+ \rightarrow \eta_c(1S)\pi^- K^+$  of  $(3.3 \pm 1.1^{+1.2}_{-1.1})\%$  from an amplitude analysis.

NODE=M240225

NODE=M240R01  
NODE=M240R01

NODE=M240R01;LINKAGE=A

 $\Gamma(\pi^\pm\psi(3770))/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	<sup>1</sup> ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^- D\bar{D}$

<sup>1</sup> From a measurement of  $\sigma(e^+e^- \rightarrow \pi^+\pi^- D\bar{D})$  between  $\sqrt{s} = 4.08$  and 4.6 GeV.

NODE=M240R00  
NODE=M240R00

NODE=M240R00;LINKAGE=A

 $X(4100)^\pm$  REFERENCES

ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18AN	EPJ C78 1019	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M240

REFID=59910  
REFID=59335

**Z<sub>c</sub>(4200)**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

*I*, *G*, *C* need confirmation.

OMITTED FROM SUMMARY TABLE  
was  $X(4200)^\pm$

This state shows properties different from a conventional  $q\bar{q}$  state.  
A candidate for an exotic structure. See the review on non- $q\bar{q}$  states.

Reported by CHILIKIN 14 in  $J/\psi\pi^+$  at a significance of  $6.2\sigma$ . Assignments of  $0^-$ ,  $1^-$ ,  $2^-$ , and  $2^+$  excluded at  $6.1\sigma$ ,  $7.4\sigma$ ,  $4.4\sigma$ , and  $7.0\sigma$  level, respectively. Needs confirmation.

NODE=M231

NODE=M231

**Z<sub>c</sub>(4200) MASS**

NODE=M231M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4196^{+31+17}_{-29-13}$	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

NODE=M231M

**Z<sub>c</sub>(4200) WIDTH**

NODE=M231W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$370 \pm 70^{+70}_{-132}$	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

NODE=M231W

**Z<sub>c</sub>(4200) DECAY MODES**

NODE=M231215;NODE=M231

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi\pi^+$	seen

DESIG=1

**Z<sub>c</sub>(4200) BRANCHING RATIOS**

NODE=M231220

$\Gamma(J/\psi\pi^+)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma$
<b>seen</b>	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$	
••• We do not use the following data for averages, fits, limits, etc. •••				
possibly seen	<sup>1</sup> AAIJ	19R	LHCB $B^0 \rightarrow K^+ \pi^- J/\psi + \text{c.c.}$	

NODE=M231R01  
NODE=M231R01

<sup>1</sup> From a model-independent analysis.

NODE=M231R01;LINKAGE=C

**Z<sub>c</sub>(4200) REFERENCES**

NODE=M231

AAIJ	19R	PRL 122 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)
CHILIKIN	14	PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)

REFID=59776  
REFID=56344

**$Z_{cs}(4220)^+$** 

$$I(J^P) = \frac{1}{2}(1^+)$$

NODE=M260

## OMITTED FROM SUMMARY TABLE

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M260

Seen by AAIJ 21E in  $B^+ \rightarrow Z_{cs}(4220)^+ \phi$  with  $Z_{cs}(4220)^+ \rightarrow J/\psi K^+$  using an amplitude analysis of  $B^+ \rightarrow J/\psi \phi K^+$  with a significance (accounting for systematic uncertainties) of  $5.9 \sigma$ . The  $J^P = 1^+$  assignment is favored over  $1^-$  with a significance of  $2 \sigma$  and other assignments are disfavored by  $4.9 \sigma$ .

 **$Z_{cs}(4220)^+$  MASS**

NODE=M260M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4216 \pm 24^{+43}_{-30}$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M260M

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.9 \sigma$ .

NODE=M260M;LINKAGE=A

 **$Z_{cs}(4220)^+$  WIDTH**

NODE=M260W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$233 \pm 52^{+97}_{-73}$	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M260W

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.9 \sigma$ .

NODE=M260W;LINKAGE=A

 **$Z_{cs}(4220)^+$  DECAY MODES**

NODE=M260215;NODE=M260

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad J/\psi K^+$	seen

DESIG=1

$\Gamma(J/\psi K^+)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	<sup>1</sup> AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M260R01  
NODE=M260R01

<sup>1</sup> From an amplitude analysis of the decay  $B^+ \rightarrow J/\psi \phi K^+$  with a significance of  $5.9 \sigma$ .

NODE=M260R01;LINKAGE=A

 **$Z_{cs}(4220)^+$  REFERENCES**

NODE=M260

AAIJ 21E PRL 127 082001 R. Aaij *et al.* (LHCb Collab.) JP

REFID=61150

**$R_{c0}(4240)$** 
 $I^G(J^{PC}) = 1^+(0^{--})$   
 I, G, C need confirmation.

 OMITTED FROM SUMMARY TABLE  
 was  $X(4240)^\pm$ 

 Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

 Spin and parity assignment  $J^P = 0^-$  is favored over  $1^-$ ,  $2^-$ , and  $2^+$  by  $8\sigma$  and over  $1^+$  by  $1\sigma$ , according to the four-dimensional amplitude analysis of AAIJ 14AG.

NODE=M216

NODE=M216

 **$R_{c0}(4240)$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4239 \pm 18^{+45}_{-10}$	<sup>1</sup> AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

<sup>1</sup>From a 4-dimensional analysis when a second, lower mass resonance is allowed in the  $Z_c(4430)$  fit, with significance  $6\sigma$  including systematic variations.

NODE=M216M

NODE=M216M

NODE=M216M;LINKAGE=AA

 **$R_{c0}(4240)$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$220 \pm 47^{+108}_{-74}$	<sup>1</sup> AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

<sup>1</sup>From a 4-dimensional analysis when a second, lower mass resonance is allowed in the  $Z_c(4430)$  fit, with significance  $6\sigma$  including systematic variations.

NODE=M216W

NODE=M216W

NODE=M216W;LINKAGE=AA

 **$R_{c0}(4240)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \pi^- \psi(2S)$	seen

NODE=M216215;NODE=M216

DESIG=1

 **$R_{c0}(4240)$  BRANCHING RATIOS**

$\Gamma(\pi^- \psi(2S))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

<sup>1</sup>From a 4-dimensional analysis when a second, lower mass resonance is allowed in the  $Z_c(4430)$  fit. No partial branching fraction quoted.

NODE=M216225

NODE=M216R01

NODE=M216R01

NODE=M216R01;LINKAGE=AA

 **$R_{c0}(4240)$  REFERENCES**

AAIJ	14AG PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	---------------------	-----------------------	----------------

NODE=M216

REFID=55896



**X(4250)<sup>±</sup>**

$$I^G(J^{PC}) = 1^-(?^{?+})$$

$I, G, C$  need confirmation.

## OMITTED FROM SUMMARY TABLE

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

Observed by MIZUK 08 in the  $\pi^+\chi_{c1}(1P)$  invariant mass distribution in  $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$  decays. Not seen by LEES 12B in this same mode after accounting for  $K\pi$  resonant mass and angular structure.

NODE=M192

NODE=M192

**X(4250)<sup>±</sup> MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4248^{+44+180}_{-29-35}$	<sup>1</sup> MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

<sup>1</sup> From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192M

NODE=M192M

NODE=M192M;LINKAGE=MI

**X(4250)<sup>±</sup> WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$177^{+54+316}_{-39-61}$	<sup>1</sup> MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

<sup>1</sup> From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192W

NODE=M192W

NODE=M192W;LINKAGE=MI

**X(4250)<sup>±</sup> DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \pi^+\chi_{c1}(1P)$	seen

NODE=M192215;NODE=M192

DESIG=1

**X(4250)<sup>±</sup> BRANCHING RATIOS**

$$\frac{\Gamma(\pi^+\chi_{c1}(1P))}{\Gamma_{\text{total}}} \quad \Gamma_1/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	<sup>1</sup> MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen <sup>2</sup> LEES 12B BABR  $B \rightarrow K\pi\chi_{c1}(1P)$

<sup>1</sup> With a product branching fraction measurement of  $B(\bar{B}^0 \rightarrow K^-X(4250)^+) \times B(X(4250)^+ \rightarrow \pi^+\chi_{c1}(1P)) = (4.0^{+2.3+19.7}_{-0.9-0.5}) \times 10^{-5}$ .

<sup>2</sup> With a product branching fraction limit of  $B(\bar{B}^0 \rightarrow X(4250)^+K^-) \times B(X(4250)^+ \rightarrow \chi_{c1}\pi^+) < 4.0 \times 10^{-5}$  at 90% CL.

NODE=M192225

NODE=M192R01  
NODE=M192R01

NODE=M192R01;LINKAGE=MI

NODE=M192R01;LINKAGE=LE

**X(4250)<sup>±</sup> REFERENCES**

LEES	12B	PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08	PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M192

REFID=54042  
REFID=52535

**Z<sub>c</sub>(4430)**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

G, C need confirmation.

was X(4430)<sup>±</sup>

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

First seen by CHOI 08 in  $B \rightarrow K\pi^+\psi(2S)$  decays, confirmed by AAIJ 14AG, and confirmed in a model-independent way by AAIJ 15BH. Also seen by CHILIKIN 14 in  $B \rightarrow K^+\pi J/\psi$  decays.

$J^P$  was determined by CHILIKIN 13 and AAIJ 14AG.

NODE=M195

NODE=M195

**Z<sub>c</sub>(4430) MASS**

NODE=M195M

NODE=M195M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**4478<sup>+15</sup><sub>-18</sub> OUR AVERAGE**

4475 ± 7 <sup>+15</sup> <sub>-25</sub>	1 AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
--	--------	-----------	------------------------------------

4485 ± 22 <sup>+28</sup> <sub>-11</sub>	1 CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
---	------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4443 <sup>+15+19</sup> <sub>-12-13</sub>	2 MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
--	---------	---------	--------------------------------

4433 ± 4 ± 2	3 CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
--------------	--------	---------	--------------------------------

<sup>1</sup> From a four-dimensional amplitude analysis.

<sup>2</sup> From a Dalitz plot analysis. Superseded by CHILIKIN 13.

<sup>3</sup> Superseded by MIZUK 09 and CHILIKIN 13.

NODE=M195M;LINKAGE=A  
NODE=M195M;LINKAGE=MI  
NODE=M195M;LINKAGE=CH

**Z<sub>c</sub>(4430) WIDTH**

NODE=M195W

NODE=M195W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

**181 ± 31 OUR AVERAGE**

172 ± 13 <sup>+37</sup> <sub>-34</sub>	1 AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
--	--------	-----------	------------------------------------

200 <sup>+41+26</sup> <sub>-46-35</sub>	1 CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
---	------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

107 <sup>+86+74</sup> <sub>-43-56</sub>	2 MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
---	---------	---------	--------------------------------

45 <sup>+18+30</sup> <sub>-13-13</sub>	3 CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
--	--------	---------	--------------------------------

<sup>1</sup> From a four-dimensional amplitude analysis.

<sup>2</sup> From a Dalitz plot analysis. Superseded by CHILIKIN 13.

<sup>3</sup> Superseded by MIZUK 09 and CHILIKIN 13.

NODE=M195W;LINKAGE=A  
NODE=M195W;LINKAGE=MI  
NODE=M195W;LINKAGE=CH

**Z<sub>c</sub>(4430) DECAY MODES**

NODE=M195215;NODE=M195

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\pi^+\psi(2S)$	seen
$\Gamma_2$ $\pi^+J/\psi$	seen

DESIG=1

DESIG=2

**Z<sub>c</sub>(4430) BRANCHING RATIOS**

NODE=M195225

$\Gamma(\pi^+\psi(2S))/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma$
---	-------------------

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	1 AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
------	--------	-----------	------------------------------------

seen	2 CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
------	------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	3 AUBERT	09AA BABR	$B \rightarrow K\pi^+\psi(2S)$
----------	----------	-----------	--------------------------------

seen	4 MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
------	---------	---------	--------------------------------

<sup>1</sup> From a four-dimensional amplitude analysis. No product of branching fractions quoted.

<sup>2</sup> From a four-dimensional amplitude analysis. Measured a product of branching fractions  $B(B^0 \rightarrow Z_c(4430)^- K^+) \times B(Z_c(4430)^- \rightarrow \psi(2S)\pi^-) = (6.0^{+1.7+2.5}_{-2.0-1.4}) \times 10^{-5}$ .

<sup>3</sup> AUBERT 09AA quotes  $B(B^+ \rightarrow \bar{K}^0 Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+\psi(2S)) < 4.7 \times 10^{-5}$  and  $B(\bar{B}^0 \rightarrow K^- Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+\psi(2S)) < 3.1 \times 10^{-5}$  at 95% CL.

<sup>4</sup> Measured a product of branching fractions  $B(\bar{B}^0 \rightarrow K^- Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+\psi(2S)) = (3.2^{+1.8+5.3}_{-0.9-1.6}) \times 10^{-5}$ . Superseded by CHILIKIN 13.

NODE=M195R01  
NODE=M195R01

NODE=M195R01;LINKAGE=AA  
NODE=M195R01;LINKAGE=A

NODE=M195R01;LINKAGE=AU

NODE=M195R01;LINKAGE=MI

$\Gamma(\pi^+ J/\psi)/\Gamma_{\text{total}}$  $\Gamma_2/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	1,2 CHILIKIN	14	BELL $\bar{B}^0 \rightarrow K^- \pi^+ J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<b>not seen</b>	3 AUBERT	09AA BABR	$B \rightarrow K \pi^+ J/\psi$
1 CHILIKIN 14 reports $B(\bar{B}^0 \rightarrow Z_c(4430)^+ K^-) \times B(Z_c(4430)^+ \rightarrow J/\psi \pi^+) = (5.4^{+4.0+1.1}_{-1.0-0.9}) \times 10^{-6}$ .			
2 A broad enhancement seen by AAIJ 19R in the decays $B^0 \rightarrow J/\psi \pi^+ K^-$ at 4600 MeV can be due to an interplay of $Z_c(4430)$ , $Z_c(4200)$ and the fitting polynomials.			
3 AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+ J/\psi) < 1.5 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- Z_c(4430)^+) \times B(Z_c(4430)^+ \rightarrow \pi^+ J/\psi) < 0.4 \times 10^{-5}$ at 95% CL.			

NODE=M195R02  
NODE=M195R02

NODE=M195R02;LINKAGE=A

NODE=M195R02;LINKAGE=E

NODE=M195R02;LINKAGE=AU

 **$Z_c(4430)$  REFERENCES**

AAIJ	19R	PRL 122 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BH	PR D92 112009	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AG	PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
CHILIKIN	14	PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)
CHILIKIN	13	PR D88 074026	K. Chilikin <i>et al.</i>	(BELLE Collab.) JP
AUBERT	09AA	PR D79 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)
MIZUK	09	PR D80 031104	R. Mizuk <i>et al.</i>	(BELLE Collab.)
CHOI	08	PRL 100 142001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)

NODE=M195

REFID=59776

REFID=57110

REFID=55896

REFID=56344

REFID=55551

REFID=52940

REFID=52960

REFID=52178

NODE=M232

 **$X(5568)^\pm$**  $I(J^P) = ?(??)$ 

## OMITTED FROM SUMMARY TABLE

Seen as a peak in the  $B_s \pi^\pm$  mass spectrum with a significance of more than  $3\sigma$  by ABAZOV 16E and ABAZOV 18A in inclusive  $p\bar{p}$  collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

NODE=M232

 **$X(5568)^\pm$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>5566.9^{+3.2+0.6}_{-3.1-1.2}</math></b>	278	1 ABAZOV	18A D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5567.8 \pm 2.9^{+0.9}_{-1.9}$	133	2 ABAZOV	16E D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$

NODE=M232M

NODE=M232M

1 From the combined analysis of  $B_s^0 \rightarrow J/\psi \phi$  and  $B_s^0 \rightarrow D_s^\pm \mu^\mp X$  decays.

2 Assumes  $X(5568)^\pm \rightarrow B_s \pi^\pm$  decay. If  $X(5568)^\pm \rightarrow B_s^* \pi^\pm$  decay is assumed, the mass shifts upward by 49 MeV.

NODE=M232M;LINKAGE=B

NODE=M232M;LINKAGE=A

 **$X(5568)^\pm$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b><math>18.6^{+7.9+3.5}_{-6.1-3.8}</math></b>	278	1 ABAZOV	18A D0	$p\bar{p} \rightarrow B_s \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$21.9 \pm 6.4^{+5.0}_{-2.5}$	133	ABAZOV	16E D0	$p\bar{p} \rightarrow B_s \pi^\pm X$

NODE=M232W

NODE=M232W

1 From the combined analysis of  $B_s^0 \rightarrow J/\psi \phi$  and  $B_s^0 \rightarrow D_s^\pm \mu^\mp X$  decays.

NODE=M232W;LINKAGE=B

 **$X(5568)^\pm$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $B_s \pi^\pm$	seen

NODE=M232215;NODE=M232

DESIG=1

 $\Gamma(B_s \pi^\pm)/\Gamma_{\text{total}}$  $\Gamma_1/\Gamma$ 

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	145	1 ABAZOV	18A D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$
<b>seen</b>	133	2 ABAZOV	16E D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$

NODE=M232R01

NODE=M232R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	<sup>3</sup> AABOUD	18L ATLS	$pp \rightarrow B_S^0 \pi^\pm X$
not seen	<sup>4</sup> AALTONEN	18A CDF	$p\bar{p} \rightarrow B_S^0 \pi^\pm X$
not seen	<sup>5</sup> SIRUNYAN	18J CMS	$pp \rightarrow B_S^0 \pi^\pm X$
not seen	<sup>6</sup> AAIJ	16AI LHCB	$pp \rightarrow B_S^0 \pi^\pm X$

OCCUR=2

<sup>1</sup> With  $B_S$  mesons reconstructed in decays to  $D_S^\pm \mu^\mp X$ .

NODE=M232R01;LINKAGE=F  
NODE=M232R01;LINKAGE=A

<sup>2</sup> Seen in  $p\bar{p}$  collisions at 1.96 TeV at a rate of  $(8.6 \pm 1.9 \pm 1.4)\%$  relative to inclusive  $B_S$  production in the kinematic region  $10 < p_T(B_S) < 30$  GeV/c, with  $B_S$  mesons reconstructed in decays to  $J/\psi\phi$ . An alternative possibility,  $X(5568)^\pm \rightarrow B_S^* \pi^\pm$  with a missing  $\gamma$ , could not be ruled out.

NODE=M232R01;LINKAGE=E

<sup>3</sup> Not seen in  $24.4 \text{ fb}^{-1}$  of  $pp$  collision data at  $\sqrt{s} = 7$  and 8 TeV with  $B_S$  mesons reconstructed in decays to  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $X(5568)^\pm \rightarrow B_S \pi^\pm$  relative to inclusive  $B_S$  production is less than 1.5% at  $p_T(B_S) > 10$  GeV/c and less than 1.6% at  $p_T(B_S) > 15$  GeV/c at 95% CL.

NODE=M232R01;LINKAGE=D

<sup>4</sup> Not seen in  $9.6 \text{ fb}^{-1}$  of  $p\bar{p}$  collision data at  $\sqrt{s} = 1.96$  TeV with  $B_S$  mesons reconstructed in decays to  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $X(5568)^\pm \rightarrow B_S \pi^\pm$  relative to inclusive  $B_S$  production is less than 6.7% at 95% CL.

NODE=M232R01;LINKAGE=C

<sup>5</sup> Not seen in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions data at  $\sqrt{s} = 8$  TeV with  $B_S$  mesons reconstructed in decays to  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $X(5568)^\pm \rightarrow B_S \pi^\pm$  relative to inclusive  $B_S$  production is less than 1.1% at  $p_T(B_S) > 10$  GeV/c and less than 1.0% at  $p_T(B_S) > 15$  GeV/c at 95% CL.

NODE=M232R01;LINKAGE=B

<sup>6</sup> Not seen in  $3 \text{ fb}^{-1}$  of  $pp$  collision data at  $\sqrt{s} = 7$  and 8 TeV in a scan over the  $X(5568)$  mass and width, with  $B_S$  mesons reconstructed in decays to  $D_S^- \pi^+$  or  $J/\psi\phi$ . An upper limit on the production rate times branching fraction for  $X(5568)^\pm \rightarrow B_S \pi^\pm$  relative to inclusive  $B_S$  production is less than 2.1% at  $p_T(B_S) > 10$  GeV/c at 90% CL.

### $X(5568)^\pm$ REFERENCES

NODE=M232

AABOUD	18L	PRL 120 202007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58829
AALTONEN	18A	PRL 120 202006	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=58828
ABAZOV	18A	PR D97 092004	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=58937
SIRUNYAN	18J	PRL 120 202005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58827
AAIJ	16AI	PRL 117 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57549
ABAZOV	16E	PRL 117 022003	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=57453

NODE=M268

**$X(6900)$**

$$J^{PC} = ???$$

OMITTED FROM SUMMARY TABLE

State incompatible with a  $q\bar{q}$  structure. See the review on "Heavy Non- $q\bar{q}$  Mesons."

NODE=M268

### $X(6900)$ MASS

NODE=M268M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>6886 \pm 11 \pm 11</math></b>	<sup>1</sup> AAIJ	20AY LHCB	$pp \rightarrow J/\psi J/\psi X$

NODE=M268M

<sup>1</sup> AAIJ 20AY value assumes the interference of a Breit-Wigner shape with non-resonant single-parton scattering. Without interference, the mass is  $6905 \pm 11 \pm 7$  MeV.

NODE=M268M;LINKAGE=A

### $X(6900)$ WIDTH

NODE=M268W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>168 \pm 33 \pm 69</math></b>	<sup>1</sup> AAIJ	20AY LHCB	$pp \rightarrow J/\psi J/\psi X$

NODE=M268W

<sup>1</sup> AAIJ 20AY value assumes the interference of a Breit-Wigner shape with non-resonant single-parton scattering. Without interference, the width is  $80 \pm 19 \pm 33$  MeV.

NODE=M268W;LINKAGE=A

### $X(6900)$ DECAY MODES

NODE=M268215;NODE=M268

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $J/\psi J/\psi$	seen

DESIG=1

$\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	20AY LHCB	$pp \rightarrow J/\psi J/\psi X$

NODE=M268R00  
NODE=M268R00

**X(6900) REFERENCES**

AAIJ

20AY SCIB 65 1983

R. Aaij *et al.*

(LHCb Collab.)

NODE=M268

REFID=61631

NODE=M207

**Z<sub>b</sub>(10610)**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

was X(10610)

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

NODE=M207

Observed by BONDAR 12 in  $\Upsilon(5S)$  decays to  $\Upsilon(nS)\pi^+\pi^-$  ( $n = 1, 2, 3$ ) and  $h_b(mP)\pi^+\pi^-$  ( $m = 1, 2$ ).  $J^P = 1^+$  is favored from angular analyses.

**Z<sub>b</sub>(10610)<sup>±</sup> MASS**

NODE=M207M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10607.2±2.0</b>	<sup>1</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10608.5±3.4 <sup>+3.7</sup> <sub>-1.4</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10608.1±1.2 <sup>+1.5</sup> <sub>-0.2</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10607.4±1.5 <sup>+0.8</sup> <sub>-0.2</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10611 ±4 ±3	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10609 ±2 ±3	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10608 ±2 ±3	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10605 ±2 <sup>+3</sup> <sub>-1</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10599 <sup>+6</sup> <sub>-3</sub> <sup>+5</sup> <sub>-4</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M207M

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

<sup>1</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>2</sup> Correlated with the corresponding result from BONDAR 12.

<sup>3</sup> Superseded by the average measurement of BONDAR 12.

NODE=M207M;LINKAGE=BO

NODE=M207M;LINKAGE=A

NODE=M207M;LINKAGE=BN

NODE=M207M0

**Z<sub>b</sub>(10610)<sup>0</sup> MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10609±4±4</b>	<sup>1</sup> KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$

NODE=M207M0

<sup>1</sup> From a simultaneous fit to the KROKOVNY 13 Dalitz analysis of  $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$  decays with fixed width  $\Gamma(Z_b(10610)^0) = 18.4$  MeV.

NODE=M207M0;LINKAGE=A

**Z<sub>b</sub>(10610)<sup>±</sup> WIDTH**

NODE=M207W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>18.4± 2.4</b>	<sup>1</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
18.5± 5.3 <sup>+6.1</sup> <sub>-2.3</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
20.8± 2.5 <sup>+0.3</sup> <sub>-2.1</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
18.7± 3.4 <sup>+2.5</sup> <sub>-1.3</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
22.3± 7.7 <sup>+3.0</sup> <sub>-4.0</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
24.2± 3.1 <sup>+2.0</sup> <sub>-3.0</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
17.6± 3.0±3.0	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
11.4 <sup>+ 4.5+2.1</sup> <sub>- 3.9-1.2</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
13 <sup>+10</sup> <sub>- 8</sub> <sup>+9</sup> <sub>-7</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M207W

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

<sup>1</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>2</sup> Correlated with the corresponding result from BONDAR 12.

<sup>3</sup> Superseded by the average measurement of BONDAR 12.

NODE=M207W;LINKAGE=BO

NODE=M207W;LINKAGE=A

NODE=M207W;LINKAGE=BN

**Z<sub>b</sub>(10610) DECAY MODES**

NODE=M207215;NODE=M207

Mode	Fraction ( $\Gamma_i/\Gamma$ )	
$\Gamma_1$ $\Upsilon(1S)\pi^+$	$(5.4^{+1.9}_{-1.5}) \times 10^{-3}$	DESIG=1
$\Gamma_2$ $\Upsilon(1S)\pi^0$	not seen	DESIG=9
$\Gamma_3$ $\Upsilon(2S)\pi^+$	$(3.6^{+1.1}_{-0.8})\%$	DESIG=2
$\Gamma_4$ $\Upsilon(2S)\pi^0$	seen	DESIG=10
$\Gamma_5$ $\Upsilon(3S)\pi^+$	$(2.1^{+0.8}_{-0.6})\%$	DESIG=3
$\Gamma_6$ $\Upsilon(3S)\pi^0$	seen	DESIG=11
$\Gamma_7$ $h_b(1P)\pi^+$	$(3.5^{+1.2}_{-0.9})\%$	DESIG=4
$\Gamma_8$ $h_b(2P)\pi^+$	$(4.7^{+1.7}_{-1.3})\%$	DESIG=5
$\Gamma_9$ $B^+\bar{B}^0$	not seen	DESIG=8
$\Gamma_{10}$ $B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	$(85.6^{+2.1}_{-2.9})\%$	DESIG=6

**Z<sub>b</sub>(10610) BRANCHING RATIOS**

NODE=M207225

$\Gamma(\Upsilon(1S)\pi^+)/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$

NODE=M207R01  
NODE=M207R01

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>5.4^{+1.6+1.1}_{-1.3-0.8}</math></b>	<sup>1</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$

<sup>1</sup> Assuming the Z<sub>b</sub>(10610) decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^+\bar{B}^{*0} + \bar{B}^0B^{*+}$ , and using the results from BONDAR 12 and MIZUK 16.

NODE=M207R01;LINKAGE=A

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_2/\Gamma$

NODE=M207R09  
NODE=M207R09

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	KROKOVNY	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0\pi^0$

$\Gamma(\Upsilon(2S)\pi^+)/\Gamma_{\text{total}}$   $\Gamma_3/\Gamma$

NODE=M207R02  
NODE=M207R02

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>3.62^{+0.76+0.79}_{-0.59-0.53}</math></b>	<sup>1</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

<sup>1</sup> Assuming the Z<sub>b</sub>(10610) decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^+\bar{B}^{*0} + \bar{B}^0B^{*+}$ , and using the results from BONDAR 12 and MIZUK 16.

NODE=M207R02;LINKAGE=A

$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$   $\Gamma_4/\Gamma$

NODE=M207R10  
NODE=M207R10

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> KROKOVNY	13	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^0\pi^0$

<sup>1</sup> Combined significance in  $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$ , including systematics, of  $6.5\sigma$ .

NODE=M207R10;LINKAGE=A

$\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$   $\Gamma_5/\Gamma$

NODE=M207R03  
NODE=M207R03

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b><math>2.15^{+0.55+0.60}_{-0.42-0.43}</math></b>	<sup>1</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

<sup>1</sup> Assuming the Z<sub>b</sub>(10610) decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^+\bar{B}^{*0} + \bar{B}^0B^{*+}$ , and using the results from BONDAR 12 and MIZUK 16.

NODE=M207R03;LINKAGE=A

$\Gamma(\Upsilon(3S)\pi^0)/\Gamma_{\text{total}}$  $\Gamma_6/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	<sup>1</sup> KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^0\pi^0$

<sup>1</sup> Combined significance in  $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$ , including systematics, of  $6.5\sigma$ .

NODE=M207R11  
NODE=M207R11

NODE=M207R11;LINKAGE=A

 $\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$  $\Gamma_7/\Gamma$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.45<sup>+0.87+0.86</sup><sub>-0.71-0.63</sub></b>	<sup>1</sup> GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$
<b>possibly seen</b>	<sup>2</sup> MIZUK 16	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
<b>seen</b>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M207R04  
NODE=M207R04

NODE=M207R04;LINKAGE=C

NODE=M207R04;LINKAGE=A  
NODE=M207R04;LINKAGE=B $\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$  $\Gamma_8/\Gamma$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.67<sup>+1.24+1.18</sup><sub>-1.00-0.89</sub></b>	<sup>1</sup> GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$
<b>possibly seen</b>	<sup>2</sup> MIZUK 16	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$
<b>seen</b>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M207R05  
NODE=M207R05

NODE=M207R05;LINKAGE=C

NODE=M207R05;LINKAGE=A  
NODE=M207R05;LINKAGE=B $\Gamma(B^+ \bar{B}^0)/\Gamma_{\text{total}}$  $\Gamma_9/\Gamma$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>not seen</b>	GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^0$

NODE=M207R08  
NODE=M207R08 $[\Gamma(B^+ \bar{B}^{*0}) + \Gamma(B^{*+} \bar{B}^0)]/\Gamma_{\text{total}}$  $\Gamma_{10}/\Gamma$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>85.6<sup>+1.5+1.5</sup><sub>-2.0-2.1</sub></b>	357	<sup>1</sup> GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- B^{*+} \bar{B}^0$

NODE=M207R00  
NODE=M207R00

NODE=M207R00;LINKAGE=A

 $[\Gamma(B^+ \bar{B}^{*0}) + \Gamma(B^{*+} \bar{B}^0)] / [\Gamma(\Upsilon(1S)\pi^+) + \Gamma(\Upsilon(2S)\pi^+) + \Gamma(\Upsilon(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)]$  $\Gamma_{10}/(\Gamma_1 + \Gamma_3 + \Gamma_5 + \Gamma_7 + \Gamma_8)$ 

VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5.93<sup>+0.99+1.01</sup><sub>-0.69-0.73</sub></b>	357	<sup>1</sup> GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

NODE=M207R07  
NODE=M207R07

NODE=M207R07;LINKAGE=A

 **$Z_b(10610)$  REFERENCES**

NODE=M207

GARMASH 16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=57446
MIZUK 16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
GARMASH 15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=56811
KROKOVNY 13	PR D88 052016	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=55588
BONDAR 12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)	REFID=53963

**Z<sub>b</sub>(10650)**

$$I^G(J^{PC}) = 1^+(1^{+-})$$

*I, G, C* need confirmation.

was  $X(10650)^\pm$ 

Properties incompatible with a  $q\bar{q}$  structure (exotic state). See the review on non- $q\bar{q}$  states.

Observed by BONDAR 12 in  $\Upsilon(5S)$  decays to  $\Upsilon(nS)\pi^+\pi^-$  ( $n = 1, 2, 3$ ) and  $h_b(mP)\pi^+\pi^-$  ( $m = 1, 2$ ).  $J^P = 1^+$  is favored from angular analyses.

NODE=M208

NODE=M208

**Z<sub>b</sub>(10650) MASS**

NODE=M208M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>10652.2±1.5</b>	<sup>1</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10656.7±5.0 <sup>+1.1</sup> <sub>-3.1</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10650.7±1.5 <sup>+0.5</sup> <sub>-0.2</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10651.2±1.0 <sup>+0.4</sup> <sub>-0.3</sub>	<sup>2</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10657 ±6 ±3	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10651 ±2 ±3	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10652 ±1 ±2	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10654 ±3 <sup>+1</sup> <sub>-2</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 <sup>+2</sup> <sub>-3</sub> <sup>+3</sup> <sub>-2</sub>	<sup>3</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M208M

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

<sup>1</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>2</sup> Correlated with the corresponding result from BONDAR 12.

<sup>3</sup> Superseded by the average measurement of BONDAR 12.

NODE=M208M;LINKAGE=BO

NODE=M208M;LINKAGE=A

NODE=M208M;LINKAGE=BN

**Z<sub>b</sub>(10650) WIDTH**

NODE=M208W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>11.5± 2.2</b>	<sup>4</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \text{hadrons}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
12.1 <sup>+11.3</sup> <sub>-4.8</sub> <sup>+2.7</sup> <sub>-0.6</sub>	<sup>5</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
14.2± 3.7 <sup>+0.9</sup> <sub>-0.4</sub>	<sup>5</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
9.3± 2.2 <sup>+0.3</sup> <sub>-0.5</sub>	<sup>5</sup> GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
16.3± 9.8 <sup>+6.0</sup> <sub>-2.0</sub>	<sup>6</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
13.3± 3.3 <sup>+4.0</sup> <sub>-3.0</sub>	<sup>6</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
8.4± 2.0± 2.0	<sup>6</sup> BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
20.9 <sup>+5.4</sup> <sub>-4.7</sub> <sup>+2.1</sup> <sub>-5.7</sub>	<sup>6</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
19 ± 7 <sup>+11</sup> <sub>-7</sub>	<sup>6</sup> BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M208W

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

<sup>4</sup> Average of the BONDAR 12 measurements in separate channels.

<sup>5</sup> Correlated with the corresponding result from BONDAR 12.

<sup>6</sup> Superseded by the average measurement of BONDAR 12.

NODE=M208W;LINKAGE=BO

NODE=M208W;LINKAGE=A

NODE=M208W;LINKAGE=BN

**Z<sub>b</sub>(10650)<sup>+</sup> DECAY MODES**

NODE=M208215;NODE=M208

$Z_b(10650)^-$  decay modes are charge conjugates of the modes below.

NODE=M208

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$ $\Upsilon(1S)\pi^+$	$(1.7^{+0.8}_{-0.6}) \times 10^{-3}$
$\Gamma_2$ $\Upsilon(2S)\pi^+$	$(1.4^{+0.6}_{-0.4}) \%$
$\Gamma_3$ $\Upsilon(3S)\pi^+$	$(1.6^{+0.7}_{-0.5}) \%$
$\Gamma_4$ $h_b(1P)\pi^+$	$(8.4^{+2.9}_{-2.4}) \%$

DESIG=1

DESIG=2

DESIG=3

DESIG=4



$\Gamma_5$	$h_b(2P)\pi^+$	(15 $\pm$ 4 ) %	DESIG=5
$\Gamma_6$	$B^+\bar{B}^0$	not seen	DESIG=8
$\Gamma_7$	$B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	not seen	DESIG=6
$\Gamma_8$	$B^{*+}\bar{B}^{*0}$	(74 $\frac{+4}{-6}$ ) %	DESIG=7

### $Z_b(10650)$ BRANCHING RATIOS

#### $\Gamma(\Upsilon(1S)\pi^+)/\Gamma_{\text{total}}$

 $\Gamma_1/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.7<math>\frac{+0.7+0.3}{-0.6-0.2}</math></b>	<sup>7</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$

<sup>7</sup> Assuming the  $Z_b(10650)$  decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

NODE=M208225

NODE=M208R01  
NODE=M208R01

NODE=M208R01;LINKAGE=A

#### $\Gamma(\Upsilon(2S)\pi^+)/\Gamma_{\text{total}}$

 $\Gamma_2/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.39<math>\frac{+0.48+0.34}{-0.38-0.23}</math></b>	<sup>8</sup> GARMASH	16	$e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

<sup>8</sup> Assuming the  $Z_b(10650)$  decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

NODE=M208R02  
NODE=M208R02

NODE=M208R02;LINKAGE=A

#### $\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$

 $\Gamma_3/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.63<math>\frac{+0.53+0.39}{-0.42-0.28}</math></b>	<sup>9</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

<sup>9</sup> Assuming the  $Z_b(10650)$  decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

NODE=M208R03  
NODE=M208R03

NODE=M208R03;LINKAGE=A

#### $\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$

 $\Gamma_4/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>8.41<math>\frac{+2.43+1.49}{-2.12-1.06}</math></b>	<sup>10</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	<sup>11</sup> MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
seen	<sup>12</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

<sup>10</sup> Assuming the  $Z_b(10650)$  decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

<sup>11</sup> Using  $e^+e^-$  energies near the  $\Upsilon(11020)$ .

<sup>12</sup> Using  $e^+e^-$  energies near the  $\Upsilon(10860)$ .

NODE=M208R04  
NODE=M208R04

NODE=M208R04;LINKAGE=C

NODE=M208R04;LINKAGE=A  
NODE=M208R04;LINKAGE=B

#### $\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$

 $\Gamma_5/\Gamma$ 

<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>14.7<math>\frac{+3.2+2.8}{-2.8-2.3}</math></b>	<sup>13</sup> GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	<sup>14</sup> MIZUK	16	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$
seen	<sup>15</sup> BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

<sup>13</sup> Assuming the  $Z_b(10650)$  decay width is saturated by the channels  $\pi^+\Upsilon(1S, 2S, 3S)$ ,  $\pi^+h_b(1P, 2P)$ , and  $B^{*+}\bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16.

<sup>14</sup> Using  $e^+e^-$  energies near the  $\Upsilon(11020)$ .

<sup>15</sup> Using  $e^+e^-$  energies near the  $\Upsilon(10860)$ .

NODE=M208R05  
NODE=M208R05

NODE=M208R05;LINKAGE=C

NODE=M208R05;LINKAGE=A  
NODE=M208R05;LINKAGE=B

$\Gamma(B^+\bar{B}^0)/\Gamma_{\text{total}}$					$\Gamma_6/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^0$	NODE=M208R08 NODE=M208R08

$[\Gamma(B^+\bar{B}^{*0}) + \Gamma(B^{*+}\bar{B}^0)]/\Gamma_{\text{total}}$					$\Gamma_7/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT		
not seen	GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$	NODE=M208R00 NODE=M208R00

$\Gamma(B^{*+}\bar{B}^{*0})/\Gamma_{\text{total}}$					$\Gamma_8/\Gamma$	
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT		
$73.7^{+3.4+2.7}_{-4.4-3.5}$	161	<sup>16</sup> GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$	NODE=M208R06 NODE=M208R06

<sup>16</sup> Assuming the  $Z_b(10650)$  decay width is saturated by the channels  $\pi^+ \Upsilon(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^{*+} \bar{B}^{*0}$ , and using the results from BONDAR 12 and MIZUK 16. Using the mass and width of the  $Z_b(10650)$  from BONDAR 12.

NODE=M208R06;LINKAGE=A

$\Gamma(B^{*+}\bar{B}^{*0})/[\Gamma(\Upsilon(1S)\pi^+) + \Gamma(\Upsilon(2S)\pi^+) + \Gamma(\Upsilon(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)]$					$\Gamma_8/(\Gamma_1+\Gamma_2+\Gamma_3+\Gamma_4+\Gamma_5)$
VALUE (units $10^{-2}$ )	EVTS	DOCUMENT ID	TECN	COMMENT	
not seen					NODE=M208R07 NODE=M208R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.80^{+0.69+0.54}_{-0.40-0.36}$	161	<sup>17</sup> GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$	NODE=M208R07;LINKAGE=A
----------------------------------	-----	-----------------------	----	------	--	------------------------

<sup>17</sup> Combined with the results of BONDAR 12 and MIZUK 16. Not independent from  $Z_b(10650)$  branching fractions to  $\pi^+ \Upsilon(1S, 2S, 3S)$ ,  $\pi^+ h_b(1P, 2P)$ , and  $B^{*+} \bar{B}^{*0}$ .

### $Z_b(10650)$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=57446
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=56811
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)	REFID=53963

## Further States

OMITTED FROM SUMMARY TABLE

This section contains states observed by a single group or states poorly established that thus need confirmation.

QUANTUM NUMBERS, MASSES, WIDTHS, AND BRANCHING RATIOS

$X(360) \quad I^G(J^{PC}) = ?^?(?^{?+})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$360 \pm 7 \pm 9$	$64 \pm 18$	2.3k	<sup>1</sup> ABRAAMYAN	09	CNTR 2.75 $dC \rightarrow \gamma\gamma X$
<sup>1</sup> Not seen in $pC \rightarrow \gamma\gamma X$ at 5.5 GeV/c.					

NODE=M300K08  
NODE=M300K08

NODE=M300K08;LINKAGE=AB

$X(1070) \quad I^G(J^{PC}) = ?^?(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	COMMENT		
$1072 \pm 1$	$3.5 \pm 0.5$	<sup>1</sup> VLADIMIRSK...08	$40 \pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$		NODE=M300J07 NODE=M300J07
<sup>1</sup> Supersedes GRIGOR'EV 05.					

NODE=M300J07;LINKAGE=VL

$X(1110) \quad I^G(J^{PC}) = 0^+(\text{even}^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1107 \pm 4$	$111 \pm 8 \pm 15$	DAFTARI	87	DBC	$0. \bar{p}n \rightarrow \rho^- \pi^+ \pi^-$

NODE=M300J30  
NODE=M300J30

$f_0(1200-1600) \quad I^G(J^{PC}) = 0^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1323 \pm 8$	$237 \pm 20$	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$	NODE=M300J98 NODE=M300J98
$1480^{+100}_{-150}$	$1030^{+80}_{-170}$	<sup>1</sup> ANISOVICH	03	SPEC	
$1530^{+90}_{-250}$	$560 \pm 40$	<sup>2</sup> ANISOVICH	03	SPEC	OCCUR=2

<sup>1</sup>K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ ,  $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$ ,  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^0$ ,  $K^+ K_S^0 \pi^-$  at rest,  $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$ ,  $K_S^0 K^- \pi^0$ ,  $K_S^0 K_S^0 \pi^-$  at rest.

NODE=M300;LINKAGE=KM

<sup>2</sup>K-matrix pole from combined analysis of  $\pi^- p \rightarrow \pi^0 \pi^0 n$ ,  $\pi^- p \rightarrow K \bar{K} n$ ,  $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$ ,  $\pi^0 \eta \eta$ ,  $\pi^0 \pi^0 \eta$  at rest.

NODE=M300;LINKAGE=MK

<b>X(1420)</b> $I^G(J^{PC}) = 2^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1420±20	160 ± 10	FILIPPI	00	OBLX 0 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$	NODE=M300J61 NODE=M300J61

<b>X(1545)</b> $I^G(J^{PC}) = ??(?^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1545±3	6.0 ± 2.5	<sup>1</sup> VLADIMIRSK...08		40 $\pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$	NODE=M300K07 NODE=M300K07

<sup>1</sup>Supersedes VLADIMIRSKII 00.

NODE=M300K07;LINKAGE=VL

<b>X(1575)</b> $I^G(J^{PC}) = ??(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1576 <sup>+49+98</sup> <sub>-55-91</sub>	818 <sup>+22+64</sup> <sub>-23-133</sub>	<sup>1</sup> ABLIKIM	06S	BES $J/\psi \rightarrow K^+ K^- \pi^0$	NODE=M300J08 NODE=M300J08

<sup>1</sup>A broad peak observed at  $K^+ K^-$  invariant mass. Mass and width above are its pole position. The observed branching ratio is  $B(J/\psi \rightarrow X \pi^0) B(X \rightarrow K^+ K^-) = (8.5 \pm 0.6^{+2.7}_{-3.6}) \times 10^{-4}$ .

NODE=M300J08;LINKAGE=AB

<b>X(1600)</b> $I^G(J^{PC}) = 2^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1600±100	400 ± 200	<sup>1</sup> ALBRECHT 91F	ARG	10.2 $e^+ e^- \rightarrow e^+ e^- 2(\pi^+ \pi^-)$	NODE=M300J99 NODE=M300J99

<sup>1</sup>Our estimate.

NODE=M300J99;LINKAGE=A

<b>X(1650)</b> $I^G(J^{PC}) = 0^-(?^{?-})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1652±7	<50	100	PROKOSHKIN 96	GAM2	32,38 $\pi p \rightarrow \omega \eta n$

NODE=M300J62  
NODE=M300J62

<b>X(1730)</b> $I^G(J^{PC}) = ??(?^{?+})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1731.0±1.2±2.0	3.2 ± 0.8 ± 1.3	58	VLADIMIRSK...07	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 X$

NODE=M300K06  
NODE=M300K06

<b>f<sub>2</sub>(1750)</b> $I^G(J^{PC}) = 0^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1755±10	67 ± 12	870	<sup>1</sup> SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M300JAM  
NODE=M300JAM

<b>Γ(K<math>\bar{K}</math>)</b>					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
17±5	870	<sup>2</sup> SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	NODE=M300JA1 NODE=M300JA1

<b>Γ(γγ)</b>					
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.13±0.04	870	<sup>2</sup> SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	NODE=M300JA2 NODE=M300JA2

<b>Γ(ππ)</b>					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.3±1.0	870	<sup>2</sup> SCHEGELSKY 06A	RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	NODE=M300JA3 NODE=M300JA3

$\Gamma(\eta\eta)$ 

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2.0±0.5	870	<sup>2</sup> SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

<sup>1</sup> From analysis of L3 data at 91 and 183–209 GeV.

<sup>2</sup> From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M300JA4  
NODE=M300JA4

NODE=M300JAM;LINKAGE=SC  
NODE=M300JA;LINKAGE=SC

**X(1775)**  $I^G(J^{PC}) = 1^-(?^-+)$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1763±20	192 ± 60	CONDO 91	SHF	$\gamma p \rightarrow (p\pi^+)(\pi^+\pi^-\pi^-)$
1787±18	118 ± 60	CONDO 91	SHF	$\gamma p \rightarrow n\pi^+\pi^+\pi^-$

NODE=M300J60  
NODE=M300J60

OCCUR=2

**X(1850 - 3100)**  $I^G(J^{PC}) = ?^?(1^{--})$ 

$\Gamma(e^+e^-) \cdot B(X \rightarrow \text{hadrons})$ (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<120	90	<sup>1</sup> ANASHIN	11	KEDR $e^+e^- \rightarrow \text{hadrons}$

NODE=M300K28  
NODE=M300K28

<sup>1</sup> This limit is center-of-mass energy dependent. We quote the most stringent one.

NODE=M300K28;LINKAGE=AN

**X(1855)**  $I^G(J^{PC}) = ?^?(???)$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1856.6±5	20 ± 5	BRIDGES	86D	SPEC $0. \bar{p}d \rightarrow \pi\pi N$

NODE=M300J31  
NODE=M300J31

**X(1870)**  $I^G(J^{PC}) = ?^?(2??)$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1870±40	250 ± 30	ALDE	86D	GAM4 $100 \pi^- p \rightarrow 2\eta X$

NODE=M300J45  
NODE=M300J45

**a<sub>3</sub>(1875)**  $I^G(J^{PC}) = 1^-(3^{++})$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1874±43±96	385 ± 121 ± 114	CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M300J95  
NODE=M300J95

**B(a<sub>3</sub>(1875) → f<sub>2</sub>(1270)π)/B(a<sub>3</sub>(1875) → ρπ)**

VALUE	DOCUMENT ID	TECN	COMMENT
0.8±0.2	<sup>1</sup> CHUNG 02	B852	$18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M300B7  
NODE=M300B7

<sup>1</sup> Using the observable fractions of 50.0% ρπ, 56.5% f<sub>2</sub>π, and 11.8% ρ<sub>3</sub>π.

NODE=M300B;LINKAGE=C1

**B(a<sub>3</sub>(1875) → ρ<sub>3</sub>(1690)π)/B(a<sub>3</sub>(1875) → ρπ)**

VALUE	DOCUMENT ID	TECN	COMMENT
0.9±0.3	<sup>1</sup> CHUNG 02	B852	$18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M300B8  
NODE=M300B8

<sup>1</sup> Using the observable fractions of 50.0% ρπ, 56.5% f<sub>2</sub>π, and 11.8% ρ<sub>3</sub>π.

NODE=M300B8;LINKAGE=C1

**a<sub>1</sub>(1930)**  $I^G(J^{PC}) = 1^-(1^{++})$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1930 <sup>+30</sup> <sub>-70</sub>	155 ± 45	ANISOVICH 01F	SPEC	$2.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J92  
NODE=M300J92

**X(1935)**  $I^G(J^{PC}) = 1^+(1^{-?})$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1935±20	215 ± 30	EVANGELIS... 79	OMEG	$10,16 \pi^- p \rightarrow \bar{p}pn$

NODE=M300J33  
NODE=M300J33

**ρ<sub>2</sub>(1940)**  $I^G(J^{PC}) = 1^+(2^{--})$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1940±40	155 ± 40	<sup>1</sup> ANISOVICH 02	SPEC	$0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J85  
NODE=M300J85

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J85;LINKAGE=AY

**ω<sub>3</sub>(1945)**  $I^G(J^{PC}) = 0^-(3^{--})$ 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1945±20	115 ± 22	<sup>1</sup> ANISOVICH 02B	SPEC	$0.6-1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J65  
NODE=M300J65

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J65;LINKAGE=AZ

<b><math>a_2(1950)</math></b>		$I^G(J^{PC}) = 1^-(2^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1950^{+30}_{-70}$	$180^{+30}_{-70}$	<sup>1</sup> ANISOVICH	01F SPEC	1.96–2.41 $p\bar{p}$	

NODE=M300K24  
NODE=M300K24

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K24;LINKAGE=AN

<b><math>\omega(1960)</math></b>		$I^G(J^{PC}) = 0^-(1^{--})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1960 \pm 25$	$195 \pm 60$	<sup>1</sup> ANISOVICH	02B SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$	

NODE=M300J79  
NODE=M300J79

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J79;LINKAGE=AZ

<b><math>b_1(1960)</math></b>		$I^G(J^{PC}) = 1^+(1^{+-})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1960 \pm 35$	$230 \pm 50$	<sup>1</sup> ANISOVICH	02 SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$	

NODE=M300J67  
NODE=M300J67

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J67;LINKAGE=AY

<b><math>h_1(1965)</math></b>		$I^G(J^{PC}) = 0^-(1^{+-})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1965 \pm 45$	$345 \pm 75$	<sup>1</sup> ANISOVICH	02B SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$	

NODE=M300J64  
NODE=M300J64

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J64;LINKAGE=AZ

<b><math>f_1(1970)</math></b>		$I^G(J^{PC}) = 0^+(1^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1971 \pm 15$	$240 \pm 45$	ANISOVICH	00J SPEC		

NODE=M300J1  
NODE=M300J1

<b><math>X(1970)</math></b>		$I^G(J^{PC}) = ??(???)$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1970 \pm 10$	$40 \pm 20$	CHLIAPNIK...	80 HBC	32 $K^+p \rightarrow 2K_S^0 2\pi X$	

NODE=M300J46  
NODE=M300J46

<b><math>X(1975)</math></b>		$I^G(J^{PC}) = ??(???)$			
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$1973 \pm 15$	80	30	CASO	70 HBC	11.2 $\pi^-p \rightarrow \rho 2\pi$

NODE=M300J47  
NODE=M300J47

<b><math>\omega_2(1975)</math></b>		$I^G(J^{PC}) = 0^-(2^{--})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
$1975 \pm 20$	$175 \pm 25$	<sup>1</sup> ANISOVICH	02B SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$	

NODE=M300J81  
NODE=M300J81

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J81;LINKAGE=AZ

<b><math>a_2(1990)</math></b>		$I^G(J^{PC}) = 1^-(2^{++})$			
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2050 \pm 10 \pm 40$	$190 \pm 22 \pm 100$	18k	<sup>1</sup> SCHEGELSKY	06 RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
$2003 \pm 10 \pm 19$	$249 \pm 23 \pm 32$		LU	05 B852	$18 \pi^-p \rightarrow \omega\pi^-\pi^0p$

NODE=M300J2  
NODE=M300J2

<sup>1</sup> From analysis of L3 data at 183–209 GeV.

NODE=M300J2;LINKAGE=SC

$\Gamma(\gamma\gamma) \Gamma(\pi^+ \pi^- \pi^0) / \Gamma(\text{total})$ 

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.11±0.04±0.05	18k	<sup>1</sup> SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

<sup>1</sup> From analysis of L3 data at 183–209 GeV.

NODE=M300J2G  
 NODE=M300J2G

NODE=M300J2G;LINKAGE=SC

$\rho(2000)$ $I^G(J^{PC}) = 1^+(1^- -)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2000±30	260 ± 45	<sup>1</sup> BUGG	04C	RVUE Compilation
~ 1988	~ 244	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J77  
 NODE=M300J77

NODE=M300;LINKAGE=AY

$f_2(2000)$ $I^G(J^{PC}) = 0^+(2^+ +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2001±10	312 ± 32	ANISOVICH	00J	SPEC
~ 1996	~ 134	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

NODE=M300J25  
 NODE=M300J25

$X(2000)$ $I^G(J^{PC}) = 1^-(?^?+)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1964±35	225 ± 50	<sup>1</sup> ARMSTRONG	93D	E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
~ 2100	~ 500	<sup>1</sup> ANTIPOV	77	CIBS	- 25 $\pi^- p \rightarrow p\pi^- \rho_3$
2214±15	355 ± 21	<sup>2</sup> BALTAY	77	HBC	0 15 $\pi^- p \rightarrow \Delta^{++} 3\pi$
2080±40	340 ± 80	KALELKAR	75	HBC	+ 15 $\pi^+ p \rightarrow p\pi^+ \rho_3$

<sup>1</sup> Cannot determine spin to be 3.  
<sup>2</sup> BALTAY 77 favors  $J^P = ,3^+$ .

NODE=M300K01  
 NODE=M300K01

NODE=M300K01;LINKAGE=AA  
 NODE=M300K01;LINKAGE=B

$X(2000)$ $I^G(J^{PC}) = ?^?(4^+ +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
1998±3±5	<15	VLADIMIRSK...03	SPEC	$\pi^- p \rightarrow K_S^0 K_S^0 M M$

NODE=M300J97  
 NODE=M300J97

$\eta(2010)$ $I^G(J^{PC}) = 0^+(0^- +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	
2010 <sup>+35</sup> <sub>-60</sub>	270 ± 60	ANISOVICH	00J	SPEC

NODE=M300J5  
 NODE=M300J5

$\pi_1(2015)$ $I^G(J^{PC}) = 1^-(1^- +)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2014±20±16	230 ± 32 ± 73	145k	LU	05	B852 18 $\pi^- p \rightarrow \omega\pi^- \pi^0 p$
2001±30±92	333 ± 52 ± 49	69k	KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$

NODE=M300J05  
 NODE=M300J05

$a_0(2020)$ $I^G(J^{PC}) = 1^-(0^+ +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	
2025±30	330 ± 75	ANISOVICH	99C	SPEC

NODE=M300J6  
 NODE=M300J6

$X(2020)$ $I^G(J^{PC}) = ?^?(?^?+)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2015±3	10 ± 4	FERRER	99	RVUE $\pi p \rightarrow p\bar{p}\pi(\pi)$

NODE=M300J34  
 NODE=M300J34

$h_3(2025)$ $I^G(J^{PC}) = 0^-(3^+ -)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2025±20	145 ± 30	<sup>1</sup> ANISOVICH	02B	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J78  
 NODE=M300J78

NODE=M300J78;LINKAGE=AZ

$h_3(2030)$ $I^G(J^{PC}) = 1^+(3^+ -)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2032±12	117 ± 11	<sup>1</sup> ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J69  
 NODE=M300J69

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J69;LINKAGE=AY

<b><math>a_2(2030)</math></b> $I^G(J^{PC}) = 1^-(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2030±20	205 ± 30	<sup>1</sup> ANISOVICH	01F	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K23  
NODE=M300K23

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K23;LINKAGE=AN

<b><math>a_3(2030)</math></b> $I^G(J^{PC}) = 1^-(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2031±12	150 ± 18	<sup>1</sup> ANISOVICH	01F	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K20  
NODE=M300K20

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K20;LINKAGE=AN

<b><math>\eta_2(2030)</math></b> $I^G(J^{PC}) = 0^+(2^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2030±5±15	205 ± 10 ± 15	ANISOVICH	00E	SPEC	

NODE=M300J8  
NODE=M300J8

**$B(a_2\pi)_{L=0}/B(a_2\pi)_{L=2}$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.05±0.03	<sup>1</sup> ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300B1  
NODE=M300B1

<sup>1</sup> Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B1;LINKAGE=AN

**$B(a_0\pi)/B(a_2\pi)_{L=2}$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.10±0.08	<sup>1</sup> ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300B2  
NODE=M300B2

<sup>1</sup> Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B2;LINKAGE=AN

**$B(f_2\eta)/B(a_2\pi)_{L=2}$**

VALUE	DOCUMENT ID	TECN	COMMENT
0.13±0.06	<sup>1</sup> ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300B3  
NODE=M300B3

<sup>1</sup> Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B3;LINKAGE=AN

<b><math>f_3(2050)</math></b> $I^G(J^{PC}) = 0^+(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2048±8	213 ± 34	ANISOVICH	00J	SPEC	2.0 $p\bar{p} \rightarrow \eta\pi^0\pi^0$

NODE=M300J7  
NODE=M300J7

<b><math>f_0(2060)</math></b> $I^G(J^{PC}) = 0^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
~ 2050	~ 120	<sup>1</sup> OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2060	~ 50	<sup>1</sup> OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$

NODE=M300J59  
NODE=M300J59

<sup>1</sup> See SEMENOV 99 and KLOET 96.

OCCUR=2

NODE=M300J;LINKAGE=A

<b><math>\pi(2070)</math></b> $I^G(J^{PC}) = 1^-(0^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2070±35	310 <sup>+100</sup> <sub>-50</sub>	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J91  
NODE=M300J91

<b><math>X(2075)</math></b> $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2075±12±5	90 ± 35 ± 9	<sup>1</sup> ABLIKIM	04J	BES2	$J/\psi \rightarrow K^- p\bar{\Lambda}$

NODE=M300J01  
NODE=M300J01

<sup>1</sup> From a fit in the region  $M_{p\bar{\Lambda}} - M_p - M_{\Lambda} < 150$  MeV. *S*-wave in the  $p\bar{\Lambda}$  system preferred.

NODE=M300J01;LINKAGE=AB

A similar near-threshold enhancement in the  $p\bar{\Lambda}$  system is observed in  $B^+ \rightarrow p\bar{\Lambda}\bar{D}^0$  by CHEN 11F.

<b><math>X(2080)</math></b> $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2080±10	110 ± 20	KREYMER	80	STRC	13 $\pi^- d \rightarrow p\bar{p}n(n_s)$

NODE=M300J35  
NODE=M300J35

<b>X(2080)</b> $I^G(J^{PC}) = ?^?(3^{-?})$						NODE=M300J37 NODE=M300J37
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2080±10	190 ± 15	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p \bar{p} n$	
<hr/>						
<b>a<sub>1</sub>(2095)</b> $I^G(J^{PC}) = 1^-(1^{++})$						NODE=M300J04 NODE=M300J04
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2096±17±121	451 ± 41 ± 81	69k	KUHN	04	B852 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$	
<b>B(a<sub>1</sub>(2095) → f<sub>1</sub>(1285)π) / B(a<sub>1</sub>(2095) → a<sub>1</sub>(1260))</b>						NODE=M300B03 NODE=M300B03
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
3.18±0.64	69k	KUHN	04	B852	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$	
<hr/>						
<b>η(2100)</b> $I^G(J^{PC}) = 0^+(0^{-+})$						NODE=M300J48 NODE=M300J48
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2050 <sup>+30+75</sup> <sub>-24-26</sub>	250 <sup>+36+181</sup> <sub>-30-164</sub>		1	ABLIKIM	16N BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
2103±50	187 ± 75	586	2	BISELLO	89B DM2 $J/\psi \rightarrow 4\pi\gamma$	
<p><sup>1</sup> From a partial wave analysis of <math>J/\psi \rightarrow \gamma \phi \phi</math>, for which the primary signal is <math>\eta(2225) \rightarrow \phi \phi</math>, and that also finds significant signals for for <math>0^{-+}</math> phase space, <math>f_0(2100)</math>, <math>f_2(2010)</math>, <math>f_2(2300)</math>, <math>f_2(2340)</math>, and a previously unseen <math>0^{-+}</math> state <math>X(2500)</math> (<math>M = 2470^{+15+101}_{-19-23}</math> MeV, <math>\Gamma = 230^{+64+56}_{-35-33}</math> MeV).</p> <p><sup>2</sup> ASTON 81B sees no peak, has 850 events in Ajinenko+Barth bins. ARESTOV 80 sees no peak.</p>						
<hr/>						
<b>X(2100)</b> $I^G(J^{PC}) = ?^?(0^{??})$						NODE=M300J49 NODE=M300J49
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2100±40	250 ± 40	ALDE	86D	GAM4	100 $\pi^- p \rightarrow 2\eta X$	
<hr/>						
<b>X(2110)</b> $I^G(J^{PC}) = 1^+(3^{-?})$						NODE=M300J36 NODE=M300J36
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2110±10	330 ± 20	EVANGELIS...	79	OMEG	10,16 $\pi^- p \rightarrow \bar{p} p n$	
<hr/>						
<b>X(2120)</b> $I^G(J^{PC}) = ?^?(0^{??})$						NODE=M300A07 NODE=M300A07
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2122.4±6.7 <sup>+4.7</sup> <sub>-2.7</sub>	83 ± 16 <sup>+31</sup> <sub>-11</sub>	647	ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$	
<hr/>						
<b>f<sub>2</sub>(2140)</b> $I^G(J^{PC}) = 0^+(2^{++})$						NODE=M300J50 NODE=M300J50
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2141±12	49 ± 28	389	GREEN	86	MPSF 400 $pA \rightarrow 4KX$	
<hr/>						
<b>X(2150)</b> $I^G(J^{PC}) = ?^?(2^{+?})$						NODE=M300J38 NODE=M300J38
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2150±10	260 ± 10	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p \bar{p} n$	
<hr/>						
<b>a<sub>2</sub>(2175)</b> $I^G(J^{PC}) = 1^-(2^{++})$						NODE=M300J88 NODE=M300J88
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2175±40	310 <sup>+90</sup> <sub>-45</sub>	ANISOVICH	01F	SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$	
<hr/>						
<b>η(2190)</b> $I^G(J^{PC}) = 0^+(0^{-+})$						NODE=M300J13 NODE=M300J13
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2190±50	850 ± 100	BUGG	99	BES		
<hr/>						
<b>ω<sub>2</sub>(2195)</b> $I^G(J^{PC}) = 0^-(2^{--})$						NODE=M300J82 NODE=M300J82
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		
2195±30	225 ± 40	1	ANISOVICH	02B	SPEC 0.6-1.9 $p\bar{p} \rightarrow \omega \eta, \omega \pi^0 \pi^0$	



<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J82;LINKAGE=AZ

<b><math>\omega(2205)</math></b>		$I^G(J^{PC}) = 0^-(1^{--})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2222 ± 7 ± 2	59 ± 30 ± 6	<sup>1</sup> ABLIKIM	22i BES3	2.0–3.8 $e^+ e^- \rightarrow \omega \pi^0 \pi^0$
2205 ± 30	350 ± 90	<sup>2</sup> ANISOVICH	02B SPEC	0.6–1.9 $p \bar{p} \rightarrow \omega \eta, \omega \pi^0 \pi^0$

NODE=M300J80  
NODE=M300J80

<sup>1</sup> From the fit to the cross section by the coherent sum of resonant component parametrized by a modified Breit-Wigner amplitude and a phase-space contribution for the continuum. The observed structure can be also due to  $\omega(2290)$  or  $\omega(2330)$ .

NODE=M300J80;LINKAGE=A

<sup>2</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J80;LINKAGE=AZ

<b><math>\Gamma(\omega(2205) \rightarrow e^+ e^-) \times \Gamma(\omega(2205) \rightarrow \omega \pi^0 \pi^0) / \Gamma(\text{total})</math></b>		
VALUE (eV)	DOCUMENT ID	TECN COMMENT
0.3 ± 0.1 ± 0.1	<sup>1</sup> ABLIKIM	22i BES3 2.0–3.8 $e^+ e^- \rightarrow \omega \pi^0 \pi^0$

NODE=M300A04  
NODE=M300A04

<sup>1</sup> From a solution of the fit to the cross section by the coherent sum of resonant component parametrized by a modified Breit-Wigner amplitude and a phase-space contribution for the continuum. The observed structure can be also due to  $\omega(2290)$  or  $\omega(2330)$ . The other solution gives  $13.8 \pm 6.6 \pm 5.2$  eV.

NODE=M300A04;LINKAGE=A

<b><math>X(2210)</math></b>		$I^G(J^{PC}) = ??(???)$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2210 <sup>+79</sup> <sub>-21</sub>	203 <sup>+437</sup> <sub>-87</sub>	EVANGELIS...	79B OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
2207 ± 22	130	CASO	70 HBC	11.2 $\pi^- p$

NODE=M300J51  
NODE=M300J51

<b><math>X_2(2210)</math></b>		$I^G(J^{PC}) = 0^+(2^{++})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2210 ± 60	360 ± 120	<sup>1</sup> KLEMP	22 RVUE	$J/\psi \rightarrow \gamma \pi^0 \pi^0,$ $\gamma K_S^0 K_S^0$

NODE=M300A05  
NODE=M300A05

<sup>1</sup> Fit of the tensor partial waves from BES3 in the multipole basis. Might be a cluster of  $J^{PC} = 2^{++}$  resonances. The ratio of decay widths  $K K^- / \pi \pi$  is  $0.23 \pm 0.05$ .

NODE=M300A05;LINKAGE=A

<b><math>h_1(2215)</math></b>		$I^G(J^{PC}) = 0^-(1^{+-})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2215 ± 40	325 ± 55	<sup>1</sup> ANISOVICH	02B SPEC	0.6–1.9 $p \bar{p} \rightarrow \omega \eta, \omega \pi^0 \pi^0$

NODE=M300J27  
NODE=M300J27

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J27;LINKAGE=AZ

<b><math>\rho_2(2225)</math></b>		$I^G(J^{PC}) = 1^+(2^{--})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2225 ± 35	335 <sup>+100</sup> <sub>-50</sub>	<sup>1</sup> ANISOVICH	02 SPEC	0.6–1.9 $p \bar{p} \rightarrow \omega \pi^0,$ $\omega \eta \pi^0, \pi^+ \pi^-$

NODE=M300J70  
NODE=M300J70

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J70;LINKAGE=AY

<b><math>\rho_4(2230)</math></b>		$I^G(J^{PC}) = 1^+(4^{--})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2230 ± 25	210 ± 30	<sup>1</sup> ANISOVICH	02 SPEC	0.6–1.9 $p \bar{p} \rightarrow \omega \pi^0,$ $\omega \eta \pi^0, \pi^+ \pi^-$

NODE=M300J74  
NODE=M300J74

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J74;LINKAGE=AY

<b><math>b_1(2240)</math></b>		$I^G(J^{PC}) = 1^+(1^{+-})$		
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2240 ± 35	320 ± 85	<sup>1</sup> ANISOVICH	02 SPEC	0.6–1.9 $p \bar{p} \rightarrow \omega \pi^0,$ $\omega \eta \pi^0, \pi^+ \pi^-$

NODE=M300J87  
NODE=M300J87

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J87;LINKAGE=AY

**$f_2(2240)$**   $I^G(J^{PC}) = 0^+(2^{++})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2240±15	241 ± 30	<sup>1</sup> ANISOVICH 00J	SPEC	1.92–2.41 $p\bar{p}$
~ 2226	~ 226	HASAN 94	RVUE	$p\bar{p} \rightarrow \pi\pi$

NODE=M300K26  
NODE=M300K26

• • • We do not use the following data for averages, fits, limits, etc. • • •

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

NODE=M300K26;LINKAGE=AN

**$b_3(2245)$**   $I^G(J^{PC}) = 1^+(3^{+-})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2245±50	320 ± 70	<sup>1</sup> BUGG 04C	RVUE	

NODE=M300K10  
NODE=M300K10

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300K10;LINKAGE=AY

**$\eta_2(2250)$**   $I^G(J^{PC}) = 0^+(2^{-+})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2248±20	280 ± 20	ANISOVICH 00I	SPEC	
2267±14	290 ± 50	ANISOVICH 00J	SPEC	

NODE=M300J17  
NODE=M300J17

**$\pi_4(2250)$**   $I^G(J^{PC}) = 1^-(4^{-+})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250±15	215 ± 25	ANISOVICH 01F	SPEC	2.0 $p\bar{p} \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J73  
NODE=M300J73

**$\omega_4(2250)$**   $I^G(J^{PC}) = 0^-(4^{--})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250±30	150 ± 50	<sup>1</sup> ANISOVICH 02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J84  
NODE=M300J84

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J84;LINKAGE=AZ

**$\omega_5(2250)$**   $I^G(J^{PC}) = 0^-(5^{--})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250±70	320 ± 95	<sup>1</sup> BUGG 04	RVUE	

NODE=M300K11  
NODE=M300K11

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300;LINKAGE=AZ

**$\omega_3(2255)$**   $I^G(J^{PC}) = 0^-(3^{--})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255±15	175 ± 30	<sup>1</sup> ANISOVICH 02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J66  
NODE=M300J66

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J66;LINKAGE=AZ

**$a_4(2255)$**   $I^G(J^{PC}) = 1^-(4^{++})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2237 ± 5	291 ± 12	UMAN 06	E835	5.2 $p\bar{p} \rightarrow \eta\eta\pi^0$
2255±40	330 <sup>+110</sup> <sub>-50</sub>	<sup>1</sup> ANISOVICH 01F	SPEC	1.96–2.41 $p\bar{p}$

NODE=M300K21  
NODE=M300K21

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K21;LINKAGE=AN

**$a_2(2255)$**   $I^G(J^{PC}) = 1^-(2^{++})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255±20	230 ± 15	<sup>1</sup> ANISOVICH 01G	SPEC	1.96–2.41 $p\bar{p}$

NODE=M300K22  
NODE=M300K22

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

NODE=M300K22;LINKAGE=AN

<b>X(2260)</b> $I^G(J^{PC}) = 0^+(4^{+?})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2260 ± 20	400 ± 100	EVANGELIS...	79	OMEG 10,16 $\pi^- p \rightarrow \bar{p} p n$

NODE=M300J40  
NODE=M300J40

<b><math>\rho(2270)</math></b> $I^G(J^{PC}) = 1^+(1^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2265 ± 40	325 ± 80	<sup>1</sup> ANISOVICH	02	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
2280 ± 50	440 ± 110	ATKINSON	85	OMEG 20–70 $\gamma p \rightarrow p\omega\pi^+\pi^-\pi^0$

NODE=M300J86  
NODE=M300J86

<sup>1</sup> From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J86;LINKAGE=AY

<b><math>a_1(2270)</math></b> $I^G(J^{PC}) = 1^-(1^{++})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2270 <sup>+55</sup> <sub>-40</sub>	305 <sup>+70</sup> <sub>-40</sub>	ANISOVICH	01F	SPEC 2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J72  
NODE=M300J72

<b><math>h_3(2275)</math></b> $I^G(J^{PC}) = 0^-(3^{+-})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2275 ± 25	190 ± 45	<sup>1</sup> ANISOVICH	02B	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J28  
NODE=M300J28

<sup>1</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J28;LINKAGE=AZ

<b><math>a_3(2275)</math></b> $I^G(J^{PC}) = 1^-(3^{++})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2275 ± 35	350 <sup>+100</sup> <sub>-50</sub>	<sup>1</sup> ANISOVICH	01G	SPEC 1.96–2.41 $\bar{p}p$

NODE=M300K19  
NODE=M300K19

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

NODE=M300K19;LINKAGE=AN

<b><math>\pi_2(2285)</math></b> $I^G(J^{PC}) = 1^-(2^{-+})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2285 ± 20 ± 25	250 ± 20 ± 25	<sup>1</sup> ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300K25  
NODE=M300K25

<sup>1</sup> Reanalysis of ADOEIT 96 and ANISOVICH 00E.

NODE=M300K25;LINKAGE=AN

<b><math>\omega_3(2285)</math></b> $I^G(J^{PC}) = 0^-(3^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2278 ± 28	224 ± 50	<sup>1</sup> BUGG	04A	RVUE
2285 ± 60	230 ± 40	<sup>2</sup> ANISOVICH	02B	SPEC 0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J83  
NODE=M300J83

<sup>1</sup> Partial wave analysis of the data on  $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$  from BARNES 00.

NODE=M300J83;LINKAGE=BU

<sup>2</sup> From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J83;LINKAGE=AZ

<b><math>\omega(2290)</math></b> $I^G(J^{PC}) = 0^-(1^{--})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2290 ± 20	275 ± 35	<sup>1</sup> BUGG	04A	RVUE

NODE=M300J02  
NODE=M300J02

<sup>1</sup> Partial wave analysis of the data on  $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$  from BARNES 00.

NODE=M300J02;LINKAGE=BU

<b><math>f_2(2295)</math></b> $I^G(J^{PC}) = 0^+(2^{++})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2293 ± 13	216 ± 37	<sup>1</sup> ANISOVICH	00J	SPEC 1.92–2.41 $p\bar{p}$

NODE=M300K27  
NODE=M300K27

<sup>1</sup> From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

NODE=M300K27;LINKAGE=AN

<b><math>f_3(2300)</math></b> $I^G(J^{PC}) = 0^+(3^{++})$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2334 ± 25	200 ± 20	<sup>1</sup> BUGG	04A	RVUE

NODE=M300J19  
NODE=M300J19

<sup>1</sup> Partial wave analysis of the data on  $\rho\bar{p} \rightarrow \bar{\Lambda}\Lambda$  from BARNES 00.

NODE=M300J19;LINKAGE=BU

<b>f<sub>1</sub>(2310)</b> $I^G(J^{PC}) = 0^+(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN		
2310±60	255 ± 70	ANISOVICH	00J	SPEC	

NODE=M300J23  
NODE=M300J23

<b>η(2320)</b> $I^G(J^{PC}) = 0^+(0^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN		
2320±15	230 ± 35	<sup>1</sup> ANISOVICH	00M	SPEC	

NODE=M300J18  
NODE=M300J18<sup>1</sup> From the combined analysis of  $\bar{p}p \rightarrow \eta\eta\eta$  from ANISOVICH 00M and  $\bar{p}p \rightarrow \eta\pi^0\pi^0$  from ANISOVICH 00J.

NODE=M300;LINKAGE=B

<b>η<sub>4</sub>(2330)</b> $I^G(J^{PC}) = 0^+(4^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2328±38	240 ± 90	ANISOVICH	00J	SPEC 2.0 $\rho\bar{p} \rightarrow \eta\pi^0\pi^0$	

NODE=M300J22  
NODE=M300J22

<b>ω(2330)</b> $I^G(J^{PC}) = 0^-(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2330±30	435 ± 75	ATKINSON	88	OMEG 25-50 $\gamma p \rightarrow \rho^\pm \rho^0 \pi^\mp$	

NODE=M300J53  
NODE=M300J53

<b>X(2340)</b> $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2340±20	180 ± 60	126	<sup>1</sup> BALTAY	75	HBC 15 $\pi^+ p \rightarrow p 5\pi$

NODE=M300J54  
NODE=M300J54<sup>1</sup> Dominant decay into  $\rho^0 \rho^0 \pi^+$ . BALTAY 78 finds confirmation in  $2\pi^+ \pi^- 2\pi^0$  events which contain  $\rho^+ \rho^0 \pi^0$  and  $2\rho^+ \pi^-$ .

NODE=M300J;LINKAGE=B1

<b>π(2360)</b> $I^G(J^{PC}) = 1^-(0^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2360±25	$300^{+100}_{-50}$	ANISOVICH	01F	SPEC 2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$	

NODE=M300J90  
NODE=M300J90

<b>X(2360)</b> $I^G(J^{PC}) = ??(4^{+?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2360±10	430 ± 30	ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p\bar{p}n$	

NODE=M300J42  
NODE=M300J42

<b>X(2440)</b> $I^G(J^{PC}) = ??(5^{-?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2440±10	310 ± 20	ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p\bar{p}n$	

NODE=M300J43  
NODE=M300J43

<b>a<sub>6</sub>(2450)</b> $I^G(J^{PC}) = 1^-(6^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2450±130	400 ± 250	CLELAND	82B	SPEC 50 $\pi p \rightarrow K_S^0 K^\pm p$	

NODE=M300K12  
NODE=M300K12

<b>X(2540)</b> $I^G(J^{PC}) = 0^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2539±14 <sup>+38</sup> <sub>-14</sub>	274 <sup>+77</sup> <sub>-61</sub> ±126 <sub>-163</sub>	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M300K30  
NODE=M300K30

<b>Γ(γγ) × B(K<math>\bar{K}</math>)</b>					
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
40 <sup>+9</sup> <sub>-7</sub> ±17 <sub>-40</sub>	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$		

NODE=M300K3G  
NODE=M300K3G

<b>X(2600)</b> $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2618.3±2.0 <sup>+16.3</sup> <sub>-1.4</sub>	195 ± 5 <sup>+26</sup> <sub>-17</sub>	ABLIKIM	22G	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$	

NODE=M300A01  
NODE=M300A01

**$B(J/\psi \rightarrow \gamma X(2600)) \times B(X(2600) \rightarrow f_0(1500)\eta') \times B(f_0(1500) \rightarrow \pi^+\pi^-)$** 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$3.09 \pm 0.21^{+1.14}_{-0.77}$	<sup>1</sup> ABLIKIM	22G BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M300A02  
 NODE=M300A02

<sup>1</sup> The  $\pi^+\pi^-$  mass spectrum is described by a coherent sum of two Breit-Wigner resonances,  $f_0(1500)$  and a new  $X(1540)$  with mass  $1540.2 \pm 7.0^{+36.3}_{-6.1}$  MeV and width  $157 \pm 19^{+11}_{-77}$  MeV.

NODE=M300A02;LINKAGE=A

 **$B(J/\psi \rightarrow \gamma X(2600)) \times B(X(2600) \rightarrow X(1540)\eta') \times B(X(1540) \rightarrow \pi^+\pi^-)$** 

VALUE (units $10^{-5}$ )	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.19^{+0.38}_{-1.21}$	<sup>1</sup> ABLIKIM	22G BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M300A03  
 NODE=M300A03

<sup>1</sup> The  $\pi^+\pi^-$  mass spectrum is described by a coherent sum of two Breit-Wigner resonances,  $f_0(1500)$  and a new  $X(1540)$  with mass  $1540.2 \pm 7.0^{+36.3}_{-6.1}$  MeV and width  $157 \pm 19^{+11}_{-77}$  MeV.

NODE=M300A03;LINKAGE=A

 **$X(2632) \quad I^G(J^{PC}) = ??(???)$** 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$2635.2 \pm 3.3$		<sup>1</sup> EVDOKIMOV 04	SELX	$X(2632) \rightarrow D_s^+ \eta$
$2631.6 \pm 2.1$	< 17	<sup>2</sup> EVDOKIMOV 04	SELX	$X(2632) \rightarrow D_s^0 K^+$

NODE=M300J03  
 NODE=M300J03

OCCUR=2

<sup>1</sup> From a mass difference to  $D_s^+$  of  $666.9 \pm 3.3$  MeV.

NODE=M300J03;LINKAGE=EV  
 NODE=M300J03;LINKAGE=ED

<sup>2</sup> From a mass difference to  $D_s^0$  of  $767.0 \pm 2.0$  MeV.

 **$B(X(2632) \rightarrow D_s^0 K^+)/B(X(2632) \rightarrow D_s^+ \eta)$** 

VALUE	DOCUMENT ID	TECN	COMMENT
$0.14 \pm 0.06$	<sup>1</sup> EVDOKIMOV 04	SELX	

NODE=M300B01  
 NODE=M300B01

<sup>1</sup> Possible interpretation of this decay pattern is discussed by YASUI 07.

NODE=M300B01;LINKAGE=YA

 **$X(2680) \quad I^G(J^{PC}) = ??(???)$** 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$2676 \pm 27$	150	CASO	70 HBC	$11.2 \pi^- p \rightarrow \rho^- \pi^+ \pi^- p$

NODE=M300J55  
 NODE=M300J55

 **$X(2710) \quad I^G(J^{PC}) = ??(6^{+?})$** 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$2710 \pm 20$	$170 \pm 40$	ROZANSKA	80 SPRK	$18 \pi^- p \rightarrow p \bar{p} n$

NODE=M300J44  
 NODE=M300J44

 **$X(2750) \quad I^G(J^{PC}) = ??(7^{-?})$** 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$2747 \pm 32$	$195 \pm 75$	DENNEY	83 LASS	$10 \pi^+ p \rightarrow K^+ K^- \pi^+ p$

NODE=M300J56  
 NODE=M300J56

 **$f_6(3100) \quad I^G(J^{PC}) = 0^+(6^{++})$** 

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$3100 \pm 100$	$700 \pm 130$	BINON	05 GAMS	$33 \pi^- p \rightarrow \eta \eta n$

NODE=M300J06  
 NODE=M300J06

 **$X(3250) \quad I^G(J^{PC}) = ??(???)$  3-Body Decays**

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$3250 \pm 8 \pm 20$	$45 \pm 18$	ALEEV	93 BIS2	$X(3250) \rightarrow \Lambda \bar{p} K^+$
$3265 \pm 7 \pm 20$	$40 \pm 18$	ALEEV	93 BIS2	$X(3250) \rightarrow \bar{\Lambda} p K^-$

NODE=M300J57  
 NODE=M300J57

OCCUR=2

 **$X(3250) \quad I^G(J^{PC}) = ??(???)$  4-Body Decays**

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$3245 \pm 8 \pm 20$	$25 \pm 11$	ALEEV	93 BIS2	$X(3250) \rightarrow \Lambda \bar{p} K^+ \pi^\pm$
$3250 \pm 9 \pm 20$	$50 \pm 20$	ALEEV	93 BIS2	$X(3250) \rightarrow \bar{\Lambda} p K^- \pi^\mp$
$3270 \pm 8 \pm 20$	$25 \pm 11$	ALEEV	93 BIS2	$X(3250) \rightarrow K_S^0 p \bar{p} K^\pm$

NODE=M300J58  
 NODE=M300J58

OCCUR=2

OCCUR=3

 **$X(3350) \quad I^G(J^{PC}) = ??(???)$** 

MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3350^{+10}_{-20} \pm 20$	$70^{+40}_{-30} \pm 40$	$50 \pm 10$	<sup>1</sup> GABYSHEV	06A BELL	$B^- \rightarrow \Lambda_c^+ \bar{p} \pi^-$

NODE=M300J09  
 NODE=M300J09

<sup>1</sup>A similar enhancement in the  $\Lambda_c^+ \bar{p}$  final state is also reported by BABAR collaboration in AUBERT 10H.

NODE=M300J09;LINKAGE=AU

## REFERENCES for Further States

NODE=M300

ABLIKIM	22G	PRL 129 042001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61642
ABLIKIM	22I	PR D105 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61644
KLEMPY	22	PL B830 137171	E. Klempt <i>et al.</i>	(BONN)	REFID=61646
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)	REFID=57512
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ANISOVICH	12	PR D85 014001	A.V. Anisovich <i>et al.</i>		REFID=53961
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ANASHIN	11	PL B703 543	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53932
ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
CHEN	11F	PR D84 071501	P. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53814
AUBERT	10H	PR D82 031102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53363
ABRAAMYAN	09	PR C80 034001	Kh.U. Abraamyan <i>et al.</i>		REFID=53100
VLADIMIRSK...	08	PAN 71 2129	V.V. Vladimirovsky <i>et al.</i>	(ITEP)	REFID=52681
VLADIMIRSK...	07	PAN 70 1706	V. Vladimirovsky <i>et al.</i>		REFID=52058
YASUI	07	PR D76 034009	S. Yasui, M. Oka		REFID=51907
ABLIKIM	06S	PRL 97 142002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51451
GABYSHEV	06A	PRL 97 242001	N. Gabyshev <i>et al.</i>	(BELLE Collab.)	REFID=51565
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
GRIGOR'EV	05	PAN 68 1271	V.K. Grigor'ev <i>et al.</i>	(ITEP)	REFID=50844
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ABLIKIM	04J	PRL 93 112002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50196
BUGG	04	PL B595 556 (errata.)	D.V. Bugg		REFID=49763
BUGG	04A	EPJ C36 161	D.V. Bugg		REFID=50158
BUGG	04C	PRPL 397 257	D.V. Bugg		REFID=50203
EVDOKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)	REFID=50337
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)	REFID=49773
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
VLADIMIRSK...	03	PAN 66 700	V.V. Vladimirovsky <i>et al.</i>		REFID=49419
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	02B	PL B542 19	A.V. Anisovich <i>et al.</i>		REFID=48829
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01C	PL B507 23	A.V. Anisovich <i>et al.</i>		REFID=48325
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352
ANISOVICH	01G	PL B517 273	A.V. Anisovich <i>et al.</i>		REFID=48353
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH	00D	PL B476 15	A.V. Anisovich <i>et al.</i>		REFID=47944
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
ANISOVICH	00I	PL B491 40	A.V. Anisovich <i>et al.</i>		REFID=47949
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
ANISOVICH	00M	PL B496 145	A.V. Anisovich <i>et al.</i>		REFID=48009
BARNES	00	PR C62 055203	P.D. Barnes <i>et al.</i>		REFID=47965
FILIPPI	00	PL B495 284	A. Filippi <i>et al.</i>	(OBELIX Experiment)	REFID=48006
VLADIMIRSKII	00	JETPL 72 486	V.V. Vladimirovskii <i>et al.</i>		REFID=47997
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>		REFID=46902
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>		REFID=46926
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>		REFID=47416
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
BUGG	99	PL B458 511	D.V. Bugg <i>et al.</i>		REFID=46938
FERRER	99	EPJ C10 249	A. Ferrer <i>et al.</i>		REFID=47404
SEMENOV	99	SPU 42 847	S.V. Semenov		REFID=47363
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
PROKOSHKIN	96	PD 41 247	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=45182
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALEEV	93	PAN 56 1358	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)	REFID=43668
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43596
ALBRECHT	91F	ZPHY C50 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41658
CONDO	91	PR D43 2787	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=41588
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
ATKINSON	88	ZPHY C38 535	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=40556
DAFTARI	87	PRL 58 859	I.K. Daftari <i>et al.</i>	(SYRA)	REFID=40412
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
BRIDGES	86D	PL B180 313	D.L. Bridges <i>et al.</i>	(SYRA, BNL, CASE+)	REFID=21984
GREEN	86	PRL 56 1639	D.R. Green <i>et al.</i>	(FNAL, ARIZ, FSU+)	REFID=21872
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=22000
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
ASTON	81B	NP B189 205	D. Aston <i>et al.</i>	(BONN, CERN, EPOL, GLAS+)	REFID=11553
ARESTOV	80	IHEP 80-165	Y.I. Arestov <i>et al.</i>	(SERP)	REFID=22312
CHLIAPNIK...	80	ZPHY C3 285	P.V. Chliapnikov <i>et al.</i>	(SERP, BRUX, MONS)	REFID=21996
KREYMER	80	PR D22 36	A.E. Kreymer <i>et al.</i>	(IND, PURD, SLAC+)	REFID=21970
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)	REFID=21774
EVANGELIS...	79	NP B153 253	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21966
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21967
BALTAY	78	PR D17 52	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21569
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU)	REFID=20847
BALTAY	75	PRL 35 891	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21994
KALELKar	75	Thesis Nevis 207	M.S. Kalelkar	(COLU)	REFID=21564
CASO	70	LCN 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20590