

LIGHT UNFLAVORED MESONS ($S = C = B = 0$)

For $l = 1$ (π, b, ρ, a): $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$;
for $l = 0$ ($\eta, \eta', h, h', \omega, \phi, f, f'$): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

NODE=MXXX005

NODE=MXXX005

NODE=M014

 $f_0(500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

also known as σ ; was $f_0(600)$, $f_0(400-1200)$
See the related review(s):

Scalar Mesons below 1 GeV

f₀(500) T-MATRIX POLE \sqrt{s}

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M014PP

NODE=M014PP

NODE=M014PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(400-550)-i(200-350) OUR ESTIMATE (see Fig. 64.3 in the review)			
$(458 \pm 7^{+4}_{-10}) - i(245 \pm 6^{+7}_{-10})$	1 DANILKIN	21	RVUE Compilation
$(410 \pm 20) - i(240 \pm 15)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(512 \pm 15) - i(188 \pm 12)$	2 ABLIKIM	17	BES3 $J/\psi \rightarrow \gamma 3\pi$
$(440 \pm 10) - i(238 \pm 10)$	3 ALBALADEJO	12	RVUE Compilation
$(445 \pm 25) - i(278^{+22}_{-18})$	4,5 GARCIA-MAR..11	RVUE	Compilation
$(457^{+14}_{-13}) - i(279^{+11}_{-7})$	4,6 GARCIA-MAR..11	RVUE	Compilation
$(442^{+5}_{-8}) - i(274^{+6}_{-5})$	7 MOUSSALLAM11	RVUE	Compilation
$(452 \pm 13) - i(259 \pm 16)$	8 MENNESSIER	10	RVUE Compilation
$(448 \pm 43) - i(266 \pm 43)$	9 MENNESSIER	10	RVUE Compilation
$(455 \pm 6^{+31}_{-13}) - i(278 \pm 6^{+34}_{-43})$	10 CAPRINI	08	RVUE Compilation
$(463 \pm 6^{+31}_{-17}) - i(259 \pm 6^{+33}_{-34})$	11 CAPRINI	08	RVUE Compilation
$(552^{+84}_{-106}) - i(232^{+81}_{-72})$	12 ABLIKIM	07A	BES2 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
$(466 \pm 18) - i(223 \pm 28)$	13 BONVICINI	07	CLEO $D^+ \rightarrow \pi^-\pi^+\pi^+$
$(472 \pm 30) - i(271 \pm 30)$	14 BUGG	07A	RVUE Compilation
$(484 \pm 17) - i(255 \pm 10)$	GARCIA-MAR..07	RVUE	Compilation
$(430) - i(325)$	15 ANISOVICH	06	RVUE Compilation
$(441^{+16}_{-8}) - i(272^{+9}_{-12.5})$	16 CAPRINI	06	RVUE $\pi\pi \rightarrow \pi\pi$
$(470 \pm 50) - i(285 \pm 25)$	17 ZHOU	05	RVUE
$(541 \pm 39) - i(252 \pm 42)$	18 ABLIKIM	04A	BES2 $J/\psi \rightarrow \omega\pi^+\pi^-$
$(528 \pm 32) - i(207 \pm 23)$	19 GALLEGOS	04	RVUE Compilation
$(533 \pm 25) - i(249 \pm 25)$	20 BUGG	03	RVUE
$517 - i240$	BLACK	01	RVUE $\pi\pi \rightarrow \pi\pi$
$(470 \pm 30) - i(295 \pm 20)$	16 COLANGELO	01	RVUE $\pi\pi \rightarrow \pi\pi$
$(535^{+48}_{-36}) - i(155^{+76}_{-53})$	21 ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
$610 \pm 14 - i(310 \pm 13)$	22 SUROVTSEV	01	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
$(540^{+36}_{-29}) - i(193^{+32}_{-40})$	ISHIDA	00B	$\rho\bar{\rho} \rightarrow \pi^0\pi^0\pi^0$
$445 - i235$	HANNAH	99	RVUE π scalar form factor
$(523 \pm 12) - i(259 \pm 7)$	KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$442 - i 227$	OLLER	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
$469 - i203$	OLLER	99B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
$445 - i221$	OLLER	99C	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$
$420 - i 212$	LOCHER	98	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
$440 - i245$	23 DOBADO	97	RVUE Compilation
$(602 \pm 26) - i(196 \pm 27)$	24 ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
$(537 \pm 20) - i(250 \pm 17)$	25 KAMINSKI	97B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, 4\pi$
$470 - i250$	26,27 TORNQVIST	96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
$387 - i305$	27,28 JANSSEN	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
$420 - i370$	29 ACHASOV	94	RVUE $\pi\pi \rightarrow \pi\pi$
$(506 \pm 10) - i(247 \pm 3)$	KAMINSKI	94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
$370 - i356$	30 ZOU	94B	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

OCCUR=2

OCCUR=2

OCCUR=2

408 - $i342$	27,30 ZOU 93	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
470 - $i208$	31 VANBEVEREN 86	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta,$
(750 \pm 50) - $i(450 \pm 50)$	32 ESTABROOKS 79	RVUE	\dots $\pi\pi \rightarrow \pi\pi, K\bar{K}$
(660 \pm 100) - $i(320 \pm 70)$	PROTOPOP... 73	HBC	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
650 - $i370$	33 BASDEVANT 72	RVUE	$\pi\pi \rightarrow \pi\pi$
1	Data driven analysis using partial-wave dispersion relations .		
2	S-matrix pole; 8595 events.		
3	Applying the chiral unitary approach at NLO to the K_{e4} data of BATLEY 10 and $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.		
4	Uses the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOPESCU 73.		
5	Analytic continuation using Roy equations.		
6	Analytic continuation using GKPY equations.		
7	Using Roy equations.		
8	Average of three variants of the analytic K-matrix model. Uses the K_{e4} data of BATLEY 08A and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73 and GRAYER 74.		
9	Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73, and GRAYER 74, partially of COHEN 80 or ETKIN 82B.		
10	From the K_{e4} data of BATLEY 08A and $\pi N \rightarrow \pi\pi N$ data of HYAMS 73.		
11	From the K_{e4} data of BATLEY 08A and $\pi N \rightarrow \pi\pi N$ data of PROTOPOPESCU 73, GRAYER 74, and ESTABROOKS 74.		
12	From a mean of three different $f_0(500)$ parameterizations. Uses 40k events.		
13	From an isobar model using 2.6k events.		
14	Reanalysis of ABLIKIM 04A, PISLAK 01, and HYAMS 73 data.		
15	Using the N/D method.		
16	From the solution of the Roy equation (ROY 71) for the isoscalar S-wave and using a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.		
17	Reanalysis of the data from PROTOPOPESCU 73, ESTABROOKS 74, GRAYER 74, ROSSELET 77, PISLAK 03, and AKHMETSHIN 04.		
18	From a mean of six different analyses and $f_0(500)$ parameterizations.		
19	Using data on $\psi(2S) \rightarrow J/\psi\pi\pi$ from BAI 00E and on $\Upsilon(nS) \rightarrow \Upsilon(mS)\pi\pi$ from BUTLER 94B and ALEXANDER 98.		
20	From a combined analysis of HYAMS 73, AUGUSTIN 89, AITALA 01B, and PISLAK 01.		
21	A similar analysis (KOMADA 01) finds $(580^{+79}_{-30}) - i(190^{+107}_{-49})$ MeV.		
22	Coupled channel reanalysis of BATON 70, BENSINGER 71, BAILLON 72, HYAMS 73, HYAMS 75, ROSSELET 77, COHEN 80, and ETKIN 82B using the uniformizing variable.		
23	Using the inverse amplitude method and data of ESTABROOKS 73, GRAYER 74, and PROTOPOPESCU 73.		
24	Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.		
25	Average and spread of 4 variants ("up" and "down") of KAMINSKI 97B 3-channel model.		
26	Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.		
27	Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.		
28	Analysis of data from FALVARD 88.		
29	Analysis of data from OCHS 73, ESTABROOKS 75, ROSSELET 77, and MUKHIN 80.		
30	Analysis of data from OCHS 73, GRAYER 74, and ROSSELET 77.		
31	Coupled-channel analysis using data from PROTOPOPESCU 73, HYAMS 73, HYAMS 75, GRAYER 74, ESTABROOKS 74, ESTABROOKS 75, FROGGATT 77, CORN DEN 79, BISWAS 81.		
32	Analysis of data from APEL 72C, GRAYER 74, CASON 76, PAWLICKI 77. Includes spread and errors of 4 solutions.		
33	Analysis of data from BATON 70, BENSINGER 71, COLTON 71, BAILLON 72, PROTOPOPESCU 73, and WALKER 67.		

NODE=M014PP;LINKAGE=P
 NODE=M014PP;LINKAGE=I
 NODE=M014PP;LINKAGE=AD

 NODE=M014PP;LINKAGE=GM

 NODE=M014PP;LINKAGE=GR
 NODE=M014PP;LINKAGE=GC
 NODE=M014PP;LINKAGE=MO
 NODE=M014PP;LINKAGE=ME

 NODE=M014PP;LINKAGE=MN

 NODE=M014PP;LINKAGE=CA
 NODE=M014PP;LINKAGE=AP

 NODE=M014PP;LINKAGE=AL
 NODE=M014PP;LINKAGE=BO
 NODE=M014PP;LINKAGE=BU
 NODE=M014PP;LINKAGE=AN
 NODE=M014PP;LINKAGE=CL

 NODE=M014PP;LINKAGE=ZH

 NODE=M014PP;LINKAGE=AB
 NODE=M014PP;LINKAGE=GA

 NODE=M014PP;LINKAGE=PS
 NODE=M014PP;LINKAGE=KI
 NODE=M014PP;LINKAGE=SU

 NODE=M014PP;LINKAGE=DO

 NODE=M014PP;LINKAGE=AA

 NODE=M014PP;LINKAGE=E
 NODE=M014PP;LINKAGE=B

 NODE=M014PP;LINKAGE=G
 NODE=M014PP;LINKAGE=C
 NODE=M014PP;LINKAGE=D
 NODE=M014PP;LINKAGE=F
 NODE=M014PP;LINKAGE=BV

 NODE=M014PP;LINKAGE=A

 NODE=M014PP;LINKAGE=J

$f_0(500)$ BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

400 to 800 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

513 \pm 32	34 MURAMATSU 02	CLEO	$e^+e^- \approx 10$ GeV
478 $^{+24}_{-23} \pm 17$	AITALA	01B E791	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
563 $^{+58}_{-29}$	35 ISHIDA 01		$\Upsilon(3S) \rightarrow \Upsilon \pi\pi$
555	36 ASNER 00	CLE2	$\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
540 \pm 36	ISHIDA 00B		$p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$
750 \pm 4	ALEKSEEV 99	SPEC	1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^- \pi^+ n$
744 \pm 5	ALEKSEEV 98	SPEC	1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^- \pi^+ n$

NODE=M014M

 NODE=M014M
 → UNCHECKED ←

759± 5	37 TROYAN	98	5.2 $np \rightarrow np\pi^+\pi^-$
780±30	ALDE	97	GAM2 450 $pp \rightarrow pp\pi^0\pi^0$
585±20	38 ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
761±12	39 SVEC	96	RVUE 6-17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 860	40,41 TORNQVIST	96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1165±50	42,43 ANISOVICH	95	RVUE $\pi^-p \rightarrow \pi^0\pi^0n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
414±20	39 AUGUSTIN	89	DM2

34 Statistical uncertainty only.

35 A similar analysis (KOMADA 01) finds 526^{+48}_{-37} MeV.

36 From the best fit of the Dalitz plot.

37 6σ effect, no PWA.

38 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

39 Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+N$ on polarized targets. The fit does not include $f_0(980)$.

40 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

41 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.

42 Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.

43 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

NODE=M014M;LINKAGE=UT

NODE=M014M;LINKAGE=KI

NODE=M014M;LINKAGE=KK

NODE=M014M;LINKAGE=TN

NODE=M014M;LINKAGE=AA

NODE=M014M;LINKAGE=E

NODE=M014M;LINKAGE=B

NODE=M014M;LINKAGE=GG

NODE=M014M;LINKAGE=F

NODE=M014M;LINKAGE=G

$f_0(500)$ BREIT-WIGNER WIDTH

NODE=M014W

VALUE (MeV) DOCUMENT ID TECN COMMENT

100 to 800 OUR ESTIMATE

• • • We do not use the following data for averages, fits, limits, etc. • • •

335± 67	44 MURAMATSU	02	CLEO $e^+e^- \approx 10$ GeV
324 $^{+42}_{-40}$ ±21	AITALA	01B	E791 $D^+ \rightarrow \pi^-\pi^+\pi^+$
372 $^{+229}_{-95}$	45 ISHIDA	01	$\Upsilon(3S) \rightarrow \Upsilon\pi\pi$
540	46 ASNER	00	CLE2 $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
372± 80	ISHIDA	00B	$p\bar{p} \rightarrow \pi^0\pi^0\pi^0$
119± 13	ALEKSEEV	99	SPEC 1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
77± 22	ALEKSEEV	98	SPEC 1.78 $\pi^- p_{\text{polar}} \rightarrow \pi^-\pi^+n$
35± 12	47 TROYAN	98	5.2 $np \rightarrow np\pi^+\pi^-$
780± 60	ALDE	97	GAM2 450 $pp \rightarrow pp\pi^0\pi^0$
385± 70	48 ISHIDA	97	$\pi\pi \rightarrow \pi\pi$
290± 54	49 SVEC	96	RVUE 6-17 $\pi N_{\text{polar}} \rightarrow \pi^+\pi^-N$
~ 880	50,51 TORNQVIST	96	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
460± 40	52,53 ANISOVICH	95	RVUE $\pi^-p \rightarrow \pi^0\pi^0n,$ $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\pi^0\eta, \pi^0\eta\eta$
494± 58	49 AUGUSTIN	89	DM2

44 Statistical uncertainty only.

45 A similar analysis (KOMADA 01) finds 301^{+145}_{-100} MeV.

46 From the best fit of the Dalitz plot.

47 6σ effect, no PWA.

48 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.

49 Breit-Wigner fit to S-wave intensity measured in $\pi N \rightarrow \pi^-\pi^+N$ on polarized targets. The fit does not include $f_0(980)$.

50 Uses data from ASTON 88, OCHS 73, HYAMS 73, ARMSTRONG 91B, GRAYER 74, CASON 83, ROSSELET 77, and BEIER 72B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

51 Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays.

52 Uses $\pi^0\pi^0$ data from ANISOVICH 94, AMSLER 94D, and ALDE 95B, $\pi^+\pi^-$ data from OCHS 73, GRAYER 74 and ROSSELET 77, and $\eta\eta$ data from ANISOVICH 94.

53 The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

NODE=M014W;LINKAGE=UT

NODE=M014W;LINKAGE=KI

NODE=M014W;LINKAGE=KK

NODE=M014W;LINKAGE=TN

NODE=M014W;LINKAGE=AA

NODE=M014W;LINKAGE=E

NODE=M014W;LINKAGE=B

NODE=M014W;LINKAGE=GG

NODE=M014W;LINKAGE=F

NODE=M014W;LINKAGE=G

NODE=M014W

NODE=M014W
→ UNCHECKED ←

$f_0(500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 $\gamma\gamma$	seen

NODE=M014215;NODE=M014

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

 $f_0(500)$ PARTIAL WIDTHS

NODE=M014220

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_2
------------------------	-------------	-------------	------	---------	------------

NODE=M014W2
NODE=M014W2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1.37 ± 13	+0.09 -0.06	54 DANILKIN	21	RVUE	Compilation	
2.05 ± 0.21		55 DAI	14A	RVUE	Compilation	
1.7 ± 0.4		56 HOFERICH... 11		RVUE	Compilation	
3.08 ± 0.82		57 MENNESSIER 11		RVUE	Compilation	
2.08 ± 0.2	+0.07 -0.04	58 MOUSSALLAM11		RVUE	Compilation	
2.08		59 MAO	09	RVUE	Compilation	
1.2 ± 0.4		60 BERNABEU 08		RVUE		
3.9 ± 0.6		57 MENNESSIER 08		RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
1.8 ± 0.4		61 OLLER 08		RVUE	Compilation	
1.68 ± 0.15		61,62 OLLER 08A		RVUE	Compilation	
3.1 ± 0.5		63,64 PENNINGTON 08		RVUE	Compilation	
2.4 ± 0.4		64,65 PENNINGTON 08		RVUE	Compilation	
4.1 ± 0.3		66 PENNINGTON 06		RVUE	$\gamma\gamma \rightarrow \pi^0\pi^0$	
3.8 ± 1.5		67,68 BOGLIONE 99		RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
5.4 ± 2.3		67 MORGAN 90		RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
10 ± 6		COURAU 86		DM1	$e^+e^- \rightarrow \pi^+\pi^-e^+e^-$	

OCCUR=2

54 Using the value of the coupled channel analysis.

55 Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.

56 Using Roy-Steiner equations with $\pi\pi$ phase shifts from an update of COLANGELO 01 and from GARCIA-MARTIN 11A.

57 Using an analytic K-matrix model.

58 Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.

59 Used dispersion theory. The value quoted used the $f_0(500)$ pole position of 457 - i276 MeV.60 Using p , n polarizabilities from PDG 06 and fitting to $\pi\pi$ phase motion from GARCIA-MARTIN 07 and σ -poles from GARCIA-MARTIN 07 and CAPRINI 06.

61 Using twice-subtracted dispersion integrals.

62 Supersedes OLLER 08.

63 Solution A (preferred solution based on χ^2 -analysis).

64 Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

65 Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

66 Using unitarity and the σ pole position from CAPRINI 06.67 This width could equally well be assigned to the $f_0(1370)$. The authors analyse data from BOYER 90 and MARSISKE 90 and report strong correlation with $\gamma\gamma$ width of $f_2(1270)$.

68 Supersedes MORGAN 90.

NODE=M014W2;LINKAGE=C
NODE=M014W2;LINKAGE=B

NODE=M014W2;LINKAGE=HO

NODE=M014W2;LINKAGE=ME
NODE=M014W2;LINKAGE=MO

NODE=M014W2;LINKAGE=MA

NODE=M014W2;LINKAGE=BE

NODE=M014W2;LINKAGE=OL
NODE=M014W2;LINKAGE=LLNODE=M014W2;LINKAGE=P1
NODE=M014W2;LINKAGE=P3

NODE=M014W2;LINKAGE=P2

NODE=M014W2;LINKAGE=PE
NODE=M014W2;LINKAGE=A

NODE=M014W2;LINKAGE=BL

 $f_0(500)$ REFERENCES

DANILKIN 21	PR D103 114023	I. Danilkin, O. Deineka, M. Vanderhaeghen (MAINZ)
SARANTSEV 21	PL B816 136227	A.V. Sarantsev <i>et al.</i> (BONN, PNPI)
ABLIKIM 17	PRL 118 012001	M. Ablikim <i>et al.</i> (BESIII Collab.)
DAI 14A	PR D90 036004	L.-Y. Dai, M.R. Pennington (CEBAF)
ALBALADEJO 12	PR D86 034003	M. Albaladejo, J.A. Oller (MURC)
GARCIA-MAR... 11	PRL 107 072001	R. Garcia-Martin <i>et al.</i> (MADR, CRAC)
GARCIA-MAR... 11A	PR D83 074004	R. Garcia-Martin <i>et al.</i> (MADR, CRAC)
HOFERICH... 11	EPJ C71 1743	M. Hoferichter, D.R. Phillips, C. Schat (BONN+)
MENNESSIER 11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang
MOUSSALLAM 11	EPJ C71 1814	B. Moussallam
BATLEY 10	PL B686 101	J.R. Batley <i>et al.</i> (CERN NA48/2 Collab.)
BATLEY 10C	EPJ C70 635	J.R. Batley <i>et al.</i> (CERN NA48/2 Collab.)
MENNESSIER 10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang
MAO 09	PR D79 116008	Y. Mao <i>et al.</i>
BATLEY 08A	EPJ C54 411	J.R. Batley <i>et al.</i> (CERN NA48/2 Collab.)
BERNABEU 08	PRL 100 241804	J. Bernabeu, J. Prades (IFIC, GRAN)
CAPRINI 08	PR D77 114019	I. Caprini
MENNESSIER 08	PL B665 205	G. Mennessier, S. Narison, W. Ochs
OLLER 08	PL B659 201	J.A. Oller, L. Roca, C. Schat (MURC, UBA)
OLLER 08A	EPJ A37 15	J.A. Oller, L. Roca (MURC)
PENNINGTON 08	EPJ C56 1	M.R. Pennington <i>et al.</i>

NODE=M014

REFID=62088

REFID=61091

REFID=57650

REFID=55923

REFID=54277

REFID=16761

REFID=54121

REFID=53966

REFID=53637

REFID=53975

REFID=53285

REFID=53567

REFID=53657

REFID=52992

REFID=52487

REFID=52271

REFID=52273

REFID=52299

REFID=53976

REFID=53977

REFID=52303

UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
ABLIKIM	07A	PL B645 19	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51614
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=51721
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>		REFID=53252
GARCIA-MAR...	07	PR D76 074034	R. Garcia-Martin, J.R. Pelaez, F.J. Yndurain		REFID=51949
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51652
ANISOVICH	06	IJMP A21 3615	V.V. Anisovich		REFID=51137
CAPRINI	06	PRL 96 132001	I. Caprini, G. Colangelo, H. Leutwyler	(BCIP+)	REFID=51076
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
PENNINGTON	06	PRL 97 011601	M.R. Pennington		REFID=51184
ZHOU	05	JHEP 0502 043	Z.Y. Zhou <i>et al.</i>		REFID=50823
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49740
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
BUETTIKER	04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam		REFID=56428; ERROR=-1
GALLEGOS	04	PR D69 074033	A. Gallegos <i>et al.</i>		REFID=49769
BUGG	03	PL B572 1	D.V. Bugg		REFID=49586
PISLAK	03	PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=49344
Also		PR D81 119903E	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=53337
MURAMATSU	02	PRL 89 251802	H. Muramatsu <i>et al.</i>	(CLEO Collab.)	REFID=49081
Also		PRL 90 059901 (err.)	H. Muramatsu <i>et al.</i>	(CLEO Collab.)	REFID=49385
AITALA	01B	PRL 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
BLACK	01	PR D64 014031	D. Black <i>et al.</i>		REFID=48314
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leutwyler		REFID=49180
ISHIDA	01	PL B518 47	M. Ishida <i>et al.</i>		REFID=48354
KOMADA	01	PL B508 31	T. Komada <i>et al.</i>		REFID=48541
PISLAK	01	PRL 87 221801	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=48433
Also		PR D67 072004	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=49344
Also		PRL 105 019901E	S. Pislak <i>et al.</i>	(BNL E865 Collab.)	REFID=53338
SUROVTSEV	01	PR D63 054024	Y.S. Surovtsev, D. Krupa, M. Nagy		REFID=48310
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
BAI	00E	PR D62 032002	J. Bai <i>et al.</i>	(BES Collab.)	REFID=47955
ISHIDA	00B	PTP 104 203	M. Ishida <i>et al.</i>		REFID=48358
ALEKSEEV	99	NP B541 3	I.G. Alekseev <i>et al.</i>		REFID=46614
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington		REFID=46931
HANNAH	99	PR D60 017502	T. Hannah		REFID=46935
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (err.)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (err.)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
ALEKSEEV	98	PAN 61 174	I.G. Alekseev <i>et al.</i>		REFID=46328
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
LOCHER	98	EPJ C4 317	M.P. Locher <i>et al.</i>	(PSI)	REFID=46372
TROYAN	98	JINRRC 5-91 33	Yu. Troyan <i>et al.</i>		REFID=46615
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
DOBADO	97	PR D56 3057	A. Dobado, J.R. Pelaez		REFID=53964
ISHIDA	97	PTP 98 1005	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45998
KAMINSKI	97B	PL B413 130	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, IPN)	REFID=45778
Also		PTP 95 745	S. Ishida <i>et al.</i>	(TOKY, MIYA, KEK)	REFID=45770
SVEC	96	PR D53 2343	M. Svec	(MCGI)	REFID=44509
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELSE)	REFID=44507
ALDE	95B	ZPHY C66 375	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=44375
ANISOVICH	95	PL B355 363	V.V. Anisovich <i>et al.</i>	(PNPI, SERP)	REFID=44442
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
ACHASOV	94	PR D49 5779	N.N. Achasov, G.N. Shestakov	(NOVM)	REFID=44087
AMSLER	94D	PL B333 277	C. Amstler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43799
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)	REFID=45771
ZOU	94B	PR D50 591	B.S. Zou, D.V. Bugg	(LOQM)	REFID=44072
ZOU	93	PR D48 3948	B.S. Zou, D.V. Bugg	(LOQM)	REFID=43672
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
COURAU	86	NP B271 1	A. Courau <i>et al.</i>	(CLER, LALO)	REFID=44510
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i>	(NIJM, BIEL)	REFID=45769
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BISWAS	81	PRL 47 1378	N.N. Biswas <i>et al.</i>	(NDAM, ANL)	REFID=21106
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL) IJP	REFID=20381
MUKHIN	80	JETPL 32 601	K.N. Mukhin <i>et al.</i>	(KIAE)	REFID=44528
		Translated from ZETFP 32 616.			
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
ESTABROOKS	79	PR D19 2678	P. Estabrooks	(CARL)	REFID=20375
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)	REFID=21072
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJ	REFID=20367
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)	REFID=11004
CASON	76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL) IJ	REFID=21064
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
SRINIVASAN	75	PR D12 681	V. Srinivasan <i>et al.</i>	(NDAM, ANL)	REFID=21062
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
ESTABROOKS	73	Tallahassee	P.G. Estabrooks <i>et al.</i>	(CERN, MPIM)	REFID=20345
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)	REFID=20349
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu <i>et al.</i>	(LBL)	REFID=20108
APEL	72C	PL 41B 542	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=21013
BAILLON	72	PL 38B 555	P.H. Baillon <i>et al.</i>	(SLAC)	REFID=20093
BASDEVANT	72	PL 41B 178	J.L. Basdevant, C.D. Froggatt, J.L. Petersen	(CERN)	REFID=20095
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)	REFID=44530
BENSINGER	71	PL 36B 134	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=21006
COLTON	71	PR D3 2028	E.P. Colton <i>et al.</i>	(LBL, FNAL, UCLA+)	REFID=44533
ROY	71	PL 36B 353	S.M. Roy		REFID=51107
BATON	70	PL 33B 528	J.P. Baton, G. Laurens, J. Reigner	(SACL)	REFID=20086
WALKER	67	RMP 39 695	W.D. Walker	(WISC)	REFID=20960

$\rho(770)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

NODE=M009

 $\rho(770)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(761–765) – i (71–74) OUR ESTIMATE			
$(763.7^{+1.7}_{-1.5}) - i(73.2^{+1.0}_{-1.1})$	¹ GARCIA-MAR..11	RVUE	Compilation
$(754 \pm 18) - i(74 \pm 10)$	² PELAEZ	04A RVUE	$\pi\pi \rightarrow \pi\pi$
$(762.4 \pm 1.8) - i(72.6 \pm 1.4)$	COLANGELO	01 RVUE	$\pi\pi \rightarrow \pi\pi$
¹ Reanalysis of the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOESCU 73 using GKPY equations.			
² Reanalysis of data from PROTOPOESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.			

NODE=M009PP

NODE=M009PP

NODE=M009PP

→ UNCHECKED ←

OCCUR=2

OCCUR=2

NODE=M009PP;LINKAGE=C

NODE=M009PP;LINKAGE=B

 $\rho(770)$ MASSWe no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NODE=M009205

NODE=M009205

NEUTRAL ONLY, e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
775.26 ± 0.23 OUR AVERAGE				
775.3 ± 0.5 ± 0.6		¹ ACHASOV	21	SND $e^+e^- \rightarrow \pi^+\pi^-$
775.02 ± 0.35		² LEES	12G	BABR $e^+e^- \rightarrow \pi^+\pi^-\gamma$
775.97 ± 0.46 ± 0.70	900k	³ AKHMETSHIN	07	$e^+e^- \rightarrow \pi^+\pi^-$
774.6 ± 0.4 ± 0.5	800k	^{4,5} ACHASOV	06	SND $e^+e^- \rightarrow \pi^+\pi^-$
775.65 ± 0.64 ± 0.50	114k	^{6,7} AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow \pi^+\pi^-$
775.9 ± 0.5 ± 0.5	1.98M	⁸ ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.8 ± 0.9 ± 2.0	500k	⁸ ACHASOV	02	SND $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ± 1.1		⁹ BARKOV	85	OLYA $e^+e^- \rightarrow \pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
763.49 ± 0.53		¹⁰ BARTOS	17	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
758.23 ± 0.46		¹¹ BARTOS	17A	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
775.8 ± 0.5 ± 0.3	1.98M	¹² ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.9 ± 0.6 ± 0.5	1.98M	¹³ ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.0 ± 0.6 ± 1.1	500k	¹⁴ ACHASOV	02	SND $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
775.1 ± 0.7 ± 5.3		¹⁵ BENAYOUN	98	RVUE $e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
770.5 ± 1.9 ± 5.1		¹⁶ GARDNER	98	RVUE $0.28-0.92 e^+e^- \rightarrow \pi^+\pi^-$
764.1 ± 0.7		¹⁷ O'CONNELL	97	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
757.5 ± 1.5		¹⁸ BERNICHA	94	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
768 ± 1		¹⁹ GESHKEN...	89	RVUE $e^+e^- \rightarrow \pi^+\pi^-$

NODE=M009M0

NODE=M009M0

OCCUR=2

OCCUR=3

OCCUR=3

NODE=M009M0;LINKAGE=D

NODE=M009M0;LINKAGE=LE

NODE=M009M0;LINKAGE=AK

NODE=M009M0;LINKAGE=AC

NODE=M009M0;LINKAGE=SN

NODE=M009M0;LINKAGE=GS

NODE=M009M0;LINKAGE=PT

NODE=M009M0;LINKAGE=CH

NODE=M009M0;LINKAGE=K

NODE=M009M0;LINKAGE=A

NODE=M009M0;LINKAGE=B

NODE=M009M0;LINKAGE=DF

NODE=M009M0;LINKAGE=WO

¹ From a fit of the cross section in the energy range $0.525 < \sqrt{s} < 0.883$ GeV parameterized by the sum of the Breit-Wigner amplitudes for the $\rho(770)$, ω and $\rho(1450)$ resonances.

² Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

³ A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

⁴ Supersedes ACHASOV 05A.

⁵ A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

⁶ Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.

⁷ Update of AKHMETSHIN 02.

⁸ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁹ From the GOUNARIS 68 parametrization of the pion form factor.

¹⁰ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

¹¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

¹² Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.

¹³ Without limitations on masses and widths.

¹⁴ Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0\pi\pi} = g_{\rho^\pm\pi\pi}$.

¹⁵ Using the data of BARKOV 85 in the hidden local symmetry model.

¹⁶ From the fit to $e^+e^- \rightarrow \pi^+\pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.

¹⁷ A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.

¹⁸ Applying the S-matrix formalism to the BARKOV 85 data.

¹⁹ Includes BARKOV 85 data. Model-dependent width definition.

NODE=M009M0;LINKAGE=HC

NODE=M009M0;LINKAGE=K2

NODE=M009M0;LINKAGE=G8

NODE=M009M0;LINKAGE=AB

NODE=M009M0;LINKAGE=AA

NODE=M009M0;LINKAGE=F

CHARGED ONLY, τ DECAYS and e^+e^-

NODE=M009M5

NODE=M009M5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
775.11±0.34 OUR AVERAGE					
774.6 ±0.2 ±0.5	5.4M	^{1,2} FUJIKAWA	08	BELL ±	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
775.5 ±0.7		^{2,3} SCHAEEL	05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
775.5 ±0.5 ±0.4	1.98M	⁴ ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
775.1 ±1.1 ±0.5	87k	^{5,6} ANDERSON	00A	CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
761.60±0.95		⁷ BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
774.8 ±0.6 ±0.4	1.98M	⁸ ALOISIO	03	KLOE -	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
776.3 ±0.6 ±0.7	1.98M	⁸ ALOISIO	03	KLOE +	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
773.9 ±2.0 ^{+0.3} -1.0		⁹ SANZ-CILLERO	03	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
774.5 ±0.7 ±1.5	500k	⁴ ACHASOV	02	SND ±	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
775.1 ±0.5		¹⁰ PICH	01	RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

OCCUR=2

OCCUR=3

OCCUR=2

¹ $|F_\pi(0)|^2$ fixed to 1.

² From the GOUNARIS 68 parametrization of the pion form factor.

³ The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.

⁴ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁵ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.

⁶ From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.

⁷ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.

⁸ Without limitations on masses and widths.

⁹ Using the data of BARATE 97M and the effective chiral Lagrangian.

¹⁰ From a fit of the model-independent parameterization of the pion form factor to the data of BARATE 97M.

NODE=M009M5;LINKAGE=FU

NODE=M009M5;LINKAGE=GO

NODE=M009M5;LINKAGE=SC

NODE=M009M5;LINKAGE=CH

NODE=M009M5;LINKAGE=A6

NODE=M009M5;LINKAGE=K1

NODE=M009M5;LINKAGE=A

NODE=M009M5;LINKAGE=WO

NODE=M009M5;LINKAGE=Z

NODE=M009M5;LINKAGE=PC

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
763.0±0.3±1.2	600k	¹ ABELE	99E	CBAR 0±	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$

¹ Assuming the equality of ρ^+ and ρ^- masses and widths.

NODE=M009M7

NODE=M009M7

NODE=M009M7;LINKAGE=LB

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
766.5±1.1 OUR AVERAGE					
763.7±3.2		ABELE	97	CBAR	$\bar{p} n \rightarrow \pi^- \pi^0 \pi^0$
768 ±9		AGUILAR...	91	EHS	400 $p p$
767 ±3	2935	¹ CAPRARO	87	SPEC -	200 $\pi^- \text{Cu} \rightarrow \pi^- \pi^0 \text{Cu}$
761 ±5	967	¹ CAPRARO	87	SPEC -	200 $\pi^- \text{Pb} \rightarrow \pi^- \pi^0 \text{Pb}$
771 ±4		HUSTON	86	SPEC +	202 $\pi^+ \text{A} \rightarrow \pi^+ \pi^0 \text{A}$
766 ±7	6500	² BYERLY	73	OSPK -	5 $\pi^- p$
766.8±1.5	9650	³ PISUT	68	RVUE -	1.7-3.2 $\pi^- p$, $t < 10$
767 ±6	900	¹ EISNER	67	HBC -	4.2 $\pi^- p$, $t < 10$

NODE=M009M2

NODE=M009M2

OCCUR=2

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

² Phase shift analysis. Systematic errors added corresponding to spread of different fits.

³ From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

NODE=M009M2;LINKAGE=Z

NODE=M009M2;LINKAGE=X

NODE=M009M2;LINKAGE=A

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
769.2± 0.9 OUR AVERAGE				
770.8± 1.3 ^{+2.3} _{-2.4}	900k	ANDREEV 20	H1	$ep \rightarrow e\pi^+\pi^-p$
771 ± 2 ⁺² ₋₁	63.5k	¹ ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^-p$
770 ± 2 ± 1	79k	² BREITWEG 98B	ZEUS	50–100 γp
767.6± 2.7		BARTALUCCI 78	CNTR	$\gamma p \rightarrow e^+e^-p$
775 ± 5		GLADDING 73	CNTR	2.9–4.7 γp
767 ± 4	1930	BALLAM 72	HBC	2.8 γp
770 ± 4	2430	BALLAM 72	HBC	4.7 γp
765 ± 10		ALVENSLEB... 70	CNTR	$\gamma A, t < 0.01$
767.7± 1.9	140k	BIGGS 70	CNTR	$< 4.1 \gamma C \rightarrow \pi^+\pi^-C$
765 ± 5	4000	ASBURY 67B	CNTR	$\gamma + Pb$

• • • We do not use the following data for averages, fits, limits, etc. • • •

771 ± 2	79k	³ BREITWEG 98B	ZEUS	50–100 γp
---------	-----	---------------------------	------	-------------------

¹ Using the KUHN 90 parametrization of the pion form factor, neglecting ρ - ω interference.
² From the parametrization according to SOEDING 66.
³ From the parametrization according to ROSS 66.

NODE=M009M0P
 NODE=M009M0P

OCCUR=2

OCCUR=2

NODE=M009M0P;LINKAGE=AB
 NODE=M009M0P;LINKAGE=B5
 NODE=M009M0P;LINKAGE=B6

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
769.0 ± 0.9 OUR AVERAGE				
Error includes scale factor of 1.4. See the ideogram below.				
765 ± 6		BERTIN 97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
773 ± 1.6		WEIDENAUER 93	ASTE	$\bar{p}p \rightarrow \pi^+\pi^-\omega$
762.6 ± 2.6		AGUILAR-... 91	EHS	400 pp
770 ± 2		¹ HEYN 81	RVUE	Pion form factor
768 ± 4		^{2,3} BOHACIK 80	RVUE	
769 ± 3		⁴ WICKLUND 78	ASPK	3,4,6 $\pi^\pm N$
768 ± 1	76k	DEUTSCH... 76	HBC	16 π^+p
767 ± 4	4100	ENGLER 74	DBC	$6 \pi^+n \rightarrow \pi^+\pi^-p$
775 ± 4	32k	² PROTOPOP... 73	HBC	$7.1 \pi^+p, t < 0.4$
764 ± 3	6.8k	⁵ RATCLIFF 72	ASPK	$15 \pi^-p, t < 0.3$
774 ± 3	1.7k	REYNOLDS 69	HBC	$2.26 \pi^-p$
769.2 ± 1.5	13.3k	⁶ PISUT 68	RVUE	$1.7\text{--}3.2 \pi^-p, t < 10$

• • • We do not use the following data for averages, fits, limits, etc. • • •

774.34±0.18±0.35	970k	⁷ ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
772.93±0.18±0.34	970k	⁸ ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
773.5 ± 2.5		⁹ COLANGELO 01	RVUE	$\pi\pi \rightarrow \pi\pi$
762.3 ± 0.5 ± 1.2	600k	¹⁰ ABELE 99E	CBAR	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
777 ± 2	4.9k	¹¹ ADAMS 97	E665	$470 \mu p \rightarrow \mu XB$
770 ± 2		¹² BOGOLYUB... 97	MIRA	$32 \bar{p}p \rightarrow \pi^+\pi^-X$
768 ± 8		¹² BOGOLYUB... 97	MIRA	$32 pp \rightarrow \pi^+\pi^-X$
761.1 ± 2.9		DUBNICKA 89	RVUE	π form factor
777.4 ± 2.0		¹³ CHABAUD 83	ASPK	$17 \pi^-p$ polarized
769.5 ± 0.7		^{2,3} LANG 79	RVUE	
770 ± 9		³ ESTABROOKS 74	RVUE	$17 \pi^-p \rightarrow \pi^+\pi^-n$
773.5 ± 1.7	11.2k	¹⁴ JACOBS 72	HBC	$2.8 \pi^-p$
775 ± 3	2.2k	¹⁵ HYAMS 68	OSPK	$11.2 \pi^-p$

¹ HEYN 81 includes all spacelike and timelike F_π values until 1978.
² From pole extrapolation.
³ From phase shift analysis of GRAYER 74 data.
⁴ Phase shift analysis. Systematic errors added corresponding to spread of different fits.
⁵ Published values contain misprints. Corrected by private communication RATCLIFF 74.
⁶ Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDBERGER 64, ABOLINS 63.
⁷ From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.
⁸ From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.
⁹ Breit-Wigner mass from a phase-shift analysis of HYAMS 73 and PROTOPOPESCU 73 data.
¹⁰ Using relativistic Breit-Wigner and taking into account ρ - ω interference.
¹¹ Systematic errors not evaluated.
¹² Systematic effects not studied.
¹³ From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P-wave intensity. CHABAUD 83 includes data of GRAYER 74.

NODE=M009M0R
 NODE=M009M0R

OCCUR=2

OCCUR=2

NODE=M009M0R;LINKAGE=B0
 NODE=M009M0R;LINKAGE=C0
 NODE=M009M0R;LINKAGE=H
 NODE=M009M0R;LINKAGE=X
 NODE=M009M0R;LINKAGE=03
 NODE=M009M0R;LINKAGE=R

NODE=M009M0R;LINKAGE=B

NODE=M009M0R;LINKAGE=C


NODE=M009M0R;LINKAGE=CL

NODE=M009M0R;LINKAGE=BL
 NODE=M009M0R;LINKAGE=A1
 NODE=M009M0R;LINKAGE=QQ
 NODE=M009M0R;LINKAGE=G

¹⁴ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

¹⁵ Of HYAMS 68 six parametrizations, this is theoretically soundest. MR

NODE=M009M0R;LINKAGE=Z
 NODE=M009M0R;LINKAGE=02



$m_{\rho(770)^0} - m_{\rho(770)^\pm}$

NODE=M009D

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-0.7 ± 0.8	OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
-2.4 ± 0.8	1	SCHAEL 05C	ALEP		$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
0.4 ± 0.7 ± 0.6	1.98M	2	ALOISIO 03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.3 ± 1.1 ± 2.0	500k	2	ACHASOV 02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
1.6 ± 0.6 ± 1.7	600k		ABELE 99E	CBAR ±0	$0.0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
-4 ± 4	3000	3	REYNOLDS 69	HBC -0	$2.26 \pi^- \rho$
-5 ± 5	3600	3	FOSTER 68	HBC ±0	$0.0 \bar{p} p$
2.4 ± 2.1	22950	4	PISUT 68	RVUE	$\pi N \rightarrow \rho N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
-3.37 ± 1.06		5	BARTOS 17A	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$, $\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$

NODE=M009D

¹ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEL 05c and $e^+ e^-$ data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

NODE=M009D;LINKAGE=SC

² Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

NODE=M009D;LINKAGE=CH

³ From quoted masses of charged and neutral modes.


NODE=M009D;LINKAGE=A

⁴ Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDHABER 64, ABOLINS 63.

NODE=M009D;LINKAGE=R

⁵ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M009D;LINKAGE=B



$m_{\rho(770)^+} - m_{\rho(770)^-}$

NODE=M009D1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M009D1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.5 \pm 0.8 \pm 0.7$	1.98M	¹ ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
-----------------------	-------	----------------------	----	------	--

NODE=M009D;LINKAGE=WO

¹Without limitations on masses and widths.

$\rho(770)$ RANGE PARAMETER

NODE=M009R

The range parameter R enters an energy-dependent correction to the width, of the form $(1 + q_r^2 R^2) / (1 + q^2 R^2)$, where q is the momentum of one of the pions in the $\pi\pi$ rest system. At resonance, $q = q_r$.

NODE=M009R

VALUE (GeV^{-1})	DOCUMENT ID	TECN	CHG	COMMENT
-----------------------------	-------------	------	-----	---------

NODE=M009R

$5.3^{+0.9}_{-0.7}$	¹ CHABAUD	83	ASPK	0	$17 \pi^- p$ polarized
---------------------	----------------------	----	------	---	------------------------

¹The old PISUT 68 value, properly corrected, was 3.2 ± 0.6 .

NODE=M009R;LINKAGE=01

$\rho(770)$ WIDTH

NODE=M009220

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

NODE=M009220

NEUTRAL ONLY, $e^+ e^-$

NODE=M009W0

NODE=M009W0

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------


147.4 ± 0.8 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.

$145.6 \pm 0.6 \pm 0.8$		¹ ACHASOV	21	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
149.59 ± 0.67		² LEES	12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
$145.98 \pm 0.75 \pm 0.50$	900k	³ AKHMETSHIN	07		$e^+ e^- \rightarrow \pi^+ \pi^-$
$146.1 \pm 0.8 \pm 1.5$	800k	^{4,5} ACHASOV	06	SND	$e^+ e^- \rightarrow \pi^+ \pi^-$
$143.85 \pm 1.33 \pm 0.80$	114k	^{6,7} AKHMETSHIN	04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^-$
$147.3 \pm 1.5 \pm 0.7$	1.98M	⁸ ALOISIO	03	KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
$151.1 \pm 2.6 \pm 3.0$	500k	⁸ ACHASOV	02	SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.5 ± 3.0		⁹ BARKOV	85	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

144.06±0.85		10	BARTOS	17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
144.56±0.80		11	BARTOS	17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
143.9 ±1.3 ±1.1	1.98M	12	ALOISIO	03	KLOE	$1.02 \frac{e^+e^- \rightarrow \pi^+\pi^-}{\pi^+\pi^-\pi^0}$	OCCUR=2
147.4 ±1.5 ±0.7	1.98M	13	ALOISIO	03	KLOE	$1.02 \frac{e^+e^- \rightarrow \pi^+\pi^-}{\pi^+\pi^-\pi^0}$	OCCUR=3
149.8 ±2.2 ±2.0	500k	14	ACHASOV	02	SND	$1.02 \frac{e^+e^- \rightarrow \pi^+\pi^-}{\pi^+\pi^-\pi^0}$	OCCUR=3
147.9 ±1.5 ±7.5		15	BENAYOUN	98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$	
153.5 ±1.3 ±4.6		16	GARDNER	98	RVUE	$0.28-0.92 \frac{e^+e^- \rightarrow \pi^+\pi^-}{\pi^+\pi^-}$	
145.0 ±1.7		17	O'CONNELL	97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
142.5 ±3.5		18	BERNICHIA	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
138 ±1		19	GESHKEN...	89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	

- 1 From a fit of the cross section in the energy range $0.525 < \sqrt{s} < 0.883$ GeV parameterized by the sum of the Breit-Wigner amplitudes for the $\rho(770)$, ω and $\rho(1450)$ resonances. NODE=M009W0;LINKAGE=D
- 2 Using the GOUNARIS 68 parametrization with the complex phase of the $\rho-\omega$ interference and leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit. NODE=M009W0;LINKAGE=LE
- 3 A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05. NODE=M009W;LINKAGE=AK
- 4 Supersedes ACHASOV 05A. NODE=M009W0;LINKAGE=AC
NODE=M009W0;LINKAGE=SN
- 5 A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A. NODE=M009W5;LINKAGE=GS
NODE=M009W5;LINKAGE=P2
- 6 Using the GOUNARIS 68 parametrization with the complex phase of the $\rho-\omega$ interference. NODE=M009W;LINKAGE=CH
NODE=M009W;LINKAGE=K
- 7 From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02. NODE=M009W0;LINKAGE=A
NODE=M009W0;LINKAGE=B
- 8 Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$. NODE=M009W;LINKAGE=DF
NODE=M009W;LINKAGE=WO
- 9 From the GOUNARIS 68 parametrization of the pion form factor. NODE=M009W;LINKAGE=HC
NODE=M009W;LINKAGE=K2
- 10 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C. NODE=M009W;LINKAGE=G8
- 11 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A. NODE=M009W;LINKAGE=AB
NODE=M009W;LINKAGE=AA
NODE=M009W;LINKAGE=F
- 12 Assuming $m_{\rho^+} = m_{\rho^-} = m_{\rho^0}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-} = \Gamma_{\rho^0}$.
- 13 Without limitations on masses and widths.
- 14 Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0\pi\pi} = g_{\rho^\pm\pi\pi}$.
- 15 Using the data of BARKOV 85 in the hidden local symmetry model.
- 16 From the fit to $e^+e^- \rightarrow \pi^+\pi^-$ data from the compilations of HEYN 81 and BARKOV 85, including the GOUNARIS 68 parametrization of the pion form factor.
- 17 A fit of BARKOV 85 data assuming the direct $\omega\pi\pi$ coupling.
- 18 Applying the S-matrix formalism to the BARKOV 85 data.
- 19 Includes BARKOV 85 data. Model-dependent width definition.



CHARGED ONLY, τ DECAYS and e^+e^-

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
149.1 ± 0.8 OUR FIT					
149.1 ± 0.8 OUR AVERAGE					
148.1 ± 0.4	± 1.7	5.4M	1,2 FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
149.0 ± 1.2			2,3 SCHAEL	05c ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
149.9 ± 2.3	± 2.0	500k	4 ACHASOV	02 SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.4 ± 1.4	± 1.4	87k	5,6 ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
139.90 ± 0.46			7 BARTOS	17A RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
143.7 ± 1.3	± 1.2	1.98M	4 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
142.9 ± 1.3	± 1.4	1.98M	8 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
144.7 ± 1.4	± 1.2	1.98M	8 ALOISIO	03 KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
150.2 ± 2.0	$+0.7$ -1.6		9 SANZ-CILLERO	03 RVUE	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
150.9 ± 2.2	± 2.0	500k	10 ACHASOV	02 SND	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M009W5
 NODE=M009W5

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

¹ $|F_\pi(0)|^2$ fixed to 1.

² From the GOUNARIS 68 parametrization of the pion form factor.

³ The error combines statistical and systematic uncertainties. Supersedes BARATE 97M.

⁴ Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

⁵ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.

⁶ From the GOUNARIS 68 parametrization of the pion form factor. The second error is a model error taking into account different parametrizations of the pion form factor.

⁷ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.

⁸ Without limitations on masses and widths.

⁹ Using the data of BARATE 97M and the effective chiral Lagrangian.

¹⁰ Assuming $m_{\rho^0} = m_{\rho^\pm}$, $g_{\rho^0 \pi\pi} = g_{\rho^\pm \pi\pi}$.

NODE=M009W5;LINKAGE=FU
 NODE=M009W5;LINKAGE=GO
 NODE=M009W5;LINKAGE=SC

NODE=M009W5;LINKAGE=CH
 NODE=M009W5;LINKAGE=A6
 NODE=M009W5;LINKAGE=K1

NODE=M009W5;LINKAGE=A

NODE=M009W5;LINKAGE=W0
 NODE=M009W5;LINKAGE=Z

NODE=M009W5;LINKAGE=HC

MIXED CHARGES, OTHER REACTIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
149.5 ± 1.3	600k	1 ABELE	99E CBAR	0 \pm	0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$

¹ Assuming the equality of ρ^+ and ρ^- masses and widths.

NODE=M009W7
 NODE=M009W7

NODE=M009W;LINKAGE=LB

CHARGED ONLY, HADROPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150.2 ± 2.4 OUR FIT					
150.2 ± 2.4 OUR AVERAGE					
152.8 ± 4.3			ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^- \pi^0 \pi^0$
155 ± 11	2.9k	1 CAPRARO	87 SPEC	-	200 $\pi^- \text{Cu} \rightarrow \pi^- \pi^0 \text{Cu}$
154 ± 20	967	1 CAPRARO	87 SPEC	-	200 $\pi^- \text{Pb} \rightarrow \pi^- \pi^0 \text{Pb}$
150 ± 5		HUSTON	86 SPEC	+	202 $\pi^+ \text{A} \rightarrow \pi^+ \pi^0 \text{A}$
146 ± 12	6.5k	2 BYERLY	73 OSPK	-	5 $\pi^- p$
148.2 ± 4.1	9.6k	3 PISUT	68 RVUE	-	1.7-3.2 $\pi^- p$, $t < 10$
146 ± 13	900	EISNER	67 HBC	-	4.2 $\pi^- p$, $t < 10$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

137.0 ± 0.4 ⁴ ABLIKIM 17 BES3 $J/\psi \rightarrow \gamma 3\pi$

¹ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

² Phase shift analysis. Systematic errors added corresponding to spread of different fits.

³ From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

⁴ S-matrix pole at a fixed ρ meson mass of 775.49 MeV.

NODE=M009W2
 NODE=M009W2

OCCUR=2

NODE=M009W2;LINKAGE=Z
 NODE=M009W2;LINKAGE=X
 NODE=M009W2;LINKAGE=A1

NODE=M009W2;LINKAGE=A

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
151.5 $^{+1.9}_{-2.1}$ OUR AVERAGE				
151.3 ± 2.2	$^{+1.6}_{-2.8}$	900k	ANDREEV 20	H1 $ep \rightarrow e\pi^+\pi^-p$
155 ± 5	± 2	63.5k	1 ABRAMOWICZ	12 ZEUS $ep \rightarrow e\pi^+\pi^-p$
146 ± 3	± 13	79k	2 BREITWEG	98B ZEUS 50-100 γp
150.9 ± 3.0			BARTALUCCI 78	CNTR $\gamma p \rightarrow e^+e^-p$

NODE=M009W0P
 NODE=M009W0P

• • • We do not use the following data for averages, fits, limits, etc. • • •

138 ± 3	79k	³ BREITWEG	98B	ZEUS	50–100 γp	OCCUR=2
147 ± 11		GLADDING	73	CNTR	2.9–4.7 γp	
155 ± 12	2430	BALLAM	72	HBC	4.7 γp	
145 ± 13	1930	BALLAM	72	HBC	2.8 γp	OCCUR=2
140 ± 5		ALVENSLEB...	70	CNTR	γA , $t < 0.01$	
146.1 ± 2.9	140k	BIGGS	70	CNTR	$< 4.1 \gamma C \rightarrow \pi^+ \pi^- C$	
160 ± 10		LANZEROTTI	68	CNTR	γp	
130 ± 5	4000	ASBURY	67B	CNTR	$\gamma + Pb$	

¹ Using the KUHN 90 parametrization of the pion form factor, neglecting ρ - ω interference.

² From the parametrization according to SOEDING 66.

³ From the parametrization according to ROSS 66.

NODE=M009W0P;LINKAGE=AB

NODE=M009W;LINKAGE=B5

NODE=M009W;LINKAGE=B6

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

150.9 ± 1.7 OUR AVERAGE Error includes scale factor of 1.1.

122 ± 20		BERTIN	97C	OBLX	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
145.7 ± 5.3		WEIDENAUER	93	ASTE	$\bar{p} p \rightarrow \pi^+ \pi^- \omega$
144.9 ± 3.7		DUBNICKA	89	RVUE	π form factor
148 ± 6		^{1,2} BOHACIK	80	RVUE	
152 ± 9		³ WICKLUND	78	ASPK	3,4,6 $\pi^\pm p N$
154 ± 2	76k	DEUTSCH...	76	HBC	16 $\pi^+ p$
157 ± 8	6.8k	⁴ RATCLIFF	72	ASPK	15 $\pi^- p$, $t < 0.3$
143 ± 8	1.7k	REYNOLDS	69	HBC	2.26 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

150.85 ± 0.55 ± 0.67	970k	⁵ ABLIKIM	18C	BES3	$\eta'(958) \rightarrow \gamma \pi^+ \pi^-$	
150.18 ± 0.55 ± 0.65	970k	⁶ ABLIKIM	18C	BES3	$\eta'(958) \rightarrow \gamma \pi^+ \pi^-$	OCCUR=2
147.0 ± 2.5	600k	⁷ ABELE	99E	CBAR	0.0 $\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$	
146 ± 3	4.9k	⁸ ADAMS	97	E665	470 $\mu p \rightarrow \mu X B$	
160.0 + 4.1 - 4.0		⁹ CHABAUD	83	ASPK	17 $\pi^- p$ polarized	
155 ± 1		¹⁰ HEYN	81	RVUE	π form factor	
148.0 ± 1.3		^{1,2} LANG	79	RVUE		
146 ± 14	4.1k	ENGLER	74	DBC	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$	
143 ± 13		² ESTABROOKS	74	RVUE	17 $\pi^- p \rightarrow \pi^+ \pi^- n$	
160 ± 10	32k	¹ PROTOPOP...	73	HBC	7.1 $\pi^+ p$, $t < 0.4$	
145 ± 12	2.2k	^{3,11} HYAMS	68	OSPK	11.2 $\pi^- p$	
163 ± 15	13.3k	¹² PISUT	68	RVUE	1.7–3.2 $\pi^- p$, $t < 10$	

¹ From pole extrapolation.

² From phase shift analysis of GRAYER 74 data.

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁴ Published values contain misprints. Corrected by private communication RATCLIFF 74.

⁵ From a fit to $\pi^+ \pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.

⁶ From a fit to $\pi^+ \pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.

⁷ Using relativistic Breit-Wigner and taking into account ρ - ω interference.

⁸ Systematic errors not evaluated.

⁹ From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P -wave intensity. CHABAUD 83 includes data of GRAYER 74.

¹⁰ HEYN 81 includes all spacelike and timelike F_π values until 1978.

¹¹ Of HYAMS 68 six parametrizations this is theoretically soundest. MR

¹² Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, GOLDHABER 64, ABOLINS 63.

NODE=M009W;LINKAGE=C

NODE=M009W;LINKAGE=H

NODE=M009W;LINKAGE=Z

NODE=M009W;LINKAGE=03

NODE=M009W0R;LINKAGE=B

NODE=M009W0R;LINKAGE=C

NODE=M009W;LINKAGE=BL

NODE=M009W;LINKAGE=A1

NODE=M009W;LINKAGE=G

NODE=M009W;LINKAGE=B

NODE=M009W;LINKAGE=02

NODE=M009W;LINKAGE=R

$\Gamma_{\rho(770)^0} - \Gamma_{\rho(770)^\pm}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.3 ± 1.3 OUR AVERAGE Error includes scale factor of 1.4.

-0.2 ± 1.0		¹ SCHAEEL	05C	ALEP	$\tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
3.6 ± 1.8 ± 1.7	1.98M	² ALOISIO	03	KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.66 ± 0.85		³ BARTOS	17A	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-, \tau^- \rightarrow \pi^- \pi^0 \nu_\tau$
-------------	--	---------------------	-----	------	--

NODE=M009W6

NODE=M009W6

¹ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEEL 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. Supersedes BARATE 97M.

² Assuming $m_{\rho^+} = m_{\rho^-}$, $\Gamma_{\rho^+} = \Gamma_{\rho^-}$.

³ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

$$\Gamma_{\rho(770)^+} = \Gamma_{\rho(770)^-}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.8 \pm 2.0 \pm 0.5$	1.98M	¹ ALOISIO	03	KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ Without limitations on masses and widths.

NODE=M009W6;LINKAGE=SC

NODE=M009W6;LINKAGE=CH
NODE=M009W6;LINKAGE=A

NODE=M009W16
NODE=M009W16

NODE=M009W16;LINKAGE=WO

NODE=M009225;NODE=M009

$\rho(770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\pi\pi$	~ 100	%
Γ_2 $K\bar{K}$		
$\rho(770)^\pm$ decays		
Γ_3 $\pi^\pm\pi^0$	~ 100	%
Γ_4 $\pi^\pm\gamma$	$(4.5 \pm 0.5) \times 10^{-4}$	S=2.2
Γ_5 $\pi^\pm\eta$	< 6	$\times 10^{-3}$ CL=84%
Γ_6 $\pi^\pm\pi^+\pi^-\pi^0$	< 2.0	$\times 10^{-3}$ CL=84%
$\rho(770)^0$ decays		
Γ_7 $\pi^+\pi^-$	~ 100	%
Γ_8 $\pi^+\pi^-\gamma$	$(9.9 \pm 1.6) \times 10^{-3}$	
Γ_9 $\pi^0\gamma$	$(4.7 \pm 0.8) \times 10^{-4}$	S=1.7
Γ_{10} $\eta\gamma$	$(3.00 \pm 0.21) \times 10^{-4}$	
Γ_{11} $\pi^0\pi^0\gamma$	$(4.5 \pm 0.8) \times 10^{-5}$	
Γ_{12} $\mu^+\mu^-$	[a] $(4.55 \pm 0.28) \times 10^{-5}$	
Γ_{13} e^+e^-	[a] $(4.72 \pm 0.05) \times 10^{-5}$	
Γ_{14} $\pi^+\pi^-\pi^0$	$(1.01^{+0.54}_{-0.36} \pm 0.34) \times 10^{-4}$	
Γ_{15} $\pi^+\pi^-\pi^+\pi^-$	$(1.8 \pm 0.9) \times 10^{-5}$	
Γ_{16} $\pi^+\pi^-\pi^0\pi^0$	$(1.6 \pm 0.8) \times 10^{-5}$	
Γ_{17} $\pi^0e^+e^-$	< 1.2	$\times 10^{-5}$ CL=90%
Γ_{18} ηe^+e^-		

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=81

NODE=M009;CLUMP=A
DESIG=11;OUR EVAL;→ UNCHECKED ←
DESIG=3
DESIG=5
DESIG=21

NODE=M009;CLUMP=B
DESIG=12;OUR EVAL;→ UNCHECKED ←
DESIG=60
DESIG=40
DESIG=8
DESIG=80
DESIG=6
DESIG=4
DESIG=7;OUR EVAL;→ UNCHECKED ←
DESIG=22
DESIG=30
DESIG=9
DESIG=10

[a] The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+\mu^-) = \Gamma(\rho^0 \rightarrow e^+e^-) \times 0.99785$.

LINKAGE=MD2

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 10 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.7$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4		-100	
Γ		15	-15
		x_3	x_4

Mode	Rate (MeV)	Scale factor
Γ_3 $\pi^\pm\pi^0$	150.2 ± 2.4	
Γ_4 $\pi^\pm\gamma$	0.068 ± 0.007	2.3

DESIG=11
DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 7 branching ratios uses 21 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 9.5$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_8	-100								
x_9	-5	0							
x_{10}	-1	0	1						
x_{11}	-1	0	0	0					
x_{12}	2	-3	0	0	0				
x_{13}	0	0	-6	-9	0	0			
x_{15}	-1	0	0	0	0	0	0		
Γ	0	0	3	5	0	0	-54	0	
	x_7	x_8	x_9	x_{10}	x_{11}	x_{12}	x_{13}	x_{15}	

Mode	Rate (MeV)	Scale factor
Γ_7 $\pi^+ \pi^-$	147.5 ± 0.9	
Γ_8 $\pi^+ \pi^- \gamma$	1.48 ± 0.24	
Γ_9 $\pi^0 \gamma$	0.070 ± 0.012	1.7
Γ_{10} $\eta \gamma$	0.0447 ± 0.0032	
Γ_{11} $\pi^0 \pi^0 \gamma$	0.0066 ± 0.0012	
Γ_{12} $\mu^+ \mu^-$	[a] 0.0068 ± 0.0004	
Γ_{13} $e^+ e^-$	[a] 0.00704 ± 0.00006	
Γ_{15} $\pi^+ \pi^- \pi^+ \pi^-$	0.0027 ± 0.0014	

DESIG=12
DESIG=60
DESIG=40
DESIG=8
DESIG=80
DESIG=6
DESIG=4
DESIG=22

$\rho(770)$ PARTIAL WIDTHS

NODE=M009230


$\Gamma(\pi^\pm \gamma)$

Γ_4

NODE=M009W3
NODE=M009W3

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
68 ± 7 OUR FIT	Error includes scale factor of 2.3.			
68 ± 7 OUR AVERAGE	Error includes scale factor of 2.2. See the ideogram below.			
81 ± 4 ± 4	CAPRARO	87	SPEC -	200 $\pi^- A \rightarrow \pi^- \pi^0 A$
59.8 ± 4.0	HUSTON	86	SPEC +	202 $\pi^+ A \rightarrow \pi^+ \pi^0 A$
71 ± 7	JENSEN	83	SPEC -	156-260 $\pi^- A \rightarrow \pi^- \pi^0 A$

WEIGHTED AVERAGE
68±7 (Error scaled by 2.2)



$\Gamma(\pi^0\gamma)$ Γ_9

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$77 \pm 17 \pm 11$	36500	¹ ACHASOV 03	SND	$0.60-0.97 e^+e^- \rightarrow \pi^0\gamma$
121 ± 31		DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

¹ Using $\Gamma_{\text{total}} = 147.9 \pm 1.3$ MeV and $B(\rho \rightarrow \pi^0\gamma)$ from ACHASOV 03.

NODE=M009W31
NODE=M009W31

NODE=M009W31;LINKAGE=AV

 $\Gamma(\eta\gamma)$ Γ_{10}

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

62 ± 17	¹ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$
-------------	--------------------------	----	---------------------------------

¹ Solution corresponding to constructive ω - ρ interference.

NODE=M009W32
NODE=M009W32

NODE=M009W32;LINKAGE=L

 $\Gamma(e^+e^-)$ Γ_{13}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

7.04 ± 0.06 OUR FIT

7.04 ± 0.06 OUR AVERAGE

$7.048 \pm 0.057 \pm 0.050$	900k	¹ AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
$7.06 \pm 0.11 \pm 0.05$	114k	^{2,3} AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
$6.77 \pm 0.10 \pm 0.30$		BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.12 \pm 0.02 \pm 0.11$	800k	⁴ ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$
6.3 ± 0.1		⁵ BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$

¹ A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

² Using the GOUNARIS 68 parametrization with the complex phase of the ρ - ω interference.

³ From a fit in the energy range 0.61 to 0.96 GeV. Update of AKHMETSHIN 02.

⁴ Supersedes ACHASOV 05A.

⁵ Using the data of BARKOV 85 in the hidden local symmetry model.

NODE=M009W4
NODE=M009W4

OCCUR=2

NODE=M009W4;LINKAGE=AK
NODE=M009W4;LINKAGE=GS
NODE=M009W4;LINKAGE=P2
NODE=M009W4;LINKAGE=AC
NODE=M009W4;LINKAGE=K2

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)$ Γ_{15}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.8 \pm 1.4 \pm 0.5$	153	AKHMETSHIN 00	CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
-----------------------	-----	---------------	------	--

NODE=M009W33
NODE=M009W33

 $\rho(770) \Gamma(e^+e^-)\Gamma(i)/\Gamma^2(\text{total})$

NODE=M009233

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ $\Gamma_{13}/\Gamma \times \Gamma_7/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.89 ± 0.04 OUR AVERAGE

$4.889 \pm 0.015 \pm 0.039$		¹ ACHASOV 21	SND	$e^+e^- \rightarrow \pi^+\pi^-$
$4.876 \pm 0.023 \pm 0.064$	800k	^{2,3} ACHASOV 06	SND	$e^+e^- \rightarrow \pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.72 ± 0.02		⁴ BENAYOUN 10	RVUE	$0.4-1.05 e^+e^-$
-----------------	--	--------------------------	------	-------------------

¹ From a fit of the cross section in the energy range $0.525 < \sqrt{s} < 0.883$ GeV parameterized by the sum of the Breit-Wigner amplitudes for the $\rho(770)$, ω and $\rho(1450)$ resonances.

² Supersedes ACHASOV 05A.

³ A fit of the SND data from 400 to 1000 MeV using parameters of the $\rho(1450)$ and $\rho(1700)$ from a fit of the data of BARKOV 85, BISELLO 89 and ANDERSON 00A.

⁴ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M009G4;LINKAGE=A

NODE=M009G4;LINKAGE=AC
NODE=M009G4;LINKAGE=SN

NODE=M009G4;LINKAGE=BE

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$ $\Gamma_{13}/\Gamma \times \Gamma_{10}/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.42 ± 0.10 OUR FIT

1.45 ± 0.12 OUR AVERAGE

$1.32 \pm 0.14 \pm 0.08$	33k	¹ ACHASOV 07B	SND	$0.6-1.38 e^+e^- \rightarrow \eta\gamma$
$1.50 \pm 0.65 \pm 0.09$	17.4k	² AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$
$1.61 \pm 0.20 \pm 0.11$	23k	^{3,4} AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.85 ± 0.49		⁵ DOLINSKY 89	ND	$e^+e^- \rightarrow \eta\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.05 ± 0.02		⁶ BENAYOUN 10	RVUE	$0.4-1.05 e^+e^-$
-----------------	--	--------------------------	------	-------------------

NODE=M009G1
NODE=M009G1

- ¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- ² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
- ³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
- ⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
- ⁵ Recalculated by us from the cross section in the peak.
- ⁶ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M009G1;LINKAGE=AH

NODE=M009G;LINKAGE=AH
 NODE=M009G;LINKAGE=AK
 NODE=M009G;LINKAGE=BQ

NODE=M009G;LINKAGE=LP
 NODE=M009G1;LINKAGE=BE

$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\pi^0\gamma)/\Gamma_{total}$ $\Gamma_{13}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-8}) EVTS DOCUMENT ID TECN COMMENT

2.2 ± 0.4 OUR FIT Error includes scale factor of 1.7.

2.21 ± 0.34 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

1.98 ± 0.22 ± 0.10 ¹ ACHASOV 16A SND 0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$

2.90 $^{+0.60}_{-0.55}$ ± 0.18 18k AKHMETSHIN 05 CMD2 0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$

3.61 ± 0.74 ± 0.49 10k ² DOLINSKY 89 ND $e^+e^- \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.875 ± 0.026 ³ BENAYOUN 10 RVUE 0.4-1.05 e^+e^-

2.37 ± 0.53 ± 0.33 36k ⁴ ACHASOV 03 SND 0.60-0.97 $e^+e^- \rightarrow \pi^0\gamma$


- ¹ From the VMD model with the $\rho(770)$, $\omega(782)$, $\phi(1020)$ resonances, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 03.
- ² Recalculated by us from the cross section in the peak.
- ³ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.
- ⁴ Using $\sigma_{\phi \rightarrow \pi^0\gamma}$ from ACHASOV 00 and $m_\rho = 775.97$ MeV in the model with the energy-independent phase of ρ - ω interference equal to $(-10.2 \pm 7.0)^\circ$.

NODE=M009G2;LINKAGE=B

NODE=M009G2;LINKAGE=LP
 NODE=M009G2;LINKAGE=BE

NODE=M009G;LINKAGE=SH

WEIGHTED AVERAGE
 2.21 ± 0.34 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(e^+e^-)/\Gamma_{total} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ $\Gamma_{13}/\Gamma \times \Gamma_{14}/\Gamma$

VALUE (units 10^{-9}) EVTS DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.903 ± 0.076 ¹ BENAYOUN 10 RVUE 0.4-1.05 e^+e^-

4.58 $^{+2.46}_{-1.64}$ ± 1.56 1.2M ² ACHASOV 03D RVUE 0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

- ¹ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.
- ² Statistical significance is less than 3σ .

NODE=M009G3
 NODE=M009G3

NODE=M009G3;LINKAGE=BE
 NODE=M009G3;LINKAGE=AC

$\rho(770)$ BRANCHING RATIOS

NODE=M009235

$\Gamma(\pi^\pm\eta)/\Gamma(\pi\pi)$ Γ_5/Γ_1

VALUE (units 10^{-4}) CL% DOCUMENT ID TECN CHG COMMENT

<60 84 FERBEL 66 HBC ± $\pi^\pm p$ above 2.5

NODE=M009R4
 NODE=M009R4

$\Gamma(\pi^\pm \pi^+ \pi^- \pi^0)/\Gamma(\pi\pi)$ Γ_6/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<20	84	FERBEL	66	HBC	\pm $\pi^\pm p$ above 2.5
• • • We do not use the following data for averages, fits, limits, etc. • • •					
35 \pm 40		JAMES	66	HBC	+ 2.1 $\pi^+ p$

NODE=M009R1
 NODE=M009R1

 $\Gamma(\pi^+ \pi^- \gamma)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0099\pm0.0016 OUR FIT				
0.0099\pm0.0016		¹ DOLINSKY	91	ND $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0111 \pm 0.0014		² VASSERMAN	88	ND $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<0.005	90	³ VASSERMAN	88	ND $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

NODE=M009R12
 NODE=M009R12

- ¹ Bremsstrahlung from a decay pion and for photon energy above 50 MeV.
² Superseded by DOLINSKY 91.
³ Structure radiation due to quark rearrangement in the decay.

OCCUR=2

NODE=M009R12;LINKAGE=J
 NODE=M009R12;LINKAGE=I
 NODE=M009R12;LINKAGE=N

 $\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.20 \pm 0.52		¹ ACHASOV	16A	SND 0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
6.21 $^{+1.28}_{-1.18}$ \pm 0.39	18k	^{2,3} AKHMETSHIN	05	CMD2 0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
5.22 \pm 1.17 \pm 0.75	36k	^{3,4} ACHASOV	03	SND 0.60-0.97 $e^+ e^- \rightarrow \pi^0 \gamma$
6.8 \pm 1.7		⁵ BENAYOUN	96	RVUE 0.54-1.04 $e^+ e^- \rightarrow \pi^0 \gamma$
7.9 \pm 2.0		³ DOLINSKY	89	ND $e^+ e^- \rightarrow \pi^0 \gamma$

NODE=M009R9
 NODE=M009R9

- ¹ Using $B(\rho \rightarrow e^+ e^-)$ from PDG 15. Supersedes ACHASOV 03.
² Using $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$.
³ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}^2$.
⁴ Using $B(\rho \rightarrow e^+ e^-) = (4.54 \pm 0.10) \times 10^{-5}$.
⁵ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

NODE=M009R9;LINKAGE=C
 NODE=M009R9;LINKAGE=AK
 NODE=M009R9;LINKAGE=BZ
 NODE=M009R9;LINKAGE=AS
 NODE=M009R9;LINKAGE=A

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3.00\pm0.21 OUR FIT					
2.90\pm0.32 OUR AVERAGE					
2.79 \pm 0.34 \pm 0.03	33k	¹ ACHASOV	07B	SND	0.6-1.38 $e^+ e^- \rightarrow \eta\gamma$
3.6 \pm 0.9		² ANDREWS	77	CNTR 0	6.7-10 γ Cu
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.21 \pm 1.39 \pm 0.20	17.4k	^{3,4} AKHMETSHIN	05	CMD2	0.60-1.38 $e^+ e^- \rightarrow \eta\gamma$
3.39 \pm 0.42 \pm 0.23		^{2,5,6} AKHMETSHIN	01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
1.9 $^{+0.6}_{-0.8}$		⁷ BENAYOUN	96	RVUE	0.54-1.04 $e^+ e^- \rightarrow \eta\gamma$
4.0 \pm 1.1		^{2,4} DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$

NODE=M009R7
 NODE=M009R7

- ¹ ACHASOV 07B reports $[\Gamma(\rho(770) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\rho(770) \rightarrow e^+ e^-)] = (1.32 \pm 0.14 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\rho(770) \rightarrow e^+ e^-) = (4.72 \pm 0.05) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.
² Solution corresponding to constructive ω - ρ interference.
³ Using $B(\rho \rightarrow e^+ e^-) = (4.67 \pm 0.09) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
⁴ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.
⁵ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
⁶ Using $B(\rho \rightarrow e^+ e^-) = (4.75 \pm 0.10) \times 10^{-5}$ from AKHMETSHIN 02 and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
⁷ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution. Constructive ρ - ω interference solution.

NODE=M009R7;LINKAGE=AO

NODE=M009R7;LINKAGE=A
 NODE=M009R;LINKAGE=AK
 NODE=M009R7;LINKAGE=AZ
 NODE=M009R;LINKAGE=BQ

NODE=M009R;LINKAGE=BX

NODE=M009R7;LINKAGE=C

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{11}/Γ NODE=M009R14
NODE=M009R14**4.5±0.8 OUR FIT****4.5^{+0.9}_{-0.8} OUR AVERAGE**5.2^{+1.5}_{-1.3}±0.6 190 ¹ AKHMETSHIN 04B CMD2 0.6–0.97 $e^+e^- \rightarrow \pi^0\pi^0\gamma$

OCCUR=2

4.1^{+1.0}_{-0.9}±0.3 295 ² ACHASOV 02F SND 0.36–0.97 $e^+e^- \rightarrow \pi^0\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.8^{+3.4}_{-1.8}±0.5 63 ³ ACHASOV 00G SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$, and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0\pi^0$ with a branching ratio $(2.0^{+1.1}_{-0.9} \pm 0.3) \times 10^{-5}$ differing from zero by 2.0 standard deviations.

NODE=M009R14;LINKAGE=AH

² This branching ratio includes the conventional VMD mechanism $\rho \rightarrow \omega\pi^0$, $\omega \rightarrow \pi^0\gamma$ and the new decay mode $\rho \rightarrow f_0(500)\gamma$, $f_0(500) \rightarrow \pi^0\pi^0$ with a branching ratio $(1.9^{+0.9}_{-0.8} \pm 0.4) \times 10^{-5}$ differing from zero by 2.4 standard deviations. Supersedes ACHASOV 00G.

NODE=M009R;LINKAGE=FF

³ Superseded by ACHASOV 02F.

NODE=M009R;LINKAGE=GF

 $\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-)$ VALUE (units 10^{-5})

DOCUMENT ID

TECN

COMMENT

 Γ_{12}/Γ_7 NODE=M009R5
NODE=M009R5**4.60±0.28 OUR FIT****4.6 ±0.2 ±0.2**

ANTIPOV

89

SIGM

 $\pi^- \text{Cu} \rightarrow \mu^+\mu^-\pi^- \text{Cu}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.2^{+1.6}_{-3.6} ¹ ROTHWELL 69 CNTR Photoproduction5.6 ±1.5 ² WEHMANN 69 OSPK 12 $\pi^- \text{C, Fe}$ 9.7^{+3.1}_{-3.3} ^{3,4} HYAMS 67 OSPK 11 $\pi^- \text{Li, H}$

¹ Possibly large ρ - ω interference leads us to increase the minus error.

NODE=M009R5;LINKAGE=R

² Result contains $11 \pm 11\%$ correction using SU(3) for central value. The error on the correction takes account of possible ρ - ω interference and the upper limit agrees with the upper limit of $\omega \rightarrow \mu^+\mu^-$ from this experiment.

NODE=M009R5;LINKAGE=W

³ But he even enlarges his error to take residual ω contamination into account. Since his value is high, seems the other experiments also can't have too many ω 's. But maybe Hyams has additional μ 's from $\rho \rightarrow \pi\pi$, decaying π 's.

NODE=M009R5;LINKAGE=01

⁴ HYAMS 67's mass resolution is 20 MeV. The ω region was excluded.

NODE=M009R5;LINKAGE=H

 $\Gamma(e^+e^-)/\Gamma(\pi\pi)$ VALUE (units 10^{-4})

DOCUMENT ID

TECN

COMMENT

 Γ_{13}/Γ_1 NODE=M009R3
NODE=M009R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.40±0.05 ^{1,2} BENAKSAS 72 OSPK $e^+e^- \rightarrow \pi^+\pi^-$

¹ The ρ' contribution is not taken into account.

NODE=M009R;LINKAGE=KS

² Barkov excludes Auslender and Benaksas for large statistical and systematic errors.

NODE=M009R3;LINKAGE=01

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

CL%

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{14}/Γ NODE=M009R10
NODE=M009R10**0.88±0.23±0.30**¹ LEES

21B

BABR

 $10.5 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.01^{+0.54}_{-0.36}±0.34 1.2M ² ACHASOV 03D RVUE 0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ <1.2 90 VASSERMAN 88B ND $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ From the cross section for $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. Statistical evidence is more than 6σ .

NODE=M009R10;LINKAGE=A

² Statistical significance is less than 3σ .

NODE=M009R;LINKAGE=NS

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi\pi)$

VALUE

CL%

DOCUMENT ID

TECN

CHG

COMMENT

 Γ_{14}/Γ_1 NODE=M009R6
NODE=M009R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

~0.01 BRAMON 86 RVUE 0 $J/\psi \rightarrow \omega\pi^0$ <0.01 84 ¹ ABRAMS 71 HBC 0 $3.7 \pi^+p$

¹ Model dependent, assumes $l = 1, 2, \text{ or } 3$ for the 3π system.

NODE=M009R6;LINKAGE=G

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.8±0.9 OUR FIT					
1.8±0.9±0.3	153		AKHMETSHIN 00	CMD2	0.6–0.97 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<20	90		KURDADZE 88	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M009R13
NODE=M009R13

$$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma(\pi\pi) \quad \Gamma_{15}/\Gamma_1$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<15	90	ERBE	69	HBC	0 2.5–5.8 γp
<20		CHUNG	68	HBC	0 3.2,4.2 $\pi^- p$
<20	90	HUSON	68	HLBC	0 16.0 $\pi^- p$
<80		JAMES	66	HBC	0 2.1 $\pi^+ p$

NODE=M009R11
NODE=M009R11

$$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
1.60±0.74±0.18		¹ ACHASOV	09A	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 4	90	AULCHENKO	87C	ND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
<20	90	KURDADZE	86	OLYA $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
¹ Assuming no interference between the ρ and ω contributions.				

NODE=M009R8
NODE=M009R8

NODE=M009R8;LINKAGE=AC

$$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	ACHASOV	08	SND 0.36–0.97 $e^+e^- \rightarrow \pi^0 e^+ e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<1.6		AKHMETSHIN 05A	CMD2	0.72–0.84 e^+e^-

NODE=M009R15
NODE=M009R15

$$\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
<0.7	AKHMETSHIN 05A	CMD2	0.72–0.84 e^+e^-

NODE=M009R16
NODE=M009R16

$\rho(770)$ REFERENCES

ACHASOV	21	JHEP 2101 113	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61040
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
ANDREEV	20	EPJ C80 1189	V. Andreev <i>et al.</i>	(H1 Collab.)	REFID=60773
ABLIKIM	18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
ABLIKIM	17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57650
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57513
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=2
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
GARCIA-MAR...	11	PRL 107 072001	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=16761
BATLEY	10C	EPJ C70 635	J.R. Batley <i>et al.</i>	(CERN NA48/2 Collab.)	REFID=53567
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53101
		Translated from ZETF 136 442.			
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52258
		Translated from ZETF 134 80.			
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AULCHENKO	06A	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51513
		Translated from ZETFP 84 491.			
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51045
		Translated from ZETF 128 1201.			
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
ALOISIO	05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
		Translated from ZETFP 82 841.			

NODE=M009

SCHAEL	05C	PRPL 421 191	S. Schael et al.	(ALEPH Collab.)	REFID=50845
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=49609
AKHMETSHIN	04B	MPL B580 119	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=49610
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
ACHASOV	03	PL B559 171	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=49187
ACHASOV	03D	PR D68 052006	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=49577
ALOISIO	03	PL B561 55	A. Aloisio et al.	(KLOE Collab.)	REFID=49404
SANZ-CILLERO	03	EPJ C27 587	J.J. Sanz-Cillero, A. Pich		REFID=49399
ACHASOV	02	PR D65 032002	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02F	PL B537 201	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=48816
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=48565
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=48167
COLANGELO	01	NP B603 125	G. Colangelo, J. Gasser, H. Leytwyler		REFID=49180
PICH	01	PR D63 093005	A. Pich, J. Portoles		REFID=48313
ACHASOV	00	EPJ C12 25	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00D	JETPL 72 282	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=47882
ACHASOV	00G	Translated from ZETFP 72 411. JETPL 71 355	M.N. Achasov et al.	(Novosibirsk SND Collab.)	REFID=47929
AKHMETSHIN	00	Translated from ZETFP 71 519. PL B475 190	R.R. Akhmetshin et al.	(Novosibirsk CMD-2 Collab.)	REFID=47421
ANDERSON	00A	PR D61 112002	S. Anderson et al.	(CLEO Collab.)	REFID=47468
ABELE	99E	PL B469 270	A. Abele et al.	(Crystal Barrel Collab.)	REFID=47414
BENAYOUN	98	EPJ C2 269	M. Benayoun et al.	(IPNP, NOVO, ADLD+)	REFID=45859
BREITWEG	98B	EPJ C2 247	J. Breitweg et al.	(ZEUS Collab.)	REFID=46354
GARDNER	98	PR D57 2716	S. Gardner, H.B. O'Connell		REFID=46366
		Also PR D62 019903 (err.)	S. Gardner, H.B. O'Connell		REFID=47981
ABELE	97	PL B391 191	A. Abele et al.	(Crystal Barrel Collab.)	REFID=45415
ADAMS	97	ZPHY C74 237	M.R. Adams et al.	(E665 Collab.)	REFID=45533
BARATE	97M	ZPHY C76 15	R. Barate et al.	(ALEPH Collab.)	REFID=45622
BERTIN	97C	PL B408 476	A. Bertin et al.	(OBELIX Collab.)	REFID=45701
BOGOLYUB...	97	PAN 60 46	M.Y. Bogolyubsky et al.	(MOSU, SERP)	REFID=45393
O'CONNELL	97	Translated from YAF 60 53. NP A623 559	H.B. O'Connell et al.	(ADLD)	REFID=45860
BENAYOUN	96	ZPHY C72 221	M. Benayoun et al.	(IPNP, NOVO)	REFID=45753
BERNICH	94	PR D50 4454	A. Bernicha, G. Lopez Castro, J. Pestieau	(LOUV+)	REFID=44097
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer et al.	(ASTERIX Collab.)	REFID=43585
AGUILAR...	91	ZPHY C50 405	M. Aguilar-Benitez et al.	(LEBC-EHS Collab.)	REFID=41637
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky et al.	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn et al.	(MPIM)	REFID=45862
ANTIPOV	89	ZPHY C42 185	Y.M. Antipov et al.	(SERP, JINR, BGNA+)	REFID=40739
BISELLO	89	PL B220 321	D. Bisello et al.	(DM2 Collab.)	REFID=40740
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky et al.	(NOVO)	REFID=41003
DUBNICKA	89	JP G15 1349	S. Dubnicka et al.	(JINR, SLOV)	REFID=44082
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
KURDADZE	88	JETPL 47 512	L.M. Kurdadze et al.	(NOVO)	REFID=41121
VASSERMAN	88	Translated from ZETFP 47 432. SJNP 47 1035	I.B. Vasserman et al.	(NOVO)	REFID=41019
VASSERMAN	88B	Translated from YAF 47 1635. SJNP 48 480	I.B. Vasserman et al.	(NOVO)	REFID=41020
AULCHENKO	87C	Translated from YAF 48 753. IYF 87-90 Preprint	V.M. Aulchenko et al.	(NOVO)	REFID=41370
CAPRARO	87	NP B288 659	L. Capraro et al.	(CLER, FRAS, MILA+)	REFID=40003
BRAMON	86	PL B173 97	A. Bramon, J. Casulleras	(BARC)	REFID=22102
HUSTON	86	PR D33 3199	J. Huston et al.	(ROCH, FNAL, MINN)	REFID=20137
KURDADZE	86	JETPL 43 643	L.M. Kurdadze et al.	(NOVO)	REFID=40287
BARKOV	85	Translated from ZETFP 43 497. NP B256 365	L.M. Barkov et al.	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin et al.	(NOVO)	REFID=20561
CHABAUD	83	NP B223 1	V. Chabaud et al.	(CERN, CRAC, MPIM)	REFID=20131
JENSEN	83	PR D27 26	T. Jensen et al.	(ROCH, FNAL, MINN)	REFID=20132
HEYN	81	ZPHY C7 169	M.F. Heyn, C.B. Lang	(GRAZ)	REFID=20129
BOHACIK	80	PR D21 1342	J. Bohacik, H. Kuhnelt	(SLOV, WIEN)	REFID=20128
COHEN	80	PR D22 2595	D. Cohen et al.	(ANL)	REFID=20381
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)	REFID=20126
BARTALUCCI	78	NC 44A 587	S. Bartalucci et al.	(DESY, FRAS)	REFID=20122
WICKLUND	78	PR D17 1197	A.B. Wicklund et al.	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews et al.	(ROCH)	REFID=20120
DEUTSCH...	76	NP B103 426	M. Deuschmann et al.	(AACH3, BERL, BONN+)	REFID=20119
ENGLER	74	PR D10 2070	A. Engler et al.	(CMU, CASE)	REFID=20110
ESTABROOKS	74	NP B79 301	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20111
GRAYER	74	NP B75 189	G. Grayer et al.	(CERN, MPIM)	REFID=20113
RATCLIFF	74	Private Comm.			REFID=40128 ERROR=3
BYERLY	73	PR D7 637	W.L. Byerly et al.	(MICH)	REFID=20104
GLADDING	73	PR D8 3721	G.E. Gladding et al.	(HARV)	REFID=20106
HYAMS	73	NP B64 134	B.D. Hyams et al.	(CERN, MPIM)	REFID=20107
PROTOPOP...	73	PR D7 1279	S.D. Protopopescu et al.	(LBL)	REFID=20108
BALLAM	72	PR D5 545	J. Ballam et al.	(SLAC, LBL, TUFTS)	REFID=20094
BENAKSAS	72	PL 39B 289	D. Benaksas et al.	(ORSAY)	REFID=20096
JACOBS	72	PR D6 1291	L.D. Jacobs	(SACL)	REFID=20101
RATCLIFF	72	PL 38B 345	B.N. Ratcliff et al.	(SLAC)	REFID=20102
ABRAMS	71	PR D4 653	G.S. Abrams et al.	(LBL)	REFID=20090
ALVENSLEB...	70	PRL 24 786	H. Alvensleben et al.	(DESY)	REFID=20085
BIGGS	70	PRL 24 1197	P.J. Biggs et al.	(DARE)	REFID=20087
ERBE	69	PR 188 2060	R. Erbe et al.	(German Bubble Chamber Collab.)	REFID=20074
MALAMUD	69	Argonne Conf. 93	E.I. Malamud, P.E. Schlein	(UCLA)	REFID=20077
REYNOLDS	69	PR 184 1424	B.G. Reynolds et al.	(FSU)	REFID=20080
ROTHWELL	69	PRL 23 1521	P.L. Rothwell et al.	(NEAS)	REFID=20082
WEHMANN	69	PR 178 2095	A.A. Wehmann et al.	(HARV, CASE, SLAC+)	REFID=20084
ARMENISE	68	NC 54A 999	N. Armenise et al.	(BARI, BGNA, FIRZ+)	REFID=20054
BATON	68	PR 176 1574	J.P. Baton, G. Laurens	(SACL)	REFID=20056
CHUNG	68	PR 165 1491	S.U. Chung et al.	(LRL)	REFID=20059
FOSTER	68	NP B6 107	M. Foster et al.	(CERN, CDEF)	REFID=20061
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054
HUSON	68	PL 28B 208	R. Huson et al.	(ORSAY, MILA, UCLA)	REFID=20062
HYAMS	68	NP B7 1	B.D. Hyams et al.	(CERN, MPIM)	REFID=20063
LANZEROTTI	68	PR 166 1365	L.J. Lanzerotti et al.	(HARV)	REFID=20068
PISUT	68	NP B6 325	J. Pisut, M. Roos	(CERN)	REFID=20070
ASBURY	67B	PRL 19 865	J.G. Asbury et al.	(DESY, COLU)	REFID=20038
BACON	67	PR 157 1263	T.C. Bacon et al.	(BNL)	REFID=20039
EISNER	67	PR 164 1699	R.L. Eisner et al.	(PURD)	REFID=20046
HUWE	67	PL 24B 252	D.O. Huwe et al.	(COLU)	REFID=20049
HYAMS	67	PL 24B 634	B.D. Hyams et al.	(CERN, MPIM)	REFID=20050
MILLER	67B	PR 153 1423	D.H. Miller et al.	(PURD)	REFID=20051
ALFF...	66	PR 145 1072	C. Alff-Steinberger et al.	(COLU, RUTG)	REFID=10762
FERBEL	66	PL 21 111	T. Ferbel	(ROCH)	REFID=20028
HAGOPIAN	66	PR 145 1128	V. Hagopian et al.	(PENN, SACL)	REFID=20030
HAGOPIAN	66B	PR 152 1183	V. Hagopian, Y.L. Pan	(PENN, LRL)	REFID=20031
JACOBS	66B	UCRL 16877	L.D. Jacobs	(LRL)	REFID=20033
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
ROSS	66	PR 149 1172	M. Ross, L. Stodolsky		REFID=46380
SOEDING	66	PL B19 702	P. Soeding		REFID=46385
WEST	66	PR 149 1089	E. West et al.	(WISC)	REFID=20035
BLIEDEN	65	PL 19 444	H.R. Blieden et al.	(CERN MMS Collab.)	REFID=20016
CARMONY	64	PRL 12 254	D.D. Carmony et al.	(UCB)	REFID=20578
GOLDHABER	64	PRL 12 336	G. Goldhaber et al.	(LRL, UCB)	REFID=20013
ABOLINS	63	PRL 11 381	M.A. Abolins et al.	(UCSD)	REFID=20006

$\omega(782)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M001

 $\omega(782)$ MASS

NODE=M001M

NODE=M001M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
782.66±0.13 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.		
777.9 ±2.2 $\begin{smallmatrix} +4.3 \\ -2.2 \end{smallmatrix}$	900k	ANDREEV	20 H1	$e p \rightarrow e \pi^+ \pi^- p$
783.20±0.13±0.16	18680	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+ e^- \rightarrow \pi^0 \gamma$
782.68±0.09±0.04	11200	¹ AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.79±0.08±0.09	1.2M	² ACHASOV 03D	RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
781.96±0.17±0.80	11k	³ AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
782.08±0.36±0.82	3463	⁴ AMSLER 94C	CBAR	0.0 $\bar{p} p \rightarrow \omega \eta \pi^0$
781.96±0.13±0.17	15k	AMSLER 93B	CBAR	0.0 $\bar{p} p \rightarrow \omega \pi^0 \pi^0$
782.4 ±0.2	270k	WEIDENAUER 93	ASTE	$\bar{p} p \rightarrow 2\pi^+ 2\pi^- \pi^0$
782.4 ±0.5	7000	⁵ KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
782.58±0.03±0.01		⁶ HOID 20	RVUE	$e^+ e^- \rightarrow \pi^0 \gamma$
781.68±0.09±0.03		⁷ COLANGELO 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-$
782.63±0.03±0.01		⁸ HOFERICH... 19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
781.91±0.24		⁹ LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
782.7 ±0.1 ±1.5	19500	¹⁰ WURZINGER 95	SPEC	1.33 $p d \rightarrow {}^3\text{He} \omega$
781.78±0.10		¹⁰ BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.2 ±0.4	1488	¹¹ KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
783.3 ±0.4	433	CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
782.5 ±0.8	33260	ROOS 80	RVUE	0.0-3.6 $\bar{p} p$
782.6 ±0.8	3000	BENKHEIRI 79	OMEG	9-12 $\pi^\pm p$
781.8 ±0.6	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p} p \rightarrow 5\pi$
782.7 ±0.9	535	VANAPEL... 78	HBC	7.2 $\bar{p} p \rightarrow \bar{p} p \omega$
783.5 ±0.8	2100	GESSAROLI 77	HBC	11 $\pi^- p \rightarrow \omega n$
782.5 ±0.8	418	AGUILAR-... 72B	HBC	3.9,4.6 $K^- p$
783.4 ±1.0	248	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K^+ K^- \omega$
781.0 ±0.6	510	BIZZARRI 71	HBC	0.0 $p \bar{p} \rightarrow K_1^+ K_1^- \omega$
783.7 ±1.0	3583	¹² COYNE 71	HBC	3.7 $\pi^+ p \rightarrow \rho \pi^+ \pi^+ \pi^- \pi^0$
784.1 ±1.2	750	ABRAMOVI... 70	HBC	3.9 $\pi^- p$
783.2 ±1.6		¹³ BIGGS 70B	CNTR	<4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
782.4 ±0.5	2400	BIZZARRI 69	HBC	0.0 $\bar{p} p$

OCCUR=2

OCCUR=2

OCCUR=2


¹ Update of AKHMETSHIN 00C.² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.³ From the $\eta \rightarrow \gamma \gamma$ decay.⁴ From the $\eta \rightarrow 3\pi^0$ decay.⁵ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.⁶ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 782.736 ± 0.024 MeV.⁷ The ω mass was extracted from a dispersively improved Breit-Wigner parameterization, the ω width fixed at 8.49 ± 0.08 MeV. The value does not include vacuum polarization which would shift the mass to $781.81 \pm 0.09 \pm 0.03$ MeV. The mixing parameter is assumed real valued.⁸ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.⁹ From the $\rho - \omega$ interference in the $\pi^+ \pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.¹⁰ Systematic uncertainties underestimated.¹¹ Systematic uncertainties not estimated.¹² From best-resolution sample of COYNE 71.¹³ From $\omega - \rho$ interference in the $\pi^+ \pi^-$ mass spectrum assuming ω width 12.6 MeV.NODE=M001M;LINKAGE=PT
NODE=M001M;LINKAGE=VHNODE=M001M;LINKAGE=S1
NODE=M001M;LINKAGE=S2
NODE=M001M;LINKAGE=B
NODE=M001M;LINKAGE=G

NODE=M001M;LINKAGE=A

NODE=M001M;LINKAGE=H

NODE=M001M;LINKAGE=LE

NODE=M001M;LINKAGE=KB
NODE=M001M;LINKAGE=I
NODE=M001M;LINKAGE=D
NODE=M001M;LINKAGE=F



$\omega(782)$ WIDTH

NODE=M001W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.68±0.13 OUR AVERAGE				
8.68±0.23±0.10	11200	1 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.68±0.04±0.15	1.2M	2 ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
8.65±0.06±0.01		3 HOID 20	RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.71±0.04±0.04		4 HOFERICH... 19	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.13±0.45		5 LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
8.2 ±0.3	19500	6 WURZINGER 95	SPEC	1.33 $p d \rightarrow {}^3\text{He}\omega$
8.4 ±0.1		7 AULCHENKO 87	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.30±0.40		6 BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.8 ±0.9	1488	8 KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0 ±0.8	433	6 CORDIER 80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
12 ±2	1430	COOPER 78B	HBC	0.7-0.8 $\bar{p}p \rightarrow 5\pi$
9.4 ±2.5	2100	GESSAROLI 77	HBC	$11 \pi^- p \rightarrow \omega n$
10.22±0.43	20000	9 KEYNE 76	CNTR	$\pi^- p \rightarrow \omega n$
13.3 ±2	418	AGUILAR... 72B	HBC	3.9,4.6 $K^- p$
9.1 ±0.8	451	6 BENAKSAS 72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
10.5 ±1.5		BORENSTEIN 72	HBC	2.18 $K^- p$
7.70±0.9 ±1.15	940	BROWN 72	MMS	2.5 $\pi^- p \rightarrow nMM$
10.3 ±1.4	510	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K_1^+ K_1^-\omega$
12.8 ±3.0	248	BIZZARRI 71	HBC	0.0 $p\bar{p} \rightarrow K^+ K^-\omega$
9.5 ±1.0	3583	COYNE 71	HBC	3.7 $\pi^+ p \rightarrow \rho\pi^+\pi^+\pi^-\pi^0$

NODE=M001W

OCCUR=2

¹ Update of AKHMETSHIN 00C.

NODE=M001W;LINKAGE=PT
NODE=M001W;LINKAGE=VH

² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

NODE=M001W;LINKAGE=E

³ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 8.63 ± 0.05 MeV.

NODE=M001W;LINKAGE=F

⁴ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

NODE=M001W;LINKAGE=LE

⁵ From the $\rho-\omega$ interference in the $\pi^+\pi^-$ mass spectrum using the Breit-Wigner for the ω and leaving its mass and width as free parameters of the fit.

NODE=M001W;LINKAGE=I

⁶ Systematic uncertainties underestimated.

NODE=M001W;LINKAGE=G

⁷ Relativistic Breit-Wigner includes radiative corrections. Systematic uncertainties not estimated.

NODE=M001W;LINKAGE=J

⁸ Systematic uncertainties not estimated.

NODE=M001W;LINKAGE=B

⁹ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

$\omega(782)$ DECAY MODES

NODE=M001215;NODE=M001

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 $\pi^+\pi^-\pi^0$	(89.2 \pm 0.7) %		DESIG=1
Γ_2 $\pi^0\gamma$	(8.35 \pm 0.27) %	S=2.2	DESIG=3
Γ_3 $\pi^+\pi^-$	(1.53 \pm 0.12) %	S=1.2	DESIG=2
Γ_4 neutrals (excluding $\pi^0\gamma$)	(7 $^{+8}_{-4}$) $\times 10^{-3}$	S=1.1	DESIG=13
Γ_5 $\eta\gamma$	(4.5 \pm 0.4) $\times 10^{-4}$	S=1.1	DESIG=6
Γ_6 $\pi^0e^+e^-$	(7.7 \pm 0.6) $\times 10^{-4}$		DESIG=14
Γ_7 $\pi^0\mu^+\mu^-$	(1.34 \pm 0.18) $\times 10^{-4}$	S=1.5	DESIG=11
Γ_8 ηe^+e^-			DESIG=18
Γ_9 e^+e^-	(7.38 \pm 0.22) $\times 10^{-5}$	S=1.9	DESIG=7
Γ_{10} $\pi^+\pi^-\pi^0\pi^0$	< 2 $\times 10^{-4}$	CL=90%	DESIG=12
Γ_{11} $\pi^+\pi^-\gamma$	< 3.6 $\times 10^{-3}$	CL=95%	DESIG=4
Γ_{12} $\pi^+\pi^-\pi^+\pi^-$	< 1 $\times 10^{-3}$	CL=90%	DESIG=15
Γ_{13} $\pi^0\pi^0\gamma$	(6.7 \pm 1.1) $\times 10^{-5}$		DESIG=5
Γ_{14} $\eta\pi^0\gamma$	< 3.3 $\times 10^{-5}$	CL=90%	DESIG=17
Γ_{15} $\mu^+\mu^-$	(7.4 \pm 1.8) $\times 10^{-5}$		DESIG=8
Γ_{16} 3γ	< 1.9 $\times 10^{-4}$	CL=95%	DESIG=10
Charge conjugation (C) violating modes			
Γ_{17} $\eta\pi^0$	C < 2.1 $\times 10^{-4}$	CL=90%	NODE=M001;CLUMP=A DESIG=9
Γ_{18} $2\pi^0$	C < 2.2 $\times 10^{-4}$	CL=90%	DESIG=193
Γ_{19} $3\pi^0$	C < 2.3 $\times 10^{-4}$	CL=90%	DESIG=16
Γ_{20} invisible	< 7 $\times 10^{-5}$	CL=90%	DESIG=194

CONSTRAINED FIT INFORMATION

An overall fit to 15 branching ratios uses 48 measurements and one constraint to determine 10 parameters. The overall fit has a $\chi^2 = 48.0$ for 39 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	23								
x_3	-18	-4							
x_4	-92	-55	1						
x_5	7	23	-1	-15					
x_6	-1	0	0	0	0				
x_7	0	0	0	0	0	0			
x_9	-24	-73	4	47	-31	0	0		
x_{13}	1	4	0	-2	1	0	0	-3	
x_{15}	0	0	0	0	0	0	0	0	0
	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_9	x_{13}

 $\omega(782)$ PARTIAL WIDTHS

NODE=M001218

$\Gamma(\pi^0\gamma)$	Γ_2					
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT		
880 \pm 50	7815	¹ ACHASOV	13	SND	1.05-2.00	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
788 \pm 12 \pm 27	36500	² ACHASOV	03	SND	0.60-0.97	$e^+e^- \rightarrow \pi^0\gamma$
764 \pm 51	10625	DOLINSKY	89	ND		$e^+e^- \rightarrow \pi^0\gamma$

NODE=M001W1
NODE=M001W1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

¹Systematic uncertainty not estimated.²Using $\Gamma_\omega = 8.44 \pm 0.09$ MeV and $B(\omega \rightarrow \pi^0\gamma)$ from ACHASOV 03.NODE=M001W1;LINKAGE=AC
NODE=M001W1;LINKAGE=AD

$\Gamma(\eta\gamma)$ Γ_5

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1±2.5	¹ DOLINSKY	89	ND	$e^+e^- \rightarrow \eta\gamma$
---------	-----------------------	----	----	---------------------------------

¹ Using $\Gamma_\omega = 8.4 \pm 0.1$ MeV and $B(\omega \rightarrow \eta\gamma)$ from DOLINSKY 89.

NODE=M001W2
NODE=M001W2

NODE=M001W2;LINKAGE=DA

 $\Gamma(e^+e^-)$ Γ_9

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.60 ± 0.02 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.591±0.015	11200	^{1,2} AKHMETSHIN	04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-------------	-------	---------------------------	----	------	--------------------------------------

0.653±0.003±0.021	1.2M	³ ACHASOV	03D	RVUE	0.44–2.00 $e^+e^- \rightarrow$
-------------------	------	----------------------	-----	------	--------------------------------

0.600±0.031	10625	DOLINSKY	89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-------------	-------	----------	----	----	--

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.891 \pm 0.007$ and $\Gamma_{\text{total}} = 8.44 \pm 0.09$ MeV.

² Update of AKHMETSHIN 00C.

³ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

NODE=M001W7
NODE=M001W7

→ UNCHECKED ←

NODE=M001W7;LINKAGE=3P
NODE=M001W7;LINKAGE=PT
NODE=M001W;LINKAGE=VF

 $\omega(782) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M001235

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_9/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

569.8±3.1±8.2	¹ LEES	21B	BABR	10.5 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
----------------------	-------------------	-----	------	---

¹ From the cross section for $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

NODE=M001R02
NODE=M001R02

NODE=M001R02;LINKAGE=A

 $\omega(782) \Gamma(e^+e^-)\Gamma(i)/\Gamma^2(\text{total})$

NODE=M001225

 $\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.59±0.19 OUR FIT Error includes scale factor of 2.1.

6.36±0.14 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

6.24±0.11±0.08	11.2k	¹ AKHMETSHIN	04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
----------------	-------	-------------------------	----	------	--------------------------------------

6.74±0.04±0.24	1.2M	^{2,3} ACHASOV	03D	RVUE	0.44–2.00 $e^+e^- \rightarrow$
----------------	------	------------------------	-----	------	--------------------------------

6.37±0.35		² DOLINSKY	89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
-----------	--	-----------------------	----	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.20±0.13		⁴ BENAYOUN	10	RVUE	0.4–1.05 e^+e^-
-----------	--	-----------------------	----	------	-------------------

6.70±0.06±0.27		⁵ AUBERT,B	04N	BABR	10.6 $e^+e^- \rightarrow$
----------------	--	-----------------------	-----	------	---------------------------

6.45±0.24		⁶ BARKOV	87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	--	---------------------	----	-----	--------------------------------------

5.79±0.42	1488	⁷ KURDADZE	83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	------	-----------------------	-----	------	--------------------------------------

5.89±0.54	433	⁶ CORDIER	80	DM1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	-----	----------------------	----	-----	--------------------------------------

7.54±0.84	451	⁶ BENAJSAS	72B	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
-----------	-----	-----------------------	-----	------	--------------------------------------

¹ Update of AKHMETSHIN 00C.

² Recalculated by us from the cross section in the peak.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.

⁵ Superseded by LEES 21B.

⁶ Recalculated by us from the cross section in the peak. Systematic uncertainties underestimated.

⁷ Recalculated by us from the cross section in the peak. Systematic uncertainties not estimated.


NODE=M001G2
NODE=M001G2

NODE=M001G;LINKAGE=PT
NODE=M001G;LINKAGE=LP
NODE=M001G;LINKAGE=VH

NODE=M001G2;LINKAGE=BE
NODE=M001G2;LINKAGE=C
NODE=M001G2;LINKAGE=A

NODE=M001G2;LINKAGE=B

WEIGHTED AVERAGE
6.36±0.14 (Error scaled by 1.3)



$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma \times \Gamma_2/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
6.16 ±0.14 OUR FIT				Error includes scale factor of 1.8.
6.34 ±0.10 OUR AVERAGE				
6.336 ±0.056 ±0.089		¹ ACHASOV 16A SND		0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
6.47 ±0.14 ±0.39	18k	AKHMETSHIN 05 CMD2		0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
6.34 ±0.21 ±0.21	10k	² DOLINSKY 89 ND		$e^+e^- \rightarrow \pi^0\gamma$
6.80 ±0.13		³ BENAYOUN 10 RVUE		0.4–1.05 e^+e^-
6.50 ±0.11 ±0.20	36k	⁴ ACHASOV 03 SND		0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$

NODE=M001G4
NODE=M001G4

¹ From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 03.

² Recalculated by us from the cross section in the peak.

³ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.

⁴ Using $\sigma(\phi \rightarrow \pi^0\gamma)$ from ACHASOV 00 and $m_\omega = 782.57$ MeV in the model with the energy-independent phase of ρ - ω interference equal to $(-10.2 \pm 7.0)^\circ$.

NODE=M001G4;LINKAGE=A

NODE=M001G4;LINKAGE=LP
NODE=M001G4;LINKAGE=BE
NODE=M001G;LINKAGE=SH

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma \times \Gamma_3/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.28 ±0.05 OUR AVERAGE				
1.318 ±0.051 ±0.021		¹ ACHASOV 21 SND		$e^+e^- \rightarrow \pi^+\pi^-$
1.225 ±0.058 ±0.041	800k	² ACHASOV 06 SND		$e^+e^- \rightarrow \pi^+\pi^-$
1.166 ±0.036		³ BENAYOUN 13 RVUE		0.4–1.05 e^+e^-
1.05 ±0.08		⁴ DAVIER 13 RVUE		$e^+e^- \rightarrow \pi^+\pi^-(\gamma)$

NODE=M001G5
NODE=M001G5

¹ From a fit of the cross section in the energy range $0.525 < \sqrt{s} < 0.883$ GeV parameterized by the sum of the Breit-Wigner amplitudes for the $\rho(770)$, ω and $\rho(1450)$ resonances. The measured phase of the $\rho(770)$ - ω interference is $(110.7 \pm 1.5 \pm 1.0)^\circ$.

² Supersedes ACHASOV 05A.

³ A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$, $K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data. Supersedes BENAYOUN 10.

⁴ From $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$ data of LEES 12G.

NODE=M001G5;LINKAGE=C

NODE=M001G5;LINKAGE=AC
NODE=M001G5;LINKAGE=B

NODE=M001G5;LINKAGE=A

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}$

$\Gamma_9/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
3.32 ±0.28 OUR FIT				Error includes scale factor of 1.1. $[(3.32 \pm 0.28) \times 10^{-8}]$ OUR 2023 FIT Scale factor = 1.1]
3.18 ±0.28 OUR AVERAGE				
3.10 ±0.31 ±0.11	33k	¹ ACHASOV 07B SND		0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
3.17 ^{+1.85} _{-1.31} ±0.21	17.4k	² AKHMETSHIN 05 CMD2		0.60–1.38 $e^+e^- \rightarrow \eta\gamma$
3.41 ±0.52 ±0.21	23k	^{3,4} AKHMETSHIN 01B CMD2		$e^+e^- \rightarrow \eta\gamma$
4.50 ±0.10		⁵ BENAYOUN 10 RVUE		0.4–1.05 e^+e^-

NODE=M001G3
NODE=M001G3

NEW

- ¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- ² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
- ³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
- ⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
- ⁵ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.

NODE=M001G3;LINKAGE=AH

NODE=M001G;LINKAGE=AH

NODE=M001G;LINKAGE=AK

NODE=M001G;LINKAGE=BQ

NODE=M001G3;LINKAGE=BE

NODE=M001G01

NODE=M001G01

NODE=M001G01;LINKAGE=A

$\Gamma(e^+e^-)/\Gamma_{\text{total}} \times \Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \Gamma_{15}/\Gamma$

VALUE (units 10^{-9})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±1.8±2.2	4.5M	¹ ANASTASI	17	KLOE $e^+e^- \rightarrow \mu^+\mu^-\gamma$

- ¹ From a fit of the real part of the vacuum polarization by a sum of the leptonic and hadronic contributions, where the hadronic contribution is parametrized as a sum of Breit-Wigner resonances $\omega(782)$, $\phi(1020)$ and using a GOUNARIS 68 parametrization for the $\rho(770)$, and a non-resonant term.

NODE=M001220

$\omega(782)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9024±0.0019		¹ AMBROSINO 08G	KLOE	1.0–1.03 $e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
0.8965±0.0016±0.0048	1.2M	^{2,3} ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.880 ±0.020 ±0.032	11200	^{3,4} AKHMETSHIN 00C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.8942±0.0062		³ DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

- ¹ Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

- ² Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

- ³ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$.

- ⁴ Using $\Gamma(e^+e^-) = 0.60 \pm 0.02$ keV.

NODE=M001R21

NODE=M001R21

NODE=M001R21

NODE=M001R21;LINKAGE=AM

NODE=M001R;LINKAGE=VF

NODE=M001R;LINKAGE=ZL

NODE=M001R;LINKAGE=KH

$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.88±0.18		¹ ACHASOV 16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
8.09±0.14		² AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
9.06±0.20±0.57	18k	^{3,4} AKHMETSHIN 05	CMD2	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
9.34±0.15±0.31	36k	⁴ ACHASOV 03	SND	0.60–0.97 $e^+e^- \rightarrow \pi^0\gamma$
8.65±0.16±0.42	1.2M	^{5,6} ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.39±0.24	9k	⁷ BENAYOUN 96	RVUE	$e^+e^- \rightarrow \pi^0\gamma$
8.88±0.62	10k	⁴ DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

- ¹ Using $B(\omega \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 03.

- ² Not independent of $\Gamma(\pi^0\gamma) / \Gamma(\pi^+\pi^-\pi^0)$ from AMBROSINO 08G.

- ³ Using $B(\omega \rightarrow e^+e^-) = (7.14 \pm 0.13) \times 10^{-5}$.

- ⁴ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

- ⁵ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+\pi^-) = (1.70 \pm 0.28)\%$.

- ⁶ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}^2$.

- ⁷ Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001R28

NODE=M001R28

NODE=M001R28;LINKAGE=A

NODE=M001R28;LINKAGE=AM

NODE=M001R;LINKAGE=AH

NODE=M001R;LINKAGE=VL

NODE=M001R28;LINKAGE=VF

NODE=M001R28;LINKAGE=ZL

NODE=M001R28;LINKAGE=A1

$\Gamma(\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0) \quad \Gamma_2/\Gamma_1$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

9.35±0.30 OUR FIT Error includes scale factor of 2.4.

9.05±0.27 OUR AVERAGE Error includes scale factor of 1.8.

8.97±0.16	AMBROSINO 08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
9.94±0.36±0.38	¹ AULCHENKO 00A	SND	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
8.4 ±1.3	KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
10.9 ±2.5	BENAKSAS 72C	OSPK	$e^+e^- \rightarrow \pi^0\gamma$
8.1 ±2.0	BALDIN 71	HLBC	$2.9 \pi^+\pi^-$
13 ±4	JACQUET 69B	HLBC	$2.05 \pi^+\pi^- \rightarrow \pi^+\pi^-\omega$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.7 ±0.2 ±0.5	^{2,3} ACHASOV 03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.9 ±0.7	² DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\gamma$

NODE=M001R3

NODE=M001R3

- 1 From $\sigma_0^{\omega\pi^0 \rightarrow \pi^0\pi^0\gamma}(m_\phi)/\sigma_0^{\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0\pi^0}(m_\phi)$ with a phase-space correction factor of 1/1.023.
- 2 Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.
- 3 Using ACHASOV 03. Based on 1.2M events.

NODE=M001R3;LINKAGE=AL

NODE=M001R3;LINKAGE=VL
NODE=M001R3;LINKAGE=VW $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_3/Γ See also $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$.NODE=M001R15
NODE=M001R15
NODE=M001R15

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
1.53±0.12 OUR FIT				Error includes scale factor of 1.2. $[(1.53^{+0.11}_{-0.13}) \times 10^{-2}$ OUR 2023 FIT Scale factor = 1.2]

NEW

1.49±0.13 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
------------------------------	--	--	--	---

1.46±0.12±0.02	900k	1 AKHMETSHIN 07		$e^+e^- \rightarrow \pi^+\pi^-$
1.30±0.24±0.05	11.2k	2 AKHMETSHIN 04	CMD2	$e^+e^- \rightarrow \pi^+\pi^-$
2.38 ^{+1.77} _{-0.90} ±0.18	5.4k	3 ACHASOV 02E	SND	1.1-1.38 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
2.3 ±0.5		BARKOV 85	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
1.6 ^{+0.9} _{-0.7}		QUENZER 78	DM1	$e^+e^- \rightarrow \pi^+\pi^-$
3.6 ±1.9		BENAKSAS 72	OSPK	$e^+e^- \rightarrow \pi^+\pi^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1.29±0.22±0.03	970k	4,5 ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
1.28±0.22±0.03	970k	6,7 ABLIKIM 18C	BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
1.52±0.08		8 HANHART 18	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1.75±0.11	4.5M	9 ACHASOV 05A	SND	$e^+e^- \rightarrow \pi^+\pi^-$
2.01±0.29		10 BENAYOUN 03	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1.9 ±0.3		11 GARDNER 99	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
2.3 ±0.4		12 BENAYOUN 98	RVUE	$e^+e^- \rightarrow \pi^+\pi^-, \mu^+\mu^-$
1.0 ±0.11		13 WICKLUND 78	ASPK	3,4,6 $\pi^\pm N$
1.22±0.30		ALVENSLEB... 71C	CNTR	Photoproduction
1.3 ^{+1.2} _{-0.9}		MOFFEIT 71	HBC	2.8,4.7 γp
0.80 ^{+0.28} _{-0.20}		14 BIGGS 70B	CNTR	4.2 $\gamma C \rightarrow \pi^+\pi^- C$

OCCUR=3

OCCUR=2

1 A combined fit of AKHMETSHIN 07, AULCHENKO 06, and AULCHENKO 05.

NODE=M001R15;LINKAGE=AK

2 Update of AKHMETSHIN 02.

NODE=M001R15;LINKAGE=PT

3 From the $m_{\pi^+\pi^-}$ spectrum taking into account the interference of the $\rho\pi$ and $\omega\pi$ amplitudes.

NODE=M001R;LINKAGE=VE

4 From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and box anomaly components.

NODE=M001R15;LINKAGE=E

5 ABLIKIM 18C reports $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.25 \pm 0.21 \pm 0.52) \times 10^{-4}$ which we divide by our best value $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M001R15;LINKAGE=H

6 From a fit to $\pi^+\pi^-$ mass using $\rho(770)$ (parametrized with the Gounaris-Sakurai approach), $\omega(782)$, and $\rho(1450)$ components.

NODE=M001R15;LINKAGE=J

7 ABLIKIM 18C reports $[\Gamma(\omega(782) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \omega\gamma)] = (3.22 \pm 0.21 \pm 0.52) \times 10^{-4}$ which we divide by our best value $B(\eta'(958) \rightarrow \omega\gamma) = (2.52 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M001R15;LINKAGE=K

8 Dispersive analysis. Value extracted from average of data from AUBERT 09AS, AKHMETSHIN 07, ACHASOV 06, AMBROSINO 11A, BABUSCI 13D, ABLIKIM 16B normalised by PDG 16 evaluation for $\Gamma(\omega \rightarrow e^+e^-)$.

NODE=M001R15;LINKAGE=D

9 Using $\Gamma(\omega \rightarrow e^+e^-)$ from the 2004 Edition of this Review (PDG 04).

NODE=M001R;LINKAGE=SN

10 Using the data of AKHMETSHIN 02 in the hidden local symmetry model.

NODE=M001R;LINKAGE=BY

11 Using the data of BARKOV 85.

NODE=M001R15;LINKAGE=H4

12 Using the data of BARKOV 85 in the hidden local symmetry model.


NODE=M001R15;LINKAGE=Q

13 From a model-dependent analysis assuming complete coherence.

NODE=M001R15;LINKAGE=F

14 Re-evaluated under $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ by BEHREND 71 using more accurate $\omega \rightarrow \rho$ photoproduction cross-section ratio.

NODE=M001R15;LINKAGE=B



$\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$

See also $\Gamma(\pi^+\pi^-)/\Gamma_{total}$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0172±0.0014 OUR FIT			Error includes scale factor of 1.2.
0.026 ±0.005 OUR AVERAGE			

0.021 +0.028 -0.009	1,2	RATCLIFF	72	ASPK	15 $\pi^- p \rightarrow n2\pi$
0.028 ±0.006	1	BEHREND	71	ASPK	Photoproduction
0.022 +0.009 -0.01	3	ROOS	70	RVUE	

¹ The fitted width of these data is 160 MeV in agreement with present average, thus the ω contribution is overestimated. Assuming ρ width 145 MeV.

² Significant interference effect observed. NB of $\omega \rightarrow 3\pi$ comes from an extrapolation.

³ ROOS 70 combines ABRAMOVICH 70 and BIZZARRI 70.

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^0\gamma)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.20±0.04	1.98M	¹ ALOISIO	03	KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$

¹ Using the data of ALOISIO 02D.

$\Gamma(\text{neutrals})/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.091±0.006 OUR FIT				
0.081±0.011 OUR AVERAGE				

0.075±0.025		BIZZARRI	71	HBC	0.0 $p\bar{p}$
0.079±0.019		DEINET	69B	OSPK	1.5 $\pi^- p$
0.084±0.015		BOLLINI	68C	CNTR	2.1 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.073±0.018	42	BASILE	72B	CNTR	1.67 $\pi^- p$

$\Gamma(\text{neutrals})/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.102±0.008 OUR FIT				
0.103^{+0.011}_{-0.010} OUR AVERAGE				

0.15 ±0.04	46	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
0.10 ±0.03	19	BARASH	67B	HBC	0.0 $\bar{p}p$
0.134±0.026	850	DIGIUGNO	66B	CNTR	1.4 $\pi^- p$
0.097±0.016	348	FLATTE	66	HBC	1.4 - 1.7 $K^- p \rightarrow \Lambda MM$
0.06 +0.05 -0.02		JAMES	66	HBC	2.1 $\pi^+ p$
0.08 ±0.03	35	KRAEMER	64	DBC	1.2 $\pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ±0.02	20	BUSCHBECK	63	HBC	1.5 $K^- p$

Γ_3/Γ_1

NODE=M001R2
NODE=M001R2
NODE=M001R2

Γ_3/Γ_2

NODE=M001R33
NODE=M001R33

$(\Gamma_2+\Gamma_4)/\Gamma$

NODE=M001R14
NODE=M001R14

$(\Gamma_2+\Gamma_4)/\Gamma_1$

NODE=M001R1
NODE=M001R1

$\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$ $\Gamma_2/(\Gamma_2+\Gamma_4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M001R18
 NODE=M001R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.78 ± 0.07		¹ DAKIN	72	OSPK	$1.4 \pi^- p \rightarrow nMM$
>0.81	90	DEINET	69B	OSPK	

¹ Error statistical only. Authors obtain good fit also assuming $\pi^0\gamma$ as the only neutral decay.

NODE=M001R18;LINKAGE=D

 $\Gamma(\text{neutrals})/\Gamma(\text{charged particles})$ $(\Gamma_2+\Gamma_4)/(\Gamma_1+\Gamma_3)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M001R9
 NODE=M001R9

0.100 ± 0.008 OUR FIT

0.124 ± 0.021

FELDMAN 67C OSPK 1.2 $\pi^- p$

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M001R19
 NODE=M001R19

4.5 ± 0.4 OUR FIT Error includes scale factor of 1.1. [(4.5 ± 0.4) × 10⁻⁴ OUR 2023 FIT Scale factor = 1.1]

NEW

6.3 ± 1.3 OUR AVERAGE Error includes scale factor of 1.2.

6.6 ± 1.7		¹ ABELE	97E	CBAR	$0.0 \bar{p} p \rightarrow 5\gamma$
8.3 ± 2.1		ALDE	93	GAM2	$38\pi^- p \rightarrow \omega n$
3.0 ^{+2.5} _{-1.8}		² ANDREWS	77	CNTR	6.7–10 γCu

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 ± 0.4 ± 0.1	33k	³ ACHASOV	07B	SND	0.6–1.38 $e^+ e^- \rightarrow \eta\gamma$
4.44 ^{+2.59} _{-1.83} ± 0.28	17.4k	^{4,5} AKHMETSHIN	05	CMD2	0.60–1.38 $e^+ e^- \rightarrow \eta\gamma$
5.10 ± 0.72 ± 0.34	23k	⁶ AKHMETSHIN	01B	CMD2	$e^+ e^- \rightarrow \eta\gamma$
0.7 to 5.5		⁷ CASE	00	CBAR	$0.0 p\bar{p} \rightarrow \eta\eta\gamma$
6.56 ^{+2.41} _{-2.55}	3525	^{2,8} BENAYOUN	96	RVUE	$e^+ e^- \rightarrow \eta\gamma$
7.3 ± 2.9		^{2,4} DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$

¹ No flat $\eta\eta\gamma$ background assumed.

² Solution corresponding to constructive $\omega\rho$ interference.

³ ACHASOV 07B reports $[\Gamma(\omega(782) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow e^+ e^-)] = (3.10 \pm 0.31 \pm 0.11) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow e^+ e^-) = (7.38 \pm 0.22) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁴ Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁵ Using $B(\omega \rightarrow e^+ e^-) = (7.14 \pm 0.13) \times 10^{-5}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁶ Using $B(\omega \rightarrow e^+ e^-) = (7.07 \pm 0.19) \times 10^{-5}$ and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$. Solution corresponding to constructive $\omega\rho$ interference. The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively). Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁷ Depending on the degree of coherence with the flat $\eta\eta\gamma$ background and using $B(\omega \rightarrow \pi^0\gamma) = (8.5 \pm 0.5) \times 10^{-2}$.

⁸ Reanalysis of DRUZHININ 84, DOLINSKY 89, DOLINSKY 91 taking into account the triangle anomaly contributions.

NODE=M001R;LINKAGE=EA
 NODE=M001R19;LINKAGE=A
 NODE=M001R19;LINKAGE=AO

NODE=M001R13;LINKAGE=WL
 NODE=M001R19;LINKAGE=AK
 NODE=M001R19;LINKAGE=TS

NODE=M001R;LINKAGE=CS

NODE=M001R19;LINKAGE=A1

 $\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$ Γ_5/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M001R11
 NODE=M001R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0098 ± 0.0024		¹ ALDE	93	GAM2	$38\pi^- p \rightarrow \omega n$
0.0082 ± 0.0033		² DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta\gamma$
0.010 ± 0.045		APEL	72B	OSPK	4–8 $\pi^- p \rightarrow n3\gamma$

¹ Model independent determination.

² Solution corresponding to constructive $\omega\rho$ interference.

NODE=M001R11;LINKAGE=A
 NODE=M001R11;LINKAGE=K

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M001R23
 NODE=M001R23

7.7 ± 0.6 OUR FIT

7.7 ± 0.6 OUR AVERAGE

7.61 ± 0.53 ± 0.64		ACHASOV	08	SND	0.36–0.97 $e^+ e^- \rightarrow \pi^0 e^+ e^-$
8.19 ± 0.71 ± 0.62		AKHMETSHIN	05A	CMD2	0.72–0.84 $e^+ e^-$
5.9 ± 1.9	43	DOLINSKY	88	ND	$e^+ e^- \rightarrow \pi^0 e^+ e^-$

$\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.34±0.18 OUR FIT Error includes scale factor of 1.5.**1.34±0.19 OUR AVERAGE** Error includes scale factor of 1.5. See the ideogram below.

1.41±0.09±0.15		ARNALDI	16	NA60	400 GeV (p -A) collisions
1.72±0.25±0.14	3k	ARNALDI	09	NA60	158A In-In collisions
0.96±0.23		DZHELADIN	81B	CNTR	25-33 $\pi^- p \rightarrow \omega n$

NODE=M001R12
 NODE=M001R12

WEIGHTED AVERAGE
 1.34±0.19 (Error scaled by 1.5)

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<1.1 AKHMETSHIN 05A CMD2 0.72-0.84 $e^+ e^-$

NODE=M001R34
 NODE=M001R34

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.738±0.022 OUR FIT Error includes scale factor of 1.9.

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.700±0.016	11200	1,2 AKHMETSHIN 04	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.752±0.004±0.024	1.2M	2,3 ACHASOV 03D	RVUE	0.44-2.00 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.714±0.036		2 DOLINSKY 89	ND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.72 ±0.03		2 BARKOV 87	CMD	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.64 ±0.04	1488	2 KURDADZE 83B	OLYA	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.675±0.069	433	2 CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.83 ±0.10	451	2 BENAKSAS 72B	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.77 ±0.06		4 AUGUSTIN 69D	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.65 ±0.13	33	5 ASTVACAT...	68	OSPK Assume SU(3)+mixing

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = 0.891 \pm 0.007$. Update of AKHMETSHIN 00C.² Not independent of the corresponding $\Gamma(e^+ e^-) \times \Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}^2$.³ Using ACHASOV 03, ACHASOV 03D and $B(\omega \rightarrow \pi^+ \pi^-) = (1.70 \pm 0.28)\%$.⁴ Rescaled by us to correspond to ω width 8.4 MeV. Systematic errors underestimated.⁵ Not resolved from ρ decay. Error statistical only.

NODE=M001R13;LINKAGE=4P
 NODE=M001R13;LINKAGE=ZL
 NODE=M001R13;LINKAGE=VF
 NODE=M001R13;LINKAGE=E
 NODE=M001R13;LINKAGE=A

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 2 90 ACHASOV 09A SND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<200 90 KURDADZE 86 OLYA $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

NODE=M001R5
 NODE=M001R5

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0036	95	WEIDENAUER 90	ASTE	$\rho\bar{p} \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.004	95	BITYUKOV 88B	SPEC	$32 \pi^- p \rightarrow \pi^+\pi^-\gamma X$

NODE=M001R22
NODE=M001R22

 $\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{11}/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.066	90	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$
<0.05	90	FLATTE 66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda\pi^+\pi^-\gamma$

NODE=M001R4
NODE=M001R4

 $\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 x 10⁻³	90	KURDADZE 88	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M001R24
NODE=M001R24

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

6.7±1.1 OUR FIT
6.5±1.2 OUR AVERAGE

$6.4^{+2.4}_{-2.0} \pm 0.8$	190	¹ AKHMETSHIN 04B	CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$
$6.6^{+1.4}_{-1.3} \pm 0.6$	295	ACHASOV 02F	SND	$0.36-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

$11.8^{+2.1}_{-1.9} \pm 1.4$	190	² AKHMETSHIN 04B	CMD2	$0.6-0.97 e^+e^- \rightarrow \pi^0\pi^0\gamma$
$7.8 \pm 2.7 \pm 2.0$	63	^{1,3} ACHASOV 00G	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$12.7 \pm 2.3 \pm 2.5$	63	^{2,3} ACHASOV 00G	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

OCCUR=2

OCCUR=2

¹ In the model assuming the $\rho \rightarrow \pi^0\pi^0\gamma$ decay via the $\omega\pi$ and $f_0(500)\gamma$ mechanisms.

² In the model assuming the $\rho \rightarrow \pi^0\pi^0\gamma$ decay via the $\omega\pi$ mechanism only.

³ Superseded by ACHASOV 02F.

NODE=M001R29;LINKAGE=A
NODE=M001R29;LINKAGE=B
NODE=M001R;LINKAGE=GF

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{13}/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.00045	90	DOLINSKY 89	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.08	95	JACQUET 69B	HLBC	$2.05 \pi^+ p \rightarrow \pi^+ p \omega$

NODE=M001R10
NODE=M001R10

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^0\gamma)$ Γ_{13}/Γ_2

VALUE (units 10 ⁻⁴)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

8.0±1.3 OUR FIT

8.5±2.9

40 ± 14		ALDE	94B	GAM2	$38\pi^- p \rightarrow \pi^0\pi^0\gamma n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 50	90	DOLINSKY 89	ND		$e^+e^- \rightarrow \pi^0\pi^0\gamma$
<1800	95	KEYNE 76	CNTR		$\pi^- p \rightarrow \omega n$
<1500	90	BENAKSAS 72C	OSPK		e^+e^-
<1400		BALDIN 71	HLBC		$2.9 \pi^+ p$
<1000	90	BARMIN 64	HLBC		$1.3-2.8 \pi^- p$

NODE=M001R7
NODE=M001R7

 $\Gamma(\pi^0\pi^0\gamma)/\Gamma(\text{neutrals})$ $\Gamma_{13}/(\Gamma_2+\Gamma_4)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.22 ± 0.07		¹ DAKIN 72	OSPK	$1.4 \pi^- p \rightarrow nMM$
<0.19	90	DEINET 69B	OSPK	

¹ See $\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$.

NODE=M001R17
NODE=M001R17

NODE=M001R17;LINKAGE=D

 $\Gamma(\eta\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

<3.3	90	AKHMETSHIN 04B	CMD2	$0.6-0.97 e^+e^- \rightarrow \eta\pi^0\gamma$
----------------	----	----------------	------	---

NODE=M001R32
NODE=M001R32

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-5})

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_{15}/Γ NODE=M001R30
NODE=M001R30**7.4±1.8 OUR FIT****7.4±1.8 OUR AVERAGE**

6.6±1.4±1.7	4.5M	¹ ANASTASI	17	KLOE	$e^+ e^- \rightarrow \mu^+ \mu^- \gamma$
9.0±2.9±1.1	18	HEISTER	02C	ALEP	$Z \rightarrow \mu^+ \mu^- + X$

¹ Assuming lepton universality in the decay $\omega \rightarrow \ell^+ \ell^-$ and correcting for different phase space between electron and muon final states.

NODE=M001R30;LINKAGE=A

 $\Gamma(\mu^+ \mu^-)/\Gamma(\pi^+ \pi^- \pi^0)$ VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{15}/Γ_1 NODE=M001R6
NODE=M001R6

<0.2	90	WILSON	69	OSPK	$12 \pi^- C \rightarrow Fe$
------	----	--------	----	------	-----------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7	74	FLATTE	66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \mu^+ \mu^-$
<1.2		BARBARO-...	65	HBC	$2.7 K^- p$

 $\Gamma(\pi^0 \mu^+ \mu^-)/\Gamma(\mu^+ \mu^-)$

VALUE

EVTS

DOCUMENT ID

TECN

COMMENT

 Γ_7/Γ_{15} NODE=M001R20
NODE=M001R20

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.2±0.6	30	¹ DZHELYADIN	79	CNTR	$25-33 \pi^- p$
---------	----	-------------------------	----	------	-----------------

¹ Superseded by DZHELYADIN 81B result above.

NODE=M001R20;LINKAGE=S

 $\Gamma(3\gamma)/\Gamma_{\text{total}}$ VALUE (units 10^{-4})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{16}/Γ NODE=M001R27
NODE=M001R27

<1.9	95	¹ ABELE	97E	CBAR	$0.0 \bar{p} p \rightarrow 5\gamma$
------	----	--------------------	-----	------	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2	90	¹ PROKOSHKIN	95	GAM2	$38 \pi^- p \rightarrow 3\gamma n$
----	----	-------------------------	----	------	------------------------------------

¹ From direct 3γ decay search.

NODE=M001R27;LINKAGE=A

 $\Gamma(\eta \pi^0)/\Gamma_{\text{total}}$

Violates C conservation.

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{17}/Γ NODE=M001R25
NODE=M001R25
NODE=M001R25

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.001	90	ALDE	94B	GAM2	$38\pi^- p \rightarrow \eta \pi^0 n$
--------	----	------	-----	------	--------------------------------------

 $[\Gamma(\eta \gamma) + \Gamma(\eta \pi^0)]/\Gamma(\pi^+ \pi^- \pi^0)$

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 $(\Gamma_5 + \Gamma_{17})/\Gamma_1$ NODE=M001R8
NODE=M001R8

<0.016	90	¹ FLATTE	66	HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \pi^+ \pi^- MM$
--------	----	---------------------	----	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.045	95	JACQUET	69B	HLBC	$2.05 \pi^+ p \rightarrow \pi^+ p \omega$
--------	----	---------	-----	------	---

¹ Restated by us using $B(\eta \rightarrow \text{charged modes}) = 29.2\%$.

NODE=M001R8;LINKAGE=A

 $\Gamma(\eta \pi^0)/\Gamma(\pi^0 \gamma)$

Violates C conservation.

VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{17}/Γ_2 NODE=M001R35
NODE=M001R35
NODE=M001R35

<2.6	90	¹ STAROSTIN	09	CRYM	$\gamma p \rightarrow \eta \pi^0 p$
------	----	------------------------	----	------	-------------------------------------

¹ STAROSTIN 09 reports $[\Gamma(\omega(782) \rightarrow \eta \pi^0)/\Gamma(\omega(782) \rightarrow \pi^0 \gamma)] \times [B(\eta \rightarrow 2\gamma)] < 1.01 \times 10^{-3}$ which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M001R35;LINKAGE=ST

 $\Gamma(2\pi^0)/\Gamma(\pi^0 \gamma)$

Violates C conservation and Bose-Einstein statistics.

VALUE (units 10^{-3})

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{18}/Γ_2 NODE=M001R36
NODE=M001R36
NODE=M001R36

<2.59	90	STAROSTIN	09	CRYM	$\gamma p \rightarrow 2\pi^0 p$
-------	----	-----------	----	------	---------------------------------

 $\Gamma(3\pi^0)/\Gamma_{\text{total}}$

Violates C conservation.

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

 Γ_{19}/Γ NODE=M001R26
NODE=M001R26
NODE=M001R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3×10^{-4}	90	PROKOSHKIN	95	GAM2	$38 \pi^- p \rightarrow 3\pi^0 n$
----------------------	----	------------	----	------	-----------------------------------

$\Gamma(3\pi^0)/\Gamma(\pi^0\gamma)$ Violates C conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.72	90	STAROSTIN 09	CRYM	$\gamma p \rightarrow 3\pi^0 p$

 Γ_{19}/Γ_2

NODE=M001R37
 NODE=M001R37
 NODE=M001R37

 $\Gamma(3\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Violates C conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	BARBERIS 01	450	$pp \rightarrow p_f 3\pi^0 p_s$

 Γ_{19}/Γ_1

NODE=M001R31
 NODE=M001R31
 NODE=M001R31

 $\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\pi^0)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.1 $\times 10^{-5}$	90	ABLIKIM 18S	BES3	$J/\psi \rightarrow \omega\eta \rightarrow \omega\pi^+\pi^-\pi^0$

 Γ_{20}/Γ_1

NODE=M001R01
 NODE=M001R01

PARAMETER Λ IN $\omega \rightarrow \pi^0 \ell^+ \ell^-$ DECAY

In the pole approximation the electromagnetic transition form factor for a resonance of mass M is given by the expression:

$$|F|^2 = (1 - M^2/\Lambda^2)^{-2},$$

where for the parameter Λ vector dominance predicts $\Lambda = M_p \approx 0.770$ GeV. The ARNALDI 09 measurement is in obvious conflict with this expectation. Note that for $\eta \rightarrow \gamma\mu^+\mu^-$ decay ARNALDI 09 and DZHELYADIN 80 obtain the value of Λ consistent with vector dominance.

NODE=M001230

NODE=M001230

PARAMETER Λ IN $\omega \rightarrow \pi^0 \mu^+ \mu^-$ DECAY

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.670 \pm 0.006 OUR AVERAGE

0.6707 \pm 0.0039 \pm 0.0056		¹ ARNALDI	16	NA60 400 GeV (p -A) collisions
0.668 \pm 0.009 \pm 0.003	3k	² ARNALDI	09	NA60 158A In-In collisions

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.65 \pm 0.03		DZHELYADIN 81B	CNTR	25-33 $\pi^- p \rightarrow \omega n$
-----------------	--	----------------	------	--------------------------------------

¹ ARNALDI 16 reports $\Lambda^{-2}(\omega) = 2.223 \pm 0.026 \pm 0.037$ GeV⁻² which we converted to the quoted Λ value.

² ARNALDI 09 reports $\Lambda^{-2}(\omega) = 2.24 \pm 0.06 \pm 0.02$ GeV⁻² which we converted to the quoted Λ value.

NODE=M001LAM
 NODE=M001LAM

NODE=M001LAM;LINKAGE=A

NODE=M001LAM;LINKAGE=B

PARAMETER Λ IN $\omega \rightarrow \pi^0 e^+ e^-$ DECAY

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.709 \pm 0.037	1.1k	¹ ADLARSON	17B	A2MM $\gamma p \rightarrow \omega p$
-------------------------------------	------	-----------------------	-----	--------------------------------------

¹ ADLARSON 17B reports $\Lambda^{-2}(\omega\pi^0) = 1.99 \pm 0.21$ GeV⁻² that we converted to the quoted Λ value.

NODE=M001A02
 NODE=M001A02

NODE=M001A02;LINKAGE=A

ENERGY DEPENDENCE OF $\omega \rightarrow \pi^+\pi^-\pi^0$ DALITZ PLOT

The following experiments fit to one or more of the coefficients α , β , γ for |matrix element|² $\propto P(1 + 2\alpha Z + 2\beta Z^{3/2} \sin(3\phi) + 2\gamma Z^2 + O(Z^{5/2}))$ where P is the P -wave phase-space factor and Z , ϕ are kinematical variables as defined in ADLARSON 17.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.133 \pm 0.008 OUR AVERAGE

0.1321 \pm 0.0067 \pm 0.0046	260k	¹ ABLIKIM	18AD	BES3 $J/\psi \rightarrow \omega\eta$
0.147 \pm 0.036	44k	ADLARSON 17	WASA	α in $pd \rightarrow {}^3\text{He } \omega$, $pp \rightarrow pp\omega$

¹ Keeping a term linear in Z only. A fit with the terms proportional to Z and $Z^{3/2}$ gives $\alpha = 0.133 \pm 0.041$ and $\beta = 0.037 \pm 0.054$.

NODE=M001A00

NODE=M001A00

NODE=M001A00

NODE=M001A00;LINKAGE=A

ω(782) REFERENCES

NODE=M001

ACHASOV	21	JHEP 2101 113	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61040
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
ANDREEV	20	EPJ C80 1189	V. Andreev <i>et al.</i>	(H1 Collab.)	REFID=60773
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)	REFID=61048
COLANGELO	19	JHEP 1902 006	G. Colangelo, M. Hoferichter, P. Stoffer	(BERN+)	REFID=59585
HOFRICHT...	19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)	REFID=60974
ABLIKIM	18AD	PR D98 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59498
ABLIKIM	18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58971
HANHART	18	EPJ C78 450	C. Hanhart <i>et al.</i>		REFID=59186
ADLARSON	17	PL B770 418	P. Adlarson <i>et al.</i>	(WASA-at-COSY Collab.)	REFID=57907
ADLARSON	17B	PR C95 035208	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)	REFID=58155
ANASTASI	17	PL B767 485	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)	REFID=57887
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57126
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57513
ARNALDI	16	PL B757 437	R. Arnaldi <i>et al.</i>	(NA60 Collab.)	REFID=57220
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977; ERROR=4
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
BABUSCI	13D	PL B720 336	D. Babusci <i>et al.</i>	(CATA, CALB, BARI)	REFID=55337
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)	REFID=55357
DAVIER	13	EPJ C73 2597	M. Davier <i>et al.</i>		REFID=55499
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54305
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
ACHASOV	09A	JETP 109 379	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=53101
		Translated from ZETF 136 442.			
ARNALDI	09	PL B677 260	R. Arnaldi <i>et al.</i>	(NA60 Collab.)	REFID=52720
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
STAROSTIN	09	PR C79 065201	A. Starostin <i>et al.</i>	(Crystal Ball Collab. at MAMI)	REFID=53001
ACHASOV	08	JETP 107 61	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=52258
		Translated from ZETF 134 80.			
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52573
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AULCHENKO	06	JETPL 84 413	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51513
		Translated from ZETFP 84 491.			
ACHASOV	05A	JETP 101 1053	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51045
		Translated from ZETF 128 1201.			
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AKHMETSHIN	05A	PL B613 29	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50508
AULCHENKO	05	JETPL 82 743	V.M. Aulchenko <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51060
		Translated from ZETFP 82 841.			
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AKHMETSHIN	04B	PL B580 119	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49610
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
ACHASOV	03	PL B559 171	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49187
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
BENAYOUN	03	EPJ C29 397	M. Benayoun <i>et al.</i>		REFID=49477
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
ACHASOV	02F	PL B537 201	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48816
AKHMETSHIN	02	PL B527 161	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48565
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48564
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48324
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
		Translated from ZETFP 72 411.			
ACHASOV	00G	JETPL 71 355	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47929
		Translated from ZETFP 71 519.			
AKHMETSHIN	00C	PL B476 33	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47423
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
		Translated from ZETF 117 1067.			
CASE	00	PR D61 032002	T. Case <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47409
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
GARDNER	99	PR D59 076002	S. Gardner, H.B. O'Connell		REFID=46919
BENAYOUN	98	EPJ C2 269	M. Benayoun <i>et al.</i>	(IPNP, NOVO, ADLD+)	REFID=45859
ABELE	97E	PL B411 361	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45755
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
PROKOSHKIN	95	PD 40 273	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=44616
		Translated from DANS 342 610.			
WURZINGER	95	PR C51 443	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=45209
ALDE	94B	PL B340 122	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=44100
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
ALDE	93	PAN 56 1229	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43603
		Translated from YAF 56 137.			
Also		ZPHY C61 35	D.M. Alde <i>et al.</i>	(SERP, LAPP, LANL, BELG+)	REFID=43790
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43602
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
WEIDENAUER	90	ZPHY C47 353	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=41368
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
BITYUKOV	88B	SJNP 47 800	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=41021
		Translated from YAF 47 1258.			
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41022
		Translated from YAF 48 442.			
KURDADZE	88	JETPL 47 512	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=41121
		Translated from ZETFP 47 432.			
AULCHENKO	87	PL B186 432	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=40007
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			

KURDADZE	86	JETPL 43 643	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=40287
		Translated from ZETFP 43 497.			
BARKOV	85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
KURDADZE	83B	JETPL 36 274	A.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20244
		Translated from ZETFP 36 221.			
DZHELYADIN	81B	PL 102B 296	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20242
CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
DZHELYADIN	80	PL 94B 548	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10831
ROOS	80	LNC 27 321	M. Roos, A. Pellinen	(HELS)	REFID=20241
BENKHEIRI	79	NP B150 268	P. Benkheiri <i>et al.</i>	(EPOL, CERN, CDEF+)	REFID=20238
DZHELYADIN	79	PL 84B 143	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=20239
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
QUENZER	78	PL 76B 512	A. Quenzer <i>et al.</i>	(LALO)	REFID=20123
VANAPEL...	78	NP B133 245	G.W. van Apeldoorn <i>et al.</i>	(ZEEM)	REFID=20234
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
KEYNE	76	PR D14 28	J. Keyne <i>et al.</i>	(LOIC, SHMP)	REFID=20226
Also		PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72B	PL 41B 234	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20206
BASILE	72B	Phil. Conf. 153	M. Basile <i>et al.</i>	(CERN)	REFID=20207
BENAKSAS	72	PL 39B 289	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20096
BENAKSAS	72B	PL 42B 507	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20209
BENAKSAS	72C	PL 42B 511	D. Benaksas <i>et al.</i>	(ORSAY)	REFID=20517
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
BROWN	72	PL 42B 117	R.M. Brown <i>et al.</i>	(ILL, ILLC)	REFID=20211
DAKIN	72	PR D6 2321	J.T. Dakin <i>et al.</i>	(PRIN)	REFID=20212
RATCLIFF	72	PL 38B 345	B.N. Ratcliff <i>et al.</i>	(SLAC)	REFID=20102
ALVENSLEB...	71C	PRL 27 888	H. Alvensleben <i>et al.</i>	(DESY)	REFID=20193
BALDIN	71	SJNP 13 758	A.B. Baldin <i>et al.</i>	(ITEP)	REFID=20195
		Translated from YAF 13 1318.			
BEHREND	71	PRL 27 61	H.J. Behrend <i>et al.</i>	(ROCH, CORN, FNAL)	REFID=20197
BIZZARRI	71	NP B27 140	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20198
COYNE	71	NP B32 333	D.G. Coyne <i>et al.</i>	(LRL)	REFID=20201
MOFFFEIT	71	NP B29 349	K.C. Moffeit <i>et al.</i>	(LRL, UCB, SLAC+)	REFID=20204
ABRAMOVI...	70	NP B20 209	M. Abramovich <i>et al.</i>	(CERN)	REFID=20180
BIGGS	70B	PRL 24 1201	P.J. Biggs <i>et al.</i>	(DARE)	REFID=20184
BIZZARRI	70	PRL 25 1385	R. Bizzarri <i>et al.</i>	(ROMA, SYRA)	REFID=20181
ROOS	70	DNPL/R7 173	M. Roos	(CERN)	REFID=20191
		Proc. Daresbury Study Weekend No. 1.			
AUGUSTIN	69D	PL 28B 513	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=20169
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)	REFID=20171
DEINET	69B	PL 30B 426	W. Deinet <i>et al.</i>	(KARL, CERN)	REFID=20173
JACQUET	69B	NC 63A 743	F. Jacquet <i>et al.</i>	(EPOL, BERG)	REFID=20176
WILSON	69	Private Comm.	R. Wilson	(HARV)	REFID=20179
Also		PR 178 2095	A.A. Wehmann <i>et al.</i>	(HARV, CASE, SLAC+)	REFID=20084
ASTVACAT...	68	PL 27B 45	R.G. Astvatsaturov <i>et al.</i>	(JINR, MOSU)	REFID=20055
BOLLINI	68C	NC 56A 531	D. Bollini <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20164
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
FELDMAN	67C	PR 159 1219	M. Feldman <i>et al.</i>	(PENN)	REFID=20161
DIGIUGNO	66B	NC 44A 1272	G. Di Giugno <i>et al.</i>	(NAPL, FRAS, TRST)	REFID=20156
FLATTE	66	PR 145 1050	S.M. Flatte <i>et al.</i>	(LRL)	REFID=20157
JAMES	66	PR 142 896	F.E. James, H.L. Kraybill	(YALE, BNL)	REFID=10770
BARBARO...	65	PRL 14 279	A. Barbaro-Galtieri, R.D. Tripp	(LRL)	REFID=20152
BARMIN	64	JETP 18 1289	V.V. Barmin <i>et al.</i>	(ITEP)	REFID=20149
		Translated from ZETF 45 1879.			
KRAEMER	64	PR 136 B496	R.W. Kraemer <i>et al.</i>	(JHU, NWES, WOOD)	REFID=10755
BUSCHBECK	63	Siena Conf. 1 166	B. Buschbeck <i>et al.</i>	(VIEN, CERN, ANIK)	REFID=20146

$\eta'(958)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

NODE=M002

 $\eta'(958)$ MASS

NODE=M002M

NODE=M002M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
957.78 ±0.06 OUR AVERAGE				
957.793±0.054±0.036	3.9k	LIBBY	08	CLEO $J/\psi \rightarrow \gamma\eta'$
957.9 ±0.2 ±0.6	4800	WURZINGER	96	SPEC 1.68 $p d \rightarrow {}^3\text{He}\eta'$
957.46 ±0.33		DUANE	74	MMS $\pi^- p \rightarrow n\text{MM}$
958.2 ±0.5	1414	DANBURG	73	HBC 2.2 $K^- p \rightarrow \Lambda\eta'$
958 ±1	400	JACOBS	73	HBC 2.9 $K^- p \rightarrow \Lambda\eta'$
956.1 ±1.1	3415	¹ BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
957.5 ±0.2		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
959 ±1	630	² BELADIDZE	92C	VES 36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
958 ±1	340	² ARMSTRONG	91B	OMEG 300 $pp \rightarrow pp\eta\pi^+\pi^-$
958.2 ±0.4	622	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
957.8 ±0.2	2420	² AUGUSTIN	90	DM2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
956.3 ±1.0	143	² GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
957.4 ±1.4	535	³ BASILE	71	CNTR 1.6 $\pi^- p \rightarrow n\eta'$
957 ±1		RITTENBERG	69	HBC 1.7-2.7 $K^- p$

OCCUR=2

OCCUR=2

¹ Using all η' decays.² Systematic uncertainty not estimated.³ Using η' decays into neutrals. Not independent of the other listed BASILE 71 η' mass measurement.

NODE=M002M;LINKAGE=BS

NODE=M002M;LINKAGE=NS

NODE=M002M;LINKAGE=BA

 $\eta'(958)$ WIDTH

NODE=M002W

NODE=M002W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.188±0.006 OUR FIT					
0.230±0.021 OUR AVERAGE					
0.226±0.017±0.014	2300	CZERWINSKI	10	MMS	$pp \rightarrow pp\eta'$
0.40 ±0.22	4800	WURZINGER	96	SPEC	1.68 $p d \rightarrow {}^3\text{He}\eta'$
0.28 ±0.10	1000	BINNIE	79	MMS	0 $\pi^- p \rightarrow n\text{MM}$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.20 ±0.04		BAI	04J	BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

 $\eta'(958)$ DECAY MODES

NODE=M002215;NODE=M002

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\pi^+\pi^-\eta$	(42.5 ±0.5)%	DESIG=1
Γ_2 $\rho^0\gamma$ (including non-resonant $\pi^+\pi^-\gamma$)	(29.5 ±0.4)%	DESIG=9
Γ_3 $\rho^0\gamma$		DESIG=213
Γ_4 $\pi^0\pi^0\eta$	(22.4 ±0.5)%	DESIG=2
Γ_5 $\omega\gamma$	(2.52 ±0.07)%	DESIG=7
Γ_6 ωe^+e^-	(2.0 ±0.4) × 10 ⁻⁴	DESIG=205
Γ_7 $\gamma\gamma$	(2.307±0.033)%	DESIG=6
Γ_8 $3\pi^0$	(2.50 ±0.17) × 10 ⁻³	DESIG=8
Γ_9 $\mu^+\mu^-\gamma$	(1.13 ±0.28) × 10 ⁻⁴	DESIG=20
Γ_{10} $\pi^+\pi^-\mu^+\mu^-$	(1.9 ±0.4) × 10 ⁻⁵	DESIG=201
Γ_{11} $\pi^+\pi^-\pi^0$	(3.61 ±0.17) × 10 ⁻³	DESIG=121
Γ_{12} $(\pi^+\pi^-\pi^0)$ S-wave	(3.8 ±0.5) × 10 ⁻³	DESIG=211
Γ_{13} $\pi^\mp\rho^\pm$	(7.4 ±2.3) × 10 ⁻⁴	DESIG=210
Γ_{14} $2(\pi^+\pi^-)$	(8.3 ±0.9) × 10 ⁻⁵	DESIG=131
Γ_{15} $\pi^+\pi^-2\pi^0$	(1.8 ±0.4) × 10 ⁻⁴	DESIG=202
Γ_{16} $2(\pi^+\pi^-)$ neutrals	< 1 %	95% DESIG=132

Γ_{17}	$2(\pi^+\pi^-\pi^0)$	< 1.8	$\times 10^{-3}$	90%	DESIG=141
Γ_{18}	$2(\pi^+\pi^-)2\pi^0$	< 1	%	95%	DESIG=15
Γ_{19}	$3(\pi^+\pi^-)$	< 3.1	$\times 10^{-5}$	90%	DESIG=203
Γ_{20}	$K^\pm\pi^\mp$	< 4	$\times 10^{-5}$	90%	DESIG=207
Γ_{21}	$\pi^+\pi^-e^+e^-$	(2.42 ± 0.10)	$\times 10^{-3}$		DESIG=10
Γ_{22}	$\pi^+e^-\nu_e + \text{c.c.}$	< 2.1	$\times 10^{-4}$	90%	DESIG=204
Γ_{23}	γe^+e^-	(4.91 ± 0.27)	$\times 10^{-4}$		DESIG=28
Γ_{24}	$\pi^0\gamma\gamma$	(3.20 ± 0.24)	$\times 10^{-3}$		DESIG=24
Γ_{25}	$\pi^0\gamma\gamma(\text{non resonant})$	(6.2 ± 0.9)	$\times 10^{-4}$		DESIG=212
Γ_{26}	$\eta\gamma\gamma$	< 1.33	$\times 10^{-4}$	90%	DESIG=214
Γ_{27}	$4\pi^0$	< 4.94	$\times 10^{-5}$	90%	DESIG=26
Γ_{28}	e^+e^-	< 5.6	$\times 10^{-9}$	90%	DESIG=150
Γ_{29}	$e^+e^-e^+e^-$	(4.5 ± 1.1)	$\times 10^{-6}$		DESIG=215
Γ_{30}	invisible	< 6	$\times 10^{-4}$	90%	DESIG=200

**Charge conjugation (C), Parity (P),
Lepton family number (LF) violating modes**

NODE=M002;CLUMP=B

Γ_{31}	$\pi^+\pi^-$	P, CP	< 1.8	$\times 10^{-5}$	90%	DESIG=111
Γ_{32}	$\pi^0\pi^0$	P, CP	< 4	$\times 10^{-4}$	90%	DESIG=25
Γ_{33}	$\pi^0e^+e^-$	C	[a] < 1.4	$\times 10^{-3}$	90%	DESIG=16
Γ_{34}	$\pi^0\rho^0$	C	< 4	%	90%	DESIG=18
Γ_{35}	ηe^+e^-	C	[a] < 2.4	$\times 10^{-3}$	90%	DESIG=17
Γ_{36}	3γ	C	< 1.0	$\times 10^{-4}$	90%	DESIG=23
Γ_{37}	$\mu^+\mu^-\pi^0$	C	[a] < 6.0	$\times 10^{-5}$	90%	DESIG=22
Γ_{38}	$\mu^+\mu^-\eta$	C	[a] < 1.5	$\times 10^{-5}$	90%	DESIG=21
Γ_{39}	$e\mu$	LF	< 4.7	$\times 10^{-4}$	90%	DESIG=27

[a] C parity forbids this to occur as a single-photon process.

LINKAGE=CS

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 20 branching ratios uses 52 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 69.5$ for 44 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-25							
x_4	-75	-43						
x_5	-7	-6	-2					
x_7	-11	-7	9	-1				
x_8	-17	-10	19	0	2			
x_{11}	-1	-1	-1	0	0	0		
x_{21}	-8	30	-14	-2	-2	-3	0	
Γ	11	-10	-1	1	-40	0	0	-3
	x_1	x_2	x_4	x_5	x_7	x_8	x_{11}	x_{21}

	Mode	Rate (MeV)	
Γ_1	$\pi^+\pi^-\eta$	0.0799 ± 0.0029	DESIG=1
Γ_2	$\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma)$	0.0554 ± 0.0019	DESIG=9
Γ_4	$\pi^0\pi^0\eta$	0.0421 ± 0.0017	DESIG=2
Γ_5	$\omega\gamma$	0.00474 ± 0.00020	DESIG=7
Γ_7	$\gamma\gamma$	0.00434 ± 0.00013	DESIG=6
Γ_8	$3\pi^0$	$(4.7 \pm 0.4) \times 10^{-4}$	DESIG=8
Γ_{11}	$\pi^+\pi^-\pi^0$	$(6.8 \pm 0.4) \times 10^{-4}$	DESIG=121
Γ_{21}	$\pi^+\pi^-e^+e^-$	$(4.54 \pm 0.23) \times 10^{-4}$	DESIG=10

$\eta'(958)$ PARTIAL WIDTHS

NODE=M002220

 $\Gamma(\gamma\gamma)$ Γ_7

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

4.34±0.14 OUR FIT**4.28±0.19 OUR AVERAGE**

4.17±0.10±0.27	2000	1 ACCIARRI	98Q L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.53±0.29±0.51	266	KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
3.61±0.13±0.48		2 BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\eta'(958)$
4.6 ±1.1 ±0.6	23	BARU	90 MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.57±0.25±0.44		BUTLER	90 MRK2	$e^+e^- \rightarrow e^+e^-\eta'(958)$
5.08±0.24±0.71	547	3 ROE	90 ASP	$e^+e^- \rightarrow e^+e^-2\gamma$
3.8 ±0.7 ±0.6	34	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.9 ±0.5 ±0.5	136	4 WILLIAMS	88 CBAL	$e^+e^- \rightarrow e^+e^-2\gamma$
4.7 ±0.6 ±0.9	143	5 GIDAL	87 MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.0 ±0.9		6 BARTEL	85E JADE	$e^+e^- \rightarrow e^+e^-2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ No non-resonant $\pi^+\pi^-$ contribution found.

² Reevaluated by us using $B(\eta' \rightarrow \rho(770)\gamma) = (30.2 \pm 1.3)\%$.

³ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁴ Reevaluated by us using $B(\eta' \rightarrow \gamma\gamma) = (2.11 \pm 0.13)\%$.

⁵ Superseded by BUTLER 90.

⁶ Systematic error not evaluated.

NODE=M002W4
NODE=M002W4NODE=M002W4;LINKAGE=AC
NODE=M002W4;LINKAGE=K1
NODE=M002W4;LINKAGE=K2
NODE=M002W4;LINKAGE=K3
NODE=M002W4;LINKAGE=C
NODE=M002W4;LINKAGE=A $\Gamma(e^+e^-)$ Γ_{28}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.1 × 10⁻³ 90 1,2 ACHASOV 15 SND 0.958 $e^+e^- \rightarrow \pi\pi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.0 × 10⁻³ 90 2 ACHASOV 15 SND 0.958 $e^+e^- \rightarrow \pi\pi\eta$

<2.4 × 10⁻³ 90 2 AKHMETSHIN 15 CMD3 0.958 $e^+e^- \rightarrow \pi^+\pi^-\eta$

¹ Combining data of ACHASOV 15 and AKHMETSHIN 15.

² Using η and η' branching fractions from PDG 14.

NODE=M002W1
NODE=M002W1

OCCUR=2

NODE=M002W1;LINKAGE=A
NODE=M002W1;LINKAGE=B $\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

NODE=M002223

NODE=M002223

 $\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_2/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1.28±0.04 OUR FIT**1.26±0.07 OUR AVERAGE** Error includes scale factor of 1.2.

1.09±0.04±0.13		BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\rho(770)^0\gamma$
1.35±0.09±0.21		AIHARA	87 TPC	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.13±0.04±0.13	867	ALBRECHT	87B ARG	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.53±0.09±0.21		ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.14±0.08±0.11	243	BERGER	84B PLUT	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.73±0.34±0.35	95	JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.49±0.13±0.027	213	BARTEL	82B JADE	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.85±0.31±0.24	43	BEHREND	82C CELL	$e^+e^- \rightarrow e^+e^-\rho\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M002G1
NODE=M002G1 $\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_4/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

0.97±0.04 OUR FIT Error includes scale factor of 1.1.**0.92±0.06±0.11** ¹ KARCH 92 CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.95±0.05±0.08 ² KARCH 90 CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$

1.00±0.08±0.10 ^{2,3} ANTREASYAN 87 CBAL $e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$

¹ Reevaluated by us using $B(\eta \rightarrow \gamma\gamma) = (39.21 \pm 0.34)\%$. Supersedes ANTREASYAN 87 and KARCH 90.

² Superseded by KARCH 92.

³ Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.

NODE=M002G2
NODE=M002G2

NODE=M002G2;LINKAGE=K4

NODE=M002G2;LINKAGE=A
NODE=M002G2;LINKAGE=D

$\eta'(958) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M002224

 $\Gamma(\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{28}/\Gamma$

VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	¹ AKHMETSHIN 15	CMD3	$0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$
¹ AKHMETSHIN 15 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta) \times \Gamma(\eta'(958) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] < 4.1 \times 10^{-4}$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.				

NODE=M002G01
NODE=M002G01

NODE=M002G01;LINKAGE=A

 $\eta'(958) \text{ BRANCHING RATIOS}$

NODE=M002230

 $\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
42.5 ± 0.5 OUR FIT	Error includes scale factor of 1.1.			
41.24 ± 0.08 ± 1.24	312k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
42.4 ± 1.1 ± 0.4	1.2k	¹ PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.				

NODE=M002R47
NODE=M002R47

NODE=M002R47;LINKAGE=PE

 $\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{\text{total}}$ **0.2804** Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1191 ± 0.0015 OUR FIT	Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.123 ± 0.014	107	RITTENBERG	69 HBC	$1.7-2.7 K^-p$
0.10 ± 0.04	10	LONDON	66 HBC	$2.24 K^-p \rightarrow \Lambda 2\pi^+ 2\pi^- \pi^0$
0.07 ± 0.04	7	BADIER	65B HBC	$3 K^-p$

NODE=M002R3
NODE=M002R3 $\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{\text{total}}$ **0.7196** Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.306 ± 0.004 OUR FIT	Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.314 ± 0.026	281	RITTENBERG	69 HBC	$1.7-2.7 K^-p$

NODE=M002R1
NODE=M002R1 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
29.5 ± 0.4 OUR FIT	Error includes scale factor of 1.1.			
29.90 ± 0.03 ± 0.55	913k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
28.7 ± 0.7 ± 0.4	0.2k	¹ PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma\eta'$
32.9 ± 3.3	298	RITTENBERG	69 HBC	$1.7-2.7 K^-p$
20 ± 10	20	LONDON	66 HBC	$2.24 K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$
34 ± 9	35	BADIER	65B HBC	$3 K^-p$
¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.				

NODE=M002R6
NODE=M002R6

NODE=M002R6;LINKAGE=PE

 $\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
33.34 ± 0.06 ± 1.60	970k	¹ ABLIKIM	18C BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
34.43 ± 0.52 ± 1.97	970k	² ABLIKIM	18C BES3	$\eta'(958) \rightarrow \gamma\pi^+\pi^-$
¹ From a fit to $\pi^+\pi^-$ mass using $\rho(770)$, $\omega(782)$, and box anomaly components.				
² From a fit to $\pi^+\pi^-$ mass using $\rho(770)$, $\omega(782)$, and $\rho(1450)$ components.				

NODE=M002R66
NODE=M002R66

OCCUR=2

NODE=M002R66;LINKAGE=A

NODE=M002R66;LINKAGE=B

 $\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))/\Gamma(\pi^+\pi^-\eta)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.694 ± 0.014 OUR FIT	Error includes scale factor of 1.1.		
0.683 ± 0.020 OUR AVERAGE			
0.677 ± 0.024 ± 0.011	PEDLAR	09 CLE3	$J/\psi \rightarrow \eta'\gamma$
0.69 ± 0.03	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta'\gamma$

NODE=M002R43
NODE=M002R43

$$\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi^+ \pi^- \eta (\text{neutral decay}))$$
 $\Gamma_2/0.714\Gamma_1$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.972 ± 0.020 OUR FIT	Error includes scale factor of 1.1.			
0.97 ± 0.09 OUR AVERAGE				
0.70 ± 0.22		AMSLER	04B CBAR	0 $\bar{p}p \rightarrow \pi^+ \pi^- \eta$
1.07 ± 0.17		BELADIDZE	92C VES	36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
0.92 ± 0.14	473	DANBURG	73 HBC	2.2 $K^- p \rightarrow \Lambda X^0$
1.11 ± 0.18	192	JACOBS	73 HBC	2.9 $K^- p \rightarrow \Lambda X^0$

NODE=M002R27
 NODE=M002R27

$$\Gamma(\pi^0 \pi^0 \eta) / \Gamma_{\text{total}}$$
 Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
22.4 ± 0.6 OUR FIT	Error includes scale factor of 1.1.			
21.36 ± 0.10 ± 0.92	52k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
23.5 ± 1.3 ± 0.4	3.2k	¹ PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma \eta'$

NODE=M002R48
 NODE=M002R48

¹ Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R48;LINKAGE=PE

$$\Gamma(\pi^0 \pi^0 \eta (3\pi^0 \text{ decay})) / \Gamma_{\text{total}}$$
 $0.321\Gamma_4/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0718 ± 0.0018 OUR FIT	Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.11 ± 0.06	4	BENSINGER	70 DBC	2.2 $\pi^+ d$

NODE=M002R26
 NODE=M002R26

$$\Gamma(\pi^0 \pi^0 \eta) / \Gamma(\pi^+ \pi^- \eta)$$
 Γ_4/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.527 ± 0.019 OUR FIT	Error includes scale factor of 1.1.		
0.555 ± 0.043 ± 0.013	PEDLAR	09 CLE3	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R45
 NODE=M002R45

$$\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / \Gamma(\pi \pi \eta)$$
 $\Gamma_2/(\Gamma_1 + \Gamma_4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.454 ± 0.009 OUR FIT	Error includes scale factor of 1.1.		
0.43 ± 0.02 ± 0.02	BARBERIS	98C OMEG	450 $p p \rightarrow p_f \eta' p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.31 ± 0.15	DAVIS	68 HBC	5.5 $K^- p$

NODE=M002R7
 NODE=M002R7

$$\Gamma(\omega \gamma) / \Gamma_{\text{total}}$$
 Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.52 ± 0.07 OUR FIT				
2.50 ± 0.07 OUR AVERAGE				
2.489 ± 0.018 ± 0.074	23k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma \eta'$
2.55 ± 0.03 ± 0.16	33.2k	¹ ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta' \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.34 ± 0.30 ± 0.04	70	² PEDLAR	09 CLEO	$J/\psi \rightarrow \gamma \eta'$

NODE=M002R49
 NODE=M002R49

¹ Using $B(J/\psi \rightarrow \eta' \gamma) = (5.15 \pm 0.16) \times 10^{-3}$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

² Not independent of other η' branching fractions and ratios in PEDLAR 09.

NODE=M002R49;LINKAGE=A
 NODE=M002R49;LINKAGE=PE

$$\Gamma(\omega \gamma) / \Gamma(\pi^+ \pi^- \eta)$$
 Γ_5/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0593 ± 0.0018 OUR FIT	Error includes scale factor of 1.1.			
0.055 ± 0.007 ± 0.001		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta' \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.068 ± 0.013	68	ZANFINO	77 ASPK	8.4 $\pi^- p$

NODE=M002R17
 NODE=M002R17

$$\Gamma(\omega \gamma) / \Gamma(\pi^0 \pi^0 \eta)$$
 Γ_5/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.113 ± 0.004 OUR FIT			
0.147 ± 0.016	ALDE	87B GAM2	38 $\pi^- p \rightarrow n 4\gamma$

NODE=M002R33
 NODE=M002R33

$$\Gamma(\omega e^+ e^-) / \Gamma(\omega \gamma)$$
 Γ_6/Γ_5

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
7.71 ± 1.34 ± 0.54	¹ ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R60
 NODE=M002R60

¹ Obtained from other ABLIKIM 15AD measurements with common systematics taken into account.

NODE=M002R60;LINKAGE=A

$\Gamma(\omega e^+ e^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.97±0.34±0.17	66	¹ ABLIKIM	15AD BES3	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R59
 NODE=M002R59

¹ Using $B(J/\psi \rightarrow \eta' \gamma) = (5.15 \pm 0.16) \times 10^{-3}$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

NODE=M002R59;LINKAGE=A

$$\frac{\Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma)) / [\Gamma(\pi^+ \pi^- \eta) + \Gamma(\pi^0 \pi^0 \eta) + \Gamma(\omega \gamma)]}{\Gamma_2 / (\Gamma_1 + \Gamma_4 + \Gamma_5)}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.437±0.008 OUR FIT			Error includes scale factor of 1.1.

NODE=M002R18
 NODE=M002R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.25 ± 0.14	DAUBER	64	HBC	1.95 $K^- p$
-------------	--------	----	-----	--------------

$$\frac{[\Gamma(\pi^0 \pi^0 \eta (\text{charged decay})) + \Gamma(\omega (\text{charged decay}) \gamma)] / \Gamma_{\text{total}}}{(0.286\Gamma_4 + 0.89\Gamma_5) / \Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0864±0.0017 OUR FIT				Error includes scale factor of 1.1.

NODE=M002R4
 NODE=M002R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.045 ± 0.029	42	RITTENBERG	69	HBC	1.7–2.7 $K^- p$
---------------	----	------------	----	-----	-----------------

$$\frac{\Gamma(\pi^+ \pi^- \text{ neutrals}) / \Gamma_{\text{total}}}{(0.714\Gamma_1 + 0.286\Gamma_4 + 0.89\Gamma_5) / \Gamma}$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3897±0.0028 OUR FIT				Error includes scale factor of 1.1.

NODE=M002R2
 NODE=M002R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 ± 0.1	39	LONDON	66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \text{ neutrals}$
0.35 ± 0.06	33	BADIER	65B	HBC	3 $K^- p$

$$\frac{\Gamma(\gamma \gamma) / \Gamma_{\text{total}}}{\Gamma_7 / \Gamma}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.307±0.035 OUR FIT				Error includes scale factor of 1.1.

NODE=M002R19
 NODE=M002R19

2.31 ± 0.06 OUR AVERAGE Error includes scale factor of 1.8.

2.331 ± 0.012 ± 0.035	71k	ABLIKIM	19T	BES	$J/\psi \rightarrow \gamma \eta'$
1.99 $^{+0.31}_{-0.27}$ ± 0.07	114	¹ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma \gamma$
2.00 ± 0.18		² STANTON	80	SPEC	$8.45 \pi^- p \rightarrow n \pi^+ \pi^- 2\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.25 ± 0.16 ± 0.03	0.3k	³ PEDLAR	09	CLEO	$J/\psi \rightarrow \gamma \eta'$
1.8 ± 0.2	6000	⁴ APEL	79	NICE	$15-40 \pi^- p \rightarrow n 2\gamma$
2.5 ± 0.7		DUANE	74	MMS	$\pi^- p \rightarrow n \text{MM}$
1.71 ± 0.33	68	DALPIAZ	72	CNTR	$1.6 \pi^- p \rightarrow n X^0$
2.0 $^{+0.8}_{-0.6}$	31	HARVEY	71	OSPK	$3.65 \pi^- p \rightarrow n X^0$

¹ WICHT 08 reports $[\Gamma(\eta'(958) \rightarrow \gamma \gamma) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta' K^+)] = (1.40^{+0.16+0.15}_{-0.15-0.12}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta' K^+) = (7.04 \pm 0.25) \times 10^{-5}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Includes APEL 79 result.

³ Not independent of other η' branching fractions and ratios in PEDLAR 09.

⁴ Data is included in STANTON 80 evaluation.

NODE=M002R19;LINKAGE=WI

NODE=M002R19;LINKAGE=S
 NODE=M002R19;LINKAGE=PE
 NODE=M002R19;LINKAGE=A

$$\frac{\Gamma(\gamma \gamma) / \Gamma(\pi^+ \pi^- \eta)}{\Gamma_7 / \Gamma_1}$$

VALUE	DOCUMENT ID	TECN	COMMENT	
0.0543±0.0012 OUR FIT			Error includes scale factor of 1.1.	
0.053 ± 0.004 ± 0.001	PEDLAR	09	CLE3	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R46
 NODE=M002R46

$$\frac{\Gamma(\gamma \gamma) / \Gamma(\rho^0 \gamma (\text{including non-resonant } \pi^+ \pi^- \gamma))}{\Gamma_7 / \Gamma_2}$$

VALUE	DOCUMENT ID	TECN	COMMENT	
0.0783±0.0016 OUR FIT			Error includes scale factor of 1.1.	
0.080 ± 0.008	ABLIKIM	06E	BES2	$J/\psi \rightarrow \eta' \gamma$

NODE=M002R42
 NODE=M002R42

$$\frac{\Gamma(\gamma \gamma) / \Gamma(\pi^0 \pi^0 \eta)}{\Gamma_7 / \Gamma_4}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.1031±0.0028 OUR FIT			
0.105 ± 0.010 OUR AVERAGE			Error includes scale factor of 1.9.

NODE=M002R38
 NODE=M002R38

0.091 ± 0.009	AMSLER	93	CBAR	0.0 $\bar{p} p$
0.112 ± 0.002 ± 0.006	ALDE	87B	GAM2	38 $\pi^- p \rightarrow n 2\gamma$

$\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta(\text{neutral decay}))$ $\Gamma_7/0.714\Gamma_4$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M002R28
 NODE=M002R28

0.144 ± 0.004 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.188 ± 0.058	16	APEL	72	OSPK	$3.8 \pi^- p \rightarrow n\chi^0$
---------------	----	------	----	------	-----------------------------------

 $\Gamma(\text{neutrals})/\Gamma_{\text{total}}$ $(0.714\Gamma_4 + 0.09\Gamma_5 + \Gamma_7)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M002R5
 NODE=M002R5

0.185 ± 0.004 OUR FIT Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.185 ± 0.022	535	BASILE	71	CNTR	$1.6 \pi^- p \rightarrow n\chi^0$
0.189 ± 0.026	123	RITTENBERG	69	HBC	$1.7-2.7 K^- p$

 $\Gamma(3\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R55
 NODE=M002R55

2.50 ± 0.17 OUR FIT**3.57 ± 0.26 OUR AVERAGE**

3.522 ± 0.082 ± 0.254	2015	ABLIKIM	17	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
4.79 ± 0.59 ± 1.14	183	¹ ABLIKIM	15P	BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.56 ± 0.22 ± 0.34	309	² ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(3\pi^0)$
--------------------	-----	----------------------	-----	------	-------------------------------------

¹We have added all systematic uncertainties in quadrature to a single value.

²Superseded by ABLIKIM 17.

NODE=M002R55;LINKAGE=A
 NODE=M002R55;LINKAGE=B

 $\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_8/Γ_4

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R32
 NODE=M002R32

112 ± 8 OUR FIT**78 ± 10 OUR AVERAGE**

86 ± 19	235	BLIK	08	GAMS	$32 \pi^- p \rightarrow \eta' n$
74 ± 15		ALDE	87B	GAM2	$38 \pi^- p \rightarrow n6\gamma$
75 ± 18		BINON	84	GAM2	$30-40 \pi^- p \rightarrow n6\gamma$

 $\Gamma(\mu^+\mu^-\gamma)/\Gamma(\gamma\gamma)$ Γ_9/Γ_7

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M002R29
 NODE=M002R29

4.9 ± 1.2	33	VIKTOROV	80	CNTR	$25,33 \pi^- p \rightarrow 2\mu\gamma$
------------------	----	----------	----	------	--

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M002R50
 NODE=M002R50

1.9 ± 0.4 OUR AVERAGE [(2.0 ± 0.4) × 10⁻⁵ OUR 2023 AVERAGE]

1.94 ± 0.37 ± 0.02	53	¹ ABLIKIM	21I	BES3	$J/\psi \rightarrow \gamma\eta'(958)$
---------------------------	----	----------------------	-----	------	---------------------------------------

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.9	90	² ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
< 24	90	³ NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$

¹ABLIKIM 21I reports $(1.97 \pm 0.33 \pm 0.19) \times 10^{-5}$ from a measurement of $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$ assuming $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.21 \pm 0.17) \times 10^{-3}$, which we rescale to our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.28 \pm 0.06) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R50;LINKAGE=E

²Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

NODE=M002R50;LINKAGE=A

³Not independent of measured value of Γ_{10}/Γ_1 from NAIK 09.

NODE=M002R50;LINKAGE=NA

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{10}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R03
 NODE=M002R03

< 0.5	90	¹ NAIK	09	CLEO	$J/\psi \rightarrow \gamma\eta'$
-----------------	----	-------------------	----	------	----------------------------------

¹NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\mu^+\mu^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 1.3 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R03;LINKAGE=NA

 $\Gamma(\pi^+\pi^-\mu^+\mu^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{10}/Γ_2

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M002R57
 NODE=M002R57

< 1.0	90	ABLIKIM	130	BES3	$J/\psi \rightarrow \gamma\eta'$
-----------------	----	---------	-----	------	----------------------------------

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.61 ± 0.18 OUR FIT				
3.61 ± 0.18 OUR AVERAGE				
3.591 ± 0.054 ± 0.174	6067	ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
4.28 ± 0.49 ± 1.11	78	¹ ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$
3.7 $^{+1.1}_{-0.9}$ ± 0.4		² NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.83 ± 0.15 ± 0.39 1014 ³ ABLIKIM 12E BES3 $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$

¹ We have added all systematic uncertainties in quadrature to a single value.

² Not independent of measured value of Γ_{11}/Γ_1 from NAIK 09.

³ Superseded by ABLIKIM 17.

NODE=M002R21
NODE=M002R21

NODE=M002R21;LINKAGE=A
NODE=M002R21;LINKAGE=NA
NODE=M002R21;LINKAGE=B

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{11}/Γ_1

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5 ± 0.4 OUR FIT				Error includes scale factor of 1.1.
8.27 $^{+2.49}_{-2.12}$ ± 0.04	20	¹ NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (21 $^{+6}_{-5}$ ± 2) × 10⁻³ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.36 ± 0.18) × 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.$

NODE=M002R01
NODE=M002R01

NODE=M002R01;LINKAGE=NA

 $\Gamma((\pi^+\pi^-\pi^0) S\text{-wave})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
37.63 ± 0.77 ± 5.00	6580	¹ ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$

¹ We have added all systematic uncertainties in quadrature .

NODE=M002R63
NODE=M002R63

NODE=M002R63;LINKAGE=A

 $\Gamma(\pi^\mp\rho^\pm)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.44 ± 0.60 ± 2.23	1231	¹ ABLIKIM	17 BES3	$J/\psi \rightarrow \gamma(\pi^\mp\rho^\pm)$

¹ We have added all systematic uncertainties in quadrature .

NODE=M002R62
NODE=M002R62

NODE=M002R62;LINKAGE=A

 $\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
8.3 ± 0.9 OUR AVERAGE					[(8.4 ± 0.9) × 10 ⁻⁵ OUR 2023 AVERAGE]
8.3 ± 0.9 ± 0.1	199	¹ ABLIKIM	14M BES3		$J/\psi \rightarrow \gamma\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 24 90 ² NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$
<1000 90 RITTENBERG 69 HBC 1.7-2.7 $K^-\rho$

¹ ABLIKIM 14M reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}}] × [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$ = (4.40 ± 0.35 ± 0.30) × 10⁻⁷ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.28 ± 0.06) × 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Not independent of measured value of Γ_{14}/Γ_1 from NAIK 09.

NODE=M002R24
NODE=M002R24

NEW

NODE=M002R24;LINKAGE=A

NODE=M002R24;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{14}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.6	90	¹ NAIK	09 CLEO	$J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ < 1.4 × 10⁻³ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.36 × 10^{-2}$.

NODE=M002R04
NODE=M002R04

NODE=M002R04;LINKAGE=NA

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.78 ± 0.38 ± 0.02	84	¹ ABLIKIM	14M BES3		$J/\psi \rightarrow \gamma\eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<27 90 ² NAIK 09 CLEO $J/\psi \rightarrow \gamma\eta'$

¹ ABLIKIM 14M reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}] × [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$ = (9.38 ± 1.79 ± 0.89) × 10⁻⁷ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.28 ± 0.06) × 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Not independent of measured value of Γ_{15}/Γ_1 from NAIK 09.

NODE=M002R51
NODE=M002R51

NODE=M002R51;LINKAGE=A

NODE=M002R51;LINKAGE=NA

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{15}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

NODE=M002R05
NODE=M002R05

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-2\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 15 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R05;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-) \text{ neutrals})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	DANBURG	73	HBC $2.2 K^-p \rightarrow \Lambda X^0$

NODE=M002R22
NODE=M002R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01	90	RITTENBERG	69	HBC $1.7-2.7 K^-p$
-------	----	------------	----	--------------------

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.002	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

NODE=M002R23
NODE=M002R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01	90	RITTENBERG	69	HBC $1.7-2.7 K^-p$
-------	----	------------	----	--------------------

¹ Not independent of measured value of Γ_{17}/Γ_1 from NAIK 09.

NODE=M002R23;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma(\pi^+\pi^-\eta)$ Γ_{17}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

NODE=M002R06
NODE=M002R06

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 11 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R06;LINKAGE=NA

 $\Gamma(2(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	KALBFLEISCH	64B	HBC $K^-p \rightarrow \Lambda 2(\pi^+\pi^-)+MM$

NODE=M002R16
NODE=M002R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.01	90	LONDON	66	HBC Compilation
-------	----	--------	----	-----------------

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	¹ ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

NODE=M002R07
NODE=M002R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 53	90	² NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
<500	95	KALBFLEISCH	64B	HBC $K^-p \rightarrow \Lambda 2(\pi^+\pi^-)$

¹ Using $B(J/\psi \rightarrow \gamma\eta'(958)) = (5.16 \pm 0.15) \times 10^{-3}$.

² Not independent of measured value of Γ_{19}/Γ_1 from NAIK 09.

NODE=M002R07;LINKAGE=A
NODE=M002R07;LINKAGE=NA $\Gamma(3(\pi^+\pi^-))/\Gamma(\pi^+\pi^-\eta)$ Γ_{19}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

NODE=M002R08
NODE=M002R08

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow 3(\pi^+\pi^-))/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 3.0 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R08;LINKAGE=NA

 $\Gamma(K^\pm\pi^\mp)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{20}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 $\times 10^{-4}$	90	ABLIKIM	16M	BES3 $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

NODE=M002R61
NODE=M002R61 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.42 ± 0.10 OUR FIT					

NODE=M002R12
NODE=M002R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.11 ± 0.12 ± 0.14	429	¹ ABLIKIM	130	BES3 $J/\psi \rightarrow \gamma\eta'$
2.5 $^{+1.2}_{-0.9}$ ± 0.5		² NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
<6	90	RITTENBERG	65	HBC $2.7 K^-p$

¹ Using $\Gamma_2/\Gamma = (29.3 \pm 0.6)\%$ from PDG 12.

² Not independent of measured value of Γ_{21}/Γ_1 from NAIK 09.

NODE=M002R12;LINKAGE=A
NODE=M002R12;LINKAGE=NA

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\pi^+\pi^-\eta)$ Γ_{21}/Γ_1

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.69±0.25 OUR FIT

5.51^{+3.00}_{-2.30}±0.03	8	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$
--	---	-------------------	----	---------------------------------------

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-e^+e^-)/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] = (14^{+7}_{-5} \pm 3) \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R02
NODE=M002R02

NODE=M002R02;LINKAGE=NA

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma(\rho^0\gamma(\text{including non-resonant } \pi^+\pi^-\gamma))$ Γ_{21}/Γ_2

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.20±0.31 OUR FIT

8.20±0.16±0.27	2584	ABLIKIM	21J	BES3 $J/\psi \rightarrow \gamma\eta'$
-----------------------	------	---------	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 0.4 ± 0.5	429	¹ ABLIKIM	130	BES3 $J/\psi \rightarrow \gamma\eta'$
-----------------	-----	----------------------	-----	---------------------------------------

¹ Superseded by ABLIKIM 21J.NODE=M002R56
NODE=M002R56

NODE=M002R56;LINKAGE=A

 $\Gamma(\pi^+e^-\nu_e + \text{c.c.})/\Gamma(\pi^+\pi^-\eta)$ Γ_{22}/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<5.0	90	ABLIKIM	13G	BES3 $J/\psi \rightarrow \phi\eta'$
----------------	----	---------	-----	-------------------------------------

NODE=M002R54
NODE=M002R54 $\Gamma(\gamma e^+e^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.9	90	BRIERE	00	CLEO $10.6 e^+e^-$
------	----	--------	----	--------------------

NODE=M002R40
NODE=M002R40 $\Gamma(\gamma e^+e^-)/\Gamma(\gamma\gamma)$ Γ_{23}/Γ_7

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.13±0.09±0.07	864	ABLIKIM	150	BES3 $J/\psi \rightarrow \gamma e^+e^-$
-----------------------	-----	---------	-----	---

NODE=M002R00
NODE=M002R00 $\Gamma(\pi^0\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.20±0.07±0.23	3.4k	ABLIKIM	17T	BES3 $J/\psi \rightarrow \gamma\eta'$
-----------------------	------	---------	-----	---------------------------------------

NODE=M002R64
NODE=M002R64 $\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_{24}/Γ_4

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<37	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n4\gamma$
---------------	----	------	-----	--

NODE=M002R35
NODE=M002R35 $\Gamma(\pi^0\gamma\gamma(\text{non resonant}))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.16±0.64±0.67	655	ABLIKIM	17T	BES3 $J/\psi \rightarrow \gamma\eta'$
-----------------------	-----	---------	-----	---------------------------------------

NODE=M002R65
NODE=M002R65 $\Gamma(\eta\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.33 × 10⁻⁴	90	ABLIKIM	19AW	BES3 $J/\psi \rightarrow \gamma\eta' \rightarrow \gamma\gamma\gamma 2\gamma$
-----------------------------------	----	---------	------	--

NODE=M002R67
NODE=M002R67 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<4.94 × 10⁻⁵	90	ABLIKIM	20E	BES3 $J/\psi \rightarrow \eta'\gamma$
-----------------------------------	----	---------	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.2 × 10 ⁻⁴	90	DONSKOV	14	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
-------------------------	----	---------	----	---

NODE=M002R58
NODE=M002R58 $\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{27}/Γ_4

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<23	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n8\gamma$
-----	----	------	-----	--

NODE=M002R37
NODE=M002R37

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-9}$	90	¹ ACHASOV	15	SND $0.958 e^+e^- \rightarrow \pi\pi\eta$
$< 12 \times 10^{-9}$	90	² AKHMETSHIN	15	CMD3 $0.958 e^+e^- \rightarrow \pi^+\pi^-\eta$
$< 2.1 \times 10^{-7}$	90	VOROBYEV	88	ND $e^+e^- \rightarrow \pi^+\pi^-\eta$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Combining data of ACHASOV 15 and AKHMETSHIN 15 and using $\Gamma(\eta') = 0.198 \pm 0.009$ MeV.

² Using $\Gamma_{\eta'(958)} = 198 \pm 9$ keV, $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.9 \pm 0.7)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.20)\%$.

NODE=M002R39
NODE=M002R39

NODE=M002R39;LINKAGE=B

NODE=M002R39;LINKAGE=A

 $\Gamma(e^+e^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-6})	EVT5	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.1 \pm 0.1$	30	¹ ABLIKIM	22E	BES3 $J/\psi \rightarrow \gamma\eta'$

¹ ABLIKIM 22E reports $(4.5 \pm 1.0 \pm 0.5) \times 10^{-6}$ from a measurement of $[\Gamma(\eta'(958) \rightarrow e^+e^-e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))]$ assuming $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.25 \pm 0.07) \times 10^{-3}$, which we rescale to our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = (5.28 \pm 0.06) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M002R68
NODE=M002R68

NODE=M002R68;LINKAGE=B

 $\Gamma(\text{invisible})/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 9.5	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

¹ Not independent of measured value of Γ_{30}/Γ_1 from NAIK 09.

NODE=M002R52
NODE=M002R52

NODE=M002R52;LINKAGE=NA

 $\Gamma(\text{invisible})/\Gamma(\pi^+\pi^-\eta)$ Γ_{30}/Γ_1

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.1	90	¹ NAIK	09	CLEO $J/\psi \rightarrow \gamma\eta'$

¹ NAIK 09 reports $[\Gamma(\eta'(958) \rightarrow \text{invisible})/\Gamma(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)] < 5.4 \times 10^{-3}$ which we multiply by our best value $B(\eta \rightarrow 2\gamma) = 39.36 \times 10^{-2}$.

NODE=M002R09
NODE=M002R09

NODE=M002R09;LINKAGE=NA

 $\Gamma(\text{invisible})/\Gamma(\gamma\gamma)$ Γ_{30}/Γ_7

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4	90	ABLIKIM	13	BES3 $J/\psi \rightarrow \phi\eta'$
< 6.69	90	ABLIKIM	06Q	BES $J/\psi \rightarrow \phi\eta'$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M002R44
NODE=M002R44

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.18	90	¹ AAIJ	17D	LHCB $D_{(s)}^+ \rightarrow \pi^+\pi^-\pi^+$
< 0.5	90	² ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-$
< 29	90	³ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+\pi^-$
< 3.3	90	⁴ MORI	07A	BELL $\gamma\gamma \rightarrow \pi^+\pi^-$
< 800	95	DANBURG	73	HBC $2.2 K^-p \rightarrow \Lambda X^0$
< 200	90	RITTENBERG	69	HBC $1.7-2.7 K^-p$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Using branching fractions of $D_{(s)}^+$ decays from PDG 15.

² ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.28 \times 10^{-3}$.

³ Taking into account interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.

⁴ Without interference with the $\gamma\gamma \rightarrow \pi^+\pi^-$ continuum.

NODE=M002R20
NODE=M002R20

OCCUR=2

NODE=M002R20;LINKAGE=A

NODE=M002R20;LINKAGE=AL

NODE=M002R20;LINKAGE=MO

NODE=M002R20;LINKAGE=MR

 $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-4}$	90	¹ ABLIKIM	11G	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ ABLIKIM 11G reports $[\Gamma(\eta'(958) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta'(958))] < 2.84 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta'(958)) = 5.28 \times 10^{-3}$.

NODE=M002R53
NODE=M002R53

NODE=M002R53;LINKAGE=AL

$\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_{32}/Γ_4

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<45	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n4\gamma$

NODE=M002R36
NODE=M002R36 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	BRIERE	00	CLEO $10.6 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<13	90	RITTENBERG	65	HBC $2.7 K^- p$

NODE=M002R8
NODE=M002R8 $\Gamma(\pi^0 \rho^0)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	RITTENBERG	65	HBC $2.7 K^- p$

NODE=M002R10
NODE=M002R10 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.4	90	BRIERE	00	CLEO $10.6 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<11	90	RITTENBERG	65	HBC $2.7 K^- p$

NODE=M002R9
NODE=M002R9 $\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_{36}/Γ_4

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.6	90	ALDE	87B	GAM2 $38 \pi^- p \rightarrow n3\gamma$

NODE=M002R34
NODE=M002R34 $\Gamma(\mu^+ \mu^- \pi^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	90	DZHELADIN	81	CNTR $30 \pi^- p \rightarrow \eta' n$

NODE=M002R31
NODE=M002R31 $\Gamma(\mu^+ \mu^- \eta)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	DZHELADIN	81	CNTR $30 \pi^- p \rightarrow \eta' n$

NODE=M002R30
NODE=M002R30 $\Gamma(e\mu)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.7	90	BRIERE	00	CLEO $10.6 e^+ e^-$

NODE=M002R41
NODE=M002R41 $\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

NODE=M002225

$$|\text{MATRIX ELEMENT}|^2 = |1 + \alpha Y|^2 + CX + DX^2$$

X and Y are Dalitz variables; α is complex and C , and D are real-valued. Parameters C and D are not necessarily equal to c and d , respectively, in the generalized parameterization following this one. May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays. Because of different initial assumptions and strong correlations of the parameters we do not average the parameters in the section below.

NODE=M002225

 $Re(\alpha)$ decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.034 \pm 0.002 \pm 0.002$	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
$-0.054 \pm 0.004 \pm 0.001$	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
$-0.033 \pm 0.005 \pm 0.003$	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
$-0.072 \pm 0.012 \pm 0.006$	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
$-0.021 \pm 0.018 \pm 0.017$	6.7k	³ BRIERE	00	CLEO $10.6 e^+ e^- \rightarrow \eta\pi^+\pi^- X$
$-0.058 \pm 0.013 \pm 0.003$	5.4k	⁴ ALDE	86	GAM2 $38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
-0.08 ± 0.03		^{4,5} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

NODE=M002A0
NODE=M002A0

OCCUR=2

¹ See ABLIKIM 11 for the full correlation matrix.² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.³ Assuming $\text{Im}(\alpha) = 0$, $C = 0$, and $D = 0$.⁴ Assuming $C = 0$.⁵ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.NODE=M002A0;LINKAGE=AB
NODE=M002A0;LINKAGE=AMNODE=M002A0;LINKAGE=BR
NODE=M002A0;LINKAGE=A
NODE=M002A0;LINKAGE=KA

$Im(\alpha)$ decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.000±0.019±0.001	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
0.000±0.038±0.002	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
0.000±0.049±0.001	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.0 ±0.1 ±0.0	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
-0.00 ±0.13 ±0.00	5.4k	³ ALDE	86	GAM2 $38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
0.0 ±0.3		^{3,4} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

¹ See ABLIKIM 11 for the full correlation matrix.

² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

³ Assuming $C = 0$.

⁴ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002IA0
NODE=M002IA0

OCCUR=2

NODE=M002IA0;LINKAGE=AB
NODE=M002IA0;LINKAGE=AM

NODE=M002IA0;LINKAGE=A
NODE=M002IA0;LINKAGE=KA

C decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.0027±0.0024±0.0015	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
0.018 ±0.009 ±0.003	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
0.020 ±0.018 ±0.004	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

NODE=M002C0
NODE=M002C0

NODE=M002C0;LINKAGE=AB
NODE=M002C0;LINKAGE=AM

D decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.053±0.004±0.004	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.061±0.009±0.005	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.059±0.012±0.004	44k	¹ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.066±0.030±0.015	7k	² AMELIN	05A	VES $28 \pi^- A \rightarrow \eta\pi^+\pi^-\pi^- A^*$
0.00 ±0.03 ±0.00	5.4k	³ ALDE	86	GAM2 $38 \pi^- p \rightarrow n\eta\pi^0\pi^0$
0		^{3,4} KALBFLEISCH	74	RVUE $\eta' \rightarrow \eta\pi^+\pi^-$

¹ See ABLIKIM 11 for the full correlation matrix.

² Superseded by DOROFEEV 07, which found this parameterization unacceptable. See below.

³ Assuming $C = 0$.

⁴ From the data of DAUBER 64, RITTENBERG 69, AGUILAR-BENITEZ 72B, JACOBS 73, and DANBURG 73.

NODE=M002D0
NODE=M002D0

NODE=M002D0;LINKAGE=AB
NODE=M002D0;LINKAGE=AM

NODE=M002D0;LINKAGE=AL
NODE=M002D0;LINKAGE=KA

 $\eta'(958) \rightarrow \eta\pi\pi$ DECAY PARAMETERS

NODE=M002227

$$|\text{MATRIX ELEMENT}|^2 \propto 1 + aY + bY^2 + cX + dX^2$$

X and Y are Dalitz variables and a, b, c, and d are real-valued parameters.

May be different for $\eta'(958) \rightarrow \eta\pi^+\pi^-$ and $\eta'(958) \rightarrow \eta\pi^0\pi^0$ decays.

We do not average measurements in the section below because parameter values from each experiment are strongly correlated.

NODE=M002227

a decay parameter

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.077±0.003±0.001		¹ ABLIKIM	23AH	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.056±0.004±0.002	351k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^+\pi^-$
-0.087±0.009±0.006	56k	ABLIKIM	18	BES3 $\eta' \rightarrow \eta\pi^0\pi^0$
-0.074±0.008±0.006	124k	ADLARSON	18A	A2MM $\eta' \rightarrow \eta\pi^0\pi^0$
-0.072±0.007±0.008		² GONZALEZ-S.	18A	RVUE $\eta' \rightarrow \eta\pi^0\pi^0$
-0.047±0.011±0.003	44k	³ ABLIKIM	11	BES3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-0.066±0.016±0.003	15k	⁴ BLIK	09	GAM4 $32.5 \pi^- p \rightarrow \eta' n$
-0.127±0.016±0.008	20k	⁵ DOROFEEV	07	VES $27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ Fit IV, ignoring noncusp terms. Supersedes ABLIKIM 18.

² Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

³ See ABLIKIM 11 for the full correlation matrix.

⁴ From $\eta' \rightarrow \eta\pi^0\pi^0$ decay.

⁵ From $\eta' \rightarrow \eta\pi^+\pi^-$ decay.

NODE=M002DPA
NODE=M002DPA

OCCUR=2

NODE=M002DPA;LINKAGE=B
NODE=M002DPA;LINKAGE=A

NODE=M002DPA;LINKAGE=AB
NODE=M002DPA;LINKAGE=BL
NODE=M002DPA;LINKAGE=DO

b decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.066 \pm 0.006 \pm 0.001$		¹ ABLIKIM 23AH	BES3	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.049 \pm 0.006 \pm 0.006$	351k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.073 \pm 0.014 \pm 0.005$	56k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.063 \pm 0.014 \pm 0.005$	124k	ADLARSON 18A	A2MM	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.052 \pm 0.001 \pm 0.002$		² GONZALEZ-S...18A	RVUE	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.069 \pm 0.019 \pm 0.009$	44k	³ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.063 \pm 0.028 \pm 0.004$	15k	⁴ BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$-0.106 \pm 0.028 \pm 0.014$	20k	⁵ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ Fit IV, ignoring noncusp terms. Supersedes ABLIKIM 18.

² Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

³ See ABLIKIM 11 for the full correlation matrix.

⁴ From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

⁵ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPB
NODE=M002DPB

OCCUR=2

NODE=M002DPB;LINKAGE=B
NODE=M002DPB;LINKAGE=A

NODE=M002DPB;LINKAGE=AB
NODE=M002DPB;LINKAGE=BL
NODE=M002DPB;LINKAGE=DO

NODE=M002DPC
NODE=M002DPC

c decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$0.0027 \pm 0.0024 \pm 0.0018$	351k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^+ \pi^-$
$0.019 \pm 0.011 \pm 0.003$	44k	¹ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$-0.107 \pm 0.096 \pm 0.003$	15k	² BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$0.015 \pm 0.011 \pm 0.014$	20k	³ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ See ABLIKIM 11 for the full correlation matrix.

² From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay.

³ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPC;LINKAGE=AB
NODE=M002DPC;LINKAGE=BL
NODE=M002DPC;LINKAGE=DO

d decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$-0.068 \pm 0.004 \pm 0.001$		¹ ABLIKIM 23AH	BES3	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.063 \pm 0.004 \pm 0.003$	351k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^+ \pi^-$
$-0.074 \pm 0.009 \pm 0.004$	56k	ABLIKIM 18	BES3	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.050 \pm 0.009 \pm 0.005$	124k	ADLARSON 18A	A2MM	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.051 \pm 0.008 \pm 0.006$		² GONZALEZ-S...18A	RVUE	$\eta' \rightarrow \eta \pi^0 \pi^0$
$-0.073 \pm 0.012 \pm 0.003$	44k	³ ABLIKIM 11	BES3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
$0.018 \pm 0.078 \pm 0.006$	15k	⁴ BLIK 09	GAM4	$32.5 \pi^- p \rightarrow \eta' n$
$-0.082 \pm 0.017 \pm 0.008$	20k	⁵ DOROFEEV 07	VES	$27 \pi^- p \rightarrow \eta' n,$ $\pi^- A \rightarrow \eta' \pi^- A^*$

¹ Fit IV, ignoring noncusp terms. Supersedes ABLIKIM 18.

² Theoretical analysis of ADLARSON 18A using resonance chiral perturbation theory to one loop.

³ See ABLIKIM 11 for the full correlation matrix.

⁴ From $\eta' \rightarrow \eta \pi^0 \pi^0$ decay. If $c \equiv 0$ from Bose-Einstein symmetry, $d = -0.067 \pm 0.020 \pm 0.003$.

⁵ From $\eta' \rightarrow \eta \pi^+ \pi^-$ decay.

NODE=M002DPD
NODE=M002DPD

OCCUR=2

NODE=M002DPD;LINKAGE=B
NODE=M002DPD;LINKAGE=A

NODE=M002DPD;LINKAGE=AB
NODE=M002DPD;LINKAGE=BL

NODE=M002DPD;LINKAGE=DO

$\eta'(958)$ β PARAMETER
 $|\text{MATRIX ELEMENT}|^2 = (1 + 2\beta Z)$

See the "Note on η Decay Parameters" in our 1994 edition Physical Review
D50 1173 (1994), p. 1454.

NODE=M002226

NODE=M002226

 β decay parameter

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.61 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.2.		
$-0.640 \pm 0.046 \pm 0.047$	1.8k	ABLIKIM 15G	BES3	$J/\psi \rightarrow \gamma (\pi^0 \pi^0 \pi^0)$
-0.59 ± 0.18	235	BLIK 08	GAMS	$32 \pi^- p \rightarrow \eta' n$
-0.1 ± 0.3		ALDE 87B	GAM2	$38 \pi^- p \rightarrow n 3\pi^0$

NODE=M002B0
NODE=M002B0

$\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Particle Listings for definition of this parameter.

NODE=M002235

NODE=M002235

DECAY ASYMMETRY PARAMETER FOR $\pi^+\pi^-\gamma$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.03 ±0.04 OUR AVERAGE				
-0.019±0.056		AIHARA 87	TPC	$2\gamma \rightarrow \pi^+\pi^-\gamma$
-0.069±0.078	295	GRIGORIAN 75	STRC	$2.1 \pi^- p$
0.00 ±0.10	103	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda \pi^+\pi^-\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.07 ±0.08	152	RITTENBERG 65	HBC	$2.1-2.7 K^- p$

NODE=M002A

NODE=M002A

 $\eta'(958) \rightarrow \gamma \ell^+ \ell^-$ TRANSITION FORM FACTOR SLOPE

Related to the effective virtual meson mass Λ , via slope $\approx \Lambda^{-2}$. See e.g. LANDSBERG 85, eq. (3.8), for a detailed definition.

NODE=M002FFL

NODE=M002FFL

VALUE (GeV^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
1.62±0.17 OUR AVERAGE				
1.60±0.17±0.08	864	¹ ABLIKIM 150	BES3	$J/\psi \rightarrow \gamma e^+ e^-$
1.7 ±0.4	33	¹ VIKTOROV 80		$25,33 \pi^- p \rightarrow 2\mu\gamma$

NODE=M002FFL

¹In the single-pole Ansatz where slope = $1/(\Lambda^2 + \gamma^2)$ with Λ , γ being a Breit-Wigner mass, width for the effective contributing vector meson.

NODE=M002FFL;LINKAGE=A

 $\eta'(958)$ REFERENCES

NODE=M002

ABLIKIM 23AH	PRL 130 081901	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62206
ABLIKIM 22E	PR D105 112010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61639
ABLIKIM 21I	PR D103 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61112
ABLIKIM 21J	PR D103 092005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61114
ABLIKIM 20E	PR D101 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60252
ABLIKIM 19AV	PR D100 052015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60000
ABLIKIM 19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59773
ABLIKIM 18	PR D97 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58711
ABLIKIM 18C	PRL 120 242003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58843
ADLARSON 18A	PR D98 012001	P. Adlarson <i>et al.</i>	(A2 Collab. at MAMI)	REFID=58957
GONZALEZ-S... 18A	EPJ C78 758	S. Gonzalez-Solis, E. Passemar	(BEIJ, IND+)	REFID=59316
AAJ 17D	PL B764 233	R. Aajj <i>et al.</i>	(LHCb Collab.)	REFID=57702
ABLIKIM 17	PRL 118 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57650
ABLIKIM 17T	PR D96 012005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58013
ABLIKIM 16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57511
ABLIKIM 15AD	PR D92 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56983
ABLIKIM 15G	PR D92 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56602
ABLIKIM 15O	PR D92 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56780
ABLIKIM 15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
ACHASOV 15	PR D91 092010	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=56788
AKHMETSHIN 15	PL B740 273	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=56386
PDG 15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=5
ABLIKIM 14M	PRL 112 251801	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55904
DONSKOV 14	MPL A29 1450213	S. Donskov <i>et al.</i>	(GAMS-4 π Collab.)	REFID=56321
PDG 14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ABLIKIM 13	PR D87 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54829
ABLIKIM 13G	PR D87 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54952
ABLIKIM 13O	PR D87 092011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55388
ABLIKIM 13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
ABLIKIM 12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54270
PDG 12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ABLIKIM 11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM 11G	PR D84 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53711
CZERWINSKI 10	PRL 105 122001	E. Czerwinski <i>et al.</i>	(COSY-11 Collab.)	REFID=53364
BLIK 09	PAN 72 231	A.M. Blik <i>et al.</i>	(IHEP (Protvino))	REFID=52727
	Translated from YAF 72 258.			
NAIK 09	PRL 102 061801	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52678
PEDLAR 09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
BLIK 08	PAN 71 2124	A. Blik <i>et al.</i>	(GAMS-4 π Collab.)	REFID=52663
	Translated from YAF 71 2161.			
LIBBY 08	PRL 101 182002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=52591
WICHT 08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
DOROFEEV 07	PL B651 22	V. Dorofeev <i>et al.</i>	(VES Collab.)	REFID=51711
MORI 07A	JPSJ 76 074102	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51691
ABLIKIM 06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM 06Q	PRL 97 202002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51487
AMELIN 05A	PAN 68 372	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=50766
	Translated from YAF 68 401.			
AMSLER 04B	EPJ C33 23	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51079
BAI 04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
BRIERE 00	PRL 84 26	R. Briere <i>et al.</i>	(CLEO Collab.)	REFID=47410
ACCIARRI 98Q	PL B418 399	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46316
BARBERIS 98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
WURZINGER 96	PL B374 283	R. Wurzinger <i>et al.</i>	(BONN, ORSAY, SACL+)	REFID=44992
PDG 94	PR D50 1173	L. Montanet <i>et al.</i>	(CERN, LBL, BOST+)	REFID=43653
AMSLER 93	ZPHY C58 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43311
BELADIDZE 92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)	REFID=43175
	Translated from YAF 55 2748.			

KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=42170
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
BEHREND	91	ZPHY C49 401	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41497
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
KARCH	90	PL B249 353	K. Karch <i>et al.</i>	(Crystal Ball Collab.)	REFID=41377
ROE	90	PR D41 17	N.A. Roe <i>et al.</i>	(ASP Collab.)	REFID=41014
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40564
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
WILLIAMS	88	Translated from YAF 48 436	D.A. Williams <i>et al.</i>	(Crystal Ball Collab.)	REFID=40567
AIHARA	87	PR D38 1365	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40009
ALBRECHT	87B	PL B199 457	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40265
ALDE	87B	ZPHY C36 603	D.M. Alde <i>et al.</i>	(LANL, BELG, SERP, LAPP)	REFID=40236
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)	REFID=40223
ALDE	86	PL B177 115	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=20310
BARTEL	85E	PL 160B 421	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10843
LANDSBERG	85	PRPL 128 301	L.G. Landsberg	(SERP)	REFID=10844
ALTHOFF	84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=20305
BERGER	84B	PL 142B 125	C. Berger	(PLUTO Collab.)	REFID=20306
BINON	84	PL 140B 264	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP+)	REFID=20307
JENNI	83	PR D27 1031	P. Jenni <i>et al.</i>	(SLAC, LBL)	REFID=20304
BARTEL	82B	PL 113B 190	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=20300
BEHREND	82C	PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (err.)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
DZHELYADIN	81	PL 105B 239	R.I. Dzhelyadin <i>et al.</i>	(SERP)	REFID=10836
STANTON	80	PL B92 353	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+)	REFID=40294
VIKTOROV	80	SJNP 32 520	V.A. Viktorov <i>et al.</i>	(SERP)	REFID=20298
APEL	79	PL 83B 131	W.D. Apel, K.H. Augenstein, E. Bertolucci	(KARLK+)	REFID=20295
BINNIE	79	PL 83B 141	D.M. Binnie <i>et al.</i>	(LOIC)	REFID=20296
ZANFINO	77	PRL 38 930	C. Zanfino <i>et al.</i>	(CARL, MCGI, OHIO+)	REFID=20293
GRIGORIAN	75	NP B91 232	A. Grigorian <i>et al.</i>	(+)	REFID=20287
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
DUANE	74	PRL 32 425	A. Duane <i>et al.</i>	(LOIC, SHMP)	REFID=20284
KALBFLEISCH	74	PR D10 916	G.R. Kalbfleisch	(BNL)	REFID=20286
DANBURG	73	PR D8 3744	J.S. Danburg <i>et al.</i>	(BNL, MICH) JP	REFID=20280
JACOBS	73	PR D8 18	S.M. Jacobs <i>et al.</i>	(BRAN, UMD, SYRA+)	REFID=20281
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
APEL	72	PL 40B 680	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA)	REFID=20275
DALPIAZ	72	PL 42B 377	P.F. Dalpiaz <i>et al.</i>	(CERN)	REFID=20278
BASILE	71	NC 3A 371	M. Basile <i>et al.</i>	(CERN, BGNA, STRB)	REFID=20270
HARVEY	71	PRL 27 885	E.H. Harvey <i>et al.</i>	(MINN, MICH)	REFID=20272
BENSINGER	70	PL 33B 505	J.R. Bensinger <i>et al.</i>	(WISC)	REFID=20268
RITTENBERG	69	Thesis UCRL 18863	A. Rittenberg	(LRL) I	REFID=20266
DAVIS	68	PL 27B 532	R. Davis <i>et al.</i>	(NWES, ANL)	REFID=20263
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IJP	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=20253
RITTENBERG	65	PRL 15 556	A. Rittenberg, G.R. Kalbfleisch	(LRL, BNL)	REFID=10761
DAUBER	64	PRL 13 449	P.M. Dauber <i>et al.</i>	(UCLA) JP	REFID=20247
KALBFLEISCH	64B	PRL 13 349	G.R. Kalbfleisch, O.I. Dahl, A. Rittenberg	(LRL) JP	REFID=20252

$f_0(980)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the related review(s):
 Scalar Mesons below 1 GeV

NODE=M003

 $f_0(980)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(980-1010) - i (20-35) OUR ESTIMATE (see Fig. 64.4 in the review)			
$(993 \pm 2^{+2}_{-1}) - i(21 \pm 3^{+2}_{-4})$	1 DANILKIN	21	RVUE Compilation
$(1014 \pm 8) - i(35 \pm 5)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(992.8 \pm 1.3) - i(30.7 \pm 2.3)$	2 ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
$(1003^{+5}_{-27}) - i(21^{+10}_{-8})$	3 GARCIA-MAR..11	RVUE	Compilation
$(996 \pm 7) - i(25^{+10}_{-6})$	4 GARCIA-MAR..11	RVUE	Compilation
$(996^{+4}_{-14}) - i(24^{+11}_{-3})$	5 MOUSSALLAM11	RVUE	Compilation
$(981 \pm 43) - i(18 \pm 11)$	6 MENNESSIER	10	RVUE Compilation
$(1030^{+30}_{-10}) - i(35^{+10}_{-16})$	7 ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(973^{+39}_{-127}) - i(11^{+189}_{-11})$	8 PELAEZ	04A	RVUE $\pi\pi \rightarrow \pi\pi$

NODE=M003PP

NODE=M003PP

NODE=M003PP

→ UNCHECKED ←

OCCUR=2

¹ Data driven analysis using partial-wave dispersion relations .² 5 poles, 5 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$), and BINON 84C ($\eta\eta'$). Based on 18.5k events. Second solution 977.8 ± 1.7 MeV.³ Reanalysis of the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOESCU 73 using Roy equations.⁴ Reanalysis of the K_{e4} data of BATLEY 10C and the $\pi N \rightarrow \pi\pi N$ data of HYAMS 73, GRAYER 74, and PROTOPOESCU 73 using GKPY equations.⁵ Uses Roy equations.⁶ Average of the analyses of three data sets in the K-matrix model. Uses the data of BATLEY 08A, HYAMS 73 , and GRAYER 74 , partially of COHEN 80 or ETKIN 82B.⁷ On sheet II in a 2-pole solution. The other pole is found on sheet III at $(850 - i 100)$ MeV.⁸ Reanalysis of data from PROTOPOESCU 73, ESTABROOKS 74, GRAYER 74, and COHEN 80 in the unitarized ChPT model.

NODE=M003PP;LINKAGE=I

NODE=M003PP;LINKAGE=D

NODE=M003PP;LINKAGE=B

NODE=M003PP;LINKAGE=C

NODE=M003PP;LINKAGE=F

NODE=M003PP;LINKAGE=G

NODE=M003PP;LINKAGE=H

NODE=M003PP;LINKAGE=A

 $f_0(980)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
990 ±20 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$992.0^{+8.5}_{-7.5} \pm 8.6$		1 AAIJ	19H LHCb	$pp \rightarrow D^\pm X$
989.4 ± 1.3	424	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
989.9 ± 0.4	706	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma 3\pi$
$977^{+11}_{-9} \pm 1$	44	2 ECKLUND	09 CLEO	$4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + \text{c.c.}$
$982.2 \pm 1.0^{+8.1}_{-8.0}$		3 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$976.8 \pm 0.3^{+10.1}_{-0.6}$	64k	4 AMBROSINO	07 KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
$984.7 \pm 0.4^{+2.4}_{-3.7}$	64k	5 AMBROSINO	07 KLOE	$1.02 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
973 ± 3	262 ± 30	6 AUBERT	07AKBABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$
970 ± 7	54 ± 9	6 AUBERT	07AKBABR	$10.6 e^+ e^- \rightarrow \phi \pi^0 \pi^0 \gamma$
953 ± 20	2.6k	7 BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
$985.6^{+1.2}_{-1.5} \pm 1.1$		8 MORI	07 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
$983.0 \pm 0.6^{+4.0}_{-3.0}$		9 AMBROSINO	06B KLOE	$1.02 e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

NODE=M003M1

NODE=M003M1

→ UNCHECKED ←

OCCUR=2

OCCUR=2

977.3 ± 0.9 ^{+3.7} _{-4.3}		10	AMBROSINO	06B	KLOE	1.02 e ⁺ e ⁻ → π ⁺ π ⁻ γ	OCCUR=2
950 ± 9	2426	11	GARMASH	06	BELL	B ⁺ → K ⁺ π ⁺ π ⁻	
965 ± 10		12	ABLIKIM	05	BES2	J/ψ → φπ ⁺ π ⁻ , φK ⁺ K ⁻	
1031 ± 8		13	ANISOVICH	03	RVUE		
1037 ± 31			TIKHOMIROV	03	SPEC	40.0 π ⁻ C → K _S ⁰ K _S ⁰ K _L ⁰ X	
973 ± 1	2438	14	ALOISIO	02D	KLOE	e ⁺ e ⁻ → π ⁰ π ⁰ γ	
977 ± 3 ± 2	848	15	AITALA	01A	E791	D _s ⁺ → π ⁻ π ⁺ π ⁺	
969.8 ± 4.5	419	16	ACHASOV	00H	SND	e ⁺ e ⁻ → π ⁰ π ⁰ γ	
985 ⁺¹⁶ ₋₁₂	419	17,18	ACHASOV	00H	SND	e ⁺ e ⁻ → π ⁰ π ⁰ γ	OCCUR=2
976 ± 5 ± 6		19	AKHMETSHIN	99B	CMD2	e ⁺ e ⁻ → π ⁺ π ⁻ γ	
977 ± 3 ± 6	268	19	AKHMETSHIN	99C	CMD2	e ⁺ e ⁻ → π ⁰ π ⁰ γ	
975 ± 4 ± 6		20	AKHMETSHIN	99C	CMD2	e ⁺ e ⁻ → π ⁰ π ⁰ γ	OCCUR=2
975 ± 4 ± 6		21	AKHMETSHIN	99C	CMD2	e ⁺ e ⁻ → π ⁺ π ⁻ γ, π ⁰ π ⁰ γ	OCCUR=3
985 ± 10			BARBERIS	99	OMEG	450 pp → p _S p _f K ⁺ K ⁻	
982 ± 3			BARBERIS	99B	OMEG	450 pp → p _S p _f π ⁺ π ⁻	
982 ± 3			BARBERIS	99C	OMEG	450 pp → p _S p _f π ⁰ π ⁰	
987 ± 6 ± 6		22	BARBERIS	99D	OMEG	450 pp → K ⁺ K ⁻ , π ⁺ π ⁻	
989 ± 15			BELLAZZINI	99	GAM4	450 pp → ppπ ⁰ π ⁰	
991 ± 3		23	KAMINSKI	99	RVUE	ππ → ππ, K ⁻ K ⁻ , σσ	
~ 980		23	OLLER	99	RVUE	ππ → ππ, K ⁻ K ⁻	
~ 993.5			OLLER	99B	RVUE	ππ → ππ, K ⁻ K ⁻	
~ 987		23	OLLER	99C	RVUE	ππ → ππ, K ⁻ K ⁻ , ηη	
957 ± 6		24	ACKERSTAFF	98Q	OPAL	Z → f ₀ X	
960 ± 10			ALDE	98	GAM4		
1015 ± 15		23	ANISOVICH	98B	RVUE	Compilation	
1008		25	LOCHER	98	RVUE	ππ → ππ, K ⁻ K ⁻	
955 ± 10		24	ALDE	97	GAM2	450 pp → ppπ ⁰ π ⁰	
994 ± 9		26	BERTIN	97C	OBLX	0.0 p̄p → π ⁺ π ⁻ π ⁰	
993.2 ± 6.5 ± 6.9		27	ISHIDA	96	RVUE	ππ → ππ, K ⁻ K ⁻	
1006			TORNQVIST	96	RVUE	ππ → ππ, K ⁻ K ⁻ , Kπ, ηπ	
997 ± 5	3k	28	ALDE	95B	GAM2	38 π ⁻ p → π ⁰ π ⁰ n	
960 ± 10	10k	29	ALDE	95B	GAM2	38 π ⁻ p → π ⁰ π ⁰ n	OCCUR=2
994 ± 5			AMSLER	95B	CBAR	0.0 p̄p → 3π ⁰	
~ 996		30	AMSLER	95D	CBAR	0.0 p̄p → π ⁰ π ⁰ π ⁰ , π ⁰ ηη, π ⁰ π ⁰ η	
987 ± 6		31	ANISOVICH	95	RVUE		
1015			JANSSEN	95	RVUE	ππ → ππ, K ⁻ K ⁻	
983		32	BUGG	94	RVUE	p̄p → η2π ⁰	
973 ± 2		33	KAMINSKI	94	RVUE	ππ → ππ, K ⁻ K ⁻	
988		34	ZOU	94B	RVUE		
988 ± 10		35	MORGAN	93	RVUE	ππ(K ⁻ K ⁻) → ππ(K ⁻ K ⁻), J/ψ → φππ(K ⁻ K ⁻), D _s → π(ππ)	
971.1 ± 4.0		24	AGUILAR-...	91	EHS	400 pp	
979 ± 4		36	ARMSTRONG	91	OMEG	300 pp → ppππ, ppK ⁻ K ⁻	
956 ± 12			BREAKSTONE	90	SFM	pp → ppπ ⁺ π ⁻	
959.4 ± 6.5		24	AUGUSTIN	89	DM2	J/ψ → ωπ ⁺ π ⁻	
978 ± 9		24	ABACHI	86B	HRS	e ⁺ e ⁻ → π ⁺ π ⁻ X	
985.0 ^{+9.0} _{-39.0}			ETKIN	82B	MPS	23 π ⁻ p → n2K _S ⁰	
974 ± 4		36	GIDAL	81	MRK2	J/ψ → π ⁺ π ⁻ X	
975		37	ACHASOV	80	RVUE		
986 ± 10		36	AGUILAR-...	78	HBC	0.7 p̄p → K _S ⁰ K _S ⁰	
969 ± 5		36	LEEPER	77	ASPK	2-2.4 π ⁻ p → π ⁺ π ⁻ n, K ⁺ K ⁻ n	
987 ± 7		36	BINNIE	73	CNTR	π ⁻ p → nMM	
1012 ± 6		38	GRAYER	73	ASPK	17 π ⁻ p → π ⁺ π ⁻ n	
1007 ± 20		38	HYAMS	73	ASPK	17 π ⁻ p → π ⁺ π ⁻ n	
997 ± 6		38	PROTOPOP...	73	HBC	7 π ⁺ p → π ⁺ pπ ⁺ π ⁻	

- 1 From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.
- 2 Using a relativistic Breit-Wigner function and taking into account the finite D_s mass.
- 3 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0} \pi \pi = 0$.
- 4 In the kaon-loop fit.
- 5 In the no-structure fit.
- 6 Systematic errors not estimated.
- 7 FLATTE 76 parameterization. $g_{f_0} \pi \pi = 329 \pm 96 \text{ MeV}/c^2$ assuming $g_{f_0} K \bar{K} / g_{f_0} \pi \pi = 2$.
- 8 Breit-Wigner mass. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0} K K / g_{f_0} \pi \pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.
- 9 In the kaon-loop fit following formalism of ACHASOV 89.
- 10 In the no-structure fit assuming a direct coupling of ϕ to $f_0 \gamma$.
- 11 FLATTE 76 parameterization. Supersedes GARMASH 05.
- 12 FLATTE 76 parameterization, $g_{f_0} K \bar{K} / g_{f_0} \pi \pi = 4.21 \pm 0.25 \pm 0.21$.
- 13 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.
- 14 From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution.
- 15 Coupled-channel Breit-Wigner, couplings $g_\pi = 0.09 \pm 0.01 \pm 0.01$, $g_K = 0.02 \pm 0.04 \pm 0.03$.
- 16 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
- 17 Supersedes ACHASOV 98I.
- 18 In the "narrow resonance" approximation.
- 19 Assuming $\Gamma(f_0) = 40 \text{ MeV}$.
- 20 From a narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.
- 21 From the combined fit of the photon spectra in the reactions $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$.
- 22 Supersedes BARBERIS 99 and BARBERIS 99B
- 23 T-matrix pole.
- 24 From invariant mass fit.
- 25 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039–93i) MeV.
- 26 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963–29i) MeV.
- 27 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
- 28 At high $|t|$.
- 29 At low $|t|$.
- 30 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953–55i) MeV and on sheet IV at (938–35i) MeV.
- 31 Combined fit of ALDE 95B, ANISOVICH 94, AMSLER 94D.
- 32 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996–103i) MeV.
- 33 From sheet II pole position.
- 34 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797–185i) MeV and can be interpreted as a shadow pole.
- 35 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978–28i) MeV.
- 36 From coupled channel analysis.
- 37 Coupled channel analysis with finite width corrections.
- 38 Included in AGUILAR-BENITEZ 78 fit.

NODE=M003M1;LINKAGE=F

NODE=M003M1;LINKAGE=EC
NODE=M003M1;LINKAGE=UE

NODE=M003M1;LINKAGE=AK
NODE=M003M1;LINKAGE=AS
NODE=M003M1;LINKAGE=NS

NODE=M003M1;LINKAGE=BO
NODE=M003M1;LINKAGE=MO

NODE=M003M1;LINKAGE=AB
NODE=M003M1;LINKAGE=AM
NODE=M003M1;LINKAGE=GR

NODE=M003M1;LINKAGE=AL
NODE=M003M1;LINKAGE=KM

NODE=M003M1;LINKAGE=KD

NODE=M003M1;LINKAGE=TL
NODE=M003M1;LINKAGE=V9
NODE=M003M1;LINKAGE=V8
NODE=M003M1;LINKAGE=AI
NODE=M003M1;LINKAGE=SM
NODE=M003M1;LINKAGE=ST

NODE=M003M1;LINKAGE=SL

NODE=M003M1;LINKAGE=BD
NODE=M003M1;LINKAGE=AN
NODE=M003M1;LINKAGE=A
NODE=M003M1;LINKAGE=LC
NODE=M003M1;LINKAGE=X
NODE=M003M1;LINKAGE=AA

NODE=M003M1;LINKAGE=LA
NODE=M003M1;LINKAGE=LB
NODE=M003M1;LINKAGE=KL

NODE=M003M1;LINKAGE=CF
NODE=M003M1;LINKAGE=C2
NODE=M003M1;LINKAGE=KM
NODE=M003M1;LINKAGE=L

NODE=M003M1;LINKAGE=K
NODE=M003M1;LINKAGE=B
NODE=M003M1;LINKAGE=B
NODE=M003M1;LINKAGE=R

$f_0(980)$ WIDTH

Width determination very model dependent. Peak width in $\pi\pi$ is about 50 MeV, but decay width can be much larger.

NODE=M003W1

NODE=M003W1

NODE=M003W1

→ UNCHECKED ←

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10 to 100 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
15.3 ± 4.7	424	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
9.5 ± 1.1	706	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma 3\pi$
$91 \begin{smallmatrix} + 30 \\ - 22 \end{smallmatrix} \pm 3$	44	1 ECKLUND	09 CLEO	$4.17 e^+ e^- \rightarrow D_s^- D_s^{*+} + \text{c.c.}$
$66.9 \pm 2.2 \begin{smallmatrix} + 17.6 \\ - 12.5 \end{smallmatrix}$		2 UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
65 ± 13	262 ± 30	3 AUBERT	07AK BABR	$10.6 e^+ e^- \rightarrow \phi \pi^+ \pi^- \gamma$

81 ± 21	54 ± 9	3	AUBERT	07AK	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\pi^0\gamma$	OCCUR=2
51.3 ⁺ ₋ 20.8 ⁺ ₋ 13.2 ⁺ ₋ 17.7 ⁻ ₋ 3.8		4	MORI	07	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$	
61 ± 9 ⁺¹⁴ ₋₈	2584	5	GARMASH	05	BELL	$B^+ \rightarrow K^+\pi^+\pi^-$	
64 ± 16		6	ANISOVICH	03	RVUE		
121 ± 23			TIKHOMIROV	03	SPEC	40.0 $\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$	
~ 70		7	BRAMON	02	RVUE	1.02 $e^+e^- \rightarrow \pi^0\pi^0\gamma$	
44 ± 2 ± 2	848	8	AITALA	01A	E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$	
201 ± 28	419	9	ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	OCCUR=2
122 ± 13	419	10,11	ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
56 ± 20		12	AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$	
65 ± 20			BARBERIS	99	OMEG	450 $pp \rightarrow p_s p_f K^+ K^-$	
80 ± 10			BARBERIS	99B	OMEG	450 $pp \rightarrow p_s p_f \pi^+ \pi^-$	
80 ± 10			BARBERIS	99C	OMEG	450 $pp \rightarrow p_s p_f \pi^0 \pi^0$	
48 ± 12 ± 8		13	BARBERIS	99D	OMEG	450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$	
65 ± 25			BELLAZZINI	99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$	
71 ± 14		14	KAMINSKI	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$	
~ 28		14	OLLER	99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~ 25			OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
~ 14		14	OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \eta\eta$	
70 ± 20			ALDE	98	GAM4		
86 ± 16		14	ANISOVICH	98B	RVUE	Compilation	
54		15	LOCHER	98	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
69 ± 15		16	ALDE	97	GAM2	450 $pp \rightarrow pp\pi^0\pi^0$	
38 ± 20		17	BERTIN	97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$	
~ 100		18	ISHIDA	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
34			TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$	
48 ± 10	3k	19	ALDE	95B	GAM2	38 $\pi^-p \rightarrow \pi^0\pi^0n$	OCCUR=2
95 ± 20	10k	20	ALDE	95B	GAM2	38 $\pi^-p \rightarrow \pi^0\pi^0n$	
26 ± 10			AMSLER	95B	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0$	
~ 112		21	AMSLER	95D	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$	
80 ± 12		22	ANISOVICH	95	RVUE		
30			JANSSEN	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
74		23	BUGG	94	RVUE	$\bar{p}p \rightarrow \eta 2\pi^0$	
29 ± 2		24	KAMINSKI	94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$	
46		25	ZOU	94B	RVUE		
48 ± 12		26	MORGAN	93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_s \rightarrow \pi(\pi\pi)$	
37.4 ± 10.6		16	AGUILAR...	91	EHS	400 pp	
72 ± 8		27	ARMSTRONG	91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$	
110 ± 30			BREAKSTONE	90	SFM	$pp \rightarrow pp\pi^+\pi^-$	
29 ± 13		16	ABACHI	86B	HRS	$e^+e^- \rightarrow \pi^+\pi^-X$	
120 ± 281 ± 20			ETKIN	82B	MPS	23 $\pi^-p \rightarrow n 2K_S^0$	
28 ± 10		27	GIDAL	81	MRK2	$J/\psi \rightarrow \pi^+\pi^-X$	
70 to 300		28	ACHASOV	80	RVUE		
100 ± 80		29	AGUILAR...	78	HBC	0.7 $\bar{p}p \rightarrow K_S^0 K_S^0$	
30 ± 8		27	LEEPER	77	ASPK	2-2.4 $\pi^-p \rightarrow \pi^+\pi^-n, K^+K^-n$	
48 ± 14		27	BINNIE	73	CNTR	$\pi^-p \rightarrow nMM$	
32 ± 10		30	GRAYER	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$	
30 ± 10		30	HYAMS	73	ASPK	17 $\pi^-p \rightarrow \pi^+\pi^-n$	
54 ± 16		30	PROTOPOP...	73	HBC	7 $\pi^+p \rightarrow \pi^+p\pi^+\pi^-$	

- 1 Using a relativistic Breit-Wigner function and taking into account the finite D_S mass.
 2 Breit-Wigner $\pi\pi$ width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0}^2 K K / g_{f_0}^2 \pi\pi = 0$.
 3 Systematic errors not estimated.
 4 Breit-Wigner $\pi\pi$ width. Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0}^2 K K / g_{f_0}^2 \pi\pi = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.
 5 Breit-Wigner, solution 1, PWA ambiguous.
 6 K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.
 7 Using the data of AKHMETSHIN 99C, ACHASOV 00H, and ALOISIO 02D.
 8 Breit-Wigner width.
 9 Supersedes ACHASOV 98I. Using the model of ACHASOV 89.
 10 Supersedes ACHASOV 98I.
 11 In the "narrow resonance" approximation.
 12 From the combined fit of the photon spectra in the reactions $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$, $\pi^0 \pi^0 \gamma$.
 13 Supersedes BARBERIS 99 and BARBERIS 99B.
 14 T-matrix pole.
 15 On sheet II in a 2 pole solution. The other pole is found on sheet III at (1039–93*i*) MeV.
 16 From invariant mass fit.
 17 On sheet II in a 2 pole solution. The other pole is found on sheet III at (963–29*i*) MeV.
 18 Reanalysis of data from HYAMS 73, GRAYER 74, SRINIVASAN 75, and ROSSELET 77 using the interfering amplitude method.
 19 At high $|t|$.
 20 At low $|t|$.
 21 On sheet II in a 4-pole solution, the other poles are found on sheet III at (953–55*i*) MeV and on sheet IV at (938–35*i*) MeV.
 22 Combined fit of ALDE 95B, ANISOVICH 94.
 23 On sheet II in a 2 pole solution. The other pole is found on sheet III at (996–103*i*) MeV.
 24 From sheet II pole position.
 25 On sheet II in a 2 pole solution. The other pole is found on sheet III at (797–185*i*) MeV and can be interpreted as a shadow pole.
 26 On sheet II in a 2 pole solution. The other pole is found on sheet III at (978–28*i*) MeV.
 27 From coupled channel analysis.
 28 Coupled channel analysis with finite width corrections.
 29 From coupled channel fit to the HYAMS 73 and PROTOPODESCU 73 data. With a simultaneous fit to the $\pi\pi$ phase-shifts, inelasticity and to the $K_S^0 K_S^0$ invariant mass.
 30 Included in AGUILAR-BENITEZ 78 fit.

NODE=M003W1;LINKAGE=EC
 NODE=M003W1;LINKAGE=UE

NODE=M003W1;LINKAGE=NS
 NODE=M003W1;LINKAGE=MO

NODE=M003W1;LINKAGE=GA
 NODE=M003W1;LINKAGE=KM

NODE=M003W1;LINKAGE=BR
 NODE=M003W1;LINKAGE=TL
 NODE=M003W1;LINKAGE=V9
 NODE=M003W1;LINKAGE=V8
 NODE=M003W1;LINKAGE=AI
 NODE=M003W1;LINKAGE=SL

NODE=M003W1;LINKAGE=BD
 NODE=M003W1;LINKAGE=AN
 NODE=M003W1;LINKAGE=LO
 NODE=M003W1;LINKAGE=A
 NODE=M003W1;LINKAGE=X
 NODE=M003W1;LINKAGE=AA

NODE=M003W1;LINKAGE=LA
 NODE=M003W1;LINKAGE=LB
 NODE=M003W1;LINKAGE=KL

NODE=M003W1;LINKAGE=CF
 NODE=M003W1;LINKAGE=C2
 NODE=M003W1;LINKAGE=KM
 NODE=M003W1;LINKAGE=L

NODE=M003W1;LINKAGE=K
 NODE=M003W1;LINKAGE=B
 NODE=M003W1;LINKAGE=B
 NODE=M003W1;LINKAGE=C

NODE=M003W1;LINKAGE=R

$f_0(980)$ DECAY MODES

NODE=M003215;NODE=M003

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 $K\bar{K}$	seen
Γ_3 $\gamma\gamma$	seen
Γ_4 $e^+ e^-$	

DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=5;OUR EVAL;→ UNCHECKED ←
 DESIG=4

$f_0(980)$ PARTIAL WIDTHS

NODE=M003220

$\Gamma(\gamma\gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT	Γ_3
------------------------	-------------	-------------	------	---------	------------

NODE=M003W4
 NODE=M003W4

0.29 $^{+0.11}_{-0.06}$ OUR AVERAGE

0.286 $^{+0.017+0.211}_{-0.070}$	1	UEHARA	08A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
0.205 $^{+0.095+0.147}_{-0.083-0.117}$	2	MORI	07	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
0.42 $\pm 0.06 \pm 0.18$	3	OEST	90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32 ±0.05	4 DAI	14A	RVUE	Compilation
0.16 ±0.01	5 MENNESSIER	11	RVUE	
0.29 ±0.21 $\begin{smallmatrix} +0.02 \\ -0.07 \end{smallmatrix}$	6 MOUSSALLAM11		RVUE	Compilation
0.42	7,8 PENNINGTON	08	RVUE	Compilation
0.10	8,9 PENNINGTON	08	RVUE	Compilation
0.28 $\begin{smallmatrix} +0.09 \\ -0.13 \end{smallmatrix}$	10 BOGLIONE	99	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$
0.29 ±0.07 ±0.12	11,12 BOYER	90	MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
0.31 ±0.14 ±0.09	11,12 MARSISKE	90	CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
0.63 ±0.14	13 MORGAN	90	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$

OCCUR=2

¹ Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0 KK}/g_{f_0 \pi\pi} = 0$.

NODE=M003W4;LINKAGE=UE

² Using finite width corrections according to FLATTE 76 and ACHASOV 05, and the ratio $g_{f_0 KK}/g_{f_0 \pi\pi} = 4.21 \pm 0.25 \pm 0.21$ from ABLIKIM 05.

NODE=M003W4;LINKAGE=MO

³ OEST 90 quote systematic errors $\begin{smallmatrix} +0.08 \\ -0.18 \end{smallmatrix}$. We use ±0.18. Observed 60 events.

NODE=M003W4;LINKAGE=H

⁴ Using dispersive analysis with phases from GARCIA-MARTIN 11A and BUETTIKER 04 as input.

NODE=M003W4;LINKAGE=D

⁵ Uses an analytic K-matrix model. Compilation.

NODE=M003W4;LINKAGE=ME

⁶ Using dispersion integral with phase input from Roy equations and data from MARSISKE 90, BOYER 90, BEHREND 92, UEHARA 08A, and MORI 07.

NODE=M003W4;LINKAGE=MU

⁷ Solution A (preferred solution based on χ^2 -analysis).

NODE=M003W4;LINKAGE=P1

⁸ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

NODE=M003W4;LINKAGE=P3

⁹ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

NODE=M003W4;LINKAGE=P2

¹⁰ Supersedes MORGAN 90.

NODE=M003W4;LINKAGE=BL

¹¹ From analysis allowing arbitrary background unconstrained by unitarity.

NODE=M003W4;LINKAGE=B

¹² Data included in MORGAN 90, BOGLIONE 99 analyses.

NODE=M003W4;LINKAGE=C

¹³ From amplitude analysis of BOYER 90 and MARSISKE 90, data corresponds to resonance parameters $m = 989$ MeV, $\Gamma = 61$ MeV.

NODE=M003W4;LINKAGE=A

$\Gamma(e^+e^-)$

Γ_4

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<8.4	90	VOROBYEV	88 ND	$e^+e^- \rightarrow \pi^0\pi^0$

NODE=M003W3
NODE=M003W3

$f_0(980)$ BRANCHING RATIOS

NODE=M003225

$\Gamma(\pi\pi)/[\Gamma(\pi\pi) + \Gamma(K\bar{K})]$

$\Gamma_1/(\Gamma_1+\Gamma_2)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M003R1
NODE=M003R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.52 ±0.12	9.9k	1 AUBERT	060	BABR	$B^\pm \rightarrow K^\pm \pi^\pm \pi^\mp$
0.75 $\begin{smallmatrix} +0.11 \\ -0.13 \end{smallmatrix}$		2 ABLIKIM	05Q	BES2	$\chi_{c0} \rightarrow 2\pi^+ 2\pi^-, \pi^+\pi^- K^+ K^-$
0.84 ±0.02		3 ANISOVICH	02D	SPEC	Combined fit
~ 0.68		OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
0.67 ±0.09		4 LOVERRE	80	HBC	$4 \pi^- p \rightarrow n 2K_S^0$
0.81 $\begin{smallmatrix} +0.09 \\ -0.04 \end{smallmatrix}$		4 CASON	78	STRC	$7 \pi^- p \rightarrow n 2K_S^0$
0.78 ±0.03		4 WETZEL	76	OSPK	$8.9 \pi^- p \rightarrow n 2K_S^0$

¹ Recalculated by us using $\Gamma(K^+ K^-) / \Gamma(\pi^+ \pi^-) = 0.69 \pm 0.32$ from AUBERT 060 and isospin relations.

NODE=M003R1;LINKAGE=AU

² Using data from ABLIKIM 04G.

NODE=M003R1;LINKAGE=AB

³ From a combined K-matrix analysis of Crystal Barrel ($p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

NODE=M003R;LINKAGE=CH

⁴ Measure $\pi\pi$ elasticity assuming two resonances coupled to the $\pi\pi$ and $K\bar{K}$ channels only.

NODE=M003R1;LINKAGE=B

f₀(980) REFERENCES

NODE=M003

DANILKIN	21	PR D103 114023	I. Danilkin, O. Deineka, M. Vanderhaeghen (MAINZ)	REFID=62088
SARANTSEV	21	PL B816 136227	A.V. Sarantsev et al. (BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht et al. (Crystal Barrel Collab.)	REFID=60439
AAJ	19H	JHEP 1904 063	R. Aaij et al. (LHCb Collab.)	REFID=59670
AOUDE	18	PR D98 056021	R.T. Aoude et al.	REFID=59987
ABLIKIM	15P	PR D92 012007	M. Ablikim et al. (BESIII Collab.)	REFID=56781
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington (CEBAF)	REFID=55923
ABLIKIM	12E	PRL 108 182001	M. Ablikim et al. (BESIII Collab.)	REFID=54270
GARCIA-MAR...	11	PRL 107 072001	R. Garcia-Martin et al. (MADR, CRAC)	REFID=16761
GARCIA-MAR...	11A	PR D83 074004	R. Garcia-Martin et al. (MADR, CRAC)	REFID=54121
MENNESSIER	11	PL B696 40	G. Mennessier, S. Narison, X.-G. Wang	REFID=53637
MOUSSALLAM	11	EPJ C71 1814	B. Moussallam	REFID=53975
BATLEY	10C	EPJ C70 635	J.R. Batley et al. (CERN NA48/2 Collab.)	REFID=53567
MENNESSIER	10	PL B688 59	G. Mennessier, S. Narison, X.-G. Wang	REFID=53657
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev (PNPI)	REFID=52719
ECKLUND	09	PR D80 052009	K.M. Ecklund et al. (CLEO Collab.)	REFID=53041
BATLEY	08A	EPJ C54 411	J.R. Batley et al. (CERN NA48/2 Collab.)	REFID=52487
PENNINGTON	08	EPJ C56 1	M.R. Pennington et al.	REFID=52303
UEHARA	08A	PR D78 052004	S. Uehara et al. (BELLE Collab.)	REFID=52309
AMBROSINO	07	EPJ C49 473	F. Ambrosino et al. (KLOE Collab.)	REFID=51618
AUBERT	07AK	PR D76 012008	B. Aubert et al. (BABAR Collab.)	REFID=51908
BONVICINI	07	PR D76 012001	G. Bonvicini et al. (CLEO Collab.)	REFID=51722
MORI	07	PR D75 051101	T. Mori et al. (BELLE Collab.)	REFID=51652
AMBROSINO	06B	PL B634 148	F. Ambrosino et al. (KLOE Collab.)	REFID=51043
AUBERT	06O	PR D74 032003	B. Aubert et al. (BABAR Collab.)	REFID=51141
GARMASH	06	PRL 96 251803	A. Garmash et al. (BELLE Collab.)	REFID=51162
ABLIKIM	05	PL B607 243	M. Ablikim et al. (BES Collab.)	REFID=50450
ABLIKIM	05Q	PR D72 092002	M. Ablikim et al. (BES Collab.)	REFID=50958
ACHASOV	05	PR D72 013006	N.N. Achasov, G.N. Shestakov	REFID=50762
GARMASH	05	PR D71 092003	A. Garmash et al. (BELLE Collab.)	REFID=50641
ABLIKIM	04G	PR D70 092002	M. Ablikim et al. (BES Collab.)	REFID=50187
BUETTIKER	04	EPJ C33 409	P. Buettiker, S. Descotes-Genon, B. Moussallam	REFID=50428; ERROR=6
PELAEZ	04A	MPL A19 2879	J.R. Pelaez (MADU)	REFID=50347
ANISOVICH	03	EPJ A16 229	V.V. Anisovich et al.	REFID=49401
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov et al.	REFID=49423
ALOISIO	02D	Translated from YAF 66 860	A. Aloisio et al. (KLOE Collab.)	REFID=48824
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich et al.	REFID=48831
BRAMON	02	EPJ C26 253	A. Bramon et al.	REFID=49178
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin (Novosibirsk SND Collab.)	REFID=48312
AITALA	01A	PRL 86 765	E.M. Aitala et al. (FNAL E791 Collab.)	REFID=48004
AITALA	01B	PRL 86 770	E.M. Aitala et al. (FNAL E791 Collab.)	REFID=48005
ACHASOV	00B	PL B485 349	M.N. Achasov et al. (Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin et al. (Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin et al. (Novosibirsk CMD-2 Collab.)	REFID=47393
BARBERIS	99	PL B453 305	D. Barberis et al. (Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis et al. (Omega Expt.)	REFID=46922
BARBERIS	99C	PL B453 325	D. Barberis et al. (Omega Expt.)	REFID=46923
BARBERIS	99D	PL B462 462	D. Barberis et al. (Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini et al.	REFID=47400
BOGLIONE	99	EPJ C9 11	M. Boglione, M.R. Pennington	REFID=46931
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau (CRAC, PARIN)	REFID=46927
OLLER	99	PR D60 099906 (errat.)	J.A. Oller et al.	REFID=46899
OLLER	99B	NP A652 407 (errat.)	J.A. Oller, E. Oset	REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset	REFID=47386
ACHASOV	98Q	PL B440 442	M.N. Achasov et al.	REFID=46600
ACKERSTAFF	98Q	EPJ C4 19	K. Ackerstaff et al. (OPAL Collab.)	REFID=46145
ALDE	98	EPJ A3 361	D. Alde et al. (GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde et al. (GAMS Collab.)	REFID=46914
ANISOVICH	98B	Translated from YAF 62 446	V.V. Anisovich et al.	REFID=46331
LOCHER	98	SPU 41 419	V.V. Anisovich et al.	REFID=46372
ALDE	97	EPJ C4 317	M.P. Locher et al. (PSI)	REFID=45392
BERTIN	97C	PL B397 350	D.M. Alde et al. (GAMS Collab.)	REFID=45701
ISHIDA	96	PL B408 476	A. Bertin et al. (OBELIX Collab.)	REFID=45770
TORNQVIST	96	PTP 95 745	S. Ishida et al. (TOKY, MIYA, KEK)	REFID=44507
ALDE	95B	PRL 76 1575	N.A. Tornqvist, M. Roos (HELS)	REFID=44375
AMSLER	95B	ZPHY C66 375	D.M. Alde et al. (GAMS Collab.)	REFID=44377
AMSLER	95D	PL B342 433	C. Amsler et al. (Crystal Barrel Collab.)	REFID=44441
ANISOVICH	95	PL B355 425	C. Amsler et al. (Crystal Barrel Collab.)	REFID=44442
JANSSEN	95	PL B355 363	V.V. Anisovich et al. (PNPI, SERP)	REFID=44508
AMSLER	94D	PR D52 2690	G. Janssen et al. (STON, ANDL, JULI)	REFID=44093
ANISOVICH	94	PL B333 277	C. Amsler et al. (Crystal Barrel Collab.)	REFID=43659
BUGG	94	PL B323 233	V.V. Anisovich et al. (Crystal Barrel Collab.)	REFID=44078
KAMINSKI	94	PR D50 4412	D.V. Bugg et al. (LOQM)	REFID=45771
ZOU	94B	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet (CRAC+)	REFID=44072
MORGAN	93	PR D48 1185	B.S. Zou, D.V. Bugg (LOQM)	REFID=43614
BEHREND	92	PR D48 1185	D. Morgan, M.R. Pennington (RAL DURH)	REFID=43172
AGUILAR...	91	ZPHY C56 381	H.J. Behrend (CELLO Collab.)	REFID=41637
ARMSTRONG	91	ZPHY C50 405	M. Aguilar-Benitez et al. (LEBC-EHS Collab.)	REFID=41744
BOYER	90	ZPHY C51 351	T.A. Armstrong et al. (ATHU, BARI, BIRM+)	REFID=41362
BREAKSTONE	90	PR D42 1350	J. Boyer et al. (Mark II Collab.)	REFID=41376
MARSISKE	90	ZPHY C48 569	A.M. Breakstone et al. (ISU, BGNA, CERN+)	REFID=41351
MORGAN	90	PR D41 3324	H. Marsiske et al. (Crystal Ball Collab.)	REFID=41583
OEST	90	ZPHY C48 623	D. Morgan, M.R. Pennington (RAL, DURH)	REFID=41358
ACHASOV	89	ZPHY C47 343	T. Oest et al. (JADE Collab.)	REFID=48021
AUGUSTIN	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko	REFID=41004
VOROBYEV	88	NP B320 1	J.E. Augustin, G. Cosme (DM2 Collab.)	REFID=41023
ABACHI	86B	JNP 48 273	P.V. Vorobiev et al. (NOVO)	REFID=20394
LONGACRE	86	Translated from YAF 48 436	S. Abachi et al. (PURD, ANL, IND, MICH+)	REFID=20768
BINON	84C	PRL 57 1990	R.S. Longacre et al. (BNL, BRAN, CUNY+)	REFID=21418
BINON	83	PL B177 223	F.G. Binon et al. (BELG, LAPP SERP+)	REFID=20750
ETKIN	82B	NC 80A 363	F.G. Binon et al. (BELG, LAPP, SERP+)	REFID=20390
GIDAL	81	NC 78A 313	A. Etkin et al. (BNL, CUNY, TUFTS, VAND)	REFID=20386
ACHASOV	80	PR D25 1786	G. Gidal et al. (SLAC, LBL)	REFID=20458
COHEN	80	PL 107B 153	N.N. Achasov, S.A. Devyanin, G.N. Shestakov (NOVM)	REFID=20381
LOVERRE	80	JNP 32 566	D. Cohen et al. (ANL) IJP	REFID=20382
AGUILAR...	78	Translated from YAF 32 1098	P.F. Loverre et al. (CERN, CDEF, MADR+) IJP	REFID=20368
CASON	78	PR D22 2595	M. Aguilar-Benitez et al. (MADR, BOMB+)	REFID=20370
LEEPER	77	ZPHY C6 187	N.M. Cason et al. (NDAM, ANL)	REFID=20365
ROSSELET	77	NP B140 73	R.J. Leeper et al. (ISU)	REFID=11004
FLATTE	76	PRL 41 271	L. Rosselet et al. (GEVA, SAFL)	REFID=20446
WETZEL	76	PR D15 574	S.M. Flatte (CERN)	REFID=20362
HYAMS	75	NP B115 208	W. Wetzel et al. (ETH, CERN, LOIC)	REFID=20355
SRINIVASAN	75	NP B100 205	B.D. Hyams et al. (CERN, MPIM)	REFID=21062
ESTABROOKS	74	PR D12 681	V. Srinivasan et al. (NDAM, ANL)	REFID=20111
GRAY	74	NP B79 301	P.G. Estabrooks, A.D. Martin (DURH)	REFID=20113
BINNIE	73	NP B75 189	G. Grayer et al. (CERN, MPIM)	REFID=20343
GRAY	73	PRL 31 1534	D.M. Binnie et al. (LOIC, SHMP)	REFID=20347
HYAMS	73	Tallahassee	G. Grayer et al. (CERN, MPIM)	REFID=20107
PROTOPOP...	73	NP B64 134	B.D. Hyams et al. (CERN, MPIM)	REFID=20108
		PR D7 1279	S.D. Protopopescu et al. (LBL)	

$a_0(980)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

See the related review(s):
 Scalar Mesons below 1 GeV

NODE=M036

 $a_0(980)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(970-1020) - i (30-70) OUR ESTIMATE (see Fig. 64.2 in the review) [(960-1030) - i (20-70) MeV OUR 2023 ESTIMATE]			
$(1002.4 \pm 1.4 \pm 6.6)$ $-i(63.5 \pm 2.9)$	¹ ALBRECHT	20	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1000.7^{+12.9}_{-0.7})$ $-i(36.6^{+12.7}_{-2.6})$	² LU	20	RVUE $\gamma\gamma \rightarrow \pi^0 \eta, K_S^0 K_S^0$
$(989 \pm 5) - i(40 \pm 5)$	³ BUGG	08A	RVUE $\bar{p}p$ annihilation data
$(1117^{+24}_{-320}) - i(12^{+43}_{-12})$	⁴ PELAEZ	04A	RVUE $\pi\pi \rightarrow \pi\pi, \pi K \rightarrow \pi K$
$(982 \pm 3) - i(46 \pm 4)$	⁵ ABELE	98	CBAR $0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$

¹ Pole mass on sheet closest to the physical axis - the more remote pole is extracted at $(1004.1 \pm 1.5 \pm 6.5) - i(48.6 \pm 1.2 \pm 3.4)$ MeV.

² T-matrix pole on sheet II.

³ T-matrix pole on sheet II. Parameterizes couplings to $\bar{K}K, \pi\eta,$ and $\pi\eta'$. Uses AMSLER 94D and ABELE 98.

⁴ Reanalysis of data from LINGLIN 73, ESTABROOKS 78, and ASTON 88 in the unitarized ChPT model.

⁵ T-matrix pole on sheet II; the pole on sheet III is at $(1006 - i 49)$ MeV.

NODE=M036PP

NODE=M036PP

NODE=M036PP

NEW; → UNCHECKED ←

NODE=M036PP;LINKAGE=E

NODE=M036PP;LINKAGE=LU

NODE=M036PP;LINKAGE=C

NODE=M036PP;LINKAGE=A

NODE=M036PP;LINKAGE=D

 $a_0(980)$ MASS

VALUE (MeV)	DOCUMENT ID
980 ± 20 OUR ESTIMATE	Mass determination very model dependent

NODE=M036205

NODE=M036MX

→ UNCHECKED ←

 $\eta\pi$ FINAL STATE ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$982.5 \pm 1.6 \pm 1.1$	16.9k	¹ AMBROSINO	09F	KLOE	$1.02 e^+ e^- \rightarrow \eta\pi^0 \gamma$
986 ± 4		ANISOVICH	09	RVUE	$0.0 \bar{p}p, \pi N$
$982.3^{+0.6}_{-0.7} \quad ^{+3.1}_{-4.7}$		² UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
$985 \pm 4 \pm 6$	318	ACHARD	02B	L3	$183-209 e^+ e^- \rightarrow e^+ e^- \eta\pi^+ \pi^-$
995^{+52}_{-10}	36	³ ACHASOV	00F	SND	$e^+ e^- \rightarrow \eta\pi^0 \gamma$
994^{+33}_{-8}	36	⁴ ACHASOV	00F	SND	$e^+ e^- \rightarrow \eta\pi^0 \gamma$
975 ± 7		BARBERIS	00H		$450 pp \rightarrow p_f \eta\pi^0 p_s$
988 ± 8		BARBERIS	00H		$450 pp \rightarrow \Delta_f^{++} \eta\pi^- p_s$
~ 1055		⁵ OLLER	99	RVUE	$\eta\pi, K\bar{K}$
~ 1009.2		⁵ OLLER	99B	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
993.1 ± 2.1		⁶ TEIGE	99	B852	$18.3 \pi^- p \rightarrow \eta\pi^+ \pi^- n$
988 ± 6		⁵ ANISOVICH	98B	RVUE	Compilation
987		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi,$ $\eta\pi$
991		JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi,$ $\eta\pi$
$984.45 \pm 1.23 \pm 0.34$		AMSLER	94C	CBAR	$0.0 \bar{p}p \rightarrow \omega\eta\pi^0$
982 ± 2		⁷ AMSLER	92	CBAR	$0.0 \bar{p}p \rightarrow \eta\eta\pi^0$
984 ± 4	1040	⁷ ARMSTRONG	91B	OMEG ±	$300 pp \rightarrow p p \eta\pi^+ \pi^-$
976 ± 6		ATKINSON	84E	OMEG ±	$25-55 \gamma p \rightarrow \eta\pi n$
986 ± 3	500	⁸ EVANGELIS...	81	OMEG ±	$12 \pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$

NODE=M036M1

NODE=M036M1

OCCUR=2

OCCUR=2

990 ± 7	145	⁸ GURTU	79	HBC ±	4.2 $K^- p \rightarrow \Lambda \eta 2\pi$
980 ±11	47	CONFORTO	78	OSPK -	4.5 $\pi^- p \rightarrow p X^-$
978 ±16	50	CORDEN	78	OMEG ±	12-15 $\pi^- p \rightarrow n \eta 2\pi$
977 ± 7		GRASSLER	77	HBC -	16 $\pi^\mp p \rightarrow p \eta 3\pi$
989 ± 4	70	WELLS	75	HBC -	3.1-6 $K^- p \rightarrow \Lambda \eta 2\pi$
972 ±10	150	DEFOIX	72	HBC ±	0.7 $\bar{p} p \rightarrow 7\pi$
970 ±15	20	BARNES	69C	HBC -	4-5 $K^- p \rightarrow \Lambda \eta 2\pi$
980 ±10		CAMPBELL	69	DBC ±	2.7 $\pi^+ d$
980 ±10	15	MILLER	69B	HBC -	4.5 $K^- N \rightarrow \eta \pi \Lambda$
980 ±10	30	AMMAR	68	HBC ±	5.5 $K^- p \rightarrow \Lambda \eta 2\pi$

OCCUR=2

- ¹ Using the model of ACHASOV 89 and ACHASOV 03B.
- ² From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.
- ³ Using the model of ACHASOV 89. Supersedes ACHASOV 98B.
- ⁴ Using the model of JAFFE 77. Supersedes ACHASOV 98B.
- ⁵ T-matrix pole.
- ⁶ Breit-Wigner fit, average between a_0^\pm and a_0^0 . The fit favors a slightly heavier a_0^\pm .
- ⁷ From a single Breit-Wigner fit.
- ⁸ From $f_1(1285)$ decay.

NODE=M036M1;LINKAGE=AM
 NODE=M036M1;LINKAGE=UE

NODE=M036M1;LINKAGE=V1
 NODE=M036M1;LINKAGE=M2
 NODE=M036M1;LINKAGE=AN

NODE=M036M1;LINKAGE=BF
 NODE=M036M1;LINKAGE=A
 NODE=M036M1;LINKAGE=R

$K\bar{K}$ ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M036M2
 NODE=M036M2

• • • We do not use the following data for averages, fits, limits, etc. • • •

947.7 ⁺ _{-5.0} ± 6.6		¹ AAIJ	19H	LHCB	$pp \rightarrow D^\pm X$
925 ± 5 ± 8	190k	² AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
~ 1053		³ OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
975 ±15		BERTIN	98B	OBLX	$0.0 \bar{p} p \rightarrow K^\pm K_S \pi^\mp$
970 ±10	316	DEBILLY	80	HBC	$1.2-2 \bar{p} p \rightarrow f_1(1285)\omega$
1016 ±10	100	⁴ ASTIER	67	HBC	$0.0 \bar{p} p$
1003.3 ± 7.0	143	^{5,6} ROSENFELD	65	RVUE	

- ¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.
- ² Using a two-channel resonance parametrization with couplings fixed to ABELE 98.
- ³ T-matrix pole.
- ⁴ ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.
- ⁵ Note on J^P . Main argument for 0^+ is small Q value. Isotropy of decay distribution in $\bar{p} p$ at rest proves nothing. See discussion by Rosenfeld (Oxford) and Butterworth (Heidelberg).
- ⁶ Plus systematic errors.

NODE=M036M2;LINKAGE=C

NODE=M036M2;LINKAGE=B
 NODE=M036M2;LINKAGE=AN
 NODE=M036M2;LINKAGE=A
 NODE=M036M2;LINKAGE=01

NODE=M036M2;LINKAGE=S

$a_0(980)$ WIDTH

NODE=M036210

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M036W1

50 to 100 OUR ESTIMATE Width determination very model dependent. Peak width in $\eta\pi$ is about 60 MeV, but decay width can be much larger.

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

75.6 ± 1.6 ^{+17.4} _{-10.0}		¹ UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
50 ±13 ± 4	318	ACHARD	02B	L3	$183-209 e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
72 ±16		BARBERIS	00H		$450 pp \rightarrow p_f \eta \pi^0 p_s$
61 ±19		BARBERIS	00H		$450 pp \rightarrow \Delta_f^{++} \eta \pi^- p_s$
~ 42		² OLLER	99	RVUE	$\eta\pi, K\bar{K}$
~ 112		² OLLER	99B	RVUE	$\pi\pi \rightarrow \eta\pi, K\bar{K}$
71 ± 7		TEIGE	99	B852	$18.3 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
92 ±20		² ANISOVICH	98B	RVUE	Compilation
65 ±10		³ BERTIN	98B	OBLX ±	$0.0 \bar{p} p \rightarrow K^\pm K_S \pi^\mp$
~ 100		TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
202		JANSSEN	95	RVUE	$\eta\pi \rightarrow \eta\pi, K\bar{K}, K\pi, \eta\pi$
54.12 ± 0.34 ± 0.12		AMSLER	94C	CBAR	$0.0 \bar{p} p \rightarrow \omega \eta \pi^0$
54 ±10		⁴ AMSLER	92	CBAR	$0.0 \bar{p} p \rightarrow \eta \eta \pi^0$

OCCUR=2

95 ±14	1040	⁴ ARMSTRONG 91B	OMEG ±	300 $\rho\rho \rightarrow \rho\rho\eta\pi^+\pi^-$
62 ±15	500	⁵ EVANGELIS...	81 OMEG ±	12 $\pi^- \rho \rightarrow \eta\pi^+\pi^-\pi^-\rho$
60 ±20	145	⁵ GURTU	79 HBC ±	4.2 $K^- \rho \rightarrow \Lambda\eta 2\pi$
60 ⁺⁵⁰ ₋₃₀	47	CONFORTO	78 OSPK -	4.5 $\pi^- \rho \rightarrow \rho X^-$
86.0 ^{+60.0} _{-50.0}	50	CORDEN	78 OMEG ±	12-15 $\pi^- \rho \rightarrow n\eta 2\pi$
44 ±22		GRASSLER	77 HBC -	16 $\pi^{\mp} \rho \rightarrow \rho\eta 3\pi$
80 to 300		⁶ FLATTE	76 RVUE -	4.2 $K^- \rho \rightarrow \Lambda\eta 2\pi$
16.0 ^{+25.0} _{-16.0}	70	⁷ WELLS	75 HBC -	3.1-6 $K^- \rho \rightarrow \Lambda\eta 2\pi$
30 ± 5	150	⁸ DEFOIX	72 HBC ±	0.7 $\bar{p}\rho \rightarrow 7\pi$
40 ±15		CAMPBELL	69 DBC ±	2.7 $\pi^+ d$
60 ±30	15	MILLER	69B HBC -	4.5 $K^- N \rightarrow \eta\pi\Lambda$
80 ±30	30	AMMAR	68 HBC ±	5.5 $K^- \rho \rightarrow \Lambda\eta 2\pi$

- ¹ From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.
- ² T-matrix pole.
- ³ The $\eta\pi$ width.
- ⁴ From a single Breit-Wigner fit.
- ⁵ From $f_1(1285)$ decay.
- ⁶ Using a two-channel resonance parametrization of GAY 76B data.
- ⁷ Weak evidence only for $a_0(980)^+$ production.
- ⁸ This number has very little meaning. Error is much too small. Vlada

NODE=M036W1;LINKAGE=UE

NODE=M036W1;LINKAGE=AN
 NODE=M036W1;LINKAGE=BE
 NODE=M036W1;LINKAGE=A
 NODE=M036W1;LINKAGE=R
 NODE=M036W1;LINKAGE=F
 NODE=M036W1;LINKAGE=W
 NODE=M036W1;LINKAGE=01

NODE=M036W2
 NODE=M036W2

K \bar{K} ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 48		¹ OLLER	99C	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
~ 25	100	² ASTIER	67	HBC ±	
57 ± 13	143	³ ROSENFELD	65	RVUE ±	

- ¹ T-matrix pole.
- ² ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.
- ³ Plus systematic errors.

NODE=M036W2;LINKAGE=AN
 NODE=M036W2;LINKAGE=A
 NODE=M036W2;LINKAGE=S

$a_0(980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi$	seen
Γ_2 $K\bar{K}$	seen
Γ_3 $\eta'\pi$	seen
Γ_4 $\rho\pi$	not seen
Γ_5 $\gamma\gamma$	seen
Γ_6 e^+e^-	

NODE=M036215;NODE=M036

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=8
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6

$a_0(980)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	DOCUMENT ID	TECN	Γ_5
------------------------	-------------	------	------------

••• We do not use the following data for averages, fits, limits, etc. •••

0.30 ± 0.10	¹ AMSLER	98	RVUE
-------------	---------------------	----	------

- ¹ Using $\Gamma_{\gamma\gamma} B(a_0(980) \rightarrow \eta\pi) = 0.24 \pm 0.08$ keV.

NODE=M036217

NODE=M036W4
 NODE=M036W4

NODE=M036W4;LINKAGE=A

$a_0(980)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_5/\Gamma$
---	-------------	------	---------	---------------------------

0.21 ^{+0.08}_{-0.04} OUR AVERAGE

0.128 ^{+0.003+0.502} _{-0.002-0.043}	¹ UEHARA	09A	BELL	$\gamma\gamma \rightarrow \pi^0\eta$
0.28 ± 0.04 ± 0.10	44	OEST	90	JADE $e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.19 ± 0.07 ^{+0.10} _{-0.07}		ANTREASYAN	86	CBAL $e^+e^- \rightarrow e^+e^-\pi^0\eta$

- ¹ From a fit with the S-wave amplitude including two interfering Breit-Wigners plus a background term.

NODE=M036220

NODE=M036G1
 NODE=M036G1

NODE=M036G1;LINKAGE=UE

$\Gamma(\eta\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_6/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.5	90	VOROBYEV	88	ND	$e^+e^- \rightarrow \pi^0\eta$

NODE=M036G2
NODE=M036G2

$a_0(980)$ BRANCHING RATIOS

NODE=M036225

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$					Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.172±0.019 OUR AVERAGE					
0.137±0.036±0.042	¹ ABLIKIM	22AH	BES3		$D_s^+ \rightarrow K_S^0 K^+ \pi^0$
0.23 ±0.05	² ABELE	98	CBAR		$0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
0.166±0.01 ±0.02	³ BARBERIS	98C	OMEG		$450 p p \rightarrow p_f f_1(1285) p_s$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.138±0.001±0.035	⁴ ALBRECHT	20	CBAR		$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
1.20 ±0.15	⁵ ANISOVICH	09	RVUE		$0.0 \bar{p}p, \pi N$
1.05 ±0.07 ±0.05	⁶ BUGG	08A	RVUE	0	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
0.57 ±0.16	⁷ BARGIOTTI	03	OBLX		$\bar{p}p$
~ 0.60	OLLER	99B	RVUE		$\pi\pi \rightarrow \eta\pi, K\bar{K}$
0.7 ±0.3	³ CORDEN	78	OMEG		$12-15 \pi^- p \rightarrow n\eta 2\pi$
0.25 ±0.08	³ DEFOIX	72	HBC	±	$0.7 \bar{p} \rightarrow 7\pi$

NODE=M036R2
NODE=M036R2

¹ Using $D_s^+ \rightarrow a_0(980)^+ \pi^0$ from ABLIKIM 19BE.

² Using $\pi^0 \pi^0 \eta$ from AMSLER 94D.

³ From the decay of $f_1(1285)$.

⁴ Residues from T-matrix pole with 2 poles, 2 channels. Solution on adjacent sheet $0.149 \pm 0.001 \pm 0.039$.

⁵ This is a ratio of couplings.

⁶ A ratio of couplings, using AMSLER 94D and ABELE 98. Supersedes BUGG 94.

⁷ Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0,$ and $K^\pm K_S^0 \pi^\mp$.

NODE=M036R2;LINKAGE=B
NODE=M036R2;LINKAGE=Q
NODE=M036R2;LINKAGE=L
NODE=M036R2;LINKAGE=A

NODE=M036R2;LINKAGE=AN
NODE=M036R2;LINKAGE=BU
NODE=M036R;LINKAGE=BG

$\Gamma(\eta'\pi)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	116k	¹ CHEN	20A	BELL	$D^0 \rightarrow K^- \pi^+ \eta$

NODE=M036R00
NODE=M036R00

¹ From an amplitude analysis of the $D^0 \rightarrow K^- \pi^+ \eta$ decay in a three-channel Flatte model with a 10.1 σ significance. Earlier observed by ABLIKIM 17K in the $\chi_{c1} \rightarrow \eta\pi^+ \pi^-$ decay with a 8.9 σ significance.

NODE=M036R00;LINKAGE=A

$\Gamma(\rho\pi)/\Gamma(\eta\pi)$					Γ_4/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
$\rho\pi$ forbidden.					
<0.25	70	¹ AMMAR	70	HBC	± 4.1,5.5 $K^- p \rightarrow \Lambda\eta 2\pi$

NODE=M036R1
NODE=M036R1
NODE=M036R1

¹ Not clear if they really observed the $a_0(980)$ 3 standard deviations.

NODE=M036R1;LINKAGE=01

$a_0(980)$ REFERENCES

NODE=M036

ABLIKIM	22AH	PRL 129 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61880
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)	REFID=60333
LU	20	EPJ C80 436	J. Lu, B. Moussallam		REFID=60436
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
ABLIKIM	19BE	PRL 123 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60055
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57953
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
BUGG	08A	PR D78 074023	D.V. Bugg	(LOQM)	REFID=52578
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
ACHASOV	03B	PR D68 014006	N.N. Achasov, A.V. Kiselev		REFID=49476
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48574
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
OLLER	99	PR D60 099906 (errat.)	J.A. Oller <i>et al.</i>		REFID=46899
OLLER	99B	NP A652 407 (errat.)	J.A. Oller, E. Oset		REFID=46924
OLLER	99C	PR D60 074023	J.A. Oller, E. Oset		REFID=47386
TEIGE	99	PR D59 012001	S. Teige <i>et al.</i>	(BNL E852 Collab.)	REFID=46613
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46317
AMSLER	98	RMP 70 1293	C. Amsler		REFID=46601
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331

Translated from UFN 168 481.

BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46346
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)	REFID=44507
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)	REFID=44508
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44091
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41862
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
Translated from YAF 48 436.					
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20624
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
DEBILLY	80	NP B176 1	L. de Billy <i>et al.</i>	(CURIN, LAUS, NEUC+)	REFID=20461
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)	REFID=20456
CONFORTO	78	LNC 23 419	B. Conforto <i>et al.</i>	(RHEL, TINTO, CHIC+)	REFID=20451
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20452
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20447
JAFFE	77	PR D15 267,281	R. Jaffe	(MIT)	REFID=43673
FLATTE	76	PL 63B 224	S.M. Flatte	(CERN)	REFID=20446
GAY	76B	PL 63B 220	J.B. Gay <i>et al.</i>	(CERN, AMST, NIJM) JP	REFID=20445
WELLS	75	NP B101 333	J. Wells <i>et al.</i>	(OXF)	REFID=20444
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)	REFID=20435
AMMAR	70	PR D2 430	R. Ammar <i>et al.</i>	(KANS, NWES, ANL, WISC)	REFID=20428
BARNES	69C	PRL 23 610	V.E. Barnes <i>et al.</i>	(BNL, SYRA)	REFID=20418
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)	REFID=20419
MILLER	69B	PL 29B 255	D.H. Miller <i>et al.</i>	(PURD)	REFID=20424
Also		PR 188 2011	W.L. Yen <i>et al.</i>	(PURD)	REFID=20425
AMMAR	68	PRL 21 1832	R. Ammar <i>et al.</i>	(NWES, ANL)	REFID=20412
ASTIER	67	PL 25B 294	A. Astier <i>et al.</i>	(CDEF, CERN, IRAD)	REFID=20405
Includes data of BARLOW 67, CONFORTO 67, and ARMENTEROS 65.					
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
CONFORTO	67	NP B3 469	G. Conforto <i>et al.</i>	(CERN, CDEF, IPNP+)	REFID=20411
ARMENTEROS	65	PL 17 344	R. Armenteros <i>et al.</i>	(CERN, CDEF)	REFID=20396
ROSENFELD	65	Oxford Conf. 58	A.H. Rosenfeld	(LRL)	REFID=20399

NODE=M004

 $\phi(1020)$

$$I^G(J^{PC}) = 0^-(1^--)$$

 $\phi(1020)$ MASS

NODE=M004M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1019.461 ± 0.016 OUR AVERAGE				
1019.463 ± 0.061	2.3M	¹ KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+K^-$, $K_S^0 K_L^0$
1019.462 ± 0.042 ± 0.056	28k	² LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1019.51 ± 0.02 ± 0.05		³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^- \gamma$
1019.30 ± 0.02 ± 0.10	105k	AKHMETSHIN 06	CMD2	0.98-1.06 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.52 ± 0.05 ± 0.05	17.4k	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.483 ± 0.011 ± 0.025	272k	⁴ AKHMETSHIN 04	CMD2	$\eta\gamma$ $e^+e^- \rightarrow K_L^0 K_S^0$
1019.42 ± 0.05	1900k	⁵ ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-$, $K_S^0 K_L^0, \pi^+\pi^-\pi^0$
1019.40 ± 0.04 ± 0.05	23k	AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1019.36 ± 0.12		⁶ ACHASOV	00B	SND $e^+e^- \rightarrow \eta\gamma$
1019.38 ± 0.07 ± 0.08	2200	⁷ AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.51 ± 0.07 ± 0.10	11169	AKHMETSHIN 98	CMD2	2γ $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1019.5 ± 0.4		BARBERIS	98	OMEG 450 $pp \rightarrow pp2K^+2K^-$
1019.42 ± 0.06	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
1019.7 ± 0.3	2012	DAVENPORT	86	MPSF 400 $pA \rightarrow 4KX$
1019.7 ± 0.1 ± 0.1	5079	ALBRECHT	85D	ARG 10 $e^+e^- \rightarrow K^+K^-X$
1019.3 ± 0.1	1500	ARENTON	82	AEMS 11.8 polar. $pp \rightarrow KK$
1019.67 ± 0.17	25080	⁸ PELLINEN	82	RVUE
1019.52 ± 0.13	3681	BUKIN	78C	OLYA $e^+e^- \rightarrow$ hadrons

NODE=M004M

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

1018.4	± 0.5	± 0.1	⁹ ALBRECHT	20	CBAR	0.9 $\bar{p}p \rightarrow$ $K^+ K^- \pi^0$		
1019.21	± 0.04	± 0.03	¹⁰ HOID	20	RVUE	$e^+ e^- \rightarrow \pi^0 \gamma$		
1019.54	± 0.10	± 0.51	¹¹ AAIJ	19H	LHCB	$p\bar{p} \rightarrow D^\pm X$	OCCUR=2	
1019.20	± 0.02	± 0.01	¹² HOFERICHT...	19	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$		
1019.469	± 0.061	1.7M	KOZYREV	18	CMD3	$e^+ e^- \rightarrow K^+ K^-$		
1019.457	± 0.061	610k	KOZYREV	16	CMD3	$e^+ e^- \rightarrow K_S^0 K_L^0$		
1019.48	± 0.01		LEES	13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$		
1019.441	± 0.008	± 0.080	542k	¹³ AKHMETSHIN	08	CMD2	1.02 $e^+ e^- \rightarrow$ $K^+ K^-$	
1019.63	± 0.07	12540	¹⁴ AUBERT,B	05J	BABR	$D^0 \rightarrow \bar{K}^0 K^+ K^-$		
1019.8	± 0.7		ARMSTRONG	86	OMEG	85 $\pi^+ / p p \rightarrow$ $\pi^+ / p^4 K p$		
1020.1	± 0.11	5526	¹⁴ ATKINSON	86	OMEG	20-70 γp		
1019.7	± 1.0		BEBEK	86	CLEO	$e^+ e^- \rightarrow \gamma(4S)$		
1019.411	± 0.008	642k	¹⁵ DIJKSTRA	86	SPEC	100-200 $\pi^\pm, \bar{p}, p,$ $K^\pm, \text{ on Be}$		
1020.9	± 0.2		¹⁴ FRAME	86	OMEG	13 $K^+ p \rightarrow \phi K^+ p$		
1021.0	± 0.2		¹⁴ ARMSTRONG	83B	OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$	OCCUR=2	
1020.0	± 0.5		¹⁴ ARMSTRONG	83B	OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$		
1019.7	± 0.3		¹⁴ BARATE	83	GOLI	190 $\pi^- \text{Be} \rightarrow 2\mu X$		
1019.8	± 0.2	± 0.5	766	IVANOV	81	OLYA	1-1.4 $e^+ e^- \rightarrow$ $K^+ K^-$	
1019.4	± 0.5	337	COOPER	78B	HBC	0.7-0.8 $\bar{p}p \rightarrow$ $K_S^0 K_L^0 \pi^+ \pi^-$		
1020	± 1	383	¹⁴ BALDI	77	CNTR	10 $\pi^- p \rightarrow \pi^- \phi p$	OCCUR=2	
1018.9	± 0.6	800	COHEN	77	ASPK	6 $\pi^\pm N \rightarrow$ $K^+ K^- N$		
1019.7	± 0.5	454	KALBFLEISCH	76	HBC	2.18 $K^- p \rightarrow \Lambda K \bar{K}$		
1019.4	± 0.8	984	BESCH	74	CNTR	2 $\gamma p \rightarrow p K^+ K^-$		
1020.3	± 0.4	100	BALLAM	73	HBC	2.8-9.3 γp		
1019.4	± 0.7		BINNIE	73B	CNTR	$\pi^- p \rightarrow \phi n$		
1019.6	± 0.5	120	¹⁶ AGUILAR-...	72B	HBC	3.9,4.6 $K^- p \rightarrow$ $\Lambda K^+ K^-$	OCCUR=2	
1019.9	± 0.5	100	¹⁶ AGUILAR-...	72B	HBC	3.9,4.6 $K^- p \rightarrow$ $K^- p K^+ K^-$		
1020.4	± 0.5	131	COLLEY	72	HBC	10 $K^+ p \rightarrow K^+ p \phi$		
1019.9	± 0.3	410	STOTTLE...	71	HBC	2.9 $K^- p \rightarrow$ $\Sigma / \Lambda K \bar{K}$		

¹ Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.

² Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

³ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.

⁴ Update of AKHMETSHIN 99D

⁵ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of $K^+ K^-$, $K_S K_L$, $\pi^+ \pi^- \pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

⁶ Using a total width of 4.43 ± 0.05 MeV. Systematic uncertainty included.

⁷ Using a total width of 4.43 ± 0.05 MeV.

⁸ PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DE-GROOT 74.

⁹ Width fixed at 4.2 MeV.

¹⁰ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization. Inclusion of vacuum polarization gives 1019.457 ± 0.020 MeV.

¹¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

¹² The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.

¹³ Strongly correlated with AKHMETSHIN 04.

¹⁴ Systematic errors not evaluated.

¹⁵ Weighted and scaled average of 12 measurements of DIJKSTRA 86.

¹⁶ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

$\phi(1020)$ WIDTH

NODE=M004W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.249±0.013 OUR AVERAGE				
4.245±0.013	2.3M	¹ KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+K^-$, $K_S^0 K_L^0$
4.205±0.103±0.067	28k	² LEES	14H	BABR $e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
4.29 ±0.04 ±0.07		³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^- \gamma$
4.30 ±0.06 ±0.17	105k	AKHMETSHIN	06	CMD2 0.98–1.06 $e^+e^- \rightarrow$ $\pi^+\pi^-\pi^0$
4.280±0.033±0.025	272k	⁴ AKHMETSHIN	04	CMD2 $e^+e^- \rightarrow K_L^0 K_S^0$
4.21 ±0.04	1900k	⁵ ACHASOV	01E	SND $e^+e^- \rightarrow K^+K^-$, $K_S K_L, \pi^+\pi^-\pi^0$
4.44 ±0.09	55600	AKHMETSHIN	95	CMD2 $e^+e^- \rightarrow$ hadrons
4.5 ±0.7	1500	ARENTON	82	AEMS 11.8 polar. $pp \rightarrow KK$
4.2 ±0.6	766	⁶ IVANOV	81	OLYA 1–1.4 $e^+e^- \rightarrow K^+K^-$
4.3 ±0.6		⁶ CORDIER	80	DM1 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.36 ±0.29	3681	⁶ BUKIN	78C	OLYA $e^+e^- \rightarrow$ hadrons
4.4 ±0.6	984	⁶ BESCH	74	CNTR $2\gamma p \rightarrow pK^+K^-$
4.67 ±0.72	681	⁶ BALAKIN	71	OSPK $e^+e^- \rightarrow$ hadrons
4.09 ±0.29		BIZOT	70	OSPK $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.07 ±0.13 ±0.01		⁷ HOID	20	RVUE $e^+e^- \rightarrow \pi^0 \gamma$
4.23 ±0.04 ±0.02		⁸ HOFERICH...	19	RVUE $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.249±0.015	1.7M	KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+K^-$
4.240±0.017	610k	KOZYREV	16	CMD3 $e^+e^- \rightarrow K_S^0 K_L^0$
4.37 ±0.02		LEES	13F	BABR $D^+ \rightarrow K^+K^-\pi^+$
4.24 ±0.02 ±0.03	542k	⁹ AKHMETSHIN	08	CMD2 1.02 $e^+e^- \rightarrow K^+K^-$
4.28 ±0.13	12540	¹⁰ AUBERT,B	05J	BABR $D^0 \rightarrow \bar{K}^0 K^+K^-$
4.45 ±0.06	271k	DIJKSTRA	86	SPEC 100 π^- Be
3.6 ±0.8	337	⁶ COOPER	78B	HBC 0.7–0.8 $\bar{p}p \rightarrow$ $K_S^0 K_L^0 \pi^+\pi^-$
4.5 ±0.50	1300	^{6,10} AKERLOF	77	SPEC 400 $pA \rightarrow K^+K^-X$
4.5 ±0.8	500	^{6,10} AYRES	74	ASPK 3–6 $\pi^- p \rightarrow$ $K^+K^-n, K^-p \rightarrow$ $K^+K^-\Lambda/\Sigma^0$
3.81 ±0.37		COSME	74B	OSPK $e^+e^- \rightarrow K_L^0 K_S^0$
3.8 ±0.7	454	⁶ BORENSTEIN	72	HBC 2.18 $K^-p \rightarrow K\bar{K}n$

- ¹ Average of KOZYREV 16 and KOZYREV 18 values taking into account the correlated uncertainties. Supersedes individual KOZYREV 16 and KOZYREV 18 results.
- ² Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.
- ³ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations.
- ⁴ Update of AKHMETSHIN 99D
- ⁵ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
- ⁶ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- ⁷ The values were extracted from a dispersively improved Breit-Wigner parameterization.
- ⁸ The values were extracted from a dispersively improved Breit-Wigner parameterization and do not include vacuum polarization.
- ⁹ Strongly correlated with AKHMETSHIN 04.
- ¹⁰ Systematic errors not evaluated.

NODE=M004W

OCCUR=2

NODE=M004W;LINKAGE=G

NODE=M004W;LINKAGE=E

NODE=M004W;LINKAGE=C

NODE=M004W;LINKAGE=GS

NODE=M004W;LINKAGE=AE

NODE=M004W;LINKAGE=D

NODE=M004W;LINKAGE=I

NODE=M004W;LINKAGE=K

NODE=M004W;LINKAGE=AH

NODE=M004W;LINKAGE=A

 $\phi(1020)$ DECAY MODES

NODE=M004215;NODE=M004

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 K^+K^-	(49.1 ±0.5)%	S=1.3
Γ_2 $K_L^0 K_S^0$	(33.9 ±0.4)%	S=1.2
Γ_3 $\rho\pi + \pi^+\pi^-\pi^0$	(15.4 ±0.4)%	S=1.2
Γ_4 $\rho\pi$		DESIG=16
Γ_5 $\pi^+\pi^-\pi^0$		DESIG=3
Γ_6 $\eta\gamma$	(1.301±0.024)%	S=1.2
Γ_7 $\pi^0\gamma$	(1.32 ±0.05) × 10 ⁻³	DESIG=7
Γ_8 $\ell^+\ell^-$	—	DESIG=256;OUR EVAL;→ UNCHECKED ←
Γ_9 e^+e^-	(2.979±0.033) × 10 ⁻⁴	S=1.2

DESIG=1

DESIG=2

DESIG=24

DESIG=16

DESIG=3

DESIG=4

DESIG=7

DESIG=256;OUR EVAL;→ UNCHECKED ←

DESIG=5

Γ_{10}	$\mu^+ \mu^-$	$(2.85 \pm 0.22) \times 10^{-4}$	S=1.2	DESIG=6
Γ_{11}	$\eta e^+ e^-$	$(1.08 \pm 0.04) \times 10^{-4}$		DESIG=17
Γ_{12}	$\pi^+ \pi^-$	$(7.3 \pm 1.3) \times 10^{-5}$		DESIG=8
Γ_{13}	$\omega \pi^0$	$(4.7 \pm 0.5) \times 10^{-5}$		DESIG=25
Γ_{14}	$\omega \gamma$	< 5	CL=84%	DESIG=10
Γ_{15}	$\rho \gamma$	$< 1.2 \times 10^{-5}$	CL=90%	DESIG=12
Γ_{16}	$\pi^+ \pi^- \gamma$	$(4.1 \pm 1.3) \times 10^{-5}$		DESIG=9
Γ_{17}	$f_0(980) \gamma$	$(3.22 \pm 0.19) \times 10^{-4}$	S=1.1	DESIG=20
Γ_{18}	$\pi^0 \pi^0 \gamma$	$(1.12 \pm 0.06) \times 10^{-4}$		DESIG=19
Γ_{19}	$\pi^+ \pi^- \pi^+ \pi^-$	$(3.9 \pm_{-2.2}^{+2.8}) \times 10^{-6}$		DESIG=15
Γ_{20}	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	$< 4.6 \times 10^{-6}$	CL=90%	DESIG=14
Γ_{21}	$\pi^0 e^+ e^-$	$(1.33 \pm_{-0.10}^{+0.07}) \times 10^{-5}$		DESIG=21
Γ_{22}	$\pi^0 \eta \gamma$	$(7.27 \pm 0.30) \times 10^{-5}$	S=1.5	DESIG=22
Γ_{23}	$a_0(980) \gamma$	$(7.6 \pm 0.6) \times 10^{-5}$		DESIG=23
Γ_{24}	$K^0 \bar{K}^0 \gamma$	$< 1.9 \times 10^{-8}$	CL=90%	DESIG=257
Γ_{25}	$\eta'(958) \gamma$	$(6.21 \pm 0.20) \times 10^{-5}$		DESIG=194
Γ_{26}	$\eta \pi^0 \pi^0 \gamma$	$< 2 \times 10^{-5}$	CL=90%	DESIG=195
Γ_{27}	$\mu^+ \mu^- \gamma$	$(1.4 \pm 0.5) \times 10^{-5}$		DESIG=196
Γ_{28}	$\rho \gamma \gamma$	$< 1.2 \times 10^{-4}$	CL=90%	DESIG=250
Γ_{29}	$\eta \pi^+ \pi^-$	$< 1.8 \times 10^{-5}$	CL=90%	DESIG=255
Γ_{30}	$\eta \mu^+ \mu^-$	$< 9.4 \times 10^{-6}$	CL=90%	DESIG=26
Γ_{31}	$\eta U \rightarrow \eta e^+ e^-$	$< 1 \times 10^{-6}$	CL=90%	DESIG=259
Γ_{32}	invisible	$< 1.7 \times 10^{-4}$	CL=90%	DESIG=260

Lepton Family number (LF) violating modes

Γ_{33}	$e^\pm \mu^\mp$	LF	$< 2 \times 10^{-6}$	CL=90%
---------------	-----------------	----	----------------------	--------

NODE=M004;CLUMP=A
DESIG=258

CONSTRAINED FIT INFORMATION

An overall fit to 30 branching ratios uses 80 measurements and one constraint to determine 14 parameters. The overall fit has a $\chi^2 = 61.8$ for 67 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-72									
x_3	-61	-12								
x_6	-19	17	2							
x_7	-11	11	1	8						
x_9	47	-50	-8	-35	-22					
x_{10}	-5	4	1	3	2	-9				
x_{12}	-3	3	0	2	1	-6	1			
x_{13}	-4	4	1	3	2	-8	1	0		
x_{17}	0	0	0	0	0	0	0	0	0	
x_{18}	-9	9	1	18	4	-18	2	1	1	0
x_{19}	-1	1	0	1	0	-2	0	0	0	0
x_{23}	0	0	0	0	0	0	0	0	0	0
x_{25}	-6	5	1	32	2	-11	1	1	1	0
	x_1	x_2	x_3	x_6	x_7	x_9	x_{10}	x_{12}	x_{13}	x_{17}
x_{19}	0									
x_{23}	0	0								
x_{25}	6	0	0							
	x_{18}	x_{19}	x_{23}							

$\phi(1020)$ PARTIAL WIDTHS

NODE=M004218

 $\Gamma(\eta\gamma)$ Γ_6

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

58.9 ± 0.5 ± 2.4	ACHASOV	00	SND $e^+e^- \rightarrow \eta\gamma$
------------------	---------	----	-------------------------------------

NODE=M004W6
NODE=M004W6 $\Gamma(\pi^0\gamma)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.40 ± 0.16 ^{+0.43} _{-0.40}	ACHASOV	00	SND $e^+e^- \rightarrow \pi^0\gamma$
---	---------	----	--------------------------------------

NODE=M004W7
NODE=M004W7 $\Gamma(\ell^+\ell^-)$ Γ_8

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.320 ± 0.017 ± 0.015	¹ AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow \mu^+\mu^-$
-----------------------	------------------------	----	---

¹ Weighted average of Γ_{ee} and $\sqrt{\Gamma_{ee}\Gamma_{\mu\mu}}$ from AMBROSINO 05 assuming lepton universality.

NODE=M004W5
NODE=M004W5

NODE=M004W5;LINKAGE=AM

 $\Gamma(e^+e^-)$ Γ_9

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1.27 ± 0.04 OUR EVALUATION**1.251 ± 0.021 OUR AVERAGE** Error includes scale factor of 1.1.

1.235 ± 0.006 ± 0.022	¹ AKHMETSHIN	11	CMD2 1.02 $e^+e^- \rightarrow \phi$
1.32 ± 0.05 ± 0.03	² AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow e^+e^-$
1.28 ± 0.05	AKHMETSHIN	95	CMD2 1.02 $e^+e^- \rightarrow \phi$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-$, $K_S^0K_L^0$, $\pi^+\pi^-\pi^0$, $\eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² From forward-backward asymmetry and using $\Gamma_{\text{total}} = 4.26 \pm 0.05$ MeV from the 2004 edition of this Review.

NODE=M004W8
NODE=M004W8

→ UNCHECKED ←

NODE=M004W8;LINKAGE=AK

NODE=M004W8;LINKAGE=AM

 $(\Gamma(e^+e^-) \times \Gamma(\mu^+\mu^-))^{1/2}$ $(\Gamma_9\Gamma_{10})^{1/2}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

1.320 ± 0.018 ± 0.017	AMBROSINO	05	KLOE 1.02 $e^+e^- \rightarrow \mu^+\mu^-$
------------------------------	-----------	----	---

NODE=M004W9
NODE=M004W9 $\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M004223

 $\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_9/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.6340 ± 0.0070 ± 0.0039		¹ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$
---------------------------------	--	-------------------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.669 ± 0.001 ± 0.023	1.7M	KOZYREV	18	CMD3 $e^+e^- \rightarrow K^+K^-$
-----------------------	------	---------	----	----------------------------------

¹ Using a phenomenological model based on KUHN 90 with a sum of Breit-Wigner resonances for $\rho(770)$, $\omega(782)$, $\phi(1020)$ and their higher mass excitations. The first error combines statistical and systematic uncertainties. The second one is due to the parametrization of the charged kaon form factor and mass calibration.

NODE=M004G01
NODE=M004G01

NODE=M004G01;LINKAGE=A

 $\Gamma(K_L^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_9/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.4200 ± 0.0033 ± 0.0123	28k	¹ LEES	14H	BABR $e^+e^- \rightarrow K_S^0K_L^0\gamma$
---------------------------------	-----	-------------------	-----	--

¹ Using a vector meson dominance model with contribution from $\phi(1020)$ and higher mass excitations of $\rho(770)$, $\omega(782)$, and $\phi(1020)$.

NODE=M004GXX
NODE=M004GXX

NODE=M004GXX;LINKAGE=A

 $[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)] \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_9/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

184.1 ± 2.1 ± 8.0	¹ LEES	21B	BABR 10.5 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
--------------------------	-------------------	-----	--

¹ From the cross section for $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ with contributions from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$.

NODE=M004R03
NODE=M004R03

NODE=M004R03;LINKAGE=A

$\phi(1020) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M004224

$$\Gamma(K^+K^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

14.64 ±0.27 OUR FIT Error includes scale factor of 1.4. $[(14.64 \pm 0.28) \times 10^{-5}]$ OUR 2023 FIT Scale factor = 1.4]

NODE=M004G10
NODE=M004G10

NEW


14.6 ±0.5 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.

15.789±0.541	1.7M	KOZYREV	18	CMD3	$e^+e^- \rightarrow K^+K^-$
14.27 ±0.05 ±0.31	542k	AKHMETSHIN	08	CMD2	$1.02 e^+e^- \rightarrow K^+K^-$
13.93 ±0.14 ±0.99	1000k	¹ ACHASOV	01E	SND	$e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$

NODE=M004G10;LINKAGE=AE

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

WEIGHTED AVERAGE
14.6±0.5 (Error scaled by 1.8)



$$\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

10.11 ±0.12 OUR FIT

10.07 ±0.13 OUR AVERAGE

10.078±0.223	610k	¹ KOZYREV	16	CMD3	$e^+e^- \rightarrow K_S^0 K_L^0$
10.01 ±0.04 ±0.17	272k	² AKHMETSHIN	04	CMD2	$e^+e^- \rightarrow K_L^0 K_S^0$
10.27 ±0.07 ±0.34	500k	³ ACHASOV	01E	SND	$e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$

NODE=M004G6
NODE=M004G6

¹ KOZYREV 16 also reports $\Gamma(e^+e^-) B(\phi \rightarrow K_S^0 K_L^0) = (0.428 \pm 0.001 \pm 0.009)$ keV.

² Update of AKHMETSHIN 99D

³ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

NODE=M004G6;LINKAGE=A
NODE=M004G;LINKAGE=GS
NODE=M004G6;LINKAGE=AE

$$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \Gamma_9/\Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.58 ±0.11 OUR FIT Error includes scale factor of 1.1.

4.51 ±0.14 OUR AVERAGE

4.51 ±0.16 ±0.11	105k	AKHMETSHIN	06	CMD2	$0.98-1.06 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.665±0.042±0.261	400k	¹ ACHASOV	01E	SND	$e^+e^- \rightarrow K^+K^-, K_S K_L, \pi^+\pi^-\pi^0$
4.35 ±0.27 ±0.08	11169	² AKHMETSHIN	98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

NODE=M004G7
NODE=M004G7

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.38 ±0.12		BENAYOUN	10	RVUE	$0.4-1.05 e^+e^-$
4.30 ±0.08 ±0.21		³ AUBERT,B	04N	BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

- ¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S K_L$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.
- ² Recalculated by us from the cross section in the peak.
- ³ Superseded by LEES 21B.

NODE=M004G7;LINKAGE=AE

NODE=M004G;LINKAGE=B
NODE=M004G7;LINKAGE=A

NODE=M004G2
NODE=M004G2

NEW

$\Gamma(\eta\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_9/\Gamma$


VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.88 ± 0.07 OUR FIT				Error includes scale factor of 1.2. $[(3.88 \pm 0.07) \times 10^{-6}]$ OUR 2023 FIT Scale factor = 1.2]
3.93 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
4.050 ± 0.067 ± 0.118	33k	¹ ACHASOV 07B	SND	0.6–1.38 $e^+e^- \rightarrow \eta\gamma$
4.093 ^{+0.040} _{-0.043} ± 0.247	17.4k	² AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \eta\gamma$
3.850 ± 0.041 ± 0.159	23k	^{3,4} AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
4.00 ± 0.04 ± 0.11		⁵ ACHASOV 00	SND	$e^+e^- \rightarrow \eta\gamma$
3.53 ± 0.08 ± 0.17	2200	^{6,7} AKHMETSHIN 99F	CMD2	$e^+e^- \rightarrow \eta\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.19 ± 0.06		⁸ BENAYOUN 10	RVUE	0.4–1.05 e^+e^-

NODE=M004G2;LINKAGE=AH

NODE=M004G2;LINKAGE=AK
NODE=M004G;LINKAGE=AK
NODE=M004G;LINKAGE=BQ

NODE=M004G2;LINKAGE=A
NODE=M004G;LINKAGE=A
NODE=M004G2;LINKAGE=C
NODE=M004G7;LINKAGE=BE

- ¹ From a combined fit of $\sigma(e^+e^- \rightarrow \eta\gamma)$ with $\eta \rightarrow 3\pi^0$ and $\eta \rightarrow \pi^+\pi^-\pi^0$, and fixing $B(\eta \rightarrow 3\pi^0) / B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.44 \pm 0.04$. Recalculated by us from the cross section at the peak. Supersedes ACHASOV 00D and ACHASOV 06A.
- ² From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.
- ³ From the $\eta \rightarrow 3\pi^0$ decay and using $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.
- ⁴ The combined fit from 600 to 1380 MeV taking into account $\rho(770)$, $\omega(782)$, $\phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).
- ⁵ From the $\eta \rightarrow 2\gamma$ decay and using $B(\eta \rightarrow 2\gamma) = (39.21 \pm 0.34) \times 10^{-2}$.
- ⁶ Recalculated by the authors from the cross section in the peak.
- ⁷ From the $\eta \rightarrow \pi^+\pi^-\pi^0$ decay and using $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.1 \pm 0.5) \times 10^{-2}$.
- ⁸ A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-$, $\pi^+\pi^-\pi^0$, $\pi^0\gamma$, $\eta\gamma$ data.



$\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
3.94 ± 0.16 OUR FIT				
3.95 ± 0.17 OUR AVERAGE				
4.04 ± 0.09 ± 0.19		¹ ACHASOV 16A	SND	0.60–1.38 $e^+e^- \rightarrow \pi^0\gamma$
3.75 ± 0.11 ± 0.29	18k	AKHMETSHIN 05	CMD2	0.60-1.38 $e^+e^- \rightarrow \pi^0\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.29 ± 0.11		² BENAYOUN 10	RVUE	0.4–1.05 e^+e^-
3.67 ± 0.10 ^{+0.27} _{-0.25}		³ ACHASOV 00	SND	$e^+e^- \rightarrow \pi^0\gamma$

NODE=M004G3
NODE=M004G3

- ¹ From the VMD model with the interfering $\rho(770)$, $\omega(782)$, $\phi(1020)$ resonances, and an additional resonance describing the total contribution of the $\rho(1450)$ and $\omega(1420)$ states. Supersedes ACHASOV 00.
² A simultaneous fit of $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma$ data.
³ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\pi^0 \rightarrow 2\gamma) = (98.798 \pm 0.032) \times 10^{-2}$.

NODE=M004G3;LINKAGE=B

NODE=M004G3;LINKAGE=BE
 NODE=M004G3;LINKAGE=A

$\Gamma(\mu^+\mu^-)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{10}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

8.5 ± 0.6 OUR FIT Error includes scale factor of 1.2. [(8.5 ± 0.6) × 10⁻⁸ OUR 2023 FIT]

NODE=M004G5
 NODE=M004G5

NEW


8.8 ± 0.9 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

8.36 ± 0.59 ± 0.37	ACHASOV	01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$
9.9 ± 1.4 ± 0.9	¹ ACHASOV	99C	SND	$e^+e^- \rightarrow \mu^+\mu^-$
14.4 ± 3.0	² VASSERMAN	81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
8.6 ± 5.9	² AUGUSTIN	73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M004G5
 NODE=M004G5

- ¹ Recalculated by the authors from the cross section in the peak.
² Recalculated by us from the cross section in the peak.

NODE=M004G5;LINKAGE=A
 NODE=M004G5;LINKAGE=B



$\Gamma(\pi^+\pi^-)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{12}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

2.2 ± 0.4 OUR FIT

2.2 ± 0.4 OUR AVERAGE

2.1 ± 0.3 ± 0.3	¹ ACHASOV	00C	SND	$e^+e^- \rightarrow \pi^+\pi^-$
1.95 ^{+1.15} _{-0.87}	² GOLUBEV	86	ND	$e^+e^- \rightarrow \pi^+\pi^-$
6.01 ^{+3.19} _{-2.51}	² VASSERMAN	81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$

NODE=M004G4
 NODE=M004G4

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.31 ± 0.99 ³ BENAYOUN 13 RVUE 0.4-1.05 e^+e^-

- ¹ Recalculated by the authors from the cross section in the peak.
² Recalculated by us from the cross section in the peak.
³ A simultaneous fit to $e^+e^- \rightarrow \pi^+\pi^-, \pi^+\pi^-\pi^0, \pi^0\gamma, \eta\gamma, K\bar{K}$, and $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$ data.

NODE=M004G4;LINKAGE=AC
 NODE=M004G4;LINKAGE=B
 NODE=M004G4;LINKAGE=A

$\Gamma(\omega\pi^0)/\Gamma_{total} \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{13}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

1.40 ± 0.15 OUR FIT

1.37 ± 0.17 ± 0.01 ^{1,2} AMBROSINO 08G KLOE $e^+e^- \rightarrow \pi^+\pi^-\pi^0, 2\pi^0\gamma$

- ¹ Recalculated by the authors from the cross section at the peak.
² AMBROSINO 08G reports $[\Gamma(\phi(1020) \rightarrow \omega\pi^0)/\Gamma_{total} \times \Gamma(\phi(1020) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (1.22 \pm 0.13 \pm 0.08) \times 10^{-8}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M004G11
 NODE=M004G11

NODE=M004G11;LINKAGE=AB
 NODE=M004G11;LINKAGE=AM

$$\frac{\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}}{\Gamma_{18}/\Gamma \times \Gamma_9/\Gamma}$$

VALUE (units 10^{-8}) DOCUMENT ID TECN COMMENT

3.34±0.17 OUR FIT

3.33^{+0.04+0.19}_{-0.09-0.20} ¹ AMBROSINO 07 KLOE $e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ Calculated by the authors from the cross section at the peak.

NODE=M004G9
NODE=M004G9

NODE=M004G9;LINKAGE=AM

$$\frac{\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}}{\Gamma_{19}/\Gamma \times \Gamma_9/\Gamma}$$

VALUE (units 10^{-9}) EVTS DOCUMENT ID TECN COMMENT

1.2^{+0.8}_{-0.7} OUR FIT

1.17±0.52±0.64 3285 ¹ AKHMETSHIN 00E CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

¹ Recalculated by the authors from the cross section in the peak.

NODE=M004G8
NODE=M004G8

NODE=M004G8;LINKAGE=A

$\phi(1020)$ BRANCHING RATIOS

$$\frac{\Gamma(K^+K^-)/\Gamma_{\text{total}}}{\Gamma_1/\Gamma}$$

VALUE EVTS DOCUMENT ID TECN COMMENT

0.491±0.005 OUR FIT Error includes scale factor of 1.3. [0.491±0.005 OUR 2023 FIT Scale factor = 1.3]

0.493±0.010 OUR AVERAGE

0.492±0.012 2913 AKHMETSHIN 95 CMD2 $e^+e^- \rightarrow K^+K^-$
0.44 ±0.05 321 KALBFLEISCH 76 HBC $2.18 K^-p \rightarrow \Lambda K^+K^-$
0.49 ±0.06 270 DEGROOT 74 HBC $4.2 K^-p \rightarrow \Lambda\phi$
0.540±0.034 565 BALAKIN 71 OSPK $e^+e^- \rightarrow K^+K^-$
0.48 ±0.04 252 LINDSEY 66 HBC $2.1-2.7 K^-p \rightarrow \Lambda K^+K^-$

••• We do not use the following data for averages, fits, limits, etc. •••

0.493±0.003±0.007 ¹ AKHMETSHIN 11 CMD2 $1.02 e^+e^- \rightarrow K^+K^-$
0.476±0.017 1000k ² ACHASOV 01E SND $e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

NODE=M004220

NODE=M004R1
NODE=M004R1
NEW

NODE=M004R1;LINKAGE=AK

NODE=M004R1;LINKAGE=B2

$$\frac{\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}}}{\Gamma_2/\Gamma}$$

VALUE EVTS DOCUMENT ID TECN COMMENT

0.339±0.004 OUR FIT Error includes scale factor of 1.2. [0.339±0.004 OUR 2023 FIT Scale factor = 1.2]

0.331±0.009 OUR AVERAGE

0.335±0.010 40644 AKHMETSHIN 95 CMD2 $e^+e^- \rightarrow K_L^0 K_S^0$
0.326±0.035 DOLINSKY 91 ND $e^+e^- \rightarrow K_L^0 K_S^0$
0.310±0.024 DRUZHININ 84 ND $e^+e^- \rightarrow K_L^0 K_S^0$

••• We do not use the following data for averages, fits, limits, etc. •••

0.336±0.002±0.006 ¹ AKHMETSHIN 11 CMD2 $1.02 e^+e^- \rightarrow K_S^0 K_L^0$
0.351±0.013 500k ² ACHASOV 01E SND $e^+e^- \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0$

0.27 ±0.03 133 KALBFLEISCH 76 HBC $2.18 K^-p \rightarrow \Lambda K_L^0 K_S^0$
0.257±0.030 95 ³ BALAKIN 71 OSPK $e^+e^- \rightarrow K_L^0 K_S^0$

0.40 ±0.04 167 LINDSEY 66 HBC $2.1-2.7 K^-p \rightarrow \Lambda K_L^0 K_S^0$

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Balakin error increased by Paul.

NODE=M004R2
NODE=M004R2
NEW

NODE=M004R2;LINKAGE=AK

NODE=M004R2;LINKAGE=B2
NODE=M004R2;LINKAGE=01

$$\frac{\Gamma(K_L^0 K_S^0)/\Gamma(K^+K^-)}{\Gamma_2/\Gamma_1}$$

VALUE EVTS DOCUMENT ID TECN COMMENT

0.690±0.014 OUR FIT Error includes scale factor of 1.3. [0.690±0.015 OUR 2023 FIT Scale factor = 1.3]

0.740±0.031 OUR AVERAGE

0.70 ±0.06 2732 BUKIN 78C OLYA $e^+e^- \rightarrow K_L^0 K_S^0$
0.82 ±0.08 LOSTY 78 HBC $4.2 K^-p \rightarrow \phi$ hyperon
0.71 ±0.05 LAVEN 77 HBC $10 K^-p \rightarrow K^+K^-\Lambda$
0.71 ±0.08 LYONS 77 HBC $3-4 K^-p \rightarrow \Lambda\phi$
0.89 ±0.10 144 AGUILAR-... 72B HBC $3.9, 4.6 K^-p$

••• We do not use the following data for averages, fits, limits, etc. •••

0.638±0.022 2.3M ¹ KOZYREV 18 CMD3 $e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$
0.68 ±0.03 ² AKHMETSHIN 95 CMD2 $e^+e^- \rightarrow K_L^0 K_S^0, K^+K^-$

NODE=M004R19
NODE=M004R19
NEW

¹ The prediction taking into account phase-space difference, radiative corrections, isospin breaking, and the Sommerfeld-Gamow-Sakharov factor gives 0.630.

² Theoretical analysis of BRAMON 00 taking into account phase-space difference, electromagnetic radiative corrections, as well as isospin breaking, predicts 0.62. FLOREZ-BAEZ 08 predicts 0.63 considering also structure-dependent radiative corrections. FIS-CHBACH 02 calculates additional corrections caused by the close threshold and predicts 0.68. See also BENAYOUN 01 and DUBYNISKIY 07. BENAYOUN 12 obtains 0.71 ± 0.01 in the HLS model.

NODE=M004R19;LINKAGE=A

NODE=M004R19;LINKAGE=KH

 $\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$ $\Gamma_2/(\Gamma_1+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.408 ± 0.005 OUR FIT	Error includes scale factor of 1.3. [0.408 ± 0.005 OUR 2023 FIT Scale factor = 1.3]			
0.45 ± 0.04 OUR AVERAGE	Error includes scale factor of 1.3.			
0.44 ± 0.07		¹ LONDON	66 HBC	2.24 $K^- p \rightarrow \Lambda K \bar{K}$
0.48 ± 0.07	52	BADIER	65B HBC	3 $K^- p$
0.40 ± 0.10	34	SCHLEIN	63 HBC	1.95 $K^- p \rightarrow \Lambda K \bar{K}$

NODE=M004R5
NODE=M004R5
NEW

¹ This is probably not affected by their controversial background subtraction; the value is from their numbers of $K_1 K_2$ vs $K^+ K^-$ events.

NODE=M004R5;LINKAGE=01

 $[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.154 ± 0.004 OUR FIT	Error includes scale factor of 1.2.			
0.151 ± 0.009 OUR AVERAGE	Error includes scale factor of 1.7.			
0.161 ± 0.008	11761	AKHMETSHIN 95	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.143 ± 0.007		DOLINSKY 91	ND	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.155 ± 0.002 ± 0.005		¹ AKHMETSHIN 11	CMD2	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.159 ± 0.008	400k	² ACHASOV 01E	SND	$e^+ e^- \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0$
0.145 ± 0.009 ± 0.003	11169	³ AKHMETSHIN 98	CMD2	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.139 ± 0.007		⁴ PARROUR 76B	OSPK	$e^+ e^-$

NODE=M004R3
NODE=M004R3

¹ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+ K^-, K_S^0 K_L^0, \pi^+ \pi^- \pi^0, \eta, \gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

² Using $B(\phi \rightarrow e^+ e^-) = (2.93 \pm 0.14) \times 10^{-4}$.

³ Using $B(\phi \rightarrow e^+ e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁴ Using $\Gamma(\phi) = 4.1$ MeV. If interference between the $\rho\pi$ and 3π modes is neglected, the fraction of the $\rho\pi$ is more than 80% at the 90% confidence level.

NODE=M004R3;LINKAGE=AK

NODE=M004R3;LINKAGE=B2

NODE=M004R;LINKAGE=8D

NODE=M004R3;LINKAGE=E

 $[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma(K^+ K^-)$ Γ_3/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.313 ± 0.010 OUR FIT	Error includes scale factor of 1.2.			
0.28 ± 0.09	34	AGUILAR-...	72B HBC	3.9, 4.6 $K^- p$

NODE=M004R20
NODE=M004R20 $[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma(K\bar{K})$ $\Gamma_3/(\Gamma_1+\Gamma_2)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.185 ± 0.005 OUR FIT	Error includes scale factor of 1.2.		
0.24 ± 0.04 OUR AVERAGE	Error includes scale factor of 1.2.		
0.237 ± 0.039	CERRADA 77B	HBC	4.2 $K^- p \rightarrow \Lambda 3\pi$
0.30 ± 0.15	LONDON 66	HBC	2.24 $K^- p \rightarrow \Lambda \pi^+ \pi^- \pi^0$

NODE=M004R6
NODE=M004R6 $[\Gamma(\rho\pi) + \Gamma(\pi^+ \pi^- \pi^0)]/\Gamma(K_L^0 K_S^0)$ Γ_3/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.453 ± 0.012 OUR FIT	Error includes scale factor of 1.1.			
0.51 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.1.			
0.56 ± 0.07	3681	BUKIN 78C	OLYA	$e^+ e^- \rightarrow K_L^0 K_S^0, \pi^+ \pi^- \pi^0$
0.47 ± 0.06	516	COSME 74	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M004R7
NODE=M004R7 $\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
≈ 0.0087		1.98M	^{1,2} ALOISIO 03	KLOE	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
< 0.0006	90		³ ACHASOV 02	SND	1.02 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
< 0.23	90		³ CORDIER 80	DM1	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
< 0.20	90		³ PARROUR 76B	OSPK	$e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

NODE=M004R46
NODE=M004R46

¹ From a fit without limitations on charged and neutral ρ masses and widths.

² Adding the direct and $\omega\pi$ contributions and considering the interference between the $\rho\pi$ and $\pi^+ \pi^- \pi^0$.

³ Neglecting the interference between the $\rho\pi$ and $\pi^+ \pi^- \pi^0$.

NODE=M004R;LINKAGE=L1

NODE=M004R;LINKAGE=L2

NODE=M004R;LINKAGE=46

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.301 ± 0.024 OUR FIT Error includes scale factor of 1.2. $[(1.301 \pm 0.025) \times 10^{-2} \text{ OUR } 2023 \text{ FIT Scale factor} = 1.2]$

1.26 ± 0.04 OUR AVERAGE

1.246 ± 0.025 ± 0.057	10k	¹ ACHASOV	98F	SND	$e^+e^- \rightarrow 7\gamma$
1.18 ± 0.11	279	² AKHMETSHIN	95	CMD2	$e^+e^- \rightarrow \pi^+\pi^-3\gamma$
1.30 ± 0.06		³ DRUZHININ	84	ND	$e^+e^- \rightarrow 3\gamma$
1.4 ± 0.2		⁴ DRUZHININ	84	ND	$e^+e^- \rightarrow 6\gamma$
0.88 ± 0.20	290	KURDADZE	83C	OLYA	$e^+e^- \rightarrow 3\gamma$
1.35 ± 0.29		ANDREWS	77	CNTR	6.7-10 γ Cu
1.5 ± 0.4	54	³ COSME	76	OSPK	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.38 ± 0.02 ± 0.02		⁵ AKHMETSHIN	11	CMD2	$1.02 e^+e^- \rightarrow \eta\gamma$
1.36 ± 0.05 ± 0.01	33k	⁶ ACHASOV	07B	SND	$0.6-1.38 e^+e^- \rightarrow \eta\gamma$
1.373 ± 0.014 ± 0.085	17.4k	^{7,8} AKHMETSHIN	05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$
1.287 ± 0.013 ± 0.063		^{9,10} AKHMETSHIN	01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.338 ± 0.012 ± 0.052		¹¹ ACHASOV	00	SND	$e^+e^- \rightarrow \eta\gamma$
1.18 ± 0.03 ± 0.06	2200	¹² AKHMETSHIN	99F	CMD2	$e^+e^- \rightarrow \eta\gamma$
1.21 ± 0.07		¹³ BENAYOUN	96	RVUE	$0.54-1.04 e^+e^- \rightarrow \eta\gamma$

¹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.2 \pm 0.4) \times 10^{-2}$.

² From $\pi^+\pi^-\pi^0$ decay mode of η .

³ From 2γ decay mode of η .

⁴ From $3\pi^0$ decay mode of η .

⁵ Combined analysis of the CMD-2 data on $\phi \rightarrow K^+K^-, K_S^0 K_L^0, \pi^+\pi^-\pi^0, \eta\gamma$ assuming that the sum of their branching fractions is 0.99741 ± 0.00007 .

⁶ ACHASOV 07B reports $[\Gamma(\phi(1020) \rightarrow \eta\gamma)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow e^+e^-)] = (4.050 \pm 0.067 \pm 0.118) \times 10^{-6}$ which we divide by our best value $B(\phi(1020) \rightarrow e^+e^-) = (2.979 \pm 0.033) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Supersedes ACHASOV 00D and ACHASOV 06A.

⁷ Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$ and $B(\eta \rightarrow \gamma\gamma) = 39.43 \pm 0.26\%$.

⁸ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\eta\gamma)/\Gamma_{\text{total}}^2$.

⁹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$ and $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$.

¹⁰ The combined fit from 600 to 1380 MeV taking into account $\rho(770), \omega(782), \phi(1020)$, and $\rho(1450)$ (mass and width fixed at 1450 MeV and 310 MeV respectively).

¹¹ From the $\eta \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹² From $\pi^+\pi^-\pi^0$ decay mode of η and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

¹³ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

NODE=M004R11
NODE=M004R11

NEW

OCCUR=2

NODE=M004R11;LINKAGE=AC
NODE=M004R11;LINKAGE=Z3
NODE=M004R11;LINKAGE=A
NODE=M004R11;LINKAGE=C
NODE=M004R11;LINKAGE=AN

NODE=M004R11;LINKAGE=AO

NODE=M004R11;LINKAGE=AH
NODE=M004R11;LINKAGE=AK
NODE=M004R;LINKAGE=AK
NODE=M004R;LINKAGE=BQ

NODE=M004R;LINKAGE=GA
NODE=M004R;LINKAGE=FF
NODE=M004R;LINKAGE=TS

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.32 ± 0.05 OUR FIT**1.31 ± 0.13 OUR AVERAGE**

1.30 ± 0.13		DRUZHININ	84	ND	$e^+e^- \rightarrow 3\gamma$
1.4 ± 0.5	32	COSME	76	OSPK	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.367 ± 0.072		¹ ACHASOV	16A	SND	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$
1.258 ± 0.037 ± 0.077	18k	^{2,3} AKHMETSHIN	05	CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$
1.226 ± 0.036 ^{+0.096} _{-0.089}		⁴ ACHASOV	00	SND	$e^+e^- \rightarrow \pi^0\gamma$
1.26 ± 0.17		⁵ BENAYOUN	96	RVUE	$0.54-1.04 e^+e^- \rightarrow \pi^0\gamma$

¹ Using $B(\phi \rightarrow e^+e^-)$ from PDG 15. Supersedes ACHASOV 00.

² Using $B(\phi \rightarrow e^+e^-) = (2.98 \pm 0.04) \times 10^{-4}$.

³ Not independent of the corresponding $\Gamma(e^+e^-) \times \Gamma(\pi^0\gamma)/\Gamma_{\text{total}}^2$.

⁴ From the $\pi^0 \rightarrow 2\gamma$ decay and using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

⁵ Reanalysis of DRUZHININ 84, DOLINSKY 89, and DOLINSKY 91 taking into account a triangle anomaly contribution.

OCCUR=3

NODE=M004R17;LINKAGE=D
NODE=M004R17;LINKAGE=AH
NODE=M004R17;LINKAGE=AK
NODE=M004R;LINKAGE=3G
NODE=M004R17;LINKAGE=TS

 $\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$ Γ_6/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.9 ± 0.3 ^{+0.7} _{-0.8}	ACHASOV	00	SND	$e^+e^- \rightarrow \eta\gamma, \pi^0\gamma$
--	---------	----	-----	--

NODE=M004R42
NODE=M004R42

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.979 ± 0.033 OUR FIT				Error includes scale factor of 1.2. $[(2.979 \pm 0.033) \times 10^{-4}$ OUR 2023 FIT Scale factor = 1.3]
2.98 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.1.
2.93 ± 0.14	1900k	¹ ACHASOV	01E SND	$e^+e^- \rightarrow K^+K^-, K_S^0K_L^0, \pi^+\pi^-\pi^0$
2.88 ± 0.09	55600	AKHMETSHIN 95	CMD2	$e^+e^- \rightarrow$ hadrons
3.00 ± 0.21	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow$ hadrons
3.10 ± 0.14		² PARROUR 76	OSPK	e^+e^-
3.3 ± 0.3		COSME 74	OSPK	$e^+e^- \rightarrow$ hadrons
2.81 ± 0.25	681	BALAKIN 71	OSPK	$e^+e^- \rightarrow$ hadrons
3.50 ± 0.27		CHATELUS 71	OSPK	e^+e^-

NODE=M004R16

NODE=M004R16

NEW

¹ From the combined fit assuming that the total $\phi(1020)$ production cross section is saturated by those of K^+K^- , $K_S^0K_L^0$, $\pi^+\pi^-\pi^0$, and $\eta\gamma$ decays modes and using ACHASOV 00B for the $\eta\gamma$ decay mode.

NODE=M004R16;LINKAGE=AE

² Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.

NODE=M004R16;LINKAGE=E

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
2.85 ± 0.22 OUR FIT			Error includes scale factor of 1.2. $[(2.85 \pm 0.19) \times 10^{-4}$ OUR 2023 FIT]
2.5 ± 0.4 OUR AVERAGE			
2.69 ± 0.46	¹ HAYES 71	CNTR	$8.3, 9.8 \gamma C \rightarrow \mu^+\mu^-X$
2.17 ± 0.60	¹ EARLES 70	CNTR	$6.0 \gamma C \rightarrow \mu^+\mu^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.87 ± 0.20 ± 0.14	² ACHASOV 01G	SND	$e^+e^- \rightarrow \mu^+\mu^-$
3.30 ± 0.45 ± 0.32	³ ACHASOV 99C	SND	$e^+e^- \rightarrow \mu^+\mu^-$
4.83 ± 1.02	⁴ VASSERMAN 81	OLYA	$e^+e^- \rightarrow \mu^+\mu^-$
2.87 ± 1.98	⁴ AUGUSTIN 73	OSPK	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M004R10

NODE=M004R10

NEW

¹ Neglecting interference between resonance and continuum.

NODE=M004R10;LINKAGE=A

² Using $B(\phi \rightarrow e^+e^-) = (2.91 \pm 0.07) \times 10^{-4}$.

NODE=M004R;LINKAGE=GZ

³ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

NODE=M004R10;LINKAGE=8D

⁴ Recalculated by us using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

NODE=M004R;LINKAGE=VA

 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.08 ± 0.04 OUR AVERAGE				
1.075 ± 0.007 ± 0.038	30k	¹ BABUSCI 15	KLOE	$1.02 e^+e^- \rightarrow \eta e^+e^-$
1.19 ± 0.19 ± 0.12	213	² ACHASOV 01B	SND	$e^+e^- \rightarrow \eta e^+e^-$
1.14 ± 0.10 ± 0.06	355	³ AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.13 ± 0.14 ± 0.07	183	⁴ AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
1.21 ± 0.14 ± 0.09	130	⁵ AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
1.04 ± 0.20 ± 0.08	42	⁶ AKHMETSHIN 01	CMD2	$e^+e^- \rightarrow \eta e^+e^-$
1.3 ^{+0.8} _{-0.6}	7	GOLUBEV 85	ND	$e^+e^- \rightarrow \eta e^+e^-$

NODE=M004R24

NODE=M004R24

OCCUR=2

OCCUR=3

OCCUR=4

¹ Using $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.23)\%$ from PDG 12.

NODE=M004R24;LINKAGE=A

² Using $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.32)\%$, $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06)\%$, and $B(\phi \rightarrow e^+e^-) = (3.00 \pm 0.06) \times 10^{-4}$.

NODE=M004R;LINKAGE=VM

³ The average of the branching ratios separately obtained from the $\eta \rightarrow \gamma\gamma$, $3\pi^0$, $\pi^+\pi^-\pi^0$ decays.

NODE=M004R;LINKAGE=H1

⁴ From $\eta \rightarrow \gamma\gamma$ decays and using $B(\eta \rightarrow \gamma\gamma) = (39.33 \pm 0.25) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

NODE=M004R;LINKAGE=H2

⁵ From $\eta \rightarrow 3\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\eta \rightarrow 3\pi^0) = (32.24 \pm 0.29) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

NODE=M004R;LINKAGE=H3

⁶ From $\eta \rightarrow \pi^+\pi^-\pi^0$ decays and using $B(\pi^0 \rightarrow \gamma\gamma) = (98.798 \pm 0.033) \times 10^{-2}$, $B(\pi^0 \rightarrow e^+e^-\gamma) = (1.198 \pm 0.032) \times 10^{-2}$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (23.0 \pm 0.4) \times 10^{-2}$, $B(\phi \rightarrow \pi^+\pi^-\pi^0) = (15.5 \pm 0.6) \times 10^{-2}$, and $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$.

NODE=M004R;LINKAGE=H4

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M004R18
 NODE=M004R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.71 \pm 0.11 \pm 0.09$		¹ ACHASOV	00C	SND	$e^+e^- \rightarrow \pi^+\pi^-$
$0.65^{+0.38}_{-0.29}$		¹ GOLUBEV	86	ND	$e^+e^- \rightarrow \pi^+\pi^-$
$2.01^{+1.07}_{-0.84}$		¹ VASSERMAN	81	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
<6.6	95	BUKIN	78B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$
<2.7	95	ALVENSLEB...	72	CNTR	$6.7 \gamma C \rightarrow C\pi^+\pi^-$

¹ Using $B(\phi \rightarrow e^+e^-) = (2.99 \pm 0.08) \times 10^{-4}$.

NODE=M004R18;LINKAGE=8D

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M004R28
 NODE=M004R28

4.7 ± 0.5 OUR FIT

5.2^{+1.3}_{-1.1}

^{1,2} AULCHENKO 00A SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.4 ± 0.6		³ AMBROSINO	08G	KLOE	$e^+e^- \rightarrow \pi^+\pi^-2\pi^0, 2\pi^0\gamma$
~ 5.4		⁴ ACHASOV	00E	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$5.5^{+1.6}_{-1.4} \pm 0.3$		^{2,5} AULCHENKO	00A	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$
$4.8^{+1.9}_{-1.7} \pm 0.8$		⁴ ACHASOV	99	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

¹ Using the 1996 and 1998 data.

² (2.3 ± 0.3)% correction for other decay modes of the $\omega(782)$ applied.

³ Not independent of the corresponding $\Gamma(\omega\pi^0) \times \Gamma(e^+e^-) / \Gamma^2(\text{total})$.

⁴ Using the 1996 data.

⁵ Using the 1998 data.

NODE=M004R28;LINKAGE=K2
 NODE=M004R28;LINKAGE=K3
 NODE=M004R28;LINKAGE=AM
 NODE=M004R28;LINKAGE=VS
 NODE=M004R28;LINKAGE=K1

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M004R14
 NODE=M004R14

<0.05 84 LINDSEY 66 HBC 2.1–2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals

 $\Gamma(\rho\gamma)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M004R15
 NODE=M004R15

< 0.12 90 ¹ AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7	90	AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<200	84	LINDSEY 66	HBC	2.1–2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals

¹ Supersedes AKHMETSHIN 97C.

NODE=M004R;LINKAGE=1N

 $\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M004R12
 NODE=M004R12

0.41 ± 0.12 ± 0.04 30175 ¹ AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.3	90	² AKHMETSHIN 97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<600	90	KALBFLEISCH 75	HBC	2.18 $K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$
< 70	90	COSME 74	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<400	90	LINDSEY 65	HBC	2.1–2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-$ neutrals

¹ For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible. Supersedes AKHMETSHIN 97C.

² For $E_\gamma > 20$ MeV and assuming that $B(\phi(1020) \rightarrow f_0(980)\gamma)$ is negligible.

NODE=M004R;LINKAGE=2N

NODE=M004R12;LINKAGE=Z3

 $\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M004R30
 NODE=M004R30

3.22 ± 0.19 OUR FIT Error includes scale factor of 1.1.

3.21 ± 0.19 OUR AVERAGE

$3.21^{+0.03}_{-0.09} \pm 0.18$		¹ AMBROSINO 07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
$2.90 \pm 0.21 \pm 1.54$		² AKHMETSHIN 99C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma, \pi^0\pi^0\gamma$

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.47±0.21	2438	³ ALOISIO	02D	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
3.5 ±0.3 ^{+1.3} _{-0.5}	419	^{4,5} ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.93±0.46±0.50	27188	⁶ AKHMETSHIN	99B	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
3.05±0.25±0.72	268	⁷ AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.5 ±0.5	268	⁸ AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
3.42±0.30±0.36	164	⁴ ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
< 1	90	⁹ AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 7	90	¹⁰ AKHMETSHIN	97C	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
< 20	90	DRUZHININ	87	ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

OCCUR=2

OCCUR=2

¹ Obtained by the authors taking into account the $\pi^+\pi^-$ decay mode. Includes a component due to $\pi\pi$ production via the $f_0(500)$ meson. Supersedes ALOISIO 02D.

NODE=M004R30;LINKAGE=MB

² From the combined fit of the photon spectra in the reactions $e^+e^- \rightarrow \pi^+\pi^-\gamma, \pi^0\pi^0\gamma$.

NODE=M004R;LINKAGE=SL

³ From the negative interference with the $f_0(500)$ meson of AITALA 01B using the ACHASOV 89 parameterization for the $f_0(980)$, a Breit-Wigner for the $f_0(500)$, and ACHASOV 01F for the $\rho\pi$ contribution. Superseded by AMBROSINO 07.

NODE=M004R;LINKAGE=KD

⁴ Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.

NODE=M004R;LINKAGE=AI

⁵ Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

NODE=M004R;LINKAGE=U8

⁶ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.

NODE=M004R;LINKAGE=3N

⁷ Neglecting other intermediate mechanisms ($\rho\pi, \sigma\gamma$).

NODE=M004R;LINKAGE=SM

⁸ A narrow pole fit taking into account $f_0(980)$ and $f_0(1200)$ intermediate mechanisms.

NODE=M004R;LINKAGE=ST

⁹ For destructive interference with the Bremsstrahlung process

NODE=M004R30;LINKAGE=A

¹⁰ For constructive interference with the Bremsstrahlung process

NODE=M004R30;LINKAGE=B

$\Gamma(f_0(980)\gamma)/\Gamma(\eta\gamma)$

 Γ_{17}/Γ_6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48±0.15 OUR FIT	Error includes scale factor of 1.1.			

NODE=M004R44
NODE=M004R44

2.6 ±0.2 ^{+0.8}_{-0.3}	419	¹ ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
--	-----	----------------------	-----	-----	---------------------------------------

¹ Assuming that the $\pi^0\pi^0\gamma$ final state is completely determined by the $f_0\gamma$ mechanism, neglecting the decay $B(\phi \rightarrow K\bar{K}\gamma)$ and using $B(f_0 \rightarrow \pi^+\pi^-) = 2B(f_0 \rightarrow \pi^0\pi^0)$.

NODE=M004R44;LINKAGE=AI

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$

 Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ±0.06 OUR AVERAGE					

NODE=M004R26
NODE=M004R26

1.07 ^{+0.01} _{-0.03} ^{+0.06} _{-0.06}		¹ AMBROSINO	07	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
--	--	------------------------	----	------	---------------------------------------

1.08 ±0.17 ±0.09	268	AKHMETSHIN	99C	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------------------	-----	------------	-----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.09 ±0.03 ±0.05	2438	ALOISIO	02D	KLOE	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
------------------	------	---------	-----	------	---------------------------------------

1.158±0.093±0.052	419	^{2,3} ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
-------------------	-----	------------------------	-----	-----	---------------------------------------

<10	90	DRUZHININ	87	ND	$e^+e^- \rightarrow 5\gamma$
-----	----	-----------	----	----	------------------------------

¹ Supersedes ALOISIO 02D.

NODE=M004R26;LINKAGE=MB

² Using the value $B(\phi \rightarrow \eta\gamma) = (1.338 \pm 0.053) \times 10^{-2}$.

NODE=M004R26;LINKAGE=U8

³ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

NODE=M004R26;LINKAGE=V8

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\eta\gamma)$

 Γ_{18}/Γ_6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86 ±0.04 OUR FIT				

NODE=M004R39
NODE=M004R39

0.865±0.070±0.017	419	¹ ACHASOV	00H	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
--------------------------	-----	----------------------	-----	-----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90 ±0.08 ±0.07	164	ACHASOV	98I	SND	$e^+e^- \rightarrow 5\gamma$
------------------	-----	---------	-----	-----	------------------------------

¹ Supersedes ACHASOV 98I. Excluding $\omega\pi^0$.

NODE=M004R39;LINKAGE=V8

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$

 Γ_{19}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.5 ±2.7 ±1.6	6.8k	¹ AKHMETSHIN	17	CMD3	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$

NODE=M004R22
NODE=M004R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.93±1.74±2.14	3.3k	AKHMETSHIN	00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
----------------	------	------------	-----	------	---

< 870	90	CORDIER	79	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
-------	----	---------	----	------	---

¹ Using the cross section at the ϕ meson peak $\sigma(\phi) = 4172 \pm 42$ nb, the nonresonant cross section $\sigma(0) = 1.263 \pm 0.027$ nb and $\text{Re}(Z) = 0.146 \pm 0.030$, $\text{Im}(Z) = -0.002 \pm 0.024$ for the complex amplitude of the $\phi \rightarrow \pi^+\pi^-\pi^+\pi^-$ transition.

NODE=M004R22;LINKAGE=A

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.6	90	AKHMETSHIN 00E	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<150	95	BARKOV 88	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

NODE=M004R27
NODE=M004R27

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.33^{+0.07}_{-0.10} OUR AVERAGE					
1.35 ± 0.05 ^{+0.05} _{-0.10}		9.5k	¹ ANASTASI	16B KLOE	$e^+e^- \rightarrow \pi^0 e^+ e^-$
$1.01 \pm 0.28 \pm 0.29$		52	² ACHASOV	02D SND	$e^+e^- \rightarrow \pi^0 e^+ e^-$
$1.22 \pm 0.34 \pm 0.21$		46	³ AKHMETSHIN	01C CMD2	$e^+e^- \rightarrow \pi^0 e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<12	90		DOLINSKY	88 ND	$e^+e^- \rightarrow \pi^0 e^+ e^-$

NODE=M004R31
NODE=M004R31

¹ Using $B(\pi^0 \rightarrow \gamma\gamma)$ from the 2014 Edition of this Review (PDG 14).

² Using various branching ratios from the 2000 Edition of this Review (PDG 00).

³ Using $B(\pi^0 \rightarrow \gamma\gamma) = 0.98798 \pm 0.00032$, $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033) \times 10^{-2}$, and $B(\eta \rightarrow \pi^+\pi^-\gamma) = (4.75 \pm 0.11) \times 10^{-2}$.

NODE=M004R31;LINKAGE=A
NODE=M004R;LINKAGE=DS
NODE=M004R;LINKAGE=5H

 $\Gamma(\pi^0 \eta\gamma)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.27\pm0.30 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.					
7.06 ± 0.22		16.9k	¹ AMBROSINO	09F KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
$8.51 \pm 0.51 \pm 0.57$		607	² ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
$7.96 \pm 0.60 \pm 0.40$		197	³ ALOISIO	02C KLOE	$e^+e^- \rightarrow \eta\pi^0\gamma$
$8.8 \pm 1.4 \pm 0.9$		36	⁴ ACHASOV	00F SND	$e^+e^- \rightarrow \eta\pi^0\gamma$
$9.0 \pm 2.4 \pm 1.0$		80	AKHMETSHIN	99C CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$7.01 \pm 0.10 \pm 0.20$		13.3k	^{2,5} AMBROSINO	09F KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
$7.12 \pm 0.13 \pm 0.22$		3.6k	^{3,6} AMBROSINO	09F KLOE	$1.02 e^+e^- \rightarrow \eta\pi^0\gamma$
$8.3 \pm 2.3 \pm 1.2$		20	ACHASOV	98B SND	$e^+e^- \rightarrow 5\gamma$
<250	90		DOLINSKY	91 ND	$e^+e^- \rightarrow \pi^0\eta\gamma$

NODE=M004R32
NODE=M004R32

OCCUR=2

OCCUR=2

OCCUR=3

¹ Combined results of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decay modes measurements.

² From the decay mode $\eta \rightarrow \gamma\gamma$.

³ From the decay mode $\eta \rightarrow \pi^+\pi^-\pi^0$.

⁴ Supersedes ACHASOV 98B.


⁵ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \gamma\gamma) = (39.31 \pm 0.20)\%$.

⁶ Using $B(\phi \rightarrow \eta\gamma) = (1.304 \pm 0.025)\%$, $B(\eta \rightarrow 3\pi^0) = (32.56 \pm 0.23)\%$, and $B(\eta \rightarrow \pi^+\pi^-\pi^0) = (22.73 \pm 0.28)\%$.

NODE=M004R32;LINKAGE=AM
NODE=M004R;LINKAGE=C1
NODE=M004R;LINKAGE=C2
NODE=M004R32;LINKAGE=AF
NODE=M004R32;LINKAGE=AB

NODE=M004R32;LINKAGE=AR

WEIGHTED AVERAGE
7.27 \pm 0.30 (Error scaled by 1.5)



$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$ Γ_{23}/Γ VALUE (units 10^{-5}) CL% EVTS

DOCUMENT ID TECN COMMENT

7.6±0.6 OUR FIT
7.6±0.6 OUR AVERAGE

7.4±0.7

¹ ALOISIO 02C KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$

8.8±1.7

36

² ACHASOV 00F SND $e^+e^- \rightarrow \eta\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11 ±2

³ GOKALP 02 RVUE $e^+e^- \rightarrow \eta\pi^0\gamma$

<500

90

DOLINSKY 91 ND $e^+e^- \rightarrow \pi^0\eta\gamma$ ¹ Using $M_{a_0(980)}=984.8$ MeV and assuming $a_0(980)\gamma$ dominance.² Assuming $a_0(980)\gamma$ dominance in the $\eta\pi^0\gamma$ final state.³ Using data of ACHASOV 00F.NODE=M004R33
NODE=M004R33NODE=M004R;LINKAGE=C3
NODE=M004R33;LINKAGE=AF
NODE=M004R;LINKAGE=GK $\Gamma(f_0(980)\gamma)/\Gamma(a_0(980)\gamma)$ Γ_{17}/Γ_{23}

VALUE

DOCUMENT ID TECN COMMENT

6.1±0.6¹ ALOISIO 02C KLOE $e^+e^- \rightarrow \eta\pi^0\gamma$ ¹ Using results of ALOISIO 02D and assuming that $f_0(980)$ decays into $\pi\pi$ only and $a_0(980)$ into $\eta\pi$ only.NODE=M004R47
NODE=M004R47

NODE=M004R;LINKAGE=C4

 $\Gamma(K^0\bar{K}^0\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE CL%

DOCUMENT ID TECN COMMENT

<1.9 × 10⁻⁸

90

AMBROSINO 09C KLOE $e^+e^- \rightarrow K_S^0 K_S^0 \gamma$ NODE=M004R48
NODE=M004R48 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$ Γ_{25}/Γ VALUE (units 10^{-5}) CL% EVTS

DOCUMENT ID TECN COMMENT

6.21±0.20 OUR FIT[(6.21 ± 0.21) × 10⁻⁵ OUR 2023 FIT]**6.21±0.29 OUR AVERAGE**[(6.21 ± 0.30) × 10⁻⁵ OUR 2023 AVERAGE]

6.21±0.27±0.11

3407

¹ AMBROSINO 07A KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^-\gamma$ 6.7 ^{+2.8} ±0.8

12

² AULCHENKO 03B SND $e^+e^- \rightarrow \eta'\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7 ^{+5.0} ±1.5

7

AULCHENKO 03B SND $e^+e^- \rightarrow 7\gamma$

6.10±0.61±0.43

120

³ ALOISIO 02E KLOE $1.02 e^+e^- \rightarrow \pi^+\pi^-\pi^-\gamma$ 8.2 ^{+2.1} ±1.1

21

⁴ AKHMETSHIN 00B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^-\gamma$ 4.9 ^{+2.2} ±0.6

9

⁵ AKHMETSHIN 00F CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\geq 2\gamma$

6.4 ±1.6

30

⁶ AKHMETSHIN 00F CMD2 $e^+e^- \rightarrow \eta'(958)\gamma$ 6.7 ^{+3.4} ±1.0

5

⁷ AULCHENKO 99 SND $e^+e^- \rightarrow \pi^+\pi^-\pi^-\gamma$

<11

90

AULCHENKO 98 SND $e^+e^- \rightarrow 7\gamma$ 12 ⁺⁷ ±2

6

⁴ AKHMETSHIN 97B CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^-\gamma$

<41

90

DRUZHININ 87 ND $e^+e^- \rightarrow \gamma\eta\pi^+\pi^-$ ¹ AMBROSINO 07A reports $[\Gamma(\phi(1020) \rightarrow \eta'(958)\gamma)/\Gamma_{\text{total}}] / [B(\phi(1020) \rightarrow \eta\gamma)] = (4.77 \pm 0.09 \pm 0.19) \times 10^{-3}$ which we multiply by our best value $B(\phi(1020) \rightarrow \eta\gamma) = (1.301 \pm 0.024) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.² Averaging AULCHENKO 03B with AULCHENKO 99.³ Using $B(\phi \rightarrow \eta\gamma) = (1.297 \pm 0.033)\%$.⁴ Using the value $B(\phi \rightarrow \eta\gamma) = (1.26 \pm 0.06) \times 10^{-2}$.⁵ Using $B(\phi \rightarrow K_L^0 K_S^0) = (33.8 \pm 0.6)\%$.⁶ Averaging AKHMETSHIN 00B with AKHMETSHIN 00F.⁷ Using the value $B(\eta' \rightarrow \eta\pi^+\pi^-) = (43.7 \pm 1.5) \times 10^{-2}$ and $B(\eta \rightarrow \gamma\gamma) = (39.25 \pm 0.31) \times 10^{-2}$.NODE=M004R25
NODE=M004R25

NEW

NEW

OCCUR=2

OCCUR=2

NODE=M004R25;LINKAGE=AM

NODE=M004R25;LINKAGE=BK
NODE=M004R;LINKAGE=E2

NODE=M004R25;LINKAGE=Q

NODE=M004R;LINKAGE=T2

NODE=M004R;LINKAGE=T3

NODE=M004R25;LINKAGE=AU

 $\Gamma(\eta'(958)\gamma)/\Gamma(K_L^0 K_S^0)$ Γ_{25}/Γ_2 VALUE (units 10^{-4}) EVTS

DOCUMENT ID TECN COMMENT

1.83±0.06 OUR FIT**1.46^{+0.64} ±0.18**

9

¹ AKHMETSHIN 00F CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\geq 2\gamma$ ¹ Using various branching ratios of K_S^0 , K_L^0 , η , η' from the 2000 edition (The European Physical Journal **C15** 1 (2000)) of this Review.NODE=M004R43
NODE=M004R43

NODE=M004R;LINKAGE=T1

$\Gamma(\eta'(958)\gamma)/\Gamma(\eta\gamma)$ Γ_{25}/Γ_6 VALUE (units 10^{-3}) EVTS

DOCUMENT ID TECN COMMENT

4.77±0.15 OUR FIT**4.78±0.20 OUR AVERAGE**

4.77±0.09±0.19 3407

AMBROSINO 07A KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-7\gamma$

4.70±0.47±0.31 120

¹ ALOISIO 02E KLOE 1.02 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ 6.5 $^{+1.7}_{-1.5}$ ±0.8 21AKHMETSHIN 00B CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5 $^{+5.2}_{-4.0}$ ±1.4 6² AKHMETSHIN 97B CMD2 $e^+e^- \rightarrow \pi^+\pi^-3\gamma$ ¹ From the decay mode $\eta' \rightarrow \eta\pi^+\pi^-$, $\eta \rightarrow \gamma\gamma$.² Superseded by AKHMETSHIN 00B.NODE=M004R34
NODE=M004R34NODE=M004R;LINKAGE=E1
NODE=M004R;LINKAGE=KS $\Gamma(\eta\pi^0\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{26}/Γ VALUE (units 10^{-5}) CL%

DOCUMENT ID TECN COMMENT

<2

90

AULCHENKO 98 SND $e^+e^- \rightarrow 7\gamma$ NODE=M004R36
NODE=M004R36 $\Gamma(\mu^+\mu^-\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ VALUE (units 10^{-5}) EVTS

DOCUMENT ID TECN COMMENT

1.43±0.45±0.14

27188

¹ AKHMETSHIN 99B CMD2 $e^+e^- \rightarrow \mu^+\mu^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ±1.0

824 ± 33

² AKHMETSHIN 97C CMD2 $e^+e^- \rightarrow \mu^+\mu^-\gamma$ ¹ For $E_\gamma > 20$ MeV. Supersedes AKHMETSHIN 97C.² For $E_\gamma > 20$ MeV.NODE=M004R35
NODE=M004R35NODE=M004R35;LINKAGE=3N
NODE=M004R35;LINKAGE=A $\Gamma(\rho\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{28}/Γ VALUE (units 10^{-4}) CL%

DOCUMENT ID TECN COMMENT

<1.2

90

AULCHENKO 08 CMD2 $\phi \rightarrow \pi^+\pi^-\gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5

90

AKHMETSHIN 98 CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ NODE=M004R37
NODE=M004R37 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ VALUE (units 10^{-5}) CL%

DOCUMENT ID TECN COMMENT

< 1.8

90

AKHMETSHIN 00E CMD2 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.1

90

AULCHENKO 08 CMD2 $\phi \rightarrow \eta\pi^+\pi^-$

<30

90

AKHMETSHIN 98 CMD2 $e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$ NODE=M004R38
NODE=M004R38 $\Gamma(\eta\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ VALUE (units 10^{-6}) CL%

DOCUMENT ID TECN COMMENT

<9.4

90

AKHMETSHIN 01 CMD2 $e^+e^- \rightarrow \eta e^+e^-$ NODE=M004R45
NODE=M004R45 $\Gamma(\eta U \rightarrow \eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE CL%

DOCUMENT ID TECN COMMENT

<1 × 10⁻⁶

90

¹ BABUSCI 13B KLOE 1.02 $e^+e^- \rightarrow \eta e^+e^-$ ¹ For a narrow vector U with mass between 5 and 470 MeV, from the combined analysis of $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow \pi^0\pi^0\pi^0$ from ARCHILLI 12. Measured 90% CL limits as a function of m_U range from 2.2×10^{-8} to 10^{-6} .NODE=M004R01
NODE=M004R01

NODE=M004R01;LINKAGE=A

 $\Gamma(\text{invisible})/\Gamma(K^+K^-)$ Γ_{32}/Γ_1

VALUE CL%

DOCUMENT ID TECN COMMENT

<3.4 × 10⁻⁴

90

ABLIKIM 18S BES3 $J/\psi \rightarrow \phi\eta \rightarrow \phi\pi^+\pi^-\pi^0$ NODE=M004R02
NODE=M004R02

————— Lepton Family number (LF) violating modes —————

NODE=M00422A

 $\Gamma(e^\pm\mu^\mp)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE CL%

DOCUMENT ID TECN COMMENT

<2 × 10⁻⁶

90

ACHASOV 10A SND $e^+e^- \rightarrow e^\pm\mu^\mp$ NODE=M004R29
NODE=M004R29 $\pi^+\pi^-\pi^0 / \rho\pi$ AMPLITUDE RATIO a_1 IN DECAY OF $\phi \rightarrow \pi^+\pi^-\pi^0$

NODE=M004D1

NIECKNIG 12 describes final-state interactions between the three pions in a dispersive framework using data on the $\pi\pi$ P -wave scattering phase shift.

NODE=M004D1

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.2 OUR AVERAGE					
10.1±4.4±1.7		80k	¹ AKHMETSHIN 06	CMD2	1.017–1.021 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0±1.1±0.6		1.98M	^{2,3} ALOISIO	03 KLOE	1.02 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$-6 < a_1 < 6$		500k	³ ACHASOV	02 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
$-16 < a_1 < 11$	90	9.8k	^{1,4} AKHMETSHIN 98	CMD2	$e^+e^- \rightarrow \pi^+\pi^-\gamma\gamma$
¹ Dalitz plot analysis taking into account interference between the contact and $\rho\pi$ amplitudes.					
² From a fit without limitations on charged and neutral ρ masses and widths.					
³ Recalculated by us to match the notations of AKHMETSHIN 98.					
⁴ Assuming zero phase for the contact term.					

NODE=M004D1

NODE=M004D1;LINKAGE=AK

NODE=M004D;LINKAGE=L1

NODE=M004D;LINKAGE=L3

NODE=M004D1;LINKAGE=KL

PARAMETER β IN $\phi \rightarrow P e^+ e^-$ DECAYS

In the one-pole approximation the electromagnetic transition form factor for $\phi \rightarrow P e^+ e^-$ ($P = \pi, \eta$) is given as a function of the $e^+ e^-$ invariant mass squared, q^2 , by the expression:

$$|F(q^2)|^2 = (1 - q^2/\Lambda^2)^{-2},$$

where vector meson dominance predicts parameter $\Lambda \approx 0.770$ GeV ($\Lambda^{-2} \approx 1.687$ GeV $^{-2}$). The slope of this form factor, $\beta = dF/dq^2(q^2=0)$, equals Λ^{-2} in this approximation.

The measurements below obtain β in the one-pole approximation.

NODE=M004230

NODE=M004230

PARAMETER β IN $\phi \rightarrow \pi^0 e^+ e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
2.02±0.11	9.5k	¹ ANASTASI	16B KLOE	1.02 $e^+e^- \rightarrow \pi^0 e^+e^-$

¹ The error combines statistical and systematic uncertainties.

NODE=M004A00
NODE=M004A00

NODE=M004A00;LINKAGE=A

PARAMETER β IN $\phi \rightarrow \eta e^+ e^-$ DECAY

VALUE (GeV $^{-2}$)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29±0.13 OUR AVERAGE				
1.28±0.10 $^{+0.09}_{-0.08}$	30k	BABUSCI	15 KLOE	1.02 $e^+e^- \rightarrow \eta e^+e^-$
3.8 ± 1.8	213	¹ ACHASOV	01B SND	1.02 $e^+e^- \rightarrow \eta e^+e^-$

¹ The uncertainty is statistical only. The systematic one is negligible, in comparison.

NODE=M004BFP
NODE=M004BFP

NODE=M004BFP;LINKAGE=A

$\phi(1020)$ REFERENCES

LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
HOID	20	EPJ C80 988	B.-L. Hoid, M. Hoferichter, B. Kubis	(BONN, BERN)	REFID=61048
AAIJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59670
HOFRICHT...	19	JHEP 1908 137	M. Hoferichter, B.-L. Hoid, B. Kubis	(WASH, BONN)	REFID=60974
ABLIKIM	18S	PR D98 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58971
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>		REFID=59987
KOZYREV	18	PL B779 64	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)	REFID=58794
AKHMETSHIN	17	PL B768 345	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=57893
ACHASOV	16A	PR D93 092001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57513
ANASTASI	16B	PL B757 362	A. Anastasi <i>et al.</i>	(KLOE-2 Collab.)	REFID=57399
KOZYREV	16	PL B760 314	E.A. Kozyrev <i>et al.</i>	(CMD-3 Collab.)	REFID=57514
BABUSCI	15	PL B742 1	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)	REFID=56374
PDG	15	RPP 2015 at pdg.lbl.gov		(PDG Collab.)	REFID=56977;ERROR=7
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
BABUSCI	13B	PL B720 111	D. Babusci <i>et al.</i>	(KLOE-2 Collab.)	REFID=55068
BENAYOUN	13	EPJ C73 2453	M. Benayoun, P. David, L. DelBuono	(PARIN, BERLIN+)	REFID=55357
LEES	13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55127
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
ARCHILLI	12	PL B706 251	F. Archilli <i>et al.</i>	(KLOE-2 Collab.)	REFID=53951
BENAYOUN	12	EPJ C72 1848	M. Benayoun <i>et al.</i>		REFID=54281
NIECKNIG	12	EPJ C72 2014	F. Niecknig, B. Kubis, S.P. Schneider	(BONN)	REFID=54305
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
AKHMETSHIN	11	PL B695 412	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=53645
ACHASOV	10A	PR D81 057102	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=53352
BENAYOUN	10	EPJ C65 211	M. Benayoun <i>et al.</i>		REFID=53212
AMBROSINO	09C	PL B679 10	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52969
AMBROSINO	09F	PL B681 5	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=53105
AKHMETSHIN	08	PL B669 217	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=52572
AMBROSINO	08G	PL B669 223	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=52573
AULCHENKO	08	JETPL 88 85	V. Aulchenko <i>et al.</i>	(CMD-2 Collab.)	REFID=52268

Translated from ZETFP 88 93.

NODE=M004

FLOREZ-BAEZ	08	PR D78 077301	F.V. Florez-Baez, G. Lopez Castro		REFID=52584
ACHASOV	07B	PR D76 077101	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51942
AMBROSINO	07	EPJ C49 473	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51616
AMBROSINO	07A	PL B648 267	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=51646
DUBYSNSKIY	07	PR D75 113001	S. Dubynskiy <i>et al.</i>		REFID=51719
ACHASOV	06A	PR D74 014016	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=51133
AKHMETSHIN	06	PL B642 203	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=51465
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AMBROSINO	05	PL B608 199	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=50453
AUBERT,B	05J	PR D72 052008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50824
AKHMETSHIN	04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ALOISIO	03	PL B561 55	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=49404
AULCHENKO	03B	JETP 97 24	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49613
		Translated from ZETF 124 28.			
ACHASOV	02	PR D65 032002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48549
ACHASOV	02D	JETPL 75 449	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48814
		Translated from ZETFP 75 539.			
ALOISIO	02C	PL B536 209	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48823
ALOISIO	02D	PL B537 21	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48824
ALOISIO	02E	PL B541 45	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=48825
FISCHBACH	02	PL B526 355	E. Fischbach, A.W. Overhauser, B. Woodahl		REFID=48575
GOKALP	02	JP G28 2783	A. Gokalp <i>et al.</i>		REFID=49167
ACHASOV	01B	PL B504 275	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48111
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
ACHASOV	01F	PR D63 094007	N.N. Achasov, V.V. Gubin	(Novosibirsk SND Collab.)	REFID=48312
ACHASOV	01G	PR L 86 1698	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48315
AITALA	01B	PR L 86 770	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48005
AKHMETSHIN	01	PL B501 191	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48110
AKHMETSHIN	01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
AKHMETSHIN	01C	PL B503 237	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48323
BENAYOUN	01	EPJ C22 503	M. Benayoun, H.B. O'Connell		REFID=48570
ACHASOV	00	EPJ C12 25	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47417
ACHASOV	00B	JETP 90 17	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47425
		Translated from ZETF 117 22.			
ACHASOV	00C	PL B474 188	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47431
ACHASOV	00D	JETPL 72 282	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47882
		Translated from ZETFP 72 411.			
ACHASOV	00E	NP B569 158	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47927
ACHASOV	00F	PL B479 53	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47928
ACHASOV	00H	PL B485 349	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47930
AKHMETSHIN	00B	PL B473 337	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47422
AKHMETSHIN	00E	PL B491 81	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47936
AKHMETSHIN	00F	PL B494 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47937
AULCHENKO	00A	JETP 90 927	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47953
		Translated from ZETF 117 1067.			
BRAMON	00	PL B486 406	A. Bramon <i>et al.</i>		REFID=47969
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ACHASOV	99	PL B449 122	M.N. Achasov <i>et al.</i>		REFID=46896
ACHASOV	99C	PL B456 304	M.N. Achasov <i>et al.</i>		REFID=46939
AKHMETSHIN	99B	PL B462 371	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47392
AKHMETSHIN	99C	PL B462 380	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47393
AKHMETSHIN	99D	PL B466 385	R.R. Akhmetshin <i>et al.</i>		REFID=47397
		Also			
		PL B508 217 (err.)	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48328
AKHMETSHIN	99F	PL B460 242	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47473
AULCHENKO	99	JETPL 69 97	V.M. Aulchenko <i>et al.</i>		REFID=46920
		Translated from ZETFP 69 87.			
ACHASOV	98B	PL B438 441	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46317
ACHASOV	98F	JETPL 68 573	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46321
ACHASOV	98I	PL B440 442	M.N. Achasov <i>et al.</i>		REFID=46600
AKHMETSHIN	98	PL B434 426	R.R. Akhmetshin <i>et al.</i>	(CMD-2 Collab.)	REFID=46325
AULCHENKO	98	PL B436 199	V.M. Aulchenko <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=46336
BARBERIS	98	PL B432 436	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46344
AKHMETSHIN	97B	PL B415 445	R.R. Akhmetshin <i>et al.</i>	(NOVO, BOST, PITT+)	REFID=45801
AKHMETSHIN	97C	PL B415 452	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=45802
BENAYOUN	96	ZPHY C72 221	M. Benayoun <i>et al.</i>	(IPNP, NOVO)	REFID=45753
AKHMETSHIN	95	PL B364 199	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=44617
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ACHASOV	89	NP B315 465	N.N. Achasov, V.N. Ivanchenko		REFID=48021
DOLINSKY	89	ZPHY C42 511	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41003
BARKOV	88	SJNP 47 248	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=41024
		Translated from YAF 47 393.			
DOLINSKY	88	SJNP 48 277	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41022
		Translated from YAF 48 442.			
DRUZHININ	87	ZPHY C37 1	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=40448
ARMSTRONG	86	PL 166B 245	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=20563
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
BEBEK	86	PRL 56 1893	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=11540
DAVENPORT	86	PR D33 2519	T.F. Davenport (TUFTS, ARIZ, FNAL, FSU, NDAM+)		REFID=20567
DIJKSTRA	86	ZPHY C31 375	H. Dijkstra <i>et al.</i>	(ANIK, BRIS, CERN+)	REFID=20568
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)	REFID=20569
GOLUBEV	86	SJNP 44 409	V.B. Golubev <i>et al.</i>	(NOVO)	REFID=40449
		Translated from YAF 44 633.			
ALBRECHT	85D	PL 153B 343	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=20562
GOLUBEV	85	SJNP 41 756	V.B. Golubev <i>et al.</i>	(NOVO)	REFID=40450
		Translated from YAF 41 1183.			
DRUZHININ	84	PL 144B 136	V.P. Druzhinin <i>et al.</i>	(NOVO)	REFID=20561
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
BARATE	83	PL 121B 449	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, IND)	REFID=12177
KURDADZE	83C	JETPL 38 366	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20560
		Translated from ZETFP 38 306.			
ARENTON	82	PR D25 2241	M.W. Arenton <i>et al.</i>	(ANL, ILL)	REFID=20556
PELLINEN	82	PS 25 599	A. Pellinen, M. Roos	(HELS)	REFID=20557
DAUM	81	PL 100B 439	C. Daum <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)	REFID=20552
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
		Also			
		Private Comm.	S.I. Eidelman	(NOVO)	REFID=20554
VASSERMAN	81	PL 99B 62	I.B. Vasserman <i>et al.</i>	(NOVO)	REFID=20555
		Also			
		SJNP 35 240	L.M. Kurdadze <i>et al.</i>		REFID=47475
		Translated from YAF 35 352.			

CORDIER	80	NP B172 13	A. Cordier <i>et al.</i>	(LALO)	REFID=20240
CORDIER	79	PL 81B 389	A. Cordier <i>et al.</i>	(LALO)	REFID=20549
BUKIN	78B	SJNP 27 521	A.D. Bukin <i>et al.</i>	(NOVO)	REFID=20545
BUKIN	78C	Translated from YAF 27 985. SJNP 27 516	A.D. Bukin <i>et al.</i>	(NOVO)	REFID=20544
COOPER	78B	NP B146 1	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=20235
LOSTY	78	NP B133 38	M.J. Losty <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=20547
AKERLOF	77	PRL 39 861	C.W. Akerlof <i>et al.</i>	(FNAL, MICH, PURD)	REFID=20534
ANDREWS	77	PRL 38 198	D.E. Andrews <i>et al.</i>	(ROCH)	REFID=20120
BALDI	77	PL 68B 381	R. Baldi <i>et al.</i>	(GEVA)	REFID=20536
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+)	REFID=20537
COHEN	77	PRL 38 269	D. Cohen <i>et al.</i>	(ANL)	REFID=20538
LAVEN	77	NP B127 43	H. Laven <i>et al.</i>	(AACH3, BERL, CERN, LOIC+)	REFID=20541
LYONS	77	NP B125 207	L. Lyons, A.M. Cooper, A.G. Clark	(OXF)	REFID=20232
COSME	76	PL 63B 352	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20529
KALBFLEISCH	76	PR D13 22	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20531
PARROUR	76	PL 63B 357	G. Parrou <i>et al.</i>	(ORSAY)	REFID=20532
PARROUR	76B	PL 63B 362	G. Parrou <i>et al.</i>	(ORSAY)	REFID=20533
KALBFLEISCH	75	PR D11 987	G.R. Kalbfleisch, R.C. Strand, J.W. Chapman	(BNL+)	REFID=20223
AYRES	74	PRL 32 1463	D.S. Ayres <i>et al.</i>	(ANL)	REFID=20522
BESCH	74	NP B70 257	H.J. Besch <i>et al.</i>	(BONN)	REFID=20523
COSME	74	PL 48B 155	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20525
COSME	74B	PL 48B 159	G. Cosme <i>et al.</i>	(ORSAY)	REFID=20526
DEGROOT	74	NP B74 77	A.J. de Groot <i>et al.</i>	(AMST, NIJM)	REFID=20527
AUGUSTIN	73	PRL 30 462	J.E. Augustin <i>et al.</i>	(ORSAY)	REFID=47515
BALLAM	73	PR D7 3150	J. Ballam <i>et al.</i>	(SLAC, LBL)	REFID=20520
BINNIE	73B	PR D8 2789	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=20216
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
ALVENSLEB...	72	PRL 28 66	H. Alvensleben <i>et al.</i>	(MIT, DESY)	REFID=20514
BORENSTEIN	72	PR D5 1559	S.R. Borenstein <i>et al.</i>	(BNL, MICH)	REFID=20215
COLLEY	72	NP B50 1	D.C. Colley <i>et al.</i>	(BIRM, GLAS)	REFID=20519
BALAKIN	71	PL 34B 328	V.E. Balakin <i>et al.</i>	(NOVO)	REFID=20507
CHATELUS	71	Thesis LAL 1247	Y. Chatelus	(STRB)	REFID=20508
Also		PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)	REFID=20501
HAYES	71	PR D4 899	S. Hayes <i>et al.</i>	(CORN)	REFID=20511
STOTTLE...	71	Thesis ORO 2504 170	A.R. Stottlemyer	(UMD)	REFID=20512
BIZOT	70	PL 32B 416	J.C. Bizot <i>et al.</i>	(ORSAY)	REFID=20501
Also		Liverpool Sym. 69	J.P. Perez-y-Jorba		REFID=20502
EARLES	70	PRL 25 1312	D.R. Earles <i>et al.</i>	(NEAS)	REFID=20504
LINDSEY	66	PR 147 913	J.S. Lindsey, G. Smith	(LRL)	REFID=20481
LONDON	66	PR 143 1034	G.W. London <i>et al.</i>	(BNL, SYRA) IGJPC	REFID=11774
BADIER	65B	PL 17 337	J. Badier <i>et al.</i>	(EPOL, SACL, AMST)	REFID=20253
LINDSEY	65	PRL 15 221	J.S. Lindsey, G.A. Smith	(LRL)	REFID=20478
LINDSEY 65 data included in LINDSEY 66.					
SCHLEIN	63	PRL 10 368	P.E. Schlein <i>et al.</i>	(UCLA) IGJP	REFID=20474

h₁(1170)

$$I^G(J^{PC}) = 0^-(1^{+-})$$

h₁(1170) MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1166 ± 5 ± 3	¹ ANDO	92	SPEC	8 π ⁻ p → π ⁺ π ⁻ π ⁰ n
1168 ± 4	ANDO	92	SPEC	8 π ⁻ p → π ⁺ π ⁻ π ⁰ n
1190 ± 60	² DANKOWY...	81	SPEC 0	8 π p → 3π n

¹ Average and spread of values using 2 variants of the model of BOWLER 75.
² Uses the model of BOWLER 75.

h₁(1170) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
375 ± 6 ± 34	³ ANDO	92	SPEC	8 π ⁻ p → π ⁺ π ⁻ π ⁰ n
345 ± 6	ANDO	92	SPEC	8 π ⁻ p → π ⁺ π ⁻ π ⁰ n
320 ± 50	⁴ DANKOWY...	81	SPEC 0	8 π p → 3π n

³ Average and spread of values using 2 variants of the model of BOWLER 75.
⁴ Uses the model of BOWLER 75.

h₁(1170) DECAY MODES

Mode	Fraction (Γ _i /Γ)
Γ ₁ ρπ	seen

NODE=M030M

NODE=M030M

OCCUR=2

NODE=M030M;LINKAGE=B
 NODE=M030M;LINKAGE=C

NODE=M030W

NODE=M030W

OCCUR=2

NODE=M030W;LINKAGE=B
 NODE=M030W;LINKAGE=C

NODE=M030215;NODE=M030

$h_1(1170)$ BRANCHING RATIOS

NODE=M030220

$\Gamma(\rho\pi)/\Gamma_{total}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen		ANDO	92	SPEC $8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
seen		ATKINSON	84	OMEG $20-70 \gamma p \rightarrow \pi^+ \pi^- \pi^0 p$	
seen		DANKOWY...	81	SPEC $8 \pi p \rightarrow 3 \pi n$	

NODE=M030R1
NODE=M030R1

$h_1(1170)$ REFERENCES

NODE=M030

ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
ATKINSON	84	NP B231 15	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

REFID=43171
REFID=20574
REFID=20572
REFID=20571

$b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

NODE=M011


$b_1(1235)$ MASS

NODE=M011M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1229.5 ± 3.2 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.			
1225 ± 5		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
1235 ± 15		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
1236 ± 16		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
1222 ± 6		ATKINSON 84E	OMEG ±		25-55 $\gamma p \rightarrow \omega \pi X$
1237 ± 7		ATKINSON 84E	OMEG 0		25-55 $\gamma p \rightarrow \omega \pi X$
1239 ± 5		EVANGELIS... 81	OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
1251 ± 8	450	GESSAROLI 77	HBC -		11 $\pi^- p \rightarrow \pi^- \omega p$
1245 ± 11	890	FLATTE 76C	HBC -		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
1222 ± 4	1400	CHALOUPKA 74	HBC -		3.9 $\pi^- p$
1220 ± 7	600	KARSHON 74B	HBC +		4.9 $\pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1190 ± 10		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
1213 ± 5		ATKINSON 84C	OMEG 0		20-70 γp
1271 ± 11		COLLICK 84	SPEC +		200 $\pi^+ Z \rightarrow Z \pi \omega$

NODE=M011M

OCCUR=2



$b_1(1235)$ WIDTH

NODE=M011W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
142 ± 9 OUR AVERAGE		Error includes scale factor of 1.2.			
113 ± 12		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
160 ± 30		ALDE	92C	GAM2	38,100 $\pi^- p \rightarrow \omega \pi^0 n$
151 ± 31		FUKUI	91	SPEC	8.95 $\pi^- p \rightarrow \omega \pi^0 n$
170 ± 15		EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow \omega \pi p$
170 ± 50	225	BALTAY	78B	HBC +	15 $\pi^+ p \rightarrow p 4\pi$
155 ± 32	450	GESSAROLI	77	HBC -	11 $\pi^- p \rightarrow \pi^- \omega p$
182 ± 45	890	FLATTE	76C	HBC -	4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
135 ± 20	1400	CHALOUPKA	74	HBC -	3.9 $\pi^- p$
156 ± 22	600	KARSHON	74B	HBC +	4.9 $\pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
210 ± 19		AUGUSTIN	89	DM2 ±	$e^+ e^- \rightarrow 5\pi$
231 ± 14		ATKINSON	84C	OMEG 0	20-70 γp
232 ± 29		COLLICK	84	SPEC +	200 $\pi^+ Z \rightarrow Z \pi \omega$

NODE=M011W

 $b_1(1235)$ DECAY MODES

NODE=M011215;NODE=M011

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\omega \pi$ [D/S amplitude ratio = 0.277 ± 0.027]	seen	
Γ_2 $\pi^\pm \gamma$	(1.6 ± 0.4) × 10 ⁻³	
Γ_3 $\eta \rho$	seen	
Γ_4 $\pi^+ \pi^+ \pi^- \pi^0$	< 50 %	84%
Γ_5 $K^*(892)^\pm K^\mp$	seen	
Γ_6 $(KK)^\pm \pi^0$	< 8 %	90%
Γ_7 $K_S^0 K_L^0 \pi^\pm$	< 6 %	90%
Γ_8 $K_S^0 K_S^0 \pi^\pm$	< 2 %	90%
Γ_9 $\phi \pi$	< 1.5 %	84%

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=74

DESIG=71;OUR EST;→ UNCHECKED ←

DESIG=73;OUR EST;→ UNCHECKED ←

DESIG=72;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

 $b_1(1235)$ PARTIAL WIDTHS

NODE=M011220

$\Gamma(\pi^\pm \gamma)$	VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2
230 ± 60		COLLICK	84	SPEC +	200 $\pi^+ Z \rightarrow Z \pi \omega$	

NODE=M011W3


NODE=M011W3

 **$b_1(1235)$ D-wave/S-wave AMPLITUDE RATIO
IN DECAY OF $b_1(1235) \rightarrow \omega \pi$**

NODE=M011DS

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.277 ± 0.027 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.			
0.269 ± 0.009 ± 0.010		NOZAR	02	MPS -	18 $\pi^- p \rightarrow \omega \pi^- p$
0.23 ± 0.03		AMSLER	94C	CBAR	0.0 $\bar{p}p \rightarrow \omega \eta \pi^0$
0.45 ± 0.04		AMSLER	93B	CBAR	0.0 $\bar{p}p \rightarrow \omega \pi^0 \pi^0$
0.235 ± 0.047		ATKINSON	84C	OMEG	20-70 γp
0.4 $\begin{smallmatrix} +0.1 \\ -0.1 \end{smallmatrix}$		GESSAROLI	77	HBC -	11 $\pi^- p \rightarrow \pi^- \omega p$
0.21 ± 0.08		CHUNG	75B	HBC +	7.1 $\pi^+ p$
0.3 ± 0.1		CHALOUPKA	74	HBC -	3.9-7.5 $\pi^- p$
0.35 ± 0.25	600	KARSHON	74B	HBC +	4.9 $\pi^+ p$

NODE=M011DS



$b_1(1235)$ D-wave/S-wave AMPLITUDE PHASE DIFFERENCE IN DECAY OF $b_1(1235) \rightarrow \omega\pi$

NODE=M011PH

VALUE (°)	DOCUMENT ID	TECN	CHG	COMMENT
10.5±2.4±3.9	NOZAR	02	MPS	- 18 $\pi^- p \rightarrow \omega\pi^- p$

NODE=M011PH

$b_1(1235)$ BRANCHING RATIOS

NODE=M011230

$\Gamma(\eta\rho)/\Gamma(\omega\pi)$

 Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.10	ATKINSON	84D	OMEG	20-70 γp

NODE=M011R9
NODE=M011R9

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\omega\pi)$

 Γ_4/Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.5	ABOLINS	63	HBC	+ 3.5 $\pi^+ p$

NODE=M011R1
NODE=M011R1

$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$

 Γ_5/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	¹ ABLIKIM	10E	BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M011R10
NODE=M011R10

¹ From a fit including ten additional resonances and energy-independent Breit-Wigner width.

NODE=M011R10;LINKAGE=AB

$\Gamma((K\bar{K})^\pm \pi^0)/\Gamma(\omega\pi)$

 Γ_6/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.08	90	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R6
NODE=M011R6

$\Gamma(K_S^0 K_L^0 \pi^\pm)/\Gamma(\omega\pi)$

 Γ_7/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.06	90	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R8
NODE=M011R8

$\Gamma(K_S^0 K_S^0 \pi^\pm)/\Gamma(\omega\pi)$

 Γ_8/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.02	90	BALTAY	67	HBC	± 0.0 $\bar{p}p$

NODE=M011R7
NODE=M011R7

$\Gamma(\phi\pi)/\Gamma(\omega\pi)$

 Γ_9/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.004	95	VIKTOROV	96	SPEC	0 32.5 $\pi^- p \rightarrow K^+ K^- \pi^0 n$

NODE=M011R4
NODE=M011R4

••• We do not use the following data for averages, fits, limits, etc. •••

<0.04	95	BIZZARRI	69	HBC	± 0.0 $\bar{p}p$
<0.015		DAHL	67	HBC	1.6-4.2 $\pi^- p$

$b_1(1235)$ REFERENCES

ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)
NOZAR	02	PL B541 35	M. Nozar <i>et al.</i>	
VIKTOROV	96	PAN 59 1184	V.A. Viktorov <i>et al.</i>	(SERP)
		Translated from YAF 59 1239.		
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	93B	PL B311 362	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)
FUKUI	91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)
ATKINSON	84C	NP B243 1	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+) JP
ATKINSON	84D	NP B242 269	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
ATKINSON	84E	PL 138B 459	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
COLLICK	84	PRL 53 2374	B. Collick <i>et al.</i>	(MINN, ROCH, FNAL)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+) JP
FLATTE	76C	PL 64B 225	S.M. Flatte <i>et al.</i>	(CERN, AMST, NIJM+) JP
CHUNG	75B	PR D11 2426	S.U. Chung <i>et al.</i>	(BNL, LBL, UCSC) JP
CHALOUPKA	74	PL 51B 407	V. Chaloupka <i>et al.</i>	(CERN) JP
KARSHON	74B	PR D10 3608	U. Karshon <i>et al.</i>	(REHO) JP
BIZZARRI	69	NP B14 169	R. Bizzarri <i>et al.</i>	(CERN, CDEF)
BALTAY	67	PRL 18 93	C. Baltay <i>et al.</i>	(COLU)
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL)
ABOLINS	63	PRL 11 381	M.A. Abolins <i>et al.</i>	(UCSD)

NODE=M011

REFID=53361
REFID=48850
REFID=45203

REFID=44091
REFID=43602
REFID=43585
REFID=41859
REFID=41581
REFID=41004
REFID=20625
REFID=20623
REFID=20624
REFID=20626
REFID=20462
REFID=21265
REFID=20230
REFID=20615
REFID=20613
REFID=20611
REFID=20612
REFID=20171
REFID=20159
REFID=20321
REFID=20006

$a_1(1260)$

$$I^G(J^{PC}) = 1^-(1^+ +)$$

See also our review under the $a_1(1260)$ in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M010

NODE=M010

$a_1(1260)$ T-MATRIX POLE \sqrt{s}

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M010PP

NODE=M010PP

NODE=M010PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$(1209 \pm 4 \pm_{-9}^{+12}) - i(288 \pm 6 \pm_{-10}^{+45})$			OUR ESTIMATE
$(1209 \pm 4 \pm_{-9}^{+12}) - i(288 \pm 6 \pm_{-10}^{+45})$	MIKHASENKO 18	RVUE	$\tau^- \rightarrow \pi^- \pi^+ \pi^- \nu_\tau$

→ UNCHECKED ←

$a_1(1260)$ MASS

NODE=M010M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1230 ± 40				OUR ESTIMATE
1299 ±₋₂₈⁺¹²	46M	1 AGHASYAN	18B	COMP 190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1195.05 ± 1.05 ± 6.33	894k	AAIJ	18A1	LHCB $D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
1225 ± 9 ± 20	7k	2 DARGENT	17	RVUE $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1255 ± 6 ± ₋₁₇ ⁺⁷	420k	3 ALEKSEEV	10	COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1243 ± 12 ± 20		4 AUBERT	07AU	BABR 10.6 $e^+ e^- \rightarrow \rho^0 \rho^\pm \pi^\mp \gamma$
1230–1270	6360	5 LINK	07A	FOCS $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1203 ± 3		6 GOMEZ-DUM..04		RVUE $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$
1330 ± 24	90k	SALVINI	04	OBLX $\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
1331 ± 10 ± 3	37k	7 ASNER	00	CLE2 10.6 $e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
1255 ± 7 ± 6	5904	8 ABREU	98G	DLPH $e^+ e^-$
1207 ± 5 ± 8	5904	9 ABREU	98G	DLPH $e^+ e^-$
1196 ± 4 ± 5	5904	10,11 ABREU	98G	DLPH $e^+ e^-$
1240 ± 10		BARBERIS	98B	450 $pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$
1262 ± 9 ± 7		8,12 ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94$, $\tau \rightarrow 3\pi \nu$
1210 ± 7 ± 2		9,12 ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94$, $\tau \rightarrow 3\pi \nu$
1211 ± 7 ± ₋₀ ⁺⁵⁰		9 ALBRECHT	93C	ARG $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1121 ± 8		13 ANDO	92	SPEC $8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1242 ± 37		14 IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1260 ± 14		15 IVANOV	91	RVUE $\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$

NODE=M010M

→ UNCHECKED ←

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

1250 ± 9	16	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
1208 ± 15		ARMSTRONG	90	OMEG	$300.0 p p \rightarrow$ $p p \pi^+ \pi^- \pi^0$	
1220 ± 15	17	ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1260 ± 25	18	BOWLER	88	RVUE		
1166 ± 18 ± 11		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1164 ± 41 ± 23		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$	OCCUR=2
1250 ± 40	17	TORNQVIST	87	RVUE		
1046 ± 11		ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1056 ± 20 ± 15		RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1194 ± 14 ± 10		SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
1255 ± 23		BELLINI	85	SPEC	$40 \pi^- A \rightarrow$ $\pi^- \pi^+ \pi^- A$	
1240 ± 80	19	DANKOWY...	81	SPEC	$8.45 \pi^- p \rightarrow n 3\pi$	
1280 ± 30	19	DAUM	81B	CNTR	$63.94 \pi^- p \rightarrow p 3\pi$	
1041 ± 13	20	GAVILLET	77	HBC	$4.2 K^- p \rightarrow \Sigma 3\pi$	

- ¹ Statistical error negligible.
- ² Reanalysis of CLEO data using Breit-Wigner parameterization.
- ³ Superseded by AGHASYAN 2018B.
- ⁴ The $\rho^\pm \pi^\mp$ state can be also due to the $\pi(1300)$.
- ⁵ Using the Breit-Wigner parameterization; strong correlation between mass and width.
- ⁶ Using the data of BARATE 98R.
- ⁷ From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.
- ⁸ Uses the model of KUHNS 90.
- ⁹ Uses the model of ISGUR 89.
- ¹⁰ Includes the effect of a possible a_1' state.
- ¹¹ Uses the model of FEINDT 90.
- ¹² Supersedes AKERS 95P.
- ¹³ Average and spread of values using 2 variants of the model of BOWLER 75.
- ¹⁴ Reanalysis of RUCKSTUHL 86.
- ¹⁵ Reanalysis of SCHMIDKE 86.
- ¹⁶ Reanalysis of ALBRECHT 86B.
- ¹⁷ From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.
- ¹⁸ From a combined reanalysis of ALBRECHT 86B and DAUM 81B.
- ¹⁹ Uses the model of BOWLER 75.
- ²⁰ Produced in K^- backward scattering.

NODE=M010M;LINKAGE=Q
 NODE=M010M;LINKAGE=V
 NODE=M010M;LINKAGE=R
 NODE=M010M;LINKAGE=AU
 NODE=M010M;LINKAGE=LI
 NODE=M010M;LINKAGE=GO
 NODE=M010M;LINKAGE=B6
 NODE=M010M;LINKAGE=KS
 NODE=M010M;LINKAGE=IM
 NODE=M010M;LINKAGE=A1
 NODE=M010M;LINKAGE=F1
 NODE=M010M;LINKAGE=X
 NODE=M010M;LINKAGE=P
 NODE=M010M;LINKAGE=I
 NODE=M010M;LINKAGE=L
 NODE=M010M;LINKAGE=M
 NODE=M010M;LINKAGE=K
 NODE=M010M;LINKAGE=G
 NODE=M010M;LINKAGE=D
 NODE=M010M;LINKAGE=F

$a_1(1260)$ WIDTH

NODE=M010W
 NODE=M010W
 → UNCHECKED ←

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
250 to 600 OUR ESTIMATE				
420 ± 35 OUR AVERAGE				
380 ± 80	46M	¹ AGHASYAN	18B COMP	$190 \pi^- p \rightarrow$ $\pi^- \pi^+ \pi^- p$
430 ± 24 ± 31		DARGENT	17 RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
422.01 ± 2.10 ± 12.72	894k	AAIJ	18A1 LHCB	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
367 ± 9 ± 28	420k	² ALEKSEEV	10 COMP	$190 \pi^- P b \rightarrow$ $\pi^- \pi^- \pi^+ P b'$
410 ± 31 ± 30		³ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow$ $\rho^0 \rho^\pm \pi^\mp \gamma$
520-680	6360	⁴ LINK	07A FOCUS	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
480 ± 20		⁵ GOMEZ-DUM...	04 RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu_\tau$
580 ± 41	90k	SALVINI	04 OBLX	$\bar{p} p \rightarrow 2\pi^+ 2\pi^-$
460 ± 85	205	⁶ DRUTSKOY	02 BELL	$B \rightarrow D(*) K^- K^* 0$
814 ± 36 ± 13	37k	⁷ ASNER	00 CLE2	$10.6 e^+ e^- \rightarrow$ $\tau^+ \tau^-, \tau^- \rightarrow$ $\pi^- \pi^0 \pi^0 \nu_\tau$
450 ± 50	22k	⁸ AKHMETSHIN	99E CMD2	$1.05-1.38 e^+ e^- \rightarrow$ $\pi^+ \pi^- \pi^0 \pi^0$
570 ± 10		⁹ BONDAR	99 RVUE	$e^+ e^- \rightarrow 4\pi, \tau \rightarrow$ $3\pi \nu_\tau$
587 ± 27 ± 21	5904	¹⁰ ABREU	98G DLPH	$e^+ e^-$
478 ± 3 ± 15	5904	¹¹ ABREU	98G DLPH	$e^+ e^-$
425 ± 14 ± 8	5904	^{12,13} ABREU	98G DLPH	$e^+ e^-$
400 ± 35		BARBERIS	98B	$450 p p \rightarrow$ $p_f \pi^+ \pi^- \pi^0 p_s$
621 ± 32 ± 58		^{10,14} ACKERSTAFF	97R OPAL	$E_{cm}^{e^+e^-} = 88-94, \tau \rightarrow$ $3\pi \nu$

OCCUR=2
 OCCUR=3

457	± 15	± 17	11,14	ACKERSTAFF	97R	OPAL	$E_{\text{cm}}^{\text{e}^{\text{e}}} = 88-94, \tau \rightarrow 3\pi\nu$	OCCUR=2
446	± 21	$+140$ -0	11	ALBRECHT	93C	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
239	± 11			ANDO	92	SPEC	$8 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^0 n$	
266	± 13	± 4	15	ANDO	92	SPEC	$8 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^0 n$	OCCUR=3
465	$+228$ -143		16	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	
298	$+40$ -34		17	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=2
488	± 32		18	IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$	OCCUR=3
430	± 50			ARMSTRONG	90	OMEG	$300.0 p p \rightarrow p p \pi^+ \pi^- \pi^0$	
420	± 40		19	ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
396	± 43		20	BOWLER	88	RVUE		
405	± 75	± 25		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
419	± 108	± 57		BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$	OCCUR=2
521	± 27			ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
476	$+132$ -120	± 54		RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
462	± 56	± 30		SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$	
292	± 40			BELLINI	85	SPEC	$40 \pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
380	± 100		21	DANKOWY...	81	SPEC	$8.45 \pi^- \rho \rightarrow n 3\pi$	
300	± 50		21	DAUM	81B	CNTR	$63,94 \pi^- \rho \rightarrow p 3\pi$	
230	± 50		22	GAVILLET	77	HBC	$4.2 K^- p \rightarrow \Sigma 3\pi$	

¹ Statistical error negligible.

² Superseded by AGHASYAN 2018B.

³ The $\rho^\pm \pi^\mp$ state can be also due to the $\pi(1300)$.

⁴ Using the Breit-Wigner parameterization; strong correlation between mass and width.

⁵ Using the data of BARATE 98R.

⁶ From a fit of the $K^- K^{*0}$ distribution assuming $m_{a_1} = 1230$ MeV and purely resonant production of the $K^- K^{*0}$ system.

⁷ From a fit to the 3π mass spectrum including the $K \bar{K}^*(892)$ threshold.

⁸ Using the $a_1(1260)$ mass of 1230 MeV.

⁹ From AKHMETSIN 99E and ASNER 00 data using the $a_1(1260)$ mass of 1230 MeV.

¹⁰ Uses the model of KUHNS 90.

¹¹ Uses the model of ISGUR 89.

¹² Includes the effect of a possible a_1' state.

¹³ Uses the model of FEINDT 90.

¹⁴ Supersedes AKERS 95P.

¹⁵ Average and spread of values using 2 variants of the model of BOWLER 75.

¹⁶ Reanalysis of RUCKSTUHL 86.

¹⁷ Reanalysis of SCHMIDKE 86.

¹⁸ Reanalysis of ALBRECHT 86B.

¹⁹ From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

²⁰ From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

²¹ Uses the model of BOWLER 75.

²² Produced in K^- backward scattering.

NODE=M010W;LINKAGE=Q
 NODE=M010W;LINKAGE=R
 NODE=M010W;LINKAGE=AU
 NODE=M010W;LINKAGE=LI
 NODE=M010W;LINKAGE=GO
 NODE=M010W;LINKAGE=DR

NODE=M010W;LINKAGE=B6
 NODE=M010W;LINKAGE=WE
 NODE=M010W;LINKAGE=WB
 NODE=M010W;LINKAGE=KS
 NODE=M010W;LINKAGE=IM
 NODE=M010W;LINKAGE=A1
 NODE=M010W;LINKAGE=F1
 NODE=M010W;LINKAGE=X
 NODE=M010W;LINKAGE=P
 NODE=M010W;LINKAGE=I
 NODE=M010W;LINKAGE=L
 NODE=M010W;LINKAGE=M
 NODE=M010W;LINKAGE=K
 NODE=M010W;LINKAGE=G
 NODE=M010W;LINKAGE=D
 NODE=M010W;LINKAGE=F

$a_1(1260)$ DECAY MODES

NODE=M010215;NODE=M010

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $(\rho\pi)S$ -wave, $\rho \rightarrow \pi\pi$	seen
Γ_3 $(\rho\pi)D$ -wave, $\rho \rightarrow \pi\pi$	seen
Γ_4 $(\rho(1450)\pi)S$ -wave, $\rho \rightarrow \pi\pi$	seen
Γ_5 $(\rho(1450)\pi)D$ -wave, $\rho \rightarrow \pi\pi$	seen
Γ_6 $f_0(500)\pi$, $f_0 \rightarrow \pi\pi$	seen
Γ_7 $f_0(980)\pi$, $f_0 \rightarrow \pi\pi$	seen
Γ_8 $f_0(1370)\pi$, $f_0 \rightarrow \pi\pi$	seen
Γ_9 $f_2(1270)\pi$, $f_2 \rightarrow \pi\pi$	seen
Γ_{10} $\pi^+ \pi^- \pi^0$	seen
Γ_{11} $\pi^0 \pi^0 \pi^0$	not seen
Γ_{12} $K K \pi$	seen
Γ_{13} $K^*(892) K$	seen
Γ_{14} $\pi \gamma$	seen

DESIG=17;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=8;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=10;OUR EST;→ UNCHECKED ←
 DESIG=16;OUR EST;→ UNCHECKED ←
 DESIG=11
 DESIG=12;OUR EST;→ UNCHECKED ←
 DESIG=13;OUR EST;→ UNCHECKED ←
 DESIG=22
 DESIG=23;OUR EST;→ UNCHECKED ←
 DESIG=18;OUR EST;→ UNCHECKED ←
 DESIG=14;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←

$a_1(1260)$ PARTIAL WIDTHS

NODE=M010220

 $\Gamma(\pi\gamma)$ Γ_{14}

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
640 ± 246	ZIELINSKI	84C SPEC	200 $\pi^+ Z \rightarrow Z 3\pi$


NODE=M010W4
NODE=M010W4 D -wave/ S -wave AMPLITUDE RATIO IN DECAY OF $a_1(1260) \rightarrow \rho\pi$

NODE=M010DS

VALUE	DOCUMENT ID	TECN	COMMENT
-0.062 ± 0.020 OUR AVERAGE	Error includes scale factor of 2.3. See the ideogram below.		

NODE=M010DS

$-0.043 \pm 0.009 \pm 0.005$	LINK	07A	FOCS $D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
$-0.14 \pm 0.04 \pm 0.07$	¹ CHUNG	02	B852 $18.3 \pi^- \rho \rightarrow \pi^+ \pi^- \pi^- \rho$
$-0.10 \pm 0.02 \pm 0.02$	^{2,3} ACKERSTAFF	97R	OPAL $E_{cm}^{ee} = 88-94, \tau \rightarrow 3\pi\nu$
-0.11 ± 0.02	² ALBRECHT	93C	ARG $\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$

¹ Deck-type background not subtracted.² Uses the model of ISGUR 89.³ Supersedes AKERS 95P.NODE=M010DS;LINKAGE=C
NODE=M010DS;LINKAGE=IM
NODE=M010DS;LINKAGE=X $a_1(1260)$ BRANCHING RATIOS

NODE=M010225

 $\Gamma((\rho\pi)_{S\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M010R5
NODE=M010R5

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
60.19	37k	¹ ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$

¹ From a fit to the Dalitz plot.

NODE=M010R5;LINKAGE=B1

 $\Gamma((\rho\pi)_{D\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M010R6
NODE=M010R6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.30 \pm 0.60 \pm 0.22$	37k	¹ ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$

¹ From a fit to the Dalitz plot.

NODE=M010R6;LINKAGE=B1

 $\Gamma((\rho(1450)\pi)_{S\text{-wave}, \rho \rightarrow \pi\pi})/\Gamma_{\text{total}}$ Γ_4/Γ NODE=M010R7
NODE=M010R7

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.56 \pm 0.84 \pm 0.32$	37k	^{1,2} ASNER	00	CLE2 $10.6 e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$

¹ From a fit to the Dalitz plot.² Assuming for $\rho(1450)$ mass and width of 1370 and 386 MeV respectively.NODE=M010R7;LINKAGE=B1
NODE=M010R7;LINKAGE=B2

$\Gamma((\rho(1450)\pi)_{D\text{-wave}}, \rho \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.04 \pm 1.20 \pm 0.28$	37k	^{1,2} ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
--------------------------	-----	----------------------	----	---

¹ From a fit to the Dalitz plot.

² Assuming for $\rho(1450)$ mass and width of 1370 and 386 MeV respectively.

NODE=M010R8
NODE=M010R8

NODE=M010R8;LINKAGE=B1
NODE=M010R8;LINKAGE=B2

 $\Gamma(f_0(500)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen		CHUNG	02	B852 18.3 $\pi^-p \rightarrow \pi^+\pi^-\pi^-p$
$18.76 \pm 4.29 \pm 1.48$	37k	^{1,2} ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$

¹ From a fit to the Dalitz plot.

² Assuming for $f_0(500)$ (σ) mass and width of 860 and 880 MeV respectively.

NODE=M010R9
NODE=M010R9

NODE=M010R9;LINKAGE=B1
NODE=M010R9;LINKAGE=B3

 $\Gamma(f_0(500)\pi, f_0 \rightarrow \pi\pi)/[\Gamma((\rho\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi) + \Gamma((\rho\pi)_{S\text{-wave}}, \rho \rightarrow \pi\pi)]$ $\Gamma_6/(\Gamma_2+\Gamma_2)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.06 ± 0.05	90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+2\pi^-$
~ 0.3	28k	AKHMETSHIN	99E	CMD2 1.05–1.38 $e^+e^- \rightarrow \pi^+\pi^-\pi^+\pi^-$
0.003 ± 0.003		¹ LONGACRE	82	RVUE

¹ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from GAVILLET 77, DAUM 80, and DANKOWYCH 81.

NODE=M010R4
NODE=M010R4

NODE=M010R4;LINKAGE=E

 $\Gamma(f_0(980)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen ¹ ALEXEEV 21 COMP $\pi^-p \rightarrow \pi^-\pi^+\pi^-p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	37k	ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
----------	-----	-------	----	---

¹ The $a_1(1260)^- \rightarrow f_0(980)\pi^-$ decay mode via the Triangle Singularity mechanism from MIKHASENKO 15 and ACETI 16 explains the $a_1(1420)^-$ signal observed by ADOLPH 15C.

NODE=M010R10
NODE=M010R10

NODE=M010R10;LINKAGE=A

 $\Gamma(f_0(1370)\pi, f_0 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.40 \pm 2.71 \pm 1.26$	37k	^{1,2} ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
--------------------------	-----	----------------------	----	---

¹ From a fit to the Dalitz plot.

² Assuming for $f_0(1370)$ mass and width of 1186 and 350 MeV respectively.

NODE=M010R11
NODE=M010R11

NODE=M010R11;LINKAGE=B1
NODE=M010R11;LINKAGE=B4

 $\Gamma(f_2(1270)\pi, f_2 \rightarrow \pi\pi)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.19 \pm 0.49 \pm 0.17$	37k	^{1,2} ASNER	00	CLE2 10.6 $e^+e^- \rightarrow \tau^+\tau^-$, $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$
--------------------------	-----	----------------------	----	---

¹ From a fit to the Dalitz plot.

² Assuming for $f_2(1270)$ mass and width of 1275 and 185 MeV respectively.

NODE=M010R12
NODE=M010R12

NODE=M010R12;LINKAGE=B1
NODE=M010R12;LINKAGE=B5

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	COMMENT
-------	-------------	---------

seen BARBERIS 98B 450 $pp \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$

NODE=M010R00
NODE=M010R00

 $\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{11}/Γ_{10}

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.008	90	¹ BARBERIS	01 450 $pp \rightarrow p_f 3\pi^0 p_s$
-----------	----	-----------------------	--

¹ Inconsistent with observations of $\sigma\pi$, $f_0(1370)\pi$, and $f_2(1270)\pi$ decay modes.

NODE=M010R15
NODE=M010R15

NODE=M010R15;LINKAGE=RB

$\Gamma(K^*(892)K)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.2±0.5	2255	1 COAN	04 CLEO	$\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$
8 to 15	205	2 DRUTSKOY	02 BELL	$B \rightarrow D^{(*)} K^- K^{*0}$
3.3±0.5±0.1	37k	3 ASNER	00 CLE2	10.6 $e^+ e^- \rightarrow \tau^+ \tau^-$, $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$
2.6±0.3		4 BARATE	99R ALEP	$\tau \rightarrow K \bar{K} \pi \nu_\tau$

¹ Using structure functions from KUHN 92 and DECKER 93A and $B(\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau) = (0.155 \pm 0.006 \pm 0.009)\%$ from BRIERE 03.

² From a comparison to ALAM 94 assuming purely resonant production of the $K^- K^{*0}$ system.

³ From a fit to the 3π mass spectrum including the $K\bar{K}^*(892)$ threshold.

⁴ Assuming $a_1(1260)$ dominance and taking $B(\tau \rightarrow a_1(1260) \nu_\tau)$ from BUSKULIC 96.

NODE=M010R13
NODE=M010R13

NODE=M010R13;LINKAGE=CO

NODE=M010R13;LINKAGE=DR

NODE=M010R13;LINKAGE=B6

NODE=M010R13;LINKAGE=BA

$a_1(1260)$ REFERENCES

NODE=M010

ALEXEEV	21	PRL 127 082501	G.D. Alexeev <i>et al.</i>	(COMPASS Collab.)	REFID=62008
AAIJ	18A1	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59187
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
MIKHASENKO	18	PR D98 096021	M. Mikhasenko <i>et al.</i>	(JPAC Collab.)	REFID=59487
DARGENT	17	JHEP 1705 143	P. dArgent <i>et al.</i>	(HEID, BRIS)	REFID=58121
ACETI	16	PR D94 096015	F. Aceti, L.R. Dai, E. Oset	(IFIC, LNUDA)	REFID=62015
ADOLPH	15C	PRL 115 082001	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56790
MIKHASENKO	15	PR D91 094015	M. Mikhasenko, B. Ketzner, A. Sarantsev	(BONN+)	REFID=56825
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
LINK	07A	PR D75 052003	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=51713
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
GOMEZ-DUM...	04	PR D69 073002	D. Gomez Dumm, A. Pich, J. Portoles		REFID=49771
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)	REFID=53226
BRIERE	03	PRL 90 181802	R. A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=49360
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
DRUTSKOY	02	PL B542 171	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=48780
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48324
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47339
AKHMETSHIN	99E	PL B466 392	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47411
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47366
BONDAR	99	PL B466 403	A.E. Bondar <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47358
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=45909
BARATE	98R	EPJ C4 409	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=46147
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
ACKERSTAFF	97R	ZPHY C75 593	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45616
BUSKULIC	96	ZPHY C70 579	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44588
AKERS	95P	ZPHY C67 45	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44366
ALAM	94	PR D50 43	M.S. Alam <i>et al.</i>	(CLEO Collab.)	REFID=43738
ALBRECHT	93C	ZPHY C58 61	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=43310
DECKER	93A	ZPHY C58 445	R. Decker <i>et al.</i>		REFID=51577
ANDO	92	PL B291 496	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)	REFID=43171
KUHN	92	ZPHY C56 661	J.H. Kuhn, E. Mirkes		REFID=51576
IVANOV	91	ZPHY C49 563	Y.P. Ivanov, A.A. Osipov, M.K. Volkov	(JINR)	REFID=41750
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch	(WA76 Coll.)	REFID=41375
FEINDT	90	ZPHY C48 681	M. Feindt	(HAMB)	REFID=45912
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ISGUR	89	PR D39 1357	N. Isgur, C. Morningstar, C. Reader	(TNTO)	REFID=40730
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)	REFID=40578
BAND	87	PL B198 297	H.R. Band <i>et al.</i>	(MAC Collab.)	REFID=40263
TORNQVIST	87	ZPHY C36 695	N.A. Tornqvist	(HELS)	REFID=40030
ALBRECHT	86B	ZPHY C33 7	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=20884
RUCKSTUHL	86	PRL 56 2132	W. Ruckstuhl <i>et al.</i>	(DELCO Collab.)	REFID=10349
SCHMIDKE	86	PRL 57 527	W.B. Schmidke <i>et al.</i>	(Mark II Collab.)	REFID=10350
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>		REFID=47490
ZIELINSKI	84C	PRL 52 1195	M. Zielinski <i>et al.</i>	(ROCH, MINN, FNAL)	REFID=20882
LONGACRE	82	PR D26 82	R.S. Longacre	(BNL)	REFID=20878
DANKOWY...	81	PRL 46 580	J.A. Dankowycz <i>et al.</i>	(TNTO, BNL, CARL+)	REFID=20572
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=20872
DAUM	80	PL B9B 281	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP	REFID=20868
GAVILLET	77	PL B9B 119	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+) JP	REFID=20852
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)	REFID=20571

$f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M005

 $f_2(1270)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M005PP

NODE=M005PP

NODE=M005PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1260–1283) – i (90–110) OUR ESTIMATE			
$(1268 \pm 8) - i (101 \pm 6)$	RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1263.3 \pm 0.2 \pm 1.5) - i$ $(96.9 \pm 0.2 \pm 0.8)$	ALBRECHT	20	RVUE $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta,$ $\pi^0K^+K^-$
$(1270 \pm 8) - i (97 \pm 8)$	¹ ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1278 \pm 5) - i (102 \pm 10)$	¹ BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$

¹ Amplitude did not include dispersive corrections.

NODE=M005PP;LINKAGE=A

 $f_2(1270)$ MASS

NODE=M005M

NODE=M005M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1275.4 ± 0.8 OUR AVERAGE				
Error includes scale factor of 1.1.				
$1275.8 \pm 1.0 \pm 0.4$		¹ BOGOLYUB...	13	SPEC $7\pi^+(K^+,p)A \rightarrow n\gamma + X$
$1262 \pm \frac{1}{2} \pm 8$		² ABLIKIM	06v	BES2 $e^+e^- \rightarrow J/\psi \rightarrow$ $\gamma\pi^+\pi^-$
1275 ± 15		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$
1283 ± 5		ALDE	98	GAM4 $100 \pi^-p \rightarrow \pi^0\pi^0n$
1272 ± 8	200k	PROKOSHKIN	94	GAM2 $38 \pi^-p \rightarrow \pi^0\pi^0n$
1269.7 ± 5.2	5730	AUGUSTIN	89	DM2 $e^+e^- \rightarrow 5\pi$
1283 ± 8	400	³ ALDE	87	GAM4 $100 \pi^-p \rightarrow 4\pi^0n$
1274 ± 5		³ AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$
1283 ± 6		⁴ LONGACRE	86	MPS $22 \pi^-p \rightarrow n2K_S^0$
1276 ± 7		COURAU	84	DLCO $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
1273.3 ± 2.3		⁵ CHABAUD	83	ASPK $17 \pi^-p$ polarized
1280 ± 4		⁶ CASON	82	STRC $8 \pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$
1281 ± 7	11600	GIDAL	81	MRK2 J/ψ decay
1282 ± 5		⁷ CORDEN	79	OMEG $12-15 \pi^-p \rightarrow n2\pi$
1269 ± 4	10k	APEL	75	NICE $40 \pi^-p \rightarrow n2\pi^0$
1272 ± 4	4600	ENGLER	74	DBC $6 \pi^+n \rightarrow \pi^+\pi^-p$
1277 ± 4	5300	FLATTE	71	HBC $7.0 \pi^+p$
1273 ± 8		³ STUNTEBECK	70	HBC $8 \pi^-p, 5.4 \pi^+d$
1265 ± 8		BOESEBECK	68	HBC $8 \pi^+p$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1257 ± 6		⁸ KLEMP	22	RVUE $J/\psi(1S) \rightarrow \gamma\pi^0\pi^0,$ $\gamma K_S^0 K_S^0$
1263 ± 12		CARVER	21	CLAS $\gamma p \rightarrow \pi^0\pi^0p$
$1259 \pm 4 \pm 4$	1.7k	^{9,10} DOBBS	15	$J/\psi \rightarrow \gamma\pi^+\pi^-$
$1267 \pm 4 \pm 3$	1.5k	^{9,10} DOBBS	15	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1277 ± 6	870	¹¹ SCHEGELSKY	06A	RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$
1251 ± 10		TIKHOMIROV	03	SPEC $40.0 \pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$
1260 ± 10		¹² ALDE	97	GAM2 $450 pp \rightarrow pp\pi^0\pi^0$
1278 ± 6		¹² GRYGOREV	96	SPEC $40 \pi^-N \rightarrow K_S^0 K_S^0 X$
1262 ± 11		AGUILAR-...	91	EHS $400 pp$
1275 ± 10		AKER	91	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$
1220 ± 10		BREAKSTONE	90	SFM $pp \rightarrow pp\pi^+\pi^-$
1288 ± 12		ABACHI	86B	HRS $e^+e^- \rightarrow \pi^+\pi^-X$
1284 ± 30	3k	BINON	83	GAM2 $38 \pi^-p \rightarrow n2\eta$
1280 ± 20	3k	APEL	82	CNTR $25 \pi^-p \rightarrow n2\pi^0$
1284 ± 10	16000	DEUTSCH...	76	HBC $16 \pi^+p$
1258 ± 10	600	TAKAHASHI	72	HBC $8 \pi^-p \rightarrow n2\pi$
1275 ± 13		ARMENISE	70	HBC $9 \pi^+n \rightarrow p\pi^+\pi^-$
1261 ± 5	1960	³ ARMENISE	68	DBC $5.1 \pi^+n \rightarrow p\pi^+MM^-$
1270 ± 10	360	³ ARMENISE	68	DBC $5.1 \pi^+n \rightarrow p\pi^0MM$
1268 ± 6		¹³ JOHNSON	68	HBC $3.7-4.2 \pi^-p$

OCCUR=2

OCCUR=2

OCCUR=2

- ¹ Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
- ² Breit-Wigner mass.
- ³ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
- ⁴ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- ⁵ From an energy-independent partial-wave analysis.
- ⁶ From an amplitude analysis of the reaction $\pi^+\pi^-\rightarrow 2\pi^0$.
- ⁷ From an amplitude analysis of $\pi^+\pi^-\rightarrow \pi^+\pi^-$ scattering data.
- ⁸ Fit of the tensor partial waves from BES3 in the multipole basis.
- ⁹ Using CLEO-c data but not authored by the CLEO Collaboration.
- ¹⁰ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 185$ MeV.
- ¹¹ From analysis of L3 data at 91 and 183–209 GeV.
- ¹² Systematic uncertainties not estimated.
- ¹³ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

NODE=M005M;LINKAGE=B
 NODE=M005M;LINKAGE=K
 NODE=M005M;LINKAGE=T
 NODE=M005M;LINKAGE=L
 NODE=M005M;LINKAGE=O
 NODE=M005M;LINKAGE=P
 NODE=M005M;LINKAGE=S
 NODE=M005M;LINKAGE=Q
 NODE=M005M;LINKAGE=C
 NODE=M005M;LINKAGE=D
 NODE=M005M;LINKAGE=SC
 NODE=M005M;LINKAGE=QQ
 NODE=M005M;LINKAGE=J

$f_2(1270)$ WIDTH

NODE=M005W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M005W

186.6^{+2.8}_{-2.2} OUR FIT Error includes scale factor of 1.5. [186.6 ± 2.3 MeV OUR 2023
 FIT Scale factor = 1.5]


NEW

185.8^{+2.8}_{-2.1} OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.

190.3 ± 1.9 ± 1.8		¹ BOGOLYUB...	13	SPEC	7π ⁺ (K ⁺ ,p)A → nγ + X	
175 ⁺⁶ ₋₄ ± 10		² ABLIKIM	06v	BES2	e ⁺ e ⁻ → J/ψ → γπ ⁺ π ⁻	
190 ± 20		ABLIKIM	05	BES2	J/ψ → φπ ⁺ π ⁻	
171 ± 10		ALDE	98	GAM4	100 π ⁻ p → π ⁰ π ⁰ n	
192 ± 5	200k	PROKOSHKIN	94	GAM2	38 π ⁻ p → π ⁰ π ⁰ n	
180 ± 24		AGUILAR-...	91	EHS	400 pp	
169 ± 9	5730	³ AUGUSTIN	89	DM2	e ⁺ e ⁻ → 5π	
150 ± 30	400	³ ALDE	87	GAM4	100 π ⁻ p → 4π ⁰ n	
186 ⁺⁹ ₋₂		⁴ LONGACRE	86	MPS	22 π ⁻ p → n2K _S ⁰	
179.2 ^{+6.9} _{-6.6}		⁵ CHABAUD	83	ASPK	17 π ⁻ p polarized	
160 ± 11		DENNEY	83	LASS	10 π ⁺ N	
196 ± 10	3k	APEL	82	CNTR	25 π ⁻ p → n2π ⁰	
152 ± 9		⁶ CASON	82	STRC	8 π ⁺ p → Δ ⁺⁺ π ⁰ π ⁰	
186 ± 27	11600	GIDAL	81	MRK2	J/ψ decay	
216 ± 13		⁷ CORDEN	79	OMEG	12–15 π ⁻ p → n2π	
190 ± 10	10k	APEL	75	NICE	40 π ⁻ p → n2π ⁰	
192 ± 16	4600	ENGLER	74	DBC	6 π ⁺ n → π ⁺ π ⁻ p	
183 ± 15	5300	FLATTE	71	HBC	7 π ⁺ p → Δ ⁺⁺ f ₂	
196 ± 30		³ STUNTEBECK	70	HBC	8 π ⁻ p, 5.4 π ⁺ d	
216 ± 20	1960	³ ARMENISE	68	DBC	5.1 π ⁺ n → pπ ⁺ MM ⁻	OCCUR=2
128 ± 27		³ BOESEBECK	68	HBC	8 π ⁺ p	
176 ± 21		^{3,8} JOHNSON	68	HBC	3.7–4.2 π ⁻ p	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
168 ± 7		⁹ KLEMP	22	RVUE	J/ψ(1S) → γπ ⁰ π ⁰ , γK _S ⁰ K _S ⁰	
183 ± 2		CARVER	21	CLAS	γp → π ⁰ π ⁰ p	
195 ± 15	870	¹⁰ SCHEGELSKY	06A	RVUE	γγ → K _S ⁰ K _S ⁰	
121 ± 26		TIKHOMIROV	03	SPEC	40.0 π ⁻ C → K _S ⁰ K _S ⁰ K _L ⁰ X	
187 ± 20		¹¹ ALDE	97	GAM2	450 pp → ppπ ⁰ π ⁰	
184 ± 10		¹¹ GRYGOREV	96	SPEC	40 π ⁻ N → K _S ⁰ K _S ⁰ X	
200 ± 10		AKER	91	CBAR	0.0 p̄p → 3π ⁰	
240 ± 40	3k	BINON	83	GAM2	38 π ⁻ p → n2η	
187 ± 30	650	³ ANTIPOV	77	CIBS	25 π ⁻ p → p3π	
225 ± 38	16000	DEUTSCH...	76	HBC	16 π ⁺ p	
166 ± 28	600	³ TAKAHASHI	72	HBC	8 π ⁻ p → n2π	
173 ± 53		³ ARMENISE	70	HBC	9 π ⁺ n → pπ ⁺ π ⁻	OCCUR=2

- 1 Averaged over six nuclear targets, no statistically significant dependence on target nucleus observed.
- 2 Breit-Wigner width
- 3 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- 4 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
- 5 From an energy-independent partial-wave analysis.
- 6 From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$.
- 7 From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.
- 8 JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.
- 9 Fit of the tensor partial waves from BES3 in the multipole basis.
- 10 From analysis of L3 data at 91 and 183–209 GeV.
- 11 Systematic uncertainties not estimated.

NODE=M005W;LINKAGE=C
 NODE=M005W;LINKAGE=D
 NODE=M005W;LINKAGE=T
 NODE=M005W;LINKAGE=L
 NODE=M005W;LINKAGE=R
 NODE=M005W;LINKAGE=Q
 NODE=M005W;LINKAGE=U
 NODE=M005W;LINKAGE=J
 NODE=M005W;LINKAGE=M
 NODE=M005W;LINKAGE=SC
 NODE=M005W;LINKAGE=QQ



$f_2(1270)$ DECAY MODES

NODE=M005215;NODE=M005

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 $\pi\pi$	(84.3 $^{+2.8}_{-1.0}$) %	S=1.2	DESIG=1
Γ_2 $\pi^+\pi^-2\pi^0$	(7.7 $^{+1.2}_{-3.1}$) %	S=1.2	DESIG=3
Γ_3 $K\bar{K}$	(4.6 ± 0.4) %	S=2.7	DESIG=4
Γ_4 $2\pi^+2\pi^-$	(2.8 ± 0.4) %	S=1.2	DESIG=2
Γ_5 $\eta\eta$	(4.0 ± 0.8) $\times 10^{-3}$	S=2.1	DESIG=7
Γ_6 $4\pi^0$	(3.0 ± 1.0) $\times 10^{-3}$		DESIG=9
Γ_7 $\gamma\gamma$	(1.42 ± 0.24) $\times 10^{-5}$	S=1.4	DESIG=8
Γ_8 $\eta\pi\pi$	< 8 $\times 10^{-3}$	CL=95%	DESIG=6
Γ_9 $K^0K^-\pi^+ + c.c.$	< 3.4 $\times 10^{-3}$	CL=95%	DESIG=5
Γ_{10} e^+e^-	< 6 $\times 10^{-10}$	CL=90%	DESIG=10

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 44 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 82.3$ for 37 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-91						
x_3	10	-39					
x_4	10	-38	1				
x_5	1	-6	0	0			
x_6	0	-7	0	0	0		
x_7	4	1	-15	0	0	0	
Γ	-72	66	-10	-7	-1	0	-6
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $\pi\pi$	157.2 $^{+5.0}_{-1.0}$	
Γ_2 $\pi^+\pi^-2\pi^0$	14.3 $^{+2.3}_{-6.0}$	1.3
Γ_3 $K\bar{K}$	8.5 ± 0.8	2.8
Γ_4 $2\pi^+2\pi^-$	5.2 ± 0.7	1.2
Γ_5 $\eta\eta$	0.75 ± 0.14	2.1
Γ_6 $4\pi^0$	0.56 ± 0.19	
Γ_7 $\gamma\gamma$	0.0026 ± 0.0005	1.4

DESIG=1

DESIG=3

DESIG=4

DESIG=2

DESIG=7

DESIG=9

DESIG=8

$f_2(1270)$ PARTIAL WIDTHS

NODE=M005220

$\Gamma(\pi\pi)$

 Γ_1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W1
NODE=M005W1

157.2 $^{+5.0}_{-1.0}$ OUR FIT

NEW

[157.2 $^{+4.0}_{-1.1}$ MeV OUR 2023 FIT]

157.0 $^{+6.0}_{-1.0}$

¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

152 ± 8 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005W1;LINKAGE=L
NODE=M005W1;LINKAGE=SC

$\Gamma(K\bar{K})$

 Γ_3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W4
NODE=M005W4

8.5 ± 0.8 OUR FIT Error includes scale factor of 2.8.

9.0 $^{+0.7}_{-0.3}$

¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.5 ± 2.0 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005W4;LINKAGE=L
NODE=M005W4;LINKAGE=SC

$\Gamma(\eta\eta)$

 Γ_5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M005W7
NODE=M005W7

0.75 ± 0.14 OUR FIT Error includes scale factor of 2.1.

1.0 ± 0.1

¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow n 2K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.4 870 ² SCHEGELSKY 06A RVUE $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M005W7;LINKAGE=L
NODE=M005W7;LINKAGE=SC

$\Gamma(\gamma\gamma)$ Γ_7

The value of this width depends on the theoretical model used. Unitary approaches with scalars typically (with exception of PENNINGTON 08) give values clustering around 2.6 keV; without an S -wave contribution, values are systematically higher (typically around 3 keV).

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ±0.5 OUR FIT	Error includes scale factor of 1.4.			
2.93 ±0.40		¹ DAI	14A	RVUE Compilation
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.14 ±0.20		^{2,3} PENNINGTON 08		RVUE Compilation
3.82 ±0.30		^{3,4} PENNINGTON 08		RVUE Compilation
2.55 ±0.15	870	⁵ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
2.84 ±0.35		BOGLIONE 99	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
2.93 ±0.23 ±0.32		⁶ YABUKI 95	VNS	
2.58 ±0.13 ^{+0.36} _{-0.27}		⁷ BEHREND 92	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.10 ±0.35 ±0.35		⁸ BLINOV 92	MD1	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.27 ±0.47 ±0.11		ADACHI 90D	TOPZ	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.15 ±0.04 ±0.39		BOYER 90	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.19 ±0.16 ^{+0.29} _{-0.28}		MARSISKE 90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
2.35 ±0.65		⁹ MORGAN 90	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^-, \pi^0 \pi^0$
3.19 ±0.09 ^{+0.22} _{-0.38}	2177	OEST 90	JADE	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
3.2 ±0.1 ±0.4		¹⁰ AIHARA 86B	TPC	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.5 ±0.1 ±0.5		BEHREND 84B	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.85 ±0.25 ±0.5		¹¹ BERGER 84	PLUT	$e^+ e^- \rightarrow e^+ e^- 2\pi$
2.70 ±0.05 ±0.20		COURAU 84	DLCO	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.52 ±0.13 ±0.38		¹² SMITH 84C	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.7 ±0.2 ±0.6		EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
2.9 ^{+0.6} _{-0.4} ±0.6		¹³ EDWARDS 82F	CBAL	$e^+ e^- \rightarrow e^+ e^- 2\pi^0$
3.2 ±0.2 ±0.6		BRANDELIK 81B	TASS	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
3.6 ±0.3 ±0.5		ROUSSARIE 81	MRK2	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$
2.3 ±0.8		¹⁴ BERGER 80B	PLUT	$e^+ e^-$

¹ Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis. Supersedes PENNINGTON 08.

² Solution A (preferred solution based on χ^2 -analysis).

³ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

⁴ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

⁵ From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

⁶ With a narrow scalar state around 1220 MeV.

⁷ Using a unitarized model with a 300 - 500 keV wide scalar at 1100 MeV.

⁸ Using the unitarized model of LYTH 85.

⁹ Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1370)$: $\Gamma(f_2) + 1/4 \Gamma(f^0) = 3.6 \pm 0.3$ KeV.

¹⁰ Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.

¹¹ Using the MENNESSIER 83 model.

¹² Superseded by BOYER 90.

¹³ If helicity = 2 assumption is not made.

¹⁴ Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.

NODE=M005W8

NODE=M005W8

NODE=M005W8

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M005W8;LINKAGE=A

NODE=M005W8;LINKAGE=P1

NODE=M005W8;LINKAGE=P3

NODE=M005W8;LINKAGE=P2

NODE=M005W8;LINKAGE=SC

NODE=M005W8;LINKAGE=YA

NODE=M005W;LINKAGE=B

NODE=M005W;LINKAGE=A

NODE=M005PW;LINKAGE=C

NODE=M005PW;LINKAGE=B

NODE=M005PW;LINKAGE=X

NODE=M005PW;LINKAGE=V

NODE=M005PW;LINKAGE=H

NODE=M005PW;LINKAGE=A

 $\Gamma(e^+ e^-)$ Γ_{10}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.11	90	ACHASOV	00K	SDN $e^+ e^- \rightarrow \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<1.7	90	VOROBYEV	88	ND $e^+ e^- \rightarrow \pi^0 \pi^0$

NODE=M005W9

NODE=M005W9

$f_2(1270) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M005223

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_7/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M005G1
NODE=M005G1**0.121±0.020 OUR FIT** Error includes scale factor of 1.3.**0.091±0.007±0.027** ¹ ALBRECHT 90G ARG $e^+e^- \rightarrow e^+e^- K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.104±0.007±0.072 ² ALBRECHT 90G ARG $e^+e^- \rightarrow e^+e^- K^+K^-$

OCCUR=2

¹ Using an incoherent background.

NODE=M005G1;LINKAGE=A

² Using a coherent background.

NODE=M005G1;LINKAGE=K

 $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_7/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M005G02
NODE=M005G02**11.5^{+1.8+4.5}_{-2.0-3.7}** ¹ UEHARA 10A BELL 10.6 $e^+e^- \rightarrow e^+e^- \eta\eta$ ¹ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_0(Y)$.

NODE=M005G02;LINKAGE=UE

Helicity-0/Helicity-2 RATIO IN $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi\pi$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M005HR0
NODE=M005HR0**3.7±0.3^{+15.9}_{-2.9}** UEHARA 08A BELL 10.6 $e^+e^- \rightarrow e^+e^- \pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.5±1.8 ¹ DAI 14A RVUE Compilation13 ^{2,3} PENNINGTON 08 RVUE Compilation26 ^{3,4} PENNINGTON 08 RVUE Compilation

OCCUR=2

OCCUR=3

¹ Based on a K -matrix analysis of BELLE data from MORI 07, UEHARA 08A, UEHARA 09 and UEHARA 13. The width is derived for the pole on the third sheet which is closest to the physical axis.

NODE=M005HR0;LINKAGE=A

² Solution A (preferred solution based on χ^2 -analysis).

NODE=M005HR0;LINKAGE=P1

³ Dispersion theory based amplitude analysis of BOYER 90, MARSISKE 90, BEHREND 92, and MORI 07.

NODE=M005HR0;LINKAGE=P3

⁴ Solution B (worse than solution A; still acceptable when systematic uncertainties are included).

NODE=M005HR0;LINKAGE=P2

 $f_2(1270)$ BRANCHING RATIOS

NODE=M005225

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R10
NODE=M005R10**0.843^{+0.028}_{-0.010} OUR FIT** Error includes scale factor of 1.2. [0.843^{+0.029}_{-0.009} OUR 2023 FIT

NEW

Scale factor = 1.2]

0.837±0.020 OUR AVERAGE0.849±0.025 CHABAUD 83 ASPK 17 $\pi^- p$ polarized0.85 ±0.05 250 BEAUPRE 71 HBC 8 $\pi^+ p \rightarrow \Delta^{++} f_2$ 0.8 ±0.04 600 OH 70 HBC 1.26 $\pi^- p \rightarrow \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.856±0.001±0.05 ¹ ALBRECHT 20 RVUE 0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0 K^+K^-$ ¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

NODE=M005R10;LINKAGE=A

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi\pi)$ Γ_2/Γ_1 Should be twice $\Gamma(2\pi^+2\pi^-)/\Gamma(\pi\pi)$ if decay is $\rho\rho$. (See ASCOLI 68D.)

NODE=M005R2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R2
NODE=M005R2**0.091^{+0.015}_{-0.040} OUR FIT** Error includes scale factor of 1.2. [0.091^{+0.014}_{-0.040} OUR 2023 FIT

NEW

Scale factor = 1.2]

0.15 ±0.06 600 EISENBERG 74 HBC 4.9 $\pi^+ p \rightarrow \Delta^{++} f_2$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 EMMS 75D DBC 4 $\pi^+ n \rightarrow p f_2$

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M005R00
 NODE=M005R00

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.033±0.001±0.005	¹ ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta,$ $\pi^0\eta\eta, \pi^0K^+K^-$
-------------------	-----------------------	----	------	--

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

NODE=M005R00;LINKAGE=A

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_3/Γ_1

We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R3

NODE=M005R3

NODE=M005R3

0.054^{+0.005}_{-0.006} OUR FIT Error includes scale factor of 2.7.

0.041^{+0.004}_{-0.005} OUR AVERAGE

0.045±0.01	¹ BARGIOTTI	03	OBLX	$\bar{p}p$	
0.037 ^{+0.008} _{-0.021}	ETKIN	82B	MPS	23 $\pi^-p \rightarrow n2K_S^0$	
0.045±0.009	CHABAUD	81	ASPK	17 π^-p polarized	
0.039±0.008	LOVERRE	80	HBC	4 $\pi^-p \rightarrow K\bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.052±0.025	ABLIKIM	04E	BES2	$J/\psi \rightarrow \omega K^+K^-$	
0.036±0.005	² COSTA	80	OMEG	1-2.2 $\pi^-p \rightarrow K^+K^-n$	
0.030±0.005	³ MARTIN	79	RVUE		
0.027±0.009	⁴ POLYCHRO...	79	STRC	7 $\pi^-p \rightarrow n2K_S^0$	
0.025±0.015	EMMS	75D	DBC	4 $\pi^+n \rightarrow pf_2$	
0.031±0.012	20	ADERHOLZ	69	HBC	8 $\pi^+p \rightarrow K^+K^-p$

¹ Coupled channel analysis of $\pi^+\pi^-\pi^0, K^+K^-\pi^0,$ and $K^\pm K_S^0\pi^\mp$.

² Re-evaluated by CHABAUD 83.

³ Includes PAWLICKI 77 data.

⁴ Takes into account the $f_2(1270)$ - $f_2'(1525)$ interference.

NODE=M005R;LINKAGE=BG

NODE=M005R3;LINKAGE=D

NODE=M005R3;LINKAGE=F

NODE=M005R3;LINKAGE=M

 $\Gamma(2\pi^+2\pi^-)/\Gamma(\pi\pi)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M005R1

NODE=M005R1

0.033±0.005 OUR FIT Error includes scale factor of 1.2.

0.033±0.004 OUR AVERAGE Error includes scale factor of 1.1.

0.024±0.006	160	EMMS	75D	DBC	4 $\pi^+n \rightarrow pf_2$
0.051±0.025	70	EISENBERG	74	HBC	4.9 $\pi^+p \rightarrow \Delta^{++}f_2$
0.043 ^{+0.007} _{-0.011}	285	¹ LOUIE	74	HBC	3.9 $\pi^-p \rightarrow nf_2$
0.037±0.007	154	ANDERSON	73	DBC	6 $\pi^+n \rightarrow pf_2$
0.047±0.013		OH	70	HBC	1.26 $\pi^-p \rightarrow \pi^+\pi^-n$

¹ LOUIE 74 was quoted as 0.065 in PDG 74. Factor 2/3 to go from $\pi^+\pi^- \rightarrow \pi\pi$ forgotten. Mike L.

NODE=M005R1;LINKAGE=02

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

NODE=M005R7

NODE=M005R7

4.0±0.8 OUR FIT Error includes scale factor of 2.1.

2.9±0.5 OUR AVERAGE

2.7±0.7	BINON	05	GAMS	33 $\pi^-p \rightarrow \eta\eta n$
2.8±0.7	ALDE	86D	GAM4	100 $\pi^-p \rightarrow 2\eta n$
5.2±1.7	BINON	83	GAM2	38 $\pi^-p \rightarrow 2\eta n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.0±1.0±2.0	¹ ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta,$ $\pi^0\eta\eta, \pi^0K^+K^-$
-------------	-----------------------	----	------	--

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

NODE=M005R7;LINKAGE=A

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$ Γ_5/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M005R6

NODE=M005R6

0.003±0.001 BARBERIS 00E 450 $pp \rightarrow pf\eta\eta p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.05	95	EDWARDS	82F	CBAL	$e^+e^- \rightarrow e^+e^-2\eta$
<0.016	95	EMMS	75D	DBC	4 $\pi^+n \rightarrow pf_2$
<0.09	95	EISENBERG	74	HBC	4.9 $\pi^+p \rightarrow \Delta^{++}f_2$

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0030 ± 0.0010 OUR FIT					
0.003 ± 0.001	400 ± 50	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$	NODE=M005R11 NODE=M005R11
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE (units 10^{-5})		DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.57 ± 0.01 ^{+1.39} _{-0.14}		UEHARA	08A	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	NODE=M005R13 NODE=M005R13
$\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$					Γ_8/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.010	95	EMMS	75D	DBC 4 $\pi^+ n \rightarrow p f_2$	NODE=M005R5 NODE=M005R5
$\Gamma(K^0 K^- \pi^+ + \text{c.c.})/\Gamma(\pi\pi)$					Γ_9/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.004	95	EMMS	75D	DBC 4 $\pi^+ n \rightarrow p f_2$	NODE=M005R4 NODE=M005R4
$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$					Γ_{10}/Γ
VALUE (units 10^{-10})	CL%	DOCUMENT ID	TECN	COMMENT	
<6	90	ACHASOV	00K	SND $e^+ e^- \rightarrow \pi^0 \pi^0$	NODE=M005R12 NODE=M005R12

 $f_2(1270)$ REFERENCES

$f_2(1270)$ REFERENCES					NODE=M005
KLEMP	22	PL B830 137171	E. Klempt <i>et al.</i>	(BONN)	REFID=61646
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
CARVER	21	PRL 126 082002	M. Carver <i>et al.</i>	(CLAS Collab.)	REFID=61097
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
DAI	14A	PR D90 036004	L.-Y. Dai, M.R. Pennington	(CEBAF)	REFID=55923
BOGOLYUB...	13	PAN 76 1324	M.Yu. Bogolyubsky <i>et al.</i>	(HYPERON-M Collab.)	REFID=55585
Translated from YAF 76 1389.					
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
PENNINGTON	08	EPJ C56 1	M.R. Pennington <i>et al.</i>		REFID=52303
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
MORI	07	PR D75 051101	T. Mori <i>et al.</i>	(BELLE Collab.)	REFID=51652
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
Translated from YAF 68 998.					
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
Translated from YAF 66 860.					
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BOGLIONE	99	EPJ C9 11	M. Boggione, M.R. Pennington		REFID=46931
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
Translated from YAF 62 446.					
ALDE	97	PL B397 350	D.M. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45392
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
GRYGOREV	96	PAN 59 2105	V.K. Grigoriev, O.N. Baloshin, B.P. Barkov	(ITEP)	REFID=45566
Translated from YAF 59 2187.					
YABUKI	95	JPSJ 64 435	F. Yabuki <i>et al.</i>	(VENUS Collab.)	REFID=46384
PROKOSHKIN	94	PD 39 420	Y.D. Prokoshkin, A.A. Kondashov	(SERP)	REFID=44094
Translated from DANS 336 613.					
BEHREND	92	ZPHY C56 381	H.J. Behrend	(CELLO Collab.)	REFID=43172
BLINOV	92	ZPHY C53 33	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41858
AGUILAR-...	91	ZPHY C50 405	M. Aguilar-Benitez <i>et al.</i>	(LEBC-EHS Collab.)	REFID=41637
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	REFID=41587
ADACHI	90D	PL B234 185	I. Adachi <i>et al.</i>	(TOPAZ Collab.)	REFID=41345
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
BOYER	90	PR D42 1350	J. Boyer <i>et al.</i>	(Mark II Collab.)	REFID=41362
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)	REFID=41376
MARSISKE	90	PR D41 3324	H. Marsiske <i>et al.</i>	(Crystal Ball Collab.)	REFID=41351
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)	REFID=41583
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
Translated from YAF 48 436.					

ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
ABACHI	86B	PRL 57 1990	S. Abachi <i>et al.</i>	(PURD, ANL, IND, MICH+)	REFID=20394
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=20764
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LANDRO	86	PL B172 445	M. Landro, K.J. Mork, H.A. Olsen	(UTRO)	REFID=20767
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
LYTH	85	JP G11 459	D.H. Lyth		REFID=42169
BEHREND	84B	ZPHY C23 223	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20757
BERGER	84	ZPHY C26 199	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20760
COURAU	84	PL 147B 227	A. Courau <i>et al.</i>	(CIT, SLAC)	REFID=20758
SMITH	84C	PR D30 851	J.R. Smith <i>et al.</i>	(SLAC, LBL, HARV)	REFID=20759
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
Also		SJNP 38 561	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20751
		Translated from YAF 38 934.			
CHABAUD	83	NP B223 1	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20131
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
MENNESSIER	83	ZPHY C16 241	G. Mennessier	(MONP)	REFID=20393
APEL	82	NP B201 197	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20745
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
BRANDELIK	81B	ZPHY C10 117	R. Brandelik <i>et al.</i>	(TASSO Collab.)	REFID=20741
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
ROUSSARIE	81	PL 105B 304	A. Roussarie <i>et al.</i>	(SLAC, LBL)	REFID=20388
BERGER	80B	PL 94B 254	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=20736
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
LOVERRE	80	ZPHY C6 187	P.F. Loverre <i>et al.</i>	(CERN, CDEF, MADR+)	REFID=20382
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
PDG	78	PL 75B 1	C. Bricman <i>et al.</i>		REFID=40124
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL)	REFID=20367
DEUTSCH...	76	NP B103 426	M. Deutschmann <i>et al.</i>	(AACH3, BERL, BONN+)	REFID=20119
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
EMMS	75D	NP B96 155	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL)	REFID=20721
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
EISENBERG	74	PL 52B 239	Y. Eisenberg <i>et al.</i>	(REHO)	REFID=20715
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
LOUIE	74	PL 48B 385	J. Louie <i>et al.</i>	(SACL, CERN)	REFID=20719
PDG	74	PL 50B 1	V. Chaloupka <i>et al.</i>		REFID=40125
ANDERSON	73	PRL 31 562	J.C. Anderson <i>et al.</i>	(CMU, CASE)	REFID=20710
TAKAHASHI	72	PR D6 1266	K. Takahashi <i>et al.</i>	(TOHOK, PENN, NDAM+)	REFID=20103
BEAUPRE	71	NP B28 77	J.V. Beaupre <i>et al.</i>	(AACH, BERL, CERN)	REFID=20698
FLATTE	71	PL 34B 551	S.M. Flatte <i>et al.</i>	(LBL)	REFID=20700
ARMENISE	70	LCN 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
OH	70	PR D1 2494	B.Y. Oh <i>et al.</i>	(WISC, TNTO) JP	REFID=20335
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)	REFID=20054
ASCOLI	68D	PRL 21 1712	G. Ascoli <i>et al.</i>	(ILL)	REFID=20681
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
JOHNSON	68	PR 176 1651	P.B. Johnson <i>et al.</i>	(NDAM, PURD, SLAC)	REFID=20065
EISNER	67	PR 164 1699	R.L. Eisner <i>et al.</i>	(PURD)	REFID=20046
DERADO	65	PRL 14 872	I. Derado <i>et al.</i>	(NDAM)	REFID=20668
LEE	64	PRL 12 342	Y.Y. Lee <i>et al.</i>	(MICH)	REFID=20663
BONDAR	63	PL 5 153	L. Bondar <i>et al.</i>	(AACH, BIRM, BONN, DESY+)	REFID=20657

NODE=M008

 $f_1(1285)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

 $f_1(1285)$ MASS

NODE=M008M

NODE=M008M

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1281.8 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below. [1281.9 ± 0.5 MeV OUR 2023 AVERAGE Scale factor = 1.8]		
1280.2 ± 0.6 ^{+1.2} _{-1.5}	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1281.0 ± 0.8		DICKSON	16 CLAS	$2.55 \gamma p \rightarrow \eta \pi^+ \pi^- p$
1287.4 ± 3.0	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
1281.16 ± 0.39 ± 0.45		¹ LEES	12X BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$
1285.1 ± 1.0 ^{+1.6} _{-0.3}		² ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
1281 ± 2 ± 1		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
1276.1 ± 8.1 ± 8.0	203	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
1274 ± 6	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
1280 ± 4		ACCIARRI	01G L3	
1288 ± 4 ± 5	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
1284 ± 6	1400	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1281 ± 1		BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
1281 ± 1		BARBERIS	97C OMEG	$450 pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
1280 ± 2		³ ANTINORI	95 OMEG	$300,450 pp \rightarrow pp2(\pi^+ \pi^-)$
1282.2 ± 1.5		LEE	94 MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
1279 ± 5		FUKUI	91C SPEC	$8.95 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
1278 ± 2	140	ARMSTRONG	89 OMEG	$300 pp \rightarrow K \bar{K} \pi pp$
1278 ± 2		ARMSTRONG	89G OMEG	$85 \pi^+ p \rightarrow 4\pi \pi p,$ $pp \rightarrow 4\pi pp$
1280.1 ± 2.1	60	RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1285 ± 1	4750	⁴ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1280 ± 1	504	BITYUKOV	88 SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
1280 ± 4		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
1277 ± 2	420	REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K K \pi X$
1285 ± 2		CHUNG	85 SPEC	$8 \pi^- p \rightarrow N K \bar{K} \pi$
1279 ± 2	604	ARMSTRONG	84 OMEG	$85 \pi^+ p \rightarrow K \bar{K} \pi \pi p,$ $pp \rightarrow K \bar{K} \pi pp$
1286 ± 1		CHAUVAT	84 SPEC	ISR 31.5 pp
1278 ± 4		EVANGELIS...	81 OMEG	$12 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
1283 ± 3	103	DIONISI	80 HBC	$4 \pi^- p \rightarrow K \bar{K} \pi n$
1282 ± 2	320	NACASCH	78 HBC	$0.7, 0.76 \bar{p} p \rightarrow K \bar{K} 3\pi$
1279 ± 5	210	GRASSLER	77 HBC	$16 \pi^\mp p$
1286 ± 3	180	DUBOC	72 HBC	$1.2 \bar{p} p \rightarrow 2K 4\pi$
1283 ± 5		DAHL	67 HBC	$1.6-4.2 \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1289.3 ± 2.8	234	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
1284.2 ± 2.2		⁵ AAIJ	14Y LHCB	$\bar{B}_s^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$
1281.9 ± 0.5		⁵ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$
1282.8 ± 0.6		⁵ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
1270 ± 10		AMELIN	95 VES	$37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
1280 ± 2		ABATZIS	94 OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
1282 ± 4		ARMSTRONG	93C E760	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1270 ± 6 ± 10		ARMSTRONG	92C OMEG	$300 pp \rightarrow pp \pi^+ \pi^- \gamma$
1281 ± 1		ARMSTRONG	89E OMEG	$300 pp \rightarrow pp2(\pi^+ \pi^-)$
1279 ± 6 ± 10	16	BECKER	87 MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$


OCCUR=2

1286 ± 9		GIDAL	87	MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
1287 ± 5	353	BITYUKOV	84B	SPEC	$32 \pi^- p \rightarrow K^+K^-\pi^0 n$
~ 1279		⁶ TORNQVIST	82B	RVUE	
1275 ± 6	31	BROMBERG	80	SPEC	$100 \pi^- p \rightarrow K\bar{K}\pi X$
1288 ± 9	200	GURTU	79	HBC	$4.2 K^- p \rightarrow n\eta 2\pi$
~ 1275.0	46	⁷ STANTON	79	CNTR	$8.5 \pi^- p \rightarrow n2\gamma 2\pi$
1271 ± 10	34	CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow K^+K^-\pi n$
1295 ± 12	85	CORDEN	78	OMEG	$12-15 \pi^- p \rightarrow n5\pi$
1292 ± 10	150	DEFOIX	72	HBC	$0.7 \bar{p} p \rightarrow 7\pi$
1280 ± 3	500	⁸ THUN	72	MMS	$13.4 \pi^- p$
1303 ± 8		BARDADIN-...	71	HBC	$8 \pi^+ p \rightarrow p6\pi$
1283 ± 6		BOESEBECK	71	HBC	$16.0 \pi p \rightarrow p5\pi$
1270 ± 10		CAMPBELL	69	DBC	$2.7 \pi^+ d$
1285 ± 7		LORSTAD	69	HBC	$0.7 \bar{p} p, 4,5\text{-body}$
1290 ± 7		D'ANDLAU	68	HBC	$1.2 \bar{p} p, 5-6 \text{ body}$

OCCUR=2

- ¹ Using the $2\pi^+2\pi^-$ and $\pi^+\pi^-\eta$ modes of $f_1(1285)$ decay.
- ² The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.
- ³ Supersedes ABATZIS 94, ARMSTRONG 89E.
- ⁴ From partial wave analysis of $K^+\bar{K}^0\pi^-$ system.
- ⁵ No systematic error given.
- ⁶ From a unitarized quark-model calculation.
- ⁷ From phase shift analysis of $\eta\pi^+\pi^-$ system.
- ⁸ Seen in the missing mass spectrum.

NODE=M008M;LINKAGE=LE
 NODE=M008M;LINKAGE=BL
 NODE=M008M;LINKAGE=B
 NODE=M008M;LINKAGE=A
 NODE=M008M;LINKAGE=N1
 NODE=M008M;LINKAGE=T
 NODE=M008M;LINKAGE=P
 NODE=M008M;LINKAGE=S



$f_1(1285)$ WIDTH

NODE=M008W

Only experiments giving width error less than 20 MeV are kept for averaging.

NODE=M008W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
23.0 ± 1.1 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
[22.7 ± 1.1 MeV OUR 2023 AVERAGE Scale factor = 1.5]				
28.2 ± 1.1 ⁺ _{-2.9}	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
18.4 ± 1.4		DICKSON	16 CLAS	$2.55 \gamma p \rightarrow \eta \pi^+ \pi^- p$
18.3 ± 6.3	87	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$
22.0 ± 3.1 ⁺ _{-1.5}		¹ ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta \pi^+ \pi^-)$
35 ± 6 ± 4		AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow f_1(1285) \pi^+ \pi^- \gamma$
40.0 ± 8.6 ± 9.3	203	BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \pi^+ \pi^-$
29 ± 12	237	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
45 ± 9 ± 7	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
55 ± 18	1400	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
24 ± 3		BARBERIS	97B OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
20 ± 2		BARBERIS	97C OMEG	$450 pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
36 ± 5		² ANTINORI	95 OMEG	$300,450 pp \rightarrow pp2(\pi^+ \pi^-)$
29.0 ± 4.1		LEE	94 MPS2	$18 \pi^- p \rightarrow K^+ \bar{K}^0 2\pi^- p$
25 ± 4	140	ARMSTRONG	89 OMEG	$300 pp \rightarrow K \bar{K} \pi pp$
22 ± 2	4750	³ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
25 ± 4	504	BITYUKOV	88 SPEC	$32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
19 ± 5		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
32 ± 8	420	REEVES	86 SPEC	$6.6 p \bar{p} \rightarrow K K \pi X$
22 ± 2		CHUNG	85 SPEC	$8 \pi^- p \rightarrow N K \bar{K} \pi$
32 ± 3	604	ARMSTRONG	84 OMEG	$85 \pi^+ p \rightarrow K \bar{K} \pi p, pp \rightarrow K \bar{K} \pi pp$
24 ± 3		CHAUVAT	84 SPEC	ISR 31.5 pp
29 ± 10	103	DIONISI	80 HBC	$4 \pi^- p \rightarrow K \bar{K} \pi n$
28.3 ± 6.7	320	NACASCH	78 HBC	$0.7, 0.76 \bar{p} p \rightarrow K \bar{K} 3\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
17.1 ± 3.4	234	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
32.4 ± 5.8		⁴ AAIJ	14Y LHCb	$\bar{B}^0(s) \rightarrow J/\psi 2(\pi^+ \pi^-)$
18.2 ± 1.2		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-)$
19.4 ± 1.5		⁴ SOSA	99 SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+)$
40 ± 5		ABATZIS	94 OMEG	$450 pp \rightarrow pp2(\pi^+ \pi^-)$
31 ± 5		ARMSTRONG	89E OMEG	$300 pp \rightarrow pp2(\pi^+ \pi^-)$
41 ± 12		ARMSTRONG	89G OMEG	$85 \pi^+ p \rightarrow 4\pi p, pp \rightarrow 4\pi pp$
17.9 ± 10.9	60	RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
14 ⁺²⁰ ₋₁₄ ± 10	16	BECKER	87 MRK3	$e^+ e^- \rightarrow \phi K \bar{K} \pi$
26 ± 12		EVANGELIS...	81 OMEG	$12 \pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
25 ± 15	200	GURTU	79 HBC	$4.2 K^- p \rightarrow n \eta 2\pi$
~ 10		⁵ STANTON	79 CNTR	$8.5 \pi^- p \rightarrow n 2\gamma 2\pi$
24 ± 18	210	GRASSLER	77 HBC	$16 \pi^\mp p$
28 ± 5	150	⁶ DEFOIX	72 HBC	$0.7 \bar{p} p \rightarrow 7\pi$
46 ± 9	180	⁶ DUBOC	72 HBC	$1.2 \bar{p} p \rightarrow 2K 4\pi$
37 ± 5	500	⁷ THUN	72 MMS	$13.4 \pi^- p$
10 ± 10		BOESEBECK	71 HBC	$16.0 \pi p \rightarrow p 5\pi$
30 ± 15		CAMPBELL	69 DBC	$2.7 \pi^+ d$
60 ± 15		⁶ LORSTAD	69 HBC	$0.7 \bar{p} p, 4,5\text{-body}$
35 ± 10		⁶ DAHL	67 HBC	$1.6\text{--}4.2 \pi^- p$

NODE=M008W

NEW

OCCUR=2

¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.

² Supersedes ABATZIS 94, ARMSTRONG 89E.

³ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.

⁴ No systematic error given.

⁵ From phase shift analysis of $\eta \pi^+ \pi^-$ system.

⁶ Resolution is not unfolded.

⁷ Seen in the missing mass spectrum.

NODE=M008W;LINKAGE=BL

NODE=M008W;LINKAGE=B


NODE=M008W;LINKAGE=A

NODE=M008W;LINKAGE=N1

NODE=M008W;LINKAGE=P

NODE=M008W;LINKAGE=R

NODE=M008W;LINKAGE=S



$f_1(1285)$ DECAY MODES

NODE=M008215;NODE=M008

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 4π	$(32.7 \pm 1.8) \%$	S=1.2	DESIG=21
Γ_2 $\pi^0 \pi^0 \pi^+ \pi^-$	$(21.8 \pm 1.2) \%$	S=1.2	DESIG=22
Γ_3 $2\pi^+ 2\pi^-$	$(10.9 \pm 0.6) \%$	S=1.2	DESIG=20
Γ_4 $\rho^0 \pi^+ \pi^-$	$(10.9 \pm 0.6) \%$	S=1.2	DESIG=191
Γ_5 $\rho^0 \rho^0$	seen		DESIG=23
Γ_6 $4\pi^0$	$< 7 \times 10^{-4}$	CL=90%	DESIG=7
Γ_7 $\eta \pi^+ \pi^-$	$(35 \pm 15) \%$		DESIG=198
Γ_8 $\eta \pi \pi$	$(52.2 \pm 1.9) \%$	S=1.2	DESIG=3
Γ_9 $a_0(980) \pi$ [ignoring $a_0(980) \rightarrow K \bar{K}$]	$(38 \pm 4) \%$		DESIG=4
Γ_{10} $\eta \pi \pi$ [excluding $a_0(980) \pi$]	$(14 \pm 4) \%$		DESIG=5
Γ_{11} $K \bar{K} \pi$	$(9.0 \pm 0.4) \%$	S=1.1	DESIG=1
Γ_{12} $K \bar{K}^*(892)$	not seen		DESIG=6
Γ_{13} $\pi^+ \pi^- \pi^0$	$(3.0 \pm 0.9) \times 10^{-3}$		DESIG=197
Γ_{14} $\rho^\pm \pi^\mp$	$< 3.1 \times 10^{-3}$	CL=95%	DESIG=199
Γ_{15} $\gamma \rho^0$	$(6.1 \pm 1.0) \%$	S=1.7	DESIG=13
Γ_{16} $\phi \gamma$	$(7.4 \pm 2.6) \times 10^{-4}$		DESIG=10
Γ_{17} $e^+ e^-$	$< 9.4 \times 10^{-9}$	CL=90%	DESIG=200
Γ_{18} $\gamma \gamma^*$			DESIG=9
Γ_{19} $\gamma \gamma$			DESIG=8

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 18 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 24.0$ for 14 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_9	-29			
x_{10}	-12	-89		
x_{11}	22	-9	-4	
x_{15}	-24	-8	-3	-27
	x_1	x_9	x_{10}	x_{11}

$f_1(1285) \Gamma(i) \Gamma(\gamma\gamma) / \Gamma(\text{total})$

NODE=M008217

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{19} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{19} / \Gamma$			
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.62	95	GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

NODE=M008G2
NODE=M008G2

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*) / \Gamma_{\text{total}}$		$\Gamma_8 \Gamma_{18} / \Gamma = (\Gamma_9 + \Gamma_{10}) \Gamma_{18} / \Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.4 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.4.			
1.18 ± 0.25 ± 0.20	26	^{1,2} AIHARA	88B	TPC	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
2.30 ± 0.61 ± 0.42		^{1,3} GIDAL	87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$

NODE=M008G3
NODE=M008G3

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 0.3 ± 0.3	420	⁴ ACHARD	02B	L3	183-209 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
-----------------	-----	---------------------	-----	----	--

¹ Assuming a ρ -pole form factor.

² Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.

³ Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49.

⁴ Published value multiplied by the $\eta\pi\pi$ branching ratio 0.52.

NODE=M008G3;LINKAGE=A
NODE=M008G3;LINKAGE=F
NODE=M008G3;LINKAGE=B
NODE=M008G3;LINKAGE=AC

$f_1(1285)$ BRANCHING RATIOS

NODE=M008220

$\Gamma(K\bar{K}\pi) / \Gamma(4\pi)$		Γ_{11} / Γ_1			
VALUE		DOCUMENT ID	TECN	COMMENT	
0.274 ± 0.017 OUR FIT		Error includes scale factor of 1.4. [0.274 ± 0.017 OUR 2023 FIT Scale factor = 1.4]			
0.271 ± 0.016 OUR AVERAGE		Error includes scale factor of 1.2.			

NODE=M008R1
NODE=M008R1
NEW

0.265 ± 0.014	¹ BARBERIS	97C	OMEG 450	$\rho\rho \rightarrow \rho\rho K_S^0 K^\pm \pi^\mp$
0.28 ± 0.05	² ARMSTRONG	89E	OMEG 300	$\rho\rho f_1(1285)$
0.37 ± 0.03 ± 0.05	³ ARMSTRONG	89G	OMEG 85	$\pi\rho \rightarrow 4\pi X$

¹ Using $2(\pi^+ \pi^-)$ data from BARBERIS 97B.

² Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.

³ 4π consistent with being entirely $\rho\pi\pi$.

NODE=M008R1;LINKAGE=B
NODE=M008R1;LINKAGE=M
NODE=M008R1;LINKAGE=A

$\Gamma(\pi^0 \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$		$\Gamma_2 / \Gamma = \frac{2}{3} \Gamma_1 / \Gamma$			
VALUE		DOCUMENT ID	TECN	COMMENT	
0.218 ± 0.012 OUR FIT		Error includes scale factor of 1.2. [0.218 ± 0.013 OUR 2023 FIT Scale factor = 1.2]			

NODE=M008R18
NODE=M008R18
NEW

$\Gamma(2\pi^+ 2\pi^-) / \Gamma_{\text{total}}$		$\Gamma_3 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$			
VALUE		DOCUMENT ID	TECN	COMMENT	
0.109 ± 0.006 OUR FIT		Error includes scale factor of 1.2. [0.109 ± 0.006 OUR 2023 FIT Scale factor = 1.2]			

NODE=M008R17
NODE=M008R17
NEW

$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$		$\Gamma_4 / \Gamma = \frac{1}{3} \Gamma_1 / \Gamma$			
VALUE		DOCUMENT ID	TECN	COMMENT	
0.109 ± 0.006 OUR FIT		Error includes scale factor of 1.2. [0.109 ± 0.006 OUR 2023 FIT Scale factor = 1.2]			

NODE=M008R19
NODE=M008R19
NEW

$$\Gamma(\rho^0 \pi^+ \pi^-) / \Gamma(2\pi^+ 2\pi^-)$$

 Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
1.0 ± 0.4	GRASSLER 77	HBC	16 GeV $\pi^\pm p$

NODE=M008R6
 NODE=M008R6

$$\Gamma(\rho^0 \rho^0) / \Gamma_{\text{total}}$$

 Γ_5/Γ

VALUE	DOCUMENT ID	COMMENT
seen	BARBERIS 00C	450 $pp \rightarrow p_f 4\pi p_s$

NODE=M008R21
 NODE=M008R21

$$\Gamma(4\pi^0) / \Gamma_{\text{total}}$$

 Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	ALDE 87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$

NODE=M008R8
 NODE=M008R8

$$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma(\eta \pi^+ \pi^-)$$

 Γ_{13}/Γ_7

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
0.86 ± 0.16 ± 0.20	2.3k	¹ DOROFEEV 11	VES	$\pi^- N \rightarrow \pi^- f_1(1285) N$

NODE=M008R02
 NODE=M008R02

¹ Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+ \pi^-$ mass spectrum.

NODE=M008R02;LINKAGE=DO

$$\Gamma(\eta \pi \pi) / \Gamma_{\text{total}}$$

 $\Gamma_8/\Gamma = (\Gamma_9 + \Gamma_{10})/\Gamma$

VALUE	DOCUMENT ID
0.522 ± 0.019 OUR FIT	Error includes scale factor of 1.2. [0.522 ± 0.020 OUR 2023 FIT Scale factor = 1.2]

NODE=M008R22
 NODE=M008R22
 NEW

$$\Gamma(4\pi) / \Gamma(\eta \pi \pi)$$

 $\Gamma_1/\Gamma_8 = \Gamma_1/(\Gamma_9 + \Gamma_{10})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.63 ± 0.06 OUR FIT	Error includes scale factor of 1.2. [0.63 ± 0.06 OUR 2023 FIT Scale factor = 1.3]		

NODE=M008R4
 NODE=M008R4
 NEW

0.41 ± 0.14 OUR AVERAGE

0.37 ± 0.11 ± 0.11	BOLTON 92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.64 ± 0.40	GURTU 79	HBC	4.2 $K^- p$
0.93 ± 0.30	¹ GRASSLER 77	HBC	16 $\pi^\mp p$

¹ Assuming $\rho \pi \pi$ and $a_0(980) \pi$ intermediate states.

NODE=M008R4;LINKAGE=M

$$\Gamma(2\pi^+ 2\pi^-) / \Gamma(\eta \pi \pi)$$

 Γ_3/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
0.28 ± 0.02 ± 0.02	¹ LEES 12X	BABR	$\tau^- \rightarrow \pi^- f_1(1285) \nu_\tau$

NODE=M008R04
 NODE=M008R04

¹ Assuming $B(f_1(1285) \rightarrow \pi \pi \eta) = 3/2 B(f_1(1285) \rightarrow \pi^+ \pi^- \eta)$.

NODE=M008R04;LINKAGE=LE

$$\Gamma(a_0(980) \pi [\text{ignoring } a_0(980) \Rightarrow K \bar{K}]) / \Gamma(\eta \pi \pi)$$

 $\Gamma_9/\Gamma_8 = \Gamma_9/(\Gamma_9 + \Gamma_{10})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.08 OUR FIT				
0.72 ± 0.07 OUR AVERAGE				
0.74 ± 0.02 ± 0.09		DICKSON 16	CLAS	$\gamma p \rightarrow f_1(1285) p$
0.72 ± 0.15		GURTU 79	HBC	4.2 $K^- p$
0.6 ^{+0.3} _{-0.2}		CORDEN 78	OMEG	12-15 $\pi^- p$

NODE=M008R3
 NODE=M008R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.69	95	ACHARD 02B	L3	183-209 $e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
0.28 ± 0.07		ALDE 97B	GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1.0 ± 0.3		GRASSLER 77	HBC	16 $\pi^\mp p$

$$\Gamma(K \bar{K} \pi) / \Gamma(\eta \pi \pi)$$

 $\Gamma_{11}/\Gamma_8 = \Gamma_{11}/(\Gamma_9 + \Gamma_{10})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.172 ± 0.011 OUR FIT			
[0.172 ± 0.012 OUR 2023 FIT Scale factor = 1.1]			
0.176 ± 0.012 OUR AVERAGE			

NODE=M008R2
 NODE=M008R2
 NEW

0.216 ± 0.010 ± 0.031	DICKSON 16	CLAS	$\gamma p \rightarrow f_1(1285) p$
0.166 ± 0.01 ± 0.008	BARBERIS 98C	OMEG	450 $pp \rightarrow p_f f_1(1285) p_s$
0.42 ± 0.15	GURTU 79	HBC	4.2 $K^- p$
0.5 ± 0.2	¹ CORDEN 78	OMEG	12-15 $\pi^- p$
0.20 ± 0.08	² DEFOIX 72	HBC	0.7 $\bar{p} p \rightarrow 7\pi$
0.16 ± 0.08	CAMPBELL 69	DBC	2.7 $\pi^+ d$

OCCUR=2

¹ CORDEN 78 assumes low-mass $\eta \pi \pi$ region is dominantly 1^{++} . See BARBERIS 98C and MANAK 00A for discussion.

NODE=M008R2;LINKAGE=CD

² $K \bar{K}$ system characterized by the $l = 1$ threshold enhancement. (See under $a_0(980)$).

NODE=M008R2;LINKAGE=K

$\Gamma(K\bar{K}^*(892))/\Gamma_{total}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	NACASCH 78	HBC	0.7,0.76 $\bar{p}p \rightarrow K\bar{K}3\pi$
seen	¹ ACHARD 07	L3	183-209 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$

¹A clear signal of 19.8 ± 4.4 events observed at high Q^2 .

NODE=M008R5
NODE=M008R5

NODE=M008R5;LINKAGE=CH

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.055±0.074	2.3k	¹ DOROFEEV 11	VES	$\pi^-N \rightarrow \pi^-\pi^-\pi^0(1285)N$

¹Value obtained selecting the region corresponding to $f_0(980)$ in the $\pi^+\pi^-$ mass spectrum. The systematic error includes the uncertainty on the partial width $f_1 \rightarrow \eta\pi\pi$ obtained from PDG 10 data.

NODE=M008R01
NODE=M008R01

NODE=M008R01;LINKAGE=DO

$\Gamma(\rho^\pm\pi^\mp)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
<0.31	95	DOROFEEV 11	VES	$\pi^-N \rightarrow \pi^-\pi^-\pi^0(1285)N$

NODE=M008R03
NODE=M008R03

$\Gamma(\gamma\rho^0)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
6.1±1.0 OUR FIT				Error includes scale factor of 1.7. $[(6.1 \pm 1.0) \times 10^{-2} \text{ OUR 2023 FIT Scale factor} = 1.7]$

••• We do not use the following data for averages, fits, limits, etc. •••

2.8±0.7±0.6		¹ AMELIN 95	VES	37 $\pi^-N \rightarrow \pi^-\pi^+\pi^-\gamma N$
<5	95	BITYUKOV 91B	SPEC	32 $\pi^-p \rightarrow \pi^+\pi^-\gamma n$

¹Not an independent measurement.

NODE=M008R15
NODE=M008R15

NEW

NODE=M008R15;LINKAGE=A

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$ $\Gamma_{15}/\Gamma_3 = \Gamma_{15}/\frac{1}{3}\Gamma_1$

VALUE	DOCUMENT ID	TECN	COMMENT
0.55±0.10 OUR FIT			Error includes scale factor of 1.5.
0.45±0.18	¹ COFFMAN 90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+ 2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.

NODE=M008R13
NODE=M008R13

NODE=M008R13;LINKAGE=E


$\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$ $\Gamma_8/\Gamma_{15} = (\Gamma_9+\Gamma_{10})/\Gamma_{15}$

VALUE	DOCUMENT ID	TECN	COMMENT
8.6±1.6 OUR FIT			Error includes scale factor of 1.9.
8.5±2.0 OUR AVERAGE			Error includes scale factor of 2.2. See the ideogram below.
21.3±4.4	DICKSON 16	CLAS	$\gamma p \rightarrow f_1(1285)p$
10.0±1.0±2.0	BARBERIS 98C	OMEG	450 $pp \rightarrow p f_1(1285) p_S$
7.5±1.0	¹ ARMSTRONG 92C	OMEG	300 $pp \rightarrow pp\pi^+\pi^-\gamma, pp\eta\pi^+\pi^-$

¹Published value multiplied by 1.5.

NODE=M008R16
NODE=M008R16

NODE=M008R16;LINKAGE=B



$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$ Γ_{15}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

>0.035	90	¹ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
--------	----	----------------------	----	--

¹ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$.

NODE=M008R12
NODE=M008R12

NODE=M008R12;LINKAGE=F

 $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{16}/Γ_{11}

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

$0.82 \pm 0.21 \pm 0.20$	19		BITYUKOV	88	SPEC $32.5 \pi^- p \rightarrow K^+ K^- \pi^0 n$
--	----	--	----------	----	---

••• We do not use the following data for averages, fits, limits, etc. •••

<0.50	95		BARBERIS	98c	OMEG 450 $pp \rightarrow p_f f_1(1285) p_s$
-------	----	--	----------	-----	---

<0.93	95		AMELIN	95	VES $37 \pi^- N \rightarrow \pi^- \pi^+ \pi^- \gamma N$
-------	----	--	--------	----	---

NODE=M008R9
NODE=M008R9

 $\Gamma(e^+e^-)/\Gamma_{total}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<9.4 \times 10^{-9}$	90	¹ ACHASOV	20	SND $e^+e^- \rightarrow \eta\pi^0\pi^0$
--	----	----------------------	----	---

¹ ACHASOV 20 reports two candidate events corresponding to a significance of 2.5σ and the branching fraction of $(5.1^{+3.7}_{-2.7}) \times 10^{-9}$.

NODE=M008R00
NODE=M008R00

OCCUR=2

NODE=M008R00;LINKAGE=B

 $f_1(1285)$ REFERENCES

ABLIKIM	23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHASOV	20	PL B800 135074	M.N. Achasov <i>et al.</i>	(SND Collab.)
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DICKSON	16	PR C93 065202	R. Dickson <i>et al.</i>	(JLab CLAS Collab.)
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAJ	14Y	PRL 112 091802	R. Aaji <i>et al.</i>	(LHCb Collab.)
LEES	12X	PR D86 092010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
DOROFEEV	11	EPJ A47 68	V. Dorofeev <i>et al.</i>	(SERP, MIPT)
PDG	10	JP G37 075021	K. Nakamura <i>et al.</i>	(PDG Collab.)
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACHARD	02B	PL B526 269	P. Achard <i>et al.</i>	(L3 Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)
SOSA	99	PRL 83 913	M. Sosa <i>et al.</i>	
BARBERIS	98C	PL B440 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
BARBERIS	97B	Translated from YAF 60 458	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	97C	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
AMELIN	95	ZPHY C66 71	D.V. Amelin <i>et al.</i>	(VES Collab.)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)
ARMSTRONG	92C	ZPHY C54 371	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)
BITYUKOV	91B	SJNP 54 318	S.I. Bityukov <i>et al.</i>	(SERP)
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
ARMSTRONG	89	PL B221 216	T.A. Armstrong <i>et al.</i>	(CERN, CDEF, BIRM+)
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ARMSTRONG	89G	ZPHY C43 55	T.A. Armstrong <i>et al.</i>	(CERN, BIRM, BARI+)
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
AIHARA	88B	PL B209 107	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD)
BITYUKOV	88	PL B203 327	S.I. Bityukov <i>et al.</i>	(SERP)
MIR	88	Photon-Photon 88, 126	R. Mir	(Mark III Collab.)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
BECKER	87	PRL 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)
GIDAL	87	PRL 59 2012	G. Gidal <i>et al.</i>	(LBL, SLAC, HARV)
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+)
REEVES	86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+)
CHUNG	85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+)
ARMSTRONG	84	PL 146B 793	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BITYUKOV	84B	PL 144B 133	S.I. Bityukov <i>et al.</i>	(SERP)
CHAUVAT	84	PL 148B 382	P. Chauvat <i>et al.</i>	(CERN, CLER, UCLA+)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)
BROMBERG	80	PR D22 1513	C.M. Bromberg <i>et al.</i>	(CIT, FNAL, ILLC+)
DIONISI	80	NP B169 1	C. Dionisi <i>et al.</i>	(CERN, MADR, CDEF+)
GURTU	79	NP B151 181	A. Gurtu <i>et al.</i>	(CERN, ZEEM, NIJM, OXF)
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+)
CORDEN	78	NP B144 253	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
NACASCH	78	NP B135 203	R. Nacasch <i>et al.</i>	(PARIS, MADR, CERN)
GRASSLER	77	NP B121 189	H. Grassler <i>et al.</i>	(AACH3, BERL, BONN+)
DEFOIX	72	NP B44 125	C. Defoix <i>et al.</i>	(CDEF, CERN)
DUBOC	72	NP B46 429	J. Duboc <i>et al.</i>	(PARIS, LIVP)
THUN	72	PRL 28 1733	R. Thun <i>et al.</i>	(STON, NEAS)
BARDADIN...	71	PR D4 2711	M. Bardadin-Otwinowska <i>et al.</i>	(WARSA)
BOESEBECK	71	PL 34B 659	K. Boesebeck	(AACH, BERL, BONN, CERN, CRAC+)
CAMPBELL	69	PRL 22 1204	J.H. Campbell <i>et al.</i>	(PURD)
LORSTAD	69	NP B14 63	B. Lorstad <i>et al.</i>	(CDEF, CERN) JP
D'ANDLAU	68	NP B5 693	C. d'Andlau <i>et al.</i>	(CDEF, CERN, IRAD+)
DAHL	67	PR 163 1377	O.I. Dahl <i>et al.</i>	(LRL) IJP

NODE=M008

REFID=62055
REFID=60144
REFID=60024
REFID=57487
REFID=56781
REFID=55837
REFID=54714
REFID=53931
REFID=16755
REFID=53229
REFID=51698
REFID=52049
REFID=50167
REFID=49548
REFID=48574
REFID=48319
REFID=49649
REFID=47959
REFID=47989
REFID=46937
REFID=46346
REFID=45396

REFID=45758
REFID=45759
REFID=44376
REFID=44437
REFID=44090
REFID=44092
REFID=43587
REFID=42097
REFID=42175
REFID=41864

REFID=41748
REFID=41350
REFID=40729
REFID=41011
REFID=40930
REFID=40924
REFID=40572
REFID=40568
REFID=40569
REFID=41574

REFID=40221
REFID=40015
REFID=40223
REFID=20891
REFID=20936
REFID=20934
REFID=20929
REFID=20468
REFID=20932
REFID=20573
REFID=20462
REFID=20922
REFID=20924
REFID=20456
REFID=20887
REFID=20452
REFID=20919
REFID=20447
REFID=20435
REFID=20339
REFID=20911
REFID=20196
REFID=20905
REFID=20419
REFID=20901
REFID=20897
REFID=20321

$\eta(1295)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M037


NODE=M037

NODE=M037M

NODE=M037M

 $\eta(1295)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1294±4 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
1302±9±8	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
1282±5	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1299±4	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1295±4		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1264±8		¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~ 1275		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$



¹ PWA analysis of AUGUSTIN 92 assigns 0^{-+} quantum numbers to this state rather than 1^{++} as before.

NODE=M037M;LINKAGE=AG

 $\eta(1295)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
55± 5 OUR AVERAGE				
57±23±21	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
66±13	9082	MANAK	00A MPS	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
53± 6		FUKUI	91C SPEC	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<40	2100	ALDE	97B GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
44±20		¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
~ 70		STANTON	79 CNTR	8.4 $\pi^- p \rightarrow n \eta 2\pi$

¹ PWA analysis of AUGUSTIN 92 assigns 0^{-+} quantum numbers to this state rather than 1^{++} as before.

NODE=M037W

NODE=M037W

NODE=M037W;LINKAGE=AG

$\eta(1295)$ DECAY MODES

NODE=M037215;NODE=M037

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi^+\pi^-$	seen
Γ_2 $a_0(980)\pi$	seen
Γ_3 $\gamma\gamma$	
Γ_4 $\eta\pi^0\pi^0$	seen
Γ_5 $\eta(\pi\pi)S\text{-wave}$	seen
Γ_6 $\sigma\eta$	seen
Γ_7 $K\bar{K}\pi$	seen

DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←
DESIG=7;OUR EST;→ UNCHECKED ←

 $\eta(1295)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M037220

 $\Gamma(\eta\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_3/\Gamma$ NODE=M037G2
NODE=M037G2

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.066	95	ACCIARRI	01G L3	183-202 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$ $e^+e^-\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6	90	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
<0.3		ANTREASYAN	87 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi\pi$

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_3/\Gamma$ NODE=M037G3
NODE=M037G3

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.014	90	1,2 AHOHE	05 CLE2	10.6 $e^+e^- \rightarrow e^+e^-K_S^0K^\pm\pi^\mp$ $e^+e^-K_S^0K^\pm\pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using $\eta(1295)$ mass and width 1294 MeV and 55 MeV, respectively.² Assuming three-body phase-space decay to $K_S^0K^\pm\pi^\mp$.NODE=M037G3;LINKAGE=AH
NODE=M037G3;LINKAGE=B3 $\eta(1295)$ BRANCHING RATIOS

NODE=M037225

 $\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M037R1
NODE=M037R1

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	BERTIN	97 OBLX	0.0 $\bar{p}p \rightarrow K^\pm(K^0)\pi^\mp\pi^+\pi^-$
seen	BIRMAN	88 MPS	8 $\pi^-p \rightarrow K^+\bar{K}^0\pi^-n$
large	ANDO	86 SPEC	8 $\pi^-p \rightarrow \eta\pi^+\pi^-n$
large	STANTON	79 CNTR	8.4 $\pi^-p \rightarrow n\eta 2\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi^0\pi^0)$ Γ_2/Γ_4 NODE=M037R2
NODE=M037R2

VALUE	DOCUMENT ID	TECN	COMMENT
0.65 ± 0.10	¹ ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

¹ Assuming that $a_0(980)$ decays only to $\eta\pi$.

NODE=M037R2;LINKAGE=A

 $\Gamma(\eta(\pi\pi)S\text{-wave})/\Gamma(\eta\pi^0\pi^0)$ Γ_5/Γ_4 NODE=M037R4
NODE=M037R4

VALUE	DOCUMENT ID	TECN	COMMENT
0.35 ± 0.10	ALDE	97B GAM4	100 $\pi^-p \rightarrow \eta\pi^0\pi^0n$

 $\Gamma(a_0(980)\pi)/\Gamma(\sigma\eta)$ Γ_2/Γ_6 NODE=M037R5
NODE=M037R5

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.48 ± 0.22	9082	MANAK	00A MPS	18 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

 $\eta(1295)$ REFERENCES

NODE=M037

AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)	REFID=50764
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48319
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=49649
MANAK	00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)	REFID=47989
ALDE	97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=45396
		Translated from YAF 60 458.			
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45417
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
FUKUI	91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41748
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40564
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
ANTREASYAN	87	PR D36 2633	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=40008
ANDO	86	PRL 57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP	REFID=20891
STANTON	79	PRL 42 346	N.R. Stanton <i>et al.</i>	(OSU, CARL, MCGI+) JP	REFID=20887

$\pi(1300)$

$$I^G(J^{PC}) = 1^-(0^-+)$$

NODE=M058

 $\pi(1300)$ MASS

NODE=M058M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M058M

1300±100 OUR ESTIMATE

→ UNCHECKED ←

- • • We do not use the following data for averages, fits, limits, etc. • • •

1128± 26±70		DARGENT	17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1345± 8±10	18k	¹ SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
1200± 40	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1343± 15±24		CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1375± 40		ABELE	01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1275± 15		BERTIN	97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
~ 1114		ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
1190± 30		ZIELINSKI	84	SPEC	$200 \pi^+ Z \rightarrow Z 3\pi$
1240± 30		BELLINI	82	SPEC	$40 \pi^- A \rightarrow A 3\pi$
1273± 50		² AARON	81	RVUE	
1342± 20		BONESINI	81	OMEG	$12 \pi^- p \rightarrow p 3\pi$
~ 1400		DAUM	81B	SPEC	$63,94 \pi^- p$

¹ From analysis of L3 data at 183–209 GeV.² Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.NODE=M058M;LINKAGE=SC
NODE=M058M;LINKAGE=E $\pi(1300)$ WIDTH

NODE=M058W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M058W

200 to 600 OUR ESTIMATE

→ UNCHECKED ←

- • • We do not use the following data for averages, fits, limits, etc. • • •

314± 39±66		DARGENT	17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
260± 20±30	18k	³ SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
470±120	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
449± 39±47		CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
268± 50		ABELE	01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
218±100		BERTIN	97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+ 2\pi^-$
~ 340		ABELE	96	CBAR	$0.0 \bar{p}p \rightarrow 5\pi^0$
440± 80		ZIELINSKI	84	SPEC	$200 \pi^+ Z \rightarrow Z 3\pi$
360±120		BELLINI	82	SPEC	$40 \pi^- A \rightarrow A 3\pi$
580±100		⁴ AARON	81	RVUE	
220± 70		BONESINI	81	OMEG	$12 \pi^- p \rightarrow p 3\pi$
~ 600		DAUM	81B	SPEC	$63,94 \pi^- p$

³ From analysis of L3 data at 183–209 GeV.⁴ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.NODE=M058W;LINKAGE=SC
NODE=M058W;LINKAGE=E $\pi(1300)$ DECAY MODES

NODE=M058215;NODE=M058

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\pi(\pi\pi)S$ -wave	seen
Γ_3 $\gamma\gamma$	

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4 $\pi(1300)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M058218

 $\Gamma(\rho\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_3/\Gamma$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

NODE=M058G1
NODE=M058G1**<0.085** 90 ACCIARRI 97T L3 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$

- • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	95	⁵ SCHEGELSKY	06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$
<0.54	90	ALBRECHT	97B	ARG	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$

⁵ From analysis of L3 data at 183–209 GeV.

NODE=M058G1;LINKAGE=SC

$\pi(1300)$ BRANCHING RATIOS $\Gamma(\pi\pi)_{S\text{-wave}}/\Gamma(\rho\pi)$ Γ_2/Γ_1

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.4		90k	SALVINI	04	OBLX $\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
seen			CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+ 2\pi^- p$
<0.15	90		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
2.12			⁶ AARON	81	RVUE

⁶ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

NODE=M058220

NODE=M058R1
NODE=M058R1

NODE=M058R1;LINKAGE=E

 $\pi(1300)$ REFERENCES

DARGENT	17	JHEP 1705 143	P. dArgent <i>et al.</i>	(HEID, BRIS)
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>	
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ACCIARRI	97T	PL B413 147	M. Acciari <i>et al.</i>	(L3 Collab.)
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
BERTIN	97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ZIELINSKI	84	PR D30 1855	M. Zielinski <i>et al.</i>	(ROCH, MINN, FNAL)
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)
AARON	81	PR D24 1207	R.A. Aaron, R.S. Longacre	(NEAS, BNL)
BONESINI	81	PL 103B 75	M. Bonesini <i>et al.</i>	(MILA, LIVP, DARE+)
DANKOWY...	81	PRL 46 580	J.A. Dankowych <i>et al.</i>	(TNTO, BNL, CARL+)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
DAUM	80	PL 89B 281	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BOWLER	75	NP B97 227	M.G. Bowler <i>et al.</i>	(OXFTP, DARE)

NODE=M058

REFID=58121
REFID=51186
REFID=53226
REFID=48837
REFID=48334
REFID=45761
REFID=45418
REFID=45763
REFID=45011
REFID=20881
REFID=21134
REFID=20870
REFID=21130
REFID=20572
REFID=20872
REFID=20868
REFID=20571

NODE=M012

 $a_2(1320)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

 $a_2(1320)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1305–1321)–i(52–58) OUR ESTIMATE			
$(1318.7 \pm 1.9 \pm 1.3) - i(53.8 \pm 2.3^{+1.7}_{-0.9})$	¹ KOPF	21	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
$(1312.5 \pm 0.7 \pm 2.6) - i(53.5 \pm 0.6 \pm 1.9)$	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1306.0 \pm 0.8 \pm 1.3) - i(57.2 \pm 0.8 \pm 0.0)$	³ RODAS	19	RVUE $91 \pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
$(1309 \pm 4) - i(55 \pm 2)$	⁴ ANISOVICH	09	RVUE $\bar{p}p, \pi N$

¹ Extraction based on a combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

² T-matrix pole with 2 poles, 2 channels ($\pi^0\eta$ and $K\bar{K}$).

³ Coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data. Supersedes JACKURA 18. Performed by JPAC.

⁴ Amplitude did not include dispersive corrections. From analysis of $\eta\pi$ mode.

NODE=M012PP

NODE=M012PP

NODE=M012PP

→ UNCHECKED ←

NODE=M012PP;LINKAGE=B

NODE=M012PP;LINKAGE=C

NODE=M012PP;LINKAGE=D

NODE=M012PP;LINKAGE=A

 $a_2(1320)$ MASS

VALUE (MeV)	DOCUMENT ID
1318.2 ± 0.6 OUR AVERAGE	Includes data from the 4 datablocks that follow this one. Error includes scale factor of 1.2.

NODE=M012M0

NODE=M012M0

3π MODE

NODE=M012M1
NODE=M012M1

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
The data in this block is included in the average printed for a previous datablock.

1318.6 ± 1.3 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

1314.5 ^{+4.0} _{-3.3}	46M	¹ AGHASYAN	18B	COMP	190 π ⁻ p → π ⁻ π ⁺ π ⁻ p	
1326 ± 2 ± 2		CHUNG	02	B852	18.3 π ⁻ p → π ⁺ π ⁻ π ⁻ p	
1317 ± 3		BARBERIS	98B		450 p p → ρ _f π ⁺ π ⁻ π ⁰ p _s	
1323 ± 4 ± 3		ACCIARRI	97T	L3	e ⁺ e ⁻ → e ⁺ e ⁻ π ⁺ π ⁻ π ⁰	
1320 ± 7		ALBRECHT	97B	ARG	e ⁺ e ⁻ → e ⁺ e ⁻ π ⁺ π ⁻ π ⁰	
1311.3 ± 1.6 ± 3.0	72.4k	AMELIN	96	VES	36 π ⁻ p → π ⁺ π ⁻ π ⁰ n	
1310 ± 5		ARMSTRONG	90	OMEG 0	300.0 p p → p p π ⁺ π ⁻ π ⁰	
1323.8 ± 2.3	4022	AUGUSTIN	89	DM2 ±	J/ψ → ρ [±] a ₂ [∓]	
1320.6 ± 3.1	3562	AUGUSTIN	89	DM2 0	J/ψ → ρ ⁰ a ₂ ⁰	OCCUR=2
1317 ± 2	25k	² DAUM	80C	SPEC -	63,94 π ⁻ p → 3π p	
1320 ± 10	1097	² BALTAY	78B	HBC +0	15 π ⁺ p → p 4π	
1306 ± 8		FERRERSORIA	78	OMEG -	9 π ⁻ p → p 3π	
1318 ± 7	1.6k	² EMMS	75	DBC 0	4 π ⁺ n → ρ(3π) ⁰	
1315 ± 5		² ANTIPOV	73C	CNTR -	25,40 π ⁻ p → ρ η π ⁻	
1306 ± 9	1580	CHALOUKKA	73	HBC -	3.9 π ⁻ p	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1321 ± 1 ⁺⁰ ₋₇	420k	³ ALEKSEEV	10	COMP	190 π ⁻ P b → π ⁻ π ⁻ π ⁺ P b'	
1300 ± 2 ± 4	18k	⁴ SCHEGELSKY	06	RVUE 0	γγ → π ⁺ π ⁻ π ⁰	
1305 ± 14		CONDO	93	SHF	γγ → n π ⁺ π ⁺ π ⁻	
1310 ± 2		² EVANGELIS...	81	OMEG -	12 π ⁻ p → 3π p	
1343 ± 11	490	BALTAY	78B	HBC 0	15 π ⁺ p → Δ 3π	OCCUR=2
1309 ± 5	5k	BINNIE	71	MMS -	π ⁻ p near a ₂ thresh- old	OCCUR=2
1299 ± 6	28k	BOWEN	71	MMS -	5 π ⁻ p	
1300 ± 6	24k	BOWEN	71	MMS +	5 π ⁺ p	OCCUR=2
1309 ± 4	17k	BOWEN	71	MMS -	7 π ⁻ p	OCCUR=3
1306 ± 4	941	ALSTON...	70	HBC +	7.0 π ⁺ p → 3π p	


¹ Statistical error negligible.

² From a fit to J^P = 2⁺ ρ π partial wave.

³ Superseded by AGHASYAN 2018B.

⁴ From analysis of L3 data at 183–209 GeV.

NODE=M012M1;LINKAGE=D
NODE=M012M1;LINKAGE=P
NODE=M012M1;LINKAGE=C
NODE=M012M1;LINKAGE=SC



$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M2
NODE=M012M2**1318.1± 0.7 OUR AVERAGE**

1319 ± 5	4700	^{1,2} CLELAND	82B	SPEC	+	50 $\pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
1324 ± 6	5200	^{1,2} CLELAND	82B	SPEC	-	50 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
1320 ± 2	4000	CHABAUD	80	SPEC	-	17 $\pi^- A \rightarrow K_S^0 K^- A$	
1312 ± 4	11000	CHABAUD	78	SPEC	-	9.8 $\pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
1316 ± 2	4730	CHABAUD	78	SPEC	-	18.8 $\pi^- p \rightarrow K^- K_S^0 p$	
1318 ± 1		^{1,3} MARTIN	78D	SPEC	-	10 $\pi^- p \rightarrow K_S^0 K^- p$	
1320 ± 2	2724	MARGULIE	76	SPEC	-	23 $\pi^- p \rightarrow K^- K_S^0 p$	
1313 ± 4	730	FOLEY	72	CNTR	-	20.3 $\pi^- p \rightarrow K^- K_S^0 p$	
1319 ± 3	1500	³ GRAYER	71	ASPK	-	17.2 $\pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1304 ± 10	870	⁴ SCHEGELSKY	06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
1330 ± 11	1000	^{1,2} CLELAND	82B	SPEC	+	30 $\pi^+ p \rightarrow K_S^0 K^+ p$	
1324 ± 5	350	HYAMS	78	ASPK	+	12.7 $\pi^+ p \rightarrow K^+ K_S^0 p$	

¹ From a fit to $J^P = 2^+$ partial wave.² Number of events evaluated by us.³ Systematic error in mass scale subtracted.⁴ From analysis of L3 data at 91 and 183–209 GeV.NODE=M012M2;LINKAGE=P
NODE=M012M2;LINKAGE=W
NODE=M012M2;LINKAGE=S
NODE=M012M2;LINKAGE=SC **$\eta\pi$ MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M3
NODE=M012M3**1317.7± 1.4 OUR AVERAGE**

1308 ± 9		BARBERIS	00H			450 $p p \rightarrow p_f \eta \pi^0 p_s$	
1316 ± 9		BARBERIS	00H			450 $p p \rightarrow \Delta_f^{++} \eta \pi^- p_s$	OCCUR=2
1317 ± 1 ± 2		THOMPSON	97	MPS		18 $\pi^- p \rightarrow \eta \pi^- p$	
1315 ± 5 ± 2		¹ AMSLER	94D	CBAR		0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta$	
1325.1 ± 5.1		AOYAGI	93	BKEI		$\pi^- p \rightarrow \eta \pi^- p$	
1317.7 ± 1.4 ± 2.0		BELADIDZE	93	VES		37 $\pi^- N \rightarrow \eta \pi^- N$	
1323 ± 8	1000	² KEY	73	OSPK	-	6 $\pi^- p \rightarrow p \pi^- \eta$	
1307 ± 1 ± 6		³ JACKURA	18	RVUE		$\pi^- p \rightarrow \eta \pi^- p$	
1315 ± 12		⁴ ADOLPH	15	COMP		191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$	
1324 ± 5		ARMSTRONG	93C	E760	0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
1336.2 ± 1.7	2561	DELFOSE	81	SPEC	+	$\pi^\pm p \rightarrow p \pi^\pm \eta$	OCCUR=2
1330.7 ± 2.4	1653	DELFOSE	81	SPEC	-	$\pi^\pm p \rightarrow p \pi^\pm \eta$	
1324 ± 8	6200	^{2,5} CONFORTO	73	OSPK	-	6 $\pi^- p \rightarrow p \pi^- \eta$	

¹ The systematic error of 2 MeV corresponds to the spread of solutions.² Error includes 5 MeV systematic mass-scale error.³ Superseded by RODAS 19.⁴ ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the $\eta\pi$ and $\rho\pi$ channels into account.⁵ Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.NODE=M012M3;LINKAGE=DD
NODE=M012M3;LINKAGE=E
NODE=M012M3;LINKAGE=B
NODE=M012M3;LINKAGE=A
NODE=M012M3;LINKAGE=M **$\eta'\pi$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012M4
NODE=M012M4**1322 ± 7 OUR AVERAGE**

1318 ± 8 $^{+3}_{-5}$		IVANOV	01	B852		18 $\pi^- p \rightarrow \eta' \pi^- p$	
1327.0 ± 10.7		BELADIDZE	93	VES		37 $\pi^- N \rightarrow \eta' \pi^- N$	

$a_2(1320)$ WIDTH

NODE=M012210

3 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

 NODE=M012W1
 NODE=M012W1
105.0 $^{+1.7}_{-1.9}$ OUR AVERAGE

106.6 $^{+3.4}_{-7.0}$	46M	¹ AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$	
108 $\pm 3 \pm 15$		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
120 ± 10		BARBERIS	98B		450 $p p \rightarrow p_f \pi^+ \pi^- \pi^0 p_s$	
105 $\pm 10 \pm 11$		ACCIARRI	97T	L3	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
120 ± 10		ALBRECHT	97B	ARG	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$	
103.0 $\pm 6.0 \pm 3.3$	72.4k	AMELIN	96	VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$	
120 ± 10		ARMSTRONG	90	OMEG 0	300.0 $p p \rightarrow p p \pi^+ \pi^- \pi^0$	
107.0 ± 9.7	4022	AUGUSTIN	89	DM2 \pm	$J/\psi \rightarrow \rho^\pm a_2^\mp$	
118.5 ± 12.5	3562	AUGUSTIN	89	DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$	OCCUR=2
97 ± 5		² EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow 3\pi p$	
96 ± 9	25k	² DAUM	80C	SPEC -	63,94 $\pi^- p \rightarrow 3\pi p$	
110 ± 15	1097	² BALTAY	78B	HBC +0	15 $\pi^+ p \rightarrow p 4\pi$	
112 ± 18	1.6k	² EMMS	75	DBC 0	4 $\pi^+ n \rightarrow p(3\pi)^0$	
122 ± 14	1.2k	^{2,3} WAGNER	75	HBC 0	7 $\pi^+ p \rightarrow \Delta^{++}(3\pi)^0$	
115 ± 15		² ANTIPOV	73C	CNTR -	25,40 $\pi^- p \rightarrow p \eta \pi^-$	
99 ± 15	1580	CHALOUKPA	73	HBC -	3.9 $\pi^- p$	
105 ± 5	28k	BOWEN	71	MMS -	5 $\pi^- p$	
99 ± 5	24k	BOWEN	71	MMS +	5 $\pi^+ p$	OCCUR=2
103 ± 5	17k	BOWEN	71	MMS -	7 $\pi^- p$	OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

110 $\pm 2 \begin{smallmatrix} +2 \\ -15 \end{smallmatrix}$	420k	⁴ ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
117 $\pm 6 \pm 20$	18k	⁵ SCHEGELSKY	06	RVUE 0	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$	
120 ± 40		CONDO	93	SHF	$\gamma p \rightarrow n \pi^+ \pi^+ \pi^-$	
115 ± 14	490	BALTAY	78B	HBC 0	15 $\pi^+ p \rightarrow \Delta 3\pi$	OCCUR=2
72 ± 16	5k	BINNIE	71	MMS -	$\pi^- p$ near a_2 thresh- old	OCCUR=2
79 ± 12	941	ALSTON-...	70	HBC +	7.0 $\pi^+ p \rightarrow 3\pi p$	

¹ Statistical error negligible.

² From a fit to $J^P = 2^+ \rho\pi$ partial wave.

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁴ Superseded by AGHASYAN 2018B.

⁵ From analysis of L3 data at 183–209 GeV.

 NODE=M012W1;LINKAGE=C
 NODE=M012W1;LINKAGE=P
 NODE=M012W1;LINKAGE=S
 NODE=M012W1;LINKAGE=E
 NODE=M012W1;LINKAGE=SC
 $K\bar{K}$ AND $\eta\pi$ MODES

VALUE (MeV)	DOCUMENT ID
-------------	-------------

 NODE=M012W0
 NODE=M012W0
107 ± 5 OUR ESTIMATE

→ UNCHECKED ←

110.4 ± 1.7 OUR AVERAGE Includes data from the 2 datablocks that follow this one. **$K\bar{K}$ MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

 NODE=M012W2
 NODE=M012W2

The data in this block is included in the average printed for a previous datablock.

109.8 ± 2.4 OUR AVERAGE

112 ± 20	4700	^{1,2} CLELAND	82B	SPEC +	50 $\pi^+ p \rightarrow K_S^0 K^+ p$	OCCUR=2
120 ± 25	5200	^{1,2} CLELAND	82B	SPEC -	50 $\pi^- p \rightarrow K_S^0 K^- p$	OCCUR=3
106 ± 4	4000	CHABAUD	80	SPEC -	17 $\pi^- A \rightarrow K_S^0 K^- A$	
126 ± 11	11000	CHABAUD	78	SPEC -	9.8 $\pi^- p \rightarrow K^- K_S^0 p$	
101 ± 8	4730	CHABAUD	78	SPEC -	18.8 $\pi^- p \rightarrow K^- K_S^0 p$	OCCUR=2
113 ± 4		^{1,3} MARTIN	78D	SPEC -	10 $\pi^- p \rightarrow K_S^0 K^- p$	
105 ± 8	2724	³ MARGULIE	76	SPEC -	23 $\pi^- p \rightarrow K^- K_S^0 p$	
113 ± 19	730	FOLEY	72	CNTR -	20.3 $\pi^- p \rightarrow K^- K_S^0 p$	
123 ± 13	1500	³ GRAYER	71	ASPK -	17.2 $\pi^- p \rightarrow K^- K_S^0 p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

120	± 15	870	⁴ SCHEGELSKY	06A	RVUE	0	$\gamma\gamma \rightarrow K_S^0 K_S^0$
121	± 51	1000	^{1,2} CLELAND	82B	SPEC	+	$30 \pi^+ p \rightarrow K_S^0 K^+ p$
110	± 18	350	HYAMS	78	ASPK	+	$12.7 \pi^+ p \rightarrow K^+ K_S^0 p$

¹ From a fit to $J^P = 2^+$ partial wave.

² Number of events evaluated by us.

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁴ From analysis of L3 data at 91 and 183–209 GeV.

NODE=M012W2;LINKAGE=P
 NODE=M012W2;LINKAGE=W
 NODE=M012W2;LINKAGE=S
 NODE=M012W2;LINKAGE=SC

$\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M012W3
 NODE=M012W3

111.1 \pm 2.4 OUR AVERAGE

115	± 20		BARBERIS	00H			$450 p p \rightarrow p_f \eta \pi^0 p_s$
112	± 14		BARBERIS	00H			$450 p p \rightarrow \Delta_f^{++} \eta \pi^- p_s$
112	$\pm 3 \pm 2$		¹ AMSLER	94D	CBAR		$0.0 \bar{p} p \rightarrow \pi^0 \pi^0 \eta$
103	$\pm 6 \pm 3$		BELADIDZE	93	VES		$37 \pi^- N \rightarrow \eta \pi^- N$
112.2 \pm 5.7	2561		DELFOSSÉ	81	SPEC	+	$\pi^\pm p \rightarrow p \pi^\pm \eta$
116.6 \pm 7.7	1653		DELFOSSÉ	81	SPEC	-	$\pi^\pm p \rightarrow p \pi^\pm \eta$
108	± 9	1000	KEY	73	OSPK	-	$6 \pi^- p \rightarrow p \pi^- \eta$

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

112	$\pm 1 \pm 8$		² JACKURA	18	RVUE		$\pi^- p \rightarrow \eta \pi^- p$
119	± 14		³ ADOLPH	15	COMP		$191 \pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
127	$\pm 2 \pm 2$		⁴ THOMPSON	97	MPS		$18 \pi^- p \rightarrow \eta \pi^- p$
118	± 10		ARMSTRONG	93C	E760	0	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
104	± 9	6200	⁵ CONFORTO	73	OSPK	-	$6 \pi^- p \rightarrow p \text{M}^-$

¹ The systematic error of 2 MeV corresponds to the spread of solutions.

² Superseded by RODAS 19.

³ ADOLPH 15 value is derived from a Breit-Wigner fit with mass-dependent width taking the $\eta\pi$ and $\rho\pi$ channels into account.

⁴ Resolution is not unfolded.

⁵ Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.

NODE=M012W3;LINKAGE=DD
 NODE=M012W3;LINKAGE=C
 NODE=M012W3;LINKAGE=B

NODE=M012W3;LINKAGE=A
 NODE=M012W3;LINKAGE=M

$\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

119 \pm 25 OUR AVERAGE

140 \pm 35 \pm 20	IVANOV	01	B852	$18 \pi^- p \rightarrow \eta' \pi^- p$
106 \pm 32	BELADIDZE	93	VES	$37 \pi^- N \rightarrow \eta' \pi^- N$

NODE=M012W4
 NODE=M012W4

$a_2(1320)$ DECAY MODES

NODE=M012215;NODE=M012

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 3π	(70.1 \pm 2.7) %	S=1.2
Γ_2 $\rho(770)\pi$		
Γ_3 $f_2(1270)\pi$		
Γ_4 $\rho(1450)\pi$		
Γ_5 $\eta\pi$	(14.5 \pm 1.2) %	
Γ_6 $\omega\pi\pi$	(10.6 \pm 3.2) %	S=1.3
Γ_7 $K\bar{K}$	(4.9 \pm 0.8) %	
Γ_8 $\eta'(958)\pi$	(5.5 \pm 0.9) $\times 10^{-3}$	
Γ_9 $\pi^\pm\gamma$	(2.91 \pm 0.27) $\times 10^{-3}$	
Γ_{10} $\gamma\gamma$	(9.4 \pm 0.7) $\times 10^{-6}$	
Γ_{11} e^+e^-	< 5 $\times 10^{-9}$	CL=90%

DESIG=1
 DESIG=11
 DESIG=12
 DESIG=13
 DESIG=3
 DESIG=4
 DESIG=2
 DESIG=8
 DESIG=7
 DESIG=9
 DESIG=10

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 18 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 9.3$ for 15 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	10		
x_6	-89	-46	
x_7	-1	-2	-24
	x_1	x_5	x_6

$a_2(1320)$ PARTIAL WIDTHS

$\Gamma(\eta\pi)$

Γ_5

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.5 ± 3.0 870 ¹ SCHEGELSKY 06A RVUE 0 $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$ keV and SU(3) relations.

NODE=M012220

NODE=M012W6
NODE=M012W6

NODE=M012W6;LINKAGE=SC

$\Gamma(K\bar{K})$

Γ_7

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0^{+2.0}_{-1.5} 870 ¹ SCHEGELSKY 06A RVUE 0 $\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(a_2(1320) \rightarrow \gamma\gamma) = 0.91$ keV and SU(3) relations.

NODE=M012W5
NODE=M012W5

NODE=M012W5;LINKAGE=SC

$\Gamma(\pi^\pm\gamma)$

Γ_9

VALUE (keV) EVTS DOCUMENT ID TECN CHG COMMENT

311 ± 25 OUR AVERAGE

358 ± 6 ± 42 ¹ ADOLPH 14 COMP - 190 $\pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$

284 ± 25 ± 25 7.1k MOLCHANOV 01 SELX 600 $\pi^- \text{A} \rightarrow \pi^+ \pi^- \pi^- \text{A}$

295 ± 60 CIHANGIR 82 SPEC + 200 $\pi^+ \text{A}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

461 ± 110 ² MAY 77 SPEC ± 9.7 γA

¹ Primakoff reaction using $a_2(1320) \rightarrow 3\pi$ branching ratio of 70.1%.

² Assuming one-pion exchange.

NODE=M012W7
NODE=M012W7

NODE=M012W7;LINKAGE=AD
NODE=M012W;LINKAGE=M2

$\Gamma(\gamma\gamma)$

Γ_{10}

VALUE (keV) EVTS DOCUMENT ID TECN CHG COMMENT

1.00 ± 0.06 OUR AVERAGE

0.98 ± 0.05 ± 0.09 ACCIARRI 97T L3 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

0.96 ± 0.03 ± 0.13 ALBRECHT 97B ARG $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

1.26 ± 0.26 ± 0.18 36 BARU 90 MD1 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

1.00 ± 0.07 ± 0.15 415 BEHREND 90C CELL 0 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

1.03 ± 0.13 ± 0.21 BUTLER 90 MRK2 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \pi^0$

1.01 ± 0.14 ± 0.22 85 OEST 90 JADE $e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$

0.90 ± 0.27 ± 0.15 56 ¹ ALTHOFF 86 TASS 0 $e^+ e^- \rightarrow e^+ e^- 3\pi$

1.14 ± 0.20 ± 0.26 ² ANTREASYAN 86 CBAL 0 $e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$

1.06 ± 0.18 ± 0.19 BERGER 84C PLUT 0 $e^+ e^- \rightarrow e^+ e^- 3\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.81 ± 0.19^{+0.42}_{-0.11} 35 ¹ BEHREND 82C CELL 0 $e^+ e^- \rightarrow e^+ e^- 3\pi$

0.77 ± 0.18 ± 0.27 22 ² EDWARDS 82F CBAL 0 $e^+ e^- \rightarrow e^+ e^- \pi^0 \eta$

¹ From $\rho\pi$ decay mode.

² From $\eta\pi^0$ decay mode.

NODE=M012W9
NODE=M012W9

NODE=M012W;LINKAGE=F
NODE=M012W;LINKAGE=G

$\Gamma(e^+e^-)$ Γ_{11}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.56	90	ACHASOV	00K SND	$e^+e^- \rightarrow \pi^0\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
<25	90	VOROBYEV	88 ND	$e^+e^- \rightarrow \pi^0\eta$

NODE=M012W10
 NODE=M012W10

 $a_2(1320) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M012223

 $\Gamma(3\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{10}/\Gamma$

VALUE (keV)	EVTs	DOCUMENT ID	TECN	COMMENT
0.65±0.02±0.02	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
¹ From analysis of L3 data at 183–209 GeV.				

NODE=M012G2
 NODE=M012G2

NODE=M012G2;LINKAGE=SC

 $\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_{10}/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.145 ^{+0.097} _{-0.034}	¹ UEHARA 09A	BELL	$e^+e^- \rightarrow e^+e^-\eta\pi^0$
¹ From the D_2 -wave. The fraction of the D_0 -wave is 3.4 ^{+2.3} _{-1.1} %.			

NODE=M012G01
 NODE=M012G01

NODE=M012G01;LINKAGE=UE

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_{10}/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.126±0.007±0.028	¹ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.081±0.006±0.027	² ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
¹ Using an incoherent background.			
² Using a coherent background.			

NODE=M012G1
 NODE=M012G1

OCCUR=2

NODE=M012G1;LINKAGE=A
 NODE=M012G1;LINKAGE=B

 $a_2(1320)$ BRANCHING RATIOS

NODE=M012225

 $[\Gamma(f_2(1270)\pi) + \Gamma(\rho(1450)\pi)]/\Gamma(\rho(770)\pi)$ $(\Gamma_3+\Gamma_4)/\Gamma_2$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.12	90	ABRAMOVI...	70B HBC	-	3.93 π^-p

NODE=M012R9
 NODE=M012R9

 $\Gamma(\rho(770)\pi)/\Gamma(f_2(1270)\pi)$ Γ_2/Γ_3

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
16.5 ^{+1.2} _{-2.4}	46M	¹ AGHASYAN 18B	COMP	190 $\pi^-p \rightarrow \pi^-\pi^+\pi^-p$

NODE=M012R00
 NODE=M012R00

¹ Statistical error negligible.

NODE=M012R00;LINKAGE=A

 $\Gamma(\eta\pi)/\Gamma(3\pi)$ Γ_5/Γ_1

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
0.207±0.018		OUR FIT			
0.213±0.020		OUR AVERAGE			
0.18 ±0.05		FORINO 76	HBC		11 π^-p
0.22 ±0.05	52	ANTIPOV 73	CNTR	-	40 π^-p
0.211±0.044	149	CHALOUKKA 73	HBC	-	3.9 π^-p
0.246±0.042	167	ALSTON-... 71	HBC	+	7.0 π^+p
0.25 ±0.09	15	BOECKMANN 70	HBC	+	5.0 π^+p
0.23 ±0.08	22	ASCOLI 68	HBC	-	5 π^-p
0.12 ±0.08		CHUNG 68	HBC	-	3.2 π^-p
0.22 ±0.09		CONTE 67	HBC	-	11.0 π^-p

NODE=M012R3
 NODE=M012R3

 $\Gamma(\omega\pi\pi)/\Gamma(3\pi)$ Γ_6/Γ_1

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
0.15±0.05		OUR FIT Error includes scale factor of 1.3.			
0.15±0.05		OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.			
0.28±0.09	60	DIAZ 74	DBC	0	6 π^+n
0.18±0.08		¹ KARSHON 74	HBC		Avg. of above two
0.10±0.05	279	² CHALOUKKA 73	HBC	-	3.9 π^-p
••• We do not use the following data for averages, fits, limits, etc. •••					
0.29±0.08	140	¹ KARSHON 74	HBC	0	4.9 π^+p
0.10±0.04	60	¹ KARSHON 74	HBC	+	4.9 π^+p
0.19±0.08		DEFOIX 73	HBC	0	0.7 $\bar{p}p$

NODE=M012R12
 NODE=M012R12

OCCUR=3


OCCUR=2

¹ KARSHON 74 suggest an additional $I = 0$ state strongly coupled to $\omega\pi\pi$ which could explain discrepancies in branching ratios and masses. We use a central value and a systematic spread.

² Decays to $b_1(1040)\pi$, $b_1 \rightarrow \omega\pi$. Error increased to account for possible systematic errors of complicated analysis.

NODE=M012R12;LINKAGE=K

NODE=M012R12;LINKAGE=01



$\Gamma(K\bar{K})/\Gamma(3\pi)$

VALUE EVTS DOCUMENT ID TECN CHG COMMENT Γ_7/Γ_1

0.070±0.012 OUR FIT

0.078±0.017

CHABAUD 78 RVUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.011±0.003		¹ BERTIN	98B	OBLX	0.0	$\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
0.056±0.014	50	² CHALOUPKA	73	HBC	-	$3.9 \pi^- p$
0.097±0.018	113	² ALSTON-...	71	HBC	+	$7.0 \pi^+ p$
0.06 ±0.03		² ABRAMOVI...	70B	HBC	-	$3.93 \pi^- p$
0.054±0.022		² CHUNG	68	HBC	-	$3.2 \pi^- p$

¹ Using 4π data from BERTIN 97D.

² Included in CHABAUD 78 review.

NODE=M012R1
NODE=M012R1

NODE=M012R1;LINKAGE=BE
NODE=M012R1;LINKAGE=C

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$

VALUE DOCUMENT ID TECN COMMENT Γ_7/Γ_5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ±0.22 ^{+0.09} / _{-0.11}	¹ KOPF	21	RVUE	0.9	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and $191 \pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
0.352±0.011±0.175	² ALBRECHT	20	RVUE	0.9	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
0.08 ±0.02	³ BERTIN	98B	OBLX	0.0	$\bar{p}p \rightarrow K^\pm K_S \pi^\mp$

¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

² Residues from T-matrix pole with 2 poles, 2 channels ($\pi^0\eta$ and $K\bar{K}$).

³ Using $\eta\pi\pi$ data from AMSLER 94D.

NODE=M012R14
NODE=M012R14

NODE=M012R14;LINKAGE=B

NODE=M012R14;LINKAGE=A
NODE=M012R14;LINKAGE=BE

$\Gamma(\eta\pi)/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$

VALUE EVTS DOCUMENT ID TECN CHG COMMENT $\Gamma_5/(\Gamma_1+\Gamma_5+\Gamma_7)$

0.162±0.012 OUR FIT

0.140±0.028 OUR AVERAGE

0.13 ±0.04		ESPIGAT	72	HBC	±	$0.0 \bar{p}p$
0.15 ±0.04	34	BARNHAM	71	HBC	+	$3.7 \pi^+ p$

NODE=M012R2
NODE=M012R2

$\Gamma(K\bar{K})/[\Gamma(3\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$ $\Gamma_7/(\Gamma_1+\Gamma_5+\Gamma_7)$

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

NODE=M012R8
NODE=M012R8

0.054±0.009 OUR FIT

0.048±0.012 OUR AVERAGE

0.05 ±0.02		TOET	73	HBC	+	5 $\pi^+ p$
0.09 ±0.04		TOET	73	HBC	0	5 $\pi^+ p$
0.03 ±0.02	8	¹ DAMERI	72	HBC	-	11 $\pi^- p$
0.06 ±0.03	17	BARNHAM	71	HBC	+	3.7 $\pi^+ p$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.020±0.004		² ESPIGAT	72	HBC	±	0.0 $\bar{p} p$
-------------	--	----------------------	----	-----	---	-----------------

¹ Montanet agrees. Vlada.² Not averaged because of discrepancy between masses from $K\bar{K}$ and $\rho\pi$ modes.

NODE=M012R8;LINKAGE=01
NODE=M012R8;LINKAGE=A

 $\Gamma(\eta'(958)\pi)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

NODE=M012R4
NODE=M012R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.006	95	ALDE	92B	GAM2		38,100 $\pi^- p \rightarrow \eta' \pi^0 n$
--------	----	------	-----	------	--	--

<0.02	97	BARNHAM	71	HBC	+	3.7 $\pi^+ p$
-------	----	---------	----	-----	---	---------------

0.004±0.004		¹ BOESEBECK	68	HBC	+	8 $\pi^+ p$
-------------	--	------------------------	----	-----	---	-------------

¹ No longer valid since $\Gamma(K\bar{K})/\Gamma(3\pi)$ value has changed (MORRISON 71).

NODE=M012R4;LINKAGE=B

 $\Gamma(\eta'(958)\pi)/\Gamma(3\pi)$ Γ_8/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

NODE=M012R5
NODE=M012R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011	90	EISENSTEIN	73	HBC	-	5 $\pi^- p$
--------	----	------------	----	-----	---	-------------

<0.04		ALSTON-...	71	HBC	+	7.0 $\pi^+ p$
-------	--	------------	----	-----	---	---------------

0.04 $\begin{smallmatrix} +0.03 \\ -0.04 \end{smallmatrix}$		BOECKMANN	70	HBC	0	5.0 $\pi^+ p$
---	--	-----------	----	-----	---	---------------

 $\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$ Γ_8/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M012R13
NODE=M012R13

0.038±0.005 OUR AVERAGE

0.05 ±0.02	ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
------------	--------	----	------	---

0.032±0.009	ABELE	97C	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \pi^0 \eta'$
-------------	-------	-----	------	---

0.047±0.010±0.004	¹ BELADIDZE	93	VES	37 $\pi^- N \rightarrow a_2^- N$
-------------------	------------------------	----	-----	----------------------------------

0.034±0.008±0.005	BELADIDZE	92	VES	36 $\pi^- C \rightarrow a_2^- C$
-------------------	-----------	----	-----	----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.046±0.015 $\begin{smallmatrix} +0.07 \\ -0.006 \end{smallmatrix}$	² KOPF	21	RVUE	0.9 $p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
---	-------------------	----	------	---

OCCUR=2

¹ Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = 0.441$, $B(\eta \rightarrow \gamma \gamma) = 0.389$ and $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 0.236$.

NODE=M012R13;LINKAGE=A

² From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi$, $\eta'\pi$ and $K\bar{K}$ systems.

NODE=M012R13;LINKAGE=C

 $\Gamma(\pi^\pm \gamma)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M012R11
NODE=M012R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.005 $\begin{smallmatrix} +0.005 \\ -0.003 \end{smallmatrix}$	¹ EISENBERG	72	HBC	4.3, 5.25, 7.5 γp
--	------------------------	----	-----	---------------------------

¹ Pion-exchange model used in this estimation.

NODE=M012R11;LINKAGE=R

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M012R15
NODE=M012R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90	ACHASOV	00K	SND $e^+ e^- \rightarrow \pi^0 \pi^0$
----	----	---------	-----	---------------------------------------

a₂(1320) REFERENCES

NODE=M012

KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)	REFID=61470
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=59554
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)	REFID=59003
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=55911
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
MOLCHANOV	01	PL B521 171	V.V. Molchanov <i>et al.</i>	(FNAL SELEX Collab.)	REFID=48559
ACHASOV	00K	PL B492 8	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47933
BARBERIS	00H	PL B488 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47964
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
ABELE	97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45531
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=45418
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=45584
AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44649
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BELADIDZE	93	PL B313 276	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43598
CONDO	93	PR D48 3045	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=43600
ALDE	92B	ZPHY C54 549	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41852
BELADIDZE	92	ZPHY C54 235	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42171
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	90	ZPHY C48 213	T.A. Armstrong, M. Benayoun, W. Beusch	(WA76 Coll.)	REFID=41375
BARU	90	ZPHY C48 581	S.E. Baru <i>et al.</i>	(MD-1 Collab.)	REFID=41366
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41356
BUTLER	90	PR D42 1368	F. Butler <i>et al.</i>	(Mark II Collab.)	REFID=41363
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)	REFID=41023
ALTHOFF	86	ZPHY C31 537	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21287
ANTREASYAN	86	PR D33 1847	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=20469
BERGER	84C	PL 149B 427	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=21286
BEHREND	82C	PL 114B 378	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20303
Also		PL 125B 518 (errat.)	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=20302
CHANGIR	82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
EDWARDS	82F	PL 110B 82	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=20747
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
CHABAUD	80	NP B175 189	V. Chabaud <i>et al.</i>	(CERN, MPIM, AMST)	REFID=21274
DAUM	80C	PL 89B 276	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21275
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
CHABAUD	78	NP B145 349	V. Chabaud <i>et al.</i>	(CERN, MPIM)	REFID=21267
FERRERSORIA	78	PL 74B 287	A. Ferrer Soria <i>et al.</i>	(ORSAY, CERN, CDEF+)	REFID=21270
HYAMS	78	NP B146 303	B.D. Hyams <i>et al.</i>	(CERN, MPIM, ATEN)	REFID=21271
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA) JP	REFID=21272
MAY	77	PR D16 1983	E.N. May <i>et al.</i>	(ROCH, CORN)	REFID=20450
FORINO	76	NC 35A 465	A. Forino <i>et al.</i>	(BGNA, FIRZ, GENO, MILA+)	REFID=21259
MARGULIE	76	PR D14 667	M. Margulies <i>et al.</i>	(BNL, CUNY)	REFID=21261
EMMS	75	PL 58B 117	M.J. Emms <i>et al.</i>	(BIRM, DURH, RHEL) JP	REFID=21254
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP	REFID=20843
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)	REFID=21248
KARSHON	74	PRL 32 852	U. Karshon <i>et al.</i>	(REHO)	REFID=21249
ANTIPOV	73	NP B63 175	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=21238
ANTIPOV	73C	NP B63 153	Y.M. Antipov <i>et al.</i>	(CERN, SERP) JP	REFID=20817
CHALOUPKA	73	PL 44B 211	V. Chaloupka <i>et al.</i>	(CERN)	REFID=21242
CONFORTO	73	PL 45B 154	G. Conforto <i>et al.</i>	(EFI, FNAL, TNTO+)	REFID=21243
DEFOIX	73	PL 43B 141	C. Defoix <i>et al.</i>	(CDEF)	REFID=21244
EISENSTEIN	73	PR D7 278	L. Eisenstein <i>et al.</i>	(ILL)	REFID=21245
KEY	73	PRL 30 503	A.W. Key <i>et al.</i>	(TNTO, EFI, FNAL, WISC)	REFID=21246
TOET	73	NP B63 248	D.Z. Toet <i>et al.</i>	(NIJM, BONN, DURH, TORI)	REFID=20714
DAMERI	72	NC 9A 1	M. Dameri <i>et al.</i>	(GENO, MILA, SACL)	REFID=20338
EISENBERG	72	PR D5 15	Y. Eisenberg <i>et al.</i>	(REHO, SLAC, TELA)	REFID=20098
ESPIGAT	72	NP B36 93	P. Espigat <i>et al.</i>	(CERN, CDEF)	REFID=21232
FOLEY	72	PR D6 747	K.J. Foley <i>et al.</i>	(BNL, CUNY)	REFID=21233
ALSTON...	71	PL 34B 156	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21214
BARNHAM	71	PRL 26 1494	K.W.J. Barnham <i>et al.</i>	(LBL)	REFID=21215
BINNIE	71	PL 36B 257	D.M. Binnie <i>et al.</i>	(LOIC, SHMP)	REFID=21217
BOWEN	71	PRL 26 1663	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21219
GRAYER	71	PL 34B 333	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=21223
ABRAMOVI...	70B	NP B23 466	M. Abramovich <i>et al.</i>	(CERN) JP	REFID=21195
ALSTON...	70	PL 33B 607	M. Alston-Garnjost <i>et al.</i>	(LRL)	REFID=21196
BOECKMANN	70	NP B16 221	K. Boeckmann <i>et al.</i>	(BONN, DURH, NIJM+)	REFID=21202
ASCOLI	68	PRL 20 1321	G. Ascoli <i>et al.</i>	(ILL) JP	REFID=21171
BOESEBECK	68	NP B4 501	K. Boesebeck <i>et al.</i>	(AACH, BERL, CERN)	REFID=20585
CHUNG	68	PR 165 1491	S.U. Chung <i>et al.</i>	(LRL)	REFID=20059
CONTE	67	NC 51A 175	F. Conte <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=21166

$f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances" and a note on "Non- $q\bar{q}$ Candidates" in PDG 06, Journal of Physics **G33** 1 (2006).

NODE=M147

NODE=M147

 $f_0(1370)$ T-MATRIX POLE \sqrt{s}

NODE=M147PP

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M147PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1250-1440) -i (60-300) OUR ESTIMATE			
$(1245 \pm 40) - i(300 \pm_{-70}^{+30})$	1 PELAEZ	23	RVUE Compilation
$(1380 \pm_{-60}^{+70}) - i(220 \pm_{-70}^{+80})$	2 PELAEZ	23	RVUE Compilation
$(1370 \pm 40) - i(195 \pm 20)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1280.6 \pm 1.6 \pm 47.4) - i(205.2 \pm 1.7 \pm 20.7)$	3 ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
$(1290 \pm 50) - i(170 \pm_{-40}^{+20})$	4 ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1373 \pm 15) - i(137 \pm 10)$	5 BARGIOTTI	03	OBLX $\bar{p}p$
$(1302 \pm 17) - i(166 \pm 18)$	6 BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS	99D	OMEG $450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1406 \pm 19) - i(80 \pm 6)$	7 KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH	98B	RVUE Compilation
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS	97B	OMEG $450 pp \rightarrow pp2(\pi^+ \pi^-)$
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1380 \pm 40) - i(180 \pm 25)$	ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
$(1300 \pm 15) - i(115 \pm 8)$	BUGG	96	RVUE
$(1330 \pm 50) - i(150 \pm 40)$	8 AMSLER	95B	CBAR $\bar{p}p \rightarrow 3\pi^0$
$(1360 \pm 35) - i(150-300)$	8 AMSLER	95C	CBAR $\bar{p}p \rightarrow \pi^0 \eta\eta$
$(1390 \pm 30) - i(190 \pm 40)$	9 AMSLER	95D	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
1346 - i249	10,11 JANSSEN	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1214 - i168	11,12 TORNQVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1364 - i139	AMSLER	94D	CBAR $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$(1365 \pm_{-55}^{+20}) - i(134 \pm 35)$	ANISOVICH	94	CBAR $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$
$(1340 \pm 40) - i(127 \pm_{-20}^{+30})$	13 BUGG	94	RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$
$(1430 \pm 5) - i(73 \pm 13)$	14 KAMINSKI	94	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$
1420 - i220	15 AU	87	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}$

NODE=M147PP

→ UNCHECKED ←

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M147PP;LINKAGE=G
 NODE=M147PP;LINKAGE=I
 NODE=M147PP;LINKAGE=E

NODE=M147PP;LINKAGE=AO
 NODE=M147PP;LINKAGE=BG
 NODE=M147PP;LINKAGE=PC
 NODE=M147PP;LINKAGE=TK
 NODE=M147PP;LINKAGE=K
 NODE=M147PP;LINKAGE=A

NODE=M147PP;LINKAGE=C
 NODE=M147PP;LINKAGE=DD

NODE=M147PP;LINKAGE=BB

NODE=M147PP;LINKAGE=C1
 NODE=M147PP;LINKAGE=KM
 NODE=M147PP;LINKAGE=H

¹ From forward dispersion relation applied to $\pi\pi$ scattering data.

² From partial-wave dispersion relation applied to $\pi\pi \rightarrow \bar{K}K$ data.

³ T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$), and BINON 84C ($\eta\eta'$).

⁴ Another pole is found at $(1510 \pm 130) - i(800 \pm_{-150}^{+100})$ MeV.

⁵ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

⁶ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$.

⁷ T-matrix pole on sheet ---.

⁸ Supersedes ANISOVICH 94.

⁹ Coupled-channel analysis of $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$, and $\pi^0 \pi^0 \eta$ on sheet IV. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

¹⁰ Analysis of data from FALVARD 88.

¹¹ The pole is on Sheet III. Demonstrates explicitly that $f_0(500)$ and $f_0(1370)$ are two different poles.

¹² Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

¹³ Reanalysis of ANISOVICH 94 data.

¹⁴ T-matrix pole on sheet III.

¹⁵ Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

$f_0(1370)$ BREIT-WIGNER MASS

NODE=M147205

VALUE (MeV)

DOCUMENT ID

1200 to 1500 OUR ESTIMATE

NODE=M147M

→ UNCHECKED ←

 $\pi\pi$ MODE

NODE=M147M1

NODE=M147M1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1400±40		¹ AUBERT 09L	BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
1470 ⁺⁶⁺⁷² ₋₇₋₂₅₅		² UEHARA 08A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1259±55	2.6k	BONVICINI 07	CLEO	$D^+ \rightarrow \pi^-\pi^+\pi^+$
1309±1±15		³ BUGG 07A	RVUE	0.0 $p\bar{p} \rightarrow 3\pi^0$
1449±13	4.3k	⁴ GARMASH 06	BELL	$B^+ \rightarrow K^+\pi^+\pi^-$
1350±50		ABLIKIM 05	BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
1265±30 ⁺²⁰ ₋₃₅		ABLIKIM 05Q	BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
1434±18±9	848	AITALA 01A	E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$
1308±10		BARBERIS 99B	OMEG	450 $pp \rightarrow p_S p_f \pi^+\pi^-$
1315±50		BELLAZZINI 99	GAM4	450 $pp \rightarrow pp\pi^0\pi^0$
1315±30		ALDE 98	GAM4	100 $\pi^-p \rightarrow \pi^0\pi^0n$
1280±55		BERTIN 98	OBLX	0.05-0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
1186		^{5,6} TORNQVIST 95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472±12		ARMSTRONG 91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275±20		BREAKSTONE90	SFM	62 $pp \rightarrow pp\pi^+\pi^-$
1420±20		AKESSON 86	SPEC	63 $pp \rightarrow pp\pi^+\pi^-$
1256		FROGGATT 77	RVUE	$\pi^+\pi^-$ channel

¹ Breit-Wigner mass.² Breit-Wigner mass. May also be the $f_0(1500)$.³ Reanalysis of ABELE 96C data.⁴ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0\pi^+\pi^-$ decays. Supersedes GARMASH 05.⁵ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.⁶ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays

NODE=M147M1;LINKAGE=BW

NODE=M147M1;LINKAGE=UE

NODE=M147M1;LINKAGE=BU

NODE=M147M1;LINKAGE=GR

NODE=M147M1;LINKAGE=BB

NODE=M147M1;LINKAGE=FF

 $K\bar{K}$ MODE

NODE=M147M2

NODE=M147M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1422±15±28		¹ AAIJ 19H	LHCB	$pp \rightarrow D^\pm X$
1360±31±28	430	^{2,3} DOBBS 15		$J/\psi \rightarrow \gamma K^+K^-$
1350±48±15	168	^{2,3} DOBBS 15		$\psi(2S) \rightarrow \gamma K^+K^-$
1440±6		VLADIMIRSK...06	SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1391±10		TIKHOMIROV 03	SPEC	40.0 $\pi^-C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440±50		BOLONKIN 88	SPEC	40 $\pi^-p \rightarrow K_S^0 K_S^0 n$
1463±9		ETKIN 82B	MPS	23 $\pi^-p \rightarrow n2K_S^0$
1425±15		WICKLUND 80	SPEC	6 $\pi N \rightarrow K^+K^-N$
~ 1300		POLYCHRO... 79	STRC	7 $\pi^-p \rightarrow n2K_S^0$

¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the isobar model A.² Using CLEO-c data but not authored by the CLEO Collaboration.³ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 346$ MeV.

NODE=M147M2;LINKAGE=F

NODE=M147M2;LINKAGE=A

NODE=M147M2;LINKAGE=B

 4π MODE $2(\pi\pi)_S + \rho\rho$

NODE=M147M3

NODE=M147M3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1395±40		ABELE 01	CBAR	0.0 $\bar{p}d \rightarrow \pi^-4\pi^0p$
1374±38		AMSLER 94	CBAR	0.0 $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
1345±12		ADAMO 93	OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-$
1386±30		GASPERO 93	DBC	0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$
~ 1410	5751	¹ BETTINI 66	DBC	0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$

¹ $\rho\rho$ dominant.

NODE=M147M3;LINKAGE=BE

$\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1262^{+51+82}_{-78-103}$	¹ UEHARA	10A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
1430	AMSLER	92	CBAR 0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
1220 ± 40	ALDE	86D	GAM4 100 $\pi^-p \rightarrow n2\eta$

¹ Breit-Wigner mass. May also be the $f_0(1500)$.

NODE=M147M4
NODE=M147M4

NODE=M147M4;LINKAGE=UE

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1330.2^{+5.9}_{-6.5} \pm 5.1$	¹ AAIJ	19H	LHCB $pp \rightarrow D^\pm X$
1306 ± 20	² ANISOVICH	03	RVUE

¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUDE 18.

² K-matrix pole from combined analysis of $\pi^-p \rightarrow \pi^0\pi^0n$, $\pi^-p \rightarrow K\bar{K}n$, $\pi^+\pi^- \rightarrow \pi^+\pi^-$, $\bar{p}p \rightarrow \pi^0\pi^0\pi^0$, $\pi^0\eta\eta$, $\pi^0\pi^0\eta$, $\pi^+\pi^-\pi^0$, $K^+K^-\pi^0$, $K_S^0K_S^0\pi^0$, $K^+K_S^0\pi^-$ at rest, $\bar{p}n \rightarrow \pi^-\pi^-\pi^+$, $K_S^0K^- \pi^0$, $K_S^0K_S^0\pi^-$ at rest.

NODE=M147M5
NODE=M147M5

NODE=M147M5;LINKAGE=A

NODE=M147M;LINKAGE=KM

 $f_0(1370)$ BREIT-WIGNER WIDTH

VALUE (MeV)	DOCUMENT ID
-------------	-------------

200 to 500 OUR ESTIMATE

NODE=M147210

NODE=M147W
→ UNCHECKED ←

 $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

300 ± 80		¹ AUBERT	09L	BABR $B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
90^{+2+50}_{-1-22}		² UEHARA	08A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
298 ± 21	2.6k	BONVICINI	07	CLEO $D^+ \rightarrow \pi^-\pi^+\pi^+$
126 ± 25	4286	³ GARMASH	06	BELL $B^+ \rightarrow K^+\pi^+\pi^-$
265 ± 40		ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-$
$350 \pm 100^{+105}_{-60}$		ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
$173 \pm 32 \pm 6$	848	AITALA	01A	E791 $D_s^+ \rightarrow \pi^-\pi^+\pi^+$
222 ± 20		BARBERIS	99B	OMEG 450 $pp \rightarrow p_S p_f \pi^+\pi^-$
255 ± 60		BELLAZZINI	99	GAM4 450 $pp \rightarrow pp\pi^0\pi^0$
190 ± 50		ALDE	98	GAM4 100 $\pi^-p \rightarrow \pi^0\pi^0n$
323 ± 13		BERTIN	98	OBLX 0.05–0.405 $\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
350		^{4,5} TORNQVIST	95	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195 ± 33		ARMSTRONG	91	OMEG 300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$
285 ± 60		BREAKSTONE	90	SFM 62 $pp \rightarrow pp\pi^+\pi^-$
460 ± 50		AKESSON	86	SPEC 63 $pp \rightarrow pp\pi^+\pi^-$
~ 400		⁶ FROGGATT	77	RVUE $\pi^+\pi^-$ channel

¹ The systematic errors are not reported.

² Breit-Wigner width. May also be the $f_0(1500)$.

³ Also observed by GARMASH 07 in $B^0 \rightarrow K_S^0\pi^+\pi^-$ decays. Supersedes GARMASH 05.

⁴ Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CA-SON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

⁵ Also observed by ASNER 00 in $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$ decays

⁶ Width defined as distance between 45 and 135° phase shift.

NODE=M147W1
NODE=M147W1

NODE=M147W1;LINKAGE=NS
NODE=M147W1;LINKAGE=UE
NODE=M147W1;LINKAGE=GR
NODE=M147W1;LINKAGE=BB

NODE=M147W1;LINKAGE=FF
NODE=M147W1;LINKAGE=E

 $K\bar{K}$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$324 \pm 38 \pm 42$	¹ AAIJ	19H	LHCB $pp \rightarrow D^\pm X$
121 ± 15	VLADIMIRSK...06	SPEC	40 $\pi^-p \rightarrow K_S^0K_S^0n$
55 ± 26	TIKHOMIROV 03	SPEC	40.0 $\pi^-C \rightarrow K_S^0K_S^0K_L^0X$
250 ± 80	BOLONKIN 88	SPEC	40 $\pi^-p \rightarrow K_S^0K_S^0n$
118^{+138}_{-16}	ETKIN 82B	MPS	23 $\pi^-p \rightarrow n2K_S^0$
160 ± 30	WICKLUND 80	SPEC	6 $\pi N \rightarrow K^+K^-N$
~ 150	POLYCHRO... 79	STRC	7 $\pi^-p \rightarrow n2K_S^0$

¹ From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the isobar model A.

NODE=M147W2
NODE=M147W2

OCCUR=3

NODE=M147W2;LINKAGE=C

4 π MODE 2($\pi\pi$) $_S$ + $\rho\rho$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
275 \pm 55		ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 \pm 61		AMSLER	94	CBAR 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
398 \pm 26		ADAMO	93	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 \pm 50		GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$
\sim 90	5751	¹ BETTINI	66	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+ 3\pi^-$

¹ $\rho\rho$ dominant.NODE=M147W3
NODE=M147W3

NODE=M147W3;LINKAGE=BE

 $\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
484 $^{+246+246}_{-170-263}$	¹ UEHARA	10A	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$
250	AMSLER	92	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \eta\eta$
320 \pm 40	ALDE	86D	GAM4 100 $\pi^- p \rightarrow n2\eta$

¹ Breit-Wigner width. May also be the $f_0(1500)$.NODE=M147W4
NODE=M147W4

NODE=M147W4;LINKAGE=UE

COUPLED CHANNEL MODE

VALUE (MeV)	DOCUMENT ID	TECN
-------------	-------------	------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●		
147 $^{+30}_{-50}$	¹ ANISOVICH	03 RVUE

¹ K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta\eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.NODE=M147W5
NODE=M147W5

NODE=M147W;LINKAGE=KM

 $f_0(1370)$ DECAY MODES

NODE=M147215;NODE=M147

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 4π	seen
Γ_3 $4\pi^0$	seen
Γ_4 $2\pi^+ 2\pi^-$	seen
Γ_5 $\pi^+ \pi^- 2\pi^0$	seen
Γ_6 $\rho\rho$	seen
Γ_7 $2(\pi\pi)_S$ -wave	seen
Γ_8 $\pi(1300)\pi$	seen
Γ_9 $a_1(1260)\pi$	seen
Γ_{10} $\eta\eta$	seen
Γ_{11} $K\bar{K}$	seen
Γ_{12} $K\bar{K}n\pi$	not seen
Γ_{13} 6π	not seen
Γ_{14} $\omega\omega$	not seen
Γ_{15} $\gamma\gamma$	seen
Γ_{16} $e^+ e^-$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=10;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←
DESIG=14;OUR EST;→ UNCHECKED ←
DESIG=15;OUR EST;→ UNCHECKED ←
DESIG=16;OUR EVAL;→ UNCHECKED ←
DESIG=17;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=11;OUR EST;→ UNCHECKED ←
DESIG=18;OUR EVAL;→ UNCHECKED ←
DESIG=19;OUR EVAL;→ UNCHECKED ←
DESIG=20;OUR EVAL;→ UNCHECKED ←
DESIG=12;OUR EST;→ UNCHECKED ←
DESIG=13;OUR EST;→ UNCHECKED ← **$f_0(1370)$ PARTIAL WIDTHS**

NODE=M147217

 $\Gamma(\gamma\gamma)$ See $\gamma\gamma$ widths under $f_0(500)$ and MORGAN 90. **Γ_{15}** NODE=M147W11
NODE=M147W11
NODE=M147W11 **$\Gamma(e^+ e^-)$**

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<20	90	VOROBYEV	88	ND $e^+ e^- \rightarrow \pi^0 \pi^0$
-----	----	----------	----	--------------------------------------

 Γ_{16} NODE=M147W12
NODE=M147W12

$f_0(1370) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M147225

 $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{10}\Gamma_{15}/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M147G01
NODE=M147G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

$121^{+133+169}_{-53-106}$	¹ UEHARA	10A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$
----------------------------	---------------------	-----	------	--

¹Including interference with the $f'_2(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

NODE=M147G01;LINKAGE=UE

 $f_0(1370) \text{ BRANCHING RATIOS}$

NODE=M147220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M147R3
NODE=M147R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10	95	OCHS	13	RVUE
0.26 ± 0.09		BUGG	96	RVUE
<0.15		¹ AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
<0.06		GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

¹Using AMSLER 95B ($3\pi^0$).

NODE=M147R3;LINKAGE=B

 $\Gamma(4\pi)/\Gamma_{\text{total}}$ $\Gamma_2/\Gamma = (\Gamma_3+\Gamma_4+\Gamma_5)/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R4
NODE=M147R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.72	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$
-------	---------	----	---

 $\Gamma(4\pi^0)/\Gamma(4\pi)$ Γ_3/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R12
NODE=M147R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ABELE	96	CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$
0.068 ± 0.005	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$

¹Model-dependent evaluation.

NODE=M147R12;LINKAGE=GA

 $\Gamma(2\pi^+2\pi^-)/\Gamma(4\pi)$ $\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3+\Gamma_4+\Gamma_5)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R5
NODE=M147R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.420 ± 0.014	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow 2\pi^+3\pi^-$
-------------------	----------------------	----	---

¹Model-dependent evaluation.

NODE=M147R5;LINKAGE=A

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(4\pi)$ $\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3+\Gamma_4+\Gamma_5)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R6
NODE=M147R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.512 ± 0.019	¹ GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow \text{hadrons}$
-------------------	----------------------	----	---

¹Model-dependent evaluation.

NODE=M147R6;LINKAGE=A

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_6/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R17
NODE=M147R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26 ± 0.07	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$
-----------------	-------	-----	--

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(\pi\pi)$ Γ_7/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R15
NODE=M147R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.6 ± 2.6	¹ ABELE	01	CBAR 0.0 $\bar{p}d \rightarrow \pi^-4\pi^0 p$
---------------	--------------------	----	---

¹From the combined data of ABELE 96 and ABELE 96C.

NODE=M147R;LINKAGE=KZ

 $\Gamma(2(\pi\pi)_{S\text{-wave}})/\Gamma(4\pi)$ Γ_7/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R16
NODE=M147R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.51 ± 0.09	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$
-----------------	-------	-----	--

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_{S\text{-wave}})$ Γ_6/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R10
 NODE=M147R10

••• We do not use the following data for averages, fits, limits, etc. •••

large	BARBERIS	00C	450 $p\rho \rightarrow \rho_f 4\pi p_S$
1.6 ± 0.2	AMSLER	94	CBAR $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
~ 0.65	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons

OCCUR=2

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_8/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R18
 NODE=M147R18

••• We do not use the following data for averages, fits, limits, etc. •••

0.17 ± 0.06	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi\rho$
-----------------	-------	-----	--

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_9/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R19
 NODE=M147R19

••• We do not use the following data for averages, fits, limits, etc. •••

0.06 ± 0.02	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi\rho$
-----------------	-------	-----	--

 $\Gamma(\eta\eta)/\Gamma(4\pi)$ $\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R14
 NODE=M147R14

••• We do not use the following data for averages, fits, limits, etc. •••

(28 ± 11) $\times 10^{-3}$	¹ ANISOVICH	02D	SPEC Combined fit
(4.7 ± 2.0) $\times 10^{-3}$	BARBERIS	00E	450 $p\rho \rightarrow \rho_f \eta\eta p_S$

¹ From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$), GAMS ($\pi\rho \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi\rho \rightarrow K\bar{K}n$) data.

NODE=M147R14;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R11
 NODE=M147R11

••• We do not use the following data for averages, fits, limits, etc. •••

0.35 ± 0.13	BUGG	96	RVUE
-----------------	------	----	------

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_{11}/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R13
 NODE=M147R13

••• We do not use the following data for averages, fits, limits, etc. •••

0.08 ± 0.08	ABLIKIM	05	BES2 $J/\psi \rightarrow \phi\pi^+\pi^-, \phi K^+ K^-$
0.91 ± 0.20	¹ BARGIOTTI	03	OBLX $\bar{p}p$
0.12 ± 0.06	² ANISOVICH	02D	SPEC Combined fit
0.46 $\pm 0.15 \pm 0.11$	BARBERIS	99D	OMEG 450 $p\rho \rightarrow K^+ K^-, \pi^+ \pi^-$

¹ Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$), GAMS ($\pi\rho \rightarrow \pi^0 \pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi\rho \rightarrow K\bar{K}n$) data.

NODE=M147R;LINKAGE=BG
 NODE=M147R;LINKAGE=CH

 $\Gamma(K\bar{K}n\pi)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R20
 NODE=M147R20

••• We do not use the following data for averages, fits, limits, etc. •••

<0.03	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	--

 $\Gamma(6\pi)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R21
 NODE=M147R21

••• We do not use the following data for averages, fits, limits, etc. •••

<0.22	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	--

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M147R22
 NODE=M147R22

••• We do not use the following data for averages, fits, limits, etc. •••

<0.13	GASPERO	93	DBC 0.0 $\bar{p}n \rightarrow$ hadrons
-------	---------	----	--

$\phi_0(1370)$ REFERENCES

NODE=M147

PELAEZ	23	PRL 130 051902	J.R. Pelaez, A. Rodas, J. Ruiz de Elvira (MADU+)	REFID=62199
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i> (BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i> (Crystal Barrel Collab.)	REFID=60439
AAJ	19H	JHEP 1904 063	R. Aaij <i>et al.</i> (LHCb Collab.)	REFID=59670
AOUDE	18	PR D98 056021	R.T. Aoude <i>et al.</i>	REFID=59987
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i> (NWES)	REFID=56805
OCHS	13	JP G40 043001	W. Ochs	REFID=55367
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i> (BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev (PNPI)	REFID=52719
AUBERT	09L	PR D79 072006	B. Aubert <i>et al.</i> (BABAR Collab.)	REFID=52723
PDG	08	PL B667 1	C. Amsler <i>et al.</i> (PDG Collab.)	REFID=52166
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i> (BELLE Collab.)	REFID=52309
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i> (CLEO Collab.)	REFID=51721
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>	REFID=53252
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i> (BELLE Collab.)	REFID=51594
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i> (BELLE Collab.)	REFID=51162
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i> (PDG Collab.)	REFID=51004
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i> (ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i> (BES Collab.)	REFID=50450
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i> (BES Collab.)	REFID=50958
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i> (BELLE Collab.)	REFID=50641
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	REFID=49401
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i> (OBELIX Collab.)	REFID=49217
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	REFID=49423
		Translated from YAF 66 860.		
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	REFID=48831
		Translated from YAF 65 1583.		
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i> (Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i> (Crystal Barrel Collab.)	REFID=48356
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)	REFID=48004
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i> (CLEO Collab.)	REFID=47339
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i> (WA 102 Collab.)	REFID=47959
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i> (WA 102 Collab.)	REFID=47961
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i> (Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i> (Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>	REFID=47400
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau (CRAC, PARIN)	REFID=46927
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i> (GAM4 Collab.)	REFID=46605
		Also PAN 62 405	D. Alde <i>et al.</i> (GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.		
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	REFID=46331
		Translated from UFN 168 481.		
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i> (OBELIX Collab.)	REFID=45782
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i> (WA 102 Collab.)	REFID=45758
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i> (OBELIX Collab.)	REFID=45701
ABELE	96	PL B380 453	A. Abele <i>et al.</i> (Crystal Barrel Collab.)	REFID=45011
ABELE	96B	PL B385 425	A. Abele <i>et al.</i> (Crystal Barrel Collab.)	REFID=45038
ABELE	96C	NP A609 562	A. Abele <i>et al.</i> (Crystal Barrel Collab.)	REFID=45076
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou (LOQM, PNPI)	REFID=45094
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=44441
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i> (STON, ADLD, JULI)	REFID=44508
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist (HEL5)	REFID=44529
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i> (Crystal Barrel Collab.) JPC	REFID=43660
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i> (Crystal Barrel Collab.) JPC	REFID=43659
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i> (LOQM)	REFID=44078
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet (CRAC+)	REFID=45771
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i> (OBELIX Collab.) JPC	REFID=43657
GASPERO	93	NP A562 407	M. Gaspero (ROMA1) JPC	REFID=43658
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i> (Crystal Barrel Collab.)	REFID=43169
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+)	REFID=41744
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+)	REFID=41862
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i> (ISU, BGNA, CERN+)	REFID=41376
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington (RAL, DURH)	REFID=41583
ASTON	88	NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)	REFID=40262
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i> (ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i> (CLER, FRAS, LALO+)	REFID=40576
VOROBYEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i> (NOVO)	REFID=41023
		Translated from YAF 48 436.		
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington (DURH, RAL)	REFID=40064
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i> (Axial Field Spec. Collab.)	REFID=21123
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i> (BELG, LAPP, SERP, CERN+)	REFID=20765
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i> (BNL, BRAN, CUNY+)	REFID=20768
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i> (BELG, LAPP, SERP+)	REFID=21418
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i> (BELG, LAPP, SERP+)	REFID=20750
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i> (NDAM, ANL)	REFID=20752
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i> (BNL, CUNY, TUFTS, VAND)	REFID=20390
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i> (ANL)	REFID=20383
BECKER	79	NP B151 46	H. Becker <i>et al.</i> (MPIM, CERN, ZEEM, CRAC)	REFID=21084
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i> (NDAM, ANL)	REFID=20378
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen (GLAS, NORD)	REFID=21072
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i> (GEVA, SACL)	REFID=11004
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i> (CERN, MPIM)	REFID=20355
GRAYR	74	NP B75 189	G. Grayer <i>et al.</i> (CERN, MPIM)	REFID=20113
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i> (CERN, MPIM)	REFID=20107
OCHS	73	Thesis	W. Ochs (MPIM, MUNI)	REFID=20349
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i> (PENN)	REFID=44530
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i> (PADO, PISA)	REFID=21361

NODE=M111

$\pi_1(1400)$

$I^G(J^{PC}) = 1^-(1^-+)$

OMITTED FROM SUMMARY TABLE

Coupled channel analyses (see e.g. ALBRECHT 20) favor the existence of only one broad 1^-+ isovector state consistent with $\pi_1(1600)$ in the 1400–1600 MeV region. See the review on "Spectroscopy of Light Meson Resonances." See also $\pi_1(1600)$.

NODE=M111

$\pi_1(1400)$ REFERENCES

ALBRECHT 20 EPJ C80 453 M. Albrecht *et al.* (Crystal Barrel Collab.)

NODE=M111

REFID=60439

NODE=M027

$\eta(1405)$

$I^G(J^{PC}) = 0^+(0^-+)$

See also the $\eta(1475)$.

NODE=M027

$\eta(1405)$ MASS


NODE=M027205

NODE=M027MX

VALUE (MeV) DOCUMENT ID

$1408.7^{+2.0}_{-1.2}$ OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below. [1408.8 ± 2.0 MeV OUR 2023 AVERAGE Scale factor = 2.2]

NEW



$\eta\pi\pi$ MODE

NODE=M027M1
NODE=M027M1

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
The data in this block is included in the average printed for a previous datablock.

1405.8 ± 2.6 OUR AVERAGE Error includes scale factor of 2.3. See the ideogram below.

1409.0 ± 1.7	743	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
1407.0 ± 3.5	198	ABLIKIM	12E BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$
1399.8 ± 2.2 ^{+2.8} _{-0.1}	1	ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
1392 ± 14	900 ± 375	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$

OCCUR=2

1394 ± 8	6.6 ± 2.0k	AMSLER	04B CBAR	$0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta$
1404 ± 6	9082	MANAK	00A MPS	$18 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
1424 ± 6	2200	ALDE	97B GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1400 ± 6		² BOLTON	92B MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1388 ± 4		FUKUI	91C SPEC	$8.95 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
1398 ± 6	261	³ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1420 ± 5		ANDO	86 SPEC	$8 \pi^- p \rightarrow \eta \pi^+ \pi^- n$


OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1404.0 ± 11.0	195	ABLIKIM	19BABES3	$e^+ e^- \rightarrow \psi(2S)$
1385 ± 7		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
1409 ± 3		⁴ AMSLER	95F CBAR	$0 \bar{p} p \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta$

NODE=M027M1;LINKAGE=BL
 NODE=M027M1;LINKAGE=J1
 NODE=M027M1;LINKAGE=A1
 NODE=M027M1;LINKAGE=A

- ¹ The selected process is $J/\psi \rightarrow \omega a_0(980) \pi$.
- ² From fit to the $a_0(980) \pi 0^-+$ partial wave.
- ³ Best fit with a single Breit Wigner.
- ⁴ Superseded by AMSLER 04B.



$K \bar{K} \pi$ MODE ($a_0(980) \pi$ or direct $K \bar{K} \pi$)

NODE=M027M4
 NODE=M027M4

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

1413.5^{+2.0}_{-0.9} OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below. [1413.9 ± 1.7 MeV OUR 2023 AVERAGE Scale factor = 1.1]

NEW

1391.7 ± 0.7 ^{+11.3} _{-0.3}	126k	¹ ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1413 ± 14	3651	² NICHITIU	02 OBLX	$0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1416 ± 4 ± 2	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
1405 ± 5		³ CICALO	99 OBLX	$0 \bar{p} p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
1407 ± 5		³ BERTIN	97 OBLX	$0 \bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
1416 ± 2		³ BERTIN	95 OBLX	$0 \bar{p} p \rightarrow K \bar{K} \pi \pi \pi$
1416 ± 8 ⁺⁷ ₋₅	700	⁴ BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1413 ± 5		⁴ RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$

OCCUR=2


OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

1459 ± 5		⁵ AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
----------	--	-----------------------	--------	---

- ¹ ABLIKIM 23M reports for this state a significance from the fit much higher than 35 σ .
- ² Decaying dominantly directly to $K^+ K^- \pi^0$.
- ³ Decaying into $(K \bar{K})_S \pi$, $(K \pi)_S \bar{K}$, and $a_0(980) \pi$.
- ⁴ From fit to the $a_0(980) \pi 0^-+$ partial wave. Cannot rule out a $a_0(980) \pi 1^++$ partial wave.
- ⁵ Excluded from averaging because averaging would be meaningless.

NODE=M027M4;LINKAGE=A
 NODE=M027M;LINKAGE=NC
 NODE=M027M4;LINKAGE=FX
 NODE=M027M4;LINKAGE=C2
 NODE=M027M4;LINKAGE=AA



$\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1403±17 OUR AVERAGE				Error includes scale factor of 1.8.
1390±12	235 ± 91	AMSLER	04B CBAR	$0\bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$
1424±10±11	547	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1401±18		^{1,2} AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$
1432± 8		² COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$

NODE=M027M2
NODE=M027M2

¹ Best fit with a single Breit Wigner.

² This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

OCCUR=4

NODE=M027M2;LINKAGE=E
NODE=M027M2;LINKAGE=X

4π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1420±20		BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
1489±12	3270	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

NODE=M027M3
NODE=M027M3

NODE=M027M3;LINKAGE=E

$K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1452.7± 3.3	191	^{1,2} ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
1437.6± 3.2	249 ± 35	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
1445.9± 5.7	62 ± 18	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
1442 ± 10	410	¹ BAI	98C BES	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1445 ± 8	693	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1433 ± 8	296	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1413 ± 8	500	¹ DUCH	89 ASTE	$\bar{p}p \rightarrow \pi^+\pi^- K^\pm \pi^\mp K^0$
1453 ± 7	170	¹ RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1419 ± 1	8800	¹ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ K^0 \pi^- n$
1424 ± 3	620	¹ REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K\bar{K}\pi X$
1421 ± 2		¹ CHUNG	85 SPEC	$8 \pi^- p \rightarrow K\bar{K}\pi n$
1440 ⁺²⁰ ₋₁₅	174	¹ EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1440 ⁺¹⁰ ₋₁₅		¹ SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1425 ± 7	800	^{1,3} BAILLON	67 HBC	$0\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

² Systematic uncertainty not evaluated.

³ From best fit of 0^-+ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.

NODE=M027M6
NODE=M027M6

NODE=M027M;LINKAGE=NP

NODE=M027M;LINKAGE=NS

NODE=M027M6;LINKAGE=H

$\eta(1405)$ WIDTH


NODE=M027210

VALUE (MeV) _____ DOCUMENT ID _____

NODE=M027WX

50.3±2.5 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.6. See the ideogram below. [50.1 ± 2.6 MeV OUR 2023 AVERAGE Scale factor = 1.7]

NEW



$\eta\pi\pi$ MODE

NODE=M027W1
NODE=M027W1

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____

The data in this block is included in the average printed for a previous datablock.

52.6± 3.2 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

48.3± 5.2	743	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^0)$
55.0±11.0	198	ABLIKIM	12E	BES3	$J/\psi \rightarrow \gamma(\pi^0\pi^0\pi^0)$
52.8± 7.6 ^{+0.1} _{-7.6}	1	ABLIKIM	11J	BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
55 ±11	900	AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$
55 ±12	6.6k	AMSLER	04B	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma$
80 ±21	9.0k	MANAK	00A	MPS	$18 \pi^-p \rightarrow \eta\pi^+\pi^-n$
85 ±18	2.2k	ALDE	97B	GAM4	$100 \pi^-p \rightarrow \eta\pi^0\pi^0n$
47 ±13		2 BOLTON	92B	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
59 ± 4		FUKUI	91C	SPEC	$8.95 \pi^-p \rightarrow \eta\pi^+\pi^-n$
53 ±11		3 AUGUSTIN	90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
31 ± 7		ANDO	86	SPEC	$8 \pi^-p \rightarrow \eta\pi^+\pi^-n$

OCCUR=2


OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

79.0±16.0	195	ABLIKIM	19BA	BES3	$e^+e^- \rightarrow \psi(2S)$
86 ±10	4	AMSLER	95F	CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$

- The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$.
- From fit to the $a_0(980)\pi 0^-+$ partial wave.
- From $\eta\pi^+\pi^-$ mass distribution - mainly $a_0(980)\pi$ - no spin-parity determination available.
- Superseded by AMSLER 04B.

NODE=M027W1;LINKAGE=BL
NODE=M027W1;LINKAGE=A1
NODE=M027W1;LINKAGE=D1
NODE=M027W1;LINKAGE=A



$K\bar{K}\pi$ MODE ($a_0(980)\pi$ or direct $K\bar{K}\pi$)

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M027W4
NODE=M027W4


49 ± 4 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.
[48 ± 4 MeV OUR 2023 AVERAGE Scale factor = 2.1]

NEW

60.8 ± 1.2 ^{+5.5} _{-12.0}	126K	ABLIKIM	23M	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$	
51 ± 6	3651	¹ NICHITIU	02	OBLX	$0 \bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
42 ± 10 ± 9	20k	ADAMS	01B	B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$	
50 ± 4		CICALO	99	OBLX	$0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$	
48 ± 5		² BERTIN	97	OBLX	$0.0 \bar{p}p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$	
50 ± 4		² BERTIN	95	OBLX	$0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$	
75 ± 9		AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$	
91 ⁺⁶⁷ ₋₃₁ ⁺¹⁵ ₋₃₈		³ BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=2
19 ± 7		³ RATH	89	MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$	OCCUR=3

- ¹Decaying dominantly directly to $K^+ K^- \pi^0$.
- ²Decaying into $(K\bar{K})_S \pi$, $(K\pi)_S \bar{K}$, and $a_0(980)\pi$.
- ³From fit to the $a_0(980)\pi 0^- +$ partial wave, but $a_0(980)\pi 1^+ +$ cannot be excluded.

NODE=M027W;LINKAGE=NC
NODE=M027W4;LINKAGE=FX
NODE=M027W4;LINKAGE=C



$\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
89 ±17	OUR AVERAGE	Error includes scale factor of 1.7.		
64 ±18	235 ± 91	AMSLER	04B CBAR	$0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
101.0 ± 8.8 ± 8.8	547	BAI	04J BES2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
174 ±44		AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$
90 ±26		¹ COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$

¹ This peak in the $\gamma\rho$ channel may not be related to the $\eta(1405)$.

NODE=M027W2
NODE=M027W2

OCCUR=2

NODE=M027W2;LINKAGE=X

4 π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
160 ±30		BUGG	95 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$
144 ±13	3270	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

NODE=M027W3
NODE=M027W3

NODE=M027W3;LINKAGE=F2

 $K\bar{K}\pi$ MODE (unresolved)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
45.9 ± 8.2	191	^{1,2} ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K K \pi$
48.9 ± 9.0	249 ± 35	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K_S^0 K^+ \pi^- + c.c.$
34.2 ± 18.5	62 ± 18	^{1,2} ABLIKIM	08E BES2	$J/\psi \rightarrow \omega K^+ K^- \pi^0$
93 ± 14	296	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
105 ± 10	693	¹ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
62 ± 16	500	¹ DUCH	89 ASTE	$\bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
100 ± 11	170	¹ RATH	89 MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
66 ± 2	8800	¹ BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
60 ± 10	620	¹ REEVES	86 SPEC	$6.6 p\bar{p} \rightarrow K K \pi X$
60 ± 10		¹ CHUNG	85 SPEC	$8 \pi^- p \rightarrow K\bar{K}\pi n$
55 ⁺²⁰ / ₋₃₀	174	¹ EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
50 ⁺³⁰ / ₋₂₀		¹ SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
80 ± 10	800	^{1,3} BAILLON	67 HBC	$0.0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

¹ These experiments identify only one pseudoscalar in the 1400–1500 range. Data could also refer to $\eta(1475)$.

² Systematic uncertainty not evaluated.

³ From best fit to 0^-+ partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.

NODE=M027W6
NODE=M027W6

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

NODE=M027W;LINKAGE=NP

NODE=M027W;LINKAGE=NS

NODE=M027W6;LINKAGE=H1

 $\eta(1405)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\bar{K}\pi$	seen	
Γ_2 $\eta\pi\pi$	seen	
Γ_3 $a_0(980)\pi$	seen	
Γ_4 $\eta(\pi\pi)_S$ -wave	seen	
Γ_5 $f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0$	not seen	
Γ_6 $f_0(980)\eta$	seen	
Γ_7 4π	seen	
Γ_8 $\rho\rho$	<58 %	99.85%
Γ_9 $\gamma\gamma$		
Γ_{10} $\rho^0\gamma$	seen	
Γ_{11} $\phi\gamma$		
Γ_{12} $K^*(892)K$	seen	

NODE=M027215;NODE=M027

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=9;OUR EST;→ UNCHECKED ←

DESIG=15

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

DESIG=12

DESIG=7

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=13

DESIG=11;OUR EST;→ UNCHECKED ←

 $\eta(1405)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M027220

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_1\Gamma_9/\Gamma$**

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.035	90	^{1,2} AHOHE	05 CLE2	$10.6 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$

¹ Using $\eta(1405)$ mass and width 1410 MeV and 51 MeV, respectively.

² Assuming three-body phase-space decay to $K_S^0 K^\pm \pi^\mp$.

NODE=M027G3
NODE=M027G3

NODE=M027G3;LINKAGE=AH

NODE=M027G3;LINKAGE=B3

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_9/\Gamma$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.095	95	ACCIARRI	01G L3	183-202 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

NODE=M027G5
NODE=M027G5

 $\Gamma(\rho^0\gamma) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{10}\Gamma_9/\Gamma$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	95	ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
------	----	---------	----------	---

NODE=M027G8
NODE=M027G8

 $\eta(1405)$ BRANCHING RATIOS

NODE=M027225

 $\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

1.09±0.48 ¹ AMSLER 04B CBAR $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.5 90 EDWARDS 83B CBAL $J/\psi \rightarrow \eta\pi\pi\gamma$

<1.1 90 SCHARRE 80 MRK2 $J/\psi \rightarrow \eta\pi\pi\gamma$

<1.5 95 FOSTER 68B HBC $0.0 \bar{p}p$

¹ Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.

NODE=M027R3
NODE=M027R3

NODE=M027R3;LINKAGE=AM

 $\Gamma(\rho^0\gamma)/\Gamma(\eta\pi\pi)$ Γ_{10}/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.111±0.064 AMSLER 04B CBAR $0 \bar{p}p$

NODE=M027R12
NODE=M027R12

 $\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$ Γ_3/Γ_1

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.15 ¹ BERTIN 95 OBLX $0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$

~ 0.8 500 ¹ DUCH 89 ASTE $\bar{p}p \rightarrow \pi^+\pi^-\pi^+\pi^-\pi^0$

~ 0.75 ¹ REEVES 86 SPEC $6.6 \bar{p}p \rightarrow KK\pi X$

¹ Assuming that the $a_0(980)$ decays only into $K\bar{K}$.

NODE=M027R4
NODE=M027R4

NODE=M027R4;LINKAGE=C

 $\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$ Γ_3/Γ_2

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29±0.10 ABELE 98E CBAR $0 \bar{p}p \rightarrow \eta\pi^0\pi^0\pi^0$

0.19±0.04 2200 ¹ ALDE 97B GAM4 $100 \pi^-p \rightarrow \eta\pi^0\pi^0n$

0.56±0.04±0.03 ¹ AMSLER 95F CBAR $0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0\pi^0\eta$

¹ Assuming that the $a_0(980)$ decays only into $\eta\pi$.

NODE=M027R2
NODE=M027R2

NODE=M027R2;LINKAGE=A

 $\Gamma(a_0(980)\pi)/\Gamma(\eta(\pi\pi)S\text{-wave})$ Γ_3/Γ_4

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.91±0.12 ANISOVICH 01 SPEC $0.0 \bar{p}p \rightarrow \eta\pi^+\pi^-\pi^+\pi^-$

0.15±0.04 9082 ¹ MANAK 00A MPS $18 \pi^-p \rightarrow \eta\pi^+\pi^-n$

0.70±0.12±0.20 ² BAI 99 BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Statistical error only.

² Assuming that the $a_0(980)$ decays only into $\eta\pi$.

NODE=M027R9
NODE=M027R9

NODE=M027R;LINKAGE=K3
NODE=M027R9;LINKAGE=BK

 $\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$ Γ_{10}/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0152±0.0038 ¹ COFFMAN 90 MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Using $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma K\bar{K}\pi)=4.2 \times 10^{-3}$ and $B(J/\psi \rightarrow \gamma\eta(1405) \rightarrow \gamma\gamma\rho^0)=6.4 \times 10^{-5}$.

NODE=M027R7
NODE=M027R7

NODE=M027R7;LINKAGE=D

 $\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$ Γ_9/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<**1.78 × 10⁻³** 90 ¹ ABLIKIM 180 BES3 $\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$

¹ Using results from BAI 00D.

NODE=M027R02
NODE=M027R02

NODE=M027R02;LINKAGE=A

 $\Gamma(\eta(\pi\pi)S\text{-wave})/\Gamma(\eta\pi\pi)$ Γ_4/Γ_2

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.81±0.04 2200 ALDE 97B GAM4 $100 \pi^-p \rightarrow \eta\pi^0\pi^0n$

NODE=M027R8
NODE=M027R8

$\Gamma(f_0(980)\eta)/\Gamma(\eta\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32±0.07 ¹ ANISOVICH 00 SPEC 0.9–1.2 $\bar{p}p \rightarrow \eta 3\pi^0$

¹ Using preliminary Crystal Barrel data.

Γ₆/Γ₂
 NODE=M027R10
 NODE=M027R10

NODE=M027R10;LINKAGE=D

 $\Gamma(f_0(980)\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ¹ ABLIKIM 17AJ BES3 $\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

¹ ABLIKIM 17AJ reports $B(\psi(2S) \rightarrow \gamma\eta(1405) \rightarrow \gamma f_0(980)\pi^0 \rightarrow \gamma\pi^+\pi^-\pi^0) < 5.0 \times 10^{-7}$.

Γ₅/Γ
 NODE=M027R00
 NODE=M027R00

NODE=M027R00;LINKAGE=A

 $\Gamma(\rho\rho)/\Gamma_{total}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.58 99.85 ^{1,2} AMSLER 04B CBAR 0 $\bar{p}p$

¹ Assuming that the $\eta(1405)$ decays are saturated by the $\pi\pi\eta$, $K\bar{K}\pi$ and $\rho\rho$ modes.

² Using the data of BAILLON 67 on $\bar{p}p \rightarrow K\bar{K}\pi$.

Γ₈/Γ
 NODE=M027R13
 NODE=M027R13

NODE=M027R13;LINKAGE=AM
 NODE=M027R13;LINKAGE=AS

 $\Gamma(K^*(892)K)/\Gamma(a_0(980)\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.084±0.024 ¹ ADAMS 01B B852 18 GeV $\pi^-p \rightarrow K^+K^-\pi^0n$

¹ Statistical error only.

Γ₁₂/Γ₃
 NODE=M027R11
 NODE=M027R11

NODE=M027R11;LINKAGE=K3

 $\Gamma(\phi\gamma)/\Gamma(\rho^0\gamma)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.09±0.03 ¹ ABLIKIM 18I BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$

0.13±0.04 ² ABLIKIM 18I BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$

<0.77 95 ³ BAI 04J BES2 $J/\psi \rightarrow \gamma\gamma K^+K^-$

¹ Constructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18I reports the inverse as 11.10 ± 3.5 .

² Destructive interference between $X(1835)$ and $\eta(1405)/\eta(1475)$ decays to $\gamma\phi$ is assumed. Also see $\eta(1475)$. ABLIKIM 18I reports the inverse as 7.53 ± 2.49 .

³ Calculated by us from $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \phi\gamma\gamma) < 0.82 \times 10^{-4}$ and $B(J/\psi \rightarrow \eta(1405)\gamma \rightarrow \rho^0\gamma\gamma) = (1.07 \pm 0.17 \pm 0.11) \times 10^{-4}$.

Γ₁₁/Γ₁₀
 NODE=M027R14
 NODE=M027R14

OCCUR=2

NODE=M027R14;LINKAGE=A

NODE=M027R14;LINKAGE=B

NODE=M027R14;LINKAGE=BA

 $\eta(1405)$ REFERENCES

ABLIKIM 23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 12E	PRL 108 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)
AHOHE 05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)
AMSLER 04B	EPJ C33 23	C. Amshler <i>et al.</i>	(Crystal Barrel Collab.)
BAI 04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)
NICHITIU 02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ACCIARRI 01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS 01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH 01	NP A690 567	A.V. Anisovich <i>et al.</i>	
ANISOVICH 00	PL B472 168	A.V. Anisovich <i>et al.</i>	
BAI 00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
MANAK 00A	PR D62 012003	J.J. Manak <i>et al.</i>	(BNL E852 Collab.)
BAI 99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
CICALO 99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)
ABELE 98E	NP B514 45	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BAI 98C	PL B440 217	J.Z. Bai <i>et al.</i>	(BES Collab.)
ALDE 97B	PAN 60 386	D. Alde <i>et al.</i>	(GAMS Collab.)
Translated from YAF 60 458.			
BERTIN 97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
AMSLER 95F	PL B358 389	C. Amshler <i>et al.</i>	(Crystal Barrel Collab.)
BERTIN 95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BUGG 95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
AUGUSTIN 92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BOLTON 92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)
FUKUI 91C	PL B267 293	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)
AUGUSTIN 90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)
BAI 90C	PR D41 1410	Z. Bai <i>et al.</i>	(Mark III Collab.)
COFFMAN 90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)
BISELLO 89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)
DUCH 89	ZPHY C45 223	K.D. Duch <i>et al.</i>	(ASTERIX Collab.) JP
RATH 89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
BIRMAN 88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASH) JP
ANDO 86	PR D57 1296	A. Ando <i>et al.</i>	(KEK, KYOT, NIRS, SAGA+) IJP
REEVES 86	PR D34 1960	D.F. Reeves <i>et al.</i>	(FLOR, BNL, IND+) JP
CHUNG 85	PRL 55 779	S.U. Chung <i>et al.</i>	(BNL, FLOR, IND+) JP
ALTHOFF 84E	PL 147B 487	M. Althoff <i>et al.</i>	(TASSO Collab.)
EDWARDS 83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
EDWARDS 82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
Also	PRL 50 219	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
SCHARRE 80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)
FOSTER 68B	NP B8 174	M. Foster <i>et al.</i>	(CERN, CDEF)
BAILLON 67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)

NODE=M027

REFID=62055
 REFID=60024
 REFID=58893
 REFID=58925
 REFID=58322
 REFID=55386
 REFID=54270
 REFID=53931
 REFID=52143
 REFID=50764
 REFID=51079
 REFID=50167
 REFID=48848
 REFID=48319
 REFID=49649
 REFID=48308
 REFID=47429
 REFID=47954
 REFID=47989
 REFID=46606
 REFID=47394
 REFID=46314
 REFID=46337
 REFID=45396

REFID=45417
 REFID=44613
 REFID=44614
 REFID=44438
 REFID=41584
 REFID=42176
 REFID=41748
 REFID=41352
 REFID=41578
 REFID=41350
 REFID=40575
 REFID=41016
 REFID=40924
 REFID=40568
 REFID=20891
 REFID=20936
 REFID=20934
 REFID=20305
 REFID=21318
 REFID=21314
 REFID=21315
 REFID=21329
 REFID=21179
 REFID=20407

NODE=M109

h₁(1415)

$$I^G(J^{PC}) = 0^-(1^{+-})$$

was h₁(1380)

h₁(1415) MASS

NODE=M109M

NODE=M109M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1409⁺⁹₋₈ OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.


1384 ± 6	⁺⁹ ₋₀	1 ABLIKIM	22C BES3	J/ψ → γη'η' → 4/5γ2(π ⁺ π ⁻)
1423 ± 2.1 ± 7.3	2.2k	2 ABLIKIM	18AB BES3	J/ψ → η'h ₁ → η'K* \bar{K}
1412 ± 4 ± 8		2 ABLIKIM	15M BES3	ψ(2S) → γχ _{c1,2} → γφ(h ₁ → K* \bar{K})
1440 ± 60		ABELE	97H CBAR	$\bar{p}p$ → K _L ⁰ K _S ⁰ π ⁰ π ⁰
1380 ± 20		ASTON	88C LASS	11 K ⁻ p → K _S ⁰ K [±] π [∓] Λ

¹ From a partial wave analysis of the systems (γX), with X → η'η', and (η'X), with X → γη' in the decay J/ψ → γη'η'. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

² Final states K⁺K⁻π⁰ and K_S⁰K[±]π[∓].

NODE=M109M;LINKAGE=B

NODE=M109M;LINKAGE=A



h₁(1415) WIDTH

NODE=M109W

NODE=M109W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

78 ± 11 OUR AVERAGE

66 ± 10	⁺¹² ₋₁₀	1 ABLIKIM	22C BES3	J/ψ → γη'η' → 4/5γ2(π ⁺ π ⁻)
90.3 ± 9.8 ± 17.5	2.2k	2 ABLIKIM	18AB BES3	J/ψ → η'h ₁ → η'K* \bar{K}
84 ± 12 ± 40		2 ABLIKIM	15M BES3	ψ(2S) → γχ _{c1,2} → γφ(h ₁ → K* \bar{K})
170 ± 80		ABELE	97H CBAR	$\bar{p}p$ → K _L ⁰ K _S ⁰ π ⁰ π ⁰
80 ± 30		ASTON	88C LASS	11 K ⁻ p → K _S ⁰ K [±] π [∓] Λ

¹ From a partial wave analysis of the systems (γX), with X → η'η', and (η'X), with X → γη' in the decay J/ψ → γη'η'. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

² Final states K⁺K⁻π⁰ and K_S⁰K[±]π[∓].

NODE=M109W;LINKAGE=B

NODE=M109W;LINKAGE=A

$h_1(1415)$ DECAY MODES

NODE=M109215;NODE=M109

Mode	
Γ_1	$K\bar{K}^*(892)+$ c.c.

DESIG=1

 $h_1(1415)$ REFERENCES

ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

NODE=M109

REFID=61637
REFID=59456
REFID=56778
REFID=45765
REFID=40282 $f_1(1420)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

NODE=M006

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M006

 $f_1(1420)$ MASS

NODE=M006M2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M006M2

1428.4 $^{+1.5}_{-1.3}$ OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below. [1426.3 \pm 0.9 MeV OUR 2023 AVERAGE Scale factor = 1.1]


NEW

1433.5 \pm 1.1 $^{+27.9}_{-0.7}$	126K	ABLIKIM	23M	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1434 \pm 5 \pm 5	133	¹ ACHARD	07	L3	$183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$
1426 \pm 6	711	ABDALLAH	03H	DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
1420 \pm 14	3651	NICHITIU	02	OBLX	$0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1428 \pm 4 \pm 2	20k	ADAMS	01B	B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
1426 \pm 1		BARBERIS	97C	OMEG	$450 p p \rightarrow p p K_S^0 K^\pm \pi^\mp$
1425 \pm 8		BERTIN	97	OBLX	$0.0 \bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
1430 \pm 4		² ARMSTRONG	92E	OMEG	$85,300 \pi^+ p, p p \rightarrow \pi^+ p, p p (K\bar{K}\pi)$
1462 \pm 20		³ AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
1443 $^{+7}_{-6}$ $^{+3}_{-2}$	1100	BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1425 \pm 10	17	BEHREND	89	CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
1442 \pm 5 $^{+10}_{-17}$	111	BECKER	87	MRK3	$e^+ e^-, \omega K\bar{K}\pi$
1423 \pm 4		GIDAL	87B	MRK2	$e^+ e^- \rightarrow e^+ e^- K\bar{K}\pi$
1417 \pm 13	13	AIHARA	86C	TPC	$e^+ e^- \rightarrow e^+ e^- K\bar{K}\pi$
1422 \pm 3		CHAUVAT	84	SPEC	ISR 31.5 $p p$
1440 \pm 10		⁴ BROMBERG	80	SPEC	$100 \pi^- p \rightarrow K\bar{K}\pi X$
1426 \pm 6	221	DIONISI	80	HBC	$4 \pi^- p \rightarrow K\bar{K}\pi n$
1420 \pm 20		DAHL	67	HBC	$1.6-4.2 \pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1430.8 \pm 0.9		⁵ SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}} (K_S^0 K^+ \pi^-) p_{\text{fast}}$
1433.4 \pm 0.8		⁵ SOSA	99	SPEC	$p p \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
1435 \pm 9		PROKOSHKIN	97B	GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
1429 \pm 3	389	ARMSTRONG	89	OMEG	$300 p p \rightarrow K\bar{K}\pi p p$
1425 \pm 2	1520	ARMSTRONG	84	OMEG	$85 \pi^+ p, p p \rightarrow (\pi^+, p)(K\bar{K}\pi)p$
~ 1420		BITYUKOV	84	SPEC	$32 K^- p \rightarrow K^+ K^- \pi^0 \gamma$

OCCUR=2

- 1 From a fit with a width fixed at 55 MeV.
- 2 This result supersedes ARMSTRONG 84, ARMSTRONG 89.
- 3 From fit to the $K^*(892)K 1^{++}$ partial wave.
- 4 Mass error increased to account for $a_0(980)$ mass cut uncertainties.
- 5 No systematic error given.

NODE=M006M2;LINKAGE=CH
 NODE=M006M2;LINKAGE=C
 NODE=M006M2;LINKAGE=B
 NODE=M006M2;LINKAGE=A
 NODE=M006M2;LINKAGE=N1



f₁(1420) WIDTH

NODE=M006W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
56.7 ± 3.3 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below. [54.5 ± 2.6 MeV OUR 2023 AVERAGE]		
95.9 ± 2.3 ^{+13.6} _{-10.9}	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
51 ± 14	711	ABDALLAH	03H DLPH	$91.2 e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp + X$
61 ± 8	3651	NICHITIU	02 OBLX	$0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
38 ± 9 ± 6	20k	ADAMS	01B B852	18 GeV $\pi^- p \rightarrow K^+ K^- \pi^0 n$
58 ± 4		BARBERIS	97C OMEG	450 $pp \rightarrow pp K_S^0 K^\pm \pi^\mp$
45 ± 10		BERTIN	97 OBLX	0.0 $\bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
58 ± 10		⁶ ARMSTRONG	92E OMEG	85,300 $\pi^+ p, pp \rightarrow \pi^+ p, pp (K \bar{K} \pi)$
129 ± 41		⁷ AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
68 ⁺²⁹ ₋₁₈ ± 8 ⁺⁸ ₋₉	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
42 ± 22	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
40 ⁺¹⁷ ₋₁₃ ± 5	111	BECKER	87 MRK3	$e^+ e^- \rightarrow \omega K \bar{K} \pi$
35 ⁺⁴⁷ ₋₂₀	13	AIHARA	86C TPC	$e^+ e^- \rightarrow e^+ e^- K \bar{K} \pi$
47 ± 10		CHAUVAT	84 SPEC	ISR 31.5 pp
62 ± 14		BROMBERG	80 SPEC	100 $\pi^- p \rightarrow K \bar{K} \pi X$
40 ± 15	221	DIONISI	80 HBC	4 $\pi^- p \rightarrow K \bar{K} \pi n$
60 ± 20		DAHL	67 HBC	1.6-4.2 $\pi^- p$

NODE=M006W
 NODE=M006W
 NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

68.7 ± 2.9

⁸ SOSA 99 SPEC $pp \rightarrow p_{slow} (K_S^0 K^+ \pi^-) p_{fast}$

58.8 ± 3.3		⁸ SOSA	99	SPEC	$pp \rightarrow p_{\text{slow}} (K_S^0 K^- \pi^+) p_{\text{fast}}$
90 ± 25		PROKOSHKIN	97B	GAM4	$100 \pi^- p \rightarrow \eta \pi^0 \pi^0 n$
58 ± 8	389	ARMSTRONG	89	OMEG	$300 pp \rightarrow K \bar{K} \pi pp$
62 ± 5	1520	ARMSTRONG	84	OMEG	$85 \pi^+ p, pp \rightarrow (\pi^+, p)(K \bar{K} \pi)p$
~ 50		BITYUKOV	84	SPEC	$32 K^- p \rightarrow K^+ K^- \pi^0 \gamma$

OCCUR=2


⁶ This result supersedes ARMSTRONG 84, ARMSTRONG 89.

⁷ From fit to the $K^*(892)K 1^{++}$ partial wave.

⁸ No systematic error given.

NODE=M006W;LINKAGE=C
 NODE=M006W;LINKAGE=B
 NODE=M006W;LINKAGE=N1

WEIGHTED AVERAGE
 56.7±3.3 (Error scaled by 1.3)



f₁(1420) DECAY MODES

NODE=M006215;NODE=M006

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K} \pi$	seen
Γ_2 $K \bar{K}^*(892) + c.c.$	seen
Γ_3 $\eta \pi \pi$	possibly seen
Γ_4 $a_0(980) \pi$	
Γ_5 $\pi \pi \rho$	
Γ_6 4π	
Γ_7 $\rho^0 \gamma$	
Γ_8 $\phi \gamma$	seen

DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=4
 DESIG=3
 DESIG=6
 DESIG=8
 DESIG=9;OUR EST;→ UNCHECKED ←

f₁(1420) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M006220

$\Gamma(K \bar{K} \pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$

NODE=M006G2
 NODE=M006G2

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.4 OUR AVERAGE					
3.2±0.6±0.7		133	9,10 ACHARD	07 L3	$183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$
3.0±0.9±0.7			11,12 BEHREND	89 CELL	$e^+ e^- \rightarrow e^+ e^- K_S^0 K \pi$
2.3 ^{+1.0} _{-0.9} ±0.8			HILL	89 JADE	$e^+ e^- \rightarrow e^+ e^- K^\pm K_S^0 \pi^\mp$
1.3±0.5±0.3			AIHARA	88B TPC	$e^+ e^- \rightarrow e^+ e^- K^\pm K_S^0 \pi^\mp$
1.6±0.7±0.3			11,13 GIDAL	87B MRK2	$e^+ e^- \rightarrow e^+ e^- K \bar{K} \pi$
<8.0	95		JENNI	83 MRK2	$e^+ e^- \rightarrow e^+ e^- K \bar{K} \pi$

••• We do not use the following data for averages, fits, limits, etc. •••

⁹ From a fit with a width fixed at 55 MeV.

¹⁰ The form factor parameter from the fit is 926 ± 78 MeV.

¹¹ Assume a ρ -pole form factor.

¹² A ϕ -pole form factor gives considerably smaller widths.

¹³ Published value divided by 2.

NODE=M006G2;LINKAGE=CH
 NODE=M006G2;LINKAGE=CR
 NODE=M006G2;LINKAGE=A
 NODE=M006G2;LINKAGE=D
 NODE=M006G2;LINKAGE=B

$f_1(1420)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892)+c.c.)/\Gamma(K\bar{K}\pi)$

Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.76 ± 0.06	BROMBERG 80	SPEC	100 $\pi^- p \rightarrow K\bar{K}\pi X$
0.86 ± 0.12	DIONISI 80	HBC	4 $\pi^- p \rightarrow K\bar{K}\pi n$

NODE=M006225

NODE=M006R1
 NODE=M006R1

$\Gamma(\pi\pi\rho)/\Gamma(K\bar{K}\pi)$

Γ_5/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.3	95	CORDEN 78	OMEG	12-15 $\pi^- p$
<2.0		DAHL 67	HBC	1.6-4.2 $\pi^- p$

NODE=M006R2
 NODE=M006R2

$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$

Γ_3/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	95	ARMSTRONG 91B	OMEG	300 $p p \rightarrow p p \eta \pi^+ \pi^-$
1.35 ± 0.75		KOPKE 89	MRK3	$J/\psi \rightarrow \omega \eta \pi \pi (K\bar{K}\pi)$
<0.6	90	GIDAL 87	MRK2	$e^+ e^- \rightarrow e^+ e^- \eta \pi^+ \pi^-$
<0.5	95	CORDEN 78	OMEG	12-15 $\pi^- p$
1.5 ± 0.8		DEFOIX 72	HBC	0.7 $\bar{p} p$

NODE=M006R3
 NODE=M006R3

$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$

Γ_4/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.1	90	PROKOSHKIN 97B	GAM4	100 $\pi^- p \rightarrow \eta \pi^0 \pi^0 n$
not seen in either mode		ANDO 86	SPEC	8 $\pi^- p$
not seen in either mode		CORDEN 78	OMEG	12-15 $\pi^- p$
0.4 ± 0.2		DEFOIX 72	HBC	0.7 $\bar{p} p \rightarrow 7\pi$

NODE=M006R4
 NODE=M006R4

$\Gamma(4\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$

Γ_6/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.90	95	DIONISI 80	HBC	4 $\pi^- p$

NODE=M006R5
 NODE=M006R5

$\Gamma(K\bar{K}\pi)/[\Gamma(K\bar{K}^*(892)+c.c.)+\Gamma(a_0(980)\pi)]$

$\Gamma_1/(\Gamma_2+\Gamma_4)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.65 ± 0.27	¹⁴ DIONISI 80	HBC	4 $\pi^- p$

NODE=M006R6
 NODE=M006R6

¹⁴ Calculated using $\Gamma(K\bar{K})/\Gamma(\eta\pi) = 0.24 \pm 0.07$ for $a_0(980)$ fractions.

NODE=M006R6;LINKAGE=C

$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}^*(892)+c.c.)$

Γ_4/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.042 ± 0.014 OUR AVERAGE				
0.44 ± 0.19		ABLIKIM 21U	BES3	$D_s^+ \rightarrow f_1(1420)\pi^+$
0.04 ± 0.01 ± 0.01		BARBERIS 98C	OMEG	450 $p p \rightarrow p_f f_1(1420) p_s$
<0.04	68	ARMSTRONG 84	OMEG	85 $\pi^+ p$

NODE=M006R7
 NODE=M006R7

$\Gamma(4\pi)/\Gamma(K\bar{K}\pi)$

Γ_6/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.62	95	ARMSTRONG 89G	OMEG	85 $\pi p \rightarrow 4\pi X$

NODE=M006R8
 NODE=M006R8

$\Gamma(\rho^0\gamma)/\Gamma_{total}$

Γ_7/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.08	95	¹⁵ ARMSTRONG 92C	SPEC	300 $p p \rightarrow p p \pi^+ \pi^- \gamma$

NODE=M006R9
 NODE=M006R9

¹⁵ Using the data on the $\bar{K}K\pi$ mode from ARMSTRONG 89.

NODE=M006R9;LINKAGE=A

$\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_1
<0.02	95	BARBERIS	98C	OMEG 450 pp \rightarrow $p_f f_1(1420) p_S$	

NODE=M006R10
 NODE=M006R10

 $\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_1
0.003 \pm 0.001 \pm 0.001	BARBERIS	98C	OMEG 450 pp \rightarrow $p_f f_1(1420) p_S$	

NODE=M006R11
 NODE=M006R11

 $f_1(1420)$ REFERENCES

Author	Year	Journal	Page	Collaboration	REFID
ABLIKIM	23M	JHEP	2303 121	M. Ablikim <i>et al.</i> (BESIII Collab.)	REFID=62055
ABLIKIM	21U	PR	D104 032011	M. Ablikim <i>et al.</i> (BESIII Collab.)	REFID=61155
ACHARD	07	JHEP	0703 018	P. Achard <i>et al.</i> (L3 Collab.)	REFID=51698
ABDALLAH	03H	PL	B569 129	J. Abdallah <i>et al.</i> (DELPHI Collab.)	REFID=49548
NICHITIU	02	PL	B545 261	F. Nichitiu <i>et al.</i> (OBELIX Collab.)	REFID=48848
ADAMS	01B	PL	B516 264	G.S. Adams <i>et al.</i> (BNL E852 Collab.)	REFID=49649
SOSA	99	PRL	83 913	M. Sosa <i>et al.</i>	REFID=46937
BARBERIS	98C	PL	B440 225	D. Barberis <i>et al.</i> (WA 102 Collab.)	REFID=46346
BARBERIS	97C	PL	B413 225	D. Barberis <i>et al.</i> (WA 102 Collab.)	REFID=45759
BERTIN	97	PL	B400 226	A. Bertin <i>et al.</i> (OBELIX Collab.)	REFID=45417
PROKOSHKIN	97B	PD	42 298	Yu.D. Prokoshkin, S.A. Sadovsky	REFID=45549
Translated from DANS 354 751.					
ARMSTRONG	92C	ZPHY	C54 371	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+)	REFID=42097
ARMSTRONG	92E	ZPHY	C56 29	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+) JPC	REFID=43173
AUGUSTIN	92	PR	D46 1951	J.E. Augustin, G. Cosme (DM2 Collab.)	REFID=41584
ARMSTRONG	91B	ZPHY	C52 389	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+)	REFID=41862
BAI	90C	PRL	65 2507	Z. Bai <i>et al.</i> (Mark III Collab.)	REFID=41578
ARMSTRONG	89	PL	B221 216	T.A. Armstrong <i>et al.</i> (CERN, CDEF, BIRM+) JPC	REFID=40729
ARMSTRONG	89G	ZPHY	C43 55	T.A. Armstrong <i>et al.</i> (CERN, BIRM, BARI+)	REFID=40930
BEHREND	89	ZPHY	C42 367	H.J. Behrend <i>et al.</i> (CELLO Collab.)	REFID=40732
HILL	89	ZPHY	C42 355	P. Hill <i>et al.</i> (JADE Collab.) JP	REFID=40741
KOPKE	89	PRPL	174 67	L. Kopke <i>et al.</i> (CERN)	REFID=41863
AIHARA	88B	PL	B209 107	H. Aihara <i>et al.</i> (TPC-2 γ Collab.)	REFID=40572
BECKER	87	PRL	59 186	J.J. Becker <i>et al.</i> (Mark III Collab.) JP	REFID=40015
GIDAL	87	PRL	59 2012	G. Gidal <i>et al.</i> (LBL, SLAC, HARV)	REFID=40223
GIDAL	87B	PRL	59 2016	G. Gidal <i>et al.</i> (LBL, SLAC, HARV)	REFID=40224
AIHARA	86C	PRL	57 2500	H. Aihara <i>et al.</i> (TPC-2 γ Collab.) JP	REFID=21326
ANDO	86	PRL	57 1296	A. Ando <i>et al.</i> (KEK, KYOT, NIRS, SAGA+)	REFID=20891
ARMSTRONG	84	PL	146B 273	T.A. Armstrong <i>et al.</i> (ATHU, BARI, BIRM+) JP	REFID=20929
BITYUKOV	84	SJNP	39 735	S. Bitukov <i>et al.</i> (SERP)	REFID=45856
Translated from YAF 39 1165.					
CHAUVAT	84	PL	148B 382	P. Chauvat <i>et al.</i> (CERN, CLER, UCLA+)	REFID=20932
JENNI	83	PR	D27 1031	P. Jenni <i>et al.</i> (SLAC, LBL)	REFID=20304
BROMBERG	80	PR	D22 1513	C.M. Bromberg <i>et al.</i> (CIT, FNAL, ILLC+)	REFID=20922
DIONISI	80	NP	B169 1	C. Dionisi <i>et al.</i> (CERN, MADR, CDEF+) IJP	REFID=20924
CORDEN	78	NP	B144 253	M.J. Corden <i>et al.</i> (BIRM, RHEL, TELA+)	REFID=20452
DEFOIX	72	NP	B44 125	C. Defoix <i>et al.</i> (CDEF, CERN)	REFID=20435
DAHL	67	PR	163 1377	O.I. Dahl <i>et al.</i> (LRL) IJP	REFID=20321
Also		PRL	14 1074	D.H. Miller <i>et al.</i> (LRL, UCB)	REFID=21291

NODE=M006

$\omega(1420)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

See also the $\omega(1650)$ particle listing.

NODE=M125

NODE=M125

NODE=M125M

NODE=M125M

→ UNCHECKED ←

 $\omega(1420)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1410± 60 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1418± 30± 10	824	¹ AKHMETSHIN 17A	CMD3	1.4-2.0 $e^+e^- \rightarrow \omega\eta$
1470± 50	13.1k	² AULCHENKO 15A	SND	1.05-1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1382± 23± 70		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1350± 20± 20		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1400± 50± 130	1.2M	³ ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1450± 10		⁴ HENNER 02	RVUE	1.2-2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1373± 70	177	⁵ AKHMETSHIN 00D	CMD2	1.2-1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
1370± 25	5095	ANISOVICH 00H	SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
1400 ⁺¹⁰⁰ ₋₂₀₀		⁶ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
~ 1400		⁷ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
~ 1460		⁸ ACHASOV 98H	RVUE	$e^+e^- \rightarrow K^+K^-$
1440± 70		⁹ CLEGG 94	RVUE	
1419± 31	315	¹⁰ ANTONELLI 92	DM2	1.34-2.4 $e^+e^- \rightarrow \rho\pi$
¹ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.				
² From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.				
³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.				
⁴ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.				
⁵ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.				
⁶ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.				
⁷ Using the data from ANTONELLI 92.				
⁸ Using the data from IVANOV 81 and BISELLO 88B.				
⁹ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.				
¹⁰ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.				

NODE=M125M;LINKAGE=F

NODE=M125M;LINKAGE=E

NODE=M125M;LINKAGE=VH

NODE=M125M;LINKAGE=AB

NODE=M125M;LINKAGE=KL

NODE=M125M;LINKAGE=L1

NODE=M125M;LINKAGE=L2

NODE=M125M;LINKAGE=L3

NODE=M125M;LINKAGE=AD

NODE=M125M;LINKAGE=B

 $\omega(1420)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
290± 190 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
440± 125	267	¹ ACHASOV 20B	SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
104± 35± 10	824	² AKHMETSHIN 17A	CMD3	1.4-2.0 $e^+e^- \rightarrow \omega\eta$
880± 170	13.1k	³ AULCHENKO 15A	SND	1.05-1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
480± 180		⁴ ACHASOV 10D	SND	1.075-2.0 $e^+e^- \rightarrow \pi^0\gamma$
130± 50± 100		AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
450± 70± 70		AUBERT,B 04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
870 ⁺⁵⁰⁰ ₋₃₀₀ ± 450	1.2M	⁵ ACHASOV 03D	RVUE	0.44-2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
199± 15		⁶ HENNER 02	RVUE	1.2-2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
188± 45	177	⁷ AKHMETSHIN 00D	CMD2	1.2-1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
360 ⁺¹⁰⁰ ₋₆₀	5095	ANISOVICH 00H	SPEC	0.0 $\rho\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$
240± 70		⁸ CLEGG 94	RVUE	
174± 59	315	⁹ ANTONELLI 92	DM2	1.34-2.4 $e^+e^- \rightarrow \rho\pi$

NODE=M125W

NODE=M125W

→ UNCHECKED ←

- ¹ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass of $\omega(1420)$ is fixed to the PDG 18 value of 1420 MeV.
- ² From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.
- ³ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.
- ⁴ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.
- ⁵ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.
- ⁶ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.
- ⁷ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.
- ⁸ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.
- ⁹ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

NODE=M125W;LINKAGE=H

NODE=M125W;LINKAGE=F

NODE=M125W;LINKAGE=E

NODE=M125W;LINKAGE=G

NODE=M125W;LINKAGE=VH

NODE=M125W;LINKAGE=AB

NODE=M125W;LINKAGE=KL

NODE=M125W;LINKAGE=AD

NODE=M125W;LINKAGE=B

 $\omega(1420)$ DECAY MODES

NODE=M125215;NODE=M125

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\omega\pi\pi$	seen
Γ_3 $\omega\eta$	
Γ_4 $b_1(1235)\pi$	seen
Γ_5 e^+e^-	seen
Γ_6 $\pi^0\gamma$	

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=7

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=6

 $\omega(1420)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M125230

$\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1/\Gamma \times \Gamma_5/\Gamma$
VALUE (units 10^{-6})	EVTS DOCUMENT ID TECN COMMENT

NODE=M125G3
NODE=M125G3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.73 ± 0.08	13.1k	¹ AULCHENKO	15A	SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.82 ± 0.05 ± 0.06		AUBERT,B	04N	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
0.65 ± 0.13 ± 0.21	1.2M	^{2,3} ACHASOV	03D	RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.625 ± 0.160		^{4,5} CLEGG	94	RVUE	
0.466 ± 0.178		^{6,7} ANTONELLI	92	DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi$

NODE=M125G3;LINKAGE=A

NODE=M125G;LINKAGE=AW

NODE=M125G;LINKAGE=VH

NODE=M125G;LINKAGE=AD

NODE=M125G;LINKAGE=SE

NODE=M125G;LINKAGE=A

NODE=M125G;LINKAGE=ES

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.

² Calculated by us from the cross section at the peak.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁵ From the partial and leptonic width given by the authors.

⁶ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

⁷ From the product of the leptonic width and partial branching ratio given by the authors.

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_2/\Gamma \times \Gamma_5/\Gamma$
VALUE (units 10^{-8})	DOCUMENT ID TECN COMMENT

NODE=M125G4
NODE=M125G4

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

19.7 ± 5.7	AUBERT	07AU	BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1.9 ± 1.9	¹ AKHMETSHIN 00D	CMD2	1.2–2.4 $e^+e^- \rightarrow \omega\pi^+\pi^-$	

¹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

NODE=M125G;LINKAGE=KL

$\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.5 ± 0.6	267	¹ ACHASOV	20B SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
$2.1^{+1.0}_{-0.8}$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
$5.0 \pm 2.6 \pm 0.3$	824	² AKHMETSHIN	17A CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$
$1.6^{+0.9}_{-0.7}$	898	³ ACHASOV	16B SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$

NODE=M125G6
 NODE=M125G6

¹ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass of $\omega(1420)$ is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of $\omega(1420)$ to the PDG 18 value of 220 MeV results in $(3.0 \pm 1.6) \times 10^{-8}$ measurement.

NODE=M125G6;LINKAGE=C

² From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1420) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1420) \rightarrow e^+e^-) = 5.3 \pm 1.6$ eV.

NODE=M125G6;LINKAGE=B

³ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass and the width of $\omega(1420)$ are fixed to the 2014 edition (PDG 14) of this review.

NODE=M125G6;LINKAGE=A

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.23 ± 0.14	¹ ACHASOV	10D SND	$1.075-2.0 e^+e^- \rightarrow \pi^0\gamma$
$2.03^{+0.70}_{-0.75}$	² AKHMETSHIN	05 CMD2	$0.60-1.38 e^+e^- \rightarrow \pi^0\gamma$

NODE=M125G5
 NODE=M125G5

¹ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.

NODE=M125G5;LINKAGE=A

² Using 1420 MeV and 220 MeV for the $\omega(1420)$ mass and width.

NODE=M125G5;LINKAGE=AK

 $\omega(1420)$ BRANCHING RATIOS

NODE=M125225

 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.301 ± 0.029 possibly seen	¹ HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
	AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$

NODE=M125R2
 NODE=M125R2

 $\Gamma(\omega\pi\pi)/\Gamma(b_1(1235)\pi)$ Γ_2/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.60 ± 0.16	5095	ANISOVICH	00H SPEC	$0.0 p\bar{p} \rightarrow \omega\pi^0\pi^0\pi^0$

NODE=M125R1
 NODE=M125R1

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
seen	ACHASOV	20A SND	$1.15-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.699 ± 0.029	¹ HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M125R3
 NODE=M125R3

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 6.6	1.2M	^{2,3} ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
23 ± 1		¹ HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

NODE=M125R4
 NODE=M125R4

¹ Assuming that the $\omega(1420)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

NODE=M125R;LINKAGE=AC

² Calculated by us from the cross section at the peak.

NODE=M125R;LINKAGE=AW

³ Assuming that the $\omega(1420)$ decays into $\rho\pi$ only.

NODE=M125R;LINKAGE=GS

$\omega(1420)$ REFERENCES

ACHASOV	20A	EPJ C80 993	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60923
ACHASOV	20B	EPJ C80 1008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60924
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59887
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=58239
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57537
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56843
		Translated from ZETF 148 34.			
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
AKHMETSHIN	05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>		REFID=49177
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
ANISOVICH	00H	PL B485 341	A.V. Anisovich <i>et al.</i>		REFID=47948
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)	REFID=21586
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553

NODE=M125

 $f_2(1430)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry lists nearby peaks observed in the D wave of the $K\bar{K}$ and $\pi^+\pi^-$ systems. Needs confirmation.

NODE=M066

 $f_2(1430)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1430 OUR ESTIMATE			
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$1440 \pm 11 \pm 3$	LEES	21A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
1453 ± 4	¹ VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1421 ± 5	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
1480 ± 50	AKESSON	86	SPEC $pp \rightarrow pp \pi^+ \pi^-$
1436^{+26}_{-16}	DAUM	84	CNTR $17-18 \pi^- p \rightarrow K^+ K^- n$
1412 ± 3	DAUM	84	CNTR $63 \pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
1439^{+5}_{-6}	² BEUSCH	67	OSPK $5,7,12 \pi^- p \rightarrow K_S^0 K_S^0 n$
¹ $J^{PC} = 0^{++}$ or 2^{++} .			
² Not seen by WETZEL 76.			

NODE=M066M1

NODE=M066M1

→ UNCHECKED ←

OCCUR=2

NODE=M066M;LINKAGE=AC

NODE=M066M;LINKAGE=C

 $f_2(1430)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$46 \pm 15 \pm 5$	LEES	21A	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
13 ± 5	³ VLADIMIRSK...01	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
30 ± 9	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
150 ± 50	AKESSON	86	SPEC $pp \rightarrow pp \pi^+ \pi^-$
81^{+56}_{-29}	DAUM	84	CNTR $17-18 \pi^- p \rightarrow K^+ K^- n$
14 ± 6	DAUM	84	CNTR $63 \pi^- p \rightarrow K_S^0 K_S^0 n, K^+ K^- n$
43^{+17}_{-18}	⁴ BEUSCH	67	OSPK $5,7,12 \pi^- p \rightarrow K_S^0 K_S^0 n$
³ $J^{PC} = 0^{++}$ or 2^{++} .			
⁴ Not seen by WETZEL 76.			

NODE=M066W1

NODE=M066W1

OCCUR=2

NODE=M066W;LINKAGE=AC

NODE=M066W;LINKAGE=C

$f_2(1430)$ DECAY MODES

NODE=M066215;NODE=M066

Mode	
Γ_1	$K\bar{K}$
Γ_2	$\pi\pi$

DESIG=1

DESIG=2

 $f_2(1430)$ REFERENCES

NODE=M066

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
VLADIMIRSK...	01	PAN 64 1895 Translated from YAF 64 1979.	V.V. Vladimisky <i>et al.</i>	
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
DAUM	84	ZPHY C23 339	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP
WETZEL	76	NP B115 208	W. Wetzel <i>et al.</i>	(ETH, CERN, LOIC)
BEUSCH	67	PL 25B 357	W. Beusch <i>et al.</i>	(ETH, CERN)

REFID=61442
REFID=48571REFID=40268
REFID=21123
REFID=21372
REFID=20362
REFID=20320

NODE=M149

 $a_0(1450)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M149

 $a_0(1450)$ T-MATRIX POLE \sqrt{s}

NODE=M149PP

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M149PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1290–1500) – i (30–140) OUR ESTIMATE			
$(1302.1 \pm 1.1 \pm 3.9) - i$ (56.2 \pm 0.7 \pm 1.7)	¹ ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1515 \pm 30) - i$ (115 \pm 18)	ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1432 \pm 13 \pm 25) - i$ (98 \pm 5 \pm 5)	² BUGG	08A	RVUE $\bar{p}p$
$(1441^{+40}_{-15}) - i$ (55 \pm 7)	³ BAKER	03	SPEC $\bar{p}p \rightarrow \omega \pi^+ \pi^- \pi^0$
$(1303 \pm 16) - i$ (46 \pm 8)	⁴ BARGIOTTI	03	OBLX $\bar{p}p$
$(1296 \pm 10) - i$ (41 \pm 11)	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$(1565 \pm 30) - i$ (146 \pm 20)	ANISOVICH	98B	RVUE Compilation
$(1470 \pm 25) - i$ (132 \pm 15)	⁵ AMSLER	95D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$

NODE=M149PP

→ UNCHECKED ←

¹ T-matrix pole, 2 poles, 2 channels ($\pi\eta, K\bar{K}$).² Using data from AMSLER 94D, ABELE 98, and BAKER 03. Supersedes BUGG 94.³ From the pole position of a fitted Breit-Wigner amplitude.⁴ Coupled channel analysis of $\pi^+ \pi^- \pi^0, K^+ K^- \pi^0,$ and $K^\pm K_S^0 \pi^\mp$.⁵ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

NODE=M149PP;LINKAGE=A

NODE=M149PP;LINKAGE=G

NODE=M149PP;LINKAGE=C

NODE=M149PP;LINKAGE=D

NODE=M149PP;LINKAGE=F

 $a_0(1450)$ MASS

NODE=M149M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1439 \pm 34 OUR AVERAGE		Error includes scale factor of 1.8.		
1480 \pm 30		ABELE	98	CBAR $0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
1410 \pm 25		ETKIN	82C	MPS $23 \pi^- p \rightarrow n 2K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1458 \pm 14 \pm 15	190k	¹ AAJ	16N	LHCB $D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
$1316.8^{+0.7+24.7}_{-1.0-4.6}$		² UEHARA	09A	BELL $\gamma\gamma \rightarrow \pi^0 \eta$
1477 \pm 10	80k	³ UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
1290 \pm 10		⁴ BERTIN	98B	OBLX $0.0 \bar{p}p \rightarrow K^\pm K_S^0 \pi^\mp$
1450 \pm 40		AMSLER	94D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \eta$
\sim 1300		MARTIN	78	SPEC $10 K^\pm p \rightarrow K_S^0 \pi p$
1255 \pm 5		⁵ CASON	76	

NODE=M149M

- ¹ Using a model with Gaussian constraints to the PDG averaged values .
² May be a different state.
³ Statistical error only.
⁴ Not confirmed by BUGG 08A.
⁵ Isospin 0 not excluded.

NODE=M149M;LINKAGE=A
 NODE=M149M;LINKAGE=UE
 NODE=M149M;LINKAGE=ST
 NODE=M149M;LINKAGE=BE
 NODE=M149M;LINKAGE=CC

$a_0(1450)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
258 ± 14	OUR AVERAGE			
265 ± 15		ABELE	98 CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
230 ± 30		ETKIN	82C MPS	23 $\pi^- p \rightarrow n 2K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
282 ± 12 ± 13	190k	¹ AAIJ	16N LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
65.0 ⁺ ₋ 2.1+99.1 5.4-32.6		² UEHARA	09A BELL	$\gamma\gamma \rightarrow \pi^0 \eta$
267 ± 11	80k	³ UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta \eta \pi^0$
80 ± 5		⁴ BERTIN	98B OBLX	0.0 $\bar{p}p \rightarrow K^\pm K_S \pi^\mp$
270 ± 40		AMSLER	94D CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
~ 250		MARTIN	78 SPEC	10 $K^\pm p \rightarrow K_S^0 \pi p$
79 ± 10		⁵ CASON	76	

NODE=M149W

NODE=M149W

- ¹ Using a model with Gaussian constraints to the PDG averaged values .
² May be a different state.
³ Statistical error only.
⁴ Not confirmed by BUGG 08A.
⁵ Isospin 0 not excluded.

NODE=M149W;LINKAGE=A
 NODE=M149W;LINKAGE=UE
 NODE=M149W;LINKAGE=ST
 NODE=M149W;LINKAGE=BE
 NODE=M149W;LINKAGE=CC

$a_0(1450)$ DECAY MODES

Branching fractions are given relative to the one **DEFINED AS 1**.

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi \eta$	0.093 ± 0.020
Γ_2 $\pi \eta'(958)$	0.033 ± 0.017
Γ_3 $K \bar{K}$	0.082 ± 0.028
Γ_4 $\omega \pi \pi$	DEFINED AS 1
Γ_5 $a_0(980) \pi \pi$	seen
Γ_6 $\gamma \gamma$	seen

NODE=M149215;NODE=M149

NODE=M149

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

$a_0(1450)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\pi\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1 \Gamma_6 / \Gamma$
432 ± 6 ⁺ ₋ 1073 256	¹ UEHARA	09A BELL	$\gamma\gamma \rightarrow \pi^0 \eta$	

NODE=M149225

NODE=M149G01
 NODE=M149G01

- ¹ May be a different state.

NODE=M149G01;LINKAGE=UE

$a_0(1450)$ BRANCHING RATIOS

$\Gamma(\pi\eta'(958))/\Gamma(\pi\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.35 ± 0.16	¹ ABELE	98 CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.43 ± 0.19	ABELE	97C CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta'$	

NODE=M149220

NODE=M149R1
 NODE=M149R1

- ¹ Using $\pi^0 \eta$ from AMSLER 94D.

NODE=M149R1;LINKAGE=A

$\Gamma(K\bar{K})/\Gamma(\pi\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.88 ± 0.23	¹ ABELE	98 CBAR	0.0 $\bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.887 ± 0.041 ± 0.97	² ALBRECHT	20 RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$	

NODE=M149R2
 NODE=M149R2

- ¹ Using $\pi^0 \eta$ from AMSLER 94D.
² Residues from T-matrix pole, 2 poles, 2 channels ($\pi\eta, K\bar{K}$).

NODE=M149R2;LINKAGE=A
 NODE=M149R2;LINKAGE=B

$\Gamma(\omega\pi\pi)/\Gamma(\pi\eta)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
10.7±2.3	35280	¹ BAKER	03 SPEC	$\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

¹ Using results on $\bar{p}p \rightarrow a_0(1450)^0\pi^0$, $a_0(1450) \rightarrow \eta\pi^0$ from ABELE 96C and assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.

NODE=M149R3
 NODE=M149R3

NODE=M149R;LINKAGE=PP

 $\Gamma(a_0(980)\pi\pi)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BUGG	08A RVUE	$\bar{p}p$

NODE=M149R01
 NODE=M149R01

 $\Gamma(a_0(980)\pi\pi)/\Gamma(\pi\eta)$ Γ_5/Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
•••	ANISOVICH	01 RVUE	0	$\bar{p}p \rightarrow \eta 2\pi^+ 2\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

≤ 4.3

NODE=M149R02
 NODE=M149R02

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ UEHARA	09A BELL	$\gamma\gamma \rightarrow \pi^0\eta$

¹ May be a different state.

NODE=M149R03
 NODE=M149R03

NODE=M149R03;LINKAGE=UE

 $a_0(1450)$ REFERENCES

ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09A	PR D80 032001	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53002
BUGG	08A	PR D78 074023	D.V. Bugg	(LOQM)	REFID=52578
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>		REFID=49414
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	01	NP A690 567	A.V. Anisovich <i>et al.</i>		REFID=48308
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BERTIN	98B	PL B434 180	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=46351
ABELE	97C	PL B404 179	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45531
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20391
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=22446
CASON	76	PRL 36 1485	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=21064

NODE=M149

$\rho(1450)$

$$I^G(J^{PC}) = 1^+(1^-)$$

NODE=M105

 $\rho(1450)$ MASS

NODE=M105205

 $\rho(1450)$ MASS

VALUE (MeV)

DOCUMENT ID

1465±25 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

NODE=M105M0

NODE=M105M0

→ UNCHECKED ←

 $\eta\rho^0$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1506±11	13.4k	¹ GRIBANOV 20	CMD3	1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1500±10	7.4k	² ACHASOV 18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
1497±14		³ AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$
1421±15		⁴ AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1470±20		ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1446±10		FUKUI 88	SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

NODE=M105M1

NODE=M105M1

¹ Mass and width of the $\rho(770)$ fixed at 775 and 149 MeV, respectively; solution 2 of model 2, $\eta \rightarrow \gamma\gamma$ decays used.

NODE=M105M1;LINKAGE=B

² From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

NODE=M105M1;LINKAGE=A

³ Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$.

NODE=M105M;LINKAGE=SW

⁴ Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

NODE=M105M1;LINKAGE=KL

 $\omega\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1510±7	10.2k	¹ ACHASOV 16D	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1544±22 ⁺¹¹ ₋₄₆	821	² MATVIENKO 15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
1491±19	7815	³ ACHASOV 13	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1582±17±25	2382	⁴ AKHMETSHIN 03B	CMD2	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1349±25 ⁺¹⁰ ₋₅	341	⁵ ALEXANDER 01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
1523±10		⁶ EDWARDS 00A	CLE2	$\tau^- \rightarrow \omega\pi^- \nu_\tau$
1463±25		⁷ CLEGG 94	RVUE	
1250		⁸ ASTON 80C	OMEG	20–70 $\gamma p \rightarrow \omega\pi^0 p$
1290±40		⁸ BARBER 80C	SPEC	3–5 $\gamma p \rightarrow \omega\pi^0 p$

NODE=M105M3

NODE=M105M3

OCCUR=2

¹ From a phenomenological model based on vector meson dominance with interfering $\rho(770)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

NODE=M105M3;LINKAGE=D

² Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.

NODE=M105M3;LINKAGE=C

³ From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M105M3;LINKAGE=AC

⁴ Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the $\omega\pi^0$ and $\pi^+\pi^-$ mass dependence of the total width. $\rho(1700)$ mass and width fixed at 1700 MeV and 240 MeV, respectively.

NODE=M105M3;LINKAGE=HK

⁵ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming the $\omega\pi^-$ mass dependence for the total width.

NODE=M105M3;LINKAGE=3Z

⁶ Mass-independent width parameterization. $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.

NODE=M105M;LINKAGE=E1

⁷ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

NODE=M105M3;LINKAGE=B

⁸ Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.

NODE=M105M3;LINKAGE=A

 4π MODE

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1435±40		ABELE 01B	CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0\pi^+$
1350±50		ACHASOV 97	RVUE	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1449±4		¹ ARMSTRONG 89E	OMEG	300 $pp \rightarrow pp2(\pi^+\pi^-)$

NODE=M105M6

NODE=M105M6

¹ Not clear whether this observation has $l=1$ or 0.

NODE=M105M6;LINKAGE=A

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1326.35 ± 3.46		¹ BARTOS 17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1342.31 ± 46.62		² BARTOS 17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1373.83 ± 11.37		³ BARTOS 17A	RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1429 ± 41	20k	⁴ LEES 17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
1350 ± 20	$\begin{matrix} +20 \\ -30 \end{matrix}$ 63.5k	⁵ ABRAMOWICZ12	ZEUS	$e p \rightarrow e\pi^+\pi^- p$
1493 ± 15		⁶ LEES 12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
1446 ± 7	± 28 5.4M	^{7,8} FUJIKAWA 08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1328 ± 15		⁹ SCHAEEL 05C	ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1406 ± 15	87k	^{7,10} ANDERSON 00A	CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
~ 1368		¹¹ ABELE 99C	CBAR	$0.0 \bar{p}d \rightarrow \pi^+\pi^-\pi^- p$
1348 ± 33		BERTIN 98	OBLX	$0.05-0.405 \bar{n}p \rightarrow$ $2\pi^+\pi^-$
1411 ± 14		¹² ABELE 97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
1370 $\begin{matrix} +90 \\ -70 \end{matrix}$		ACHASOV 97	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1359 ± 40		¹⁰ BERTIN 97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1282 ± 37		BERTIN 97D	OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+2\pi^-$
1424 ± 25		BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
1265.5 ± 75.3		DUBNICKA 89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
1292 ± 17		¹³ KURDADZE 83	OLYA	$0.64-1.4 e^+e^- \rightarrow$ $\pi^+\pi^-$

¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

² Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

³ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.

⁴ From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.

⁵ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.

⁶ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

⁷ From the GOUNARIS 68 parametrization of the pion form factor.

⁸ $|F_\pi(0)|^2$ fixed to 1.

⁹ From the combined fit of the τ^- data from ANDERSON 00A and SCHAEEL 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. $\rho(1700)$ mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.

¹⁰ $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV, respectively.

¹¹ $\rho(1700)$ mass and width fixed at 1780 MeV and 275 MeV respectively.

¹² T-matrix pole.

¹³ Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

 $K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1208 ± 8 ± 9	190k	¹ AAJ 16N	LHCB		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
1422.8 ± 6.5	27k	² ABELE 99D	CBAR ±		$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$

¹ Using the GOUNARIS 68 parameterization with fixed width.

² K-matrix pole. Isospin not determined, could be $\omega(1420)$.

 $K\bar{K}^*(892) + c.c.$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1505 ± 19 ± 7	AUBERT 08S	BABR 10.6	$e^+e^- \rightarrow K\bar{K}^*(892)\gamma$

 $m_{\rho(1450)^0} - m_{\rho(1450)^\pm}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
-31.53 ± 47.99	¹ BARTOS 17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M105M5
NODE=M105M5

OCCUR=2

OCCUR=2

NODE=M105M5;LINKAGE=D

NODE=M105M5;LINKAGE=E

NODE=M105M5;LINKAGE=F

NODE=M105M5;LINKAGE=B

NODE=M105M5;LINKAGE=AB

NODE=M105M5;LINKAGE=LE

NODE=M105M5;LINKAGE=1K

NODE=M105M5;LINKAGE=FU

NODE=M105M5;LINKAGE=SC

NODE=M105M5;LINKAGE=A

NODE=M105M5;LINKAGE=C5

NODE=M105M5;LINKAGE=QQ

NODE=M105M5;LINKAGE=KD

NODE=M105M7

NODE=M105M7

NODE=M105M7;LINKAGE=A

NODE=M105M7;LINKAGE=AN

NODE=M105M8

NODE=M105M8

NODE=M105DM

NODE=M105DM

NODE=M105DM;LINKAGE=A

$\rho(1450)$ WIDTH

NODE=M105210

 $\rho(1450)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

400± 60 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

••• We do not use the following data for averages, fits, limits, etc. •••

480±180	¹ ACHASOV	10D	SND	1.075–2.0 $e^+e^- \rightarrow \pi^0\gamma$
---------	----------------------	-----	-----	--

¹From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.

NODE=M105W0

NODE=M105W0

→ UNCHECKED ←

NODE=M105W0;LINKAGE=A

 $\eta\rho^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

321±27	13.4k	¹ GRIBANOV	20	CMD3	1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
280±20	7.4k	² ACHASOV	18	SND	1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
226±44		³ AKHMETSHIN 01B	CMD2		$e^+e^- \rightarrow \eta\gamma$
211±31		⁴ AKHMETSHIN 00D	CMD2		$e^+e^- \rightarrow \eta\pi^+\pi^-$
230±30		ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
60±15		FUKUI	88	SPEC	8.95 $\pi^-p \rightarrow \eta\pi^+\pi^-n$

¹Mass and width of the $\rho(770)$ fixed at 775 and 149 MeV, respectively; solution 2 of model 2, $\eta \rightarrow \gamma\gamma$ decays used.

²From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

³Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$.

⁴Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

NODE=M105W1

NODE=M105W1

NODE=M105W1;LINKAGE=B

NODE=M105W1;LINKAGE=A

NODE=M105W;LINKAGE=SW

NODE=M105W1;LINKAGE=KL

 $\omega\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

440± 40	10.2k	¹ ACHASOV	16D	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
303^{+31+69}_{-52-7}	821	² MATVIENKO	15	BELL	$\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
429± 42±10	2382	³ AKHMETSHIN 03B	CMD2		$e^+e^- \rightarrow \pi^0\pi^0\gamma$
547± 86 ⁺⁴⁶ ₋₄₅	341	⁴ ALEXANDER	01B	CLE2	$B \rightarrow D^{(*)}\omega\pi^-$
400± 35		⁵ EDWARDS	00A	CLE2	$\tau^- \rightarrow \omega\pi^-\nu_\tau$
311± 62		⁶ CLEGG	94	RVUE	
300		⁷ ASTON	80C	OMEG	20–70 $\gamma p \rightarrow \omega\pi^0 p$
320±100		⁷ BARBER	80C	SPEC	3–5 $\gamma p \rightarrow \omega\pi^0 p$

¹From a phenomenological model based on vector meson dominance with interfering $\rho(770)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

²Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.

³Using the data of AKHMETSHIN 03B and BISELLO 91B assuming the $\omega\pi^0$ and $\pi^+\pi^-$ mass dependence of the total width. $\rho(1700)$ mass and width fixed at 1700 MeV and 240 MeV, respectively.

⁴Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming the $\omega\pi^-$ mass dependence for the total width.

⁵Mass-independent width parameterization. $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV respectively.

⁶Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

⁷Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.

NODE=M105W3

NODE=M105W3

OCCUR=3

NODE=M105W3;LINKAGE=D

NODE=M105W3;LINKAGE=C

NODE=M105W3;LINKAGE=HK

NODE=M105W3;LINKAGE=3Z

NODE=M105W;LINKAGE=E1

NODE=M105W3;LINKAGE=B

NODE=M105W3;LINKAGE=A

4 π MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

325±100	ABELE	01B	CBAR	0.0 $\bar{p}n \rightarrow 2\pi^- 2\pi^0\pi^+$
---------	-------	-----	------	---

NODE=M105W66

NODE=M105W66

$\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
324.13 ± 12.01		1 BARTOS	17 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
492.17 ± 138.38		2 BARTOS	17A RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
340.87 ± 23.84		3 BARTOS	17A RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
576 ± 29	20k	4 LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
460 ± 30	$\begin{matrix} +40 \\ -45 \end{matrix}$ 63.5k	5 ABRAMOWICZ12	ZEUS	$ep \rightarrow e\pi^+\pi^-p$
427 ± 31		6 LEES	12G BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
434 ± 16	±60 5.4M	7,8 FUJIKAWA	08 BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
468 ± 41		9 SCHAEEL	05C ALEP	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
455 ± 41	87k	7,10 ANDERSON	00A CLE2	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
~ 374		11 ABELE	99C CBAR	$0.0 \bar{p}d \rightarrow \pi^+\pi^-\pi^-p$
275 ± 10		BERTIN	98 OBLX	$0.05-0.405 \bar{n}p \rightarrow \pi^+\pi^+\pi^-$
343 ± 20		12 ABELE	97 CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
310 ± 40		10 BERTIN	97C OBLX	$0.0 \bar{p}p \rightarrow \pi^+\pi^-\pi^0$
236 ± 36		BERTIN	97D OBLX	$0.05 \bar{p}p \rightarrow 2\pi^+2\pi^-$
269 ± 31		BISELLO	89 DM2	$e^+e^- \rightarrow \pi^+\pi^-$
391 ± 70		DUBNICKA	89 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
218 ± 46		13 KURDADZE	83 OLYA	$0.64-1.4 e^+e^- \rightarrow \pi^+\pi^-$

NODE=M105W5
 NODE=M105W5

OCCUR=2

- 1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
- 2 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
- 3 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.
- 4 From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.
- 5 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- 6 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
- 7 From the GOUNARIS 68 parametrization of the pion form factor.
- 8 $|F_\pi(0)|^2$ fixed to 1.
- 9 From the combined fit of the τ^- data from ANDERSON 00A and SCHAEEL 05C and e^+e^- data from the compilation of BARKOV 85, AKHMETSHIN 04, and ALOISIO 05. $\rho(1700)$ mass and width fixed at 1713 MeV and 235 MeV, respectively. Supersedes BARATE 97M.
- 10 $\rho(1700)$ mass and width fixed at 1700 MeV and 235 MeV, respectively.
- 11 $\rho(1700)$ mass and width fixed at 1780 MeV and 275 MeV respectively.
- 12 T-matrix pole.
- 13 Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.

NODE=M105W5;LINKAGE=C

NODE=M105W5;LINKAGE=D

NODE=M105W5;LINKAGE=E

NODE=M105W5;LINKAGE=B

NODE=M105W5;LINKAGE=AB

NODE=M105W5;LINKAGE=LE

NODE=M105W5;LINKAGE=1K

NODE=M105W5;LINKAGE=FU

NODE=M105W5;LINKAGE=SC

NODE=M105W5;LINKAGE=A

NODE=M105W5;LINKAGE=C5

NODE=M105W5;LINKAGE=QQ

NODE=M105W5;LINKAGE=KD

 $K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
410 ± 19 ± 35	190k	1 AAIJ	16N LHCB		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
146.5 ± 10.5	27k	2 ABELE	99D CBAR	±	$0.0 \bar{p}p \rightarrow K^+ K^- \pi^0$

- 1 Using the GOUNARIS 68 parameterization with fixed mass.
- 2 K-matrix pole. Isospin not determined, could be $\omega(1420)$.

NODE=M105W7
 NODE=M105W7

NODE=M105W7;LINKAGE=A

NODE=M105W7;LINKAGE=AN

 $K\bar{K}^*(892) + c.c.$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
418 ± 25 ± 4	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K\bar{K}^*(892)\gamma$

NODE=M105W8
 NODE=M105W8

 $\Gamma_{\rho(1450)^0} = \Gamma_{\rho(1450)^\pm}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
151.30 ± 140.42	1 BARTOS	17A RVUE	$e^+e^- \rightarrow \pi^+\pi^-$, $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$

- • • We do not use the following data for averages, fits, limits, etc. • • •

- 1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M105DW
 NODE=M105DW

NODE=M105DW;LINKAGE=A

$\rho(1450)$ DECAY MODES

NODE=M105215;NODE=M105

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 $\pi^+\pi^-$	seen
Γ_3 4π	seen
Γ_4 $\omega\pi$	
Γ_5 $a_1(1260)\pi$	
Γ_6 $h_1(1170)\pi$	
Γ_7 $\pi(1300)\pi$	
Γ_8 $\rho\rho$	
Γ_9 $\rho(\pi\pi)$ S-wave	
Γ_{10} e^+e^-	seen
Γ_{11} $\eta\rho$	seen
Γ_{12} $a_2(1320)\pi$	not seen
Γ_{13} $K\bar{K}$	seen
Γ_{14} K^+K^-	seen
Γ_{15} $K\bar{K}^*(892)+c.c.$	possibly seen
Γ_{16} $\pi^0\gamma$	seen
Γ_{17} $\eta\gamma$	seen
Γ_{18} $f_0(500)\gamma$	not seen
Γ_{19} $f_0(980)\gamma$	not seen
Γ_{20} $f_0(1370)\gamma$	not seen
Γ_{21} $f_2(1270)\gamma$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=20;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=6
DESIG=10
DESIG=11
DESIG=12
DESIG=13
DESIG=14
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=3
DESIG=8;OUR EST;→ UNCHECKED ←
DESIG=7;OUR EVAL;→ UNCHECKED ←
DESIG=21;OUR EVAL;→ UNCHECKED ←
DESIG=15;OUR EST;→ UNCHECKED ←
DESIG=23;OUR EST;→ UNCHECKED ←
DESIG=9
DESIG=16;OUR EST;→ UNCHECKED ←
DESIG=17;OUR EST;→ UNCHECKED ←
DESIG=18;OUR EST;→ UNCHECKED ←
DESIG=19;OUR EST;→ UNCHECKED ←

 $\rho(1450) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M105220

 $\Gamma(\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{10}/\Gamma$ NODE=M105G3
NODE=M105G3

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.12	¹ DIEKMAN 88	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
$0.027^{+0.015}_{-0.010}$	² KURDADZE 83	OLYA	$0.64-1.4 e^+e^- \rightarrow \pi^+\pi^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

¹ Using total width = 235 MeV.² Using for $\rho(1700)$ mass and width 1600 ± 20 and 300 ± 10 MeV respectively.NODE=M105G3;LINKAGE=B
NODE=M105G3;LINKAGE=KD **$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{11}\Gamma_{10}/\Gamma$** NODE=M105G4
NODE=M105G4

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$335 \pm 27 \pm 20$	13.4k	¹ GRIBANOV 20	CMD3	$1.1-2.0 e^+e^- \rightarrow \eta\pi^+\pi^-$
$210 \pm 24 \pm 10$		² LEES 18	BABR	$e^+e^- \rightarrow \eta\pi^+\pi^-$
74 ± 20		³ AKHMETSHIN 00D	CMD2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
91 ± 19		ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$

¹ Mass and width of the $\rho(770)$ fixed at 775 and 149 MeV, respectively; solution 2 of model 2, $\eta \rightarrow \gamma\gamma$ decays used.² Includes non-resonant contribution. The selected fit model includes three ρ excited states. Model uncertainty is 20%.³ Using the data of ANTONELLI 88, DOLINSKY 91, and AKHMETSHIN 00D. The energy-independent width of the $\rho(1450)$ and $\rho(1700)$ mesons assumed.

NODE=M105G4;LINKAGE=B

NODE=M105G4;LINKAGE=A

NODE=M105G4;LINKAGE=KL

 $\Gamma(K\bar{K}^*(892)+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_{10}/\Gamma$ NODE=M105G8
NODE=M105G8

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
$127 \pm 15 \pm 6$	AUBERT 08S	BABR	$10.6 e^+e^- \rightarrow K\bar{K}^*(892)\gamma$

 $\Gamma(\eta\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_{10}/\Gamma$ NODE=M105G6
NODE=M105G6

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
<16.4	¹ AKHMETSHIN 05	CMD2	$0.60-1.38 e^+e^- \rightarrow \eta\gamma$
$2.2 \pm 0.5 \pm 0.3$	² AKHMETSHIN 01B	CMD2	$e^+e^- \rightarrow \eta\gamma$

¹ From 2γ decay mode of η using 1465 MeV and 310 MeV for the $\rho(1450)$ mass and width. Recalculated by us.² Using the data of AKHMETSHIN 01B on $e^+e^- \rightarrow \eta\gamma$, AKHMETSHIN 00D and ANTONELLI 88 on $e^+e^- \rightarrow \eta\pi^+\pi^-$. Recalculated by us using width of 226 MeV.

NODE=M105G6;LINKAGE=AK

NODE=M105G;LINKAGE=SW

$\rho(1450) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M105230

$$\Gamma(\omega\pi)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R05
NODE=M105R05

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ± 0.4	10.2k	¹ ACHASOV	16D SND	$1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
5.3 ± 0.4	7815	² ACHASOV	13 SND	$1.05-2.00 e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$

OCCUR=3

¹ From a phenomenological model based on vector meson dominance with interfering $\rho(770)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainties not estimated. Supersedes ACHASOV 13.

NODE=M105R05;LINKAGE=A

² From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M105R05;LINKAGE=AC

$$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R00
NODE=M105R00

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.3 ± 0.3	7.4k	¹ ACHASOV	18 SND	$1.22-2.00 e^+ e^- \rightarrow \eta \pi^+ \pi^-$
$4.3^{+1.1}_{-0.9} \pm 0.2$	4.9k	² AULCHENKO	15 SND	$1.22-2.00 e^+ e^- \rightarrow \eta \pi^+ \pi^-$

¹ From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

NODE=M105R00;LINKAGE=B

² From a fit to the $e^+ e^- \rightarrow \eta \pi^+ \pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta \rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

NODE=M105R00;LINKAGE=A

$$\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R17
NODE=M105R17

VALUE (units 10^{-9})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ± 1.4	¹ ACHASOV	10D SND	$1.075-2.0 e^+ e^- \rightarrow \pi^0 \gamma$
---------------	----------------------	---------	--

¹ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. Systematic errors not evaluated.

NODE=M105R17;LINKAGE=A

$$\Gamma(f_0(500)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R01
NODE=M105R01

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<4.0	90	ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	---------	--------	--

$$\Gamma(f_0(980)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{19}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R02
NODE=M105R02

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.6	90	ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	---------	--------	--

$$\Gamma(f_0(1370)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R03
NODE=M105R03

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<3.5	90	ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	---------	--------	--

$$\Gamma(f_2(1270)\gamma)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}} \quad \Gamma_{21}/\Gamma \times \Gamma_{10}/\Gamma$$

NODE=M105R04
NODE=M105R04

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.8	90	¹ ACHASOV	11 SND	$e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
--------	----	----------------------	--------	--

¹ Using Breit-Wigner parametrization of the $\rho(1450)$ with mass and width of 1465 MeV and 400 MeV, respectively.

NODE=M105R01;LINKAGE=AC

 $\rho(1450)$ BRANCHING RATIOS

NODE=M105225

$$\Gamma(\pi\pi)/\Gamma(4\pi) \quad \Gamma_1/\Gamma_3$$

NODE=M105R15
NODE=M105R15

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.37 ± 0.10	^{1,2} ABELE	01B CBAR	$0.0 \bar{p} n \rightarrow 5\pi$
-----------------	----------------------	----------	----------------------------------

¹ $\omega\pi$ not included.

² Using ABELE 97.

NODE=M105R;LINKAGE=BL

NODE=M105R;LINKAGE=LK

$\Gamma(K^+K^-)/\Gamma(\pi^+\pi^-)$ Γ_{14}/Γ_2

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
30.7±8.4±8.2	20k	¹ LEES	17C	BABR $J/\psi \rightarrow h^+ h^- \pi^0$

NODE=M105R08
 NODE=M105R08

¹ From Dalitz plot analyses in isobar models.

NODE=M105R08;LINKAGE=A

 $\Gamma(\omega\pi)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

seen	821	¹ MATVIENKO	15	BELL $\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$
seen	1.6k	ACHASOV	12	SND $e^+ e^- \rightarrow \pi^0 \pi^0 \gamma$
~ 0.21		CLEGG	94	RVUE

NODE=M105R5
 NODE=M105R5

¹ Using Breit-Wigner parameterization of the $\rho(1450)$ and assuming equal probabilities of the $\rho(1450) \rightarrow \pi\pi$ and $\rho(1450) \rightarrow \omega\pi$ decays.

NODE=M105R5;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(\omega\pi)$ Γ_1/Γ_4

VALUE	DOCUMENT ID	TECN
-------	-------------	------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 0.32	CLEGG	94	RVUE
--------	-------	----	------

NODE=M105R6
 NODE=M105R6

 $\Gamma(\omega\pi)/\Gamma(4\pi)$ Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN
-------	-------------	------

••• We do not use the following data for averages, fits, limits, etc. •••

< 0.14	CLEGG	88	RVUE
--------	-------	----	------

NODE=M105R3
 NODE=M105R3

 $\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_5/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.27±0.08	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

¹ $\omega\pi$ not included.

NODE=M105R10
 NODE=M105R10

NODE=M105R10;LINKAGE=BL

 $\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$ Γ_6/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.08±0.04	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

¹ $\omega\pi$ not included.

NODE=M105R11
 NODE=M105R11

NODE=M105R11;LINKAGE=BL

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_7/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.37±0.13	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

¹ $\omega\pi$ not included.

NODE=M105R12
 NODE=M105R12

NODE=M105R12;LINKAGE=BL

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_8/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.11±0.05	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

¹ $\omega\pi$ not included.

NODE=M105R13
 NODE=M105R13

NODE=M105R13;LINKAGE=BL

 $\Gamma(\rho(\pi\pi)_{\text{S-wave}})/\Gamma(4\pi)$ Γ_9/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.17±0.09	¹ ABELE	01B	CBAR 0.0 $\bar{p}n \rightarrow 5\pi$
-----------	--------------------	-----	--------------------------------------

¹ $\omega\pi$ not included.

NODE=M105R14
 NODE=M105R14

NODE=M105R14;LINKAGE=BL

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

seen	35	¹ ACHASOV	14	SND 1.15–2.00 $e^+ e^- \rightarrow \eta\gamma$
------	----	----------------------	----	--

••• We do not use the following data for averages, fits, limits, etc. •••

< 0.04		DONNACHIE	87B	RVUE
--------	--	-----------	-----	------

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

NODE=M105R2
 NODE=M105R2

NODE=M105R2;LINKAGE=A

$\Gamma(\eta\rho)/\Gamma(\omega\pi)$ Γ_{11}/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R4
 NODE=M105R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.081 ± 0.020	^{1,2} AULCHENKO	15	SND	$1.22-2.00 e^+ e^- \rightarrow \eta\pi^+\pi^-$
~ 0.24	³ DONNACHIE	91	RVUE	
>2	FUKUI	91	SPEC	$8.95 \pi^- \rho \rightarrow \omega\pi^0 n$

¹ From a fit to the $e^+ e^- \rightarrow \eta\pi^+\pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

NODE=M105R4;LINKAGE=A

² Reports the inverse of the quoted value as 12.3 ± 3.1 .

NODE=M105R4;LINKAGE=B
 NODE=M105R;LINKAGE=A

³ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

 $\Gamma(\pi\pi)/\Gamma(\eta\rho)$ Γ_1/Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R07
 NODE=M105R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.4	¹ AULCHENKO	15	SND	$1.22-2.00 e^+ e^- \rightarrow \eta\pi^+\pi^-$
---------------	------------------------	----	-----	--

¹ From a fit to the $e^+ e^- \rightarrow \eta\pi^+\pi^-$ cross section with vector meson dominance model including $\rho(770)$, $\rho(1450)$, and $\rho(1700)$ decaying exclusively via $\eta\rho(770)$. Masses and widths of vector states are fixed to PDG 14. Coupling constants are assumed to be real.

NODE=M105R07;LINKAGE=A

 $\Gamma(a_2(1320)\pi)/\Gamma_{total}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R9
 NODE=M105R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMELIN	00	YES	$37 \pi^- \rho \rightarrow \eta\pi^+\pi^- n$
----------	--------	----	-----	--

 $\Gamma(K\bar{K})/\Gamma(\omega\pi)$ Γ_{13}/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R8
 NODE=M105R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.08	¹ DONNACHIE	91	RVUE	
---------	------------------------	----	------	--

¹ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

NODE=M105R8;LINKAGE=A

 $\Gamma(K\bar{K}^*(892) + c.c.)/\Gamma_{total}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M105R16
 NODE=M105R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	COAN	04	CLEO	$\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$
---------------	------	----	------	---

 $\Gamma(\eta\gamma)/\Gamma_{total}$ Γ_{17}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M105R06
 NODE=M105R06

seen	35	¹ ACHASOV	14	SND	$1.15-2.00 e^+ e^- \rightarrow \eta\gamma$
------	----	----------------------	----	-----	--

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

NODE=M105R06;LINKAGE=A

 $\rho(1450)$ REFERENCES

NODE=M105

GRIBANOV	20	JHEP 2001 112	S.S. Gribanov <i>et al.</i>	(CMD-3 Collab.)	REFID=60620
ACHASOV	18	PR D97 012008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58785
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16D	PR D94 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57618
AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55912
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)	REFID=55687
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
		Translated from ZETFP 94 796.			
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
ACHASOV	11	JETP 113 75	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=16721
		Translated from ZETF 140 87.			
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
		Translated from ZETF 130 437.			

AKHMETSHIN 05	PL B605 26	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=50330
ALOISIO 05	PL B606 12	A. Aloisio <i>et al.</i>	(KLOE Collab.)	REFID=51112
SCHAEEL 05C	PRPL 421 191	S. Schaeel <i>et al.</i>	(ALEPH Collab.)	REFID=50845
AKHMETSHIN 04	PL B578 285	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49609
COAN 04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
AKHMETSHIN 03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE 01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
AKHMETSHIN 01B	PL B509 217	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=48167
ALEXANDER 01B	PR D64 092001	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48391
AKHMETSHIN 00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN 00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANDERSON 00A	PR D61 112002	S. Anderson <i>et al.</i>	(CLEO Collab.)	REFID=47468
EDWARDS 00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE 99C	PL B450 275	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46916
ABELE 99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BERTIN 98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
ABELE 97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV 97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BARATE 97M	ZPHY C76 15	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45622
BERTIN 97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
BERTIN 97D	PL B414 220	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45763
CLEGG 94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
BISELLO 91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=41752
DOLINSKY 91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
DONNACHIE 91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41632
FUKUI 91	PL B257 241	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=41581
KUHN 90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
ARMSTRONG 89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41011
BISELLO 89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA 89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
ANTONELLI 88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
CLEGG 88	ZPHY C40 313	A.B. Clegg, A. Donnachie	(MCHS, LANC)	REFID=40922
DIEKMANN 88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI 88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
ALBRECHT 87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40418
DONNACHIE 87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
DOLINSKY 86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246
BARKOV 85	NP B256 365	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=20134
KURDADZE 83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
ASTON 80C	Translated from ZETFP 37 613.	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=20652
BARBER 80C	PL 92B 211	D.P. Barber <i>et al.</i>	(DARE, LANC, SHEF)	REFID=20653
GOUNARIS 68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

NODE=M175

$\eta(1475)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See the $\eta(1405)$ and the related review on "Spectroscopy of Light Meson Resonances."

NODE=M175

$\eta(1475)$ MASS

NODE=M175M5

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
1476 ± 4 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.
 [1475 ± 4 MeV OUR 2023 AVERAGE Scale factor = 1.4]

NODE=M175M5

NEW

1507.6 ± 1.6	$^{+15.5}_{-32.2}$	126K	¹ ABLIKIM	23M	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1469 ± 14	± 13	74	ACHARD	07	L3	$183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K^\pm \pi^\mp$
1460 ± 12		3651	NICHITIU	02	OBLX	$0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1485 ± 8	± 5	20k	ADAMS	01B	B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
1500 ± 10			CICALO	99	OBLX	$0 \bar{p} p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
1464 ± 10			BERTIN	97	OBLX	$0 \bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
1460 ± 10			BERTIN	95	OBLX	$0 \bar{p} p \rightarrow K \bar{K} \pi \pi$
1490	$^{+14}_{-8}$	$^{+3}_{-16}$	1100	BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1475 ± 4			RATH	89	MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$

OCCUR=2

OCCUR=2


• • • We do not use the following data for averages, fits, limits, etc. • • •

1477 ± 7	± 13		² ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
1565 ± 8	$^{+0}_{-63}$		³ ABLIKIM	15T	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
1421 ± 14			AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$

OCCUR=2

- 1 ABLIKIM 23M reports for this state a significance from the fit much higher than 35σ .
- 2 From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^{-+}$. Other J^{PC} not excluded.
- 3 Could also be the $\eta(1405)$.

NODE=M175M5;LINKAGE=C
 NODE=M175M5;LINKAGE=B
 NODE=M175M5;LINKAGE=A



eta(1475) WIDTH

NODE=M175W5
 NODE=M175W5
 NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
96 ± 9	OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below. [90 ± 9 MeV OUR 2023 AVERAGE Scale factor = 1.6]		
115.8 ± 2.4 ^{+14.8} _{-10.9}	126K	ABLIKIM	23M BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
67 ± 18 ± 7	74	ACHARD	07 L3	$183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^\pm \pi^\mp$
120 ± 15	3651	NICHITIU	02 OBLX	$0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
98 ± 18 ± 3	20k	ADAMS	01B B852	$18 \text{ GeV } \pi^- p \rightarrow K^+ K^- \pi^0 n$
100 ± 20		CICALO	99 OBLX	$0 \bar{p} p \rightarrow K^\pm K_S^0 \pi^\mp \pi^+ \pi^-$
105 ± 15		BERTIN	97 OBLX	$0.0 \bar{p} p \rightarrow K^\pm (K^0) \pi^\mp \pi^+ \pi^-$
105 ± 15		BERTIN	95 OBLX	$0 \bar{p} p \rightarrow K \bar{K} \pi \pi \pi$
54 ⁺³⁷ ₋₂₁ ⁺¹³ ₋₂₄		BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
51 ± 13		RATH	89 MPS	$21.4 \pi^- p \rightarrow n K_S^0 K_S^0 \pi^0$


••• We do not use the following data for averages, fits, limits, etc. •••

118 ± 22 ± 17		1 ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
45 ⁺¹⁴ ₋₁₃ ⁺²¹ ₋₂₈		2 ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
63 ± 18		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$

OCCUR=2
 OCCUR=2

OCCUR=2
 NODE=M175W5;LINKAGE=B
 NODE=M175W5;LINKAGE=A

- 1 From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^{-+}$. Other J^{PC} not excluded.
- 2 Could also be the $\eta(1405)$.



η(1475) DECAY MODES

NODE=M175215;NODE=M175

Mode	Fraction (Γ _i /Γ)
Γ ₁ K K̄ π	seen
Γ ₂ K K̄*(892) + c.c.	seen
Γ ₃ a ₀ (980) π	seen
Γ ₄ γ γ	seen
Γ ₅ K _S ⁰ K _S ⁰ η	possibly seen
Γ ₆ γ φ(1020)	possibly seen

DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=7;OUR EST;→ UNCHECKED ←
DESIG=8;OUR EVAL;→ UNCHECKED ←
DESIG=9

η(1475) Γ(i)Γ(γγ)/Γ(total)

NODE=M175220

Γ(K K̄ π) × Γ(γγ)/Γ_{total} Γ₁Γ₄/Γ

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.23±0.05±0.05		74	¹ ACHARD	07 L3	183-209 e ⁺ e ⁻ → e ⁺ e ⁻ K _S ⁰ K [±] π [∓]

NODE=M175G2
NODE=M175G2

••• We do not use the following data for averages, fits, limits, etc. •••
< 0.089 90 ^{2,3}AHOHE 05 CLE2 10.6 e⁺e⁻ → e⁺e⁻K_S⁰K[±]π[∓]

¹Supersedes ACCIARRI 01G. Using B(K_S⁰ → π⁺π⁻) = 0.6895.
²Using η(1475) mass of 1481 MeV and width of 48 MeV. The upper limit increases to 0.140 keV if the world average value, 87 MeV, of the width is used.
³Assuming three-body phase-space decay to K_S⁰K[±]π[∓].

NODE=M175G2;LINKAGE=CH
NODE=M175G2;LINKAGE=AH
NODE=M175G2;LINKAGE=B3

η(1475) BRANCHING RATIOS

NODE=M175225

Γ(K K̄*(892) + c.c.)/Γ(K K̄ π) Γ₂/Γ₁

VALUE	DOCUMENT ID	TECN	COMMENT
0.50±0.10	¹ BAILLON	67 HBC	0.0 p̄p → K K̄ π π π

NODE=M175R1
NODE=M175R1

¹Data could also refer to η(1405).

NODE=M175R;LINKAGE=BL

Γ(K K̄*(892) + c.c.)/[Γ(K K̄*(892) + c.c.) + Γ(a₀(980)π)] Γ₂/(Γ₂+Γ₃)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.25	90	EDWARDS	82E CBAL	J/ψ → K ⁺ K ⁻ π ⁰ γ

NODE=M175R6
NODE=M175R6

••• We do not use the following data for averages, fits, limits, etc. •••

$\Gamma(\gamma\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
$<1.27 \times 10^{-3}$	90	¹ ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma$	

¹ Using results from BAI 00D.NODE=M175R01
NODE=M175R01

NODE=M175R01;LINKAGE=A

 $\Gamma(\gamma\phi(1020))/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
possibly seen	¹ ABLIKIM	181	BES3 $J/\psi \rightarrow \gamma \gamma \phi(1020)$	

¹ Seen as a peak in $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^{-+}$. Other J^{PC} not excluded. Also see $\eta(1405)$.NODE=M175R00
NODE=M175R00

NODE=M175R00;LINKAGE=A

 $\eta(1475)$ REFERENCES

ABLIKIM	23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	181	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	180	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ACHARD	07	JHEP 0703 018	P. Achard <i>et al.</i>	(L3 Collab.)
AHOHE	05	PR D71 072001	R. Ahohe <i>et al.</i>	(CLEO Collab.)
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)
ACCIARRI	01G	PL B501 1	M. Acciarri <i>et al.</i>	(L3 Collab.)
ADAMS	01B	PL B516 264	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)
CICALO	99	PL B462 453	C. Cicalo <i>et al.</i>	(OBELIX Collab.)
BERTIN	97	PL B400 226	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BERTIN	95	PL B361 187	A. Bertin <i>et al.</i>	(OBELIX Collab.)
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)
RATH	89	PR D40 693	M.G. Rath <i>et al.</i>	(NDAM, BRAN, BNL, CUNY+)
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)
BAILLON	67	NC 50A 393	P.H. Baillon <i>et al.</i>	(CERN, CDEF, IRAD)

NODE=M175

REFID=62055
REFID=58893
REFID=58925
REFID=56785
REFID=51698
REFID=50764
REFID=48848
REFID=48319
REFID=49649
REFID=47954
REFID=47394
REFID=45417
REFID=44614
REFID=41584
REFID=41578
REFID=40924
REFID=21314
REFID=20407 $f_0(1500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M152

NODE=M152

 $f_0(1500)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M152PP

NODE=M152PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1430–1530) – i (40–90) OUR ESTIMATE			
$(1450 \pm 10) - i (53 \pm 8)$	¹ RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1483 \pm 15) - i (58 \pm 6)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1496 \pm 1.2^+_{-26.4}) - i (40.4 \pm 0.3^+_{-2.5})$	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
$(1465 \pm 18) - i (50 \pm 9)$	³ ROPERTZ	18	RVUE $\bar{B}_s^0 \rightarrow J/\psi(\pi^+ \pi^- / K^+ K^-)$
$(1486 \pm 10) - i (57 \pm 5)$	ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
$(1489 \pm 8) - i (51 \pm 5)$	⁴ ANISOVICH	03	RVUE
$(1515 \pm 12) - i (55 \pm 12)$	BARBERIS	00A	$450 \bar{p}p \rightarrow p_f(\eta\eta', \eta'\eta')p_s$
$(1511 \pm 9) - i (51 \pm 9)$	⁵ BARBERIS	00C	$450 \bar{p}p \rightarrow p_f 4\pi p_s$
$(1510 \pm 8) - i (55 \pm 8)$	BARBERIS	00E	$450 \bar{p}p \rightarrow p_f \eta\eta p_s$
$(1502 \pm 12 \pm 10) - i (49 \pm 9 \pm 8)$	⁶ BARBERIS	99D	OMEG $450 \bar{p}p \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1447 \pm 27) - i (54 \pm 23)$	⁷ KAMINSKI	99	RVUE $\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$(1499 \pm 8) - i (65 \pm 10)$	ANISOVICH	98B	RVUE Compilation.
$(1510 \pm 20) - i (60 \pm 18)$	BARBERIS	97B	OMEG $450 \bar{p}p \rightarrow p_p 2(\pi^+ \pi^-)$
$(1449 \pm 20) - i (57 \pm 15)$	BERTIN	97C	OBLX $0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1515 \pm 20) - i (53 \pm 8)$	ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
$(1500 \pm 8) - i (66 \pm 8)$	ABELE	96C	RVUE Compilation.
$(1500 \pm 10) - i (77 \pm 15)$	⁸ AMSLER	95D	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
$(1520 \pm 25) - i (74^+_{-13})$	⁹ ANISOVICH	94	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$
$(1505 \pm 20) - i (75 \pm 10)$	¹⁰ BUGG	94	RVUE $\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0\pi^0$

NODE=M152PP

→ UNCHECKED ←

- ¹ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA). NODE=M152PP;LINKAGE=E
- ² T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$), and BINON 84C ($\eta\eta'$). NODE=M152PP;LINKAGE=A
- ³ T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants. NODE=M152PP;LINKAGE=F
- ⁴ Pole position from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K\bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest. NODE=M152PP;LINKAGE=G
- ⁵ Average between $\pi^+ \pi^- 2\pi^0$ and $2(\pi^+ \pi^-)$. NODE=M152PP;LINKAGE=C
- ⁶ Supersedes BARBERIS 99 and BARBERIS 99B. NODE=M152PP;LINKAGE=H
- ⁷ T-matrix pole on sheet $--+$. NODE=M152PP;LINKAGE=I
- ⁸ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D. NODE=M152PP;LINKAGE=D
- ⁹ From a simultaneous analysis of the annihilations $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$. NODE=M152PP;LINKAGE=J
- ¹⁰ Reanalysis of ANISOVICH 94 data. NODE=M152PP;LINKAGE=K

$f_0(1500)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1522 ± 25		¹ BERTIN	98	OBLX 0.05–0.405 $\bar{p} p \rightarrow \pi^+ \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1492.5 ± 3.6 ⁺ _{–20.5}		² ABLIKIM	22G	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
1447 ± 16 ± 13	163	^{3,4} DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1442 ± 9 ± 4	261	^{3,4} DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
1460.9 ± 2.9		⁵ AAIJ	14BR	LHCB $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$
1468 ⁺¹⁴ _{–15} ⁺²³ _{–74}	5.5k	⁶ ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1470 ± 60	568	⁷ KLEMP	08	E791 $D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
1470 ⁺⁶ _{–7} ⁺⁷² _{–255}		⁸ UEHARA	08A	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1466 ± 6 ± 20		⁹ ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1495 ± 4		AMSLER	06	CBAR 0.9 $\bar{p} p \rightarrow K^+ K^- \pi^0$
1539 ± 20	9.9k	AUBERT	06O	BABR $B^+ \rightarrow K^+ K^+ K^-$
1473 ± 5	80k	^{9,10} UMAN	06	E835 5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
1478 ± 6		VLADIMIRSK...	06	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1493 ± 7		⁹ BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
1524 ± 14	1400	¹¹ GARMASH	05	BELL $B^+ \rightarrow K^+ K^+ K^-$
1490 ± 30		⁹ ABELE	01	CBAR 0.0 $\bar{p} d \rightarrow \pi^- 4\pi^0 p$
1497 ± 10		⁹ BARBERIS	99	OMEG 450 $pp \rightarrow p_s p_f K^+ K^-$
1502 ± 10		⁹ BARBERIS	99B	OMEG 450 $pp \rightarrow p_s p_f \pi^+ \pi^-$
1530 ± 45		⁹ BELLAZZINI	99	GAM4 450 $pp \rightarrow pp \pi^0 \pi^0$
1505 ± 18		⁹ FRENCH	99	300 $pp \rightarrow p_f (K^+ K^-) p_s$
1580 ± 80		⁹ ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
~ 1520		REYES	98	SPEC 800 $pp \rightarrow p_s p_f K_S^0 K_S^0$
~ 1475		FRABETTI	97D	E687 $D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 1505		ABELE	96	CBAR 0.0 $\bar{p} p \rightarrow 5\pi^0$
1460 ± 20	120	⁹ AMELIN	96B	VES 37 $\pi^- A \rightarrow \eta \eta \pi^- A$
1500 ± 8		BUGG	96	RVUE
1500 ± 15		¹² AMSLER	95B	CBAR 0.0 $\bar{p} p \rightarrow 3\pi^0$
1505 ± 15		¹³ AMSLER	95C	CBAR 0.0 $\bar{p} p \rightarrow \eta \eta \pi^0$
1445 ± 5		¹⁴ ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1497 ± 30		⁹ ANTINORI	95	OMEG 300,450 $pp \rightarrow pp\pi^+ \pi^-$
~ 1505		BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1446 ± 5		⁹ ABATZIS	94	OMEG 450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1545 ± 25		⁹ AMSLER	94E	CBAR 0.0 $\bar{p} p \rightarrow \pi^0 \eta \eta'$
1560 ± 25		⁹ AMSLER	92	CBAR 0.0 $\bar{p} p \rightarrow \pi^0 \eta \eta$
1550 ± 45 ± 30		⁹ BELADIDZE	92C	VES 36 $\pi^- Be \rightarrow \pi^- \eta' \eta Be$
1449 ± 4		⁹ ARMSTRONG	89E	OMEG 300 $pp \rightarrow pp2(\pi^+ \pi^-)$
1610 ± 20		⁹ ALDE	88	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$

NODE=M152M

NODE=M152M

OCCUR=2

OCCUR=2

OCCUR=2

~ 1525		ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1570 ± 20	600	⁹ ALDE	87	GAM4	100	$\pi^- p \rightarrow 4\pi^0 n$
1575 ± 45		¹⁵ ALDE	86D	GAM4	100	$\pi^- p \rightarrow 2\eta n$
1568 ± 33		⁹ BINON	84C	GAM2	38	$\pi^- p \rightarrow \eta\eta' n$
1592 ± 25		⁹ BINON	83	GAM2	38	$\pi^- p \rightarrow 2\eta n$
1525 ± 5		⁹ GRAY	83	DBC	0.0	$\bar{p}N \rightarrow 3\pi$

¹ Breit-Wigner mass.

² The $\pi^+ \pi^-$ mass spectrum is described by a coherent sum of two Breit-Wigner resonances, $f_0(1500)$ and a new $X(1540)$ with mass $1540.2 \pm 7.0^{+36.3}_{-6.1}$ MeV and width $157 \pm 19^{+11}_{-77}$ MeV.

³ Using CLEO-c data but not authored by the CLEO Collaboration.

⁴ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 109$ MeV.

⁵ Solution I, statistical error only.

⁶ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

⁷ Reanalysis of AITALA 01A data. This state could also be $f_0(1370)$.

⁸ Breit-Wigner mass. May also be the $f_0(1370)$.

⁹ Breit-Wigner mass.

¹⁰ Statistical error only.

¹¹ Breit-Wigner, solution 1, PWA ambiguous.

¹² T-matrix pole, supersedes ANISOVICH 94.

¹³ T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

¹⁴ Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

¹⁵ From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152M;LINKAGE=PP
NODE=M152M;LINKAGE=M

NODE=M152M;LINKAGE=F
NODE=M152M;LINKAGE=G
NODE=M152M;LINKAGE=I
NODE=M152M;LINKAGE=C

NODE=M152M;LINKAGE=KL
NODE=M152M;LINKAGE=UE
NODE=M152M;LINKAGE=E
NODE=M152M;LINKAGE=ST
NODE=M152M;LINKAGE=GA
NODE=M152M;LINKAGE=D
NODE=M152M;LINKAGE=D1
NODE=M152M;LINKAGE=B
NODE=M152M;LINKAGE=AZ

$f_0(1500)$ WIDTH

NODE=M152W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
108 ± 33		¹ BERTIN	98	OBLX $0.05\text{--}0.405 \bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
107 ± 9^{+21}_{-7}		² ABLIKIM	22G	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
124 ± 7		³ AAIJ	14BR	LHCB $\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$
$136^{+41+28}_{-26-100}$	5.5k	⁴ ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
90^{+2+50}_{-1-22}		⁵ UEHARA	08A	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
$108^{+14}_{-11} \pm 25$		⁶ ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
121 ± 8		AMSLER	06	CBAR $0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
257 ± 33	9.9k	AUBERT	06O	BABR $B^+ \rightarrow K^+ K^+ K^-$
108 ± 9	80k	^{6,7} UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
119 ± 10		VLADIMIRSK...	06	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
90 ± 15		⁶ BINON	05	GAMS $33 \pi^- p \rightarrow \eta \eta n$
136 ± 23	1400	⁸ GARMASH	05	BELL $B^+ \rightarrow K^+ K^+ K^-$
140 ± 40		⁶ ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
104 ± 25		⁶ BARBERIS	99	OMEG $450 pp \rightarrow p_s p_f K^+ K^-$
131 ± 15		⁶ BARBERIS	99B	OMEG $450 pp \rightarrow p_s p_f \pi^+ \pi^-$
160 ± 50		⁶ BELLAZZINI	99	GAM4 $450 pp \rightarrow pp \pi^0 \pi^0$
100 ± 33		⁶ FRENCH	99	$300 pp \rightarrow p_f (K^+ K^-) p_s$
280 ± 100		⁶ ALDE	98	GAM4 $100 \pi^- p \rightarrow \pi^0 \pi^0 n$
~ 100		FRABETTI	97D	E687 $D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$
~ 169		ABELE	96	CBAR $0.0 \bar{p}p \rightarrow 5\pi^0$
100 ± 30	120	⁶ AMELIN	96B	VES $37 \pi^- A \rightarrow \eta \eta \pi^- A$
132 ± 15		BUGG	96	RVUE
120 ± 25		⁹ AMSLER	95B	CBAR $0.0 \bar{p}p \rightarrow 3\pi^0$
120 ± 30		¹⁰ AMSLER	95C	CBAR $0.0 \bar{p}p \rightarrow \eta \eta \pi^0$
65 ± 10		¹¹ ANTINORI	95	OMEG $300,450 pp \rightarrow pp2(\pi^+ \pi^-)$

NODE=M152W

OCCUR=2

199 ± 30		⁶ ANTINORI	95	OMEG	300,450 $p p \rightarrow p p \pi^+ \pi^-$
56 ± 12		⁶ ABATZIS	94	OMEG	450 $p p \rightarrow p p 2(\pi^+ \pi^-)$
100 ± 40		⁶ AMSLER	94E	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \eta \eta'$
245 ± 50		⁶ AMSLER	92	CBAR	0.0 $\bar{p} p \rightarrow \pi^0 \eta \eta$
153 ± 67 ± 50		⁶ BELADIDZE	92C	VES	36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
78 ± 18		⁶ ARMSTRONG	89E	OMEG	300 $p p \rightarrow p p 2(\pi^+ \pi^-)$
170 ± 40		⁶ ALDE	88	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$
150 ± 20	600	⁶ ALDE	87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
265 ± 65		¹² ALDE	86D	GAM4	100 $\pi^- p \rightarrow 2\eta n$
260 ± 60		⁶ BINON	84C	GAM2	38 $\pi^- p \rightarrow \eta \eta' n$
210 ± 40		⁶ BINON	83	GAM2	38 $\pi^- p \rightarrow 2\eta n$
101 ± 13		⁶ GRAY	83	DBC	0.0 $\bar{p} N \rightarrow 3\pi$

OCCUR=2

¹ Breit-Wigner width.

² The $\pi^+ \pi^-$ mass spectrum is described by a coherent sum of two Breit-Wigner resonances, $f_0(1500)$ and a new $X(1540)$ with mass $1540.2 \pm 7.0^{+36.3}_{-6.1}$ MeV and width $157 \pm 19^{+11}_{-77}$ MeV.

³ Solution I, statistical error only.

⁴ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

⁵ Breit-Wigner width. May also be the $f_0(1370)$.

⁶ Breit-Wigner width.

⁷ Statistical error only.

⁸ Breit-Wigner, solution 1, PWA ambiguous.

⁹ T-matrix pole, supersedes ANISOVICH 94.

¹⁰ T-matrix pole, supersedes ANISOVICH 94 and AMSLER 92.

¹¹ Supersedes ABATZIS 94, ARMSTRONG 89E. Breit-Wigner mass.

¹² From central value and spread of two solutions. Breit-Wigner mass.

NODE=M152W;LINKAGE=PP
NODE=M152W;LINKAGE=J

NODE=M152W;LINKAGE=G
NODE=M152W;LINKAGE=C

NODE=M152W;LINKAGE=UE
NODE=M152W;LINKAGE=E
NODE=M152W;LINKAGE=ST
NODE=M152W;LINKAGE=GA
NODE=M152W;LINKAGE=D
NODE=M152W;LINKAGE=D1
NODE=M152W;LINKAGE=B
NODE=M152W;LINKAGE=AZ

$f_0(1500)$ DECAY MODES

NODE=M152215;NODE=M152

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 $\pi \pi$	(34.5 ± 2.2) %	1.2
Γ_2 $\pi^+ \pi^-$	seen	
Γ_3 $2\pi^0$	seen	
Γ_4 4π	(48.9 ± 3.3) %	1.2
Γ_5 $4\pi^0$	seen	
Γ_6 $2\pi^+ 2\pi^-$	seen	
Γ_7 $2(\pi\pi)_{S\text{-wave}}$	seen	
Γ_8 $\rho\rho$	seen	
Γ_9 $\pi(1300)\pi$	seen	
Γ_{10} $a_1(1260)\pi$	seen	
Γ_{11} $\eta\eta$	(6.0 ± 0.9) %	1.1
Γ_{12} $\eta\eta'(958)$	(2.2 ± 0.8) %	1.4
Γ_{13} $K\bar{K}$	(8.5 ± 1.0) %	1.1
Γ_{14} $\gamma\gamma$	not seen	

DESIG=8
DESIG=9
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=7
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←
DESIG=11;OUR EST;→ UNCHECKED ←
DESIG=12;OUR EST;→ UNCHECKED ←
DESIG=13;OUR EST;→ UNCHECKED ←
DESIG=14;OUR EST;→ UNCHECKED ←
DESIG=1
DESIG=2
DESIG=4
DESIG=10;OUR EST;→ UNCHECKED ←

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 10 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 5.6$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-88			
x_{11}	27	-56		
x_{12}	3	-32	26	
x_{13}	43	-64	20	2
	x_1	x_4	x_{11}	x_{12}

$f_0(1500) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M152217

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{14}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M152G1
NODE=M152G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$33^{+12}_{-6} + 1809_{-21}$		¹ UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
not seen		ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$
<460	95	BARATE	00E ALEP	$\gamma\gamma \rightarrow \pi^+ \pi^-$

¹ May also be the $f_0(1370)$. Multiplied by us by 3 to obtain the $\pi\pi$ value.

NODE=M152G1;LINKAGE=UE

 $f_0(1500) \text{ BRANCHING RATIOS}$

NODE=M152220

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN
-------	-------------	------

NODE=M152R8
NODE=M152R8

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.454±0.104	BUGG	96 RVUE
-------------	------	---------

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R10
NODE=M152R10seen BERTIN 98 OBLX 0.05-0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen FRABETTI 97D E687 $D_s^\pm \rightarrow \pi^\mp \pi^\pm \pi^\pm$ $\Gamma(4\pi)/\Gamma(\pi\pi)$ Γ_4/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R6
NODE=M152R6**1.42±0.18 OUR FIT** Error includes scale factor of 1.2.**1.42±0.18 OUR AVERAGE** Error includes scale factor of 1.2.1.37±0.16 BARBERIS 00D 450 $p\bar{p} \rightarrow p_f 4\pi p_s$ 2.1 ±0.6 ¹ AMSLER 98 RVUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.1 ±0.2 ² ANISOVICH 02D SPEC Combined fit3.4 ±0.8 ¹ ABELE 96 CBAR 0.0 $\bar{p}p \rightarrow 5\pi^0$ ¹ Excluding $\rho\rho$ contribution to 4π .² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0, \pi^0 \eta \eta, \pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n, \eta \eta n, \eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.NODE=M152R6;LINKAGE=C
NODE=M152R6;LINKAGE=CH $\Gamma(2(\pi\pi)\text{s-wave})/\Gamma(\pi\pi)$ Γ_7/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R14
NODE=M152R14

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.42±0.26 ¹ ABELE 01 CBAR 0.0 $\bar{p}d \rightarrow \pi^- 4\pi^0 p$ ¹ From the combined data of ABELE 96 and ABELE 96C.

NODE=M152R;LINKAGE=KZ

 $\Gamma(2(\pi\pi)\text{s-wave})/\Gamma(4\pi)$ Γ_7/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R15
NODE=M152R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.26±0.07 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$ $\Gamma(\rho\rho)/\Gamma(4\pi)$ Γ_8/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R16
NODE=M152R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13±0.08 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$ $\Gamma(\rho\rho)/\Gamma(2(\pi\pi)\text{s-wave})$ Γ_8/Γ_7

VALUE	DOCUMENT ID	COMMENT
-------	-------------	---------

NODE=M152R11
NODE=M152R11**2.87±0.34 OUR AVERAGE** Error includes scale factor of 1.1.3.3 ±0.5 BARBERIS 00C 450 $p\bar{p} \rightarrow p_f \pi^+ \pi^- 2\pi^0 p_s$ 2.6 ±0.4 BARBERIS 00C 450 $p\bar{p} \rightarrow p_f 2(\pi^+ \pi^-) p_s$

OCCUR=2

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ Γ_9/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R17
NODE=M152R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.50±0.25 ABELE 01B CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$ Γ_{10}/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R18
 NODE=M152R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.12±0.05	ABELE	01B	CBAR 0.0 $\bar{p}d \rightarrow 5\pi p$
-----------	-------	-----	--

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R1
 NODE=M152R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

large	ALDE	88	GAM4 300 $\pi^- N \rightarrow \eta\eta\pi^- N$
large	BINON	83	GAM2 38 $\pi^- p \rightarrow 2\eta n$

 $\Gamma(\eta\eta)/\Gamma(\pi\pi)$ Γ_{11}/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R13
 NODE=M152R13

0.173±0.024 OUR FIT Error includes scale factor of 1.1.

0.175±0.027 OUR AVERAGE

0.18 ±0.03	BARBERIS	00E	450 $p\bar{p} \rightarrow p_f\eta\eta p_s$
0.157±0.060	¹ AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.080±0.033	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
0.11 ±0.03	² ANISOVICH	02D	SPEC Combined fit
0.078±0.013	³ ABELE	96C	RVUE Compilation
0.230±0.097	⁴ AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$

¹ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

³ 2π width determined to be 60 ± 12 MeV.

⁴ Using AMSLER 95B ($3\pi^0$).

NODE=M152R3;LINKAGE=AB
 NODE=M152R;LINKAGE=CH

NODE=M152R3;LINKAGE=CM
 NODE=M152R3;LINKAGE=A

 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$ Γ_5/Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R5
 NODE=M152R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.8±0.3	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$
---------	------	----	---

 $\Gamma(\eta\eta'(958))/\Gamma(\pi\pi)$ Γ_{12}/Γ_1

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M152R12
 NODE=M152R12

6.4±2.2 OUR FIT Error includes scale factor of 1.4.

9.5±2.6	BARBERIS	00A	450 $p\bar{p} \rightarrow p_f\eta\eta p_s$
----------------	----------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

16.6 ^{+4.2} _{-4.0}	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
0.5±0.3	² ANISOVICH	02D	SPEC Combined fit

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.

² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

NODE=M152R12;LINKAGE=A

NODE=M152R12;LINKAGE=CH

 $\Gamma(\eta\eta'(958))/\Gamma(\eta\eta)$ Γ_{12}/Γ_{11}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R2
 NODE=M152R2

0.37±0.13 OUR FIT Error includes scale factor of 1.5.

0.29±0.10	¹ AMSLER	95C	CBAR 0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
------------------	---------------------	-----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.05±0.03	² ANISOVICH	02D	SPEC Combined fit
0.84±0.23	ABELE	96C	RVUE Compilation
2.7 ±0.8	BINON	84C	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

¹ Using AMSLER 94E ($\eta\eta'\pi^0$).

² From a combined K-matrix analysis of Crystal Barrel (0. $p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

NODE=M152R2;LINKAGE=A

NODE=M152R2;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M152R9
 NODE=M152R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.044±0.021	BUGG	96	RVUE
-------------	------	----	------

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_{13}/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.246 ± 0.025 OUR FIT**0.236 ± 0.026 OUR AVERAGE**

0.25 ± 0.03	¹ BARGIOTTI	03	OBLX $\bar{p}p$
0.19 ± 0.07	² ABELE	98	CBAR $0.0 \bar{p}p \rightarrow K_L^0 K^\pm \pi^\mp$
0.20 ± 0.08	³ ABELE	96B	CBAR $0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.16 ± 0.05	⁴ ANISOVICH	02D	SPEC Combined fit
0.33 ± 0.03 ± 0.07	BARBERIS	99D	OMEG $450 p\bar{p} \rightarrow K^+ K^-, \pi^+ \pi^-$

¹ Coupled channel analysis of $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, and $K^\pm K_S^0 \pi^\mp$.

² Using $\pi^0 \pi^0$ from AMSLER 95B.

³ Using AMSLER 95B ($3\pi^0$), AMSLER 94C ($2\pi^0 \eta$) and SU(3).

⁴ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta \eta n$, $\eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.

NODE=M152R7
NODE=M152R7

NODE=M152R;LINKAGE=BG
NODE=M152R7;LINKAGE=A
NODE=M152R7;LINKAGE=D
NODE=M152R7;LINKAGE=CH

 $\Gamma(K\bar{K})/\Gamma(\eta\eta)$ Γ_{13}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

1.43 ± 0.24 OUR FIT Error includes scale factor of 1.1.**1.85 ± 0.41** BARBERIS 00E 450 $p\bar{p} \rightarrow p_f \eta \eta p_S$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ± 0.6	¹ ANISOVICH	02D	SPEC Combined fit
<0.4	90	² PROKOSHKIN	91 GAM4 $300 \pi^- p \rightarrow \pi^- p \eta \eta$
<0.6	³ BINON	83	GAM2 $38 \pi^- p \rightarrow 2\eta n$

¹ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$), GAMS ($\pi p \rightarrow \pi^0 \pi^0 n$, $\eta \eta n$, $\eta \eta' n$), and BNL ($\pi p \rightarrow K \bar{K} n$) data.

² Combining results of GAM4 with those of WA76 on $K \bar{K}$ central production.

³ Using ETKIN 82B and COHEN 80.

NODE=M152R4
NODE=M152R4

NODE=M152R4;LINKAGE=CH

NODE=M152R4;LINKAGE=BZ
NODE=M152R4;LINKAGE=A

 $\phi_0(1500)$ REFERENCES

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
Also		PR D107 079901 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62033
ABLIKIM	22G	PRL 129 042001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61642
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ROPERTZ	18	EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis	(BONN, JULI)	REFID=59332
AAIJ	17V	JHEP 1708 037	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57828
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
AAIJ	14BR	PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56035
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
KLEMP	08	EPJ C55 39	E. Klemp, M. Matveev, A.V. Sarantsev	(BONN+)	REFID=52286
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52309
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
AUBERT	06O	PR D74 032003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51141
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
		Translated from YAF 68 998.			
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=50641
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)	REFID=49217
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48334
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)	REFID=48004
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00D	PL B474 423	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47960
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>		REFID=47400
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)	REFID=46927
ABELE	98	PR D57 3860	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45863
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
		Translated from YAF 62 446.			
AMSLER	98	RMP 70 1293	C. Amsler		REFID=46601
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			

NODE=M152

BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
REYES	98	PRL 81 4079	M.A. Reyes <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
FRABETTI	97D	PL B407 79	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AMELIN	96B	PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
Translated from YAF 59 1021.				
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+)
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)
ABATZIS	94	PL B324 509	S. Abatzis <i>et al.</i>	(ATHU, BARI, BIRM+)
AMSLER	94C	PL B327 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	94E	PL B340 259	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bitjukov, G.V. Borisov	(SERP+)
Translated from YAF 55 2748.				
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)
Translated from DANS 316 900.				
ARMSTRONG	89E	PL B228 536	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)
ALDE	88	PL B201 160	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
Also				
SJNP 38 561				
Translated from YAF 38 934.				
GRAY	83	PR D27 307	L. Gray <i>et al.</i>	(SYRA)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
COHEN	80	PR D22 2595	D. Cohen <i>et al.</i>	(ANL)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)

REFID=45782
 REFID=46378
 REFID=45758
 REFID=45701
 REFID=45554
 REFID=45011
 REFID=45038
 REFID=45076
 REFID=44725
 REFID=45094
 REFID=44377
 REFID=44440
 REFID=44441
 REFID=44437
 REFID=44438
 REFID=44090
 REFID=44091
 REFID=44093
 REFID=44080
 REFID=43659
 REFID=44078
 REFID=43169
 REFID=43175
 REFID=41719
 REFID=41011
 REFID=40283
 REFID=40330
 REFID=40221
 REFID=20765
 REFID=20768
 REFID=21418
 REFID=20750
 REFID=20751
 REFID=21370
 REFID=20390
 REFID=20381
 REFID=20355


$f_1(1510)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

OMITTED FROM SUMMARY TABLE
 See the review on "Spectroscopy of Light Meson Resonances."

$f_1(1510)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1518 ± 5 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
1530 ± 10		ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
1512 ± 4	600	¹ BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1526 ± 6	271	GAVILLET	82 HBC	4.2 $K^- p \rightarrow \Lambda K K \pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 1525		² BAUER	93B	$\gamma \gamma^* \rightarrow \pi^+ \pi^- \pi^0 \pi^0$
¹ From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.				
² Not seen by AIHARA 88C in the $K_S^0 K^\pm \pi^\mp$ final state.				



NODE=M084
 NODE=M084
 NODE=M084M;LINKAGE=A
 NODE=M084M;LINKAGE=C

$f_1(1510)$ WIDTH


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
73±25 OUR AVERAGE	Error includes scale factor of 2.5. See the ideogram below.			
100±40		ASTON	88C LASS	11 $K^- p \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$
35±15	600	³ BIRMAN	88 MPS	8 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
107±15	271	GAVILLET	82 HBC	4.2 $K^- p \rightarrow \Lambda K K \pi$

³From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ state.

NODE=M084W

NODE=M084W

NODE=M084W;LINKAGE=A



$f_1(1510)$ DECAY MODES

NODE=M084215;NODE=M084

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}^*(892) + c.c.$	seen
Γ_2 $\pi^+ \pi^- \eta'$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2

$f_1(1510)$ BRANCHING RATIOS

NODE=M084225

$\Gamma(\pi^+ \pi^- \eta')/\Gamma_{total}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	230	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$	

NODE=M084R01
NODE=M084R01

$f_1(1510)$ REFERENCES

NODE=M084

ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
BAUER	93B	PR D48 3976	D.A. Bauer <i>et al.</i>	(SLAC)	REFID=43678
AIHARA	88C	PR D38 1	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=40564
ASTON	88C	PL B201 573	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP	REFID=40282
BIRMAN	88	PRL 61 1557	A. Birman <i>et al.</i>	(BNL, FSU, IND, MASD) JP	REFID=40568
GAVILLET	82	ZPHY C16 119	P. Gavillet <i>et al.</i>	(CERN, CDEF, PADO+)	REFID=20877

NODE=M013

$f'_2(1525)$


$$I^G(J^{PC}) = 0^+(2^{++})$$

$f'_2(1525)$ MASS

NODE=M013MX

VALUE (MeV) DOCUMENT ID
1517.3±2.4 OUR AVERAGE Includes data from the 6 datablocks that follow this one.
 Error includes scale factor of 2.8. See the ideogram below. [1517.4 ± 2.5 MeV OUR 2023
 AVERAGE Scale factor = 2.8]

NODE=M013MX
 NEW



PRODUCED BY PION BEAM

NODE=M013M1
 NODE=M013M1

VALUE (MeV) EVTs DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

• • • We do not use the following data for averages, fits, limits, etc. • • •

1521±13		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1547 ⁺¹⁰ ₋₂	1	LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1496 ⁺⁹ ₋₈	2	CHABAUD 81	ASPK	6 $\pi^- p \rightarrow K^+ K^- n$
1497 ⁺⁸ ₋₉		CHABAUD 81	ASPK	18.4 $\pi^- p \rightarrow K^+ K^- n$
1492±29		GORLICH 80	ASPK	17 $\pi^- p$ polarized $\rightarrow K^+ K^- n$
1502±25	3	CORDEN 79	OMEG	12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
1480	14	CRENNELL 66	HBC	6.0 $\pi^- p \rightarrow K_S^0 K_S^0 n$

OCCUR=2

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
² CHABAUD 81 is a reanalysis of PAWLICKI 77 data.
³ From an amplitude analysis where the $f'_2(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.

NODE=M013M;LINKAGE=L
 NODE=M013M;LINKAGE=D
 NODE=M013M;LINKAGE=N

PRODUCED BY K^\pm BEAM

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

NODE=M013M2
 NODE=M013M2

1518.0 ± 2.7 OUR AVERAGE Includes data from the datablock that follows this one.
 Error includes scale factor of 2.9. See the ideogram below. [1518.1 ± 2.8 MeV OUR 2023
 AVERAGE Scale factor = 3.0]

NEW


1526.8 ± 4.3	ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1504 ± 12	BOLONKIN	86	SPEC	40	$K^- p \rightarrow K_S^0 K_S^0 \Upsilon$
1529 ± 3	ARMSTRONG	83B	OMEG	18.5	$K^- p \rightarrow K^- K^+ \Lambda$
1521 ± 6	650	AGUILAR-...	81B	HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
1521 ± 3	572	ALHARRAN	81	HBC	8.25 $K^- p \rightarrow \Lambda K \bar{K}$
1522 ± 6	123	BARREIRO	77	HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
1528 ± 7	166	EVANGELIS...	77	OMEG	10 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1527 ± 3	120	BRANDENB...	76c	ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1519 ± 7	100	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1514 ± 8	61	BINON	07	GAMS	32.5 $K^- p \rightarrow \eta \eta (\Lambda / \Sigma^0)$
1513 ± 10	1	BARKOV	99	SPEC	40 $K^- p \rightarrow K_S^0 K_S^0 \Upsilon$

¹Systematic errors not estimated.

NODE=M013M2;LINKAGE=SK



PRODUCED BY K^\pm BEAM (MeV)

PRODUCED IN e^+e^- ANNIHILATION AND PARTICLE DECAYS

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

NODE=M013M3
 NODE=M013M3

1514 ± 4 OUR AVERAGE Error includes scale factor of 3.6. See the ideogram below.
 [1514⁺⁵₋₄ MeV OUR 2023 AVERAGE Scale factor = 3.8]


NEW

1515.4 ± 2.5 ^{+3.2} _{-7.6}	126K	ABLIKIM	23M	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
1522.2 ± 2.8 ^{+5.3} _{-2.0}		AAIJ	13AN	LHCB	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
1513 ± 5 ⁺⁴ ₋₁₀	5.5k	¹ ABLIKIM	13N	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1525.3 ^{+1.2} _{-1.4} ^{+3.7} _{-2.1}		UEHARA	13	BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
1521 ± 5		ABLIKIM	05	BES2	$J/\psi \rightarrow \phi K^+ K^-$
1518 ± 1 ± 3		ABE	04	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1519 ± 2 ⁺¹⁵ ₋₅		BAI	03G	BES	$J/\psi \rightarrow \gamma K \bar{K}$

1523 ± 6	331	² ACCIARRI	01H L3	91, 183-209	$e^+e^- \rightarrow e^+e^-K_S^0K_S^0$
1535 ± 5 ± 4		ABREU	96C DLPH	$Z^0 \rightarrow K^+K^- + X$	
1516 ± 5 ⁺⁹ / ₋₁₅		BAI	96C BES	$J/\psi \rightarrow \gamma K^+K^-$	
1531.6 ± 10.0		AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+K^-$	
1496 ± 2		³ FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+K^-$	OCCUR=2
1525 ± 10 ± 10		BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+K^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1518 ± 3		⁴ KLEMPPT	22 RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0, \gamma K_S^0 K_S^0$	
1503 ± 11		⁵ RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$	
1532 ± 3 ± 6	644	^{6,7} DOBBS	15	$J/\psi \rightarrow \gamma K^+K^-$	
1557 ± 9 ± 3	113	^{6,7} DOBBS	15	$\psi(2S) \rightarrow \gamma K^+K^-$	OCCUR=2
1526 ± 7	29	⁸ LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+K^- \gamma$	
1523 ± 5	870	⁹ SCHEGELSKY	06A RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
1515 ± 5		¹⁰ FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+K^-$	

- ¹ From partial wave analysis including all possible combinations of $0^{++}, 2^{++},$ and 4^{++} resonances.
- ² Supersedes ACCIARRI 95J.
- ³ From an analysis including interference with $f_0(1710)$.
- ⁴ Fit of the tensor partial waves from BES3 in the multipole basis.
- ⁵ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).
- ⁶ Using CLEO-c data but not authored by the CLEO Collaboration.
- ⁷ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 73$ MeV.
- ⁸ From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated.
- ⁹ From analysis of L3 data at 91 and 183-209 GeV.
- ¹⁰ From an analysis ignoring interference with $f_0(1710)$.

NODE=M013M3;LINKAGE=A
 NODE=M013M;LINKAGE=HA
 NODE=M013M;LINKAGE=F2
 NODE=M013M3;LINKAGE=F
 NODE=M013M3;LINKAGE=E
 NODE=M013M3;LINKAGE=B
 NODE=M013M3;LINKAGE=C
 NODE=M013M3;LINKAGE=D
 NODE=M013M3;LINKAGE=SC
 NODE=M013M;LINKAGE=F1



PRODUCED IN $\bar{p}p$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.			

NODE=M013M9
 NODE=M013M9

1512 ± 4 OUR AVERAGE

1513 ± 4	AMSLER	06	CBAR	0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
1508 ± 9	¹ AMSLER	02	CBAR	0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1495.0 ± 1.1 ± 8.1	² ALBRECHT	20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
1530 ± 12	³ ANISOVICH	09	RVUE	0.0 $\bar{p}p, \pi N$

¹ T-matrix pole.

² T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

³ 4-poles, 5-channel K matrix fit.

NODE=M013M;LINKAGE=TT

NODE=M013M9;LINKAGE=A

NODE=M013M9;LINKAGE=AN

NODE=M013M4

NODE=M013M4

CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

1515±15

BARBERIS 99 OMEG 450 $p p \rightarrow p_s p_f K^+ K^-$

PRODUCED IN $e p$ COLLISIONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M013M10

NODE=M013M10

1512±3^{+1.4}_{-0.5} ¹ CHEKANOV 08 ZEUS $e p \rightarrow K_S^0 K_S^0 X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1537⁺⁹₋₈ 84 ² CHEKANOV 04 ZEUS $e p \rightarrow K_S^0 K_S^0 X$

¹ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

² Systematic errors not estimated.

NODE=M013M10;LINKAGE=HE

NODE=M013M10;LINKAGE=CH

$f_2'(1525)$ WIDTH

VALUE (MeV)	DOCUMENT ID	COMMENT
-------------	-------------	---------

86 ±5 OUR FIT Error includes scale factor of 2.2.

86.9^{+2.3}_{-2.1} PDG 18 Average of width measurements

NODE=M013WX

NODE=M013WX

PRODUCED BY PION BEAM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

84.4± 2.7 OUR AVERAGE Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.7. See the ideogram below. [86.9^{+2.3}_{-2.1} MeV OUR 2023 AVERAGE Scale factor = 1.4]

• • • We do not use the following data for averages, fits, limits, etc. • • •

102 ±42 TIKHOMIROV 03 SPEC 40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$

108 ⁺⁵₋₂ ¹ LONGACRE 86 MPS 22 $\pi^- p \rightarrow K_S^0 K_S^0 n$

69 ⁺²²₋₁₆ ² CHABAUD 81 ASPK 6 $\pi^- p \rightarrow K^+ K^- n$

137 ⁺²³₋₂₁ CHABAUD 81 ASPK 18.4 $\pi^- p \rightarrow K^+ K^- n$

OCCUR=2

150 ⁺⁸³₋₅₀ GORLICH 80 ASPK 17 $\pi^- p$ polarized $\rightarrow K^+ K^- n$

165 ±42 ³ CORDEN 79 OMEG 12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$

92 ⁺³⁹₋₂₂ ⁴ POLYCHRO... 79 STRC 7 $\pi^- p \rightarrow n K_S^0 K_S^0$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

² CHABAUD 81 is a reanalysis of PAWLICKI 77 data.

³ From an amplitude analysis where the $f_2'(1525)$ width and elasticity are in complete disagreement with the values obtained from $K\bar{K}$ channel, making the solution dubious.


⁴ From a fit to the D with $f_2(1270)$ - $f_2'(1525)$ interference. Mass fixed at 1516 MeV.

NODE=M013W;LINKAGE=L

NODE=M013W;LINKAGE=D

NODE=M013W;LINKAGE=N

NODE=M013W;LINKAGE=M



PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M013W2
NODE=M013W2

82 ± 6 OUR AVERAGE

90 ± 12		ASTON	88D	LASS	11	$K^- p \rightarrow K_S^0 K_S^0 \Lambda$
73 ± 18		BOLONKIN	86	SPEC	40	$K^- p \rightarrow K_S^0 K_S^0 \gamma$
83 ± 15		ARMSTRONG	83B	OMEG	18.5	$K^- p \rightarrow K^- K^+ \Lambda$
85 ± 16	650	AGUILAR-...	81B	HBC	4.2	$K^- p \rightarrow \Lambda K^+ K^-$
80 ⁺¹⁴ ₋₁₁	572	ALHARRAN	81	HBC	8.25	$K^- p \rightarrow \Lambda K \bar{K}$
72 ± 25	166	EVANGELIS...	77	OMEG	10	$K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
69 ± 22	100	AGUILAR-...	72B	HBC	3.9,4.6	$K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

92 ⁺²⁵ ₋₁₆	61	BINON	07	GAMS	32.5	$K^- p \rightarrow \eta \eta (\Lambda / \Sigma^0)$
75 ± 20		¹ BARKOV	99	SPEC	40	$K^- p \rightarrow K_S^0 K_S^0 \gamma$
62 ⁺¹⁹ ₋₁₄	123	BARREIRO	77	HBC	4.15	$K^- p \rightarrow \Lambda K_S^0 K_S^0$
61 ± 8	120	BRANDENB...	76C	ASPK	13	$K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$

¹ Systematic errors not estimated.

NODE=M013W2;LINKAGE=SK

PRODUCED IN $e^+ e^-$ ANNIHILATION AND PARTICLE DECAYS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M013W3
NODE=M013W3

86 ± 4 OUR AVERAGE Error includes scale factor of 2.4. See the ideogram below.


[89.2^{+3.4}_{-3.0} MeV OUR 2023 AVERAGE Scale factor = 1.8]

NEW

64.0 ± 4.3 ^{+2.0} _{-6.1}	126K	ABLIKIM	23M	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \pi^0$
84 ± 6 ⁺¹⁰ ₋₅		AAIJ	13AN	LHCB	$\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$
75 ⁺¹² ₋₁₀ ⁺¹⁶ ₋₈	5.5k	¹ ABLIKIM	13N	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
82.9 ^{+2.1} _{-2.2} ^{+3.3} _{-2.0}		UEHARA	13	BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
77 ± 15		ABLIKIM	05	BES2	$J/\psi \rightarrow \phi K^+ K^-$
82 ± 2 ± 3		ABE	04	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$

75 ± 4 ⁺¹⁵ / ₋₅		BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$	
100 ± 15	331	² ACCIARRI	01H L3	$91, 183-209 e^+ e^- \rightarrow e^+ e^- K_S^0 K_S^0$	
60 ± 20 ± 19		ABREU	96C DLPH	$Z^0 \rightarrow K^+ K^- + X$	
60 ± 23 ⁺¹³ / ₋₂₀		BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$	
103 ± 30		AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$	
100 ± 3		³ FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$	OCUR=2
85 ± 35		BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
78 ± 6		⁴ KLEMP	22 RVUE	$J/\psi(1S) \rightarrow \gamma \pi^0 \pi^0, \gamma K_S^0 K_S^0$	
84 ± 15		⁵ RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K})$	
37 ± 12	29	⁶ LEES	14H BABR	$e^+ e^- \rightarrow K_S^0 K_S^0 K^+ K^- \gamma$	
104 ± 10	870	⁷ SCHEGELSKY	06A RVUE	$\gamma \gamma \rightarrow K_S^0 K_S^0$	
62 ± 10		⁸ FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$	
¹ From partial wave analysis including all possible combinations of $0^{++}, 2^{++},$ and 4^{++} resonances. ² Supersedes ACCIARRI 95J. ³ From an analysis including interference with $f_0(1710)$. ⁴ Fit of the tensor partial waves from BES3 in the multipole basis. ⁵ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA). ⁶ From a fit to a Breit-Wigner line shape plus a second-order polynomial function. Systematic errors not evaluated. ⁷ From analysis of L3 data at 91 and 183–209 GeV. ⁸ From an analysis ignoring interference with $f_0(1710)$.					

NODE=M013W3;LINKAGE=A
 NODE=M013W;LINKAGE=HA
 NODE=M013W;LINKAGE=F2
 NODE=M013W3;LINKAGE=D
 NODE=M013W3;LINKAGE=C
 NODE=M013W3;LINKAGE=B
 NODE=M013W3;LINKAGE=SC
 NODE=M013W;LINKAGE=F1



PRODUCED IN $\bar{p}p$ ANNIHILATION

VALUE (MeV) DOCUMENT ID TECN COMMENT
 The data in this block is included in the average printed for a previous datablock.

77 ± 5 OUR AVERAGE

76 ± 6		AMSLER	06 CBAR	$0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$	
79 ± 8		¹ AMSLER	02 CBAR	$0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
104.8 ± 0.9 ± 9.8		² ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$	
128 ± 20		³ ANISOVICH	09 RVUE	$0.0 \bar{p}p, \pi N$	

¹ T-matrix pole.
² T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi \pi$), LONGACRE 86 ($K \bar{K}$), BINON 83 ($\eta \eta$).
³ K-matrix, 4-poles, 5-channel fit.

NODE=M013W9;LINKAGE=TT
 NODE=M013W9;LINKAGE=A
 NODE=M013W9;LINKAGE=AN

CENTRAL PRODUCTION

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ COMMENT _____
 The data in this block is included in the average printed for a previous datablock.

NODE=M013W4
 NODE=M013W4

70±25

BARBERIS 99 OMEG 450 $p p \rightarrow p_s p_f K^+ K^-$ **PRODUCED IN $e p$ COLLISIONS**

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____
 The data in this block is included in the average printed for a previous datablock.

NODE=M013W10
 NODE=M013W10

83± $\begin{smallmatrix} 9^{+5} \\ -4 \end{smallmatrix}$ ¹CHEKANOV 08 ZEUS $e p \rightarrow K_S^0 K_S^0 X$

••• We do not use the following data for averages, fits, limits, etc. •••

50 $\begin{smallmatrix} +34 \\ -22 \end{smallmatrix}$ 84 ²CHEKANOV 04 ZEUS $e p \rightarrow K_S^0 K_S^0 X$

¹In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.

²Systematic errors not estimated.

NODE=M013W10;LINKAGE=HE

NODE=M013W10;LINKAGE=CH

 $f_2'(1525)$ DECAY MODES

NODE=M013215;NODE=M013

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 $K \bar{K}$	(87.6±2.2) %	1.1
Γ_2 $\eta\eta$	(11.6±2.2) %	1.1
Γ_3 $\pi\pi$	(8.3±1.6) × 10 ⁻³	
Γ_4 $K \bar{K}^*(892) + \text{c.c.}$		
Γ_5 $\pi K \bar{K}$		
Γ_6 $\pi\pi\eta$		
Γ_7 $\pi^+\pi^+\pi^-\pi^-$		
Γ_8 $\gamma\gamma$	(9.5±1.1) × 10 ⁻⁷	1.1

DESIG=2

DESIG=4

DESIG=1

DESIG=3

DESIG=6

DESIG=5

DESIG=7

DESIG=8

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 2 partial widths, a combination of partial widths obtained from integrated cross sections, and 3 branching ratios uses 17 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 18.2$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100			
x_3	-6	-1		
x_8	-19	19	1	
Γ	-4	4	0	-44
	x_1	x_2	x_3	x_8

Mode	Rate (MeV)	Scale factor
Γ_1 $K \bar{K}$	75 ±4	1.8
Γ_2 $\eta\eta$	9.9 ±1.9	1.1
Γ_3 $\pi\pi$	0.71±0.14	1.1
Γ_8 $\gamma\gamma$	(8.2 ±0.9) × 10 ⁻⁵	

DESIG=2

DESIG=4

DESIG=1

DESIG=8

 $f_2'(1525)$ PARTIAL WIDTHS

NODE=M013220

 $\Gamma(K \bar{K})$ Γ_1

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ COMMENT _____
75±4 OUR FIT Error includes scale factor of 1.8.

NODE=M013W6
 NODE=M013W6

63± $\begin{smallmatrix} 6 \\ -5 \end{smallmatrix}$ ¹LONGACRE 86 MPS 22 $\pi^- p \rightarrow K_S^0 K_S^0 n$

¹From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

NODE=M013PW;LINKAGE=L

$\Gamma(\eta\eta)$ Γ_2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.9±0.19 OUR FIT Error includes scale factor of 1.1.

••• We do not use the following data for averages, fits, limits, etc. •••

5.0±0.8	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
24 $\begin{smallmatrix} +3 \\ -1 \end{smallmatrix}$		² LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.² From a partial-wave analysis of data using a K-matrix formalism with 5 poles.NODE=M013W7
NODE=M013W7

NODE=M013W7;LINKAGE=SC

NODE=M013PW7;LINKAGE=L

 $\Gamma(\pi\pi)$ Γ_3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.71±0.14 OUR FIT Error includes scale factor of 1.1.

1.4 $\begin{smallmatrix} +1.0 \\ -0.5 \end{smallmatrix}$		¹ LONGACRE 86	MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
--	--	--------------------------	-----	--

••• We do not use the following data for averages, fits, limits, etc. •••

0.2 $\begin{smallmatrix} +1.0 \\ -0.2 \end{smallmatrix}$	870	² SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
--	-----	-----------------------------	------	--

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.² From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.NODE=M013W5
NODE=M013W5

NODE=M013PW5;LINKAGE=L

NODE=M013W5;LINKAGE=SC

 $\Gamma(\gamma\gamma)$ Γ_8

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.082±0.009 OUR FIT

••• We do not use the following data for averages, fits, limits, etc. •••

0.13 ±0.03	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$
------------	-----	-----------------------------	------	--

¹ From analysis of L3 data at 91 and 183–209 GeV, using $\Gamma(f_2'(1525) \rightarrow K\bar{K}) = 68$ MeV and SU(3) relations.NODE=M013W8
NODE=M013W8

NODE=M013W8;LINKAGE=SC

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.746±0.002 $\begin{smallmatrix} +0.166 \\ -0.162 \end{smallmatrix}$	¹ ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
--	--------------------------	------	---

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).NODE=M013R00
NODE=M013R00

NODE=M013R00;LINKAGE=A

 $f_2'(1525) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M013223

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_8/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

0.072 ±0.007 OUR FIT**0.072 ±0.007 OUR AVERAGE**

0.048 $\begin{smallmatrix} +0.067 \\ -0.008 \end{smallmatrix}$ $\begin{smallmatrix} +0.108 \\ -0.012 \end{smallmatrix}$	UEHARA 13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
0.0564±0.0048±0.0116	ABE 04	BELL	10.6 $e^+e^- \rightarrow e^+e^- K^+K^-$
0.076 ±0.006 ±0.011	331 ACCIARRI 01H	L3	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.067 ±0.008 ±0.015	² ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
0.11 $\begin{smallmatrix} +0.03 \\ -0.02 \end{smallmatrix}$ ±0.02	BEHREND 89c	CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.10 $\begin{smallmatrix} +0.04 \\ -0.03 \end{smallmatrix}$ $\begin{smallmatrix} +0.03 \\ -0.02 \end{smallmatrix}$	BERGER 88	PLUT	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.12 ±0.07 ±0.04	² AIHARA 86B	TPC	$e^+e^- \rightarrow e^+e^- K^+K^-$
0.11 ±0.02 ±0.04	² ALTHOFF 83	TASS	$e^+e^- \rightarrow e^+e^- K\bar{K}$

••• We do not use the following data for averages, fits, limits, etc. •••

0.0314±0.0050±0.0077	³ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^- K^+K^-$
----------------------	---------------------------	-----	------------------------------------

¹ Supersedes ACCIARRI 95J. From analysis of L3 data at 91 and 183–209 GeV,² Using an incoherent background.³ Using a coherent background.NODE=M013G1
NODE=M013G1

OCCUR=2

NODE=M013G;LINKAGE=HA

NODE=M013G1;LINKAGE=A

NODE=M013G1;LINKAGE=B

$f'_2(1525)$ BRANCHING RATIOS $\Gamma(\eta\eta)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.059 ± 0.003 ± 0.026	¹ ALBRECHT 20	RVUE	0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
seen	UEHARA 10A	BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \eta \eta$
0.10 ± 0.03	² PROKOSHKIN 91	GAM4 300	$\pi^- p \rightarrow \pi^- p \eta \eta$

¹ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

² Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.

NODE=M013225

NODE=M013R8
NODE=M013R8

NODE=M013R8;LINKAGE=A

NODE=M013R8;LINKAGE=B

 $\Gamma(\eta\eta)/\Gamma(K\bar{K})$ Γ_2/Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.132 ± 0.028 OUR FIT**0.115 ± 0.028 OUR AVERAGE**

0.119 ± 0.015 ± 0.036	61	¹ BINON	07	GAMS	32.5 $K^- p \rightarrow \eta\eta(\Lambda/\Sigma^0)$
0.11 ± 0.04		² PROKOSHKIN 91	GAM4 300		$\pi^- p \rightarrow \pi^- p \eta \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.14	90	BARBERIS	00E		450 $pp \rightarrow p_f \eta \eta p_s$
< 0.50		BARNES	67	HBC	4.6, 5.0 $K^- p$

¹ Using the compilation of the cross sections for $f'_2(1525)$ production in $K^- p$ collisions from ASTON 88D.

² Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma\eta\eta$.

NODE=M013R3
NODE=M013R3

NODE=M013R3;LINKAGE=BI

NODE=M013R3;LINKAGE=B

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

0.83 ± 0.16 OUR FIT**0.75 ± 0.16 OUR AVERAGE**

0.7 ± 0.2		COSTA	80	OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
2.7 ^{+7.1} _{-1.3}		¹ GORLICH	80	ASPK	17, 18 $\pi^- p$
0.75 ± 0.25		^{1,2} MARTIN	79	RVUE	
3.4 ± 1.5 ± 1.0		³ ALBRECHT 20	RVUE		0.9 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
< 6	95	AGUILAR-...	81B	HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
19 ± 3		CORDEN	79	OMEG	12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
< 4.5	95	BARREIRO	77	HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
1.2 ± 0.4		¹ PAWLICKI	77	SPEC	6 $\pi N \rightarrow K^+ K^- N$
< 6.3	90	BRANDENB...	76C	ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
< 0.86		¹ BEUSCH	75B	OSPK	8.9 $\pi^- p \rightarrow K^0 \bar{K}^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assuming that the $f'_2(1525)$ is produced by an one-pion exchange production mechanism.

² MARTIN 79 uses the PAWLICKI 77 data with different input value of the $f'_2(1525) \rightarrow K\bar{K}$ branching ratio.

³ Residue from T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

NODE=M013R1
NODE=M013R1

NODE=M013R1;LINKAGE=C

NODE=M013R1;LINKAGE=D

NODE=M013R1;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$ Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0094 ± 0.0018 OUR FIT**0.075 ± 0.035**AUGUSTIN 87 DM2 $J/\psi \rightarrow \gamma\pi^+\pi^-$ NODE=M013R7
NODE=M013R7 $[\Gamma(K\bar{K}^*(892) + \text{c.c.}) + \Gamma(\pi K\bar{K})]/\Gamma(K\bar{K})$ $(\Gamma_4 + \Gamma_5)/\Gamma_1$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.35	95	AGUILAR-...	72B	HBC	3.9, 4.6 $K^- p$
< 0.4	67	AMMAR	67	HBC	

NODE=M013R5
NODE=M013R5

$\Gamma(\pi\pi\eta)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_1
-------	-----	-------------	------	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.41	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
<0.3	67	AMMAR	67	HBC	

NODE=M013R4
NODE=M013R4

 $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_1
-------	-----	-------------	------	---------	---------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.32	95	AGUILAR-...	72B	HBC	3.9,4.6 $K^- p$
-------	----	-------------	-----	-----	-----------------

NODE=M013R6
NODE=M013R6

 $f'_2(1525)$ REFERENCES

NODE=M013

ABLIKIM	23M	JHEP 2303 121	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62055
KLEMP	22	PL B830 137171	E. Klempt <i>et al.</i>	(BONN)	REFID=61646
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
AAIJ	13AN	PR D87 072004	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55137
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
BINON	07	PAN 70 1713	F. Binon <i>et al.</i>	(GAMS Collab.)	REFID=52057
		Translated from YAF 70 1758.			
AMSLER	06	PL B639 165	C. Amstler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
CHEKANOV	04	PL B578 33	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=49672
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
AMSLER	02	EPJ C23 29	C. Amstler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARKOV	99	JETPL 70 248	B.P. Barkov <i>et al.</i>		REFID=47379
		Translated from ZETFP 70 242.			
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
ACCIARRI	95J	PL B363 118	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44615
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
ASTON	88D	NP B301 525	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40330
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BERGER	88	ZPHY C37 329	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=40566
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAITIS...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
AIHARA	86B	PRL 57 404	H. Aihara <i>et al.</i>	(TPC-2 γ Collab.)	REFID=20764
BOLONKIN	86	SJNP 43 776	B.V. Bolonkin <i>et al.</i>	(ITEP) JP	REFID=44646
		Translated from YAF 43 1211.			
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	83	PL 121B 216	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21408
ARMSTRONG	83B	NP B224 193	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=20558
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
AGUILAR-...	81B	ZPHY C8 313	M. Aguilar-Benitez <i>et al.</i>	(CERN, CDEF+)	REFID=21104
ALHARRAN	81	NP B191 26	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=21403
CHABAUD	81	APP B12 575	V. Chabaud <i>et al.</i>	(CERN, CRAC, MPIM)	REFID=20742
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
MARTIN	79	NP B158 520	A.D. Martin, E.N. Ozmutlu	(DURH)	REFID=20377
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)	REFID=20378
BARREIRO	77	NP B121 237	F. Barreiro <i>et al.</i>	(CERN, AMST, NIJM+)	REFID=21392
EVANGELIS...	77	NP B127 384	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20540
PAWLICKI	77	PR D15 3196	A.J. Pawlicki <i>et al.</i>	(ANL) IJP	REFID=20367
BRANDENB...	76C	NP B104 413	G.W. Brandenburg <i>et al.</i>	(SLAC)	REFID=20225
BEUSCH	75B	PL 60B 101	W. Beusch <i>et al.</i>	(CERN, ETH)	REFID=21390
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
AGUILAR-...	72B	PR D6 29	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=20205
AMMAR	67	PRL 19 1071	R. Ammar <i>et al.</i>	(NWES, ANL) JP	REFID=21382
BARNES	67	PRL 19 964	V.E. Barnes <i>et al.</i>	(BNL, SYRA) IJPC	REFID=21383
CRENNELL	66	PRL 16 1025	D.J. Crennell <i>et al.</i>	(BNL) I	REFID=20317

$f_2(1565)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Seen mostly in antinucleon-nucleon annihilation. Needs confirmation in other channels. See the review on "Spectroscopy of Light Meson Resonances."

NODE=M123

NODE=M123

 $f_2(1565)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M123PP

NODE=M123PP

NODE=M123PP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1495–1560) – i (40–110) OUR ESTIMATE			
$(1560 \pm 15) - i (140 \pm 20)$	¹ ANISOVICH	09	RVUE 0.0 $\bar{p}p, \pi N$
$(1552 \pm 13) - i (57 \pm 12)$	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta,$ $\pi^0 \pi^0 \pi^0$
$(1507 \pm 15) - i (65 \pm 10)$	BERTIN	97C	OBLX 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1534 \pm 20) - i (90 \pm 30)$	² ABELE	96C	RVUE Compilation
$(\sim 1552) - i (\sim 71)$	³ AMSLER	95D	CBAR 0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0,$ $\pi^0 \eta \eta, \pi^0 \pi^0 \eta$

¹ On sheet II in a two-pole solution.² T-matrix pole, large coupling to $\rho\rho$ and $\omega\omega$, could be $f_2(1640)$.³ Coupled-channel analysis of AMSLER 95B, AMSLER 95C, and AMSLER 94D.

NODE=M123PP;LINKAGE=A

NODE=M123PP;LINKAGE=D

NODE=M123PP;LINKAGE=E

 $f_2(1565)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1575±18	⁴ BERTIN	98	OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1590±10	^{5,6} AMELIN	06	VES 36 $\pi^- p \rightarrow \omega \omega n$
1550±10±20	⁶ AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1598±11±9	BAKER	99B	SPEC 0 $\bar{p}p \rightarrow \omega \omega \pi^0$
1598±72	BALOSHIN	95	SPEC 40 $\pi^- C \rightarrow K_S^0 K_S^0 X$
1566 ⁺⁸⁰ ₋₅₀	⁷ ANISOVICH	94	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0, \eta \eta \pi^0$
1502±9	ADAMO	93	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1488±10	⁸ ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1508±10	⁸ ARMSTRONG	93D	E760 $\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
1525±10	⁸ ARMSTRONG	93D	E760 $\bar{p}p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 1504	⁹ WEIDENAUER	93	ASTE 0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$
1540±15	⁸ ADAMO	92	OBLX $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1515±10	¹⁰ AKER	91	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0$
1565±20	MAY	90	ASTE 0.0 $\bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
1477±5	BRIDGES	86C	DBC 0.0 $\bar{p}N \rightarrow 3\pi^- 2\pi^+$

⁴ Breit-Wigner mass.⁵ Supersedes the $\omega\omega$ state of BELADIDZE 92B earlier assigned to the $f_2(1640)$.⁶ Breit-Wigner width.⁷ From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta \eta$ including AKER 91 data.⁸ J^P not determined, could be partly $f_0(1500)$.⁹ J^P not determined.¹⁰ Superseded by AMSLER 95B.

NODE=M123M

NODE=M123M

OCCUR=2

NODE=M123M;LINKAGE=G

NODE=M123M;LINKAGE=AM

NODE=M123M;LINKAGE=D

NODE=M123M;LINKAGE=C

NODE=M123M;LINKAGE=E

NODE=M123M;LINKAGE=F

NODE=M123M;LINKAGE=BA

 $f_2(1565)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
119±24	¹¹ BERTIN	98	OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$

NODE=M123W

NODE=M123W

- • • We do not use the following data for averages, fits, limits, etc. • • •

140± 11	^{12,13} AMELIN	06	VES	36	$\pi^- p \rightarrow \omega \omega n$
130± 20±40	¹³ AMELIN	00	VES	37	$\pi^- p \rightarrow \eta \pi^+ \pi^- n$
263±101	BALOSHIN	95	SPEC	40	$\pi^- C \rightarrow K_S^0 K_S^0 X$
166 ⁺ ₋₂₀	¹⁴ ANISOVICH	94	CBAR	0.0	$\bar{p} p \rightarrow 3\pi^0, \eta \eta \pi^0$
130± 10	¹⁵ ADAMO	93	OBLX		$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
148± 27	¹⁶ ARMSTRONG	93C	E760		$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
103± 15	¹⁶ ARMSTRONG	93D	E760		$\bar{p} p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111± 10	¹⁶ ARMSTRONG	93D	E760		$\bar{p} p \rightarrow \eta \pi^0 \pi^0 \rightarrow 6\gamma$
~ 206	¹⁷ WEIDENAUER	93	ASTE	0.0	$\bar{p} N \rightarrow 3\pi^- 2\pi^+$
132± 37	¹⁶ ADAMO	92	OBLX		$\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
120± 10	¹⁸ AKER	91	CBAR	0.0	$\bar{p} p \rightarrow 3\pi^0$
170± 40	MAY	90	ASTE	0.0	$\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$
116± 9	BRIDGES	86C	DBC	0.0	$\bar{p} N \rightarrow 3\pi^- 2\pi^+$

¹¹ Breit-Wigner width.

¹² Supersedes the $\omega \omega$ state of BELADIDZE 92B earlier assigned to the $f_2(1640)$.

¹³ Breit-Wigner width.

¹⁴ From a simultaneous analysis of the annihilations $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta \eta$ including AKER 91 data.

¹⁵ Supersedes ADAMO 92.

¹⁶ J^P not determined, could be partly $f_0(1500)$.

¹⁷ J^P not determined.

¹⁸ Superseded by AMSLER 95B.

OCCUR=2

NODE=M123W;LINKAGE=G
 NODE=M123W;LINKAGE=AM
 NODE=M123W;LINKAGE=H
 NODE=M123W;LINKAGE=D

NODE=M123W;LINKAGE=C
 NODE=M123W;LINKAGE=E
 NODE=M123W;LINKAGE=F
 NODE=M123W;LINKAGE=BA

$f_2(1565)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi \pi$	seen
Γ_2 $\pi^+ \pi^-$	seen
Γ_3 $\pi^0 \pi^0$	seen
Γ_4 $\rho^0 \rho^0$	seen
Γ_5 $2\pi^+ 2\pi^-$	seen
Γ_6 $\eta \eta$	seen
Γ_7 $\omega \omega$	seen
Γ_8 $K \bar{K}$	seen
Γ_9 $\gamma \gamma$	seen

NODE=M123215;NODE=M123

DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=10;OUR EST;→ UNCHECKED ←

$f_2(1565)$ PARTIAL WIDTHS

$\Gamma(\eta \eta)$

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT Γ_6

- • • We do not use the following data for averages, fits, limits, etc. • • •

1.2±0.3 870 ¹⁹ SCHEGELSKY 06A RVUE $\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M123225

NODE=M123W3
 NODE=M123W3

$\Gamma(K \bar{K})$

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT Γ_8

- • • We do not use the following data for averages, fits, limits, etc. • • •

2.0±1.0 870 ¹⁹ SCHEGELSKY 06A RVUE $\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M123W1
 NODE=M123W1

$\Gamma(\gamma \gamma)$

VALUE (keV) EVTS DOCUMENT ID TECN COMMENT Γ_9

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.70±0.14 870 ¹⁹ SCHEGELSKY 06A RVUE $\gamma \gamma \rightarrow K_S^0 K_S^0$

NODE=M123W2
 NODE=M123W2

¹⁹ From analysis of L3 data at 91 and 183–209 GeV, using $f_2(1565)$ mass of 1570 MeV, width of 160 MeV, $\Gamma(\pi \pi) = 25$ MeV, and SU(3) relations.

NODE=M123W1;LINKAGE=SC

$f_2(1565)$ BRANCHING RATIOS

$\Gamma(\pi \pi)/\Gamma_{\text{total}}$

VALUE DOCUMENT ID TECN COMMENT Γ_1/Γ

- • • We do not use the following data for averages, fits, limits, etc. • • •

seen BAKER 99B SPEC $0 \bar{p} p \rightarrow \omega \omega \pi^0$

NODE=M123220

NODE=M123R5
 NODE=M123R5

$\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R1
 NODE=M123R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BERTIN	98	OBLX 0.05–0.405 $\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
not seen	²⁰ ANISOVICH	94B	RVUE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
seen	MAY	89	ASTE $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$

²⁰ ANISOVICH 94B is from a reanalysis of MAY 90.

NODE=M123R1;LINKAGE=A

 $\Gamma(\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R3
 NODE=M123R3

seen	AMSLER	95B	CBAR 0.0 $\bar{p}p \rightarrow 3\pi^0$
------	--------	-----	--

 $\Gamma(\pi^+\pi^-)/\Gamma(\rho^0\rho^0)$ Γ_2/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R2
 NODE=M123R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042±0.013	BRIDGES	86B	DBC $\bar{p}N \rightarrow 3\pi^-2\pi^+$
-------------	---------	-----	---

 $\Gamma(\eta\eta)/\Gamma(\pi^0\pi^0)$ Γ_6/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R4
 NODE=M123R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.024±0.005±0.012	²¹ ARMSTRONG	93C	E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
-------------------	-------------------------	-----	---

²¹ J^P not determined, could be partly $f_0(1500)$.

NODE=M123R4;LINKAGE=E

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M123R6
 NODE=M123R6

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	BAKER	99B	SPEC 0 $\bar{p}p \rightarrow \omega\omega\pi^0$
------	-------	-----	---

 $f_2(1565)$ REFERENCES

NODE=M123

ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=51574
		Translated from YAF 69 715.			
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
BAKER	99B	PL B467 147	C.A. Baker <i>et al.</i>		REFID=47398
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45782
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45076
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44377
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44440
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44441
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=44621
		Translated from YAF 58 50.			
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=44093
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.)	REFID=43659
ANISOVICH	94B	PR D50 1972	V.V. Anisovich <i>et al.</i>	(LOQM)	REFID=44071
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.)	REFID=43657
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43596
WEIDENAUER	93	ZPHY C59 387	P. Weidenauer <i>et al.</i>	(ASTERIX Collab.)	REFID=43585
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)	REFID=42177
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42172
AKER	91	PL B260 249	E. Aker <i>et al.</i>	(Crystal Barrel Collab.)	REFID=41587
MAY	90	ZPHY C46 203	B. May <i>et al.</i>	(ASTERIX Collab.)	REFID=41365
MAY	89	PL B225 450	B. May <i>et al.</i>	(ASTERIX Collab.) IJP	REFID=40921
BRIDGES	86B	PRL 56 215	D.L. Bridges <i>et al.</i>	(SYRA, CASE)	REFID=21376
BRIDGES	86C	PRL 57 1534	D.L. Bridges <i>et al.</i>	(SYRA)	REFID=21377

$\rho(1570)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

May be an OZI-violating decay mode of $\rho(1700)$. See the review on "Spectroscopy of Light Meson Resonances."

NODE=M188

NODE=M188

 $\rho(1570)$ MASS

NODE=M188M

NODE=M188M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1570±36±62	54	¹ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••				
1614±2		² ACHASOV	23A SND	$e^+e^- \rightarrow \omega\pi^0$
1585±15		³ ACHASOV	20C SND	1.3–2.0 $e^+e^- \rightarrow K^+K^-\pi^0$
1480±40		⁴ BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$

¹ From the fit with two resonances.² From a vector dominance fit to the Born cross section between 1.05 and 2.0 GeV with $\rho(770)$, $\rho(1570)$, $\rho(1700)$, $\rho(2150)$. The fit also uses SND data from the VEPP-2M collider below 1.02 GeV and from LEES 17H and ABLIKIM 21A above 1.5 GeV.³ From a fit using a two resonance model in which the mass and width of the other resonance are fixed at the $\rho(1700)$ values from PDG 20.⁴ Systematic errors not estimated.NODE=M188M;LINKAGE=AU
NODE=M188M;LINKAGE=B

NODE=M188M;LINKAGE=A

NODE=M188M;LINKAGE=BI

 $\rho(1570)$ WIDTH

NODE=M188W

NODE=M188W

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
140±90 OUR AVERAGE		[144 ± 90 MeV OUR 2023 AVERAGE]		
144±75±43	54	¹ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
••• We do not use the following data for averages, fits, limits, etc. •••				
492±4		² ACHASOV	23A SND	$e^+e^- \rightarrow \omega\pi^0$
75±30		³ ACHASOV	20C SND	1.3–2.0 $e^+e^- \rightarrow K^+K^-\pi^0$
130±60		⁴ BITYUKOV	87 SPEC	32.5 $\pi^-p \rightarrow \phi\pi^0n$

¹ From the fit with two resonances.² From a vector dominance fit to the Born cross section between 1.05 and 2.0 GeV with $\rho(770)$, $\rho(1570)$, $\rho(1700)$, $\rho(2150)$. The fit also uses SND data from the VEPP-2M collider below 1.02 GeV and from LEES 17H and ABLIKIM 21A above 1.5 GeV.³ From a fit using a two resonance model in which the mass and width of the other resonance are fixed at the $\rho(1700)$ values from PDG 20.⁴ Systematic errors not estimated.NODE=M188W;LINKAGE=AU
NODE=M188W;LINKAGE=B

NODE=M188W;LINKAGE=A

NODE=M188W;LINKAGE=BI

 $\rho(1570)$ DECAY MODES

NODE=M188215;NODE=M188

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	
Γ_2 $\phi\pi$	not seen
Γ_3 $\omega\pi$	

DESIG=1

DESIG=2

DESIG=3

 $\rho(1570)$ $\Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M188225

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
3.5±0.9±0.3		54	¹ AUBERT	08S BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$	
••• We do not use the following data for averages, fits, limits, etc. •••						
<70	90		² AULCHENKO	87B ND	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$	

¹ From the fit with two resonances.² Using mass and width of BITYUKOV 87.NODE=M188G01;LINKAGE=AU
NODE=M188G01;LINKAGE=AL **$\rho(1570)$ BRANCHING RATIOS**

NODE=M188220

$\Gamma(\phi\pi)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
not seen		ABELE	97H CBAR	$\bar{p}p \rightarrow K_L^0 K_S^0 \pi^0 \pi^0$	
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.01		¹ DONNACHIE	91 RVUE		

¹ Using data from BISELLO 91B, DOLINSKY 86, and ALBRECHT 87L.NODE=M188R01
NODE=M188R01

NODE=M188R01;LINKAGE=DO

$\Gamma(\phi\pi)/\Gamma(\omega\pi)$ Γ_2/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M188R02
 NODE=M188R02

••• We do not use the following data for averages, fits, limits, etc. •••

>0.5	95	BITYUKOV	87	SPEC	32.5 $\pi^- p \rightarrow \phi\pi^0 n$
------	----	----------	----	------	--

 $\rho(1570)$ REFERENCES

NODE=M188

ACHASOV	23A	PR D108 092012	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=62428
ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61028
ACHASOV	20C	EPJ C80 1139	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60935
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
LEES	17H	PR D96 092009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58311
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
ABELE	97H	PL B415 280	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45765
BISELLO	91B	NPBPS B21 111	D. Bisello	(DM2 Collab.)	REFID=41752
DONNACHIE	91	ZPHY C51 689	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=41632
ALBRECHT	87L	PL B185 223	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40418
AULCHENKO	87B	JETPL 45 145	V.M. Aulchenko <i>et al.</i>	(NOVO)	REFID=41373
		Translated from ZETFP	45 118.		
BITYUKOV	87	PL B188 383	S.I. Bityukov <i>et al.</i>	(SERP)	REFID=40011
DOLINSKY	86	PL B174 453	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=20246

NODE=M166

 $h_1(1595)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

OMITTED FROM SUMMARY TABLE

Seen in a partial-wave analysis of the $\omega\eta$ system produced in the reaction $\pi^- p \rightarrow \omega\eta n$ at 18 GeV/c.

NODE=M166

 $h_1(1595)$ MASS

NODE=M166M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$1594 \pm 15^{+10}_{-60}$	EUGENIO	01	SPEC 18 $\pi^- p \rightarrow \omega\eta n$

NODE=M166M

 $h_1(1595)$ WIDTH

NODE=M166W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
380^{+90}_{-120} OUR AVERAGE	[384 ⁺ ₋₁₂₀ MeV OUR 2023 AVERAGE]		
$384 \pm 60^{+70}_{-100}$	EUGENIO	01	SPEC 18 $\pi^- p \rightarrow \omega\eta n$

NODE=M166W

NEW

 $h_1(1595)$ DECAY MODES

NODE=M166215;NODE=M166

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \omega\eta$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

 $h_1(1595)$ REFERENCES

NODE=M166

EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>
---------	----	-------------	--------------------------

REFID=48010

$\pi_1(1600)$

$$I^G(J^{PC}) = 1^-(1^-+)$$

Coupled channel analyses favor the existence of only one broad 1^-+ isovector state consistent with $\pi_1(1600)$ in the 1400–1600 MeV region. Accordingly, the $\pi_1(1400)$ entries of the previous Reviews have been moved into this section. See the review on "Spectroscopy of Light Meson Resonances."

NODE=M164

NODE=M164

 $\pi_1(1600)$ T-Matrix Pole \sqrt{s}

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M164TMP

NODE=M164TMP

NODE=M164TMP

→ UNCHECKED ←

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1480–1680) – i (150–300) OUR ESTIMATE			
$(1623 \pm 47^{+24}_{-75}) - i(228 \pm 44^{+72}_{-88})$	¹ KOPF	21	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
$(1564 \pm 24 \pm 86) - i(246 \pm 27 \pm 51)$	² RODAS	19	RVUE 191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$(1405 \pm 4^{+15}_{-18}) - i(314 \pm 14^{+18}_{-69})$	³ ALBRECHT	20	RVUE $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.			
² The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data.			
³ Superseded by KOPF 21.			

NODE=M164TMP;LINKAGE=B

NODE=M164TMP;LINKAGE=A
NODE=M164TMP;LINKAGE=AL **$\pi_1(1600)$ MASS ($\eta\pi$ mode)**

Not seen by PROKOSHKIN 95B, BUGG 94, APEL 81, BOUTEMEUR 90, and AGHASYAN 18B.

NODE=M164MEP

NODE=M164MEP


NODE=M164MEP

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1354 ±25	OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			
1257 ±20 ±25	23.5k	ADAMS	07B	B852	18 $\pi^- p \rightarrow \eta \pi^0 n$
1384 ±20 ±35	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
1360 ±25		ABELE	99	CBAR	0.0 $\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
1400 ±20 ±20		ABELE	98B	CBAR	0.0 $\bar{p}n \rightarrow \pi^- \pi^0 \eta$
1370 ±16 $^{+50}_{-30}$		¹ THOMPSON	97	MPS	18 $\pi^- p \rightarrow \eta \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1323.1 ± 4.6		² AOYAGI	93	BKEI	$\pi^- p \rightarrow \eta \pi^- p$
1406 ±20		³ ALDE	88B	GAM4 0	100 $\pi^- p \rightarrow \eta \pi^0 n$
¹ Natural parity exchange, questioned by DZIERBA 03.					
² Unnatural parity exchange.					
³ Seen in the P_0 -wave intensity of the $\eta\pi^0$ system, unnatural parity exchange.					

NODE=M164MEP;LINKAGE=B

NODE=M164MEP;LINKAGE=C

NODE=M164MEP;LINKAGE=A



NODE=M164M

NODE=M164M

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1645⁺⁴⁰₋₁₇ OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

[1661⁺¹⁵₋₁₁ MeV OUR 2023 AVERAGE Scale factor = 1.2]

1600 ⁺¹¹⁰ ₋₆₀	46M	1 AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1709 ± 24 ± 41	69k	2 KUHN	04	B852	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$
1597 ± 10 ⁺⁴⁵ ₋₁₀		2 IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1660 ± 10 ⁺⁰ ₋₆₄	420k	3 ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
1664 ± 8 ± 10	145k	4 LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1593 ± 8 ⁺²⁹ ₋₄₇		2,5 ADAMS	98B	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

¹ Statistical error negligible. See also the review ALEXEEV 22.


² Natural parity exchange.

³ Superseded by AGHASYAN 2018B.

⁴ May be a different state: natural and unnatural parity exchanges.

⁵ Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with 2.6 M events of $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data.

NODE=M164M;LINKAGE=B
 NODE=M164M;LINKAGE=A
 NODE=M164M;LINKAGE=C
 NODE=M164M;LINKAGE=LU
 NODE=M164M;LINKAGE=DZ



$\pi_1(1600)$ WIDTH ($\eta\pi$ mode)

Not seen by PROKOSHKIN 95B, BUGG 94, APEL 81, BOUTEMEUR 90, and AGHASYAN 18B.

NODE=M164WEP

NODE=M164WEP

NODE=M164WEP

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
330 ± 35	OUR AVERAGE				
354 ± 64 ± 58	23.5k	ADAMS	07B	B852	18 $\pi^- p \rightarrow \eta\pi^0 n$
378 ± 50 ± 50	90k	SALVINI	04	OBLX	$\bar{p}p \rightarrow 2\pi^+ 2\pi^-$
220 ± 90		ABELE	99	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\pi^0\eta$
310 ± 50 + 50 - 30		ABELE	98B	CBAR	0.0 $\bar{p}n \rightarrow \pi^- \pi^0\eta$
385 ± 40 + 65 - 105		¹ THOMPSON	97	MPS	18 $\pi^- p \rightarrow \eta\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

143.2 ± 12.5		² AOYAGI	93	BKEI	$\pi^- p \rightarrow \eta\pi^- p$
180 ± 20		³ ALDE	88B	GAM4 0	100 $\pi^- p \rightarrow \eta\pi^0 n$

¹ Resolution is not unfolded, natural parity exchange, questioned by DZIERBA 03.

² Unnatural parity exchange.

³ Seen in the P_0 -wave intensity of the $\eta\pi^0$ system, unnatural parity exchange.

NODE=M164WEP;LINKAGE=QQ

NODE=M164WEP;LINKAGE=C

NODE=M164WEP;LINKAGE=A

NODE=M164W

$\pi_1(1600)$ WIDTH (non- $\eta\pi$ mode)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
370⁺₋₆₀	OUR AVERAGE			

[240 ± 50 MeV OUR 2023 AVERAGE Scale factor = 1.7]

580 ⁺ ₋₂₃₀	46M	¹ AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
403 ± 80 ± 115	69k	² KUHN	04	B852	18 $\pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$
340 ± 40 ± 50		² IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

269 ± 21 ⁺ ₋₆₄	420k	³ ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
185 ± 25 ± 28	145k	⁴ LU	05	B852	18 $\pi^- p \rightarrow \omega\pi^- \pi^0 p$
168 ± 20 ⁺ ₋₁₂		^{2,5} ADAMS	98B	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$

¹ Statistical error negligible. See also the review ALEXEEV 22.

² Natural parity exchange.

³ Superseded by AGHASYAN 2018B.

⁴ May be a different state: natural and unnatural parity exchanges.

⁵ Superseded by DZIERBA 06 excluding this state in a more refined PWA analysis, with 2.6 M events of $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data.

NODE=M164W

NEW

NODE=M164W;LINKAGE=B

NODE=M164W;LINKAGE=A

NODE=M164W;LINKAGE=C

NODE=M164W;LINKAGE=LU

NODE=M164W;LINKAGE=DZ

$\pi_1(1600)$ DECAY MODES

NODE=M164215;NODE=M164

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi\pi$	seen
Γ_2 $\rho^0\pi^-$	seen
Γ_3 $f_2(1270)\pi^-$	not seen
Γ_4 $b_1(1235)\pi$	seen
Γ_5 $\eta'(958)\pi^-$	seen
Γ_6 $\eta\pi$	seen
Γ_7 $f_1(1285)\pi$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2
DESIG=4
DESIG=5
DESIG=3
DESIG=7;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←

 $\pi_1(1600)$ BRANCHING RATIOS

NODE=M164220

$\Gamma(\rho^0\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen ALEKSEEV 10 COMP 190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen NOZAR 09 CLAS $\gamma p \rightarrow 2\pi^+ \pi^- n$

not seen ¹ DZIERBA 06 B852 18 $\pi^- p$

¹ From the PWA analysis of 2.6 M $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data. Supersedes ADAMS 98B.

NODE=M164R1
NODE=M164R1

NODE=M164R1;LINKAGE=DZ

$\Gamma(f_2(1270)\pi^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ¹ DZIERBA 06 B852 18 $\pi^- p$

¹ From the PWA analysis of 2.6 M $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$ and 3 M events of $\pi^- p \rightarrow \pi^- \pi^0 \pi^0 p$ of E852 data. Supersedes CHUNG 02.

NODE=M164R3
NODE=M164R3

NODE=M164R3;LINKAGE=DZ

$\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 35280 ¹ BAKER 03 SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 145k LU 05 B852 18 $\pi^- p \rightarrow \omega\pi^-\pi^0 p$

¹ $B((b_1\pi)_{D\text{-wave}})/B((b_1\pi)_{S\text{-wave}})=0.3 \pm 0.1$.

NODE=M164R4
NODE=M164R4

NODE=M164R;LINKAGE=RB

$\Gamma(\eta'(958)\pi^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen IVANOV 01 B852 18 $\pi^- p \rightarrow \eta'\pi^- p$

NODE=M164R2
NODE=M164R2

$\Gamma(\eta'(958)\pi^-)/\Gamma(\eta\pi)$ Γ_5/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.54 \pm 1.1^{+1.8}_{-0.27}$ ¹ KOPF 21 RVUE 0.9 $p\bar{p} \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta,$
 $\pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow$
 $\pi^- \pi^- \pi^+ p$

¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

NODE=M164R00
NODE=M164R00

NODE=M164R00;LINKAGE=A

$\Gamma(f_1(1285)\pi)/\Gamma(\eta'(958)\pi^-)$ Γ_7/Γ_5

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

3.80 ± 0.78 69k ¹ KUHN 04 B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

¹ Using $\eta'(958)\pi$ data from IVANOV 01.

NODE=M164R5
NODE=M164R5

NODE=M164R;LINKAGE=KU

$\pi_1(1600)$ REFERENCES

ALEXEEV	22	PR D105 012005	G.D. Alexeev <i>et al.</i>	(COMPASS Collab.)	NODE=M164
KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)	REFID=61491
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=61470
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=60439
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59554
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=56385
NOZAR	09	PRL 102 102002	M. Nozar <i>et al.</i>	(JLab CLAS Collab.)	REFID=53356
ADAMS	07B	PL B657 27	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=52758
DZIERBA	06	PR D73 072001	A.R. Dzierba <i>et al.</i>	(BNL E852 Collab.)	REFID=52048
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=51077
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
SALVINI	04	EPJ C35 21	P. Salvini <i>et al.</i>	(OBELIX Collab.)	REFID=49773
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>		REFID=53226
DZIERBA	03	PR D67 094015	A.R. Dzierba <i>et al.</i>		REFID=49414
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=49412
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ABELE	99	PL B446 349	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48317
ABELE	98B	PL B423 175	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46602
ADAMS	98B	PRL 81 5760	G.S. Adams <i>et al.</i>	(BNL E852 Collab.)	REFID=45864
THOMPSON	97	PRL 79 1630	D.R. Thompson <i>et al.</i>	(BNL E852 Collab.)	REFID=46610
PROKOSHKIN	95B	PAN 58 606	Y.D. Prokoshkin, S.A. Sadovsky	(SERP)	REFID=45584
		Translated from YAF 58 662.			REFID=44619
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)	REFID=44078
AOYAGI	93	PL B314 246	H. Aoyagi <i>et al.</i>	(BKEI Collab.)	REFID=43599
BOUTEMEUR	90	Hadron 89 Conf. p 119	M. Boutemur, M. Poulet	(SERP, BELG, LANL+)	REFID=41751
ALDE	88B	PL B205 397	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=40558
APEL	81	NP B193 269	W.D. Apel <i>et al.</i>	(SERP, CERN)	REFID=22913

 $a_1(1640)$

$$I^G(J^{PC}) = 1^-(1^{++})$$

Possibly seen in the study of the hadronic structure in decay $\tau \rightarrow 3\pi\nu_\tau$ (ABREU 98G and ASNER 00).

 $a_1(1640)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1655 ± 16 OUR AVERAGE		Error includes scale factor of 1.2.		
1700 ⁺³⁵ ₋₁₃₀	46M	¹ AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1691 ± 18 ± 30		DARGENT 17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
1630 ± 20	35k	² BAKER 03	SPEC	$\bar{p} p \rightarrow \omega \pi^+ \pi^- \pi^0$
1714 ± 9 ± 36		CHUNG 02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
1640 ± 12 ± 30		BAKER 99	SPEC	1.94 $\bar{p} p \rightarrow 4\pi^0$
1670 ± 90		BELLINI 85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$

¹Statistical error negligible.²Using the $a_1(1260)$ mass and width results of BOWLER 88. $a_1(1640)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
250 ± 40 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		
[254 ± 40 MeV OUR 2023 AVERAGE Scale factor = 1.8]				
510 ⁺¹⁷⁰ ₋₉₀	46M	¹ AGHASYAN 18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
171 ± 33 ± 40		DARGENT 17	RVUE	$D^0 \rightarrow \pi^- \pi^+ \pi^- \pi^+$
225 ± 30	35k	² BAKER 03	SPEC	$\bar{p} p \rightarrow \omega \pi^+ \pi^- \pi^0$
308 ± 37 ± 62		CHUNG 02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
300 ± 22 ± 40		BAKER 99	SPEC	1.94 $\bar{p} p \rightarrow 4\pi^0$
300 ± 100		BELLINI 85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M161

NODE=M161

NODE=M161M

NODE=M161M

NODE=M161M;LINKAGE=A

NODE=M161M;LINKAGE=KB


NODE=M161W

NODE=M161W

NEW

- ¹ Statistical error negligible.
- ² Using the $a_1(1260)$ mass and width results of BOWLER 88.

NODE=M161W;LINKAGE=A
 NODE=M161W;LINKAGE=KB



$a_1(1640)$ DECAY MODES

NODE=M161215;NODE=M161

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi\pi$	seen
Γ_2 $f_2(1270)\pi$	seen
Γ_3 $\sigma\pi$	seen
Γ_4 $\rho\pi S\text{-wave}$	seen
Γ_5 $\rho\pi D\text{-wave}$	seen
Γ_6 $\omega\pi\pi$	seen
Γ_7 $f_1(1285)\pi$	seen
Γ_8 $a_1(1260)\eta$	not seen

DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=8

$a_1(1640)$ BRANCHING RATIOS

NODE=M161220

$\Gamma(f_2(1270)\pi)/\Gamma(\sigma\pi)$	Γ_2/Γ_3		
VALUE	DOCUMENT ID	TECN	COMMENT
0.24±0.07	BAKER	99	SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$

NODE=M161R1
 NODE=M161R1

$\Gamma(\rho\pi D\text{-wave})/\Gamma_{\text{total}}$	Γ_5/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M161R2
 NODE=M161R2

- • • We do not use the following data for averages, fits, limits, etc. • • •
- seen CHUNG 02 B852 18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
- seen AMELIN 95B VES 36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$

$\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$	Γ_6/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT

NODE=M161R3
 NODE=M161R3

- • • We do not use the following data for averages, fits, limits, etc. • • •
- seen 35280 ¹ BAKER 03 SPEC $\bar{p}p \rightarrow \omega\pi^+\pi^-\pi^0$

¹ Assuming the $\omega\rho$ mechanism for the $\omega\pi\pi$ state.

NODE=M161R;LINKAGE=KB

$\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$	Γ_7/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M161R4
 NODE=M161R4

- • • We do not use the following data for averages, fits, limits, etc. • • •
- not seen KUHN 04 B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
- seen LEE 94 MPS2 18 $\pi^- p \rightarrow K^+ \bar{K}^0 \pi^- \pi^- p$

$\Gamma(a_1(1260)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	KUHN	04 B852	18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

NODE=M161R5
 NODE=M161R5

 $a_1(1640)$ REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
DARGENT	17	JHEP 1705 143	P. d'Argent <i>et al.</i>	(HEID, BRIS)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
BAKER	03	PL B563 140	C.A. Baker <i>et al.</i>	
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>	
ABREU	98G	PL B426 411	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
LEE	94	PL B323 227	J.H. Lee <i>et al.</i>	(BNL, IND, KYUN, MASD+)
BOWLER	88	PL B209 99	M.G. Bowler	(OXF)
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>	
Translated from YAF 41 1223.				

NODE=M161

REFID=59471
 REFID=58121
 REFID=49773
 REFID=49414
 REFID=48837
 REFID=47339
 REFID=46888
 REFID=45909
 REFID=44433
 REFID=44092
 REFID=40578
 REFID=47490

NODE=M117

 $f_2(1640)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

 $f_2(1640)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1639 ± 6 OUR AVERAGE	Error includes scale factor of 1.2.		
1620 ± 16	BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1647 ± 7	ADAMO	92 OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
1635 ± 7	ALDE	90 GAM2	$38 \pi^- p \rightarrow \omega \omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1640 ± 5	AMSLER	06 CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$
1659 ± 6	VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1643 ± 7	¹ ALDE	89B GAM2	$38 \pi^- p \rightarrow \omega \omega n$
¹ Superseded by ALDE 90.			

NODE=M117M

NODE=M117M

NODE=M117M;LINKAGE=BB

 $f_2(1640)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
100⁺⁶⁰₋₄₀ OUR AVERAGE		Error includes scale factor of 2.9.		[99 ⁺⁶⁰ ₋₄₀ MeV OUR 2023
AVERAGE Scale factor = 2.9]				
140 ⁺⁶⁰ ₋₂₀		BUGG	95 MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
58 ± 20		ADAMO	92 OBLX	$\bar{n} p \rightarrow 3\pi^+ 2\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
44 ± 9		AMSLER	06 CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$
152 ± 18		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
< 70	90	ALDE	90 GAM2	$38 \pi^- p \rightarrow \omega \omega n$

NODE=M117W

NODE=M117W

NEW

 $f_2(1640)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega \omega$	seen
Γ_2 4π	seen
Γ_3 $K \bar{K}$	seen

NODE=M117215;NODE=M117

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=3

 $f_2(1640)$ BRANCHING RATIOS $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AMSLER	06 CBAR	$0.9 \bar{p} p \rightarrow K^+ K^- \pi^0$

NODE=M117220

NODE=M117R2
 NODE=M117R2

 $f_2(1640)$ REFERENCES

AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
VLADIMIRSK...06		PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)
Translated from YAF 69 515				
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH) JP
ADAMO	92	PL B287 368	A. Adamo <i>et al.</i>	(OBELIX Collab.)
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)
ALDE	89B	PL B216 451	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+) IGJPC

NODE=M117

REFID=51136
 REFID=51191

REFID=44438
 REFID=42177
 REFID=40935
 REFID=40735

$\eta_2(1645)$

$$I^G(J^{PC}) = 0^+(2^-+)$$

NODE=M154

 $\eta_2(1645)$ MASS

NODE=M154M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1617± 5 OUR AVERAGE				
1613± 8	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1617± 8	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
1620±20	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1645±14±15	ADOMEIT	96	CBAR 0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1645± 6±20	ANISOVICH	00E	SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M154M

 $\eta_2(1645)$ WIDTH

NODE=M154W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
181±11 OUR AVERAGE				
185±17	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
177±18	BARBERIS	00C		450 $pp \rightarrow p_f 4\pi p_s$
180±25	BARBERIS	97B	OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
180 ⁺⁴⁰ ₋₂₁ ±25	ADOMEIT	96	CBAR 0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200±25	ANISOVICH	00E	SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M154W

 $\eta_2(1645)$ DECAY MODES

NODE=M154215;NODE=M154

Mode	Fraction (Γ_i/Γ)
Γ_1 $a_2(1320)\pi$	seen
Γ_2 $K\bar{K}\pi$	seen
Γ_3 $K^*\bar{K}$	seen
Γ_4 $\eta\pi^+\pi^-$	seen
Γ_5 $a_0(980)\pi$	seen
Γ_6 $f_2(1270)\eta$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←

 $\eta_2(1645)$ BRANCHING RATIOS

NODE=M154220

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.07±0.03	¹ BARBERIS	97C	OMEG 450 $pp \rightarrow ppK\bar{K}\pi$	

NODE=M154R1
NODE=M154R1

¹ Using $2(\pi^+\pi^-)$ data from BARBERIS 97B.

NODE=M154R1;LINKAGE=A

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_5
13.1±2.3 OUR AVERAGE				
13.5±4.6	² ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$	
13.0±2.7	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M154R3
NODE=M154R3

² Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M154R3;LINKAGE=AN

VALUE	DOCUMENT ID	COMMENT	Γ_6/Γ
not seen	BARBERIS	00B 450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$	

NODE=M154R4
NODE=M154R4

 $\eta_2(1645)$ REFERENCES

NODE=M154

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47958
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
BARBERIS	97C	PL B413 225	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45759
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202

$\omega(1650)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

See also the $\omega(1420)$ particle listing.

NODE=M126

NODE=M126

NODE=M126M

NODE=M126M

→ UNCHECKED ←

 $\omega(1650)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1670 ± 30 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1660.0 ± 8.4		¹ LICHARD	23 RVUE	$e^+e^- \rightarrow \omega\eta$
1698 ± 10	267	² ACHASOV	20B SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
1651 ± 3 $^{+16}_{-6}$	183k	³ ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+K^-\pi^0$
1673 $^{+6}_{-7}$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
1671 ± 6 ± 10	824	⁴ AKHMETSHIN	17A CMD3	$1.4-2.0 e^+e^- \rightarrow \omega\eta$
1660 ± 10	898	⁵ ACHASOV	16B SND	$1.34-2.00 e^+e^- \rightarrow \omega\eta$
1680 ± 10	13.1k	⁶ AULCHENKO	15A SND	$1.05-1.80 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1667 ± 13 ± 6		AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
1645 ± 8	13	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
1660 ± 10 ± 2		AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1770 ± 50 ± 60	1.2M	⁷ ACHASOV	03D RVUE	$0.44-2.00 e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1619 ± 5		⁸ HENNER	02 RVUE	$1.2-2.0 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1700 ± 20		EUGENIO	01 SPEC	$18 \pi^-p \rightarrow \omega\eta n$
1705 ± 26	612	⁹ AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1820 $^{+190}_{-150}$		¹⁰ ACHASOV	98H RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1840 $^{+100}_{-70}$		¹¹ ACHASOV	98H RVUE	$e^+e^- \rightarrow \omega\pi^+\pi^-$
1780 $^{+170}_{-300}$		¹² ACHASOV	98H RVUE	$e^+e^- \rightarrow K^+K^-$
~ 2100		¹³ ACHASOV	98H RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1606 ± 9		¹⁴ CLEGG	94 RVUE	
1662 ± 13	750	¹⁵ ANTONELLI	92 DM2	$1.34-2.4 e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
1670 ± 20		ATKINSON	83B OMEG	$20-70 \gamma p \rightarrow 3\pi X$
1657 ± 13		CORDIER	81 DM1	$e^+e^- \rightarrow \omega 2\pi$
1679 ± 34	21	ESPOSITO	80 FRAM	$e^+e^- \rightarrow 3\pi$
1652 ± 17		COSME	79 OSPK	$e^+e^- \rightarrow 3\pi$

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=4

NODE=M126M;LINKAGE=J

NODE=M126M;LINKAGE=H

NODE=M126M;LINKAGE=G

NODE=M126M;LINKAGE=F

NODE=M126M;LINKAGE=E

NODE=M126M;LINKAGE=A

NODE=M126M;LINKAGE=VH

NODE=M126M;LINKAGE=AB

NODE=M126M;LINKAGE=KI

NODE=M126M;LINKAGE=L1

NODE=M126M;LINKAGE=L2

NODE=M126M;LINKAGE=L3

NODE=M126M;LINKAGE=L4

NODE=M126M;LINKAGE=AD

¹ From a VDM fit to AKHMETSHIN 17A $\omega\eta$ data with two additional resonances of low statistical evidences.² From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass of $\omega(1420)$ is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of $\omega(1420)$ to the PDG 18 value of 220 MeV results in 1694 ± 9 MeV measurement.³ Could also be $\rho(1700)$. Branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (5.3 \pm 0.3^{+0.6}_{-0.5}) \times 10^{-5}$.⁴ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.⁵ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.⁶ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.⁷ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.⁸ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.⁹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.¹⁰ Using data from BARKOV 87, DOLINSKY 91, and ANTONELLI 92.¹¹ Using the data from ANTONELLI 92.¹² Using the data from IVANOV 81 and BISELLO 88B.¹³ Using the data from BISELLO 91C.¹⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹⁵ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

NODE=M126M;LINKAGE=AE

$\omega(1650)$ WIDTH

NODE=M126W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
315 ± 35 OUR ESTIMATE				
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
106 ± 15		¹ LICHARD	23 RVUE	$e^+e^- \rightarrow \omega\eta$
110 ± 16	267	² ACHASOV	20B SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
194 ± 8 ⁺ ₇ ¹⁵	183k	³ ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+K^-\pi^0$
95 ± 11		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
113 ± 9 ± 10	824	⁴ AKHMETSHIN	17A CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
110 ± 20	898	⁵ ACHASOV	16B SND	1.34–2.00 $e^+e^- \rightarrow \omega\eta$
310 ± 30	13.1k	⁶ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
222 ± 25 ± 20		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
114 ± 14	13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
230 ± 30 ± 20		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
490 ⁺ ₁₅₀ ²⁰⁰ ± 130	1.2M	⁷ ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
250 ± 14		⁸ HENNER	02 RVUE	1.2–2.0 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
250 ± 50		EUGENIO	01 SPEC	18 $\pi^-p \rightarrow \omega\eta n$
370 ± 25	612	⁹ AKHMETSHIN	00D CMD2	$e^+e^- \rightarrow \omega\pi^+\pi^-$
113 ± 20		¹⁰ CLEGG	94 RVUE	
280 ± 24	750	¹¹ ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$
160 ± 20		ATKINSON	83B OMEG	20–70 $\gamma p \rightarrow 3\pi X$
136 ± 46		CORDIER	81 DM1	$e^+e^- \rightarrow \omega 2\pi$
99 ± 49	21	ESPOSITO	80 FRAM	$e^+e^- \rightarrow 3\pi$
42 ± 17		COSME	79 OSPK	$e^+e^- \rightarrow 3\pi$

NODE=M126W

→ UNCHECKED ←

OCCUR=2

OCCUR=5

OCCUR=4

¹ From a VDM fit to AKHMETSHIN 17A $\omega\eta$ data with two additional resonances of low statistical evidences.

² From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass of $\omega(1420)$ is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of $\omega(1420)$ to the PDG 18 value of 220 MeV results in 94 ± 13 MeV measurement.

³ Could also be $\rho(1700)$. Branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+K^-\pi^0 = (5.3 \pm 0.3^{+0.6}_{-0.5}) \times 10^{-5}$.

⁴ From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating.

⁵ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

⁶ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.

⁷ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁸ Using results of CORDIER 81 and preliminary data of DOLINSKY 91 and ANTONELLI 92.

⁹ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

¹⁰ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

¹¹ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

NODE=M126W;LINKAGE=J

NODE=M126W;LINKAGE=H

NODE=M126W;LINKAGE=G

NODE=M126W;LINKAGE=F

NODE=M126W;LINKAGE=E

NODE=M126W;LINKAGE=A

NODE=M126W;LINKAGE=VH

NODE=M126W;LINKAGE=AB

NODE=M126W;LINKAGE=KI

NODE=M126W;LINKAGE=AD

NODE=M126W;LINKAGE=AE

NODE=M126215;NODE=M126

$\omega(1650)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\rho(1450)\pi$	seen
Γ_3 $\omega\pi\pi$	seen
Γ_4 $\omega\eta$	seen
Γ_5 e^+e^-	seen
Γ_6 $\pi^0\gamma$	not seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=6

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=5

$\omega(1650) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M126230

 $\Gamma(\rho\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma \times \Gamma_5/\Gamma$ NODE=M126G3
NODE=M126G3

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.56 ± 0.23	13.1k	¹ AULCHENKO	15A SND	1.05–1.80 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
1.3 ± 0.1 ± 0.1		AUBERT,B	04N BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
1.2 $^{+0.4}_{-0.1}$ ± 0.8	1.2M	^{2,3} ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.921 ± 0.230		^{4,5} CLEGG	94 RVUE	
0.479 ± 0.050	750	^{6,7} ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ From a fit with contributions from $\omega(782)$, $\phi(1020)$, $\omega(1420)$, and $\omega(1650)$. See ACHASOV 20A for a further analysis of the $\pi^+\pi^-\pi^0$ data.

² Calculated by us from the cross section at the peak.

³ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁵ From the partial and leptonic width given by the authors.

⁶ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

⁷ From the product of the leptonic width and partial branching ratio given by the authors.

NODE=M126G3;LINKAGE=A

NODE=M126G;LINKAGE=AW
NODE=M126G;LINKAGE=VH

NODE=M126G;LINKAGE=AD

NODE=M126G;LINKAGE=SE
NODE=M126G;LINKAGE=AE
NODE=M126G;LINKAGE=ES $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_5/\Gamma$ NODE=M126G4
NODE=M126G4

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
7.0 ± 0.5		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
4.1 ± 0.9 ± 1.3	1.2M	^{1,2} ACHASOV	03D RVUE	0.44–2.00 $e^+e^- \rightarrow \pi^+\pi^-\pi^0$
5.40 ± 0.95		³ AKHMETSHIN	00D CMD2	1.2–1.38 $e^+e^- \rightarrow \omega\pi^+\pi^-$
3.18 ± 0.80		^{4,5} CLEGG	94 RVUE	
6.07 ± 0.61	750	^{6,7} ANTONELLI	92 DM2	1.34–2.4 $e^+e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ Calculated by us from the cross section at the peak.

² From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+\pi^-\pi^0$ and ANTONELLI 92 on the $\omega\pi^+\pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

³ Using the data of AKHMETSHIN 00D and ANTONELLI 92. The $\rho\pi$ dominance for the energy dependence of the $\omega(1420)$ and $\omega(1650)$ width assumed.

⁴ From a fit to two Breit-Wigner functions and using the data of DOLINSKY 91 and ANTONELLI 92.

⁵ From the partial and leptonic width given by the authors.

⁶ From the combined fit of the $\rho\pi$ and $\omega\pi\pi$ final states.

⁷ From the product of the leptonic width and partial branching ratio given by the authors.

NODE=M126G4;LINKAGE=AW
NODE=M126G4;LINKAGE=VH

NODE=M126G4;LINKAGE=KL

NODE=M126G4;LINKAGE=AD

NODE=M126G4;LINKAGE=SE
NODE=M126G4;LINKAGE=AE
NODE=M126G4;LINKAGE=ES $\Gamma(\omega\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_5/\Gamma$ NODE=M126G5
NODE=M126G5

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
6.4 ± 0.9	267	¹ ACHASOV	20B SND	$e^+e^- \rightarrow \omega\eta \rightarrow \eta\pi^0\gamma$
5.62 $^{+0.45}_{-0.42}$		ACHASOV	19 SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
4.5 ± 0.3 ± 0.3	824	² AKHMETSHIN	17A CMD3	1.4–2.0 $e^+e^- \rightarrow \omega\eta$
4.4 ± 0.5	898	³ ACHASOV	16B SND	1.34–2.00 $e^+e^- \rightarrow \omega\eta$
5.7 ± 0.6	13	AUBERT	06D BABR	10.6 $e^+e^- \rightarrow \omega\eta\gamma$
< 60 at 90% CL		⁴ AKHMETSHIN	03B CMD2	$e^+e^- \rightarrow \eta\pi^0\gamma$

¹ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$. The mass of $\omega(1420)$ is fixed to the PDG 18 value of 1420 MeV. Fixing also the width of $\omega(1420)$ to the PDG 18 value of 220 MeV results in $(5.4 \pm 0.6) \times 10^{-7}$ measurement.

² From a fit of the interfering $\omega(1420)$ and $\omega(1650)$ with a relative phase of π and other parameters floating. From an alternative fit $\Gamma(\omega(1650) \rightarrow \omega\eta)/\Gamma_{\text{total}} \times \Gamma(\omega(1650) \rightarrow e^+e^-) = 51 \pm 3$ eV.

³ From a fit with contributions from $\omega(1420)$, $\omega(1650)$, and $\phi(1680)$.

⁴ $\omega(1650)$ mass and width fixed at 1700 MeV and 250 MeV, respectively.

NODE=M126G5;LINKAGE=C

NODE=M126G5;LINKAGE=B

NODE=M126G5;LINKAGE=A
NODE=M126G5;LINKAGE=KH

$\omega(1650)$ BRANCHING RATIOS $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.65	1.2M	¹ ACHASOV	03D	RVUE $0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.380 ± 0.014		² HENNER	02	RVUE $1.2-2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

² Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

NODE=M126225

NODE=M126R3
NODE=M126R3

NODE=M126R;LINKAGE=VH

NODE=M126R;LINKAGE=AC

 $\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	ACHASOV	20A	SND $1.15-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
-------------	---------	-----	---

NODE=M126R02
NODE=M126R02 $\Gamma(\omega\pi\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.35	1.2M	¹ ACHASOV	03D	RVUE $0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
0.620 ± 0.014		² HENNER	02	RVUE $1.2-2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ From the combined fit of ANTONELLI 92, ACHASOV 01E, ACHASOV 02E, and ACHASOV 03D data on the $\pi^+ \pi^- \pi^0$ and ANTONELLI 92 on the $\omega \pi^+ \pi^-$ final states. Supersedes ACHASOV 99E and ACHASOV 02E.

² Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

NODE=M126R2
NODE=M126R2

NODE=M126R2;LINKAGE=VH

NODE=M126R2;LINKAGE=AC

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 18	1.2M	^{1,2} ACHASOV	03D	RVUE $0.44-2.00 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$
32 ± 1		² HENNER	02	RVUE $1.2-2.0 e^+ e^- \rightarrow \rho\pi, \omega\pi\pi$

¹ Calculated by us from the cross section at the peak.

² Assuming that the $\omega(1650)$ decays into $\rho\pi$ and $\omega\pi\pi$ only.

NODE=M126R4
NODE=M126R4

NODE=M126R;LINKAGE=AW

NODE=M126R4;LINKAGE=AC

 $\Gamma(\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	¹ ACHASOV	10D	SND $1.075-2.0 e^+ e^- \rightarrow \pi^0 \gamma$
-----------------	----------------------	-----	--

¹ From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. The width of the highest mass effective resonance is fixed at 315 MeV.

NODE=M126R00
NODE=M126R00

NODE=M126R00;LINKAGE=A

 $\omega(1650)$ REFERENCES

NODE=M126

LICHARD	23	PR D108 092005	P. Lichard	(OPAV, CTUP)	REFID=62426
ACHASOV	20A	EPJ C80 993	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60923
ACHASOV	20B	EPJ C80 1008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60924
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59887
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AKHMETSHIN	17A	PL B773 150	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)	REFID=58239
ACHASOV	16B	PR D94 092002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57537
AULCHENKO	15A	JETP 121 27	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56843
		Translated from ZETF 148 34.			
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
ACHASOV	03D	PR D68 052006	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=49577
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ACHASOV	02E	PR D66 032001	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48815
HENNER	02	EPJ C26 3	V.K. Henner <i>et al.</i>		REFID=49177
ACHASOV	01E	PR D63 072002	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=48311
EUGENIO	01	PL B497 190	P. Eugenio <i>et al.</i>		REFID=48010
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
ACHASOV	99E	PL B462 365	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47391
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41867
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			
ATKINSON	83B	PL 127B 132	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21502
CORDIER	81	PL 106B 155	A. Cordier <i>et al.</i>	(ORSAY)	REFID=21586
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
ESPOSITO	80	LCN 28 195	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=21584
COSME	79	NP B152 215	G. Cosme <i>et al.</i>	(IPN)	REFID=21475

$\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

NODE=M045

 $\omega_3(1670)$ MASS

NODE=M045M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1667 ± 4 OUR AVERAGE				
1665.3 ± 5.2 ± 4.5	23400	AMELIN	96 VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1685 ± 20	60	BAUBILLIER	79 HBC	8.2 $K^- p$ backward
1673 ± 12	430	^{1,2} BALTAY	78E HBC	15 $\pi^+ p \rightarrow \Delta 3\pi$
1650 ± 12		CORDEN	78B OMEG	8-12 $\pi^- p \rightarrow N 3\pi$
1669 ± 11	600	² WAGNER	75 HBC	7 $\pi^+ p \rightarrow \Delta^{++} 3\pi$
1678 ± 14	500	DIAZ	74 DBC	6 $\pi^+ n \rightarrow p 3\pi^0$
1660 ± 13	200	DIAZ	74 DBC	6 $\pi^+ n \rightarrow p \omega \pi^0 \pi^0$
1679 ± 17	200	MATTHEWS	71D DBC	7.0 $\pi^+ n \rightarrow p 3\pi^0$
1670 ± 20		KENYON	69 DBC	8 $\pi^+ n \rightarrow p 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1700	110	¹ CERRADA	77B HBC	4.2 $K^- p \rightarrow \Lambda 3\pi$
1695 ± 20		BARNES	69B HBC	4.6 $K^- p \rightarrow \omega 2\pi X$
1636 ± 20		ARMENISE	68B DBC	5.1 $\pi^+ n \rightarrow p 3\pi^0$

NODE=M045M

OCCUR=2

¹ Phase rotation seen for $J^P = 3^- \rho\pi$ wave.² From a fit to $I(J^P) = 0(3^-) \rho\pi$ partial wave.NODE=M045M;LINKAGE=E
NODE=M045M;LINKAGE=P $\omega_3(1670)$ WIDTH

NODE=M045W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
168 ± 10 OUR AVERAGE				
149 ± 19 ± 7	23400	AMELIN	96 VES	36 $\pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
160 ± 80	60	³ BAUBILLIER	79 HBC	8.2 $K^- p$ backward
173 ± 16	430	^{4,5} BALTAY	78E HBC	15 $\pi^+ p \rightarrow \Delta 3\pi$
253 ± 39		CORDEN	78B OMEG	8-12 $\pi^- p \rightarrow N 3\pi$
173 ± 28	600	^{3,5} WAGNER	75 HBC	7 $\pi^+ p \rightarrow \Delta^{++} 3\pi$
167 ± 40	500	DIAZ	74 DBC	6 $\pi^+ n \rightarrow p 3\pi^0$
122 ± 39	200	DIAZ	74 DBC	6 $\pi^+ n \rightarrow p \omega \pi^0 \pi^0$
155 ± 40	200	³ MATTHEWS	71D DBC	7.0 $\pi^+ n \rightarrow p 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
90 ± 20		BARNES	69B HBC	4.6 $K^- p \rightarrow \omega 2\pi$
100 ± 40		KENYON	69 DBC	8 $\pi^+ n \rightarrow p 3\pi^0$
112 ± 60		ARMENISE	68B DBC	5.1 $\pi^+ n \rightarrow p 3\pi^0$

NODE=M045W

OCCUR=2

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.⁴ Phase rotation seen for $J^P = 3^- \rho\pi$ wave.⁵ From a fit to $I(J^P) = 0(3^-) \rho\pi$ partial wave.NODE=M045W;LINKAGE=S
NODE=M045W;LINKAGE=E
NODE=M045W;LINKAGE=P $\omega_3(1670)$ DECAY MODES

NODE=M045215;NODE=M045

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \rho\pi$	seen
$\Gamma_2 \quad \omega\pi\pi$	seen
$\Gamma_3 \quad b_1(1235)\pi$	possibly seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

 $\omega_3(1670)$ BRANCHING RATIOS

NODE=M045220

$\Gamma(\omega\pi\pi)/\Gamma(\rho\pi)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.71 ± 0.27	100	DIAZ	74 DBC	6 $\pi^+ n \rightarrow p 5\pi^0$	
$\Gamma(b_1(1235)\pi)/\Gamma(\rho\pi)$					Γ_3/Γ_1
VALUE		DOCUMENT ID	TECN	COMMENT	
possibly seen		DIAZ	74 DBC	6 $\pi^+ n \rightarrow p 5\pi^0$	

NODE=M045R3
NODE=M045R3NODE=M045R4
NODE=M045R4

$\Gamma(b_1(1235)\pi)/\Gamma(\omega\pi\pi)$ Γ_3/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

>0.75	68	BAUBILLIER 79	HBC	8.2 $K^- p$ backward
-------	----	---------------	-----	----------------------

NODE=M045R5
NODE=M045R5

 $\omega_3(1670)$ REFERENCES

AMELIN 96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BAUBILLIER 79	PL 89B 131	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BALTAY 78E	PRL 40 87	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
CORDEN 78B	NP B138 235	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
CERRADA 77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+) JP
WAGNER 75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP
DIAZ 74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)
MATTHEWS 71D	PR D3 2561	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC)
BARNES 69B	PRL 23 142	V.E. Barnes <i>et al.</i>	(BNL)
KENYON 69	PRL 23 146	I.R. Kenyon <i>et al.</i>	(BNL, UCND, ORNL)
ARMENISE 68B	PL 26B 336	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)

NODE=M045

REFID=44649
REFID=21522
REFID=21520
REFID=21269
REFID=20537
REFID=20843
REFID=21248
REFID=21515
REFID=21512
REFID=20800
REFID=20783

NODE=M034

 $\pi_2(1670)$

$$I^{G(J^{PC})} = 1^-(2^-+)$$

 $\pi_2(1670)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

1670.6^{+2.9}_{-1.2} OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

1642	$\begin{smallmatrix} +12 \\ -1 \end{smallmatrix}$	46M	1	AGHASYAN	18B	COMP	190	$\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$		
1749	± 10	± 100	145k	LU	05	B852	18	$\pi^- p \rightarrow \omega \pi^- \pi^0 p$		
1676	± 3	± 8		2	CHUNG	02	B852	18.3	$\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
1685	± 10	± 30		BARBERIS	01		450	$pp \rightarrow p_f 3\pi^0 p_s$		
1687	± 9	± 15		AMELIN	99	VES	37	$\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$		
1669	± 4			BARBERIS	98B		450	$pp \rightarrow p_f \rho \pi p_s$		
1670	± 4			BARBERIS	98B		450	$pp \rightarrow p_f f_2(1270) \pi p_s$		
1690	± 14			3	BERDNIKOV	94	VES	37	$\pi^- A \rightarrow K^+ K^- \pi^- A$	
1710	± 20	700		ANTIPOV	87	SIGM	-	50	$\pi^- \text{Cu} \rightarrow \mu^+ \mu^- \pi^- \text{Cu}$	
1676	± 6			3	EVANGELIS...	81	OMEG	-	12	$\pi^- p \rightarrow 3\pi p$
1657	± 14			3,4	DAUM	80D	SPEC	-	63-94	$\pi p \rightarrow 3\pi X$
1662	± 10	2000		3	BALTAY	77	HBC	+	15	$\pi^+ p \rightarrow p 3\pi$

NODE=M034M

NODE=M034M

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

1658	± 3	$\begin{smallmatrix} +24 \\ -8 \end{smallmatrix}$	420k	5	ALEKSEEV	10	COMP	190	$\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$	
1730	± 20			6	AMELIN	95B	VES	36	$\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
1742	± 31	± 49		ANTREASYAN	90	CBAL		$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$		
1624	± 21			2	BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1622	± 35			7	BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1693	± 28			8	BELLINI	85	SPEC	40	$\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$	
1710	± 20			9	DAUM	81B	SPEC	-	63,94	$\pi^- p$
1660	± 10			3	ASCOLI	73	HBC	-	5-25	$\pi^- p \rightarrow p \pi_2$

OCCUR=2

OCCUR=3

¹ Statistical error negligible.

² From $f_2(1270)\pi$ decay.

³ From a fit to $J^P = 2^- S$ -wave $f_2(1270)\pi$ partial wave.

⁴ Clear phase rotation seen in $2^- S$, $2^- P$, $2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.

⁵ Superseded by AGHASYAN 2018B.

⁶ J^{PC} ambiguous.

⁷ From $\rho\pi$ decay.

⁸ From $\sigma\pi$ decay.

NODE=M034M;LINKAGE=A

NODE=M034M;LINKAGE=F2

NODE=M034M;LINKAGE=P

NODE=M034M;LINKAGE=D

NODE=M034M;LINKAGE=B


NODE=M034M;LINKAGE=AX

NODE=M034M;LINKAGE=R2

NODE=M034M;LINKAGE=S2

⁹ From a two-resonance fit to four 2^-0^+ waves. This should not be averaged with all the single resonance fits.

NODE=M034M;LINKAGE=L



$\pi_2(1670)$ WIDTH

NODE=M034W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
258⁺⁸₋₉	OUR AVERAGE	Error includes scale factor of 1.2.			
311 ⁺¹² ₋₂₃	46M	10 AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
408 \pm 60 \pm 250	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
254 \pm 3 \pm 31		11 CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
265 \pm 30 \pm 40		BARBERIS	01		450 $pp \rightarrow p_f 3\pi^0 p_s$
168 \pm 43 \pm 53		AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
268 \pm 15		BARBERIS	98B		450 $pp \rightarrow p_f p \pi p_s$
256 \pm 15		BARBERIS	98B		450 $pp \rightarrow p_f f_2(1270) \pi p_s$
190 \pm 50		12 BERDNIKOV	94	VES	37 $\pi^- A \rightarrow K^+ K^- \pi^- A$
170 \pm 80	700	ANTIPOV	87	SIGM	- 50 $\pi^- Cu \rightarrow \mu^+ \mu^- \pi^- Cu$
260 \pm 20		12 EVANGELIS...	81	OMEG	- 12 $\pi^- p \rightarrow 3\pi p$
219 \pm 20		12,13 DAUM	80D	SPEC	- 63-94 $\pi p \rightarrow 3\pi X$
285 \pm 60	2000	12 BALTAY	77	HBC	+ 15 $\pi^+ p \rightarrow p 3\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
271 \pm 9 ⁺²² ₋₂₄	420k	14 ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
310 \pm 20		15 AMELIN	95B	VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
236 \pm 49 \pm 36		ANTREASYAN	90	CBAL	$e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0 \pi^0$
304 \pm 22		11 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
404 \pm 108		16 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
330 \pm 90		17 BELLINI	85	SPEC	40 $\pi^- A \rightarrow \pi^- \pi^+ \pi^- A$
312 \pm 50		18 DAUM	81B	SPEC	- 63,94 $\pi^- p$
270 \pm 60		12 ASCOLI	73	HBC	- 5-25 $\pi^- p \rightarrow p \pi_2$

NODE=M034W

OCCUR=2

OCCUR=2

OCCUR=3

- ¹⁰ Statistical error negligible.
¹¹ From $f_2(1270)\pi$ decay.
¹² From a fit to $J^P = 2^- f_2(1270)\pi$ partial wave.
¹³ Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.
¹⁴ Superseded by AGHASYAN 2018B.
¹⁵ J^{PC} ambiguous.
¹⁶ From $\rho\pi$ decay.
¹⁷ From $\sigma\pi$ decay.
¹⁸ From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.

NODE=M034W;LINKAGE=A
 NODE=M034W;LINKAGE=F2
 NODE=M034W;LINKAGE=P
 NODE=M034W;LINKAGE=D

 NODE=M034W;LINKAGE=B
 NODE=M034W;LINKAGE=AX
 NODE=M034W;LINKAGE=R2
 NODE=M034W;LINKAGE=S2
 NODE=M034W;LINKAGE=L

$\pi_2(1670)$ DECAY MODES

NODE=M034215;NODE=M034

Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1 3π	(95.8±1.4) %		DESIG=20
Γ_2 $\pi^+\pi^-\pi^0$			DESIG=22
Γ_3 $\pi^0\pi^0\pi^0$			DESIG=23
Γ_4 $f_2(1270)\pi$	(56.3±3.2) %		DESIG=8
Γ_5 $\rho\pi$	(31 ±4) %		DESIG=2
Γ_6 $\sigma\pi$	(10 ±4) %		DESIG=13
Γ_7 $\pi(\pi\pi)_S$ -wave	(8.7±3.4) %		DESIG=11
Γ_8 $\pi^\pm\pi^+\pi^-$	(53 ±4) %		DESIG=10
Γ_9 $K\bar{K}^*(892)+$ c.c.	(4.2±1.4) %		DESIG=5
Γ_{10} $\omega\rho$	(2.7±1.1) %		DESIG=14
Γ_{11} $\pi^\pm\gamma$	(7.0±1.2) × 10 ⁻⁴		DESIG=27
Γ_{12} $\gamma\gamma$	< 2.8 × 10 ⁻⁷	90%	DESIG=12
Γ_{13} $\eta\pi$	< 5 %		DESIG=3
Γ_{14} $\pi^\pm 2\pi^+ 2\pi^-$	< 5 %		DESIG=4
Γ_{15} $\rho(1450)\pi$	< 3.6 × 10 ⁻³	97.7%	DESIG=15
Γ_{16} $b_1(1235)\pi$	< 1.9 × 10 ⁻³	97.7%	DESIG=16
Γ_{17} $\eta 3\pi$			DESIG=24
Γ_{18} $f_1(1285)\pi$	possibly seen		DESIG=25
Γ_{19} $a_2(1320)\pi$	not seen		DESIG=26

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.9$ for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_5	-53		
x_7	-29	-59	
x_9	-8	-21	-9
	x_4	x_5	x_7

$\pi_2(1670)$ PARTIAL WIDTHS

NODE=M034217

$\Gamma(\pi^\pm\gamma)$						Γ_{11}
VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT		
181±11±27	19 ADOLPH	14	COMP	-	190 $\pi^- \text{Pb} \rightarrow \pi^+ \pi^- \pi^- \text{Pb}'$	NODE=M034W2 NODE=M034W2

- ¹⁹ Primakoff reaction. Assumes incoherent $f_2(1270)\pi$ contribution to 3π final state and uses $B(\pi_2(1670) \rightarrow f_2\pi) = 56\%$.

NODE=M034W2;LINKAGE=AD

$\Gamma(\gamma\gamma)$ Γ_{12}

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.072	90	20 ACCIARRI	97T	L3	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.19	90	20 ALBRECHT	97B	ARG	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.41 ±0.23±0.28		ANTREASYAN 90	CBAL	0	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0\pi^0$
0.8 ±0.3 ±0.12		21 BEHREND	90C	CELL	0
1.3 ±0.3 ±0.2		22 BEHREND	90C	CELL	0

NODE=M034W1
 NODE=M034W1

²⁰ Decaying into $f_2(1270)\pi$ and $\rho\pi$.

²¹ Constructive interference between $f_2(1270)\pi, \rho\pi$ and background.

²² Incoherent Ansatz.

OCCUR=2

NODE=M034W1;LINKAGE=QQ

NODE=M034W1;LINKAGE=C

NODE=M034W1;LINKAGE=G

 $\pi_2(1670) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M034230

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_{12}/\Gamma$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.1	95	23 SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$

NODE=M034G01
 NODE=M034G01

²³ From analysis of L3 data at 183–209 GeV.

NODE=M034G01;LINKAGE=SC

 $\pi_2(1670) \text{ BRANCHING RATIOS}$

NODE=M034220

 $\Gamma(3\pi)/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma = (\Gamma_4 + \Gamma_5 + \Gamma_7)/\Gamma$

VALUE	DOCUMENT ID
0.958 ± 0.014 OUR FIT	

NODE=M034R20
 NODE=M034R20

 $\Gamma(\pi^0\pi^0\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_3/Γ_2

VALUE	DOCUMENT ID	COMMENT
0.29 ± 0.03 ± 0.05	BARBERIS 01	450 $\rho\rho \rightarrow \rho_f 3\pi^0 \rho_s$

NODE=M034R21
 NODE=M034R21

 $\Gamma(\rho\pi)/0.565\Gamma(f_2(1270)\pi)$ $\Gamma_5/0.565\Gamma_4$ (With $f_2(1270) \rightarrow \pi^+\pi^-$.)

VALUE	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.09 OUR AVERAGE	Error includes scale factor of 1.9.		
0.76 ± 0.07 ± 0.10	CHUNG 02	B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$
1.01 ± 0.05	BARBERIS 98B		450 $\rho\rho \rightarrow \rho_f \pi^+\pi^-\pi^0 \rho_s$

NODE=M034R16
 NODE=M034R16
 NODE=M034R16

 $\Gamma(\sigma\pi)/\Gamma(f_2(1270)\pi)$ Γ_6/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.02 ± 0.07	CHUNG 02	B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M034R15
 NODE=M034R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.24 ± 0.10 ^{24,25} BAKER 99 SPEC 1.94 $\bar{p}p \rightarrow 4\pi^0$

 $\frac{1}{2}\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$ $\frac{1}{2}\Gamma_5/\Gamma_8 = \frac{1}{2}\Gamma_5/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.29 ± 0.04 OUR FIT				
0.29 ± 0.05	²⁶ DAUM	81B	SPEC	63.94 $\pi^- p$

NODE=M034R2
 NODE=M034R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.3 BARTSCH 68 HBC + 8 $\pi^+ p \rightarrow 3\pi p$

 $0.565\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$ $0.565\Gamma_4/\Gamma_8 = 0.565\Gamma_4/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$ (With $f_2(1270) \rightarrow \pi^+\pi^-$.)

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.604 ± 0.035 OUR FIT				
0.60 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.3.			

NODE=M034R3

NODE=M034R3

NODE=M034R3

0.61 ± 0.04 ²⁶ DAUM 81B SPEC 63.94 $\pi^- p$

0.76 $\begin{matrix} +0.24 \\ -0.34 \end{matrix}$ ARMENISE 69 DBC + 5.1 $\pi^+ d \rightarrow d3\pi$

0.35 ± 0.20 BALTAY 68 HBC + 7–8.5 $\pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.59 BARTSCH 68 HBC + 8 $\pi^+ p \rightarrow 3\pi p$

$0.624\Gamma(\pi(\pi\pi)_{S\text{-wave}})/\Gamma(\pi^\pm\pi^+\pi^-)$

$0.624\Gamma_7/\Gamma_8 = 0.624\Gamma_7/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$

(With $(\pi\pi)_{S\text{-wave}} \rightarrow \pi^+\pi^-$.)

VALUE	DOCUMENT ID	TECN	COMMENT
0.10±0.04 OUR FIT			
0.10±0.05	26 DAUM	81B SPEC	63,94 $\pi^- p$

NODE=M034R11

NODE=M034R11

NODE=M034R11

 $\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(f_2(1270)\pi)$ **Γ_9/Γ_4**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.075±0.025 OUR FIT				
0.075±0.025	27 ARMSTRONG	82B OMEG	-	16 $\pi^- p \rightarrow K^+ K^- \pi^- p$

NODE=M034R13

NODE=M034R13

 $\Gamma(\omega\rho)/\Gamma_{\text{total}}$ **Γ_{10}/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
0.027±0.004±0.010	28 AMELIN	99 VES	37 $\pi^- A \rightarrow \omega\pi^- \pi^0 A^*$

NODE=M034R17

NODE=M034R17

 $\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$

$\Gamma_{13}/\Gamma_8 = \Gamma_{13}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$

(All η decays.)

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.09	BALTAY	68 HBC	+	7-8.5 $\pi^+ p$
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.10	CRENNELL	70 HBC	-	6 $\pi^- p \rightarrow f_2\pi^- N$

NODE=M034R5

NODE=M034R5

NODE=M034R5

 $\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$

$\Gamma_{14}/\Gamma_8 = \Gamma_{14}/(0.565\Gamma_4 + \frac{1}{2}\Gamma_5 + 0.624\Gamma_7)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.10	CRENNELL	70 HBC	-	6 $\pi^- p \rightarrow f_2\pi^- N$
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.1	BALTAY	68 HBC	+	7,8.5 $\pi^+ p$

NODE=M034R6

NODE=M034R6

 $\Gamma(\rho(1450)\pi)/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0036	97.7	AMELIN	99 VES	37 $\pi^- A \rightarrow \omega\pi^- \pi^0 A^*$

NODE=M034R18

NODE=M034R18

 $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0019	97.7	AMELIN	99 VES	37 $\pi^- A \rightarrow \omega\pi^- \pi^0 A^*$

NODE=M034R19

NODE=M034R19

 $\Gamma(f_1(1285)\pi)/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
possibly seen	69k	KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M034R23

NODE=M034R23

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ **Γ_{19}/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
not seen	69k	KUHN	04 B852	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$

NODE=M034R24

NODE=M034R24

D-wave/S-wave RATIO FOR $\pi_2(1670) \rightarrow f_2(1270)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.18±0.06	24 BAKER	99 SPEC	1.94 $\bar{p}p \rightarrow 4\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.22±0.10	26 DAUM	81B SPEC	63,94 $\pi^- p$

NODE=M034R14

NODE=M034R14

F-wave/P-wave RATIO FOR $\pi_2(1670) \rightarrow \rho\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.72±0.07±0.14	CHUNG	02 B852	18.3 $\pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M034R22

NODE=M034R22

24 Using preliminary CBAR data.

25 With the $\sigma\pi$ in $L=2$ and the $f_2(1270)\pi$ in $L=0$.26 From a two-resonance fit to four 2^-0^+ waves.27 From a partial-wave analysis of $K^+K^-\pi^-$ system.28 Normalized to the $B(\pi_2(1670) \rightarrow f_2\pi)$.

NODE=M034R;LINKAGE=BK

NODE=M034R15;LINKAGE=A

NODE=M034R;LINKAGE=L

NODE=M034R13;LINKAGE=M

NODE=M034R;LINKAGE=DM

$\pi_2(1670)$ REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ADOLPH	14	EPJ A50 79	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=55911
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)	REFID=49773
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
BARBERIS	01	PL B507 14	D. Barberis <i>et al.</i>		REFID=48324
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
		Translated from YAF 62 487.			
BAKER	99	PL B449 114	C.A. Baker <i>et al.</i>		REFID=46888
BARBERIS	98B	PL B422 399	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=46345
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761
ALBRECHT	97B	ZPHY C74 469	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=45418
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44433
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)	REFID=44073
ANTREASYAN	90	ZPHY C48 561	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=41372
BEHREND	90C	ZPHY C46 583	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41356
ANTIPOV	87	EPL 4 403	Y.M. Antipov <i>et al.</i>	(SERP, JINR, INRM+)	REFID=40004
BELLINI	85	SJNP 41 781	D. Bellini <i>et al.</i>		REFID=47490
		Translated from YAF 41 1223.			
ARMSTRONG	82B	NP B202 1	T.A. Armstrong, B. Baccari	(AACH3, BARI, BONN+)	REFID=20874
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=20872
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
		Also NP B186 594	C. Evangelista		REFID=21576
DAUM	80D	PL B89B 285	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+) JP	REFID=21573
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP	REFID=20847
ASCOLI	73	PR D7 669	G. Ascoli	(ILL, TINTO, GENO, HAMB, MILA+) JP	REFID=21553
CRENNELL	70	PRL 24 781	D.J. Crennell <i>et al.</i>	(BNL)	REFID=20805
ARMENISE	69	LCN 2 501	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20689
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I	REFID=21531
BARTSCH	68	NP B7 345	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN) JP	REFID=21532

NODE=M067

 $\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

 $\phi(1680)$ MASS

NODE=M067205

 e^+e^- PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1680 ± 20				OUR ESTIMATE
• • •				We do not use the following data for averages, fits, limits, etc. • • •
1656.8 ± 4.9		¹ LICHARD 23	RVUE	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
1683 ± 7 ± 9		² ZHU 23	BELL	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
1678 $\pm \frac{5}{3}$ ± 7		³ ZHU 23A	RVUE	$e^+e^- \rightarrow \eta\phi$
1673 ± 5		⁴ ABLIKIM 22L	BES3	$2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$
1680 $\pm \frac{12}{13}$ ± 21	1.8k	⁵ ABLIKIM 20F	BES3	$\psi(2S) \rightarrow K^+K^-\eta$
1662 ± 20		⁶ ACHASOV 20C	SND	$1.3-2.0 e^+e^- \rightarrow K^+K^-\pi^0$
1641 $\pm \frac{24}{18}$		ACHASOV 19	SND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
1667 ± 5 ± 11	3k	⁷ IVANOV 19A	CMD3	$1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$
1700 ± 23	2k	⁸ ACHASOV 18A	SND	$1.3-2.0 e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$
1674 ± 12 ± 6	6.2k	⁹ LEES 14H	BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
1733 ± 10 ± 10		¹⁰ LEES 12F	BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
1689 ± 7 ± 10	4.8k	¹¹ SHEN 09	BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
1709 ± 20 ± 43		¹² AUBERT 08S	BABR	$10.6 e^+e^- \rightarrow \text{hadrons}$
1623 ± 20	948	¹³ AKHMETSHIN 03	CMD2	$1.05-1.38 e^+e^- \rightarrow K_L^0 K_S^0$
~ 1500		¹⁴ ACHASOV 98H	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0, \omega\pi^+\pi^-, K^+K^-$
~ 1900		¹⁵ ACHASOV 98H	RVUE	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1700 ± 20		¹⁶ CLEGG 94	RVUE	$e^+e^- \rightarrow K^+K^-, K_S^0 K \pi$
1657 ± 27	367	BISELLO 91C	DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655 ± 17		¹⁷ BISELLO 88B	DM2	$e^+e^- \rightarrow K^+K^-$
1680 ± 10		¹⁸ BUON 82	DM1	$e^+e^- \rightarrow \text{hadrons}$
1677 ± 12		¹⁹ MANE 82	DM1	$e^+e^- \rightarrow K_S^0 K \pi$

NODE=M067M1

NODE=M067M1

→ UNCHECKED ←

OCCUR=4

- ¹ From a VDM fit to ZHU 23 $\eta\phi\gamma$ data with two resonances, $\phi(1680)$, $\phi(2170)$, and a third resonance with mass 1850.7 ± 5.3 MeV and width 25 ± 35 MeV of 1.7σ statistical evidence. NODE=M067M1;LINKAGE=I
- ² From a fit using a vector meson dominance model with contributions from $\phi(1680)$, $\phi(2170)$ and non resonant contribution. NODE=M067M1;LINKAGE=H
- ³ From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. NODE=M067M1;LINKAGE=J
- ⁴ From a partial wave amplitude analysis at $\sqrt{s} = 2.125$ GeV which includes all the possible intermediate states that match J^{PC} conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only. NODE=M067M1;LINKAGE=G
- ⁵ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+K^-\eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$. NODE=M067M1;LINKAGE=E
- ⁶ From a fit using a vector meson dominance model with contribution from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, $\rho(1450)$. NODE=M067M1;LINKAGE=F
- ⁷ From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution. NODE=M067M1;LINKAGE=D
- ⁸ Assuming the $K\bar{K}^*(892) + c.c.$ dynamics. Systematic uncertainties not estimated. NODE=M067M1;LINKAGE=C
- ⁹ Using a vector meson dominance model with contribution from $\phi(1020)$, $\phi(1680)$, and higher mass excitations of $\rho(770)$ and $\omega(782)$. NODE=M067M1;LINKAGE=B
- ¹⁰ Using events with $\pi\pi$ invariant mass less than 0.85 GeV. NODE=M067M1;LINKAGE=A
- ¹¹ From a fit with two incoherent Breit-Wigners. NODE=M067M1;LINKAGE=SH
- ¹² From the simultaneous fit to the $K\bar{K}^*(892) + c.c.$ and $\phi\eta$ data from AUBERT 08s using the results of AUBERT 07AK. NODE=M067M1;LINKAGE=AU
- ¹³ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known. NODE=M067M;LINKAGE=HK
- ¹⁴ Using data from IVANOV 81, BARKOV 87, BISELLO 88B, DOLINSKY 91, and ANTONELLI 92. NODE=M067M1;LINKAGE=L1
- ¹⁵ Using the data from BISELLO 91C. NODE=M067M1;LINKAGE=L4
- ¹⁶ Using BISELLO 88B and MANE 82 data. NODE=M067M;LINKAGE=A
- ¹⁷ From global fit including ρ , ω , ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation. NODE=M067M;LINKAGE=E
- ¹⁸ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0K_L^0$, $K_S^0K_L^\pm\pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation. NODE=M067M;LINKAGE=C
- ¹⁹ Fit to one channel only, neglecting interference with ω , $\rho(1700)$. NODE=M067M;LINKAGE=D

 $p\bar{p}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1700 ± 8	¹ AMSLER	06	CBAR 0.9 $p\bar{p} \rightarrow K^+K^-\pi^0$
----------	---------------------	----	---

¹ Could also be $\rho(1700)$.

NODE=M067M3
NODE=M067M3

NODE=M067M3;LINKAGE=AM

 $\phi(1680)$ WIDTH

NODE=M067210

 e^+e^- PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

150 ± 50 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

• • • We do not use the following data for averages, fits, limits, etc. • • •

150.8 ± 7.0		¹ LICHARD	23	RVUE $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
149 ± 12 ± 13		² ZHU	23	BELL $e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
156 ± 5 ± 9		³ ZHU	23A	RVUE $e^+e^- \rightarrow \eta\phi$
172 ± 8		⁴ ABLIKIM	22L	BES3 $2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$
185 $^{+30}_{-26}$ $^{+25}_{-47}$ 1.8k		⁵ ABLIKIM	20F	BES3 $\psi(2S) \rightarrow K^+K^-\eta$
159 ± 32		⁶ ACHASOV	20C	SND $1.3-2.0 e^+e^- \rightarrow K^+K^-\pi^0$
103 $^{+26}_{-24}$		ACHASOV	19	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
176 ± 23 ± 38 3k		⁷ IVANOV	19A	CMD3 $1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$
300 ± 50 2k		⁸ ACHASOV	18A	SND $1.3-2.0 e^+e^- \rightarrow K_S^0K_L^0\pi^0$
165 ± 38 ± 70 6.2k		⁹ LEES	14H	BABR $e^+e^- \rightarrow K_S^0K_L^0\gamma$
300 ± 15 ± 37		¹⁰ LEES	12F	BABR $10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
211 ± 14 ± 19 4.8k		¹¹ SHEN	09	BELL $10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

NODE=M067W1
NODE=M067W1

→ UNCHECKED ←

322	± 77	± 160	12	AUBERT	08S	BABR	10.6	$e^+e^- \rightarrow$	hadrons
139	± 60	948	13	AKHMETSHIN	03	CMD2	1.05–1.38	$e^+e^- \rightarrow$	$K_L^0 K_S^0$
300	± 60		14	CLEGG	94	RVUE		$e^+e^- \rightarrow$	$K^+ K^-, K_S^0 K \pi$
146	± 55	367		BISELLO	91C	DM2		$e^+e^- \rightarrow$	$K_S^0 K^\pm \pi^\mp$
207	± 45		15	BISELLO	88B	DM2		$e^+e^- \rightarrow$	$K^+ K^-$
185	± 22		16	BUON	82	DM1		$e^+e^- \rightarrow$	hadrons
102	± 36		17	MANE	82	DM1		$e^+e^- \rightarrow$	$K_S^0 K \pi$

¹ From a VDM fit to ZHU 23 $\eta\phi\gamma$ data with two resonances, $\phi(1680)$, $\phi(2170)$, and a third resonance with mass 1850.7 ± 5.3 MeV and width 25 ± 35 MeV of 1.7 σ statistical evidence.

² From a fit using a vector meson dominance model with contributions from $\phi(1680)$, $\phi(2170)$ and non resonant contribution.

³ From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3.

⁴ From a partial wave amplitude analysis at $\sqrt{s} = 2.125$ GeV which includes all the possible intermediate states that match J^{PC} conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

⁵ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (12.0 \pm 1.3^{+6.5}_{-6.9}) \times 10^{-6}$.

⁶ From a fit using a vector meson dominance model with contribution from $\rho(770)$, $\omega(782)$, $\phi(1020)$, $\omega(1420)$, $\rho(1450)$.

⁷ From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.

⁸ Assuming the $K\bar{K}^*(892) + c.c.$ dynamics. Systematic uncertainties not estimated.

⁹ Using a vector meson dominance model with contribution from $\phi(1020)$, $\phi(1680)$, and higher mass excitations of $\rho(770)$ and $\omega(782)$.

¹⁰ Using events with $\pi\pi$ invariant mass less than 0.85 GeV.

¹¹ From a fit with two incoherent Breit-Wigners.

¹² From the simultaneous fit to the $K\bar{K}^*(892) + c.c.$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

¹³ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known.

¹⁴ Using BISELLO 88B and MANE 82 data.

¹⁵ From global fit including ρ , ω , ϕ and $\rho(1700)$

¹⁶ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

¹⁷ Fit to one channel only, neglecting interference with ω , $\rho(1700)$.

NODE=M067W1;LINKAGE=I

NODE=M067W1;LINKAGE=H

NODE=M067W1;LINKAGE=J

NODE=M067W1;LINKAGE=G

NODE=M067W1;LINKAGE=E

NODE=M067W1;LINKAGE=F

NODE=M067W1;LINKAGE=D

NODE=M067W1;LINKAGE=C

NODE=M067W1;LINKAGE=B

NODE=M067W1;LINKAGE=A

NODE=M067W1;LINKAGE=SH

NODE=M067W1;LINKAGE=AU

NODE=M067W;LINKAGE=HK

NODE=M067W;LINKAGE=A

NODE=M067W;LINKAGE=E

NODE=M067W;LINKAGE=C

NODE=M067W;LINKAGE=D

NODE=M067W3

NODE=M067W3

$\rho\bar{\rho}$ ANNIHILATION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

143 \pm 24	¹ AMSLER	06	CBAR	$0.9 \bar{\rho}\rho \rightarrow K^+ K^- \pi^0$
--------------	---------------------	----	------	--

¹ Could also be $\rho(1700)$.

NODE=M067W3;LINKAGE=AM

$\phi(1680)$ DECAY MODES

NODE=M067215;NODE=M067

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}^*(892) + c.c.$	seen
Γ_2 $K_S^0 K \pi$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $K_L^0 K_S^0$	
Γ_5 e^+e^-	seen
Γ_6 $\omega\pi\pi$	not seen
Γ_7 $\phi\pi\pi$	
Γ_8 $K^+ K^- \pi^+ \pi^-$	seen
Γ_9 $\eta\phi$	seen
Γ_{10} $K^+ K^- \eta$	
Γ_{11} $\eta\gamma$	seen
Γ_{12} $K^+ K^- \pi^0$	
Γ_{13} $f'_2(1525)\gamma$	not seen

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=11

DESIG=12;OUR EVAL;→ UNCHECKED ←

DESIG=10

DESIG=14

DESIG=13

DESIG=2

DESIG=15

$\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel (I) in e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(I)$ or the branching ratio $\Gamma(I)/\text{total}$.

NODE=M067220

NODE=M067220

 $\Gamma(K_L^0 K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4 \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.3±2.4±6.2	6.2k	¹ LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_L^0 \gamma$
--------------	------	-------------------	----------	---

¹ Using a vector meson dominance model with contribution from $\phi(1020)$, $\phi(1680)$, and higher mass excitations of $\rho(770)$ and $\omega(782)$.

NODE=M067A00
NODE=M067A00

NODE=M067A00;LINKAGE=A

 $\Gamma(\phi\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7 \Gamma_5 / \Gamma$

VALUE (10 ⁻² keV)	DOCUMENT ID	TECN	COMMENT
------------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2±0.2±0.3	LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
-------------	------	----------	--

NODE=M067G02
NODE=M067G02 $\Gamma(\eta\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_9 \Gamma_5 / \Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

122± 6±13		¹ ZHU	23 BELL	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
-----------	--	------------------	---------	--

65 ⁺ ₄ ±13 to		² ZHU	23A RVUE	$e^+e^- \rightarrow \eta\phi$
-------------------------------------	--	------------------	----------	-------------------------------

215 ⁺ ₅ ±11				
-----------------------------------	--	--	--	--

94±13±15	3k	³ IVANOV	19A CMD3	$1.59-2.007 e^+e^- \rightarrow K^+K^-\eta$
----------	----	---------------------	----------	--

¹ From a solution of the fit using a vector meson dominance model with contributions from $\phi(1680)$, $\phi(2170)$ and non resonant contribution. Other solutions with equal fit quality give (219 ± 15 ± 18) eV, (163 ± 11 ± 13) eV and (203 ± 12 ± 18) eV.

² From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. Four solutions are found, with equal fit quality: (79 ± 4 ± 16) eV, (127 ± 5 ± 12) eV, (65⁺₄ ± 13) eV, (215⁺₅ ± 11) eV.

³ From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.

NODE=M067R00
NODE=M067R00

NODE=M067R00;LINKAGE=B

NODE=M067R00;LINKAGE=C

NODE=M067R00;LINKAGE=A

 $\phi(1680) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$

This combination of a branching ratio into channel (i) and branching ratio into e^+e^- is directly measured and obtained from the cross section at the peak. We list only data that have not been used to determine the branching ratio into (i) or e^+e^- .

NODE=M067223

NODE=M067223

 $\Gamma(K_L^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.131±0.059	948	¹ AKHMETSHIN 03	CMD2	$1.05-1.38 e^+e^- \rightarrow K_L^0 K_S^0$
-------------	-----	----------------------------	------	--

¹ From the combined fit of AKHMETSHIN 03 and MANE 81 also including ρ , ω , and ϕ . Neither isospin nor flavor structure known. Recalculated by us.

NODE=M067G5
NODE=M067G5

NODE=M067G;LINKAGE=GK

 $\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.15±0.16±0.01		¹ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow K\bar{K}^*(892)\gamma + \text{c.c.}$
----------------	--	---------------------	----------	---

3.29±1.57	367	² BISELLO	91C DM2	$1.35-2.40 e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
-----------	-----	----------------------	---------	--

¹ From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

² Recalculated by us with the published value of $B(K\bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)$.

NODE=M067G6
NODE=M067G6

NODE=M067G6;LINKAGE=AU

NODE=M067G;LINKAGE=GL

 $\Gamma(\phi\pi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10 ⁻⁷)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.86±0.14±0.21	4.8k	¹ SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
----------------	------	-------------------	---------	--

¹ Multiplied by 3/2 to take into account the $\phi\pi^0\pi^0$ mode. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.

NODE=M067G01
NODE=M067G01

NODE=M067G01;LINKAGE=SH

$\Gamma(\eta\phi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_9/\Gamma \times \Gamma_5/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.64^{+1.74}_{-1.80}$		ACHASOV	19	SND $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta$
$5.3 \pm 0.6 \pm 0.9$	3k	¹ IVANOV	19A	CMD3 $e^+e^- \rightarrow K^+K^-\eta$
$4.3 \pm 1.0 \pm 0.9$		² AUBERT	08S	BABR $10.6 e^+e^- \rightarrow \phi\eta\gamma$

¹ From a fit with coherent interference of the $\phi(1680)$ with a non-resonant contribution.

² From the simultaneous fit to the $K\bar{K}^*(892) + \text{c.c.}$ and $\phi\eta$ data from AUBERT 08S using the results of AUBERT 07AK.

NODE=M067G7
NODE=M067G7

NODE=M067G7;LINKAGE=A
NODE=M067G7;LINKAGE=AU

 $\phi(1680)$ BRANCHING RATIOS $\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K\pi)$ Γ_1/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

dominant	MANE	82	DM1 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
----------	------	----	--

NODE=M067225

NODE=M067R3
NODE=M067R3

 $\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892) + \text{c.c.})$ Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.07 ± 0.01	BUON	82	DM1 e^+e^-
-----------------	------	----	--------------

NODE=M067R2
NODE=M067R2

 $\Gamma(\omega\pi\pi)/\Gamma(K\bar{K}^*(892) + \text{c.c.})$ Γ_6/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

< 0.10	BUON	82	DM1 e^+e^-
----------	------	----	--------------

NODE=M067R1
NODE=M067R1

 $\Gamma(\eta\phi)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	35	¹ ACHASOV	14	SND $1.15-2.00 e^+e^- \rightarrow \eta\gamma$
------	----	----------------------	----	---

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

NODE=M067R01
NODE=M067R01

NODE=M067R01;LINKAGE=A

 $\Gamma(\eta\phi)/\Gamma(K\bar{K}^*(892) + \text{c.c.})$ Γ_9/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

≈ 0.37	¹ AUBERT	08S	BABR $10.6 e^+e^- \rightarrow \text{hadrons}$
----------------	---------------------	-----	---

¹ From the fit including data from AUBERT 07AK.

NODE=M067R5
NODE=M067R5

NODE=M067R5;LINKAGE=AU

 $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	35	¹ ACHASOV	14	SND $1.15-2.00 e^+e^- \rightarrow \eta\gamma$
------	----	----------------------	----	---

¹ From a phenomenological model based on vector meson dominance with $\rho(1450)$ and $\phi(1680)$ masses and widths from the PDG 12.

NODE=M067R02
NODE=M067R02

NODE=M067R02;LINKAGE=A

 $\Gamma(f_2'(1525)\gamma)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	¹ ACHASOV	22	SND $1.17-2.00 e^+e^- \rightarrow \eta\eta\gamma$
----------	----------------------	----	---

¹ The 90% CL upper limit on the Born cross sections $\sigma(e^+e^- \rightarrow \phi(1680) \rightarrow f_2'(1525)\gamma \rightarrow \eta\eta\gamma)$ and $\sigma(e^+e^- \rightarrow \rho(1700) \rightarrow f_0(1500)\gamma \rightarrow \eta\eta\gamma)$ is 10.6 pb.

NODE=M067R03
NODE=M067R03

NODE=M067R03;LINKAGE=A

 $\phi(1680)$ REFERENCES

LICHARD	23	PR D108 092005	P. Lichard	(OPAV, CTUP)	REFID=62426
ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)	REFID=61911
ZHU	23A	CP C47 113003	W. Zhu, X. Wang	(RVUE)	REFID=62439
ABLIKIM	22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61649
ACHASOV	22	EPJ C82 168	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61655
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
ACHASOV	20C	EPJ C80 1139	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=60935
ACHASOV	19	PR D99 112004	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59887
IVANOV	19A	PL B798 134946	V.L. Ivanov <i>et al.</i>	(CMD-3 Collab.)	REFID=60133
ACHASOV	18A	PR D97 032011	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58883
ACHASOV	14	PR D90 032002	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55912
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
PDG	12	PR D86 010001	J. Beringer <i>et al.</i>	(PDG Collab.)	REFID=54066
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000

NODE=M067

AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
AKHMETSHIN	03	PL B551 27	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49172
Also		PAN 65 1222	E.V. Anashkin, V.M. Aulchenko, R.R. Akhmetshin		REFID=48827
		Translated from YAF 65 1255.			
ACHASOV	98H	PR D57 4334	N.N. Achasov, A.A. Kozhevnikov		REFID=46323
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
ANTONELLI	92	ZPHY C56 15	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=43168
BISELLO	91C	ZPHY C52 227	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41867
DOLINSKY	91	PRPL 202 99	S.I. Dolinsky <i>et al.</i>	(NOVO)	REFID=41369
BISELLO	88B	ZPHY C39 13	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40581
BARKOV	87	JETPL 46 164	L.M. Barkov <i>et al.</i>	(NOVO)	REFID=40280
		Translated from ZETFP 46 132.			
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)	REFID=21494
MANE	82	PL 112B 178	F. Mane <i>et al.</i>	(LALO)	REFID=21590
IVANOV	81	PL 107B 297	P.M. Ivanov <i>et al.</i>	(NOVO)	REFID=20553
MANE	81	PL 99B 261	F. Mane <i>et al.</i>	(ORSAY)	REFID=21588

NODE=M015

 $\rho_3(1690)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

 $\rho_3(1690)$ MASS

NODE=M015205

VALUE (MeV) _____ DOCUMENT ID _____
1688.8±2.1 OUR AVERAGE Includes data from the 5 datablocks that follow this one.

NODE=M015M

2 π MODE

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____
 The data in this block is included in the average printed for a previous datablock.

NODE=M015M1
NODE=M015M1**1686± 4 OUR AVERAGE**

1677±14		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow 2\pi p$
1679±11	476	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow \pi^+ \pi^- n$
1678±12	175	¹ ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
1690± 7	600	¹ ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
1693± 8		² GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1678±12		MATTHEWS	71C	DBC	0	7 $\pi^+ N$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
1734±10		³ CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow n2\pi$
1692±12		^{2,4} ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1737±23		ARMENISE	70	DBC	0	9 $\pi^+ N$
1650±35	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
1687±21		STUNTEBECK	70	HDBC	0	8 $\pi^- p, 5.4 \pi^+ d$
1683±13		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
1670±30		GOLDBERG	65	HBC	0	6 $\pi^+ d, 8 \pi^- p$

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.² Uses same data as HYAMS 75.³ From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K\bar{K}$ result.⁴ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.NODE=M015M1;LINKAGE=E
NODE=M015M1;LINKAGE=G
NODE=M015M1;LINKAGE=M

NODE=M015M1;LINKAGE=I

 $K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____
 The data in this block is included in the average printed for a previous datablock.

NODE=M015M2
NODE=M015M2**1696± 4 OUR AVERAGE**

1699± 5		ALPER	80	CNTR	0	62 $\pi^- p \rightarrow K^+ K^- n$
1698±12	6k	^{5,6} MARTIN	78D	SPEC		10 $\pi p \rightarrow K_S^0 K^- p$
1692± 6		BLUM	75	ASPK	0	18.4 $\pi^- p \rightarrow nK^+ K^-$
1690±16		ADERHOLZ	69	HBC	+	8 $\pi^+ p \rightarrow K\bar{K}\pi$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1694± 8		⁷ COSTA	80	OMEG		10 $\pi^- p \rightarrow K^+ K^- n$
---------	--	--------------------	----	------	--	------------------------------------

⁵ From a fit to $J^P = 3^-$ partial wave.⁶ Systematic error on mass scale subtracted.⁷ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.NODE=M015M2;LINKAGE=P
NODE=M015M2;LINKAGE=S
NODE=M015M2;LINKAGE=L

(4 π) \pm MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015M3
 NODE=M015M3

1686 \pm 5 OUR AVERAGE Error includes scale factor of 1.1.

1694 \pm 6		⁸ EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow \rho 4\pi$
1665 \pm 15	177	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow \rho 4\pi$
1670 \pm 10		THOMPSON	74	HBC	+	13 $\pi^+ p$
1687 \pm 20		CASON	73	HBC	-	8,18.5 $\pi^- p$
1685 \pm 14		⁹ CASON	73	HBC	-	8,18.5 $\pi^- p$
1680 \pm 40	144	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N 4\pi$
1689 \pm 20	102	⁹ BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N 2\rho$
1705 \pm 21		CASO	70	HBC	-	11.2 $\pi^- p \rightarrow n \rho 2\pi$

OCCUR=2

OCCUR=3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1718 \pm 10		¹⁰ EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow \rho 4\pi$
1673 \pm 9		¹¹ EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow \rho 4\pi$
1733 \pm 9	66	⁹ KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow \rho 4\pi$
1630 \pm 15		HOLMES	72	HBC	+	10-12 $K^+ p$
1720 \pm 15		BALTAY	68	HBC	+	7, 8.5 $\pi^+ p$

OCCUR=2

OCCUR=3

⁸ From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.⁹ From $\rho^\pm \rho^0$ mode.¹⁰ From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.¹¹ From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

NODE=M015M3;LINKAGE=A

NODE=M015M3;LINKAGE=F

NODE=M015M3;LINKAGE=B

NODE=M015M3;LINKAGE=C

 $\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015M5
 NODE=M015M5

1681 \pm 7 OUR AVERAGE

1670 \pm 25		¹² ALDE	95	GAM2		38 $\pi^- p \rightarrow \omega \pi^0 n$
1690 \pm 15		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow \omega \pi p$
1666 \pm 14		GESSAROLI	77	HBC		11 $\pi^- p \rightarrow \omega \pi p$
1686 \pm 9		THOMPSON	74	HBC	+	13 $\pi^+ p$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1654 \pm 24		BARNHAM	70	HBC	+	10 $K^+ p \rightarrow \omega \pi X$
---------------	--	---------	----	-----	---	-------------------------------------

¹² Supersedes ALDE 92C.

NODE=M015M5;LINKAGE=A

 $\eta\pi^+\pi^-$ MODE(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M015M6

NODE=M015M6

NODE=M015M6

1682 \pm 12 OUR AVERAGE

1685 \pm 10 \pm 20	AMELIN	00	VES		37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
1680 \pm 15	FUKUI	88	SPEC	0	8.95 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1700 \pm 47	¹³ ANDERSON	69	MMS	-	16 $\pi^- p$ backward
1632 \pm 15	^{13,14} FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow \rho MM$
1700 \pm 15	^{13,14} FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow \rho MM$
1748 \pm 15	^{13,14} FOCACCI	66	MMS	-	7-12 $\pi^- p \rightarrow \rho MM$

OCCUR=2

OCCUR=3

¹³ Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+ 2\pi^-$, with 0, 1, 2 $\pi^+ \pi^-$ pairs in ρ band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1976)¹⁴ Not seen by BOWEN 72.

NODE=M015M6;LINKAGE=R

NODE=M015M6;LINKAGE=N

 $\rho_3(1690)$ WIDTH


NODE=M015210

2 π , $K\bar{K}$, AND $K\bar{K}\pi$ MODES

VALUE (MeV)	DOCUMENT ID
-------------	-------------

161 \pm 10 OUR AVERAGE Includes data from the 5 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.

NODE=M015W
 NODE=M015W



2π MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W1
NODE=M015W1

186±14 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

220±29		DENNEY	83	LASS		10 $\pi^+ N$
246±37		EVANGELIS...	81	OMEG	-	12 $\pi^- p \rightarrow 2\pi p$
116±30	476	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow \pi^+ \pi^- n$
162±50	175	15 ANTIPOV	77	CIBS	0	25 $\pi^- p \rightarrow p3\pi$
167±40	600	ENGLER	74	DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
200±18		16 GRAYER	74	ASPK	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
156±36		MATTHEWS	71C	DBC	0	7 $\pi^+ N$
171±65		ARMENISE	70	DBC	0	9 $\pi^+ d$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
322±35		17 CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow n2\pi$
240±30		16,18 ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
180±30	122	BARTSCH	70B	HBC	+	8 $\pi^+ p \rightarrow N2\pi$
267 ⁺⁷² ₋₄₆		STUNTEBECK	70	HDBC	0	8 $\pi^- p$, 5.4 $\pi^+ d$
188±49		ARMENISE	68	DBC	0	5.1 $\pi^+ d$
180±40		GOLDBERG	65	HBC	0	6 $\pi^+ d$, 8 $\pi^- p$


¹⁵Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹⁶Uses same data as HYAMS 75 and BECKER 79.

¹⁷From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K\bar{K}$ result.

¹⁸From phase-shift analysis. Error takes account of spread of different phase-shift solutions.

NODE=M015W1;LINKAGE=T
NODE=M015W1;LINKAGE=G
NODE=M015W1;LINKAGE=M
NODE=M015W1;LINKAGE=I



K \bar{K} AND K \bar{K} π MODES

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W2
NODE=M015W2

204±18 OUR AVERAGE

199±40	6000	¹⁹ MARTIN	78D	SPEC	10 $\pi^- p \rightarrow K_S^0 K^- p$
205±20		BLUM	75	ASPK 0	18.4 $\pi^- p \rightarrow n K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
219±4		ALPER	80	CNTR 0	62 $\pi^- p \rightarrow K^+ K^- n$
186±11		²⁰ COSTA	80	OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
112±60		ADERHOLZ	69	HBC +	8 $\pi^+ p \rightarrow K \bar{K} \pi$

¹⁹ From a fit to $J^P = 3^-$ partial wave.

²⁰ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

NODE=M015W2;LINKAGE=P
NODE=M015W2;LINKAGE=L

(4 π) $^\pm$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M015W3
NODE=M015W3

129±10 OUR AVERAGE

123±13		²¹ EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow p 4\pi$
105±30	177	BALTAY	78B	HBC +	15 $\pi^+ p \rightarrow p 4\pi$
169 ⁺⁷⁰ ₋₄₈		CASON	73	HBC -	8,18.5 $\pi^- p$
135±30	144	BARTSCH	70B	HBC +	8 $\pi^+ p \rightarrow N 4\pi$
160±30	102	BARTSCH	70B	HBC +	8 $\pi^+ p \rightarrow N 2\rho$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
230±28		²² EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow p 4\pi$
184±33		²³ EVANGELIS...	81	OMEG -	12 $\pi^- p \rightarrow p 4\pi$
150	66	²⁴ KLIGER	74	HBC -	4.5 $\pi^- p \rightarrow p 4\pi$
106±25		THOMPSON	74	HBC +	13 $\pi^+ p$
125 ⁺⁸³ ₋₃₅		²⁴ CASON	73	HBC -	8,18.5 $\pi^- p$
130±30		HOLMES	72	HBC +	10-12 $K^+ p$
180±30	90	²⁴ BARTSCH	70B	HBC +	8 $\pi^+ p \rightarrow N a_2 \pi$
100±35		BALTAY	68	HBC +	7, 8.5 $\pi^+ p$

²¹ From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.

²² From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.

²³ From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.

²⁴ From $\rho^\pm \rho^0$ mode.

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

NODE=M015W3;LINKAGE=A
NODE=M015W3;LINKAGE=B
NODE=M015W3;LINKAGE=C
NODE=M015W3;LINKAGE=F

$\omega\pi$ MODE

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____
 The data in this block is included in the average printed for a previous datablock.

190±40 OUR AVERAGE

230±65 ²⁵ ALDE 95 GAM2 38 $\pi^- p \rightarrow \omega\pi^0 n$
 190±65 EVANGELIS... 81 OMEG - 12 $\pi^- p \rightarrow \omega\pi p$
 160±56 GESSAROLI 77 HBC 11 $\pi^- p \rightarrow \omega\pi p$
 ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●
 89±25 THOMPSON 74 HBC + 13 $\pi^+ p$
 130⁺⁷³₋₄₃ BARNHAM 70 HBC + 10 $K^+ p \rightarrow \omega\pi X$

²⁵ Supersedes ALDE 92C.

NODE=M015W5
 NODE=M015W5

NODE=M015W5;LINKAGE=A

 $\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____
 The data in this block is included in the average printed for a previous datablock.

130±40 OUR AVERAGE Error includes scale factor of 1.8. [126 ± 40 MeV OUR
 2023 AVERAGE Scale factor = 1.8]

220±30±50 AMELIN 00 VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
 106±27 FUKUI 88 SPEC 0 8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
 ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●
 195 ²⁶ ANDERSON 69 MMS - 16 $\pi^- p$ backward
 < 21 26,27 FOCACCI 66 MMS - 7-12 $\pi^- p \rightarrow pMM$
 < 30 26,27 FOCACCI 66 MMS - 7-12 $\pi^- p \rightarrow pMM$
 < 38 26,27 FOCACCI 66 MMS - 7-12 $\pi^- p \rightarrow pMM$

²⁶ Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+2\pi^-$, with 0, 1, 2 $\pi^+\pi^-$ pairs in ρ^0 band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1979)

²⁷ Not seen by BOWEN 72.

NODE=M015W6

NODE=M015W6

NODE=M015W6

NEW

OCCUR=2

OCCUR=3

NODE=M015W6;LINKAGE=R

NODE=M015W6;LINKAGE=N

 $\rho_3(1690)$ DECAY MODES

NODE=M015215;NODE=M015

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 4π	(71.1 ± 1.9) %	
Γ_2 $\pi^\pm\pi^+\pi^-\pi^0$	(67 ± 22) %	
Γ_3 $\omega\pi$	(16 ± 6) %	
Γ_4 $\pi\pi$	(23.6 ± 1.3) %	
Γ_5 $K\bar{K}\pi$	(3.8 ± 1.2) %	
Γ_6 $K\bar{K}$	(1.58± 0.26) %	1.2
Γ_7 $\eta\pi^+\pi^-$	seen	
Γ_8 $\rho(770)\eta$	seen	
Γ_9 $\pi\pi\rho$	seen	
Γ_{10} $a_2(1320)\pi$	seen	
Γ_{11} $\rho\rho$	seen	
Γ_{12} $\phi\pi$		
Γ_{13} $\eta\pi$		
Γ_{14} $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

DESIG=2

DESIG=11

DESIG=7

DESIG=1

DESIG=3

DESIG=4

DESIG=13

DESIG=14;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=10

DESIG=12

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 14.7$ for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-77		
x_5	-74	17	
x_6	-15	2	0
	x_1	x_4	x_5

$\rho_3(1690)$ BRANCHING RATIOS

NODE=M015220

$\Gamma(\pi\pi)/\Gamma_{total}$ Γ_4/Γ

VALUE DOCUMENT ID TECN CHG COMMENT

0.236±0.013 OUR FIT

0.243±0.013 OUR AVERAGE

0.259 ^{+0.018} _{-0.019}	BECKER	79	ASPK	0	17 $\pi^- p$ polarized
0.23 ±0.02	CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow n2\pi$
0.22 ±0.04	²⁸ MATTHEWS	71C	HDBC	0	7 $\pi^+ n \rightarrow \pi^- p$
0.245±0.006	²⁹ ESTABROOKS	75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$

²⁸ One-pion-exchange model used in this estimation.

²⁹ From phase-shift analysis of HYAMS 75 data.

NODE=M015R1
NODE=M015R1

NODE=M015R1;LINKAGE=P
NODE=M015R1;LINKAGE=G

$\Gamma(\pi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$ Γ_4/Γ_2

VALUE DOCUMENT ID TECN CHG COMMENT

0.35±0.11 CASON 73 HBC - 8,18.5 $\pi^- p$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.2	HOLMES	72	HBC	+	10-12 $K^+ p$
<0.12	BALLAM	71B	HBC	-	16 $\pi^- p$

NODE=M015R2
NODE=M015R2

$\Gamma(\pi\pi)/\Gamma(4\pi)$ Γ_4/Γ_1

VALUE DOCUMENT ID TECN CHG COMMENT

0.332±0.026 OUR FIT Error includes scale factor of 1.1.

0.30 ±0.10 BALTAY 78B HBC 0 15 $\pi^+ p \rightarrow p4\pi$

NODE=M015R3
NODE=M015R3

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_6/Γ_4


VALUE DOCUMENT ID TECN CHG COMMENT

0.067±0.011 OUR FIT Error includes scale factor of 1.2.

0.118^{+0.040}_{-0.032} OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.

0.191 ^{+0.040} _{-0.037}	GORLICH	80	ASPK	0	17,18 $\pi^- p$ polarized
0.08 ±0.03	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.08 ^{+0.08} _{-0.03}	CRENNELL	68B	HBC		6.0 $\pi^- p$

NODE=M015R4
NODE=M015R4



$\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$ Γ_5/Γ_4

VALUE DOCUMENT ID TECN CHG COMMENT

0.16±0.05 OUR FIT

0.16±0.05 ³⁰BARTSCH 70B HBC + 8 $\pi^+ p$

³⁰ Increased by us to correspond to $B(\rho_3(1690) \rightarrow \pi\pi)=0.24$.

NODE=M015R5
NODE=M015R5

NODE=M015R5;LINKAGE=A

$$\frac{\Gamma(\pi^+\pi^0) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{(\Gamma_9 + \Gamma_{10} + \Gamma_{11})}{\Gamma_2}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.94 ± 0.09 OUR AVERAGE				
0.96 ± 0.21	BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
0.88 ± 0.15	BALLAM	71B	HBC	- 16 $\pi^- p$
1 ± 0.15	BARTSCH	70B	HBC	+ 8 $\pi^+ p$
consistent with 1	CASO	68	HBC	- 11 $\pi^- p$

NODE=M015R6
NODE=M015R6

$$\frac{\Gamma(\rho\rho)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{\Gamma_{11}}{\Gamma_2}$$

VALUE	EVT5	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.12 ± 0.11		BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
0.56	66	KLIGER	74	HBC	- 4.5 $\pi^- p \rightarrow p4\pi$
0.13 ± 0.09	31	THOMPSON	74	HBC	+ 13 $\pi^+ p$
0.7 ± 0.15		BARTSCH	70B	HBC	+ 8 $\pi^+ p$

NODE=M015R7
NODE=M015R7

³¹ $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

NODE=M015R7;LINKAGE=T

$$\frac{\Gamma(\rho\rho)}{[\Gamma(\pi^+\pi^0) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]} \quad \frac{\Gamma_{11}}{(\Gamma_9 + \Gamma_{10} + \Gamma_{11})}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.48 ± 0.16	CASO	68	HBC	- 11 $\pi^- p$

NODE=M015R8
NODE=M015R8

$$\frac{\Gamma(a_2(1320)\pi)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{\Gamma_{10}}{\Gamma_2}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.66 ± 0.08	BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
0.36 ± 0.14	32 THOMPSON	74	HBC	+ 13 $\pi^+ p$
not seen	CASON	73	HBC	- 8,18.5 $\pi^- p$
0.6 ± 0.15	BARTSCH	70B	HBC	+ 8 $\pi^+ p$
0.6	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R9
NODE=M015R9

³² $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

NODE=M015R9;LINKAGE=T

$$\frac{\Gamma(\omega\pi)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{\Gamma_3}{\Gamma_2}$$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
0.23 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.2.			
0.33 ± 0.07		THOMPSON	74	HBC	+ 13 $\pi^+ p$
0.12 ± 0.07		BALLAM	71B	HBC	- 16 $\pi^- p$
0.25 ± 0.10		BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$
0.25 ± 0.10		JOHNSTON	68	HBC	- 7.0 $\pi^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 0.11	95	BALTAY	78B	HBC	+ 15 $\pi^+ p \rightarrow p4\pi$
< 0.09		KLIGER	74	HBC	- 4.5 $\pi^- p \rightarrow p4\pi$

NODE=M015R10
NODE=M015R10

$$\frac{\Gamma(\phi\pi)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{\Gamma_{12}}{\Gamma_2}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 0.11	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R11
NODE=M015R11

$$\frac{\Gamma(\pi^+2\pi^+2\pi^-\pi^0)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{\Gamma_{14}}{\Gamma_2}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 0.15	BALTAY	68	HBC	+ 7,8.5 $\pi^+ p$

NODE=M015R12
NODE=M015R12

$$\frac{\Gamma(\eta\pi)}{\Gamma(\pi^+\pi^+\pi^-\pi^0)} \quad \frac{\Gamma_{13}}{\Gamma_2}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 0.02	THOMPSON	74	HBC	+ 13 $\pi^+ p$

NODE=M015R13
NODE=M015R13

$$\frac{\Gamma(K\bar{K})}{\Gamma_{\text{total}}} \quad \frac{\Gamma_6}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.0158 ± 0.0026 OUR FIT	Error includes scale factor of 1.2.			
0.0130 ± 0.0024 OUR AVERAGE				

NODE=M015R14
NODE=M015R14

0.013 ± 0.003	COSTA	80	OMEG 0	10 $\pi^- p \rightarrow K^+ K^- n$
0.013 ± 0.004	33 MARTIN	78B	SPEC	- 10 $\pi p \rightarrow K_S^0 K^- p$

³³ From $(\Gamma_4\Gamma_6)^{1/2} = 0.056 \pm 0.034$ assuming $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.

NODE=M015R14;LINKAGE=B

$$\frac{\Gamma(\omega\pi)}{[\Gamma(\omega\pi) + \Gamma(\rho\rho)]} \qquad \frac{\Gamma_3}{(\Gamma_3 + \Gamma_{11})}$$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22±0.08	CASON	73	HBC	-	8,18.5 $\pi^- p$
-----------	-------	----	-----	---	------------------

NODE=M015R16
NODE=M015R16

$$\frac{\Gamma(\eta\pi^+\pi^-)}{\Gamma_{\text{total}}} \qquad \frac{\Gamma_7}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	FUKUI	88	SPEC	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
------	-------	----	------	---

NODE=M015R17
NODE=M015R17

$$\frac{\Gamma(a_2(1320)\pi)}{\Gamma(\rho(770)\eta)} \qquad \frac{\Gamma_{10}}{\Gamma_8}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

5.5±2.0	AMELIN	00	VES	37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$
---------	--------	----	-----	---

NODE=M015R18
NODE=M015R18

$\rho_3(1690)$ REFERENCES

NODE=M015

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP	REFID=44371
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41859
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
EVANGELIS...	81	NP B178 197	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=20462
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21665
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
GORLICH	80	NP B174 16	L. Gorlich <i>et al.</i>	(CRAC, MPIM, CERN+)	REFID=20738
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)	REFID=21084
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20374
BALTAY	78B	PR D17 62	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21265
MARTIN	78B	NP B140 158	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=21273
MARTIN	78D	PL 74B 417	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=21272
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
GESSAROLI	77	NP B126 382	R. Gessaroli <i>et al.</i>	(BGNA, FIRZ, GENO+)	REFID=20230
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP	REFID=21651
ESTABROOKS	75	NP B95 322	P.G. Estabrooks, A.D. Martin	(DURH)	REFID=20642
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355
ENGLER	74	PR D10 2070	A. Engler <i>et al.</i>	(CMU, CASE)	REFID=20110
GRAYR	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)	REFID=20113
KLIGER	74	SJNP 19 428	G.K. Kliger <i>et al.</i>	(ITEP)	REFID=21648
		Translated from YAF 19 839.			
OREN	74	NP B71 189	Y. Oren <i>et al.</i>	(ANL, OXF)	REFID=20221
THOMPSON	74	NP B69 220	G. Thompson <i>et al.</i>	(PURD)	REFID=21650
CASON	73	PR D7 1971	N.M. Cason <i>et al.</i>	(NDAM)	REFID=20606
BOWEN	72	PRL 29 890	D.R. Bowen <i>et al.</i>	(NEAS, STON)	REFID=21711
HOLMES	72	PR D6 3336	R. Holmes <i>et al.</i>	(ROCH)	REFID=21639
BALLAM	71B	PR D3 2606	J. Ballam <i>et al.</i>	(SLAC)	REFID=21630
MATTHEWS	71C	NP B33 1	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC) JP	REFID=21633
ARMENISE	70	LCN 4 199	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ)	REFID=20693
BARNHAM	70	PRL 24 1083	K.W.J. Barnham <i>et al.</i>	(BIRM)	REFID=21624
BARTSCH	70B	NP B22 109	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN)	REFID=21625
CASO	70	LCN 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SAFL)	REFID=20590
STUNTEBECK	70	PL 32B 391	P.H. Stuntebeck <i>et al.</i>	(NDAM)	REFID=20696
ADERHOLZ	69	NP B11 259	M. Aderholz <i>et al.</i>	(AACH3, BERL, CERN+)	REFID=20687
ANDERSON	69	PRL 22 1390	E.W. Anderson <i>et al.</i>	(BNL, CMU)	REFID=20795
ARMENISE	68	NC 54A 999	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+) I	REFID=20054
BALTAY	68	PRL 20 887	C. Baltay <i>et al.</i>	(COLU, ROCH, RUTG, YALE) I	REFID=21531
CASO	68	NC 54A 983	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SAFL)	REFID=20586
CRENNELL	68B	PL 28B 136	D.J. Crennell <i>et al.</i>	(BNL)	REFID=21616
JOHNSTON	68	PRL 20 1414	T.F. Johnston <i>et al.</i>	(TNTO, WISC) IJP	REFID=21617
FOCACCI	66	PRL 17 890	M.N. Focacci <i>et al.</i>	(CERN)	REFID=20402
GOLDBERG	65	PL 17 354	M. Goldberg <i>et al.</i>	(CERN, EPOL, ORSAY+)	REFID=21601

$\rho(1700)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

NODE=M065

 $\rho(1700)$ MASS

NODE=M065205

 $\eta\rho^0$ AND $\pi^+\pi^-$ MODES

VALUE (MeV)

DOCUMENT ID

1720±20 OUR ESTIMATE

NODE=M065M0

NODE=M065M0

→ UNCHECKED ←

 $\eta\rho^0$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M065M6

NODE=M065M6

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1834±12	13.4k	¹ GRIBANOV	20	CMD3	1.1–2.0	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1840±10	7.4k	² ACHASOV	18	SND	1.22–2.00	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1740±20		ANTONELLI	88	DM2		$e^+e^- \rightarrow \eta\pi^+\pi^-$
1701±15		³ FUKUI	88	SPEC	8.95	$\pi^-p \rightarrow \eta\pi^+\pi^-n$

¹ Mass and width of the $\rho(770)$ fixed at 775 and 149 MeV, respectively; solution 2 of model 2, $\eta \rightarrow \gamma\gamma$ decays used.

NODE=M065M6;LINKAGE=B

² From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

NODE=M065M6;LINKAGE=A

³ Assuming $\rho^+f_0(1370)$ decay mode interferes with $a_1(1260)^+\pi$ background. From a two Breit-Wigner fit.

NODE=M065M;LINKAGE=B

 $\pi\pi$ MODE

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

NODE=M065M1

NODE=M065M1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1770.54±5.49		¹ BARTOS	17	RVUE		$e^+e^- \rightarrow \pi^+\pi^-$
1718.50±65.44		² BARTOS	17A	RVUE		$e^+e^- \rightarrow \pi^+\pi^-$
1766.80±52.36		³ BARTOS	17A	RVUE		$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1644±36	20k	⁴ LEES	17C	BABR		$J/\psi \rightarrow \pi^+\pi^-\pi^0$
1780±20	⁺¹⁵ ₋₂₀	63.5k	⁵ ABRAMOWICZ12	ZEUS		$ep \rightarrow e\pi^+\pi^-p$
1861±17		⁶ LEES	12G	BABR		$e^+e^- \rightarrow \pi^+\pi^-\gamma$
1728±17	±89	5.4M	^{7,8} FUJIKAWA	08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
1780	⁺³⁷ ₋₂₉		⁹ ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
1719±15		⁹ BERTIN	97C	OBLX	0.0	$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
1730±30		CLEGG	94	RVUE		$e^+e^- \rightarrow \pi^+\pi^-$
1768±21		BISELLO	89	DM2		$e^+e^- \rightarrow \pi^+\pi^-$
1745.7±91.9		DUBNICKA	89	RVUE		$e^+e^- \rightarrow \pi^+\pi^-$
1546±26		GESHKEN...	89	RVUE		
1650		¹⁰ ERKAL	85	RVUE	20–70	$\gamma p \rightarrow \gamma\pi$
1550±70		ABE	84B	HYBR	20	$\gamma p \rightarrow \pi^+\pi^-p$
1590±20		¹¹ ASTON	80	OMEG	20–70	$\gamma p \rightarrow p2\pi$
1600±10		¹² ATIYA	79B	SPEC	50	$\gamma C \rightarrow C2\pi$
1598	⁺²⁴ ₋₂₂	BECKER	79	ASPK	17	π^-p polarized
1659±25		¹⁰ LANG	79	RVUE		
1575		¹⁰ MARTIN	78C	RVUE	17	$\pi^-p \rightarrow \pi^+\pi^-n$
1610±30		¹⁰ FROGGATT	77	RVUE	17	$\pi^-p \rightarrow \pi^+\pi^-n$
1590±20		¹³ HYAMS	73	ASPK	17	$\pi^-p \rightarrow \pi^+\pi^-n$

¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.

NODE=M065M1;LINKAGE=C

² Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.

NODE=M065M1;LINKAGE=D

³ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUBNICKA 10 to analyze the data of FUJIKAWA 08.

NODE=M065M1;LINKAGE=E

⁴ From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.

NODE=M065M1;LINKAGE=A

⁵ Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.

NODE=M065M1;LINKAGE=AB

⁶ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

NODE=M065M1;LINKAGE=LE

⁷ $|F_\pi(0)|^2$ fixed to 1.

⁸ From the GOUNARIS 68 parametrization of the pion form factor.

⁹ T-matrix pole.

¹⁰ From phase shift analysis of HYAMS 73 data.

¹¹ Simple relativistic Breit-Wigner fit with constant width.

¹² An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

¹³ Included in BECKER 79 analysis.

NODE=M065M1;LINKAGE=FU
 NODE=M065M1;LINKAGE=GO
 NODE=M065M;LINKAGE=QQ
 NODE=M065M;LINKAGE=P
 NODE=M065M;LINKAGE=M
 NODE=M065M;LINKAGE=RR

NODE=M065M;LINKAGE=H

NODE=M065M8
 NODE=M065M8

$\pi\omega$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1723 ± 2		¹ ACHASOV 23A	SND	$e^+e^- \rightarrow \omega\pi^0$
1708 ± 41	7815	² ACHASOV 13	SND	1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1550 to 1620		³ ACHASOV 00i	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1580 to 1710		⁴ ACHASOV 00i	SND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
1710 ± 90		ACHASOV 97	RVUE	$e^+e^- \rightarrow \omega\pi^0$

OCCUR=2

NODE=M065M8;LINKAGE=A

NODE=M065M8;LINKAGE=AC

NODE=M065M;LINKAGE=I1

NODE=M065M;LINKAGE=I2

¹ From a vector dominance fit to the Born cross section between 1.05 and 2.0 GeV with $\rho(770)$, $\rho(1570)$, $\rho(1700)$, $\rho(2150)$. The fit also uses SND data from the VEPP-2M collider below 1.02 GeV and from LEES 17H and ABLIKIM 21A above 1.5 GeV.

² From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

³ Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00i on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.

⁴ Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00i on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$.

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1688.7 ± 3.1 ^{+141.1} _{1.3}		¹ ALBRECHT 20	RVUE		0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$
1541 ± 12 ± 33	190k	² AAIJ 16N	LHCB		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
1740.8 ± 22.2	27k	³ ABELE 99D	CBAR ±		0.0 $\bar{p}p \rightarrow K^+K^-\pi^0$
1582 ± 36	1600	CLELAND 82B	SPEC ±		50 $\pi p \rightarrow K_S^0 K^\pm p$

¹ T-matrix pole, 2 poles, 3 channels, including $\pi\pi$ scattering data from HYAMS 75.

² Using the GOUNARIS 68 parameterization with a fixed width. Value is average using different $K\pi$ S-wave parameterizations in fit.

³ K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

NODE=M065M2;LINKAGE=D

NODE=M065M2;LINKAGE=A

NODE=M065M2;LINKAGE=AN

2 ($\pi^+\pi^-$) MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1851 ⁺²⁷ ₂₄		ACHASOV 97	RVUE	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1570 ± 20		¹ CORDIER 82	DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1520 ± 30		² ASTON 81E	OMEG	20–70 $\gamma p \rightarrow p4\pi$
1654 ± 25		³ DIBIANCA 81	DBC	$\pi^+d \rightarrow pp2(\pi^+\pi^-)$
1666 ± 39		¹ BACCI 80	FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1780	34	KILLIAN 80	SPEC	11 $e^-p \rightarrow 2(\pi^+\pi^-)$
1500		⁴ ATIYA 79B	SPEC	50 $\gamma C \rightarrow C4\pi^\pm$
1570 ± 60	65	⁵ ALEXANDER 75	HBC	7.5 $\gamma p \rightarrow p4\pi$
1550 ± 60		² CONVERSI 74	OSPK	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1550 ± 50	160	SCHACHT 74	STRC	5.5–9 $\gamma p \rightarrow p4\pi$
1450 ± 100	340	SCHACHT 74	STRC	9–18 $\gamma p \rightarrow p4\pi$
1430 ± 50	400	BINGHAM 72B	HBC	9.3 $\gamma p \rightarrow p4\pi$

NODE=M065M4
 NODE=M065M4

OCCUR=2

¹ Simple relativistic Breit-Wigner fit with model dependent width.

² Simple relativistic Breit-Wigner fit with constant width.

³ One peak fit result.

⁴ Parameters roughly estimated, not from a fit.

⁵ Skew mass distribution compensated by Ross-Stodolsky factor.

NODE=M065M;LINKAGE=A

NODE=M065M4;LINKAGE=M

NODE=M065M;LINKAGE=O

NODE=M065M;LINKAGE=C

NODE=M065M;LINKAGE=D

$\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1660 ± 30	ATKINSON 85B	OMEG	20–70 γp

NODE=M065M5
 NODE=M065M5

• • • We do not use the following data for averages, fits, limits, etc. • • •

3($\pi^+\pi^-$) AND 2($\pi^+\pi^-\pi^0$) MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1730±34	¹ FRABETTI	04	E687 $\gamma p \rightarrow 3\pi^+ 3\pi^- p$
1783±15	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

¹ From a fit with two resonances with the JACOB 72 continuum.

NODE=M065M7
NODE=M065M7

NODE=M065M;LINKAGE=PI

 $m_{\rho(1700)^0} - m_{\rho(1700)\pm}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

-48.30±83.81	¹ BARTOS	17A	RVUE $e^+e^- \rightarrow \pi^+\pi^-$, $\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
--------------	---------------------	-----	---

¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M065DM

NODE=M065DM

NODE=M065DM;LINKAGE=A

 $\rho(1700)$ WIDTH **$\eta\rho^0$ AND $\pi^+\pi^-$ MODES**

VALUE (MeV)	DOCUMENT ID
-------------	-------------

250±100 OUR ESTIMATE

NODE=M065210

NODE=M065W0
NODE=M065W0

→ UNCHECKED ←

 $\eta\rho^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M065W6
NODE=M065W6

• • • We do not use the following data for averages, fits, limits, etc. • • •

47±19	13.4k	¹ GRIBANOV	20	CMD3	1.1-2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
132±40	7.4k	² ACHASOV	18	SND	1.22-2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
150±30		ANTONELLI	88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
282±44		³ FUKUI	88	SPEC	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

¹ Mass and width of the $\rho(770)$ fixed at 775 and 149 MeV, respectively; solution 2 of model 2, $\eta \rightarrow \gamma\gamma$ decays used.

² From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

³ Assuming $\rho^+ f_0(1370)$ decay mode interferes with $a_1(1260)^+\pi$ background. From a two Breit-Wigner fit.

NODE=M065W6;LINKAGE=B

NODE=M065W6;LINKAGE=A

NODE=M065W;LINKAGE=B

 $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M065W1
NODE=M065W1

• • • We do not use the following data for averages, fits, limits, etc. • • •

268.98±11.40		¹ BARTOS	17	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
489.58±16.95		² BARTOS	17A	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
414.71±119.48		³ BARTOS	17A	RVUE	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
109 ± 19	20k	⁴ LEES	17C	BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
310 ± 30	$\frac{+25}{-35}$ 63.5k	⁵ ABRAMOWICZ12		ZEUS	$ep \rightarrow e\pi^+\pi^-p$
316 ± 26		⁶ LEES	12G	BABR	$e^+e^- \rightarrow \pi^+\pi^-\gamma$
164 ± 21	$\frac{+89}{-26}$ 5.4M	^{7,8} FUJIKAWA	08	BELL	$\tau^- \rightarrow \pi^-\pi^0\nu_\tau$
275 ± 45		⁹ ABELE	97	CBAR	$\bar{p}n \rightarrow \pi^-\pi^0\pi^0$
310 ± 40		⁹ BERTIN	97C	OBLX	0.0 $\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
400 ± 100		CLEGG	94	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
224 ± 22		BISELLO	89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
242.5 ± 163.0		DUBNICKA	89	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$
620 ± 60		GESHKEN...	89	RVUE	
<315		¹⁰ ERKAL	85	RVUE	20-70 $\gamma p \rightarrow \gamma\pi$
280 $\frac{+30}{-80}$		ABE	84B	HYBR	20 $\gamma p \rightarrow \pi^+\pi^-p$

OCCUR=2

230 ± 80		11 ASTON	80 OMEG	20-70	$\gamma p \rightarrow p 2\pi$
283 ± 14		12 ATIYA	79B SPEC	50	$\gamma C \rightarrow C 2\pi$
175 + 98 - 53		BECKER	79 ASPK	17	$\pi^- p$ polarized
232 ± 34		10 LANG	79 RVUE		
340		10 MARTIN	78C RVUE	17	$\pi^- p \rightarrow \pi^+ \pi^- n$
300 ± 100		10 FROGGATT	77 RVUE	17	$\pi^- p \rightarrow \pi^+ \pi^- n$
180 ± 50		13 HYAMS	73 ASPK	17	$\pi^- p \rightarrow \pi^+ \pi^- n$

- 1 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of LEES 12G and ABLIKIM 16C.
- 2 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, and AMBROSINO 11A.
- 3 Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of FUJIKAWA 08.
- 4 From a Dalitz plot analysis in an isobar model with $\rho(1450)$ and $\rho(1700)$ masses and widths floating.
- 5 Using the KUHN 90 parametrization of the pion form factor, neglecting $\rho-\omega$ interference.
- 6 Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.
- 7 $|F_\pi(0)|^2$ fixed to 1.
- 8 From the GOUNARIS 68 parametrization of the pion form factor.
- 9 T-matrix pole.
- 10 From phase shift analysis of HYAMS 73 data.
- 11 Simple relativistic Breit-Wigner fit with constant width.
- 12 An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.
- 13 Included in BECKER 79 analysis.

NODE=M065W1;LINKAGE=C
 NODE=M065W1;LINKAGE=D
 NODE=M065W1;LINKAGE=E
 NODE=M065W1;LINKAGE=A
 NODE=M065W1;LINKAGE=AB
 NODE=M065W1;LINKAGE=LE
 NODE=M065W1;LINKAGE=FU
 NODE=M065W1;LINKAGE=GO
 NODE=M065W;LINKAGE=QQ
 NODE=M065W;LINKAGE=P
 NODE=M065W;LINKAGE=M
 NODE=M065W;LINKAGE=R
 NODE=M065W;LINKAGE=H

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

150.9 ± 2.5 ⁺⁶⁰ _{-10.6}		1 ALBRECHT	20 RVUE		0.9 $\bar{p} p \rightarrow K^+ K^- \pi^0$
187.2 ± 26.7	27k	2 ABELE	99D CBAR	±	0.0 $\bar{p} p \rightarrow K^+ K^- \pi^0$
265 ± 120	1600	CLELAND	82B SPEC	±	50 $\pi p \rightarrow K_S^0 K^\pm p$

- 1 T-matrix pole, 2 poles, 3 channels, including $\pi\pi$ scattering data from HYAMS 75.
- 2 K-matrix pole. Isospin not determined, could be $\omega(1650)$ or $\phi(1680)$.

NODE=M065W2;LINKAGE=A
 NODE=M065W2;LINKAGE=AN

$2(\pi^+\pi^-)$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

510 ± 40		1 CORDIER	82 DM1		$e^+ e^- \rightarrow 2(\pi^+\pi^-)$
400 ± 50		2 ASTON	81E OMEG	20-70	$\gamma p \rightarrow p 4\pi$
400 ± 146		3 DIBIANCA	81 DBC		$\pi^+ d \rightarrow p p 2(\pi^+\pi^-)$
700 ± 160		1 BACCI	80 FRAG		$e^+ e^- \rightarrow 2(\pi^+\pi^-)$
100	34	KILLIAN	80 SPEC	11	$e^- p \rightarrow 2(\pi^+\pi^-)$
600		4 ATIYA	79B SPEC	50	$\gamma C \rightarrow C 4\pi^\pm$
340 ± 160	65	5 ALEXANDER	75 HBC	7.5	$\gamma p \rightarrow p 4\pi$
360 ± 100		2 CONVERSI	74 OSPK		$e^+ e^- \rightarrow 2(\pi^+\pi^-)$
400 ± 120	160	6 SCHACHT	74 STRC	5.5-9	$\gamma p \rightarrow p 4\pi$
850 ± 200	340	6 SCHACHT	74 STRC	9-18	$\gamma p \rightarrow p 4\pi$
650 ± 100	400	BINGHAM	72B HBC	9.3	$\gamma p \rightarrow p 4\pi$

- 1 Simple relativistic Breit-Wigner fit with model-dependent width.
- 2 Simple relativistic Breit-Wigner fit with constant width.
- 3 One peak fit result.
- 4 Parameters roughly estimated, not from a fit.
- 5 Skew mass distribution compensated by Ross-Stodolsky factor.
- 6 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M065W4
 NODE=M065W4

OCCUR=2

NODE=M065W;LINKAGE=A
 NODE=M065W4;LINKAGE=M
 NODE=M065W;LINKAGE=O
 NODE=M065W;LINKAGE=C
 NODE=M065W;LINKAGE=D
 NODE=M065W;LINKAGE=E

$\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

300 ± 50	ATKINSON	85B OMEG	20-70 γp
----------	----------	----------	------------------

NODE=M065W5
 NODE=M065W5

$\omega\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
371±3	¹ ACHASOV	23A	SND $e^+e^- \rightarrow \omega\pi^0$
350 to 580	² ACHASOV	00I	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
490 to 1040	³ ACHASOV	00I	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ From a vector dominance fit to the Born cross section between 1.05 and 2.0 GeV with $\rho(770)$, $\rho(1570)$, $\rho(1700)$, $\rho(2150)$. The fit also uses SND data from the VEPP-2M collider below 1.02 GeV and from LEES 17H and ABLIKIM 21A above 1.5 GeV.

² Taking into account both $\rho(1450)$ and $\rho(1700)$ contributions. Using the data of ACHASOV 00I on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$. $\rho(1450)$ mass and width fixed at 1400 MeV and 500 MeV respectively.

³ Taking into account the $\rho(1700)$ contribution only. Using the data of ACHASOV 00I on $e^+e^- \rightarrow \omega\pi^0$ and of EDWARDS 00A on $\tau^- \rightarrow \omega\pi^- \nu_\tau$.

NODE=M065W9
NODE=M065W9

OCCUR=2

NODE=M065W9;LINKAGE=A

NODE=M065W;LINKAGE=I1

NODE=M065W;LINKAGE=I2

 $3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
315±100	¹ FRABETTI	04	E687 $\gamma p \rightarrow 3\pi^+3\pi^-p$
285±20	CLEGG	90	RVUE $e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

¹ From a fit with two resonances with the JACOB 72 continuum.

NODE=M065W7
NODE=M065W7

NODE=M065W;LINKAGE=PI

$$\Gamma_{\rho(1700)^0} = \Gamma_{\rho(1700)^\pm}$$

NODE=M065DW

NODE=M065DW

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
74.87±120.67	¹ BARTOS	17A	RVUE $e^+e^- \rightarrow \pi^+\pi^-, \tau^- \rightarrow \pi^-\pi^0\nu_\tau$

¹ Applies the Unitary & Analytic Model of the pion electromagnetic form factor of DUB-NICKA 10 to analyze the data of ACHASOV 06, AKHMETSHIN 07, AUBERT 09AS, AMBROSINO 11A, and FUJIKAWA 08.

NODE=M065DW;LINKAGE=A

 $\rho(1700)$ DECAY MODES

NODE=M065215;NODE=M065

Mode	Fraction (Γ_i/Γ)
Γ_1 4π	
Γ_2 $2(\pi^+\pi^-)$	seen
Γ_3 $\rho\pi\pi$	seen
Γ_4 $\rho^0\pi^+\pi^-$	seen
Γ_5 $\rho^0\pi^0\pi^0$	
Γ_6 $\rho^\pm\pi^\mp\pi^0$	seen
Γ_7 $a_1(1260)\pi$	seen
Γ_8 $h_1(1170)\pi$	seen
Γ_9 $\pi(1300)\pi$	seen
Γ_{10} $\rho\rho$	seen
Γ_{11} $\pi^+\pi^-$	seen
Γ_{12} $\pi\pi$	seen
Γ_{13} $K\bar{K}^*(892)+c.c.$	seen
Γ_{14} $\eta\rho$	seen
Γ_{15} $a_2(1320)\pi$	not seen
Γ_{16} $K\bar{K}$	seen
Γ_{17} e^+e^-	seen
Γ_{18} $\pi^0\omega$	seen
Γ_{19} $\pi^0\gamma$	not seen
Γ_{20} $f_0(1500)\gamma$	not seen

DESIG=20

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=12;OUR EST;→ UNCHECKED ←

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=7

DESIG=9;OUR EST;→ UNCHECKED ←

DESIG=15;OUR EST;→ UNCHECKED ←

DESIG=16;OUR EST;→ UNCHECKED ←

DESIG=17;OUR EST;→ UNCHECKED ←

DESIG=18;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=13;OUR EST;→ UNCHECKED ←

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=11;OUR EST;→ UNCHECKED ←

DESIG=14;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=194

DESIG=195

$$\rho(1700) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$$

NODE=M065225

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the cross-section into channel i in e^+e^- annihilation.

NODE=M065225

 $\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$

$\Gamma_2\Gamma_{17}/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.6 ±0.2	DELCOURT	81B	DM1 $e^+e^- \rightarrow 2(\pi^+\pi^-)$
2.83±0.42	BACCI	80	FRAG $e^+e^- \rightarrow 2(\pi^+\pi^-)$

NODE=M065G2
NODE=M065G2

$$\Gamma(\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}\Gamma_{17}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13	¹ DIEKMAN	88	RVUE $e^+e^- \rightarrow \pi^+\pi^-$
0.029 ^{+0.016} -0.012	KURDADZE	83	OLYA 0.64–1.4 $e^+e^- \rightarrow \pi^+\pi^-$

¹ Using total width = 220 MeV.

NODE=M065G4
NODE=M065G4

NODE=M065G4;LINKAGE=B

$$\Gamma(K\bar{K}^*(892)+c.c.) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}\Gamma_{17}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.305±0.071	¹ BIZOT	80	DM1 e^+e^-
-------------	--------------------	----	--------------

¹ Model dependent.

NODE=M065G10
NODE=M065G10

NODE=M065G;LINKAGE=M

$$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}\Gamma_{17}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35±0.53±0.08	13.4k	¹ GRIBANOV	20	CMD3 1.1–2.0 $e^+e^- \rightarrow \eta\pi^+\pi^-$
84 ±26 ±4		² LEES	18	BABR $e^+e^- \rightarrow \eta\pi^+\pi^-$
7 ±3		ANTONELLI	88	DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$

¹ Mass and width of the $\rho(770)$ fixed at 775 and 149 MeV, respectively; solution 2 of model 2, $\eta \rightarrow \gamma\gamma$ decays used.

² Includes non-resonant contribution. The selected fit model includes three ρ excited states. Model uncertainty is 80%.

NODE=M065G11
NODE=M065G11

NODE=M065G11;LINKAGE=B

NODE=M065G11;LINKAGE=A

$$\Gamma(K\bar{K}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}\Gamma_{17}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.035±0.029	¹ BIZOT	80	DM1 e^+e^-
-------------	--------------------	----	--------------

¹ Model dependent.

NODE=M065G5
NODE=M065G5

NODE=M065G5;LINKAGE=M

$$\Gamma(\rho\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_{17}/\Gamma$$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.510±0.090	¹ BIZOT	80	DM1 e^+e^-
-------------	--------------------	----	--------------

¹ Model dependent.

NODE=M065G12
NODE=M065G12

NODE=M065G12;LINKAGE=M

$$\rho(1700) \Gamma(i)/\Gamma(\text{total}) \times \Gamma(e^+e^-)/\Gamma(\text{total})$$

NODE=M065240

$$\Gamma(\pi^0\omega)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{17}/\Gamma$$

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.09±0.05	10.2k	¹ ACHASOV	16D	SND 1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
1.7 ±0.4	7815	² ACHASOV	13	SND 1.05–2.00 $e^+e^- \rightarrow \pi^0\pi^0\gamma$

¹ From a phenomenological model based on vector meson dominance with interfering $\rho(700)$, $\rho(1450)$, and $\rho(1700)$. The $\rho(1700)$ mass and width are fixed at 1720 MeV and 250 MeV, respectively. Systematic uncertainty not estimated. Supersedes ACHASOV 13.

² From a phenomenological model based on vector meson dominance with the interfering $\rho(1450)$ and $\rho(1700)$ and their widths fixed at 400 and 250 MeV, respectively. Systematic uncertainty not estimated.

NODE=M065R01
NODE=M065R01

OCCUR=4

NODE=M065R01;LINKAGE=B

NODE=M065R01;LINKAGE=AC

$$\Gamma(\eta\rho)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma \times \Gamma_{17}/\Gamma$$

VALUE (units 10 ⁻⁸)	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.3 ^{+3.8} -3.1	7.4k	¹ ACHASOV	18	SND 1.22–2.00 $e^+e^- \rightarrow \eta\pi^+\pi^-$
-----------------------------	------	----------------------	----	---

¹ From the combined fit of AULCHENKO 15 and ACHASOV 18 in the model with the interfering $\rho(1450)$, $\rho(1700)$ and $\rho(2150)$ with the parameters of the $\rho(1450)$ and $\rho(1700)$ floating and the mass and width of the $\rho(2150)$ fixed at 2155 MeV and 320 MeV, respectively. The phases of the resonances are π , 0 and π , respectively.

NODE=M065R00
NODE=M065R00

NODE=M065R00;LINKAGE=A

$\rho(1700)$ BRANCHING RATIOS

NODE=M065230

 $\Gamma(\rho\pi\pi)/\Gamma(4\pi)$ **Γ_3/Γ_1**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.28±0.06 ¹ ABELE 01B CBAR 0.0 $\bar{p}n \rightarrow 5\pi$ ¹ $\omega\pi$ not included.NODE=M065R19
NODE=M065R19

NODE=M065R;LINKAGE=BL

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2(\pi^+\pi^-))$ **Γ_4/Γ_2**

VALUE	EVS	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

~ 1.0 DELCOURT 81B DM1 $e^+e^- \rightarrow 2(\pi^+\pi^-)$ 0.7 ±0.1 500 SCHACHT 74 STRC 5.5-18 $\gamma p \rightarrow p4\pi$ 0.80 ¹ BINGHAM 72B HBC 9.3 $\gamma p \rightarrow p4\pi$ ¹ The $\pi\pi$ system is in S-wave.NODE=M065R1
NODE=M065R1

NODE=M065R1;LINKAGE=S

 $\Gamma(\rho^0\pi^0\pi^0)/\Gamma(\rho^\pm\pi^\mp\pi^0)$ **Γ_5/Γ_6**

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

<0.10 ATKINSON 85B OMEG 20-70 γp <0.15 ATKINSON 82 OMEG 0 20-70 $\gamma p \rightarrow p4\pi$ NODE=M065R6
NODE=M065R6 **$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$** **$\Gamma_7/\Gamma_1$**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.16±0.05 ¹ ABELE 01B CBAR 0.0 $\bar{p}n \rightarrow 5\pi$ ¹ $\omega\pi$ not included.NODE=M065R15
NODE=M065R15

NODE=M065R15;LINKAGE=BL

 $\Gamma(h_1(1170)\pi)/\Gamma(4\pi)$ **Γ_8/Γ_1**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.17±0.06 ¹ ABELE 01B CBAR 0.0 $\bar{p}n \rightarrow 5\pi$ ¹ $\omega\pi$ not included.NODE=M065R16
NODE=M065R16

NODE=M065R16;LINKAGE=BL

 $\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$ **Γ_9/Γ_1**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.30±0.10 ¹ ABELE 01B CBAR 0.0 $\bar{p}n \rightarrow 5\pi$ ¹ $\omega\pi$ not included.NODE=M065R17
NODE=M065R17

NODE=M065R17;LINKAGE=BL

 $\Gamma(\rho\rho)/\Gamma(4\pi)$ **Γ_{10}/Γ_1**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.09±0.03 ¹ ABELE 01B CBAR 0.0 $\bar{p}n \rightarrow 5\pi$ ¹ $\omega\pi$ not included.NODE=M065R18
NODE=M065R18

NODE=M065R18;LINKAGE=BL

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{11}/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.108±0.017^{+0.162}_{-0.004} ¹ ALBRECHT 20 RVUE 0.9 $\bar{p}p \rightarrow K^+K^-\pi^0$ 0.287^{+0.043}_{-0.042} BECKER 79 ASPK 17 π^-p polarized0.15 to 0.30 ² MARTIN 78C RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$ <0.20 ³ COSTA... 77B RVUE $e^+e^- \rightarrow 2\pi, 4\pi$ 0.30 ±0.05 ² FROGGATT 77 RVUE 17 $\pi^-p \rightarrow \pi^+\pi^-n$ <0.15 ⁴ EISENBERG 73 HBC 5 $\pi^+p \rightarrow \Delta^{++}2\pi$ 0.25 ±0.05 ⁵ HYAMS 73 ASPK 17 $\pi^-p \rightarrow \pi^+\pi^-n$ ¹ Residue from T-matrix pole, 2 poles, 3 channels, Chew-Mandelstam functions and simplified analytic continuation for the 4π channel. Includes scattering data from HYAMS 75 and model-independent calculation of GARCIA-MARTIN 11A.² From phase shift analysis of HYAMS 73 data.³ Estimate using unitarity, time reversal invariance, Breit-Wigner.⁴ Estimated using one-pion-exchange model.⁵ Included in BECKER 79 analysis.NODE=M065R5
NODE=M065R5

NODE=M065R5;LINKAGE=A

NODE=M065R5;LINKAGE=P

NODE=M065R5;LINKAGE=C

NODE=M065R5;LINKAGE=E

NODE=M065R5;LINKAGE=H

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.007 \pm 0.006^{+0.041}_{-0.002}$	¹ ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow K^+ K^- \pi^0$
-------------------------------------	-----------------------	----	---

¹ Residue from T-matrix pole, 2 poles, 3 channels, Chew-Mandelstam functions and simplified analytic continuation for the 4π channel. Includes scattering data from HYAMS 75 and model-independent calculation of GARCIA-MARTIN 11A.

 Γ_{16}/Γ

NODE=M065R03
NODE=M065R03

NODE=M065R03;LINKAGE=A

 $\Gamma(\pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ± 0.05	ASTON	80	OMEG $20-70 \gamma p \rightarrow p 2\pi$
< 0.14	¹ DAVIER	73	STRC $6-18 \gamma p \rightarrow p 4\pi$
< 0.2	² BINGHAM	72B	HBC $9.3 \gamma p \rightarrow p 2\pi$

¹ Upper limit is estimate.

² 2σ upper limit.

 Γ_{11}/Γ_2

NODE=M065R3
NODE=M065R3

NODE=M065R3;LINKAGE=E
NODE=M065R3;LINKAGE=S

 $\Gamma(\pi\pi)/\Gamma(4\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.16 ± 0.04	^{1,2} ABELE	01B	CBAR $0.0 \bar{p}n \rightarrow 5\pi$
-----------------	----------------------	-----	--------------------------------------

¹ Using ABELE 97.

² $\omega\pi$ not included.

 Γ_{12}/Γ_1

NODE=M065R20
NODE=M065R20

NODE=M065R;LINKAGE=LK
NODE=M065R20;LINKAGE=BL

 $\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	COAN	04	CLEO $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$
---------------	------	----	--

 Γ_{13}/Γ

NODE=M065R21
NODE=M065R21

 $\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(2(\pi^+ \pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.15 ± 0.03	¹ DELCOURT	81B	DM1 $e^+ e^- \rightarrow \bar{K} K \pi$
-----------------	-----------------------	-----	---

¹ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

 Γ_{13}/Γ_2

NODE=M065R9
NODE=M065R9

NODE=M065R9;LINKAGE=D

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen		AKHMETSHIN 00D	CMD2	$e^+ e^- \rightarrow \eta \pi^+ \pi^-$
< 0.04		DONNACHIE 87B	RVUE	
< 0.02	58	ATKINSON 86B	OMEG	$20-70 \gamma p$

 Γ_{14}/Γ

NODE=M065R12
NODE=M065R12

 $\Gamma(\eta\rho)/\Gamma(2(\pi^+ \pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.123 ± 0.027	DELCOURT	82	DM1 $e^+ e^- \rightarrow \pi^+ \pi^- \text{MM}$
~ 0.1	ASTON	80	OMEG $20-70 \gamma p$

 Γ_{14}/Γ_2

NODE=M065R8
NODE=M065R8

 $\Gamma(\pi^+ \pi^- \text{ neutrals})/\Gamma(2(\pi^+ \pi^-))$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.4	¹ BALLAM	74	HBC $9.3 \gamma p$
---------------	---------------------	----	--------------------

¹ Upper limit. Background not subtracted.

 $(\Gamma_5 + \Gamma_6 + 0.714\Gamma_{14})/\Gamma_2$

NODE=M065R7
NODE=M065R7

NODE=M065R7;LINKAGE=U

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	AMELIN	00	VES $37 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
----------	--------	----	---

 Γ_{15}/Γ

NODE=M065R14
NODE=M065R14

 $\Gamma(K\bar{K})/\Gamma(2(\pi^+ \pi^-))$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.015 ± 0.010		¹ DELCOURT	81B	DM1	$e^+ e^- \rightarrow \bar{K} K$
< 0.04	95	BINGHAM	72B	HBC	0 $9.3 \gamma p$

¹ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

 Γ_{16}/Γ_2

NODE=M065R4
NODE=M065R4

NODE=M065R4;LINKAGE=D

$\Gamma(K\bar{K})/\Gamma(K\bar{K}^*(892)+c.c.)$ Γ_{16}/Γ_{13}

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.052 ± 0.026	BUON	82	DM1 $e^+e^- \rightarrow$ hadrons
---------------	------	----	----------------------------------

NODE=M065R10
NODE=M065R10

 $\Gamma(\pi^0\omega)/\Gamma_{total}$ Γ_{18}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		MATVIENKO	15	BELL $\bar{B}^0 \rightarrow D^{*+}\omega\pi^-$
seen	1.6k	ACHASOV	12	SND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
not seen	2382	AKHMETSHIN	03B	CMD2 $e^+e^- \rightarrow \pi^0\pi^0\gamma$
seen		ACHASOV	97	RVUE $e^+e^- \rightarrow \omega\pi^0$

NODE=M065R13
NODE=M065R13

 $\Gamma(\pi^0\gamma)/\Gamma_{total}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ¹ACHASOV 10D SND 1.075–2.0 $e^+e^- \rightarrow \pi^0\gamma$

¹From a fit of a VMD model with two effective resonances with masses of 1450 MeV and 1700 MeV to describe the excited vector states $\omega(1420)$, $\rho(1450)$, $\omega(1650)$, and $\rho(1700)$. The width of the highest mass effective resonance is fixed at 315 MeV.

NODE=M065R02
NODE=M065R02

NODE=M065R02;LINKAGE=A

 $\Gamma(f_0(1500)\gamma)/\Gamma_{total}$ Γ_{20}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ¹ACHASOV 22 SND 1.17–2.00 $e^+e^- \rightarrow \eta\eta\gamma$

¹The 90% CL upper limit on the Born cross sections $\sigma(e^+e^- \rightarrow \phi(1680) \rightarrow f_2'(1525)\gamma \rightarrow \eta\eta\gamma)$ and $\sigma(e^+e^- \rightarrow \rho(1700) \rightarrow f_0(1500)\gamma \rightarrow \eta\eta\gamma)$ is 10.6 pb.

NODE=M065R04
NODE=M065R04

NODE=M065R04;LINKAGE=A

 $\rho(1700)$ REFERENCES

NODE=M065

ACHASOV	23A	PR D108 092012	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=62428
ACHASOV	22	EPJ C82 168	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=61655
ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61028
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
GRIBANOV	20	JHEP 2001 112	S.S. Gribov <i>et al.</i>	(CMD-3 Collab.)	REFID=60620
ACHASOV	18	PR D97 012008	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=58785
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
BARTOS	17	PR D96 113004	E. Bartos <i>et al.</i>		REFID=58325
BARTOS	17A	IJMP A32 1750154	E. Bartos <i>et al.</i>		REFID=58367
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
LEES	17H	PR D96 092009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58311
AAIJ	16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM	16C	PL B753 629	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57138
ACHASOV	16D	PR D94 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=57618
AULCHENKO	15	PR D91 052013	V.M. Aulchenko <i>et al.</i>	(SND Collab.)	REFID=56793
MATVIENKO	15	PR D92 012013	D. Matvienko <i>et al.</i>	(BELLE Collab.)	REFID=56601
ACHASOV	13	PR D88 054013	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=55584
ABRAMOWICZ	12	EPJ C72 1869	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54274
ACHASOV	12	JETPL 94 734	M.N. Achasov <i>et al.</i>		REFID=54275
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
AMBROSINO	11A	PL B700 102	F. Ambrosino <i>et al.</i>	(KLOE Collab.)	REFID=16683
GARCIA-MAR...	11A	PR D83 074004	R. Garcia-Martin <i>et al.</i>	(MADR, CRAC)	REFID=54121
ACHASOV	10D	PR D98 112001	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=59494
DUBNICKA	10	APS 60 1	S. Dubnicka, A.Z. Dubnickova		REFID=58797
AUBERT	09AS	PRL 103 231801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53136
FUJIKAWA	08	PR D78 072006	M. Fujikawa <i>et al.</i>	(BELLE Collab.)	REFID=52536
AKHMETSHIN	07	PL B648 28	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=51615
ACHASOV	06	JETP 103 380	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=51113
COAN	04	PRL 92 232001	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=49945
FRABETTI	04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=49614
AKHMETSHIN	03B	PL B562 173	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=49406
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48356
ACHASOV	00I	PL B486 29	M.N. Achasov <i>et al.</i>	(Novosibirsk SND Collab.)	REFID=47931
AKHMETSHIN	00D	PL B489 125	R.R. Akhmetshin <i>et al.</i>	(Novosibirsk CMD-2 Collab.)	REFID=47935
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
EDWARDS	00A	PR D61 072003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=47465
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
ABELE	97	PL B391 191	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45415
ACHASOV	97	PR D55 2663	N.N. Achasov <i>et al.</i>	(NOVM)	REFID=45382
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)	REFID=45701
CLEGG	94	ZPHY C62 455	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=44081
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=41355
KUHN	90	ZPHY C48 445	J.H. Kuhn <i>et al.</i>	(MPIM)	REFID=45862
BISELLO	89	PL B220 321	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=40740
DUBNICKA	89	JP G15 1349	S. Dubnicka <i>et al.</i>	(JINR, SLOV)	REFID=44082
GESHKEN...	89	ZPHY C45 351	B.V. Geshkenbein	(ITEP)	REFID=41017
ANTONELLI	88	PL B212 133	A. Antonelli <i>et al.</i>	(DM2 Collab.)	REFID=40583
DIEKMAN	88	PRPL 159 99	B. Diekmann	(BONN)	REFID=40272
FUKUI	88	PL B202 441	S. Fukui <i>et al.</i>	(SUGI, NAGO, KEK, KYOT+)	REFID=40273
DONNACHIE	87B	ZPHY C34 257	A. Donnachie, A.B. Clegg	(MCHS, LANC)	REFID=40920
ATKINSON	86B	ZPHY C30 531	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21508
ATKINSON	85B	ZPHY C26 499	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21506
ERKAL	85	ZPHY C29 485	C. Erkal, M.G. Olsson	(WISC)	REFID=20136
ABE	84B	PRL 53 751	K. Abe <i>et al.</i>	(SLAC HFP Collab.)	REFID=21503
KURDADZE	83	JETPL 37 733	L.M. Kurdadze <i>et al.</i>	(NOVO)	REFID=20133
Translated from ZETFP 37 613.					

ATKINSON	82	PL 108B 55	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)
BUON	82	PL 118B 221	J. Buon <i>et al.</i>	(LALO, MONP)
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)
CORDIER	82	PL 109B 129	A. Cordier <i>et al.</i>	(LALO)
DEL COURT	82	PL 113B 93	B. Delcourt <i>et al.</i>	(LALO)
ASTON	81E	NP B189 15	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
DEL COURT	81B	Bonn Conf. 205	B. Delcourt	(ORSAY)
Also		PL 109B 129	A. Cordier <i>et al.</i>	(LALO)
DIBIANCA	81	PR D23 595	F.A. di Bianca <i>et al.</i>	(CASE, CMU)
ASTON	80	PL 92B 215	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)
BACCI	80	PL 95B 139	C. Bacci <i>et al.</i>	(ROMA, FRAS)
BIZOT	80	Madison Conf. 546	J.C. Bizot <i>et al.</i>	(LALO, MONP)
KILLIAN	80	PR D21 3005	T.J. Killian <i>et al.</i>	(CORN)
ATIYA	79B	PRL 43 1691	M.S. Atiya <i>et al.</i>	(COLU, ILL, FNAL)
BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
LANG	79	PR D19 956	C.B. Lang, A. Mas-Parareda	(GRAZ)
MARTIN	78C	ANP 114 1	A.D. Martin, M.R. Pennington	(CERN)
COSTA...	77B	PL 71B 345	B. Costa de Beauregard, B. Pire, T.N. Truong	(EPOL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ALEXANDER	75	PL 57B 487	G. Alexander <i>et al.</i>	(TELA)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
BALLAM	74	NP B76 375	J. Ballam <i>et al.</i>	(SLAC, LBL, MPIM)
CONVERSI	74	PL 52B 493	M. Conversi <i>et al.</i>	(ROMA, FRAS)
SCHACHT	74	NP B81 205	P. Schacht <i>et al.</i>	(MPIM)
DAVIER	73	NP B58 31	M. Davier <i>et al.</i>	(SLAC)
EISENBERG	73	PL 43B 149	Y. Eisenberg <i>et al.</i>	(REHO)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
BINGHAM	72B	PL 41B 635	H.H. Bingham <i>et al.</i>	(LBL, UCB, SLAC) IJGP
JACOB	72	PR D5 1847	M. Jacob, R. Slansky	
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai	

REFID=21493
REFID=21494
REFID=21281
REFID=21495
REFID=21496
REFID=21487
REFID=21490
REFID=21495
REFID=21492
REFID=21478
REFID=21481
REFID=21482
REFID=21484
REFID=21470
REFID=21084
REFID=20126
REFID=21661
REFID=21465
REFID=21072
REFID=21450
REFID=20355
REFID=20610
REFID=20637
REFID=21449
REFID=21434
REFID=21435
REFID=20107
REFID=21426
REFID=49668
REFID=48054

NODE=M162

 $a_2(1700)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

 $a_2(1700)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1630–1780) – i (60–250) OUR ESTIMATE			
$(1686 \pm 22^{+19}_{-7}) - i(211 \pm 38^{+32}_{-29})$	1 KOPF	21	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$ and 191 $\pi^- p \rightarrow \pi^- \pi^- \pi^+ p$
$(1638.9 \pm 2.3^{+57.4}_{-0.1}) - i(112.0 \pm 1.3^{+0.9}_{-24.2})$	2 ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta, \pi^0 K^+ K^-$
$(1722 \pm 15 \pm 67) - i(124 \pm 9 \pm 32)$	3 RODAS	19	RVUE $191 \pi^- p \rightarrow \eta' \pi^- p$
$(1698 \pm 44) - i(133 \pm 28)$	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta$

¹Based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi$, $\eta'\pi$ and $K\bar{K}$ systems.

²Based on 2 poles, 2 channels ($\pi\eta$, $K\bar{K}$).

³The coupled-channel analysis of both the $\eta\pi$ and $\eta'\pi$ systems using ADOLPH 15 data.

NODE=M162PP

NODE=M162PP

NODE=M162PP

→ UNCHECKED ←

NODE=M162PP;LINKAGE=A

NODE=M162PP;LINKAGE=B

NODE=M162PP;LINKAGE=C

 $a_2(1700)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1706 ± 14 OUR AVERAGE				
1681^{+22}_{-35}	46M	1,2 AGHASYAN	18B	COMP $190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$1726 \pm 12 \pm 25$		2 ABLIKIM	17K	BES3 $\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
$1722 \pm 9 \pm 15$	18k	3 SCHEGELSKY	06	RVUE $\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$
1660 ± 40		2 ABELE	99B	CBAR $1.94 \bar{p}p \rightarrow \pi^0 \eta \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$1720 \pm 10 \pm 60$		4 JACKURA	18	RVUE $\pi^- p \rightarrow \eta \pi^- p$
1675 ± 25		ANISOVICH	09	RVUE $0.0 \bar{p}p, \pi N$
1702 ± 7	80k	5 UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta \eta \pi^0$
$1721 \pm 13 \pm 44$	145k	LU	05	B852 $18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$
$1737 \pm 5 \pm 7$		ABE	04	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
1767 ± 14	221	6 ACCIARRI	01H	L3 $\gamma \gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} = 91, 183-209 \text{ GeV}$
~ 1775		7 GRYGOREV	99	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
$1752 \pm 21 \pm 4$		ACCIARRI	97T	L3 $\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

NODE=M162M

NODE=M162M

- ¹ Statistical error negligible.
- ² Breit-Wigner mass.
- ³ From analysis of L3 data at 183–209 GeV.
- ⁴ Superseded by RODAS 19.
- ⁵ Statistical error only.
- ⁶ Spin 2 dominant, isospin not determined, could also be $I=1$.
- ⁷ Possibly two $J^P = 2^+$ resonances with isospins 0 and 1.

NODE=M162M;LINKAGE=B
 NODE=M162M;LINKAGE=E
 NODE=M162M;LINKAGE=SC
 NODE=M162M;LINKAGE=D
 NODE=M162M;LINKAGE=ST
 NODE=M162M;LINKAGE=HA
 NODE=M162M;LINKAGE=GR

$a_2(1700)$ WIDTH

NODE=M162W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M162W

380⁺₋₅₀ OUR AVERAGE Error includes scale factor of 3.9. See the ideogram below.

NEW


[378⁺₋₅₀ MeV OUR 2023 AVERAGE Scale factor = 3.9]

436 ⁺ ₋₁₆	46M	1,2 AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
190 \pm 18 \pm 30		² ABLIKIM	17K	BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
336 \pm 20 \pm 20	18k	³ SCHEGELSKY	06	RVUE	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$
280 \pm 70		² ABELE	99B	CBAR	1.94 $\bar{p} p \rightarrow \pi^0 \eta \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
280 \pm 10 \pm 70		⁴ JACKURA	18	RVUE	$\pi^- p \rightarrow \eta \pi^- p$
270 ⁺ ₋₂₀		ANISOVICH	09	RVUE	0.0 $\bar{p} p, \pi N$
417 \pm 19	80k	⁵ UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
279 \pm 49 \pm 66	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
151 \pm 22 \pm 24		ABE	04	BELL	10.6 $e^+ e^- \rightarrow$ $e^+ e^- K^+ K^-$
187 \pm 60	221	⁶ ACCIARRI	01H	L3	$\gamma \gamma \rightarrow K_S^0 K_S^0, E_{cm}^{ee} =$ 91, 183–209 GeV
150 \pm 110 \pm 34		ACCIARRI	97T	L3	$\gamma \gamma \rightarrow \pi^+ \pi^- \pi^0$

NODE=M162W;LINKAGE=B
 NODE=M162W;LINKAGE=E
 NODE=M162W;LINKAGE=SC
 NODE=M162W;LINKAGE=D
 NODE=M162W;LINKAGE=ST
 NODE=M162W;LINKAGE=HA

- ¹ Statistical error negligible.
- ² Breit-Wigner width.
- ³ From analysis of L3 data at 183–209 GeV.
- ⁴ Superseded by RODAS 19.
- ⁵ Statistical error only.
- ⁶ Spin 2 dominant, isospin not determined, could also be $I=1$.

WEIGHTED AVERAGE
 380+60-50 (Error scaled by 3.9)



$a_2(1700)$ DECAY MODES

NODE=M162215;NODE=M162

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi$	(2.5±0.6) %
Γ_2 $\eta'\pi$	seen
Γ_3 $\gamma\gamma$	(7.9±1.7) × 10 ⁻⁷
Γ_4 $\rho\pi$	seen
Γ_5 $f_2(1270)\pi$	seen
Γ_6 $K\bar{K}$	(1.3±0.8) %
Γ_7 $\omega\pi^-\pi^0$	seen
Γ_8 $\omega\rho$	seen

DESIG=4

DESIG=8;OUR EVAL;→ UNCHECKED ←

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=5

DESIG=6;OUR EVAL;→ UNCHECKED ←

DESIG=7;OUR EVAL;→ UNCHECKED ←

 $a_2(1700)$ PARTIAL WIDTHS

NODE=M162220

$\Gamma(\eta\pi)$					Γ_1
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
9.5±2.0	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
¹ From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.					

NODE=M162W3
NODE=M162W3

NODE=M162W3;LINKAGE=SC

$\Gamma(\gamma\gamma)$					Γ_3
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.30±0.05	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
¹ From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.					

NODE=M162W2
NODE=M162W2

NODE=M162W2;LINKAGE=SC

$\Gamma(K\bar{K})$					Γ_6
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
5.0±3.0	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
¹ From analysis of L3 data at 91 and 183–209 GeV, using $a_2(1700)$ mass of 1730 MeV and width of 340 MeV, and SU(3) relations.					

NODE=M162W1
NODE=M162W1

NODE=M162W1;LINKAGE=SC

 $a_2(1700)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M162225

$[\Gamma(\rho\pi) + \Gamma(f_2(1270)\pi)] \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$(\Gamma_4 + \Gamma_5)\Gamma_3/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.29±0.04±0.02		ACCIARRI	97T L3	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.37 ^{+0.12} _{-0.08} ±0.10	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	
¹ From analysis of L3 data at 183–209 GeV.					

NODE=M162G1
NODE=M162G1

NODE=M162G1;LINKAGE=SC

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_6\Gamma_3/\Gamma$
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
20.6 ± 4.2 ± 4.6	¹ ABE	04	BELL	10.6 e ⁺ e ⁻ → e ⁺ e ⁻ K ⁺ K ⁻	
49 ±11 ±13	² ACCIARRI	01H L3		$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{ee} = 91,$ 183–209 GeV	
¹ Assuming spin 2.					
² Spin 2 dominant, isospin not determined, could also be I=1.					

NODE=M162G2
NODE=M162G2NODE=M162G2;LINKAGE=AB
NODE=M162G2;LINKAGE=HA **$a_2(1700)$ BRANCHING RATIOS**

NODE=M162235

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$					Γ_4/Γ_5
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.4±0.4±0.1	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$	
¹ From analysis of L3 data at 183–209 GeV.					

NODE=M162R01
NODE=M162R01

NODE=M162R01;LINKAGE=SC

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$ Γ_6/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.029 ± 0.04	$^{+0.011}_{-0.012}$	¹ KOPF	21 RVUE	$0.9 p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and $191 \pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
4.134 ± 0.106	$^{+4.909}_{-2.988}$	² ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$

¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

² Residues from T-matrix pole, 2 poles, 2 channels ($\pi\eta, K\bar{K}$).

NODE=M162R00
NODE=M162R00

NODE=M162R00;LINKAGE=B

NODE=M162R00;LINKAGE=A

 $\Gamma(\eta'\pi)/\Gamma(\eta\pi)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.035 ± 0.044	$^{+0.069}_{-0.012}$	¹ KOPF	21 RVUE	$0.9 p\bar{p} \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$ and $191 \pi^- p \rightarrow$ $\pi^- \pi^- \pi^+ p$
-------------------	----------------------	-------------------	---------	---

¹ From T-matrix pole based on combined fit of Crystal Barrel and $\pi\pi$ scattering data (ALBRECHT 20), and COMPASS data (ADOLPH 15), using a coupled-channel model of $\eta\pi, \eta'\pi$ and $K\bar{K}$ systems.

NODE=M162R02
NODE=M162R02

NODE=M162R02;LINKAGE=A

 $a_2(1700)$ REFERENCES

KOPF	21	EPJ C81 1056	B. Kopf <i>et al.</i>	(BOCH)	REFID=61470
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
RODAS	19	PRL 122 042002	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=59554
AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
JACKURA	18	PL B779 464	A. Jackura <i>et al.</i>	(JPAC and COMPASS Collab.)	REFID=59003
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57953
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48321
ABELE	99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46904
GRYGOREV	99	PAN 62 470	V.K. Grygorev <i>et al.</i>		REFID=46909
ACCIARRI	97T	PL B413 147	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45761

NODE=M162

$a_0(1710)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

NODE=M263

OMITTED FROM SUMMARY TABLE

Evidence for this state is also inferred from the interference of the $K^+ K^-$ and $K_S^0 K_S^0$ decays of the $f_0(1710)$ in $D_s^+ \rightarrow f_0(1710) \pi^+$, leading to a relative branching ratio an order of magnitude larger than expected from isospin symmetry (ABLIKIM 22F). See also the review on "Spectroscopy of Light Meson Resonances."

NODE=M263

 $a_0(1710)$ MASS

NODE=M263M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1713±19 OUR AVERAGE	Error includes scale factor of 3.8. See the ideogram below. [1711 ± 27 MeV OUR 2023 AVERAGE Scale factor = 5.1]		
1736±10±12	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$
1817± 8±20	² ABLIKIM	22AH BES3	$D_s^+ \rightarrow K_S^0 K^+ \pi^0$
1704± 5± 2	LEES	21A BABR	$\eta_c(1S) \rightarrow \pi^+ \pi^- \eta$

NODE=M263M


NEW

¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

² Observed to decay into $K_S^0 K^+$ in a Breit-Wigner amplitude analysis involving D_s^+ decays into $\bar{K}^*(892)^0 K^+$, $\bar{K}^*(892)^+ K_S^0$, $\bar{K}^*(1410)^0 K^+$, $a_0(980)^+ \pi^0$, and $a_0(1817)^+ \pi^0$.

NODE=M263M;LINKAGE=B

NODE=M263M;LINKAGE=A

 **$a_0(1710)$ WIDTH**

NODE=M263W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
107±15 OUR AVERAGE	[106 ± 15 MeV OUR 2023 AVERAGE]		
134±17±61	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K \pi)$
97±22±15	² ABLIKIM	22AH BES3	$D_s^+ \rightarrow K_S^0 K^+ \pi^0$
110±15±11	LEES	21A BABR	$\eta_c(1S) \rightarrow \pi^+ \pi^- \eta$

NODE=M263W

NEW

¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

² Observed to decay into $K_S^0 K^+$ in a Breit-Wigner amplitude analysis involving D_s^+ decays into $\bar{K}^*(892)^0 K^+$, $\bar{K}^*(892)^+ K_S^0$, $\bar{K}^*(1410)^0 K^+$, $a_0(980)^+ \pi^0$, and $a_0(1817)^+ \pi^0$.

NODE=M263W;LINKAGE=B

NODE=M263W;LINKAGE=A

$a_0(1710)$ DECAY MODES

NODE=M263215;NODE=M263

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\eta$	seen
Γ_2 K^+K^-	
Γ_3 $K_S^0 K_S^0$	
Γ_4 $K_S^0 K^+$	seen

DESIG=1
DESIG=2
DESIG=3
DESIG=4

$\Gamma(\pi\eta)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A	BABR $\eta_c(1S) \rightarrow \pi^+ \pi^- \eta$

NODE=M263R01
NODE=M263R01

$\Gamma(K^+K^-)/\Gamma(K_S^0 K_S^0)$	Γ_2/Γ_3		
VALUE	DOCUMENT ID	TECN	COMMENT
0.32±0.12	¹ ABLIKIM	22F	BES3 $D_s^+ \rightarrow K_S^0 K_S^0 \pi^+$

NODE=M263R00
NODE=M263R00
OCCUR=2

¹ Using $D_s^+ \rightarrow K^+ K^- \pi^+$ from ABLIKIM 21AE. The apparent violation of isospin symmetry may be due to a destructive interference with the $f_0(1710)$ in the $K^+ K^-$ channel, and a constructive interference in the $K_S^0 K_S^0$ channel.

NODE=M263R00;LINKAGE=B

$\Gamma(K_S^0 K^+)/\Gamma_{\text{total}}$	Γ_4/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	22AH	BES3 $D_s^+ \rightarrow K_S^0 K^+ \pi^0$

NODE=M263R02
NODE=M263R02

 $a_0(1710)$ REFERENCES

AAJ	23AH PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	22AH PRL 129 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22F PR D105 L051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AE PR D104 012016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21A PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)

NODE=M263

REFID=62349
REFID=61880
REFID=61641
REFID=61367
REFID=61442

$f_0(1710)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M068

NODE=M068

 $f_0(1710)$ T-MATRIX POLE \sqrt{s}

Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M068PP

NODE=M068PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1680–1820) – i (50–180) OUR ESTIMATE			
$(1769 \pm 8) - i(78 \pm 6)$	¹ RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
$(1700 \pm 18) - i(127 \pm 12)$	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1803 \pm 3.5^{+45.5}_{-10.4}) - i(145 \pm 2.5^{+16.3}_{-9.6})$	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
$(1732 \pm 15) - i(160^{+25}_{-10})$	³ ANISOVICH	03	RVUE $\pi\pi, K\bar{K}, \eta\eta, \eta\eta', \pi\pi\pi\pi$
$(1698 \pm 18) - i(60 \pm 13)$	BARBERIS	00E	OMEG 450 $pp \rightarrow p_f \eta \eta p_S$
$(1770 \pm 12) - i(110 \pm 20)$	⁴ ANISOVICH	99B	SPEC 0.6–1.2 $p\bar{p} \rightarrow \eta\eta\pi^0$
$(1727 \pm 12 \pm 11) - i(63 \pm 8 \pm 9)$	BARBERIS	99D	OMEG 450 $pp \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1750 \pm 30) - i(125 \pm 70)$	ANISOVICH	98B	RVUE Compilation

NODE=M068PP

→ UNCHECKED ←

OCCUR=3

¹ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).

NODE=M068PP;LINKAGE=O

² T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$), and BINON 84C ($\eta\eta'$).

NODE=M068PP;LINKAGE=H

³ Solution I.

NODE=M068PP;LINKAGE=A

⁴ Not seen by AMSLER 02.

NODE=M068PP;LINKAGE=AV

$f_0(1710)$ Breit-Wigner MASS

NODE=M068M

NODE=M068M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1733^{+8}_{-7}	OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
1757 ± 24	± 9	LEES	21A	BABR $\eta_c(1S) \rightarrow \eta' K^+ K^-$
1759 ± 6	$^{+14}_{-25}$ 5.5k	1 ABLIKIM	13N	BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
1750^{+6}_{-7}	$^{+29}_{-18}$	2 UEHARA	13	BELL $\gamma \gamma \rightarrow K_S^0 K_S^0$
1701 ± 5	$^{+9}_{-2}$ 4k	3 CHEKANOV	08	ZEUS $e p \rightarrow K_S^0 K_S^0 X$
1765^{+4}_{-3}	± 13	ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
1738 ± 30		ABLIKIM	04E	BES2 $J/\psi \rightarrow \omega K^+ K^-$
1740 ± 4	$^{+10}_{-25}$	BAI	03G	BES $J/\psi \rightarrow \gamma K \bar{K}$
1740^{+30}_{-25}		BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
1710 ± 25		4 FRENCH	99	300 $p p \rightarrow p_f(K^+ K^-) p_s$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1744 ± 7	± 5 381	5,6 DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
1705 ± 11	± 5 237	5,6 DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
1706 ± 4	± 5 1.0k	5,6 DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
1690 ± 8	± 3 349	5,6 DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
1750 ± 13		AMSLER	06	CBAR $1.64 \bar{p} p \rightarrow K^+ K^- \pi^0$
1747 ± 5	80k	7 UMAN	06	E835 $5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
1776 ± 15		VLADIMIRSK...	06	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1670 ± 20		BINON	05	GAMS $33 \pi^- p \rightarrow \eta \eta n$
1682 ± 16		TIKHOMIROV	03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1670 ± 26	3.6k	8 NICHITIU	02	OBLX $0 \bar{p} p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1730 ± 15		BARBERIS	99	OMEG $450 p p \rightarrow p_s p_f K^+ K^-$
1750 ± 20		BARBERIS	99B	OMEG $450 p p \rightarrow p_s p_f \pi^+ \pi^-$
1720 ± 39		BAI	98H	BES $J/\psi \rightarrow \gamma \pi^0 \pi^0$
1775 ± 1.5	57	9 BARKOV	98	$\pi^- p \rightarrow K_S^0 K_S^0 n$
1690 ± 11		10 ABREU	96C	DLPH $Z^0 \rightarrow K^+ K^- + X$
1696 ± 5	$^{+9}_{-34}$	11 BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$
1781 ± 8	$^{+10}_{-31}$	BAI	96C	BES $J/\psi \rightarrow \gamma K^+ K^-$
1768 ± 14		BALOSHIN	95	SPEC $40 \pi^- C \rightarrow K_S^0 K_S^0 X$
1750 ± 15		12 BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1620 ± 16		11 BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
1748 ± 10		13 ARMSTRONG	93C	E760 $\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6 \gamma$
~ 1750		BREAKSTONE	93	SFM $p p \rightarrow p p \pi^+ \pi^- \pi^+ \pi^-$
1744 ± 15		14 ALDE	92D	GAM2 $38 \pi^- p \rightarrow \eta \eta n$
1713 ± 10		15 ARMSTRONG	89D	OMEG $300 p p \rightarrow p p K^+ K^-$
1706 ± 10		15 ARMSTRONG	89D	OMEG $300 p p \rightarrow p p K_S^0 K_S^0$
1707 ± 10		13 AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$
1700 ± 15		11 BOLONKIN	88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1720 ± 60		BOLONKIN	88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1638 ± 10		16 FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
1690 ± 4		17 FALVARD	88	DM2 $J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$
1698 ± 15		13 AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
1720 ± 10	± 10	11 BALTRUSAIT..	87	MRK3 $J/\psi \rightarrow \gamma K^+ K^-$
1755 ± 8		18 ALDE	86C	GAM2 $38 \pi^- p \rightarrow n 2 \eta$
1730^{+2}_{-10}		19 LONGACRE	86	RVUE $22 \pi^- p \rightarrow n 2 K_S^0$
1742 ± 15		13 WILLIAMS	84	MPSF $200 \pi^- N \rightarrow 2 K_S^0 X$
1670 ± 50		BLOOM	83	CBAL $J/\psi \rightarrow \gamma 2 \eta$
1650 ± 50		BURKE	82	MRK2 $J/\psi \rightarrow \gamma 2 \rho$
1640 ± 50		20,21 EDWARDS	82D	CBAL $J/\psi \rightarrow \gamma 2 \eta$
1730 ± 10	± 20	22 ETKIN	82C	MPS $23 \pi^- p \rightarrow n 2 K_S^0$

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2


OCCUR=2

- 1 From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.
- 2 Spin 0 favored over spin 2.
- 3 In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.
- 4 $J^P = 0^+$, supersedes ARMSTRONG 89D.
- 5 Using CLEO-c data but not authored by the CLEO Collaboration.
- 6 From a fit to a Breit-Wigner line shape with fixed $\Gamma = 135$ MeV.
- 7 Systematic errors not estimated.
- 8 Decaying to $f_0(1370)\pi\pi$.
- 9 No J^{PC} determination.
- 10 No J^{PC} determination, width not determined.
- 11 $J^P = 2^+$.
- 12 From a fit to the 0^+ partial wave.
- 13 No J^{PC} determination.
- 14 ALDE 92D combines all the GAMS-2000 data.
- 15 $J^P = 2^+$, superseded by FRENCH 99.
- 16 From an analysis ignoring interference with $f_2'(1525)$.
- 17 From an analysis including interference with $f_2'(1525)$.
- 18 Superseded by ALDE 92D.
- 19 Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.
- 20 $J^P = 2^+$ preferred.
- 21 From fit neglecting nearby $f_2'(1525)$. Replaced by BLOOM 83.
- 22 Superseded by LONGACRE 86.

NODE=M068M;LINKAGE=D
 NODE=M068M;LINKAGE=H
 NODE=M068M;LINKAGE=HE

 NODE=M068M;LINKAGE=C3
 NODE=M068M;LINKAGE=F
 NODE=M068M;LINKAGE=G
 NODE=M068M;LINKAGE=CH
 NODE=M068M;LINKAGE=NC
 NODE=M068M;LINKAGE=4A
 NODE=M068M;LINKAGE=A4
 NODE=M068M;LINKAGE=A3
 NODE=M068M;LINKAGE=Q0
 NODE=M068M;LINKAGE=A1
 NODE=M068M;LINKAGE=AA
 NODE=M068M;LINKAGE=C
 NODE=M068M;LINKAGE=A
 NODE=M068M;LINKAGE=B
 NODE=M068M;LINKAGE=BB
 NODE=M068M;LINKAGE=A9

 NODE=M068M;LINKAGE=B2
 NODE=M068M;LINKAGE=E
 NODE=M068M;LINKAGE=B1




$f_0(1710)$ Breit-Wigner WIDTH

NODE=M068W
 NODE=M068W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
150 \pm 12 $-$ 10	OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
175 \pm 23 \pm 4		LEES	21A BABR	$\eta_c(1S) \rightarrow \eta' K^+ K^-$
172 \pm 10 $\begin{smallmatrix} +32 \\ -16 \end{smallmatrix}$	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
139 \pm 11 $\begin{smallmatrix} +96 \\ -50 \end{smallmatrix}$		² UEHARA	13 BELL	$\gamma \gamma \rightarrow K_S^0 K_S^0$
100 \pm 24 $\begin{smallmatrix} +7 \\ -22 \end{smallmatrix}$	4k	³ CHEKANOV	08 ZEUS	$e p \rightarrow K_S^0 K_S^0 X$
145 \pm 8 \pm 69		ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
125 \pm 20		ABLIKIM	04E BES2	$J/\psi \rightarrow \omega K^+ K^-$
166 $\begin{smallmatrix} +5 \\ -8 \end{smallmatrix}$ $\begin{smallmatrix} +15 \\ -10 \end{smallmatrix}$		BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
120 $\begin{smallmatrix} +50 \\ -40 \end{smallmatrix}$		BAI	00A BES	$J/\psi \rightarrow \gamma (\pi^+ \pi^- \pi^+ \pi^-)$
105 \pm 34		⁴ FRENCH	99	300 pp $\rightarrow p_f(K^+ K^-) p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

148	$\begin{matrix} + 40 \\ - 30 \end{matrix}$		AMSLER	06	CBAR	1.64	$\bar{p}p \rightarrow K^+ K^- \pi^0$	
188	± 13	80k	⁵ UMAN	06	E835	5.2	$\bar{p}p \rightarrow \eta\eta\pi^0$	
250	± 30		VLADIMIRSK...	06	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
260	± 50		BINON	05	GAMS	33	$\pi^- p \rightarrow \eta\eta n$	
102	± 26		TIKHOMIROV	03	SPEC	40.0	$\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
267	± 44	3651	⁶ NICHITIU	02	OBLX	0	$\bar{p}p \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
100	± 25		BARBERIS	99	OMEG	450	$pp \rightarrow p_S p_f K^+ K^-$	
160	± 30		BARBERIS	99B	OMEG	450	$pp \rightarrow p_S p_f \pi^+ \pi^-$	
30	± 7	57	⁷ BARKOV	98			$\pi^- p \rightarrow K_S^0 K_S^0 n$	
103	± 18	$\begin{matrix} +30 \\ -11 \end{matrix}$	⁸ BAI	96C	BES		$J/\psi \rightarrow \gamma K^+ K^-$	
85	± 24	$\begin{matrix} +22 \\ -19 \end{matrix}$	BAI	96C	BES		$J/\psi \rightarrow \gamma K^+ K^-$	OCCUR=2
56	± 19		BALOSHIN	95	SPEC	40	$\pi^- C \rightarrow K_S^0 K_S^0 X$	
160	± 40		⁹ BUGG	95	MRK3		$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$	
160	$\begin{matrix} + 60 \\ - 20 \end{matrix}$		⁸ BUGG	95	MRK3		$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$	OCCUR=2
264	± 25		¹⁰ ARMSTRONG	93C	E760		$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	
200	to 300		BREAKSTONE	93	SFM		$pp \rightarrow pp \pi^+ \pi^- \pi^+ \pi^-$	
< 80	90% CL		¹¹ ALDE	92D	GAM2	38	$\pi^- p \rightarrow \eta\eta N^*$	
181	± 30		¹² ARMSTRONG	89D	OMEG	300	$pp \rightarrow pp K^+ K^-$	
104	± 30		¹² ARMSTRONG	89D	OMEG	300	$pp \rightarrow pp K_S^0 K_S^0$	OCCUR=2
166.4	± 33.2		¹⁰ AUGUSTIN	88	DM2		$J/\psi \rightarrow \gamma K^+ K^-, K_S^0 K_S^0$	
30	± 20		⁸ BOLONKIN	88	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
350	± 150		BOLONKIN	88	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	OCCUR=2
148	± 17		¹³ FALVARD	88	DM2		$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	
184	± 6		¹⁴ FALVARD	88	DM2		$J/\psi \rightarrow \phi K^+ K^-, K_S^0 K_S^0$	OCCUR=2
136	± 28		¹⁰ AUGUSTIN	87	DM2		$J/\psi \rightarrow \gamma \pi^+ \pi^-$	
130	± 20		⁸ BALTRUSAIT...	87	MRK3		$J/\psi \rightarrow \gamma K^+ K^-$	
122	$\begin{matrix} + 74 \\ - 15 \end{matrix}$		¹⁵ LONGACRE	86	RVUE	22	$\pi^- p \rightarrow n 2K_S^0$	
57	± 38		¹⁶ WILLIAMS	84	MPSF	200	$\pi^- N \rightarrow 2K_S^0 X$	
160	± 80		BLOOM	83	CBAL		$J/\psi \rightarrow \gamma 2\eta$	
200	± 100		BURKE	82	MRK2		$J/\psi \rightarrow \gamma 2\rho$	
220	$\begin{matrix} + 100 \\ - 70 \end{matrix}$		^{17,18} EDWARDS	82D	CBAL		$J/\psi \rightarrow \gamma 2\eta$	
200	$\begin{matrix} + 156 \\ - 9 \end{matrix}$		¹⁹ ETKIN	82B	MPS	23	$\pi^- p \rightarrow n 2K_S^0$	
			¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.					NODE=M068W;LINKAGE=F
			² Spin 0 favored over spin 2.					NODE=M068W;LINKAGE=G
			³ In the SU(3) based model with a specific interference pattern of the $f_2(1270)$, $a_2^0(1320)$, and $f_2'(1525)$ mesons incoherently added to the $f_0(1710)$ and non-resonant background.					NODE=M068W;LINKAGE=HE
			⁴ $J^P = 0^+$, supersedes ARMSTRONG 89D.					NODE=M068W;LINKAGE=C3
			⁵ Systematic errors not estimated.					NODE=M068W;LINKAGE=CH
			⁶ Decaying to $f_0(1370)\pi\pi$.					NODE=M068W;LINKAGE=NC
			⁷ No J^{PC} determination.					NODE=M068W;LINKAGE=4A
			⁸ $J^P = 2^+$.					NODE=M068W;LINKAGE=A3
			⁹ From a fit to the 0^+ partial wave.					NODE=M068W;LINKAGE=Q0
			¹⁰ No J^{PC} determination.					NODE=M068W;LINKAGE=A1
			¹¹ ALDE 92D combines all the GAMS-2000 data.					NODE=M068W;LINKAGE=AA
			¹² $J^P = 2^+$, (0^+ excluded).					NODE=M068W;LINKAGE=B
			¹³ From an analysis ignoring interference with $f_2'(1525)$.					NODE=M068W;LINKAGE=C
			¹⁴ From an analysis including interference with $f_2'(1525)$.					NODE=M068W;LINKAGE=D
			¹⁵ Uses MRK3 data. From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.					NODE=M068W;LINKAGE=A9
			¹⁶ No J^{PC} determination.					NODE=M068M;LINKAGE=WI
			¹⁷ $J^P = 2^+$ preferred.					NODE=M068W;LINKAGE=B2
			¹⁸ From fit neglecting nearby $f_2'(1525)$. Replaced by BLOOM 83.					NODE=M068W;LINKAGE=E
			¹⁹ From an amplitude analysis of the $K_S^0 K_S^0$ system, superseded by LONGACRE 86.					NODE=M068W;LINKAGE=A



$f_0(1710)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\eta\eta$	seen
Γ_3 $\eta\eta'$	
Γ_4 $\pi\pi$	seen
Γ_5 $\gamma\gamma$	seen
Γ_6 $\omega\omega$	seen

NODE=M068215;NODE=M068

DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=7
DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=6;OUR EST;→ UNCHECKED ←
DESIG=4

$f_0(1710)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_5/\Gamma$
$12^{+3}_{-2} + 227_8$		UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M068220

NODE=M068G2
NODE=M068G2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<480	95	ALBRECHT	90G	ARG	$\gamma\gamma \rightarrow K^+ K^-$
<110	95	¹ BEHREND	89C	CELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
<280	95	¹ ALTHOFF	85B	TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

¹ Assuming helicity 2.

NODE=M068G2;LINKAGE=F

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_4\Gamma_5/\Gamma$
<0.82	95	¹ BARATE	00E	ALEP $\gamma\gamma \rightarrow \pi^+ \pi^-$	

¹ Assuming spin 0.

NODE=M068G3
NODE=M068G3

NODE=M068G;LINKAGE=Z

$f_0(1710)$ BRANCHING RATIOS

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
-------	------	-------------	------	---------	-------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	1004	¹ DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$
seen	349	¹ DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$
0.36 ± 0.12		ALBALADEJO	08	RVUE	
$0.38^{+0.09}_{-0.19}$		² LONGACRE	86	MPS	$22 \pi^- p \rightarrow n 2K_S^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

NODE=M068225

NODE=M068R2
NODE=M068R2

OCCUR=2

NODE=M068R2;LINKAGE=A
NODE=M068R;LINKAGE=L

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.22±0.12	ALBALADEJO 08	RVUE
0.18 ^{+0.03} _{-0.13}	¹ LONGACRE 86	RVUE

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

 Γ_2/Γ

NODE=M068R1
NODE=M068R1

NODE=M068R1;LINKAGE=L

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	381	¹ DOBBS 15		$J/\psi \rightarrow \gamma\pi^+\pi^-$
seen	237	¹ DOBBS 15		$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
not seen		AMSLER 02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
0.039 ^{+0.002} _{-0.024}		² LONGACRE 86	RVUE	

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2. Fit with constrained inelasticity.

 Γ_4/Γ

NODE=M068R5
NODE=M068R5

OCCUR=2

NODE=M068R5;LINKAGE=A
NODE=M068R5;LINKAGE=L

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.23±0.05 OUR AVERAGE Error includes scale factor of 1.2.

0.64±0.27 ±0.18		LEES 18A	BABR	$\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.41 ^{+0.11} _{-0.17}		ABLIKIM 06V	BES2	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
0.2 ±0.024±0.036		BARBERIS 99D	OMEG 450	$p\bar{p} \rightarrow K^+K^-, \pi^+\pi^-$
0.39±0.14		ARMSTRONG 91	OMEG 300	$p\bar{p} \rightarrow p\bar{p}\pi\pi, p\bar{p}K\bar{K}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.32±0.14		ALBALADEJO 08	RVUE	
< 0.11	95	¹ ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+K^-$
5.8 ^{+9.1} _{-5.5}		² ANISOVICH 02D	SPEC	Combined fit

¹ Using data from ABLIKIM 04A.

² From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

 Γ_4/Γ_1

NODE=M068R6
NODE=M068R6

NODE=M068R;LINKAGE=AB
NODE=M068R;LINKAGE=CH

 $\Gamma(\eta\eta)/\Gamma(K\bar{K})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

0.48±0.15 BARBERIS 00E 450 $p\bar{p} \rightarrow p_f\eta\eta p_s$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.46 ^{+0.70} _{-0.38}		¹ ANISOVICH 02D	SPEC	Combined fit
<0.02	90	² PROKOSHKIN 91	GA24	$300 \pi^- p \rightarrow \pi^- p\eta\eta$

¹ From a combined K-matrix analysis of Crystal Barrel ($0. p\bar{p} \rightarrow \pi^0\pi^0\pi^0, \pi^0\eta\eta, \pi^0\pi^0\eta$), GAMS ($\pi p \rightarrow \pi^0\pi^0 n, \eta\eta n, \eta\eta' n$), and BNL ($\pi p \rightarrow K\bar{K} n$) data.

² Combining results of GAM4 with those of ARMSTRONG 89D.

 Γ_2/Γ_1

NODE=M068R7
NODE=M068R7

NODE=M068R7;LINKAGE=CH

NODE=M068R;LINKAGE=A

 $\Gamma(\eta\eta')/\Gamma(\pi\pi)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.87 × 10⁻³ 90 ¹ ABLIKIM 22AS BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.

 Γ_3/Γ_4

NODE=M068R00
NODE=M068R00

NODE=M068R00;LINKAGE=A

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 180 ABLIKIM 06H BES $J/\psi \rightarrow \gamma\omega\omega$

 Γ_6/Γ

NODE=M068R3
NODE=M068R3

f₀(1710) REFERENCES

NODE=M068

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
Also		PR D107 079901 (err.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62033
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61442
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58950
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ALBALADEJO	08	PRL 101 252002	M. Albaladejo, J.A. Oller		REFID=52656
CHEKANOV	08	PRL 101 112003	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52275
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
		Translated from YAF 68 998.			
ABLIKIM	04A	PL B598 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49740
ABLIKIM	04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>		REFID=48831
		Translated from YAF 65 1583.			
NICHITIU	02	PL B545 261	F. Nichitiu <i>et al.</i>	(OBELIX Collab.)	REFID=48848
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
BARATE	00E	PL B472 189	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47428
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>		REFID=46886
BARBERIS	99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46922
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=47395
FRENCH	99	PL B460 213	B. French <i>et al.</i>	(WA76 Collab.)	REFID=47491
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>		REFID=46331
		Translated from UFN 168 481.			
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BARKOV	98	JETPL 68 764	B.P. Barkov <i>et al.</i>		REFID=46616
ABREU	96C	PL B379 309	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44671
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BALOSHIN	95	PAN 58 46	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=44621
		Translated from YAF 58 50.			
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ARMSTRONG	93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
BREAKSTONE	93	ZPHY C58 251	A.M. Breakstone <i>et al.</i>	(IOWA, CERN, DORT+)	REFID=43312
ALDE	92D	PL B284 457	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=41591
Also		SJNP 54 451	D.M. Alde <i>et al.</i>	(GAM2 Collab.)	REFID=44696
		Translated from YAF 54 745.			
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)	REFID=41744
PROKOSHKIN	91	SPD 36 155	Y.D. Prokoshkin	(GAM2 and GAM4 Collab.)	REFID=41719
		Translated from DANS 316 900.			
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41374
ARMSTRONG	89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41010
BEHREND	89C	ZPHY C43 91	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40915
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
ALDE	86C	PL B182 105	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=21694
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=21418
WILLIAMS	84	PR D30 877	E.G.H. Williams <i>et al.</i>	(VAND, NDAM, TUFTS+)	REFID=21693
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
ETKIN	82C	PR D25 2446	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20391
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355

X(1750)

$$I^G(J^{PC}) = ?^-(1^{--})$$

OMITTED FROM SUMMARY TABLE

The X(1750) was separated from the $\phi(1680)$ in the 2022 listings due to its incompatible mass and incompatible pattern of $\bar{K}K$ and $\bar{K}^*(892)K$ branching fractions.

NODE=M255

NODE=M255

X(1750) MASS

NODE=M255M

NODE=M255M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1753.8 ± 2.7 OUR AVERAGE			
1784 ± 12 $\begin{smallmatrix} +0 \\ -27 \end{smallmatrix}$	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
1753.5 ± 1.5 ± 2.3	LINK	02K FOCS	20-160 $\gamma p \rightarrow K^+ K^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1726 ± 22	BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
1760 ± 20	ATKINSON	85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$
1690 ± 10	ASTON	81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$

X(1750) WIDTH

NODE=M255W

NODE=M255W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120 ± 10 OUR AVERAGE			
106 $\begin{smallmatrix} +22 \\ -19 \end{smallmatrix}$ $\begin{smallmatrix} +8 \\ -36 \end{smallmatrix}$	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
122.2 ± 6.2 ± 8.0	LINK	02K FOCS	20-160 $\gamma p \rightarrow K^+ K^- p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
121 ± 47	BUSENITZ	89 TPS	$\gamma p \rightarrow K^+ K^- X$
80 ± 40	ATKINSON	85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$
100 ± 40	ASTON	81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$

X(1750) DECAY MODES

NODE=M255215;NODE=M255

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^+ K^-$	seen
Γ_2 $\bar{K}^*(892)^0 K_S^0$	not seen
Γ_3 $K^*(892)^\pm K^\mp$	not seen
Γ_4 $\eta \phi$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4

$$\Gamma(\bar{K}^*(892)^0 K_S^0)/\Gamma(K^+ K^-) \quad \Gamma_2/\Gamma_1$$

NODE=M255R01
NODE=M255R01

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.065	90	LINK	02K FOCS	$\gamma p \rightarrow K^+ K^- p$

$$\Gamma(K^*(892)^\pm K^\mp)/\Gamma(K^+ K^-) \quad \Gamma_3/\Gamma_1$$

NODE=M255R02
NODE=M255R02

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.183	90	LINK	02K FOCS	$\gamma p \rightarrow K^+ K^- p$

$$\Gamma(\eta \phi)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

NODE=M255R00
NODE=M255R00

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ZHU	23A RVUE	$e^+ e^- \rightarrow \eta \phi$

¹ Reported with a 2 σ significance in the fit and an upper limit of $\Gamma(e^+ e^-) B(X(1750) \rightarrow \eta \phi)$ in the range 136–322 eV.

NODE=M255R00;LINKAGE=A

X(1750) REFERENCES

NODE=M255

ZHU	23A	CP C47 113003	W. Zhu, X. Wang	(RVUE)	REFID=62439
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
LINK	02K	PL B545 50	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=48845
BUSENITZ	89	PR D40 1	J.K. Busenitz <i>et al.</i>	(ILL, FNAL)	REFID=40927
ATKINSON	85C	ZPHY C27 233	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=21596
ASTON	81F	PL 104B 231	D. Aston	(BONN, CERN, EPOL, GLAS, LANC+)	REFID=21585

$\eta(1760)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

NODE=M114

OMITTED FROM SUMMARY TABLE

Seen by DM2 in the $\rho\rho$ system (BISELLO 89B). Structure in this region has been reported before in the same system (BALTRUSAITIS 86B) and in the $\omega\omega$ system (BALTRUSAITIS 85C, BISELLO 87).

NODE=M114

 $\eta(1760)$ MASS

NODE=M114M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1751±15 OUR AVERAGE				
$1768^{+24}_{-25} \pm 10$	465	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$1744 \pm 10 \pm 15$	1045	² ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
••• We do not use the following data for averages, fits, limits, etc. •••				
$1703^{+12}_{-11} \pm 2$		³ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
1760 ± 11	320	⁴ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

NODE=M114M

OCCUR=2

¹ From a single-resonance fit.² From a partial wave analysis including $\eta(1760)$, $f_0(1710)$, $f_2(1640)$, and $f_2(1910)$.³ From a two-resonance fit.⁴ Estimated by us from various fits. Systematic uncertainties not estimated.

NODE=M114M;LINKAGE=ZA

NODE=M114M;LINKAGE=MA

NODE=M114M;LINKAGE=ZH

NODE=M114M;LINKAGE=A

 $\eta(1760)$ WIDTH

NODE=M114W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
240±30 OUR AVERAGE				
$224^{+62}_{-56} \pm 25$	465	⁵ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$244^{+24}_{-21} \pm 25$	1045	⁶ ABLIKIM	06H BES	$J/\psi \rightarrow \gamma\omega\omega$
••• We do not use the following data for averages, fits, limits, etc. •••				
$42^{+36}_{-22} \pm 15$		⁷ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
60 ± 16	320	⁸ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

NODE=M114W

OCCUR=2

⁵ From a single-resonance fit.⁶ From a partial wave analysis including $\eta(1760)$, $f_0(1710)$, $f_2(1640)$, and $f_2(1910)$.⁷ From a two-resonance fit.⁸ Estimated by us from various fits. Systematic uncertainties not estimated.

NODE=M114W;LINKAGE=ZA

NODE=M114W;LINKAGE=MA

NODE=M114W;LINKAGE=ZH

NODE=M114W;LINKAGE=B

 $\eta(1760)$ DECAY MODES

NODE=M114215;NODE=M114

Mode	Fraction (Γ_i/Γ)
Γ_1 4π	
Γ_2 $2\pi^+2\pi^-$	seen
Γ_3 $\pi^+\pi^-2\pi^0$	seen
Γ_4 $\rho^0\rho^0$	seen
Γ_5 $\rho^+\rho^-$	seen
Γ_6 $\omega\omega$	seen
Γ_7 $\eta'\pi^+\pi^-$	seen
Γ_8 $\gamma\gamma$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=7

DESIG=8;OUR EVAL;→ UNCHECKED ←

DESIG=9;OUR EVAL;→ UNCHECKED ←

 $\eta(1760)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M114225

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				$\Gamma_7\Gamma_8/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$28.2^{+7.9}_{-7.5} \pm 3.7$	465	⁹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••				
$3.0^{+2.0}_{-1.2} \pm 0.8$	52	¹⁰ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
$18^{+13}_{-10} \pm 5$	315	¹¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$

NODE=M114G01

NODE=M114G01

OCCUR=2

OCCUR=3

⁹ From a single-resonance fit.¹⁰ From a two-resonance fit. For constructive interference with the X(1835).¹¹ From a two-resonance fit. For destructive interference with the X(1835).

NODE=M114G01;LINKAGE=ZH

NODE=M114G01;LINKAGE=ZA

NODE=M114G01;LINKAGE=ZN

$\eta(1760)$ BRANCHING RATIOS

$\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$		Γ_2/Γ			NODE=M114210
VALUE		DOCUMENT ID	TECN	COMMENT	NODE=M114R01 NODE=M114R01
seen		BISELLO	89B DM2	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$	
$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma_{\text{total}}$		Γ_3/Γ			NODE=M114R02 NODE=M114R02
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		BISELLO	89B DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^- 2\pi^0$	
$\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$		Γ_4/Γ			NODE=M114R03 NODE=M114R03
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		BISELLO	89B DM2	$J/\psi \rightarrow \gamma \rho^0 \rho^0$	
seen		BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \gamma \rho^0 \rho^0$	
$\Gamma(\rho^+ \rho^-)/\Gamma_{\text{total}}$		Γ_5/Γ			NODE=M114R04 NODE=M114R04
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		BISELLO	89B DM2	$J/\psi \rightarrow \gamma \rho^+ \rho^-$	
seen		BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \gamma \rho^+ \rho^-$	
$\Gamma(\omega\omega)/\Gamma_{\text{total}}$		Γ_6/Γ			NODE=M114R06 NODE=M114R06
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		BISELLO	87 DM2	$J/\psi \rightarrow \omega\omega$	
seen		BALTRUSAIT...85C	MRK3	$J/\psi \rightarrow \gamma\omega\omega$	
$\Gamma(\gamma\gamma)/\Gamma(\omega\omega)$		Γ_8/Γ_6			NODE=M114R08 NODE=M114R08
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.48 \times 10^{-3}$	90	¹² ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$	
¹² Using results from ABLIKIM 06H.					NODE=M114R08;LINKAGE=A

 $\eta(1760)$ REFERENCES

ABLIKIM	180	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)	REFID=54763
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
BALTRUSAIT... 86	PR D33 629		R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BALTRUSAIT... 86B	PR D33 1222		R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT... 85C	PRL 55 1723		R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095

NODE=M264

 $f_0(1770)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M264

 $f_0(1770)$ Breit-Wigner MASS

NODE=M264M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	NODE=M264M
1784^{+16}_{-14}	OUR AVERAGE	Error includes scale factor of 1.1.			
1814 ± 31	7.2k	¹ KHOLODENK..21	VES	$29 \pi^- p \rightarrow n\omega\phi$	
$1795 \pm 7^{+23}_{-20}$		ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma\omega\phi$	
$1760 \pm 15^{+15}_{-10}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1765 ± 15		SARANTSEV	21 RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	
1814 ± 18	2,3	AAIJ	14BR LHCB	$\bar{B}_s^0 \rightarrow J/\psi \pi^+ \pi^-$	
$1812^{+19}_{-26} \pm 18$	4	ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma\omega\phi$	
1790^{+40}_{-30}		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$	

- ¹ From partial wave analysis of $\omega\phi$ invariant mass including 0^{++} , 2^{++} , and 0^{-+} resonances.
- ² Second solution: 1800 ± 22 MeV. The fit favors $f_0(1770)$ to $f_0(1710)$.
- ³ Statistical error only.
- ⁴ Not seen by LIU 09 in $B^\pm \rightarrow K^\pm \omega\phi$.

NODE=M264M;LINKAGE=M

NODE=M264M;LINKAGE=A
 NODE=M264M;LINKAGE=B
 NODE=M264M;LINKAGE=L

$f_0(1770)$ Breit-Wigner WIDTH

NODE=M264W


NODE=M264W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
161±21 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
182±19	7.2k	¹ KHOLODENK..21	VES	29 $\pi^- p \rightarrow n\omega\phi$
95±10 ⁺⁷⁸ ₋₈₂		ABLIKIM	13J BES3	$J/\psi \rightarrow \gamma\omega\phi$
125±25 ⁺¹⁰ ₋₁₅		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
180±20		SARANTSEV	21 RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
328±34		^{2,3} AAIJ	14BR LHCB	$\bar{B}_s^0 \rightarrow J/\psi\pi^+\pi^-$
105±20±28		⁴ ABLIKIM	06J BES2	$J/\psi \rightarrow \gamma\omega\phi$
270 ⁺⁶⁰ ₋₃₀		⁵ ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$

- ¹ From partial wave analysis of $\omega\phi$ invariant mass including 0^{++} , 2^{++} , and 0^{-+} resonances.
- ² Second solution: 263 ± 30 MeV. The fit favors $f_0(1770)$ to $f_0(1710)$.
- ³ Statistical error only.
- ⁴ Not seen by LIU 09 in $B^\pm \rightarrow K^\pm \omega\phi$.
- ⁵ $f_0(1710)$ width fixed to PDG value.

NODE=M264W;LINKAGE=J

NODE=M264W;LINKAGE=A
 NODE=M264W;LINKAGE=B
 NODE=M264W;LINKAGE=I
 NODE=M264W;LINKAGE=AB



$f_0(1770)$ DECAY MODES

NODE=M264215;NODE=M264

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 $K\bar{K}$	seen
Γ_3 $\eta\eta$	seen
Γ_4 $\omega\phi$	seen

DESIG=1
 DESIG=2
 DESIG=3
 DESIG=4

$\Gamma(\pi\pi)/\Gamma_{total}$

Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	SARANTSEV	21 RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
seen	AAIJ	14BR LHCB	$\bar{B}_s^0 \rightarrow J/\psi\pi^+\pi^-$
seen	ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$

NODE=M264R01
 NODE=M264R01

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

NODE=M264R02
 NODE=M264R02

 $\Gamma(\eta\eta)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen	SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

NODE=M264R03
 NODE=M264R03

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
seen	7.2k	KHOLODENK..21	VES	$29 \pi^- p \rightarrow n\omega\phi$	
seen		SARANTSEV 21	RVUE	$J/\psi \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$	

NODE=M264R04
 NODE=M264R04

 $f_0(1770)$ REFERENCES

KHOLODENK...21	PAN 83 1602	M.S. Kholodenko	(VES Collab.)	REFID=61410
SARANTSEV 21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
AAIJ 14BR	PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56035
ABLIKIM 13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54955
LIU 09	PR D79 071102	C. Liu <i>et al.</i>	(BELLE Collab.)	REFID=52752
ABLIKIM 06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM 05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM 05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958

NODE=M264

$\pi(1800)$

$$J^{PC} = 1^-(0^-+)$$

See the review on "Non- $q\bar{q}$ Mesons."

NODE=M075

NODE=M075

 $\pi(1800)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M075M

NODE=M075M

1810^{+9}_{-11} OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.

1804^{+6}_{-9}	46M	1 AGHASYAN	18B	COMP	$190 \pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$1876 \pm 18 \pm 16$	4k	2 EUGENIO	08	B852	$18 \pi^- p \rightarrow \eta\eta\pi^- p$
$1774 \pm 18 \pm 20$		3 CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
$1863 \pm 9 \pm 10$		4 CHUNG	02	B852	$18.3 \pi^- p \rightarrow \pi^+ \pi^- \pi^- p$
$1840 \pm 10 \pm 10$	1.2k	AMELIN	96B	VES	$37 \pi^- A \rightarrow \eta\eta\pi^- A$
$1775 \pm 7 \pm 10$		5 AMELIN	95B	VES	$36 \pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
1790 ± 14		6 BERDNIKOV	94	VES	$37 \pi^- A \rightarrow K^+ K^- \pi^- A$
$1873 \pm 33 \pm 20$		BELADIDZE	92C	VES	$36 \pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
$1814 \pm 10 \pm 23$	426	BITYUKOV	91	VES	$36 \pi^- C \rightarrow \pi^- \eta\eta C$
1770 ± 30	1.1k	BELLINI	82	SPEC	$40 \pi^- A \rightarrow 3\pi A$

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

$1785 \pm 9^{+12}_{-6}$	420k	7 ALEKSEEV	10	COMP	$190 \pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
$1737 \pm 5 \pm 15$		AMELIN	99	VES	$37 \pi^- A \rightarrow \omega \pi^- \pi^0 A^*$

¹ Statistical error negligible.

² From a single-pole fit.

³ In the $f_0(980)\pi$ wave.


⁴ In the $f_0(500)\pi$ wave.

⁵ From a fit to $J^{PC} = 0^-+ f_0(980)\pi, f_0(1370)\pi$ waves.

⁶ From a fit to $J^{PC} = 0^-+ K_0^*(1430)K^-$ and $f_0(980)\pi^-$ waves.

⁷ Superseded by AGHASYAN 2018B.

NODE=M075M;LINKAGE=B
 NODE=M075M;LINKAGE=SP
 NODE=M075M;LINKAGE=C1
 NODE=M075M;LINKAGE=C2
 NODE=M075M;LINKAGE=AX
 NODE=M075M;LINKAGE=A
 NODE=M075M;LINKAGE=C



π(1800) WIDTH

NODE=M075W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
215⁺⁷₋₈ OUR AVERAGE					
220 ⁺⁸ ₋₁₁	46M	⁸ AGHASYAN	18B	COMP	190 π ⁻ p → π ⁻ π ⁺ π ⁻ p
221 ± 26 ± 38	4k	⁹ EUGENIO	08	B852	- 18 π ⁻ p → η η π ⁻ p
223 ± 48 ± 50		¹⁰ CHUNG	02	B852	18.3 π ⁻ p → π ⁺ π ⁻ π ⁻ p
191 ± 21 ± 20		¹¹ CHUNG	02	B852	18.3 π ⁻ p → π ⁺ π ⁻ π ⁻ p
210 ± 30 ± 30	1.2k	AMELIN	96B	VES	- 37 π ⁻ A → η η π ⁻ A
190 ± 15 ± 15		¹² AMELIN	95B	VES	- 36 π ⁻ A → π ⁺ π ⁻ π ⁻ A
210 ± 70		¹³ BERDNIKOV	94	VES	- 37 π ⁻ A → K ⁺ K ⁻ π ⁻ A
225 ± 35 ± 20		BELADIDZE	92C	VES	- 36 π ⁻ Be → π ⁻ η' η Be
205 ± 18 ± 32	426	BITYUKOV	91	VES	- 36 π ⁻ C → π ⁻ η η C
310 ± 50	1.1k	BELLINI	82	SPEC	- 40 π ⁻ A → 3π A
208 ± 22 ⁺²¹ ₋₃₇	420k	¹⁴ ALEKSEEV	10	COMP	190 π ⁻ Pb → π ⁻ π ⁻ π ⁺ Pb'
259 ± 19 ± 6		AMELIN	99	VES	37 π ⁻ A → ω π ⁻ π ⁰ A*

NODE=M075W

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

- ⁸Statistical error negligible.
- ⁹From a single-pole fit.
- ¹⁰In the $f_0(980)\pi$ wave.
- ¹¹In the $f_0(500)\pi$ wave.
- ¹²From a fit to $J^{PC} = 0^{-+} f_0(980)\pi, f_0(1370)\pi$ waves.
- ¹³From a fit to $J^{PC} = 0^{-+} K_0^*(1430)K^-$ and $f_0(980)\pi^-$ waves.
- ¹⁴Superseded by AGHASYAN 2018B.

NODE=M075W;LINKAGE=B
 NODE=M075W;LINKAGE=SP
 NODE=M075W;LINKAGE=C1
 NODE=M075W;LINKAGE=C2
 NODE=M075W;LINKAGE=AX
 NODE=M075W;LINKAGE=A
 NODE=M075W;LINKAGE=C

π(1800) DECAY MODES

NODE=M075215;NODE=M075

Mode	Fraction (Γ_i/Γ)
Γ_1 π ⁺ π ⁻ π ⁻	seen
Γ_2 $f_0(500)\pi^-$	seen
Γ_3 $f_0(980)\pi^-$	seen
Γ_4 $f_0(1370)\pi^-$	seen
Γ_5 $f_0(1500)\pi^-$	not seen
Γ_6 ρ π ⁻	not seen
Γ_7 η η π ⁻	seen

DESIG=10;OUR EST;→ UNCHECKED ←
 DESIG=11;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=1
 DESIG=12
 DESIG=2
 DESIG=7;OUR EST;→ UNCHECKED ←

Γ_8	$a_0(980)\eta$	seen	DESIG=5;OUR EST;→ UNCHECKED ←
Γ_9	$a_2(1320)\eta$	not seen	DESIG=13
Γ_{10}	$f_2(1270)\pi$	not seen	DESIG=14
Γ_{11}	$f_0(1370)\pi^-$	not seen	DESIG=15
Γ_{12}	$f_0(1500)\pi^-$	seen	DESIG=6;OUR EST;→ UNCHECKED ←
Γ_{13}	$\eta\eta'(958)\pi^-$	seen	DESIG=8;OUR EST;→ UNCHECKED ←
Γ_{14}	$K_0^*(1430)K^-$	seen	DESIG=4
Γ_{15}	$K^*(892)K^-$	not seen	DESIG=9

$\pi(1800)$ BRANCHING RATIOS

$\Gamma(f_0(980)\pi^-)/\Gamma(f_0(500)\pi^-)$					Γ_3/Γ_2	NODE=M075220
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		NODE=M075R11 NODE=M075R11
0.44±0.08±0.38	15 CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$		

$\Gamma(f_0(980)\pi^-)/\Gamma(f_0(1370)\pi^-)$					Γ_3/Γ_4	NODE=M075R5 NODE=M075R5
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.7±1.3	16 AMELIN	95B	VES	-	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$	
---------	-----------	-----	-----	---	--	--

$\Gamma(f_0(1370)\pi^-)/\Gamma_{total}$					Γ_4/Γ	NODE=M075R1 NODE=M075R1
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
seen	BELLINI	82	SPEC	-	40 $\pi^- A \rightarrow 3\pi A$	

$\Gamma(f_0(1500)\pi^-)/\Gamma_{total}$					Γ_5/Γ	NODE=M075R12 NODE=M075R12
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen	CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$		

$\Gamma(\rho\pi^-)/\Gamma_{total}$					Γ_6/Γ	NODE=M075R2 NODE=M075R2
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen	BELLINI	82	SPEC	-	40 $\pi^- A \rightarrow 3\pi A$	

$\Gamma(\rho\pi^-)/\Gamma(f_0(980)\pi^-)$					Γ_6/Γ_3	NODE=M075R6 NODE=M075R6
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.25		CHUNG	02	B852	18.3 $\pi^- p \rightarrow \pi^+ \pi^- \pi^- p$	
<0.14	90	AMELIN	95B	VES	-	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$

$\Gamma(\eta\eta\pi^-)/\Gamma(\pi^+ \pi^- \pi^-)$					Γ_7/Γ_1	NODE=M075R8 NODE=M075R8
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.5±0.1	1200	16 AMELIN	96B	VES	-	37 $\pi^- A \rightarrow \eta\eta\pi^- A$
---------	------	-----------	-----	-----	---	--

$\Gamma(a_2(1320)\eta)/\Gamma_{total}$					Γ_9/Γ	NODE=M075R13 NODE=M075R13
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen	EUGENIO	08	B852	18 $\pi^- p \rightarrow \eta\eta\pi^- p$		

$\Gamma(f_2(1270)\pi)/\Gamma_{total}$					Γ_{10}/Γ	NODE=M075R14 NODE=M075R14
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen	EUGENIO	08	B852	18 $\pi^- p \rightarrow \eta\eta\pi^- p$		

$\Gamma(f_0(1370)\pi^-)/\Gamma_{total}$					Γ_{11}/Γ	NODE=M075R15 NODE=M075R15
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>		
not seen	EUGENIO	08	B852	18 $\pi^- p \rightarrow \eta\eta\pi^- p$		

$\Gamma(f_0(1500)\pi^-)/\Gamma(a_0(980)\eta)$					Γ_{12}/Γ_8	NODE=M075R7 NODE=M075R7
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.48 ±0.17	4k	16,17 EUGENIO	08	B852	-	18 $\pi^- p \rightarrow \eta\eta\pi^- p$
0.030 ^{+0.014} _{-0.011}		16 ANISOVICH	01B	SPEC	0	0.6-1.94 $p\bar{p} \rightarrow \eta\eta\pi^0\pi^0$
0.08 ±0.03	1200	16,18 AMELIN	96B	VES	-	37 $\pi^- A \rightarrow \eta\eta\pi^- A$

$\Gamma(\eta\eta'(958)\pi^-)/\Gamma(\eta\eta\pi^-)$ Γ_{13}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29 ± 0.07		¹⁶ BELADIDZE	92C	VES	−	36 π^- Be → $\pi^- \eta' \eta$ Be
0.3 ± 0.1	426 ± 57	¹⁶ BITYUKOV	91	VES	−	36 π^- C → $\pi^- \eta \eta$ C

NODE=M075R10
NODE=M075R10 $\Gamma(K_0^*(1430)K^-)/\Gamma_{total}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

seen	BERDNIKOV	94	VES	−	37 π^- A → $K^+ K^- \pi^-$ A
------	-----------	----	-----	---	----------------------------------

NODE=M075R4
NODE=M075R4 $\Gamma(K^*(892)K^-)/\Gamma_{total}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

not seen	BERDNIKOV	94	VES	−	37 π^- A → $K^+ K^- \pi^-$ A
----------	-----------	----	-----	---	----------------------------------

NODE=M075R9
NODE=M075R9¹⁵ Assuming that $f_0(980)$ decays only to $\pi\pi$.¹⁶ Systematic errors not estimated.¹⁷ From a single-pole fit.¹⁸ Assuming that $f_0(1500)$ decays only to $\eta\eta$ and $a_0(980)$ decays only to $\eta\pi$.NODE=M075R;LINKAGE=CK
NODE=M075R5;LINKAGE=NS
NODE=M075R7;LINKAGE=SP
NODE=M075R7;LINKAGE=A $\pi(1800)$ REFERENCES

NODE=M075

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)	REFID=52160
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>		REFID=48318
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
		Translated from YAF 62 487.			
AMELIN	96B	PAN 59 976	D.V. Amelin <i>et al.</i>	(SERP, TBIL) IGJPC	REFID=44725
		Translated from YAF 59 1021.			
AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)	REFID=44433
BERDNIKOV	94	PL B337 219	E.B. Berdnikov <i>et al.</i>	(SERP, TBIL)	REFID=44073
BELADIDZE	92C	SJNP 55 1535	G.M. Beladidze, S.I. Bityukov, G.V. Borisov	(SERP+)	REFID=43175
		Translated from YAF 55 2748.			
BITYUKOV	91	PL B268 137	S.I. Bityukov <i>et al.</i>	(SERP, TBIL)	REFID=41749
BELLINI	82	PRL 48 1697	G. Bellini <i>et al.</i>	(MILA, BGNA, JINR)	REFID=21134

NODE=M038

 $f_2(1810)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M038

 $f_2(1810)$ MASS

NODE=M038M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1815 ± 12	OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.			
------------------	--------------------	---	--	--	--

1822	+29 −24	+66 −57	5.5k	¹ ABLIKIM	13N	BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
------	------------	------------	------	----------------------	-----	------	---

1737	± 9	+198 −65		² UEHARA	10A	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta \eta$
------	-----	-------------	--	---------------------	-----	------	--

1800	± 30		40	ALDE	88D	GAM4	$300 \pi^- p \rightarrow \pi^- p 4\pi^0$
------	------	--	----	------	-----	------	--

1806	± 10		1600	ALDE	87	GAM4	$100 \pi^- p \rightarrow 4\pi^0 n$
------	------	--	------	------	----	------	------------------------------------

1870	± 40			³ ALDE	86D	GAM4	$100 \pi^- p \rightarrow \eta \eta n$
------	------	--	--	-------------------	-----	------	---------------------------------------

1857	+35 −24			⁴ COSTA	80	OMEG	$10 \pi^- p \rightarrow K^+ K^- n$
------	------------	--	--	--------------------	----	------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1845.0 ± 2.2 ⁺ _−	1.6 7.2			⁵ ALBRECHT	20	RVUE	$0.9 \bar{p} p \rightarrow \pi^0 \pi^0 \eta,$ $\pi^0 \eta \eta, \pi^0 K^+ K^-$
--	------------	--	--	-----------------------	----	------	---

NODE=M038M

NODE=M038M

1858	+18 −71			⁶ LONGACRE	86	RVUE	Compilation
------	------------	--	--	-----------------------	----	------	-------------

NODE=M038M;LINKAGE=B

1799	± 15			⁷ CASON	82	STRC	$8 \pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
------	------	--	--	--------------------	----	------	---

NODE=M038M;LINKAGE=UE

NODE=M038M;LINKAGE=F

NODE=M038M;LINKAGE=A


NODE=M038M;LINKAGE=C

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.² Breit-Wigner mass. Could also be the $f_2(1910)$.³ Seen in only one solution.⁴ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.⁵ T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).⁶ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038M;LINKAGE=L

⁷ From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

NODE=M038M;LINKAGE=P1



$f_2(1810)$ WIDTH

NODE=M038W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
197 ± 22	OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
229 +52 -42 +88 -155	5.5k	1 ABLIKIM	13N BES3	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\eta\eta$
228 +21 -20 +234 -153		2 UEHARA	10A BELL	$10.6 e^+e^- \rightarrow e^+e^-\eta\eta$
160 ± 30	40	ALDE	88D GAM4	$300 \pi^-p \rightarrow \pi^-p4\pi^0$
190 ± 20	1600	ALDE	87 GAM4	$100 \pi^-p \rightarrow 4\pi^0n$
250 ± 30		3 ALDE	86D GAM4	$100 \pi^-p \rightarrow \eta\eta n$
185 +102 -139		4 COSTA	80 OMEG	$10 \pi^-p \rightarrow K^+K^-n$

NODE=M038W

• • • We do not use the following data for averages, fits, limits, etc. • • •

260.9 ± 3.9 +199.9 -38.2		5 ALBRECHT	20 RVUE	$0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta, \pi^0K^+K^-$
388 +15 -21		6 LONGACRE	86 RVUE	Compilation
280 +42 -35		7 CASON	82 STRC	$8 \pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$

NODE=M038W;LINKAGE=B

¹ From partial wave analysis including all possible combinations of $0^{++}, 2^{++},$ and 4^{++} resonances.

² Breit-Wigner width. Could also be the $f_2(1910)$.

³ Seen in only one solution.

⁴ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

⁵ T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).

⁶ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

⁷ From an amplitude analysis of the reaction $\pi^+\pi^- \rightarrow 2\pi^0$. The resonance in the $2\pi^0$ final state is not confirmed by PROKOSHKIN 97.

NODE=M038W;LINKAGE=UE


NODE=M038W;LINKAGE=F

NODE=M038W;LINKAGE=A

NODE=M038W;LINKAGE=C

NODE=M038W;LINKAGE=L

NODE=M038W;LINKAGE=P1



$f_2(1810)$ DECAY MODES

NODE=M038215;NODE=M038

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen
Γ_2 $\eta\eta$	seen
Γ_3 $4\pi^0$	seen
Γ_4 K^+K^-	seen
Γ_5 $\gamma\gamma$	seen

DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=5;OUR EST;→ UNCHECKED ←

$f_2(1810)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M038225

$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_5/\Gamma$
5.2 ^{+0.9} +37.3 -0.8-4.5	¹ UEHARA	10A	BELL	10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

NODE=M038G01
NODE=M038G01

¹ Including interference with the $f_2'(1525)$ (parameters fixed to the values from the 2008 edition of this review, PDG 08) and $f_2(1270)$. May also be the $f_0(1500)$.

NODE=M038G01;LINKAGE=UE

$f_2(1810)$ BRANCHING RATIOS

NODE=M038220

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
not seen	AMSLER	02	CBAR	$0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
not seen	PROKOSHKIN	97	GAM2	$38 \pi^-p \rightarrow \pi^0\pi^0n$
0.21 ^{+0.02} -0.03	¹ LONGACRE	86	RVUE	Compilation
0.44±0.03	² CASON	82	STRC	$8 \pi^+p \rightarrow \Delta^{++}\pi^0\pi^0$

NODE=M038R2
NODE=M038R2

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R2;LINKAGE=L

² Included in LONGACRE 86 global analysis.

NODE=M038R;LINKAGE=C

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	ABLIKIM	13N	BES3	PWA of $J/\psi \rightarrow \gamma\eta\eta$

NODE=M038R3
NODE=M038R3

••• We do not use the following data for averages, fits, limits, etc. •••

0.008 ^{+0.028} -0.003	¹ LONGACRE	86	RVUE	Compilation
-----------------------------------	-----------------------	----	------	-------------

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R3;LINKAGE=L

$\Gamma(\pi\pi)/\Gamma(4\pi^0)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_3
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.75	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$	

NODE=M038R4
NODE=M038R4

 $\Gamma(4\pi^0)/\Gamma(\eta\eta)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
••• We do not use the following data for averages, fits, limits, etc. •••				
0.8 ± 0.3	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$	

NODE=M038R5
NODE=M038R5

 $\Gamma(K^+K^-)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
••• We do not use the following data for averages, fits, limits, etc. •••				
$0.003^{+0.019}_{-0.002}$	¹ LONGACRE	86	RVUE Compilation	
seen	COSTA	80	OMEG 10 $\pi^- p \rightarrow K^+K^- n$	

NODE=M038R1
NODE=M038R1

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.

NODE=M038R1;LINKAGE=L

 $f_2(1810)$ REFERENCES

ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
PROKOSHKIN	97	PD 42 117	Y.D. Prokoshkin <i>et al.</i>	(SERP)	REFID=45386
ALDE	88D	Translated from DANS 353 323. SJNP 47 810	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=44652
ALDE	87	Translated from YAF 47 1273. PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
COSTA	80	NP B175 402	G. Costa <i>et al.</i>	(BARI, BONN, CERN, GLAS+)	REFID=20737
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355

NODE=M038

 $X(1835)$

$$J^{PC} = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Could be a superposition of two states, one with small width appearing as threshold enhancement in $p\bar{p}$, the other one with a larger width. For the former ABLIKIM 12D determine $J^{PC} = 0^-+.$

NODE=M085

NODE=M085

 $X(1835)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$1826.5^{+13.0}_{-3.4}$ OUR AVERAGE				
$1825.3 \pm 2.4^{+17.3}_{-2.4}$		¹ ABLIKIM	16J	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
$1844 \pm 9^{+16}_{-25}$		² ABLIKIM	15T	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
••• We do not use the following data for averages, fits, limits, etc. •••				
$1839 \pm 26 \pm 26$		³ ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
$1909.5 \pm 15.9^{+9.4}_{-27.5}$		⁴ ABLIKIM	16J	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
$1842.2 \pm 4.2^{+7.1}_{-2.6}$	0.6k	ABLIKIM	13U	BES3 $J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
$1832^{+19}_{-5} \pm 26$		⁵ ABLIKIM	12D	BES3 $J/\psi \rightarrow \gamma p\bar{p}$
$1836.5 \pm 3.0^{+5.6}_{-2.1}$	4265	⁶ ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
$1877.3 \pm 6.3^{+3.4}_{-7.4}$		⁷ ABLIKIM	11J	BES3 $J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
$1837^{+10}_{-12} \pm 9$	231	^{8,9} ALEXANDER	10	CLEO $J/\psi \rightarrow \gamma p\bar{p}$
$1833.7 \pm 6.1 \pm 2.7$	264	ABLIKIM	05R	BES2 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
1831 ± 7		^{9,10} ABLIKIM	05R	BES2 $J/\psi \rightarrow \gamma p\bar{p}$
$1859^{+3}_{-10} \pm 5$		⁹ BAI	03F	BES2 $J/\psi \rightarrow \gamma p\bar{p}$

NODE=M085M

NODE=M085M

OCCUR=2

OCCUR=2

- ¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors. NODE=M085M;LINKAGE=B
- ² Decay dominated by $f_0(980)\eta$ hence $I^G(J^{PC}) = 0^+(0^-+)$. NODE=M085M;LINKAGE=D
- ³ From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded. NODE=M085M;LINKAGE=C
- ⁴ Pole mass from a fit of the measured $\pi^+\pi^-\eta'$ lineshape to a Flatte formula that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold; the fit also includes known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. NODE=M085M;LINKAGE=A
- ⁵ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G. NODE=M085M;LINKAGE=AK
- ⁶ From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma X(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$. NODE=M085M;LINKAGE=AI
- ⁷ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$. NODE=M085M;LINKAGE=BL
- ⁸ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV. NODE=M085M;LINKAGE=AE
- ⁹ Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, DEL-AMO-SANCHEZ 12 in $B \rightarrow D(D^*)p\bar{p}(\pi)$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$. NODE=M085M;LINKAGE=HF
- ¹⁰ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Systematic errors not estimated. NODE=M085M;LINKAGE=AB

X(1835) WIDTH

NODE=M085W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	-----	------	-------------	------	---------

NODE=M085W

242 $^{+14}_{-15}$ OUR AVERAGE

245.2 ± 13.1 $^{+4.6}_{-9.6}$			1 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
192 $^{+20}_{-17}$ $^{+62}_{-43}$			2 ABLIKIM	15T BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
175 ± 57 ± 25			3 ABLIKIM	18I BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
273.5 ± 21.4 $^{+6.1}_{-64.0}$			4 ABLIKIM	16J BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
83 ± 14 ± 11	90	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$
< 76			5 ABLIKIM	12D BES3	$J/\psi \rightarrow \gamma p\bar{p}$
190 ± 9 $^{+38}_{-36}$		4265	6 ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
57 ± 12 $^{+19}_{-4}$			7 ABLIKIM	11J BES3	$J/\psi \rightarrow \omega(\eta\pi^+\pi^-)$
0 $^{+44}_{-0}$		231	8,9 ALEXANDER	10 CLEO	$J/\psi \rightarrow \gamma p\bar{p}$
67.7 ± 20.3 ± 7.7		264	ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
< 153	90		9,10 ABLIKIM	05R BES2	$J/\psi \rightarrow \gamma p\bar{p}$
< 30			9 BAI	03F BES2	$J/\psi \rightarrow \gamma p\bar{p}$

OCCUR=2

OCCUR=2

- ¹ From a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold through interference with a second previously unseen narrow resonance near 1870 MeV. The fit uses Breit-Wigner functions for the signal shapes and includes known backgrounds and contributors. NODE=M085W;LINKAGE=B
- ² Decay dominated by $f_0(980)\eta$ hence $I^G(J^{PC}) = 0^+(0^-+)$. NODE=M085W;LINKAGE=D
- ³ From a fit to $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^-+$. Other J^{PC} not excluded. NODE=M085W;LINKAGE=C
- ⁴ Pole width from a fit of the measured $\pi^+\pi^-\eta'$ lineshape to a Flatte formula that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold; the fit also includes known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner function ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. NODE=M085W;LINKAGE=A
- ⁵ From the fit including final state interaction effects in isospin 0 *S*-wave according to SIBIRTSEV 05A. Supersedes ABLIKIM 10G. NODE=M085W;LINKAGE=AK
- ⁶ From a fit of the $\pi^+\pi^-\eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma X(2370)$, for $M(\pi^+\pi^-\eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0\pi^+\pi^-\eta'$. NODE=M085W;LINKAGE=AI
- ⁷ The selected process is $J/\psi \rightarrow \omega a_0(980)\pi$. This state may be due also to $\eta_2(1870)$ or to a combination of $X(1835)$ and $\eta_2(1870)$. NODE=M085W;LINKAGE=BL
- ⁸ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV. NODE=M085W;LINKAGE=AE

⁹Evidence for a threshold enhancement in the $p\bar{p}$ mass spectrum was also reported by ABE 02K, AUBERT,B 05L, and WANG 05A in $B^+ \rightarrow p\bar{p}K^+$, WANG 05A in $B^0 \rightarrow p\bar{p}K_S^0$, ABE 02W in $\bar{B}^0 \rightarrow p\bar{p}D^0$, DEL-AMO-SANCHEZ 12 in $B \rightarrow D(D^*)p\bar{p}(\pi)$, and WEI 08 in $B^+ \rightarrow p\bar{p}\pi^+$ decays. Not seen by ATHAR 06 in $\Upsilon(1S) \rightarrow p\bar{p}\gamma$.

¹⁰From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A. Systematic errors not estimated.

NODE=M085W;LINKAGE=HF

NODE=M085W;LINKAGE=AB

X(1835) DECAY MODES

NODE=M085215;NODE=M085

Mode	Fraction (Γ_i/Γ)
Γ_1 $p\bar{p}$	seen
Γ_2 $\eta'\pi^+\pi^-$	seen
Γ_3 $\gamma\gamma$	
Γ_4 $K_S^0 K_S^0 \eta$	seen
Γ_5 $\gamma\phi(1020)$	possibly seen
Γ_6 $3(\pi^+\pi^-)$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2

DESIG=4

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6

DESIG=7

X(1835) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M085225

$\Gamma(\eta'\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_3/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M085G01

NODE=M085G01

••• We do not use the following data for averages, fits, limits, etc. •••

<35.6	90	¹ ZHANG	12A	BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
-------	----	--------------------	-----	------	--

OCCUR=2

<83	90	² ZHANG	12A	BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
-----	----	--------------------	-----	------	--

NODE=M085G01;LINKAGE=ZH

¹From a two-resonance fit and constructive interference of the $\eta(1760)$ and $X(1835)$, a significance of 2.8σ .

²From a two-resonance fit and destructive interference of the $\eta(1760)$ and $X(1835)$, a significance of 2.8σ .

NODE=M085G01;LINKAGE=ZA

X(1835) BRANCHING RATIOS

NODE=M085220

$\Gamma(p\bar{p})/\Gamma(\eta'\pi^+\pi^-)$ Γ_1/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M085R01

NODE=M085R01

••• We do not use the following data for averages, fits, limits, etc. •••

0.333	ABLIKIM	05R	BES2	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
-------	---------	-----	------	--

$\Gamma(\eta'\pi^+\pi^-)/\Gamma(K_S^0 K_S^0 \eta)$ Γ_2/Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M085R00

NODE=M085R00

••• We do not use the following data for averages, fits, limits, etc. •••

6.7 ± 1.8	¹ ABLIKIM	15T	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
---------------	----------------------	-----	------	--

NODE=M085R00;LINKAGE=A

¹Using results from ABLIKIM 05R.

$\Gamma(\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M085R03

NODE=M085R03

seen	¹ ABLIKIM	16J	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$
------	----------------------	-----	------	--

NODE=M085R03;LINKAGE=A

¹ABLIKIM 16J quotes $B(J/\psi \rightarrow \gamma X(1835)) \times B(X(1835) \rightarrow \pi^+\pi^-\eta') = (3.93 \pm 0.38_{-0.84}^{+0.31}) \times 10^{-4}$ from a fit of the measured $\pi^+\pi^-\eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

$\Gamma(\gamma\phi(1020))/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M085R04

NODE=M085R04

possibly seen	¹ ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma\gamma\phi(1020)$
---------------	----------------------	-----	------	---

NODE=M085R04;LINKAGE=A

¹Seen as a peak in $\gamma\phi$ invariant mass. Angular analysis consistent with $J^{PC} = 0^{-+}$. Other J^{PC} not excluded.

$\Gamma(\gamma\gamma)/\Gamma(\eta'\pi^+\pi^-)$ Γ_3/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M085R05

NODE=M085R05

$<9.80 \times 10^{-3}$	90	¹ ABLIKIM	18O	BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma$
------------------------	----	----------------------	-----	------	---

NODE=M085R05;LINKAGE=A

¹Using results from ABLIKIM 16J.

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	0.6k	ABLIKIM	13U BES3	$J/\psi \rightarrow \gamma 3(\pi^+\pi^-)$

NODE=M085R06
 NODE=M085R06

X(1835) REFERENCES

ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.) JPC
DEL-AMO-SA...	12	PR D85 092017	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
ZHANG	12A	PR D86 052002	C.C. Zhang <i>et al.</i>	(BELLE Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11J	PRL 107 182001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	10G	CP C34 421	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
WEI	08	PL B659 80	J.-T. Wei <i>et al.</i>	(BELLE Collab.)
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)
AUBERT,B	05L	PR D72 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer	
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)
ABE	02K	PRL 88 181803	K. Abe <i>et al.</i>	(BELLE Collab.)
ABE	02W	PRL 89 151802	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M085

REFID=58893
 REFID=58925
 REFID=57454
 REFID=56785
 REFID=55582
 REFID=54269
 REFID=54286
 REFID=54763
 REFID=53684
 REFID=53931
 REFID=55685
 REFID=53525
 REFID=52086
 REFID=50993
 REFID=50985
 REFID=50827
 REFID=51038
 REFID=50651
 REFID=49473
 REFID=48690
 REFID=48980

 $\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

NODE=M054

 $\phi_3(1850)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1854 ± 7 OUR AVERAGE				
1855 ± 10		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
1870 ⁺³⁰ ₋₂₀	430	ARMSTRONG	82 OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
1850 ± 10	123	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \Lambda$

NODE=M054M

NODE=M054M

 $\phi_3(1850)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
87⁺²⁸₋₂₃ OUR AVERAGE	Error includes scale factor of 1.2.			
64 ± 31		ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
160 ⁺⁹⁰ ₋₅₀	430	ARMSTRONG	82 OMEG	18.5 $K^- p \rightarrow$ $K^- K^+ \Lambda$
80 ⁺⁴⁰ ₋₃₀	123	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \Lambda$

NODE=M054W

NODE=M054W

 $\phi_3(1850)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}$	seen
Γ_2 $K K^*(892) + \text{c.c.}$	seen

NODE=M054215;NODE=M054

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

 $\phi_3(1850)$ BRANCHING RATIOS

$\Gamma(K \bar{K}^*(892) + \text{c.c.})/\Gamma(K \bar{K})$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.55^{+0.85}_{-0.45}	ASTON	88E LASS	11 $K^- p \rightarrow K^- K^+ \Lambda$, $K_S^0 K^\pm \pi^\mp \Lambda$
0.8 ± 0.4	ALHARRAN	81B HBC	8.25 $K^- p \rightarrow K \bar{K} \pi \Lambda$

NODE=M054220

NODE=M054R1

NODE=M054R1

••• We do not use the following data for averages, fits, limits, etc. •••

 $\phi_3(1850)$ REFERENCES

ASTON	88E	PL B208 324	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) IGJPC
ARMSTRONG	82	PL 110B 77	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+) JP
ALHARRAN	81B	PL 101B 357	S. Al-Harran <i>et al.</i>	(BIRM, CERN, GLAS+)

NODE=M054

REFID=40577
 REFID=21405
 REFID=21702

$\eta_1(1855)$

$$I^G(J^{PC}) = 0^+(1^{-+})$$

NODE=M267

OMITTED FROM SUMMARY TABLE

Meson with exotic (non- $q\bar{q}$) quantum numbers. A state decaying into $\eta\eta'$ with possible quantum numbers 1^{-+} was reported earlier in this mass region BARBERIS 00A in high energy central pp production and by ALDE 91B in $\pi^- p$ interactions, see the $f_2(1910)$, and the review on "Spectroscopy of Light Meson Resonances."

NODE=M267

 $\eta_1(1855)$ MASS

NODE=M267M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$1855 \pm 9^{+6}_{-1}$	¹ ABLIKIM	22A1	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$

NODE=M267M

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave. For analysis details see ABLIKIM 22AS.

NODE=M267M;LINKAGE=A

 $\eta_1(1855)$ WIDTH

NODE=M267W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$188 \pm 18^{+3}_{-8}$	¹ ABLIKIM	22A1	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$

NODE=M267W

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P -wave. For analysis details see ABLIKIM 22AS.

NODE=M267W;LINKAGE=A

 $\eta_1(1855)$ DECAY MODES

NODE=M267215;NODE=M267

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \eta\eta'$	seen

DESIG=1

$\Gamma(\eta\eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	ABLIKIM	22A1	BES3 $J/\psi \rightarrow \gamma\eta\eta'$	
seen	BARBERIS	00A	450 $pp \rightarrow p_f \eta\eta' p_s$	
seen	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$	

NODE=M267R00
NODE=M267R00 **$\eta_1(1855)$ REFERENCES**

NODE=M267

ABLIKIM	22A1	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61881
Also		PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62033
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
ALDE	91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41844
		Translated from YAF 54 751.			

$\eta_2(1870)$

$$I^G(J^{PC}) = 0^+(2^-+)$$

NODE=M101

 $\eta_2(1870)$ MASS

NODE=M101M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1842 ± 8 OUR AVERAGE				
1835 ± 12		BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
1844 ± 13		BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
1840 ± 25		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
1875 ± 20 ± 35		ADOMEIT	96 CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
1881 ± 32 ± 40	26	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1860 ± 5 ± 15		ANISOVICH	00E SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
1840 ± 15		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M101M

 $\eta_2(1870)$ WIDTH

NODE=M101W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
225 ± 14 OUR AVERAGE				
235 ± 22		BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
228 ± 23		BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
200 ± 40		BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+ \pi^-)$
200 ± 25 ± 45		ADOMEIT	96 CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
221 ± 92 ± 44	26	KARCH	92 CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
250 ± 25 $^{+50}_{-35}$		ANISOVICH	00E SPEC	0.9–1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
170 ± 40		BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M101W

 $\eta_2(1870)$ DECAY MODES

NODE=M101225;NODE=M101

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta \pi \pi$	seen
Γ_2 $a_2(1320) \pi$	seen
Γ_3 $f_2(1270) \eta$	seen
Γ_4 $a_0(980) \pi$	seen
Γ_5 $\gamma \gamma$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=8;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=9

 $\eta_2(1870)$ BRANCHING RATIOS

NODE=M101230

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_3
$\Gamma(a_2(1320)\pi)/\Gamma(f_2(1270)\eta)$				
1.7 ± 0.4 OUR AVERAGE				
1.60 ± 0.40	¹ ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$
20.4 ± 6.6	BARBERIS	00B		450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
4.1 ± 2.3	ADOMEIT	96	CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.				

NODE=M101R2
NODE=M101R2

NODE=M101R2;LINKAGE=AN

VALUE	DOCUMENT ID	COMMENT	Γ_2/Γ_4
$\Gamma(a_2(1320)\pi)/\Gamma(a_0(980)\pi)$			
32.6 ± 12.6			
	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$

NODE=M101R4
NODE=M101R4

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3
$\Gamma(a_0(980)\pi)/\Gamma(f_2(1270)\eta)$				
0.48 ± 0.45				
	¹ ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$
¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.				

NODE=M101R01
NODE=M101R01

NODE=M101R01;LINKAGE=AN

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				
seen	KARCH	92	CBAL	$e^+ e^- \rightarrow e^+ e^- \eta \pi^0 \pi^0$

NODE=M101R02
NODE=M101R02

$\eta_2(1870)$ REFERENCES

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>	
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)
KARCH	92	ZPHY C54 33	K. Karch <i>et al.</i>	(Crystal Ball Collab.)

NODE=M101

REFID=53631
REFID=47945
REFID=47958
REFID=47959
REFID=46606
REFID=45758
REFID=45202
REFID=42170

$\pi_2(1880)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

NODE=M185

$\pi_2(1880)$ MASS

NODE=M185M

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M185M

1874^{+26}_{-5} OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.


1847^{+20}_{-3}	46M	¹ AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$1929 \pm 24 \pm 18$	4k	EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
$1876 \pm 11 \pm 67$	145k	LU	05	B852	- 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
$2003 \pm 88 \pm 148$	69k	KUHN	04	B852	- 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1880 ± 20		ANISOVICH	01B	SPEC	0 0.6-1.94 $\bar{p} p \rightarrow \eta \eta \pi^0 \pi^0$
---------------	--	-----------	-----	------	--

NODE=M185M;LINKAGE=A

¹Statistical error negligible.



$\pi_2(1880)$ WIDTH

NODE=M185W

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M185W

237^{+33}_{-30} OUR AVERAGE Error includes scale factor of 1.2.

246^{+33}_{-28}	46M	² AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$323 \pm 87 \pm 43$	4k	EUGENIO	08	B852	- 18 $\pi^- p \rightarrow \eta \eta \pi^- p$
$146 \pm 17 \pm 62$	145k	LU	05	B852	- 18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
$306 \pm 132 \pm 121$	69k	KUHN	04	B852	- 18 $\pi^- p \rightarrow \eta \pi^+ \pi^- \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

255 ± 45		ANISOVICH	01B	SPEC	0 0.6-1.94 $\bar{p} p \rightarrow \eta \eta \pi^0 \pi^0$
--------------	--	-----------	-----	------	--

²Statistical error negligible.

NODE=M185W;LINKAGE=A

$\pi_2(1880)$ DECAY MODES

NODE=M185215;NODE=M185

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\eta\pi^-$	seen
Γ_2 $a_0(980)\eta$	seen
Γ_3 $a_2(1320)\eta$	seen
Γ_4 $f_0(1500)\pi$	seen
Γ_5 $f_1(1285)\pi$	seen
Γ_6 $\omega\pi^-\pi^0$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=5;OUR EVAL;→ UNCHECKED ←
DESIG=6;OUR EVAL;→ UNCHECKED ←

 $\Gamma(a_2(1320)\eta)/\Gamma(f_1(1285)\pi)$ Γ_3/Γ_5

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------	------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

22.7±7.3	69k	KUHN	04	B852	-	18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^- p$
----------	-----	------	----	------	---	--

NODE=M185R01
NODE=M185R01

 $\Gamma(f_0(1500)\pi)/\Gamma(a_0(980)\eta)$ Γ_4/Γ_2

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

0.28 ^{+0.20} _{-0.15}	³ ANISOVICH	01B	SPEC	0	0.6-1.94 $\bar{p}p \rightarrow \eta\eta\pi^0\pi^0$
--	------------------------	-----	------	---	--

³Systematic errors not estimated.

NODE=M185R02
NODE=M185R02

NODE=M185R02;LINKAGE=NS

 $\pi_2(1880)$ REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)
EUGENIO	08	PL B660 466	P. Eugenio <i>et al.</i>	(BNL E852 Collab.)
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)
ANISOVICH	01B	PL B500 222	A.V. Anisovich <i>et al.</i>	

NODE=M185

REFID=59471
REFID=52160
REFID=50459
REFID=49773
REFID=48318

$\rho(1900)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M170

NODE=M170

 $\rho(1900)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

1880±10		¹ ABLIKIM	22L	BES3	2.0-3.08 $e^+e^- \rightarrow K^+K^-\pi^0$
1909±17±25	54	² AUBERT	08S	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
1880±30		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
1860±20		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
1910±10		^{3,4} FRABETTI	04	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
1870±10		ANTONELLI	96	SPEC	$e^+e^- \rightarrow$ hadrons

NODE=M170M

NODE=M170M

OCCUR=2

¹ From a partial wave amplitude analysis at $\sqrt{s} = 2.125$ GeV which includes all the possible intermediate states that match J^{PC} conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

² From the fit with two resonances.

³ From a fit with two resonances with the JACOB 72 continuum.

⁴ Supersedes FRABETTI 01.

NODE=M170M;LINKAGE=A

NODE=M170M;LINKAGE=AU
NODE=M170M;LINKAGE=PI
NODE=M170M;LINKAGE=RS

 $\rho(1900)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

69±15		¹ ABLIKIM	22L	BES3	2.0-3.08 $e^+e^- \rightarrow K^+K^-\pi^0$
48±17±2	54	² AUBERT	08S	BABR	10.6 $e^+e^- \rightarrow \phi\pi^0\gamma$
130±30		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 3\pi^+3\pi^-\gamma$
160±20		AUBERT	06D	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$
37±13		^{3,4} FRABETTI	04	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$
10±5		ANTONELLI	96	SPEC	$e^+e^- \rightarrow$ hadrons

NODE=M170W

NODE=M170W

OCCUR=2

- ¹ From a partial wave amplitude analysis at $\sqrt{s} = 2.125$ GeV which includes all the possible intermediate states that match J^{PC} conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.
- ² From the fit with two resonances.
- ³ From a fit with two resonances with the JACOB 72 continuum.
- ⁴ Supersedes FRABETTI 01.

NODE=M170W;LINKAGE=A

NODE=M170W;LINKAGE=AU
NODE=M170W;LINKAGE=PI
NODE=M170W;LINKAGE=RS $\rho(1900) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$ $\Gamma(\phi\pi)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_6/\Gamma$

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.2 \pm 1.2 \pm 0.8$	54	¹ AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\pi^0\gamma$
-----------------------	----	---------------------	----------	---

¹ From the fit with two resonances.

NODE=M170215

NODE=M170B01
NODE=M170B01

NODE=M170B01;LINKAGE=AU

 $\rho(1900)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 6π	seen
Γ_2 $3\pi^+3\pi^-$	seen
Γ_3 $2\pi^+2\pi^-2\pi^0$	
Γ_4 $\phi\pi$	seen
Γ_5 hadrons	seen
Γ_6 e^+e^-	seen
Γ_7 NN	not seen

NODE=M170225;NODE=M170

DESIG=5;OUR EST;→ UNCHECKED ←
DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=6
DESIG=7;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ← $\rho(1900)$ BRANCHING RATIOS

$\Gamma(6\pi)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	8k	AKHMETSHIN 13	CMD3	$e^+e^- \rightarrow 3\pi^+3\pi^-$	
not seen		AGNELLO 02	OBLX	$\bar{n}p \rightarrow 3\pi^+2\pi^-\pi^0$	
seen		FRABETTI 01	E687	$\gamma p \rightarrow 3\pi^+3\pi^-p$	
seen		ANTONELLI 96	SPEC	$e^+e^- \rightarrow \text{hadrons}$	

NODE=M170230

NODE=M170R1
NODE=M170R1 $\rho(1900)$ REFERENCES

ABLIKIM 22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AKHMETSHIN 13	PL B723 82	R.R. Akhmetshin <i>et al.</i>	(CMD-3 Collab.)
AUBERT 08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT 06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)
FRABETTI 04	PL B578 290	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
AGNELLO 02	PL B527 39	M. Agnello <i>et al.</i>	(OBELIX Collab.)
FRABETTI 01	PL B514 240	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)
ANTONELLI 96	PL B365 427	A. Antonelli <i>et al.</i>	(FENICE Collab.)
JACOB 72	PR D5 1847	M. Jacob, R. Slansky	

NODE=M170

REFID=61649
REFID=55370
REFID=52242
REFID=51047
REFID=49614
REFID=48576
REFID=48350
REFID=44633
REFID=49668

$f_2(1910)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

We list here three different peaks with close masses and widths seen in the mass distributions of $\omega\omega$, $\eta\eta'$, and K^+K^- final states. ALDE 91B argues that they are of different nature.

NODE=M142

NODE=M142

 $f_2(1910)$ MASS

NODE=M142205


NODE=M142MX

 $f_2(1910)$ $\omega\omega$ MODENODE=M142M2
NODE=M142M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 ± 9 OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
1890 ± 10	¹ AMELIN	06	VES 36 $\pi^- p \rightarrow \omega\omega n$
1897 ± 11	BARBERIS	00F	450 $p p \rightarrow p_f \omega\omega p_S$
1924 ± 14	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$

¹ Supersedes BELADIDZE 92B.

NODE=M142M2;LINKAGE=AM

 **$f_2(1910)$ $\eta\eta'$ MODE**NODE=M142M3
NODE=M142M3

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1934 ± 16	¹ BARBERIS	00A	450 $p p \rightarrow p_f \eta\eta' p_S$
1934 ± 20	² ANISOVICH	00J	SPEC
1911 ± 10	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta\eta' n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Also compatible with $J^{PC} = 1^-+$.² Combined fit with $\eta\eta$, $\pi\pi$, and $\eta\pi\pi$.NODE=M142M3;LINKAGE=KS
NODE=M142M3;LINKAGE=AN **$f_2(1910)$ K^+K^- MODE**NODE=M142M4
NODE=M142M4

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1941 ± 18	¹ AMSLER	06	CBAR 1.64 $\bar{p} p \rightarrow K^+ K^- \pi^0$

¹ Tentative, could be $f_2(1950)$.

NODE=M142M4;LINKAGE=A

$f_2(1910)$ WIDTH

NODE=M142210


NODE=M142WX

NODE=M142W2
NODE=M142W2 **$f_2(1910)$ $\omega\omega$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
167±21 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
165±19	¹ AMELIN	06	VES 36 $\pi^- p \rightarrow \omega\omega n$
202±32	BARBERIS	00F	450 $pp \rightarrow p_f \omega \omega p_S$
91±50	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$

¹ Supersedes BELADIDZE 92B.

NODE=M142W2;LINKAGE=AM

 **$f_2(1910)$ $\eta\eta'$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
140±40 OUR AVERAGE	[141 ± 40 MeV OUR 2023 AVERAGE]		
141±41	¹ BARBERIS	00A	450 $pp \rightarrow p_f \eta \eta' p_S$
271±25	² ANISOVICH	00J	SPEC
90±35	ALDE	91B	GAM2 38 $\pi^- p \rightarrow \eta \eta' n$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

¹ Also compatible with $J^{PC}=1^-+$.² Combined fit with $\eta\eta$, $\pi\pi$, and $\eta\pi\pi$.NODE=M142W3
NODE=M142W3

NEW

NODE=M142W3;LINKAGE=KS
NODE=M142W3;LINKAGE=AN **$f_2(1910)$ $K^+ K^-$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120±40	AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+ K^- \pi^0$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

NODE=M142W4
NODE=M142W4 **$f_2(1910)$ DECAY MODES**

NODE=M142215;NODE=M142

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^0 \pi^0$	
Γ_2 $K^+ K^-$	seen
Γ_3 $K_S^0 K_S^0$	
Γ_4 $\eta\eta$	seen
Γ_5 $\omega\omega$	seen
Γ_6 $\eta\eta'$	seen
Γ_7 $\eta'\eta'$	
Γ_8 $\rho\rho$	seen
Γ_9 $a_2(1320)\pi$	seen
Γ_{10} $f_2(1270)\eta$	seen

DESIG=6

DESIG=11

DESIG=8

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=9

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=12;OUR EST;→ UNCHECKED ←

DESIG=13;OUR EST;→ UNCHECKED ←

$f_2(1910)$ BRANCHING RATIOS $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AMSLER	06	CBAR 1.64 $\bar{p}p \rightarrow K^+K^-\pi^0$

¹ Tentative, could be $f_2(1950)$.

NODE=M142225

NODE=M142R11
NODE=M142R11

NODE=M142R11;LINKAGE=A

 $\Gamma(\pi^0\pi^0)/\Gamma(\eta\eta')$ Γ_1/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
<0.1	ALDE	89	GAM2 $38\pi^-p \rightarrow \eta\eta'n$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M142R4
NODE=M142R4 $\Gamma(K_S^0K_S^0)/\Gamma(\eta\eta')$ Γ_3/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.066	90	BALOSHIN	86	SPEC $40\pi p \rightarrow K_S^0K_S^0n$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M142R7
NODE=M142R7 $\Gamma(\eta\eta)/\Gamma(\eta\eta')$ Γ_4/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	ALDE	91B	GAM2 $38\pi^-p \rightarrow \eta\eta'n$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M142R6
NODE=M142R6 $\Gamma(\omega\omega)/\Gamma(\eta\eta')$ Γ_5/Γ_6

VALUE	DOCUMENT ID	COMMENT
2.6 ± 0.6	BARBERIS 00F	$450 pp \rightarrow p_f\omega\omega p_S$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M142R10
NODE=M142R10 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
probably not seen	BARBERIS	00A	$450 pp \rightarrow p_f\eta'\eta' p_S$
possibly seen	BELADIDZE	92D	VES $37\pi^-p \rightarrow \eta'\eta'n$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M142R8
NODE=M142R8 $\Gamma(\rho\rho)/\Gamma(\omega\omega)$ Γ_8/Γ_5

VALUE	DOCUMENT ID	COMMENT
2.6 ± 0.4	BARBERIS 00F	$450 pp \rightarrow p_f\omega\omega p_S$

••• We do not use the following data for averages, fits, limits, etc. •••

NODE=M142R9
NODE=M142R9 $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$ Γ_{10}/Γ_9

VALUE	DOCUMENT ID	TECN	COMMENT
0.09 ± 0.05	¹ ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.NODE=M142R12
NODE=M142R12

NODE=M142R12;LINKAGE=AN

 $f_2(1910)$ REFERENCES

NODE=M142

ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=51574
AMSLER	06	PL B639 165	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=51136
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BARBERIS	00A	PL B471 429	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47957
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47962
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
BELADIDZE	92B	ZPHY C54 367	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=42172
BELADIDZE	92D	ZPHY C57 13	G.M. Beladidze <i>et al.</i>	(VES Collab.)	REFID=43309
ALDE	91B	SJNP 54 455	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=41844
		Translated from YAF 54 751.			
Also		PL B276 375	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41911
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=40935
ALDE	89	PL B216 447	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)	REFID=40727
Also		SJNP 48 1035	D.M. Alde <i>et al.</i>	(BELG, SERP, LANL, LAPP)	REFID=44697
		Translated from YAF 48 1724.			
BALOSHIN	86	SJNP 43 959	O.N. Baloshin <i>et al.</i>	(ITEP)	REFID=40734
		Translated from YAF 43 1487.			

$a_0(1950)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. Seen in $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$ by LEES 16A with significance 2.5σ in $K_S^0 K^\pm \pi^\mp$ and 4.2σ in $K^+ K^- \pi^0$.

NODE=M227

NODE=M227

 $a_0(1950)$ MASS

NODE=M227M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1931±14±22	12k	^{1,2} LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
1949±32±76	8k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
1927±15±23	4k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$

NODE=M227M

OCCUR=3

••• We do not use the following data for averages, fits, limits, etc. •••

¹ From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating width.

OCCUR=2

NODE=M227M;LINKAGE=A

² Weighted average of the $K_S^0 K^\pm$ and $K^+ K^-$ decay modes.

NODE=M227M;LINKAGE=B

 $a_0(1950)$ WIDTH

NODE=M227W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
270±40 OUR AVERAGE		[271 ± 40 MeV OUR 2023 AVERAGE]		
271±22±29	12k	^{1,2} LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
265±36±110	8k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$
274±28±30	4k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K^+ K^- \pi^0$

NODE=M227W

NEW

OCCUR=3

••• We do not use the following data for averages, fits, limits, etc. •••

¹ From a model-independent partial wave analysis fit to a relativistic Breit-Wigner function with a floating mass.

OCCUR=2

NODE=M227W;LINKAGE=A

² Weighted average of the $K_S^0 K^\pm$ and $K^+ K^-$ decay modes.

NODE=M227W;LINKAGE=B

 $a_0(1950)$ DECAY MODES

NODE=M227215;NODE=M227

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad K\bar{K}$	seen

DESIG=1

 $a_0(1950)$ BRANCHING RATIOS

NODE=M227225

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

NODE=M227R01

NODE=M227R01

¹ From a model-independent partial wave analysis.

NODE=M227R01;LINKAGE=A

 $a_0(1950)$ REFERENCES

NODE=M227

LEES 16A PR D93 012005 J.P. Lees *et al.* (BABAR Collab.)

REFID=57125

$f_2(1950)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M135

 $f_2(1950)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

NODE=M135PP

NODE=M135PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1830–2020) – i (110–220) OUR ESTIMATE			
$(1955 \pm 75) - i (175 \pm 57)$	¹ RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1978.2 \pm 1.8^{+28.4}_{-16.9}) - i$ $(118.8 \pm 0.8^{+20.8}_{-7.8})$	² ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta \eta,$ $\pi^0 K^+ K^-$
$(1867 \pm 46) - i (193 \pm 29)$	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$
¹ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).			
² T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ($\pi\pi$), LONGACRE 86 ($K\bar{K}$), BINON 83 ($\eta\eta$).			

NODE=M135PP

→ UNCHECKED ←

NODE=M135PP;LINKAGE=A

NODE=M135PP;LINKAGE=B

 $f_2(1950)$ MASS

NODE=M135M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1936 ± 12 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.			
1940 ± 50	BAI	00A	BES $J/\psi \rightarrow \gamma (\pi^+ \pi^- \pi^+ \pi^-)$
1980 ± 22	¹ BARBERIS	00C	450 $pp \rightarrow pp4\pi$
1940 ± 22	² BARBERIS	00C	450 $pp \rightarrow pp2\pi2\pi^0$
1960 ± 30	BARBERIS	97B	OMEG $450 pp \rightarrow pp2(\pi^+ \pi^-)$
1918 ± 12	ANTINORI	95	OMEG $300,450 pp \rightarrow pp2(\pi^+ \pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2038^{+13+12}_{-11-73}	³ UEHARA	09	BELL $10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1930 ± 25	⁴ BINON	05	GAMS $33 \pi^- p \rightarrow \eta \eta n$
1980 ± 2 ± 14	ABE	04	BELL $10.6 e^+ e^- \rightarrow e^+ e^- K^+ K^-$
2010 ± 25	ANISOVICH	00J	SPEC
1980 ± 50	ANISOVICH	99B	SPEC $1.35-1.94 p\bar{p} \rightarrow \eta \eta \pi^0$
~ 1990	⁵ OAKDEN	94	RVUE $0.36-1.55 \bar{p}p \rightarrow \pi\pi$
1950 ± 15	⁶ ASTON	91	LASS $11 K^- p \rightarrow \Lambda K \bar{K} \pi \pi$
¹ Decaying into $\pi^+ \pi^- 2\pi^0$.			
² Decaying into $2(\pi^+ \pi^-)$.			
³ Taking into account $f_4(2050)$.			
⁴ First solution, PWA is ambiguous.			
⁵ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.			
⁶ Cannot determine spin to be 2.			

NODE=M135M

OCCUR=2

NODE=M135M;LINKAGE=A4


NODE=M135M;LINKAGE=B4

NODE=M135M;LINKAGE=UE

NODE=M135M;LINKAGE=BI

NODE=M135M;LINKAGE=BB

NODE=M135M;LINKAGE=A



f₂(1950) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
464 ± 24 OUR AVERAGE			
380 ⁺¹²⁰ ₋₉₀	BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
520 ± 50	¹ BARBERIS	00C	450 $pp \rightarrow pp4\pi$
485 ± 55	² BARBERIS	00C	450 $pp \rightarrow pp4\pi$
460 ± 40	BARBERIS	97B OMEG	450 $pp \rightarrow pp2(\pi^+\pi^-)$
390 ± 60	ANTINORI	95 OMEG	300,450 $pp \rightarrow pp2(\pi^+\pi^-)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

441 ⁺²⁷ ₋₂₅ ⁺²⁸ ₋₁₉₂	³ UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
450 ± 50	⁴ BINON	05 GAMS	33 $\pi^-p \rightarrow \eta\eta n$
297 ± 12 ± 6	ABE	04 BELL	10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
495 ± 35	ANISOVICH	00J SPEC	
500 ± 100	ANISOVICH	99B SPEC	1.35-1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
~ 100	⁵ OAKDEN	94 RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
250 ± 50	⁶ ASTON	91 LASS	11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

¹ Decaying into $\pi^+\pi^-2\pi^0$.

² Decaying into $2(\pi^+\pi^-)$.

³ Taking into account $f_4(2050)$.

⁴ First solution, PWA is ambiguous.

⁵ From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

⁶ Cannot determine spin to be 2.

NODE=M135W

NODE=M135W

OCCUR=2

NODE=M135W;LINKAGE=A4

NODE=M135W;LINKAGE=B4

NODE=M135W;LINKAGE=UE

NODE=M135W;LINKAGE=BI

NODE=M135W;LINKAGE=BB

NODE=M135W;LINKAGE=A

f₂(1950) DECAY MODES

NODE=M135215;NODE=M135

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\bar{K}^*(892)$	seen
Γ_2 $\pi\pi$	
Γ_3 $\pi^+\pi^-$	seen
Γ_4 $\pi^0\pi^0$	seen
Γ_5 4π	seen
Γ_6 $\pi^+\pi^-\pi^+\pi^-$	
Γ_7 $a_2(1320)\pi$	
Γ_8 $f_2(1270)\pi\pi$	
Γ_9 $\eta\eta$	seen
Γ_{10} $K\bar{K}$	seen
Γ_{11} $\gamma\gamma$	seen
Γ_{12} $p\bar{p}$	seen

DESIG=1

DESIG=11

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=10;OUR EST;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

DESIG=3

DESIG=4

DESIG=5

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=9;OUR EST;→ UNCHECKED ←

DESIG=12

$f_2(1950) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M135225

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{10}\Gamma_{11}/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M135G1
NODE=M135G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$122 \pm 4 \pm 26$	¹ ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
--------------------	------------------	----	---

¹ Assuming spin 2.

NODE=M135G1;LINKAGE=AB

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_{11}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M135G2
NODE=M135G2

• • • We do not use the following data for averages, fits, limits, etc. • • •

$162^{+69+1137}_{-42-204}$	¹ UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
----------------------------	---------------------	----	---

¹ Taking into account $f_4(2050)$.

NODE=M135G2;LINKAGE=UE

 $f_2(1950)$ BRANCHING RATIOS

NODE=M135220

 $\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

NODE=M135R1
NODE=M135R1

seen	ASTON	91	LASS 0	11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$
------	-------	----	--------	--

 $\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M135R3
NODE=M135R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	BARBERIS	00B	450 $pp \rightarrow p_f \eta \pi^+ \pi^- p_s$
----------	----------	-----	---

not seen	BARBERIS	00C	450 $pp \rightarrow p_f 4\pi p_s$
----------	----------	-----	-----------------------------------

possibly seen	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$
---------------	----------	-----	---

 $\Gamma(\eta\eta)/\Gamma(4\pi)$ Γ_9/Γ_5

VALUE	CL%	DOCUMENT ID	COMMENT
-------	-----	-------------	---------

NODE=M135R5
NODE=M135R5

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 5.0 \times 10^{-3}$	90	BARBERIS	00E 450 $pp \rightarrow p_f \eta \eta p_s$
------------------------	----	----------	--

 $\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$ Γ_9/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M135R6
NODE=M135R6

0.14 ± 0.05	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
--------------------	--------	----	--

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M135R07
NODE=M135R07

seen	111	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p\bar{p}$
------	-----	-----------	----	---

 $f_2(1950)$ REFERENCES

NODE=M135

RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
		Translated from YAF 68 998.			
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)	REFID=48580
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47958
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>		REFID=46886
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP	REFID=44437
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)	REFID=41746
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)	REFID=20768
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)	REFID=20750
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)	REFID=20355

NODE=M017

$a_4(1970)$

$$I^G(J^{PC}) = 1^-(4^{++})$$

was $a_4(2040)$

$a_4(1970)$ MASS

NODE=M017M

NODE=M017M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1967±16 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.					
1935 ⁺¹¹ ₋₁₃	46M	¹ AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1900 ⁺⁸⁰ ₋₂₀		ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
1985±10±13	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
1996±25±43		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi p$
2000±40 ⁺⁶⁰ ₋₂₀		IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
2010±20		² ALDE	96	GAM2 0	38 $\pi^- p \rightarrow \eta \pi^0 n$
2040±30		³ CLELAND	82B	SPEC ±	50 $\pi p \rightarrow K_S^0 K^\pm p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1885±13 ⁺⁵⁰ ₋₂	420k	⁴ ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ Pb'$
2004± 6	80k	⁵ UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
2005 ⁺²⁵ ₋₄₅		⁶ ANISOVICH	01F	SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
1944± 8±50		⁷ AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
1903±10		⁸ BALDI	78	SPEC -	10 $\pi^- p \rightarrow p K_S^0 K^-$
2030±50		⁹ CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi n$

- ¹ Statistical error negligible.
- ² From a simultaneous fit to the G_+ and G_0 wave intensities.
- ³ From an amplitude analysis.
- ⁴ Superseded by AGHASYAN 2018B.
- ⁵ Statistical error only.
- ⁶ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.
- ⁷ May be a different state.
- ⁸ From a fit to the Y_8^0 moment. Limited by phase space.
- ⁹ $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.

NODE=M017M;LINKAGE=B
 NODE=M017M;LINKAGE=A
 NODE=M017M;LINKAGE=C
 NODE=M017M;LINKAGE=D
 NODE=M017M;LINKAGE=ST
 NODE=M017M;LINKAGE=AN
 NODE=M017M;LINKAGE=DM
 NODE=M017M;LINKAGE=Y
 NODE=M017M;LINKAGE=M



$a_4(1970)$ WIDTH

NODE=M017W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
324⁺₋₁₈	OUR AVERAGE				
333 ⁺ ₋₂₁	46M	¹ AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
300 ⁺ ₋₁₀₀		ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$
231 \pm 30 \pm 46	145k	LU	05	B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
298 \pm 81 \pm 85		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi p$
350 \pm 100 ⁺ ₋₅₀		IVANOV	01	B852	18 $\pi^- p \rightarrow \eta' \pi^- p$
370 \pm 80		² ALDE	96	GAM2 0	38 $\pi^- p \rightarrow \eta \pi^0 n$
380 \pm 150		³ CLELAND	82B	SPEC \pm	50 $\pi p \rightarrow K_S^0 K^\pm p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

294 \pm 25 ⁺ ₋₁₉	420k	⁴ ALEKSEEV	10	COMP	190 $\pi^- Pb \rightarrow \pi^- \pi^- \pi^+ P_b'$
401 \pm 16	80k	⁵ UMAN	06	E835	5.2 $\bar{p} p \rightarrow \eta \eta \pi^0$
180 \pm 30		⁶ ANISOVICH	01F	SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
324 \pm 26 \pm 75		⁷ AMELIN	99	VES	37 $\pi^- A \rightarrow \omega \pi^- \pi^0 A^*$
166 \pm 43		⁸ BALDI	78	SPEC $-$	10 $\pi^- p \rightarrow p K_S^0 K^-$
510 \pm 200		⁹ CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi n$

¹ Statistical error negligible.

² From a simultaneous fit to the G_+ and G_0 wave intensities.

³ From an amplitude analysis.

⁴ Superseded by AGHASYAN 2018B.

⁵ Statistical error only.

⁶ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

⁷ May be a different state.

⁸ From a fit to the Y_8^0 moment. Limited by phase space.

⁹ $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.

NODE=M017W

NODE=M017W;LINKAGE=B
 NODE=M017W;LINKAGE=A
 NODE=M017W;LINKAGE=C
 NODE=M017W;LINKAGE=D
 NODE=M017W;LINKAGE=ST
 NODE=M017W;LINKAGE=AN
 NODE=M017W;LINKAGE=DM
 NODE=M017W;LINKAGE=Y
 NODE=M017W;LINKAGE=M

$a_4(1970)$ DECAY MODES

NODE=M017215;NODE=M017

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\pi^+ \pi^- \pi^0$	seen
Γ_3 $\rho\pi$	seen
Γ_4 $f_2(1270)\pi$	seen
Γ_5 $\omega \pi^- \pi^0$	seen
Γ_6 $\omega\rho$	seen
Γ_7 $\eta\pi$	seen
Γ_8 $\eta'(958)\pi$	seen

DESIG=1
 DESIG=2
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6;OUR EST;→ UNCHECKED ←
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=8
 DESIG=3
 DESIG=4;OUR EST;→ UNCHECKED ←

$a_4(1970)$ BRANCHING RATIOS

NODE=M017220

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	BALDI	78	SPEC \pm	10 $\pi^- p \rightarrow K_S^0 K^- p$

NODE=M017R1
 NODE=M017R1

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	CORDEN	78C	OMEG 0	15 $\pi^- p \rightarrow 3\pi n$

NODE=M017R2
 NODE=M017R2

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$	Γ_3/Γ_4		
VALUE	DOCUMENT ID	TECN	COMMENT

NODE=M017R4
 NODE=M017R4

1.7⁺_{-0.8} OUR AVERAGE Error includes scale factor of 3.7.

2.9 ⁺ _{-0.4}	46M	¹ AGHASYAN	18B	COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1.1 \pm 0.2 \pm 0.2		CHUNG	02	B852	18.3 $\pi^- p \rightarrow 3\pi p$

¹ Statistical error negligible.

NODE=M017R4;LINKAGE=A

$\Gamma(\eta\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_7/Γ
seen	ALDE	96	GAM2	0	$38 \pi^- p \rightarrow \eta \pi^0 n$

NODE=M017R3
NODE=M017R3

 $\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_7
0.23±0.07	ADOLPH	15	COMP	191 $\pi^- p \rightarrow \eta^{(\prime)} \pi^- p$

NODE=M017R01
NODE=M017R01

 $\Gamma(\omega\rho)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
seen	145k	LU	05	B852	$18 \pi^- p \rightarrow \omega \pi^- \pi^0 p$

NODE=M017R5
NODE=M017R5

 $a_4(1970)$ REFERENCES

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
ADOLPH	15	PL B740 303	M. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56385
ALEKSEEV	10	PRL 104 241803	M.G. Alekseev <i>et al.</i>	(COMPASS Collab.)	REFID=53356
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352
IVANOV	01	PRL 86 3977	E.I. Ivanov <i>et al.</i>	(BNL E852 Collab.)	REFID=48317
AMELIN	99	PAN 62 445	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=46910
ANISOVICH	99C	Translated from YAF 62 487. PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>		REFID=46902
ALDE	96	PAN 59 982	S.V. Donskov <i>et al.</i>	(GAMS Collab.) IGJPC	REFID=45207
CLELAND	82B	Translated from YAF 59 1027. NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
BALDI	78	PL 74B 413	R. Baldi <i>et al.</i>	(GEVA) JP	REFID=21783
CORDEN	78C	NP B136 77	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+) JP	REFID=20859

NODE=M017

 $\rho_3(1990)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

OMITTED FROM SUMMARY TABLE

NODE=M167

 $\rho_3(1990)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1982±14	¹ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2007	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167M

NODE=M167M

NODE=M167M;LINKAGE=AY

 $\rho_3(1990)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

188±24	² ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 287	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$

² From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M167W

NODE=M167W

NODE=M167W;LINKAGE=AY

 $\rho_3(1990)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103

NODE=M167

$\pi_2(2005)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

NODE=M239

OMITTED FROM SUMMARY TABLE

 $\pi_2(2005)$ MASS

NODE=M239M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1963⁺¹⁷₋₂₇ OUR AVERAGE				
1962 ⁺¹⁷ ₋₂₉	46M	¹ AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
1974 ± 14 ± 83	145k	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2005 ± 15		ANISOVICH	01F SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
¹ Statistical uncertainty negligible.				

NODE=M239M

NODE=M239M;LINKAGE=A

 $\pi_2(2005)$ WIDTH

NODE=M239W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
370⁺¹⁶₋₉₀ OUR AVERAGE				
371 ⁺¹⁶ ₋₁₂₀	46M	¹ AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
341 ± 61 ± 139	145k	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
200 ± 40		ANISOVICH	01F SPEC	2.0 $\bar{p} p \rightarrow 3\pi^0, \pi^0 \eta, \pi^0 \eta'$
¹ Statistical uncertainty negligible.				

NODE=M239W

NODE=M239W;LINKAGE=A

 $\pi_2(2005)$ DECAY MODES

NODE=M239215;NODE=M239

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^- \pi^+ \pi^-$	seen
Γ_2 $\omega \pi^0 \pi^-$	seen

DESIG=1

DESIG=2

 $\pi_2(2005)$ BRANCHING RATIOS

NODE=M239220

$\Gamma(\pi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AGHASYAN	18B COMP	190 $\pi^- p \rightarrow \pi^- \pi^+ \pi^- p$
$\Gamma(\omega \pi^0 \pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	LU	05 B852	18 $\pi^- p \rightarrow \omega \pi^- \pi^0 p$

NODE=M239R00
NODE=M239R00NODE=M239R01
NODE=M239R01 $\pi_2(2005)$ REFERENCES

NODE=M239

AGHASYAN	18B	PR D98 092003	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59471
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352

$f_2(2010)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M106

 $f_2(2010)$ MASS

NODE=M106M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2010 \pm \begin{smallmatrix} 60 \\ 80 \end{smallmatrix}$ OUR AVERAGE	[2011 $\pm \begin{smallmatrix} 60 \\ 80 \end{smallmatrix}$ MeV OUR 2023 AVERAGE]		
$2011 \pm \begin{smallmatrix} 62 \\ 76 \end{smallmatrix}$	¹ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi \phi n$
$2062 \pm \begin{smallmatrix} 6 \\ 7 \end{smallmatrix}$	² ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
2005 ± 12	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1980 ± 20	³ BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
$2050 \pm \begin{smallmatrix} 90 \\ 50 \end{smallmatrix}$	ETKIN	85	MPS 22 $\pi^- p \rightarrow 2\phi n$
$2120 \pm \begin{smallmatrix} 20 \\ 120 \end{smallmatrix}$	LINDENBAUM	84	RVUE
2160 ± 50	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M106M

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

- ¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi \phi 2^{++} S_2$, D_2 , and D_0 is $98 \pm 1_{-3}$, $0 \pm 1_{-0}$, and $2 \pm 2_{-1}$, respectively.
- ² From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.
- ³ Statistically very weak, only 1.4 s.d.

NODE=M106M;LINKAGE=C

NODE=M106M;LINKAGE=A

NODE=M106M;LINKAGE=E

 $f_2(2010)$ WIDTH

NODE=M106W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200 ± 60 OUR AVERAGE	[202 ± 60 MeV OUR 2023 AVERAGE]		
$202 \pm \begin{smallmatrix} 67 \\ 62 \end{smallmatrix}$	⁴ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi \phi n$
$165 \pm \begin{smallmatrix} 17 \\ 5 \end{smallmatrix}$	⁵ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
209 ± 32	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
145 ± 50	⁶ BOLONKIN	88	SPEC 40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
$200 \pm \begin{smallmatrix} 160 \\ 50 \end{smallmatrix}$	ETKIN	85	MPS 22 $\pi^- p \rightarrow 2\phi n$
$300 \pm \begin{smallmatrix} 150 \\ 50 \end{smallmatrix}$	LINDENBAUM	84	RVUE
310 ± 70	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$

NODE=M106W

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

- ⁴ Includes data of ETKIN 85.
- ⁵ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.
- ⁶ Statistically very weak, only 1.4 s.d.

NODE=M106W;LINKAGE=C

NODE=M106W;LINKAGE=A

NODE=M106W;LINKAGE=E

 $f_2(2010)$ DECAY MODES

NODE=M106215;NODE=M106

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi \phi$	seen
Γ_2 $K \bar{K}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2

 $f_2(2010)$ BRANCHING RATIOS

NODE=M106230

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$	

NODE=M106R01

NODE=M106R01

 $f_2(2010)$ REFERENCES

NODE=M106

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
Also		PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62033
VLADIMIRSK...06		PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
		Translated from YAF 69 515.			
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)	REFID=40580
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=40285
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21871
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)	REFID=21869
ETKIN	82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21866
Also		Brighton Conf. 351	S.J. Lindenbaum	(BNL, CUNY)	REFID=21867

$f_0(2020)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

Needs confirmation.

 $f_0(2020)$ T-MATRIX POLE \sqrt{s} Note that $\Gamma = -2 \text{Im}(\sqrt{s})$.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(1870–2080) – i (120–240) OUR ESTIMATE			
(2038 ± 48) – i (156 ± 41)	¹ RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K})$
(1925 ± 25) – i (160 ± 18)	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
(1910 ± 50) – i (199 ± 40)	² ROPERTZ	18	RVUE $\bar{B}_S^0 \rightarrow J/\psi(\pi^+\pi^-/K^+K^-)$
(1992 ± 16) – i (221 ± 30)	³ BARBERIS	00C	450 $p\bar{p} \rightarrow p_f 4\pi p_S$
(2020 ± 35) – i (205 ± 25)	BARBERIS	97B	OMEG 450 $p\bar{p} \rightarrow p\bar{p} 2(\pi^+\pi^-)$
¹ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma\pi^0\pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).			
² T-matrix pole of 3 channel unitary model fit to data from AAIJ 14BR and AAIJ 17V extracted using Pade approximants.			
³ Average between $\pi^+\pi^- 2\pi^0$ and $2(\pi^+\pi^-)$.			

NODE=M156
 NODE=M156
 NODE=M156PP
 NODE=M156PP
 NODE=M156PP
 → UNCHECKED ←

NODE=M156PP;LINKAGE=A
 NODE=M156PP;LINKAGE=C
 NODE=M156PP;LINKAGE=B

 $f_0(2020)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1982 ± 3⁺⁵⁴₋₀		¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2010 ± 6 ⁺⁶ ₋₄		² ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
2037 ± 8	80k	³ UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
2040 ± 38		ANISOVICH	00J	SPEC
2010 ± 60		ALDE	98	GAM4 $100 \pi^- p \rightarrow \pi^0\pi^0 n$
¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta' X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.				
² From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.				
³ Statistical error only.				

NODE=M156M
 NODE=M156M
 NODE=M156M;LINKAGE=C
 NODE=M156M;LINKAGE=D
 NODE=M156M;LINKAGE=ST

 $f_0(2020)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
440 ± 50 OUR AVERAGE		[436 ± 50 MeV OUR 2023 AVERAGE]		
436 ± 4⁺⁴⁶₋₄₉		¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
203 ± 9 ⁺¹³ ₋₁₁		² ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta'$
296 ± 17	80k	³ UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta\eta\pi^0$
405 ± 40		ANISOVICH	00J	SPEC
240 ± 100		ALDE	98	GAM4 $100 \pi^- p \rightarrow \pi^0\pi^0 n$
¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta' X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.				
² From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta\eta'$ P-wave.				
³ Statistical error only.				

NODE=M156W
 NODE=M156W
 NEW
 NODE=M156W;LINKAGE=C
 NODE=M156W;LINKAGE=D
 NODE=M156W;LINKAGE=ST

$f_0(2020)$ DECAY MODES

NODE=M156215;NODE=M156

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi\pi$	seen
Γ_2 $\pi^0\pi^0$	seen
Γ_3 $\rho\rho$	seen
Γ_4 $\omega\omega$	seen
Γ_5 $\eta\eta$	seen
Γ_6 $\eta'\eta'$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←
DESIG=5
DESIG=6

 $f_0(2020)$ BRANCHING RATIOS

NODE=M156220

$\Gamma(\rho\rho)/\Gamma(\omega\omega)$	DOCUMENT ID	COMMENT	Γ_3/Γ_4
VALUE			

NODE=M156R1
NODE=M156R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 3 BARBERIS 00F 450 $\rho\rho \rightarrow p_f\omega p_s$

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
VALUE				

NODE=M156R01
NODE=M156R01

seen UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
VALUE				

NODE=M156R00
NODE=M156R00

seen ¹ ABLIKIM 22C BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M156R00;LINKAGE=A

 $f_0(2020)$ REFERENCES

NODE=M156

ABLIKIM 22AS PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61891
Also PR D107 079901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62033
ABLIKIM 22C PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61637
RODAS 22 EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)	REFID=61610
SARANTSEV 21 PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ABLIKIM 18AA PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ROPERTZ 18 EPJ C78 1000	S. Ropertz, C. Hanhart, B. Kubis	(BONN, JULI)	REFID=59332
AAIJ 17V JHEP 1708 037	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57828
ABLIKIM 15AE PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
AAIJ 14BR PR D89 092006	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56035
UMAN 06 PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
ANISOVICH 00J PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BARBERIS 00C PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47959
BARBERIS 00F PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47962
ALDE 98 EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
Translated from YAF 62 446.			
BARBERIS 97B PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=45758

$f_4(2050)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

NODE=M016

 $f_4(2050)$ MASS

NODE=M016M

NODE=M016M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2018±11 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
1960±15		AMELIN	06	VES 36 $\pi^- p \rightarrow \omega \omega n$
2005±10		¹ BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
1998±15		ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
2060±20		ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega \omega n$
2038±30		AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
2086±15		BALTRUSAIT..	87	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
2000±60		ALDE	86D	GAM4 100 $\pi^- p \rightarrow n 2 \eta$
2020±20	40k	² BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2 \pi^0$
2015±28		³ CASON	82	STRC 8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
2031 ⁺²⁵ ₋₃₆		ETKIN	82B	MPS 23 $\pi^- p \rightarrow n 2 K_S^0$
2020±30	700	APEL	75	NICE 40 $\pi^- p \rightarrow n 2 \pi^0$
2050±25		BLUM	75	ASPK 18.4 $\pi^- p \rightarrow n K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1966±25		⁴ ANISOVICH	09	RVUE 0.0 $\bar{p} p, \pi N$
1885 ⁺¹⁴⁺²¹⁸ ₋₁₃₋₂₅		⁵ UEHARA	09	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
2018± 6		ANISOVICH	00J	SPEC 2.0 $\bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$ $\eta \eta, \eta \eta', \pi \pi$
~ 2000		⁶ MARTIN	98	RVUE $\bar{N} N \rightarrow \pi \pi$
~ 2010		⁷ MARTIN	97	RVUE $\bar{N} N \rightarrow \pi \pi$
~ 2040		⁸ OAKDEN	94	RVUE 0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
~ 1990		⁹ OAKDEN	94	RVUE 0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
1978± 5		¹⁰ ALPER	80	CNTR 62 $\pi^- p \rightarrow K^+ K^- n$
2040±10		¹⁰ ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p \bar{p} n$
1935±13		¹⁰ CORDEN	79	OMEG 12-15 $\pi^- p \rightarrow n 2 \pi$
1988± 7		EVANGELIS...	79B	OMEG 10 $\pi^- p \rightarrow K^+ K^- n$
1922±14		¹¹ ANTIPOV	77	CIBS 25 $\pi^- p \rightarrow p 3 \pi$


OCCUR=2

¹ From the first PWA solution.² From a partial-wave analysis of the data.³ From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2 \pi^0$.⁴ K matrix pole.⁵ Taking into account the $f_2(1950)$. Helicity-2 production favored.⁶ Energy-dependent analysis.⁷ Single energy analysis.⁸ From solution A of amplitude analysis of data on $\bar{p} p \rightarrow \pi \pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.⁹ From solution B of amplitude analysis of data on $\bar{p} p \rightarrow \pi \pi$. See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.¹⁰ $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.¹¹ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M016M;LINKAGE=BI
 NODE=M016M;LINKAGE=N
 NODE=M016M;LINKAGE=NN
 NODE=M016M;LINKAGE=KM
 NODE=M016M;LINKAGE=UE
 NODE=M016M;LINKAGE=RB
 NODE=M016M;LINKAGE=BR
 NODE=M016M;LINKAGE=B

NODE=M016M;LINKAGE=BB

NODE=M016M;LINKAGE=M
 NODE=M016M;LINKAGE=T



$f_4(2050)$ WIDTH

NODE=M016W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
237± 18 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.		
290± 20		AMELIN	06	VES 36 $\pi^- p \rightarrow \omega \omega n$
340± 80		12 BINON	05	GAMS 33 $\pi^- p \rightarrow \eta \eta n$
395± 40		ALDE	98	GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
170± 60		ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega \omega n$
304± 60		AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
210± 63		BALTRUSAIT..	87	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
400±100		ALDE	86D	GAM4 100 $\pi^- p \rightarrow n 2 \eta$
240± 40	40k	13 BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2 \pi^0$
190± 14		DENNEY	83	LASS 10 $\pi^+ n/\pi^+ p$
186 ⁺¹⁰³ ₋₅₈		14 CASON	82	STRC 8 $\pi^+ p \rightarrow \Delta^{++} \pi^0 \pi^0$
305 ⁺³⁶ ₋₁₁₉		ETKIN	82B	MPS 23 $\pi^- p \rightarrow n 2 K_S^0$
180± 60	700	APEL	75	NICE 40 $\pi^- p \rightarrow n 2 \pi^0$
225 ⁺¹²⁰ ₋₇₀		BLUM	75	ASPK 18.4 $\pi^- p \rightarrow n K^+ K^-$

NODE=M016W

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 260± 40 15 ANISOVICH 09 RVUE 0.0 $\bar{p} p, \pi N$
- 453± 20⁺³¹₋₁₂₉ 16 UEHARA 09 BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
- 182± 7 ANISOVICH 00J SPEC 2.0 $\bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0,$
 $\eta \eta, \eta \eta', \pi \pi$
- ~ 170 17 MARTIN 98 RVUE $N \bar{N} \rightarrow \pi \pi$
- ~ 200 18 MARTIN 97 RVUE $\bar{N} N \rightarrow \pi \pi$
- ~ 60 19 OAKDEN 94 RVUE 0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
- ~ 80 20 OAKDEN 94 RVUE 0.36-1.55 $\bar{p} p \rightarrow \pi \pi$
- 243± 16 21 ALPER 80 CNTR 62 $\pi^- p \rightarrow K^+ K^- n$
- 140± 15 21 ROZANSKA 80 SPRK 18 $\pi^- p \rightarrow p \bar{p} n$
- 263± 57 21 CORDEN 79 OMEG 12-15 $\pi^- p \rightarrow n 2 \pi$
- 100± 28 EVANGELIS... 79B OMEG 10 $\pi^- p \rightarrow K^+ K^- n$
- 107± 56 22 ANTIPOV 77 CIBS 25 $\pi^- p \rightarrow p 3 \pi$

OCCUR=2

12 From the first PWA solution.
 13 From a partial-wave analysis of the data.
 14 From an amplitude analysis of the reaction $\pi^+ \pi^- \rightarrow 2 \pi^0$.
 15 K matrix pole.
 16 Taking into account the $f_2(1950)$. Helicity-2 production favored.
 17 Energy-dependent analysis.
 18 Single energy analysis.


NODE=M016W;LINKAGE=BI
 NODE=M016W;LINKAGE=N
 NODE=M016W;LINKAGE=NN
 NODE=M016W;LINKAGE=KM
 NODE=M016W;LINKAGE=UE
 NODE=M016W;LINKAGE=RB
 NODE=M016W;LINKAGE=BR

- 19 From solution A of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
- 20 From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$. See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
- 21 $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.
- 22 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

NODE=M016W;LINKAGE=BW

NODE=M016W;LINKAGE=BB

NODE=M016W;LINKAGE=M
NODE=M016W;LINKAGE=T



$f_4(2050)$ DECAY MODES

NODE=M016215;NODE=M016

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega\omega$	seen
Γ_2 $\pi\pi$	(17.0±1.5) %
Γ_3 $K\bar{K}$	(6.8 ^{+3.4} _{-1.8}) × 10 ⁻³
Γ_4 $\eta\eta$	(2.1±0.8) × 10 ⁻³
Γ_5 $4\pi^0$	< 1.2 %
Γ_6 $\gamma\gamma$	seen
Γ_7 $a_2(1320)\pi$	seen

DESIG=6
DESIG=1
DESIG=2
DESIG=3
DESIG=5
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=7

$f_4(2050)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M016220

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_3\Gamma_6/\Gamma$**

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.29	95	ALTHOFF	85B TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

NODE=M016G2
NODE=M016G2

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ **$\Gamma_2\Gamma_6/\Gamma$**

VALUE (eV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
23.1 ^{+3.6} _{-3.3} + 70.5			23 UEHARA	09 BELL	10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
<1100	95	13 ± 4	OEST	90 JADE	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$

NODE=M016G3
NODE=M016G3

²³ Taking into account the $f_2(1950)$. Helicity-2 production favored.

NODE=M016G3;LINKAGE=UE

$f_4(2050)$ BRANCHING RATIOS

NODE=M016225

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$ **Γ_1/Γ**

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AMELIN	06 VES	36 $\pi^-p \rightarrow \omega\omega n$
••• We do not use the following data for averages, fits, limits, etc. •••			
not seen	BARBERIS	00F	450 $pp \rightarrow p_f\omega\omega p_s$

NODE=M016R7
NODE=M016R7

$\Gamma(\omega\omega)/\Gamma(\pi\pi)$				Γ_1/Γ_2	
VALUE	DOCUMENT ID	TECN	COMMENT		
1.5±0.3	ALDE	90	GAM2 38 $\pi^- p \rightarrow \omega\omega n$		NODE=M016R5 NODE=M016R5
$\Gamma(\pi\pi)/\Gamma_{total}$				Γ_2/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.170±0.015 OUR AVERAGE					NODE=M016R1 NODE=M016R1
0.18 ±0.03	24 BINON	83C	GAM2 38 $\pi^- p \rightarrow n4\gamma$		
0.16 ±0.03	24 CASON	82	STRC 8 $\pi^+ p \rightarrow \Delta^{++}\pi^0\pi^0$		
0.17 ±0.02	24 CORDEN	79	OMEG 12-15 $\pi^- p \rightarrow n2\pi$		
²⁴ Assuming one pion exchange.					NODE=M016R1;LINKAGE=A
$\Gamma(K\bar{K})/\Gamma(\pi\pi)$				Γ_3/Γ_2	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.04^{+0.02}_{-0.01}	ETKIN	82B	MPS 23 $\pi^- p \rightarrow n2K_S^0$		NODE=M016R2 NODE=M016R2
$\Gamma(\eta\eta)/\Gamma_{total}$				Γ_4/Γ	
VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT		
2.1±0.8	ALDE	86D	GAM4 100 $\pi^- p \rightarrow n4\gamma$		NODE=M016R3 NODE=M016R3
$\Gamma(4\pi^0)/\Gamma_{total}$				Γ_5/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
<0.012	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$		NODE=M016R4 NODE=M016R4
$\Gamma(a_2(1320)\pi)/\Gamma_{total}$				Γ_7/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
seen	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta\pi^+\pi^- n$		NODE=M016R6 NODE=M016R6

f₄(2050) REFERENCES

					NODE=M016
ANISOVICH	09	IJMP A24 2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)	REFID=52719
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
AMELIN	06	PAN 69 690	D.V. Amelin <i>et al.</i>	(VES Collab.)	REFID=51574
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)	REFID=47432
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47962
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
MARTIN	98	PR C57 3492	B.R. Martin <i>et al.</i>		REFID=46373
MARTIN	97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)	REFID=45685
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALDE	90	PL B241 600	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP+)	REFID=40935
OEST	90	ZPHY C47 343	T. Oest <i>et al.</i>	(JADE Collab.)	REFID=41358
ALDE	87	PL B198 286	D.M. Alde <i>et al.</i>	(LANL, BRUX, SERP, LAPP)	REFID=40221
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BINON	84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP)	REFID=21780
BINON	83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+)	REFID=40288
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
CASON	82	PRL 48 1316	N.M. Cason <i>et al.</i>	(NDAM, ANL)	REFID=20746
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)	REFID=20390
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=21665
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)	REFID=21774
CORDEN	79	NP B157 250	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)	REFID=20374
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21967
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
APEL	75	PL 57B 398	W.D. Apel <i>et al.</i>	(KARLK, KARLE, PISA, SERP+)	REFID=20720
BLUM	75	PL 57B 403	W. Blum <i>et al.</i>	(CERN, MPIM) JP	REFID=21651

$\pi_2(2100)$

$$I^G(J^{PC}) = 1^-(2^-+)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M020

NODE=M020

NODE=M020M

NODE=M020M

NODE=M020M;LINKAGE=AX
NODE=M020M;LINKAGE=L

NODE=M020W

NODE=M020W
NEWNODE=M020W;LINKAGE=AX
NODE=M020W;LINKAGE=L

NODE=M020215;NODE=M020

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EST;→ UNCHECKED ←
DESIG=4;OUR EST;→ UNCHECKED ←

NODE=M020220

NODE=M020R1
NODE=M020R1NODE=M020R2
NODE=M020R2NODE=M020R3
NODE=M020R3NODE=M020R4
NODE=M020R4

NODE=M020R;LINKAGE=L

NODE=M020

REFID=44433
REFID=20872 $\pi_2(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2090± 29 OUR AVERAGE			
2090± 30	¹ AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
2100± 150	² DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$

¹ From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, (\pi\pi)_S\pi$ waves.
² From a two-resonance fit to four 2^-0^+ waves.

 $\pi_2(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
620± 50 OUR AVERAGE	Error includes scale factor of 1.2. [625 ± 50 MeV OUR 2023 AVERAGE Scale factor = 1.2]		
520± 100	³ AMELIN	95B VES	36 $\pi^- A \rightarrow \pi^+ \pi^- \pi^- A$
651± 50	⁴ DAUM	81B CNTR	63,94 $\pi^- p \rightarrow 3\pi X$

³ From a fit to $J^{PC} = 2^-+ f_2(1270)\pi, (\pi\pi)_S\pi$ waves.
⁴ From a two-resonance fit to four 2^-0^+ waves.

 $\pi_2(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $\rho\pi$	seen
Γ_3 $f_2(1270)\pi$	seen
Γ_4 $(\pi\pi)_S\pi$	seen

 $\pi_2(2100)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.19±0.05	⁵ DAUM	81B CNTR	63,94 $\pi^- p$	
$\Gamma(f_2(1270)\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.36±0.09	⁵ DAUM	81B CNTR	63,94 $\pi^- p$	
$\Gamma((\pi\pi)_S\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
0.45±0.07	⁵ DAUM	81B CNTR	63,94 $\pi^- p$	
D-wave/S-wave RATIO FOR $\pi_2(2100) \rightarrow f_2(1270)\pi$	DOCUMENT ID	TECN	COMMENT	
0.39±0.23	⁵ DAUM	81B CNTR	63,94 $\pi^- p$	

⁵ From a two-resonance fit to four 2^-0^+ waves.

 $\pi_2(2100)$ REFERENCES

AMELIN	95B	PL B356 595	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
DAUM	81B	NP B182 269	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

$f_0(2100)$

$$J^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M168

NODE=M168

NODE=M168M

NODE=M168M

 $f_0(2100)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2095⁺¹⁷₋₁₉ OUR AVERAGE				
2116±27±17		LEES	21A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
2081±13 ⁺²⁴ ₋₃₆	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
2090±30		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2075±20		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
2090±10±6	529	^{2,3} DOBBS	15	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2099±17±8	283	^{2,3} DOBBS	15	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$
2105±8	80k	⁴ UMAN	06 E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
2102±13		⁵ ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0, \eta \eta, \eta \eta', \pi^+ \pi^-$
2105±10		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p} p \rightarrow \eta \eta, \eta \eta'$
~ 2104		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
~ 2122		HASAN	94 RVUE	$\bar{p} p \rightarrow \pi \pi$

OCCUR=2

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 209$ MeV.

⁴ Statistical error only.

⁵ Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

NODE=M168M;LINKAGE=A

NODE=M168M;LINKAGE=B

NODE=M168M;LINKAGE=C

NODE=M168M;LINKAGE=ST

NODE=M168M;LINKAGE=AN

 $f_0(2100)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
287⁺³²₋₂₄ OUR AVERAGE				
289±34±15		LEES	21A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' \pi^+ \pi^-$
273 ⁺²⁷ ₋₂₄ ±70±23	5.5k	¹ ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
330±100		BAI	00A BES	$J/\psi \rightarrow \gamma(\pi^+ \pi^- \pi^+ \pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
260±25		SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
236±14	80k	² UMAN	06 E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
211±29		³ ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \eta \pi^0 \pi^0, \pi^0 \pi^0, \eta \eta, \eta \eta', \pi^+ \pi^-$
200±25		ANISOVICH	99K SPEC	$0.6-1.94 \bar{p} p \rightarrow \eta \eta, \eta \eta'$
~ 203		BUGG	95	$J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
~ 273		HASAN	94 RVUE	$\bar{p} p \rightarrow \pi \pi$

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

² Statistical error only.

³ Includes the data of ANISOVICH 00B indicating to exotic decay pattern.

NODE=M168W

NODE=M168W

NODE=M168W;LINKAGE=A

NODE=M168W;LINKAGE=ST

NODE=M168W;LINKAGE=AN

 $f_0(2100)$ REFERENCES

LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61442
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47426
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103

NODE=M168

$f_2(2150)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called T_0 .

NODE=M042

NODE=M042

NODE=M042205

NODE=M042M

NODE=M042M

 $f_2(2150)$ MASS **$f_2(2150)$ MASS, COMBINED MODES (MeV)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

2157±12 OUR AVERAGE Includes data from the datablock that follows this one.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2170±6 80k ¹UMAN 06 E835 5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$ ¹Statistical error only.

NODE=M042M;LINKAGE=ST

 $\eta\eta$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

NODE=M042M3

NODE=M042M3

2157±12 OUR AVERAGE

2151±16	BARBERIS	00E	450 $pp \rightarrow p_f\eta\eta p_S$
2175±20	PROKOSHKIN	95D	GAM4 300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $pp \rightarrow pp 2\eta$
2130±35	SINGOVSKI	94	GAM4 450 $pp \rightarrow pp 2\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2140±30 ²ABELE 99B CBAR 1.94 $\bar{p}p \rightarrow \pi^0\eta\eta$ 2104±20 ³ARMSTRONG 93C E760 $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$ ²Spin not determined.³No J^{PC} determination.

NODE=M042M3;LINKAGE=K3

NODE=M042M3;LINKAGE=A

 $\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2135±20±45 ⁴ADOMEIT 96 CBAR 0 1.94 $\bar{p}p \rightarrow \eta 3\pi^0$ ⁴ANISOVICH 00E recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

NODE=M042M4;LINKAGE=AD

 $\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2090 ⁵OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ ~ 2120 ⁶OAKDEN 94 RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$ ~ 2170 ⁷MARTIN 80B RVUE~ 2150 ⁷MARTIN 80C RVUE~ 2150 ⁸DULUDE 78B OSPK 1–2 $\bar{p}p \rightarrow \pi^0\pi^0$

NODE=M042M1

NODE=M042M1

OCCUR=2

NODE=M042M1;LINKAGE=B

⁵OAKDEN 94 makes an amplitude analysis of LEAR data on $\bar{p}p \rightarrow \pi\pi$ using a method based on Barrelet zeros. This is solution A. The amplitude analysis of HASAN 94 includes earlier data as well, and assume that the data can be parametrized in terms of towers of nearly degenerate resonances on the leading Regge trajectory. See also KLOET 96 and MARTIN 97 who make related analyses.⁶From solution B of amplitude analysis of data on $\bar{p}p \rightarrow \pi\pi$.⁷ $I(J^P) = 0(2^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.⁸ $I^G(J^P) = 0^+(2^+)$ from partial-wave amplitude analysis.

NODE=M042M1;LINKAGE=BB

NODE=M042M1;LINKAGE=P

NODE=M042M1;LINKAGE=L

S-CHANNEL $\bar{p}p$, $\bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2139⁺⁸₋₉ ⁹EVANGELIS... 97 SPEC 0.6–2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$ ~ 2190 ⁹CUTTS 78B CNTR 0.97–3 $\bar{p}p \rightarrow \bar{N}N$ 2155±15 ^{9,10}COUPLAND 77 CNTR 0 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$ 2193±2 ^{9,11}ALSPECTOR 73 CNTR $\bar{p}p$ S channel⁹Isospins 0 and 1 not separated.¹⁰From a fit to the total elastic cross section.¹¹Referred to as T or T region by ALSPECTOR 73.


NODE=M042M2

NODE=M042M2

NODE=M042M2;LINKAGE=I

NODE=M042M2;LINKAGE=E

NODE=M042M2;LINKAGE=M



ηππ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
250±25±45	¹⁵ ADOMEIT 96	CBAR	0	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$
	¹⁵ ANISOVICH 00E			recommends to withdraw ADOMEIT 96 that assumed a single $J^P = 2^+$ resonance.

NODE=M042W4
NODE=M042W4

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
250 OUR ESTIMATE			
~ 70	¹⁶ OAKDEN 94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	¹⁷ MARTIN 80B	RVUE	
~ 250	¹⁷ MARTIN 80C	RVUE	
~ 250	¹⁸ DULUDE 78B	OSPK	1-2 $\bar{p}p \rightarrow \pi^0\pi^0$
	¹⁶ See however KLOET 96		who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.
	¹⁷ $I(J^P) = 0(2^+)$		from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.
	¹⁸ $I^G(J^P) = 0^+(2^+)$		from partial-wave amplitude analysis.

NODE=M042W4;LINKAGE=AD

NODE=M042W1
NODE=M042W1
→ UNCHECKED ←

S-CHANNEL $\bar{p}p, \bar{N}N$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
56 ⁺³¹ ₋₁₆	¹⁹ EVANGELIS... 97	SPEC		0.6-2.4 $\bar{p}p \rightarrow K_S^0 K_S^0$
135±75	^{20,21} COUPLAND 77	CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
98± 8	²¹ ALSPECTOR 73	CNTR		$\bar{p}p$ S channel
	¹⁹ Isospin 0 and 2 not separated.			
	²⁰ From a fit to the total elastic cross section.			
	²¹ Isospins 0 and 1 not separated.			

NODE=M042W1;LINKAGE=CC

NODE=M042W1;LINKAGE=P
NODE=M042W1;LINKAGE=L

NODE=M042W2
NODE=M042W2

NODE=M042W2;LINKAGE=F
NODE=M042W2;LINKAGE=E
NODE=M042W2;LINKAGE=I

K \bar{K} MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
91±62	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
150±30	ABLIKIM 04E	BES2	$J/\psi \rightarrow \omega K^+ K^-$
270±50	BARBERIS 99	OMEG	450 $p\bar{p} \rightarrow p_S p_f K^+ K^-$

NODE=M042W5
NODE=M042W5

$f_2(2150)$ DECAY MODES

NODE=M042215;NODE=M042

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	
Γ_2 $\eta\eta$	seen
Γ_3 $K\bar{K}$	seen
Γ_4 $f_2(1270)\eta$	seen
Γ_5 $a_2(1320)\pi$	seen
Γ_6 $\rho\bar{\rho}$	seen

DESIG=1

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=5;OUR EST;→ UNCHECKED ←

DESIG=6

 $f_2(2150)$ BRANCHING RATIOS

NODE=M042220

 $\Gamma(K\bar{K})/\Gamma(\eta\eta)$ Γ_3/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
1.28±0.23		BARBERIS	00E	450 $p\bar{p} \rightarrow p_f \eta \eta p_s$

NODE=M042R1

NODE=M042R1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<0.1	95	²² PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\rho 2\eta$
------	----	------------------------------	------	---

²² Using data from ARMSTRONG 89D.

NODE=M042R1;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(\eta\eta)$ Γ_1/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.33	95	²³ PROKOSHKIN 95D	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$, 450 $p\bar{p} \rightarrow p\rho 2\eta$

NODE=M042R2

NODE=M042R2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

²³ Derived from a $\pi^0\pi^0/\eta\eta$ limit.

NODE=M042R2;LINKAGE=A

 $\Gamma(f_2(1270)\eta)/\Gamma(a_2(1320)\pi)$ Γ_4/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
0.79±0.11	²⁴ ADOMEIT 96	CBAR	1.94 $\bar{p}p \rightarrow \eta 3\pi^0$

NODE=M042R3

NODE=M042R3

²⁴ Using $B(a_2(1320) \rightarrow \eta\pi) = 0.145$

NODE=M042R3;LINKAGE=A

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	73	ALEXANDER 10	CLEO	$\psi(2S) \rightarrow \gamma\rho\bar{\rho}$

NODE=M042R04

NODE=M042R04

 $f_2(2150)$ REFERENCES

NODE=M042

ALEXANDER 10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
UMAN 06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK... 06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
	Translated from YAF 69 515.			
ABLIKIM 04E	PL B603 138	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50174
ANISOVICH 00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
BARBERIS 00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)	REFID=47961
ABELE 99B	EPJ C8 67	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=46904
BARBERIS 99	PL B453 305	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46921
EVANGELIS... 97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)	REFID=45687
MARTIN 97	PR C56 1114	B.R. Martin, G.C. Oades	(LOUC, AARH)	REFID=45685
ADOMEIT 96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
KLOET 96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
PROKOSHKIN 95D	PD 40 495	Y.D. Prokoshkin	(SERP) IGJPC	REFID=44647
	Translated from DANS 344 469.			
HASAN 94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN 94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
SINGOVSKI 94	NC A107 1911	A.V. Singovsky	(SERP)	REFID=44648
ARMSTRONG 93C	PL B307 394	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43587
ARMSTRONG 89D	PL B227 186	T.A. Armstrong, M. Benayoun	(ATHU, BARI, BIRM+)	REFID=41010
MARTIN 80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP	REFID=21838
MARTIN 80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP	REFID=21837
CUTTS 78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)	REFID=21733
DULUDE 78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP	REFID=21850
COUPLAND 77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)	REFID=21830
ALSPECTOR 73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)	REFID=21813

$\rho(2150)$

$$I^G(J^{PC}) = 1^+(1^-)$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $T_1(2190)$. See the review on "Spectroscopy of Light Meson Resonances."

NODE=M032

NODE=M032

 $\rho(2150)$ MASS

NODE=M032205

 e^+e^- PRODUCED

NODE=M032M3

NODE=M032M3

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2044 ± 31 ± 4		¹ ABLIKIM	23BQ BES3	$e^+e^- \rightarrow a_2(1320)^+ \pi^- +$ c.c. $\rightarrow \eta \pi^+ \pi^-$
2095 ± 4		² ACHASOV	23A SND	$e^+e^- \rightarrow \omega \pi^0$
2034 ± 13 ± 9		³ ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega \pi^0$
2111 ± 43 ± 25		⁴ ABLIKIM	21X BES3	$e^+e^- \rightarrow \eta' \pi^+ \pi^-$
2255 $\begin{smallmatrix} +17 \\ -18 \end{smallmatrix}$ $\begin{smallmatrix} +50 \\ -41 \end{smallmatrix}$	1.8k	⁵ ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
2201 ± 19		⁶ LEES	20 BABR	$e^+e^- \rightarrow K^+ K^- \gamma$
2227 ± 9 ± 9		⁷ LEES	20 RVUE	$e^+e^- \rightarrow K^+ K^-$
2039 ± 8 $\begin{smallmatrix} +36 \\ -18 \end{smallmatrix}$		⁸ ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
2239.2 ± 7.1 ± 11.3		⁹ ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+ K^-$
2254 ± 22		¹⁰ LEES	12G BABR	$e^+e^- \rightarrow \pi^+ \pi^- \gamma$
2150 ± 40 ± 50		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow$ $f_1(1285) \pi^+ \pi^- \gamma$
1990 ± 80		AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \eta' \pi^+ \pi^- \gamma$
2153 ± 37		BIAGINI	91 RVUE	$e^+e^- \rightarrow \pi^+ \pi^-, K^+ K^-$
2110 ± 50		¹¹ CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+ \pi^-),$ $2(\pi^+ \pi^- \pi^0)$

OCCUR=2

OCCUR=2

¹ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of a Breit-Wigner resonance and a non-resonant contribution. Could be another state.

NODE=M032M3;LINKAGE=J

² From a vector dominance fit to the Born cross section between 1.05 and 2.0 GeV with $\rho(770)$, $\rho(1570)$, $\rho(1700)$, $\rho(2150)$. The fit also uses SND data from the VEPP-2M collider below 1.02 GeV and from LEES 17H and ABLIKIM 21A above 1.5 GeV.

NODE=M032M3;LINKAGE=K

³ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\rho(770)$, $\rho(1450)$ and $\rho(1700)$. Could be another state.

NODE=M032M3;LINKAGE=G

⁴ From a Breit-Wigner fit to the Born cross section, including an s -dependent continuum amplitude.

NODE=M032M3;LINKAGE=H

⁵ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X \eta \rightarrow K^+ K^- \eta = (21.7 \pm 1.9^{+7.7}_{-8.3}) \times 10^{-6}$.

NODE=M032M3;LINKAGE=F

⁶ From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is 3.5σ .

NODE=M032M3;LINKAGE=C

⁷ From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.

NODE=M032M3;LINKAGE=D

⁸ Could also be another state. Seen in J/ψ decay with branching ratio $J/\psi \rightarrow X \pi^0 \rightarrow K^+ K^- \pi^0 = (6.7 \pm 1.1^{+2.2}_{-1.8}) \times 10^{-6}$.

NODE=M032M3;LINKAGE=B

⁹ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

NODE=M032M3;LINKAGE=E

¹⁰ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

NODE=M032M3;LINKAGE=LE

¹¹ Includes ATKINSON 85.

NODE=M032M3;LINKAGE=A

 $\bar{p}p \rightarrow \pi\pi$

NODE=M032M1

NODE=M032M1

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
~ 2191	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2070	¹ OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2170	² MARTIN	80B	RVUE
~ 2100	² MARTIN	80C	RVUE

¹ See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

NODE=M032M1;LINKAGE=CC

² $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

NODE=M032M;LINKAGE=P

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2110 ± 35	¹ ANISOVICH 02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
~ 2190	² CUTTS 78B	CNTR	0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{2,3} COUPLAND 77	CNTR	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{2,4} ALSPECTOR 73	CNTR	$\bar{p}p$ S channel
2190 ± 10	⁵ ABRAMS 70	CNTR	S channel $\bar{p}N$

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

² Isospins 0 and 1 not separated.

³ From a fit to the total elastic cross section.

⁴ Referred to as T or T region by ALSPECTOR 73.

⁵ Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M032M2
NODE=M032M2

NODE=M032M;LINKAGE=AY

NODE=M032M;LINKAGE=I
NODE=M032M;LINKAGE=E
NODE=M032M;LINKAGE=M
NODE=M032M;LINKAGE=B

 $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2140 ± 30	ALDE 95	GAM2	38 $\pi^- p \rightarrow \omega\pi^0 n$
2170 ± 30	ALDE 92C	GAM4	100 $\pi^- p \rightarrow \omega\pi^0 n$

NODE=M032M4
NODE=M032M4

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\rho(2150)$ WIDTH

NODE=M032210

 $e^+ e^-$ PRODUCED

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
163 ± 69 ± 24		¹ ABLIKIM 23BQ	BES3	$e^+ e^- \rightarrow a_2(1320)^+ \pi^- +$ c.c. $\rightarrow \eta\pi^+ \pi^-$
270 ± 3		² ACHASOV 23A	SND	$e^+ e^- \rightarrow \omega\pi^0$
234 ± 30 ± 25		³ ABLIKIM 21A	BES3	$e^+ e^- \rightarrow \omega\pi^0$
135 ± 34 ± 30		⁴ ABLIKIM 21X	BES3	$e^+ e^- \rightarrow \eta' \pi^+ \pi^-$
460 ⁺⁵⁴ ₋₄₈ ⁺¹⁶⁰ ₋₉₀	1.8k	⁵ ABLIKIM 20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
70 ± 38		⁶ LEES 20	BABR	$e^+ e^- \rightarrow K^+ K^- \gamma$
127 ± 14 ± 4		⁷ LEES 20	RVUE	$e^+ e^- \rightarrow K^+ K^-$
196 ± 23 ⁺²⁵ ₋₂₇		⁸ ABLIKIM 19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$
139.8 ± 12.3 ± 20.6		⁹ ABLIKIM 19L	BES3	$e^+ e^- \rightarrow K^+ K^-$
109 ± 76		¹⁰ LEES 12G	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
350 ± 40 ± 50		AUBERT 07AU	BABR	10.6 $e^+ e^- \rightarrow$ $f_1(1285) \pi^+ \pi^- \gamma$
310 ± 140		AUBERT 07AU	BABR	10.6 $e^+ e^- \rightarrow \eta' \pi^+ \pi^- \gamma$
389 ± 79		BIAGINI 91	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^-, K^+ K^-$
410 ± 100		¹¹ CLEGG 90	RVUE	$e^+ e^- \rightarrow 3(\pi^+ \pi^-),$ $2(\pi^+ \pi^- \pi^0)$

NODE=M032W3
NODE=M032W3

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of a Breit-Wigner resonance and a non-resonant contribution. Could be another state.

² From a vector dominance fit to the Born cross section between 1.05 and 2.0 GeV with $\rho(770)$, $\rho(1570)$, $\rho(1700)$, $\rho(2150)$. The fit also uses SND data from the VEPP-2M collider below 1.02 GeV and from LEES 17H and ABLIKIM 21A above 1.5 GeV.

³ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\rho(770)$, $\rho(1450)$ and $\rho(1700)$. Could be another state.

⁴ From a Breit-Wigner fit to the Born cross section, including an s -dependent continuum amplitude.

⁵ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (21.7 \pm 1.9 ^{+7.7} _{-8.3}) \times 10^{-6}$.

⁶ From the fit to the BABAR data of LEES 13Q assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution. The resonance significance is 3.5σ .

⁷ From the fit to the BABAR data of LEES 13Q and BESIII data of ABLIKIM 19L assuming a coherent sum of a single Breit-Wigner resonance and a nonresonant contribution.

⁸ Could also be another state. Seen in J/ψ decay with branching ratio $J/\psi \rightarrow X\pi^0 \rightarrow K^+ K^- \pi^0 = (6.7 \pm 1.1 ^{+2.2} _{-1.8}) \times 10^{-6}$.

⁹ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

¹⁰ Using the GOUNARIS 68 parametrization of the pion form factor leaving the masses and widths of the $\rho(1450)$, $\rho(1700)$, and $\rho(2150)$ resonances as free parameters of the fit.

¹¹ Includes ATKINSON 85.

NODE=M032W3;LINKAGE=J

NODE=M032W3;LINKAGE=K

NODE=M032W3;LINKAGE=G

NODE=M032W3;LINKAGE=H

NODE=M032W3;LINKAGE=F

NODE=M032W3;LINKAGE=C

NODE=M032W3;LINKAGE=D

NODE=M032W3;LINKAGE=B

NODE=M032W3;LINKAGE=E

NODE=M032W3;LINKAGE=LE

NODE=M032W3;LINKAGE=A

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 296	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 40	¹ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250	² MARTIN	80B	RVUE	
~ 200	² MARTIN	80C	RVUE	

¹See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

² $I(J^P) = 1(1^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

NODE=M032W1
NODE=M032W1

NODE=M032W1;LINKAGE=CC

NODE=M032W;LINKAGE=P

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

230±50	¹ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
135±75	^{2,3} COUPLAND	77	CNTR	0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
98±8	³ ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
~ 85	⁴ ABRAMS	70	CNTR	S channel $\bar{p}N$

¹From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

²From a fit to the total elastic cross section.

³Isospins 0 and 1 not separated.

⁴Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M032W2
NODE=M032W2

NODE=M032W;LINKAGE=AY

NODE=M032W;LINKAGE=E

NODE=M032W;LINKAGE=I

NODE=M032W;LINKAGE=B

 $\pi^-p \rightarrow \omega\pi^0n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

320±70	ALDE	95	GAM2	38 $\pi^-p \rightarrow \omega\pi^0n$
~ 300	ALDE	92C	GAM4	100 $\pi^-p \rightarrow \omega\pi^0n$

NODE=M032W4
NODE=M032W4

 $\rho(2150)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	
Γ_2 $\pi^+\pi^-$	seen
Γ_3 K^+K^-	seen
Γ_4 $3(\pi^+\pi^-)$	seen
Γ_5 $2(\pi^+\pi^-\pi^0)$	seen
Γ_6 $\eta'\pi^+\pi^-$	seen
Γ_7 $f_1(1285)\pi^+\pi^-$	seen
Γ_8 $\omega\pi^0$	seen
Γ_9 $\omega\pi^0\eta$	seen
Γ_{10} $a_2(1320)\pi$	
Γ_{11} $p\bar{p}$	

NODE=M032215;NODE=M032

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

DESIG=7;OUR EVAL;→ UNCHECKED ←

DESIG=8;OUR EVAL;→ UNCHECKED ←

DESIG=9;OUR EVAL;→ UNCHECKED ←

DESIG=12

DESIG=10

 $\rho(2150) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_6\Gamma_1/\Gamma$
------------	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

23.3±5.3±3.3	¹ ABLIKIM	21X	BES3	$e^+e^- \rightarrow \eta'\pi^+\pi^-$
--------------	----------------------	-----	------	--------------------------------------

¹From a Breit-Wigner fit to the Born cross section interfering constructively with the continuum. For destructive interference the value is $0.64 \pm 0.49 \pm 0.42$ eV.

NODE=M032220

NODE=M032R02

NODE=M032R02

NODE=M032R02;LINKAGE=A

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_8\Gamma_1/\Gamma$
------------	-------------	------	---------	---------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

34±11±16	ABLIKIM	21A	BES3	$e^+e^- \rightarrow \omega\pi^0$
----------	---------	-----	------	----------------------------------

NODE=M032R00

NODE=M032R00

$$\Gamma(a_2(1320)\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{10}\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

34.6 ± 17.1 ± 6.0	¹ ABLIKIM	23BQ BES3	$e^+e^- \rightarrow a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$
-------------------	----------------------	-----------	--

¹ From a solution of the fit to the cross section between 2.00 and 3.08 GeV using a coherent sum of a single Breit-Wigner resonance and a non-resonant contribution. $B(a_2(1320) \rightarrow \eta\pi) \times B(\eta \rightarrow \gamma\gamma)$ fixed to 0.057. Another solution with equal fit quality gives 137.1 ± 73.3 ± 2.1 eV.

NODE=M032R04
NODE=M032R04

NODE=M032R04;LINKAGE=A

$$\rho(2150) \Gamma(i)\Gamma(e^+e^-)/\Gamma^2(\text{total})$$

NODE=M032230

$$\Gamma(f_1(1285)\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma \times \Gamma_1/\Gamma$$

VALUE (units 10 ⁻⁷)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

3.1 ± 0.6 ± 0.5	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow f_1(1285)\pi^+\pi^-\gamma$
-----------------	---------------------	-----------	---

¹ Calculated by us from the reported value of cross section at the peak.

NODE=M032G01
NODE=M032G01

NODE=M032G01;LINKAGE=AU

$$\Gamma(\eta'\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma \times \Gamma_1/\Gamma$$

VALUE (units 10 ⁻⁸)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.9 ± 1.9	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \eta'\pi^+\pi^-\gamma$
-----------	---------------------	-----------	---

¹ Calculated by us from the reported value of cross section at the peak.

NODE=M032G02
NODE=M032G02

NODE=M032G02;LINKAGE=AU

$\rho(2150)$ REFERENCES

NODE=M032

ABLIKIM	23BQ	PR D108 L111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62510
ACHASOV	23A	PR D108 092012	M.N. Achasov <i>et al.</i>	(SND Collab.)	REFID=62428
ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61028
ABLIKIM	21X	PR D103 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61231
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
LEES	20	PR D101 012011	J.P. Lees <i>et al.</i>	(BESIII Collab.)	REFID=60211
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59612
LEES	17H	PR D96 092009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58311
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	12G	PR D86 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54299
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP	REFID=44371
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALDE	92C	ZPHY C54 553	D.M. Alde <i>et al.</i>	(BELG, SERP, KEK, LANL+)	REFID=41859
BIAGINI	91	NC 104A 363	M.E. Biagini <i>et al.</i>	(FRAS, PRAG)	REFID=41894
CLEGG	90	ZPHY C45 677	A.B. Clegg, A. Donnachie	(LANC, MCHS)	REFID=41355
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=22000
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP	REFID=21838
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP	REFID=21837
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)	REFID=21733
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)	REFID=21830
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)	REFID=21824
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)	REFID=21813
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)	REFID=21807
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)	REFID=21805
GOUNARIS	68	PRL 21 244	G.J. Gounaris, J.J. Sakurai		REFID=48054

$\phi(2170)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

See the review on "Spectroscopy of Light Meson Resonances."

NODE=M103

NODE=M103

NODE=M103M

NODE=M103M
NEW $\phi(2170)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2164 ± 6 OUR AVERAGE				
[2163 ± 7 MeV OUR 2023 AVERAGE Scale factor = 1.1]				
2178 ± 20 ± 5		¹ ABLIKIM	23AX BES3	$e^+e^- \rightarrow \phi\pi^+\pi^-$
2190 ± 19 ± 37		² ABLIKIM	22L BES3	$2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$
2176 ± 24 ± 3		³ ABLIKIM	21A BES3	$e^+e^- \rightarrow \omega\eta$
2163.5 ± 6.2 ± 3.0		⁴ ABLIKIM	21T BES3	$e^+e^- \rightarrow \phi\eta$
2177.5 ± 4.8 ± 19.5		⁵ ABLIKIM	20M BES3	$e^+e^- \rightarrow \eta'\phi$
2126.5 ± 16.8 ± 12.4		⁶ ABLIKIM	20S BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2215.7 ± 8.3		⁷ LICHARD	23 RVUE	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$
2169 ± 5 ± 6		⁸ ZHU	23A RVUE	$e^+e^- \rightarrow \eta\phi$
2273.7 ± 5.7 ± 19.3		⁹ ABLIKIM	21AP BES3	$e^+e^- \rightarrow K_S^0 K_L^0$
2135 ± 8 ± 9	95	ABLIKIM	19I BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$
2239.2 ± 7.1 ± 11.3		¹⁰ ABLIKIM	19L BES3	$e^+e^- \rightarrow K^+K^-$
2200 ± 6 ± 5	471	ABLIKIM	15H BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$
2180 ± 8 ± 8		^{11,12} LEES	12F BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
2079 ± 13 ⁺⁷⁹ / ₋₂₈	4.8k	¹³ SHEN	09 BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2186 ± 10 ± 6	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$
2125 ± 22 ± 10	483	AUBERT	08S BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$
2192 ± 14	116	¹⁴ AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
2169 ± 20	149	¹⁴ AUBERT	07AK BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
2175 ± 10 ± 15	201	^{12,15} AUBERT, BE	06D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

OCCUR=2

NODE=M103M;LINKAGE=J

NODE=M103M;LINKAGE=I

NODE=M103M;LINKAGE=F

NODE=M103M;LINKAGE=G

NODE=M103M;LINKAGE=D

NODE=M103M;LINKAGE=E

NODE=M103M;LINKAGE=K

NODE=M103M;LINKAGE=L

NODE=M103M;LINKAGE=H

NODE=M103M;LINKAGE=C

NODE=M103M;LINKAGE=A

NODE=M103M;LINKAGE=AB

NODE=M103M;LINKAGE=SH

NODE=M103M;LINKAGE=AU

NODE=M103M;LINKAGE=B

¹ From a fit to the e^+e^- cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a non-resonant contribution.

² By a simultaneous fit of the $K_2^*(1430)^+K^-$ and $K^*(892)^+K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

³ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\omega(1420)$ and $\omega(1650)/\phi(1680)$.

⁴ From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ($\phi(1680)$ and $\phi(2170)$) and a nonresonant term.

⁵ From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

⁶ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

⁷ From a VDM fit to ZHU 23 $\eta\phi\gamma$ data with two resonances, $\phi(1680)$, $\phi(2170)$, and a third resonance with mass 1850.7 ± 5.3 MeV and width 25 ± 35 MeV of 1.7σ statistical evidence.

⁸ From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. The statistical significance for $\phi(2170)$ is 7.2σ .

⁹ From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to $\rho(2150)$.

¹⁰ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

¹¹ Fit includes interference with the $\phi(1680)$.

¹² From the $\phi f_0(980)$ component.

¹³ From a fit with two incoherent Breit-Wigners.

¹⁴ From the $K^+K^-f_0(980)$ component.

¹⁵ Superseded by LEES 12F.

 $\phi(2170)$ WIDTH

NODE=M103W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M103W

106	$\begin{matrix} +24 \\ -18 \end{matrix}$	OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.	
------------	--	--------------------	---	--

NEW

[103	$\begin{matrix} +28 \\ -21 \end{matrix}$	MeV OUR 2023 AVERAGE	Scale factor = 2.2]	
------	--	----------------------	---------------------	--

140	± 36	± 16	1	ABLIKIM	23AX	BES3	$e^+e^- \rightarrow \phi\pi^+\pi^-$
191	± 28	± 60	2	ABLIKIM	22L	BES3	$2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$
89	± 50	± 5	3	ABLIKIM	21A	BES3	$e^+e^- \rightarrow \omega\eta$
31.1	$\begin{matrix} +21.1 \\ -11.6 \end{matrix}$	± 1.1	4	ABLIKIM	21T	BES3	$e^+e^- \rightarrow \phi\eta$
149.0	± 15.6	± 8.9	5	ABLIKIM	20M	BES3	$e^+e^- \rightarrow \eta'\phi$
106.9	± 32.1	± 28.1	6	ABLIKIM	20S	BES3	$e^+e^- \rightarrow K^+K^-\pi^0\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

35	± 23		7	LICHARD	23	RVUE	$e^+e^- \rightarrow \Upsilon(nS) \rightarrow \phi\eta\gamma$	
96	$\begin{matrix} +17 \\ -14 \end{matrix}$	± 9	8	ZHU	23A	RVUE	$e^+e^- \rightarrow \eta\phi$	
86	± 44	± 51	9	ABLIKIM	21AP	BES3	$e^+e^- \rightarrow K_S^0 K_L^0$	
104	± 24	± 12	95	ABLIKIM	19I	BES3	$e^+e^- \rightarrow \eta\phi f_0(980)$	
139.8	± 12.3	± 20.6	10	ABLIKIM	19L	BES3	$e^+e^- \rightarrow K^+K^-$	
104	± 15	± 15	471	ABLIKIM	15H	BES3	$J/\psi \rightarrow \eta\phi\pi^+\pi^-$	
77	± 15	± 10	11,12	LEES	12F	BABR	$10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$	
192	± 23	$\begin{matrix} +25 \\ -61 \end{matrix}$	4.8k	13	SHEN	09	BELL	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
65	± 23	± 17	52	ABLIKIM	08F	BES	$J/\psi \rightarrow \eta\phi f_0(980)$	
61	± 50	± 13	483	AUBERT	08S	BABR	$10.6 e^+e^- \rightarrow \phi\eta\gamma$	
71	± 21		116	14	AUBERT	07AK	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
102	± 27		149	14	AUBERT	07AK	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
58	± 16	± 20	201	12,15	AUBERT,BE	06D	BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

OCCUR=2

¹ From a fit to the e^+e^- cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a non-resonant contribution.

NODE=M103W;LINKAGE=J

² By a simultaneous fit of the $K_2^*(1430)^+K^-$ and $K^*(892)^+K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103W;LINKAGE=I

³ From a fit to the cross section between 2.00 and 3.08 GeV with a coherent sum of Breit-Wigner amplitudes, including contributions from $\omega(1420)$ and $\omega(1650)/\phi(1680)$.

NODE=M103W;LINKAGE=F

⁴ From a fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ($\phi(1680)$ and $\phi(2170)$) and a nonresonant term.

NODE=M103W;LINKAGE=G

⁵ From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

NODE=M103W;LINKAGE=D

⁶ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103W;LINKAGE=E

⁷ From a VDM fit to ZHU 23 $\eta\phi\gamma$ data with two resonances, $\phi(1680)$, $\phi(2170)$, and a third resonance with mass 1850.7 ± 5.3 MeV and width 25 ± 35 MeV of 1.7σ statistical evidence.

NODE=M103W;LINKAGE=K

⁸ From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. The statistical significance for $\phi(2170)$ is 7.2σ .

NODE=M103W;LINKAGE=L

⁹ From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to $\rho(2150)$.

NODE=M103W;LINKAGE=H

¹⁰ The observed structure can be due to both the $\phi(2170)$ and $\rho(2150)$.

NODE=M103W;LINKAGE=C

¹¹ Fit includes interference with the $\phi(1680)$.

NODE=M103W;LINKAGE=A

¹² From the $\phi f_0(980)$ component.

NODE=M103W;LINKAGE=AB

¹³ From a fit with two incoherent Breit-Wigners.


NODE=M103W;LINKAGE=SH

¹⁴ From the $K^+K^-f_0(980)$ component.

NODE=M103W;LINKAGE=AU

¹⁵ Superseded by LEES 12F.

NODE=M103W;LINKAGE=B



$\phi(2170)$ DECAY MODES

NODE=M103215;NODE=M103

Mode	Fraction (Γ_i/Γ)
Γ_1 $e^+ e^-$	seen
Γ_2 $\phi\eta$	seen
Γ_3 $\omega\eta$	seen
Γ_4 $\phi\eta'$	seen
Γ_5 $\phi\pi\pi$	seen
Γ_6 $\phi f_0(980)$	seen
Γ_7 $K_S^0 K_L^0$	
Γ_8 $K^+ K^- \pi^+ \pi^-$	
Γ_9 $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-$	seen
Γ_{10} $K^+ K^- \pi^0 \pi^0$	
Γ_{11} $K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0$	seen
Γ_{12} $K^{*0} K^\pm \pi^\mp$	not seen
Γ_{13} $K^*(892)^0 \bar{K}^*(892)^0$	not seen
Γ_{14} $K^*(892)^+ K^*(892)^-$	
Γ_{15} $K^*(892)^+ K^- + c.c.$	
Γ_{16} $K(1460)^+ K^- + c.c.$	
Γ_{17} $K_1(1270)^+ K^- + c.c.$	
Γ_{18} $K_1(1400)^+ K^- + c.c.$	
Γ_{19} $K_2^*(1430)^+ K^- + c.c.$	

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=5;OUR EVAL;→ UNCHECKED ←
 DESIG=16;OUR EVAL;→ UNCHECKED ←
 DESIG=11;OUR EVAL;→ UNCHECKED ←
 DESIG=9
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=17
 DESIG=3
 DESIG=6
 DESIG=4
 DESIG=7
 DESIG=8
 DESIG=10
 DESIG=15
 DESIG=19
 DESIG=12
 DESIG=14
 DESIG=13
 DESIG=18

$\phi(2170) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M103230

$\Gamma(\phi\eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$						$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
0.17	90		1 ZHU	23	BELL $e^+ e^- \rightarrow \gamma(nS) \rightarrow \phi\eta\gamma$	
$0.36^{+0.05}_{-0.03} \pm 0.07$ to $41 \pm 2 \pm 6$			2 ZHU	23A	RVUE $e^+ e^- \rightarrow \eta\phi$	
$0.24^{+0.12}_{-0.07}$			3 ABLIKIM	21T	BES3 $e^+ e^- \rightarrow \phi\eta$	
$1.7 \pm 0.7 \pm 1.3$		483	AUBERT	08S	BABR $10.6 e^+ e^- \rightarrow \phi\eta\gamma$	

NODE=M103G2
 NODE=M103G2

¹ From a solution of the fit using a vector meson dominance model with contributions from $\phi(1680)$, $\phi(2170)$ and non resonant contribution with mass and width of $\phi(2170)$ fixed at 2163.5 MeV and 31.1 MeV respectively. Four solutions are found with equal fit quality giving 0.17 eV (solution I and II) and 18.6 eV (III and IV) at 90% CL.

NODE=M103G2;LINKAGE=B

² From the analysis of the combined measurements of $\sigma(e^+e^- \rightarrow \eta\phi)$ from BaBar, Belle, BESIII, CMD3. The statistical significance for $\phi(2170)$ is 7.2σ . Four solutions are found, with equal fit quality: $(0.56^{+0.03}_{-0.02} \pm 0.07)$ eV, $(0.36^{+0.05}_{-0.03} \pm 0.07)$ eV, $(38 \pm 1 \pm 5)$ eV, $(41 \pm 2 \pm 6)$ eV.

NODE=M103G2;LINKAGE=C

³ From a solution of the fit to the cross section below 3.5 GeV measured by BaBar and BESIII with a coherent sum of two modified Breit-Wigner amplitudes ($\phi(1680)$ and $\phi(2170)$) and a nonresonant term. The other solution gives $10.11^{+3.87}_{-3.13}$ eV.

NODE=M103G2;LINKAGE=A

$\Gamma(\omega\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
0.43±0.15±0.04	¹ ABLIKIM	21A	BES3 $e^+e^- \rightarrow \omega\eta$

NODE=M103R09
NODE=M103R09

¹ For constructive interference with $\omega(1420)$ and $\omega(1650)/\phi(1680)$. For destructive interference: $1.25 \pm 0.48 \pm 0.18$ eV.

NODE=M103R09;LINKAGE=A

$\Gamma(\phi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
7.1±0.7±0.7	¹ ABLIKIM	20M	BES3 $e^+e^- \rightarrow \eta'\phi$

NODE=M103R00
NODE=M103R00

¹ From a fit using a coherent sum of a phase-space modified Breit-Wigner function and a phase-space term.

NODE=M103R00;LINKAGE=A

$\Gamma(\phi f_0(980)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M103G1
NODE=M103G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.3 \pm 0.3 \pm 0.3$		^{1,2} LEES	12F	BABR $10.6 e^+e^- \rightarrow \phi\pi^+\pi^-\gamma$
$2.5 \pm 0.8 \pm 0.4$	201	^{2,3} AUBERT, BE	06D	BABR $10.6 e^+e^- \rightarrow K^+K^-\pi\pi\gamma$

¹ From a fit with constructive interference with the $\phi(1680)$. In a fit with destructive interference, the value is larger by a factor of 12.

NODE=M103G1;LINKAGE=A

² For $f_0(980) \rightarrow \pi\pi$.

³ Superseded by LEES 12F.

NODE=M103G1;LINKAGE=AB
NODE=M103G1;LINKAGE=B

$\Gamma(K_S^0 K_L^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M103R10
NODE=M103R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.9 \pm 0.6 \pm 0.7$		¹ ABLIKIM	21AP	BES3 $e^+e^- \rightarrow K_S^0 K_L^0$
-----------------------	--	----------------------	------	---------------------------------------

¹ From a fit to the cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a nonresonant contribution. The observed structure can be also due to $\rho(2150)$.

NODE=M103R10;LINKAGE=A

$\Gamma(K^*(892)^+ K^*(892)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{14}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	¹ ABLIKIM	20S	BES3 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0$

NODE=M103R08
NODE=M103R08

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103R08;LINKAGE=A

$\Gamma(K^*(892)^+ K^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{15}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M103R11
NODE=M103R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.0 ± 0.3		¹ ABLIKIM	22L	BES3 $2.0-3.08 e^+e^- \rightarrow K^+K^-\pi^0$
---------------	--	----------------------	-----	--

¹ From a solution of a simultaneous fit of the $K_2^*(1430)^+ K^-$ and $K^*(892)^+ K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. The other solution gives 7.1 ± 0.9 eV. Significance 3.7σ .

NODE=M103R11;LINKAGE=A

$\Gamma(K(1460)^+ K^- + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M103R05
NODE=M103R05

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.0 ± 3.8		¹ ABLIKIM	20S	BES3 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0$
---------------	--	----------------------	-----	--

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function.

NODE=M103R05;LINKAGE=A

$$\Gamma(K_1(1270)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{17} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<12.5	90	¹ ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

NODE=M103R06
NODE=M103R06

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives an upper limit value of 297.6 eV.

NODE=M103R06;LINKAGE=A

$$\Gamma(K_1(1400)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{18} \Gamma_1 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
4.7 ± 3.3	¹ ABLIKIM	20S BES3	$e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0$

NODE=M103R07
NODE=M103R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ By a simultaneous fit of the intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. A second solution of the fit with equal fit quality gives a value of 98.8 ± 7.8 eV.

NODE=M103R07;LINKAGE=A

$$\Gamma(K_2^*(1430)^+ K^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{19} \Gamma_1 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
12.6 ± 2.4	¹ ABLIKIM	22L BES3	2.0–3.08 $e^+ e^- \rightarrow K^+ K^- \pi^0$

NODE=M103R12
NODE=M103R12

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a solution of a simultaneous fit of the $K_2^*(1430)^+ K^-$ and $K^*(892)^+ K^-$ intermediate channels in a partial-wave analysis, assuming the same structure, modelled with a coherent sum of a nonresonant component and a resonant component by a Breit-Wigner function. The other solution gives 161.1 ± 20.6 eV.

NODE=M103R12;LINKAGE=A

$$\phi(2170) \Gamma(i) \Gamma(e^+ e^-) / \Gamma^2(\text{total})$$

NODE=M103220

$$\Gamma(\phi \pi \pi) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_5 / \Gamma \times \Gamma_1 / \Gamma$$

VALUE (units 10 ⁻⁷)	EVTs	DOCUMENT ID	TECN	COMMENT
1.65 ± 0.15 ± 0.18	4.8k	¹ SHEN	09 BELL	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

NODE=M103G01
NODE=M103G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Multiplied by 3/2 to take into account the $\phi \pi^0 \pi^0$ mode. Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.6)\%$.

NODE=M103G01;LINKAGE=SH

$\phi(2170)$ BRANCHING RATIOS

NODE=M103225

$$\Gamma(\phi \pi \pi) / \Gamma_{\text{total}} \quad \Gamma_5 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	23AX BES3	$e^+ e^- \rightarrow \phi \pi^+ \pi^-$

NODE=M103R13
NODE=M103R13

¹ From a fit to the $e^+ e^-$ cross section between 2.00 and 3.08 GeV with a sum of Breit-Wigner amplitude and a non-resonant contribution.

NODE=M103R13;LINKAGE=A

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_9 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

NODE=M103R01
NODE=M103R01

$$\Gamma(K^+ K^- f_0(980) \rightarrow K^+ K^- \pi^0 \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{11} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^0 \pi^0 \gamma$

NODE=M103R02
NODE=M103R02

$$\Gamma(K^{*0} K^\pm \pi^\mp) / \Gamma_{\text{total}} \quad \Gamma_{12} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	AUBERT	07AK BABR	10.6 GeV $e^+ e^-$

NODE=M103R03
NODE=M103R03

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) / \Gamma_{\text{total}} \quad \Gamma_{13} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	10C BES2	$J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$

NODE=M103R04
NODE=M103R04

$\phi(2170)$ REFERENCES

ABLIKIM	23AX	PR D108 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62350
LICHARD	23	PR D108 092005	P. Lichard	(OPAV, CTUP)	REFID=62426
ZHU	23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)	REFID=61911
ZHU	23A	CP C47 113003	W. Zhu, X. Wang	(RVUE)	REFID=62439
ABLIKIM	22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61649
ABLIKIM	21A	PL B813 136059	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61028
ABLIKIM	21AP	PR D104 092014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61448
ABLIKIM	21T	PR D104 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61154
ABLIKIM	20M	PR D102 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60337
ABLIKIM	20S	PRL 124 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60542
ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59605
ABLIKIM	19L	PR D99 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59612
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56773
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511

NODE=M103

 $f_0(2200)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in $K_S^0 K_S^0$ (AUGUSTIN 88), $K^+ K^-$ (ABLIKIM 05Q) and $\eta\eta$ (BINON 05) system. Not seen in $\Upsilon(1S)$ radiative decays (BARU 89).

NODE=M112

NODE=M112

 $f_0(2200)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2187±14 OUR AVERAGE				
2170±20 ⁺¹⁰ ₋₁₅		ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
2197±17		¹ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
2200±25		SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
2206±12± 8	381	^{2,3} DOBBS	15	$J/\psi \rightarrow \gamma K^+ K^-$
2188±17±16	203	^{2,3} DOBBS	15	$\psi(2S) \rightarrow \gamma K^+ K^-$
2210±50		⁴ BINON	05	GAMS $33 \pi^- p \rightarrow \eta\eta n$
~ 2122		HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2321		HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

NODE=M112M

NODE=M112M

¹ Cannot determine spin to be 0.² Using CLEO-c data but not authored by the CLEO Collaboration.³ From a fit to a Breit-Wigner line shape with fixed $\Gamma = 238$ MeV.⁴ First solution, PWA is ambiguous.

OCCUR=2

OCCUR=2

NODE=M112M;LINKAGE=A
 NODE=M112M;LINKAGE=B
 NODE=M112M;LINKAGE=C
 NODE=M112M;LINKAGE=BI

 $f_0(2200)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
210±40 OUR AVERAGE			
[207±40 MeV OUR 2023 AVERAGE]			
220±60 ⁺⁴⁰ ₋₄₅	ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
201±51	⁵ AUGUSTIN	88	DM2 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
••• We do not use the following data for averages, fits, limits, etc. •••			
150±30	SARANTSEV	21	RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
380±90	⁶ BINON	05	GAMS $33 \pi^- p \rightarrow \eta\eta n$
~ 273	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 223	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

NODE=M112W

NODE=M112W

NEW

⁵ Cannot determine spin to be 0.⁶ First solution, PWA is ambiguous.

OCCUR=2

NODE=M112W;LINKAGE=A
 NODE=M112W;LINKAGE=BI

 $f_0(2200)$ REFERENCES

SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
BARU	89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574

NODE=M112

$f_J(2220)$

$$I^G(J^{PC}) = 0^+(2^{++} \text{ or } 4^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation. See our mini-review in the 2004 edition of this Review, PDG 04.

NODE=M082

NODE=M082

 $f_J(2220)$ MASS

NODE=M082M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2231.1 ± 3.5 OUR AVERAGE				
2235 ± 4 ± 6	74	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
2230 $^{+6}_{-7}$ ±16	46	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
2232 $^{+8}_{-7}$ ±15	23	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
2235 ± 4 ± 5	32	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
2209 $^{+17}_{-15}$ ±10		ASTON	88F LASS	$11 K^-p \rightarrow K^+K^- \Lambda$
2230 ± 20		BOLONKIN	88 SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n$
2220 ± 10	41	¹ ALDE	86B GA24	$38-100 \pi p \rightarrow n\eta\eta'$
2230 ± 6 ± 14	93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
2232 ± 7 ± 7	23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0 K_S^0$

NODE=M082M

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

2223.9 ± 2.5		² VLADIMIRSK...08	SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n + m\pi^0$
2246 ± 36		BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ALDE 86B uses data from both the GAMS-2000 and GAMS-4000 detectors.² $J^{PC} = 2^{++}$. Systematic uncertainties not evaluatedNODE=M082M;LINKAGE=A
NODE=M082M;LINKAGE=VL $f_J(2220)$ WIDTH

NODE=M082W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
23 $^{+8}_{-7}$ OUR AVERAGE					
19 $^{+13}_{-11}$ ±12		74	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
20 $^{+20}_{-15}$ ±17		46	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+K^-$
20 $^{+25}_{-16}$ ±14		23	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
15 $^{+12}_{-9}$ ± 9		32	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
60 $^{+107}_{-57}$			ASTON	88F LASS	$11 K^-p \rightarrow K^+K^- \Lambda$
80 ± 30			BOLONKIN	88 SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n$
26 $^{+20}_{-16}$ ±17		93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
18 $^{+23}_{-15}$ ±10		23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0 K_S^0$

NODE=M082W

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.6 ± 2.5		¹ VLADIMIRSK...08	SPEC	$40 \pi^-p \rightarrow K_S^0 K_S^0 n + m\pi^0$
<80	90	ALDE	87C GAM2	$38 \pi^-p \rightarrow \eta'\eta n$

¹ $J^{PC} = 2^{++}$. Systematic uncertainties not evaluated

NODE=M082W;LINKAGE=VL

 $f_J(2220)$ DECAY MODES

NODE=M082215;NODE=M082

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	not seen
Γ_2 $\pi^+\pi^-$	not seen
Γ_3 $K\bar{K}$	not seen
Γ_4 $p\bar{p}$	not seen
Γ_5 $\gamma\gamma$	not seen
Γ_6 $\eta\eta'(958)$	seen
Γ_7 $\phi\phi$	not seen
Γ_8 $\eta\eta$	not seen

DESIG=5

DESIG=6;OUR EST;→ UNCHECKED ←

DESIG=1

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=7;OUR EST;→ UNCHECKED ←

DESIG=8;OUR EST;→ UNCHECKED ←

$f_J(2220) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M082220

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_5/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	95	¹ ACCIARRI	01H L3	$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}^{\text{ee}} = 91, 183-209 \text{ GeV}$

NODE=M082G1
NODE=M082G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.6	95	¹ GODANG	97 CLE2	$\gamma\gamma \rightarrow K_S^0 K_S^0$
< 86	95	¹ ALBRECHT	90G ARG	$\gamma\gamma \rightarrow K^+ K^-$
<1000	95	² ALTHOFF	85B TASS	$\gamma\gamma, K\bar{K}\pi$

 $\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	95	ALAM	98C CLE2	$\gamma\gamma \rightarrow \pi^+ \pi^-$

NODE=M082G3
NODE=M082G3¹ Assuming $J^P = 2^+$.² True for $J^P = 0^+$ and $J^P = 2^+$.NODE=M082G1;LINKAGE=D
NODE=M082G1;LINKAGE=C $f_J(2220) \Gamma(i)\Gamma(\rho\bar{\rho})/\Gamma^2(\text{total})$

NODE=M082223

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<18	95	¹ AMSLER	01 CBAR	$1.4-1.5 \rho\bar{\rho} \rightarrow \pi^0 \pi^0$

NODE=M082GG1
NODE=M082GG1

• • • We do not use the following data for averages, fits, limits, etc. • • •

<(11-42)	99	² HASAN	96 SPEC	$1.35-1.55 \rho\bar{\rho} \rightarrow \pi^+ \pi^-$
----------	----	--------------------	---------	--

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_7/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6	95	³ EVANGELIS...	98 SPEC	$1.1-2.0 \rho\bar{\rho} \rightarrow \phi\phi$

NODE=M082GG2
NODE=M082GG2 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<4	95	¹ AMSLER	01 CBAR	$1.4-1.5 \rho\bar{\rho} \rightarrow \eta\eta$

NODE=M082GG3
NODE=M082GG3¹ For $J^P = 2^+$ in the mass range 2222-2240 MeV and the total width between 10 and 20 MeV.² For $J^P = 2^+$ and $J^P = 4^+$ in the mass range 2220-2245 MeV and the total width of 15 MeV.³ For $J^P = 2^+$, the mass of 2235 MeV and the total width of 15 MeV.

NODE=M082GG;LINKAGE=A

NODE=M082GG;LINKAGE=B

NODE=M082GG;LINKAGE=C

 $f_J(2220) \text{ BRANCHING RATIOS}$

NODE=M082225

 $\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	COMMENT
not seen	¹ DOBBS 15	$J/\psi \rightarrow \gamma\pi\pi$
not seen	¹ DOBBS 15	$\psi(2S) \rightarrow \gamma\pi\pi$

NODE=M082R00
NODE=M082R00

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M082R00;LINKAGE=A

 $\Gamma(K\bar{K})/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	COMMENT
not seen	¹ DOBBS 15	$J/\psi \rightarrow \gamma K\bar{K}$
not seen	¹ DOBBS 15	$\psi(2S) \rightarrow \gamma K\bar{K}$

NODE=M082R01
NODE=M082R01

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M082R01;LINKAGE=A

 $\Gamma(\pi\pi)/\Gamma(K\bar{K})$ Γ_1/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
1.0±0.5	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma 2\pi, K\bar{K}$

NODE=M082R2
NODE=M082R2 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M082R1
NODE=M082R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		¹ AUBERT	07AV BABR	$B \rightarrow \rho\bar{\rho}K^*$
not seen		WANG	05A BELL	$B^+ \rightarrow \bar{\rho}\rho K^+$
<3.0	95	² EVANGELIS...	97 SPEC	$1.96-2.40 \rho\bar{\rho} \rightarrow K_S^0 K_S^0$
<1.1	99.7	³ BARNES	93 SPEC	$1.3-1.57 \rho\bar{\rho} \rightarrow K_S^0 K_S^0$
<2.6	99.7	³ BARDIN	87 CNTR	$1.3-1.5 \rho\bar{\rho} \rightarrow K^+ K^-$
<3.6	99.7	³ SCULLI	87 CNTR	$1.29-1.55 \rho\bar{\rho} \rightarrow K^+ K^-$

- ¹ Assuming $\Gamma < 30$ MeV.
² Assuming $\Gamma \sim 20$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.
³ Assuming $\Gamma = 30\text{-}35$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.

NODE=M082R1;LINKAGE=AU
 NODE=M082R1;LINKAGE=C
 NODE=M082R1;LINKAGE=B

 $\Gamma(p\bar{p})/\Gamma(K\bar{K})$ Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.09	BAI	96B	BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}, K\bar{K}$

NODE=M082R3
 NODE=M082R3

 $f_J(2220)$ REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
VLADIMIRSK...	08	PAN 71 2129	V.V. Vladimirovsky <i>et al.</i>	(ITEP)
		Translated from YAF 71 2166.		
AUBERT	07AV	PR D76 092004	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
AMSLER	01	PL B520 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ALAM	98C	PRL 81 3328	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
EVANGELIS...	98	PR D57 5370	C. Evangelista <i>et al.</i>	(JETSET Collab.)
EVANGELIS...	97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
GODANG	97	PRL 79 3829	R. Godang <i>et al.</i>	(CLEO Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASTON	88F	PL B215 199	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ALDE	87C	SJNP 45 255	D. Alde <i>et al.</i>	
		Translated from YAF 45 405.		
BARDIN	87	PL B195 292	G. Bardin <i>et al.</i>	(SACL, FERR, CERN, PADO+)
SCULLI	87	PRL 58 1715	J. Sculli <i>et al.</i>	(NYU, BNL)
ALDE	86B	PL B177 120	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)

NODE=M082

REFID=56805
 REFID=52681

REFID=51990
 REFID=50651
 REFID=49653
 REFID=48321
 REFID=48558
 REFID=46326
 REFID=46342
 REFID=46365
 REFID=45687
 REFID=45760
 REFID=44736
 REFID=45197
 REFID=43601
 REFID=41374
 REFID=40585
 REFID=40580
 REFID=47474

OTHER RELATED PAPERS

DEL-AMO-SA...	100	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
---------------	-----	----------------	----------------------------------	-----------------

REFID=53533

NODE=M270

 $\omega(2220)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

OMITTED FROM SUMMARY TABLE

 $\omega(2220)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2232±19±27	¹ ABLIKIM	23G	BES3 $2.0\text{-}3.1 e^+e^- \rightarrow \omega\pi\pi$
••• We do not use the following data for averages, fits, limits, etc. •••			
2250±25±27	² ABLIKIM	23G	BES3 $2.0\text{-}3.1 e^+e^- \rightarrow \omega\pi^+\pi^-$
2222±7±2	³ ABLIKIM	22I	BES3 $2.0\text{-}3.8 e^+e^- \rightarrow \omega\pi^0\pi^0$
2205±30	⁴ ANISOVICH	02B	SPEC $0.6\text{-}1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M270M

NODE=M270M

OCCUR=2

¹ From a fit to $\omega\pi^+\pi^-$ and $\omega\pi^0\pi^0$ with a Breit-Wigner resonance interfering with the continuum. Supersedes ABLIKIM 22I.

NODE=M270M;LINKAGE=A

² From a fit to $\omega\pi^+\pi^-$ with a Breit-Wigner resonance interfering with the continuum.

NODE=M270M;LINKAGE=B

³ From the fit to the cross section by the coherent sum of resonant component parametrized by a modified Breit-Wigner amplitude and a phase-space contribution for the continuum.

NODE=M270M;LINKAGE=C

⁴ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M270M;LINKAGE=D

 $\omega(2220)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
93±53±20	⁵ ABLIKIM	23G	BES3 $2.0\text{-}3.1 e^+e^- \rightarrow \omega\pi\pi$
••• We do not use the following data for averages, fits, limits, etc. •••			
125±43±15	⁶ ABLIKIM	23G	BES3 $2.0\text{-}3.1 e^+e^- \rightarrow \omega\pi^+\pi^-$
59±30±6	⁷ ABLIKIM	22I	BES3 $2.0\text{-}3.8 e^+e^- \rightarrow \omega\pi^0\pi^0$
350±90	⁸ ANISOVICH	02B	SPEC $0.6\text{-}1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M270W

NODE=M270W

OCCUR=2

⁵ From a fit to $\omega\pi^+\pi^-$ and $\omega\pi^0\pi^0$ with a Breit-Wigner resonance interfering with the continuum. Supersedes ABLIKIM 22I.

NODE=M270W;LINKAGE=A

⁶ From a fit to $\omega\pi^+\pi^-$ with a Breit-Wigner resonance interfering with the continuum.

NODE=M270W;LINKAGE=B

⁷ From the fit to the cross section by the coherent sum of resonant component parametrized by a modified Breit-Wigner amplitude and a phase-space contribution for the continuum.

NODE=M270W;LINKAGE=C

⁸ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M270W;LINKAGE=D

$\omega(2220)$ DECAY MODES

NODE=M270215;NODE=M270

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega \pi \pi$	seen
Γ_2 $\omega \pi^+ \pi^-$	seen
Γ_3 $\omega \pi^0 \pi^0$	seen
Γ_4 $e^+ e^-$	seen

DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EVAL;→ UNCHECKED ←

 $\omega(2220) \Gamma(i)\Gamma(e^+ e^-)/\Gamma(\text{total})$

NODE=M270235

$\Gamma(\omega \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_3 \Gamma_4/\Gamma$
VALUE (eV) DOCUMENT ID TECN COMMENT

NODE=M270G00
NODE=M270G00

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.3±0.1±0.1 ⁹ ABLIKIM 22I BES3 2.0–3.8 $e^+ e^- \rightarrow \omega \pi^0 \pi^0$

⁹ Superseded by ABLIKIM 23G.

NODE=M270G00;LINKAGE=A

$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_2 \Gamma_4/\Gamma$
VALUE (eV) DOCUMENT ID TECN COMMENT

NODE=M270R01
NODE=M270R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9±0.4±0.4 ¹⁰ ABLIKIM 23G BES3 2.0–3.1 $e^+ e^- \rightarrow \omega \pi^+ \pi^-$

¹⁰ From a fit to $\omega \pi^+ \pi^-$ with a Breit-Wigner resonance interfering with the continuum. Solution with constructive interference: 52.9 ± 17.0 ± 13.1 eV.

NODE=M270R01;LINKAGE=A

$\Gamma(\omega \pi \pi) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$ $\Gamma_1 \Gamma_4/\Gamma$
VALUE (eV) DOCUMENT ID TECN COMMENT

NODE=M270R00
NODE=M270R00

0.9±0.5±0.2 ¹¹ ABLIKIM 23G BES3 2.0–3.1 $e^+ e^- \rightarrow \omega \pi \pi$

¹¹ From a fit to $\omega \pi^+ \pi^-$ and $\omega \pi^0 \pi^0$ with a Breit-Wigner resonance interfering with the continuum. Solution with constructive interference: 61.1 ± 32.1 ± 15.4 eV. Supersedes ABLIKIM 22I.

NODE=M270R00;LINKAGE=A

 $\omega(2220)$ REFERENCES

NODE=M270

ABLIKIM	23G	JHEP 2301 111	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62049
Also		JHEP 2303 093 (errat.)	M. Ablikim, <i>et. al.</i>	(BESIII Collab.)	REFID=62054
ABLIKIM	22I	PR D105 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61644
ANISOVICH	02B	PL B542 19	A.V. Anisovich <i>et al.</i>		REFID=48829
ANISOVICH	01C	PL B507 23	A.V. Anisovich <i>et al.</i>		REFID=48325
ANISOVICH	00D	PL B476 15	A.V. Anisovich <i>et al.</i>		REFID=47944

NODE=M115

$\eta(2225)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Seen in $J/\psi \rightarrow \gamma \phi \phi$. Possibly seen in $B \rightarrow \phi \phi K$ by LEES 11A.

NODE=M115

 $\eta(2225)$ MASS

NODE=M115M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M115M

2221⁺¹³₋₁₀ OUR AVERAGE

2216⁺⁴⁺²¹₋₅₋₁₁ ¹ ABLIKIM 16N BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

2240⁺³⁰⁺³⁰₋₂₀₋₂₀ 196 ± 19 ABLIKIM 08I BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

2230 ± 25 ± 15 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

2214 ± 20 ± 13 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2220 BISELLO 86B DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^{-+} phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^{-+} state $X(2500)$ ($M = 2470_{-19}^{+15+101}_{-23}$ MeV, $\Gamma = 230_{-35}^{+64+56}_{-33}$ MeV).

NODE=M115M;LINKAGE=B

 $\eta(2225)$ WIDTH

NODE=M115W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M115W

185⁺₋₂₀ OUR AVERAGE

185 ⁺ ₋₁₄₋₁₇		¹ ABLIKIM	16N BES3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
190 [±] 30 ⁺⁶⁰ ₋₄₀	196 ± 19	ABLIKIM	08I BES	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
150 ⁺ ₋₆₀ ± 60		BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 80 BISELLO 86B DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^{-+} phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^{-+} state $X(2500)$ ($M = 2470_{-19}^{+15+101}_{-23}$ MeV, $\Gamma = 230_{-35}^{+64+56}_{-33}$ MeV).

NODE=M115W;LINKAGE=A

 $\eta(2225)$ REFERENCES

NODE=M115

ABLIKIM 16N PR D93 112011	M. Ablikim (BESIII Collab.)
LEES 11A PR D84 012001	J.P. Lees <i>et al.</i> (BABAR Collab.)
ABLIKIM 08I PL B662 330	M. Ablikim <i>et al.</i> (BES Collab.)
BAI 90B PRL 65 1309	Z. Bai <i>et al.</i> (Mark III Collab.)
BISELLO 86B PL B179 294	D. Bisello <i>et al.</i> (DM2 Collab.)

REFID=57512
REFID=16595
REFID=52255
REFID=41354
REFID=22101 **$\rho_3(2250)$**

$$I^G(J^{PC}) = 1^+(3^{--})$$

NODE=M044

OMITTED FROM SUMMARY TABLE

Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $f_4(2300)$, $\rho_5(2350)$.

NODE=M044

 $\rho_3(2250)$ MASS

NODE=M044205

 $\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M044M1
NODE=M044M1

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2248 ± 17 ⁺⁵⁹ ₋₅	1.8k	¹ ABLIKIM	20F BES3		$\psi(2S) \rightarrow K^+ K^- \eta$
~ 2232		HASAN	94 RVUE		$\bar{p}p \rightarrow \pi\pi$
~ 2090		² OAKDEN	94 RVUE		0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2250		³ MARTIN	80B RVUE		
~ 2300		³ MARTIN	80C RVUE		
~ 2140		⁴ CARTER	78B CNTR 0		0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2150		⁵ CARTER	77 CNTR 0		0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

¹ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (1.9 \pm 0.4_{-1.3}^{+0.5}) \times 10^{-6}$.

NODE=M044M1;LINKAGE=A

² See however KLOET 96 who fit $\pi^+\pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

NODE=M044M1;LINKAGE=CC

³ $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.

NODE=M044M1;LINKAGE=P

⁴ $I = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.

NODE=M044M1;LINKAGE=K

⁵ $I(J^P) = 1(3^-)$ from amplitude analysis.

NODE=M044M1;LINKAGE=J

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

NODE=M044M2
NODE=M044M2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2260 ± 20	⁶ ANISOVICH	02 SPEC		0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0$, $\omega\eta\pi^0, \pi^+\pi^-$
~ 2190	⁷ CUTTS	78B CNTR		0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2155 ± 15	^{7,8} COUPLAND	77 CNTR 0		0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{7,9} ALSPECTOR	73 CNTR		$\bar{p}p$ S channel
2190 ± 10	¹⁰ ABRAMS	70 CNTR		S channel $\bar{p}N$

⁶From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

⁷Isospins 0 and 1 not separated.

⁸From a fit to the total elastic cross section.

⁹Referred to as T or T region by ALSPECTOR 73.

¹⁰Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M044M;LINKAGE=AY

NODE=M044M2;LINKAGE=I

NODE=M044M2;LINKAGE=E

NODE=M044M2;LINKAGE=M

NODE=M044M2;LINKAGE=B

Other processes

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2290±20±30	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
------------	--------	----	---

NODE=M044M3

NODE=M044M3

$\rho_3(2250)$ WIDTH

$\bar{p}p \rightarrow \pi\pi$ or $K\bar{K}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$185^{+31}_{-26} + 17_{-103}$	1.8k	¹¹ ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
~ 220		HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 60		¹² OAKDEN	94	RVUE	0.36-1.55 $\bar{p}p \rightarrow \pi\pi$
~ 250		¹³ MARTIN	80B	RVUE	
~ 200		¹³ MARTIN	80C	RVUE	
~ 150		¹⁴ CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 200		¹⁵ CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

NODE=M044210

NODE=M044W1

NODE=M044W1

¹¹Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow X\eta \rightarrow K^+ K^- \eta = (1.9 \pm 0.4^{+0.5}_{-1.3}) \times 10^{-6}$.

NODE=M044W1;LINKAGE=A

¹²See however KLOET 96 who fit $\pi^+ \pi^-$ only and find waves only up to $J = 3$ to be important but not significantly resonant.

NODE=M044W1;LINKAGE=CC

¹³ $I(J^P) = 1(3^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

NODE=M044W1;LINKAGE=P

¹⁴ $l = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.

NODE=M044W1;LINKAGE=K

¹⁵ $I(J^P) = 1(3^-)$ from amplitude analysis.

NODE=M044W1;LINKAGE=J

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

160±25	¹⁶ ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega \pi^0$, $\omega \eta \pi^0, \pi^+ \pi^-$
135±75	^{17,18} COUPLAND	77	CNTR 0	0.7-2.4 $p\bar{p} \rightarrow \bar{p}p$
98± 8	¹⁸ ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
~ 85	¹⁹ ABRAMS	70	CNTR	S channel $\bar{p}N$

NODE=M044W2

NODE=M044W2

¹⁶From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M044W;LINKAGE=AY

¹⁷From a fit to the total elastic cross section.

NODE=M044W2;LINKAGE=E

¹⁸Isospins 0 and 1 not separated.

NODE=M044W2;LINKAGE=I

¹⁹Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

NODE=M044W2;LINKAGE=B

Other processes

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

230±50±80	AMELIN	00	VES 37 $\pi^- p \rightarrow \eta \pi^+ \pi^- n$
-----------	--------	----	---

NODE=M044W3

NODE=M044W3

$\rho_3(2250)$ REFERENCES

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
PEASLEE	75	PL 57B 189	D.C. Peaslee <i>et al.</i>	(CANB, BARI, BROW+)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
COOPER	68	PRL 20 1059	W.A. Cooper <i>et al.</i>	(ANL)

NODE=M044

REFID=60256

REFID=48828

REFID=48327

REFID=48349

REFID=47432

REFID=47950

REFID=45212

REFID=44103

REFID=45210

REFID=21838

REFID=21837

REFID=21964

REFID=21733

REFID=21963

REFID=21830

REFID=21824

REFID=21813

REFID=21807

REFID=21805

$f_2(2300)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M107

 $f_2(2300)$ MASS

NODE=M107M

NODE=M107M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2297 ± 28	¹ ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2262 ± 4 ± 28	² ABLIKIM	21AI BES3	3.51–4.60 $e^+ e^- \rightarrow \phi \Lambda \bar{\Lambda}$
2243 ⁺⁷ ₋₆ ⁺³ ₋₂₉	³ UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
2270 ± 12	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2327 ± 9 ± 6	ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
2231 ± 10	BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
2220 ⁺⁹⁰ ₋₂₀	LINDENBAUM	84 RVUE	
2320 ± 40	ETKIN	82 MPS	22 $\pi^- p \rightarrow 2\phi n$

¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi 2^{++} S_2$, D_2 , and D_0 is 6^{+15}_-5 , 25^{+18}_-14 , and 69^{+16}_-27 , respectively.

² Threshold enhancement in $\Lambda\bar{\Lambda}$, preferred J^{PC} are 2^{++} , 2^{-+} , or 1^{++} . Could be another state.

³ Spin 2 preferred, tentatively assigned to $f_2(2300)$.

NODE=M107M;LINKAGE=C

NODE=M107M;LINKAGE=B

NODE=M107M;LINKAGE=A

 $f_2(2300)$ WIDTH

NODE=M107W

NODE=M107W

NEW

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 ± 40 OUR AVERAGE	[149 ± 40 MeV OUR 2023 AVERAGE]		
149 ± 41	¹ ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
72 ± 5 ± 43	² ABLIKIM	21AI BES3	3.51–4.60 $e^+ e^- \rightarrow \phi \Lambda \bar{\Lambda}$
145 ± 12 ⁺²⁷ ₋₃₄	³ UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
90 ± 29	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
275 ± 36 ± 20	ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$
133 ± 50	BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
200 ± 50	LINDENBAUM	84 RVUE	
220 ± 70	ETKIN	82 MPS	22 $\pi^- p \rightarrow 2\phi n$

¹ Includes data of ETKIN 85.

² Threshold enhancement in $\Lambda\bar{\Lambda}$, preferred J^{PC} are 2^{++} , 2^{-+} , or 1^{++} . Could be another state.

³ Spin 2 preferred, tentatively assigned to $f_2(2300)$.

NODE=M107W;LINKAGE=C

NODE=M107W;LINKAGE=B

NODE=M107W;LINKAGE=A

 $f_2(2300)$ DECAY MODES

NODE=M107215;NODE=M107

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi\phi$	seen
Γ_2 $K\bar{K}$	seen
Γ_3 $\gamma\gamma$	seen
Γ_4 $\Lambda\bar{\Lambda}$	seen

DESIG=1

DESIG=2

DESIG=3

DESIG=4

 $f_2(2300)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M107225

 $\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_2\Gamma_3/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.2 ^{+0.5} _{-0.4} ^{+1.3} _{-2.2}	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
44 ± 6 ± 12	¹ ABE	04 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$

¹ Assuming spin 2.

NODE=M107G1

NODE=M107G1

NODE=M107G1;LINKAGE=AB

$f_2(2340)$ BRANCHING RATIOS

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	BOOTH	86	OMEG 85 $\pi^- \text{Be} \rightarrow 2\phi \text{Be}$	
seen	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$	

NODE=M107220

NODE=M107R01
NODE=M107R01

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	VLADIMIRSK...06	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$	
seen	ABE	04	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$	

NODE=M107R02
NODE=M107R02

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen	UEHARA	13	BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$	

NODE=M107R03
NODE=M107R03

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
seen	¹ ABLIKIM	21AI	BES3 3.51–4.60 $e^+ e^- \rightarrow \phi\Lambda\bar{\Lambda}$	

NODE=M107R00
NODE=M107R00

¹ Threshold enhancement in $\Lambda\bar{\Lambda}$, preferred J^{PC} are 2^{++} , 2^{-+} , or 1^{++} . Could be another state.

NODE=M107R00;LINKAGE=A

 $f_2(2300)$ REFERENCES

ABLIKIM	21AI	PR D104 052006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61440
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49650
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=40285
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIVP, GLAS, CERN)	REFID=21870
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21871
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)	REFID=21869
ETKIN	82	PRL 49 1620	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21866

NODE=M107

NODE=M041

 $f_4(2300)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_0(2350)$. Contains results mostly from formation experiments. For further production experiments see the Further States entry. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $\rho_5(2350)$.

NODE=M041

 $f_4(2300)$ MASS **$\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 2314	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 2300	¹ MARTIN	80B	RVUE
~ 2300	¹ MARTIN	80C	RVUE
~ 2340	² CARTER	78B	CNTR 0.7–2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2330	DULUDE	78B	OSPK 1–2 $\bar{p}p \rightarrow \pi^0 \pi^0$
~ 2310	³ CARTER	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \pi\pi$

¹ $I(J^P) = 0(4^+)$ from simultaneous analysis of $\bar{p}p \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

² $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.

³ $I(J^P) = 0(4^+)$ from amplitude analysis.

NODE=M041205

NODE=M041M1
NODE=M041M1NODE=M041M1;LINKAGE=P
NODE=M041M1;LINKAGE=K
NODE=M041M1;LINKAGE=J**S-CHANNEL $\bar{p}p$ or $\bar{N}N$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2283±17	⁴ ANISOVICH	00J	SPEC
~ 2380	⁵ CUTTS	78B	CNTR 0.97–3 $\bar{p}p \rightarrow \bar{N}N$
2345±15	^{5,6} COUPLAND	77	CNTR 0.7–2.4 $\bar{p}p \rightarrow \bar{p}p$
2359± 2	^{5,7} ALSPECTOR	73	CNTR $\bar{p}p$ S channel
2375±10	ABRAMS	70	CNTR S channel $\bar{N}N$

⁴ From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on $\bar{p}p \rightarrow \eta\pi^0\pi^0$, $\pi^0\pi^0$, $\eta\eta$, $\eta\eta'$, $\pi^+\pi^-$.

⁵ Isospins 0 and 1 not separated.

⁶ From a fit to the total elastic cross section.

⁷ Referred to as U or U region by ALSPECTOR 73.

NODE=M041M2
NODE=M041M2

NODE=M041M2;LINKAGE=AN

NODE=M041M2;LINKAGE=I
NODE=M041M2;LINKAGE=E
NODE=M041M2;LINKAGE=M

$\pi^- p \rightarrow \eta \pi \pi n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2330 \pm 20 \pm 40$	AMELIN	00	VES $37 \pi^- p \rightarrow \eta \pi^+ \pi^- n$

NODE=M041M3
 NODE=M041M3

 $\rho\rho$ CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	COMMENT
2320 ± 60 OUR ESTIMATE		
2332 ± 15	BARBERIS	00F $450 \rho\rho \rightarrow p_f \omega \omega p_s$

NODE=M041M4
 NODE=M041M4
 → UNCHECKED ←

 $f_4(2300)$ WIDTH

NODE=M041210

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
~ 278	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$
~ 200	⁸ MARTIN	80C	RVUE
~ 150	⁹ CARTER	78B	CNTR $0.7-2.4 \bar{p}p \rightarrow K^- K^+$
~ 210	¹⁰ CARTER	77	CNTR $0.7-2.4 \bar{p}p \rightarrow \pi\pi$
⁸ $I(J^P) = 0(4^+)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.			
⁹ $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.			
¹⁰ $I(J^P) = 0(4^+)$ from amplitude analysis.			

NODE=M041W1
 NODE=M041W1

NODE=M041W1;LINKAGE=P
 NODE=M041W1;LINKAGE=K
 NODE=M041W1;LINKAGE=J

S-CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
310 ± 25	¹¹ ANISOVICH	00J	SPEC
135^{+150}_{-65}	^{12,13} COUPLAND	77	CNTR $0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
165^{+18}_{-8}	¹³ ALSPECTOR	73	CNTR $\bar{p}p$ S channel
~ 190	ABRAMS	70	CNTR S channel $\bar{N}N$
¹¹ From the combined analysis of ANISOVICH 99C and ANISOVICH 99F on $\bar{p}p \rightarrow \eta \pi^0 \pi^0$, $\pi^0 \pi^0$, $\eta\eta$, $\eta\eta'$, $\pi^+ \pi^-$.			
¹² From a fit to the total elastic cross section.			
¹³ Isospins 0 and 1 not separated.			

NODE=M041W2
 NODE=M041W2

NODE=M041W2;LINKAGE=AN

NODE=M041W2;LINKAGE=E
 NODE=M041W2;LINKAGE=I

 $\pi^- p \rightarrow \eta \pi \pi n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$235 \pm 50 \pm 40$	AMELIN	00	VES $37 \pi^- p \rightarrow \eta \pi^+ \pi^- n$

NODE=M041W3
 NODE=M041W3

 $\rho\rho$ CENTRAL PRODUCTION

VALUE (MeV)	DOCUMENT ID	COMMENT
250 ± 80 OUR ESTIMATE		
260 ± 57	BARBERIS	00F $450 \rho\rho \rightarrow p_f \omega \omega p_s$

NODE=M041W4
 NODE=M041W4
 → UNCHECKED ←

 $f_4(2300)$ DECAY MODES

NODE=M041215;NODE=M041

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\rho$	seen
Γ_2 $\omega\omega$	seen
Γ_3 $\eta\pi\pi$	seen
Γ_4 $\pi\pi$	seen
Γ_5 $K\bar{K}$	seen
Γ_6 $N\bar{N}$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=3;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=5;OUR EST;→ UNCHECKED ←
 DESIG=6;OUR EST;→ UNCHECKED ←

 $f_4(2300)$ BRANCHING RATIOS

NODE=M041220

 $\Gamma(\rho\rho)/\Gamma(\omega\omega)$ **Γ_1/Γ_2**

VALUE	DOCUMENT ID	COMMENT
2.8 ± 0.5	BARBERIS	00F $450 \rho\rho \rightarrow p_f \omega \omega p_s$

NODE=M041R1
 NODE=M041R1

$f_4(2300)$ REFERENCES

AMELIN	00	NP A668 83	D. Amelin <i>et al.</i>	(VES Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARBERIS	00F	PL B484 198	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99C	PL B452 173	A.V. Anisovich <i>et al.</i>	
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
DULUDE	78B	PL 79B 335	R.S. Dulude <i>et al.</i>	(BROW, MIT, BARI) JP
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)

NODE=M041

REFID=47432
 REFID=47950
 REFID=47962
 REFID=46903
 REFID=46926
 REFID=44103
 REFID=21838
 REFID=21837
 REFID=21964
 REFID=21733
 REFID=21850
 REFID=21963
 REFID=21830
 REFID=21813
 REFID=21807

 $f_0(2330)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

NODE=M169

 $f_0(2330)$ MASS

NODE=M169M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$2312 \pm 7_{-3}^{+7}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
$2312 \pm 2_{-0}^{+10}$	² ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
2419 ± 64	³ RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K})$
2340 ± 20	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
2314 ± 25	⁴ BUGG	04A RVUE	
2337 ± 14	ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \pi \pi, \eta \eta$
~ 2321	HASAN	94 RVUE	$\bar{p} p \rightarrow \pi \pi$

NODE=M169M

- ¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.
² From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.
³ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).
⁴ Partial wave analysis of the data on $p \bar{p} \rightarrow \bar{\Lambda} \Lambda$ from BARNES 00.

NODE=M169M;LINKAGE=C

NODE=M169M;LINKAGE=B

NODE=M169M;LINKAGE=A

NODE=M169M;LINKAGE=BU

 $f_0(2330)$ WIDTH

NODE=M169W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$65 \pm 10_{-12}^{+3}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
$134 \pm 5_{-9}^{+30}$	² ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
274 ± 94	³ RODAS	22 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K})$
165 ± 25	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
144 ± 20	⁴ BUGG	04A RVUE	
217 ± 33	ANISOVICH	00J SPEC	$2.0 \bar{p} p \rightarrow \pi \pi, \eta \eta$
~ 223	HASAN	94 RVUE	$\bar{p} p \rightarrow \pi \pi$

NODE=M169W

- ¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.
² From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.
³ T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma \pi^0 \pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).
⁴ Partial wave analysis of the data on $p \bar{p} \rightarrow \bar{\Lambda} \Lambda$ from BARNES 00.

NODE=M169W;LINKAGE=C

NODE=M169W;LINKAGE=B

NODE=M169W;LINKAGE=A

NODE=M169W;LINKAGE=BU

$f_0(2330)$ REFERENCES

ABLIKIM	22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also		PR D107 079901 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BUGG	04A	EPJ C36 161	D.V. Bugg	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BARNES	00	PR C62 055203	P.D. Barnes <i>et al.</i>	
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)

NODE=M169

REFID=61891
 REFID=62033
 REFID=61637
 REFID=61610
 REFID=61091
 REFID=59455
 REFID=56984
 REFID=50158
 REFID=47950
 REFID=47965
 REFID=44103

 $f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M108

 $f_2(2340)$ MASS

NODE=M108M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2346⁺²¹₋₁₀ OUR AVERAGE				
2346 ± 8 ⁺²² ₋₆		¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
2362 ⁺³¹⁺¹⁴⁰ ₋₃₀₋₆₃	5.5k	² ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
2339 ± 55		³ ETKIN	88 MPS	$22 \pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2350 ± 7	80k	⁴ UMAN	06 E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
2392 ± 10		BOOTH	86 OMEG	$85 \pi^- \text{Be} \rightarrow 2\phi \text{Be}$
2360 ± 20		LINDENBAUM	84 RVUE	

NODE=M108M

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M108M;LINKAGE=B

² From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M108M;LINKAGE=A

³ Includes data of ETKIN 85. The percentage of the resonance going into $\phi \phi 2^{++} S_2$, D_2 , and D_0 is 37 ± 19 , 4^{+12}_{-4} , and 59^{+21}_{-19} , respectively.

NODE=M108M;LINKAGE=C

⁴ Statistical error only.

NODE=M108M;LINKAGE=ST

 $f_2(2340)$ WIDTH

NODE=M108W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
331⁺²⁷₋₁₈ OUR AVERAGE				
332 ± 14 ⁺²⁶ ₋₁₂		¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
334 ⁺⁶²⁺¹⁶⁵ ₋₅₄₋₁₀₀	5.5k	² ABLIKIM	13N BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \eta \eta$
319 ⁺⁸¹ ₋₆₉		³ ETKIN	88 MPS	$22 \pi^- p \rightarrow \phi \phi n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
218 ± 16	80k	⁴ UMAN	06 E835	$5.2 \bar{p} p \rightarrow \eta \eta \pi^0$
198 ± 50		BOOTH	86 OMEG	$85 \pi^- \text{Be} \rightarrow 2\phi \text{Be}$
150 ⁺¹⁵⁰ ₋₅₀		LINDENBAUM	84 RVUE	

NODE=M108W

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta' \eta'$, and ($\eta' X$), with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M108W;LINKAGE=B

² From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M108W;LINKAGE=A

³ Includes data of ETKIN 85.

NODE=M108W;LINKAGE=C

⁴ Statistical error only.

NODE=M108W;LINKAGE=ST

$f_2(2340)$ DECAY MODES

NODE=M108215;NODE=M108

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi\phi$	seen
Γ_2 $\eta\eta$	seen
Γ_3 $\eta'\eta'$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2
DESIG=3

 $f_2(2340)$ BRANCHING RATIOS

NODE=M108220

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	UMAN	06	E835 $5.2 \bar{p}p \rightarrow \eta\eta\pi^0$

NODE=M108R01
NODE=M108R01

$\Gamma(\eta'\eta')/\Gamma_{\text{total}}$	Γ_3/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M108R00
NODE=M108R00

¹ From a partial wave analysis of the systems (γX), with $X \rightarrow \eta'\eta'$, and ($\eta' X$), with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M108R00;LINKAGE=A

 $f_2(2340)$ REFERENCES

NODE=M108

ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61637
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
ETKIN	88	PL B201 568	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=40285
BOOTH	86	NP B273 677	P.S.L. Booth <i>et al.</i>	(LIVP, GLAS, CERN)	REFID=21870
ETKIN	85	PL 165B 217	A. Etkin <i>et al.</i>	(BNL, CUNY)	REFID=21871
LINDENBAUM	84	CNPP 13 285	S.J. Lindenbaum	(CUNY)	REFID=21869

NODE=M033

 $\rho_5(2350)$

$$I^G(J^{PC}) = 1^+(5^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_1(2400)$. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $f_4(2300)$.

NODE=M033

 $\rho_5(2350)$ MASS

NODE=M033205

 $\pi^- p \rightarrow \omega\pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2330±35	ALDE	95	GAM2 $38 \pi^- p \rightarrow \omega\pi^0 n$

NODE=M033M3
NODE=M033M3

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

NODE=M033M1
NODE=M033M1

••• We do not use the following data for averages, fits, limits, etc. •••

~ 2303	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
~ 2300	¹ MARTIN	80B	RVUE	
~ 2250	¹ MARTIN	80C	RVUE	
~ 2500	² CARTER	78B	CNTR 0	$0.7-2.4 \bar{p}p \rightarrow K^- K^+$
~ 2480	³ CARTER	77	CNTR 0	$0.7-2.4 \bar{p}p \rightarrow \pi\pi$

S-CHANNEL $\bar{N}N$

NODE=M033M2
NODE=M033M2

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

••• We do not use the following data for averages, fits, limits, etc. •••

2300±45	⁴ ANISOVICH	02	SPEC	$0.6-1.9 \bar{p}p \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
2295±30	ANISOVICH	00J	SPEC	
~ 2380	⁵ CUTTS	78B	CNTR	$0.97-3 \bar{p}p \rightarrow \bar{N}N$
2345±15	^{5,6} COUPLAND	77	CNTR 0	$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
2359± 2	^{5,7} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2350±10	⁸ ABRAMS	70	CNTR	S channel $\bar{N}N$
2360±25	⁹ OH	70B	HDBC -0	$\bar{p}(pn), K^* K 2\pi$

$\pi^- p \rightarrow K^+ K^- n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2307 ± 6	ALPER	80	CNTR	0	62 $\pi^- p \rightarrow K^+ K^- n$
----------	-------	----	------	---	------------------------------------

¹ $I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

² $I = 0(1); J^P = 5^-$ from Barrelet-zero analysis.

³ $I(J^P) = 1(5^-)$ from amplitude analysis.

⁴ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

⁵ Isospins 0 and 1 not separated.

⁶ From a fit to the total elastic cross section.

⁷ Referred to as U or U region by ALSPECTOR 73.

⁸ For $I = 1 \bar{N}N$.

⁹ No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.

NODE=M033M4
NODE=M033M4

NODE=M033M1;LINKAGE=P
NODE=M033M1;LINKAGE=K
NODE=M033M1;LINKAGE=J
NODE=M033M2;LINKAGE=AY

NODE=M033M2;LINKAGE=I
NODE=M033M2;LINKAGE=E
NODE=M033M2;LINKAGE=M
NODE=M033M2;LINKAGE=A
NODE=M033M2;LINKAGE=N

 $\rho_5(2350)$ WIDTH

NODE=M033210

 $\pi^- p \rightarrow \omega \pi^0 n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

400 ± 100	ALDE	95	GAM2	38 $\pi^- p \rightarrow \omega \pi^0 n$
-----------	------	----	------	---

NODE=M033W3
NODE=M033W3

 $\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 169	HASAN	94	RVUE	$\bar{p}p \rightarrow \pi\pi$
-------	-------	----	------	-------------------------------

~ 250	10 MARTIN	80B	RVUE	
-------	-----------	-----	------	--

~ 300	10 MARTIN	80C	RVUE	
-------	-----------	-----	------	--

~ 150	11 CARTER	78B	CNTR	0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
-------	-----------	-----	------	---	--

~ 210	12 CARTER	77	CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$
-------	-----------	----	------	---	---------------------------------------

NODE=M033W1
NODE=M033W1

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

260 ± 75	13 ANISOVICH	02	SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega \pi^0, \omega \eta \pi^0, \pi^+ \pi^-$
----------	--------------	----	------	---

235 ⁺ ₄₀	ANISOVICH	00J	SPEC	
--------------------------------	-----------	-----	------	--

135 ⁺ ₆₅ 150 ⁺ ₆₅	14,15 COUPLAND	77	CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
--	----------------	----	------	---	---

165 ⁺ ₈ 18 ⁺ ₈	15 ALSPECTOR	73	CNTR		$\bar{p}p$ S channel
---	--------------	----	------	--	----------------------

< 60	16 OH	70B	HDBC	-0	$\bar{p}(pn), K^* K 2\pi$
------	-------	-----	------	----	---------------------------

~ 140	ABRAMS	67C	CNTR		S channel $\bar{p}N$
-------	--------	-----	------	--	----------------------

NODE=M033W2
NODE=M033W2

 $\pi^- p \rightarrow K^+ K^- n$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

245 ± 20	ALPER	80	CNTR	0	62 $\pi^- p \rightarrow K^+ K^- n$
----------	-------	----	------	---	------------------------------------

¹⁰ $I(J^P) = 1(5^-)$ from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.

¹¹ $I = 0(1); J^P = 5^-$ from Barrelet-zero analysis.

¹² $I(J^P) = 1(5^-)$ from amplitude analysis.

¹³ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

¹⁴ From a fit to the total elastic cross section.

¹⁵ Isospins 0 and 1 not separated.

¹⁶ No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.

NODE=M033W4
NODE=M033W4

OCCUR=2

NODE=M033W1;LINKAGE=P
NODE=M033W1;LINKAGE=K
NODE=M033W1;LINKAGE=J
NODE=M033W2;LINKAGE=AY

NODE=M033W2;LINKAGE=E
NODE=M033W2;LINKAGE=I
NODE=M033W2;LINKAGE=N

$\rho_5(2350)$ REFERENCES

ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>	
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>	
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
ALDE	95	ZPHY C66 379	D.M. Alde <i>et al.</i>	(GAMS Collab.) JP
HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)
ALPER	80	PL 94B 422	B. Alper <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
MARTIN	80B	NP B176 355	B.R. Martin, D. Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	A.D. Martin, M.R. Pennington	(DURH) JP
CARTER	78B	NP B141 467	A.A. Carter	(LOQM)
CUTTS	78B	PR D17 16	D. Cutts <i>et al.</i>	(STON, WISC)
CARTER	77	PL 67B 117	A.A. Carter <i>et al.</i>	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	M. Coupland <i>et al.</i>	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	J. Alspector <i>et al.</i>	(RUTG, UPNJ)
OH	73	NP B51 57	B.Y. Oh <i>et al.</i>	(MSU)
CHAPMAN	71B	PR D4 1275	J.W. Chapman <i>et al.</i>	(MICH)
ABRAMS	70	PR D1 1917	R.J. Abrams <i>et al.</i>	(BNL)
OH	70B	PRL 24 1257	B.Y. Oh <i>et al.</i>	(MSU)
ABRAMS	67C	PRL 18 1209	R.J. Abrams <i>et al.</i>	(BNL)

NODE=M033

REFID=48828
REFID=48327
REFID=48349
REFID=47950
REFID=44371
REFID=44103
REFID=21665
REFID=21838
REFID=21837
REFID=21964
REFID=21733
REFID=21963
REFID=21830
REFID=21813
REFID=21931
REFID=21926
REFID=21807
REFID=21925
REFID=21804

X(2370)

$$J^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

NODE=M247

X(2370) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2357 ±17 OUR AVERAGE		Error includes scale factor of 2.7.		
2341.6 ± 6.5 ± 5.7		¹ ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K \bar{K} \eta'$
2376.3 ± 8.7 ^{+3.2} _{-4.3}	565	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M247M

NODE=M247M

¹ The state observed by ABLIKIM 11C at 2120 MeV is not observed with 90% CL upper limit of 1.49×10^{-5} for $J/\psi \rightarrow \gamma X(2120) \rightarrow \gamma K^+ K^- \eta'$ and 6.38×10^{-6} for $K_S^0 K_S^0 \eta'$.

NODE=M247M;LINKAGE=A

X(2370) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
114⁺¹²₋₁₀ OUR AVERAGE			
117 ± 10 ± 8	¹ ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K \bar{K} \eta'$
83 ± 17 ⁺⁴⁴ ₋₆	ABLIKIM	11C BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M247W

NODE=M247W

¹ The state observed by ABLIKIM 11C at 2120 MeV is not observed with 90% CL upper limit of 1.49×10^{-5} for $J/\psi \rightarrow \gamma X(2120) \rightarrow \gamma K^+ K^- \eta'$ and 6.38×10^{-6} for $K_S^0 K_S^0 \eta'$.

NODE=M247W;LINKAGE=A

X(2370) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^+ K^- \eta'$	seen
Γ_2 $K_S^0 K_S^0 \eta'$	seen
Γ_3 $\pi^+ \pi^- \eta'$	seen
Γ_4 $\eta \eta \eta'$	not seen

NODE=M247215;NODE=M247

DESIG=1

DESIG=2

DESIG=3

DESIG=4

X(2370) BRANCHING RATIOS

$\Gamma(K^+ K^- \eta')/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K^+ K^- \eta'$
$\Gamma(K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	20Q BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$

NODE=M247225

NODE=M247R01

NODE=M247R01

NODE=M247R02

NODE=M247R02

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

NODE=M247R00
 NODE=M247R00

 $\Gamma(\eta\eta\eta')/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	21C	BES3 $J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$

NODE=M247R03
 NODE=M247R03

¹ ABLIKIM 21C measured $B(J/\psi(1S) \rightarrow \gamma X(2370) \rightarrow \gamma\eta\eta\eta') < 9.2 \times 10^{-6}$.

NODE=M247R03;LINKAGE=A

X(2370) REFERENCES

ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)

NODE=M247

REFID=61030
 REFID=60457
 REFID=53684

 $f_0(2470)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

Seen by ABLIKIM 22C with a significance of 5.2σ in a partial-wave analysis of the systems (γX) , $X \rightarrow \eta'\eta'$ and $(\eta'X)$, $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$.

NODE=M266

NODE=M266

 $f_0(2470)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2470 \pm 4_{-6}^{+4}$	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M266M

NODE=M266M

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta'X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M266M;LINKAGE=A

 $f_0(2470)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$75 \pm 9_{-8}^{+11}$	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M266W

NODE=M266W

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta'X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M266W;LINKAGE=C

 $f_0(2470)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \eta'\eta'$	seen

NODE=M266215;NODE=M266

DESIG=1

 $\Gamma(\eta'\eta')/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma\eta'\eta' \rightarrow 4/5\gamma 2(\pi^+\pi^-)$

NODE=M266R00
 NODE=M266R00

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta'\eta'$, and $(\eta'X)$, with $X \rightarrow \gamma\eta'$ in the decay $J/\psi \rightarrow \gamma\eta'\eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M266R00;LINKAGE=A

 $f_0(2470)$ REFERENCES

ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
---------	-----	----------------	--------------------------	------------------

NODE=M266

REFID=61637

$f_6(2510)$

$$J^G(J^{PC}) = 0^+(6^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M089

NODE=M089

 $f_6(2510)$ MASS

NODE=M089M

VALUE (MeV) DOCUMENT ID TECN COMMENT
2470±50 OUR AVERAGE Error includes scale factor of 2.1. [2465 ± 50 MeV OUR 2023 AVERAGE Scale factor = 2.1]

NODE=M089M

NEW

2420±30 ALDE 98 GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
 2510±30 BINON 84B GAM2 38 $\pi^- p \rightarrow n 2\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2485±40 ¹ ANISOVICH 00J SPEC 1.92–2.41 $p\bar{p}$

NODE=M089M;LINKAGE=AN

¹From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B.

 $f_6(2510)$ WIDTH

NODE=M089W

VALUE (MeV) DOCUMENT ID TECN COMMENT
260±40 OUR AVERAGE
 [255 ± 40 MeV OUR 2023 AVERAGE]

NODE=M089W

NEW

270±60 ALDE 98 GAM4 100 $\pi^- p \rightarrow \pi^0 \pi^0 n$
 240±60 BINON 84B GAM2 38 $\pi^- p \rightarrow n 2\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

410±90 ² ANISOVICH 00J SPEC 1.92–2.41 $p\bar{p}$

NODE=M089W;LINKAGE=AN

²From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B.

 $f_6(2510)$ DECAY MODES

NODE=M089215;NODE=M089

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	(6.0±1.0) %

DESIG=1

 $f_6(2510)$ BRANCHING RATIOS

NODE=M089220

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$ Γ_1/Γ
 VALUE DOCUMENT ID TECN COMMENT
0.06±0.01 ³ BINON 83C GAM2 38 $\pi^- p \rightarrow n 4\gamma$

NODE=M089R1

NODE=M089R1

³Assuming one pion exchange and using data of BOLOTOV 74.

NODE=M089R1;LINKAGE=A

 $f_6(2510)$ REFERENCES

NODE=M089

ANISOVICH 00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH 00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
ANISOVICH 99C	PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH 99F	NP A651 253	A.V. Anisovich <i>et al.</i>		REFID=46926
ANISOVICH 99J	PL B471 271	A.V. Anisovich <i>et al.</i>		REFID=47416
ANISOVICH 99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
ALDE 98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)	REFID=46605
Also	PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)	REFID=46914
	Translated from YAF 62 446.			
BINON 84B	LNC 39 41	F.G. Binon <i>et al.</i>	(SERP, BELG, LAPP) JP	REFID=21780
BINON 83C	SJNP 38 723	F.G. Binon <i>et al.</i>	(SERP, BRUX+)	REFID=40288
	Translated from YAF 38 1199.			
BOLOTOV 74	PL 52B 489	V.N. Bolotov <i>et al.</i>	(SERP)	REFID=44705

STRANGE MESONS

($S = \pm 1, C = B = 0$)

$K^+ = u\bar{s}, K^0 = d\bar{s}, \bar{K}^0 = \bar{d}s, K^- = \bar{u}s$, similarly for K^{*} 's

$K_0^*(700)$

$$I(J^P) = \frac{1}{2}(0^+)$$

also known as κ ; was $K_0^*(800)$

See the related review(s):

Scalar Mesons below 1 GeV

$K_0^*(700)$ T-Matrix Pole \sqrt{s}

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(630-730) - i (260-340) OUR ESTIMATE (see Fig. 64.1 in the review)			
$(702 \pm 12^{+4}_{-5}) - i (285 \pm 16^{+8}_{-13})$	¹ DANILKIN	21	RVUE Compilation
$(648 \pm 7) - i (280 \pm 16)$	² PELAEZ	20	RVUE $\pi K \rightarrow \pi K$
$(670 \pm 18) - i (295 \pm 28)$	³ PELAEZ	17	RVUE $\pi K \rightarrow \pi K$
$(764 \pm 63^{+71}_{-54}) - i (306 \pm 149^{+143}_{-85})$	⁴ ABLIKIM	11B	BES2 $1.3k J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
$(665 \pm 9) - i (268^{+21}_{-6})$	⁵ GUO	11B	RVUE
$(849 \pm 77^{+18}_{-14}) - i (256 \pm 40^{+46}_{-22})$	⁴ ABLIKIM	10E	BES2 $1.4k J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
$(663 \pm 8 \pm 34) - i (329 \pm 5 \pm 22)$	⁶ BUGG	10	RVUE S-matrix pole
$(706.0 \pm 1.8 \pm 22.8) - i (319.4 \pm 2.2 \pm 20.2)$	⁷ BONVICINI	08A	CLEO $141k D^+ \rightarrow K^- \pi^+ \pi^+$
$(841 \pm 30^{+81}_{-73}) - i (309 \pm 45^{+48}_{-72})$	⁴ ABLIKIM	06C	BES2 $25k J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
$(750^{+30}_{-55}) - i (342 \pm 60)$	⁸ BUGG	06	RVUE
$(658 \pm 13) - i (279 \pm 12)$	⁹ DESCOTES-G.	06	RVUE $\pi K \rightarrow \pi K$
$(757 \pm 33) - i (279 \pm 41)$	¹⁰ GUO	06	RVUE
$(694 \pm 53) - i (303 \pm 30)$	¹¹ ZHOU	06	RVUE $K p \rightarrow K^- \pi^+ n$
$(594 \pm 79) - i (362 \pm 166)$	¹¹ ZHENG	04	RVUE $K^- p \rightarrow K^- \pi^+ n$
$(722 \pm 60) - i (386 \pm 50)$	¹¹ BUGG	03	RVUE $11 K^- p \rightarrow K^- \pi^+ n$
$(875 \pm 75) - i (335 \pm 110)$	¹² ISHIDA	97B	RVUE $11 K^- p \rightarrow K^- \pi^+ n$
$727 - i 263$	¹³ VANBEVEREN	86	RVUE

¹ Data driven analysis using partial-wave dispersion relations .

² Extracted employing πK partial wave analysis from ESTABROOKS 78 and ASTON 88, Roy-Steiner equations and once subtracted forward dispersion relations.

³ Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

⁴ Extracted from Breit-Wigner parameters.

⁵ Fit to scattering phase shifts using UChPT amplitudes with explicit resonances.

⁶ Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.

⁷ From a complex pole included in the fit. Using parameters from the model that fits data best.

⁸ Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C using for the κ an s-dependent width with an Adler zero near threshold.

⁹ Using Roy-Steiner equations (ROY 71) consistent with unitarity, analyticity and crossing symmetry constraints.

¹⁰ From UChPT fitted to MERCER 71, BINGHAM 72 and ESTABROOKS 78. Amplitude shown to be consistent with data of ABLIKIM 06C.

¹¹ Reanalysis of ASTON 88 data.

¹² Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes. Extracted from Breit-Wigner parameters.

¹³ Unitarized Quark Model.

NODE=MXXX020

NODE=MXXX020

NODE=M174

NODE=M174TMP

NODE=M174TMP

→ UNCHECKED ←

NODE=M174TMP;LINKAGE=P

NODE=M174TMP;LINKAGE=J

NODE=M174TMP;LINKAGE=N

NODE=M174TMP;LINKAGE=A

NODE=M174TMP;LINKAGE=D

NODE=M174TMP;LINKAGE=H

NODE=M174TMP;LINKAGE=B

NODE=M174TMP;LINKAGE=G

NODE=M174TMP;LINKAGE=I

NODE=M174TMP;LINKAGE=O

NODE=M174TMP;LINKAGE=F

NODE=M174TMP;LINKAGE=M

NODE=M174TMP;LINKAGE=E

$K_0^*(700)$ Breit-Wigner Mass

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
845 ±17	OUR AVERAGE			
826 ±49 $^{+49}_{-34}$	1.3k	¹ ABLIKIM	11B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
810 ±68 $^{+15}_{-24}$	1.4k	² ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
856 ±17 ±13	54k	LINK	07B FOCUS	$D^+ \rightarrow K^- \pi^+ \pi^+$
878 ±23 $^{+64}_{-55}$	25k	³ ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
797 ±19 ±43	15k	^{4,5} AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
888.0 ± 1.9	141k	⁶ BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
855 ±15	0.6k	⁷ CAWLFIELD	06A CLEO	$D^0 \rightarrow K^+ K^- \pi^0$
905 $^{+65}_{-30}$		⁸ ISHIDA	97B RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M174M
 NODE=M174M

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M174M;LINKAGE=LI

NODE=M174M;LINKAGE=BL

NODE=M174M;LINKAGE=EP

NODE=M174M;LINKAGE=A

NODE=M174M;LINKAGE=AU

NODE=M174M;LINKAGE=C

NODE=M174M;LINKAGE=CA

NODE=M174M;LINKAGE=IS

¹ The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$ MeV.

² From a fit including ten additional resonances and energy-independent Breit-Wigner width.

³ A fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model with mass and width of the $K_0^*(700)$ from ABLIKIM 06C well describes the left slope of the $K_S^0 \pi^-$ invariant mass spectrum in $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$ decay studied by EPIFANOV 07. Averaged value from different parameterizations.

⁴ Not seen by KOPP 01 using 7070 events of $D^0 \rightarrow K^- \pi^+ \pi^0$. LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than $K_0^*(700)$ in their high statistics analysis of $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$.

⁵ AUBERT 07T does not find evidence for the charged $K_0^*(700)$ using 11k events of $D^0 \rightarrow K^- K^+ \pi^0$.

⁶ Using parameters from the model that fits data best.

⁷ Breit-Wigner parameters. A significant S-wave can be also modeled as a non-resonant contribution.

⁸ Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.

 $K_0^*(700)$ Breit-Wigner Width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
468 ± 30	OUR AVERAGE			
449 ±156 $^{+144}_{-81}$	1.3k	¹ ABLIKIM	11B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$
536 ± 87 $^{+106}_{-47}$	1.4k	² ABLIKIM	10E BES2	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
464 ± 28 ± 22	54k	LINK	07B FOCUS	$D^+ \rightarrow K^- \pi^+ \pi^+$
499 ± 52 $^{+55}_{-87}$	25k	³ ABLIKIM	06C BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
410 ± 43 ± 87	15k	^{4,5} AITALA	02 E791	$D^+ \rightarrow K^- \pi^+ \pi^+$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
550.4 ± 11.8	141k	⁶ BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
251 ± 48	0.6k	⁷ CAWLFIELD	06A CLEO	$D^0 \rightarrow K^+ K^- \pi^0$
545 $^{+235}_{-110}$		⁸ ISHIDA	97B RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M174W

NODE=M174W

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M174W;LINKAGE=LI

NODE=M174W;LINKAGE=BL

NODE=M174W;LINKAGE=EP

NODE=M174W;LINKAGE=A

NODE=M174W;LINKAGE=AU

NODE=M174W;LINKAGE=C

NODE=M174W;LINKAGE=CA

NODE=M174W;LINKAGE=IS

¹ The Breit-Wigner parameters from a fit with seven intermediate resonances. The S-matrix pole position is $(764 \pm 63^{+71}_{-54}) - i(306 \pm 149^{+143}_{-85})$ MeV.

² From a fit including ten additional resonances and energy-independent Breit-Wigner width.

³ A fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model with mass and width of the $K_0^*(700)$ from ABLIKIM 06C well describes the left slope of the $K_S^0 \pi^-$ invariant mass spectrum in $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$ decay studied by EPIFANOV 07. Averaged value from different parameterizations.

⁴ Not seen by KOPP 01 using 7070 events of $D^0 \rightarrow K^- \pi^+ \pi^0$. LINK 02E and LINK 05I show clear evidence for a constant non-resonant scalar amplitude rather than $K_0^*(700)$ in their high statistics analysis of $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$.

⁵ AUBERT 07T does not find evidence for the charged $K_0^*(700)$ using 11k events of $D^0 \rightarrow K^- K^+ \pi^0$.

⁶ Using parameters from the model that fits data best.

⁷ Statistical error only. A fit to the Dalitz plot including the $K_0^*(700)^\pm$, $K^*(892)^\pm$, and ϕ resonances modeled as Breit-Wigners. A significant S-wave can be also modeled as a non-resonant contribution.

⁸ Reanalysis of ASTON 88 using interfering Breit-Wigner amplitudes.

$K^*(700)$ DECAY MODES

NODE=M174215;NODE=M174

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	100 %

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $K^*(700)$ REFERENCES

NODE=M174

DANILKIN	21	PR D103 114023	I. Danilkin, O. Deineka, M. Vanderhaeghen (MAINZ)
PELAEZ	20	PRL 124 172001	J.R. Pelaez <i>et al.</i>
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira
ABLIKIM	11B	PL B698 183	M. Ablikim <i>et al.</i> (BES II Collab.)
GUO	11B	PR D84 034005	Z.-H. Guo, J.A. Oller
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i> (BES II Collab.)
BUGG	10	PR D81 014002	D.V. Bugg (LOQM)
LINK	09	PL B681 14	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i> (CLEO Collab.)
AUBERT	07T	PR D76 011102	B. Aubert <i>et al.</i> (BABAR Collab.)
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i> (BELLE Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i> (BES Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
Also		PR D74 059901 (errat.)	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
BUGG	06	PL B632 471	D.V. Bugg (LOQM)
CAWLFIELD	06A	PR D74 031108	C. Cawfield <i>et al.</i> (CLEO Collab.)
DESCOTES-G...	06	EPJ C48 553	S. Descotes-Genon, B. Moussallam
GUO	06	NP A773 78	F.K. Guo <i>et al.</i>
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng
LINK	05I	PL B621 72	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>
BUGG	03	PL B572 1	D.V. Bugg
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i> (FNAL E791 Collab.)
LINK	02E	PL B535 43	J.M. Link <i>et al.</i> (FNAL FOCUS Collab.)
KOPP	01	PR D63 092001	S. Kopp <i>et al.</i> (CLEO Collab.)
ISHIDA	97B	PTP 98 621	S. Ishida <i>et al.</i>
ASTON	88	NP B296 493	D. Aston <i>et al.</i> (SLAC, NAGO, CINC, INUS)
VANBEVEREN	86	ZPHY C30 615	E. van Beveren <i>et al.</i> (NIJM, BIEL)
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i> (MCGI, CARL, DURH+)
BINGHAM	72	NP B41 1	H.H. Bingham <i>et al.</i> (International K^+ Collab.)
MERCER	71	NP B32 381	R. Mercer <i>et al.</i> (JHU)
ROY	71	PL 36B 353	S.M. Roy

REFID=62088
 REFID=60559
 REFID=57836
 REFID=53683
 REFID=58808
 REFID=53361
 REFID=53213
 REFID=53056
 REFID=52426
 REFID=51726
 REFID=51929
 REFID=51875
 REFID=51037
 REFID=51051
 REFID=51458
 REFID=50996
 REFID=51153
 REFID=51518
 REFID=51164
 REFID=51198
 REFID=50679
 REFID=50165
 REFID=49586
 REFID=48807
 REFID=48728
 REFID=48134
 REFID=48655
 REFID=40262
 REFID=45769
 REFID=22443
 REFID=22415
 REFID=22412
 REFID=51107

 $K^*(892)$

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M018

 $K^*(892)$ T-Matrix Pole \sqrt{s}

NODE=M018TMP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
(890 ± 14) - i (26 ± 6) OUR ESTIMATE			
(890 ± 2) - i (25.6 ± 1.2)	¹ PELAEZ 20	RVUE	$\pi K \rightarrow \pi K$
(892 ± 1) - i (29 ± 1)	² PELAEZ 17	RVUE	$\pi K \rightarrow \pi K$
(889 ± 13) - i (24 ± 4)	³ PELAEZ 04A	RVUE	$\pi K \rightarrow \pi K$

NODE=M018TMP

→ UNCHECKED ←

OCCUR=2

¹ Extracted employing πK partial wave analysis from ESTABROOKS 78 and ASTON 88, Roy-Steiner equations and once subtracted forward dispersion relations.

NODE=M018TMP;LINKAGE=A

² Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M018TMP;LINKAGE=E

³ Reanalysis of data from ESTABROOKS 78 and ASTON 88 in the unitarized ChPT model.

NODE=M018TMP;LINKAGE=B

 $K^*(892)$ MASS

NODE=M018205

CHARGED ONLY, HADROPRODUCED

NODE=M018M1
 NODE=M018M1

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
891.67 ± 0.26 OUR AVERAGE					
892.2 ± 0.5 ± 1.7		ALBRECHT 20	CBAR		0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
892.6 ± 0.5	5840	BAUBILLIER 84B	HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
888 ± 3		NAPIER 84	SPEC	+	200 $\pi^- p \rightarrow 2K_S^0 X$
891 ± 1		NAPIER 84	SPEC	-	200 $\pi^- p \rightarrow 2K_S^0 X$
891.7 ± 2.1	3700	BARTH 83	HBC	+	70 $K^+ p \rightarrow K^0 \pi^+ X$
891 ± 1	4100	TOAFF 81	HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892.8 ± 1.6		AJINENKO 80	HBC	+	32 $K^+ p \rightarrow K^0 \pi^+ X$
890.7 ± 0.9	1800	AGUILAR-... 78B	HBC	±	0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
886.6 ± 2.4	1225	BALAND 78	HBC	±	12 $\bar{p}p \rightarrow (K\pi)^\pm X$

OCCUR=2

891.7 ±0.6	6706	COOPER	78	HBC	±	0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$	
891.9 ±0.7	9000	¹ PALER	75	HBC	-	14.3 $K^- p \rightarrow (K\pi)^- X$	
892.2 ±1.5	4404	AGUILAR...	71B	HBC	-	3.9,4.6 $K^- p \rightarrow (K\pi)^- p$	
891 ±2	1000	CRENNELL	69D	DBC	-	3.9 $K^- N \rightarrow K^0 \pi^- X$	
890 ±3.0	720	BARLOW	67	HBC	±	1.2 $\bar{p}p \rightarrow (K^0 \pi)^\pm K^\mp$	
889 ±3.0	600	BARLOW	67	HBC	±	1.2 $\bar{p}p \rightarrow (K^0 \pi)^\pm K \pi$	OCCUR=2
891 ±2.3	620	² DEBAERE	67B	HBC	+	3.5 $K^+ p \rightarrow K^0 \pi^+ p$	
891.0 ±1.2	1700	³ WOJCICKI	64	HBC	-	1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

893.6 ±0.1 ^{+0.2} _{-0.3}	183k	ABLIKIM	19AQ	BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$	
895.6 ±0.8	4k	⁴ LEES	17C	BABR		$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	
893.2 ±0.1 ±1.0	190k	⁵ AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$	
893.5 ±1.1	27k	⁶ ABELE	99D	CBAR	±	0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$	
890.4 ±0.2 ±0.5	80k	⁷ BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
890.0 ±2.3	800	^{2,3} CLELAND	82	SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$	
896.0 ±1.1	3200	^{2,3} CLELAND	82	SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$	OCCUR=2
893 ±1	3600	^{2,3} CLELAND	82	SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$	OCCUR=3
896.0 ±1.9	380	DELFOSSÉ	81	SPEC	+	50 $K^\pm p \rightarrow K^\pm \pi^0 p$	
886.0 ±2.3	187	DELFOSSÉ	81	SPEC	-	50 $K^\pm p \rightarrow K^\pm \pi^0 p$	OCCUR=2
894.2 ±2.0	765	² CLARK	73	HBC	-	3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
894.3 ±1.5	1150	^{2,3} CLARK	73	HBC	-	3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$	OCCUR=2
892.0 ±2.6	341	² SCHWEING...	68	HBC	-	5.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$	OCCUR=2

¹ Inclusive reaction. Complicated background and phase-space effects.

² Mass errors enlarged by us to Γ/\sqrt{N} . See note.

³ Number of events in peak reevaluated by us.

⁴ From a Dalitz plot analysis in an isobar model with charged and neutral K^* (892) masses and widths floating.

⁵ Average of fit results with different parametrizations for the $K\pi$ S-wave.

⁶ K-matrix pole.

⁷ From a partial wave amplitude analysis.

NODE=M018M;LINKAGE=I
 NODE=M018M;LINKAGE=D
 NODE=M018M;LINKAGE=W
 NODE=M018M1;LINKAGE=B

NODE=M018M1;LINKAGE=A
 NODE=M018M1;LINKAGE=AN
 NODE=M018M1;LINKAGE=F

NODE=M018MCT
 NODE=M018MCT

CHARGED ONLY, PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
895.47 ±0.20 ±0.74	53k	¹ EPIFANOV	07	BELL $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

892.0 ±0.5		² BOITO	10	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
892.0 ±0.9		^{3,4} BOITO	09	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
895.3 ±0.2		^{4,5} JAMIN	08	RVUE $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
896.4 ±0.9	12k	⁶ BONVICINI	02	CLEO $\tau^- \rightarrow K^- \pi^0 \nu_\tau$
895 ±2		⁷ BARATE	99R	ALEP $\tau^- \rightarrow K^- \pi^0 \nu_\tau$

¹ From a fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model.

² From the pole position of the $K\pi$ vector form factor using EPIFANOV 07 and constraints from K_{J3} decays in ANTONELLI 10.

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

⁴ Systematic uncertainties not estimated.

⁵ Reanalysis of EPIFANOV 07 using resonance chiral theory.

⁶ Calculated by us from the shift by 4.7 ± 0.9 MeV (statistical uncertainty only) reported in BONVICINI 02 with respect to the world average value from PDG 00.

⁷ With mass and width of the $K^*(1410)$ fixed at 1412 MeV and 227 MeV, respectively.

NODE=M018MCT;LINKAGE=EF
 NODE=M018MCT;LINKAGE=BT

NODE=M018MCT;LINKAGE=BI

NODE=M018MCT;LINKAGE=NS
 NODE=M018MCT;LINKAGE=JA
 NODE=M018MCT;LINKAGE=BO

NODE=M018MCT;LINKAGE=BA

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
895.55 ±0.20 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
894.68 ±0.25 ±0.05		¹ ABLIKIM	16F	BES3 $D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.4 ±0.2 ±0.2	243k	² DEL-AMO-SA..11I	BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
895.7 ±0.2 ±0.3	141k	³ BONVICINI	08A	CLEO $D^+ \rightarrow K^- \pi^+ \pi^+$
895.41 ±0.32 ^{+0.35} _{-0.43}	18k	⁴ LINK	05I	FOCS $D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
896 ±2		BARBERIS	98E	OMEG 450 $pp \rightarrow p_f p_s K^* \bar{K}^*$
895.9 ±0.5 ±0.2		ASTON	88	LASS 11 $K^- p \rightarrow K^- \pi^+ n$
894.52 ±0.63	25k	⁵ ATKINSON	86	OMEG 20-70 γp

NODE=M018M2
 NODE=M018M2

894.63±0.76	20k	⁵ ATKINSON	86	OMEG	20-70	γp		OCCUR=2
897 ±1	28k	EVANGELIS...	80	OMEG	10	$\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$		
898.4 ±1.4	1180	AGUILAR-...	78B	HBC	0.76	$\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$		
894.9 ±1.6		WICKLUND	78	ASPK	3,4,6	$K^\pm N \rightarrow (K\pi)^0 N$		
897.6 ±0.9		BOWLER	77	DBC	5.4	$K^+ d \rightarrow K^+ \pi^- p p$		
895.5 ±1.0	3600	MCCUBBIN	75	HBC	3.6	$K^- p \rightarrow K^- \pi^+ n$		
897.1 ±0.7	22k	⁵ PALER	75	HBC	14.3	$K^- p \rightarrow (K\pi)^0 X$		
896.0 ±0.6	10k	FOX	74	RVUE	2	$K^- p \rightarrow K^- \pi^+ n$		
896.0 ±0.6		FOX	74	RVUE	2	$K^+ n \rightarrow K^+ \pi^- p$		OCCUR=2
896 ±2		⁶ MATISON	74	HBC	12	$K^+ p \rightarrow K^+ \pi^- \Delta$		
896 ±1	3186	LEWIS	73	HBC	2.1-2.7	$K^+ p \rightarrow K \pi \pi p$		
894.0 ±1.3		⁶ LINGLIN	73	HBC	2-13	$K^+ p \rightarrow$ $K^+ \pi^- \pi^+ p$		
898.4 ±1.3	1700	⁷ BUCHNER	72	DBC	4.6	$K^+ n \rightarrow K^+ \pi^- p$		
897.9 ±1.1	2934	⁷ AGUILAR-...	71B	HBC	3.9,4.6	$K^- p \rightarrow K^- \pi^+ n$		
898.0 ±0.7	5362	⁷ AGUILAR-...	71B	HBC	3.9,4.6	$K^- p \rightarrow$ $K^- \pi^+ \pi^- p$		OCCUR=2
895 ±1	4300	⁸ HABER	70	DBC	3	$K^- N \rightarrow K^- \pi^+ X$		
893.7 ±2.0	10k	DAVIS	69	HBC	12	$K^+ p \rightarrow K^+ \pi^- \pi^+ p$		
894.7 ±1.4	1040	⁷ DAUBER	67B	HBC	2.0	$K^- p \rightarrow K^- \pi^+ \pi^- p$		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●								
895.50±0.92±2.6		⁹ ADUSZKIEW...	20A	NA61	158	$p p$		
898.1 ±1.0	4k	¹⁰ LEES	17C	BABR	J/ψ	$\rightarrow K_S^0 K^\pm \pi^\mp$		
895.53±0.17		LEES	13F	BABR	D^+	$\rightarrow K^+ K^- \pi^+$		
894.9 ±0.5 ±0.7	14.4k	¹¹ MITCHELL	09A	CLEO	D_s^+	$\rightarrow K^+ K^- \pi^+$		
896.2 ±0.3	20k	¹² AUBERT	07AK	BABR	10.6	$e^+ e^- \rightarrow$ $K^{*0} K^\pm \pi^\mp \gamma$		
900.7 ±1.1	5900	BARTH	83	HBC	70	$K^+ p \rightarrow K^+ \pi^- X$		

¹ Taking also into account the $K_0^*(1430)^0$ and $K_2^*(1430)^0$.

² Taking into account the $K^*(892)^0$, S-wave and P-wave ($K^*(1410)^0$).

³ From the isobar model with a complex pole for the κ .

⁴ Fit to $K \pi$ mass spectrum includes a non-resonant scalar component.

⁵ Inclusive reaction. Complicated background and phase-space effects.

⁶ From pole extrapolation.

⁷ Mass errors enlarged by us to Γ/\sqrt{N} . See note.

⁸ Number of events in peak reevaluated by us.


⁹ For transverse momenta between 0.6 and 0.8 GeV/c and rapidity $0 < y < 0.5$.

¹⁰ From a Dalitz plot analysis in an isobar model with charged and neutral $K^*(892)$ masses and widths floating.

¹¹ This value comes from a fit with χ^2 of 178/117.

¹² Systematic uncertainties not estimated.

NODE=M018M2;LINKAGE=B
 NODE=M018M2;LINKAGE=DE
 NODE=M018M2;LINKAGE=BO
 NODE=M018M2;LINKAGE=LI
 NODE=M018M2;LINKAGE=I
 NODE=M018M;LINKAGE=C
 NODE=M018M2;LINKAGE=D
 NODE=M018M2;LINKAGE=W
 NODE=M018M2;LINKAGE=C
 NODE=M018M2;LINKAGE=A
 NODE=M018M2;LINKAGE=MI
 NODE=M018M2;LINKAGE=NS



K*(892) MASSES AND MASS DIFFERENCES

NODE=M018209

Unrealistically small errors have been reported by some experiments. We use simple “realistic” tests for the minimum errors on the determination of a mass and width from a sample of N events:

$$\delta_{\min}(m) = \frac{\Gamma}{\sqrt{N}}, \quad \delta_{\min}(\Gamma) = 4 \frac{\Gamma}{\sqrt{N}} . \quad (1)$$

We consistently increase unrealistic errors before averaging. For a detailed discussion, see the 1971 edition of this Note.

$m_{K^*(892)^0} - m_{K^*(892)^\pm}$					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
6.7±1.2 OUR AVERAGE					
7.7±1.7	2980	AGUILAR-... 78B	HBC	±0	0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
5.7±1.7	7338	AGUILAR-... 71B	HBC	-0	3.9,4.6 $K^- p$
6.3±4.1	283	¹ BARASH 67B	HBC		0.0 $\bar{p}p$

¹Number of events in peak reevaluated by us.

NODE=M018D

NODE=M018D

NODE=M018D;LINKAGE=W

K*(892) RANGE PARAMETER

All from partial wave amplitude analyses.

NODE=M018R

NODE=M018R

NODE=M018R

VALUE (GeV ⁻¹)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2.1 ±0.5 ±0.5	243k	¹ DEL-AMO-SA.11I	BABR	0	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$
3.96±0.54 ^{+1.31} _{-0.90}	18k	² LINK 05I	FOCS	0	$D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$
3.4 ±0.7		ASTON 88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
••• We do not use the following data for averages, fits, limits, etc. •••					
12.1 ±3.2 ±3.0		BIRD 89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

¹ Taking into account the $K^*(892)^0$, S -wave and P -wave ($K^*(1410)^0$).

² Fit to $K\pi$ mass spectrum includes a non-resonant scalar component.

NODE=M018R;LINKAGE=DE
NODE=M018R;LINKAGE=LI

$K^*(892)$ WIDTH

NODE=M018215

CHARGED ONLY, HADROPRODUCED

NODE=M018W1
NODE=M018W1

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

51.4±0.8 OUR FIT

51.4±0.8 OUR AVERAGE

54.4±0.9±1.7		ALBRECHT	20	CBAR	0.9 $\bar{p}p \rightarrow K^+ K^- \pi^0$
49 ±2	5840	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
56 ±4		NAPIER	84	SPEC	- 200 $\pi^- p \rightarrow 2K_S^0 X$
51 ±2	4100	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
50.5±5.6		AJINENKO	80	HBC	+ 32 $K^+ p \rightarrow K^0 \pi^+ X$
45.8±3.6	1800	AGUILAR-...	78B	HBC	± 0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
52.0±2.5	6706	¹ COOPER	78	HBC	± 0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$
52.1±2.2	9000	² PALER	75	HBC	- 14.3 $K^- p \rightarrow (K\pi)^- X$
46.3±6.7	765	¹ CLARK	73	HBC	- 3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$
48.2±5.7	1150	^{1,3} CLARK	73	HBC	- 3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$
54.3±3.3	4404	¹ AGUILAR-...	71B	HBC	- 3.9,4.6 $K^- p \rightarrow (K\pi)^- p$
46 ±5	1700	^{1,3} WOJCICKI	64	HBC	- 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.7±0.2 ^{+0.1} _{-0.2}	183k	ABLIKIM	19AQ	BES	± $J/\psi \rightarrow K^+ K^- \pi^0$
43.6±1.3	4k	⁴ LEES	17C	BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
47.2±0.3±2.3	190k	⁵ AAIJ	16N	LHCB	$D^0 \rightarrow K_S^0 K^\pm \pi^\mp$
54.8±1.7	27k	⁶ ABELE	99D	CBAR	± 0.0 $\bar{p}p \rightarrow K^+ K^- \pi^0$
45.2±1 ±2	80k	⁷ BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
42.8±7.1	3700	BARTH	83	HBC	+ 70 $K^+ p \rightarrow K^0 \pi^+ X$
64.0±9.2	800	^{1,3} CLELAND	82	SPEC	+ 30 $K^+ p \rightarrow K_S^0 \pi^+ p$
62.0±4.4	3200	^{1,3} CLELAND	82	SPEC	+ 50 $K^+ p \rightarrow K_S^0 \pi^+ p$
55 ±4	3600	^{1,3} CLELAND	82	SPEC	- 50 $K^+ p \rightarrow K_S^0 \pi^- p$
62.6±3.8	380	DELFOSSÉ	81	SPEC	+ 50 $K^\pm p \rightarrow K^\pm \pi^0 p$
50.5±3.9	187	DELFOSSÉ	81	SPEC	- 50 $K^\pm p \rightarrow K^\pm \pi^0 p$

OCCUR=2

OCCUR=3

OCCUR=2

¹ Width errors enlarged by us to $4 \times \Gamma/\sqrt{N}$; see note.

² Inclusive reaction. Complicated background and phase-space effects.

³ Number of events in peak reevaluated by us.

⁴ From a Dalitz plot analysis in an isobar model with charged and neutral $K^*(892)$ masses and widths floating.

⁵ Average of fit results with different parametrizations for the $K\pi$ S -wave.

⁶ K -matrix pole.

⁷ From a partial wave amplitude analysis.

NODE=M018W;LINKAGE=D
NODE=M018W;LINKAGE=I
NODE=M018W;LINKAGE=W
NODE=M018W1;LINKAGE=B

NODE=M018W1;LINKAGE=A
NODE=M018W1;LINKAGE=AN
NODE=M018W1;LINKAGE=F

CHARGED ONLY, PRODUCED IN τ LEPTON DECAYS

NODE=M018W5
NODE=M018W5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

46.2±0.6±1.2 53k ¹ EPIFANOV 07 BELL $\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$

• • • We do not use the following data for averages, fits, limits, etc. • • •

46.5±1.1		² BOITO	10	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
46.2±0.4		^{3,4} BOITO	09	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
47.5±0.4		^{4,5} JAMIN	08	RVUE	$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
55 ±8		⁶ BARATE	99R	ALEP	$\tau^- \rightarrow K^- \pi^0 \nu_\tau$

¹ From a fit in the $K_0^*(700) + K^*(892) + K^*(1410)$ model.

² From the pole position of the $K\pi$ vector form factor using EPIFANOV 07 and constraints from K_{J3} decays in ANTONELLI 10.

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

⁴ Systematic uncertainties not estimated.

⁵ Reanalysis of EPIFANOV 07 using resonance chiral theory.

⁶ With mass and width of the $K^*(1410)$ fixed at 1412 MeV and 227 MeV, respectively.

NODE=M018W5;LINKAGE=EF
NODE=M018W5;LINKAGE=BT

NODE=M018W5;LINKAGE=BI

NODE=M018W5;LINKAGE=NS
NODE=M018W5;LINKAGE=JA
NODE=M018W5;LINKAGE=BA

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

47.3 ±0.5 OUR FIT Error includes scale factor of 2.0. [47.3 ± 0.5 MeV OUR 2023 FIT Scale factor = 1.9]

47.3 ±0.5 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.

46.53 ±0.56 ±0.31		¹ ABLIKIM	16F	BES3	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$	
46.5 ±0.3 ±0.2	243k	² DEL-AMO-SA..11I	BABR	BABR	$D^+ \rightarrow K^- \pi^+ e^+ \nu_e$	
45.3 ±0.5 ±0.6	141k	³ BONVICINI	08A	CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$	
47.79 ±0.86 $^{+1.32}_{-1.06}$	18k	⁴ LINK	05I	FOCS	$D^+ \rightarrow K^- \pi^+ \mu^+ \nu_\mu$	
54 ±3		BARBERIS	98E	OMEG	450 $pp \rightarrow p_f p_s K^* \bar{K}^*$	
50.8 ±0.8 ±0.9		ASTON	88	LASS	11 $K^- p \rightarrow K^- \pi^+ n$	
46.5 ±4.3	5900	BARTH	83	HBC	70 $K^+ p \rightarrow K^+ \pi^- X$	
54 ±2	28k	EVANGELIS...	80	OMEG	10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$	
45.9 ±4.8	1180	AGUILAR-...	78B	HBC	0.76 $\bar{p} p \rightarrow K^\mp K_S^0 \pi^\pm$	
51.2 ±1.7		WICKLUND	78	ASPK	3,4,6 $K^\pm N \rightarrow (K\pi)^0 N$	
48.9 ±2.5		BOWLER	77	DBC	5.4 $K^+ d \rightarrow K^+ \pi^- pp$	
48 $^{+3}_{-2}$	3600	MCCUBBIN	75	HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$	
50.6 ±2.5	22k	⁵ PALER	75	HBC	14.3 $K^- p \rightarrow (K\pi)^0 X$	
47 ±2	10k	FOX	74	RVUE	2 $K^- p \rightarrow K^- \pi^+ n$	
51 ±2		FOX	74	RVUE	2 $K^+ n \rightarrow K^+ \pi^- p$	
46.0 ±3.3	3186	⁶ LEWIS	73	HBC	2.1-2.7 $K^+ p \rightarrow K\pi\pi p$	OCCUR=2
51.4 ±5.0	1700	⁶ BUCHNER	72	DBC	4.6 $K^+ n \rightarrow K^+ \pi^- p$	
55.8 $^{+4.2}_{-3.4}$	2934	⁶ AGUILAR-...	71B	HBC	3.9,4.6 $K^- p \rightarrow K^- \pi^+ n$	
48.5 ±2.7	5362	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p \rightarrow K^- \pi^+ \pi^- p$	OCCUR=2
54.0 ±3.3	4300	^{6,7} HABER	70	DBC	3 $K^- N \rightarrow K^- \pi^+ X$	
53.2 ±2.1	10k	⁶ DAVIS	69	HBC	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$	
44 ±5.5	1040	⁶ DAUBER	67B	HBC	2.0 $K^- p \rightarrow K^- \pi^+ \pi^- p$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
48.8 ±1.8 ±2.0		⁸ ADUSZKIEW...20A	NA61	NA61	158 pp	
52.6 ±1.7	4k	⁹ LEES	17C	BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$	
44.90 ±0.30		LEES	13F	BABR	$D^+ \rightarrow K^+ K^- \pi^+$	
45.7 ±1.1 ±0.5	14.4k	¹⁰ MITCHELL	09A	CLEO	$D_s^+ \rightarrow K^+ K^- \pi^+$	
50.6 ±0.9	20k	¹¹ AUBERT	07AK	BABR	10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$	

¹ Taking also into account the $K_0^*(1430)^0$ and $K_2^*(1430)^0$.

² Taking into account the $K^*(892)^0$, S -wave and P -wave ($K^*(1410)^0$).

³ From the isobar model with a complex pole for the κ .

⁴ Fit to $K\pi$ mass spectrum includes a non-resonant scalar component.

⁵ Inclusive reaction. Complicated background and phase-space effects.

⁶ Width errors enlarged by us to $4 \times \Gamma/\sqrt{N}$; see note.

⁷ Number of events in peak reevaluated by us.

⁸ For transverse momenta between 0.6 and 0.8 GeV/c and rapidity $0 < y < 0.5$.

⁹ From a Dalitz plot analysis in an isobar model with charged and neutral $K^*(892)$ masses and widths floating.

¹⁰ This value comes from a fit with χ^2 of 178/117.

¹¹ Systematic uncertainties not estimated.

NODE=M018W2

NODE=M018W2

NEW

OCCUR=2

OCCUR=2

NODE=M018W2;LINKAGE=B

NODE=M018W2;LINKAGE=DE

NODE=M018W2;LINKAGE=BO

NODE=M018W2;LINKAGE=LN

NODE=M018W2;LINKAGE=I

NODE=M018W2;LINKAGE=D

NODE=M018W2;LINKAGE=W


NODE=M018W2;LINKAGE=C

NODE=M018W2;LINKAGE=A

NODE=M018W2;LINKAGE=MI

NODE=M018W2;LINKAGE=NS

WEIGHTED AVERAGE
47.3±0.5 (Error scaled by 2.0)



K*(892) DECAY MODES

NODE=M018220;NODE=M018

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\pi$	~ 100	%
Γ_2 $(K\pi)^\pm$	(99.902 ± 0.009) %	
Γ_3 $(K\pi)^0$	(99.754 ± 0.021) %	
Γ_4 $K^0\gamma$	$(2.46 \pm 0.21) \times 10^{-3}$	
Γ_5 $K^\pm\gamma$	$(9.8 \pm 0.9) \times 10^{-4}$	
Γ_6 $K\pi\pi$	< 7	$\times 10^{-4}$ 95%

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=11
DESIG=12
DESIG=4
DESIG=3
DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 14 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.7$ for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c}
 x_5 \\
 \Gamma
 \end{array}
 \begin{array}{|c|}
 \hline
 -100 \\
 \hline
 17 \quad -17 \\
 \hline
 \end{array}
 \begin{array}{c}
 x_2 \\
 x_5
 \end{array}$$

Mode	Rate (MeV)
Γ_2 $(K\pi)^\pm$	51.4 ± 0.8
Γ_5 $K^\pm\gamma$	0.050 ± 0.005

DESIG=11
DESIG=3

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 23 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 68.4$ for 21 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_4	-100	
Γ	13	-13
	x_3	x_4

Mode	Rate (MeV)	Scale factor
Γ_3 ($K\pi^0$)	47.2 \pm 0.5	2.0
Γ_4 ($K^0\gamma$)	0.117 \pm 0.010	

DESIG=12

DESIG=4

$K^*(892)$ PARTIAL WIDTHS

NODE=M018225

$\Gamma(K^0\gamma)$						Γ_4
VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
116 \pm 10 OUR FIT						
116.5 \pm 9.9	584	CARLSMITH	86	SPEC	0	$K_L^0 A \rightarrow K_S^0 \pi^0 A$

NODE=M018W4
NODE=M018W4

$\Gamma(K^\pm\gamma)$						Γ_5
VALUE (keV)		DOCUMENT ID	TECN	CHG	COMMENT	
50 \pm 5 OUR FIT						
50 \pm 5 OUR AVERAGE						
48 \pm 11		BERG	83	SPEC	-	156 $K^- A \rightarrow \bar{K} \pi A$
51 \pm 5		CHANDLEE	83	SPEC	+	200 $K^+ A \rightarrow K \pi A$

NODE=M018W3
NODE=M018W3

$K^*(892)$ BRANCHING RATIOS

NODE=M018230

$\Gamma(K^0\gamma)/\Gamma_{\text{total}}$						Γ_4/Γ
VALUE (units 10^{-3})		DOCUMENT ID	TECN	CHG	COMMENT	
2.46 \pm 0.21 OUR FIT						
1.5 \pm 0.7		CARITHERS	75B	CNTR	0	8-16 $\bar{K}^0 A$

NODE=M018R3
NODE=M018R3

$\Gamma(K^\pm\gamma)/\Gamma_{\text{total}}$						Γ_5/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
0.98 \pm 0.09 OUR FIT						
<1.6	95	BEMPORAD	73	CNTR	+	10-16 $K^+ A$

NODE=M018R2
NODE=M018R2

$\Gamma(K\pi\pi)/\Gamma((K\pi)^\pm)$						Γ_6/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
< 7 \times 10⁻⁴	95	JONGEJANS	78	HBC	4	$K^- p \rightarrow p \bar{K}^0 2\pi$
<20 \times 10 ⁻⁴		WOJCICKI	64	HBC	-	1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NODE=M018R1
NODE=M018R1

$K^*(892)$ REFERENCES

NODE=M018

ADUSZKIEW... 20A	EPJ C80 460	A. Aduszkiewicz <i>et al.</i>	(CERN NA61 Collab.)	REFID=60631
ALBRECHT 20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)	REFID=60439
PELAEZ 20	PRL 124 172001	J.R. Pelaez <i>et al.</i>		REFID=60559
ABLIKIM 19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
LEES 17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
PELAEZ 17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira		REFID=57836
AAIJ 16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57273
ABLIKIM 16F	PR D94 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57307
LEES 13F	PR D87 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55127
DEL-AMO-SA... 11I	PR D83 072001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16493
ANTONELLI 10	EPJ C69 399	M. Antonelli <i>et al.</i>	(FlaviaNet Working Group)	REFID=53448
BOITO 10	JHEP 1009 031	D.R. Boito, R. Escribano, M. Jamin	(BARC)	REFID=53632
BOITO 09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin		REFID=52728

MITCHELL	09A	PR D79 072008	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52756
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=52426
JAMIN	08	PL B664 78	M. Jamin, A. Pich, J. Portoles		REFID=52285
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
EPIFANOV	07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)	REFID=51929
LINK	05I	PL B621 72	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)	REFID=50679
PELAEZ	04A	MPL A19 2879	J.R. Pelaez	(MADU)	REFID=50347
BONVICINI	02	PRL 88 111803	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=48701
PDG	00	EPJ C15 1	D.E. Groom <i>et al.</i>	(PDG Collab.)	REFID=47469
ABELE	99D	PL B468 178	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)	REFID=47401
BARATE	99R	EPJ C11 599	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47366
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)	REFID=46348
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ATKINSON	86	ZPHY C30 521	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=20564
CARLSMITH	86	PRL 56 18	D. Carlsmith <i>et al.</i>	(EFI, SACL)	REFID=22461
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
NAPIER	84	PL 149B 514	A. Napier <i>et al.</i>	(TUFTS, ARIZ, FNAL, FLOR+)	REFID=22460
BARTH	83	NP B223 296	M. Barth <i>et al.</i>	(BRUX, CERN, GENO, MONS+)	REFID=22456
BERG	83	Thesis UMI 83-21652	D.M. Berg	(ROCH)	REFID=22457
CHANDLEE	83	PRL 51 168	C. Handlee <i>et al.</i>	(ROCH, FNAL, MINN)	REFID=22458
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
DELFOSSÉ	81	NP B183 349	A. Delfosse <i>et al.</i>	(GEVA, LAUS)	REFID=21277
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
AJINENKO	80	ZPHY C5 177	I.V. Ajinenko <i>et al.</i>	(SERP, BRUX, MONS+)	REFID=22449
EVANGELIS...	80	NP B165 383	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=22450
AGUILAR-...	78B	NP B141 101	M. Aguilar-Benitez <i>et al.</i>	(MADR, TATA+)	REFID=22438
BALAND	78	NP B140 220	J.F. Baland <i>et al.</i>	(MONS, BELG, CERN+)	REFID=20369
COOPER	78	NP B136 365	A.M. Cooper <i>et al.</i>	(TATA, CERN, CDEF+)	REFID=22441
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
JONGEJANS	78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
WICKLUND	78	PR D17 1197	A.B. Wicklund <i>et al.</i>	(ANL)	REFID=20124
BOWLER	77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
CARITHERS	75B	PRL 35 349	W.C.J. Carithers <i>et al.</i>	(ROCH, MCGI)	REFID=22433
MCCUBBIN	75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
PALER	75	NP B96 1	K. Paler <i>et al.</i>	(RHEL, SACL, EPOL)	REFID=22435
FOX	74	NP B80 403	G.C. Fox, M.L. Griss	(CIT)	REFID=22430
MATISON	74	PR D9 1872	M.J. Matison <i>et al.</i>	(LBL)	REFID=22431
BEMPORAD	73	NP B51 1	C. Bemporad <i>et al.</i>	(CERN, ETH, LOIC)	REFID=22416
CLARK	73	NP B54 432	A.G. Clark, L. Lyons, D. Radojicic	(OXF)	REFID=22426
LEWIS	73	NP B60 283	P.H. Lewis <i>et al.</i>	(LOWC, LOIC, CDEF)	REFID=22427
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
BUCHNER	72	NP B45 333	K. Buchner <i>et al.</i>	(MPIM, CERN, BRUX)	REFID=22418
AGUILAR-...	71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
HABER	70	NP B17 289	B. Haber <i>et al.</i>	(REHO, SACL, BGNA, EPOL)	REFID=22406
CRENNELL	69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS	69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
SCHWEING...	68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
BARASH	67B	PR 156 1399	N. Barash <i>et al.</i>	(COLU)	REFID=20160
BARLOW	67	NC 50A 701	J. Barlow <i>et al.</i>	(CERN, CDEF, IRAD, LIVP)	REFID=20041
DAUBER	67B	PR 153 1403	P.M. Dauber <i>et al.</i>	(UCLA)	REFID=22389
DEBAERE	67B	NC 51A 401	W. de Baere <i>et al.</i>	(BRUX, CERN)	REFID=22390
WOJCICKI	64	PR 135 B484	S.G. Wojcicki	(LRL)	REFID=22379

$K_1(1270)$

$$I(J^P) = \frac{1}{2}(1^+)$$


NODE=M028

 $K_1(1270)$ MASS

NODE=M028MX

NODE=M028MX

VALUE (MeV) _____ DOCUMENT ID _____

1253±7 OUR AVERAGE Includes data from the 4 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.**PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE**NODE=M028M2
NODE=M028M2

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____

The data in this block is included in the average printed for a previous datablock.

1275±10 700 GAVILLET 78 HBC + 4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$ **PRODUCED BY K BEAMS**NODE=M028M3
NODE=M028M3

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ CHG _____ COMMENT _____

The data in this block is included in the average printed for a previous datablock.

1270±10 ¹ DAUM 81C CNTR - 63 $K^- p \rightarrow K^- 2\pi p$

••• We do not use the following data for averages, fits, limits, etc. •••

~ 1276 ² TORNQVIST 82B RVUE

~ 1300 VERGEEST 79 HBC - 4.2 $K^- p \rightarrow (\bar{K}\pi\pi)^- p$

1289±25 ³ CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

~ 1300 BRANDENB... 76 ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

~ 1270 OTTER 76 HBC - 10,14,16 $K^- p \rightarrow (\bar{K}\pi\pi)^- p$

1260 DAVIS 72 HBC + 12 $K^+ p$

1234±12 FIRESTONE 72B DBC + 12 $K^+ d$

¹ Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.² From a unitarized quark-model calculation.³ From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

NODE=M028M3;LINKAGE=DA

NODE=M028M3;LINKAGE=T

NODE=M028M3;LINKAGE=E

PRODUCED BY BEAMS OTHER THAN K MESONSNODE=M028M1
NODE=M028M1

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____

The data in this block is included in the average printed for a previous datablock.

1248.1 ± 3.3 ± 1.4 GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1289.81 ± 0.56 ± 1.66	894k	AAIJ	18A	LHCB	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
1279 ± 10	25k	¹ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
1294 ± 10	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
1300	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
1242 + 9 - 10		² ASTIER	69	HBC	$\bar{p} p$
1300	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

¹ Systematic errors not estimated.
² This was called the C meson.

NODE=M028M1;LINKAGE=AB
 NODE=M028M1;LINKAGE=A

PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M028MT
 NODE=M028MT

The data in this block is included in the average printed for a previous datablock.

1250 ± 50 OUR AVERAGE [1254 ± 50 MeV OUR 2023 AVERAGE]

NEW

1254 ± 33 ± 34	7k	ASNER	00B	CLEO	$\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
-----------------------	----	-------	-----	------	---

$K_1(1270)$ WIDTH


NODE=M028WX

VALUE (MeV)	DOCUMENT ID
-------------	-------------

NODE=M028WX
 → UNCHECKED ←

90 ± 20 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

101 ± 12 OUR AVERAGE Includes data from the 4 datablocks that follow this one. Error includes scale factor of 2.2. See the ideogram below.



PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE

NODE=M028W2
 NODE=M028W2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

75 ± 15	700	GAVILLET	78	HBC	$4.2 K^- p \rightarrow \Xi^- K \pi \pi$
----------------	-----	----------	----	-----	---

PRODUCED BY K BEAMS

NODE=M028W3
 NODE=M028W3

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

90 ± 8	¹ DAUM	81C	CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
~ 150	VERGEEST	79	HBC	-	4.2 $K^- p \rightarrow (\bar{K} \pi \pi)^- p$
150 ± 71	² CARNEGIE	77	ASPK	±	13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$
~ 200	BRANDENB...	76	ASPK	±	13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$
120	DAVIS	72	HBC	+	12 $K^+ p$
188 ± 21	FIRESTONE	72B	DBC	+	12 $K^+ d$

¹ Well described in the chiral unitary approach of GENG 07 with two poles at 1195 and 1284 MeV and widths of 246 and 146 MeV, respectively.

NODE=M028W3;LINKAGE=DA

² From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

NODE=M028W3;LINKAGE=E

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

The data in this block is included in the average printed for a previous datablock.

119.5 ± 5.2 ± 6.7		GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$
--------------------------	--	-------	----	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

116.11 ± 1.65 ± 2.96	894k	AAIJ	18A1	LHCB	$D^0 \rightarrow K^\mp \pi^\pm \pi^\pm \pi^\mp$
131 ± 21	25k	¹ ABLIIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
66 ± 15	310	RODEBACK	81	HBC	$4 \pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72	HBC	$4.5 \pi^- p \rightarrow \Lambda K 2\pi$
127 ⁺⁷ / ₋₂₅		ASTIER	69	HBC	$\bar{p} p$
60	45	CRENNELL	67	HBC	$6 \pi^- p \rightarrow \Lambda K 2\pi$

¹Systematic errors not estimated.NODE=M028W1
NODE=M028W1

NODE=M028W1;LINKAGE=AB

PRODUCED IN τ LEPTON DECAYS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

260 ⁺⁹⁰/₋₇₀ ± 80	7k	ASNER	00B	CLEO	$\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
---	----	-------	-----	------	---

NODE=M028WT
NODE=M028WT **$K_1(1270)$ DECAY MODES**

NODE=M028215;NODE=M028

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 $K\rho$	(38 ± 13) %	2.2
Γ_2 $K_0^*(1430)\pi$	(28 ± 4) %	
Γ_3 $K^*(892)\pi$	(21 ± 10) %	2.2
Γ_4 $K\omega$	(11.0 ± 2.0) %	
Γ_5 $K f_0(1370)$	(3.0 ± 2.0) %	
Γ_6 γK^0	seen	

DESIG=2
DESIG=7
DESIG=1
DESIG=5
DESIG=8
DESIG=9;OUR EST;→ UNCHECKED ← **$K_1(1270)$ PARTIAL WIDTHS**

NODE=M028220

 $\Gamma(K\rho)$ **Γ_1**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

57 ± 5	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
75 ± 6	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W5
NODE=M028W5 **$\Gamma(K_0^*(1430)\pi)$** **Γ_2**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

26 ± 6	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
--------	----------	-----	------	---	--

NODE=M028W7
NODE=M028W7 **$\Gamma(K^*(892)\pi)$** **Γ_3**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 ± 11	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
2 ± 2	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W4
NODE=M028W4 **$\Gamma(K\omega)$** **Γ_4**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 4	MAZZUCATO	79	HBC	+	4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
24 ± 3	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

NODE=M028W6
NODE=M028W6 **$\Gamma(K f_0(1370))$** **Γ_5**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 5	CARNEGIE	77B	ASPK	±	13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
--------	----------	-----	------	---	--

NODE=M028W8
NODE=M028W8

$\Gamma(\gamma K^0)$ Γ_6

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

 $73.2 \pm 6.1 \pm 28.3$

ALAVI-HARATI02B

KTEV

 $K + A \rightarrow K^* + A$ NODE=M028W9
NODE=M028W9 $K_1(1270)$ BRANCHING RATIOS

NODE=M028225

 $\Gamma(K\rho)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE

DOCUMENT ID

TECN

COMMENT

 0.38 ± 0.13 OUR FIT

Error includes scale factor of 2.2.

 0.42 ± 0.06 ¹ DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.584 ± 0.043

² GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

dominant

RODEBACK 81 HBC $4 \pi^- p \rightarrow \Lambda K 2\pi$ NODE=M028R2
NODE=M028R2 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE

DOCUMENT ID

TECN

COMMENT

 0.28 ± 0.04 ¹ DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0201 ± 0.0064

² GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$ NODE=M028R4
NODE=M028R4 $\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE

DOCUMENT ID

TECN

COMMENT

 0.21 ± 0.10 OUR FIT

Error includes scale factor of 2.2.

 0.16 ± 0.05 ¹ DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.171 ± 0.023

² GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$ NODE=M028R1
NODE=M028R1 $\Gamma(K^*(892)\pi)/\Gamma(K\rho)$ Γ_3/Γ_1

VALUE

DOCUMENT ID

TECN

COMMENT

 0.56 ± 0.29 OUR FIT

Error includes scale factor of 2.2.

 $0.99 \pm 0.15 \pm 0.18$

ABLIKIM

21U

BES3

 $D_s^+ \rightarrow \bar{K}_1(1270)^0 K^+$ NODE=M028R00
NODE=M028R00 $\Gamma(K\omega)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE

DOCUMENT ID

TECN

COMMENT

 0.11 ± 0.02 ¹ DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.225 ± 0.052

² GULER 11 BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$ NODE=M028R3
NODE=M028R3 $\Gamma(K\omega)/\Gamma(K\rho)$ Γ_4/Γ_1

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.30

95

RODEBACK

81

HBC

 $4 \pi^- p \rightarrow \Lambda K 2\pi$ NODE=M028R6
NODE=M028R6 $\Gamma(K f_0(1370))/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE

DOCUMENT ID

TECN

COMMENT

 0.03 ± 0.02 ¹ DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$ NODE=M028R5
NODE=M028R5D-wave/S-wave RATIO FOR $K_1(1270) \rightarrow K^*(892)\pi$ NODE=M028R9
NODE=M028R9

VALUE

DOCUMENT ID

TECN

COMMENT

 1.0 ± 0.7 ¹ DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$ ¹ Average from low and high t data.² Assuming that decays are saturated by the $K\rho$, $K_0^*(1430)\pi$, $K^*(892)\pi$, $K\omega$ decay modes and neglecting interference between them. The values $B(\omega \rightarrow \pi^+ \pi^-) = (1.53^{+0.11}_{-0.13})\%$ and $B(K_0^*(1430) \rightarrow K\pi) = (93 \pm 10)\%$ are used. Systematic uncertainties not estimated.NODE=M028R;LINKAGE=F
NODE=M028R1;LINKAGE=GU

$K_1(1270)$ REFERENCES

ABLIKIM	21U	PR D104 032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18A1	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
GENG	07	PR D75 014017	L.S. Geng <i>et al.</i>	
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
RODEBACK	81	ZPHY C9 9	S. Rodeback <i>et al.</i>	(CERN, CDEF, MADR+)
MAZZUCATO	79	NP B156 532	M. Mazzucato <i>et al.</i>	(CERN, ZEEM, NIJM+)
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)
GAVILLET	78	PL 76B 517	P. Gavillet <i>et al.</i>	(AMST, CERN, NIJM+)
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)
CARNEGIE	77B	PL 68B 287	R.K. Carnegie <i>et al.</i>	(SLAC)
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP
OTTER	76	NP B106 77	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP
CRENNELL	72	PR D6 1220	D.J. Crennell <i>et al.</i>	(BNL)
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)
ASTIER	69	NP B10 65	A. Astier <i>et al.</i>	(CDEF, CERN, IPNP, LIVP) JP
CRENNELL	67	PRL 19 44	D.J. Crennell <i>et al.</i>	(BNL) I

NODE=M028

REFID=61155
 REFID=59187
 REFID=53668
 REFID=51623
 REFID=51037
 REFID=48822
 REFID=47766
 REFID=20573
 REFID=22548
 REFID=22550
 REFID=20867
 REFID=22542
 REFID=22538
 REFID=22535
 REFID=22536
 REFID=22532
 REFID=22533
 REFID=22419
 REFID=22505
 REFID=22506
 REFID=22482
 REFID=22473

NODE=M064

 $K_1(1400)$

$$I(J^P) = \frac{1}{2}(1^+)$$

 $K_1(1400)$ MASS

NODE=M064M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1403 ± 7 OUR AVERAGE					
1463 ± 64 ± 68	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
1373 ± 14 ± 18		¹ ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1392 ± 18		BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
1410 ± 25		DAUM	81C	CNTR	- 63 $K^- p \rightarrow K^- 2\pi p$
1415 ± 15		ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1404 ± 10		² CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1418 ± 8	25k	³ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 1350		⁴ TORNQVIST	82B	RVUE	
~ 1400		VERGEEST	79	HBC	- 4.2 $K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 1400		BRANDENB...	76	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
1420		DAVIS	72	HBC	+ 12 $K^+ p$
1368 ± 18		FIRESTONE	72B	DBC	+ 12 $K^+ d$

NODE=M064M

NODE=M064M

¹ From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.² From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.³ Systematic errors not estimated.⁴ From a unitarized quark-model calculation.

NODE=M064M;LINKAGE=P
 NODE=M064M;LINKAGE=E
 NODE=M064M;LINKAGE=AB
 NODE=M064M;LINKAGE=T

 $K_1(1400)$ WIDTH

NODE=M064W


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
174 ± 13 OUR AVERAGE Error includes scale factor of 1.6. See the ideogram below.					
$300^{+370}_{-110} \pm 140$	7k	ASNER	00B	CLEO	± $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$
188 ± 54 ± 60		¹ ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
276 ± 65		BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
195 ± 25		DAUM	81C	CNTR	- 63 $K^- p \rightarrow K^- 2\pi p$
180 ± 10		ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
142 ± 16		² CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
152 ± 16	25k	³ ABLIKIM	06C	BES2	$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$
~ 200		VERGEEST	79	HBC	- 4.2 $K^- p \rightarrow (\bar{K}\pi\pi)^- p$
~ 160		BRANDENB...	76	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
80		DAVIS	72	HBC	+ 12 $K^+ p$
241 ± 30		FIRESTONE	72B	DBC	+ 12 $K^+ d$

NODE=M064W

NODE=M064W

- ¹ From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.
- ² From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.
- ³ Systematic errors not estimated.

NODE=M064W;LINKAGE=P
 NODE=M064W;LINKAGE=E
 NODE=M064W;LINKAGE=AB



$K_1(1400)$ DECAY MODES

NODE=M064215;NODE=M064

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	(94 \pm 6) %
Γ_2 $K\rho$	(3.0 \pm 3.0) %
Γ_3 $K f_0(1370)$	(2.0 \pm 2.0) %
Γ_4 $K\omega$	(1.0 \pm 1.0) %
Γ_5 $K_0^*(1430)\pi$	not seen
Γ_6 γK^0	seen
Γ_7 $K\phi$	seen

DESIG=1
 DESIG=2
 DESIG=8
 DESIG=5
 DESIG=7;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=10

$K_1(1400)$ PARTIAL WIDTHS

NODE=M064220

$\Gamma(K^*(892)\pi)$					Γ_1
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
117\pm10	CARNEGIE	77	ASPK	\pm 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	
$\Gamma(K\rho)$					Γ_2
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2\pm1	CARNEGIE	77	ASPK	\pm 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	
$\Gamma(K\omega)$					Γ_4
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
23\pm12	CARNEGIE	77	ASPK	\pm 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$	
$\Gamma(\gamma K^0)$					Γ_6
VALUE (keV)	DOCUMENT ID	TECN	COMMENT		
280 \pm 50 OUR AVERAGE [281 \pm 50 keV OUR 2023 AVERAGE]					
280.8\pm23.2\pm40.4	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$		

NODE=M064W1
 NODE=M064W1
 NODE=M064W2
 NODE=M064W2
 NODE=M064W5
 NODE=M064W5
 NODE=M064W6
 NODE=M064W6
 NEW

$K_1(1400)$ BRANCHING RATIOS

NODE=M064225

$\Gamma(K^*(892)\pi)/\Gamma_{total}$				Γ_1/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.94\pm0.06	¹ DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$

NODE=M064R1
 NODE=M064R1

¹ Average from low and high t data.

NODE=M064R1;LINKAGE=F

$\Gamma(K\rho)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.03 ± 0.03	¹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

¹ Average from low and high t data. Γ_2/Γ NODE=M064R2
NODE=M064R2

NODE=M064R2;LINKAGE=F

 $\Gamma(K f_0(1370))/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.02 ± 0.02	¹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

¹ Average from low and high t data. Γ_3/Γ NODE=M064R5
NODE=M064R5

NODE=M064R5;LINKAGE=F

 $\Gamma(K\omega)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.01 ± 0.01	¹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

¹ Average from low and high t data. Γ_4/Γ NODE=M064R3
NODE=M064R3

NODE=M064R3;LINKAGE=F

 $\Gamma(K\phi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	¹ AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 9.2 σ . Γ_7/Γ NODE=M064R00
NODE=M064R00

NODE=M064R00;LINKAGE=A

 $\Gamma(K_0^*(1430)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

¹ Average from low and high t data. Γ_5/Γ NODE=M064R4
NODE=M064R4

NODE=M064R4;LINKAGE=F

D-wave/S-wave RATIO FOR $K_1(1400) \rightarrow K^*(892)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.04 ± 0.01	¹ DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

¹ Average from low and high t data.NODE=M064R9
NODE=M064R9

NODE=M064R9;LINKAGE=F

 $K_1(1400)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)	REFID=48822
ASNER	00B	PR D62 072006	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=47766
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
TORNQVIST	82B	NP B203 268	N.A. Tornqvist	(HELS)	REFID=20573
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
VERGEEST	79	NP B158 265	J.S.M. Vergeest <i>et al.</i>	(NIJM, AMST, CERN+)	REFID=22542
CARNEGIE	77	NP B127 509	R.K. Carnegie <i>et al.</i>	(SLAC)	REFID=22535
BRANDENB...	76	PRL 36 703	G.W. Brandenburg <i>et al.</i>	(SLAC) JP	REFID=22532
DAVIS	72	PR D5 2688	P.J. Davis <i>et al.</i>	(LBL)	REFID=22505
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)	REFID=22506

NODE=M064

$K^*(1410)$

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M094

 $K^*(1410)$ T-MATRIX POLE \sqrt{s}

NODE=M094PP

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M094PP

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(1368 \pm 38) - i(106^{+48}_{-59})$	¹ PELAEZ	17	RVUE $\pi K \rightarrow \pi K$
--------------------------------------	---------------------	----	--------------------------------

OCCUR=2

¹ Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M094PP;LINKAGE=B

 $K^*(1410)$ MASS

NODE=M094M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M094M

1414 ± 15 OUR AVERAGE Error includes scale factor of 1.3.

$1380 \pm 21 \pm 19$		ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
$1420 \pm 7 \pm 10$		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1437 \pm 8 \pm 16$	190k	¹ AAIJ	16N	LHCB		$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
---------------------	------	-------------------	-----	------	--	---

OCCUR=2

$1426 \pm 8 \pm 24$	190k	² AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
---------------------	------	-------------------	-----	------	--	---

1276^{+72}_{-77}		^{3,4} BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
--------------------	--	----------------------	----	------	--	---

1367 ± 54		BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
---------------	--	------	----	------	---	--

1474 ± 25		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
---------------	--	------------	-----	-----	---	---

1500 ± 30		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
---------------	--	-------	----	-----	---	---

¹ Using a parametrization for the $K\pi$ S -wave similar to ASTON 88 with fixed resonance width.

NODE=M094M;LINKAGE=A

² Using a $K\pi$ S -wave parametrization with resonant and non-resonant contributions.

NODE=M094M;LINKAGE=C

³ From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

NODE=M094M;LINKAGE=BI

⁴ Systematic uncertainties not estimated.

NODE=M094M;LINKAGE=NS

 $K^*(1410)$ WIDTH

NODE=M094W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

NODE=M094W

232 ± 21 OUR AVERAGE Error includes scale factor of 1.1.

$176 \pm 52 \pm 22$		ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
---------------------	--	-------	----	------	---	------------------------------------

$240 \pm 18 \pm 12$		ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
---------------------	--	-------	----	------	---	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$210 \pm 20 \pm 60$	190k	¹ AAIJ	16N	LHCB		$D^0 \rightarrow (K_S^0 \pi^\mp) K^\pm$
---------------------	------	-------------------	-----	------	--	---

OCCUR=2

$270 \pm 20 \pm 40$	190k	¹ AAIJ	16N	LHCB		$D^0 \rightarrow K_S^0 (K^\pm \pi^\mp)$
---------------------	------	-------------------	-----	------	--	---

198^{+61}_{-87}		^{2,3} BOITO	09	RVUE		$\tau^- \rightarrow K_S^0 \pi^- \nu_\tau$
-------------------	--	----------------------	----	------	--	---

114 ± 101		BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
---------------	--	------	----	------	---	--

275 ± 65		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
--------------	--	------------	-----	-----	---	---

500 ± 100		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
---------------	--	-------	----	-----	---	---

¹ Using a $K\pi$ S -wave parametrization with resonant and non-resonant contributions.

NODE=M094W;LINKAGE=A

² From the pole position of the $K\pi$ vector form factor in the complex s -plane and using EPIFANOV 07 data.

NODE=M094W;LINKAGE=BI

³ Systematic uncertainties not estimated.

NODE=M094W;LINKAGE=NS

 $K^*(1410)$ DECAY MODES

NODE=M094215;NODE=M094

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K^*(892)\pi$	> 40 %	95%
Γ_2 $K\pi$	(6.6 ± 1.3) %	
Γ_3 $K\rho$	< 7 %	95%
Γ_4 γK^0	< 2.3 $\times 10^{-4}$	90%
Γ_5 $K\phi$	seen	

DESIG=2

DESIG=1

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

DESIG=5

$K^*(1410)$ PARTIAL WIDTHS $\Gamma(\gamma K^0)$ Γ_4

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<52.9	90	ALAVI-HARATI02B	KTEV	$K + A \rightarrow K^* + A$

NODE=M094217

NODE=M094W1
NODE=M094W1 **$K^*(1410)$ BRANCHING RATIOS** $\Gamma(K\rho)/\Gamma(K^*(892)\pi)$ Γ_3/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.17	95	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M094220

NODE=M094R1
NODE=M094R1 $\Gamma(K\pi)/\Gamma(K^*(892)\pi)$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.16	95	ASTON	84	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

NODE=M094R2
NODE=M094R2 $\Gamma(K\pi)/\Gamma_{total}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.066 ± 0.010 ± 0.008	ASTON 88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M094R3
NODE=M094R3 $\Gamma(K\phi)/\Gamma_{total}$ Γ_5/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	¹ AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M094R00
NODE=M094R00¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 7.7 σ .

NODE=M094R00;LINKAGE=A

 $K^*(1410)$ REFERENCES

AAIJ 21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
PELAEZ 17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira	
AAIJ 16N	PR D93 052018	R. Aaij <i>et al.</i>	(LHCb Collab.)
BOITO 09	EPJ C59 821	D.R. Boito, R. Escribano, M. Jamin	
EPIFANOV 07	PL B654 65	D. Epifanov <i>et al.</i>	(BELLE Collab.)
ALAVI-HARATI 02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)
BIRD 89	SLAC-332	P.F. Bird	(SLAC)
ASTON 88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON 87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON 84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP
BAUBILLIER 82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ETKIN 80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP
ESTABROOKS 78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)

NODE=M094

REFID=61150
REFID=57836
REFID=57273
REFID=52728
REFID=51929
REFID=48822
REFID=41002
REFID=40262
REFID=40234
REFID=22689
REFID=22551
REFID=22545
REFID=22443 $K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

NODE=M019

 $K_0^*(1430)$ T-MATRIX POLE \sqrt{s}

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M019PP

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

(1431 ± 6) - i (110 ± 19) ¹ PELAEZ 17 RVUE $\pi K \rightarrow \pi K$

NODE=M019PP

¹ Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M019PP;LINKAGE=A

 $K_0^*(1430)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M019M

1425 ± 50 OUR ESTIMATE

NODE=M019M

→ UNCHECKED ←

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1493 ± 4 ± 7	¹ AAIJ	23AH LHCb	$B^+ \rightarrow K^+(K_S^0 K\pi)$
1449 ± 17 ± 2	² LEES	21A BABR	$\eta_c(1S) \rightarrow \eta' K^+ K^-$
1438 ± 8 ± 4 5.4k	³ LEES	14E BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
1427 ± 4 ± 13	⁴ BUGG	10 RVUE	S-matrix pole
1466.6 ± 0.7 ± 3.4 141k	⁵ BONVICINI	08A CLEO	$D^+ \rightarrow K^- \pi^+ \pi^+$
~ 1412	⁶ LINK	07 FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
1461.0 ± 4.0 ± 2.1 54k	⁷ LINK	07B FOCS	$D^+ \rightarrow K^- \pi^+ \pi^+$
1406 ± 29	⁸ BUGG	06 RVUE	
1435 ± 6	⁹ ZHOU	06 RVUE	$K p \rightarrow K^- \pi^+ n$

1455 ± 20 ± 15	ABLIKIM	05Q	BES2	$\psi(2S) \rightarrow$ $\gamma\pi^+\pi^-K^+K^-$
1456 ± 8	10 ZHENG	04	RVUE	$K^-p \rightarrow K^-\pi^+n$
~ 1419	11 BUGG	03	RVUE	11 $K^-p \rightarrow K^-\pi^+n$
~ 1440	12 LI	03	RVUE	11 $K^-p \rightarrow K^-\pi^+n$
1459 ± 9	13 AITALA	02	E791	$D^+ \rightarrow K^-\pi^+\pi^+$
~ 1440	14 JAMIN	00	RVUE	$Kp \rightarrow Kp$
1436 ± 8	15 BARBERIS	98E	OMEG	450 $pp \rightarrow$ $p_f p_s K^+ K^-\pi^+\pi^-$
1415 ± 25	11 ANISOVICH	97C	RVUE	11 $K^-p \rightarrow K^-\pi^+n$
~ 1450	16 TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi$
1412 ± 6	17 ASTON	88	LASS	11 $K^-p \rightarrow K^-\pi^+n$
~ 1430	BAUBILLIER	84B	HBC	8.25 $K^-p \rightarrow \bar{K}^0\pi^-p$
~ 1425	18 ESTABROOKS	78	ASPK	13 $K^\pm p \rightarrow K^\pm\pi^\pm(n, \Delta)$
~ 1450.0	MARTIN	78	SPEC	10 $K^\pm p \rightarrow K_S^0\pi p$

¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c.$

² Using a $K\pi - K\eta'$ coupled channel Breit-Wigner function.

³ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed width $\Gamma(K_0^*(1430)) = 210$ MeV.

⁴ S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.

⁵ From the isobar model with a complex pole for the κ .

⁶ From a non-parametric analysis.

⁷ A Breit-Wigner mass and width.

⁸ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s-dependent width and an Adler zero near threshold.

⁹ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1950)$.

¹⁰ Using ASTON 88 and assuming $K_0^*(700)$.

¹¹ T-matrix pole. Reanalysis of ASTON 88 data.

¹² Breit-Wigner fit. Using ASTON 88.

¹³ Assuming a low-mass scalar $K\pi$ resonance, $\kappa(700)$.

¹⁴ T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.

¹⁵ J^P not determined, could be $K_2^*(1430)$.

¹⁶ T-matrix pole.

¹⁷ Uses a model for the background, without this background they get a mass 1340 MeV, where the phase shift passes 90° .

¹⁸ Mass defined by pole position. From elastic $K\pi$ partial-wave analysis.

NODE=M019M;LINKAGE=F
NODE=M019M;LINKAGE=C
NODE=M019M;LINKAGE=LE

NODE=M019M;LINKAGE=BG

NODE=M019M;LINKAGE=BO
NODE=M019M;LINKAGE=LI
NODE=M019M;LINKAGE=BW
NODE=M019M;LINKAGE=BU

NODE=M019M;LINKAGE=ZU
NODE=M019M;LINKAGE=ZH
NODE=M019M;LINKAGE=A1
NODE=M019M;LINKAGE=E
NODE=M019M;LINKAGE=A0
NODE=M019M;LINKAGE=JM
NODE=M019M;LINKAGE=JP
NODE=M019M;LINKAGE=TT
NODE=M019M;LINKAGE=D

NODE=M019M;LINKAGE=A

$K_0^*(1430)$ WIDTH

NODE=M019W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
270 ± 80				OUR ESTIMATE

NODE=M019W

→ UNCHECKED ←

• • • We do not use the following data for averages, fits, limits, etc. • • •

215 ± 7 ± 4	1 AAIJ	23AH	LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$
210 ± 20 ± 12	2 LEES	14E	BABR	$\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$
270 ± 10 ± 40	3 BUGG	10	RVUE	S-matrix pole
174.2 ± 1.9 ± 3.2	4 BONVICINI	08A	CLEO	$D^+ \rightarrow K^-\pi^+\pi^+$
~ 500	5 LINK	07	FOCS	$D^+ \rightarrow K^- K^+ \pi^+$
177.0 ± 8.0 ± 3.4	6 LINK	07B	FOCS	$D^+ \rightarrow K^-\pi^+\pi^+$
350 ± 40	7 BUGG	06	RVUE	
288 ± 22	8 ZHOU	06	RVUE	$Kp \rightarrow K^-\pi^+n$
270 ± 45 ⁺³⁰ -35	ABLIKIM	05Q	BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
217 ± 31	9 ZHENG	04	RVUE	$K^-p \rightarrow K^-\pi^+n$
~ 316	10 BUGG	03	RVUE	11 $K^-p \rightarrow K^-\pi^+n$
~ 350	11 LI	03	RVUE	11 $K^-p \rightarrow K^-\pi^+n$
175 ± 17	12 AITALA	02	E791	$D^+ \rightarrow K^-\pi^+\pi^+$
~ 300	13 JAMIN	00	RVUE	$Kp \rightarrow Kp$
196 ± 45	14 BARBERIS	98E	OMEG	450 $pp \rightarrow$ $p_f p_s K^+ K^-\pi^+\pi^-$
330 ± 50	10 ANISOVICH	97C	RVUE	11 $K^-p \rightarrow K^-\pi^+n$
~ 320	15 TORNQVIST	96	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi$
294 ± 23	ASTON	88	LASS	11 $K^-p \rightarrow K^-\pi^+n$
~ 200	BAUBILLIER	84B	HBC	8.25 $K^-p \rightarrow \bar{K}^0\pi^-p$
200 to 300	16 ESTABROOKS	78	ASPK	13 $K^\pm p \rightarrow K^\pm\pi^\pm(n, \Delta)$

- ¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c.$
- ² Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a likelihood scan in the presence of several interfering scalar-meson resonances with fixed mass $M(K_0^*(1430)) = 1435$ MeV.
- ³ S-Matrix pole. Supersedes BUGG 06. Combined analysis of ASTON 88, ABLIKIM 06C, AITALA 06, and LINK 09 using an s-dependent width with couplings to $K\pi$ and $K\eta'$, and the Adler zero near thresholds.
- ⁴ From the isobar model with a complex pole for the κ .
- ⁵ From a non-parametric analysis.
- ⁶ A Breit-Wigner mass and width.
- ⁷ S-matrix pole. Reanalysis of ASTON 88, AITALA 02, and ABLIKIM 06C including the κ with an s-dependent width and an Adler zero near threshold.
- ⁸ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1950)$.
- ⁹ Using ASTON 88 and assuming $K_0^*(700)$.
- ¹⁰ T-matrix pole. Reanalysis of ASTON 88 data.
- ¹¹ Breit-Wigner fit. Using ASTON 88.
- ¹² Assuming a low-mass scalar $K\pi$ resonance, $\kappa(700)$.
- ¹³ T-matrix pole. Using data from ESTABROOKS 78 and ASTON 88.
- ¹⁴ J^P not determined, could be $K_2^*(1430)$.
- ¹⁵ T-matrix pole.
- ¹⁶ From elastic $K\pi$ partial-wave analysis.

NODE=M019W;LINKAGE=A
 NODE=M019W;LINKAGE=LE
 NODE=M019W;LINKAGE=BG
 NODE=M019W;LINKAGE=BO
 NODE=M019W;LINKAGE=LI
 NODE=M019W;LINKAGE=BW
 NODE=M019W;LINKAGE=BU
 NODE=M019W;LINKAGE=ZU
 NODE=M019W;LINKAGE=ZH
 NODE=M019W;LINKAGE=A1
 NODE=M019W;LINKAGE=E
 NODE=M019W;LINKAGE=A0
 NODE=M019W;LINKAGE=JM
 NODE=M019W;LINKAGE=JP
 NODE=M019W;LINKAGE=TT
 NODE=M019W;LINKAGE=C

$K_0^*(1430)$ DECAY MODES

NODE=M019215;NODE=M019

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(93 \pm 10) %
Γ_2 $K\eta$	(8.6 $^{+2.7}_{-3.4}$) %
Γ_3 $K\eta'(958)$	seen

DESIG=1
 DESIG=2
 DESIG=3

$K_0^*(1430)$ BRANCHING RATIOS

NODE=M019220

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.93\pm0.04\pm0.09	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M019R1
 NODE=M019R1

$\Gamma(K\eta)/\Gamma(K\pi)$	Γ_2/Γ_1			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.2\pm2.5$^{+1.0}_{-2.5}$	5.4k	¹ LEES	14E	BABR $\eta_c(1S) \rightarrow K^+ K^- \eta/\pi^0$

NODE=M019R01
 NODE=M019R01

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$. From a Dalitz analysis in the presence of several interfering scalar-meson resonances.

NODE=M019R01;LINKAGE=LE

$\Gamma(K\eta'(958))/\Gamma_{\text{total}}$	Γ_3/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M019R00
 NODE=M019R00

$\Gamma(K\eta'(958))/\Gamma(K\pi)$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.397\pm0.064\pm0.054	¹ LEES	21A	BABR $\eta_c(1S) \rightarrow \eta' K^+ K^-$

NODE=M019R02
 NODE=M019R02

¹ Using $K\pi$ data from LEES 14E.

NODE=M019R02;LINKAGE=A

$K_0^*(1430)$ REFERENCES

NODE=M019

AAIJ	23AH	PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	21A	PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira	
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	14E	PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)
BUGG	10	PR D81 014002	D.V. Bugg	(LOQM)
LINK	09	PL B681 14	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
BONVICINI	08A	PR D78 052001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
LINK	07	PL B648 156	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
LINK	07B	PL B653 1	J.M. Link <i>et al.</i>	(FNAL FOCUS Collab.)
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)
AITALA	06	PR D73 032004	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
Also		PR D74 059901 (errata)	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
BUGG	06	PL B632 471	D.V. Bugg	(LOQM)
ZHOU	06	NP A775 212	Z.Y. Zhou, H.Q. Zheng	

REFID=62349
 REFID=61442
 REFID=57836
 REFID=55901
 REFID=55937
 REFID=53213
 REFID=53056
 REFID=52426
 REFID=51702
 REFID=51875
 REFID=51037
 REFID=51051
 REFID=51458
 REFID=50996
 REFID=51198

ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
ZHENG	04	NP A733 235	H.Q. Zheng <i>et al.</i>	
BUGG	03	PL B572 1	D.V. Bugg	
LI	03	PR D67 034025	L. Li, B. Zou, G. Li	
AITALA	02	PRL 89 121801	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
JAMIN	00	NP B587 331	M. Jamin <i>et al.</i>	
BARBERIS	98E	PL B436 204	D. Barberis <i>et al.</i>	(Omega Expt.)
ANISOVICH	97C	PL B413 137	A.V. Anisovich, A.V. Sarantsev	
TORNQVIST	96	PRL 76 1575	N.A. Tornqvist, M. Roos	(HELS)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)

REFID=50958
REFID=50165
REFID=49586
REFID=49192
REFID=48807
REFID=47983
REFID=46348
REFID=45815
REFID=44507
REFID=40262
REFID=22459
REFID=22443
REFID=22446

NODE=M022

$K_2^*(1430)$

$$I(J^P) = \frac{1}{2}(2^+)$$

We consider that phase-shift analyses provide more reliable determinations of the mass and width.

NODE=M022

$K_2^*(1430)$ T-MATRIX POLE \sqrt{s}

NODE=M022PP

VALUE (MeV) DOCUMENT ID TECN COMMENT

NODE=M022PP

••• We do not use the following data for averages, fits, limits, etc. •••

(1424 ± 4) - i(66 ± 2) ¹PELAEZ 17 RVUE $\pi K \rightarrow \pi K$

¹Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M022PP;LINKAGE=A

$K_2^*(1430)$ MASS

NODE=M022205

CHARGED ONLY, WITH FINAL STATE $K\pi$

NODE=M022M1
NODE=M022M1

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

1427.3 ± 1.5 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

1432.7 ± 0.7	$\frac{+2.2}{-2.3}$	183k	ABLIKIM	19AQ	BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
1420 ± 4		1587	BAUBILLIER	84B	HBC	-	$8.25 K^- p \rightarrow \bar{K}^0 \pi^- p$
1436 ± 5.5		400	^{1,2} CLELAND	82	SPEC	+	$30 K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2		1500	^{1,2} CLELAND	82	SPEC	+	$50 K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2		1200	^{1,2} CLELAND	82	SPEC	-	$50 K^+ p \rightarrow K_S^0 \pi^- p$
1423 ± 5		935	TOAFF	81	HBC	-	$6.5 K^- p \rightarrow \bar{K}^0 \pi^- p$
1428.0 ± 4.6			³ MARTIN	78	SPEC	+	$10 K^\pm p \rightarrow K_S^0 \pi p$
1423.8 ± 4.6			³ MARTIN	78	SPEC	-	$10 K^\pm p \rightarrow K_S^0 \pi p$
1420.0 ± 3.1		1400	AGUILAR-...	71B	HBC	-	$3.9, 4.6 K^- p$
1425 ± 8.0		225	^{1,2} BARNHAM	71C	HBC	+	$K^+ p \rightarrow K^0 \pi^+ p$
1416 ± 10		220	CRENNELL	69D	DBC	-	$3.9 K^- N \rightarrow \bar{K}^0 \pi^- N$
1414 ± 13.0		60	¹ LIND	69	HBC	+	$9 K^+ p \rightarrow K^0 \pi^+ p$
1427 ± 12		63	¹ SCHWEING...	68	HBC	-	$5.5 K^- p \rightarrow \bar{K} \pi N$
1423 ± 11.0		39	¹ BASSANO	67	HBC	-	$4.6-5.0 K^- p \rightarrow \bar{K}^0 \pi^- p$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

1428 ± 2		4300	⁴ ABLIKIM	22L	BES3		$2.0-3.08 e^+ e^- \rightarrow K^+ K^- \pi^0$
1423.4 ± 2 ± 3		24809 ± 820	⁵ BIRD	89	LASS	-	$11 K^- p \rightarrow \bar{K}^0 \pi^- p$

¹Errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

²Number of events in peak re-evaluated by us.

³Systematic error added by us.

⁴From a partial wave amplitude analysis at $\sqrt{s} = 2.125$ GeV which includes all the possible intermediate states that match J^{PC} conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.

⁵From a partial wave amplitude analysis.


NODE=M022M;LINKAGE=D

NODE=M022M;LINKAGE=W

NODE=M022M;LINKAGE=B

NODE=M022M1;LINKAGE=A

NODE=M022M;LINKAGE=F



NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1432.4 ± 1.3 OUR AVERAGE				
1431.2 ± 1.8 ± 0.7	1	ASTON 88	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 4 ± 6	1	ASTON 87	LASS	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1433 ± 6 ± 10	1	ASTON 84B	LASS	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
1471 ± 12	1	BAUBILLIER 82B	HBC	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
1428 ± 3	1	ASTON 81C	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 2	1	ESTABROOKS 78	ASPK	13 $K^\pm p \rightarrow p K \pi$
1440 ± 10	1	BOWLER 77	DBC	5.5 $K^+ d \rightarrow K \pi p p$
1428.5 ± 3.9	1786 ± 127	2 AUBERT 07AK	BABR	10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
1420 ± 7	300	HENDRICK 76	DBC	8.25 $K^+ N \rightarrow K^+ \pi N$
1421.6 ± 4.2	800	MCCUBBIN 75	HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
1420.1 ± 4.3		3 LINGLIN 73	HBC	2-13 $K^+ p \rightarrow K^+ \pi^- X$
1419.1 ± 3.7	1800	AGUILAR-... 71B	HBC	3.9,4.6 $K^- p$
1416 ± 6	600	CORDS 71	DBC	9 $K^+ n \rightarrow K^+ \pi^- p$
1421.1 ± 2.6	2200	DAVIS 69	HBC	12 $K^+ p \rightarrow K^+ \pi^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From phase shift or partial-wave analysis.

² Systematic errors not estimated.

³ From pole extrapolation, using world $K^+ p$ data summary tape.

NODE=M022M4
NODE=M022M4

NODE=M022M;LINKAGE=P
NODE=M022M4;LINKAGE=NS
NODE=M022M;LINKAGE=C

$K_2^*(1430)$ WIDTH

CHARGED ONLY, WITH FINAL STATE $K \pi$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
100.0 ± 2.2 OUR FIT					Error includes scale factor of 1.1. [100.0 ± 2.1 MeV OUR 2023 FIT]
100.0 ± 2.2 OUR AVERAGE					Error includes scale factor of 1.1.
102.5 ± 1.6 ^{+3.1} _{-2.8}	183k	ABLIKIM 19AQ	BES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
109 ± 22	400	1,2 CLELAND 82	SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
124 ± 12.8	1500	1,2 CLELAND 82	SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
113 ± 12.8	1200	1,2 CLELAND 82	SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$
85 ± 16	935	TOAFF 81	HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
96.5 ± 3.8		MARTIN 78	SPEC	+	10 $K^\pm p \rightarrow K_S^0 \pi p$
97.7 ± 4.0		MARTIN 78	SPEC	-	10 $K^\pm p \rightarrow K_S^0 \pi p$
94.7 ^{+15.1} _{-12.5}	1400	AGUILAR-... 71B	HBC	-	3.9,4.6 $K^- p$
107 ± 4	4300	3 ABLIKIM 22L	BES3		2.0-3.08 $e^+ e^- \rightarrow K^+ K^- \pi^0$
98 ± 4 ± 4	25k	4 BIRD 89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M022210

NODE=M022W1
NODE=M022W1
NEW

OCCUR=2
OCCUR=3

OCCUR=2

- ¹ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- ² Number of events in peak re-evaluated by us.
- ³ From a partial wave amplitude analysis at $\sqrt{s} = 2.125$ GeV which includes all the possible intermediate states that match J^{PC} conservation in the subsequent two-body decay. The intermediate states are parameterized with the relativistic Breit-Wigner functions. Statistical error only.
- ⁴ From a partial wave amplitude analysis.

NODE=M022W;LINKAGE=D
 NODE=M022W;LINKAGE=W
 NODE=M022W1;LINKAGE=A

NODE=M022W;LINKAGE=F


NODE=M022W4
 NODE=M022W4

NEUTRAL ONLY

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
109 ± 5 OUR AVERAGE				Error includes scale factor of 1.9. See the ideogram below.
116.5 ± 3.6 ± 1.7		¹ ASTON 88	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
129 ± 15 ± 15		¹ ASTON 87	LASS	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
131 ± 24 ± 20		¹ ASTON 84B	LASS	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
143 ± 34		¹ BAUBILLIER 82B	HBC	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
98 ± 8		¹ ASTON 81C	LASS	11 $K^- p \rightarrow K^- \pi^+ n$
140 ± 30		¹ ETKIN 80	SPEC	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
98 ± 5		¹ ESTABROOKS 78	ASPK	13 $K^\pm p \rightarrow pK\pi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
113.7 ± 9.2	1786 ± 127	² AUBERT 07AK	BABR	10.6 $e^+ e^- \rightarrow K^{*0} K^\pm \pi^\mp \gamma$
125 ± 29	300	³ HENDRICK 76	DBC	8.25 $K^+ N \rightarrow K^+ \pi N$
116 ± 18	800	MCCUBBIN 75	HBC	3.6 $K^- p \rightarrow K^- \pi^+ n$
61 ± 14		⁴ LINGLIN 73	HBC	2-13 $K^+ p \rightarrow K^+ \pi^- X$
116.6 ^{+10.3} _{-15.5}	1800	AGUILAR-...	71B	HBC 3.9,4.6 $K^- p$
144 ± 24.0	600	³ CORDS 71	DBC	9 $K^+ n \rightarrow K^+ \pi^- p$
101 ± 10	2200	DAVIS 69	HBC	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$

NODE=M022W;LINKAGE=P
 NODE=M022W4;LINKAGE=NS
 NODE=M022W4;LINKAGE=D
 NODE=M022W;LINKAGE=C

- ¹ From phase shift or partial-wave analysis.
- ² Systematic errors not estimated.
- ³ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- ⁴ From pole extrapolation, using world $K^+ p$ data summary tape.



$K_2^*(1430)$ DECAY MODES

NODE=M022215;NODE=M022

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $K\pi$	(49.9 ± 1.2) %	
Γ_2 $K^*(892)\pi$	(24.7 ± 1.5) %	
Γ_3 $K^*(892)\pi\pi$	(13.4 ± 2.2) %	
Γ_4 $K\rho$	(8.7 ± 0.8) %	S=1.2

DESIG=1
 DESIG=2
 DESIG=6
 DESIG=3

Γ_5	$K\omega$	$(2.9 \pm 0.8) \%$		DESIG=4
Γ_6	$K^+\gamma$	$(2.4 \pm 0.5) \times 10^{-3}$	S=1.1	DESIG=8
Γ_7	$K\eta$	$(1.5^{+3.4}_{-1.0}) \times 10^{-3}$	S=1.3	DESIG=5
Γ_8	$K\omega\pi$	$< 7.2 \times 10^{-4}$	CL=95%	DESIG=7
Γ_9	$K^0\gamma$	$< 9 \times 10^{-4}$	CL=90%	DESIG=10;OUR EVAL;→ UNCHECKED ←

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 10 branching ratios uses 32 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 21.1$ for 25 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-9						
x_3	-40	-73					
x_4	-8	36	-52				
x_5	-11	-3	-26	-7			
x_6	-1	-1	-1	-1	0		
x_7	-4	-7	-5	-5	-2	0	
Γ	0	0	0	0	0	-11	0
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $K\pi$	49.9 ± 1.6	DESIG=1
Γ_2 $K^*(892)\pi$	24.7 ± 1.6	DESIG=2
Γ_3 $K^*(892)\pi\pi$	13.5 ± 2.3	DESIG=6
Γ_4 $K\rho$	8.7 ± 0.8	1.2 DESIG=3
Γ_5 $K\omega$	2.9 ± 0.8	DESIG=4
Γ_6 $K^+\gamma$	0.24 ± 0.05	1.1 DESIG=8
Γ_7 $K\eta$	$0.15^{+0.34}_{-0.10}$	1.3 DESIG=5

$K_2^*(1430)$ PARTIAL WIDTHS

NODE=M022220

$\Gamma(K^+\gamma)$

 Γ_6

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
240 ± 50 OUR FIT				Error includes scale factor of 1.1. [241 ± 50 keV OUR 2023 FIT
Scale factor = 1.1]				NEW

240 ± 45	CIHANGIR	82	SPEC	+	200 $K^+Z \rightarrow ZK^+\pi^0$, $ZK_S^0\pi^+$
--------------------------------	----------	----	------	---	---

NODE=M022W8
NODE=M022W8
NEW

$\Gamma(K^0\gamma)$

 Γ_9

VALUE (keV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 5.4	90	ALAVI-HARATI02B	KTEV		$K + A \rightarrow K^* + A$
••• We do not use the following data for averages, fits, limits, etc. •••					
< 84	90	CARLSMITH	87	SPEC	0 $60-200 K_L^0 A \rightarrow$ $K_S^0\pi^0 A$

NODE=M022W9
NODE=M022W9

$K_2^*(1430)$ BRANCHING RATIOS

NODE=M022225

$\Gamma(K\pi) / \Gamma_{\text{total}}$

 Γ_1 / Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.499 ± 0.012 OUR FIT				
0.488 ± 0.014 OUR AVERAGE				
$0.485 \pm 0.006 \pm 0.020$	¹ ASTON	88	LASS	0 $11 K^- p \rightarrow K^- \pi^+ n$
0.49 ± 0.02	¹ ESTABROOKS	78	ASPK	\pm $13 K^\pm p \rightarrow pK\pi$

NODE=M022R1
NODE=M022R1¹ From phase shift analysis.

NODE=M022R;LINKAGE=P

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$

Γ_2/Γ_1

NODE=M022R4
NODE=M022R4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.496±0.034 OUR FIT				
0.47 ±0.04 OUR AVERAGE				
0.44 ±0.09	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
0.62 ±0.19	LAUSCHER	75	HBC	0 10,16 $K^- p \rightarrow K^- \pi^+ n$
0.54 ±0.16	DEHM	74	DBC	0 4.6 $K^+ N$
0.47 ±0.08	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
0.47 ±0.10	BASSANO	67	HBC	-0 4.6,5.0 $K^- p$
0.45 ±0.13	BADIER	65C	HBC	- 3 $K^- p$

$\Gamma(K\omega)/\Gamma(K\pi)$

Γ_5/Γ_1

NODE=M022R5
NODE=M022R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.059±0.017 OUR FIT				
0.070±0.035 OUR AVERAGE				
0.05 ±0.04	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
0.13 ±0.07	BASSOMPIE...	69	HBC	0 5 $K^+ p$

OCCUR=2

$\Gamma(K\rho)/\Gamma(K\pi)$

Γ_4/Γ_1

NODE=M022R6
NODE=M022R6

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.174±0.017 OUR FIT Error includes scale factor of 1.2.				
0.150^{+0.029}_{-0.017} OUR AVERAGE				
0.18 ±0.05	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$
0.02 ^{+0.10} _{-0.02}	DEHM	74	DBC	0 4.6 $K^+ N$
0.16 ±0.05	AGUILAR-...	71B	HBC	3.9,4.6 $K^- p$
0.14 ±0.10	BASSANO	67	HBC	-0 4.6,5.0 $K^- p$
0.14 ±0.07	BADIER	65C	HBC	- 3 $K^- p$


$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$

Γ_4/Γ_2

NODE=M022R7
NODE=M022R7

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.350±0.031 OUR FIT Error includes scale factor of 1.4.				
0.354±0.033 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.				
0.293±0.032±0.020	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
0.38 ±0.09	BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
0.39 ±0.03	DAUM	81C	CNTR	63 $K^- p \rightarrow K^- 2\pi p$

WEIGHTED AVERAGE
0.354±0.033 (Error scaled by 1.4)



$\Gamma(K\omega)/\Gamma(K^*(892)\pi)$

Γ_5/Γ_2

NODE=M022R8
NODE=M022R8

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.118±0.034 OUR FIT				
0.10 ±0.04	FIELD	67	HBC	- 3.8 $K^- p$

$\Gamma(K\eta)/\Gamma(K^*(892)\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.006^{+0.014}_{-0.004} OUR FIT Error includes scale factor of 1.2.

0.07 ± 0.04 FIELD 67 HBC - 3.8 $K^- p$

 Γ_7/Γ_2

NODE=M022R9
NODE=M022R9

 $\Gamma(K\eta)/\Gamma(K\pi)$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
-------	-----	-------------	------	-----	---------

0.0030^{+0.0070}_{-0.0020} OUR FIT Error includes scale factor of 1.3.

0 ± 0.0056 ¹ ASTON 88B LASS - 11 $K^- p \rightarrow K^- \eta p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04 95 AGUILAR-... 71B HBC 3.9,4.6 $K^- p$

<0.065 ² BASSOMPIE... 69 HBC 5.0 $K^+ p$

<0.02 BISHOP 69 HBC 3.5 $K^+ p$

¹ ASTON 88B quote < 0.0092 at CL=95%. We convert this to a central value and 1 sigma error in order to be able to use it in our constrained fit.

² Restated by us.

 Γ_7/Γ_1

NODE=M022R10
NODE=M022R10

NODE=M022R;LINKAGE=PQ

NODE=M022R;LINKAGE=R

 $\Gamma(K^*(892)\pi\pi)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.134 ± 0.022 OUR FIT

0.12 ± 0.04 ¹ GOLDBERG 76 HBC - 3 $K^- p \rightarrow p \bar{K}^0 \pi \pi \pi$

¹ Assuming $\pi\pi$ system has isospin 1, which is supported by the data.

 Γ_3/Γ

NODE=M022R11
NODE=M022R11

NODE=M022R;LINKAGE=T

 $\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

0.27 ± 0.05 OUR FIT

0.21 ± 0.08 ^{1,2} JONGEJANS 78 HBC - 4 $K^- p \rightarrow p \bar{K}^0 \pi \pi \pi$

¹ Restated by us.

² Assuming $\pi\pi$ system has isospin 1, which is supported by the data.

 Γ_3/Γ_1

NODE=M022R12
NODE=M022R12

NODE=M022R12;LINKAGE=R

NODE=M022R12;LINKAGE=T

 $\Gamma(K\omega\pi)/\Gamma_{total}$

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

<0.72 95 0 JONGEJANS 78 HBC 4 $K^- p \rightarrow p \bar{K}^0 4\pi$

 Γ_8/Γ

NODE=M022R13
NODE=M022R13

 $K_2^*(1430)$ REFERENCES

ABLIKIM	22L	JHEP 2207 045	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61649
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira		REFID=57836
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
ALAVI-HARATI	02B	PRL 89 072001	A. Alavi-Harati <i>et al.</i>	(FNAL KTeV Collab.)	REFID=48822
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40281
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
CARLSMITH	87	PR D36 3502	D. Carlsmith <i>et al.</i>	(EFI, SAFL)	REFID=40557
ASTON	84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)	REFID=22763
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
CIHANGIR	82	PL 117B 123	S. Cihangir <i>et al.</i>	(FNAL, MINN, ROCH)	REFID=21280
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22821
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
JONGEJANS	78	NP B139 383	B. Jongejans <i>et al.</i>	(ZEEM, CERN, NIJM+)	REFID=22445
MARTIN	78	NP B134 392	A.D. Martin <i>et al.</i>	(DURH, GEVA)	REFID=22446
BOWLER	77	NP B126 31	M.G. Bowler <i>et al.</i>	(OXF)	REFID=22437
GOLDBERG	76	LNC 17 253	J. Goldberg	(HAIF)	REFID=22742
HENDRICK	76	NP B112 189	K. Hendrickx <i>et al.</i>	(MONS, SAFL, PARIS+)	REFID=22743
LAUSCHER	75	NP B86 189	P. Lauscher <i>et al.</i>	(ABCLV Collab.) JP	REFID=22582
MCCUBBIN	75	NP B86 13	N.A. McCubbin, L. Lyons	(OXF)	REFID=22434
DEHM	74	NP B75 47	G. Dehm <i>et al.</i>	(MPIM, BRUX, MONS, CERN)	REFID=22736
LINGLIN	73	NP B55 408	D. Linglin	(CERN)	REFID=22428
AGUILAR-...	71B	PR D4 2583	M. Aguilar-Benitez, R.L. Eisner, J.B. Kinson	(BNL)	REFID=22408
BARNHAM	71C	NP B28 171	K.W.J. Barnham <i>et al.</i>	(BIRM, GLAS)	REFID=22409
CORDS	71	PR D4 1974	D. Cords <i>et al.</i>	(PURD, UCD, IUPU)	REFID=22411
BASSOMPIE...	69	NP B13 189	G. Bassompierre <i>et al.</i>	(CERN, BRUX) JP	REFID=22710
BISHOP	69	NP B9 403	J.M. Bishop <i>et al.</i>	(WISC)	REFID=22485
CRENNELL	69D	PRL 22 487	D.J. Crennell <i>et al.</i>	(BNL)	REFID=22399
DAVIS	69	PRL 23 1071	P.J. Davis <i>et al.</i>	(LRL)	REFID=22400
LIND	69	NP B14 1	V.G. Lind <i>et al.</i>	(LRL) JP	REFID=22404
SCHWEING...	68	PR 166 1317	F. Schweingruber <i>et al.</i>	(ANL, NWES)	REFID=22398
Also		Thesis	F.L. Schweingruber	(NWES, NWES)	REFID=22709
BASSANO	67	PRL 19 968	D. Bassano <i>et al.</i>	(BNL, SYRA)	REFID=22695
FIELD	67	PL 24B 638	J.H. Field <i>et al.</i>	(UCSD)	REFID=22701
BADIER	65C	PL 19 612	J. Badier <i>et al.</i>	(EPOL, SAFL, AMST)	REFID=22690

NODE=M022

K(1460)

$$I(J^P) = \frac{1}{2}(0^-)$$

Observed in $K\pi\pi$ partial-wave analysis.

NODE=M021

NODE=M021

NODE=M021M

NODE=M021M

NODE=M021M;LINKAGE=A

NODE=M021W

NODE=M021W

NODE=M021W;LINKAGE=A

NODE=M021215;NODE=M021

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=4

NODE=M021220

NODE=M021W1
NODE=M021W1NODE=M021W2
NODE=M021W2NODE=M021W3
NODE=M021W3NODE=M021R00
NODE=M021R00

NODE=M021R00;LINKAGE=A

NODE=M021

REFID=61150
REFID=59187
REFID=22548
REFID=22767**K(1460) MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1482.40 ± 3.58 ± 15.22	894k	AAIJ	18AI	LHCB	$D^0 \rightarrow K^\mp 2\pi^\pm \pi^\mp$
~ 1460	63	DAUM	81C	CNTR	$K^- p \rightarrow K^- 2\pi p$
~ 1400	13	¹ BRANDENB...	76B	ASPK	$K^\pm p \rightarrow K^+ 2\pi p$

¹ Coupled mainly to $K f_0(1370)$. Decay into $K^*(892)\pi$ seen.**K(1460) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

335.60 ± 6.20 ± 8.65	894k	AAIJ	18AI	LHCB	$D^0 \rightarrow K^\mp 2\pi^\pm \pi^\mp$
~ 260	63	DAUM	81C	CNTR	$K^- p \rightarrow K^- 2\pi p$
~ 250	15	¹ BRANDENB...	76B	ASPK	$K^\pm p \rightarrow K^+ 2\pi p$

¹ Coupled mainly to $K f_0(1370)$. Decay into $K^*(892)\pi$ seen.**K(1460) DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K\rho$	seen
Γ_3 $K_0^*(1430)\pi$	seen
Γ_4 $K\phi$	seen

K(1460) PARTIAL WIDTHS **$\Gamma(K^*(892)\pi)$** **Γ_1**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 109	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-------	------	-----	--

 $\Gamma(K\rho)$ **Γ_2**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 34	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
------	------	-----	--

 $\Gamma(K_0^*(1430)\pi)$ **Γ_3**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

~ 117	DAUM	81C	CNTR 63 $K^- p \rightarrow K^- 2\pi p$
-------	------	-----	--

 $\Gamma(K\phi)/\Gamma_{\text{total}}$ **Γ_4/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	24k	¹ AAIJ	21E	LHCB $B^+ \rightarrow J/\psi\phi K^+$
------	-----	-------------------	-----	---------------------------------------

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 12 σ .**K(1460) REFERENCES**

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AI	EPJ C78 443	R. Aaij <i>et al.</i>	(LHCb Collab.)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)
BRANDENB...	76B	PRL 36 1239	G.W. Brandenburg <i>et al.</i>	(SLAC) JP

$K_2(1580)$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+ \pi^-$ system. Needs confirmation.

NODE=M039

NODE=M039

 $K_2(1580)$ MASS

NODE=M039M

VALUE (MeV)	DOCUMENT ID	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1580	OTTER	79	10,14,16 $K^- p$

NODE=M039M

 $K_2(1580)$ WIDTH

NODE=M039W

VALUE (MeV)	DOCUMENT ID	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 110	OTTER	79	10,14,16 $K^- p$

NODE=M039W

 $K_2(1580)$ DECAY MODES

NODE=M039215;NODE=M039

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K_2^*(1430)\pi$	possibly seen

DESIG=1

DESIG=2

 $K_2(1580)$ BRANCHING RATIOS

NODE=M039220

$\Gamma(K^*(892)\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
seen	OTTER	79	HBC	10,14,16 $K^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
possibly seen	GULER	11	BELL	$B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$	

NODE=M039R1
NODE=M039R1

$\Gamma(K_2^*(1430)\pi)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
possibly seen	OTTER	79	HBC	10,14,16 $K^- p$	

NODE=M039R2
NODE=M039R2 **$K_2(1580)$ REFERENCES**

NODE=M039

GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)
OTTER	79	NP B147 1	G. Otter <i>et al.</i>	(AACH3, BERL, CERN, LOIC+) JP

REFID=53668
REFID=22772

K(1630)

$$I(J^P) = \frac{1}{2}(?^?)$$

OMITTED FROM SUMMARY TABLE

Seen as a narrow peak, compatible with the experimental resolution, in the invariant mass of the $K_S^0 \pi^+ \pi^-$ system produced in $\pi^- p$ interactions at high momentum transfers.

NODE=M160

NODE=M160

K(1630) MASS

NODE=M160M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1629±7	~ 75	KARNAUKHOV98	BC	16.0 $\pi^- p \rightarrow (K_S^0 \pi^+ \pi^-)$ $X^+ \pi^- X^0$

NODE=M160M

K(1630) WIDTH

NODE=M160W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
16⁺¹⁹₋₁₆	~ 75	¹ KARNAUKHOV98	BC	16.0 $\pi^- p \rightarrow (K_S^0 \pi^+ \pi^-)$ $X^+ \pi^- X^0$

NODE=M160W

¹ Compatible with an experimental resolution of 14 ± 1 MeV.

NODE=M160W;LINKAGE=A

K(1630) DECAY MODES

NODE=M160215;NODE=M160

Mode

 $\Gamma_1 \quad K_S^0 \pi^+ \pi^-$

DESIG=1

K(1630) REFERENCES

KARNAUKHOV 98 PAN 61 203 V.M. Karnaukhov, C. Coca, V.I. Moroz
Translated from YAF 61 252.

NODE=M160

REFID=46371

NODE=M099

K₁(1650)

$$I(J^P) = \frac{1}{2}(1^+)$$

This entry contains various peaks in strange meson systems ($K^+ \phi$, $K \pi \pi$) reported in partial-wave analysis in the 1600–1900 mass region.

NODE=M099

K₁(1650) MASS

NODE=M099M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1650±50		FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1861±10 ⁺¹⁶ ₋₄₆	24k	¹ AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1911±37 ⁺¹²⁴ ₋₄₈	24k	¹ AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1793±59 ⁺¹⁵³ ₋₁₀₁	4289	^{2,3} AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
~ 1840		ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 1800		DAUM	81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

NODE=M099M

OCCUR=2

¹ One of two K_1 states reported by AAIJ 21E. From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 4.5σ .

NODE=M099M;LINKAGE=C

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 7.6σ .

NODE=M099M;LINKAGE=A

³ Superseded by AAIJ 21E.

NODE=M099M;LINKAGE=B

K₁(1650) WIDTH

NODE=M099W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
150± 50		FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$

NODE=M099W

• • • We do not use the following data for averages, fits, limits, etc. • • •

149 ± 41 ⁺²³¹ ₋₂₃	24k	1 AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
276 ± 50 ⁺³¹⁹ ₋₁₅₉	24k	1 AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
365 ± 157 ⁺¹³⁸ ₋₂₁₅	4289	2,3 AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
~ 250		DAUM	81C	CNTR	- 63 $K^- p \rightarrow K^- 2\pi p$

OCCUR=2

¹ One of two K_1 states reported by AAIJ 21E. From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 4.5 σ .

NODE=M099W;LINKAGE=C

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 7.6 σ .

NODE=M099W;LINKAGE=A

³ Superseded by AAIJ 21E.

NODE=M099W;LINKAGE=B

$K_1(1650)$ DECAY MODES

NODE=M099215;NODE=M099

Mode	
Γ_1	$K \pi \pi$
Γ_2	$K \phi$

DESIG=1

DESIG=2

$K_1(1650)$ REFERENCES

NODE=M099

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

REFID=61150
REFID=57657
REFID=57636
REFID=20569
REFID=22801
REFID=22548

$K^*(1680)$

$$I(J^P) = \frac{1}{2}(1^-)$$

NODE=M095

$K^*(1680)$ MASS

NODE=M095M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1718 ± 18 OUR AVERAGE					
1722 ± 20 ⁺³³ ₋₁₀₉	4289	¹ AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1677 ± 10 ± 32		ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
1735 ± 10 ± 20		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M095M

• • • We do not use the following data for averages, fits, limits, etc. • • •

1678 ± 64		BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1800 ± 70		ETKIN	80	MPS 0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
~ 1650		ESTABROOKS	78	ASPK 0	13 $K^\pm p \rightarrow K^\pm \pi^\pm n$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5 σ .

NODE=M095M;LINKAGE=A

$K^*(1680)$ WIDTH

NODE=M095W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
320 ± 110 OUR AVERAGE Error includes scale factor of 4.2. [322 ± 110 MeV OUR 2023 AVERAGE Scale factor = 4.2]					
354 ± 75 ⁺¹⁴⁰ ₋₁₈₁	4289	² AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
205 ± 16 ± 34		ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
423 ± 18 ± 30		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

NODE=M095W

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

454 ± 270		BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
170 ± 30		ETKIN	80	MPS 0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
250 to 300		ESTABROOKS	78	ASPK 0	13 $K^\pm p \rightarrow K^\pm \pi^\pm n$

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5 σ .

NODE=M095W;LINKAGE=A

K*(1680) DECAY MODES

NODE=M095215;NODE=M095

Mode	Fraction (Γ_i/Γ)	
Γ_1 $K\pi$	(38.7 \pm 2.5) %	DESIG=1
Γ_2 $K\rho$	(31.4 $^{+5.0}_{-2.1}$) %	DESIG=3
Γ_3 $K^*(892)\pi$	(29.9 $^{+2.2}_{-5.0}$) %	DESIG=2
Γ_4 $K\phi$	seen	DESIG=4
Γ_5 $K\eta$	(1.4 $^{+1.0}_{-0.8}$) %	DESIG=6

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 2.9$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-36	
x_3	-39	-72
	x_1	x_2

K*(1680) BRANCHING RATIOS

NODE=M095220

$\Gamma(K\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.387\pm0.026 OUR FIT						NODE=M095R4
0.388\pm0.014\pm0.022	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	NODE=M095R4

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$						Γ_1/Γ_3
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
1.30$^{+0.23}_{-0.14}$ OUR FIT						NODE=M095R2
2.8 \pm1.1	ASTON	84	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$	NODE=M095R2

$\Gamma(K\rho)/\Gamma(K\pi)$						Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
0.81$^{+0.14}_{-0.09}$ OUR FIT						NODE=M095R3
1.2 \pm0.4	ASTON	84	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$	NODE=M095R3

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$						Γ_2/Γ_3
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
1.05$^{+0.27}_{-0.11}$ OUR FIT						NODE=M095R1
0.97\pm0.09$^{+0.30}_{-0.10}$	ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	NODE=M095R1

$\Gamma(K\phi)/\Gamma_{\text{total}}$						Γ_4/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
seen	24k	³ AAIJ	21E	LHCB $B^+ \rightarrow J/\psi \phi K^+$		NODE=M095R00
• • • We do not use the following data for averages, fits, limits, etc. • • •						NODE=M095R00
seen	4289	^{4,5} AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$		NODE=M095R00

³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 4.7 σ .

⁴ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.5 σ .

⁵ Superseded by AAIJ 21E.

NODE=M095R00;LINKAGE=B
 NODE=M095R00;LINKAGE=A
 NODE=M095R00;LINKAGE=C

$\Gamma(K\eta)/\Gamma(K\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_1
$0.037 \pm 0.007^{+0.024}_{-0.018}$	116k	⁶ CHEN	20A BELL	$D^0 \rightarrow K^- \pi^+ \eta$	

NODE=M095R02
NODE=M095R02

⁶CHEN 20A quotes the ratio $\Gamma(K^*(1680)^- \rightarrow K^- \eta)/\Gamma(K^*(1680)^- \rightarrow K^- \pi^0) = 0.11 \pm 0.02^{+0.06}_{-0.04} \pm 0.04(\text{BPDG})$ where the last uncertainty comes from $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.20)\%$. We divide it by 3 taking into account that $\Gamma(K^*(1680)^- \rightarrow K^- \pi^0)/\Gamma(K^*(1680)^- \rightarrow (K\pi)^-) = 1/3$.

NODE=M095R02;LINKAGE=A

$\Gamma(K\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
$1.44 \pm 0.21^{+0.96}_{-0.73}$	116k	⁷ CHEN	20A BELL	$D^0 \rightarrow K^- \pi^+ \eta$	

NODE=M095R01
NODE=M095R01

••• We do not use the following data for averages, fits, limits, etc. •••

⁷From an amplitude analysis of the decay $D^0 \rightarrow K^- \pi^+ \eta$ with a significance of 16σ . Not independent of the CHEN 20A measurement of $\Gamma(K^*(1680) \rightarrow K\eta)/\Gamma(K^*(1680) \rightarrow K\pi)$.

NODE=M095R01;LINKAGE=A

$K^*(1680)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)	
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)	
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	
ASTON	84	PL 149B 258	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP	

NODE=M095

REFID=61150
REFID=60333
REFID=57657
REFID=57636
REFID=41002
REFID=40262
REFID=40234
REFID=22689
REFID=22545
REFID=22443

$K_2(1770)$

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this Review, PDG 04.

NODE=M023

NODE=M023

$K_2(1770)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1773 ± 8 OUR AVERAGE					
$1777 \pm 35^{+122}_{-77}$	4289	¹ AAIJ	17C LHCb		$B^+ \rightarrow J/\psi \phi K^+$
1773 ± 8		² ASTON	93 LASS		$11K^- p \rightarrow K^- \omega p$
1743 ± 15		TIKHOMIROV	03 SPEC		$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1810 ± 20		FRAME	86 OMEG +		$13 K^+ p \rightarrow \phi K^+ p$
~ 1730		ARMSTRONG	83 OMEG -		$18.5 K^- p \rightarrow 3K p$
~ 1780		³ DAUM	81C CNTR -		$63 K^- p \rightarrow K^- 2\pi p$
1710 ± 15	60	CHUNG	74 HBC -		$7.3 K^- p \rightarrow K^- \omega p$
1767 ± 6		BLIEDEN	72 MMS -		$11-16 K^- p$
1730 ± 20	306	⁴ FIRESTONE	72B DBC +		$12 K^+ d$
1765 ± 40		⁵ COLLEY	71 HBC +		$10 K^+ p \rightarrow K 2\pi N$
1740		DENEGRI	71 DBC -		$12.6 K^- d \rightarrow K 2\pi d$
1745 ± 20		AGUILAR-...	70C HBC -		$4.6 K^- p$
1780 ± 15		BARTSCH	70C HBC -		$10.1 K^- p$
1760 ± 15		LUDLAM	70 HBC -		$12.6 K^- p$

NODE=M023M

NODE=M023M

- ¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.0σ .
- ² From a partial wave analysis of the $K^- \omega$ system.
- ³ From a partial wave analysis of the $K^- 2\pi$ system.
- ⁴ Produced in conjunction with excited deuteron.
- ⁵ Systematic errors added correspond to spread of different fits.

NODE=M023M;LINKAGE=C
NODE=M023M;LINKAGE=A
NODE=M023M;LINKAGE=B
NODE=M023M;LINKAGE=P
NODE=M023M;LINKAGE=X

$K_2(1770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
186 ± 14 OUR AVERAGE					
$217 \pm 116^{+221}_{-154}$	4289	⁶ AAIJ	17C LHCb		$B^+ \rightarrow J/\psi \phi K^+$
186 ± 14		⁷ ASTON	93 LASS		$11K^- p \rightarrow K^- \omega p$

NODE=M023W

NODE=M023W

• • • We do not use the following data for averages, fits, limits, etc. • • •

147 ± 70		TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
140 ± 40		FRAME 86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
~ 220		ARMSTRONG 83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 210		⁸ DAUM 81C	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$
110 ± 50	60	CHUNG 74	HBC -	7.3 $K^- p \rightarrow K^- \omega p$
100 ± 26		BLIEDEN 72	MMS -	11-16 $K^- p$
210 ± 30	306	⁹ FIRESTONE 72B	DBC +	12 $K^+ d$
90 ± 70		¹⁰ COLLEY 71	HBC +	10 $K^+ p \rightarrow K 2\pi N$
130		DENEGRI 71	DBC -	12.6 $K^- d \rightarrow \bar{K} 2\pi d$
100 ± 50		AGUILAR-... 70C	HBC -	4.6 $K^- p$
138 ± 40		BARTSCH 70C	HBC -	10.1 $K^- p$
50 ⁺ ₋ 40 20		LUDLAM 70	HBC -	12.6 $K^- p$

⁶ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.0 σ .

⁷ From a partial wave analysis of the $K^- \omega$ system.

⁸ From a partial wave analysis of the $K^- 2\pi$ system.

⁹ Produced in conjunction with excited deuteron.

¹⁰ Systematic errors added correspond to spread of different fits.

NODE=M023W;LINKAGE=A
 NODE=M023W;LINKAGE=B
 NODE=M023W;LINKAGE=C
 NODE=M023W;LINKAGE=P
 NODE=M023W;LINKAGE=X

$K_2(1770)$ DECAY MODES

NODE=M023215;NODE=M023

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi \pi$	
Γ_2 $K_2^*(1430)\pi$	seen
Γ_3 $K^*(892)\pi$	seen
Γ_4 $K f_2(1270)$	seen
Γ_5 $K f_0(980)$	
Γ_6 $K \phi$	seen
Γ_7 $K \omega$	seen

DESIG=1;OUR EST;→ UNCHECKED ←
 DESIG=2;OUR EST;→ UNCHECKED ←
 DESIG=4;OUR EST;→ UNCHECKED ←
 DESIG=9;OUR EST;→ UNCHECKED ←
 DESIG=11
 DESIG=10
 DESIG=8

$K_2(1770)$ BRANCHING RATIOS

NODE=M023220

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$ Γ_2/Γ_1

NODE=M023R1

$(K_2^*(1430) \rightarrow K\pi)$

NODE=M023R1
 NODE=M023R1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-------	-------------	------	-----	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.03	DAUM 81C	CNTR		63 $K^- p \rightarrow K^- 2\pi p$
~ 1.0	¹¹ FIRESTONE 72B	DBC +		12 $K^+ d$
< 1.0	COLLEY 71	HBC		10 $K^+ p$
0.2 ± 0.2	AGUILAR-... 70C	HBC -		4.6 $K^- p$
< 1.0	BARTSCH 70C	HBC -		10.1 $K^- p$
1.0	BARBARO-... 69	HBC +		12.0 $K^+ p$

¹¹ Produced in conjunction with excited deuteron.

NODE=M023R1;LINKAGE=P

$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$ Γ_3/Γ_1

NODE=M023R3
 NODE=M023R3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.23	DAUM 81C	CNTR		63 $K^- p \rightarrow K^- 2\pi p$
--------	----------	------	--	-----------------------------------

$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$ Γ_4/Γ_1

NODE=M023R4

$(f_2(1270) \rightarrow \pi\pi)$

NODE=M023R4
 NODE=M023R4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 0.74	DAUM 81C	CNTR		63 $K^- p \rightarrow K^- 2\pi p$
--------	----------	------	--	-----------------------------------

$\Gamma(K f_0(980))/\Gamma_{total}$ Γ_5/Γ

NODE=M023R6
 NODE=M023R6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
---------------	---------------	------	--	--

$\Gamma(K\phi)/\Gamma_{\text{total}}$						Γ_6/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	24k	12 AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi\phi K^+$	
seen		ARMSTRONG	83	OMEG	$18.5 K^- p \rightarrow K^- \phi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
seen	4289	13,14 AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi\phi K^+$	

¹² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 7.9 σ .

¹³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.0 σ .

¹⁴ Superseded by AAIJ 21E.

NODE=M023R5
NODE=M023R5

NODE=M023R5;LINKAGE=C
NODE=M023R5;LINKAGE=A
NODE=M023R5;LINKAGE=B

$\Gamma(K\omega)/\Gamma_{\text{total}}$						Γ_7/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		OTTER	81	HBC	\pm	8.25,10,16 $K^\pm p$
seen		CHUNG	74	HBC	$-$	7.3 $K^- p \rightarrow K^- \omega p$

NODE=M023R2
NODE=M023R2

$K_2(1770)$ REFERENCES

NODE=M023

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=43597
FRAME	86	NP B276 667	D. Frame <i>et al.</i>	(GLAS)	REFID=20569
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)	REFID=22801
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)	REFID=22548
OTTER	81	NP B181 1	G. Otter	(AACH3, BERL, LOIC, VIEN, BIRM+)	REFID=22549
CHUNG	74	PL 51B 413	S.U. Chung <i>et al.</i>	(BNL)	REFID=22735
BLIEDEN	72	PL 39B 668	H.R. Blieden <i>et al.</i>	(STON, NEAS)	REFID=22788
FIRESTONE	72B	PR D5 505	A. Firestone <i>et al.</i>	(LBL)	REFID=22506
COLLEY	71	NP B26 71	D.C. Colley <i>et al.</i>	(BIRM, GLAS)	REFID=22785
DENEGRI	71	NP B28 13	D. Denegri <i>et al.</i>	(JHU) JP	REFID=22497
AGUILAR...	70C	PRL 25 54	M. Aguilar-Benitez <i>et al.</i>	(BNL)	REFID=22782
BARTSCH	70C	PL 33B 186	J. Bartsch <i>et al.</i>	(AACH, BERL, CERN+)	REFID=22783
LUDLAM	70	PR D2 1234	T. Ludlam, J. Sandweiss, A.J. Slaughter	(YALE)	REFID=22784
BARBARO...	69	PRL 22 1207	A. Barbaro-Galtieri <i>et al.</i>	(LRL)	REFID=22483

NODE=M060

$K_3^*(1780)$

$$I(J^P) = \frac{1}{2}(3^-)$$

$K_3^*(1780)$ T-MATRIX POLE \sqrt{s}

NODE=M060PP

VALUE (MeV)		DOCUMENT ID	TECN	COMMENT	
(1754 \pm 13) - i (119 \pm 14)		¹ PELAEZ	17	RVUE	$\pi K \rightarrow \pi K$

NODE=M060PP

¹ Reanalysis of ESTABROOKS 78 and ASTON 88 satisfying Forward Dispersion Relations and using sequences of Pade approximants.

NODE=M060PP;LINKAGE=A

$K_3^*(1780)$ MASS

NODE=M060M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
1779 \pm 8 OUR AVERAGE		Error includes scale factor of 1.2.					
1813 \pm 15 $^{+65}_{-16}$	18k	¹ ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$		
1781 \pm 8 \pm 4		² ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	
1740 \pm 14 \pm 15		² ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
1779 \pm 11		³ BALDI	76	SPEC	+	10 $K^+ p \rightarrow K^0 \pi^+ p$	
1776 \pm 26		⁴ BRANDENB...	76D	ASPK	0	13 $K^\pm p \rightarrow K^\pm \pi^\mp N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
1720 \pm 10 \pm 15	6111	⁵ BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
1749 \pm 10		ASTON	88B	LASS	-	11 $K^- p \rightarrow K^- \eta p$	
1780 \pm 9	300	BAUBILLIER	84B	HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
1790 \pm 15		BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow K_S^0 2\pi N$	
1784 \pm 9	2060	CLELAND	82	SPEC	\pm	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$	
1786 \pm 15		⁶ ASTON	81D	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	
1762 \pm 9	190	TOAFF	81	HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
1850 \pm 50		ETKIN	80	MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$	
1812 \pm 28		BEUSCH	78	OMEG		10 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
1786 \pm 8		CHUNG	78	MPS	0	6 $K^- p \rightarrow K^- \pi^+ n$	

NODE=M060M

¹ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (2.0 \pm 0.4^{+1.9}_{-0.4}) \times 10^{-6}$.

² From energy-independent partial-wave analysis.

³ From a fit to Y_6^2 moment. $J^P = 3^-$ found.

⁴ Confirmed by phase shift analysis of ESTABROOKS 78, yields $J^P = 3^-$.

⁵ From a partial wave amplitude analysis.

⁶ From a fit to the Y_6^0 moment.

NODE=M060M;LINKAGE=D

NODE=M060M;LINKAGE=K

NODE=M060M;LINKAGE=M

NODE=M060M;LINKAGE=A

NODE=M060M;LINKAGE=F

NODE=M060M;LINKAGE=J

$K_3^*(1780)$ WIDTH

NODE=M060W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
161±17 OUR AVERAGE		Error includes scale factor of 1.1.			
191 ⁺⁴³⁺³ ₋₃₇₋₈₁	1.8k	¹ ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
203±30±8		² ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
171±42±20		² ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
135±22		³ BALDI	76	SPEC	+ 10 $K^+ p \rightarrow K^0 \pi^+ p$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
187±31±20	6111	⁴ BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
193 ⁺⁵¹ ₋₃₇		ASTON	88B	LASS	- 11 $K^- p \rightarrow K^- \eta p$
99±30	300	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 130		BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 2\pi N$
191±24	2060	CLELAND	82	SPEC	± 50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
225±60		⁵ ASTON	81D	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
~ 80	190	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
240±50		ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
181±44		⁶ BEUSCH	78	OMEG	10 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
96±31		CHUNG	78	MPS	0 6 $K^- p \rightarrow K^- \pi^+ n$
270±70		⁷ BRANDENB...	76D	ASPK	0 13 $K^\pm p \rightarrow K^\pm \pi^\mp N$

NODE=M060W

NODE=M060W;LINKAGE=A

NODE=M060W;LINKAGE=K

NODE=M060W;LINKAGE=M

NODE=M060W;LINKAGE=F

NODE=M060W;LINKAGE=J

NODE=M060W;LINKAGE=D

NODE=M060W;LINKAGE=E

¹ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (2.0 \pm 0.4^{+1.9}_{-0.4}) \times 10^{-6}$.

² From energy-independent partial-wave analysis.

³ From a fit to Y_6^2 moment. $J^P = 3^-$ found.

⁴ From a partial wave amplitude analysis.

⁵ From a fit to Y_6^0 moment.

⁶ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

⁷ ESTABROOKS 78 find that BRANDENBURG 76D data are consistent with 175 MeV width. Not averaged.

$K_3^*(1780)$ DECAY MODES

NODE=M060215;NODE=M060

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\rho$	(31 ± 9) %	DESIG=3
Γ_2 $K^*(892)\pi$	(20 ± 5) %	DESIG=2
Γ_3 $K\pi$	(18.8± 1.0) %	DESIG=1
Γ_4 $K\eta$	(30 ± 13) %	DESIG=6
Γ_5 $K_2^*(1430)\pi$	< 16 %	95% DESIG=4

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 4 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	85		
x_3	18	21	
x_4	-98	-94	-27
	x_1	x_2	x_3

$K_3^*(1780)$ BRANCHING RATIOS

NODE=M060220

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$

 Γ_1/Γ_2 NODE=M060R5
NODE=M060R5

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.52±0.23 OUR FIT				
1.52±0.21±0.10	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$

 Γ_2/Γ_3 NODE=M060R7
NODE=M060R7

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.09±0.26 OUR FIT				
1.09±0.26	ASTON	84B	LASS	0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma_{\text{total}}$

 Γ_3/Γ NODE=M060R4
NODE=M060R4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.188±0.010 OUR FIT				
0.188±0.010 OUR AVERAGE				
0.187±0.008±0.008	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
0.19 ±0.02	ESTABROOKS	78	ASPK	0 13 $K^\pm p \rightarrow K\pi N$

$\Gamma(K\eta)/\Gamma(K\pi)$

 Γ_4/Γ_3 NODE=M060R8
NODE=M060R8

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
1.6 ±0.7 OUR FIT				

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.41±0.050	¹ BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
0.50±0.18	ASTON	88B	LASS	- 11 $K^- p \rightarrow K^- \eta p$

¹ This result supersedes ASTON 88B.

NODE=M060R8;LINKAGE=H

$\Gamma(K_2^*(1430)\pi)/\Gamma(K^*(892)\pi)$

 Γ_5/Γ_2 NODE=M060R6
NODE=M060R6

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.78	95	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$K_3^*(1780)$ REFERENCES

NODE=M060

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
PELAEZ	17	EPJ C77 91	J.R. Pelaez, A.Rodas, J.R. de Elvira		REFID=57836
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	88B	PL B201 169	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP	REFID=40281
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234
ASTON	84B	NP B247 261	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA)	REFID=22763
BAUBILLIER	84B	ZPHY C26 37	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22459
BAUBILLIER	82B	NP B202 21	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22551
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81D	PL 99B 502	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22820
TOAFF	81	PR D23 1500	S. Toaff <i>et al.</i>	(ANL, KANS)	REFID=22454
ETKIN	80	PR D22 42	A. Etkin <i>et al.</i>	(BNL, CUNY) JP	REFID=22545
BEUSCH	78	PL 74B 282	W. Beusch <i>et al.</i>	(CERN, AACH3, ETH) JP	REFID=22537
CHUNG	78	PRL 40 355	S.U. Chung <i>et al.</i>	(BNL, BRAN, CUNY+) JP	REFID=22814
ESTABROOKS	78	NP B133 490	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+) JP	REFID=22443
Also		PR D17 658	P.G. Estabrooks <i>et al.</i>	(MCGI, CARL, DURH+)	REFID=22444
BALDI	76	PL 63B 344	R. Baldi <i>et al.</i>	(GEVA) JP	REFID=22807
BRANDENB...	76D	PL 60B 478	G.W. Brandenburg <i>et al.</i>	(SLAC) JP	REFID=22808

$K_2(1820)$

$$I(J^P) = \frac{1}{2}(2^-)$$

See our mini-review in the 2004 edition of this *Review* (PDG 04) under $K_2(1770)$.

NODE=M146

NODE=M146

 $K_2(1820)$ MASS

NODE=M146M

NODE=M146M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1819±12 OUR AVERAGE				
$1853 \pm 27^{+18}_{-35}$	4289	¹ AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
1816 ± 13		² ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 1840		³ DAUM	81C CNTR	$63 K^- p \rightarrow K^- 2\pi p$
¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 σ .				
² From a partial wave analysis of the $K^- \omega$ system.				
³ From a partial wave analysis of the $K^- 2\pi$ system.				

NODE=M146M;LINKAGE=B

NODE=M146M;LINKAGE=A

NODE=M146M;LINKAGE=C

 $K_2(1820)$ WIDTH

NODE=M146W

NODE=M146W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
264±34 OUR AVERAGE				
$167 \pm 58^{+82}_{-72}$	4289	¹ AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
276 ± 35		² ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 230		³ DAUM	81C CNTR	$63 K^- p \rightarrow K^- 2\pi p$
¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 3.0 σ .				
² From a partial wave analysis of the $K^- \omega$ system.				
³ From a partial wave analysis of the $K^- 2\pi$ system.				

NODE=M146W;LINKAGE=A

NODE=M146W;LINKAGE=B

NODE=M146W;LINKAGE=C

 $K_2(1820)$ DECAY MODES

NODE=M146215;NODE=M146

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi \pi$	seen
Γ_2 $K_2^*(1430) \pi$	seen
Γ_3 $K^*(892) \pi$	seen
Γ_4 $K f_2(1270)$	seen
Γ_5 $K \omega$	seen
Γ_6 $K \phi$	seen

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

DESIG=7

 $K_2(1820)$ BRANCHING RATIOS

NODE=M146220

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 0.77	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K} 2\pi p$
$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 0.05	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K} 2\pi p$
$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$	Γ_4/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 0.18	DAUM	81C CNTR	$63K^- p \rightarrow \bar{K} 2\pi p$

NODE=M146R1

NODE=M146R1

NODE=M146R2

NODE=M146R2

NODE=M146R3

NODE=M146R3

$\Gamma(K\phi)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
seen	4289	^{2,3} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi\phi K^+$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.8 σ .
² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 3.0 σ .
³ Superseded by AAIJ 21E.

NODE=M146R00
 NODE=M146R00

NODE=M146R00;LINKAGE=B
 NODE=M146R00;LINKAGE=A
 NODE=M146R00;LINKAGE=C

$K_2(1820)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ASTON	93	PL B308 186	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
DAUM	81C	NP B187 1	C. Daum <i>et al.</i>	(AMST, CERN, CRAC, MPIM+)

NODE=M146

REFID=61150
 REFID=57657
 REFID=57636
 REFID=49653
 REFID=43597
 REFID=22548

NODE=M088

$K(1830)$

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of $K\phi$ system. Needs confirmation.

NODE=M088

$K(1830)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$1874 \pm 43^{+59}_{-115}$	4289	^{1,2} AAIJ	17C LHCB		$B^+ \rightarrow J/\psi\phi K^+$
~ 1830		ARMSTRONG 83	OMEG	-	$18.5 K^- p \rightarrow 3K p$

- ¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 3.5 σ .
² A subsequent amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ by AAIJ 21E did not confirm this measurement.

NODE=M088M;LINKAGE=A
 NODE=M088M;LINKAGE=B

NODE=M088M

NODE=M088M

$K(1830)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$168 \pm 90^{+280}_{-104}$	4289	^{3,4} AAIJ	17C LHCB		$B^+ \rightarrow J/\psi\phi K^+$
~ 250		ARMSTRONG 83	OMEG	-	$18.5 K^- p \rightarrow 3K p$

- ³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 3.5 σ .
⁴ A subsequent amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ by AAIJ 21E did not confirm this measurement.

NODE=M088W

NODE=M088W

NODE=M088W;LINKAGE=A
 NODE=M088W;LINKAGE=B

$K(1830)$ DECAY MODES

NODE=M088215;NODE=M088

Mode

Γ_1 $K\phi$

DESIG=1

$K(1830)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
ARMSTRONG	83	NP B221 1	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)JP

NODE=M088

REFID=61150
 REFID=57657
 REFID=57636
 REFID=22801

$K_0^*(1950)$

$$I(J^P) = \frac{1}{2}(0^+)$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+$ system. Needs confirmation.

NODE=M134

NODE=M134

 $K_0^*(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1957±14 OUR AVERAGE				
[1944 ± 18 MeV OUR 2023 AVERAGE]				
1980±14±19	¹ AAIJ	23AH	LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
1942±22±21	LEES	21A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' K^+ K^-$
1945±10±20	² ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
1917±12	³ ZHOU	06	RVUE	$K p \rightarrow K^- \pi^+ n$
1820±40	⁴ ANISOVICH	97C	RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$ ² We take the central value of the two solutions and the larger error given.³ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.⁴ T-matrix pole. Reanalysis of ASTON 88 data.

NODE=M134M

NODE=M134M

NEW

NODE=M134M;LINKAGE=B

NODE=M134M;LINKAGE=A

NODE=M134M;LINKAGE=ZU

NODE=M134M;LINKAGE=A1

 $K_0^*(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
170±50 OUR AVERAGE				Error includes scale factor of 2.2. See the ideogram below.
[100 ± 40 MeV OUR 2023 AVERAGE Scale factor = 1.3]				
229±26±16	¹ AAIJ	23AH	LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
80±32±20	LEES	21A	BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow \eta' K^+ K^-$
201±34±79	² ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
145±38	³ ZHOU	06	RVUE	$K p \rightarrow K^- \pi^+ n$
250±100	⁴ ANISOVICH	97C	RVUE	11 $K^- p \rightarrow K^- \pi^+ n$

¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$ ² We take the central value of the two solutions and the larger error given.³ S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.⁴ T-matrix pole. Reanalysis of ASTON 88 data.

NODE=M134W

NODE=M134W


NEW

NODE=M134W;LINKAGE=B

NODE=M134W;LINKAGE=A

NODE=M134W;LINKAGE=ZU

NODE=M134W;LINKAGE=A1



$K_0^*(1950)$ DECAY MODES

NODE=M134215;NODE=M134

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^- \pi^+$	(52±14) %

DESIG=1

 $K_0^*(1950)$ BRANCHING RATIOS

NODE=M134220

$\Gamma(K^- \pi^+)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.52±0.08±0.12	¹ ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
~ 0.60	² ZHOU	06	RVUE		$K p \rightarrow K^- \pi^+ n$
¹ We take the central value of the two solutions and the larger error given.					
² S-matrix pole. Using ASTON 88 and assuming $K_0^*(700)$, $K_0^*(1430)$.					

NODE=M134R1
NODE=M134R1NODE=M134R1;LINKAGE=A
NODE=M134R1;LINKAGE=ZU **$K_0^*(1950)$ REFERENCES**

NODE=M134

AAIJ	23AH PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	21A PR D104 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ZHOU	06 NP A775 212	Z.Y. Zhou, H.Q. Zheng	
ANISOVICH	97C PL B413 137	A.V. Anisovich, A.V. Sarantsev	
ASTON	88 NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

REFID=62349
REFID=61442
REFID=51198
REFID=45815
REFID=40262

NODE=M104

 $K_2^*(1980)$

$$I(J^P) = \frac{1}{2}(2^+)$$

Needs confirmation.

NODE=M104

 $K_2^*(1980)$ MASS

NODE=M104M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1990⁺⁶⁰₋₅₀	OUR AVERAGE	Error includes scale factor of 2.8. See the ideogram below.			
[1994 ⁺⁶⁰ ₋₅₀	MeV	OUR 2023 AVERAGE	Scale factor = 2.8]		
2046 ⁺¹⁷⁺⁶⁷ ₋₁₆₋₁₅	1.8k	¹ ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
1868± 8 ⁺⁴⁰ ₋₅₇	183k	ABLIKIM	19AQBES	±	$J/\psi \rightarrow K^+ K^- \pi^0$
1973± 8± 25		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2073±94 ⁺²⁴⁵ ₋₂₄₀	4289	^{2,3} AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
2020±20		TIKHOMIROV	03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1978±40	241	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
¹ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (7.0 \pm 0.5^{+3.7}_{-0.6}) \times 10^{-6}$.					
² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 σ .					
³ A reanalysis by AAIJ 21E using a larger data sample did not confirm this measurement, the new result having a significance of only 1.6 σ .					

NODE=M104M

NEW

NODE=M104M;LINKAGE=C

NODE=M104M;LINKAGE=B
NODE=M104M;LINKAGE=E



$K_2^*(1980)$ WIDTH

NODE=M104W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
348⁺⁵⁰₋₃₀	OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
408 ⁺³⁸ ₋₃₄	72	1 ABLIKIM	20F	BES3	$\psi(2S) \rightarrow K^+ K^- \eta$
272 \pm 24 ⁺⁵⁰ ₋₁₅	183k	ABLIKIM	19AQ	BES	$J/\psi \rightarrow K^+ K^- \pi^0$
373 \pm 33 \pm 60		ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
678 \pm 311 ⁺¹¹⁵³ ₋₅₅₉	4289	2,3 AAIJ	17C	LHCB	$B^+ \rightarrow J/\psi \phi K^+$
180 \pm 70		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
398 \pm 47	241	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NODE=M104W

NODE=M104W;LINKAGE=C


¹ Seen in $\psi(2S)$ decay with branching ratio $\psi(2S) \rightarrow K^\pm X \rightarrow K^+ K^- \eta = (7.0 \pm 0.5^{+3.7}_{-0.6}) \times 10^{-6}$.

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4 σ .

³ A reanalysis by AAIJ 21E using a larger data sample did not confirm this measurement, the new result having a significance of only 1.6 σ .

NODE=M104W;LINKAGE=B

NODE=M104W;LINKAGE=E



$K_2^*(1980)$ DECAY MODES

NODE=M104215;NODE=M104

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	possibly seen
Γ_2 $K\rho$	possibly seen
Γ_3 $K f_2(1270)$	possibly seen
Γ_4 $K\phi$	seen
Γ_5 $K\eta$	seen

DESIG=2
DESIG=3
DESIG=4
DESIG=5
DESIG=6

$K_2^*(1980)$ BRANCHING RATIOS

NODE=M104220

$\Gamma(K^*(892)\pi)/\Gamma_{total}$ Γ_1/Γ

NODE=M104R01
NODE=M104R01

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

$\Gamma(K\rho)/\Gamma_{total}$ Γ_2/Γ

NODE=M104R02
NODE=M104R02

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	GULER	11	BELL $B^+ \rightarrow J/\psi K^+ \pi^+ \pi^-$

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$ Γ_2/Γ_1

NODE=M104R1
NODE=M104R1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
$1.49 \pm 0.24 \pm 0.09$	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$\Gamma(K f_2(1270))/\Gamma_{total}$ Γ_3/Γ

NODE=M104R3
NODE=M104R3

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	TIKHOMIROV 03	SPEC	40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$

$\Gamma(K\phi)/\Gamma_{total}$ Γ_4/Γ

NODE=M104R00
NODE=M104R00

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	4289	^{1,2} AAIJ	17C	LHCB $B^+ \rightarrow J/\psi \phi K^+$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.4σ .

NODE=M104R00;LINKAGE=A
NODE=M104R00;LINKAGE=C

² A reanalysis by AAIJ 21E using a larger data sample did not confirm this measurement, the new result having a significance of only 1.6σ .

$\Gamma(K\eta)/\Gamma_{total}$ Γ_5/Γ

NODE=M104R03
NODE=M104R03

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1.8k	¹ ABLIKIM	20F	BES3 $\psi(2S) \rightarrow K^+ K^- \eta$
seen	116k	² CHEN	20A	BELL $D^0 \rightarrow K^- \pi^+ \eta$

¹ Seen decaying to $K\eta$ in an amplitude analysis of $\psi(2S) \rightarrow K^+ K^- \eta$.

NODE=M104R03;LINKAGE=A
NODE=M104R03;LINKAGE=B

² From an amplitude analysis of the decay $D^0 \rightarrow K^- \pi^+ \eta$ with a significance of 17σ .

$K_2^*(1980)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
CHEN	20A	PR D102 012002	Y.Q. Chen <i>et al.</i>	(BELLE Collab.)	REFID=60333
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636
GULER	11	PR D83 032005	H. Guler <i>et al.</i>	(BELLE Collab.)	REFID=53668
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>		REFID=49423
		Translated from YAF 66 860.			
BIRD	89	SLAC-332	P.F. Bird	(SLAC)	REFID=41002
ASTON	87	NP B292 693	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40234

NODE=M104

REFID=61150
REFID=60256
REFID=60333
REFID=59909
REFID=57657
REFID=57636
REFID=53668
REFID=49423REFID=41002
REFID=40234

NODE=M035

 $K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

 $K_4^*(2045)$ MASS

NODE=M035M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2048⁺⁸₋₉ OUR AVERAGE		Error includes scale factor of 1.1.			
2090 [±] 9 ⁺¹¹ ₋₂₉	183k	ABLIKIM	19AQ	BES	± $J/\psi \rightarrow K^+ K^- \pi^0$
2062 [±] 14 [±] 13		¹ ASTON	86	LASS	0 $11 K^- p \rightarrow K^- \pi^+ n$
2039 [±] 10	400	^{2,3} CLELAND	82	SPEC	± $50 K^+ p \rightarrow K_S^0 \pi^\pm p$
2070 ⁺¹⁰⁰ ₋₄₀		⁴ ASTON	81C	LASS	0 $11 K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2079 [±] 7	431	TORRES	86	MPSF	400 $pA \rightarrow 4KX$
2088 [±] 20	650	BAUBILLIER	82	HBC	- $8.25 K^- p \rightarrow K_S^0 \pi^- p$
2115 [±] 46	488	CARMONY	77	HBC	0 $9 K^+ d \rightarrow K^+ \pi^+ s X$

NODE=M035M

¹ From a fit to all moments.² From a fit to 8 moments.³ Number of events evaluated by us.⁴ From energy-independent partial-wave analysis.NODE=M035M;LINKAGE=E
NODE=M035M;LINKAGE=B
NODE=M035M;LINKAGE=W
NODE=M035M;LINKAGE=D **$K_4^*(2045)$ WIDTH**

NODE=M035W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
199⁺²⁷₋₁₉ OUR AVERAGE					
201 [±] 19 ⁺⁵⁷ ₋₁₇	183k	ABLIKIM	19AQ	BES	± $J/\psi \rightarrow K^+ K^- \pi^0$
221 [±] 48 [±] 27		⁵ ASTON	86	LASS	0 $11 K^- p \rightarrow K^- \pi^+ n$
189 [±] 35	400	^{6,7} CLELAND	82	SPEC	± $50 K^+ p \rightarrow K_S^0 \pi^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
61 [±] 58	431	TORRES	86	MPSF	400 $pA \rightarrow 4KX$
170 ⁺¹⁰⁰ ₋₅₀	650	BAUBILLIER	82	HBC	- $8.25 K^- p \rightarrow K_S^0 \pi^- p$
240 ⁺⁵⁰⁰ ₋₁₀₀		⁸ ASTON	81C	LASS	0 $11 K^- p \rightarrow K^- \pi^+ n$
300 [±] 200		CARMONY	77	HBC	0 $9 K^+ d \rightarrow K^+ \pi^+ s X$

NODE=M035W

⁵ From a fit to all moments.⁶ From a fit to 8 moments.⁷ Number of events evaluated by us.⁸ From energy-independent partial-wave analysis.NODE=M035W;LINKAGE=E
NODE=M035W;LINKAGE=B
NODE=M035W;LINKAGE=W
NODE=M035W;LINKAGE=D **$K_4^*(2045)$ DECAY MODES**

NODE=M035215;NODE=M035

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(9.9 \pm 1.2) %
Γ_2 $K^*(892)\pi\pi$	(9 \pm 5) %
Γ_3 $K^*(892)\pi\pi\pi$	(7 \pm 5) %
Γ_4 $\rho K\pi$	(5.7 \pm 3.2) %
Γ_5 $\omega K\pi$	(5.0 \pm 3.0) %
Γ_6 $\phi K\pi$	(2.8 \pm 1.4) %
Γ_7 $\phi K^*(892)$	(1.4 \pm 0.7) %

DESIG=1

DESIG=2

DESIG=5

DESIG=3

DESIG=4

DESIG=6

DESIG=7

$K_4^*(2045)$ BRANCHING RATIOS

NODE=M035220

$\Gamma(K\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.099±0.012	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$	

NODE=M035R1
NODE=M035R1

$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$					Γ_2/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.89±0.53	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow p K_S^0 3\pi$	

NODE=M035R2
NODE=M035R2

$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$					Γ_3/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.75±0.49	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow p K_S^0 3\pi$	

NODE=M035R5
NODE=M035R5

$\Gamma(\rho K\pi)/\Gamma(K\pi)$					Γ_4/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.58±0.32	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow p K_S^0 3\pi$	

NODE=M035R3
NODE=M035R3

$\Gamma(\omega K\pi)/\Gamma(K\pi)$					Γ_5/Γ_1
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.50±0.30	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow p K_S^0 3\pi$	

NODE=M035R4
NODE=M035R4

$\Gamma(\phi K\pi)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.028±0.014	⁹ TORRES	86	MPSF	400 $pA \rightarrow 4KX$	

NODE=M035R6
NODE=M035R6

$\Gamma(\phi K^*(892))/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	DOCUMENT ID	TECN	COMMENT		
0.014±0.007	⁹ TORRES	86	MPSF	400 $pA \rightarrow 4KX$	

NODE=M035R7
NODE=M035R7⁹ Error determination is model dependent.

NODE=M035R;LINKAGE=A

 $K_4^*(2045)$ REFERENCES

NODE=M035

ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=40262
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)	REFID=22462
TORRES	86	PR D34 707	S. Torres <i>et al.</i>	(VPI, ARIZ, FNAL, FSU+)	REFID=22845
BAUBILLIER	82	PL 118B 447	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)	REFID=22842
CLELAND	82	NP B208 189	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=22455
ASTON	81C	PL 106B 235	D. Aston <i>et al.</i>	(SLAC, CARL, OTTA) JP	REFID=22821
CARMONY	77	PR D16 1251	D.D. Carmony <i>et al.</i>	(PURD, UCD, IUPU)	REFID=22811

$K_2(2250)$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems reported in the 2150–2260 MeV region, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the $J^P = 2^-$ wave.

NODE=M040

NODE=M040

 $K_2(2250)$ MASS

NODE=M040M

NODE=M040M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
2247±17 OUR AVERAGE						
2200±40		¹ ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$	
2235±50		¹ BAUBILLIER 81	HBC	-	8 $K^- p \rightarrow \Lambda \bar{p} X$	
2260±20		¹ CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
2280±20		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
2147±4	37	CHLIAPNIK...	79	HBC	+	32 $K^+ p \rightarrow \bar{\Lambda} p X$
2240±20	20	LISSAUER 70	HBC		9 $K^+ p$	
¹ $J^P = 2^-$ from moments analysis.						

NODE=M040M;LINKAGE=Q

 $K_2(2250)$ WIDTH

NODE=M040W

NODE=M040W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
180±30 OUR AVERAGE						
Error includes scale factor of 1.4.						
150±30		² ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$	
210±30		² CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
180±60		TIKHOMIROV 03	SPEC		40.0 $\pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$	
~ 200		² BAUBILLIER 81	HBC	-	8 $K^- p \rightarrow \Lambda \bar{p} X$	
~ 40	37	CHLIAPNIK...	79	HBC	+	32 $K^+ p \rightarrow \bar{\Lambda} p X$
80±20	20	LISSAUER 70	HBC		9 $K^+ p$	
² $J^P = 2^-$ from moments analysis.						

NODE=M040W;LINKAGE=Q

 $K_2(2250)$ DECAY MODES

NODE=M040215;NODE=M040

Mode	
Γ_1	$K \pi \pi$
Γ_2	$K f_2(1270)$
Γ_3	$K^*(892) f_0(980)$
Γ_4	$\rho \bar{\Lambda}$

DESIG=1

DESIG=3

DESIG=4

DESIG=2

 $K_2(2250)$ REFERENCES

NODE=M040

REFID=49423

TIKHOMIROV 03	PAN 66 828 Translated from YAF 66 860.	G.D. Tikhomirov <i>et al.</i>	
ARMSTRONG 83C	NP B227 365	T.A. Armstrong <i>et al.</i>	(BARI, BIRM, CERN+)
BAUBILLIER 81	NP B183 1	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+) JP
CLELAND 81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+) JP
CHLIAPNIK...	NP B158 253	P.V. Chliapnikov <i>et al.</i>	(CERN, BELG, MONS)
LISSAUER 70	NP B18 491	D. Lissauer <i>et al.</i>	(LBL)

REFID=22852

REFID=22850

REFID=22851

REFID=22849

REFID=22847

$K_3(2320)$

$$I(J^P) = \frac{1}{2}(3^+)$$

OMITTED FROM SUMMARY TABLE

Seen in the $J^P = 3^+$ wave of the antihyperon-nucleon system.
Needs confirmation.

NODE=M090

NODE=M090

 $K_3(2320)$ MASS

NODE=M090M

NODE=M090M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2324 ± 24 OUR AVERAGE				
2330 ± 40	¹ ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
2320 ± 30	¹ CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
¹ $J^P = 3^+$ from moments analysis.				

NODE=M090M;LINKAGE=P

 $K_3(2320)$ WIDTH

NODE=M090W

NODE=M090W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
150 ± 30	² ARMSTRONG 83C	OMEG	-	18 $K^- p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 250	² CLELAND 81	SPEC	±	50 $K^+ p \rightarrow \Lambda \bar{p} X$
² $J^P = 3^+$ from moments analysis.				

NODE=M090W;LINKAGE=P

 $K_3(2320)$ DECAY MODES

NODE=M090215;NODE=M090

Mode

 $\Gamma_1 \quad p \bar{\Lambda}$

DESIG=1

 $K_3(2320)$ REFERENCES

NODE=M090

ARMSTRONG 83C NP B227 365 T.A. Armstrong *et al.* (BARI, BIRM, CERN+)
CLELAND 81 NP B184 1 W.E. Cleland *et al.* (PITT, GEVA, LAUS+)

REFID=22852
REFID=22851

NODE=M098

 $K_5^*(2380)$

$$I(J^P) = \frac{1}{2}(5^-)$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

NODE=M098

 $K_5^*(2380)$ MASS

NODE=M098M

NODE=M098M

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2382 ± 14 ± 19	¹ ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
¹ From a fit to all the moments.				

NODE=M098M;LINKAGE=E

 $K_5^*(2380)$ WIDTH

NODE=M098W

NODE=M098W

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
180 ± 50 OUR AVERAGE	[178 ± 50 MeV OUR 2023 AVERAGE]			
178 ± 37 ± 32	² ASTON 86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
² From a fit to all the moments.				

NEW

NODE=M098W;LINKAGE=E

 $K_5^*(2380)$ DECAY MODES

NODE=M098215;NODE=M098

Mode

Fraction (Γ_i/Γ) $\Gamma_1 \quad K \pi$

(6.1 ± 1.2) %

DESIG=1

$K_5^*(2380)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.061 ± 0.012	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

NODE=M098220

NODE=M098R1
NODE=M098R1 **$K_5^*(2380)$ REFERENCES**

ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)

NODE=M098

REFID=40262
REFID=22462

NODE=M091

 $K_4(2500)$

$$I(J^P) = \frac{1}{2}(4^-)$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

NODE=M091

 $K_4(2500)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2490 ± 20	¹ CLELAND	81	SPEC \pm	50 $K^+ p \rightarrow \Lambda \bar{p}$
¹ $J^P = 4^-$ from moments analysis.				

NODE=M091M

NODE=M091M

NODE=M091M;LINKAGE=R

 $K_4(2500)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
~ 250	² CLELAND	81	SPEC \pm	50 $K^+ p \rightarrow \Lambda \bar{p}$
² $J^P = 4^-$ from moments analysis.				

NODE=M091W

NODE=M091W

NODE=M091W;LINKAGE=R

 $K_4(2500)$ DECAY MODES

NODE=M091215;NODE=M091

Mode

 $\Gamma_1 \quad p \bar{\Lambda}$

DESIG=1

 $K_4(2500)$ REFERENCES

CLELAND	81	NP B184 1	W.E. Cleland <i>et al.</i>	(PITT, GEVA, LAUS+)
---------	----	-----------	----------------------------	---------------------

NODE=M091

REFID=22851

K(3100)

$$I^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE
also known as $K_J^?(3100)$

Narrow peak observed in several ($\Lambda\bar{p} +$ pions) and ($\bar{\Lambda}p +$ pions) states in Σ^- Be reactions by BOURQUIN 86 and in np and nA reactions by ALEEVEV 93. Not seen by BOEHNLEIN 91. If due to strong decays, this state has exotic quantum numbers ($B=0, Q=+1, S=-1$ for $\Lambda\bar{p}\pi^+\pi^+$ and $I \geq 3/2$ for $\Lambda\bar{p}\pi^-$). Needs confirmation.

NODE=M129

NODE=M129

K(3100) MASS

NODE=M129205

VALUE (MeV)
≈ 3100 OUR ESTIMATE

DOCUMENT ID

NODE=M129M
→ UNCHECKED ←

3-BODY DECAYS

NODE=M129M1
NODE=M129M1

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3054 ± 11 OUR AVERAGE			
3060 ± 7 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+$
3056 ± 7 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-$
3055 ± 8 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^-$
3045 ± 8 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^+$

OCCUR=2
OCCUR=3
OCCUR=4

4-BODY DECAYS

NODE=M129M2
NODE=M129M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3059 ± 11 OUR AVERAGE			
3067 ± 6 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
3060 ± 8 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$
3055 ± 7 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$
3052 ± 8 ± 20	¹ ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3105 ± 30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
3115 ± 30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

OCCUR=2
OCCUR=3
OCCUR=4

OCCUR=2

5-BODY DECAYS

NODE=M129M3
NODE=M129M3

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3095 ± 30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$
¹ Supersedes ALEEVEV 90.			

NODE=M129M;LINKAGE=A

K(3100) WIDTH

NODE=M129210

3-BODY DECAYS

NODE=M129W1
NODE=M129W1

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
42 ± 16	² ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+$
36 ± 15	² ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-$
50 ± 18	² ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^-$
30 ± 15	² ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^+$

OCCUR=2
OCCUR=3
OCCUR=4

4-BODY DECAYS

NODE=M129W2
NODE=M129W2

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22 ± 8		² ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
28 ± 12		² ALEEVEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$
32 ± 15		² ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$
30 ± 15		² ALEEVEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$
<30	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
<80	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

OCCUR=2
OCCUR=3
OCCUR=4

OCCUR=2

5-BODY DECAYS

NODE=M129W3
NODE=M129W3

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<30	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$
² Supersedes ALEEVEV 90.				

NODE=M129W;LINKAGE=A

K(3100) DECAY MODES

NODE=M129215;NODE=M129

Mode		
Γ_1	$K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+$	DESIG=1
Γ_2	$K(3100)^{--} \rightarrow \Lambda \bar{p} \pi^-$	DESIG=2
Γ_3	$K(3100)^- \rightarrow \Lambda \bar{p} \pi^+ \pi^-$	DESIG=3
Γ_4	$K(3100)^+ \rightarrow \Lambda \bar{p} \pi^+ \pi^+$	DESIG=4
Γ_5	$K(3100)^0 \rightarrow \Lambda \bar{p} \pi^+ \pi^+ \pi^-$	DESIG=5
Γ_6	$K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$	DESIG=6

$\Gamma(\Sigma(1385)^+ \bar{p})/\Gamma(\Lambda \bar{p} \pi^+)$					Γ_6/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.04	90	ALEEV	93	BIS2	$K(3100)^0 \rightarrow \Sigma(1385)^+ \bar{p}$

NODE=M129R1
NODE=M129R1**K(3100) REFERENCES**

ALEEV	93	PAN 56 1358 Translated from YAF 56 100.	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
BOEHNLEIN	91	NPBPS B21 174	A. Boehnlein <i>et al.</i>	(FLOR, BNL, IND+)
ALEEV	90	ZPHY C47 533	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)
BOURQUIN	86	PL B172 113	M.H. Bourquin <i>et al.</i>	(GEVA, RAL, HEIDP+)

NODE=M129

REFID=43668
REFID=41743
REFID=42173
REFID=22928

CHARMED MESONS

(C = ±1)

$$D^+ = c\bar{d}, D^0 = c\bar{u}, \bar{D}^0 = \bar{c}u, D^- = \bar{c}d, \text{ similarly for } D^{*'}\text{'s}$$

NODE=MXXX035

NODE=MXXX035

 $D^*(2007)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

 $J^P = 1^-$ established by ABLIKIM 23AZ.

NODE=M061

NODE=M061

 $D^*(2007)^0$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M061M

NODE=M061M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

2006.85 ± 0.05 OUR FIT Error includes scale factor of 1.1.

• • • We do not use the following data for averages, fits, limits, etc. • • •

2006 ± 1.5	¹ GOLDHABER 77	MRK1	$e^+ e^-$
------------	---------------------------	------	-----------

¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 .

NODE=M061M

NODE=M061M;LINKAGE=G

 $m_{D^*(2007)^0} - m_{D^0}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M061DM

NODE=M061DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

142.014 ± 0.030 OUR FIT Error includes scale factor of 1.5.

142.016 ± 0.030 OUR AVERAGE Error includes scale factor of 1.5.

142.007 ± 0.015 ± 0.014	10k	¹ TOMARADZE 15	CLEO	$e^+ e^- \rightarrow$ hadrons
-------------------------	-----	---------------------------	------	-------------------------------

142.2 ± 0.3 ± 0.2	145	ALBRECHT 95F	ARG	$e^+ e^- \rightarrow$ hadrons
-------------------	-----	--------------	-----	-------------------------------

142.12 ± 0.05 ± 0.05	1176	BORTOLETTO92B	CLE2	$e^+ e^- \rightarrow$ hadrons
----------------------	------	---------------	------	-------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

142.2 ± 2.0		SADROZINSKI 80	CBAL	$D^{*0} \rightarrow D^0 \pi^0$
-------------	--	----------------	------	--------------------------------

142.7 ± 1.7		² GOLDHABER 77	MRK1	$e^+ e^-$
-------------	--	---------------------------	------	-----------

NODE=M061DM

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration . This value comes from the average of the results for two decay modes, $D^0 \rightarrow K^- \pi^+$ and $D^0 \rightarrow K^- \pi^+ \pi^- \pi^+$.

² From simultaneous fit to $D^*(2010)^+$, $D^*(2007)^0$, D^+ , and D^0 .

NODE=M061DM;LINKAGE=A

NODE=M061DM;LINKAGE=G

$D^*(2007)^0$ WIDTH

NODE=M061W

NODE=M061W

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	¹ ABACHI	88B HRS	$D^{*0} \rightarrow D^+ \pi^-$

¹ Assuming $m_{D^{*0}} = 2007.2 \pm 2.1 \text{ MeV}/c^2$.

NODE=M061W;LINKAGE=A

$D^*(2007)^0$ DECAY MODES

NODE=M061220;NODE=M061

NODE=M061

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $D^0 \pi^0$	(64.7 \pm 0.9) %	
Γ_2 $D^0 \gamma$	(35.3 \pm 0.9) %	
Γ_3 $D^0 e^+ e^-$	(3.91 \pm 0.33) $\times 10^{-3}$	
Γ_4 $\mu^+ \mu^-$	< 2.5 $\times 10^{-8}$	90%
Γ_5 $e^+ e^-$	< 1.7 $\times 10^{-6}$	90%

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 5 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 2.5$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_2 \begin{vmatrix} -100 \\ x_1 \end{vmatrix}$$

$D^*(2007)^0$ BRANCHING RATIOS

NODE=M061225

$\Gamma(D^0 \pi^0) / \Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
---	-------	------	-------------	------	---------	-------------------

NODE=M061R2
NODE=M061R2
NEW

0.647 \pm 0.009 OUR FIT

[0.647 \pm 0.009 OUR 2023 FIT]

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.655 \pm 0.008 \pm 0.005	3.2k	¹ ABLIKIM	15B BES3	$e^+ e^- \rightarrow$ hadrons
0.635 \pm 0.003 \pm 0.017	69k	¹ AUBERT, BE	05G BABR	10.6 $e^+ e^- \rightarrow$ hadrons
0.596 \pm 0.035 \pm 0.028	858	² ALBRECHT	95F ARG	$e^+ e^- \rightarrow$ hadrons
0.636 \pm 0.023 \pm 0.033	1097	² BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons

¹ Derived from the ratio $\Gamma(D^0 \pi^0) / \Gamma(D^0 \gamma)$ assuming that the branching fractions of $D^{*0} \rightarrow D^0 \pi^0$ and $D^{*0} \rightarrow D^0 \gamma$ decays sum to 100%.

NODE=M061R2;LINKAGE=AU

² The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

NODE=M061R2;LINKAGE=A

$\Gamma(D^0 \gamma) / \Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
--	-------	------	-------------	------	---------	-------------------

NODE=M061R1
NODE=M061R1
NEW

0.353 \pm 0.009 OUR FIT

[0.353 \pm 0.009 OUR 2023 FIT]

0.381 \pm 0.029 OUR AVERAGE

0.404 \pm 0.035 \pm 0.028	456	¹ ALBRECHT	95F ARG	$e^+ e^- \rightarrow$ hadrons
0.364 \pm 0.023 \pm 0.033	621	¹ BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons
0.37 \pm 0.08 \pm 0.08		ADLER	88D MRK3	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.345 ± 0.008 ± 0.005	1.8k	² ABLIKIM	15B	BES3	$e^+e^- \rightarrow$	hadrons
0.365 ± 0.003 ± 0.017	68k	² AUBERT, BE	05G	BABR	10.6 $e^+e^- \rightarrow$	hadrons
0.47 ± 0.23		LOW	87	HRS	29 GeV e^+e^-	
0.53 ± 0.13		BARTEL	85G	JADE	e^+e^- ,	hadrons
0.47 ± 0.12		COLES	82	MRK2	e^+e^-	
0.45 ± 0.15		GOLDHABER	77	MRK1	e^+e^-	

¹ The BUTLER 92 and ALBRECHT 95F branching ratios are not independent, they have been constrained by the authors to sum to 100%.

² Derived from the ratio $\Gamma(D^0\pi^0) / \Gamma(D^0\gamma)$ assuming that the branching fractions of $D^{*0} \rightarrow D^0\pi^0$ and $D^{*0} \rightarrow D^0\gamma$ decays sum to 100%.

NODE=M061R;LINKAGE=A

NODE=M061R;LINKAGE=AU

$\Gamma(D^0\pi^0)/\Gamma(D^0\gamma)$

Γ_1/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

1.83 ± 0.07 OUR FIT

[1.83 ± 0.07 OUR 2023 FIT Scale factor = 1.1]

1.85 ± 0.07 OUR AVERAGE

1.90 ± 0.07 ± 0.05	4.9k	ABLIKIM	15B	BES3	10.6 $e^+e^- \rightarrow$	hadrons
1.74 ± 0.02 ± 0.13		AUBERT, BE	05G	BABR	10.6 $e^+e^- \rightarrow$	hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.789 ± 0.082		¹ AAIJ	22N	LHCB	$B^0, B_S^0 \rightarrow \bar{D}^{*0}(K\pi, \pi\pi)$	
---------------	--	-------------------	-----	------	---	--

¹ Statistical error only.

NODE=M061R3
NODE=M061R3
NEW

NODE=M061R3;LINKAGE=A

$\Gamma(D^0e^+e^-)/\Gamma(D^0\gamma)$

Γ_3/Γ_2

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

11.08 ± 0.76 ± 0.49 421 ABLIKIM 21BD BES3 4.178 GeV e^+e^-

NODE=M061R00
NODE=M061R00

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 2.5 × 10⁻⁸ 90 ¹ AAIJ 23D LHCB $B^- \rightarrow \pi^- \mu^+ \mu^-$

¹ AAIJ 23D reports $< 2.6 \times 10^{-8}$ from a measurement of $[\Gamma(D^*(2007)^0 \rightarrow \mu^+\mu^-) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \bar{D}^*(2007)^0\pi^+)]$ assuming $B(B^+ \rightarrow \bar{D}^*(2007)^0\pi^+) = (4.90 \pm 0.17) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow \bar{D}^*(2007)^0\pi^+) = 5.17 \times 10^{-3}$. The reported value also assumes $B(B^- \rightarrow J/\psi(1S)K^-) = (10.20 \pm 0.19) \times 10^{-4}$ and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (59.61 \pm 0.33) \times 10^{-3}$ for the normalization mode $B^- \rightarrow J/\psi(1S)K^-$.

NODE=M061R01
NODE=M061R01

NODE=M061R01;LINKAGE=A

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$

Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.7 × 10⁻⁶ 90 SHEMYAKIN 20 CMD3 $e^+e^- \rightarrow D^0\pi^0, D^0\gamma$

NODE=M061R02
NODE=M061R02

$D^*(2007)^0$ REFERENCES

NODE=M061

AAIJ	23D	EPJ C83 666	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62095
ABLIKIM	23AZ	PL B846 138245	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=62420
AAIJ	22N	PR D105 072005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61737
ABLIKIM	21BD	PR D104 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61535
SHEMYAKIN	20	PAN 83 954	D.N. Shemyakin	(CMD-3 Collab.)	REFID=62315
ABLIKIM	15B	PR D91 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56375
TOMARADZE	15	PR D91 011102	A. Tomaradze <i>et al.</i>	(NWES)	REFID=57142
AUBERT, BE	05G	PR D72 091101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50942
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44374
BORTOLETTO	92B	PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)	REFID=43116
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43170
ABACHI	88B	PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)	REFID=40584
ADLER	88D	PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40579
LOW	87	PL B183 232	E.H. Low <i>et al.</i>	(HRS Collab.)	REFID=40017
BARTEL	85G	PL 161B 197	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=22880
COLES	82	PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)	REFID=22866
SADROZINSKI	80	Madison Conf. 681	H.F.W. Sadrozinski <i>et al.</i>	(PRIN, CIT+)	REFID=22877
GOLDHABER	77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)	REFID=11434

$D^*(2010)^\pm$

$$I(J^P) = \frac{1}{2}(1^-)$$

 $J^P = 1^-$ established by ABLIKIM 23AZ.

NODE=M062

NODE=M062

NODE=M062M

NODE=M062M

NODE=M062M

 $D^*(2010)^\pm$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
-------------	-------------	------	-----	---------

2010.26 ± 0.05 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

2008 ± 3	¹ GOLDHABER 77	MRK1	±	$e^+ e^-$
2008.6 ± 1.0	² PERUZZI 77	LGW	±	$e^+ e^-$

¹ From simultaneous fit to $D^*(2010)^+$, $D^*(2007)^0$, D^+ , and D^0 ; not independent of FELDMAN 77B mass difference below.

² PERUZZI 77 mass not independent of FELDMAN 77B mass difference below and PERUZZI 77 D^0 mass value.

NODE=M062M;LINKAGE=G

NODE=M062M;LINKAGE=P

 $m_{D^*(2010)^+} - m_{D^+}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M062MD

NODE=M062MD

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

140.603 ± 0.015 OUR FIT**140.602 ± 0.014 OUR AVERAGE**

140.6010 ± 0.0068 ± 0.0129	151k	LEES	17F BABR	$e^+ e^- \rightarrow$ hadrons
140.64 ± 0.08 ± 0.06	620	BORTOLETTO92B	CLE2	$e^+ e^- \rightarrow$ hadrons

NODE=M062MD

 $m_{D^*(2010)^+} - m_{D^0}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M062DM

NODE=M062DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

145.4258 ± 0.0017 OUR FIT

[145.4258 ± 0.0017 MeV OUR 2023 FIT]

145.4258 ± 0.0020 OUR AVERAGE Error includes scale factor of 1.2.

145.4259 ± 0.0004 ± 0.0017	312.8k	LEES	13X BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K\pi, K3\pi)\pi^\pm$	OCCUR=3
145.412 ± 0.002 ± 0.012		ANASTASSOV 02	CLE2	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K\pi)\pi^\pm$	
145.54 ± 0.08	611	¹ ADINOLFI 99	BEAT	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.45 ± 0.02		¹ BREITWEG 99	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K\pi)\pi^\pm$	
145.42 ± 0.05		¹ BREITWEG 99	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- 3\pi)\pi^\pm$	OCCUR=2
145.5 ± 0.15	103	² ADLOFF 97B	H1	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.44 ± 0.08	152	² BREITWEG 97	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm, D^0 \rightarrow K^- 3\pi$	
145.42 ± 0.11	199	² BREITWEG 97	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm, D^0 \rightarrow K^- \pi^+$	OCCUR=2
145.4 ± 0.2	48	² DERRICK 95	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.39 ± 0.06 ± 0.03		BARLAG 92B	ACCM	$\pi^- 230$ GeV	
145.5 ± 0.2	115	² ALEXANDER 91B	OPAL	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.30 ± 0.06		² DECAMP 91J	ALEP	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.40 ± 0.05 ± 0.10		ABACHI 88B	HRS	$D^{*\pm} \rightarrow D^0 \pi^\pm$	
145.46 ± 0.07 ± 0.03		ALBRECHT 85F	ARG	$D^{*\pm} \rightarrow D^0 \pi^+$	
145.5 ± 0.3	28	BAILEY 83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$	OCCUR=2
145.5 ± 0.3	60	FITCH 81	SPEC	$\pi^- A$	
145.3 ± 0.5	30	FELDMAN 77B	MRK1	$D^{*+} \rightarrow D^0 \pi^+$	

NODE=M062DM

NEW

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

145.4256 ± 0.0006 ± 0.0017	138.5k	LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- \pi^+) \pi^\pm$
145.4266 ± 0.0005 ± 0.0019	174.3k	LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- 2\pi^+ \pi^-) \pi^\pm$
145.44 ± 0.09	122	² BREITWEG	97B	ZEUS	$D^{*\pm} \rightarrow D^0 \pi^\pm, D^0 \rightarrow K^- \pi^+$
145.8 ± 1.5	16	AHLEN	83	HRS	$D^{*+} \rightarrow D^0 \pi^+$
145.1 ± 1.8	12	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.1 ± 0.5	14	BAILEY	83	SPEC	$D^{*\pm} \rightarrow D^0 \pi^\pm$
145.5 ± 0.5	14	YELTON	82	MRK2	$29 e^+ e^- \rightarrow K^- \pi^+$
~ 145.5		AVERY	80	SPEC	γA
145.2 ± 0.6	2	BLIETSCHAU	79	BEBC	νp

OCCUR=2

OCCUR=3

¹ Statistical errors only.

² Systematic error not evaluated.

NODE=M062DM;LINKAGE=AV
NODE=M062DM;LINKAGE=A

$m_{D^*(2010)^+} - m_{D^*(2007)^0}$

NODE=M062EM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M062EM

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 1.8	¹ PERUZZI	77	LGW	$e^+ e^-$
-----------	----------------------	----	-----	-----------

¹ Not independent of FELDMAN 77B mass difference above, PERUZZI 77 D^0 mass, and GOLDHABER 77 $D^*(2007)^0$ mass.

NODE=M062EM;LINKAGE=P

$D^*(2010)^\pm$ WIDTH

NODE=M062W

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	-----	------	-------------	------	---------

NODE=M062W

83.4 ± 1.8 OUR AVERAGE

83.3 ± 1.2 ± 1.4	1.4	312.8k	¹ LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K \pi, K 3\pi) \pi^\pm$
96 ± 4 ± 22			¹ ANASTASSOV	02	CLE2	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K \pi) \pi^\pm$

OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

83.4 ± 1.7 ± 1.5	1.5	138.5k	¹ LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- \pi^+) \pi^\pm$
83.2 ± 1.5 ± 2.6	2.6	174.3k	¹ LEES	13X	BABR	$D^{*\pm} \rightarrow D^0 \pi^\pm \rightarrow (K^- 2\pi^+ \pi^-) \pi^\pm$
<131	90	110	BARLAG	92B	ACCM	π^- 230 GeV

OCCUR=2

¹ Ignoring the electromagnetic contribution from $D^{*\pm} \rightarrow D^\pm \gamma$.

NODE=M062W;LINKAGE=LE

$D^*(2010)^\pm$ DECAY MODES

NODE=M062225;NODE=M062

$D^*(2010)^-$ modes are charge conjugates of the modes below.

NODE=M062

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 \pi^+$	(67.7 ± 0.5) %
Γ_2 $D^+ \pi^0$	(30.7 ± 0.5) %
Γ_3 $D^+ \gamma$	(1.6 ± 0.4) %

DESIG=1

DESIG=3

DESIG=2

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 6 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.3$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-62	
x_3	-43	-44
	x_1	x_2

$D^*(2010)^+$ BRANCHING RATIOS

$\Gamma(D^0\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.677 ±0.005 OUR FIT					
0.677 ±0.006 OUR AVERAGE					
0.6759 ±0.0029 ±0.0064		^{1,2,3} BARTELT	98	CLE2	e^+e^-
0.688 ±0.024 ±0.013		ALBRECHT	95F	ARG	$e^+e^- \rightarrow$ hadrons
0.681 ±0.010 ±0.013		¹ BUTLER	92	CLE2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.57 ±0.04 ±0.04		ADLER	88D	MRK3	e^+e^-
0.44 ±0.10		COLES	82	MRK2	e^+e^-
0.6 ±0.15		³ GOLDBABER	77	MRK1	e^+e^-

¹ The branching ratios are not independent, they have been constrained by the authors to sum to 100%.

² Systematic error includes theoretical error on the prediction of the ratio of hadronic modes.

³ Assuming that isospin is conserved in the decay.

$\Gamma(D^+\pi^0)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.307 ±0.005 OUR FIT						
0.3073 ±0.0013 ±0.0062						
		^{1,2,3}	BARTELT	98	CLE2	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.312 ±0.011 ±0.008	1404		ALBRECHT	95F	ARG	$e^+e^- \rightarrow$ hadrons
0.308 ±0.004 ±0.008	410		¹ BUTLER	92	CLE2	$e^+e^- \rightarrow$ hadrons
0.26 ±0.02 ±0.02			ADLER	88D	MRK3	e^+e^-
0.34 ±0.07			COLES	82	MRK2	e^+e^-

¹ The branching ratios are not independent, they have been constrained by the authors to sum to 100%.

² Systematic error includes theoretical error on the prediction of the ratio of hadronic modes.

³ Assuming that isospin is conserved in the decay.

$\Gamma(D^+\gamma)/\Gamma_{\text{total}}$	VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.016 ±0.004 OUR FIT							
0.016 ±0.005 OUR AVERAGE							
0.0168 ±0.0042 ±0.0029				^{1,2} BARTELT	98	CLE2	e^+e^-
0.011 ±0.014 ±0.016			12	¹ BUTLER	92	CLE2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<0.052		90		ALBRECHT	95F	ARG	$e^+e^- \rightarrow$ hadrons
0.17 ±0.05 ±0.05				ADLER	88D	MRK3	e^+e^-
0.22 ±0.12				³ COLES	82	MRK2	e^+e^-

¹ The branching ratios are not independent, they have been constrained by the authors to sum to 100%.

² Systematic error includes theoretical error on the prediction of the ratio of hadronic modes.

³ Not independent of $\Gamma(D^0\pi^+)/\Gamma_{\text{total}}$ and $\Gamma(D^+\pi^0)/\Gamma_{\text{total}}$ measurement.

 $D^*(2010)^\pm$ REFERENCES

ABLIKIM	23AZ	PL B846 138245	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=62420
LEES	17F	PRL 119 202003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58277
LEES	13X	PRL 111 111801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55564
Also		PR D88 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55547
Also		PR D88 079902 (err.)	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55695
ANASTASSOV	02	PR D65 032003	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=48550
ADINOLFI	99	NP B547 3	M. Adinolfi <i>et al.</i>	(Beatrice Collab.)	REFID=46925
BREITWEG	99	EPJ C6 67	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=46604
BARTELT	98	PRL 80 3919	J. Bartelt <i>et al.</i>	(CLEO Collab.)	REFID=46349
ADLOFF	97B	ZPHY C72 593	C. Adloff <i>et al.</i>	(H1 Collab.)	REFID=45421
BREITWEG	97	PL B401 192	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45520
BREITWEG	97B	PL B407 402	J. Breitweg <i>et al.</i>	(ZEUS Collab.)	REFID=45699
ALBRECHT	95F	ZPHY C66 63	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44374
DERRICK	95	PL B349 225	M. Derrick <i>et al.</i>	(ZEUS Collab.)	REFID=44373
BARLAG	92B	PL B278 480	S. Barlag <i>et al.</i>	(ACCMOR Collab.)	REFID=42174
BORTOLETTO	92B	PRL 69 2046	D. Bortoletto <i>et al.</i>	(CLEO Collab.)	REFID=43116
BUTLER	92	PRL 69 2041	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43170

NODE=M062230

NODE=M062R1
NODE=M062R1

NODE=M062R1;LINKAGE=A

NODE=M062R1;LINKAGE=B

NODE=M062R1;LINKAGE=G

NODE=M062R3
NODE=M062R3

NODE=M062R3;LINKAGE=A

NODE=M062R3;LINKAGE=B

NODE=M062R3;LINKAGE=G

NODE=M062R2
NODE=M062R2

NODE=M062R2;LINKAGE=A

NODE=M062R2;LINKAGE=B

NODE=M062R2;LINKAGE=C

NODE=M062

ALEXANDER 91B PL B262 341	G. Alexander <i>et al.</i>	(OPAL Collab.)
DECAMP 91J PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)
ABACHI 88B PL B212 533	S. Abachi <i>et al.</i>	(ANL, IND, MICH, PURD+)
ADLER 88D PL B208 152	J. Adler <i>et al.</i>	(Mark III Collab.)
ALBRECHT 85F PL 150B 235	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
AHLEN 83 PRL 51 1147	S.P. Ahlen <i>et al.</i>	(ANL, IND, LBL+)
BAILEY 83 PL 132B 230	R. Bailey <i>et al.</i>	(AMST, BRIS, CERN, CRAC+)
COLES 82 PR D26 2190	M.W. Coles <i>et al.</i>	(LBL, SLAC)
YELTON 82 PRL 49 430	J.M. Yelton <i>et al.</i>	(SLAC, LBL, UCB+)
FITCH 81 PRL 46 761	V.L. Fitch <i>et al.</i>	(PRIN, SACL, TORI+)
AVERY 80 PRL 44 1309	P. Avery <i>et al.</i>	(ILL, FNAL, COLU)
BLIETSCHAU 79 PL 86B 108	J. Blietschau <i>et al.</i>	(AACH3, BONN, CERN+)
FELDMAN 77B PRL 38 1313	G.J. Feldman <i>et al.</i>	(Mark I Collab.)
GOLDHABER 77 PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)
PERUZZI 77 PRL 39 1301	I. Peruzzi <i>et al.</i>	(LGW Collab.)

REFID=41553
 REFID=41614
 REFID=40584
 REFID=40579
 REFID=11527
 REFID=22868
 REFID=22870
 REFID=22866
 REFID=22867
 REFID=22863
 REFID=11498
 REFID=22861
 REFID=22858
 REFID=11434
 REFID=11435

NODE=M252

$D_0^*(2300)$

$$I(J^P) = \frac{1}{2}(0^+)$$

was $D_0^*(2400)$

There is a strong evidence that recent data on $B \rightarrow D\pi\pi$ (AAIJ 15Y, AAIJ 16AH) and $B \rightarrow D\pi K$ (AAIJ 14BH, AAIJ 15V, AAIJ 15X) call for two poles in the scalar $I = 1/2 \pi D$ amplitude in this mass range. The data are consistent with a lower pole at $(2105^{+6}_{-8}) - i(102^{+10}_{-11})$ MeV and a higher pole at $(2451^{+35}_{-26}) - i(134^{+7}_{-8})$ MeV (DU 18A, DU 19, DU 21). For details see review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M252

$D_0^*(2300)$ MASS

NODE=M252M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2343±10 OUR AVERAGE	Error	includes scale factor of 1.5. See the ideogram below.			
2360±15±30		1 AAIJ	15X LHC B	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2349± 6± 4		2 AAIJ	15Y LHC B	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2297± 8±20	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
2308±17±32		ABE	04D BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2354± 7±11		3 AAIJ	15Y LHC B	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2403±14±35	18.8k	4 LINK	04A FOCS	+	γA
2407±21±35	9.8k	4 LINK	04A FOCS	0	γA

NODE=M252M

OCCUR=2

OCCUR=2


- ¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi S$ -wave and the $D\pi S$ - and P -waves.
- ² Modeling the $\pi^+\pi^- S$ -wave with the Isobar formalism.
- ³ Modeling the $\pi^+\pi^- S$ -wave with the K-matrix formalism.
- ⁴ Possibly the feed-down from another state.

NODE=M252M;LINKAGE=A

NODE=M252M;LINKAGE=B

NODE=M252M;LINKAGE=C

NODE=M252M;LINKAGE=D



$D_0^*(2300)$ WIDTH

NODE=M252W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
229±16 OUR AVERAGE						
255±26±51		1 AAIJ	15X	LHCB	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
217±13±13		2 AAIJ	15Y	LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
273±12±48	3.4k	AUBERT	09AB	BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
276±21±63		ABE	04D	BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
230±15±21		3 AAIJ	15Y	LHCB	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
283±24±34	18.8k	4 LINK	04A	FOCS	+	γA
240±55±59	9.8k	4 LINK	04A	FOCS	0	γA

NODE=M252W

OCCUR=2

OCCUR=2

¹ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S -wave and the $D\pi$ S - and P -waves.

² Modeling the $\pi^+ \pi^-$ S -wave with the Isobar formalism.

³ Modeling the $\pi^+ \pi^-$ S -wave with the K-matrix formalism.

⁴ Possibly the feed-down from another state.

NODE=M252W;LINKAGE=A

NODE=M252W;LINKAGE=B

NODE=M252W;LINKAGE=C

NODE=M252W;LINKAGE=D

 $D_0^*(2300)$ DECAY MODES

NODE=M252215;NODE=M252

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\pi^\pm$	seen

DESIG=1

$\Gamma(D\pi^\pm)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen		AAIJ	15X	LHCB	+	$D^*(2300)^+ \rightarrow D^0 \pi^+$
seen		AAIJ	15Y	LHCB	+	$D^*(2300)^+ \rightarrow D^0 \pi^+$
seen	3.4k	AUBERT	09AB	BABR	0	$D^*(2300)^0 \rightarrow D^+ \pi^-$
seen		ABE	04D	BELL	0	$D^*(2300)^0 \rightarrow D^+ \pi^-$
seen	18.8k	LINK	04A	FOCS	+	$D^*(2300)^+ \rightarrow D^0 \pi^+$

NODE=M252R01

NODE=M252R01

 $D_0^*(2300)$ REFERENCES

NODE=M252

DU	21	PRL 126 192001	M.-L. Du <i>et al.</i>			REFID=61096
DU	19	PR D99 114002	M.-L. Du, F.-K. Guo, U.-G. Meissner			REFID=59893
DU	18A	PR D98 094018	M.-L. Du <i>et al.</i>			REFID=61424
AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>		(LHCb Collab.)	REFID=57518
AAIJ	15V	PR D91 092002	R. Aaij <i>et al.</i>		(LHCb Collab.)	REFID=56575
Also		PR D93 119901 (errata.)	R. Aaij <i>et al.</i>		(LHCb Collab.)	REFID=57289
AAIJ	15X	PR D92 012012	R. Aaij <i>et al.</i>		(LHCb Collab.)	REFID=56588
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>		(LHCb Collab.)	REFID=56609
AAIJ	14BH	PR D90 072003	R. Aaij <i>et al.</i>		(LHCb Collab.)	REFID=56208
AUBERT	09AB	PR D79 112004	B. Aubert <i>et al.</i>		(BABAR Collab.)	REFID=52941
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>		(BELLE Collab.)	REFID=50011
LINK	04A	PL B586 11	J.M. Link <i>et al.</i>		(FOCUS Collab.)	REFID=49775

$D_1(2420)$

$$I(J^P) = \frac{1}{2}(1^+)$$

NODE=M253

 $D_1(2420)$ MASS

NODE=M253M

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , D_s^{*0} , $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M253M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2422.1±0.6 OUR FIT	Error includes scale factor of 1.7. [2422.1 ± 0.6 MeV OUR 2023 FIT Scale factor = 1.7]				
2422.1±0.8 OUR AVERAGE	Error includes scale factor of 2.1. See the ideogram below.				
2424.8±0.1±0.7	79k	¹ AAIJ	20D	LHCB	0 $B^- \rightarrow D^{*+} \pi^- \pi^-$
2427.2±1.0±1.2	4207	ABLIKIM	20P	BES3	+ $e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$
2419.6±0.1±0.7	210k	AAIJ	13CC	LHCB	0 $p p \rightarrow D^{*+} \pi^- X$
2423.1±1.5 ^{+0.4} _{-1.0}	2.7k	² ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2421.9±4.7 ^{+3.4} _{-1.2}	759	³ ABRAMOWICZ13	ZEUS	+	$e^\pm p \rightarrow D^{*0} \pi^+ X$
2420.1±0.1±0.8	103k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2426 ±3 ±1	151	ABE	05A	BELL	0 $B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
2421 ±2 ±1	124	ABE	05A	BELL	+ $\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
2421.4±1.5±0.9		⁴ ABE	04D	BELL	0 $B^- \rightarrow D^{*+} \pi^- \pi^-$
2421 ⁺¹ ₋₂ ±2	286	AVERY	94C	CLE2	0 $e^+ e^- \rightarrow D^{*+} \pi^- X$
2425 ±2 ±2	146	BERGFELD	94B	CLE2	+ $e^+ e^- \rightarrow D^{*0} \pi^+ X$
2422 ±2 ±2	51	FRABETTI	94B	E687	0 $\gamma Be \rightarrow D^{*+} \pi^- X$
2428 ±3 ±2	279	AVERY	90	CLEO	0 $e^+ e^- \rightarrow D^{*+} \pi^- X$
2414 ±2 ±5	171	ALBRECHT	89H	ARG	0 $e^+ e^- \rightarrow D^{*+} \pi^- X$
2428 ±8 ±5	171	ANJOS	89C	TPS	0 $\gamma N \rightarrow D^{*+} \pi^- X$
2443 ±7 ±5	190	ANJOS	89C	TPS	+ $\gamma N \rightarrow D^0 \pi^+ X^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2420.5±2.1±0.9	3.1k	⁵ CHEKANOV	09	ZEUS	0 $e^\pm p \rightarrow D^{*+} \pi^- X$
2421.7±0.7±0.6	7.5k	ABULENCIA	06A	CDF	0 $1900 p \bar{p} \rightarrow D^{*+} \pi^- X$
2425 ±3	235	⁶ ABREU	98M	DLPH	0 $e^+ e^-$

NODE=M253M

NEW

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M253M;LINKAGE=B

NODE=M253M;LINKAGE=AR


NODE=M253M;LINKAGE=BA

NODE=M253M;LINKAGE=AB

NODE=M253M;LINKAGE=CH

NODE=M253M;LINKAGE=K

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.² From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 .³ From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^+ are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1 , respectively.⁴ Fit includes the contribution from $D_1^*(2430)^0$.⁵ Calculated using the mass difference $m(D_1^0) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of 0.9 MeV.⁶ No systematic error given.



$m_{D_1(2420)^0} - m_{D^{*+}}$

NODE=M253DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$ and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M253DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
411.8±0.6 OUR FIT		Error includes scale factor of 1.7. [411.8 ± 0.6 MeV OUR 2023 FIT Scale factor = 1.7]		
411.5±0.8 OUR AVERAGE				
410.2±2.1±0.9	3.1k	CHEKANOV 09	ZEUS	$e^\pm p \rightarrow D^{*+} \pi^- X$
411.7±0.7±0.4	7.5k	ABULENCIA 06A	CDF	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$

NODE=M253DM
NEW

$m_{D_1(2420)^\pm} - m_{D_1(2420)^0}$

NODE=M253DMC

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4₋₃⁺²±3	BERGFELD 94B	CLE2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M253DMC

$D_1(2420)$ WIDTH

NODE=M253W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
31.3± 1.9 OUR AVERAGE		Error includes scale factor of 2.8. See the ideogram below.			
33.6± 0.3± 2.7	79k	¹ AAIJ 20D	LHCB	0	$B^- \rightarrow D^{*+} \pi^- \pi^-$
23.2± 2.3± 2.3	4207	ABLIKIM 20P	BES3	+	$e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$
35.2± 0.4± 0.9	210k	AAIJ 13CC	LHCB	0	$p p \rightarrow D^{*+} \pi^- X$
38.8± 5.0 ⁺ _{-5.4}	2.7k	² ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
31.4± 0.5± 1.3	103k	DEL-AMO-SA...10P	BABR	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
20.0± 1.7± 1.3	7.5k	ABULENCIA 06A	CDF	0	1900 $p\bar{p} \rightarrow D^{*+} \pi^- X$
24 ± 7 ± 8	151	ABE 05A	BELL	0	$B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
21 ± 5 ± 8	124	ABE 05A	BELL	+	$\bar{B}^0 \rightarrow D^+ \pi^+ \pi^- \pi^-$
23.7± 2.7± 4.0		³ ABE 04D	BELL	0	$B^- \rightarrow D^{*+} \pi^- \pi^-$
20 ⁺ ₋₅ ± 6 ± 3	286	AVERY 94C	CLE2	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$

NODE=M253W


OCCUR=2

OCCUR=2

26	$\begin{matrix} +8 \\ -7 \end{matrix}$	± 4	146	BERGFELD	94B	CLE2	+	$e^+e^- \rightarrow D^{*0}\pi^+X$	OCCUR=2
15	± 8	± 4	51	FRABETTI	94B	E687	0	$\gamma Be \rightarrow D^{*+}\pi^-X$	
23	$\begin{matrix} +8 \\ -6 \end{matrix}$	$\begin{matrix} +10 \\ -3 \end{matrix}$	279	AVERY	90	CLEO	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	
13	± 6	$\begin{matrix} +10 \\ -5 \end{matrix}$	171	ALBRECHT	89H	ARG	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	
41	± 19	± 8	190	ANJOS	89C	TPS	+	$\gamma N \rightarrow D^0\pi^+X^0$	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●									
53.2	± 7.2	$\begin{matrix} +3.3 \\ -4.9 \end{matrix}$	3.1k	CHEKANOV	09	ZEUS	0	$e^\pm p \rightarrow D^{*+}\pi^-X$	
58	± 14	± 10	171	ANJOS	89C	TPS	0	$\gamma N \rightarrow D^{*+}\pi^-X$	

- ¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+}\pi^-\pi^-$ decay.
- ² From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 .
- ³ Fit includes the contribution from $D_1^*(2430)^0$.

NODE=M253W;LINKAGE=B
 NODE=M253W;LINKAGE=AR
 NODE=M253W;LINKAGE=AB



$D_1(2420)$ DECAY MODES

NODE=M253215;NODE=M253
 NODE=M253

$\bar{D}_1(2420)$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2007)^0\pi$	seen
Γ_2 $D\pi^+\pi^-$	
Γ_3 $D\rho^0$	
Γ_4 $Df_0(500)$	
Γ_5 $D_0^*(2300)^0\pi$	
Γ_6 $D^0\pi$	
Γ_7 $D^*\pi^+\pi^-$	

DESIG=1
 DESIG=3
 DESIG=4
 DESIG=5
 DESIG=6
 DESIG=2
 DESIG=7

$D_1(2420)$ BRANCHING RATIOS

NODE=M253220

$\Gamma(D^*(2007)^0\pi)/\Gamma_{total}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ACKERSTAFF	97W	OPAL	0 $e^+e^- \rightarrow D^{*+}\pi^-X$
seen	AVERY	90	CLEO	0 $e^+e^- \rightarrow D^{*+}\pi^-X$
seen	ALBRECHT	89H	ARG	0 $e^+e^- \rightarrow D^*\pi^-X$
seen	ANJOS	89C	TPS	0 $\gamma N \rightarrow D^{*+}\pi^-X$
seen	ANJOS	89C	TPS	+ $\gamma N \rightarrow D^0\pi^+X^0$

NODE=M253R01
 NODE=M253R01

OCCUR=2

$\Gamma(D^0\pi)/\Gamma(D^*(2007)^0\pi)$ Γ_6/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.18	90	BERGFELD	94B	CLE2	+	$e^+e^- \rightarrow \text{hadrons}$
<0.24	90	AVERY	90	CLEO	0	$e^+e^- \rightarrow D^+\pi^-X$

NODE=M253R02
 NODE=M253R02
 OCCUR=2

 $D_1(2420)$ POLARIZATION AMPLITUDE A_{D_1}

NODE=M253PAH

A polarization amplitude A_{D_1} is a parameter that depends on the initial polarization of the D_1 and is sensitive to a possible S -wave contribution to its decay. For D_1 decays the helicity angle, θ_h , distribution varies like $1 + A_{D_1} \cos^2 \theta_h$, where θ_h is the angle in the D^* rest frame between the two pions emitted by the $D_1 \rightarrow D^*\pi$ and the $D^* \rightarrow D\pi$.

NODE=M253PAH

Unpolarized D_1 decaying purely via D -wave is predicted to give $A_{D_1} = 3$.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
5.73±0.25 OUR AVERAGE						
7.8 ^{+6.7} _{-2.7} ^{+4.6} _{-1.8}	2.7k	¹ ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$	
5.72±0.25	103k	DEL-AMO-SA...10P	BABR	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	
5.9 ^{+3.0} _{-1.7} ^{+2.4} _{-1.0}		CHEKANOV	09	ZEUS	0	$e^\pm p \rightarrow D^{*+}\pi^-X$
3.30±0.48	210k	² AAIJ	13CC	LHCB	0	$pp \rightarrow D^{*+}\pi^-X$
3.8 ±0.6 ±0.8		³ AUBERT	09Y	BABR	0	$B^+ \rightarrow D_1^0 \ell^+ \nu_\ell$
3.8 ±0.6 ±0.8		³ AUBERT	09Y	BABR	+	$B^0 \rightarrow D_1^- \ell^+ \nu_\ell$
2.74 ^{+1.40} _{-0.93}		⁴ AVERY	94C	CLE2	0	$e^+e^- \rightarrow D^{*+}\pi^-X$

NODE=M253PAH

••• We do not use the following data for averages, fits, limits, etc. •••

OCCUR=2

¹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 . A pure D -wave not excluded although some \bar{S} -wave mixing possible.

NODE=M253PAH;LINKAGE=AR

² Systematic uncertainty not estimated. Resonance parameters fixed.

NODE=M253PAH;LINKAGE=A

³ Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths and helicity angle distributions for charged and neutral D_1 mesons.

NODE=M253PAH;LINKAGE=AU

⁴ Systematic uncertainties not estimated.

NODE=M253PAH;LINKAGE=AV

 $D_1(2420)$ REFERENCES

NODE=M253

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60253
ABLIKIM	20P	PL B804 135395	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60395
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
ABRAMOWICZ	13	NP B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54743
DEL-AMO-SA...	10P	PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52929
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)	REFID=51054
ABE	05A	PRL 94 221805	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50755
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50011
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46315
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)	REFID=44096
BERGFELD	94B	PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)	REFID=44099
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=41001
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)	REFID=40737

$D_1(2430)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$

 $J^P = 1^+$ determined by AAIJ 20D.

NODE=M180

NODE=M180

NODE=M180M

NODE=M180M

 $D_1(2430)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2412 ± 9 OUR AVERAGE				
2411 ± 3 ± 9	79k	¹ AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2427 ± 26 ± 25		ABE	04D BELLE	$B^- \rightarrow D^{*+} \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2477 ± 28		² AUBERT	06L BABR	$\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.² Systematic errors not estimated.NODE=M180M;LINKAGE=A
NODE=M180M;LINKAGE=AU **$D_1(2430)^0$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
314 ± 29 OUR AVERAGE				
309 ± 9 ± 28	79k	¹ AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
384 ⁺¹⁰⁷ ₋₇₅ ± 74		ABE	04D BELLE	$B^- \rightarrow D^{*+} \pi^- \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
266 ± 97		² AUBERT	06L BABR	$\bar{B}^0 \rightarrow D^{*+} \omega \pi^-$

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.² Systematic errors not estimated.

NODE=M180W

NODE=M180W

NODE=M180W;LINKAGE=A
NODE=M180W;LINKAGE=AU **$D_1(2430)^0$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2010)^+ \pi^-$	seen

NODE=M180215;NODE=M180

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D_1(2430)^0$ REFERENCES

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AUBERT	06L	PR D74 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M180

REFID=60253
REFID=51140
REFID=50011

$D_2^*(2460)$

$$I(J^P) = \frac{1}{2}(2^+)$$

NODE=M254

 $D_2^*(2460)$ MASS

NODE=M254M

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M254M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2461.1±0.8 OUR FIT		Error includes scale factor of 6.3. [2461.1 ^{+0.7} _{-0.8} MeV OUR 2023 FIT Scale factor = 6.2]			
2461.1±0.7 OUR AVERAGE		Error includes scale factor of 5.2. See the ideogram below.			
2463.7±0.4±0.7	28k	1 AAIJ	16AH LHCb	0	$B^- \rightarrow D^+ \pi^- \pi^-$
2464.0±1.4±0.5	2k	2 AAIJ	15V LHCb	0	$B^- \rightarrow D^+ K^- \pi^-$
2465.6±1.8±1.3		3 AAIJ	15X LHCb	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
2468.6±0.6±0.3		4 AAIJ	15Y LHCb	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2460.4±0.4±1.2	82k	AAIJ	13CC LHCb	0	$pp \rightarrow D^{*+} \pi^- X$
2460.4±0.1±0.1	675k	AAIJ	13CC LHCb	0	$pp \rightarrow D^+ \pi^- X$
2463.1±0.2±0.6	342k	AAIJ	13CC LHCb	+	$pp \rightarrow D^0 \pi^+ X$
2462.5±2.4 ^{+1.3} _{-1.1}	2.3k	5 ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2460.6±4.4 ^{+3.6} _{-0.8}	1371	6 ABRAMOWICZ13	ZEUS	+	$e^\pm p \rightarrow D^{(*)0} \pi^+ X$
2462.2±0.1±0.8	243k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2465.4±0.2±1.1	111k	7 DEL-AMO-SA..10P	BABR	+	$e^+ e^- \rightarrow D^0 \pi^+ X$
2460.4±1.2±2.2	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
2465.7±1.8 ^{+1.4} _{-4.8}	2909	KUZMIN	07 BELL	+	$e^+ e^- \rightarrow \text{hadrons}$
2461.6±2.1±3.3		8 ABE	04D BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
2464.5±1.1±1.9	5.8k	8 LINK	04A FOCS	0	γA
2465 ±3 ±3	486	AVERY	94C CLE2	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2463 ±3 ±3	310	BERGFELD	94B CLE2	+	$e^+ e^- \rightarrow D^0 \pi^+ X$
2453 ±3 ±2	128	FRABETTI	94B E687	0	$\gamma Be \rightarrow D^+ \pi^- X$
2453 ±3 ±2	185	FRABETTI	94B E687	+	$\gamma Be \rightarrow D^0 \pi^+ X$
2461 ±3 ±1	440	AVERY	90 CLEO	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2455 ±3 ±5	337	ALBRECHT	89B ARG	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2469 ±4 ±6		ALBRECHT	89F ARG	+	$e^+ e^- \rightarrow D^0 \pi^+ X$
2459 ±3 ±2	153	ANJOS	89C TPS	0	$\gamma N \rightarrow D^+ \pi^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
2468.1±0.6±0.5		9 AAIJ	15Y LHCb	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2469.1±3.7 ^{+1.2} _{-1.3}	1.5k	10 CHEKANOV	09 ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
2463.3±0.6±0.8	20k	ABULENCIA	06A CDF	0	1900 $p\bar{p} \rightarrow D^+ \pi^- X$
2467.6±1.5±0.8	3.5k	11 LINK	04A FOCS	+	γA
2461 ±6	126	12 ABREU	98M DLPH	0	$e^+ e^-$
2466 ±7	1	ASRATYAN	95 BEBC	0	53,40 $\nu(\bar{\nu}) \rightarrow pX, dX$

NODE=M254M

NEW

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

NODE=M254M;LINKAGE=B

NODE=M254M;LINKAGE=A

NODE=M254M;LINKAGE=CA

NODE=M254M;LINKAGE=BC

NODE=M254M;LINKAGE=AR

NODE=M254M;LINKAGE=AB

NODE=M254M;LINKAGE=DE

NODE=M254M;LINKAGE=LI

NODE=M254M;LINKAGE=CC

NODE=M254M;LINKAGE=CH

¹ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

² From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, nonresonant spin-0 and spin-1 components as well as the $D_0^*(2400)^0$, $D_2^*(2460)^0$ and $D_1^*(2760)^0$ resonances.

³ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S -wave and the $D\pi$ S - and P -waves.

⁴ Modeling the $\pi^+ \pi^-$ S -wave with the Isobar formalism.

⁵ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 .

⁶ From the fit of the $M(D^0 \pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1 , respectively.

⁷ At a fixed width of 50.5 MeV.

⁸ Fit includes the contribution from $D_0^*(2400)^0$.

⁹ Modeling the $\pi^+ \pi^-$ S -wave with the K-matrix formalism.


¹⁰ Calculated using the mass difference $m(D_2^{*0}) - m(D^{*+})_{PDG}$ reported below and $m(D^{*+})_{PDG} = 2010.27 \pm 0.17$ MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of $^{+1.2}_{-1.3}$ MeV.

¹¹ Fit includes the contribution from $D_0^*(2400)^\pm$. Not independent of the corresponding mass difference measurement, $(m_{D_2^*(2460)^\pm}) - (m_{D_2^*(2460)^0})$.

NODE=M254M;LINKAGE=LC

¹² No systematic error given.

NODE=M254M;LINKAGE=K



$m_{D_2^*(2460)^0} - m_{D^+}$

NODE=M254DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M254DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
591.5±0.8 OUR FIT	Error includes scale factor of 6.0.			[591.5 ^{+0.7} _{-0.8} MeV OUR 2023 FIT Scale factor = 5.9]
593.9±0.6±0.5	20k	ABULENCIA	06A CDF	1900 $p\bar{p} \rightarrow D^+ \pi^- X$

NODE=M254DM

NEW

$m_{D_2^*(2460)^0} - m_{D^{*+}}$

NODE=M254DM2

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M254DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
450.9±0.8 OUR FIT	Error includes scale factor of 6.0.			[450.9 ^{+0.7} _{-0.8} MeV OUR 2023 FIT Scale factor = 5.9]
458.8±3.7^{+1.2}_{-1.3}	1.5k	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{(*)+} \pi^- X$

NODE=M254DM2

NEW

$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0}$

NODE=M254DMC

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2.4±1.7 OUR AVERAGE			
3.1±1.9±0.9	LINK	04A FOCS	γA
- 2 ±4 ±4	BERGFELD	94B CLE2	$e^+ e^- \rightarrow \text{hadrons}$
0 ±4	FRABETTI	94B E687	$\gamma Be \rightarrow D\pi X$
14 ±5 ±8	ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ X$

NODE=M254DMC

 $D_2^*(2460)$ WIDTH

NODE=M254W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
47.3± 0.8 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.			
47.0± 0.8± 1.0	28k	¹ AAIJ	16AH LHCb	0	$B^- \rightarrow D^+ \pi^- \pi^-$
43.8± 2.9± 1.8	2k	² AAIJ	15V LHCb	0	$B^- \rightarrow D^+ K^- \pi^-$
46.0± 3.4± 3.2		³ AAIJ	15X LHCb	+	$B^0 \rightarrow \bar{D}^0 K^+ \pi^-$
47.3± 1.5± 0.7		⁴ AAIJ	15Y LHCb	+	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
43.2± 1.2± 3.0	82k	AAIJ	13CC LHCb	0	$pp \rightarrow D^{*+} \pi^- X$
45.6± 0.4± 1.1	675k	AAIJ	13CC LHCb	0	$pp \rightarrow D^+ \pi^- X$
48.6± 1.3± 1.9	342k	AAIJ	13CC LHCb	+	$pp \rightarrow D^0 \pi^+ X$
46.6± 8.1 ⁺ ₋ 5.9 3.8	2.3k	⁵ ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+} \pi^- X$
50.5± 0.6± 0.7	243k	DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^+ \pi^- X$
41.8± 2.5± 2.9	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+ \pi^- \pi^-$
49.7± 3.8± 6.4	2909	KUZMIN	07 BELL	+	$e^+ e^- \rightarrow \text{hadrons}$
49.2± 2.3± 1.3	20k	ABULENCIA	06A CDF	0	1900 $p\bar{p} \rightarrow D^+ \pi^- X$
45.6± 4.4± 6.7		⁶ ABE	04D BELL	0	$B^- \rightarrow D^+ \pi^- \pi^-$
38.7± 5.3± 2.9	5.8k	⁶ LINK	04A FOCS	0	γA
34.1± 6.5± 4.2	3.5k	⁷ LINK	04A FOCS	+	γA
28 ⁺ ₋ 8 7 ± 6	486	AVERY	94C CLE2	0	$e^+ e^- \rightarrow D^+ \pi^- X$
27 ⁺ ₋ 11 8 ± 5	310	BERGFELD	94B CLE2	+	$e^+ e^- \rightarrow D^0 \pi^+ X$
25 ±10 ± 5	128	FRABETTI	94B E687	0	$\gamma Be \rightarrow D^+ \pi^- X$
23 ± 9 ± 5	185	FRABETTI	94B E687	+	$\gamma Be \rightarrow D^0 \pi^+ X$
20 ⁺ ₋ 9 ⁺ 12 ⁻ -10 ± 440		AVERY	90 CLEO	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
15 ⁺ ₋ 13 ⁺ 10 ⁻ -10 ± 337		ALBRECHT	89B ARG	0	$e^+ e^- \rightarrow D^+ \pi^- X$
20 ±10 ± 5	153	ANJOS	89C TPS	0	$\gamma N \rightarrow D^+ \pi^- X$

NODE=M254W

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

46.0± 1.4± 1.8 ⁸ AAIJ 15Y LHCb + $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$

NODE=M254W;LINKAGE=D

¹ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

² From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, nonresonant spin-0 and spin-1 components as well as the $D_0^*(2400)^0$, $D_2^*(2460)^0$ and $D_1^*(2760)^0$ resonances.

NODE=M254W;LINKAGE=A

³ From the Dalitz plot analysis including various K^* and D^{**} mesons as well as broad structures in the $K\pi$ S-wave and the $D\pi$ S- and P-waves.

NODE=M254W;LINKAGE=AC

⁴ Modeling the $\pi^+ \pi^-$ S-wave with the Isobar formalism.

NODE=M254W;LINKAGE=B

⁵ From the combined fit of the $M(D^+ \pi^-)$ and $M(D^{*+} \pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1.

NODE=M254W;LINKAGE=AR

⁶ Fit includes the contribution from $D_0^*(2400)^0$.

NODE=M254W;LINKAGE=LI


⁷ Fit includes the contribution from $D_0^*(2400)^\pm$.

NODE=M254W;LINKAGE=LC

⁸ Modeling the $\pi^+ \pi^-$ S-wave with the K-matrix formalism.

NODE=M254W;LINKAGE=C

WEIGHTED AVERAGE
47.3±0.8 (Error scaled by 1.5)



$D_2^*(2460)$ DECAY MODES

NODE=M254215;NODE=M254

$\bar{D}_2^*(2460)$ modes are charge conjugates of modes below.

NODE=M254

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\pi^-$	seen
Γ_2 $D^*(2010)\pi^-$	seen
Γ_3 $D\pi^+\pi^-$	
Γ_4 $D^*\pi^+\pi^-$	

DESIG=1
DESIG=2
DESIG=3
DESIG=4

$D_2^*(2460)$ BRANCHING RATIOS

NODE=M254220

$\Gamma(D\pi^-)/\Gamma_{total}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	3.4k	AUBERT	09AB BABR	0	$B^- \rightarrow D^+\pi^-\pi^-$	
seen	337	ALBRECHT	89B ARG	0	$e^+e^- \rightarrow D^+\pi^-X$	
seen		ALBRECHT	89F ARG	+	$e^+e^- \rightarrow D^0\pi^+X$	
seen		ANJOS	89C TPS	0	$\gamma N \rightarrow D^+\pi^-X$	

NODE=M254R1
NODE=M254R1

$\Gamma(D^*(2010)\pi^-)/\Gamma_{total}$						Γ_2/Γ
VALUE	DOCUMENT ID	TECN	CHG	COMMENT		
seen	ACKERSTAFF 97W	OPAL	0	$e^+e^- \rightarrow D^{*+}\pi^-X$		
seen	AVERY 90	CLEO	0	$e^+e^- \rightarrow D^{*+}\pi^-X$		
seen	ALBRECHT 89H	ARG	0	$e^+e^- \rightarrow D^*\pi^-X$		

NODE=M254R2
NODE=M254R2

$\Gamma(D\pi^-)/\Gamma(D^*(2010)\pi^-)$						Γ_1/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
1.52±0.14 OUR AVERAGE						
1.4 ±0.3 ±0.3	2.3k	1 ABRAMOWICZ13	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$	
1.1 ±0.4 $\begin{smallmatrix} +0.3 \\ -0.2 \end{smallmatrix}$	1371	2 ABRAMOWICZ13	ZEUS	+	$e^\pm p \rightarrow D^{(*)0}\pi^+X$	OCCUR=2
1.47±0.03±0.16	379k	DEL-AMO-SA...10P	BABR	0	$e^+e^- \rightarrow D^{(*)+}\pi^-X$	
2.8 ±0.8 $\begin{smallmatrix} +0.5 \\ -0.6 \end{smallmatrix}$	1.5k	CHEKANOV 09	ZEUS	0	$e^\pm p \rightarrow D^{(*)+}\pi^-X$	
2.2 ±0.7 ±0.6		AVERY 94C	CLE2	0	$e^+e^- \rightarrow D^{*+}\pi^-X$	

NODE=M254R3
NODE=M254R3

1.9 $\pm 1.1 \pm 0.3$ BERGFELD 94B CLE2 + $e^+e^- \rightarrow \text{hadrons}$
 2.3 ± 0.8 AVERY 90 CLEO 0 e^+e^-
 3.0 $\pm 1.1 \pm 1.5$ ALBRECHT 89H ARG 0 $e^+e^- \rightarrow D^*\pi^-X$
 ●●● We do not use the following data for averages, fits, limits, etc. ●●●

OCCUR=2

1.9 ± 0.5 ABE 04D BELL 0 $B^- \rightarrow D^{(*)+}\pi^-\pi^-$
¹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions. and A_{D_2} fixed to the theoretical prediction of -1 .
² From the fit of the $M(D^0\pi^+)$ distribution. The widths of the D_1^+ and D_2^{*+} are fixed to 25 MeV and 37 MeV, and A_{D_1} and A_{D_2} are fixed to the theoretical predictions of 3 and -1 , respectively.

NODE=M254R3;LINKAGE=AR

NODE=M254R3;LINKAGE=AB

$\Gamma(D\pi^-)/[\Gamma(D\pi^-) + \Gamma(D^*(2010)\pi^-)]$ $\Gamma_1/(\Gamma_1+\Gamma_2)$

VALUE EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M254R01
 NODE=M254R01

●●● We do not use the following data for averages, fits, limits, etc. ●●●

0.62 $\pm 0.03 \pm 0.02$ 8414 ¹ AUBERT 09Y BABR 0 $B^+ \rightarrow D_2^{*0}\ell^+\nu_\ell$
 0.62 $\pm 0.03 \pm 0.02$ 3361 ¹ AUBERT 09Y BABR + $\bar{B}^0 \rightarrow D_2^{*+}\ell^-\nu_\ell$

OCCUR=2

¹ Assuming $\Gamma(\Upsilon(4S) \rightarrow B^+B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 1.065 \pm 0.026$ and equal partial widths for charged and neutral D_2^* mesons.

NODE=M254R01;LINKAGE=AU

$D_2^*(2460)$ POLARIZATION AMPLITUDE A_{D_2}

NODE=M254PAM

A polarization amplitude A_{D_2} is a parameter that depends on the initial polarization of the D_2 . For D_2 decays the helicity angle, θ_H , distribution varies like $1 + A_{D_2} \cos^2(\theta_H)$, where θ_H is the angle in the D^* rest frame between the two pions emitted by the $D_2 \rightarrow D^*\pi$ and $D^* \rightarrow D\pi$.

NODE=M254PAM

VALUE EVTS DOCUMENT ID TECN CHG COMMENT

NODE=M254PAM

●●● We do not use the following data for averages, fits, limits, etc. ●●●

-1.16 ± 0.35 2.3k ¹ ABRAMOWICZ13 ZEUS 0 $e^\pm p \rightarrow D^{(*)+}\pi^-X$
 consistent with -1 243k DEL-AMO-SA...10P BABR 0 $e^+e^- \rightarrow D^+\pi^-X$
 $-0.74^{+0.49}_{-0.38}$ ² AVERY 94C CLE2 0 $e^+e^- \rightarrow D^{*+}\pi^-X$

¹ From the combined fit of the $M(D^+\pi^-)$ and $M(D^{*+}\pi^-)$ distributions.

NODE=M254PAM;LINKAGE=AB

² Systematic uncertainties not estimated.

NODE=M254PAM;LINKAGE=AV

$D_2^*(2460)$ REFERENCES

NODE=M254

AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57518
AAIJ	15V	PR D91 092002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56575
Also		PR D93 119901 (errata.)	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57289
AAIJ	15X	PR D92 012012	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56588
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56609
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
ABRAMOWICZ 13	NP	B866 229	H. Abramowicz <i>et al.</i>	(ZEUS Collab.)	REFID=54743
DEL-AMO-SA...10P	PR	D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534
AUBERT	09AB	PR D79 112004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52941
AUBERT	09Y	PRL 103 051803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52929
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
KUZMIN	07	PR D76 012006	A. Kuzmin <i>et al.</i>	(BELLE Collab.)	REFID=51854
ABULENCIA	06A	PR D73 051104	A. Abulencia <i>et al.</i>	(CDF Collab.)	REFID=51054
ABE	04D	PR D69 112002	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50011
LINK	04A	PL B586 11	J.M. Link <i>et al.</i>	(FOCUS Collab.)	REFID=49775
ABREU	98M	PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46315
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN	95	ZPHY C68 43	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)	REFID=44439
AVERY	94C	PL B331 236	P. Avery <i>et al.</i>	(CLEO Collab.)	REFID=44096
BERGFELD	94B	PL B340 194	T. Bergfeld <i>et al.</i>	(CLEO Collab.)	REFID=44099
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT	89B	PL B221 422	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=40736
ALBRECHT	89F	PL B231 208	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40931
ALBRECHT	89H	PL B232 398	H. Albrecht <i>et al.</i>	(ARGUS Collab.) JP	REFID=41001
ANJOS	89C	PRL 62 1717	J.C. Anjos <i>et al.</i>	(FNAL E691 Collab.)	REFID=40737

REFID=40737

$D_0(2550)^0$

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

 $J^P = 0^-$ determined by AAIJ 20D.

NODE=M198

NODE=M198

NODE=M198M


NODE=M198M

 $D_0(2550)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2549 ±19 OUR AVERAGE		Error includes scale factor of 4.6. See the ideogram below.		
2518 ± 2 ±7	79k	¹ AAIJ	20D LHCb	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2579.5 ± 3.4 ±5.5	60k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$
2539.4 ± 4.5 ±6.8	34k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.

NODE=M198M;LINKAGE=A

 $D_0(2550)^0$ WIDTH


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
165 ±24 OUR AVERAGE		Error includes scale factor of 2.0. See the ideogram below.		
199 ± 5 ±17	79k	¹ AAIJ	20D LHCb	$B^- \rightarrow D^{*+} \pi^- \pi^-$
177.5 ±17.8 ±46.0	60k	AAIJ	13CC LHCb	$pp \rightarrow D^{*+} \pi^- X$
130 ±12 ±13	34k	DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{*+} \pi^- X$

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.

NODE=M198W

NODE=M198W

NODE=M198W;LINKAGE=A



$D_0(2550)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^{*+} \pi^-$	seen

NODE=M198215;NODE=M198

DESIG=1;OUR EVAL;→ UNCHECKED ←

$D_0(2550)^0$ POLARIZATION AMPLITUDE A_{D_J}

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$ decays.

NODE=M198PAM

NODE=M198PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M198PAM

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●
4.2±1.3 60k ¹ AAIJ 13CC LHCb $pp \rightarrow D^{*+} \pi^- X$

¹Systematic uncertainty not estimated.

NODE=M198PAM;LINKAGE=A

$D_0(2550)^0$ REFERENCES

NODE=M198

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA... 10P	PR D82	111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

REFID=60253
REFID=55581
REFID=53534

$D_1^*(2600)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

OMITTED FROM SUMMARY TABLE

was $D_j^*(2600)$ $J^P = 1^-$ determined by AAIJ 20D.

NODE=M199

NODE=M199

NODE=M199M

NODE=M199M

OCCUR=2


NODE=M199M;LINKAGE=B

NODE=M199M;LINKAGE=A

NODE=M199M;LINKAGE=DE

 $D_1^*(2600)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2627 ±10	OUR AVERAGE	Error includes scale factor of 4.4. See the ideogram below.			
2641.9 ± 1.8 ± 4.5	79k	¹ AAIJ	20D	LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2681.1 ± 5.6 ± 14.0	28k	² AAIJ	16AH	LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
2649.2 ± 3.5 ± 3.5	51k	AAIJ	13CC	LHCB	$pp \rightarrow D^{*+} \pi^- X$
2608.7 ± 2.4 ± 2.5	26k	DEL-AMO-SA...10P	BABR	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2621.3 ± 3.7 ± 4.2	13k	³ DEL-AMO-SA...10P	BABR	+	$e^+ e^- \rightarrow D^0 \pi^+ X$

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.² From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.³ At a fixed width of 93 MeV. $D_1^*(2600)^0$ WIDTH

NODE=M199W


NODE=M199W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
141 ±23	OUR AVERAGE	Error includes scale factor of 2.7. See the ideogram below.		
149 ± 4 ± 20	79k	¹ AAIJ	20D	LHCB $B^- \rightarrow D^{*+} \pi^- \pi^-$
186.7 ± 8.5 ± 11.9	28k	² AAIJ	16AH	LHCB $B^- \rightarrow D^+ \pi^- \pi^-$
140.2 ± 17.1 ± 18.6	51k	AAIJ	13CC	LHCB $pp \rightarrow D^{*+} \pi^- X$
93 ± 6 ± 13	26k	DEL-AMO-SA...10P	BABR	$e^+ e^- \rightarrow D^+ \pi^- X$

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.² From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

NODE=M199W;LINKAGE=B

NODE=M199W;LINKAGE=A



$D_1^*(2600)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\pi$	seen
Γ_2 $D^+\pi^-$	seen
Γ_3 $D^0\pi^\pm$	seen
Γ_4 $D^*\pi$	seen
Γ_5 $D^{*+}\pi^-$	seen

NODE=M199215;NODE=M199

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=3;OUR EVAL;→ UNCHECKED ←
 DESIG=4;OUR EVAL;→ UNCHECKED ←
 DESIG=5;OUR EVAL;→ UNCHECKED ←

$D_1^*(2600)^0$ BRANCHING RATIOS

$\Gamma(D^+\pi^-)/\Gamma(D^{*+}\pi^-)$					Γ_2/Γ_5
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.32±0.02±0.09	76k	DEL-AMO-SA...10P	BABR	$e^+e^- \rightarrow D^{(*)+}\pi^- X$	

NODE=M199220

NODE=M199R01
 NODE=M199R01

$D_1^*(2600)^0$ REFERENCES

AAIJ 20D PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ 16AH PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ 13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)
DEL-AMO-SA...10P PR D82 111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)

NODE=M199

REFID=60253
 REFID=57518
 REFID=55581
 REFID=53534

$D^*(2640)^\pm$

$$I(J^P) = \frac{1}{2}(??)$$

OMITTED FROM SUMMARY TABLE

Seen in Z decays by ABREU 98M. Not seen by ABBIENDI 01N and CHEKANOV 09. Needs confirmation.

NODE=M158

NODE=M158

 $D^*(2640)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2637 ± 2 ± 6	66 ± 14	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

NODE=M158M

NODE=M158M

 $D^*(2640)^\pm$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<15	95	ABREU	98M DLPH	$e^+ e^- \rightarrow D^{*+} \pi^+ \pi^- X$

NODE=M158W

NODE=M158W

 $D^*(2640)^+$ DECAY MODES $D^*(2640)^-$ modes are charge conjugates of modes below.

NODE=M158215;NODE=M158

NODE=M158

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2010)^+ \pi^+ \pi^-$	seen

DESIG=1;OUR EST;→ UNCHECKED ←

 $D^*(2640)^\pm$ REFERENCES

CHEKANOV 09 EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)
ABBIENDI 01N EPJ C20 445	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
ABREU 98M PL B426 231	P. Abreu <i>et al.</i>	(DELPHI Collab.)

NODE=M158

REFID=52733

REFID=48296

REFID=46315

NODE=M228

 $D_2(2740)^0$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

was $D(2740)^0$ $J^P = 2^-$ determined by (AAIJ 20D).

NODE=M228

 $D_2(2740)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2747 ± 6 OUR AVERAGE				
2751 ± 3 ± 7	79k	¹ AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2737.0 ± 3.5 ± 11.2	7.7k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$

NODE=M228M

NODE=M228M

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.

NODE=M228M;LINKAGE=A

 $D_2(2740)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
88 ± 19 OUR AVERAGE				
102 ± 6 ± 26	79k	¹ AAIJ	20D LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
73.2 ± 13.4 ± 25.0	7.7k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$

NODE=M228W

NODE=M228W

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.

NODE=M228W;LINKAGE=A

$D_2(2740)^0$ DECAY MODES

NODE=M228215;NODE=M228

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^{*+} \pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D_2(2740)^0$ POLARIZATION AMPLITUDE A_{D_J}

NODE=M228PAM

NODE=M228PAM

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$ decays.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M228PAM

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

3.1±2.2 7.7k ¹ AAIJ 13CC LHCB $pp \rightarrow D^{*+} \pi^- X$ ¹ Systematic uncertainty not estimated.

NODE=M228PAM;LINKAGE=A

 $D_2(2740)^0$ REFERENCES

NODE=M228

AAIJ	20D PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	13CC JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=60253
REFID=55581

NODE=M203

 $D_3^*(2750)$

$$I(J^P) = \frac{1}{2}(3^-)$$

J^P determined by AAIJ 15Y from the Dalitz plot analysis of $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$ decays.

NODE=M203

 $D_3^*(2750)$ MASS

NODE=M203M

NODE=M203M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2763.1± 3.2 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.			
2753 ± 4 ± 6	79k	¹ AAIJ	20D	LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
2775.5± 4.5± 6.5	28k	² AAIJ	16AH	LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
2798 ± 7 ± 7		³ AAIJ	15Y	LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
2761.1± 5.1± 6.5	14k	AAIJ	13CC	LHCB 0	$pp \rightarrow D^{*+} \pi^- X$
2760.1± 1.1± 3.7	56k	AAIJ	13CC	LHCB 0	$pp \rightarrow D^+ \pi^- X$
2771.7± 1.7± 3.8	20k	AAIJ	13CC	LHCB +	$pp \rightarrow D^0 \pi^+ X$
2752.4± 1.7± 2.7	23.5k	⁴ DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2763.3± 2.3± 2.3	11.3k	⁴ DEL-AMO-SA..10P	BABR	0	$e^+ e^- \rightarrow D^+ \pi^- X$
2769.7± 3.8± 1.5	5.7k	^{4,5} DEL-AMO-SA..10P	BABR	+	$e^+ e^- \rightarrow D^0 \pi^+ X$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

2802 ±11 ±10 ⁶ AAIJ 15Y LHCB $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$

OCCUR=2

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.

NODE=M203M;LINKAGE=D

² From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

NODE=M203M;LINKAGE=C


³ Modeling the $\pi^+ \pi^-$ S-wave with the Isobar formalism.⁴ The states observed in the $D^* \pi$ and $D \pi$ final states are not necessarily the same.⁵ At a fixed width of 60.9 MeV.⁶ Modeling the $\pi^+ \pi^-$ S-wave with the K-matrix formalism.

NODE=M203M;LINKAGE=A

NODE=M203M;LINKAGE=DE

NODE=M203M;LINKAGE=DA

NODE=M203M;LINKAGE=B



$D_3^*(2750)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
66 ± 5	OUR AVERAGE				
66 ± 10 ± 14	79k	1 AAIJ	20D	LHCB	$B^- \rightarrow D^{*+} \pi^- \pi^-$
95.3 ± 9.6 ± 34.0	28k	2 AAIJ	16AH	LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
105 ± 18 ± 24		3 AAIJ	15Y	LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
74.4 ± 3.4 ± 37.0	14k	AAIJ	13CC	LHCB 0	$pp \rightarrow D^{*+} \pi^- X$
74.4 ± 3.4 ± 19.1	56k	AAIJ	13CC	LHCB 0	$pp \rightarrow D^+ \pi^- X$
66.7 ± 6.6 ± 10.5	20k	AAIJ	13CC	LHCB +	$pp \rightarrow D^0 \pi^+ X$
71 ± 6 ± 11	23.5k	4 DEL-AMO-SA..10P	BABR		$e^+ e^- \rightarrow D^{*+} \pi^- X$
60.9 ± 5.1 ± 3.6	11.3k	4 DEL-AMO-SA..10P	BABR		$e^+ e^- \rightarrow D^+ \pi^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

154 ± 27 ± 16		5 AAIJ	15Y	LHCB	$B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$
---------------	--	--------	-----	------	---

¹ From a full four-body amplitude analysis of the $B^- \rightarrow D^{*+} \pi^- \pi^-$ decay.

² From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

³ Modeling the $\pi^+ \pi^-$ S-wave with the Isobar formalism.

⁴ The states observed in the $D^* \pi$ and $D \pi$ final states are not necessarily the same.

⁵ Modeling the $\pi^+ \pi^-$ S-wave with the K-matrix formalism.

$D_3^*(2750)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D \pi$	seen
Γ_2 $D^+ \pi^-$	seen
Γ_3 $D^0 \pi^\pm$	seen
Γ_4 $D^* \pi$	seen
Γ_5 $D^{*+} \pi^-$	seen

$D_3^*(2750)$ BRANCHING RATIOS

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_5
0.42 ± 0.05 ± 0.11	34.8k	1 DEL-AMO-SA..10P	BABR	$e^+ e^- \rightarrow D^{(*)+} \pi^- X$	

NODE=M203W

NODE=M203W

OCCUR=2

OCCUR=4

OCCUR=2

OCCUR=2

NODE=M203W;LINKAGE=D

NODE=M203W;LINKAGE=C

NODE=M203W;LINKAGE=A

NODE=M203W;LINKAGE=DE

NODE=M203W;LINKAGE=B

NODE=M203215;NODE=M203

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=5;OUR EVAL;→ UNCHECKED ←

NODE=M203220

NODE=M203R01

NODE=M203R01

¹ The states observed in the $D^*\pi$ and $D\pi$ final states are not necessarily the same.

NODE=M203R01;LINKAGE=DE

$D_3^*(2750)$ POLARIZATION AMPLITUDE A_D

NODE=M203PAM

NODE=M203PAM

A polarization amplitude A_D is a parameter that depends on the initial polarization of the $D_3^*(2750)$. For $D_3^*(2750)$ decays the helicity angle, θ_H , distribution varies like $1 + A_D \cos(\theta_H)$, where θ_H is the angle in the D^* rest frame between the two pions emitted by the $D_3^*(2750) \rightarrow D^*\pi$ and $D^* \rightarrow D\pi$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M203PAM

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.33±0.28	23.5k	¹ DEL-AMO-SA..10P	BABR	$e^+e^- \rightarrow D^{*+}\pi^- X$
------------	-------	------------------------------	------	------------------------------------

NODE=M203PAM;LINKAGE=DE

¹ Systematic uncertainties not estimated. The states observed in the $D^*\pi$ and $D\pi$ final states are not necessarily the same.

$D_3^*(2750)$ REFERENCES

NODE=M203

AAIJ	20D	PR D101 032005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60253
AAIJ	16AH	PR D94 072001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57518
AAIJ	15Y	PR D92 032002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=56609
AAIJ	13CC	JHEP 1309 145	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55581
DEL-AMO-SA...10P	PR D82	111101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53534

NODE=M249

$D_1^*(2760)^0$

$$I(J^P) = \frac{1}{2}(1^-)$$

OMITTED FROM SUMMARY TABLE

J^P determined by AAIJ 15V from the Dalitz plot analysis of $B^- \rightarrow D^+ K^- \pi^-$ decays.

NODE=M249

$D_1^*(2760)^0$ MASS

NODE=M249M

NODE=M249M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2781±18±13	2k	¹ AAIJ	15V LHCB	$B^- \rightarrow D^+ K^- \pi^-$

¹ From the amplitude analysis in the model describing the $D^+\pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, nonresonant spin-0 and spin-1 components as well as the $D_0^*(2400)^0$, $D_2^*(2460)^0$ and $D_1^*(2760)^0$ resonances.

NODE=M249M;LINKAGE=A

$D_1^*(2760)^0$ WIDTH

NODE=M249W

NODE=M249W

NEW

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
180±40 OUR AVERAGE	[177 ± 40 MeV OUR 2023 AVERAGE]			
177±32±21	2k	¹ AAIJ	15V LHCB	$B^- \rightarrow D^+ K^- \pi^-$

¹ From the amplitude analysis in the model describing the $D^+\pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, nonresonant spin-0 and spin-1 components as well as the $D_0^*(2400)^0$, $D_2^*(2460)^0$ and $D_1^*(2760)^0$ resonances.

NODE=M249W;LINKAGE=A

$D_1^*(2760)^0$ DECAY MODES

NODE=M249215;NODE=M249

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^+\pi^-$	seen

DESIG=1

$D_1^*(2760)^0$ BRANCHING RATIOS

NODE=M249225

$\Gamma(D^+\pi^-)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		¹ AAIJ	15V LHCB	$B^- \rightarrow D^+ K^- \pi^-$	

NODE=M249R01

NODE=M249R01

OCCUR=2

¹ From the amplitude analysis in the model describing the $D^+\pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, nonresonant spin-0 and spin-1 components as well as the $D_0^*(2400)^0$, $D_2^*(2460)^0$ and $D_1^*(2760)^0$ resonances.

NODE=M249R01;LINKAGE=A

$D_1^*(2760)^0$ REFERENCES

AAIJ 15V PR D91 092002 R. Aaij *et al.* (LHCb Collab.) JP
 Also PR D93 119901 (errat.) R. Aaij *et al.* (LHCb Collab.)

NODE=M249

REFID=56575
REFID=57289

NODE=M229

 $D(3000)^0$

$$I(J^P) = \frac{1}{2}(?)$$

OMITTED FROM SUMMARY TABLE

Both natural- and unnatural-parity components observed depending on the decay mode (AAIJ 13CC).

NODE=M229

 $D(3000)^0$ MASS

NODE=M229M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3210 ±60 OUR AVERAGE		[3214 ± 60 MeV OUR 2023 AVERAGE]		
3214 ±29 ±49	28k	¹ AAIJ	16AH LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2971.8 ± 8.7	9.5k	^{2,3} AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
3008.1 ± 4.0	17.6k	^{2,4} AAIJ	13CC LHCB	$pp \rightarrow D^+ \pi^- X$

NODE=M229M

NEW

OCCUR=2

NODE=M229M;LINKAGE=D

¹ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

² Systematic uncertainty not estimated.

³ Unnatural parity preferred.

⁴ Natural parity state. A state $D(3000)^+$ is possibly seen in $D^0 \pi^+$ final state.

NODE=M229M;LINKAGE=A

NODE=M229M;LINKAGE=B

NODE=M229M;LINKAGE=C

 $D(3000)^0$ WIDTH

NODE=M229W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
190 ±80 OUR AVERAGE		[186 ± 80 MeV OUR 2023 AVERAGE]		
186 ±38 ±72	28k	⁵ AAIJ	16AH LHCB	$B^- \rightarrow D^+ \pi^- \pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
188.1 ±44.8	9.5k	^{6,7} AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$
110.5 ±11.5	17.6k	^{6,8} AAIJ	13CC LHCB	$pp \rightarrow D^+ \pi^- X$

NODE=M229W

NEW

OCCUR=2

NODE=M229W;LINKAGE=D

⁵ From the amplitude analysis in the model describing the $D^+ \pi^-$ wave together with virtual contributions from the $D^*(2007)^0$ and B^{*0} states, and components corresponding to the $D_2^*(2460)^0$, $D_1^*(2680)^0$, $D_3^*(2760)^0$, and $D_2^*(3000)^0$ resonances.

⁶ Systematic uncertainty not estimated.

⁷ Unnatural parity preferred.

⁸ Natural parity state. A state $D(3000)^+$ is possibly seen in $D^0 \pi^+$ final state.

NODE=M229W;LINKAGE=A

NODE=M229W;LINKAGE=C

NODE=M229W;LINKAGE=B

 $D(3000)^0$ DECAY MODES

NODE=M229215;NODE=M229

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^{*+} \pi^-$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←

 $D(3000)^0$ POLARIZATION AMPLITUDE A_{D_J}

NODE=M229PAM

A polarization amplitude A_{D_J} is a parameter that depends on the initial polarization of the D_J . For D_J decays the helicity angle, θ_H , distribution varies like $1 + A_{D_J} \cos^2(\theta_H)$, where θ_H is the angle in the D_J rest frame between the two pions emitted in the $D_J \rightarrow D^* \pi$ and $D^* \rightarrow D \pi$ decays.

NODE=M229PAM

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.5 ±0.9	9.5k	⁹ AAIJ	13CC LHCB	$pp \rightarrow D^{*+} \pi^- X$

NODE=M229PAM

⁹ Systematic uncertainty not estimated.

NODE=M229PAM;LINKAGE=A

 $D(3000)^0$ REFERENCES

NODE=M229

AAIJ 16AH PR D94 072001 R. Aaij *et al.* (LHCb Collab.)
 AAIJ 13CC JHEP 1309 145 R. Aaij *et al.* (LHCb Collab.)

REFID=57518
REFID=55581

CHARMED, STRANGE MESONS

($C = \pm 1, S = \pm 1$)

(including possibly non- $q\bar{q}$ states)

$$D_s^+ = c\bar{s}, D_s^- = \bar{c}s, \text{ similarly for } D_s^{*\pm}$$

 $D_s^{*\pm}$

$$I(J^P) = 0(1^-)$$

 $J^P = 1^-$ established by ABLIKIM 23AZ.

NODE=MXXX040

NODE=MXXX040

NODE=S074

NODE=S074

NODE=S074M

NODE=S074M

NODE=S074M

NODE=S074M;LINKAGE=E

NODE=S074DM

NODE=S074DM

NODE=S074DM

NODE=S074DM;LINKAGE=A

NODE=S074W

NODE=S074W

NODE=S074215;NODE=S074

NODE=S074

$D_s^{*\pm}$ MASS

The fit includes $D_s^\pm, D_s^0, D_s^{\pm*}, D_s^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2112.2 ± 0.4 OUR FIT			
2106.6 ± 2.1 ± 2.7	¹ BLAYLOCK	87	MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

¹ Assuming D_s^\pm mass = 1968.7 ± 0.9 MeV.

$m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes $D_s^\pm, D_s^0, D_s^{\pm*}, D_s^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
143.8 ± 0.4 OUR FIT				
143.9 ± 0.4 OUR AVERAGE				
143.76 ± 0.39 ± 0.40		GRONBERG	95	CLE2 e^+e^-
144.22 ± 0.47 ± 0.37		BROWN	94	CLE2 e^+e^-
142.5 ± 0.8 ± 1.5		² ALBRECHT	88	ARG $e^+e^- \rightarrow D_s^\pm \gamma X$
139.5 ± 8.3 ± 9.7	60	AIHARA	84D	TPC $e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
143.0 ± 18.0	8	ASRATYAN	85	HLBC FNAL 15-ft, ν - ² H
110 ± 46		BRANDELIK	79	DASP $e^+e^- \rightarrow D_s^\pm \gamma X$

² Result includes data of ALBRECHT 84B.

$D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.9	90	GRONBERG	95	CLE2 e^+e^-
< 4.5	90	ALBRECHT	88	ARG $E_{cm}^{ee} = 10.2$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 4.9	90	BROWN	94	CLE2 e^+e^-
< 22	90	BLAYLOCK	87	MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

D_s^{*+} DECAY MODES

D_s^{*-} modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D_s^+ \gamma$	(93.6 ± 0.4) %
Γ_2 $D_s^+ \pi^0$	(5.77 ± 0.35) %
Γ_3 $D_s^+ e^+ e^-$	(6.7 ± 1.6) × 10 ⁻³
Γ_4 $e^+ \nu_e$	(2.1 $^{+1.2}_{-0.9}$) × 10 ⁻⁵

DESIG=1

DESIG=2

DESIG=3

DESIG=4

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.0$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-91	
x_3	-38	-4
	x_1	x_2

D_s^{*+} BRANCHING RATIOS

NODE=S074220

$\Gamma(D_s^{*+} \gamma) / \Gamma_{\text{total}}$ Γ_1 / Γ
 VALUE (%) DOCUMENT ID TECN COMMENT

NODE=S074R1
NODE=S074R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ASRATYAN	91	HLBC	$\bar{\nu}_\mu \text{Ne}$
seen	ALBRECHT	88	ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$
seen	AIHARA	84D		
seen	ALBRECHT	84B		
seen	BRANDELIK	79		

$\Gamma(D_s^{*+} \pi^0) / \Gamma(D_s^{*+} \gamma)$ Γ_2 / Γ_1
 VALUE (units 10^{-2}) DOCUMENT ID TECN COMMENT

NODE=S074R2
NODE=S074R2**6.2 ± 0.4 OUR FIT**

[0.062 ± 0.008 OUR 2023 FIT]

NEW

6.2 ± 0.4 OUR AVERAGE

[0.062 ± 0.008 OUR 2023 AVERAGE]

NEW

6.16 ± 0.43 ± 0.18	ABLIKIM	23P	BES3	$e^+ e^-$
6.2 ± 0.5 ± 0.6	AUBERT, BE	05G	BABR	10.6 $e^+ e^- \rightarrow$ hadrons
6.2 $^{+2.0}_{-1.8}$ ± 2.2	GRONBERG	95	CLE2	$e^+ e^-$

$\Gamma(D_s^{*+} e^+ e^-) / \Gamma(D_s^{*+} \gamma)$ Γ_3 / Γ_1
 VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

NODE=S074R01
NODE=S074R01**7.2 ± 1.7 OUR FIT**

7.2 $^{+1.5}_{-1.3}$ ± 1.0	38	CRONIN-HEN..12	CLEO	4.17 $e^+ e^- \rightarrow$ hadrons
--	----	----------------	------	------------------------------------

$\Gamma(e^+ \nu_e) / \Gamma_{\text{total}}$ Γ_4 / Γ
 VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

NODE=S074R02
NODE=S074R02

2.1 $^{+1.2}_{-0.9}$ ± 0.2	ABLIKIM	23BF	BES3	$e^+ e^- \rightarrow D_s^- D_s^{*+}$
--	---------	------	------	--------------------------------------

$D_s^{*\pm}$ REFERENCES

NODE=S074

ABLIKIM	23AZ	PL B846	138245	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=62420
ABLIKIM	23BF	PRL 131	141802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62430
ABLIKIM	23P	PR D107	032011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62059
CRONIN-HEN...12	PR D86	072005		D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=54627
AUBERT, BE	05G	PR D72	091101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50942
GRONBERG	95	PRL 75	3232	J. Gronberg <i>et al.</i>	(CLEO Collab.)	REFID=44568
BROWN	94	PR D50	1884	D. Brown <i>et al.</i>	(CLEO Collab.)	REFID=43868
ASRATYAN	91	PL B257	525	A.E. Asratyan <i>et al.</i>	(ITEP, BELG, SACL+)	REFID=41582
ALBRECHT	88	PL B207	349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40269
BLAYLOCK	87	PRL 58	2171	G.T. Blaylock <i>et al.</i>	(Mark III Collab.)	REFID=40005
ASRATYAN	85	PL 156B	441	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=22887
AIHARA	84D	PRL 53	2465	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=11561
ALBRECHT	84B	PL 146B	111	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22886
BRANDELIK	79	PL 80B	412	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=11442

$$D_{s0}^*(2317)^\pm$$

$$I(J^P) = 0(0^+)$$

J, P need confirmation.

AUBERT 06P and CHOI 15A do not observe neutral and doubly charged partners of the $D_{s0}^*(2317)^\pm$. See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M172

NODE=M172

$D_{s0}^*(2317)^\pm$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M172M

NODE=M172M

NODE=M172M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2317.8±0.5 OUR FIT				
2318.0±0.7 OUR AVERAGE				
2318.3±1.2±1.2	115	¹ ABLIKIM	18J BES3	4.6 $e^+e^- \rightarrow D_s^{*\pm} D_{s0}^*(2317)^\mp$
2319.6±0.2±1.4	3.1k	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow D_s^+ \pi^0 X$
2317.3±0.4±0.8	1.0k	² AUBERT	04E BABR	10.6 e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2317.2±1.3	88	³ AUBERT,B	04S BABR	$B \rightarrow D_{s0}^{(*)}(2317)^+ \bar{D}^{(*)}$
2317.2±0.5±0.9	761	⁴ MIKAMI	04 BELL	10.6 e^+e^-
2316.8±0.4±3.0	1.2k	^{4,5} AUBERT	03G BABR	10.6 e^+e^-
2317.6±1.3	273	^{4,6} AUBERT	03G BABR	10.6 e^+e^-
2319.8±2.1±2.0	24	⁴ KROKOVNY	03B BELL	10.6 e^+e^-

OCCUR=2

NODE=M172M;LINKAGE=A

NODE=M172M;LINKAGE=AU

NODE=M172M;LINKAGE=AB

NODE=M172M;LINKAGE=B1

NODE=M172M;LINKAGE=A1

NODE=M172M;LINKAGE=A2

¹ From a fit of the D_s^* recoil mass where the $D_{s0}^*(2317)$ signal is described with a Crystal Ball function convolved with a Gaussian function.

² Supersedes AUBERT 03G.

³ Systematic errors not evaluated.

⁴ Not independent of the corresponding $m_{D_{s0}^*(2317)} - m_{D_s}$.

⁵ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.

⁶ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.

$m_{D_{s0}^*(2317)^\pm} - m_{D_s^\pm}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M172DM

NODE=M172DM

NODE=M172DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
349.4±0.5 OUR FIT				
349.2±0.7 OUR AVERAGE				
348.7±0.5±0.7	761	MIKAMI	04 BELL	10.6 e^+e^-
350.0±1.2±1.0	135	BESSION	03 CLE2	10.6 e^+e^-
351.3±2.1±1.9	24	⁷ KROKOVNY	03B BELL	10.6 e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
349.6±0.4±3.0	1267	^{8,9} AUBERT	03G BABR	10.6 e^+e^-
350.2±1.3	273	^{10,11} AUBERT	03G BABR	10.6 e^+e^-
⁷ Recalculated by us using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV.				
⁸ From $D_s^+ \rightarrow K^+ K^- \pi^+$ decay.				
⁹ Recalculated by us using $m_{D_s^+} = 1967.20 \pm 0.03$ MeV.				
¹⁰ From $D_s^+ \rightarrow K^+ K^- \pi^+ \pi^0$ decay.				
¹¹ Recalculated by us using $m_{D_s^+} = 1967.4 \pm 0.2$ MeV. Systematic errors not estimated.				

OCCUR=2

NODE=M172DM;LINKAGE=K3

NODE=M172DM;LINKAGE=A1

NODE=M172DM;LINKAGE=C1

NODE=M172DM;LINKAGE=A2

NODE=M172DM;LINKAGE=C2

$D_{s0}^*(2317)^\pm$ WIDTH

NODE=M172W

NODE=M172W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.8	95	3180	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow D_s^+ \pi^0 X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 4.6	90	761	MIKAMI	04 BELL	10.6 e^+e^-
< 10			AUBERT	03G BABR	10.6 e^+e^-
< 7	90	135	BESSION	03 CLE2	10.6 e^+e^-

$D_{s0}^*(2317)^\pm$ DECAY MODES

NODE=M172215;NODE=M172

 $D_{s0}^*(2317)^-$ modes are charge conjugates of modes below.

NODE=M172

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $D_s^+ \pi^0$	$(100^{+0}_{-20})\%$	
Γ_2 $D_s^+ \gamma$	$< 5\%$	90%
Γ_3 $D_s^*(2112)^+ \gamma$	$< 6\%$	90%
Γ_4 $D_s^+ \gamma \gamma$	$< 18\%$	95%
Γ_5 $D_s^*(2112)^+ \pi^0$	$< 11\%$	90%
Γ_6 $D_s^+ \pi^+ \pi^-$	$< 4 \times 10^{-3}$	90%
Γ_7 $D_s^+ \pi^0 \pi^0$	not seen	

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7;OUR EVAL;→ UNCHECKED ←

 $D_{s0}^*(2317)^\pm$ BRANCHING RATIOS

NODE=M172220

$\Gamma(D_s^+ \pi^0)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
$1.00^{+0.00+0.00}_{-0.14-0.14}$	47	ABLIKIM	18J	BES3	$4.6 e^+ e^- \rightarrow D_s^{*\pm} D_{s0}^*(2317)^\mp$	

NODE=M172R1
NODE=M172R1

• • • We do not use the following data for averages, fits, limits, etc. • • •
seen 1.5k AUBERT 03G BABR $10.6 e^+ e^-$

$\Gamma(D_s^+ \gamma)/\Gamma(D_s^+ \pi^0)$						Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.05	90	MIKAMI	04	BELL	$10.6 e^+ e^-$	
<0.14	95	AUBERT	06P	BABR	$10.6 e^+ e^-$	
<0.052	90	BESSION	03	CLE2	$10.6 e^+ e^-$	

NODE=M172R5
NODE=M172R5

$\Gamma(D_s^*(2112)^+ \gamma)/\Gamma(D_s^+ \pi^0)$						Γ_3/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.059	90	BESSION	03	CLE2	$10.6 e^+ e^-$	
<0.16	95	AUBERT	06P	BABR	$10.6 e^+ e^-$	
<0.18	90	MIKAMI	04	BELL	$10.6 e^+ e^-$	

NODE=M172R6
NODE=M172R6

$\Gamma(D_s^+ \gamma \gamma)/\Gamma(D_s^+ \pi^0)$						Γ_4/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.18	95	AUBERT	06P	BABR	$10.6 e^+ e^-$	
not seen		AUBERT	03G	BABR	$10.6 e^+ e^-$	

NODE=M172R7
NODE=M172R7

$\Gamma(D_s^*(2112)^+ \pi^0)/\Gamma(D_s^+ \pi^0)$						Γ_5/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.11	90	BESSION	03	CLE2	$10.6 e^+ e^-$	

NODE=M172R8
NODE=M172R8

$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma(D_s^+ \pi^0)$						Γ_6/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.004	90	MIKAMI	04	BELL	$10.6 e^+ e^-$	
<0.005	95	AUBERT	06P	BABR	$10.6 e^+ e^-$	
<0.019	90	BESSION	03	CLE2	$10.6 e^+ e^-$	

NODE=M172R9
NODE=M172R9

$\Gamma(D_s^+ \pi^0 \pi^0)/\Gamma(D_s^+ \pi^0)$						Γ_7/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.25	95	AUBERT	06P	BABR	$10.6 e^+ e^-$	

NODE=M172R10
NODE=M172R10

$D_{s0}^*(2317)^\pm$ REFERENCES

ABLIKIM	18J	PR D97 051103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58895
CHOI	15A	PR D91 092011	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=56577
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
AUBERT	03G	PRL 90 242001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49417
BESSION	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

NODE=M172

 $D_{s1}(2460)^\pm$

$$I(J^P) = 0(1^+)$$

See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M173

NODE=M173

 $D_{s1}(2460)^\pm$ MASS

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M173M

NODE=M173M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M173M

2459.5±0.6 OUR FIT Error includes scale factor of 1.1.**2459.6±0.9 OUR AVERAGE** Error includes scale factor of 1.3.2460.1±0.2±0.8 ¹ AUBERT 06P BABR 10.6 e^+e^- 2458.0±1.0±1.0 195 AUBERT 04E BABR 10.6 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

2459.5±1.2±3.7 920 AUBERT 06P BABR 10.6 $e^+e^- \rightarrow D_s^+ \gamma X$

OCCUR=2

2458.6±1.0±2.5 560 AUBERT 06P BABR 10.6 $e^+e^- \rightarrow D_s^+ \pi^0 \gamma X$

OCCUR=3

2460.2±0.2±0.8 123 AUBERT 06P BABR 10.6 $e^+e^- \rightarrow D_s^+ \pi^+ \pi^- X$

OCCUR=4

2458.9±1.5 112 ² AUBERT,B 04S BABR $B \rightarrow D_{s1}(2460)^+ \bar{D}^{(*)}$ 2461.1±1.6 139 ³ AUBERT,B 04S BABR $B \rightarrow D_{s1}(2460)^+ \bar{D}^{(*)}$

OCCUR=2

2456.5±1.3±1.3 126 ^{4,5} MIKAMI 04 BELL 10.6 e^+e^- 2459.5±1.3±2.0 152 ^{6,7} MIKAMI 04 BELL 10.6 e^+e^-

OCCUR=2

2459.9±0.9±1.6 60 ^{6,7} MIKAMI 04 BELL 10.6 e^+e^-

OCCUR=3

2459.2±1.6±2.0 57 KROKOVNY 03B BELL 10.6 e^+e^- ¹ The average of the values obtained from the $D_s^+ \gamma, D_s^+ \pi^0 \gamma, D_s^+ \pi^+ \pi^-$ final state.

NODE=M173M;LINKAGE=UB

² Systematic errors not evaluated. From the decay to $D_s^{*+} \pi^0$.

NODE=M173M;LINKAGE=AU

³ Systematic errors not evaluated. From the decay to $D_s^+ \gamma$.

NODE=M173M;LINKAGE=AB

⁴ Not independent of the corresponding $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$.

NODE=M173M;LINKAGE=B1

⁵ Using $m_{D_s^{*+}} = 2112.4 \pm 0.7$ MeV.

NODE=M173M;LINKAGE=B2

⁶ Not independent of the corresponding $m_{D_{s1}(2460)^\pm} - m_{D_s^\pm}$.

NODE=M173M;LINKAGE=B3

⁷ Using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV.

NODE=M173M;LINKAGE=B4

 $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$

NODE=M173MD

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M173MD


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M173MD

347.3±0.7 OUR FIT Error includes scale factor of 1.2.**347.1±2.2 OUR AVERAGE** Error includes scale factor of 1.9. See the ideogram below.344.1±1.3±1.1 126 MIKAMI 04 BELL 10.6 e^+e^- 351.2±1.7±1.0 41 BESSION 03 CLE2 10.6 e^+e^- 346.8±1.6±1.9 57 ⁸ KROKOVNY 03B BELL 10.6 e^+e^-

⁸ Recalculated by us using $m_{D_s^{*+}} = 2112.4 \pm 0.7$ MeV.

NODE=M173MD;LINKAGE=K3



$m_{D_{s1}(2460)^{\pm}} - m_{D_s^{\pm}}$

NODE=M173DM

The fit includes $D^{\pm}, D^0, D_s^{\pm}, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0,$ and $D_{s1}(2536)^{\pm}$ mass and mass difference measurements.

NODE=M173DM

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M173DM

491.1±0.6 OUR FIT Error includes scale factor of 1.1.

491.3±1.4 OUR AVERAGE

491.0±1.3±1.9	152	⁹ MIKAMI	04	BELL	10.6 $e^+ e^-$
491.4±0.9±1.5	60	¹⁰ MIKAMI	04	BELL	10.6 $e^+ e^-$

OCCUR=2

⁹ From the decay to $D_s^{\pm} \gamma$.

NODE=M173DM;LINKAGE=M1

¹⁰ From the decay to $D_s^{\pm} \pi^+ \pi^-$.

NODE=M173DM;LINKAGE=M2

$D_{s1}(2460)^{\pm}$ WIDTH

NODE=M173W

VALUE (MeV) CL% EVTS DOCUMENT ID TECN COMMENT

NODE=M173W

< 3.5 95 123 AUBERT 06P BABR 10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.3 95 560 AUBERT 06P BABR 10.6 $e^+ e^- \rightarrow D_s^+ \pi^0 \gamma X$

OCCUR=2

<10 195 AUBERT 04E BABR 10.6 $e^+ e^-$

< 5.5 90 126 MIKAMI 04 BELL 10.6 $e^+ e^-$

< 7 90 41 BESSON 03 CLE2 10.6 $e^+ e^-$

$D_{s1}(2460)^+$ DECAY MODES

NODE=M173215;NODE=M173

$D_{s1}(2460)^-$ modes are charge conjugates of the modes below.

NODE=M173

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $D_s^{*+} \pi^0$	(48 ± 11) %	
Γ_2 $D_s^+ \gamma$	(18 ± 4) %	
Γ_3 $D_s^+ \pi^+ \pi^-$	(4.3± 1.3) %	S=1.1
Γ_4 $D_s^{*+} \gamma$	< 8 %	CL=90%

DESIG=1

DESIG=2

DESIG=3

DESIG=4

Γ_5	$D_{s0}^*(2317)^+\gamma$	$(3.7^{+5.0}_{-2.4})\%$	DESIG=5
Γ_6	$D_s^+\pi^0$		DESIG=7
Γ_7	$D_s^+\pi^0\pi^0$		DESIG=8
Γ_8	$D_s^+\gamma\gamma$		DESIG=9

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 8 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 3.4$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	80		
x_3	68	62	
x_5	-3	25	26
	x_1	x_2	x_3

$D_{s1}(2460)^\pm$ BRANCHING RATIOS

NODE=M173220

$\Gamma(D_s^{*+}\pi^0)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE EVTS DOCUMENT ID TECN COMMENT

0.48±0.11 OUR FIT

0.56±0.13±0.09 ¹¹ AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^-\bar{D}^*$

••• We do not use the following data for averages, fits, limits, etc. •••

seen 41 BESSON 03 CLE2 $10.6 e^+ e^-$

¹¹ Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.

NODE=M173R1
NODE=M173R1

NODE=M173R1;LINKAGE=AU

$\Gamma(D_s^+\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE DOCUMENT ID TECN COMMENT

0.18±0.04 OUR FIT

0.16±0.04±0.03 ¹² AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^-\bar{D}^*$

¹² Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.

NODE=M173R6
NODE=M173R6

NODE=M173R6;LINKAGE=AU

$\Gamma(D_s^+\gamma)/\Gamma(D_s^{*+}\pi^0)$ Γ_2/Γ_1

VALUE CL% EVTS DOCUMENT ID TECN COMMENT

0.38 ±0.05 OUR FIT

0.44 ±0.09 OUR AVERAGE

0.55 ±0.13 ±0.08 152 MIKAMI 04 BELL $10.6 e^+ e^-$

0.38 ±0.11 ±0.04 38 KROKOVNY 03B BELL $10.6 e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

0.274±0.045±0.020 251 ¹³ AUBERT,B 04S BABR $B \rightarrow D_{s1}(2460)^+\bar{D}^*$

< 0.49 90 BESSON 03 CLE2 $10.6 e^+ e^-$

¹³ Used by AUBERT 06N in their measurement of $B(D_s^{*-}\pi^0)$ and $B(D_s^-\gamma)$.

NODE=M173R2
NODE=M173R2

NODE=M173R2;LINKAGE=AU

$\Gamma(D_s^+\pi^+\pi^-)/\Gamma(D_s^{*+}\pi^0)$ Γ_3/Γ_1

VALUE CL% EVTS DOCUMENT ID TECN COMMENT

0.090±0.020 OUR FIT Error includes scale factor of 1.2.

0.14 ±0.04 ±0.02 60 MIKAMI 04 BELL $10.6 e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.08 90 BESSON 03 CLE2 $10.6 e^+ e^-$

NODE=M173R3
NODE=M173R3

$\Gamma(D_s^{*+}\gamma)/\Gamma(D_s^{*+}\pi^0)$ Γ_4/Γ_1

VALUE CL% DOCUMENT ID TECN COMMENT

<0.16 90 BESSON 03 CLE2 $10.6 e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.31 90 MIKAMI 04 BELL $10.6 e^+ e^-$

NODE=M173R4
NODE=M173R4

$\Gamma(D_{s0}^*(2317)^+\gamma)/\Gamma(D_s^{*+}\pi^0)$					Γ_5/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.22	95	AUBERT	04E	BABR	10.6 e ⁺ e ⁻	NODE=M173R5 NODE=M173R5
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.58	90	BESSION	03	CLE2	10.6 e ⁺ e ⁻	
$\Gamma(D_s^{*+}\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_1/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
0.93±0.09 OUR FIT						
0.97±0.09±0.05		AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R7 NODE=M173R7
$\Gamma(D_s^+\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_2/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
0.35 ±0.04 OUR FIT						
0.337±0.036±0.038		AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R8 NODE=M173R8
$\Gamma(D_s^+\pi^+\pi^-)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_3/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
0.083±0.017 OUR FIT Error includes scale factor of 1.2.						
0.077±0.013±0.008		AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R9 NODE=M173R9
$\Gamma(D_s^{*+}\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_4/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.24	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R10 NODE=M173R10
$\Gamma(D_{s0}^*(2317)^+\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_5/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.25	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R11 NODE=M173R11
$\Gamma(D_s^+\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_6/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.042	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R12 NODE=M173R12
$\Gamma(D_s^+\pi^0\pi^0)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_7/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.68	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R13 NODE=M173R13
$\Gamma(D_s^+\gamma\gamma)/[\Gamma(D_s^{*+}\pi^0) + \Gamma(D_{s0}^*(2317)^+\gamma)]$					$\Gamma_8/(\Gamma_1+\Gamma_5)$	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.33	95	AUBERT	06P	BABR	10.6 e ⁺ e ⁻	NODE=M173R14 NODE=M173R14

$D_{s1}(2460)^\pm$ REFERENCES

					NODE=M173
AUBERT	06N	PR D74 031103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51142
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
AUBERT	04E	PR D69 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49747
AUBERT,B	04S	PRL 93 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50195
MIKAMI	04	PRL 92 012002	Y. Mikami <i>et al.</i>	(BELLE Collab.)	REFID=49629
BESSION	03	PR D68 032002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=49583
KROKOVNY	03B	PRL 91 262002	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=49615

$D_{s1}(2536)^\pm$ $I(J^P) = 0(1^+)$
 J, P need confirmation.Seen in $D^*(2010)^+ K^0$, $D^*(2007)^0 K^+$, and $D_s^+ \pi^+ \pi^-$. Not seen in $D^+ K^0$ or $D^0 K^+$. $J^P = 1^+$ assignment strongly favored.

NODE=M121

NODE=M121

 $D_{s1}(2536)^\pm$ MASS

NODE=M121M

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121M

NODE=M121M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2535.11 ± 0.06 OUR FIT				
2535.21 ± 0.28 OUR AVERAGE				
2537.7 ± 0.5 ± 3.1	24	¹ ABLIKIM	19P BES3	4.6 $e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
2535.7 ± 0.6 ± 0.5	46	² ABAZOV	09G D0	$B_s^0 \rightarrow D_{s1}^- \mu^+ \nu_\mu X$
2534.78 ± 0.31 ± 0.40	182	AUBERT	08B BABR	$B \rightarrow \bar{D}^{(*)} D^* K$
2534.6 ± 0.3 ± 0.7	193	AUBERT	06P BABR	10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$
2535.3 ± 0.7	92	³ HEISTER	02B ALEP	$e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2534.2 ± 1.2	9	ASRATYAN	94 BEBC	$\nu N \rightarrow D^* K^0 X, D^{*0} K^\pm X$
2535 ± 0.6 ± 1	75	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2535.2 ± 0.5 ± 1.5	28	ALBRECHT	92R ARG	10.4 $e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2536.6 ± 0.7 ± 0.4		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.9 ± 0.6 ± 2.0		ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2534.1 ± 0.6	116	⁴ AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2535.08 ± 0.01 ± 0.15	8038	⁵ LEES	11B BABR	10.6 $e^+ e^- \rightarrow D^{*+} K_S^0 X$
2535.57 ^{+0.44} _{-0.41} ± 0.10	236	⁶ CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
2535.3 ± 0.2 ± 0.5	134	⁷ ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*0} K^+ X$
2534.8 ± 0.6 ± 0.6	44	⁸ ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535 ± 28		⁹ ASRATYAN	88 HLBC	$\nu N \rightarrow D_s \gamma \gamma X$

OCCUR=2

NODE=M121M;LINKAGE=D

NODE=M121M;LINKAGE=AB

NODE=M121M;LINKAGE=HI

NODE=M121M;LINKAGE=AU

NODE=M121M;LINKAGE=LE

NODE=M121M;LINKAGE=CH

NODE=M121M;LINKAGE=A

NODE=M121M;LINKAGE=C

NODE=M121M;LINKAGE=B

 $m_{D_{s1}(2536)^\pm} - m_{D_s^*(2111)}$

NODE=M121DM

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DM

NODE=M121DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
422.9 ± 0.4 OUR FIT			
424 ± 28	ASRATYAN	88 HLBC	$D_s^{*\pm} \gamma$

 $m_{D_{s1}(2536)^\pm} - m_{D^*(2010)^\pm}$

NODE=M121DN

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DN

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
524.85±0.04 OUR FIT				
524.84±0.04 OUR AVERAGE				
524.83±0.01±0.04	8038	¹⁰ LEES	11B BABR	10.6 $e^+e^- \rightarrow D^{*+} K_S^0 X$
525.30 ^{+0.44} _{-0.41} ±0.10	236 ± 30	CHEKANOV	09 ZEUS	$e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
525.3 ±0.6 ±0.1	41	HEISTER	02B ALEP	$e^+e^- \rightarrow D^{*+} K^0 X$
524.7 ±0.6 ±0.2	44	ALEXANDER93	CLE2	$e^+e^- \rightarrow D^{*+} K_S^0 X$
¹⁰ Assuming S-wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$, using a Breit-Wigner line shape corresponding to L=0.				

NODE=M121DN

OCCUR=2

NODE=M121DN;LINKAGE=LE

$m_{D_{s1}(2536)^\pm} - m_{D^*(2007)^0}$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}, D_s^{*\pm}, D_1(2420)^0, D_2^*(2460)^0$,
and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

NODE=M121DP

NODE=M121DP

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
528.26±0.05 OUR FIT				
Error includes scale factor of 1.1.				
528.68±0.28 OUR AVERAGE				
528.7 ±1.9 ±0.5	51	HEISTER	02B ALEP	$e^+e^- \rightarrow D^{*0} K^+ X$
527.3 ±2.2	29	ACKERSTAFF	97W OPAL	$e^+e^- \rightarrow D^{*0} K^+ X$
528.7 ±0.2 ±0.2	134	ALEXANDER	93 CLE2	$e^+e^- \rightarrow D^{*0} K^+ X$

NODE=M121DP

OCCUR=2

$D_{s1}(2536)^\pm$ WIDTH

NODE=M121W

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.92±0.05 OUR AVERAGE				
1.7 ±1.2 ±0.6	24	¹¹ ABLIKIM	19P BES3	4.6 $e^+e^- \rightarrow D_s^+ \bar{D}^0 K^-$
0.92±0.03±0.04	8038	¹² LEES	11B BABR	10.6 $e^+e^- \rightarrow D^{*+} K_S^0 X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.75±0.23	116	¹³ AUSHEV	11 BELL	$B \rightarrow D_{s1}(2536)^+ D^{(*)}$
< 2.5	95 193	AUBERT	06P BABR	10.6 $e^+e^- \rightarrow$ $D_s^+ \pi^+ \pi^- X$
< 3.2	90 75	FRABETTI	94B E687	$\gamma Be \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
< 2.3	90	ALEXANDER	93 CLEO	$e^+e^- \rightarrow D^{*0} K^+ X$
< 3.9	90	ALBRECHT	92R ARG	10.4 $e^+e^- \rightarrow D^{*0} K^+ X$
< 5.44	90	AVERY	90 CLEO	$e^+e^- \rightarrow D^{*+} K^0 X$
< 4.6	90	ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$

NODE=M121W

¹¹ From a fit of the D_s^+ recoil mass distribution with an incoherent sum of the S-wave and S-wave Breit-Wigner line shapes.

¹² Assuming S-wave decay of the $D_{s1}(2536)$ to $D^{*+} K_S^0$, using a Breit-Wigner line shape corresponding to L=0.

¹³ Systematic uncertainties not evaluated.

NODE=M121W;LINKAGE=A

NODE=M121W;LINKAGE=LE

NODE=M121W;LINKAGE=AU

$D_{s1}(2536)^+$ DECAY MODES

NODE=M121215;NODE=M121

Branching fractions are given relative to the one **DEFINED AS 1**.
 $D_{s1}(2536)^-$ modes are charge conjugates of the modes below.

NODE=M121

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $D^*(2010)^+ K^0$	0.85 ±0.12	DESIG=1
Γ_2 $(D^*(2010)^+ K^0)_{S-wave}$	0.61 ±0.09	DESIG=7
Γ_3 $(D^*(2010)^+ K^0)_{D-wave}$		DESIG=9
Γ_4 $K_S^0 D^*(2010)^+$	0.48 ±0.07	DESIG=10
Γ_5 $D^+ \pi^- K^+$	0.028±0.005	DESIG=8
Γ_6 $D^*(2007)^0 K^+$	DEFINED AS 1	DESIG=4
Γ_7 $D^+ K^0$	<0.34	90%
Γ_8 $D^0 K^+$	<0.12	90%
Γ_9 $D_s^{*+} \gamma$	possibly seen	DESIG=3
Γ_{10} $D_s^+ \pi^+ \pi^-$	seen	DESIG=6

$D_{s1}(2536)^+$ BRANCHING RATIOS

NODE=M121220

$$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^*(2010)^+ K^0)$$

 Γ_6/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.18±0.16 OUR AVERAGE				
0.88±0.24±0.08	116	AUSHEV	11	BELL $B \rightarrow D_{s1}(2536)^+ D^{(*)}$
2.3 ±0.6 ±0.3	236 ± 30	CHEKANOV	09	ZEUS $e^\pm p \rightarrow D^{*+} K_S^0 X,$ $D^{*0} K^+ X$
1.32±0.47±0.23	92	¹⁴ HEISTER	02B	ALEP $e^+ e^- \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
1.9 $^{+1.1}_{-0.9}$ ±0.4	35	¹⁴ ACKERSTAFF	97W	OPAL $e^+ e^- \rightarrow D^{*0} K^+ X,$ $D^{*+} K^0 X$
1.1 ±0.3		ALEXANDER	93	CLEO $e^+ e^- \rightarrow$ $D^{*0} K^+ X, D^{*+} K^0 X$
1.4 ±0.3 ±0.2		¹⁵ ALBRECHT	92R	ARG 10.4 $e^+ e^- \rightarrow$ $D^{*0} K^+ X, D^{*+} K^0 X$

NODE=M121R6
NODE=M121R6¹⁴ Ratio of the production rates measured in Z^0 decays.¹⁵ Evaluated by us from published inclusive cross-sections.NODE=M121R6;LINKAGE=6A
NODE=M121R6;LINKAGE=A

$$\Gamma(K_S^0 D^*(2010)^+)/\Gamma(D^*(2007)^0 K^+)$$

 Γ_4/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.48±0.07±0.02	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

NODE=M121R00
NODE=M121R00

$$\Gamma((D^*(2010)^+ K^0)_{S-wave})/\Gamma(D^*(2010)^+ K^0)$$

 Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.72±0.05±0.01	5485	BALAGURA	08	BELL 10.6 $e^+ e^- \rightarrow D^{*+} K^0 X$

NODE=M121R8
NODE=M121R8

$$\Gamma(D^+ \pi^- K^+)/\Gamma(D^*(2010)^+ K^0)$$

 Γ_5/Γ_1

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
3.27±0.18±0.37	1264	BALAGURA	08	BELL 10.6 $e^+ e^- \rightarrow D^+ \pi^- K^+ X$

NODE=M121R9
NODE=M121R9

$$\Gamma(D^+ K^0)/\Gamma(D^*(2010)^+ K^0)$$

 Γ_7/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.40	90	ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*+} K^0 X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.43	90	ALBRECHT	89E	ARG $D_{s1}^* \rightarrow D^*(2010) K^0$

NODE=M121R1
NODE=M121R1

$$\Gamma(D^0 K^+)/\Gamma(D^*(2007)^0 K^+)$$

 Γ_8/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.12	90	ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121R4
NODE=M121R4

$$\Gamma(D_s^{*+} \gamma)/\Gamma_{total}$$

 Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	ASRATYAN	88	HLBC $\nu N \rightarrow D_s \gamma \gamma X$

NODE=M121R3
NODE=M121R3

$$\Gamma(D_s^{*+} \gamma)/\Gamma(D^*(2007)^0 K^+)$$

 Γ_9/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42	90	ALEXANDER	93	CLEO $e^+ e^- \rightarrow D^{*0} K^+ X$

NODE=M121R5
NODE=M121R5

$$\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma_{total}$$

 Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	06P	BABR 10.6 $e^+ e^- \rightarrow D_s^+ \pi^+ \pi^- X$

NODE=M121R7
NODE=M121R7 $D_{s1}(2536)^\pm$ REFERENCES

NODE=M121

GAO	23	PR D108 112015	B.S. Gao <i>et al.</i>	(BELLE Collab.)	REFID=62519
ABLIKIM	19P	CP C43 031001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59617
AUSHEV	11	PR D83 051102	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=16505
LEES	11B	PR D83 072003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16773
ABAZOV	09G	PRL 102 051801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52652
CHEKANOV	09	EPJ C60 25	S. Chekanov <i>et al.</i>	(ZEUS Collab.)	REFID=52733
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BALAGURA	08	PR D77 032001	V. Balagura <i>et al.</i>	(BELLE Collab.)	REFID=52133
AUBERT	06P	PR D74 032007	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51144
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ASRATYAN	94	ZPHY C61 563	A.E. Asratyan <i>et al.</i>	(BIRM, BELG, CERN+)	REFID=43667
FRABETTI	94B	PRL 72 324	P.L. Frabetti <i>et al.</i>	(FNAL E687 Collab.)	REFID=43687
ALEXANDER	93	PL B303 377	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=43316
ALBRECHT	92R	PL B297 425	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=43179
AVERY	90	PR D41 774	P. Avery, D. Besson	(CLEO Collab.)	REFID=41013
ALBRECHT	89E	PL B230 162	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40914
ASRATYAN	88	ZPHY C40 483	A.E. Asratyan <i>et al.</i>	(ITEP, SERP)	REFID=40916

$D_{s2}^*(2573)$

$$I(J^P) = 0(2^+)$$

NODE=M148

 $D_{s2}^*(2573)$ MASS

NODE=M148M

NODE=M148M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2569.1 ±0.8 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.		
2570.7 ±2.0 ±1.7	62	¹ ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
2568.39 ±0.29 ±0.26		AAIJ	14AW LHCb	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
2569.4 ±1.6 ±0.5	82	AAIJ	11A LHCb	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$
2572.2 ±0.3 ±1.0		AUBERT,BE	06E BABR	$e^+ e^- \rightarrow D K X$
2574.5 ±3.3 ±1.6		ALBRECHT	96 ARG	$e^+ e^- \rightarrow D^0 K^+ X$
2573.2 $^{+1.7}_{-1.6}$ ±0.9	217	KUBOTA	94 CLE2	$e^+ e^- \sim 10.5$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2570.0 ±4.3	25	² EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$
2568.6 ±3.2	64	³ HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$

¹ From a fit of the D_s^+ recoil mass distribution .² Not independent of the mass difference below.³ Calculated using $m_{D^0} = 1864.5 \pm 0.5$ MeV and the mass difference below.

NODE=M148M;LINKAGE=A

NODE=M148M;LINKAGE=EV

NODE=M148M;LINKAGE=HI

 $m_{D_{s2}^*(2573)} - m_{D^0}$

NODE=M148DM

NODE=M148DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
704 ±3 ±1	64	HEISTER	02B ALEP	$e^+ e^- \rightarrow D^0 K^+ X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
705.4 ±4.3	25	¹ EVDOKIMOV	04 SELX	$600 \Sigma^- A \rightarrow D^0 K^+ X$
¹ Systematic errors not estimated.				

NODE=M148DM;LINKAGE=EV

 $D_{s2}^*(2573)$ WIDTH

NODE=M148W

NODE=M148W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.9 ±0.7 OUR AVERAGE				
17.2 ±3.6 ±1.1	62	¹ ABLIKIM	19P BES3	$4.6 e^+ e^- \rightarrow D_s^+ \bar{D}^0 K^-$
16.9 ±0.5 ±0.6		AAIJ	14AW LHCb	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
12.1 ±4.5 ±1.6	82	AAIJ	11A LHCb	$B_s \rightarrow D_{s2}^*(2573) \mu \bar{\nu} X$

27.1±0.6±5.6	AUBERT,BE	06E	BABR	$e^+e^- \rightarrow DKX$
10.4±8.3±3.0	ALBRECHT	96	ARG	$e^+e^- \rightarrow D^0K^+X$
16 $\begin{smallmatrix} +5 \\ -4 \end{smallmatrix}$ ±3	217	KUBOTA	94	CLE2 $e^+e^- \sim 10.5$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

14 $\begin{smallmatrix} +9 \\ -6 \end{smallmatrix}$	25	² EVDOKIMOV	04	SELX	600 $\Sigma^- A \rightarrow D^0K^+X$
---	----	------------------------	----	------	--------------------------------------

¹ From a fit of the D_s^+ recoil mass distribution .

² Systematic errors not estimated.

NODE=M148W;LINKAGE=A
NODE=M148W;LINKAGE=EV

$D_{s2}^*(2573)^+$ DECAY MODES

$D_{s2}^*(2573)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 D^0K^+	seen
Γ_2 $D^*(2007)^0K^+$	not seen
Γ_3 $D^+K_S^0$	seen
Γ_4 $D^{*+}K_S^0$	seen

NODE=M148215;NODE=M148

NODE=M148

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=4;OUR EVAL;→ UNCHECKED ←

DESIG=5;OUR EVAL;→ UNCHECKED ←

$D_{s2}^*(2573)^+$ BRANCHING RATIOS

NODE=M148220

$\Gamma(D^0K^+)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen	217	KUBOTA	94	CLE2	\pm $e^+e^- \sim 10.5$ GeV	

NODE=M148R2
NODE=M148R2

$\Gamma(D^*(2007)^0K^+)/\Gamma(D^0K^+)$						Γ_2/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.33	90	KUBOTA	94	CLE2	$+$ $e^+e^- \sim 10.5$ GeV	

NODE=M148R1
NODE=M148R1

$\Gamma(D^+K_S^0)/\Gamma(D^0K^+)$						Γ_3/Γ_1
VALUE	DOCUMENT ID	TECN	COMMENT			
0.49±0.10±0.02	GAO	23	BELL	e^+e^- at 10.52 GeV		

NODE=M148R01
NODE=M148R01

$\Gamma(D^{*+}K_S^0)/\Gamma(D^+K_S^0)$						Γ_4/Γ_3
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
0.044±0.005±0.011	2000	¹ AAIJ	16AW	LHCB	$pp \rightarrow D^{*+}K_S^0X$ at 7, 8 TeV	

NODE=M148R00
NODE=M148R00

¹ First observation of the $D_{s2}^*(2573)^+ \rightarrow D^{*+}K_S^0$ decay with a significance of 6.9 σ .

NODE=M148R00;LINKAGE=A

$D_{s2}^*(2573)$ REFERENCES

NODE=M148

GAO	23	PR D108 112015	B.S. Gao <i>et al.</i>	(BELLE Collab.)	REFID=62519
ABLIKIM	19P	CP C43 031001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59617
AAIJ	16AW	JHEP 1602 133	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60464
AAIJ	14AW	PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=56105
AAIJ	11A	PL B698 14	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=16665
AUBERT,BE	06E	PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51512
EVDOKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)	REFID=50337
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
ALBRECHT	96	ZPHY C69 405	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44631
KUBOTA	94	PRL 72 1972	Y. Kubota <i>et al.</i>	(CLEO Collab.)	REFID=43781

$D_{s0}(2590)^+$

$I(J^P) = 0(0^-)$

OMITTED FROM SUMMARY TABLE

 $D_{s0}(2590)^+$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2591±6±7	444	¹ AAIJ	21A LHCb	$B^0 \rightarrow D^-(D^+ K^+ \pi^-)$

¹ The mass is calculated from the position of the T-matrix pole

NODE=M256M

NODE=M256M

NODE=M256M;LINKAGE=B

 $D_{s0}(2590)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
89±16±12	444	¹ AAIJ	21A LHCb	$B^0 \rightarrow D^-(D^+ K^+ \pi^-)$

¹ The width is calculated from the position of the T-matrix pole

NODE=M256W

NODE=M256W

NODE=M256W;LINKAGE=B

 $D_{s0}(2590)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^+ K^+ \pi^-$	seen

NODE=M256215;NODE=M256

DESIG=1

$\Gamma(D^+ K^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	444

NODE=M256R01
NODE=M256R01 $D_{s0}(2590)^+$ REFERENCESAAIJ 21A PRL 126 122002 R. Aaij *et al.* (LHCb Collab.)

NODE=M256

REFID=61092

NODE=M182

 $D_{s1}^*(2700)^\pm$

$I(J^P) = 0(1^-)$

 $D_{s1}^*(2700)^+$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2714 ± 5 OUR AVERAGE				Error includes scale factor of 1.5. See the ideogram below.
2732.3± 4.3± 5.8	15.7k	AAIJ	16AW LHCb	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
2699 $^{+14}_{-7}$		¹ LEES	15C BABR	$B \rightarrow D D^0 K^+$
2709.2± 1.9± 4.5	52k	² AAIJ	12AU LHCb	$pp \rightarrow (DK)^+ X$ at 7 TeV
2710 ± 2 $^{+12}_{-7}$	10.4k	³ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$
2708 ± 9 $^{+11}_{-10}$	182	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2694 ± 8 $^{+13}_{-3}$		LEES	15C BABR	$B^0 \rightarrow D^- D^0 K^+$
2707 ± 8 ± 8		LEES	15C BABR	$B^+ \rightarrow \bar{D}^0 D^0 K^+$
2688 ± 4 ± 3		⁴ AUBERT, BE	06E BABR	10.6 $e^+ e^- \rightarrow DKX$

NODE=M182M

NODE=M182M

OCCUR=2

OCCUR=3


¹ From a combined analysis of $B^0 \rightarrow D^- D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 D^0 K^+$.² From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{sJ}^*(2860)^+$.³ From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.⁴ Superseded by AUBERT 09AR.

NODE=M182M;LINKAGE=B

NODE=M182M;LINKAGE=AA

NODE=M182M;LINKAGE=AB

NODE=M182M;LINKAGE=AU



$D_{s1}^{*+}(2700)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
122 ±10 OUR AVERAGE				
136 ±19 ±24	15.7k	AAIJ	16AW LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
127 ⁺²⁴ / ₋₁₉		¹ LEES	15C BABR	$B \rightarrow D D^0 K^+$
115.8 ± 7.3 ±12.1	52k	² AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
149 ± 7 ⁺³⁹ / ₋₅₂	10.4k	³ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$
108 ±23 ⁺³⁶ / ₋₃₁	182	BRODZICKA	08 BELL	$B^+ \rightarrow D^0 \bar{D}^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

145 ±24 ⁺²² / ₋₁₄		LEES	15C BABR	$B^0 \rightarrow D^- D^0 K^+$
113 ±21 ⁺²⁰ / ₋₁₆		LEES	15C BABR	$B^+ \rightarrow \bar{D}^0 D^0 K^+$
112 ± 7 ±36		⁴ AUBERT,BE	06E BABR	10.6 $e^+ e^- \rightarrow D K X$

¹ From a combined analysis of $B^0 \rightarrow D^- D^0 K^+$ and $B^+ \rightarrow \bar{D}^0 D^0 K^+$.

² From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^{*+}(2573)^+$, $D_{s1}^{*+}(2700)^+$ and spin-0 $D_{sJ}^{*+}(2860)^+$.

³ From simultaneous fits to the two DK mass spectra and to the total $D^* K$ mass spectrum.

⁴ Superseded by AUBERT 09AR.

$D_{s1}^{*+}(2700)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 DK	
Γ_2 $D^0 K^+$	seen
Γ_3 $D^+ K_S^0$	seen
Γ_4 $D^* K$	
Γ_5 $D^{*0} K^+$	seen
Γ_6 $D^{*+} K_S^0$	seen

$D_{s1}^{*+}(2700)^+$ BRANCHING RATIOS

$\Gamma(D^* K)/\Gamma(DK)$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
0.91±0.13±0.12	10.4k	¹ AUBERT	09AR BABR	$e^+ e^- \rightarrow D^{(*)} K X$	

¹ From the average of the corresponding ratios with $D^{(*)0} K^+$ and $D^{(*)+} K_S^0$.

NODE=M182W

NODE=M182W

OCCUR=2

OCCUR=3

NODE=M182W;LINKAGE=A

NODE=M182W;LINKAGE=AA

NODE=M182W;LINKAGE=AB

NODE=M182W;LINKAGE=AU

NODE=M182215;NODE=M182

DESIG=2

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

NODE=M182225

NODE=M182R01

NODE=M182R01

NODE=M182R01;LINKAGE=AU

$\Gamma(D^{*0}K^+)/\Gamma(D^0K^+)$

VALUE	EVT5	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
0.88±0.14±0.14	7716	¹ AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$	

¹ From the $D^{*0}K^+$ and D^0K^+ , where $D^{*0} \rightarrow D^0\pi^0$.

NODE=M182R02
NODE=M182R02

NODE=M182R02;LINKAGE=AU

 $\Gamma(D^{*+}K_S^0)/\Gamma(D^+K_S^0)$

VALUE	EVT5	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
1.14±0.39±0.23	2700	¹ AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$	

¹ From the $D^{*+}K_S^0$ and $D^+K_S^0$, where $D^{*+} \rightarrow D^+\pi^0$.

NODE=M182R03
NODE=M182R03

NODE=M182R03;LINKAGE=AU

 $D_{s1}^*(2700)^\pm$ REFERENCES

AAIJ	16AW JHEP 1602 133	R. Aaij <i>et al.</i>	(LHCb Collab.)
LEES	15C PR D91 052002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AAIJ	12AU JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
BRODZICKA	08 PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)
AUBERT,BE	06E PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M182

REFID=60464
REFID=56412
REFID=54735
REFID=53135
REFID=52144
REFID=51512

NODE=M196

$D_{s1}^*(2860)^\pm$

$$I(J^P) = 0(1^-)$$

OMITTED FROM SUMMARY TABLE

was $D_{s,J}^*(2860)$

J^P consistent with 1^- from angular analysis of AAIJ 14AW.

NODE=M196

 $D_{s1}^*(2860)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2859±12±24	¹ AAIJ	14AW LHCB	$B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$

¹ Separated from the spin-3 component $D_{s3}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M196M

NODE=M196M

NODE=M196M;LINKAGE=A

 $D_{s1}^*(2860)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
160±80 OUR AVERAGE	[159 ± 80 MeV OUR 2023 AVERAGE]		
159±23±77	¹ AAIJ	14AW LHCB	$B_S^0 \rightarrow \bar{D}^0 K^- \pi^+$

¹ Separated from the spin-3 component $D_{s3}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M196W

NODE=M196W

NEW

NODE=M196W;LINKAGE=A

 $D_{s1}^*(2860)^\pm$ REFERENCES

AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
------	---------------------	-----------------------	-------------------

NODE=M196

REFID=56105

NODE=M226

 $D_{s3}^*(2860)^\pm$

$$I(J^P) = 0(3^-)$$

J^P consistent with 3^- from angular analysis of AAIJ 14AW. Observed by AUBERT, BE 06E and AUBERT 09AR in inclusive production of DK and D^*K in e^+e^- annihilation.

NODE=M226

 $D_{s3}^*(2860)^+$ MASS

NODE=M226M

NODE=M226M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2860.5±2.6±6.5		¹ AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2867.1±4.3±1.9	3.1k	AAIJ	16AW LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
2866.1±1.0±6.3	36k	^{2,3} AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
2862 ±2 $\begin{smallmatrix} +5 \\ -2 \end{smallmatrix}$	3122	^{2,4} AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)} K X$
2856.6±1.5±5.0		⁵ AUBERT, BE	06E BABR	$e^+e^- \rightarrow DK X$

¹ Separated from the spin-1 component $D_{s1}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M226M;LINKAGE=A

² Possible contribution from the $D_{s1}^*(2860)$ state.

NODE=M226M;LINKAGE=D

³ From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{s,J}^*(2860)^+$.

NODE=M226M;LINKAGE=E

⁴ From simultaneous fits to the two DK mass spectra and to the total D^*K mass spectrum.

NODE=M226M;LINKAGE=C

⁵ Superseded by AUBERT 09AR.

NODE=M226M;LINKAGE=B

 $D_{s3}^*(2860)^+$ WIDTH

NODE=M226W

NODE=M226W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
53 ± 7 ± 7		¹ AAIJ	14AW LHCB	$B_s^0 \rightarrow \bar{D}^0 K^- \pi^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
50 ±11 ±13	3.1k	AAIJ	16AW LHCB	$pp \rightarrow D^{*+} K_S^0 X$ at 7, 8 TeV
69.9± 3.2± 6.6	36k	^{2,3} AAIJ	12AU LHCB	$pp \rightarrow (DK)^+ X$ at 7 TeV
48 ± 3 ± 6	3122	^{2,4} AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)} K X$
47 ± 7 ±10		⁵ AUBERT, BE	06E BABR	$e^+e^- \rightarrow DK X$

¹ Separated from the spin-1 component $D_{s1}^*(2860)^-$ by a fit of the helicity angle of the $\bar{D}^0 K^-$ system, with a statistical significance of the spin-3 and spin-1 components in excess of 10σ .

NODE=M226W;LINKAGE=A

² Possible contribution from the $D_{s1}^*(2860)$ state.

NODE=M226W;LINKAGE=D

³ From the combined fit of the $D^+ K_S^0$ and $D^0 K^+$ modes in the model including the $D_{s2}^*(2573)^+$, $D_{s1}^*(2700)^+$ and spin-0 $D_{s,J}^*(2860)^+$.

NODE=M226W;LINKAGE=E

⁴ From simultaneous fits to the two DK mass spectra and to the total D^*K mass spectrum.

NODE=M226W;LINKAGE=C

⁵ Superseded by AUBERT 09AR.

NODE=M226W;LINKAGE=B

 $D_{s3}^*(2860)^\pm$ DECAY MODES

NODE=M226215;NODE=M226

Mode	Fraction (Γ_i/Γ)
Γ_1 DK	
Γ_2 $D^0 K^+$	seen
Γ_3 $D^+ K_S^0$	seen
Γ_4 $D^* K$	
Γ_5 $D^{*0} K^+$	seen
Γ_6 $D^{*+} K_S^0$	seen

DESIG=1

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

DESIG=4

DESIG=5;OUR EVAL;→ UNCHECKED ←

DESIG=6;OUR EVAL;→ UNCHECKED ←

$D_{s3}^*(2860)^\pm$ BRANCHING RATIOS $\Gamma(D^*K)/\Gamma(DK)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.10 \pm 0.15 \pm 0.19$	3122	¹ AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$

¹ From the average of the corresponding ratios with $D^{(*)0}K^+$ and $D^{(*)+}K_S^0$.

NODE=M226225

NODE=M226R01
NODE=M226R01

NODE=M226R01;LINKAGE=AU

 $\Gamma(D^{*0}K^+)/\Gamma(D^0K^+)$ Γ_5/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.04 \pm 0.17 \pm 0.20$	2241	¹ AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$

¹ From the $D^{*0}K^+$ and D^0K^+ , where $D^{*0} \rightarrow D^0\pi^0$.

NODE=M226R02
NODE=M226R02

NODE=M226R02;LINKAGE=AU

 $\Gamma(D^{*+}K_S^0)/\Gamma(D^+K_S^0)$ Γ_6/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.38 \pm 0.35 \pm 0.49$	881	¹ AUBERT	09AR BABR	$e^+e^- \rightarrow D^{(*)}KX$

¹ From the $D^{*+}K_S^0$ and $D^+K_S^0$, where $D^{*+} \rightarrow D^+\pi^0$.

NODE=M226R03
NODE=M226R03

NODE=M226R03;LINKAGE=AU

 $D_{s3}^*(2860)^\pm$ REFERENCES

AAIJ	16AW JHEP 1602 133	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14AW PRL 113 162001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
AAIJ	12AU JHEP 1210 151	R. Aaij <i>et al.</i>	(LHCb Collab.)
AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
AUBERT,BE	06E PRL 97 222001	B. Aubert <i>et al.</i>	(BABAR Collab.)

NODE=M226

REFID=60464
REFID=56105
REFID=54735
REFID=53135
REFID=51512

NODE=M197

 $D_{sJ}(3040)^\pm$

$$I(J^P) = 0(?^?)$$

OMITTED FROM SUMMARY TABLE

Observed by AUBERT 09AR in inclusive production of D^*K in e^+e^- annihilation.

NODE=M197

 $D_{sJ}(3040)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$3044 \pm 8 \pm 30 \pm 5$	AUBERT	09AR BABR	$e^+e^- \rightarrow D^*KX$

NODE=M197M

NODE=M197M

 $D_{sJ}(3040)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
240 ± 60 OUR AVERAGE	[239 \pm 60 MeV OUR 2023 AVERAGE]		
$239 \pm 35 \pm 46 \pm 42$	AUBERT	09AR BABR	$e^+e^- \rightarrow D^*KX$

NODE=M197W

NODE=M197W

NEW

 $D_{sJ}(3040)^\pm$ DECAY MODES

Mode			
Γ_1	D^*K		DESIG=1
Γ_2	$D^{*0}K^+$		DESIG=2
Γ_3	$D^{*+}K_S^0$		DESIG=3

NODE=M197215;NODE=M197

 $D_{sJ}(3040)^\pm$ REFERENCES

AUBERT	09AR PR D80 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)
--------	--------------------	-------------------------	-----------------

NODE=M197

REFID=53135

BOTTOM MESONS

($B = \pm 1$)

$B^+ = u\bar{b}$, $B^0 = d\bar{b}$, $\bar{B}^0 = \bar{d}b$, $B^- = \bar{u}b$, similarly for B^{*} 's

$B_1(5721)$

$I(J^P) = \frac{1}{2}(1^+)$
I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

B₁(5721) MASS

B₁(5721)⁺ mass

OUR FIT uses $m_{B^{*0}}$ and $m_{B_1^+} - m_{B^{*0}}$ to determine $m_{B_1(5721)^+}$.

VALUE (MeV) _____ DOCUMENT ID _____

5726.0^{+2.5}_{-2.7} OUR FIT

[5725.9^{+2.5}_{-2.7} MeV OUR 2023 FIT]

$m_{B_1^+} - m_{B^{*0}}$

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____

401.2^{+2.4}_{-2.7} OUR FIT

401.2^{+2.4}_{-2.7} OUR AVERAGE

400.5 ± 1.8 ± 3.1 8k ¹ AAIJ 15AB LHCB $p\bar{p}$ at 7, 8 TeV

402 ± 3 ⁺¹₋₃ ² AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_1^+} - m_{B^0}] - (m_{B^{*0}} - m_{B^0}) - m_{\pi^+} = 260.9 \pm 1.8 \pm 3.1$

MeV which we adjust by the π^+ mass and assume $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_1(5721)^+} - m_{B^{*0}} - m_{\pi^+} = 262 \pm 3^{+1}_{-3}$ MeV which we adjusted by the π^+ mass.

B₁(5721)⁰ mass

OUR FIT uses mass differences measurements listed below to determine the mass

$m_{B_1(5721)^0}$.

VALUE (MeV) _____ DOCUMENT ID _____

5726.1 ± 1.2 OUR FIT Error includes scale factor of 1.2. [5726.1 ± 1.3 MeV OUR 2023 FIT Scale factor = 1.2]

$m_{B_1^0} - m_{B^+}$

VALUE (MeV) _____ DOCUMENT ID _____ TECN _____ COMMENT _____

446.7 ± 1.2 OUR FIT Error includes scale factor of 1.2. [446.7 ± 1.3 MeV OUR 2023 FIT Scale factor = 1.2]

441.5 ± 2.4 ± 1.3 ¹ ABAZOV 07T D0 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

446.2^{+1.9+1.0}_{-2.1-1.2} ¹ AALTONEN 09D CDF Repl. by AALTONEN 14I

¹ Observed in $B_1^0 \rightarrow B^{*+} \pi^-$.

$m_{B_1^0} - m_{B^{*+}}$

VALUE (MeV) _____ EVTS _____ DOCUMENT ID _____ TECN _____ COMMENT _____

401.4 ± 1.2 OUR FIT Error includes scale factor of 1.2.

402.8 ± 1.1 OUR AVERAGE

403.4 ± 0.7 ± 1.5 35k ¹ AAIJ 15AB LHCB $p\bar{p}$ at 7, 8 TeV

402.3 ± 0.9^{+1.1}_{-1.2} ² AALTONEN 14I CDF $p\bar{p}$ at 1.96 TeV

NODE=MXXX045

NODE=MXXX045

NODE=M244

NODE=M244

NODE=M244205

NODE=M244M+

NODE=M244M+
NODE=M244M+

NEW

NODE=M244DM+
NODE=M244DM+

NODE=M244DM+;LINKAGE=A

NODE=M244DM+;LINKAGE=AA

NODE=M244M0

NODE=M244M0

NODE=M244M0

NEW

NODE=M244DM0

NODE=M244DM0

NEW

NODE=M244DM0;LINKAGE=AA

NODE=M244DM1

NODE=M244DM1

¹ AAIJ 15AB reports $[m_{B_1^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 263.9 \pm 0.7 \pm 1.4$

MeV which we adjust by the π^- mass and $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_1(5721)^0} - m_{B^{*+}} - m_{\pi^-} = 262.7 \pm 0.9^{+1.1}_{-1.2}$ MeV which we adjusted by the π^- mass.

NODE=M244DM1;LINKAGE=B

NODE=M244DM1;LINKAGE=AA

$B_1(5721)$ WIDTH

NODE=M244210

$B_1(5721)^+$ width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31 ± 6 OUR AVERAGE				Error includes scale factor of 1.1.
29.1 ± 3.6 ± 4.3	8k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
49 $^{+12}_{-10}$ $^{+2}_{-13}$		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M244W+
NODE=M244W+

$B_1(5721)^0$ width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.5 ± 3.4 OUR AVERAGE				Error includes scale factor of 1.1.
30.1 ± 1.5 ± 3.5	35k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
23 ± 3 ± 4		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M244W0
NODE=M244W0

$B_1(5721)$ DECAY MODES

NODE=M244215;NODE=M244

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^* \pi$	seen

DESIG=1

$\Gamma(B^* \pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
seen	AAIJ	15AB LHCB	±0	$p\bar{p}$ at 7, 8 TeV	
seen	AALTONEN	14I CDF	±	$p\bar{p}$ at 1.96 TeV	
seen	AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV	
seen	¹ ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV	

NODE=M244R01
NODE=M244R01

OCCUR=2

¹ Observed in $B_1^0 \rightarrow B^{*+} \pi^-$ with $B^{*+} \rightarrow B^+ \gamma$ and $B^+ \rightarrow J/\psi \pi^+$.

NODE=M244R01;LINKAGE=AB

$B_1(5721)$ REFERENCES

NODE=M244

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=56628
REFID=56029
REFID=52700
REFID=52014

$B_J^*(5732)$

$$I(J^P) = ?(??)$$

OMITTED FROM SUMMARY TABLE
also known as B^{**}

Signal can be interpreted as stemming from several narrow and broad resonances.

NODE=M151

NODE=M151

 $B_J^*(5732)$ MASS

NODE=M151M

NODE=M151M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5698 ± 8 OUR AVERAGE	Error includes scale factor of 1.2.			
5710 ± 20		¹ AFFOLDER	01F CDF	$p\bar{p}$ at 1.8 TeV
5695 ⁺¹⁷ ₋₁₉		² BARATE	98L ALEP	$e^+e^- \rightarrow Z$
5704 ± 4 ± 10	1944	³ BUSKULIC	96D ALEP	$E_{cm}^{ee} = 88-94$ GeV
5732 ± 5 ± 20	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88-94$ GeV
5681 ± 11	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5713 ± 2		⁴ ACCIARRI	99N L3	$e^+e^- \rightarrow Z$

NODE=M151M;LINKAGE=MF

¹ AFFOLDER 01F uses the reconstructed B meson through semileptonic decay channels. The fraction of light B mesons that are produced at $L=1$ B^{**} states is measured to be $0.28 \pm 0.06 \pm 0.03$.

² BARATE 98L uses fully reconstructed B mesons to search for B^{**} production in the $B\pi^\pm$ system. In the framework of heavy quark symmetry (HQSS), they also measured the mass of B_2^* to be 5739^{+8+6}_{-11-4} MeV/ c^2 and the relative production rate of $B(b \rightarrow B_2^* \rightarrow B^{(*)}\pi)/B(b \rightarrow B_{u,d}) = (31 \pm 9^{+6}_{-5})\%$.

NODE=M151M;LINKAGE=B

³ Using $m_{B\pi} - m_B = 424 \pm 4 \pm 10$ MeV.

NODE=M151M;LINKAGE=A

⁴ ACCIARRI 99N uses inclusive reconstructed B mesons to search for B^{**} production in the $B^{(*)}\pi^\pm$ system. In the framework of HQET, they measured the mass of B_1^* and B_2^* to be $5670 \pm 10 \pm 13$ MeV and $5768 \pm 5 \pm 6$ with the $B(b \rightarrow B^{**}) = (32 \pm 3 \pm 6) \times 10^{-2}$. They also reported the evidence for the existence of an excited B -meson state or mixture of states in the region 5.9–6.0 GeV.

NODE=M151M;LINKAGE=N

 $B_J^*(5732)$ WIDTH

NODE=M151W

NODE=M151W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
128 ± 18 OUR AVERAGE				
145 ± 28	2157	ABREU	95B DLPH	$E_{cm}^{ee} = 88-94$ GeV
116 ± 24	1738	AKERS	95E OPAL	$E_{cm}^{ee} = 88-94$ GeV

 $B_J^*(5732)$ DECAY MODES

NODE=M151215;NODE=M151

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^*\pi + B\pi$	seen
Γ_2 $B^*\pi(X)$	[a] (85 ± 29) %

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=2

[a] X refers to decay modes with or without additional accompanying decay particles.

LINKAGE=151

 $B_J^*(5732)$ BRANCHING RATIOS

NODE=M151220

NODE=M151220

X refers to decay modes with or without additional accompanying decay particles.

$\Gamma(B^*\pi(X))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.85^{+0.26}_{-0.27} ± 0.12	ABBIENDI	02E OPAL	$e^+e^- \rightarrow Z$

NODE=M151R1
NODE=M151R1

$B_2^*(5732)$ REFERENCES

ABBIENDI	02E	EPJ C23 437	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AFFOLDER	01F	PR D64 072002	T. Affolder <i>et al.</i>	(CDF Collab.)
ACCIARRI	99N	PL B465 323	M. Acciarri <i>et al.</i>	(L3 Collab.)
BARATE	98L	PL B425 215	R. Barate <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96D	ZPHY C69 393	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABREU	95B	PL B345 598	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	95E	ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)

NODE=M151

REFID=48742
REFID=48369
REFID=47247
REFID=46082
REFID=44677
REFID=44131
REFID=44182

NODE=M245

 $B_2^*(5747)$

$$I(J^P) = \frac{1}{2}(2^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M245

 $B_2^*(5747)$ MASS

NODE=M245205

 $B_2^*(5747)^+$ mass

NODE=M245M+

OUR FIT uses m_{B^0} and $m_{B_2^{*+}} - m_{B^0}$ to determine $m_{B_2^*(5747)^+}$.NODE=M245M+
NODE=M245M+
NEW

VALUE (MeV)	DOCUMENT ID
5737.3±0.7 OUR FIT	
[5737.2 ± 0.7 MeV OUR 2023 FIT]	

 $m_{B_2^{*+}} - m_{B^0}$ NODE=M245DM+
NODE=M245DM+

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
457.5 ± 0.7 OUR FIT				
457.5 ± 0.7 OUR AVERAGE				
457.62±0.72±0.40	4k	¹ AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
457.3 ± 1.3 $\begin{smallmatrix} +0.3 \\ -0.9 \end{smallmatrix}$		² AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_2^{*+}} - m_{B^0}] - m_{\pi^+} = 318.1 \pm 0.7 \pm 0.4$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_2^*(5747)^+} - m_{B^0} - m_{\pi^+} = 317.7 \pm 1.2^{+0.3}_{-0.9}$ MeV which we adjusted by the π^+ mass.

NODE=M245DM+;LINKAGE=B

NODE=M245DM+;LINKAGE=A

 $B_2^*(5747)^0$ mass

NODE=M245M0

OUR FIT uses m_{B^+} , $m_{B_1^0} - m_{B^+}$, and mass differences below to determine $m_{B_2^*(5747)^0}$. The -0.659 correlation between statistical uncertainties of $m_{B_1^0} - m_{B^+}$ and $m_{B_2^{*0}} - m_{B_1^0}$ measurements reported by ABAZOV 07T is taken into account.

NODE=M245M0

VALUE (MeV)	DOCUMENT ID
5739.6±0.7 OUR FIT	Error includes scale factor of 1.4. [5739.5 ± 0.7 MeV OUR 2023 FIT Scale factor = 1.4]

NODE=M245M0
NEW **$m_{B_2^{*0}} - m_{B_1^0}$** NODE=M245DM0
NODE=M245DM0
NEW

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.5±1.4 OUR FIT	Error includes scale factor of 1.3. [13.4 ± 1.4 MeV OUR 2023 FIT Scale factor = 1.3]		

26.2±3.1±0.9 ¹ ABAZOV 07T D0 $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

14.9 $\begin{smallmatrix} +2.2+1.2 \\ -2.5-1.4 \end{smallmatrix}$ ¹ AALTONEN 09D CDF Repl. by AALTONEN 14I

¹ Observed in $B_2^{*0} \rightarrow B^{*+}\pi^-$ and $B_2^{*0} \rightarrow B^+\pi^-$.

NODE=M245DM0;LINKAGE=AB

 $m_{B_2^{*0}} - m_{B^+}$ NODE=M245DM2
NODE=M245DM2

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
460.2 ± 0.6 OUR FIT	Error includes scale factor of 1.4.			
459.9 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.8.			
460.18±0.37±0.33	17k	¹ AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
457.5 ± 1.2 $\begin{smallmatrix} +0.8 \\ -0.9 \end{smallmatrix}$		² AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

¹ AAIJ 15AB reports $[m_{B_2^{*0}} - m_{B^+}] - m_{\pi^-} = 320.6 \pm 0.4 \pm 0.3$ MeV which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event.

² AALTONEN 14I reports $m_{B_2^*(5747)^0} - m_{B^+} - m_{\pi^-} = 317.9 \pm 1.2^{+0.8}_{-0.9}$ MeV which we adjusted by the π^- mass.

NODE=M245DM2;LINKAGE=A

NODE=M245DM2;LINKAGE=AA

$B_2^*(5747)$ WIDTH

NODE=M245210

 $B_2^*(5747)^+$ width

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 ±5 OUR AVERAGE	Error includes scale factor of 2.2.			
23.6±2.0±2.1	4k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
11 ^{+4 +3} _{-3 -4}		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV

NODE=M245W+
NODE=M245W+ **$B_2^*(5747)^0$ width**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
24.2±1.7 OUR AVERAGE				
24.5±1.0± 1.5	17k	AAIJ	15AB LHCB	$p\bar{p}$ at 7, 8 TeV
22 ^{+3 +4} _{-2 -5}		AALTONEN	14I CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22.7 ^{+3.8 + 3.2} _{-3.2 -10.2}		AALTONEN	09D CDF	Repl. by AALTONEN 14I

NODE=M245W0
NODE=M245W0 **$B_2^*(5747)$ DECAY MODES**

NODE=M245215;NODE=M245

Mode	Fraction (Γ_i/Γ)
Γ_1 $B\pi$	seen
Γ_2 $B^*\pi$	seen

DESIG=1

DESIG=2

 $\Gamma(B\pi)/\Gamma_{total}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	4k,17k	AAIJ	15AB LHCB	±0	$p\bar{p}$ at 7, 8 TeV
seen		AALTONEN	14I CDF	±	$p\bar{p}$ at 1.96 TeV
seen		AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV
seen		ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV

NODE=M245R01
NODE=M245R01 **$\Gamma(B^*\pi)/\Gamma_{total}$** Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
seen	4k,17k	AAIJ	15AB LHCB	±0	$p\bar{p}$ at 7, 8 TeV
seen		AALTONEN	09D CDF	0	$p\bar{p}$ at 1.96 TeV
seen		ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV

NODE=M245R02
NODE=M245R02 **$\Gamma(B^*\pi)/\Gamma(B\pi)$** Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.84±0.27 OUR AVERAGE					
0.71±0.14±0.30	17k	AAIJ	15AB LHCB	0	$p\bar{p}$ at 7, 8 TeV
1.0 ±0.5 ±0.8	4k	AAIJ	15AB LHCB	±	$p\bar{p}$ at 7, 8 TeV
1.10±0.42±0.31		¹ ABAZOV	07T D0	0	$p\bar{p}$ at 1.96 TeV

NODE=M245R03
NODE=M245R03

¹ Converted from measured ratio of $R = B(B_2^{*0} \rightarrow B^{*+}\pi^-) / B(B_2^{*0} \rightarrow B^{(*)+}\pi^-) = 0.475 \pm 0.095 \pm 0.069$.

NODE=M245R03;LINKAGE=AB

 $B_2^*(5747)$ REFERENCES

NODE=M245

AAIJ	15AB	JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I	PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09D	PRL 102 102003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07T	PRL 99 172001	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=56628
REFID=56029
REFID=52700
REFID=52014

$B_J(5840)$

$$I(J^P) = \frac{1}{2}(??)$$

I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M246

NODE=M246

 $B_J(5840)$ MASS

NODE=M246205

 $B_J(5840)^+$ MASS

NODE=M246M+

OUR FIT uses m_{B^0} and $m_{B_J(5840)^+} - m_{B^0}$ to determine $m_{B_J(5840)^+}$.NODE=M246M+
NODE=M246M+

VALUE (MeV)	DOCUMENT ID
-------------	-------------

5851±19 OUR FIT **$m_{B_J(5840)^+} - m_{B^0}$** NODE=M246DM+
NODE=M246DM+

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

571±19 OUR FIT**571±13±14** 7k 1 AAIJ 15AB LHCB pp at 7, 8 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

595±26±14 7k 2 AAIJ 15AB LHCB pp at 7, 8 TeV

OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 431 \pm 13 \pm 14$ MeV which we adjust bythe π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.² AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 455 \pm 26 \pm 14$ MeV which we adjust bythe π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM+;LINKAGE=A

NODE=M246DM+;LINKAGE=B

 $m_{B_J(5840)^+} - m_{B^{*0}}$ NODE=M246DM1
NODE=M246DM1

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

565±15±14 7k 1 AAIJ 15AB LHCB pp at 7, 8 TeV¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 425 \pm 15 \pm 14$ MeV which we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM1;LINKAGE=A

 $B_J(5840)^0$ MASS

NODE=M246M0

OUR FIT uses m_{B^+} and $m_{B_J(5840)^0} - m_{B^+}$ to determine $m_{B_J(5840)^0}$.NODE=M246M0
NODE=M246M0

VALUE (MeV)	DOCUMENT ID
-------------	-------------

5863±9 OUR FIT **$m_{B_J(5840)^0} - m_{B^+}$** NODE=M246DM0
NODE=M246DM0

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

584±9 OUR FIT**584±5±7** 12k 1 AAIJ 15AB LHCB pp at 7, 8 TeV

••• We do not use the following data for averages, fits, limits, etc. •••

610±22±7 12k 2 AAIJ 15AB LHCB pp at 7, 8 TeV

OCCUR=2

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV which we adjust bythe π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.² AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 471 \pm 22 \pm 7$ MeV which we adjust bythe π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM0;LINKAGE=A

NODE=M246DM0;LINKAGE=B

$m_{B_J(5840)^0} - m_{B^{*+}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$584 \pm 5 \pm 7$	12k	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV
-------------------	-----	-------------------	-----------	------------------

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 444 \pm 5 \pm 7$ MeV

which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246DM2
NODE=M246DM2

NODE=M246DM2;LINKAGE=A

 $B_J(5840)$ WIDTH

NODE=M246210

 $B_J(5840)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

220 ± 80 OUR AVERAGE [224 ± 80 MeV OUR 2023 AVERAGE]

224 ± 24 ± 80	7k	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV
----------------------	----	-------------------	-----------	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

215 ± 27 ± 80	7k	² AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------	----	-------------------	-----------	------------------

229 ± 27 ± 80	7k	³ AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------	----	-------------------	-----------	------------------

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246W+
NODE=M246W+

NEW

OCCUR=2

OCCUR=3

NODE=M246W+;LINKAGE=A

NODE=M246W+;LINKAGE=B

NODE=M246W+;LINKAGE=C

 $B_J(5840)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

130 ± 40 OUR AVERAGE [127 ± 40 MeV OUR 2023 AVERAGE]

127 ± 17 ± 34	12k	¹ AAIJ	15AB LHCB	pp at 7, 8 TeV
----------------------	-----	-------------------	-----------	------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

107 ± 20 ± 34	12k	² AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------	-----	-------------------	-----------	------------------

119 ± 17 ± 34	12k	³ AAIJ	15AB LHCB	pp at 7, 8 TeV
---------------	-----	-------------------	-----------	------------------

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M246W0
NODE=M246W0

NEW

OCCUR=2

OCCUR=3

NODE=M246W0;LINKAGE=A

NODE=M246W0;LINKAGE=B

NODE=M246W0;LINKAGE=C

 $B_J(5840)$ DECAY MODES

NODE=M246215;NODE=M246

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^* \pi$	seen
Γ_2 $B \pi$	possibly seen

DESIG=1

DESIG=2

 $B_J(5840)$ BRANCHING RATIOS

NODE=M246220

$\Gamma(B^* \pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
---	-------	------	-------------	------	-----	---------	-------------------

seen	7k	AAIJ	15AB LHCB	±		pp at 7, 8 TeV	
------	----	------	-----------	---	--	------------------	--

seen	12k	AAIJ	15AB LHCB	0		pp at 7, 8 TeV	
------	-----	------	-----------	---	--	------------------	--

NODE=M246R01
NODE=M246R01

OCCUR=2

$\Gamma(B \pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ
---------------------------------------	-------	------	-------------	------	-----	---------	-------------------

possibly seen	7k	¹ AAIJ	15AB LHCB	±		pp at 7, 8 TeV	
---------------	----	-------------------	-----------	---	--	------------------	--

possibly seen		¹ AAIJ	15AB LHCB	0		pp at 7, 8 TeV	
---------------	--	-------------------	-----------	---	--	------------------	--

NODE=M246R02
NODE=M246R02

OCCUR=2

¹ A $B \pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^* \pi$ is allowed.

NODE=M246R02;LINKAGE=A

 $B_J(5840)$ REFERENCES

NODE=M246

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	--------------------	-----------------------	----------------

REFID=56628

$B_J(5970)$

$$I(J^P) = \frac{1}{2}(??)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M248

NODE=M248

NODE=M248205

NODE=M248M+

NODE=M248M+

NODE=M248M+

NEW

NODE=M248DM+

NODE=M248DM+

OCCUR=2

NODE=M248DM+;LINKAGE=B

NODE=M248DM+;LINKAGE=A

NODE=M248DM+;LINKAGE=C

NODE=M248DM1

NODE=M248DM1

NODE=M248DM1;LINKAGE=A

NODE=M248M0

NODE=M248M0

NODE=M248M0

NODE=M248DM0

NODE=M248DM0

OCCUR=2

NODE=M248DM0;LINKAGE=B

NODE=M248DM0;LINKAGE=A

NODE=M248DM0;LINKAGE=C

 $B_J(5970)$ MASS **$B_J(5970)^+$ MASS**OUR FIT uses m_{B^0} and $m_{B_J(5970)^+} - m_{B^0}$ to determine $m_{B_J(5970)^+}$.

VALUE (MeV)

DOCUMENT ID

5965 ± 5 OUR FIT

[5964 ± 5 MeV OUR 2023 FIT]

 $m_{B_J(5970)^+} - m_{B^0}$

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

685 ± 5 OUR FIT**685 ± 5 OUR AVERAGE**

685.3 ± 4.1 ± 2.5

2k

1 AAIJ

15AB LHCB pp at 7, 8 TeV

681 ± 5 ± 12

1.4k

2 AALTONEN

14l CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.8 ± 4.5 ± 2.5

2k

3 AAIJ

15AB LHCB pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 545.8 \pm 4.1 \pm 2.5$ MeV which we adjust by

the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AALTONEN 14l reports $m_{B_J(5970)^+} - m_{B^0} - m_{\pi^+} = 541 \pm 5 \pm 12$ MeV which we adjusted by the π^+ mass.

³ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - m_{\pi^+} = 547 \pm 5 \pm 3$ MeV which we adjust by

the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

 $m_{B_J(5970)^+} - m_{B^{*0}}$

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

686.0 ± 4.0 ± 2.5

2k

1 AAIJ

15AB LHCB pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^+} - m_{B^0}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^+} = 547 \pm 4 \pm 3$ MeV which

we adjust by the π^+ mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*0}} - m_{B^0}) = (m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

 $B_J(5970)^0$ MASSOUR FIT uses m_{B^+} and $m_{B_J(5970)^0} - m_{B^+}$ to determine $m_{B_J(5970)^0}$.

VALUE (MeV)

DOCUMENT ID

5971 ± 5 OUR FIT **$m_{B_J(5970)^0} - m_{B^+}$**

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

691 ± 5 OUR FIT**691 ± 5 OUR AVERAGE**

689.9 ± 2.9 ± 5.1

10k

1 AAIJ

15AB LHCB pp at 7, 8 TeV

698 ± 5 ± 12

2.6k

2 AALTONEN

14l CDF $p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

714.3 ± 6.4 ± 5.1

10k

3 AAIJ

15AB LHCB pp at 7, 8 TeV

¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 550.4 \pm 2.9 \pm 5.1$ MeV which we adjust by

the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses two relativistic Breit-Wigner functions in the fit for mass difference.

² AALTONEN 14l reports $m_{B_J(5970)^0} - m_{B^+} - m_{\pi^-} = 558 \pm 5 \pm 12$ MeV which we adjusted by the π^- mass.

³ AAIJ 15AB reports $[m_{B_J^0} - m_{B^+}] - m_{\pi^-} = 575 \pm 6 \pm 5$ MeV which we adjust by

the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = (-1)^J$ and uses three relativistic Breit-Wigner functions in the fit for mass difference.

$m_{B_J(5970)^0} - m_{B^{*+}}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

691.6 ± 3.7 ± 5.1	10k	¹ AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV
¹ AAIJ 15AB reports $[m_{B_J^0} - m_{B^{*+}}] - (m_{B^{*+}} - m_{B^+}) - m_{\pi^-} = 552 \pm 4 \pm 5$ MeV				

which we adjust by the π^- mass. The masses inside the square brackets were measured for each candidate event. The result assumes $P = -(-1)^J$, $(m_{B^{*+}} - m_{B^+}) = 45.01 \pm 0.30 \pm 0.23$ MeV, and uses three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M248DM2
NODE=M248DM2

NODE=M248DM2;LINKAGE=A

 $B_J(5970)$ WIDTH

NODE=M248210

 $B_J(5970)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

62 ± 20 OUR AVERAGE

63 ± 15 ± 17	2k	¹ AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV
60 ⁺³⁰ ₋₂₀ ± 40	1.4k	AALTONEN	14I CDF	$\rho\bar{\rho}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

61 ± 14 ± 17	2k	² AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV
61 ± 15 ± 17	2k	³ AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M248W+
NODE=M248W+

OCCUR=2
OCCUR=3

NODE=M248W+;LINKAGE=A

NODE=M248W+;LINKAGE=B

NODE=M248W+;LINKAGE=C

 $B_J(5970)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

81 ± 12 OUR AVERAGE

82 ± 8 ± 9	10k	¹ AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV
70 ⁺³⁰ ₋₂₀ ± 30	2.6k	AALTONEN	14I CDF	$\rho\bar{\rho}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

56 ± 7 ± 9	10k	² AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV
82 ± 10 ± 9	10k	³ AAIJ	15AB LHCB	$\rho\rho$ at 7, 8 TeV

¹ Assuming $P = (-1)^J$ and using two relativistic Breit-Wigner functions in the fit for mass difference.

² Assuming $P = (-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

³ Assuming $P = -(-1)^J$ and using three relativistic Breit-Wigner functions in the fit for mass difference.

NODE=M248W0
NODE=M248W0

OCCUR=2
OCCUR=3

NODE=M248W0;LINKAGE=A

NODE=M248W0;LINKAGE=B

NODE=M248W0;LINKAGE=C

 $B_J(5970)$ DECAY MODES

NODE=M248215;NODE=M248

Mode	Fraction (Γ_i/Γ)
Γ_1 $B\pi$	possibly seen
Γ_2 $B^*\pi$	seen

DESIG=1
DESIG=2

 $B_J(5970)$ BRANCHING RATIOS

NODE=M248220

$\Gamma(B\pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
--------------------------------------	-------	------	-------------	------	-----	---------	-------------------

possibly seen		2k	¹ AAIJ	15AB LHCB	±	$\rho\rho$ at 7, 8 TeV	
possibly seen		10k	¹ AAIJ	15AB LHCB	0	$\rho\rho$ at 7, 8 TeV	
possibly seen		2.6k	AALTONEN	14I CDF	0	$\rho\bar{\rho}$ at 1.96 TeV	
possibly seen		1.4k	AALTONEN	14I CDF	±	$\rho\bar{\rho}$ at 1.96 TeV	

¹ A $B\pi$ decay is forbidden from a $P = -(-1)^J$ parent, whereas $B^*\pi$ is allowed.

NODE=M248R01
NODE=M248R01

OCCUR=2

OCCUR=2

NODE=M248R01;LINKAGE=A

$\Gamma(B^*\pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ
--	-------	------	-------------	------	-----	---------	-------------------

seen		10k	AAIJ	15AB LHCB	0	$\rho\rho$ at 7, 8 TeV	
seen		2k	AAIJ	15AB LHCB	±	$\rho\rho$ at 7, 8 TeV	
seen		2.6k	AALTONEN	14I CDF	0	$\rho\bar{\rho}$ at 1.96 TeV	
seen		1.4k	AALTONEN	14I CDF	±	$\rho\bar{\rho}$ at 1.96 TeV	

NODE=M248R02
NODE=M248R02

OCCUR=2

OCCUR=2

 $B_J(5970)$ REFERENCES

NODE=M248

AAIJ	15AB JHEP 1504 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)

REFID=56628
REFID=56029

BOTTOM, STRANGE MESONS

($B = \pm 1, S = \mp 1$)

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \text{ similarly for } B_s^{*\prime}s$$

$B_{s1}(5830)^0$

$$I(J^P) = 0(1^+)$$

I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

$B_{s1}(5830)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5828.73 ± 0.20 OUR FIT			
[5828.70 ± 0.20 MeV OUR 2023 FIT]			
5828.65 ± 0.24 OUR AVERAGE			
5828.78 ± 0.09 ± 0.29	¹ SIRUNYAN	18DF CMS	pp at 8 TeV
5828.40 ± 0.04 ± 0.41	¹ AAIJ	130 LHCb	pp at 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5829.4 ± 0.7	² AALTONEN	08k CDF	Repl. by AALTONEN 14i
¹ Uses $B_{s1}(5830)^0 \rightarrow B^{*+} K^-$ decay.			
² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.			

$$m_{B_{s1}^0} - m_{B^{*+}}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
503.98 ± 0.17 OUR FIT			
[503.99 ± 0.17 MeV OUR 2023 FIT]			
504.03 ± 0.12 ± 0.15	¹ AALTONEN	14i CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
504.41 ± 0.21 ± 0.14	² AALTONEN	08k CDF	Repl. by AALTONEN 14i
¹ AALTONEN 14i reports $m_{B_{s1}(5830)^0} - m_{B^{*+}} - m_{K^-} = 10.35 \pm 0.12 \pm 0.15$ MeV which we adjusted by the K^- mass.			
² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.			

$B_{s1}(5830)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
0.5 ± 0.3 ± 0.3	AALTONEN	14i CDF	$p\bar{p}$ at 1.96 TeV

$B_{s1}(5830)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^{*+} K^-$	seen
Γ_2 $B^{*0} K_S^0$	

$B_{s1}(5830)^0$ BRANCHING RATIOS

$\Gamma(B^{*+} K^-)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AALTONEN	08k CDF	$p\bar{p}$ at 1.96 TeV
$\Gamma(B^{*0} K_S^0)/\Gamma(B^{*+} K^-)$	Γ_2/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.49 ± 0.12 ± 0.07	¹ SIRUNYAN	18DF CMS	pp at 8 TeV
¹ With the branching fractions $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$ and $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$.			

NODE=MXXX046

NODE=MXXX046

NODE=M187

NODE=M187

NODE=M187M

NODE=M187M

NEW

NODE=M187M;LINKAGE=AI

NODE=M187M;LINKAGE=AA

NODE=M187DM

NODE=M187DM

NEW

NODE=M187DM;LINKAGE=AL

NODE=M187DM;LINKAGE=AA

NODE=M187W

NODE=M187W

NODE=M187215;NODE=M187

DESIG=1

DESIG=2

NODE=M187220

NODE=M187R01

NODE=M187R01

NODE=M187R00

NODE=M187R00

NODE=M187R00;LINKAGE=A

$B_{s1}(5830)^0$ REFERENCES

SIRUNYAN	18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AALTONEN	14l PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AAIJ	130 PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)

NODE=M187

REFID=59328
REFID=56029
REFID=54968
REFID=52235 **$B_{s2}^*(5840)^0$** $I(J^P) = 0(2^+)$
 I, J, P need confirmation.

Quantum numbers shown are quark-model predictions.

NODE=M186

NODE=M186

 $B_{s2}^*(5840)^0$ MASS

NODE=M186M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5839.88 ± 0.12 OUR FIT			
[5839.86 ± 0.12 MeV OUR 2023 FIT]			
5839.92 ± 0.14 OUR AVERAGE			
5839.86 ± 0.09 ± 0.17	SIRUNYAN	18DF CMS	pp at 8 TeV
5839.99 ± 0.05 ± 0.20	AAIJ	130 LHCb	pp at 7 TeV
5839.6 ± 1.1 ± 0.7	¹ ABAZOV	08E D0	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5839.7 ± 0.7	² AALTONEN	08K CDF	Repl. by AALTONEN 14l
¹ Observed in $B_{s2}^{*0} \rightarrow B^+ K^-$. Measured production rate of B_{s2}^{*0} relative to B^+ to be $(1.15 \pm 0.23 \pm 0.13)\%$.			
² Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.			

NODE=M186M

NEW

NODE=M186M;LINKAGE=AB

NODE=M186M;LINKAGE=AA

$$m_{B_{s2}^{*0}} - m_{B_{s1}^0}$$

NODE=M186DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.5 ± 0.6	¹ AALTONEN	08K CDF	Repl. by AALTONEN 14l
¹ Uses two-body decays into K^- and B^+ mesons reconstructed as $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$ or $B^+ \rightarrow \bar{D}^0 \pi^+$, $\bar{D}^0 \rightarrow K^+ \pi^-$.			
$m_{B_{s2}^{*0}} - m_{B^+}$			

NODE=M186DM

NODE=M186DM;LINKAGE=AA

NODE=M186DM2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
560.48 ± 0.12 OUR FIT			
[560.52 ± 0.14 MeV OUR 2023 FIT]			
560.41 ± 0.13 ± 0.14	¹ AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV
¹ AALTONEN 14l reports $m_{B_{s2}^*(5840)^0} - m_{B^+} - m_{K^-} = 66.73 \pm 0.13 \pm 0.14$ MeV which we adjusted by the K^- mass.			

NODE=M186DM2

NEW

NODE=M186DM2;LINKAGE=AL

 $B_{s2}^*(5840)^0$ WIDTH

NODE=M186W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1.49 ± 0.27 OUR AVERAGE			
1.52 ± 0.34 ± 0.30	SIRUNYAN	18DF CMS	pp at 8 TeV
1.4 ± 0.4 ± 0.2	AALTONEN	14l CDF	$p\bar{p}$ at 1.96 TeV
1.56 ± 0.13 ± 0.47	¹ AAIJ	130 LHCb	pp at 7 TeV
¹ Uses $B_{s2}^*(5840)^0 \rightarrow B^{*+} K^-$ decays.			

NODE=M186W

NODE=M186W;LINKAGE=AI

 $B_{s2}^*(5840)^0$ DECAY MODES

NODE=M186215;NODE=M186

Branching fractions are given relative to the one **DEFINED AS 1**.

NODE=M186

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^+ K^-$	DEFINED AS 1
Γ_2 $B^{*+} K^-$	0.093 ± 0.018
Γ_3 $B^0 K_S^0$	0.43 ± 0.11
Γ_4 $B^{*0} K_S^0$	0.04 ± 0.04

DESIG=1

DESIG=2

DESIG=4

DESIG=3

$B_{s2}^*(5840)^0$ BRANCHING RATIOS $\Gamma(B^+ K^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AALTONEN	08K CDF	$p\bar{p}$ at 1.96 TeV
seen	¹ ABAZOV	08E D0	$p\bar{p}$ at 1.96 TeV

¹ Measured production rate of B_{s2}^{*0} relative to B^+ to be $(1.15 \pm 0.23 \pm 0.13)\%$.

NODE=M186220

NODE=M186R01
NODE=M186R01

NODE=M186R01;LINKAGE=AB

 $\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.093 ± 0.013 ± 0.012	AAIJ	130 LHCB	pp at 7 TeV

NODE=M186R02
NODE=M186R02 $\Gamma(B^{*0} K_S^0)/\Gamma(B^0 K_S^0)$ Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.093 ± 0.086 ± 0.014	¹ SIRUNYAN	18DF CMS	pp at 8 TeV

NODE=M186R00
NODE=M186R00

¹ With the branching fraction $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$.

NODE=M186R00;LINKAGE=A

 $\Gamma(B^0 K_S^0)/\Gamma(B^+ K^-)$ Γ_3/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.432 ± 0.077 ± 0.078	¹ SIRUNYAN	18DF CMS	pp at 8 TeV

NODE=M186R03
NODE=M186R03

¹ With the branching fractions $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$ and $B(B^0 \rightarrow J/\psi K^{*0}) = (1.28 \pm 0.05) \times 10^{-3}$.

NODE=M186R03;LINKAGE=A

 $\Gamma(B^{*+} K^-)/\Gamma(B^+ K^-)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.081 ± 0.021 ± 0.015	¹ SIRUNYAN	18DF CMS	pp at 8 TeV

NODE=M186R04
NODE=M186R04

¹ With the branching fraction $B(B^+ \rightarrow J/\psi K^+) = (1.026 \pm 0.031) \times 10^{-3}$.

NODE=M186R04;LINKAGE=A

 $B_{s2}^*(5840)^0$ REFERENCES

SIRUNYAN	18DF EPJ C78 939	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59328
AALTONEN	14I PR D90 012013	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=56029
AAIJ	130 PRL 110 151803	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54968
AALTONEN	08K PRL 100 082001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52235
ABAZOV	08E PRL 100 082002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52232

NODE=M186

 $B_{sJ}^*(5850)$ $I(J^P) = ?(??)$

OMITTED FROM SUMMARY TABLE

Signal can be interpreted as coming from $\bar{b}s$ states. Needs confirmation.

NODE=M153

NODE=M153

 $B_{sJ}^*(5850)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5853 ± 15	141	AKERS	95E OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M153M

NODE=M153M

 $B_{sJ}^*(5850)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
47 ± 22	141	AKERS	95E OPAL	$E_{\text{cm}}^{ee} = 88-94$ GeV

NODE=M153W

NODE=M153W

 $B_{sJ}^*(5850)$ REFERENCES

AKERS	95E ZPHY C66 19	R. Akers <i>et al.</i>	(OPAL Collab.)
-------	-----------------	------------------------	----------------

NODE=M153

REFID=44182

$B_{sJ}(6063)^0$

$I(J^P) = 0(?^?)$

OMITTED FROM SUMMARY TABLE

 $B_{sJ}(6063)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$6063.5 \pm 1.2 \pm 0.8$	¹ AAIJ	21D LHCB	pp at 7, 8, 13 TeV

¹ Seen in the decay channel $B^\pm K^\mp$. Integrated luminosity = 9 fb⁻¹.

NODE=M257M

NODE=M257M

NODE=M257M;LINKAGE=A

 $B_{sJ}(6063)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$26 \pm 4 \pm 4$	¹ AAIJ	21D LHCB	pp at 7, 8, 13 TeV

¹ Seen in the decay channel $B^\pm K^\mp$. Integrated luminosity = 9 fb⁻¹.

NODE=M257W

NODE=M257W

NODE=M257W;LINKAGE=A

 $B_{sJ}(6063)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^+ K^-$	seen

NODE=M257215;NODE=M257

DESIG=1

$\Gamma(B^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	AAIJ	21D LHCB	$B^\pm K^\mp$ mass spectrum	

NODE=M257R01
NODE=M257R01 $B_{sJ}(6063)^0$ REFERENCESAAIJ 21D EPJ C81 601 R. Aaij *et al.* (LHCb Collab.)

NODE=M257

REFID=61130

NODE=M258

 $B_{sJ}(6114)^0$

$I(J^P) = 0(?^?)$

OMITTED FROM SUMMARY TABLE

 $B_{sJ}(6114)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$6114 \pm 3 \pm 5$	¹ AAIJ	21D LHCB	pp at 7, 8, 13 TeV

¹ Seen in the decay channel $B^\pm K^\mp$. Integrated luminosity = 9 fb⁻¹.

NODE=M258M

NODE=M258M

NODE=M258M;LINKAGE=A

 $B_{sJ}(6114)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$66 \pm 18 \pm 21$	¹ AAIJ	21D LHCB	pp at 7, 8, 13 TeV

¹ Seen in the decay channel $B^\pm K^\mp$. Integrated luminosity = 9 fb⁻¹.

NODE=M258W

NODE=M258W

NODE=M258W;LINKAGE=B

 $B_{sJ}(6114)^0$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $B^+ K^-$	seen

NODE=M258215;NODE=M258

DESIG=1

$\Gamma(B^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	AAIJ	21D LHCB	$B^\pm K^\mp$ mass spectrum	

NODE=M258R01
NODE=M258R01 $B_{sJ}(6114)^0$ REFERENCESAAIJ 21D EPJ C81 601 R. Aaij *et al.* (LHCb Collab.)

NODE=M258

REFID=61130

BOTTOM, CHARMED MESONS ($B = C = \pm 1$)

$$B_c^+ = c\bar{b}, B_c^- = \bar{c}b, \text{ similarly for } B_c^{*'}s$$

$B_c(2S)^\pm$

$$I(J^P) = 0(0^-)$$

Quantum numbers neither measured nor confirmed.

$B_c(2S)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
6871.2±1.0 OUR AVERAGE				
6871.7±1.3±0.3	24	^{1,2} AAIJ	19Y LHCB	pp at 7, 8, 13 TeV
6870.6±1.4±0.3	51	^{3,4} SIRUNYAN	19M CMS	pp at 13 TeV
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
not seen		⁵ AAIJ	18AL LHCB	pp at 8 TeV
6842 ±4 ±5	57	^{6,7} AAD	14AQ ATLS	pp at 7, 8 TeV

¹ AAIJ 19Y observed $B_c(2S)^+$ in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 2.2 (3.2) global (local) standard deviations significance.

² AAIJ 19Y reports mass difference measurement of $M(B_c(2S)^+) - M(B_c^+) = 597.2 \pm 1.3 \pm 0.1$ MeV. We have adjusted this measurement with our best value of $M(B_c^+) = 6274.47 \pm 0.32$ MeV. The first uncertainty of the $M(B_c(2S)^+)$ value is a total of uncertainties reported by the experiment and the second one comes from our best value of $M(B_c^+)$.

³ SIRUNYAN 19M observed $B_c(2S)^+$ in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 6.5 standard deviations significance.

⁴ SIRUNYAN 19M reports mass difference measurement of $M(B_c(2S)^+) - M(B_c^+) = 596.1 \pm 1.2 \pm 0.8$ MeV. We have adjusted this measurement with our best value of $M(B_c^+) = 6274.47 \pm 0.32$ MeV. The first uncertainty of the $M(B_c(2S)^+)$ value is a total of uncertainties reported by the experiment and the second one comes from our best value of $M(B_c^+)$.

⁵ AAIJ 18AL reports an upper limit on the ratio of production cross sections for $[\sigma(B_c(2S)^+)/\sigma(B_c^+)]: B(B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-) < 0.04-0.09$ at 95% CL for the mass value reported by AAD 14AQ.

⁶ Observed in the decay mode $B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-$ ($B_c^+ \rightarrow J/\psi \pi^+$) with 5.2 standard deviations significance.

⁷ Might be the $B_c^*(2S)$.

$B_c(2S)^\pm$ DECAY MODES $\times B(\bar{b} \rightarrow B_c(2S))$

The following quantities are not pure branching ratios; rather the fractions $\Gamma_i/\Gamma \times B(\bar{b} \rightarrow B_c(2S))$.

Mode	Fraction (Γ_i/Γ)
Γ_1 $B_c^+ \pi^+ \pi^-$	seen

$B_c(2S)^\pm$ BRANCHING RATIOS

$\Gamma(B_c^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	57	¹ AAD	14AQ ATLS	pp at 7, 8 TeV
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
not seen		² AAIJ	18AL LHCB	pp at 8 TeV

¹ Observed with 5.2 standard deviations significance.

² AAIJ 18AL reports an upper limit on the ratio of production cross sections for $[\sigma(B_c(2S)^+)/\sigma(B_c^+)]: B(B_c(2S)^+ \rightarrow B_c^+ \pi^+ \pi^-) < 0.04-0.09$ at 95% CL for the mass value reported by AAD 14AQ.

NODE=MXXX049

NODE=MXXX049

NODE=M217

NODE=M217

NODE=M217M

NODE=M217M

SYCLP=A

SYCLP=A

NODE=M217M;LINKAGE=E

NODE=M217M;LINKAGE=F

NODE=M217M;LINKAGE=B

NODE=M217M;LINKAGE=D

NODE=M217M;LINKAGE=A

NODE=M217M;LINKAGE=AA

NODE=M217M;LINKAGE=C

NODE=M217215;NODE=M217

NODE=M217

DESIG=1

NODE=M217225

NODE=M217R01

NODE=M217R01

NODE=M217R01;LINKAGE=AA

NODE=M217R01;LINKAGE=A

$B_c(2S)^\pm$ REFERENCES

AAJ	19Y	PRL 122 232001	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	19M	PRL 122 132001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAJ	18AL	JHEP 1801 138	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	14AQ	PRL 113 212004	G. Aad <i>et al.</i>	(ATLAS Collab.)

NODE=M217

REFID=59795
REFID=59639
REFID=59246
REFID=56117

See the related review(s):
Spectroscopy of Mesons Containing Two Heavy Quarks


$c\bar{c}$ MESONS

(including possibly non- $q\bar{q}$ states)

NODE=MXXX025

NODE=M826

Updated March 2024.



The level scheme of meson states containing a minimal quark content of $c\bar{c}$ and having $S = C = B = 0$. The name of a state is determined by its quantum numbers $I^G J^{PC}$ (see the review “Naming Scheme for Hadrons”). States included in the Summary Tables are shown with solid lines; selected states not in the Summary Tables, but with assigned quantum numbers, are shown with dotted lines. The arrows indicate the most dominant hadronic transitions. Single photon transitions, including $\psi(nS) \rightarrow \gamma\eta_c(mS)$, $\psi(nS) \rightarrow \gamma\chi_{cJ}(1P)$, and $\chi_{cJ}(1P) \rightarrow \gamma J/\psi$, are omitted for clarity. For orientation, the location of the thresholds related to a pair of ground state open charm mesons is indicated in the figure. The J^P of $T_{c\bar{c}}(4020)$ is not established. Decays to open flavour final states are not shown in the figure.

$\eta_c(1S)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

NODE=M026

 $\eta_c(1S)$ MASS

NODE=M026M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2984.1 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.2. [2983.9 ± 0.4 MeV OUR 2023 AVERAGE Scale factor = 1.2]		
2985.01 ± 0.17 ± 0.89	35k	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
2983.9 ± 0.7 ± 0.1		¹ AAIJ	20H LHCB	$p p \rightarrow b X \rightarrow p \bar{p} X$
2985.9 ± 0.7 ± 2.1	1705	ABLIKIM	19AV BES3	$J/\psi \rightarrow \gamma \omega \omega$
2984.6 ± 0.7 ± 2.2	2673	XU	18 BELL	$e^+ e^- \rightarrow$ $e^+ e^- \eta' \pi^+ \pi^-$
2986.7 ± 0.5 ± 0.9	11k	² AAIJ	17AD LHCB	$p p \rightarrow B^+ X \rightarrow$ $p \bar{p} K^+ X$
2982.8 ± 1.0 ± 0.5	6.4k	³ AAIJ	17BB LHCB	$p p \rightarrow b \bar{b} X \rightarrow$ $2(K^+ K^-) X$
2982.2 ± 1.5 ± 0.1	2.0k	⁴ AAIJ	15BI LHCB	$p p \rightarrow \eta_c(1S) X$
2983.5 ± 1.4 $\begin{smallmatrix} + 1.6 \\ - 3.6 \end{smallmatrix}$		⁵ ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma \eta_c$
2979.8 ± 0.8 ± 3.5	4.5k	^{6,7} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$
2984.1 ± 1.1 ± 2.1	900	^{6,7,8} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$
2984.3 ± 0.6 ± 0.6		^{9,10} ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma \eta_c$
2984.49 ± 1.16 ± 0.52	832	⁶ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
2982.7 ± 1.8 ± 2.2	486	ZHANG	12A BELL	$e^+ e^- \rightarrow$ $e^+ e^- \eta' \pi^+ \pi^-$
2984.5 ± 0.8 ± 3.1	11k	DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow$ $K^+ K^- \pi^+ \pi^- \pi^0$
2985.4 ± 1.5 $\begin{smallmatrix} + 0.5 \\ - 2.0 \end{smallmatrix}$	920	¹⁰ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm \pi^\mp)$
2982.2 ± 0.4 ± 1.6	14k	¹¹ LEES	10 BABR	$10.6 e^+ e^- \rightarrow$ $e^+ e^- K_S^0 K^\pm \pi^\mp$
2985.8 ± 1.5 ± 3.1	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S) K^(*) \rightarrow$ $K \bar{K} \pi K^(*)$
2986.1 ± 1.0 ± 2.5	7.5k	UEHARA	08 BELL	$\gamma \gamma \rightarrow \eta_c \rightarrow$ hadrons
2970 ± 5 ± 6	501	¹² ABE	07 BELL	$e^+ e^- \rightarrow J/\psi(c\bar{c})$
2971 ± 3 $\begin{smallmatrix} + 2 \\ - 1 \end{smallmatrix}$	195	WU	06 BELL	$B^+ \rightarrow p \bar{p} K^+$
2974 ± 7 $\begin{smallmatrix} + 2 \\ - 1 \end{smallmatrix}$	20	WU	06 BELL	$B^+ \rightarrow \Lambda \bar{\Lambda} K^+$
2981.8 ± 1.3 ± 1.5	592	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta_c' \rightarrow$ $K_S^0 K^\pm \pi^\mp$
2984.1 ± 2.1 ± 1.0	190	¹³ AMBROGIANI	03 E835	$\bar{p} p \rightarrow \eta_c \rightarrow \gamma \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2982.5 ± 0.4 ± 1.4	12k	¹⁴ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
2982.2 ± 0.6		¹⁵ MITCHELL	09 CLEO	$e^+ e^- \rightarrow \gamma X$
2982 ± 5	270	¹⁶ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
2982.5 ± 1.1 ± 0.9	2.5k	¹⁷ AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(1S) \rightarrow K \bar{K} \pi$
2977.5 ± 1.0 ± 1.2		^{15,18} BAI	03 BES	$J/\psi \rightarrow \gamma \eta_c$
2979.6 ± 2.3 ± 1.6	180	¹⁹ FANG	03 BELL	$B \rightarrow \eta_c K$
2976.3 ± 2.3 ± 1.2		^{15,20} BAI	00F BES	$J/\psi, \psi(2S) \rightarrow \gamma \eta_c$
2976.6 ± 2.9 ± 1.3	140	^{15,21} BAI	00F BES	$J/\psi \rightarrow \gamma \eta_c$
2980.4 ± 2.3 ± 0.6		²² BRANDENB...	00B CLE2	$\gamma \gamma \rightarrow \eta_c \rightarrow$ $K^\pm K_S^0 \pi^\mp$
2975.8 ± 3.9 ± 1.2		²¹ BAI	99B BES	Sup. by BAI 00F
2999 ± 8	25	ABREU	98O DLPH	$e^+ e^- \rightarrow e^+ e^-$ +hadrons
2988.3 $\begin{smallmatrix} + 3.3 \\ - 3.1 \end{smallmatrix}$		ARMSTRONG	95F E760	$\bar{p} p \rightarrow \gamma \gamma$
2974.4 ± 1.9		^{15,23} BISELLO	91 DM2	$J/\psi \rightarrow \eta_c \gamma$
2969 ± 4 ± 4	80	¹⁵ BAI	90B MRK3	$J/\psi \rightarrow$ $\gamma K^+ K^- K^+ K^-$
2956 ± 12 ± 12		¹⁵ BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
2982.6 $\begin{smallmatrix} + 2.7 \\ - 2.3 \end{smallmatrix}$	12	BAGLIN	87B SPEC	$\bar{p} p \rightarrow \gamma \gamma$
2980.2 ± 1.6		^{15,23} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

NODE=M026M

NODE=M026M
NEW

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

2984	$\pm 2.3 \pm 4.0$	¹⁵ GAISER	86	CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
2976	± 8	^{15,24} BALTRUSAIT.	.84	MRK3	$J/\psi \rightarrow 2\phi\gamma$
2982	± 8	¹⁸ ²⁵ HIMEL	80B	MRK2	e^+e^-
2980	± 9	²⁵ PARTRIDGE	80B	CBAL	e^+e^-

¹ AAIJ 20H report $m_{J/\psi} - m_{\eta_c(1S)} = 113.0 \pm 0.7 \pm 0.1$ MeV. We use the current value $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to obtain the quoted mass.

NODE=M026M;LINKAGE=H

² AAIJ 17AD report $m_{J/\psi} - m_{\eta_c(1S)} = 110.2 \pm 0.5 \pm 0.9$ MeV. We use the current value $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to obtain the quoted mass.

NODE=M026M;LINKAGE=F

³ From a fit of the $\phi\phi$ invariant mass with the mass and width of $\eta_c(1S)$ as free parameters.

NODE=M026M;LINKAGE=G

⁴ AAIJ 15BI reports $m_{J/\psi} - m_{\eta_c(1S)} = 114.7 \pm 1.5 \pm 0.1$ MeV from a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. We have used current value of $m_{J/\psi} = 3096.900 \pm 0.006$ MeV to arrive at the quoted $m_{\eta_c(1S)}$ result.

NODE=M026M;LINKAGE=D

⁵ Taking into account an asymmetric photon lineshape.

NODE=M026M;LINKAGE=E

⁶ With floating width.

NODE=M026M;LINKAGE=AL

⁷ Ignoring possible interference with the non-resonant 0^- amplitude.

NODE=M026M;LINKAGE=LS

⁸ Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+\pi^-\pi^0$ decays.

NODE=M026M;LINKAGE=EL

⁹ From a simultaneous fit to six decay modes of the η_c .

NODE=M026M;LINKAGE=BL

¹⁰ Accounts for interference with non-resonant continuum.

NODE=M026M;LINKAGE=VA

¹¹ Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.

NODE=M026M;LINKAGE=LE

¹² From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

NODE=M026M;LINKAGE=EB

¹³ Using mass of $\psi(2S) = 3686.00$ MeV.

NODE=M026M;LINKAGE=BG

¹⁴ Not independent from the measurements reported by LEES 10.

NODE=M026M;LINKAGE=DE

¹⁵ MITCHELL 09 observes a significant asymmetry in the lineshapes of $\psi(2S) \rightarrow \gamma\eta_c$ and $J/\psi \rightarrow \gamma\eta_c$ transitions. If ignored, this asymmetry could lead to significant bias whenever the mass and width are measured in $\psi(2S)$ or J/ψ radiative decays.

NODE=M026M;LINKAGE=MI

¹⁶ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

NODE=M026M;LINKAGE=AU

¹⁷ Superseded by LEES 10.

NODE=M026M;LINKAGE=UB

¹⁸ From a simultaneous fit of five decay modes of the η_c .

NODE=M026M;LINKAGE=AK

¹⁹ Superseded by VINOKUROVA 11.

NODE=M026M;LINKAGE=FA

²⁰ Weighted average of the $\psi(2S)$ and $J/\psi(1S)$ samples. Using an η_c width of 13.2 MeV.

NODE=M026M;LINKAGE=KZ

²¹ Average of several decay modes. Using an η_c width of 13.2 MeV.

NODE=M026M;LINKAGE=C1

²² Superseded by ASNER 04.

NODE=M026M;LINKAGE=NN

²³ Average of several decay modes.

NODE=M026M;LINKAGE=A

²⁴ $\eta_c \rightarrow \phi\phi$.

NODE=M026M;LINKAGE=B

²⁵ Mass adjusted by us to correspond to $J/\psi(1S)$ mass = 3097 MeV.

NODE=M026M;LINKAGE=M

$\eta_c(1S)$ WIDTH

NODE=M026W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
30.5\pm 0.5 OUR FIT	Error includes scale factor of 1.2.	[32.0 \pm 0.7 MeV OUR 2023 FIT]		
30.5\pm 0.5 OUR AVERAGE	Error includes scale factor of 1.1.	[32.1 \pm 0.8 MeV OUR 2023 AVERAGE Scale factor = 1.1]		
29.7 \pm 0.5 \pm 0.2	35k	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$
33.8 \pm 1.6 \pm 4.1	1705	ABLIKIM	19AV BES3	$J/\psi \rightarrow \gamma\omega\omega$
30.8 $^{+2.3}_{-2.2}$ \pm 2.9	2673	XU	18 BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
34.0 \pm 1.9 \pm 1.3	11k	AAIJ	17AD LHCB	$pp \rightarrow B^+X \rightarrow p\bar{p}K^+X$
31.4 \pm 3.5 \pm 2.0	6.4k	¹ AAIJ	17BB LHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$
27.2 \pm 3.1 $^{+5.4}_{-2.6}$		² ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma\eta_c$
25.2 \pm 2.6 \pm 2.4	4.5k	^{3,4} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\pi^0$
34.8 \pm 3.1 \pm 4.0	900	^{3,4,5} LEES	14E BABR	$\gamma\gamma \rightarrow K^+K^-\eta$
32.0 \pm 1.2 \pm 1.0		^{6,7} ABLIKIM	12F BES3	$\psi(2S) \rightarrow \gamma\eta_c$
36.4 \pm 3.2 \pm 1.7	832	³ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma$ hadrons
37.8 $^{+5.8}_{-5.3}$ \pm 3.1	486	ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
36.2 \pm 2.8 \pm 3.0	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$
35.1 \pm 3.1 $^{+1.0}_{-1.6}$	920	⁷ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm(K_S^0 K^\pm\pi^\mp)$
31.7 \pm 1.2 \pm 0.8	14k	⁸ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^-K_S^0 K^\pm\pi^\mp$
36.3 $^{+3.7}_{-3.6}$ \pm 4.4	0.9k	AUBERT	08AB BABR	$B \rightarrow \eta_c(1S)K^(*) \rightarrow K\bar{K}\pi K^(*)$
28.1 \pm 3.2 \pm 2.2	7.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \eta_c \rightarrow$ hadrons
48 $^{+8}_{-7}$ \pm 5	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$

NODE=M026W

NEW

NEW

OCCUR=2

40 ±19 ±5	20	WU	06	BELL	$B^+ \rightarrow \Lambda \bar{\Lambda} K^+$	OCCUR=2
24.8 ± 3.4 ± 3.5	592	ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$	
20.4 ⁺ ₋ 7.7 ± 2.0	190	AMBROGIANI	03	E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$	
23.9 ⁺ ₋ 12.6 ± 7.1		ARMSTRONG	95F	E760	$\bar{p}p \rightarrow \gamma\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

32.1 ± 1.1 ± 1.3	12k	⁹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	OCCUR=2
34.3 ± 2.3 ± 0.9	2.5k	¹⁰ AUBERT	04D	BABR $\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K \bar{K} \pi$	
17.0 ± 3.7 ± 7.4		¹¹ BAI	03	BES $J/\psi \rightarrow \gamma \eta_c$	
29 ± 8 ± 6	180	¹² FANG	03	BELL $B \rightarrow \eta_c K$	
11.0 ± 8.1 ± 4.1		¹³ BAI	00F	BES $J/\psi \rightarrow \gamma \eta_c$ and $\psi(2S) \rightarrow \gamma \eta_c$	
27.0 ± 5.8 ± 1.4		¹⁴ BRANDENB...	00B	CLE2 $\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$	
7.0 ⁺ ₋ 7.5 ± 7.0	12	BAGLIN	87B	SPEC $\bar{p}p \rightarrow \gamma\gamma$	
10.1 ⁺ ₋ 33.0 ± 8.2	23	¹⁵ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma p \bar{p}$	
11.5 ± 4.5		GAISER	86	CBAL $J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$	
< 40 90% CL	18	HIMEL	80B	MRK2 $e^+ e^-$	
< 20 90% CL		PARTRIDGE	80B	CBAL $e^+ e^-$	

¹ From a fit of the $\phi\phi$ invariant mass with the mass and width of $\eta_c(1S)$ as free parameters.

² Taking into account an asymmetric photon lineshape.

³ With floating mass.

⁴ Ignoring possible interference with the non-resonant 0^- amplitude.

⁵ Using both, $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶ From a simultaneous fit to six decay modes of the η_c .

⁷ Accounts for interference with non-resonant continuum.

⁸ Taking into account interference with the non-resonant $J^P = 0^-$ amplitude.

⁹ Not independent from the measurements reported by LEES 10.

¹⁰ Superseded by LEES 10.

¹¹ From a simultaneous fit of five decay modes of the η_c .

¹² Superseded by VINOKUROVA 11.

¹³ From a fit to the 4-prong invariant mass in $\psi(2S) \rightarrow \gamma \eta_c$ and $J/\psi(1S) \rightarrow \gamma \eta_c$ decays.

¹⁴ Superseded by ASNER 04.

¹⁵ Positive and negative errors correspond to 90% confidence level.

NODE=M026W;LINKAGE=B
 NODE=M026W;LINKAGE=A
 NODE=M026W;LINKAGE=AL
 NODE=M026W;LINKAGE=LS
 NODE=M026W;LINKAGE=EL
 NODE=M026W;LINKAGE=BL
 NODE=M026W;LINKAGE=VA
 NODE=M026W;LINKAGE=LE
 NODE=M026W;LINKAGE=DE
 NODE=M026W;LINKAGE=UB
 NODE=M026W;LINKAGE=AK
 NODE=M026W;LINKAGE=FA
 NODE=M026W;LINKAGE=KZ
 NODE=M026W;LINKAGE=NN
 NODE=M026W;LINKAGE=L

$\eta_c(1S)$ DECAY MODES

NODE=M026215;NODE=M026

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Decays involving hadronic resonances		
Γ_1 $\eta'(958) \pi\pi$	(2.0 ± 0.4) %	S=1.4
Γ_2 $\eta'(958) K \bar{K}$	(1.73 ± 0.35) %	
Γ_3 $\eta'(958) \eta\eta$	(3.4 ± 0.6) × 10 ⁻³	
Γ_4 $\rho\rho$	(1.8 ± 0.4) %	
Γ_5 $K^*(892)^0 K^- \pi^+ + \text{c.c.}$	(1.8 ± 0.5) %	
Γ_6 $K^*(892) \bar{K}^*(892)$	(7.0 ± 1.2) × 10 ⁻³	
Γ_7 $K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-$	(1.4 ± 0.6) %	
Γ_8 $\phi K^+ K^-$	(3.3 ⁺ ₋ 1.2 ± 1.1) × 10 ⁻³	
Γ_9 $\phi\phi$	(1.8 ± 0.4) × 10 ⁻³	S=2.3
Γ_{10} $\phi 2(\pi^+ \pi^-)$	< 4 × 10 ⁻³	CL=90%
Γ_{11} $a_0(980) \pi$	seen	
Γ_{12} $a_2(1320) \pi$	seen	
Γ_{13} $K^*(892) \bar{K} + \text{c.c.}$	< 1.28 %	CL=90%
Γ_{14} $f_2(1270) \eta$	seen	
Γ_{15} $f_2(1270) \eta'$	seen	
Γ_{16} $\omega\omega$	(2.7 ± 0.9) × 10 ⁻³	S=2.1
Γ_{17} $\omega\phi$	< 2.5 × 10 ⁻⁴	CL=90%
Γ_{18} $f_2(1270) f_2(1270)$	(1.08 ± 0.27) %	
Γ_{19} $f_2(1270) f_2'(1525)$	(9.7 ± 3.2) × 10 ⁻³	

NODE=M026;CLUMP=A

DESIG=24

DESIG=85

DESIG=93

DESIG=19

DESIG=26

DESIG=18

DESIG=57

DESIG=28

DESIG=17

DESIG=58

DESIG=21

DESIG=22

DESIG=40

DESIG=23

DESIG=92

DESIG=20

DESIG=47

DESIG=46

DESIG=59

Γ ₂₀	$f_0(500)\eta$	seen	DESIG=86
Γ ₂₁	$f_0(500)\eta'$	seen	DESIG=87
Γ ₂₂	$f_0(980)\eta$	seen	DESIG=70
Γ ₂₃	$f_0(980)\eta'$	seen	DESIG=88
Γ ₂₄	$f_0(1500)\eta$	seen	DESIG=71
Γ ₂₅	$f_0(1710)\eta'$	seen	DESIG=90
Γ ₂₆	$f_0(2100)\eta'$	seen	DESIG=91
Γ ₂₇	$f_0(2200)\eta$	seen	DESIG=72
Γ ₂₈	$a_0(1320)\pi$	seen	DESIG=74
Γ ₂₉	$a_0(1450)\pi$	seen	DESIG=75
Γ ₃₀	$a_2(1700)\pi$	seen	DESIG=94
Γ ₃₁	$a_0(1710)\pi$	seen	DESIG=97
Γ ₃₂	$a_0(1950)\pi$	seen	DESIG=79
Γ ₃₃	$K_0^*(1430)\bar{K} + \text{c.c.}$	seen	DESIG=76
Γ ₃₄	$K_2^*(1430)\bar{K} + \text{c.c.}$	seen	DESIG=77
Γ ₃₅	$K_0^*(1950)\bar{K} + \text{c.c.}$	seen	DESIG=78
Γ ₃₆	$K_0^*(2600)\bar{K} + \text{c.c.}$	seen	DESIG=95

Decays into stable hadrons

			NODE=M026;CLUMP=B	
Γ ₃₇	$K\bar{K}\pi$	(7.1 ± 0.4) %	S=1.1	DESIG=14
Γ ₃₈	$K\bar{K}\eta$	(1.32 ± 0.15) %		DESIG=25
Γ ₃₉	$\eta\pi^+\pi^-$	(1.6 ± 0.4) %		DESIG=16
Γ ₄₀	$\eta 2(\pi^+\pi^-)$	(4.3 ± 1.3) %		DESIG=61
Γ ₄₁	$K^+K^-\pi^+\pi^-$	(8.3 ± 1.8) × 10 ⁻³	S=1.9	DESIG=15
Γ ₄₂	$K^+K^-\pi^+\pi^-\pi^0$	(3.4 ± 0.6) %		DESIG=60
Γ ₄₃	$K^0K^-\pi^+\pi^-\pi^+ + \text{c.c.}$	(5.4 ± 1.5) %		DESIG=62
Γ ₄₄	$K^+K^- 2(\pi^+\pi^-)$	(8.4 ± 2.4) × 10 ⁻³		DESIG=55
Γ ₄₅	$2(K^+K^-)$	(1.4 ± 0.4) × 10 ⁻³	S=1.4	DESIG=27
Γ ₄₆	$\pi^+\pi^-\pi^0$	< 4 × 10 ⁻⁴	CL=90%	DESIG=81
Γ ₄₇	$\pi^+\pi^-\pi^0\pi^0$	(4.6 ± 1.0) %		DESIG=63
Γ ₄₈	$2(\pi^+\pi^-)$	(9.6 ± 1.5) × 10 ⁻³	S=1.4	DESIG=11
Γ ₄₉	$2(\pi^+\pi^-\pi^0)$	(15.9 ± 2.0) %		DESIG=64
Γ ₅₀	$3(\pi^+\pi^-)$	(1.89 ± 0.34) %		DESIG=56
Γ ₅₁	$p\bar{p}$	(1.33 ± 0.11) × 10 ⁻³	S=1.1	DESIG=12
Γ ₅₂	$p\bar{p}\pi^0$	(3.4 ± 1.3) × 10 ⁻³		DESIG=65
Γ ₅₃	$p\bar{p}\pi^+\pi^-$	(3.7 ± 0.5) × 10 ⁻³		DESIG=13
Γ ₅₄	$\Lambda\bar{\Lambda}$	(1.10 ± 0.28) × 10 ⁻³	S=1.5	DESIG=45
Γ ₅₅	$K^+\bar{p}\Lambda + \text{c.c.}$	(2.5 ± 0.4) × 10 ⁻³		DESIG=82
Γ ₅₆	$\bar{\Lambda}(1520)\Lambda + \text{c.c.}$	(3.0 ± 1.3) × 10 ⁻³		DESIG=83
Γ ₅₇	$\Sigma^+\bar{\Sigma}^-$	(2.6 ± 0.5) × 10 ⁻³		DESIG=66
Γ ₅₈	$\Xi^-\bar{\Xi}^+$	(1.07 ± 0.24) × 10 ⁻³		DESIG=67

Radiative decays

Γ ₅₉	$\gamma\gamma$	(1.66 ± 0.13) × 10 ⁻⁴	S=1.2	NODE=M026;CLUMP=C DESIG=31
-----------------	----------------	------------------------------------	-------	-------------------------------

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

			NODE=M026;CLUMP=D	
Γ ₆₀	$\pi^+\pi^-$	$P, CP < 1.3 \times 10^{-4}$	CL=90%	DESIG=51
Γ ₆₁	$\pi^0\pi^0$	$P, CP < 4 \times 10^{-5}$	CL=90%	DESIG=52
Γ ₆₂	K^+K^-	$P, CP < 7 \times 10^{-4}$	CL=90%	DESIG=53
Γ ₆₃	$K_S^0K_S^0$	$P, CP < 4 \times 10^{-4}$	CL=90%	DESIG=54

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_6	14									
x_9	11	13								
x_{16}	7	8	8							
x_{18}	9	11	11	7						
x_{37}	25	25	22	12	17					
x_{38}	13	13	11	6	9	51				
x_{41}	7	7	6	4	5	15	8			
x_{45}	5	5	5	2	3	12	6	4		
x_{48}	13	17	17	10	15	26	13	8	5	
x_{51}	19	20	20	11	16	39	20	11	11	24
x_{53}	7	7	8	4	5	22	11	5	10	8
x_{54}	5	7	7	4	6	12	6	3	4	10
x_{59}	-38	-35	-27	-16	-22	-63	-32	-17	-12	-31
Γ	-1	-1	-1	0	-1	-2	-1	0	0	-1
	x_1	x_6	x_9	x_{16}	x_{18}	x_{37}	x_{38}	x_{41}	x_{45}	x_{48}
x_{53}	21									
x_{54}	13	9								
x_{59}	-47	-17	-11							
Γ	1	0	0	-20						
	x_{51}	x_{53}	x_{54}	x_{59}						

$\eta_c(1S)$ PARTIAL WIDTHS

NODE=M026217

$\Gamma(\gamma\gamma)$

Γ_{59}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M026W1
NODE=M026W1
NEW

5.1 ± 0.4 OUR FIT	Error includes scale factor of 1.2. [5.4 ± 0.4 keV OUR 2023 FIT]			
••• We do not use the following data for averages, fits, limits, etc. •••				
5.8 ± 1.1	486	¹ ZHANG	12A BELL	$e^+e^- \rightarrow e^+e^-\eta'\pi^+\pi^-$
5.2 ± 1.2	273 ± 43	^{2,3} AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
5.5 ± 1.2 ± 1.8	157 ± 33	⁴ KUO	05 BELL	$\gamma\gamma \rightarrow p\bar{p}$
7.4 ± 0.4 ± 2.3		⁵ ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$
13.9 ± 2.0 ± 3.0	41	⁶ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \eta_c$
3.8 ⁺ _{-1.0} 1.1 ⁺ _{-1.0} 1.9 _{-1.0}	190	⁷ AMBROGIANI	03 E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
7.6 ± 0.8 ± 2.3		^{5,8} BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
6.9 ± 1.7 ± 2.1	76	⁹ ACCIARRI	99T L3	$e^+e^- \rightarrow e^+e^-\eta_c$
27 ± 16 ± 10	5	⁵ SHIRAI	98 AMY	58 e^+e^-
6.7 ⁺ _{-1.7} 2.4 ± 2.3		⁴ ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$
11.3 ± 4.2		¹⁰ ALBRECHT	94H ARG	$e^+e^- \rightarrow e^+e^-\eta_c$
8.0 ± 2.3 ± 2.4	17	¹¹ ADRIANI	93N L3	$e^+e^- \rightarrow e^+e^-\eta_c$
5.9 ⁺ _{-1.8} 2.1 ± 1.9		⁷ CHEN	90B CLEO	$e^+e^- \rightarrow e^+e^-\eta_c$
6.4 ⁺ _{-3.4} 5.0 ± 2.4		¹² AIHARA	88D TPC	$e^+e^- \rightarrow e^+e^-X$
4.3 ⁺ _{-3.7} 3.4 ± 2.4		⁴ BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
28 ± 15		^{5,13} BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

- ¹ Assuming there is no interference with the non-resonant background.
² Calculated by us using $\Gamma(\eta_c \rightarrow K\bar{K}\pi) \times \Gamma(\eta_c \rightarrow \gamma\gamma) / \Gamma = 0.44 \pm 0.05$ keV from PDG 06 and $B(\eta_c \rightarrow K\bar{K}\pi) = (8.5 \pm 1.8)\%$ from AUBERT 06E.
³ Systematic errors not evaluated.
⁴ Normalized to $B(\eta_c \rightarrow \rho\bar{\rho}) = (1.3 \pm 0.4) \times 10^{-3}$.
⁵ Normalized to $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$.
⁶ Average of $K_S^0 K^\pm \pi^\mp$, $\pi^+ \pi^- K^+ K^-$, and $2(K^+ K^-)$ decay modes.
⁷ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.
⁸ Superseded by ASNER 04.
⁹ Normalized to the sum of 9 branching ratios.
¹⁰ Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow \phi\phi)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.
¹¹ Superseded by ACCIARRI 99T.
¹² Normalized to the sum of $B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp)$, $B(\eta_c \rightarrow 2K^+ 2K^-)$, $B(\eta_c \rightarrow K^+ K^- \pi^+ \pi^-)$, and $B(\eta_c \rightarrow 2\pi^+ 2\pi^-)$.
¹³ Re-evaluated by AIHARA 88D.

NODE=M026W1;LINKAGE=ZH
 NODE=M026W1;LINKAGE=AU
 NODE=M026W1;LINKAGE=NS
 NODE=M026W1;LINKAGE=N3
 NODE=M026W1;LINKAGE=N2
 NODE=M026W;LINKAGE=FF
 NODE=M026W1;LINKAGE=N6
 NODE=M026W1;LINKAGE=NN
 NODE=M026W1;LINKAGE=N1
 NODE=M026W1;LINKAGE=N5
 NODE=M026W1;LINKAGE=WD
 NODE=M026W1;LINKAGE=N4
 NODE=M026W1;LINKAGE=A

$\eta_c(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M026220

$\Gamma(\eta'(958)\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_{59}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
102 ± 18 OUR FIT		Error includes scale factor of 1.5.	[101 ± 12 eV OUR 2023 FIT]	
98.1 ± 3.9 ± 11.7	2673	XU	18	BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$75.8^{+6.3}_{-6.2} \pm 8.4$	486	¹ ZHANG	12A	BELL $e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$

NODE=M026G10
 NODE=M026G10
 NEW

¹ Superseded by XU 18.

NODE=M026G10;LINKAGE=A

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_{59}/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<39	90	< 1556	UEHARA	08	BELL $\gamma\gamma \rightarrow 2(\pi^+ \pi^-)$

NODE=M026G09
 NODE=M026G09

$\Gamma(K^*(892)\bar{K}^*(892)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_{59}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35 ± 6 OUR FIT		[34 ± 6 eV OUR 2023 FIT]		
32.4 ± 4.2 ± 5.8	882 ± 115	UEHARA	08	BELL $\gamma\gamma \rightarrow \pi^+ \pi^- K^+ K^-$

NODE=M026G08
 NODE=M026G08
 NEW

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_9\Gamma_{59}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.2 ± 2.2 OUR FIT		Error includes scale factor of 2.7.	[8.5 ± 0.9 eV OUR 2023 FIT]	
7.75 ± 0.66 ± 0.62	386 ± 31	¹ LIU	12B	BELL $\gamma\gamma \rightarrow 2(K^+ K^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$6.8 \pm 1.2 \pm 1.3$	132 ± 23	UEHARA	08	BELL $\gamma\gamma \rightarrow 2(K^+ K^-)$

NODE=M026G07
 NODE=M026G07
 NEW

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.

NODE=M026G07;LINKAGE=LI

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_{59}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13 ± 5 OUR FIT		Error includes scale factor of 2.2.	[11.0 ± 2.5 eV OUR 2023 FIT]	
8.67 ± 2.86 ± 0.96	85 ± 29	¹ LIU	12B	BELL $\gamma\gamma \rightarrow 2(\pi^+ \pi^- \pi^0)$

NODE=M026G03
 NODE=M026G03
 NEW

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

NODE=M026G03;LINKAGE=LI

$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_{59}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.49	90	¹ LIU	12B	BELL $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.				

NODE=M026G04
 NODE=M026G04

NODE=M026G04;LINKAGE=LI

$\Gamma(f_2(1270)f_2(1270)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{18}\Gamma_{59}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
55 ± 14 OUR FIT		[52 ± 13 eV OUR 2023 FIT]		
69 ± 17 ± 12	3182 ± 766	UEHARA	08	BELL $\gamma\gamma \rightarrow 2(\pi^+ \pi^-)$

NODE=M026G19
 NODE=M026G19
 NEW

$$\Gamma(f_2(1270)f_2'(1525)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{19}\Gamma_{59}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
49 ± 9 ± 13	1128 ± 206	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

NODE=M026G20
NODE=M026G20

$$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{37}\Gamma_{59}/\Gamma$$

VALUE (keV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.360 ± 0.022 OUR FIT				Error includes scale factor of 1.5. [0.378 ± 0.021 keV OUR 2023 FIT]

NODE=M026G14
NODE=M026G14

NEW

0.396 ± 0.016 OUR AVERAGE

[0.407 ± 0.027 keV OUR 2023 AVERAGE Scale factor = 1.2]

0.386 ± 0.008 ± 0.021	12k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
0.374 ± 0.009 ± 0.031	14k	¹ LEES	10 BABR	$10.6 e^+e^- \rightarrow e^+e^- K_S^0 K^\pm \pi^\mp$
0.407 ± 0.022 ± 0.028		^{2,3} ASNER	04 CLEO	$\gamma\gamma \rightarrow \eta_c' \rightarrow K_S^0 K^\pm \pi^\mp$
0.60 ± 0.12 ± 0.09	41	^{3,4} ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
1.47 ± 0.87 ± 0.27		³ SHIRAI	98 AMY	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
0.84 ± 0.21		³ ALBRECHT	94H ARG	$\gamma\gamma \rightarrow K^\pm K_S^0 \pi^\mp$
0.60 ^{+0.23} _{-0.20}		³ CHEN	90B CLEO	$\gamma\gamma \rightarrow \eta_c K^\pm K_S^0 \pi^\mp$
1.06 ± 0.41 ± 0.27	11	³ BRAUNSCH...	89 TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$
1.5 ^{+0.60} _{-0.45} ± 0.3	7	³ BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.418 ± 0.044 ± 0.022		^{3,5} BRANDENB...	00B CLE2	$\gamma\gamma \rightarrow \eta_c \rightarrow K^\pm K_S^0 \pi^\mp$
<0.63	95	³ BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
<4.4	95	ALTHOFF	85B TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$

¹ From the corrected and unfolded mass spectrum.

² Calculated by us from the value reported in ASNER 04 that assumes $B(\eta_c \rightarrow K\bar{K}\pi) = 5.5 \pm 1.7\%$

³ We have multiplied $K^\pm K_S^0 \pi^\mp$ measurement by 3 to obtain $K\bar{K}\pi$.

⁴ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (1.5 \pm 0.4)\%$.

⁵ Superseded by ASNER 04.

NODE=M026G14;LINKAGE=LE
NODE=M026G14;LINKAGE=AA

NODE=M026G14;LINKAGE=C
NODE=M026G;LINKAGE=BB

NODE=M026G14;LINKAGE=NN

$$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{41}\Gamma_{59}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
42 ± 9 OUR FIT				Error includes scale factor of 2.1. [35 ± 5 eV OUR 2023 FIT]
27 ± 6 OUR AVERAGE				

NODE=M026G15
NODE=M026G15

NEW

25.7 ± 3.2 ± 4.9	2019 ± 248	UEHARA	08 BELL	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
280 ± 100 ± 60	42	¹ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$
170 ± 80 ± 20	13.9 ± 6.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow \pi^+\pi^-K^+K^-$

¹ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow \pi^+\pi^-K^+K^-) = (2.0 \pm 0.7)\%$.

NODE=M026G;LINKAGE=CC

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{59}/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.190 ± 0.006 ± 0.028	11k	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M026G02
NODE=M026G02

$$\Gamma(2(K^+K^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{45}\Gamma_{59}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 2.1 OUR FIT				Error includes scale factor of 1.5. [7.4 ± 1.5 eV OUR 2023 FIT]
5.8 ± 1.9 OUR AVERAGE				

NODE=M026G27
NODE=M026G27

NEW

5.6 ± 1.1 ± 1.6	216 ± 42	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$
350 ± 90 ± 60	46	¹ ABDALLAH	03J DLPH	$\gamma\gamma \rightarrow 2(K^+K^-)$
231 ± 90 ± 23	9.1 ± 3.3	² ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(K^+K^-)$

¹ Calculated by us from the value reported in ABDALLAH 03J, which uses $B(\eta_c \rightarrow 2(K^+K^-)) = (2.1 \pm 1.2)\%$.

NODE=M026G;LINKAGE=DD

² Includes all topological modes except $\eta_c \rightarrow \phi\phi$.

NODE=M026G;LINKAGE=EE

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{48}\Gamma_{59}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
48 ± 7 OUR FIT	Error includes scale factor of 1.5. [47 ± 5 eV OUR 2023 FIT]			
42 ± 6 OUR AVERAGE				
40.7 ± 3.7 ± 5.3	5381 ± 492	UEHARA	08 BELL	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$
180 ± 70 ± 20	21.4 ± 8.6	ALBRECHT	94H ARG	$\gamma\gamma \rightarrow 2(\pi^+\pi^-)$

NODE=M026G11
 NODE=M026G11
 NEW

$$\Gamma(\rho\bar{\rho}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_{59}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7 ± 0.6 OUR FIT	Error includes scale factor of 1.1. [7.3 ± 0.7 eV OUR 2023 FIT]			
6.2 $\begin{smallmatrix} +1.1 \\ -1.0 \end{smallmatrix}$ OUR AVERAGE	Error includes scale factor of 1.1. [7.2 ± 1.7 eV OUR 2023 AVERAGE]			
7.20 ± 1.53 $\begin{smallmatrix} +0.67 \\ -0.75 \end{smallmatrix}$	157 ± 33	¹ KUO	05 BELL	$\gamma\gamma \rightarrow \rho\bar{\rho}$
4.6 $\begin{smallmatrix} +1.3 \\ -1.1 \end{smallmatrix}$ ± 0.4	190	AMBROGIANI	03 E835	$\bar{p}p \rightarrow \gamma\gamma$
8.1 $\begin{smallmatrix} +2.9 \\ -2.0 \end{smallmatrix}$		ARMSTRONG	95F E760	$\bar{p}p \rightarrow \gamma\gamma$

NODE=M026G01
 NODE=M026G01
 NEW
 NEW

¹ Not independent from the $\Gamma_{\gamma\gamma}$ reported by the same experiment.

NODE=M026G01;LINKAGE=GG

$\eta_c(1S)$ BRANCHING RATIOS

HADRONIC DECAYS

$$\Gamma(\eta'(958)K\bar{K})/\Gamma(\eta'(958)\pi\pi) \quad \Gamma_2/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.859 ± 0.052 ± 0.043	¹ LEES	21A BABR	$\gamma\gamma \rightarrow \eta' K^+ K^-$, $\eta' \pi^+ \pi^-$

NODE=M026R55
 NODE=M026R55

¹ Based on Dalitz-plot analysis of the $\eta_c \rightarrow \eta' K^+ K^-$, $\eta' \pi^+ \pi^-$ final states where the fit fractions and relative phases are determined for numerous two-body intermediate states.

NODE=M026R55;LINKAGE=A

$$\Gamma(\eta'(958)\eta\eta)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.4 ± 0.5 ± 0.3	¹ ABLIKIM	21C BES3	$J/\psi(1S) \rightarrow \gamma\eta\eta\eta'$

NODE=M026R63
 NODE=M026R63

¹ ABLIKIM 21C reports $[\Gamma(\eta_c(1S) \rightarrow \eta'(958)\eta\eta)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (4.86 \pm 0.62 \pm 0.45) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R63;LINKAGE=A

$$\Gamma(\rho\rho)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.1 ± 0.5 ± 0.1		72	¹ ABLIKIM	05L BES2	$J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$
2.3 ± 0.5 ± 0.2		113	^{2,3} BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^0\rho^0$
2.1 ± 1.0 ± 0.2		32	^{4,5} BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^+\rho^-$
<14		90	⁶ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$

NODE=M026R9
 NODE=M026R9

OCCUR=2

¹ ABLIKIM 05L reports $[\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.6 \pm 0.6 \pm 0.4) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R9;LINKAGE=F

² BISELLO 91 reports $[\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.30 \pm 0.30 \pm 0.60) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R9;LINKAGE=A

³ The value reported by BISELLO 91 has been multiplied by 3 to account for isospin symmetry.

NODE=M026R9;LINKAGE=B

⁴ BISELLO 91 reports $[\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (3.0 \pm 1.3 \pm 0.6) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R9;LINKAGE=C

⁵ The value reported by BISELLO 91 has been multiplied by 3/2 to account for isospin symmetry.

NODE=M026R9;LINKAGE=D

⁶ Using $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R9;LINKAGE=G

$\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.8 \pm 0.4 \pm 0.2$	63	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-----------------------	----	-----------------------------	------	------------------------------------

¹BALTRUSAITIS 86 reports $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (2.6 \pm 0.6) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R16
NODE=M026R16

NODE=M026R16;LINKAGE=A

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$135 \pm 57 \pm 13$	45	¹ ABLIKIM	06A BES2	$J/\psi \rightarrow K^{*0} \bar{K}^{*0} \pi^+ \pi^- \gamma$
---------------------	----	----------------------	----------	---

¹ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 \bar{K}^*(892)^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.91 \pm 0.64 \pm 0.48) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R25
NODE=M026R25

NODE=M026R25;LINKAGE=AB

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.9^{+0.9}_{-0.8} \pm 1.1$	$14.1^{+4.4}_{-3.7}$	¹ HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
-----------------------------	----------------------	--------------------	---------	--------------------------------------

¹Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12^{+0.10}_{-0.12}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R21
NODE=M026R21

NODE=M026R;LINKAGE=BB

 $\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<40	90	¹ ABLIKIM	06A BES2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-) \gamma$
-----	----	----------------------	----------	---

¹ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow \phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] < 0.603 \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R26
NODE=M026R26

NODE=M026R26;LINKAGE=AB

 $\Gamma(a_0(980) \pi)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

seen		AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
------	--	------	-----------	------------------------------------

seen		LEES	21A BABR	Dalitz anal. of $\eta_c \rightarrow \pi^+ \pi^- \eta$
------	--	------	----------	---

seen		LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$
------	--	------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02	90	^{1,2} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-------	----	-------------------------------	------	------------------------------------

¹The quoted branching ratio uses $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

²We are assuming $B(a_0(980) \rightarrow \eta \pi) > 0.5$.

NODE=M026R11
NODE=M026R11

NODE=M026R11;LINKAGE=E
NODE=M026R11;LINKAGE=F

 $\Gamma(a_2(1320) \pi)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

seen	[<0.02 (CL = 90%)]	OUR 2022 BEST LIMIT]		
------	--------------------	----------------------	--	--

seen		LEES	21A BABR	Dalitz anal. of $\eta_c \rightarrow \pi^+ \pi^- \eta$
------	--	------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02	90	¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-------	----	-----------------------------	------	------------------------------------

¹The quoted branching ratio uses $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R12
NODE=M026R12

NODE=M026R12;LINKAGE=E

 $\Gamma(K^*(892) \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.0128	90	BISELLO	91 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
---------	----	---------	--------	---

<0.0132	90	¹ BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
---------	----	----------------------	--------	---

¹The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R17
NODE=M026R17

OCCUR=2

NODE=M026R17;LINKAGE=E

$\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
seen		LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta$

NODE=M026R13
NODE=M026R13

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011 90 ¹ BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_C \gamma$

¹ The quoted branching ratio uses $B(J/\psi(1S) \rightarrow \gamma \eta_C(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R13;LINKAGE=E

 $\Gamma(f_2(1270)\eta')/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta'$; $K^+K^-\eta'$

NODE=M026R60
NODE=M026R60

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5×10^{-4}	90	¹ ABLIKIM	17P BES3	$J/\psi \rightarrow \pi^+\pi^-\pi^0 K^+K^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<17 $\times 10^{-4}$ 90 ² ABLIKIM 05L BES2 $J/\psi \rightarrow \pi^+\pi^-\pi^0 K^+K^-\gamma$

¹ Using $B(J/\psi \rightarrow \gamma \eta_C) = 0.017 \pm 0.004$.

² The quoted branching ratio uses $B(J/\psi(1S) \rightarrow \gamma \eta_C(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R22
NODE=M026R22

NODE=M026R22;LINKAGE=A
NODE=M026R22;LINKAGE=E

 $\Gamma(f_0(500)\eta)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta$

NODE=M026R57
NODE=M026R57

 $\Gamma(f_0(500)\eta')/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C(1S) \rightarrow \pi^+\pi^-\eta'$

NODE=M026R58
NODE=M026R58

 $\Gamma(f_0(980)\eta)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta$
seen	LEES	14E BABR	Dalitz anal. of $\eta_C \rightarrow K^+K^-\eta$

NODE=M026R41
NODE=M026R41

 $\Gamma(f_0(980)\eta')/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta'$; $K^+K^-\eta'$

NODE=M026R56
NODE=M026R56

 $\Gamma(f_0(1500)\eta)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta$
seen	LEES	14E BABR	Dalitz anal. of $\eta_C \rightarrow K^+K^-\eta$

NODE=M026R42
NODE=M026R42

 $\Gamma(f_0(1710)\eta')/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow K^+K^-\eta'$

NODE=M026R59
NODE=M026R59

 $\Gamma(a_2(1700)\pi)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M026P10
NODE=M026P10

 $\Gamma(f_0(2100)\eta')/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	21A BABR	Dalitz anal. of $\eta_C \rightarrow \pi^+\pi^-\eta$

NODE=M026R61
NODE=M026R61

 $\Gamma(f_0(2200)\eta)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	LEES	14E BABR	Dalitz anal. of $\eta_C \rightarrow K^+K^-\eta$

NODE=M026R43
NODE=M026R43

 $\Gamma(a_0(1320)\pi)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
seen	LEES	14E BABR	Dalitz anal. of $\eta_C \rightarrow K^+K^-\pi^0$

NODE=M026R45
NODE=M026R45

$\Gamma(a_0(1450)\pi)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$
seen	LEES	21A BABR	Dalitz anal. of $\eta_c \rightarrow \pi^+ \pi^- \eta$
seen	LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

NODE=M026R46
 NODE=M026R46

 $\Gamma(a_0(1710)\pi)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$
seen	LEES	21A BABR	Dalitz anal. of $\eta_c \rightarrow \pi^+ \pi^- \eta'$

NODE=M026P13
 NODE=M026P13

 $\Gamma(a_0(1950)\pi)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	EPTS	DOCUMENT ID	TECN	COMMENT
seen		LEES	21A BABR	Dalitz anal. of $\eta_c(1S) \rightarrow \pi^+ \pi^- \eta'$
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$

NODE=M026R00
 NODE=M026R00

¹ From a model-independant partial wave analysis.

NODE=M026R00;LINKAGE=A

 $\Gamma(K_0^*(1430)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	EPTS	DOCUMENT ID	TECN	COMMENT
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
seen		LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta/\pi^0$

NODE=M026R47
 NODE=M026R47

¹ From a model-independant partial wave analysis.

NODE=M026R47;LINKAGE=A

 $\Gamma(K_2^*(1430)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$
seen	LEES	21A BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta'$
seen	LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \pi^0$

NODE=M026R48
 NODE=M026R48

 $\Gamma(K_0^*(1950)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	EPTS	DOCUMENT ID	TECN	COMMENT
seen		AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$
seen		LEES	21A BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta'$
seen	12k	¹ LEES	16A BABR	$\gamma\gamma \rightarrow \eta_c(1S) \rightarrow K\bar{K}\pi$
seen		LEES	14E BABR	Dalitz anal. of $\eta_c \rightarrow K^+ K^- \eta/\pi^0$

NODE=M026R49
 NODE=M026R49

¹ From a Dalitz plot analysis using an isobar model.

NODE=M026R49;LINKAGE=A

 $\Gamma(K_0^*(2600)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K\pi)$

NODE=M026P11
 NODE=M026P11

 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-2})	EPTS	DOCUMENT ID	TECN	COMMENT
7.1±0.4 OUR FIT				Error includes scale factor of 1.1. [(7.0 ± 0.4) × 10 ⁻² OUR 2023 FIT]

NODE=M026R4
 NODE=M026R4

NEW

7.4±0.6 OUR AVERAGE

[(6.7 ± 0.5) × 10⁻² OUR 2023 AVERAGE Scale factor = 1.1]

NEW

6.9±0.7±0.6	146	¹ ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$
7.8±0.6±0.6	267	² ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$

OCCUR=2

••• We do not use the following data for averages, fits, limits, etc. •••

6.1±1.2±0.6	55	^{3,4} ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^0$
7.6±1.3±0.8	107	^{5,6} ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\pm$
8.5±1.8		⁷ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c\bar{c}}$
4.7±1.2±0.5	0.6k	^{8,9} BAI	04 BES	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$
6.2±1.7±0.6	33	^{10,11} BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
4.9±1.2±0.5	68	^{12,13} BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$
4.8±1.7	95	^{14,15} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
5.5±2.1±0.5	32	^{16,17} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
4.0±1.1±0.4	63	^{18,19} BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

13 $^{+7}_{-5}$ ±2	20	HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$
--------------------	----	-------	----------	--------------------------------------

< 10.7 90% CL	15	PARTRIDGE	80B CBAL	$J/\psi \rightarrow \eta_c \gamma$
---------------	----	-----------	----------	------------------------------------

- 1 ABLIKIM 19AP quotes $B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.15 \pm 0.12 \pm 0.10) \times 10^{-2}$ which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=C
- 2 ABLIKIM 19AP quotes $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (2.60 \pm 0.21 \pm 0.20) \times 10^{-2}$ which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=F
- 3 ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (4.54 \pm 0.76 \pm 0.48) \times 10^{-6}$ which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=BK
- 4 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (27.24 \pm 4.56 \pm 2.88) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. NODE=M026R4;LINKAGE=CK
- 5 ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (11.35 \pm 1.25 \pm 1.50) \times 10^{-6}$ which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=BL
- 6 ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (34.05 \pm 3.75 \pm 4.50) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values. NODE=M026R4;LINKAGE=CL
- 7 Determined from the ratio of $B(B^\pm \rightarrow K^\pm \eta_c) B(\eta_c \rightarrow K \bar{K} \pi) = (7.4 \pm 0.5 \pm 0.7) \times 10^{-5}$ reported in AUBERT, B 04B and $B(B^\pm \rightarrow K^\pm \eta_c) = (8.7 \pm 1.5) \times 10^{-3}$ reported in AUBERT 06E. NODE=M026R4;LINKAGE=AB
- 8 BAI 04 reports $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (2.2 \pm 0.3 \pm 0.5) \times 10^{-4}$ which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=H
- 9 BAI 04 reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (6.6 \pm 0.9 \pm 1.5) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=I
- 10 BISELLO 91 reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (8.76 \pm 1.80 \pm 1.68) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=J
- 11 BISELLO 91 reports $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.46 \pm 0.30 \pm 0.28) \times 10^{-4}$ which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=M
- 12 BISELLO 91 reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (6.9 \pm 1.2 \pm 1.2) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=L
- 13 BISELLO 91 reports $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (2.3 \pm 0.4 \pm 0.4) \times 10^{-4}$ which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=N
- 14 Average from $K^+ K^- \pi^0$ and $K^\pm K_S^0 \pi^\mp$ decay channels. NODE=M026R4;LINKAGE=D
- 15 The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages. NODE=M026R4;LINKAGE=E
- 16 BALTRUSAITIS 86 reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (7.8 \pm 3.0) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=Q
- 17 BALTRUSAITIS 86 reports $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.3 \pm 0.5) \times 10^{-4}$ which we multiply by 6 to account for isospin symmetry. NODE=M026R4;LINKAGE=S
- 18 BALTRUSAITIS 86 reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (5.7 \pm 1.5) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=R
- 19 BALTRUSAITIS 86 reports $B(J/\psi \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (1.9 \pm 0.5) \times 10^{-4}$ which we multiply by 3 to account for isospin symmetry. NODE=M026R4;LINKAGE=T
- 20 HIMEL 80B reports $[\Gamma(\eta_c(1S) \rightarrow K \bar{K} \pi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \eta_c(1S))] = (4.5_{-1.8}^{+2.4}) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = (3.6 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. NODE=M026R4;LINKAGE=G

 $\Gamma(\phi K^+ K^-) / \Gamma(K \bar{K} \pi)$ Γ_8 / Γ_{37}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.052_{-0.014}^{+0.016} \pm 0.014$	7	1 HUANG	03 BELL	$B^\pm \rightarrow K^\pm \phi \phi$

NODE=M026R02
NODE=M026R02

- 1 Using $B(B^+ \rightarrow \eta_c K^+) = (1.25 \pm 0.12_{-0.12}^{+0.10}) \times 10^{-3}$ from FANG 03 and $B(\eta_c \rightarrow K \bar{K} \pi) = (5.5 \pm 1.7) \times 10^{-2}$.

NODE=M026R02;LINKAGE=BB

$\Gamma(K\bar{K}\eta)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M026R15
NODE=M026R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.9 \pm 0.5 \pm 0.1$ <3.1	7	1,2	ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta K^+ K^-$
	90	3	BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

OCCUR=2

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \eta) = (2.11 \pm 1.01 \pm 0.32) \times 10^{-6}$ which we multiply by 2 to account for isospin symmetry.

NODE=M026R15;LINKAGE=AK

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (4.22 \pm 2.02 \pm 0.64) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R15;LINKAGE=AM

³ The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R15;LINKAGE=E

 $\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$ Γ_{38}/Γ_{37}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M026R40
NODE=M026R40
NEW

0.186 ± 0.018 OUR FIT

[0.187 ± 0.018 OUR 2023 FIT]

0.190 ± 0.008 ± 0.017	5.4k	1	LEES	14E BABR	$\gamma\gamma \rightarrow K^+ K^- \eta/\pi^0$
------------------------------	------	---	------	----------	---

NODE=M026R40;LINKAGE=LE

¹ LEES 14E reports $B(\eta_c(1S) \rightarrow K^+ K^- \eta)/B(\eta_c(1S) \rightarrow K^+ K^- \pi^0) = 0.571 \pm 0.025 \pm 0.051$, which we divide by 3 to account for isospin symmetry. It uses both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M026R6
NODE=M026R6

1.6 ± 0.4 OUR AVERAGE [(1.7 ± 0.5) × 10⁻² OUR 2023 AVERAGE]

NEW

1.6 ± 0.4 ± 0.2	33	1	ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta \pi^+ \pi^-$
------------------------	----	---	---------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.4 ± 2.0	75	2	BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
-----------	----	---	----------------	------	------------------------------------

3.7 ± 1.3 ± 2.0	18	2	PARTRIDGE	80B CBAL	$J/\psi \rightarrow \eta \pi^+ \pi^- \gamma$
-----------------	----	---	-----------	----------	--

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (7.22 \pm 1.47 \pm 1.11) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R6;LINKAGE=AB

² The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

NODE=M026R6;LINKAGE=E

 $\Gamma(\eta 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M026R05
NODE=M026R05

4.3 ± 1.3 OUR AVERAGE [(4.6 ± 1.4) × 10⁻² OUR 2023 AVERAGE]

NEW

4.3 ± 1.2 ± 0.4	39	1	ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta 2(\pi^+ \pi^-)$
------------------------	----	---	---------	----------	---

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (19.17 \pm 3.77 \pm 3.72) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R05;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ Γ_{42}/Γ_{37}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M026R01
NODE=M026R01

0.477 ± 0.017 ± 0.070	11k	1	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
------------------------------	-----	---	-----------------	------	--

¹ We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M026R01;LINKAGE=DE

 $\Gamma(K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M026R06
NODE=M026R06

5.4 ± 1.5 OUR AVERAGE [(5.7 ± 1.6) × 10⁻² OUR 2023 AVERAGE]

NEW

5.4 ± 1.4 ± 0.5	43	1,2	ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\mp 2\pi^\pm$
------------------------	----	-----	---------	----------	--

- ¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^- \pi^- 2\pi^+)$
 $= (12.01 \pm 2.22 \pm 2.04) \times 10^{-6}$ which we multiply by 2 to take c.c. into account.
- ² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + c.c.) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (24.02 \pm 4.44 \pm 4.08) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R06;LINKAGE=AA

NODE=M026R06;LINKAGE=AB

$\Gamma(K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{44} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.4±2.4 OUR AVERAGE

[(7.6 ± 2.4) × 10⁻³ OUR 2023 AVERAGE]

8 ± 4 ± 1	10	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- 2(\pi^+ \pi^-)$
8.6±2.8±0.8	100	² ABLIKIM	06A BES2	$J/\psi \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

NODE=M026R23

NODE=M026R23

NEW

- ¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (3.60 \pm 1.71 \pm 0.64) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R23;LINKAGE=AL

- ² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.21 \pm 0.32 \pm 0.24) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R23;LINKAGE=AB

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{46} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<4 × 10⁻⁴ (CL = 90%) [$<5 \times 10^{-4}$ (CL = 90%) OUR 2023 BEST LIMIT]

<4 × 10 ⁻⁴	90	¹ ABLIKIM	17AJ BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$
-----------------------	----	----------------------	-----------	---

- ¹ ABLIKIM 17AJ reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \eta_c(1S))] < 1.6 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \eta_c(1S)) = 3.6 \times 10^{-3}$.

NODE=M026R51

NODE=M026R51

NODE=M026R51;LINKAGE=A

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{47} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.6±1.0 OUR AVERAGE [(4.8 ± 1.1) × 10⁻² OUR 2023 AVERAGE]

4.6±0.9±0.5	118	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \pi^+ \pi^- 2\pi^0$
-------------	-----	----------------------	----------	--

- ¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (20.31 \pm 2.20 \pm 3.33) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R07

NODE=M026R07

NEW

NODE=M026R07;LINKAGE=AB

$\Gamma(2(\pi^+ \pi^- \pi^0)) / \Gamma_{\text{total}}$ Γ_{49} / Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

15.9±2.0 OUR AVERAGE

[(16.2 ± 2.1) × 10⁻² OUR 2023 AVERAGE]

15.3±1.8±1.8	333	ABLIKIM	19AP BES3	$h_c \rightarrow \gamma \eta_c$
16.8±2.8±1.7	175	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^- \pi^0)$

- ¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^- \pi^0)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma \eta_c(1S))] = (75.13 \pm 7.42 \pm 9.99) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma \eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R08

NODE=M026R08

NEW

NODE=M026R08;LINKAGE=AB

$\Gamma(3(\pi^+ \pi^-)) / \Gamma_{\text{total}}$ Γ_{50} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

18.9±3.4 OUR AVERAGE

[(18 ± 4) × 10⁻³ OUR 2023 AVERAGE]

20 ± 5 ± 2	51	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 3(\pi^+ \pi^-)$
18 ± 4 ± 2	479	² ABLIKIM	06A BES2	$J/\psi \rightarrow 3(\pi^+ \pi^-) \gamma$

NODE=M026R24

NODE=M026R24

NEW

- ¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (8.82 \pm 1.57 \pm 1.59) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.
- ² ABLIKIM 06A reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (2.59 \pm 0.32 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R24;LINKAGE=AL

NODE=M026R24;LINKAGE=AB

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
13.3± 1.1 OUR FIT		Error includes scale factor of 1.1.		$[(13.5 \pm 1.3) \times 10^{-4}$ OUR 2023 FIT]

NODE=M026R2
NODE=M026R2

NEW

12.0± 3.0 OUR AVERAGE		$[(11.5 \pm 1.9) \times 10^{-4}$ OUR 2023 AVERAGE]		
------------------------------	--	--	--	--

NEW

12.0± 2.6±1.5	34	ABLIKIM	19AP BES3	$h_c \rightarrow \gamma\eta_c$
----------------------	----	---------	-----------	--------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

15 ± 5 ± 1	15	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma p\bar{p}$
12.9 ⁺ ₋ 1.8 _{2.1} ± 0.8	195	² WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
13.5 ± 3.0 ± 1.3	213	³ BAI	04 BES	$J/\psi \rightarrow \gamma p\bar{p}$
9.2 ± 3.5 ± 0.9	18	⁴ BISELLO	91 DM2	$J/\psi \rightarrow \gamma p\bar{p}$
10 ± 5 ± 1	23	⁵ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$
22 ⁺ ₋ 11 ± 3		⁶ HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$

- ¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (0.65 \pm 0.19 \pm 0.10) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R2;LINKAGE=AB

- ² WU 06 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (1.42 \pm 0.11⁺_{-0.20}) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.10 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=WU

- ³ BAI 04 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.9 \pm 0.3 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=C

- ⁴ BISELLO 91 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (0.13 \pm 0.04 \pm 0.03) \times 10^{-4}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=D

- ⁵ BALTRUSAITIS 86 reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] = (1.4 \pm 0.7) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=F

- ⁶ HIMEL 80B reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\eta_c(1S))] = (8⁺₋₄) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = (3.6 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R2;LINKAGE=G

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$ $\Gamma_{51}/\Gamma \times \Gamma_9/\Gamma$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
0.24±0.07 OUR FIT			Error includes scale factor of 1.9. $[(0.214 \pm 0.035) \times 10^{-5}$ OUR 2023 FIT]

NODE=M026R33
NODE=M026R33

NEW

4.0⁺₋ 3.5_{3.2}	BAGLIN	89	SPEC $\bar{p}p \rightarrow K^+K^-K^+K^-$
--	--------	----	--

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.12 ± 0.03	14	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0\gamma p\bar{p}\pi^0$

NODE=M026R09
NODE=M026R09

• • • We do not use the following data for averages, fits, limits, etc. • • •

- ¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] \times [B(h_c(1P) \rightarrow \gamma\eta_c(1S))] = (1.53 \pm 0.49 \pm 0.23) \times 10^{-6}$ which we divide by our best values $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$, $B(h_c(1P) \rightarrow \gamma\eta_c(1S)) = (60 \pm 4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M026R09;LINKAGE=AB

$\Gamma(K^+ \bar{p} \Lambda + c.c.) / \Gamma_{\text{total}}$ Γ_{55} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$2.46^{+0.33}_{-0.32} \pm 0.16$	157	¹ LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$
---------------------------------	-----	-----------------	----	--

¹ LU 19 reports $(2.83^{+0.36}_{-0.34} \pm 0.35) \times 10^{-3}$ from a measurement of $[\Gamma(\eta_c(1S) \rightarrow K^+ \bar{p} \Lambda + c.c.) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)]$ assuming $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \eta_c K^+) = (1.10 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R53
NODE=M026R53

NODE=M026R53;LINKAGE=A

 $\Gamma(\bar{\Lambda}(1520) \Lambda + c.c.) / \Gamma_{\text{total}}$ Γ_{56} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.0 ± 1.3 OUR AVERAGE		[(3.1 ± 1.3) × 10 ⁻³ OUR 2023 AVERAGE]		
------------------------------	--	---	--	--

3.0 ± 1.3 ± 0.2	43	¹ LU	19	BELL $B^+ \rightarrow \bar{p} \Lambda K^+ K^+$
------------------------	----	-----------------	----	--

¹ LU 19 reports $(3.48 \pm 1.48 \pm 0.46) \times 10^{-3}$ from a measurement of $[\Gamma(\eta_c(1S) \rightarrow \bar{\Lambda}(1520) \Lambda + c.c.) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)]$ assuming $B(B^+ \rightarrow \eta_c K^+) = (9.6 \pm 1.1) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \eta_c K^+) = (1.10 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R54
NODE=M026R54

NEW

NODE=M026R54;LINKAGE=A

 $\Gamma(\Sigma^+ \bar{\Sigma}^-) / \Gamma_{\text{total}}$ Γ_{57} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.6 \pm 0.4 \pm 0.2$	112	¹ ABLIKIM	13C	BES3 $J/\psi \rightarrow \gamma p \bar{p} \pi^0 \pi^0$
-----------------------	-----	----------------------	-----	--

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Sigma^+ \bar{\Sigma}^-) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (3.60 \pm 0.48 \pm 0.31) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R28
NODE=M026R28

NODE=M026R28;LINKAGE=AB

 $\Gamma(\Xi^- \bar{\Xi}^+) / \Gamma_{\text{total}}$ Γ_{58} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.07 \pm 0.22 \pm 0.10$	78	¹ ABLIKIM	13C	BES3 $J/\psi \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$
--------------------------	----	----------------------	-----	--

¹ ABLIKIM 13C reports $[\Gamma(\eta_c(1S) \rightarrow \Xi^- \bar{\Xi}^+) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.51 \pm 0.27 \pm 0.14) \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R29
NODE=M026R29

NODE=M026R29;LINKAGE=AB

————— RADIATIVE DECAYS —————

 $\Gamma(\gamma\gamma) / \Gamma_{\text{total}}$ Γ_{59} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.66 ± 0.13 OUR FIT			Error includes scale factor of 1.2. [(1.68 ± 0.12) × 10 ⁻⁴ OUR 2023 FIT]		
----------------------------	--	--	---	--	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

$3.2 \pm 1.0 \pm 0.3$		¹ ABLIKIM	13i	BES3	
$0.9^{+1.9}_{-0.8} \pm 0.1$	$1.2^{+2.8}_{-1.1}$	² ADAMS	08	CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
$2.0^{+0.9}_{-0.7} \pm 0.1$	13	³ WICHT	08	BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$
$2.80^{+0.67}_{-0.58} \pm 1.0$		⁴ ARMSTRONG	95F	E760	$\bar{p} p \rightarrow \gamma\gamma$
< 9	90	⁵ BISELLO	91	DM2	$J/\psi \rightarrow \gamma\gamma\gamma$
$6^{+4}_{-3} \pm 4$		⁴ BAGLIN	87B	SPEC	$\bar{p} p \rightarrow \gamma\gamma$
< 18	90	⁶ BLOOM	83	CBAL	$J/\psi \rightarrow \eta_c \gamma$

¹ ABLIKIM 13i reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (4.5 \pm 1.2 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ADAMS 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) / \Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma \eta_c(1S))] = (1.2^{+2.7}_{-1.1} \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) = (1.41 \pm 0.14) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026310

NODE=M026R31
NODE=M026R31

NEW

NODE=M026R31;LINKAGE=AL

NODE=M026R31;LINKAGE=AD

³WICHT 08 reports $[\Gamma(\eta_c(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c K^+)] = (2.2^{+0.9+0.4}_{-0.7-0.2}) \times 10^{-7}$ which we divide by our best value $B(B^+ \rightarrow \eta_c K^+) = (1.10 \pm 0.07) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴Not independent from the values of the total and two-photon width quoted by the same experiment.

⁵The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

⁶Using $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

NODE=M026R31;LINKAGE=W1

NODE=M026R31;LINKAGE=AB

NODE=M026R31;LINKAGE=E

NODE=M026R31;LINKAGE=C

 $\Gamma(\rho\rho)/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{51}/\Gamma \times \Gamma_{59}/\Gamma$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
0.221 ± 0.019 OUR FIT				Error includes scale factor of 1.2. $[(0.228 \pm 0.022) \times 10^{-6}$ OUR 2023 FIT]
0.26 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.4.

NODE=M026R32

NODE=M026R32

NEW

0.224 ^{+0.038} _{-0.037} ± 0.020	190	AMBROGIANI 03	E835	$\bar{p}p \rightarrow \eta_c \rightarrow \gamma\gamma$
---	-----	---------------	------	--

0.336 ^{+0.080} _{-0.070}		ARMSTRONG 95F	E760	$\bar{p}p \rightarrow \gamma\gamma$
---	--	---------------	------	-------------------------------------

0.68 ^{+0.42} _{-0.31}	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$
--	----	--------	----------	-------------------------------------

NODE=M026R32

NODE=M026R32

NEW

————— Charge conjugation (C), Parity (P), —————
 ————— Lepton family number (LF) violating modes —————

NODE=M026320

 $\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<13 (CL = 90%)				$[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.82 \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R34

NODE=M026R34

<13	90	¹ ABLIKIM	11G BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-$
---------------	----	----------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<80	90	² ABLIKIM	06B BES2	$J/\psi \rightarrow \pi^+\pi^-\gamma$
---------------	----	----------------------	----------	---------------------------------------

¹ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.82 \times 10^{-6}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R34;LINKAGE=AL

²ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 1.1 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R34;LINKAGE=AB

 $\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 4	90	¹ ABLIKIM	11G BES3	$J/\psi \rightarrow \gamma\pi^0\pi^0$

NODE=M026R35

NODE=M026R35

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	90	² ABLIKIM	06B BES2	$J/\psi \rightarrow \pi^0\pi^0\gamma$
---------------	----	----------------------	----------	---------------------------------------

¹ABLIKIM 11G reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 6.0 \times 10^{-7}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R35;LINKAGE=AL

²ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow \pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.71 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R35;LINKAGE=AB

 $\Gamma(K^+ K^-)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<70 (CL = 90%)				$[\Gamma(\eta_c(1S) \rightarrow K^+ K^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.96 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R36

NODE=M026R36

<70	90	¹ ABLIKIM	06B BES2	$J/\psi \rightarrow K^+ K^- \gamma$
---------------	----	----------------------	----------	-------------------------------------

¹ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.96 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R36;LINKAGE=AB

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<40 (CL = 90%)				$[\Gamma(\eta_c(1S) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.53 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R37

NODE=M026R37

<40	90	¹ ABLIKIM	06B BES2	$J/\psi \rightarrow K_S^0 K_S^0 \gamma$
---------------	----	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<32	90	^{2,3} UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
---------------	----	-----------------------	---------	--

< 5.6	90	^{4,5} UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
-----------------	----	-----------------------	---------	--

OCCUR=2

¹ABLIKIM 06B reports $[\Gamma(\eta_c(1S) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \gamma\eta_c(1S))] < 0.53 \times 10^{-5}$ which we divide by our best value $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 1.41 \times 10^{-2}$.

NODE=M026R37;LINKAGE=AB

²Using $\Gamma(\gamma\gamma)(\eta_c) = 5.3 \pm 0.5$ keV. UEHARA 13 reports $\Gamma(\gamma\gamma) \times B(K_S^0 K_S^0) < 1.6$ eV.

NODE=M026R37;LINKAGE=A

³Taking into account interference with the non-resonant continuum.

NODE=M026R37;LINKAGE=U1

⁴Using $\Gamma(\gamma\gamma)(\eta_c) = 5.3 \pm 0.5$ keV. UEHARA 13 reports $\Gamma(\gamma\gamma) \times B(K_S^0 K_S^0) < 0.29$ eV.

NODE=M026R37;LINKAGE=B

⁵Neglecting interference with the non-resonant continuum.

NODE=M026R37;LINKAGE=U2

$\eta_c(1S)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M026230

$$\Gamma(\eta_c(1S) \rightarrow \eta'(958)\pi\pi)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R64
NODE=M026R64

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.8 ±0.5 OUR FIT Error includes scale factor of 1.4.**5.25±1.65** 14 ¹BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c\gamma$ ¹The value reported by BALTRUSAITIS 86 has been multiplied by 3/2 to account for isospin symmetry.

NODE=M026R64;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow \rho\rho)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_4/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R65
NODE=M026R65

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.6 ±0.6 OUR AVERAGE Error includes scale factor of 1.2.1.6 ±0.6 ±0.4 72 ABLIKIM 05L BES2 $J/\psi \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$ 3.30±0.30±0.60 113 ¹BISELLO 91 DM2 $J/\psi \rightarrow \gamma\rho^0\rho^0$ 3.0 ±1.3 ±0.6 32 ²BISELLO 91 DM2 $J/\psi \rightarrow \gamma\rho^+\rho^-$ ¹The value reported by BISELLO 91 has been multiplied by 3 to account for isospin symmetry.

OCCUR=2

NODE=M026R65;LINKAGE=A

²The value reported by BISELLO 91 has been multiplied by 3/2 to account for isospin symmetry.

NODE=M026R65;LINKAGE=B

$$\Gamma(\eta_c(1S) \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_5/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R66
NODE=M026R66

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.6±0.6 63 BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c\gamma$

$$\Gamma(\eta_c(1S) \rightarrow K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_6/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R67
NODE=M026R67

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.99±0.17 OUR FIT**1.17±0.29 OUR AVERAGE**1.4 ±0.3 ±0.5 60 ABLIKIM 05L BES2 $J/\psi \rightarrow K^+K^-\pi^+\pi^-\gamma$ 1.04±0.36±0.18 14 ¹BISELLO 91 DM2 $e^+e^- \rightarrow \gamma K^+K^-\pi^+\pi^-$ 1.2 ±0.6 9 ¹BALTRUSAIT..86 MRK3 $J/\psi \rightarrow \eta_c\gamma$ ¹The reported value has been multiplied by 2 to account for isospin symmetry.

NODE=M026R67;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow K^*(892)^0\bar{K}^*(892)^0\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_7/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R68
NODE=M026R68

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.91±0.64±0.48 45 ABLIKIM 06A BES2 $J/\psi \rightarrow K^{*0}\bar{K}^{*0}\pi^+\pi^-\gamma$

$$\Gamma(\eta_c(1S) \rightarrow \phi K^+ K^-)/\Gamma_{\text{total}} \times \Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}} \Gamma_8/\Gamma \times \Gamma_{253}^{B^\pm}/\Gamma^{B^\pm}$$

NODE=M026R69
NODE=M026R69

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.6^{+1.1}_{-0.9}±0.8 14.1^{+4.4}_{-3.7} HUANG 03 BELL $B^+ \rightarrow (\phi K^+ K^-) K^+$

$$\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_9/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R80
NODE=M026R80

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.6±0.6 OUR FIT Error includes scale factor of 2.2.**4.1±0.6 OUR AVERAGE** Error includes scale factor of 1.2.4.3±0.5^{+0.5}_{-1.2} 1.2k ABLIKIM 17P BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ 3.3±0.6±0.6 72 ABLIKIM 05L BES2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ 3.9±0.9±0.7 19 BISELLO 91 DM2 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ 3.8^{+2.3}_{-1.5}±0.7 5 BISELLO 91 DM2 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

OCCUR=2

9.3±2.0±1.6 80 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ 8.5±2.7±1.8 BAI 90B MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$

OCCUR=3

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

3.3±0.6±0.6 357 ¹BAI 04 BES $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$ ¹Superseded by ABLIKIM 05L.

NODE=M026R80;LINKAGE=E

$$\Gamma(\eta_c(1S) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}} \quad \Gamma_9/\Gamma \times \Gamma_{253}^{B^\pm}/\Gamma^{B^\pm}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT

2.0±0.5 OUR FIT Error includes scale factor of 2.2.

3.3^{+1.2}_{-1.0} OUR AVERAGE Error includes scale factor of 1.5.

4.7±1.2±0.5		AUBERT,B	04B	BABR	$B^\pm \rightarrow K^\pm \eta_c$
2.2 ^{+1.0} _{-0.7} ±0.5	7	HUANG	03	BELL	$B^\pm \rightarrow K^\pm \phi\phi$

NODE=M026R70
NODE=M026R70

$$\Gamma(\eta_c(1S) \rightarrow \omega\omega)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{16}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

3.7 ±1.2 OUR FIT Error includes scale factor of 2.1.

4.90±0.17±0.77 1705 ABLIKIM 19AV BES3 $J/\psi \rightarrow \gamma\omega\omega$

NODE=M026R71
NODE=M026R71

$$\Gamma(\eta_c(1S) \rightarrow f_2(1270) f_2(1270))/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

1.5±0.4 OUR FIT

1.3±0.3^{+0.3}_{-0.4} 91.2 ± 19.8 ABLIKIM 04M BES $J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M026R72
NODE=M026R72

$$\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

10.1 ±0.9 OUR FIT Error includes scale factor of 1.5.

6.7 ±0.8 OUR AVERAGE

6.6 ±0.9 ±1.5	0.6k	¹ BAI	04	BES	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$
8.76±1.80±1.68	33	² BISELLO	91	DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
6.9 ±1.2 ±1.2	68	³ BISELLO	91	DM2	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$
7.8 ±3.0	32	⁴ BALTRUSAIT..86	MRK3		$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
5.7 ±1.5	63	⁵ BALTRUSAIT..86	MRK3		$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$

NODE=M026R73
NODE=M026R73

OCCUR=2

OCCUR=2

¹ BAI 04 reports $B(J/\psi \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (2.2 \pm 0.3 \pm 0.5) \times 10^{-4}$ which we multiply by 3 to account for isospin symmetry.

NODE=M026R73;LINKAGE=A

² BISELLO 91 reports $B(J/\psi \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.46 \pm 0.30 \pm 0.28) \times 10^{-4}$ which we multiply by 6 to account for isospin symmetry.

NODE=M026R73;LINKAGE=B

³ BISELLO 91 reports $B(J/\psi \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (2.3 \pm 0.4 \pm 0.4) \times 10^{-4}$ which we multiply by 3 to account for isospin symmetry.

NODE=M026R73;LINKAGE=C

⁴ BALTRUSAITIS 86 reports $B(J/\psi \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (1.3 \pm 0.5) \times 10^{-4}$ which we multiply by 6 to account for isospin symmetry.

NODE=M026R73;LINKAGE=D

⁵ BALTRUSAITIS 86 reports $B(J/\psi \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^\pm K_S^0 \pi^\mp) = (1.9 \pm 0.5) \times 10^{-4}$ which we multiply by 3 to account for isospin symmetry.

NODE=M026R73;LINKAGE=E

$$\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} \times \Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma \times \Gamma_{253}^{B^\pm}/\Gamma^{B^\pm}$$

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

7.9 ±0.5 OUR FIT Error includes scale factor of 1.1.

7.5 ±0.8 OUR AVERAGE

8.01±0.42^{+1.71}_{-1.65} ¹ VINOKUROVA 11 BELL e⁺e⁻ → $\Upsilon(4S)$

7.4 ±0.5 ±0.7 AUBERT,B 04B BABR $B^\pm \rightarrow K^\pm \eta_c$

¹ VINOKUROVA 11 reports $B(B^+ \rightarrow \eta_c K^+, \eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (26.7 \pm 1.4 \pm 2.9 \pm 4.9) \times 10^{-6}$, where the first uncertainty is statistical, the second is due to systematics, and the third comes from interference of $\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp$ with nonresonant $K_S^0 K^\pm \pi^\mp$. We combined both systematic uncertainties to single values. We multiply the reported result by 3 to account for isospin symmetry.

NODE=M026R74
NODE=M026R74

NODE=M026R74;LINKAGE=D

$$\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{37}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

2.6±0.4 OUR FIT Error includes scale factor of 1.3.

4.5^{+2.4}_{-1.8} HIMEL 80B MRK2 $\psi(2S) \rightarrow \eta_c \gamma$

NODE=M026R75
NODE=M026R75

$$\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_{37}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R76
NODE=M026R76

VALUE (units 10⁻²) EVTS DOCUMENT ID TECN COMMENT

4.28±0.34 OUR FIT

4.1 ±0.6 OUR AVERAGE

3.7 ±0.7 ±0.3	55	^{1,2} ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^0$
4.6 ±0.8 ±0.3	107	^{3,4} ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\pm$

OCCUR=2

NODE=M026R76;LINKAGE=A

NODE=M026R76;LINKAGE=B

NODE=M026R76;LINKAGE=C

NODE=M026R76;LINKAGE=D

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \pi^0) = (4.54 \pm 0.76 \pm 0.48) \times 10^{-6}$ which we multiply by 6 to account for isospin symmetry.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\pi)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (27.24 \pm 4.56 \pm 2.88) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) = (11.35 \pm 1.25 \pm 1.50) \times 10^{-6}$ which we multiply by 3 to account for isospin symmetry.

⁴ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (34.05 \pm 3.75 \pm 4.50) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_{38}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R77
NODE=M026R77

VALUE (units 10⁻³) EVTS DOCUMENT ID TECN COMMENT

7.9±1.0 OUR FIT

7.9±2.9±0.4	7	^{1,2} ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta K^+ K^-$
-------------	---	------------------------	-----	------	--

¹ ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma\eta_c) \cdot B(\eta_c \rightarrow K^+ K^- \eta) = (2.11 \pm 1.01 \pm 0.32) \times 10^{-6}$ which we multiply by 2 to account for isospin symmetry.

² ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.22 \pm 2.02 \pm 0.64) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R77;LINKAGE=A

NODE=M026R77;LINKAGE=B

$$\Gamma(\eta_c(1S) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_{39}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R78
NODE=M026R78

VALUE (units 10⁻³) EVTS DOCUMENT ID TECN COMMENT

9.7±2.5±0.7	33	¹ ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta \pi^+ \pi^-$
-------------	----	----------------------	-----	------	--

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (7.22 \pm 1.47 \pm 1.11) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R78;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_{39}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R79
NODE=M026R79

VALUE (units 10⁻⁴) EVTS DOCUMENT ID TECN COMMENT

4.2±0.9 OUR AVERAGE

4.6±1.1	75	BALTRUSAIT..86	MRK3	J/ψ → η _c γ
3.1±1.1±1.5	18	PARTRIDGE	80B CBAL	J/ψ → ηπ ⁺ π ⁻ γ

$$\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}$$

$$\Gamma_{40}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R81
NODE=M026R81

VALUE (units 10⁻²) EVTS DOCUMENT ID TECN COMMENT

2.6±0.7±0.2	39	¹ ABLIKIM	12N	BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta 2(\pi^+\pi^-)$
-------------	----	----------------------	-----	------	--

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \eta 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (19.17 \pm 3.77 \pm 3.72) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R81;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{41} / \Gamma \times \Gamma_{237}^{J/\psi(1S)} / \Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.17±0.26 OUR FIT Error includes scale factor of 2.0.

1.9 ±0.6 OUR AVERAGE Error includes scale factor of 2.4.

1.5 ±0.2 ±0.2	0.4k	BAI	04	BES $J/\psi \rightarrow \gamma K^+ K^- \pi^+ \pi^-$
2.7 ±0.4	110	BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

NODE=M026R84
NODE=M026R84

$$\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{41} / \Gamma \times \Gamma_{182}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

3.0±0.8 OUR FIT Error includes scale factor of 1.7.

4.0^{+6.0}_{-2.5} HIMEL 80B MRK2 $\psi(2S) \rightarrow \eta_c \gamma$

NODE=M026R85
NODE=M026R85

$$\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{41} / \Gamma \times \Gamma_{25}^{h_c(1P)} / \Gamma_{h_c(1P)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.0±1.0 OUR FIT Error includes scale factor of 1.7.

5.6±1.3±0.4 38 ¹ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- \pi^+ \pi^-$

¹ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] = (4.16 \pm 0.76 \pm 0.59) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R83
NODE=M026R83

NODE=M026R83;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{43} / \Gamma \times \Gamma_{25}^{h_c(1P)} / \Gamma_{h_c(1P)}$$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

3.2±0.8±0.2 ^{1,2}ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K_S^0 K^\mp \pi^\mp 2\pi^\pm$

¹ABLIKIM 12N quotes $B(\psi(2S) \rightarrow \pi^0 h_c) \cdot B(h_c \rightarrow \gamma \eta_c) \cdot B(\eta_c \rightarrow K_S^0 K^- \pi^- 2\pi^+) = (12.01 \pm 2.22 \pm 2.04) \times 10^{-6}$ which we multiply by 2 to take c.c. into account.

²ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^0 K^- \pi^+ \pi^- \pi^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] = (24.02 \pm 4.44 \pm 4.08) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R86
NODE=M026R86

NODE=M026R86;LINKAGE=A

NODE=M026R86;LINKAGE=B

$$\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{44} / \Gamma \times \Gamma_{237}^{J/\psi(1S)} / \Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.21±0.32±0.24 100 ABLIKIM 06A BES2 $J/\psi \rightarrow K^+ K^- 2(\pi^+ \pi^-) \gamma$

NODE=M026R88
NODE=M026R88

$$\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}$$

$$\Gamma_{44} / \Gamma \times \Gamma_{25}^{h_c(1P)} / \Gamma_{h_c(1P)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.8±2.5±0.3 10 ¹ABLIKIM 12N BES3 $\psi(2S) \rightarrow \pi^0 \gamma K^+ K^- 2(\pi^+ \pi^-)$

¹ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow K^+ K^- 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P) \pi^0)] = (3.60 \pm 1.71 \pm 0.64) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P) \pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R87
NODE=M026R87

NODE=M026R87;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-)) / \Gamma_{\text{total}} \times \Gamma(B^+ \rightarrow \eta_c K^+) / \Gamma_{\text{total}}$$

$$\Gamma_{45} / \Gamma \times \Gamma_{253}^{B^\pm} / \Gamma_{B^\pm}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.6±0.4 OUR FIT Error includes scale factor of 1.4.

1.8^{+0.6}_{-0.5} 14.5^{+4.6}_{-3.0} HUANG 03 BELL $B^+ \rightarrow 2(K^+ K^-) K^+$

NODE=M026R90
NODE=M026R90

$$\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{45}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R89
NODE=M026R89

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
0.85±0.24 OUR FIT		Error includes scale factor of 1.3.		
1.3 ±0.5 ±0.1	7	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(K^+ K^-)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(K^+ K^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (0.94 \pm 0.37 \pm 0.14) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R89;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{47}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R91
NODE=M026R91

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
2.7±0.5±0.2	118	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma \pi^+ \pi^- 2\pi^0$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (20.31 \pm 2.20 \pm 3.33) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R91;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{48}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R93
NODE=M026R93

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
1.35±0.19 OUR FIT		Error includes scale factor of 1.3.		
1.36±0.23 OUR AVERAGE				
1.3 ±0.2 ±0.4	0.5k	BAI	04 BES	$J/\psi \rightarrow \gamma 2(\pi^+ \pi^-)$
1.33±0.22±0.20	137	BISELLO	91 DM2	$J/\psi \rightarrow \gamma 2(\pi^+ \pi^-)$
1.6 ±0.6	25	BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma \eta_c$

$$\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{48}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

NODE=M026R94
NODE=M026R94

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
3.4±0.7 OUR FIT		Error includes scale factor of 1.3.		
5.7+3.9 -2.4		HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$

$$\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{48}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R92
NODE=M026R92

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
0.57±0.09 OUR FIT		Error includes scale factor of 1.3.		
1.01±0.19±0.07	100	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^-)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (7.51 \pm 0.85 \pm 1.11) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R92;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{49}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

NODE=M026R95
NODE=M026R95

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
10.1±1.7±0.7	175	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 2(\pi^+ \pi^- \pi^0)$

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 2(\pi^+ \pi^- \pi^0))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (75.13 \pm 7.42 \pm 9.99) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R95;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \times \Gamma_{50}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

NODE=M026R97
NODE=M026R97

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
2.59±0.32±0.47	471	ABLIKIM	06A BES2	$J/\psi \rightarrow 3(\pi^+ \pi^-) \gamma$

$$\Gamma(\eta_c(1S) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{50}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.19±0.30±0.08	51	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma 3(\pi^+ \pi^-)$

NODE=M026R96
NODE=M026R96

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (8.82 \pm 1.57 \pm 1.59) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R96;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{51}/\Gamma \times \Gamma_{237}^{J/\psi(1S)}/\Gamma_{J/\psi(1S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.88±0.18 OUR FIT				Error includes scale factor of 1.2.
1.61±0.29 OUR AVERAGE				
1.9 ±0.3 ±0.3	213	BAI	04 BES	$J/\psi \rightarrow \gamma p\bar{p}$
1.3 ±0.4 ±0.3	18	BISELLO	91 DM2	$J/\psi \rightarrow \gamma p\bar{p}$
1.4 ±0.7	23	BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$

NODE=M026R99
NODE=M026R99

$$\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{51}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±0.8 OUR FIT				
8.7±2.9±0.6	15	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}$

NODE=M026R98
NODE=M026R98

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (0.65 \pm 0.19 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026R98;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{51}/\Gamma \times \Gamma_{182}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.8±0.7 OUR FIT			Error includes scale factor of 1.2.
8 ⁺⁸/₋₄	HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c \gamma$

NODE=M026P00
NODE=M026P00

$$\Gamma(\eta_c(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(B^+ \rightarrow \eta_c K^+)/\Gamma_{\text{total}} \quad \Gamma_{51}/\Gamma \times \Gamma_{253}^{B^+}/\Gamma_{B^+}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.12 OUR FIT				Error includes scale factor of 1.1.
1.54±0.19 OUR AVERAGE				Error includes scale factor of 1.1.
1.42±0.11 ^{+0.16} / _{-0.20}	195	WU	06 BELL	$B^+ \rightarrow p\bar{p}K^+$
1.8 ^{+0.3} / _{-0.2} ±0.2		AUBERT,B	05L BABR	$e^+ e^- \rightarrow \Upsilon(4S)$

NODE=M026P01
NODE=M026P01

$$\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{52}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.1±0.7±0.1	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}\pi^0$

NODE=M026P02
NODE=M026P02

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (1.53 \pm 0.49 \pm 0.23) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026P02;LINKAGE=A

$$\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}} \quad \Gamma_{53}/\Gamma \times \Gamma_{25}^{h_c(1P)}/\Gamma_{h_c(1P)}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.19±0.30 OUR FIT				
3.1 ±1.0 ±0.2	19	¹ ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma p\bar{p}\pi^+\pi^-$

NODE=M026P07
NODE=M026P07

¹ ABLIKIM 12N reports $[\Gamma(\eta_c(1S) \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(h_c(1P) \rightarrow \gamma \eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (2.30 \pm 0.65 \pm 0.36) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M026P07;LINKAGE=A

DEL-AMO-SA... 11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
VINOKUROVA 11	PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=53927
LEES 10	PR D81 052010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53236
MITCHELL 09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
ADAMS 08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT 08AB	PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52267
UEHARA 08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT 08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE 07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ABLIKIM 06A	PL B633 19	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50987
ABLIKIM 06B	EPJ C45 337	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50988
AUBERT 06E	PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
PDG 06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
WU 06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM 05L	PR D72 072005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50837
AUBERT,B 05L	PR D72 051101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50827
KUO 05	PL B621 41	C.C. Kuo <i>et al.</i>	(BELLE Collab.)	REFID=50801
ABE 04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM 04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
ASNER 04	PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=49745
AUBERT 04D	PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49746
AUBERT,B 04B	PR D70 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50043
BAI 04	PL B578 16	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49620
ABDALLAH 03J	EPJ C31 481	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49625
AMBROGIANI 03	PL B566 45	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=49465
BAI 03	PL B555 174	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49185
FANG 03	PRL 90 071801	F. Fang <i>et al.</i>	(BELLE Collab.)	REFID=49206
HUANG 03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
ABE,K 02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAI 00F	PR D62 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48546
BRANDENB... 00B	PRL 85 3095	G. Brandenburg <i>et al.</i>	(CLEO Collab.)	REFID=48553
ACCIARRI 99T	PL B461 155	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47476
BAI 99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ABREU 98O	PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46553
SHIRAI 98	PL B424 405	M. Shirai <i>et al.</i>	(AMY Collab.)	REFID=46381
ARMSTRONG 95F	PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=44623
ALBRECHT 94H	PL B338 390	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44098
ADRIANI 93N	PL B318 575	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=43670
BISELLO 91	NP B350 1	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41668
BAI 90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
CHEN 90B	PL B243 169	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=41360
BAGLIN 89	PL B231 557	C. Baglin, S. Baird, G. Bassompierre	(R704 Collab.)	REFID=41015
BEHREND 89	ZPHY C42 367	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40732
BRAUNSCH... 89	ZPHY C41 533	W. Braunschweig <i>et al.</i>	(TASSO Collab.)	REFID=40728
AIHARA 88D	PRL 60 2355	H. Aihara <i>et al.</i>	(TPC Collab.)	REFID=40588
BAGLIN 87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BALTRUSAIT... 86	PR D33 629	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22009
BERGER 86	PL 167B 120	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22010
GAISER 86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
ALTHOFF 85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)	REFID=21349
BALTRUSAIT... 84	PRL 52 2126	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+) JP	REFID=22006
BLOOM 83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
HIMEL 80B	PRL 45 1146	T.M. Himel <i>et al.</i>	(SLAC, LBL, UCB)	REFID=22003
PARTRIDGE 80B	PRL 45 1150	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22004

J/ψ(1S)

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M070

J/ψ(1S) MASS

NODE=M070M

NODE=M070M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3096.900 ± 0.006 OUR AVERAGE				
3096.900 ± 0.002 ± 0.006		¹ ANASHIN 15	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3096.89 ± 0.09	502	² ARTAMONOV 00	OLYA	$e^+e^- \rightarrow \text{hadrons}$
3096.91 ± 0.03 ± 0.01		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
3096.95 ± 0.1 ± 0.3	193	BAGLIN 87	SPEC	$\bar{p}p \rightarrow e^+e^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3096.66 ± 0.19 ± 0.02	6.1k	⁴ AAIJ 15Bl	LHCB	$pp \rightarrow J/\psi X$
3096.917 ± 0.010 ± 0.007		AULCHENKO 03	KEDR	$e^+e^- \rightarrow \text{hadrons}$
3097.5 ± 0.3		GRIBUSHIN 96	FMPS	$515 \pi^- \text{Be} \rightarrow 2\mu X$
3098.4 ± 2.0	38k	LEMOIGNE 82	GOLI	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3096.93 ± 0.09	502	⁵ ZHOLENTZ 80	REDE	e^+e^-
3097.0 ± 1		⁶ BRANDELIK 79c	DASP	e^+e^-

¹Supersedes AULCHENKO 03.²Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).³Mass central value and systematic error recalculated by us according to Eq.(16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.⁴From a sample of $\eta_c(1S)$ and J/ψ produced in b -hadron decays. Systematic uncertainties not estimated.⁵Superseded by ARTAMONOV 00.⁶From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.NODE=M070M;LINKAGE=A
NODE=M070M;LINKAGE=AR

NODE=M070M;LINKAGE=NW

NODE=M070M;LINKAGE=B

NODE=M070M;LINKAGE=RZ
NODE=M070M;LINKAGE=F**J/ψ(1S) WIDTH**

NODE=M070W

NODE=M070W

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
92.6 ± 1.7 OUR AVERAGE				Error includes scale factor of 1.1.
92.45 ± 1.40 ± 1.48		¹ ANASHIN 20	KEDR	e^+e^-
96.1 ± 3.2	13k	² ADAMS 06A	CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
84.4 ± 8.9		BAI 95B	BES	e^+e^-
91 ± 11 ± 6		³ ARMSTRONG 93B	E760	$\bar{p}p \rightarrow e^+e^-$
85.5 ^{+6.1} _{-5.8}		⁴ HSUEH 92	RVUE	See Υ mini-review
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
92.94 ± 1.83		^{5,6} ANASHIN 18A	KEDR	e^+e^-
94.1 ± 2.7		⁷ ANASHIN 10	KEDR	$3.097 e^+e^- \rightarrow e^+e^-, \mu^+\mu^-$
93.7 ± 3.5	7.8k	² AUBERT 04	BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$

¹Based on the same dataset as ANASHIN 18A and correlated to the values reported there²Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(e^+e^-) = (5.94 \pm 0.06)\%$ and $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.³The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].⁴Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75B, BRANDELIK 79C.⁵Using $\Gamma(e^+e^-)$ from ANASHIN 18A and $B(J/\psi(1S) \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ from PDG 16.⁶Superseded by ANASHIN 20 that is based on the same dataset.⁷Assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$ and using $\Gamma(e^+e^-)/\Gamma_{\text{total}} = (5.94 \pm 0.06)\%$.

NODE=M070W;LINKAGE=C

NODE=M070W;LINKAGE=AA

NODE=M070W;LINKAGE=AN
NODE=M070W;LINKAGE=A

NODE=M070W;LINKAGE=B

NODE=M070W;LINKAGE=D
NODE=M070W;LINKAGE=AS**J/ψ(1S) DECAY MODES**

NODE=M070215;NODE=M070

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(87.7 ± 0.5) %	DESIG=3
Γ_2 virtual $\gamma \rightarrow \text{hadrons}$	(13.46 ± 0.07) %	DESIG=4
Γ_3 ggg	(64.1 ± 1.0) %	DESIG=249
Γ_4 γgg	(8.8 ± 1.1) %	DESIG=250
Γ_5 e^+e^-	(5.971 ± 0.032) %	DESIG=1
Γ_6 $e^+e^-\gamma$	[a] (8.8 ± 1.4) × 10 ⁻³	DESIG=5
Γ_7 $\mu^+\mu^-$	(5.961 ± 0.033) %	DESIG=2

Decays involving hadronic resonances

				NODE=M070;CLUMP=A
Γ_8	$\rho\pi$	$(1.88 \pm 0.12) \%$	S=2.6	DESIG=20
Γ_9	$\rho^0\pi^0$	$(6.2 \pm 0.6) \times 10^{-3}$		DESIG=21
Γ_{10}	$a_2(1320)^0\pi^+\pi^- \rightarrow$ $2(\pi^+\pi^-)\pi^0$	$(2.8 \pm 0.6) \times 10^{-3}$		DESIG=442
Γ_{11}	$a_2(1320)^+\pi^-\pi^0 + \text{c.c.} \rightarrow$ $2(\pi^+\pi^-)\pi^0$	$(3.7 \pm 0.7) \times 10^{-3}$		DESIG=443
Γ_{12}	$a_2(1320)\rho$	$(1.09 \pm 0.22) \%$		DESIG=43
Γ_{13}	$\eta\pi^+\pi^-$	$(3.8 \pm 0.7) \times 10^{-4}$		DESIG=239
Γ_{14}	$\eta\pi^+\pi^-\pi^0$	$(1.17 \pm 0.20) \%$		DESIG=420
Γ_{15}	$\eta\pi^+\pi^-3\pi^0$	$(4.9 \pm 1.0) \times 10^{-3}$		DESIG=422
Γ_{16}	$\eta\rho$	$(1.93 \pm 0.23) \times 10^{-4}$		DESIG=22
Γ_{17}	$\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow$ $\eta\phi\pi^+\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$		DESIG=287
Γ_{18}	$\eta\phi(2170) \rightarrow$ $\eta K^*(892)^0\bar{K}^*(892)^0$	$< 2.52 \times 10^{-4}$	CL=90%	DESIG=253
Γ_{19}	ηK^+K^-	$(8.6 \pm 3.0) \times 10^{-4}$		DESIG=455
Γ_{20}	$\eta K^\pm K_S^0\pi^\mp$	[b] $(2.2 \pm 0.4) \times 10^{-3}$		DESIG=230
Γ_{21}	$\eta K^*(892)^0\bar{K}^*(892)^0$	$(1.15 \pm 0.26) \times 10^{-3}$		DESIG=252
Γ_{22}	$\rho\eta'(958)$	$(8.1 \pm 0.8) \times 10^{-5}$	S=1.6	DESIG=23
Γ_{23}	$\rho^\pm\pi^\mp\pi^+\pi^-2\pi^0$	$(2.8 \pm 0.8) \%$		DESIG=415
Γ_{24}	$\rho^+\rho^-\pi^+\pi^-\pi^0$	$(6 \pm 4) \times 10^{-3}$		DESIG=416
Γ_{25}	$\rho^+K^+K^-\pi^- + \text{c.c.} \rightarrow$ $K^+K^-\pi^+\pi^-\pi^0$	$(3.5 \pm 0.8) \times 10^{-3}$		DESIG=444
Γ_{26}	$\rho^\mp K^\pm K_S^0$	$(1.9 \pm 0.4) \times 10^{-3}$		DESIG=342
Γ_{27}	$h_1(1415)\eta' \rightarrow \gamma\eta\eta'$			DESIG=435
Γ_{28}	$h_1(1595)\eta' \rightarrow \gamma\eta\eta'$			DESIG=437
Γ_{29}	$\rho(1450)\pi$			DESIG=310
Γ_{30}	$\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0$	$(2.2 \pm 1.1) \times 10^{-4}$		DESIG=328
Γ_{31}	$\rho(1450)^\pm\pi^\mp \rightarrow K_S^0 K^\pm\pi^\mp$	$(3.3 \pm 0.6) \times 10^{-4}$		DESIG=329
Γ_{32}	$\rho(1450)^0\pi^0 \rightarrow K^+K^-\pi^0$	$(2.7 \pm 0.6) \times 10^{-4}$		DESIG=312
Γ_{33}	$\rho(1450)\eta'(958) \rightarrow$ $\pi^+\pi^-\eta'(958)$	$(3.3 \pm 0.7) \times 10^{-6}$		DESIG=345
Γ_{34}	$\rho(1700)\pi$			DESIG=325
Γ_{35}	$\rho(1700)\pi \rightarrow \pi^+\pi^-\pi^0$	$(1.6 \pm 1.1) \times 10^{-4}$		DESIG=313
Γ_{36}	$\rho(2150)\pi$			DESIG=326
Γ_{37}	$\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$	$(10 \pm 40) \times 10^{-6}$		DESIG=314
Γ_{38}	$\rho_3(1690)\pi \rightarrow \pi^+\pi^-\pi^0$			DESIG=316
Γ_{39}	$\omega\pi^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.4	DESIG=32
Γ_{40}	$\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0$	$(1.6 \pm 0.7) \times 10^{-5}$		DESIG=327
Γ_{41}	$\omega\pi^+\pi^-$	$(8.5 \pm 1.0) \times 10^{-3}$	S=1.3	DESIG=24
Γ_{42}	$\omega\pi^0\pi^0$	$(3.4 \pm 0.8) \times 10^{-3}$		DESIG=140
Γ_{43}	$\omega 3\pi^0$	$(1.9 \pm 0.6) \times 10^{-3}$		DESIG=421
Γ_{44}	$\omega f_2(1270)$	$(4.3 \pm 0.6) \times 10^{-3}$		DESIG=28
Γ_{45}	$\omega\eta$	$(1.74 \pm 0.20) \times 10^{-3}$	S=1.6	DESIG=30
Γ_{46}	$\omega\pi^+\pi^-\pi^0$	$(4.0 \pm 0.7) \times 10^{-3}$		DESIG=211
Γ_{47}	$\omega\pi^0\eta$	$(3.4 \pm 1.7) \times 10^{-4}$		DESIG=360
Γ_{48}	$\omega\pi^+\pi^+\pi^-\pi^-$	$(8.5 \pm 3.4) \times 10^{-3}$		DESIG=26
Γ_{49}	$\omega\pi^+\pi^-2\pi^0$	$(3.3 \pm 0.5) \%$		DESIG=412
Γ_{50}	$\omega\eta'\pi^+\pi^-$	$(1.12 \pm 0.13) \times 10^{-3}$		DESIG=385
Γ_{51}	$\omega\eta'(958)$	$(1.89 \pm 0.18) \times 10^{-4}$		DESIG=31
Γ_{52}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$		DESIG=150
Γ_{53}	$\omega f_0(1710) \rightarrow \omega K\bar{K}$	$(4.8 \pm 1.1) \times 10^{-4}$		DESIG=130
Γ_{54}	$\omega f_1(1420)$	$(6.8 \pm 2.4) \times 10^{-4}$		DESIG=105
Γ_{55}	$\omega f_2'(1525)$	$< 2.2 \times 10^{-4}$	CL=90%	DESIG=29
Γ_{56}	$\omega X(1835) \rightarrow \omega p\bar{p}$	$< 3.9 \times 10^{-6}$	CL=95%	DESIG=263
Γ_{57}	$\omega X(1835), X \rightarrow \eta'\pi^+\pi^-$	$< 6.2 \times 10^{-5}$		DESIG=386
Γ_{58}	ωK^+K^-	$(1.52 \pm 0.31) \times 10^{-3}$		DESIG=441
Γ_{59}	$\omega K^\pm K_S^0\pi^\mp$	[b] $(3.4 \pm 0.5) \times 10^{-3}$		DESIG=101
Γ_{60}	$\omega K\bar{K}$	$(1.9 \pm 0.4) \times 10^{-3}$		DESIG=27
Γ_{61}	$\omega K^*(892)\bar{K} + \text{c.c.}$	$(6.1 \pm 0.9) \times 10^{-3}$		DESIG=102

Γ ₆₂	$\eta' K^{*\pm} K^\mp$	$(1.48 \pm 0.13) \times 10^{-3}$		DESIG=355
Γ ₆₃	$\eta' K^{*0} \bar{K}^0 + \text{c.c.}$	$(1.66 \pm 0.21) \times 10^{-3}$		DESIG=357
Γ ₆₄	$\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.}$	$(2.16 \pm 0.31) \times 10^{-4}$		DESIG=353
Γ ₆₅	$\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp$	$(1.51 \pm 0.23) \times 10^{-4}$		DESIG=354
Γ ₆₆	$\eta' h_1(1415) \rightarrow \gamma \eta' \eta'$	$(4.7 \pm 1.1 / 2.0) \times 10^{-7}$		DESIG=430
Γ ₆₇	$\bar{K} K^*(892) + \text{c.c.}$			DESIG=331
Γ ₆₈	$\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(4.8 \pm 0.5) \times 10^{-3}$		DESIG=332
Γ ₆₉	$K^+ K^*(892)^- + \text{c.c.}$	$(6.0 \pm 0.8 / 1.0) \times 10^{-3}$	S=2.9	DESIG=121
Γ ₇₀	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.13 / 0.20) \times 10^{-3}$		DESIG=231
Γ ₇₁	$K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.0 \pm 0.4) \times 10^{-3}$		DESIG=232
Γ ₇₂	$K^0 \bar{K}^*(892)^0 + \text{c.c.}$	$(4.2 \pm 0.4) \times 10^{-3}$		DESIG=122
Γ ₇₃	$K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}$	$(3.2 \pm 0.4) \times 10^{-3}$		DESIG=233
Γ ₇₄	$\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}$	$(5.7 \pm 0.8) \times 10^{-3}$		DESIG=214
Γ ₇₅	$K^*(892)^\pm K^\mp \pi^0$	$(4.1 \pm 1.3) \times 10^{-3}$		DESIG=343
Γ ₇₆	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(2.0 \pm 0.5) \times 10^{-3}$		DESIG=299
Γ ₇₇	$K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-$	$(6.7 \pm 2.2) \times 10^{-4}$		DESIG=300
Γ ₇₈	$K^*(892)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(3.8 \pm 0.5) \times 10^{-3}$		DESIG=445
Γ ₇₉	$K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(6.3 \pm 0.6 / 0.5) \times 10^{-6}$		DESIG=376
Γ ₈₀	$K^*(892)^0 K_S^0 \pi^0$	$(7 \pm 4) \times 10^{-4}$		DESIG=344
Γ ₈₁	$K^*(892)^\pm K^*(700)^\mp$	$(1.1 \pm 1.0 / 0.6) \times 10^{-3}$		DESIG=257
Γ ₈₂	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.3 \pm 0.6) \times 10^{-4}$		DESIG=46
Γ ₈₃	$K^*(892)^\pm K^*(892)^\mp$	$(1.00 \pm 0.22 / 0.40) \times 10^{-3}$		DESIG=256
Γ ₈₄	$K_1(1400)^\pm K^\mp$	$(3.8 \pm 1.4) \times 10^{-3}$		DESIG=132
Γ ₈₅	$K^*(1410) \bar{K} + \text{c.c.}$			DESIG=317
Γ ₈₆	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(7 \pm 4) \times 10^{-5}$		DESIG=330
Γ ₈₇	$K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(8 \pm 5) \times 10^{-5}$		DESIG=318
Γ ₈₈	$K_2^*(1430) \bar{K} + \text{c.c.}$			DESIG=319
Γ ₈₉	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0$	$(1.0 \pm 0.5) \times 10^{-4}$		DESIG=321
Γ ₉₀	$K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp$	$(3.8 \pm 1.0) \times 10^{-4}$		DESIG=320
Γ ₉₁	$\bar{K}_2^*(1430) K + \text{c.c.}$	$< 4.0 \times 10^{-3}$	CL=90%	DESIG=45
Γ ₉₂	$K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(2.69 \pm 0.25 / 0.19) \times 10^{-4}$		DESIG=381
Γ ₉₃	$K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-$	$(2.6 \pm 0.9) \times 10^{-3}$		DESIG=446
Γ ₉₄	$K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}$	$(3.6 \pm 1.8) \times 10^{-3}$		DESIG=301
Γ ₉₅	$\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}$	$(4.67 \pm 0.29) \times 10^{-3}$		DESIG=48
Γ ₉₆	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.}$	$(3.4 \pm 2.9) \times 10^{-3}$		DESIG=303
Γ ₉₇	$K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}$	$(4 \pm 4) \times 10^{-4}$		DESIG=304
Γ ₉₈	$K_2^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3}$	CL=90%	DESIG=47
Γ ₉₉	$\bar{K}_2^*(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(6.9 \pm 0.9) \times 10^{-4}$		DESIG=235
Γ ₁₀₀	$K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(1.10 \pm 0.60 / 0.14) \times 10^{-5}$		DESIG=382

Γ ₁₀₁	$K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0$	$(6.2 \pm_{-1.6}^{+2.9}) \times 10^{-6}$		DESIG=383
Γ ₁₀₂	$K_1(1270)^\pm K^\mp$	$< 3.0 \times 10^{-3}$	CL=90%	DESIG=131
Γ ₁₀₃	$K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0$	$(8.5 \pm 2.5) \times 10^{-7}$		DESIG=377
Γ ₁₀₄	$a_2(1320)^\pm \pi^\mp$	$[b] < 4.3 \times 10^{-3}$	CL=90%	DESIG=42
Γ ₁₀₅	$\phi \pi^0$	$3 \times 10^{-6} \text{ or } 1 \times 10^{-7}$		DESIG=33;OUR EVAL;→ UNCHECKED ←
Γ ₁₀₆	$\phi \pi^+ \pi^-$	$(9.4 \pm 1.5) \times 10^{-4}$	S=1.7	DESIG=34
Γ ₁₀₇	$\phi \pi^0 \pi^0$	$(5.0 \pm 1.0) \times 10^{-4}$		DESIG=76
Γ ₁₀₈	$\phi 2(\pi^+ \pi^-)$	$(1.60 \pm 0.32) \times 10^{-3}$		DESIG=35
Γ ₁₀₉	$\phi \eta$	$(7.4 \pm 0.6) \times 10^{-4}$	S=1.2	DESIG=37
Γ ₁₁₀	$\phi \eta'(958)$	$(4.6 \pm 0.5) \times 10^{-4}$	S=2.2	DESIG=38
Γ ₁₁₁	$\phi \eta \eta'$	$(2.32 \pm 0.17) \times 10^{-4}$		DESIG=387
Γ ₁₁₂	$\phi f_0(980)$	$(3.2 \pm 0.9) \times 10^{-4}$	S=1.9	DESIG=41
Γ ₁₁₃	$\phi f_0(980) \rightarrow \phi \pi^+ \pi^-$	$(2.60 \pm 0.34) \times 10^{-4}$		DESIG=236
Γ ₁₁₄	$\phi f_0(980) \rightarrow \phi \pi^0 \pi^0$	$(1.8 \pm 0.5) \times 10^{-4}$		DESIG=237
Γ ₁₁₅	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(4.5 \pm 1.0) \times 10^{-6}$		DESIG=278
Γ ₁₁₆	$\phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \rho^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-6}$		DESIG=279
Γ ₁₁₇	$\phi f_0(980) \eta \rightarrow \eta \phi \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-4}$		DESIG=229
Γ ₁₁₈	$\phi a_0(980)^0 \rightarrow \phi \eta \pi^0$	$(4.4 \pm 1.4) \times 10^{-6}$		DESIG=258
Γ ₁₁₉	$\phi f_2(1270)$	$(3.2 \pm 0.6) \times 10^{-4}$		DESIG=39
Γ ₁₂₀	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$		DESIG=106
Γ ₁₂₁	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-$	$(9.4 \pm 2.8) \times 10^{-7}$		DESIG=280
Γ ₁₂₂	$\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi 3\pi^0$	$(2.1 \pm 2.2) \times 10^{-7}$		DESIG=281
Γ ₁₂₃	$\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-$	$(2.0 \pm 1.0) \times 10^{-5}$		DESIG=128
Γ ₁₂₄	$\phi f_2'(1525)$	$(8 \pm 4) \times 10^{-4}$	S=2.7	DESIG=40
Γ ₁₂₅	$\phi X(1835) \rightarrow \phi p \bar{p}$	$< 2.1 \times 10^{-7}$	CL=90%	DESIG=291
Γ ₁₂₆	$\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-$	$< 2.8 \times 10^{-4}$	CL=90%	DESIG=288
Γ ₁₂₇	$\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-$	$< 6.13 \times 10^{-5}$	CL=90%	DESIG=289
Γ ₁₂₈	$\phi K \bar{K}$	$(1.77 \pm 0.16) \times 10^{-3}$	S=1.3	DESIG=36
Γ ₁₂₉	$\phi f_0(1710) \rightarrow \phi K \bar{K}$	$(3.6 \pm 0.6) \times 10^{-4}$		DESIG=129
Γ ₁₃₀	$\phi K^+ K^-$	$(8.3 \pm 1.1) \times 10^{-4}$		DESIG=295
Γ ₁₃₁	$\phi K_S^0 K_S^0$	$(5.9 \pm 1.5) \times 10^{-4}$		DESIG=305
Γ ₁₃₂	$\phi K^\pm K_S^0 \pi^\mp$	$[b] (7.2 \pm 0.8) \times 10^{-4}$		DESIG=103
Γ ₁₃₃	$\phi K^*(892) \bar{K} + \text{c.c.}$	$(2.18 \pm 0.23) \times 10^{-3}$		DESIG=104
Γ ₁₃₄	$b_1(1235)^\pm \pi^\mp$	$[b] (3.0 \pm 0.5) \times 10^{-3}$		DESIG=49
Γ ₁₃₅	$b_1(1235)^0 \pi^0$	$(2.3 \pm 0.6) \times 10^{-3}$		DESIG=160
Γ ₁₃₆	$f_2'(1525) K^+ K^-$	$(1.06 \pm 0.35) \times 10^{-3}$		DESIG=308
Γ ₁₃₇	$\Delta(1232)^+ \bar{p}$	$< 1 \times 10^{-4}$	CL=90%	DESIG=112
Γ ₁₃₈	$\Delta(1232)^{++} \bar{p} \pi^-$	$(1.6 \pm 0.5) \times 10^{-3}$		DESIG=70
Γ ₁₃₉	$\Delta(1232)^{++} \Delta(1232)^{--}$	$(1.10 \pm 0.29) \times 10^{-3}$		DESIG=66
Γ ₁₄₀	$\bar{\Sigma}(1385)^0 p K^-$	$(5.1 \pm 3.2) \times 10^{-4}$		DESIG=74
Γ ₁₄₁	$\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.}$	$< 8.2 \times 10^{-6}$	CL=90%	DESIG=111
Γ ₁₄₂	$\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.}$	$[b] (3.0 \pm 0.7) \times 10^{-4}$		DESIG=68
Γ ₁₄₃	$\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.}$	$(3.3 \pm 0.8) \times 10^{-4}$		DESIG=450
Γ ₁₄₄	$\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.}$	$[b] (1.08 \pm 0.06) \times 10^{-3}$		DESIG=67
Γ ₁₄₅	$\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.}$	$(1.25 \pm 0.07) \times 10^{-3}$		DESIG=451
Γ ₁₄₆	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(1.07 \pm 0.08) \times 10^{-3}$		DESIG=309
Γ ₁₄₇	$\Lambda(1520) \bar{\Lambda} + \text{c.c.} \rightarrow \gamma \Lambda \bar{\Lambda}$	$< 4.1 \times 10^{-6}$	CL=90%	DESIG=260
Γ ₁₄₈	$\bar{\Lambda}(1520) \Lambda + \text{c.c.}$	$< 1.80 \times 10^{-3}$	CL=90%	DESIG=364
Γ ₁₄₉	$\Xi^0 \Xi^0$	$(1.17 \pm 0.04) \times 10^{-3}$		DESIG=248
Γ ₁₅₀	$\Xi(1530)^- \Xi^+ + \text{c.c.}$	$(3.18 \pm 0.08) \times 10^{-4}$		DESIG=107
Γ ₁₅₁	$\Xi(1530)^0 \Xi^0$	$(3.2 \pm 1.4) \times 10^{-4}$		DESIG=108
Γ ₁₅₂	$\Theta(1540) \bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.}$	$[c] < 1.1 \times 10^{-5}$	CL=90%	DESIG=205
Γ ₁₅₃	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	$[c] < 2.1 \times 10^{-5}$	CL=90%	DESIG=206
Γ ₁₅₄	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	$[c] < 1.6 \times 10^{-5}$	CL=90%	DESIG=207
Γ ₁₅₅	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	$[c] < 5.6 \times 10^{-5}$	CL=90%	DESIG=208
Γ ₁₅₆	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	$[c] < 1.1 \times 10^{-5}$	CL=90%	DESIG=209

Decays into stable hadrons

NODE=M070;CLUMP=B

Γ ₁₅₇	$2(\pi^+\pi^-\pi^0)$	(4.2 ± 0.4) %	S=2.1	DESIG=9
Γ ₁₅₈	$3(\pi^+\pi^-\pi^0)$	(2.9 ± 0.6) %		DESIG=11
Γ ₁₅₉	$\pi^+\pi^-\pi^0$	(1.9 ± 0.9) %		DESIG=358
Γ ₁₆₀	$\pi^+\pi^-\pi^0$	(6.5 ± 1.3) × 10 ⁻³		DESIG=419
Γ ₁₆₁	$\rho^\pm\pi^\mp\pi^0\pi^0$	(1.41 ± 0.22) %		DESIG=362
Γ ₁₆₂	$\rho^+\rho^-\pi^0$	(6.0 ± 1.1) × 10 ⁻³		DESIG=363
Γ ₁₆₃	$\pi^+\pi^-\pi^0$	(2.00 ± 0.07) %	S=2.0	DESIG=7
Γ ₁₆₄	$2(\pi^+\pi^-\pi^0)$	(1.61 ± 0.20) %		DESIG=210
Γ ₁₆₅	$\pi^+\pi^-\pi^0K^+K^-$	(1.52 ± 0.27) %	S=1.4	DESIG=18
Γ ₁₆₆	$\pi^+\pi^-$	(1.47 ± 0.14) × 10 ⁻⁴		DESIG=6
Γ ₁₆₇	$2(\pi^+\pi^-)$	(3.20 ± 0.25) × 10 ⁻³	S=1.2	DESIG=8
Γ ₁₆₈	$3(\pi^+\pi^-)$	(4.3 ± 0.4) × 10 ⁻³		DESIG=10
Γ ₁₆₉	$2(\pi^+\pi^-)3\pi^0$	(6.2 ± 0.9) %		DESIG=411
Γ ₁₇₀	$4(\pi^+\pi^-)\pi^0$	(9.0 ± 3.0) × 10 ⁻³		DESIG=12
Γ ₁₇₁	$2(\pi^+\pi^-)\eta$	(2.29 ± 0.28) × 10 ⁻³		DESIG=201
Γ ₁₇₂	$3(\pi^+\pi^-)\eta$	(7.2 ± 1.5) × 10 ⁻⁴		DESIG=202
Γ ₁₇₃	$2(\pi^+\pi^-\pi^0)\eta$	(1.6 ± 0.5) × 10 ⁻³		DESIG=418
Γ ₁₇₄	$\pi^+\pi^-\pi^0\pi^0\eta$	(2.4 ± 0.5) × 10 ⁻³		DESIG=359
Γ ₁₇₅	$\rho^\pm\pi^\mp\pi^0\eta$	(1.9 ± 0.8) × 10 ⁻³		DESIG=361
Γ ₁₇₆	K^+K^-	(2.86 ± 0.21) × 10 ⁻⁴		DESIG=13
Γ ₁₇₇	$K_S^0K_L^0$	(1.95 ± 0.11) × 10 ⁻⁴	S=2.4	DESIG=75
Γ ₁₇₈	$K_S^0K_S^0$	< 1.4 × 10 ⁻⁸	CL=95%	DESIG=14
Γ ₁₇₉	$K\bar{K}\pi$	(6.1 ± 1.0) × 10 ⁻³		DESIG=15
Γ ₁₈₀	$K^+K^-\pi^0$	(2.88 ± 0.12) × 10 ⁻³		DESIG=334
Γ ₁₈₁	$K_S^0K^\pm\pi^\mp$	(5.3 ± 0.5) × 10 ⁻³		DESIG=335
Γ ₁₈₂	$K_S^0K_L^0\pi^0$	(2.06 ± 0.26) × 10 ⁻³		DESIG=336
Γ ₁₈₃	$K^*(892)^0\bar{K}^0 + c.c. \rightarrow K_S^0K_L^0\pi^0$	(1.21 ± 0.18) × 10 ⁻³		DESIG=339
Γ ₁₈₄	$K_2^*(1430)^0\bar{K}^0 + c.c. \rightarrow K_S^0K_L^0\pi^0$	(4.3 ± 1.3) × 10 ⁻⁴		DESIG=338
Γ ₁₈₅	$K^+K^-\pi^+\pi^-$	(7.0 ± 1.0) × 10 ⁻³		DESIG=16
Γ ₁₈₆	$K^+K^-\pi^0\pi^0$	(2.13 ± 0.22) × 10 ⁻³		DESIG=234
Γ ₁₈₇	$K^+K^-\pi^0\pi^0\pi^0$	(1.61 ± 0.29) × 10 ⁻³		DESIG=452
Γ ₁₈₈	$K_S^0K^\pm\pi^\mp\pi^0\pi^0$	(5.3 ± 0.7) × 10 ⁻³		DESIG=453
Γ ₁₈₉	$K_S^0K^\pm\pi^\mp\pi^+\pi^-$	(6.3 ± 0.4) × 10 ⁻³		DESIG=454
Γ ₁₉₀	$K_S^0K^\pm\rho(770)^\pm\pi^0$	(2.9 ± 0.8) × 10 ⁻³		DESIG=463
Γ ₁₉₁	$K_S^0K_L^0\pi^+\pi^-$	(3.8 ± 0.6) × 10 ⁻³		DESIG=296
Γ ₁₉₂	$K_S^0K_L^0\pi^0\pi^0$	(1.9 ± 0.4) × 10 ⁻³		DESIG=337
Γ ₁₉₃	$K_S^0K_L^0\eta$	(1.45 ± 0.33) × 10 ⁻³		DESIG=340
Γ ₁₉₄	$K_S^0K_S^0\pi^+\pi^-$	(1.68 ± 0.19) × 10 ⁻³		DESIG=297
Γ ₁₉₅	$K^\mp K_S^0\pi^\pm\pi^0$	(5.7 ± 0.5) × 10 ⁻³		DESIG=341
Γ ₁₉₆	$K_S^0K^\pm\pi^\mp\rho(770)^0$	(3.1 ± 0.5) × 10 ⁻³		DESIG=456
Γ ₁₉₇	$K^+K^-2(\pi^+\pi^-)$	(3.1 ± 1.3) × 10 ⁻³		DESIG=17
Γ ₁₉₈	$K^+K^-\pi^+\pi^-$	(4.7 ± 0.7) × 10 ⁻³		DESIG=238
Γ ₁₉₉	$2(K^+K^-)$	(7.2 ± 0.8) × 10 ⁻⁴		DESIG=19
Γ ₂₀₀	$K^+K^-K_S^0K_S^0$	(4.2 ± 0.7) × 10 ⁻⁴		DESIG=298
Γ ₂₀₁	$K_S^0K^*(892)^0\pi^+\pi^-$	(1.7 ± 0.6) × 10 ⁻³		DESIG=457
Γ ₂₀₂	$K_S^0K^*(892)^0\pi^0\pi^0$	(1.01 ± 0.18) × 10 ⁻³		DESIG=462
Γ ₂₀₃	$K^\mp K^*(892)^\pm\pi^+\pi^-$	(3.4 ± 1.2) × 10 ⁻³		DESIG=458
Γ ₂₀₄	$K^*(892)^\pm K^*(892)^0\pi^\mp$	(4.8 ± 1.0) × 10 ⁻³		DESIG=459
Γ ₂₀₅	$K^\mp K^*(892)^\pm\pi^0\pi^0$	(1.57 ± 0.32) × 10 ⁻³		DESIG=461
Γ ₂₀₆	$K^*(892)^+K^*(892)^-\pi^0$	(1.12 ± 0.23) %		DESIG=460
Γ ₂₀₇	$\rho\bar{\rho}$	(2.120 ± 0.029) × 10 ⁻³		DESIG=50
Γ ₂₀₈	$\rho\bar{\rho}\pi^0$	(1.19 ± 0.08) × 10 ⁻³	S=1.1	DESIG=52
Γ ₂₀₉	$\rho\bar{\rho}\pi^+\pi^-$	(6.0 ± 0.5) × 10 ⁻³	S=1.3	DESIG=54
Γ ₂₁₀	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	[d] (2.3 ± 0.9) × 10 ⁻³	S=1.9	DESIG=55

Γ ₂₁₁	$p\bar{p}\eta$	$(2.00 \pm 0.12) \times 10^{-3}$		DESIG=56
Γ ₂₁₂	$p\bar{p}\rho$	$< 3.1 \times 10^{-4}$	CL=90%	DESIG=57
Γ ₂₁₃	$p\bar{p}\omega$	$(9.8 \pm 1.0) \times 10^{-4}$	S=1.3	DESIG=58
Γ ₂₁₄	$p\bar{p}\eta'(958)$	$(1.29 \pm 0.14) \times 10^{-4}$	S=2.0	DESIG=59
Γ ₂₁₅	$p\bar{p}a_0(980) \rightarrow p\bar{p}\pi^0\eta$	$(6.8 \pm 1.8) \times 10^{-5}$		DESIG=276
Γ ₂₁₆	$p\bar{p}\phi$	$(5.19 \pm 0.33) \times 10^{-5}$		DESIG=127
Γ ₂₁₇	$p\bar{n}\pi^-$	$(2.12 \pm 0.09) \times 10^{-3}$		DESIG=53
Γ ₂₁₈	$n\bar{n}$	$(2.09 \pm 0.16) \times 10^{-3}$		DESIG=64
Γ ₂₁₉	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$		DESIG=65
Γ ₂₂₀	$nN(1440)$	seen		DESIG=215;OUR EST;→ UNCHECKED ←
Γ ₂₂₁	$nN(1520)$	seen		DESIG=216;OUR EST;→ UNCHECKED ←
Γ ₂₂₂	$nN(1535)$	seen		DESIG=217;OUR EST;→ UNCHECKED ←
Γ ₂₂₃	$\Lambda\bar{\Lambda}$	$(1.88 \pm 0.08) \times 10^{-3}$	S=2.6	DESIG=60
Γ ₂₂₄	$\Lambda\bar{\Lambda}\pi^0$	$(3.8 \pm 0.4) \times 10^{-5}$		DESIG=109
Γ ₂₂₅	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(4.3 \pm 1.0) \times 10^{-3}$		DESIG=261
Γ ₂₂₆	$\Lambda\bar{\Lambda}\eta$	$(1.62 \pm 0.17) \times 10^{-4}$		DESIG=228
Γ ₂₂₇	$\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.}$	[b] $(1.26 \pm 0.05) \times 10^{-3}$	S=1.2	DESIG=71
Γ ₂₂₈	$\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.}$	$(1.21 \pm 0.07) \times 10^{-3}$	S=1.8	DESIG=449
Γ ₂₂₉	$pK^-\bar{\Lambda} + \text{c.c.}$	$(8.6 \pm 1.1) \times 10^{-4}$		DESIG=72
Γ ₂₃₀	$pK^-\Sigma^0$	$(2.9 \pm 0.8) \times 10^{-4}$		DESIG=73
Γ ₂₃₁	$\bar{\Lambda}nK_S^0 + \text{c.c.}$	$(6.5 \pm 1.1) \times 10^{-4}$		DESIG=225
Γ ₂₃₂	$\Lambda\bar{\Sigma} + \text{c.c.}$	$(2.83 \pm 0.23) \times 10^{-5}$		DESIG=61
Γ ₂₃₃	$\Sigma^+\Sigma^-$	$(1.07 \pm 0.04) \times 10^{-3}$		DESIG=247
Γ ₂₃₄	$\Sigma^0\Sigma^0$	$(1.172 \pm 0.032) \times 10^{-3}$	S=1.4	DESIG=63
Γ ₂₃₅	$\Sigma^+\Sigma^-\eta$	$(6.3 \pm 0.4) \times 10^{-5}$		DESIG=448
Γ ₂₃₆	$\Xi^-\Xi^+$	$(9.7 \pm 0.8) \times 10^{-4}$	S=1.4	DESIG=62

Radiative decays

Γ ₂₃₇	$\gamma\eta_c(1S)$	$(1.41 \pm 0.14) \%$	S=1.3	NODE=M070;CLUMP=C DESIG=85
Γ ₂₃₈	$\gamma\eta_c(1S) \rightarrow 3\gamma$			DESIG=246
Γ ₂₃₉	$\gamma\eta_c(1S) \rightarrow \gamma\eta\eta\eta'$			DESIG=391
Γ ₂₄₀	3γ	$(1.16 \pm 0.22) \times 10^{-5}$		DESIG=81
Γ ₂₄₁	4γ	$< 9 \times 10^{-6}$	CL=90%	DESIG=244
Γ ₂₄₂	5γ	$< 1.5 \times 10^{-5}$	CL=90%	DESIG=245
Γ ₂₄₃	$\gamma\pi^0$	$(3.39 \pm 0.08) \times 10^{-5}$		DESIG=82
Γ ₂₄₄	$\gamma\pi^0\pi^0$	$(1.15 \pm 0.05) \times 10^{-3}$		DESIG=283
Γ ₂₄₅	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9	DESIG=95
Γ ₂₄₆	$\gamma f_2(1270) f_2(1270)$	$(9.5 \pm 1.7) \times 10^{-4}$		DESIG=203
Γ ₂₄₇	$\gamma f_2(1270) f_2(1270)$ (non resonant)	$(8.2 \pm 1.9) \times 10^{-4}$		DESIG=204
Γ ₂₄₈	$\gamma\pi^+\pi^- 2\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$		DESIG=99
Γ ₂₄₉	$\gamma K_S^0 K_S^0$	$(8.1 \pm 0.4) \times 10^{-4}$		DESIG=378
Γ ₂₅₀	$\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}]$	$(7 \pm 4) \times 10^{-4}$	S=2.1	DESIG=176
Γ ₂₅₁	$\gamma K^+ K^- \pi^+ \pi^-$	$(2.1 \pm 0.6) \times 10^{-3}$		DESIG=143
Γ ₂₅₂	$\gamma K^*(892) \bar{K}^*(892)$	$(4.0 \pm 1.3) \times 10^{-3}$		DESIG=145
Γ ₂₅₃	$\gamma\eta$	$(1.090 \pm 0.013) \times 10^{-3}$		DESIG=83
Γ ₂₅₄	$\gamma\eta\pi^0$	$(2.14 \pm 0.31) \times 10^{-5}$		DESIG=292
Γ ₂₅₅	$\gamma f_0(500) \rightarrow \gamma\pi\pi$			DESIG=398
Γ ₂₅₆	$\gamma f_0(500) \rightarrow \gamma K\bar{K}$			DESIG=399
Γ ₂₅₇	$\gamma f_0(500) \rightarrow \gamma\eta\eta$			DESIG=400
Γ ₂₅₈	$\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0$	$< 2.5 \times 10^{-6}$	CL=95%	DESIG=293
Γ ₂₅₉	$\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0$	$< 6.6 \times 10^{-6}$	CL=95%	DESIG=294
Γ ₂₆₀	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$		DESIG=96
Γ ₂₆₁	$\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-$	$(6.2 \pm 2.4) \times 10^{-4}$		DESIG=142
Γ ₂₆₂	$\gamma\eta'(958)$	$(5.28 \pm 0.06) \times 10^{-3}$	S=1.3	DESIG=84
Γ ₂₆₃	$\gamma f_0(980) \rightarrow \gamma\pi\pi$			DESIG=393
Γ ₂₆₄	$\gamma f_0(980) \rightarrow \gamma K\bar{K}$			DESIG=394
Γ ₂₆₅	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$		DESIG=94
Γ ₂₆₆	$\gamma\rho\omega$	$< 5.4 \times 10^{-4}$	CL=90%	DESIG=226
Γ ₂₆₇	$\gamma\rho\phi$	$< 8.8 \times 10^{-5}$	CL=90%	DESIG=227
Γ ₂₆₈	$\gamma\omega\omega$	$(1.61 \pm 0.33) \times 10^{-3}$		DESIG=97

Γ ₂₆₉	$\gamma\phi\phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1	DESIG=98
Γ ₂₇₀	$\gamma\eta(1405/1475) \rightarrow \gamma K\bar{K}\pi$	$(2.8 \pm 0.6) \times 10^{-3}$	S=1.6	DESIG=89
Γ ₂₇₁	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0$	$(7.8 \pm 2.0) \times 10^{-5}$	S=1.8	DESIG=171
Γ ₂₇₂	$\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=170
Γ ₂₇₃	$\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0$	$(1.7 \pm 0.4) \times 10^{-3}$	S=1.3	DESIG=124
Γ ₂₇₄	$\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi$	$< 8.2 \times 10^{-5}$	CL=95%	DESIG=212
Γ ₂₇₅	$\gamma\eta(1405) \rightarrow \gamma\gamma\gamma$	$< 2.63 \times 10^{-6}$	CL=90%	DESIG=348
Γ ₂₇₆	$\gamma\eta(1475) \rightarrow \gamma\gamma\gamma$	$< 1.86 \times 10^{-6}$	CL=90%	DESIG=349
Γ ₂₇₇	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$		DESIG=125
Γ ₂₇₈	$\gamma\eta(1760) \rightarrow \gamma\omega\omega$	$(1.98 \pm 0.33) \times 10^{-3}$		DESIG=224
Γ ₂₇₉	$\gamma\eta(1760) \rightarrow \gamma\gamma\gamma$	$< 4.80 \times 10^{-6}$	CL=90%	DESIG=347
Γ ₂₈₀	$\gamma\eta(2225)$	$(3.14 \pm 0.50) \times 10^{-4}$		DESIG=126
Γ ₂₈₁	$\gamma f_2(1270)$	$(1.63 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=86
Γ ₂₈₂	$\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0$	$(2.58 \pm 0.60) \times 10^{-5}$		DESIG=373
Γ ₂₈₃	$\gamma f_1(1285)$	$(6.1 \pm 0.8) \times 10^{-4}$		DESIG=88
Γ ₂₈₄	$\gamma f_0(1370) \rightarrow \gamma\pi\pi$			DESIG=395
Γ ₂₈₅	$\gamma f_0(1370) \rightarrow \gamma K\bar{K}$	$(4.2 \pm 1.5) \times 10^{-4}$		DESIG=284
Γ ₂₈₆	$\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0$	$(1.1 \pm 0.4) \times 10^{-5}$		DESIG=368
Γ ₂₈₇	$\gamma f_0(1370) \rightarrow \gamma\eta\eta$			DESIG=396
Γ ₂₈₈	$\gamma f_0(1370) \rightarrow \gamma\eta\eta'$			DESIG=397
Γ ₂₈₉	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(7.9 \pm 1.3) \times 10^{-4}$		DESIG=175
Γ ₂₉₀	$\gamma f_0(1500) \rightarrow \gamma\pi\pi$	$(1.09 \pm 0.24) \times 10^{-4}$		DESIG=172
Γ ₂₉₁	$\gamma f_0(1500) \rightarrow \gamma\eta\eta$	$(1.7 \pm 0.6) \times 10^{-5}$		DESIG=265
Γ ₂₉₂	$\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0$	$(1.59 \pm 0.24) \times 10^{-5}$		DESIG=369
Γ ₂₉₃	$\gamma f_0(1500) \rightarrow \gamma\eta\eta'$			DESIG=401
Γ ₂₉₄	$\gamma f_1(1510) \rightarrow \gamma\eta\pi^+\pi^-$	$(4.5 \pm 1.2) \times 10^{-4}$		DESIG=141
Γ ₂₉₅	$\gamma f_2'(1525)$	$(5.7 \pm 0.8) \times 10^{-4}$	S=1.5	DESIG=87
Γ ₂₉₆	$\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0$	$(8.0 \pm 0.7) \times 10^{-5}$		DESIG=374
Γ ₂₉₇	$\gamma f_2'(1525) \rightarrow \gamma\eta\eta$	$(3.4 \pm 1.4) \times 10^{-5}$		DESIG=268
Γ ₂₉₈	$\gamma f_2(1565) \rightarrow \gamma\eta\eta'$			DESIG=432
Γ ₂₉₉	$\gamma f_2(1640) \rightarrow \gamma\omega\omega$	$(2.8 \pm 1.8) \times 10^{-4}$		DESIG=222
Γ ₃₀₀	$\gamma f_0(1710) \rightarrow \gamma\pi\pi$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=135
Γ ₃₀₁	$\gamma f_0(1710) \rightarrow \gamma K\bar{K}$	$(9.5 \pm 1.0) \times 10^{-4}$	S=1.5	DESIG=91
Γ ₃₀₂	$\gamma f_0(1710) \rightarrow \gamma\omega\omega$	$(3.1 \pm 1.0) \times 10^{-4}$		DESIG=221
Γ ₃₀₃	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	$(2.4 \pm 1.2) \times 10^{-4}$		DESIG=266
Γ ₃₀₄	$\gamma f_0(1710) \rightarrow \gamma\eta\eta'$			DESIG=402
Γ ₃₀₅	$\gamma f_0(1710) \rightarrow \gamma\omega\phi$	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=262
Γ ₃₀₆	$\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0$	$(1.11 \pm 0.20) \times 10^{-5}$		DESIG=370
Γ ₃₀₇	$\gamma f_2(1810) \rightarrow \gamma\eta\eta$	$(5.4 \pm 3.5) \times 10^{-5}$		DESIG=269
Γ ₃₀₈	$\gamma\eta_1(1855) \rightarrow \gamma\eta\eta'$	$(2.7 \pm 0.4) \times 10^{-6}$		DESIG=447
Γ ₃₀₉	$\gamma f_0(1770) \rightarrow \gamma\eta\eta'$			DESIG=431
Γ ₃₁₀	$\gamma f_2(1910) \rightarrow \gamma\omega\omega$	$(2.0 \pm 1.4) \times 10^{-4}$		DESIG=223
Γ ₃₁₁	$\gamma f_2(1950) \rightarrow \gamma K^*(892)\bar{K}^*(892)$	$(7.0 \pm 2.2) \times 10^{-4}$		DESIG=144
Γ ₃₁₂	$\gamma f_2(2010) \rightarrow \gamma\eta\eta'$			DESIG=440
Γ ₃₁₃	$\gamma f_0(2020) \rightarrow \gamma\pi\pi$			DESIG=403
Γ ₃₁₄	$\gamma f_0(2020) \rightarrow \gamma K\bar{K}$			DESIG=404
Γ ₃₁₅	$\gamma f_0(2020) \rightarrow \gamma\eta\eta$			DESIG=405
Γ ₃₁₆	$\gamma f_0(2020) \rightarrow \gamma\eta'\eta'$	$(2.63 \pm 0.32) \times 10^{-4}$		DESIG=426
Γ ₃₁₇	$\gamma f_0(2020) \rightarrow \gamma\eta\eta'$			DESIG=438

Γ ₃₁₈	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$		DESIG=100
Γ ₃₁₉	$\gamma f_4(2050) \rightarrow \gamma \eta \eta'$			DESIG=433
Γ ₃₂₀	$\gamma f_0(2100) \rightarrow \gamma \eta \eta$	$(1.13 \pm_{-0.30}^{0.60}) \times 10^{-4}$		DESIG=267
Γ ₃₂₁	$\gamma f_0(2100) \rightarrow \gamma K \bar{K}$			DESIG=406
Γ ₃₂₂	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(6.2 \pm 1.0) \times 10^{-4}$		DESIG=286
Γ ₃₂₃	$\gamma f_0(2200)$			DESIG=123
Γ ₃₂₄	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(5.9 \pm 1.3) \times 10^{-4}$		DESIG=285
Γ ₃₂₅	$\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0$	$(2.72 \pm_{-0.50}^{0.19}) \times 10^{-4}$		DESIG=371
Γ ₃₂₆	$\gamma f_0(2200) \rightarrow \gamma \pi \pi$			DESIG=407
Γ ₃₂₇	$\gamma f_0(2200) \rightarrow \gamma \eta \eta$			DESIG=408
Γ ₃₂₈	$\gamma f_J(2220)$			DESIG=92
Γ ₃₂₉	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	< 3.9	$\times 10^{-5}$	CL=90% DESIG=136
Γ ₃₃₀	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	< 4.1	$\times 10^{-5}$	CL=90% DESIG=137
Γ ₃₃₁	$\gamma f_J(2220) \rightarrow \gamma p \bar{p}$	$(1.5 \pm 0.8) \times 10^{-5}$		DESIG=138
Γ ₃₃₂	$\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0$	$(4.9 \pm 0.7) \times 10^{-5}$		DESIG=372
Γ ₃₃₃	$\gamma f_0(2330) \rightarrow \gamma \pi \pi$			DESIG=409
Γ ₃₃₄	$\gamma f_0(2330) \rightarrow \gamma \eta \eta$			DESIG=410
Γ ₃₃₅	$\gamma f_0(2330) \rightarrow \gamma \eta' \eta'$	$(6.1 \pm_{-1.8}^{4.0}) \times 10^{-6}$		DESIG=427
Γ ₃₃₆	$\gamma f_0(2330) \rightarrow \gamma \eta \eta'$			DESIG=439
Γ ₃₃₇	$\gamma f_2(2340) \rightarrow \gamma \eta \eta$	$(5.6 \pm_{-2.2}^{2.4}) \times 10^{-5}$		DESIG=270
Γ ₃₃₈	$\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0$	$(5.5 \pm_{-1.5}^{4.0}) \times 10^{-5}$		DESIG=375
Γ ₃₃₉	$\gamma f_2(2340) \rightarrow \gamma \eta' \eta'$	$(8.7 \pm_{-1.8}^{0.9}) \times 10^{-6}$		DESIG=428
Γ ₃₄₀	$\gamma f_0(2470) \rightarrow \gamma \eta' \eta'$	$(8.2 \pm_{-2.8}^{4.0}) \times 10^{-7}$		DESIG=429
Γ ₃₄₁	$\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta'$	$(2.7 \pm_{-0.8}^{0.6}) \times 10^{-4}$	S=1.6	DESIG=213
Γ ₃₄₂	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(7.7 \pm_{-0.9}^{1.5}) \times 10^{-5}$		DESIG=254
Γ ₃₄₃	$\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta$	$(3.3 \pm_{-1.3}^{2.0}) \times 10^{-5}$		DESIG=282
Γ ₃₄₄	$\gamma X(1835) \rightarrow \gamma \gamma \phi(1020)$			DESIG=346
Γ ₃₄₅	$\gamma X(1835) \rightarrow \gamma \gamma \gamma$	< 3.56	$\times 10^{-6}$	CL=90% DESIG=350
Γ ₃₄₆	$\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-)$	$(2.4 \pm_{-0.8}^{0.7}) \times 10^{-5}$		DESIG=264
Γ ₃₄₇	$\gamma X(2370) \rightarrow \gamma K^+ K^- \eta'$	$(1.8 \pm 0.7) \times 10^{-5}$		DESIG=388
Γ ₃₄₈	$\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta'$	$(1.2 \pm 0.5) \times 10^{-5}$		DESIG=389
Γ ₃₄₉	$\gamma X(2370) \rightarrow \gamma \eta \eta \eta'$	< 9.2	$\times 10^{-6}$	CL=90% DESIG=390
Γ ₃₅₀	$\gamma p \bar{p}$	$(3.8 \pm 1.0) \times 10^{-4}$		DESIG=90
Γ ₃₅₁	$\gamma p \bar{p} \pi^+ \pi^-$	< 7.9	$\times 10^{-4}$	CL=90% DESIG=93
Γ ₃₅₂	$\gamma \Lambda \bar{\Lambda}$	< 1.3	$\times 10^{-4}$	CL=90% DESIG=200
Γ ₃₅₃	$\gamma A^0 \rightarrow \gamma \text{invisible}$	[e] < 1.7	$\times 10^{-6}$	CL=90% DESIG=251
Γ ₃₅₄	$\gamma A^0 \rightarrow \gamma \mu^+ \mu^-$	[f] < 7.8	$\times 10^{-7}$	CL=90% DESIG=259

Dalitz decays

Γ ₃₅₅	$\pi^0 e^+ e^-$	$(7.6 \pm 1.4) \times 10^{-7}$		NODE=M070;CLUMP=G DESIG=271
Γ ₃₅₆	$\eta e^+ e^-$	$(1.42 \pm 0.08) \times 10^{-5}$		DESIG=272
Γ ₃₅₇	$\eta'(958) e^+ e^-$	$(6.59 \pm 0.18) \times 10^{-5}$		DESIG=273
Γ ₃₅₈	$X(1835) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta'$	$(3.58 \pm 0.25) \times 10^{-6}$		DESIG=423
Γ ₃₅₉	$X(2120) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta'$	$(8.2 \pm 1.3) \times 10^{-7}$		DESIG=425
Γ ₃₆₀	$X(2370) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta'$	$(1.08 \pm 0.17) \times 10^{-6}$		DESIG=424
Γ ₃₆₁	$\eta U \rightarrow \eta e^+ e^-$	[g] < 9.11	$\times 10^{-7}$	CL=90% DESIG=352
Γ ₃₆₂	$\eta'(958) U \rightarrow \eta'(958) e^+ e^-$	[g] < 2.0	$\times 10^{-7}$	CL=90% DESIG=366
Γ ₃₆₃	$\phi e^+ e^-$	< 1.2	$\times 10^{-7}$	CL=90% DESIG=384

Weak decays

Γ_{364}	$D^- e^+ \nu_e + \text{c.c.}$	< 7.1	$\times 10^{-8}$	CL=90%	NODE=M070;CLUMP=E DESIG=218
Γ_{365}	$\bar{D}^0 e^+ e^- + \text{c.c.}$	< 8.5	$\times 10^{-8}$	CL=90%	DESIG=219
Γ_{366}	$D_s^- e^+ \nu_e + \text{c.c.}$	< 1.3	$\times 10^{-6}$	CL=90%	DESIG=220
Γ_{367}	$D_s^{*-} e^+ \nu_e + \text{c.c.}$	< 1.8	$\times 10^{-6}$	CL=90%	DESIG=290
Γ_{368}	$D^- \pi^+ + \text{c.c.}$	< 7.5	$\times 10^{-5}$	CL=90%	DESIG=241
Γ_{369}	$\bar{D}^0 \bar{K}^0 + \text{c.c.}$	< 1.7	$\times 10^{-4}$	CL=90%	DESIG=242
Γ_{370}	$\bar{D}^0 \bar{K}^{*0} + \text{c.c.}$	< 2.5	$\times 10^{-6}$	CL=90%	DESIG=275
Γ_{371}	$D_s^- \pi^+ + \text{c.c.}$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=243
Γ_{372}	$D_s^- \rho^+ + \text{c.c.}$	< 1.3	$\times 10^{-5}$	CL=90%	DESIG=274

**Charge conjugation (C), Parity (P),
Lepton Family number (LF) violating modes**

Γ_{373}	$\gamma\gamma$	C	< 2.7	$\times 10^{-7}$	CL=90%	NODE=M070;CLUMP=D DESIG=80
Γ_{374}	$\gamma\phi$	C	< 1.4	$\times 10^{-6}$	CL=90%	DESIG=277
Γ_{375}	$e^\pm \mu^\mp$	LF	< 1.6	$\times 10^{-7}$	CL=90%	DESIG=177
Γ_{376}	$e^\pm \tau^\mp$	LF	< 7.5	$\times 10^{-8}$	CL=90%	DESIG=178
Γ_{377}	$\mu^\pm \tau^\mp$	LF	< 2.0	$\times 10^{-6}$	CL=90%	DESIG=179
Γ_{378}	$\Lambda_c^+ e^- + \text{c.c.}$		< 6.9	$\times 10^{-8}$	CL=90%	DESIG=379

Other decays

Γ_{379}	invisible	< 7	$\times 10^{-4}$	CL=90%	NODE=M070;CLUMP=F DESIG=240
----------------	-----------	-----	------------------	--------	--------------------------------

[a] For $E_\gamma > 100$ MeV.

LINKAGE=EGM

[b] The value is for the sum of the charge states or particle/antiparticle states indicated.

LINKAGE=SG

[c] $\Theta(1540)$ is a hypothetical pentaquark state of $1.54 \text{ GeV}/c^2$ mass and a width of less than $25 \text{ MeV}/c^2$.

LINKAGE=THT

[d] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta, p\bar{p}\omega, p\bar{p}\eta'$.

LINKAGE=MF

[e] For a narrow state A with mass less than 960 MeV.

LINKAGE=NSA

[f] For a narrow scalar or pseudoscalar A^0 with mass 0.21–3.0 GeV.

LINKAGE=NA0

[g] For a dark photon U with mass between 100 and 2100 MeV.

LINKAGE=DPH

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

 $J/\psi(1S)$ PARTIAL WIDTHS

NODE=M070220

 $\Gamma(\text{hadrons})$ **Γ_1**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$81.37 \pm 1.36 \pm 1.30$	¹ ANASHIN	20	KEDR $e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
74.1 \pm 8.1	BAI	95B	BES $e^+ e^-$
59 \pm 24	BALDINI-...	75	FRAG $e^+ e^-$
59 \pm 14	BOYARSKI	75	MRK1 $e^+ e^-$
50 \pm 25	ESPOSITO	75B	FRAM $e^+ e^-$

NODE=M070W3
NODE=M070W3

¹ Based on the same dataset as ANASHIN 18A and correlated to the values reported there

NODE=M070W3;LINKAGE=A

$\Gamma(e^+e^-)$ Γ_5

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.53 ±0.10 OUR AVERAGE				
5.550 ±0.056 ±0.089		1,2 ANASHIN	18A KEDR	e^+e^-
5.36 $^{+0.29}_{-0.28}$		3 HSUEH	92 RVUE	See Υ mini-review
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.58 ±0.05 ±0.08		4 ABLIKIM	16Q BES3	$3.773 e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.71 ±0.16	13k	5 ADAMS	06A CLEO	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.57 ±0.19	7.8k	5 AUBERT	04 BABR	$e^+e^- \rightarrow \mu^+\mu^-\gamma$
5.14 ±0.39		BAI	95B BES	e^+e^-
4.72 ±0.35		ALEXANDER	89 RVUE	See Υ mini-review
4.4 ±0.6		3 BRANDELIK	79C DASP	e^+e^-
4.6 ±0.8		6 BALDINI-...	75 FRAG	e^+e^-
4.8 ±0.6		BOYARSKI	75 MRK1	e^+e^-
4.6 ±1.0		ESPOSITO	75B FRAM	e^+e^-

NODE=M070W1
NODE=M070W1

OCCUR=4

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

⁴ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.973 \pm 0.007 \pm 0.037)\%$ from ABLIKIM 13R.

⁵ Calculated by us from the reported values of $\Gamma(e^+e^-) \times B(\mu^+\mu^-)$ using $B(\mu^+\mu^-) = (5.93 \pm 0.06)\%$.

⁶ Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

NODE=M070W1;LINKAGE=D

NODE=M070W1;LINKAGE=E
NODE=M070W1;LINKAGE=FNODE=M070W1;LINKAGE=A
NODE=M070W1;LINKAGE=AA

NODE=M070W1;LINKAGE=B

 $\Gamma(\mu^+\mu^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.13 ±0.52	BAI	95B BES	e^+e^-
4.8 ±0.6	BOYARSKI	75 MRK1	e^+e^-
5 ±1	ESPOSITO	75B FRAM	e^+e^-

NODE=M070W2
NODE=M070W2 $\Gamma(\gamma\gamma)$ Γ_{373}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5.4	90	BRANDELIK	79C DASP	e^+e^-

NODE=M070W70
NODE=M070W70 $J/\psi(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M070225

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation.

NODE=M070225

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.884 ±0.048 ±0.078	1,2 ANASHIN	18A KEDR	e^+e^-
4 ±0.8	3 BALDINI-...	75 FRAG	e^+e^-
3.9 ±0.8	3 ESPOSITO	75B FRAM	e^+e^-

NODE=M070G3
NODE=M070G3

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

NODE=M070G3;LINKAGE=A

NODE=M070G3;LINKAGE=B
NODE=M070G3;LINKAGE=S $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
333.1 ± 6.6 ±4.0	1,2 ANASHIN	18A KEDR	e^+e^-
332.3 ± 6.4 ±4.8	ANASHIN	10 KEDR	$3.097 e^+e^- \rightarrow e^+e^-$
350 ± 20	BRANDELIK	79C DASP	e^+e^-
320 ± 70	3 BALDINI-...	75 FRAG	e^+e^-
340 ± 90	3 ESPOSITO	75B FRAM	e^+e^-
360 ±100	3 FORD	75 SPEC	e^+e^-

NODE=M070G1
NODE=M070G1

OCCUR=2

¹ From the cross sections of $e^+e^- \rightarrow e^+e^-$ and $e^+e^- \rightarrow$ hadrons near the $J/\psi(1S)$ peak.

² Based on the same dataset as ANASHIN 20 and correlated to the values reported there.

³ Data redundant with branching ratios or partial widths above.

NODE=M070G1;LINKAGE=A

NODE=M070G1;LINKAGE=B
NODE=M070G1;LINKAGE=S

$$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_7\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
333 ± 4 OUR AVERAGE				
333.4 ± 2.5 ± 4.4		ABLIKIM	16Q BES3	3.773 e ⁺ e ⁻ → μ ⁺ μ ⁻ γ
331.8 ± 5.2 ± 6.3		ANASHIN	10 KEDR	3.097 e ⁺ e ⁻ → μ ⁺ μ ⁻
338.4 ± 5.8 ± 7.1	13k	ADAMS	06A CLEO	e ⁺ e ⁻ → μ ⁺ μ ⁻ γ
330.1 ± 7.7 ± 7.3	7.8k	AUBERT	04 BABR	e ⁺ e ⁻ → μ ⁺ μ ⁻ γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
510 ± 90		DASP	75 DASP	e ⁺ e ⁻
380 ± 50		¹ ESPOSITO	75B FRAM	e ⁺ e ⁻

NODE=M070G2
NODE=M070G2

¹Data redundant with branching ratios or partial widths above.

NODE=M070G2;LINKAGE=S

$$\Gamma(\eta\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{13}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 OUR AVERAGE				
2.34 ± 0.43 ± 0.16	49	LEES	18 BABR	e ⁺ e ⁻ → ηπ ⁺ π ⁻ γ
2.22 ± 0.96 ± 0.02	9	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → ηπ ⁺ π ⁻ γ

NODE=M070G25
NODE=M070G25

¹AUBERT 07AU reports [Γ(J/ψ(1S) → ηπ⁺π⁻) × Γ(J/ψ(1S) → e⁺e⁻)/Γ_{total}] × [B(η → π⁺π⁻π⁰)] = 0.51 ± 0.22 ± 0.03 eV which we divide by our best value B(η → π⁺π⁻π⁰) = (23.02 ± 0.25) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G25;LINKAGE=AU

$$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{14}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
64.8 ± 11.1 ± 0.4	200	¹ LEES	21C BABR	e ⁺ e ⁻ → γ _{ISR} (π ⁺ π ⁻ 4π ⁰)

NODE=M070Q20
NODE=M070Q20

¹LEES 21C reports [Γ(J/ψ(1S) → ηπ⁺π⁻π⁰) × Γ(J/ψ(1S) → e⁺e⁻)/Γ_{total}] × [B(η → 3π⁰)] = 21.1 ± 1.7 ± 3.2 eV which we divide by our best value B(η → 3π⁰) = (32.57 ± 0.21) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q20;LINKAGE=A

$$\Gamma(\eta\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{15}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
26.9 ± 5.7 ± 0.1	101	¹ LEES	21C BABR	e ⁺ e ⁻ → γ _{ISR} (π ⁺ π ⁻ 3π ⁰ γγ)

NODE=M070Q19
NODE=M070Q19

¹LEES 21C reports [Γ(J/ψ(1S) → ηπ⁺π⁻3π⁰) × Γ(J/ψ(1S) → e⁺e⁻)/Γ_{total}] × [B(η → 2γ)] = 10.6 ± 1.6 ± 1.6 eV which we divide by our best value B(η → 2γ) = (39.36 ± 0.18) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q19;LINKAGE=A

$$\Gamma(\eta K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{19}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
4.76 ± 1.64 ± 0.03	¹ LEES	23 BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M070Q54
NODE=M070Q54

¹LEES 23 reports [Γ(J/ψ(1S) → ηK⁺K⁻) × Γ(J/ψ(1S) → e⁺e⁻)/Γ_{total}] × [B(η → 3π⁰)] = 1.55 ± 0.51 ± 0.16 eV which we divide by our best value B(η → 3π⁰) = (32.57 ± 0.21) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q54;LINKAGE=A

$$\Gamma(\eta K_S^0 K_S^0 \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{20}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
7.3 ± 1.4 ± 0.4	44	LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

NODE=M070G38
NODE=M070G38

$$\Gamma(K_S^0 K^\pm \pi^\mp \rho(770)^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{196}\Gamma_5/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
17.3 ± 2.1 ± 1.7	LEES	23 BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M070Q55
NODE=M070Q55

$$\Gamma(\rho^\pm \pi^\mp \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{23}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
160 ± 40 OUR AVERAGE		[155 ± 40 eV OUR 2023 AVERAGE]		
155 ± 26 ± 36	14k	LEES	21 BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

NODE=M070Q13
NODE=M070Q13

NEW

$$\Gamma(\rho^+ \rho^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$$

 $\Gamma_{24}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
32 ± 13 ± 15	14k	LEES	21 BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

NODE=M070Q14
NODE=M070Q14

$$\Gamma(\rho^\mp K^\pm K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{26} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.4±1.0±1.9	130	LEES	17D BABR	$e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$

NODE=M070G31
NODE=M070G31

$$\Gamma(\omega \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{41} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
53.6±5.0±0.4	788	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \gamma$

NODE=M070G24
NODE=M070G24

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^+ \pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 47.8 \pm 3.1 \pm 3.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G24;LINKAGE=AU

$$\Gamma(\omega \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{42} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
27.8±3.5±0.2	398	¹ LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- 3\pi^0 \gamma$

NODE=M070P54
NODE=M070P54

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 24.8 \pm 1.8 \pm 2.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P54;LINKAGE=A

$$\Gamma(\omega 3\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{43} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±3.1±0.1	89	¹ LEES	21C BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

NODE=M070Q18
NODE=M070Q18

¹ LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega 3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 9.4 \pm 2.3 \pm 1.5$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q18;LINKAGE=A

$$\Gamma(\omega \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{46} \Gamma_5 / \Gamma$$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.3±0.2	170	AUBERT	06D BABR	$10.6 e^+ e^- \rightarrow \omega \pi^+ \pi^- \pi^0 \gamma$

NODE=M070G8
NODE=M070G8

$$\Gamma(\omega \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{45} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
16.9±7.6±0.2	¹ LEES	21C BABR	$e^+ e^- \rightarrow \gamma_{ISR}(\pi^+ \pi^- 4\pi^0)$

NODE=M070Q21
NODE=M070Q21

¹ Different final state as in AUBERT 06. LEES 21C reports $[\Gamma(J/\psi(1S) \rightarrow \omega \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 4.9 \pm 2.1 \pm 0.7$ eV which we divide by our best values $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$, $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070Q21;LINKAGE=A

$$\Gamma(\omega \pi^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{47} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.96±0.01	27	¹ LEES	18E BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \pi^0 \eta \gamma$

NODE=M070P58
NODE=M070P58

¹ LEES 18E reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^0 \eta) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 1.7 \pm 0.8 \pm 0.3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P58;LINKAGE=A

$$\Gamma(\omega \pi^+ \pi^- 2\pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{49} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
185±30±1	14k	¹ LEES	21 BABR	$10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$

NODE=M070Q11
NODE=M070Q11

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \omega \pi^+ \pi^- 2\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 165 \pm 9 \pm 25$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q11;LINKAGE=A

$$\Gamma(\omega K \bar{K}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{60} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.70±1.98±0.03	24	¹ AUBERT	07AU BABR	$10.6 e^+ e^- \rightarrow \omega K^+ K^- \gamma$

NODE=M070G29
NODE=M070G29

¹ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \omega K \bar{K}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0)] = 3.3 \pm 1.3 \pm 1.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G29;LINKAGE=AU

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{69} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
29.0 ± 1.7 ± 1.3	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁺ K [*] (892) ⁻ γ

NODE=M070G18
NODE=M070G18

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^+ K^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{70} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.96 ± 0.85 ± 0.70	155	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁰ γ

NODE=M070G20
NODE=M070G20

$$\Gamma(K^+ K^*(892)^- + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{71} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
16.76 ± 1.70 ± 1.00	89	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ

NODE=M070G21
NODE=M070G21

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{72} \Gamma_5 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
26.6 ± 2.5 ± 1.5	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K ⁰ $\bar{K}^*(892)^0$ γ

NODE=M070G19
NODE=M070G19

$$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.} \rightarrow K^0 K^\pm \pi^\mp + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{73} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
17.70 ± 1.70 ± 1.00	94	AUBERT	08S BABR	10.6 e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] γ

NODE=M070G22
NODE=M070G22

$$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{74} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
42.6 ± 4.8 ± 7.2	99	¹ LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

NODE=M070G39
NODE=M070G39

¹ Dividing by 1/6 to account for B(K^{*}(892)⁰ → K_S⁰ π⁰) = 1/6.

NODE=M070G39;LINKAGE=A

$$\Gamma(K^*(892)^\pm K^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{75} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.8 ± 2.8 ± 6.8	80	¹ LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

NODE=M070G32
NODE=M070G32

¹ Dividing by 1/4 to account for B(K^{*}(892)[±] → K_S⁰ π[±]) = 1/4.

NODE=M070G32;LINKAGE=A

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{76} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.0 ± 2.8 OUR AVERAGE				
9.2 ± 1.2 ± 3.2	64	¹ LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ
14.8 ± 4.8 ± 1.2	53	² LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _S ⁰ γ

NODE=M070GY4
NODE=M070GY4

¹ Dividing by 1/2 to take into account B(K^{*}(892)[±] → K[±] π[∓]) = 1/2.

² Dividing by 1/4 to take into account B(K^{*}(892) → K_S⁰ π) = 1/4.

NODE=M070GY4;LINKAGE=B
NODE=M070GY4;LINKAGE=A

$$\Gamma(K^*(892)^+ K_S^0 \pi^- + \text{c.c.} \rightarrow K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{77} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.7 ± 1.2 ± 0.3	53	LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _S ⁰ γ

NODE=M070GY5
NODE=M070GY5

$$\Gamma(K^*(892)^0 K_S^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{80} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.60 ± 0.75 ± 2.25	34	¹ LEES	17D BABR	e ⁺ e ⁻ → K _S ⁰ K [±] π [∓] π ⁰ γ

NODE=M070G33
NODE=M070G33

¹ Dividing by 2/3 to account for B(K^{*}(892)⁰ → K⁺ π⁻) = 2/3.

NODE=M070G33;LINKAGE=A

$$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{82} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.28 ± 0.34 ± 0.07	47 ± 12	¹ LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

NODE=M070G01
NODE=M070G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.28 ± 0.40 ± 0.11	25 ± 8	^{1,2} AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
--------------------	--------	-----------------------	-----------	--

¹ Dividing by (2/3)² to take twice into account that B(K^{*}0 → K⁺ π⁻) = 2/3 B(K^{*}0 → K π).

NODE=M070G01;LINKAGE=AE

² Superseded by LEES 12F.

NODE=M070G01;LINKAGE=A

$$\Gamma(K^*(892)^\pm K^*(892)^\mp) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{83} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.80 ± 0.48 ± 0.32	1 ± 5	¹ LEES	14H BABR	e ⁺ e ⁻ → π ⁺ π ⁻ K _S ⁰ K _S ⁰ γ

NODE=M070GY8
NODE=M070GY8

¹ Dividing by (1/4)² to take twice into account B(K^{*}(892) → K_S⁰ π) = 1/4.

NODE=M070GY8;LINKAGE=A

$$\Gamma(K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{94} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.1 ± 9.8 ± 0.5	35	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GY6
NODE=M070GY6

¹ Dividing by 1/4 to take into account $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$.

NODE=M070GY6;LINKAGE=A

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 10.0 \pm 4.8 \pm 0.8$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GY6;LINKAGE=B

$$\Gamma(K_2^*(1430)^0 K^- \pi^+ + \text{c.c.} \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{\text{total}} \quad \Gamma_{93} / \Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.65 ± 0.80 ± 0.44	1094	ANASHIN	22 KEDR	$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M070Q43
NODE=M070Q43

$$\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{95} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.8 ± 1.4 ± 0.6	710	1,2,3 LEES	12F BABR	$10.6 e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M070G02
NODE=M070G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

33 ± 4 ± 1 317 2,4 AUBERT 07AK BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 12.89 \pm 0.54 \pm 0.41$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G02;LINKAGE=A

² Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3 B(K^{*0} \rightarrow K \pi)$.

NODE=M070G02;LINKAGE=AE

³ The $K_2^*(1430)$ cannot be distinguished from the $K_0^*(1430)$.

NODE=M070G02;LINKAGE=B

⁴ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 16.4 \pm 1.1 \pm 1.4$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G02;LINKAGE=UB

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{96} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
18.6 ± 16.1 ± 0.4	8 ± 8	1,2 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GY9
NODE=M070GY9

¹ Dividing by $(1/4)^2$ to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$ and $B(K^*(1430) \rightarrow K_S^0 \pi) = 1/4 B(K^*(1430) \rightarrow K \pi)$.

NODE=M070GY9;LINKAGE=A

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow K_2^*(1430)^- K^*(892)^+ + \text{c.c.}) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}] \times [B(K_2^*(1430) \rightarrow K \pi)] = 9.28 \pm 8.0 \pm 0.32$ eV which we divide by our best value $B(K_2^*(1430) \rightarrow K \pi) = (49.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GY9;LINKAGE=B

$$\Gamma(K_2^*(1430)^- K^*(892)^+ + \text{c.c.} \rightarrow K^*(892)^+ K_S^0 \pi^- + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{97} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.32 ± 2.00 ± 0.08	8 ± 8	1 LEES	14H BABR	$e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$

NODE=M070GZ0
NODE=M070GZ0

¹ Dividing by 1/4 to take into account $B(K^*(892) \rightarrow K_S^0 \pi) = 1/4$.

NODE=M070GZ0;LINKAGE=A

$$\Gamma(\bar{K}_2^*(1770)^0 K^*(892)^0 + \text{c.c.} \rightarrow K^*(892)^0 K^- \pi^+ + \text{c.c.}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{99} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.4 ± 0.3	110 ± 14	1 AUBERT 07AK BABR	10.6	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

NODE=M070G03
NODE=M070G03

¹ Dividing by 2/3 to take into account that $B(K^{*0} \rightarrow K^+ \pi^-) = 2/3$.

NODE=M070G03;LINKAGE=AE

$$\Gamma(\phi \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{106} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.48 ± 0.35 OUR AVERAGE				

NODE=M070G14
NODE=M070G14

4.46 ± 0.49 ± 0.05 181 1 LEES 12F BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

4.51 ± 0.48 ± 0.05 254 ± 23 2 SHEN 09 BELL 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.3 ± 0.7 ± 0.1 103 3 AUBERT, BE 06D BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$

- ¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.19 \pm 0.23 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² SHEN 09 reports $4.50 \pm 0.41 \pm 0.26$ eV from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)]$ assuming $B(\phi(1020) \rightarrow K^+K^-) = (49.2 \pm 0.6) \times 10^{-2}$, which we rescale to our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 2.61 \pm 0.30 \pm 0.18$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G14;LINKAGE=B

NODE=M070G14;LINKAGE=SH

NODE=M070G14;LINKAGE=AU

$\Gamma(\phi\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{107}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.77±0.57±0.03	45	¹ LEES 12F	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
3.13±0.88±0.03	23	² AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\pi^0\gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.36 \pm 0.27 \pm 0.07$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Superseded by LEES 12F. AUBERT, BE 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi\pi^0\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 1.54 \pm 0.40 \pm 0.16$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G15
NODE=M070G15

NODE=M070G15;LINKAGE=A

NODE=M070G15;LINKAGE=AU

$\Gamma(\phi 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{108}\Gamma_5/\Gamma$

VALUE (10^{-2} keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.96±0.19±0.01	35	¹ AUBERT 06D	BABR	10.6 $e^+e^- \rightarrow \phi 2(\pi^+\pi^-)\gamma$
¹ AUBERT 06D reports $[\Gamma(J/\psi(1S) \rightarrow \phi 2(\pi^+\pi^-)) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = (0.47 \pm 0.09 \pm 0.03) \times 10^{-2}$ keV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G10
NODE=M070G10

NODE=M070G10;LINKAGE=AU

$\Gamma(\phi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{109}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.6±1.4 OUR AVERAGE				
[6.1 ± 2.7 eV OUR 2023 AVERAGE]				
4.1±1.6±0.4		¹ LEES 23	BABR	$e^+e^- \rightarrow \gamma_{ISR} \text{ hadrons}$
6.1±2.7±0.4	6	² AUBERT 07AU	BABR	10.6 $e^+e^- \rightarrow \phi\eta\gamma$
¹ LEES 23 quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi^0) = 0.64 \pm 0.26 \pm 0.06$ eV.				
² AUBERT 07AU quotes $\Gamma_{ee}^{J/\psi} \cdot B(J/\psi \rightarrow \phi\eta) \cdot B(\phi \rightarrow K^+K^-) \cdot B(\eta \rightarrow 3\pi) = 0.84 \pm 0.37 \pm 0.05$ eV.				

NODE=M070G28
NODE=M070G28
NEW

OCCUR=2

NODE=M070G28;LINKAGE=A

NODE=M070G28;LINKAGE=AU

$\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{113}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.44±0.19 OUR AVERAGE				
1.40±0.25±0.01	57 ± 9	¹ LEES 12F	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
1.48±0.27±0.09	60 ± 11	² SHEN 09	BELL	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
●●● We do not use the following data for averages, fits, limits, etc. ●●●				
1.02±0.24±0.01	20 ± 5	³ AUBERT 07AK	BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
¹ LEES 12F reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.69 \pm 0.11 \pm 0.05$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				
² Multiplied by 2/3 to take into account the $\phi\pi^+\pi^-$ mode only. Using $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.6)\%$.				
³ Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] = 0.50 \pm 0.11 \pm 0.04$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M070G05
NODE=M070G05

NODE=M070G05;LINKAGE=A

NODE=M070G05;LINKAGE=SH

NODE=M070G05;LINKAGE=UB

$$\Gamma(\phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{114} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

0.98 ± 0.26 ± 0.01	16 ± 4	¹ LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ
---------------------------	--------	-------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.96 ± 0.40 ± 0.01	7.0 ± 2.8	² AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ
--------------------	-----------	---------------------	-----------	--

¹ LEES 12F reports [$\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+ K^-$)] = 0.48 ± 0.12 ± 0.05 eV which we divide by our best value B($\phi(1020) \rightarrow K^+ K^-$) = (49.1 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(J/\psi(1S) \rightarrow \phi f_0(980) \rightarrow \phi \pi^0 \pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+ K^-$)] = 0.47 ± 0.19 ± 0.05 eV which we divide by our best value B($\phi(1020) \rightarrow K^+ K^-$) = (49.1 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G06
NODE=M070G06

NODE=M070G06;LINKAGE=A

NODE=M070G06;LINKAGE=UB

$$\Gamma(\phi f_2(1270)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{119} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

1.79 ± 0.32^{+0.02}_{-0.06}	61	^{1,2,3} LEES	12F BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
--	----	-----------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.08 ± 0.73 ^{+0.05} _{-0.14}	44	^{2,4} AUBERT	07AK BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ
---	----	-----------------------	-----------	--

¹ LEES 12F reports [$\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($f_2(1270) \rightarrow \pi\pi$)] = 1.51 ± 0.25 ± 0.10 eV which we divide by our best value B($f_2(1270) \rightarrow \pi\pi$) = (84.3^{+2.8}_{-1.0}) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using B($\phi \rightarrow K^+ K^-$) = (48.9 ± 0.5)%.

³ Using π⁺π⁻ invariant mass between 1.1 and 1.5 GeV. May include other sources such as f₀(1370).

⁴ Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(J/\psi(1S) \rightarrow \phi f_2(1270)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($f_2(1270) \rightarrow \pi\pi$)] = 3.44 ± 0.55 ± 0.28 eV which we divide by our best value B($f_2(1270) \rightarrow \pi\pi$) = (84.3^{+2.8}_{-1.0}) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G07
NODE=M070G07

NODE=M070G07;LINKAGE=A

NODE=M070G07;LINKAGE=AE

NODE=M070G07;LINKAGE=B

NODE=M070G07;LINKAGE=UB

$$\Gamma(\phi f'_2(1525)) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{124} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

8.2 ± 3.2 ± 0.2	11	^{1,2} LEES	14H BABR	e ⁺ e ⁻ → K _S ⁰ K _S ⁰ K ⁺ K ⁻ γ
------------------------	----	---------------------	----------	---

¹ Dividing by 1/4 to take into account B($f'_2(1525) \rightarrow K_S^0 K_S^0$) = 1/4 B($f'_2(1525) \rightarrow K \bar{K}$) and using B($\phi \rightarrow K^+ K^-$) = (48.9 ± 0.5)%.

² LEES 14H reports [$\Gamma(J/\psi(1S) \rightarrow \phi f'_2(1525)) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($f'_2(1525) \rightarrow K \bar{K}$)] = 7.2 ± 2.8 ± 0.3 eV which we divide by our best value B($f'_2(1525) \rightarrow K \bar{K}$) = (87.6 ± 2.2) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GZ2
NODE=M070GZ2

NODE=M070GZ2;LINKAGE=A

NODE=M070GZ2;LINKAGE=B

$$\Gamma(\phi K^+ K^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{130} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

4.60 ± 0.62 ± 0.05	163	¹ LEES	12F BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ K ⁺ K ⁻ γ
---------------------------	-----	-------------------	----------	--

¹ LEES 12F reports [$\Gamma(J/\psi(1S) \rightarrow \phi K^+ K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+ K^-$)] = 2.26 ± 0.26 ± 0.16 eV which we divide by our best value B($\phi(1020) \rightarrow K^+ K^-$) = (49.1 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G09
NODE=M070G09

NODE=M070G09;LINKAGE=A

$$\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{131} \Gamma_5 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

3.26 ± 0.84 ± 0.03	29	¹ LEES	14H BABR	e ⁺ e ⁻ → K _S ⁰ K _S ⁰ K ⁺ K ⁻ γ
---------------------------	----	-------------------	----------	---

¹ LEES 14H reports [$\Gamma(J/\psi(1S) \rightarrow \phi K_S^0 K_S^0) \times \Gamma(J/\psi(1S) \rightarrow e^+ e^-) / \Gamma_{\text{total}}$] × [B($\phi(1020) \rightarrow K^+ K^-$)] = 1.6 ± 0.4 ± 0.1 eV which we divide by our best value B($\phi(1020) \rightarrow K^+ K^-$) = (49.1 ± 0.5) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GZ1
NODE=M070GZ1

NODE=M070GZ1;LINKAGE=A

$\Gamma(f'_2(1525)K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{136}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8±1.9±0.1	16	1,2 LEES	14H BABR	$e^+e^- \rightarrow K_S^0 K_S^0 K^+K^-\gamma$

NODE=M070GZ4
NODE=M070GZ4

¹ Dividing by 1/4 to take into account $B(f'_2(1525) \rightarrow K_S^0 K_S^0) = 1/4 B(f'_2(1525) \rightarrow K\bar{K})$.

NODE=M070GZ4;LINKAGE=A

² LEES 14H reports $[\Gamma(J/\psi(1S) \rightarrow f'_2(1525)K^+K^-) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K\bar{K})] = 5.12 \pm 1.68 \pm 0.20$ eV which we divide by our best value $B(f'_2(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070GZ4;LINKAGE=B

 $\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{157}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
303±5±18	4990	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

NODE=M070G23
NODE=M070G23

 $\Gamma(\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{159}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
100 ±50 OUR AVERAGE				Error includes scale factor of 4.3. [104 ± 50 eV OUR 2023 AVERAGE Scale factor = 4.3]

NODE=M070P53
NODE=M070P53

NEW

55 ±16 ± 1	14k	¹ LEES	21 BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)3\pi^0\gamma$
150.0± 4.0±15.0	2.3k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-3\pi^0) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] = 19.2 \pm 4.5 \pm 3.2$ eV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}} = 0.3469 \pm 0.0034$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P53;LINKAGE=A

 $\Gamma(\pi^+\pi^-4\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{160}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.8±4.4±5.4	340	LEES	21C BABR	$e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

NODE=M070Q17
NODE=M070Q17

 $\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{161}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.0±9.0±8.0	1.2k	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

NODE=M070P55
NODE=M070P55

 $\Gamma(\rho^+\rho^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{162}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33.0±5.0±3.3	529	LEES	18E BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

NODE=M070P56
NODE=M070P56

 $\Gamma(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{163}\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.1248±0.0019±0.0026	LEES	21B BABR	$10.5 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$

NODE=M070G5
NODE=M070G5

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.122 ±0.005 ±0.008	AUBERT,B	04N BABR	$10.6 e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma$
---------------------	----------	----------	---

 $\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{164}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.9±0.5±1.0	761	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

NODE=M070G7
NODE=M070G7

 $\Gamma(\pi^+\pi^-\pi^0K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{165}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
107.0±4.3±6.4	768	AUBERT	07AU BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0\gamma$

NODE=M070G27
NODE=M070G27

 $\Gamma(2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{167}\Gamma_5/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.4±0.9±0.4		LEES	12E BABR	$10.6 e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$

NODE=M070G11
NODE=M070G11

• • • We do not use the following data for averages, fits, limits, etc. • • •

19.5±1.4±1.3	270	¹ AUBERT	05D BABR	$10.6 e^+e^- \rightarrow 2(\pi^+\pi^-)\gamma$
--------------	-----	---------------------	----------	---

¹ Superseded by LEES 12E.

NODE=M070G11;LINKAGE=AU

 $\Gamma(3(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{168}\Gamma_5/\Gamma$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.37±0.16±0.14	496	AUBERT	06D BABR	$10.6 e^+e^- \rightarrow 3(\pi^+\pi^-)\gamma$

NODE=M070G6
NODE=M070G6

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{169}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
350±50 OUR AVERAGE		[345 ± 50 eV OUR 2023 AVERAGE]		
345±10±50	14k	LEES 21	BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

NODE=M070Q12
NODE=M070Q12
NEW

$$\Gamma(2(\pi^+\pi^-)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{171}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1±2.4±0.1	85	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ

NODE=M070G26
NODE=M070G26

¹AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → 2γ)] = 5.16 ± 0.85 ± 0.39 eV which we divide by our best value B(η → 2γ) = (39.36 ± 0.18) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G26;LINKAGE=AU

$$\Gamma(2(\pi^+\pi^-\pi^0)\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{173}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±2.6±1.4	14k	LEES 21	BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

NODE=M070Q15
NODE=M070Q15

$$\Gamma(\pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{174}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
13.1± 2.7 OUR AVERAGE				

26.1±17.9±0.3	14k	¹ LEES 21	BABR	10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ
12.8± 1.8±2.0	203	LEES 18E	BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ π ⁰ ηγ

¹LEES 21 reports [$\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] × [B(η → π⁺π⁻π⁰)] = 6 ± 4 ± 1 eV which we divide by our best value B(η → π⁺π⁻π⁰) = (23.02 ± 0.25) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070P57
NODE=M070P57

NODE=M070P57;LINKAGE=A

$$\Gamma(\rho^\pm\pi^\mp\pi^0\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{175}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.5±4.1±1.6	168	LEES 18E	BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ π ⁰ ηγ

NODE=M070P59
NODE=M070P59

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{176}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78±0.11±0.05	462	¹ LEES 15J	BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.94±0.11±0.05	462	² LEES 15J	BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ
1.42±0.23±0.08	51	³ LEES 13Q	BABR	e ⁺ e ⁻ → K ⁺ K ⁻ γ

NODE=M070G08
NODE=M070G08

OCCUR=2

¹ sin φ > 0.

² sin φ < 0.

³ Interference with non-resonant K⁺K⁻ production not taken into account.

NODE=M070G08;LINKAGE=A
NODE=M070G08;LINKAGE=B
NODE=M070G08;LINKAGE=BA

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{182}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.4±1.3±0.6	182	LEES 17A	BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ

NODE=M070G41
NODE=M070G41

$$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{183}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.9±0.4	106	LEES 17A	BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ

NODE=M070G42
NODE=M070G42

$$\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K_L^0 \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{184}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.7±0.1	37	LEES 17A	BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ

NODE=M070G43
NODE=M070G43

$$\Gamma(K_S^0 K^*(892)^0 \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{201}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
9.45±3.15±0.90	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M070Q56
NODE=M070Q56

$$\Gamma(K_S^0 K^*(892)^0 \pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{202}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
5.59±0.79±0.55	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M070Q61
NODE=M070Q61

$$\Gamma(K^\mp K^*(892)^\pm \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{203}\Gamma_5/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
18.6±6.3±1.8	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M070Q57
NODE=M070Q57

$\Gamma(K^\mp K^*(892)^\pm \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{205}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
8.67±1.56±0.84		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q60 NODE=M070Q60
$\Gamma(K^*(892)^\pm K^*(892)^0 \pi^\mp) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{204}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
26.6±4.5±2.7		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q58 NODE=M070Q58
$\Gamma(K^*(892)^+ K^*(892)^- \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{206}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
62.1±10.8±6.30		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q59 NODE=M070Q59
$\Gamma(K^+ K^- \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{185}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
37.94±0.81±1.10	3.1k	LEES	12F	BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$		NODE=M070G12 NODE=M070G12
• • • We do not use the following data for averages, fits, limits, etc. • • •						
36.3 ±1.3 ±2.1	1.5k	¹ AUBERT	07AK	BABR 10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$		
33.6 ±2.7 ±2.7	233	² AUBERT	05D	BABR 10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \gamma$		
¹ Superseded by LEES 12F.						
² Superseded by AUBERT 07AK.						
$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{186}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
11.75±0.81±0.90	388	LEES	12F	BABR 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$		NODE=M070G04 NODE=M070G04
• • • We do not use the following data for averages, fits, limits, etc. • • •						
13.6 ±1.1 ±1.3	203	¹ AUBERT	07AK	BABR 10.6 $e^+ e^- \rightarrow \pi^0 \pi^0 K^+ K^- \gamma$		
¹ Superseded by LEES 12F.						
$\Gamma(K^+ K^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{187}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
8.9±1.3±0.9		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q51 NODE=M070Q51
$\Gamma(K_S^0 K_L^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{188}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
29.3±2.6±2.9		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q52 NODE=M070Q52
$\Gamma(K_S^0 K_L^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{189}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
34.6±1.4±1.8		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q53 NODE=M070Q53
$\Gamma(K_S^0 K_L^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{191}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
20.8±2.3±2.1	248	LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_L^0 \gamma$		NODE=M070GY1 NODE=M070GY1
$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{192}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
10.3±2.3±0.5	47	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \pi^0 \pi^0 \gamma$		NODE=M070G40 NODE=M070G40
$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{193}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
8.0±1.8±0.4	45	LEES	17A	BABR $e^+ e^- \rightarrow K_S^0 K_L^0 \eta \gamma$		NODE=M070G35 NODE=M070G35
$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{194}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
9.3±0.9±0.5	133	LEES	14H	BABR $e^+ e^- \rightarrow \pi^+ \pi^- K_S^0 K_S^0 \gamma$		NODE=M070GY2 NODE=M070GY2
$\Gamma(K^\mp K_S^0 \pi^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{195}\Gamma_5/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT		
31.7±1.9±1.8	393	LEES	17D	BABR $e^+ e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0 \gamma$		NODE=M070G34 NODE=M070G34
$\Gamma(K_S^0 K^\pm \rho(770)^\pm \pi^0) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_{190}\Gamma_5/\Gamma$	
VALUE (eV)		DOCUMENT ID	TECN	COMMENT		
16.0±4.1±1.6		LEES	23	BABR $e^+ e^- \rightarrow \gamma_{ISR}$ hadrons		NODE=M070Q62 NODE=M070Q62

$$\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{197}\Gamma_5/\Gamma$$

VALUE (10 ⁻² keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.75±0.23±0.17	205	AUBERT	06D BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ 2(π ⁺ π ⁻)γ

NODE=M070G9
NODE=M070G9

$$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{198}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
25.9±3.9±0.1	73	¹ AUBERT	07AU BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ ηγ

NODE=M070G30
NODE=M070G30

¹AUBERT 07AU reports [$\Gamma(J/\psi(1S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(J/\psi(1S) \rightarrow e^+e^-) / \Gamma_{\text{total}} \times [B(\eta \rightarrow 2\gamma)] = 10.2 \pm 1.3 \pm 0.8$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070G30;LINKAGE=AU

$$\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{199}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.00±0.33±0.29	287 ± 24	LEES	12F BABR	10.6 e ⁺ e ⁻ → 2(K ⁺ K ⁻)γ

NODE=M070G13
NODE=M070G13

• • • We do not use the following data for averages, fits, limits, etc. • • •
4.11±0.39±0.30 156 ± 15 ¹AUBERT 07AK BABR 10.6 e⁺e⁻ → 2(K⁺K⁻)γ
4.0 ±0.7 ±0.6 38 ²AUBERT 05D BABR 10.6 e⁺e⁻ → 2(K⁺K⁻)γ

NODE=M070G13;LINKAGE=A
NODE=M070G13;LINKAGE=AU

¹Superseded by LEES 12F.
²Superseded by AUBERT 07AK.

$$\Gamma(K^+K^-K_S^0K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{200}\Gamma_5/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.3±0.4±0.1	29	LEES	14H BABR	e ⁺ e ⁻ → K _S ⁰ K _S ⁰ K ⁺ K ⁻ γ

NODE=M070GY3
NODE=M070GY3

$$\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{207}\Gamma_5/\Gamma$$


VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.9±0.6 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.			

NODE=M070G4
NODE=M070G4

11.3±0.4±0.3 821 ¹LEES 13O BABR e⁺e⁻ → p \bar{p} γ
12.9±0.4±0.4 918 ²LEES 13Y BABR e⁺e⁻ → p \bar{p} γ
9.7±1.7 ³ARMSTRONG 93B E760 p \bar{p} → e⁺e⁻
• • • We do not use the following data for averages, fits, limits, etc. • • •
12.0±0.6±0.5 438 ⁴AUBERT 06B BABR e⁺e⁻ → p \bar{p} γ

NODE=M070G4;LINKAGE=B
NODE=M070G4;LINKAGE=C
NODE=M070G;LINKAGE=A
NODE=M070G4;LINKAGE=A

¹ISR photon reconstructed in the detector
²ISR photon undetected
³Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.
⁴Superseded by LEES 13O



$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{223}\Gamma_5/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
10.7±0.9±0.7	AUBERT	07BD	BABR	10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$	NODE=M070G16 NODE=M070G16

$\Gamma(\Sigma^+\bar{\Sigma}^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{233}\Gamma_5/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.8±1.5±0.8	GONG	23	BELL	$e^+e^- \rightarrow \Sigma^+\bar{\Sigma}^-$	NODE=M070Q50 NODE=M070Q50

$\Gamma(\Sigma^0\bar{\Sigma}^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$				$\Gamma_{234}\Gamma_5/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
5.2±1.5±0.6	GONG	23	BELL	$e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0$	NODE=M070G17 NODE=M070G17
6.4±1.2±0.6	AUBERT	07BD	BABR	10.6 $e^+e^- \rightarrow \Sigma^0\bar{\Sigma}^0\gamma$	OCCUR=2

$J/\psi(1S)$ BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ above.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				Γ_1/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.877±0.005 OUR AVERAGE					NODE=M070R3 NODE=M070R3
0.878±0.005	BAI	95B	BES	e^+e^-	
0.86 ±0.02	BOYARSKI	75	MRK1	e^+e^-	

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$				Γ_2/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.1346±0.0007 OUR AVERAGE	[0.135 ± 0.003 OUR 2023 AVERAGE]				NODE=M070R4 NODE=M070R4 NEW
0.1346±0.0007	¹ LIAO	23	RVUE	e^+e^-	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.135 ±0.003	^{2,3} SETH	04	RVUE	e^+e^-	
0.17 ±0.02	² BOYARSKI	75	MRK1	e^+e^-	
¹ Using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (5.967 \pm 0.023)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.					NODE=M070R4;LINKAGE=A
² Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.					NODE=M070R4;LINKAGE=C
³ Using $B(J/\psi \rightarrow \ell^+\ell^-) = (5.90 \pm 0.09)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.					NODE=M070R4;LINKAGE=SE

$\Gamma(ggg)/\Gamma_{\text{total}}$				Γ_3/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
64.1±1.0	6 M	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- + \text{hadrons}$	NODE=M070S65 NODE=M070S65
¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the PDG 08 values of $B(\ell^+\ell^-)$, $B(\text{virtual } \gamma \rightarrow \text{hadrons})$, and $B(\gamma\eta_c)$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 08.					NODE=M070S65;LINKAGE=BE

$\Gamma(\gamma gg)/\Gamma_{\text{total}}$				Γ_4/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
8.79±1.05	200 k	¹ BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^-\gamma + \text{hadrons}$	NODE=M070S66 NODE=M070S66
¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = 0.137 \pm 0.001 \pm 0.016 \pm 0.004$ from BESSON 08 and the value of $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 08.					NODE=M070S66;LINKAGE=BE

$\Gamma(\gamma gg)/\Gamma(ggg)$				Γ_4/Γ_3	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
13.7±0.1±0.7	6 M	BESSON	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$	NODE=M070S67 NODE=M070S67

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_5/Γ VALUE (units 10^{-2}) EVTS**5.971 ± 0.032 OUR AVERAGE**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.983 ± 0.007 ± 0.037	720k	ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.945 ± 0.067 ± 0.042	15k	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.90 ± 0.05 ± 0.10		BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.09 ± 0.33		BAI	95B BES	$e^+ e^-$
5.92 ± 0.15 ± 0.20		COFFMAN	92 MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75 MRK1	$e^+ e^-$

NODE=M070R1
NODE=M070R1 $\Gamma(e^+e^-\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ VALUE (units 10^{-3})**8.8 ± 1.3 ± 0.4**¹ For $E_\gamma > 100$ MeV.

DOCUMENT ID	TECN	COMMENT
¹ ARMSTRONG 96	E760	$\bar{p} p \rightarrow e^+ e^- \gamma$

NODE=M070S33
NODE=M070S33

NODE=M070S33;LINKAGE=A

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_7/Γ VALUE (units 10^{-2}) EVTS**5.961 ± 0.033 OUR AVERAGE**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.973 ± 0.007 ± 0.038	770k	ABLIKIM	13R BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.960 ± 0.065 ± 0.050	17k	LI	05C CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
5.84 ± 0.06 ± 0.10		BAI	98D BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.08 ± 0.33		BAI	95B BES	$e^+ e^-$
5.90 ± 0.15 ± 0.19		COFFMAN	92 MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
6.9 ± 0.9		BOYARSKI	75 MRK1	$e^+ e^-$

NODE=M070R2
NODE=M070R2 $\Gamma(e^+e^-)/\Gamma(\mu^+\mu^-)$ Γ_5/Γ_7

VALUE

1.0016 ± 0.0031 OUR AVERAGE

VALUE	DOCUMENT ID	TECN	COMMENT
1.0022 ± 0.0044 ± 0.0048	¹ AULCHENKO 14	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
1.0017 ± 0.0017 ± 0.0033	² ABLIKIM 13R	BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.002 ± 0.021 ± 0.013	³ ANASHIN 10	KEDR	3.097 $e^+ e^- \rightarrow e^+ e^-, \mu^+ \mu^-$
0.997 ± 0.012 ± 0.006	LI 05C	CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.011 ± 0.013 ± 0.016	BAI 98D	BES	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
1.00 ± 0.07	BAI 95B	BES	$e^+ e^-$
1.00 ± 0.05	BOYARSKI 75	MRK1	$e^+ e^-$
0.91 ± 0.15	ESPOSITO 75B	FRAM	$e^+ e^-$
0.93 ± 0.10	FORD 75	SPEC	$e^+ e^-$

NODE=M070R5
NODE=M070R5¹ From 235.3k $J/\psi \rightarrow e^+ e^-$ and 156.6k $J/\psi \rightarrow \mu^+ \mu^-$ observed events.² Not independent of the corresponding measurements of $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$.³ Not independent of the corresponding measurements of $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ and $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$.NODE=M070R5;LINKAGE=A
NODE=M070R5;LINKAGE=AB

NODE=M070R5;LINKAGE=AN

HADRONS DECAYS

NODE=M070305

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_8/Γ VALUE (units 10^{-2}) EVTS**1.88 ± 0.12 OUR AVERAGE**

Error includes scale factor of 2.6. See the ideogram below.

[(1.69 ± 0.15) × 10⁻² OUR 2023 AVERAGE Scale factor = 2.4]

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.072 ± 0.017 ± 0.062	19.8k	¹ ANASHIN 23	KEDR	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.18 ± 0.19		^{2,3} AUBERT,B 04N	BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma$
2.184 ± 0.005 ± 0.201	220k	^{3,4} BAI 04H	BES	$e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$
2.091 ± 0.021 ± 0.116		^{3,5} BAI 04H	BES	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$
1.21 ± 0.20		BAI 96D	BES	$e^+ e^- \rightarrow \rho\pi$
1.42 ± 0.01 ± 0.19		COFFMAN 88	MRK3	$e^+ e^-$
1.3 ± 0.3	150	FRANKLIN 83	MRK2	$e^+ e^-$
1.6 ± 0.4	183	ALEXANDER 78	PLUT	$e^+ e^-$
1.33 ± 0.21		BRANDELIK 78B	DASP	$e^+ e^-$
1.0 ± 0.2	543	BARTEL 76	CNTR	$e^+ e^-$
1.3 ± 0.3	153	JEAN-MARIE 76	MRK1	$e^+ e^-$

NODE=M070R20
NODE=M070R20


NEW

OCCUR=2

- ¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.
- ² From the ratio of $\Gamma(e^+e^-)B(\pi^+\pi^-\pi^0)$ and $\Gamma(e^+e^-)B(\mu^+\mu^-)$ (AUBERT 04).
- ³ Not independent of their $B(\pi^+\pi^-\pi^0)$.
- ⁴ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.
- ⁵ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.

NODE=M070R20;LINKAGE=A

NODE=M070R20;LINKAGE=AU
 NODE=M070R20;LINKAGE=BU
 NODE=M070R20;LINKAGE=BA
 NODE=M070R20;LINKAGE=BI

 $\Gamma(\rho\pi)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_8/Γ_{163}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$1.142 \pm 0.011 \pm 0.026$	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.331 ± 0.033	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

NODE=M070P18
 NODE=M070P18

OCCUR=2

¹ From a Dalitz plot analysis in an isobar model.² From a Dalitz plot analysis in a Veneziano model.

NODE=M070P18;LINKAGE=A
 NODE=M070P18;LINKAGE=B

 $\Gamma(\rho^0\pi^0)/\Gamma(\rho\pi)$ Γ_9/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
$0.328 \pm 0.005 \pm 0.027$	COFFMAN 88	MRK3	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.35 ± 0.08	ALEXANDER 78	PLUT	e^+e^-
0.32 ± 0.08	BRANDELIK 78B	DASP	e^+e^-
0.39 ± 0.11	BARTEL 76	CNTR	e^+e^-
0.37 ± 0.09	JEAN-MARIE 76	MRK1	e^+e^-

NODE=M070R21
 NODE=M070R21

 $\Gamma(\rho^+K^+K^-\pi^- + c.c \rightarrow K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.53 \pm 0.16 \pm 0.81$	485	ANASHIN 22	KEDR	$J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M070Q41
 NODE=M070Q41

 $\Gamma(a_2(1320)^0\pi^+\pi^- \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.84 \pm 0.08 \pm 0.60$	1317	ANASHIN 22	KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M070Q39
 NODE=M070Q39

 $\Gamma(a_2(1320)^+\pi^-\pi^0 + c.c \rightarrow 2(\pi^+\pi^-\pi^0))/\Gamma_{total}$ Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.67 \pm 0.09 \pm 0.73$	1628	ANASHIN 22	KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M070Q40
 NODE=M070Q40

$\Gamma(a_2(1320)\rho)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS**10.9±2.2 OUR AVERAGE**

11.7±0.7±2.5 7584

8.4±4.5 36

DOCUMENT ID TECN COMMENT

AUGUSTIN 89 DM2 $J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$ VANNUCCI 77 MRK1 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0$ Γ_{12}/Γ NODE=M070R43
NODE=M070R43 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**3.78±0.68**

471

DOCUMENT ID TECN COMMENT

¹ ABLIKIM 19Q BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$ ¹ From an energy scan of $e^+ e^- \rightarrow J/\psi \rightarrow \eta\pi^+\pi^-$ assuming PDG 16 values for $\Gamma(e^+ e^-)$, $\Gamma(\mu^+ \mu^-)$, and $\Gamma(\text{total})$. Γ_{13}/Γ NODE=M070P81
NODE=M070P81

NODE=M070P81;LINKAGE=A

 $\Gamma(\eta\rho)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS**0.193±0.023 OUR AVERAGE**

0.194±0.017±0.029 299

0.193±0.013±0.029

DOCUMENT ID TECN COMMENT

JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$ COFFMAN 88 MRK3 $e^+ e^- \rightarrow \pi^+ \pi^- \eta$ Γ_{16}/Γ NODE=M070R22
NODE=M070R22 $\Gamma(\eta\phi(2170) \rightarrow \eta\phi f_0(980) \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**1.20±0.14±0.37**

471

DOCUMENT ID TECN COMMENT

ABLIKIM 15H BES3 $e^+ e^- \rightarrow J/\psi \rightarrow \phi\eta\pi^+\pi^-$ Γ_{17}/Γ NODE=M070B12
NODE=M070B12 $\Gamma(\eta\phi(2170) \rightarrow \eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$

VALUE CL%

<2.52 × 10⁻⁴

90

DOCUMENT ID TECN COMMENT

ABLIKIM 10C BES2 $J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$ Γ_{18}/Γ NODE=M070S70
NODE=M070S70 $\Gamma(\eta K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**21.8±2.2±3.4**

232 ± 23

DOCUMENT ID TECN COMMENT

ABLIKIM 08E BES2 $e^+ e^- \rightarrow J/\psi$ Γ_{20}/Γ NODE=M070S57
NODE=M070S57 $\Gamma(\eta K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-3}) EVTS**1.15±0.13±0.22**

209

DOCUMENT ID TECN COMMENT

ABLIKIM 10C BES2 $J/\psi \rightarrow \eta K^+ \pi^- K^- \pi^+$ Γ_{21}/Γ NODE=M070S69
NODE=M070S69 $\Gamma(\rho\eta'(958))/\Gamma_{\text{total}}$ VALUE (units 10^{-5}) EVTS**8.1 ± 0.8 OUR AVERAGE** Error includes scale factor of 1.6.

7.90±0.19±0.49 3476

8.3 ± 3.0 ± 1.2 19

11.4 ± 1.4 ± 1.6

DOCUMENT ID TECN COMMENT

¹ ABLIKIM 17AK BES3 $J/\psi \rightarrow \pi^+ \pi^- \eta'$ JOUSSET 90 DM2 $J/\psi \rightarrow \text{hadrons}$ COFFMAN 88 MRK3 $J/\psi \rightarrow \pi^+ \pi^- \eta'$ Γ_{22}/Γ NODE=M070R23
NODE=M070R23

NODE=M070R23;LINKAGE=A

 $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ VALUE (units 10^{-4}) EVTS**2.2±0.2±1.1**

19.8k

DOCUMENT ID TECN COMMENT

¹ ANASHIN 23 KEDR $e^+ e^- \rightarrow J/\psi \rightarrow \pi^+ \pi^- \pi^0$ ¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0 \pi^0$, $J/\psi \rightarrow \rho^+ \pi^-$, $J/\psi \rightarrow \rho^- \pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered. Γ_{30}/Γ NODE=M070Q49
NODE=M070Q49

NODE=M070Q49;LINKAGE=A

 $\Gamma(\rho(1450)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$

VALUE (%) EVTS

10.9 ± 1.7 ± 2.7 20k

0.80±0.27 20k

DOCUMENT ID TECN COMMENT

¹ LEES 17C BABR $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ ² LEES 17C BABR $J/\psi \rightarrow \pi^+ \pi^- \pi^0$ Γ_{30}/Γ_{163} NODE=M070P25
NODE=M070P25

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a Dalitz plot analysis in an isobar model.² From a Dalitz plot analysis in a Veneziano model.

OCCUR=2

NODE=M070P25;LINKAGE=A
NODE=M070P25;LINKAGE=B $\Gamma(\rho(1450)^\pm \pi^\mp \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$

VALUE (%) EVTS

6.3±0.8±0.6

4k

DOCUMENT ID TECN COMMENT

¹ LEES 17C BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$ Γ_{31}/Γ_{181} NODE=M070P31
NODE=M070P31¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P31;LINKAGE=A

$$\Gamma(\rho(1450)^0 \pi^0 \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0) \quad \Gamma_{32} / \Gamma_{180}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
9.3 ± 2.0 ± 0.6	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P27
NODE=M070P27

NODE=M070P27;LINKAGE=A

$$\Gamma(\rho(1450) \eta'(958) \rightarrow \pi^+ \pi^- \eta'(958)) / \Gamma_{\text{total}} \quad \Gamma_{33} / \Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
3.28 ± 0.55 ± 0.44	119	¹ ABLIKIM	17AK BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta'$

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+ \pi^- \eta'$.

NODE=M070P36
NODE=M070P36

NODE=M070P36;LINKAGE=A

$$\Gamma(\rho(1700) \pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{35} / \Gamma_{163}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 2 ± 5	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
22 ± 6	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

NODE=M070P21
NODE=M070P21

OCCUR=2

NODE=M070P21;LINKAGE=A

NODE=M070P21;LINKAGE=B

$$\Gamma(\rho(2150) \pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{37} / \Gamma_{163}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4 ± 1 ± 20	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
600 ± 250	20k	² LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in an isobar model.

² From a Dalitz plot analysis in a Veneziano model.

NODE=M070P22
NODE=M070P22

OCCUR=2

NODE=M070P22;LINKAGE=A

NODE=M070P22;LINKAGE=B

$$\Gamma(\rho_3(1690) \pi \rightarrow \pi^+ \pi^- \pi^0) / \Gamma(\pi^+ \pi^- \pi^0) \quad \Gamma_{38} / \Gamma_{163}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
• • •				We do not use the following data for averages, fits, limits, etc. • • •
4.0 ± 0.8	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+ \pi^- \pi^0$

¹ From a Dalitz plot analysis in a Veneziano model.

NODE=M070P24
NODE=M070P24

NODE=M070P24;LINKAGE=A


$$\Gamma(\omega \pi^0) / \Gamma_{\text{total}} \quad \Gamma_{39} / \Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.538 ± 0.012 ± 0.065	2090	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega \pi^0$
0.360 ± 0.028 ± 0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
0.482 ± 0.019 ± 0.064		COFFMAN	88 MRK3	$e^+ e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

¹ Using $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.1 \pm 0.7)\%$.

NODE=M070R32
NODE=M070R32

NODE=M070R32;LINKAGE=BL



$\Gamma(\omega\pi^0 \rightarrow \pi^+\pi^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ Γ_{40}/Γ_{163}

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
8±3±2	20k	¹ LEES	17C BABR	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

¹ From a Dalitz plot analysis in an isobar model and significance 4.9 σ .

NODE=M070P23
NODE=M070P23

NODE=M070P23;LINKAGE=A

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{total}$ Γ_{41}/Γ


VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±1.0 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
10.6±1.2±0.1	3531	¹ ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.0±1.6	18058	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$
7.8±1.6	215	BURMESTER	77D PLUT	e^+e^-
6.8±1.9	348	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega\pi^+\pi^-)/\Gamma_{total}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.946 \pm 0.016 \pm 0.108) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R24
NODE=M070R24

NODE=M070R24;LINKAGE=A

WEIGHTED AVERAGE
8.5±1.0 (Error scaled by 1.3)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\omega\pi^0\pi^0)/\Gamma_{total}$ Γ_{42}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.4±0.3±0.7	509	AUGUSTIN	89 DM2	$J/\psi \rightarrow \pi^+\pi^-\pi^0$

NODE=M070S26
NODE=M070S26

$\Gamma(\omega f_2(1270))/\Gamma_{total}$ Γ_{44}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±0.6 OUR AVERAGE				
4.3±0.2±0.6	5860	AUGUSTIN	89 DM2	e^+e^-
4.0±1.6	70	BURMESTER	77D PLUT	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1.9±0.8	81	VANNUCCI	77 MRK1	$e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M070R28
NODE=M070R28

$\Gamma(\omega\eta)/\Gamma_{total}$ Γ_{45}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.74 ±0.20 OUR AVERAGE		Error includes scale factor of 1.6. See the ideogram below.		
2.352±0.273	5k	¹ ABLIKIM	06F BES2	$J/\psi \rightarrow \omega\eta$
1.44 ±0.40 ±0.14	13	² AUBERT	06D BABR	$10.6 e^+e^- \rightarrow \omega\eta\gamma$
1.43 ±0.10 ±0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.71 ±0.08 ±0.20		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta$


NODE=M070R30
NODE=M070R30

¹ Using $B(\eta \rightarrow 2\gamma) = (39.43 \pm 0.26)\%$, $B(\eta \rightarrow \pi^+\pi^-\pi^0) = 22.6 \pm 0.4\%$, $B(\eta \rightarrow \pi^+\pi^-\gamma) = 4.68 \pm 0.11\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

² Using $\Gamma(J/\psi \rightarrow e^+e^-) = 5.52 \pm 0.14 \pm 0.04$ keV.

NODE=M070R30;LINKAGE=BL

NODE=M070R30;LINKAGE=EE



$\Gamma(\omega\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
85±34	140	VANNUCCI	77	MRK1 $e^+e^- \rightarrow 3(\pi^+\pi^-)\pi^0$

NODE=M070R26
NODE=M070R26

$\Gamma(\omega\eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.12±0.02±0.13	14k	¹ ABLIKIM	19AC	BES3 $J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

NODE=M070P83
NODE=M070P83

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.

NODE=M070P83;LINKAGE=A

$\Gamma(\omega\eta'(958))/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89±0.18 OUR AVERAGE				
2.08±0.30±0.14	137	¹ ABLIKIM	17AK	BES3 $J/\psi \rightarrow \pi^+\pi^-\eta'$
2.26±0.43	218	² ABLIKIM	06F	BES2 $J/\psi \rightarrow \omega\eta'$
1.8 $^{+1.0}_{-0.8}$ ±0.3	6	JOUSSET	90	DM2 $J/\psi \rightarrow$ hadrons
1.66±0.17±0.19		COFFMAN	88	MRK3 $e^+e^- \rightarrow 3\pi\eta'$

NODE=M070R31
NODE=M070R31

¹ From a partial wave analysis of the decay $J/\psi \rightarrow \pi^+\pi^-\eta'$.

² Using $B(\eta' \rightarrow \pi^+\pi^-\eta) = (44.3 \pm 1.5)\%$, $B(\eta' \rightarrow \pi^+\pi^-\gamma) = 29.5 \pm 1.0\%$, $B(\eta \rightarrow 2\gamma) = 39.43 \pm 0.26\%$, and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.1 \pm 0.7)\%$.

NODE=M070R31;LINKAGE=A
NODE=M070R31;LINKAGE=BL

$\Gamma(\omega f_0(980))/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.41±0.27±0.47	¹ AUGUSTIN	89	DM2 $J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M070S27
NODE=M070S27

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

NODE=M070S27;LINKAGE=K

$\Gamma(\omega f_0(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
4.8±1.1±0.3	^{1,2} FALVARD	88	DM2 $J/\psi \rightarrow$ hadrons

NODE=M070S25
NODE=M070S25

¹ Includes unknown branching fraction $f_0(1710) \rightarrow K\bar{K}$.

² Addition of $f_0(1710) \rightarrow K^+K^-$ and $f_0(1710) \rightarrow K^0\bar{K}^0$ branching ratios.

NODE=M070S25;LINKAGE=F
NODE=M070S25;LINKAGE=G

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8$^{+1.9}_{-1.6}$±1.7	111 $^{+31}_{-26}$	BECKER	87	MRK3 $e^+e^- \rightarrow$ hadrons

NODE=M070S5
NODE=M070S5

$\Gamma(\omega f'_2(1525))/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-4}$	90	¹ VANNUCCI 77	MRK1	$e^+e^- \rightarrow \pi^+\pi^-\pi^0 K^+K^-$

••• We do not use the following data for averages, fits, limits, etc. •••

$<2.8 \times 10^{-4}$	90	¹ FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
-----------------------	----	-------------------------	-----	-------------------------------------

¹ Re-evaluated assuming $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

NODE=M070R29
NODE=M070R29

NODE=M070R29;LINKAGE=C

 $\Gamma(\omega X(1835) \rightarrow \omega p\bar{p})/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-6}$	95	ABLIKIM 13P	BES3	$J/\psi \rightarrow \gamma\pi^0 p\bar{p}$

NODE=M070S81
NODE=M070S81

 $\Gamma(\omega X(1835), X \rightarrow \eta'\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-5}$	¹ ABLIKIM 19AC	BES3	$J/\psi \rightarrow \omega\eta'\pi^+\pi^-$

¹ Using the decays $\omega \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$.

NODE=M070P84
NODE=M070P84

NODE=M070P84;LINKAGE=A

 $\Gamma(\omega K^+K^-)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.52 \pm 0.30 \pm 0.01$	276	¹ ANASHIN 22	KEDR	$J/\psi \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹ ANASHIN 22 reports $[\Gamma(J/\psi(1S) \rightarrow \omega K^+K^-)/\Gamma_{\text{total}}] \times [B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)] = (0.136 \pm 0.008 \pm 0.026) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070Q38
NODE=M070Q38

NODE=M070Q38;LINKAGE=A

 $\Gamma(\omega K^\pm K_S^0 \pi^\mp)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
34 ± 5 OUR AVERAGE				
$37.7 \pm 0.8 \pm 5.8$	1972 ± 41	ABLIKIM 08E	BES2	$e^+e^- \rightarrow J/\psi$
$29.5 \pm 1.4 \pm 7.0$	879 ± 41	BECKER 87	MRK3	$e^+e^- \rightarrow \text{hadrons}$

NODE=M070S1
NODE=M070S1

 $\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19 ± 4 OUR AVERAGE				
$19.8 \pm 2.1 \pm 3.9$		¹ FALVARD 88	DM2	$J/\psi \rightarrow \text{hadrons}$
16 ± 10	22	FELDMAN 77	MRK1	e^+e^-

¹ Addition of ωK^+K^- and $\omega K^0\bar{K}^0$ branching ratios.

NODE=M070R27
NODE=M070R27

NODE=M070R27;LINKAGE=B

 $\Gamma(\omega K^*(892)\bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
61 ± 9 OUR AVERAGE				
$62.0 \pm 6.8 \pm 10.6$	899 ± 98	ABLIKIM 08E	BES2	$J/\psi \rightarrow \omega K_S^0 K^\pm \pi^\mp$
$65.3 \pm 10.2 \pm 13.5$	176 ± 28	ABLIKIM 08E	BES2	$J/\psi \rightarrow \omega K^+K^-\pi^0$
$53 \pm 14 \pm 14$	530 ± 140	BECKER 87	MRK3	$e^+e^- \rightarrow \text{hadrons}$

NODE=M070S2
NODE=M070S2

OCCUR=2

 $\Gamma(\eta' K^{*\pm} K^\mp)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.48 ± 0.13 OUR AVERAGE			
$1.50 \pm 0.02 \pm 0.19$	¹ ABLIKIM 18AB	BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$
$1.47 \pm 0.03 \pm 0.17$	² ABLIKIM 18AB	BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+K^-\pi^0$.

² From $\eta' K_S^0 K^\pm \pi^\mp$.

NODE=M070P48
NODE=M070P48

OCCUR=2

NODE=M070P48;LINKAGE=A

NODE=M070P48;LINKAGE=B

 $\Gamma(\eta' K^{*0} \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.66 ± 0.03 ± 0.21	¹ ABLIKIM 18AB	BES3	$J/\psi \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

NODE=M070P49
NODE=M070P49

NODE=M070P49;LINKAGE=A

 $\Gamma(\eta' h_1(1415) \rightarrow \eta' K^* \bar{K} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.16 ± 0.12 ± 0.29	1.1k	¹ ABLIKIM 18AB	BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K_S^0 K^\pm \pi^\mp$.

NODE=M070P52
NODE=M070P52

NODE=M070P52;LINKAGE=A

$$\Gamma(\eta' h_1(1415) \rightarrow \eta' K^{*\pm} K^\mp) / \Gamma_{\text{total}} \quad \Gamma_{65} / \Gamma$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.51 \pm 0.09 \pm 0.21$	1.0k	¹ ABLIKIM	18AB BES3	$J/\psi \rightarrow \eta' h_1 \rightarrow \eta' K^* \bar{K}$

¹ From $\eta' K^+ K^- \pi^0$.

NODE=M070P44
NODE=M070P44

NODE=M070P44;LINKAGE=A

$$\Gamma(\eta' h_1(1415) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}} \quad \Gamma_{66} / \Gamma$$

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$4.69 \pm 0.80^{+0.74}_{-1.82}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q28
NODE=M070Q28

NODE=M070Q28;LINKAGE=A

$$\Gamma(h_1(1415) \eta' \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}} \quad \Gamma_{27} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.08 \pm 0.01^{+0.01}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

NODE=M070Q33
NODE=M070Q33

NODE=M070Q33;LINKAGE=A

$$\Gamma(h_1(1595) \eta' \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}} \quad \Gamma_{28} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.16 \pm 0.02^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

NODE=M070Q34
NODE=M070Q34

NODE=M070Q34;LINKAGE=A

$$\Gamma(\bar{K} K^*(892) + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp) / \Gamma(K_S^0 K^\pm \pi^\mp) \quad \Gamma_{68} / \Gamma_{181}$$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$90.5 \pm 0.9 \pm 3.8$	4k	¹ LEES	17C BABR	$J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P30
NODE=M070P30

NODE=M070P30;LINKAGE=A

$$\Gamma(K^+ K^*(892)^- + \text{c.c.}) / \Gamma_{\text{total}} \quad \Gamma_{69} / \Gamma$$


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.0 $^{+0.8}_{-1.0}$ OUR AVERAGE Error includes scale factor of 2.9. See the ideogram below.

$8.07 \pm 0.04^{+0.38}_{-0.61}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$
$4.57 \pm 0.17 \pm 0.70$	2285	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$5.26 \pm 0.13 \pm 0.53$		COFFMAN	88 MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp,$ $K^+ K^- \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.6 ± 0.6	24	FRANKLIN	83 MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77 MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76 DASP	$J/\psi \rightarrow K^\pm X$



$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma_{total}$ Γ_{70} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.69 \pm 0.01 \pm 0.20$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P79
NODE=M070P79

$\Gamma(K^+ K^*(892)^- + c.c. \rightarrow K^+ K^- \pi^0) / \Gamma(K^+ K^- \pi^0)$ $\Gamma_{70} / \Gamma_{180}$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$92.4 \pm 1.5 \pm 3.4$	2k	¹ LEES	17C BABR	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P26
NODE=M070P26

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P26;LINKAGE=A

$\Gamma(K^0 \bar{K}^*(892)^0 + c.c.) / \Gamma_{total}$ Γ_{72} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.2 ± 0.4 OUR AVERAGE				
$3.96 \pm 0.15 \pm 0.60$	1192	JOUSSET 90 DM2		$J/\psi \rightarrow \text{hadrons}$
$4.33 \pm 0.12 \pm 0.45$		COFFMAN 88 MRK3		$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
2.7 ± 0.6	45	VANNUCCI 77 MRK1		$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

NODE=M070S16
NODE=M070S16

$\Gamma(\bar{K}^*(892)^0 K^+ \pi^- + c.c.) / \Gamma_{total}$ Γ_{74} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
5.73 ± 0.14 ± 0.82	¹ ANASHIN 22 KEDR		$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$
seen	² ABLIKIM 06C BES2		$J/\psi \rightarrow \bar{K}^*(892)^0 K^+ \pi^-$

NODE=M070S52
NODE=M070S52

¹ Obtained from $J/\psi \rightarrow K^*(892) K^- \pi^+ + c.c. \rightarrow K^+ K^- \pi^+ \pi^-$ taking the value 2/3 for the probability of the $K^*(892)^0 \rightarrow K^+ \pi^-$ decay.

NODE=M070S52;LINKAGE=A

² A $K_0^*(700)$ is observed by ABLIKIM 06C in the $K^+ \pi^-$ mass spectrum of the $\bar{K}^*(892)^0 K^+ \pi^-$ final state against the $\bar{K}^*(892)$. A corresponding branching fraction of the $J/\psi(1S)$ is not presented.

NODE=M070S52;LINKAGE=AB

$\Gamma(K^*(892)^0 K^- \pi^+ + c.c. \rightarrow K^+ K^- \pi^+ \pi^-) / \Gamma_{total}$ Γ_{78} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.81 ± 0.10 ± 0.54	1559	ANASHIN 22 KEDR		$J/\psi \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M070Q42
NODE=M070Q42

$\Gamma(K^*(892)^0 K_S^0 \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{total}$ Γ_{79} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
6.28^{+0.16+0.59}_{-0.17-0.52}	ABLIKIM 18AA BES3		$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P71
NODE=M070P71

$\Gamma(K^*(892)^\pm K^*(700)^\mp) / \Gamma_{total}$ Γ_{81} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09 ± 0.18^{+0.94}_{-0.54}	655	ABLIKIM 10E BES2		$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$

NODE=M070S74
NODE=M070S74

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
----	----	----------	----	--

NODE=M070R46
NODE=M070R46

 $\Gamma(K^*(892)^\pm K^*(892)^\mp)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.00 \pm 0.19^{+0.11}_{-0.32}$	323	ABLIKIM	10E	BES2 $J/\psi \rightarrow K^\pm K_S^0 \pi^\mp \pi^0$
---------------------------------	-----	---------	-----	---

NODE=M070S73
NODE=M070S73

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$3.8 \pm 0.8 \pm 1.2$	¹ BAI	99C	BES $e^+ e^-$
-----------------------	------------------	-----	---------------

NODE=M070S35
NODE=M070S35

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

NODE=M070S35;LINKAGE=M3

 $\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{86}/Γ_{180}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$2.3 \pm 1.1 \pm 0.7$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$
-----------------------	----	-------------------	-----	---

NODE=M070P28
NODE=M070P28

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P28;LINKAGE=A

 $\Gamma(K^*(1410) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{87}/Γ_{181}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$1.5 \pm 0.5 \pm 0.9$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
-----------------------	----	-------------------	-----	---

NODE=M070P32
NODE=M070P32

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P32;LINKAGE=A

 $\Gamma(K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K^\pm K^\mp \pi^0)/\Gamma(K^+ K^- \pi^0)$ Γ_{89}/Γ_{180}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$3.5 \pm 1.3 \pm 0.9$	2k	¹ LEES	17C	BABR $J/\psi \rightarrow K^+ K^- \pi^0$
-----------------------	----	-------------------	-----	---

NODE=M070P29
NODE=M070P29

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P29;LINKAGE=A

 $\Gamma(K_2^*(1430) \bar{K} + \text{c.c.} \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma(K_S^0 K^\pm \pi^\mp)$ Γ_{90}/Γ_{181}

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

$7.1 \pm 1.3 \pm 1.2$	4k	¹ LEES	17C	BABR $J/\psi \rightarrow K_S^0 K^\pm \pi^\mp$
-----------------------	----	-------------------	-----	---

NODE=M070P33
NODE=M070P33

¹ From a Dalitz plot analysis in an isobar model.

NODE=M070P33;LINKAGE=A

 $\Gamma(\bar{K}_2^*(1430) K + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<40 \times 10^{-4}$	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow K^0 \bar{K}_2^{*0}$
----------------------	----	----------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<66 \times 10^{-4}$	90	BRAUNSCH...	76	DASP $e^+ e^- \rightarrow K^\pm \bar{K}_2^{*\mp}$
----------------------	----	-------------	----	---

NODE=M070R45
NODE=M070R45

 $\Gamma(K_2^*(1430)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$2.69 \pm 0.04^{+0.25}_{-0.19}$	183k	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+ K^- \pi^0$
---------------------------------	------	---------	------	--

NODE=M070P76
NODE=M070P76

 $\Gamma(\bar{K}_2^*(1430)^0 K^*(892)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.7 ± 2.6	40	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
---------------	----	----------	----	--

NODE=M070R48
NODE=M070R48

 $\Gamma(K_2^*(1430)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<29 \times 10^{-4}$	90	VANNUCCI	77	MRK1 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$
----------------------	----	----------	----	--

NODE=M070R47
NODE=M070R47

 $\Gamma(K_2^*(1980)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.1 \pm 0.1^{+0.6}_{-0.1}$	183k	ABLIKIM	19AQ	BES $J/\psi \rightarrow K^+ K^- \pi^0$
-----------------------------	------	---------	------	--

NODE=M070P77
NODE=M070P77

$\Gamma(K_4^*(2045)^+ K^- + \text{c.c.} \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.2 \pm 0.7^{+2.8}_{-1.4}$	183k	ABLIKIM	19AQ BES	$J/\psi \rightarrow K^+ K^- \pi^0$

NODE=M070P78
NODE=M070P78

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.0 \times 10^{-3}$	90	¹ BAI	99C BES	$e^+ e^-$

NODE=M070S34
NODE=M070S34

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

NODE=M070S34;LINKAGE=M2

 $\Gamma(K_1(1270) K_S^0 \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
$8.54^{+1.07+2.35}_{-1.20-2.13}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P72
NODE=M070P72

 $\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 43 \times 10^{-4}$	90	BRAUNSCH... 76	DASP	$e^+ e^-$

NODE=M070R42
NODE=M070R42

 $\Gamma(\phi \pi^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

The two different fit values of ABLIKIM 15K below have the same statistical significance of 6.4 σ and cannot be distinguished at this moment.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.94 \pm 0.16 \pm 0.16$		0.8k	¹ ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$
$0.124 \pm 0.033 \pm 0.030$		35 ± 9	² ABLIKIM	15K BES3	$e^+ e^- \rightarrow J/\psi \rightarrow K^+ K^- \gamma \gamma$

NODE=M070R33
NODE=M070R33

NODE=M070R33

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.4	90	³ ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \gamma \gamma$
< 6.8	90	COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \pi^0$

¹ Corresponding to one of the two fit solutions with $\delta = (-95.9 \pm 1.5)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

NODE=M070R33;LINKAGE=A

² Corresponding to one of the two fit solutions with $\delta = (-152.1 \pm 7.7)^\circ$ for the phase angle between the resonant $J/\psi \rightarrow \phi \pi^0$ and non-phi $J/\psi \rightarrow K^+ K^- \pi^0$ contributions.

NODE=M070R33;LINKAGE=C

³ Superseded by ABLIKIM 15K.

NODE=M070R33;LINKAGE=B

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94 ± 0.15 OUR AVERAGE		Error includes scale factor of 1.7.		
$1.09 \pm 0.02 \pm 0.13$		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
$0.78 \pm 0.03 \pm 0.12$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.1 ± 0.9	23	FELDMAN	77 MRK1	$e^+ e^-$

NODE=M070R34
NODE=M070R34

 $\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$16.0 \pm 1.0 \pm 3.0$	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070R35
NODE=M070R35

 $\Gamma(\phi \eta)/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.74 ± 0.06 OUR AVERAGE		Error includes scale factor of 1.2.		
$0.71 \pm 0.10 \pm 0.05$	99 ± 14	¹ ZHU	23 BELL	$e^+ e^- \rightarrow \mathcal{T}(\text{nS}) \rightarrow \phi \eta \gamma$
$0.898 \pm 0.024 \pm 0.089$		ABLIKIM	05B BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \text{hadr}$
$0.64 \pm 0.04 \pm 0.11$	346	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
$0.661 \pm 0.045 \pm 0.078$		COFFMAN	88 MRK3	$e^+ e^- \rightarrow K^+ K^- \eta$

NODE=M070R37
NODE=M070R37


¹ From a fit to the combined $\phi \eta$ invariant mass spectrum with a Gaussian function for the J/ψ signals and a second-order polynomial function for the backgrounds.

NODE=M070R37;LINKAGE=A

$\Gamma(\phi\eta'(958))/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-4})	CL% EVTS	DOCUMENT ID	TECN	COMMENT
4.6 ± 0.5 OUR AVERAGE		Error includes scale factor of 2.2. See the ideogram below.		
5.10 ± 0.03 ± 0.32	31k	ABLIKIM	16M BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
5.46 ± 0.31 ± 0.56		ABLIKIM	05B BES2	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$
4.1 ± 0.3 ± 0.8	167	JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
3.08 ± 0.34 ± 0.36		COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+K^-\eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 13	90	VANNUCCI	77 MRK1	e^+e^-

NODE=M070R38
 NODE=M070R38

 $\Gamma(\phi\eta\eta')/\Gamma_{\text{total}}$ Γ_{111}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.32 ± 0.06 ± 0.16	2.2k	¹ ABLIKIM	19AN BES3	$e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$

NODE=M070P85
 NODE=M070P85

¹ Including contributions from intermediate resonances. Evidence for an intermediate resonance at $M \approx 2$ GeV and $\Gamma \approx 150$ MeV decaying to $\phi\eta'$ with $J^P = 1^+$ or $J^P = 1^-$, and $B(J/\psi \rightarrow \eta X) \times B(X \rightarrow \phi\eta') \approx 10^{-4}$.

NODE=M070P85;LINKAGE=A

 $\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$ Γ_{112}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
4.6 ± 0.4 ± 0.8		¹ FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
2.6 ± 0.6	50	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+K^-K^+K^-$

NODE=M070R41
 NODE=M070R41

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

NODE=M070R41;LINKAGE=A

 $\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
4.50 ± 0.80 ± 0.61	355	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$

NODE=M070S97
 NODE=M070S97

 $\Gamma(\phi\pi^0 f_0(980) \rightarrow \phi\pi^0\rho^0\pi^0)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.67 ± 0.50 ± 0.24	70	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+K^-3\pi$

NODE=M070S98
 NODE=M070S98

 $\Gamma(\phi f_0(980)\eta \rightarrow \eta\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.23 ± 0.75 ± 0.73	52	ABLIKIM	08F BES	$J/\psi \rightarrow \eta\phi f_0(980)$

NODE=M070R08
 NODE=M070R08

$\Gamma(\phi a_0(980)^0 \rightarrow \phi \eta \pi^0) / \Gamma_{\text{total}}$ Γ_{118} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
4.37 ± 1.35	1 ABLIKIM	18D BES3	$J/\psi \rightarrow \phi \eta \pi^0$
5.0 ± 2.7 ± 2.5	2 ABLIKIM	11D BES3	$J/\psi \rightarrow \phi \eta \pi^0$

NODE=M070S75
 NODE=M070S75

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assuming constructive interference between $a_0(980) - f_0(980)$ mixing and electromagnetic decay. Destructive interference gives a value of $(4.93 \pm 1.77) \times 10^{-6}$ for this branching fraction.

NODE=M070S75;LINKAGE=A

² Assuming $a_0(980) - f_0(980)$ mixing and isospin breaking via γ^* and $K^* K$ loops.

NODE=M070S75;LINKAGE=AB

 $\Gamma(\phi f_2(1270)) / \Gamma_{\text{total}}$ Γ_{119} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.45	90	FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
< 0.37	90	VANNUCCI	77 MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

NODE=M070R39
 NODE=M070R39

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\phi f_1(1285)) / \Gamma_{\text{total}}$ Γ_{120} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR AVERAGE				
3.4 ± 1.8 ± 1.5	1.1k	1 ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
3.2 ± 0.6 ± 0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
2.1 ± 0.5 ± 0.4	25	2 JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070S6
 NODE=M070S6

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2 ± 0.1 16 BECKER 87 MRK3 $J/\psi \rightarrow \phi K \bar{K} \pi$

OCCUR=2

¹ ABLIKIM 15H reports $[\Gamma(J/\psi(1S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.20 \pm 0.6 \pm 0.14) \times 10^{-4}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070S6;LINKAGE=A

² We attribute to the $f_1(1285)$ the signal observed in the $\pi^+ \pi^- \eta$ invariant mass distribution at 1297 MeV.

NODE=M070S6;LINKAGE=Q

 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi \pi^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{121} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
9.36 ± 2.31 ± 1.54	78	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S99
 NODE=M070S99

 $\Gamma(\phi f_1(1285) \rightarrow \phi \pi^0 f_0(980) \rightarrow \phi 3\pi^0) / \Gamma_{\text{total}}$ Γ_{122} / Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
2.08 ± 1.63 ± 1.47	9	ABLIKIM	15P BES3	$J/\psi \rightarrow K^+ K^- 3\pi$

NODE=M070S00
 NODE=M070S00

 $\Gamma(\phi \eta(1405) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{123} / Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.01 ± 0.58 ± 0.82		172	1 ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070S23
 NODE=M070S23

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 17 90 ² FALVARD 88 DM2 $J/\psi \rightarrow \text{hadrons}$

¹ With 3.6 σ significance.

NODE=M070S23;LINKAGE=B

² Includes unknown branching fraction $\eta(1405) \rightarrow \eta \pi \pi$.

NODE=M070S23;LINKAGE=A

 $\Gamma(\phi f'_2(1525)) / \Gamma_{\text{total}}$ Γ_{124} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE				Error includes scale factor of 2.7.
12.3 ± 0.6 ± 2.0		^{1,2} FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
4.8 ± 1.8	46	¹ GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

NODE=M070R40
 NODE=M070R40

¹ Re-evaluated using $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$.

NODE=M070R40;LINKAGE=B

² Including interference with $f_0(1710)$.

NODE=M070R40;LINKAGE=C

 $\Gamma(\phi X(1835) \rightarrow \phi \rho \bar{\rho}) / \Gamma_{\text{total}}$ Γ_{125} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.1 × 10⁻⁷	90	1 ABLIKIM	16K BES3	$J/\psi \rightarrow \rho \bar{\rho} K_S^0 K_L^0$ $\rho \bar{\rho} K^+ K^-$

NODE=M070P00
 NODE=M070P00

OCCUR=2

¹ Upper limit applies to any $\rho \bar{\rho}$ mass enhancement near threshold.

NODE=M070P00;LINKAGE=A

$\Gamma(\phi X(1835) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{126} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-4}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070B10
NODE=M070B10

 $\Gamma(\phi X(1870) \rightarrow \phi \eta \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{127} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.13 \times 10^{-5}$	90	ABLIKIM	15H BES3	$e^+ e^- \rightarrow J/\psi \rightarrow \phi \eta \pi^+ \pi^-$

NODE=M070B11
NODE=M070B11

 $\Gamma(\phi K \bar{K}) / \Gamma_{\text{total}}$ Γ_{128} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17.7 ± 1.6 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
16.6 ± 1.9 ± 1.2	163 ± 19	LEES	12F BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
21.4 ± 0.4 ± 2.2		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
48 $\begin{smallmatrix} +20 \\ -16 \end{smallmatrix} \pm 6$	9.0 $\begin{smallmatrix} +3.7 \\ -3.0 \end{smallmatrix}$	1,2 HUANG	03 BELL	$B^+ \rightarrow (\phi K^+ K^-) K^+$
14.6 ± 0.8 ± 2.1		3 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
18 ± 8	14	FELDMAN	77 MRK1	$e^+ e^-$


NODE=M070R36
NODE=M070R36

¹ We have multiplied $K^+ K^-$ measurement by 2 to obtain $K \bar{K}$.

² Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.

³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

NODE=M070R36;LINKAGE=AA
NODE=M070R36;LINKAGE=CC
NODE=M070R36;LINKAGE=A

 $\Gamma(\phi f_0(1710) \rightarrow \phi K \bar{K}) / \Gamma_{\text{total}}$ Γ_{129} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.6 ± 0.2 ± 0.6	1,2 FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S24
NODE=M070S24

¹ Including interference with $f_2'(1525)$.

² Includes unknown branching fraction $f_0(1710) \rightarrow K \bar{K}$.

NODE=M070S24;LINKAGE=D
NODE=M070S24;LINKAGE=E

 $\Gamma(\phi K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$ Γ_{132} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 0.8 OUR AVERAGE				
7.4 ± 0.6 ± 1.4	227 ± 19	ABLIKIM	08E BES2	$e^+ e^- \rightarrow J/\psi$
7.4 ± 0.9 ± 1.1		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
7 ± 0.6 ± 1.0	163 ± 15	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S3
NODE=M070S3

 $\Gamma(\phi K^*(892) \bar{K} + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{133} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
21.8 ± 2.3 OUR AVERAGE				
20.8 ± 2.7 ± 3.9	195 ± 25	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K_S^0 K^\pm \pi^\mp$
29.6 ± 3.7 ± 4.7	238 ± 30	ABLIKIM	08E BES2	$J/\psi \rightarrow \phi K^+ K^- \pi^0$
20.7 ± 2.4 ± 3.0		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87 MRK3	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M070S4
NODE=M070S4

OCCUR=2

$\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$					Γ_{134}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
30±5 OUR AVERAGE						
31±6	4600	AUGUSTIN	89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$	NODE=M070R49
29±7	87	BURMESTER	77D	PLUT	$e^+ e^-$	NODE=M070R49
$\Gamma(b_1(1235)^0 \pi^0)/\Gamma_{\text{total}}$					Γ_{135}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
23±3±5						
	229	AUGUSTIN	89	DM2	$e^+ e^-$	NODE=M070S28
						NODE=M070S28
$\Gamma(\Delta(1232)^+ \bar{p})/\Gamma_{\text{total}}$					Γ_{137}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.1 × 10⁻³						
	90	HENRARD	87	DM2	$e^+ e^-$	NODE=M070S14
						NODE=M070S14
$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$					Γ_{138}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.58±0.23±0.40						
	332	EATON	84	MRK2	$e^+ e^-$	NODE=M070R70
						NODE=M070R70
$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$					Γ_{139}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.10±0.09±0.28						
	233	EATON	84	MRK2	$e^+ e^-$	NODE=M070R66
						NODE=M070R66
$\Gamma(\bar{\Sigma}(1385)^0 p K^-)/\Gamma_{\text{total}}$					Γ_{140}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.51±0.26±0.18						
	89	EATON	84	MRK2	$e^+ e^-$	NODE=M070R74
						NODE=M070R74
$\Gamma(\Sigma(1385)^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{141}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.82 × 10⁻⁵						
	90	ABLIKIM	13F	BES3	$J/\psi \rightarrow p \bar{p} \pi^+ \pi^- \gamma \gamma$	NODE=M070S13
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	90	HENRARD	87	DM2	$e^+ e^-$	NODE=M070S13
$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{142}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.30±0.07 OUR AVERAGE						
[(0.31 ± 0.05) × 10 ⁻³ OUR 2023 AVERAGE]						
0.30±0.03±0.08	74 ± 8	HENRARD	87	DM2	$e^+ e^-$	NODE=M070R68
0.29±0.11±0.10	26	EATON	84	MRK2	$e^+ e^-$	NODE=M070R68
						NEW
$\Gamma(\Sigma(1385)^+ \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{143}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.33±0.08 OUR AVERAGE						
0.34±0.04±0.08	77	HENRARD	87	DM2	$e^+ e^-$	NODE=M070Q47
0.31±0.11±0.11	28	EATON	84	MRK2	$e^+ e^-$	NODE=M070Q47
$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{144}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.08 ±0.06 OUR AVERAGE						
[(1.16 ± 0.05) × 10 ⁻³ OUR 2023 AVERAGE]						
1.096±0.012±0.071	43k	ABLIKIM	16L	BES3	$e^+ e^-$	NODE=M070R67
1.23 ±0.07 ±0.30	0.8k	ABLIKIM	12P	BES2	$e^+ e^-$	NODE=M070R67
1.00 ±0.04 ±0.21	0.6k	HENRARD	87	DM2	$e^+ e^-$	
0.86 ±0.18 ±0.22	56	EATON	84	MRK2	$e^+ e^-$	NEW
$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{145}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.25 ±0.07 OUR AVERAGE						
1.258±0.014±0.078	53k	ABLIKIM	16L	BES3	$e^+ e^-$	NODE=M070Q48
1.50 ±0.08 ±0.38	1k	ABLIKIM	12P	BES2	$e^+ e^-$	NODE=M070Q48
1.19 ±0.04 ±0.25	0.7k	HENRARD	87	DM2	$e^+ e^-$	
1.03 ±0.24 ±0.25	68	EATON	84	MRK2	$e^+ e^-$	

$\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$					Γ_{146}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.071 ± 0.009 ± 0.082	103k	ABLIKIM	17E BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons		NODE=M070P17 NODE=M070P17 OCCUR=2

$\Gamma(\Lambda(1520)\bar{\Lambda} + \text{c.c.} \rightarrow \gamma\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$					Γ_{147}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<4.1 × 10⁻⁶	90	ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Lambda}\gamma$		NODE=M070S77 NODE=M070S77

$\Gamma(\bar{\Lambda}(1520)\Lambda + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{148}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.80 × 10⁻³	90	LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$		NODE=M070P60 NODE=M070P60

$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$					Γ_{149}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.17 ± 0.04 OUR AVERAGE						
1.165 ± 0.004 ± 0.043	135k	ABLIKIM	17E BES3	$e^+ e^- \rightarrow J/\psi \rightarrow$ hadrons		NODE=M070S64 NODE=M070S64
1.20 ± 0.12 ± 0.21	206	ABLIKIM	080 BES2	$e^+ e^- \rightarrow J/\psi$		

$\Gamma(\Xi(1530)^- \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{150}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.318 ± 0.008 OUR AVERAGE						
0.317 ± 0.002 ± 0.008	70k	ABLIKIM	20 BES3	$e^+ e^- \rightarrow J/\psi$		NODE=M070S9 NODE=M070S9
0.59 ± 0.09 ± 0.12	75	HENRARD	87 DM2	$e^+ e^-$		

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$					Γ_{151}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.32 ± 0.12 ± 0.07	24 ± 9	HENRARD	87 DM2	$e^+ e^-$		NODE=M070S10 NODE=M070S10

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 \rho K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{152}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.1 × 10⁻⁵	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S47 NODE=M070S47

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{153}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.1 × 10⁻⁵	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S48 NODE=M070S48

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{154}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.6 × 10⁻⁵	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S49 NODE=M070S49

$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$					Γ_{155}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<5.6 × 10⁻⁵	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S50 NODE=M070S50

$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 \rho K^- \bar{n})/\Gamma_{\text{total}}$					Γ_{156}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.1 × 10⁻⁵	90	BAI	04G BES2	$e^+ e^-$		NODE=M070S51 NODE=M070S51


STABLE HADRONS

$\Gamma(2(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$					Γ_{157}/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		
4.2 ± 0.4 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.		NODE=M070307
5.44 ± 0.07 ± 0.33	23K	ANASHIN	22 KEDR	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$		NODE=M070R9 NODE=M070R9
3.5 ± 0.8 ± 0.1	14k	¹ LEES	21 BABR	$10.6 e^+ e^- \rightarrow$ $2(\pi^+ \pi^-)3\pi^0 \gamma$		
4.73 ± 0.44	228k	² ABLIKIM	19Q BES3	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$		
3.25 ± 0.49	46055	AUGUSTIN	89 DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-)\pi^0$		
3.17 ± 0.42	147	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow$ hadrons		
3.64 ± 0.52	1500	BURMESTER	77D PLUT	$e^+ e^-$		
4 ± 1	675	JEAN-MARIE	76 MRK1	$e^+ e^-$		

- ¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}})] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0)] = (14.8 \pm 2.6 \pm 2.2) \times 10^{-3}$ keV which we divide by our best values $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) = (18.2 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.
- ² From an energy scan of $e^+e^- \rightarrow J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$, assuming PDG 16 values for $\Gamma(e^+e^-)$, $\Gamma(\mu^+\mu^-)$, and $\Gamma(\text{total})$, and for a phase difference between strong and electromagnetic amplitudes of $(84.9 \pm 3.6)^\circ$. An alternative solution is $(4.85 \pm 0.45)\%$ with a phase of $(-84.7 \pm 3.1)^\circ$.

NODE=M070R9;LINKAGE=B

NODE=M070R9;LINKAGE=A



$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$				Γ_{158}/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.029 ± 0.006	OUR AVERAGE				
0.028 ± 0.009	11	FRANKLIN	83	MRK2	$e^+e^- \rightarrow \text{hadrons}$
0.029 ± 0.007	181	JEAN-MARIE	76	MRK1	e^+e^-

NODE=M070R11
NODE=M070R11

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$				Γ_{163}/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
20.0 ± 0.7	OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.			
[(21.0 ± 0.8) × 10 ⁻³ OUR 2023 AVERAGE Scale factor = 1.6]					
18.78 ± 0.13 ± 0.51	19.8k	¹ ANASHIN	23	KEDR	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
21.37 ± 0.04 ^{+0.64} _{-0.62}	1.8M	² ABLIKIM	12H	BES3	$e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0$
23.0 ± 2.0 ± 0.4	256	³ AUBERT	07AU	BABR	10.6 $e^+e^- \rightarrow J/\psi \pi^+\pi^-\gamma$
21.84 ± 0.05 ± 2.01	220k	^{4,5} BAI	04H	BES	e^+e^-
20.91 ± 0.21 ± 1.16		^{5,6} BAI	04H	BES	e^+e^-
15 ± 2	168	FRANKLIN	83	MRK2	e^+e^-

NODE=M070R7
NODE=M070R7
NEW

OCCUR=2

- ¹ By a simultaneous fit of the $\pi\pi$ invariant mass distribution over the decay modes $J/\psi \rightarrow \rho^0\pi^0$, $J/\psi \rightarrow \rho^+\pi^-$, $J/\psi \rightarrow \rho^-\pi^+$. In the fit only the intermediate states $\rho(770)\pi$ and $\rho(1450)\pi$ are considered.
- ² The quoted systematic error includes a contribution of 1.23% (added in quadrature) from the uncertainty on the number of J/ψ events.
- ³ AUBERT 07AU reports $[\Gamma(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] = (18.6 \pm 1.2 \pm 1.1) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}} = 0.808 \pm 0.014$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ From $J/\psi \rightarrow \pi^+\pi^-\pi^0$ events directly.
- ⁵ Mostly $\rho\pi$, see also $\rho\pi$ subsection.
- ⁶ Obtained comparing the rates for $\pi^+\pi^-\pi^0$ and $\mu^+\mu^-$, using J/ψ events produced via $\psi(2S) \rightarrow \pi^+\pi^-J/\psi$ and with $B(J/\psi \rightarrow \mu^+\mu^-) = 5.88 \pm 0.10\%$.


NODE=M070R7;LINKAGE=B

NODE=M070R7;LINKAGE=AB

NODE=M070R7;LINKAGE=AU

NODE=M070R;LINKAGE=BA
NODE=M070R;LINKAGE=BU
NODE=M070R;LINKAGE=BI

WEIGHTED AVERAGE
20.0±0.7 (Error scaled by 2.0)



$\Gamma(\pi^+ \pi^- \pi^0 K^+ K^-) / \Gamma_{total}$ **Γ_{165} / Γ**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.52±0.27 OUR AVERAGE		Error includes scale factor of 1.4.		
1.74±0.08±0.24	2616	ANASHIN	22	KEDR $J/\psi \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.2 ±0.3	309	VANNUCCI	77	MRK1 $e^+ e^-$

NODE=M070R18
NODE=M070R18

$\Gamma(\pi^+ \pi^-) / \Gamma_{total}$ **Γ_{166} / Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.47±0.14 OUR AVERAGE				
1.47±0.13±0.13	140	¹ METREVELI	12	$\psi(2S) \rightarrow 2(\pi^+ \pi^-)$
1.58±0.20±0.15	84	BALTRUSAIT..85D	MRK3	$e^+ e^-$
1.0 ±0.5	5	BRANDELIK	78B	DASP $e^+ e^-$
1.6 ±1.6	1	VANNUCCI	77	MRK1 $e^+ e^-$

NODE=M070R6
NODE=M070R6

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070R6;LINKAGE=ME

$\Gamma(2(\pi^+ \pi^-)) / \Gamma_{total}$ **Γ_{167} / Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.20±0.25 OUR AVERAGE		Error includes scale factor of 1.2.		
2.88±0.14±0.24	2654	ANASHIN	22	KEDR $J/\psi \rightarrow 2(\pi^+ \pi^-)$
3.53±0.12±0.29	1107	¹ ABLIKIM	05H	BES2 $e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \pi^+ \pi^-, J/\psi \rightarrow 2(\pi^+ \pi^-)$
4.0 ±1.0	76	JEAN-MARIE	76	MRK1 $e^+ e^-$

NODE=M070R8
NODE=M070R8

¹ Computed using $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

NODE=M070R8;LINKAGE=AB

$\Gamma(3(\pi^+ \pi^-)) / \Gamma_{total}$ **Γ_{168} / Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
40±20	32	JEAN-MARIE	76	MRK1 $e^+ e^-$

NODE=M070R10
NODE=M070R10

$\Gamma(4(\pi^+ \pi^-) \pi^0) / \Gamma_{total}$ **Γ_{170} / Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
90±30	13	JEAN-MARIE	76	MRK1 $e^+ e^-$

NODE=M070R12
NODE=M070R12

$\Gamma(2(\pi^+ \pi^-) \eta) / \Gamma_{total}$ **Γ_{171} / Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29±0.28 OUR AVERAGE				
3.1 ±1.5 ±0.1	14k	¹ LEES	21	BABR $10.6 e^+ e^- \rightarrow 2(\pi^+ \pi^-) 3\pi^0 \gamma$
2.26±0.08±0.27	4.8k	ABLIKIM	05C	BES2 $e^+ e^- \rightarrow 2(\pi^+ \pi^-) \eta$

NODE=M070S42
NODE=M070S42

¹ LEES 21 reports $[\Gamma(J/\psi(1S) \rightarrow 2(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}] \times [\Gamma(J/\psi(1S) \rightarrow e^+e^-)] \times [B(\eta \rightarrow 3\pi^0)] = (5.6 \pm 2.6 \pm 0.8) \times 10^{-3}$ keV which we divide by our best values $\Gamma(J/\psi(1S) \rightarrow e^+e^-) = 5.53 \pm 0.10$ keV, $B(\eta \rightarrow 3\pi^0) = (32.57 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070S42;LINKAGE=A

$\Gamma(3(\pi^+\pi^-)\eta)/\Gamma_{\text{total}}$				Γ_{172}/Γ
VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
7.24±0.96±1.11	616	ABLIKIM	05C BES2	$e^+e^- \rightarrow 3(\pi^+\pi^-)\eta$

NODE=M070S43
NODE=M070S43

$\Gamma(K^+K^-)/\Gamma_{\text{total}}$				Γ_{176}/Γ
VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
2.86±0.09±0.19	1k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$
2.39±0.24±0.22	107	² BALTRUSAIT..85D	MRK3	e^+e^-
2.2 ±0.9	6	² BRANDELIK	79C DASP	e^+e^-

NODE=M070R13
NODE=M070R13

• • • We do not use the following data for averages, fits, limits, etc. • • •
¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
² Interference with non-resonant K^+K^- production not taken into account.


NODE=M070R13;LINKAGE=ME
NODE=M070R13;LINKAGE=BA

$\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$				Γ_{177}/Γ
VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
1.95±0.11 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.		
1.93±0.01±0.05	110k	ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$
2.62±0.15±0.14	0.3k	¹ METREVELI	12	$\psi(2S) \rightarrow \pi^+\pi^-K_S^0 K_L^0$
1.82±0.04±0.13	2.1k	² BAI	04A BES2	$J/\psi \rightarrow K_S^0 K_L^0 \rightarrow \pi^+\pi^-X$
1.18±0.12±0.18		JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$
1.01±0.16±0.09	74	BALTRUSAIT..85D	MRK3	e^+e^-

NODE=M070R75
NODE=M070R75

• • • We do not use the following data for averages, fits, limits, etc. • • •
¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6868 \pm 0.0027$.

NODE=M070R75;LINKAGE=ME
NODE=M070R;LINKAGE=HZ



$\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$				Γ_{178}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10⁻⁸	95	¹ ABLIKIM	17AH BES3	$J/\psi \rightarrow K_S^0 K_S^0 \rightarrow \pi^+\pi^-\pi^+\pi^-$
<1 × 10 ⁻⁶	95	¹ BAI	04D BES	e^+e^-
<5.2 × 10 ⁻⁶	90	¹ BALTRUSAIT..85C	MRK3	e^+e^-

NODE=M070R14
NODE=M070R14

¹ Forbidden by CP.

NODE=M070R14;LINKAGE=C

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$ Γ_{179}/Γ VALUE (units 10^{-4}) EVTS**61 ± 10 OUR AVERAGE**

55.2 ± 12.0

25

DOCUMENT ID TECN COMMENT

FRANKLIN 83 MRK2 $e^+e^- \rightarrow K^+K^-\pi^0$

78.0 ± 21.0

126

VANNUCCI 77 MRK1 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$ NODE=M070R15
NODE=M070R15 $\Gamma(K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{180}/Γ VALUE (units 10^{-3}) EVTS**2.88 ± 0.01 ± 0.12**

183k

DOCUMENT ID TECN COMMENT

ABLIKIM 19AQ BES $J/\psi \rightarrow K^+K^-\pi^0$ NODE=M070P80
NODE=M070P80 $\Gamma(K^+K^-\pi^0)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{180}/\Gamma_{163}$

VALUE (%) EVTS

12.0 ± 0.3 ± 0.9

23k

DOCUMENT ID TECN COMMENT

LEES 17C BABR $J/\psi \rightarrow h^+h^-\pi^0$ NODE=M070P34
NODE=M070P34 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma(\pi^+\pi^-\pi^0)$ $\Gamma_{181}/\Gamma_{163}$

VALUE (%) EVTS

26.5 ± 0.5 ± 2.1

24k

DOCUMENT ID TECN COMMENT

LEES 17C BABR $J/\psi \rightarrow h^0 h^+ h^-$ NODE=M070P35
NODE=M070P35 $\Gamma(K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{185}/Γ VALUE (units 10^{-3}) EVTS**7.04 ± 0.26 ± 0.92**

2671

DOCUMENT ID TECN COMMENT

ANASHIN 22 KEDR $J/\psi \rightarrow K^+K^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.2 ± 2.3

205

VANNUCCI 77 MRK1 e^+e^- NODE=M070R16
NODE=M070R16 $\Gamma(K^+K^-2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{197}/Γ VALUE (units 10^{-4}) EVTS**31 ± 13**

30

DOCUMENT ID TECN COMMENT

VANNUCCI 77 MRK1 e^+e^- NODE=M070R17
NODE=M070R17 $\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$ Γ_{199}/Γ VALUE (units 10^{-3}) EVTS $1.4^{+0.5}_{-0.4} \pm 0.2$ $11.0^{+4.3}_{-3.5}$ ¹ HUANG 03 BELL $B^+ \rightarrow 2(K^+K^-) K^+$

0.7 ± 0.3

VANNUCCI 77 MRK1 e^+e^- ¹ Using $B(B^+ \rightarrow J/\psi K^+) = (1.01 \pm 0.05) \times 10^{-3}$.NODE=M070R19
NODE=M070R19

NODE=M070R19;LINKAGE=CC

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{207}/Γ VALUE (units 10^{-3}) EVTS**2.120 ± 0.029 OUR AVERAGE**

2.112 ± 0.004 ± 0.031

314k

DOCUMENT ID TECN COMMENT

ABLIKIM 12C BES3 e^+e^-

2.17 ± 0.16 ± 0.04

317

¹ WU 06 BELL $B^+ \rightarrow p\bar{p}K^+$

2.26 ± 0.01 ± 0.14

63316

BAI 04E BES2 $e^+e^- \rightarrow J/\psi$

1.97 ± 0.22

99

BALDINI 98 FENI e^+e^-

1.91 ± 0.04 ± 0.30

PALLIN 87 DM2 e^+e^-

2.16 ± 0.07 ± 0.15

1420

EATON 84 MRK2 e^+e^-

2.5 ± 0.4

133

BRANDELIK 79C DASP e^+e^-

2.0 ± 0.5

BESCH 78 BONA e^+e^-

2.2 ± 0.2

331

² PERUZZI 78 MRK1 e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.3

48

ANTONELLI 93 SPEC e^+e^- ¹ WU 06 reports $[\Gamma(J/\psi(1S) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)] = (2.21 \pm 0.13 \pm 0.10) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.² Assuming angular distribution $(1 + \cos^2\theta)$.NODE=M070R50
NODE=M070R50

NODE=M070R50;LINKAGE=WU

NODE=M070R50;LINKAGE=A

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{208}/Γ VALUE (units 10^{-3}) EVTS**1.19 ± 0.08 OUR AVERAGE** Error includes scale factor of 1.1.

1.33 ± 0.02 ± 0.11

11k

ABLIKIM 09B BES2 e^+e^-

1.13 ± 0.09 ± 0.09

685

EATON 84 MRK2 e^+e^-

1.4 ± 0.4

BRANDELIK 79C DASP e^+e^-

1.00 ± 0.15

109

PERUZZI 78 MRK1 e^+e^- NODE=M070R52
NODE=M070R52


$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{total}$

Γ_{209}/Γ

NODE=M070R54
NODE=M070R54

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.5 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
6.46 ± 0.17 ± 0.43	1435	EATON	84 MRK2	e^+e^-
3.8 ± 1.6	48	BESCH	81 BONA	e^+e^-
5.5 ± 0.6	533	PERUZZI	78 MRK1	e^+e^-

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{total}$

Γ_{210}/Γ

NODE=M070R55
NODE=M070R55
NODE=M070R55

Including $p\bar{p}\pi^+\pi^-\gamma$ and excluding ω, η, η'

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.9 OUR AVERAGE		Error includes scale factor of 1.9.		
3.36 ± 0.65 ± 0.28	364	EATON	84 MRK2	e^+e^-
1.6 ± 0.6	39	PERUZZI	78 MRK1	e^+e^-

$\Gamma(p\bar{p}\eta)/\Gamma_{total}$

Γ_{211}/Γ

NODE=M070R56
NODE=M070R56

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00 ± 0.12 OUR AVERAGE				
1.91 ± 0.02 ± 0.17	13k	¹ ABLIKIM	09 BES2	e^+e^-
2.03 ± 0.13 ± 0.15	826	EATON	84 MRK2	e^+e^-
2.5 ± 1.2		BRANDELIK	79C DASP	e^+e^-
2.3 ± 0.4	197	PERUZZI	78 MRK1	e^+e^-

¹ From the combination of $p\bar{p}\eta \rightarrow p\bar{p}\gamma\gamma$ and $p\bar{p}\eta \rightarrow p\bar{p}\pi^+\pi^-\pi^0$ channels.

NODE=M070R56;LINKAGE=AB

$\Gamma(p\bar{p}\rho)/\Gamma_{total}$

Γ_{212}/Γ

NODE=M070R57
NODE=M070R57


VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.31 × 10⁻³	90	EATON	84 MRK2	$e^+e^- \rightarrow \text{hadrons}\gamma$

$\Gamma(p\bar{p}\omega)/\Gamma_{total}$

Γ_{213}/Γ

NODE=M070R58
NODE=M070R58

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.10 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
0.90 ± 0.02 ± 0.09	2670	ABLIKIM	13P BES3	e^+e^-
0.98 ± 0.03 ± 0.14	2449	ABLIKIM	08 BES2	e^+e^-
1.10 ± 0.17 ± 0.18	486	EATON	84 MRK2	e^+e^-
1.6 ± 0.3	77	PERUZZI	78 MRK1	e^+e^-



$\Gamma(\rho\bar{\rho}\eta(958))/\Gamma_{\text{total}}$

Γ_{214}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.129 ± 0.014 OUR AVERAGE				Error includes scale factor of 2.0.
$0.126 \pm 0.002 \pm 0.007$	16k	¹ ABLIKIM	19N BES3	e^+e^-
$0.200 \pm 0.023 \pm 0.028$	265 ± 31	² ABLIKIM	09 BES2	e^+e^-
$0.68 \pm 0.23 \pm 0.17$	19	EATON	84 MRK2	e^+e^-
1.8 ± 0.6	19	PERUZZI	78 MRK1	e^+e^-

NODE=M070R59
 NODE=M070R59

¹ From the combination of $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\pi^+\pi^-\eta$ and $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\pi^+\pi^-\gamma$ channels.

² From the combination of $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\pi^+\pi^-\eta$ and $\rho\bar{\rho}\eta' \rightarrow \rho\bar{\rho}\gamma\rho^0$ channels.

NODE=M070R59;LINKAGE=A
 NODE=M070R59;LINKAGE=AB

$\Gamma(\rho\bar{\rho}a_0(980) \rightarrow \rho\bar{\rho}\pi^0\eta)/\Gamma_{\text{total}}$

Γ_{215}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$6.8 \pm 1.2 \pm 1.3$	ABLIKIM	14N BES3	$e^+e^- \rightarrow J/\psi$

NODE=M070S94
 NODE=M070S94

$\Gamma(\rho\bar{\rho}\phi)/\Gamma_{\text{total}}$

Γ_{216}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.519 ± 0.033 OUR AVERAGE				
$0.523 \pm 0.006 \pm 0.033$	14k	ABLIKIM	16K BES3	$J/\psi \rightarrow \rho\bar{\rho}K_S^0 K_L^0$, $\rho\bar{\rho}K^+ K^-$
$0.45 \pm 0.13 \pm 0.07$		FALVARD	88 DM2	$J/\psi \rightarrow \text{hadrons}$

NODE=M070S22
 NODE=M070S22

$\Gamma(\rho\bar{\rho}\pi^-)/\Gamma_{\text{total}}$

Γ_{217}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.12 ± 0.09 OUR AVERAGE				
$2.36 \pm 0.02 \pm 0.21$	59k	ABLIKIM	06K BES2	$J/\psi \rightarrow \rho\pi^-\bar{n}$
$2.47 \pm 0.02 \pm 0.24$	55k	ABLIKIM	06K BES2	$J/\psi \rightarrow \bar{\rho}\pi^+n$
$2.02 \pm 0.07 \pm 0.16$	1288	EATON	84 MRK2	$e^+e^- \rightarrow \rho\pi^-$
$1.93 \pm 0.07 \pm 0.16$	1191	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\rho}\pi^+$
1.7 ± 0.7	32	BESCH	81 BONA	$e^+e^- \rightarrow \rho\pi^-$
1.6 ± 1.2	5	BESCH	81 BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$
2.16 ± 0.29	194	PERUZZI	78 MRK1	$e^+e^- \rightarrow \rho\pi^-$
2.04 ± 0.27	204	PERUZZI	78 MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$

NODE=M070R53
 NODE=M070R53

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

$\Gamma(n\bar{n})/\Gamma_{\text{total}}$

Γ_{218}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.09 ± 0.16 OUR AVERAGE				
$2.07 \pm 0.01 \pm 0.17$	36k	ABLIKIM	12C BES3	e^+e^-
2.31 ± 0.49	79	BALDINI	98 FENI	e^+e^-
1.8 ± 0.9		BESCH	78 BONA	e^+e^-
1.90 ± 0.55	40	ANTONELLI	93 SPEC	e^+e^-

NODE=M070R64
 NODE=M070R64

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{total}$

Γ_{219}/Γ

NODE=M070R65
NODE=M070R65

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8±3.6	5	BESCH	81	BONA e^+e^-

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{total}$

Γ_{223}/Γ

NODE=M070R60
NODE=M070R60

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.88 ±0.08 OUR AVERAGE				Error includes scale factor of 2.6. See the ideogram below. [(1.89 ± 0.09) × 10 ⁻³ OUR 2023 AVERAGE Scale factor = 2.8]


NEW

1.64 ±0.12 ±0.09		ABLIKIM	23S	BES3	$e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$
1.943±0.003±0.033	441k	ABLIKIM	17L	BES3	e^+e^-
2.03 ±0.03 ±0.15	8887	ABLIKIM	06	BES2	$J/\psi \rightarrow \Lambda\bar{\Lambda}$
1.96 ^{+0.47} _{-0.44} ±0.04	46	¹ WU	06	BELL	$B^+ \rightarrow \Lambda\bar{\Lambda}K^+$
1.08 ±0.06 ±0.24	631	BAI	98G	BES	e^+e^-
1.38 ±0.05 ±0.20	1847	PALLIN	87	DM2	e^+e^-
1.58 ±0.08 ±0.19	365	EATON	84	MRK2	e^+e^-
2.6 ±1.6	5	BESCH	81	BONA	e^+e^-
1.1 ±0.2	196	PERUZZI	78	MRK1	e^+e^-

NODE=M070R60;LINKAGE=WU

¹WU 06 reports [$\Gamma(J/\psi(1S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{total}$] × [B($B^+ \rightarrow J/\psi(1S)K^+$)] = (2.00 ^{+0.34}_{-0.29} ± 0.34) × 10⁻⁶ which we divide by our best value B($B^+ \rightarrow J/\psi(1S)K^+$) = (1.020 ± 0.019) × 10⁻³. Our first error is their experiment's error and our second error is the systematic error from using our best value.

WEIGHTED AVERAGE
1.88±0.08 (Error scaled by 2.6)



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{total}$

Γ_{224}/Γ

NODE=M070S11
NODE=M070S11

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.78±0.27±0.30		323	¹ ABLIKIM	13F	BES3 $J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

••• We do not use the following data for averages, fits, limits, etc. •••

< 6.4	90	² ABLIKIM	07H	BES2	$e^+e^- \rightarrow \psi(2S)$
23 ±7 ±8	11	BAI	98G	BES	e^+e^-
22 ±5 ±5	19	HENRRARD	87	DM2	e^+e^-

¹Using B($\Lambda \rightarrow \pi^- p$) = 63.9% and B($\pi^0 \rightarrow \gamma\gamma$) = 98.8%.

²Using B($\Lambda \rightarrow \pi^- p$) = 63.9%.

NODE=M070S11;LINKAGE=AL
NODE=M070S11;LINKAGE=AB

$\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{total}$

Γ_{225}/Γ

NODE=M070S78
NODE=M070S78

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.30±0.13±0.99	2.4k	ABLIKIM	12P	BES2 J/ψ

$\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{226}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.2±1.7 OUR AVERAGE				
15.7±0.80±1.54	454	¹ ABLIKIM	13F BES3	$J/\psi \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
26.2±6.0 ±4.4	44	² ABLIKIM	07H BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M070R07
 NODE=M070R07

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

NODE=M070R07;LINKAGE=AL
 NODE=M070R07;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{227}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.26 ±0.05 OUR AVERAGE				
Error includes scale factor of 1.2. $[(0.83 \pm 0.07) \times 10^{-3}]$ OUR 2023 AVERAGE Scale factor = 1.2]				
1.244±0.002±0.045	2.6M	ABLIKIM	23BU BES3	e^+e^-
1.52 ±0.08 ±0.16	589	¹ ABLIKIM	07H BES2	e^+e^-
1.11 ±0.06 ±0.20	342 ± 18	HENRARD	87 DM2	e^+e^-
1.38 ±0.21 ±0.35	118	EATON	84 MRK2	e^+e^-

NODE=M070R71
 NODE=M070R71

NEW

OCCUR=3
 OCCUR=2
 OCCUR=2

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\Sigma^+ \rightarrow \pi^0 p) = 51.6\%$.

NODE=M070R71;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{228}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.21 ±0.07 OUR AVERAGE				
Error includes scale factor of 1.8.				
1.221±0.002±0.038	2.7M	ABLIKIM	23BU BES3	e^+e^-
0.90 ±0.06 ±0.16	225	HENRARD	87 DM2	e^+e^-
1.53 ±0.17 ±0.38	135	EATON	84 MRK2	e^+e^-

NODE=M070Q46
 NODE=M070Q46

 $\Gamma(pK^-\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{229}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.11 OUR AVERAGE				
0.84 ^{+0.17} _{-0.15} ±0.02	45	¹ LU	19 BELL	$B^+ \rightarrow \bar{p}\Lambda K^+ K^+$
0.89±0.07±0.14	307	EATON	84 MRK2	e^+e^-

NODE=M070R72
 NODE=M070R72

¹ LU 19 reports $(8.32^{+1.63}_{-1.45} \pm 0.49) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow pK^-\bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S)K^+)]$ assuming $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.026 \pm 0.031) \times 10^{-3}$, which we rescale to our best value $B(B^+ \rightarrow J/\psi(1S)K^+) = (1.020 \pm 0.019) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M070R72;LINKAGE=A

 $\Gamma(pK^-\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{230}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.06±0.05	90	EATON	84 MRK2	e^+e^-

NODE=M070R73
 NODE=M070R73

 $\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{231}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.46±0.20±1.07	1058	¹ ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070S56
 NODE=M070S56

¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.

NODE=M070S56;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Sigma} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{232}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.83±0.23 OUR AVERAGE					
2.74±0.24±0.22		234 ± 21	¹ ABLIKIM	12B BES3	$J/\psi \rightarrow \Lambda\bar{\Sigma}^0$
2.92±0.22±0.24		308 ± 24	² ABLIKIM	12B BES3	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$

NODE=M070R61
 NODE=M070R61

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18			² HENRARD	87 DM2	$J/\psi \rightarrow \bar{\Lambda}\Sigma^0$
<15	90		PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$

¹ ABLIKIM 12B quotes $B(J/\psi \rightarrow \Lambda\bar{\Sigma}^0)$ which we multiply by 2.

² ABLIKIM 12B and HENRARD 87 quote results for $B(J/\psi \rightarrow \bar{\Lambda}\Sigma^0)$ which we multiply by 2.

NODE=M070R61;LINKAGE=AB
 NODE=M070R61;LINKAGE=AC

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{233}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.07 ±0.04 OUR AVERAGE				
1.061±0.004±0.036	87k	ABLIKIM	21AT BES3	$J/\psi \rightarrow p\pi^0\bar{p}\pi^0$
1.50 ±0.10 ±0.22	399	ABLIKIM	08O BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070S09
 NODE=M070S09

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{total}$

Γ_{234}/Γ

NODE=M070R63
NODE=M070R63

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.172 ± 0.032 OUR AVERAGE		Error includes scale factor of 1.4.		
1.164 ± 0.004 ± 0.023	111k	ABLIKIM	17L BES3	$J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0$
1.33 ± 0.04 ± 0.11	1.7k	ABLIKIM	06 BES2	$J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0$
1.06 ± 0.04 ± 0.23	884	PALLIN	87 DM2	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
1.58 ± 0.16 ± 0.25	90	EATON	84 MRK2	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
1.3 ± 0.4	52	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.4 ± 2.6	3	BESCH	81 BONA	$e^+ e^- \rightarrow \Sigma^+ \bar{\Sigma}^-$

$\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{total}$

Γ_{235}/Γ

NODE=M070Q45
NODE=M070Q45


VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.34 ± 0.21 ± 0.37	1821	ABLIKIM	22AY BES3	$J/\psi \rightarrow \Sigma^+ \bar{\Sigma}^- \eta$

$\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{total}$

Γ_{236}/Γ

NODE=M070R62
NODE=M070R62

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.08 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
1.040 ± 0.006 ± 0.074	43k	ABLIKIM	16L BES3	$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
0.90 ± 0.03 ± 0.18	961	ABLIKIM	12P BES2	$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$
0.70 ± 0.06 ± 0.12	132	HENRARD	87 DM2	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.14 ± 0.08 ± 0.20	194	EATON	84 MRK2	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$
1.4 ± 0.5	51	PERUZZI	78 MRK1	$e^+ e^- \rightarrow \Xi^- \bar{\Xi}^+$



RADIATIVE DECAYS

NODE=M070310

$\Gamma(\gamma \eta_c(1S))/\Gamma_{total}$

Γ_{237}/Γ

NODE=M070R85
NODE=M070R85

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.14 OUR FIT	Error includes scale factor of 1.3.		
1.7 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.5.		
2.00 ± 0.31 ± 0.02	¹ MITCHELL	09 CLEO	$e^+ e^- \rightarrow \gamma X$
1.27 ± 0.36	GAISER	86 CBAL	$J/\psi \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
3.40 ± 0.33	² ANASHIN	14 KEDR	$J/\psi \rightarrow \gamma \eta_c$

¹ MITCHELL 09 reports $(1.98 \pm 0.09 \pm 0.30) \times 10^{-2}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma \eta_c(1S))/\Gamma_{total}] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (35.04 \pm 0.07 \pm 0.77) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Statistical uncertainty only.

NODE=M070R85;LINKAGE=M1

NODE=M070R85;LINKAGE=A

$\Gamma(3\gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
11.6±2.2 OUR AVERAGE					
11.3±1.8±2.0		113 ± 18	ABLIKIM	13I	BES3 $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$
12 ±3 ±2		24.2 ^{+7.2} _{-6.0}	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 Γ_{240}/Γ

NODE=M070R81
NODE=M070R81

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55 90 PARTRIDGE 80 CBAL e^+e^-

 $\Gamma(4\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9 × 10 ⁻⁶	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 Γ_{241}/Γ

NODE=M070S06
NODE=M070S06

 $\Gamma(5\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<15 × 10 ⁻⁶	90	ADAMS	08	CLEO $\psi(2S) \rightarrow \pi^+\pi^- J/\psi$

 Γ_{242}/Γ

NODE=M070S07
NODE=M070S07

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.39±0.08 OUR AVERAGE				

 Γ_{243}/Γ

NODE=M070R82
NODE=M070R82
NEW

[(3.56 ± 0.17) × 10⁻⁵ OUR 2023 AVERAGE]

3.34±0.02±0.09	176k	ABLIKIM	23BD	BES3 $J/\psi \rightarrow \pi^0\gamma$
3.59±0.20±0.04	1.6k	¹ ABLIKIM	180	BES3 $\psi(2S) \rightarrow \pi^+\pi^- \gamma\gamma\gamma$
3.63±0.36±0.13		PEDLAR	09	CLE3 $J/\psi \rightarrow \pi^0\gamma$
3.13 ^{+0.65} _{-0.47}	586	ABLIKIM	06E	BES2 $J/\psi \rightarrow \pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.6 ±1.1 ±0.7 BLOOM 83 CBAL e^+e^-
7.3 ±4.7 10 BRANDELIK 79C DASP e^+e^-

¹ ABLIKIM 180 reports [$\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}$] × [B($\pi^0 \rightarrow 2\gamma$)] = (3.57 ± 0.12 ± 0.16) × 10⁻⁵ from a measurement of [$\Gamma(J/\psi(1S) \rightarrow \gamma\pi^0)/\Gamma_{\text{total}}$] × [B($\pi^0 \rightarrow 2\gamma$)] × [B($\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-$)] assuming B($\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-$) = (34.49 ± 0.30) × 10⁻², which we rescale to our best values B($\pi^0 \rightarrow 2\gamma$) = (98.823 ± 0.034) × 10⁻², B($\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-$) = (34.69 ± 0.34) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R82;LINKAGE=A

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.15±0.05	¹ ABLIKIM	15AE	BES3 $J/\psi \rightarrow \gamma\pi^0\pi^0$

 Γ_{244}/Γ

NODE=M070B00
NODE=M070B00

¹ The uncertainty is systematic as statistical is negligible.

NODE=M070B00;LINKAGE=A

 $\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ±0.5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.		

 Γ_{245}/Γ


NODE=M070R95
NODE=M070R95

4.32±0.14±0.73	¹ BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
2.08±0.13±0.35	² BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$
3.05±0.08±0.45	² BALTRUSAIT..	86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$
4.85±0.45±1.20	³ BURKE	82	MRK2 e^+e^-

OCCUR=2

¹ 4 π mass less than 3.0 GeV.
² 4 π mass less than 2.0 GeV.
³ 4 π mass less than 2.5 GeV.

NODE=M070R95;LINKAGE=A
NODE=M070R95;LINKAGE=B
NODE=M070R95;LINKAGE=M



$\Gamma(\gamma f_2(1270) f_2(1270))/\Gamma_{total}$ Γ_{246}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.5±0.7±1.6	646 ± 45	ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070S45
NODE=M070S45

$\Gamma(\gamma f_2(1270) f_2(1270) (\text{non resonant}))/\Gamma_{total}$ Γ_{247}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.2±0.8±1.7	¹ ABLIKIM	04M BES	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M070S46
NODE=M070S46

¹ Subtracting contribution from intermediate $\eta_c(1S)$ decays.

NODE=M070S46;LINKAGE=AB

$\Gamma(\gamma \pi^+ \pi^- 2\pi^0)/\Gamma_{total}$ Γ_{248}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.3±0.2±3.1	¹ BALTRUSAIT..86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$

NODE=M070R99
NODE=M070R99

¹ 4π mass less than 2.0 GeV.

NODE=M070R99;LINKAGE=M

$\Gamma(\gamma K_S^0 K_S^0)/\Gamma_{total}$ Γ_{249}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.1±0.4	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P73
NODE=M070P73

$\Gamma(\gamma(K\bar{K}\pi) [J^{PC} = 0^{-+}])/ \Gamma_{total}$ Γ_{250}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.7 ±0.4 OUR AVERAGE	Error includes scale factor of 2.1.		
0.58±0.03±0.20	¹ BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
2.1 ±0.1 ±0.7	² BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$

NODE=M070S38
NODE=M070S38

OCCUR=2

¹ For a broad structure around 1800 MeV.

² For a broad structure around 2040 MeV.

NODE=M070S38;LINKAGE=BD
NODE=M070S38;LINKAGE=BE

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{total}$ Γ_{251}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.1±0.6	1516	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B05
NODE=M070B05

$\Gamma(\gamma K^*(892) \bar{K}^*(892))/\Gamma_{total}$ Γ_{252}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0±0.3±1.3	320	¹ BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$

NODE=M070B07
NODE=M070B07

¹ Summed over all charges.

NODE=M070R;LINKAGE=B7

$\Gamma(\gamma\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.090±0.013 OUR AVERAGE[(1.085 ± 0.018) × 10⁻³ OUR 2023 AVERAGE]

1.096±0.001±0.019	2.2M	ABLIKIM	23BD BES3	$J/\psi \rightarrow \eta\gamma$
1.067±0.005±0.023	87.9k	ABLIKIM	21AMBES3	$e^+e^- \rightarrow J/\psi$
1.12 ±0.05 ±0.01	18.6k	¹ ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
1.101±0.029±0.022		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta\gamma$
1.123±0.089	11k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

0.88 ±0.08 ±0.11		BLOOM	83 CBAL	e^+e^-
0.82 ±0.10		BRANDELIK	79C DASP	e^+e^-
1.3 ±0.4	21	BARTEL	77 CNTR	e^+e^-

¹ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = (4.42 \pm 0.04 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(\gamma\eta\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

21.4±1.8±2.5

21.4±1.8±2.5	596	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$
--------------	-----	---------	----------	------------------------------

 $\Gamma(\gamma a_0(980)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.5 × 10⁻⁶

<2.5 × 10 ⁻⁶	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$
-------------------------	----	---------	----------	------------------------------

 $\Gamma(\gamma a_2(1320)^0 \rightarrow \gamma\eta\pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<6.6 × 10⁻⁶

<6.6 × 10 ⁻⁶	95	ABLIKIM	16P BES3	$J/\psi \rightarrow 5\gamma$
-------------------------	----	---------	----------	------------------------------

 $\Gamma(\gamma\eta\pi\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.1 ±1.0 OUR AVERAGE

5.85±0.3±1.05	¹ EDWARDS	83B CBAL	$J/\psi \rightarrow \eta\pi^+\pi^-$
7.8 ±1.2±2.4	¹ EDWARDS	83B CBAL	$J/\psi \rightarrow \eta 2\pi^0$

¹Broad enhancement at 1700 MeV. $\Gamma(\gamma\eta_2(1870) \rightarrow \gamma\eta\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.2±2.2±0.9

6.2±2.2±0.9	BAI	99 BES	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
-------------	-----	--------	---

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.28±0.06 OUR AVERAGE

Error includes scale factor of 1.3. See the ideogram below.

[(5.25 ± 0.07) × 10⁻³ OUR 2023 AVERAGE Scale factor = 1.3]

5.40±0.01±0.11	638k	ABLIKIM	23BD BES3	$J/\psi \rightarrow \gamma\eta'$
5.27±0.03±0.05	36k	ABLIKIM	19T BES	$J/\psi \rightarrow \gamma\eta'$
5.43±0.23±0.09	5.0k	¹ ABLIKIM	180 BES3	$\psi(2S) \rightarrow \pi^+\pi^-\gamma\gamma\gamma$
4.77±0.22±0.06		² ABLIKIM	11 BES3	$J/\psi \rightarrow \eta'\gamma$
5.24±0.12±0.11		PEDLAR	09 CLE3	$J/\psi \rightarrow \eta'\gamma$
5.55±0.44	35k	ABLIKIM	06E BES2	$J/\psi \rightarrow \eta'\gamma$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

4.50±0.14±0.53		BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$
----------------	--	--------	----------	--

4.30±0.31±0.71		BOLTON	92B MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \pi^+\pi^-\pi^0$
----------------	--	--------	----------	---

4.04±0.16±0.85	622	AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
----------------	-----	----------	--------	---

4.39±0.09±0.66	2420	AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
----------------	------	----------	--------	---

4.1 ±0.3 ±0.6		BLOOM	83 CBAL	$e^+e^- \rightarrow 3\gamma + \text{hadrons}$
---------------	--	-------	---------	---

2.9 ±1.1	6	BRANDELIK	79C DASP	$e^+e^- \rightarrow 3\gamma$
----------	---	-----------	----------	------------------------------

2.4 ±0.7	57	BARTEL	76 CNTR	$e^+e^- \rightarrow 2\gamma\rho$
----------	----	--------	---------	----------------------------------

 Γ_{253}/Γ

NODE=M070R83

NODE=M070R83

NEW

 Γ_{254}/Γ

NODE=M070P01

NODE=M070P01

 Γ_{258}/Γ

NODE=M070P02

NODE=M070P02

 Γ_{259}/Γ

NODE=M070P03

NODE=M070P03

 Γ_{260}/Γ

NODE=M070R96

NODE=M070R96

 Γ_{261}/Γ

NODE=M070S37

NODE=M070S37

 Γ_{262}/Γ

NODE=M070R84

NODE=M070R84


NEW

¹ ABLIKIM 180 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] = (1.26 \pm 0.02 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] \times [B(\eta'(958) \rightarrow \gamma\gamma)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.49 \pm 0.30) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \gamma\gamma) = (2.307 \pm 0.033) \times 10^{-2}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R84;LINKAGE=A

² ABLIKIM 11 reports $(4.84 \pm 0.03 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(J/\psi(1S) \rightarrow \gamma\eta'(958))/\Gamma_{\text{total}}] / [B(\eta'(958) \rightarrow \pi^+\pi^-\eta)] / [B(\eta \rightarrow 2\gamma)]$ assuming $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (43.2 \pm 0.7) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.31 \pm 0.20) \times 10^{-2}$, which we rescale to our best values $B(\eta'(958) \rightarrow \pi^+\pi^-\eta) = (42.5 \pm 0.5) \times 10^{-2}$, $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M070R84;LINKAGE=AB



$\Gamma(\gamma f_0(500) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{255}/Γ

NODE=M070P95
NODE=M070P95

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

..... We do not use the following data for averages, fits, limits, etc.

10.5 ± 2.0 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{256}/Γ

NODE=M070P97
NODE=M070P97

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

..... We do not use the following data for averages, fits, limits, etc.

5 ± 5 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(500) \rightarrow \gamma\eta\eta)/\Gamma_{\text{total}}$ Γ_{257}/Γ

NODE=M070P98
NODE=M070P98

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

..... We do not use the following data for averages, fits, limits, etc.

4 ± 3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma\pi\pi)/\Gamma_{\text{total}}$ Γ_{263}/Γ

NODE=M070P90
NODE=M070P90

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

..... We do not use the following data for averages, fits, limits, etc.

1.3 ± 0.2 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma f_0(980) \rightarrow \gamma K\bar{K})/\Gamma_{\text{total}}$ Γ_{264}/Γ

NODE=M070P91
NODE=M070P91

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT

..... We do not use the following data for averages, fits, limits, etc.

0.8 ± 0.3 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

$\Gamma(\gamma\rho\rho)/\Gamma_{total}$ Γ_{265}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
4.5 ± 0.8 OUR AVERAGE				
4.7 ± 0.3 ± 0.9		¹ BALTRUSAIT...86B	MRK3	$J/\psi \rightarrow 4\pi\gamma$
3.75 ± 1.05 ± 1.20		² BURKE	82 MRK2	$J/\psi \rightarrow 4\pi\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.09	90	³ BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
¹ 4π mass less than 2.0 GeV.				
² 4π mass less than 2.0 GeV. We have multiplied $2\rho^0$ measurement by 3 to obtain 2ρ .				
³ 4π mass in the range 2.0–25 GeV.				

NODE=M070R94
NODE=M070R94

NODE=M070R94;LINKAGE=N
NODE=M070R94;LINKAGE=M
NODE=M070R94;LINKAGE=A

$\Gamma(\gamma\rho\omega)/\Gamma_{total}$ Γ_{266}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10⁻⁴	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070R05
NODE=M070R05

$\Gamma(\gamma\rho\phi)/\Gamma_{total}$ Γ_{267}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.8 × 10⁻⁵	90	ABLIKIM	08A BES2	$e^+e^- \rightarrow J/\psi$

NODE=M070R06
NODE=M070R06

$\Gamma(\gamma\omega\omega)/\Gamma_{total}$ Γ_{268}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61 ± 0.33 OUR AVERAGE				
6.0 ± 4.8 ± 1.8		ABLIKIM	08A BES2	$J/\psi \rightarrow \gamma\omega\pi^+\pi^-$
1.41 ± 0.2 ± 0.42	120 ± 17	BISELLO	87 SPEC	e^+e^- , hadrons γ
1.76 ± 0.09 ± 0.45		BALTRUSAIT...85C	MRK3	$e^+e^- \rightarrow$ hadrons γ


NODE=M070R97
NODE=M070R97

$\Gamma(\gamma\phi\phi)/\Gamma_{total}$ Γ_{269}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.2 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
7.5 ± 0.6 ± 1.2	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
3.4 ± 0.8 ± 0.6	33 ± 7	¹ BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
3.1 ± 0.7 ± 0.4		¹ BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
¹ $\phi\phi$ mass less than 2.9 GeV, η_c excluded.				

NODE=M070R98
NODE=M070R98

NODE=M070R98;LINKAGE=C



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{total}$ Γ_{270}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.		
1.66 ± 0.1 ± 0.58	^{1,2} BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
3.8 ± 0.3 ± 0.6	³ AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
4.0 ± 0.7 ± 1.0	³ EDWARDS	82E CBAL	$J/\psi \rightarrow K^+ K^- \pi^0 \gamma$
4.3 ± 1.7	^{3,4} SCHARRE	80 MRK2	e^+e^-

NODE=M070R89
NODE=M070R89

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78 ± 0.21 ± 0.33	3,5,6	AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$	
0.83 ± 0.13 ± 0.18	3,7,8	AUGUSTIN	92	DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$	OCCUR=2
0.66 ^{+0.17+0.24} _{-0.16-0.15}	3,6,9	BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
1.03 ^{+0.21+0.26} _{-0.18-0.19}	3,8,10	BAI	90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	OCCUR=2

¹ Interference with the $J/\psi(1S)$ radiative transition to the broad $K \bar{K} \pi$ pseudoscalar state around 1800 is $(0.15 \pm 0.01 \pm 0.05) \times 10^{-3}$.

² Interference with $J/\psi \rightarrow \gamma f_1(1420)$ is $(-0.03 \pm 0.01 \pm 0.01) \times 10^{-3}$.

³ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

⁴ Corrected for spin-zero hypothesis for $\eta(1405)$.

⁵ From fit to the $a_0(980) \pi 0^-+$ partial wave.

⁶ $a_0(980) \pi$ mode.

⁷ From fit to the $K^*(892) K 0^-+$ partial wave.

⁸ $K^* K$ mode.

⁹ From $a_0(980) \pi$ final state.

¹⁰ From $K^*(890) K$ final state.

NODE=M070R89;LINKAGE=BD

NODE=M070R89;LINKAGE=BE

NODE=M070R89;LINKAGE=B

NODE=M070R89;LINKAGE=C

NODE=M070R89;LINKAGE=H


NODE=M070R89;LINKAGE=K9

NODE=M070R89;LINKAGE=J

NODE=M070R89;LINKAGE=K8

NODE=M070R89;LINKAGE=D

NODE=M070R89;LINKAGE=E



$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\rho^0) / \Gamma_{\text{total}}$ Γ_{271}/Γ

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

0.78 ± 0.20 OUR AVERAGE Error includes scale factor of 1.8.

1.07 ± 0.17 ± 0.11 ¹ BAI 04J BES2 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

0.64 ± 0.12 ± 0.07 ¹ COFFMAN 90 MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

¹ Includes unknown branching fraction $\eta(1405) \rightarrow \gamma\rho^0$.

NODE=M070S30

NODE=M070S30

NODE=M070S30;LINKAGE=C

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\eta\pi^+\pi^-) / \Gamma_{\text{total}}$ Γ_{272}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

3.0 ± 0.5 OUR AVERAGE

2.6 ± 0.7 ± 0.4 BAI 99 BES $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

3.38 ± 0.33 ± 0.64 ¹ BOLTON 92B MRK3 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

7.0 ± 0.6 ± 1.1 261 ² AUGUSTIN 90 DM2 $J/\psi \rightarrow \gamma\eta\pi^+\pi^-$

¹ Via $a_0(980) \pi$.

² Includes unknown branching fraction to $\eta\pi^+\pi^-$.

NODE=M070S29

NODE=M070S29

NODE=M070S29;LINKAGE=RR

NODE=M070S29;LINKAGE=R

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\rho^0\rho^0) / \Gamma_{\text{total}}$ Γ_{273}/Γ

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT

1.7 ± 0.4 OUR AVERAGE Error includes scale factor of 1.3.

2.1 ± 0.4 BUGG 95 MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\pi^+\pi^-$

1.36 ± 0.38 ^{1,2} BISELLO 89B DM2 $J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits.

² Includes unknown branching fraction to $\rho^0\rho^0$.

NODE=M070S19

NODE=M070S19

NODE=M070S19;LINKAGE=A

NODE=M070S19;LINKAGE=B

$\Gamma(\gamma\eta(1405/1475) \rightarrow \gamma\gamma\phi)/\Gamma_{\text{total}}$ Γ_{274}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<82	95		BAI	04J	BES2 $J/\psi \rightarrow \gamma\gamma K^+ K^-$

NODE=M070R77
 NODE=M070R77

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.03 \pm 0.92 \pm 0.91$	1.3k	1	ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$
$10.36 \pm 1.51 \pm 1.54$	1.9k	2	ABLIKIM	18I	BES3 $J/\psi \rightarrow \gamma\gamma\phi(1020)$

OCCUR=2

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

NODE=M070R77;LINKAGE=B

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

NODE=M070R77;LINKAGE=A

 $\Gamma(\gamma\eta(1405) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{275}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.63 $\times 10^{-6}$	90	ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

NODE=M070P38
 NODE=M070P38

 $\Gamma(\gamma\eta(1475) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{276}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.86 $\times 10^{-6}$	90	ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

NODE=M070P39
 NODE=M070P39

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{\text{total}}$ Γ_{277}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.09	1,2 BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

NODE=M070S20
 NODE=M070S20

¹ Estimated by us from various fits.

NODE=M070S20;LINKAGE=A

² Includes unknown branching fraction to $\rho^0\rho^0$.

NODE=M070S20;LINKAGE=B

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\omega\omega)/\Gamma_{\text{total}}$ Γ_{278}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.98 $\pm 0.08 \pm 0.32$	1045	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma\omega\omega$

NODE=M070R04
 NODE=M070R04

 $\Gamma(\gamma\eta(1760) \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{279}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.80 $\times 10^{-6}$	90	ABLIKIM	18O	BES3 $\psi(2S) \rightarrow \pi^+ \pi^- \gamma\gamma\gamma$

NODE=M070P40
 NODE=M070P40

 $\Gamma(\gamma\eta(2225))/\Gamma_{\text{total}}$ Γ_{280}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S21
 NODE=M070S21

3.14 $^{+0.50}_{-0.19}$ OUR AVERAGE

2.40 ± 0.10 $^{+2.47}_{-0.18}$		1,2	ABLIKIM	16N	BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
4.4 $\pm 0.4 \pm 0.8$	196	2	ABLIKIM	08I	BES $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
3.3 $\pm 0.8 \pm 0.5$		2	BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
2.7 $\pm 0.6 \pm 0.6$		2	BAI	90B	MRK3 $J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
2.4 $^{+1.5}_{-1.0}$		3,4	BISELLO	89B	DM2 $J/\psi \rightarrow 4\pi\gamma$

OCCUR=2

¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$ that also finds significant signals for for $\eta(2100)$, 0^-+ phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^-+ state $X(2500)$ ($M = 2470^{+15+101}_{-19-23}$ MeV, $\Gamma = 230^{+64+56}_{-35-33}$ MeV).

NODE=M070S21;LINKAGE=C

² Includes unknown branching fraction to $\phi\phi$.

NODE=M070S21;LINKAGE=U

³ Estimated by us from various fits.

NODE=M070S21;LINKAGE=A

⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

NODE=M070S21;LINKAGE=B

 $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{281}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R86
 NODE=M070R86


1.63 ± 0.12 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

2.07 ± 0.16 $^{+0.02}_{-0.07}$	2.4k	1,2	DOBBS	15	$J/\psi \rightarrow \gamma\pi\pi$
1.63 ± 0.26 $^{+0.02}_{-0.05}$		3	ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+ \pi^-$
1.42 ± 0.21 $^{+0.02}_{-0.05}$		4	ABLIKIM	06V	BES2 $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^0 \pi^0$
1.33 $\pm 0.05 \pm 0.20$		5	AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma\pi^+ \pi^-$
1.36 $\pm 0.09 \pm 0.23$		5	BALTRUSAIT.	87	MRK3 $J/\psi \rightarrow \gamma\pi^+ \pi^-$
1.48 $\pm 0.25 \pm 0.30$	178		EDWARDS	82B	CBAL $e^+ e^- \rightarrow 2\pi^0 \gamma$
2.0 ± 0.7	35		ALEXANDER	78	PLUT $e^+ e^-$
1.2 ± 0.6	30	6	BRANDELIK	78B	DASP $e^+ e^- \rightarrow \pi^+ \pi^- \gamma$

OCCUR=2

- ¹ Using CLEO-c data but not authored by the CLEO Collaboration.
- ² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.744 \pm 0.052 \pm 0.122) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.371 \pm 0.010 \pm 0.222) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ ABLIKIM 06V reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (1.200 \pm 0.027 \pm 0.174) \times 10^{-3}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.
- ⁶ Restated by us to take account of spread of E1, M2, E3 transitions.

NODE=M070R86;LINKAGE=A
 NODE=M070R86;LINKAGE=DO
 NODE=M070R86;LINKAGE=AI
 NODE=M070R86;LINKAGE=AL
 NODE=M070R86;LINKAGE=X
 NODE=M070R86;LINKAGE=T



$\Gamma(\gamma f_2(1270) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$	Γ_{282}/Γ		
VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$2.58^{+0.08+0.59}_{-0.09-0.20}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P68
 NODE=M070P68

$\Gamma(\gamma f_1(1285))/\Gamma_{\text{total}}$	Γ_{283}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.61 ± 0.08 OUR AVERAGE			
0.69 $\pm 0.16 \pm 0.20$	1 BAI	04J BES2	$J/\psi \rightarrow \gamma \gamma \rho^0$
0.61 $\pm 0.04 \pm 0.21$	2 BAI	00D BES	$J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
0.45 $\pm 0.09 \pm 0.17$	3 BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.625 $\pm 0.063 \pm 0.103$	4 BOLTON	92 MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.70 $\pm 0.08 \pm 0.16$	5 BOLTON	92B MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M070R88
 NODE=M070R88

- ¹ Assuming $B(f_1(1285) \rightarrow \rho^0 \gamma) = 0.055 \pm 0.013$.
- ² Assuming $\Gamma(f_1(1285) \rightarrow K \bar{K} \pi)/\Gamma_{\text{total}} = 0.090 \pm 0.004$.
- ³ Assuming $\Gamma(f_1(1285) \rightarrow \eta \pi \pi)/\Gamma_{\text{total}} = 0.5 \pm 0.18$.
- ⁴ Obtained summing the sequential decay channels
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow a_0(980) \pi, a_0(980) \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.
- ⁵ Using $B(f_1(1285) \rightarrow a_0(980) \pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta \pi$.

NODE=M070R88;LINKAGE=BI
 NODE=M070R88;LINKAGE=BD
 NODE=M070R88;LINKAGE=BA
 NODE=M070R88;LINKAGE=B

NODE=M070R88;LINKAGE=A

$\Gamma(\gamma f_0(1370) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{284} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

38 ± 10	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	--------------	------	---

NODE=M070P92
NODE=M070P92

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{285} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$4.19 \pm 0.73 \pm 1.34$	478	¹ DOBBS 15		$J/\psi \rightarrow \gamma K \bar{K}$
--------------------------	-----	-----------------------	--	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.3 ± 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------	------	---

NODE=M070R00
NODE=M070R00

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070R00;LINKAGE=A

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{286} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.07^{+0.08+0.36}_{-0.07-0.34}$	ABLIKIM 18AA	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	--------------	------	---

NODE=M070P63
NODE=M070P63

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{287} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5 ± 1.0	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------	------	---

NODE=M070P93
NODE=M070P93

 $\Gamma(\gamma f_0(1370) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{288} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9 ± 0.3	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------	------	---

NODE=M070P94
NODE=M070P94

 $\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{\text{total}}$ Γ_{289} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.79 ± 0.13 OUR AVERAGE

$0.68 \pm 0.04 \pm 0.24$	BAI	00D	BES $J/\psi \rightarrow \gamma K^\pm K_S^0 \pi^\mp$
--------------------------	-----	-----	---

$0.76 \pm 0.15 \pm 0.21$	^{1,2} AUGUSTIN	92	DM2 $J/\psi \rightarrow \gamma K \bar{K} \pi$
--------------------------	-------------------------	----	---

$0.87 \pm 0.14^{+0.14}_{-0.11}$	¹ BAI	90C	MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
---------------------------------	------------------	-----	--

NODE=M070S31
NODE=M070S31

OCCUR=2

¹ Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.

NODE=M070S31;LINKAGE=A

² From fit to the $K^*(892)K 1^{++}$ partial wave.

NODE=M070S31;LINKAGE=D

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{290} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.09 ± 0.24 OUR AVERAGE

$1.21 \pm 0.29 \pm 0.24$	174	¹ DOBBS 15		$J/\psi \rightarrow \gamma \pi \pi$
--------------------------	-----	-----------------------	--	-------------------------------------

$1.00 \pm 0.03 \pm 0.45$		² ABLIKIM 06V	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
--------------------------	--	--------------------------	------	---

$1.02 \pm 0.09 \pm 0.45$		² ABLIKIM 06V	BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$
--------------------------	--	--------------------------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.90 ± 0.17	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------------	--------------	------	---

5.7 ± 0.8	^{3,4} BUGG	95	MRK3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \pi^+ \pi^-$
---------------	---------------------	----	--

NODE=M070S32
NODE=M070S32

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070S32;LINKAGE=C

² Including unknown branching fraction to $\pi\pi$.

NODE=M070S32;LINKAGE=AB

³ Including unknown branching ratio for $f_0(1500) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$.

NODE=M070S32;LINKAGE=A

⁴ Assuming that $f_0(1500)$ decays only to two S-wave dipions.

NODE=M070S32;LINKAGE=B

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{291} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.65^{+0.26+0.51}_{-0.31-1.40}$	5.5k	¹ ABLIKIM 13N	BES3	$J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	--------------------------	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.1 ± 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	--------------	------	---

NODE=M070S83
NODE=M070S83

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S83;LINKAGE=A

$\Gamma(\gamma f_0(1500) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{total}$ Γ_{292} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.16^{+0.18}_{-0.56}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.7 ± 0.3	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P64
NODE=M070P64

$\Gamma(\gamma f_0(1500) \rightarrow \gamma \eta \eta') / \Gamma_{total}$ Γ_{293} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
$18.1 \pm 1.1^{+1.9}_{-1.3}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
12 ± 5	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.			

NODE=M070P99
NODE=M070P99

NODE=M070P99;LINKAGE=A

$\Gamma(\gamma f_1(1510) \rightarrow \gamma \eta \pi^+ \pi^-) / \Gamma_{total}$ Γ_{294} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$4.5 \pm 1.0 \pm 0.7$	BAI	99 BES	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M070S36
NODE=M070S36

$\Gamma(\gamma f_2'(1525)) / \Gamma_{total}$ Γ_{295} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7^{+0.8}_{-0.5}$			OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.		
$8.1 \pm 0.9 \pm 0.2$	750	1,2	DOBBS	15	$J/\psi \rightarrow \gamma K\bar{K}$
$3.85 \pm 0.17^{+1.91}_{-0.73}$		3	BAI	03G BES	$J/\psi \rightarrow \gamma K\bar{K}$
$3.6 \pm 0.4^{+1.4}_{-0.4}$		3	BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$5.6 \pm 1.4 \pm 0.9$		3	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$4.5 \pm 0.4 \pm 0.9$		3	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.8 \pm 1.6 \pm 1.4$		3	BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<3.4	90	4	BRANDELIK	79C DASP	$e^+ e^- \rightarrow \pi^+ \pi^- \gamma$
<2.3	90	3	ALEXANDER	78 PLUT	$e^+ e^- \rightarrow K^+ K^- \gamma$


NODE=M070R87
NODE=M070R87

OCCUR=3
OCCUR=4
OCCUR=2

- ¹ Using CLEO-c data but not authored by the CLEO Collaboration.
- ² DOBBS 15 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma f_2'(1525)) / \Gamma_{total}] \times [B(f_2'(1525) \rightarrow K\bar{K})] = (7.09 \pm 0.46 \pm 0.67) \times 10^{-4}$ which we divide by our best value $B(f_2'(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ Using $B(f_2'(1525) \rightarrow K\bar{K}) = 0.888$.
- ⁴ Assuming isotropic production and decay of the $f_2'(1525)$ and isospin.

NODE=M070R87;LINKAGE=B
NODE=M070R87;LINKAGE=D0

NODE=M070R87;LINKAGE=A1
NODE=M070R87;LINKAGE=I



$\Gamma(\gamma f_2(1565) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{298} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.32 \pm 0.05^{+0.12}_{-0.02}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.

NODE=M070Q31
NODE=M070Q31

NODE=M070Q31;LINKAGE=A

 $\Gamma(\gamma f_2'(1525) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{296} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$7.99^{+0.03+0.69}_{-0.04-0.50}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
----------------------------------	---------	-----------	---

NODE=M070P69
NODE=M070P69

 $\Gamma(\gamma f_2'(1525) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{297} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.42^{+0.43+1.37}_{-0.51-1.30}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	----------	---------------------------------------

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S86
NODE=M070S86

NODE=M070S86;LINKAGE=A

 $\Gamma(\gamma f_2(1640) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ Γ_{299} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.28 \pm 0.05 \pm 0.17$	141	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
--------------------------	-----	---------	---------	---

NODE=M070R02
NODE=M070R02

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{300} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.8 ± 0.5 OUR AVERAGE

$3.72 \pm 0.30 \pm 0.43$	483	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
$3.96 \pm 0.06 \pm 1.12$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$
$3.99 \pm 0.15 \pm 2.64$		² ABLIKIM	06V BES2	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma \pi^0 \pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.2		³ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
$2.5 \pm 1.6 \pm 0.8$		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Including unknown branching fraction to $\pi \pi$.

³ There is a further $(2.4 \pm 0.8) \times 10^{-4}$ scalar contribution at 1765 MeV.

NODE=M070B01
NODE=M070B01

OCCUR=2

NODE=M070B01;LINKAGE=A
NODE=M070B01;LINKAGE=A
NODE=M070B01;LINKAGE=B

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{301} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

9.5 ± 1.0 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.

$8.00^{+0.12+1.24}_{-0.08-0.40}$		¹ ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$11.76 \pm 0.54 \pm 0.94$	1.2k	² DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
$9.62 \pm 0.29^{+3.51}_{-1.86}$		³ BAI	03G BES	$J/\psi \rightarrow \gamma K \bar{K}$
$5.0 \pm 0.8^{+1.8}_{-0.4}$		^{1,4} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
$9.2 \pm 1.4 \pm 1.4$		¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
$10.4 \pm 1.2 \pm 1.6$		¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$9.6 \pm 1.2 \pm 1.8$		¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.3 ± 0.8		⁵ SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma (\pi \pi, K \bar{K}, \eta \eta, \omega \phi)$
$1.6 \pm 0.2^{+0.6}_{-0.2}$		^{1,6} BAI	96C BES	$J/\psi \rightarrow \gamma K^+ K^-$
< 0.8	90	⁷ BISELLO	89B	$J/\psi \rightarrow 4\pi \gamma$
$1.6 \pm 0.4 \pm 0.3$		⁸ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
3.8 ± 1.6		⁹ EDWARDS	82D CBAL	$e^+ e^- \rightarrow \eta \eta \gamma$

NODE=M070R91
NODE=M070R91


OCCUR=2

OCCUR=2

OCCUR=2

- 1 Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied $K^+ K^-$ measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.
- 2 Using CLEO-c data but not authored by the CLEO Collaboration.
- 3 Includes unknown branching ratio to $K^+ K^-$ or $K_S^0 K_S^0$.
- 4 Assuming $J^P = 2^+$ for $f_0(1710)$.
- 5 There is a further $(6 \pm 2) \times 10^{-4}$ scalar contribution at 1765 MeV.
- 6 Assuming $J^P = 0^+$ for $f_0(1710)$.
- 7 Includes unknown branching fraction to $\rho^0 \rho^0$.
- 8 Includes unknown branching fraction to $\pi^+ \pi^-$.
- 9 Includes unknown branching fraction to $\eta\eta$.

NODE=M070R91;LINKAGE=B
 NODE=M070R91;LINKAGE=D
 NODE=M070R91;LINKAGE=K9
 NODE=M070R91;LINKAGE=A1
 NODE=M070R91;LINKAGE=E
 NODE=M070R91;LINKAGE=A2
 NODE=M070R91;LINKAGE=C
 NODE=M070R91;LINKAGE=Z
 NODE=M070R91;LINKAGE=A



$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}}$ **Γ_{302} / Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.31 \pm 0.06 \pm 0.08$	180	ABLIKIM	06H	BES $J/\psi \rightarrow \gamma \omega \omega$

NODE=M070R01
 NODE=M070R01

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ **Γ_{303} / Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.35^{+0.13+1.24}_{-0.11-0.74}$	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$

NODE=M070S84
 NODE=M070S84

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 1.2 \pm 0.4 ² SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
- ¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.
- ² There is a further $(0.7 \pm 0.1) \times 10^{-4}$ scalar contribution at 1765 MeV.

NODE=M070S84;LINKAGE=A
 NODE=M070S84;LINKAGE=B

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ **Γ_{304} / Γ**

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
6.5 ± 2.5	¹ SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

¹ There is a further $(2.5 \pm 1.1) \times 10^{-5}$ scalar contribution at 1765 MeV.

NODE=M070Q00
 NODE=M070Q00

NODE=M070Q00;LINKAGE=A

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \omega \phi) / \Gamma_{\text{total}}$ **Γ_{305} / Γ**

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.6 OUR AVERAGE				
$2.00 \pm 0.08^{+1.38}_{-1.64}$	1.3k	ABLIKIM	13J	BES3 $J/\psi \rightarrow \gamma \omega \phi$
$2.61 \pm 0.27 \pm 0.65$	95	ABLIKIM	06J	BES2 $J/\psi \rightarrow \gamma \omega \phi$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 0.1 \pm 0.1 ¹ SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
- ¹ There is a further $(2.2 \pm 0.4) \times 10^{-4}$ scalar contribution at 1765 MeV.

NODE=M070S79
 NODE=M070S79

NODE=M070S79;LINKAGE=A

$$\Gamma(\gamma f_0(1770) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}} \quad \Gamma_{306} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$1.11 \pm 0.06^{+0.19}_{-0.32}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P65
NODE=M070P65

$$\Gamma(\gamma f_0(1770) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}} \quad \Gamma_{309} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q30
NODE=M070Q30

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.11 \pm 0.01^{+0.04}_{-0.03}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.

NODE=M070Q30;LINKAGE=A

$$\Gamma(\gamma f_2(1810) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}} \quad \Gamma_{307} / \Gamma$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
--------------------------	------	-------------	---------

NODE=M070S87
NODE=M070S87

$5.40^{+0.60+3.42}_{-0.67-2.35}$	5.5k	¹ ABLIKIM	13N $J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S87;LINKAGE=A

$$\Gamma(\gamma \eta_1(1855) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}} \quad \Gamma_{308} / \Gamma$$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q44
NODE=M070Q44

$2.70 \pm 0.41^{+0.16}_{-0.35}$	¹ ABLIKIM	22AI BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and the resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave. For analysis details see ABLIKIM 22AS.

NODE=M070Q44;LINKAGE=A

$$\Gamma(\gamma f_2(1910) \rightarrow \gamma \omega \omega) / \Gamma_{\text{total}} \quad \Gamma_{310} / \Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070R03
NODE=M070R03

$0.20 \pm 0.04 \pm 0.13$	151	ABLIKIM	06H BES	$J/\psi \rightarrow \gamma \omega \omega$
--------------------------	-----	---------	---------	---

$$\Gamma(\gamma f_2(1950) \rightarrow \gamma K^*(892) \bar{K}^*(892)) / \Gamma_{\text{total}} \quad \Gamma_{311} / \Gamma$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070B06
NODE=M070B06

$0.7 \pm 0.1 \pm 0.2$	BAI	00B BES	$J/\psi \rightarrow \gamma K^+ K^0 \pi^+ \pi^-$
-----------------------	-----	---------	---

$$\Gamma(\gamma f_2(2010) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}} \quad \Gamma_{312} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q37
NODE=M070Q37

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.71 \pm 0.06^{+0.10}_{-0.06}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.

NODE=M070Q37;LINKAGE=A

$$\Gamma(\gamma f_0(2020) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}} \quad \Gamma_{313} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q01
NODE=M070Q01

• • • We do not use the following data for averages, fits, limits, etc. • • •

42 ± 10	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

$$\Gamma(\gamma f_0(2020) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}} \quad \Gamma_{314} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q02
NODE=M070Q02

• • • We do not use the following data for averages, fits, limits, etc. • • •

55 ± 25	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

$$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}} \quad \Gamma_{315} / \Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q03
NODE=M070Q03

• • • We do not use the following data for averages, fits, limits, etc. • • •

10 ± 10	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

$$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}} \quad \Gamma_{316} / \Gamma$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070Q25
NODE=M070Q25

$2.63 \pm 0.06^{+0.31}_{-0.46}$	¹ ABLIKIM	22C BES3	$J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
---------------------------------	----------------------	----------	---

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q25;LINKAGE=A

$\Gamma(\gamma f_0(2020) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{317}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.28 \pm 0.12^{+0.29}_{-0.20}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.

NODE=M070Q35
NODE=M070Q35

NODE=M070Q35;LINKAGE=A

 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{318}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.7 \pm 0.5 \pm 0.5$	¹ BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
-----------------------	-----------------------------	------	---

¹ Assuming branching fraction $f_4(2050) \rightarrow \pi \pi / \text{total} = 0.167$.

NODE=M070S7
NODE=M070S7

NODE=M070S7;LINKAGE=V

 $\Gamma(\gamma f_4(2050) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{319}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06 \pm 0.01^{+0.03}_{-0.01}$	¹ ABLIKIM	22AS BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta'$
---------------------------------	----------------------	-----------	--

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P-wave.

NODE=M070Q32
NODE=M070Q32

NODE=M070Q32;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{320}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.13^{+0.09+0.64}_{-0.10-0.28}$	5.5k	¹ ABLIKIM	13N BES3	$J/\psi \rightarrow \gamma \eta \eta$
----------------------------------	------	----------------------	----------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.8 ± 1.5	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	---------	---

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S85
NODE=M070S85

NODE=M070S85;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{322}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$6.24 \pm 0.48 \pm 0.87$	744	¹ DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
--------------------------	-----	--------------------	----	-------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.8	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	---------	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070B08
NODE=M070B08

NODE=M070B08;LINKAGE=A

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{321}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

32 ± 20	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-------------	-----------	---------	---

NODE=M070Q04
NODE=M070Q04

 $\Gamma(\gamma f_0(2200)) / \Gamma_{\text{total}}$ Γ_{323}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5	¹ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
-----	-----------------------	--------	---

¹ Includes unknown branching fraction to $K_S^0 K_S^0$.

NODE=M070S18
NODE=M070S18

NODE=M070S18;LINKAGE=A

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma \pi \pi) / \Gamma_{\text{total}}$ Γ_{326}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5 ± 2	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	-----------	---------	---

NODE=M070Q05
NODE=M070Q05

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K}) / \Gamma_{\text{total}}$ Γ_{324}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$5.86 \pm 0.49 \pm 1.20$	490	¹ DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
--------------------------	-----	--------------------	----	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.5 ± 0.5	SARANTSEV	21 RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
---------------	-----------	---------	---

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M070B09
NODE=M070B09

NODE=M070B09;LINKAGE=A

$\Gamma(\gamma f_0(2200) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{325}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
$2.72^{+0.08+0.17}_{-0.06-0.47}$	ABLIKIM	18AA BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070P66
 NODE=M070P66

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma \eta \eta)/\Gamma_{\text{total}}$ Γ_{327}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.7 ± 0.4 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070Q06
 NODE=M070Q06

 $\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{328}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>300			¹ BAI	96B BES	$e^+e^- \rightarrow \gamma \bar{p}p, K\bar{K}$
>250	99.9		² HASAN	96 SPEC	$\bar{p}p \rightarrow \pi^+\pi^-$
< 2.3	95		³ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95		³ AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$12.4^{+6.4}_{-5.2} \pm 2.8$		23	³ BALTRUSAIT...86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
$8.4^{+3.4}_{-2.8} \pm 1.6$		93	³ BALTRUSAIT...86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

NODE=M070R92
 NODE=M070R92

OCCUR=2

OCCUR=2

¹ Using BARNES 93.

² Using BAI 96B.

³ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

NODE=M070R92;LINKAGE=A
 NODE=M070R92;LINKAGE=M
 NODE=M070R92;LINKAGE=W

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{329}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.9	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma \pi \pi$
$14 \pm 8 \pm 4$		BAI	98H BES	$J/\psi \rightarrow \gamma \pi^0 \pi^0$
$8.4 \pm 2.6 \pm 3.0$		BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma \pi^+ \pi^-$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $2.6/5.2 \times 10^{-5}$ and $1.3/1.9 \times 10^{-5}$, respectively.

NODE=M070B02
 NODE=M070B02

NODE=M070B02;LINKAGE=A
 NODE=M070B02;LINKAGE=D0

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{330}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.1	90	^{1,2} DOBBS	15	$J/\psi \rightarrow \gamma K \bar{K}$
< 3.6		³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
< 2.9		³ DEL-AMO-SA..100	BABR	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
$6.6 \pm 2.9 \pm 2.4$		BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
$10.8 \pm 4.0 \pm 3.2$		BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$

NODE=M070B03
 NODE=M070B03

OCCUR=2

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $1.7/3.1 \times 10^{-5}$ and $1.2/2.0 \times 10^{-5}$, respectively.

³ For spin 2 and helicity 0; other combinations lead to more stringent upper limits.

NODE=M070B03;LINKAGE=A
 NODE=M070B03;LINKAGE=D0

NODE=M070B03;LINKAGE=DE

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \rho \bar{\rho})/\Gamma_{\text{total}}$ Γ_{331}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.5 \pm 0.6 \pm 0.5$ BAI 96B BES $e^+e^- \rightarrow J/\psi \rightarrow \gamma \rho \bar{\rho}$

NODE=M070B04
 NODE=M070B04

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{332}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$4.95 \pm 0.21^{+0.66}_{-0.72}$ ABLIKIM 18AA BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 ± 0.1 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070P67
 NODE=M070P67

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{333}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 2 SARANTSEV 21 RVUE $J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$

NODE=M070Q07
 NODE=M070Q07

$\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{334} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ± 0.4	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma(\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
-----------	--------------	------	---

NODE=M070Q08
NODE=M070Q08

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$ Γ_{335} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

6.09 ± 0.64^{+4.00}_{-1.68}	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
--	----------------------	-----	--

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q26
NODE=M070Q26

NODE=M070Q26;LINKAGE=A

 $\Gamma(\gamma f_0(2330) \rightarrow \gamma \eta \eta') / \Gamma_{\text{total}}$ Γ_{336} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.10 ± 0.02 ^{+0.01} _{-0.02}	¹ ABLIKIM	22AS	BES3 $J/\psi(1S) \rightarrow \gamma \eta \eta'$
---	----------------------	------	---

¹ From a Breit-Wigner fit involving 9 resonances and a resonating exotic $\eta_1(1855) \rightarrow \eta \eta'$ P -wave.

NODE=M070Q36
NODE=M070Q36

NODE=M070Q36;LINKAGE=A

 $\Gamma(\gamma f_0(2470) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$ Γ_{340} / Γ

VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

8.18 ± 1.77^{+3.73}_{-2.23}	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
--	----------------------	-----	--

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q27
NODE=M070Q27

NODE=M070Q27;LINKAGE=A

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{337} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.60^{+0.62+2.37}_{-0.65-2.07}	5.5k	¹ ABLIKIM	13N	BES3 $J/\psi \rightarrow \gamma \eta \eta$
---	------	----------------------	-----	--

¹ From partial wave analysis including all possible combinations of 0^{++} , 2^{++} , and 4^{++} resonances.

NODE=M070S88
NODE=M070S88

NODE=M070S88;LINKAGE=A

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma \eta' \eta') / \Gamma_{\text{total}}$ Γ_{339} / Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

8.67 ± 0.70^{+0.61}_{-1.67}	¹ ABLIKIM	22C	BES3 $J/\psi \rightarrow \gamma \eta' \eta' \rightarrow 4/5 \gamma 2(\pi^+ \pi^-)$
--	----------------------	-----	--

¹ From a partial wave analysis of the systems (γX) , with $X \rightarrow \eta' \eta'$, and $(\eta' X)$, with $X \rightarrow \gamma \eta'$ in the decay $J/\psi \rightarrow \gamma \eta' \eta'$. The intermediate resonance X is parametrized by a constant-width, relativistic Breit-Wigner.

NODE=M070Q29
NODE=M070Q29

NODE=M070Q29;LINKAGE=A

 $\Gamma(\gamma f_2(2340) \rightarrow \gamma K_S^0 K_S^0) / \Gamma_{\text{total}}$ Γ_{338} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

5.54^{+0.34+3.82}_{-0.40-1.49}	ABLIKIM	18AA	BES3 $J/\psi \rightarrow \gamma K_S^0 K_S^0$
---	---------	------	--

NODE=M070P70
NODE=M070P70

 $\Gamma(\gamma X(1835) \rightarrow \gamma \pi^+ \pi^- \eta') / \Gamma_{\text{total}}$ Γ_{341} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.7^{+0.6}_{-0.8}		OUR AVERAGE		Error includes scale factor of 1.6.
--	--	--------------------	--	-------------------------------------

3.93 ± 0.38 ^{+0.31} _{-0.84}	¹ ABLIKIM	16J	BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
---	----------------------	-----	------	---

2.2 ± 0.4 ± 0.4	264	ABLIKIM	05R	BES2 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
-----------------	-----	---------	-----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.87 ± 0.09 ^{+0.49} _{-0.52}	4265	² ABLIKIM	11C	BES3 $J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
---	------	----------------------	-----	--

¹ From a fit of the measured $\pi^+ \pi^- \eta'$ lineshape that accounts for the abrupt distortion observed at the $p\bar{p}$ threshold with a Flatte formula in addition to known backgrounds and contributors, as well as an *ad hoc* Breit-Wigner ($M \approx 1919$ MeV; $\Gamma \approx 51$ MeV) that is required for a good fit. Another explanation for the distortion provided by ABLIKIM 16J is that a second resonance near 1870 MeV interferes with the $X(1835)$; fits to this possibility yield product branching fraction values compatible with that shown within the respective systematic uncertainties.

² From a fit of the $\pi^+ \pi^- \eta'$ mass distribution to a combination of $\gamma f_1(1510)$, $\gamma X(1835)$, and two states $\gamma X(2120)$ and $\gamma X(2370)$, for $M(\pi^+ \pi^- \eta') < 2.8$ GeV, and accounting for backgrounds from non- η' events and $J/\psi \rightarrow \pi^0 \pi^+ \pi^- \eta'$.

NODE=M070R78
NODE=M070R78

NODE=M070R78;LINKAGE=A

NODE=M070R78;LINKAGE=A

$\Gamma(\gamma X(1835) \rightarrow \gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{342}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S71
 NODE=M070S71

0.77^{+0.15}_{-0.09} OUR AVERAGE

0.90 ^{+0.04+0.27} _{-0.11-0.55}		1 ABLIKIM	12D	BES3	$J/\psi \rightarrow \gamma p \bar{p}$
1.14 ^{+0.43+0.42} _{-0.30-0.26}	231	2 ALEXANDER	10	CLEO	$J/\psi \rightarrow \gamma p \bar{p}$
0.70 \pm 0.04 ^{+0.19} _{-0.08}		BAI	03F	BES2	$J/\psi \rightarrow \gamma p \bar{p}$

¹ From the fit including final state interaction effects in isospin 0 S -wave according to SIBIRTSEV 05A.

² From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma X(1835)$, γR with $M(R) = 2100$ MeV and $\Gamma(R) = 160$ MeV, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV.

NODE=M070S71;LINKAGE=AK

NODE=M070S71;LINKAGE=AL

 $\Gamma(\gamma X(1835) \rightarrow \gamma K_S^0 K_S^0 \eta)/\Gamma_{\text{total}}$ Γ_{343}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070S96
 NODE=M070S96

3.31 ^{+0.33+1.96} _{-0.30-1.29}	ABLIKIM	15T	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta$
--	---------	-----	------	--

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \phi(1020))/\Gamma_{\text{total}}$ Γ_{344}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070P37
 NODE=M070P37

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.77 \pm 0.35 \pm 0.25	305	1 ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$
8.09 \pm 1.99 \pm 1.36	1.3k	2 ABLIKIM	18I	BES3	$J/\psi \rightarrow \gamma \gamma \phi(1020)$

OCCUR=2

¹ Constructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

NODE=M070P37;LINKAGE=A

² Destructive interference between the $X(1835)$ and $\eta(1405)/\eta(1475)$ is assumed in a fit to the $\gamma\phi$ invariant mass.

NODE=M070P37;LINKAGE=B

 $\Gamma(\gamma X(1835) \rightarrow \gamma \gamma \gamma)/\Gamma_{\text{total}}$ Γ_{345}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070P41
 NODE=M070P41

$<3.56 \times 10^{-6}$	90	ABLIKIM	180	BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \gamma \gamma \gamma$
------------------------	----	---------	-----	------	---

 $\Gamma(\gamma X(1835) \rightarrow \gamma 3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{346}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M070S82
 NODE=M070S82

2.44 \pm 0.36 ^{+0.60} _{-0.74}	0.6k	ABLIKIM	13U	BES3	$J/\psi \rightarrow \gamma 3(\pi^+ \pi^-)$
---	------	---------	-----	------	--

 $\Gamma(\gamma X(2370) \rightarrow \gamma K^+ K^- \eta')/\Gamma_{\text{total}}$ Γ_{347}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070P86
 NODE=M070P86

1.79 \pm 0.23 \pm 0.65	ABLIKIM	20Q	BES3	$J/\psi \rightarrow \gamma K^+ K^- \eta'$
----------------------------	---------	-----	------	---

 $\Gamma(\gamma X(2370) \rightarrow \gamma K_S^0 K_S^0 \eta')/\Gamma_{\text{total}}$ Γ_{348}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M070P87
 NODE=M070P87

1.18 \pm 0.32 \pm 0.39	ABLIKIM	20Q	BES3	$J/\psi \rightarrow \gamma K_S^0 K_S^0 \eta'$
----------------------------	---------	-----	------	---

 $\Gamma(\gamma X(2370) \rightarrow \gamma \eta \eta \eta')/\Gamma_{\text{total}}$ Γ_{349}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070P88
 NODE=M070P88

$<9.2 \times 10^{-6}$ (CL = 90%)		[$<9.2 \times 10^{-6}$ (CL = 90%) OUR 2023 BEST LIMIT]		
----------------------------------	--	---	--	--

$<9.2 \times 10^{-6}$	90	ABLIKIM	21C	BES3	$J/\psi(1S) \rightarrow \gamma \eta \eta \eta'$
-----------------------	----	---------	-----	------	---

 $\Gamma(\gamma p \bar{p})/\Gamma_{\text{total}}$ Γ_{350}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

NODE=M070R90
 NODE=M070R90

0.38 \pm 0.07 \pm 0.07		49	EATON	84	MRK2 $e^+ e^-$
----------------------------	--	----	-------	----	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.11	90	PERUZZI	78	MRK1 $e^+ e^-$
---------	----	---------	----	----------------

 $\Gamma(\gamma p \bar{p} \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{351}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070R93
 NODE=M070R93

$<0.79 \times 10^{-3}$	90	EATON	84	MRK2 $e^+ e^-$
------------------------	----	-------	----	----------------

 $\Gamma(\gamma \Lambda \bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{352}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M070S8
 NODE=M070S8

$<0.13 \times 10^{-3}$	90	HENRARD	87	DM2 $e^+ e^-$
------------------------	----	---------	----	---------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.16 \times 10^{-3}$	90	BAI	98G	BES $e^+ e^-$
------------------------	----	-----	-----	---------------

$\Gamma(\gamma A^0 \rightarrow \gamma \text{invisible})/\Gamma_{\text{total}}$ Γ_{353}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

$<1.7 \times 10^{-6}$	90	88M	¹ ABLIKIM	20K BES3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
-----------------------	----	-----	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.3 \times 10^{-6}$	90	3.7M	² INSLEER	10 CLEO	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
-----------------------	----	------	----------------------	---------	---

¹ For a narrow state, A^0 , with mass $m_{A^0} < 1.2$ GeV. The limit varies with m_{A^0} , reaching its largest value of 1.7×10^{-6} at 1.2 GeV and being 7.0×10^{-7} for $m_{A^0} = 0$.

² The limit varies with mass m_{A^0} of a narrow state A^0 and is 4.3×10^{-6} for $m_{A^0} = 0$, reaches its largest value of 6.3×10^{-6} at $m_{A^0} = 500$ MeV, and is 3.6×10^{-6} at $m_{A^0} = 960$ MeV.

NODE=M070S68
NODE=M070S68

NODE=M070S68;LINKAGE=A

NODE=M070S68;LINKAGE=IN

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{354}/Γ (narrow state A^0 with $0.2 \text{ GeV} < m_{A^0} < 3 \text{ GeV}$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$<7.8 \times 10^{-7}$	90	¹ ABLIKIM	22H BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
-----------------------	----	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.5 \times 10^{-5}$	90	² ABLIKIM	16E BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
-----------------------	----	----------------------	----------	---

$<2.1 \times 10^{-5}$	90	³ ABLIKIM	12 BES3	$J/\psi \rightarrow \gamma \mu^+ \mu^-$
-----------------------	----	----------------------	---------	---

¹ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3.0 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(1.2\text{--}778.0) \times 10^{-9}$.

² For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.212–3 GeV. The measured 90% CL limit as a function of m_{A^0} is in the range $(2.8\text{--}495.3) \times 10^{-8}$.

³ For a narrow scalar or pseudoscalar, A^0 , with a mass in the range 0.21–3.00 GeV. The measured 90% CL limit as a function of m_{A^0} ranges from 4×10^{-7} to 2.1×10^{-5} .

NODE=M070S76
NODE=M070S76

NODE=M070S76;LINKAGE=B

NODE=M070S76;LINKAGE=A

NODE=M070S76;LINKAGE=AB

———— DALITZ DECAYS ————

 $\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{355}/Γ

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$7.56 \pm 1.32 \pm 0.50$	39	ABLIKIM	14I BES3	$J/\psi \rightarrow \pi^0 e^+ e^-$
--------------------------	----	---------	----------	------------------------------------

NODE=M070S89
NODE=M070S89

 $\Gamma(\eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{356}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.42 \pm 0.04 \pm 0.07$	2.47k	^{1,2} ABLIKIM	19A BES3	$J/\psi \rightarrow \eta e^+ e^-$
--------------------------	-------	------------------------	----------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.16 \pm 0.07 \pm 0.06$	320	¹ ABLIKIM	14I BES3	$J/\psi \rightarrow \eta e^+ e^-$
--------------------------	-----	----------------------	----------	-----------------------------------

¹ Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

² Approximation of the transition form factor squared as an incoherent sum of the ρ -meson and one-pole non-resonant amplitudes gives the pole mass $m(\Lambda) = 2.56 \pm 0.04 \pm 0.03$ GeV. Supersedes ABLIKIM 14I.

NODE=M070S90
NODE=M070S90

NODE=M070S90;LINKAGE=A
NODE=M070S90;LINKAGE=C

 $\Gamma(\eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{357}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$6.59 \pm 0.07 \pm 0.17$	8.9k	¹ ABLIKIM	19H BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$
--------------------------	------	----------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.81 \pm 0.16 \pm 0.31$	1.4k	^{1,2} ABLIKIM	14I BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$
--------------------------	------	------------------------	----------	---

¹ Using both $\eta' \rightarrow \gamma \pi^+ \pi^-$ and $\eta' \rightarrow \pi^+ \pi^- \eta$ decays.

² Superseded by ABLIKIM 19H.

NODE=M070S91
NODE=M070S91

NODE=M070S91;LINKAGE=A
NODE=M070S91;LINKAGE=B

 $\Gamma(X(1835) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{358}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$3.58 \pm 0.19 \pm 0.16$	1364	¹ ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$
--------------------------	------	----------------------	----------	--

¹ Assuming constructive interference. Destructive interference gives a value of $(4.43 \pm 0.23 \pm 0.19) \times 10^{-6}$ for this branching fraction.

NODE=M070Q22
NODE=M070Q22

NODE=M070Q22;LINKAGE=A

 $\Gamma(X(2120) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{359}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$0.82 \pm 0.12 \pm 0.06$	310	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$
--------------------------	-----	---------	----------	--

NODE=M070Q24
NODE=M070Q24

 $\Gamma(X(2370) e^+ e^-, X \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}$ Γ_{360}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

$1.08 \pm 0.14 \pm 0.10$	397	ABLIKIM	22B BES3	$J/\psi \rightarrow \pi^+ \pi^- \eta' e^+ e^-$
--------------------------	-----	---------	----------	--

NODE=M070Q23
NODE=M070Q23

$\Gamma(\eta U \rightarrow \eta e^+ e^-)/\Gamma_{\text{total}}$ Γ_{361}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.11 \times 10^{-7}$	90	¹ ABLIKIM 19A	BES3	$J/\psi \rightarrow \eta e^+ e^-$

NODE=M070P42
NODE=M070P42

¹ For a dark photon U with mass between 10 and 2400 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.9×10^{-8} to 91.1×10^{-8} .

NODE=M070P42;LINKAGE=A

 $\Gamma(\eta'(958) U \rightarrow \eta'(958) e^+ e^-)/\Gamma_{\text{total}}$ Γ_{362}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-7}$	90	¹ ABLIKIM 19H	BES3	$J/\psi \rightarrow \eta'(958) e^+ e^-$

NODE=M070P61
NODE=M070P61

¹ For a dark photon U with mass between 100 and 2100 MeV. Obtained 90% C.L. limits as a function of m_U range from 1.8×10^{-8} to 2.0×10^{-7} . The corresponding limits on the branching fraction $J/\psi \rightarrow \eta' U$ range from 5.7×10^{-8} to 7.4×10^{-7} .

NODE=M070P61;LINKAGE=A

 $\Gamma(\phi e^+ e^-)/\Gamma_{\text{total}}$ Γ_{363}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ ABLIKIM 19AB	BES3	$J/\psi \rightarrow \phi e^+ e^-$

NODE=M070P82
NODE=M070P82

¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) = (34.49 \pm 0.30)\%$.

NODE=M070P82;LINKAGE=A

———— WEAK DECAYS ————

 $\Gamma(D^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{364}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.1 \times 10^{-8}$	90	ABLIKIM 21Q	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S53
NODE=M070S53

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.2 \times 10^{-5}$	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	-------------	------	------------------------------

 $\Gamma(\bar{D}^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{365}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.5 \times 10^{-8}$	90	¹ ABLIKIM 17AF	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S54
NODE=M070S54

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1.1 \times 10^{-5}$	90	ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	-------------	------	------------------------------

¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.

NODE=M070S54;LINKAGE=A

 $\Gamma(D_s^- e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{366}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S55
NODE=M070S55

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<3.6 \times 10^{-5}$	90	¹ ABLIKIM 06M	BES2	$e^+ e^- \rightarrow J/\psi$
-----------------------	----	--------------------------	------	------------------------------

¹ Using $B(D_s^- \rightarrow \phi \pi^-) = 4.4 \pm 0.5\%$.

NODE=M070S55;LINKAGE=AB

 $\Gamma(D_s^{*-} e^+ \nu_e + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{367}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	ABLIKIM 14R	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070B13
NODE=M070B13

 $\Gamma(D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{368}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.5 \times 10^{-5}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S61
NODE=M070S61

 $\Gamma(\bar{D}^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{369}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S62
NODE=M070S62

 $\Gamma(\bar{D}^0 \bar{K}^{*0} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{370}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-6}$	90	ABLIKIM 14K	BES3	$e^+ e^- \rightarrow J/\psi$

NODE=M070S93
NODE=M070S93

 $\Gamma(D_s^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{371}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90	ABLIKIM 08J	BES2	$e^+ e^- \rightarrow J/\psi$

NODE=M070S63
NODE=M070S63

$\Gamma(D_s^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{372}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-5}$	90	ABLIKIM	14K BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S92 NODE=M070S92

———— CHARGE CONJUGATION (C), PARITY (P), ————
 ———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_{373}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$< 2.7 \times 10^{-7}$	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	NODE=M070R80 NODE=M070R80
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$< 0.5 \times 10^{-5}$	90	ADAMS	08 CLEO	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	
$< 1.6 \times 10^{-4}$	90	¹ WICHT	08 BELL	$B^\pm \rightarrow K^\pm \gamma\gamma$	
$< 2.2 \times 10^{-5}$	90	ABLIKIM	07J BES2	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$	
$< 50 \times 10^{-5}$	90	BARTEL	77 CNTR	$e^+ e^-$	

¹ WICHT 08 reports $[\Gamma(J/\psi(1S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow J/\psi(1S) K^+)] < 0.16 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow J/\psi(1S) K^+) = 1.020 \times 10^{-3}$.

NODE=M070R80;LINKAGE=WI

$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$					Γ_{374}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-6}$	90	ABLIKIM	14Q BES3	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	NODE=M070S95 NODE=M070S95

$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$					Γ_{375}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-7}$	90	ABLIKIM	13L BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S39 NODE=M070S39
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<1.1 \times 10^{-6}$	90	BAI	03D BES	$e^+ e^- \rightarrow J/\psi$	

$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$					Γ_{376}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<7.5 \times 10^{-8}$	90	ABLIKIM	21M BES3	$e^+ e^- \rightarrow J/\psi$	NODE=M070S40 NODE=M070S40
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$<8.3 \times 10^{-6}$	90	¹ ABLIKIM	04 BES	$e^+ e^- \rightarrow J/\psi$	
¹ Superseded by ABLIKIM 21M.					

NODE=M070S40;LINKAGE=A

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$					Γ_{377}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.0 \times 10^{-6}$	90	ABLIKIM	04 BES	$e^+ e^- \rightarrow J/\psi$	NODE=M070S41 NODE=M070S41

$\Gamma(\Lambda_c^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{378}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.9 \times 10^{-8}$	90	ABLIKIM	19AF BES3	$e^+ e^- \rightarrow J/\psi \rightarrow p K^- \pi^+ e^- (+ \text{c.c.})$	NODE=M070P74 NODE=M070P74

———— OTHER DECAYS ————

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$					Γ_{379}/Γ_5
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<6.6 \times 10^{-2}$	90	LEES	13I BABR	$B \rightarrow K^{(*)} J/\psi$	NODE=M070S80 NODE=M070S80

$\Gamma(\text{invisible})/\Gamma(\mu^+ \mu^-)$					Γ_{379}/Γ_7
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.2 \times 10^{-2}$	90	ABLIKIM	08G BES2	$\psi(2S) \rightarrow \pi^+ \pi^- J/\psi$	NODE=M070S60 NODE=M070S60

J/ψ(1S) REFERENCES

DOCUMENT ID	TECN	COMMENT	REFID
ABLIKIM 23BD	PR D108 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23BU	PR D108 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 23S	PR D107 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ANASHIN 23	JHEP 2306 196	V.V. Anashin <i>et al.</i>	(KEDR Collab.)
GONG 23	PR D107 072008	G. Gong <i>et al.</i>	(BELLE Collab.)
LEES 23	PR D107 072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LIAO 23	PR D107 112007	L. Liao <i>et al.</i>	
ZHU 23	PR D107 012006	W. Zhu <i>et al.</i>	(BELLE Collab.)
ABLIKIM 22AI	PRL 129 192002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 22AS	PR D106 072012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
Also	PR D107 079901 (err.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 22B	PRL 129 022002	M. Ablikim <i>et al.</i>	(BESIII Collab.)

NODE=M070

REFID=62425
REFID=62517
REFID=62063
REFID=62057
REFID=62073
REFID=62061
REFID=62279
REFID=61911
REFID=61881
REFID=61891
REFID=61891
REFID=62033
REFID=61902
REFID=61636

ABLIKIM	22C	PR D105 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61637
ABLIKIM	22H	PR D105 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61643
ANASHIN	22	EPJ C82 938	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=61894
ABLIKIM	21AM	PR D104 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61445
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61463
ABLIKIM	21C	PR D103 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61030
ABLIKIM	21M	PR D103 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61118
ABLIKIM	21Q	JHEP 2106 157	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61125
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61113
LEES	21B	PR D104 112003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61450
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61451
SARANTSEV	21	PL B816 136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)	REFID=61091
ABLIKIM	20	PR D101 012004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60048
ABLIKIM	20K	PR D101 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60316
ABLIKIM	20Q	EPJ C80 746	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60457
ANASHIN	20	JHEP 2007 112	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=60512
ABLIKIM	19A	PR D99 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59517
Also		PR D104 099901 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61449
ABLIKIM	19AB	PR D99 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59846
ABLIKIM	19AC	PR D99 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59850
ABLIKIM	19AF	PR D99 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59857
ABLIKIM	19AN	PR D99 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59890
ABLIKIM	19AQ	PR D100 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59909
ABLIKIM	19H	PR D99 012013	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59604
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59615
ABLIKIM	19Q	PL B791 375	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59747
ABLIKIM	19T	PRL 122 142002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59773
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)	REFID=59614
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59455
ABLIKIM	18AB	PR D98 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59456
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58849
ABLIKIM	18I	PR D97 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58893
ABLIKIM	18O	PR D97 072014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58925
ANASHIN	18A	JHEP 1805 119	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=59102
LEES	18	PR D97 052007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58900
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58315
ABLIKIM	17AH	PR D96 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58317
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58324
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57903
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57967
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57966
LEES	17C	PR D95 072007	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57981
LEES	17D	PR D95 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57990
ABLIKIM	16E	PR D93 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57265
ABLIKIM	16J	PRL 117 042002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57454
ABLIKIM	16K	PR D93 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57509
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57510
ABLIKIM	16M	PR D93 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57511
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)	REFID=57512
ABLIKIM	16P	PR D94 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57522
ABLIKIM	16Q	PL B761 98	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57566
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
AAIJ	15BI	EPJ C75 311	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57147
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56984
ABLIKIM	15H	PR D91 052017	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56773
ABLIKIM	15K	PR D91 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56776
ABLIKIM	15P	PR D92 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56781
ABLIKIM	15T	PRL 115 091803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56785
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14I	PR D89 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55900
ABLIKIM	14K	PR D89 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55902
ABLIKIM	14N	PR D90 052009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55905
ABLIKIM	14Q	PR D90 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56238
ABLIKIM	14R	PR D90 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56388
ANASHIN	14	PL B738 391	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56130
AULCHENKO	14	PL B731 227	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=55655
LEES	14H	PR D89 092002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55940
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54920
ABLIKIM	13I	PR D87 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54954
ABLIKIM	13J	PR D87 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54955
ABLIKIM	13L	PR D87 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55300
ABLIKIM	13N	PR D87 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55387
ABLIKIM	13P	PR D87 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55392
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55402
ABLIKIM	13U	PR D88 091502	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55582
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54265
ABLIKIM	12B	PR D86 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54267
ABLIKIM	12C	PR D86 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54268
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54269
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54273
ABLIKIM	12P	CP C36 1031	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54863
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
ABLIKIM	11	PR D83 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53646
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16715
ABLIKIM	10C	PL B685 27	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53349
ABLIKIM	10E	PL B693 88	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53361
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
ANASHIN	10	PL B685 134	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53220
DEL-AMO-SA...	10O	PRL 105 172001	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53533
INSLER	10	PR D81 091101	J. Insler <i>et al.</i>	(CLEO Collab.)	REFID=53359

ABLIKIM	09	PL B676 25	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52718
ABLIKIM	09B	PR D80 052004	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53099
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
SHEN	09	PR D80 031101	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53000
ABLIKIM	08	EPJ C53 15	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52047
ABLIKIM	08A	PR D77 012001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52128
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
ABLIKIM	08E	PR D77 032005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52143
ABLIKIM	08F	PRL 100 102003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52154
ABLIKIM	08G	PRL 100 192001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52253
ABLIKIM	08I	PL B662 330	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52255
ABLIKIM	08J	PL B663 297	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52256
ABLIKIM	08O	PR D78 092005	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52571
ADAMS	08	PRL 101 101801	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=52261
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
BESSON	08	PR D78 032012	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=52685
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ABLIKIM	07J	PR D76 117101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52072
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
Also		PR D77 119902E (err.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
ABLIKIM	06	PL B632 181	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50986
ABLIKIM	06C	PL B633 681	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51037
ABLIKIM	06E	PR D73 052008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51057
ABLIKIM	06F	PR D73 052007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51058
ABLIKIM	06H	PR D73 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51125
ABLIKIM	06J	PRL 96 162002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51127
ABLIKIM	06K	PRL 97 062001	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51128
ABLIKIM	06M	PL B639 418	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51130
ABLIKIM	06V	PL B642 441	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51507
ADAMS	06A	PR D73 051103	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51036
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
WU	06	PRL 97 162003	C.-H. Wu <i>et al.</i>	(BELLE Collab.)	REFID=51472
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50450
ABLIKIM	05B	PR D71 032003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50496
ABLIKIM	05C	PL B610 192	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50507
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05R	PRL 95 262001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50985
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
LI	05C	PR D71 111103	Z. Li <i>et al.</i>	(CLEO Collab.)	REFID=50802
SIBIRTSEV	05A	PR D71 054010	A. Sibirtsev, J. Haidenbauer		REFID=51038
ABLIKIM	04	PL B598 172	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49739
ABLIKIM	04M	PR D70 112008	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50329
AUBERT	04	PR D69 011103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49611
AUBERT,B	04N	PR D70 072004	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50184
BAI	04A	PR D69 012003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49607
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04E	PL B591 42	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49751
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04H	PR D70 012005	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49754
BAI	04J	PL B594 47	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50167
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03D	PL B561 49	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49403
BAI	03F	PRL 91 022001	J.Z. Bai <i>et al.</i>	(BES II Collab.)	REFID=49473
BAI	03G	PR D68 052003	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49580
HUANG	03	PRL 91 241802	H.-C. Huang <i>et al.</i>	(BELLE Collab.)	REFID=49621
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	00B	PL B472 200	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47427
BAI	00D	PL B476 25	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47954
BAI	99	PL B446 356	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46606
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98G	PL B424 213	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46341
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46342
BALDINI	98	PL B444 111	R. Baldini <i>et al.</i>	(FENICE Collab.)	REFID=46608
ARMSTRONG	96	PR D54 7067	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45146
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44736
BAI	96C	PRL 77 3959	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45169
BAI	96D	PR D54 1221	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=45198
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)	REFID=45197
BAI	95B	PL B355 374	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=44434
BUGG	95	PL B353 378	D.V. Bugg <i>et al.</i>	(LOQM, PNPI, WASH)	REFID=44438
ANTONELLI	93	PL B301 317	A. Antonelli <i>et al.</i>	(FENICE Collab.)	REFID=43314
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)	REFID=43601
AUGUSTIN	92	PR D46 1951	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41584
BOLTON	92	PL B278 495	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42175
BOLTON	92B	PRL 69 1328	T. Bolton <i>et al.</i>	(Mark III Collab.)	REFID=42176
COFFMAN	92	PRL 68 282	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41866
HSUEH	92	PR D45 2181	S. Hsueh, S. Palestini	(FNAL, TORI)	REFID=41899
AUGUSTIN	90	PR D42 10	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=41352

BAI	90B	PRL 65 1309	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41354
BAI	90C	PRL 65 2507	Z. Bai <i>et al.</i>	(Mark III Collab.)	REFID=41578
BISELLO	90	PL B241 617	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=41359
COFFMAN	90	PR D41 1410	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=41350
JOUSSET	90	PR D41 1389	J. Jousset <i>et al.</i>	(DM2 Collab.)	REFID=41349
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
AUGUSTIN	89	NP B320 1	J.E. Augustin, G. Cosme	(DM2 Collab.)	REFID=41004
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
AUGUSTIN	88	PRL 60 2238	J.E. Augustin <i>et al.</i>	(DM2 Collab.)	REFID=40574
COFFMAN	88	PR D38 2695	D.M. Coffman <i>et al.</i>	(Mark III Collab.)	REFID=40346
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40576
AUGUSTIN	87	ZPHY C36 369	J.E. Augustin <i>et al.</i>	(LALO, CLER, FRAS+)	REFID=40268
BAGLIN	87	NP B286 592	C. Baglin <i>et al.</i>	(LAPP, CERN, GENO, LYON+)	REFID=40002
BALTRUSAIT...	87	PR D35 2077	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=40010
BECKER	87	PR 59 186	J.J. Becker <i>et al.</i>	(Mark III Collab.)	REFID=40015
BISELLO	87	PL B192 239	D. Bisello <i>et al.</i>	(PADO, CLER, FRAS+)	REFID=40012
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
HENRARD	87	NP B292 670	P. Henrard <i>et al.</i>	(CLER, FRAS, LALO+)	REFID=40261
PALLIN	87	NP B292 653	D. Pallin <i>et al.</i>	(CLER, FRAS, LALO, PADO)	REFID=40243
BALTRUSAIT... 86B	86B	PR D33 1222	R.M. Baltrusaitis <i>et al.</i>	(Mark III Collab.)	REFID=22100
BALTRUSAIT... 86D	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)	REFID=21865
BISELLO	86B	PL B179 294	D. Bisello <i>et al.</i>	(DM2 Collab.)	REFID=22101
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
BALTRUSAIT... 85C	85C	PRL 55 1723	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22095
BALTRUSAIT... 85D	85D	PR D32 566	R.M. Baltrusaitis <i>et al.</i>	(CIT, UCSC+)	REFID=22097
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
EATON	84	PR D29 804	M.W. Eaton <i>et al.</i>	(LBL, SLAC)	REFID=22092
BLOOM	83	ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS	83B	PRL 51 859	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21318
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
BURKE	82	PRL 49 632	D.L. Burke <i>et al.</i>	(LBL, SLAC)	REFID=21676
EDWARDS	82B	PR D25 3065	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22080
EDWARDS	82D	PRL 48 458	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21677
Also		ARNS 33 143	E.D. Bloom, C. Peck	(SLAC, CIT)	REFID=21682
EDWARDS	82E	PRL 49 259	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=21314
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
BESCH	81	ZPHY C8 1	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)	REFID=22077
GIDAL	81	PL 107B 153	G. Gidal <i>et al.</i>	(SLAC, LBL)	REFID=20386
PARTRIDGE	80	PRL 44 712	R. Partridge <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22073
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
Translated from YAF 34 1471.					
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
ALEXANDER	78	PL 72B 493	G. Alexander <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22065
BESCH	78	PL 78B 347	H.J. Besch <i>et al.</i>	(BONN, DESY, MAINZ)	REFID=22066
BRANDELIK	78B	PL 74B 292	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22067
PERUZZI	78	PR D17 2901	I. Peruzzi <i>et al.</i>	(SLAC, LBL)	REFID=22068
BARTEL	77	PL 66B 489	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22058
BURMESTER	77D	PL 72B 135	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22060
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
VANNUCCI	77	PR D15 1814	F. Vannucci <i>et al.</i>	(SLAC, LBL)	REFID=22063
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
BRAUNSCH...	76	PL 63B 487	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22054
JEAN-MARIE	76	PRL 36 291	B. Jean-Marie <i>et al.</i>	(SLAC, LBL) IG	REFID=22056
BALDINI-...	75	PL 58B 471	R. Baldini-Celio <i>et al.</i>	(FRAS, ROMA)	REFID=22026
BOYARSKI	75	PRL 34 1357	A.M. Boyarski <i>et al.</i>	(SLAC, LBL) JPC	REFID=22030
DASP	75	PL 56B 491	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22036
ESPOSITO	75B	LNC 14 73	B. Esposito <i>et al.</i>	(FRAS, NAPL, PADO+)	REFID=22038
FORD	75	PRL 34 604	R.L. Ford <i>et al.</i>	(SLAC, PENN)	REFID=22039

BRANCHING RATIOS OF $\psi(2S)$, $\chi_{c0,1,2}$ AND $\eta_C(1S)$

Updated March 2024 by J.J. Hernández-Rey (IFIC, Valencia), R. Mitchell (Indiana U.), S. Navas (U. of Granada), and C. Patrignani (Bologna Univ., INFN)

Since 2002, the treatment of the branching ratios of the $\psi(2S)$ and $\chi_{c0,1,2}$ has undergone an important restructuring. Since the 2023 online edition, the $\eta_c(1S)$ is now treated in a similar way (see the end of this note).

When measuring a branching ratio experimentally, it is not always possible to normalize the number of events observed in the corresponding decay mode to the total number of particles produced. Therefore, the experimenters sometimes report the number of observed decays with respect to another decay mode of the same or another particle in the relevant decay chain. This is actually equivalent to measuring combinations of branching fractions of several decay modes.

To extract the branching ratio of a given decay mode, the collaborations use some previously reported measurements of the required branching ratios. However, the values are frequently taken from the *Review of Particle Physics* (RPP), which in turn uses the branching ratio reported by the experiment in the following edition, giving rise either to correlations or to plain vicious circles, as discussed in more detail in earlier editions of this review [1,2].

The way to avoid these dependencies and correlations is to extract the branching ratios through a fit that uses the truly measured combinations of branching fractions and partial widths. This fit, in fact, should involve decays from the four concerned particles, $\psi(2S)$, χ_{c0} , χ_{c1} , and χ_{c2} , and occasionally some combinations of branching ratios of more than one of them. This is what is done since the 2002 edition [3].

The PDG policy is to quote the results of the collaborations in a manner as close as possible to what appears in their original publications. However, in order to avoid the problems mentioned above, we had in some cases to work out the values originally measured, using the number of events and detection efficiencies given by the collaborations, or rescaling back the published results. The information was sometimes spread over several articles, and some articles referred to papers still unpublished, which in turn contained the relevant numbers in footnotes.

Even though the experimental collaborations are entitled to extract whatever branching ratios they consider appropriate by

using other published results, we would like to encourage them to also quote explicitly in their articles the actual quantities measured, so that they can be used directly in averages and fits of different experimental determinations.

To inform the reader how we computed some of the values used in this edition of RPP, we use footnotes to indicate the branching ratios actually given by the experiments and the quantities they use to derive them from the true combination of branching ratios actually measured.

None of the branching ratios of the $\chi_{c0,1,2}$ are measured independently of the $\psi(2S)$ radiative decays. We tried to identify those branching ratios which can be correlated in a non-trivial way, and although we cannot preclude the existence of other cases, we are confident that the most relevant correlations have already been removed. Nevertheless, correlations in the errors of different quantities measured by the same experiment have not been taken into account.

The $\eta_c(1S)$ can be produced in $\gamma\gamma$ collisions, in the radiative decays of vector and axial-vector charmonia (J/ψ , $\psi(2S)$ and $h_c(1P)$) or in B meson decays. As for the $\psi(2S)$ and $\chi_{c0,1,2}$, correlations can be introduced if the derived branching ratios are not properly extracted. We now obtain the corresponding branching ratios using the products of branching ratios originally measured by the experiments, sometimes among different particles (cross-particle branching ratios) and performing an overall fit. In some cases, we were obliged to infer the original product from the information given by the collaborations. However, for recent experiments our policy is not to use anymore extracted branching ratios, so that if the original measured quantity is missing in the relevant publication, the value will not be used in our averages and fits.

1. Fit Information for the $\psi(2S)$ and $\chi_{c0,1,2}$

This is an overall fit to 4 total widths, 1 partial width, 25 combinations of partial widths, 10 branching fractions, and 76 branching ratios or combinations thereof. Of the latter, 58 involve decays of more than one particle.

The overall fit uses 253 measurements to determine 49 parameters and gives a χ^2 of 389.6 for 204 degrees of freedom. We have applied scaling factors in the final errors of the fitted values following the procedure described in section 5.2.3 of our Introduction. After rescaling, the total χ^2 is 149.6.

In the listing we provide the inter-particle correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the corresponding parameter x_i .

2. Fit Information for the $\eta_c(1S)$

This is an overall fit to 1 total width, 10 combinations of partial widths, 6 branching fractions and 32 branching ratios or combinations thereof. Of the latter, 31 involve decays of more than one particle.

The overall fit uses 113 measurements to determine 19 parameters and has a χ^2 of 184.6 for 94 degrees of freedom. We have applied scaling factors in the final errors of the fitted values following the procedure described in section 5.2.3 of our Introduction. After rescaling, the total χ^2 is 74.1.

In the listing we provide the inter-particle correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the corresponding parameter x_i .

References

1. Y.F. Gu and X.H. Li, Phys. Lett. **B449**, 361 (1999).
2. C. Patrignani, Phys. Rev. **D64**, 034017 (2001).
3. K. Hagiwara *et al.* (Particle Data Group), Phys. Rev. **D68**, 010001 (2002).

$\chi_{c0}(1P)$	$I^G(J^{PC}) = 0^+(0^{++})$	NODE=M056
-----------------	-----------------------------	-----------

$\chi_{c0}(1P)$ MASS					NODE=M056M
<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M056M
3414.71 ± 0.30 OUR AVERAGE					
3413.0 ± 1.9 ± 0.6	933	¹ AAIJ	17BB LHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+K^-)X$	
3414.2 ± 0.5 ± 2.3	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow \text{hadrons}$	
3406 ± 7 ± 6	230	² ABE	07 BELL	$e^+e^- \rightarrow J/\psi(c\bar{c})$	
3414.21 ± 0.39 ± 0.27		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$	
3414.7 ⁺ 0.7 _{-0.6} ± 0.2		³ ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$	
3415.5 ± 0.4 ± 0.4	392	⁴ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$	
3417.4 ⁺ 1.8 _{-1.9} ± 0.2		³ AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$	
3414.1 ± 0.6 ± 0.8		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$	
3417.8 ± 0.4 ± 4		³ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$	
3416 ± 3 ± 4		⁵ TANENBAUM	78 MRK1	e^+e^-	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3414.6 ± 1.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
3416.5 ± 3.0		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^-\chi_{c0}$	
3422 ± 10		⁵ BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$	
3415 ± 9		⁵ BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$	

- ¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c0}(1P)$ fixed to the PDG 16 value.
² From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.
³ Using mass of $\psi(2S) = 3686.0$ MeV.
⁴ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.
⁵ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

NODE=M056M;LINKAGE=A

NODE=M056M;LINKAGE=EB

NODE=M056M;LINKAGE=C

NODE=M056M;LINKAGE=NW

NODE=M056M;LINKAGE=D

 $\chi_{c0}(1P)$ WIDTH

NODE=M056W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10.7±0.6 OUR FIT	Error includes scale factor of 1.1.		[10.8 ± 0.6 MeV OUR 2023 FIT]	
10.5±0.8 OUR AVERAGE	Error includes scale factor of 1.1.			
10.6±1.9±2.6	5.4k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow$ hadrons
12.6 ^{+1.5+0.9} _{-1.6-1.1}		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
8.6 ^{+1.7} _{-1.3} ±0.1		ANDREOTTI	03 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$
9.7±1.0	392	¹ BAGNASCO	02 E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
16.6 ^{+5.2} _{-3.7} ±0.1		AMBROGIANI	99B E835	$\bar{p}p \rightarrow e^+e^-\gamma$
14.3±2.0±3.0		BAI	981 BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
13.5±3.3±4.2		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X, \gamma\pi^0\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
13.2±2.1	266	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M056W

NEW

¹ Recalculated by ANDREOTTI 05A.

NODE=M056W;LINKAGE=AN

 $\chi_{c0}(1P)$ DECAY MODES

NODE=M056215;NODE=M056

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	(2.3 ± 0.4) %	S=2.0
Γ_2 $\rho^0\pi^+\pi^-$	(9.1 ± 3.1) × 10 ⁻³	S=1.1
Γ_3 $\rho^0\rho^0$		
Γ_4 $f_0(980)f_0(980)$	(6.7 ± 2.1) × 10 ⁻⁴	
Γ_5 $\pi^+\pi^-\pi^0\pi^0$	(3.3 ± 0.4) %	
Γ_6 $\rho^+\pi^-\pi^0 + \text{c.c.}$	(2.9 ± 0.4) %	
Γ_7 $4\pi^0$	(3.3 ± 0.4) × 10 ⁻³	
Γ_8 $\pi^+\pi^-K^+K^-$	(1.82±0.16) %	S=1.2
Γ_9 $K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-$	(9.9 ^{+4.0} _{-2.8}) × 10 ⁻⁴	
Γ_{10} $K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	(8.0 ^{+2.0} _{-2.4}) × 10 ⁻⁴	
Γ_{11} $K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	(6.3 ± 1.9) × 10 ⁻³	
Γ_{12} $K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-$	< 2.7 × 10 ⁻³	CL=90%
Γ_{13} $f_0(980)f_0(980)$	(1.6 ^{+1.0} _{-0.9}) × 10 ⁻⁴	
Γ_{14} $f_0(980)f_0(2200)$	(7.9 ^{+2.0} _{-2.5}) × 10 ⁻⁴	
Γ_{15} $f_0(1370)f_0(1370)$	< 2.7 × 10 ⁻⁴	CL=90%
Γ_{16} $f_0(1370)f_0(1500)$	< 1.7 × 10 ⁻⁴	CL=90%
Γ_{17} $f_0(1370)f_0(1710)$	(6.7 ^{+3.5} _{-2.3}) × 10 ⁻⁴	
Γ_{18} $f_0(1500)f_0(1370)$	< 1.3 × 10 ⁻⁴	CL=90%
Γ_{19} $f_0(1500)f_0(1500)$	< 5 × 10 ⁻⁵	CL=90%
Γ_{20} $f_0(1500)f_0(1710)$	< 7 × 10 ⁻⁵	CL=90%
Γ_{21} $K^+K^-\pi^+\pi^-\pi^0$	(8.6 ± 0.9) × 10 ⁻³	
Γ_{22} $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	(4.2 ± 0.4) × 10 ⁻³	
Γ_{23} $K^+K^-\pi^0\pi^0$	(5.6 ± 0.9) × 10 ⁻³	
Γ_{24} $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(2.49±0.33) %	
Γ_{25} $\rho^+K^-K^0 + \text{c.c.}$	(1.21±0.21) %	

NODE=M056;CLUMP=A

DESIG=3

DESIG=9

DESIG=54

DESIG=20

DESIG=61

DESIG=62

DESIG=70

DESIG=5

DESIG=31

DESIG=32

DESIG=33

DESIG=34

DESIG=23

DESIG=24

DESIG=25

DESIG=26

DESIG=27

DESIG=28

DESIG=29

DESIG=30

DESIG=75

DESIG=87

DESIG=63

DESIG=65

DESIG=66

Г26	$K^*(892)^- K^+ \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	$(4.6 \pm 1.2) \times 10^{-3}$		DESIG=67
Г27	$K_S^0 K_S^0 \pi^+ \pi^-$	$(5.7 \pm 1.1) \times 10^{-3}$		DESIG=41
Г28	$K^+ K^- \eta \pi^0$	$(3.0 \pm 0.7) \times 10^{-3}$		DESIG=68
Г29	$3(\pi^+ \pi^-)$	$(1.95 \pm 0.22) \%$	S=3.3	DESIG=4
Г30	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(7.5 \pm 1.6) \times 10^{-3}$		DESIG=10
Г31	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.7 \pm 0.6) \times 10^{-3}$		DESIG=21
Г32	$\pi \pi$	$(8.5 \pm 0.4) \times 10^{-3}$	S=1.2	DESIG=18
Г33	$\pi^0 \eta$	$< 1.8 \times 10^{-4}$		DESIG=35
Г34	$\pi^0 \eta'$	$< 1.1 \times 10^{-3}$		DESIG=36
Г35	$\pi^0 \eta_c$	$< 1.6 \times 10^{-3}$	CL=90%	DESIG=86
Г36	$\eta \eta$	$(3.01 \pm 0.25) \times 10^{-3}$	S=1.3	DESIG=13
Г37	$\eta \eta'$	$(9.1 \pm 1.1) \times 10^{-5}$		DESIG=37
Г38	$\eta' \eta'$	$(2.17 \pm 0.12) \times 10^{-3}$		DESIG=46
Г39	$\omega \omega$	$(9.7 \pm 1.1) \times 10^{-4}$		DESIG=22
Г40	$\omega \phi$	$(1.42 \pm 0.13) \times 10^{-4}$		DESIG=76
Г41	$\omega K^+ K^-$	$(1.94 \pm 0.21) \times 10^{-3}$		DESIG=88
Г42	$K^+ K^-$	$(6.07 \pm 0.33) \times 10^{-3}$	S=1.1	DESIG=2
Г43	$K_S^0 K_S^0$	$(3.17 \pm 0.19) \times 10^{-3}$	S=1.1	DESIG=15
Г44	$\pi^+ \pi^- \eta$	$< 2.0 \times 10^{-4}$	CL=90%	DESIG=50
Г45	$\pi^+ \pi^- \eta'$	$< 4 \times 10^{-4}$	CL=90%	DESIG=53
Г46	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$< 9 \times 10^{-5}$	CL=90%	DESIG=17
Г47	$K^+ K^- \pi^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=47
Г48	$K^+ K^- \eta$	$< 2.3 \times 10^{-4}$	CL=90%	DESIG=51
Г49	$K^+ K^- K_S^0 K_S^0$	$(1.4 \pm 0.5) \times 10^{-3}$		DESIG=42
Г50	$K_S^0 K_S^0 K_S^0 K_S^0$	$(5.8 \pm 0.5) \times 10^{-4}$		DESIG=94
Г51	$K^+ K^- K^+ K^-$	$(2.8 \pm 0.4) \times 10^{-3}$	S=1.5	DESIG=14
Г52	$K^+ K^- \phi$	$(9.7 \pm 2.5) \times 10^{-4}$		DESIG=44
Г53	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(3.7 \pm 0.6) \times 10^{-3}$		DESIG=91
Г54	$K^+ K^- \pi^0 \phi$	$(1.90 \pm 0.35) \times 10^{-3}$		DESIG=92
Г55	$\phi \pi^+ \pi^- \pi^0$	$(1.18 \pm 0.15) \times 10^{-3}$		DESIG=89
Г56	$\phi \phi$	$(8.48 \pm 0.31) \times 10^{-4}$		DESIG=16
Г57	$\phi \phi \eta$	$(8.4 \pm 1.0) \times 10^{-4}$		DESIG=96
Г58	$\rho \bar{\rho}$	$(2.21 \pm 0.14) \times 10^{-4}$	S=1.6	DESIG=11
Г59	$\rho \bar{\rho} \pi^0$	$(7.0 \pm 0.7) \times 10^{-4}$	S=1.3	DESIG=48
Г60	$\rho \bar{\rho} \eta$	$(3.5 \pm 0.4) \times 10^{-4}$		DESIG=52
Г61	$\rho \bar{\rho} \omega$	$(5.3 \pm 0.6) \times 10^{-4}$		DESIG=69
Г62	$\rho \bar{\rho} \phi$	$(6.0 \pm 1.4) \times 10^{-5}$		DESIG=74
Г63	$\rho \bar{\rho} \pi^+ \pi^-$	$(2.1 \pm 0.7) \times 10^{-3}$	S=1.4	DESIG=8
Г64	$\rho \bar{\rho} \pi^0 \pi^0$	$(1.04 \pm 0.28) \times 10^{-3}$		DESIG=64
Г65	$\rho \bar{\rho} K^+ K^- (\text{non-resonant})$	$(1.22 \pm 0.26) \times 10^{-4}$		DESIG=71
Г66	$\rho \bar{\rho} K_S^0 K_S^0$	$< 8.8 \times 10^{-4}$	CL=90%	DESIG=40
Г67	$\rho \bar{n} \pi^-$	$(1.27 \pm 0.11) \times 10^{-3}$		DESIG=43
Г68	$\bar{\rho} n \pi^+$	$(1.37 \pm 0.12) \times 10^{-3}$		DESIG=82
Г69	$\rho \bar{n} \pi^- \pi^0$	$(2.34 \pm 0.21) \times 10^{-3}$		DESIG=83
Г70	$\bar{\rho} n \pi^+ \pi^0$	$(2.21 \pm 0.19) \times 10^{-3}$		DESIG=84
Г71	$\Lambda \bar{\Lambda}$	$(3.60 \pm 0.17) \times 10^{-4}$	S=1.1	DESIG=19
Г72	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(1.18 \pm 0.13) \times 10^{-3}$		DESIG=38
Г73	$\Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant})$	$< 5 \times 10^{-4}$	CL=90%	DESIG=77
Г74	$\Lambda \bar{\Lambda} \eta$	$(2.3 \pm 0.4) \times 10^{-4}$		DESIG=102
Г75	$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	$< 5 \times 10^{-4}$	CL=90%	DESIG=78
Г76	$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	$< 5 \times 10^{-4}$	CL=90%	DESIG=79
Г77	$K^+ \bar{\rho} \Lambda + \text{c.c.}$	$(1.25 \pm 0.12) \times 10^{-3}$	S=1.3	DESIG=49
Г78	$n K_S^0 \bar{\Lambda} + \text{c.c.}$	$(6.7 \pm 0.5) \times 10^{-4}$		DESIG=101
Г79	$K^*(892)^+ \bar{\rho} \Lambda + \text{c.c.}$	$(4.8 \pm 0.9) \times 10^{-4}$		DESIG=98
Г80	$K^+ \bar{\rho} \Lambda(1520) + \text{c.c.}$	$(3.0 \pm 0.8) \times 10^{-4}$		DESIG=72
Г81	$\Lambda(1520) \bar{\Lambda}(1520)$	$(3.1 \pm 1.2) \times 10^{-4}$		DESIG=73
Г82	$\Sigma^0 \bar{\Sigma}^0$	$(4.69 \pm 0.32) \times 10^{-4}$		DESIG=58

Γ ₈₃	$\Sigma^+ \bar{p} K_S^0 + c.c.$	$(3.53 \pm 0.27) \times 10^{-4}$		DESIG=97
Γ ₈₄	$\Sigma^0 \bar{p} K^+ + c.c.$	$(3.04 \pm 0.20) \times 10^{-4}$		DESIG=100
Γ ₈₅	$\Sigma^+ \bar{\Sigma}^-$	$(4.7 \pm 0.8) \times 10^{-4}$	S=2.6	DESIG=59
Γ ₈₆	$\Sigma^- \bar{\Sigma}^+$	$(5.1 \pm 0.5) \times 10^{-4}$		DESIG=99
Γ ₈₇	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$(1.6 \pm 0.6) \times 10^{-4}$		DESIG=80
Γ ₈₈	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$(2.3 \pm 0.7) \times 10^{-4}$		DESIG=81
Γ ₈₉	$K^- \Lambda \bar{\Xi}^+ + c.c.$	$(1.95 \pm 0.35) \times 10^{-4}$		DESIG=85
Γ ₉₀	$\Xi^0 \bar{\Xi}^0$	$(4.5 \pm 0.5) \times 10^{-4}$	S=1.7	DESIG=60
Γ ₉₁	$\Xi^- \bar{\Xi}^+$	$(4.47 \pm 0.20) \times 10^{-4}$		DESIG=39
Γ ₉₂	$\Omega^- \bar{\Omega}^+$	$(3.5 \pm 0.6) \times 10^{-5}$		DESIG=103
Γ ₉₃	$\eta_c \pi^+ \pi^-$	$< 7 \times 10^{-4}$	CL=90%	DESIG=90

Radiative decays

Γ ₉₄	$\gamma J/\psi(1S)$	$(1.41 \pm 0.09) \%$	S=1.7	NODE=M056;CLUMP=B DESIG=6
Γ ₉₅	$\gamma \rho^0$	$< 9 \times 10^{-6}$	CL=90%	DESIG=55
Γ ₉₆	$\gamma \omega$	$< 8 \times 10^{-6}$	CL=90%	DESIG=56
Γ ₉₇	$\gamma \phi$	$< 6 \times 10^{-6}$	CL=90%	DESIG=57
Γ ₉₈	$\gamma \gamma$	$(2.04 \pm 0.10) \times 10^{-4}$	S=1.1	DESIG=7
Γ ₉₉	$e^+ e^- J/\psi(1S)$	$(1.34 \pm 0.30) \times 10^{-4}$		DESIG=93
Γ ₁₀₀	$\mu^+ \mu^- J/\psi(1S)$	$< 1.9 \times 10^{-5}$	CL=90%	DESIG=95

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 86 branching ratios uses 253 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 389.6$ for 204 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x ₂	47									
x ₈	8	4								
x ₃₀	5	2	33							
x ₃₂	2	1	8	3						
x ₃₆	1	0	4	1	14					
x ₄₂	3	1	8	3	21	12				
x ₄₃	2	1	7	3	20	12	18			
x ₅₁	4	2	5	2	6	4	6	6		
x ₅₆	2	1	3	1	4	3	4	4	2	
x ₅₈	0	0	2	1	1	1	6	6	2	1
x ₇₁	1	1	8	2	26	16	23	22	6	5
x ₉₄	0	0	2	0	9	5	6	5	2	1
x ₉₈	-10	-5	-4	-4	17	11	14	14	0	1
Γ	-22	-10	-22	-12	-15	-7	-16	-14	-12	-7
	x ₁	x ₂	x ₈	x ₃₀	x ₃₂	x ₃₆	x ₄₂	x ₄₃	x ₅₁	x ₅₆
x ₇₁	7									
x ₉₄	-37	7								
x ₉₈	4	20	8							
Γ	-3	-14	-4	-37						
	x ₅₈	x ₇₁	x ₉₄	x ₉₈						

$\chi_{c0}(1P)$ PARTIAL WIDTHS

———— $\chi_{c0}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$ ————

NODE=M056217

NODE=M056223

$$\chi_{c0}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$$

NODE=M056224

$$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
51 ± 8 OUR FIT					Error includes scale factor of 1.9. [52 ± 4 eV OUR 2023 FIT]
49 ± 10 OUR AVERAGE					Error includes scale factor of 1.8.
44.7 ± 3.6 ± 4.9		3.6k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+\pi^-)$
75 ± 13 ± 8			EISENSTEIN	01	CLE2 $e^+e^- \rightarrow e^+e^-\chi_{c0}$

NODE=M056G2
 NODE=M056G2
 NEW

$$\Gamma(\rho^0\rho^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<12		90	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+\pi^-)$

NODE=M056G07
 NODE=M056G07

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<12		90	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(\pi^+\pi^-)$

$$\Gamma(\pi^+\pi^-K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
40 ± 4 OUR FIT					Error includes scale factor of 1.1. [40.0 ± 3.5 eV OUR 2023 FIT]
38.8 ± 3.7 ± 4.7		1.7k	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M056G08
 NODE=M056G08
 NEW

$$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{21}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
26 ± 4 ± 4		1094	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-$

NODE=M056G01
 NODE=M056G01

$$\Gamma(K^+\bar{K}^*(892)^0\pi^- + c.c.) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
16 ± 4 OUR FIT					
16.7 ± 6.1 ± 3.0		495 ± 182	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M056G09
 NODE=M056G09

$$\Gamma(K^*(892)^0\bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<6		90	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M056G10
 NODE=M056G10

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<6		90	UEHARA	08	BELL $\gamma\gamma \rightarrow \chi_{c0} \rightarrow K^+K^-\pi^+\pi^-$

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
18.7 ± 1.5 OUR FIT					Error includes scale factor of 1.2. [18.8 ± 1.3 eV OUR 2023 FIT]
23 ± 5 OUR AVERAGE					
29.7 ^{+17.4} _{-12.0} ± 4.8		103 ⁺⁶⁰ ₋₄₂	1 UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
22.7 ± 3.2 ± 3.5		129 ± 18	2 NAKAZAWA	05	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

NODE=M056G3
 NODE=M056G3
 NEW

¹We multiplied the measurement by 3 to convert from $\pi^0\pi^0$ to $\pi\pi$. Interference with the continuum included.

²We have multiplied $\pi^+\pi^-$ measurement by 3/2 to obtain $\pi\pi$.

NODE=M056G3;LINKAGE=UE

NODE=M056G;LINKAGE=NA

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.4 ± 2.3 ± 1.2		22	1 UEHARA	10A	BELL 10.6 $e^+e^- \rightarrow e^+e^-\eta\eta$

NODE=M056G06
 NODE=M056G06

¹Interference with the continuum not included.

NODE=M056G06;LINKAGE=UE

$$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{39}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.9		90	1 LIU	12B	BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M056G02
 NODE=M056G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.9		90	1 LIU	12B	BELL $\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

¹Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M056G02;LINKAGE=LI

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{40}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.34		90	1 LIU	12B	BELL $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M056G03
 NODE=M056G03

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.34		90	1 LIU	12B	BELL $\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

¹Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M056G03;LINKAGE=LI

$$\Gamma(K^+K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
13.3 ± 1.1 OUR FIT					Error includes scale factor of 1.1. [13.4 ± 1.0 eV OUR 2023 FIT]
14.3 ± 1.6 ± 2.3		153 ± 17	NAKAZAWA	05	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$

NODE=M056G4
 NODE=M056G4
 NEW

$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{43}\Gamma_{98}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
7.0 ± 0.6 OUR FIT				Error includes scale factor of 1.2. [7.0 ± 0.5 eV OUR 2023 FIT]	
8.7 ± 1.7 ± 0.9	266	¹ UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	NEW
• • • We do not use the following data for averages, fits, limits, etc. • • •					
7.00 ± 0.65 ± 0.71	134 ± 12	CHEN	07B BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c0}$	
¹ Supersedes CHEN 07B.					

NODE=M056G5
NODE=M056G5

NODE=M056G5;LINKAGE=UE

$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{51}\Gamma_{98}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
6.2 ± 1.0 OUR FIT				Error includes scale factor of 1.5. [6.2 ± 0.7 eV OUR 2023 FIT]	
7.9 ± 1.3 ± 1.1	215 ± 36	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+ K^-)$	NEW

NODE=M056G11
NODE=M056G11

NEW

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{56}\Gamma_{98}/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.86 ± 0.13 OUR FIT				Error includes scale factor of 1.1. [1.76 ± 0.18 eV OUR 2023 FIT]	
1.72 ± 0.33 ± 0.14	56 ± 11	¹ LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+ K^-)$	NEW
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.3 ± 0.9 ± 0.4	23.6 ± 9.6	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c0} \rightarrow 2(K^+ K^-)$	
¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$.					

NODE=M056G12
NODE=M056G12

NEW

NODE=M056G12;LINKAGE=LI

$\chi_{c0}(1P)$ BRANCHING RATIOS

NODE=M056220

HADRONIC DECAYS

NODE=M056305

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-))$					Γ_2/Γ_1
VALUE		DOCUMENT ID	TECN	COMMENT	
0.39 ± 0.12 OUR FIT					
0.39 ± 0.12		TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$	

NODE=M056R54
NODE=M056R54

$\Gamma(f_0(980) f_0(980))/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
6.7 ± 2.1 OUR AVERAGE				[(6.6 ± 2.1) × 10 ⁻⁴ OUR 2023 AVERAGE]	
6.7 ± 2.1 ± 0.2	36 ± 9	¹ ABLIKIM	04G BES	$\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$	NEW
¹ ABLIKIM 04G reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980) f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (6.5 \pm 1.6 \pm 1.3) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M056R24
NODE=M056R24

NEW

NODE=M056R24;LINKAGE=AB

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$					Γ_5/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
3.3 ± 0.4 ± 0.1	1751.4	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	
¹ HE 08B reports $3.54 \pm 0.10 \pm 0.43 \pm 0.18 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

NODE=M056R62
NODE=M056R62

NODE=M056R62;LINKAGE=HE

$\Gamma(\rho^+ \pi^- \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
2.9 ± 0.4 ± 0.1	1358.5	^{1,2} HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$	
¹ HE 08B reports $3.04 \pm 0.18 \pm 0.42 \pm 0.16 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ \pi^- \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² Calculated by us. We have added the values from HE 08B for $\rho^+ \pi^- \pi^0$ and $\rho^- \pi^+ \pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.					

NODE=M056R63
NODE=M056R63

NODE=M056R63;LINKAGE=HE

NODE=M056R63;LINKAGE=OC

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.3±0.4±0.1	3296	¹ ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

NODE=M056R71
 NODE=M056R71

¹ ABLIKIM 11A reports $(3.34 \pm 0.06 \pm 0.44) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R71;LINKAGE=AB

 $\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma(\pi^+\pi^-K^+K^-)$ Γ_{30}/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
0.41±0.09 OUR FIT			
0.41±0.10	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$

NODE=M056R55
 NODE=M056R55

 $\Gamma(K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.9^{+4.0}_{-2.8} OUR AVERAGE				$[(9.8^{+4.0}_{-2.8}) \times 10^{-4} \text{ OUR 2023 AVERAGE}]$
9.9^{+3.6}_{-2.8}±0.2	83	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R36
 NODE=M056R36

NEW

¹ ABLIKIM 05Q reports $(10.44 \pm 2.37^{+3.05}_{-1.90}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0\bar{K}_0^*(1430)^0 \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R36;LINKAGE=AB

 $\Gamma(K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.0^{+2.0}_{-2.4}±0.2	62	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R37
 NODE=M056R37

¹ ABLIKIM 05Q reports $(8.49 \pm 1.66^{+1.32}_{-1.99}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_0^*(1430)^0\bar{K}_2^*(1430)^0 + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R37;LINKAGE=AB

 $\Gamma(K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.3±1.9±0.1	68	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R38
 NODE=M056R38

¹ ABLIKIM 05Q reports $(6.66 \pm 1.31^{+1.60}_{-1.51}) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1270)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The measurement assumes $B(K_1(1270) \rightarrow K\rho(770)) = 42 \pm 6\%$.

NODE=M056R38;LINKAGE=AB

 $\Gamma(K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R39
 NODE=M056R39

¹ ABLIKIM 05Q reports $< 2.85 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K_1(1400)^+K^- + \text{c.c.} \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$. The measurement assumes $B(K_1(1400) \rightarrow K^*(892)\pi) = 94 \pm 6\%$.

NODE=M056R39;LINKAGE=AB

 $\Gamma(f_0(980)f_0(980))/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.3^{+10.5}_{-9.0}±0.4	28	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R28
 NODE=M056R28

¹ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(980))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.59 \pm 0.50^{+0.89}_{-0.72}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. One of the $f_0(980)$ mesons is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay.

NODE=M056R28;LINKAGE=AB

$\Gamma(f_0(980)f_0(2200))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.9^{+2.0}_{-2.5} \pm 0.2$	77	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R29
NODE=M056R29

¹ABLIKIM 05Q reports $(8.42 \pm 1.42^{+1.65}_{-2.29}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(980)f_0(2200))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(980) \rightarrow \pi^+\pi^-$ and $f_0(2200) \rightarrow K^+K^-$ decays.

NODE=M056R29;LINKAGE=AB

$\Gamma(f_0(1370)f_0(1370))/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R30
NODE=M056R30

¹ABLIKIM 05Q reports $< 2.9 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$. One of the $f_0(1370)$ mesons is identified via decay to $\pi^+\pi^-$ while the other via K^+K^- decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R30;LINKAGE=AB

$\Gamma(f_0(1370)f_0(1500))/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R31
NODE=M056R31

¹ABLIKIM 05Q reports $< 1.8 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+\pi^-$ and $f_0(1500) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R31;LINKAGE=AB

$\Gamma(f_0(1370)f_0(1710))/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.7^{+3.5}_{-2.3} \pm 0.2$	61	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R32
NODE=M056R32

¹ABLIKIM 05Q reports $(7.12 \pm 1.85^{+3.28}_{-1.68}) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1370)f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The f_0 mesons are identified via $f_0(1370) \rightarrow \pi^+\pi^-$ and $f_0(1710) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R32;LINKAGE=AB

$\Gamma(f_0(1500)f_0(1370))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$

NODE=M056R33
NODE=M056R33

¹ABLIKIM 05Q reports $< 1.4 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500)f_0(1370))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+\pi^-$ and $f_0(1370) \rightarrow K^+K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.

NODE=M056R33;LINKAGE=AB

$\Gamma(f_0(1500) f_0(1500))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
¹ ABLIKIM 05Q reports $< 0.55 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500) f_0(1500))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$. One of the $f_0(1500)$ is identified via decay to $\pi^+ \pi^-$ while the other via $K^+ K^-$ decay. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

NODE=M056R34
NODE=M056R34

NODE=M056R34;LINKAGE=AB

 $\Gamma(f_0(1500) f_0(1710))/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	¹ ABLIKIM	05Q	BES2 $\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
¹ ABLIKIM 05Q reports $< 0.73 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow f_0(1500) f_0(1710))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$. The f_0 mesons are identified via $f_0(1500) \rightarrow \pi^+ \pi^-$ and $f_0(1710) \rightarrow K^+ K^-$ decays. Both branching fractions for these f_0 decays are implicitly included in the quoted result.				

NODE=M056R35
NODE=M056R35

NODE=M056R35;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.61±0.13±0.94	9.0k	¹ ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.				

NODE=M056R85
NODE=M056R85

NODE=M056R85;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.22±0.10±0.43	2.7k	¹ ABLIKIM	13B	BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma) = (9.68 \pm 0.31)\%$.				

NODE=M056R86
NODE=M056R86

NODE=M056R86;LINKAGE=A

 $\Gamma(K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.56±0.09±0.01	213.5	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $0.59 \pm 0.05 \pm 0.08 \pm 0.03 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R64
NODE=M056R64

NODE=M056R64;LINKAGE=HE

 $\Gamma(K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.49±0.33±0.06	401.7	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $2.64 \pm 0.15 \pm 0.31 \pm 0.14 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R66
NODE=M056R66

NODE=M056R66;LINKAGE=HE

 $\Gamma(\rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.21±0.21±0.03	179.7	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$
¹ HE 08B reports $1.28 \pm 0.16 \pm 0.15 \pm 0.07 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho^+ K^- K^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M056R67
NODE=M056R67

NODE=M056R67;LINKAGE=HE

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.46±0.12±0.01	64.1	¹ HE	08B	CLEO $e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M056R68
NODE=M056R68

¹ HE 08B reports $0.49 \pm 0.10 \pm 0.07 \pm 0.03 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R68;LINKAGE=HE

$\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ **Γ_{27}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.7 \pm 1.1 \pm 0.1$	152 ± 14	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R47
NODE=M056R47

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.558 \pm 0.051 \pm 0.089) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R47;LINKAGE=AB

$\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ **Γ_{28}/Γ**

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.30 \pm 0.07 \pm 0.01$	56.4	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M056R69
NODE=M056R69

¹ HE 08B reports $0.32 \pm 0.05 \pm 0.05 \pm 0.02 \%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R69;LINKAGE=HE

$\Gamma(3(\pi^+ \pi^-))/\Gamma_{\text{total}}$ **Γ_{29}/Γ**

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
19.5 ± 2.2 OUR AVERAGE		Error includes scale factor of 3.3. See the ideogram below. [(19.6 ± 2.1) × 10 ⁻³ OUR 2023 AVERAGE Scale factor = 3.3]		

NODE=M056R4
NODE=M056R4

NEW


$20.8 \pm 0.5 \pm 0.5$	145K	¹ ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
$11.7 \pm 1.0 \pm 1.9$		² BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c0}$
$12.5 \pm 2.9 \pm 0.5$		² TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R4;LINKAGE=A

¹ ABLIKIM 22Q reports $(2.080 \pm 0.006 \pm 0.068) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow 3(\pi^+ \pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.79 \pm 0.2) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

NODE=M056R;LINKAGE=X1



$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.72^{+0.60}_{-0.54} \pm 0.04$	64	¹ ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$

NODE=M056R26
 NODE=M056R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.57 \pm 0.40 \pm 0.04$ 30 \pm 6 ^{2,3} ABLIKIM 04H BES Repl. by ABLIKIM 05Q

¹ ABLIKIM 05Q reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(0.168 \pm 0.035^{+0.047}_{-0.040}) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R26;LINKAGE=AI

² Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

NODE=M056R;LINKAGE=AL

³ ABLIKIM 04H reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = $(1.53 \pm 0.29 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R26;LINKAGE=AB

 $\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-3}$	90	¹ ABLIKIM	15N BES3	$\psi(2S) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$

NODE=M056R00
 NODE=M056R00

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

NODE=M056R00;LINKAGE=A

 $\Gamma(\eta \eta')/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$9.1 \pm 1.1 \pm 0.2$		85	¹ ABLIKIM	17AI BES3	$\psi(2S) \rightarrow \gamma \eta' \eta$

NODE=M056R03
 NODE=M056R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 24 90 35 \pm 13 ² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \eta' \eta$

< 50 90 ³ ADAMS 07 CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ABLIKIM 17AI reports $(8.92 \pm 0.84 \pm 0.65) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R03;LINKAGE=A

² ASNER 09 reports $< 0.25 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R03;LINKAGE=AS

³ Superseded by ASNER 09. ADAMS 07 reports $< 0.5 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R03;LINKAGE=AD

 $\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.17 ± 0.12 OUR AVERAGE				

NODE=M056R04
 NODE=M056R04

$2.24 \pm 0.13 \pm 0.05$ 2.5k ¹ ABLIKIM 17AI BES3 $\psi(2S) \rightarrow \gamma \eta' \eta'$

$2.00 \pm 0.21 \pm 0.05$ 0.4k ² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \eta' \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.60 \pm 0.41 \pm 0.04$ 23 ³ ADAMS 07 CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$

¹ ABLIKIM 17AI reports $(2.19 \pm 0.03 \pm 0.14) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04;LINKAGE=A

² ASNER 09 reports $(2.12 \pm 0.13 \pm 0.21) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04;LINKAGE=AS

³ Superseded by ASNER 09. ADAMS 07 reports $(1.7 \pm 0.4 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.0922 \pm 0.0011 \pm 0.0046$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R04;LINKAGE=AD

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.11 OUR AVERAGE				
0.94±0.11±0.02	991	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
2.2 ±0.7 ±0.1	38.1±9.6	² ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma 6\pi$

 Γ_{39}/Γ NODE=M056R27
NODE=M056R27

¹ ABLIKIM 11K reports $(0.95 \pm 0.03 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R27;LINKAGE=AL

² ABLIKIM 05N reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (0.212 \pm 0.053 \pm 0.037) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R27;LINKAGE=AB

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.42±0.13 OUR AVERAGE				[(1.41 ± 0.13) × 10 ⁻⁴ OUR 2023 AVERAGE]
1.42±0.13±0.03	486	¹ ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons

 Γ_{40}/Γ NODE=M056R76
NODE=M056R76

• • • We do not use the following data for averages, fits, limits, etc. • • •

NEW

1.18±0.22±0.03 76 ^{2,3} ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (13.83 \pm 0.70 \pm 1.01) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R76;LINKAGE=A

² ABLIKIM 11K reports $(1.2 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R76;LINKAGE=AL

³ Superseded by ABLIKIM 19J.

NODE=M056R76;LINKAGE=B

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.94±0.06±0.20	1.4k	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

 Γ_{41}/Γ NODE=M056R87
NODE=M056R87

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$.

NODE=M056R87;LINKAGE=A

 $\Gamma(\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
<1.0	90	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$

 Γ_{44}/Γ NODE=M056R08
NODE=M056R08

¹ ATHAR 07 reports $< 0.21 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R08;LINKAGE=AT

² ABLIKIM 06R reports $< 1.1 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R08;LINKAGE=AB

 $\Gamma(\pi^+ \pi^- \eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.4	90	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

 Γ_{45}/Γ NODE=M056R51
NODE=M056R51

¹ ATHAR 07 reports $< 0.38 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \pi^+ \pi^- \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R51;LINKAGE=AT

 $\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.09	90	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
<0.7	90	^{2,3} ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c0}$
<0.7	90	^{3,4} BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c0}$

 Γ_{46}/Γ NODE=M056R17
NODE=M056R17

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ATHAR 07 reports $< 0.10 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R17;LINKAGE=AT

² ABLIKIM 06R reports $< 0.70 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R17;LINKAGE=AB

³ We have multiplied the $K_S^0 K^+ \pi^-$ measurement by a factor of 2 to convert to $K^0 K^+ \pi^-$.

NODE=M056R17;LINKAGE=BA

⁴ Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (32.6 \pm 0.5)\%$.

NODE=M056R17;LINKAGE=X1

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.06	90	¹ ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M056R05
NODE=M056R05

¹ ATHAR 07 reports $< 0.06 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R05;LINKAGE=AT

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.23	90	¹ ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M056R09
NODE=M056R09

¹ ATHAR 07 reports $< 0.24 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R09;LINKAGE=AT

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.41 ± 0.47 ± 0.03	16.8 ± 4.8	¹ ABLIKIM 050	BES2	$\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R48
NODE=M056R48

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (0.138 \pm 0.039 \pm 0.025) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R48;LINKAGE=AB

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.8 ± 0.5 ± 0.1	319	¹ ABLIKIM 19AA	BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$

NODE=M056R95
NODE=M056R95

¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))] = (5.64 \pm 0.33 \pm 0.37) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

NODE=M056R95;LINKAGE=A

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.25 ± 0.02	38	¹ ABLIKIM 06T	BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M056R01
NODE=M056R01

¹ ABLIKIM 06T reports $(1.03 \pm 0.22 \pm 0.15) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R01;LINKAGE=AB

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.68 ± 0.30 ± 0.50	ABLIKIM 15M	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R90
NODE=M056R90 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.90 ± 0.14 ± 0.32	ABLIKIM 15M	BES3	$\psi(2S) \rightarrow \gamma \chi_{c0}$

NODE=M056R91
NODE=M056R91

$\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.18±0.07±0.13	538	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.68 \pm 0.31)\%$.

NODE=M056R88
NODE=M056R88

NODE=M056R88;LINKAGE=A

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.848±0.031 OUR FIT				

[(0.80 ± 0.07) × 10⁻³ OUR 2023 FIT]

0.859±0.034 OUR AVERAGE [(0.9 ± 0.5) × 10⁻³ OUR 2006 AVERAGE]

0.859±0.027±0.020 2701 ¹ ABLIKIM 23N BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ Measured using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ and $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ from PDG 22.

NODE=M056R16
NODE=M056R16

NEW

NEW

NODE=M056R16;LINKAGE=A

 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.4±1.0±0.2	186.6	¹ ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma\phi\phi\eta$

¹ ABLIKIM 20B reports $(8.41 \pm 0.74 \pm 0.62) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R98
NODE=M056R98

NODE=M056R98;LINKAGE=A

 $\Gamma(\rho\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.70±0.07 OUR AVERAGE			

Error includes scale factor of 1.3. [(0.70 ± 0.07) × 10⁻³ OUR 2023 AVERAGE Scale factor = 1.3]

0.73±0.06±0.02 ¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma\rho\bar{p}X$
0.56±0.12±0.01 ² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(7.76 \pm 0.37 \pm 0.51 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.59 \pm 0.10 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R06
NODE=M056R06

NEW

NODE=M056R06;LINKAGE=ON

NODE=M056R06;LINKAGE=AT

 $\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.35±0.04 OUR AVERAGE			

0.35±0.04±0.01 ¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma\rho\bar{p}X$
0.37±0.11±0.01 ² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ONYISI 10 reports $(3.73 \pm 0.38 \pm 0.28 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ATHAR 07 reports $(0.39 \pm 0.11 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R50
NODE=M056R50

NODE=M056R50;LINKAGE=ON

NODE=M056R50;LINKAGE=AT

 $\Gamma(\rho\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.53±0.06 OUR AVERAGE			

[(0.52 ± 0.06) × 10⁻³ OUR 2023 AVERAGE]

0.53±0.06±0.01 ¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma\rho\bar{p}X$

¹ ONYISI 10 reports $(5.57 \pm 0.48 \pm 0.42 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \rho\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R70
NODE=M056R70

NEW

NODE=M056R70;LINKAGE=ON

$\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0±1.4±0.1	42 ± 8	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M056R75
 NODE=M056R75

¹ ABLIKIM 11F reports $(6.12 \pm 1.18 \pm 0.86) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R75;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.1 ± 0.7 OUR EVALUATION	Error includes scale factor of 1.4. Treating systematic error as correlated.		

NODE=M056R7
 NODE=M056R7

→ UNCHECKED ←

2.1 ± 1.0 OUR AVERAGE Error includes scale factor of 2.0.

1.57 ± 0.21 ± 0.53	¹ BAI	99B BES	$\psi(2S) \rightarrow \gamma\chi_{c0}$
4.20 ± 1.15 ± 0.18	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.4 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M056R7;LINKAGE=X1

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.104±0.028±0.002	39.5	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

NODE=M056R65
 NODE=M056R65

¹ HE 08B reports $0.11 \pm 0.02 \pm 0.02 \pm 0.01\%$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R65;LINKAGE=HE

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.22±0.26±0.03	48 ± 8	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M056R72
 NODE=M056R72

¹ ABLIKIM 11F reports $(1.24 \pm 0.20 \pm 0.18) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R72;LINKAGE=AB

 $\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8.8	90	¹ ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

NODE=M056R46
 NODE=M056R46

¹ Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

NODE=M056R;LINKAGE=AB

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.7±1.1 OUR AVERAGE				

NODE=M056R49
 NODE=M056R49

12.9 ± 1.1 ± 0.3	5150	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
11.3 ± 3.1 ± 0.3		² ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.26 \pm 0.02 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R49;LINKAGE=AL

² ABLIKIM 06I reports $[\Gamma(\chi_{c0}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.10 \pm 0.24 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R49;LINKAGE=AB

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
13.7±1.2±0.3	5808	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

NODE=M056R82
 NODE=M056R82

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (1.34 \pm 0.03 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R82;LINKAGE=AL

$\Gamma(\rho\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
23.4±2.0±0.5	2480	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\rho\bar{n}\pi^-\pi^0$

NODE=M056R83
 NODE=M056R83

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \rho\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.29 \pm 0.08 \pm 0.18) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R83;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
22.1±1.9 OUR AVERAGE [(22.1 ± 1.8) × 10 ⁻⁴ OUR 2023 AVERAGE]				
22.1±1.8±0.5	2757	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

NODE=M056R84
 NODE=M056R84

NEW

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c0}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (2.16 \pm 0.07 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R84;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
118±12±3		426	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••					
<400	90		² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c0}\gamma$

NODE=M056R44
 NODE=M056R44

¹ ABLIKIM 12I reports $(119.0 \pm 6.4 \pm 11.4) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R44;LINKAGE=AL

² Using $B(\psi(2S) \rightarrow \chi_{c0}\gamma) = (9.2 \pm 0.5)\%$

NODE=M056R44;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M056R77
 NODE=M056R77

¹ ABLIKIM 12I reports $< 54 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R77;LINKAGE=AL

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

NODE=M056R78
 NODE=M056R78

¹ ABLIKIM 12I reports $< 55 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R78;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<50	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

NODE=M056R79
 NODE=M056R79

¹ ABLIKIM 12I reports $< 50 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.

NODE=M056R79;LINKAGE=AL

 $\Gamma(K^+\bar{p}\Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.25±0.12 OUR AVERAGE Error includes scale factor of 1.3. [(1.25 ± 0.12) × 10 ⁻³ OUR 2023 AVERAGE Scale factor = 1.3]				
1.31±0.09±0.03	9k	^{1,2} ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$
1.01±0.19±0.02		³ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M056R07
 NODE=M056R07

NEW

¹ ABLIKIM 13D reports $(1.32 \pm 0.03 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R07;LINKAGE=AB

² Using $B(\Lambda \rightarrow p \pi^-) = 63.9\%$.

NODE=M056R07;LINKAGE=LB

³ ATHAR 07 reports $(1.07 \pm 0.17 \pm 0.12) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R07;LINKAGE=AT

$\Gamma(K^*(892)^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{79}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.8±0.9±0.1	254	¹ ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+} \bar{p} \Lambda$

NODE=M056R99
NODE=M056R99

¹ ABLIKIM 19AU reports $[(\Gamma(\chi_{c0}(1P) \rightarrow K^*(892)^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}) \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]] = (4.7 \pm 0.7 \pm 0.5) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R99;LINKAGE=A

$\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{80}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0±0.8 OUR AVERAGE				[(2.9 ± 0.7) × 10 ⁻⁴ OUR 2023 AVERAGE]

NODE=M056R73
NODE=M056R73

3.0±0.7±0.1 62 ± 12 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NEW

¹ ABLIKIM 11F reports $(3.00 \pm 0.58 \pm 0.50) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R73;LINKAGE=AB

$\Gamma(n K_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±0.5 OUR AVERAGE				[(6.6 ± 0.5) × 10 ⁻⁴ OUR 2023 AVERAGE]

NODE=M056P02
NODE=M056P02

6.7±0.5±0.2 1284 ¹ ABLIKIM 21AV BES3 $\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$

NEW

¹ ABLIKIM 21AV reports $(6.65 \pm 0.26 \pm 0.41) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow n K_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.0979 \pm 0.0020$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Also uses $B(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$.

NODE=M056P02;LINKAGE=A

$\Gamma(\Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}$ Γ_{81}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±1.2±0.1	28 ± 10	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

NODE=M056R74
NODE=M056R74

¹ ABLIKIM 11F reports $(3.18 \pm 1.11 \pm 0.53) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R74;LINKAGE=AB

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.69±0.32 OUR AVERAGE				[(4.68 ± 0.32) × 10 ⁻⁴ OUR 2023 AVERAGE]

NODE=M056R59
NODE=M056R59

4.83±0.34±0.11 1046 ¹ ABLIKIM 18V BES3 $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

NEW

4.2 ± 0.7 ± 0.1 78 ± 10 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.7 ± 0.5 ± 0.1 243 ^{3,4} ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

- ¹ ABLIKIM 18V reports $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (4.72 \pm 0.18 \pm 0.28) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² NAIK 08 reports $(4.41 \pm 0.56 \pm 0.47) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ABLIKIM 13H reports $(4.78 \pm 0.34 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ Superseded by ABLIKIM 18V

NODE=M056R59;LINKAGE=A

NODE=M056R59;LINKAGE=NA

NODE=M056R59;LINKAGE=AB

NODE=M056R59;LINKAGE=B

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$

Γ_{85}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

4.7 ± 0.8 OUR AVERAGE Error includes scale factor of 2.6. $[(4.6 \pm 0.8) \times 10^{-4}$ OUR 2023 AVERAGE Scale factor = 2.6]

NODE=M056R60
NODE=M056R60
NEW

5.11 ± 0.35 ± 0.12 747 ¹ ABLIKIM 18V BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

3.1 ± 0.7 ± 0.1 39 ± 7 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.5 ± 0.5 ± 0.1 148 ^{3,4} ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

- ¹ ABLIKIM 18V reports $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))] = (4.99 \pm 0.24 \pm 0.24) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² NAIK 08 reports $(3.25 \pm 0.57 \pm 0.43) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.22 \pm 0.11 \pm 0.46) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ABLIKIM 13H reports $(4.54 \pm 0.42 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ Superseded by ABLIKIM 18V

NODE=M056R60;LINKAGE=A

NODE=M056R60;LINKAGE=NA

NODE=M056R60;LINKAGE=AB

NODE=M056R60;LINKAGE=B

$\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$

Γ_{86}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

5.1 ± 0.5 ± 0.1 2143 ¹ ABLIKIM 20I BES3 $\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

NODE=M056P00
NODE=M056P00

- ¹ ABLIKIM 20I reports $(5.13 \pm 0.24 \pm 0.41) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.79 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056P00;LINKAGE=A

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

Γ_{87}/Γ

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

16.3 ± 5.8 ± 0.4 27 ¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M056R80
NODE=M056R80

- ¹ ABLIKIM 12I reports $(16.4 \pm 5.7 \pm 1.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R80;LINKAGE=AL

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

Γ_{88}/Γ

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

23 ± 7 ± 1 33 ¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M056R81
NODE=M056R81

- ¹ ABLIKIM 12I reports $(23.5 \pm 6.2 \pm 2.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.68 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R81;LINKAGE=AL

$\Gamma(K^- \Lambda_{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{89}/Γ VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT**1.95 ± 0.35 OUR AVERAGE** [(1.94 ± 0.35) × 10⁻⁴ OUR 2023 AVERAGE]**1.95 ± 0.35 ± 0.05** 57 ¹ ABLIKIM 15i BES3 $\psi(2S) \rightarrow \gamma K^- \Lambda_{\Xi}^+ + \text{c.c.}$

¹ ABLIKIM 15i reports $[\Gamma(\chi_{c0}(1P) \rightarrow K^- \Lambda_{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ = (1.90 ± 0.30 ± 0.16) × 10⁻⁵ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.77 ± 0.23) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R92
NODE=M056R92

NEW

NODE=M056R92;LINKAGE=A

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{90}/Γ VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT**4.5 ± 0.5 OUR AVERAGE** Error includes scale factor of 1.7. [(4.5 ± 0.5) × 10⁻⁴ OUR 2023 AVERAGE Scale factor = 1.7]4.68 ± 0.31 ± 0.11 1741 ¹ ABLIKIM 220 BES3 $\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$
3.2 ± 0.8 ± 0.1 23.3 ± 4.9 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$

¹ ABLIKIM 220 reports (4.67 ± 0.19 ± 0.26) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.79 ± 0.2) × 10⁻², which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.77 ± 0.23) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R61
NODE=M056R61

NEW

NODE=M056R61;LINKAGE=A

² NAIK 08 reports (3.34 ± 0.70 ± 0.48) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.22 ± 0.11 ± 0.46) × 10⁻², which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.77 ± 0.23) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R61;LINKAGE=NA

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{91}/Γ VALUE (units 10^{-4}) CL% EVTS DOCUMENT ID TECN COMMENT**4.47 ± 0.20 OUR AVERAGE**[(4.45 ± 0.19) × 10⁻⁴ OUR 2023 AVERAGE]4.44 ± 0.18 ± 0.10 4932 ¹ ABLIKIM 220 BES3 $\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$
4.9 ± 0.7 ± 0.1 95 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<10.3 90 ³ ABLIKIM 06D BES2 $\psi(2S) \rightarrow \chi_{c0} \gamma$

¹ ABLIKIM 220 reports (4.43 ± 0.08 ± 0.18) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.79 ± 0.2) × 10⁻², which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.77 ± 0.23) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R45
NODE=M056R45

NEW

NODE=M056R45;LINKAGE=A

² NAIK 08 reports (5.14 ± 0.60 ± 0.47) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.22 ± 0.11 ± 0.46) × 10⁻², which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))$ = (9.77 ± 0.23) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M056R45;LINKAGE=NA

³ Using $B(\psi(2S) \rightarrow \chi_{c0} \gamma)$ = (9.2 ± 0.5)%

NODE=M056R45;LINKAGE=AB

 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT**3.51 ± 0.54 ± 0.29** 284 ABLIKIM 23T BES3 $\chi_{cJ} \rightarrow \Omega^- \bar{\Omega}^+$ NODE=M056P04
NODE=M056P04 $\Gamma(\eta_c \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE CL% DOCUMENT ID TECN COMMENT

< 7 × 10⁻⁴ 90 ^{1,2} ABLIKIM 13B BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 41 × 10⁻⁴ 90 ^{1,3} ABLIKIM 13B BES3 $e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

¹ Using 1.06 × 10⁸ $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c0} \gamma)$ = (9.68 ± 0.31)%.

² From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

³ From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

NODE=M056R89
NODE=M056R89

OCCUR=2

NODE=M056R89;LINKAGE=A

NODE=M056R89;LINKAGE=B

NODE=M056R89;LINKAGE=C

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{32}/\Gamma$	
VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT		
18.9±1.5 OUR FIT	Error includes scale factor of 1.5. $[(18.8 \pm 1.0) \times 10^{-7}$ OUR 2023 FIT]				NEW
15.3±2.4±0.8	¹ ANDREOTTI	03	E835 $\bar{p}p \rightarrow \chi_{c0} \rightarrow \pi^0\pi^0$		NODE=M056R;LINKAGE=AD
¹ We have multiplied $B(p\bar{p}) \cdot B(\pi^0\pi^0)$ measurement by 3 to obtain $B(p\bar{p}) \cdot B(\pi\pi)$.					
$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0\eta)/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{33}/\Gamma$	
VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT		
<0.4	ANDREOTTI	05C	E835 $\bar{p}p \rightarrow \pi^0\eta$		NODE=M056R41 NODE=M056R41
$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi^0\eta')/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{34}/\Gamma$	
VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT		
<2.5	ANDREOTTI	05C	E835 $\bar{p}p \rightarrow \pi^0\eta$		NODE=M056R42 NODE=M056R42
$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{36}/\Gamma$	
VALUE (units 10^{-7})	DOCUMENT ID	TECN	COMMENT		
6.7±0.7 OUR FIT	Error includes scale factor of 1.4. $[(6.7 \pm 0.5) \times 10^{-7}$ OUR 2023 FIT]				NEW
4.0±1.2^{+0.5}_{-0.3}	ANDREOTTI	05C	E835 $\bar{p}p \rightarrow \eta\eta$		NODE=M056R40 NODE=M056R40
$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta')/\Gamma_{\text{total}}$				$\Gamma_{58}/\Gamma \times \Gamma_{37}/\Gamma$	
VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT		
2.1^{+2.3}_{-1.5}	ANDREOTTI	05C	E835 $\bar{p}p \rightarrow \pi^0\eta$		NODE=M056R43 NODE=M056R43
• • • We do not use the following data for averages, fits, limits, etc. • • •					

————— RADIATIVE DECAYS —————

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$				Γ_{95}/Γ	
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 9	90	1.2 ± 4.5	¹ BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<10	90	6 ± 12	² ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\rho^0$
¹ BENNETT 08A reports $< 9.6 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.					
² ABLIKIM 11E reports $< 10.5 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.					
NODE=M056R56;LINKAGE=BE					
NODE=M056R56;LINKAGE=AB					
$\Gamma(\gamma\omega)/\Gamma_{\text{total}}$				Γ_{96}/Γ	
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8	90	0.0 ± 2.8	¹ BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\omega$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<13	90	5 ± 11	² ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\omega$
¹ BENNETT 08A reports $< 8.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.					
² ABLIKIM 11E reports $< 12.9 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.					
NODE=M056R57;LINKAGE=BE					
NODE=M056R57;LINKAGE=AB					
$\Gamma(\gamma\phi)/\Gamma_{\text{total}}$				Γ_{97}/Γ	
VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 6	90	0.1 ± 1.6	¹ BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma\gamma\phi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<16	90	15 ± 7	² ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma\gamma\phi$
¹ BENNETT 08A reports $< 6.4 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.					
² ABLIKIM 11E reports $< 16.2 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow \gamma\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = 9.77 \times 10^{-2}$.					
NODE=M056R58;LINKAGE=BE					
NODE=M056R58;LINKAGE=AB					

$\Gamma(e^+e^- J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R93
 NODE=M056R93

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.54 ± 0.33 ± 0.04	56	^{1,2} ABLIKIM	17I	BES3	$\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$
--------------------	----	------------------------	-----	------	--

NODE=M056R93;LINKAGE=B

¹ ABLIKIM 17I reports $(1.51 \pm 0.30 \pm 0.13) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c0}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Not independent from other measurements reported by ABLIKIM 17I

NODE=M056R93;LINKAGE=C

 $\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ Γ_{99}/Γ_{94}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R94
 NODE=M056R94

9.5 ± 1.9 ± 0.7	56	¹ ABLIKIM	17I	BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$
------------------------	----	----------------------	-----	------	--

NODE=M056R94;LINKAGE=A

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) \times B(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)) = (15.8 \pm 0.3 \pm 0.6) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

 $\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S))$ Γ_{100}/Γ_{99}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

NODE=M056R97
 NODE=M056R97

<0.14	90	<9.5	ABLIKIM	19Z	BES3	$\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma(\mu^+ \mu^- J/\psi)$
-----------------	----	------	---------	-----	------	---

 $\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S))$ Γ_{98}/Γ_{94}

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M056R18
 NODE=M056R18

1.45 ± 0.13 OUR FIT Error includes scale factor of 1.6. $[(1.45 \pm 0.08) \times 10^{-2}$ OUR 2023 FIT]

NEW

2.0 ± 0.4 OUR AVERAGE

2.2 ± 0.4	^{+0.1} _{-0.2}	¹ ANDREOTTI	04	E835	$p\bar{p} \rightarrow \chi_{c0} \rightarrow \gamma\gamma$
-----------	------------------------------------	------------------------	----	------	---

1.45 ± 0.74	² AMBROGIANI	00B	E835	$\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$
-------------	-------------------------	-----	------	--

¹ The values of $B(p\bar{p})B(\gamma\gamma)$ and $B(\gamma\gamma)B(\gamma J/\psi)$ measured by ANDREOTTI 04 are not independent. The latter is used in the fit because of smaller systematics.

NODE=M056R;LINKAGE=AN

² Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

NODE=M056R;LINKAGE=7A

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$ $\Gamma_{58}/\Gamma \times \Gamma_{94}/\Gamma$

VALUE (units 10^{-7})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056R19
 NODE=M056R19

31.1 ± 2.2 OUR FIT Error includes scale factor of 1.4. $[(31.1 \pm 1.5) \times 10^{-7}$ OUR 2023 FIT]

NEW

28.2 ± 2.1 OUR AVERAGE

28.0 ± 1.9 ± 1.3	392	^{1,2,3} BAGNASCO	02	E835	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
------------------	-----	---------------------------	----	------	---

29.3 ^{+5.7} _{-4.7} ± 1.5	89	^{1,2} AMBROGIANI	99B	$\bar{p}p \rightarrow \chi_{c0} \rightarrow J/\psi\gamma$
--	----	---------------------------	-----	---

¹ Values in $(\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$ and $(\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}})$ are not independent. The latter is used in the fit since it is less correlated to the total width.

NODE=M056R;LINKAGE=KS

² Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

NODE=M056R19;LINKAGE=7A
 NODE=M056R19;LINKAGE=AN

³ Recalculated by ANDREOTTI 05A.

 $\chi_{c0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M056230

 $\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$ $\Gamma_{58}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{\psi(2S)}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M056B6
 NODE=M056B6

21.6 ± 1.3 OUR FIT Error includes scale factor of 1.5. $[(21.7 \pm 0.9) \times 10^{-6}$ OUR 2023 FIT]

NEW

23.7 ± 1.0 OUR AVERAGE

23.7 ± 0.8 ± 0.9	1222	ABLIKIM	13v	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
------------------	------	---------	-----	------	--

23.7 ± 1.4 ± 1.4	383 ± 22	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
------------------	----------	-------------------	----	------	--

23.6 ^{+3.7} _{-3.4} ± 3.4	89.5 ⁺¹⁴ ₋₁₃	BAI	04F	BES	$\psi(2S) \rightarrow \gamma \chi_{c0}(1P) \rightarrow \gamma p\bar{p}$
--	------------------------------------	-----	-----	-----	---

¹ Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow p\bar{p}) = (25.7 \pm 1.5 \pm 1.5 \pm 1.3) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.

NODE=M056B6;LINKAGE=NA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{58}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5}) DOCUMENT ID TECN COMMENT
6.2±0.4 OUR FIT Error includes scale factor of 1.5. [(6.25 ± 0.26) × 10⁻⁵ OUR 2023 FIT]

NODE=M056B1
 NODE=M056B1
 NEW

4.6±1.9 ¹BAI 98I BES $\psi(2S) \rightarrow \gamma\chi_{c0} \rightarrow \gamma\bar{p}p$
¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B;LINKAGE=B1

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{71}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT
35.2±1.3 OUR FIT
35.1±1.4 OUR AVERAGE Error includes scale factor of 1.1.

NODE=M056B20
 NODE=M056B20

35.6±1.0±1.0 1486 ABLIKIM 21L BES3 $\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+$
 31.2±3.3±2.0 131 ¹NAIK 08 CLEO $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
 ●●● We do not use the following data for averages, fits, limits, etc. ●●●
 32.0±1.9±2.2 369 ^{2,3}ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
¹ Calculated by us. NAIK 08 reports $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.8 \pm 3.6 \pm 2.2 \pm 1.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$.
² Superseded by ABLIKIM 21L
³ Calculated by us. ABLIKIM 13H reports $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) = (33.3 \pm 2.0 \pm 2.6) \times 10^{-5}$ from a measurement of $B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c0})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.62 \pm 0.31)\%$.

NODE=M056B20;LINKAGE=NA

NODE=M056B20;LINKAGE=A
 NODE=M056B20;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{71}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT
10.1±0.4 OUR FIT

NODE=M056B21
 NODE=M056B21

13.0^{+3.6}_{-3.5}±2.5 15.2^{+4.2}_{-4.0} ¹BAI 03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
¹ BAI 03E reports [$B(\chi_{c0} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c0}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^-p) / B(J/\psi \rightarrow p\bar{p})] = (2.45^{+0.68}_{-0.65} ± 0.46)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^-p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

NODE=M056B21;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{74}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT
2.26±0.30±0.20 67 ABLIKIM 22A0 BES3 $\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+\gamma\gamma$

NODE=M056P03
 NODE=M056P03

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{94}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT
0.137±0.009 OUR FIT Error includes scale factor of 1.7. [(0.138 ± 0.005) × 10⁻² OUR 2023 FIT]

NODE=M056B2
 NODE=M056B2

NEW

0.147±0.029 OUR AVERAGE Error includes scale factor of 4.6.
 0.158±0.003±0.006 4.8k ¹ABLIKIM 17N BES3 $\psi(2S) \rightarrow \gamma\gamma J/\psi$
 0.024±0.015±0.205 12k ABLIKIM 17U BES3 $e^+e^- \rightarrow \gamma X$
 0.069±0.018 ²OREGLIA 82 CBAL $\psi(2S) \rightarrow \gamma\chi_{c0}$
 0.4 ±0.3 ³BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma\chi_{c0}$
 0.16 ±0.11 ³BARTEL 78B CNTR $\psi(2S) \rightarrow \gamma\chi_{c0}$
 3.3 ±1.7 ⁴BIDDICK 77 CNTR $e^+e^- \rightarrow \gamma X$
 ●●● We do not use the following data for averages, fits, limits, etc. ●●●
 0.151±0.003±0.010 4.3k ⁵ABLIKIM 12O BES3 $\psi(2S) \rightarrow \gamma\chi_{c0}$
 0.125±0.007±0.013 560 ⁶MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma\chi_{c0}$
 0.18 ±0.01 ±0.02 ⁷ADAM 05A CLEO Repl. by MENDEZ 08

NODE=M056B2;LINKAGE=A

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

NODE=M056B;LINKAGE=3Q
 NODE=M056B;LINKAGE=2Q
 NODE=M056B;LINKAGE=EA
 NODE=M056B2;LINKAGE=B
 NODE=M056B2;LINKAGE=ME
 NODE=M056B2;LINKAGE=AD

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S))}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \quad \Gamma_{94}/\Gamma \times \frac{\Gamma_{179}^{\psi(2S)}}{\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT
0.396±0.025 OUR FIT Error includes scale factor of 1.7. [(0.397 ± 0.015) × 10⁻² OUR 2023 FIT]

NODE=M056B8
 NODE=M056B8

NEW

0.358±0.020±0.037 560 MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma \chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.55 ± 0.04 ± 0.06 172 ¹ ADAM 05A CLEO Repl. by MENDEZ 08

¹ Not independent from other values reported by ADAM 05A.

NODE=M056B;LINKAGE=AD

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \gamma \gamma)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \quad \Gamma_{98}/\Gamma \times \frac{\Gamma_{179}^{\psi(2S)}}{\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT
2.00±0.09 OUR FIT
 [(2.00 ± 0.08) × 10⁻⁵ OUR 2023 FIT]

NODE=M056B3
 NODE=M056B3

NEW

1.95±0.09 OUR AVERAGE

1.93±0.08±0.05 3.5k ABLIKIM 17AE BES3 $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$

2.17±0.32±0.10 0.2k ECKLUND 08A CLEO $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$

3.7 ± 1.8 ± 1.0 LEE 85 CBAL $\psi(2S) \rightarrow \gamma \chi_{c0}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.17±0.17±0.12 0.8k ¹ ABLIKIM 12A BES3 $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow 3\gamma$

¹ Superseded by ABLIKIM 17AE.

NODE=M056B3;LINKAGE=A

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi \pi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma_{\text{total}}} \quad \Gamma_{32}/\Gamma \times \frac{\Gamma_{179}^{\psi(2S)}}{\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
8.34±0.34 OUR FIT Error includes scale factor of 1.2. [(8.34 ± 0.29) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B22
 NODE=M056B22

NEW

8.80±0.34 OUR AVERAGE

9.11±0.08±0.65 17k ¹ ABLIKIM 10A BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c0}$

8.81±0.11±0.43 8.9k ² ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \pi^+ \pi^-$

8.13±0.19±0.89 2.8k ³ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma \pi^0 \pi^0$

OCCUR=2

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c0} \rightarrow \pi^0 \pi^0) = (3.23 \pm 0.03 \pm 0.23 \pm 0.14) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.4 \pm 0.4)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

NODE=M056B22;LINKAGE=AB

² Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^+ \pi^-) = (6.37 \pm 0.08 \pm 0.31 \pm 0.32) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi \pi$.

NODE=M056B22;LINKAGE=AS

³ Calculated by us. ASNER 09 reports $B(\chi_{c0} \rightarrow \pi^0 \pi^0) = (2.94 \pm 0.07 \pm 0.32 \pm 0.15) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c0}) = (9.22 \pm 0.11 \pm 0.46)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

NODE=M056B22;LINKAGE=AN

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi \pi)}{\Gamma_{\text{total}}} \times \frac{\Gamma(\psi(2S) \rightarrow \gamma \chi_{c0}(1P))}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \quad \Gamma_{32}/\Gamma \times \frac{\Gamma_{179}^{\psi(2S)}}{\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
24.0±1.0 OUR FIT Error includes scale factor of 1.2. [(24.0 ± 0.8) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B5
 NODE=M056B5

NEW

20.7±1.7 OUR AVERAGE

23.9±2.7±4.1 97 ± 11 ¹ BAI 03C BES $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma \pi^0 \pi^0$

20.2±1.1±1.5 720 ± 32 ² BAI 98i BES $\psi(2S) \rightarrow \gamma \chi_{c0} \rightarrow \gamma \pi^+ \pi^-$

¹ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

NODE=M056B;LINKAGE=D1

² Calculated by us. The value for $B(\chi_{c0} \rightarrow \pi^+ \pi^-)$ reported in BAI 98i is derived using $B(\psi^f \rightarrow \gamma \chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi^f \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi \pi$.

NODE=M056B;LINKAGE=D2

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{36}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
2.94±0.22 OUR FIT Error includes scale factor of 1.2. [(2.95 ± 0.18) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B11
 NODE=M056B11
 NEW

3.12±0.19 OUR AVERAGE

3.23±0.09±0.23 2132 ¹ ABLIKIM 10A BES3 e⁺e⁻ → ψ(2S) → γχ_{c0}
 2.93±0.12±0.29 0.9k ² ASNER 09 CLEO ψ(2S) → γηη

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.86±0.46±0.37 48 ³ ADAMS 07 CLEO ψ(2S) → γχ_{c0}

¹ Calculated by us. ABLIKIM 10A reports B(χ_{c0} → ηη) = (3.44 ± 0.10 ± 0.24 ± 0.13) × 10⁻³ using B(ψ(2S) → γχ_{c0}) = (9.4 ± 0.4)%.

NODE=M056B11;LINKAGE=AB

² Calculated by us. ASNER 09 reports B(χ_{c0} → ηη) = (3.18 ± 0.13 ± 0.31 ± 0.16) × 10⁻³ using B(ψ(2S) → γχ_{c0}) = (9.22 ± 0.11 ± 0.46)%.

NODE=M056B11;LINKAGE=AS

³ Superseded by ASNER 09. Calculated by us. The value of B(χ_{c0}(1P) → ηη) reported by ADAMS 07 was derived using B(ψ(2S) → γχ_{c0}(1P)) = (9.22 ± 0.11 ± 0.46)% (ATHAR 04).

NODE=M056B11;LINKAGE=AD

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \eta\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{36}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-3}) DOCUMENT ID TECN COMMENT
0.85 ± 0.06 OUR FIT Error includes scale factor of 1.2. [(0.85 ± 0.05) × 10⁻³ OUR 2023 FIT]

NODE=M056B10
 NODE=M056B10

0.578±0.241±0.158 BAI 03C BES ψ(2S) → γηη

NEW

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{42}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
5.93±0.28 OUR FIT [(5.92 ± 0.28) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B23
 NODE=M056B23

5.97±0.07±0.32 8.1k ¹ ASNER 09 CLEO ψ(2S) → γK⁺K⁻

NEW

¹ Calculated by us. ASNER 09 reports B(χ_{c0} → K⁺K⁻) = (6.47 ± 0.08 ± 0.35 ± 0.32) × 10⁻³ using B(ψ(2S) → γχ_{c0}) = (9.22 ± 0.11 ± 0.46)%.

NODE=M056B23;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{42}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT
1.71±0.08 OUR FIT

NODE=M056B9
 NODE=M056B9

1.63±0.10±0.15 774 ± 38 ¹ BAI 98I BES ψ(2S) → γK⁺K⁻

¹ Calculated by us. The value for B(χ_{c0} → K⁺K⁻) reported by BAI 98I is derived using B(ψ(2S) → γχ_{c0}) = (9.3 ± 0.8)% and B(ψ(2S) → J/ψπ⁺π⁻) = (32.4 ± 2.6)% [BAI 98D].

NODE=M056B9;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{43}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma\psi(2S)}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
3.10±0.16 OUR FIT [(3.10 ± 0.16) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B12
 NODE=M056B12

3.18±0.17 OUR AVERAGE

3.22±0.07±0.17 2.1k ¹ ASNER 09 CLEO ψ(2S) → γK_S⁰K_S⁰
 3.02±0.19±0.33 322 ABLIKIM 050 BES2 ψ(2S) → γK_S⁰K_S⁰

NEW

¹ Calculated by us. ASNER 09 reports B(χ_{c0} → K_S⁰K_S⁰) = (3.49 ± 0.08 ± 0.18 ± 0.17) × 10⁻³ using B(ψ(2S) → γχ_{c0}) = (9.22 ± 0.11 ± 0.46)%.

NODE=M056B12;LINKAGE=AS

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{43}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT
8.9±0.5 OUR FIT [(8.9 ± 0.5) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B13
 NODE=M056B13

5.6±0.8±1.3 ¹ BAI 99B BES ψ(2S) → γK_S⁰K_S⁰

NEW

¹ Calculated by us. The value of B(χ_{c0} → K_S⁰K_S⁰) reported by BAI 99B was derived using B(ψ(2S) → γχ_{c0}(1P)) = (9.3 ± 0.8)% and B(ψ(2S) → J/ψπ⁺π⁻) = (32.4 ± 2.6)% [BAI 98D].

NODE=M056B13;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow 2(\pi^+\pi^-))/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_1/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10⁻³) DOCUMENT ID TECN COMMENT

6.6±1.0 OUR FIT Error includes scale factor of 2.0. [(6.6 ± 0.5) × 10⁻³ OUR 2023 FIT]

NODE=M056B4
NODE=M056B4

NEW

6.9±2.4 OUR AVERAGE Error includes scale factor of 3.8.

4.4±0.1±0.9 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c0}$
9.3±0.9 ² TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c0}$

NODE=M056B;LINKAGE=B2

¹ Calculated by us. The value for $B(\chi_{c0} \rightarrow 2\pi^+2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98b].

² The value $B(\psi(1S) \rightarrow \gamma\chi_{c0}) \times B(\chi_{c0} \rightarrow 2\pi^+2\pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

NODE=M056B;LINKAGE=J1

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_8/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10⁻³) DOCUMENT ID TECN COMMENT

1.78±0.15 OUR FIT Error includes scale factor of 1.1. [(1.78 ± 0.14) × 10⁻³ OUR 2023 FIT]

NODE=M056B18
NODE=M056B18

NEW

1.64±0.05±0.2 ABLIKIM 05Q BES2 $\psi(2S) \rightarrow \gamma\chi_{c0}$

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \pi^+\pi^-K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_8/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10⁻³) DOCUMENT ID TECN COMMENT

5.1 ±0.4 OUR FIT Error includes scale factor of 1.1. [(5.1 ± 0.4) × 10⁻³ OUR 2023 FIT]

NODE=M056B19
NODE=M056B19

NEW

5.8 ±1.6 OUR AVERAGE Error includes scale factor of 2.3.

4.22±0.20±0.97 BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c0}$
7.4 ±1.0 ¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c0}$

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+\pi^-J/\psi) \times B(J/\psi \rightarrow \ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

NODE=M056B19;LINKAGE=TA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{51}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10⁻⁴) EVTS DOCUMENT ID TECN COMMENT

2.8 ±0.4 OUR FIT Error includes scale factor of 1.5. [(2.76 ± 0.28) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B14
NODE=M056B14

NEW

3.20±0.11±0.41 278 ¹ ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

NODE=M056B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow K^+K^-K^+K^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{51}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10⁻⁴) DOCUMENT ID TECN COMMENT

8.0±1.2 OUR FIT Error includes scale factor of 1.5. [(8.0 ± 0.8) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B15
NODE=M056B15

NEW

6.1±0.8±0.9 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow 2K^+2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98b].

NODE=M056B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{56}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10⁻⁴) EVTS DOCUMENT ID TECN COMMENT

0.829±0.034 OUR FIT [(0.78 ± 0.07) × 10⁻⁴ OUR 2023 FIT]

NODE=M056B16
NODE=M056B16

NEW

0.78 ±0.08 OUR AVERAGE

0.77 ±0.03 ±0.08 612 ¹ ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons
0.86 ±0.19 ±0.12 26 ² ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.62 \pm 0.31)\%$.

² Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.2 \pm 0.4)\%$.

NODE=M056B16;LINKAGE=AL

NODE=M056B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{56}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

2.39±0.10 OUR FIT

[(2.25 ± 0.21) × 10^{-4} OUR 2023 FIT]

2.6 ±1.0 ±1.1 ¹BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

¹ Calculated by us. The value of $B(\chi_{c0} \rightarrow \phi\phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.3 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M056B17

NODE=M056B17

NEW

NODE=M056B17;LINKAGE=BA

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{83}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

3.45±0.17±0.19 493 ¹ABLIKIM 19BB BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_c^0 \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (3.52 \pm 0.19 \pm 0.21) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma\chi_c^0) = (9.79 \pm 0.20)\%$ and other branching fractions from PDG 18.

NODE=M056B24

NODE=M056B24

NODE=M056B24;LINKAGE=A

$$\frac{\Gamma(\chi_{c0}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))/\Gamma_{\text{total}}}{\Gamma_{84}/\Gamma \times \Gamma_{179}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

2.97±0.12±0.14 871 ¹ABLIKIM 20AE BES3 $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$

¹ Calculated by us. ABLIKIM 20AE reports $B(\chi_c^0 \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (3.03 \pm 0.12 \pm 0.15) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma\chi_c^0) = (9.79 \pm 0.20)\%$ and other branching fractions from PDG 20.

NODE=M056P01

NODE=M056P01

NODE=M056P01;LINKAGE=A

$\chi_{c0}(1P)$ REFERENCES

ABLIKIM	23N	JHEP 2305 069	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62056
ABLIKIM	23T	PR D107 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62064
ABLIKIM	22AO	PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61887
ABLIKIM	22O	JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61652
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)	REFID=61634
ABLIKIM	21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61465
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61117
ABLIKIM	20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60733
ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60212
ABLIKIM	20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60303
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59844
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60026
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59606
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59837
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58990
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AAJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58310
ABLIKIM	17AI	PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58321
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56778
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56779
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761

NODE=M056

BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
ABE	07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50958
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
ANDREOTTI	05C	PR D72 112002	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50991
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABE	04G	PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ABLIKIM	04G	PR D70 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50187
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ANDREOTTI	04	PL B584 16	M. Andreotti <i>et al.</i>	(E835 Collab.)	REFID=49744
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
ANDREOTTI	03	PRL 91 091801	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=49578
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE,K	02	PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
BAGNASCO	02	PL B533 237	S. Bagnasco <i>et al.</i>	(FNAL E835 Collab.)	REFID=48833
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
AMBROGIANI	99B	PRL 83 2902	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47389
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059

$\chi_{c1}(1P)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See the Review on "Branching Ratios of $\psi(2S)$, $\chi_{c0,1,2}$ and $\eta_c(1S)$ " before the $\chi_{c0}(1P)$ Listings.

NODE=M055

NODE=M055

 $\chi_{c1}(1P)$ MASS

NODE=M055M

NODE=M055M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3510.67 ± 0.05 OUR AVERAGE		Error includes scale factor of 1.2.		
3509.84 ± 0.69 ± 0.64	2.8k	AAIJ	23AH LHCb	$B^+ \rightarrow K^+ (K_S^0 K \pi)$
3508.4 ± 1.9 ± 0.7	460	¹ AAIJ	17BB LHCb	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+ K^-)X$
3510.71 ± 0.04 ± 0.09	4.8k	² AAIJ	17BI LHCb	$\chi_{c1} \rightarrow J/\psi \mu^+ \mu^-$
3510.30 ± 0.14 ± 0.16		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3510.719 ± 0.051 ± 0.019		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
3509.4 ± 0.9		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3510.60 ± 0.087 ± 0.019	513	³ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
3511.3 ± 0.4 ± 0.4	30	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
3512.3 ± 0.3 ± 4.0		⁴ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7	91	⁵ LEMOIGNE	82 GOLI	$185 \pi^- Be \rightarrow \gamma \mu^+ \mu^- A$
3510.4 ± 0.6		OREGLIA	82 CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1	254	⁶ HIMEL	80 MRK2	$e^+ e^- \rightarrow J/\psi 2\gamma$
3509 ± 11	21	BRANDELIK	79B DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
3507 ± 3		⁶ BARTEL	78B CNTR	$e^+ e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4		^{6,7} TANENBAUM	78 MRK1	$e^+ e^-$
3513 ± 7	367	⁶ BIDDICK	77 CNTR	$\psi(2S) \rightarrow \gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3500 ± 10	40	TANENBAUM	75 MRK1	Hadrons γ

OCCUR=2

NODE=M055M;LINKAGE=A

NODE=M055M;LINKAGE=B

NODE=M055M;LINKAGE=NW

NODE=M055M;LINKAGE=C

NODE=M055M;LINKAGE=P

NODE=M055M;LINKAGE=D

NODE=M055M;LINKAGE=M

¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c1}(1P)$ fixed to the PDG 16 value.

² AAIJ 17BI reports also $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$ MeV.

³ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

⁴ Using mass of $\psi(2S) = 3686.0$ MeV.

⁵ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁶ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁷ From a simultaneous fit to radiative and hadronic decay channels.

 $\chi_{c1}(1P)$ WIDTH

NODE=M055W

NODE=M055W

NEW

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.84 ± 0.04 OUR FIT			Error includes scale factor of 1.1. [0.84 ± 0.04 MeV OUR 2023 FIT]		
0.88 ± 0.05 OUR AVERAGE					
1.39 ^{+0.40} _{-0.38} ^{+0.26} _{-0.77}			ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
0.876 ± 0.045 ± 0.026			ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
0.87 ± 0.11 ± 0.08	513	¹	ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<1.3	95		BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
<3.8	90		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

¹ Recalculated by ANDREOTTI 05A.

NODE=M055W;LINKAGE=AN

 $\chi_{c1}(1P)$ DECAY MODES

NODE=M055215;NODE=M055

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons		
Γ_2 $e^+ e^-$	$(1.4^{+1.5}_{-1.0}) \times 10^{-7}$	

DESIG=112

DESIG=110

Hadronic decays

NODE=M055;CLUMP=A

Г ₃	$3(\pi^+\pi^-)$	(1.04±0.16) %	S=4.6	DESIG=6
Г ₄	$2(\pi^+\pi^-)$	(7.6 ±2.6) × 10 ⁻³		DESIG=5
Г ₅	$\pi^+\pi^-\pi^0\pi^0$	(1.19±0.15) %		DESIG=51
Г ₆	$\rho^+\pi^-\pi^0 + \text{c.c.}$	(1.45±0.24) %		DESIG=52
Г ₇	$\rho^0\pi^+\pi^-$	(3.9 ±3.5) × 10 ⁻³		DESIG=9
Г ₈	$4\pi^0$	(5.4 ±0.8) × 10 ⁻⁴		DESIG=60
Г ₉	$\pi^+\pi^-K^+K^-$	(4.5 ±1.0) × 10 ⁻³		DESIG=7
Г ₁₀	$K^+K^-\pi^0\pi^0$	(1.12±0.27) × 10 ⁻³		DESIG=53
Г ₁₁	$K^+K^-\pi^+\pi^-\pi^0$	(1.15±0.13) %		DESIG=79
Г ₁₂	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	(7.5 ±0.8) × 10 ⁻³		DESIG=84
Г ₁₃	$K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(8.6 ±1.4) × 10 ⁻³		DESIG=55
Г ₁₄	$\rho^-K^+\bar{K}^0 + \text{c.c.}$	(5.0 ±1.2) × 10 ⁻³		DESIG=56
Г ₁₅	$K^*(892)^0\bar{K}^0\pi^0 \rightarrow$ $K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.}$	(2.3 ±0.6) × 10 ⁻³		DESIG=57
Г ₁₆	$K^+K^-\eta\pi^0$	(1.12±0.34) × 10 ⁻³		DESIG=58
Г ₁₇	$\pi^+\pi^-K_S^0K_S^0$	(6.9 ±2.9) × 10 ⁻⁴		DESIG=28
Г ₁₈	$K^+K^-\eta$	(3.2 ±1.0) × 10 ⁻⁴		DESIG=42
Г ₁₉	$\bar{K}^0K^+\pi^- + \text{c.c.}$	(7.0 ±0.6) × 10 ⁻³	S=1.1	DESIG=17
Г ₂₀	$K^*(892)^0\bar{K}^0 + \text{c.c.}$	(1.03±0.15) × 10 ⁻³		DESIG=32
Г ₂₁	$K^*(892)^+K^- + \text{c.c.}$	(1.21±0.23) × 10 ⁻³		DESIG=33
Г ₂₂	$K_J^*(1430)^0\bar{K}^0 + \text{c.c.} \rightarrow$ $K_S^0K^+\pi^- + \text{c.c.}$	< 8 × 10 ⁻⁴	CL=90%	DESIG=34
Г ₂₃	$K_J^*(1430)^+K^- + \text{c.c.} \rightarrow$ $K_S^0K^+\pi^- + \text{c.c.}$	< 2.1 × 10 ⁻³	CL=90%	DESIG=35
Г ₂₄	$K^+K^-\pi^0$	(1.81±0.24) × 10 ⁻³		DESIG=38
Г ₂₅	$\eta\pi^+\pi^-$	(4.62±0.24) × 10 ⁻³		DESIG=31
Г ₂₆	$a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	(3.2 ±0.4) × 10 ⁻³	S=2.1	DESIG=36
Г ₂₇	$a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	(1.76±0.24) × 10 ⁻⁴		DESIG=93
Г ₂₈	$a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	(4.6 ±0.7) × 10 ⁻⁵		DESIG=96
Г ₂₉	$f_2(1270)\eta \rightarrow \eta\pi^+\pi^-$	(3.5 ±0.6) × 10 ⁻⁴		DESIG=94
Г ₃₀	$f_4(2050)\eta \rightarrow \eta\pi^+\pi^-$	(2.5 ±0.9) × 10 ⁻⁵		DESIG=95
Г ₃₁	$\pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	< 5 × 10 ⁻⁵	CL=90%	DESIG=97
Г ₃₂	$\pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	< 1.5 × 10 ⁻⁵	CL=90%	DESIG=98
Г ₃₃	$\pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-$	< 8 × 10 ⁻⁶	CL=90%	DESIG=99
Г ₃₄	$f_2(1270)\eta$	(6.7 ±1.1) × 10 ⁻⁴		DESIG=37
Г ₃₅	$\pi^+\pi^-\eta'$	(2.2 ±0.4) × 10 ⁻³		DESIG=44
Г ₃₆	$K^+K^-\eta'(958)$	(8.8 ±0.9) × 10 ⁻⁴		DESIG=85
Г ₃₇	$K_0^*(1430)^+K^- + \text{c.c.}$	(6.4 $\begin{smallmatrix} +2.2 \\ -2.8 \end{smallmatrix}$) × 10 ⁻⁴		DESIG=86
Г ₃₈	$K_2^*(1430)^+K^- + \text{c.c.}$	(1.61±0.31) × 10 ⁻³		DESIG=115
Г ₃₉	$K_2^*(1430)\bar{K}^0 + \text{c.c.}$	(1.17±0.20) × 10 ⁻³		DESIG=116
Г ₄₀	$f_0(980)\eta'(958)$	(1.6 $\begin{smallmatrix} +1.4 \\ -0.7 \end{smallmatrix}$) × 10 ⁻⁴		DESIG=87
Г ₄₁	$f_0(1710)\eta'(958)$	(7 $\begin{smallmatrix} +7 \\ -5 \end{smallmatrix}$) × 10 ⁻⁵		DESIG=88
Г ₄₂	$f_2'(1525)\eta'(958)$	(9 ±6) × 10 ⁻⁵		DESIG=89
Г ₄₃	$\pi^0 f_0(980) \rightarrow \pi^0\pi^+\pi^-$	(3.5 ±0.9) × 10 ⁻⁷		DESIG=61
Г ₄₄	$K^+\bar{K}^*(892)^0\pi^- + \text{c.c.}$	(3.2 ±2.1) × 10 ⁻³		DESIG=10
Г ₄₅	$K^*(892)^0\bar{K}^*(892)^0$	(1.4 ±0.4) × 10 ⁻³		DESIG=21
Г ₄₆	$K^+K^-K_S^0K_S^0$	< 4 × 10 ⁻⁴	CL=90%	DESIG=29
Г ₄₇	$K_S^0K_S^0K_S^0K_S^0$	(3.5 ±1.0) × 10 ⁻⁵		DESIG=102
Г ₄₈	$K^+K^-K^+K^-$	(5.4 ±1.1) × 10 ⁻⁴		DESIG=14
Г ₄₉	$K^+K^-\phi$	(4.1 ±1.5) × 10 ⁻⁴		DESIG=30
Г ₅₀	$\bar{K}^0K^+\pi^-\phi + \text{c.c.}$	(3.3 ±0.5) × 10 ⁻³		DESIG=90
Г ₅₁	$K^+K^-\pi^0\phi$	(1.62±0.30) × 10 ⁻³		DESIG=91
Г ₅₂	$\phi\pi^+\pi^-\pi^0$	(7.5 ±1.0) × 10 ⁻⁴		DESIG=82
Г ₅₃	$\omega\omega$	(5.7 ±0.7) × 10 ⁻⁴		DESIG=66
Г ₅₄	ωK^+K^-	(7.8 ±0.9) × 10 ⁻⁴		DESIG=81

Г55	$\omega\phi$	$(2.7 \pm 0.4) \times 10^{-5}$		DESIG=67
Г56	$\phi\phi$	$(4.26 \pm 0.21) \times 10^{-4}$		DESIG=68
Г57	$\phi\phi\eta$	$(3.0 \pm 0.5) \times 10^{-4}$		DESIG=104
Г58	$\rho\bar{\rho}$	$(7.6 \pm 0.4) \times 10^{-5}$	S=1.2	DESIG=11
Г59	$\rho\bar{\rho}\pi^0$	$(1.55 \pm 0.18) \times 10^{-4}$		DESIG=39
Г60	$\rho\bar{\rho}\eta$	$(1.45 \pm 0.25) \times 10^{-4}$		DESIG=43
Г61	$\rho\bar{\rho}\omega$	$(2.12 \pm 0.31) \times 10^{-4}$		DESIG=59
Г62	$\rho\bar{\rho}\phi$	$< 1.7 \times 10^{-5}$	CL=90%	DESIG=65
Г63	$\rho\bar{\rho}\pi^+\pi^-$	$(5.0 \pm 1.9) \times 10^{-4}$		DESIG=8
Г64	$\rho\bar{\rho}\pi^0\pi^0$	$< 5 \times 10^{-4}$	CL=90%	DESIG=54
Г65	$\rho\bar{\rho}K^+K^-$ (non-resonant)	$(1.27 \pm 0.22) \times 10^{-4}$		DESIG=62
Г66	$\rho\bar{\rho}K_S^0 K_S^0$	$< 4.5 \times 10^{-4}$	CL=90%	DESIG=25
Г67	$\rho\bar{n}\pi^-$	$(3.8 \pm 0.5) \times 10^{-4}$		DESIG=74
Г68	$\bar{\rho}n\pi^+$	$(3.9 \pm 0.5) \times 10^{-4}$		DESIG=75
Г69	$\rho\bar{n}\pi^-\pi^0$	$(1.03 \pm 0.12) \times 10^{-3}$		DESIG=76
Г70	$\bar{\rho}n\pi^+\pi^0$	$(1.01 \pm 0.12) \times 10^{-3}$		DESIG=77
Г71	$\Lambda\bar{\Lambda}$	$(1.27 \pm 0.09) \times 10^{-4}$	S=1.1	DESIG=19
Г72	$\Lambda\bar{\Lambda}\pi^+\pi^-$	$(2.9 \pm 0.5) \times 10^{-4}$		DESIG=24
Г73	$\Lambda\bar{\Lambda}\pi^+\pi^-$ (non-resonant)	$(2.5 \pm 0.6) \times 10^{-4}$		DESIG=69
Г74	$\Lambda\bar{\Lambda}\eta$	$(5.9 \pm 1.5) \times 10^{-5}$		DESIG=111
Г75	$\Sigma(1385)^+\bar{\Lambda}\pi^- + c.c.$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=70
Г76	$\Sigma(1385)^-\bar{\Lambda}\pi^+ + c.c.$	$< 1.3 \times 10^{-4}$	CL=90%	DESIG=71
Г77	$K^+\bar{p}\Lambda + c.c.$	$(4.2 \pm 0.4) \times 10^{-4}$	S=1.2	DESIG=40
Г78	$nK_S^0\bar{\Lambda} + c.c.$	$(1.66 \pm 0.17) \times 10^{-4}$		DESIG=109
Г79	$K^*(892)^+\bar{p}\Lambda + c.c.$	$(4.9 \pm 0.7) \times 10^{-4}$		DESIG=106
Г80	$K^+\bar{p}\Lambda(1520) + c.c.$	$(1.7 \pm 0.4) \times 10^{-4}$		DESIG=63
Г81	$\Lambda(1520)\bar{\Lambda}(1520)$	$< 9 \times 10^{-5}$	CL=90%	DESIG=64
Г82	$\Sigma^0\bar{\Sigma}^0$	$(4.2 \pm 0.6) \times 10^{-5}$		DESIG=48
Г83	$\Sigma^+\bar{p}K_S^0 + c.c.$	$(1.53 \pm 0.12) \times 10^{-4}$		DESIG=105
Г84	$\Sigma^0\bar{p}K^+ + c.c.$	$(1.46 \pm 0.10) \times 10^{-4}$		DESIG=108
Г85	$\Sigma^+\bar{\Sigma}^-$	$(3.6 \pm 0.7) \times 10^{-5}$		DESIG=49
Г86	$\Sigma^-\bar{\Sigma}^+$	$(5.7 \pm 1.5) \times 10^{-5}$		DESIG=107
Г87	$\Sigma(1385)^+\bar{\Sigma}(1385)^-$	$< 9 \times 10^{-5}$	CL=90%	DESIG=72
Г88	$\Sigma(1385)^-\bar{\Sigma}(1385)^+$	$< 5 \times 10^{-5}$	CL=90%	DESIG=73
Г89	$K^-\Lambda\bar{\Xi}^+ + c.c.$	$(1.35 \pm 0.24) \times 10^{-4}$		DESIG=92
Г90	$\Xi^0\bar{\Xi}^0$	$(7.5 \pm 1.3) \times 10^{-5}$		DESIG=50
Г91	$\Xi^-\bar{\Xi}^+$	$(6.0 \pm 0.6) \times 10^{-5}$		DESIG=26
Г92	$\Omega^-\bar{\Omega}^+$	$(1.49 \pm 0.25) \times 10^{-5}$		DESIG=113
Г93	$\pi^+\pi^- + K^+K^-$	$< 2.1 \times 10^{-3}$		DESIG=23
Г94	$K_S^0 K_S^0$	$< 6 \times 10^{-5}$	CL=90%	DESIG=27
Г95	$\eta_c\pi^+\pi^-$	$< 3.2 \times 10^{-3}$	CL=90%	DESIG=83
Radiative decays				
Г96	$\gamma J/\psi(1S)$	$(34.3 \pm 1.3) \%$	S=1.3	NODE=M055;CLUMP=B DESIG=1
Г97	$\gamma\rho^0$	$(2.16 \pm 0.17) \times 10^{-4}$		DESIG=45
Г98	$\gamma\omega$	$(6.8 \pm 0.8) \times 10^{-5}$		DESIG=46
Г99	$\gamma\phi$	$(2.4 \pm 0.5) \times 10^{-5}$		DESIG=47
Г100	$\gamma\gamma$	$< 6.3 \times 10^{-6}$	CL=90%	DESIG=4
Г101	$e^+e^- J/\psi(1S)$	$(3.46 \pm 0.24) \times 10^{-3}$		DESIG=100
Г102	$\mu^+\mu^- J/\psi(1S)$	$(2.33 \pm 0.29) \times 10^{-4}$		DESIG=103

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 86 branching ratios uses 253 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 389.6$ for 204 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_{48}	4				
x_{58}	-1	0			
x_{71}	12	5	-1		
x_{96}	20	9	-25	23	
Γ	-9	-4	-60	-11	-31
	x_{19}	x_{48}	x_{58}	x_{71}	x_{96}

$\chi_{c1}(1P)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$					Γ_2
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
$0.12^{+0.13}_{-0.08}$	250	¹ ABLIKIM	22AF BES3	$e^+e^- \rightarrow \chi_{c1} \rightarrow \gamma J/\psi$	NODE=M055220 NODE=M055W1 NODE=M055W1
¹ Assuming $\Gamma(\chi_{c1} \rightarrow \gamma J/\psi) = 0.28$ MeV.					NODE=M055W1;LINKAGE=A

$\chi_{c1}(1P) \Gamma(i)\Gamma(\gamma J/\psi(1S))/\Gamma(\text{total})$

$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$					$\Gamma_{58}\Gamma_{96}/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
21.9 ± 0.8 OUR FIT					NODE=M055G1 NODE=M055G1 NEW
[21.9 ± 0.8 eV OUR 2023 FIT]					
21.4 ± 0.9 OUR AVERAGE					
21.5 ± 0.5 ± 0.8		¹ ANDREOTTI 05A	E835	$p\bar{p} \rightarrow e^+e^-\gamma$	
21.4 ± 1.5 ± 2.2		^{1,2} ARMSTRONG 92	E760	$p\bar{p} \rightarrow e^+e^-\gamma$	
19.9 ^{+4.4} _{-4.0}		¹ BAGLIN 86B	SPEC	$p\bar{p} \rightarrow e^+e^-X$	
¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.					NODE=M055G;LINKAGE=7A
² Recalculated by ANDREOTTI 05A.					NODE=M055G;LINKAGE=AN

$\chi_{c1}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
10.4 ± 1.6 OUR AVERAGE				Error includes scale factor of 4.6. $[(10.5 \pm 1.5) \times 10^{-3}]$	NODE=M055R6 NODE=M055R6 NEW
OUR 2023 AVERAGE Scale factor = 4.6]					
10.92 ± 0.23 ± 0.30	84K	¹ ABLIKIM 22Q	BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+\pi^-)$	
5.4 ± 0.7 ± 0.9		² BAI 99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c1}$	
16.0 ± 5.9 ± 0.8		² TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$	
¹ ABLIKIM 22Q reports $(1.092 \pm 0.004 \pm 0.035) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					NODE=M055R6;LINKAGE=A
² Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.					NODE=M055R;LINKAGE=X2

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

7.6±2.6 OUR EVALUATION Treating systematic error as correlated.NODE=M055R4
NODE=M055R4

→ UNCHECKED ←

8 ±4 OUR AVERAGE Error includes scale factor of 1.5.4.6±2.1±2.6 ¹BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c1}$ 12.5±4.2±0.6 ¹TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$ ¹Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R4;LINKAGE=X2

 $\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.19±0.15±0.03 604.7 ¹HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ NODE=M055R35
NODE=M055R35¹HE 08B reports $1.28 \pm 0.06 \pm 0.15 \pm 0.08\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R35;LINKAGE=HE

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

1.45±0.24±0.04 712.3 ^{1,2}HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ NODE=M055R36
NODE=M055R36¹HE 08B reports $1.56 \pm 0.13 \pm 0.22 \pm 0.10\%$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R36;LINKAGE=HE

²Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M055R36;LINKAGE=OC

 $\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

3.9±3.5 ¹TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$ NODE=M055R8
NODE=M055R8¹Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R;LINKAGE=T

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.4±0.8±0.1 608 ¹ABLIKIM 11A BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$ NODE=M055R44
NODE=M055R44¹ABLIKIM 11A reports $(0.57 \pm 0.03 \pm 0.08) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R44;LINKAGE=AB

 $\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

4.5±1.0 OUR EVALUATION Treating systematic error as correlated.NODE=M055R5
NODE=M055R5

→ UNCHECKED ←

4.5±0.9 OUR AVERAGE4.2±0.4±0.9 ¹BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c1}$ 7.3±3.0±0.4 ¹TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$ ¹Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R5;LINKAGE=X2

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.12±0.27±0.03 45.1 ¹HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$ NODE=M055R37
NODE=M055R37¹HE 08B reports $(0.12 \pm 0.02 \pm 0.02 \pm 0.01) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R37;LINKAGE=HE

$\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
11.46±0.12±1.29	12k	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R00
NODE=M055R00

NODE=M055R00;LINKAGE=A

 $\Gamma(K_S^0K^\pm\pi^\mp\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.52±0.11±0.79	5.1k	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c1}$

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R60
NODE=M055R60

NODE=M055R60;LINKAGE=A

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.13±0.02	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.92 \pm 0.09 \pm 0.11 \pm 0.06$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R39
NODE=M055R39

NODE=M055R39;LINKAGE=HE

 $\Gamma(\rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.50±0.12±0.01	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.54 \pm 0.11 \pm 0.07 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R40
NODE=M055R40

NODE=M055R40;LINKAGE=HE

 $\Gamma(K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.23±0.06±0.01	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.25 \pm 0.06 \pm 0.03 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0\bar{K}^0\pi^0 \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R41
NODE=M055R41

NODE=M055R41;LINKAGE=HE

 $\Gamma(K^+K^-\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.112±0.034±0.003	141.3	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

¹ HE 08B reports $0.12 \pm 0.03 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\eta\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R42
NODE=M055R42

NODE=M055R42;LINKAGE=HE

 $\Gamma(\pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.9±2.9±0.2	19.8 ± 7.7	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1}\gamma$

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ = $(0.67 \pm 0.26 \pm 0.11) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R05
NODE=M055R05

NODE=M055R05;LINKAGE=AB

 $\Gamma(K^+K^-\eta)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.2±1.0±0.1	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(0.34 \pm 0.10 \pm 0.04) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+K^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R25
NODE=M055R25

NODE=M055R25;LINKAGE=AT

$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.03±0.15 OUR AVERAGE[(1.0 ± 0.4) × 10⁻³ OUR 2023 AVERAGE]

1.04±0.13±0.10	262	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
0.98±0.37±0.03	22	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ AAIJ 23AH reports $(1.04 \pm 0.13 \pm 0.04 \pm 0.09) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

² ABLIKIM 06R reports $(1.1 \pm 0.4 \pm 0.1) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R09
NODE=M055R09

NEW

NODE=M055R09;LINKAGE=B

NODE=M055R09;LINKAGE=AB

 $\Gamma(K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.21±0.23 OUR AVERAGE[(1.4 ± 0.6) × 10⁻³ OUR 2023 AVERAGE]

1.18±0.17±0.17	288	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
1.43±0.65±0.04	27	² ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ AAIJ 23AH reports $(1.18 \pm 0.17 \pm 0.14 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

² ABLIKIM 06R reports $(1.6 \pm 0.7 \pm 0.2) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R10
NODE=M055R10

NEW

NODE=M055R10;LINKAGE=C

NODE=M055R10;LINKAGE=AB

 $\Gamma(K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 8 \times 10^{-4}$	90	¹ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
----------------------	----	----------------------	----------	---

¹ ABLIKIM 06R reports $< 0.9 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^0 \bar{K}^0 + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R12
NODE=M055R12

NODE=M055R12;LINKAGE=AB

 $\Gamma(K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 2.1 \times 10^{-3}$	90	¹ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$
------------------------	----	----------------------	----------	---

¹ ABLIKIM 06R reports $< 2.4 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_J^*(1430)^+ K^- + \text{c.c.} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R13
NODE=M055R13

NODE=M055R13;LINKAGE=AB

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.81±0.24±0.05	¹ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
-----------------------	--------------------	---------	---

¹ ATHAR 07 reports $(1.95 \pm 0.16 \pm 0.23) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R20
NODE=M055R20

NODE=M055R20;LINKAGE=AT

 $\Gamma(\eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

4.62±0.24 OUR AVERAGE[(4.62 ± 0.23) × 10⁻³ OUR 2023 AVERAGE]

4.58±0.23±0.13		^{1,2} ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$
4.7 ± 0.5 ± 0.1		³ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$
5.3 ± 0.9 ± 0.1	222	⁴ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R08
NODE=M055R08

NEW

- ¹ From an amplitude analysis using an isobar model.
- ² ABLIKIM 17K reports $(4.67 \pm 0.03 \pm 0.23 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ATHAR 07 reports $(5.0 \pm 0.3 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ⁴ ABLIKIM 06R reports $(5.9 \pm 0.7 \pm 0.8) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R08;LINKAGE=A
 NODE=M055R08;LINKAGE=B

NODE=M055R08;LINKAGE=AT

NODE=M055R08;LINKAGE=AB

$\Gamma(a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.4 OUR AVERAGE				Error includes scale factor of 2.1. $[(3.2 \pm 0.4) \times 10^{-3}$ OUR 2023 AVERAGE Scale factor = 2.2]
3.33±0.19±0.09		^{1,2} ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$
1.79±0.63±0.05	58	³ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R15
 NODE=M055R15

NEW

- ¹ From an amplitude analysis using an isobar model.
- ² ABLIKIM 17K reports $(3.40 \pm 0.03 \pm 0.19 \pm 0.11) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ABLIKIM 06R reports $(2.0 \pm 0.5 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_0(980)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R15;LINKAGE=A
 NODE=M055R15;LINKAGE=B

NODE=M055R15;LINKAGE=AB

$\Gamma(a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.176 ± 0.023 ± 0.005	^{1,2} ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R72
 NODE=M055R72

- ¹ From an amplitude analysis using an isobar model.
- ² ABLIKIM 17K reports $(0.18 \pm 0.01 \pm 0.02 \pm 0.01) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1320)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R72;LINKAGE=A
 NODE=M055R72;LINKAGE=B

$\Gamma(a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.6 ± 0.7 ± 0.1	^{1,2} ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R75
 NODE=M055R75

- ¹ From an amplitude analysis using an isobar model.
- ² ABLIKIM 17K reports $(4.7 \pm 0.4 \pm 0.6 \pm 0.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow a_2(1700)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R75;LINKAGE=A
 NODE=M055R75;LINKAGE=B

$\Gamma(f_2(1270)\eta \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
3.5 ± 0.6 ± 0.1	^{1,2} ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R73
 NODE=M055R73

OCCUR=2

- ¹ From an amplitude analysis using an isobar model.
- ² ABLIKIM 17K reports $(0.36 \pm 0.01 \pm 0.06 \pm 0.01) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R73;LINKAGE=A
 NODE=M055R73;LINKAGE=D

$\Gamma(f_4(2050)\eta \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.5±0.9±0.1	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R74
NODE=M055R74

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $(2.6 \pm 0.4 \pm 0.8 \pm 0.1) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_4(2050)\eta \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R74;LINKAGE=A
NODE=M055R74;LINKAGE=B

 $\Gamma(\pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5 × 10⁻⁵	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R76
NODE=M055R76

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $< 4.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1400)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R76;LINKAGE=A
NODE=M055R76;LINKAGE=B

 $\Gamma(\pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.5 × 10⁻⁵	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R77
NODE=M055R77

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $< 1.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(1600)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R77;LINKAGE=A
NODE=M055R77;LINKAGE=B

 $\Gamma(\pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8 × 10⁻⁶	90	1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$

NODE=M055R78
NODE=M055R78

¹ From an amplitude analysis using an isobar model.

² ABLIKIM 17K reports $< 8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi_1(2015)^+\pi^- + \text{c.c.} \rightarrow \eta\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R78;LINKAGE=A
NODE=M055R78;LINKAGE=B

 $\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.67±0.11 OUR AVERAGE				

NODE=M055R16
NODE=M055R16

0.63±0.11±0.02		1,2 ABLIKIM	17K BES3	$\psi(2S) \rightarrow \gamma\eta\pi^+\pi^-$
2.7 ±0.8 ±0.1	53	³ ABLIKIM	06R BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

¹ ABLIKIM 17K reports $(6.4 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R16;LINKAGE=B

² From an amplitude analysis using an isobar model.

³ ABLIKIM 06R reports $(3.0 \pm 0.7 \pm 0.5) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow f_2(1270)\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R16;LINKAGE=D
NODE=M055R16;LINKAGE=C

 $\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.2±0.4±0.1	1 ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M055R28
NODE=M055R28

¹ ATHAR 07 reports $(2.4 \pm 0.4 \pm 0.3) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.0907 \pm 0.0011 \pm 0.0054$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R28;LINKAGE=AT

$\Gamma(K^+ K^- \eta'(958))/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.75±0.87	310	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R64
NODE=M055R64

¹ Derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.2 \pm 0.4)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.

NODE=M055R64;LINKAGE=A

 $\Gamma(K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.61±0.19±0.24	351	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+ (K_S^0 K \pi)$

NODE=M055R93
NODE=M055R93

¹ AAIJ 23AH reports $(1.61 \pm 0.19 \pm 0.19 \pm 0.14) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

NODE=M055R93;LINKAGE=B

 $\Gamma(K_2^*(1430) \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.17±0.16±0.11	278	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+ (K_S^0 K \pi)$

NODE=M055R94
NODE=M055R94

¹ AAIJ 23AH reports $(1.17 \pm 0.16 \pm 0.05 \pm 0.10) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K_2^*(1430) \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.})]$ assuming $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

NODE=M055R94;LINKAGE=B

 $\Gamma(K_0^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
6.41±0.57^{+2.09}_{-2.71}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R65
NODE=M055R65

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

NODE=M055R65;LINKAGE=A

 $\Gamma(f_0(980) \eta'(958))/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.65±0.47^{+1.32}_{-0.56}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R66
NODE=M055R66

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

NODE=M055R66;LINKAGE=A

 $\Gamma(f_0(1710) \eta'(958))/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.71±0.22^{+0.68}_{-0.48}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R67
NODE=M055R67

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

NODE=M055R67;LINKAGE=A

 $\Gamma(f_2'(1525) \eta'(958))/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.92±0.23^{+0.55}_{-0.51}	¹ ABLIKIM	14J BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$

NODE=M055R68
NODE=M055R68

¹ Normalized to $B(\chi_{c1} \rightarrow K^+ K^- \eta'(958))$ branching fraction.

NODE=M055R68;LINKAGE=A

 $\Gamma(\pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
0.35±0.09		ABLIKIM	18D BES3	$\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

NODE=M055R18
NODE=M055R18

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 ¹ ABLIKIM 11D BES3 $\psi(2S) \rightarrow \gamma \pi^0 \pi^+ \pi^-$

¹ ABLIKIM 11D reports $[\Gamma(\chi_{c1}(1P) \rightarrow \pi^0 f_0(980) \rightarrow \pi^0 \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ < 6.0×10^{-7} which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R18;LINKAGE=BR

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
32±21	¹ TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R9
NODE=M055R9

¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R9;LINKAGE=T

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.44±0.36±0.04	28.4 ± 5.5	^{1,2} ABLIKIM	04H BES	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

NODE=M055R26
 NODE=M055R26

¹ ABLIKIM 04H reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.40 \pm 0.27 \pm 0.22) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R26;LINKAGE=AB

² Assumes $B(K^*(892)^0 \rightarrow K^- \pi^+) = 2/3$.

NODE=M055R26;LINKAGE=AL

 $\Gamma(K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4 × 10⁻⁴	90	3.2 ± 2.4	¹ ABLIKIM	050 BES2	$\psi(2S) \rightarrow \chi_{c1} \gamma$

NODE=M055R06
 NODE=M055R06

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] < 4.2 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R06;LINKAGE=AB

 $\Gamma(K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.35±0.10±0.01	22	¹ ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$

NODE=M055R82
 NODE=M055R82

¹ Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0 K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (3.4 \pm 0.9 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

NODE=M055R82;LINKAGE=A

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.41±0.15±0.01	17	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M055R07
 NODE=M055R07

¹ ABLIKIM 06T reports $(0.46 \pm 0.16 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R07;LINKAGE=AB

 $\Gamma(\bar{K}^0 K^+ \pi^- \phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
3.27±0.28±0.46	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R69
 NODE=M055R69

 $\Gamma(K^+ K^- \pi^0 \phi)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
1.62±0.12±0.28	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R70
 NODE=M055R70

 $\Gamma(\phi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75±0.06±0.08	373	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R62
 NODE=M055R62

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R62;LINKAGE=A

 $\Gamma(\omega \omega)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.7±0.7±0.2	597	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons

NODE=M055R49
 NODE=M055R49

¹ ABLIKIM 11K reports $(6.0 \pm 0.3 \pm 0.7) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R49;LINKAGE=AL

 $\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.78±0.04±0.08	628	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R61
 NODE=M055R61

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$.

NODE=M055R61;LINKAGE=A

$\Gamma(\omega\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
0.27±0.04±0.01	105	¹ ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons

 Γ_{55}/Γ

NODE=M055R50
NODE=M055R50

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.21±0.06±0.01 15 ^{2,3} ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (2.67 \pm 0.31 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R50;LINKAGE=A

² ABLIKIM 11K reports $(0.22 \pm 0.06 \pm 0.02) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R50;LINKAGE=AL

³ Superseded by ABLIKIM 19J.

NODE=M055R50;LINKAGE=B

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
4.26±0.21 OUR AVERAGE				[(4.2 ± 0.5) × 10 ⁻⁴ OUR 2023 AVERAGE]

 Γ_{56}/Γ

NODE=M055R51
NODE=M055R51

4.26±0.17±0.12 1529 ^{1,2} ABLIKIM 23N BES3 $\psi(2S) \rightarrow \gamma$ hadrons

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.2 ± 0.5 ± 0.1 366 ³ ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons

¹ Using $B(\phi \rightarrow K^+ K^-) = (49.2 \pm 0.5) \times 10^{-2}$ from PDG 22.

NODE=M055R51;LINKAGE=C

² ABLIKIM 23N reports $(4.26 \pm 0.13 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R51;LINKAGE=D

³ ABLIKIM 11K reports $(4.4 \pm 0.3 \pm 0.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R51;LINKAGE=AL

 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
3.0±0.5±0.1	83.6	¹ ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma\phi\phi\eta$

 Γ_{57}/Γ

NODE=M055R85
NODE=M055R85

¹ ABLIKIM 20B reports $(2.96 \pm 0.43 \pm 0.22) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R85;LINKAGE=A

 $\Gamma(\rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.155±0.018 OUR AVERAGE			
0.163±0.019±0.004	¹ ONYISI 10	CLE3	$\psi(2S) \rightarrow \gamma\rho\bar{\rho}X$
0.112±0.047±0.003	² ATHAR 07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

 Γ_{59}/Γ

NODE=M055R21
NODE=M055R21

¹ ONYISI 10 reports $(1.75 \pm 0.16 \pm 0.13 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R21;LINKAGE=ON

² ATHAR 07 reports $(1.2 \pm 0.5 \pm 0.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \rho\bar{\rho}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R21;LINKAGE=AT

 $\Gamma(\rho\bar{\rho}\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
0.145±0.024±0.004		¹ ONYISI 10	CLE3	$\psi(2S) \rightarrow \gamma\rho\bar{\rho}X$

 Γ_{60}/Γ

NODE=M055R27
NODE=M055R27

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.15 90 ² ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

- ¹ ONYISI 10 reports $(1.56 \pm 0.22 \pm 0.14 \pm 0.10) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² ATHAR 07 reports $< 0.16 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R27;LINKAGE=ON

NODE=M055R27;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.212±0.030±0.006	¹ ONYISI 10	CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$

NODE=M055R43
NODE=M055R43

- ¹ ONYISI 10 reports $(2.28 \pm 0.28 \pm 0.16 \pm 0.14) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R43;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7 × 10⁻⁵	90	¹ ABLIKIM 11F	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M055R48
NODE=M055R48

- ¹ ABLIKIM 11F reports $< 1.82 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R48;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.50±0.19 OUR EVALUATION	Treating systematic error as correlated.		
0.50±0.19 OUR AVERAGE			

NODE=M055R7
NODE=M055R7

→ UNCHECKED ←

0.46±0.12±0.15

¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c1}$

1.08±0.77±0.05

¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$

- ¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (8.8 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$.

NODE=M055R7;LINKAGE=X2

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5 × 10⁻⁴	90	¹ HE 08B	CLEO	$e^+e^- \rightarrow \gamma h^+h^-h^0h^0$

NODE=M055R38
NODE=M055R38

- ¹ HE 08B reports $< 0.05 \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R38;LINKAGE=HE

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.27±0.22±0.04	82 ± 9	¹ ABLIKIM 11F	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NODE=M055R45
NODE=M055R45

- ¹ ABLIKIM 11F reports $(1.35 \pm 0.15 \pm 0.19) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R45;LINKAGE=AB

 $\Gamma(p\bar{p}K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	¹ ABLIKIM 06D	BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$

NODE=M055R02
NODE=M055R02

- ¹ Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) = (9.1 \pm 0.6)\%$.

NODE=M055R;LINKAGE=AB

 $\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8±0.5±0.1	1412	¹ ABLIKIM 12J	BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$

NODE=M055R56
NODE=M055R56

- ¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.37 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R56;LINKAGE=AL

$\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±0.5±0.1	1625	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$

NODE=M055R57
 NODE=M055R57

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.38 \pm 0.02 \pm 0.04) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R57;LINKAGE=AL

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.3±1.1±0.3	1082	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$

NODE=M055R58
 NODE=M055R58

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (1.00 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R58;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.1±1.1±0.3	1261	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$

NODE=M055R59
 NODE=M055R59

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c1}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))] = (0.98 \pm 0.05 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R59;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
29±5±1		105	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M055R01
 NODE=M055R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

<150	90	² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \gamma\chi_{c1}$
------	----	----------------------	----------	--

¹ ABLIKIM 12I reports $(31.1 \pm 3.4 \pm 3.9) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R01;LINKAGE=AL

² Using $B(\psi(2S) \rightarrow \chi_{c1}\gamma) (9.1 \pm 0.6)\%$.

NODE=M055R01;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
25±6±1	13	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$

NODE=M055R19
 NODE=M055R19

¹ ABLIKIM 12I reports $(26.2 \pm 5.5 \pm 3.3) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R19;LINKAGE=AL

 $\Gamma(\Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10⁻⁴	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^+\bar{\Lambda}\pi^-$

NODE=M055R52
 NODE=M055R52

¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+\bar{\Lambda}\pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R52;LINKAGE=AL

 $\Gamma(\Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<13	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Sigma(1385)^-\bar{\Lambda}\pi^+$

NODE=M055R53
 NODE=M055R53

¹ ABLIKIM 12I reports $< 14 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^-\bar{\Lambda}\pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R53;LINKAGE=AL

$\Gamma(K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{π} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R22
 NODE=M055R22

4.2 ± 0.4 OUR AVERAGE Error includes scale factor of 1.2. $[(4.2 \pm 0.4) \times 10^{-4}]$ OUR 2023 AVERAGE Scale factor = 1.2]

NEW

$9.2^{+2.8}_{-2.4} \pm 0.4$	24	¹ LU	19	BELL	$B^+ \rightarrow \bar{p} \Lambda K^+ K^+$
$4.2 \pm 0.4 \pm 0.1$	3k	^{2,3} ABLIKIM	13D	BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$
$3.1 \pm 0.9 \pm 0.1$		⁴ ATHAR	07	CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ LU 19 reports $(9.15^{+2.63}_{-2.25} \pm 0.86) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(1P) K^+)]$ assuming $B(B^+ \rightarrow \chi_{c1}(1P) K^+) = (4.79 \pm 0.23) \times 10^{-4}$, which we rescale to our best value $B(B^+ \rightarrow \chi_{c1}(1P) K^+) = (4.74 \pm 0.22) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R22;LINKAGE=A

² ABLIKIM 13D reports $(4.5 \pm 0.2 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R22;LINKAGE=AB

³ Using $B(\Lambda \rightarrow p \pi^-) = 63.9\%$.

NODE=M055R22;LINKAGE=LB

⁴ ATHAR 07 reports $(3.3 \pm 0.9 \pm 0.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R22;LINKAGE=AT

 $\Gamma(n K_S^0 \bar{\Lambda} + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{78} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R89
 NODE=M055R89

1.66 ± 0.16 ± 0.05 399 ¹ ABLIKIM 21AV BES3 $\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$

¹ ABLIKIM 21AV reports $(1.66 \pm 0.12 \pm 0.12) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow n K_S^0 \bar{\Lambda} + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.0975 \pm 0.0024$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Also uses $B(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$.

NODE=M055R89;LINKAGE=A

 $\Gamma(K^*(892)^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{79} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R86
 NODE=M055R86

4.9 ± 0.7 ± 0.1 328 ¹ ABLIKIM 19AU BES3 $\psi(2S) \rightarrow \gamma K^{*+} \bar{p} \Lambda$

¹ ABLIKIM 19AU reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^*(892)^+ \bar{p} \Lambda + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (4.8 \pm 0.5 \pm 0.4) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R86;LINKAGE=F

 $\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{80} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R46
 NODE=M055R46

1.71 ± 0.44 ± 0.05 48 ± 10 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports $(1.81 \pm 0.38 \pm 0.28) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.}) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R46;LINKAGE=AB

 $\Gamma(\Lambda(1520) \bar{\Lambda}(1520)) / \Gamma_{\text{total}}$ Γ_{81} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M055R47
 NODE=M055R47

< 9 × 10⁻⁵ 90 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports $< 1.00 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Lambda(1520) \bar{\Lambda}(1520)) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R47;LINKAGE=AB

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 0.6 \pm 0.1$		103	¹ ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

NODE=M055R32
 NODE=M055R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$
<4	90	3.8 ± 2.5	³ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.41 \pm 0.05 \pm 0.03) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R32;LINKAGE=B

² ABLIKIM 13H reports $< 0.62 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R32;LINKAGE=AB

³ NAIK 08 reports $< 0.44 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R32;LINKAGE=NA

 $\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$3.6 \pm 0.6 \pm 0.1$		59	¹ ABLIKIM	18V BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

NODE=M055R33
 NODE=M055R33

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8	90		² ABLIKIM	13H BES3	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$
<6	90	4.3 ± 2.3	³ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (0.35 \pm 0.06 \pm 0.02) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R33;LINKAGE=B

² ABLIKIM 13H reports $< 0.87 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R33;LINKAGE=AB

³ NAIK 08 reports $< 0.65 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R33;LINKAGE=NA

 $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	COMMENT
$5.7 \pm 1.5 \pm 0.2$	214	¹ ABLIKIM	20I BES3	$\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

NODE=M055R87
 NODE=M055R87

¹ ABLIKIM 20I reports $(5.7 \pm 1.4 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R87;LINKAGE=A

 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-5}$	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M055R54
 NODE=M055R54

¹ ABLIKIM 12I reports $< 10 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R54;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5 \times 10^{-5}$	90	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{\Lambda} \pi^+ \pi^-$

NODE=M055R55
 NODE=M055R55

¹ ABLIKIM 12I reports $< 5.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R55;LINKAGE=AL

$\Gamma(K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.35±0.24±0.04	49	¹ ABLIKIM	15i	BES3 $\psi(2S) \rightarrow \gamma K^- \Lambda \Xi^+ + \text{c.c.}$

NODE=M055R71
 NODE=M055R71

¹ ABLIKIM 15i reports $[\Gamma(\chi_{c1}(1P) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))] = (1.32 \pm 0.20 \pm 0.12) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R71;LINKAGE=A

 $\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.5±1.2±0.2		325	¹ ABLIKIM	220	BES3 $\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$

NODE=M055R34
 NODE=M055R34

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 1.7 ± 2.4 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Xi^0 \Xi^0$

¹ ABLIKIM 220 reports $(0.75 \pm 0.11 \pm 0.06) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R34;LINKAGE=A

² NAIK 08 reports $< 0.60 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^0 \Xi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R34;LINKAGE=NA

 $\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.60±0.06 OUR AVERAGE					

NODE=M055R03
 NODE=M055R03

0.58±0.06±0.02 692 ¹ ABLIKIM 220 BES3 $\psi(2S) \rightarrow \gamma \Xi^- \Xi^+$

0.80±0.21±0.02 16.4 ± 4.3 ² NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Xi^+ \Xi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.4 90 ³ ABLIKIM 06D BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ ABLIKIM 220 reports $(0.58 \pm 0.04 \pm 0.05) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R03;LINKAGE=A

² NAIK 08 reports $(0.86 \pm 0.22 \pm 0.08) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \Xi^- \Xi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R03;LINKAGE=NA

³ Using $B(\psi(2S) \rightarrow \chi_{c1} \gamma) (9.1 \pm 0.6)\%$.

NODE=M055R03;LINKAGE=AB

 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.49±0.23±0.10	277	ABLIKIM	23T	BES3 $\chi_{cJ} \rightarrow \Omega^- \bar{\Omega}^+$

NODE=M055R91
 NODE=M055R91

 $[\Gamma(\pi^+ \pi^-) + \Gamma(K^+ K^-)]/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<21 × 10⁻⁴		¹ FELDMAN	77	MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

NODE=M055R2
 NODE=M055R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<38 × 10⁻⁴ 90 ¹ BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

NODE=M055R2;LINKAGE=T

 $\Gamma(K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6 × 10⁻⁵		¹ ABLIKIM	050	BES2 $\psi(2S) \rightarrow \chi_{c1} \gamma$

NODE=M055R04
 NODE=M055R04

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c1}(1P) \rightarrow K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ $< 0.6 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.

NODE=M055R04;LINKAGE=AB

$\Gamma(\eta_c \pi^+ \pi^-) / \Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{95} / Γ
$< 3.2 \times 10^{-3}$	90	1,2 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$	NODE=M055R63 NODE=M055R63
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$< 4.4 \times 10^{-3}$	90	1,3 ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c1}$	OCCUR=2
¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c1} \gamma) = (9.2 \pm 0.4)\%$. ² Using the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays. ³ Using the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.					

RADIATIVE DECAYS

 $\Gamma(\gamma \rho^0) / \Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{97} / Γ
216 ± 17 OUR AVERAGE					
215 ± 22 ± 6	432 ± 25	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$	NODE=M055R29 NODE=M055R29
217 ± 24 ± 6	186 ± 15	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$	NODE=M055R29;LINKAGE=AB
¹ ABLIKIM 11E reports $(228 \pm 13 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \rho^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ² BENNETT 08A reports $(243 \pm 19 \pm 22) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \rho^0) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\gamma \omega) / \Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{98} / Γ
68 ± 8 OUR AVERAGE					
66 ± 9 ± 2	136 ± 14	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \omega$	NODE=M055R30 NODE=M055R30
74 ± 17 ± 2	39 ± 7	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \omega$	NODE=M055R30;LINKAGE=AB
¹ ABLIKIM 11E reports $(69.7 \pm 7.2 \pm 6.6) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \omega) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ² BENNETT 08A reports $(83 \pm 15 \pm 12) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \omega) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					

 $\Gamma(\gamma \phi) / \Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{99} / Γ
24 ± 5 ± 1						
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 23	90	5.2 ± 3.1	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \phi$	NODE=M055R31 NODE=M055R31
¹ ABLIKIM 11E reports $(25.8 \pm 5.2 \pm 2.3) \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \phi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.2 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ² BENNETT 08A reports $< 26 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow \gamma \phi) / \Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 9.75 \times 10^{-2}$.						

 $\Gamma(\gamma \gamma) / \Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{100} / Γ
< 6.3 × 10⁻⁶					
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 3.5 × 10 ⁻⁵	90	ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1} \rightarrow 3\gamma$	NODE=M055R3 NODE=M055R3
< 150 × 10 ⁻⁵	90	¹ YAMADA	77 DASP	$e^+ e^- \rightarrow 3\gamma$	NODE=M055R;LINKAGE=T1
¹ Estimated using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.					

$\Gamma(e^+e^- J/\psi(1S))/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R79
NODE=M055R79

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.65±0.23±0.10	1.9k	^{1,2} ABLIKIM	17I	BES3	$\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$
----------------	------	------------------------	-----	------	--

¹ ABLIKIM 17I reports $(3.73 \pm 0.09 \pm 0.25) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c1}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M055R79;LINKAGE=B

² Not independent from other measurements reported by ABLIKIM 17I

NODE=M055R79;LINKAGE=C

 $\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S))$ Γ_{101}/Γ_{96}

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R80
NODE=M055R80

10.1±0.3±0.5	1.9k	¹ ABLIKIM	17I	BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$
---------------------	------	----------------------	-----	------	--

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) \times B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (351.8 \pm 1.0 \pm 12.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

NODE=M055R80;LINKAGE=A

 $\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S))$ $\Gamma_{102}/\Gamma_{101}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055R84
NODE=M055R84

6.73±0.51±0.50	222	ABLIKIM	19Z	BES3	$\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma(\mu^+\mu^- J/\psi)$
-----------------------	-----	---------	-----	------	--

 $\chi_{c1}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M055230

 $\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$ $\Gamma_{58}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M055B1
NODE=M055B1

2.13±0.13 OUR FIT	Error includes scale factor of 1.3. [(2.14 ± 0.10) × 10 ⁻⁵ OUR 2023 FIT]		
--------------------------	---	--	--

NEW

1.1 ± 1.0	¹ BAI	98I	BES	$\psi(2S) \rightarrow \gamma \chi_{c1} \rightarrow \gamma p\bar{p}$
------------------	------------------	-----	-----	---

¹ Calculated by us. The value for $B(\chi_{c1} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M055B;LINKAGE=J2

 $\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma_{\text{total}}$ $\Gamma_{71}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055B10
NODE=M055B10

12.4±0.9 OUR FIT	Error includes scale factor of 1.1. [(12.4 ± 0.8) × 10 ⁻⁶ OUR 2023 FIT]		
-------------------------	--	--	--

NEW

12.3±0.9 OUR AVERAGE Error includes scale factor of 1.2.

12.8±0.6±0.6	528	ABLIKIM	21L	BES3	$\psi(2S) \rightarrow \gamma p\pi^- \bar{p}\pi^+$
10.5±1.6±0.6	46	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

11.2±1.0±0.9	136	^{2,3} ABLIKIM	13H	BES3	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$
--------------	-----	------------------------	-----	------	--

¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (11.6 \pm 1.8 \pm 0.7 \pm 0.7) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

NODE=M055B10;LINKAGE=NA

² Superseded by ABLIKIM 21L

NODE=M055B10;LINKAGE=A

³ Calculated by us. ABLIKIM 13H reports $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) = (12.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ from a measurement of $B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma \chi_{c1})$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (9.2 \pm 0.4)\%$.

NODE=M055B10;LINKAGE=AB

 $\Gamma(\chi_{c1}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$ $\Gamma_{71}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M055B11
NODE=M055B11

3.58±0.25 OUR FIT	Error includes scale factor of 1.1. [(3.58 ± 0.22) × 10 ⁻⁵ OUR 2023 FIT]		
--------------------------	---	--	--

NEW

7.1 $^{+2.8}_{-2.4}$ ± 1.3	$9.0^{+3.5}_{-3.1}$	¹ BAI	03E	BES	$\psi(2S) \rightarrow \gamma \Lambda\bar{\Lambda}$
--	---------------------	------------------	-----	-----	--

¹ BAI 03E reports $[B(\chi_{c1} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma \chi_{c1}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.52}_{-0.46} \pm 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

NODE=M055B11;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Lambda \bar{\Lambda} \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{74} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10 ⁻⁶)	EVTS	DOCUMENT ID	TECN	COMMENT
5.72 ± 1.34 ± 0.65	21	ABLIKIM	22AO BES3	$\psi(2S) \rightarrow \gamma p \pi^- \bar{p} \pi^+ \gamma \gamma$

NODE=M055R90
NODE=M055R90

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{96} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
3.34 ± 0.10 OUR FIT				Error includes scale factor of 1.7. [(3.34 ± 0.06) × 10 ⁻² OUR 2023 FIT]

NODE=M055B2
NODE=M055B2
NEW

3.24 ± 0.16 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

3.518 ± 0.010 ± 0.120	143k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$
3.442 ± 0.010 ± 0.132	1.9M	ABLIKIM	17U BES3	$e^+ e^- \rightarrow \gamma X$
2.81 ± 0.05 ± 0.23	13k	BAI	04I BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
2.56 ± 0.12 ± 0.20		GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
2.78 ± 0.30		² OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma \chi_{c1}$
2.2 ± 0.5		³ BRANDELIK	79B DASP	$\psi(2S) \rightarrow \gamma \chi_{c1}$
2.9 ± 0.5		³ BARTEL	78B CNTR	$\psi(2S) \rightarrow \gamma \chi_{c1}$
5.0 ± 1.5		⁴ BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$
2.8 ± 0.9		² WHITAKER	76 MRK1	$e^+ e^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.377 ± 0.009 ± 0.183	142k	⁵ ABLIKIM	12O BES3	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3.56 ± 0.03 ± 0.12	24.9k	⁶ MENDEZ	08 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c1}$
3.44 ± 0.06 ± 0.13	3.7k	⁷ ADAM	05A CLEO	Repl. by MENDEZ 08

NODE=M055B2;LINKAGE=A

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

NODE=M055B;LINKAGE=3Q

NODE=M055B;LINKAGE=2Q


NODE=M055B;LINKAGE=EA

NODE=M055B2;LINKAGE=B

NODE=M055B2;LINKAGE=ME

NODE=M055B;LINKAGE=AD

WEIGHTED AVERAGE
3.24 ± 0.16 (Error scaled by 2.1)



$$\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}} \text{ (units } 10^{-2})$$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{96} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

9.63 ± 0.29 OUR FIT Error includes scale factor of 1.7. [(9.63 ± 0.17) × 10^{-2} OUR 2023 FIT]

NODE=M055B3
NODE=M055B3

NEW

10.15 ± 0.28 OUR AVERAGE

10.17 ± 0.07 ± 0.27 24.9k MENDEZ 08 CLEO $\psi(2S) \rightarrow \gamma \chi_{c1}$
 12.6 ± 0.3 ± 3.8 3k 1 ABLIKIM 04B BES $\psi(2S) \rightarrow J/\psi X$
 8.5 ± 2.1 2 HIMEL 80 MRK2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.24 ± 0.17 ± 0.23 3.7k 3 ADAM 05A CLEO Repl. by MENDEZ 08

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c1}) \times B(\chi_{c1} \rightarrow \gamma J/\psi(1S))$ quoted in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

³ Not independent from other values reported by ADAM 05A.

NODE=M055B;LINKAGE=AB
NODE=M055B;LINKAGE=J3

NODE=M055B3;LINKAGE=AD

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{19} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

6.8 ± 0.6 OUR FIT Error includes scale factor of 1.1. [(6.8 ± 0.5) × 10^{-4} OUR 2023 FIT]

NODE=M055B16
NODE=M055B16

NEW

7.2 ± 0.6 OUR AVERAGE

7.3 ± 0.5 ± 0.5 1 ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$
 7.0 ± 0.5 ± 0.9 2 ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma \chi_{c1}$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^- + \text{c.c.})$ reported by ATHAR 07 was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (9.07 \pm 0.11 \pm 0.54)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-) = (4.0 \pm 0.3 \pm 0.5) \times 10^{-3}$. We use $B(\psi(2S) \rightarrow \gamma \chi_{c1}) = (8.7 \pm 0.4) \times 10^{-2}$.

NODE=M055B16;LINKAGE=AT

NODE=M055B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{19} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

19.6 ± 1.6 OUR FIT Error includes scale factor of 1.1. [(19.6 ± 1.6) × 10^{-4} OUR 2023 FIT]

NODE=M055B17
NODE=M055B17

NEW

13.2 ± 2.4 ± 3.2

1 BAI 99B BES $\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow K_S^0 K^+ \pi^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M055B17;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma_{\text{total}}}{\Gamma_{48} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

0.53 ± 0.11 OUR FIT

NODE=M055B14
NODE=M055B14

0.61 ± 0.11 ± 0.08 54 1 ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$.

NODE=M055B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{48} / \Gamma \times \Gamma_{180}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT

1.52 ± 0.31 OUR FIT

NODE=M055B15
NODE=M055B15

1.13 ± 0.40 ± 0.29 1 BAI 99B BES $\psi(2S) \rightarrow \gamma K^+ K^+ K^- K^-$

¹ Calculated by us. The value of $B(\chi_{c1} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c1}(1P)) = (8.7 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M055B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{58}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT
7.4±0.4 OUR FIT Error includes scale factor of 1.3. $[(7.41 \pm 0.35) \times 10^{-6}$ OUR 2023 FIT]

NODE=M055B6
 NODE=M055B6

NEW


7.8±0.6 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.9 \pm 0.4 \pm 0.3$	453	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
$8.2 \pm 0.7 \pm 0.4$	141 ± 13	¹ NAIK	08 CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
$4.8^{+1.4}_{-1.3} \pm 0.6$	$18.2^{+5.5}_{-4.9}$	BAI	04F BES	$\psi(2S) \rightarrow \gamma\chi_{c1}(1P) \rightarrow \gamma p\bar{p}$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c1} \rightarrow p\bar{p}) = (9.0 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.07 \pm 0.11 \pm 0.54)\%$.

NODE=M055B6;LINKAGE=NA

WEIGHTED AVERAGE
 7.8 ± 0.6 (Error scaled by 1.4)



$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{58}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{\psi(2S)}} \text{ (units } 10^{-6}\text{)}$$

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{83}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT
1.49±0.09±0.07 258 ¹ ABLIKIM 19BB BES3 $\psi(2S) \rightarrow \gamma\Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

NODE=M055B01
 NODE=M055B01

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_{c1} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (1.53 \pm 0.10 \pm 0.08) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = (9.75 \pm 0.24)\%$ and other branching fractions from PDG 18.

NODE=M055B01;LINKAGE=A

$$\frac{\Gamma(\chi_{c1}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))/\Gamma_{\text{total}}}{\Gamma_{84}/\Gamma \times \Gamma_{180}^{\psi(2S)}/\Gamma_{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT
1.42±0.07±0.06 493 ¹ ABLIKIM 20AE BES3 $\psi(2S) \rightarrow \gamma\Sigma^0 \bar{p} K^+ + \text{c.c.}$

NODE=M055R88
 NODE=M055R88

¹ Calculated by us. ABLIKIM 20AE reports $B(\chi_{c1} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (1.46 \pm 0.07 \pm 0.07) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c1}^0) = (9.75 \pm 0.24)\%$ and other branching fractions from PDG 20.

NODE=M055R88;LINKAGE=A

MULTIPOLE AMPLITUDES IN $\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$

NODE=M055240

$a_2 = M_2/\sqrt{E_1^2 + M_2^2}$ Magnetic quadrupole fractional transition amplitude


NODE=M055A1
 NODE=M055A1

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT
-6.7 ± 0.9 OUR AVERAGE Error includes scale factor of 2.6. See the ideogram below.

$-7.40 \pm 0.33 \pm 0.34$	164k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
$-6.26 \pm 0.63 \pm 0.24$	39k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
$0.2 \pm 3.2 \pm 0.4$	2090	AMBROGIANI	02 E835	$p\bar{p} \rightarrow \chi_{c1} \rightarrow J/\psi\gamma$
$-0.2^{+0.8}_{-2.0}$	921	OREGLIA	82 CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$

¹ Correlated with b_2 with correlation coefficient $\rho_{a_2 b_2} = 0.133$.

NODE=M055A1;LINKAGE=A



MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$ RADIATIVE DECAY

NODE=M055250

$b_2 = M2/\sqrt{E1^2 + M2^2}$ Magnetic quadrupole fractional transition amplitude

NODE=M055QB2
NODE=M055QB2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.5 ± 0.4 OUR AVERAGE				
2.29 ± 0.39 ± 0.27	164k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.76 ± 0.73 ± 0.23	39k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
7.7 ^{+5.0} _{-4.5}	921	OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Correlated with a_2 with correlation coefficient $\rho_{a_2 b_2} = 0.133$.

NODE=M055QB2;LINKAGE=A

**MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS
 $\psi(2S) \rightarrow \gamma\chi_{c1}(1S)$ and $\chi_{c1} \rightarrow \gamma J/\psi(1S)$**

NODE=M055260

a_2/b_2 Magnetic quadrupole transition amplitude ratio

NODE=M055QAR
NODE=M055QAR

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-2.27^{+0.57}_{-0.99}	39k	¹ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

¹ Statistical and systematic errors combined. Not independent of $a_2(\chi_{c1})$ and $b_2(\chi_{c1})$ values from ARTUSO 09.

NODE=M055QAR;LINKAGE=AR

$\chi_{c1}(1P)$ REFERENCES

NODE=M055

AAIJ	23AH	PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62349
ABLIKIM	23N	JHEP 2305 069	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62056
ABLIKIM	23T	PR D107 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62064
ABLIKIM	22AF	PRL 129 122001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61878
ABLIKIM	22AO	PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61887
ABLIKIM	22O	JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61652
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)	REFID=61634
ABLIKIM	21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61465
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61117
ABLIKIM	20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60733
ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60212
ABLIKIM	20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60303
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59844
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60026
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59606
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59837
LU	19	PR D99 032003	P.-C. Lu <i>et al.</i>	(BELLE Collab.)	REFID=59614
ABLIKIM	18D	PRL 121 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58849
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58990
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
AAIJ	17BI	PRL 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58278
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58310

ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM	17K	PR D95 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57953
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56778
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55901
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11D	PR D83 032003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16715
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48552
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41865
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41907
BAGLIN	86B	PL B172 455	C. Baglin (LAPP, CERN, GENO, LYON, OSLO+)	(Crystal Ball Collab.)	REFID=22145
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
Also		Private Comm.	M.J. Oreglia	(EFI)	REFID=22143
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)	REFID=22064
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151
TANENBAUM	75	PRL 35 1323	W.M. Tanenbaum <i>et al.</i>	(LBL, SLAC)	REFID=22106

$h_c(1P)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Quantum numbers are quark model prediction, $C = -$ established by $\eta_c \gamma$ decay.

NODE=M144

NODE=M144

 $h_c(1P)$ MASS

NODE=M144M

NODE=M144M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3525.37±0.14 OUR AVERAGE		Error includes scale factor of 1.2.		
3525.32±0.06±0.15	23k	ABLIKIM	22AQ BES3	$\psi(2S) \rightarrow \pi^0$ hadrons; $\pi^0 \gamma(\eta_c)$
3525.20±0.18±0.12	1282	¹ DOBBS	08A CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3525.8 ±0.2 ±0.2	13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3525.31±0.11±0.14	832	^{2,3} ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
3525.40±0.13±0.18	3679	² ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
3525.6 ±0.5	92	ADAMS	09 CLEO	$\psi(2S) \rightarrow 2(\pi^+ \pi^- \pi^0)$
3524.4 ±0.6 ±0.4	168	⁴ ROSNER	05 CLEO	$\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
3527 ±8	42	ANTONIAZZI	94 E705	$300 \pi^\pm, pLi \rightarrow$ $J/\psi \pi^0 X$
3526.28±0.18±0.19	59	⁵ ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$
3525.4 ±0.8 ±0.4	5	BAGLIN	86 SPEC	$\bar{p}p \rightarrow J/\psi X$

¹ Combination of exclusive and inclusive analyses for the reaction $\psi(2S) \rightarrow \pi^0 h_c \rightarrow \pi^0 \eta_c \gamma$. This result is the average of DOBBS 08A and ROSNER 05.

² Superseded by ABLIKIM 22AQ

³ With floating width.

⁴ Superseded by DOBBS 08A.

⁵ Mass central value and systematic error recalculated by us according to Eq.(16) in ARMSTRONG 93B, using the value for the $\psi(2S)$ mass from AULCHENKO 03.

NODE=M144M;LINKAGE=DO

NODE=M144M;LINKAGE=A

NODE=M144M;LINKAGE=AB

NODE=M144M;LINKAGE=RO

NODE=M144M;LINKAGE=NW

 $h_c(1P)$ WIDTH

NODE=M144W

NODE=M144W

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.78^{+0.27}_{-0.24}±0.12		23k	ABLIKIM	22AQ BES3	$\psi(2S) \rightarrow \pi^0$ hadrons; $\pi^0 \gamma(\eta_c)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.70±0.28±0.22		832	^{1,2} ABLIKIM	12N BES3	$\psi(2S) \rightarrow \pi^0 \gamma$ hadrons
< 1.44	90	3679	³ ABLIKIM	10B BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
< 1		13	ANDREOTTI	05B E835	$\bar{p}p \rightarrow \eta_c \gamma$
< 1.1	90	59	ARMSTRONG	92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$

¹ Superseded by ABLIKIM 22AQ

² With floating mass.

³ The central value is $\Gamma = 0.73 \pm 0.45 \pm 0.28$ MeV.

NODE=M144W;LINKAGE=A

NODE=M144W;LINKAGE=AL

NODE=M144W;LINKAGE=AB

 $h_c(1P)$ DECAY MODES

NODE=M144215;NODE=M144

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S)\pi^0$	< 5 × 10 ⁻⁴	90%
Γ_2 $J/\psi(1S)\pi\pi$	not seen	
Γ_3 $J/\psi(1S)\pi^+\pi^-$	< 2.7 × 10 ⁻³	90%
Γ_4 $p\bar{p}$	< 1.7 × 10 ⁻⁴	90%
Γ_5 $p\bar{p}\pi^0$	< 8 × 10 ⁻⁴	90%
Γ_6 $p\bar{p}\pi^+\pi^-$	(3.3±0.6) × 10 ⁻³	
Γ_7 $p\bar{p}\pi^0\pi^0$	< 6 × 10 ⁻⁴	90%
Γ_8 $p\bar{p}\pi^+\pi^-\pi^0$	(4.4±1.3) × 10 ⁻³	
Γ_9 $p\bar{p}\eta$	(7.4±2.2) × 10 ⁻⁴	
Γ_{10} $\pi^+\pi^-\pi^0$	(1.9±0.5) × 10 ⁻³	
Γ_{11} $\pi^+\pi^-\pi^0\eta$	(8.3±2.4) × 10 ⁻³	

DESIG=1

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=10

DESIG=3

DESIG=24

DESIG=11

DESIG=13

DESIG=25

DESIG=23

DESIG=5

DESIG=14

Γ_{12}	$2\pi^+ 2\pi^- \pi^0$	$(9.4 \pm 1.7) \times 10^{-3}$			DESIG=6
Γ_{13}	$3\pi^+ 3\pi^- \pi^0$	< 1.0	%	90%	DESIG=7
Γ_{14}	$K^+ K^- \pi^+ \pi^-$	< 7	$\times 10^{-4}$	90%	DESIG=12
Γ_{15}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.8 \pm 0.8) \times 10^{-3}$			DESIG=15
Γ_{16}	$K^+ K^- \pi^+ \pi^- \eta$	< 2.7	$\times 10^{-3}$	90%	DESIG=16
Γ_{17}	$K^+ K^- \pi^0$	< 6	$\times 10^{-4}$	90%	DESIG=17
Γ_{18}	$K^+ K^- \pi^0 \eta$	< 2.4	$\times 10^{-3}$	90%	DESIG=18
Γ_{19}	$K^+ K^- \eta$	< 1.0	$\times 10^{-3}$	90%	DESIG=19
Γ_{20}	$2K^+ 2K^- \pi^0$	< 2.8	$\times 10^{-4}$	90%	DESIG=20
Γ_{21}	$K_S^0 K^\pm \pi^\mp$	< 6	$\times 10^{-4}$	90%	DESIG=21
Γ_{22}	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(3.2 \pm 1.0) \times 10^{-3}$			DESIG=22

Radiative decays

Γ_{23}	$\gamma\eta$	$(4.7 \pm 2.1) \times 10^{-4}$			NODE=M144;CLUMP=R DESIG=9
Γ_{24}	$\gamma\eta'(958)$	$(1.5 \pm 0.4) \times 10^{-3}$			DESIG=8
Γ_{25}	$\gamma\eta_c(1S)$	$(60 \pm 4) \%$			DESIG=4

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

$h_c(1P)$ PARTIAL WIDTHS

NODE=M144220

$h_c(1P) \Gamma(i)\Gamma(\bar{p}p)/\Gamma(\text{total})$

NODE=M144223

$\Gamma(\gamma\eta_c(1S)) \times \Gamma(p\bar{p})/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_4/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
12.0 ± 4.5	13	¹ ANDREOTTI 05B E835	$\bar{p}p \rightarrow \eta_c \gamma$	

NODE=M144G1
NODE=M144G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.0 ± 4.5 13 ¹ ANDREOTTI 05B E835 $\bar{p}p \rightarrow \eta_c \gamma$

¹ Assuming $\Gamma = 1$ MeV.

NODE=M144G1;LINKAGE=AN

$h_c(1P)$ BRANCHING RATIOS

NODE=M144225

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(\gamma\eta_c(1S))$ Γ_1/Γ_{25}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-4}$	90	¹ ABLIKIM 22N BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$	

NODE=M144R24
NODE=M144R24

¹ ABLIKIM 22N reports $[\Gamma(h_c(1P) \rightarrow J/\psi(1S)\pi^0)/\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))]/[B(\eta_c \rightarrow K^+ K^- \pi^0)] < 7.5 \times 10^{-2}$ which we multiply by our best value $B(\eta_c \rightarrow K^+ K^- \pi^0) = 1/6 B(\eta_c(1S) \rightarrow K\bar{K}\pi) = 1/6 (7.1 \times 10^{-2})$.

NODE=M144R24;LINKAGE=B

$\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.18	90	ARMSTRONG 92D E760	$\bar{p}p \rightarrow J/\psi \pi^0$	

NODE=M144R1
NODE=M144R1

$\Gamma(J/\psi(1S)\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.7 \times 10^{-3}$	90	¹ ABLIKIM 18M BES3	$\psi(2S) \rightarrow \pi^0 \pi^+ \pi^- J/\psi$	

NODE=M144R07
NODE=M144R07

OCCUR=2

¹ ABLIKIM 18M reports $[\Gamma(h_c(1P) \rightarrow J/\psi(1S)\pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.0 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R07;LINKAGE=B

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.7 \times 10^{-4}$	90	¹ ABLIKIM 13V BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$	

NODE=M144R20
NODE=M144R20

¹ ABLIKIM 13V reports $[\Gamma(h_c(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 1.3 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R20;LINKAGE=A

$\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8 \times 10^{-4}$	90	¹ ABLIKIM	22M BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R22
 NODE=M144R22

¹ ABLIKIM 22M reports $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 5.67 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R22;LINKAGE=A

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.3 \pm 0.5 \pm 0.2$	230	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R08
 NODE=M144R08

¹ ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (2.49 \pm 0.27 \pm 0.28) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R08;LINKAGE=A

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-4}$	90	12	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R10
 NODE=M144R10

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 4.4 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R10;LINKAGE=A

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.4 \pm 1.2 \pm 0.3$	86	¹ ABLIKIM	22M BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R23
 NODE=M144R23

¹ ABLIKIM 22M reports $[\Gamma(h_c(1P) \rightarrow p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (3.30 \pm 0.71 \pm 0.59) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R23;LINKAGE=A

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.4 \pm 2.1 \pm 0.5$	20	¹ ABLIKIM	22M BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R21
 NODE=M144R21

¹ ABLIKIM 22M reports $[\Gamma(h_c(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (5.51 \pm 1.50 \pm 0.46) \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R21;LINKAGE=A

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.5 \pm 0.1$		101	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R01
 NODE=M144R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.6	90	² ADAMS	09 CLEO	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
--------	----	--------------------	---------	--

¹ ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (1.38 \pm 0.35 \pm 0.17) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R01;LINKAGE=A

² ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 0.19 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R01;LINKAGE=A

 $\Gamma(\pi^+\pi^-\pi^0\eta)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$8.3 \pm 2.3 \pm 0.6$	35	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R11
 NODE=M144R11

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow \pi^+\pi^-\pi^0\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (6.2 \pm 1.6 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R11;LINKAGE=A

 $\Gamma(2\pi^+2\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.94 ± 0.17 OUR AVERAGE				

NODE=M144R02
 NODE=M144R02

$0.86 \pm 0.16 \pm 0.06$	254	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$
--------------------------	-----	----------------------	-----------	--------------------------------------

$2.5^{+0.9}_{-0.7} \pm 0.2$	92	² ADAMS	09 CLEO	$\psi(2S) \rightarrow \pi^0 \gamma \eta_c$
-----------------------------	----	--------------------	---------	--

¹ ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow 2\pi^+ 2\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (6.40 \pm 0.81 \pm 0.87) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R02;LINKAGE=A

² ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 2\pi^+ 2\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (1.88^{+0.48+0.47}_{-0.45-0.30}) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R02;LINKAGE=AD

 $\Gamma(3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}$ **Γ_{13}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.010	90	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R03
NODE=M144R03

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.034 90 ² ADAMS 09 CLEO $\psi(2S) \rightarrow \pi^0 \gamma \eta_c$

¹ ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 7.5 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R03;LINKAGE=A

² ADAMS 09 reports $[\Gamma(h_c(1P) \rightarrow 3\pi^+ 3\pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R03;LINKAGE=AD

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7 × 10⁻⁴	90	¹ ABLIKIM	19AG BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R09
NODE=M144R09

¹ ABLIKIM 19AG reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 0.5 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R09;LINKAGE=A

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ **Γ_{15}/Γ**

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.8 ± 0.3	80	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R12
NODE=M144R12

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (2.8 \pm 0.5 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R12;LINKAGE=A

 $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ **Γ_{16}/Γ**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.7 × 10⁻³	90	24	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R13
NODE=M144R13

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.0 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R13;LINKAGE=A

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ **Γ_{17}/Γ**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<6 × 10⁻⁴	90	20	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R14
NODE=M144R14

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 4.8 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R14;LINKAGE=A

 $\Gamma(K^+ K^- \pi^0 \eta)/\Gamma_{\text{total}}$ **Γ_{18}/Γ**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.4 × 10⁻³	90	20	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R15
NODE=M144R15

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \pi^0 \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 1.8 \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R15;LINKAGE=A

 $\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ **Γ_{19}/Γ**

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.0 × 10⁻³	90	18	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R16
NODE=M144R16

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 7.5 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R16;LINKAGE=A

$\Gamma(2K^+2K^-\pi^0)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-4}$	90	11	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R17
 NODE=M144R17

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow 2K^+2K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 2.1 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R17;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-4}$	90	17	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R18
 NODE=M144R18

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K_S^0 K^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] < 4.8 \times 10^{-7}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = 7.4 \times 10^{-4}$.

NODE=M144R18;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.2 \pm 1.0 \pm 0.2$	41	¹ ABLIKIM	20AH BES3	$\psi(2S) \rightarrow \pi^0 h_c(1P)$

NODE=M144R19
 NODE=M144R19

¹ ABLIKIM 20AH reports $[\Gamma(h_c(1P) \rightarrow K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (2.4 \pm 0.7 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R19;LINKAGE=A

————— RADIATIVE DECAYS —————

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 1.5 \pm 1.4$	18	ABLIKIM	16I BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta$

NODE=M144230

NODE=M144R06
 NODE=M144R06

 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.52 \pm 0.27 \pm 0.29$	44	ABLIKIM	16I BES3	$\psi(2S) \rightarrow \pi^0 \gamma \eta'(958)$

NODE=M144R00
 NODE=M144R00

 $\Gamma(\gamma\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M144R2
 NODE=M144R2

60 ± 4 OUR FIT**57 ± 5 OUR AVERAGE**

57 ± 4 ± 4 23k ¹ ABLIKIM 22AQ BES3 $\psi(2S) \rightarrow \pi^0$ hadrons;
 $\pi^0 \gamma(\eta_c)$

56 ± 6 ± 4 ² DOBBS 08A CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

62 ± 9 ± 4 3679 ^{3,4} ABLIKIM 10B BES3 $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

56 ± 7 ± 4 1282 ⁵ DOBBS 08A CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

54 ± 14 ± 4 168 ⁶ ROSNER 05 CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

¹ ABLIKIM 22AQ reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.22_{-0.26}^{+0.27} \pm 0.19) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=C

² Average of DOBBS 08A and ROSNER 05. DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.16 \pm 0.30 \pm 0.37) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=DB

³ ABLIKIM 10B reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.58 \pm 0.40 \pm 0.50) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=A

⁴ Superseded by ABLIKIM 22AQ

NODE=M144R2;LINKAGE=B

⁵ DOBBS 08A reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.19 \pm 0.32 \pm 0.45) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=DO

⁶ ROSNER 05 reports $[\Gamma(h_c(1P) \rightarrow \gamma\eta_c(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow h_c(1P)\pi^0)] = (4.0 \pm 0.8 \pm 0.7) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow h_c(1P)\pi^0) = (7.4 \pm 0.5) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M144R2;LINKAGE=RO

$h_c(1P)$ REFERENCES

ABLIKIM	22AQ	PR D106 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61889
ABLIKIM	22M	JHEP 2205 108	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61650
	Also	JHEP 2303 022 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62053
ABLIKIM	22N	JHEP 2205 003	M. Ablikim	(BESIII Collab.)	REFID=61651
ABLIKIM	20AH	PR D102 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60750
ABLIKIM	19AG	PR D99 072008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59858
ABLIKIM	18M	PR D97 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58901
ABLIKIM	16I	PRL 116 251802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57450
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
ABLIKIM	12N	PR D86 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54741
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53348
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
ANDREOTTI	05B	PR D72 032001	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50768
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ARMSTRONG	92D	PRL 69 2337	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43174
BAGLIN	86	PL B171 135	C. Baglin <i>et al.</i>	(LAPP, CERN, TORI, STRB+)	REFID=43180

NODE=M144

 $\chi_{c2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See the Review on “Branching Ratios of $\psi(2S)$, $\chi_{c0,1,2}$ and $\eta_c(1S)$ ” before the $\chi_{c0}(1P)$ Listings.

NODE=M057

NODE=M057

 $\chi_{c2}(1P)$ MASS

NODE=M057M

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
3556.17 ± 0.07	OUR AVERAGE			
3557.3 ± 1.7 ± 0.7	611	¹ AAIJ	17BB LHCB	$pp \rightarrow b\bar{b}X \rightarrow 2(K^+ K^-)X$
3556.10 ± 0.06 ± 0.11	4.0k	² AAIJ	17BI LHCB	$\chi_{c2} \rightarrow J/\psi \mu^+ \mu^-$
3555.3 ± 0.6 ± 2.2	2.5k	UEHARA	08 BELL	$\gamma\gamma \rightarrow \text{hadrons}$
3555.70 ± 0.59 ± 0.39		ABLIKIM	05G BES2	$\psi(2S) \rightarrow \gamma\chi_{c2}$
3556.173 ± 0.123 ± 0.020		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
3559.9 ± 2.9		EISENSTEIN	01 CLE2	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
3556.4 ± 0.7		BAI	99B BES	$\psi(2S) \rightarrow \gamma X$
3556.22 ± 0.131 ± 0.020	585	³ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
3556.9 ± 0.4 ± 0.5	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
3557.8 ± 0.2 ± 4		⁴ GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3553.4 ± 2.2	66	⁵ LEMOIGNE	82 GOLJ	$185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- A$
3555.9 ± 0.7		⁶ OREGLIA	82 CBAL	$e^+ e^- \rightarrow J/\psi 2\gamma$
3557 ± 1.5	69	⁷ HIMEL	80 MRK2	$e^+ e^- \rightarrow J/\psi 2\gamma$
3551 ± 11	15	BRANDELIK	79B DASP	$e^+ e^- \rightarrow J/\psi 2\gamma$
3553 ± 4		⁷ BARTEL	78B CNTR	$e^+ e^- \rightarrow J/\psi 2\gamma$
3553 ± 4 ± 4		^{7,8} TANENBAUM	78 MRK1	$e^+ e^-$
3563 ± 7	360	⁷ BIDDICK	77 CNTR	$e^+ e^- \rightarrow \gamma X$
3555.4 ± 1.3	53	UEHARA	13 BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$
3543 ± 10	4	WHITAKER	76 MRK1	$e^+ e^- \rightarrow J/\psi 2\gamma$

NODE=M057M

NODE=M057M

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a fit of the $\phi\phi$ invariant mass with the width of $\chi_{c2}(1P)$ fixed to the PDG 16 value.

² AAIJ 17BI reports also $m(\chi_{c2}) - m(\chi_{c1}) = 45.39 \pm 0.07 \pm 0.03$ MeV.

³ Recalculated by ANDREOTTI 05A, using the value of $\psi(2S)$ mass from AULCHENKO 03.

⁴ Using mass of $\psi(2S) = 3686.0$ MeV.

⁵ $J/\psi(1S)$ mass constrained to 3097 MeV.

⁶ Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁷ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁸ From a simultaneous fit to radiative and hadronic decay channels.

NODE=M057M;LINKAGE=A

NODE=M057M;LINKAGE=B

NODE=M057M;LINKAGE=NW

NODE=M057M;LINKAGE=C

NODE=M057M;LINKAGE=P

NODE=M057M;LINKAGE=E

NODE=M057M;LINKAGE=D

NODE=M057M;LINKAGE=M

 $\chi_{c2}(1P)$ WIDTH

NODE=M057W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.98 ±0.09 OUR FIT	Error includes scale factor of 1.1.		[1.97 ± 0.09 MeV OUR 2023 FIT]	
2.00 ±0.11 OUR AVERAGE				
2.10 ±0.20 ±0.02	4.0k	AAIJ	17BI LHCB	$\chi_{c2} \rightarrow J/\psi \mu^+ \mu^-$
1.915 ±0.188 ±0.013		ANDREOTTI	05A E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
1.96 ±0.17 ±0.07	585	¹ ARMSTRONG	92 E760	$\bar{p}p \rightarrow e^+ e^- \gamma$
2.6 ^{+1.4} _{-1.0}	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+ e^- X$
2.8 ^{+2.1} _{-2.0}		² GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$

¹Recalculated by ANDREOTTI 05A.

²Errors correspond to 90% confidence level; authors give only width range.

NODE=M057W

NEW

NODE=M057W;LINKAGE=AN

NODE=M057W;LINKAGE=E

$\chi_{c2}(1P)$ DECAY MODES

NODE=M057215;NODE=M057

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic decays		
Γ_1 $2(\pi^+ \pi^-)$	(1.00 ± 0.13) %	S=1.4
Γ_2 $\rho\rho$		
Γ_3 $\pi^+ \pi^- \pi^0 \pi^0$	(1.86 ± 0.24) %	
Γ_4 $\rho^+ \pi^- \pi^0 + \text{c.c.}$	(2.22 ± 0.35) %	
Γ_5 $4\pi^0$	(1.13 ± 0.15) × 10 ⁻³	
Γ_6 $K^+ K^- \pi^0 \pi^0$	(2.1 ± 0.4) × 10 ⁻³	
Γ_7 $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(1.41 ± 0.20) %	
Γ_8 $\rho^- K^+ \bar{K}^0 + \text{c.c.}$	(4.2 ± 1.3) × 10 ⁻³	
Γ_9 $K^*(892)^0 K^- \pi^+ \rightarrow$ $K^- \pi^+ K^0 \pi^0 + \text{c.c.}$	(3.0 ± 0.8) × 10 ⁻³	
Γ_{10} $K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(3.9 ± 0.9) × 10 ⁻³	
Γ_{11} $K^*(892)^- K^+ \pi^0 \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(3.8 ± 0.8) × 10 ⁻³	
Γ_{12} $K^*(892)^+ \bar{K}^0 \pi^- \rightarrow$ $K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.}$	(3.0 ± 0.8) × 10 ⁻³	
Γ_{13} $K^+ K^- \eta \pi^0$	(1.3 ± 0.4) × 10 ⁻³	
Γ_{14} $K^+ K^- \pi^+ \pi^-$	(8.3 ± 1.1) × 10 ⁻³	S=1.2
Γ_{15} $K^+ K^- \pi^+ \pi^- \pi^0$	(1.17 ± 0.13) %	
Γ_{16} $K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	(7.3 ± 0.8) × 10 ⁻³	
Γ_{17} $K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	(2.1 ± 1.0) × 10 ⁻³	
Γ_{18} $K^*(892)^0 \bar{K}^*(892)^0$	(2.2 ± 0.9) × 10 ⁻³	S=2.3
Γ_{19} $3(\pi^+ \pi^-)$	(1.53 ± 0.19) %	S=3.8
Γ_{20} $\phi\phi$	(1.23 ± 0.07) × 10 ⁻³	S=1.9
Γ_{21} $\phi\phi\eta$	(5.4 ± 0.7) × 10 ⁻⁴	
Γ_{22} $\omega\omega$	(8.6 ± 1.0) × 10 ⁻⁴	
Γ_{23} $\omega K^+ K^-$	(7.3 ± 0.9) × 10 ⁻⁴	
Γ_{24} $\omega\phi$	(9.7 ± 2.8) × 10 ⁻⁶	
Γ_{25} $\pi\pi$	(2.27 ± 0.10) × 10 ⁻³	
Γ_{26} $\rho^0 \pi^+ \pi^-$	(3.6 ± 1.5) × 10 ⁻³	
Γ_{27} $\pi^+ \pi^- \pi^0$ (non-resonant)	(2.0 ± 0.4) × 10 ⁻⁵	
Γ_{28} $\rho(770)^\pm \pi^\mp$	(6 ± 4) × 10 ⁻⁶	
Γ_{29} $\pi^+ \pi^- \eta$	(4.9 ± 1.3) × 10 ⁻⁴	
Γ_{30} $\pi^+ \pi^- \eta'$	(5.1 ± 1.9) × 10 ⁻⁴	
Γ_{31} $\eta\eta$	(5.5 ± 0.5) × 10 ⁻⁴	
Γ_{32} $K^+ K^-$	(1.02 ± 0.15) × 10 ⁻³	S=2.3
Γ_{33} $K_S^0 K_S^0$	(5.3 ± 0.4) × 10 ⁻⁴	
Γ_{34} $K^*(892)^\pm K^\mp$	(1.46 ± 0.21) × 10 ⁻⁴	
Γ_{35} $K^*(892)^0 \bar{K}^0 + \text{c.c.}$	(1.27 ± 0.27) × 10 ⁻⁴	
Γ_{36} $K_2^*(1430)^\pm K^\mp$	(1.51 ± 0.13) × 10 ⁻³	
Γ_{37} $K_2^*(1430)^0 \bar{K}^0 + \text{c.c.}$	(1.27 ± 0.17) × 10 ⁻³	
Γ_{38} $K_3^*(1780)^\pm K^\mp$	(5.3 ± 0.8) × 10 ⁻⁴	
Γ_{39} $K_3^*(1780)^0 \bar{K}^0 + \text{c.c.}$	(5.7 ± 2.1) × 10 ⁻⁴	
Γ_{40} $a_2(1320)^0 \pi^0$	(1.31 ± 0.35) × 10 ⁻³	

NODE=M057;CLUMP=A

DESIG=3

DESIG=43

DESIG=50

DESIG=51

DESIG=62

DESIG=52

DESIG=54

DESIG=55

DESIG=60

DESIG=56

DESIG=57

DESIG=58

DESIG=59

DESIG=5

DESIG=67

DESIG=78

DESIG=10

DESIG=21

DESIG=4

DESIG=16

DESIG=99

DESIG=25

DESIG=79

DESIG=68

DESIG=22

DESIG=9

DESIG=95

DESIG=96

DESIG=39

DESIG=42

DESIG=14

DESIG=2

DESIG=15

DESIG=87

DESIG=88

DESIG=89

DESIG=90

DESIG=91

DESIG=92

DESIG=93

Γ ₄₁	$a_2(1320)^\pm \pi^\mp$	$(1.8 \pm 0.6) \times 10^{-3}$		DESIG=94
Γ ₄₂	$\bar{K}^0 K^+ \pi^- + \text{c.c.}$	$(1.30 \pm 0.19) \times 10^{-3}$		DESIG=17
Γ ₄₃	$K^+ K^- \pi^0$	$(3.1 \pm 0.8) \times 10^{-4}$		DESIG=36
Γ ₄₄	$K^+ K^- \eta$	$< 3.3 \times 10^{-4}$	CL=90%	DESIG=40
Γ ₄₅	$K^+ K^- \eta'(958)$	$(1.94 \pm 0.34) \times 10^{-4}$		DESIG=82
Γ ₄₆	$\eta \eta'$	$(2.2 \pm 0.5) \times 10^{-5}$		DESIG=34
Γ ₄₇	$\eta' \eta'$	$(4.6 \pm 0.6) \times 10^{-5}$		DESIG=35
Γ ₄₈	$\pi^+ \pi^- K_S^0 K_S^0$	$(2.2 \pm 0.5) \times 10^{-3}$		DESIG=29
Γ ₄₉	$K^+ K^- K_S^0 K_S^0$	$< 4 \times 10^{-4}$	CL=90%	DESIG=30
Γ ₅₀	$K_S^0 K_S^0 K_S^0 K_S^0$	$(1.15 \pm 0.18) \times 10^{-4}$		DESIG=97
Γ ₅₁	$K^+ K^- K^+ K^-$	$(1.67 \pm 0.22) \times 10^{-3}$	S=1.1	DESIG=24
Γ ₅₂	$K^+ K^- \phi$	$(1.45 \pm 0.30) \times 10^{-3}$		DESIG=32
Γ ₅₃	$\bar{K}^0 K^+ \pi^- \phi + \text{c.c.}$	$(4.8 \pm 0.7) \times 10^{-3}$		DESIG=83
Γ ₅₄	$K^+ K^- \pi^0 \phi$	$(2.7 \pm 0.5) \times 10^{-3}$		DESIG=84
Γ ₅₅	$\phi \pi^+ \pi^- \pi^0$	$(9.3 \pm 1.2) \times 10^{-4}$		DESIG=80
Γ ₅₆	$p \bar{p}$	$(7.3 \pm 0.4) \times 10^{-5}$	S=1.1	DESIG=11
Γ ₅₇	$p \bar{p} \pi^0$	$(4.7 \pm 0.4) \times 10^{-4}$		DESIG=37
Γ ₅₈	$p \bar{p} \eta$	$(1.77 \pm 0.25) \times 10^{-4}$		DESIG=41
Γ ₅₉	$p \bar{p} \omega$	$(3.7 \pm 0.4) \times 10^{-4}$		DESIG=61
Γ ₆₀	$p \bar{p} \phi$	$(2.8 \pm 0.9) \times 10^{-5}$		DESIG=66
Γ ₆₁	$p \bar{p} \pi^+ \pi^-$	$(1.32 \pm 0.34) \times 10^{-3}$		DESIG=8
Γ ₆₂	$p \bar{p} \pi^0 \pi^0$	$(8.0 \pm 2.4) \times 10^{-4}$		DESIG=53
Γ ₆₃	$p \bar{p} K^+ K^- (\text{non-resonant})$	$(1.94 \pm 0.33) \times 10^{-4}$		DESIG=63
Γ ₆₄	$p \bar{p} K_S^0 K_S^0$	$< 7.9 \times 10^{-4}$	CL=90%	DESIG=28
Γ ₆₅	$p \bar{n} \pi^-$	$(8.7 \pm 1.0) \times 10^{-4}$		DESIG=31
Γ ₆₆	$\bar{p} n \pi^+$	$(9.1 \pm 0.8) \times 10^{-4}$		DESIG=75
Γ ₆₇	$p \bar{n} \pi^- \pi^0$	$(2.21 \pm 0.18) \times 10^{-3}$		DESIG=76
Γ ₆₈	$\bar{p} n \pi^+ \pi^0$	$(2.15 \pm 0.19) \times 10^{-3}$		DESIG=77
Γ ₆₉	$\Lambda \bar{\Lambda}$	$(1.86 \pm 0.16) \times 10^{-4}$		DESIG=19
Γ ₇₀	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(1.28 \pm 0.16) \times 10^{-3}$		DESIG=27
Γ ₇₁	$\Lambda \bar{\Lambda} \pi^+ \pi^- (\text{non-resonant})$	$(6.7 \pm 1.5) \times 10^{-4}$		DESIG=70
Γ ₇₂	$\Lambda \bar{\Lambda} \eta$	$(1.07 \pm 0.26) \times 10^{-4}$		DESIG=105
Γ ₇₃	$\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.}$	$< 4 \times 10^{-4}$	CL=90%	DESIG=71
Γ ₇₄	$\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.}$	$< 6 \times 10^{-4}$	CL=90%	DESIG=72
Γ ₇₅	$K^+ \bar{p} \Lambda + \text{c.c.}$	$(7.9 \pm 0.6) \times 10^{-4}$		DESIG=38
Γ ₇₆	$n K_S^0 \bar{\Lambda} + \text{c.c.}$	$(3.64 \pm 0.29) \times 10^{-4}$		DESIG=104
Γ ₇₇	$K^*(892)^+ \bar{p} \Lambda + \text{c.c.}$	$(8.3 \pm 1.2) \times 10^{-4}$		DESIG=101
Γ ₇₈	$K^+ \bar{p} \Lambda(1520) + \text{c.c.}$	$(2.9 \pm 0.7) \times 10^{-4}$		DESIG=64
Γ ₇₉	$\Lambda(1520) \bar{\Lambda}(1520)$	$(4.7 \pm 1.5) \times 10^{-4}$		DESIG=65
Γ ₈₀	$\Sigma^0 \bar{\Sigma}^0$	$(3.7 \pm 0.6) \times 10^{-5}$		DESIG=47
Γ ₈₁	$\Sigma^+ \bar{p} K_S^0 + \text{c.c.}$	$(8.4 \pm 1.0) \times 10^{-5}$		DESIG=100
Γ ₈₂	$\Sigma^0 \bar{p} K^+ + \text{c.c.}$	$(9.3 \pm 0.8) \times 10^{-5}$		DESIG=103
Γ ₈₃	$\Sigma^+ \bar{\Sigma}^-$	$(3.4 \pm 0.7) \times 10^{-5}$		DESIG=48
Γ ₈₄	$\Sigma^- \bar{\Sigma}^+$	$(4.5 \pm 1.8) \times 10^{-5}$		DESIG=102
Γ ₈₅	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$< 1.6 \times 10^{-4}$	CL=90%	DESIG=73
Γ ₈₆	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$< 8 \times 10^{-5}$	CL=90%	DESIG=74
Γ ₈₇	$K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$	$(1.80 \pm 0.32) \times 10^{-4}$		DESIG=85
Γ ₈₈	$\Xi^0 \bar{\Xi}^0$	$(1.86 \pm 0.22) \times 10^{-4}$		DESIG=49
Γ ₈₉	$\Xi^- \bar{\Xi}^+$	$(1.46 \pm 0.12) \times 10^{-4}$		DESIG=26
Γ ₉₀	$\Omega^- \bar{\Omega}^+$	$(4.52 \pm 0.30) \times 10^{-5}$		DESIG=106
Γ ₉₁	$J/\psi(1S) \pi^+ \pi^- \pi^0$	$< 1.5 \%$	CL=90%	DESIG=12
Γ ₉₂	$\pi^0 \eta_c$	$< 3.2 \times 10^{-3}$	CL=90%	DESIG=81
Γ ₉₃	$\eta_c(1S) \pi^+ \pi^-$	$< 5.4 \times 10^{-3}$	CL=90%	DESIG=69
Radiative decays				
Γ ₉₄	$\gamma J/\psi(1S)$	$(19.5 \pm 0.8) \%$	S=1.5	NODE=M057;CLUMP=B DESIG=6
Γ ₉₅	$\gamma \rho^0$	$< 1.9 \times 10^{-5}$	CL=90%	DESIG=44
Γ ₉₆	$\gamma \omega$	$< 6 \times 10^{-6}$	CL=90%	DESIG=45
Γ ₉₇	$\gamma \phi$	$< 8 \times 10^{-6}$	CL=90%	DESIG=46
Γ ₉₈	$\gamma \gamma$	$(2.92 \pm 0.12) \times 10^{-4}$	S=1.3	DESIG=7
Γ ₉₉	$e^+ e^- J/\psi(1S)$	$(2.20 \pm 0.15) \times 10^{-3}$		DESIG=86
Γ ₁₀₀	$\mu^+ \mu^- J/\psi(1S)$	$(2.07 \pm 0.34) \times 10^{-4}$		DESIG=98

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 86 branching ratios uses 253 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 389.6$ for 204 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x14	7									
x17	2	26								
x18	2	2	0							
x20	4	4	1	1						
x25	4	4	1	2	5					
x26	26	2	1	1	1	1				
x31	2	2	0	1	3	16	1			
x32	3	3	1	1	2	7	1	3		
x33	3	3	1	1	3	16	1	9	4	
x42	1	1	0	1	2	8	0	5	2	5
x51	3	3	1	1	3	9	1	5	3	5
x56	6	6	1	2	4	12	2	6	4	7
x69	2	2	0	1	3	15	0	8	3	8
x94	12	12	3	3	9	25	4	13	8	15
x98	-10	-10	-3	-1	-3	19	-3	12	1	9
Γ	-18	-18	-5	-5	-12	-26	-6	-13	-10	-16
	x1	x14	x17	x18	x20	x25	x26	x31	x32	x33
x51	3									
x56	4	5								
x69	4	4	6							
x94	7	10	-12	12						
x98	6	2	27	11	11					
Γ	-8	-12	-42	-13	-48	-47				
	x42	x51	x56	x69	x94	x98				

$\chi_{c2}(1P)$ PARTIAL WIDTHS

$$\chi_{c2}(1P) \Gamma(i) \Gamma(\gamma J/\psi(1S)) / \Gamma(\text{total})$$

NODE=M057220

NODE=M057223

$$\Gamma(p\bar{p}) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{56} \Gamma_{94} / \Gamma$$

NODE=M057G1
NODE=M057G1

VALUE (eV) DOCUMENT ID TECN COMMENT

28.0 ± 1.3 OUR FIT Error includes scale factor of 1.1. [27.5 ± 1.2 eV OUR 2023 FIT]

NEW

27.5 ± 1.5 OUR AVERAGE

27.0 ± 1.5 ± 1.1	¹ ANDREOTTI	05A	E835	$p\bar{p} \rightarrow e^+ e^- \gamma$
27.7 ± 1.5 ± 2.0	^{1,2} ARMSTRONG	92	E760	$p\bar{p} \rightarrow e^+ e^- \gamma$
36 ± 8	¹ BAGLIN	86B	SPEC	$p\bar{p} \rightarrow e^+ e^- X$

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

² Recalculated by ANDREOTTI 05A.

NODE=M057G;LINKAGE=7A
NODE=M057G;LINKAGE=AN

$$\Gamma(\gamma\gamma) \times \Gamma(\gamma J/\psi(1S)) / \Gamma_{\text{total}} \quad \Gamma_{98} \Gamma_{94} / \Gamma$$

NODE=M057G2
NODE=M057G2

VALUE (eV) EVTS DOCUMENT ID TECN COMMENT

113 ± 5 OUR FIT Error includes scale factor of 1.3. [107 ± 5 eV OUR 2023 FIT]

NEW

123 ± 6 OUR AVERAGE

NEW

[117 ± 10 eV OUR 2023 AVERAGE]

124.1 ± 2.5 ± 5.9	4960	¹ SEINO	23	BELL	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
111 ± 12 ± 9	147	² DOBBS	06	CLE3	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
139 ± 55 ± 21		^{2,3} ACCIARRI	99E	L3	$e^+ e^- \rightarrow e^+ e^- \chi_{c2}$

242 ± 65 ± 51	2,4 ACKER.,K...	98	OPAL	$e^+e^- \rightarrow e^+e^- \chi_{c2}$
150 ± 42 ± 36	2,5 DOMINICK	94	CLE2	$e^+e^- \rightarrow e^+e^- \chi_{c2}$
470 ± 240 ± 120	2,6 BAUER	93	TPC	$e^+e^- \rightarrow e^+e^- \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

114 ± 11 ± 9	136	2,7 ABE	02T BELL	$e^+e^- \rightarrow e^+e^- \chi_{c2}$
--------------	-----	---------	----------	---------------------------------------

¹ Calculated from the measured $\Gamma_{\gamma\gamma} \times B(\chi_{c2}(1S) \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 14.8 \pm 0.3 \pm 0.7$ eV, using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 11.93 \pm 0.05\%$.

² Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1187 \pm 0.0008$.

³ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACCIARRI 99E is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) \times B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.0162 \pm 0.0014$.

⁴ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in ACKERSTAFF,K 98 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1203 \pm 0.0038$.

⁵ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in DOMINICK 94 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

⁶ The value for $\Gamma(\chi_{c2} \rightarrow \gamma\gamma)$ reported in BAUER 93 is derived using $B(\chi_{c2} \rightarrow \gamma J/\psi(1S)) = 0.135 \pm 0.011$, $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0627 \pm 0.0020$, and $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0597 \pm 0.0025$.

⁷ All systematic errors added in quadrature. Superseded by SEINO 23.

NODE=M057G2;LINKAGE=A

NODE=M057G;LINKAGE=LL

NODE=M057G;LINKAGE=J4

NODE=M057G;LINKAGE=J5

NODE=M057G;LINKAGE=J6

NODE=M057G;LINKAGE=J7

NODE=M057G;LINKAGE=GT

NODE=M057224

NODE=M057G3
NODE=M057G3

NEW

$\chi_{c2}(1P) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$				
$\Gamma(2(\pi^+\pi^-)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$			$\Gamma_1\Gamma_{98}/\Gamma$	
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
5.8 ± 0.7 OUR FIT		Error includes scale factor of 1.4.	[5.7 ± 0.5 eV OUR 2023 FIT]	
5.2 ± 0.7 OUR AVERAGE				
5.01 ± 0.44 ± 0.55	1597 ± 138	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$
6.4 ± 1.8 ± 0.8		EISENSTEIN	01 CLE2	$e^+e^- \rightarrow e^+e^- \chi_{c2}$

NODE=M057G08
NODE=M057G08

$\Gamma(\rho\rho) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					
VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<7.8		90 <598	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+\pi^-)$

$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.8 ± 0.6 OUR FIT		Error includes scale factor of 1.2.	[4.7 ± 0.5 eV OUR 2023 FIT]	
4.42 ± 0.42 ± 0.53	780 ± 74	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M057G09
NODE=M057G09

NEW

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.5 ± 0.9 ± 1.5	1250	DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M057G02
NODE=M057G02

$\Gamma(K^*(892)^0 \bar{K}^*(892)^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 ± 0.5 OUR FIT		Error includes scale factor of 2.3.	[1.26 ± 0.24 eV OUR 2023 FIT]	
0.8 ± 0.17 ± 0.27	151 ± 30	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow K^+K^-\pi^+\pi^-$

NODE=M057G10
NODE=M057G10

NEW

$\Gamma(\phi\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.71 ± 0.05 OUR FIT		Error includes scale factor of 1.5.	[0.60 ± 0.05 eV OUR 2023 FIT]	
0.62 ± 0.07 ± 0.05	89 ± 11	¹ LIU	12B BELL	$\gamma\gamma \rightarrow 2(K^+K^-)$

NODE=M057G12
NODE=M057G12

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.58 ± 0.18 ± 0.16	26.5 ± 8.1	UEHARA	08 BELL	$\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+K^-)$
--------------------	------------	--------	---------	--

¹ Supersedes UEHARA 08. Using $B(\phi \rightarrow K^+K^-) = (48.9 \pm 0.5)\%$.

NODE=M057G12;LINKAGE=LI

$\Gamma(\omega\omega) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$				
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.64		90	¹ LIU	12B BELL
				$\gamma\gamma \rightarrow 2(\pi^+\pi^-\pi^0)$

NODE=M057G03
NODE=M057G03

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using $B(\omega \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7)\%$.

NODE=M057G03;LINKAGE=LI

$$\Gamma(\omega\phi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M057G04
NODE=M057G04

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04	90	¹ LIU	12B	BELL $\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
-------	----	------------------	-----	---

¹ Using $B(\phi \rightarrow K^+ K^-) = (48.9 \pm 0.5)\%$ and $B(\omega \rightarrow \pi^+ \pi^- \pi^0) = (89.2 \pm 0.7)\%$.

NODE=M057G04;LINKAGE=LI

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G4
NODE=M057G4

1.31±0.08 OUR FIT Error includes scale factor of 1.2. [1.25 ± 0.07 eV OUR 2023 FIT]

NEW

1.18±0.25 OUR AVERAGE

1.44±0.54±0.47	34 ± 13	¹ UEHARA	09	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
1.14±0.21±0.17	54 ± 10	² NAKAZAWA	05	BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^-$

¹ We multiplied the measurement by 3 to convert from $\pi^0 \pi^0$ to $\pi\pi$. Interference with the continuum included.

NODE=M057G4;LINKAGE=UE

² We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi\pi$.

NODE=M057G;LINKAGE=NA

$$\Gamma(\rho^0 \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{26}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G07
NODE=M057G07

2.1±0.9 OUR FIT

3.2±1.9±0.5 986 ± 578 UEHARA 08 BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(\pi^+ \pi^-)$

$$\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{31}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G13
NODE=M057G13

0.53±0.22±0.09 8 ¹ UEHARA 10A BELL 10.6 $e^+ e^- \rightarrow e^+ e^- \eta\eta$

¹ Interference with the continuum not included.

NODE=M057G13;LINKAGE=UE

$$\Gamma(K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{32}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G5
NODE=M057G5

0.59±0.09 OUR FIT Error includes scale factor of 2.1. [0.56 ± 0.04 eV OUR 2023 FIT]

NEW

0.44±0.11±0.07 33 ± 8 NAKAZAWA 05 BELL 10.6 $e^+ e^- \rightarrow e^+ e^- K^+ K^-$

$$\Gamma(K_S^0 K_S^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G6
NODE=M057G6

0.307±0.026 OUR FIT Error includes scale factor of 1.1. [0.294 ± 0.025 eV OUR 2023 FIT]

NEW

0.27 $\begin{smallmatrix} +0.07 \\ -0.06 \end{smallmatrix} \pm 0.03$ 53 ¹ UEHARA 13 BELL $\gamma\gamma \rightarrow K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.31 ± 0.05 ± 0.03	38 ± 7	CHEN	07B	BELL $e^+ e^- \rightarrow e^+ e^- \chi_{c2}$
--------------------	--------	------	-----	--

¹ Supersedes CHEN 07B.

NODE=M057G6;LINKAGE=UE

$$\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G01
NODE=M057G01

0.75±0.11 OUR FIT

[0.72 ± 0.11 eV OUR 2023 FIT]

NEW

1.20±0.33±0.13 126 ¹ DEL-AMO-SA..11M BABR $\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

¹ We have multiplied $\bar{K}^0 K \pi$ by 2/3 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$

NODE=M057G01;LINKAGE=DE

$$\Gamma(K^+ K^- K^+ K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_{98}/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M057G11
NODE=M057G11

0.97±0.13 OUR FIT Error includes scale factor of 1.1. [0.93 ± 0.11 eV OUR 2023 FIT]

NEW

1.10±0.21±0.15 126 ± 24 UEHARA 08 BELL $\gamma\gamma \rightarrow \chi_{c2} \rightarrow 2(K^+ K^-)$

$$\Gamma(\eta_c(1S)\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{93}\Gamma_{98}/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M057G05
NODE=M057G05

<15.7 90 LEES 12AE BABR $e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

$\chi_{c2}(1P)$ BRANCHING RATIOS

NODE=M057225

HADRONIC DECAYS

NODE=M057305

$$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma(2(\pi^+ \pi^-)) \quad \Gamma_{26}/\Gamma_1$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M057R38
NODE=M057R38

0.36±0.15 OUR FIT

[0.36 ± 0.15 eV OUR 2023 FIT]

NEW

0.31±0.17 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c2}$

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.86±0.24 OUR AVERAGE				[(1.83 ± 0.23)% OUR 2023 AVERAGE]
1.86±0.23±0.05	903.5	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R46
 NODE=M057R46
 NEW

¹ HE 08B reports $1.87 \pm 0.07 \pm 0.22 \pm 0.13$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R46;LINKAGE=HE

 $\Gamma(\rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
2.22±0.35 OUR AVERAGE				[(2.19 ± 0.34)% OUR 2023 AVERAGE]
2.22±0.34±0.05	1031.9	^{1,2} HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R47
 NODE=M057R47
 NEW

¹ HE 08B reports $2.23 \pm 0.11 \pm 0.32 \pm 0.16$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^+\pi^-\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R47;LINKAGE=HE

² Calculated by us. We have added the values from HE 08B for $\rho^+\pi^-\pi^0$ and $\rho^-\pi^+\pi^0$ decays assuming uncorrelated statistical and fully correlated systematic uncertainties.

NODE=M057R47;LINKAGE=OC

 $\Gamma(4\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.13±0.15 OUR AVERAGE				[(1.11 ± 0.15) $\times 10^{-3}$ OUR 2023 AVERAGE]
1.13±0.15±0.03	1164	¹ ABLIKIM	11A BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R58
 NODE=M057R58

NEW

¹ ABLIKIM 11A reports $(1.21 \pm 0.05 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 4\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R58;LINKAGE=AB

 $\Gamma(K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.21±0.04±0.01	76.9	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R48
 NODE=M057R48

NODE=M057R48;LINKAGE=HE

¹ HE 08B reports $0.21 \pm 0.03 \pm 0.03 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.41±0.20 OUR AVERAGE				[(1.38 ± 0.20)% OUR 2023 AVERAGE]
1.41±0.20±0.03	211.6	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R50
 NODE=M057R50

NEW

¹ HE 08B reports $1.41 \pm 0.11 \pm 0.16 \pm 0.10$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+\pi^-\bar{K}^0\pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R50;LINKAGE=HE

 $\Gamma(\rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.42±0.13 OUR AVERAGE				[(0.41 ± 0.12)% OUR 2023 AVERAGE]
0.42±0.13±0.01	62.9	¹ HE	08B CLEO	$e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R51
 NODE=M057R51

NEW

¹ HE 08B reports $0.42 \pm 0.11 \pm 0.06 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho^-K^+\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R51;LINKAGE=HE

$\Gamma(K^*(892)^0 K^- \pi^+ \rightarrow K^- \pi^+ K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.08 OUR AVERAGE		[(0.29 ± 0.08)% OUR 2023 AVERAGE]		
0.30±0.08±0.01	38.7	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R57
 NODE=M057R57
 NEW

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 K^- \pi^+ \rightarrow K^- \pi^+ K^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R57;LINKAGE=HE

 $\Gamma(K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.39±0.09 OUR AVERAGE		[(0.38 ± 0.09)% OUR 2023 AVERAGE]		
0.39±0.09±0.01	63.0	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R52
 NODE=M057R52
 NEW

¹ HE 08B reports $0.39 \pm 0.07 \pm 0.05 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R52;LINKAGE=HE

 $\Gamma(K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.38±0.08 OUR AVERAGE		[(0.37 ± 0.08)% OUR 2023 AVERAGE]		
0.38±0.08±0.01	51.1	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R53
 NODE=M057R53
 NEW

¹ HE 08B reports $0.38 \pm 0.07 \pm 0.04 \pm 0.03$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^- K^+ \pi^0 \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R53;LINKAGE=HE

 $\Gamma(K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.30±0.08 OUR AVERAGE		[(0.29 ± 0.08)% OUR 2023 AVERAGE]		
0.30±0.08±0.01	39.3	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R54
 NODE=M057R54
 NEW

¹ HE 08B reports $0.30 \pm 0.07 \pm 0.04 \pm 0.02$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{K}^0 \pi^- \rightarrow K^+ \pi^- \bar{K}^0 \pi^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R54;LINKAGE=HE

 $\Gamma(K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.130±0.045±0.003	22.9	¹ HE	08B CLEO	$e^+ e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NODE=M057R55
 NODE=M057R55

¹ HE 08B reports $0.13 \pm 0.04 \pm 0.02 \pm 0.01$ % from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R55;LINKAGE=HE

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
11.69±0.13±1.31	11k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R00
 NODE=M057R00

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R00;LINKAGE=A

 $\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.30±0.11±0.75	4.5k	¹ ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R73
 NODE=M057R73

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R73;LINKAGE=A

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma(K^+ K^- \pi^+ \pi^-)$ Γ_{17}/Γ_{14}

VALUE	DOCUMENT ID	TECN	COMMENT
0.25±0.13 OUR FIT			
0.25±0.13	TANENBAUM 78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

NODE=M057R39
 NODE=M057R39

$\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

15.3±1.9 OUR AVERAGE Error includes scale factor of 3.8. $[(15.2 \pm 1.7) \times 10^{-3}$ OUR 2023 AVERAGE Scale factor = 3.7]

15.9±0.4±0.4	112K	¹ ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+\pi^-)$
8.6±0.9±1.6		² BAI	99B BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
8.7±5.9±0.4		² TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ ABLIKIM 22Q reports $(1.565 \pm 0.005 \pm 0.048) \times 10^{-2}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow 3(\pi^+\pi^-))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Rescaled by us using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+\pi^-$ to $K^0 K^+\pi^-$ decay.

NODE=M057R4
NODE=M057R4

NEW

NODE=M057R4;LINKAGE=A

NODE=M057R;LINKAGE=X3

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.23 ±0.07 OUR FIT Error includes scale factor of 1.9. $[(1.06 \pm 0.09) \times 10^{-3}$ OUR 2023 FIT]

1.27 ±0.04 OUR AVERAGE $[(1.9 \pm 0.7) \times 10^{-3}$ OUR 2006 AVERAGE]

1.267±0.028±0.033	4247	¹ ABLIKIM	23N BES3	$\psi(2S) \rightarrow \gamma$ hadrons
--------------------------	------	----------------------	----------	---------------------------------------

¹ Measured using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.75 \pm 0.24) \times 10^{-2}$ and $B(\phi \rightarrow K^+K^-) = (49.2 \pm 0.5) \times 10^{-2}$ from PDG 22.

NODE=M057R20
NODE=M057R20

NEW

NEW

NODE=M057R20;LINKAGE=A

 $\Gamma(\phi\phi\eta)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.4±0.7 OUR AVERAGE $[(5.3 \pm 0.6) \times 10^{-4}$ OUR 2023 AVERAGE]

5.4±0.7±0.1	143.6	¹ ABLIKIM	20B BES3	$\psi(2S) \rightarrow \gamma \phi\phi\eta$
--------------------	-------	----------------------	----------	--

¹ ABLIKIM 20B reports $(5.33 \pm 0.52 \pm 0.39) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \phi\phi\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R97
NODE=M057R97

NEW

NODE=M057R97;LINKAGE=A

 $\Gamma(\omega\omega)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.86±0.10 OUR AVERAGE $[(0.84 \pm 0.10) \times 10^{-3}$ OUR 2023 AVERAGE]

0.83±0.10±0.02	762	¹ ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
1.76±0.58±0.04	27.7±7.4	² ABLIKIM	05N BES2	$\psi(2S) \rightarrow \gamma \chi_{c2} \rightarrow \gamma 6\pi$

¹ ABLIKIM 11K reports $(8.9 \pm 0.3 \pm 1.1) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 05N reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.165 \pm 0.044 \pm 0.032) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R28
NODE=M057R28

NEW

NODE=M057R28;LINKAGE=AL

NODE=M057R28;LINKAGE=AB

 $\Gamma(\omega K^+K^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.73±0.04±0.08	512	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$
-----------------------	-----	----------------------	----------	--

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R74
NODE=M057R74

NODE=M057R74;LINKAGE=A

 $\Gamma(\omega\phi)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

9.7±2.8 OUR AVERAGE $[(9.6 \pm 2.7) \times 10^{-6}$ OUR 2023 AVERAGE]

9.7±2.8±0.2		33	¹ ABLIKIM	19J BES3	$\psi(2S) \rightarrow \gamma$ hadrons
--------------------	--	----	----------------------	----------	---------------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

<19		90	^{2,3} ABLIKIM	11K BES3	$\psi(2S) \rightarrow \gamma$ hadrons
-----	--	----	------------------------	----------	---------------------------------------

NODE=M057R63
NODE=M057R63

NEW

¹ ABLIKIM 19J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.91 \pm 0.23 \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R63;LINKAGE=A

² ABLIKIM 11K reports $< 2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \omega\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R63;LINKAGE=AL

³ Superseded by ABLIKIM 19J.

NODE=M057R63;LINKAGE=B

$\Gamma(\pi^+\pi^-\pi^0(\text{non-resonant}))/\Gamma_{\text{total}}$

 Γ_{27}/Γ

VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	COMMENT
2.0±0.4±0.1	64	¹ ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

NODE=M057R84
NODE=M057R84

OCCUR=2

¹ ABLIKIM 17AG reports $(2.1 \pm 0.4 \pm 0.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\pi^0(\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R84;LINKAGE=B

$\Gamma(\rho(770)^\pm\pi^\mp)/\Gamma_{\text{total}}$

 Γ_{28}/Γ

VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	COMMENT
0.62±0.39±0.02	15	¹ ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\pi^0$

NODE=M057R85
NODE=M057R85

¹ ABLIKIM 17AG reports $(0.64 \pm 0.39 \pm 0.07) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \rho(770)^\pm\pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R85;LINKAGE=A

$\Gamma(\pi^+\pi^-\eta)/\Gamma_{\text{total}}$

 Γ_{29}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
0.49±0.13 OUR AVERAGE				$[(0.48 \pm 0.13) \times 10^{-3}]$ OUR 2023 AVERAGE]

NODE=M057R08
NODE=M057R08

NEW

0.49±0.13±0.01 ¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

••• We do not use the following data for averages, fits, limits, etc. •••

< 1.5 90 ² ABLIKIM 06R BES2 $\psi(2S) \rightarrow \gamma\chi_{c2}$

¹ ATHAR 07 reports $(0.49 \pm 0.12 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R08;LINKAGE=AT

² ABLIKIM 06R reports $< 1.7 \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R08;LINKAGE=AB

$\Gamma(\pi^+\pi^-\eta')/\Gamma_{\text{total}}$

 Γ_{30}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.51±0.19 OUR AVERAGE			$[(0.50 \pm 0.18) \times 10^{-3}]$ OUR 2023 AVERAGE]

NODE=M057R35
NODE=M057R35

NEW

0.51±0.19±0.01 ¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports $(0.51 \pm 0.18 \pm 0.06) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+\pi^-\eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R35;LINKAGE=AT

$\Gamma(K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}$

 Γ_{34}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.46±0.21 OUR AVERAGE			$[(1.44 \pm 0.21) \times 10^{-4}]$ OUR 2023 AVERAGE]

NODE=M057R86
NODE=M057R86

NEW

1.46±0.21±0.04 ¹ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K \bar{K} \pi$

••• We do not use the following data for averages, fits, limits, etc. •••

$1.75 \pm 0.27 \pm 0.04$ ² ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

OCCUR=2

$1.36 \pm 0.27 \pm 0.03$ ³ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

OCCUR=3

- ¹ ABLIKIM 17AG reports $(1.5 \pm 0.1 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² ABLIKIM 17AG reports $(1.8 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ABLIKIM 17AG reports $(1.4 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R86;LINKAGE=A

NODE=M057R86;LINKAGE=B

NODE=M057R86;LINKAGE=C

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{35}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.27±0.27 OUR AVERAGE [(1.24 ± 0.27) × 10 ⁻⁴ OUR 2023 AVERAGE]			
1.27±0.27±0.03	¹ ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

NODE=M057R87
NODE=M057R87

NEW

- ¹ ABLIKIM 17AG reports $(1.3 \pm 0.2 \pm 0.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R87;LINKAGE=A

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$ **Γ_{36}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
15.1±1.3 OUR AVERAGE [(14.8 ± 1.2) × 10 ⁻⁴ OUR 2023 AVERAGE]			
15.1±1.2±0.4	¹ ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K \bar{K} \pi$
17.7±1.6±0.4	² ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
13.2±1.5±0.3	³ ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

NODE=M057R88
NODE=M057R88

NEW

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

- ¹ ABLIKIM 17AG reports $(15.5 \pm 0.6 \pm 1.2) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² ABLIKIM 17AG reports $(18.2 \pm 0.8 \pm 1.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ³ ABLIKIM 17AG reports $(13.6 \pm 0.8 \pm 1.4) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R88;LINKAGE=A

NODE=M057R88;LINKAGE=B

NODE=M057R88;LINKAGE=C

 $\Gamma(K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ **Γ_{37}/Γ**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
12.7±1.7 OUR AVERAGE [(12.4 ± 1.7) × 10 ⁻⁴ OUR 2023 AVERAGE]			
12.7±1.7±0.3	¹ ABLIKIM	17AG BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

NODE=M057R89
NODE=M057R89

NEW

- ¹ ABLIKIM 17AG reports $(13.0 \pm 1.0 \pm 1.5) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_2^*(1430)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R89;LINKAGE=A

$\Gamma(K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{38}/Γ VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

5.3±0.8 OUR AVERAGE [(5.2 ± 0.8) × 10⁻⁴ OUR 2023 AVERAGE]NODE=M057R90
NODE=M057R90

NEW

5.3±0.8±0.1¹ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K \bar{K} \pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.2±1.0±0.1

² ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

OCCUR=2

5.7±1.8±0.1

³ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

OCCUR=3

¹ ABLIKIM 17AG reports (5.4 ± 0.5 ± 0.7) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R90;LINKAGE=A

² ABLIKIM 17AG reports (5.3 ± 0.5 ± 0.9) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R90;LINKAGE=B

³ ABLIKIM 17AG reports (5.9 ± 1.1 ± 1.5) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^\pm K^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R90;LINKAGE=C

 $\Gamma(K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{39}/Γ VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

5.7±2.1 OUR AVERAGE [(5.6 ± 2.1) × 10⁻⁴ OUR 2023 AVERAGE]NODE=M057R91
NODE=M057R91

NEW

5.7±2.1±0.1¹ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ ABLIKIM 17AG reports (5.9 ± 1.6 ± 1.5) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K_3^*(1780)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R91;LINKAGE=A

 $\Gamma(a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

13.1±3.5 OUR AVERAGE [(12.9 ± 3.4) × 10⁻⁴ OUR 2023 AVERAGE]NODE=M057R92
NODE=M057R92

NEW

13.1±3.5±0.3¹ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

¹ ABLIKIM 17AG reports (13.5 ± 1.6 ± 3.2) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R92;LINKAGE=A

 $\Gamma(a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{41}/Γ VALUE (units 10^{-4})

DOCUMENT ID TECN COMMENT

17.9±6.2±0.4¹ ABLIKIM 17AG BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ ABLIKIM 17AG reports (18.4 ± 3.3 ± 5.5) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow a_2(1320)^\pm \pi^\mp)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R93
NODE=M057R93

NODE=M057R93;LINKAGE=A

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{43}/Γ VALUE (units 10^{-3})

DOCUMENT ID TECN COMMENT

0.31±0.08 OUR AVERAGE [(0.30 ± 0.08) × 10⁻³ OUR 2023 AVERAGE]NODE=M057R05
NODE=M057R05

NEW

0.31±0.08±0.01¹ ATHAR 07 CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ATHAR 07 reports (0.31 ± 0.07 ± 0.04) × 10⁻³ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R05;LINKAGE=AT

$\Gamma(K^+ K^- \eta)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.33 (CL = 90%)				[<0.32 $\times 10^{-3}$ (CL = 90%) OUR 2023 BEST LIMIT]
<0.33	90	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma h^+ h^- h^0$
¹ ATHAR 07 reports < 0.33 $\times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.				

NODE=M057R09
NODE=M057R09

NODE=M057R09;LINKAGE=AT

 $\Gamma(K^+ K^- \eta'(958))/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.94\pm0.34	107	¹ ABLIKIM	14J	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \eta'(958)$
¹ Derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.72 \pm 0.34)\%$. Uncertainty includes both statistical and systematic contributions combined in quadrature.				

NODE=M057R78
NODE=M057R78

NODE=M057R78;LINKAGE=A

 $\Gamma(\eta\eta')/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.2\pm0.5\pm0.1		20	¹ ABLIKIM	17AI	BES3 $\psi(2S) \rightarrow \gamma \eta' \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 6	90	3.3 \pm 8.0	² ASNER	09	CLEO $\psi(2S) \rightarrow \gamma \eta \eta'$
< 23	90		³ ADAMS	07	CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$
¹ ABLIKIM 17AI reports $(2.27 \pm 0.43 \pm 0.25) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² ASNER 09 reports < 0.6 $\times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.					
³ Superseded by ASNER 09. ADAMS 07 reports < 2.3 $\times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.					

NODE=M057R03
NODE=M057R03

NODE=M057R03;LINKAGE=A

NODE=M057R03;LINKAGE=AS

NODE=M057R03;LINKAGE=AD

 $\Gamma(\eta' \eta')/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
4.6\pm0.6\pm0.1		60	¹ ABLIKIM	17AI	BES3 $\psi(2S) \rightarrow \gamma \eta' \eta'$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 10	90	12 \pm 7	² ASNER	09	CLEO $\psi(2S) \rightarrow \gamma \eta' \eta'$
< 31	90		³ ADAMS	07	CLEO $\psi(2S) \rightarrow \gamma \chi_{c2}$
¹ ABLIKIM 17AI reports $(4.76 \pm 0.56 \pm 0.38) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² ASNER 09 reports < 1.0 $\times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.					
³ Superseded by ASNER 09. ADAMS 07 reports < 3.1 $\times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \eta' \eta')/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0933 \pm 0.0014 \pm 0.0061$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.					

NODE=M057R04
NODE=M057R04

NODE=M057R04;LINKAGE=A

NODE=M057R04;LINKAGE=AS

NODE=M057R04;LINKAGE=AD

 $\Gamma(\pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2\pm0.5\pm0.1	57 \pm 11	¹ ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma \chi_{c2}$
¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow \pi^+ \pi^- K_S^0 K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ = $(0.207 \pm 0.039 \pm 0.033) \times 10^{-3}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M057R31
NODE=M057R31

NODE=M057R31;LINKAGE=AB

$\Gamma(K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4	90	2.3 ± 2.2	¹ ABLIKIM	050 BES2	$e^+e^- \rightarrow \chi_{c2}\gamma$

NODE=M057R32
NODE=M057R32

¹ ABLIKIM 050 reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] < 3.5 \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R32;LINKAGE=AB

 $\Gamma(K_S^0K_S^0K_S^0K_S^0)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.15 ± 0.18 OUR AVERAGE				[(1.13 ± 0.18) × 10 ⁻⁴ OUR 2023 AVERAGE]
1.15 ± 0.18 ± 0.03	68	¹ ABLIKIM	19AA BES3	$\psi(2S) \rightarrow \gamma 4K_S^0$

NODE=M057R94
NODE=M057R94

NEW

¹ Using $B(K_S^0 \rightarrow \pi^+\pi^-) = (69.20 \pm 0.05)\%$. ABLIKIM 19AA reports $[\Gamma(\chi_{c2}(1P) \rightarrow K_S^0K_S^0K_S^0K_S^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (10.8 \pm 1.5 \pm 0.8) \times 10^{-6}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value..

NODE=M057R94;LINKAGE=A

 $\Gamma(K^+K^-\phi)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.45 ± 0.30 OUR AVERAGE				[(1.42 ± 0.29) × 10 ⁻³ OUR 2023 AVERAGE]
1.45 ± 0.30 ± 0.04	52	¹ ABLIKIM	06T BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$

NODE=M057R01
NODE=M057R01

NEW

¹ ABLIKIM 06T reports $(1.67 \pm 0.26 \pm 0.24) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+K^-\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R01;LINKAGE=AB

 $\Gamma(\bar{K}^0K^+\pi^-\phi + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
4.83 ± 0.32 ± 0.66	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R79
NODE=M057R79

 $\Gamma(K^+K^-\pi^0\phi)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
2.74 ± 0.16 ± 0.44	ABLIKIM	15M BES3	$\psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R80
NODE=M057R80

 $\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.93 ± 0.06 ± 0.10	408	¹ ABLIKIM	13B BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$

NODE=M057R75
NODE=M057R75

¹ Using 1.06×10^8 $\psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (8.72 \pm 0.34)\%$.

NODE=M057R75;LINKAGE=A

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.47 ± 0.04 OUR AVERAGE			
0.48 ± 0.04 ± 0.01	¹ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
0.44 ± 0.09 ± 0.01	² ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M057R06
NODE=M057R06

¹ ONYISI 10 reports $(4.83 \pm 0.25 \pm 0.35 \pm 0.31) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=ON

² ATHAR 07 reports $(0.44 \pm 0.08 \pm 0.05) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R06;LINKAGE=AT

 $\Gamma(p\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.177 ± 0.025 OUR AVERAGE			
			[(0.174 ± 0.025) × 10 ⁻³ OUR 2023 AVERAGE]
0.175 ± 0.027 ± 0.004	¹ ONYISI	10 CLE3	$\psi(2S) \rightarrow \gamma p\bar{p}X$
0.189 ± 0.071 ± 0.005	² ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

NODE=M057R34
NODE=M057R34

NEW

¹ ONYISI 10 reports $(1.76 \pm 0.23 \pm 0.14 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R34;LINKAGE=ON

² ATHAR 07 reports $(0.19 \pm 0.07 \pm 0.02) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\eta)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R34;LINKAGE=AT

 $\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

0.37±0.04 OUR AVERAGE [(0.36 ± 0.04) × 10⁻³ OUR 2023 AVERAGE]NODE=M057R56
NODE=M057R56**0.37±0.04±0.01** ¹ ONYISI 10 CLE3 $\psi(2S) \rightarrow \gamma p\bar{p}X$

NEW

¹ ONYISI 10 reports $(3.68 \pm 0.35 \pm 0.26 \pm 0.24) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R56;LINKAGE=ON

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.8±0.9±0.1 24 ± 7 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$ NODE=M057R62
NODE=M057R62

¹ ABLIKIM 11F reports $(3.04 \pm 0.85 \pm 0.43) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R62;LINKAGE=AB

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.32±0.34 OUR EVALUATION Treating systematic error as correlated.NODE=M057R6
NODE=M057R6**1.3 ± 0.4 OUR AVERAGE** Error includes scale factor of 1.3.

→ UNCHECKED ←

1.17±0.19±0.30 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma\chi_{c2}$ 2.64±1.03±0.14 ¹ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$

¹ Rescaled by us using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.3 \pm 0.4)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.6 \pm 0.5)\%$. Multiplied by a factor of 2 to convert from $K_S^0 K^+ \pi^-$ to $K^0 K^+ \pi^-$ decay.

NODE=M057R6;LINKAGE=X3

 $\Gamma(p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
-----------	------	-------------	------	---------

0.080±0.024 OUR AVERAGE [(0.078 ± 0.023)% OUR 2023 AVERAGE]NODE=M057R49
NODE=M057R49**0.080±0.024±0.002** 29.2 ¹ HE 08B CLEO $e^+e^- \rightarrow \gamma h^+ h^- h^0 h^0$

NEW

¹ HE 08B reports $0.08 \pm 0.02 \pm 0.01 \pm 0.01\%$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}\pi^0\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R49;LINKAGE=HE

 $\Gamma(p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.94±0.33 OUR AVERAGE [(1.91 ± 0.32) × 10⁻⁴ OUR 2023 AVERAGE]NODE=M057R59
NODE=M057R59**1.94±0.32±0.05** 131 ± 12 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p\bar{p}K^+K^-$

NEW

¹ ABLIKIM 11F reports $(2.08 \pm 0.19 \pm 0.30) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p}K^+K^- \text{ (non-resonant)})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R59;LINKAGE=AB

 $\Gamma(p\bar{p}K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<7.9 90 ¹ ABLIKIM 06D BES2 $\psi(2S) \rightarrow \chi_{c2}\gamma$ NODE=M057R30
NODE=M057R30

¹ Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

NODE=M057R;LINKAGE=AB

$\Gamma(p\bar{n}\pi^-)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

8.7±1.0 OUR AVERAGE[(8.5 ± 0.9) × 10⁻⁴ OUR 2023 AVERAGE]

8.5±1.0±0.2	3309	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-$
10.4±3.5±0.3		² ABLIKIM	06I BES2	$\psi(2S) \rightarrow \gamma p\pi^- X$

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.80 \pm 0.02 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 06I reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.97 \pm 0.20 \pm 0.26) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R33
 NODE=M057R33
 NEW

NODE=M057R33;LINKAGE=AL

NODE=M057R33;LINKAGE=AB

 $\Gamma(\bar{p}n\pi^+)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.1±0.8 OUR AVERAGE [(8.9 ± 0.8) × 10⁻⁴ OUR 2023 AVERAGE]

9.1±0.8±0.2	3732	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+$
-------------	------	----------------------	----------	--

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (0.85 \pm 0.02 \pm 0.07) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R70
 NODE=M057R70
 NEW

NODE=M057R70;LINKAGE=AL

 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

22.1±1.8 OUR AVERAGE [(21.7 ± 1.8) × 10⁻⁴ OUR 2023 AVERAGE]

22.1±1.7±0.5	2128	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma p\bar{n}\pi^-\pi^0$
--------------	------	----------------------	----------	--

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.07 \pm 0.06 \pm 0.15) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R71
 NODE=M057R71
 NEW

NODE=M057R71;LINKAGE=AL

 $\Gamma(\bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

21.5±1.9 OUR AVERAGE [(21.1 ± 1.8) × 10⁻⁴ OUR 2023 AVERAGE]

21.5±1.8±0.5	2352	¹ ABLIKIM	12J BES3	$\psi(2S) \rightarrow \gamma\bar{p}n\pi^+\pi^0$
--------------	------	----------------------	----------	---

¹ ABLIKIM 12J reports $[\Gamma(\chi_{c2}(1P) \rightarrow \bar{p}n\pi^+\pi^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))] = (2.01 \pm 0.06 \pm 0.16) \times 10^{-4}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R72
 NODE=M057R72
 NEW

NODE=M057R72;LINKAGE=AL

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

128±16 OUR AVERAGE [(125 ± 15) × 10⁻⁵ OUR 2023 AVERAGE]

128±15±3		371	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
----------	--	-----	----------------------	----------	---

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

<350		90	² ABLIKIM	06D BES2	$\psi(2S) \rightarrow \chi_{c2}\gamma$
------	--	----	----------------------	----------	--

¹ ABLIKIM 12I reports $(137.0 \pm 7.6 \pm 15.7) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\psi(2S) \rightarrow \chi_{c2}\gamma) = (9.3 \pm 0.6)\%$.

NODE=M057R29
 NODE=M057R29
 NEW

NODE=M057R29;LINKAGE=AL

NODE=M057R29;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

67±15 OUR AVERAGE [(66 ± 15) × 10⁻⁵ OUR 2023 AVERAGE]

67±15±2		36	¹ ABLIKIM	12I BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}\pi^+\pi^-$
---------	--	----	----------------------	----------	---

¹ ABLIKIM 12I reports $(71.8 \pm 14.5 \pm 8.2) \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\pi^+\pi^- (\text{non-resonant}))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R65
 NODE=M057R65

NEW

NODE=M057R65;LINKAGE=AL

$\Gamma(\Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<40	90	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Lambda} \pi^-$
¹ ABLIKIM 12I reports $< 42 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Lambda} \pi^- + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.				

NODE=M057R66
NODE=M057R66

NODE=M057R66;LINKAGE=AL

 $\Gamma(\Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<60	90	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Lambda} \pi^+$
¹ ABLIKIM 12I reports $< 61 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Lambda} \pi^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.				

NODE=M057R67
NODE=M057R67

NODE=M057R67;LINKAGE=AL

 $\Gamma(K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.9±0.6 OUR AVERAGE				
[(7.8 ± 0.5) × 10 ⁻⁴ OUR 2023 AVERAGE]				

NODE=M057R07
NODE=M057R07

NEW

7.8±0.5±0.2	5k	^{1,2} ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma \Lambda \bar{p} K^+$
8.5±1.6±0.2		³ ATHAR	07 CLEO	$\psi(2S) \rightarrow \gamma h^+ h^- h^0$

¹ ABLIKIM 13D reports $(8.4 \pm 0.3 \pm 0.6) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R07;LINKAGE=AB

² Using $B(\Lambda \rightarrow p \pi^-) = 63.9\%$.

³ ATHAR 07 reports $(8.5 \pm 1.4 \pm 1.0) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R07;LINKAGE=LB
NODE=M057R07;LINKAGE=AT $\Gamma(n K_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{76}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.64±0.29 OUR AVERAGE				
[(3.58 ± 0.28) × 10 ⁻⁴ OUR 2023 AVERAGE]				

NODE=M057P01
NODE=M057P01

NEW

3.64±0.27±0.09	879	¹ ABLIKIM	21AV BES3	$\psi(2S) \rightarrow \gamma n K_S^0 \bar{\Lambda} + \text{c.c.}$
¹ ABLIKIM 21AV reports $(3.58 \pm 0.16 \pm 0.23) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow n K_S^0 \bar{\Lambda} + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 0.0952 \pm 0.0020$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Also uses $B(\bar{\Lambda} \rightarrow \bar{p} \pi^+) = (63.9 \pm 0.5)\%$ and $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$.				

NODE=M057P01;LINKAGE=B

 $\Gamma(K^*(892)^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{77}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.3±1.2 OUR AVERAGE				
[(8.2 ± 1.1) × 10 ⁻⁴ OUR 2023 AVERAGE]				

NODE=M057R98
NODE=M057R98

NEW

8.3±1.2±0.2	476	¹ ABLIKIM	19AU BES3	$\psi(2S) \rightarrow \gamma K^{*+} \bar{p} \Lambda$
¹ ABLIKIM 19AU reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^+ \bar{p} \Lambda + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ = $(7.8 \pm 0.9 \pm 0.6) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M057R98;LINKAGE=F

 $\Gamma(K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{78}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9±0.7 OUR AVERAGE				
[(2.8 ± 0.7) × 10 ⁻⁴ OUR 2023 AVERAGE]				

NODE=M057R60
NODE=M057R60

NEW

2.9±0.7±0.1	79 ± 13	¹ ABLIKIM	11F BES3	$\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$
¹ ABLIKIM 11F reports $(3.06 \pm 0.50 \pm 0.54) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow K^+ \bar{p} \Lambda(1520) + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M057R60;LINKAGE=AB

$\Gamma(\Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{total}$ Γ_{79}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

4.7±1.5 OUR AVERAGE [(4.6 ± 1.5) × 10⁻⁴ OUR 2023 AVERAGE]

4.7±1.5±0.1 29 ± 7 ¹ ABLIKIM 11F BES3 $\psi(2S) \rightarrow \gamma p \bar{p} K^+ K^-$

¹ ABLIKIM 11F reports (5.05 ± 1.29 ± 0.93) × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Lambda(1520)\bar{\Lambda}(1520))/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R61
NODE=M057R61

NEW

NODE=M057R61;LINKAGE=AB

$\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{total}$ Γ_{80}/Γ

VALUE (units 10^{-5}) CL% EVTS DOCUMENT ID TECN COMMENT

3.7±0.6±0.1 91 ¹ ABLIKIM 18V BES3 $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6 90 ² ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

<7 90 7.5 ± 3.4 ³ NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{\Sigma}^0$

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.35 \pm 0.05 \pm 0.02) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R43
NODE=M057R43

NODE=M057R43;LINKAGE=A

² ABLIKIM 13H reports < 0.65 × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R43;LINKAGE=AB

³ NAIK 08 reports < 0.75 × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{\Sigma}^0)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R43;LINKAGE=NA

$\Gamma(\Sigma^+ \bar{\Sigma}^-)/\Gamma_{total}$ Γ_{83}/Γ

VALUE (units 10^{-5}) CL% EVTS DOCUMENT ID TECN COMMENT

3.4±0.7±0.1 55 ¹ ABLIKIM 18V BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<8 90 ² ABLIKIM 13H BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

<7 90 4.0 ± 3.5 ³ NAIK 08 CLEO $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{\Sigma}^-$

¹ ABLIKIM 18V reports $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))] = (0.32 \pm 0.06 \pm 0.03) \times 10^{-5}$ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R44
NODE=M057R44

NODE=M057R44;LINKAGE=A

² ABLIKIM 13H reports < 0.88 × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R44;LINKAGE=AB

³ NAIK 08 reports < 0.67 × 10⁻⁴ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{\Sigma}^-)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R44;LINKAGE=NA

$\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{total}$ Γ_{84}/Γ

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT

4.5±1.8 OUR AVERAGE [(4.4 ± 1.8) × 10⁻⁵ OUR 2023 AVERAGE]

4.5±1.8±0.1 131 ¹ ABLIKIM 20I BES3 $\psi(2S) \rightarrow \gamma \Sigma^- \bar{\Sigma}^+$

¹ ABLIKIM 20I reports (4.4 ± 1.7 ± 0.5) × 10⁻⁵ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^- \bar{\Sigma}^+)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M057R99
NODE=M057R99

NEW

NODE=M057R99;LINKAGE=A

$\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{total}$ Γ_{85}/Γ

VALUE (units 10^{-5}) CL% DOCUMENT ID TECN COMMENT

<16 90 ¹ ABLIKIM 12I BES3 $\psi(2S) \rightarrow \gamma \Sigma(1385)^+ \bar{\Sigma}(1385)^-$

¹ ABLIKIM 12I reports < 17 × 10⁻⁵ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{total}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R68
NODE=M057R68

NODE=M057R68;LINKAGE=AL

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	¹ ABLIKIM 12I	BES3	$\psi(2S) \rightarrow \gamma \Sigma(1385)^- \bar{\Sigma}(1385)^+$
¹ ABLIKIM 12I reports $< 8.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.72 \pm 0.34) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.				

NODE=M057R69
NODE=M057R69

NODE=M057R69;LINKAGE=AL

 $\Gamma(K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.80±0.32 OUR AVERAGE				[(1.76 ± 0.32) × 10 ⁻⁴ OUR 2023 AVERAGE]
1.80±0.32±0.04	51	¹ ABLIKIM 15I	BES3	$\psi(2S) \rightarrow \gamma K^- \Lambda \bar{\Xi}^+ + \text{c.c.}$
¹ ABLIKIM 15I reports $[\Gamma(\chi_{c2}(1P) \rightarrow K^- \Lambda \bar{\Xi}^+ + \text{c.c.})/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ = (1.68 ± 0.26 ± 0.15) × 10 ⁻⁵ which we divide by our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M057R81
NODE=M057R81

NEW

NODE=M057R81;LINKAGE=A

 $\Gamma(\Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.86±0.22 OUR AVERAGE					[(1.83 ± 0.22) × 10 ⁻⁴ OUR 2023 AVERAGE]
1.86±0.22±0.05		804	¹ ABLIKIM 220	BES3	$\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.1	90	3	² NAIK 08	CLEO	$\psi(2S) \rightarrow \gamma \Xi^0 \bar{\Xi}^0$
¹ ABLIKIM 220 reports $(1.83 \pm 0.15 \pm 0.16) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² NAIK 08 reports $< 1.06 \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^0 \bar{\Xi}^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.					

NODE=M057R45
NODE=M057R45

NEW

NODE=M057R45;LINKAGE=A

NODE=M057R45;LINKAGE=NA

 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.46±0.12 OUR AVERAGE					[(1.44 ± 0.12) × 10 ⁻⁴ OUR 2023 AVERAGE]
1.47±0.12±0.04		1691	¹ ABLIKIM 220	BES3	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
1.45±0.32±0.04		29 ± 5	² NAIK 08	CLEO	$\psi(2S) \rightarrow \gamma \Xi^- \bar{\Xi}^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.7	90		³ ABLIKIM 06D	BES2	$\psi(2S) \rightarrow \chi_{c2} \gamma$
¹ ABLIKIM 220 reports $(1.44 \pm 0.06 \pm 0.11) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.52 \pm 0.20) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
² NAIK 08 reports $(1.45 \pm 0.30 \pm 0.15) \times 10^{-4}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.33 \pm 0.14 \pm 0.61) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.					
³ Using $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (9.3 \pm 0.6)\%$.					

NODE=M057R17
NODE=M057R17

NEW

NODE=M057R17;LINKAGE=A

NODE=M057R17;LINKAGE=NA

NODE=M057R17;LINKAGE=AB

 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.52±0.24±0.18	1038	ABLIKIM 23T	BES3	$\chi_{cJ} \rightarrow \Omega^- \bar{\Omega}^+$

NODE=M057P03
NODE=M057P03 $\Gamma(J/\psi(1S) \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	90	BARATE 81	SPEC	190 GeV $\pi^- \text{Be} \rightarrow 2\pi 2\mu$

NODE=M057R13
NODE=M057R13

$\Gamma(\pi^0 \eta_c)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-3}$	90	¹ ABLIKIM	15N BES3	$\psi(2S) e^+ e^- \rightarrow \gamma \pi^0 \eta_c$

¹ Using $B(\eta_c \rightarrow K_S^0 K^\pm \pi^\mp) \times B(K_S^0 \rightarrow \pi^+ \pi^-) \times B(\pi^0 \rightarrow \gamma \gamma) = (1.66 \pm 0.11) \times 10^{-2}$.

NODE=M057R77
NODE=M057R77

NODE=M057R77;LINKAGE=A

 $\Gamma(\eta_c(1S) \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.54 \times 10^{-2}$	90	^{1,2} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$
$<1.2 \times 10^{-2}$	90	^{1,3} ABLIKIM	13B BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using $1.06 \times 10^8 \psi(2S)$ mesons and $B(\psi(2S) \rightarrow \chi_{c2} \gamma) = (8.72 \pm 0.34)\%$.

² From the $\eta_c \rightarrow K_S^0 K^\pm \pi^\mp$ decays.

³ From the $\eta_c \rightarrow K^+ K^- \pi^0$ decays.

NODE=M057R76
NODE=M057R76

OCCUR=2

NODE=M057R76;LINKAGE=A
NODE=M057R76;LINKAGE=B
NODE=M057R76;LINKAGE=C $\Gamma(\eta_c(1S) \pi^+ \pi^-)/\Gamma(\bar{K}^0 K^+ \pi^- + \text{c.c.})$ Γ_{93}/Γ_{42}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<16.4	90	¹ LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$

¹ We divided the reported limit by 2 to take into account the $K_L^0 K^+ \pi^-$ mode.

NODE=M057R64
NODE=M057R64

NODE=M057R64;LINKAGE=LE

NODE=M057310

RADIATIVE DECAYS

 $\Gamma(\gamma \rho^0)/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<19	90	13 ± 11	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \rho^0$
<40	90	17.2 ± 6.8	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \rho^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABLIKIM 11E reports $<20.8 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

² BENNETT 08A reports $<50 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \rho^0)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R40
NODE=M057R40

NODE=M057R40;LINKAGE=AB

NODE=M057R40;LINKAGE=BE

 $\Gamma(\gamma \omega)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<6	90	1 ± 6	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \omega$
<6	90	0.0 ± 1.8	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \omega$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABLIKIM 11E reports $<6.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

² BENNETT 08A reports $<7.0 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \omega)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R41
NODE=M057R41

NODE=M057R41;LINKAGE=AB

NODE=M057R41;LINKAGE=BE

 $\Gamma(\gamma \phi)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8 (CL = 90%)		$[<7 \times 10^{-6}$ (CL = 90%) OUR 2023 BEST LIMIT]			
<8	90	5 ± 5	¹ ABLIKIM	11E BES3	$\psi(2S) \rightarrow \gamma \gamma \phi$
<11	90	1.3 ± 2.5	² BENNETT	08A CLEO	$\psi(2S) \rightarrow \gamma \gamma \phi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABLIKIM 11E reports $<8.1 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

² BENNETT 08A reports $<13 \times 10^{-6}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow \gamma \phi)/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = 9.36 \times 10^{-2}$.

NODE=M057R42
NODE=M057R42

NODE=M057R42;LINKAGE=AB

NODE=M057R42;LINKAGE=BE

$$\Gamma(e^+e^- J/\psi(1S))/\Gamma_{\text{total}} \quad \Gamma_{99}/\Gamma$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M057R82
NODE=M057R82

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.41 ± 0.15 ± 0.06	1.3k	^{1,2} ABLIKIM	17I	BES3	$\psi(2S) \rightarrow \gamma e^+ e^- J/\psi$
--------------------	------	------------------------	-----	------	--

NODE=M057R82;LINKAGE=B

¹ ABLIKIM 17I reports $(2.48 \pm 0.08 \pm 0.16) \times 10^{-3}$ from a measurement of $[\Gamma(\chi_{c2}(1P) \rightarrow e^+ e^- J/\psi(1S))/\Gamma_{\text{total}}] \times [B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.11 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (9.36 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Not independent from other measurements reported by ABLIKIM 17I

NODE=M057R82;LINKAGE=C

$$\Gamma(e^+e^- J/\psi(1S))/\Gamma(\gamma J/\psi(1S)) \quad \Gamma_{99}/\Gamma_{94}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M057R83
NODE=M057R83

11.3 ± 0.4 ± 0.5	1.3k	¹ ABLIKIM	17I	BES3	$\psi(2S) \rightarrow e^+ e^- \gamma J/\psi$
-------------------------	------	----------------------	-----	------	--

NODE=M057R83;LINKAGE=A

¹ Uses $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) \times B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (199.6 \pm 0.8 \pm 7.0) \times 10^{-4}$ from ABLIKIM 17N and accounts for common systematic errors.

$$\Gamma(\mu^+\mu^- J/\psi(1S))/\Gamma(e^+e^- J/\psi(1S)) \quad \Gamma_{100}/\Gamma_{99}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M057R96
NODE=M057R96

9.40 ± 0.79 ± 1.15	219	ABLIKIM	19Z	BES3	$\psi(2S) \rightarrow \gamma \chi_c \rightarrow \gamma(\mu^+\mu^- J/\psi)$
---------------------------	-----	---------	-----	------	--

$$\Gamma(\gamma\gamma)/\Gamma(\gamma J/\psi(1S)) \quad \Gamma_{98}/\Gamma_{94}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057R23
NODE=M057R23

1.50 ± 0.08 OUR FIT	Error includes scale factor of 1.5. [(1.50 ± 0.05) × 10 ⁻³ OUR 2023 FIT]		
----------------------------	---	--	--

NEW

0.99 ± 0.18	¹ AMBROGIANI 00B	E835	$\bar{p}p \rightarrow \chi_{c2} \rightarrow \gamma\gamma, \gamma J/\psi$
--------------------	-----------------------------	------	--

¹ Calculated by us using $B(J/\psi(1S) \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$.

NODE=M057R;LINKAGE=7A

$$\Gamma(\gamma\gamma)/\Gamma_{\text{total}} \times \Gamma(\rho\bar{\rho})/\Gamma_{\text{total}} \quad \Gamma_{98}/\Gamma \times \Gamma_{56}/\Gamma$$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057R24
NODE=M057R24

2.12 ± 0.15 OUR FIT	Error includes scale factor of 1.2. [(2.09 ± 0.13) × 10 ⁻⁸ OUR 2023 FIT]		
----------------------------	---	--	--

NEW

1.7 ± 0.4 OUR AVERAGE

1.60 ± 0.42	ARMSTRONG	93	E760	$\bar{p}p \rightarrow \gamma\gamma X$
9.9 ± 4.5	BAGLIN	87B	SPEC	$\bar{p}p \rightarrow \gamma\gamma X$

$\chi_{c2}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M057230

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-) \quad \Gamma_{14}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057B18
NODE=M057B18

2.24 ± 0.30 OUR FIT	Error includes scale factor of 1.2. [(2.31 ± 0.26) × 10 ⁻³ OUR 2023 FIT]		
----------------------------	---	--	--

NEW

2.5 ± 0.9 OUR AVERAGE Error includes scale factor of 2.3.

1.90 ± 0.14 ± 0.44	BAI	99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
--------------------	-----	-----	-----	---

3.8 ± 0.67	¹ TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$
------------	------------------------	----	------	---

¹ The reported value is derived using $B(\psi(2S) \rightarrow \pi^+ \pi^- J/\psi) \times B(J/\psi \rightarrow \ell^+ \ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

NODE=M057B18;LINKAGE=TA

$$\Gamma(\chi_{c2}(1P) \rightarrow K^*(892)^0 \bar{K}^*(892)^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P))/\Gamma_{\text{total}} \quad \Gamma_{18}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M057B19
NODE=M057B19

2.1 ± 0.9 OUR FIT	Error includes scale factor of 2.3. [(2.1 ± 0.4) × 10 ⁻⁴ OUR 2023 FIT]		
--------------------------	---	--	--

NEW

3.11 ± 0.36 ± 0.48	ABLIKIM	04H	BES2	$\psi(2S) \rightarrow \gamma \chi_{c2}$
---------------------------	---------	-----	------	---

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{56}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.96±0.10 OUR FIT				Error includes scale factor of 1.1. [(2.01 ± 0.09) × 10 ⁻⁵ OUR 2023 FIT]

NODE=M057B1
NODE=M057B1

NEW

1.4 ± 1.1 ¹ BAI 98I BES $\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow \gamma\bar{p}p$

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow p\bar{p})$ reported in BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M057B;LINKAGE=J8

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{56}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
6.79±0.34 OUR FIT				Error includes scale factor of 1.1. [(6.98 ± 0.32) × 10 ⁻⁶ OUR 2023 FIT]

NODE=M057B6
NODE=M057B6

NEW

7.1 ± 0.5 OUR AVERAGE Error includes scale factor of 1.2.

7.3 ± 0.4 ± 0.3	405	ABLIKIM	13V	BES3	$\psi(2S) \rightarrow \gamma p\bar{p}$
7.2 ± 0.7 ± 0.4	121 ± 12	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$
4.4 ^{+1.6} _{-1.4} ± 0.6	14.3 ^{+5.2} _{-4.7}	BAI	04F	BES	$\psi(2S) \rightarrow \gamma\chi_{c2}(1P) \rightarrow \gamma\bar{p}p$

¹ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow p\bar{p}) = (7.7 \pm 0.8 \pm 0.4 \pm 0.5) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

NODE=M057B6;LINKAGE=NA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{69}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
17.4±1.4 OUR FIT				
17.3±1.5 OUR AVERAGE				

NODE=M057B10
NODE=M057B10

18.2±0.8±1.7	670	ABLIKIM	21L	BES3	$\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+$
15.9±2.1±1.0	71	¹ NAIK	08	CLEO	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.2±1.4±0.9	207	^{2,3} ABLIKIM	13H	BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$
--------------	-----	------------------------	-----	------	---

¹ Calculated by us. NAIK 08 reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (17.0 \pm 2.2 \pm 1.1 \pm 1.1) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

NODE=M057B10;LINKAGE=NA

² Superseded by ABLIKIM 21L

NODE=M057B10;LINKAGE=A
NODE=M057B10;LINKAGE=AB

³ Calculated by us. ABLIKIM 13H reports $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) = (20.8 \pm 1.6 \pm 2.3) \times 10^{-5}$ from a measurement of $B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) \times B(\psi(2S) \rightarrow \gamma\chi_{c2})$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = (8.74 \pm 0.35)\%$.

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)}{\Gamma_{69}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.0±0.4 OUR FIT				

NODE=M057B11
NODE=M057B11

7.1^{+3.1} _{-2.9} ± 1.3 8.3^{+3.7} _{-3.4} ¹ BAI 03E BES $\psi(2S) \rightarrow \gamma\Lambda\bar{\Lambda}$

¹ BAI 03E reports $[B(\chi_{c2} \rightarrow \Lambda\bar{\Lambda}) B(\psi(2S) \rightarrow \gamma\chi_{c2}) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)] \times [B^2(\Lambda \rightarrow \pi^- p) / B(J/\psi \rightarrow p\bar{p})] = (1.33^{+0.59} _{-0.55} ± 0.25)\%$. We calculate from this measurement the presented value using $B(\Lambda \rightarrow \pi^- p) = (63.9 \pm 0.5)\%$ and $B(J/\psi \rightarrow p\bar{p}) = (2.17 \pm 0.07) \times 10^{-3}$.

NODE=M057B11;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{72}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.00±0.20±0.14	32	ABLIKIM	22A0	BES3	$\psi(2S) \rightarrow \gamma p\pi^-\bar{p}\pi^+\gamma\gamma$

NODE=M057P02
NODE=M057P02

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi\pi)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{\text{total}}}{\Gamma_{25}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.12±0.08 OUR FIT				[(2.12 ± 0.08) × 10 ⁻⁴ OUR 2023 FIT]
2.17±0.09 OUR AVERAGE				

NODE=M057B02
NODE=M057B02

NEW

2.19±0.05±0.15	4.5k	¹ ABLIKIM	10A	BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \gamma\chi_{c2}$
2.23±0.06±0.10	2.5k	² ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$
1.90±0.08±0.20	0.8k	³ ASNER	09	CLEO	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$

OCCUR=2

- ¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \pi^0 \pi^0) = (0.88 \pm 0.02 \pm 0.06 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.
- ² Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^+ \pi^-) = (1.59 \pm 0.04 \pm 0.07 \pm 0.10) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi \pi$.
- ³ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow \pi^0 \pi^0) = (0.68 \pm 0.03 \pm 0.07 \pm 0.04) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$. We have multiplied the $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.

NODE=M057B02;LINKAGE=AB

NODE=M057B02;LINKAGE=AS

NODE=M057B02;LINKAGE=AN

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \pi \pi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{25} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.611 ± 0.024 OUR FIT[(0.612 ± 0.023) × 10⁻³ OUR 2023 FIT]**0.54 ± 0.06 OUR AVERAGE**

0.66 ± 0.18 ± 0.37	21 ± 6	¹ BAI	03C BES	$\psi(2S) \rightarrow \gamma \pi^0 \pi^0$
0.54 ± 0.05 ± 0.04	185 ± 16	² BAI	98I BES	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$

¹ We have multiplied $\pi^0 \pi^0$ measurement by 3 to obtain $\pi \pi$.² Calculated by us. The value for $B(\chi_{c2} \rightarrow \pi^+ \pi^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D]. We have multiplied $\pi^+ \pi^-$ measurement by 3/2 to obtain $\pi \pi$.NODE=M057B9
NODE=M057B9

NEW

NODE=M057B;LINKAGE=BM

NODE=M057B;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \eta \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{31} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

0.52 ± 0.04 OUR FIT**0.52 ± 0.04 OUR AVERAGE**

0.54 ± 0.03 ± 0.04		386	¹ ABLIKIM	10A BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \gamma \chi_{c2}$
0.47 ± 0.05 ± 0.05		156	ASNER	09 CLEO	$\psi(2S) \rightarrow \gamma \eta \eta$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

< 0.44	90		² ADAMS	07 CLEO	$\psi(2S) \rightarrow \gamma \chi_{c2}$
< 3	90		BAI	03C BES	$\psi(2S) \rightarrow \gamma \eta \eta \rightarrow 5\gamma$
0.62 ± 0.31 ± 0.19			LEE	85 CBAL	$\psi(2S) \rightarrow \text{photons}$

¹ Calculated by us. ABLIKIM 10A reports $B(\chi_{c2} \rightarrow \eta \eta) = (0.65 \pm 0.04 \pm 0.05 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.3 \pm 0.4)\%$.² Superseded by ASNER 09.NODE=M057B04
NODE=M057B04

NODE=M057B04;LINKAGE=AB

NODE=M057B04;LINKAGE=AD

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{32} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

9.5 ± 1.4 OUR FIT Error includes scale factor of 2.4. [(9.6 ± 0.6) × 10⁻⁵ OUR 2023 FIT]**10.5 ± 0.3 ± 0.6** 1.6k ¹ ASNER 09 CLEO $\psi(2S) \rightarrow \gamma K^+ K^-$ ¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K^+ K^-) = (1.13 \pm 0.03 \pm 0.06 \pm 0.07) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.NODE=M057B03
NODE=M057B03

NEW

NODE=M057B03;LINKAGE=AS

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)}{\Gamma_{32} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.27 ± 0.04 OUR FIT Error includes scale factor of 2.4. [(0.276 ± 0.017) × 10⁻³ OUR 2023 FIT]**0.190 ± 0.034 ± 0.019** 115 ± 13 ¹ BAI 98I BES $\psi(2S) \rightarrow \gamma K^+ K^-$ ¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow K^+ K^-)$ reported by BAI 98I is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].NODE=M057B8
NODE=M057B8

NEW

NODE=M057B;LINKAGE=BI

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{33} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

5.0 ± 0.4 OUR FIT**5.0 ± 0.4 OUR AVERAGE**

4.9 ± 0.3 ± 0.3	373 ± 20	¹ ASNER	09	CLEO $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
5.72 ± 0.76 ± 0.63	65	ABLIKIM	050	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

¹ Calculated by us. ASNER 09 reports $B(\chi_{c2} \rightarrow K_S^0 K_S^0) = (0.53 \pm 0.03 \pm 0.03 \pm 0.03) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

NODE=M057B12
NODE=M057B12

NODE=M057B12;LINKAGE=AS

$$\Gamma(\chi_{c2}(1P) \rightarrow K_S^0 K_S^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_{33} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

14.3 ± 1.1 OUR FIT[(14.4 ± 1.1) × 10⁻⁵ OUR 2023 FIT]**14.7 ± 4.1 ± 3.3**

	¹ BAI	99B	BES	$\psi(2S) \rightarrow \gamma K_S^0 K_S^0$
--	------------------	-----	-----	---

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow K_S^0 K_S^0)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M057B13
NODE=M057B13

NEW

NODE=M057B13;LINKAGE=BA

$$\Gamma(\chi_{c2}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{42} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.22 ± 0.17 OUR FIT[(1.22 ± 0.17) × 10⁻⁴ OUR 2023 FIT]**1.15 ± 0.18 OUR AVERAGE**

1.21 ± 0.19 ± 0.09	37	¹ ATHAR	07	CLEO $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
0.97 ± 0.32 ± 0.13	28	² ABLIKIM	06R	BES2 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

¹ Calculated by us. ATHAR 07 reports $B(\chi_{c2} \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (1.3 \pm 0.2 \pm 0.1 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.33 \pm 0.14 \pm 0.61)\%$.

² Calculated by us. ABLIKIM 06R reports $B(\chi_{c2} \rightarrow K_S^0 K^\pm \pi^\mp) = (0.6 \pm 0.2 \pm 0.1) \times 10^{-3}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (8.1 \pm 0.6)\%$. We have multiplied by 2 to obtain $\bar{K}^0 K^+ \pi^- + \text{c.c.}$ from $K_S^0 K^\pm \pi^\mp$.

NODE=M057B05
NODE=M057B05

NEW

NODE=M057B05;LINKAGE=AT

NODE=M057B05;LINKAGE=AB

$$\Gamma(\chi_{c2}(1P) \rightarrow 2(\pi^+ \pi^-)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)$$

$$\Gamma_1 / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

2.7 ± 0.4 OUR FIT Error includes scale factor of 1.4. [(2.79 ± 0.26) × 10⁻³ OUR 2023 FIT]**3.1 ± 1.0 OUR AVERAGE** Error includes scale factor of 2.5.

2.3 ± 0.1 ± 0.5	¹ BAI	99B	BES	$\psi(2S) \rightarrow \gamma \chi_{c2}$
4.3 ± 0.6	² TANENBAUM	78	MRK1	$\psi(2S) \rightarrow \gamma \chi_{c2}$

¹ Calculated by us. The value for $B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in BAI 99B is derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

² The value for $B(\psi(2S) \rightarrow \gamma \chi_{c2}) \times B(\chi_{c2} \rightarrow 2\pi^+ 2\pi^-)$ reported in TANENBAUM 78 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times B(J/\psi(1S)\ell^+\ell^-) = (4.6 \pm 0.7)\%$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

NODE=M057B5
NODE=M057B5

NEW

NODE=M057B;LINKAGE=K1

NODE=M057B;LINKAGE=K2

$$\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}$$

$$\Gamma_{51} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.56 ± 0.21 OUR FIT Error includes scale factor of 1.1. [(1.57 ± 0.19) × 10⁻⁴ OUR 2023 FIT]**1.76 ± 0.16 ± 0.24**

	¹ ABLIKIM	06T	BES2	$\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
--	----------------------	-----	------	---

¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

NODE=M057B14
NODE=M057B14

NEW

NODE=M057B14;LINKAGE=AB

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow K^+ K^- K^+ K^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \quad \frac{\Gamma_{51} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}{\Gamma_{20} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT
4.5 ± 0.6 OUR FIT Error includes scale factor of 1.1. $[(4.5 \pm 0.5) \times 10^{-4}$ OUR 2023 FIT]

NODE=M057B15
 NODE=M057B15

NEW

3.6 ± 0.6 ± 0.6 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow 2K^+ 2K^-)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M057B15;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{20} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT
1.15 ± 0.07 OUR FIT Error includes scale factor of 1.8. $[(1.01 \pm 0.08) \times 10^{-4}$ OUR 2023 FIT]

NODE=M057B16
 NODE=M057B16

NEW

0.98 ± 0.13 OUR AVERAGE Error includes scale factor of 1.3.
 0.94 ± 0.03 ± 0.10 849 ¹ ABLIKIM 11K BES3 $\psi(2S) \rightarrow \gamma$ hadrons
 1.38 ± 0.24 ± 0.23 41 ² ABLIKIM 06T BES2 $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi \phi)$ reported by ABLIKIM 11K was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.74 \pm 0.35)\%$.
² Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi \phi)$ reported by ABLIKIM 06T was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (8.1 \pm 0.4)\%$.

NODE=M057B16;LINKAGE=AL

NODE=M057B16;LINKAGE=AB

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \phi \phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma(\psi(2S) \rightarrow J/\psi(1S) \pi^+ \pi^-)} \quad \frac{\Gamma_{20} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}{\Gamma_{20} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-4}) DOCUMENT ID TECN COMMENT
3.31 ± 0.21 OUR FIT Error includes scale factor of 1.7. $[(2.92 \pm 0.24) \times 10^{-4}$ OUR 2023 FIT]

NODE=M057B17
 NODE=M057B17

NEW

4.8 ± 1.3 ± 1.3 ¹ BAI 99B BES $\psi(2S) \rightarrow \gamma 2K^+ 2K^-$
¹ Calculated by us. The value of $B(\chi_{c2} \rightarrow \phi \phi)$ reported by BAI 99B was derived using $B(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) = (7.8 \pm 0.8)\%$ and $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = (32.4 \pm 2.6)\%$ [BAI 98D].

NODE=M057B17;LINKAGE=BA

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{81} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-6}) EVTS DOCUMENT ID TECN COMMENT
7.85 ± 0.77 ± 0.44 129 ¹ ABLIKIM 19BB BES3 $\psi(2S) \rightarrow \gamma \Sigma^+ \bar{p} K_S^0 + \text{c.c.}$

NODE=M057B07
 NODE=M057B07

¹ Calculated by us. ABLIKIM 19BB reports $B(\chi_{c2} \rightarrow \Sigma^+ \bar{p} K_S^0 + \text{c.c.}) = (8.25 \pm 0.83 \pm 0.49) \times 10^{-5}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$ and other branching fractions from PDG 18.

NODE=M057B07;LINKAGE=A

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{82} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-5}) EVTS DOCUMENT ID TECN COMMENT
0.87 ± 0.06 ± 0.04 271 ¹ ABLIKIM 20AE BES3 $\psi(2S) \rightarrow \gamma \Sigma^0 \bar{p} K^+ + \text{c.c.}$

NODE=M057P00
 NODE=M057P00

¹ Calculated by us. ABLIKIM 20AE reports $B(\chi_{c2} \rightarrow \Sigma^0 \bar{p} K^+ + \text{c.c.}) = (0.91 \pm 0.06 \pm 0.05) \times 10^{-4}$ using $B(\psi(2S) \rightarrow \gamma \chi_{c2}) = (9.52 \pm 0.20)\%$ and other branching fractions from PDG 20.

NODE=M057P00;LINKAGE=A

$$\frac{\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \chi_{c2}(1P)) / \Gamma_{\text{total}}}{\Gamma_{94} / \Gamma \times \Gamma_{181}^{\psi(2S)} / \Gamma_{12}^{\psi(2S)}}$$

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT
1.82 ± 0.07 OUR FIT Error includes scale factor of 1.9. $[(1.81 \pm 0.04) \times 10^{-2}$ OUR 2023 FIT]

NODE=M057B2
 NODE=M057B2

NEW

1.69 ± 0.16 OUR AVERAGE Error includes scale factor of 3.4. See the ideogram below.
 1.996 ± 0.008 ± 0.070 81k ¹ ABLIKIM 17N BES3 $\psi(2S) \rightarrow \gamma \gamma J/\psi$
 1.793 ± 0.008 ± 0.163 1.0M ABLIKIM 17U BES3 $e^+ e^- \rightarrow \gamma X$
 1.62 ± 0.04 ± 0.12 5.8k BAI 04I BES2 $\psi(2S) \rightarrow J/\psi \gamma \gamma$
 0.99 ± 0.10 ± 0.08 GAISER 86 CBAL $\psi(2S) \rightarrow \gamma X$
 1.47 ± 0.17 ² OREGLIA 82 CBAL $\psi(2S) \rightarrow \gamma \chi_{c2}$
 1.8 ± 0.5 ³ BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma \chi_{c2}$
 1.2 ± 0.2 ³ BARTEL 78B CNTR $\psi(2S) \rightarrow \gamma \chi_{c2}$
 2.2 ± 1.2 ⁴ BIDDICK 77 CNTR $e^+ e^- \rightarrow \gamma X$
 1.2 ± 0.7 ² WHITAKER 76 MRK1 $e^+ e^-$

••• We do not use the following data for averages, fits, limits, etc. •••

1.874±0.007±0.102	76k	5 ABLIKIM	120	BES3	$\psi(2S) \rightarrow \gamma\chi_{c2}$
1.95 ±0.02 ±0.07	12.4k	6 MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2}$
1.85 ±0.04 ±0.07	1.9k	7 ADAM	05A	CLEO	Repl. by MENDEZ 08

¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$.

² Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

³ Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.

⁴ Assumes isotropic gamma distribution.

⁵ Superseded by ABLIKIM 17N.

⁶ Not independent from other measurements of MENDEZ 08.

⁷ Not independent from other values reported by ADAM 05A.

NODE=M057B2;LINKAGE=A

NODE=M057B;LINKAGE=3Q

NODE=M057B;LINKAGE=2Q


NODE=M057B;LINKAGE=EA

NODE=M057B2;LINKAGE=B

NODE=M057B2;LINKAGE=ME

NODE=M057B;LINKAGE=AD

WEIGHTED AVERAGE
1.69±0.16 (Error scaled by 3.4)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{total} \text{ (units } 10^{-2}\text{)}$$

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma_{94}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.26±0.19 OUR FIT	Error includes scale factor of 1.8.			[(5.22 ± 0.11) × 10 ⁻² OUR 2023 FIT]

NODE=M057B3
NODE=M057B3

NEW

5.53±0.17 OUR AVERAGE

5.56±0.05±0.16	12.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2}$
6.0 ±2.8	1.3k	1 ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
3.9 ±1.2		2 HIMEL	80	MRK2	$\psi(2S) \rightarrow \gamma\chi_{c2}$

••• We do not use the following data for averages, fits, limits, etc. •••

5.52±0.13±0.13	1.9k	3 ADAM	05A	CLEO	Repl. by MENDEZ 08
----------------	------	--------	-----	------	--------------------

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow \gamma\chi_{c2}) \times B(\chi_{c2} \rightarrow \gamma J/\psi(1S))$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

NODE=M057B;LINKAGE=AB

NODE=M057B;LINKAGE=H8

NODE=M057B3;LINKAGE=AD

$$\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma)/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma\chi_{c2}(1P))/\Gamma_{total} \times \Gamma_{98}/\Gamma \times \Gamma_{181}^{\psi(2S)}/\Gamma_{12}^{\psi(2S)}$$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.74±0.10 OUR FIT	Error includes scale factor of 1.3.			[(2.71 ± 0.08) × 10 ⁻⁵ OUR 2023 FIT]

NODE=M057B4
NODE=M057B4

NEW

2.82±0.10 OUR AVERAGE

2.83±0.08±0.06	5k	1 ABLIKIM	17AE	BES3	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
2.68±0.28±0.15	0.3k	ECKLUND	08A	CLEO	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
7.0 ±2.1 ±2.0		LEE	85	CBAL	$\psi(2S) \rightarrow \gamma\chi_{c2}$

••• We do not use the following data for averages, fits, limits, etc. •••

2.81±0.17±0.15	1.1k	2 ABLIKIM	12A	BES3	$\psi(2S) \rightarrow \gamma\chi_{c2} \rightarrow 3\gamma$
----------------	------	-----------	-----	------	--

¹ ABLIKIM 17AE measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0} / \Gamma_{\gamma\gamma}^{\lambda=2} = 0.000 \pm 0.006 \pm 0.012$.

NODE=M057B4;LINKAGE=A

² ABLIKIM 12A measures the ratio of two-photon partial widths for the helicity $\lambda = 0$ and helicity $\lambda = 2$ components to be $f_{0/2} = \Gamma_{\gamma\gamma}^{\lambda=0} / \Gamma_{\gamma\gamma}^{\lambda=2} = 0.00 \pm 0.02 \pm 0.02$. Superseded by ABLIKIM 17AE.

NODE=M057B4;LINKAGE=AB

$\Gamma(\chi_{c2}(1P) \rightarrow \gamma\gamma) / \Gamma(\chi_{c0}(1P) \rightarrow \gamma\gamma)$ $\Gamma_{98} / \Gamma_{98}^{\chi_{c0}(1P)}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M057B06
NODE=M057B06**0.292 ± 0.028 OUR AVERAGE**

0.295 ± 0.014 ± 0.028	8k	¹ ABLIKIM	17AE BES3	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
-----------------------	----	----------------------	-----------	--

0.278 ± 0.050 ± 0.036	0.5k	¹ ECKLUND	08A CLEO	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
-----------------------	------	----------------------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.271 ± 0.029 ± 0.030	1.9k	^{1,2} ABLIKIM	12A BES3	$\psi(2S) \rightarrow \gamma\chi_{cJ} \rightarrow 3\gamma$
-----------------------	------	------------------------	----------	--

¹ Not independent from the values of $\Gamma(\chi_{c0}, \chi_{c2})$ and $B(\psi(2S) \rightarrow \chi_{c0}, \chi_{c2})$.

NODE=M057B06;LINKAGE=AB
NODE=M057B06;LINKAGE=A

² Superseded by ABLIKIM 17AE.

MULTIPOLE AMPLITUDES IN $\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$ RADIATIVE DECAY

NODE=M057240

$a_2 = M_2 / \sqrt{E_1^2 + M_2^2 + E_3^2}$ Magnetic quadrupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M057A1
NODE=M057A1**-11.0 ± 1.0 OUR AVERAGE**

-12.0 ± 1.3 ± 0.4	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-------------------	-----	----------------------	----------	---

-9.3 ± 1.6 ± 0.3	19.8k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
------------------	-------	---------------------	---------	---

-9.3 ⁺ ₋ 3.9 _{4.1} ± 0.6	5.9k	³ AMBROGIANI	02 E835	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
---	------	-------------------------	---------	---

-14 ± 6	1.9k	³ ARMSTRONG	93E E760	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
---------	------	------------------------	----------	---

-33.3 ⁺ ₋ 11.6 _{29.2}	441	³ OREGLIA	82 CBAL	$\psi(2S) \rightarrow \chi_{c1}\gamma \rightarrow J/\psi\gamma\gamma$
--	-----	----------------------	---------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

-7.9 ± 1.9 ± 0.3	19.8k	⁴ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
------------------	-------	---------------------	---------	---

OCCUR=2

¹ Correlated with a_3 , b_2 , and b_3 with correlation coefficients $\rho_{a_2 a_3} = 0.733$, $\rho_{a_2 b_2} = -0.605$, and $\rho_{a_2 b_3} = -0.095$.

NODE=M057A1;LINKAGE=B

² From a fit with floating M_2 amplitudes a_2 and b_2 , and fixed E_3 amplitudes $a_3 = b_3 = 0$.

NODE=M057A1;LINKAGE=AR

³ Assuming $a_3 = 0$.

NODE=M057A1;LINKAGE=A

⁴ From a fit with floating M_2 and E_3 amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057A1;LINKAGE=AT

$a_3 = E_3 / \sqrt{E_1^2 + M_2^2 + E_3^2}$ Electric octupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M057A2
NODE=M057A2

-0.3 ± 1.0 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

-1.3 ± 0.9 ± 0.4	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
------------------	-----	----------------------	----------	---

1.7 ± 1.4 ± 0.3	19.8k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-----------------	-------	---------------------	---------	---

2.0 ⁺ ₋ 5.5 _{4.4} ± 0.9	5908	AMBROGIANI	02 E835	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
--	------	------------	---------	---


0 ⁺ ₋ 6 ₅	1904	ARMSTRONG	93E E760	$\rho\bar{\rho} \rightarrow \chi_{c2} \rightarrow J/\psi\gamma$
--	------	-----------	----------	---

¹ Correlated with a_2 , b_2 , and b_3 with correlation coefficients $\rho_{a_2 a_3} = 0.733$, $\rho_{a_3 b_2} = -0.422$, and $\rho_{a_3 b_3} = -0.024$.

NODE=M057A2;LINKAGE=A

² From a fit with floating M_2 and E_3 amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057A2;LINKAGE=AR



$a_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude (units 10^{-2})

MULTIPOLE AMPLITUDES IN $\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ RADIATIVE DECAY

NODE=M057250

$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude

NODE=M057QB2
NODE=M057QB2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.9 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
1.7±0.8±0.2	89k	1 ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
4.6±1.0±1.3	13.8k	2 ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
0.2±1.5±0.4	19.8k	3 ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
- 5.1 ^{+5.4} _{-3.6}	721	2 ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
13.2 ^{+9.8} _{-7.5}	441	4 OREGLIA	82 CBAL	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.0±1.3±0.3 19.8k 4 ARTUSO 09 CLEO $\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M057QB2;LINKAGE=A

¹ Correlated with a_2 , a_3 , and b_3 with correlation coefficients $\rho_{a_2 b_2} = -0.605$, $\rho_{a_3 b_2} = -0.422$, and $\rho_{b_2 b_3} = 0.384$.

² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .


NODE=M057QB2;LINKAGE=AB

³ From a fit with floating $M2$ and $E3$ amplitudes a_2 , b_2 , and a_3 , and b_3 .

NODE=M057QB2;LINKAGE=AT

⁴ From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$.

NODE=M057QB2;LINKAGE=AR



$b_2 = M2/\sqrt{E1^2 + M2^2 + E3^2}$ Magnetic quadrupole fractional transition amplitude (units 10^{-2})

$b_3 = E3/\sqrt{E1^2 + M2^2 + E3^2}$ Electric octupole fractional transition amplitude

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-1.0±0.6 OUR AVERAGE				
-1.4±0.7±0.4	89k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.5±0.8±1.8	13.8k	² ABLIKIM	11I BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$
-0.8±1.2±0.2	19.8k	ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$
-2.7 ^{+4.3} _{-2.9}	721	² ABLIKIM	04I BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-, \gamma K^+K^-$

¹ Correlated with a_2 , a_3 , and b_2 with correlation coefficients $\rho_{a_2 b_3} = -0.095$, $\rho_{a_3 b_3} = -0.024$, and $\rho_{b_2 b_3} = 0.384$.

² From a fit with floating $M2$ and $E3$ amplitudes b_2 and b_3 .

NODE=M057QB3
NODE=M057QB3

NODE=M057QB3;LINKAGE=A

NODE=M057QB3;LINKAGE=AB

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

$\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$ and $\chi_{c2} \rightarrow \gamma J/\psi(1S)$

NODE=M057260

b_2/a_2 Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
-11⁺¹⁴ -15	19.8k	¹ ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma\gamma\ell^+\ell^-$

NODE=M057QAR
NODE=M057QAR

¹ Statistical and systematic errors combined. From a fit with floating $M2$ amplitudes a_2 and b_2 , and fixed $E3$ amplitudes $a_3=b_3=0$. Not independent of values for $a_2(\chi_{c2}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

NODE=M057QAR;LINKAGE=AR

$\chi_{c2}(1P)$ REFERENCES

NODE=M057

ABLIKIM	23N	JHEP 2305 069	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62056
ABLIKIM	23T	PR D107 092004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62064
SEINO	23	JHEP 2301 160	Y. Seino <i>et al.</i>	(BELLE Collab.)	REFID=62051
ABLIKIM	22AO	PR D106 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61887
ABLIKIM	22O	JHEP 2206 074	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61652
ABLIKIM	22Q	PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)	REFID=61634
ABLIKIM	21AV	JHEP 2111 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61465
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61117
ABLIKIM	20AE	PR D102 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60733
ABLIKIM	20B	PR D101 012012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60212
ABLIKIM	20I	PR D101 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60303
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19AA	PR D99 052008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59844
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM	19BB	PR D100 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60026
ABLIKIM	19J	PR D99 012015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59606
ABLIKIM	19Z	PR D99 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59837
ABLIKIM	18V	PR D97 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58990
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)	REFID=59304
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
AAIJ	17BI	PRL 119 221801	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58278
ABLIKIM	17AE	PR D96 092007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58310

ABLIKIM	17AG	PR D96 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58316
ABLIKIM	17AI	PR D96 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58321
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15M	PR D91 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56778
ABLIKIM	15N	PR D91 112018	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56779
ABLIKIM	14J	PR D89 074030	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55901
ABLIKIM	13B	PR D87 012002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54877
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13H	PR D87 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54953
ABLIKIM	13V	PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ABLIKIM	12A	PR D85 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54266
ABLIKIM	12I	PR D86 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54736
ABLIKIM	12J	PR D86 052011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54737
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
LIU	12B	PRL 108 232001	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=54303
ABLIKIM	11A	PR D83 012006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53647
ABLIKIM	11E	PR D83 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16717
ABLIKIM	11F	PR D83 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=16719
ABLIKIM	11I	PR D84 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53930
ABLIKIM	11K	PRL 107 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53940
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
ABLIKIM	10A	PR D81 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53347
ONYISI	10	PR D82 011103	P.U.E. Onyisi <i>et al.</i>	(CLEO Collab.)	REFID=53360
UEHARA	10A	PR D82 114031	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=53641
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
ASNER	09	PR D79 072007	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52721
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52761
BENNETT	08A	PRL 101 151801	J.V. Bennett <i>et al.</i>	(CLEO Collab.)	REFID=52575
ECKLUND	08A	PR D78 091501	K.M. Ecklund <i>et al.</i>	(CLEO Collab.)	REFID=52583
HE	08B	PR D78 092004	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52588
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
NAIK	08	PR D78 031101	P. Naik <i>et al.</i>	(CLEO Collab.)	REFID=52301
UEHARA	08	EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
ADAMS	07	PR D75 071101	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=51651
ATHAR	07	PR D75 032002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51618
CHEN	07B	PL B651 15	W.T. Chen <i>et al.</i>	(BELLE Collab.)	REFID=51710
ABLIKIM	06D	PR D73 052006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51049
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06T	PL B642 197	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51453
DOBBS	06	PR D73 071101	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51062
ABLIKIM	05G	PR D71 092002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50756
ABLIKIM	05N	PL B630 7	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50847
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05A	NP B717 34	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50769
NAKAZAWA	05	PL B615 39	H. Nakazawa <i>et al.</i>	(BELLE Collab.)	REFID=50807
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04H	PR D70 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50188
ABLIKIM	04I	PR D70 092004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50189
BAI	04F	PR D69 092001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49752
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
BAI	03E	PR D67 112001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49416
ABE	02T	PL B540 33	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48813
AMBROGIANI	02	PR D65 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48552
EISENSTEIN	01	PRL 87 061801	B.I. Eisenstein <i>et al.</i>	(CLEO Collab.)	REFID=48344
AMBROGIANI	00B	PR D62 052002	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47940
ACCIARRI	99E	PL B453 73	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46943
BAI	99B	PR D60 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47385
ACKER...K...	98	PL B439 197	K. Akerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46324
BAI	98D	PR D58 092006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46338
BAI	98I	PRL 81 3091	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46343
DOMINICK	94	PR D50 4265	J. Dominick <i>et al.</i>	(CLEO Collab.)	REFID=44077
ARMSTRONG	93	PRL 70 2988	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43306
ARMSTRONG	93E	PR D48 3037	T.A. Armstrong <i>et al.</i>	(FNAL-E760 Collab.)	REFID=48616
BAUER	93	PL B302 345	D.A. Bauer <i>et al.</i>	(TPC Collab.)	REFID=43315
ARMSTRONG	92	NP B373 35	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41865
Also		PRL 68 1468	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=41907
BAGLIN	87B	PL B187 191	C. Baglin <i>et al.</i>	(R704 Collab.)	REFID=40018
BAGLIN	86B	PL B172 455	C. Baglin (LAPP, CERN, GENO, LYON, OSLO+)		REFID=22145
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
LEE	85	SLAC 282	R.A. Lee	(SLAC)	REFID=40589
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
OREGLIA	82	PR D25 2259	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22120
Also		Private Comm.	M.J. Oreglia	(EFI)	REFID=22143
BARATE	81	PR D24 2994	R. Barate <i>et al.</i>	(SACL, LOIC, SHMP, CERN+)	REFID=22164
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
Also		Private Comm.	G. Trilling	(LBL, UCB)	REFID=22113
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151

$$I^G(J^{PC}) = 0^+(0^-+)$$

Quantum numbers are quark model predictions.

NODE=M059

NODE=M059

NODE=M059M

NODE=M059M

NEW

$\eta_c(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3637.7 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.2. [3637.7 ± 1.1 MeV OUR 2023 AVERAGE Scale factor = 1.2]
3637.90 ± 0.54 ± 1.40	3.7k	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
3643.4 ± 2.3 ± 4.4	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
3635.1 ± 3.7 ± 2.9	106	XU	18 BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
3633.6 ± 1.7 ± 0.6	106	¹ AAIJ	17AD LHCB	$pp \rightarrow B^+ X \rightarrow p \bar{p} K^+ X$
3636.4 ± 4.1 ± 0.7	365	² AAIJ	17BB LHCB	$pp \rightarrow b \bar{b} X \rightarrow 2(K^+ K^-) X$
3637.0 ± 5.7 ± 3.4	178	^{3,4} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^0$
3635.1 ± 5.8 ± 2.1	47	^{3,5} LEES	14E BABR	$\gamma \gamma \rightarrow K^+ K^- \eta$
3646.9 ± 1.6 ± 3.6	57	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
3637.6 ± 2.9 ± 1.6	127	⁶ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
3638.5 ± 1.5 ± 0.8	624	³ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
3640.5 ± 3.2 ± 2.5	1201	³ DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
3636.1 ^{+3.9 +0.7} ^{-4.2 -2.0}	128	⁷ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
3626 ± 5 ± 6	311	⁸ ABE	07 BELL	$e^+ e^- \rightarrow J/\psi (c \bar{c})$
3645.0 ± 5.5 ^{+4.9} ^{-7.8}	121	AUBERT	05C BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
3642.9 ± 3.1 ± 1.5	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

OCCUR=2

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

3639 ± 7	98	⁹ AUBERT	06E BABR	$B^\pm \rightarrow K^\pm X_{c \bar{c}}$
3630.8 ± 3.4 ± 1.0	112	¹⁰ AUBERT	04D BABR	$\gamma \gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
3654 ± 6 ± 8	39	¹¹ CHOI	02 BELL	$B \rightarrow K K_S K^- \pi^+$
3594 ± 5		¹² EDWARDS	82C CBAL	$e^+ e^- \rightarrow \gamma X$

NODE=M059M;LINKAGE=B

¹AAIJ 17AD report $m_{\psi(2S)} - m_{\eta_c(2S)} = 52.5 \pm 1.7 \pm 0.6$ MeV. We use the current value $m_{\psi(2S)} = 3686.097 \pm 0.025$ MeV to obtain the quoted mass.

²From a fit of the $\phi\phi$ invariant mass with the width of $\eta_c(2S)$ fixed to the PDG 16 value.

³Ignoring possible interference with continuum.

⁴With a width fixed to 11.3 MeV.

⁵With a width fixed to 11.3 MeV. Using both $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow \pi^+ \pi^- \pi^0$ decays.

⁶From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

⁷Accounts for interference with non-resonant continuum.

⁸From a fit of the J/ψ recoil mass spectrum. Supersedes ABE,K 02 and ABE 04G.

⁹From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

¹⁰Superseded by DEL-AMO-SANCHEZ 11M.

¹¹Superseded by VINOKUROVA 11.

¹²Assuming mass of $\psi(2S) = 3686$ MeV.

NODE=M059M;LINKAGE=C

NODE=M059M;LINKAGE=DE

NODE=M059M;LINKAGE=LE

NODE=M059M;LINKAGE=LS

NODE=M059M;LINKAGE=AB

NODE=M059M;LINKAGE=VA

NODE=M059M;LINKAGE=EB

NODE=M059M;LINKAGE=AU

NODE=M059M;LINKAGE=AR

NODE=M059M;LINKAGE=CH

NODE=M059M;LINKAGE=A

$\eta_c(2S)$ WIDTH

NODE=M059W

NODE=M059W

NEW

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
11.8 ± 1.6 OUR AVERAGE				[13.9 ± 2.6 MeV OUR 2023 AVERAGE]
10.77 ± 1.62 ± 1.08	3.7k	AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
19.8 ± 3.9 ± 3.1	569	ABLIKIM	22Q BES3	$\psi(2S) \rightarrow \gamma 3(\pi^+ \pi^-)$
9.9 ± 4.8 ± 2.9	57	ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$
16.9 ± 6.4 ± 4.8	127	¹ ABLIKIM	12G BES3	$\psi(2S) \rightarrow \gamma K^0 K \pi, K K \pi^0$
13.4 ± 4.6 ± 3.2	624	² DEL-AMO-SA..11M	BABR	$\gamma \gamma \rightarrow K_S^0 K^\pm \pi^\mp$
6.6 ^{+8.4 +2.6} ^{-5.1 -0.9}	128	³ VINOKUROVA	11 BELL	$B^\pm \rightarrow K^\pm (K_S^0 K^\pm \pi^\mp)$
6.3 ± 12.4 ± 4.0	61	ASNER	04 CLEO	$\gamma \gamma \rightarrow \eta'_c \rightarrow K_S^0 K^\pm \pi^\mp$

••• We do not use the following data for averages, fits, limits, etc. •••

< 23	90	98	⁴ AUBERT	06E	BABR	$B^\pm \rightarrow K^\pm X_c \bar{c}$
22 ± 14		121	AUBERT	05C	BABR	$e^+ e^- \rightarrow J/\psi c \bar{c}$
17.0 ± 8.3 ± 2.5		112	⁵ AUBERT	04D	BABR	$\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K \bar{K} \pi$
< 55	90	39	⁶ CHOI	02	BELL	$B \rightarrow K K_S K^- \pi^+$
< 8.0	95		⁷ EDWARDS	82C	CBAL	$e^+ e^- \rightarrow \gamma X$

¹ From a simultaneous fit to $K_S^0 K^\pm \pi^\mp$ and $K^+ K^- \pi^0$ decay modes.

² Ignoring possible interference with continuum.

³ Accounts for interference with non-resonant continuum.

⁴ From the fit of the kaon momentum spectrum. Systematic errors not evaluated.

⁵ Superseded by DEL-AMO-SANCHEZ 11M.

⁶ For a mass value of 3654 ± 6 MeV. Superseded by VINOKUROVA 11.

⁷ For a mass value of 3594 ± 5 MeV

NODE=M059W;LINKAGE=AB
 NODE=M059W;LINKAGE=DE
 NODE=M059W;LINKAGE=VA
 NODE=M059W;LINKAGE=AU
 NODE=M059W;LINKAGE=AR
 NODE=M059W;LINKAGE=W2
 NODE=M059W;LINKAGE=W

$\eta_c(2S)$ DECAY MODES

NODE=M059215;NODE=M059

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	not seen	
Γ_2 $K \bar{K} \pi$	(1.9±1.2) %	
Γ_3 $K \bar{K} \eta$	(5 ± 4) × 10 ⁻³	
Γ_4 $2\pi^+ 2\pi^-$	< 2.1 %	90%
Γ_5 $\rho^0 \rho^0$	< 1.9 × 10 ⁻³	90%
Γ_6 $3\pi^+ 3\pi^-$	(1.3±0.9) %	
Γ_7 $K^+ K^- \pi^+ \pi^-$	< 1.4 %	90%
Γ_8 $K^{*0} \bar{K}^{*0}$	< 2.9 × 10 ⁻³	90%
Γ_9 $K^+ K^- \pi^+ \pi^- \pi^0$	(1.4±1.0) %	
Γ_{10} $K^+ K^- 2\pi^+ 2\pi^-$	< 1.4 %	90%
Γ_{11} $K_S^0 K^- 2\pi^+ \pi^- + c.c.$	(1.0±0.8) %	
Γ_{12} $2K^+ 2K^-$	< 1.3 × 10 ⁻³	90%
Γ_{13} $\phi \phi$	< 1.1 × 10 ⁻³	90%
Γ_{14} $\rho \bar{\rho}$	< 2.0 × 10 ⁻³	90%
Γ_{15} $\rho \bar{\rho} \pi^+ \pi^-$	seen	
Γ_{16} $\gamma\gamma$	(1.8±1.2) × 10 ⁻⁴	
Γ_{17} $\gamma J/\psi(1S)$	< 1.4 %	90%
Γ_{18} $\pi^+ \pi^- \eta$	(4.3±3.2) × 10 ⁻³	
Γ_{19} $\pi^+ \pi^- \eta'$	(2.6±1.9) × 10 ⁻³	
Γ_{20} $K_2^*(1430) \bar{K} + c.c.$	seen	
Γ_{21} $K_0^*(1950) \bar{K} + c.c.$	seen	
Γ_{22} $a_0(1710) \pi$	seen	
Γ_{23} $a_0(1450) \pi$	seen	
Γ_{24} $a_2(1700) \pi$	seen	
Γ_{25} $K_0^*(2600) \bar{K} + c.c.$	seen	
Γ_{26} $\pi^+ \pi^- \eta_c(1S)$	< 25 %	90%

DESIG=1
 DESIG=4
 DESIG=20
 DESIG=5
 DESIG=16
 DESIG=8
 DESIG=6
 DESIG=17
 DESIG=9
 DESIG=10
 DESIG=11
 DESIG=7
 DESIG=18
 DESIG=3
 DESIG=22
 DESIG=2
 DESIG=21
 DESIG=12
 DESIG=13
 DESIG=24
 DESIG=25
 DESIG=26
 DESIG=27
 DESIG=28
 DESIG=29
 DESIG=15

$\eta_c(2S)$ PARTIAL WIDTHS

NODE=M059216

$\Gamma(\gamma\gamma)$

Γ_{16}

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M059W1
 NODE=M059W1

••• We do not use the following data for averages, fits, limits, etc. •••

0.44±0.14	106	¹ XU	18	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
1.3 ± 0.6		² ASNER	04	CLEO	$\gamma\gamma \rightarrow \eta_c' \rightarrow K_S^0 K^\pm \pi^\mp$

¹ Assuming that the branching fraction into $\eta' \pi^+ \pi^-$ is the same as for $\eta_c(1S)$.

² They measure $\Gamma(\eta_c(2S)\gamma\gamma) B(\eta_c(2S) \rightarrow K \bar{K} \pi) = (0.18 \pm 0.05 \pm 0.02) \Gamma(\eta_c(1S)\gamma\gamma) B(\eta_c(1S) \rightarrow K \bar{K} \pi)$. The value for $\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)$ is derived assuming that the branching fractions for $\eta_c(2S)$ and $\eta_c(1S)$ decays to $K_S K \pi$ are equal and using $\Gamma(\eta_c(1S) \rightarrow \gamma\gamma) = 7.4 \pm 0.4 \pm 2.3$ keV.

NODE=M059W1;LINKAGE=A
 NODE=M059W1;LINKAGE=AS

$\Gamma(\gamma\gamma) \times \Gamma(\pi^+ \pi^- \eta')/\Gamma_{total}$

$\Gamma_{16}\Gamma_{19}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M059R29
 NODE=M059R29

5.6^{+1.2}_{-1.1} ± 1.1	106	XU	18	BELL	$e^+ e^- \rightarrow e^+ e^- \eta' \pi^+ \pi^-$
--	-----	----	----	------	---

$\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M059218

 $\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_{16}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$41 \pm 4 \pm 6$	624	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$

NODE=M059G04
NODE=M059G04¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059G04;LINKAGE=DE

 $\Gamma(2\pi^+2\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 6.5	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(\pi^+\pi^-)$

NODE=M059G01
NODE=M059G01 $\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_7\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 5.0	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-$

NODE=M059G02
NODE=M059G02 $\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_9\Gamma_{16}/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$30 \pm 6 \pm 5$	1201	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M059G05
NODE=M059G05¹ Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059G05;LINKAGE=DE

 $\Gamma(2K^+2K^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{12}\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S) \rightarrow 2(K^+K^-)$

NODE=M059G03
NODE=M059G03 $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{26}\Gamma_{16}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 133	90	LEES	12AE	BABR $e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

NODE=M059G06
NODE=M059G06 $\eta_c(2S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma^2(\text{total})$

NODE=M059217

 $\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma \times \Gamma_{16}/\Gamma$

VALUE (units 10^{-8})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.6	90	^{1,2,3} AMBROGIANI 01	E835	$\bar{p}p \rightarrow \gamma\gamma$

NODE=M059G1
NODE=M059G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

 < 8.0 90 ^{1,2,4} AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$

OCCUR=2

 < 12.0 90 ^{2,4} AMBROGIANI 01 E835 $\bar{p}p \rightarrow \gamma\gamma$

OCCUR=3

¹ Including the measurements of of ARMSTRONG 95F in the AMBROGIANI 01 analysis.

NODE=M059G1;LINKAGE=A

² For a total width $\Gamma=5$ MeV.

NODE=M059G1;LINKAGE=B

³ For the resonance mass region 3589–3599 MeV/ c^2 .

NODE=M059G1;LINKAGE=C1

⁴ For the resonance mass region 3575–3660 MeV/ c^2 .

NODE=M059G1;LINKAGE=C2

 $\eta_c(2S) \text{ BRANCHING RATIOS}$

NODE=M059220

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABREU	980	DLPH $e^+e^- \rightarrow e^+e^- + \text{hadrons}$

NODE=M059R1
NODE=M059R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen ¹ EDWARDS 82C CBAL $e^+e^- \rightarrow \gamma X$ ¹ For a mass value of 3594 ± 5 MeV

NODE=M059R;LINKAGE=W

 $\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.9 \pm 0.4 \pm 1.1$	59 ± 12	¹ AUBERT	08AB	BABR $B \rightarrow \eta_c(2S)K \rightarrow K\bar{K}\pi K$

NODE=M059R3
NODE=M059R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 127 ± 18 ABLIKIM 12G BES3 $\psi(2S) \rightarrow \gamma K\bar{K}\pi$ seen 39 ± 11 ² CHOI 02 BELL $B \rightarrow K K_S K^-\pi^+$ ¹ Derived from a measurement of $[B(B^+ \rightarrow \eta_c(2S)K^+) \times B(\eta_c(2S) \rightarrow K\bar{K}\pi)] / [B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (9.6^{+2.0}_{-1.9} \pm 2.5)\%$ and using $B(B^+ \rightarrow \eta_c(2S)K^+) = (3.4 \pm 1.8) \times 10^{-4}$, and $[B(B^+ \rightarrow \eta_c K^+) \times B(\eta_c \rightarrow K\bar{K}\pi)] = (6.88 \pm 0.77^{+0.55}_{-0.66}) \times 10^{-5}$.

NODE=M059R3;LINKAGE=AU

² For a mass value of 3654 ± 6 MeV

NODE=M059R;LINKAGE=W2

$\Gamma(K\bar{K}\eta)/\Gamma(K\bar{K}\pi)$ Γ_3/Γ_2

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
27.3±7.0±9.0	225	¹ LEES	14E	BABR $\gamma\gamma \rightarrow K^+ K^- \gamma\gamma$

NODE=M059R26
 NODE=M059R26

¹ LEES 14E reports $B(\eta_c(2S) \rightarrow K^+ K^- \eta)/B(\eta_c(2S) \rightarrow K^+ K^- \pi^0) = 0.82 \pm 0.21 \pm 0.27$, which we divide by 3 to account for isospin symmetry.

NODE=M059R26;LINKAGE=LE

 $\Gamma(2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059R01
 NODE=M059R01

 $\Gamma(\rho^0 \rho^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma 2\pi^+ 2\pi^-$

NODE=M059R15
 NODE=M059R15

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059R02
 NODE=M059R02

 $\Gamma(K^*0 \bar{K}^*0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^-$

NODE=M059R16
 NODE=M059R16

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$ Γ_9/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.73±0.17±0.17	1201	¹ DEL-AMO-SA..11M	BABR	$\gamma\gamma \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M059R21
 NODE=M059R21

¹ We have multiplied the value of $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K_S^0 K^\pm \pi^\mp)$ reported in DEL-AMO-SANCHEZ 11M by a factor 1/3 to obtain $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma(K\bar{K}\pi)$. Not independent from other measurements reported in DEL-AMO-SANCHEZ 11M.

NODE=M059R21;LINKAGE=DE

 $\Gamma(K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	57 ± 17	ABLIKIM	13K	BES3 $\psi(2S) \rightarrow \gamma K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$

NODE=M059R22
 NODE=M059R22

 $\Gamma(2K^+ 2K^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	UEHARA	08	BELL $\gamma\gamma \rightarrow \eta_c(2S)$

NODE=M059R03
 NODE=M059R03

 $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ABLIKIM	11H	BES3 $\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M059R17
 NODE=M059R17

 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	106	¹ AAIJ	17AD	LHCB $pp \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$

NODE=M059R04
 NODE=M059R04

¹ AAIJ 17AD report a 6.4 standard deviation signal, with $B(B^+ \rightarrow \eta_c(2S) K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) = (1.58 \pm 0.33 \pm 0.09) \times 10^{-2}$.

OCCUR=2

NODE=M059R04;LINKAGE=A

 $\Gamma(p\bar{p}\pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	110	¹ CHILIKIN	19	BELL $e^+ e^- \rightarrow \Upsilon(4S)$

NODE=M059R30
 NODE=M059R30

¹ CHILIKIN 19 reports signals in $B^+ \rightarrow \eta_c(2S) K^+$ and $B^0 \rightarrow \eta_c(2S) K_S^0$ with 12.3 and 5.9 standard deviations, respectively.

NODE=M059R30;LINKAGE=A

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R2
 NODE=M059R2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4 × 10 ⁻⁴	90	¹ WICHT	08	BELL B [±] → K [±] γγ
not seen		AMBROGIANI	01	E835 $\bar{p}p \rightarrow \gamma\gamma$
<0.01	90	LEE	85	CBAL $\psi' \rightarrow \text{photons}$

¹ WICHT 08 reports $[\Gamma(\eta_c(2S) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \eta_c(2S)K^+)] < 0.18 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \eta_c(2S)K^+) = 4.4 \times 10^{-4}$.

NODE=M059R2;LINKAGE=WI

 $\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma(K\bar{K}\pi)$ Γ_{26}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R23
 NODE=M059R23

<3.33	90	¹ LEES	12AE BABR	e ⁺ e ⁻ → e ⁺ e ⁻ π ⁺ π ⁻ η _c
-------	----	-------------------	-----------	--

¹We divided the reported limit by 3 to take into account isospin relations.

NODE=M059R23;LINKAGE=LE

η_c(2S) CROSS-PARTICLE BRANCHING RATIOS

NODE=M059230

 $\Gamma(\eta_c(2S) \rightarrow K\bar{K}\eta)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_3/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R25
 NODE=M059R25

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11.8 × 10 ⁻⁶	90	¹ CRONIN-HEN..10	CLEO	ψ(2S) → γK ⁺ K ⁻ η
--------------------------	----	-----------------------------	------	--

¹ CRONIN-HENNESSY 10 reports a limit of < 5.9 × 10⁻⁶ for the decay η_c(2S) → K⁺K⁻η which we multiply by 2 account for isospin symmetry. It assumes Γ(η_c(2S)) = 14 MeV. It also gives the analytic dependence of limits on width.

NODE=M059R25;LINKAGE=CR

 $\Gamma(\eta_c(2S) \rightarrow 2\pi^+2\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_4/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R05
 NODE=M059R05

<14.6 × 10 ⁻⁶	90	¹ CRONIN-HEN..10	CLEO	ψ(2S) → γ2π ⁺ 2π ⁻
--------------------------	----	-----------------------------	------	--

¹ Assuming Γ(η_c(2S)) = 14 MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R05;LINKAGE=CR

 $\Gamma(\eta_c(2S) \rightarrow \rho^0\rho^0)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R18
 NODE=M059R18

<12.7 × 10 ⁻⁷	90	ABLIKIM	11H BES3	ψ(2S) → γ2π ⁺ 2π ⁻
--------------------------	----	---------	----------	--

 $\Gamma(\eta_c(2S) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE (units 10 ⁻⁶)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	------	-------------	------	---------

NODE=M059R06
 NODE=M059R06

9.2±1.0±1.2	569		ABLIKIM	22Q BES3	ψ(2S) → γ3(π ⁺ π ⁻)
-------------	-----	--	---------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.2	90	¹ CRONIN-HEN..10	CLEO	ψ(2S) → γ3π ⁺ 3π ⁻
-------	----	-----------------------------	------	--

¹ Assuming Γ(η_c(2S)) = 14 MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R06;LINKAGE=CR

 $\Gamma(\eta_c(2S) \rightarrow K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_7/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R07
 NODE=M059R07

<9.6 × 10 ⁻⁶	90	¹ CRONIN-HEN..10	CLEO	ψ(2S) → γK ⁺ K ⁻ π ⁺ π ⁻
-------------------------	----	-----------------------------	------	--

¹ Assuming Γ(η_c(2S)) = 14 MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R07;LINKAGE=CR

 $\Gamma(\eta_c(2S) \rightarrow K^{*0}\bar{K}^{*0})/\Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}$ $\Gamma_8/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma\psi(2S)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M059R19
 NODE=M059R19

<19.6 × 10 ⁻⁷	90	ABLIKIM	11H BES3	ψ(2S) → γK ⁺ K ⁻ π ⁺ π ⁻
--------------------------	----	---------	----------	--

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_9 / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<43.0 \times 10^{-6}$	90	¹ CRONIN-HEN...10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M059R08
NODE=M059R08

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R08;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K^+ K^- 2\pi^+ 2\pi^-) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{10} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K^+ K^- 2\pi^+ 2\pi^-$

NODE=M059R09
NODE=M059R09

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R09;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{11} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$7.03 \pm 2.10 \pm 0.7$	60		ABLIKIM	13K BES3	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$

NODE=M059R10
NODE=M059R10

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15.2	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma K_S^0 K^- 2\pi^+ \pi^- + \text{c.c.}$
---------	----	-----------------------------	------	--

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R10;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \phi\phi) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{13} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.8 \times 10^{-7}$	90	ABLIKIM	11H BES3	$\psi(2S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M059R20
NODE=M059R20

$$\Gamma(\eta_c(2S) \rightarrow \rho\bar{\rho}) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{14} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-6}$	90	ABLIKIM	13V BES3	$\psi(2S) \rightarrow \gamma \rho\bar{\rho}$

NODE=M059R24
NODE=M059R24

$$\Gamma(\eta_c(2S) \rightarrow \gamma J/\psi(1S)) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{17} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<9.7 \times 10^{-6}$	90	33	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma J/\psi$

NODE=M059R27
NODE=M059R27

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033)\%$.

NODE=M059R27;LINKAGE=A

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta) / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{18} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.97 \pm 0.81 \pm 0.26$		106	ABLIKIM	23Q BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.3	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta$
--------	----	-----------------------------	------	--

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R11;LINKAGE=CR

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta') / \Gamma_{\text{total}} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S)) / \Gamma_{\text{total}}$$

$$\Gamma_{19} / \Gamma \times \Gamma_{183}^{\psi(2S)} / \Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<14.2 \times 10^{-6}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta'$

NODE=M059R12
NODE=M059R12

¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R12;LINKAGE=CR

$\Gamma(K_2^*(1430)\bar{K} + c.c.)/\Gamma_{total}$ Γ_{20}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M059R34
 NODE=M059R34

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

NODE=M059R34;LINKAGE=A

 $\Gamma(a_0(1710)\pi)/\Gamma_{total}$ Γ_{22}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M059R37
 NODE=M059R37

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

NODE=M059R37;LINKAGE=A

 $\Gamma(a_0(1450)\pi)/\Gamma_{total}$ Γ_{23}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M059R38
 NODE=M059R38

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

NODE=M059R38;LINKAGE=A

 $\Gamma(a_2(1700)\pi)/\Gamma_{total}$ Γ_{24}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M059R39
 NODE=M059R39

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

NODE=M059R39;LINKAGE=A

 $\Gamma(K_0^*(2600)\bar{K} + c.c.)/\Gamma_{total}$ Γ_{25}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M059R40
 NODE=M059R40

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

NODE=M059R40;LINKAGE=A

 $\Gamma(K_0^*(1950)\bar{K} + c.c.)/\Gamma_{total}$ Γ_{21}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$

NODE=M059R35
 NODE=M059R35

¹ From a Dalitz plot analysis of $\eta_c(2S) \rightarrow K_S^0 K^+ \pi^- + c.c..$

NODE=M059R35;LINKAGE=A

$$\Gamma(\eta_c(2S) \rightarrow \pi^+ \pi^- \eta_c(1S))/\Gamma_{total} \times \Gamma(\psi(2S) \rightarrow \gamma \eta_c(2S))/\Gamma_{total}$$

$$\Gamma_{26}/\Gamma \times \Gamma_{183}^{\psi(2S)}/\Gamma_{\psi(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	¹ CRONIN-HEN..10	CLEO	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \eta_c(1S)$

NODE=M059R14
 NODE=M059R14


¹ Assuming $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M059R14;LINKAGE=CR

 $\eta_c(2S)$ REFERENCES

NODE=M059

AAIJ	23AH PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62349
ABLIKIM	23Q PR D107 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62060
ABLIKIM	22Q PR D106 032014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61663
CHILIKIN	19 PR D100 012001	K. Chilikin <i>et al.</i>	(BELLE Collab.)	REFID=59899
XU	18 PR D98 072001	Q.N. Xu <i>et al.</i>	(BELLE Collab.)	REFID=59453
AAIJ	17AD PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
AAIJ	17BB EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
ABLIKIM	17N PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
PDG	16 CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
LEES	14E PR D89 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55937
ABLIKIM	13K PR D87 052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54956
ABLIKIM	13V PR D88 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55583
ABLIKIM	12G PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54272
LEES	12AE PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
ABLIKIM	11H PR D84 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53929
DEL-AMO-SA...	11M PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
VINOKUROVA	11 PL B706 139	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=53927
CRONIN-HEN...	10 PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=53233
AUBERT	08AB PR D78 012006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52267
UEHARA	08 EPJ C53 1	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=52064
WICHT	08 PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
ABE	07 PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=51627
AUBERT	06E PRL 96 052002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51059
AUBERT	05C PR D72 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50773
ABE	04G PR D70 071102	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=50182
ASNER	04 PRL 92 142001	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=49745
AUBERT	04D PRL 92 142002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49746
ABE,K	02 PRL 89 142001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=49188
CHOI	02 PRL 89 102001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=48760
AMBROGIANI	01 PR D64 052003	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=48340
ABREU	98O PL B441 479	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46553
ARMSTRONG	95F PR D52 4839	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=44623
LEE	85 SLAC 282	R.A. Lee	(SLAC)	REFID=40589
EDWARDS	82C PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22173



$$I^G(J^{PC}) = 0^-(1^{--})$$

See the Review on "Branching Ratios of $\psi(2S)$, $\chi_{c0,1,2}$ and $\eta_c(1S)$ " before the $\chi_{c0}(1P)$ Listings.

NODE=M071

NODE=M071

$\psi(2S)$ MASS

NODE=M071M

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

NODE=M071M

NODE=M071M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.10 ± 0.06 OUR FIT		Error includes scale factor of 5.9.		
3686.097 ± 0.010 OUR AVERAGE				
3686.099 ± 0.004 ± 0.009		¹ ANASHIN	15	KEDR $e^+e^- \rightarrow$ hadrons
3686.12 ± 0.06 ± 0.10	4k	AAIJ	12H	LHCB $p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$
3685.95 ± 0.10	413	² ARTAMONOV	00	OLYA $e^+e^- \rightarrow$ hadrons
3685.98 ± 0.09 ± 0.04		³ ARMSTRONG	93B	E760 $\bar{p}p \rightarrow e^+e^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3686.08 ± 0.07	1301	⁴ AAIJ	23AP	LHCB $B_S^0 \rightarrow J/\psi 2(\pi^+ \pi^-)$
3686.114 ± 0.007 ^{+0.011} _{-0.016}		⁵ ANASHIN	12	KEDR $e^+e^- \rightarrow$ hadrons
3686.111 ± 0.025 ± 0.009		AULCHENKO	03	KEDR $e^+e^- \rightarrow$ hadrons
3686.00 ± 0.10	413	⁶ ZHOLENTZ	80	OLYA e^+e^-

¹ Supersedes AULCHENKO 03 and ANASHIN 12.

² Reanalysis of ZHOLENTZ 80 using new electron mass (COHEN 87) and radiative corrections (KURAEV 85).

³ Mass central value and systematic error recalculated by us according to Eq. (16) in ARMSTRONG 93B, using the value for the $J/\psi(1S)$ mass from AULCHENKO 03.

⁴ From a fit of a relativistic S -wave Breit-Wigner convolved with the detector resolution. The width of $\psi(2S)$ is constrained to the PDG 22 value. Systematic errors not evaluated.

⁵ From the scans in 2004 and 2006. ANASHIN 12 reports the value $3686.114 \pm 0.007 \pm 0.011^{+0.002}_{-0.012}$ MeV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

⁶ Superseded by ARTAMONOV 00.

NODE=M071M;LINKAGE=A

NODE=M071M;LINKAGE=AR

NODE=M071M;LINKAGE=NW

NODE=M071M;LINKAGE=B

NODE=M071M;LINKAGE=AN

NODE=M071M;LINKAGE=RZ

$m_{\psi(2S)} - m_{J/\psi(1S)}$

NODE=M071DM

NODE=M071DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.188 ± 0.028 OUR AVERAGE			
589.194 ± 0.027 ± 0.011	¹ AULCHENKO	03	KEDR $e^+e^- \rightarrow$ hadrons
589.7 ± 1.2	LEMOIGNE	82	GOLI $185 \pi^- \text{Be} \rightarrow \gamma \mu^+ \mu^- \text{A}$
589.07 ± 0.13	¹ ZHOLENTZ	80	OLYA e^+e^-
588.7 ± 0.8	LUTH	75	MRK1
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
588 ± 1	² BAI	98E	BES e^+e^-

¹ Redundant with data in mass above.

² Systematic errors not evaluated.

NODE=M071DM;LINKAGE=R

NODE=M071DM;LINKAGE=BD

$\psi(2S)$ WIDTH

NODE=M071W

NODE=M071W

NEW

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
293 ± 9 OUR FIT		Error includes scale factor of 1.2. [294 ± 8 keV OUR 2023 FIT]		
286 ± 16 OUR AVERAGE				
358 ± 88 ± 4		ABLIKIM	08B	BES2 $e^+e^- \rightarrow$ hadrons
290 ± 25 ± 4	2.7k	ANDREOTTI	07	E835 $p\bar{p} \rightarrow e^+e^-, J/\psi X$
331 ± 58 ± 2		ABLIKIM	06L	BES2 $e^+e^- \rightarrow$ hadrons
264 ± 27		¹ BAI	02B	BES2 e^+e^-
287 ± 37 ± 16		² ARMSTRONG	93B	E760 $\bar{p}p \rightarrow e^+e^-$

¹ From a simultaneous fit to the hadronic and $\mu^+ \mu^-$ cross section, assuming $\Gamma = \Gamma_h + \Gamma_e + \Gamma_\mu + \Gamma_\tau$ and lepton universality. Does not include vacuum polarization correction.

² The initial-state radiation correction reevaluated by ANDREOTTI 07 in its Ref. [4].

NODE=M071W;LINKAGE=BC

NODE=M071W;LINKAGE=AN

$\psi(2S)$ DECAY MODES

NODE=M071220;NODE=M071

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Γ_1 hadrons	(97.85 \pm 0.13) %		DESIG=3
Γ_2 virtual $\gamma \rightarrow$ hadrons	(1.79 \pm 0.04) %		DESIG=4
Γ_3 ggg	(10.6 \pm 1.6) %		DESIG=255
Γ_4 γgg	(1.03 \pm 0.29) %		DESIG=256
Γ_5 light hadrons	(15.4 \pm 1.5) %		DESIG=226
Γ_6 K_S^0 anything	(16.0 \pm 1.1) %		DESIG=325
Γ_7 $e^+ e^-$	(7.94 \pm 0.22) $\times 10^{-3}$	S=1.3	DESIG=1
Γ_8 $\mu^+ \mu^-$	(8.0 \pm 0.6) $\times 10^{-3}$		DESIG=2
Γ_9 $\tau^+ \tau^-$	(3.1 \pm 0.4) $\times 10^{-3}$		DESIG=68
Decays into $J/\psi(1S)$ and anything			
Γ_{10} $J/\psi(1S)$ anything	(61.5 \pm 0.7) %	S=1.3	NODE=M071;CLUMP=A DESIG=11
Γ_{11} $J/\psi(1S)$ neutrals	(25.4 \pm 0.5) %	S=1.6	DESIG=12
Γ_{12} $J/\psi(1S) \pi^+ \pi^-$	(34.69 \pm 0.34) %	S=1.1	DESIG=13
Γ_{13} $J/\psi(1S) \pi^0 \pi^0$	(18.2 \pm 0.5) %	S=1.6	DESIG=14
Γ_{14} $J/\psi(1S) \eta$	(3.37 \pm 0.06) %	S=1.2	DESIG=15
Γ_{15} $J/\psi(1S) \pi^0$	(1.268 \pm 0.032) $\times 10^{-3}$		DESIG=18
Hadronic decays			
Γ_{16} $\pi^+ \pi^-$	(7.8 \pm 2.6) $\times 10^{-6}$		NODE=M071;CLUMP=B DESIG=21
Γ_{17} $\pi^+ \pi^- \pi^0$	(2.01 \pm 0.17) $\times 10^{-4}$	S=1.7	DESIG=36
Γ_{18} $\rho(770) \pi \rightarrow \pi^+ \pi^- \pi^0$	(3.2 \pm 1.2) $\times 10^{-5}$	S=1.8	DESIG=22
Γ_{19} $\rho(2150) \pi \rightarrow \pi^+ \pi^- \pi^0$	(1.9 $^{+1.2}_{-0.4}$) $\times 10^{-4}$		DESIG=201
Γ_{20} $2(\pi^+ \pi^-)$	(2.4 \pm 0.6) $\times 10^{-4}$	S=2.2	DESIG=24
Γ_{21} $\rho^0 \pi^+ \pi^-$	(2.2 \pm 0.6) $\times 10^{-4}$	S=1.4	DESIG=33
Γ_{22} $2(\pi^+ \pi^-) \pi^0$	(2.9 \pm 1.0) $\times 10^{-3}$	S=4.7	DESIG=25
Γ_{23} $\rho a_2(1320)$	(2.6 \pm 0.9) $\times 10^{-4}$		DESIG=65
Γ_{24} $\pi^+ \pi^- \pi^0 \pi^0 \pi^0$	(5.3 \pm 1.0) $\times 10^{-3}$		DESIG=312
Γ_{25} $\pi^+ \pi^- 4\pi^0$	(1.4 \pm 1.0) $\times 10^{-3}$		DESIG=332
Γ_{26} $\rho^\pm \pi^\mp \pi^0 \pi^0$	< 2.7 $\times 10^{-3}$	CL=90%	DESIG=315
Γ_{27} $3(\pi^+ \pi^-)$	(3.5 \pm 2.0) $\times 10^{-4}$	S=2.8	DESIG=32
Γ_{28} $2(\pi^+ \pi^- \pi^0)$	(4.8 \pm 1.5) $\times 10^{-3}$		DESIG=221
Γ_{29} $3(\pi^+ \pi^-) \pi^0$	(3.5 \pm 1.6) $\times 10^{-3}$		DESIG=37
Γ_{30} $2(\pi^+ \pi^-) 3\pi^0$	(1.42 \pm 0.31) %		DESIG=329
Γ_{31} $\eta \pi^+ \pi^-$	< 1.6 $\times 10^{-4}$	CL=90%	DESIG=202
Γ_{32} $\eta \pi^+ \pi^- \pi^0$	(9.5 \pm 1.7) $\times 10^{-4}$		DESIG=203
Γ_{33} $\eta 2(\pi^+ \pi^-)$	(1.2 \pm 0.6) $\times 10^{-3}$		DESIG=251
Γ_{34} $\eta \pi^+ \pi^- \pi^0 \pi^0$	< 4 $\times 10^{-4}$	CL=90%	DESIG=313
Γ_{35} $\eta \pi^+ \pi^- 3\pi^0$	< 2.1 $\times 10^{-3}$	CL=90%	DESIG=334
Γ_{36} $\eta 2(\pi^+ \pi^- \pi^0)$	< 2.1 $\times 10^{-3}$	CL=90%	DESIG=328
Γ_{37} $\rho \eta$	(2.2 \pm 0.6) $\times 10^{-5}$	S=1.1	DESIG=94
Γ_{38} $\eta' \pi^+ \pi^- \pi^0$	(4.5 \pm 2.1) $\times 10^{-4}$		DESIG=204
Γ_{39} $\eta' \rho$	(1.9 $^{+1.7}_{-1.2}$) $\times 10^{-5}$		DESIG=93
Γ_{40} $\omega \pi^0$	(2.1 \pm 0.6) $\times 10^{-5}$		DESIG=92
Γ_{41} $\omega \pi^+ \pi^-$	(7.3 \pm 1.2) $\times 10^{-4}$	S=2.1	DESIG=75
Γ_{42} $\omega \pi^+ \pi^- 2\pi^0$	(8.7 \pm 2.4) $\times 10^{-3}$		DESIG=327
Γ_{43} $b_1^\pm \pi^\mp$	(4.0 \pm 0.6) $\times 10^{-4}$	S=1.1	DESIG=40
Γ_{44} $\omega f_2(1270)$	(2.2 \pm 0.4) $\times 10^{-4}$		DESIG=64
Γ_{45} $\omega \pi^0 \pi^0$	(1.11 \pm 0.35) $\times 10^{-3}$		DESIG=314
Γ_{46} $\omega 3\pi^0$	< 8 $\times 10^{-4}$	CL=90%	DESIG=333
Γ_{47} $b_1^0 \pi^0$	(2.4 \pm 0.6) $\times 10^{-4}$		DESIG=193
Γ_{48} $\omega \eta$	< 1.1 $\times 10^{-5}$	CL=90%	DESIG=95
Γ_{49} $\omega \eta'$	(3.2 $^{+2.5}_{-2.1}$) $\times 10^{-5}$		DESIG=91
Γ_{50} $\phi \pi^0$	< 4 $\times 10^{-7}$	CL=90%	DESIG=96
Γ_{51} $\phi \pi^+ \pi^-$	(1.18 \pm 0.26) $\times 10^{-4}$	S=1.5	DESIG=78
Γ_{52} $\phi f_0(980) \rightarrow \pi^+ \pi^-$	(7.5 \pm 3.3) $\times 10^{-5}$	S=1.6	DESIG=81
Γ_{53} $\phi \eta$	(3.10 \pm 0.31) $\times 10^{-5}$		DESIG=89

Г54	$\eta\phi(2170), \phi(2170) \rightarrow$ $\phi f_0(980), f_0 \rightarrow \pi^+\pi^-$	$< 2.2 \times 10^{-6}$	CL=90%	DESIG=316
Г55	$\phi\eta'$	$(1.54 \pm 0.20) \times 10^{-5}$		DESIG=90
Г56	$\phi f_1(1285)$	$(3.0 \pm 1.3) \times 10^{-5}$		DESIG=319
Г57	$\phi\eta(1405) \rightarrow \phi\pi^+\pi^-\eta$	$(8.5 \pm 1.7) \times 10^{-6}$		DESIG=320
Г58	$\phi f'_2(1525)$	$(4.4 \pm 1.6) \times 10^{-5}$		DESIG=67
Г59	K^+K^-	$(7.5 \pm 0.5) \times 10^{-5}$		DESIG=23
Г60	$K^+K^-\pi^+$	$(7.3 \pm 0.5) \times 10^{-4}$		DESIG=26
Г61	$K^+K^-\pi^0$	$(4.07 \pm 0.31) \times 10^{-5}$		DESIG=38
Г62	$K_S^0 K_S^0$	$< 4.6 \times 10^{-6}$		DESIG=86
Г63	$K_S^0 K_L^0$	$(5.34 \pm 0.33) \times 10^{-5}$		DESIG=85
Г64	$K_S^0 K_L^0 \pi^0$	$< 3.0 \times 10^{-4}$	CL=90%	DESIG=303
Г65	$K^+K^-\pi^0\pi^0$	$(2.6 \pm 1.3) \times 10^{-4}$		DESIG=298
Г66	$K^+K^-\pi^0\pi^0\pi^0$	$(6.6 \pm 2.8) \times 10^{-4}$		DESIG=341
Г67	$K_S^0 K^\pm \pi^\mp \pi^0 \pi^0$	$(1.7 \pm 0.6) \times 10^{-3}$		DESIG=342
Г68	$K_S^0 K^\pm \pi^\mp \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-3}$		DESIG=343
Г69	$K^+K^-\pi^+\pi^-\pi^0$	$(1.26 \pm 0.09) \times 10^{-3}$		DESIG=206
Г70	$\omega f_0(1710) \rightarrow \omega K^+K^-$	$(5.9 \pm 2.2) \times 10^{-5}$		DESIG=216
Г71	$K^*(892)^0 K^-\pi^+\pi^0 + \text{c.c.}$	$(8.6 \pm 2.2) \times 10^{-4}$		DESIG=217
Г72	$K^*(892)^+ K^-\pi^+\pi^- + \text{c.c.}$	$(9.6 \pm 2.8) \times 10^{-4}$		DESIG=218
Г73	$K^*(892)^+ K^-\rho^0 + \text{c.c.}$	$(7.3 \pm 2.6) \times 10^{-4}$		DESIG=219
Г74	$K^*(892)^0 K^-\rho^+ + \text{c.c.}$	$(6.1 \pm 1.8) \times 10^{-4}$		DESIG=220
Г75	$K_S^0 K_S^0 \pi^+ \pi^-$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=225
Г76	$K_S^0 K_L^0 \pi^0 \pi^0$	$(1.3 \pm 0.6) \times 10^{-3}$		DESIG=304
Г77	$K_S^0 K^*(892)^0 \pi^0 \pi^0$	$(3.0 \pm 1.3) \times 10^{-4}$		DESIG=348
Г78	$K_S^0 K^\pm \rho(770)^\mp \pi^0$	$< 7 \times 10^{-4}$	CL=90%	DESIG=352
Г79	$K_S^0 K^\pm \pi^\mp \rho(770)^0$	$< 7 \times 10^{-4}$	CL=90%	DESIG=353
Г80	$K^\mp K^*(892)^\pm \pi^0 \pi^0$	$(7.0 \pm 2.9) \times 10^{-4}$		DESIG=349
Г81	$K^*(892)^+ K^*(892)^- \pi^0$	$(3.6 \pm 1.8) \times 10^{-3}$		DESIG=350
Г82	$K_S^0 K_L^0 \eta$	$(1.3 \pm 0.5) \times 10^{-3}$		DESIG=305
Г83	$K^+K^-\rho^0$	$(2.2 \pm 0.4) \times 10^{-4}$		DESIG=205
Г84	$K^*(892)^0 \bar{K}_2^*(1430)^0$	$(1.9 \pm 0.5) \times 10^{-4}$		DESIG=66
Г85	$K^+K^-\pi^+\pi^-\eta$	$(1.3 \pm 0.7) \times 10^{-3}$		DESIG=252
Г86	$K^+K^-2(\pi^+\pi^-)$	$(1.9 \pm 0.9) \times 10^{-3}$		DESIG=222
Г87	$K^+K^-2(\pi^+\pi^-)\pi^0$	$(1.00 \pm 0.31) \times 10^{-3}$		DESIG=240
Г88	$K^+K^*(892)^- + \text{c.c.}$	$(2.9 \pm 0.4) \times 10^{-5}$	S=1.2	DESIG=39
Г89	$2(K^+K^-)$	$(6.3 \pm 1.3) \times 10^{-5}$		DESIG=208
Г90	$2(K^+K^-)\pi^0$	$(1.10 \pm 0.28) \times 10^{-4}$		DESIG=209
Г91	$K^+K^-\phi$	$(7.0 \pm 1.6) \times 10^{-5}$		DESIG=79
Г92	$K_S^0 K_S^0 \phi$	$(3.53 \pm 0.29) \times 10^{-5}$		DESIG=347
Г93	$K_1(1270)^\pm K^\mp$	$(1.00 \pm 0.28) \times 10^{-3}$		DESIG=41
Г94	$K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.}$	$(6.7 \pm 2.5) \times 10^{-4}$		DESIG=34
Г95	$\eta K^+K^-, \text{ no } \eta\phi$	$(3.49 \pm 0.17) \times 10^{-5}$		DESIG=207
Г96	ηK^+K^-	$< 2.6 \times 10^{-4}$	CL=90%	DESIG=351
Г97	$X(1750)\eta \rightarrow K^+K^-\eta$	$(4.8 \pm 2.8) \times 10^{-6}$		DESIG=324
Г98	$K_1(1400)^\pm K^\mp$	$< 3.1 \times 10^{-4}$	CL=90%	DESIG=42
Г99	$K_2^*(1430)^\pm K^\mp$	$(7.1 \pm 1.3 \mp 0.9) \times 10^{-5}$		DESIG=265
Г100	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	$(1.09 \pm 0.20) \times 10^{-4}$		DESIG=194
Г101	ωK^+K^-	$(1.62 \pm 0.11) \times 10^{-4}$	S=1.1	DESIG=76
Г102	$\omega K_S^0 K_S^0$	$(7.0 \pm 0.5) \times 10^{-5}$		DESIG=330
Г103	$\omega K^*(892)^+ K^- + \text{c.c.}$	$(2.07 \pm 0.26) \times 10^{-4}$		DESIG=276
Г104	$\omega K_2^*(1430)^+ K^- + \text{c.c.}$	$(6.1 \pm 1.2) \times 10^{-5}$		DESIG=277
Г105	$\omega \bar{K}^*(892)^0 K^0$	$(1.68 \pm 0.30) \times 10^{-4}$		DESIG=278
Г106	$\omega \bar{K}_2^*(1430)^0 K^0$	$(5.8 \pm 2.2) \times 10^{-5}$		DESIG=279
Г107	$\omega X(1440) \rightarrow \omega K_S^0 K^-\pi^+ +$ c.c.	$(1.6 \pm 0.4) \times 10^{-5}$		DESIG=282

Γ_{108}	$\omega X(1440) \rightarrow \omega K^+ K^- \pi^0$	$(1.09 \pm 0.26) \times 10^{-5}$		DESIG=283
Γ_{109}	$\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ +$ c.c.	$(3.0 \pm 1.0) \times 10^{-6}$		DESIG=284
Γ_{110}	$\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0$	$(1.2 \pm 0.7) \times 10^{-6}$		DESIG=285
Γ_{111}	$\rho \bar{\rho}$	$(2.94 \pm 0.09) \times 10^{-4}$	S=1.3	DESIG=27
Γ_{112}	$n \bar{n}$	$(3.06 \pm 0.15) \times 10^{-4}$		DESIG=309
Γ_{113}	$\rho \bar{\rho} \pi^0$	$(1.53 \pm 0.07) \times 10^{-4}$		DESIG=35
Γ_{114}	$N(940) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(6.4 \pm_{-1.3}^{+1.8}) \times 10^{-5}$		DESIG=267
Γ_{115}	$N(1440) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(7.3 \pm_{-1.5}^{+1.7}) \times 10^{-5}$	S=2.5	DESIG=261
Γ_{116}	$N(1520) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(6.4 \pm_{-1.8}^{+2.3}) \times 10^{-6}$		DESIG=268
Γ_{117}	$N(1535) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(2.5 \pm 1.0) \times 10^{-5}$		DESIG=269
Γ_{118}	$N(1650) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(3.8 \pm_{-1.7}^{+1.4}) \times 10^{-5}$		DESIG=270
Γ_{119}	$N(1720) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(1.79 \pm_{-0.70}^{+0.26}) \times 10^{-5}$		DESIG=271
Γ_{120}	$N(2300) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(2.6 \pm_{-0.7}^{+1.2}) \times 10^{-5}$		DESIG=272
Γ_{121}	$N(2570) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \pi^0$	$(2.13 \pm_{-0.31}^{+0.40}) \times 10^{-5}$		DESIG=273
Γ_{122}	$\rho \bar{\rho} \pi^+ \pi^-$	$(6.0 \pm 0.4) \times 10^{-4}$		DESIG=31
Γ_{123}	$\rho \bar{\rho} K^+ K^-$	$(2.7 \pm 0.7) \times 10^{-5}$		DESIG=212
Γ_{124}	$\rho \bar{\rho} \eta$	$(6.0 \pm 0.4) \times 10^{-5}$		DESIG=200
Γ_{125}	$N(1535) \bar{\rho} +$ c.c. $\rightarrow \rho \bar{\rho} \eta$	$(4.5 \pm_{-0.6}^{+0.7}) \times 10^{-5}$		DESIG=264
Γ_{126}	$\rho \bar{\rho} \pi^+ \pi^- \pi^0$	$(7.3 \pm 0.7) \times 10^{-4}$		DESIG=211
Γ_{127}	$\rho \bar{\rho} \rho^0$	$(5.0 \pm 2.2) \times 10^{-5}$		DESIG=210
Γ_{128}	$\rho \bar{\rho} \omega$	$(6.9 \pm 2.1) \times 10^{-5}$		DESIG=77
Γ_{129}	$\rho \bar{\rho} \eta'$	$(1.10 \pm 0.13) \times 10^{-5}$		DESIG=317
Γ_{130}	$\rho \bar{\rho} \phi$	$(6.1 \pm 0.6) \times 10^{-6}$		DESIG=80
Γ_{131}	$\phi X(1835) \rightarrow \rho \bar{\rho} \phi$	$< 1.82 \times 10^{-7}$	CL=90%	DESIG=318
Γ_{132}	$\rho \bar{n} \pi^-$ or c.c.	$(2.48 \pm 0.17) \times 10^{-4}$		DESIG=227
Γ_{133}	$\rho \bar{n} \pi^- \pi^0$	$(3.2 \pm 0.7) \times 10^{-4}$		DESIG=228
Γ_{134}	$\Lambda \bar{\Lambda}$	$(3.81 \pm 0.13) \times 10^{-4}$	S=1.4	DESIG=28
Γ_{135}	$\Lambda \bar{\Lambda} \pi^0$	$(1.4 \pm 0.7) \times 10^{-6}$		DESIG=238
Γ_{136}	$\Lambda \bar{\Lambda} \eta$	$(2.43 \pm 0.32) \times 10^{-5}$		DESIG=239
Γ_{137}	$\Lambda \bar{\Lambda} \eta'$	$(7.3 \pm 1.0) \times 10^{-6}$		DESIG=346
Γ_{138}	$\Lambda(1670) \bar{\Lambda} \rightarrow \Lambda \bar{\Lambda} \eta$	$(1.3 \pm 0.7) \times 10^{-5}$		DESIG=336
Γ_{139}	$\Lambda \bar{\Lambda} \omega(782)$	$(3.3 \pm 0.4) \times 10^{-5}$		DESIG=340
Γ_{140}	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	$(2.8 \pm 0.6) \times 10^{-4}$		DESIG=213
Γ_{141}	$\Lambda \bar{\rho} K^+$	$(1.00 \pm 0.14) \times 10^{-4}$		DESIG=214
Γ_{142}	$\Lambda \bar{\rho} K^*(892)^+ +$ c.c.	$(6.3 \pm 0.7) \times 10^{-5}$		DESIG=321
Γ_{143}	$\Lambda \bar{\rho} K^+ \pi^+ \pi^-$	$(1.8 \pm 0.4) \times 10^{-4}$		DESIG=215
Γ_{144}	$\bar{\Lambda} n K_S^0 +$ c.c.	$(8.1 \pm 1.8) \times 10^{-5}$		DESIG=237
Γ_{145}	$\Delta^{++} \bar{\Delta}^{--}$	$(1.28 \pm 0.35) \times 10^{-4}$		DESIG=70
Γ_{146}	$\Lambda \bar{\Sigma}^+ \pi^- +$ c.c.	$(1.40 \pm 0.13) \times 10^{-4}$		DESIG=280
Γ_{147}	$\Lambda \bar{\Sigma}^- \pi^+ +$ c.c.	$(1.54 \pm 0.14) \times 10^{-4}$		DESIG=281
Γ_{148}	$\Lambda \bar{\Sigma}^0 +$ c.c.	$(1.6 \pm 0.7) \times 10^{-6}$		DESIG=326
Γ_{149}	$\Lambda \bar{\Sigma}^0$			DESIG=307
Γ_{150}	$\Sigma^0 \bar{\rho} K^+ +$ c.c.	$(1.67 \pm 0.18) \times 10^{-5}$		DESIG=274
Γ_{151}	$\Sigma^+ \bar{\Sigma}^-$	$(2.43 \pm 0.10) \times 10^{-4}$	S=1.4	DESIG=223
Γ_{152}	$\Sigma^0 \bar{\Sigma}^0$	$(2.35 \pm 0.09) \times 10^{-4}$	S=1.1	DESIG=71
Γ_{153}	$\Sigma^- \bar{\Sigma}^+$	$(2.82 \pm 0.09) \times 10^{-4}$		DESIG=335
Γ_{154}	$\Sigma^+ \bar{\Sigma}^- \eta$	$(9.6 \pm 2.4) \times 10^{-6}$		DESIG=339
Γ_{155}	$\Sigma^+ \bar{\Sigma}^- \omega$	$(1.89 \pm 0.28) \times 10^{-5}$		DESIG=344
Γ_{156}	$\Sigma^+ \bar{\Sigma}^- \phi$	$(3.0 \pm 0.7) \times 10^{-6}$		DESIG=345
Γ_{157}	$\Sigma(1385)^+ \bar{\Sigma}(1385)^-$	$(8.5 \pm 0.7) \times 10^{-5}$		DESIG=72
Γ_{158}	$\Sigma(1385)^- \bar{\Sigma}(1385)^+$	$(8.5 \pm 0.8) \times 10^{-5}$		DESIG=297
Γ_{159}	$\Sigma(1385)^0 \bar{\Sigma}(1385)^0$	$(6.9 \pm 0.7) \times 10^{-5}$		DESIG=299

Γ ₁₆₀	$\Xi^- \Xi^+$	$(2.87 \pm 0.11) \times 10^{-4}$	S=1.1	DESIG=29
Γ ₁₆₁	$\Xi^0 \Xi^0$	$(2.3 \pm 0.4) \times 10^{-4}$	S=4.2	DESIG=224
Γ ₁₆₂	$\Xi(1530)^0 \Xi(1530)^0$	$(6.8 \pm 0.4) \times 10^{-5}$		DESIG=73
Γ ₁₆₃	$\Lambda \Xi^+ K^- + \text{c.c.}$	$(3.9 \pm 0.4) \times 10^{-5}$		DESIG=293
Γ ₁₆₄	$\Xi(1530)^- \Xi(1530)^+$	$(1.15 \pm 0.07) \times 10^{-4}$		DESIG=322
Γ ₁₆₅	$\Xi(1530)^- \Xi^+$	$(7.0 \pm 1.2) \times 10^{-6}$		DESIG=323
Γ ₁₆₆	$\Xi(1530)^0 \Xi^0$	$(5.3 \pm 0.5) \times 10^{-6}$		DESIG=331
Γ ₁₆₇	$\Xi(1690)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ +$ c.c.	$(5.2 \pm 1.6) \times 10^{-6}$		DESIG=294
Γ ₁₆₈	$\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ +$ c.c.	$(1.20 \pm 0.32) \times 10^{-5}$		DESIG=295
Γ ₁₆₉	$\Sigma^0 \Xi^+ K^- + \text{c.c.}$	$(3.7 \pm 0.4) \times 10^{-5}$		DESIG=296
Γ ₁₇₀	$\Omega^- \bar{\Omega}^+$	$(5.66 \pm 0.30) \times 10^{-5}$	S=1.3	DESIG=74
Γ ₁₇₁	$\eta_c \pi^+ \pi^- \pi^0$	$< 1.0 \times 10^{-3}$	CL=90%	DESIG=229
Γ ₁₇₂	$h_c(1P) \pi^0$	$(7.4 \pm 0.5) \times 10^{-4}$		DESIG=254
Γ ₁₇₃	$\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.}$	$< 1.7 \times 10^{-6}$	CL=90%	DESIG=310
Γ ₁₇₄	$\Theta(1540) \bar{\Theta}(1540) \rightarrow$ $K_S^0 p K^- \bar{n} + \text{c.c.}$	[a] $< 8.8 \times 10^{-6}$	CL=90%	DESIG=195
Γ ₁₇₅	$\Theta(1540) K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n}$	[a] $< 1.0 \times 10^{-5}$	CL=90%	DESIG=196
Γ ₁₇₆	$\Theta(1540) K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n$	[a] $< 7.0 \times 10^{-6}$	CL=90%	DESIG=197
Γ ₁₇₇	$\bar{\Theta}(1540) K^+ n \rightarrow K_S^0 \bar{p} K^+ n$	[a] $< 2.6 \times 10^{-5}$	CL=90%	DESIG=198
Γ ₁₇₈	$\bar{\Theta}(1540) K_S^0 p \rightarrow K_S^0 p K^- \bar{n}$	[a] $< 6.0 \times 10^{-6}$	CL=90%	DESIG=199

Radiative decays

Γ ₁₇₉	$\gamma \chi_{c0}(1P)$	$(9.77 \pm 0.23) \%$	S=1.1	DESIG=56
Γ ₁₈₀	$\gamma \chi_{c1}(1P)$	$(9.75 \pm 0.27) \%$	S=1.1	DESIG=58
Γ ₁₈₁	$\gamma \chi_{c2}(1P)$	$(9.36 \pm 0.23) \%$	S=1.2	DESIG=59
Γ ₁₈₂	$\gamma \eta_c(1S)$	$(3.6 \pm 0.5) \times 10^{-3}$	S=1.3	DESIG=61
Γ ₁₈₃	$\gamma \eta_c(2S)$	$(7 \pm 5) \times 10^{-4}$		DESIG=63
Γ ₁₈₄	$\gamma \pi^0$	$(1.04 \pm 0.22) \times 10^{-6}$	S=1.4	DESIG=52
Γ ₁₈₅	$\gamma 2(\pi^+ \pi^-)$	$(4.0 \pm 0.6) \times 10^{-4}$		DESIG=241
Γ ₁₈₆	$\gamma 3(\pi^+ \pi^-)$	$< 1.7 \times 10^{-4}$	CL=90%	DESIG=249
Γ ₁₈₇	$\gamma \eta'(958)$	$(1.24 \pm 0.04) \times 10^{-4}$		DESIG=54
Γ ₁₈₈	$\gamma f_2(1270)$	$(2.73 \pm_{-0.25}^{+0.29}) \times 10^{-4}$	S=1.8	DESIG=82
Γ ₁₈₉	$\gamma f_0(1370) \rightarrow \gamma K \bar{K}$	$(3.1 \pm 1.7) \times 10^{-5}$		DESIG=286
Γ ₁₉₀	$\gamma f_0(1500)$	$(9.3 \pm 1.9) \times 10^{-5}$		DESIG=287
Γ ₁₉₁	$\gamma f_2'(1525)$	$(3.3 \pm 0.8) \times 10^{-5}$		DESIG=288
Γ ₁₉₂	$\gamma f_0(1710)$			DESIG=236
Γ ₁₉₃	$\gamma f_0(1710) \rightarrow \gamma \pi \pi$	$(3.5 \pm 0.6) \times 10^{-5}$		DESIG=83
Γ ₁₉₄	$\gamma f_0(1710) \rightarrow \gamma K \bar{K}$	$(6.6 \pm 0.7) \times 10^{-5}$		DESIG=84
Γ ₁₉₅	$\gamma f_0(2100) \rightarrow \gamma \pi \pi$	$(4.8 \pm 1.0) \times 10^{-6}$		DESIG=289
Γ ₁₉₆	$\gamma f_0(2200) \rightarrow \gamma K \bar{K}$	$(3.2 \pm 1.0) \times 10^{-6}$		DESIG=290
Γ ₁₉₇	$\gamma f_J(2220) \rightarrow \gamma \pi \pi$	$< 5.8 \times 10^{-6}$	CL=90%	DESIG=291
Γ ₁₉₈	$\gamma f_J(2220) \rightarrow \gamma K \bar{K}$	$< 9.5 \times 10^{-6}$	CL=90%	DESIG=292
Γ ₁₉₉	$\gamma \eta$	$(9.2 \pm 1.8) \times 10^{-7}$		DESIG=53
Γ ₂₀₀	$\gamma \eta \pi^+ \pi^-$	$(8.7 \pm 2.1) \times 10^{-4}$		DESIG=230
Γ ₂₀₁	$\gamma \eta(1405)$			DESIG=231
Γ ₂₀₂	$\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi$	$< 9 \times 10^{-5}$	CL=90%	DESIG=62
Γ ₂₀₃	$\gamma \eta(1405) \rightarrow \gamma \eta \pi^+ \pi^-$	$(3.6 \pm 2.5) \times 10^{-5}$		DESIG=232
Γ ₂₀₄	$\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow$ $\gamma \pi^+ \pi^- \pi^0$	$< 5.0 \times 10^{-7}$	CL=90%	DESIG=308
Γ ₂₀₅	$\gamma \eta(1475)$			DESIG=233
Γ ₂₀₆	$\gamma \eta(1475) \rightarrow \gamma K \bar{K} \pi$	$< 1.4 \times 10^{-4}$	CL=90%	DESIG=234
Γ ₂₀₇	$\gamma \eta(1475) \rightarrow \gamma \eta \pi^+ \pi^-$	$< 8.8 \times 10^{-5}$	CL=90%	DESIG=235
Γ ₂₀₈	$\gamma K^{*0} K^+ \pi^- + \text{c.c.}$	$(3.7 \pm 0.9) \times 10^{-4}$		DESIG=242
Γ ₂₀₉	$\gamma K^{*0} \bar{K}^{*0}$	$(2.4 \pm 0.7) \times 10^{-4}$		DESIG=243

NODE=M071;CLUMP=C

Γ_{210}	$\gamma K_S^0 K^+ \pi^- + \text{c.c.}$	$(2.6 \pm 0.5) \times 10^{-4}$		DESIG=244
Γ_{211}	$\gamma K^+ K^- \pi^+ \pi^-$	$(1.9 \pm 0.5) \times 10^{-4}$		DESIG=245
Γ_{212}	$\gamma K^+ K^- 2(\pi^+ \pi^-)$	$< 2.2 \times 10^{-4}$	CL=90%	DESIG=248
Γ_{213}	$\gamma 2(K^+ K^-)$	$< 4 \times 10^{-5}$	CL=90%	DESIG=250
Γ_{214}	$\gamma p \bar{p}$	$(3.9 \pm 0.5) \times 10^{-5}$	S=2.0	DESIG=246
Γ_{215}	$\gamma f_2(1950) \rightarrow \gamma p \bar{p}$	$(1.20 \pm 0.22) \times 10^{-5}$		DESIG=257
Γ_{216}	$\gamma f_2(2150) \rightarrow \gamma p \bar{p}$	$(7.2 \pm 1.8) \times 10^{-6}$		DESIG=258
Γ_{217}	$\gamma X(1835) \rightarrow \gamma p \bar{p}$	$(4.6 \pm_{-4.0}^{+1.8}) \times 10^{-6}$		DESIG=259
Γ_{218}	$\gamma X \rightarrow \gamma p \bar{p}$	[b] $< 2 \times 10^{-6}$	CL=90%	DESIG=260
Γ_{219}	$\gamma p \bar{p} \pi^+ \pi^-$	$(2.8 \pm 1.4) \times 10^{-5}$		DESIG=247
Γ_{220}	$\gamma \gamma$	$< 1.5 \times 10^{-4}$	CL=90%	DESIG=51
Γ_{221}	$\gamma \gamma J/\psi$	$(3.1 \pm_{-1.2}^{+1.0}) \times 10^{-4}$		DESIG=266
Γ_{222}	$e^+ e^- \eta'$	$(1.90 \pm 0.26) \times 10^{-6}$		DESIG=311
Γ_{223}	$e^+ e^- \eta_c(1S)$	$(3.8 \pm 0.4) \times 10^{-5}$		DESIG=338
Γ_{224}	$e^+ e^- \chi_{c0}(1P)$	$(1.06 \pm 0.25) \times 10^{-3}$		DESIG=300
Γ_{225}	$e^+ e^- \chi_{c1}(1P)$	$(8.5 \pm 0.7) \times 10^{-4}$		DESIG=301
Γ_{226}	$e^+ e^- \chi_{c2}(1P)$	$(6.8 \pm 0.8) \times 10^{-4}$		DESIG=302
Weak decays				
Γ_{227}	$D^0 e^+ e^- + \text{c.c.}$	$< 1.4 \times 10^{-7}$	CL=90%	NODE=M071;CLUMP=E DESIG=306
Γ_{228}	$\Lambda_c^+ \bar{\Sigma}^- + \text{c.c.}$	$< 1.4 \times 10^{-5}$	CL=90%	DESIG=337
Other decays				
Γ_{229}	invisible	$< 1.6 \%$	CL=90%	NODE=M071;CLUMP=D DESIG=275
[a] $\Theta(1540)$ is a hypothetical pentaquark state of $1.54 \text{ GeV}/c^2$ mass and a width of less than $25 \text{ MeV}/c^2$.				
[b] For a narrow resonance in the range $2.2 < M(X) < 2.8 \text{ GeV}$.				

CONSTRAINED FIT INFORMATION

A multiparticle fit to $\chi_{c1}(1P)$, $\chi_{c0}(1P)$, $\chi_{c2}(1P)$, and $\psi(2S)$ with 4 total widths, a partial width, 25 combinations of partial widths obtained from integrated cross section, and 86 branching ratios uses 253 measurements to determine 49 parameters. The overall fit has a $\chi^2 = 389.6$ for 204 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \cdot \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$.

x_8	3									
x_9	0	0								
x_{12}	21	13	2							
x_{13}	22	5	1	28						
x_{14}	11	6	1	44	11					
x_{111}	0	0	0	3	2	1				
x_{179}	0	0	0	2	1	1	0			
x_{180}	1	0	0	2	0	1	0	0		
x_{181}	1	1	0	4	0	2	0	0	0	
Γ	-85	-4	-1	-29	-31	-15	-4	-1	-1	-2
	x_7	x_8	x_9	x_{12}	x_{13}	x_{14}	x_{111}	x_{179}	x_{180}	x_{181}

FIT INFORMATION

A multiparticle fit to $\eta_c(1S)$, $J/\psi(1S)$, $\psi(2S)$, $h_c(1P)$, and B^\pm with the total width, 10 combinations of partial widths obtained from integrated cross section, and 38 branching ratios uses 113 measurements to determine 19 parameters. The overall fit has a $\chi^2 = 184.6$ for 94 degrees of freedom.

$\psi(2S)$ PARTIAL WIDTHS

NODE=M071225

 $\Gamma(\text{hadrons})$ Γ_1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
258±26	BAI	02B	BES2 e^+e^-
224±56	LUTH	75	MRK1 e^+e^-

NODE=M071W3
NODE=M071W3 $\Gamma(e^+e^-)$ Γ_7

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.33 ±0.04 OUR FIT	Error includes scale factor of 1.1. [2.33 ± 0.04 keV OUR 2023 FIT]		
2.29 ±0.06 OUR AVERAGE			
2.23 ±0.10 ±0.02	¹ ABLIKIM	15V	BES3 4.0–4.4 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
2.338±0.037±0.096	ABLIKIM	08B	BES2 $e^+e^- \rightarrow \text{hadrons}$
2.330±0.036±0.110	ABLIKIM	06L	BES2 $e^+e^- \rightarrow \text{hadrons}$
2.44 ±0.21	² BAI	02B	BES2 e^+e^-
2.14 ±0.21	ALEXANDER	89	RVUE See \mathcal{T} mini-review
2.279±0.015±0.042	³ ANASHIN	18	KEDR e^+e^-
2.282±0.015±0.042	⁴ ANASHIN	18	KEDR e^+e^-
2.0 ±0.3	BRANDELIK	79C	DASP e^+e^-
2.1 ±0.3	⁵ LUTH	75	MRK1 e^+e^-

NODE=M071W1
NODE=M071W1
NEW

OCCUR=2

¹ ABLIKIM 15V reports $2.213 \pm 0.018 \pm 0.099$ keV from a measurement of $[\Gamma(\psi(2S) \rightarrow e^+e^-)] \times [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)]$ assuming $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.95 \pm 0.45) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071W1;LINKAGE=A

² From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channel, assuming $\Gamma_e = \Gamma_\mu = \Gamma_\tau/0.38847$.

NODE=M071W;LINKAGE=BB

³ Combining $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$ from ANASHIN 18 with $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$ from ANASHIN 12 and assuming lepton universality.

NODE=M071W1;LINKAGE=B

⁴ From the sum of $\Gamma_{e^+e^-} \cdot B(\text{hadrons})$ from ANASHIN 12, $\Gamma_{e^+e^-} \cdot B(e^+e^-)$ and $\Gamma_{e^+e^-} \cdot B(\mu^+\mu^-)$ from ANASHIN 18, and $\Gamma_{e^+e^-} \cdot B(\tau^+\tau^-)$ from ANASHIN 07.

NODE=M071W1;LINKAGE=C

⁵ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

NODE=M071W1;LINKAGE=F

 $\Gamma(\gamma\gamma)$ Γ_{220}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<43	90	BRANDELIK	79C	DASP e^+e^-

NODE=M071W51
NODE=M071W51 $\psi(2S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M071230

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel(i) in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(i)$ or the branching ratio $\Gamma(i)/\text{total}$.

NODE=M071230

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_7/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.233±0.015±0.042	¹ ANASHIN	12	KEDR $e^+e^- \rightarrow \text{hadrons}$
2.2 ±0.4	ABRAMS	75	MRK1 e^+e^-

NODE=M071G3
NODE=M071G3

¹ ANASHIN 12 reports the value $2.233 \pm 0.015 \pm 0.037 \pm 0.020$ keV, where the third uncertainty is due to assumptions on the interference between the resonance and hadronic continuum. We combined the two systematic uncertainties.

NODE=M071G3;LINKAGE=AN

 $\Gamma(K_S^0 \text{ anything}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_6\Gamma_7/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.3738±0.0067±0.0200	ABLIKIM	21S	BES3 $e^+e^- \rightarrow K_S^0 \text{ anything}$

NODE=M071P30
NODE=M071P30 $\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_7/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
21.2±0.7±1.2	¹ ANASHIN	18	KEDR e^+e^-

NODE=M071P14
NODE=M071P14

¹ From the average of nine scans of the $\psi(2S)$.

NODE=M071P14;LINKAGE=A

$$\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_8\Gamma_7/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
19.3±0.3±0.5	¹ ANASHIN	18	KEDR $\psi(2S) \rightarrow \mu^+\mu^-$

¹ From the average of nine scans of the $\psi(2S)$.

NODE=M071P13
NODE=M071P13

NODE=M071P13;LINKAGE=A

$$\Gamma(\tau^+\tau^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_9\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.0±2.6	79	¹ ANASHIN	07	KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9.0±2.6 79 ¹ ANASHIN 07 KEDR $e^+e^- \rightarrow \psi(2S) \rightarrow \tau^+\tau^-$

¹ Using $\psi(2S)$ total width of 337 ± 13 keV. Systematic errors not evaluated.

NODE=M071G9
NODE=M071G9

NODE=M071G9;LINKAGE=AN

$$\Gamma(J/\psi(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{12}\Gamma_7/\Gamma$$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.808±0.014 OUR FIT				Error includes scale factor of 1.1. [0.808±0.013 keV OUR 2023 FIT]
0.836±0.025 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below. [0.837±0.025 keV OUR 2023 AVERAGE Scale factor = 1.3]
0.78 ±0.12 ±0.07		¹ LEES	23	BABR $e^+e^- \rightarrow \gamma_{ISR}$ hadrons
0.837±0.028±0.005		² LEES	12E	BABR $10.6 e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$
0.852±0.010±0.026	19.5k	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$
0.68 ±0.09		³ BAI	98E	BES e^+e^-
0.93 ±0.08 ±0.03	256	⁴ AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
0.755±0.048±0.004	544	⁵ AUBERT	05D	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$

0.808±0.014 OUR FIT Error includes scale factor of 1.1. [0.808±0.013 keV OUR 2023 FIT]

0.836±0.025 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below. [0.837±0.025 keV OUR 2023 AVERAGE Scale factor = 1.3]

0.78 ±0.12 ±0.07 ¹ LEES 23 BABR $e^+e^- \rightarrow \gamma_{ISR}$ hadrons

0.837±0.028±0.005 ² LEES 12E BABR $10.6 e^+e^- \rightarrow 2\pi^+2\pi^-\gamma$

0.852±0.010±0.026 19.5k ADAM 06 CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$

0.68 ±0.09 ³ BAI 98E BES e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.93 ±0.08 ±0.03 256 ⁴ AUBERT 07AU BABR $10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

0.755±0.048±0.004 544 ⁵ AUBERT 05D BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-\gamma$

¹ LEES 23 reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow K_S^0 K^\pm \pi^\mp)] = (4.14 \pm 0.55 \pm 0.29) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² LEES 12E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = (49.9 \pm 1.3 \pm 1.0) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ The value of $\Gamma(e^+e^-)$ quoted in BAI 98E is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (32.4 \pm 2.6) \times 10^{-2}$ and $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1203 \pm 0.0038$. Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.

⁴ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0186 \pm 0.0012 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.00 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ AUBERT 05D reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(J/\psi(1S) \rightarrow \mu^+\mu^-)] = 0.0450 \pm 0.0018 \pm 0.0022$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by LEES 12E.

NODE=M071G1
NODE=M071G1

NEW

NEW


NODE=M071G1;LINKAGE=B

NODE=M071G1;LINKAGE=LE

NODE=M071G1;LINKAGE=A

NODE=M071G1;LINKAGE=UB

NODE=M071G1;LINKAGE=AU



$\Gamma(J/\psi(1S)\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{13}\Gamma/\Gamma$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.425±0.012 OUR FIT				Error includes scale factor of 1.4. [0.425±0.009 keV OUR 2023 FIT]

NODE=M071G6
NODE=M071G6
NEW
NEW

0.413±0.019 OUR AVERAGE

[0.411 ± 0.020 keV OUR 2023 AVERAGE]

0.45 ±0.13 ±0.02	1	LEES	23	BABR	$e^+e^- \rightarrow \gamma_{ISR} \text{hadrons}$
0.45 ±0.12 ±0.04	2	LEES	23	BABR	$e^+e^- \rightarrow \gamma_{ISR} \text{hadrons}$
0.411±0.008±0.018 3.6k		ADAM	06	CLEO	$3.773 e^+e^- \rightarrow \gamma\psi(2S)$

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.51 ±0.09 ±0.02	142	3 LEES	18E	BABR	$10.6 e^+e^- \rightarrow J/\psi\pi^0\pi^0\gamma$
------------------	-----	--------	-----	------	--

¹ LEES 23 reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow K^+K^-\pi^0)] = (1.31 \pm 0.35 \pm 0.13) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow K^+K^-\pi^0) = (2.88 \pm 0.12) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G6;LINKAGE=C

² LEES 23 reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow K_S^0 K^\pm\pi^\mp)] = (2.36 \pm 0.59 \pm 0.24) \times 10^{-3}$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow K_S^0 K^\pm\pi^\mp) = (5.3 \pm 0.5) \times 10^{-3}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G6;LINKAGE=D

³ LEES 18E reports $[\Gamma(\psi(2S) \rightarrow J/\psi(1S)\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{total}] \times [B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0)] = 0.0101 \pm 0.0015 \pm 0.0011$ keV which we divide by our best value $B(J/\psi(1S) \rightarrow \pi^+\pi^-\pi^0) = (2.00 \pm 0.07) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G6;LINKAGE=A

$\Gamma(J/\psi(1S)\eta) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{14}\Gamma/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
78.6± 1.8 OUR FIT				Error includes scale factor of 1.1. [78.6 ± 1.6 eV OUR 2023 FIT]
87 ± 9 OUR AVERAGE				

NODE=M071G7
NODE=M071G7
NEW

83 ±25 ±5	14	1 AUBERT	07AU	BABR	$10.6 e^+e^- \rightarrow J/\psi\pi^+\pi^-\pi^0\gamma$
88 ± 6 ±7	291 ± 24	ADAM	06	CLEO	$3.773 e^+e^- \rightarrow \gamma\psi(2S)$

¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow J/\psi\eta) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot B(\eta \rightarrow \pi^+\pi^-\pi^0) = 1.11 \pm 0.33 \pm 0.07$ eV.

NODE=M071G7;LINKAGE=UB

$\Gamma(J/\psi(1S)\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{15}\Gamma/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8	90	<37	ADAM	06	CLEO $3.773 e^+e^- \rightarrow \gamma\psi(2S)$

NODE=M071G8
NODE=M071G8

$\Gamma(2(\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_{22}\Gamma/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
29.7±2.2±1.8	410	AUBERT	07AU	BABR $10.6 e^+e^- \rightarrow 2(\pi^+\pi^-\pi^0)\gamma$

NODE=M071G01
NODE=M071G01

$$\Gamma(\pi^+\pi^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{24}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±1.8±1.2	177	LEES	18E	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ

NODE=M071P16
NODE=M071P16

$$\Gamma(\pi^+\pi^-4\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{25}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.3±2.3±0.5	18	LEES	21C	BABR e ⁺ e ⁻ → γ _{ISR} (π ⁺ π ⁻ 4π ⁰)

NODE=M071P36
NODE=M071P36

$$\Gamma(\rho^\pm\pi^\mp\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{26}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<6.2	90	LEES	18E	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ 3π ⁰ γ

NODE=M071P18
NODE=M071P18

$$\Gamma(2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{28}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
11.2±3.3±1.3	43	AUBERT	06D	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻ π ⁰)γ

NODE=M071G4
NODE=M071G4

$$\Gamma(2(\pi^+\pi^-)3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{30}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
33±5±5	14k	LEES	21	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

NODE=M071P31
NODE=M071P31

$$\Gamma(\eta 2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{33}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.87±1.41±0.01		16	¹ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)ηγ

NODE=M071G03
NODE=M071G03

••• We do not use the following data for averages, fits, limits, etc. •••

<7	90	14k	² LEES	21	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ
----	----	-----	-------------------	----	--

¹AUBERT 07AU reports [Γ(ψ(2S) → η2(π⁺π⁻)) × Γ(ψ(2S) → e⁺e⁻)/Γ_{total}] × [B(η → 2γ)] = 1.13 ± 0.55 ± 0.08 eV which we divide by our best value B(η → 2γ) = (39.36 ± 0.18) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G03;LINKAGE=UB

²LEES 21 reports [Γ(ψ(2S) → η2(π⁺π⁻)) × Γ(ψ(2S) → e⁺e⁻)/Γ_{total}] × [B(η → 3π⁰)] < 2.3 eV which we divide by our best value B(η → 3π⁰) = 32.57 × 10⁻².

NODE=M071G03;LINKAGE=A

$$\Gamma(\eta\pi^+\pi^-\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{34}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.85	90	LEES	18E	BABR 10.6 e ⁺ e ⁻ → π ⁺ π ⁻ π ⁰ π ⁰ ηγ

NODE=M071P15
NODE=M071P15

$$\Gamma(\eta\pi^+\pi^-3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{35}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ LEES	21C	BABR e ⁺ e ⁻ → γ _{ISR} (π ⁺ π ⁻ 3π ⁰ γγ)

NODE=M071P40
NODE=M071P40

¹LEES 21C reports [Γ(ψ(2S) → ηπ⁺π⁻3π⁰) × Γ(ψ(2S) → e⁺e⁻)/Γ_{total}] × [B(η → 2γ)] < 1.9 eV which we divide by our best value B(η → 2γ) = 39.36 × 10⁻².

NODE=M071P40;LINKAGE=A

$$\Gamma(\eta 2(\pi^+\pi^-\pi^0)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{36}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<5	90	14k	¹ LEES	21	BABR 10.6 e ⁺ e ⁻ → 2(π ⁺ π ⁻)3π ⁰ γ

NODE=M071P33
NODE=M071P33

¹LEES 21 reports [Γ(ψ(2S) → η2(π⁺π⁻π⁰)) × Γ(ψ(2S) → e⁺e⁻)/Γ_{total}] × [B(η → 2γ)] < 1.9 eV which we divide by our best value B(η → 2γ) = 39.36 × 10⁻².

NODE=M071P33;LINKAGE=A

$$\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{41}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.01±0.84±0.02	37	¹ AUBERT	07AU	BABR 10.6 e ⁺ e ⁻ → ωπ ⁺ π ⁻ γ

NODE=M071G02
NODE=M071G02

¹AUBERT 07AU reports [Γ(ψ(2S) → ωπ⁺π⁻) × Γ(ψ(2S) → e⁺e⁻)/Γ_{total}] × [B(ω(782) → π⁺π⁻π⁰)] = 2.69 ± 0.73 ± 0.16 eV which we divide by our best value B(ω(782) → π⁺π⁻π⁰) = (89.2 ± 0.7) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G02;LINKAGE=UB

$$\Gamma(\omega\pi^+\pi^-2\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{42}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
20.2±5.6±0.1	14k	¹ LEES	21	BABR 10.6 $e^+e^- \rightarrow \frac{2(\pi^+\pi^-)3\pi^0\gamma$

NODE=M071P32
NODE=M071P32

¹ LEES 21 reports [$\Gamma(\psi(2S) \rightarrow \omega\pi^+\pi^-2\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$] = $18 \pm 4 \pm 3$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P32;LINKAGE=A

$$\Gamma(\omega\pi^0\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{45}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.58±0.82±0.02	33	¹ LEES	18E	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-3\pi^0\gamma$

NODE=M071P17
NODE=M071P17

¹ LEES 18E reports [$\Gamma(\psi(2S) \rightarrow \omega\pi^0\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$] = $2.3 \pm 0.7 \pm 0.2$ eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = (89.2 \pm 0.7) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P17;LINKAGE=A

$$\Gamma(\omega3\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{46}\Gamma_7/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	¹ LEES	21C	BABR $e^+e^- \rightarrow \gamma_{ISR}(\pi^+\pi^-4\pi^0)$

NODE=M071P39
NODE=M071P39

¹ LEES 21C reports [$\Gamma(\psi(2S) \rightarrow \omega3\pi^0) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\omega(782) \rightarrow \pi^+\pi^-\pi^0)$] < 1.6 eV which we divide by our best value $B(\omega(782) \rightarrow \pi^+\pi^-\pi^0) = 89.2 \times 10^{-2}$.

NODE=M071P39;LINKAGE=A

$$\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{51}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.55±0.19±0.01	19	¹ LEES	12F	BABR 10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$

NODE=M071G10
NODE=M071G10

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.57±0.23±0.01	10	² AUBERT, BE 06D	BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
----------------	----	-----------------------------	------	--

¹ LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\phi(1020) \rightarrow K^+K^-)$] = $0.27 \pm 0.09 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G10;LINKAGE=A

² Superseded by LEES 12F. AUBERT, BE 06D reports [$\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\phi(1020) \rightarrow K^+K^-)$] = $0.28 \pm 0.11 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G10;LINKAGE=AU

$$\Gamma(\phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{52}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.346±0.129±0.004	12	¹ LEES	12F	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

NODE=M071G13
NODE=M071G13

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.346±0.168±0.004	6±3	² AUBERT	07AK	BABR 10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
-------------------	-----	---------------------	------	---

¹ LEES 12F reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\phi(1020) \rightarrow K^+K^-)$] = $0.17 \pm 0.06 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G13;LINKAGE=A

² Superseded by LEES 12F. AUBERT 07AK reports [$\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+\pi^-) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}$] \times [$B(\phi(1020) \rightarrow K^+K^-)$] = $0.17 \pm 0.08 \pm 0.02$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = (49.1 \pm 0.5) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G13;LINKAGE=AU

$$\Gamma(K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{59}\Gamma_7/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.147±0.035±0.005	66	¹ LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.197±0.035±0.005	66	² LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.35±0.14±0.03	11	³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

NODE=M071G06
NODE=M071G06

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.147±0.035±0.005	66	¹ LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.197±0.035±0.005	66	² LEES	15J	BABR $e^+e^- \rightarrow K^+K^-\gamma$
0.35±0.14±0.03	11	³ LEES	13Q	BABR $e^+e^- \rightarrow K^+K^-\gamma$

OCCUR=2

¹ $\sin\phi > 0$.

² $\sin\phi < 0$.

³ Interference with non-resonant K^+K^- production not taken into account.

NODE=M071G06;LINKAGE=A
NODE=M071G06;LINKAGE=B
NODE=M071G06;LINKAGE=BA

$$\Gamma(K^+ K^- \pi^+) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{60} \Gamma_7 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.92±0.30±0.06	133	LEES 12F	BABR	10.6 e ⁺ e ⁻ → π ⁺ π ⁻ K ⁺ K ⁻ γ

NODE=M071G12
NODE=M071G12

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.56±0.42±0.16 85 ¹AUBERT 07AK BABR 10.6 e⁺e⁻ → π⁺π⁻K⁺K⁻γ

¹ Superseded by LEES 12F.

NODE=M071G12;LINKAGE=A

$$\Gamma(K_S^0 K_L^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{64} \Gamma_7 / \Gamma$$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.7	90	8	LEES 17A	BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ γ

NODE=M071G15
NODE=M071G15

$$\Gamma(K^+ K^- \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{65} \Gamma_7 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.60±0.31±0.03	17	LEES 12F	BABR	10.6 e ⁺ e ⁻ → π ⁰ π ⁰ K ⁺ K ⁻ γ

NODE=M071G08
NODE=M071G08

$$\Gamma(K^+ K^- \pi^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{66} \Gamma_7 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.54±0.63±0.15	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P47
NODE=M071P47

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{67} \Gamma_7 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
4.0±1.4±0.4	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P48
NODE=M071P48

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{68} \Gamma_7 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
5.1±0.7±0.4	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P49
NODE=M071P49

$$\Gamma(K^+ K^- \pi^+ \pi^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{69} \Gamma_7 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±1.3±0.3	32	AUBERT 07AU	BABR	10.6 e ⁺ e ⁻ → K ⁺ K ⁻ π ⁺ π ⁻ π ⁰ γ

NODE=M071G04
NODE=M071G04

$$\Gamma(K_S^0 K_L^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{76} \Gamma_7 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.92±1.27±0.15	14	LEES 17A	BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ π ⁰ π ⁰ γ

NODE=M071G14
NODE=M071G14

$$\Gamma(K_S^0 K^*(892)^0 \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{77} \Gamma_7 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
0.71±0.29±0.07	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P54
NODE=M071P54

$$\Gamma(K_S^0 K^\pm \rho(770)^\mp \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{78} \Gamma_7 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P58
NODE=M071P58

$$\Gamma(K_S^0 K^\pm \pi^\mp \rho(770)^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{79} \Gamma_7 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P59
NODE=M071P59

$$\Gamma(K^*(892)^+ K^*(892)^- \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{81} \Gamma_7 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
8.46±4.05±0.90	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P56
NODE=M071P56

$$\Gamma(K^\mp K^*(892)^\pm \pi^0 \pi^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{80} \Gamma_7 / \Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.62±0.66±0.15	LEES 23	BABR	e ⁺ e ⁻ → γ _{ISR} hadrons

NODE=M071P55
NODE=M071P55

$$\Gamma(K_S^0 K_L^0 \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{82} \Gamma_7 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.14±1.08±0.16	16	LEES 17A	BABR	e ⁺ e ⁻ → K _S ⁰ K _L ⁰ ηγ

NODE=M071G16
NODE=M071G16

$\Gamma(K^+K^-\pi^+\pi^-\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{85}\Gamma/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
3.05±1.80±0.01	7	¹ AUBERT 07AU BABR	10.6	$e^+e^- \rightarrow K^+K^-\pi^+\pi^-\eta\gamma$

NODE=M071G05
NODE=M071G05

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+K^-\pi^+\pi^-\eta) \times \Gamma(\psi(2S) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 2\gamma)] = 1.2 \pm 0.7 \pm 0.1$ eV which we divide by our best value $B(\eta \rightarrow 2\gamma) = (39.36 \pm 0.18) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071G05;LINKAGE=UB

 $\Gamma(K^+K^-2(\pi^+\pi^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{86}\Gamma/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.4±2.1±0.3	26	AUBERT 06D BABR	10.6	$e^+e^- \rightarrow K^+K^-2(\pi^+\pi^-)\gamma$

NODE=M071G5
NODE=M071G5

 $\Gamma(2(K^+K^-)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{89}\Gamma/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.22±0.10±0.02	13	LEES 12F BABR	10.6	$e^+e^- \rightarrow K^+K^-K^+K^-\gamma$

NODE=M071G07
NODE=M071G07

 $\Gamma(p\bar{p}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{111}\Gamma/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.686±0.024 OUR FIT				Error includes scale factor of 1.2. [0.686 ± 0.019 eV OUR 2023 FIT]

NODE=M071G2
NODE=M071G2

NEW

0.63 ± 0.05 OUR AVERAGE Error includes scale factor of 1.2.

0.67 ± 0.12 ± 0.02	43	¹ LEES 130 BABR		$e^+e^- \rightarrow p\bar{p}\gamma$
0.74 ± 0.07 ± 0.04	142	² LEES 13Y BABR		$e^+e^- \rightarrow p\bar{p}\gamma$
0.579 ± 0.038 ± 0.036	2.7k	ANDREOTTI 07 E835		$p\bar{p} \rightarrow e^+e^-, J/\psi X$
0.70 ± 0.17 ± 0.03	22	³ AUBERT 06B BABR		$e^+e^- \rightarrow p\bar{p}\gamma$

¹ ISR photon reconstructed in the detector

² ISR photon undetected

³ Superseded by LEES 130

NODE=M071G2;LINKAGE=C
NODE=M071G2;LINKAGE=B
NODE=M071G2;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{134}\Gamma/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
1.5±0.4±0.1	AUBERT 07BD BABR	10.6	$e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$

NODE=M071G11
NODE=M071G11

 $\psi(2S)$ BRANCHING RATIOS

NODE=M071235

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.9785±0.0013 OUR AVERAGE			
0.9779±0.0015	¹ BAI 02B BES2		e^+e^-
0.981 ± 0.003	¹ LUTH 75 MRK1		e^+e^-

NODE=M071R3
NODE=M071R3

¹ Includes cascade decay into $J/\psi(1S)$.

NODE=M071R;LINKAGE=P

 $\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0179±0.0004 OUR AVERAGE	[0.0173 ± 0.0014 OUR 2023 AVERAGE Scale factor = 1.5]		

NODE=M071R5
NODE=M071R5

NEW

0.0179±0.0004
• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0166±0.0010	^{2,3} SETH 04 RVUE		e^+e^-
0.0199±0.0019	² BAI 02B BES2		e^+e^-
0.029 ± 0.004	² LUTH 75 MRK1		e^+e^-

¹ Using $B(\psi(2S) \rightarrow \ell^+\ell^-) = (0.794 \pm 0.017)\%$ and $R = 2.26 \pm 0.01$ determined by a fit to data from Mark-I, DM2, BESII, KEDR, and BESIII.

² Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.

³ Using $B(\psi(2S) \rightarrow \ell^+\ell^-) = (0.73 \pm 0.04)\%$ from RPP-2002 and $R = 2.28 \pm 0.04$ determined by a fit to data from BAI 00 and BAI 02C. Superseded by LIAO 23.

NODE=M071R5;LINKAGE=A

NODE=M071R;LINKAGE=Z

NODE=M071R5;LINKAGE=SE

 $\Gamma(gg)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
10.58±1.62	2.9 M	¹ LIBBY 09 CLEO		$\psi(2S) \rightarrow \text{hadrons}$

NODE=M071S43
NODE=M071S43

¹ Calculated using $\Gamma(\gamma gg)/\Gamma(gg) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09, $B(\psi(2S) \rightarrow XJ/\psi)$ relative and absolute branching fractions from MENDEZ 08, $B(\psi(2S) \rightarrow \gamma\eta_c)$ from MITCHELL 09, and $B(\psi(2S) \rightarrow \text{virtual } \gamma \rightarrow \text{hadrons})$, $B(\psi(2S) \rightarrow \gamma\chi_{cJ})$, and $B(\psi(2S) \rightarrow \ell^+\ell^-)$ from PDG 08. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

NODE=M071S43;LINKAGE=LI

$\Gamma(\gamma g g)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.025 ± 0.288	200 k	¹ LIBBY	09	CLEO $\psi(2S) \rightarrow \gamma + \text{hadrons}$

NODE=M071S44
 NODE=M071S44

¹ Calculated using $\Gamma(\gamma g g)/\Gamma(g g g) = 0.097 \pm 0.026 \pm 0.016$ from LIBBY 09. The statistical error is negligible and the systematic error is largely uncorrelated with that of $\Gamma(g g g)/\Gamma_{\text{total}}$ LIBBY 09 measurement.

NODE=M071S44;LINKAGE=LI

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_4/Γ_3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.7 ± 2.6 ± 1.6	2.9 M	LIBBY	09	CLEO $\psi(2S) \rightarrow (\gamma +) \text{hadrons}$

NODE=M071S45
 NODE=M071S45

 $\Gamma(\text{light hadrons})/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.154 ± 0.015	¹ MENDEZ	08	CLEO $e^+ e^- \rightarrow \psi(2S)$

NODE=M071S27
 NODE=M071S27

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.169 ± 0.026 ² ADAM 05A CLEO $e^+ e^- \rightarrow \psi(2S)$

¹ Uses $B(\psi(2S) \rightarrow J/\psi X)$ from MENDEZ 08 and other branching fractions from PDG 07.

NODE=M071S27;LINKAGE=ME

² Uses $B(J/\psi X)$ from ADAM 05A, $B(\chi_{c,J} \gamma)$, $B(\eta_c \gamma)$ from ATHAR 04 and $B(\ell^+ \ell^-)$ from PDG 04. Superseded by MENDEZ 08.

NODE=M071S27;LINKAGE=AD

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
79.4 ± 2.2 OUR FIT	Error includes scale factor of 1.3. $[(79.3 \pm 1.7) \times 10^{-4}]$ OUR 2023 FIT		

NODE=M071R1
 NODE=M071R1

• • • We do not use the following data for averages, fits, limits, etc. • • •

88 ± 13 ¹ FELDMAN 77 RVUE $e^+ e^-$

NEW

¹ From an overall fit assuming equal partial widths for $e^+ e^-$ and $\mu^+ \mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.

NODE=M071R;LINKAGE=L

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	DOCUMENT ID
80 ± 6 OUR FIT	

NODE=M071R2
 NODE=M071R2

 $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$ Γ_8/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
1.00 ± 0.08 OUR FIT	[1.00 ± 0.08 OUR 2023 FIT]		

NODE=M071R4
 NODE=M071R4
 NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.89 ± 0.16 BOYARSKI 75C MRK1 $e^+ e^-$

 $\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
31 ± 4 OUR FIT			
30.8 ± 2.1 ± 3.8	¹ ABLIKIM	06W	BES $e^+ e^- \rightarrow \psi(2S)$

NODE=M071R75
 NODE=M071R75

¹ Computed using PDG 02 value of $B(\psi(2S) \rightarrow \text{hadrons}) = 0.9810 \pm 0.0030$ to estimate the total number of $\psi(2S)$ events.

NODE=M071R75;LINKAGE=AB

————— **DECAYS INTO $J/\psi(1S)$ AND ANYTHING** —————

NODE=M071305

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$

$$\Gamma_{10}/\Gamma = (\Gamma_{12} + \Gamma_{13} + \Gamma_{14} + 0.343\Gamma_{180} + 0.195\Gamma_{181})/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.615 ± 0.007 OUR FIT	Error includes scale factor of 1.3. $[0.614 \pm 0.006]$ OUR 2023 FIT			

NODE=M071R10
 NODE=M071R10
 NEW

0.55 ± 0.07 OUR AVERAGE

0.51 ± 0.12	BRANDELIK	79C	DASP	$e^+ e^- \rightarrow \mu^+ \mu^- X$
0.57 ± 0.08	ABRAMS	75B	MRK1	$e^+ e^- \rightarrow \mu^+ \mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.644 ± 0.006 ± 0.016	¹ ABLIKIM	21Z	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- X$
0.6254 ± 0.0016 ± 0.0155	1.1M	² MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+ \ell^- X$
0.5950 ± 0.0015 ± 0.0190	151k	ADAM	05A	CLEO Repl. by MENDEZ 08

¹ From a fit to the $e^+ e^- \rightarrow J/\psi X$ cross section between 3.645 and 3.891 GeV, with $\Gamma(ee)$ and Γ fixed to the PDG 20 values of the cross particle fit which are correlated to "OUR FIT" value for $B(\psi(2S) \rightarrow J/\psi X)$.

NODE=M071R10;LINKAGE=A

² Not independent from other measurements of MENDEZ 08.

NODE=M071R10;LINKAGE=ME

$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\text{anything})$


Γ_7/Γ_{10}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.291 ± 0.035 OUR FIT				Error includes scale factor of 1.3. [(1.291 ± 0.026) × 10 ⁻² OUR 2023 FIT]
1.28 ± 0.04 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
1.22 ± 0.02 ± 0.05	5097 ± 73	¹ ANDREOTTI 05	E835	$p\bar{p} \rightarrow \psi(2S) \rightarrow e^+e^-$
1.28 ± 0.03 ± 0.02		¹ AMBROGIANI 00A	E835	$p\bar{p} \rightarrow \psi(2S)$
1.44 ± 0.08 ± 0.02		¹ ARMSTRONG 97	E760	$\bar{p}p \rightarrow \psi(2S)$
¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.				

NODE=M071R72
NODE=M071R72

NEW

NODE=M071R;LINKAGE=7A



$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\text{anything})$

Γ_8/Γ_{10}

VALUE	DOCUMENT ID	TECN	COMMENT
0.0130 ± 0.0010 OUR FIT			
0.014 ± 0.003	HILGER	75	SPEC e^+e^-

NODE=M071R74
NODE=M071R74

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$

Γ_{11}/Γ

VALUE	DOCUMENT ID
0.254 ± 0.005 OUR FIT	
	Error includes scale factor of 1.6. [0.2538 ± 0.0032 OUR 2023 FIT]

NODE=M071R18
NODE=M071R18

NEW

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$


Γ_{12}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.3469 ± 0.0034 OUR FIT				Error includes scale factor of 1.1. [0.3468 ± 0.0030 OUR 2023 FIT]
0.348 ± 0.005 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.3498 ± 0.0002 ± 0.0045	20M	ABLIKIM	13R	BES3 $\psi(2S) \rightarrow J/\psi\pi^+\pi^-$
0.3504 ± 0.0007 ± 0.0077	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.323 ± 0.014		BAI	02B	BES2 e^+e^-
0.32 ± 0.04		ABRAMS	75B	MRK1 $e^+e^- \rightarrow J/\psi\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.3354 ± 0.0014 ± 0.0110	60k	¹ ADAM	05A	CLEO Repl. by MENDEZ 08
¹ Not independent from other values reported by ADAM 05A.				

NODE=M071R12
NODE=M071R12

NEW

NODE=M071R;LINKAGE=AD



$\Gamma(e^+e^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_7/Γ_{12}

VALUE	DOCUMENT ID	TECN	COMMENT
0.0229±0.0006 OUR FIT			Error includes scale factor of 1.3. [0.0229 ± 0.0005 OUR 2023 FIT]
0.0252±0.0028±0.0011	¹ AUBERT	02B	BABR e^+e^-

¹ Using $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0593 \pm 0.0010$.

NODE=M071R73
NODE=M071R73
NEW

NODE=M071R73;LINKAGE=7A

$\Gamma(\mu^+\mu^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_8/Γ_{12}

VALUE	DOCUMENT ID	TECN	COMMENT
0.0230±0.0017 OUR FIT			
0.0228±0.0018 OUR AVERAGE			
0.0230±0.0020±0.0012	¹ AAIJ	16Y	LHCB $\Lambda_b^0 \rightarrow \psi(2S)X$
0.0216±0.0026±0.0014	² AUBERT	02B	BABR e^+e^-
0.0327±0.0077±0.0072	² GRIBUSHIN	96	FMPS $515 \pi^- Be \rightarrow 2\mu X$

¹ Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033) \times 10^{-2}$.
² Using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10) \times 10^{-2}$.

NODE=M071R63
NODE=M071R63

NODE=M071R63;LINKAGE=A
NODE=M071R;LINKAGE=Q2

$\Gamma(\tau^+\tau^-)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_9/Γ_{12}

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
8.8 ±1.1 OUR FIT			
8.73±1.39±1.57	BAI	02	BES e^+e^-

NODE=M071R76
NODE=M071R76

$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{12}/Γ_{10}


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.564 ±0.004 OUR FIT				Error includes scale factor of 1.7. [0.5645 ± 0.0026 OUR 2023 FIT]
0.554 ±0.008 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.5604±0.0009±0.0062	565k	MENDEZ	08	CLEO $\psi(2S) \rightarrow \ell^+\ell^-\pi^+\pi^-$
0.525 ±0.009 ±0.022	4k	ANDREOTTI	05	E835 $\psi(2S) \rightarrow J/\psi X$
0.536 ±0.007 ±0.016	20k	^{1,2} ABLIKIM	04B	BES $\psi(2S) \rightarrow J/\psi X$
0.496 ±0.037		ARMSTRONG	97	E760 $\bar{p}p \rightarrow \psi(2S)$
0.5637±0.0027±0.0046	60k	ADAM	05A	CLEO Repl. by MENDEZ 08

NODE=M071R70
NODE=M071R70
NEW

¹ From a fit to the J/ψ recoil mass spectra.

² ABLIKIM 04B quotes $B(\psi(2S) \rightarrow J/\psi X) / B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-)$.

NODE=M071R;LINKAGE=AB
NODE=M071R;LINKAGE=AL



$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_{11}/\Gamma_{12} = (0.9761\Gamma_{13} + 0.719\Gamma_{14} + 0.343\Gamma_{180} + 0.195\Gamma_{181})/\Gamma_{12}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.732 ± 0.013 OUR FIT	Error includes scale factor of 1.7.	[0.732 ± 0.008 OUR 2023 FIT]	

NODE=M071R11
NODE=M071R11
NEW

0.73 ± 0.09 TANENBAUM 76 MRK1 e^+e^-

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\text{anything})$ Γ_{13}/Γ_{10}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.297 ± 0.005 OUR FIT		Error includes scale factor of 1.7.	[0.2968 ± 0.0031 OUR 2023 FIT]	

NODE=M071R69
NODE=M071R69
NEW

0.320 ± 0.012 OUR AVERAGE

0.300 ± 0.008 ± 0.022	1655 ± 44	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.328 ± 0.013 ± 0.008		AMBROGIANI	00A	E835	$p\bar{p} \rightarrow \psi(2S)$
0.323 ± 0.033		ARMSTRONG	97	E760	$p\bar{p} \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.2829 ± 0.0012 ± 0.0056	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.2776 ± 0.0025 ± 0.0043	13.4k	ADAM	05A	CLEO	Repl. by MENDEZ 08

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{13}/Γ_{12}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.526 ± 0.013 OUR FIT		Error includes scale factor of 1.7.	[0.526 ± 0.008 OUR 2023 FIT]	

NODE=M071R14
NODE=M071R14
NEW

0.513 ± 0.022 OUR AVERAGE Error includes scale factor of 2.2.

0.5047 ± 0.0022 ± 0.0102	61k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\pi^0$
0.570 ± 0.009 ± 0.026	14k	¹ ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.4924 ± 0.0047 ± 0.0086	73k	^{2,3} ADAM	05A	CLEO	Repl. by MENDEZ 08
0.571 ± 0.018 ± 0.044		⁴ ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.53 ± 0.06		TANENBAUM	76	MRK1	e^+e^-
0.64 ± 0.15		⁵ HILGER	75	SPEC	e^+e^-

¹ From a fit to the J/ψ recoil mass spectra.

² Not independent from other values reported by ADAM 05A.

³ Using 13,217 $J/\psi\pi^0\pi^0$ and 60,010 $J/\psi\pi^+\pi^-$ events.

⁴ Not independent from other values reported by ANDREOTTI 05.

⁵ Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.

NODE=M071R14;LINKAGE=AB
NODE=M071R14;LINKAGE=AD
NODE=M071R14;LINKAGE=AM
NODE=M071R;LINKAGE=AN
NODE=M071R;LINKAGE=I

$\Gamma(J/\psi(1S)\eta)/\Gamma_{total}$

Γ_{14}/Γ

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

33.7 ± 0.6 OUR FIT Error includes scale factor of 1.2. [(33.7 ± 0.5) × 10⁻³ OUR 2023 FIT]

32.9 ± 1.7 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.

33.75 ± 0.17 ± 0.86	68.2k	ABLIKIM	12M	BES3	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$
29.8 ± 0.9 ± 2.3	5.7k	BAI	04I	BES2	$\psi(2S) \rightarrow J/\psi\gamma\gamma$
25.5 ± 2.9	386	¹ OREGLIA	80	CBAL	$e^+e^- \rightarrow J/\psi2\gamma$
45 ± 12	17	² BRANDELIK	79B	DASP	$e^+e^- \rightarrow J/\psi2\gamma$
42 ± 6	164	² BARTEL	78B	CNTR	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •


34.3 ± 0.4 ± 0.9	18.4k	³ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-\eta$
32.5 ± 0.6 ± 1.1	2.8k	⁴ ADAM	05A	CLEO	Repl. by MENDEZ 08
43 ± 8	44	TANENBAUM	76	MRK1	e^+e^-

- ¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+\ell^-) = 0.1181 \pm 0.0020$.
- ² Recalculated by us using $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$.
- ³ Not independent from other measurements of MENDEZ 08.
- ⁴ Not independent from other values reported by ADAM 05A.

NODE=M071R15
NODE=M071R15

NEW

NODE=M071R;LINKAGE=3Q
NODE=M071R;LINKAGE=2Q
NODE=M071R15;LINKAGE=ME
NODE=M071R15;LINKAGE=AD



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\text{anything})$

Γ_{14}/Γ_{10}

VALUE EVTS DOCUMENT ID TECN COMMENT

0.0549 ± 0.0009 OUR FIT Error includes scale factor of 1.2. [0.0549 ± 0.0008 OUR 2023 FIT]

0.058 ± 0.007 OUR AVERAGE Error includes scale factor of 1.4. See the ideogram below.

0.050 ± 0.006 ± 0.003	298 ± 20	ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$
0.072 ± 0.009		AMBROGIANI	00A	E835	$\rho\bar{\rho} \rightarrow \psi(2S)$
0.061 ± 0.015		ARMSTRONG	97	E760	$\bar{p}p \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0549 ± 0.0006 ± 0.0009	18.4k	¹ MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+\ell^-\eta$
0.0546 ± 0.0010 ± 0.0007	2.8k	ADAM	05A	CLEO	Repl. by MENDEZ 08


- ¹ Not independent from other measurements of MENDEZ 08.

NODE=M071R68
NODE=M071R68

NEW

OCCUR=2

NODE=M071R68;LINKAGE=ME



$\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$

Γ_{14}/Γ_{12}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0972±0.0016 OUR FIT				Error includes scale factor of 1.1. [0.0972 ± 0.0014 OUR 2023 FIT]

NODE=M071R71
NODE=M071R71
NEW

0.0979±0.0018 OUR AVERAGE

0.0979±0.0010±0.0015	18.4k	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- \eta$
0.098 ±0.005 ±0.010	2k	¹ ABLIKIM	04B	BES	$\psi(2S) \rightarrow J/\psi X$
0.091 ±0.021		² HIMEL	80	MRK2	$e^+ e^- \rightarrow \psi(2S) X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.0968±0.0019±0.0013	2.8k	³ ADAM	05A	CLEO	Repl. by MENDEZ 08
0.095 ±0.007 ±0.007		⁴ ANDREOTTI	05	E835	$\psi(2S) \rightarrow J/\psi X$

¹ From a fit to the J/ψ recoil mass spectra.

² The value for $B(\psi(2S) \rightarrow J/\psi(1S)\eta)$ reported in HIMEL 80 is derived using $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (33 \pm 3)\%$ and $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.138 \pm 0.018$. Calculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = (0.1181 \pm 0.0020)$.

³ Not independent from other values reported by ADAM 05A.

⁴ Not independent from other values reported by ANDREOTTI 05.

NODE=M071R71;LINKAGE=AB
NODE=M071R;LINKAGE=8H

NODE=M071R71;LINKAGE=AD
NODE=M071R71;LINKAGE=AN

$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{total}$

Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12.68±0.32 OUR AVERAGE				

NODE=M071R16
NODE=M071R16

12.6 ±0.2 ±0.3	4.1k	ABLIKIM	12M	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$
13.3 ±0.8 ±0.3	530	MENDEZ	08	CLEO	$\psi(2S) \rightarrow \ell^+ \ell^- 2\gamma$
14.3 ±1.4 ±1.2	280	BAI	04i	BES2	$\psi(2S) \rightarrow J/\psi \gamma \gamma$
14 ±6	7	HIMEL	80	MRK2	$e^+ e^-$
9 ±2 ±1	23	¹ OREGLIA	80	CBAL	$\psi(2S) \rightarrow J/\psi 2\gamma$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
13 ±1 ±1	88	ADAM	05A	CLEO	Repl. by MENDEZ 08

¹ Recalculated by us using $B(J/\psi(1S) \rightarrow \ell^+ \ell^-) = 0.1181 \pm 0.0020$.

NODE=M071R16;LINKAGE=3Q

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)anything)$

$\Gamma_{15}/\Gamma_{10} = \Gamma_{15}/(\Gamma_{12}+\Gamma_{13}+\Gamma_{14}+0.343\Gamma_{180}+0.195\Gamma_{181})$

NODE=M071S25
NODE=M071S25

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				

0.213±0.012±0.003	527	¹ MENDEZ	08	CLEO	$e^+ e^- \rightarrow J/\psi \gamma \gamma$
0.22 ±0.02 ±0.01		² ADAM	05A	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow J/\psi \gamma \gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.

² Not independent from other values reported by ADAM 05A.

NODE=M071S25;LINKAGE=ME
NODE=M071S25;LINKAGE=AD

$\Gamma(J/\psi(1S)\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_{15}/Γ_{12}

VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.380 \pm 0.022 \pm 0.005$	527	¹ MENDEZ	08	CLEO	$e^+e^- \rightarrow J/\psi\gamma\gamma$
$0.39 \pm 0.04 \pm 0.01$		² ADAM	05A	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow J/\psi\gamma\gamma$

¹ Not independent from other values reported by MENDEZ 08. Supersedes ADAM 05A.
² Not independent from other values reported by ADAM 05A.

NODE=M071S26
 NODE=M071S26

NODE=M071S26;LINKAGE=ME
 NODE=M071S26;LINKAGE=AD

NODE=M071310

———— HADRONIC DECAYS ————

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ Γ_{16}/Γ

VALUE (units 10^{-5}) CL% EVTS DOCUMENT ID TECN COMMENT

0.78 ± 0.26 OUR AVERAGE

$0.76 \pm 0.25 \pm 0.06$	30	¹ METREVELI	12		$\psi(2S) \rightarrow \pi^+\pi^-$
8 ± 5		BRANDELIK	79C	DASP	e^+e^-
<2.1	90	DOBBS	06A	CLEO	$e^+e^- \rightarrow \psi(2S)$
<5	90	FELDMAN	77	MRK1	e^+e^-

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration. Using $\psi(3770) \rightarrow \pi^+\pi^-$ for continuum subtraction.

NODE=M071R20
 NODE=M071R20

NODE=M071R20;LINKAGE=ME

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{17}/Γ

VALUE (units 10^{-4}) EVTS DOCUMENT ID TECN COMMENT

2.01 ± 0.17 OUR AVERAGE Error includes scale factor of 1.7. See the ideogram below.


$2.14 \pm 0.03^{+0.12}_{-0.11}$	7k	¹ ABLIKIM	12H	BES3	$e^+e^- \rightarrow \psi(2S)$
$1.81 \pm 0.18 \pm 0.19$	260 ± 19	² ABLIKIM	05J	BES2	$e^+e^- \rightarrow \psi(2S)$
$1.88^{+0.16}_{-0.15} \pm 0.28$	194	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$
0.85 ± 0.46	4	FRANKLIN	83	MRK2	$e^+e^- \rightarrow$ hadrons

¹ From $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$ events directly. The quoted systematic error includes a contribution of 4% (added in quadrature) from the uncertainty on the number of $\psi(2S)$ events.
² From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

NODE=M071R36
 NODE=M071R36

NODE=M071R36;LINKAGE=AB

NODE=M071R;LINKAGE=AK



$\Gamma(\rho(770)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{18}/Γ

VALUE (units 10^{-4}) CL% EVTS DOCUMENT ID TECN COMMENT

0.32 ± 0.12 OUR AVERAGE Error includes scale factor of 1.8.

$0.51 \pm 0.07 \pm 0.11$		¹ ABLIKIM	05J	BES2	$\psi(2S) \rightarrow \rho(770)\pi \rightarrow \pi^+\pi^-\pi^0$
$0.24^{+0.08}_{-0.07} \pm 0.02$	22	ADAM	05	CLEO	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R26
 NODE=M071R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.83	90	1	FRANKLIN	83	MRK2	e^+e^-
<10	90		BARTEL	76	CNTR	e^+e^-
<10	90	2	ABRAMS	75	MRK1	e^+e^-

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

² Final state $\rho^0\pi^0$.

NODE=M071R26;LINKAGE=AK
NODE=M071R;LINKAGE=N

$\Gamma(\rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{19}/Γ

<u>VALUE (units 10^{-4})</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	--------------------	-------------	----------------

1.94 ± 0.25^{+1.15}_{-0.34} ¹ ABLIKIM 05J BES2 $\psi(2S) \rightarrow \rho(2150)\pi \rightarrow \pi^+\pi^-\pi^0$

NODE=M071R57
NODE=M071R57

¹ From a PW analysis of $\psi(2S) \rightarrow \pi^+\pi^-\pi^0$.

NODE=M071R57;LINKAGE=AK

$\Gamma(2(\pi^+\pi^-))/\Gamma_{total}$ Γ_{20}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.4 ± 0.6 OUR AVERAGE Error includes scale factor of 2.2.

2.2 ± 0.2 ± 0.2	308	BRIERE	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$
4.5 ± 1.0		TANENBAUM	78	MRK1	e^+e^-

NODE=M071R27
NODE=M071R27

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{21}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

2.2 ± 0.6 OUR AVERAGE Error includes scale factor of 1.4.

2.0 ± 0.2 ± 0.4	285.5	BRIERE	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)$
4.2 ± 1.5		TANENBAUM	78	MRK1	e^+e^-

NODE=M071R33
NODE=M071R33

$\Gamma(2(\pi^+\pi^-)\pi^0)/\Gamma_{total}$ Γ_{22}/Γ

<u>VALUE (units 10^{-4})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---	-------------	--------------------	-------------	----------------

29 ± 10 OUR AVERAGE Error includes scale factor of 4.7. See the ideogram below.

[(29 ± 10) × 10⁻⁴ OUR 2023 AVERAGE Scale factor = 4.7]


24.9 ± 0.7 ± 3.6	2173	ABLIKIM	07D	BES2	$e^+e^- \rightarrow \psi(2S)$
127 ± 12 ± 2	410	¹ AUBERT	07AU	BABR	10.6 $e^+e^- \rightarrow 2(\pi^+\pi^-)\pi^0\gamma$
26.1 ± 0.7 ± 3.0	1703	BRIERE	05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
30 ± 8	42	FRANKLIN	83	MRK2	e^+e^-

NODE=M071R22
NODE=M071R22

NEW

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0)/\Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (297 \pm 22 \pm 18) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R22;LINKAGE=UB



$\Gamma(\rho\pi^0)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.55±0.73±0.47		112 ± 31	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3		90	BAI	98J BES	e^+e^-

NODE=M071R65
NODE=M071R65 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ±2.0 OUR AVERAGE		Error includes scale factor of 2.8.		
5.45±0.42±0.87	671	ABLIKIM	05H BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow 3(\pi^+\pi^-)$
1.5 ±1.0		¹ TANENBAUM	78 MRK1	e^+e^-

NODE=M071R32
NODE=M071R32¹ Assuming entirely strong decay.

NODE=M071R32;LINKAGE=K

 $\Gamma(3(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
35±16	6	FRANKLIN	83 MRK2	$e^+e^- \rightarrow \text{hadrons}$

NODE=M071R37
NODE=M071R37 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M071S06
NODE=M071S06 $\Gamma(\eta\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
9.5±0.7±1.5		¹ BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
10.3±0.8±1.4	201.7	² BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow \gamma\gamma)$	
8.1±1.4±1.6	50.0	² BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \eta 3\pi(\eta \rightarrow 3\pi)$	

NODE=M071S07
NODE=M071S07¹ Average of $\eta \rightarrow \gamma\gamma$ and $\eta \rightarrow 3\pi$.² Not independent from other values reported by BRIERE 05.NODE=M071S07;LINKAGE=BR
NODE=M071S07;LINKAGE=BI $\Gamma(\rho\eta)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ±0.6 OUR AVERAGE		Error includes scale factor of 1.1.		
3.0 $^{+1.1}_{-0.9}$ ±0.2	18	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
1.78 $^{+0.67}_{-0.62}$ ±0.17	13	ABLIKIM	04L BES	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R94
NODE=M071R94 $\Gamma(\eta'\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.5±1.6±1.3	12.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadr}$

NODE=M071S08
NODE=M071S08 $\Gamma(\eta'\rho)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.87 $^{+1.64}_{-1.11}$ ±0.33	2	ABLIKIM	04L BES	$e^+e^- \rightarrow \psi(2S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.02 ±0.11 ±0.24	143	¹ ABLIKIM	17AK BES3	$e^+e^- \rightarrow \psi(2S)$	
0.569±0.128±0.236	80	² ABLIKIM	17AK BES3	$e^+e^- \rightarrow \psi(2S)$	

NODE=M071R93
NODE=M071R93¹ Destructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

NODE=M071R93;LINKAGE=A

² Constructive-interference solution of a partial wave analysis of the decay $\psi(2S) \rightarrow \pi^+\pi^-\eta'$.

NODE=M071R93;LINKAGE=B

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1 ±0.6 OUR AVERAGE				
2.5 $^{+1.2}_{-1.0}$ ±0.2	14	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
1.87 $^{+0.68}_{-0.62}$ ±0.28	14	ABLIKIM	04L BES	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R92
NODE=M071R92

$\Gamma(\omega\pi^+\pi^-)/\Gamma_{total}$

Γ_{41}/Γ


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
8.4±0.5±1.2	386	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
12.2±2.2±0.7	37	¹ AUBERT	07AU BABR	10.6 $e^+e^- \rightarrow \omega\pi^+\pi^-\gamma$
8.2±0.5±0.7	391	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
4.8±0.6±0.7	100 ± 22	² BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow \omega\pi^+\pi^-) \cdot B(\omega \rightarrow 3\pi) = 2.69 \pm 0.73 \pm 0.16$ eV.				
² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

NODE=M071R77
NODE=M071R77

NODE=M071R77;LINKAGE=UB

NODE=M071R77;LINKAGE=B3

WEIGHTED AVERAGE
7.3±1.2 (Error scaled by 2.1)



$\Gamma(b_1^\pm\pi^\mp)/\Gamma_{total}$

Γ_{43}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ±0.6 OUR AVERAGE				Error includes scale factor of 1.1.
5.1 ±0.6 ±0.8	202	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
4.18 ^{+0.43} _{-0.42} ±0.92	170	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$
3.2 ±0.6 ±0.5	61 ± 11	^{1,2} BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5.2 ±0.8 ±1.0		¹ BAI	99C BES	Repl. by BAI 03B
¹ Assuming $B(b_1 \rightarrow \omega\pi)=1$.				
² Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.				

NODE=M071R40
NODE=M071R40

NODE=M071R;LINKAGE=M1
NODE=M071R40;LINKAGE=B3

$\Gamma(\omega f_2(1270))/\Gamma_{total}$

Γ_{44}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.2 ±0.4 OUR AVERAGE					
2.3 ±0.5 ±0.4		57	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$
2.05±0.41±0.38		62±12	BAI	04C BES2	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<1.5		90	¹ BAI	03B BES	$\psi(2S) \rightarrow 2(\pi^+\pi^-)\pi^0$
<1.7		90	BAI	98J BES	Repl. by BAI 03B
¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.					

NODE=M071R64
NODE=M071R64

NODE=M071R64;LINKAGE=B3

$\Gamma(b_1^0\pi^0)/\Gamma_{total}$

Γ_{47}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35^{+0.47}_{-0.42} ±0.40	45	ADAM	05 CLEO	$e^+e^- \rightarrow \psi(2S)$

NODE=M071R21
NODE=M071R21

$\Gamma(\omega\eta)/\Gamma_{total}$ Γ_{48}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<3.1	90	ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$

NODE=M071R95
NODE=M071R95

$\Gamma(\omega\eta')/\Gamma_{total}$ Γ_{49}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.2^{+2.4}_{-2.0} \pm 0.7$	4	¹ ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$

NODE=M071R91
NODE=M071R91

¹ Calculated combining $\eta' \rightarrow \gamma\rho$ and $\eta\pi^+\pi^-$ channels.

NODE=M071R91;LINKAGE=AI

$\Gamma(\phi\pi^0)/\Gamma_{total}$ Γ_{50}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	ABLIKIM	12L	BES3 $e^+e^- \rightarrow \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.7	90	ADAM	05	CLEO $e^+e^- \rightarrow \psi(2S)$
<0.4	90	ABLIKIM	04K	BES $e^+e^- \rightarrow \psi(2S)$

NODE=M071R96
NODE=M071R96

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{total}$ Γ_{51}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.18 ± 0.26 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
2.3 ± 0.8 ± 0.1	19 ± 6	LEES	12F	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
0.9 ± 0.2 ± 0.1	47.6	BRIERE	05	CLEO $e^+e^- \rightarrow \psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
1.5 ± 0.2 ± 0.2	51.5 ± 8.3	¹ BAI	03B	BES $\psi(2S) \rightarrow K^+K^-\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.45 ± 0.96 ± 0.04	10 ± 4	^{2,3} AUBERT	07AK	BABR $10.6 e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$

NODE=M071R80
NODE=M071R80


¹ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R80;LINKAGE=B3
NODE=M071R80;LINKAGE=BE

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi\pi^+\pi^-)/\Gamma_{total}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (0.57 \pm 0.22 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+K^-) = (49.3 \pm 0.6)\%$.

NODE=M071R80;LINKAGE=UB



$\Gamma(\phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{52} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.75 ± 0.33 OUR AVERAGE		Error	includes scale factor of 1.6.	
1.5 ± 0.5 ± 0.1	12 ± 4	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$
0.6 ± 0.2 ± 0.1	18.4 ± 6.4	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.46 ± 0.71 ± 0.02	6 ± 3	^{2,3} AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow \phi f_0(980) \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (0.34 \pm 0.16 \pm 0.04) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(\phi \rightarrow K^+ K^-) = (49.3 \pm 0.6)\%$.

NODE=M071R83
NODE=M071R83

NODE=M071R83;LINKAGE=B3
NODE=M071R83;LINKAGE=BE

NODE=M071R83;LINKAGE=UB

 $\Gamma(\phi \eta) / \Gamma_{\text{total}}$ Γ_{53} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.10 ± 0.31 OUR AVERAGE				
3.14 ± 0.23 ± 0.23	0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.0 $^{+1.5}_{-1.1}$ ± 0.4	6	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
3.3 ± 1.1 ± 0.5	17	ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R89
NODE=M071R89

 $\Gamma(\eta \phi(2170), \phi(2170) \rightarrow \phi f_0(980), f_0 \rightarrow \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{54} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2 × 10⁻⁶	90	ABLIKIM	19I BES3	$e^+ e^- \rightarrow \eta \phi f_0(980)$

NODE=M071P19
NODE=M071P19

 $\Gamma(\phi \eta') / \Gamma_{\text{total}}$ Γ_{55} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.54 ± 0.20 OUR AVERAGE				
1.51 ± 0.16 ± 0.12	201	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$
3.1 ± 1.4 ± 0.7	8	¹ ABLIKIM	04K BES	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R90
NODE=M071R90

¹ Calculated combining $\eta' \rightarrow \gamma \rho$ and $\eta \pi^+ \pi^-$ channels.

NODE=M071R;LINKAGE=AI

 $\Gamma(\phi f_1(1285)) / \Gamma_{\text{total}}$ Γ_{56} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.0 ± 0.4 ± 1.3	234	¹ ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071P22
NODE=M071P22

¹ ABLIKIM 19BA reports $[\Gamma(\psi(2S) \rightarrow \phi f_1(1285)) / \Gamma_{\text{total}}] \times [B(f_1(1285) \rightarrow \eta \pi^+ \pi^-)] = (1.03 \pm 0.10 \pm 0.09) \times 10^{-5}$ which we divide by our best value $B(f_1(1285) \rightarrow \eta \pi^+ \pi^-) = (35 \pm 15) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P22;LINKAGE=B

 $\Gamma(\phi \eta(1405) \rightarrow \phi \pi^+ \pi^- \eta) / \Gamma_{\text{total}}$ Γ_{57} / Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
8.46 ± 1.37 ± 0.92	195	ABLIKIM	19BA BES3	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071P24
NODE=M071P24

 $\Gamma(\phi f'_2(1525)) / \Gamma_{\text{total}}$ Γ_{58} / Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.44 ± 0.12 ± 0.11	20 ± 6	BAI	04C		$\psi(2S) \rightarrow 2(K^+ K^-)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.45	90	BAI	98J BES		$e^+ e^- \rightarrow 2(K^+ K^-)$

NODE=M071R67
NODE=M071R67

 $\Gamma(K^+ K^-) / \Gamma_{\text{total}}$ Γ_{59} / Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.48 ± 0.23 ± 0.39	1.3k	¹ METREVELI	12		$\psi(2S) \rightarrow K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
6.2 ± 1.5 ± 0.2	66	^{2,3} LEES	15J BABR		$e^+ e^- \rightarrow K^+ K^- \gamma$
8.3 ± 1.5 ± 0.2	66	^{3,4} LEES	15J BABR		$e^+ e^- \rightarrow K^+ K^- \gamma$
6.3 ± 0.6 ± 0.3		⁵ DOBBS	06A CLEO		$e^+ e^-$
10 ± 7		⁵ BRANDELIK	79C DASP		$e^+ e^-$
< 5	90	FELDMAN	77 MRK1		$e^+ e^-$

NODE=M071R23
NODE=M071R23

OCCUR=2

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² $\sin \phi > 0$.

³ Using $\Gamma(\psi(2S) \rightarrow e^+ e^-) = (2.37 \pm 0.04)$ keV.

⁴ $\sin \phi < 0$.

⁵ Interference with non-resonant $K^+ K^-$ production not taken into account.

NODE=M071R23;LINKAGE=ME
NODE=M071R23;LINKAGE=A
NODE=M071R23;LINKAGE=B
NODE=M071R23;LINKAGE=C
NODE=M071R23;LINKAGE=BA

$\Gamma(K^+ K^- \pi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{60}/Γ
7.3±0.5 OUR AVERAGE					
8.1±1.3±0.3	133	LEES	12F BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	
7.1±0.3±0.4	817.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$	
16 ±4		¹ TANENBAUM	78 MRK1	$e^+ e^-$	
11.0±1.9±0.2	85	² AUBERT	07AK BABR	10.6 $e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^- \gamma$	

NODE=M071R24
NODE=M071R24

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Assuming entirely strong decay.

² Superseded by LEES 12F. AUBERT 07AK reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (2.56 \pm 0.42 \pm 0.16) \times 10^{-3}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R24;LINKAGE=K
NODE=M071R24;LINKAGE=BE

 $\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{61}/Γ
4.07±0.16±0.26						
<8.9	90	1	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$	

NODE=M071R38
NODE=M071R38

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{62}/Γ
<0.046				
	¹ BAI	04D BES	$e^+ e^-$	

NODE=M071R88
NODE=M071R88

¹ Forbidden by CP.

NODE=M071R;LINKAGE=BA

 $\Gamma(K_S^0 K_L^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{63}/Γ
5.34±0.33 OUR AVERAGE					
5.28±0.25±0.34	478 ± 23	¹ METREVELI	12	$\psi(2S) \rightarrow K_S^0 K_L^0$	
5.8 ±0.8 ±0.4		DOBBS	06A CLEO	$e^+ e^-$	
5.24±0.47±0.48	156 ± 14	² BAI	04B BES2	$\psi(2S) \rightarrow K_S^0 K_L^0 \rightarrow \pi^+ \pi^- X$	

NODE=M071R87
NODE=M071R87

¹ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.

² Using $B(K_S^0 \rightarrow \pi^+ \pi^-) = 0.6860 \pm 0.0027$.

NODE=M071R87;LINKAGE=ME
NODE=M071R;LINKAGE=KZ

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{69}/Γ
12.6±0.9 OUR AVERAGE					
18.9±5.7±0.3	32	¹ AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \pi^0 \gamma$	
11.7±1.0±1.5	597	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	
12.7±0.5±1.0	711.6	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

NODE=M071S10
NODE=M071S10

¹ AUBERT 07AU reports $[\Gamma(\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+ e^-)] = (44 \pm 13 \pm 3) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+ e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071S10;LINKAGE=UB

 $\Gamma(\omega f_0(1710) \rightarrow \omega K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{70}/Γ
5.9±2.0±0.9					
	19	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

NODE=M071S20
NODE=M071S20

 $\Gamma(K^*(892)^0 K^- \pi^+ \pi^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{71}/Γ
8.6±1.3±1.8					
	238	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

NODE=M071S21
NODE=M071S21

 $\Gamma(K^*(892)^+ K^- \pi^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{72}/Γ
9.6±2.2±1.7					
	133	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

NODE=M071S22
NODE=M071S22

$\Gamma(K^*(892)^+ K^- \rho^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{73}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.3±2.2±1.4	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S23
NODE=M071S23

 $\Gamma(K^*(892)^0 K^- \rho^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{74}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±1.3±1.2	125	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

NODE=M071S24
NODE=M071S24

 $\Gamma(K_S^0 K_S^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{75}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±0.25±0.37	83 ± 9	ABLIKIM	050 BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R49
NODE=M071R49

 $\Gamma(K^+ K^- \rho^0)/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.2±0.2±0.4	223.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M071S09
NODE=M071S09

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0)/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.86±0.32±0.43		93 ± 16	BAI	04C	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^-$
<1.2		90	BAI	98J BES	$e^+ e^-$

NODE=M071R66
NODE=M071R66

 $\Gamma(K^+ K^- \pi^+ \pi^- \eta)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.3±0.7±0.1	7	¹ AUBERT	07AU BABR	10.6 $e^+ e^- \rightarrow K^+ K^- \pi^+ \pi^- \eta \gamma$
		¹ AUBERT 07AU quotes $\Gamma_{ee}^{\psi(2S)} \cdot B(\psi(2S) \rightarrow 2(\pi^+ \pi^-) \eta) \cdot B(\eta \rightarrow \gamma \gamma) = 1.2 \pm 0.7 \pm 0.1 \text{ eV}$.		

NODE=M071S39
NODE=M071S39

NODE=M071S39;LINKAGE=UB

 $\Gamma(K^+ K^- 2(\pi^+ \pi^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.0±2.5±1.8	65	ABLIKIM	07D BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071R09
NODE=M071R09

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ±0.4 OUR AVERAGE		Error includes scale factor of 1.2.			
3.18±0.30 ^{+0.26} _{-0.31}		0.2k	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$
2.9 ^{+1.3} _{-1.7} ±0.4		9.6 ± 4.2	ABLIKIM	05i BES2	$e^+ e^- \rightarrow \psi(2S)$
1.3 ^{+1.0} _{-0.7} ±0.3		7	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$
<5.4		90	FRANKLIN	83 MRK2	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M071R39
NODE=M071R39

 $\Gamma(2(K^+ K^-))/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.63±0.13 OUR AVERAGE				
0.9 ±0.4 ±0.1	13	LEES	12F BABR	10.6 $e^+ e^- \rightarrow 2(K^+ K^-) \gamma$
0.6 ±0.1 ±0.1	59.2	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$

NODE=M071S12
NODE=M071S12

 $\Gamma(2(K^+ K^-) \pi^0)/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.2±0.2	44.7	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-) \pi^0$

NODE=M071S13
NODE=M071S13

 $\Gamma(K^+ K^- \phi)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.16 OUR AVERAGE				
0.8 ±0.2 ±0.1	36.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow 2(K^+ K^-)$
0.6 ±0.2 ±0.1	16.1 ± 5.0	¹ BAI	03B BES	$\psi(2S) \rightarrow 2(K^+ K^-)$
		¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.		

NODE=M071R81
NODE=M071R81

NODE=M071R81;LINKAGE=B3

$\Gamma(K_S^0 K_S^0 \phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{92}/Γ
0.353 ± 0.020 ± 0.021	687	¹ ABLIKIM	23BA BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K_S^0 K_S^0 K^+ K^-$	NODE=M071P53 NODE=M071P53

¹ Solution with a constructive interference of the signal with the continuum background.

NODE=M071P53;LINKAGE=A

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{93}/Γ
10.0 ± 1.8 ± 2.1	¹ BAI	99C BES	$e^+ e^-$	NODE=M071R41 NODE=M071R41

¹ Assuming $B(K_1(1270) \rightarrow K \rho) = 0.42 \pm 0.06$

NODE=M071R;LINKAGE=M2

 $\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT	Γ_{94}/Γ
6.7 ± 2.5	TANENBAUM 78	MRK1	$e^+ e^-$	NODE=M071R34 NODE=M071R34

 $\Gamma(\eta K^+ K^-, \text{ no } \eta\phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{95}/Γ
3.49 ± 0.09 ± 0.15	1.8k		¹ ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$	NODE=M071S11 NODE=M071S11
• • • We do not use the following data for averages, fits, limits, etc. • • •						
3.08 ± 0.29 ± 0.25	0.3k		^{1,2} ABLIKIM	12L BES3	$\psi(2S) \rightarrow K^+ K^- \gamma\gamma$	
<13	90		BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$	

¹ Excluding $\eta\phi$.

² Superseded by ABLIKIM 20F.

NODE=M071S11;LINKAGE=AB
NODE=M071S11;LINKAGE=A

 $\Gamma(\eta K^+ K^-) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{96}\Gamma_7/\Gamma$
<0.6	90	¹ LEES	23 BABR	$e^+ e^- \rightarrow \gamma_{ISR} \text{ hadrons}$	NODE=M071P57 NODE=M071P57

¹ LEES 23 reports $[\Gamma(\psi(2S) \rightarrow \eta K^+ K^-) \times \Gamma(\psi(2S) \rightarrow e^+ e^-)/\Gamma_{\text{total}}] \times [B(\eta \rightarrow 3\pi^0)] < 0.2 \text{ eV}$ which we divide by our best value $B(\eta \rightarrow 3\pi^0) = 32.57 \times 10^{-2}$.

NODE=M071P57;LINKAGE=A

 $\Gamma(X(1750)\eta \rightarrow K^+ K^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT	Γ_{97}/Γ
4.8 ± 1.0 ± 2.6	ABLIKIM	20F BES3	$\psi(2S) \rightarrow K^+ K^- \eta$	NODE=M071P28 NODE=M071P28

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{98}/Γ
<3.1	90	¹ BAI	99C BES	$e^+ e^-$	NODE=M071R45 NODE=M071R45

¹ Assuming $B(K_1(1400) \rightarrow K^* \pi) = 0.94 \pm 0.06$

NODE=M071R;LINKAGE=M3

 $\Gamma(K_2^*(1430)^\pm K^\mp)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
7.12 ± 0.62 ± 0.61	251 ± 22	ABLIKIM	12L BES3	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071S54 NODE=M071S54

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ
10.9 ± 2.0 OUR AVERAGE					NODE=M071R30 NODE=M071R30
13.3 ^{+2.4} _{-2.8} ± 1.7	65.6 ± 9.0	ABLIKIM	05i BES2	$e^+ e^- \rightarrow \psi(2S)$	
9.2 ^{+2.7} _{-2.2} ± 0.9	25	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$	

 $\Gamma(K^+ K^*(892)^- + \text{c.c.})/\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{88}/Γ_{100}
0.16 ± 0.06 OUR AVERAGE				NODE=M071R46 NODE=M071R46
0.22 ^{+0.10} _{-0.14}	ABLIKIM	05i BES2	$e^+ e^- \rightarrow \psi(2S)$	
0.14 ^{+0.08} _{-0.06}	ADAM	05 CLEO	$e^+ e^- \rightarrow \psi(2S)$	

$\Gamma(\omega K^+ K^-)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.62±0.11 OUR AVERAGE				Error includes scale factor of 1.1.
1.56±0.04±0.11	2.8k	ABLIKIM	14G BES3	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
2.38±0.37±0.29	78	ABLIKIM	06G BES2	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.9 ±0.3 ±0.3	76.8	BRIERE	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$
1.5 ±0.3 ±0.2	23	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+ K^- \pi^+ \pi^- \pi^0$

¹ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.NODE=M071R78
NODE=M071R78

NODE=M071R78;LINKAGE=B3

 $\Gamma(\omega K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
7.04±0.39±0.36	1.5k	ABLIKIM	21AL BES3	$\psi(2S) \rightarrow \pi^+ \pi^- \pi^0 K_S^0 K_S^0$

NODE=M071P34
NODE=M071P34 $\Gamma(\omega K^*(892)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
20.7±2.6 OUR AVERAGE				
18.9±2.9±2.2	396	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
22.6±3.0±2.4	535	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S67
NODE=M071S67

OCCUR=2

 $\Gamma(\omega K_2^*(1430)^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1 ±1.2 OUR AVERAGE				
6.39±1.50±0.78	128	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$
5.86±1.61±0.83	143	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

NODE=M071S68
NODE=M071S68

OCCUR=2

 $\Gamma(\omega \bar{K}^*(892)^0 K^0)/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
16.8±2.5±1.6	356	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

NODE=M071S69
NODE=M071S69 $\Gamma(\omega \bar{K}_2^*(1430)^0 K^0)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
5.82±2.08±0.72	116	ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

NODE=M071S70
NODE=M071S70 $\Gamma(\omega X(1440) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.60±0.27±0.24	109	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.NODE=M071S71
NODE=M071S71

NODE=M071S71;LINKAGE=AB

 $\Gamma(\omega X(1440) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.09±0.20±0.16	82	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ X(1440) compatible with $\eta(1405)$ and $\eta(1475)$. A $f_1(1420)$ is also possible.NODE=M071S72
NODE=M071S72

NODE=M071S72;LINKAGE=AB

 $\Gamma(\omega f_1(1285) \rightarrow \omega K_S^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.302±0.098±0.027	22	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K_S^0 K^- \pi^+$

¹ Statistical significance 4.5σ . This measurement is equivalent to a limit of $< 0.478 \times 10^{-5}$ at 90% C.L.NODE=M071S73
NODE=M071S73

NODE=M071S73;LINKAGE=AB

 $\Gamma(\omega f_1(1285) \rightarrow \omega K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{110}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.125±0.070±0.013	10	¹ ABLIKIM	13M BES3	$\psi(2S) \rightarrow \omega K^+ K^- \pi^0$

¹ Statistical significance 3.2σ . This measurement is equivalent to a limit of $< 0.221 \times 10^{-5}$ at 90% C.L.NODE=M071S74
NODE=M071S74

NODE=M071S74;LINKAGE=AB

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.94 ± 0.09 OUR FIT Error includes scale factor of 1.3. $[(2.94 \pm 0.08) \times 10^{-4} \text{ OUR } 2023 \text{ FIT}]$

3.02 ± 0.08 OUR AVERAGE

3.05 ± 0.02 ± 0.12	19k	ABLIKIM	18T BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.08 ± 0.05 ± 0.18	4.5k	¹ DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
3.36 ± 0.09 ± 0.25	1.6k	ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.87 ± 0.12 ± 0.15	557	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
1.4 ± 0.8	4	BRANDELIK	79C DASP	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
2.3 ± 0.7		FELDMAN	77 MRK1	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

 Γ_{111}/Γ

NODE=M071R25

NODE=M071R25

NEW

NODE=M071R25;LINKAGE=A

 $\Gamma(p\bar{p})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

8.49 ± 0.28 OUR FIT Error includes scale factor of 1.3. $[(8.49 \pm 0.23) \times 10^{-4} \text{ OUR } 2023 \text{ FIT}]$

6.98 ± 0.49 ± 0.97

BAI	01	BES	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}$
-----	----	-----	--

 Γ_{111}/Γ_{12}

NODE=M071S40

NODE=M071S40

NEW

 $\Gamma(n\bar{n})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.06 ± 0.06 ± 0.14 6k ABLIKIM 18T BES3 $e^+e^- \rightarrow \psi(2S) \rightarrow n\bar{n}$

 Γ_{112}/Γ

NODE=M071P10

NODE=M071P10

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.53 ± 0.07 OUR AVERAGE

1.65 ± 0.03 ± 0.15	4.5k	ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
1.54 ± 0.06 ± 0.06	948	ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$
1.32 ± 0.10 ± 0.15	256	¹ ABLIKIM	05E BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\gamma\gamma$
1.4 ± 0.5	9	FRANKLIN	83 MRK2	e^+e^-

¹ Computed using $B(\pi^0 \rightarrow \gamma\gamma) = (98.80 \pm 0.03)\%$.

 Γ_{113}/Γ

NODE=M071R35

NODE=M071R35

NODE=M071R35;LINKAGE=AB

 $\Gamma(N(940)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

6.42 ± 0.20 $\begin{smallmatrix} +1.78 \\ -1.28 \end{smallmatrix}$ 1.9k ¹ ABLIKIM 13A BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{114}/Γ

NODE=M071S56

NODE=M071S56

NODE=M071S56;LINKAGE=AB

 $\Gamma(N(1440)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

7.3 $\begin{smallmatrix} +1.7 \\ -1.5 \end{smallmatrix}$ OUR AVERAGE Error includes scale factor of 2.5.

3.58 ± 0.25 $\begin{smallmatrix} +1.59 \\ -0.84 \end{smallmatrix}$	1.1k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow p\bar{p}\pi^0$
8.1 ± 0.7 ± 0.3	474	² ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \pi^0 p\bar{p}$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

² From a fit of the $p\bar{p}$ and $p\pi^0$ mass distributions to a combination of $N(1440)\bar{p}$, a broad $p\bar{p}$ enhancement around 2100 MeV, and two other broad, unestablished resonances.

 Γ_{115}/Γ

NODE=M071S50

NODE=M071S50

NODE=M071S50;LINKAGE=AB

NODE=M071S50;LINKAGE=AL

 $\Gamma(N(1520)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

0.64 ± 0.05 $\begin{smallmatrix} +0.22 \\ -0.17 \end{smallmatrix}$ 0.2k ¹ ABLIKIM 13A BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{116}/Γ

NODE=M071S57

NODE=M071S57

NODE=M071S57;LINKAGE=AB

 $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.47 ± 0.28 $\begin{smallmatrix} +0.99 \\ -0.97 \end{smallmatrix}$ 0.7k ¹ ABLIKIM 13A BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{117}/Γ

NODE=M071S58

NODE=M071S58

NODE=M071S58;LINKAGE=AB

 $\Gamma(N(1650)\bar{p} + \text{c.c.} \rightarrow p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

3.76 ± 0.28 $\begin{smallmatrix} +1.37 \\ -1.66 \end{smallmatrix}$ 1.1k ¹ ABLIKIM 13A BES3 $\psi(2S) \rightarrow p\bar{p}\pi^0$

¹ From a fit of $\pi^0 p\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

 Γ_{118}/Γ

NODE=M071S59

NODE=M071S59

NODE=M071S59;LINKAGE=AB

$\Gamma(N(1720)\bar{p} + \text{c.c.} \rightarrow \rho\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.79 \pm 0.10^{+0.24}_{-0.71}$	0.5k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow \rho\bar{p}\pi^0$

NODE=M071S60
NODE=M071S60¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

NODE=M071S60;LINKAGE=AB

 $\Gamma(N(2300)\bar{p} + \text{c.c.} \rightarrow \rho\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.62 \pm 0.28^{+1.12}_{-0.64}$	0.9k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow \rho\bar{p}\pi^0$

NODE=M071S61
NODE=M071S61¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

NODE=M071S61;LINKAGE=AB

 $\Gamma(N(2570)\bar{p} + \text{c.c.} \rightarrow \rho\bar{p}\pi^0)/\Gamma_{\text{total}}$ Γ_{121}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.13 \pm 0.08^{+0.40}_{-0.30}$	0.8k	¹ ABLIKIM	13A BES3	$\psi(2S) \rightarrow \rho\bar{p}\pi^0$

NODE=M071S62
NODE=M071S62¹ From a fit of $\pi^0 \rho\bar{p}$ data to eight distinct intermediate $N\bar{p}$ resonant states.

NODE=M071S62;LINKAGE=AB

 $\Gamma(\rho\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{122}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.4 OUR AVERAGE				
$5.9 \pm 0.2 \pm 0.4$	904.5	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-$
8 ± 2		¹ TANENBAUM	78 MRK1	e^+e^-

NODE=M071R31
NODE=M071R31¹ Assuming entirely strong decay.

NODE=M071R;LINKAGE=K

 $\Gamma(\rho\bar{p}K^+K^-)/\Gamma_{\text{total}}$ Γ_{123}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.7 \pm 0.6 \pm 0.4$	30.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}K^+K^-$

NODE=M071S16
NODE=M071S16 $\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{124}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
6.0 ± 0.4 OUR AVERAGE				
$6.4 \pm 0.2 \pm 0.6$	679	¹ ABLIKIM	13S BES3	$\psi(2S) \rightarrow \eta\rho\bar{p}$
$5.6 \pm 0.6 \pm 0.3$	154	¹ ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta\rho\bar{p}$
$5.8 \pm 1.1 \pm 0.7$	44.8 ± 8.5	² ABLIKIM	05E BES2	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\gamma\gamma$
$8 \pm 3 \pm 3$	9.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$

NODE=M071R56
NODE=M071R56¹ With $N(1535)$ decaying to $\rho\eta$.² Computed using $B(\eta \rightarrow \gamma\gamma) = (39.43 \pm 0.26)\%$.NODE=M071R56;LINKAGE=A
NODE=M071R56;LINKAGE=AB $\Gamma(N(1535)\bar{p} + \text{c.c.} \rightarrow \rho\bar{p}\eta)/\Gamma_{\text{total}}$ Γ_{125}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
4.5^{+0.7}_{-0.6} OUR AVERAGE				
$5.2 \pm 0.3^{+3.2}_{-1.2}$	527	¹ ABLIKIM	13S BES3	$\psi(2S) \rightarrow \eta\rho\bar{p}$
$4.4 \pm 0.6 \pm 0.3$	123	² ALEXANDER	10 CLEO	$\psi(2S) \rightarrow \eta\rho\bar{p}$

NODE=M071S53
NODE=M071S53¹ With $N(1535)$ decaying to $\rho\eta$.² From a fit of the $\rho\bar{p}$ and $\rho\eta$ distributions to a combination of $N^*(1535)\bar{p}$ and a broad $\rho\bar{p}$ enhancement around 2100 MeV.NODE=M071S53;LINKAGE=A
NODE=M071S53;LINKAGE=AL $\Gamma(\rho\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{126}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.3 \pm 0.4 \pm 0.6$	434.9	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-\pi^0$

NODE=M071S15
NODE=M071S15 $\Gamma(\rho\bar{p}\rho^0)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.5 \pm 0.1 \pm 0.2$	61.1	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}\pi^+\pi^-$

NODE=M071S14
NODE=M071S14

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.69±0.21 OUR AVERAGE				
0.6 ±0.2 ±0.2	21.2	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$
0.8 ±0.3 ±0.1	14.9 ± 0.1	¹ BAI	03B BES	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

¹Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R79
NODE=M071R79

NODE=M071R;LINKAGE=B3

 $\Gamma(p\bar{p}\eta')/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10±0.10±0.08	491	¹ ABLIKIM	19N BES3	$\psi(2S) \rightarrow \eta' p\bar{p}$

¹From the combination of $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\eta$ and $p\bar{p}\eta' \rightarrow p\bar{p}\pi^+\pi^-\gamma$ channels.

NODE=M071P20
NODE=M071P20

NODE=M071P20;LINKAGE=A

 $\Gamma(p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.06±0.38±0.48		753	ABLIKIM	19A0 BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M071R82
NODE=M071R82

<24	90	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$
<26	90	¹ BAI	03B BES	$\psi(2S) \rightarrow K^+K^-p\bar{p}$

¹Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

NODE=M071R82;LINKAGE=B3

 $\Gamma(\phi X(1835) \rightarrow p\bar{p}\phi)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.82 × 10⁻⁷	90	ABLIKIM	19A0 BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}K^+K^-$

NODE=M071P21
NODE=M071P21 $\Gamma(p\bar{n}\pi^- \text{ or c.c.})/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48±0.17 OUR AVERAGE				
2.45±0.11±0.21	851	ABLIKIM	06I BES2	$e^+e^- \rightarrow p\pi^-X$
2.52±0.12±0.22	849	ABLIKIM	06I BES2	$e^+e^- \rightarrow \bar{p}\pi^+X$

NODE=M071R01
NODE=M071R01

OCCUR=2


 $\Gamma(p\bar{n}\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.18±0.50±0.50	135 ± 21	ABLIKIM	06I BES2	$e^+e^- \rightarrow p\pi^-\pi^0X$

NODE=M071R02
NODE=M071R02 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{134}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.81±0.13 OUR AVERAGE					Error includes scale factor of 1.4. See the ideogram below.
3.97±0.02±0.12	31k	ABLIKIM	17L BES3		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
3.71±0.05±0.15	6.5k	¹ DOBBS	17		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
3.39±0.20±0.32	337	ABLIKIM	07C BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
6.4 ±1.8 ±0.1		² AUBERT	07BD BABR		10.6 $e^+e^- \rightarrow \Lambda\bar{\Lambda}\gamma$
3.28±0.23±0.25	208	PEDLAR	05 CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.75±0.09±0.23	1.9k	^{1,3} DOBBS	14		$e^+e^- \rightarrow \Lambda\bar{\Lambda}$
1.81±0.20±0.27	80	⁴ BAI	01 BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 4	90	FELDMAN	77 MRK1		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

NODE=M071R28
NODE=M071R28NODE=M071R28;LINKAGE=A
NODE=M071R28;LINKAGE=A¹Using CLEO-c data but not authored by the CLEO Collaboration.²AUBERT 07BD reports $[\Gamma(\psi(2S) \rightarrow \Lambda\bar{\Lambda})/\Gamma_{\text{total}}] \times [\Gamma(\psi(2S) \rightarrow e^+e^-)] = (15 \pm 4 \pm 1) \times 10^{-4}$ keV which we divide by our best value $\Gamma(\psi(2S) \rightarrow e^+e^-) = 2.33 \pm 0.04$ keV. Our first error is their experiment's error and our second error is the systematic error from using our best value.³Superseded by DOBBS 17.⁴Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.NODE=M071R28;LINKAGE=B
NODE=M071R28;LINKAGE=PP

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{135}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.42±0.39±0.59		23	¹ ABLIKIM 22AP	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
< 2.9	90		² ABLIKIM 13F	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
< 120	90		³ ABLIKIM 07H	BES2	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ With a significance of 3.7 σ . The corresponding 90% CL upper limit is 2.47×10^{-6} .

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\pi^0 \rightarrow \gamma\gamma) = 98.8\%$.

³ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.4\%$.

NODE=M071R6
NODE=M071R6

NODE=M071R6;LINKAGE=A
NODE=M071R6;LINKAGE=AL
NODE=M071R6;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{136}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.43±0.32 OUR AVERAGE					
2.34±0.18±0.52		218	ABLIKIM 22AP	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
2.48±0.34±0.19		60	¹ ABLIKIM 13F	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$
< 4.9	90		² ABLIKIM 07H	BES2	$e^+e^- \rightarrow \psi(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$ and $B(\eta \rightarrow \gamma\gamma) = 39.31\%$.

² Using $B(\Lambda \rightarrow \pi^- p) = 63.9\%$.

NODE=M071R7
NODE=M071R7

NODE=M071R7;LINKAGE=AL
NODE=M071R7;LINKAGE=AB

 $\Gamma(\Lambda\bar{\Lambda}\eta')/\Gamma_{\text{total}}$ Γ_{137}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
7.34±0.94±0.43	218	ABLIKIM 23BV	BES3	$\psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-\gamma(\eta))$

NODE=M071P52
NODE=M071P52

 $\Gamma(\Lambda\bar{\Lambda}\omega(782))/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.30±0.34±0.29	207	¹ ABLIKIM 22AZ	BES3	$e^+e^- \rightarrow \psi(2S)$

¹ Using $B(\Lambda \rightarrow \pi^- p) = 0.639$ and $B(\omega \rightarrow \pi^+\pi^-\pi^0) = 0.893$.

NODE=M071P46
NODE=M071P46

OCCUR=2

NODE=M071P46;LINKAGE=B

 $\Gamma(\Lambda(1670)\bar{\Lambda} \rightarrow \Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.29±0.31±0.62	116	¹ ABLIKIM 22AP	BES3	$\psi(2S) \rightarrow p\bar{p}\pi^+\pi^-\gamma\gamma$

¹ From a partial wave analysis of the $\Lambda\eta$ system.

NODE=M071P42
NODE=M071P42

NODE=M071P42;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{140}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.8±0.4±0.5	73.4	BRIERE 05	CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow p\bar{p}2(\pi^+\pi^-)$

NODE=M071S17
NODE=M071S17

$\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$					Γ_{141}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.0±0.1±0.1	74.0	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}K^+\pi^-$		NODE=M071S18 NODE=M071S18

$\Gamma(\Lambda\bar{p}K^*(892)^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{142}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
6.3±0.5±0.5	1011	ABLIKIM	19AU BES3	$e^+e^- \rightarrow \psi(2S)$		NODE=M071P25 NODE=M071P25

$\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{143}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.8±0.3±0.3	45.8	BRIERE	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \rho\bar{p}K^+\pi^+\pi^-\pi^-$		NODE=M071S19 NODE=M071S19

$\Gamma(\bar{\Lambda}nK_S^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{144}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.81±0.11±0.14	50	¹ ABLIKIM	08C BES2	$e^+e^- \rightarrow J/\psi$		NODE=M071R08 NODE=M071R08
¹ Using $B(\bar{\Lambda} \rightarrow \bar{p}\pi^+) = 63.9\%$ and $B(K_S^0 \rightarrow \pi^+\pi^-) = 69.2\%$.						
						NODE=M071R08;LINKAGE=AB

$\Gamma(\Delta^{++}\bar{\Delta}^{--})/\Gamma_{\text{total}}$					Γ_{145}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
12.8±1.0±3.4	157	¹ BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$		NODE=M071R50 NODE=M071R50
¹ Estimated using $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.310 \pm 0.028$.						
						NODE=M071R50;LINKAGE=PP

$\Gamma(\Lambda\bar{\Sigma}^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{146}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.40±0.03±0.13	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$		NODE=M071S65 NODE=M071S65

$\Gamma(\Lambda\bar{\Sigma}^-\pi^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{147}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.54±0.04±0.13	2.8k	ABLIKIM	13W BES3	$\psi(2S) \rightarrow \text{hadrons}$		NODE=M071S66 NODE=M071S66

$\Gamma(\Lambda\bar{\Sigma}^0 + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{148}/Γ	
VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.60±0.31±0.59	60	ABLIKIM	21L BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$		NODE=M071P29 NODE=M071P29

$\Gamma(\Lambda\bar{\Sigma}^0)/\Gamma_{\text{total}}$					Γ_{149}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.23±0.23±0.08	30	¹ DOBBS	17	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$		NODE=M071P08 NODE=M071P08

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071P08;LINKAGE=A

$\Gamma(\Sigma^0\bar{p}K^+ + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{150}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.67±0.13±0.12	276	¹ ABLIKIM	13D BES3	$\psi(2S) \rightarrow \gamma\Lambda\bar{p}K^+$		NODE=M071S63 NODE=M071S63

¹ Using $B(\Lambda \rightarrow p\pi^-) = 63.9\%$, and $B(\Sigma^0 \rightarrow \Lambda\gamma) = 100\%$.

NODE=M071S63;LINKAGE=AB

$\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$					Γ_{151}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
2.43±0.10 OUR AVERAGE	Error includes scale factor of 1.4.					
2.52±0.04±0.09	5.4k	ABLIKIM	21AT BES3	$\psi(2S) \rightarrow p\pi^0\bar{p}\pi^0$		
2.31±0.06±0.10	1.9k	¹ DOBBS	17	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$		
2.57±0.44±0.68	35	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$		

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.51±0.15±0.16 281 ^{1,2} DOBBS 14 $e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² Superseded by DOBBS 17.

NODE=M071R47;LINKAGE=A

NODE=M071R47;LINKAGE=B

$\Gamma(\Sigma^0 \bar{\Sigma}^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.35±0.09 OUR AVERAGE		Error includes scale factor of 1.1.		
2.44±0.03±0.11	7k	ABLIKIM	17L BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.22±0.05±0.11	2.6k	¹ DOBBS	17	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.35±0.36±0.32	59	ABLIKIM	07C BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.63±0.35±0.21	58	PEDLAR	05 CLEO	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.25±0.11±0.16	439	^{1,2} DOBBS	14	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
1.2 ±0.4 ±0.4	8	³ BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17.³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. Γ_{152}/Γ NODE=M071R51
NODE=M071R51 $\Gamma(\Sigma^- \bar{\Sigma}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.82±0.04±0.08	6.6k	ABLIKIM	22AV BES3	$\psi(2S) \rightarrow n\pi^- \bar{n}\pi^+$

 Γ_{153}/Γ NODE=M071P41
NODE=M071P41 $\Gamma(\Sigma^+ \bar{\Sigma}^- \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
9.59±2.37±0.61	21	ABLIKIM	22AY BES3	$\psi(2S) \rightarrow \Sigma^+ \bar{\Sigma}^- \eta$

 Γ_{154}/Γ NODE=M071P45
NODE=M071P45 $\Gamma(\Sigma^+ \bar{\Sigma}^- \omega)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.89±0.18±0.21	199	ABLIKIM	23BE BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{155}/Γ NODE=M071P50
NODE=M071P50 $\Gamma(\Sigma^+ \bar{\Sigma}^- \phi)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
2.96±0.54±0.41	55	ABLIKIM	23BE BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{156}/Γ NODE=M071P51
NODE=M071P51 $\Gamma(\Sigma(1385)^+ \bar{\Sigma}(1385)^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±0.7 OUR AVERAGE		Error includes scale factor of 1.1.		
8.4±0.5±0.5	1.5k	ABLIKIM	16L BES3	$\psi(2S) \rightarrow \Sigma(1385)^+ \bar{\Sigma}(1385)^-$
11 ±3 ±3	14	¹ BAI	01 BES	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{157}/Γ NODE=M071R52
NODE=M071R52¹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

OCCUR=2

NODE=M071R52;LINKAGE=PP

 $\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±0.6±0.6	1.4k	ABLIKIM	16L BES3	$\psi(2S) \rightarrow \Sigma(1385)^- \bar{\Sigma}(1385)^+$

 Γ_{158}/Γ NODE=M071R00
NODE=M071R00 $\Gamma(\Sigma(1385)^0 \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.69±0.05±0.05	2.2k	ABLIKIM	17E BES3	$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{159}/Γ NODE=M071P00
NODE=M071P00 $\Gamma(\Xi^- \bar{\Xi}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.87±0.11 OUR AVERAGE			Error includes scale factor of 1.1.		
3.03±0.05±0.14	3.6k	¹ DOBBS	17		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.78±0.05±0.14	5k	ABLIKIM	16L BES3		$\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$
3.03±0.40±0.32	67	ABLIKIM	07C BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.38±0.30±0.21	63	PEDLAR	05 CLEO		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.66±0.12±0.20	548	^{1,2} DOBBS	14		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
0.94±0.27±0.15	12	³ BAI	01 BES		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
<2	90	FELDMAN	77 MRK1		$e^+e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{160}/Γ NODE=M071R29
NODE=M071R29¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17.³ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.NODE=M071R29;LINKAGE=A
NODE=M071R29;LINKAGE=B
NODE=M071R29;LINKAGE=PP

$\Gamma(\Xi^0 \Xi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.3 ± 0.4 OUR AVERAGE		Error includes scale factor of 4.2.		
2.73 ± 0.03 ± 0.13	11k	ABLIKIM	17E BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
1.97 ± 0.06 ± 0.11	1.2k	¹ DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
2.75 ± 0.64 ± 0.61	19	PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.02 ± 0.19 ± 0.15	112	^{1,2} DOBBS	14	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.² Superseded by DOBBS 17. Γ_{161}/Γ NODE=M071R48
NODE=M071R48 $\Gamma(\Xi(1530)^0 \Xi(1530)^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.77 ± 0.14 ± 0.39		2951	ABLIKIM	21A0 BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 32	90		PEDLAR	05 CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$
< 8.1	90		¹ BAI	01 BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

¹ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$. Γ_{162}/Γ NODE=M071R53
NODE=M071R53 $\Gamma(\Lambda \Xi^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.86 ± 0.27 ± 0.32	236	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.}$

 Γ_{163}/Γ NODE=M071S82
NODE=M071S82 $\Gamma(\Xi(1530)^- \Xi(1530)^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
11.45 ± 0.40 ± 0.59	5k	ABLIKIM	19AT BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{164}/Γ NODE=M071P26
NODE=M071P26 $\Gamma(\Xi(1530)^- \Xi^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
7.0 ± 1.1 ± 0.4	399	ABLIKIM	19AT BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{165}/Γ NODE=M071P27
NODE=M071P27 $\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
0.53 ± 0.04 ± 0.03	278	ABLIKIM	21A0 BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$

 Γ_{166}/Γ NODE=M071P35
NODE=M071P35 $\Gamma(\Xi(1690)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
5.21 ± 1.48 ± 0.57	74	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.}$

 Γ_{167}/Γ NODE=M071S83
NODE=M071S83 $\Gamma(\Xi(1820)^- \Xi^+ \rightarrow K^- \Lambda \Xi^+ + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
12.03 ± 2.94 ± 1.22	136	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Lambda \Xi^+ + \text{c.c.}$

 Γ_{168}/Γ NODE=M071S84
NODE=M071S84 $\Gamma(\Sigma^0 \Xi^+ K^- + \text{c.c.})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.67 ± 0.33 ± 0.28	142	ABLIKIM	15I BES3	$e^+ e^- \rightarrow \psi(2S) \rightarrow K^- \Sigma^0 \Xi^+ + \text{c.c.}$

 Γ_{169}/Γ NODE=M071S85
NODE=M071S85 $\Gamma(\Omega^- \bar{\Omega}^+)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.66 ± 0.30 OUR AVERAGE			Error includes scale factor of 1.3.		
5.85 ± 0.12 ± 0.25	4k	¹ ABLIKIM	21E BES3	$\psi(2S) \rightarrow \Omega^- \bar{\Omega}^+ \rightarrow \Lambda K^- \bar{\Lambda} K^+$	
5.2 ± 0.3 ± 0.3	326	^{1,2} DOBBS	17	$e^+ e^- \rightarrow \psi(2S) \rightarrow \text{hadrons}$	

 Γ_{170}/Γ NODE=M071R54
NODE=M071R54

• • • We do not use the following data for averages, fits, limits, etc. • • •

$4.7 \pm 0.9 \pm 0.5$	27	^{1,2,3} DOBBS	14		$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
<15	90	ABLIKIM	12Q	BES2	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
<16	90	PEDLAR	05	CLEO	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons
< 7.3	90	⁴ BAI	01	BES	$e^+ e^- \rightarrow \psi(2S) \rightarrow$ hadrons

¹ Using $B(\Omega^- \rightarrow \Lambda K^-) = (67.8 \pm 0.7)\%$ and $B(\Lambda \rightarrow p \pi^-) = (63.9 \pm 0.5)\%$.

² Using CLEO-c data but not authored by the CLEO Collaboration.

³ Superseded by DOBBS 17.

⁴ Estimated using $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.310 \pm 0.028$.

NODE=M071R54;LINKAGE=D
NODE=M071R54;LINKAGE=A
NODE=M071R54;LINKAGE=B
NODE=M071R54;LINKAGE=PP

$\Gamma(\eta_c \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

Γ_{171}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	PEDLAR	07	CLEO $e^+ e^- \rightarrow \psi(2S)$

NODE=M071R03
NODE=M071R03

$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$

Γ_{172}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.4 ± 0.5 OUR AVERAGE				
$7.32 \pm 0.34 \pm 0.41$	46k	ABLIKIM	22AQ	BES3 $\psi(2S) \rightarrow \pi^0$ hadrons
$9.0 \pm 1.5 \pm 1.3$	3k	¹ GE	11	CLEO $\psi(2S) \rightarrow \pi^0$ anything

NODE=M071S42
NODE=M071S42

• • • We do not use the following data for averages, fits, limits, etc. • • •

$8.4 \pm 1.3 \pm 1.0$	11k	² ABLIKIM	10B	BES3 $\psi(2S) \rightarrow \pi^0 h_c$
seen	92^{+23}_{-22}	ADAMS	09	CLEO $\psi(2S) \rightarrow 2\pi^+ 2\pi^- 2\pi^0$
seen	1282	DOBBS	08A	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$
seen	168 ± 40	ROSNER	05	CLEO $\psi(2S) \rightarrow \pi^0 \eta_c \gamma$

¹ Assuming a width $\Gamma(h_c(1P)) = 0.86 \text{ MeV} \equiv \Gamma_0$, a measured dependence of the central value of $B = (7.6 + 1.4 \times \Gamma(h_c(1P))/\Gamma_0) \times 10^{-4}$, and with a systematic error that accounts for the width variation range 0.43–1.29 MeV.

² Superseded by ABLIKIM 22AQ

NODE=M071S42;LINKAGE=GE

NODE=M071S42;LINKAGE=A

$\Gamma(\Lambda_c^+ \bar{p} e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{173}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.7 × 10 ⁻⁶	90	450M	ABLIKIM	18Q	BES3 $e^+ e^- \rightarrow \psi(2S)$

NODE=M071P11
NODE=M071P11

$\Gamma(\Theta(1540)\bar{\Theta}(1540) \rightarrow K_S^0 p K^- \bar{n} + \text{c.c.})/\Gamma_{\text{total}}$

Γ_{174}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.88	90	BAI	04G	BES2 $e^+ e^-$

NODE=M071S01
NODE=M071S01

$\Gamma(\Theta(1540)K^- \bar{n} \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

Γ_{175}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.0	90	BAI	04G	BES2 $e^+ e^-$

NODE=M071S02
NODE=M071S02

$\Gamma(\Theta(1540)K_S^0 \bar{p} \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

Γ_{176}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.70	90	BAI	04G	BES2 $e^+ e^-$

NODE=M071S03
NODE=M071S03

$\Gamma(\bar{\Theta}(1540)K^+ n \rightarrow K_S^0 \bar{p} K^+ n)/\Gamma_{\text{total}}$

Γ_{177}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	BAI	04G	BES2 $e^+ e^-$

NODE=M071S04
NODE=M071S04

$\Gamma(\bar{\Theta}(1540)K_S^0 p \rightarrow K_S^0 p K^- \bar{n})/\Gamma_{\text{total}}$

Γ_{178}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<0.60	90	BAI	04G	BES2 $e^+ e^-$

NODE=M071S05
NODE=M071S05

RADIATIVE DECAYS

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma_{total}$

Γ_{179}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.77 ±0.23 OUR FIT				Error includes scale factor of 1.1. [(9.79 ± 0.20) × 10 ⁻² OUR 2023 FIT]
9.33 ±0.26 OUR AVERAGE				
9.389 ±0.014 ±0.332	4.7M	ABLIKIM	17U	BES3 e ⁺ e ⁻ → γX
9.22 ±0.11 ±0.46	72k	ATHAR	04	CLEO e ⁺ e ⁻ → γX
9.9 ±0.5 ±0.8		¹ GAISER	86	CBAL e ⁺ e ⁻ → γX
7.2 ±2.3		¹ BIDDICK	77	CNTR e ⁺ e ⁻ → γX
7.5 ±2.6		¹ WHITAKER	76	MRK1 e ⁺ e ⁻

¹ Angular distribution (1+cos²θ) assumed.

NODE=M071315

NODE=M071R55
NODE=M071R55

NEW

NODE=M071R;LINKAGE=A

$\Gamma(\gamma\chi_{c1}(1P))/\Gamma_{total}$

Γ_{180}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.75 ±0.27 OUR FIT				Error includes scale factor of 1.1. [(9.75 ± 0.24) × 10 ⁻² OUR 2023 FIT]
9.54 ±0.29 OUR AVERAGE				
9.905 ±0.011 ±0.353	5.0M	ABLIKIM	17U	BES3 e ⁺ e ⁻ → γX
9.07 ±0.11 ±0.54	76k	ATHAR	04	CLEO e ⁺ e ⁻ → γX
9.0 ±0.5 ±0.7		¹ GAISER	86	CBAL e ⁺ e ⁻ → γX
7.1 ±1.9		² BIDDICK	77	CNTR e ⁺ e ⁻ → γX

¹ Angular distribution (1-0.189 cos²θ) assumed.

² Valid for isotropic distribution of the photon.

NODE=M071R58
NODE=M071R58

NEW

NODE=M071R;LINKAGE=G
NODE=M071R;LINKAGE=B

$\Gamma(\gamma\chi_{c0}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

$\Gamma_{179}/\Gamma_{180}$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.02 ±0.01 ±0.07	¹ ATHAR	04	CLEO e ⁺ e ⁻ → γX

¹ Not independent from ATHAR 04 measurements of B(γχ_{cJ}).

NODE=M071R97
NODE=M071R97

NODE=M071R97;LINKAGE=AH

$\Gamma(\gamma\chi_{c2}(1P))/\Gamma_{total}$

Γ_{181}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
9.36 ±0.23 OUR FIT				Error includes scale factor of 1.2. [(9.52 ± 0.20) × 10 ⁻² OUR 2023 FIT]
9.42 ±0.31 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
9.621 ±0.013 ±0.272	4.2M	ABLIKIM	17U	BES3 e ⁺ e ⁻ → γX
9.33 ±0.14 ±0.61	79k	ATHAR	04	CLEO e ⁺ e ⁻ → γX
8.0 ±0.5 ±0.7		¹ GAISER	86	CBAL e ⁺ e ⁻ → γX
7.0 ±2.0		² BIDDICK	77	CNTR e ⁺ e ⁻ → γX

¹ Angular distribution (1-0.052 cos²θ) assumed.


² Valid for isotropic distribution of the photon.

NODE=M071R59
NODE=M071R59

NEW

NODE=M071R;LINKAGE=F
NODE=M071R59;LINKAGE=B

WEIGHTED AVERAGE
9.42±0.31 (Error scaled by 1.3)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$$\frac{\Gamma(\gamma\chi_{c0}(1P)) + \Gamma(\gamma\chi_{c1}(1P)) + \Gamma(\gamma\chi_{c2}(1P))}{\Gamma_{\text{total}}} \quad \frac{\Gamma_{179} + \Gamma_{180} + \Gamma_{181}}{\Gamma}$$

VALUE	DOCUMENT ID	TECN	COMMENT
27.6 ± 0.3 ± 2.0	¹ ATHAR 04	CLEO	e ⁺ e ⁻ → γX

¹ Not independent from ATHAR 04 measurements of B(γχ_{cJ}).

NODE=M071R19
NODE=M071R19

NODE=M071R;LINKAGE=AH

$$\frac{\Gamma(\gamma\chi_{c0}(1P))}{\Gamma(\gamma\chi_{c2}(1P))} \quad \frac{\Gamma_{179}}{\Gamma_{181}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
0.99 ± 0.02 ± 0.08	¹ ATHAR 04	CLEO	e ⁺ e ⁻ → γX

¹ Not independent from ATHAR 04 measurements of B(γχ_{cJ}).

NODE=M071R99
NODE=M071R99

NODE=M071R99;LINKAGE=AH

$$\frac{\Gamma(\gamma\chi_{c2}(1P))}{\Gamma(\gamma\chi_{c1}(1P))} \quad \frac{\Gamma_{181}}{\Gamma_{180}}$$

VALUE	DOCUMENT ID	TECN	COMMENT
1.03 ± 0.02 ± 0.03	¹ ATHAR 04	CLEO	e ⁺ e ⁻ → γX

¹ Not independent from ATHAR 04 measurements of B(γχ_{cJ}).

NODE=M071R98
NODE=M071R98

NODE=M071R98;LINKAGE=AH

$$\frac{\Gamma(\gamma\eta_c(1S))}{\Gamma_{\text{total}}} \quad \frac{\Gamma_{182}}{\Gamma}$$

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
0.36 ± 0.05 OUR FIT				Error includes scale factor of 1.3.
0.34 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.432 ± 0.016 ± 0.060		MITCHELL 09	CLEO	e ⁺ e ⁻ → γX
0.32 ± 0.04 ± 0.06	2.5k	¹ ATHAR 04	CLEO	e ⁺ e ⁻ → γX
0.28 ± 0.06		² GAISER 86	CBAL	e ⁺ e ⁻ → γX


¹ ATHAR 04 used Γ_{η_c(1S)} = 24.8 ± 4.9 MeV to obtain this result.

² GAISER 86 used Γ_{η_c(1S)} = 11.5 ± 4.5 MeV to obtain this result.

NODE=M071R60
NODE=M071R60

NODE=M071R60;LINKAGE=AT

NODE=M071R60;LINKAGE=GA



$$\frac{\Gamma(\gamma\eta_c(2S))}{\Gamma_{\text{total}}} \quad \frac{\Gamma_{183}}{\Gamma}$$

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
7 ± 2 ± 4		¹ ABLIKIM 12G	BES3	ψ(2S) → γK ⁰ Kπ, KKπ ⁰
< 8	90	² CRONIN-HEN..10	CLEO	ψ(2S) → γK ⁰ Kπ
< 20	90	ATHAR 04	CLEO	e ⁺ e ⁻ → γX
20-130	95	EDWARDS 82C	CBAL	e ⁺ e ⁻ → γX

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M071R62
NODE=M071R62

¹ ABLIKIM 12G reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] = (1.30 \pm 0.20 \pm 0.30) \times 10^{-5}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = (1.9 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R62;LINKAGE=AB

² CRONIN-HENNESSY 10 reports $[\Gamma(\psi(2S) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K\bar{K}\pi)] < 14.5 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$. This measurement assumes $\Gamma(\eta_c(2S)) = 14$ MeV. CRONIN-HENNESSY 10 gives the analytic dependence of limits on width.

NODE=M071R62;LINKAGE=CR

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{184}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.04 ± 0.22 OUR AVERAGE			Error includes scale factor of 1.4.		
0.95 ± 0.16 ± 0.05		423	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^0$
1.58 ± 0.40 ± 0.13		37	ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma\pi^0$

NODE=M071R42
NODE=M071R42

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5	90		PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
< 5400	95		¹ LIBERMAN	75 SPEC	e^+e^-
< 1×10^4	90		WIJK	75 DASP	e^+e^-

¹ Restated by us using $B(\psi(2S) \rightarrow \mu^+\mu^-) = 0.0077$.

NODE=M071R;LINKAGE=U

 $\Gamma(\gamma 2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{185}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
39.6 ± 2.8 ± 5.0	583	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S28
NODE=M071S28 $\Gamma(\gamma 3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{186}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 17	90	ABLIKIM	07D BES2	$e^+e^- \rightarrow \psi(2S)$

NODE=M071S36
NODE=M071S36 $\Gamma(\gamma\eta'(958))/\Gamma_{\text{total}}$ Γ_{187}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.04 OUR AVERAGE					
1.251 ± 0.022 ± 0.062		56k	ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma\pi^+\pi^-\eta,$ $\gamma\pi^0\pi^0\eta$
1.26 ± 0.03 ± 0.08		2226	¹ ABLIKIM	10F BES3	$\psi(2S) \rightarrow 3\gamma\pi^+\pi^-,$ $2\gamma\pi^+\pi^-$
1.19 ± 0.08 ± 0.03			PEDLAR	09 CLE3	$\psi(2S) \rightarrow \gamma X$
1.24 ± 0.27 ± 0.15	23		ABLIKIM	06R BES2	$e^+e^- \rightarrow \psi(2S)$
1.54 ± 0.31 ± 0.20	~ 43		BAI	98F BES	$\psi(2S) \rightarrow \pi^+\pi^-2\gamma,$ $\pi^+\pi^-3\gamma$

NODE=M071R44
NODE=M071R44

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 60	90		² BRAUNSCH...	77 DASP	e^+e^-
< 11	90		³ BARTEL	76 CNTR	e^+e^-

¹ Combining the results from $\eta' \rightarrow \pi^+\pi^-\eta$ and $\eta' \rightarrow \pi^+\pi^-\gamma$ decay modes.

² Restated by us using total decay width 228 keV.

³ The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.

NODE=M071R44;LINKAGE=AB
NODE=M071R;LINKAGE=R
NODE=M071R;LINKAGE=C $\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$ Γ_{188}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.73^{+0.29}_{-0.25} OUR AVERAGE				Error includes scale factor of 1.8.

NODE=M071R84
NODE=M071R84

2.84 ± 0.15 ^{+0.03} _{-0.10}	1.9k	1,2 DOBBS	15	$\psi(2S) \rightarrow \gamma\pi\pi$
2.12 ± 0.19 ± 0.32		3,4 BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.08 ± 0.19 ± 0.33	200.6 ± 18.8	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^+\pi^-$	OCCUR=2
2.90 ± 1.08 ± 1.07	29.9 ± 11.1	³ BAI	03C BES	$\psi(2S) \rightarrow \gamma\pi^0\pi^0$	OCCUR=3

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_2(1270))/\Gamma_{\text{total}}] \times [B(f_2(1270) \rightarrow \pi\pi)] = (2.39 \pm 0.09 \pm 0.09) \times 10^{-4}$ which we divide by our best value $B(f_2(1270) \rightarrow \pi\pi) = (84.3^{+2.8}_{-1.0}) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071R84;LINKAGE=A
NODE=M071R84;LINKAGE=B

³ Normalized to $B(\psi(2S) \rightarrow J/\psi\pi^+\pi^-) = 0.305 \pm 0.016$.

⁴ Combining the results from $\pi^+\pi^-$ and $\pi^0\pi^0$ decay modes.

NODE=M071R;LINKAGE=3B
NODE=M071R;LINKAGE=B9

$\Gamma(\gamma f_0(1370) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{189}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
3.1±1.0±1.4	175	¹ DOBBS 15	$\psi(2S) \rightarrow \gamma K \bar{K}$

NODE=M071S75
 NODE=M071S75

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S75;LINKAGE=A

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$ Γ_{190}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
9.3±1.8±0.6	274	^{1,2} DOBBS 15	$\psi(2S) \rightarrow \gamma \pi \pi$

NODE=M071S76
 NODE=M071S76

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f_0(1500))/\Gamma_{\text{total}}] \times [B(f_0(1500) \rightarrow \pi \pi)] = (3.2 \pm 0.6 \pm 0.2) \times 10^{-5}$ which we divide by our best value $B(f_0(1500) \rightarrow \pi \pi) = (34.5 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071S76;LINKAGE=A

² Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S76;LINKAGE=B

 $\Gamma(\gamma f'_2(1525))/\Gamma_{\text{total}}$ Γ_{191}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	COMMENT
3.3±0.8±0.1	136	^{1,2} DOBBS 15	$\psi(2S) \rightarrow \gamma K \bar{K}$

NODE=M071S77
 NODE=M071S77

¹ DOBBS 15 reports $[\Gamma(\psi(2S) \rightarrow \gamma f'_2(1525))/\Gamma_{\text{total}}] \times [B(f'_2(1525) \rightarrow K \bar{K})] = (2.9 \pm 0.6 \pm 0.3) \times 10^{-5}$ which we divide by our best value $B(f'_2(1525) \rightarrow K \bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071S77;LINKAGE=A

² Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S77;LINKAGE=B

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{193}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5 ±0.6 OUR AVERAGE				
3.6 ±0.4 ±0.5	290	¹ DOBBS 15		$\psi(2S) \rightarrow \gamma \pi \pi$
3.01±0.41±1.24	35.6 ± 4.8	² BAI 03C	BES	$\psi(2S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M071R85
 NODE=M071R85

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071R85;LINKAGE=A

² Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R85;LINKAGE=3B

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{194}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.6 ±0.7 OUR AVERAGE					
6.7 ±0.6 ±0.6		375	¹ DOBBS 15		$\psi(2S) \rightarrow \gamma K \bar{K}$
6.04±0.90±1.32		39.6 ± 5.9	^{2,3} BAI 03C	BES	$\psi(2S) \rightarrow \gamma K^+ K^-$

NODE=M071R86
 NODE=M071R86

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 15.6 90 6.8 ± 3.1 ^{2,3} BAI 03C BES $\psi(2S) \rightarrow \gamma K_S^0 K_S^0$

OCCUR=2

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071R86;LINKAGE=A

² Includes unknown branching fractions to $K^+ K^-$ or $K_S^0 K_S^0$. We have multiplied the $K^+ K^-$ result by a factor of 2 and the $K_S^0 K_S^0$ result by a factor of 4 to obtain the $K \bar{K}$ result.

NODE=M071R;LINKAGE=CK

³ Normalized to $B(\psi(2S) \rightarrow J/\psi \pi^+ \pi^-) = 0.305 \pm 0.016$.

NODE=M071R86;LINKAGE=3B

 $\Gamma(\gamma f_0(2100) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{195}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	COMMENT
4.8±0.5±0.9	373	¹ DOBBS 15	$\psi(2S) \rightarrow \gamma \pi \pi$

NODE=M071S78
 NODE=M071S78

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S78;LINKAGE=A

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{196}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	COMMENT
3.2±0.6±0.8	207	¹ DOBBS 15	$\psi(2S) \rightarrow \gamma K \bar{K}$

NODE=M071S79
 NODE=M071S79

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S79;LINKAGE=A

 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi \pi)/\Gamma_{\text{total}}$ Γ_{197}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<5.8 × 10⁻⁶	90	^{1,2} DOBBS 15	$\psi(2S) \rightarrow \gamma \pi \pi$

NODE=M071S80
 NODE=M071S80

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

NODE=M071S80;LINKAGE=A

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $\pi^+ \pi^-$ and $\pi^0 \pi^0$ are $3.2/4.3 \times 10^{-6}$ and $2.6/4.0 \times 10^{-6}$, respectively.

NODE=M071S80;LINKAGE=D0

$\Gamma(\gamma f_J(2220) \rightarrow \gamma K \bar{K})/\Gamma_{\text{total}}$ Γ_{198}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.5 \times 10^{-6}$	90	1,2 DOBBS 15		$\psi(2S) \rightarrow \gamma K \bar{K}$

NODE=M071S81
 NODE=M071S81

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

² For $\Gamma = 20/50$ MeV, the 90% CL upper limits for $K^+ K^-$ and $K_S^0 K_S^0$ are $2.1/4.3 \times 10^{-6}$ and $3.7/5.5 \times 10^{-6}$, respectively.

NODE=M071S81;LINKAGE=A
 NODE=M071S81;LINKAGE=DO

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$ Γ_{199}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.92 ± 0.18 OUR AVERAGE					
$0.85 \pm 0.18 \pm 0.04$		382	¹ ABLIKIM	17X BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$
$1.38 \pm 0.48 \pm 0.09$		13	¹ ABLIKIM	10F BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$

NODE=M071R43
 NODE=M071R43

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2	90	PEDLAR 09	CLE3	$\psi(2S) \rightarrow \gamma X$
< 90	90	BAI 98F	BES	$\psi(2S) \rightarrow \pi^+ \pi^- 3\gamma$
< 200	90	YAMADA 77	DASP	$e^+ e^- \rightarrow 3\gamma$

¹ Combining the results from $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow 3\pi^0$ decay modes.

NODE=M071R43;LINKAGE=AB

 $\Gamma(\gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{200}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.71 ± 1.25 ± 1.64	418	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M071R04
 NODE=M071R04

 $\Gamma(\gamma \eta(1405) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{202}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$

NODE=M071R61
 NODE=M071R61

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3	90	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$
< 1.2	90	¹ SCHARRE 80	MRK1	$e^+ e^-$

OCCUR=2

¹ Includes unknown branching fraction $\eta(1405) \rightarrow K \bar{K} \pi$.

NODE=M071R;LINKAGE=E

 $\Gamma(\gamma \eta(1405) \rightarrow \gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{203}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.36 ± 0.25 ± 0.05	10	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M071R05
 NODE=M071R05

 $\Gamma(\gamma \eta(1405) \rightarrow \gamma f_0(980) \pi^0 \rightarrow \gamma \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{204}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.0 × 10⁻⁷	90	ABLIKIM 17AJ	BES3	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- \pi^0$

NODE=M071P09
 NODE=M071P09

 $\Gamma(\gamma \eta(1475) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$ Γ_{206}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma K^+ K^- \pi^0$

NODE=M071R06
 NODE=M071R06

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.5	90	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma K_S^0 K^+ \pi^- + \text{c.c.}$
---------	----	-------------	------	---

OCCUR=2

 $\Gamma(\gamma \eta(1475) \rightarrow \gamma \eta \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{207}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.88	90	ABLIKIM 06R	BES2	$\psi(2S) \rightarrow \gamma \eta \pi^+ \pi^-$

NODE=M071R07
 NODE=M071R07

 $\Gamma(\gamma K^* K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{208}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
37.0 ± 6.1 ± 7.2	237	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S29
 NODE=M071S29

 $\Gamma(\gamma K^* \bar{K}^* \pi^0)/\Gamma_{\text{total}}$ Γ_{209}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
24.0 ± 4.5 ± 5.0	41	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S30
 NODE=M071S30

 $\Gamma(\gamma K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{210}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
25.6 ± 3.6 ± 3.6	115	ABLIKIM 07D	BES2	$e^+ e^- \rightarrow \psi(2S)$

NODE=M071S31
 NODE=M071S31

$\Gamma(\gamma K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_{211}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
19.1±2.7±4.3	132	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071S32 NODE=M071S32
$\Gamma(\gamma K^+ K^- 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$					Γ_{212}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<22	90	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071S35 NODE=M071S35
$\Gamma(\gamma 2(K^+ K^-))/\Gamma_{\text{total}}$					Γ_{213}/Γ	
VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT		
<4	90	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071S37 NODE=M071S37
$\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{214}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
3.9 ±0.5 OUR AVERAGE					Error includes scale factor of 2.0.	
4.18±0.26±0.18	348	¹ ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$	
2.9 ±0.4 ±0.4	142	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$	
¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.						NODE=M071S33;LINKAGE=AL
$\Gamma(\gamma f_2(1950) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{215}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
1.2±0.2±0.1	111	¹ ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$	NODE=M071S46 NODE=M071S46
¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.						NODE=M071S46;LINKAGE=AL
$\Gamma(\gamma f_2(2150) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{216}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.72±0.18±0.03	73	¹ ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$	NODE=M071S47 NODE=M071S47
¹ From a fit of the $p\bar{p}$ mass distribution to a combination of $\gamma f_2(1950)$, $\gamma f_2(2150)$, and $\gamma p\bar{p}$ phase space, for $M(p\bar{p}) < 2.85$ GeV, and accounting for backgrounds from $\psi(2S) \rightarrow \pi^0 p\bar{p}$ and continuum.						NODE=M071S47;LINKAGE=AL
$\Gamma(\gamma X(1835) \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{217}/Γ	
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
4.57±0.36^{+1.77}_{-4.26}		ABLIKIM	12D	BES3	$J/\psi \rightarrow \gamma p\bar{p}$	NODE=M071S48 NODE=M071S48
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<1.6	90	ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$	
<5.4	90	ABLIKIM	07D	BES	$\psi(2S) \rightarrow \gamma p\bar{p}$	
$\Gamma(\gamma X \rightarrow \gamma p\bar{p})/\Gamma_{\text{total}}$					Γ_{218}/Γ	
For a narrow resonance in the range $2.2 < M(X) < 2.8$ GeV.						
VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT		
<2	90	ALEXANDER	10	CLEO	$\psi(2S) \rightarrow \gamma p\bar{p}$	NODE=M071S49 NODE=M071S49 NODE=M071S49
$\Gamma(\gamma p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$					Γ_{219}/Γ	
VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT		
2.8±1.2±0.7	17	ABLIKIM	07D	BES2	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071S34 NODE=M071S34
$\Gamma(\gamma\gamma J/\psi)/\Gamma_{\text{total}}$					Γ_{221}/Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
3.1±0.6^{+0.8}_{-1.0}	1.1k	ABLIKIM	120	BES3	$e^+ e^- \rightarrow \psi(2S)$	NODE=M071S55 NODE=M071S55
• • • We do not use the following data for averages, fits, limits, etc. • • •						
3.2±0.6	1.1k	¹ ABLIKIM	17N	BES3	$\psi(2S) \rightarrow \gamma\gamma J/\psi$	
¹ Uses $B(J/\psi \rightarrow e^+e^-) = (5.971 \pm 0.032)\%$ and $B(J/\psi \rightarrow \mu^+\mu^-) = (5.961 \pm 0.033)\%$. No systematic error estimation.						NODE=M071S55;LINKAGE=A

$\Gamma(e^+e^-\eta')/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.26 OUR AVERAGE				
1.99±0.33±0.12	57	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$, $\eta' \rightarrow \gamma \pi^+ \pi^-$
1.79±0.38±0.11	20	ABLIKIM	18Z BES3	$\psi(2S) \rightarrow \eta' e^+ e^-$, $\eta' \rightarrow \eta \pi^+ \pi^-$

 Γ_{222}/Γ NODE=M071P12
NODE=M071P12

OCCUR=2

 $\Gamma(e^+e^-\eta_c(1S))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.77±0.40±0.18	3k	¹ ABLIKIM	22AX BES3	$e^+e^- \rightarrow \psi(2S)$

 Γ_{223}/Γ NODE=M071P44
NODE=M071P44¹ From a fit to the recoil mass distribution of e^+e^- with inclusive $\eta_c(1S)$ decays.

NODE=M071P44;LINKAGE=A

 $\Gamma(e^+e^-\chi_{c0}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.6±2.5 OUR AVERAGE				[(10.6 ± 2.4) × 10 ⁻⁴ OUR 2023 AVERAGE]
10.6±2.4±0.7	48	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

 Γ_{224}/Γ NODE=M071P01
NODE=M071P01

NEW

¹ ABLIKIM 17I reports (11.7 ± 2.5 ± 1.0) × 10⁻⁴ from a measurement of [$\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c0}(1P))/\Gamma_{\text{total}}$] × [B($\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)$)] assuming B($\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)$) = (1.27 ± 0.06) × 10⁻², which we rescale to our best value B($\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)$) = (1.41 ± 0.09) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P01;LINKAGE=B

 $\Gamma(e^+e^-\chi_{c0}(1P))/\Gamma(\gamma\chi_{c0}(1P))$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
9.4±1.9±0.6	48	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

 $\Gamma_{224}/\Gamma_{179}$ NODE=M071P04
NODE=M071P04¹ Uses B($\psi(2S) \rightarrow \gamma\chi_{c0}(1P)$) × B($\chi_{c0}(1P) \rightarrow \gamma J/\psi(1S)$) = (15.8 ± 0.3 ± 0.6) × 10⁻⁴ from ABLIKIM 17N and accounts for common systematic errors.

NODE=M071P04;LINKAGE=A

 $\Gamma(e^+e^-\chi_{c1}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.5±0.7 OUR AVERAGE				[(8.5 ± 0.6) × 10 ⁻⁴ OUR 2023 AVERAGE]
8.5±0.6±0.3	873	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

 Γ_{225}/Γ NODE=M071P02
NODE=M071P02

NEW

¹ ABLIKIM 17I reports (8.6 ± 0.3 ± 0.6) × 10⁻⁴ from a measurement of [$\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c1}(1P))/\Gamma_{\text{total}}$] × [B($\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$)] assuming B($\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$) = (33.9 ± 1.2) × 10⁻², which we rescale to our best value B($\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$) = (34.3 ± 1.3) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P02;LINKAGE=B

 $\Gamma(e^+e^-\chi_{c1}(1P))/\Gamma(\gamma\chi_{c1}(1P))$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
8.3±0.3±0.4	873	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

 $\Gamma_{225}/\Gamma_{180}$ NODE=M071P05
NODE=M071P05¹ Uses B($\psi(2S) \rightarrow \gamma\chi_{c1}(1P)$) × B($\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)$) = (351.8 ± 1.0 ± 12.0) × 10⁻⁴ from ABLIKIM 17N and accounts for common systematic errors.

NODE=M071P05;LINKAGE=A

 $\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8±0.8 OUR AVERAGE				[(7.0 ± 0.8) × 10 ⁻⁴ OUR 2023 AVERAGE]
6.8±0.7±0.3	227	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

 Γ_{226}/Γ NODE=M071P03
NODE=M071P03

NEW

¹ ABLIKIM 17I reports (6.9 ± 0.5 ± 0.6) × 10⁻⁴ from a measurement of [$\Gamma(\psi(2S) \rightarrow e^+e^-\chi_{c2}(1P))/\Gamma_{\text{total}}$] × [B($\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$)] assuming B($\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$) = (19.2 ± 0.7) × 10⁻², which we rescale to our best value B($\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$) = (19.5 ± 0.8) × 10⁻². Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M071P03;LINKAGE=B

 $\Gamma(e^+e^-\chi_{c2}(1P))/\Gamma(\gamma\chi_{c2}(1P))$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
6.6±0.5±0.4	227	¹ ABLIKIM	17I BES3	$\psi(2S) \rightarrow e^+e^-\gamma J/\psi$

 $\Gamma_{226}/\Gamma_{181}$ NODE=M071P06
NODE=M071P06¹ Uses B($\psi(2S) \rightarrow \gamma\chi_{c2}(1P)$) × B($\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)$) = (199.6 ± 0.8 ± 7.0) × 10⁻⁴ from ABLIKIM 17N and accounts for common systematic errors.

NODE=M071P06;LINKAGE=A

WEAK DECAYS

$\Gamma(D^0 e^+ e^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{227}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-7}$	90	¹ ABLIKIM	17AF BES3	$e^+ e^- \rightarrow \psi(2S)$	
¹ Using D^0 decays to $K^- \pi^+$, $K^- \pi^+ \pi^0$, and $K^- \pi^+ \pi^+ \pi^-$.					

NODE=M071330

NODE=M071P07
NODE=M071P07

NODE=M071P07;LINKAGE=A

$\Gamma(\Lambda_c^+ \bar{\Sigma}^- + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{228}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-5}$	90	¹ ABLIKIM	23 BES3	$e^+ e^- \rightarrow \psi(2S)$	
¹ Using $\Lambda_c^+ \rightarrow p K^- \pi^+$ and $\bar{\Sigma}^- \rightarrow \bar{p} \pi^0$.					

NODE=M071P43
NODE=M071P43

NODE=M071P43;LINKAGE=A

NODE=M071320

$\Gamma(\text{invisible})/\Gamma(e^+ e^-)$					Γ_{229}/Γ_7
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.0	90	LEES	13I BABR	$B \rightarrow K^{(*)} \psi(2S)$	

NODE=M071S64
NODE=M071S64 $\psi(2S)$ CROSS-PARTICLE BRANCHING RATIOS

For measurements involving $B(\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)) \times B(\chi_{cJ}(1P) \rightarrow X)$ see the corresponding entries in the $\chi_{cJ}(1P)$ sections.

NODE=M071240

NODE=M071240

MULTIPOLE AMPLITUDE RATIOS IN RADIATIVE DECAYS

 $\psi(2S) \rightarrow \gamma \chi_{cJ}(1P)$ and $\chi_{cJ} \rightarrow \gamma J/\psi(1S)$

NODE=M071250

 $a_2(\chi_{c1})/a_2(\chi_{c2})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
63 ± 7 OUR AVERAGE				
61.7 ± 8.3	253k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
67 ⁺¹⁹ ₋₁₃	59k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M071QAR
NODE=M071QAR¹ Statistical and systematic errors combined.² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $a_2(\chi_{c1}(1P))$ and $a_2(\chi_{c2}(1P))$ from ARTUSO 09.NODE=M071QAR;LINKAGE=A
NODE=M071QAR;LINKAGE=AR $b_2(\chi_{c2})/b_2(\chi_{c1})$ Magnetic quadrupole transition amplitude ratio

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
60 ± 31 OUR AVERAGE				
74 ± 40	253k	¹ ABLIKIM	17N BES3	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
37 ⁺⁵³ ₋₄₇	59k	² ARTUSO	09 CLEO	$\psi(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M071QBR
NODE=M071QBR¹ Statistical and systematic errors combined. Derived from the reported measurement of $b_2(\chi_{c1})/b_2(\chi_{c2}) = 1.35 \pm 0.72$.² Statistical and systematic errors combined. Using values from fits with floating $M2$ amplitudes $a_2(\chi_{c1})$, $a_2(\chi_{c2})$, $b_2(\chi_{c1})$, $b_2(\chi_{c2})$ and fixed $E3$ amplitudes of $a_3(\chi_{c2}) = b_3(\chi_{c2}) = 0$. Not independent of values for $b_2(\chi_{c1}(1P))$ and $b_2(\chi_{c2}(1P))$ from ARTUSO 09.

NODE=M071QBR;LINKAGE=A

NODE=M071QBR;LINKAGE=AR

 $\psi(2S)$ REFERENCES

AAIJ	23AP	JHEP 2307 084	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23	CP C47 013002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BA	PR D108 052001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BE	PR D108 092011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	23BV	PR D108 112014	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	23	PR D107 072001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LIAO	23	PR D107 112007	L. Liao <i>et al.</i>	
ABLIKIM	22AP	PR D106 072006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AQ	PR D106 072007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AV	JHEP 2212 016	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AX	PR D106 112002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AY	PR D106 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AZ	PR D106 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	21AL	PR D104 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AO	PR D104 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21AT	JHEP 2111 226	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21E	PRL 126 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21L	PR D103 112004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21S	PL B820 136576	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	21Z	PRL 127 082002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	21	PR D103 092001	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	21C	PR D104 112004	J.P. Lees <i>et al.</i>	(BABAR Collab.)

NODE=M071

REFID=62412
REFID=61897
REFID=62421
REFID=62427
REFID=62518
REFID=62061
REFID=62279
REFID=61888
REFID=61889
REFID=61899
REFID=61901
REFID=61902
REFID=61903
REFID=61634
REFID=61444
REFID=61447
REFID=61463
REFID=61033
REFID=61117
REFID=61152
REFID=61265
REFID=61113
REFID=61451

ABLIKIM	20F	PR D101 032008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60256
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19AO	PR D99 112010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59892
ABLIKIM	19AT	PR D100 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59989
ABLIKIM	19AU	PR D100 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59996
ABLIKIM	19BA	PR D100 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60024
ABLIKIM	19I	PR D99 012014	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59605
ABLIKIM	19N	PR D99 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59615
ABLIKIM	18Q	PR D97 091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58933
ABLIKIM	18T	PR D98 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58975
ABLIKIM	18Z	PL B783 452	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59038
ANASHIN	18	PL B781 174	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=59013
LEES	18E	PR D98 112015	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=59505
ABLIKIM	17AF	PR D96 111101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58315
ABLIKIM	17AJ	PR D96 112008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58322
ABLIKIM	17AK	PR D96 112012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58324
ABLIKIM	17E	PL B770 217	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57903
ABLIKIM	17I	PRL 118 221802	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57931
ABLIKIM	17L	PR D95 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57967
ABLIKIM	17N	PR D95 072004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57978
ABLIKIM	17U	PR D96 032001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58026
ABLIKIM	17X	PR D96 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58216
DOBBS	17	PR D96 092004	S. Dobbs <i>et al.</i>	(NWES, WAYN)	REFID=58670
LEES	17A	PR D95 052001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=57966
AAJ	16Y	JHEP 1605 132	R. Aaj <i>et al.</i>	(LHCb Collab.)	REFID=57333
ABLIKIM	16L	PR D93 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57510
ABLIKIM	15I	PR D91 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56774
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56787
ANASHIN	15	PL B749 50	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=56792
DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)	REFID=56805
LEES	15J	PR D92 072008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56988
ABLIKIM	14G	PR D89 112006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55898
DOBBS	14	PL B739 90	S. Dobbs <i>et al.</i>	(NWES, WAYN)	REFID=56333
ABLIKIM	13A	PRL 110 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54834
ABLIKIM	13D	PR D87 012007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54879
ABLIKIM	13F	PR D87 052007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54920
ABLIKIM	13M	PR D87 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55386
ABLIKIM	13R	PR D88 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55402
ABLIKIM	13S	PR D88 032010	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55403
ABLIKIM	13W	PR D88 112007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55634
LEES	13I	PR D87 112005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55161
LEES	13O	PR D87 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55293
LEES	13Q	PR D88 032013	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55404
LEES	13Y	PR D88 072009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55589
AAJ	12H	EPJ C72 1972	R. Aaj <i>et al.</i>	(LHCb Collab.)	REFID=54056
ABLIKIM	12D	PRL 108 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54269
ABLIKIM	12G	PRL 109 042003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54272
ABLIKIM	12H	PL B710 594	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54273
ABLIKIM	12L	PR D86 072011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54739
ABLIKIM	12M	PR D86 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54740
ABLIKIM	12O	PRL 109 172002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54742
ABLIKIM	12Q	CP C36 1040	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=54864
ANASHIN	12	PL B711 280	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54038
LEES	12E	PR D85 112009	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54297
LEES	12F	PR D86 012008	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54298
METREVELI	12	PR D85 092007	Z. Metreveli <i>et al.</i>	(NWES, FLOR, WAYN+)	REFID=54304
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)	REFID=53960
ABLIKIM	10B	PRL 104 132002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53348
ABLIKIM	10F	PRL 105 261801	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53630
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=53525
CRONIN-HENNESSY	10	PR D81 052002	D. Cronin-Hennessey <i>et al.</i>	(CLEO Collab.)	REFID=53233
ADAMS	09	PR D80 051106	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=53103
ARTUSO	09	PR D80 112003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=53206
LIBBY	09	PR D80 072002	J. Libby <i>et al.</i>	(CLEO Collab.)	REFID=53124
MITCHELL	09	PRL 102 011801	R.E. Mitchell <i>et al.</i>	(CLEO Collab.)	REFID=52676
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08C	PL B659 789	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52130
DOBBS	08A	PRL 101 182003	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52579
MENDEZ	08	PR D78 011102	H. Mendez <i>et al.</i>	(CLEO Collab.)	REFID=52684
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ABLIKIM	07C	PL B648 149	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51636
ABLIKIM	07D	PRL 99 011802	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=51725
ABLIKIM	07H	PR D76 092003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52046
ANASHIN	07	JETPL 85 347	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=51655
		Translated from ZETFP 85 429.			
ANDREOTTI	07	PL B654 74	M. Andreotti <i>et al.</i>	(Femilab E835 Collab.)	REFID=51944
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07AU	PR D76 092005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52049
		Also PR D77 119902E (err.)	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52266
AUBERT	07BD	PR D76 092006	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52050
PDG	07	Unofficial 2007 WWW edition		(PDG Collab.)	REFID=52717; ERROR=8
PEDLAR	07	PR D75 011102	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=51630
ABLIKIM	06G	PR D73 052004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51048
ABLIKIM	06I	PR D74 012004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51126
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06R	PR D74 072001	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51447
ABLIKIM	06W	PR D74 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51560
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT	06D	PR D73 052003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51047
AUBERT, BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511

DOBBS	06A	PR D74 011105	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=51158
ABLIKIM	05E	PR D71 072006	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50757
ABLIKIM	05H	PR D72 012002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50759
ABLIKIM	05I	PL B614 37	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50758
ABLIKIM	05J	PL B619 247	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50760
ABLIKIM	05O	PL B630 21	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50846
ADAM	05	PRL 94 012005	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50451
ADAM	05A	PRL 94 232002	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50763
ANDREOTTI	05	PR D71 032006	M. Andreotti <i>et al.</i>	(FNAL E835 Collab.)	REFID=50497
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
BRIERE	05	PRL 95 062001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50785
PEDLAR	05	PR D72 051108	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=50808
ROSNER	05	PRL 95 102003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=50812
ABLIKIM	04B	PR D70 012003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=49741
ABLIKIM	04K	PR D70 112003	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50327
ABLIKIM	04L	PR D70 112007	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50328
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
BAI	04B	PRL 92 052001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49608
BAI	04C	PR D69 072001	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49749
BAI	04D	PL B589 7	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49750
BAI	04G	PR D70 012004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49753
BAI	04I	PR D70 012006	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49755
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
SETH	04	PR D69 097503	K.K. Seth		REFID=49779
AULCHENKO	03	PL B573 63	V.M. Aulchenko <i>et al.</i>	(KEDR Collab.)	REFID=49579
BAI	03B	PR D67 052002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49186
BAI	03C	PR D67 032004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49190
AUBERT	02B	PR D65 031101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48548
BAI	02	PR D65 052004	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48578
BAI	02B	PL B550 24	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=49171
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
PDG	02	PR D66 010001	K. Hagiwara <i>et al.</i>	(PDG Collab.)	REFID=48632
BAI	01	PR D63 032002	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=48003
AMBROGIANI	00A	PR D62 032004	M. Ambrogiani <i>et al.</i>	(FNAL E835 Collab.)	REFID=47939
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
BAI	99C	PRL 83 1918	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=47420
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
BAI	98F	PR D58 097101	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46340
BAI	98J	PRL 81 5080	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46554
ARMSTRONG	97	PR D55 1153	T.A. Armstrong <i>et al.</i>	(E760 Collab.)	REFID=45416
GRIBUSHIN	96	PR D53 4723	A. Gribushin <i>et al.</i>	(E672 and E706 Collab.)	REFID=44739
ARMSTRONG	93B	PR D47 772	T.A. Armstrong <i>et al.</i>	(FNAL E760 Collab.)	REFID=43307
ALEXANDER	89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
GAISER	86	PR D34 711	J. Gaiser <i>et al.</i>	(Crystal Ball Collab.)	REFID=22012
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41 733.			
FRANKLIN	83	PRL 51 963	M.E.B. Franklin <i>et al.</i>	(LBL, SLAC)	REFID=22216
EDWARDS	82C	PRL 48 70	C. Edwards <i>et al.</i>	(CIT, HARV, PRIN+)	REFID=22173
LEMOIGNE	82	PL 113B 509	Y. Lemoigne <i>et al.</i>	(SACL, LOIC, SHMP+)	REFID=22084
HIMEL	80	PRL 44 920	T. Himel <i>et al.</i>	(LBL, SLAC)	REFID=22119
OREGLIA	80	PRL 45 959	M.J. Oreglia <i>et al.</i>	(SLAC, CIT, HARV+)	REFID=22207
SCHARRE	80	PL 97B 329	D.L. Scharre <i>et al.</i>	(SLAC, LBL)	REFID=21329
ZHOLENTZ	80	PL 96B 214	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10320
Also		SJNP 34 814	A.A. Zholents <i>et al.</i>	(NOVO)	REFID=10321
		Translated from YAF 34 1471.			
BRANDELIK	79B	NP B160 426	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22115
BRANDELIK	79C	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22114
BARTEL	78B	PL 79B 492	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22111
TANENBAUM	78	PR D17 1731	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL)	REFID=22112
BIDDICK	77	PRL 38 1324	C.J. Biddick <i>et al.</i>	(UCSD, UMD, PAVI+)	REFID=22059
BRAUNSCH...	77	PL 67B 249	W. Braunschweig <i>et al.</i>	(DASP Collab.)	REFID=22197
BURMESTER	77	PL 66B 395	J. Burmester <i>et al.</i>	(DESY, HAMB, SIEG+)	REFID=22198
FELDMAN	77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)	REFID=22062
YAMADA	77	Hamburg Conf. 69	S. Yamada	(DASP Collab.)	REFID=22064
BARTEL	76	PL 64B 483	W. Bartel <i>et al.</i>	(DESY, HEIDP)	REFID=22192
TANENBAUM	76	PRL 36 402	W.M. Tanenbaum <i>et al.</i>	(SLAC, LBL) IG	REFID=22194
WHITAKER	76	PRL 37 1596	J.S. Whitaker <i>et al.</i>	(SLAC, LBL)	REFID=22151
ABRAMS	75	Stanford Symp. 25	G.S. Abrams	(LBL)	REFID=22176
ABRAMS	75B	PRL 34 1181	G.S. Abrams <i>et al.</i>	(LBL, SLAC)	REFID=22177
BOYARSKI	75C	Palermo Conf. 54	A.M. Boyarski <i>et al.</i>	(SLAC, LBL)	REFID=22179
HILGER	75	PRL 35 625	E. Hilger <i>et al.</i>	(STAN, PENN)	REFID=22186
LIBERMAN	75	Stanford Symp. 55	A.D. Liberman	(STAN)	REFID=22046
LUTH	75	PRL 35 1124	V. Luth <i>et al.</i>	(SLAC, LBL) JPC	REFID=22188
WIJK	75	Stanford Symp. 69	B.H. Wiik	(DESY)	REFID=22050

$\psi(3770)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M053

 $\psi(3770)$ MASS (MeV)

NODE=M053M

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

NODE=M053M

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
3773.7±0.7 OUR FIT Error includes scale factor of 2.5. [3773.7 ± 0.4 MeV OUR 2023 FIT Scale factor = 1.4]

NODE=M053M

NEW

3778.1±0.7 OUR AVERAGE

3778.1±0.7±0.6		¹ AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$
3779.2 ^{+1.8+0.6} _{-1.7-0.8}		² ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
3775.5±2.4±0.5	57	AUBERT	08B BABR	$B \rightarrow D\bar{D}K$
3776 ±5 ±4	68	BRODZICKA	08 BELL	$B^+ \rightarrow D^0\bar{D}^0K^+$
3778.8±1.9±0.9		AUBERT	07BE BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3779.8±0.6		³ SHAMOV	17 RVUE	$e^+e^- \rightarrow D\bar{D}$, hadrons
3772.0±1.9		^{4,5} ABLIKIM	08D BES2	$e^+e^- \rightarrow \text{hadrons}$
3778.4±3.0±1.3	34	CHISTOV	04 BELL	Sup. by BRODZICKA 08

¹ Measured in prompt hadroproduction.

NODE=M053M;LINKAGE=B

² Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

NODE=M053M;LINKAGE=AN

³ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.

NODE=M053M;LINKAGE=A

⁴ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

NODE=M053M;LINKAGE=AB

⁵ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

NODE=M053M;LINKAGE=NI

 $m_{\psi(3770)} = m_{\psi(2S)}$

NODE=M053DM

OUR FIT includes measurements of $m_{\psi(2S)}$, $m_{\psi(3770)}$, and $m_{\psi(3770)} - m_{\psi(2S)}$.

NODE=M053DM

VALUE (MeV) DOCUMENT ID TECN COMMENT
87.6±0.7 OUR FIT Error includes scale factor of 2.4. [87.6 ± 0.4 MeV OUR 2023 FIT Scale factor = 1.4]

NODE=M053DM

NEW

86.6±0.7 OUR AVERAGE Error includes scale factor of 2.0. See the ideogram below.


86.9±0.4		¹ ABLIKIM	07E BES2	$e^+e^- \rightarrow \text{hadrons}$
86.7±0.7		ABLIKIM	06L BES2	$e^+e^- \rightarrow \text{hadrons}$
80 ±2		SCHINDLER	80 MRK2	e^+e^-
86 ±2		² BACINO	78 DLCO	e^+e^-
88 ±3		RAPIDIS	77 LGW	e^+e^-

¹ BES-II $\psi(2S)$ mass subtracted (see ABLIKIM 06L).

NODE=M053DM;LINKAGE=AK

² SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).

NODE=M053DM;LINKAGE=S



$\psi(3770)$ WIDTH

NODE=M053W

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
27.2± 1.0 OUR FIT				
27.5± 0.9 OUR AVERAGE				
24.9 ⁺ 4.6 ⁺ 0.5 - 4.0 ⁻ 1.1		1 ANASHIN 12A	KEDR	$e^+e^- \rightarrow D\bar{D}$
30.4± 8.5		2,3 ABLIKIM 08D	BES2	$e^+e^- \rightarrow$ hadrons
27 ±10 ±5	68	BRODZICKA 08	BELL	$B^+ \rightarrow D^0\bar{D}^0 K^+$
28.5± 1.2±0.2		3 ABLIKIM 07E	BES2	$e^+e^- \rightarrow$ hadrons
23.5± 3.7±0.9		AUBERT 07BE	BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
26.9± 2.4±0.3		3 ABLIKIM 06L	BES2	$e^+e^- \rightarrow$ hadrons
24 ± 5		3 SCHINDLER 80	MRK2	e^+e^-
24 ± 5		3 BACINO 78	DLCO	e^+e^-
28 ± 5		3 RAPIDIS 77	LGW	e^+e^-

NODE=M053W

• • • We do not use the following data for averages, fits, limits, etc. • • •

25.8± 1.3		4 SHAMOV 17	RVUE	$e^+e^- \rightarrow D\bar{D},$ hadrons
-----------	--	-------------	------	--

¹ Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.

² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.

³ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.

⁴ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.

NODE=M053W;LINKAGE=AN

NODE=M053W;LINKAGE=AB

NODE=M053W;LINKAGE=NI

NODE=M053W;LINKAGE=A

$\psi(3770)$ DECAY MODES

NODE=M053220;NODE=M053

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $D\bar{D}$	(93 ⁺⁸ / ₋₉) %	S=2.0
Γ_2 $D^0\bar{D}^0$	(52 ⁺⁴ / ₋₅) %	S=2.0
Γ_3 D^+D^-	(41 ±4) %	S=2.0
Γ_4 $J/\psi X$	(5.0 ±2.2) × 10 ⁻³	
Γ_5 $J/\psi \pi^+\pi^-$	(1.93±0.28) × 10 ⁻³	
Γ_6 $J/\psi \pi^0\pi^0$	(8.0 ±3.0) × 10 ⁻⁴	
Γ_7 $J/\psi \eta$	(8.7 ±1.2) × 10 ⁻⁴	
Γ_8 $J/\psi \pi^0$	< 2.8 × 10 ⁻⁴	CL=90%
Γ_9 e^+e^-	(9.6 ±0.7) × 10 ⁻⁶	S=1.3

DESIG=2

DESIG=5

DESIG=6

DESIG=235

DESIG=4

DESIG=46

DESIG=47

DESIG=48

DESIG=1

Decays to light hadrons

NODE=M053;CLUMP=H

Γ ₁₀	$b_1(1235)\pi$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=20
Γ ₁₁	$\phi\eta'$	< 2.3	$\times 10^{-5}$	CL=90%	DESIG=17
Γ ₁₂	$\omega\eta'$	< 4	$\times 10^{-4}$	CL=90%	DESIG=16
Γ ₁₃	$\rho^0\eta'$	< 6	$\times 10^{-4}$	CL=90%	DESIG=15
Γ ₁₄	$\phi\eta$	(3.1 ±0.7)	$\times 10^{-4}$		DESIG=8
Γ ₁₅	$\omega\eta$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=14
Γ ₁₆	$\rho^0\eta$	< 5	$\times 10^{-4}$	CL=90%	DESIG=13
Γ ₁₇	$\phi\pi^0$	< 3	$\times 10^{-5}$	CL=90%	DESIG=12
Γ ₁₈	$\omega\pi^0$	< 6	$\times 10^{-4}$	CL=90%	DESIG=11
Γ ₁₉	$\pi^+\pi^-\pi^0$	< 5	$\times 10^{-6}$	CL=90%	DESIG=9
Γ ₂₀	$\rho\pi$	< 5	$\times 10^{-6}$	CL=90%	DESIG=10
Γ ₂₁	K^+K^-				DESIG=234
Γ ₂₂	$K^*(892)^+K^- + \text{c.c.}$	< 1.4	$\times 10^{-5}$	CL=90%	DESIG=19
Γ ₂₃	$K^*(892)^0K^0 + \text{c.c.}$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=18
Γ ₂₄	$K_S^0K_L^0$	< 1.2	$\times 10^{-5}$	CL=90%	DESIG=3
Γ ₂₅	$2(\pi^+\pi^-)$	< 1.12	$\times 10^{-3}$	CL=90%	DESIG=21
Γ ₂₆	$2(\pi^+\pi^-)\pi^0$	< 1.06	$\times 10^{-3}$	CL=90%	DESIG=22
Γ ₂₇	$2(\pi^+\pi^-\pi^0)$	< 5.85	%	CL=90%	DESIG=208
Γ ₂₈	$\omega\pi^+\pi^-$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=24
Γ ₂₉	$3(\pi^+\pi^-)$	< 9.1	$\times 10^{-3}$	CL=90%	DESIG=52
Γ ₃₀	$3(\pi^+\pi^-)\pi^0$	< 1.37	%	CL=90%	DESIG=55
Γ ₃₁	$3(\pi^+\pi^-)2\pi^0$	< 11.74	%	CL=90%	DESIG=210
Γ ₃₂	$\eta\pi^+\pi^-$	< 1.24	$\times 10^{-3}$	CL=90%	DESIG=23
Γ ₃₃	$\pi^+\pi^-2\pi^0$	< 8.9	$\times 10^{-3}$	CL=90%	DESIG=206
Γ ₃₄	$\rho^0\pi^+\pi^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=64
Γ ₃₅	$\eta3\pi$	< 1.34	$\times 10^{-3}$	CL=90%	DESIG=25
Γ ₃₆	$\eta2(\pi^+\pi^-)$	< 2.43	%	CL=90%	DESIG=53
Γ ₃₇	$\eta\rho^0\pi^+\pi^-$	< 1.45	%	CL=90%	DESIG=221
Γ ₃₈	$\eta'3\pi$	< 2.44	$\times 10^{-3}$	CL=90%	DESIG=26
Γ ₃₉	$K^+K^-\pi^+\pi^-$	< 9.0	$\times 10^{-4}$	CL=90%	DESIG=27
Γ ₄₀	$\phi\pi^+\pi^-$	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=28
Γ ₄₁	$K^+K^-2\pi^0$	< 4.2	$\times 10^{-3}$	CL=90%	DESIG=207
Γ ₄₂	$4(\pi^+\pi^-)$	< 1.67	%	CL=90%	DESIG=62
Γ ₄₃	$4(\pi^+\pi^-)\pi^0$	< 3.06	%	CL=90%	DESIG=63
Γ ₄₄	$\phi f_0(980)$	< 4.5	$\times 10^{-4}$	CL=90%	DESIG=29
Γ ₄₅	$K^+K^-\pi^+\pi^-\pi^0$	< 2.36	$\times 10^{-3}$	CL=90%	DESIG=30
Γ ₄₆	$K^+K^-\rho^0\pi^0$	< 8	$\times 10^{-4}$	CL=90%	DESIG=67
Γ ₄₇	$K^+K^-\rho^+\pi^-$	< 1.46	%	CL=90%	DESIG=68
Γ ₄₈	ωK^+K^-	< 3.4	$\times 10^{-4}$	CL=90%	DESIG=32
Γ ₄₉	$\phi\pi^+\pi^-\pi^0$	< 3.8	$\times 10^{-3}$	CL=90%	DESIG=69
Γ ₅₀	$K^{*0}K^-\pi^+\pi^0 + \text{c.c.}$	< 1.62	%	CL=90%	DESIG=70
Γ ₅₁	$K^{*+}K^-\pi^+\pi^- + \text{c.c.}$	< 3.23	%	CL=90%	DESIG=71
Γ ₅₂	$K^+K^-\pi^+\pi^-2\pi^0$	< 2.67	%	CL=90%	DESIG=209
Γ ₅₃	$K^+K^-2(\pi^+\pi^-)$	< 1.03	%	CL=90%	DESIG=57
Γ ₅₄	$K^+K^-2(\pi^+\pi^-)\pi^0$	< 3.60	%	CL=90%	DESIG=58
Γ ₅₅	ηK^+K^-	< 4.1	$\times 10^{-4}$	CL=90%	DESIG=31
Γ ₅₆	$\eta K^+K^-\pi^+\pi^-$	< 1.24	%	CL=90%	DESIG=222
Γ ₅₇	$\rho^0 K^+K^-$	< 5.0	$\times 10^{-3}$	CL=90%	DESIG=65
Γ ₅₈	$2(K^+K^-)$	< 6.0	$\times 10^{-4}$	CL=90%	DESIG=33
Γ ₅₉	ϕK^+K^-	< 7.5	$\times 10^{-4}$	CL=90%	DESIG=34
Γ ₆₀	$2(K^+K^-)\pi^0$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=35
Γ ₆₁	$2(K^+K^-)\pi^+\pi^-$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=59
Γ ₆₂	$K_S^0K^-\pi^+$	< 3.2	$\times 10^{-3}$	CL=90%	DESIG=200
Γ ₆₃	$K_S^0K^-\pi^+\pi^0$	< 1.33	%	CL=90%	DESIG=201
Γ ₆₄	$K_S^0K^-\rho^+$	< 6.6	$\times 10^{-3}$	CL=90%	DESIG=214
Γ ₆₅	$K_S^0K^-2\pi^+\pi^-$	< 8.7	$\times 10^{-3}$	CL=90%	DESIG=202
Γ ₆₆	$K_S^0K^-\pi^+\rho^0$	< 1.6	%	CL=90%	DESIG=215

Γ ₆₇	$K_S^0 K^- \pi^+ \eta$	< 1.3	%	CL=90%	DESIG=216
Γ ₆₈	$K_S^0 K^- 2\pi^+ \pi^- \pi^0$	< 4.18	%	CL=90%	DESIG=203
Γ ₆₉	$K_S^0 K^- 2\pi^+ \pi^- \eta$	< 4.8	%	CL=90%	DESIG=217
Γ ₇₀	$K_S^0 K^- \pi^+ 2(\pi^+ \pi^-)$	< 1.22	%	CL=90%	DESIG=204
Γ ₇₁	$K_S^0 K^- \pi^+ 2\pi^0$	< 2.65	%	CL=90%	DESIG=205
Γ ₇₂	$K_S^0 K^- K^+ K^- \pi^+$	< 4.9	$\times 10^{-3}$	CL=90%	DESIG=218
Γ ₇₃	$K_S^0 K^- K^+ K^- \pi^+ \pi^0$	< 3.0	%	CL=90%	DESIG=219
Γ ₇₄	$K_S^0 K^- K^+ K^- \pi^+ \eta$	< 2.2	%	CL=90%	DESIG=220
Γ ₇₅	$K_S^0 K^- \pi^+ + \text{c.c.}$	< 9.7	$\times 10^{-3}$	CL=90%	DESIG=60
Γ ₇₆	$\rho \bar{\rho}$				DESIG=233
Γ ₇₇	$\rho \bar{\rho} \pi^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=54
Γ ₇₈	$\rho \bar{\rho} \pi^+ \pi^-$	< 5.8	$\times 10^{-4}$	CL=90%	DESIG=36
Γ ₇₉	$\Lambda \bar{\Lambda}$	< 1.2	$\times 10^{-4}$	CL=90%	DESIG=42
Γ ₈₀	$\rho \bar{\rho} \pi^+ \pi^- \pi^0$	< 1.85	$\times 10^{-3}$	CL=90%	DESIG=37
Γ ₈₁	$\omega \rho \bar{\rho}$	< 2.9	$\times 10^{-4}$	CL=90%	DESIG=39
Γ ₈₂	$\Lambda \bar{\Lambda} \pi^0$	< 7	$\times 10^{-5}$	CL=90%	DESIG=72
Γ ₈₃	$\rho \bar{\rho} 2(\pi^+ \pi^-)$	< 2.6	$\times 10^{-3}$	CL=90%	DESIG=61
Γ ₈₄	$\eta \rho \bar{\rho}$	< 5.4	$\times 10^{-4}$	CL=90%	DESIG=38
Γ ₈₅	$\eta \rho \bar{\rho} \pi^+ \pi^-$	< 3.3	$\times 10^{-3}$	CL=90%	DESIG=223
Γ ₈₆	$\rho^0 \rho \bar{\rho}$	< 1.7	$\times 10^{-3}$	CL=90%	DESIG=66
Γ ₈₇	$\rho \bar{\rho} K^+ K^-$	< 3.2	$\times 10^{-4}$	CL=90%	DESIG=40
Γ ₈₈	$\eta \rho \bar{\rho} K^+ K^-$	< 6.9	$\times 10^{-3}$	CL=90%	DESIG=224
Γ ₈₉	$\pi^0 \rho \bar{\rho} K^+ K^-$	< 1.2	$\times 10^{-3}$	CL=90%	DESIG=225
Γ ₉₀	$\phi \rho \bar{\rho}$	< 1.3	$\times 10^{-4}$	CL=90%	DESIG=41
Γ ₉₁	$\Lambda \bar{\Lambda} \pi^+ \pi^-$	< 2.5	$\times 10^{-4}$	CL=90%	DESIG=43
Γ ₉₂	$\Lambda \bar{\rho} K^+$	< 2.8	$\times 10^{-4}$	CL=90%	DESIG=44
Γ ₉₃	$\Lambda \bar{\rho} K^+ \pi^+ \pi^-$	< 6.3	$\times 10^{-4}$	CL=90%	DESIG=45
Γ ₉₄	$\Lambda \bar{\Lambda} \eta$	< 1.9	$\times 10^{-4}$	CL=90%	DESIG=226
Γ ₉₅	$\Sigma^+ \bar{\Sigma}^-$	< 1.0	$\times 10^{-4}$	CL=90%	DESIG=227
Γ ₉₆	$\Sigma^0 \bar{\Sigma}^0$	< 4	$\times 10^{-5}$	CL=90%	DESIG=228
Γ ₉₇	$\Xi^+ \bar{\Xi}^-$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=229
Γ ₉₈	$\Xi^0 \bar{\Xi}^0$	< 1.4	$\times 10^{-4}$	CL=90%	DESIG=230
Γ ₉₉	$\Xi^- \bar{\Xi}^+$	(1.4 ± 0.4)	$\times 10^{-4}$		DESIG=236

Radiative decays

Γ ₁₀₀	$\gamma \chi_{c2}$	< 6.4	$\times 10^{-4}$	CL=90%	DESIG=51
Γ ₁₀₁	$\gamma \chi_{c1}$	(2.49 ± 0.23)	$\times 10^{-3}$		DESIG=50
Γ ₁₀₂	$\gamma \chi_{c0}$	(6.9 ± 0.6)	$\times 10^{-3}$		DESIG=49
Γ ₁₀₃	$\gamma \eta_c$	< 7	$\times 10^{-4}$	CL=90%	DESIG=231
Γ ₁₀₄	$\gamma \eta_c(2S)$	< 9	$\times 10^{-4}$	CL=90%	DESIG=232
Γ ₁₀₅	$\gamma \eta'$	< 1.8	$\times 10^{-4}$	CL=90%	DESIG=213
Γ ₁₀₆	$\gamma \eta$	< 1.5	$\times 10^{-4}$	CL=90%	DESIG=212
Γ ₁₀₇	$\gamma \pi^0$	< 2	$\times 10^{-4}$	CL=90%	DESIG=211

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 3 branching ratios uses 23 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 20.1$ for 19 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	99		
x_9	0	0	
Γ	0	0	-44
	x_2	x_3	x_9

	Mode	Rate (MeV)	Scale factor	
Γ ₂	$D^0 \bar{D}^0$	14.0 ± 1.4	1.8	DESIG=5
Γ ₃	$D^+ D^-$	11.2 ± 1.1	1.7	DESIG=6
Γ ₉	$e^+ e^-$	(2.62 ± 0.18) $\times 10^{-4}$	1.4	DESIG=1

$\psi(3770)$ PARTIAL WIDTHS

NODE=M053225

 $\Gamma(e^+e^-)$ Γ_9

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.262±0.018 OUR FIT				Error includes scale factor of 1.4.
0.256±0.016 OUR AVERAGE				Error includes scale factor of 1.2.
0.154 ^{+0.079+0.021} _{-0.058-0.027}		1,2 ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
0.22 ±0.05		3,4 ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
0.277±0.011±0.013		4 ABLIKIM	07E BES2	$e^+e^- \rightarrow$ hadrons
0.203±0.003 ^{+0.041} _{-0.027}	1.4M	4,5 BESSON	06 CLEO	$e^+e^- \rightarrow$ hadrons
0.276±0.050		4 SCHINDLER	80 MRK2	e^+e^-
0.18 ±0.06		4 BACINO	78 DLCO	e^+e^-
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.196±0.018		6 SHAMOV	17 RVUE	$e^+e^- \rightarrow D\bar{D}$, hadrons
0.414 ^{+0.072+0.093} _{-0.080-0.028}		2,7 ANASHIN	12A KEDR	$e^+e^- \rightarrow D\bar{D}$
0.37 ±0.09		8 RAPIDIS	77 LGW	e^+e^-

NODE=M053W1
NODE=M053W1

OCCUR=2

OCCUR=2

NODE=M053W1;LINKAGE=A1
NODE=M053W1;LINKAGE=AN
NODE=M053W1;LINKAGE=AB

NODE=M053W1;LINKAGE=NI

NODE=M053W1;LINKAGE=BE

NODE=M053W1;LINKAGE=B

NODE=M053W1;LINKAGE=A2

NODE=M053W1;LINKAGE=R

¹ Solution I of the two solutions.² Taking into account interference between the resonant and non-resonant $D\bar{D}$ production.³ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = 0^\circ$.⁴ Interference between the resonant and non-resonant $D\bar{D}$ production not taken into account.⁵ BESSON 06 (as corrected in BESSON 10) measure $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow$ hadrons) = $6.36 \pm 0.08^{+0.41}_{-0.30}$ nb at $\sqrt{s} = 3773 \pm 1$ MeV, and obtain Γ_{ee} from the Born-level cross section calculated using $\psi(3770)$ mass and width from our 2004 edition, PDG 04.⁶ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.⁷ Solution II of the two solutions.⁸ See also $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ below. $\psi(3770) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M053235

 $\Gamma(\Xi^- \Xi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{99}\Gamma_9/\Gamma$

VALUE (10^{-2} eV)	DOCUMENT ID	TECN	COMMENT
3.55±0.92	1 ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(3770)$

NODE=M053P01
NODE=M053P01¹ From a fit to $e^+e^- \rightarrow \Xi^- \Xi^+$ cross sections. Signal significance is 4.5σ .

NODE=M053P01;LINKAGE=A

 $\psi(3770)$ BRANCHING RATIOS

NODE=M053230

 $\Gamma(D\bar{D})/\Gamma_{\text{total}}$ $\Gamma_1/\Gamma = (\Gamma_2+\Gamma_3)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.93^{+0.08}_{-0.09} OUR FIT				Error includes scale factor of 2.0.
0.93^{+0.08}_{-0.09} OUR AVERAGE				Error includes scale factor of 2.1.
0.849±0.056±0.018		1 ABLIKIM	08B BES2	$e^+e^- \rightarrow$ non- $D\bar{D}$
1.033±0.014 ^{+0.048} _{-0.066}	1.427M	2 BESSON	06 CLEO	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
0.836±0.049		3 SHAMOV	17 RVUE	$e^+e^- \rightarrow D\bar{D}$, hadrons
0.866±0.050±0.036		4,5 ABLIKIM	07K BES2	$e^+e^- \rightarrow$ non- $D\bar{D}$
0.836±0.073±0.042		5 ABLIKIM	06L BES2	$e^+e^- \rightarrow D\bar{D}$
0.855±0.017±0.058		5,6 ABLIKIM	06N BES2	$e^+e^- \rightarrow D\bar{D}$

NODE=M053R1
NODE=M053R1¹ Neglecting interference.² Obtained by comparing a measurement of the total cross section (corrected in BESSON 10) with that of $D\bar{D}$ reported by CLEO in DOBBS 07.³ From the joint analysis of the data on the $D\bar{D}$ and inclusive hadronic cross sections in the $\psi(3770)$ region from BaBar, Belle, BES-II, CLEO and KEDR.⁴ Using $\sigma^{obs} = 7.07 \pm 0.58$ nb and neglecting interference.⁵ Not independent of ABLIKIM 08B.⁶ From a measurement of $\sigma(e^+e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.NODE=M053R1;LINKAGE=AI
NODE=M053R1;LINKAGE=BE

NODE=M053R1;LINKAGE=A

NODE=M053R1;LINKAGE=AL
NODE=M053R1;LINKAGE=SU
NODE=M053R1;LINKAGE=AB

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

 Γ_2/Γ NODE=M053R46
NODE=M053R46**0.52^{+0.04}_{-0.05} OUR FIT** Error includes scale factor of 2.0.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.467±0.047±0.023	ABLIKIM	06L	BES2	$e^+e^- \rightarrow D^0\bar{D}^0$
0.499±0.013±0.038	¹ ABLIKIM	06N	BES2	$e^+e^- \rightarrow D^0\bar{D}^0$

¹ From a measurement of $\sigma(e^+e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

NODE=M053R46;LINKAGE=AB

 $\Gamma(D^+D^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

 Γ_3/Γ NODE=M053R47
NODE=M053R47**0.41 ± 0.04 OUR FIT** Error includes scale factor of 2.0.

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.369±0.037±0.028	ABLIKIM	06L	BES2	$e^+e^- \rightarrow D^+D^-$
0.357±0.011±0.034	¹ ABLIKIM	06N	BES2	$e^+e^- \rightarrow D^+D^-$

¹ From a measurement of $\sigma(e^+e^- \rightarrow D\bar{D})$ at $\sqrt{s} = 3773$ MeV, using the $\psi(3770)$ resonance parameters measured by ABLIKIM 06L.

NODE=M053R47;LINKAGE=AB

 $\Gamma(D^0\bar{D}^0)/\Gamma(D^+D^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

 Γ_2/Γ_3 NODE=M053R5
NODE=M053R5**1.253±0.016 OUR FIT****1.253±0.016 OUR AVERAGE**

1.252±0.009±0.013	5.3M	BONVICINI	14	CLEO	$e^+e^- \rightarrow D\bar{D}$
1.39 ± 0.31 ± 0.12		PAKHLOVA	08	BELL	10.6 $e^+e^- \rightarrow D\bar{D}\gamma$
1.78 ± 0.33 ± 0.24		AUBERT	07BE	BABR	$e^+e^- \rightarrow D\bar{D}\gamma$
1.27 ± 0.12 ± 0.08		ABLIKIM	06L	BES2	$e^+e^- \rightarrow D\bar{D}$
2.43 ± 1.50 ± 0.43	34	¹ CHISTOV	04	BELL	$B^+ \rightarrow \psi(3770)K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.258±0.016±0.014		² DOBBS	07	CLEO	$e^+e^- \rightarrow D\bar{D}$
-------------------	--	--------------------	----	------	-------------------------------

¹ See ADLER 88C for older measurements of this quantity.² Superseded by BONVICINI 14.NODE=M053R5;LINKAGE=CH
NODE=M053R5;LINKAGE=DO $\Gamma(J/\psi X)/\Gamma_{\text{total}}$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

 Γ_4/Γ NODE=M053P00
NODE=M053P00**0.5±0.2±0.1** ¹ ABLIKIM 21Z BES3 $e^+e^- \rightarrow \ell^+\ell^-X$ ¹ From a fit to the $e^+e^- \rightarrow J/\psi X$ cross section between 3.645 and 3.891 GeV, with $\psi(2S)$ and $\psi(3770)$ masses, total widths and leptonic widths fixed to the values from the PDG 20. An alternative fit with an improved χ^2 , corresponding to a significance of 5.3 σ , uses an additional resonance with a mass of $3766.2 \pm 3.8 \pm 0.4$ MeV/ c^2 , a total width of $22.2 \pm 5.9 \pm 1.4$ MeV, and $\Gamma(e^+e^-)B(J/\psi X) = 79.4 \pm 85.5 \pm 11.7$ eV, possibly compatible with the results of ABLIKIM 08H.

NODE=M053P00;LINKAGE=A

 $\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

 Γ_5/Γ NODE=M053R4
NODE=M053R4**1.93±0.28 OUR AVERAGE**

1.89±0.20±0.20	231 ± 33	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$
3.4 ± 1.4 ± 0.9	17.8 ± 4.8	BAI	05	BES2	$e^+e^- \rightarrow \psi(3770)$

 $\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

 Γ_6/Γ NODE=M053R7
NODE=M053R7**0.080±0.025±0.016** 39 ± 14 ADAM 06 CLEO $e^+e^- \rightarrow \psi(3770)$ $\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

 Γ_7/Γ NODE=M053R8
NODE=M053R8**8.7±1.2 OUR AVERAGE**[(90 ± 40) × 10⁻⁵ OUR 2023 AVERAGE]

8.7±1.0±0.8	232 ± 23	¹ ABLIKIM	23v	BES3	$e^+e^- \rightarrow \psi(3770)$
8.7±3.3±2.2	22 ± 10	ADAM	06	CLEO	$e^+e^- \rightarrow \psi(3770)$

¹ Incoherent fit. Alternate fits that include interference with background yield results between $(11.2 \pm 5.8 \pm 1.1) \times 10^{-4}$ and $(11.6 \pm 6.0 \pm 1.1) \times 10^{-4}$.

NODE=M053R8;LINKAGE=A

 $\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

 Γ_8/Γ NODE=M053R9
NODE=M053R9**<28** 90 <10 ADAM 06 CLEO $e^+e^- \rightarrow \psi(3770)$

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_9/Γ VALUE (units 10^{-5})

DOCUMENT ID

TECN

COMMENT

NODE=M053R2
NODE=M053R2**0.96±0.07 OUR FIT** Error includes scale factor of 1.3.**1.3 ±0.2**

RAPIDIS

77

LGW

 e^+e^-

————— DECAYS TO LIGHT HADRONS —————

NODE=M053250

 $\Gamma(b_1(1235)\pi)/\Gamma_{\text{total}}$ Γ_{10}/Γ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R82
NODE=M053R82**<1.4**

90

1

ADAMS

06

CLEO

 $e^+e^- \rightarrow \psi(3770)$ ¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R82;LINKAGE=AD

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R83
NODE=M053R83**<2.3 × 10⁻⁵ (CL = 90%)** [$<7 \times 10^{-4}$ (CL = 90%) OUR 2023 BEST LIMIT]**<2.3 × 10⁻⁵**

90

1

ABLIKIM

23BC

BES3

 $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $<7 \times 10^{-4}$

90

2

ADAMS

06

CLEO

 $e^+e^- \rightarrow \psi(3770)$ ¹ ABLIKIM 23BC fit to $e^+e^- \rightarrow \phi\eta'$ cross sections between 3.508 and 4.951 GeV considering interference between continuum and $\psi(3770)$ amplitudes.

NODE=M053R83;LINKAGE=A

² ADAMS 06 compare cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R83;LINKAGE=AD

 $\Gamma(\omega\eta')/\Gamma_{\text{total}}$ Γ_{12}/Γ VALUE (units 10^{-4})

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R84
NODE=M053R84**<4**

90

1

ADAMS

06

CLEO

 $e^+e^- \rightarrow \psi(3770)$ ¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R84;LINKAGE=AD

 $\Gamma(\rho^0\eta')/\Gamma_{\text{total}}$ Γ_{13}/Γ VALUE (units 10^{-4})

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R85
NODE=M053R85**<6**

90

1

ADAMS

06

CLEO

 $e^+e^- \rightarrow \psi(3770)$ ¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R85;LINKAGE=AD

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{14}/Γ VALUE (units 10^{-4})

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R6
NODE=M053R6**3.1±0.6±0.3**

1

ADAMS

06

CLEO

 $3.773 e^+e^- \rightarrow \phi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19

90

2

ABLIKIM

07B

BES2

 $e^+e^- \rightarrow \psi(3770)$ ¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R6;LINKAGE=AD

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R6;LINKAGE=AK

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{15}/Γ VALUE (units 10^{-5})

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R86
NODE=M053R86**<1.4**

90

1

ADAMS

06

CLEO

 $e^+e^- \rightarrow \psi(3770)$ ¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R86;LINKAGE=AD

 $\Gamma(\rho^0\eta)/\Gamma_{\text{total}}$ Γ_{16}/Γ VALUE (units 10^{-4})

CL%

DOCUMENT ID

TECN

COMMENT

NODE=M053R87
NODE=M053R87**<5**

90

1

ADAMS

06

CLEO

 $e^+e^- \rightarrow \psi(3770)$ ¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R87;LINKAGE=AD

$\Gamma(\phi\pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 3	90	¹ ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R11
 NODE=M053R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R11;LINKAGE=AD

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R11;LINKAGE=AK

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	¹ ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R88
 NODE=M053R88

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R88;LINKAGE=AD

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	^{1,2} ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R89
 NODE=M053R89

¹ Data suggest possible destructive interference with continuum.

NODE=M053R89;LINKAGE=AD

² Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R89;LINKAGE=AS

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	^{1,2} ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R90
 NODE=M053R90

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R90;LINKAGE=AD

² Data suggest possible destructive interference with continuum.

NODE=M053R90;LINKAGE=AS

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$\sim 10^{-5}$	¹ DRUZHININ	15 RVUE	$e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\sim 10^{-5}$

¹ DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+e^- \rightarrow K^+K^-$ and $e^+e^- \rightarrow K_S^0 K_L^0$.

NODE=M053R00
 NODE=M053R00

NODE=M053R00;LINKAGE=A

 $\Gamma(K^*(892)^+K^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	¹ ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R91
 NODE=M053R91

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R91;LINKAGE=AS

 $\Gamma(K^*(892)^0\bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ ADAMS	06 CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R92
 NODE=M053R92

¹ Comparing cross sections at $\sqrt{s} = 3.773$ GeV and $\sqrt{s} = 3.671$ GeV, neglecting interference, and using $\sigma(\psi(3770) \rightarrow D\bar{D}) = 6.39 \pm 0.20$ nb.

NODE=M053R92;LINKAGE=AD

 $\Gamma(K_S^0 K_L^0)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	¹ CRONIN-HEN..06	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R3
 NODE=M053R3

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	² ABLIKIM	04F BES	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	---------	---------------------------------

¹ Using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = (6.38 \pm 0.08_{-0.30}^{+0.41})$ nb from BESSON 06 and $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6895 \pm 0.0014$.

NODE=M053R3;LINKAGE=CR

² Using $B(K_S^0 \rightarrow \pi^+\pi^-) = 0.6860 \pm 0.0027$.

NODE=M053R3;LINKAGE=AB

$\Gamma(2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<11.2	90	¹ HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R21
 NODE=M053R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48	90	² ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R21;LINKAGE=HU
 NODE=M053R21;LINKAGE=AK

 $\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10.6	90	¹ HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R22
 NODE=M053R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

<62	90	² ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R22;LINKAGE=HU
 NODE=M053R22;LINKAGE=AK

 $\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<58.5	90	305	ABLIKIM 08N	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R72
 NODE=M053R72

 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.0	90	¹ HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R24
 NODE=M053R24

• • • We do not use the following data for averages, fits, limits, etc. • • •

<55	90	² ABLIKIM 07I	BES2	$3.77 e^+e^-$
-----	----	--------------------------	------	---------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R24;LINKAGE=HU
 NODE=M053R24;LINKAGE=AK

 $\Gamma(3(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<91	90	¹ ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R07
 NODE=M053R07

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R07;LINKAGE=AK

 $\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<137	90	¹ ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R10
 NODE=M053R10

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R10;LINKAGE=AK

 $\Gamma(3(\pi^+\pi^-)2\pi^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<117.4	90	59	ABLIKIM 08N	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R74
 NODE=M053R74

 $\Gamma(\eta\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.24	90	¹ HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R23
 NODE=M053R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.3	90	² ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$
------	----	--------------------------	------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R23;LINKAGE=HU
 NODE=M053R23;LINKAGE=AK

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<8.9	90	218	ABLIKIM 08N	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R70
 NODE=M053R70

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R53
NODE=M053R53

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R53;LINKAGE=AK

 $\Gamma(\eta 3\pi)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<13.4	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R25
NODE=M053R25

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R25;LINKAGE=HU

 $\Gamma(\eta 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<243	90	¹ ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R08
NODE=M053R08

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R08;LINKAGE=AK

 $\Gamma(\eta \rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<1.45	90	¹ ABLIKIM	10D BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R77
NODE=M053R77

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R77;LINKAGE=AK

 $\Gamma(\eta' 3\pi)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<24.4	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R26
NODE=M053R26

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R26;LINKAGE=HU

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 9.0	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R27
NODE=M053R27

• • • We do not use the following data for averages, fits, limits, etc. • • •

<48	90	² ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	----------------------------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R27;LINKAGE=HU

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R27;LINKAGE=AK

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	¹ HUANG	06A CLEO	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R28
NODE=M053R28

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	² ABLIKIM	07B BES2	$e^+ e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	----------------------------------

¹ Using $\sigma_{tot}(e^+ e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R28;LINKAGE=HU

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R28;LINKAGE=AK

 $\Gamma(K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<4.2	90	14	ABLIKIM	08N BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R71
NODE=M053R71 $\Gamma(4(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<16.7	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R50
NODE=M053R50

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R50;LINKAGE=AK

 $\Gamma(4(\pi^+ \pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<30.6	90	¹ ABLIKIM	07F BES2	$e^+ e^- \rightarrow \psi(3770)$

NODE=M053R52
NODE=M053R52

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R52;LINKAGE=AK

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4.5	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R29
 NODE=M053R29

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R29;LINKAGE=HU

 $\Gamma(K^+K^-\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 23.6	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R30
 NODE=M053R30

• • • We do not use the following data for averages, fits, limits, etc. • • •

<111	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R30;LINKAGE=HU
 NODE=M053R30;LINKAGE=AK

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(K^+K^-\rho^0\pi^0)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	¹ ABLIKIM	07I BES2	$3.77 e^+e^-$

NODE=M053R58
 NODE=M053R58

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R58;LINKAGE=AK

 $\Gamma(K^+K^-\rho^+\pi^-)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<146	90	¹ ABLIKIM	07I BES2	$3.77 e^+e^-$

NODE=M053R59
 NODE=M053R59

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R59;LINKAGE=AK

 $\Gamma(\omega K^+K^-)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.4	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R32
 NODE=M053R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

<66	90	² ABLIKIM	07I BES2	$3.77 e^+e^-$
-----	----	----------------------	----------	---------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R32;LINKAGE=HU
 NODE=M053R32;LINKAGE=AK

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\phi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<38	90	¹ ABLIKIM	07I BES2	$3.77 e^+e^-$

NODE=M053R60
 NODE=M053R60

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R60;LINKAGE=AK

 $\Gamma(K^{*0}K^-\pi^+\pi^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<162	90	¹ ABLIKIM	07I BES2	$3.77 e^+e^-$

NODE=M053R61
 NODE=M053R61

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R61;LINKAGE=AK

 $\Gamma(K^{*+}K^-\pi^+\pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<323	90	¹ ABLIKIM	07I BES2	$3.77 e^+e^-$

NODE=M053R62
 NODE=M053R62

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R62;LINKAGE=AK

 $\Gamma(K^+K^-\pi^+\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<26.7	90	24	ABLIKIM	08N BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R73
 NODE=M053R73

 $\Gamma(K^+K^-2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<10.3	90	¹ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R57
 NODE=M053R57

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R57;LINKAGE=AK

$\Gamma(K^+K^-2(\pi^+\pi^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<36.0	90	¹ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R51
 NODE=M053R51

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R51;LINKAGE=AK

 $\Gamma(\eta K^+K^-)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.1	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R31
 NODE=M053R31

••• We do not use the following data for averages, fits, limits, etc. •••

<31	90	² ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R31;LINKAGE=HU

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R31;LINKAGE=AK

 $\Gamma(\eta K^+K^-\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
<1.24	90	¹ ABLIKIM	10D BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R78
 NODE=M053R78

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R78;LINKAGE=AK

 $\Gamma(\rho^0 K^+K^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<5.0	90	¹ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R54
 NODE=M053R54

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R54;LINKAGE=AK

 $\Gamma(2(K^+K^-))/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 6.0	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R33
 NODE=M053R33

••• We do not use the following data for averages, fits, limits, etc. •••

<17	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R33;LINKAGE=HU

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R33;LINKAGE=AK

 $\Gamma(\phi K^+K^-)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 7.5	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R34
 NODE=M053R34

••• We do not use the following data for averages, fits, limits, etc. •••

<24	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R34;LINKAGE=HU

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R34;LINKAGE=AK

 $\Gamma(2(K^+K^-)\pi^0)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.9	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R35
 NODE=M053R35

••• We do not use the following data for averages, fits, limits, etc. •••

<46	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R35;LINKAGE=HU

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R35;LINKAGE=AK

 $\Gamma(2(K^+K^-)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	¹ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R48
 NODE=M053R48

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R48;LINKAGE=AK

$\Gamma(K_S^0 K^- \pi^+)/\Gamma_{\text{total}}$			Γ_{62}/Γ				
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
<3.2	90	18	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R64 NODE=M053R64
$\Gamma(K_S^0 K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$			Γ_{63}/Γ				
<13.3	90	40	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R65 NODE=M053R65
$\Gamma(K_S^0 K^- \rho^+)/\Gamma_{\text{total}}$			Γ_{64}/Γ				
<6.6	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R15 NODE=M053R15
$\Gamma(K_S^0 K^- 2\pi^+ \pi^-)/\Gamma_{\text{total}}$			Γ_{65}/Γ				
<8.7	90	39	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R66 NODE=M053R66
$\Gamma(K_S^0 K^- \pi^+ \rho^0)/\Gamma_{\text{total}}$			Γ_{66}/Γ				
<1.6	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R16 NODE=M053R16
$\Gamma(K_S^0 K^- \pi^+ \eta)/\Gamma_{\text{total}}$			Γ_{67}/Γ				
<1.3	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R17 NODE=M053R17
$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$			Γ_{68}/Γ				
<41.8	90	23	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R67 NODE=M053R67
$\Gamma(K_S^0 K^- 2\pi^+ \pi^- \eta)/\Gamma_{\text{total}}$			Γ_{69}/Γ				
<4.8	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R18 NODE=M053R18
$\Gamma(K_S^0 K^- \pi^+ 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$			Γ_{70}/Γ				
<12.2	90	4	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R68 NODE=M053R68
$\Gamma(K_S^0 K^- \pi^+ 2\pi^0)/\Gamma_{\text{total}}$			Γ_{71}/Γ				
<26.5	90	17	ABLIKIM	08M	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R69 NODE=M053R69
$\Gamma(K_S^0 K^- K^+ K^- \pi^+)/\Gamma_{\text{total}}$			Γ_{72}/Γ				
<4.9	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R19 NODE=M053R19
$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$			Γ_{73}/Γ				
<3.0	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R75 NODE=M053R75
$\Gamma(K_S^0 K^- K^+ K^- \pi^+ \eta)/\Gamma_{\text{total}}$			Γ_{74}/Γ				
<2.2	90		ABLIKIM	09C	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R76 NODE=M053R76
$\Gamma(K^{*0} K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$			Γ_{75}/Γ				
<9.7	90		¹ ABLIKIM	07F	BES2	$e^+ e^- \rightarrow \psi(3770)$	NODE=M053R55 NODE=M053R55

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma_{obs}(e^+ e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38 \text{ nb}$.

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	EVT5	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		¹ AAIJ	17AD LHCB	$p\bar{p} \rightarrow B^+ X \rightarrow p\bar{p}K^+ X$
----------	--	-------------------	-----------	--

$7.1^{+8.6}_{-2.9}$	684	² ABLIKIM	14L BES3	$e^+e^- \rightarrow \psi(3770)$
---------------------	-----	----------------------	----------	---------------------------------

310 ± 30	684	³ ABLIKIM	14L BES3	$e^+e^- \rightarrow \psi(3770)$
--------------	-----	----------------------	----------	---------------------------------

¹ AAIJ 17AD reports $B(B^+ \rightarrow \psi(3770)K^+ \rightarrow p\bar{p}K^+)/B(B^+ \rightarrow J/\psi K^+ \rightarrow p\bar{p}K^+) < 0.09$ (0.10) at 90% (95%) CL.

² Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

³ Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

Γ_{76}/Γ
 NODE=M053R98
 NODE=M053R98

OCCUR=2

NODE=M053R98;LINKAGE=C

NODE=M053R98;LINKAGE=A

NODE=M053R98;LINKAGE=B

 $\Gamma(p\bar{p}\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 0.4 90 ^{1,2} ABLIKIM 14O BES3 $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$59^{+3}_{-2} \pm 5$		^{1,3} ABLIKIM	14O BES3	$e^+e^- \rightarrow \psi(3770)$
----------------------	--	------------------------	----------	---------------------------------

<12	90	⁴ ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Calculated by the authors using $\sigma(e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}) = 6.36 \pm 0.08^{+0.41}_{-0.30}$ nb from BESSON 10.

² Solution I of two equivalent solutions in a fit with a resonance interfering with continuum.

³ Solution II of two equivalent solutions in a fit with a resonance interfering with continuum.

⁴ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

Γ_{77}/Γ
 NODE=M053R09
 NODE=M053R09

OCCUR=2

NODE=M053R09;LINKAGE=A

NODE=M053R09;LINKAGE=B

NODE=M053R09;LINKAGE=C

NODE=M053R09;LINKAGE=AK

 $\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 5.8 90 ¹ HUANG 06A CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<16	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

Γ_{78}/Γ
 NODE=M053R36
 NODE=M053R36

NODE=M053R36;LINKAGE=HU

NODE=M053R36;LINKAGE=AK

 $\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 1.2×10^{-4} 90 ¹ HUANG 06A CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.8×10^{-4}	90	² ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(3770)$
------------------------	----	----------------------	-----------	---------------------------------

< 4×10^{-4}	90	³ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$
----------------------	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV. At a 90% CL the lower bound is $> 2.4 \times 10^{-6}$.

³ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

Γ_{79}/Γ
 NODE=M053R42
 NODE=M053R42

NODE=M053R42;LINKAGE=HU

NODE=M053R42;LINKAGE=A

NODE=M053R42;LINKAGE=AK

 $\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 18.5 90 ¹ HUANG 06A CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<73	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

Γ_{80}/Γ
 NODE=M053R37
 NODE=M053R37

NODE=M053R37;LINKAGE=HU

NODE=M053R37;LINKAGE=AK

 $\Gamma(\omega p\bar{p})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 2.9 90 ¹ HUANG 06A CLEO $e^+e^- \rightarrow \psi(3770)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<30	90	² ABLIKIM	07I BES2	$3.77 e^+e^-$
-----	----	----------------------	----------	---------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Using $\sigma^{obs} = 7.15 \pm 0.27 \pm 0.27$ nb and neglecting interference.

Γ_{81}/Γ
 NODE=M053R39
 NODE=M053R39

NODE=M053R39;LINKAGE=HU

NODE=M053R39;LINKAGE=AB

$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{82}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 0.7	90	¹ ABLIKIM 13Q	BES3	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R63
 NODE=M053R63

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12	90	² ABLIKIM 07I	BES2	$3.77 e^+e^-$
-----	----	--------------------------	------	---------------

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R63;LINKAGE=A
 NODE=M053R63;LINKAGE=AK

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\rho\bar{\rho}2(\pi^+\pi^-))/\Gamma_{\text{total}}$ Γ_{83}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2.6	90	¹ ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R49
 NODE=M053R49

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R49;LINKAGE=AK

 $\Gamma(\eta\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{84}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.4	90	¹ HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R38
 NODE=M053R38

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11	90	² ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R38;LINKAGE=HU
 NODE=M053R38;LINKAGE=AK

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{85}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	¹ ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R79
 NODE=M053R79

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R79;LINKAGE=AK

 $\Gamma(\rho^0\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{86}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ ABLIKIM 07F	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R56
 NODE=M053R56

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R56;LINKAGE=AK

 $\Gamma(\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{87}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	¹ HUANG 06A	CLEO	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R40
 NODE=M053R40

• • • We do not use the following data for averages, fits, limits, etc. • • •

<11	90	² ABLIKIM 07B	BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	--------------------------	------	---------------------------------

¹ Using $\sigma_{tot}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R40;LINKAGE=HU
 NODE=M053R40;LINKAGE=AK

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

 $\Gamma(\eta\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{88}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<6.9	90	¹ ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R80
 NODE=M053R80

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R80;LINKAGE=AK

 $\Gamma(\pi^0\rho\bar{\rho}K^+K^-)/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	90	¹ ABLIKIM 10D	BES2	$e^+e^- \rightarrow \psi(3770)$

NODE=M053R81
 NODE=M053R81

¹ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{obs}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R81;LINKAGE=AK

$\Gamma(\phi\rho\bar{p})/\Gamma_{\text{total}}$ Γ_{90}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.3	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
------	----	--------------------	----------	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<9	90	² ABLIKIM	07B BES2	$e^+e^- \rightarrow \psi(3770)$
----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R41
NODE=M053R41

NODE=M053R41;LINKAGE=HU
NODE=M053R41;LINKAGE=AK

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{91}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 2.5	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
-------	----	--------------------	----------	---------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 4.7	90	² ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
-------	----	----------------------	----------	---------------------------------

<39	90	³ ABLIKIM	07F BES2	$e^+e^- \rightarrow \psi(3770)$
-----	----	----------------------	----------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

² Assuming that interference effects between resonance and continuum can be neglected.

³ Assuming that interference effects between resonance and continuum can be neglected and using $\sigma^{\text{obs}}(e^+e^- \rightarrow \psi(3770)) = 7.15 \pm 0.38$ nb.

NODE=M053R43
NODE=M053R43

NODE=M053R43;LINKAGE=HU
NODE=M053R43;LINKAGE=A
NODE=M053R43;LINKAGE=AK

 $\Gamma(\Lambda\bar{p}K^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<2.8	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
------	----	--------------------	----------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R44
NODE=M053R44

NODE=M053R44;LINKAGE=HU

 $\Gamma(\Lambda\bar{p}K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<6.3	90	¹ HUANG	06A CLEO	$e^+e^- \rightarrow \psi(3770)$
------	----	--------------------	----------	---------------------------------

¹ Using $\sigma_{\text{tot}}(e^+e^- \rightarrow \psi(3770)) = 7.9 \pm 0.6$ nb at the resonance.

NODE=M053R45
NODE=M053R45

NODE=M053R45;LINKAGE=HU

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.9	90	¹ ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R93
NODE=M053R93

NODE=M053R93;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.0	90	¹ ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R94
NODE=M053R94

NODE=M053R94;LINKAGE=A

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<0.4	90	¹ ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R95
NODE=M053R95

NODE=M053R95;LINKAGE=A

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.5	90	¹ ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R96
NODE=M053R96

NODE=M053R96;LINKAGE=A

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.4	90	¹ ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(3770)$
------	----	----------------------	----------	---------------------------------

¹ Assuming that interference effects between resonance and continuum can be neglected.

NODE=M053R97
NODE=M053R97

NODE=M053R97;LINKAGE=A

———— RADIATIVE DECAYS ————

$\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$					Γ_{100}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<0.64	90	¹ ABLIKIM	15J BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.0	90	² BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
<0.9	90	³ COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

¹ This limit is equivalent to $(0.25 \pm 0.21 \pm 0.18) \times 10^{-3}$ branching fraction value.

² Uses $B(\psi(2S) \rightarrow \gamma\chi_{c2}) = 9.22 \pm 0.11 \pm 0.46\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

³ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+e^- \rightarrow \psi(3770))$.

NODE=M053240

NODE=M053R03
NODE=M053R03NODE=M053R03;LINKAGE=A
NODE=M053R03;LINKAGE=BR

NODE=M053R03;LINKAGE=CO

$\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$					Γ_{101}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.49±0.23 OUR AVERAGE					
2.0 ±0.8 ±0.1	202	¹ ABLIKIM	16B BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
2.48±0.15±0.23	0.6k	ABLIKIM	15J BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	
2.4 ±0.8 ±0.2		² ABLIKIM	14H BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow K_S^0 K^\pm \pi^\mp$	
2.9 ±0.5 ±0.4		³ BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}, \gamma\gamma J/\psi$	OCCUR=2
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
3.9 ±1.4 ±0.6	54	⁴ BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
2.8 ±0.5 ±0.4	53	⁵ COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

NODE=M053R02
NODE=M053R02

OCCUR=2

NODE=M053R02;LINKAGE=A

NODE=M053R02;LINKAGE=AB

NODE=M053R02;LINKAGE=BI
NODE=M053R02;LINKAGE=BR

NODE=M053R02;LINKAGE=CO

¹ ABLIKIM 16B reports $(1.94 \pm 0.42 \pm 0.64) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.55 \pm 0.31) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = (9.75 \pm 0.27) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c1})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)] = (8.51 \pm 2.39 \pm 1.42) \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp) = 0.00349 \pm 0.00031$. Our first error is their experiment's error and our second error is the systematic error from using our best value. We have calculated the best value of $B(\chi_{c1}(1P) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/2 of $B(\chi_{c1}(1P) \rightarrow \bar{K}^0 K^+ \pi^- + \text{c.c.}) = (7.0 \pm 0.6) \times 10^{-3}$.

³ Averages the two measurements from COAN 06A and BRIERE 06.

⁴ Uses $B(\psi(2S) \rightarrow \gamma\chi_{c1}) = 9.07 \pm 0.11 \pm 0.54\%$ from ATHAR 04, $\psi(2S)$ mass and width from PDG 04, and $\Gamma_{ee}(\psi(2S)) = 2.54 \pm 0.03 \pm 0.11$ keV from ADAM 06.

⁵ Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+e^- \rightarrow \psi(3770))$.

$\Gamma(\gamma\chi_{c1})/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{101}/Γ_5
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
1.49±0.31±0.26	53 ± 10	¹ COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

¹ Using $B(\psi(3770) \rightarrow J/\psi\pi^+\pi^-) = (1.89 \pm 0.20 \pm 0.20) \times 10^{-3}$ from ADAM 06.

NODE=M053R04
NODE=M053R04

NODE=M053R04;LINKAGE=CO

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$					Γ_{102}/Γ
VALUE (units 10^{-3})	CL% EVTS	DOCUMENT ID	TECN	COMMENT	
6.9±0.6 OUR AVERAGE					
6.7±0.7±0.2	2.2k	¹ ABLIKIM	16B BES3	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
7.3±0.7±0.6	274	BRIERE	06 CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma + \text{hadrons}$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 44	90	² COAN	06A CLEO	$e^+e^- \rightarrow \psi(3770) \rightarrow \gamma\gamma J/\psi$	

NODE=M053R01
NODE=M053R01

- ¹ ABLIKIM 16B reports $(6.88 \pm 0.28 \pm 0.67) \times 10^{-3}$ from a measurement of $[\Gamma(\psi(3770) \rightarrow \gamma\chi_{c0})/\Gamma_{\text{total}}] / [B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P))]$ assuming $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.99 \pm 0.27) \times 10^{-2}$, which we rescale to our best value $B(\psi(2S) \rightarrow \gamma\chi_{c0}(1P)) = (9.77 \pm 0.23) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.
- ² Using $\Gamma_{ee}(\psi(2S)) = (2.54 \pm 0.03 \pm 0.11)$ keV from ADAM 06 and taking $\sigma(e^+e^- \rightarrow D\bar{D})$ from HE 05 for $\sigma(e^+e^- \rightarrow \psi(3770))$.

NODE=M053R01;LINKAGE=B

 $\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c2})$ $\Gamma_{102}/\Gamma_{100}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M053R06
NODE=M053R06

••• We do not use the following data for averages, fits, limits, etc. •••

>8	90	¹ BRIERE	06	CLEO $e^+e^- \rightarrow \psi(3770)$
----	----	---------------------	----	--------------------------------------

¹ Not independent of other results in BRIERE 06.

NODE=M053R06;LINKAGE=BR

 $\Gamma(\gamma\chi_{c0})/\Gamma(\gamma\chi_{c1})$ $\Gamma_{102}/\Gamma_{101}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M053R05
NODE=M053R05

••• We do not use the following data for averages, fits, limits, etc. •••

2.5±0.6		¹ BRIERE	06	CLEO $e^+e^- \rightarrow \psi(3770)$
---------	--	---------------------	----	--------------------------------------

¹ Not independent of other results in BRIERE 06.

NODE=M053R05;LINKAGE=BR

 $\Gamma(\gamma\eta_c)/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M053R99
NODE=M053R99

$<7 \times 10^{-4}$	90	¹ ABLIKIM	14H	BES3
---------------------	----	----------------------	-----	------

¹ ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c)/\Gamma_{\text{total}}] \times [B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 16 \times 10^{-6}$ which we divide by our best value $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp) = 2.38 \times 10^{-2}$. We have calculated the best value of $B(\eta_c(1S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(1S) \rightarrow K\bar{K}\pi) = 7.1 \times 10^{-2}$.

NODE=M053R99;LINKAGE=AB

 $\Gamma(\gamma\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{104}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M053R20
NODE=M053R20

$<9 \times 10^{-4}$	90	¹ ABLIKIM	14H	BES3
---------------------	----	----------------------	-----	------

¹ ABLIKIM 14H reports $[\Gamma(\psi(3770) \rightarrow \gamma\eta_c(2S))/\Gamma_{\text{total}}] \times [B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)] < 5.6 \times 10^{-6}$ which we divide by our best value $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp) = 6 \times 10^{-3}$. We have calculated the best value of $B(\eta_c(2S) \rightarrow K_S^0 K^\pm \pi^\mp)$ as 1/3 of $B(\eta_c(2S) \rightarrow K\bar{K}\pi) = 1.9 \times 10^{-2}$.

NODE=M053R20;LINKAGE=AB

 $\Gamma(\gamma\eta')/\Gamma_{\text{total}}$ Γ_{105}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M053R14
NODE=M053R14

<1.8	90	¹ PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
--------	----	---------------------	----	--------------------------------------

¹ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.

NODE=M053R14;LINKAGE=PE

 $\Gamma(\gamma\eta)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M053R13
NODE=M053R13

<1.5	90	¹ PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
--------	----	---------------------	----	--------------------------------------

¹ Assuming maximal destructive interference between $\psi(3770)$ and continuum sources.

NODE=M053R13;LINKAGE=PE

 $\Gamma(\gamma\pi^0)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M053R12
NODE=M053R12

<2	90	PEDLAR	09	CLE3 $\psi(2S) \rightarrow \gamma X$
------	----	--------	----	--------------------------------------

 $\psi(3770)$ REFERENCES

NODE=M053

ABLIKIM	23BC	PR D108 052015	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62423
ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62437
ABLIKIM	23V	PR D107 L091101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62269
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21Z	PRL 127 082002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61265
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59697
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
SHAMOV	17	PL B769 187	A.G. Shamov, K.Yu. Todyshv		REFID=57900
ABLIKIM	16B	PL B753 103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57126
ABLIKIM	15J	PR D91 092009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56775
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)	REFID=56962
ABLIKIM	14H	PR D89 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55899
ABLIKIM	14L	PL B735 101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55903
ABLIKIM	14O	PR D90 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55906
BONVICINI	14	PR D89 072002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=55798

ABLIKIM	13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55393
ANASHIN	12A	PL B711 292	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=54055
ABLIKIM	10D	EPJ C66 11	M. Ablikim <i>et al.</i>	(BES II Collab.)	REFID=53350
BESSION	10	PRL 104 159901 (err.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
ABLIKIM	09C	EPJ C64 243	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=53134
PEDLAR	09	PR D79 111101	T.K. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=52998
ABLIKIM	08B	PL B659 74	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52129
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
ABLIKIM	08H	PRL 101 102004	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52254
ABLIKIM	08M	PL B670 179	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52569
ABLIKIM	08N	PL B670 184	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52570
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
BRODZICKA	08	PRL 100 092001	J. Brodzicka <i>et al.</i>	(BELLE Collab.)	REFID=52144
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
ABLIKIM	07B	PL B650 111	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51704
ABLIKIM	07E	PL B652 238	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51882
ABLIKIM	07F	PL B656 30	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51940
ABLIKIM	07I	EPJ C52 805	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52045
ABLIKIM	07K	PR D76 122002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52073
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
DOBBS	07	PR D76 112001	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=52075
ABLIKIM	06L	PRL 97 121801	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51129
ABLIKIM	06N	PL B641 145	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51131
ADAM	06	PRL 96 082004	N.E. Adam <i>et al.</i>	(CLEO Collab.)	REFID=50989
ADAMS	06	PR D73 012002	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50990
BESSION	06	PRL 96 092002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51041
Also		PRL 104 159901 (err.)	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=53245
BRIERE	06	PR D74 031106	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=51149
COAN	06A	PRL 96 182002	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51155
CRONIN-HEN...	06	PR D74 012005	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=51156
HUANG	06A	PRL 96 032003	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=50999
BAI	05	PL B605 63	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50332
HE	05	PRL 95 121801	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=50924
Also		PRL 96 199903 (err.)	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51211
ABLIKIM	04F	PR D70 077101	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50185
ATHAR	04	PR D70 112002	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50331
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
ADLER	88C	PRL 60 89	J. Adler <i>et al.</i>	(Mark III Collab.)	REFID=40361
SCHINDLER	80	PR D21 2716	R.H. Schindler <i>et al.</i>	(Mark II Collab.)	REFID=22222
BACINO	78	PRL 40 671	W.J. Bacino <i>et al.</i>	(SLAC, UCLA, UCI)	REFID=11437
RAPIDIS	77	PRL 39 526	P.A. Rapidis <i>et al.</i>	(LGW Collab.)	REFID=22220

NODE=M212

 $\psi_2(3823)$

$$I^G(J^{PC}) = 0^-(2^{--})$$

I, J, P need confirmation.

was $\psi(3823)$, $X(3823)$

Seen by BHARDWAJ 13 in $B \rightarrow \chi_{c1} \gamma K$ and ABLIKIM 15S in $e^+ e^- \rightarrow \pi^+ \pi^- \gamma \chi_{c1}$ decays as a narrow peak in the invariant mass distribution of the $\chi_{c1} \gamma$ system. Properties consistent with the $\psi_2(1^3D_2) c\bar{c}$ state.

NODE=M212

 $\psi_2(3823)$ MASS

NODE=M212M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3823.51 ± 0.34 OUR AVERAGE				

NODE=M212M

[3823.5 ± 0.5 MeV OUR 2023 AVERAGE Scale factor = 1.4]

NEW

3824.5 ± 2.4 ± 1.0	30	¹ ABLIKIM 23J	BES3	$e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$
3823.12 ± 0.43 ± 0.13	120	ABLIKIM 22R	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
3824.08 ± 0.53 ± 0.14	137	² AAIJ 20S	LHCB	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
3823.1 ± 1.8 ± 0.7	33 ± 10	³ BHARDWAJ 13	BELL	$B^\pm \rightarrow \chi_{c1} \gamma K^\pm$
3821.7 ± 1.3 ± 0.7	19 ± 5	⁴ ABLIKIM 15S	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$

¹ Using the measured $m_{\psi_2(3823)} - m_{\psi(2S)}$ and assuming $m_{\psi(2S)} = 3686.097$ MeV from PDG 22.

NODE=M212M;LINKAGE=D

² Using the measured $m_{\psi_2(3823)} - m_{\psi(2S)} = 137.98 \pm 0.53 \pm 0.14$ MeV.

NODE=M212M;LINKAGE=C

³ From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics. Corrected for the measured $\psi(2S)$ mass using $B \rightarrow \psi(2S) K \rightarrow (\gamma \chi_{c1}) K$ decays.

NODE=M212M;LINKAGE=A

⁴ From a simultaneous unbinned maximum likelihood fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1} \gamma$ and $\psi_2(3823) \rightarrow \chi_{c1} \gamma$ together, with floating mass scale offset for $\psi(2S)$, floating $\psi_2(3823)$ mass, and zero $\psi_2(3823)$ width, resulting in a significance of 5.9σ when including systematic uncertainties. Superseded by ABLIKIM 22R.

NODE=M212M;LINKAGE=B

$m_{\psi_2(3823)} - m_{\psi(2S)}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

137.98 ± 0.53 ± 0.14	137	¹ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
----------------------	-----	-------------------	----------	--

¹AAIJ 20S also reports $m_{\chi_{c1}(3872)} - m_{\psi_2(3823)} = 47.50 \pm 0.53 \pm 0.13$ MeV.

NODE=M212A00
NODE=M212A00

NODE=M212A00;LINKAGE=A

 $\psi_2(3823)$ WIDTH

VALUE (MeV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	-----	------	-------------	------	---------

< 2.9	90	120	ABLIKIM	22R BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
-------	----	-----	---------	----------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 18.8	90	30	¹ ABLIKIM	23J BES3	$e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$
--------	----	----	----------------------	----------	--

< 5.2	90		² AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
-------	----	--	-------------------	----------	--

< 16	90		³ ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
------	----	--	----------------------	----------	--

< 24	90		⁴ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1} \gamma K^\pm$
------	----	--	-----------------------	---------	--

¹From a fit of $e^+ e^- \rightarrow \pi^0 \pi^0 \chi_{c1} \gamma$ data at \sqrt{s} values from 4.23 to 4.70 GeV to a Breit-Wigner function with floating width, using the Bayesian approach.

²AAIJ 20S also provides a limit of < 6.6 MeV with 95% CL.

³From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$ data (the $\pi^+ \pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to a Breit-Wigner function with the mass fixed from the likelihood fit above, Gaussian resolution smearing, and floating width.

⁴From a simultaneous fit to $B^\pm \rightarrow (\chi_{c1} \gamma) K^\pm$ and $B^0 \rightarrow (\chi_{c1} \gamma) K_S^0$ with significance 4.0σ including systematics.

NODE=M212W

NODE=M212W

NODE=M212W;LINKAGE=D

NODE=M212W;LINKAGE=C

NODE=M212W;LINKAGE=B

NODE=M212W;LINKAGE=A

 $\psi_2(3823)$ DECAY MODES

Branching fractions are given relative to the one **DEFINED AS 1**.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $J/\psi(1S) \pi^+ \pi^-$	<0.06	90%
Γ_2 $J/\psi(1S) \pi^0 \pi^0$	<0.11	90%
Γ_3 $J/\psi(1S) \pi^0$	<0.030	90%
Γ_4 $J/\psi(1S) \eta$	<0.14	90%
Γ_5 $\chi_{c0} \gamma$	<0.24	90%
Γ_6 $\chi_{c1} \gamma$	DEFINED AS 1	
Γ_7 $\chi_{c2} \gamma$	0.28 $^{+0.14}_{-0.11}$	

NODE=M212215;NODE=M212

NODE=M212

 $\psi_2(3823)$ BRANCHING RATIOS

$\Gamma(J/\psi(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen		¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$
----------	--	----------------------	----------	-------------------------------------

seen	137 ± 26	AAIJ	20S LHCb	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$
------	----------	------	----------	--

¹From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

NODE=M212225

NODE=M212R00
NODE=M212R00

NODE=M212R00;LINKAGE=A

$\Gamma(J/\psi(1S) \pi^+ \pi^-) / \Gamma(\chi_{c1} \gamma)$ Γ_1/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.06	90	¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$
-------	----	----------------------	----------	-------------------------------------

¹From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

NODE=M212R06
NODE=M212R06

NODE=M212R06;LINKAGE=A

$\Gamma(J/\psi(1S) \pi^0 \pi^0) / \Gamma(\chi_{c1} \gamma)$ Γ_2/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.11	90	¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$
-------	----	----------------------	----------	-------------------------------------

¹From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

NODE=M212R07
NODE=M212R07

NODE=M212R07;LINKAGE=A

$\Gamma(J/\psi(1S) \pi^0) / \Gamma(\chi_{c1} \gamma)$ Γ_3/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.03	90	¹ ABLIKIM	210 BES3	$e^+ e^- \rightarrow \pi^+ \pi^- X$
-------	----	----------------------	----------	-------------------------------------

¹From a simultaneous unbinned maximum likelihood fit of the $\pi^+ \pi^-$ recoil mass distributions of seven decay channels in the process $e^+ e^- \rightarrow \pi^+ \pi^- X$.

NODE=M212R08
NODE=M212R08

NODE=M212R08;LINKAGE=A

$\Gamma(J/\psi(1S)\eta)/\Gamma(\chi_{c1}\gamma)$ Γ_4/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.14	90	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M212R09
 NODE=M212R09

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R09;LINKAGE=A

 $\Gamma(J/\psi(1S)\eta)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_4/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
$4.4^{+2.5}_{-1.9} \pm 0.9$	¹ AAIJ	22D LHCB	$B^+ \rightarrow J/\psi(1S)\eta K^+$

NODE=M212R10
 NODE=M212R10

¹ Using the branching ratio for $B^+ \rightarrow \psi_2(3823)K^+$ with $\psi_2(3823) \rightarrow J/\psi(1S)\pi^+\pi^-$ from AAIJ 20S.

NODE=M212R10;LINKAGE=A

 $\Gamma(\chi_{c0}\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M212R04
 NODE=M212R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R04;LINKAGE=A

 $\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	120	¹ ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
seen	63 ± 9	² ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
seen	16 ± 5	³ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^0\pi^0X$
seen	33 ± 10	⁴ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$

NODE=M212R01
 NODE=M212R01

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak R with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV, which leads to $\Gamma(e^+e^-)B(R \rightarrow \pi^+\pi^-\psi_2(3823))B(\psi_2(3823) \rightarrow \chi_{c1}\gamma) = 0.57 \pm 0.08$ eV.

NODE=M212R01;LINKAGE=D

² From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$. Signal has a 11.8 σ significance.

NODE=M212R01;LINKAGE=B

³ From a fit of the invariant $\pi^0\pi^0$ recoil-mass distribution. Signal has a 4.3 σ significance.

NODE=M212R01;LINKAGE=C

⁴ BHARDWAJ 13 reports $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c1}) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$ with statistical significance 3.8 σ .

NODE=M212R01;LINKAGE=A

 $\Gamma(\chi_{c0}\gamma)/\Gamma(\chi_{c1}\gamma)$ Γ_5/Γ_6

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.24	90	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M212R05
 NODE=M212R05

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

NODE=M212R05;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-X$
not seen	² ABLIKIM	15S BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$
not seen	³ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2}\gamma K^\pm$

NODE=M212R02
 NODE=M212R02

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$. Signal has a 3.2 σ significance.

NODE=M212R02;LINKAGE=C

² From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c2}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c2}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.

NODE=M212R02;LINKAGE=B

³ BHARDWAJ 13 reports $B(B^\pm \rightarrow \psi_2(3823)K^\pm) \times B(\psi_2(3823) \rightarrow \gamma\chi_{c2}) < 3.6 \times 10^{-6}$ at 90% CL.

NODE=M212R02;LINKAGE=A

$\Gamma(\chi_{c2}\gamma)/\Gamma(\chi_{c1}\gamma)$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_6
$0.28^{+0.14}_{-0.11} \pm 0.02$		9 ± 4	¹ ABLIKIM	210 BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$	
●●● We do not use the following data for averages, fits, limits, etc. ●●●						
<0.42		90	² ABLIKIM	15s BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c2}\gamma$	
<0.41		90	BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}/c2\gamma K^\pm$	

¹ From a simultaneous unbinned maximum likelihood fit of the $\pi^+\pi^-$ recoil mass distributions of seven decay channels in the process $e^+e^- \rightarrow \pi^+\pi^-X$.

² From a simultaneous unbinned maximum likelihood fit of $e^+e^- \rightarrow \pi^+\pi^-\chi_{c1(2)}\gamma$ data (the $\pi^+\pi^-$ recoil mass) taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to simulated events including both $\psi(2S) \rightarrow \chi_{c1(2)}\gamma$ and $\psi_2(3823) \rightarrow \chi_{c1(2)}\gamma$ together, with floating mass scale offset for $\psi(2S)$, $\psi_2(3823)$ mass floating (fixed to that above), and zero $\psi_2(3823)$ width.

NODE=M212R03
NODE=M212R03

NODE=M212R03;LINKAGE=B

NODE=M212R03;LINKAGE=A

 $\psi_2(3823)$ REFERENCES

ABLIKIM	23J	JHEP 2302 171	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	22D	JHEP 2204 046	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
ABLIKIM	21O	PR D103 L091102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)

NODE=M212

REFID=62052
REFID=61647
REFID=61664
REFID=61634
REFID=61121
REFID=60526
REFID=56784
REFID=55412

 $\psi_3(3842)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

J, P need confirmation.

J^P has not been measured, 3^- is the quark model prediction.

NODE=M241

NODE=M241

 $\psi_3(3842)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$3842.71 \pm 0.16 \pm 0.12$	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$

NODE=M241M

NODE=M241M

 $\psi_3(3842)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$2.79 \pm 0.51 \pm 0.35$	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$

NODE=M241W

NODE=M241W

 $\psi_3(3842)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 D^+D^-	seen
Γ_2 $D^0\bar{D}^0$	seen

NODE=M241215;NODE=M241

DESIG=1

DESIG=2

 $\psi_3(3842)$ BRANCHING RATIOS $\Gamma(D^+D^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$	

NODE=M241225

NODE=M241R01
NODE=M241R01

●●● We do not use the following data for averages, fits, limits, etc. ●●●

possibly seen ¹ ABLIKIM 22AL BES3 $e^+e^- \rightarrow \pi^+\pi^-D^+D^-$

¹ From a fit to the $\pi^+\pi^-$ recoil mass for $e^+e^- \rightarrow D^+D^-\pi^+\pi^-$.

NODE=M241R01;LINKAGE=A

 $\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$	

NODE=M241R02
NODE=M241R02

 $\psi_3(3842)$ REFERENCES

ABLIKIM	22AL	PR D106 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M241

REFID=61884
REFID=59697

$\chi_{c0}(3860)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

NODE=M237

OMITTED FROM SUMMARY TABLE

The assignment $J^P = 0^+$ is preferred over 2^+ by 2.5 sigma.

NODE=M237

Observed by CHILIKIN 17 using full amplitude analysis of the process $e^+ e^- \rightarrow J/\psi D \bar{D}$, where $D = D^0, D^+$. Not seen by AAIJ 20AI in the decay $B^+ \rightarrow D^+ D^- K^+$.

 $\chi_{c0}(3860)$ MASS

NODE=M237M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3862^{+26+40}_{-32-13}	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D \bar{D}$

NODE=M237M

 $\chi_{c0}(3860)$ WIDTH

NODE=M237W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200^{+180}_{-110} OUR AVERAGE	[201 ⁺¹⁸⁰ ₋₁₁₀ MeV OUR 2023 AVERAGE]		
$201^{+154+88}_{-67-82}$	CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D \bar{D}$

NODE=M237W

NEW

 $\chi_{c0}(3860)$ DECAY MODES

NODE=M237215;NODE=M237

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad D^0 \bar{D}^0$	seen
$\Gamma_2 \quad D^+ D^-$	seen

DESIG=1

DESIG=2

 $\chi_{c0}(3860)$ BRANCHING RATIOS

NODE=M237220

$\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D^0 \bar{D}^0$	

NODE=M237R00
NODE=M237R00

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen		CHILIKIN	17	BELL $e^+ e^- \rightarrow J/\psi D^+ D^-$	

NODE=M237R01
NODE=M237R01 $\chi_{c0}(3860)$ REFERENCES

NODE=M237

AAIJ	20AI	PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60739
CHILIKIN	17	PR D95 112003	K. Chilikin <i>et al.</i>	(BELLE Collab.) JPC	REFID=57995

$\chi_{c1}(3872)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

also known as $X(3872)$

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

First observed by CHOI 03 in $B \rightarrow K\pi^+\pi^- J/\psi(1S)$ decays as a narrow peak in the invariant mass distribution of the $\pi^+\pi^- J/\psi(1S)$ final state. Isovector hypothesis excluded by AUBERT 05B and CHOI 11.

AAIJ 13Q perform a full five-dimensional amplitude analysis of the angular correlations between the decay products in $B^+ \rightarrow \chi_{c1}(3872)K^+$ decays, where $\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-$ and $J/\psi \rightarrow \mu^+\mu^-$, which unambiguously gives the $J^{PC} = 1^{++}$ assignment under the assumption that the $\pi^+\pi^-$ and J/ψ are in an S -wave. AAIJ 15AO extend this analysis with more data to limit D -wave contributions to $< 4\%$ at 95% CL.

See the review on "Spectroscopy of Mesons Containing Two Heavy Quarks."

NODE=M176

NODE=M176

 $\chi_{c1}(3872)$ MASS FROM $J/\psi X$ MODE

NODE=M176M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3871.64 ± 0.06 OUR AVERAGE				
[3871.65 ± 0.06 MeV OUR 2023 AVERAGE]				
3870.2 ± 0.7 ± 0.3	24.6	ABLIKIM	23W BES3	$e^+e^- \rightarrow J/\psi(1S)\pi^+\pi^-\omega$
3871.64 ± 0.06 ± 0.01	19.8k	¹ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
3871.9 ± 0.7 ± 0.2	20	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$
3871.95 ± 0.48 ± 0.12	0.6k	AAIJ	12H LHCb	$pp \rightarrow J/\psi\pi^+\pi^-X$
3871.85 ± 0.27 ± 0.19	170	² CHOI	11 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3873 ⁺ 1.8 _{-1.6} ± 1.3	27	³ DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3871.61 ± 0.16 ± 0.19	6k	^{3,4} AALTONEN	09AU CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3871.4 ± 0.6 ± 0.1	93.4	AUBERT	08Y BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3868.7 ± 1.5 ± 0.4	9.4	AUBERT	08Y BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.8 ± 3.1 ± 3.0	522	^{3,5} ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3871.57 ± 0.09	155	⁶ AAIJ	23AP LHCb	$B_s^0 \rightarrow J/\psi 2(\pi^+\pi^-)$
3871.695 ± 0.067 ± 0.068	15.6k	⁷ AAIJ	20AD LHCb	$pp \rightarrow J/\psi\pi^+\pi^-X$
3871.59 ± 0.06 ± 0.03	4.2k	⁸ AAIJ	20S LHCb	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
3873.3 ± 1.1 ± 1.0	45	⁹ ABLIKIM	19V BES	$e^+e^- \rightarrow \gamma\omega J/\psi$
3860.0 ± 10.4	13.6	^{3,10} AGHASYAN	18A COMP	$\gamma^* N \rightarrow X\pi^\pm N'$
3868.6 ± 1.2 ± 0.2	8	¹¹ AUBERT	06 BABR	$B^0 \rightarrow K_S^0 J/\psi\pi^+\pi^-$
3871.3 ± 0.6 ± 0.1	61	¹¹ AUBERT	06 BABR	$B^- \rightarrow K^- J/\psi\pi^+\pi^-$
3873.4 ± 1.4	25	¹² AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi\pi^+\pi^-$
3871.3 ± 0.7 ± 0.4	730	^{3,13} ACOSTA	04 CDF2	$p\bar{p} \rightarrow J/\psi\pi^+\pi^-X$
3872.0 ± 0.6 ± 0.5	36	¹⁴ CHOI	03 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$
3836 ± 13	58	^{3,15} ANTONIAZZI	94 E705	$300 \pi^\pm Li \rightarrow J/\psi\pi^+\pi^-X$

NODE=M176M

NODE=M176M
NEW

OCCUR=2

OCCUR=2

OCCUR=2

- ¹ Calculated from $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.54 \pm 0.06$ MeV obtained by combining the data with $\chi_{c1}(3872)$ produced in B^+ decays from AAIJ 20S and inclusive b -hadron decays from AAIJ 20AD and using $m_{\psi(2S)} = 3686.097$ MeV. Breit-Wigner parametrization.
- ² The mass difference for the $\chi_{c1}(3872)$ produced in B^+ and B^0 decays is $(-0.71 \pm 0.96 \pm 0.19)$ MeV.
- ³ Width consistent with detector resolution.
- ⁴ A possible equal mixture of two states with a mass difference greater than 3.6 MeV/ c^2 is excluded at 95% CL.
- ⁵ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{J/\psi}$ using $m_{J/\psi} = 3096.916$ MeV.
- ⁶ From a fit of a relativistic S -wave Breit-Wigner convolved with the detector resolution. The width of $\chi_{c1}(3872)$ is constrained to the PDG 22 value. Systematic errors not evaluated.
- ⁷ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays and $m_{\psi(2S)} = 3686.097 \pm 0.010$ MeV. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.
- ⁸ Using Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.
- ⁹ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.
- ¹⁰ Could be a different state.
- ¹¹ Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3686.093$ MeV. Superseded by AUBERT 08Y.
- ¹² Calculated from the corresponding $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$ using $m_{\psi(2S)} = 3685.96$ MeV. Superseded by AUBERT 06.
- ¹³ Superseded by AALTONEN 09AU.
- ¹⁴ Superseded by CHOI 11.
- ¹⁵ A lower mass value can be due to an incorrect momentum scale for soft pions.

NODE=M176M;LINKAGE=F

NODE=M176M;LINKAGE=CO

NODE=M176M;LINKAGE=AC
NODE=M176M;LINKAGE=AANODE=M176M;LINKAGE=AB
NODE=M176M;LINKAGE=H

NODE=M176M;LINKAGE=D

NODE=M176M;LINKAGE=E
NODE=M176M;LINKAGE=B
NODE=M176M;LINKAGE=A

NODE=M176M;LINKAGE=AE

NODE=M176M;LINKAGE=AU

NODE=M176M;LINKAGE=AT
NODE=M176M;LINKAGE=CH
NODE=M176M;LINKAGE=AN

NODE=M176MD0

NODE=M176MD0

 $\chi_{c1}(3872)$ MASS FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$3873.71^{+0.56}_{-0.50} \pm 0.13$		¹ HIRATA	23 BELL	$B^0 \rightarrow D^0 \bar{D}^{*0} K^0$, $B^+ \rightarrow D^0 \bar{D}^{*0} K^+$
$3872.9^{+0.6}_{-0.4} \pm 0.4_{-0.5}$	50	^{2,3} AUSHEV	10 BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
$3875.1^{+0.7}_{-0.5} \pm 0.5$	33 ± 6	³ AUBERT	08B BABR	$B \rightarrow \bar{D}^{*0} D^0 K$
$3875.2 \pm 0.7^{+0.9}_{-1.8}$	24 ± 6	^{3,4} GOKHROO	06 BELL	$B \rightarrow D^0 \bar{D}^0 \pi^0 K$

- ¹ From a fit of a Breit-Wigner function with energy dependent width.
- ² Calculated from the measured $m_{\chi_{c1}(3872)} - m_{D^{*0}} - m_{D^0} = 1.1^{+0.6+0.1}_{-0.4-0.3}$ MeV.
- ³ Experiments report $D^{*0} \bar{D}^0$ invariant mass above $D^{*0} \bar{D}^0$ threshold because D^{*0} decay products are kinematically constrained to the D^{*0} mass, even though the D^{*0} may decay off-shell.
- ⁴ Superseded by AUSHEV 10.

NODE=M176MD0;LINKAGE=A
NODE=M176MD0;LINKAGE=AS
NODE=M176MD0;LINKAGE=AU

NODE=M176MD0;LINKAGE=GO

 $m_{\chi_{c1}(3872)} - m_{J/\psi}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$774.9 \pm 3.1 \pm 3.0$	522	ABAZOV	04F D0	$p\bar{p} \rightarrow J/\psi \pi^+ \pi^- X$

NODE=M176DM

NODE=M176DM

 $m_{\chi_{c1}(3872)} - m_{\psi(2S)}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$185.598 \pm 0.067 \pm 0.068$	15.6k	¹ AAIJ	20AD LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
185.54 ± 0.06	19.8k	² AAIJ	20S LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
187.4 ± 1.4	25	³ AUBERT	05R BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

NODE=M176DM2

NODE=M176DM2

- ¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization. Superseded by the combined value in AAIJ 20S.
- ² Combining $m_{\chi_{c1}(3872)} - m_{\psi(2S)} = 185.49 \pm 0.06 \pm 0.03$ MeV from AAIJ 20S and the measured mass difference from AAIJ 20AD. Breit-Wigner parametrization.
- ³ Superseded by AUBERT 06.

NODE=M176DM2;LINKAGE=A

NODE=M176DM2;LINKAGE=E

NODE=M176DM2;LINKAGE=AU

 $\chi_{c1}(3872)$ WIDTH

VALUE (MeV)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
1.19 ± 0.21 OUR AVERAGE		Error includes scale factor of 1.1.		
$1.39 \pm 0.24 \pm 0.10$	15.6k	¹ AAIJ	20AD LHCB	$pp \rightarrow J/\psi \pi^+ \pi^- X$
$0.96^{+0.19}_{-0.18} \pm 0.21$	4.2k	² AAIJ	20S LHCB	$B^+ \rightarrow J/\psi \pi^+ \pi^- K^+$

NODE=M176W

NODE=M176W

OCCUR=3

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.4	90	ABLIKIM	14	BES3	$e^+e^- \rightarrow J/\psi \pi^+ \pi^- \gamma$
<1.2	90	CHOI	11	BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$
<3.3	90	AUBERT	08Y	BABR	$B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$
<4.1	90	69 AUBERT	06	BABR	$B \rightarrow K \pi^+ \pi^- J/\psi$
<2.3	90	36 ³ CHOI	03	BELL	$B \rightarrow K \pi^+ \pi^- J/\psi$

¹ Using $\chi_{c1}(3872)$ produced in inclusive b -hadron decays. Breit-Wigner parametrization.

² Using Breit-Wigner parametrization. Partially overlapping dataset with that of AAIJ 20AD.

³ Superseded by CHOI 11.

NODE=M176W;LINKAGE=E

NODE=M176W;LINKAGE=F

NODE=M176W;LINKAGE=CH

$\chi_{c1}(3872)$ WIDTH FROM $\bar{D}^{*0} D^0$ MODE

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M176WD0

NODE=M176WD0

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.2^{+2.2}_{-1.5} \pm 0.4$		¹ HIRATA	23	BELL	$B^0 \rightarrow D^0 \bar{D}^{*0} K^0$, $B^+ \rightarrow D^0 \bar{D}^{*0} K^+$
$3.9^{+2.8+0.2}_{-1.4-1.1}$	50	² AUSHEV	10	BELL	$B \rightarrow \bar{D}^{*0} D^0 K$
$3.0^{+1.9}_{-1.4} \pm 0.9$	33 ± 6	AUBERT	08B	BABR	$B \rightarrow \bar{D}^{*0} D^0 K$

¹ From a fit of a Breit-Wigner function with energy dependent width.

² With a measured value of $B(B \rightarrow \chi_{c1}(3872)K) \times B(\chi_{c1}(3872) \rightarrow D^{*0} \bar{D}^0) = (0.80 \pm 0.20 \pm 0.10) \times 10^{-4}$, assumed to be equal for both charged and neutral modes.

NODE=M176WD0;LINKAGE=B

NODE=M176WD0;LINKAGE=AU

$\chi_{c1}(3872)$ DECAY MODES

NODE=M176215;NODE=M176

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 e^+e^-	< 2.7 $\times 10^{-7}$	90%
Γ_2 $\pi^+ \pi^- \pi^0$	< 8 $\times 10^{-3}$	90%
Γ_3 $\pi^+ \pi^- J/\psi(1S)$	(3.5 \pm 0.9) %	
Γ_4 $\pi^+ \pi^- \pi^0 J/\psi(1S)$	not seen	
Γ_5 $\omega \eta_c(1S)$	< 30 %	90%
Γ_6 $\rho(770)^0 J/\psi(1S)$	(2.8 \pm 0.7) %	
Γ_7 $\omega J/\psi(1S)$	(4.1 \pm 1.4) %	
Γ_8 $\phi \phi$	not seen	
Γ_9 $D^0 \bar{D}^0 \pi^0$	(45 \pm 21) %	
Γ_{10} $\bar{D}^{*0} D^0$	(34 \pm 12) %	
Γ_{11} $\gamma \gamma$	< 11 %	90%
Γ_{12} $D^0 \bar{D}^0$	< 26 %	90%
Γ_{13} $D^+ D^-$	< 18 %	90%
Γ_{14} $\pi^0 \chi_{c2}$	< 4 %	90%
Γ_{15} $\pi^0 \chi_{c1}$	(3.1 $^{+1.5}_{-1.3}$) %	
Γ_{16} $\pi^0 \chi_{c0}$	< 13 %	90%
Γ_{17} $\pi^+ \pi^- \eta_c(1S)$	< 13 %	90%
Γ_{18} $\pi^0 \pi^0 \chi_{c0}$	< 6 %	90%
Γ_{19} $\pi^+ \pi^- \chi_{c0}$	< 2.0 %	90%
Γ_{20} $\pi^+ \pi^- \chi_{c1}$	< 7 $\times 10^{-3}$	90%
Γ_{21} $\rho \bar{\rho}$	< 2.2 $\times 10^{-5}$	95%

Radiative decays

Γ_{22} $\gamma D^+ D^-$	< 3.5 %	90%
Γ_{23} $\gamma \bar{D}^0 D^0$	< 6 %	90%
Γ_{24} $\gamma J/\psi$	(7.8 \pm 2.9) $\times 10^{-3}$	
Γ_{25} $\gamma \chi_{c1}$	< 8 $\times 10^{-3}$	90%
Γ_{26} $\gamma \chi_{c2}$	< 2.9 %	90%
Γ_{27} $\gamma \psi(2S)$	(4.2 \pm 1.6) %	

NODE=M176;CLUMP=B

DESIG=21

DESIG=23

DESIG=9

DESIG=3

DESIG=15

DESIG=11

C-violating decays

Γ_{28} $\eta J/\psi$	< 1.7 %	90%
-----------------------------	---------	-----

NODE=M176;CLUMP=A

DESIG=4

$\chi_{c1}(3872)$ PARTIAL WIDTHS

NODE=M176220

 $\Gamma(e^+e^-)$ Γ_1

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.32	90	¹ ABLIKIM	230 BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
••• We do not use the following data for averages, fits, limits, etc. •••				
< 4.3	90	² ABLIKIM	15V BES3	$4.0-4.4 e^+e^- \rightarrow \pi^+\pi^- J/\psi$
<280	90	³ YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

NODE=M176W1
NODE=M176W1

¹ Fit to cross section using a total width value of 1.19 ± 0.21 MeV and $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = (3.8 \pm 1.2)\%$ from PDG 20.

NODE=M176W1;LINKAGE=C

² ABLIKIM 15V reports this limit from the measurement of $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) \times \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-)/\Gamma < 0.13$ eV using $\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))/\Gamma = 3\%$.

NODE=M176W1;LINKAGE=B

³ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. Assuming that $\Gamma(\pi^+\pi^- J/\psi)$ of $\chi_{c1}(3872)$ is the same as that of $\psi(2S)$ (85.4 keV).

NODE=M176W1;LINKAGE=A

 $\chi_{c1}(3872) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M176230

 $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 7.5×10^{-3} (CL = 90%)		[<0.13 eV (CL = 90%) OUR 2023 BEST LIMIT]		
< 7.5×10^{-3}	90	¹ ABLIKIM	230 BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
••• We do not use the following data for averages, fits, limits, etc. •••				
< 0.13	90	ABLIKIM	15V BES3	$4.0-4.4 e^+e^- \rightarrow \pi^+\pi^- J/\psi$
< 6.2	90	^{2,3} AUBERT	05D BABR	$10.6 e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
< 8.3	90	³ DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
<10	90	⁴ YUAN	04 RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

NODE=M176G1
NODE=M176G1

¹ Fit to cross section using a total width value of 1.19 ± 0.21 MeV from PDG 20.

NODE=M176G1;LINKAGE=B

² Using $B(\chi_{c1}(3872) \rightarrow J/\psi\pi^+\pi^-) \cdot B(J/\psi \rightarrow \mu^+\mu^-) \cdot \Gamma(\chi_{c1}(3872) \rightarrow e^+e^-) < 0.37$ eV from AUBERT 05D and $B(J/\psi \rightarrow \mu^+\mu^-) = 0.0588 \pm 0.0010$ from the PDG 04.

NODE=M176G1;LINKAGE=AU

³ Assuming $\chi_{c1}(3872)$ has $J^{PC} = 1^{--}$.

NODE=M176G1;LINKAGE=DO

⁴ Using BAI 98E data on $e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$. From theoretical calculation of the production cross section and using $B(J/\psi \rightarrow \mu^+\mu^-) = (5.88 \pm 0.10)\%$.

NODE=M176G1;LINKAGE=A

 $\chi_{c1}(3872) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M176232

 $\Gamma(\pi^+\pi^- J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_{11}/\Gamma$

VALUE (eV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
$5.5^{+4.1}_{-3.8} \pm 0.7$		3	¹ TERAMOTO	21 BELL	$e^+e^- \rightarrow \gamma^*\gamma$ at $\Upsilon(nS)$
<12.9	90		² DOBBS	05 CLE3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi\gamma$

NODE=M176H1
NODE=M176H1

¹ Measured in single-tag two-photon production assuming Q^2 dependence of a $c\bar{c}$ meson model. Here, $\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)$ is the reduced two-photon decay width, $\tilde{\Gamma}_{\gamma\gamma}$.

NODE=M176H1;LINKAGE=A

² Assuming $\chi_{c1}(3872)$ has positive C parity and spin 0.

NODE=M176H1;LINKAGE=DO

 $\Gamma(\omega J/\psi(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_{11}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.7	90	¹ LEES	12AD BABR	$e^+e^- \rightarrow e^+e^-\omega J/\psi$

NODE=M176G01
NODE=M176G01

¹ Assuming $\chi_{c1}(3872)$ has spin 2.

NODE=M176G01;LINKAGE=LE

 $\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_{11}/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<11.1	90	LEES	12AE BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

NODE=M176G02
NODE=M176G02

$\chi_{c1}(3872)$ BRANCHING RATIOS

NODE=M176235

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

<0.8 (CL = 90%) [$<0.9\%$ (CL = 90%) OUR 2023 BEST LIMIT]

<0.8	90	1,2 YIN	23 BELL	$B^+ \rightarrow \chi_{c1}(3872)K^+$
----------------	----	---------	---------	--------------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

<1.1	90	2,3 YIN	23 BELL	$B^0 \rightarrow \chi_{c1}(3872)K^0$
------	----	---------	---------	--------------------------------------

OCCUR=2

¹ YIN 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.

NODE=M176R30;LINKAGE=A

² Assuming the decay products, $\pi^+\pi^-\pi^0$, are uniformly distributed in phase space. The limit is the 90% "credible" upper limit (i.e. Bayesian).

NODE=M176R30;LINKAGE=E

³ YIN 23 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow \chi_{c1}(3872)K^0)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^0 \rightarrow \chi_{c1}(3872)K^0) = 1.3 \times 10^{-4}$.

NODE=M176R30;LINKAGE=D

 $\Gamma(\pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.035 ± 0.009 OUR AVERAGE

[0.038 ± 0.012 OUR 2023 AVERAGE]

0.035 ± 0.002 ± 0.009		¹ AAIJ	20S LHCB	$B^+ \rightarrow J/\psi\pi^+\pi^-K^+$
-----------------------	--	-------------------	----------	---------------------------------------

SYCLP=A

0.038 ± 0.004 ± 0.010		² CHOI	11 BELL	$B^+ \rightarrow \pi^+\pi^-J/\psi K^+$
-----------------------	--	-------------------	---------	--

SYCLP=A

0.037 ± 0.007 ± 0.010	93	^{3,4} AUBERT	08Y BABR	$B \rightarrow \chi_{c1}(3872)K$
-----------------------	----	-----------------------	----------	----------------------------------

SYCLP=A

••• We do not use the following data for averages, fits, limits, etc. •••

seen	151	⁵ BALA	15 BELL	$B \rightarrow \chi_{c1}(3872)K\pi$
------	-----	-------------------	---------	-------------------------------------

0.056 ± 0.018 ± 0.015	30	⁶ AUBERT	05R BABR	$B^+ \rightarrow K^+\pi^+\pi^-J/\psi$
-----------------------	----	---------------------	----------	---------------------------------------

SYCLP=A

0.060 ± 0.013 ± 0.016	36	⁷ CHOI	03 BELL	$B^+ \rightarrow K^+\pi^+\pi^-J/\psi$
-----------------------	----	-------------------	---------	---------------------------------------

¹ AAIJ 20S reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (7.95 \pm 0.15 \pm 0.33) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R6;LINKAGE=E

² CHOI 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.63 \pm 0.82 \pm 0.52) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R6;LINKAGE=F

³ AUBERT 08Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (8.4 \pm 1.5 \pm 0.7) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R6;LINKAGE=AB

⁴ superseded by LEES 20C

NODE=M176R6;LINKAGE=C

⁵ BALA 15 reports $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^0 \rightarrow \chi_{c1}(3872)K^+\pi^-) = (7.9 \pm 1.3 \pm 0.4) \times 10^{-6}$ and $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi) \times B(B^+ \rightarrow \chi_{c1}(3872)K^0\pi^+) = (10.6 \pm 3.0 \pm 0.9) \times 10^{-6}$.

NODE=M176R6;LINKAGE=A

⁶ Superseded by AUBERT 08Y. AUBERT 05R reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.28 \pm 0.41) \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R6;LINKAGE=AE

⁷ CHOI 03 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] / [B(B^+ \rightarrow \psi(2S)K^+)] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = 0.063 \pm 0.012 \pm 0.007$ which we multiply or divide by our best values $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$, $B(B^+ \rightarrow \psi(2S)K^+) = (6.24 \pm 0.21) \times 10^{-4}$, $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.69 \pm 0.34) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best values.

NODE=M176R6;LINKAGE=CH

 $\Gamma(\pi^+\pi^-\pi^0J/\psi(1S))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	¹ WANG	11B BELL	$\Upsilon(2S) \rightarrow \gamma X$
----------	-------------------	----------	-------------------------------------

NODE=M176R25

not seen	² SHEN	10A BELL	$\Upsilon(1S) \rightarrow \gamma X$
-----------------	-------------------	----------	-------------------------------------

NODE=M176R25

¹ WANG 11B reports $B(\Upsilon(2S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0J/\psi) < 2.4 \times 10^{-6}$ at 95% CL.

NODE=M176R25;LINKAGE=B

² SHEN 10A reports $B(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \pi^+\pi^-\pi^0J/\psi) < 2.8 \times 10^{-6}$ at 95% CL.

NODE=M176R25;LINKAGE=A

$\Gamma(\omega\eta_c(1S))/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.30 (CL = 90%) [**<0.33 (CL = 90%) OUR 2023 BEST LIMIT**]**<0.30** 90 ¹ VINOKUROVA 15 BELL $B^+ \rightarrow \omega\eta_c K^+$

¹ VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega\eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.9 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.

NODE=M176R24
NODE=M176R24

NODE=M176R24;LINKAGE=A

 $\Gamma(\rho(770)^0 J/\psi(1S))/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_6/Γ_3

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

78.6±2.3±2.0 ¹ AAIJ 23s LHCb $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$

¹ Assuming pure ρ contribution only, i.e. excluding the contribution from $\rho\omega$ interference. Using $B(\rho^0 \rightarrow \pi^+ \pi^-) = 100\%$.

NODE=M176R33
NODE=M176R33

NODE=M176R33;LINKAGE=A

 $\Gamma(\omega J/\psi(1S))/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.026±0.010±0.007 21±7 ¹ DEL-AMO-SA..10B BABR $B^+ \rightarrow \omega J/\psi K^+$

¹ DEL-AMO-SANCHEZ 10B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (6 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. DEL-AMO-SANCHEZ 10B also reports $B(B^0 \rightarrow \chi_{c1}(3872)K^0) \times B(\chi_{c1}(3872) \rightarrow J/\psi\omega) = (6 \pm 3 \pm 1) \times 10^{-6}$.

NODE=M176R14
NODE=M176R14

NODE=M176R14;LINKAGE=DE

 $\Gamma(\omega J/\psi(1S))/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_7/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.16±0.24 OUR AVERAGE Error includes scale factor of 1.2. [1.1 ± 0.4 OUR 2023 AVERAGE Scale factor = 1.7]1.24±0.33±0.10 ^{1,2} AAIJ 23s LHCb $B^+ \rightarrow K^+ J/\psi \pi^+ \pi^-$ 1.6 ^{+0.4} _{-0.3} ±0.2 ³ ABLIKIM 19v BES $e^+ e^- \rightarrow \gamma\omega J/\psi$ 0.8 ±0.3 ⁴ DEL-AMO-SA..10B BABR $B \rightarrow \omega J/\psi K$

¹ AAIJ 23s reports $[\Gamma(\chi_{c1}(3872) \rightarrow \omega J/\psi(1S))/\Gamma(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] \times [B(\omega(782) \rightarrow \pi^+ \pi^-)] = (1.9 \pm 0.4 \pm 0.3) \times 10^{-2}$ which we divide by our best value $B(\omega(782) \rightarrow \pi^+ \pi^-) = (1.53 \pm 0.12) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Excluding $\rho\omega$ interference effects.

³ Fit with fixed width and including two resonances, $\chi_{c0}(3915)$ and $X(3960)$.

⁴ Statistical and systematic errors added in quadrature. Uses the values of $B(B \rightarrow \chi_{c1}(3872)K) \times B(\chi_{c1}(3872) \rightarrow J/\psi \pi^+ \pi^-)$ reported in AUBERT 08Y, taking into account the common systematics.

NODE=M176R15
NODE=M176R15

NEW

OCCUR=3

NODE=M176R15;LINKAGE=G

NODE=M176R15;LINKAGE=H
NODE=M176R15;LINKAGE=A
NODE=M176R15;LINKAGE=DE $\Gamma(\phi\phi)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen ¹ AAIJ 17BB LHCb pp at 7, 8 TeV

¹ AAIJ 17BB reports $B(b \rightarrow \chi_{c1}(3872)\text{anything}) \times B(\chi_{c1}(3872) \rightarrow \phi\phi) < 4.5 \times 10^{-7}$ at 95% CL.

NODE=M176R26
NODE=M176R26

NODE=M176R26;LINKAGE=A

 $\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
-------	-----	------	-------------	------	---------

0.45^{+0.16}_{-0.19}±0.12 17 ¹ GOKHROO 06 BELL $B^+ \rightarrow D^0 \bar{D}^0 \pi^0 K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.26 90 ² CHISTOV 04 BELL Sup. by GOKHROO 06

¹ GOKHROO 06 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.02 \pm 0.31^{+0.21}_{-0.29}) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0 \pi^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 0.6 \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.

NODE=M176R12
NODE=M176R12

NODE=M176R12;LINKAGE=GO

NODE=M176R12;LINKAGE=A

 $\Gamma(D^0 \bar{D}^0 \pi^0)/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_9/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.16 90 ABLIKIM 20w BES3 $e^+ e^- \rightarrow \gamma\chi_{c1}(3872)$ NODE=M176R17
NODE=M176R17

$\Gamma(\bar{D}^{*0} D^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.34±0.08±0.09	41 ⁺⁹ ₋₈	¹ AUSHEV	10 BELL	$B^+ \rightarrow D^{*0} \bar{D}^0 K^+$

NODE=M176R13
 NODE=M176R13
 SYCLP=A

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.73±0.26±0.19 27 ± 6 ² AUBERT 08B BABR $B^+ \rightarrow \bar{D}^{*0} D^0 K^+$

SYCLP=A

¹ AUSHEV 10 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (0.77 \pm 0.16 \pm 0.10) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R13;LINKAGE=AS

² AUBERT 08B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \bar{D}^{*0} D^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] = (1.67 \pm 0.36 \pm 0.47) \times 10^{-4}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R13;LINKAGE=AU

 $\Gamma(\bar{D}^{*0} D^0)/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{10}/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
11.77±3.09	50	ABLIKIM	20W BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

NODE=M176R16
 NODE=M176R16

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.11	90	¹ WICHT	08 BELL	$e^+ e^- \rightarrow \gamma(4S)$

NODE=M176R09
 NODE=M176R09

¹ WICHT 08 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\gamma)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 2.4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.3 \times 10^{-4}$.

NODE=M176R09;LINKAGE=A

 $\Gamma(D^0 \bar{D}^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.26 (CL = 90%)	[<0.29 (CL = 90%) OUR 2023 BEST LIMIT]			
<0.26	90	¹ CHISTOV	04 BELL	$B \rightarrow K D^0 \bar{D}^0$

NODE=M176R3
 NODE=M176R3

¹ CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^0 \bar{D}^0)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 6 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.3 \times 10^{-4}$.

NODE=M176R3;LINKAGE=A

 $\Gamma(D^+ D^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.18 (CL = 90%)	[<0.19 (CL = 90%) OUR 2023 BEST LIMIT]			
<0.18	90	¹ CHISTOV	04 BELL	$B \rightarrow K D^+ D^-$

NODE=M176R4
 NODE=M176R4

¹ CHISTOV 04 reports $[\Gamma(\chi_{c1}(3872) \rightarrow D^+ D^-)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 4 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.3 \times 10^{-4}$.

NODE=M176R4;LINKAGE=A

 $\Gamma(\pi^0 \chi_{c2})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{14}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	ABLIKIM	19U BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

NODE=M176R06
 NODE=M176R06

 $\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	¹ BHARDWAJ	19 BELL	$B^\pm \rightarrow \pi^0 \chi_{c1} K^\pm$

NODE=M176R23
 NODE=M176R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ BHARDWAJ 19 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^0 \chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872) K^+)] < 8.1 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872) K^+) = 2.3 \times 10^{-4}$.

NODE=M176R23;LINKAGE=A

 $\Gamma(\pi^0 \chi_{c1})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{15}/Γ_3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
88⁺³³₋₂₇ ± 10	10.8	ABLIKIM	19U BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

NODE=M176R05
 NODE=M176R05

 $\Gamma(\pi^0 \chi_{c0})/\Gamma(\pi^+ \pi^- J/\psi(1S))$ Γ_{16}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 3.6	90	ABLIKIM	22D BES3	$e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

NODE=M176R04
 NODE=M176R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19 90 ABLIKIM 19U BES3 $e^+ e^- \rightarrow \gamma \chi_{c1}(3872)$

$\Gamma(\pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.13 (CL = 90%)				[<0.14 (CL = 90%) OUR 2023 BEST LIMIT]
<0.13	90	¹ VINOKUROVA 15	BELL	$B^+ \rightarrow \pi^+\pi^-\eta_c K^+$
		¹ VINOKUROVA 15 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\eta_c(1S))/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 3.0 \times 10^{-5}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.		

NODE=M176R22
NODE=M176R22

NODE=M176R22;LINKAGE=A

 $\Gamma(\pi^0\pi^0\chi_{c0})/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{18}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	ABLIKIM	22D	BES3 $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R28
NODE=M176R28 $\Gamma(\pi^+\pi^-\chi_{c0})/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{19}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.56	90	ABLIKIM	22D	BES3 $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R29
NODE=M176R29 $\Gamma(\pi^+\pi^-\chi_{c1})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7 × 10⁻³	90	¹ BHARDWAJ 16	BELL	$B^+ \rightarrow \pi^+\pi^-\chi_{c1}K^+$
		¹ BHARDWAJ 16 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \pi^+\pi^-\chi_{c1})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.5 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.		

NODE=M176R00
NODE=M176R00

NODE=M176R00;LINKAGE=A

 $\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10⁻⁵ (CL = 95%)				[<2.4 × 10 ⁻⁵ (CL = 95%) OUR 2023 BEST LIMIT]
<2.2 × 10⁻⁵	95	¹ AAIJ	17AD	LHCB $B^+ \rightarrow \rho\bar{\rho}K^+$
		••• We do not use the following data for averages, fits, limits, etc. •••		
<7 × 10⁻⁵	95	² AAIJ	13s	LHCB $B^+ \rightarrow \rho\bar{\rho}K^+$
		¹ AAIJ 17AD reports $[\Gamma(\chi_{c1}(3872) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 0.5 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.		
		² AAIJ 13s reports $[\Gamma(\chi_{c1}(3872) \rightarrow \rho\bar{\rho})/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.7 \times 10^{-8}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.		

NODE=M176R03
NODE=M176R03

SYCLP=A

SYCLP=A

NODE=M176R03;LINKAGE=C

NODE=M176R03;LINKAGE=B

Radiative decays

 $\Gamma(\gamma D^+D^-)/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{22}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.99	90	ABLIKIM	20W	BES3 $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176410

NODE=M176R20
NODE=M176R20 $\Gamma(\gamma \bar{D}^0D^0)/\Gamma(\pi^+\pi^-J/\psi(1S))$ Γ_{23}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.58	90	ABLIKIM	20W	BES3 $e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R21
NODE=M176R21 $\Gamma(\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
7.8 ± 2.9 OUR AVERAGE				[0.008 ± 0.004 OUR 2023 AVERAGE]
7.8^{+2.2}_{-2.0} ± 2.0		¹ BHARDWAJ 11	BELL	$B^\pm \rightarrow \gamma J/\psi K^\pm$

NODE=M176R7
NODE=M176R7NEW
SYCLP=A

••• We do not use the following data for averages, fits, limits, etc. •••

12.3 ± 3.5 ± 3.2	20	² AUBERT	09B	BABR $B^+ \rightarrow \gamma J/\psi K^+$
15 ± 5 ± 4	19	³ AUBERT, BE	06M	BABR $B^+ \rightarrow \gamma J/\psi K^+$

SYCLP=A

SYCLP=A

¹ BHARDWAJ 11 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (1.78^{+0.48}_{-0.44} \pm 0.12) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R7;LINKAGE=BA

² AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (2.8 \pm 0.8 \pm 0.1) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R7;LINKAGE=AB

³ Superseded by AUBERT 09B. AUBERT, BE 06M reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma J/\psi)/\Gamma_{\text{total}}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (3.3 \pm 1.0 \pm 0.3) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R7;LINKAGE=AU

$\Gamma(\gamma J/\psi)/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{24}/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.79±0.28	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R18
 NODE=M176R18

 $\Gamma(\gamma\chi_{c1})/\Gamma_{total}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8 × 10⁻³ (CL = 90%)	[<9 × 10 ⁻³ (CL = 90%)	OUR 2023	BEST LIMIT]	
<8 × 10⁻³	90	¹ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c1}\gamma K^\pm$
¹ BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\chi_{c1})/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 1.9 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.				

NODE=M176R08
 NODE=M176R08

NODE=M176R08;LINKAGE=B

 $\Gamma(\gamma\chi_{c1})/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{25}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.89	90	CHOI	03 BELL	$B \rightarrow K\pi^+\pi^- J/\psi$

NODE=M176R1
 NODE=M176R1

 $\Gamma(\gamma\chi_{c2})/\Gamma_{total}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.029 (CL = 90%)	[<0.032 (CL = 90%)	OUR 2023	BEST LIMIT]	
<0.029	90	¹ BHARDWAJ	13 BELL	$B^\pm \rightarrow \chi_{c2}\gamma K^\pm$
¹ BHARDWAJ 13 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\chi_{c2})/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 6.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.				

NODE=M176R01
 NODE=M176R01

NODE=M176R01;LINKAGE=B

 $\Gamma(\gamma\psi(2S))/\Gamma_{total}$ Γ_{27}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.042±0.016 OUR AVERAGE	[0.045 ± 0.020	OUR 2023	AVERAGE]	
0.042±0.012±0.011	25 ± 7	¹ AUBERT	09B BABR	$B^+ \rightarrow \gamma\psi(2S)K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	36 ± 9	² AAIJ	14AH LHCB	$B^+ \rightarrow \gamma\psi(2S)K^+$
not seen		³ BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$

NODE=M176R10
 NODE=M176R10
 NEW

¹ AUBERT 09B reports $[\Gamma(\chi_{c1}(3872) \rightarrow \gamma\psi(2S))/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] = (9.5 \pm 2.7 \pm 0.6) \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = (2.3 \pm 0.6) \times 10^{-4}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M176R10;LINKAGE=AU

² From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ.

NODE=M176R10;LINKAGE=A

³ BHARDWAJ 11 reports $B(B^+ \rightarrow K^+\chi_{c1}(3872)) \times B(\chi_{c1} \rightarrow \gamma\psi(2S)) < 3.45 \times 10^{-6}$ at 90% CL.

NODE=M176R10;LINKAGE=BH

 $\Gamma(\gamma\psi(2S))/\Gamma(\pi^+\pi^- J/\psi(1S))$ Γ_{27}/Γ_3

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.42	90	ABLIKIM	20W BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$

NODE=M176R19
 NODE=M176R19

 $\Gamma(\gamma\psi(2S))/\Gamma(\gamma J/\psi)$ Γ_{27}/Γ_{24}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.6 OUR AVERAGE					
2.46 ± 0.64 ± 0.29	36 ± 9	¹ AAIJ	14AH LHCB	$B^+ \rightarrow \gamma\psi(2S)K^+$	
3.4 ± 1.4		AUBERT	09B BABR	$B^+ \rightarrow \gamma c\bar{c}K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.1	90	BHARDWAJ	11 BELL	$B^+ \rightarrow \gamma\psi(2S)K^+$	

NODE=M176R11
 NODE=M176R11

¹ From 36.4 ± 9.0 events of $\chi_{c1}(3872) \rightarrow J/\psi\gamma$ decays with a statistical significance of 4.4σ.

NODE=M176R11;LINKAGE=A

C-violating decays

NODE=M176405

 $\Gamma(\eta J/\psi)/\Gamma_{total}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.017 (CL = 90%)	[<0.018 (CL = 90%)	OUR 2023	BEST LIMIT]	
<0.017	90	^{1,2} IWASHITA	14 BELL	$B \rightarrow K\eta J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.034	90	³ AUBERT	04Y BABR	$B \rightarrow K\eta J/\psi$

NODE=M176R2
 NODE=M176R2

¹ IWASHITA 14 reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 3.8 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.

NODE=M176R2;LINKAGE=A

² IWASHITA 14 also scans the $\eta J/\psi$ mass range 3.8–4.75 GeV and sets upper limits for $B(B^\pm \rightarrow \chi_{c1}(3872)K^\pm) \times B(\chi_{c1}(3872) \rightarrow \eta J/\psi)$ in 5 MeV intervals.

NODE=M176R2;LINKAGE=C

³ AUBERT 04Y reports $[\Gamma(\chi_{c1}(3872) \rightarrow \eta J/\psi)/\Gamma_{total}] \times [B(B^+ \rightarrow \chi_{c1}(3872)K^+)] < 7.7 \times 10^{-6}$ which we divide by our best value $B(B^+ \rightarrow \chi_{c1}(3872)K^+) = 2.3 \times 10^{-4}$.

NODE=M176R2;LINKAGE=D

χ_{c1} (3872) REFERENCES

NODE=M176

AAIJ	23AP	JHEP 2307 084	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62412
AAIJ	23S	PR D108 L011103	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62287
ABLIKIM	23O	PR D107 032007	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62058
ABLIKIM	23W	PRL 130 151904	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62068
HIRATA	23	PR D107 112011	H. Hirata <i>et al.</i>	(BELLE Collab.)	REFID=62281
YIN	23	PR D107 052004	J.H. Yin <i>et al.</i>	(BELLE Collab.)	REFID=61909
ABLIKIM	22D	PR D105 072009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61638
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)	REFID=61634
TERAMOTO	21	PRL 126 122001	Y. Teramoto <i>et al.</i>	(BELLE Collab.)	REFID=61098
AAIJ	20AD	PR D102 092005	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60673
AAIJ	20S	JHEP 2008 123	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60526
ABLIKIM	20W	PRL 124 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60570
LEES	20C	PRL 124 152001	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=60554
PDG	20	PTEP 2020 083C01	P.A. Zyla <i>et al.</i>	(PDG Collab.)	REFID=60676
ABLIKIM	19U	PRL 122 202001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59786
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=59884
AGHASYAN	18A	PL B783 334	M. Aghasyan <i>et al.</i>	(COMPASS Collab.)	REFID=59036
AAIJ	17AD	PL B769 305	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57896
AAIJ	17BB	EPJ C77 609	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=58191
BHARDWAJ	16	PR D93 052016	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=57272
AAIJ	15AO	PR D92 011102	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56771
ABLIKIM	15V	PL B749 414	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56787
BALA	15	PR D91 051101	A. Bala <i>et al.</i>	(BELLE Collab.)	REFID=56408
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (err.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)	REFID=57795
AAIJ	14AH	NP B886 665	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55897
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55647
IWASHITA	14	PTEP 2014 043C01	T. Iwashita <i>et al.</i>	(BELLE Collab.)	REFID=55925
AAIJ	13Q	PRL 110 222001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=54985
AAIJ	13S	EPJ C73 2462	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55008
BHARDWAJ	13	PRL 111 032001	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=55412
AAIJ	12H	EPJ C72 1972	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54056
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54751
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
BHARDWAJ	11	PRL 107 091803	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)	REFID=53779
CHOI	11	PR D84 052004	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=53934
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=53939
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)	REFID=53225
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53362
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545
AALTONEN	09AU	PRL 103 152001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53098
AUBERT	09B	PRL 102 132001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52722
AUBERT	08B	PR D77 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52120
AUBERT	08Y	PR D77 111101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52265
WICHT	08	PL B662 323	J. Wicht <i>et al.</i>	(BELLE Collab.)	REFID=52204
AUBERT	06	PR D73 011101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51017
AUBERT_BE	06M	PR D74 071101	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51454
GOKHROO	06	PRL 97 162002	G. Gokhroo <i>et al.</i>	(BELLE Collab.)	REFID=51432
AUBERT	05B	PR D71 031501	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50498
AUBERT	05D	PR D71 052001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50509
AUBERT	05R	PR D71 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50627
DOBBS	05	PRL 94 032004	S. Dobbs <i>et al.</i>	(CLEO Collab.)	REFID=50458
ABAZOV	04F	PRL 93 162002	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=50200
ACOSTA	04	PRL 93 072001	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=49742
AUBERT	04Y	PRL 93 041801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=49997
CHISTOV	04	PRL 93 051803	R. Chistov <i>et al.</i>	(BELLE Collab.)	REFID=50002
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)	REFID=49653
YUAN	04	PL B579 74	C.Z. Yuan <i>et al.</i>		REFID=49677
CHOI	03	PRL 91 262001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=49628
BAI	98E	PR D57 3854	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=46339
ANTONIAZZI	94	PR D50 4258	L. Antoniazzi <i>et al.</i>	(E705 Collab.)	REFID=44074

NODE=M159

$\chi_{c0}(3915)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

was $X(3915)$

The $\chi_{c0}(3915)$ was originally seen by BELLE in its $\omega J/\psi$ decay mode and was produced in both B decays in CHOI 05 and $\gamma\gamma$ collisions in UEHARA 10. The J^{PC} was determined to be 0^{++} by BABAR in LEES 12AD but this assignment was questioned by ZHOU 15C. In AAIJ 20AI LHCb found the $D^+ D^-$ decay mode of the $\chi_{c0}(3915)$ using B decays and determined its J^{PC} to be 0^{++} . Based on their compatible mass, width, and J^{PC} , we assume the state decaying to $\omega J/\psi$ and the state decaying to $D^+ D^-$ are both the $\chi_{c0}(3915)$. See also the $\chi_{c2}(3930)$.

NODE=M159

$\chi_{c0}(3915)$ MASS

NODE=M159M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3922.1 ± 1.8 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
[3921.7 ± 1.8 MeV OUR 2023 AVERAGE		Scale factor = 1.5]		
3956 ± 5 ± 10	360	¹ AAIJ	23AA LHCb	$B^+ \rightarrow D_s^+ D_s^- K^+$
3923.8 ± 1.5 ± 0.4	1.2k	² AAIJ	20AI LHCb	$B^+ \rightarrow D^+ D^- K^+$
3919.4 ± 2.2 ± 1.6	59 ± 10	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
3919.1 ⁺ ₋ $\frac{3.8}{3.4} \pm 2.0$		DEL-AMO-SA..10B	BABR	$B \rightarrow \omega J/\psi K$
3915 ± 3 ± 2	49 ± 15	UEHARA	10 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
3943 ± 11 ± 13	58 ± 11	³ CHOI	05 BELL	$B \rightarrow \omega J/\psi K$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
3922.4 ± 6.5 ± 2.0		⁴ WANG	22A BELL	$\gamma\gamma \rightarrow \gamma\psi(2S)$
3926.4 ± 2.2 ± 1.2		⁵ ABLIKIM	19v BES	$e^+ e^- \rightarrow \gamma\omega J/\psi$
3914.6 ⁺ ₋ $\frac{3.8}{3.4} \pm 2.0$		³ AUBERT	08W BABR	Superseded by DEL-AMO-SANCHEZ 10B

NODE=M159M
NEW

- ¹ $D_s^+ D_s^-$ near-threshold enhancement parameterized with a Flatte-like function .
- ² Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape.
- ³ $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.
- ⁴ Not distinguished from the $\chi_{c2}(3930)$.
- ⁵ Could also be $X(3940)$. Significance 3.1σ . Fit with additional resonance at 3963.7 ± 5.7 MeV, significance 3.4σ .

NODE=M159M;LINKAGE=D
NODE=M159M;LINKAGE=B

NODE=M159M;LINKAGE=CH
NODE=M159M;LINKAGE=C
NODE=M159M;LINKAGE=A



$\chi_{c0}(3915)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
20 ± 4 OUR AVERAGE		Error includes scale factor of 1.1. [18.8 ± 3.5 MeV OUR 2023 AVERAGE]		
43 ±13 ± 8	360	¹ AAIJ	23AA LHCb	$B^+ \rightarrow D_s^+ D_s^- K^+$
17.4 ± 5.1 ± 0.8	1.2k	² AAIJ	20Al LHCb	$B^+ \rightarrow D^+ D^- K^+$
13 ± 6 ± 3	59	LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
31 $\frac{+10}{-8}$ ± 5		DEL-AMO-SA...10B	BABR	$B \rightarrow \omega J/\psi K$
17 ±10 ± 3	49	UEHARA	10 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
87 ±22 ±26	58	³ CHOI	05 BELL	$B \rightarrow \omega J/\psi K$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
22 ±17 ± 4		⁴ WANG	22A BELL	$\gamma\gamma \rightarrow \gamma\psi(2S)$
3.8 ± 7.5 ± 2.6		⁵ ABLIKIM	19V BES	$e^+ e^- \rightarrow \gamma\omega J/\psi$
34 $\frac{+12}{-8}$ ± 5		³ AUBERT	08W BABR	Superseded by DEL-AMO-SANCHEZ 10B

¹ $D_s^+ D_s^-$ near-threshold enhancement parameterized with a Flatté-like function .

² Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape.

³ $\omega J/\psi$ threshold enhancement fitted as an S-wave Breit-Wigner resonance.

⁴ Not distinguished from the $\chi_{c2}(3930)$.

⁵ Could also be $X(3940)$. Significance 3.1σ . Fit with additional resonance at 3963.7 ± 5.7 MeV, significance 3.4σ .

NODE=M159W

NODE=M159W

NEW

NODE=M159W;LINKAGE=D
NODE=M159W;LINKAGE=BNODE=M159W;LINKAGE=CH
NODE=M159W;LINKAGE=C
NODE=M159W;LINKAGE=A $\chi_{c0}(3915)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega J/\psi$	seen
Γ_2 $\bar{D}^{*0} D^0$	not seen
Γ_3 $D^+ D^-$	seen
Γ_4 $D_s^+ D_s^-$	
Γ_5 $\pi^+ \pi^- \eta_c(1S)$	not seen
Γ_6 $\eta_c \eta$	not seen
Γ_7 $\eta_c \pi^0$	not seen
Γ_8 $K \bar{K}$	not seen
Γ_9 $\gamma\gamma$	seen
Γ_{10} $\gamma\psi(2S)$	
Γ_{11} $\pi^0 \chi_{c1}$	not seen

NODE=M159215;NODE=M159

DESIG=1;OUR EST;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=9
DESIG=11
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=6
DESIG=7
DESIG=5;OUR EVAL;→ UNCHECKED ←
DESIG=2
DESIG=10
DESIG=8 $\chi_{c0}(3915)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\omega J/\psi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_1 \Gamma_9/\Gamma$
VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	
54 ± 9 OUR AVERAGE					
52 ±10 ±3	59 ± 10	¹ LEES	12AD BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$	
61 ±17 ±8	49 ± 15	¹ UEHARA	10 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
18 ± 5 ±2	49 ± 15	² UEHARA	10 BELL	10.6 $e^+ e^- \rightarrow e^+ e^- \omega J/\psi$	
¹ For $J^P = 0^+$.					
² For $J^P = 2^+$, helicity-2.					

NODE=M159220

NODE=M159G01
NODE=M159G01

OCCUR=2

NODE=M159G01;LINKAGE=UH
NODE=M159G01;LINKAGE=UR

$\Gamma(\gamma\psi(2S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_{10} \Gamma_9/\Gamma$
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
9.8 ±3.6 ±1.3		¹ WANG	22A BELL	$\gamma\gamma \rightarrow \gamma\psi(2S)$	
¹ Not distinguished from the $\chi_{c2}(3930)$.					

NODE=M159R07
NODE=M159R07

NODE=M159R07;LINKAGE=A

$\Gamma(\pi^+ \pi^- \eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_5 \Gamma_9/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<16	90	LEES	12AE BABR	$e^+ e^- \rightarrow e^+ e^- \pi^+ \pi^- \eta_c$	

NODE=M159G02
NODE=M159G02

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_9/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<1.96	90	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M159G03
NODE=M159G03

$\chi_{c0}(3915)$ BRANCHING RATIOS

NODE=M159225

$\Gamma(\omega J/\psi)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		¹ DEL-AMO-SA...10B	BABR	$B \rightarrow \omega J/\psi K$	NODE=M159R03
seen		² CHOI	05	BELL	$B \rightarrow \omega J/\psi K$

NODE=M159R03
NODE=M159R03

¹ DEL-AMO-SANCHEZ 10B reports $B(B^{\pm} \rightarrow \chi_{c0}(3915) K^{\pm}) \times B(\chi_{c0}(3915) \rightarrow J/\psi \omega)$
 $= (3.0^{+0.7+0.5}_{-0.6-0.3}) \times 10^{-5}$ and $B(B^0 \rightarrow \chi_{c0}(3915) K^0) \times B(\chi_{c0}(3915) \rightarrow J/\psi \omega)$
 $= (2.1 \pm 0.9 \pm 0.3) \times 10^{-5}$.

NODE=M159R03;LINKAGE=DE

² CHOI 05 reports $B(B \rightarrow \chi_{c0}(3915) K) \times B(\chi_{c0}(3915) \rightarrow J/\psi \omega) = (7.1 \pm 1.3 \pm 3.1) \times 10^{-5}$.

NODE=M159R03;LINKAGE=CH

$\Gamma(\omega J/\psi)/\Gamma(\bar{D}^{*0} D^0)$					Γ_1/Γ_2
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
>0.71	90	¹ AUSHEV	10	BELL	$B \rightarrow \bar{D}^{*0} D^0 K$

NODE=M159R02
NODE=M159R02

¹ By combining the upper limit $B(B \rightarrow \chi_{c0}(3915) K) \times B(\chi_{c0}(3915) \rightarrow D^{*0} \bar{D}^0)$
 $< 0.67 \times 10^{-4}$ from AUSHEV 10 with the average of CHOI 05 and AUBERT 08w
measurements $B(B \rightarrow \chi_{c0}(3915) K) \times B(\chi_{c0}(3915) \rightarrow \omega J/\psi) = (0.51 \pm 0.11) \times 10^{-4}$.

NODE=M159R02;LINKAGE=AU

$\Gamma(D^+ D^-)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE		DOCUMENT ID	TECN	COMMENT	
seen		AAIJ	20AI	LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M159R06
NODE=M159R06

$\Gamma(D^+ D^-)/\Gamma(D_s^+ D_s^-)$					Γ_3/Γ_4
VALUE		DOCUMENT ID	TECN	COMMENT	
$0.29 \pm 0.09 \pm 0.10 \pm 0.08$		¹ AAIJ	23AA	LHCB	$B^+ \rightarrow D_s^+ D_s^- K^+$

NODE=M159R08
NODE=M159R08

¹ Assuming that AAIJ 20AI reporting on $B^+ \rightarrow D^+ D^- K^+$ also refers to $\chi_{c0}(3915)$.
The last uncertainty is due to the values of $B(D^+ \rightarrow K^- \pi^+ \pi^+)$ and $B(D_s^+ \rightarrow K^- K^+ \pi^+)$ from PDG 22.

NODE=M159R08;LINKAGE=A

$\Gamma(\eta_c \eta)/\Gamma_{\text{total}}$					Γ_6/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
not seen	90	¹ VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \eta$	NODE=M159R00

NODE=M159R00
NODE=M159R00

OCCUR=2

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ \chi_{c0}(3915)) \times B(\chi_{c0}(3915) \rightarrow \eta_c \eta) < 4.7 \times 10^{-5}$ at 90% CL.

NODE=M159R00;LINKAGE=A

$\Gamma(\eta_c \pi^0)/\Gamma_{\text{total}}$					Γ_7/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
not seen	90	¹ VINOKUROVA 15	BELL	$B^+ \rightarrow K^+ \eta_c \pi^0$	NODE=M159R04

NODE=M159R04
NODE=M159R04

OCCUR=2

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ \chi_{c0}(3915)^0) \times B(\chi_{c0}(3915) \rightarrow \eta_c \pi^0) < 1.7 \times 10^{-5}$ at 90% CL.

NODE=M159R04;LINKAGE=A

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	59 ± 10	LEES	12AD	BABR	$e^+ e^- \rightarrow e^+ e^- \omega J/\psi$
seen		UEHARA	10	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \omega J/\psi$

NODE=M159R01
NODE=M159R01

$\Gamma(\pi^0 \chi_{c1})/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
not seen	42 ± 14	¹ BHARDWAJ 19	BELL	$B^{\pm} \rightarrow \chi_{c1} \pi^0 K^{\pm}$	NODE=M159R05

NODE=M159R05
NODE=M159R05

¹ BHARDWAJ 19 reports $B(B^+ \rightarrow K^+ \chi_{c0}(3915)) \times B(\chi_{c0}(3915) \rightarrow \chi_{c1} \pi^0) < 3.8 \times 10^{-5}$ at 90% CL. A signal significance 2.3 standard deviations.

NODE=M159R05;LINKAGE=A

$\chi_{c0}(3915)$ REFERENCES

AAIJ	23AA	PRL 131 071901	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	22	PTEP 2022 083C01	R.L. Workman <i>et al.</i>	(PDG Collab.)
WANG	22A	PR D105 112011	X.L. Wang <i>et al.</i>	(BELLE Collab.)
AAIJ	20AI	PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.) JPC
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
BHARDWAJ	19	PR D99 111101	V. Bhardwaj <i>et al.</i>	(BELLE Collab.)
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also		JHEP 1702 088 (errat.)	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
ZHOU	15C	PRL 115 022001	Z.-Y. Zhou, Z. Xiao, H.-Q. Zhou	(BEIJT, NANJ)
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)
LEES	12AD	PR D86 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)
AUSHEV	10	PR D81 031103	T. Aushev <i>et al.</i>	(BELLE Collab.)
DEL-AMO-SA...	10B	PR D82 011101	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)
UEHARA	10	PRL 104 092001	S. Uehara <i>et al.</i>	(BELLE Collab.)
AUBERT	08W	PRL 101 082001	B. Aubert <i>et al.</i>	(BABAR Collab.)
CHOI	05	PRL 94 182002	S.-K. Choi <i>et al.</i>	(BELLE Collab.)

NODE=M159

REFID=62314
REFID=61634
REFID=61640
REFID=60739
REFID=59796
REFID=59884
REFID=56706
REFID=57795
REFID=56842
REFID=55592
REFID=54751
REFID=54752
REFID=53225
REFID=53362
REFID=53232
REFID=52263
REFID=50737

$\chi_{c2}(3930)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

NODE=M050

$\chi_{c2}(3930)$ MASS

NODE=M050M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3922.5 ± 1.0 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
3926.8 ± 2.4 ± 0.8	1.2k	¹ AAIJ	20AI LHCb	$B^+ \rightarrow D^+ D^- K^+$
3921.9 ± 0.6 ± 0.2		² AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$
3926.7 ± 2.7 ± 1.1	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
3929 ± 5 ± 2	64	UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$


NODE=M050M

¹ Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Previous measurements assumed a single state in this region. This analysis revealed the presence of $\chi_{c0}(3930)$ with the same mass.

NODE=M050M;LINKAGE=B

² Measured in prompt hadroproduction.

NODE=M050M;LINKAGE=A



$\chi_{c2}(3930)$ WIDTH

NODE=M050W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.2 ± 2.2 OUR AVERAGE		Error includes scale factor of 1.2.		
34.2 ± 6.6 ± 1.1	1.2k	¹ AAIJ	20AI LHCb	$B^+ \rightarrow D^+ D^- K^+$
36.6 ± 1.9 ± 0.9		² AAIJ	19M LHCb	$pp \rightarrow D\bar{D} + \text{anything}$
21.3 ± 6.8 ± 3.6	76 ± 17	AUBERT	10G BABR	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$
29 ± 10 ± 2	64	UEHARA	06 BELL	$10.6 e^+ e^- \rightarrow e^+ e^- D\bar{D}$

NODE=M050W

¹ Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Previous measurements assumed a single state in this region. This analysis revealed the presence of $\chi_{c0}(3930)$ with the same mass.

NODE=M050W;LINKAGE=B

² Measured in prompt hadroproduction.

NODE=M050W;LINKAGE=A

$\chi_{c2}(3930)$ DECAY MODES

NODE=M050215;NODE=M050

Mode	Fraction (Γ_i/Γ)
Γ_1 $\gamma\gamma$	seen
Γ_2 $K\bar{K}\pi$	
Γ_3 $K^+K^-\pi^+\pi^-\pi^0$	
Γ_4 $D\bar{D}$	seen
Γ_5 D^+D^-	seen
Γ_6 $D^0\bar{D}^0$	seen
Γ_7 $\pi^+\pi^-\eta_c(1S)$	not seen
Γ_8 $K\bar{K}$	not seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=5
DESIG=6
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4;OUR EVAL;→ UNCHECKED ←
DESIG=7;OUR EVAL;→ UNCHECKED ←
DESIG=8;OUR EVAL;→ UNCHECKED ←

 $\chi_{c2}(3930)$ PARTIAL WIDTHS

NODE=M050220

 $\chi_{c2}(3930)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M050222

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.1	90	DEL-AMO-SA...11M	BABR	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	

NODE=M050G01
NODE=M050G01

$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<3.4	90	DEL-AMO-SA...11M	BABR	$\gamma\gamma \rightarrow K^+K^-\pi^+\pi^-\pi^0$	

NODE=M050G02
NODE=M050G02

$\Gamma(D\bar{D}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_1/\Gamma$
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.21±0.04 OUR AVERAGE					
0.24±0.05±0.04	76 ± 17	AUBERT	10G	BABR	10.6 $e^+e^- \rightarrow e^+e^-D\bar{D}$
0.18±0.05±0.03	64	¹ UEHARA	06	BELL	10.6 $e^+e^- \rightarrow e^+e^-D\bar{D}$
¹ Assuming $B(D^+D^-) = 0.89 B(D^0\bar{D}^0)$.					

NODE=M050G1
NODE=M050G1

NODE=M050G1;LINKAGE=UE

$\Gamma(\pi^+\pi^-\eta_c(1S)) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_7\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<18	90	LEES	12AE	BABR	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\eta_c$

NODE=M050G03
NODE=M050G03

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$					$\Gamma_8\Gamma_1/\Gamma$
VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	
<0.256	90	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M050G04
NODE=M050G04

 $\chi_{c2}(3930)$ BRANCHING RATIOS

NODE=M050225

$\Gamma(D^+D^-)/\Gamma(D^0\bar{D}^0)$					Γ_5/Γ_6
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.74±0.43±0.16	64	UEHARA	06	BELL	10.6 $e^+e^- \rightarrow e^+e^-D\bar{D}$

NODE=M050R01
NODE=M050R01

 $\chi_{c2}(3930)$ REFERENCES

NODE=M050

AAIJ	20AI	PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60739
AAIJ	19M	JHEP 1907 035	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59697
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
LEES	12AE	PR D86 092005	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54752
DEL-AMO-SA...	11M	PR D84 012004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16751
AUBERT	10G	PR D81 092003	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53357
UEHARA	06	PRL 96 082003	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=51039

X(3940)

$$I^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Reported by ABE 07, observed in $e^+e^- \rightarrow J/\psi X$.

NODE=M029

NODE=M029

NODE=M029M

NODE=M029M

X(3940) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$3942^{+7}_{-6} \pm 6$	52	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$3943 \pm 6 \pm 6$	25	¹ ABE	07	BELL $e^+e^- \rightarrow J/\psi X$
3936 ± 14	266	² ABE	07	BELL $e^+e^- \rightarrow J/\psi(c\bar{c})$
¹ From a fit to $D^{*+}D^-$ and $D^{*0}\bar{D}^0$ events.				
² From the inclusive fit. Not independent of the exclusive measurement by ABE 07.				

OCCUR=2

NODE=M029M;LINKAGE=EB

NODE=M029M;LINKAGE=EM

X(3940) WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$37^{+26}_{-15} \pm 8$		52	PAKHLOV 08	BELL	$e^+e^- \rightarrow J/\psi X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<52	90	25	ABE	07	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M029W

NODE=M029W

X(3940) DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\bar{D}^* + c.c.$	seen
Γ_2 $D\bar{D}$	not seen
Γ_3 $J/\psi\omega$	not seen

NODE=M029215;NODE=M029

DESIG=1;OUR EVAL;→ UNCHECKED ←

DESIG=2;OUR EVAL;→ UNCHECKED ←

DESIG=3;OUR EVAL;→ UNCHECKED ←

X(3940) BRANCHING RATIOS

$\Gamma(D\bar{D}^* + c.c.)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.45	90	25	^{1,2} ABE	07	BELL $e^+e^- \rightarrow J/\psi X$
¹ For X(3940) decaying to final states with more than two tracks.					
² PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c ² may consist of several states.					

NODE=M029225

NODE=M029R01

NODE=M029R01

NODE=M029R01;LINKAGE=AB

NODE=M029R01;LINKAGE=AE

$\Gamma(D\bar{D})/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.41	90	^{1,2} ABE	07	BELL $e^+e^- \rightarrow J/\psi X$	
¹ For X(3940) decaying to final states with more than two tracks.					
² PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c ² may consist of several states.					

NODE=M029R02

NODE=M029R02

NODE=M029R02;LINKAGE=AB

NODE=M029R02;LINKAGE=AE

$\Gamma(J/\psi\omega)/\Gamma_{\text{total}}$					Γ_3/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.26	90	^{1,2} ABE	07	BELL $e^+e^- \rightarrow J/\psi X$	
¹ For X(3940) decaying to final states with more than two tracks.					
² PAKHLOV 08 finds that the inclusive peak near 3940 MeV/c ² may consist of several states.					

NODE=M029R03

NODE=M029R03

NODE=M029R03;LINKAGE=AB

NODE=M029R03;LINKAGE=AE

X(3940) REFERENCES

PAKHLOV 08	PRL 100 202001	P. Pakhlov <i>et al.</i>	(BELLE Collab.)
ABE 07	PRL 98 082001	K. Abe <i>et al.</i>	(BELLE Collab.)

NODE=M029

REFID=52302

REFID=51627

$\psi(4040)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M072

 $\psi(4040)$ MASS

NODE=M072M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4039 ± 1 OUR ESTIMATE			
4039.6 ± 4.3	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4034 ± 6	² MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4037 ± 2	³ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4040 ± 1	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4040 ± 10	BRANDELIK	78C DASP	e^+e^-
¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
³ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁴ From a fit to BES (BAI 02C) data.			

NODE=M072M
→ UNCHECKED ←

OCCUR=2

NODE=M072M;LINKAGE=AB

NODE=M072M;LINKAGE=MO

NODE=M072M;LINKAGE=ST
NODE=M072M;LINKAGE=SE $\psi(4040)$ WIDTH

NODE=M072W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 ± 10 OUR ESTIMATE			
84.5 ± 12.3	⁵ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87 ± 11	⁶ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
85 ± 10	⁷ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
89 ± 6	⁸ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
52 ± 10	BRANDELIK	78C DASP	e^+e^-
⁵ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (130 \pm 46)^\circ$.			
⁶ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁷ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁸ From a fit to BES (BAI 02C) data.			

NODE=M072W
→ UNCHECKED ←

OCCUR=2

NODE=M072W;LINKAGE=AB

NODE=M072W;LINKAGE=MO

NODE=M072W;LINKAGE=ST
NODE=M072W;LINKAGE=SE $\psi(4040)$ DECAY MODES

NODE=M072215;NODE=M072

NODE=M072

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 e^+e^-	$(1.07 \pm 0.16) \times 10^{-5}$	DESIG=5
Γ_2 $D\bar{D}$	seen	DESIG=17;OUR EST;→ UNCHECKED ←
Γ_3 $D^0\bar{D}^0$	seen	DESIG=1
Γ_4 D^+D^-	seen	DESIG=18
Γ_5 $D^*\bar{D} + \text{c.c.}$	seen	DESIG=19;OUR EST;→ UNCHECKED ←
Γ_6 $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen	DESIG=2
Γ_7 $D^*(2010)^+D^- + \text{c.c.}$	seen	DESIG=20
Γ_8 $D^*\bar{D}^*$	seen	DESIG=21;OUR EST;→ UNCHECKED ←
Γ_9 $D^*(2007)^0\bar{D}^*(2007)^0$	seen	DESIG=3
Γ_{10} $D^*(2010)^+D^*(2010)^-$	seen	DESIG=22
Γ_{11} $D\bar{D}\pi$ (excl. $D^*\bar{D}$)		DESIG=23
Γ_{12} $D^0D^-\pi^+ + \text{c.c.}$ (excl. $D^*(2007)^0\bar{D}^0 + \text{c.c.}$, $D^*(2010)^+D^- + \text{c.c.}$)	not seen	DESIG=24

Γ ₁₃	$D\bar{D}^*\pi$ (excl. $D^*\bar{D}^*$)	not seen			DESIG=25
Γ ₁₄	$D^0\bar{D}^{*-}\pi^+ + \text{c.c.}$ (excl. $D^*(2010)^+D^*(2010)^-$)	seen			DESIG=26
Γ ₁₅	$D_s^+D_s^-$	seen			DESIG=27
Γ ₁₆	$\pi^+\pi^+\pi^-\pi^-\pi^0$				DESIG=37
Γ ₁₇	$J/\psi(1S)$ hadrons				DESIG=4
Γ ₁₈	$J/\psi\pi^+\pi^-$	< 4	$\times 10^{-3}$	90%	DESIG=7
Γ ₁₉	$J/\psi\pi^0\pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=8
Γ ₂₀	$J/\psi\eta$	(5.2 ± 0.7)	$\times 10^{-3}$		DESIG=9
Γ ₂₁	$J/\psi\pi^0$	< 2.8	$\times 10^{-4}$	90%	DESIG=10
Γ ₂₂	$J/\psi\pi^+\pi^-\pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=11
Γ ₂₃	$\chi_{c1}\gamma$	< 3.4	$\times 10^{-3}$	90%	DESIG=12
Γ ₂₄	$\chi_{c2}\gamma$	< 5	$\times 10^{-3}$	90%	DESIG=13
Γ ₂₅	$\chi_{c1}\pi^+\pi^-\pi^0$	< 1.1	%	90%	DESIG=14
Γ ₂₆	$\chi_{c2}\pi^+\pi^-\pi^0$	< 3.2	%	90%	DESIG=15
Γ ₂₇	$h_c(1P)\pi^+\pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=28
Γ ₂₈	$\phi\pi^+\pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=16
Γ ₂₉	$\Lambda\bar{\Lambda}\pi^+\pi^-$	< 2.9	$\times 10^{-4}$	90%	DESIG=29
Γ ₃₀	$\Lambda\bar{\Lambda}\pi^0$	< 9	$\times 10^{-5}$	90%	DESIG=30
Γ ₃₁	$\Lambda\bar{\Lambda}\eta$	< 3.0	$\times 10^{-4}$	90%	DESIG=31
Γ ₃₂	$\Lambda\bar{\Lambda}$	< 6	$\times 10^{-6}$	90%	DESIG=36
Γ ₃₃	$\Sigma^+\bar{\Sigma}^-$	< 1.3	$\times 10^{-4}$	90%	DESIG=32
Γ ₃₄	$\Sigma^0\bar{\Sigma}^0$	< 7	$\times 10^{-5}$	90%	DESIG=33
Γ ₃₅	$\Xi^+\bar{\Xi}^-$	< 1.6	$\times 10^{-4}$	90%	DESIG=34
Γ ₃₆	$\Xi^0\bar{\Xi}^0$	< 1.8	$\times 10^{-4}$	90%	DESIG=35
Γ ₃₇	$\Xi^-\bar{\Xi}^+$	< 6	$\times 10^{-5}$	90%	DESIG=38
Γ ₃₈	$\mu^+\mu^-$	(9 ± 6)	$\times 10^{-6}$		DESIG=6

ψ(4040) PARTIAL WIDTHS

NODE=M072220

Γ(e⁺e⁻)**Γ₁**

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

0.86±0.07 OUR ESTIMATE**0.83±0.20**⁹ ABLIKIM 08D BES2 e⁺e⁻ → hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.6 to 1.4

¹⁰ MO 10 RVUE e⁺e⁻ → hadrons

0.88±0.11

¹¹ SETH 05A RVUE e⁺e⁻ → hadrons

0.91±0.13

¹² SETH 05A RVUE e⁺e⁻ → hadrons

0.75±0.15

BRANDELIK 78C DASP e⁺e⁻NODE=M072W5
NODE=M072W5

→ UNCHECKED ←

OCCUR=2

⁹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the ψ(3770), ψ(4040), ψ(4160), and ψ(4415) resonances. Phase angle fixed in the fit to δ = (130 ± 46)^o.

NODE=M072W5;LINKAGE=AB

¹⁰ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the ψ(4040), ψ(4160) and ψ(4415) resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e⁺e⁻ partial widths. We quote only the range of values.

NODE=M072W5;LINKAGE=MO

¹¹ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M072W5;LINKAGE=ST

¹² From a fit to BES (BAI 02C) data.

NODE=M072W5;LINKAGE=SE

Γ(μ⁺μ⁻)**Γ₃₈**

VALUE (keV)

DOCUMENT ID

TECN

COMMENT

0.73±0.48±0.12^{13,14} ABLIKIM 20AG BES3 e⁺e⁻ → μ⁺μ⁻¹³ From a fit to the e⁺e⁻ → μ⁺μ⁻ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming Γ(μ⁺μ⁻) = Γ(e⁺e⁻).

NODE=M072W2;LINKAGE=A

¹⁴ From solution 1 of 8 with equal fit quality. Other solutions range from 0.58 ± 0.52 ± 0.10 to 0.80 ± 0.48 ± 0.13 keV.

NODE=M072W2;LINKAGE=B

$\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

NODE=M072235

 $\Gamma(J/\psi\eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{20}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M072R00
NODE=M072R00

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5±0.3	15 ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
1.4±0.3	16 ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
7.0±0.6	17 ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$

OCCUR=2
OCCUR=3

15 Solution 1 of three equivalent fit solutions using three resonant structures.

16 Solution 2 of three equivalent fit solutions using three resonant structures.

17 Solution 3 of three equivalent fit solutions using three resonant structures.

NODE=M072R00;LINKAGE=A
NODE=M072R00;LINKAGE=B
NODE=M072R00;LINKAGE=C $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M072G01
NODE=M072G01

<2.9	90	18 HAN	15 BELL	10.58 $e^+ e^- \rightarrow \chi_{c1}\gamma$
------	----	--------	---------	---

18 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M072G01;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M072G02
NODE=M072G02

<4.6	90	19 HAN	15 BELL	10.58 $e^+ e^- \rightarrow \chi_{c2}\gamma$
------	----	--------	---------	---

19 Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M072G02;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{32}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M072R33
NODE=M072R33

<5.5 × 10 ⁻³	90	20 ABLIKIM	21AS BES3	$e^+ e^- \rightarrow \psi(4040)$
-------------------------	----	------------	-----------	----------------------------------

20 From a measurement of the $e^+ e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

NODE=M072R33;LINKAGE=A

 $\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{37}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M072R35
NODE=M072R35

<0.0519	90	21 ABLIKIM	23BK BES3	$e^+ e^- \rightarrow \psi(4040)$
---------	----	------------	-----------	----------------------------------

21 From a fit to $e^+ e^- \rightarrow \Xi^-\bar{\Xi}^+$ cross sections.

NODE=M072R35;LINKAGE=A

 $\psi(4040) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma^2(\text{total})$

NODE=M072230

 $\Gamma(J/\psi\eta) / \Gamma_{\text{total}} \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$

VALUE (units 10 ⁻⁸)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

NODE=M072R25
NODE=M072R25

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.1±1.4±1.5	22 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$
12.8±2.1±1.9	23 WANG	13B BELL	$e^+ e^- \rightarrow J/\psi\eta\gamma$

OCCUR=2

22 Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

NODE=M072R25;LINKAGE=A

23 Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4039 MeV and 80 MeV, respectively.

NODE=M072R25;LINKAGE=B

 $\psi(4040) \text{ BRANCHING RATIOS}$

NODE=M072225

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10 ⁻⁵)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

NODE=M072R4
NODE=M072R4

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 1.0	FELDMAN	77 MRK1	$e^+ e^-$
-------	---------	---------	-----------

 $\Gamma(D^0\bar{D}^0) / \Gamma_{\text{total}}$ Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M072R14
NODE=M072R14

seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$
------	--------	----------	--

seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^0\bar{D}^0$
------	----------------	------	------------------------------------

seen	PAKHLOVA	08 BELL	$e^+ e^- \rightarrow D^0\bar{D}^0\gamma$
------	----------	---------	--

 $\Gamma(D^+ D^-) / \Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M072R15
NODE=M072R15

seen	AUBERT	09M BABR	$e^+ e^- \rightarrow D^+ D^- \gamma$
------	--------	----------	--------------------------------------

seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^+ D^-$
------	----------------	------	-------------------------------

seen	PAKHLOVA	08 BELL	$e^+ e^- \rightarrow D^+ D^- \gamma$
------	----------	---------	--------------------------------------

$$\Gamma(D\bar{D})/\Gamma(D^*\bar{D} + \text{c.c.})$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_5
0.24±0.05±0.12	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}$

NODE=M072R12
NODE=M072R12

$$\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_6
0.05±0.03	²⁴ GOLDHABER	77	MRK1	e^+e^-

NODE=M072R1
NODE=M072R1

²⁴ Phase-space factor (p^3) explicitly removed.

NODE=M072R;LINKAGE=P

$$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*0}\bar{D}^0$

NODE=M072R16
NODE=M072R16

$$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
seen	²⁵ ZHUKOVA	18	BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*+}D^-$

NODE=M072R17
NODE=M072R17
OCCUR=3

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen PAKHLOVA 07 BELL $e^+e^- \rightarrow D^{*+}D^-\gamma$

²⁵ Supersedes PAKHLOVA 07.

NODE=M072R17;LINKAGE=C

$$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_6
0.95±0.09±0.10	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}$

NODE=M072R11
NODE=M072R11

$$\Gamma(D^*\bar{D}^*)/\Gamma(D^*\bar{D} + \text{c.c.})$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_5
0.18±0.14±0.03	AUBERT	09M	BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

NODE=M072R13
NODE=M072R13

$$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$

NODE=M072R18
NODE=M072R18

$$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_6
32.0±12.0	²⁶ GOLDHABER	77	MRK1	e^+e^-

NODE=M072R2
NODE=M072R2

²⁶ Phase-space factor (p^3) explicitly removed.

NODE=M072R2;LINKAGE=P

$$\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
seen	²⁷ ZHUKOVA	18	BELL	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
seen	AUBERT	09M	BABR	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$
seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D^{*+}D^{*-}$

NODE=M072R19
NODE=M072R19
OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen PAKHLOVA 07 BELL $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$

²⁷ Supersedes PAKHLOVA 07.

NODE=M072R19;LINKAGE=B

$$\Gamma(D^0D^-\pi^+ + \text{c.c. (excl. } D^*(2007)^0\bar{D}^0 + \text{c.c., } D^*(2010)^+D^- + \text{c.c.}))/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
not seen	PAKHLOVA	08A	BELL	$e^+e^- \rightarrow D^0D^-\pi^+\gamma$

NODE=M072R20
NODE=M072R20

$$\Gamma(D\bar{D}^*\pi(\text{excl. } D^*\bar{D}^*))/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
not seen	CRONIN-HEN..09	CLEO		$e^+e^- \rightarrow D\bar{D}^*\pi$

NODE=M072R21
NODE=M072R21

$$\Gamma(D^0\bar{D}^{*-}\pi^+ + \text{c.c. (excl. } D^*(2010)^+D^*(2010)^-))/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
seen	PAKHLOVA	09	BELL	$e^+e^- \rightarrow D^0D^{*-}\pi^+\gamma$

NODE=M072R22
NODE=M072R22

$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	NODE=M072R23 NODE=M072R23
seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$	

 $\Gamma(\pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				NODE=M072R34 NODE=M072R34
$(3.51 \pm 1.89 \pm 1.24) \times 10^{-5}$	²⁸ ABLIKIM	21AWBES3	$e^+ e^- \rightarrow 2\pi^+ 2\pi^- \pi^0$	
$(2.41 \pm 0.05 \pm 0.79) \times 10^{-2}$	²⁹ ABLIKIM	21AWBES3	$e^+ e^- \rightarrow 2\pi^+ 2\pi^- \pi^0$	OCCUR=2
²⁸ Solution 1 of two solutions with equal fit quality. The significance of the $\psi(4040)$ signal is 3.6σ .				NODE=M072R34;LINKAGE=A
²⁹ Solution 2 of two solutions with equal fit quality. The significance of the $\psi(4040)$ signal is 3.6σ .				NODE=M072R34;LINKAGE=B

 $\Gamma(J/\psi \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
<4	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	NODE=M072R01 NODE=M072R01

 $\Gamma(J/\psi \pi^0 \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{19}/Γ
<2	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	NODE=M072R02 NODE=M072R02

 $\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ
5.2 ± 0.5 ± 0.5		³⁰ ABLIKIM 12K	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$	NODE=M072R03 NODE=M072R03
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<7	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	
³⁰ ABLIKIM 12K measure $\sigma(e^+ e^- \rightarrow J/\psi \eta) = 32.1 \pm 2.8 \pm 1.3$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.					NODE=M072R03;LINKAGE=AB

 $\Gamma(J/\psi \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ
<0.28	90	³¹ ABLIKIM 12K	BES3	$e^+ e^- \rightarrow \ell^+ \ell^- 2\gamma$	NODE=M072R04 NODE=M072R04
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	
³¹ ABLIKIM 12K measure $\sigma(e^+ e^- \rightarrow J/\psi \pi^0) < 1.6$ pb. They assume the $\eta J/\psi$ fully originates from $\psi(4040)$ decays.					NODE=M072R04;LINKAGE=AB

 $\Gamma(J/\psi \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
<2	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	NODE=M072R05 NODE=M072R05

 $\Gamma(\chi_{c1} \gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					NODE=M072R06 NODE=M072R06
<11	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	

 $\Gamma(\chi_{c2} \gamma)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{24}/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •					NODE=M072R07 NODE=M072R07
<17	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	

 $\Gamma(\chi_{c1} \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{25}/Γ
<11	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	NODE=M072R08 NODE=M072R08

 $\Gamma(\chi_{c2} \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{26}/Γ
<32	90	COAN 06	CLEO	$3.97-4.06 e^+ e^- \rightarrow \text{hadrons}$	NODE=M072R09 NODE=M072R09

$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	32 PEDLAR	11 CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$

NODE=M072R24
 NODE=M072R24

³²From several values of \sqrt{s} near the peak of the $\psi(4040)$, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 1.0 \pm 8.0 \pm 5.4 \pm 0.2$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M072R24;LINKAGE=PE

 $\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	COAN	06 CLEO	$3.97-4.06 e^+e^- \rightarrow \text{hadrons}$

NODE=M072R10
 NODE=M072R10

 $\Gamma(\Lambda\bar{\Lambda}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.9	90	33 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R26
 NODE=M072R26

³³Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R26;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	34 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R27
 NODE=M072R27

³⁴Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R27;LINKAGE=A

 $\Gamma(\Lambda\bar{\Lambda}\eta)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	35 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R28
 NODE=M072R28

³⁵Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R28;LINKAGE=A

 $\Gamma(\Sigma^+\bar{\Sigma}^-)/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	36 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R29
 NODE=M072R29

³⁶Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R29;LINKAGE=A

 $\Gamma(\Sigma^0\bar{\Sigma}^0)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	37 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R30
 NODE=M072R30

³⁷Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R30;LINKAGE=A

 $\Gamma(\Xi^+\bar{\Xi}^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	38 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R31
 NODE=M072R31

³⁸Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R31;LINKAGE=A

 $\Gamma(\Xi^0\bar{\Xi}^0)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	39 ABLIKIM	13Q BES3	$e^+e^- \rightarrow \psi(4040)$

NODE=M072R32
 NODE=M072R32

³⁹Assuming that interference effects between resonance and continuum can be neglected.

NODE=M072R32;LINKAGE=A

 $\psi(4040)$ REFERENCES

ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62437
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21AW	PR D104 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61479
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	200	PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60344
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
ABLIKIM	13Q	PR D87 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55393
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
ABLIKIM	12K	PR D86 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54738
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724

NODE=M072

CRONIN-HEN...09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
PAKHLOVA 09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
ABLIKIM 08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)
PAKHLOVA 08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
PAKHLOVA 07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)
COAN 06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)
SETH 05A	PR D72 017501	K.K. Seth	
BAI 02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)
BAI 00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)
OSTERHELD 86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)
BRANDELIK 78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)
Also	ZPHY C1 233	R. Brandelik <i>et al.</i>	(DASP Collab.)
FELDMAN 77	PRPL 33C 285	G.J. Feldman, M.L. Perl	(LBL, SLAC)
GOLDHABER 77	PL 69B 503	G. Goldhaber <i>et al.</i>	(Mark I Collab.)

REFID=53114
 REFID=53143
 REFID=52142
 REFID=52132
 REFID=52134
 REFID=51628
 REFID=51075
 REFID=50813
 REFID=50506
 REFID=50503
 REFID=51064
 REFID=22232
 REFID=22114
 REFID=22062
 REFID=11434

NODE=M193

$\chi_{c1}(4140)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

was $X(4140)$

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

NODE=M193

Seen by AALTONEN 09AH, ABAZOV 14A, CHATRCHYAN 14M, AAIJ 17C in $B^+ \rightarrow \chi_{c1} K^+$, $\chi_{c1} \rightarrow J/\psi\phi$, and by ABAZOV 15M separately in both prompt (4.7σ) and non-prompt (5.6σ) production in $p\bar{p} \rightarrow J/\psi\phi + \text{anything}$. Not seen by SHEN 10 in $\gamma\gamma \rightarrow J/\psi\phi$ and ABLIKIM 15 in $e^+e^- \rightarrow \gamma J/\psi\phi$ at $\sqrt{s} = 4.23, 4.26, 4.36$ GeV.

$\chi_{c1}(4140)$ MASS

NODE=M193M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4146.5 ± 3.0 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
4118 ±11 ⁺¹⁹ / ₋₃₆	24k	1 AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
4143.4 ⁺ _{-3.0} ± 0.6	19	2 AALTONEN	17 CDF	$B^+ \rightarrow J/\psi\phi K^+$
4152.5 ± 1.7 ⁺ _{-5.4}	616	3 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi\phi + \text{anything}$
4159.0 ± 4.3 ± 6.6	52	4 ABAZOV	14A D0	$B^+ \rightarrow J/\psi\phi K^+$
4148.0 ± 2.4 ± 6.3	0.3k	5 CHATRCHYAN 14M	CMS	$B^+ \rightarrow J/\psi\phi K^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4146.5 ± 4.5 ⁺ _{-2.8}	4289	6,7 AAIJ	17C LHCB	$B^+ \rightarrow J/\psi\phi K^+$
4143.0 ± 2.9 ± 1.2	14	8,9 AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M193M

- ¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 13σ .
- ² Statistical significance of more than 5σ .
- ³ Statistical significance of more than 6σ .
- ⁴ Statistical significance of 3.1σ .
- ⁵ From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5σ .
- ⁶ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 8.4σ .
- ⁷ Superseded by AAIJ 21E.
- ⁸ Statistical significance of 3.8σ .
- ⁹ Superseded by AALTONEN 17.

NODE=M193M;LINKAGE=G
 NODE=M193M;LINKAGE=E
 NODE=M193M;LINKAGE=C
 NODE=M193M;LINKAGE=A
 NODE=M193M;LINKAGE=B

NODE=M193M;LINKAGE=D
 NODE=M193M;LINKAGE=H
 NODE=M193M;LINKAGE=AA
 NODE=M193M;LINKAGE=F



chi_c1(4140) WIDTH

NODE=M193W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
19 \pm 7 $\frac{+7}{-5}$ OUR AVERAGE				
162 \pm 21 $\frac{+24}{-49}$	24k	1 AAIJ	21E LHCb	$B^+ \rightarrow J/\psi \phi K^+$
15.3 $\frac{+10.4}{-6.1} \pm 2.5$	19	2 AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$
16.3 $\pm 5.6 \pm 11.4$	616	3 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi \phi + \text{anything}$
20 $\pm 13 \frac{+3}{-8}$	52	4 ABAZOV	14A D0	$B^+ \rightarrow J/\psi \phi K^+$
28 $\frac{+15}{-11} \pm 19$	0.3k	5 CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi \phi K^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
83 $\pm 21 \frac{+21}{-14}$	4289	6,7 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi \phi K^+$
11.7 $\frac{+8.3}{-5.0} \pm 3.7$	14	8,9 AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M193W

1 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 13 σ .
 2 Statistical significance of more than 5 σ .
 3 Statistical significance of more than 6 σ .
 4 Statistical significance of 3.1 σ .
 5 From a fit assuming an S-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5 σ .
 6 From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 8.4 σ .
 7 Superseded by AAIJ 21E.
 8 Statistical significance of 3.8 σ .
 9 Superseded by AALTONEN 17.

NODE=M193W;LINKAGE=G
 NODE=M193W;LINKAGE=E
 NODE=M193W;LINKAGE=C
 NODE=M193W;LINKAGE=A
 NODE=M193W;LINKAGE=B

NODE=M193W;LINKAGE=D
 NODE=M193W;LINKAGE=H
 NODE=M193W;LINKAGE=AA
 NODE=M193W;LINKAGE=F

chi_c1(4140) DECAY MODES

NODE=M193215;NODE=M193

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi \phi$	seen
Γ_2 $\gamma\gamma$	not seen

DESIG=1
 DESIG=2

chi_c1(4140) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M193220

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
<41	90	1 SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi \phi$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 6	90	2 SHEN	10 BELL	10.6 $e^+e^- \rightarrow e^+e^- J/\psi \phi$	
1 For $J^P = 0^+$. 2 For $J^P = 2^+$.					

NODE=M193G01
 NODE=M193G01

OCCUR=2

NODE=M193G01;LINKAGE=S0
 NODE=M193G01;LINKAGE=S2

$\chi_{c1}(4140)$ BRANCHING RATIOS

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$		DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	24k	1 AAIJ	21E LHCb	$B^+ \rightarrow J/\psi\phi K^+$	
seen	616	2 ABAZOV	15M D0	$p\bar{p} \rightarrow J/\psi\phi + \text{anything}$	
seen	52	3 ABAZOV	14A D0	$B^+ \rightarrow J/\psi\phi K^+$	
seen	0.3k	4 CHATRCHYAN	14M CMS	$B^+ \rightarrow J/\psi\phi K^+$	
seen	14	5 AALTONEN	09AH CDF	$B^+ \rightarrow J/\psi\phi K^+$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	4289	6,7 AAIJ	17C LHCb	$B^+ \rightarrow J/\psi\phi K^+$
not seen		8 ABLIKIM	15 BES3	$e^+e^- \rightarrow \gamma\phi J/\psi$
not seen		9 AAIJ	12AA LHCb	$pp \rightarrow B^+ X$ at 7 TeV

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 13 σ .

² Statistical significance of more than 6 σ .

³ ABAZOV 14A reports $B(B^+ \rightarrow \chi_{c1}(4140)K^+ \rightarrow J/\psi\phi K^+)/B(B^+ \rightarrow J/\psi\phi K^+) = (19 \pm 7 \pm 4)\%$ with 3.1 σ significance.

⁴ From a fit assuming an *S*-wave relativistic Breit-Wigner shape above a three-body phase-space non-resonant component with statistical significance of more than 5 σ .

⁵ Statistical significance of 3.8 σ .

⁶ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 8.4 σ .

⁷ Superseded by AAIJ 21E.

⁸ Reported $\sigma(e^+e^- \rightarrow \gamma\chi_{c1}(4140)) \cdot B(\chi_{c1}(4140) \rightarrow J/\psi\phi) < 0.35, 0.28,$ and 0.33 pb at 4.23, 4.26, and 4.36 GeV, respectively, at 90% CL.

⁹ Reported $B(B^+ \rightarrow \chi_{c1}(4140)K^+) \cdot B(\chi_{c1}(4140) \rightarrow J/\psi\phi)/B(B^+ \rightarrow J/\psi\phi K^+) < 0.07$ at 90% CL.

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$		DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
not seen		SHEN	10 BELL	$10.6 e^+e^- \rightarrow e^+e^- J/\psi\phi$	

 $\chi_{c1}(4140)$ REFERENCES

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636
AALTONEN	17	MPL A32 1750139	T. Altonen <i>et al.</i>	(CDF Collab.)	REFID=58161
ABAZOV	15M	PRL 115 232001	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=56957
ABLIKIM	15	PR D91 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56368
ABAZOV	14A	PR D89 012004	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55650
CHATRCHYAN	14M	PL B734 261	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55753
AAIJ	12AA	PR D85 091103	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=54263
SHEN	10	PRL 104 112004	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53235
AALTONEN	09AH	PRL 102 242002	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52968

NODE=M193225

NODE=M193R01
NODE=M193R01NODE=M193R01;LINKAGE=F
NODE=M193R01;LINKAGE=D
NODE=M193R01;LINKAGE=A

NODE=M193R01;LINKAGE=C

NODE=M193R01;LINKAGE=AA
NODE=M193R01;LINKAGE=E
NODE=M193R01;LINKAGE=G
NODE=M193R01;LINKAGE=B

NODE=M193R01;LINKAGE=AI

NODE=M193R02
NODE=M193R02

NODE=M193

$\psi(4160)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M025

 $\psi(4160)$ MASS

NODE=M025M

NODE=M025M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4191 ± 5 OUR AVERAGE			
4186.8 ± 8.7 ± 30	¹ ABLIKIM	23BH BES3	$e^+e^- \rightarrow D_s^{*+}D_s^{*-}$
4191 $\begin{smallmatrix} +9 \\ -8 \end{smallmatrix}$	AAIJ	13BC LHCB	$B^+ \rightarrow K^+\mu^+\mu^-$
4191.7 ± 6.5	² ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4193 ± 7	³ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4151 ± 4	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4155 ± 5	⁵ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4159 ± 20	BRANDELIK	78C DASP	e^+e^-
¹ Could also be the $\psi(4230)$.			
² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.			
³ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁴ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁵ From a fit to BES (BAI 02C) data.			

OCCUR=2

NODE=M025M;LINKAGE=B
NODE=M025M;LINKAGE=AB

NODE=M025M;LINKAGE=MO

NODE=M025M;LINKAGE=ST
NODE=M025M;LINKAGE=SE $\psi(4160)$ WIDTH

NODE=M025W

NODE=M025W
NEW

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
69 ± 10 OUR AVERAGE			
[70 ± 10 MeV OUR 2023 AVERAGE]			
55 ± 15 ± 53	¹ ABLIKIM	23BH BES3	$e^+e^- \rightarrow D_s^{*+}D_s^{*-}$
65 $\begin{smallmatrix} +22 \\ -16 \end{smallmatrix}$	AAIJ	13BC LHCB	$B^+ \rightarrow K^+\mu^+\mu^-$
71.8 ± 12.3	² ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
79 ± 14	³ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
107 ± 10	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
107 ± 16	⁵ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
78 ± 20	BRANDELIK	78C DASP	e^+e^-
¹ Could also be the $\psi(4230)$.			
² Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.			
³ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁴ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁵ From a fit to BES (BAI 02C) data.			

OCCUR=2

NODE=M025W;LINKAGE=B
NODE=M025W;LINKAGE=AB

NODE=M025W;LINKAGE=MO

NODE=M025W;LINKAGE=ST
NODE=M025W;LINKAGE=SE $\psi(4160)$ DECAY MODES

NODE=M025215;NODE=M025

NODE=M025

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more (less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 e^+e^-	$(6.9 \pm 3.3) \times 10^{-6}$	DESIG=1
Γ_2 $\mu^+\mu^-$	seen	DESIG=33
Γ_3 $D\bar{D}$	seen	DESIG=15;OUR EVAL;→ UNCHECKED ←
Γ_4 $D^0\bar{D}^0$	seen	DESIG=16
Γ_5 D^+D^-	seen	DESIG=17
Γ_6 $D^*\bar{D} + c.c.$	seen	DESIG=18;OUR EVAL;→ UNCHECKED ←

Γ_7	$D^*(2007)^0 \bar{D}^0 + \text{c.c.}$	seen			DESIG=19
Γ_8	$D^*(2010)^+ D^- + \text{c.c.}$	seen			DESIG=20
Γ_9	$D^* \bar{D}^*$	seen			DESIG=21;OUR EVAL;→ UNCHECKED ←
Γ_{10}	$D^*(2007)^0 \bar{D}^*(2007)^0$	seen			DESIG=22
Γ_{11}	$D^*(2010)^+ D^*(2010)^-$	seen			DESIG=23
Γ_{12}	$D^0 D^- \pi^+ + \text{c.c. (excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c., } D^*(2010)^+ D^- + \text{c.c.)}$	not seen			DESIG=24
Γ_{13}	$D \bar{D}^* \pi + \text{c.c. (excl. } D^* \bar{D}^*)$	seen			DESIG=25
Γ_{14}	$D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-)$	not seen			DESIG=26
Γ_{15}	$D_s^+ D_s^-$	not seen			DESIG=27
Γ_{16}	$D_s^{*+} D_s^- + \text{c.c.}$	seen			DESIG=28
Γ_{17}	$J/\psi \pi^+ \pi^-$	< 3	$\times 10^{-3}$	90%	DESIG=2
Γ_{18}	$J/\psi \pi^0 \pi^0$	< 3	$\times 10^{-3}$	90%	DESIG=3
Γ_{19}	$J/\psi K^+ K^-$	< 2	$\times 10^{-3}$	90%	DESIG=4
Γ_{20}	$J/\psi \eta$	< 8	$\times 10^{-3}$	90%	DESIG=5
Γ_{21}	$J/\psi \pi^0$	< 1	$\times 10^{-3}$	90%	DESIG=6
Γ_{22}	$J/\psi \eta'$	< 5	$\times 10^{-3}$	90%	DESIG=7
Γ_{23}	$J/\psi \pi^+ \pi^- \pi^0$	< 1	$\times 10^{-3}$	90%	DESIG=8
Γ_{24}	$\psi(2S) \pi^+ \pi^-$	< 4	$\times 10^{-3}$	90%	DESIG=9
Γ_{25}	$\chi_{c1} \gamma$	< 5	$\times 10^{-3}$	90%	DESIG=10
Γ_{26}	$\chi_{c2} \gamma$	< 1.3	%	90%	DESIG=11
Γ_{27}	$\chi_{c1} \pi^+ \pi^- \pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=12
Γ_{28}	$\chi_{c2} \pi^+ \pi^- \pi^0$	< 8	$\times 10^{-3}$	90%	DESIG=13
Γ_{29}	$h_c(1P) \pi^+ \pi^-$	< 5	$\times 10^{-3}$	90%	DESIG=29
Γ_{30}	$h_c(1P) \pi^0 \pi^0$	< 2	$\times 10^{-3}$	90%	DESIG=30
Γ_{31}	$h_c(1P) \eta$	< 2	$\times 10^{-3}$	90%	DESIG=31
Γ_{32}	$h_c(1P) \pi^0$	< 4	$\times 10^{-4}$	90%	DESIG=32
Γ_{33}	$\omega \pi^+ \pi^-$				DESIG=49
Γ_{34}	$\phi \pi^+ \pi^-$	< 2	$\times 10^{-3}$	90%	DESIG=14
Γ_{35}	$\gamma \chi_{c1}(3872)$	< 1.9	$\times 10^{-3}$	90%	DESIG=44
Γ_{36}	$\gamma \chi_{c0}(3915) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.36	$\times 10^{-4}$	90%	DESIG=35
Γ_{37}	$\gamma X(3930) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.18	$\times 10^{-4}$	90%	DESIG=36
Γ_{38}	$\gamma X(3940) \rightarrow \gamma J/\psi \pi^+ \pi^-$	< 1.47	$\times 10^{-4}$	90%	DESIG=37
Γ_{39}	$\gamma \chi_{c0}(3915) \rightarrow \gamma \gamma J/\psi$	< 1.26	$\times 10^{-4}$	90%	DESIG=39
Γ_{40}	$\gamma X(3930) \rightarrow \gamma \gamma J/\psi$	< 8.8	$\times 10^{-5}$	90%	DESIG=40
Γ_{41}	$\gamma X(3940) \rightarrow \gamma \gamma J/\psi$	< 1.79	$\times 10^{-4}$	90%	DESIG=41
Γ_{42}	$\omega \pi^0$	not seen			DESIG=47
Γ_{43}	$\omega \eta$	not seen			DESIG=48
Γ_{44}	$K^+ K^-$				DESIG=42
Γ_{45}	$K_S^0 K^\pm \pi^\mp$				DESIG=43
Γ_{46}	$p \bar{p} p \bar{p}$	not seen			DESIG=45
Γ_{47}	$\Lambda \bar{\Lambda}$	< 1.5	$\times 10^{-6}$	90%	DESIG=46
Γ_{48}	$\Xi^- \Xi^+$	< 8	$\times 10^{-5}$	90%	DESIG=50
Γ_{49}	$p K^- \bar{\Lambda} + \text{c.c.}$	< 6	$\times 10^{-6}$	90%	DESIG=51

 $\psi(4160)$ PARTIAL WIDTHS

NODE=M025220

 $\Gamma(e^+ e^-)$ **Γ_1** NODE=M025W1
NODE=M025W1

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.48±0.22	1 ABLIKIM	08D BES2	$e^+ e^- \rightarrow$ hadrons
••• We do not use the following data for averages, fits, limits, etc. •••			
0.4 to 1.1	2 MO	10 RVUE	$e^+ e^- \rightarrow$ hadrons
0.83±0.08	3 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.84±0.13	4 SETH	05A RVUE	$e^+ e^- \rightarrow$ hadrons
0.77±0.23	BRANDELIK	78C DASP	$e^+ e^-$

OCCUR=2

- ¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (293 \pm 57)^\circ$.
- ² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.
- ³ From a fit to Crystal Ball (OSTERHELD 86) data.
- ⁴ From a fit to BES (BAI 02C) data.

NODE=M025W1;LINKAGE=AB

NODE=M025W1;LINKAGE=MO

NODE=M025W1;LINKAGE=ST
NODE=M025W1;LINKAGE=SE $\Gamma(\mu^+\mu^-)$ Γ_2

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.45 ± 1.24 ± 0.94	^{1,2} ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M025W2
NODE=M025W2

¹ From a fit to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$.

NODE=M025W2;LINKAGE=A

² From solution 1 of 8 with equal fit quality. Other solutions range from $2.08 \pm 0.99 \pm 0.80$ to $2.45 \pm 1.24 \pm 0.94$ keV.

NODE=M025W2;LINKAGE=B

 $\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M025235

 $\Gamma(J/\psi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_1/\Gamma$

VALUE (eV)	EVT5	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

NODE=M025R42
NODE=M025R42

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.17 ± 0.04	86	^{1,2} ABLIKIM	20A BES3	$e^+e^- \rightarrow \eta' J/\psi$
1.07 ± 0.09	86	^{1,3} ABLIKIM	20A BES3	$e^+e^- \rightarrow \eta' J/\psi$

OCCUR=2

¹ Based on a fit to $\sigma(e^+e^- \rightarrow \eta' J/\psi)$ from $\sqrt{s} = 4.18$ to 4.60 GeV assuming interfering $\psi(4160)$ and $\psi(4260)$ contributions. At $\sqrt{s} = 4.18$ GeV, $\sigma(e^+e^- \rightarrow \eta' J/\psi) = 2.4 \pm 0.3 \pm 0.2$ pb.

NODE=M025R42;LINKAGE=A

² Solution I of the fit, corresponding to a phase of -0.03 ± 0.44 rad.

NODE=M025R42;LINKAGE=B

³ Solution II of the fit, corresponding to a phase of 2.54 ± 0.04 rad.

NODE=M025R42;LINKAGE=C

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	¹ HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

NODE=M025G01
NODE=M025G01

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M025G01;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M025G02
NODE=M025G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	¹ HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$
------	----	------------------	---------	--

NODE=M025G02;LINKAGE=A

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

 $\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{33}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M025R48
NODE=M025R48

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0011 ± 0.0008 ± 0.0001	^{1,2} ABLIKIM	23AQ BES3	fit to cross sections
0.651 ± 0.012 ± 0.040	^{2,3} ABLIKIM	23AQ BES3	fit to cross sections

OCCUR=3

OCCUR=4

¹ Solution I of the fit.

NODE=M025R48;LINKAGE=A

² From a fit to $e^+e^- \rightarrow \omega\pi^+\pi^-$ cross sections between 4 and 4.6 GeV. Recalculated from $12 \pi \Gamma(e^+e^-) B(\psi(4230) \rightarrow \omega\pi^+\pi^-)$. First uncertainty is from statistical and uncommon systematic uncertainties, and the second is a 6.2% common systematic uncertainty quoted in the paper.

NODE=M025R48;LINKAGE=C

³ Solution II of the fit.

NODE=M025R48;LINKAGE=B

 $\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{45}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M025R00
NODE=M025R00

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.71 ± 0.13 ± 0.12	¹ ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0095 ± 0.0088 ± 0.0004	² ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

OCCUR=5

¹ Solution I of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

NODE=M025R00;LINKAGE=A

² Solution II of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

NODE=M025R00;LINKAGE=D

$$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{47}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<0.7 \times 10^{-3}$	90	¹ ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4160)$

¹ From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

NODE=M025R45
NODE=M025R45

NODE=M025R45;LINKAGE=A

$$\Gamma(\Xi^-\bar{\Xi}^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{48}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<3.72 \times 10^{-2}$	90	¹ ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4160)$

¹ From a fit to $e^+e^- \rightarrow \Xi^-\bar{\Xi}^+$ cross sections.

NODE=M025R49
NODE=M025R49

NODE=M025R49;LINKAGE=A

$$\Gamma(pK^-\bar{\Lambda} + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{49}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	¹ ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4160)$

¹ From a fit to $e^+e^- \rightarrow pK^-\bar{\Lambda} + \text{c.c.}$ cross sections.

NODE=M025R51
NODE=M025R51

NODE=M025R51;LINKAGE=A

$\psi(4160) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M025230

$$\Gamma(J/\psi\eta)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{20}/\Gamma \times \Gamma_1/\Gamma$$

VALUE (units 10^{-8})	DOCUMENT ID	TECN	COMMENT
$2.8 \pm 0.9 \pm 0.9$	¹ WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\eta$
$12.8 \pm 1.7 \pm 2.0$	² WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.8 \pm 0.9 \pm 0.9$ ¹ WANG 13B BELL $e^+e^- \rightarrow J/\psi\eta\eta$
 $12.8 \pm 1.7 \pm 2.0$ ² WANG 13B BELL $e^+e^- \rightarrow J/\psi\eta\eta$

NODE=M025R32
NODE=M025R32

OCCUR=2

¹ Solution I of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

NODE=M025R32;LINKAGE=A

² Solution II of two equivalent solutions in a fit using two interfering resonances. Mass and width fixed at 4153 MeV and 103 MeV, respectively.

NODE=M025R32;LINKAGE=B

$\psi(4160)$ BRANCHING RATIOS

NODE=M025225

$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ AAIJ	13BC LHCB	$B^+ \rightarrow K^+\mu^+\mu^-$

¹ AAIJ 13BC report $B(B^+ \rightarrow K^+\psi(4160)) B(\psi(4160) \rightarrow \mu^+\mu^-) = (3.5^{+0.9}_{-0.8}) \times 10^{-9}$.

NODE=M025R31
NODE=M025R31

NODE=M025R31;LINKAGE=A

$$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*) \quad \Gamma_3/\Gamma_9$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.02 \pm 0.03 \pm 0.02$	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$

NODE=M025R14
NODE=M025R14

$$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$
seen	PAKHLOVA 08	BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$

NODE=M025R16
NODE=M025R16

$$\Gamma(D^+D^-)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$
seen	PAKHLOVA 08	BELL	$e^+e^- \rightarrow D^+D^-\gamma$
not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^+D^-\gamma$

NODE=M025R17
NODE=M025R17

$$\Gamma(D^*(2007)^0\bar{D}^0 + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$

NODE=M025R18
NODE=M025R18

$$\Gamma(D^*(2010)^+D^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_8/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ZHUKOVA	18 BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^-$
seen	PAKHLOVA	07 BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$

NODE=M025R19
NODE=M025R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Supersedes PAKHLOVA 07.

NODE=M025R19;LINKAGE=A

$$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_9
$0.34 \pm 0.14 \pm 0.05$	AUBERT	09M	BABR $e^+e^- \rightarrow \gamma D^{(*)}\bar{D}^{(*)}$	NODE=M025R15 NODE=M025R15

$$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$	NODE=M025R20 NODE=M025R20
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$	

$$\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
seen	¹ ZHUKOVA	18	BELL $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	NODE=M025R21 NODE=M025R21
seen	AUBERT	09M	BABR $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^{*-}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	PAKHLOVA	07	BELL $e^+e^- \rightarrow D^{*+}D^{*-}\gamma$	

¹Supersedes PAKHLOVA 07.

NODE=M025R21;LINKAGE=A

$$\Gamma(D^0D^-\pi^+ + \text{c.c. (excl. } D^*(2007)^0\bar{D}^0 + \text{c.c., } D^*(2010)^+D^- + \text{c.c.})/$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
not seen	PAKHLOVA	08A	BELL $e^+e^- \rightarrow D^0D^-\pi^+\gamma$	NODE=M025R22 NODE=M025R22

$$\Gamma(D\bar{D}^*\pi + \text{c.c. (excl. } D^*\bar{D}^*)/ \Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D\bar{D}^*\pi$	NODE=M025R23 NODE=M025R23

$$\Gamma(D^0D^{*-}\pi^+ + \text{c.c. (excl. } D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
not seen	PAKHLOVA	09	BELL $e^+e^- \rightarrow D^0D^{*-}\pi^+\gamma$	NODE=M025R24 NODE=M025R24

$$\Gamma(D_s^+D_s^-)/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
not seen	PAKHLOVA	11	BELL $e^+e^- \rightarrow D_s^+D_s^-\gamma$	NODE=M025R25 NODE=M025R25
not seen	DEL-AMO-SA..10N	BABR	$e^+e^- \rightarrow D_s^+D_s^-\gamma$	
not seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D_s^+D_s^-$	

$$\Gamma(D_s^{*+}D_s^- + \text{c.c.})/\Gamma_{\text{total}}$$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
seen	PAKHLOVA	11	BELL $e^+e^- \rightarrow D_s^{*+}D_s^-\gamma$	NODE=M025R26 NODE=M025R26
seen	DEL-AMO-SA..10N	BABR	$e^+e^- \rightarrow D_s^{*+}D_s^-\gamma$	
seen	CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D_s^{*+}D_s^-$	

$$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
<3	90	COAN	06	CLEO 4.12-4.2 $e^+e^- \rightarrow$ hadrons	NODE=M025R01 NODE=M025R01

$$\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
<3	90	COAN	06	CLEO 4.12-4.2 $e^+e^- \rightarrow$ hadrons	NODE=M025R02 NODE=M025R02

$$\Gamma(J/\psi K^+K^-)/\Gamma_{\text{total}}$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{19}/Γ
<2	90	COAN	06	CLEO 4.12-4.2 $e^+e^- \rightarrow$ hadrons	NODE=M025R03 NODE=M025R03

$$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ
<8	90	COAN	06	CLEO 4.12-4.2 $e^+e^- \rightarrow$ hadrons	NODE=M025R04 NODE=M025R04

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	¹ ABLIKIM	15L	BES3 $e^+e^- \rightarrow J/\psi\eta$
seen	WANG	13B	BELL $e^+e^- \rightarrow J/\psi\eta\gamma$

¹An enhancement around 4.2 GeV is observed.

NODE=M025R04;LINKAGE=A

$\Gamma(J/\psi\pi^0)/\Gamma_{\text{total}}$						Γ_{21}/Γ	NODE=M025R05 NODE=M025R05
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<1	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$						Γ_{22}/Γ	NODE=M025R06 NODE=M025R06
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<5	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(J/\psi\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{23}/Γ	NODE=M025R07 NODE=M025R07
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<1	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{24}/Γ	NODE=M025R08 NODE=M025R08
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<4	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(\chi_{c1}\gamma)/\Gamma_{\text{total}}$						Γ_{25}/Γ	NODE=M025R09 NODE=M025R09
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<7	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(\chi_{c2}\gamma)/\Gamma_{\text{total}}$						Γ_{26}/Γ	NODE=M025R10 NODE=M025R10
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<13	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(\chi_{c1}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{27}/Γ	NODE=M025R11 NODE=M025R11
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<2	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(\chi_{c2}\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{28}/Γ	NODE=M025R12 NODE=M025R12
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<8	90	COAN	06	CLEO	4.12–4.2 $e^+e^- \rightarrow$ hadrons		
$\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_{29}/Γ	NODE=M025R27 NODE=M025R27
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<5	90	¹ PEDLAR	11	CLEO	$e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$		
		¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 15.6 \pm 2.3 \pm 1.9 \pm 3.0$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.					NODE=M025R27;LINKAGE=PE
$\Gamma(h_c(1P)\pi^0\pi^0)/\Gamma_{\text{total}}$						Γ_{30}/Γ	NODE=M025R28 NODE=M025R28
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<2	90	¹ PEDLAR	11	CLEO	$e^+e^- \rightarrow h_c(1P)\pi^0\pi^0$		
		¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0\pi^0) = 3.0 \pm 3.3 \pm 1.1 \pm 0.6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.					NODE=M025R28;LINKAGE=PE
$\Gamma(h_c(1P)\eta)/\Gamma_{\text{total}}$						Γ_{31}/Γ	NODE=M025R29 NODE=M025R29
VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT		
<2	90		¹ PEDLAR	11	CLEO	$e^+e^- \rightarrow h_c(1P)\eta$	
			• • • We do not use the following data for averages, fits, limits, etc. • • •				
		possibly seen	41	² ABLIKIM	17R	BES3	$e^+e^- \rightarrow h_c(1P)\eta$
		¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\eta) = 4.7 \pm 1.7 \pm 1.0 \pm 0.9$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.					NODE=M025R29;LINKAGE=PE
		² An enhancement around 4.2 GeV is observed.					NODE=M025R29;LINKAGE=A
$\Gamma(h_c(1P)\pi^0)/\Gamma_{\text{total}}$						Γ_{32}/Γ	NODE=M025R30 NODE=M025R30
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT			
<0.4	90	¹ PEDLAR	11	CLEO	$e^+e^- \rightarrow h_c(1P)\pi^0$		
		¹ At $\sqrt{s} = 4170$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^0) = -0.7 \pm 1.8 \pm 0.7 \pm 0.1$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.					NODE=M025R30;LINKAGE=PE

$\Gamma(\phi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<2	90	COAN	06	CLEO 4.12-4.2 $e^+e^- \rightarrow$ hadrons

NODE=M025R13
NODE=M025R13

 $\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.9 $\times 10^{-3}$ (CL = 90%)		[<1.8 $\times 10^{-3}$ (CL = 90%)	OUR 2023 BEST LIMIT]

NODE=M025R43
NODE=M025R43

<1.9 $\times 10^{-3}$ 90 1,2 XIAO 13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.013 90 1,3 XIAO 13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

OCCUR=2

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R43;LINKAGE=A

² XIAO 13 reports $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 0.68 \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$.

NODE=M025R43;LINKAGE=B

³ XIAO 13 reports $[\Gamma(\psi(4160) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \gamma J/\psi)] < 1.05 \times 10^{-4}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \gamma J/\psi) = 7.8 \times 10^{-3}$.

NODE=M025R43;LINKAGE=C

 $\Gamma(\gamma\chi_{c0}(3915) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.36 $\times 10^{-4}$	90	¹ XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R35
NODE=M025R35

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R35;LINKAGE=A

 $\Gamma(\gamma X(3930) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.18 $\times 10^{-4}$	90	¹ XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R36
NODE=M025R36

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R36;LINKAGE=A

 $\Gamma(\gamma X(3940) \rightarrow \gamma J/\psi\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.47 $\times 10^{-4}$	90	¹ XIAO	13 $\psi(4160) \rightarrow \gamma J/\psi\pi^+\pi^-$

NODE=M025R37
NODE=M025R37

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R37;LINKAGE=A

 $\Gamma(\gamma\chi_{c0}(3915) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.26 $\times 10^{-4}$	90	¹ XIAO	13 $\psi(4160) \rightarrow \gamma\gamma J/\psi$

NODE=M025R39
NODE=M025R39

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R39;LINKAGE=A

 $\Gamma(\gamma X(3930) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<0.88 $\times 10^{-4}$	90	¹ XIAO	13 $\psi(4160) \rightarrow \gamma\gamma J/\psi$

NODE=M025R40
NODE=M025R40

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R40;LINKAGE=A

 $\Gamma(\gamma X(3940) \rightarrow \gamma\gamma J/\psi)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	COMMENT
<1.79 $\times 10^{-4}$	90	¹ XIAO	13 $\psi(4160) \rightarrow \gamma\gamma J/\psi$

NODE=M025R41
NODE=M025R41

¹ Obtained by analyzing CLEO data but not authored by the CLEO Collaboration.

NODE=M025R41;LINKAGE=A

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	22K	BES3 $e^+e^- \rightarrow \omega\pi^0$

NODE=M025R46
NODE=M025R46

 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	22K	BES3 $e^+e^- \rightarrow \omega\eta$

NODE=M025R47
NODE=M025R47

 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2 $\times 10^{-5}$	90	¹ DRUZHININ	15	RVUE $e^+e^- \rightarrow \psi(3770)$

NODE=M025R33
NODE=M025R33

••• We do not use the following data for averages, fits, limits, etc. •••

¹ DRUZHININ 15 uses BABAR and CLEO data taking into account interference of the processes $e^+e^- \rightarrow K^+K^-$ and $e^+e^- \rightarrow K_S^0 K_L^0$.

NODE=M025R33;LINKAGE=A

$\Gamma(p\bar{p}p\bar{p})/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	21D	BES3 4.0-4.6 $e^+e^- \rightarrow p\bar{p}p\bar{p}$

NODE=M025R44
 NODE=M025R44

 $\psi(4160)$ REFERENCES

ABLIKIM	23AQ	JHEP 2308 159	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62313
ABLIKIM	23BH	PRL 131 151903	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62433
ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62437
ABLIKIM	23BL	JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62438
ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	21D	PR D103 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61031
ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60210
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59856
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
ABLIKIM	17R	PR D96 012001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58009
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan		REFID=57991
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15L	PR D91 112005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56777
DRUZHININ	15	PR D92 054024	V.P. Druzhinin	(NOVO)	REFID=56962
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
AAIJ	13BC	PRL 111 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55229
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
XIAO	13	PR D87 057501	T. Xiao <i>et al.</i>	(NWES, WAYN)	REFID=55381
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
COAN	06	PRL 96 162003	T.E. Coan <i>et al.</i>	(CLEO Collab.)	REFID=51075
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232

NODE=M025

NODE=M190

X(4160)

$$J^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Seen by PAKHLOV 08 in $e^+e^- \rightarrow J/\psi X$, $X \rightarrow D^*\bar{D}^*$

NODE=M190

A state with consistent mass and width is seen by AAIJ 21E in $B^+ \rightarrow X(4160)K^+$ with $X(4160) \rightarrow J/\psi\phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ with a significance (accounting for systematic uncertainties) of 4.8σ . The $J^{PC} = 2^{-+}$ assignment is favored over other assignments with a significance of more than 4σ .

X(4160) MASS

NODE=M190M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4153^{+23}_{-21} OUR AVERAGE				
$4146 \pm 18 \pm 33$	24k	¹ AAIJ	21E	LHCB $B^+ \rightarrow J/\psi\phi K^+$
$4156^{+25}_{-20} \pm 15$	24	PAKHLOV	08	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M190M

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 4.8σ .

NODE=M190M;LINKAGE=A

X(4160) WIDTH

NODE=M190W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
136^{+60}_{-35} OUR AVERAGE				
$135 \pm 28^{+59}_{-30}$	24k	¹ AAIJ	21E	LHCB $B^+ \rightarrow J/\psi\phi K^+$
$139^{+111}_{-61} \pm 21$	24	PAKHLOV	08	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M190W

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 4.8σ .

NODE=M190W;LINKAGE=A

X(4160) DECAY MODES

NODE=M190215;NODE=M190

Mode	Fraction (Γ_i/Γ)
Γ_1 $D\bar{D}$	not seen
Γ_2 $D^*\bar{D} + \text{c.c.}$	not seen
Γ_3 $D^*\bar{D}^*$	seen
Γ_4 $J/\psi\phi$	seen

DESIG=1;OUR EVAL;→ UNCHECKED ←
DESIG=2;OUR EVAL;→ UNCHECKED ←
DESIG=3;OUR EVAL;→ UNCHECKED ←
DESIG=4

X(4160) BRANCHING RATIOS

NODE=M190225

$\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$					Γ_1/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.09	90	PAKHLOV	08	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M190R01
NODE=M190R01

$\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$					Γ_2/Γ_3
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.22	90	PAKHLOV	08	BELL	$e^+e^- \rightarrow J/\psi X$

NODE=M190R02
NODE=M190R02

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$					Γ_4/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	24k	¹ AAIJ	21E	LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M190R00
NODE=M190R00

¹From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 4.8 σ .

NODE=M190R00;LINKAGE=A

X(4160) REFERENCES

AAIJ 21E PRL 127 082001 R. Aaij *et al.* (LHCb Collab.)
PAKHLOV 08 PRL 100 202001 P. Pakhlov *et al.* (BELLE Collab.)

NODE=M190

REFID=61150
REFID=52302

NODE=M074

 $\psi(4230)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $Y(4230)$; was $\psi(4260)$

The original $\psi(4260)$ (also known as $Y(4260)$) was observed by AUBERT, B 05I as a peak in the energy dependence of the $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ cross section and was confirmed by HE 06B, YUAN 07, LEES 12AC, and LIU 13B in the same process. A higher-statistics analysis by ABLIKIM 17B revealed an asymmetry in the cross section and resulted in a shift of the peak position to a lower mass. The $\psi(4260)$ was therefore renamed $\psi(4230)$. The energy-dependent cross sections for e^+e^- to other channels also exhibit peaks in the same mass region. The parameters corresponding to those peaks are also listed here, but the number of states in this region remains to be determined.

NODE=M074

For details see the review on "Spectroscopy of mesons containing two heavy quarks."

 $\psi(4230)$ MASS

NODE=M074M


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4222.1 ± 2.3 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.		
4222.5 ± 2.4 MeV	OUR	2023 AVERAGE Scale factor = 1.7]		
$4226.9 \pm 6.6 \pm 22.0$		¹ ABLIKIM	23U	BES3 $e^+e^- \rightarrow K_S^0 K_S^0 J/\psi$
$4209.6 \pm 4.7 \pm 5.9$		² ABLIKIM	23X	BES3 $e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$
$4221.4 \pm 1.5 \pm 2.0$		³ ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
$4225.3 \pm 2.3 \pm 21.5$		⁴ ABLIKIM	22AU	BES3 $e^+e^- \rightarrow K^+ K^- J/\psi$
$4234.4 \pm 3.2 \pm 0.2$		⁵ ABLIKIM	21AJ	BES3 $e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
$4216.7 \pm 8.9 \pm 4.1$		⁶ ABLIKIM	20AG	BES3 $e^+e^- \rightarrow \mu^+\mu^-$
$4220.4 \pm 2.4 \pm 2.3$		⁷ ABLIKIM	20N	BES3 $e^+e^- \rightarrow \pi^0 \pi^0 J/\psi$
$4218.6 \pm 3.8 \pm 2.5$		⁷ ABLIKIM	20O	BES3 $e^+e^- \rightarrow \eta J/\psi$
$4218.5 \pm 1.6 \pm 4.0$		⁸ ABLIKIM	19AI	BES3 $e^+e^- \rightarrow \omega \chi_{c0}$
$4228.6 \pm 4.1 \pm 6.3$		ABLIKIM	19R	BES3 $e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$
$4200.6^{+7.9}_{-13.3} \pm 3.0$		⁹ ABLIKIM	19V	BES3 $e^+e^- \rightarrow \gamma \chi_{c1}(3872)$
$4218^{+5.5}_{-4.5} \pm 0.9$		ABLIKIM	17G	BES3 $e^+e^- \rightarrow \pi^+\pi^- h_c$

NODE=M074M

NEW

• • • We do not use the following data for averages, fits, limits, etc. • • •

4231.9 ± 5.3 ± 4.9	ABLIKIM	20N	BES3	$e^+e^- \rightarrow \pi^0 T_{c\bar{c}1}(3900)^0,$ $T_{c\bar{c}1}^0 \rightarrow \pi^0 J/\psi$	OCCUR=2
4222.0 ± 3.1 ± 1.4	¹⁰ ABLIKIM	17B	BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$	
4209.5 ± 7.4 ± 1.4	¹¹ ABLIKIM	17V	BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
4209.1 ± 6.8 ± 7.0	¹² ZHANG	17B	RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
4223.3 ± 1.6 ± 2.5	¹³ ZHANG	17C	RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$	OCCUR=2
4230 ± 8 ± 6 180	¹⁴ ABLIKIM	15C	BES3	$e^+e^- \rightarrow \omega\chi_{c0}$	
4258.6 ± 8.3 ± 12.1	¹⁵ LIU	13B	BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$	
4245 ± 5 ± 4	¹⁶ LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$	
4247 ± 12 $\begin{smallmatrix} +17 \\ -32 \end{smallmatrix}$	^{15,17} YUAN	07	BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$	
4284 $\begin{smallmatrix} +17 \\ -16 \end{smallmatrix}$ ± 413.6	HE	06B	CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$	
4259 ± 8 $\begin{smallmatrix} + \\ - \end{smallmatrix} \frac{2}{6} 125$	¹⁸ AUBERT,B	05I	BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$	
¹ From a three-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.128$ – 4.950 GeV.					NODE=M074M;LINKAGE=I
² From a cross-section measurement of $e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. The two other resonances have masses (widths) 4675.3 ± 29.7 (218.3 ± 73.5) MeV and 4469.1 ± 26.4 (246.3 ± 37.9) MeV.					NODE=M074M;LINKAGE=J
³ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730$ – 4.7008 GeV.					NODE=M074M;LINKAGE=F
⁴ From a two-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.127$ – 4.600 GeV. The second resonance has a mass of $4484.7 \pm 13.3 \pm 24.1$ MeV and a total width of $111.1 \pm 30.1 \pm 15.2$ MeV.					NODE=M074M;LINKAGE=H
⁵ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008$ – 4.698 GeV.					NODE=M074M;LINKAGE=C
⁶ Solution 1 of 8 with equal fit quality to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes. Other solutions range from $4212.8 \pm 7.2 \pm 4.0$ to $4219.4 \pm 11.2 \pm 4.1$ MeV.					NODE=M074M;LINKAGE=HP
⁷ From a fit of the measured cross section in the range $\sqrt{s} = 3.808$ – 4.600 GeV.					NODE=M074M;LINKAGE=GP
⁸ From a fit of the measured cross section from $\sqrt{s} = 4.178$ – 4.278 GeV. Supersedes ABLIKIM 15C.					NODE=M074M;LINKAGE=CP
⁹ Simultaneous fit to $\chi_{c1} \rightarrow \omega J/\psi$ and $\chi_{c1} \rightarrow \pi^+\pi^- J/\psi$.					NODE=M074M;LINKAGE=FP
¹⁰ From a three-resonance fit. Superseded by ABLIKIM 22AM.					NODE=M074M;LINKAGE=G
¹¹ From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} . Superseded by ABLIKIM 21AJ.					NODE=M074M;LINKAGE=BP
¹² From a three-resonance fit.					NODE=M074M;LINKAGE=A
¹³ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.					NODE=M074M;LINKAGE=CA
¹⁴ From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21$ – 4.42 GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.					NODE=M074M;LINKAGE=AP
¹⁵ From a two-resonance fit.					NODE=M074M;LINKAGE=YU
¹⁶ From a single-resonance fit. Supersedes AUBERT,B 05I.					NODE=M074M;LINKAGE=LE
¹⁷ Superseded by LIU 13B.					NODE=M074M;LINKAGE=YN
¹⁸ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.					NODE=M074M;LINKAGE=AU



psi(4230) WIDTH

NODE=M074W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
49 ± 7 OUR AVERAGE Error includes scale factor of 3.4. See the ideogram below.
 [48 ± 8 MeV OUR 2023 AVERAGE Scale factor = 3.6]

NODE=M074W
NEW

71.7±16.2±32.8	1	ABLIKIM	23U	BES3	$e^+e^- \rightarrow K_S^0 K_S^0 J/\psi$
81.6±17.8± 9.0	2	ABLIKIM	23X	BES3	$e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$
41.8± 2.9± 2.7	3	ABLIKIM	22AM	BES3	$e^+e^- \rightarrow \pi^+ \pi^- J/\psi$
72.9± 6.1±30.8	4	ABLIKIM	22AU	BES3	$e^+e^- \rightarrow K^+ K^- J/\psi$
17.6±18.1± 0.9	5	ABLIKIM	21AJ	BES3	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$
47.2±22.8±10.5	6	ABLIKIM	20AG	BES3	$e^+e^- \rightarrow \mu^+ \mu^-$
46.2± 4.7± 2.1	7	ABLIKIM	20N	BES3	$e^+e^- \rightarrow \pi^0 \pi^0 J/\psi$
82.0± 5.7± 0.4	7	ABLIKIM	20O	BES3	$e^+e^- \rightarrow \eta J/\psi$
28.2± 3.9± 1.6	8	ABLIKIM	19AI	BES3	$e^+e^- \rightarrow \omega \chi_{c0}$
77.0± 6.8± 6.3		ABLIKIM	19R	BES3	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + c.c.$
115 $\begin{smallmatrix} +38 \\ -26 \end{smallmatrix}$ ±12	9	ABLIKIM	19V	BES3	$e^+e^- \rightarrow \gamma \chi_{c1}(3872)$
66.0 $\begin{smallmatrix} +12.3 \\ -8.3 \end{smallmatrix}$ ± 0.4		ABLIKIM	17G	BES3	$e^+e^- \rightarrow \pi^+ \pi^- h_c$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
41.2±16.0±16.4		ABLIKIM	20N	BES3	$e^+e^- \rightarrow \pi^0 T_{c\bar{c}1}(3900)^0,$ $T_{c\bar{c}1}^0 \rightarrow \pi^0 J/\psi$
44.1± 4.3± 2.0	10	ABLIKIM	17B	BES3	$e^+e^- \rightarrow \pi^+ \pi^- J/\psi$
80.1±24.6± 2.9	11	ABLIKIM	17V	BES3	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$
76.6±14.2± 2.4	12	ZHANG	17B	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- \psi(2S)$
54.2± 2.6± 1.0	13	ZHANG	17C	RVUE	$e^+e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$
38 ±12 ± 2 180	14	ABLIKIM	15C	BES3	$e^+e^- \rightarrow \omega \chi_{c0}$
134.1±16.4± 5.5	15	LIU	13B	BELL	$e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
114 $\begin{smallmatrix} +16 \\ -15 \end{smallmatrix}$ ± 7	16	LEES	12AC	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
108 ±19 ±10	15,17	YUAN	07	BELL	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
73 $\begin{smallmatrix} +39 \\ -25 \end{smallmatrix}$ ± 5 13.6		HE	06B	CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$
88 ±23 $\begin{smallmatrix} +6 \\ -4 \end{smallmatrix}$ 125	18	AUBERT,B	05I	BABR	10.58 $e^+e^- \rightarrow \gamma \pi^+ \pi^- J/\psi$

OCCUR=2

¹ From a three-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.128\text{--}4.950$ GeV.
² From a cross-section measurement of $e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. The two other resonances have masses (widths) 4675.3 ± 29.7 (218.3 ± 73.5) MeV and 4469.1 ± 26.4 (246.3 ± 37.9) MeV.
³ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730\text{--}4.7008$ GeV.

NODE=M074W;LINKAGE=H

NODE=M074W;LINKAGE=I

NODE=M074W;LINKAGE=F

- 4 From a two-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.127\text{--}4.600$ GeV. The second resonance has a mass of $4484.7 \pm 13.3 \pm 24.1$ MeV and a total width of $111.1 \pm 30.1 \pm 15.2$ MeV.
- 5 From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.
- 6 Solution 1 of 8 with equal fit quality to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes. Other solutions range from $36.4 \pm 16.8 \pm 8.1$ to $49.6 \pm 22.6 \pm 11.0$ MeV.
- 7 From a fit of the measured cross section in the range $\sqrt{s} = 3.808\text{--}4.600$ GeV.
- 8 From a fit of the measured cross section from $\sqrt{s} = 4.178\text{--}4.278$ GeV. Supersedes ABLIKIM 15C.
- 9 Simultaneous fit to $\chi_{c1} \rightarrow \omega J/\psi$ and $\chi_{c1} \rightarrow \pi^+\pi^- J/\psi$.
- 10 From a three-resonance fit. Superseded by ABLIKIM 22AM.
- 11 From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} . Superseded by ABLIKIM 21AJ.
- 12 From a three-resonance fit.
- 13 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.
- 14 From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.
- 15 From a two-resonance fit.
- 16 From a single-resonance fit. Supersedes AUBERT,B 05I.
- 17 Superseded by LIU 13B.
- 18 From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074W;LINKAGE=G

NODE=M074W;LINKAGE=D

NODE=M074W;LINKAGE=GP

NODE=M074W;LINKAGE=FP

NODE=M074W;LINKAGE=CP

NODE=M074W;LINKAGE=EP

NODE=M074W;LINKAGE=E

NODE=M074W;LINKAGE=BP

NODE=M074W;LINKAGE=C

NODE=M074W;LINKAGE=B


NODE=M074W;LINKAGE=AP

NODE=M074W;LINKAGE=YU

NODE=M074W;LINKAGE=LE

NODE=M074W;LINKAGE=YN

NODE=M074W;LINKAGE=AU



$\psi(4230)$ DECAY MODES

NODE=M074215;NODE=M074

Mode	Fraction (Γ_i/Γ)	
Γ_1 e^+e^-		DESIG=1
Γ_2 $\mu^+\mu^-$	$(3.1 \pm 2.8) \times 10^{-5}$	DESIG=63
Γ_3 $\eta_c(1S)\pi^+\pi^-$	not seen	DESIG=65
Γ_4 $\eta_c(1S)\pi^+\pi^-\pi^0$	seen	DESIG=64
Γ_5 $J/\psi\pi^+\pi^-$	seen	DESIG=2
Γ_6 $J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-$	seen	DESIG=41;OUR EVAL;→ UNCHECKED ←
Γ_7 $T_{c\bar{c}1}(3900)^\pm \pi^\mp, T_{c\bar{c}1}^\pm \rightarrow$	seen	DESIG=43;OUR EVAL;→ UNCHECKED ←
Γ_8 $J/\psi\pi^0\pi^0$	seen	DESIG=4
Γ_9 $J/\psi K^+K^-$	seen	DESIG=5;OUR EVAL;→ UNCHECKED ←
Γ_{10} $J/\psi K_S^0 K_S^0$	not seen	DESIG=44
Γ_{11} $J/\psi\eta$	seen	DESIG=6
Γ_{12} $J/\psi\pi^0$	not seen	DESIG=7;OUR EVAL;→ UNCHECKED ←

Γ ₁₃	$J/\psi\eta'$	seen	DESIG=8;OUR EVAL;→ UNCHECKED ←
Γ ₁₄	$J/\psi\pi^+\pi^-\pi^0$	not seen	DESIG=9;OUR EVAL;→ UNCHECKED ←
Γ ₁₅	$J/\psi\eta\pi^0$	not seen	DESIG=45
Γ ₁₆	$J/\psi\eta\eta$	not seen	DESIG=10;OUR EVAL;→ UNCHECKED ←
Γ ₁₇	$\psi(2S)\pi^+\pi^-$	seen	DESIG=11
Γ ₁₈	$\psi(2S)\eta$	not seen	DESIG=12;OUR EVAL;→ UNCHECKED ←
Γ ₁₉	$\chi_{c0}\omega$	seen	DESIG=13
Γ ₂₀	$\chi_{c1}\pi^+\pi^-\pi^0$	not seen	DESIG=16;OUR EVAL;→ UNCHECKED ←
Γ ₂₁	$\chi_{c2}\pi^+\pi^-\pi^0$	not seen	DESIG=17;OUR EVAL;→ UNCHECKED ←
Γ ₂₂	$h_c(1P)\pi^+\pi^-$	seen	DESIG=40
Γ ₂₃	$\phi\pi^+\pi^-$	not seen	DESIG=18;OUR EVAL;→ UNCHECKED ←
Γ ₂₄	$\phi f_0(980) \rightarrow \phi\pi^+\pi^-$	not seen	DESIG=22;OUR EVAL;→ UNCHECKED ←
Γ ₂₅	ϕK^+K^-		DESIG=72
Γ ₂₆	$\phi K_S^0 K_S^0$		DESIG=73
Γ ₂₇	$\phi\eta$	not seen	DESIG=76
Γ ₂₈	$\phi\eta'$	not seen	DESIG=70;OUR EVAL;→ UNCHECKED ←
Γ ₂₉	$D\bar{D}$	not seen	DESIG=19;OUR EVAL;→ UNCHECKED ←
Γ ₃₀	$D^0\bar{D}^0$	not seen	DESIG=31
Γ ₃₁	D^+D^-	not seen	DESIG=32
Γ ₃₂	$D^*\bar{D}^+ + c.c.$	not seen	DESIG=23;OUR EVAL;→ UNCHECKED ←
Γ ₃₃	$D^*(2007)^0\bar{D}^0 + c.c.$	not seen	DESIG=33
Γ ₃₄	$D^*(2010)^+D^- + c.c.$	not seen	DESIG=34
Γ ₃₅	$D^*\bar{D}^*$		DESIG=24
Γ ₃₆	$D^*(2007)^0\bar{D}^*(2007)^0$	not seen	DESIG=35
Γ ₃₇	$D^*(2010)^+D^*(2010)^-$	not seen	DESIG=36
Γ ₃₈	$D\bar{D}\pi + c.c.$		DESIG=37
Γ ₃₉	$D^0D^-\pi^+ + c.c. (excl. D^*(2007)^0\bar{D}^{*0} + c.c., D^*(2010)^+D^- + c.c.)$	not seen	DESIG=38
Γ ₄₀	$D\bar{D}^*\pi + c.c. (excl. D^*\bar{D}^*)$	not seen	DESIG=25
Γ ₄₁	$D^0D^{*-}\pi^+ + c.c. (excl. D^*(2010)^+D^*(2010)^-)$	not seen	DESIG=39
Γ ₄₂	$D^0D^*(2010)^-\pi^+ + c.c.$	seen	DESIG=30
Γ ₄₃	$D_1(2420)\bar{D} + c.c.$	not seen	DESIG=50
Γ ₄₄	$D^*\bar{D}^*\pi$	seen	DESIG=26
Γ ₄₅	$D^{*0}D^{*-}\pi^+$	seen	DESIG=74;OUR EVAL;→ UNCHECKED ←
Γ ₄₆	$D_s^+D_s^-$	not seen	DESIG=27
Γ ₄₇	$D_s^{*+}D_s^- + c.c.$	not seen	DESIG=28
Γ ₄₈	$D_s^{*+}D_s^{*-}$	not seen	DESIG=29
Γ ₄₉	$\rho\bar{\rho}$	not seen	DESIG=3;OUR EVAL;→ UNCHECKED ←
Γ ₅₀	$\rho\bar{\rho}\pi^0$	not seen	DESIG=46;OUR EVAL;→ UNCHECKED ←
Γ ₅₁	$\rho\bar{\rho}\eta$	not seen	DESIG=61
Γ ₅₂	$\omega\pi^+\pi^-$		DESIG=71
Γ ₅₃	$\rho\bar{\rho}\omega$	not seen	DESIG=62
Γ ₅₄	$\Xi^-\Xi^+$	not seen	DESIG=51;OUR EVAL;→ UNCHECKED ←
Γ ₅₅	$\pi^+\pi^+\pi^-\pi^-$	not seen	DESIG=53;OUR EVAL;→ UNCHECKED ←
Γ ₅₆	$\pi^+\pi^+\pi^-\pi^-\pi^0$	not seen	DESIG=54;OUR EVAL;→ UNCHECKED ←
Γ ₅₇	$\omega\pi^0$	not seen	DESIG=68
Γ ₅₈	$\omega\eta$	not seen	DESIG=69
Γ ₅₉	$K_S^0 K^\pm\pi^\mp$	not seen	DESIG=20;OUR EVAL;→ UNCHECKED ←
Γ ₆₀	$K_S^0 K^\pm\pi^\mp\pi^0$	not seen	DESIG=48;OUR EVAL;→ UNCHECKED ←
Γ ₆₁	$K_S^0 K^\pm\pi^\mp\eta$	not seen	DESIG=49;OUR EVAL;→ UNCHECKED ←
Γ ₆₂	$K^+K^-\pi^0$	not seen	DESIG=21;OUR EVAL;→ UNCHECKED ←
Γ ₆₃	$K^+K^-\pi^+\pi^-$	not seen	DESIG=55;OUR EVAL;→ UNCHECKED ←
Γ ₆₄	$K^+K^-\pi^+\pi^-\pi^0$	not seen	DESIG=56;OUR EVAL;→ UNCHECKED ←
Γ ₆₅	$K^+K^+K^-K^-$	not seen	DESIG=57;OUR EVAL;→ UNCHECKED ←
Γ ₆₆	$K^+K^+K^-K^-\pi^0$	not seen	DESIG=58;OUR EVAL;→ UNCHECKED ←
Γ ₆₇	$\rho\bar{\rho}\pi^+\pi^-$	not seen	DESIG=59;OUR EVAL;→ UNCHECKED ←
Γ ₆₈	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	not seen	DESIG=60;OUR EVAL;→ UNCHECKED ←
Γ ₆₉	$\rho\bar{\rho}\rho\bar{\rho}$	not seen	DESIG=67
Γ ₇₀	$\Lambda\bar{\Lambda}$	not seen	DESIG=52;OUR EVAL;→ UNCHECKED ←
Γ ₇₁	$\rho K^-\bar{\Lambda} + c.c.$		DESIG=75

Radiative decays

Γ_{72}	$\eta_c(1S)\gamma$	possibly seen
Γ_{73}	$\eta_c(1S)\pi^0\gamma$	not seen
Γ_{74}	$\chi_{c1}\gamma$	not seen
Γ_{75}	$\chi_{c2}\gamma$	not seen
Γ_{76}	$\chi_{c1}(3872)\gamma$	seen

NODE=M074;CLUMP=C
 DESIG=47
 DESIG=66
 DESIG=14;OUR EVAL;→ UNCHECKED ←
 DESIG=15;OUR EVAL;→ UNCHECKED ←
 DESIG=42

 $\psi(4230)$ PARTIAL WIDTHS $\Gamma(\mu^+\mu^-)$ Γ_2

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.53±1.26±0.54	1,2 ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

¹ From a fit to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$.

² From solution 1 of 8 with equal fit quality. Other solutions range from $1.09 \pm 0.84 \pm 0.39$ to $1.53 \pm 1.26 \pm 0.54$ keV.

NODE=M074235

NODE=M074W01
 NODE=M074W01

NODE=M074W01;LINKAGE=A

NODE=M074W01;LINKAGE=B

 $\psi(4230) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M074230

 $\Gamma(J/\psi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_5\Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.2±1.0 OUR AVERAGE				

9.2±0.8±0.7		¹ LEES	12AC BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
8.9 ^{+3.9} _{-3.1} ±1.8	8.1	HE	06B CLEO	9.4–10.6 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.4±0.8±0.6		² LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
20.5±1.4±2.0		³ LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
6.0±1.2 ^{+4.7} _{-0.5}		^{2,4} YUAN	07 BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
20.6±2.3 ^{+9.1} _{-1.7}		^{3,4} YUAN	07 BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
5.5±1.0 ^{+0.8} _{-0.7}	125	⁵ AUBERT,B	05I BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$

¹ From a single-resonance fit. Supersedes AUBERT,B 05I.

² Solution I of two equivalent solutions in a fit using two interfering resonances.

³ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁴ Superseded by LIU 13B.

⁵ From a single-resonance fit. Two interfering resonances are not excluded. Superseded by LEES 12AC.

NODE=M074G1
 NODE=M074G1

OCCUR=2

OCCUR=2

NODE=M074G1;LINKAGE=LE
 NODE=M074G1;LINKAGE=YU
 NODE=M074G1;LINKAGE=YA
 NODE=M074G1;LINKAGE=YN
 NODE=M074G1;LINKAGE=AU

 $\Gamma(J/\psi K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_9\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.42±0.04±0.15		¹ ABLIKIM	22AU BES3	$e^+e^- \rightarrow K^+K^- J/\psi$
0.29±0.02±0.10		² ABLIKIM	22AU BES3	$e^+e^- \rightarrow K^+K^- J/\psi$
<1.7	90	³ SHEN	14 BELL	9.4–10.9 $e^+e^- \rightarrow \gamma K^+K^- J/\psi$
<1.2	90	⁴ YUAN	08 BELL	$e^+e^- \rightarrow \gamma K^+K^- J/\psi$

¹ Solution I from a two-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.127\text{--}4.600$ GeV. The second resonance has a mass of $4484.7 \pm 13.3 \pm 24.1$ MeV, a total width of $111.1 \pm 30.1 \pm 15.2$ MeV, and $\Gamma_{ee} \cdot B = 1.35 \pm 0.14 \pm 0.07$ eV. The phase difference is $1.72 \pm 0.09 \pm 0.52$ rad.

² Solution II from a two-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.127\text{--}4.600$ GeV. The second resonance has a mass of $4484.7 \pm 13.3 \pm 24.1$ MeV, a total width of $111.1 \pm 30.1 \pm 15.2$ MeV, and $\Gamma_{ee} \cdot B = 0.41 \pm 0.08 \pm 0.13$ eV. The phase difference is $5.49 \pm 0.35 \pm 0.58$ rad.

³ From a fit of the broad $K^+K^- J/\psi$ enhancement including a coherent $\psi(4260)$ amplitude with mass and width from LIU 13B. Supersedes YUAN 08. The shape of the cross section observed by ABLIKIM 18N between 4.2 and 4.3 GeV is incompatible with that of $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ in ABLIKIM 13T and ABLIKIM 17B. They also observe a broad enhancement around 4.5 GeV.

⁴ From a fit of the broad $K^+K^- J/\psi$ enhancement including a coherent $\psi(4260)$ amplitude with mass and width from YUAN 07.

NODE=M074G3
 NODE=M074G3

OCCUR=2

NODE=M074G3;LINKAGE=B

NODE=M074G3;LINKAGE=C

NODE=M074G3;LINKAGE=A

NODE=M074G3;LINKAGE=YU

$$\Gamma(J/\psi K_S^0 K_S^0) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{10} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.85	90	¹ SHEN	14 BELL	9.4–10.9 $e^+ e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$

NODE=M074G02
NODE=M074G02

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13±0.02±0.05		^{2,3} ABLIKIM	23U BES3	$e^+ e^- \rightarrow K_S^0 K_S^0 J/\psi$
0.14±0.03±0.06		^{2,4} ABLIKIM	23U BES3	$e^+ e^- \rightarrow K_S^0 K_S^0 J/\psi$
0.18±0.05±0.07		^{2,5} ABLIKIM	23U BES3	$e^+ e^- \rightarrow K_S^0 K_S^0 J/\psi$
0.20±0.04±0.07		^{2,6} ABLIKIM	23U BES3	$e^+ e^- \rightarrow K_S^0 K_S^0 J/\psi$

OCCUR=2

OCCUR=3

OCCUR=4

¹ From a fit of the $K_S^0 K_S^0 J/\psi$ mass range from 4.4 to 5.5 GeV including a coherent $\psi(4260)$ amplitude with mass and width from LIU 13B.

² A three-resonance fit to the dressed cross section in the range $\sqrt{s} = 4.128\text{--}4.950$ GeV.

³ Solution I.

⁴ Solution II.

⁵ Solution III.

⁶ Solution IV.

NODE=M074G02;LINKAGE=A

NODE=M074G02;LINKAGE=B

NODE=M074G02;LINKAGE=F

NODE=M074G02;LINKAGE=C

NODE=M074G02;LINKAGE=D

NODE=M074G02;LINKAGE=E

$$\Gamma(J/\psi \eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{11} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
8.0±1.7		¹ ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
4.8±1.0		² ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
7.0±1.5		³ ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
<14.2	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

NODE=M074G01
NODE=M074G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.0±1.7		¹ ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
4.8±1.0		² ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$
7.0±1.5		³ ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$

OCCUR=2

OCCUR=3

¹ Solution 1 of three equivalent fit solutions using three resonant structures.

² Solution 2 of three equivalent fit solutions using three resonant structures.

³ Solution 3 of three equivalent fit solutions using three resonant structures.

NODE=M074G01;LINKAGE=A

NODE=M074G01;LINKAGE=B

NODE=M074G01;LINKAGE=C

$$\Gamma(J/\psi \eta') \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{13} \Gamma_1 / \Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.06±0.03	46	^{1,2} ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$
1.38±0.11	46	^{1,3} ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$

NODE=M074R34
NODE=M074R34

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.06±0.03		^{1,2} ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$
1.38±0.11		^{1,3} ABLIKIM	20A BES3	$e^+ e^- \rightarrow \eta' J/\psi$

OCCUR=2

¹ Based on a fit to $\sigma(e^+ e^- \rightarrow \eta' J/\psi)$ from $\sqrt{s} = 4.18$ to 4.60 GeV assuming interfering $\psi(4160)$ and $\psi(4260)$ contributions. At $\sqrt{s} = 4.23$ GeV, $\sigma(e^+ e^- \rightarrow \eta' J/\psi) = 3.6 \pm 0.6 \pm 0.3$ pb.

² Solution I of the fit, corresponding to a phase of -0.03 ± 0.44 rad.

³ Solution II of the fit, corresponding to a phase of 2.54 ± 0.04 rad.

NODE=M074R34;LINKAGE=A

NODE=M074R34;LINKAGE=B

NODE=M074R34;LINKAGE=C

$$\Gamma(\psi(2S) \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}} \quad \Gamma_{17} \Gamma_1 / \Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
1.59±0.75		¹ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
1.63±0.78		² ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
0.02±0.01		³ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
1.6 ±1.3		⁴ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
1.8 ±1.4		⁵ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
<4.3	90	⁶ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^- \gamma$
7.4 ^{+2.1} -1.7		⁷ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \psi(2S) \pi^+ \pi^- \gamma$

NODE=M074G7
NODE=M074G7

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.59±0.75		¹ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
1.63±0.78		² ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
0.02±0.01		³ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
1.6 ±1.3		⁴ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
1.8 ±1.4		⁵ ABLIKIM	19K BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=2

¹ Solution I of four equivalent solutions in a fit using three interfering resonances.

² Solution II of four equivalent solutions in a fit using three interfering resonances

³ Solutions III and IV of four equivalent solutions in a fit using three interfering resonances.

⁴ Solution I of two equivalent solutions in a fit using two interfering resonances.

⁵ Solution II of two equivalent solutions in a fit using two interfering resonances.

⁶ For constructive interference with the $\psi(4360)$ in a combined fit of AUBERT 07S and WANG 07D data with three resonances.

⁷ For destructive interference with the $\psi(4360)$ in a combined fit of AUBERT 07S and WANG 07D data with three resonances.

NODE=M074G7;LINKAGE=A

NODE=M074G7;LINKAGE=B

NODE=M074G7;LINKAGE=C

NODE=M074G7;LINKAGE=AA

NODE=M074G7;LINKAGE=BB

NODE=M074G7;LINKAGE=LI

NODE=M074G7;LINKAGE=LU

$\Gamma(\chi_{c0}\omega) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_1/\Gamma$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	------	-------------	------	---------

2.5±0.2±0.3		¹ ABLIKIM	19A BES3	$e^+e^- \rightarrow \omega\chi_{c0}$
--------------------	--	----------------------	----------	--------------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

2.7±0.5±0.4	180	² ABLIKIM	15C BES3	$e^+e^- \rightarrow \omega\chi_{c0}$
-------------	-----	----------------------	----------	--------------------------------------

¹ From a fit of the measured cross section from $\sqrt{s} = 4.178\text{--}4.278$ GeV. Supersedes ABLIKIM 15C.

² From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.

NODE=M074G05
NODE=M074G05

NODE=M074G05;LINKAGE=B

NODE=M074G05;LINKAGE=A

 $\Gamma(h_c(1P)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{22}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

4.6^{+2.9}_{-1.4}±0.8	ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^-h_c$
--	---------	----------	------------------------------------

NODE=M074R47
NODE=M074R47

 $\Gamma(\phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{23}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.4	90	AUBERT,BE	06D BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^+\pi^-\gamma$
----------------	----	-----------	----------	--

NODE=M074G2
NODE=M074G2

 $\Gamma(\phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{24}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.28	90	¹ AUBERT	07AK BABR	10.6 $e^+e^- \rightarrow \pi^+\pi^-K^+K^-\gamma$
-----------------	----	---------------------	-----------	--

¹ AUBERT 07AK reports $[\Gamma(\psi(4230) \rightarrow \phi f_0(980) \rightarrow \phi\pi^+\pi^-) \times \Gamma(\psi(4230) \rightarrow e^+e^-)/\Gamma_{\text{total}}] \times [B(\phi(1020) \rightarrow K^+K^-)] < 0.14$ eV which we divide by our best value $B(\phi(1020) \rightarrow K^+K^-) = 49.1 \times 10^{-2}$.

NODE=M074G6
NODE=M074G6

NODE=M074G6;LINKAGE=AU

 $\Gamma(\phi K^+K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{25}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<1.75	90	ABLIKIM	23AE BES3	$e^+e^- \rightarrow \phi K^+K^-$
-----------------	----	---------	-----------	----------------------------------

NODE=M074R63
NODE=M074R63

 $\Gamma(\phi K_S^0 K_S^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{26}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.47	90	ABLIKIM	23AE BES3	$e^+e^- \rightarrow \phi K_S^0 K_S^0$
-----------------	----	---------	-----------	---------------------------------------

NODE=M074R64
NODE=M074R64

 $\Gamma(\phi\eta') \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{28}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.53	90	ABLIKIM	23R BES3	$e^+e^- \rightarrow \phi\eta'$
-----------------	----	---------	----------	--------------------------------

NODE=M074R61
NODE=M074R61

 $\Gamma(D^{*0}D^{*-}\pi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{45}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

5 to 22	¹ ABLIKIM	23X BES3	$e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$
---------	----------------------	----------	--

••• We do not use the following data for averages, fits, limits, etc. •••

¹ From a cross-section measurement of $e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. Depending on solutions I – VIII with the same fit qualities.

NODE=M074R65
NODE=M074R65

NODE=M074R65;LINKAGE=A

 $\Gamma(\omega\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{52}\Gamma_1/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

0.487 ± 0.008 ± 0.030	^{1,2} ABLIKIM	23AQ BES3	fit to cross sections
0.0005 ± 0.0003 ± 0.0001	^{2,3} ABLIKIM	23AQ BES3	fit to cross sections

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Solution I of the fit.

² From a fit to $e^+e^- \rightarrow \omega\pi^+\pi^-$ cross sections between 4 and 4.6 GeV. Recalculated from $12\pi\Gamma(e^+e^-)B(\psi(4230) \rightarrow \omega\pi^+\pi^-)$. First uncertainty is from statistical and uncommon systematic uncertainties, and the second is a 6.2% common systematic uncertainty quoted in the paper.

³ Solution II of the fit.

NODE=M074R62
NODE=M074R62

OCCUR=2

NODE=M074R62;LINKAGE=A
NODE=M074R62;LINKAGE=C

NODE=M074R62;LINKAGE=B

 $\Gamma(\Xi^-\Xi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{54}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

<0.0251 (CL = 90%)	$[<2.7 \times 10^{-4}$ eV (CL = 90%) OUR 2023 BEST LIMIT]			
------------------------------	---	--	--	--

<0.0251	90	¹ ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4230)$
-------------------	----	----------------------	-----------	---------------------------------

••• We do not use the following data for averages, fits, limits, etc. •••

$<2.7 \times 10^{-4}$	90	² ABLIKIM	20C BES3	$e^+e^- \rightarrow \Xi^-\Xi^+$
-----------------------	----	----------------------	----------	---------------------------------

¹ From a fit to $e^+e^- \rightarrow \Xi^-\Xi^+$ cross sections.

² Superseded by ABLIKIM 23BK.

NODE=M074R35
NODE=M074R35

NODE=M074R35;LINKAGE=A
NODE=M074R35;LINKAGE=B

$$\Gamma(\pi^+\pi^+\pi^-\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{55}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<32	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow 2\pi^+2\pi^-$

NODE=M074R37
NODE=M074R37

$$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{56}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow 2\pi^+2\pi^-\pi^0$

NODE=M074R38
NODE=M074R38

$$\Gamma(K_S^0 K^\pm \pi^\mp) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{59}\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M074G4
NODE=M074G4

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.04 ± 0.19 ± 0.09	¹ ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
0.0027 ± 0.0023 ± 0.0001	² ABLIKIM	19AE BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
< 0.5 at 90% CL	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \gamma$

OCCUR=2

¹ Solution I of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

NODE=M074G4;LINKAGE=A

² Solution II of the fit including the $\psi(4160)$ with mass 4191 ± 5 MeV and width 70 ± 10 MeV from PDG 16 and the $\psi(4230)$ with mass $4219.6 \pm 3.3 \pm 5.1$ MeV and width $56.0 \pm 3.6 \pm 6.9$ MeV from GAO 17.

NODE=M074G4;LINKAGE=B

$$\Gamma(K_S^0 K^\pm \pi^\mp \pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{60}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	ABLIKIM	19 BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \pi^0$

NODE=M074R31
NODE=M074R31

$$\Gamma(K_S^0 K^\pm \pi^\mp \eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{61}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.19	90	ABLIKIM	19 BES3	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp \eta$

NODE=M074R32
NODE=M074R32

$$\Gamma(K^+K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{62}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

NODE=M074G5
NODE=M074G5

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.6	90	AUBERT	08S BABR	10.6 $e^+e^- \rightarrow K^+K^-\pi^0\gamma$
------	----	--------	----------	---

$$\Gamma(K^+K^-\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{63}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow K^+K^-\pi^+\pi^-$

NODE=M074R39
NODE=M074R39

$$\Gamma(K^+K^-\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{64}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<43	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow K^+K^-\pi^+\pi^-\pi^0$

NODE=M074R40
NODE=M074R40

$$\Gamma(K^+K^+K^-K^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{65}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<3.8	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow 2K^+2K^-$

NODE=M074R41
NODE=M074R41

$$\Gamma(K^+K^+K^-K^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{66}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow 2K^+2K^-\pi^0$

NODE=M074R42
NODE=M074R42

$$\Gamma(p\bar{p}\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{67}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<7.2	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow p\bar{p}\pi^+\pi^-$

NODE=M074R43
NODE=M074R43

$$\Gamma(p\bar{p}\pi^+\pi^-\pi^0) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{68}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<15	90	ABLIKIM	21AW BES3	$e^+e^- \rightarrow p\bar{p}\pi^+\pi^-\pi^0$

NODE=M074R44
NODE=M074R44

$$\Gamma(\Lambda\bar{\Lambda}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{70}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.8 × 10 ⁻³	90	¹ ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4260)$

NODE=M074R36
NODE=M074R36

¹ From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

NODE=M074R36;LINKAGE=A

$$\Gamma(pK^-\bar{\Lambda} + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{71}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-3}$	90	¹ ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4230)$

¹ From a fit to $e^+e^- \rightarrow pK^-\bar{\Lambda} + \text{c.c.}$ cross sections.

NODE=M074R67
NODE=M074R67

NODE=M074R67;LINKAGE=A

$$\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{74}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.4	90	¹ HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M074G03
NODE=M074G03

NODE=M074G03;LINKAGE=A

$$\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{75}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<4.0	90	¹ HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M074G04
NODE=M074G04

NODE=M074G04;LINKAGE=A

$$\psi(4230) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$$

NODE=M074245

$$\Gamma(D^0 D^{*0}(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{42}/\Gamma \times \Gamma_1/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.42 \times 10^{-6}$	90	¹ PAKHLOVA	09 BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$

¹ Using 4263^{+8}_{-9} MeV for the mass of $\psi(4260)$.

NODE=M074R11
NODE=M074R11

NODE=M074R11;LINKAGE=PA

$\psi(4230)$ BRANCHING RATIOS

NODE=M074225

$$\Gamma(\eta_c(1S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	21B BES3	$e^+e^- \rightarrow \pi^+\pi^-\eta_c$

¹ Not seen in $e^+e^- \rightarrow \pi^+\pi^-\eta_c$ at $\sqrt{s} = 4.226$ GeV with a 90% C.L. upper limit on the cross section of 16.8 pb.

NODE=M074R56
NODE=M074R56

NODE=M074R56;LINKAGE=A

$$\Gamma(\eta_c(1S)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	21B BES3	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c$

¹ Seen as a peak in the $e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c$ cross section with a peak value of $46.1^{+9.5}_{-9.4} \pm 6.6$ pb at $\sqrt{s} = 4.226$ GeV.

NODE=M074R55
NODE=M074R55

NODE=M074R55;LINKAGE=A

$$\Gamma(J/\psi\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	DOCUMENT ID	TECN	COMMENT
seen	² ABLIKIM	17B BES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$

¹ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730-4.7008$ GeV.

² From a three-resonance fit. Superseded by ABLIKIM 22AM.

NODE=M074R51
NODE=M074R51

NODE=M074R51;LINKAGE=A

NODE=M074R51;LINKAGE=B

$$\Gamma(J/\psi f_0(980), f_0(980) \rightarrow \pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-) \quad \Gamma_6/\Gamma_5$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.13	¹ LEES	12AC BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$

¹ Systematic uncertainties not estimated.

NODE=M074R02
NODE=M074R02

NODE=M074R02;LINKAGE=LE

$$\Gamma(T_{cc1}(3900)^\pm \pi^\mp, T_{cc1}^\pm \rightarrow J/\psi\pi^\pm)/\Gamma(J/\psi\pi^+\pi^-) \quad \Gamma_7/\Gamma_5$$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.215 \pm 0.033 \pm 0.075$	¹ ABLIKIM	13T BES3	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	DOCUMENT ID	TECN	COMMENT
0.29 ± 0.08	² LIU	13B BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^-J/\psi$

¹ Assuming that the cross section of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ is fully due to the $\psi(4260)$.

² Systematic error not evaluated.

NODE=M074R01
NODE=M074R01

NODE=M074R01;LINKAGE=AB
NODE=M074R01;LINKAGE=A

$\Gamma(J/\psi\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	¹ ABLIKIM	20N	BES3 $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
------	----------------------	-----	---

¹ From a fit to the cross section $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$ at center-of-mass energies between 3.808 and 4.600 GeV.

NODE=M074R50
NODE=M074R50

NODE=M074R50;LINKAGE=A

 $\Gamma(J/\psi K_S^0 K_S^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	SHEN	14	BELL 9.4–10.9 $e^+e^- \rightarrow \gamma K_S^0 K_S^0 J/\psi$
----------	------	----	--

NODE=M074R27
NODE=M074R27 $\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	ABLIKIM	20O	BES3 $e^+e^- \rightarrow \eta J/\psi$
------	---------	-----	---------------------------------------

NODE=M074R52
NODE=M074R52 $\Gamma(J/\psi\eta\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	ABLIKIM	15Q	BES3 4.0–4.6 $e^+e^- \rightarrow J/\psi\eta\pi^0$
----------	---------	-----	---

NODE=M074R28
NODE=M074R28 $\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	¹ ABLIKIM	17V	BES3 $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
------	----------------------	-----	--

¹ From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^-\psi(2S) \rightarrow 2(\pi^+\pi^-\ell^+\ell^-)$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} .

NODE=M074R53
NODE=M074R53

NODE=M074R53;LINKAGE=A

 $\Gamma(\psi(2S)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{17}/Γ_5

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

($0.11 \pm 0.03 \pm 0.03$) to ($0.55 \pm 0.18 \pm 0.19$)	¹ ZHANG	17C	RVUE $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
--	--------------------	-----	---

¹ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

NODE=M074R30
NODE=M074R30

NODE=M074R30;LINKAGE=A

 $\Gamma(\chi_{c0}\omega)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen	180	¹ ABLIKIM	15C	BES3 $e^+e^- \rightarrow \omega\chi_{c0}$
------	-----	----------------------	-----	---

¹ From a 3-parameter fit of measured cross sections from $\sqrt{s} = 4.21\text{--}4.42$ GeV to a phase-space modified Breit-Wigner function, using the decays $\chi_{c0} \rightarrow \pi^+\pi^-$, $\chi_{c0} \rightarrow K^+K^-$, and $\omega \rightarrow \pi^+\pi^-\pi^0$.

NODE=M074R48
NODE=M074R48

NODE=M074R48;LINKAGE=A

 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	ABLIKIM	17G	BES3 $e^+e^- \rightarrow \pi^+\pi^- h_c$
------	---------	-----	--

NODE=M074R49
NODE=M074R49 $\Gamma(h_c(1P)\pi^+\pi^-)/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{22}/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.0	90	¹ PEDLAR	11	CLEO $e^+e^- \rightarrow h_c(1P)\pi^+\pi^-$
------	----	---------------------	----	---

¹ At $\sqrt{s} = 4260$ MeV, PEDLAR 11 measures $\sigma(e^+e^- \rightarrow h_c(1P)\pi^+\pi^-) = 32 \pm 17 \pm 6 \pm 6$ pb, where the errors are statistical, systematic, and due to uncertainty in $B(\psi(2S) \rightarrow \pi^0 h_c(1P))$, respectively.

NODE=M074R25
NODE=M074R25

NODE=M074R25;LINKAGE=PE

 $\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	ABLIKIM	23BT	BES3 $e^+e^- \rightarrow \phi\eta$
----------	---------	------	------------------------------------

NODE=M074R68
NODE=M074R68 $\Gamma(D\bar{D})/\Gamma(J/\psi\pi^+\pi^-)$ Γ_{29}/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.0	90	¹ AUBERT	07BE	BABR $e^+e^- \rightarrow D\bar{D}\gamma$
------	----	---------------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.0	90	CRONIN-HEN..09	CLEO	e^+e^-
------	----	----------------	------	----------

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$.

NODE=M074R2
NODE=M074R2

NODE=M074R2;LINKAGE=AU

$\Gamma(D^0\bar{D}^0)/\Gamma_{\text{total}}$					Γ_{30}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^0\bar{D}^0$		NODE=M074R12 NODE=M074R12
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		AUBERT	09M BABR	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$		
not seen		PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^0\bar{D}^0\gamma$		
$\Gamma(D^+D^-)/\Gamma_{\text{total}}$					Γ_{31}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^+D^-$		NODE=M074R13 NODE=M074R13
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		AUBERT	09M BABR	$e^+e^- \rightarrow D^+D^-\gamma$		
not seen		PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^+D^-\gamma$		
$\Gamma(D^*\bar{D}^0+c.c.)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{32}/Γ_5	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<34	90	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}^0$		NODE=M074R03 NODE=M074R03
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<45	90	CRONIN-HEN..09	CLEO	e^+e^-		
$\Gamma(D^*(2007)^0\bar{D}^0+c.c.)/\Gamma_{\text{total}}$					Γ_{33}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^0$		NODE=M074R14 NODE=M074R14
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^0\gamma$		
$\Gamma(D^*(2010)^+D^-+c.c.)/\Gamma_{\text{total}}$					Γ_{34}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^-$		NODE=M074R15 NODE=M074R15
not seen		PAKHLOVA	07 BELL	$e^+e^- \rightarrow D^{*+}D^-\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		AUBERT	09M BABR	$e^+e^- \rightarrow D^{*+}D^-\gamma$		
$\Gamma(D^*\bar{D}^*)/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{35}/Γ_5	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<11	90	CRONIN-HEN..09	CLEO	e^+e^-		NODE=M074R04 NODE=M074R04
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<40	90	AUBERT	09M BABR	$e^+e^- \rightarrow \gamma D^*\bar{D}^*$		
$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma_{\text{total}}$					Γ_{36}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}$		NODE=M074R17 NODE=M074R17
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0}\bar{D}^{*0}\gamma$		
$\Gamma(D^*(2010)^+D^*(2010)^-)/\Gamma_{\text{total}}$					Γ_{37}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^{*+}D^{*-}$		NODE=M074R18 NODE=M074R18
not seen		PAKHLOVA	07 BELL	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
not seen		AUBERT	09M BABR	$e^+e^- \rightarrow D^{*+}D^{*-}\gamma$		
$\Gamma(D^0D^-\pi^++c.c. (\text{excl. } D^*(2007)^0\bar{D}^{*0}+c.c., D^*(2010)^+D^-+c.c.))/\Gamma_{\text{total}}$					Γ_{39}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0D^-\pi^+\gamma$		NODE=M074R16 NODE=M074R16
$\Gamma(D\bar{D}^*\pi+c.c. (\text{excl. } D^*\bar{D}^*))/\Gamma_{\text{total}}$					Γ_{40}/Γ	
VALUE		DOCUMENT ID	TECN	COMMENT		
not seen		CRONIN-HEN..09	CLEO	$e^+e^- \rightarrow D^*\bar{D}^*\pi$		NODE=M074R22 NODE=M074R22
$\Gamma(D\bar{D}^*\pi+c.c. (\text{excl. } D^*\bar{D}^*))/\Gamma(J/\psi\pi^+\pi^-)$					Γ_{40}/Γ_5	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<15	90	CRONIN-HEN..09	CLEO	e^+e^-		NODE=M074R05 NODE=M074R05

$$\Gamma(D^0 D^{*-} \pi^+ + \text{c.c. (excl. } D^*(2010)^+ D^*(2010)^-))/\Gamma_{\text{total}} \quad \Gamma_{41}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+ \gamma$

NODE=M074R23
NODE=M074R23

$$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{42}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	ABLIKIM 19R	BES3	$e^+ e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$

NODE=M074R54
NODE=M074R54

$$\Gamma(D^0 D^*(2010)^- \pi^+ + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{42}/\Gamma_5$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9	90	PAKHLOVA 09	BELL	$e^+ e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M074R10
NODE=M074R10

$$\Gamma(D_1(2420) \bar{D} + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{43}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM 19AR	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$

NODE=M074R33
NODE=M074R33

¹ Results from a measurement of $\sigma(e^+ e^- \rightarrow D_1(2420) \bar{D} + \text{c.c.})$ between $\sqrt{s} = 4.3$ and 4.6 GeV.

NODE=M074R33;LINKAGE=A

$$\Gamma(D^* \bar{D}^* \pi)/\Gamma_{\text{total}} \quad \Gamma_{44}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen [not seen OUR 2023 BEST LIMIT]			

NODE=M074R24
NODE=M074R24

seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D^* \bar{D}^* \pi$
------	----------------	------	---

$$\Gamma(D^* \bar{D}^* \pi)/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{44}/\Gamma_5$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R06
NODE=M074R06

$$\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}} \quad \Gamma_{46}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

NODE=M074R19
NODE=M074R19

not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^+ D_s^-$
----------	----------------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
----------	-------------	------	--

$$\Gamma(D_s^+ D_s^-)/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{46}/\Gamma_5$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$

NODE=M074R07
NODE=M074R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3	90	CRONIN-HEN..09	CLEO	$e^+ e^-$
------	----	----------------	------	-----------

$$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_{47}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

NODE=M074R20
NODE=M074R20

not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^-$
----------	----------------	------	--------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
----------	-------------	------	---

$$\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma(J/\psi \pi^+ \pi^-) \quad \Gamma_{47}/\Gamma_5$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.8	90	CRONIN-HEN..09	CLEO	$e^+ e^-$

NODE=M074R08
NODE=M074R08

• • • We do not use the following data for averages, fits, limits, etc. • • •

<44	95	DEL-AMO-SA..10N	BABR	$10.6 e^+ e^-$
-----	----	-----------------	------	----------------

$$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}} \quad \Gamma_{48}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	CRONIN-HEN..09	CLEO	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$

NODE=M074R21
NODE=M074R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
----------	-------------	------	--

not seen	DEL-AMO-SA..10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
----------	-----------------	------	--

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma(J/\psi\pi^+\pi^-)$

 Γ_{48}/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 9.5	90	CRONIN-HEN..09	CLEO	e^+e^-
••• We do not use the following data for averages, fits, limits, etc. •••				
<30	95	DEL-AMO-SA..10N	BABR	$10.6 e^+e^-$

NODE=M074R09
 NODE=M074R09

$\Gamma(p\bar{p})/\Gamma(J/\psi\pi^+\pi^-)$

 Γ_{49}/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.13	90	¹ AUBERT	06B BABR	$e^+e^- \rightarrow p\bar{p}\gamma$

NODE=M074R1
 NODE=M074R1

¹ Using 4259 ± 10 MeV for the mass and 88 ± 24 MeV for the width of $\psi(4260)$.

NODE=M074R1;LINKAGE=AU

$\Gamma(p\bar{p}\pi^0)/\Gamma(J/\psi\pi^+\pi^-)$

 Γ_{50}/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2 × 10 ⁻⁴	90	ABLIKIM	17F BES3	$e^+e^- \rightarrow \psi(4260) \rightarrow$ hadrons

NODE=M074R00
 NODE=M074R00
 OCCUR=2

$\Gamma(p\bar{p}\eta)/\Gamma_{total}$

 Γ_{51}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	21AN BES3	$e^+e^- \rightarrow p\bar{p}\eta$

NODE=M074R45
 NODE=M074R45

$\Gamma(p\bar{p}\omega)/\Gamma_{total}$

 Γ_{53}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	21AN BES3	$e^+e^- \rightarrow p\bar{p}\omega$

NODE=M074R46
 NODE=M074R46

$\Gamma(\omega\pi^0)/\Gamma_{total}$

 Γ_{57}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	22K BES3	$e^+e^- \rightarrow \omega\pi^0$

NODE=M074R59
 NODE=M074R59

$\Gamma(\omega\eta)/\Gamma_{total}$

 Γ_{58}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	22K BES3	$e^+e^- \rightarrow \omega\eta$

NODE=M074R60
 NODE=M074R60

$\Gamma(p\bar{p}p\bar{p})/\Gamma_{total}$

 Γ_{69}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	21D BES3	$4.0-4.6 e^+e^- \rightarrow p\bar{p}p\bar{p}$

NODE=M074R58
 NODE=M074R58

Radiative decays

$\Gamma(\eta_c(1S)\gamma)/\Gamma_{total}$

 Γ_{72}/Γ

VALUE	DOCUMENT ID	COMMENT
possibly seen	¹ ABLIKIM	17W $e^+e^- \rightarrow \gamma\eta_c(1S)$

NODE=M074R29
 NODE=M074R29

¹ Significance ranges from 4.2σ to as low as 1.5σ for a flat component plus $\psi(4260)$ spectrum. Needs confirmation.

NODE=M074R29;LINKAGE=A

$\Gamma(\eta_c(1S)\pi^0\gamma)/\Gamma_{total}$

 Γ_{73}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	21B BES3	$e^+e^- \rightarrow \gamma\pi^0\eta_c$

NODE=M074R57
 NODE=M074R57

¹ Not seen in $e^+e^- \rightarrow \gamma\pi^0\eta_c$ at $\sqrt{s} = 4.226$ GeV with a 90% C.L. upper limit on the cross section of 11.2 pb.

NODE=M074R57;LINKAGE=A

$\Gamma(\chi_{c1}(3872)\gamma)/\Gamma_{total}$

 Γ_{76}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen		ABLIKIM	19v BES3	$e^+e^- \rightarrow \gamma\chi_{c1}(3872)$
seen	20 ± 5	ABLIKIM	14 BES3	$e^+e^- \rightarrow J/\psi\pi^+\pi^-\gamma$

NODE=M074R26
 NODE=M074R26

$\psi(4230)$ REFERENCES

ABLIKIM	23AE PR D108 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62092
ABLIKIM	23AQ JHEP 2308 159	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62313
ABLIKIM	23BK JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62437
ABLIKIM	23BL JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62438
ABLIKIM	23BT PR D108 112011	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62516
ABLIKIM	23R PR D107 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62062
ABLIKIM	23U PR D107 092005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62065
ABLIKIM	23X PRL 130 121901	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62072
ABLIKIM	22AM PR D106 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61885
ABLIKIM	22AU CP C46 111002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61896
ABLIKIM	22K JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	21AJ PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61441
ABLIKIM	21AN PR D104 092008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61446
ABLIKIM	21AS PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454

NODE=M074

ABLIKIM	21AW	PR D104 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61479
ABLIKIM	21B	PR D103 032006	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61029
ABLIKIM	21D	PR D103 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61031
ABLIKIM	20A	PR D101 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60210
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	20C	PRL 124 032002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60215
ABLIKIM	20N	PR D102 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60338
ABLIKIM	200	PR D102 031101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60344
ABLIKIM	19	PR D99 012003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59515
ABLIKIM	19AE	PR D99 072005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59856
ABLIKIM	19AI	PR D99 091103	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59871
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ABLIKIM	19K	PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	19R	PRL 122 102002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59765
ABLIKIM	19V	PRL 122 232002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59796
ABLIKIM	18N	PR D97 071101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58914
ABLIKIM	17B	PRL 118 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57755
ABLIKIM	17F	PL B771 45	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57909
ABLIKIM	17G	PRL 118 092002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57915
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58029
Also		PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
ABLIKIM	17W	PR D96 051101	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58212
GAO	17	PR D95 092007	X.Y. Gao, C.P. Shen, C.Z. Yuan		REFID=57991
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang		REFID=58219
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan		REFID=58463
PDG	16	CP C40 100001	C. Patrignani <i>et al.</i>	(PDG Collab.)	REFID=57140
ABLIKIM	15C	PRL 114 092003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56401
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56782
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
ABLIKIM	14	PRL 112 092001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55647
SHEN	14	PR D89 072015	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55944
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55409
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
LEES	12AC	PR D86 051102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54750
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
PEDLAR	11	PRL 107 041803	T. Pedlar <i>et al.</i>	(CLEO Collab.)	REFID=16787
DEL-AMO-SA...	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
CRONIN-HEN...	09	PR D80 072001	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=53114
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
AUBERT	08S	PR D77 092002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52242
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=52296
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
YUAN	08	PR D77 011105	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=52135
AUBERT	07AK	PR D76 012008	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51908
AUBERT	07BE	PR D76 111105	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52074
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51724
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51959
YUAN	07	PRL 99 182004	C.Z. Yuan <i>et al.</i>	(BELLE Collab.)	REFID=51960
AUBERT	06B	PR D73 012005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51026
AUBERT,BE	06D	PR D74 091103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51511
HE	06B	PR D74 091104	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=51523
AUBERT,B	05I	PRL 95 142001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50776

$\chi_{c1}(4274)$

$$J^G(J^{PC}) = 0^+(1^{++})$$

was $X(4274)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen by AAIJ 17C in $B^+ \rightarrow \chi_{c1} K^+$, $\chi_{c1} \rightarrow J/\psi \phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ with a significance (accounting for systematic uncertainties) of 6.0σ .

NODE=M233

NODE=M233

 $\chi_{c1}(4274)$ MASS

NODE=M233M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4286 $^{+8}_{-9}$ OUR AVERAGE		Error includes scale factor of 1.7.		
4294 $\pm 4 \pm \frac{3}{6}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
4274.4 $^{+8.4}_{-6.7} \pm 1.9$	22	² AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4273.3 $\pm 8.3 \pm \frac{17.2}{3.6}$	4289	^{3,4} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 18σ .				
² From a fit to the invariant mass spectrum with a significance of 3.1σ .				
³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.0σ .				
⁴ Superseded by AAIJ 21E.				

NODE=M233M

NODE=M233M;LINKAGE=C
NODE=M233M;LINKAGE=B
NODE=M233M;LINKAGE=A
NODE=M233M;LINKAGE=D

 $\chi_{c1}(4274)$ WIDTH

NODE=M233W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
51 ± 7 OUR AVERAGE				
53 $\pm 5 \pm 5$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
32.3 $^{+21.9}_{-15.3} \pm 7.6$	22	² AALTONEN	17 CDF	$B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
56 $\pm 11 \pm \frac{8}{11}$	4289	^{3,4} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 18σ .				
² From a fit to the invariant mass spectrum with a significance of 3.1σ .				
³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.0σ .				
⁴ Superseded by AAIJ 21E.				

NODE=M233W

NODE=M233W;LINKAGE=C
NODE=M233W;LINKAGE=B
NODE=M233W;LINKAGE=A
NODE=M233W;LINKAGE=D

 $\chi_{c1}(4274)$ DECAY MODES

NODE=M233215;NODE=M233

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi \phi$	seen

DESIG=1

 $\chi_{c1}(4274)$ BRANCHING RATIOS

NODE=M233220

$\Gamma(J/\psi \phi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	4289	^{2,3} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$
¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 18σ .				
² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.0σ .				
³ Superseded by AAIJ 21E.				

NODE=M233R01
NODE=M233R01

NODE=M233R01;LINKAGE=B
NODE=M233R01;LINKAGE=A
NODE=M233R01;LINKAGE=C

 $\chi_{c1}(4274)$ REFERENCES

NODE=M233

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	17	MPL A32 1750139	T. Altonen <i>et al.</i>	(CDF Collab.)

REFID=61150
REFID=57657
REFID=57636
REFID=58161

X(4350)

$$I^G(J^{PC}) = 0^+(?^{?+})$$

OMITTED FROM SUMMARY TABLE

Seen by SHEN 10 in the $\gamma\gamma \rightarrow J/\psi\phi$. Needs confirmation.

NODE=M194

NODE=M194

NODE=M194M

NODE=M194M

NODE=M194M;LINKAGE=SH

NODE=M194W

NODE=M194W

NODE=M194W;LINKAGE=SH

NODE=M194215;NODE=M194

DESIG=1

DESIG=2

NODE=M194220

NODE=M194G01
NODE=M194G01

OCCUR=2

NODE=M194G01;LINKAGE=S0
NODE=M194G01;LINKAGE=S2

NODE=M194225

NODE=M194R01
NODE=M194R01

NODE=M194R01;LINKAGE=SH

NODE=M194R02
NODE=M194R02

NODE=M194R02;LINKAGE=SH

NODE=M194

REFID=53235

X(4350) MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4350.6^{+4.6}_{-5.1} \pm 0.7$	$8.8^{+4.2}_{-3.2}$	¹ SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

¹ Statistical significance of 3.2 σ .**X(4350) WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$13^{+18}_{-9} \pm 4$	$8.8^{+4.2}_{-3.2}$	¹ SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$

¹ Statistical significance of 3.2 σ .**X(4350) DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen
Γ_2 $\gamma\gamma$	seen

X(4350) $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_1/\Gamma$
$6.7^{+3.2}_{-2.4} \pm 1.1$	$8.8^{+4.2}_{-3.2}$	¹ SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.5^{+0.7}_{-0.6} \pm 0.3$	$8.8^{+4.2}_{-3.2}$	² SHEN	10	BELL 10.6 $e^+e^- \rightarrow e^+e^- J/\psi\phi$	
-----------------------------	---------------------	-------------------	----	--	--

¹ For $J^P = 0^+$. Statistical significance of 3.2 σ .² For $J^P = 2^+$. Statistical significance of 3.2 σ .

NODE=M194220

NODE=M194G01
NODE=M194G01

OCCUR=2

NODE=M194G01;LINKAGE=S0
NODE=M194G01;LINKAGE=S2**X(4350) BRANCHING RATIOS**

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	

¹ Statistical significance of 3.2 σ .NODE=M194R01
NODE=M194R01

NODE=M194R01;LINKAGE=SH

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_2/Γ
seen	

¹ Statistical significance of 3.2 σ .NODE=M194R02
NODE=M194R02

NODE=M194R02;LINKAGE=SH

X(4350) REFERENCESSHEN 10 PRL 104 112004 C.P. Shen *et al.* (BELLE Collab.)

NODE=M194

REFID=53235

$\psi(4360)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $Y(4360)$; was $X(4360)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by AUBERT 07S, WANG 07D, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

NODE=M181

NODE=M181

 $\psi(4360)$ MASS

NODE=M181M

NODE=M181M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4374 ± 7	OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.		
4371.6 ± 2.5 ± 9.2		1 ABLIKIM	22AL BES3	$e^+e^- \rightarrow \pi^+\pi^- D^+D^-$
4298 ± 12 ± 26		2 ABLIKIM	22AMBES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
4390.3 ± 6.0 ± 0.7		3 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
4371.7 ± 7.5 ± 1.8		4 ABLIKIM	21AK BES3	$e^+e^- \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi$
4382.0 ± 13.3 ± 1.7		5 ABLIKIM	20O BES3	$e^+e^- \rightarrow \eta J/\psi$
4391.5 ^{+6.3} _{-6.8} ± 1.0		ABLIKIM	17G BES3	$e^+e^- \rightarrow \pi^+\pi^- h_c$
4347 ± 6 ± 3	279	6 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
4340 ± 16 ± 9	37	7 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4406.9 ± 17.2 ± 4.5		8 ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^- \chi_{c1}\gamma$
4320.0 ± 10.4 ± 7.0		9 ABLIKIM	17B BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
4383.8 ± 4.2 ± 0.8		10 ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
4383.7 ± 2.9 ± 6.2		11 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
4386.4 ± 2.1 ± 6.4		12 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
4355 ⁺⁹ ₋₁₀ ± 9	74	13 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
4324 ± 24		14 AUBERT	07S BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
4361 ± 9 ± 9	47	7 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$

NODE=M181M;LINKAGE=M

NODE=M181M;LINKAGE=L

NODE=M181M;LINKAGE=H

NODE=M181M;LINKAGE=G

NODE=M181M;LINKAGE=BA

NODE=M181M;LINKAGE=A

NODE=M181M;LINKAGE=WA

NODE=M181M;LINKAGE=J

NODE=M181M;LINKAGE=K

NODE=M181M;LINKAGE=C

NODE=M181M;LINKAGE=E

NODE=M181M;LINKAGE=D

NODE=M181M;LINKAGE=LI

NODE=M181M;LINKAGE=AU

¹ From a fit to the cross section for $e^+e^- \rightarrow D^+D^-\pi^+\pi^-$ in the range $\sqrt{s} = 4.190\text{--}4.946$ GeV.

² From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730\text{--}4.7008$ GeV. Parameters depend on the existence or non-existence of a state near 4.5 GeV.

³ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.

⁴ From a five-resonance fit to the cross section for $e^+e^- \rightarrow \gamma\gamma J/\psi \rightarrow \gamma\gamma\ell^+\ell^-$.

⁵ From a fit of the measured cross section in the range $\sqrt{s} = 3.808\text{--}4.600$ GeV.

⁶ From a two-resonance fit. Supersedes WANG 07D.

⁷ From a two-resonance fit.

⁸ From a fit to the $e^+e^- \rightarrow \pi^+\pi^- \psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.

⁹ From a three-resonance fit. Superseded by ABLIKIM 22AM.


¹⁰ From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^- \psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} . Superseded by ABLIKIM 21AJ.

¹¹ From a three-resonance fit.

¹² From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$ data.

¹³ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

¹⁴ From a single-resonance fit. Systematic errors not estimated.



psi(4360) WIDTH

NODE=M181W

NODE=M181W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
118 ± 12 OUR AVERAGE		Error includes scale factor of 2.1. See the ideogram below.		
167 ± 4 ± 29		1 ABLIKIM 22AL BES3		$e^+e^- \rightarrow \pi^+\pi^- D^+D^-$
127 ± 17 ± 10		2 ABLIKIM 22AMBES3		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
143.3 ± 10.0 ± 0.5		3 ABLIKIM 21AJ BES3		$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
51.1 ± 17.6 ± 1.9		4 ABLIKIM 21AK BES3		$e^+e^- \rightarrow \gamma\chi_{c2} \rightarrow \gamma\gamma J/\psi$
135.8 ± 60.8 ± 22.5		5 ABLIKIM 20O BES3		$e^+e^- \rightarrow \eta J/\psi$
139.5 ^{+16.2} _{-20.6} ± 0.6		ABLIKIM 17G BES3		$e^+e^- \rightarrow \pi^+\pi^- h_c$
103 ± 9 ± 5	279	6 WANG 15A BELL		10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
94 ± 32 ± 13	37	7 LEES 14F BABR		10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
128.1 ± 37.2 ± 2.3		8 ABLIKIM 22R BES3		$e^+e^- \rightarrow \pi^+\pi^- \chi_{c1}\gamma$
101.4 ^{+25.3} _{-19.7} ± 10.2		9 ABLIKIM 17B BES3		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
84.2 ± 12.5 ± 2.1		10 ABLIKIM 17V BES3		$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
94.2 ± 7.3 ± 2.0		11 ZHANG 17B RVUE		$e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$
96.0 ± 6.7 ± 2.7		12 ZHANG 17C RVUE		$e^+e^- \rightarrow \pi^+\pi^- J/\psi$ or $\psi(2S)$
103 ⁺¹⁷ ₋₁₅ ± 11	74	13 LIU 08H RVUE		10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
172 ± 33		14 AUBERT 07S BABR		10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$
74 ± 15 ± 10	47	7 WANG 07D BELL		10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^- \psi(2S)$

NODE=M181W;LINKAGE=L

NODE=M181W;LINKAGE=K

NODE=M181W;LINKAGE=G

NODE=M181W;LINKAGE=F

NODE=M181W;LINKAGE=BA

NODE=M181W;LINKAGE=A

NODE=M181W;LINKAGE=WA

NODE=M181W;LINKAGE=I

NODE=M181W;LINKAGE=J

NODE=M181W;LINKAGE=C

NODE=M181W;LINKAGE=E


NODE=M181W;LINKAGE=D

NODE=M181W;LINKAGE=LI

- 1 From a fit to the cross section for $e^+e^- \rightarrow D^+D^-\pi^+\pi^-$ in the range $\sqrt{s} = 4.190-4.946$ GeV.
- 2 From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 3.7730-4.7008$ GeV. Parameters depend on the existence or non-existence of a state near 4.5 GeV.
- 3 From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008-4.698$ GeV.
- 4 From a five-resonance fit to the cross section for $e^+e^- \rightarrow \gamma\gamma J/\psi \rightarrow \gamma\gamma\ell^+\ell^-$.
- 5 From a fit of the measured cross section in the range $\sqrt{s} = 3.808-4.600$ GeV.
- 6 From a two-resonance fit. Supersedes WANG 07D.
- 7 From a two-resonance fit.
- 8 From a fit to the $e^+e^- \rightarrow \pi^+\pi^- \psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.
- 9 From a three-resonance fit. Superseded by ABLIKIM 22AM.
- 10 From a fit to the cross section for $e^+e^- \rightarrow \pi^+\pi^- \psi(2S) \rightarrow 2(\pi^+\pi^-)\ell^+\ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} . Superseded by ABLIKIM 21AJ.
- 11 From a three-resonance fit.
- 12 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^- \psi(2S)$ data.
- 13 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

¹⁴ From a single-resonance fit. Systematic errors not estimated.

NODE=M181W;LINKAGE=AU



psi(4360) DECAY MODES

NODE=M181215;NODE=M181

Mode	Fraction (Γ_i/Γ)	
Γ_1 $e^+ e^-$		DESIG=1
Γ_2 $h_c \pi^+ \pi^-$	seen	DESIG=12
Γ_3 $J/\psi \pi^+ \pi^-$		DESIG=8
Γ_4 $\psi(2S) \pi^+ \pi^-$	seen	DESIG=2
Γ_5 $\psi(3770) \pi^+ \pi^-$	possibly seen	DESIG=11
Γ_6 $\psi_2(3823) \pi^+ \pi^-$	seen	DESIG=5
Γ_7 $J/\psi \eta$	seen	DESIG=4
Γ_8 $D^0 D^{*-} \pi^+$		DESIG=3
Γ_9 $D^+ D^- \pi^+ \pi^-$	seen	DESIG=17
Γ_{10} $D_1(2420) \bar{D} + c.c.$	possibly seen	DESIG=10
Γ_{11} $\phi \eta$	not seen	DESIG=20
Γ_{12} $\omega \pi^0$	not seen	DESIG=15
Γ_{13} $\omega \eta$	not seen	DESIG=16
Γ_{14} $\rho \bar{\rho} \eta$	not seen	DESIG=13
Γ_{15} $\rho \bar{\rho} \omega$	not seen	DESIG=14
Γ_{16} $\chi_{c1} \gamma$		DESIG=6
Γ_{17} $\chi_{c2} \gamma$		DESIG=7
Γ_{18} $\Xi^- \Xi^+$		DESIG=18
Γ_{19} $\rho K^- \bar{\Lambda} + c.c.$		DESIG=19

psi(4360) $\Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

NODE=M181230

$\Gamma(h_c \pi^+ \pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$	$\Gamma_2 \Gamma_1 / \Gamma$	
VALUE (eV)	DOCUMENT ID	TECN COMMENT
$11.6^{+5.0}_{-4.4} \pm 1.9$	ABLIKIM	17G BES3 $e^+ e^- \rightarrow \pi^+ \pi^- h_c$

NODE=M181R11
NODE=M181R11

$\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_4\Gamma_1/\Gamma$

VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
10.7±4.1		1 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
20.7±2.5		2 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
9.9±4.1		3 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
19.4±2.0		4 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
7.3±2.8		5 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
11.0±3.8		6 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
9.2±0.6±0.6	279	7 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
10.9±0.6±0.7	279	8 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
6.0±1.0±0.5	37	5 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
7.2±1.0±0.6	37	6 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
11.1 ^{+1.3} _{-1.2}	74	9 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
12.3±1.2	74	10 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
10.4±1.7±1.5	47	5 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
11.8±1.8±1.4	47	6 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

NODE=M181G1
 NODE=M181G1

• • • We do not use the following data for averages, fits, limits, etc. • • •

10.7±4.1		1 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
20.7±2.5		2 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=2
9.9±4.1		3 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=3
19.4±2.0		4 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=4
7.3±2.8		5 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	
11.0±3.8		6 ABLIKIM	19K BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$	OCCUR=2
9.2±0.6±0.6	279	7 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
10.9±0.6±0.7	279	8 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2
6.0±1.0±0.5	37	5 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
7.2±1.0±0.6	37	6 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2
11.1 ^{+1.3} _{-1.2}	74	9 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
12.3±1.2	74	10 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2
10.4±1.7±1.5	47	5 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	
11.8±1.8±1.4	47	6 WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$	OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

¹ Solution I of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

NODE=M181G1;LINKAGE=E

² Solution II of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

NODE=M181G1;LINKAGE=F

³ Solution III of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

NODE=M181G1;LINKAGE=G

⁴ Solution IV of four equivalent solutions in a fit using three interfering resonances. Supersedes ABLIKIM 19K.

NODE=M181G1;LINKAGE=H

⁵ Solution I of two equivalent solutions in a fit using two interfering resonances.

NODE=M181G1;LINKAGE=WA

⁶ Solution II of two equivalent solutions in a fit using two interfering resonances.

NODE=M181G1;LINKAGE=WN

⁷ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

NODE=M181G1;LINKAGE=A

⁸ Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.

NODE=M181G1;LINKAGE=B

⁹ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M181G1;LINKAGE=LI

¹⁰ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M181G1;LINKAGE=LU

 $\Gamma(J/\psi\eta) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_7\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
3.4±2.2		1 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$
1.5±1.0		2 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$
1.7±1.1		3 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$
<6.8	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

NODE=M181G01
 NODE=M181G01

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.4±2.2		1 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$	
1.5±1.0		2 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$	OCCUR=2
1.7±1.1		3 ABLIKIM	200 BES3	$e^+e^- \rightarrow \eta J/\psi$	OCCUR=3
<6.8	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$	

¹ Solution 1 of three equivalent fit solutions using three resonant structures.

NODE=M181G01;LINKAGE=A

² Solution 2 of three equivalent fit solutions using three resonant structures.

NODE=M181G01;LINKAGE=B

³ Solution 3 of three equivalent fit solutions using three resonant structures.

NODE=M181G01;LINKAGE=C

 $\Gamma(\chi_{c1}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{16}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.57	90	1 HAN	15 BELL	10.58 $e^+e^- \rightarrow \chi_{c1}\gamma$

NODE=M181G02
 NODE=M181G02

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M181G02;LINKAGE=A

 $\Gamma(\chi_{c2}\gamma) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{17}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.9	90	1 HAN	15 BELL	10.58 $e^+e^- \rightarrow \chi_{c2}\gamma$

NODE=M181G03
 NODE=M181G03

¹ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M181G03;LINKAGE=A

 $\Gamma(\Xi^-\Xi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{18}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.0448	90	1 ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4360)$

NODE=M181R15
 NODE=M181R15

¹ From a fit to $e^+e^- \rightarrow \Xi^-\Xi^+$ cross sections.

NODE=M181R15;LINKAGE=A

 $\Gamma(pK^-\bar{\Lambda} + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{19}\Gamma_1/\Gamma$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<4.7 × 10 ⁻³	90	1 ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4360)$

NODE=M181R17
 NODE=M181R17

¹ From a fit to $e^+e^- \rightarrow pK^-\bar{\Lambda} + \text{c.c.}$ cross sections.

NODE=M181R17;LINKAGE=A

$\psi(4360)$ BRANCHING RATIOS

					NODE=M181225	
$\Gamma(h_c \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_2/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R08	
seen	ABLIKIM	17G BES3	$e^+ e^- \rightarrow \pi^+ \pi^- h_c$		NODE=M181R08	
$\Gamma(\psi(2S) \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_4/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R00	
seen	¹ ABLIKIM	17V BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$		NODE=M181R00	
¹ From a fit to the cross section for $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S) \rightarrow 2(\pi^+ \pi^-) \ell^+ \ell^-$ obtained from 16 center-of-mass energies between 4.008 and 4.600 GeV and comprising 5.1 fb^{-1} .					NODE=M181R00;LINKAGE=A	
$\Gamma(\psi(2S) \pi^+ \pi^-)/\Gamma(J/\psi \pi^+ \pi^-)$					Γ_4/Γ_3	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R04	
• • • We do not use the following data for averages, fits, limits, etc. • • •					NODE=M181R04	
($0.81 \pm 0.12 \pm 0.13$) to ($42 \pm 15 \pm 15$)	¹ ZHANG	17C RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$			
¹ From a combined fit of BELLE, BABAR and BES3 $e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ and $e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$ data.					NODE=M181R04;LINKAGE=A	
$\Gamma(\psi(3770) \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_5/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R06	
possibly seen	¹ ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$		NODE=M181R06	
¹ Observe $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$ at $\sqrt{s} = 4.26, 4.36, \text{ and } 4.42 \text{ GeV}$ but cannot establish if continuum or resonant.					NODE=M181R06;LINKAGE=A	
$\Gamma(\psi_2(3823) \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_6/Γ	
<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M181R03	
seen	¹ ABLIKIM	22R BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$		NODE=M181R03	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
possibly seen	19	² ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$		
¹ From a fit to the $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances.					NODE=M181R03;LINKAGE=C	
² From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \psi_2(3823), \psi_2(3823) \rightarrow \chi_{c1} \gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4360)$ line shape.					NODE=M181R03;LINKAGE=A	
$\Gamma(J/\psi \eta)/\Gamma_{\text{total}}$					Γ_7/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R07	
seen	¹ ABLIKIM	200 BES3	$e^+ e^- \rightarrow \eta J/\psi$		NODE=M181R07	
¹ With a significance of 6.0σ .					NODE=M181R07;LINKAGE=A	
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$					$\Gamma_8/\Gamma \times \Gamma_1/\Gamma$	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M181R02	
$<0.72 \times 10^{-6}$	90	¹ PAKHLOVA	09 BELL	$e^+ e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-} \pi^+$	NODE=M181R02	
¹ Using $4355_{-10}^{+9} \pm 9 \text{ MeV}$ for the mass of $\psi(4360)$.					NODE=M181R02;LINKAGE=PA	
$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S) \pi^+ \pi^-)$					Γ_8/Γ_4	
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>	NODE=M181R01	
<8	90	PAKHLOVA	09 BELL	$e^+ e^- \rightarrow \psi(4360) \rightarrow D^0 D^{*-} \pi^+$	NODE=M181R01	
$\Gamma(D^+ D^- \pi^+ \pi^-)/\Gamma_{\text{total}}$					Γ_9/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R14	
seen	¹ ABLIKIM	22AL BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D^+ D^-$		NODE=M181R14	
¹ From a fit to the cross section for $e^+ e^- \rightarrow D^+ D^- \pi^+ \pi^-$ in the range $\sqrt{s} = 4.190\text{--}4.946 \text{ GeV}$.					NODE=M181R14;LINKAGE=A	
$\Gamma(D_1(2420) \bar{D} + \text{c.c.})/\Gamma_{\text{total}}$					Γ_{10}/Γ	
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		NODE=M181R05	
possibly seen	¹ ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$		NODE=M181R05	
¹ Evidence for $e^+ e^- \rightarrow D_1(2420) \bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV , not necessarily resonant.					NODE=M181R05;LINKAGE=A	

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$	Γ_{11}/Γ
VALUE	
not seen	
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$	Γ_{12}/Γ
VALUE	
not seen	
$\Gamma(\omega\eta)/\Gamma_{\text{total}}$	Γ_{13}/Γ
VALUE	
not seen	
$\Gamma(\rho\bar{p}\eta)/\Gamma_{\text{total}}$	Γ_{14}/Γ
VALUE	
not seen	
$\Gamma(\rho\bar{p}\omega)/\Gamma_{\text{total}}$	Γ_{15}/Γ
VALUE	
not seen	

$\psi(4360)$ REFERENCES

DOCUMENT ID	TECN	COMMENT	NODE=M181
ABLIKIM 23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	REFID=62437
ABLIKIM 23BL	JHEP 2312 027	M. Ablikim <i>et al.</i>	REFID=62438
ABLIKIM 23BT	PR D108 112011	M. Ablikim <i>et al.</i>	REFID=62516
ABLIKIM 22AL	PR D106 052012	M. Ablikim <i>et al.</i>	REFID=61884
ABLIKIM 22AM	PR D106 072001	M. Ablikim <i>et al.</i>	REFID=61885
ABLIKIM 22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	REFID=61648
ABLIKIM 22R	PRL 129 102003	M. Ablikim <i>et al.</i>	REFID=61664
ABLIKIM 21AJ	PR D104 052012	M. Ablikim <i>et al.</i>	REFID=61441
ABLIKIM 21AK	PR D104 092001	M. Ablikim <i>et al.</i>	REFID=61443
ABLIKIM 21AN	PR D104 092008	M. Ablikim <i>et al.</i>	REFID=61446
ABLIKIM 200	PR D102 031101	M. Ablikim <i>et al.</i>	REFID=60344
ABLIKIM 19AR	PR D100 032005	M. Ablikim <i>et al.</i>	REFID=59910
ABLIKIM 19K	PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	REFID=59611
ABLIKIM 17B	PRL 118 092001	M. Ablikim <i>et al.</i>	REFID=57755
ABLIKIM 17G	PRL 118 092002	M. Ablikim <i>et al.</i>	REFID=57915
ABLIKIM 17V	PR D96 032004	M. Ablikim <i>et al.</i>	REFID=58029
Also	PR D99 019903 (errata.)	M. Ablikim <i>et al.</i>	REFID=59611
ZHANG 17B	PR D96 054008	J. Zhang, J. Zhang	REFID=58219
ZHANG 17C	EPJ C77 727	J. Zhang, L. Yuan	REFID=58463
ABLIKIM 15S	PRL 115 011803	M. Ablikim <i>et al.</i>	REFID=56784
HAN 15	PR D92 012011	Y.L. Han <i>et al.</i>	REFID=56816
WANG 15A	PR D91 112007	X.L. Wang <i>et al.</i>	REFID=56839
LEES 14F	PR D89 111103	J.P. Lees <i>et al.</i>	REFID=55938
WANG 13B	PR D87 051101	X.L. Wang <i>et al.</i>	REFID=55377
PAKHLOVA 09	PR D80 091101	G. Pakhlova <i>et al.</i>	REFID=53143
LIU 08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan	REFID=52296
AUBERT 07S	PRL 98 212001	B. Aubert <i>et al.</i>	REFID=51724
WANG 07D	PRL 99 142002	X.L. Wang <i>et al.</i>	REFID=51959

$\psi(4415)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M073

 $\psi(4415)$ MASS

NODE=M073M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4421 ± 4 OUR ESTIMATE			
4415 ± 5 OUR AVERAGE			
[4415 ± 8 MeV OUR 2023 AVERAGE]			
4414.6 ± 3.4 ± 6.1	ABLIKIM	23BH BES3	$e^+e^- \rightarrow D_s^{*+} D_s^{*-}$
4415.1 ± 7.9	¹ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
4412 ± 15	² MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
4411 ± 7	³ PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
4425 ± 6	⁴ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4429 ± 9	⁵ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
4417 ± 10	BRANDELIK	78C DASP	e^+e^-
4414 ± 7	SIEGRIST	76 MRK1	e^+e^-
¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.			
² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
³ Systematic uncertainties not estimated.			
⁴ From a fit to Crystal Ball (OSTERHELD 86) data.			
⁵ From a fit to BES (BAI 02C) data.			

NODE=M073M
 → UNCHECKED ←
 NEW

OCCUR=2

NODE=M073M;LINKAGE=AB

NODE=M073M;LINKAGE=MO

NODE=M073M;LINKAGE=NS
 NODE=M073M;LINKAGE=ST
 NODE=M073M;LINKAGE=SE

 $\psi(4415)$ WIDTH

NODE=M073W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
62 ± 20 OUR ESTIMATE			
110 ± 22 OUR AVERAGE			
Error includes scale factor of 2.3. [72 ± 19 MeV OUR 2023 AVERAGE]			
122.5 ± 7.5 ± 8.1	ABLIKIM	23BH BES3	$e^+e^- \rightarrow D_s^{*+} D_s^{*-}$
71.5 ± 19.0	⁶ ABLIKIM	08D BES2	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
118 ± 32	⁷ MO	10 RVUE	$e^+e^- \rightarrow$ hadrons
77 ± 20	⁸ PAKHLOVA	08A BELL	$10.6 e^+e^- \rightarrow D^0 D^- \pi^+ \gamma$
119 ± 16	⁹ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
118 ± 35	¹⁰ SETH	05A RVUE	$e^+e^- \rightarrow$ hadrons
66 ± 15	BRANDELIK	78C DASP	e^+e^-
33 ± 10	SIEGRIST	76 MRK1	e^+e^-
⁶ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.			
⁷ Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects.			
⁸ Systematic uncertainties not estimated.			
⁹ From a fit to Crystal Ball (OSTERHELD 86) data.			
¹⁰ From a fit to BES (BAI 02C) data.			

NODE=M073W
 → UNCHECKED ←
 NEW

OCCUR=2

NODE=M073W;LINKAGE=AB

NODE=M073W;LINKAGE=MO

NODE=M073W;LINKAGE=NS
 NODE=M073W;LINKAGE=ST
 NODE=M073W;LINKAGE=SE

 $\psi(4415)$ DECAY MODES

NODE=M073215;NODE=M073

Due to the complexity of the $c\bar{c}$ threshold region, in this listing, “seen” (“not seen”) means that a cross section for the mode in question has been measured at effective \sqrt{s} near this particle’s central mass value, more

NODE=M073

(less) than 2σ above zero, without regard to any peaking behavior in \sqrt{s} or absence thereof. See mode listing(s) for details and references.

Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1 $D\bar{D}$	seen		DESIG=7;OUR EVAL;→ UNCHECKED ←
Γ_2 $D^0\bar{D}^0$	seen		DESIG=8
Γ_3 D^+D^-	seen		DESIG=9
Γ_4 $D^*\bar{D} + \text{c.c.}$	seen		DESIG=10;OUR EVAL;→ UNCHECKED ←
Γ_5 $D^*(2007)^0\bar{D}^0 + \text{c.c.}$	seen		DESIG=11
Γ_6 $D^*(2010)^+D^- + \text{c.c.}$	seen		DESIG=12
Γ_7 $D^*\bar{D}^*$	seen		DESIG=13;OUR EVAL;→ UNCHECKED ←
Γ_8 $D^*(2007)^0\bar{D}^*(2007)^0 + \text{c.c.}$	seen		DESIG=14
Γ_9 $D^*(2010)^+D^*(2010)^- + \text{c.c.}$	seen		DESIG=15
Γ_{10} $D^0D^-\pi^+$ (excl. $D^*(2007)^0\bar{D}^0$ +c.c., $D^*(2010)^+D^-$ +c.c.)	< 2.3 %	90%	DESIG=4
Γ_{11} $D\bar{D}_2^*(2460) \rightarrow D^0D^-\pi^+ + \text{c.c.}$	(10 ± 4) %		DESIG=5
Γ_{12} $D^0D^{*-}\pi^+ + \text{c.c.}$	< 11 %	90%	DESIG=6
Γ_{13} $D_1(2420)\bar{D} + \text{c.c.}$	possibly seen		DESIG=25
Γ_{14} $D_s^+D_s^-$	not seen		DESIG=16
Γ_{15} $\omega\chi_{c2}$	possibly seen		DESIG=20
Γ_{16} $D_s^{*+}D_s^- + \text{c.c.}$	seen		DESIG=17
Γ_{17} $D_s^{*+}D_s^{*-}$	seen		DESIG=18
Γ_{18} $\psi_2(3823)\pi^+\pi^-$	possibly seen		DESIG=21
Γ_{19} $\psi(3770)\pi^+\pi^-$	possibly seen		DESIG=24
Γ_{20} $J/\psi\eta$	< 6 × 10 ⁻³	90%	DESIG=19
Γ_{21} $\chi_{c1}\gamma$	< 8 × 10 ⁻⁴	90%	DESIG=22
Γ_{22} $\chi_{c2}\gamma$	< 4 × 10 ⁻³	90%	DESIG=23
Γ_{23} $\Lambda\bar{\Lambda}$	< 3.1 × 10 ⁻⁶	90%	DESIG=27
Γ_{24} $\Xi^-\bar{\Xi}^+$	< 4 × 10 ⁻⁵	90%	DESIG=30
Γ_{25} $pK^-\bar{\Lambda} + \text{c.c.}$	< 6 × 10 ⁻⁶	90%	DESIG=31
Γ_{26} $\omega\pi^0$	not seen		DESIG=28
Γ_{27} $\omega\eta$	not seen		DESIG=29
Γ_{28} e^+e^-	(9.4 ± 3.2) × 10 ⁻⁶		DESIG=1
Γ_{29} $\mu^+\mu^-$	(2.0 ± 1.0) × 10 ⁻⁵		DESIG=26

$\psi(4415)$ PARTIAL WIDTHS

NODE=M073220

$\Gamma(e^+e^-)$

Γ_{28}

VALUE (keV) DOCUMENT ID TECN COMMENT

NODE=M073W1

NODE=M073W1

0.58 ± 0.07 OUR ESTIMATE

→ UNCHECKED ←

0.35 ± 0.12

¹¹ ABLIKIM 08D BES2 $e^+e^- \rightarrow$ hadrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.4 to 0.8 ¹² MO 10 RVUE $e^+e^- \rightarrow$ hadrons

0.72 ± 0.11 ¹³ SETH 05A RVUE $e^+e^- \rightarrow$ hadrons

0.64 ± 0.23 ¹⁴ SETH 05A RVUE $e^+e^- \rightarrow$ hadrons

OCCUR=2

0.49 ± 0.13 BRANDELIK 78C DASP e^+e^-

0.44 ± 0.14 SIEGRIST 76 MRK1 e^+e^-

¹¹ Reanalysis of data presented in BAI 02C. From a global fit over the center-of-mass energy region 3.7–5.0 GeV covering the $\psi(3770)$, $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ resonances. Phase angle fixed in the fit to $\delta = (234 \pm 88)^\circ$.

NODE=M073W1;LINKAGE=AB

¹² Reanalysis of data presented in BAI 00 and BAI 02C. From a global fit over the center-of-mass energy 3.8–4.8 GeV covering the $\psi(4040)$, $\psi(4160)$ and $\psi(4415)$ resonances and including interference effects. Four sets of solutions are obtained with the same fit quality, mass and total width, but with different e^+e^- partial widths. We quote only the range of values.

NODE=M073W1;LINKAGE=MO

¹³ From a fit to Crystal Ball (OSTERHELD 86) data.

NODE=M073W1;LINKAGE=ST

¹⁴ From a fit to BES (BAI 02C) data.

NODE=M073W1;LINKAGE=SE

$\Gamma(\mu^+\mu^-)$ Γ_{29}

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
$1.25 \pm 0.28 \pm 0.35$	15,16 ABLIKIM	20AG BES3	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M073W2
NODE=M073W2

¹⁵ From a fit to the $e^+e^- \rightarrow \mu^+\mu^-$ cross section between 3.8 and 4.6 GeV to the coherent sum of four resonant amplitudes assuming $\Gamma(\mu^+\mu^-) = \Gamma(e^+e^-)$.

NODE=M073W2;LINKAGE=A

¹⁶ From solution 1 of 8 with equal fit quality. Other solutions range from $1.24 \pm 0.28 \pm 0.35$ to $1.27 \pm 0.41 \pm 0.36$ keV.

NODE=M073W2;LINKAGE=B

 $\psi(4415) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M073230

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<3.6	90	WANG	13B BELL	$e^+e^- \rightarrow J/\psi\eta\gamma$

 $\Gamma_{20}\Gamma_{28}/\Gamma$ NODE=M073G01
NODE=M073G01

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.47	90	17 HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c1}\gamma$

 $\Gamma_{21}\Gamma_{28}/\Gamma$ NODE=M073G02
NODE=M073G02

¹⁷ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M073G02;LINKAGE=A

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	18 HAN	15 BELL	$10.58 e^+e^- \rightarrow \chi_{c2}\gamma$

 $\Gamma_{22}\Gamma_{28}/\Gamma$ NODE=M073G03
NODE=M073G03

¹⁸ Using $B(\eta \rightarrow \gamma\gamma) = (39.41 \pm 0.21)\%$.

NODE=M073G03;LINKAGE=A

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-3}$	90	19 ABLIKIM	21AS BES3	$e^+e^- \rightarrow \psi(4415)$

 $\Gamma_{23}\Gamma_{28}/\Gamma$ NODE=M073R16
NODE=M073R16

¹⁹ From a measurement of the $e^+e^- \rightarrow \Lambda\bar{\Lambda}$ cross section between 3.5 and 4.6 GeV.

NODE=M073R16;LINKAGE=A

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.0217	90	20 ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4415)$

 $\Gamma_{24}\Gamma_{28}/\Gamma$ NODE=M073R19
NODE=M073R19

²⁰ From a fit to $e^+e^- \rightarrow \Xi^-\Xi^+$ cross sections.

NODE=M073R19;LINKAGE=A

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-3}$	90	21 ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4415)$

 $\Gamma_{25}\Gamma_{28}/\Gamma$ NODE=M073R21
NODE=M073R21

²¹ From a fit to $e^+e^- \rightarrow pK^-\bar{\Lambda} + \text{c.c.}$ cross sections.

NODE=M073R21;LINKAGE=A

 $\psi(4415) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$

NODE=M073235

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.99 \times 10^{-6}$	90	22 PAKHLOVA	09 BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$

 $\Gamma_{12}/\Gamma \times \Gamma_{28}/\Gamma$ NODE=M073R01
NODE=M073R01

²² Using 4421 ± 4 MeV for the mass of $\psi(4415)$.

NODE=M073R01;LINKAGE=PA

 $\psi(4415) \text{ BRANCHING RATIOS}$

NODE=M073225

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^0 \bar{D}^0 \gamma$

 Γ_2/Γ NODE=M073R04
NODE=M073R04

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^0 \bar{D}^0 \gamma$
----------	--------	----------	---

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA	08 BELL	$e^+e^- \rightarrow D^+ D^- \gamma$

 Γ_3/Γ NODE=M073R05
NODE=M073R05

••• We do not use the following data for averages, fits, limits, etc. •••

not seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^+ D^- \gamma$
----------	--------	----------	-------------------------------------

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT	09M BABR	$e^+e^- \rightarrow D^{*0} \bar{D}^0 \gamma$

 Γ_5/Γ NODE=M073R06
NODE=M073R06

$\Gamma(D^*(2010)^+ D^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	²³ ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^- \gamma$
²³ Supersedes PAKHLOVA 07.			

NODE=M073R07
NODE=M073R07

NODE=M073R07;LINKAGE=A

 $\Gamma(D\bar{D})/\Gamma(D^*\bar{D}^*)$ Γ_1/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
0.14±0.12±0.03	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

NODE=M073R02
NODE=M073R02 $\Gamma(D^*\bar{D} + \text{c.c.})/\Gamma(D^*\bar{D}^*)$ Γ_4/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.25±0.03	AUBERT 09M	BABR	$e^+ e^- \rightarrow \gamma D^{(*)} \bar{D}^{(*)}$

NODE=M073R03
NODE=M073R03 $\Gamma(D^*(2007)^0 \bar{D}^*(2007)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*0} \bar{D}^{*0} \gamma$

NODE=M073R08
NODE=M073R08 $\Gamma(D^*(2010)^+ D^*(2010)^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	²⁴ ZHUKOVA 18	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
seen	AUBERT 09M	BABR	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	PAKHLOVA 07	BELL	$e^+ e^- \rightarrow D^{*+} D^{*-} \gamma$
²⁴ Supersedes PAKHLOVA 07.			

NODE=M073R09
NODE=M073R09

NODE=M073R09;LINKAGE=A

 $\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
10.5±2.4±3.8	²⁵ PAKHLOVA 08A	BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
²⁵ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.			

NODE=M073R3
NODE=M073R3

NODE=M073R3;LINKAGE=PA

 $\Gamma(D^0 D^- \pi^+ (\text{excl. } D^*(2007)^0 \bar{D}^0 + \text{c.c.}, D^*(2010)^+ D^- + \text{c.c.})/$ $\Gamma(D\bar{D}_2^*(2460) \rightarrow D^0 D^- \pi^+ + \text{c.c.})$ Γ_{10}/Γ_{11}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.22	90	²⁶ PAKHLOVA 08A	BELL	$10.6 e^+ e^- \rightarrow D^0 D^- \pi^+ \gamma$
²⁶ Using 4421 ± 4 MeV for the mass and 62 ± 20 MeV for the width of $\psi(4415)$.				

NODE=M073R4
NODE=M073R4

NODE=M073R4;LINKAGE=PA

 $\Gamma(D_1(2420)\bar{D} + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	²⁷ ABLIKIM 19AR	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$
²⁷ Evidence for $e^+ e^- \rightarrow D_1(2420)\bar{D} + \text{c.c.}$ between $\sqrt{s} = 4.3$ and 4.6 GeV, not necessarily resonant.			

NODE=M073R15
NODE=M073R15

NODE=M073R15;LINKAGE=A

 $\Gamma(D_s^+ D_s^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$
not seen	DEL-AMO-SA...10N	BABR	$e^+ e^- \rightarrow D_s^+ D_s^- \gamma$

NODE=M073R10
NODE=M073R10 $\Gamma(\omega \chi_{c2})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	ABLIKIM 16A	BES3	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- \pi^0 \ell^+ \ell^-$

NODE=M073R00
NODE=M073R00 $\Gamma(D_s^{*+} D_s^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	PAKHLOVA 11	BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$
seen	DEL-AMO-SA...10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^- \gamma$

NODE=M073R11
NODE=M073R11

$\Gamma(D_s^{*+} D_s^{*-})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen [not seen OUR 2023 BEST LIMIT]

seen	ABLIKIM	23BH BES3	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-}$
------	---------	-----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	PAKHLOVA	11 BELL	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
----------	----------	---------	--

not seen	DEL-AMO-SA...10N	BABR	$e^+ e^- \rightarrow D_s^{*+} D_s^{*-} \gamma$
----------	------------------	------	--

NODE=M073R12
NODE=M073R12 $\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

possibly seen	19	28 ABLIKIM	15S BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$
---------------	----	------------	----------	--

²⁸ From a fit of $e^+ e^- \rightarrow \pi^+ \pi^- \psi_2(3823)$, $\psi_2(3823) \rightarrow \chi_{c1} \gamma$ cross sections taken at \sqrt{s} values of 4.23, 4.26, 4.36, 4.42, and 4.60 GeV to the $\psi(4415)$ line shape.

NODE=M073R13
NODE=M073R13

NODE=M073R13;LINKAGE=A

 $\Gamma(\psi(3770)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

possibly seen	29	ABLIKIM	19AR BES3	$e^+ e^- \rightarrow \pi^+ \pi^- D \bar{D}$
---------------	----	---------	-----------	---

²⁹ Observe $e^+ e^- \rightarrow \pi^+ \pi^- \psi(3770)$ at $\sqrt{s} = 4.26, 4.36, \text{ and } 4.42$ GeV but cannot establish if continuum or resonant.

NODE=M073R14
NODE=M073R14

NODE=M073R14;LINKAGE=A

 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	ABLIKIM	22K BES3	$e^+ e^- \rightarrow \omega \pi^0$
----------	---------	----------	------------------------------------

NODE=M073R17
NODE=M073R17 $\Gamma(\omega\eta)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

not seen	ABLIKIM	22K BES3	$e^+ e^- \rightarrow \omega \eta$
----------	---------	----------	-----------------------------------

NODE=M073R18
NODE=M073R18 $\psi(4415)$ REFERENCES

NODE=M073

ABLIKIM	23BH	PRL 131 151903	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62433
ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62437
ABLIKIM	23BL	JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62438
ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	21AS	PR D104 L091104	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61454
ABLIKIM	20AG	PR D102 112009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60744
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ZHUKOVA	18	PR D97 012002	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=58710
ABLIKIM	16A	PR D93 011102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57122
ABLIKIM	15S	PRL 115 011803	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56784
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	11	PR D83 011101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53638
DEL-AMO-SA...10N	10N	PR D82 052004	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53532
MO	10	PR D82 077501	X.H. Mo, C.Z. Yuan, P. Wang	(BHEP)	REFID=53540
AUBERT	09M	PR D79 092001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52724
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
ABLIKIM	08D	PL B660 315	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=52142
PAKHLOVA	08	PR D77 011103	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52132
PAKHLOVA	08A	PRL 100 062001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52134
PAKHLOVA	07	PRL 98 092001	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=51628
SETH	05A	PR D72 017501	K.K. Seth		REFID=50813
BAI	02C	PRL 88 101802	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50506
BAI	00	PRL 84 594	J.Z. Bai <i>et al.</i>	(BES Collab.)	REFID=50503
OSTERHELD	86	SLAC-PUB-4160	A. Osterheld <i>et al.</i>	(SLAC Crystal Ball Collab.)	REFID=51064
BRANDELIK	78C	PL 76B 361	R. Brandelik <i>et al.</i>	(DASP Collab.)	REFID=22232
SIEGRIST	76	PRL 36 700	J.L. Siegrist <i>et al.</i>	(LBL, SLAC)	REFID=22243

NODE=M234

 $\chi_{c0}(4500)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE
was $X(4500)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

NODE=M234

Seen by AAIJ 17C in $B^+ \rightarrow \chi_{c0} K^+$, $\chi_{c0} \rightarrow J/\psi \phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ with a significance (accounting for systematic uncertainties) of 6.1σ .

 $\chi_{c0}(4500)$ MASS

NODE=M234M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4474 \pm 3 \pm 3$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$4506 \pm 11^{+12}_{-15}$	4289	^{2,3} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M234M

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 20σ .

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.1σ .

³ Superseded by AAIJ 21E.

NODE=M234M;LINKAGE=B

NODE=M234M;LINKAGE=A

NODE=M234M;LINKAGE=C

 $\chi_{c0}(4500)$ WIDTH

NODE=M234W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$77 \pm 6^{+10}_{-8}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$92 \pm 21^{+21}_{-20}$	4289	^{2,3} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M234W

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 20σ .

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.1σ .

³ Superseded by AAIJ 21E.

NODE=M234W;LINKAGE=B

NODE=M234W;LINKAGE=A

NODE=M234W;LINKAGE=C

 $\chi_{c0}(4500)$ DECAY MODES

NODE=M234215;NODE=M234

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi \phi$	seen

DESIG=1

 $\chi_{c0}(4500)$ BRANCHING RATIOS

NODE=M234220

$\Gamma(J/\psi \phi)/\Gamma_{\text{total}}$	Γ_1/Γ
seen	seen
• • • We do not use the following data for averages, fits, limits, etc. • • •	
seen	seen

NODE=M234R01

NODE=M234R01

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 20σ .

² From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 6.1σ .

³ Superseded by AAIJ 21E.

NODE=M234R01;LINKAGE=B

NODE=M234R01;LINKAGE=A

NODE=M234R01;LINKAGE=C

 $\chi_{c0}(4500)$ REFERENCES

NODE=M234

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=61150
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=57657
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57636

X(4630)

$$I^G(J^{PC}) = 0^+(?^{?+})$$

NODE=M262

OMITTED FROM SUMMARY TABLE

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M262

Seen by AAIJ 21E in $B^+ \rightarrow X(4630)K^+$ with $X(4630) \rightarrow J/\psi\phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ with a significance (accounting for systematic uncertainties) of 5.5σ . The $J^P = 1^-$ assignment is favored over 2^- with a significance of 3σ and other assignments are disfavored by more than 5σ .

X(4630) MASS

NODE=M262M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4626 \pm 16^{+18}_{-110}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M262M

¹From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.5σ .

NODE=M262M;LINKAGE=A

X(4630) WIDTH

NODE=M262W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
170^{+140}_{-80} OUR AVERAGE		[174 ⁺¹⁴⁰ ₋₈₀ MeV OUR 2023 AVERAGE]		
$174 \pm 27^{+134}_{-73}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M262W

NEW

¹From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.5σ .

NODE=M262W;LINKAGE=A

X(4630) DECAY MODES

NODE=M262215;NODE=M262

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen

DESIG=1

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$	

NODE=M262R01
NODE=M262R01

¹From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.5σ .

NODE=M262R01;LINKAGE=A

X(4630) REFERENCES

NODE=M262

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	-----	----------------	-----------------------	----------------

REFID=61150

$\psi(4660)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $Y(4660)$; was $X(4660)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen in radiative return from e^+e^- collisions at $\sqrt{s} = 9.54\text{--}10.58$ GeV by WANG 07D. Also obtained in a combined fit of WANG 07D, AUBERT 07S, and LEES 14F. See also the review on "Spectroscopy of mesons containing two heavy quarks."

 $\psi(4660)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4641 ± 10	OUR AVERAGE	Error includes scale factor of 2.7. See the ideogram below. [4630 ± 6 MeV OUR 2023 AVERAGE Scale factor = 1.4]		
4708 $^{+17}_{-15}$ ± 21		1 ABLIKIM	23BI BES3	$e^+e^- \rightarrow K^+K^-J/\psi$
4701.8 ± 10.9 ± 2.7		2 ABLIKIM	23H BES3	$e^+e^- \rightarrow \phi\chi_{c2}$
4675.3 ± 29.5 ± 3.5		3 ABLIKIM	23X BES3	$e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$
4651.0 ± 37.8 ± 2.1		4 ABLIKIM	21AJ BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4619.8 $^{+8.9}_{-8.0}$ ± 2.3	66	5 JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^{*-}(2573)^-$
4625.9 $^{+6.2}_{-6.0}$ ± 0.4	89	6 JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
4652 ± 10 ± 11	279	7 WANG	15A BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4669 ± 21 ± 3	37	8 LEES	14F BABR	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4634 $^{+8}_{-7}$ $^{+5}_{-8}$ ± 6	142	9 PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4647.9 ± 8.6 ± 0.8		10 ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$
4652.5 ± 3.4 ± 1.1		11 DAI	17 RVUE	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
4645.2 ± 9.5 ± 6.0		12 ZHANG	17B RVUE	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$
4646.4 ± 9.7 ± 4.8		13 ZHANG	17C RVUE	$e^+e^- \rightarrow \pi^+\pi^-J/\psi$ or $\psi(2S)$
4661 $^{+9}_{-8}$ ± 6	44	14 LIU	08H RVUE	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4664 ± 11 ± 5	44	WANG	07D BELL	10.58 $e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$

¹ Seen as a peak in the c.m. energy dependence of the $e^+e^- \rightarrow K^+K^-J/\psi$ cross section using 5.85 fb⁻¹ of data at c.m. energies 4.61–4.95 GeV. Statistical significance is over 5 σ .

² Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term.

³ From a cross-section measurement of $e^+e^- \rightarrow D^{*0}D^{*-}\pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. The two other resonances have masses (widths) 4209.6 ± 7.5 (81.6 ± 19.9) MeV and 4469.1 ± 26.4 (246.3 ± 37.9) MeV.

⁴ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008\text{--}4.698$ GeV.

⁵ Using $D_{s2}^{*-}(2573)^- \rightarrow \bar{D}^0 K^-$ decays.

⁶ From a fit of a Breit-Wigner convolved with a Gaussian.

⁷ From a two-resonance fit. Supersedes WANG 07D.

⁸ From a two-resonance fit.

⁹ The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

¹⁰ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass 4417.5 ± 26.2 ± 3.5 MeV and width 245 ± 48 ± 13 MeV.

¹¹ The pole parameters are extracted from the speed plot.

¹² From a three-resonance fit.

¹³ From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.

¹⁴ From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189

NODE=M189

NODE=M189M

NODE=M189M

NEW

NODE=M189M;LINKAGE=M

NODE=M189M;LINKAGE=J

NODE=M189M;LINKAGE=L

NODE=M189M;LINKAGE=G

NODE=M189M;LINKAGE=F

NODE=M189M;LINKAGE=E

NODE=M189M;LINKAGE=A

NODE=M189M;LINKAGE=LE

NODE=M189M;LINKAGE=PA


NODE=M189M;LINKAGE=I

NODE=M189M;LINKAGE=C

NODE=M189M;LINKAGE=D

NODE=M189M;LINKAGE=B

NODE=M189M;LINKAGE=LI



psi(4660) WIDTH

NODE=M189W

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT

NODE=M189W

73 $^{+13}_{-11}$ **OUR AVERAGE** Error includes scale factor of 1.7. See the ideogram below.

NEW

$[72^{+14}_{-12}$ MeV OUR 2023 AVERAGE Scale factor = 1.7]

126	$^{+27}_{-23} \pm 30$	1	ABLIKIM	23BI	BES3	$e^+ e^- \rightarrow K^+ K^- J/\psi$	
	$30.5 \pm 22.3 \pm 14.6$	2	ABLIKIM	23H	BES3	$e^+ e^- \rightarrow \phi \chi_{c2}$	
	$218.3 \pm 72.9 \pm 9.3$	3	ABLIKIM	23X	BES3	$e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$	
	$155.4 \pm 24.8 \pm 0.8$	4	ABLIKIM	21AJ	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
	$47.0^{+31.3}_{-14.8} \pm 4.6$	66	5	JIA	20	BELL	$e^+ e^- \rightarrow \gamma D_s^+ D_{s2}^{*-}(2573)^-$
	$49.8^{+13.9}_{-11.5} \pm 4.0$	89	6	JIA	19A	BELL	$e^+ e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$
	$68 \pm 11 \pm 5$	279	7	WANG	15A	BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
	$104 \pm 48 \pm 10$	37	8	LEES	14F	BABR	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
	$92^{+40}_{-24} \pm 10$	142	9	PAKHLOVA	08B	BELL	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●							
	$33.1 \pm 18.6 \pm 4.1$	10	ABLIKIM	22R	BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \chi_{c1} \gamma$	
	$62.6 \pm 5.6 \pm 4.3$	11	DAI	17	RVUE	$e^+ e^- \rightarrow \Lambda_c^+ \Lambda_c^-$	
	$113.8 \pm 18.1 \pm 3.4$	12	ZHANG	17B	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$	
	$103.5 \pm 15.6 \pm 4.0$	13	ZHANG	17C	RVUE	$e^+ e^- \rightarrow \pi^+ \pi^- J/\psi$ or $\psi(2S)$	
	$42^{+17}_{-12} \pm 6$	44	14	LIU	08H	RVUE	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
	$48 \pm 15 \pm 3$	44	WANG	07D	BELL	$10.58 e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$	

NODE=M189W;LINKAGE=L

¹ Seen as a peak in the c.m. energy dependence of the $e^+ e^- \rightarrow K^+ K^- J/\psi$ cross section using 5.85 fb^{-1} of data at c.m. energies 4.61–4.95 GeV. Statistical significance is over 5σ .

NODE=M189W;LINKAGE=J

² Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term.

NODE=M189W;LINKAGE=K

³ From a cross-section measurement of $e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. The two other resonances have masses (widths) 4209.6 ± 7.5 (81.6 ± 19.9) MeV and 4469.1 ± 26.4 (246.3 ± 37.9) MeV.

NODE=M189W;LINKAGE=H

⁴ From a three-resonance fit to the Born cross section in the range $\sqrt{s} = 4.008$ – 4.698 GeV.

⁵ Using $D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-$ decays.

NODE=M189W;LINKAGE=G

⁶ From a fit of a Breit-Wigner convolved with a Gaussian.

NODE=M189W;LINKAGE=F

⁷ From a two-resonance fit. Supersedes WANG 07D.


NODE=M189W;LINKAGE=A

- 8 From a two-resonance fit.
- 9 The $\pi^+\pi^-\psi(2S)$ and $\Lambda_c^+\Lambda_c^-$ states are not necessarily the same.
- 10 From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances. The data is also consistent with a single peak with mass $4417.5 \pm 26.2 \pm 3.5$ MeV and width $245 \pm 48 \pm 13$ MeV.
- 11 The pole parameters are extracted from the speed plot.
- 12 From a three-resonance fit.
- 13 From a combined fit of BELLE, BABAR and BES3 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ and $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ data.
- 14 From a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189W;LINKAGE=LE
 NODE=M189W;LINKAGE=B
 NODE=M189W;LINKAGE=I

 NODE=M189W;LINKAGE=D
 NODE=M189W;LINKAGE=E
 NODE=M189W;LINKAGE=C

 NODE=M189W;LINKAGE=LI



$\psi(4660)$ DECAY MODES

NODE=M189215;NODE=M189

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	not seen
Γ_2 $\psi(2S)\pi^+\pi^-$	seen
Γ_3 $J/\psi\eta$	not seen
Γ_4 $D^0D^{*-}\pi^+$	not seen
Γ_5 $D^{*0}D^{*-}\pi^+$	seen
Γ_6 $\psi_2(3823)\pi^+\pi^-$	seen
Γ_7 $\chi_{c1}\gamma$	not seen
Γ_8 $\chi_{c1}\phi$	not seen
Γ_9 $\chi_{c2}\gamma$	not seen
Γ_{10} $\chi_{c2}\phi$	not seen
Γ_{11} $\Lambda_c^+\Lambda_c^-$	seen
Γ_{12} $D_s^+D_{s1}^-(2536)^-$	seen
Γ_{13} $D_s^+D_{s2}^*(2573)^-$	
Γ_{14} $\omega\pi^0$	not seen
Γ_{15} $\omega\eta$	not seen
Γ_{16} $\Xi^-\Xi^+$	
Γ_{17} $pK^-\bar{\Lambda} + c.c.$	

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=4;OUR EVAL;→ UNCHECKED ←
 DESIG=3;OUR EVAL;→ UNCHECKED ←
 DESIG=15;OUR EVAL;→ UNCHECKED ←
 DESIG=10
 DESIG=6;OUR EVAL;→ UNCHECKED ←
 DESIG=13;OUR EVAL;→ UNCHECKED ←
 DESIG=7;OUR EVAL;→ UNCHECKED ←
 DESIG=14;OUR EVAL;→ UNCHECKED ←
 DESIG=5;OUR EVAL;→ UNCHECKED ←
 DESIG=8;OUR EVAL;→ UNCHECKED ←
 DESIG=9
 DESIG=11
 DESIG=12
 DESIG=16
 DESIG=17

$\psi(4660) \Gamma(i) \times \Gamma(e^+ e^-) / \Gamma(\text{total})$

NODE=M189230

 $\Gamma(\psi(2S)\pi^+\pi^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_2 \Gamma_1 / \Gamma$ NODE=M189G1
NODE=M189G1

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
4.7±3.8		¹ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.2±3.2		² ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
4.7±4.2		³ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
11.3±3.3		⁴ ABLIKIM	21AJ BES3	$e^+ e^- \rightarrow \pi^+ \pi^- \psi(2S)$
2.0±0.3±0.2	279	⁵ WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
8.1±1.1±1.0	279	⁶ WANG	15A BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.7±1.3±0.5	37	⁷ LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.5±1.7±0.7	37	⁸ LEES	14F BABR	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
2.2 ^{+0.7} _{-0.6}	44	⁹ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
5.9±1.6	44	¹⁰ LIU	08H RVUE	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
3.0±0.9±0.3	44	⁷ WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$
7.6±1.8±0.8	44	⁸ WANG	07D BELL	10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=2

¹ Solution I of four equivalent solutions in a fit using three interfering resonances.² Solution II of four equivalent solutions in a fit using three interfering resonances.³ Solution III of four equivalent solutions in a fit using three interfering resonances.⁴ Solution IV of four equivalent solutions in a fit using three interfering resonances.⁵ Solution I of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.⁶ Solution II of two equivalent solutions from a fit using two interfering resonances. Supersedes WANG 07D.⁷ Solution I of two equivalent solutions in a fit using two interfering resonances.⁸ Solution II of two equivalent solutions in a fit using two interfering resonances.⁹ Solution I in a combined fit of AUBERT 07S and WANG 07D data with two resonances.¹⁰ Solution II in a combined fit of AUBERT 07S and WANG 07D data with two resonances.

NODE=M189G1;LINKAGE=C

NODE=M189G1;LINKAGE=D

NODE=M189G1;LINKAGE=E

NODE=M189G1;LINKAGE=F

NODE=M189G1;LINKAGE=A

NODE=M189G1;LINKAGE=B

NODE=M189G1;LINKAGE=WA

NODE=M189G1;LINKAGE=WN

NODE=M189G1;LINKAGE=LI

NODE=M189G1;LINKAGE=LU

 $\Gamma(J/\psi\eta) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_3 \Gamma_1 / \Gamma$ NODE=M189G01
NODE=M189G01

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.94	90	WANG	13B BELL	$e^+ e^- \rightarrow J/\psi \eta \gamma$

 $\Gamma(D^{*0} D^{*-} \pi^+) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_5 \Gamma_1 / \Gamma$ NODE=M189R10
NODE=M189R10

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
19 to 2005	¹ ABLIKIM	23X BES3	$e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$

¹ From a cross-section measurement of $e^+ e^- \rightarrow D^{*0} D^{*-} \pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. Depending on solutions I – VIII with same fit qualities.

NODE=M189R10;LINKAGE=A

 $\Gamma(\chi_{c1} \gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_7 \Gamma_1 / \Gamma$ NODE=M189G02
NODE=M189G02

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.45	90	¹ HAN	15 BELL	10.58 $e^+ e^- \rightarrow \chi_{c1} \gamma$

¹ Using $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$.

NODE=M189G02;LINKAGE=A

 $\Gamma(\chi_{c1} \phi) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_8 \Gamma_1 / \Gamma$ NODE=M189R08
NODE=M189R08

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
<0.04	90	¹ ABLIKIM	23H BES3	$e^+ e^- \rightarrow \phi \chi_{c1}$

¹ Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term.

NODE=M189R08;LINKAGE=A

 $\Gamma(\chi_{c2} \gamma) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ $\Gamma_9 \Gamma_1 / \Gamma$ NODE=M189G03
NODE=M189G03

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	¹ HAN	15 BELL	10.58 $e^+ e^- \rightarrow \chi_{c2} \gamma$

¹ Using $B(\eta \rightarrow \gamma \gamma) = (39.41 \pm 0.21)\%$.

NODE=M189G03;LINKAGE=A

$$\Gamma(\chi_{c2}\phi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{10}\Gamma_1/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.13	¹ ABLIKIM	23H BES3	$e^+e^- \rightarrow \phi\chi_{c2}$

NODE=M189R09
NODE=M189R09

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.13 ± 0.13 ¹ ABLIKIM 23H BES3 $e^+e^- \rightarrow \phi\chi_{c2}$

¹ Fit model parameterized as the coherent sum of a Breit-Wigner resonance and a continuum amplitude term. Constructive solution of the interference. Destructive solution gives 0.66 ± 0.41 eV.

NODE=M189R09;LINKAGE=A

$$\Gamma(D_s^+ D_{s1}(2536)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{12}\Gamma_1/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$14.3^{+2.8}_{-2.6} \pm 1.5$	89	¹ JIA	19A BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s1}(2536)^-$

NODE=M189R00
NODE=M189R00

¹ Assuming $B(D_{s1}(2536)^- \rightarrow \bar{D}^{*0} K^-) = 1$.

NODE=M189R00;LINKAGE=A

$$\Gamma(D_s^+ D_{s2}^*(2573)^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{13}\Gamma_1/\Gamma$$

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
$14.7^{+5.9}_{-4.5} \pm 3.6$	66	¹ JIA	20 BELL	$e^+e^- \rightarrow \gamma D_s^+ D_{s2}^*(2573)^-$

NODE=M189R04
NODE=M189R04

¹ Assuming $B(D_{s2}^*(2573)^- \rightarrow \bar{D}^0 K^-) = 1$.

NODE=M189R04;LINKAGE=A

$$\Gamma(\Xi^- \Xi^+) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{16}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0199	90	¹ ABLIKIM	23BK BES3	$e^+e^- \rightarrow \psi(4660)$

NODE=M189R11
NODE=M189R11

¹ From a fit to $e^+e^- \rightarrow \Xi^- \Xi^+$ cross sections.

NODE=M189R11;LINKAGE=A

$$\Gamma(pK^- \bar{\Lambda} + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{17}\Gamma_1/\Gamma$$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-3}$	90	¹ ABLIKIM	23BL BES3	$e^+e^- \rightarrow \psi(4660)$

NODE=M189R13
NODE=M189R13

¹ From a fit to $e^+e^- \rightarrow pK^- \bar{\Lambda} + \text{c.c.}$ cross sections.

NODE=M189R13;LINKAGE=A

$$\psi(4660) \Gamma(i) \times \Gamma(e^+e^-)/\Gamma^2(\text{total})$$

NODE=M189235

$$\Gamma(D^0 D^{*-} \pi^+)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma \times \Gamma_1/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.37 \times 10^{-6}$	90	¹ PAKHLOVA	09 BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M189R02
NODE=M189R02

¹ Using $4664 \pm 11 \pm 5$ MeV for the mass of $\psi(4660)$.

NODE=M189R02;LINKAGE=PA

$$\Gamma(\Lambda_c^+ \Lambda_c^-)/\Gamma_{\text{total}} \times \Gamma(e^+e^-)/\Gamma_{\text{total}} \quad \Gamma_{11}/\Gamma \times \Gamma_1/\Gamma$$

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.68^{+0.16+0.29}_{-0.15-0.30}$	142	¹ PAKHLOVA	08B BELL	$e^+e^- \rightarrow \Lambda_c^+ \Lambda_c^-$

NODE=M189R03
NODE=M189R03

¹ The $\pi^+ \pi^- \psi(2S)$ and $\Lambda_c^+ \Lambda_c^-$ states are not necessarily the same.

NODE=M189R03;LINKAGE=A

$\psi(4660)$ BRANCHING RATIOS

NODE=M189225

$$\Gamma(D^0 D^{*-} \pi^+)/\Gamma(\psi(2S)\pi^+\pi^-) \quad \Gamma_4/\Gamma_2$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 10	90	PAKHLOVA	09 BELL	$e^+e^- \rightarrow D^0 D^{*-} \pi^+$

NODE=M189R01
NODE=M189R01

$$\Gamma(\psi_2(3823)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ ABLIKIM	22R BES3	$e^+e^- \rightarrow \pi^+\pi^-\chi_{c1}\gamma$

NODE=M189R05
NODE=M189R05

¹ From a fit to the $e^+e^- \rightarrow \pi^+\pi^-\psi(3823)$ cross section between 4.23 and 4.70 GeV with two coherent Breit-Wigner resonances.

NODE=M189R05;LINKAGE=A

$$\Gamma(\omega\pi^0)/\Gamma_{\text{total}} \quad \Gamma_{14}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	22K BES3	$e^+e^- \rightarrow \omega\pi^0$

NODE=M189R06
NODE=M189R06

$$\Gamma(\omega\eta)/\Gamma_{\text{total}} \quad \Gamma_{15}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	ABLIKIM	22K BES3	$e^+e^- \rightarrow \omega\eta$

NODE=M189R07
NODE=M189R07

$\psi(4660)$ REFERENCES

ABLIKIM	23BI	PRL 131 211902	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62434
ABLIKIM	23BK	JHEP 2311 228	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62437
ABLIKIM	23BL	JHEP 2312 027	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62438
ABLIKIM	23H	JHEP 2301 132	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62050
ABLIKIM	23X	PRL 130 121901	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62072
ABLIKIM	22K	JHEP 2207 064	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61648
ABLIKIM	22R	PRL 129 102003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61664
ABLIKIM	21AJ	PR D104 052012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61441
JIA	20	PR D101 091101	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=60301
JIA	19A	PR D100 111103	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=60037
DAI	17	PR D96 116001	L.-Y. Dai, J. Haidenbauer, U.-G. Meissner	(JULI+)	REFID=58704
ZHANG	17B	PR D96 054008	J. Zhang, J. Zhang		REFID=58219
ZHANG	17C	EPJ C77 727	J. Zhang, L. Yuan		REFID=58463
HAN	15	PR D92 012011	Y.L. Han <i>et al.</i>	(BELLE Collab.)	REFID=56816
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=56839
LEES	14F	PR D89 111103	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55938
WANG	13B	PR D87 051101	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=55377
PAKHLOVA	09	PR D80 091101	G. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=53143
LIU	08H	PR D78 014032	Z.Q. Liu, X.S. Qin, C.Z. Yuan		REFID=52296
PAKHLOVA	08B	PRL 101 172001	C. Pakhlova <i>et al.</i>	(BELLE Collab.)	REFID=52596
AUBERT	07S	PRL 98 212001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51724
WANG	07D	PRL 99 142002	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=51959

NODE=M189

REFID=62434
REFID=62437
REFID=62438
REFID=62050
REFID=62072
REFID=61648
REFID=61664
REFID=61441
REFID=60301
REFID=60037
REFID=58704
REFID=58219
REFID=58463
REFID=56816
REFID=56839
REFID=55938
REFID=55377
REFID=53143
REFID=52296
REFID=52596
REFID=51724
REFID=51959 $\chi_{c1}(4685)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

OMITTED FROM SUMMARY TABLE

This state shows properties different from a conventional $q\bar{q}$ state. A candidate for an exotic structure. See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M261

NODE=M261

Seen by AAIJ 21E in $B^+ \rightarrow \chi_{c1}(4685)K^+$ with $\chi_{c1}(4685) \rightarrow J/\psi\phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ with a significance (accounting for systematic uncertainties) of 15σ . The $J^P = 1^+$ assignment is favored with high significance.

 $\chi_{c1}(4685)$ MASS

NODE=M261M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4684 \pm 7^{+13}_{-16}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M261M

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 15σ .

NODE=M261M;LINKAGE=A

 $\chi_{c1}(4685)$ WIDTH

NODE=M261W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
130 ± 40 OUR AVERAGE	[126 \pm 40 MeV OUR 2023 AVERAGE]			
$126 \pm 15^{+37}_{-41}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M261W

NEW

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 15σ .

NODE=M261W;LINKAGE=A

 $\chi_{c1}(4685)$ DECAY MODES

NODE=M261215;NODE=M261

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen

DESIG=1

 $\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$

NODE=M261R01
NODE=M261R01

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 15σ .

NODE=M261R01;LINKAGE=A

 $\chi_{c1}(4685)$ REFERENCES

NODE=M261

AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=61150
------	-----	----------------	-----------------------	-------------------	-------------

REFID=61150

$\chi_{c0}(4700)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE
was $X(4700)$

This state shows properties different from a conventional $q\bar{q}$ state.
A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

Seen by AAIJ 17C in $B^+ \rightarrow \chi_{c0} K^+$, $\chi_{c0} \rightarrow J/\psi\phi$ using an amplitude analysis of $B^+ \rightarrow J/\psi\phi K^+$ with a significance (accounting for systematic uncertainties) of 5.6σ .

NODE=M235

NODE=M235

 $\chi_{c0}(4700)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$4694 \pm 4^{+16}_{-3}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
$4741 \pm 6 \pm 6$	175	² AAIJ	21C LHCB	$B_s^0 \rightarrow J/\psi\phi\pi^+\pi^-$
$4704 \pm 10^{+14}_{-24}$	4289	^{3,4} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi\phi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 17σ .

² From a 1D fit to the $J/\psi\phi$ mass distribution with a significance of 5.3σ . The identification of this structure as the $\chi_{c0}(4700)$ needs confirmation.

³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.6σ .

⁴ Superseded by AAIJ 21E.

NODE=M235M

NODE=M235M

NODE=M235M;LINKAGE=C
NODE=M235M;LINKAGE=BNODE=M235M;LINKAGE=A
NODE=M235M;LINKAGE=D **$\chi_{c0}(4700)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$87 \pm 8^{+16}_{-6}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
$53 \pm 15 \pm 11$	175	² AAIJ	21C LHCB	$B_s^0 \rightarrow J/\psi\phi\pi^+\pi^-$
$120 \pm 31^{+42}_{-33}$	4289	^{3,4} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi\phi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 17σ .

² From a 1D fit to the $J/\psi\phi$ mass distribution with a significance of 5.3σ . The identification of this structure as the $\chi_{c0}(4700)$ needs confirmation.

³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.6σ .

⁴ Superseded by AAIJ 21E.

NODE=M235W

NODE=M235W

NODE=M235W;LINKAGE=C
NODE=M235W;LINKAGE=BNODE=M235W;LINKAGE=A
NODE=M235W;LINKAGE=D **$\chi_{c0}(4700)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\phi$	seen

NODE=M235215;NODE=M235

DESIG=1

 $\chi_{c0}(4700)$ BRANCHING RATIOS

$\Gamma(J/\psi\phi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi\phi K^+$
seen	175	² AAIJ	21C LHCB	$B_s^0 \rightarrow J/\psi\phi\pi^+\pi^-$
seen	4289	^{3,4} AAIJ	17C LHCB	$B^+ \rightarrow J/\psi\phi K^+$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 17σ .

² From a 1D fit to the $J/\psi\phi$ mass distribution with a significance of 5.3σ . The identification of this structure as the $\chi_{c0}(4700)$ needs confirmation.

³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi\phi K^+$ with a significance of 5.6σ .

⁴ Superseded by AAIJ 21E.

NODE=M235220

NODE=M235R01
NODE=M235R01NODE=M235R01;LINKAGE=C
NODE=M235R01;LINKAGE=BNODE=M235R01;LINKAGE=A
NODE=M235R01;LINKAGE=D **$\chi_{c0}(4700)$ REFERENCES**

AAIJ	21C	JHEP 2102 024	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17C	PRL 118 022003	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
Also		PR D95 012002	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M235


REFID=61124
REFID=61150
REFID=57657
REFID=57636

$b\bar{b}$ MESONS (including possibly non- $q\bar{q}$ states)

NODE=MXXX030

NODE=M849

Updated March 2024.



The level scheme of meson states containing a minimal quark content of $b\bar{b}$ and having $S = C = B = 0$. The name of a state is determined by its quantum numbers $I^G J^{PC}$ (see the review “Naming Scheme for Hadrons”). States included in the Summary Tables are shown with solid lines; those requiring confirmation are shown with dotted lines. The arrows indicate the most dominant hadronic transitions. Single photon transitions, including $\Upsilon(nS) \rightarrow \gamma\eta_b(mS)$, $\Upsilon(nS) \rightarrow \gamma\chi_{bJ}(mP)$, and $\chi_{bJ}(nP) \rightarrow \gamma\Upsilon(mS)$, are omitted for clarity. For orientation, the location of the thresholds related to a pair of ground state open bottom mesons is indicated in the figure. Decays to open flavour final states are not shown in the figure. $\Upsilon(10753)$ ω -transitions to $\chi_{b1}(1P)$ and $\chi_{b2}(1P)$ have also been reported.

NODE=M849

WIDTH DETERMINATIONS OF THE Υ STATES

As is the case for the $J/\psi(1S)$ and $\psi(2S)$, the full widths of the $b\bar{b}$ states $\Upsilon(1S)$, $\Upsilon(2S)$, and $\Upsilon(3S)$ are not directly measurable, since they are much narrower than the energy resolution of the e^+e^- storage rings where these states are

produced. The common indirect method to determine Γ starts from

$$\Gamma = \Gamma_{\ell\ell}/B_{\ell\ell} , \quad (2.1)$$

where $\Gamma_{\ell\ell}$ is one leptonic partial width and $B_{\ell\ell}$ is the corresponding branching fraction ($\ell = e, \mu, \text{ or } \tau$). One then assumes e - μ - τ universality and uses

$$\begin{aligned} \Gamma_{\ell\ell} &= \Gamma_{ee} \\ B_{\ell\ell} &= \text{average of } B_{ee}, B_{\mu\mu}, \text{ and } B_{\tau\tau} . \end{aligned} \quad (2.2)$$

The electronic partial width Γ_{ee} is also not directly measurable at e^+e^- storage rings, only in the combination $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$, where Γ_{had} is the hadronic partial width and

$$\Gamma_{\text{had}} + 3\Gamma_{ee} = \Gamma . \quad (2.3)$$

This combination is obtained experimentally from the energy-integrated hadronic cross section

$$\begin{aligned} &\int_{\text{resonance}} \sigma(e^+e^- \rightarrow \Upsilon \rightarrow \text{hadrons})dE \\ &= \frac{6\pi^2}{M^2} \frac{\Gamma_{ee}\Gamma_{\text{had}}}{\Gamma} C_r = \frac{6\pi^2}{M^2} \frac{\Gamma_{ee}^{(0)}\Gamma_{\text{had}}}{\Gamma} C_r^{(0)} , \end{aligned} \quad (2.4)$$

where M is the Υ mass, and C_r and $C_r^{(0)}$ are radiative correction factors. C_r is used for obtaining Γ_{ee} as defined in Eq. (2.1), and contains corrections from all orders of QED for describing $(b\bar{b}) \rightarrow e^+e^-$. The lowest order QED value $\Gamma_{ee}^{(0)}$, relevant for comparison with potential-model calculations, is defined by the lowest order QED graph (Born term) alone, and is about 7% lower than Γ_{ee} .

The Listings give experimental results on B_{ee} , $B_{\mu\mu}$, $B_{\tau\tau}$, and $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$. The entries of the last quantity have been re-evaluated consistently using the correction procedure of KURA EV 85 [1]. The partial width Γ_{ee} is obtained from the average values for $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$ and $B_{\ell\ell}$ using

$$\Gamma_{ee} = \frac{\Gamma_{ee}\Gamma_{\text{had}}}{\Gamma(1 - 3B_{\ell\ell})} . \quad (2.5)$$

The total width Γ is then obtained from Eq. (2.1). We do not list Γ_{ee} and Γ values of individual experiments. The Γ_{ee} values in the Meson Summary Table are also those defined in Eq. (2.1).

References

1. E.A. Kuraev, V.S. Fadin, Sov. J. Nucl. Phys. **41**, 466 (1985).

$\eta_b(1S)$

$I^G(J^{PC}) = 0^+(0^{-+})$

Quantum numbers shown are quark-model predictions. Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$.

NODE=M171

NODE=M171

NODE=M171M


NODE=M171M

$\eta_b(1S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9398.7 ± 2.0 OUR AVERAGE Error includes scale factor of 1.5. See the ideogram below.				
9394.8 ^{+2.7} _{-3.1} ± 4.5 _{-2.7}	29k	FULSOM	18 BELL	$\Upsilon(2S) \rightarrow \gamma X$
9400.7 ± 1.7 ± 1.6	33.1k	TAMPONI	15 BELL	$e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
9402.4 ± 1.5 ± 1.8	34k	¹ MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
9391.8 ± 6.6 ± 2.0	2.3k	² BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
9394.2 ^{+4.8} _{-4.9} ± 2.0	13k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
9388.9 ^{+3.1} _{-2.3} ± 2.7	19k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
9393.2 ± 3.4 ± 2.3	10	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$
9300 ± 20 ± 20		HEISTER	02D ALEP	181-209 e^+e^-

- ¹ With floating width. Not independent of the corresponding mass difference measurement.
- ² Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass difference measurements.
- ³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M;LINKAGE=MI
 NODE=M171M;LINKAGE=AU
 NODE=M171M;LINKAGE=DO



$m_{\Upsilon(1S)} - m_{\eta_b}$


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
62.3 ± 3.2 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.				
57.9 ± 1.5 ± 1.8	34k	¹ MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$
68.5 ± 6.6 ± 2.0	2.3 ± 0.5k	² BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
66.1 ^{+4.8} _{-4.9} ± 2.0	13 ± 5k	² AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$
71.4 ^{+2.3} _{-3.1} ± 2.7	19 ± 3k	² AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
67.1 ± 3.4 ± 2.3	10 ⁺⁵ ₋₄	^{2,3} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma \text{hadrons}$

NODE=M171M2

NODE=M171M2

- ¹With floating width. Not independent of the corresponding mass measurement.
- ²Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding γ energy or mass measurements.
- ³Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M171M2;LINKAGE=MI
 NODE=M171M2;LINKAGE=AU
 NODE=M171M2;LINKAGE=DO



γ ENERGY IN $\Upsilon(3S)$ DECAY

NODE=M171DM
 NODE=M171DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
920.6^{+2.8}_{-3.2} OUR AVERAGE				
918.6 ± 6.0 ± 1.9	2.3 ± 0.5k	¹ BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
921.2 ^{+2.1} _{-2.8} ± 2.4	19 ± 3k	¹ AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$

¹Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171DM;LINKAGE=BO

γ ENERGY IN $\Upsilon(2S)$ DECAY

NODE=M171U2S
 NODE=M171U2S

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
609.3^{+4.6}_{-4.5} ± 1.9				
	13 ± 5k	¹ AUBERT	09AQ BABR	$\Upsilon(2S) \rightarrow \gamma X$

¹Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV. Not independent of the corresponding mass or mass difference measurements.

NODE=M171U2S;LINKAGE=AU

$\eta_b(1S)$ WIDTH

NODE=M171W
 NODE=M171W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10⁺⁵₋₄ OUR AVERAGE				
8 ⁺⁶ ₋₅ ± 5	33.1k	¹ TAMPONI	15 BELL	$e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
10.8 ^{+4.0+4.5} _{-3.7-2.0}	34k	¹ MIZUK	12 BELL	$e^+e^- \rightarrow \gamma\pi^+\pi^- + \text{hadrons}$

¹With floating mass.

NODE=M171W;LINKAGE=MI

$\eta_b(1S)$ DECAY MODES

NODE=M171225;NODE=M171

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 hadrons	seen	
Γ_2 $3h^+3h^-$	not seen	
Γ_3 $2h^+2h^-$	not seen	
Γ_4 $4h^+4h^-$	not seen	
Γ_5 $\gamma\gamma$	not seen	
Γ_6 $\mu^+\mu^-$	$<9 \times 10^{-3}$	90%
Γ_7 $\tau^+\tau^-$	$<8\%$	90%

DESIG=7

DESIG=1;OUR EST;→ UNCHECKED ←

DESIG=2;OUR EST;→ UNCHECKED ←

DESIG=4;OUR EST;→ UNCHECKED ←

DESIG=3;OUR EST;→ UNCHECKED ←

DESIG=5

DESIG=6

 $\eta_b(1S) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

NODE=M171230

$$\Gamma(3h^+3h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_2\Gamma_5/\Gamma$$

NODE=M171G1

NODE=M171G1

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<470 95 ABDALLAH 06 DLPH 161-209 e^+e^- <132 95 HEISTER 02D ALEP 181-209 e^+e^-

$$\Gamma(2h^+2h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_3\Gamma_5/\Gamma$$

NODE=M171G2

NODE=M171G2

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<190 95 ABDALLAH 06 DLPH 161-209 e^+e^- < 48 95 HEISTER 02D ALEP 181-209 e^+e^-

$$\Gamma(4h^+4h^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \quad \Gamma_4\Gamma_5/\Gamma$$

NODE=M171G3

NODE=M171G3

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<660 95 ABDALLAH 06 DLPH 161-209 e^+e^- $\eta_b(1S)$ BRANCHING RATIOS

NODE=M171235

$$\Gamma(\text{hadrons})/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

NODE=M171R03

NODE=M171R03

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 34k MIZUK 12 BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- +$
hadrons

$$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

NODE=M171R01

NODE=M171R01

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 9×10^{-3} 90 ¹AUBERT 09Z BABR $e^+e^- \rightarrow \Upsilon(2S, 3S) \rightarrow \gamma\eta_b$ ¹Obtained using $B(\Upsilon(2S) \rightarrow \gamma\eta_b) = (4.2^{+1.1}_{-1.0} \pm 0.9) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma\eta_b)$ $= (4.8 \pm 0.5 \pm 0.6) \times 10^{-4}$. This limit is equivalent to $B(\eta_b \rightarrow \mu^+\mu^-) = (-0.25 \pm 0.51 \pm 0.33)\%$ measurement.

NODE=M171R01;LINKAGE=AU

$$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

NODE=M171R02

NODE=M171R02

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

< 8×10^{-2} 90 AUBERT 09P BABR $e^+e^- \rightarrow \gamma\tau^+\tau^-$ $\eta_b(1S)$ REFERENCES

NODE=M171

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)	REFID=59535
TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=56996
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>		REFID=54288
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=54718
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53062
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52262
ABDALLAH	06	PL B634 340	J.M. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=51042
HEISTER	02D	PL B530 56	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48577

$\Upsilon(1S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M049

 $\Upsilon(1S)$ MASS

NODE=M049M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9460.40±0.09±0.04	¹ SHAMOV	23	RVUE $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9460.11±0.11±0.07	² SHAMOV	23	RVUE $e^+e^- \rightarrow$ hadrons
9460.51±0.09±0.05	^{3,4} ARTAMONOV	00	MD1 $e^+e^- \rightarrow$ hadrons
9460.60±0.09±0.05	^{5,6} BARU	92B	MD1 $e^+e^- \rightarrow$ hadrons
9460.59±0.12	BARU	86	MD1 $e^+e^- \rightarrow$ hadrons
9460.6 ±0.4	^{6,7} ARTAMONOV	84	MD1 $e^+e^- \rightarrow$ hadrons
9459.97±0.11±0.07	⁸ MACKAY	84	CUSB $e^+e^- \rightarrow$ hadrons

NODE=M049M

OCCUR=2

¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.

NODE=M049M;LINKAGE=D

² Obtained by reanalysing CUSB data (MACKAY 84), but not authored by the CUSB collaboration.

NODE=M049M;LINKAGE=E

³ Reanalysis of BARU 92B and ARTAMONOV 84 using new electron mass (COHEN 87).

NODE=M049M;LINKAGE=AR

⁴ Superseded by SHAMOV 23.

NODE=M049M;LINKAGE=B

⁵ Supersedes BARU 86.

NODE=M049M;LINKAGE=A

⁶ Superseded by ARTAMONOV 00.

NODE=M049M;LINKAGE=RZ

⁷ Value includes data of ARTAMONOV 82.

NODE=M049M;LINKAGE=G

⁸ Reanalysed by SHAMOV 23.

NODE=M049M;LINKAGE=C

 $\Upsilon(1S)$ WIDTH

NODE=M049W

VALUE (keV)	DOCUMENT ID
54.02±1.25 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"

NODE=M049W

→ UNCHECKED ←

 $\Upsilon(1S)$ DECAY MODES

NODE=M049215;NODE=M049

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\tau^+ \tau^-$	(2.60 ±0.10) %	
Γ_2 $e^+ e^-$	(2.39 ±0.08) %	
Γ_3 $\mu^+ \mu^-$	(2.48 ±0.04) %	

DESIG=3

DESIG=2

DESIG=1

Hadronic decays

NODE=M049;CLUMP=A

Γ_4 $g g g$	(81.7 ±0.7) %	
Γ_5 $\gamma g g$	(2.2 ±0.6) %	
Γ_6 $\eta'(958)$ anything	(2.94 ±0.24) %	
Γ_7 $J/\psi(1S)$ anything	(5.4 ±0.4) × 10 ⁻⁴	S=1.4
Γ_8 $J/\psi(1S)\eta_c$	< 2.2	× 10 ⁻⁶ CL=90%
Γ_9 $J/\psi(1S)\chi_{c0}$	< 3.4	× 10 ⁻⁶ CL=90%
Γ_{10} $J/\psi(1S)\chi_{c1}$	(3.9 ±1.2) × 10 ⁻⁶	
Γ_{11} $J/\psi(1S)\chi_{c2}$	< 1.4	× 10 ⁻⁶ CL=90%
Γ_{12} $J/\psi(1S)\eta_c(2S)$	< 2.2	× 10 ⁻⁶ CL=90%
Γ_{13} $J/\psi(1S)X(3940)$	< 5.4	× 10 ⁻⁶ CL=90%
Γ_{14} $J/\psi(1S)X(4160)$	< 5.4	× 10 ⁻⁶ CL=90%
Γ_{15} $X(4350)$ anything, $X \rightarrow J/\psi(1S)\phi$	< 8.1	× 10 ⁻⁶ CL=90%
Γ_{16} $T_{c\bar{c}1}(3900)^\pm$ anything, $T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm$	< 1.3	× 10 ⁻⁵ CL=90%
Γ_{17} $T_{c\bar{c}1}(4200)^\pm$ anything, $Z_c \rightarrow J/\psi(1S)\pi^\pm$	< 6.0	× 10 ⁻⁵ CL=90%
Γ_{18} $T_{c\bar{c}1}(4430)^\pm$ anything, $T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm$	< 4.9	× 10 ⁻⁵ CL=90%
Γ_{19} X_{cs}^\pm anything, $X \rightarrow J/\psi K^\pm$	< 5.7	× 10 ⁻⁶ CL=90%
Γ_{20} $\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)\pi^+\pi^-$	< 3.8	× 10 ⁻⁵ CL=90%
Γ_{21} $\psi(4230)$ anything, $\psi \rightarrow J/\psi(1S)K^+K^-$	< 7.5	× 10 ⁻⁶ CL=90%

DESIG=117

DESIG=118

DESIG=73

DESIG=12

DESIG=146

DESIG=147

DESIG=148

DESIG=149

DESIG=150

DESIG=151

DESIG=152

DESIG=167

DESIG=168

DESIG=169

DESIG=170

DESIG=173

DESIG=161

DESIG=165

Γ ₂₂	$\chi_{c1}(4140)$ anything, $\chi_{c1} \rightarrow J/\psi(1S)\phi$	< 5.2	$\times 10^{-6}$	CL=90%	DESIG=166
Γ ₂₃	χ_{c0} anything	< 4	$\times 10^{-3}$	CL=90%	DESIG=5
Γ ₂₄	χ_{c1} anything	(1.90 ± 0.35)	$\times 10^{-4}$		DESIG=6
Γ ₂₅	$\chi_{c1}(1P)X_{tetra}$	< 3.78	$\times 10^{-5}$	CL=90%	DESIG=175
Γ ₂₆	χ_{c2} anything	(2.8 ± 0.8)	$\times 10^{-4}$		DESIG=7
Γ ₂₇	$\psi(2S)$ anything	(1.23 ± 0.20)	$\times 10^{-4}$		DESIG=8
Γ ₂₈	$\psi(2S)\eta_c$	< 3.6	$\times 10^{-6}$	CL=90%	DESIG=153
Γ ₂₉	$\psi(2S)\chi_{c0}$	< 6.5	$\times 10^{-6}$	CL=90%	DESIG=154
Γ ₃₀	$\psi(2S)\chi_{c1}$	< 4.5	$\times 10^{-6}$	CL=90%	DESIG=155
Γ ₃₁	$\psi(2S)\chi_{c2}$	< 2.1	$\times 10^{-6}$	CL=90%	DESIG=156
Γ ₃₂	$\psi(2S)\eta_c(2S)$	< 3.2	$\times 10^{-6}$	CL=90%	DESIG=157
Γ ₃₃	$\psi(2S)X(3940)$	< 2.9	$\times 10^{-6}$	CL=90%	DESIG=158
Γ ₃₄	$\psi(2S)X(4160)$	< 2.9	$\times 10^{-6}$	CL=90%	DESIG=159
Γ ₃₅	$\psi(4230)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 7.9	$\times 10^{-5}$	CL=90%	DESIG=162
Γ ₃₆	$\psi(4360)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 5.2	$\times 10^{-5}$	CL=90%	DESIG=163
Γ ₃₇	$\psi(4660)$ anything, $\psi \rightarrow \psi(2S)\pi^+\pi^-$	< 2.2	$\times 10^{-5}$	CL=90%	DESIG=164
Γ ₃₈	$T_{c\bar{c}}(4050)^\pm$ anything, $X \rightarrow \psi(2S)\pi^\pm$	< 8.8	$\times 10^{-5}$	CL=90%	DESIG=171
Γ ₃₉	$T_{c\bar{c}1}(4430)^\pm$ anything, $T_{c\bar{c}1} \rightarrow \psi(2S)\pi^\pm$	< 6.7	$\times 10^{-5}$	CL=90%	DESIG=172
Γ ₄₀	$\chi_{c1}(3872)$ anything	< 2.6	$\times 10^{-4}$	CL=90%	DESIG=194
Γ ₄₁	$T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-$	< 2.23	$\times 10^{-5}$	CL=90%	DESIG=178
Γ ₄₂	$T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp$	< 8.1	$\times 10^{-6}$	CL=90%	DESIG=179
Γ ₄₃	$T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-$	< 1.8	$\times 10^{-6}$	CL=90%	DESIG=180
Γ ₄₄	$T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$	< 1.58	$\times 10^{-5}$	CL=90%	DESIG=181
Γ ₄₅	$T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$	< 2.66	$\times 10^{-5}$	CL=90%	DESIG=182
Γ ₄₆	$T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp$	< 4.42	$\times 10^{-5}$	CL=90%	DESIG=183
Γ ₄₇	$T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%	DESIG=184
Γ ₄₈	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp$	< 2.33	$\times 10^{-5}$	CL=90%	DESIG=186
Γ ₄₉	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp$	< 4.55	$\times 10^{-5}$	CL=90%	DESIG=189
Γ ₅₀	$\rho\pi$	< 3.68	$\times 10^{-6}$	CL=90%	DESIG=11
Γ ₅₁	$\omega\pi^0$	< 3.90	$\times 10^{-6}$	CL=90%	DESIG=131
Γ ₅₂	$\pi^+\pi^-$	< 5	$\times 10^{-4}$	CL=90%	DESIG=23
Γ ₅₃	K^+K^-	< 5	$\times 10^{-4}$	CL=90%	DESIG=24
Γ ₅₄	$\rho\bar{\rho}$	< 5	$\times 10^{-4}$	CL=90%	DESIG=25
Γ ₅₅	$\pi^+\pi^-\pi^0$	(2.1 ± 0.8)	$\times 10^{-6}$		DESIG=72
Γ ₅₆	ϕK^+K^-	(2.4 ± 0.5)	$\times 10^{-6}$		DESIG=136
Γ ₅₇	$\omega\pi^+\pi^-$	(4.5 ± 1.0)	$\times 10^{-6}$		DESIG=137
Γ ₅₈	$K^*(892)^0 K^-\pi^+ + \text{c.c.}$	(4.4 ± 0.8)	$\times 10^{-6}$		DESIG=138
Γ ₅₉	$\phi f'_2(1525)$	< 1.63	$\times 10^{-6}$	CL=90%	DESIG=139
Γ ₆₀	$\omega f_2(1270)$	< 1.79	$\times 10^{-6}$	CL=90%	DESIG=140
Γ ₆₁	$\rho(770) a_2(1320)$	< 2.24	$\times 10^{-6}$	CL=90%	DESIG=141
Γ ₆₂	$K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}$	(3.0 ± 0.8)	$\times 10^{-6}$		DESIG=142
Γ ₆₃	$K_1(1270)^\pm K^\mp$	< 2.41	$\times 10^{-6}$	CL=90%	DESIG=143
Γ ₆₄	$K_1(1400)^\pm K^\mp$	(1.0 ± 0.4)	$\times 10^{-6}$		DESIG=144
Γ ₆₅	$b_1(1235)^\pm \pi^\mp$	< 1.25	$\times 10^{-6}$	CL=90%	DESIG=145
Γ ₆₆	$\pi^+\pi^-\pi^0\pi^0$	(1.28 ± 0.30)	$\times 10^{-5}$		DESIG=132
Γ ₆₇	$K_S^0 K^+\pi^- + \text{c.c.}$	(1.6 ± 0.4)	$\times 10^{-6}$		DESIG=133
Γ ₆₈	$K^*(892)^0 \bar{K}^0 + \text{c.c.}$	(2.9 ± 0.9)	$\times 10^{-6}$		DESIG=134
Γ ₆₉	$K^*(892)^- K^+ + \text{c.c.}$	< 1.11	$\times 10^{-6}$	CL=90%	DESIG=135
Γ ₇₀	$f_1(1285)$ anything	(4.6 ± 3.1)	$\times 10^{-3}$		DESIG=174
Γ ₇₁	$D^*(2010)^\pm$ anything	(2.52 ± 0.20)	%		DESIG=30
Γ ₇₂	$\frac{f_1(1285)}{2} X_{tetra}$	< 6.24	$\times 10^{-5}$	CL=90%	DESIG=176
Γ ₇₃	2H anything	(2.85 ± 0.25)	$\times 10^{-5}$		DESIG=107
Γ ₇₄	Sum of 100 exclusive modes	(1.200 ± 0.017)	%		DESIG=128

Radiative decays

				NODE=M049;CLUMP=B
Γ ₇₅	$\gamma\pi^+\pi^-$	(6.3 ±1.8)	$\times 10^{-5}$	DESIG=70
Γ ₇₆	$\gamma\pi^0\pi^0$	(1.7 ±0.7)	$\times 10^{-5}$	DESIG=71
Γ ₇₇	$\gamma\pi\pi$ (S-wave)	(4.6 ±0.7)	$\times 10^{-5}$	DESIG=190
Γ ₇₈	$\gamma\pi^0\eta$	< 2.4	$\times 10^{-6}$	CL=90% DESIG=111
Γ ₇₉	$\gamma K^+ K^-$	[a] (1.14 ±0.13)	$\times 10^{-5}$	DESIG=102
Γ ₈₀	$\gamma p\bar{p}$	[b] < 6	$\times 10^{-6}$	CL=90% DESIG=103
Γ ₈₁	$\gamma 2h^+ 2h^-$	(7.0 ±1.5)	$\times 10^{-4}$	DESIG=20
Γ ₈₂	$\gamma 3h^+ 3h^-$	(5.4 ±2.0)	$\times 10^{-4}$	DESIG=21
Γ ₈₃	$\gamma 4h^+ 4h^-$	(7.4 ±3.5)	$\times 10^{-4}$	DESIG=22
Γ ₈₄	$\gamma\pi^+\pi^- K^+ K^-$	(2.9 ±0.9)	$\times 10^{-4}$	DESIG=14
Γ ₈₅	$\gamma 2\pi^+ 2\pi^-$	(2.5 ±0.9)	$\times 10^{-4}$	DESIG=13
Γ ₈₆	$\gamma 3\pi^+ 3\pi^-$	(2.5 ±1.2)	$\times 10^{-4}$	DESIG=17
Γ ₈₇	$\gamma 2\pi^+ 2\pi^- K^+ K^-$	(2.4 ±1.2)	$\times 10^{-4}$	DESIG=18
Γ ₈₈	$\gamma\pi^+\pi^- p\bar{p}$	(1.5 ±0.6)	$\times 10^{-4}$	DESIG=15
Γ ₈₉	$\gamma 2\pi^+ 2\pi^- p\bar{p}$	(4 ±6)	$\times 10^{-5}$	DESIG=19
Γ ₉₀	$\gamma 2K^+ 2K^-$	(2.0 ±2.0)	$\times 10^{-5}$	DESIG=16
Γ ₉₁	$\gamma\eta'(958)$	< 1.9	$\times 10^{-6}$	CL=90% DESIG=55
Γ ₉₂	$\gamma\eta$	< 1.0	$\times 10^{-6}$	CL=90% DESIG=54
Γ ₉₃	$\gamma f_0(980)$	< 3	$\times 10^{-5}$	CL=90% DESIG=105
Γ ₉₄	$\gamma f_2'(1525)$	(2.9 ±0.6)	$\times 10^{-5}$	DESIG=52
Γ ₉₅	$\gamma f_2(1270)$	(1.01 ±0.06)	$\times 10^{-4}$	DESIG=51
Γ ₉₆	$\gamma\eta(1405)$	< 8.2	$\times 10^{-5}$	CL=90% DESIG=65
Γ ₉₇	$\gamma f_0(1500)$	< 1.5	$\times 10^{-5}$	CL=90% DESIG=108
Γ ₉₈	$\gamma f_0(1500) \rightarrow \gamma K^+ K^-$	(1.0 ±0.4)	$\times 10^{-5}$	DESIG=192
Γ ₉₉	$\gamma f_0(1710)$	< 2.6	$\times 10^{-4}$	CL=90% DESIG=53
Γ ₁₀₀	$\gamma f_0(1710) \rightarrow \gamma K^+ K^-$	(1.01 ±0.32)	$\times 10^{-5}$	DESIG=112
Γ ₁₀₁	$\gamma f_0(1710) \rightarrow \gamma\pi^+\pi^-$	(5.3 ±2.0)	$\times 10^{-6}$	DESIG=191
Γ ₁₀₂	$\gamma f_0(1710) \rightarrow \gamma\pi^0\pi^0$	< 1.4	$\times 10^{-6}$	CL=90% DESIG=109
Γ ₁₀₃	$\gamma f_0(1710) \rightarrow \gamma\eta\eta$	< 1.8	$\times 10^{-6}$	CL=90% DESIG=110
Γ ₁₀₄	$\gamma f_4(2050)$	< 5.3	$\times 10^{-5}$	CL=90% DESIG=104
Γ ₁₀₅	$\gamma f_0(2200) \rightarrow \gamma K^+ K^-$	< 2	$\times 10^{-4}$	CL=90% DESIG=69
Γ ₁₀₆	$\gamma f_J(2220) \rightarrow \gamma K^+ K^-$	< 8	$\times 10^{-7}$	CL=90% DESIG=60
Γ ₁₀₇	$\gamma f_J(2220) \rightarrow \gamma\pi^+\pi^-$	< 6	$\times 10^{-7}$	CL=90% DESIG=61
Γ ₁₀₈	$\gamma f_J(2220) \rightarrow \gamma p\bar{p}$	< 1.1	$\times 10^{-6}$	CL=90% DESIG=62
Γ ₁₀₉	$\gamma\eta(2225) \rightarrow \gamma\phi\phi$	< 3	$\times 10^{-3}$	CL=90% DESIG=68
Γ ₁₁₀	$\gamma\eta_c(1S)$	< 2.9	$\times 10^{-5}$	CL=90% DESIG=119
Γ ₁₁₁	$\gamma\eta_c(2S)$	< 4	$\times 10^{-4}$	CL=90% DESIG=193
Γ ₁₁₂	$\gamma\chi_{c0}$	< 6.6	$\times 10^{-5}$	CL=90% DESIG=120
Γ ₁₁₃	$\gamma\chi_{c1}$	(4.7 ^{+2.4} _{-1.9})	$\times 10^{-5}$	DESIG=121
Γ ₁₁₄	$\gamma\chi_{c2}$	< 7.6	$\times 10^{-6}$	CL=90% DESIG=122
Γ ₁₁₅	$\gamma\chi_{c1}(3872)$	< 4	$\times 10^{-5}$	CL=90% DESIG=195
Γ ₁₁₆	$\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi$	< 2.8	$\times 10^{-6}$	CL=90% DESIG=124
Γ ₁₁₇	$\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi$	< 3.0	$\times 10^{-6}$	CL=90% DESIG=125
Γ ₁₁₈	$\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi$	< 2.2	$\times 10^{-6}$	CL=90% DESIG=126
Γ ₁₁₉	$\gamma X\bar{X} (m_X < 3.1 \text{ GeV})$	[c] < 1	$\times 10^{-3}$	CL=90% DESIG=67
Γ ₁₂₀	$\gamma X\bar{X} (m_X < 4.5 \text{ GeV})$	[d] < 2.4	$\times 10^{-4}$	CL=90% DESIG=127
Γ ₁₂₁	$\gamma X \rightarrow \gamma + \geq 4 \text{ prongs}$	[e] < 1.78	$\times 10^{-4}$	CL=95% DESIG=113
Γ ₁₂₂	γA^0	[f]		DESIG=66
Γ ₁₂₃	$\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	[g] < 9	$\times 10^{-6}$	CL=90% DESIG=114
Γ ₁₂₄	$\gamma A^0 \rightarrow \gamma\tau^+\tau^-$	[a] < 1.30	$\times 10^{-4}$	CL=90% DESIG=115
Γ ₁₂₅	$\gamma A^0 \rightarrow \gamma g g$	[h] < 1	%	CL=90% DESIG=129
Γ ₁₂₆	$\gamma A^0 \rightarrow \gamma s\bar{s}$	[h] < 1	$\times 10^{-3}$	CL=90% DESIG=130

Lepton Family number (LF) violating modes

				NODE=M049;CLUMP=C
Γ ₁₂₇	$e^\pm \mu^\mp$	LF	< 3.9	$\times 10^{-7}$ CL=90% DESIG=196
Γ ₁₂₈	$\mu^\pm \tau^\mp$	LF	< 2.7	$\times 10^{-6}$ CL=90% DESIG=116
Γ ₁₂₉	$e^\pm \tau^\mp$	LF	< 2.7	$\times 10^{-6}$ CL=90% DESIG=197
Γ ₁₃₀	$\gamma e^\pm \mu^\mp$	LF	< 4.2	$\times 10^{-7}$ CL=90% DESIG=198
Γ ₁₃₁	$\gamma \mu^\pm \tau^\mp$	LF	< 6.1	$\times 10^{-6}$ CL=90% DESIG=199
Γ ₁₃₂	$\gamma e^\pm \tau^\mp$	LF	< 6.5	$\times 10^{-6}$ CL=90% DESIG=200

Other decays

Γ_{133} invisible $< 3.0 \times 10^{-4}$ CL=90%
 Γ_{134} hadrons $(96 \pm 4) \%$

NODE=M049;CLUMP=D
 DESIG=106
 DESIG=101

[a] $2m_\tau < M(\tau^+\tau^-) < 9.2$ GeV

LINKAGE=E49

[b] 2 GeV $< m_{K^+K^-} < 3$ GeV

LINKAGE=G49

[c] $X\bar{X}$ = vectors with $m < 3.1$ GeV

LINKAGE=B49

[d] X and \bar{X} = zero spin with $m < 4.5$ GeV

LINKAGE=F49

[e] 1.5 GeV $< m_X < 5.0$ GeV

LINKAGE=C49

[f] A^0 = scalar with $m < 8.0$ GeV

LINKAGE=A49

[g] 201 MeV $< M(\mu^+\mu^-) < 3565$ MeV

LINKAGE=D49

[h] 0.5 GeV $< m_X < 9.0$ GeV, where m_X is the invariant mass of the hadronic final state.

LINKAGE=I49

 $\Upsilon(1S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M049218

 $\Gamma(\mu^+\mu^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_3\Gamma_2/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
31.2±1.6±1.7	KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$

NODE=M049G1
 NODE=M049G1

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{134}\Gamma_2/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
1.240±0.016 OUR AVERAGE			
1.252±0.004±0.019	¹ ROSNER	06	CLEO $9.5 e^+e^- \rightarrow \text{hadrons}$
1.187±0.023±0.031	¹ BARU	92B	MD1 $e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.02 ±0.05	¹ JAKUBOWSKI	88	CBAL $e^+e^- \rightarrow \text{hadrons}$
1.37 ±0.06 ±0.09	² GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$
1.23 ±0.08 ±0.04	² ALBRECHT	82	DASP $e^+e^- \rightarrow \text{hadrons}$
1.13 ±0.07 ±0.11	² NICZYPORUK	82	LENA $e^+e^- \rightarrow \text{hadrons}$
1.09 ±0.25	² BOCK	80	CNTR $e^+e^- \rightarrow \text{hadrons}$
1.35 ±0.14	³ BERGER	79	PLUT $e^+e^- \rightarrow \text{hadrons}$

NODE=M049G2
 NODE=M049G2

¹ Radiative corrections evaluated following KURAEV 85.

² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

³ Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

NODE=M049G2;LINKAGE=B
 NODE=M049G2;LINKAGE=R
 NODE=M049G2;LINKAGE=P

 $\Upsilon(1S)$ PARTIAL WIDTHS

NODE=M049220

 $\Gamma(e^+e^-)$ Γ_2

VALUE (keV)	DOCUMENT ID
1.340±0.018 OUR EVALUATION	

NODE=M049W2
 NODE=M049W2
 → UNCHECKED ←

 $\Upsilon(1S)$ BRANCHING RATIOS

NODE=M049225

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.60±0.10 OUR AVERAGE				
2.53±0.13±0.04	60k	¹ BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S) \rightarrow \tau^+\tau^-$
2.61±0.12 ^{+0.09} _{-0.13}	25k	CINABRO	94B	CLE2 $e^+e^- \rightarrow \tau^+\tau^-$
2.7 ±0.4 ±0.2		² ALBRECHT	85C	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$
3.4 ±0.4 ±0.4		GILES	83	CLEO $e^+e^- \rightarrow \tau^+\tau^-$

NODE=M049R3
 NODE=M049R3

¹ BESSON 07 reports $[\Gamma(\Upsilon(1S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = 1.02 \pm 0.02 \pm 0.05$ which we multiply by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M049R3;LINKAGE=BE

² Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.

NODE=M049R3;LINKAGE=A

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.39±0.08 OUR AVERAGE				
2.40±0.01±0.12	191k	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.29±0.08±0.11		ALEXANDER	98	CLE2 $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.42±0.14±0.14	307	ALBRECHT	87	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
2.8 ±0.3 ±0.2	826	BESSON	84	CLEO $\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
5.1 ±3.0		BERGER	80C	PLUT $e^+e^- \rightarrow e^+e^-$

NODE=M049R2
 NODE=M049R2

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M049R1
NODE=M049R1

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.48±0.04 OUR AVERAGE				
2.46±0.01±0.11	246k	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
2.49±0.02±0.07	345k	ADAMS	05	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
2.49±0.08±0.13		ALEXANDER	98	CLE2 $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
2.12±0.20±0.10		¹ BARU	92	MD1 $e^+e^- \rightarrow \mu^+\mu^-$
2.31±0.12±0.10		¹ KOBEL	92	CBAL $e^+e^- \rightarrow \mu^+\mu^-$
2.52±0.07±0.07		CHEN	89B	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
2.61±0.09±0.11		KAARSBERG	89	CSB2 $e^+e^- \rightarrow \mu^+\mu^-$
2.30±0.25±0.13	86	ALBRECHT	87	ARG $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
2.9 ±0.3 ±0.2	864	BESSON	84	CLEO $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
2.7 ±0.3 ±0.3		ANDREWS	83	CLEO $e^+e^- \rightarrow \mu^+\mu^-$
3.2 ±1.3 ±0.3		ALBRECHT	82	DASP $e^+e^- \rightarrow \mu^+\mu^-$
3.8 ±1.5 ±0.2		NICZYPORUK	82	LENA $e^+e^- \rightarrow \mu^+\mu^-$
1.4 ^{+3.4} _{-1.4}		BOCK	80	CNTR $e^+e^- \rightarrow \mu^+\mu^-$
2.2 ±2.0		BERGER	79	PLUT $e^+e^- \rightarrow \mu^+\mu^-$

¹ Taking into account interference between the resonance and continuum.

NODE=M049R1;LINKAGE=G

 $\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_1/Γ_3 NODE=M049R43
NODE=M049R43

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.008±0.023 OUR AVERAGE				
1.005±0.013±0.022	0.7M	¹ DEL-AMO-SA..10c	BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^- \Upsilon(1S)$
1.02 ±0.02 ±0.05	60k	BESSON	07	CLEO $e^+e^- \rightarrow \Upsilon(1S)$

¹ Allows any number of extra photons with total energy < 500 MeV.

NODE=M049R43;LINKAGE=DE

 $\Gamma(ggg)/\Gamma_{\text{total}}$ Γ_4/Γ NODE=M049R35
NODE=M049R35

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
81.7±0.7	20M	¹ BESSON	06A	CLEO $\Upsilon(1S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ from BESSON 06A and PDG 08 values of $B(\mu^+\mu^-) = (2.48 \pm 0.05)\%$ and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

NODE=M049R35;LINKAGE=BE

 $\Gamma(\gamma gg)/\Gamma_{\text{total}}$ Γ_5/Γ NODE=M049R36
NODE=M049R36

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.20±0.60	400k	¹ BESSON	06A	CLEO $\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$

¹ Calculated using BESSON 06A values of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.70 \pm 0.01 \pm 0.13 \pm 0.24)\%$ and $\Gamma(ggg)/\Gamma_{\text{total}}$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(ggg)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

NODE=M049R36;LINKAGE=BE

 $\Gamma(\gamma gg)/\Gamma(ggg)$ Γ_5/Γ_4 NODE=M049R37
NODE=M049R37

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.70±0.01±0.27	20M	BESSON	06A	CLEO $\Upsilon(1S) \rightarrow (\gamma +) \text{hadrons}$

 $\Gamma(\eta'(958) \text{ anything})/\Gamma_{\text{total}}$ Γ_6/Γ NODE=M049R73
NODE=M049R73

VALUE	DOCUMENT ID	TECN	COMMENT
0.0294±0.0024 OUR AVERAGE			
0.030 ±0.002 ±0.002	AQUINES	06A	CLE3 $\Upsilon(1S) \rightarrow \eta' \text{ anything}$
0.028 ±0.004 ±0.002	ARTUSO	03	CLE2 $\Upsilon(1S) \rightarrow \eta' \text{ anything}$

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_7/Γ NODE=M049R12
NODE=M049R12

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.4 ±0.4 OUR FIT Error includes scale factor of 1.4.					
5.4 ±0.4 OUR AVERAGE Error includes scale factor of 1.5.					
5.25±0.13±0.25		3k	SHEN	16	BELL $e^+e^- \rightarrow J/\psi X$
6.4 ±0.4 ±0.6		730	BRIERE	04	CLEO $e^+e^- \rightarrow J/\psi X$
11 ±4 ±2			¹ FULTON	89	CLEO $e^+e^- \rightarrow \mu^+\mu^- X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8	90	ALBRECHT	92J	ARG	$e^+e^- \rightarrow e^+e^-X$, $\mu^+\mu^-X$
<17	90	MASCHMANN	90	CBAL	$e^+e^- \rightarrow \text{hadrons}$
<200	90	NICZYPORUK	83	LENA	

¹ Using $B((J/\psi) \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.

NODE=M049R12;LINKAGE=K

$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10 ⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R85
NODE=M049R85

$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.4 × 10 ⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R86
NODE=M049R86

$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10 ⁻⁶)	EVT%	DOCUMENT ID	TECN	COMMENT
3.90 ± 1.21 ± 0.23	20	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R87
NODE=M049R87

$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.4 × 10 ⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R88
NODE=M049R88

$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.2 × 10 ⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R89
NODE=M049R89

$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10 ⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R90
NODE=M049R90

$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.4 × 10 ⁻⁶	90	YANG	14	BELL $e^+e^- \rightarrow J/\psi X$

NODE=M049R91
NODE=M049R91

$\Gamma(X(4350) \text{ anything}, X \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.1 × 10 ⁻⁶	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

NODE=M049P05
NODE=M049P05

$\Gamma(T_{c\bar{c}1}(3900)^\pm \text{ anything}, T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.3 × 10 ⁻⁵	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

NODE=M049P06
NODE=M049P06

$\Gamma(T_{c\bar{c}1}(4200)^\pm \text{ anything}, Z_c \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.0 × 10 ⁻⁵	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

NODE=M049P07
NODE=M049P07

$\Gamma(T_{c\bar{c}1}(4430)^\pm \text{ anything}, T_{c\bar{c}1} \rightarrow J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.9 × 10 ⁻⁵	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$

NODE=M049P08
NODE=M049P08

$\Gamma(X_{cs}^\pm \text{ anything}, X \rightarrow J/\psi K^\pm)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.7 × 10 ⁻⁶	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^- X$

NODE=M049P11
NODE=M049P11

$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.8 × 10 ⁻⁵	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi \pi^+ \pi^- X$

NODE=M049R99
NODE=M049R99

$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow J/\psi(1S)K^+K^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.5 × 10 ⁻⁶	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

NODE=M049P03
NODE=M049P03

$\Gamma(\chi_{c1}(4140) \text{ anything}, \chi_{c1} \rightarrow J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-6}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow J/\psi K^+ K^- X$

NODE=M049P04
 NODE=M049P04

 $\Gamma(\chi_{c0} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{23}/Γ_7

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.4	90	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$

NODE=M049R25
 NODE=M049R25

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.90±0.35 OUR FIT				
1.90±0.43±0.14	215	JIA	17	BELL $\Upsilon(1S) \rightarrow \gamma J/\psi(1S)$

NODE=M049P13
 NODE=M049P13

 $\Gamma(\chi_{c1} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{24}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.35±0.07 OUR FIT				
0.35±0.08±0.06	52 ± 12	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$

NODE=M049R26
 NODE=M049R26

 $\Gamma(\chi_{c1}(1P)\chi_{tetra})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<37.8 \times 10^{-6}$	90	¹ JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

NODE=M049P15
 NODE=M049P15

¹ For a tetraquark state χ_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of χ_{tetra} mass and width range from 4.4×10^{-6} to 37.8×10^{-6} .

NODE=M049P15;LINKAGE=A

 $\Gamma(\chi_{c2} \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{26}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.52±0.12±0.09	47 ± 11	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi X$

NODE=M049R27
 NODE=M049R27

 $\Gamma(\psi(2S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.23±0.17±0.11	215	SHEN	16	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049P12
 NODE=M049P12

 $\Gamma(\psi(2S) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ Γ_{27}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.41±0.11±0.08	42 ± 11	BRIERE	04	CLEO $e^+ e^- \rightarrow J/\psi \pi^+ \pi^- X$

NODE=M049R28
 NODE=M049R28

 $\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R92
 NODE=M049R92

 $\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.5 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R93
 NODE=M049R93

 $\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.5 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R94
 NODE=M049R94

 $\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.1 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R95
 NODE=M049R95

 $\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R96
 NODE=M049R96

 $\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R97
 NODE=M049R97

 $\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-6}$	90	YANG	14	BELL $e^+ e^- \rightarrow \psi(2S) X$

NODE=M049R98
 NODE=M049R98

$\Gamma(\psi(4230) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_{35}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.9 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^- X$

NODE=M049P00
NODE=M049P00 **$\Gamma(\psi(4360) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$** **$\Gamma_{36}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^- X$

NODE=M049P01
NODE=M049P01 **$\Gamma(\psi(4660) \text{ anything}, \psi \rightarrow \psi(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$** **$\Gamma_{37}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^+\pi^- X$

NODE=M049P02
NODE=M049P02 **$\Gamma(T_{c\bar{c}}(4050)^\pm \text{ anything}, X \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}$** **$\Gamma_{38}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.8 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

NODE=M049P09
NODE=M049P09 **$\Gamma(T_{c\bar{c}1}(4430)^\pm \text{ anything}, T_{c\bar{c}1} \rightarrow \psi(2S)\pi^\pm)/\Gamma_{\text{total}}$** **$\Gamma_{39}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.7 \times 10^{-5}$	90	SHEN	16	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

NODE=M049P10
NODE=M049P10 **$\Gamma(\chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}$** **$\Gamma_{40}/\Gamma$**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6 \times 10^{-4}$ (CL = 90%)				$[<2.5 \times 10^{-4}$ (CL = 90%) OUR 2023 BEST LIMIT]
$<2.7 \times 10^{-4}$	90	¹ SHEN	16	BELL $\Upsilon(1S) \rightarrow$ $J/\psi\pi^+\pi^- X$

NODE=M049P31
NODE=M049P31

¹ SHEN 16 reports $[\Gamma(\Upsilon(1S) \rightarrow \chi_{c1}(3872) \text{ anything})/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] < 9.5 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$.

NODE=M049P31;LINKAGE=A

 $\Gamma(T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-)/\Gamma_{\text{total}}$ **Γ_{41}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<22.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$

NODE=M049P17
NODE=M049P17

¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1$.

NODE=M049P17;LINKAGE=A

 $\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}}$ **Γ_{42}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.1 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$

NODE=M049P18
NODE=M049P18

¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi\pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi\pi^\pm)$.

NODE=M049P18;LINKAGE=A

 $\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$ **Γ_{43}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow J/\psi\pi^\pm X$

NODE=M049P19
NODE=M049P19

¹ Assuming $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi\pi^\pm) = 1$

NODE=M049P19;LINKAGE=A

 $\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}}$ **Γ_{44}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<15.8 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P)\pi^\pm X$

NODE=M049P20
NODE=M049P20

¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1$

NODE=M049P20;LINKAGE=A

 $\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}}$ **Γ_{45}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<26.6 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P)\pi^\pm X$

NODE=M049P21
NODE=M049P21

¹ Assuming $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1$

NODE=M049P21;LINKAGE=A

 $\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}}$ **Γ_{46}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<44.2 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \chi_{c1}(1P)\pi^\pm X$

NODE=M049P22
NODE=M049P22

¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P)\pi^\pm)$

NODE=M049P22;LINKAGE=A

 $\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}}$ **Γ_{47}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<20.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(2S) \rightarrow \psi(2S)\pi^\pm X$

NODE=M049P23
NODE=M049P23

¹ Assuming $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S)\pi^\pm) = 1$

NODE=M049P23;LINKAGE=A

$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<23.3 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S)\pi^\pm) = 1$ NODE=M049P25
NODE=M049P25

NODE=M049P25;LINKAGE=A

 $\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4430)^\mp)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<45.5 \times 10^{-6}$	90	¹ JIA	18	BELL $\Upsilon(1S) \rightarrow \psi(2S)\pi^\pm X$

¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S)\pi^\pm) = 1 = B(T_{c\bar{c}}(4430)^\pm \rightarrow \psi(2S)\pi^\pm)$ NODE=M049P26
NODE=M049P26

NODE=M049P26;LINKAGE=A

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.68	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<1 \times 10^3$	90	BLINOV	90	MD1 $\Upsilon(1S) \rightarrow \rho^0\pi^0$
$<2 \times 10^2$	90	FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0\pi^0$
$<2.1 \times 10^3$	90	NICZYPORUK	83	LENA $\Upsilon(1S) \rightarrow \rho^0\pi^0$

NODE=M049R11
NODE=M049R11 $\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.90	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0\pi^0$

NODE=M049R05
NODE=M049R05 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow \pi^+\pi^-$

NODE=M049R57
NODE=M049R57 $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	BARU	92	MD1 $\Upsilon(1S) \rightarrow K^+K^-$

NODE=M049R58
NODE=M049R58 $\Gamma(p\bar{p})/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ BARU	96	MD1 $\Upsilon(1S) \rightarrow p\bar{p}$

¹ Supersedes BARU 92 in this node.NODE=M049R59
NODE=M049R59

NODE=M049R59;LINKAGE=A

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$2.14 \pm 0.72 \pm 0.34$		26 ± 9	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<18.4	90	ANASTASSOV	99	CLE2 $e^+e^- \rightarrow \text{hadrons}$
---------	----	------------	----	--

NODE=M049R72
NODE=M049R72 $\Gamma(\phi K^+K^-)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.36 \pm 0.37 \pm 0.29$	56	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+K^-)$

NODE=M049R75
NODE=M049R75 $\Gamma(\omega\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.46 \pm 0.67 \pm 0.72$	64	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M049R76
NODE=M049R76 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.42 \pm 0.50 \pm 0.58$	173	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+K^-\pi^+\pi^-$

NODE=M049R77
NODE=M049R77 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.63	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(K^+K^-)$

NODE=M049R78
NODE=M049R78 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.79	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+\pi^-)\pi^0$

NODE=M049R79
NODE=M049R79

$\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.24	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M049R80
NODE=M049R80

 $\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.02 \pm 0.68 \pm 0.34$	42	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R81
NODE=M049R81

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.41	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R82
NODE=M049R82

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.02 \pm 0.35 \pm 0.22$	24	SHEN	12A	BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

NODE=M049R83
NODE=M049R83

 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{65}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.25	90	SHEN	12A	BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

NODE=M049R84
NODE=M049R84

 $\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{66}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$12.8 \pm 2.0 \pm 2.3$	143 \pm 22	SHEN	13	BELL $\Upsilon(1S) \rightarrow \pi^+ \pi^- \pi^0 \pi^0$

NODE=M049R06
NODE=M049R06

 $\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{67}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$1.59 \pm 0.33 \pm 0.18$		37 \pm 8	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R07
NODE=M049R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.4	90	¹ DOBBS	12A	$\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$
------	----	--------------------	-----	--

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M049R07;LINKAGE=DO

 $\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{68}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.92 \pm 0.85 \pm 0.37$	16 \pm 5	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R08
NODE=M049R08

 $\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{69}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.11	90	SHEN	13	BELL $\Upsilon(1S) \rightarrow K_S^0 K^- \pi^+$

NODE=M049R09
NODE=M049R09

 $\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$ Γ_{70}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 2.8 \pm 1.3$	3.1k	JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

NODE=M049P14
NODE=M049P14

 $\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{71}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$25.2 \pm 1.3 \pm 1.5$		\approx 2k	¹ AUBERT	10C	BABR $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

NODE=M049R32
NODE=M049R32

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	² ALBRECHT	92J	ARG $e^+ e^- \rightarrow D^0 \pi^\pm X$
-----	----	-----------------------	-----	---

¹ For $x_p > 0.1$.

² For $x_p > 0.2$.

NODE=M049R32;LINKAGE=AU

NODE=M049R32;LINKAGE=B

 $\Gamma(f_1(1285) X_{tetra})/\Gamma_{\text{total}}$ Γ_{72}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<62.4 \times 10 ⁻⁶	90	¹ JIA	17A	BELL $e^+ e^- \rightarrow \text{hadrons}$

NODE=M049P16
NODE=M049P16

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.6×10^{-6} to 62.4×10^{-6} .

NODE=M049P16;LINKAGE=A

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{73}/Γ
2.85±0.25 OUR AVERAGE					
$2.81 \pm 0.49^{+0.20}_{-0.24}$		LEES	14G	BABR $e^+ e^- \rightarrow \overline{2H} X$	OCCUR=2
$2.86 \pm 0.19 \pm 0.21$	455	ASNER	07	CLEO $e^+ e^- \rightarrow \overline{2H} X$	

NODE=M049R33
NODE=M049R33

 $\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	DOCUMENT ID	COMMENT	Γ_{74}/Γ
1.200±0.017	1,2 DOBBS	12A $\Upsilon(1S) \rightarrow \text{hadrons}$	

NODE=M049R02
NODE=M049R02

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

NODE=M049R02;LINKAGE=DO

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M049R02;LINKAGE=NC

 $\Gamma(ggg, \gamma gg \rightarrow \bar{d} \text{ anything})/\Gamma(ggg, \gamma gg \rightarrow \text{anything})$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{75}/Γ
3.36±0.23±0.25	455	ASNER	07	CLEO $e^+ e^- \rightarrow \bar{d} X$	

NODE=M049R34
NODE=M049R34

 $\Gamma(\gamma\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_{75}/Γ
6.3±1.2±1.3	¹ ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R70
NODE=M049R70

¹ For $m_{\pi\pi} > 1$ GeV.

NODE=M049R70;LINKAGE=A

 $\Gamma(\gamma\pi^0\pi^0)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_{76}/Γ
1.7±0.6±0.3	¹ ANASTASSOV 99	CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R71
NODE=M049R71

¹ For $m_{\pi\pi} > 1$ GeV.

NODE=M049R71;LINKAGE=A

 $\Gamma(\gamma\pi\pi(\text{S-wave}))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_{77}/Γ
4.63±0.56±0.48	LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma\pi^+\pi^-$	

NODE=M049P27
NODE=M049P27

 $\Gamma(\gamma\pi^0\eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{78}/Γ
<2.4	90	¹ BESSON 07A	CLEO	$e^+ e^- \rightarrow \Upsilon(1S)$	

NODE=M049R47
NODE=M049R47

¹ BESSON 07A obtained this limit for $0.7 < m_{\pi^0\eta} < 3$ GeV.

NODE=M049R47;LINKAGE=BE

 $\Gamma(\gamma K^+ K^-)/\Gamma_{\text{total}}$

($2 < m_{K^+ K^-} < 3$ GeV)

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{79}/Γ
1.14±0.08±0.10	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

NODE=M049R24
NODE=M049R24

NODE=M049R24

 $\Gamma(\gamma p\bar{p})/\Gamma_{\text{total}}$

($2 < m_{p\bar{p}} < 3$ GeV)

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{80}/Γ
<0.6	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma p\bar{p}$	

NODE=M049R29

NODE=M049R29
NODE=M049R29

 $\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{81}/Γ
7.0±1.1±1.0	80 ± 12	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R20
NODE=M049R20

 $\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{82}/Γ
5.4±1.5±1.3	39 ± 11	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R21
NODE=M049R21

 $\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{83}/Γ
7.4±2.5±2.5	36 ± 12	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R22
NODE=M049R22

 $\Gamma(\gamma\pi^+\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{84}/Γ
2.9±0.7±0.6	29 ± 8	FULTON	90B	CLEO $e^+ e^- \rightarrow \text{hadrons}$	

NODE=M049R14
NODE=M049R14

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{85}/Γ
$2.5 \pm 0.7 \pm 0.5$	26 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M049R13 NODE=M049R13

 $\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{86}/Γ
$2.5 \pm 0.9 \pm 0.8$	17 ± 5	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M049R17 NODE=M049R17

 $\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{87}/Γ
$2.4 \pm 0.9 \pm 0.8$	18 ± 7	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M049R18 NODE=M049R18

 $\Gamma(\gamma \pi^+ \pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{88}/Γ
$1.5 \pm 0.5 \pm 0.3$	22 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M049R15 NODE=M049R15

 $\Gamma(\gamma 2\pi^+ 2\pi^- \rho \bar{\rho})/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{89}/Γ
$0.4 \pm 0.4 \pm 0.4$	7 ± 6	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M049R19 NODE=M049R19

 $\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{90}/Γ
0.2 ± 0.2	2 ± 2	FULTON	90B CLEO	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M049R16 NODE=M049R16

 $\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{91}/Γ
< 1.9	90	ATHAR 07A	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \pi^+ \pi^- \eta, \gamma \rho$	NODE=M049R55 NODE=M049R55
••• We do not use the following data for averages, fits, limits, etc. •••					
< 16	90	RICHICHI 01B	CLE2	$\Upsilon(1S) \rightarrow \gamma \eta' \rightarrow \gamma \eta \pi^+ \pi^-$	

 $\Gamma(\gamma \eta)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{92}/Γ
< 1.0	90	ATHAR 07A	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta \rightarrow \gamma \gamma \gamma, \gamma \pi^+ \pi^- \pi^0, \gamma 3\pi^0$	NODE=M049R54 NODE=M049R54
••• We do not use the following data for averages, fits, limits, etc. •••					
< 21	90	MASEK 02	CLEO	$\Upsilon(1S) \rightarrow \gamma \eta$	

 $\Gamma(\gamma f_0(980))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{93}/Γ
< 3	90	¹ ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	NODE=M049R31 NODE=M049R31

¹ Assuming $B(f_0(980) \rightarrow \pi\pi) = 1$.

NODE=M049R31;LINKAGE=AT

 $\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{94}/Γ
2.9 ± 0.6			OUR AVERAGE			NODE=M049R52 NODE=M049R52
$2.13 \pm 0.28 \pm 0.72$			¹ LEES 18A	BABR	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
$4.1 \pm 1.4 \pm 0.1$		17	² BESSON 11	CLEO	$\Upsilon(1S) \rightarrow K_S^0 K_S^0$	
$3.7 \pm 0.9 \pm 0.8$			ATHAR 06	CLE3	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

••• We do not use the following data for averages, fits, limits, etc. •••

< 14	90	³ FULTON 90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
< 19.4	90	³ ALBRECHT 89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

¹ Using $B(f_2'(1525) \rightarrow K\bar{K}) = 0.887 \pm 0.022$ and $B(K^0 \bar{K}^0) = 1/2 B(K\bar{K})$.

² BESSON 11 reports $(4.0 \pm 1.3 \pm 0.6) \times 10^{-5}$ from a measurement of $[\Gamma(\Upsilon(1S) \rightarrow \gamma f_2'(1525))/\Gamma_{\text{total}}] \times [B(f_2'(1525) \rightarrow K\bar{K})]$ assuming $B(f_2'(1525) \rightarrow K\bar{K}) = (88.8 \pm 3.1) \times 10^{-2}$, which we rescale to our best value $B(f_2'(1525) \rightarrow K\bar{K}) = (87.6 \pm 2.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. The result also assumes $B(K_S^0 \rightarrow \pi^+ \pi^-) = (69.20 \pm 0.05)\%$ and $B(f_2'(1525) \rightarrow K\bar{K}) = 4 B(f_2'(1525) \rightarrow K_S^0 K_S^0)$.

³ Assuming $B(f_2'(1525) \rightarrow K\bar{K}) = 0.71$.

NODE=M049R52;LINKAGE=A
NODE=M049R52;LINKAGE=BE

NODE=M049R52;LINKAGE=D

$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{95}/Γ
10.1 ± 0.6				OUR AVERAGE	
10.15 ± 0.59 ^{+0.54} _{-0.43}		1 LEES	18A BABR	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
10.5 ± 1.6 ^{+1.9} _{-1.8}		2 BESSON	07A CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$	
10.2 ± 0.8 ± 0.7		ATHAR	06 CLE3	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
8.1 ± 2.3 ^{+2.9} _{-2.7}		3 ANASTASSOV	99 CLE2	$e^+ e^- \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21	90	3 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<13	90	3 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<81	90	SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	
¹ Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = 1/3 B(f_2(1270) \rightarrow \pi \pi)$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.2^{+2.9}_{-0.9})\%$.					
² Using $B(f_2(1270) \rightarrow \pi^0 \pi^0) = B(f_2(1270) \rightarrow \pi \pi)/3$ and $B(f_2(1270) \rightarrow \pi \pi) = (84.7^{+2.5}_{-1.2})\%$.					
³ Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$.					

NODE=M049R51
NODE=M049R51

NODE=M049R51;LINKAGE=A

NODE=M049R51;LINKAGE=BE

NODE=M049R51;LINKAGE=C

 $\Gamma(\gamma \eta(1405))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{96}/Γ
<8.2	90	1 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^\pm \pi^\mp K_S^0$	

¹ Includes unknown branching ratio of $\eta(1405) \rightarrow K^\pm \pi^\mp K_S^0$.

NODE=M049R23
NODE=M049R23

NODE=M049R23;LINKAGE=J

 $\Gamma(\gamma f_0(1500))/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{97}/Γ
<1.5	90	1 BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.1	90	2 BESSON	07A CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$	
¹ Using $B(f_0(1500) \rightarrow \pi^0 \pi^0) = B(f_0(1500) \rightarrow \pi \pi)/3$ and $B(f_0(1500) \rightarrow \pi \pi) = (0.349 \pm 0.023)\%$.					
² Calculated by us using $B(f_0(1500) \rightarrow \eta \eta) = (5.1 \pm 0.9)\%$.					

NODE=M049R44
NODE=M049R44

OCCUR=2

NODE=M049R44;LINKAGE=BE

NODE=M049R44;LINKAGE=BS

 $\Gamma(\gamma f_0(1500) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT	Γ_{98}/Γ
1.04 ± 0.14 ± 0.33	1 LEES	18A BABR	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	

¹ LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1500) \rightarrow \gamma K \bar{K}) = (2.08 \pm 0.27 \pm 0.65) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$.

NODE=M049P29
NODE=M049P29

NODE=M049P29;LINKAGE=A

 $\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}}$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
< 2.6	90	1 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 6.3	90	1 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$	
<19	90	1 FULTON	90B CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$	
< 8	90	2 ALBRECHT	89 ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$	
<24	90	3 SCHMITT	88 CBAL	$\Upsilon(1S) \rightarrow \gamma X$	

¹ Assuming $B(f_0(1710) \rightarrow K \bar{K}) = 0.38$.

² Assuming $B(f_0(1710) \rightarrow \pi \pi) = 0.04$.

³ Assuming $B(f_0(1710) \rightarrow \eta \eta) = 0.18$.

NODE=M049R53
NODE=M049R53

OCCUR=2

OCCUR=2

NODE=M049R53;LINKAGE=E

NODE=M049R53;LINKAGE=F

NODE=M049R53;LINKAGE=A

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma K^+ K^-)/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ
1.01 ± 0.26 ± 0.18		1 LEES	18A BABR	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.7	90	ATHAR	06 CLEO	$e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma K^+ K^-$	
¹ LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma K \bar{K}) = (2.02 \pm 0.51 \pm 0.35) \times 10^{-5}$ assuming $B(K^0 \bar{K}^0) = 1/2 B(K \bar{K})$.					

NODE=M049R50
NODE=M049R50

NODE=M049R50;LINKAGE=A

$\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{101} / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$0.53 \pm 0.17 \pm 0.11$	¹ LEES	18A	BABR $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
¹ LEES 18A quotes $B(\Upsilon(1S) \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi \pi) = (0.79 \pm 0.26 \pm 0.17) \times 10^{-5}$ assuming $B(\pi^0 \pi^0) = 1/3 B(\pi \pi)$.			

NODE=M049P28
NODE=M049P28

NODE=M049P28;LINKAGE=A

 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{102} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.4	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \pi^0 \pi^0$

NODE=M049R45
NODE=M049R45 $\Gamma(\gamma f_0(1710) \rightarrow \gamma \eta \eta) / \Gamma_{\text{total}}$ Γ_{103} / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.8	90	BESSON	07A	CLEO $e^+ e^- \rightarrow \Upsilon(1S) \rightarrow \gamma \eta \eta$

NODE=M049R46
NODE=M049R46 $\Gamma(\gamma f_4(2050)) / \Gamma_{\text{total}}$ Γ_{104} / Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.3	90	¹ ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
¹ Assuming $B(f_4(2050) \rightarrow \pi \pi) = 0.17$.				

NODE=M049R30
NODE=M049R30

NODE=M049R30;LINKAGE=AT

 $\Gamma(\gamma f_0(2200) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{105} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0002	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R63
NODE=M049R63 $\Gamma(\gamma f_J(2220) \rightarrow \gamma K^+ K^-) / \Gamma_{\text{total}}$ Γ_{106} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 8	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 150	90	FULTON	90B	CLEO $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 290	90	ALBRECHT	89	ARG $\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 2000	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^-$

NODE=M049R56
NODE=M049R56 $\Gamma(\gamma f_J(2220) \rightarrow \gamma \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{107} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 6	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 120	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M049R41
NODE=M049R41 $\Gamma(\gamma f_J(2220) \rightarrow \gamma p \bar{p}) / \Gamma_{\text{total}}$ Γ_{108} / Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
< 11	90	ATHAR	06	CLE3 $\Upsilon(1S) \rightarrow \gamma p \bar{p}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 160	90	MASEK	02	CLEO $\Upsilon(1S) \rightarrow \gamma p \bar{p}$

NODE=M049R42
NODE=M049R42 $\Gamma(\gamma \eta(2225) \rightarrow \gamma \phi \phi) / \Gamma_{\text{total}}$ Γ_{109} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.003	90	BARU	89	MD1 $\Upsilon(1S) \rightarrow \gamma K^+ K^- K^+ K^-$

NODE=M049R62
NODE=M049R62 $\Gamma(\gamma \eta_c(1S)) / \Gamma_{\text{total}}$ Γ_{110} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.9 \times 10^{-5}$	90	¹ KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 5.7 \times 10^{-5}$	90	SHEN	10A	BELL $\Upsilon(1S) \rightarrow \gamma X$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

NODE=M049R38
NODE=M049R38

NODE=M049R38;LINKAGE=A

 $\Gamma(\gamma \eta_c(2S)) / \Gamma_{\text{total}}$ Γ_{111} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-4}$	90	¹ KATRENKO	20	BELL $e^+ e^- \rightarrow \gamma + \text{hadrons}$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^-$ decays.				

NODE=M049P30
NODE=M049P30

NODE=M049P30;LINKAGE=A

$\Gamma(\gamma\chi_{c0})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.6 \times 10^{-5}$	90	¹ KATRENKO 20	BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.5 \times 10^{-4}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ decays.				

 Γ_{112}/Γ NODE=M049R39
NODE=M049R39

NODE=M049R39;LINKAGE=A

 $\Gamma(\gamma\chi_{c1})/\Gamma_{\text{total}}$

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7^{+2.4+0.4}_{-1.8-0.5}$		5	¹ KATRENKO 20	BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<2.3	90		SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ decays.					

 Γ_{113}/Γ NODE=M049R40
NODE=M049R40

NODE=M049R40;LINKAGE=A

 $\Gamma(\gamma\chi_{c2})/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.6 \times 10^{-6}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<3.3 \times 10^{-5}$	90	¹ KATRENKO 20	BELL	$\Upsilon(1S) \rightarrow \gamma + \text{hadrons}$
¹ Using $\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-$ decays.				

 Γ_{114}/Γ NODE=M049R48
NODE=M049R48

NODE=M049R48;LINKAGE=A

 $\Gamma(\gamma\chi_{c1}(3872))/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5 \times 10^{-5}$	90	¹ SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$
¹ SHEN 10A reports $[\Gamma(\Upsilon(1S) \rightarrow \gamma\chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S))] < 1.6 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^-J/\psi(1S)) = 3.5 \times 10^{-2}$.				

 Γ_{115}/Γ NODE=M049P32
NODE=M049P32

NODE=M049P32;LINKAGE=A

 $\Gamma(\gamma\chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+\pi^-\pi^0 J/\psi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.8 \times 10^{-6}$	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 Γ_{116}/Γ NODE=M049R68
NODE=M049R68 $\Gamma(\gamma\chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 Γ_{117}/Γ NODE=M049R69
NODE=M049R69 $\Gamma(\gamma\chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	SHEN 10A	BELL	$\Upsilon(1S) \rightarrow \gamma X$

 Γ_{118}/Γ NODE=M049R74
NODE=M049R74 $\Gamma(\gamma X \bar{X} (m_X < 3.1 \text{ GeV}))/\Gamma_{\text{total}}$ $(X \bar{X} = \text{vectors with } m < 3.1 \text{ GeV})$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1	90	¹ BALEST 95	CLEO	$e^+e^- \rightarrow \gamma + X \bar{X}$

 Γ_{119}/Γ NODE=M049R61
NODE=M049R61
NODE=M049R61

NODE=M049R61;LINKAGE=A

 $\Gamma(\gamma X \bar{X} (m_X < 4.5 \text{ GeV}))/\Gamma_{\text{total}}$ $X \text{ and } \bar{X} = \text{zero spin with } m < 4.5 \text{ GeV}$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	¹ DEL-AMO-SA..11J	BABR	$e^+e^- \rightarrow \gamma + X \bar{X}$

 Γ_{120}/Γ NODE=M049R01
NODE=M049R01
NODE=M049R01

NODE=M049R01;LINKAGE=DA

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ $(1.5 \text{ GeV} < m_X < 5.0 \text{ GeV})$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.78	95	ROSNER 07A	CLEO	$e^+e^- \rightarrow \gamma X$

 Γ_{121}/Γ NODE=M049R64
NODE=M049R64
NODE=M049R64

$\Gamma(\gamma A^0)/\Gamma_{\text{total}}$ $(A^0 = \text{scalar with } m < 8.0 \text{ GeV})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

••• We do not use the following data for averages, fits, limits, etc. •••

$< 4.5 \times 10^{-6}$	90	¹ DEL-AMO-SA...11J	BABR	$e^+e^- \rightarrow \gamma + X$
$< 3 \times 10^{-5}$	90	² BALEST	95 CLEO	$e^+e^- \rightarrow \gamma + X$
$< 5.6 \times 10^{-5}$	90	² ANTREASYAN 90c	CBAL	$e^+e^- \rightarrow \gamma + X$

¹ For a non-interacting scalar or pseudoscalar, A^0 , with mass $m_{A^0} < 8.0 \text{ GeV}$. 90% CL upper limits range from 1.9×10^{-6} to 4.5×10^{-6} .

² For any non-interacting long-lived particle with mass $< 7.2 \text{ GeV}$.

 Γ_{122}/Γ

NODE=M049R60

NODE=M049R60

NODE=M049R60

NODE=M049R60;LINKAGE=DA

NODE=M049R60;LINKAGE=A

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ $(201 < M(\mu^+ \mu^-) < 3565 \text{ MeV})$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 9	90	¹ LOVE	08 CLEO	$e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$
-------	----	-------------------	---------	--

••• We do not use the following data for averages, fits, limits, etc. •••

< 16	90	² JIA	22 BELL	$\Upsilon(2S) \rightarrow \gamma \mu^+ \mu^- \pi^+ \pi^-$
< 9.7	90	³ LEES	13C BABR	$e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with $201 < M(\mu^+ \mu^-) < 3565 \text{ MeV}$, excluding J/ψ . Measured 90% CL limits as a function of $M(\mu^+ \mu^-)$ range from $1-9 \times 10^{-6}$.

² For a narrow scalar or pseudoscalar, A^0 , with $0.22 < M(A^0) < 9.2 \text{ GeV}$, resulting in 90% CL upper limits ranging from 3.1×10^{-7} at $M(A^0) = 0.22 \text{ GeV}$ to 1.6×10^{-5} at $M(A^0) = 9.2 \text{ GeV}$.

³ For a narrow scalar or pseudoscalar, A^0 , with mass in the range $0.212-9.2 \text{ GeV}$, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} are in the range $0.28-9.7 \times 10^{-6}$.

 Γ_{123}/Γ

NODE=M049R65

NODE=M049R65

NODE=M049R65

NODE=M049R65;LINKAGE=LO

NODE=M049R65;LINKAGE=A

NODE=M049R65;LINKAGE=LE

 $\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}}$ $(2m_\tau < M(\tau^+ \tau^-) < 9.2 \text{ GeV})$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 130	90	¹ LEES	13R BABR	$\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
---------	----	-------------------	----------	---

••• We do not use the following data for averages, fits, limits, etc. •••

< 150	90	² JIA	22 BELL	$\Upsilon(2S) \rightarrow \gamma \tau^+ \tau^- \pi^+ \pi^-$
< 50	90	³ LOVE	08 CLEO	$e^+e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with $2m_\tau < M(A^0) < 9.2 \text{ GeV}$, resulting in 90% CL upper limits of 0.9×10^{-5} at $M(A^0) = 2m_\tau$, $\approx 1.5 \times 10^{-5}$ at $M(A^0) = 7.5 \text{ GeV}$, and 13×10^{-5} at $M(A^0) = 9.2 \text{ GeV}$.

² For a narrow scalar or pseudoscalar, A^0 , with $2m_\tau < M(A^0) < 9.2 \text{ GeV}$, resulting in 90% CL upper limits ranging from 3.8×10^{-6} at $M(A^0) = 2m_\tau$ to 1.5×10^{-4} at $M(A^0) = 9.2 \text{ GeV}$.

³ For a narrow scalar or pseudoscalar, A^0 , with $2m_\tau < M(A^0) < 7.5 \text{ GeV}$, resulting in 90% CL limits ranging from 1×10^{-5} at $M(A^0) = 2m_\tau$ to 5×10^{-5} at $M(A^0) = 7.5 \text{ GeV}$.

 Γ_{124}/Γ

NODE=M049R66

NODE=M049R66

NODE=M049R66

NODE=M049R66;LINKAGE=A

NODE=M049R66;LINKAGE=B

NODE=M049R66;LINKAGE=LO

 $\Gamma(\gamma A^0 \rightarrow \gamma g g)/\Gamma_{\text{total}}$ $(0.5 \text{ GeV} < m < 9.0 \text{ GeV})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 1 \times 10^{-2}$	90	¹ LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$
----------------------	----	-------------------	----------	-------------------------------------

¹ For a narrow, CP -odd pseudoscalar, A^0 , searched for in 26 hadronic decay modes with invariant mass $0.5 \text{ GeV} < m_{A^0} < 9.0 \text{ GeV}$. Measured 90% CL limits as a function of m_{A^0} range from 10^{-6} to 10^{-2} .

 Γ_{125}/Γ

NODE=M049R03

NODE=M049R03

NODE=M049R03

NODE=M049R03;LINKAGE=A

 $\Gamma(\gamma A^0 \rightarrow \gamma s \bar{s})/\Gamma_{\text{total}}$ $(0.5 \text{ GeV} < m < 9.0 \text{ GeV})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

$< 1 \times 10^{-3}$	90	¹ LEES	13L BABR	$\Upsilon(1S) \rightarrow \gamma X$
----------------------	----	-------------------	----------	-------------------------------------

¹ For a narrow, CP -odd pseudoscalar, A^0 , searched for in 14 hadronic decay modes with invariant mass $1.5 \text{ GeV} < m_{A^0} < 9.0 \text{ GeV}$. Measured 90% CL limits as a function of m_{A^0} range from 10^{-5} to 10^{-3} .

 Γ_{126}/Γ

NODE=M049R04

NODE=M049R04

NODE=M049R04

NODE=M049R04;LINKAGE=A

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

NODE=M049230

 $\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{127}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

< 3.9	90	PATRA	22 BELL	$\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \mu^\mp$
---------	----	-------	---------	--

NODE=M049P33

NODE=M049P33

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{128}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.7 \times 10^{-6}$	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \mu^\pm \tau^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.0 \times 10^{-6}$	95	LOVE	08A	CLEO $e^+ e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M049R67
 NODE=M049R67

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{129}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- e^\pm \tau^\mp$

NODE=M049P34
 NODE=M049P34

 $\Gamma(\gamma e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{130}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<4.2	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \mu^\mp$

NODE=M049P35
 NODE=M049P35

 $\Gamma(\gamma \mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{131}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.1	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma \mu^\pm \tau^\mp$

NODE=M049P36
 NODE=M049P36

 $\Gamma(\gamma e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<6.5	90	PATRA	22	BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \gamma e^\pm \tau^\mp$

NODE=M049P37
 NODE=M049P37

OTHER DECAYS

 $\Gamma(\text{invisible})/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3.0	90	AUBERT	09AX	BABR $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<39	90	RUBIN	07	CLEO $\Upsilon(2S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$
<25	90	TAJIMA	07	BELL $\Upsilon(3S) \rightarrow \pi^+ \pi^- \Upsilon(1S)$

NODE=M049235

NODE=M049R10
 NODE=M049R10

 $\Upsilon(1S)$ REFERENCES

SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVO)	REFID=62012
JIA	22	PRL 128 081804	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=61657
PATRA	22	JHEP 2205 095	S. Patra <i>et al.</i>	(BELLE Collab.)	REFID=61653
KATRENKO	20	PRL 124 122001	P. Katrenko <i>et al.</i>	(BELLE Collab.)	REFID=60544
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58949
LEES	18A	PR D97 112006	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=58950
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=57635
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
SHEN	16	PR D93 112013	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=57515
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345
LEES	13C	PR D87 031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54949
LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55167
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55451
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>		REFID=54746
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314
BESSION	11	PR D83 037101	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=16737
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16495
AUBERT	10C	PR D81 011102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53211
DEL-AMO-SA...	10C	PRL 104 191801	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=53354
SHEN	10A	PR D82 051504	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=53545
AUBERT	09AX	PRL 103 251801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53201
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52565
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
ATHAR	07A	PR D76 072003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=51945
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
BESSION	07A	PR D75 072001	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51638
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
RUBIN	07	PR D75 031104	P. Rubin <i>et al.</i>	(CLEO Collab.)	REFID=51629
TAJIMA	07	PRL 98 132001	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=51645
AQUINES	06A	PR D74 092006	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51510
ATHAR	06	PR D73 032001	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=50993
BESSION	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
BRIERE	04	PR D70 072001	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=50183
ARTUSO	03	PR D67 052003	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=49395
MASEK	02	PR D65 072002	G. Masek <i>et al.</i>	(CLEO Collab.)	REFID=48846
RICHICHI	01B	PRL 87 141801	S.J. Richichi <i>et al.</i>	(CLEO Collab.)	REFID=48345
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424

NODE=M049

ANASTASSOV 99	PRL 82 286	A. Anastassov <i>et al.</i>	(CLEO Collab.)	REFID=46609
ALEXANDER 98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU 96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
BALEST 95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)	REFID=44146
CINABRO 94B	PL B340 129	D. Cinabro <i>et al.</i>	(CLEO Collab.)	REFID=44102
ALBRECHT 92J	ZPHY C55 25	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=42167
BARU 92	ZPHY C54 229	S.E. Baru <i>et al.</i>	(NOVO)	REFID=41860
BARU 92B	ZPHY C56 547	S.E. Baru <i>et al.</i>	(NOVO)	REFID=42168
KOBEL 92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
ANTREASYAN 90C	PL B251 204	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=41455
BLINOV 90	PL B245 311	A.E. Blinov <i>et al.</i>	(NOVO)	REFID=41361
FULTON 90B	PR D41 1401	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=41012
MASCHMANN 90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT 89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
ALEXANDER 89	NP B320 45	J.P. Alexander <i>et al.</i>	(LBL, MICH, SLAC)	REFID=40345
BARU 89	ZPHY C42 505	S.E. Baru <i>et al.</i>	(NOVO)	REFID=40917
CHEN 89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
FULTON 89	PL B224 445	R. Fulton <i>et al.</i>	(CLEO Collab.)	REFID=40918
KAARSBERG 89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUELLER... 88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
JAKUBOWSKI 88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.)	REFID=40742
SCHMITT 88	ZPHY C40 199	P. Schmitt <i>et al.</i>	(Crystal Ball Collab.)	REFID=40582
ALBRECHT 87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN 87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU 86	ZPHY C30 551	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22284
ALBRECHT 85C	PL 154B 452	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22282
KURAEV 85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.				
ARTAMONOV 84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BESSON 84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
GILES 84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
MACKAY 84	PR D29 2483	W.W. MacKay <i>et al.</i>	(CUSB Collab.)	REFID=22281
ANDREWS 83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)	REFID=22273
GILES 83	PRL 50 877	R. Giles <i>et al.</i>	(HARV, OSU, ROCH, RUTG+)	REFID=22274
NICZYPORUK 83	ZPHY C17 197	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=12488
ALBRECHT 82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
ARTAMONOV 82	PL 118B 225	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22271
NICZYPORUK 82	ZPHY C15 299	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22272
BERGER 80C	PL 93B 497	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22263
BOCK 80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264
BERGER 79	ZPHY C1 343	C. Berger <i>et al.</i>	(PLUTO Collab.)	REFID=22259

NODE=M076

$$\chi_{b0}(1P)$$

$$I^G(J^{PC}) = 0^+(0^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M076

$\chi_{b0}(1P)$ MASS

NODE=M076M

VALUE (MeV) DOCUMENT ID
9859.44 ± 0.42 ± 0.31 OUR EVALUATION From average γ energy below, using $\Upsilon(2S)$
 mass = 10023.26 ± 0.31 MeV

NODE=M076M
 → UNCHECKED ←

$m_{\chi_{b1}(1P)} - m_{\chi_{b0}(1P)}$

NODE=M076M2
 NODE=M076M2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
32.49 ± 0.93	LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma\gamma\mu^+\mu^-$

γ ENERGY IN $\Upsilon(2S)$ DECAY

NODE=M076DM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
162.5 ± 0.4 OUR AVERAGE			
162.56 ± 0.19 ± 0.42	ARTUSO	05 CLEO	$\Upsilon(2S) \rightarrow \gamma X$
162.0 ± 0.8 ± 1.2	EDWARDS	99 CLE2	$\Upsilon(2S) \rightarrow \gamma\chi(1P)$
162.1 ± 0.5 ± 1.4	ALBRECHT	85E ARG	$\Upsilon(2S) \rightarrow \text{conv.}\gamma X$
163.8 ± 1.6 ± 2.7	NERNST	85 CBAL	$\Upsilon(2S) \rightarrow \gamma X$
158.0 ± 7 ± 1	HAAS	84 CLEO	$\Upsilon(2S) \rightarrow \text{conv.}\gamma X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
149.4 ± 0.7 ± 5.0	KLOPFEN...	83 CUSB	$\Upsilon(2S) \rightarrow \gamma X$

NODE=M076DM

$\chi_{b0}(1P)$ DECAY MODES

NODE=M076215;NODE=M076

Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1 $\gamma \Upsilon(1S)$	(1.94±0.27) %		DESIG=1
Γ_2 $D^0 X$	< 10.4 %	90%	DESIG=2
Γ_3 $\pi^+ \pi^- K^+ K^- \pi^0$	< 1.6 $\times 10^{-4}$	90%	DESIG=3
Γ_4 $2\pi^+ \pi^- K^- K_S^0$	< 5 $\times 10^{-5}$	90%	DESIG=4
Γ_5 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 5 $\times 10^{-4}$	90%	DESIG=5
Γ_6 $2\pi^+ 2\pi^- 2\pi^0$	< 2.1 $\times 10^{-4}$	90%	DESIG=6
Γ_7 $2\pi^+ 2\pi^- K^+ K^-$	(1.1 ±0.6) $\times 10^{-4}$		DESIG=7
Γ_8 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.7 $\times 10^{-4}$	90%	DESIG=8
Γ_9 $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 5 $\times 10^{-4}$	90%	DESIG=9
Γ_{10} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 1.6 $\times 10^{-4}$	90%	DESIG=10
Γ_{11} $3\pi^+ 3\pi^-$	< 8 $\times 10^{-5}$	90%	DESIG=11
Γ_{12} $3\pi^+ 3\pi^- 2\pi^0$	< 6 $\times 10^{-4}$	90%	DESIG=12
Γ_{13} $3\pi^+ 3\pi^- K^+ K^-$	(2.4 ±1.2) $\times 10^{-4}$		DESIG=13
Γ_{14} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 1.0 $\times 10^{-3}$	90%	DESIG=14
Γ_{15} $4\pi^+ 4\pi^-$	< 8 $\times 10^{-5}$	90%	DESIG=15
Γ_{16} $4\pi^+ 4\pi^- 2\pi^0$	< 2.1 $\times 10^{-3}$	90%	DESIG=16
Γ_{17} $J/\psi J/\psi$	< 7 $\times 10^{-5}$	90%	DESIG=17
Γ_{18} $J/\psi \psi(2S)$	< 1.2 $\times 10^{-4}$	90%	DESIG=18
Γ_{19} $\psi(2S) \psi(2S)$	< 3.1 $\times 10^{-5}$	90%	DESIG=19
Γ_{20} $J/\psi(1S)$ anything	< 2.3 $\times 10^{-3}$	90%	DESIG=20

 $\chi_{b0}(1P)$ BRANCHING RATIOS

NODE=M076220

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M076R1
NODE=M076R1

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.94 ± 0.27 OUR AVERAGE					
2.07 ± 0.24 ± 0.21			^{1,2} LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
1.76 ± 0.30 ± 0.18		87	^{3,4} KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 4.6	90		⁵ LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$
< 6	90		WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
< 11	90		PAUSS	83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ LEES 14M quotes $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (7.75 \pm 0.91) \times 10^{-4}$ combining the results from samples of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and without converted photons. Assumes $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

NODE=M076R1;LINKAGE=A

² LEES 14M reports $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (7.75 \pm 0.91) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M076R1;LINKAGE=B

³ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

NODE=M076R1;LINKAGE=KA
NODE=M076R1;LINKAGE=KR

⁴ KORNICER 11 reports $[\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (6.59 \pm 0.96 \pm 0.60) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6^{+3.7}_{-2.6}) \times 10^{-4}$.

NODE=M076R1;LINKAGE=LE

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M076R01
NODE=M076R01

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<10.4 $\times 10^{-2}$	90	^{6,7} BRIERE	08	CLEO $\Upsilon(2S) \rightarrow \gamma D^0 X$

⁶ For $p_{D^0} > 2.5$ GeV/c.

NODE=M076R01;LINKAGE=BR
NODE=M076R01;LINKAGE=RI

⁷ The authors also present their result as $(5.6 \pm 3.6 \pm 0.5) \times 10^{-2}$.

 $\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M076R02
NODE=M076R02

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	⁸ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

⁸ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M076R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.5	90	⁹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$
⁹ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R03
NODE=M076R03

NODE=M076R03;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹⁰ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$
¹⁰ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 18 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R04
NODE=M076R04

NODE=M076R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	¹¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$
¹¹ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 8 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R05
NODE=M076R05

NODE=M076R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.1 \pm 0.6 \pm 0.1$	7	¹² ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$
¹² ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] = (4 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M076R06
NODE=M076R06

NODE=M076R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	¹³ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
¹³ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 10 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R07
NODE=M076R07

NODE=M076R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹⁴ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
¹⁴ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 20 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R08
NODE=M076R08

NODE=M076R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	¹⁵ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
¹⁵ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R09
NODE=M076R09

NODE=M076R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	¹⁶ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
¹⁶ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R10
NODE=M076R10

NODE=M076R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	¹⁷ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
¹⁷ ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 22 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.				

NODE=M076R11
NODE=M076R11

NODE=M076R11;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 1.2 \pm 0.2$	9	18 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
<p>18 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ = $(9 \pm 4 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.</p>				

NODE=M076R12
NODE=M076R12

NODE=M076R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<10	90	19 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
<p>19 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ < 37×10^{-6} which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R13
NODE=M076R13

NODE=M076R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	20 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
<p>20 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ < 3×10^{-6} which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R14
NODE=M076R14

NODE=M076R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<21	90	21 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
<p>21 ASNER 08A reports $[\Gamma(\chi_{b0}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ < 77×10^{-6} which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.</p>				

NODE=M076R15
NODE=M076R15

NODE=M076R15;LINKAGE=AS

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	22 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
<p>22 SHEN 12 reports < 7.1×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.</p>				

NODE=M076R16
NODE=M076R16

NODE=M076R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	23 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
<p>23 SHEN 12 reports < 12×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.</p>				

NODE=M076R17
NODE=M076R17

NODE=M076R17;LINKAGE=SH

 $\Gamma(\psi(2S) \psi(2S))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	24 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
<p>24 SHEN 12 reports < 3.1×10^{-5} from a measurement of $[\Gamma(\chi_{b0}(1P) \rightarrow \psi(2S) \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4) \times 10^{-2}$.</p>				

NODE=M076R18
NODE=M076R18

NODE=M076R18;LINKAGE=SH

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 2.3×10^{-3}	90	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M076R00
NODE=M076R00 $\chi_{b0}(1P)$ CROSS-PARTICLE BRANCHING RATIOS

NODE=M076230

$$\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_{73}^{\Upsilon(2S)}/\Gamma_{\Upsilon(2S)}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.7×10^{-3}	90	25 LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$
<p>25 LEES 11J quotes a central value of $\Gamma(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}} = (8.3 \pm 5.6_{-2.6}^{+3.7}) \times 10^{-4}$ and derives a 90% CL upper limit of $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}} < 4.6\%$ using $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = (3.8 \pm 0.4)\%$.</p>				

NODE=M076B02
NODE=M076B02

NODE=M076B02;LINKAGE=LE

$B(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

1.67±0.28 OUR AVERAGE

2.9	+1.7 +0.1 -1.4 -0.8	26	LEES	14M BABR $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
-----	------------------------	----	------	---

1.63±0.24±0.15	87	KORNICER	11	CLEO $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
----------------	----	----------	----	--

²⁶ From a sample of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.

NODE=M076B01
NODE=M076B01

NODE=M076B01;LINKAGE=A

 $[B(\chi_{b0}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
-----------	-------------	------	---------

3.28±0.37

27	LEES	14M	BABR $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
----	------	-----	---

²⁷ From a sample of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ without converted photons.

NODE=M076A01
NODE=M076A01

NODE=M076A01;LINKAGE=A

 $\chi_{b0}(1P)$ REFERENCES

JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54313
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)	REFID=22290
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)	REFID=22286

NODE=M076

$\chi_{b1}(1P)$

$$J^G(J^{PC}) = 0^+(1^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 1$ from SKWARNICKI 87.

NODE=M077

NODE=M077

 $\chi_{b1}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID
9892.78±0.26±0.31 OUR EVALUATION	From average γ energy below, using $\Upsilon(2S)$
mass = 10023.26 ± 0.31 MeV	

NODE=M077M


NODE=M077M
→ UNCHECKED ←

 γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
129.63±0.33 OUR AVERAGE	Error includes scale factor of 1.3.		See the ideogram below.
129.58±0.09±0.29	ARTUSO	05	CLEO $\Upsilon(2S) \rightarrow \gamma X$
128.8 ±0.4 ±0.6	EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
131.7 ±0.9 ±1.3	WALK	86	CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
131.7 ±0.3 ±1.1	ALBRECHT	85E	ARG $\Upsilon(2S) \rightarrow \text{conv. } \gamma X$
130.6 ±0.8 ±2.4	NERNST	85	CBAL $\Upsilon(2S) \rightarrow \gamma X$
129 ±0.8 ±1	HAAS	84	CLEO $\Upsilon(2S) \rightarrow \text{conv. } \gamma X$
128.1 ±0.4 ±3.0	KLOPFEN...	83	CUSB $\Upsilon(2S) \rightarrow \gamma X$
130.6 ±3.0	PAUSS	83	CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M077DM

NODE=M077DM



chi_b1(1P) DECAY MODES

NODE=M077215;NODE=M077

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\gamma \Upsilon(1S)$	(35.2 ± 2.0) %	
Γ_2 $D^0 X$	(12.6 ± 2.2) %	
Γ_3 $\pi^+ \pi^- K^+ K^- \pi^0$	(2.0 ± 0.6) × 10 ⁻⁴	
Γ_4 $2\pi^+ \pi^- K^- K_S^0$	(1.3 ± 0.5) × 10 ⁻⁴	
Γ_5 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 6 × 10 ⁻⁴	90%
Γ_6 $2\pi^+ 2\pi^- 2\pi^0$	(8.0 ± 2.5) × 10 ⁻⁴	
Γ_7 $2\pi^+ 2\pi^- K^+ K^-$	(1.5 ± 0.5) × 10 ⁻⁴	
Γ_8 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	(3.5 ± 1.2) × 10 ⁻⁴	
Γ_9 $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(8.6 ± 3.2) × 10 ⁻⁴	
Γ_{10} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	(9.3 ± 3.3) × 10 ⁻⁴	
Γ_{11} $3\pi^+ 3\pi^-$	(1.9 ± 0.6) × 10 ⁻⁴	
Γ_{12} $3\pi^+ 3\pi^- 2\pi^0$	(1.7 ± 0.5) × 10 ⁻³	
Γ_{13} $3\pi^+ 3\pi^- K^+ K^-$	(2.6 ± 0.8) × 10 ⁻⁴	
Γ_{14} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	(7.5 ± 2.6) × 10 ⁻⁴	
Γ_{15} $4\pi^+ 4\pi^-$	(2.6 ± 0.9) × 10 ⁻⁴	
Γ_{16} $4\pi^+ 4\pi^- 2\pi^0$	(1.4 ± 0.6) × 10 ⁻³	
Γ_{17} ω anything	(4.9 ± 1.4) %	
Γ_{18} ωX_{tetra}	< 4.44 × 10 ⁻⁴	90%
Γ_{19} $J/\psi J/\psi$	< 2.7 × 10 ⁻⁵	90%
Γ_{20} $J/\psi \psi(2S)$	< 1.7 × 10 ⁻⁵	90%
Γ_{21} $\psi(2S) \psi(2S)$	< 6 × 10 ⁻⁵	90%
Γ_{22} $J/\psi(1S)$ anything	< 1.1 × 10 ⁻³	90%
Γ_{23} $J/\psi(1S) X_{tetra}$	< 2.27 × 10 ⁻⁴	90%

DESIG=1
DESIG=2
DESIG=3
DESIG=4
DESIG=5
DESIG=6
DESIG=7
DESIG=8
DESIG=9
DESIG=10
DESIG=11
DESIG=12
DESIG=13
DESIG=14
DESIG=15
DESIG=16
DESIG=21
DESIG=22
DESIG=17
DESIG=18
DESIG=19
DESIG=20
DESIG=23

chi_b1(1P) BRANCHING RATIOS

NODE=M077220

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{total}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.352 ± 0.020 OUR AVERAGE				
0.356 ^{+0.016} _{-0.022} ± 0.019	964k	¹ FULSOM	18 BELL	$\Upsilon(2S) \rightarrow \gamma X$
0.364 ± 0.017 ± 0.019		^{2,3,4} LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.331 ± 0.018 ± 0.017	3222	^{4,5} KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
0.350 ± 0.023 ± 0.018	13k	⁶ LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$
0.34 ± 0.07 ± 0.02	53	^{4,7,8} WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
0.47 ± 0.18		KLOPFEN...	83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M077R1
NODE=M077R1

OCCUR=4

¹ FULSOM 18 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (2.45 \pm 0.02^{+0.11}_{-0.15}) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=J

² LEES 14M quotes $\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}} = (2.51 \pm 0.12) \%$ combining the results from samples of $\Upsilon(2S) \rightarrow \gamma \mu^+ \mu^-$ with and without converted photons.

NODE=M077R1;LINKAGE=B

³ LEES 14M reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (2.51 \pm 0.12) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=D

⁴ Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

NODE=M077R1;LINKAGE=KA

⁵ KORNICER 11 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (22.8 \pm 0.4 \pm 1.2) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=KR

⁶ LEES 11J reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (24.1 \pm 0.6 \pm 1.5) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=LE

⁷ WALK 86 quotes $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (5.8 \pm 0.9 \pm 0.7) \%$.

NODE=M077R1;LINKAGE=A

⁸ WALK 86 reports $[\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (23.4 \pm 3.63 \pm 2.82) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R1;LINKAGE=C

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
12.6±1.9±1.1	2310	¹ BRIERE	08	CLEO $\Upsilon(2S) \rightarrow \gamma D^0 X$

NODE=M077R01
NODE=M077R01

¹ For $p_{D^0} > 2.5 \text{ GeV}/c$.

NODE=M077R01;LINKAGE=BR

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.0±0.6±0.1	18	¹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

NODE=M077R02
NODE=M077R02

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (14 \pm 3 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.3±0.5±0.1	11	¹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

NODE=M077R03
NODE=M077R03

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (9 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R03;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	¹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

NODE=M077R04
NODE=M077R04

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] < 42 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

NODE=M077R04;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.0±2.4±0.4	46	¹ ASNER	08A	CLEO $\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

NODE=M077R05
NODE=M077R05

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] = (55 \pm 9 \pm 14) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M077R05;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.5±0.5±0.1	18	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(10 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R06
NODE=M077R06

NODE=M077R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±1.2±0.2	22	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(24 \pm 6 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R07
NODE=M077R07

NODE=M077R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6±3.2±0.4	26	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(59 \pm 14 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R08
NODE=M077R08

NODE=M077R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.3±3.3±0.5	21	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(64 \pm 16 \pm 16) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R09
NODE=M077R09

NODE=M077R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.9±0.6±0.1	25	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^-$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(13 \pm 3 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R10
NODE=M077R10

NODE=M077R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
17±5±1	56	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(119 \pm 18 \pm 32) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R11
NODE=M077R11

NODE=M077R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6±0.8±0.1	21	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(18 \pm 4 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R12
NODE=M077R12

NODE=M077R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.5±2.6±0.4	28	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ = $(52 \pm 11 \pm 14) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M077R13
NODE=M077R13

NODE=M077R13;LINKAGE=AS

$\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6±0.9±0.1	24	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$

NODE=M077R14
NODE=M077R14

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (18 \pm 4 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) =$
 $(6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is
the systematic error from using our best value.

NODE=M077R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
14±5±1	26	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$

NODE=M077R15
NODE=M077R15

¹ ASNER 08A reports $[\Gamma(\chi_{b1}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$
 $= (96 \pm 24 \pm 29) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) =$
 $(6.9 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is
the systematic error from using our best value.

NODE=M077R15;LINKAGE=AS

 $\Gamma(\omega \text{ anything})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.9±1.3±0.6	51k	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M077R19
NODE=M077R19 $\Gamma(\omega X_{\text{tetra}})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<44.4 × 10⁻⁵	90	¹ JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M077R23
NODE=M077R23

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the
range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range
from 3.3×10^{-5} to 44.4×10^{-5} .

NODE=M077R23;LINKAGE=A

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<2.7	90	¹ SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$

NODE=M077R16
NODE=M077R16

¹ SHEN 12 reports $< 2.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}]$
 $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times 10^{-2}$.

NODE=M077R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	¹ SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$

NODE=M077R17
NODE=M077R17

¹ SHEN 12 reports $< 1.7 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}]$
 $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times$
 10^{-2} .

NODE=M077R17;LINKAGE=SH

 $\Gamma(\psi(2S) \psi(2S))/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	¹ SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$

NODE=M077R18
NODE=M077R18

¹ SHEN 12 reports $< 6.2 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b1}(1P) \rightarrow \psi(2S) \psi(2S))/\Gamma_{\text{total}}]$
 $\times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = (6.9 \pm 0.4) \times$
 10^{-2} .

NODE=M077R18;LINKAGE=SH

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 × 10⁻³	90	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M077R00
NODE=M077R00 $\Gamma(J/\psi(1S) X_{\text{tetra}})/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<22.7 × 10⁻⁵	90	¹ JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M077R22
NODE=M077R22

¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the
range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range
from 1.8×10^{-5} to 22.7×10^{-5} .

NODE=M077R22;LINKAGE=A

 $\chi_{b1}(1P)$ Cross-Particle Branching Ratios

NODE=M077230

$$\Gamma(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_{71}^{\Upsilon(2S)}/\Gamma \Upsilon(2S)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
24.1±0.6±1.5	13k	LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$

NODE=M077B03
NODE=M077B03


$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.90±0.34 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
6.86 ^{+0.47+0.44} _{-0.45-0.35}		¹ LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
5.65±0.11±0.27	3222	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
5.8 ±0.9 ±0.7	53	WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ From a sample of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.

NODE=M077B01
NODE=M077B01

NODE=M077B01;LINKAGE=A



$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$
(units 10^{-4})

$B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
1.30±0.34 OUR AVERAGE				
1.16 ^{+0.78+0.14} _{-0.67-0.16}		¹ LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
1.33±0.30±0.23	50	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with converted photons.

NODE=M077B02
NODE=M077B02

NODE=M077B02;LINKAGE=A

$B(\chi_{b2}(1P) \rightarrow \rho X + \bar{\rho} X) / B(\chi_{b1}(1P) \rightarrow \rho X + \bar{\rho} X)$

VALUE	DOCUMENT ID	TECN	COMMENT
1.068±0.010±0.040	BRIERE	07 CLEO	$\Upsilon(2S) \rightarrow \gamma \chi_{b,J}(1P)$

NODE=M077R20
NODE=M077R20

$B(\chi_{b0}(1P) \rightarrow \rho X + \bar{\rho} X) / B(\chi_{b1}(1P) \rightarrow \rho X + \bar{\rho} X)$

VALUE	DOCUMENT ID	TECN	COMMENT
1.11±0.15±0.20	BRIERE	07 CLEO	$\Upsilon(2S) \rightarrow \gamma \chi_{b,J}(1P)$

NODE=M077R21
NODE=M077R21

$\chi_{b1}(1P)$ REFERENCES

FULSOM 18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)
JIA 17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)
LEES 14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
SHEN 12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)
KORNICER 11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)
LEES 11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER 08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE 08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE 07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO 05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
EDWARDS 99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)
SKWARNICKI 87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)
WALK 86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)
ALBRECHT 85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
NERNST 85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)
HAAS 84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)
KLOPFEN... 83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)
PAUSS 83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)

NODE=M077

REFID=59535
REFID=58318
REFID=56343
REFID=54313
REFID=16769
REFID=53936
REFID=52574
REFID=52577
REFID=51887
REFID=50454
REFID=46612
REFID=40019
REFID=22290
REFID=22288
REFID=22289
REFID=22287
REFID=22285
REFID=22286

$h_b(1P)$

$$J^{PC} = 0^-(1^{+-})$$

Quantum numbers are quark model predictions, $C = -$ established by $\eta_b\gamma$ decay.

NODE=M204

NODE=M204

 $h_b(1P)$ MASS

NODE=M204M

NODE=M204M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9899.3±0.8 OUR AVERAGE				
9899.3±0.4±1.0	112k	TAMPONI	15	BELL $e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
9899.1±0.4±1.0	70k	MIZUK	12	BELL $e^+e^- \rightarrow \pi^+\pi^-\text{ hadrons}$
9902 ±4 ±2	10.8k	LEES	11k	BABR $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

9898.2 ^{+1.1+1.0} _{-1.0-1.1}	50.0k	¹ ADACHI	12	BELL 10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM
--	-------	---------------------	----	---

¹Superseded by MIZUK 12.

NODE=M204M;LINKAGE=AD

NODE=M204215;NODE=M204

 $h_b(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta_b(1S)\gamma$	(52 ⁺⁶ ₋₅) %

DESIG=1

 $h_b(1P)$ BRANCHING RATIOS

NODE=M204225

$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT

NODE=M204R01
NODE=M204R01**52⁺⁶₋₅ OUR AVERAGE**

56 ±8 ±4	33.1k	¹ TAMPONI	15	BELL $e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
49.2±5.7 ^{+5.6} _{-3.3}	24k	MIZUK	12	BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-\text{ hadrons}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	10.8k	LEES	11k	BABR $\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$
------	-------	------	-----	---

¹Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.

NODE=M204R01;LINKAGE=A

 $h_b(1P)$ REFERENCES

NODE=M204

TAMPONI	15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)
ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)

REFID=56996
REFID=53962
REFID=54718
REFID=53937

$\chi_{b2}(1P)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 2$ from SKWARNICKI 87.

NODE=M078

NODE=M078

 $\chi_{b2}(1P)$ MASS

NODE=M078M

VALUE (MeV) DOCUMENT ID
9912.21 ± 0.26 ± 0.31 OUR EVALUATION From average γ energy below, using $\Upsilon(2S)$
 mass = 10023.26 ± 0.31 MeV

NODE=M078M
 → UNCHECKED ←

 $m_{\chi_{b2}(1P)} - m_{\chi_{b1}(1P)}$

NODE=M078DM2

VALUE (MeV) DOCUMENT ID TECN COMMENT
19.10 ± 0.25 OUR AVERAGE Error includes scale factor of 1.1.
 19.81 ± 0.65 ± 0.20 ¹ AAIJ 14BG LHCB $pp \rightarrow \gamma \mu^+ \mu^- X$
 19.01 ± 0.24 LEES 14M BABR $\Upsilon(2S) \rightarrow \gamma \mu^+ \mu^-$

NODE=M078DM2

¹ From the $\chi_{bj}(1P) \rightarrow \Upsilon(1S)\gamma$ transition.

NODE=M078DM2;LINKAGE=A

 γ ENERGY IN $\Upsilon(2S)$ DECAY

NODE=M078DM

VALUE (MeV) DOCUMENT ID TECN COMMENT
110.44 ± 0.29 OUR AVERAGE Error includes scale factor of 1.1.
 110.58 ± 0.08 ± 0.30 ARTUSO 05 CLEO $\Upsilon(2S) \rightarrow \gamma X$
 110.8 ± 0.3 ± 0.6 EDWARDS 99 CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
 107.0 ± 1.1 ± 1.3 WALK 86 CBAL $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
 110.6 ± 0.3 ± 0.9 ALBRECHT 85E ARG $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
 110.4 ± 0.8 ± 2.2 NERNST 85 CBAL $\Upsilon(2S) \rightarrow \gamma X$
 109.5 ± 0.7 ± 1.0 HAAS 84 CLEO $\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
 108.2 ± 0.3 ± 2.0 KLOPFEN... 83 CUSB $\Upsilon(2S) \rightarrow \gamma X$
 108.8 ± 4.0 PAUSS 83 CUSB $\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M078DM

 $\chi_{b2}(1P)$ DECAY MODES

NODE=M078215;NODE=M078

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\gamma \Upsilon(1S)$	(18.0 ± 1.0) %	
Γ_2 $D^0 X$	< 7.9 %	90%
Γ_3 $\pi^+ \pi^- K^+ K^- \pi^0$	(8 ± 5) × 10 ⁻⁵	
Γ_4 $2\pi^+ \pi^- K^- K_S^0$	< 1.0 × 10 ⁻⁴	90%
Γ_5 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	(5.3 ± 2.4) × 10 ⁻⁴	
Γ_6 $2\pi^+ 2\pi^- 2\pi^0$	(3.5 ± 1.4) × 10 ⁻⁴	
Γ_7 $2\pi^+ 2\pi^- K^+ K^-$	(1.1 ± 0.4) × 10 ⁻⁴	
Γ_8 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.1 ± 0.9) × 10 ⁻⁴	
Γ_9 $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(3.9 ± 1.8) × 10 ⁻⁴	
Γ_{10} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 5 × 10 ⁻⁴	90%
Γ_{11} $3\pi^+ 3\pi^-$	(7.0 ± 3.1) × 10 ⁻⁵	
Γ_{12} $3\pi^+ 3\pi^- 2\pi^0$	(1.0 ± 0.4) × 10 ⁻³	
Γ_{13} $3\pi^+ 3\pi^- K^+ K^-$	< 8 × 10 ⁻⁵	90%
Γ_{14} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	(3.6 ± 1.5) × 10 ⁻⁴	
Γ_{15} $4\pi^+ 4\pi^-$	(8 ± 4) × 10 ⁻⁵	
Γ_{16} $4\pi^+ 4\pi^- 2\pi^0$	(1.8 ± 0.7) × 10 ⁻³	
Γ_{17} $J/\psi J/\psi$	< 4 × 10 ⁻⁵	90%
Γ_{18} $J/\psi \psi(2S)$	< 5 × 10 ⁻⁵	90%
Γ_{19} $\psi(2S) \psi(2S)$	< 1.6 × 10 ⁻⁵	90%
Γ_{20} $J/\psi(1S)$ anything	(1.5 ± 0.4) × 10 ⁻³	

DESIG=1

DESIG=2

DESIG=3

DESIG=4

DESIG=5

DESIG=6

DESIG=7

DESIG=8

DESIG=9

DESIG=10

DESIG=11

DESIG=12

DESIG=13

DESIG=14

DESIG=15

DESIG=16

DESIG=17

DESIG=18

DESIG=19

DESIG=20

$\chi_{b2}(1P)$ BRANCHING RATIOS

NODE=M078220

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT		
0.180 ± 0.010 OUR AVERAGE						
$0.164^{+0.009}_{-0.010} \pm 0.008$	503k	¹ FULSOM	18	BELL	$\Upsilon(2S) \rightarrow \gamma X$	
$0.185 \pm 0.008 \pm 0.009$		^{2,3,4} LEES	14M	BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
$0.186 \pm 0.011 \pm 0.009$	1770	^{4,5} KORNICER	11	CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$	
$0.194^{+0.014}_{-0.017} \pm 0.009$	8k	⁶ LEES	11J	BABR	$\Upsilon(2S) \rightarrow X \gamma$	
$0.25 \pm 0.06 \pm 0.01$	35	^{4,7,8} WALK	86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
0.20 ± 0.05		KLOPFEN...	83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

NODE=M078R1
NODE=M078R1

OCCUR=2

¹ FULSOM 18 reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.17 \pm 0.01^{+0.06}_{-0.07}) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=H

² LEES 14M quotes $\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}} = (1.32 \pm 0.06)\%$ combining the results from samples of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and without converted photons.

NODE=M078R1;LINKAGE=B

³ LEES 14M reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.32 \pm 0.06) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=F

⁴ Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

NODE=M078R1;LINKAGE=KA

⁵ KORNICER 11 reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (1.33 \pm 0.04 \pm 0.07) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=KR

⁶ LEES 11J reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (13.9 \pm 0.5^{+0.9}_{-1.1}) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=LE

⁷ WALK 86 quotes $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (4.4 \pm 0.9 \pm 0.5) \%$.

NODE=M078R1;LINKAGE=A

⁸ WALK 86 reports $[\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (17.7 \pm 3.6 \pm 2.0) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R1;LINKAGE=E

$\Gamma(D^0 X)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$< 7.9 \times 10^{-2}$	90	^{1,2} BRIERE	08	CLEO	$\Upsilon(2S) \rightarrow \gamma D^0 X$	

NODE=M078R01
NODE=M078R01

¹ For $p_{D^0} > 2.5$ GeV/c.

NODE=M078R01;LINKAGE=BR

² The authors also present their result as $(5.4 \pm 1.9 \pm 0.5) \times 10^{-2}$.

NODE=M078R01;LINKAGE=RI

$\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT		
0.84 ± 0.50 ± 0.04	8	¹ ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$	

NODE=M078R02
NODE=M078R02

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 3 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R02;LINKAGE=AS

$\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$						Γ_4/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT		
< 1.0	90	¹ ASNER	08A	CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$	

NODE=M078R03
NODE=M078R03

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 7 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

NODE=M078R03;LINKAGE=AS

$\Gamma(2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.3±2.4±0.3	11	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M078R04
 NODE=M078R04

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+\pi^-K^-K_S^02\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (38 \pm 14 \pm 10) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R04;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.5±1.4±0.2	19	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-2\pi^0$

NODE=M078R05
 NODE=M078R05

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (25 \pm 8 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R05;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.4±0.1	14	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-$

NODE=M078R06
 NODE=M078R06

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (8 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R06;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.9±0.1	13	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-\pi^0$

NODE=M078R07
 NODE=M078R07

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (15 \pm 5 \pm 4) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R07;LINKAGE=AS

 $\Gamma(2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.9±1.8±0.2	11	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 2\pi^+2\pi^-K^+K^-2\pi^0$

NODE=M078R08
 NODE=M078R08

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 2\pi^+2\pi^-K^+K^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (28 \pm 11 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R08;LINKAGE=AS

 $\Gamma(3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+2\pi^-K^-K_S^0\pi^0$

NODE=M078R09
 NODE=M078R09

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+2\pi^-K^-K_S^0\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] < 36 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

NODE=M078R09;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.70±0.31±0.03	9	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-$

NODE=M078R10
 NODE=M078R10

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (5 \pm 2 \pm 1) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R10;LINKAGE=AS

 $\Gamma(3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
10.2±3.6±0.5	34	¹ ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+3\pi^-2\pi^0$

NODE=M078R11
 NODE=M078R11

¹ ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+3\pi^-2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P))] = (73 \pm 16 \pm 20) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma\chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M078R11;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.8	90	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] < 6 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.				

NODE=M078R12
NODE=M078R12

NODE=M078R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.6 \pm 1.5 \pm 0.2$	14	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (26 \pm 8 \pm 7) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R13
NODE=M078R13

NODE=M078R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.84 \pm 0.40 \pm 0.04$	7	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^-$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (6 \pm 2 \pm 2) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R14
NODE=M078R14

NODE=M078R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$18 \pm 7 \pm 1$	29	1 ASNER	08A CLEO	$\Upsilon(2S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
1 ASNER 08A reports $[\Gamma(\chi_{b2}(1P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] = (132 \pm 31 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M078R15
NODE=M078R15

NODE=M078R15;LINKAGE=AS

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	1 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 4.5 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi J/\psi)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078R16
NODE=M078R16

NODE=M078R16;LINKAGE=SH

 $\Gamma(J/\psi \psi(2S))/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<5	90	1 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 4.9 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow J/\psi \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078R17
NODE=M078R17

NODE=M078R17;LINKAGE=SH

 $\Gamma(\psi(2S) \psi(2S))/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	1 SHEN	12 BELL	$\Upsilon(2S) \rightarrow \gamma \psi X$
1 SHEN 12 reports $< 1.6 \times 10^{-5}$ from a measurement of $[\Gamma(\chi_{b2}(1P) \rightarrow \psi(2S) \psi(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$ assuming $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = (7.15 \pm 0.35) \times 10^{-2}$.				

NODE=M078R18
NODE=M078R18

NODE=M078R18;LINKAGE=SH

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.50 \pm 0.34 \pm 0.22$	462	JIA	17A BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M078R00
NODE=M078R00 $\chi_{b2}(1P)$ Cross-Particle Branching Ratios

NODE=M078230

$$\Gamma(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_{72}^{\Upsilon(2S)}/\Gamma \Upsilon(2S)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$13.9 \pm 0.5^{+0.9}_{-1.1}$	8k	LEES	11J BABR	$\Upsilon(2S) \rightarrow X \gamma$

NODE=M078B03
NODE=M078B03

$B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.38 ± 0.16				OUR AVERAGE
$3.63^{+0.36+0.18}_{-0.34-0.19}$		¹ LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$3.29 \pm 0.09 \pm 0.16$	1770	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
$4.4 \pm 0.9 \pm 0.5$	35	WALK	86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ From a sample of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with converted photons.

NODE=M078B01
NODE=M078B01

NODE=M078B01;LINKAGE=A

 $[B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))] / [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))]$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
55.6 ± 1.6	¹ LEES	14M BABR	$\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$

¹ From a sample of $\Upsilon(2S) \rightarrow \gamma \gamma \mu^+ \mu^-$ events without converted photons.

NODE=M078A00
NODE=M078A00

NODE=M078A00;LINKAGE=A

 $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5				OUR AVERAGE
$4.68^{+0.99}_{-0.92} \pm 0.37$		¹ LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$3.56 \pm 0.40 \pm 0.41$	126	KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with converted photons.

NODE=M078B02
NODE=M078B02

NODE=M078B02;LINKAGE=A

 $\chi_{b2}(1P)$ REFERENCES

FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)	REFID=59535
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56199
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
SHEN	12	PR D85 071102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54313
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
SKWARNICKI	87	PRL 58 972	T. Skwarnicki <i>et al.</i>	(Crystal Ball Collab.)	REFID=40019
WALK	86	PR D34 2611	W.S. Walk <i>et al.</i>	(Crystal Ball Collab.)	REFID=22290
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
NERNST	85	PRL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
PAUSS	83	PL 130B 439	F. Pauss <i>et al.</i>	(MPIM, COLU, CORN, LSU+)	REFID=22286

NODE=M078

REFID=59535
REFID=58318
REFID=56199
REFID=56343
REFID=54313
REFID=16769
REFID=53936
REFID=52574
REFID=52577
REFID=50454
REFID=46612
REFID=40019
REFID=22290
REFID=22288
REFID=22289
REFID=22287
REFID=22285
REFID=22286

NODE=M200

$\eta_b(2S)$

$$I^G(J^{PC}) = 0^+(0^-+)$$

OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

NODE=M200

 $\eta_b(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$9999.0 \pm 3.5^{+2.8}_{-1.9}$	26k	¹ MIZUK	12 BELL	$e^+ e^- \rightarrow \gamma \pi^+ \pi^- +$ hadrons

••• We do not use the following data for averages, fits, limits, etc. •••

9974.6 \pm 2.3 \pm 2.1 11 \pm 4 ^{2,3,4} DOBBS 12 $\Upsilon(2S) \rightarrow \gamma$ hadrons

¹ Assuming $\Gamma_{\eta_b(2S)} = 4.9$ MeV. Not independent of the corresponding mass difference measurement.

² SANDILYA 13 (Belle Collab.) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of $(157.8 \pm 3.6) \times 10^6 \Upsilon(2S)$ decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction $B(\Upsilon(2S) \rightarrow \eta_b(2S) \gamma) \times \sum_i B(\eta_b(2S) \rightarrow X_i) < 4.9 \times 10^{-6}$, summed over the exclusive hadronic final states X_i , is an order of magnitude smaller than that reported by DOBBS 12.

³ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

⁴ Assuming $\Gamma_{\eta_b(2S)} = 5$ MeV. Not independent of the corresponding mass difference measurement.

NODE=M200M

NODE=M200M

NODE=M200M;LINKAGE=MI

NODE=M200M;LINKAGE=A

NODE=M200M;LINKAGE=DO

NODE=M200M;LINKAGE=NI

 $m_{\Upsilon(2S)} - m_{\eta_b(2S)}$

NODE=M200DM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$24.3 \pm 3.5^{+2.8}_{-1.9}$	26k	⁵ MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^- +$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$48.7 \pm 2.3 \pm 2.1$	11 ± 4	^{6,7,8} DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
⁵ Assuming $\Gamma_{\eta_b(2S)} = 4.9$ MeV. Not independent of the corresponding mass measurement.				
⁶ SANDILYA 13 (Belle Collab.) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of $(157.8 \pm 3.6) \times 10^6$ $\Upsilon(2S)$ decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction $B(\Upsilon(2S) \rightarrow \eta_b(2S)\gamma) \times \sum_i B(\eta_b(2S) \rightarrow X_i) < 4.9 \times 10^{-6}$, summed over the exclusive hadronic final states X_i , is an order of magnitude smaller than that reported by DOBBS 12.				
⁷ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.				
⁸ Assuming $\Gamma_{\eta_b(2S)} = 5$ MeV. Not independent of the corresponding mass measurement.				

NODE=M200DM

NODE=M200DM;LINKAGE=MI

NODE=M200DM;LINKAGE=A

NODE=M200DM;LINKAGE=DO

NODE=M200DM;LINKAGE=NI

 $\eta_b(2S)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<24	90	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^-$ hadrons

NODE=M200W

NODE=M200W

 $\eta_b(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	seen

NODE=M200215;NODE=M200

DESIG=1

 $\eta_b(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	26k	MIZUK	12	BELL $e^+e^- \rightarrow \gamma\pi^+\pi^-$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	^{9,10}	DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons
⁹ SANDILYA 13 (Belle Collab.) search for such a state reconstructed in the same 26 exclusive hadronic final states as DOBBS 12 using a sample of $(157.8 \pm 3.6) \times 10^6$ $\Upsilon(2S)$ decays or about 17 times larger and find no evidence for a signal. Their 90% C.L. upper limit on the branching fraction $B(\Upsilon(2S) \rightarrow \eta_b(2S)\gamma) \times \sum_i B(\eta_b(2S) \rightarrow X_i) < 4.9 \times 10^{-6}$, summed over the exclusive hadronic final states X_i , is an order of magnitude smaller than that reported by DOBBS 12.				
¹⁰ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.				

NODE=M200225

NODE=M200R01
NODE=M200R01

NODE=M200R01;LINKAGE=A

NODE=M200R01;LINKAGE=DO

 $\eta_b(2S)$ REFERENCES

SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>	
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M200

REFID=55590
REFID=54288
REFID=54718

$\Upsilon(2S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M052

 $\Upsilon(2S)$ MASS

NODE=M052M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10023.4±0.5	¹ SHAMOV	23	RVUE $e^+e^- \rightarrow$ hadrons
••• We do not use the following data for averages, fits, limits, etc. •••			
10022.7±0.4	² SHAMOV	23	RVUE $e^+e^- \rightarrow$ hadrons
10023.5±0.5	^{3,4} ARTAMONOV	00	MD1 $e^+e^- \rightarrow$ hadrons
10023.6±0.5	^{5,6} BARU	86B	MD1 $e^+e^- \rightarrow$ hadrons
10023.1±0.4	⁷ BARBER	84	ARG $e^+e^- \rightarrow$ hadrons

NODE=M052M

OCCUR=2

¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.

NODE=M052M;LINKAGE=A

² Obtained by reanalysing ARGUS and Crystal Ball data (BARBER 84), but not authored by the ARGUS and Crystal Ball collaboration.

NODE=M052M;LINKAGE=B

³ Reanalysis of BARU 86B using new electron mass (COHEN 87).

NODE=M052M;LINKAGE=AR

⁴ Superseded by SHAMOV 23.

NODE=M052M;LINKAGE=E

⁵ Reanalysis of ARTAMONOV 84.

NODE=M052M;LINKAGE=C

⁶ Superseded by ARTAMONOV 00.

NODE=M052M;LINKAGE=RZ

⁷ Reanalysed by SHAMOV 23.

NODE=M052M;LINKAGE=D

 $m\Upsilon(3S) - m\Upsilon(2S)$

NODE=M052DM3

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M052DM3

 $\Upsilon(2S)$ WIDTH

NODE=M052W

VALUE (keV)	DOCUMENT ID	COMMENT
31.98±2.63 OUR EVALUATION	See the Note on "Width Determinations of the Υ States"	

NODE=M052W

→ UNCHECKED ←

 $\Upsilon(2S)$ DECAY MODES

NODE=M052215;NODE=M052

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(1S)\pi^+\pi^-$	(17.85±0.26) %	
Γ_2 $\Upsilon(1S)\pi^0\pi^0$	(8.6±0.4) %	
Γ_3 $\tau^+\tau^-$	(2.00±0.21) %	
Γ_4 $\mu^+\mu^-$	(1.93±0.17) %	S=2.2
Γ_5 e^+e^-	(1.91±0.16) %	
Γ_6 $\Upsilon(1S)\pi^0$	< 4	× 10 ⁻⁵ CL=90%
Γ_7 $\Upsilon(1S)\eta$	(2.9±0.4) × 10 ⁻⁴	S=2.0
Γ_8 $J/\psi(1S)$ anything	< 6	× 10 ⁻³ CL=90%
Γ_9 $J/\psi(1S)\eta_c$	< 5.4	× 10 ⁻⁶ CL=90%
Γ_{10} $J/\psi(1S)\chi_{c0}$	< 3.4	× 10 ⁻⁶ CL=90%
Γ_{11} $J/\psi(1S)\chi_{c1}$	< 1.2	× 10 ⁻⁶ CL=90%
Γ_{12} $J/\psi(1S)\chi_{c2}$	< 2.0	× 10 ⁻⁶ CL=90%
Γ_{13} $J/\psi(1S)\eta_c(2S)$	< 2.5	× 10 ⁻⁶ CL=90%
Γ_{14} $J/\psi(1S)X(3940)$	< 2.0	× 10 ⁻⁶ CL=90%
Γ_{15} $J/\psi(1S)X(4160)$	< 2.0	× 10 ⁻⁶ CL=90%
Γ_{16} χ_{c1} anything	(2.2±0.5) × 10 ⁻⁴	
Γ_{17} $\chi_{c1}(1P)^0 X_{tetra}$	< 3.67	× 10 ⁻⁵ CL=90%
Γ_{18} χ_{c2} anything	(2.3±0.8) × 10 ⁻⁴	
Γ_{19} $\psi(2S)\eta_c$	< 5.1	× 10 ⁻⁶ CL=90%
Γ_{20} $\psi(2S)\chi_{c0}$	< 4.7	× 10 ⁻⁶ CL=90%
Γ_{21} $\psi(2S)\chi_{c1}$	< 2.5	× 10 ⁻⁶ CL=90%
Γ_{22} $\psi(2S)\chi_{c2}$	< 1.9	× 10 ⁻⁶ CL=90%
Γ_{23} $\psi(2S)\eta_c(2S)$	< 3.3	× 10 ⁻⁶ CL=90%
Γ_{24} $\psi(2S)X(3940)$	< 3.9	× 10 ⁻⁶ CL=90%
Γ_{25} $\psi(2S)X(4160)$	< 3.9	× 10 ⁻⁶ CL=90%
Γ_{26} $T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-$	< 1.0	× 10 ⁻⁶ CL=90%

DESIG=4

DESIG=5

DESIG=3

DESIG=1

DESIG=2

DESIG=10

DESIG=6

DESIG=20

DESIG=143

DESIG=144

DESIG=145

DESIG=146

DESIG=147

DESIG=148

DESIG=149

DESIG=157

DESIG=160

DESIG=158

DESIG=150

DESIG=151

DESIG=152

DESIG=153

DESIG=154

DESIG=155

DESIG=156

DESIG=162

Γ ₂₇	$T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-$	< 1.67	$\times 10^{-5}$	CL=90%	DESIG=163
Γ ₂₈	$T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp$	< 7.3	$\times 10^{-6}$	CL=90%	DESIG=164
Γ ₂₉	$T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-$	< 1.35	$\times 10^{-5}$	CL=90%	DESIG=165
Γ ₃₀	$T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-$	< 2.67	$\times 10^{-5}$	CL=90%	DESIG=166
Γ ₃₁	$T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp$	< 2.72	$\times 10^{-5}$	CL=90%	DESIG=167
Γ ₃₂	$T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-$	< 2.03	$\times 10^{-5}$	CL=90%	DESIG=168
Γ ₃₃	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp$	< 1.11	$\times 10^{-5}$	CL=90%	DESIG=170
Γ ₃₄	$T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp$	< 2.11	$\times 10^{-5}$	CL=90%	DESIG=171
Γ ₃₅	${}^2\overline{H}$ anything	$(2.78^{+0.30}_{-0.26}) \times 10^{-5}$		S=1.2	DESIG=16
Γ ₃₆	hadrons	$(94 \pm 11) \%$			DESIG=101
Γ ₃₇	ggg	$(58.8 \pm 1.2) \%$			DESIG=105
Γ ₃₈	γgg	$(1.87 \pm 0.28) \%$			DESIG=106
Γ ₃₉	$\phi K^+ K^-$	$(1.6 \pm 0.4) \times 10^{-6}$			DESIG=133
Γ ₄₀	$\omega \pi^+ \pi^-$	< 2.58	$\times 10^{-6}$	CL=90%	DESIG=134
Γ ₄₁	$K^*(892)^0 K^- \pi^+ + \text{c.c.}$	$(2.3 \pm 0.7) \times 10^{-6}$			DESIG=135
Γ ₄₂	$\phi f'_2(1525)$	< 1.33	$\times 10^{-6}$	CL=90%	DESIG=136
Γ ₄₃	$\omega f_2(1270)$	< 5.7	$\times 10^{-7}$	CL=90%	DESIG=137
Γ ₄₄	$\rho(770) a_2(1320)$	< 8.8	$\times 10^{-7}$	CL=90%	DESIG=138
Γ ₄₅	$K^*(892)^0 \overline{K}_2^*(1430)^0 + \text{c.c.}$	$(1.5 \pm 0.6) \times 10^{-6}$			DESIG=139
Γ ₄₆	$K_1(1270)^\pm K^\mp$	< 3.22	$\times 10^{-6}$	CL=90%	DESIG=140
Γ ₄₇	$K_1(1400)^\pm K^\mp$	< 8.3	$\times 10^{-7}$	CL=90%	DESIG=141
Γ ₄₈	$b_1(1235)^\pm \pi^\mp$	< 4.0	$\times 10^{-7}$	CL=90%	DESIG=142
Γ ₄₉	$\rho \pi$	< 1.16	$\times 10^{-6}$	CL=90%	DESIG=126
Γ ₅₀	$\pi^+ \pi^- \pi^0$	< 8.0	$\times 10^{-7}$	CL=90%	DESIG=127
Γ ₅₁	$\omega \pi^0$	< 1.63	$\times 10^{-6}$	CL=90%	DESIG=128
Γ ₅₂	$\pi^+ \pi^- \pi^0 \pi^0$	$(1.30 \pm 0.28) \times 10^{-5}$			DESIG=129
Γ ₅₃	$K_S^0 K^+ \pi^- + \text{c.c.}$	$(1.14 \pm 0.33) \times 10^{-6}$			DESIG=130
Γ ₅₄	$K^*(892)^0 \overline{K}^0 + \text{c.c.}$	< 4.22	$\times 10^{-6}$	CL=90%	DESIG=131
Γ ₅₅	$K^*(892)^- K^+ + \text{c.c.}$	< 1.45	$\times 10^{-6}$	CL=90%	DESIG=132
Γ ₅₆	$f_1(1285)$ anything	$(2.2 \pm 1.6) \times 10^{-3}$			DESIG=159
Γ ₅₇	$f_1(1285) X_{tetra}$	< 6.47	$\times 10^{-5}$	CL=90%	DESIG=161
Γ ₅₈	$D_s^+ D_{s1}(2536)^-$				DESIG=177
Γ ₅₉	$D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0$	$(1.6 \pm 0.4) \times 10^{-5}$			DESIG=178
Γ ₆₀	$D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-$	$(8.4 \pm 2.3) \times 10^{-6}$			DESIG=179
Γ ₆₁	$D_s^{*+} D_{s1}(2536)^-$				DESIG=180
Γ ₆₂	$D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0$	$(1.4 \pm 0.4) \times 10^{-5}$			DESIG=181
Γ ₆₃	$D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-$	$(8.2 \pm 3.1) \times 10^{-6}$			DESIG=182
Γ ₆₄	$D_s^+ D_{s2}^*(2573)^-$				DESIG=183
Γ ₆₅	$D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0$	$(1.4 \pm 0.4) \times 10^{-5}$			DESIG=184
Γ ₆₆	$D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-$	$(6.9 \pm 3.0) \times 10^{-6}$			DESIG=185
Γ ₆₇	$D_s^{*+} D_{s2}^*(2573)^-$				DESIG=186
Γ ₆₈	$D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0$	$(9 \pm 5) \times 10^{-6}$			DESIG=187
Γ ₆₉	$D_s^{*+} D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-$	$(5 \pm 6) \times 10^{-6}$			DESIG=188
Γ ₇₀	Sum of 100 exclusive modes	$(2.90 \pm 0.30) \times 10^{-3}$			DESIG=121

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_{36}\Gamma_5/\Gamma$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.577 ± 0.009 OUR AVERAGE			
0.581 ± 0.004 ± 0.009	¹ ROSNER	06	CLEO $e^+e^- \rightarrow \text{hadrons}$
0.552 ± 0.031 ± 0.017	¹ BARU	96	MD1 $e^+e^- \rightarrow \text{hadrons}$
0.54 ± 0.04 ± 0.02	¹ JAKUBOWSKI	88	CBAL $e^+e^- \rightarrow \text{hadrons}$
0.58 ± 0.03 ± 0.04	² GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$
0.60 ± 0.12 ± 0.07	² ALBRECHT	82	DASP $e^+e^- \rightarrow \text{hadrons}$
0.54 ± 0.07 ^{+0.09} _{-0.05}	² NICZYPORUK	81C	LENA $e^+e^- \rightarrow \text{hadrons}$
0.41 ± 0.18	² BOCK	80	CNTR $e^+e^- \rightarrow \text{hadrons}$

¹ Radiative corrections evaluated following KURAEV 85.² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.NODE=M052G2
NODE=M052G2NODE=M052G2;LINKAGE=P
NODE=M052G2;LINKAGE=R $\Upsilon(2S)$ PARTIAL WIDTHS $\Gamma(e^+e^-)$ Γ_5

VALUE (keV)	DOCUMENT ID
0.612 ± 0.011 OUR EVALUATION	

NODE=M052220

NODE=M052W2
NODE=M052W2
→ UNCHECKED ← $\Upsilon(2S)$ BRANCHING RATIOS $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_1/Γ

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
17.85 ± 0.26 OUR FIT				
17.92 ± 0.26 OUR AVERAGE				
16.8 ± 1.1 ± 1.3	906k	¹ LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$
17.80 ± 0.05 ± 0.37	170k	² LEES	11L	BABR $\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
18.02 ± 0.02 ± 0.61	851k	³ BHARI	09	CLEO $e^+e^- \rightarrow \pi^+\pi^-MM$
17.22 ± 0.17 ± 0.75	11.8k	⁴ AUBERT	08BP	BABR $e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
19.2 ± 0.2 ± 1.0	52.6k	⁵ ALEXANDER	98	CLE2 $\pi^+\pi^-\ell^+\ell^-, \pi^+\pi^-MM$
18.1 ± 0.5 ± 1.0	11.6k	ALBRECHT	87	ARG $e^+e^- \rightarrow \pi^+\pi^-MM$
16.9 ± 4.0		GELPHMAN	85	CBAL $e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
19.1 ± 1.2 ± 0.6		BESSION	84	CLEO $\pi^+\pi^-MM$
18.9 ± 2.6		FONSECA	84	CUSB $e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$
21 ± 7	7	NICZYPORUK	81B	LENA $e^+e^- \rightarrow \ell^+\ell^-\pi^+\pi^-$

NODE=M052225

NODE=M052R4
NODE=M052R4
NODE=M052R4¹ LEES 11c reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything})] = (1.78 \pm 0.02 \pm 0.11) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \Upsilon(2S)\text{anything}) = (10.6 \pm 0.8) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.² Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.³ A weighted average of the inclusive and exclusive results.⁴ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$ and, $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.⁵ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.

NODE=M052R4;LINKAGE=ES

NODE=M052R4;LINKAGE=LE
NODE=M052R4;LINKAGE=BH
NODE=M052R4;LINKAGE=AU

NODE=M052R4;LINKAGE=T

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.6 ± 0.4 OUR AVERAGE				
8.43 ± 0.16 ± 0.42	38k	¹ BHARI	09	CLEO $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.2 ± 0.6 ± 0.8	275	² ALEXANDER	98	CLE2 $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
9.5 ± 1.9 ± 1.9	25	ALBRECHT	87	ARG $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
8.0 ± 1.5		GELPHMAN	85	CBAL $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
10.3 ± 2.3		FONSECA	84	CUSB $e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

¹ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.52 \pm 0.17)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.07)\%$.NODE=M052R5
NODE=M052R5NODE=M052R5;LINKAGE=BH
NODE=M052R5;LINKAGE=T $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_1

VALUE	DOCUMENT ID	TECN	COMMENT
0.462 ± 0.037	¹ BHARI	09	CLEO $e^+e^- \rightarrow \Upsilon(2S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ Not independent of other values reported by BHARI 09.NODE=M052R21
NODE=M052R21

NODE=M052R21;LINKAGE=BH

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.00±0.21 OUR AVERAGE				
2.00±0.12±0.18	22k	¹ BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S) \rightarrow \tau^+\tau^-$
1.7 ±1.5 ±0.6		HAAS 84B	CLEO	$e^+e^- \rightarrow \tau^+\tau^-$

NODE=M052R3
NODE=M052R3

¹ BESSON 07 reports $[\Gamma(\Upsilon(2S) \rightarrow \tau^+\tau^-)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \mu^+\mu^-)] = 1.04 \pm 0.04 \pm 0.05$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M052R3;LINKAGE=BE

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0193±0.0017 OUR AVERAGE Error includes scale factor of 2.2. See the ideogram below.					
0.0203±0.0003±0.0008		120k	ADAMS 05	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0122±0.0028±0.0019			¹ KOBEL 92	CBAL	$e^+e^- \rightarrow \mu^+\mu^-$
0.0138±0.0025±0.0015			KAARSBERG 89	CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.009 ±0.006 ±0.006			² ALBRECHT 85	ARG	$e^+e^- \rightarrow \mu^+\mu^-$
0.018 ±0.008 ±0.005			HAAS 84B	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M052R1
NODE=M052R1


• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.038	90	NICZYPORUK 81C	LENA	$e^+e^- \rightarrow \mu^+\mu^-$
--------	----	----------------	------	---------------------------------

NODE=M052R1;LINKAGE=A
NODE=M052R1;LINKAGE=R

¹ Taking into account interference between the resonance and continuum.

² Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 0.026$.



$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$

Γ_3/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.04±0.04±0.05				
	22k	BESSON 07	CLEO	$e^+e^- \rightarrow \Upsilon(2S)$

NODE=M052R17
NODE=M052R17

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$

Γ_6/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 4	90	¹ TAMPONI 13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^0$
< 18	90	² HE 08A	CLEO	$e^+e^- \rightarrow l^+l^-\gamma\gamma$
<110	90	ALEXANDER 98	CLE2	$e^+e^- \rightarrow l^+l^-\gamma\gamma$
<800	90	LURZ 87	CBAL	$e^+e^- \rightarrow l^+l^-\gamma\gamma$

NODE=M052R10
NODE=M052R10

¹ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0)/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] < 2.3 \times 10^{-4}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = 17.85 \times 10^{-2}$.

NODE=M052R10;LINKAGE=TA

² Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M052R10;LINKAGE=HE

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_6/Γ_1

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.3	90	TAMPONI	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0$

NODE=M052R09
NODE=M052R09

$\Gamma(\Upsilon(1S)\eta)/\Gamma_{total}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

2.9 ± 0.4 OUR FIT Error includes scale factor of 2.0.
2.9 ± 0.4 OUR AVERAGE Error includes scale factor of 1.9. See the ideogram below.

2.39 ± 0.31 ± 0.14	112	¹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow \ell^+\ell^-\eta$
2.1 ^{+0.7} / _{-0.6} ± 0.3	14	² HE	08A	CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$

NODE=M052R6
NODE=M052R6

• • • We use the following data for averages but not for fits. • • •

3.55 ± 0.32 ± 0.05	241	³ TAMPONI	13	BELL	$e^+e^- \rightarrow \Upsilon(1S)\eta$
--------------------	-----	----------------------	----	------	---------------------------------------

NOTFITTED

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	^{1,4} AUBERT	08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
< 28	90	ALEXANDER98		CLE2	$e^+e^- \rightarrow \ell^+\ell^-\eta$
< 50	90	ALBRECHT	87	ARG	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-MM$
< 70	90	LURZ	87	CBAL	$e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, 3\pi^0)$
< 100	90	BESSON	84	CLEO	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-MM$
< 20	90	FONSECA	84	CUSB	$e^+e^- \rightarrow \ell^+\ell^-(\gamma\gamma, \pi^+\pi^-\pi^0)$

¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M052R6;LINKAGE=AU

² Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M052R6;LINKAGE=HE


³ TAMPONI 13 reports $[\Gamma(\Upsilon(2S) \rightarrow \Upsilon(1S)\eta)/\Gamma_{total}] / [B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)] = (1.99 \pm 0.14 \pm 0.11) \times 10^{-3}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (17.85 \pm 0.26) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M052R6;LINKAGE=TA

⁴ Using $\Gamma_{ee}(\Upsilon(2S)) = 0.612 \pm 0.011$ keV.

NODE=M052R6;LINKAGE=UB

WEIGHTED AVERAGE
2.9±0.4 (Error scaled by 1.9)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our 'best' values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_1

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

1.64 ± 0.25 OUR FIT Error includes scale factor of 2.0.
1.99 ± 0.14 ± 0.11 241 TAMPONI 13 BELL $e^+e^- \rightarrow \Upsilon(1S)\eta$

NODE=M052R22
NODE=M052R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.35 ± 0.17 ± 0.08		¹ LEES	11L	BABR	$\Upsilon(2S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\mu^+\mu^-$
< 5.2	90	² AUBERT	08BP	BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$

¹ Not independent of other values reported by LEES 11L.

NODE=M052R22;LINKAGE=LE

² Not independent of other values reported by AUBERT 08BP.

NODE=M052R22;LINKAGE=AU

$\Gamma(\Upsilon(1S)\pi^0)/\Gamma(\Upsilon(1S)\eta)$					Γ_6/Γ_7	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.13	90	TAMPONI	13	BELL	$e^+ e^- \rightarrow \Upsilon(1S)\pi^0$	NODE=M052R23 NODE=M052R23
$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$					Γ_8/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<0.006	90	MASCHMANN	90	CBAL	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M052R16 NODE=M052R16
$\Gamma(J/\psi(1S)\eta_c)/\Gamma_{\text{total}}$					Γ_9/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<5.4 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R53 NODE=M052R53
$\Gamma(J/\psi(1S)\chi_{c0})/\Gamma_{\text{total}}$					Γ_{10}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<3.4 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R54 NODE=M052R54
$\Gamma(J/\psi(1S)\chi_{c1})/\Gamma_{\text{total}}$					Γ_{11}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<1.2 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R55 NODE=M052R55
$\Gamma(J/\psi(1S)\chi_{c2})/\Gamma_{\text{total}}$					Γ_{12}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.0 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R56 NODE=M052R56
$\Gamma(J/\psi(1S)\eta_c(2S))/\Gamma_{\text{total}}$					Γ_{13}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.5 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R57 NODE=M052R57
$\Gamma(J/\psi(1S)X(3940))/\Gamma_{\text{total}}$					Γ_{14}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.0 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R58 NODE=M052R58
$\Gamma(J/\psi(1S)X(4160))/\Gamma_{\text{total}}$					Γ_{15}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.0 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow J/\psi X$	NODE=M052R59 NODE=M052R59
$\Gamma(\chi_{c1} \text{ anything})/\Gamma_{\text{total}}$					Γ_{16}/Γ	
VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT		
2.24 ± 0.44 ± 0.20	376	JIA	17	BELL	$\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$	NODE=M052R00 NODE=M052R00
$\Gamma(\chi_{c1}(1P)^0 X_{\text{tetra}})/\Gamma_{\text{total}}$					Γ_{17}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<36.7 × 10 ⁻⁶	90	¹ JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$	NODE=M052R69 NODE=M052R69
¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 4.4×10^{-6} to 36.7×10^{-6} .						
$\Gamma(\chi_{c2} \text{ anything})/\Gamma_{\text{total}}$					Γ_{18}/Γ	
VALUE (units 10 ⁻⁴)		DOCUMENT ID	TECN	COMMENT		
2.28 ± 0.73 ± 0.34		JIA	17	BELL	$\Upsilon(2S) \rightarrow \gamma J/\psi(1S)$	NODE=M052R67 NODE=M052R67
$\Gamma(\psi(2S)\eta_c)/\Gamma_{\text{total}}$					Γ_{19}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<5.1 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R60 NODE=M052R60
$\Gamma(\psi(2S)\chi_{c0})/\Gamma_{\text{total}}$					Γ_{20}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<4.7 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R61 NODE=M052R61
$\Gamma(\psi(2S)\chi_{c1})/\Gamma_{\text{total}}$					Γ_{21}/Γ	
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
<2.5 × 10 ⁻⁶	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	NODE=M052R62 NODE=M052R62

NODE=M052R69;LINKAGE=A

$\Gamma(\psi(2S)\chi_{c2})/\Gamma_{\text{total}}$					Γ_{22}/Γ	NODE=M052R63 NODE=M052R63
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<1.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	
$\Gamma(\psi(2S)\eta_c(2S))/\Gamma_{\text{total}}$					Γ_{23}/Γ	NODE=M052R64 NODE=M052R64
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.3 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	
$\Gamma(\psi(2S)X(3940))/\Gamma_{\text{total}}$					Γ_{24}/Γ	NODE=M052R65 NODE=M052R65
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	
$\Gamma(\psi(2S)X(4160))/\Gamma_{\text{total}}$					Γ_{25}/Γ	NODE=M052R66 NODE=M052R66
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<3.9 \times 10^{-6}$	90	YANG	14	BELL	$e^+ e^- \rightarrow \psi(2S) X$	
$\Gamma(T_{c\bar{c}1}(3900)^+ T_{c\bar{c}1}(3900)^-)/\Gamma_{\text{total}}$					Γ_{26}/Γ	NODE=M052R71 NODE=M052R71
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<1.0 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm) = 1$.				
						NODE=M052R71;LINKAGE=A
$\Gamma(T_{c\bar{c}1}(4200)^+ T_{c\bar{c}1}(4200)^-)/\Gamma_{\text{total}}$					Γ_{27}/Γ	NODE=M052R72 NODE=M052R72
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<16.7 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(1S) \rightarrow J/\psi \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1$				
						NODE=M052R72;LINKAGE=A
$\Gamma(T_{c\bar{c}1}(3900)^\pm T_{c\bar{c}1}(4200)^\mp)/\Gamma_{\text{total}}$					Γ_{28}/Γ	NODE=M052R73 NODE=M052R73
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<7.3 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow J/\psi \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}1}(4200)^\pm \rightarrow J/\psi \pi^\pm) = 1 = B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi \pi^\pm)$.				
						NODE=M052R73;LINKAGE=A
$\Gamma(T_{c\bar{c}}(4050)^+ T_{c\bar{c}}(4050)^-)/\Gamma_{\text{total}}$					Γ_{29}/Γ	NODE=M052R74 NODE=M052R74
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<13.5 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$				
						NODE=M052R74;LINKAGE=A
$\Gamma(T_{c\bar{c}}(4250)^+ T_{c\bar{c}}(4250)^-)/\Gamma_{\text{total}}$					Γ_{30}/Γ	NODE=M052R75 NODE=M052R75
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<26.7 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1$				
						NODE=M052R75;LINKAGE=A
$\Gamma(T_{c\bar{c}}(4050)^\pm T_{c\bar{c}}(4250)^\mp)/\Gamma_{\text{total}}$					Γ_{31}/Γ	NODE=M052R76 NODE=M052R76
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<27.2 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow \chi_{c1}(1P) \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}}(4050)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm) = 1 = B(T_{c\bar{c}}(4250)^\pm \rightarrow \chi_{c1}(1P) \pi^\pm)$				
						NODE=M052R76;LINKAGE=A
$\Gamma(T_{c\bar{c}1}(4430)^+ T_{c\bar{c}1}(4430)^-)/\Gamma_{\text{total}}$					Γ_{32}/Γ	NODE=M052R77 NODE=M052R77
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<20.3 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2P) \pi^\pm) = 1$				
						NODE=M052R77;LINKAGE=A
$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}}(4055)^\mp)/\Gamma_{\text{total}}$					Γ_{33}/Γ	NODE=M052R79 NODE=M052R79
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<11.1 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1$				
						NODE=M052R79;LINKAGE=A
$\Gamma(T_{c\bar{c}}(4055)^\pm T_{c\bar{c}1}(4430)^\mp)/\Gamma_{\text{total}}$					Γ_{34}/Γ	NODE=M052R80 NODE=M052R80
VALUE	CL%	DOCUMENT ID	TECN	COMMENT		
$<21.1 \times 10^{-6}$	90	1 JIA	18	BELL	$\Upsilon(2S) \rightarrow \psi(2S) \pi^\pm X$	
		¹ Assuming $B(T_{c\bar{c}}(4055)^\pm \rightarrow \psi(2S) \pi^\pm) = 1 = B(T_{c\bar{c}1}(4430)^\pm \rightarrow \psi(2S) \pi^\pm)$				
						NODE=M052R80;LINKAGE=A

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M052R18
NODE=M052R18

2.78^{+0.30}_{-0.26} OUR AVERAGE Error includes scale factor of 1.2.

2.64 ± 0.11 ^{+0.26} _{-0.21}		LEES	14G	BABR	$e^+ e^- \rightarrow \overline{2H} X$
3.37 ± 0.50 ± 0.25	58	ASNER	07	CLEO	$e^+ e^- \rightarrow \overline{2H} X$

 $\Gamma(g g g)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M052R01
NODE=M052R01

58.8 ± 1.2 6M ¹ BESSON 06A CLEO $\Upsilon(2S) \rightarrow \text{hadrons}$

¹ Calculated using the value $\Gamma(\gamma g g)/\Gamma(g g g) = (3.18 \pm 0.04 \pm 0.22 \pm 0.41)\%$ from BESSON 06A and PDG 08 values of $B(\pi^+ \pi^- \Upsilon(1S)) = (18.1 \pm 0.4)\%$, $B(\pi^0 \pi^0 \Upsilon(1S)) = (8.6 \pm 0.4)\%$, $B(\mu^+ \mu^-) = (1.93 \pm 0.17)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with that of $\Gamma(\gamma g g)/\Gamma_{\text{total}}$ measurement of BESSON 06A.

NODE=M052R01;LINKAGE=BE

 $\Gamma(\gamma g g)/\Gamma(g g g)$ Γ_{38}/Γ_{37}

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M052R03
NODE=M052R03

3.18 ± 0.04 ± 0.47 6M BESSON 06A CLEO $\Upsilon(2S) \rightarrow (\gamma +) \text{hadrons}$

 $\Gamma(\phi K^+ K^-)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M052R43
NODE=M052R43

1.58 ± 0.33 ± 0.18 58 SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R44
NODE=M052R44

<2.58 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M052R45
NODE=M052R45

2.32 ± 0.40 ± 0.54 135 SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R46
NODE=M052R46

<1.33 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(K^+ K^-)$

 $\Gamma(\omega f_2(1270))/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R47
NODE=M052R47

<0.57 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho(770) a_2(1320))/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R48
NODE=M052R48

<0.88 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(K^*(892)^0 \overline{K}_2^*(1430)^0 + \text{c.c.})/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (units 10^{-6})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M052R49
NODE=M052R49

1.53 ± 0.52 ± 0.19 32 SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1270)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R50
NODE=M052R50

<3.22 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(K_1(1400)^\pm K^\mp)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R51
NODE=M052R51

<0.83 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow K^+ K^- \pi^+ \pi^-$

 $\Gamma(b_1(1235)^\pm \pi^\mp)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R52
NODE=M052R52

<0.40 90 SHEN 12A BELL $\Upsilon(1S) \rightarrow 2(\pi^+ \pi^-) \pi^0$

 $\Gamma(\rho\pi)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M052R27
NODE=M052R27

<1.16 90 SHEN 13 BELL $\Upsilon(2S) \rightarrow \pi^+ \pi^- \pi^0$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$						Γ_{50}/Γ	NODE=M052R28 NODE=M052R28
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<0.80	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0$		
$\Gamma(\omega\pi^0)/\Gamma_{\text{total}}$						Γ_{51}/Γ	NODE=M052R29 NODE=M052R29
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<1.63	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$		
$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$						Γ_{52}/Γ	NODE=M052R30 NODE=M052R30
<u>VALUE (units 10^{-6})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
13.0±1.9±2.1	261 ± 37	SHEN	13	BELL	$\Upsilon(2S) \rightarrow \pi^+\pi^-\pi^0\pi^0$		
$\Gamma(K_S^0 K^+ \pi^- + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{53}/Γ	NODE=M052R40 NODE=M052R40
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.14±0.30±0.13	40 ± 10	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<3.2	90	¹ DOBBS	12A		$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$		
¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.							
$\Gamma(K^*(892)^0 \bar{K}^0 + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{54}/Γ	NODE=M052R41 NODE=M052R41
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<4.22	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$		
$\Gamma(K^*(892)^- K^+ + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{55}/Γ	NODE=M052R42 NODE=M052R42
<u>VALUE (units 10^{-6})</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<1.45	90	SHEN	13	BELL	$\Upsilon(2S) \rightarrow K_S^0 K^- \pi^+$		
$\Gamma(f_1(1285) \text{ anything})/\Gamma_{\text{total}}$						Γ_{56}/Γ	NODE=M052R68 NODE=M052R68
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
2.20±1.50±0.63	2.9k	JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$		
$\Gamma(f_1(1285) X_{\text{tetra}})/\Gamma_{\text{total}}$						Γ_{57}/Γ	NODE=M052R70 NODE=M052R70
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
<64.7 × 10 ⁻⁶	90	¹ JIA	17A	BELL	$e^+ e^- \rightarrow \text{hadrons}$		
¹ For a tetraquark state X_{tetra} , with mass in the range 1.16–2.46 GeV and width in the range 0–0.3 GeV. Measured 90% CL limits as a function of X_{tetra} mass and width range from 7.8×10^{-6} to 64.7×10^{-6} .							
$\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{\text{total}}$						Γ_{59}/Γ	NODE=M052R86 NODE=M052R86
<u>VALUE (units 10^{-5})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
1.6±0.3±0.2		GAO	23	BELL	$e^+ e^-$ at 10.52 GeV		
$\Gamma(D_s^+ D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{\text{total}}$						Γ_{60}/Γ	NODE=M052R87 NODE=M052R87
<u>VALUE (units 10^{-5})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
0.84±0.18±0.15		GAO	23	BELL	$e^+ e^-$ at 10.52 GeV		
$\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K^- D^*(2007)^0)/\Gamma_{\text{total}}$						Γ_{62}/Γ	NODE=M052R88 NODE=M052R88
<u>VALUE (units 10^{-5})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
1.4±0.4±0.2		GAO	23	BELL	$e^+ e^-$ at 10.52 GeV		
$\Gamma(D_s^{*+} D_{s1}(2536)^-, D_{s1}^- \rightarrow K_S^0 D^*(2010)^-)/\Gamma_{\text{total}}$						Γ_{63}/Γ	NODE=M052R89 NODE=M052R89
<u>VALUE (units 10^{-5})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
0.82±0.25±0.19		GAO	23	BELL	$e^+ e^-$ at 10.52 GeV		
$\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{\text{total}}$						Γ_{65}/Γ	NODE=M052R90 NODE=M052R90
<u>VALUE (units 10^{-5})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
1.4±0.4±0.2		GAO	23	BELL	$e^+ e^-$ at 10.52 GeV		
$\Gamma(D_s^+ D_{s2}^*(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{\text{total}}$						Γ_{66}/Γ	NODE=M052R91 NODE=M052R91
<u>VALUE (units 10^{-5})</u>		<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
0.69±0.20±0.22		GAO	23	BELL	$e^+ e^-$ at 10.52 GeV		

$$\Gamma(D_s^{*+} D_{s2}^{*-}(2573)^-, D_{s2}^{*-} \rightarrow K^- D^0)/\Gamma_{\text{total}} \quad \Gamma_{68}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
0.9 ± 0.5 ± 0.2	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

NODE=M052R92
NODE=M052R92

$$\Gamma(D_s^{*+} D_{s2}^{*-}(2573)^-, D_{s2}^{*-} \rightarrow K_S^0 D^-)/\Gamma_{\text{total}} \quad \Gamma_{69}/\Gamma$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
0.54 ± 0.31 ± 0.47	GAO	23	BELL $e^+ e^-$ at 10.52 GeV

NODE=M052R93
NODE=M052R93

$$\Gamma(\text{Sum of 100 exclusive modes})/\Gamma_{\text{total}} \quad \Gamma_{70}/\Gamma$$

VALUE (units 10^{-2})	DOCUMENT ID	COMMENT
0.29 ± 0.03	1,2 DOBBS 12A	$\Upsilon(2S) \rightarrow \text{hadrons}$

NODE=M052R08
NODE=M052R08

¹ DOBBS 12A presents individual exclusive branching fractions or upper limits for 100 modes of four to ten pions, kaons, or protons.

NODE=M052R08;LINKAGE=DO

² Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M052R08;LINKAGE=NC

$$\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}} \quad \Gamma_{71}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.069 ± 0.004 OUR AVERAGE				
0.0693 ± 0.0012 ± 0.0041	407k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.069 ± 0.005 ± 0.009		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
0.091 ± 0.018 ± 0.022		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.065 ± 0.007 ± 0.012		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
0.080 ± 0.017 ± 0.016		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.059 ± 0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

NODE=M052R8
NODE=M052R8

$$\Gamma(\gamma \chi_{b2}(1P))/\Gamma_{\text{total}} \quad \Gamma_{72}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0715 ± 0.0035 OUR AVERAGE				
0.0724 ± 0.0011 ± 0.0040	410k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.074 ± 0.005 ± 0.008		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
0.098 ± 0.021 ± 0.024		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.058 ± 0.007 ± 0.010		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
0.102 ± 0.018 ± 0.021		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.061 ± 0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$

NODE=M052R7
NODE=M052R7

$$\Gamma(\gamma \chi_{b0}(1P))/\Gamma_{\text{total}} \quad \Gamma_{73}/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.038 ± 0.004 OUR AVERAGE				
0.0375 ± 0.0012 ± 0.0047	198k	ARTUSO	05	CLEO $e^+ e^- \rightarrow \gamma X$
0.034 ± 0.005 ± 0.006		EDWARDS	99	CLE2 $\Upsilon(2S) \rightarrow \gamma \chi(1P)$
0.064 ± 0.014 ± 0.016		ALBRECHT	85E	ARG $e^+ e^- \rightarrow \gamma \text{conv. } X$
0.036 ± 0.008 ± 0.009		NERNST	85	CBAL $e^+ e^- \rightarrow \gamma X$
0.044 ± 0.023 ± 0.009		HAAS	84	CLEO $e^+ e^- \rightarrow \gamma \text{conv. } X$

NODE=M052R9
NODE=M052R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.035 ± 0.014		KLOPFEN...	83	CUSB $e^+ e^- \rightarrow \gamma X$
---------------	--	------------	----	-------------------------------------

$$\Gamma(\gamma f_0(1710))/\Gamma_{\text{total}} \quad \Gamma_{74}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 59	90	¹ ALBRECHT	89	ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$

NODE=M052R13
NODE=M052R13

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.9	90	² ALBRECHT	89	ARG $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$
-------	----	-----------------------	----	---

OCCUR=2

¹ Re-evaluated assuming $B(f_0(1710) \rightarrow K^+ K^-) = 0.19$.

NODE=M052R13;LINKAGE=M

² Includes unknown branching ratio of $f_0(1710) \rightarrow \pi^+ \pi^-$.

NODE=M052R13;LINKAGE=N

$$\Gamma(\gamma f_2'(1525))/\Gamma_{\text{total}} \quad \Gamma_{75}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 53	90	¹ ALBRECHT	89	ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$

NODE=M052R12
NODE=M052R12

¹ Re-evaluated assuming $B(f_2'(1525) \rightarrow K \bar{K}) = 0.71$.

NODE=M052R12;LINKAGE=L

$$\Gamma(\gamma f_2(1270))/\Gamma_{\text{total}} \quad \Gamma_{76}/\Gamma$$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
< 24.1	90	¹ ALBRECHT	89	ARG $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$

NODE=M052R11
NODE=M052R11

¹ Using $B(f_2(1270) \rightarrow \pi \pi) = 0.84$.

NODE=M052R11;LINKAGE=K

$\Gamma(\gamma f_J(2220))/\Gamma_{\text{total}}$ Γ_{77}/Γ NODE=M052R14
 VALUE (units 10^{-5}) CL% DOCUMENT ID TECN COMMENT NODE=M052R14

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.8 90 ¹ ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$

¹ Includes unknown branching ratio of $f_J(2220) \rightarrow K^+ K^-$.

NODE=M052R14;LINKAGE=S

$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$ Γ_{78}/Γ NODE=M052R31
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R31
 <2.7 $\times 10^{-5}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \chi_{c0})/\Gamma_{\text{total}}$ Γ_{79}/Γ NODE=M052R32
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R32
 <1.0 $\times 10^{-4}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \chi_{c1})/\Gamma_{\text{total}}$ Γ_{80}/Γ NODE=M052R33
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R33
 <3.6 $\times 10^{-6}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \chi_{c2})/\Gamma_{\text{total}}$ Γ_{81}/Γ NODE=M052R34
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R34
 <1.5 $\times 10^{-5}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \chi_{c1}(3872))/\Gamma_{\text{total}}$ Γ_{82}/Γ NODE=M052R81
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R81
 <2.2 $\times 10^{-5}$ (CL = 90%) [$<2.1 \times 10^{-5}$ (CL = 90%) OUR 2023 BEST LIMIT]
 <2.3 $\times 10^{-5}$ 90 ¹ WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

¹ WANG 11B reports $[\Gamma(\Upsilon(2S) \rightarrow \gamma \chi_{c1}(3872))/\Gamma_{\text{total}}] \times [B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S))] < 0.8 \times 10^{-6}$ which we divide by our best value $B(\chi_{c1}(3872) \rightarrow \pi^+ \pi^- J/\psi(1S)) = 3.5 \times 10^{-2}$.

NODE=M052R81;LINKAGE=A

$\Gamma(\gamma \chi_{c1}(3872), \chi_{c1} \rightarrow \pi^+ \pi^- \pi^0 J/\psi)/\Gamma_{\text{total}}$ Γ_{83}/Γ NODE=M052R36
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R36
 <2.4 $\times 10^{-6}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \chi_{c0}(3915) \rightarrow \omega J/\psi)/\Gamma_{\text{total}}$ Γ_{84}/Γ NODE=M052R37
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R37
 <2.8 $\times 10^{-6}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \chi_{c1}(4140) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{85}/Γ NODE=M052R38
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R38
 <1.2 $\times 10^{-6}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma X(4350) \rightarrow \phi J/\psi)/\Gamma_{\text{total}}$ Γ_{86}/Γ NODE=M052R39
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R39
 <1.3 $\times 10^{-6}$ 90 WANG 11B BELL $\Upsilon(2S) \rightarrow \gamma X$

$\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$ Γ_{87}/Γ NODE=M052R15
 VALUE (units 10^{-4}) CL% EVTS DOCUMENT ID TECN COMMENT NODE=M052R15

5.5^{+1.1}_{-0.9} OUR AVERAGE Error includes scale factor of 1.2.

6.1^{+0.6+0.9}_{-0.7-0.6} 29k FULSOM 18 BELL $\Upsilon(2S) \rightarrow \gamma X$

3.9 \pm 1.1^{+1.1}_{-0.9} 13 \pm 5k ¹ AUBERT 09AQ BABR $\Upsilon(2S) \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<21 90 LEES 11J BABR $\Upsilon(2S) \rightarrow X \gamma$

< 8.4 90 ¹ BONVICINI 10 CLEO $\Upsilon(2S) \rightarrow \gamma X$

< 5.1 90 ² ARTUSO 05 CLEO $e^+ e^- \rightarrow \gamma X$

¹ Assuming $\Gamma_{\eta_b(1S)} = 10$ MeV.

² Superseded by BONVICINI 10.

NODE=M052R15;LINKAGE=BO
 NODE=M052R15;LINKAGE=SU

$\Gamma(\gamma \eta_b(1S) \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{88}/Γ NODE=M052R25
 VALUE CL% DOCUMENT ID TECN COMMENT NODE=M052R25
 <3.7 $\times 10^{-6}$ 90 SANDILYA 13 BELL $\Upsilon(2S) \rightarrow \gamma$ hadrons

$\Gamma(\gamma X_{b\bar{b}} \rightarrow \gamma \text{Sum of 26 exclusive modes})/\Gamma_{\text{total}}$ Γ_{89}/Γ

VALUE (units 10^{-6})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
< 4.9	90		SANDILYA	13 BELL	$\Upsilon(2S) \rightarrow \gamma$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$46.2^{+29.7}_{-14.2} \pm 10.6$	10		¹ DOBBS	12	$\Upsilon(2S) \rightarrow \gamma$ hadrons

¹ Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

NODE=M052R26
NODE=M052R26

NODE=M052R26;LINKAGE=DO

 $\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}}$ Γ_{90}/Γ

(1.5 GeV < m_X < 5.0 GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.95	95	ROSNER	07A CLEO	$e^+e^- \rightarrow \gamma X$

NODE=M052R19
NODE=M052R19
NODE=M052R19

 $\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}}$ Γ_{91}/Γ

(0.3 GeV < m_{A^0} < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 8×10^{-5}	90	¹ LEES	11H BABR	$\Upsilon(2S) \rightarrow \gamma$ hadrons

¹ For a narrow scalar or pseudoscalar, A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

NODE=M052R06
NODE=M052R06
NODE=M052R06

NODE=M052R06;LINKAGE=LE

 $\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<8.3	90	¹ AUBERT	09Z BABR	$e^+e^- \rightarrow A^0 \rightarrow \gamma \mu^+ \mu^-$

¹ For a narrow scalar or pseudoscalar, A^0 , with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} range from 0.26–8.3 $\times 10^{-6}$.

NODE=M052R24
NODE=M052R24

NODE=M052R24;LINKAGE=AU

LEPTON FAMILY NUMBER (LF) VIOLATING MODES

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.2	90	LEES	10B BABR	$e^+e^- \rightarrow e^\pm \tau^\mp$

NODE=M052230

NODE=M052R04
NODE=M052R04

 $\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.3	90	LEES	10B BABR	$e^+e^- \rightarrow \mu^\pm \tau^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<14.4	95	LOVE	08A CLEO	$e^+e^- \rightarrow \mu^\pm \tau^\mp$
-------	----	------	----------	---------------------------------------

NODE=M052R20
NODE=M052R20

 $\Upsilon(2S)$ Cross-Particle Branching Ratios

NODE=M052240

 $B(\Upsilon(2S) \rightarrow \pi^+ \pi^-) \times B(\Upsilon(3S) \rightarrow \Upsilon(2S) X)$

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	COMMENT
1.78 ± 0.02 ± 0.11	906k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+ \pi^- X$

NODE=M052R05
NODE=M052R05

 $\Upsilon(2S)$ REFERENCES

NODE=M052

GAO	23	PR D108 112015	B.S. Gao <i>et al.</i>	(BELLE Collab.)	REFID=62519
SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVOU)	REFID=62012
FULSOM	18	PRL 121 232001	B.G. Fulsom <i>et al.</i>	(BELLE Collab.)	REFID=59535
JIA	18	PR D97 112004	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58949
JIA	17	PR D95 012001	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=57635
JIA	17A	PR D96 112002	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=58318
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
YANG	14	PR D90 112008	S.D. Yang <i>et al.</i>	(BELLE Collab.)	REFID=56345
SANDILYA	13	PRL 111 112001	S. Sandilya <i>et al.</i>	(BELLE Collab.)	REFID=55590
SHEN	13	PR D88 011102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55395
TAMPONI	13	PR D87 011104	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=54919
DOBBS	12	PRL 109 082001	S. Dobbs <i>et al.</i>		REFID=54288
DOBBS	12A	PR D86 052003	S. Dobbs <i>et al.</i>		REFID=54746
SHEN	12A	PR D86 031102	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=54314
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53938
WANG	11B	PR D84 071107	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=53939
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662

AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER	07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079
BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
EDWARDS	99	PR D59 032003	K.W. Edwards <i>et al.</i>	(CLEO Collab.)	REFID=46612
ALEXANDER	98	PR D58 052004	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=46329
BARU	96	PRPL 267 71	S.E. Baru <i>et al.</i>	(NOVO)	REFID=44651
KOBEL	92	ZPHY C53 193	M. Kobel <i>et al.</i>	(Crystal Ball Collab.)	REFID=41861
MASCHMANN	90	ZPHY C46 555	W.S. Maschmann <i>et al.</i>	(Crystal Ball Collab.)	REFID=41224
ALBRECHT	89	ZPHY C42 349	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40731
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding, World Scientific, Singapore					
JAKUBOWSKI	88	ZPHY C40 49	Z. Jakubowski <i>et al.</i>	(Crystal Ball Collab.) IGJPC	REFID=40742
ALBRECHT	87	ZPHY C35 283	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=40016
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
LURZ	87	ZPHY C36 383	B. Lurz <i>et al.</i>	(Crystal Ball Collab.)	REFID=40021
BARU	86B	ZPHY C32 622 (err.)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
ALBRECHT	85	ZPHY C28 45	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22334
ALBRECHT	85E	PL 160B 331	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=22288
GELPHMAN	85	PR D32 2893	D. Gelpman <i>et al.</i>	(Crystal Ball Collab.)	REFID=22336
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
Translated from YAF 41 733.					
NERNST	85	PL 54 2195	R. Nernst <i>et al.</i>	(Crystal Ball Collab.)	REFID=22289
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
BARBER	84	PL 135B 498	D.P. Barber <i>et al.</i>		REFID=22327;ERROR=9
BESSON	84	PR D30 1433	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22279
FONSECA	84	NP B242 31	V. Fonseca <i>et al.</i>	(CUSB Collab.)	REFID=22329
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
HAAS	84	PRL 52 799	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22287
HAAS	84B	PR D30 1996	J. Haas <i>et al.</i>	(CLEO Collab.)	REFID=22332
KLOPFEN...	83	PRL 51 160	C. Klopfenstein <i>et al.</i>	(CUSB Collab.)	REFID=22285
ALBRECHT	82	PL 116B 383	H. Albrecht <i>et al.</i>	(DESY, DORT, HEIDH+)	REFID=22270
NICZYPORUK	81B	PL 100B 95	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22319
NICZYPORUK	81C	PL 99B 169	B. Niczyporuk <i>et al.</i>	(LENA Collab.)	REFID=22318
BOCK	80	ZPHY C6 125	P. Bock <i>et al.</i>	(HEIDP, MPIM, DESY, HAMB)	REFID=22264

$\Upsilon_2(1D)$

$$J^G(J^{PC}) = 0^-(2^{--})$$

was $\Upsilon(1D)$

First observed by BONVICINI 04 in the decay to $\gamma\gamma \Upsilon(1S)$ and confirmed by DEL-AMO-SANCHEZ 10R in the decay to $\pi^+\pi^- \Upsilon(1S)$.

Data consistent with $J^P = 2^-$. The states with $J = 1$ and 3 also possibly seen, but need confirmation.

$\Upsilon_2(1D)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10163.7 ± 1.4 OUR AVERAGE				Error includes scale factor of 1.7.
10164.5 ± 0.8 ± 0.5		DEL-AMO-SA..10R	BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$
10161.1 ± 0.6 ± 1.6	38	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

$\Upsilon_2(1D)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma\gamma \Upsilon(1S)$	seen
$\Gamma_2 \quad \gamma\chi_{bJ}(1P)$	seen
$\Gamma_3 \quad \eta \Upsilon(1S)$	not seen
$\Gamma_4 \quad \pi^+\pi^- \Upsilon(1S)$	$(6.6 \pm 1.6) \times 10^{-3}$

$\Upsilon_2(1D)$ BRANCHING RATIOS

$\Gamma(\eta \Upsilon(1S))/\Gamma(\gamma\gamma \Upsilon(1S))$	Γ_3/Γ_1			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.25	90	BONVICINI 04	CLE3	$\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$

NODE=M177

NODE=M177

NODE=M177M

NODE=M177M

NODE=M177215;NODE=M177

DESIG=1;OUR EVAL;→ UNCHECKED ←
 DESIG=2;OUR EVAL;→ UNCHECKED ←
 DESIG=3;OUR EVAL;→ UNCHECKED ←
 DESIG=4

NODE=M177225

NODE=M177R01
 NODE=M177R01

$\Gamma(\pi^+\pi^-\Upsilon(1S))/\Gamma_{\text{total}}$ Γ_4/Γ VALUE (units 10^{-2})

DOCUMENT ID TECN COMMENT

 $0.66^{+0.15}_{-0.14} \pm 0.06$ ¹ DEL-AMO-SA...10R BABR $\Upsilon(3S) \rightarrow \gamma\gamma\pi^+\pi^-\ell^+\ell^-$ ¹ Using theoretical predictions for $B(\chi_{bJ}(2P) \rightarrow \gamma\Upsilon_2(1D))$.NODE=M177R03
NODE=M177R03

NODE=M177R03;LINKAGE=DE

 $\Gamma(\pi^+\pi^-\Upsilon(1S))/\Gamma(\gamma\gamma\Upsilon(1S))$ Γ_4/Γ_1

VALUE

CL%

DOCUMENT ID

TECN

COMMENT

<1.2

90

² BONVICINI 04 CLE3 $\Upsilon(3S) \rightarrow 4\gamma\ell^+\ell^-$ ² Assuming $J = 2$.NODE=M177R02
NODE=M177R02

NODE=M177R02;LINKAGE=BO

$\Upsilon_2(1D)$ REFERENCES

DEL-AMO-SA... 10R PR D82 111102
BONVICINI 04 PR D70 032001P. del Amo Sanchez *et al.*
G. Bonvicini *et al.*(BABAR Collab.)
(CLEO Collab.)

NODE=M177

REFID=53634
REFID=49759

NODE=M079

 $\chi_{b0}(2P)$ $I^G(J^{PC}) = 0^+(0^{++})$
 J needs confirmation.Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M079

$\chi_{b0}(2P)$ MASS

VALUE (MeV)

DOCUMENT ID

10232.5 ± 0.4 ± 0.5 OUR EVALUATION From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

NODE=M079M

NODE=M079M

→ UNCHECKED ←

 $m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

23.8 ± 1.7LEES 14M BABR $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ NODE=M079M2
NODE=M079M2

γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

121.9 ± 0.4 OUR EVALUATION Treating systematic errors as correlated**122.2 ± 0.5 OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

121.55 ± 0.16 ± 0.46

ARTUSO 05 CLEO $\Upsilon(3S) \rightarrow \gamma X$

123.0 ± 0.8

4959

¹ HEINTZ 92 CSB2 $e^+e^- \rightarrow \gamma X$

124.6 ± 1.4

17

² HEINTZ 92 CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

122.3 ± 0.3 ± 0.6

9903

MORRISON 91 CLE2 $e^+e^- \rightarrow \gamma X$

NODE=M079DM

NODE=M079DM

→ UNCHECKED ←


OCCUR=2

¹ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

NODE=M079DM;LINKAGE=A

² A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M079DM;LINKAGE=B



$\chi_{b0}(2P)$ DECAY MODES

NODE=M079215;NODE=M079

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\gamma \Upsilon(2S)$	(1.38±0.30) %	
Γ_2 $\gamma \Upsilon(1S)$	(3.8 ±1.7) × 10 ⁻³	
Γ_3 $D^0 X$	< 8.2 %	90%
Γ_4 $\pi^+ \pi^- K^+ K^- \pi^0$	< 3.4 × 10 ⁻⁵	90%
Γ_5 $2\pi^+ \pi^- K^- K_S^0$	< 5 × 10 ⁻⁵	90%
Γ_6 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 2.2 × 10 ⁻⁴	90%
Γ_7 $2\pi^+ 2\pi^- 2\pi^0$	< 2.4 × 10 ⁻⁴	90%
Γ_8 $2\pi^+ 2\pi^- K^+ K^-$	< 1.5 × 10 ⁻⁴	90%
Γ_9 $2\pi^+ 2\pi^- K^+ K^- \pi^0$	< 2.2 × 10 ⁻⁴	90%
Γ_{10} $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	< 1.1 × 10 ⁻³	90%
Γ_{11} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 7 × 10 ⁻⁴	90%
Γ_{12} $3\pi^+ 3\pi^-$	< 7 × 10 ⁻⁵	90%
Γ_{13} $3\pi^+ 3\pi^- 2\pi^0$	< 1.2 × 10 ⁻³	90%
Γ_{14} $3\pi^+ 3\pi^- K^+ K^-$	< 1.5 × 10 ⁻⁴	90%
Γ_{15} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	< 7 × 10 ⁻⁴	90%
Γ_{16} $4\pi^+ 4\pi^-$	< 1.7 × 10 ⁻⁴	90%
Γ_{17} $4\pi^+ 4\pi^- 2\pi^0$	< 6 × 10 ⁻⁴	90%

DESIG=2
DESIG=1
DESIG=3
DESIG=4
DESIG=5
DESIG=6
DESIG=7
DESIG=8
DESIG=9
DESIG=10
DESIG=11
DESIG=12
DESIG=13
DESIG=14
DESIG=15
DESIG=16
DESIG=17

$\chi_{b0}(2P)$ BRANCHING RATIOS

NODE=M079220

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{total}$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
1.38±0.30 OUR AVERAGE					
1.31±0.27 ^{+0.13} _{-0.12}		3,4 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$	
3.6 ±1.6 ±0.3		3,5 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<2.8	90	6 LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$	
<8.9	90	7 CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	

NODE=M079R2
NODE=M079R2

³ Assuming $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

⁴ LEES 14M reports $[\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{total}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] = (7.7 \pm 1.6) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91.

NODE=M079R2;LINKAGE=D
NODE=M079R2;LINKAGE=E

NODE=M079R2;LINKAGE=C

⁶ LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$.

NODE=M079R2;LINKAGE=LE

⁷ Using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) < 1.19 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P)\gamma) = 0.049$.

NODE=M079R2;LINKAGE=B

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ **Γ_2/Γ**

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
-----------	-----	-------------	------	---------

NODE=M079R1
NODE=M079R1**0.38±0.17 OUR AVERAGE**

0.36±0.17±0.03		8,9,10 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.9 ± 0.7 ± 0.1		9,11 HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2	90	12 LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
------	----	---------	----------	-------------------------------------

<2.5	90	13 CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
------	----	-------------	----------	---

⁸ LEES 14M quotes $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (2.1 \pm 1.0) \times 10^{-4}$ combining the results from $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ samples with and without photon conversions.

NODE=M079R1;LINKAGE=D

⁹ Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

NODE=M079R1;LINKAGE=E

¹⁰ LEES 14M reports $[\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] = (2.1 \pm 1.0) \times 10^{-4}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M079R1;LINKAGE=F

¹¹ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91.

NODE=M079R1;LINKAGE=C

¹² LEES 11J quotes a central value of $\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$.

NODE=M079R1;LINKAGE=LE

¹³ Using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) < 0.63 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P)\gamma) = 0.049$.

NODE=M079R1;LINKAGE=B

 $\Gamma(D^0 X)/\Gamma_{\text{total}}$ **Γ_3/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M079R01
NODE=M079R01

<8.2 × 10 ⁻²	90	14,15 BRIERE	08 CLEO	$\Upsilon(3S) \rightarrow \gamma D^0 X$
-------------------------	----	--------------	---------	---

¹⁴ For $p_{D^0} > 2.5$ GeV/c.

NODE=M079R01;LINKAGE=BR

¹⁵ The authors also present their result as $(4.1 \pm 3.0 \pm 0.4) \times 10^{-2}$.

NODE=M079R01;LINKAGE=RI

 $\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ **Γ_4/Γ**

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M079R02
NODE=M079R02

<0.34	90	16 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$
-------	----	----------	----------	---

¹⁶ ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 2 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R02;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ **Γ_5/Γ**

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M079R03
NODE=M079R03

<0.5	90	17 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$
------	----	----------	----------	--

¹⁷ ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 3 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R03;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ **Γ_6/Γ**

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M079R04
NODE=M079R04

<2.2	90	18 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$
------	----	----------	----------	---

¹⁸ ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ **Γ_7/Γ**

VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M079R05
NODE=M079R05

<2.4	90	19 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$
------	----	----------	----------	--

¹⁹ ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.

NODE=M079R05;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	20 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$
20 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 9 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R06
NODE=M079R06

NODE=M079R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<2.2	90	21 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$
21 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 13 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R07
NODE=M079R07

NODE=M079R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<11	90	22 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$
22 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 63 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R08
NODE=M079R08

NODE=M079R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
23 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 39 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R09
NODE=M079R09

NODE=M079R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.7	90	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
24 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 4 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R10
NODE=M079R10

NODE=M079R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<12	90	25 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
25 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 72 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R11
NODE=M079R11

NODE=M079R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	26 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
26 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 9 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R12
NODE=M079R12

NODE=M079R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	27 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
27 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 43 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R13
NODE=M079R13

NODE=M079R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	28 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
28 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] < 10 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R14
NODE=M079R14

NODE=M079R14;LINKAGE=AS

$$\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$$

$$\Gamma_{17}/\Gamma$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<6	90	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
29 ASNER 08A reports $[\Gamma(\chi_{b0}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))]$ < 38×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = 5.9 \times 10^{-2}$.				

NODE=M079R15
NODE=M079R15

NODE=M079R15;LINKAGE=AS

$$\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_2/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<8.2	90	30 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
30 LEES 11J quotes a central value of $(\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)))/\Gamma_{\text{total}} = (3.9 \pm 2.2^{+1.2}_{-0.6}) \times 10^{-4}$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) < 1.2\%$ using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.				

NODE=M079B01
NODE=M079B01

NODE=M079B01;LINKAGE=LE

$$B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
1.4 ± 0.9 OUR AVERAGE			

$1.7^{+1.5+0.1}_{-1.4-1.2}$	31 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$1.3 \pm 1.0 \pm 0.3$	32 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
31 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.			
32 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.05 \pm 0.04 \pm 0.01)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$.			

NODE=M079A02
NODE=M079A02

NODE=M079A02;LINKAGE=A
NODE=M079A02;LINKAGE=K

$$[B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
1.71 ± 0.80	33 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$

33 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ without converted photons.

NODE=M079A00
NODE=M079A00

NODE=M079A00;LINKAGE=A

$$\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_1/\Gamma \times \Gamma_{22}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	34 LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
34 LEES 11J quotes a central value of $(\Gamma(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)))/\Gamma_{\text{total}} = (-0.3 \pm 0.2^{+0.5}_{-0.4})\%$ and derives a 90% CL upper limit of $B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) < 2.8\%$ using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) = (5.9 \pm 0.6)\%$.				

NODE=M079B02
NODE=M079B02

NODE=M079B02;LINKAGE=LE

$$B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.4 ± 1.6 OUR AVERAGE			

$6.6^{+4.9+2.0}_{-4.0-0.3}$	35 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$4.0 \pm 1.7 \pm 0.3$	36 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
35 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.			
36 Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P)) \times B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) = (0.28 \pm 0.12 \pm 0.03)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$.			

NODE=M079A03
NODE=M079A03

NODE=M079A03;LINKAGE=A
NODE=M079A03;LINKAGE=B

$$[B(\chi_{b0}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
3.31 ± 0.56	37 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$

37 From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ without converted photons.

NODE=M079A01
NODE=M079A01

NODE=M079A01;LINKAGE=A

$\chi_{b0}(2P)$ REFERENCES

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

NODE=M079

REFID=56343
REFID=53936
REFID=52574
REFID=52577
REFID=50454
REFID=43177
REFID=43604
REFID=41580
REFID=41634
REFID=41586

$\chi_{b1}(2P)$

$I^G(J^{PC}) = 0^+(1^{++})$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M080

NODE=M080

$\chi_{b1}(2P)$ MASS

NODE=M080M

VALUE (MeV) DOCUMENT ID
10255.46 ± 0.22 ± 0.50 OUR EVALUATION From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

NODE=M080M
 → UNCHECKED ←

$m_{\chi_{b1}(2P)} - m_{\chi_{b0}(2P)}$

NODE=M080M2

VALUE (MeV) DOCUMENT ID TECN COMMENT
23.5 ± 0.7 ± 0.7 ¹ HEINTZ 92 CSB2 $e^+e^- \rightarrow \gamma X, \ell^+\ell^-\gamma\gamma$

NODE=M080M2

¹ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

NODE=M080M2;LINKAGE=A

γ ENERGY IN $\Upsilon(3S)$ DECAY

NODE=M080DM

VALUE (MeV) EVTS DOCUMENT ID TECN COMMENT
99.26 ± 0.22 OUR EVALUATION Treating systematic errors as correlated

NODE=M080DM
 → UNCHECKED ←

99.53 ± 0.23 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.

99.15 ± 0.07 ± 0.25		ARTUSO	05	CLEO	$\Upsilon(3S) \rightarrow \gamma X$
99 ± 1	169	CRAWFORD	92B	CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
100.1 ± 0.4	11147	² HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma X$
100.2 ± 0.5	223	³ HEINTZ	92	CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
99.5 ± 0.1 ± 0.5	25759	MORRISON	91	CLE2	$e^+e^- \rightarrow \gamma X$


OCCUR=2

² A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

NODE=M080DM;LINKAGE=A

³ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M080DM;LINKAGE=B



$\chi_{b1}(2P)$ DECAY MODES

NODE=M080215;NODE=M080

Mode	Fraction (Γ_i/Γ)	
Γ_1 $\omega \Upsilon(1S)$	$(1.63^{+0.40}_{-0.34})\%$	DESIG=3
Γ_2 $\gamma \Upsilon(2S)$	$(18.1 \pm 1.9)\%$	DESIG=2
Γ_3 $\gamma \Upsilon(1S)$	$(9.9 \pm 1.0)\%$	DESIG=1
Γ_4 $\pi\pi\chi_{b1}(1P)$	$(9.1 \pm 1.3) \times 10^{-3}$	DESIG=4
Γ_5 $D^0 X$	$(8.8 \pm 1.7)\%$	DESIG=5
Γ_6 $\pi^+\pi^-K^+K^-\pi^0$	$(3.1 \pm 1.0) \times 10^{-4}$	DESIG=6
Γ_7 $2\pi^+\pi^-K^-K_S^0$	$(1.1 \pm 0.5) \times 10^{-4}$	DESIG=7
Γ_8 $2\pi^+\pi^-K^-K_S^0 2\pi^0$	$(7.7 \pm 3.2) \times 10^{-4}$	DESIG=8
Γ_9 $2\pi^+ 2\pi^- 2\pi^0$	$(5.9 \pm 2.0) \times 10^{-4}$	DESIG=9
Γ_{10} $2\pi^+ 2\pi^- K^+ K^-$	$(10 \pm 4) \times 10^{-5}$	DESIG=10
Γ_{11} $2\pi^+ 2\pi^- K^+ K^- \pi^0$	$(5.5 \pm 1.8) \times 10^{-4}$	DESIG=11
Γ_{12} $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	$(10 \pm 4) \times 10^{-4}$	DESIG=12
Γ_{13} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	$(6.7 \pm 2.6) \times 10^{-4}$	DESIG=13
Γ_{14} $3\pi^+ 3\pi^-$	$(1.2 \pm 0.4) \times 10^{-4}$	DESIG=14
Γ_{15} $3\pi^+ 3\pi^- 2\pi^0$	$(1.2 \pm 0.4) \times 10^{-3}$	DESIG=15
Γ_{16} $3\pi^+ 3\pi^- K^+ K^-$	$(2.0 \pm 0.8) \times 10^{-4}$	DESIG=16
Γ_{17} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	$(6.1 \pm 2.2) \times 10^{-4}$	DESIG=17
Γ_{18} $4\pi^+ 4\pi^-$	$(1.7 \pm 0.6) \times 10^{-4}$	DESIG=18
Γ_{19} $4\pi^+ 4\pi^- 2\pi^0$	$(1.9 \pm 0.7) \times 10^{-3}$	DESIG=19

 $\chi_{b1}(2P)$ BRANCHING RATIOS

NODE=M080220

 $\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M080R3
NODE=M080R3

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.63^{+0.35+0.16}_{-0.31-0.15}$	$32.6^{+6.9}_{-6.1}$	⁴ CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma\omega \Upsilon(1S)$

⁴Using $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (11.3 \pm 0.6)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = 2$
 $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 2 (2.48 \pm 0.06)\%$.

NODE=M080R3;LINKAGE=CR

 $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M080R2
NODE=M080R2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.181 ± 0.019 OUR AVERAGE				
$0.211 \pm 0.017 \pm 0.019$	^{5,6,7}	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
$0.190 \pm 0.018 \pm 0.017$	⁸	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$
$0.206 \pm 0.035 \pm 0.019$	^{5,9}	CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
$0.132 \pm 0.018 \pm 0.012$	^{5,10}	HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

⁵Assuming $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$.

⁶LEES 14M quotes $\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))/\Gamma_{\text{total}} = (2.66 \pm 0.22)\%$ combining the results from $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ samples with and without photon conversions.

⁷LEES 14M reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (2.66 \pm 0.22) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁸LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (2.4 \pm 0.1 \pm 0.2) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁹CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2$
 $B(\Upsilon(2S) \rightarrow \ell^+\ell^-) = (10.23 \pm 1.20 \pm 1.26) 10^{-4}$.

¹⁰Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$. Supersedes HEINTZ 91.

NODE=M080R2;LINKAGE=D
NODE=M080R2;LINKAGE=E

NODE=M080R2;LINKAGE=F

NODE=M080R2;LINKAGE=LE

NODE=M080R2;LINKAGE=B

NODE=M080R2;LINKAGE=C

$\Gamma(\gamma\Upsilon(1S))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.099±0.010 OUR AVERAGE				
0.107±0.006±0.010	11,12,13	LEES	14M	BABR $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
0.098±0.005±0.009	15k	LEES	11J	BABR $\Upsilon(3S) \rightarrow X\gamma$
0.103±0.023±0.009	11,15	CRAWFORD	92B	CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
0.075±0.010±0.007	11,16	HEINTZ	92	CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M080R1
 NODE=M080R1

¹¹ Assuming $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

¹² LEES 14M quotes $\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))/\Gamma_{\text{total}} = (13.48 \pm 0.72) \times 10^{-3}$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with and without converted photons.

NODE=M080R1;LINKAGE=D
 NODE=M080R1;LINKAGE=E

¹³ LEES 14M reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (13.48 \pm 0.72) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R1;LINKAGE=G

¹⁴ LEES 11J reports $[\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (12.4 \pm 0.3 \pm 0.6) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R1;LINKAGE=LE

¹⁵ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \ell^+\ell^-) = (6.47 \pm 1.12 \pm 0.82) 10^{-4}$.

NODE=M080R1;LINKAGE=B

¹⁶ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$. Supersedes HEINTZ 91.

NODE=M080R1;LINKAGE=F

 $\Gamma(\pi\pi\chi_{b1}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
9.1±1.3 OUR AVERAGE				
9.2±1.1±0.8	31k	17 LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$
8.6±2.3±2.1		18 CAWLFIELD	06	CLE3 $\Upsilon(3S) \rightarrow 2(\gamma\pi\ell)$

NODE=M080R4
 NODE=M080R4

¹⁷ LEES 11C measures $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) \times B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) = (1.16 \pm 0.07 \pm 0.12) \times 10^{-3}$. We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X) = B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)\gamma) = (12.6 \pm 1.2) \times 10^{-2}$.

NODE=M080R4;LINKAGE=LE

¹⁸ CAWLFIELD 06 quote $\Gamma(\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV assuming l-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

NODE=M080R4;LINKAGE=CA

 $\Gamma(D^0X)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
8.8±1.5±0.8				
	2243	19 BRIERE	08	CLEO $\Upsilon(3S) \rightarrow \gamma D^0X$

NODE=M080R01
 NODE=M080R01

¹⁹ For $p_{D^0} > 2.5$ GeV/c.

NODE=M080R01;LINKAGE=BR

 $\Gamma(\pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±1.0±0.3				
	30	20 ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma\pi^+\pi^-K^+K^-\pi^0$

NODE=M080R02
 NODE=M080R02

²⁰ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow \pi^+\pi^-K^+K^-\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (39 \pm 8 \pm 9) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R02;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.5±0.1				
	10	21 ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-K_S^0$

NODE=M080R03
 NODE=M080R03

²¹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (14 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R03;LINKAGE=AS

 $\Gamma(2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.7±3.1±0.7				
	15	22 ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+\pi^-K^-2\pi^0$

NODE=M080R04
 NODE=M080R04

²² ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+\pi^-K^-K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P))] = (97 \pm 30 \pm 26) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R04;LINKAGE=AS

$\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.9±2.0±0.5	36	23 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

NODE=M080R05
 NODE=M080R05

²³ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (74 \pm 16 \pm 19) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.4±0.1	12	24 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

NODE=M080R06
 NODE=M080R06

²⁴ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
5.5±1.7±0.5	38	25 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

NODE=M080R07
 NODE=M080R07

²⁵ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (69 \pm 13 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9.6±3.5±0.9	27	26 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

NODE=M080R08
 NODE=M080R08

²⁶ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (121 \pm 29 \pm 33) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R08;LINKAGE=AS

 $\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7±2.5±0.6	17	27 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$

NODE=M080R09
 NODE=M080R09

²⁷ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (85 \pm 23 \pm 22) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.2±0.4±0.1	18	28 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$

NODE=M080R10
 NODE=M080R10

²⁸ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (15 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
12±4±1	44	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$

NODE=M080R11
 NODE=M080R11

²⁹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (150 \pm 30 \pm 40) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.0±0.7±0.2	16	30 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$

NODE=M080R12
 NODE=M080R12

³⁰ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$
 $= (25 \pm 7 \pm 6) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))$
 $= (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M080R12;LINKAGE=AS

$\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.1±2.1±0.6	25	³¹ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
³¹ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(77 \pm 17 \pm 21) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M080R13
NODE=M080R13

NODE=M080R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.7±0.6±0.2	16	³² ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
³² ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(22 \pm 6 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M080R14
NODE=M080R14

NODE=M080R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19±7±2	41	³³ ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
³³ ASNER 08A reports $[\Gamma(\chi_{b1}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$ = $(241 \pm 47 \pm 72) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (12.6 \pm 1.2) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M080R15
NODE=M080R15

NODE=M080R15;LINKAGE=AS

 $\chi_{b1}(2P)$ Cross-Particle Branching Ratios

NODE=M080230

$$\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))/\Gamma_{\text{total}}$$

$$\Gamma_3/\Gamma \times \Gamma_{21}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$$


VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
12.4±0.3±0.6	15k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M080B01
NODE=M080B01 $B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$ NODE=M080A00
NODE=M080A00

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9 ±0.4 OUR AVERAGE		Error includes scale factor of 1.9. See the ideogram below.		
$3.52^{+0.28+0.17}_{-0.27-0.18}$		³⁴ LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
$3.24 \pm 0.56 \pm 0.41$	58	³⁵ CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
$2.34 \pm 0.28 \pm 0.15$		³⁶ HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
³⁴ From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.				
³⁵ CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS))$ $B(\Upsilon(nS) \rightarrow \ell^+ \ell^-)$.				
³⁶ Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.91 \pm 0.11 \pm 0.06)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$.				

NODE=M080A00;LINKAGE=A
NODE=M080A00;LINKAGE=C

NODE=M080A00;LINKAGE=B



$$B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+ \ell^-)$$

(units 10^{-4})

$$\frac{\Gamma(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) / \Gamma_{total} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) / \Gamma_{total}}{\Gamma_2 / \Gamma \times \Gamma_{21}^{\Upsilon(3S)} / \Gamma \Upsilon(3S)}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±0.1±0.2	4.3k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M080B02
NODE=M080B02

$$B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$$

3.8 ±0.6 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.


VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.95 ^{+0.75} _{-0.70} +1.01 -0.24		37 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
5.12±0.60±0.63	111	38 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$
3.30±0.33±0.20		39 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma \gamma \ell^+ \ell^-$

NODE=M080A01
NODE=M080A01

- ³⁷ From a sample of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with one converted photon.
- ³⁸ CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+ \ell^-)$.
- ³⁹ Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) = (2.29 \pm 0.23 \pm 0.21) \%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10) \%$.

NODE=M080A01;LINKAGE=A
NODE=M080A01;LINKAGE=C

NODE=M080A01;LINKAGE=B



$$B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+ \ell^-)$$

(units 10^{-4})

$B(\chi_{b1}(2P) \rightarrow \chi_{b1}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b1}(2P)X)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.16 \pm 0.07 \pm 0.12$	31k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M080R16
 NODE=M080R16

 $B(\chi_{b2}(2P) \rightarrow \rho X + \bar{\rho} X)/B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.109 \pm 0.007 \pm 0.040$	BRIERE	07 CLEO	$\Upsilon(3S) \rightarrow \gamma\chi_{b,J}(2P)$

NODE=M080R20
 NODE=M080R20

 $B(\chi_{b0}(2P) \rightarrow \rho X + \bar{\rho} X)/B(\chi_{b1}(2P) \rightarrow \rho X + \bar{\rho} X)$

VALUE	DOCUMENT ID	TECN	COMMENT
$1.082 \pm 0.025 \pm 0.060$	BRIERE	07 CLEO	$\Upsilon(3S) \rightarrow \gamma\chi_{b,J}(2P)$

NODE=M080R21
 NODE=M080R21

 $\chi_{b1}(2P)$ REFERENCES

LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)
BRIERE	07	PR D76 012005	R.A. Briere <i>et al.</i>	(CLEO Collab.)
CRAWFIELD	06	PR D73 012003	C. Cawfield <i>et al.</i>	(CLEO Collab.)
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)
CRONIN-HENNESSY	04	PRL 92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)

NODE=M080

REFID=56343
 REFID=16775
 REFID=53936
 REFID=52574
 REFID=52577
 REFID=51887
 REFID=50997
 REFID=50454
 REFID=49766
 REFID=43177
 REFID=43604
 REFID=41580
 REFID=41634
 REFID=41586

 $h_b(2P)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Quantum numbers are quark model predictions. $C = -$ established by $\eta_b\gamma$ decay.

NODE=M205

NODE=M205

 $h_b(2P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$10259.8 \pm 0.5 \pm 1.1$	90k	¹ MIZUK	12 BELL	$e^+e^- \rightarrow \pi^+\pi^-$ hadrons
$10259.8 \pm 0.6^{+1.4}_{-1.0}$	83.9k	² ADACHI	12 BELL	$10.86 e^+e^- \rightarrow \pi^+\pi^-$ MM

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹Observed with 9 standard deviations significance.

²Superseded by MIZUK 12.

NODE=M205M

NODE=M205M

NODE=M205M;LINKAGE=A
 NODE=M205M;LINKAGE=AD

 $h_b(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	not seen
Γ_2 $\eta_b(1S)\gamma$	$(22 \pm 5) \%$
Γ_3 $\eta_b(2S)\gamma$	$(48 \pm 13) \%$

NODE=M205215;NODE=M205

DESIG=1

DESIG=2

DESIG=3

 $h_b(2P)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	Γ_1/Γ
VALUE EVTS DOCUMENT ID TECN COMMENT	
not seen 83.9k ADACHI 12 BELL 10.86 $e^+e^- \rightarrow \pi^+\pi^-$ MM	

NODE=M205225

NODE=M205R01
 NODE=M205R01

$\Gamma(\eta_b(1S)\gamma)/\Gamma_{\text{total}}$	Γ_2/Γ
VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT	
$22.3 \pm 3.8^{+3.1}_{-3.3}$ 10k MIZUK 12 BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons	

NODE=M205R02
 NODE=M205R02

$\Gamma(\eta_b(2S)\gamma)/\Gamma_{\text{total}}$	Γ_3/Γ
VALUE (units 10^{-2}) EVTS DOCUMENT ID TECN COMMENT	
$47.5 \pm 10.5^{+6.8}_{-7.7}$ 26k MIZUK 12 BELL $e^+e^- \rightarrow (\gamma)\pi^+\pi^-$ hadrons	

NODE=M205R03
 NODE=M205R03

 $h_b(2P)$ REFERENCES

ADACHI	12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)
MIZUK	12	PRL 109 232002	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M205

REFID=53962
 REFID=54718

$\chi_{b2}(2P)$

$$J^G(J^{PC}) = 0^+(2^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

NODE=M081

NODE=M081

 $\chi_{b2}(2P)$ MASS

NODE=M081M

VALUE (MeV)

DOCUMENT ID

10268.65 ± 0.22 ± 0.50 OUR EVALUATION From γ energy below, using $\Upsilon(3S)$ mass = 10355.2 ± 0.5 MeV

NODE=M081M

→ UNCHECKED ←

 $m_{\chi_{b2}(2P)} - m_{\chi_{b1}(2P)}$

NODE=M081M2

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

13.10 ± 0.24 OUR AVERAGE

NODE=M081M2

12.3 ± 2.6 ± 0.6

¹ AAIJ

14BG LHCB

 $pp \rightarrow \gamma \mu^+ \mu^- X$

13.04 ± 0.26

LEES

14M BABR

 $\Upsilon(3S) \rightarrow \gamma \mu^+ \mu^-$

13.5 ± 0.4 ± 0.5

² HEINTZ

92 CSB2

 $e^+ e^- \rightarrow \gamma X, \ell^+ \ell^- \gamma \gamma$ ¹ From the $\chi_{bj}(2P) \rightarrow \Upsilon(1S) \gamma$ transition.² From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

NODE=M081M2;LINKAGE=B

NODE=M081M2;LINKAGE=A

 γ ENERGY IN $\Upsilon(3S)$ DECAY

NODE=M081DM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

86.19 ± 0.22 OUR EVALUATION

Treating systematic errors as correlated

NODE=M081DM

86.40 ± 0.18 OUR AVERAGE

→ UNCHECKED ←

86.04 ± 0.06 ± 0.27

ARTUSO

05

CLEO

 $\Upsilon(3S) \rightarrow \gamma X$

86 ± 1

101

CRAWFORD

92B

CLE2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

86.7 ± 0.4

10319

³ HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \gamma X$

86.9 ± 0.4

157

⁴ HEINTZ

92

CSB2

 $e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

86.4 ± 0.1 ± 0.4

30741

MORRISON

91

CLE2

 $e^+ e^- \rightarrow \gamma X$

OCCUR=2

³ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

NODE=M081DM;LINKAGE=A

⁴ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

NODE=M081DM;LINKAGE=B

 $\chi_{b2}(2P)$ DECAY MODES

NODE=M081215;NODE=M081

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\omega \Upsilon(1S)$	(1.10 ^{+0.34} _{-0.30}) %	DESIG=3
Γ_2 $\gamma \Upsilon(2S)$	(8.9 ± 1.2) %	DESIG=2
Γ_3 $\gamma \Upsilon(1S)$	(6.6 ± 0.8) %	DESIG=1
Γ_4 $\pi \pi \chi_{b2}(1P)$	(5.1 ± 0.9) × 10 ⁻³	DESIG=4
Γ_5 $D^0 X$	< 2.4 %	90% DESIG=5
Γ_6 $\pi^+ \pi^- K^+ K^- \pi^0$	< 1.1 × 10 ⁻⁴	90% DESIG=6
Γ_7 $2\pi^+ \pi^- K^- K_S^0$	< 9 × 10 ⁻⁵	90% DESIG=7
Γ_8 $2\pi^+ \pi^- K^- K_S^0 2\pi^0$	< 7 × 10 ⁻⁴	90% DESIG=8
Γ_9 $2\pi^+ 2\pi^- 2\pi^0$	(3.9 ± 1.6) × 10 ⁻⁴	DESIG=9
Γ_{10} $2\pi^+ 2\pi^- K^+ K^-$	(9 ± 4) × 10 ⁻⁵	DESIG=10
Γ_{11} $2\pi^+ 2\pi^- K^+ K^- \pi^0$	(2.4 ± 1.1) × 10 ⁻⁴	DESIG=11
Γ_{12} $2\pi^+ 2\pi^- K^+ K^- 2\pi^0$	(4.7 ± 2.3) × 10 ⁻⁴	DESIG=12
Γ_{13} $3\pi^+ 2\pi^- K^- K_S^0 \pi^0$	< 4 × 10 ⁻⁴	90% DESIG=13
Γ_{14} $3\pi^+ 3\pi^-$	(9 ± 4) × 10 ⁻⁵	DESIG=14
Γ_{15} $3\pi^+ 3\pi^- 2\pi^0$	(1.2 ± 0.4) × 10 ⁻³	DESIG=15
Γ_{16} $3\pi^+ 3\pi^- K^+ K^-$	(1.4 ± 0.7) × 10 ⁻⁴	DESIG=16
Γ_{17} $3\pi^+ 3\pi^- K^+ K^- \pi^0$	(4.2 ± 1.7) × 10 ⁻⁴	DESIG=17
Γ_{18} $4\pi^+ 4\pi^-$	(9 ± 5) × 10 ⁻⁵	DESIG=18
Γ_{19} $4\pi^+ 4\pi^- 2\pi^0$	(1.3 ± 0.5) × 10 ⁻³	DESIG=19

$\chi_{b2}(2P)$ BRANCHING RATIOS

NODE=M081220

 $\Gamma(\omega \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10^{+0.32+0.11}_{-0.28-0.10}	20.1 ^{+5.8} _{-5.1}	5 CRONIN-HEN..04	CLE3	$\Upsilon(3S) \rightarrow \gamma \omega \Upsilon(1S)$

NODE=M081R3
NODE=M081R3

⁵ Using $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.4 \pm 0.8)\%$ and $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = 2$
 $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 2 (2.48 \pm 0.06)\%$.

NODE=M081R3;LINKAGE=CR

 $\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.089\pm0.012 OUR AVERAGE				
0.085 \pm 0.010 \pm 0.010	6,7,8	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.084 \pm 0.011 \pm 0.010	2.5k	⁹ LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
0.096 \pm 0.022 \pm 0.012	7,10	CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
0.106 \pm 0.016 \pm 0.013	7,11	HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

NODE=M081R2
NODE=M081R2

⁶ LEES 14M quotes $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$
 $= (1.12 \pm 0.13)\%$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$ with and
without converted photons.

NODE=M081R2;LINKAGE=D

⁷ Assuming $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

NODE=M081R2;LINKAGE=E

⁸ LEES 14M reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$
 $(1.12 \pm 0.13) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$
 $(13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is
the systematic error from using our best value.

NODE=M081R2;LINKAGE=F

⁹ LEES 11J reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$
 $(1.1 \pm 0.1 \pm 0.1) \times 10^{-2}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$
 $(13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error
is the systematic error from using our best value.

NODE=M081R2;LINKAGE=LE

¹⁰ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times 2$
 $B(\Upsilon(2S) \rightarrow \ell^+ \ell^-) = (4.98 \pm 0.94 \pm 0.62) 10^{-4}$.

NODE=M081R2;LINKAGE=B

¹¹ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow$
 $\gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18) \%$ using $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.44 \pm 0.10)\%$.
Supersedes HEINTZ 91.

NODE=M081R2;LINKAGE=C

 $\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.066\pm0.008 OUR AVERAGE				
0.061 \pm 0.004 \pm 0.007	12,13,14	LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
0.070 \pm 0.004 \pm 0.008	11k	¹⁵ LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
0.077 \pm 0.018 \pm 0.009	13,16	CRAWFORD	92B CLE2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$
0.061 \pm 0.009 \pm 0.007	13,17	HEINTZ	92 CSB2	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$

NODE=M081R1
NODE=M081R1

¹² LEES 14M quotes $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$
 $= (8.03 \pm 0.50) \times 10^{-3}$ combining the results from samples of $\Upsilon(3S) \rightarrow \gamma \gamma \mu^+ \mu^-$
with and without converted photons.

NODE=M081R1;LINKAGE=D

¹³ Assuming $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$.

NODE=M081R1;LINKAGE=E

¹⁴ LEES 14M reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$
 $(8.03 \pm 0.50) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$
 $(13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is
the systematic error from using our best value.

NODE=M081R1;LINKAGE=F

¹⁵ LEES 11J reports $[\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] =$
 $(9.2 \pm 0.3 \pm 0.4) \times 10^{-3}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) =$
 $(13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error
is the systematic error from using our best value.

NODE=M081R1;LINKAGE=LE

¹⁶ CRAWFORD 92B quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times 2$
 $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (5.03 \pm 0.94 \pm 0.63) 10^{-4}$.

NODE=M081R1;LINKAGE=B

¹⁷ Recalculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow$
 $\gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.57 \pm 0.05)\%$.
Supersedes HEINTZ 91.

NODE=M081R1;LINKAGE=C

 $\Gamma(\pi \pi \chi_{b2}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
5.1\pm0.9 OUR AVERAGE				
4.9 \pm 0.7 \pm 0.6	17k	¹⁸ LEES	11C BABR	$e^+ e^- \rightarrow \pi^+ \pi^- X$
6.0 \pm 1.6 \pm 1.4		¹⁹ CAWLFIELD	06 CLE3	$\Upsilon(3S) \rightarrow 2(\gamma \pi \ell)$

NODE=M081R4
NODE=M081R4

¹⁸ $(0.64 \pm 0.05 \pm 0.08) \times 10^{-3}$. We derive the value assuming $B(\Upsilon(3S) \rightarrow \chi_{b2}(2P) X)$
 $= B(\Upsilon(3S) \rightarrow \chi_{b2}(2P) \gamma) = (13.1 \pm 1.6) \times 10^{-2}$.

NODE=M081R4;LINKAGE=LE

¹⁹ CAWLFIELD 06 quote $\Gamma(\chi_{b2}(2P) \rightarrow \pi \pi \chi_{b1}(1P)) = 0.83 \pm 0.22 \pm 0.08 \pm 0.19$ keV
assuming l-spin conservation, no D-wave contribution, $\Gamma(\chi_{b1}(2P)) = 96 \pm 16$ keV, and
 $\Gamma(\chi_{b2}(2P)) = 138 \pm 19$ keV.

NODE=M081R4;LINKAGE=CA

$\Gamma(D^0 X)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.4 \times 10^{-2}$	90	20,21 BRIERE	08	CLEO $\Upsilon(3S) \rightarrow \gamma D^0 X$

NODE=M081R01
 NODE=M081R01

²⁰ For $p_{D^0} > 2.5$ GeV/c.

²¹ The authors also present their result as $(0.2 \pm 1.4 \pm 0.1) \times 10^{-2}$.

NODE=M081R01;LINKAGE=BR
 NODE=M081R01;LINKAGE=RI

 $\Gamma(\pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<1.1	90	²² ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma \pi^+ \pi^- K^+ K^- \pi^0$

NODE=M081R02
 NODE=M081R02

²² ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow \pi^+ \pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 14 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

NODE=M081R02;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<0.9	90	²³ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- K_S^0$

NODE=M081R03
 NODE=M081R03

²³ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 12 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

NODE=M081R03;LINKAGE=AS

 $\Gamma(2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	²⁴ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ \pi^- K^- 2\pi^0$

NODE=M081R04
 NODE=M081R04

²⁴ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ \pi^- K^- K_S^0 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] < 87 \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.

NODE=M081R04;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$3.9 \pm 1.6 \pm 0.5$	23	²⁵ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- 2\pi^0$

NODE=M081R05
 NODE=M081R05

²⁵ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (51 \pm 16 \pm 13) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R05;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.9 \pm 0.4 \pm 0.1$	11	²⁶ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^-$

NODE=M081R06
 NODE=M081R06

²⁶ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R06;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$2.4 \pm 1.0 \pm 0.3$	16	²⁷ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- \pi^0$

NODE=M081R07
 NODE=M081R07

²⁷ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (32 \pm 11 \pm 8) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R07;LINKAGE=AS

 $\Gamma(2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.7 \pm 2.2 \pm 0.6$	14	²⁸ ASNER	08A	CLEO $\Upsilon(3S) \rightarrow \gamma 2\pi^+ 2\pi^- K^+ K^- 2\pi^0$

NODE=M081R08
 NODE=M081R08

²⁸ ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 2\pi^+ 2\pi^- K^+ K^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] = (62 \pm 23 \pm 17) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M081R08;LINKAGE=AS

$\Gamma(3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<4	90	29 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 2\pi^- K^- K_S^0 \pi^0$
29 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 2\pi^- K^- K_S^0 \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ < 58×10^{-6} which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 13.1 \times 10^{-2}$.				

NODE=M081R09
NODE=M081R09

NODE=M081R09;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^-)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	14	30 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^-$
30 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ = $(12 \pm 4 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R10
NODE=M081R10

NODE=M081R10;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
12±4±1	45	31 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- 2\pi^0$
31 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ = $(159 \pm 33 \pm 43) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R11
NODE=M081R11

NODE=M081R11;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
1.4±0.7±0.2	12	32 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^-$
32 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ = $(19 \pm 7 \pm 5) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R12
NODE=M081R12

NODE=M081R12;LINKAGE=AS

 $\Gamma(3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
4.2±1.7±0.5	16	33 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 3\pi^+ 3\pi^- K^+ K^- \pi^0$
33 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 3\pi^+ 3\pi^- K^+ K^- \pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ = $(55 \pm 16 \pm 15) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R13
NODE=M081R13

NODE=M081R13;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
0.9±0.4±0.1	9	34 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^-$
34 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ = $(12 \pm 5 \pm 3) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R14
NODE=M081R14

NODE=M081R14;LINKAGE=AS

 $\Gamma(4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
13±5±2	27	35 ASNER	08A CLEO	$\Upsilon(3S) \rightarrow \gamma 4\pi^+ 4\pi^- 2\pi^0$
35 ASNER 08A reports $[\Gamma(\chi_{b2}(2P) \rightarrow 4\pi^+ 4\pi^- 2\pi^0)/\Gamma_{\text{total}}] \times [B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))]$ = $(165 \pm 46 \pm 50) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (13.1 \pm 1.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M081R15
NODE=M081R15

NODE=M081R15;LINKAGE=AS

 $\chi_{b2}(2P)$ Cross-Particle Branching Ratios

NODE=M081230

 $\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S))/\Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))/\Gamma_{\text{total}}$
 $\Gamma_3/\Gamma \times \Gamma_{20}^{\Upsilon(3S)}/\Gamma \Upsilon(3S)$

VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT
9.2±0.3±0.4	11k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M081B01
NODE=M081B01

$$\Gamma(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) / \Gamma_{\text{total}} \times \Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) / \Gamma_{\text{total}}$$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.1±0.1±0.1	2.5k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M081B02
NODE=M081B02

$$B(\chi_{b2}(2P) \rightarrow \chi_{b2}(1P)\pi^+\pi^-) \times B(\Upsilon(3S) \rightarrow \chi_{b2}(2P)X)$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.64±0.05±0.08	17k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M081R16
NODE=M081R16

$$B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(1S) \rightarrow \ell^+\ell^-)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.02±0.18 OUR AVERAGE				

NODE=M081A01
NODE=M081A01

1.95 ^{+0.22+0.10} _{-0.21-0.16}		36 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.52±0.47±0.32	48	37 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
1.98±0.28±0.12		38 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

³⁶ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.

³⁷ CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.

³⁸ Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) = (0.77 \pm 0.11 \pm 0.05)\%$ using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.05)\%$.

NODE=M081A01;LINKAGE=A
NODE=M081A01;LINKAGE=C

NODE=M081A01;LINKAGE=B

$$[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(1S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
66.6±3.0	39 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M081A00
NODE=M081A00

³⁹ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ events without converted photons.

NODE=M081A00;LINKAGE=A

$$B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) \times B(\Upsilon(2S) \rightarrow \ell^+\ell^-)$$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.74±0.29 OUR AVERAGE				

NODE=M081A02
NODE=M081A02

3.22 ^{+0.58+0.16} _{-0.53-0.71}		40 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$
2.49±0.47±0.31	53	41 CRAWFORD	92B CLE2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$
2.74±0.33±0.18		42 HEINTZ	92 CSB2	$\Upsilon(3S) \rightarrow \gamma\gamma\ell^+\ell^-$

⁴⁰ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ with converted photons.

⁴¹ CRAWFORD 92B quotes $2 \times B(\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)) B(\chi_{bJ}(2P) \rightarrow \gamma \Upsilon(nS)) B(\Upsilon(nS) \rightarrow \ell^+\ell^-)$.

⁴² Calculated by us. HEINTZ 92 quotes $B(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(2P)) \times B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) = (1.90 \pm 0.23 \pm 0.18)\%$ using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$.

NODE=M081A02;LINKAGE=A
NODE=M081A02;LINKAGE=C

NODE=M081A02;LINKAGE=B

$$[B(\chi_{b2}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P))] / [B(\chi_{b1}(2P) \rightarrow \gamma \Upsilon(2S)) \times B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P))]$$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
46.9±2.0	43 LEES	14M BABR	$\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$

NODE=M081A03
NODE=M081A03

⁴³ From a sample of $\Upsilon(3S) \rightarrow \gamma\gamma\mu^+\mu^-$ without converted photons.

NODE=M081A03;LINKAGE=A

$\chi_{b2}(2P)$ REFERENCES

AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=56199
LEES	14M	PR D90 112010	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56343
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
BRIERE	08	PR D78 092007	R.A. Briere <i>et al.</i>	(CLEO Collab.)	REFID=52577
CAWLFIELD	06	PR D73 012003	C. Cawfield <i>et al.</i>	(CLEO Collab.)	REFID=50997
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
CRONIN-HEN...04	PRL	92 222002	D. Cronin-Hennessy <i>et al.</i>	(CLEO Collab.)	REFID=49766
CRAWFORD	92B	PL B294 139	G. Crawford <i>et al.</i>	(CLEO Collab.)	REFID=43177
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586

NODE=M081

$\Upsilon(3S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M048

 $\Upsilon(3S)$ MASS

NODE=M048M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10355.1±0.5	¹ SHAMOV	23	RVUE $e^+e^- \rightarrow$ hadrons
10355.2±0.5	^{2,3} ARTAMONOV	00	MD1 $e^+e^- \rightarrow$ hadrons
10355.3±0.5	^{4,5} BARU	86B	MD1 $e^+e^- \rightarrow$ hadrons

NODE=M048M

¹ Reanalysis of MD1 data using the electron mass from COHEN 87, the radiative corrections from KURAEV 85 and interference effects.

² Reanalysis of BARU 86B using new electron mass (COHEN 87).

³ Superseded by SHAMOV 23.

⁴ Reanalysis of ARTAMONOV 84.

⁵ Superseded by ARTAMONOV 00.

NODE=M048M;LINKAGE=A

NODE=M048M;LINKAGE=AR

NODE=M048M;LINKAGE=B

NODE=M048M;LINKAGE=C

NODE=M048M;LINKAGE=RZ

 $m\Upsilon(3S) - m\Upsilon(2S)$

NODE=M048DM2

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
331.50±0.02±0.13	LEES	11C	BABR $e^+e^- \rightarrow \pi^+\pi^-X$

NODE=M048DM2

 $\Upsilon(3S)$ WIDTH

NODE=M048W

VALUE (keV)	DOCUMENT ID	COMMENT
20.32±1.85 OUR EVALUATION		See the Note on "Width Determinations of the Υ States"

NODE=M048W

→ UNCHECKED ←

 $\Upsilon(3S)$ DECAY MODES

NODE=M048215;NODE=M048

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\Upsilon(2S)$ anything	(10.6 ± 0.8) %	
Γ_2 $\Upsilon(2S)\pi^+\pi^-$	(2.82± 0.18) %	S=1.6
Γ_3 $\Upsilon(2S)\pi^0\pi^0$	(1.85± 0.14) %	
Γ_4 $\Upsilon(2S)\gamma\gamma$	(5.0 ± 0.7) %	
Γ_5 $\Upsilon(2S)\pi^0$	< 5.1 × 10 ⁻⁴	CL=90%
Γ_6 $\Upsilon(1S)\pi^+\pi^-$	(4.37± 0.08) %	
Γ_7 $\Upsilon(1S)\pi^0\pi^0$	(2.20± 0.13) %	
Γ_8 $\Upsilon(1S)\eta$	< 1 × 10 ⁻⁴	CL=90%
Γ_9 $\Upsilon(1S)\pi^0$	< 7 × 10 ⁻⁵	CL=90%
Γ_{10} $h_b(1P)\pi^0$	< 1.2 × 10 ⁻³	CL=90%
Γ_{11} $h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0$	(4.3 ± 1.4) × 10 ⁻⁴	
Γ_{12} $h_b(1P)\pi^+\pi^-$	< 1.2 × 10 ⁻⁴	CL=90%
Γ_{13} $\tau^+\tau^-$	(2.29± 0.30) %	
Γ_{14} $\mu^+\mu^-$	(2.18± 0.21) %	S=2.1
Γ_{15} e^+e^-	(2.18± 0.20) %	
Γ_{16} hadrons	(93 ± 12) %	
Γ_{17} ggg	(35.7 ± 2.6) %	
Γ_{18} $\gamma g g$	(9.7 ± 1.8) × 10 ⁻³	
Γ_{19} 2H anything	(2.33± 0.33) × 10 ⁻⁵	

DESIG=8

DESIG=4

DESIG=10

DESIG=12

DESIG=107

DESIG=3

DESIG=11

DESIG=9

DESIG=106

DESIG=112

DESIG=113

DESIG=114

DESIG=16

DESIG=1

DESIG=2

DESIG=101

DESIG=109

DESIG=110

DESIG=117

Radiative decays

NODE=M048;CLUMP=B

Γ_{20} $\gamma\chi_{b2}(2P)$	(13.1 ± 1.6) %	S=3.4
Γ_{21} $\gamma\chi_{b1}(2P)$	(12.6 ± 1.2) %	S=2.4
Γ_{22} $\gamma\chi_{b0}(2P)$	(5.9 ± 0.6) %	S=1.4
Γ_{23} $\gamma\chi_{b2}(1P)$	(10.0 ± 1.0) × 10 ⁻³	S=1.7
Γ_{24} $\gamma\chi_{b1}(1P)$	(9 ± 5) × 10 ⁻⁴	S=1.8
Γ_{25} $\gamma\chi_{b0}(1P)$	(2.7 ± 0.4) × 10 ⁻³	
Γ_{26} $\gamma\eta_b(2S)$	< 6.2 × 10 ⁻⁴	CL=90%
Γ_{27} $\gamma\eta_b(1S)$	(5.1 ± 0.7) × 10 ⁻⁴	
Γ_{28} $\gamma A^0 \rightarrow \gamma$ hadrons	< 8 × 10 ⁻⁵	CL=90%
Γ_{29} $\gamma X \rightarrow \gamma + \geq 4$ prongs	[a] < 2.2 × 10 ⁻⁴	CL=95%
Γ_{30} $\gamma A^0 \rightarrow \gamma\mu^+\mu^-$	< 5.5 × 10 ⁻⁶	CL=90%
Γ_{31} $\gamma A^0 \rightarrow \gamma\tau^+\tau^-$	[b] < 1.6 × 10 ⁻⁴	CL=90%

DESIG=5

DESIG=6

DESIG=7

DESIG=103

DESIG=104

DESIG=13

DESIG=14

DESIG=15

DESIG=115

DESIG=102

DESIG=116

DESIG=108

Lepton Family number (LF) violating modes

Γ_{32}	$e^\pm \tau^\mp$	LF	< 4.2	$\times 10^{-6}$	CL=90%
Γ_{33}	$e^\pm \mu^\mp$	LF	< 3.6	$\times 10^{-7}$	CL=90%
Γ_{34}	$\mu^\pm \tau^\mp$	LF	< 3.1	$\times 10^{-6}$	CL=90%

NODE=M048;CLUMP=C

DESIG=111

DESIG=119

DESIG=105

[a] $1.5 \text{ GeV} < m_\chi < 5.0 \text{ GeV}$

LINKAGE=C48

[b] For $m_{\tau^+\tau^-}$ in the ranges 4.03–9.52 and 9.61–10.10 GeV.

LINKAGE=MRG

 $\Upsilon(3S) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$

NODE=M048218

 $\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_{16}\Gamma_{15}/\Gamma$**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.414±0.007 OUR AVERAGE			
0.413±0.004±0.006	ROSNER	06	CLEO $10.4 e^+e^- \rightarrow \text{hadrons}$
0.45 ±0.03 ±0.03	⁶ GILES	84B	CLEO $e^+e^- \rightarrow \text{hadrons}$

NODE=M048G2

NODE=M048G2

⁶ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

NODE=M048G2;LINKAGE=R

 $\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ **$\Gamma_6\Gamma_{15}/\Gamma$**

VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT
18.46±0.27±0.77	6.4k	⁷ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$

NODE=M048G01

NODE=M048G01

⁷ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M048G01;LINKAGE=AU

 $\Upsilon(3S)$ PARTIAL WIDTHS

NODE=M048220

 $\Gamma(e^+e^-)$ **Γ_{15}**

VALUE (keV)	DOCUMENT ID
0.443±0.008 OUR EVALUATION	

NODE=M048W2

NODE=M048W2

→ UNCHECKED ←

 $\Upsilon(3S)$ BRANCHING RATIOS

NODE=M048225

 $\Gamma(\Upsilon(2S)\text{anything})/\Gamma_{\text{total}}$ **Γ_1/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.106 ±0.008 OUR AVERAGE				
0.1023±0.0105	4625	^{8,9,10} BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-X$
0.111 ±0.012	4891	^{9,10,11} BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

NODE=M048R8

NODE=M048R8

⁸ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

NODE=M048R;LINKAGE=A

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

NODE=M048R;LINKAGE=B

¹⁰ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$.

NODE=M048R;LINKAGE=D

¹¹ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

NODE=M048R;LINKAGE=C

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ **Γ_2/Γ**

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.82±0.18 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.

NODE=M048R4

NODE=M048R4

3.00±0.02±0.14	543k	LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^-X$
2.40±0.10±0.26	800	¹² AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-e^+e^-$
3.12±0.49	980	^{13,14} BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-\ell^+\ell^-$
2.13±0.38	974	¹⁵ BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

4.82±0.65±0.53 138 ¹⁵ WU 93 CUSB $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ 3.1 ±2.0 5 MAGERAS 82 CUSB $\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$ ¹² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008 \text{ keV}$.

NODE=M048R4;LINKAGE=AU

¹³ From the exclusive mode.


NODE=M048R;LINKAGE=M

¹⁴ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

NODE=M048R4;LINKAGE=A

¹⁵ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

NODE=M048R4;LINKAGE=C



$\Gamma(\Upsilon(2S)\pi^0\pi^0)/\Gamma_{total}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
1.85±0.14 OUR AVERAGE				
1.82±0.09±0.12	4391	¹⁶ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.16±0.39		^{17,18} BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.7 ±0.5 ±0.2	10	¹⁹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

NODE=M048R10
NODE=M048R10

¹⁶ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.
¹⁷ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.
¹⁸ From the exclusive mode.
¹⁹ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M048R10;LINKAGE=BH
NODE=M048R;LINKAGE=K
NODE=M048R10;LINKAGE=M
NODE=M048R;LINKAGE=G

$\Gamma(\Upsilon(2S)\gamma\gamma)/\Gamma_{total}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0502±0.0069	²⁰ BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+\ell^-2\gamma$

NODE=M048R12
NODE=M048R12

²⁰ From the exclusive mode.

NODE=M048R12;LINKAGE=M

$\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{total}$ Γ_5/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<0.51	90	²¹ HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$

NODE=M048R25
NODE=M048R25

²¹ Authors assume $B(\Upsilon(2S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.06\%$.

NODE=M048R25;LINKAGE=HE

$\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{total}$ Γ_6/Γ

Abbreviation MM in the COMMENT field below stands for missing mass.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.37±0.08 OUR AVERAGE				
4.32±0.07±0.13	90k	²² LEES	11L BABR	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.01±0.13	190k	²³ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^+\pi^-$ MM
4.17±0.06±0.19	6.4k	²⁴ AUBERT	08BP BABR	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
4.52±0.35	11830	²⁵ BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$
4.46±0.34±0.50	451	²⁵ WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
4.46±0.30	11221	²⁵ BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^-X,$ $\pi^+\pi^-\ell^+\ell^-$

NODE=M048R3
NODE=M048R3
NODE=M048R3

••• We do not use the following data for averages, fits, limits, etc. •••

4.9 ±1.0	22	GREEN	82 CLEO	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
3.9 ±1.3	26	MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+\pi^-\ell^+\ell^-$

²² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

NODE=M048R3;LINKAGE=LE

²³ A weighted average of the inclusive and exclusive results.

NODE=M048R3;LINKAGE=BH
NODE=M048R3;LINKAGE=AU

²⁴ Using $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$, and $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

²⁵ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

NODE=M048R3;LINKAGE=B

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_2/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M048R28
 NODE=M048R28

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.577±0.026±0.060	800	²⁶ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\ell^+\ell^-$
-------------------	-----	----------------------	-----------	---

²⁶ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

NODE=M048R28;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

NODE=M048R11
 NODE=M048R11

2.20±0.13 OUR AVERAGE

2.24±0.09±0.11	6584	²⁷ BHARI	09 CLEO	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
1.99±0.34	56	²⁸ BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$
2.2 ±0.4 ±0.3	33	²⁹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \pi^0\pi^0\ell^+\ell^-$

²⁷ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

²⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$ and assuming $e\mu$ universality.

²⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

NODE=M048R11;LINKAGE=BH
 NODE=M048R11;LINKAGE=B
 NODE=M048R;LINKAGE=I

 $\Gamma(\Upsilon(1S)\pi^0\pi^0)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_7/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M048R26
 NODE=M048R26

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.501±0.043	³⁰ BHARI	09 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$
-------------	---------------------	---------	-----------------------------------

³⁰ Not independent of other values reported by BHARI 09.

NODE=M048R26;LINKAGE=BH

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R9
 NODE=M048R9

<0.1	90	³¹ LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
----------------	----	--------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.8	90	^{31,32} AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-\pi^0\ell^+\ell^-$
------	----	-------------------------	-----------	--

<0.18	90	³³ HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
-------	----	------------------	----------	---------------------------------------

<2.2	90	BROCK	91 CLEO	$e^+e^- \rightarrow \ell^+\ell^-\eta$
------	----	-------	---------	---------------------------------------

³¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³² Using $\Gamma_{ee}(\Upsilon(3S)) = 0.443 \pm 0.008$ keV.

³³ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M048R9;LINKAGE=LE
 NODE=M048R9;LINKAGE=AU
 NODE=M048R9;LINKAGE=HE

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_8/Γ_6

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R27
 NODE=M048R27

<0.23	90	³⁴ LEES	11L BABR	$\Upsilon(3S) \rightarrow (\pi^+\pi^-)(\gamma\gamma)\ell^+\ell^-$
-----------------	----	--------------------	----------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9	90	³⁵ AUBERT	08BP BABR	$e^+e^- \rightarrow \gamma\pi^+\pi^-(\pi^0)\ell^+\ell^-$
------	----	----------------------	-----------	--

³⁴ Not independent of other values reported by LEES 11L.

³⁵ Not independent of other values reported by AUBERT 08BP.

NODE=M048R27;LINKAGE=LE
 NODE=M048R27;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

NODE=M048R24
 NODE=M048R24

<0.07	90	³⁶ HE	08A CLEO	$e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
-----------------	----	------------------	----------	---

³⁶ Authors assume $B(\Upsilon(1S) \rightarrow e^+e^-) + B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = 4.96\%$.

NODE=M048R24;LINKAGE=HE

 $\Gamma(h_b(1P)\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M048R03
 NODE=M048R03

<1.2 × 10⁻³	90	³⁷ GE	11 CLEO	$\Upsilon(3S) \rightarrow \pi^0$ anything
----------------------------------	----	------------------	---------	---

³⁷ Assuming $M(h_b(1P)) = 9900$ MeV and $\Gamma(h_b(1P)) = 0$ MeV, and allowing $B(h_b(1P) \rightarrow \gamma\eta_b(1S))$ to vary from 0–100%.

NODE=M048R03;LINKAGE=GE

 $\Gamma(h_b(1P)\pi^0 \rightarrow \gamma\eta_b(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=M048R33
 NODE=M048R33

4.3±1.1±0.9	LEES	11K BABR	$\Upsilon(3S) \rightarrow \eta_b\gamma\pi^0$
--------------------	------	----------	--

$\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{total}$ Γ_{12}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2	90	38 LEES	11C BABR	$e^+e^- \rightarrow \pi^+\pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<18		38 BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+\pi^- X$
<15		38 BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+\pi^- X$
³⁸ For $M(h_b(1P)) = 9900$ MeV.				

NODE=M048R34
NODE=M048R34

NODE=M048R34;LINKAGE=MH

$\Gamma(\tau^+\tau^-)/\Gamma_{total}$ Γ_{13}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.29±0.21±0.22	15k	³⁹ BESSON	07 CLEO	$e^+e^- \rightarrow \Upsilon(3S) \rightarrow \tau^+\tau^-$
³⁹ BESSON 07 reports $[\Gamma(\Upsilon(3S) \rightarrow \tau^+\tau^-)/\Gamma_{total}] / [B(\Upsilon(3S) \rightarrow \mu^+\mu^-)] = 1.05 \pm 0.08 \pm 0.05$ which we multiply by our best value $B(\Upsilon(3S) \rightarrow \mu^+\mu^-) = (2.18 \pm 0.21) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.				

NODE=M048R18
NODE=M048R18

NODE=M048R18;LINKAGE=BE

$\Gamma(\tau^+\tau^-)/\Gamma(\mu^+\mu^-)$ Γ_{13}/Γ_{14}


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.968±0.016 OUR AVERAGE				
0.966±0.008±0.014	2.2M	LEES	20E BABR	$e^+e^- \rightarrow \Upsilon(3S)$
1.05 ±0.08 ±0.05	15k	BESSON	07 CLEO	$e^+e^- \rightarrow \Upsilon(3S)$

NODE=M048R19
NODE=M048R19

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ Γ_{14}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0218±0.0021 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.				
0.0239±0.0007±0.0010	81k	ADAMS	05 CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0202±0.0019±0.0033		CHEN	89B CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.0173±0.0015±0.0011		KAARSBERG	89 CSB2	$e^+e^- \rightarrow \mu^+\mu^-$
0.033 ±0.013 ±0.007	1096	ANDREWS	83 CLEO	$e^+e^- \rightarrow \mu^+\mu^-$

NODE=M048R1
NODE=M048R1



$\Gamma(ggg)/\Gamma_{total}$ Γ_{17}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
35.7±2.6	3M	⁴⁰ BESSON	06A CLEO	$\Upsilon(3S) \rightarrow \text{hadrons}$

NODE=M048R30
NODE=M048R30

NODE=M048R30;LINKAGE=BE

⁴⁰ Calculated using BESSON 06A value of $\Gamma(\gamma gg)/\Gamma(ggg) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and the PDG 08 values of $B(\Upsilon(2S) + \text{anything}) = (10.6 \pm 0.8)\%$, $B(\pi^+\pi^-\Upsilon(1S)) = (4.40 \pm 0.10)\%$, $B(\pi^0\pi^0\Upsilon(1S)) = (2.20 \pm 0.13)\%$, $B(\gamma\chi_{b2}(2P)) = (13.1 \pm 1.6)\%$, $B(\gamma\chi_{b1}(2P)) = (12.6 \pm 1.2)\%$, $B(\gamma\chi_{b0}(2P)) = (5.9 \pm 0.6)\%$, $B(\gamma\chi_{b0}(1P)) = (0.30 \pm 0.11)\%$, $B(\mu^+\mu^-) = (2.18 \pm 0.21)\%$, and $R_{\text{hadrons}} = 3.51$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(\gamma gg)/\Gamma_{total}$ BESSON 06A value.

$\Gamma(\gamma g g)/\Gamma_{total}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.18	60k	⁴¹ BESSON	06A CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

Γ_{18}/Γ

NODE=M048R31
NODE=M048R31

⁴¹ Calculated using BESSON 06A values of $\Gamma(\gamma g g)/\Gamma(g g g) = (2.72 \pm 0.06 \pm 0.32 \pm 0.37)\%$ and $\Gamma(g g g)/\Gamma_{total}$. The statistical error is negligible and the systematic error is partially correlated with $\Gamma(g g g)/\Gamma_{total}$ BESSON 06A value.

NODE=M048R31;LINKAGE=BE

$\Gamma(\gamma g g)/\Gamma(g g g)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.72±0.06±0.49	3M	BESSON	06A CLEO	$\Upsilon(3S) \rightarrow (\gamma +) \text{hadrons}$

Γ_{18}/Γ_{17}

NODE=M048R32
NODE=M048R32

$\Gamma(\overline{2H} \text{ anything})/\Gamma_{total}$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
2.33±0.15^{+0.31}_{-0.28}	LEES	14G BABR	$e^+ e^- \rightarrow \overline{2H} X$

Γ_{19}/Γ

NODE=M048R00
NODE=M048R00

$\Gamma(\gamma \chi_{b2}(2P))/\Gamma_{total}$


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.131 ±0.016 OUR AVERAGE				Error includes scale factor of 3.4. See the ideogram below.
0.1579±0.0017±0.0073	568k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.111 ±0.005 ±0.004	10319	⁴² HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.135 ±0.003 ±0.017	30741	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

Γ_{20}/Γ

NODE=M048R5
NODE=M048R5

⁴²Supersedes NARAIN 91.

NODE=M048R;LINKAGE=H



$\Gamma(\gamma \chi_{b1}(2P))/\Gamma_{total}$


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.126 ±0.012 OUR AVERAGE				Error includes scale factor of 2.4. See the ideogram below.
0.1454±0.0018±0.0073	537k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
0.115 ±0.005 ±0.005	11147	⁴³ HEINTZ	92 CSB2	$e^+ e^- \rightarrow \gamma X$
0.105 ^{+0.003} _{-0.002} ±0.013	25759	MORRISON	91 CLE2	$e^+ e^- \rightarrow \gamma X$

Γ_{21}/Γ

NODE=M048R6
NODE=M048R6

⁴³Supersedes NARAIN 91.

NODE=M048R6;LINKAGE=H




$\Gamma(\gamma\chi_{b0}(2P))/\Gamma_{total}$ **Γ_{22}/Γ**

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.059 ±0.006 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.0677±0.0020±0.0065	225k	ARTUSO	05	CLEO $e^+e^- \rightarrow \gamma X$
0.060 ±0.004 ±0.006	4959	⁴⁴ HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma X$
0.049 ^{+0.003} _{-0.004} ±0.006	9903	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma X$

NODE=M048R7
NODE=M048R7

⁴⁴Supersedes NARAIN 91.

NODE=M048R7;LINKAGE=H



$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{total}$ **Γ_{23}/Γ**

VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.0±1.0 OUR AVERAGE					Error includes scale factor of 1.7.
8.0±1.3±0.4	126	^{45,46} KORNICER	11	CLEO	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$
10.5±0.3 ^{+0.7} _{-0.6}	9.7k	LEES	11J	BABR	$\Upsilon(3S) \rightarrow X\gamma$

NODE=M048R21
NODE=M048R21

• • • We do not use the following data for averages, fits, limits, etc. • • •

<19	90	⁴⁷ ASNER	08A	CLEO	$\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
seen		⁴⁸ HEINTZ	92	CSB2	$e^+e^- \rightarrow \gamma\gamma\ell^+\ell^-$

⁴⁵ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

⁴⁶ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S))]$
 $= (1.435 \pm 0.162 \pm 0.169) \times 10^{-3}$ which we divide by our best value $B(\chi_{b2}(1P) \rightarrow \gamma \Upsilon(1S)) = (18.0 \pm 1.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴⁷ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P))]$
 $< 27.1 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b2}(1P)) = 7.15 \times 10^{-2}$.

⁴⁸ HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma \Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0, 1, 2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1, 2$ using $\Upsilon(1S) \rightarrow \ell^+ \ell^-$.

NODE=M048R21;LINKAGE=KA
 NODE=M048R21;LINKAGE=KR

NODE=M048R21;LINKAGE=AS

NODE=M048R21;LINKAGE=HE

$\Gamma(\gamma \chi_{b1}(1P))/\Gamma_{\text{total}}$

Γ_{24}/Γ

VALUE (units 10^{-3})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.9±0.5 OUR AVERAGE Error includes scale factor of 1.8.					
$1.5 \pm 0.4 \pm 0.1$		50	^{49,50} KORNICER	11 CLEO	$e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$
$0.5 \pm 0.3^{+0.2}_{-0.1}$			LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$

NODE=M048R22
 NODE=M048R22

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.7 90 ⁵¹ ASNER 08A CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$
 seen ⁵² HEINTZ 92 CSB2 $e^+ e^- \rightarrow \gamma \gamma \ell^+ \ell^-$

⁴⁹ Assuming $B(\Upsilon(1S) \rightarrow \ell^+ \ell^-) = (2.48 \pm 0.05)\%$.

⁵⁰ KORNICER 11 reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}] \times [B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S))]$
 $= (5.38 \pm 1.20 \pm 0.95) \times 10^{-4}$ which we divide by our best value $B(\chi_{b1}(1P) \rightarrow \gamma \Upsilon(1S)) = (35.2 \pm 2.0) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁵¹ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P))] < 2.5 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b1}(1P)) = 6.9 \times 10^{-2}$.

⁵² HEINTZ 92, while unable to distinguish between different J states, measures $\sum_J B(\Upsilon(3S) \rightarrow \gamma \chi_{bJ}) \times B(\chi_{bJ} \rightarrow \gamma \Upsilon(1S)) = (1.7 \pm 0.4 \pm 0.6) \times 10^{-3}$ for $J = 0, 1, 2$ using inclusive $\Upsilon(1S)$ decays and $(1.2^{+0.4}_{-0.3} \pm 0.09) \times 10^{-3}$ for $J = 1, 2$ using $\Upsilon(1S) \rightarrow \ell^+ \ell^-$.

NODE=M048R22;LINKAGE=KA
 NODE=M048R22;LINKAGE=KR

NODE=M048R22;LINKAGE=AS

NODE=M048R22;LINKAGE=HE

$\Gamma(\gamma \chi_{b0}(1P))/\Gamma_{\text{total}}$

Γ_{25}/Γ

VALUE (units 10^{-2})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.27±0.04 OUR AVERAGE					
$0.27 \pm 0.04 \pm 0.02$		2.3k	LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
$0.30 \pm 0.04 \pm 0.10$		8.7k	ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$

NODE=M048R15
 NODE=M048R15

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.8 90 ⁵³ ASNER 08A CLEO $\Upsilon(3S) \rightarrow \gamma + \text{hadrons}$

⁵³ ASNER 08A reports $[\Gamma(\Upsilon(3S) \rightarrow \gamma \chi_{b0}(1P))/\Gamma_{\text{total}}] / [B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P))] < 21.9 \times 10^{-2}$ which we multiply by our best value $B(\Upsilon(2S) \rightarrow \gamma \chi_{b0}(1P)) = 3.8 \times 10^{-2}$.

NODE=M048R15;LINKAGE=AS

$\Gamma(\gamma \eta_b(2S))/\Gamma_{\text{total}}$

Γ_{26}/Γ

VALUE (units 10^{-4})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
< 6.2	90		ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$
< 19	90		LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$

NODE=M048R16
 NODE=M048R16

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\gamma \eta_b(1S))/\Gamma_{\text{total}}$

Γ_{27}/Γ

VALUE (units 10^{-4})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
5.1±0.7 OUR AVERAGE					
$7.1 \pm 1.8 \pm 1.3$		$2.3 \pm 0.5k$	⁵⁴ BONVICINI	10 CLEO	$\Upsilon(3S) \rightarrow \gamma X$
$4.8 \pm 0.5 \pm 0.6$		$19 \pm 3k$	⁵⁴ AUBERT	09AQ BABR	$\Upsilon(3S) \rightarrow \gamma X$
< 8.5	90		LEES	11J BABR	$\Upsilon(3S) \rightarrow X \gamma$
$4.8 \pm 0.5 \pm 1.2$		$19 \pm 3k$	^{54,55} AUBERT	08V BABR	$\Upsilon(3S) \rightarrow \gamma X$
< 4.3	90		⁵⁶ ARTUSO	05 CLEO	$e^+ e^- \rightarrow \gamma X$

NODE=M048R17
 NODE=M048R17

⁵⁴ Assuming $\Gamma_{\eta_b(1S)} = 10 \text{ MeV}$.

⁵⁵ Systematic error re-evaluated by AUBERT 09AQ.

⁵⁶ Superseded by BONVICINI 10.

NODE=M048R17;LINKAGE=BO
 NODE=M048R17;LINKAGE=AU
 NODE=M048R17;LINKAGE=SU

$$\Gamma(\gamma A^0 \rightarrow \gamma \text{hadrons})/\Gamma_{\text{total}} \quad \Gamma_{28}/\Gamma$$

(0.3 GeV < m_{A^0} < 7 GeV)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 8 \times 10^{-5}$	90	57 LEES	11H BABR	$\Upsilon(3S) \rightarrow \gamma \text{ hadrons}$

NODE=M048R02

NODE=M048R02
NODE=M048R02

⁵⁷ For a narrow scalar or pseudoscalar, A^0 , excluding known resonances, with mass in the range 0.3–7 GeV. Measured 90% CL limits as a function of m_{A^0} range from 1×10^{-6} to 8×10^{-5} .

NODE=M048R02;LINKAGE=LE

$$\Gamma(\gamma X \rightarrow \gamma + \geq 4 \text{ prongs})/\Gamma_{\text{total}} \quad \Gamma_{29}/\Gamma$$

(1.5 GeV < m_X < 5.0 GeV)

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2	95	ROSNER	07A CLEO	$e^+ e^- \rightarrow \gamma X$

NODE=M048R20

NODE=M048R20
NODE=M048R20
$$\Gamma(\gamma A^0 \rightarrow \gamma \mu^+ \mu^-)/\Gamma_{\text{total}} \quad \Gamma_{30}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 5.5	90	58 AUBERT	09Z BABR	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \mu^+ \mu^-$

NODE=M048R04
NODE=M048R04

⁵⁸ For a narrow scalar or pseudoscalar, A^0 , with mass in the range 212–9300 MeV, excluding J/ψ and $\psi(2S)$. Measured 90% CL limits as a function of m_{A^0} range from 0.27–5.5 $\times 10^{-6}$.

NODE=M048R04;LINKAGE=AU

$$\Gamma(\gamma A^0 \rightarrow \gamma \tau^+ \tau^-)/\Gamma_{\text{total}} \quad \Gamma_{31}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-4}$	90	59 AUBERT	09P BABR	$e^+ e^- \rightarrow \gamma A^0 \rightarrow \gamma \tau^+ \tau^-$

NODE=M048R29
NODE=M048R29

⁵⁹ For a narrow scalar or pseudoscalar, A^0 , with $M(\tau^+ \tau^-)$ in the ranges 4.03–9.52 and 9.61–10.10 GeV. Measured 90% CL limits as a function of $M(\tau^+ \tau^-)$ range from 1.5–16 $\times 10^{-5}$.

NODE=M048R29;LINKAGE=AU

———— LEPTON FAMILY NUMBER (LF) VIOLATING MODES ————

NODE=M048230

$$\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}} \quad \Gamma_{32}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.2	90	LEES	10B BABR	$e^+ e^- \rightarrow e^\pm \tau^\mp$

NODE=M048R01
NODE=M048R01
$$\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}} \quad \Gamma_{33}/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.6 \times 10^{-7}$	90	LEES	22A BABR	$e^+ e^- \rightarrow e^\pm \mu^\mp$

NODE=M048R07
NODE=M048R07
$$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}} \quad \Gamma_{34}/\Gamma$$

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 3.1	90	LEES	10B BABR	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$

NODE=M048R23
NODE=M048R23

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 20.3	95	LOVE	08A CLEO	$e^+ e^- \rightarrow \mu^\pm \tau^\mp$
----------	----	------	----------	--

$\Upsilon(3S)$ REFERENCES

NODE=M048

SHAMOV	23	PL B839 137766	A.G. Shamov, O.L. Rezanova	(NOVO, NOVO)	REFID=62012
LEES	22A	PRL 128 091804	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=61659
LEES	20E	PRL 125 241801	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=60700
LEES	14G	PR D89 111102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55939
GE	11	PR D84 032008	J.Y. Ge <i>et al.</i>	(CLEO Collab.)	REFID=53960
KORNICER	11	PR D83 054003	M. Kornicer <i>et al.</i>	(CLEO Collab.)	REFID=16769
LEES	11C	PR D84 011104	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=16775
LEES	11H	PRL 107 221803	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
LEES	11J	PR D84 072002	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53936
LEES	11K	PR D84 091101	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53937
LEES	11L	PR D84 092003	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53938
BONVICINI	10	PR D81 031104	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=53231
LEES	10B	PRL 104 151802	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53355
AUBERT	09AQ	PRL 103 161801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53106
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53062
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
BHARI	09	PR D79 011103	S.R. Bhari <i>et al.</i>	(CLEO Collab.)	REFID=52662
ASNER	08A	PR D78 091103	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=52574
AUBERT	08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
AUBERT	08V	PRL 101 071801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52262
HE	08A	PRL 101 192001	Q. He <i>et al.</i>	(CLEO Collab.)	REFID=52587
LOVE	08A	PRL 101 201601	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52592
PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
BESSION	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
ROSNER	07A	PR D76 117102	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=52079

BESSON	06A	PR D74 012003	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51147
ROSNER	06	PRL 96 092003	J.L. Rosner <i>et al.</i>	(CLEO Collab.)	REFID=51035
ADAMS	05	PRL 94 012001	G.S. Adams <i>et al.</i>	(CLEO Collab.)	REFID=50452
ARTUSO	05	PRL 94 032001	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50454
ARTAMONOV	00	PL B474 427	A.S. Artamonov <i>et al.</i>		REFID=47424
BUTLER	94B	PR D49 40	F. Butler <i>et al.</i>	(CLEO Collab.)	REFID=43799
WU	93	PL B301 307	Q.W. Wu <i>et al.</i>	(CUSB Collab.)	REFID=43313
HEINTZ	92	PR D46 1928	U. Heintz <i>et al.</i>	(CUSB II Collab.)	REFID=43604
BROCK	91	PR D43 1448	I.C. Brock <i>et al.</i>	(CLEO Collab.)	REFID=41579
HEINTZ	91	PRL 66 1563	U. Heintz <i>et al.</i>	(CUSB Collab.)	REFID=41580
MORRISON	91	PRL 67 1696	R.J. Morrison <i>et al.</i>	(CLEO Collab.)	REFID=41634
NARAIN	91	PRL 66 3113	M. Narain <i>et al.</i>	(CUSB Collab.)	REFID=41586
CHEN	89B	PR D39 3528	W.Y. Chen <i>et al.</i>	(CLEO Collab.)	REFID=40919
KAARSBERG	89	PRL 62 2077	T.M. Kaarsberg <i>et al.</i>	(CUSB Collab.)	REFID=40733
BUCHMUEL...	88	HE e^+e^- Physics 412	W. Buchmueller, S. Cooper	(HANN, DESY, MIT)	REFID=40034
Editors: A. Ali and P. Soeding,		World Scientific, Singapore			
COHEN	87	RMP 59 1121	E.R. Cohen, B.N. Taylor	(RISC, NBS)	REFID=11616
BARU	86B	ZPHY C32 622 (errat.)	S.E. Baru <i>et al.</i>	(NOVO)	REFID=22338
KURAEV	85	SJNP 41 466	E.A. Kuraev, V.S. Fadin	(NOVO)	REFID=40033
		Translated from YAF 41 733.			
ARTAMONOV	84	PL 137B 272	A.S. Artamonov <i>et al.</i>	(NOVO)	REFID=22278
GILES	84B	PR D29 1285	R. Giles <i>et al.</i>	(CLEO Collab.)	REFID=22280
ANDREWS	83	PRL 50 807	D.E. Andrews <i>et al.</i>	(CLEO Collab.)	REFID=22273
GREEN	82	PRL 49 617	J. Green <i>et al.</i>	(CLEO Collab.)	REFID=22321
MAGERAS	82	PL 118B 453	G. Mageras <i>et al.</i>	(COLU, CORN, LSU+)	REFID=22359

NODE=M206

 $\chi_{b1}(3P)$

$$I^G(J^{PC}) = 0^+(1^+ +)$$

Observed in the radiative decay to $\Upsilon(1S, 2S, 3S)$, therefore $C = +$.
 J needs confirmation.

NODE=M206

$\chi_{b1}(3P)$ MASS

NODE=M206M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
10513.42 ± 0.41 ± 0.53		¹ SIRUNYAN	18N CMS	$pp \rightarrow \gamma \mu^+ \mu^- X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
10515.7	+ 2.2 + 1.5 - 3.9 - 2.1	2 AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10512.1	± 2.1 ± 0.9	3 AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10511.3	± 1.7 ± 2.5	4 AAIJ	14BI LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$
10530	± 5 ± 9	5 AAD	12A ATLS	$pp \rightarrow \gamma \mu^+ \mu^- X$
10551	±14 ±17	5 ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma \mu^+ \mu^- X$

NODE=M206M

¹ Systematic error includes an additional 0.5 MeV for the uncertainty on the $\Upsilon(3S)$ mass. Also measures $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$ MeV. A total of 372 $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ events was observed.

NODE=M206M;LINKAGE=D

² From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

NODE=M206M;LINKAGE=A

³ The mass of the $\chi_{b1}(3P)$ state obtained by combining the results of AAIJ 14BG with that of AAIJ 14BI. The first uncertainty is experimental and the second attributable to the unknown mass splitting, assumed to be $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

NODE=M206M;LINKAGE=B

⁴ From $\chi_{b1}(3P) \rightarrow \Upsilon(3S)\gamma$ transition assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV.

NODE=M206M;LINKAGE=C

⁵ The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.

NODE=M206M;LINKAGE=AA

$\chi_{b1}(3P)$ DECAY MODES

NODE=M206215;NODE=M206

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Upsilon(1S)\gamma$	seen
Γ_2 $\Upsilon(2S)\gamma$	seen
Γ_3 $\Upsilon(3S)\gamma$	seen

DESIG=1

DESIG=2

DESIG=3

$\chi_{b1}(3P)$ BRANCHING RATIOS

NODE=M206225

$\Gamma(\Upsilon(1S)\gamma)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		169	¹ AAIJ	14BG LHCB	$pp \rightarrow \gamma \mu^+ \mu^- X$	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
seen			AAD	12A ATLS	$pp \rightarrow \gamma \mu^+ \mu^- X$	
seen			ABAZOV	12Q D0	$p\bar{p} \rightarrow \gamma \mu^+ \mu^- X$	

NODE=M206R01

NODE=M206R01

¹ From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

NODE=M206R01;LINKAGE=A

$\Gamma(\Upsilon(2S)\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	169	¹ AAIJ	14BG LHCb	$pp \rightarrow \gamma\mu^+\mu^-X$
seen		AAD	12A ATLAS	$pp \rightarrow \gamma\mu^+\mu^-X$

¹ From $\chi_{b1}(3P) \rightarrow \Upsilon(1S, 2S)\gamma$ transitions assuming $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.5 \pm 1.5$ MeV and allowing for $\pm 30\%$ variation in the $\chi_{b2}(3P)$ production rate relative to that of $\chi_{b1}(3P)$.

NODE=M206R02
NODE=M206R02

NODE=M206R02;LINKAGE=A

 $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen		SIRUNYAN	18N CMS	$pp \rightarrow \gamma\mu^+\mu^-X$
seen	182	AAIJ	14BI LHCb	$pp \rightarrow \gamma\mu^+\mu^-X$

NODE=M206R03
NODE=M206R03

 $\chi_{b1}(3P)$ REFERENCES

SIRUNYAN	18N	PRL 121 092002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	14BG	JHEP 1410 088	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14BI	EPJ C74 3092	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	12A	PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)
ABAZOV	12Q	PR D86 031103	V.M. Abazov <i>et al.</i>	(D0 Collab.)

NODE=M206

REFID=58873
REFID=56199
REFID=56235
REFID=54037
REFID=54264

$\chi_{b2}(3P)$

$$J^{PC} = 0^+(2^{++})$$

Observed in the radiative decay to $\Upsilon(3S)$, therefore $C = +$. J needs confirmation.

NODE=M238

NODE=M238

 $\chi_{b2}(3P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10524.02 ± 0.57 ± 0.53	¹ SIRUNYAN	18N CMS	$pp \rightarrow \gamma\mu^+\mu^-X$
10530 ± 5 ± 9	² AAD	12A ATLAS	$pp \rightarrow \gamma\mu^+\mu^-X$

¹ Systematic error includes an additional 0.5 MeV for the uncertainty on the $\Upsilon(3S)$ mass. Also measures $m_{\chi_{b2}(3P)} - m_{\chi_{b1}(3P)} = 10.60 \pm 0.64 \pm 0.17$ MeV. A total of 372 $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ events was observed.

² The mass barycenter of the merged lineshapes from the $J = 1$ and 2 states.

NODE=M238M

NODE=M238M

NODE=M238M;LINKAGE=A

NODE=M238M;LINKAGE=AA

 $\chi_{b2}(3P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Upsilon(3S)\gamma$	seen

NODE=M238215;NODE=M238

DESIG=1

 $\chi_{b2}(3P)$ BRANCHING RATIOS $\Gamma(\Upsilon(3S)\gamma)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	SIRUNYAN	18N CMS	$pp \rightarrow \gamma\mu^+\mu^-X$

NODE=M238225

NODE=M238R01
NODE=M238R01

 $\chi_{b2}(3P)$ REFERENCES

SIRUNYAN	18N	PRL 121 092002	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAD	12A	PRL 108 152001	G. Aad <i>et al.</i>	(ATLAS Collab.)

NODE=M238

REFID=58873
REFID=54037

$\Upsilon(4S)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

also known as $\Upsilon(10580)$

NODE=M047

 $\Upsilon(4S)$ MASS

NODE=M047M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10579.4±1.2 OUR AVERAGE			
10579.3±0.4±1.2	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
10580.0±3.5	¹ BEBEK	87 CLEO	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10577.4±1.0	² LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BESSON 85.			
² No systematic error given.			

NODE=M047M

NODE=M047M;LINKAGE=C
NODE=M047M;LINKAGE=B **$\Upsilon(4S)$ WIDTH**

NODE=M047W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
20.5±2.5 OUR AVERAGE			
20.7±1.6±2.5	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
20 ±2 ±4	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
25 ±2.5	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

NODE=M047W

 $\Upsilon(4S)$ DECAY MODES

NODE=M047215;NODE=M047

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $B\bar{B}$	> 96 %	95%
Γ_2 B^+B^-	(51.4 ±0.6) %	
Γ_3 D^+ anything + c.c.	(17.8 ±2.6) %	
Γ_4 $B^0\bar{B}^0$	(48.6 ±0.6) %	
Γ_5 $J/\psi K_S^0 + (J/\psi, \eta_c) K_S^0$	< 4 × 10 ⁻⁷	90%
Γ_6 non- $B\bar{B}$	< 4 %	95%
Γ_7 e^+e^-	(1.57±0.08) × 10 ⁻⁵	
Γ_8 $\rho^+\rho^-$	< 5.7 × 10 ⁻⁶	90%
Γ_9 $K^*(892)^0\bar{K}^0$	< 2.0 × 10 ⁻⁶	90%
Γ_{10} $J/\psi(1S)$ anything	< 1.9 × 10 ⁻⁴	95%
Γ_{11} D^{*+} anything + c.c.	< 7.4 %	90%
Γ_{12} ϕ anything	(7.1 ±0.6) %	
Γ_{13} $\phi\eta$	< 1.8 × 10 ⁻⁶	90%
Γ_{14} $\phi\eta'$	< 4.3 × 10 ⁻⁶	90%
Γ_{15} $\rho\eta$	< 1.3 × 10 ⁻⁶	90%
Γ_{16} $\rho\eta'$	< 2.5 × 10 ⁻⁶	90%
Γ_{17} $\Upsilon(1S)$ anything	< 4 × 10 ⁻³	90%
Γ_{18} $\Upsilon(1S)\pi^+\pi^-$	(8.2 ±0.4) × 10 ⁻⁵	
Γ_{19} $\Upsilon(1S)\eta$	(1.81±0.18) × 10 ⁻⁴	
Γ_{20} $\Upsilon(1S)\eta'$	(3.4 ±0.9) × 10 ⁻⁵	
Γ_{21} $\Upsilon(2S)\pi^+\pi^-$	(8.2 ±0.8) × 10 ⁻⁵	
Γ_{22} $h_b(1P)\pi^+\pi^-$	not seen	
Γ_{23} $h_b(1P)\eta$	(2.18±0.21) × 10 ⁻³	
Γ_{24} $\eta_b(1S)\omega$	< 1.8 × 10 ⁻⁴	90%
Γ_{25} 2H anything	< 1.3 × 10 ⁻⁵	90%

DESIG=8;OUR EST;→ UNCHECKED ←

DESIG=10

DESIG=12

DESIG=11

DESIG=15

DESIG=6

DESIG=1

DESIG=16

DESIG=22

DESIG=2

DESIG=3

DESIG=4

DESIG=13

DESIG=18

DESIG=19

DESIG=20

DESIG=5

DESIG=7

DESIG=17

DESIG=26

DESIG=9

DESIG=21

DESIG=23

DESIG=27

DESIG=14

Double Radiative Decays

Γ_{26} $\gamma\gamma\Upsilon(D) \rightarrow \gamma\gamma\eta\Upsilon(1S)$	< 2.3 × 10 ⁻⁵	90%
--	--------------------------	-----

NODE=M047;CLUMP=B

DESIG=24

$\Upsilon(4S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$

Γ_7


VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.272 ± 0.029 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
0.321 ± 0.017 ± 0.029	AUBERT	05Q BABR	$e^+e^- \rightarrow$ hadrons
0.28 ± 0.05 ± 0.01	¹ ALBRECHT	95E ARG	$e^+e^- \rightarrow$ hadrons
0.192 ± 0.007 ± 0.038	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
0.283 ± 0.037	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

¹Using LEYAOUANC 77 parametrization of $\Gamma(s)$.

NODE=M047220

NODE=M047W1
NODE=M047W1

NODE=M047W1;LINKAGE=A



$\Upsilon(4S)$ BRANCHING RATIOS

———— $B\bar{B}$ DECAYS ————

The ratio of branching fraction to charged and neutral B mesons is often derived assuming isospin invariance in the decays, and relies on the knowledge of the B^+/B^0 lifetime ratio. "OUR EVALUATION" is obtained based on averages of rescaled data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <https://hflav.web.cern.ch/>. The averaging/rescaling procedure takes into account the common dependence of the measurement on the value of the lifetime ratio.

NODE=M047230

NODE=M047BBD

NODE=M047BBD

$\Gamma(B^+B^-)/\Gamma_{total}$

Γ_2/Γ

VALUE	DOCUMENT ID	COMMENT
0.514 ± 0.006 OUR EVALUATION	(Produced by HFLAV) Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$	

NODE=M047R11
NODE=M047R11
→ UNCHECKED ←

$\Gamma(D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{total}$

Γ_3/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.178 ± 0.021 ± 0.016	¹ ARTUSO	05B CLE3	$e^+e^- \rightarrow D_s^+ X$

NODE=M047R13
NODE=M047R13

¹ARTUSO 05B reports $[\Gamma(\Upsilon(4S) \rightarrow D_s^+ \text{ anything} + \text{c.c.})/\Gamma_{total}] \times [B(D_s^+ \rightarrow \phi\pi^+)] = (8.0 \pm 0.2 \pm 0.9) \times 10^{-3}$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R13;LINKAGE=AR

$\Gamma(B^0\bar{B}^0)/\Gamma_{total}$

Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.486 ± 0.006 OUR EVALUATION	Assuming $B(\Upsilon(4S) \rightarrow B\bar{B}) = 1$		

NODE=M047R12
NODE=M047R12
→ UNCHECKED ←

••• We do not use the following data for averages, fits, limits, etc. •••

0.487 ± 0.010 ± 0.008	¹ AUBERT,B	05H BABR	$\Upsilon(4S) \rightarrow \bar{B}B \rightarrow D^* \ell \nu_\ell$
-----------------------	-----------------------	----------	---

¹ Direct measurement. This value is averaged with the value extracted from the $\Gamma(B^+ B^-) / \Gamma(B^0 \bar{B}^0)$ measurements.

NODE=M047R12;LINKAGE=AU

 $\Gamma(B^+ B^-) / \Gamma(B^0 \bar{B}^0)$ Γ_2 / Γ_4

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

1.058 ± 0.024 OUR EVALUATION

Average is meaningless. [1.031 ± 0.033 OUR 2023 AVERAGE]

1.065 ± 0.012 ± 0.051	¹ CHOUDHURY 23	BELL	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K$
1.006 ± 0.036 ± 0.031	² AUBERT 04F	BABR	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K$
1.01 ± 0.03 ± 0.09	² HASTINGS 03	BELL	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow \text{dileptons}$
1.058 ± 0.084 ± 0.136	³ ATHAR 02	CLEO	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow D^* \ell \nu$
1.10 ± 0.06 ± 0.05	⁴ AUBERT 02	BABR	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow (c\bar{c}) K^*$
1.04 ± 0.07 ± 0.04	⁵ ALEXANDER 01	CLEO	$\Upsilon(4S) \rightarrow B\bar{B} \rightarrow J/\psi K^*$

NODE=M047R10
 NODE=M047R10
 → UNCHECKED ←

¹ CHOUDHURY 23 includes uncertainty due to the isospin symmetry assumption in $B \rightarrow J/\psi K$ decays.

² HASTINGS 03 and AUBERT 04F assume $\tau(B^+) / \tau(B^0) = 1.083 \pm 0.017$.

³ ATHAR 02 assumes $\tau(B^+) / \tau(B^0) = 1.074 \pm 0.028$. Supersedes BARISH 95.

⁴ AUBERT 02 assumes $\tau(B^+) / \tau(B^0) = 1.062 \pm 0.029$.

⁵ ALEXANDER 01 assumes $\tau(B^+) / \tau(B^0) = 1.066 \pm 0.024$.

NODE=M047R10;LINKAGE=A

NODE=M047R10;LINKAGE=F

NODE=M047R10;LINKAGE=D

NODE=M047R10;LINKAGE=E

NODE=M047R10;LINKAGE=C

 $[\Gamma(J/\psi K_S^0) + \Gamma((J/\psi, \eta_c) K_S^0)] / \Gamma_{\text{total}}$ Γ_5 / Γ

Forbidden by CP invariance.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<4	90	¹ TAJIMA 07A	BELL	$\Upsilon(4S) \rightarrow B^0 \bar{B}^0$
----	----	-------------------------	------	--

¹ $\Upsilon(4S)$ with CP = +1 decays to the final state with CP = -1.

NODE=M047R16

NODE=M047R16

NODE=M047R16

NODE=M047R16;LINKAGE=TA

NODE=M047NBB

non- $B\bar{B}$ DECAYS $\Gamma(\text{non-}B\bar{B}) / \Gamma_{\text{total}}$ Γ_6 / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.04	95	BARISH 96B	CLEO	$e^+ e^-$
-------	----	------------	------	-----------

NODE=M047R6

NODE=M047R6

 $\Gamma(e^+ e^-) / \Gamma_{\text{total}}$ Γ_7 / Γ

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

1.57 ± 0.08 OUR AVERAGE

1.55 ± 0.04 ± 0.07	AUBERT 05Q	BABR	$e^+ e^- \rightarrow \text{hadrons}$
--------------------	------------	------	--------------------------------------

2.77 ± 0.50 ± 0.49	¹ ALBRECHT 95E	ARG	$e^+ e^- \rightarrow \text{hadrons}$
--------------------	---------------------------	-----	--------------------------------------

¹ Using LEYAOUANC 77 parametrization of $\Gamma(s)$.

NODE=M047R5

NODE=M047R5

NODE=M047R5;LINKAGE=A

 $\Gamma(\rho^+ \rho^-) / \Gamma_{\text{total}}$ Γ_8 / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<5.7 × 10 ⁻⁶	90	AUBERT 08BO	BABR	$e^+ e^- \rightarrow \pi^+ \pi^- 2\pi^0$
-------------------------	----	-------------	------	--

NODE=M047R17

NODE=M047R17

 $\Gamma(K^*(892)^0 \bar{K}^0) / \Gamma_{\text{total}}$ Γ_9 / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.0 × 10 ⁻⁶	90	SHEN 13A	BELL	$e^+ e^- \rightarrow K^*(892)^0 \bar{K}^0$
-------------------------	----	----------	------	--

NODE=M047R02

NODE=M047R02

 $\Gamma(J/\psi(1S) \text{ anything}) / \Gamma_{\text{total}}$ Γ_{10} / Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.9	95	¹ ABE 02D	BELL	$e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$
------	----	----------------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<4.7	90	¹ AUBERT 01C	BABR	$e^+ e^- \rightarrow J/\psi X \rightarrow \ell^+ \ell^- X$
------	----	-------------------------	------	--

¹ Uses $B(J/\psi \rightarrow e^+ e^-) = 0.0593 \pm 0.0010$ and $B(J/\psi \rightarrow \mu^+ \mu^-) = 0.0588 \pm 0.0010$.

NODE=M047R1

NODE=M047R1

NODE=M047R;LINKAGE=AC

 $\Gamma(D^{*+} \text{ anything} + \text{c.c.}) / \Gamma_{\text{total}}$ Γ_{11} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.074	90	¹ ALEXANDER 90C	CLEO	$e^+ e^-$
--------	----	----------------------------	------	-----------

¹ For $x > 0.473$.

NODE=M047R2

NODE=M047R2

NODE=M047R2;LINKAGE=A

 $\Gamma(\phi \text{ anything}) / \Gamma_{\text{total}}$ Γ_{12} / Γ

VALUE (units 10^{-2})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

7.1 ± 0.1 ± 0.6		HUANG 07	CLEO	$\Upsilon(4S) \rightarrow \phi X$
------------------------	--	----------	------	-----------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.23	90	¹ ALEXANDER 90C	CLEO	$e^+ e^-$
-------	----	----------------------------	------	-----------

¹ For $x > 0.52$.

NODE=M047R3

NODE=M047R3

NODE=M047R3;LINKAGE=A

$\Gamma(\phi\eta)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.8	90	¹ BELOUS 09	BELL	$e^+e^- \rightarrow \phi\eta$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.5	90	AUBERT, BE 06F	BABR	$e^+e^- \rightarrow \phi\eta$
------	----	----------------	------	-------------------------------

¹ Using all intermedite branching fraction values from PDG 08.

NODE=M047R14
NODE=M047R14

NODE=M047R14;LINKAGE=BE

 $\Gamma(\phi\eta')/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<4.3	90	¹ BELOUS 09	BELL	$e^+e^- \rightarrow \phi\eta'$

¹ Using all intermedite branching fraction values from PDG 08.

NODE=M047R21
NODE=M047R21

NODE=M047R21;LINKAGE=BE

 $\Gamma(\rho\eta)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<1.3	90	¹ BELOUS 09	BELL	$e^+e^- \rightarrow \rho\eta$

¹ Using all intermedite branching fraction values from PDG 08.

NODE=M047R22
NODE=M047R22

NODE=M047R22;LINKAGE=BE

 $\Gamma(\rho\eta')/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.5	90	¹ BELOUS 09	BELL	$e^+e^- \rightarrow \rho\eta'$

¹ Using all intermedite branching fraction values from PDG 08.

NODE=M047R23
NODE=M047R23

NODE=M047R23;LINKAGE=BE

 $\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	90	ALEXANDER 90C	CLEO	e^+e^-

NODE=M047R4
NODE=M047R4

 $\Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
8.2 ± 0.4	OUR AVERAGE				

8.2 ± 0.5 ± 0.4	515	GUIDO 17	BELL	$\Upsilon(4S) \rightarrow \pi^+\pi^-\mu^+\mu^-$
-----------------	-----	----------	------	---

8.5 ± 1.3 ± 0.1	113 ± 16	¹ SOKOLOV 09	BELL	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
-----------------	----------	-------------------------	------	---

8.00 ± 0.64 ± 0.27	430	² AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\ell^+\ell^-$
--------------------	-----	--------------------------	------	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

17.8 ± 4.0 ± 0.3		^{3,4} SOKOLOV 07	BELL	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
------------------	--	---------------------------	------	---

9.0 ± 1.5 ± 0.2	167 ± 19	⁵ AUBERT 06R	BABR	$e^+e^- \rightarrow \pi^+\pi^-\mu^+\mu^-$
-----------------	----------	-------------------------	------	---

<12	90	GLENN 99	CLE2	e^+e^-
-----	----	----------	------	----------

¹ SOKOLOV 09 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (0.211 \pm 0.030 \pm 0.014) \times 10^{-5}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

² Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

³ SOKOLOV 07 reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (4.42 \pm 0.81 \pm 0.56) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

⁴ According to the authors, systematic errors were underestimated.

⁵ Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(1S) \rightarrow \mu^+\mu^-)] = (2.23 \pm 0.25 \pm 0.27) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.04) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R7
NODE=M047R7

NODE=M047R7;LINKAGE=SK

NODE=M047R7;LINKAGE=UB

NODE=M047R7;LINKAGE=SO

NODE=M047R7;LINKAGE=US

NODE=M047R7;LINKAGE=AU

 $\Gamma(\Upsilon(1S)\eta)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.81 ± 0.18	OUR AVERAGE				

1.70 ± 0.23 ± 0.08	49	GUIDO 17	BELL	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\mu^+\mu^-$
--------------------	----	----------	------	--

1.96 ± 0.26 ± 0.09	56	¹ AUBERT 08BP	BABR	$\Upsilon(4S) \rightarrow \pi^+\pi^-\pi^0\ell^+\ell^-$
--------------------	----	--------------------------	------	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.7	90	² TAMPONI 15	BELL	$e^+e^- \rightarrow \gamma\eta + \text{hadrons}$
------	----	-------------------------	------	--

¹ Using $B(\Upsilon(1S) \rightarrow e^+e^-) = (2.38 \pm 0.11)\%$ and $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.05)\%$.

² Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.

NODE=M047R18
NODE=M047R18

NODE=M047R18;LINKAGE=UB

NODE=M047R18;LINKAGE=A

$\Gamma(\Upsilon(1S)\eta')/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
3.43±0.88±0.21	27	GUIDO	18	BELL $\Upsilon(4S) \rightarrow (\rho^0 \gamma, \pi^+ \pi^- \eta) \mu^+ \mu^-$

NODE=M047R03
NODE=M047R03

 $\Gamma(\Upsilon(1S)\eta)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_{19}/Γ_{18}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M047R19
NODE=M047R19

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.41±0.40±0.12 56 ¹ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+ \pi^- (\pi^0) \ell^+ \ell^-$

¹ Not independent of other values reported by AUBERT 08BP.

NODE=M047R19;LINKAGE=UB

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	------	-------------	------	---------

8.2±0.8 OUR AVERAGE

7.9±1.0±0.4 181 GUIDO 17 BELL $\Upsilon(4S) \rightarrow \pi^+ \pi^- \mu^+ \mu^-$

8.6±1.1±0.7 220 ¹ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.8±1.7±0.8 97 ± 15 ² AUBERT 06R BABR $e^+ e^- \rightarrow \pi^+ \pi^- \mu^+ \mu^-$

<3.9 90 GLENN 99 CLE2 $e^+ e^-$

¹ Using $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$ and $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$.

NODE=M047R9
NODE=M047R9

NODE=M047R9;LINKAGE=UB

² Superseded by AUBERT 08BP. AUBERT 06R reports $[\Gamma(\Upsilon(4S) \rightarrow \Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}] \times [B(\Upsilon(2S) \rightarrow \mu^+ \mu^-)] = (1.69 \pm 0.26 \pm 0.20) \times 10^{-6}$ which we divide by our best value $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

NODE=M047R9;LINKAGE=AU

 $\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma(\Upsilon(1S)\pi^+\pi^-)$ Γ_{21}/Γ_{18}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.16±0.16±0.14 220 ¹ AUBERT 08BP BABR $\Upsilon(4S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$

¹ Using $B(\Upsilon(1S) \rightarrow e^+ e^-) = (2.38 \pm 0.11)\%$, $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = (2.48 \pm 0.05)\%$, $B(\Upsilon(2S) \rightarrow e^+ e^-) = (1.91 \pm 0.16)\%$, and $B(\Upsilon(2S) \rightarrow \mu^+ \mu^-) = (1.93 \pm 0.17)\%$. Not independent of other values reported by AUBERT 08BP.

NODE=M047R20
NODE=M047R20

NODE=M047R20;LINKAGE=UB

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

not seen (35⁺³²₋₂₆)k ¹ ADACHI 12 BELL 10.58 $e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-$

¹ From the upper limit on the ratio of $\sigma(e^+ e^- \rightarrow h_b(1P)\pi^+\pi^-)$ at the $\Upsilon(4S)$ to that at the $\Upsilon(5S)$ of 0.27.

NODE=M047R01
NODE=M047R01

NODE=M047R01;LINKAGE=AD

 $\Gamma(h_b(1P)\eta)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

2.18±0.11±0.18 112k ¹ TAMPONI 15 BELL $e^+ e^- \rightarrow h_b(1P)\eta$

¹ Using $B(\eta \rightarrow 2\gamma) = (39.41 \pm 0.20)\%$.

NODE=M047R00
NODE=M047R00

NODE=M047R00;LINKAGE=A

 $\Gamma(\eta_b(1S)\omega)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<1.8 × 10⁻⁴ 90 OSKIN 20 BELL $e^+ e^- \rightarrow \omega X$

NODE=M047R04
NODE=M047R04

 $\Gamma(\eta_b(1S)\omega)/\Gamma(h_b(1P)\eta)$ Γ_{24}/Γ_{23}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<8.4 × 10⁻² 90 ¹ OSKIN 20 BELL $e^+ e^- \rightarrow \omega X$

NODE=M047R05
NODE=M047R05

NODE=M047R05;LINKAGE=A

¹ Using $B(\Upsilon(4S) \rightarrow h_b(1P)\eta) = (2.18 \pm 0.11 \pm 0.18) \times 10^{-3}$ from TAMPONI 15.

 $\Gamma(\overline{2H} \text{ anything})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
--------------------------	-----	-------------	------	---------

<1.3 90 ASNER 07 CLEO $e^+ e^- \rightarrow \overline{d} X$

NODE=M047R15
NODE=M047R15

————— Double Radiative Decays —————

NODE=M047240

 $\Gamma(\gamma\gamma \Upsilon(D) \rightarrow \gamma\gamma\eta \Upsilon(1S))/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<2.3 × 10⁻⁵ 90 GUIDO 17 BELL $\Upsilon(4S) \rightarrow \gamma\gamma\pi^+\pi^-\pi^0\mu^+\mu^-$

NODE=M047R24
NODE=M047R24

$\Upsilon(4S)$ REFERENCES

CHOUDHURY 23	PR D107 L031102	S. Choudhury <i>et al.</i>	(BELLE Collab.)	REFID=62074
OSKIN 20	PR D102 092011	P. Oskin <i>et al.</i>	(BELLE Collab.)	REFID=60735
GUIDO 18	PRL 121 062001	E. Guido <i>et al.</i>	(BELLE Collab.)	REFID=58860
GUIDO 17	PR D96 052005	E. Guido <i>et al.</i>	(BELLE Collab.)	REFID=58218
TAMPONI 15	PRL 115 142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=56996
SHEN 13A	PR D88 052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI 12	PRL 108 032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
BELOUS 09	PL B681 400	K. Belous <i>et al.</i>	(BELLE Collab.)	REFID=53107
SOKOLOV 09	PR D79 051103	A. Sokolov <i>et al.</i>	(BELLE Collab.)	REFID=52760
AUBERT 08BO	PR D78 071103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52659
AUBERT 08BP	PR D78 112002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52660
PDG 08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)	REFID=52166
ASNER 07	PR D75 012009	D.M. Asner <i>et al.</i>	(CLEO Collab.)	REFID=51617
HUANG 07	PR D75 012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
SOKOLOV 07	PR D75 071103	A. Sokolov <i>et al.</i>	(BELLE Collab.)	REFID=51715
TAJIMA 07A	PRL 99 231601	O. Tajima <i>et al.</i>	(BELLE Collab.)	REFID=52066
AUBERT 06R	PRL 96 232001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51143
AUBERT,BE 06F	PR D74 111103	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=51563
ARTUSO 05B	PRL 95 261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
AUBERT 05Q	PR D72 032005	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50774
AUBERT,B 05H	PRL 95 042001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=50777
AUBERT 04F	PR D69 071101	B. Aubert <i>et al.</i>		REFID=49748
HASTINGS 03	PR D67 052004	N.C. Hastings <i>et al.</i>	(BELLE Collab.)	REFID=49209
ABE 02D	PRL 88 052001	K. Abe <i>et al.</i>	(BELLE Collab.)	REFID=48557
ATHAR 02	PR D66 052003	S.B. Athar <i>et al.</i>	(CLEO Collab.)	REFID=48832
AUBERT 02	PR D65 032001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48514
ALEXANDER 01	PRL 86 2737	J.P. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=48316
AUBERT 01C	PRL 87 162002	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=48346
GLENN 99	PR D59 052003	S. Glenn <i>et al.</i>		REFID=46890
BARISH 96B	PRL 76 1570	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44693
ALBRECHT 95E	ZPHY C65 619	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=44372
BARISH 95	PR D51 1014	B.C. Barish <i>et al.</i>	(CLEO Collab.)	REFID=44139
ALEXANDER 90C	PRL 64 2226	J. Alexander <i>et al.</i>	(CLEO Collab.)	REFID=41346
BEBEK 87	PR D36 1289	C. Bebek <i>et al.</i>	(CLEO Collab.)	REFID=40270
BESSION 85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK 85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369
LEYAOUANC 77	PL B71 397	A. Le Yaouanc <i>et al.</i>	(ORSAY)	REFID=44695

NODE=M243

 $\Upsilon(10753)$

$$I^G(J^{PC}) = ?(1^{--})$$

OMITTED FROM SUMMARY TABLE

A candidate for $\Upsilon(3D)$ state or an exotic structure.

NODE=M243

Seen by MIZUK 19 in $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n=1,2,3$) with a significance of 5.2σ .

 $\Upsilon(10753)$ MASS

NODE=M243M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$10752.7 \pm 5.9^{+0.7}_{-1.1}$	¹ MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

NODE=M243M

• • • We do not use the following data for averages, fits, limits, etc. • • •

10761 ± 2	² DONG	20A	$e^+e^- \rightarrow b\bar{b}$
---------------	-------------------	-----	-------------------------------

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M243M;LINKAGE=A

² From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M243M;LINKAGE=B

 $\Upsilon(10753)$ WIDTH

NODE=M243W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$35.5^{+17.6+3.9}_{-11.3-3.3}$	¹ MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

NODE=M243W

• • • We do not use the following data for averages, fits, limits, etc. • • •

48.5 ± 3.0	² DONG	20A	$e^+e^- \rightarrow b\bar{b}$
----------------	-------------------	-----	-------------------------------

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M243W;LINKAGE=A

² From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M243W;LINKAGE=B

$\Upsilon(10753)$ DECAY MODES

NODE=M243215;NODE=M243

Mode					
Γ_1	$\Upsilon(1S)\pi^+\pi^-$			DESIG=3	
Γ_2	$\Upsilon(2S)\pi^+\pi^-$			DESIG=4	
Γ_3	$\Upsilon(3S)\pi^+\pi^-$			DESIG=5	
Γ_4	$\omega\chi_{b1}(1P)$			DESIG=6	
Γ_5	$\omega\chi_{b2}(1P)$			DESIG=7	
Γ_6	e^+e^-			DESIG=2	

$\Upsilon(10753) \Gamma(i)\Gamma(e^+e^-)/\Gamma(\text{total})$						
$\Gamma(\Upsilon(1S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_6/\Gamma$	NODE=M243225
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			NODE=M243R00 NODE=M243R00
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.295±0.175	1,2 MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$		NODE=M243R00;LINKAGE=A
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.						
² Reported as the range 0.12–0.47 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.						
$\Gamma(\Upsilon(2S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_2\Gamma_6/\Gamma$	NODE=M243R02 NODE=M243R02
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.875±0.345	1,2 MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$		NODE=M243R02;LINKAGE=A
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.						
² Reported as the range 0.53–1.22 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.						
$\Gamma(\Upsilon(3S)\pi^+\pi^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_3\Gamma_6/\Gamma$	NODE=M243R03 NODE=M243R03
<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.235±0.025	1,2 MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$		NODE=M243R03;LINKAGE=A
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.63\text{--}11.02$ GeV, including the initial-state radiation at $\Upsilon(10860)$.						
² Reported as the range 0.21–0.26 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.						
$\Gamma(\omega\chi_{b1}(1P)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_4\Gamma_6/\Gamma$	NODE=M243R05 NODE=M243R05
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.63±0.39±0.20	68	1 ADACHI	23	BELL	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	NODE=M243R05;LINKAGE=A
¹ A fit solution with constructive interference. The other solution corresponding to destructive interference gives a value of $2.01 \pm 0.38 \pm 0.76$ eV.						
$\Gamma(\omega\chi_{b2}(1P)) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$					$\Gamma_5\Gamma_6/\Gamma$	NODE=M243R06 NODE=M243R06
<u>VALUE (eV)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.53±0.46±0.15	68	1 ADACHI	23	BELL	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$	NODE=M243R06;LINKAGE=A
¹ A fit solution with constructive interference. The other solution corresponding to destructive interference gives a value of $1.32 \pm 0.44 \pm 0.55$ eV.						

 $\Upsilon(10753)$ REFERENCES

NODE=M243

ADACHI	23	PRL 130 091902	I. Adachi <i>et al.</i>	(BELLE II Collab.)	REFID=62210
DONG	20A	CP C44 083001	X.-K. Dong <i>et al.</i>		REFID=60595
MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60090
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661

$\Upsilon(10860)$

$$J^{PC} = 0^{-}(1^{-})^{-}$$

NODE=M092

 $\Upsilon(10860)$ MASS

NODE=M092M

NODE=M092M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10885.2^{+2.6}_{-1.6} OUR AVERAGE			
10885.3 ^{±1.5} _{-0.9}	1 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
10884.7 ^{+3.6} _{-3.4} ^{+8.9} _{-1.0}	2 MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P,2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10882 ± 1	3 DONG	20A	$e^+e^- \rightarrow b\bar{b}$
10881.8 ^{+1.0} _{-1.1} ± 1.2	4,5 SANTEL	16	BELL $e^+e^- \rightarrow$ hadrons
10891.1 ± 3.2 ^{+1.2} _{-2.0}	6,7 SANTEL	16	BELL $e^+e^- \rightarrow \Upsilon(1S,2S,3S)\pi^+\pi^-$
10879 ± 3	8,9 CHEN	10	BELL $e^+e^- \rightarrow$ hadrons
10888.4 ^{+2.7} _{-2.6} ± 1.2	10 CHEN	10	BELL $e^+e^- \rightarrow \Upsilon(1S,2S,3S)\pi^+\pi^-$
10876 ± 2	8 AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
10869 ± 2	11 AUBERT	09E	BABR $e^+e^- \rightarrow$ hadrons
10868 ± 6 ± 5	12 BESSON	85	CLEO $e^+e^- \rightarrow$ hadrons
10845 ± 20	13 LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons

OCCUR=2

OCCUR=2

OCCUR=2

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M092M;LINKAGE=E

² From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

NODE=M092M;LINKAGE=D

³ From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M092M;LINKAGE=H

⁴ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).

NODE=M092M;LINKAGE=A

⁵ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

NODE=M092M;LINKAGE=B

⁶ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M092M;LINKAGE=C

⁷ Superseded by MIZUK 19.

NODE=M092M;LINKAGE=F

⁸ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M092M;LINKAGE=AU

⁹ The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.

NODE=M092M;LINKAGE=CH

¹⁰ In a model where a flat nonresonant $\Upsilon(1S,2S,3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.

NODE=M092M;LINKAGE=CE

¹¹ In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.

NODE=M092M;LINKAGE=UB

¹² Assuming four Gaussians with radiative tails and a single step in R .

NODE=M092M;LINKAGE=BE

¹³ In a coupled-channel model with three resonances and a smooth step in R .

NODE=M092M;LINKAGE=LO

 $\Upsilon(10860)$ WIDTH

NODE=M092W

NODE=M092W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
37 ± 4 OUR AVERAGE			
36.6 ^{+4.5} _{-3.9} ± 0.5 ± 1.1	1 MIZUK	19	BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
40.6 ^{+12.7} _{-8.0} ± 1.1 ± 19.1	2 MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P,2P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

49.5 ± 1.5	³ DONG	20A	$e^+e^- \rightarrow b\bar{b}$	
48.5 ^{+1.9+2.0} _{-1.8-2.8}	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow \text{hadrons}$	
53.7 ^{+7.1+1.3} _{-5.6-5.4}	6,7 SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$	OCCUR=2
46 ⁺⁹ ₋₇	8,9 CHEN	10 BELL	$e^+e^- \rightarrow \text{hadrons}$	
30.7 ^{+8.3} _{-7.0} ± 3.1	¹⁰ CHEN	10 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$	OCCUR=2
43 ± 4	⁸ AUBERT	09E BABR	$e^+e^- \rightarrow \text{hadrons}$	
74 ± 4	¹¹ AUBERT	09E BABR	$e^+e^- \rightarrow \text{hadrons}$	OCCUR=2
112 ± 17 ± 23	¹² BESSON	85 CLEO	$e^+e^- \rightarrow \text{hadrons}$	
110 ± 15	¹³ LOVELOCK	85 CUSB	$e^+e^- \rightarrow \text{hadrons}$	
¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.				NODE=M092W;LINKAGE=F
² From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.				NODE=M092W;LINKAGE=D
³ From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.				NODE=M092W;LINKAGE=G
⁴ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).				NODE=M092W;LINKAGE=A
⁵ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.				NODE=M092W;LINKAGE=B
⁶ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.				NODE=M092W;LINKAGE=C
⁷ Superseded by MIZUK 19.				NODE=M092W;LINKAGE=E
⁸ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.				NODE=M092W;LINKAGE=AU
⁹ The parameters of the $\Upsilon(11020)$ are fixed to those in AUBERT 09E.				NODE=M092W;LINKAGE=CH
¹⁰ In a model where a flat nonresonant $\Upsilon(1S, 2S, 3S)\pi^+\pi^-$ continuum interferes with a single Breit-Wigner resonance.				NODE=M092W;LINKAGE=CE
¹¹ In a model where a non-resonant $b\bar{b}$ -continuum represented by a threshold function at $\sqrt{s}=2m_B$ is incoherently added to a flat component interfering with two Breit-Wigner resonances. Not independent of other AUBERT 09E results. Systematic uncertainties not estimated.				NODE=M092W;LINKAGE=UB
¹² Assuming four Gaussians with radiative tails and a single step in R .				NODE=M092W;LINKAGE=BE
¹³ In a coupled-channel model with three resonances and a smooth step in R .				NODE=M092W;LINKAGE=LO

$\Upsilon(10860)$ DECAY MODES

NODE=M092215;NODE=M092

Mode	Fraction (Γ_i/Γ)	Confidence level	
Γ_1 $B\bar{B}X$	(76.2 ^{+2.7} _{-4.0}) %		DESIG=9
Γ_2 $B\bar{B}$	(5.5 ± 1.0) %		DESIG=2
Γ_3 $B\bar{B}^* + \text{c.c.}$	(13.7 ± 1.6) %		DESIG=3
Γ_4 $B^*\bar{B}^*$	(38.1 ± 3.4) %		DESIG=4
Γ_5 $B\bar{B}^{(*)}\pi$	< 19.7 %	90%	DESIG=10
Γ_6 $B\bar{B}\pi$	(0.0 ± 1.2) %		DESIG=23
Γ_7 $B^*\bar{B}\pi + B\bar{B}^*\pi$	(7.3 ± 2.3) %		DESIG=24
Γ_8 $B^*\bar{B}^*\pi$	(1.0 ± 1.4) %		DESIG=25
Γ_9 $B\bar{B}\pi\pi$	< 8.9 %	90%	DESIG=11
Γ_{10} $B_s^{(*)}\bar{B}_s^{(*)}$	(20.1 ± 3.1) %		DESIG=16
Γ_{11} $B_s\bar{B}_s$	(5 ± 5) × 10 ⁻³		DESIG=5
Γ_{12} $B_s\bar{B}_s^* + \text{c.c.}$	(1.35 ± 0.32) %		DESIG=7
Γ_{13} $B_s^*\bar{B}_s^*$	(17.6 ± 2.7) %		DESIG=8
Γ_{14} no open-bottom	(3.8 ^{+5.0} _{-0.5}) %		DESIG=28

Γ_{15}	$e^+ e^-$	$(8.3 \pm 2.1) \times 10^{-6}$		DESIG=1
Γ_{16}	$K^*(892)^0 \bar{K}^0$	$< 1.0 \times 10^{-5}$	90%	DESIG=29
Γ_{17}	$\Upsilon(1S) \pi^+ \pi^-$	$(5.3 \pm 0.6) \times 10^{-3}$		DESIG=17
Γ_{18}	$\Upsilon(1S) \eta$	$(8.5 \pm 1.7) \times 10^{-4}$		DESIG=44
Γ_{19}	$\Upsilon(1S) \eta'$	$< 6.9 \times 10^{-5}$	90%	DESIG=45
Γ_{20}	$\Upsilon(2S) \pi^+ \pi^-$	$(7.8 \pm 1.3) \times 10^{-3}$		DESIG=18
Γ_{21}	$\Upsilon(2S) \eta$	$(4.1 \pm 0.6) \times 10^{-3}$		DESIG=46
Γ_{22}	$\Upsilon(3S) \pi^+ \pi^-$	$(4.8 \begin{smallmatrix} +1.9 \\ -1.7 \end{smallmatrix}) \times 10^{-3}$		DESIG=19
Γ_{23}	$\Upsilon(1S) K^+ K^-$	$(6.1 \pm 1.8) \times 10^{-4}$		DESIG=20
Γ_{24}	$\eta \Upsilon_J(1D)$	$(4.8 \pm 1.1) \times 10^{-3}$		DESIG=40
Γ_{25}	$h_b(1P) \pi^+ \pi^-$	$(3.5 \begin{smallmatrix} +1.0 \\ -1.3 \end{smallmatrix}) \times 10^{-3}$		DESIG=26
Γ_{26}	$h_b(2P) \pi^+ \pi^-$	$(5.7 \begin{smallmatrix} +1.7 \\ -2.1 \end{smallmatrix}) \times 10^{-3}$		DESIG=27
Γ_{27}	$\chi_{bJ}(1P) \pi^+ \pi^- \pi^0$	$(2.5 \pm 2.3) \times 10^{-3}$		DESIG=41
Γ_{28}	$\chi_{b0}(1P) \pi^+ \pi^- \pi^0$	$< 6.3 \times 10^{-3}$	90%	DESIG=30
Γ_{29}	$\chi_{b0}(1P) \omega$	$< 3.9 \times 10^{-3}$	90%	DESIG=31
Γ_{30}	$\chi_{b0}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	$< 4.8 \times 10^{-3}$	90%	DESIG=32
Γ_{31}	$\chi_{b1}(1P) \pi^+ \pi^- \pi^0$	$(1.85 \pm 0.33) \times 10^{-3}$		DESIG=33
Γ_{32}	$\chi_{b1}(1P) \omega$	$(1.57 \pm 0.30) \times 10^{-3}$		DESIG=34
Γ_{33}	$\chi_{b1}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	$(5.2 \pm 1.9) \times 10^{-4}$		DESIG=35
Γ_{34}	$\chi_{b2}(1P) \pi^+ \pi^- \pi^0$	$(1.17 \pm 0.30) \times 10^{-3}$		DESIG=36
Γ_{35}	$\chi_{b2}(1P) \omega$	$(6.0 \pm 2.7) \times 10^{-4}$		DESIG=37
Γ_{36}	$\chi_{b2}(1P) (\pi^+ \pi^- \pi^0)_{\text{non-}\omega}$	$(6 \pm 4) \times 10^{-4}$		DESIG=38
Γ_{37}	$\gamma X_b \rightarrow \gamma \Upsilon(1S) \omega$	$< 3.8 \times 10^{-5}$	90%	DESIG=39
Γ_{38}	$\eta_b(1S) \omega$	$< 1.3 \times 10^{-3}$	90%	DESIG=42
Γ_{39}	$\eta_b(2S) \omega$	$< 5.6 \times 10^{-3}$	90%	DESIG=43

Inclusive Decays.

NODE=M092;CLUMP=I

These decay modes are submodes of one or more of the decay modes above.

NODE=M092

Γ_{40}	ϕ anything	$(13.8 \begin{smallmatrix} +2.4 \\ -1.7 \end{smallmatrix}) \%$		DESIG=12
Γ_{41}	D^0 anything + c.c.	$(112 \pm 6) \%$		DESIG=13
Γ_{42}	D_s anything + c.c.	$(44.7 \pm 2.6) \%$		DESIG=6
Γ_{43}	J/ψ anything	$(2.06 \pm 0.21) \%$		DESIG=14
Γ_{44}	B^0 anything + c.c.	$(77 \pm 8) \%$		DESIG=21
Γ_{45}	B^+ anything + c.c.	$(72 \pm 6) \%$		DESIG=22

 $\Upsilon(10860)$ PARTIAL WIDTHS

NODE=M092220

 $\Gamma(e^+ e^-)$ **Γ_{15}**

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.07 OUR AVERAGE	Error includes scale factor of 1.3.		
0.22 ± 0.05 ± 0.07	BESSION	85	CLEO $e^+ e^- \rightarrow$ hadrons
0.365 ± 0.070	LOVELOCK	85	CUSB $e^+ e^- \rightarrow$ hadrons

NODE=M092W1
NODE=M092W1 **$\Gamma(e^+ e^-) \times \Gamma(\Upsilon(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}$** **$\Gamma_{15} \Gamma_{17} / \Gamma$**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M092R50
NODE=M092R50

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.09 ± 0.34	^{1,2} MIZUK	19	BELL $e^+ e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-$
-------------	----------------------	----	---

¹ From a simultaneous fit to the $\Upsilon(nS) \pi^+ \pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6$ –11.05 GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M092R50;LINKAGE=A

² Reported as the range 0.75–1.43 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M092R50;LINKAGE=B

 $\Gamma(e^+ e^-) \times \Gamma(\Upsilon(2S) \pi^+ \pi^-) / \Gamma_{\text{total}}$ **$\Gamma_{15} \Gamma_{20} / \Gamma$**

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M092R51
NODE=M092R51

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.58 ± 1.22	^{1,2} MIZUK	19	BELL $e^+ e^- \rightarrow \Upsilon(nS) \pi^+ \pi^-$
-------------	----------------------	----	---

¹ From a simultaneous fit to the $\Upsilon(nS) \pi^+ \pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6$ –11.05 GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M092R51;LINKAGE=A

² Reported as the range 1.35–3.80 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M092R51;LINKAGE=B

$$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_{15}\Gamma_{22}/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M092R52
NODE=M092R52

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.73±0.30	1,2 MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
-----------	-----------	----	------	---

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M092R52;LINKAGE=A

² Reported as the range 0.43–1.03 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M092R52;LINKAGE=B

$\Upsilon(10860)$ BRANCHING RATIOS

NODE=M092230

"OUR EVALUATION" is obtained based on averages of rescaled data listed below. The averages and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <https://hflav.web.cern.ch/>.

NODE=M092230

$$\Gamma(B\bar{B}X)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M092R13
NODE=M092R13

0.762^{+0.027}_{-0.043} OUR EVALUATION (Produced by HFLAV)

→ UNCHECKED ←

0.71 ± 0.06 OUR AVERAGE

0.737±0.032±0.051	1063	¹ DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
-------------------	------	-----------------------	----	------	---------------------------------------

NODE=M092R13;LINKAGE=DR
NODE=M092R13;LINKAGE=HU

0.589±0.100±0.092		² HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------------------	--	--------------------	----	------	---

¹ Not independent of DRUTSKOY 10 values for $\Upsilon(5S) \rightarrow B^{\pm,0}$ anything.

² Using measurements or limits from AQUINES 06.

$$\Gamma(B\bar{B})/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE (units 10 ⁻²)	CL%	DOCUMENT ID	TECN	COMMENT
---------------------------------	-----	-------------	------	---------

NODE=M092R16
NODE=M092R16

5.5^{+1.0}_{-0.9} ± 0.4		¹ DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
--	--	-----------------------	----	------	---------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<13.8	90	² HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------	----	--------------------	----	------	---

¹ Assuming isospin conservation.

² Using measurements or limits from AQUINES 06.

NODE=M092R16;LINKAGE=DR
NODE=M092R16;LINKAGE=HU

$$\Gamma(B\bar{B})/\Gamma(B\bar{B}X) \quad \Gamma_2/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M092R05
NODE=M092R05

<0.22	90	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$
-----------------	----	---------	----	------	---

$$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M092R15
NODE=M092R15

0.137±0.016 OUR AVERAGE

0.137±0.013±0.011		¹ DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
-------------------	--	-----------------------	----	------	---------------------------------------

NODE=M092R15;LINKAGE=DR
NODE=M092R15;LINKAGE=HU

0.143±0.053±0.027		² HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------------------	--	--------------------	----	------	---

¹ Assuming isospin conservation.

² Using measurements or limits from AQUINES 06.

$$\Gamma(B\bar{B}^* + \text{c.c.})/\Gamma(B\bar{B}X) \quad \Gamma_3/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M092R06
NODE=M092R06

0.24±0.09±0.03	10	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$
-----------------------	----	---------	----	------	---

$$\Gamma(B^*\bar{B}^*)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M092R14
NODE=M092R14

0.381±0.034 OUR AVERAGE

0.375 ^{+0.021} _{-0.019} ± 0.030		¹ DRUTSKOY	10	BELL	$\Upsilon(5S) \rightarrow B^+X, B^0X$
---	--	-----------------------	----	------	---------------------------------------

NODE=M092R14;LINKAGE=DR
NODE=M092R14;LINKAGE=HU

0.436±0.083±0.072		² HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
-------------------	--	--------------------	----	------	---

¹ Assuming isospin conservation.

² Using measurements or limits from AQUINES 06.

$$\Gamma(B^*\bar{B}^*)/\Gamma(B\bar{B}X) \quad \Gamma_4/\Gamma_1$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

NODE=M092R07
NODE=M092R07

0.74±0.15±0.08	31	AQUINES	06	CLE3	$\Upsilon(5S) \rightarrow \text{hadrons}$
-----------------------	----	---------	----	------	---

$$\Gamma(B\bar{B}^{(*)}\pi)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

NODE=M092R17
NODE=M092R17

<0.197	90	¹ HUANG	07	CLEO	$\Upsilon(5S) \rightarrow \text{hadrons}$
------------------	----	--------------------	----	------	---

¹ Using measurements or limits from AQUINES 06.

NODE=M092R17;LINKAGE=HU

$\Gamma(B\bar{B}^*\pi)/\Gamma(B\bar{B}X)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_1
<0.32	90	AQUINES	06	CLE3 $\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092R08
NODE=M092R08

$\Gamma(B\bar{B}\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
$0.0 \pm 1.2 \pm 0.3$	0	¹ DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+0}\pi^- X$	

NODE=M092R28
NODE=M092R28

¹ Assuming isospin conservation.

NODE=M092R28;LINKAGE=DR

$[\Gamma(B^*\bar{B}\pi) + \Gamma(B\bar{B}^*\pi)]/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
$7.3^{+2.3}_{-2.1} \pm 0.8$	38	¹ DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+0}\pi^- X$	

NODE=M092R29
NODE=M092R29

¹ Assuming isospin conservation.

NODE=M092R29;LINKAGE=DR

$\Gamma(B^*\bar{B}^*\pi)/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
$1.0^{+1.4}_{-1.3} \pm 0.4$	5	¹ DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^{+0}\pi^- X$	

NODE=M092R30
NODE=M092R30

¹ Assuming isospin conservation.

NODE=M092R30;LINKAGE=DR

$\Gamma(B\bar{B}\pi\pi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
<0.089	90	¹ HUANG	07	CLEO $\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092R18
NODE=M092R18

¹ Using measurements or limits from AQUINES 06.

NODE=M092R18;LINKAGE=HU

$\Gamma(B\bar{B}\pi\pi)/\Gamma(B\bar{B}X)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_1
<0.14	90	AQUINES	06	CLE3 $\Upsilon(5S) \rightarrow$ hadrons	

NODE=M092R09
NODE=M092R09

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_{10}/\Gamma = (\Gamma_{11} + \Gamma_{12} + \Gamma_{13})/\Gamma$
-------	-------------	------	---------	---

NODE=M092R01
NODE=M092R01

$0.201^{+0.030}_{-0.031}$ OUR EVALUATION

→ UNCHECKED ←

$0.189^{+0.027}_{-0.021}$ OUR AVERAGE

0.172 ± 0.030	¹ ESEN	13	BELL	$\Upsilon(5S) \rightarrow D^0 X, D_s X$
$0.21^{+0.06}_{-0.03}$	² HUANG	07	CLEO	$\Upsilon(5S) \rightarrow D_s X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.180 \pm 0.013 \pm 0.032$	³ DRUTSKOY	07	BELL	$\Upsilon(5S) \rightarrow D^0 X, D_s X$
$0.160 \pm 0.026 \pm 0.058$	⁴ ARTUSO	05B	CLEO	$e^+ e^- \rightarrow D_s X$

¹ Supersedes DRUTSKOY 07.

² Supersedes ARTUSO 05B. Combining inclusive ϕ , D_s , and B measurements. Using $B(D_s^+ \rightarrow \phi\pi^+) = 4.4 \pm 0.6\%$ from PDG 06.

³ Using $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6)\%$ from PDG 06.

⁴ Uses a model-dependent estimate $B(B_s \rightarrow D_s X) = (92 \pm 11)\%$.

NODE=M092R01;LINKAGE=ES
NODE=M092R01;LINKAGE=HU

NODE=M092R01;LINKAGE=DR
NODE=M092R01;LINKAGE=AR

$\Gamma(B_s^{(*)}\bar{B}_s^{(*)})/\Gamma(B\bar{B}X)$

VALUE	DOCUMENT ID	Γ_{10}/Γ_1
-------	-------------	------------------------

NODE=M092R34
NODE=M092R34

$0.264^{+0.052}_{-0.045}$ OUR EVALUATION

→ UNCHECKED ←

$\Gamma(B_s^*\bar{B}_s^*)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{13}/\Gamma_{10} = \Gamma_{13}/(\Gamma_{11} + \Gamma_{12} + \Gamma_{13})$
--------------------------	------	-------------	------	---------	---

NODE=M092R19
NODE=M092R19

87.8 ± 1.5 OUR AVERAGE

87.0 ± 1.7	^{1,2} ESEN	13	BELL	$B_s^0 \rightarrow D_s^- \pi^+$
$90.5 \pm 3.2 \pm 0.1$	^{2,3} LI	12	BELL	$B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$90.1^{+3.8}_{-4.0} \pm 0.2$	⁴ LOUVOT	09	BELL	$10.86 e^+ e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$
$93^{+7}_{-9} \pm 1$	⁴ DRUTSKOY	07A	BELL	Superseded by LOUVOT 09

¹Supersedes LOUVOT 09.

²With $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$.

³The ratios $N(B_s^{(*)}\bar{B}_s^{(*)}) / N(B_s^{(*)}\bar{B}_s^{(*)})$ and $N(B_s^{(*)}\bar{B}_s^{(*)}) / N(B_s^{(*)}\bar{B}_s^{(*)})$ are measured with a correlation coefficient of -0.72 .

⁴From a measurement of $\sigma(e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}) / \sigma(e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)})$ at $\sqrt{s} = 10.86$ GeV.

NODE=M092R19;LINKAGE=ES

NODE=M092R19;LINKAGE=IL

NODE=M092R19;LINKAGE=L1

NODE=M092R19;LINKAGE=DR

$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$		$\Gamma_{11}/\Gamma_{10} = \Gamma_{11}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$		
VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	
$2.6^{+2.6}_{-2.5}$	LOUVOT	09	BELL	$10.86 e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$

NODE=M092R24
NODE=M092R24

$\Gamma(B_s\bar{B}_s)/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$		Γ_{11}/Γ_{13}		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	BONVICINI	06	CLE3 e^+e^-

NODE=M092R03
NODE=M092R03

$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$		$\Gamma_{12}/\Gamma_{10} = \Gamma_{12}/(\Gamma_{11}+\Gamma_{12}+\Gamma_{13})$		
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
6.7 ± 1.2 OUR AVERAGE				

NODE=M092R25
NODE=M092R25

7.3 ± 1.4		^{1,2} ESEN	13	BELL	$B_s^0 \rightarrow D_s^- \pi^+$
$4.9 \pm 2.5 \pm 0.0$	227	^{2,3} LI	12	BELL	$B_s^0 \rightarrow J/\psi \eta^{(\prime)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.3^{+3.3}_{-3.0} \pm 0.1$		LOUVOT	09	BELL	$10.86 e^+e^- \rightarrow B_s^{(*)}\bar{B}_s^{(*)}$
-----------------------------	--	--------	----	------	---

¹Supersedes LOUVOT 09.

²With $N(B_s^{(*)}\bar{B}_s^{(*)}) = (7.11 \pm 1.30) \times 10^6$.

³The ratios $N(B_s^{(*)}\bar{B}_s^{(*)}) / N(B_s^{(*)}\bar{B}_s^{(*)})$ and $N(B_s^{(*)}\bar{B}_s^{(*)}) / N(B_s^{(*)}\bar{B}_s^{(*)})$ are measured with a correlation coefficient of -0.72 .

NODE=M092R25;LINKAGE=ES

NODE=M092R25;LINKAGE=IL

NODE=M092R25;LINKAGE=L1

$\Gamma(B_s\bar{B}_s^* + \text{c.c.})/\Gamma(B_s^{(*)}\bar{B}_s^{(*)})$		Γ_{12}/Γ_{13}		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	BONVICINI	06	CLE3 e^+e^-

NODE=M092R04
NODE=M092R04

$\Gamma(\text{no open-bottom})/\Gamma_{\text{total}}$		Γ_{14}/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.038^{+0.051}_{-0.005}$ OUR EVALUATION				

NODE=M092R33
NODE=M092R33

→ UNCHECKED ←

$\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma_{\text{total}}$		Γ_{16}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-5}$	90	SHEN	13A	BELL $e^+e^- \rightarrow K^*(892)^0 \bar{K}^0$

NODE=M092R35
NODE=M092R35

$\Gamma(\eta \mathcal{T}_J(1D))/\Gamma_{\text{total}}$		Γ_{24}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	
$4.82 \pm 0.92 \pm 0.67$	¹ TAMPONI	18	BELL	$e^+e^- \rightarrow \mathcal{T}(5S) \rightarrow \eta X$

NODE=M092R48
NODE=M092R48

¹Mainly $J = 2$, assumes no continuum contribution under $\mathcal{T}(5S)$.

NODE=M092R48;LINKAGE=A

$\Gamma(\mathcal{T}(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$		Γ_{17}/Γ		
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$5.3 \pm 0.3 \pm 0.5$	325	¹ CHEN	08	BELL $10.87 e^+e^- \rightarrow \mathcal{T}(1S)\pi^+\pi^-$

NODE=M092R20
NODE=M092R20

¹Assuming that the observed events are solely due to the $\mathcal{T}(5S)$ resonance.

NODE=M092R20;LINKAGE=CH

$\Gamma(\mathcal{T}(1S)\eta)/\Gamma_{\text{total}}$		Γ_{18}/Γ		
VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT	
$0.85 \pm 0.15 \pm 0.08$	^{1,2} KOVALENKO	21	BELL	$e^+e^- \rightarrow \mathcal{T}(5S)$

NODE=M092R55
NODE=M092R55

¹Assuming that the observed events are solely due to the $\mathcal{T}(5S)$ resonance.

NODE=M092R55;LINKAGE=A

²Using a data sample of 118.3 fb^{-1} of e^+e^- collisions at $\sqrt{s} = 10.866$ GeV.

NODE=M092R55;LINKAGE=B

$\Gamma(\mathcal{T}(1S)\eta')/\Gamma_{\text{total}}$		Γ_{19}/Γ		
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.9 \times 10^{-5}$	90	^{1,2} KOVALENKO	21	BELL $e^+e^- \rightarrow \mathcal{T}(5S)$

NODE=M092R56
NODE=M092R56

¹Assuming that the observed events are solely due to the $\mathcal{T}(5S)$ resonance.

NODE=M092R56;LINKAGE=A

²Using a data sample of 118.3 fb^{-1} of e^+e^- collisions at $\sqrt{s} = 10.866$ GeV.

NODE=M092R56;LINKAGE=B

$\Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$7.8 \pm 0.6 \pm 1.1$	186	¹ CHEN 08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

NODE=M092R21
NODE=M092R21

¹ Assuming that the observed events are solely due to the $\Upsilon(5S)$ resonance.

NODE=M092R21;LINKAGE=CH

 $\Gamma(\Upsilon(2S)\eta)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$4.13 \pm 0.41 \pm 0.37$	^{1,2} KOVALENKO 21	BELL	$e^+ e^- \rightarrow \Upsilon(5S)$

NODE=M092R57
NODE=M092R57

¹ Assuming that the observed events are solely due to the $\Upsilon(5S)$ resonance.

NODE=M092R57;LINKAGE=A

² Using a data sample of 118.3 fb^{-1} of $e^+ e^-$ collisions at $\sqrt{s} = 10.866 \text{ GeV}$.

NODE=M092R57;LINKAGE=B

 $\Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$4.8 \pm 1.8 \pm 0.7$	10	¹ CHEN 08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

NODE=M092R22
NODE=M092R22

¹ Assuming that the observed events are solely due to the $\Upsilon(5S)$ resonance.

NODE=M092R22;LINKAGE=CH

 $\Gamma(\Upsilon(1S)K^+K^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
$6.1 \pm 1.6 \pm 1.0$	20	¹ CHEN 08	BELL	$10.87 e^+ e^- \rightarrow \Upsilon(1S)K^+K^-$

NODE=M092R23
NODE=M092R23

¹ Assuming that the observed events are solely due to the $\Upsilon(5S)$ resonance.

NODE=M092R23;LINKAGE=CH

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$ Γ_{25}/Γ_{20}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.45 \pm 0.08 \pm 0.07$ -0.12	ADACHI 12	BELL	$10.86 e^+ e^- \rightarrow \text{hadrons}$

NODE=M092R31
NODE=M092R31

 $\Gamma(h_b(2P)\pi^+\pi^-)/\Gamma(\Upsilon(2S)\pi^+\pi^-)$ Γ_{26}/Γ_{20}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.77 \pm 0.08 \pm 0.22$ -0.17	ADACHI 12	BELL	$10.86 e^+ e^- \rightarrow \text{hadrons}$

NODE=M092R32
NODE=M092R32

 $\Gamma(h_b(1P)\pi^+\pi^-)/\Gamma(h_b(2P)\pi^+\pi^-)$ Γ_{25}/Γ_{26}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.616 \pm 0.052 \pm 0.017$	MIZUK 16	BELL	$e^+ e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$

NODE=M092R00
NODE=M092R00

 $\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$2.5 \pm 0.6 \pm 2.2$	YIN 18	BELL	$e^+ e^- \rightarrow \text{hadrons}$

NODE=M092R49
NODE=M092R49

 $\Gamma(\chi_{b0}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-3}$	90	¹ HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R36
NODE=M092R36

¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$ from ESEN 13. Correlated with other results from HE 14.

NODE=M092R36;LINKAGE=A

 $\Gamma(\chi_{b0}(1P)\omega)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.9 \times 10^{-3}$	90	¹ HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R37
NODE=M092R37

¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$ from ESEN 13. Correlated with other results from HE 14.

NODE=M092R37;LINKAGE=A

 $\Gamma(\chi_{b0}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.8 \times 10^{-3}$	90	¹ HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R38
NODE=M092R38

¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$ from ESEN 13. Correlated with other results from HE 14.

NODE=M092R38;LINKAGE=A

 $\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
$1.85 \pm 0.23 \pm 0.23$	80	¹ HE 14	BELL	$\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$

NODE=M092R39
NODE=M092R39

¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+ e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016 \text{ nb}$ from ESEN 13. Correlated with other results from HE 14.

NODE=M092R39;LINKAGE=A

$\Gamma(\chi_{b1}(1P)\omega)/\Gamma_{\text{total}}$					Γ_{32}/Γ	NODE=M092R40 NODE=M092R40
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.57±0.22±0.21	60	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.						
$\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$					Γ_{33}/Γ	NODE=M092R41 NODE=M092R41
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.52±0.15±0.11	24	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.						
$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$					Γ_{34}/Γ	NODE=M092R42 NODE=M092R42
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
1.17±0.27±0.14	29	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.						
$\Gamma(\chi_{b2}(1P)\omega)/\Gamma_{\text{total}}$					Γ_{35}/Γ	NODE=M092R43 NODE=M092R43
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.60±0.23±0.15	13	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.						
$\Gamma(\chi_{b2}(1P)\omega)/\Gamma(\chi_{b1}(1P)\omega)$					Γ_{35}/Γ_{32}	NODE=M092R44 NODE=M092R44
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.38±0.16±0.09	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$			
¹ Accounting for correlated systematics.						
$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma_{\text{total}}$					Γ_{36}/Γ	NODE=M092R45 NODE=M092R45
<u>VALUE (units 10^{-3})</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
0.61±0.22±0.28	16	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14.						
$\Gamma(\chi_{b2}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})/\Gamma(\chi_{b1}(1P)(\pi^+\pi^-\pi^0)_{\text{non-}\omega})$					Γ_{36}/Γ_{33}	NODE=M092R46 NODE=M092R46
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>			
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1.20±0.55±0.65	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$			
¹ Accounting for correlated systematics.						
$\Gamma(\eta_b(1S)\omega)/\Gamma_{\text{total}}$					Γ_{38}/Γ	NODE=M092R53 NODE=M092R53
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<1.3 × 10⁻³	90	¹ OSKIN	20	BELL $e^+e^- \rightarrow \omega X$		
¹ Using $\sigma_{b\bar{b}} = 0.340 \pm 0.016$ nb from TAMPONI 15.						
$\Gamma(\eta_b(2S)\omega)/\Gamma_{\text{total}}$					Γ_{39}/Γ	NODE=M092R54 NODE=M092R54
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<5.6 × 10⁻³	90	¹ OSKIN	20	BELL $e^+e^- \rightarrow \omega X$		
¹ Using $\sigma_{b\bar{b}} = 0.340 \pm 0.016$ nb from TAMPONI 15.						
$\Gamma(\gamma X_b \rightarrow \gamma \Upsilon(1S)\omega)/\Gamma_{\text{total}}$					Γ_{37}/Γ	NODE=M092R47 NODE=M092R47
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>		
<3.8 × 10⁻⁵	90	¹ HE	14	BELL $\Upsilon(5S) \rightarrow \pi^+\pi^-\pi^0\gamma\Upsilon(1S)$		
¹ Assuming that all the $b\bar{b}$ events are from $\Upsilon(5S)$ resonance decays and using $\sigma(e^+e^- \rightarrow b\bar{b}) = 0.340 \pm 0.016$ nb from ESEN 13. Correlated with other results from HE 14. For a state X_b with mass between 10.55 GeV/ c^2 and 10.65 GeV/ c^2 , the obtained 90% upper limit as a function of m_{X_b} varies from 2.6×10^{-5} to 3.8×10^{-5} .						

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{40}/Γ
$0.138 \pm 0.007^{+0.023}_{-0.015}$	HUANG	07	CLEO $\Upsilon(5S) \rightarrow \phi X$	NODE=M092R12 NODE=M092R12

 $\Gamma(D^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{41}/Γ
1.12 ± 0.06 OUR AVERAGE [1.08 ± 0.08 OUR 2023 AVERAGE]				NODE=M092R10 NODE=M092R10 NEW

1.117 ± 0.005 ± 0.060 ¹ ZHUKOVA 23 BELL $\Upsilon(5S) \rightarrow D^0 X, \bar{D}^0 X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.076 ± 0.040 ± 0.068 DRUTSKOY 07 BELL $\Upsilon(5S) \rightarrow D^0 X, \bar{D}^0 X$

¹ Supersedes DRUTSKOY 07.

NODE=M092R10;LINKAGE=A

 $\Gamma(D_s \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{42}/Γ
0.447 ± 0.026 OUR AVERAGE [0.46 ± 0.06 OUR 2023 AVERAGE]				NODE=M092R02 NODE=M092R02 NEW

0.447 ± 0.003 ± 0.027 ¹ ZHUKOVA 23 BELL $\Upsilon(5S) \rightarrow D_s^\pm X$

0.44 ± 0.09 ± 0.04 ² ARTUSO 05B CLE3 $\Upsilon(5S) \rightarrow D_s^\pm X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.472 ± 0.024 ± 0.072 ³ DRUTSKOY 07 BELL $\Upsilon(5S) \rightarrow D_s^\pm X$

¹ Supersedes DRUTSKOY 07.

² ARTUSO 05B reports $[\Upsilon(10860) \rightarrow D_s \text{ anything} + \text{c.c.}]/\Gamma_{\text{total}} \times [B(D_s^+ \rightarrow \phi\pi^+)] = 0.0198 \pm 0.0019 \pm 0.0038$ which we divide by our best value $B(D_s^+ \rightarrow \phi\pi^+) = (4.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

³ Using $B(D_s^+ \rightarrow \phi\pi^+) = (4.4 \pm 0.6)\%$ from PDG 06.

NODE=M092R02;LINKAGE=A
NODE=M092R02;LINKAGE=AR

NODE=M092R02;LINKAGE=DR

 $\Gamma(J/\psi \text{ anything})/\Gamma_{\text{total}}$

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT	Γ_{43}/Γ
2.060 ± 0.160 ± 0.134	DRUTSKOY	07	BELL $\Upsilon(5S) \rightarrow J/\psi X$	NODE=M092R11 NODE=M092R11

 $\Gamma(B^0 \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{44}/Γ
$0.770^{+0.058}_{-0.056} \pm 0.061$	352	DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^0 X$	NODE=M092R26 NODE=M092R26

 $\Gamma(B^+ \text{ anything} + \text{c.c.})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{45}/Γ
$0.721^{+0.039}_{-0.038} \pm 0.050$	711	DRUTSKOY	10	BELL $\Upsilon(5S) \rightarrow B^+ X$	NODE=M092R27 NODE=M092R27

 $\Upsilon(10860)$ REFERENCES

Author	Year	Journal	Volume	Page	Collab.	REFID	
ZHUKOVA	23	JHEP	2308	131	V. Zhukova <i>et al.</i>	(BELLE Collab.)	REFID=62312
KOVALENKO	21	PR	D104	112006	E. Kovalenko <i>et al.</i>	(BELLE Collab.)	REFID=61452
DONG	20A	CP	C44	083001	X.-K. Dong <i>et al.</i>		REFID=60595
OSKIN	20	PR	D102	092011	P. Oskin <i>et al.</i>	(BELLE Collab.)	REFID=60735
MIZUK	19	JHEP	1910	220	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=60090
TAMPONI	18	EPJ	C78	633	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=59195
YIN	18	PR	D98	091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)	REFID=59468
MIZUK	16	PRL	117	142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
SANTEL	16	PR	D93	011101	D. Santel <i>et al.</i>	(BELLE Collab.)	REFID=57121
TAMPONI	15	PRL	115	142001	U. Tamponi <i>et al.</i>	(BELLE Collab.)	REFID=56996
HE	14	PRL	113	142001	X.H. He <i>et al.</i>	(BELLE Collab.)	REFID=55927
ESEN	13	PR	D87	031101	S. Esen <i>et al.</i>	(BELLE Collab.)	REFID=54894
SHEN	13A	PR	D88	052019	C.P. Shen <i>et al.</i>	(BELLE Collab.)	REFID=55591
ADACHI	12	PRL	108	032001	I. Adachi <i>et al.</i>	(BELLE Collab.)	REFID=53962
LI	12	PRL	108	181808	J. Li <i>et al.</i>	(BELLE Collab.)	REFID=54116
CHEN	10	PR	D82	091106	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53531
DRUTSKOY	10	PR	D81	112003	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=53358
AUBERT	09E	PRL	102	012001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52661
LOUVOT	09	PRL	102	021801	R. Louvot <i>et al.</i>	(BELLE Collab.)	REFID=52646
CHEN	08	PRL	100	112001	K.-F. Chen <i>et al.</i>	(BELLE Collab.)	REFID=52153
DRUTSKOY	07	PRL	98	052001	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51621
DRUTSKOY	07A	PR	D76	012002	A. Drutskoy <i>et al.</i>	(BELLE Collab.)	REFID=51852
HUANG	07	PR	D75	012002	G.S. Huang <i>et al.</i>	(CLEO Collab.)	REFID=51624
AQUINES	06	PRL	96	152001	O. Aquines <i>et al.</i>	(CLEO Collab.)	REFID=51106
BONVICINI	06	PRL	96	022002	G. Bonvicini <i>et al.</i>	(CLEO Collab.)	REFID=50995
PDG	06	JP	G33	1	W.-M. Yao <i>et al.</i>	(PDG Collab.)	REFID=51004
ARTUSO	05B	PRL	95	261801	M. Artuso <i>et al.</i>	(CLEO Collab.)	REFID=50992
BESSION	85	PRL	54	381	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=22368
LOVELOCK	85	PRL	54	377	D.M.J. Lovelock <i>et al.</i>	(CUSB Collab.)	REFID=22369

NODE=M092

$\Upsilon(11020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

NODE=M093

 $\Upsilon(11020)$ MASS

NODE=M093M

NODE=M093M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
11000 ± 4 OUR AVERAGE			
11000.0 ^{+4.0} _{-4.5} ^{+1.0} _{-1.3}	1 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10999.0 ^{+7.3} _{-7.8} ^{+16.9} _{-1.0}	2 MIZUK	16 BELL	$e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
11001 ± 1	3 DONG	20A	$e^+e^- \rightarrow b\bar{b}$
11003.0 ± 1.1 ⁺ _{-1.0} ^{+0.9} _{-1.0}	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow$ hadrons
10987.5 ^{+6.4} _{-2.5} ^{+9.1} _{-2.3}	6,7 SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
10996 ± 2	8 AUBERT	09E BABR	$e^+e^- \rightarrow$ hadrons
11019 ± 5 ± 7	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
11020 ± 30	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

OCCUR=2

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M093M;LINKAGE=E

² From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude.

NODE=M093M;LINKAGE=D

³ From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths.

NODE=M093M;LINKAGE=G

⁴ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient).

NODE=M093M;LINKAGE=A

⁵ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions.

NODE=M093M;LINKAGE=B

⁶ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero.

NODE=M093M;LINKAGE=C

⁷ Superseded by MIZUK 19.

NODE=M093M;LINKAGE=F

⁸ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated.

NODE=M093M;LINKAGE=AU

 $\Upsilon(11020)$ WIDTH

NODE=M093W

NODE=M093W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
24 ⁺⁸₋₆ OUR AVERAGE			
23.8 ^{+8.0} _{-6.8} ^{+0.7} _{-1.8}	1 MIZUK	19 BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
27 ⁺²⁷ ₋₁₁ ⁺⁵ ₋₁₂	2 MIZUK	16 BELL	$e^+e^- \rightarrow h_b(1P, 2P)\pi^+\pi^-$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
35.1 ± 1.2	3 DONG	20A	$e^+e^- \rightarrow b\bar{b}$
39.3 ^{+1.7} _{-1.6} ^{+1.3} _{-2.4}	4,5 SANTEL	16 BELL	$e^+e^- \rightarrow$ hadrons
61 ⁺⁹ ₋₁₉ ⁺² ₋₂₀	6,7 SANTEL	16 BELL	$e^+e^- \rightarrow \Upsilon(1S, 2S, 3S)\pi^+\pi^-$
37 ± 3	8 AUBERT	09E BABR	$e^+e^- \rightarrow$ hadrons
61 ± 13 ± 22	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
90 ± 20	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

OCCUR=2

- ¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$. NODE=M093W;LINKAGE=E
- ² From a simultaneous fit to the $h_b(nP)\pi^+\pi^-$, $n = 1, 2$ cross sections at 22 energy points within $\sqrt{s} = 10.77\text{--}11.02$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with eight resonance parameters (a mass and width for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single relative phase, a single relative amplitude, and two overall normalization factors, one for each n). The systematic error estimate is dominated by possible interference with a small nonresonant continuum amplitude. NODE=M093W;LINKAGE=D
- ³ From a fit to the dressed cross sections of AUBERT 09E by BaBar and SANTEL 16 by Belle above 10.68 GeV with a coherent sum of a continuum amplitude and three Breit-Wigner functions with constant widths. NODE=M093W;LINKAGE=G
- ⁴ From a fit to the total hadronic cross sections measured at 60 energy points within $\sqrt{s} = 10.82\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes and two floating continuum amplitudes with $1/\sqrt{s}$ dependence, one coherent with the resonances and one incoherent, with six resonance parameters (a mass, width, and an amplitude for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, one relative phase, and one decoherence coefficient). NODE=M093W;LINKAGE=A
- ⁵ Not including uncertain and potentially large systematic errors due to assumed continuum amplitude $1/\sqrt{s}$ dependence and related interference contributions. NODE=M093W;LINKAGE=B
- ⁶ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n=1, 2, 3$, cross sections at 25 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV to a pair of interfering Breit-Wigner amplitudes modified by phase space factors, with fourteen resonance parameters (a mass, width, and three amplitudes for each of $\Upsilon(10860)$ and $\Upsilon(11020)$, a single universal relative phase, and three decoherence coefficients, one for each n). Continuum contributions were measured (and therefore fixed) to be zero. NODE=M093W;LINKAGE=C
- ⁷ Superseded by MIZUK 19. NODE=M093W;LINKAGE=F
- ⁸ In a model where a flat non-resonant $b\bar{b}$ -continuum is incoherently added to a second flat component interfering with two Breit-Wigner resonances. Systematic uncertainties not estimated. NODE=M093W;LINKAGE=AU

$\Upsilon(11020)$ DECAY MODES

NODE=M093215;NODE=M093

Mode	Fraction (Γ_i/Γ)	
Γ_1 e^+e^-	$(5.4^{+1.9}_{-2.1}) \times 10^{-6}$	DESIG=1
Γ_2 $\Upsilon(1S)\pi^+\pi^-$		DESIG=5
Γ_3 $\Upsilon(2S)\pi^+\pi^-$		DESIG=6
Γ_4 $\Upsilon(3S)\pi^+\pi^-$		DESIG=7
Γ_5 $\chi_{bJ}(1P)\pi^+\pi^-\pi^0$	$(9^{+9}_{-8}) \times 10^{-3}$	DESIG=2
Γ_6 $\chi_{b1}(1P)\pi^+\pi^-\pi^0$	seen	DESIG=3
Γ_7 $\chi_{b2}(1P)\pi^+\pi^-\pi^0$	seen	DESIG=4

$\Upsilon(11020)$ PARTIAL WIDTHS

NODE=M093220

$\Gamma(e^+e^-)$					Γ_1	
VALUE (keV)	DOCUMENT ID	TECN	COMMENT			
0.130 ± 0.030 OUR AVERAGE						NODE=M093W1 NODE=M093W1
0.095 ± 0.03 ± 0.035	BESSION	85	CLEO $e^+e^- \rightarrow$ hadrons			
0.156 ± 0.040	LOVELOCK	85	CUSB $e^+e^- \rightarrow$ hadrons			

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_2/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN	COMMENT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.46 ± 0.08 ^{1,2} MIZUK 19 BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$. NODE=M093R04;LINKAGE=A

² Reported as the range 0.38–0.54 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions. NODE=M093R04;LINKAGE=B

$\Gamma(e^+e^-) \times \Gamma(\Upsilon(2S)\pi^+\pi^-)/\Gamma_{\text{total}}$					$\Gamma_1\Gamma_3/\Gamma$	
VALUE (eV)	DOCUMENT ID	TECN	COMMENT			

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.65 ± 0.52 ^{1,2} MIZUK 19 BELL $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$. NODE=M093R05;LINKAGE=A

² Reported as the range 0.13–1.16 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions. NODE=M093R05;LINKAGE=B

$$\Gamma(e^+e^-) \times \Gamma(\Upsilon(3S)\pi^+\pi^-)/\Gamma_{\text{total}} \quad \Gamma_1\Gamma_4/\Gamma$$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

NODE=M093R06
NODE=M093R06

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.33±0.16	^{1,2} MIZUK	19	BELL	$e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$
-----------	----------------------	----	------	---

¹ From a simultaneous fit to the $\Upsilon(nS)\pi^+\pi^-$, $n = 1, 2, 3$, cross sections at 28 energy points within $\sqrt{s} = 10.6\text{--}11.05$ GeV, including the initial-state radiation at $\Upsilon(10860)$.

NODE=M093R06;LINKAGE=A

² Reported as the range 0.17–0.49 eV obtained from multiple solutions of an amplitude fit within a model composed as a sum of Breit-Wigner functions.

NODE=M093R06;LINKAGE=B

$$\Gamma(\chi_{bJ}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
---------------------------------	-------------	------	---------

NODE=M093R00
NODE=M093R00

$8.7 \pm 4.3^{+7.6}_{-6.6}$	YIN	18	BELL	$e^+e^- \rightarrow \text{hadrons}$
-----------------------------	-----	----	------	-------------------------------------

$$\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M093R01
NODE=M093R01

seen	YIN	18	BELL	$e^+e^- \rightarrow \text{hadrons}$
------	-----	----	------	-------------------------------------

$$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma_{\text{total}} \quad \Gamma_7/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M093R02
NODE=M093R02

seen	YIN	18	BELL	$e^+e^- \rightarrow \text{hadrons}$
------	-----	----	------	-------------------------------------

$$\Gamma(\chi_{b2}(1P)\pi^+\pi^-\pi^0)/\Gamma(\chi_{b1}(1P)\pi^+\pi^-\pi^0) \quad \Gamma_7/\Gamma_6$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=M093R03
NODE=M093R03

0.4 ± 0.2	YIN	18	BELL	$e^+e^- \rightarrow \text{hadrons}$
---------------	-----	----	------	-------------------------------------

$\Upsilon(11020)$ REFERENCES

DONG	20A	CP C44 083001	X.-K. Dong <i>et al.</i>	
MIZUK	19	JHEP 1910 220	R. Mizuk <i>et al.</i>	(BELLE Collab.)
YIN	18	PR D98 091102	J.H. Yin <i>et al.</i>	(BELLE Collab.)
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)
SANTEL	16	PR D93 011101	D. Santel <i>et al.</i>	(BELLE Collab.)
AUBERT	09E	PRL 102 012001	B. Aubert <i>et al.</i>	(BABAR Collab.)
BESSION	85	PRL 54 381	D. Besson <i>et al.</i>	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	D.M.J. Lovelock <i>et al.</i>	(CUSP Collab.)

NODE=M093

REFID=60595
REFID=60090
REFID=59468
REFID=57465
REFID=57121
REFID=52661
REFID=22368
REFID=22369

OTHER MESONS

NODE=MXXX050

$$T_{cs0}^*(2870)^0$$

$$I(J^P) = ?(0^+)$$

NODE=M250

OMITTED FROM SUMMARY TABLE
was $X_0(2900)$

An exotic state with minimal quark content $\bar{c}d\bar{s}u$. Observed by AAIJ 20A1 using full amplitude analysis of $B^+ \rightarrow D^+ D^- K^+$ decays.

NODE=M250

$T_{cs0}^*(2870)^0$ MASS

NODE=M250M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M250M

$2866 \pm 7 \pm 2$	1.2k	¹ AAIJ	20A1	LHCB $B^+ \rightarrow D^+ D^- K^+$
--------------------	------	-------------------	------	------------------------------------

¹ Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M250M;LINKAGE=A

$T_{cs0}^*(2870)^0$ WIDTH

NODE=M250W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M250W

$57 \pm 12 \pm 4$	1.2k	¹ AAIJ	20A1	LHCB $B^+ \rightarrow D^+ D^- K^+$
-------------------	------	-------------------	------	------------------------------------

¹ Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M250W;LINKAGE=A

$T_{cs0}^*(2870)^0$ DECAY MODES

NODE=M250215;NODE=M250

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^- K^+$	seen

DESIG=1

 $T_{cs0}^*(2870)^0$ BRANCHING RATIOS

NODE=M250225

$\Gamma(D^- K^+)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M250R01
NODE=M250R01 $T_{cs0}^*(2870)^0$ REFERENCES

NODE=M250

AAIJ	20AF PRL 125 242001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20AI PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=60702
REFID=60739 $T_{cs1}^*(2900)^0$

$I(J^P) = ?(1^-)$

NODE=M251

OMITTED FROM SUMMARY TABLE
was $X_1(2900)$

An exotic state with minimal quark content $\bar{c}d\bar{s}u$. Observed by AAIJ 20AI using full amplitude analysis of $B^+ \rightarrow D^+ D^- K^+$ decays.

NODE=M251

 $T_{cs1}^*(2900)^0$ MASS

NODE=M251M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2904 \pm 5 \pm 1$	1.2k	¹ AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M251M

¹Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M251M;LINKAGE=B

 $T_{cs1}^*(2900)^0$ WIDTH

NODE=M251W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$110 \pm 11 \pm 4$	1.2k	¹ AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M251W

¹Obtained from the full amplitude analysis. Parameterized with the relativistic Breit-Wigner line shape. Also confirmed by the model-independent analysis of AAIJ 20AF.

NODE=M251W;LINKAGE=A

 $T_{cs1}^*(2900)^0$ DECAY MODES

NODE=M251215;NODE=M251

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^- K^+$	seen

DESIG=1

 $T_{cs1}^*(2900)^0$ BRANCHING RATIOS

NODE=M251225

$\Gamma(D^- K^+)/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAIJ	20AI LHCB	$B^+ \rightarrow D^+ D^- K^+$

NODE=M251R01
NODE=M251R01 $T_{cs1}^*(2900)^0$ REFERENCES

NODE=M251

AAIJ	20AF PRL 125 242001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20AI PR D102 112003	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=60702
REFID=60739

$T_{c\bar{s}0}^*(2900)$

$I(J^P) = 1(0^+)$

NODE=M269

OMITTED FROM SUMMARY TABLE

Observed by LHCb in AAIJ 23B using a simultaneous amplitude analysis of $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^+ \rightarrow D^- D_s^+ \pi^+$. The $T_{c\bar{s}0}^*(2900)^0 \rightarrow D_s^+ \pi^-$ and $T_{c\bar{s}0}^*(2900)^{++} \rightarrow D_s^+ \pi^+$ decays are observed with 8.0 and 6.5 σ significance, respectively.

NODE=M269

 $T_{c\bar{s}0}^*(2900)^0$ MASS

NODE=M269M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2892 ± 14 ± 15	¹ AAIJ	23C LHCB	$B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$

NODE=M269M

¹ From an amplitude analysis of $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$. A simultaneous fit to $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^- \rightarrow D^- D_s^+ \pi^+$ assuming isospin symmetry yields a mass of $2908 \pm 11 \pm 20$ MeV.

NODE=M269M;LINKAGE=A

 $T_{c\bar{s}0}^*(2900)^{++}$ MASS

NODE=M269M++

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2921 ± 17 ± 20	² AAIJ	23C LHCB	$B^- \rightarrow D^- D_s^+ \pi^+$

NODE=M269M++

² From an amplitude analysis of $B^- \rightarrow D^- D_s^+ \pi^+$. A simultaneous fit to $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^- \rightarrow D^- D_s^+ \pi^+$ assuming isospin symmetry yields a mass of $2908 \pm 11 \pm 20$ MeV.

NODE=M269M++;LINKAGE=A

 $T_{c\bar{s}0}^*(2900)^0$ WIDTH

NODE=M269W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
119 ± 26 ± 13	³ AAIJ	23C LHCB	$B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$

NODE=M269W

³ From an amplitude analysis of $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$. A simultaneous fit to $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^- \rightarrow D^- D_s^+ \pi^+$ assuming isospin symmetry yields a width of $136 \pm 23 \pm 13$ MeV.

NODE=M269W;LINKAGE=A

 $T_{c\bar{s}0}^*(2900)^{++}$ WIDTH

NODE=M269W++

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
137 ± 32 ± 17	⁴ AAIJ	23C LHCB	$B^- \rightarrow D^- D_s^+ \pi^+$

NODE=M269W++

⁴ From an amplitude analysis of $B^- \rightarrow D^- D_s^+ \pi^+$. A simultaneous fit to $B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$ and $B^- \rightarrow D^- D_s^+ \pi^+$ assuming isospin symmetry yields a width of $136 \pm 23 \pm 13$ MeV.

NODE=M269W++;LINKAGE=A

Mode	Fraction (Γ_i/Γ)
Γ_1 $D_s^+ \pi^-$	seen
Γ_2 $D_s^+ \pi^+$	seen

NODE=M269215;DESIG=1

DESIG=2

 $T_{c\bar{s}0}^*(2900)$ BRANCHING RATIOS

NODE=M269225

$\Gamma(D_s^+ \pi^-)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		AAIJ	23C LHCB	$B^0 \rightarrow \bar{D}^0 D_s^+ \pi^-$	

NODE=M269R01
NODE=M269R01

$\Gamma(D_s^+ \pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen		AAIJ	23C LHCB	$B^- \rightarrow D^- D_s^+ \pi^+$	

NODE=M269R02
NODE=M269R02

$T_{cc}(3875)^+$

$I(J^P) = ?(??)$

NODE=M265

OMITTED FROM SUMMARY TABLE

Observed with large significance by AAIJ 22E in the doubly-charmed ($C = 2$) decay mode $D^0 D^0 \pi^+$ using inclusive pp collisions at 7, 8, and 13 TeV.

NODE=M265

 $T_{cc}(3875)^+$ MASS

NODE=M265M

OUR FIT value comes from the measurement of $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$ below and $m_{D^{*+}} + m_{D^0}$ values.

NODE=M265M

VALUE (MeV)

DOCUMENT ID

NODE=M265M

3874.83 ± 0.11 OUR FIT $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$ NODE=M265DM
NODE=M265DM

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

= 0.27 ± 0.06 OUR FIT**= 0.273 ± 0.061^{+0.012}_{-0.015}** 117 1 AAIJ 22E LHCB $pp \rightarrow D^0 D^0 \pi^+ X$

¹ The fit assumes a relativistic P -wave Breit Wigner function modified by Blatt-Weisskopf form factor with radius 3.5 GeV⁻¹. The significance for $m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0}) < 0$ is 4.3 σ .

NODE=M265DM;LINKAGE=A

 $T_{cc}(3875)^+$ WIDTH

NODE=M265W

VALUE (MeV)

EVTS

DOCUMENT ID

TECN

COMMENT

NODE=M265W

0.410 ± 0.165^{+0.047}_{-0.057} 117 1 AAIJ 22E LHCB $pp \rightarrow D^0 D^0 \pi^+ X$

¹ The fit assumes a relativistic P -wave Breit Wigner function modified by Blatt-Weisskopf form factor with radius 3.5 GeV⁻¹.

NODE=M265W;LINKAGE=A

 $T_{cc}(3875)^+$ DECAY MODES

NODE=M265215;NODE=M265

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 D^0 \pi^+$	seen

DESIG=1

 $T_{cc}(3875)^+$ BRANCHING RATIOS

NODE=M265225

$\Gamma(D^0 D^0 \pi^+)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	117	AAIJ	22E LHCB	$pp \rightarrow D^0 D^0 \pi^+ X$

NODE=M265R01
NODE=M265R01 $T_{cc}(3875)^+$ REFERENCES

NODE=M265

AAIJ 22E NATP 18 751 R. Aaij *et al.* (LHCb Collab.)

REFID=61658

$T_{c\bar{c}1}(3900)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

was $Z_c(3900)$, $X(3900)$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

Charged $T_{c\bar{c}1}(3900)^\pm$ seen as a peak in the invariant mass distribution of the $J/\psi\pi^\pm$ system by BES III (ABLIKIM 13T) in $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ at c.m. energy of 4.26 GeV and by radiative return from e^+e^- collisions at \sqrt{s} from 9.46 to 10.86 GeV at Belle (LIU 13B). Partial wave analysis of ABLIKIM 17J determines $J^P = 1^+$ with more than 7σ significance. Neutral $T_{c\bar{c}1}(3900)^0$ seen in the $J/\psi\pi^0$ invariant mass distribution in $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$ at c.m. energies of 4.23, 4.26, and 4.36 GeV by BES III (ABLIKIM 15U) and at 4.17 GeV by XIAO 13A. Peaks in $(D\bar{D}^*)^{0,\pm}$ reported by BES III (ABLIKIM 14A, ABLIKIM 15AB) are assumed to be related.

NODE=M210

NODE=M210

 $T_{c\bar{c}1}(3900)$ MASS

NODE=M210M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3887.1±2.6 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.			
3893.1±2.2± 3.0		¹ ABLIKIM	20N BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
3902.6 ^{+5.2+ 3.3} _{-5.0- 1.4}		^{2,3} ABAZOV	19 D0	±	1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3881.2±4.2±52.7	6k	⁴ ABLIKIM	17J BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
3885.7 ^{+4.3} _{-5.7} ± 8.4		^{2,4} ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
3881.7±1.6± 1.6	1.2k	^{2,4} ABLIKIM	15AC BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3883.9±1.5± 4.2	1.2k	^{2,4} ABLIKIM	14A BES3	±	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
3894.5±6.6± 4.5	159	² LIU	13B BELL	±	$e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
3886 ± 4 ± 2	81	^{2,5} XIAO	13A	±	4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
3904 ± 9 ± 5	25	^{2,5} XIAO	13A	0	4.17 $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$

NODE=M210M

OCCUR=2

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

3895.0±5.2 ^{+ 4.0} _{- 2.7}	502	^{2,6} ABAZOV	18B D0	±	1.96 TeV $p\bar{p} \rightarrow J/\psi\pi^+\pi^- X$
3894.8±2.3± 3.2	356	^{2,7} ABLIKIM	15U BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
3899.0±3.6± 4.9	307	^{2,8} ABLIKIM	13T BES3	±	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ Pole mass obtained from a fit to a relativistic Breit-Wigner.² Neglecting interference between the $T_{c\bar{c}1}(3900)$ and other processes.³ Measured in weak decays of b -flavored hadrons (nonprompt).⁴ Pole mass obtained from a fit to a Flatte-like formula.⁵ For $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.⁶ The signal of the $T_{c\bar{c}1}(3900)$ is correlated with a parent $J/\psi\pi^+\pi^-$ system in the invariant mass range 4.2–4.7 GeV. Superseded by ABAZOV 19.⁷ Superseded by ABLIKIM 20N.⁸ Superseded by ABLIKIM 17J.

NODE=M210M;LINKAGE=L

NODE=M210M;LINKAGE=A

NODE=M210M;LINKAGE=G


NODE=M210M;LINKAGE=E

NODE=M210M;LINKAGE=B

NODE=M210M;LINKAGE=F

NODE=M210M;LINKAGE=J

NODE=M210M;LINKAGE=K



$T_{c\bar{c}1}(3900)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
28.4 ± 2.6 OUR AVERAGE					
44.4 ± 5.2 ± 14.0		¹ ABLIKIM	20N	BES3	0 $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
32 ⁺²⁸ ₋₂₁ ⁺²⁶ ₋₇		^{2,3} ABAZOV	19	D0	± 1.96 TeV $p\bar{p} \rightarrow \pi^+\pi^- J/\psi X$ (non-prompt)
51.8 ± 4.6 ± 36.0	6 k	⁴ ABLIKIM	17J	BES3	± $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
35 ⁺¹¹ ₋₁₂ ± 15		^{2,4} ABLIKIM	15AB	BES3	0 $e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$
26.6 ± 2.0 ± 2.1	1248	^{2,4} ABLIKIM	15AC	BES3	± $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
24.8 ± 3.3 ± 11.0	1212	^{2,4} ABLIKIM	14A	BES3	± $e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$
63 ± 24 ± 26	159	² LIU	13B	BELL	± $e^+e^- \rightarrow \gamma\pi^+\pi^- J/\psi$
37 ± 4 ± 8	81	^{2,5} XIAO	13A		± 4.17 $e^+e^- \rightarrow \pi^+\pi^- J/\psi$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
29.6 ± 8.2 ± 8.2	356	^{2,6} ABLIKIM	15U	BES3	0 $e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
46 ± 10 ± 20	307	^{2,7} ABLIKIM	13T	BES3	± $e^+e^- \rightarrow \pi^+\pi^- J/\psi$

- ¹ Pole width obtained from a fit to a relativistic Breit-Wigner.
- ² Neglecting interference between the $T_{c\bar{c}1}(3900)$ and other processes.
- ³ Measured in weak decays of b -flavored hadrons (nonprompt).
- ⁴ Pole width obtained from a fit to a Flatte-like formula.
- ⁵ For $M^2(\pi^+\pi^-) < 0.65 \text{ GeV}^2$. Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.
- ⁶ Superseded by ABLIKIM 20N.
- ⁷ Superseded by ABLIKIM 17J.

NODE=M210W

NODE=M210W

NODE=M210W;LINKAGE=G
 NODE=M210W;LINKAGE=A
 NODE=M210W;LINKAGE=F
 NODE=M210W;LINKAGE=E
 NODE=M210W;LINKAGE=B

NODE=M210W;LINKAGE=H
 NODE=M210W;LINKAGE=I

$T_{c\bar{c}1}(3900)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\pi$	seen
Γ_2 $h_c\pi^\pm$	not seen
Γ_3 $\eta_c\pi^+\pi^-$	not seen
Γ_4 $\eta_c(1S)\rho(770)^\pm$	seen
Γ_5 $(D\bar{D}^*)^\pm$	seen
Γ_6 $D^0D^{*-} + \text{c.c.}$	seen
Γ_7 $D^-D^{*0} + \text{c.c.}$	seen
Γ_8 $\omega\pi^\pm$	not seen
Γ_9 $J/\psi\eta$	not seen
Γ_{10} $D^+D^{*-} + \text{c.c.}$	seen
Γ_{11} $D^0\bar{D}^{*0} + \text{c.c.}$	seen

NODE=M210215;NODE=M210

DESIG=1
 DESIG=2
 DESIG=10
 DESIG=11;OUR EVAL;→ UNCHECKED ←
 DESIG=3;OUR EVAL;→ UNCHECKED ←
 DESIG=8
 DESIG=9
 DESIG=4
 DESIG=5
 DESIG=6
 DESIG=7

$T_{c\bar{c}1}(3900)$ BRANCHING RATIOS

NODE=M210225

$\Gamma(J/\psi\pi)/\Gamma_{\text{total}}$						Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	20N	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
seen		¹ ABAZOV	19	D0	\pm	$1.96 \text{ TeV } p\bar{p} \rightarrow \pi^+\pi^- J/\psi X \text{ (prompt)}$
seen		ABLIKIM	17J	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
seen	356	ABLIKIM	15U	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 J/\psi$
not seen		² ADOLPH	15D	COMP	\pm	$\gamma N \rightarrow J/\psi\pi^\pm N$
seen	307	ABLIKIM	13T	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$
seen	25	³ XIAO	13A		0	$4.17 e^+e^- \rightarrow \pi^0\pi^0 J/\psi$

NODE=M210R01
NODE=M210R01¹ But not seen in the "prompt" sample (no b-hadron enhancement).² ADOLPH 15D measure $B(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi\pi^\pm) \sigma(\gamma N \rightarrow T_{c\bar{c}1}(3900)^\pm N) / \sigma(\gamma N \rightarrow J/\psi N) < 3.7 \times 10^{-3}$ at 90% CL.³ Obtained by analyzing CLEO-c data but not authored by the CLEO Collaboration.NODE=M210R01;LINKAGE=C
NODE=M210R01;LINKAGE=A

NODE=M210R01;LINKAGE=XI

$\Gamma(h_c\pi^\pm)/\Gamma_{\text{total}}$						Γ_2/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	13X	BES3	\pm	$e^+e^- \rightarrow h_c\pi^+\pi^-$

NODE=M210R02
NODE=M210R02

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_3/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		¹ VINOKUROVA 15	BELL	0	$B^+ \rightarrow K^+\eta_c\pi^+\pi^-$	

NODE=M210R11
NODE=M210R11¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+ Z_c(3900)^0) \times B(X \rightarrow \eta_c\pi^+\pi^-) < 4.7 \times 10^{-5}$ at 90% CL.

NODE=M210R11;LINKAGE=VI

$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(J/\psi\pi)$						Γ_4/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
2.3 ± 0.8	332	¹ ABLIKIM	19BC	BES3	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$	

NODE=M210R12
NODE=M210R12¹ Using $e^+e^- \rightarrow \pi^\mp(T_{c\bar{c}1}(3900)^\pm \rightarrow J/\psi\pi^\pm)$ cross section at 4.23 and 4.26 GeV from ABLIKIM 17J.

NODE=M210R12;LINKAGE=A

$\Gamma((D\bar{D}^*)^\pm)/\Gamma(J/\psi\pi)$						Γ_5/Γ_1
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
$6.2 \pm 1.1 \pm 2.7$		¹ ABLIKIM	14A	BES3	\pm	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$

NODE=M210R03
NODE=M210R03¹ Assuming the same origin of the $(D\bar{D}^*)^\pm$ and $\pi^\pm J/\psi$ decay modes.

NODE=M210R03;LINKAGE=A

$\Gamma(D^0 D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_6/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	15AC	BES3	\pm	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$
seen		ABLIKIM	14A	BES3	\pm	$e^+e^- \rightarrow \pi^+ D^0 D^{*-} + \text{c.c.}$

NODE=M210R09
NODE=M210R09

$\Gamma(D^- D^{*0} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_7/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	15AC	BES3	\pm	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$
seen		ABLIKIM	14A	BES3	\pm	$e^+e^- \rightarrow \pi^+ D^- D^{*0} + \text{c.c.}$

NODE=M210R10
NODE=M210R10

$\Gamma(\omega\pi^\pm)/\Gamma_{\text{total}}$						Γ_8/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	15R	BES3	\pm	$e^+e^- \rightarrow \omega\pi^+\pi^-$

NODE=M210R00
NODE=M210R00

$\Gamma(J/\psi\eta)/\Gamma_{\text{total}}$						Γ_9/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
not seen		ABLIKIM	15Q	BES3	0	$4.0\text{--}4.6 e^+e^- \rightarrow J/\psi\eta\pi^0$

NODE=M210R04
NODE=M210R04

$\Gamma(J/\psi\eta)/\Gamma(J/\psi\pi)$						Γ_9/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	
<0.15	90	ABLIKIM	15Q	BES3	0	$4.226 e^+e^- \rightarrow J/\psi\eta\pi^0$

NODE=M210R05
NODE=M210R05

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.65	90	ABLIKIM	15Q	BES3	0	$4.257 e^+e^- \rightarrow J/\psi\eta\pi^0$
-------	----	---------	-----	------	---	--

OCCUR=2

$\Gamma(D^+ D^{*-} + \text{c.c.})/\Gamma_{\text{total}}$						Γ_{10}/Γ
VALUE		DOCUMENT ID	TECN	CHG	COMMENT	
seen		ABLIKIM	15AB	BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

NODE=M210R06
NODE=M210R06

$\Gamma(D^0\bar{D}^{*0} + c.c.)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

NODE=M210R07
NODE=M210R07 $\Gamma(D^+D^{*-} + c.c.)/\Gamma(D^0\bar{D}^{*0} + c.c.)$ Γ_{10}/Γ_{11}

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.96±0.18±0.12	ABLIKIM	15AB BES3	0	$e^+e^- \rightarrow \pi^0(D\bar{D}^*)^0$

NODE=M210R08
NODE=M210R08 **$T_{c\bar{c}s1}(3900)$ REFERENCES**

NODE=M210

ABLIKIM	20N	PR D102 012009	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=60338
ABAZOV	19	PR D100 012005	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=59903
ABLIKIM	19BC	PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=60036
ABAZOV	18B	PR D98 052010	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=59442
ABLIKIM	17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=57950
ABLIKIM	15AB	PRL 115 222002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56954
ABLIKIM	15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=56967
ABLIKIM	15Q	PR D92 012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56782
ABLIKIM	15R	PR D92 032009	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56783
ABLIKIM	15U	PRL 115 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=56786
ADOLPH	15D	PL B742 330	C. Adolph <i>et al.</i>	(COMPASS Collab.)	REFID=56791
VINOKUROVA	15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)	REFID=56706
Also		JHEP 1702 088 (errata)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)	REFID=57795
ABLIKIM	14A	PRL 112 022001	M. Ablikim <i>et al.</i>	(BESIII Collab.) JP	REFID=55648
ABLIKIM	13T	PRL 110 252001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55409
ABLIKIM	13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=55635
LIU	13B	PRL 110 252002	Z.Q. Liu <i>et al.</i>	(BELLE Collab.)	REFID=55410
XIAO	13A	PL B727 366	T. Xiao <i>et al.</i>	(NWES)	REFID=55593

NODE=M259

 $T_{c\bar{c}s1}(4000)$

$$I(J^P) = \frac{1}{2}(1^+)$$

OMITTED FROM SUMMARY TABLE

was $Z_{cs}(4000)$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M259

Seen by AAIJ 21E in $B^+ \rightarrow T_{c\bar{c}s1}(4000)^+ \phi$ with $T_{c\bar{c}s1}(4000)^+ \rightarrow J/\psi K^+$ using an amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ with a significance (accounting for systematic uncertainties) of 15σ . The $J^P = 1^+$ assignment is favored with high significance. ABLIKIM 21G also reports a $J^P = 1^+ Z_{cs}$ state in this mass region using $e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$ with a significance of 5.3σ . The incompatible values for the widths reported by AAIJ 21E and ABLIKIM 21G could either indicate the existence of two separate states or possibly be explained in a coupled channel model (see ORTEGA 21).

 $T_{c\bar{c}s1}(4000)$ MASS

NODE=M259M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3980-4010 OUR EVALUATION				
3988 ± 5 OUR AVERAGE Error includes scale factor of 2.7.				
3991 $^{+12}_{-10}$ $^{+9}_{-17}$		1 AAIJ	23AQ LHCB	$B^0 \rightarrow J/\psi \phi K_S^0$
3992.2 ± 1.7 ± 1.6		2 ABLIKIM	22AE BES3	$e^+e^- \rightarrow K_S^0(D_s^- D^{*+} + D_s^{*-} D^+)$
4003 ± 6 $^{+4}_{-14}$	24k	3 AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$
3982.5 $^{+1.8}_{-2.6}$ ± 2.1		4 ABLIKIM	21G BES3	$e^+e^- \rightarrow K^+(D_s^- D^{*0} + D_s^{*-} D^0)$

NODE=M259M
→ UNCHECKED ←

¹ From an amplitude analysis of the decay $B^0 \rightarrow J/\psi \phi K_S^0$ with a significance of 4.0σ . The mass difference with respect to the charged partner in AAIJ 21E is -12^{+11+6}_{-10-4} MeV.

NODE=M259M;LINKAGE=E

² Pole mass for a mass-, width-dependent Breit-Wigner fit to the mass spectrum recoiling against K_S^0 at center of mass energies between 4.628 and 4.699 GeV, with a significance of 4.6σ .

NODE=M259M;LINKAGE=D

³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 15σ .

NODE=M259M;LINKAGE=A

⁴ Pole mass for a mass-dependent Breit-Wigner fit to the mass spectrum recoiling against K^+ at center of mass energies between 4.628 and 4.698 GeV, with a significance of 5.3 σ .

NODE=M259M;LINKAGE=B

 $T_{c\bar{c}s1}(4000)$ WIDTH

NODE=M259W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

NODE=M259W

5-150 OUR EVALUATION

→ UNCHECKED ←
NEW

14 ± 4 OUR AVERAGE Error includes scale factor of 1.1. [13 ± 4 MeV OUR 2023 AVERAGE]

105	$+29$ -25	$+17$ -23	1 AAIJ	23AQ LHCB	$B^0 \rightarrow J/\psi \phi K_S^0$
	7.7^+ $-$	4.1 3.8	± 4.3	2 ABLIKIM	$e^+ e^- \rightarrow K_S^0 (D_s^- D^{*+} + D_s^{*-} D^+)$
131	±15	±26	24k	3 AAIJ	21E LHCB $B^+ \rightarrow J/\psi \phi K^+$
	12.8^+ $-$	5.3 4.4	± 3.0	4 ABLIKIM	$e^+ e^- \rightarrow K^+ (D_s^- D^{*0} + D_s^{*-} D^0)$

¹ From an amplitude analysis of the decay $B^0 \rightarrow J/\psi \phi K_S^0$ with a significance of 4.0 σ .

NODE=M259W;LINKAGE=D

² Pole width for a mass-, width-dependent Breit-Wigner fit to the mass spectrum recoiling against K_S^0 at center of mass energies between 4.628 and 4.699 GeV, with a significance of 4.6 σ .

NODE=M259W;LINKAGE=C

³ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 15 σ .

NODE=M259W;LINKAGE=A

⁴ Pole width for a mass-dependent Breit-Wigner fit to the mass spectrum recoiling against K^+ at center of mass energies between 4.628 and 4.698 GeV, with a significance of 5.3 σ .

NODE=M259W;LINKAGE=B

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi K^+$	seen
Γ_2 $J/\psi K_S^0$	seen
Γ_3 $D_s^+ \bar{D}^{*0}$ or $D_s^{*+} \bar{D}^0$	seen
Γ_4 $D_s^+ D^{*-}$ or $D_s^{*+} D^-$	seen

NODE=M259215;DESIG=1

DESIG=4

DESIG=2

DESIG=3

 $T_{c\bar{c}s1}(4000)$ DECAY MODES

NODE=M259225

$\Gamma(J/\psi K^+)/\Gamma_{\text{total}}$ Γ_1/Γ

NODE=M259R01
NODE=M259R01

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

seen 24k ¹ AAIJ 21E LHCB $B^+ \rightarrow J/\psi \phi K^+$

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 15 σ .

NODE=M259R01;LINKAGE=A

$\Gamma(J/\psi K_S^0)/\Gamma_{\text{total}}$ Γ_2/Γ

NODE=M259R02
NODE=M259R02

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen ¹ AAIJ 23AQ LHCB $B^0 \rightarrow J/\psi \phi K_S^0$

¹ From an amplitude analysis of the decay $B^0 \rightarrow J/\psi \phi K_S^0$ with a significance of 4.0 σ .

NODE=M259R02;LINKAGE=A

$\Gamma(D_s^+ \bar{D}^{*0}$ or $D_s^{*+} \bar{D}^0)/\Gamma_{\text{total}}$ Γ_3/Γ

NODE=M259R00
NODE=M259R00

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen ¹ ABLIKIM 21G BES3 $e^+ e^- \rightarrow K^+ (D_s^- D^{*0} + D_s^{*-} D^0)$

¹ Seen in the spectrum recoiling against K^+ in $e^+ e^- \rightarrow K^+ (D_s^- D^{*0} + D_s^{*-} D^0)$ collisions at center of mass energies between 4.628 and 4.698 GeV, with a significance of 5.3 σ .

NODE=M259R00;LINKAGE=A

$\Gamma(J/\psi K^+)/\Gamma(D_s^+ \bar{D}^{*0}$ or $D_s^{*+} \bar{D}^0)$ Γ_1/Γ_3

NODE=M259R04
NODE=M259R04

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

<0.03 90 ABLIKIM 23BI BES3 $e^+ e^- \rightarrow K^+ K^- J/\psi$

$\Gamma(D_s^+ D^{*-}$ or $D_s^{*+} D^-)/\Gamma_{\text{total}}$ Γ_4/Γ

NODE=M259R03
NODE=M259R03

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen ¹ ABLIKIM 22AE BES3 $e^+ e^- \rightarrow K_S^0 (D_s^- D^{*+} + D_s^{*-} D^+)$

¹ Seen in the mass spectrum recoiling against K_S^0 at center of mass energies between 4.628 and 4.699 GeV, with a significance of 4.6 σ .

NODE=M259R03;LINKAGE=A

$T_{c\bar{c}s1}(4000)$ REFERENCES

AAIJ	23AQ	PRL 131 131901	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABLIKIM	23BI	PRL 131 211902	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	22AE	PRL 129 112003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	21E	PRL 127 082001	R. Aaij <i>et al.</i>	(LHCb Collab.) JP
ABLIKIM	21G	PRL 126 102001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ORTEGA	21	PL B818 136382	P.G. Ortega, D.R. Entem, F. Fernandez	

NODE=M259

REFID=62429
REFID=62434
REFID=61877
REFID=61150
REFID=61065
REFID=61108

NODE=M213

 $T_{c\bar{c}}(4020)$

$$I^G(J^{PC}) = 1^+(?^-)$$

was $X(4020)$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

NODE=M213

Charged $T_{c\bar{c}}(4020)$ seen by ABLIKIM 13X from $e^+e^- \rightarrow \pi^+\pi^-h_c(1P)$ at c.m. energy from 3.90 to 4.42 GeV as a peak in the invariant mass distribution of the $\pi^\pm h_c(1P)$ system, and by ABLIKIM 14B from $e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$ events in $(D^*\bar{D}^*)^\pm$ mass. A neutral $T_{c\bar{c}}(4020)$ seen by ABLIKIM 14P at three c.m. energies in the same range in $e^+e^- \rightarrow \pi^0\pi^0 h_c(1P)$ as a peak in the larger of the two masses recoiling against a π^0 . ABLIKIM 15AA observes a 5.9σ signal in $(D^*\bar{D}^*)^0$ in $e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$ events using collisions at two c.m. energies. Production rates and mass values support grouping neutral and charged $T_{c\bar{c}}(4020)$ together as manifestations of a single $I = 1$ particle.

 $T_{c\bar{c}}(4020)$ MASS

NODE=M213M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
4024.1 ± 1.9 OUR AVERAGE					
$4025.5^{+2.0}_{-4.7} \pm 3.1$	116	¹ ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
$4026.3 \pm 2.6 \pm 3.7$	401	¹ ABLIKIM 14B	BES3	±	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
$4023.9 \pm 2.2 \pm 3.8$	61	^{1,2} ABLIKIM 14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 h_c$
$4022.9 \pm 0.8 \pm 2.7$	253	¹ ABLIKIM 13X	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-h_c$

NODE=M213M

¹ Neglecting interference between the $T_{c\bar{c}}(4020)$ and non-resonant continuum.² Assuming $J^P = 1^+$ and width of 7.9 ± 2.6 MeV.

NODE=M213M;LINKAGE=AB

NODE=M213M;LINKAGE=B

 $T_{c\bar{c}}(4020)$ WIDTH


NODE=M213W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
13 ± 5 OUR AVERAGE					
Error includes scale factor of 1.7. See the ideogram below.					
$23.0 \pm 6.0 \pm 1.0$	116	¹ ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$
$24.8 \pm 5.6 \pm 7.7$	401	¹ ABLIKIM 14B	BES3	±	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$
$7.9 \pm 2.7 \pm 2.6$	253	¹ ABLIKIM 13X	BES3	±	$e^+e^- \rightarrow \pi^+\pi^-h_c$

NODE=M213W

¹ Neglecting interference between the $T_{c\bar{c}}(4020)$ and non-resonant continuum.

NODE=M213W;LINKAGE=AB



$T_{c\bar{c}}(4020)$ DECAY MODES

NODE=M213215;NODE=M213

Mode	Fraction (Γ_i/Γ)	
Γ_1 $h_c(1P)\pi$	seen	DESIG=1
Γ_2 $D^*\bar{D}^*$	seen	DESIG=2
Γ_3 $D\bar{D}^* + \text{c.c.}$	not seen	DESIG=4
Γ_4 $\eta_c\pi^+\pi^-$	not seen	DESIG=3
Γ_5 $\eta_c(1S)\rho(770)^\pm$		DESIG=6
Γ_6 $J/\psi(1S)\pi^\pm$	not seen	DESIG=5

$T_{c\bar{c}}(4020)$ BRANCHING RATIOS

NODE=M213225

$\Gamma(h_c(1P)\pi)/\Gamma_{\text{total}}$						Γ_1/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
seen	61	ABLIKIM 14P	BES3	0	$e^+e^- \rightarrow \pi^0\pi^0 h_c$		NODE=M213R01
seen	253	ABLIKIM 13X	BES3	\pm	$e^+e^- \rightarrow \pi^+\pi^- h_c$		NODE=M213R01

$\Gamma(D^*\bar{D}^*)/\Gamma_{\text{total}}$						Γ_2/Γ	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
seen	116	¹ ABLIKIM 15AA	BES3	0	$e^+e^- \rightarrow (D^*\bar{D}^*)^0\pi^0$		NODE=M213R02
seen	401	¹ ABLIKIM 14B	BES3	\pm	$e^+e^- \rightarrow (D^*\bar{D}^*)^\pm\pi^\mp$		NODE=M213R02

¹ Neglecting interference between the $T_{c\bar{c}}(4020)$ and non-resonant continuum.

NODE=M213R02;LINKAGE=A

$\Gamma(D\bar{D}^* + \text{c.c.})/\Gamma_{\text{total}}$						Γ_3/Γ	
VALUE		DOCUMENT ID	TECN	CHG	COMMENT		
not seen		ABLIKIM 15AC	BES3	\pm	$e^+e^- \rightarrow \pi^\pm(D\bar{D}^*)^\mp$		NODE=M213R03
							NODE=M213R03

$\Gamma(\eta_c\pi^+\pi^-)/\Gamma_{\text{total}}$						Γ_4/Γ	
VALUE		DOCUMENT ID	TECN		COMMENT		
not seen		¹ VINOKUROVA 15	BELL		$B^+ \rightarrow K^+\eta_c\pi^+\pi^-$		NODE=M213R00
							NODE=M213R00

¹ VINOKUROVA 15 reports $B(B^+ \rightarrow K^+T_{c\bar{c}}(4020)^0) \times B(T_{c\bar{c}} \rightarrow \eta_c\pi^+\pi^-) < 1.6 \times 10^{-5}$ at 90% CL.

NODE=M213R00;LINKAGE=VI

$\Gamma(\eta_c(1S)\rho(770)^\pm)/\Gamma(h_c(1P)\pi)$						Γ_5/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN		COMMENT		
<1.2	90	¹ ABLIKIM 19BC	BES3		$e^+e^- \rightarrow \pi^+\pi^-\pi^0\eta_c(1S)$		NODE=M213R05
							NODE=M213R05

¹ Using $e^+e^- \rightarrow \pi^\mp(Z_c(4020)^\pm \rightarrow h_c(1P)\pi^\pm)$ cross section at 4.23, 4.26 and 4.36 GeV from ABLIKIM 13X.

NODE=M213R05;LINKAGE=A

$\Gamma(J/\psi(1S)\pi^\pm)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM 17J	BES3	$e^+e^- \rightarrow \pi^+\pi^- J/\psi$

¹ From Partial Wave Analysis assuming $J^P = 1^+$.NODE=M213R04
NODE=M213R04

NODE=M213R04;LINKAGE=A

 $T_{c\bar{c}}(4020)$ REFERENCES

ABLIKIM 19BC	PR D100 111102	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 17J	PRL 119 072001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 15AA	PRL 115 182002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 15AC	PR D92 092006	M. Ablikim <i>et al.</i>	(BESIII Collab.)
VINOKUROVA 15	JHEP 1506 132	A. Vinokurova <i>et al.</i>	(BELLE Collab.)
Also	JHEP 1702 088 (errata.)	A. Vinokurava <i>et al.</i>	(BELLE Collab.)
ABLIKIM 14B	PRL 112 132001	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 14P	PRL 113 212002	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM 13X	PRL 111 242001	M. Ablikim <i>et al.</i>	(BESIII Collab.)

NODE=M213

REFID=60036
REFID=57950
REFID=56951
REFID=56967
REFID=56706
REFID=57795
REFID=55654
REFID=56118
REFID=55635

NODE=M191

 $T_{c\bar{c}}(4050)^+$ $I^G(J^{PC}) = 1^-(?^?+)$
 I, G, C need confirmation.OMITTED FROM SUMMARY TABLE
was $X(4050)$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

NODE=M191

Observed by MIZUK 08 in the $\pi^+\chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure. $T_{c\bar{c}}(4050)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4051 \pm 14^{+20}_{-41}$	¹ MIZUK 08	BELL	$\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M191M

NODE=M191M

NODE=M191M;LINKAGE=MI

 $T_{c\bar{c}}(4050)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
82^{+21+47}_{-17-22}	¹ MIZUK 08	BELL	$\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M191W

NODE=M191W

NODE=M191W;LINKAGE=MI

 $T_{c\bar{c}}(4050)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+\chi_{c1}(1P)$	seen
$\Gamma_2 \pi^\pm\psi(3770)$	not seen
$\Gamma_3 \pi^\pm\chi_{c0}(1P)$	not seen
$\Gamma_4 \pi^\pm\chi_{c2}(1P)$	not seen

NODE=M191215;NODE=M191

DESIG=1

DESIG=2

DESIG=3

DESIG=4

 $T_{c\bar{c}}(4050)^+$ BRANCHING RATIOS $\Gamma(\pi^+\chi_{c1}(1P))/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen		¹ MIZUK 08	BELL	$\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

••• We do not use the following data for averages, fits, limits, etc. •••

not seen 16 ² ABLIKIM 21W BES3 $e^+e^- \rightarrow \chi_{cJ}\pi^+\pi^+$ not seen ³ LEES 12B BABR $B \rightarrow K\pi\chi_{c1}(1P)$ ¹ With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- T_{c\bar{c}}(4050)^+) \times B(T_{c\bar{c}}(4050)^+ \rightarrow \pi^+\chi_{c1}(1P)) = (3.0^{+1.5+3.7}_{-0.8-1.6}) \times 10^{-5}$.² ABLIKIM 21W measurement is limited by statistics.³ With a product branching fraction limit of $B(\bar{B}^0 \rightarrow T_{c\bar{c}}(4050)^+ K^-) \times B(T_{c\bar{c}}(4050)^+ \rightarrow \chi_{c1}\pi^+) < 1.8 \times 10^{-5}$ at 90% CL.

NODE=M191225

NODE=M191R01
NODE=M191R01

NODE=M191R01;LINKAGE=MI

NODE=M191R01;LINKAGE=A

NODE=M191R01;LINKAGE=LE

$\Gamma(\pi^\pm \chi_{c0}(1P))/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE	EVT	DOCUMENT ID	TECN	COMMENT
not seen	18	¹ ABLIKIM	21W BES3	$e^+e^- \rightarrow \chi_{cJ}\pi^+\pi^+$

¹ ABLIKIM 21W measurement is limited by statistics.

NODE=M191R02
NODE=M191R02

NODE=M191R02;LINKAGE=A

 $\Gamma(\pi^\pm \chi_{c2}(1P))/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	EVT	DOCUMENT ID	TECN	COMMENT
not seen	14	¹ ABLIKIM	21W BES3	$e^+e^- \rightarrow \chi_{cJ}\pi^+\pi^+$

¹ ABLIKIM 21W measurement is limited by statistics.

NODE=M191R03
NODE=M191R03

NODE=M191R03;LINKAGE=A

 $\Gamma(\pi^\pm \psi(3770))/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^-D\bar{D}$

¹ From a measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.

NODE=M191R00
NODE=M191R00

NODE=M191R00;LINKAGE=A

 $T_{c\bar{c}}(4050)^+$ REFERENCES

ABLIKIM	21W	PR D103 052010	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
LEES	12B	PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08	PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

NODE=M191

REFID=61221
REFID=59910
REFID=54042
REFID=52535

NODE=M223

$T_{c\bar{c}}(4055)^+$

$I^G(J^{PC}) = 1^+(?^-)$
 I, G, C need confirmation.

OMITTED FROM SUMMARY TABLE
was $X(4055)^\pm$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

NODE=M223

Needs confirmation. Seen by WANG 15A in the $\psi(2S)\pi^+$ invariant mass distribution in $\psi(4360) \rightarrow \psi(2S)\pi^+\pi^-$ decay.

 $T_{c\bar{c}}(4055)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4054 ± 3 ± 1	¹ WANG	15A BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
4039.3 ± 6.0	² ABLIKIM	18K BES3	$e^+e^- \rightarrow \pi^0\pi^0\psi(2S)$
4032.1 ± 2.4	³ ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Statistical significance of 3.5σ .

² Statistical error only, with significance of 5.9σ (from a fit with a 19% CL). Identified as the same structure observed in ABLIKIM 17V in $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ decays.

³ Statistical error only, with significance of 9.2σ . From an unbinned maximum likelihood fit of the $\pi^+\pi^-\psi(2S)$ Dalitz plot from data collected at $\sqrt{s} = 4.416$ GeV for a $J^C = 1^+$ state. The fit does not match the detailed structure of the data, having a C.L. of only 8%.

NODE=M223M

NODE=M223M

OCCUR=2

NODE=M223M;LINKAGE=A

NODE=M223M;LINKAGE=C

NODE=M223M;LINKAGE=B

 $T_{c\bar{c}}(4055)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
45 ± 11 ± 6	¹ WANG	15A BELL	$10.58 e^+e^- \rightarrow \gamma\pi^+\pi^-\psi(2S)$
31.9 ± 14.8	² ABLIKIM	18K BES3	$e^+e^- \rightarrow \pi^0\pi^0\psi(2S)$
26.1 ± 5.3	³ ABLIKIM	17V BES3	$e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$

••• We do not use the following data for averages, fits, limits, etc. •••

¹ Statistical significance of 3.5σ .

² Statistical error only, with significance of 5.9σ (from a fit with a 19% CL). Identified as the same structure observed in ABLIKIM 17V in $e^+e^- \rightarrow \pi^+\pi^-\psi(2S)$ decays.

³ Statistical error only, with significance of 9.2σ . From an unbinned maximum likelihood fit of the $\pi^+\pi^-\psi(2S)$ Dalitz plot from data collected at $\sqrt{s} = 4.416$ GeV for a $J^C = 1^+$ state. The fit does not match the detailed structure of the data, having a C.L. of only 8%.

NODE=M223W

NODE=M223W

NODE=M223W;LINKAGE=A

NODE=M223W;LINKAGE=C

NODE=M223W;LINKAGE=B

$T_{c\bar{c}}(4055)^+$ DECAY MODES

NODE=M223215;NODE=M223

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi^+ \psi(2S)$	seen
Γ_2 $\pi^\pm \psi(3770)$	not seen

DESIG=1

DESIG=2

 $T_{c\bar{c}}(4055)^+$ BRANCHING RATIOS

NODE=M223225

$\Gamma(\pi^+ \psi(2S))/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ WANG	15A	BELL 10.58 $e^+ e^- \rightarrow \gamma \pi^+ \pi^- \psi(2S)$

¹ Statistical significance of 3.5 σ .

NODE=M223R01

NODE=M223R01

NODE=M223R01;LINKAGE=A

$\Gamma(\pi^\pm \psi(3770))/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
not seen	¹ ABLIKIM	19AR	BES3 $e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D}$

¹ From a measurement of $\sigma(e^+ e^- \rightarrow \pi^+ \pi^- D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.

NODE=M223R00

NODE=M223R00

NODE=M223R00;LINKAGE=A

 $T_{c\bar{c}}(4055)^+$ REFERENCES

NODE=M223

ABLIKIM	19AR	PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59910
ABLIKIM	18K	PR D97 052001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58896
ABLIKIM	17V	PR D96 032004	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=58029
Also		PR D99 019903 (errat.)	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=59611
WANG	15A	PR D91 112007	X.L. Wang <i>et al.</i>	(BELLE Collab.)	REFID=56839

NODE=M240

 $T_{c\bar{c}}(4100)^+$

$$J^G(J^PC) = 1^-(?^?+)$$

OMITTED FROM SUMMARY TABLE

was $X(4100)^\pm$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

NODE=M240

Reported by AAIJ 18AN in the $\eta_c(1S)\pi^-$ invariant mass distribution in $B^0 \rightarrow \eta_c(1S)K^+\pi^-$ decays with a significance of 3.4 σ . $J^P = 0^+$ or 1^- assignment consistent with data.

 $T_{c\bar{c}}(4100)^+$ MASS

NODE=M240M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4096 \pm 20^{+18}_{-22}$	AAIJ	18AN	LHCB $B^0 \rightarrow \eta_c(1S)K^+\pi^-$

NODE=M240M

 $T_{c\bar{c}}(4100)^+$ WIDTH

NODE=M240W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150^{+80}_{-70} OUR AVERAGE	[152 ⁺⁸⁰ ₋₇₀ MeV OUR 2023 AVERAGE]		
$152 \pm 58^{+60}_{-35}$	AAIJ	18AN	LHCB $B^0 \rightarrow \eta_c(1S)K^+\pi^-$

NODE=M240W

NEW

 $T_{c\bar{c}}(4100)^+$ DECAY MODES

NODE=M240215;NODE=M240

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta_c(1S)\pi^-$	seen
Γ_2 $\pi^\pm \psi(3770)$	not seen

DESIG=1

DESIG=2

$T_{c\bar{c}}(4100)^+$ BRANCHING RATIOS

$\Gamma(\eta_c(1S)\pi^-)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		¹ AAIJ	18AN LHCB	$B^0 \rightarrow \eta_c(1S)K^+\pi^-$	
¹ AAIJ 18AN quotes a fit fraction for $B^0 \rightarrow T_{c\bar{c}}(4100)^- K^+ \rightarrow \eta_c(1S)\pi^- K^+$ of $(3.3 \pm 1.1^{+1.2}_{-1.1})\%$ from an amplitude analysis.					

NODE=M240225

NODE=M240R01
NODE=M240R01

NODE=M240R01;LINKAGE=A

$\Gamma(\pi^\pm\psi(3770))/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
not seen		¹ ABLIKIM	19AR BES3	$e^+e^- \rightarrow \pi^+\pi^-D\bar{D}$	
¹ From a measurement of $\sigma(e^+e^- \rightarrow \pi^+\pi^-D\bar{D})$ between $\sqrt{s} = 4.08$ and 4.6 GeV.					

NODE=M240R00
NODE=M240R00

NODE=M240R00;LINKAGE=A

 $T_{c\bar{c}}(4100)^+$ REFERENCES

ABLIKIM	19AR PR D100 032005	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18AN EPJ C78 1019	R. Aaij <i>et al.</i>	(LHCb Collab.)

NODE=M240

REFID=59910
REFID=59335

NODE=M231

 $T_{c\bar{c}1}(4200)^+$
 $I^G(J^{PC}) = 1^+(1^{+-})$
I, G, C need confirmation.

OMITTED FROM SUMMARY TABLE
 was $Z_c(4200)$, $X(4200)^\pm$

This state shows properties different from a conventional $q\bar{q}$ state.
 A candidate for an exotic structure. See the review on non- $q\bar{q}$ states.

NODE=M231

Reported by CHILIKIN 14 in $J/\psi\pi^+$ at a significance of 6.2σ . Assignments of 0^- , 1^- , 2^- , and 2^+ excluded at 6.1σ , 7.4σ , 4.4σ , and 7.0σ level, respectively. Needs confirmation.

 $T_{c\bar{c}1}(4200)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4196^{+31+17}_{-29-13}	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

NODE=M231M

NODE=M231M

 $T_{c\bar{c}1}(4200)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$370 \pm 70^{+70}_{-132}$	CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$

NODE=M231W

NODE=M231W

 $T_{c\bar{c}1}(4200)^+$ DECAY MODES

NODE=M231215;NODE=M231

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi\pi^+$	seen

DESIG=1

 $T_{c\bar{c}1}(4200)^+$ BRANCHING RATIOS

NODE=M231220

$\Gamma(J/\psi\pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		CHILIKIN	14	BELL $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
possibly seen		¹ AAIJ	19R LHCB	$B^0 \rightarrow K^+\pi^- J/\psi + \text{c.c.}$	
¹ From a model-independent analysis.					

NODE=M231R01
NODE=M231R01

NODE=M231R01;LINKAGE=C

 $T_{c\bar{c}1}(4200)^+$ REFERENCES

NODE=M231

AAIJ	19R PRL 122 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)
CHILIKIN	14 PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)

REFID=59776
REFID=56344

NODE=M260

 $T_{c\bar{c}s1}(4220)^+$

$$I(J^P) = \frac{1}{2}(1^+)$$

OMITTED FROM SUMMARY TABLE

was $Z_{cs}(4220)^+$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M260

Seen by AAIJ 21E in $B^+ \rightarrow T_{c\bar{c}s1}(4220)^+ \phi$ with $T_{c\bar{c}s1}(4220)^+ \rightarrow J/\psi K^+$ using an amplitude analysis of $B^+ \rightarrow J/\psi \phi K^+$ with a significance (accounting for systematic uncertainties) of 5.9σ . The $J^P = 1^+$ assignment is favored over 1^- with a significance of 2σ and other assignments are disfavored by 4.9σ .

 $T_{c\bar{c}s1}(4220)^+$ MASS

NODE=M260M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4220^{+50}_{-40} OUR AVERAGE		[4216 ⁺⁵⁰ ₋₄₀ MeV OUR 2023 AVERAGE]		
$4216 \pm 24^{+43}_{-30}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M260M

NEW

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.9σ .

NODE=M260M;LINKAGE=A

 $T_{c\bar{c}s1}(4220)^+$ WIDTH

NODE=M260W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
230^{+110}_{-90} OUR AVERAGE		[233 ⁺¹¹⁰ ₋₉₀ MeV OUR 2023 AVERAGE]		
$233 \pm 52^{+97}_{-73}$	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$

NODE=M260W

NEW

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.9σ .

NODE=M260W;LINKAGE=A

 $T_{c\bar{c}s1}(4220)^+$ DECAY MODES

NODE=M260215;NODE=M260

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi K^+$	seen

DESIG=1

$\Gamma(J/\psi K^+)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	24k	¹ AAIJ	21E LHCB	$B^+ \rightarrow J/\psi \phi K^+$	

NODE=M260R01
NODE=M260R01

¹ From an amplitude analysis of the decay $B^+ \rightarrow J/\psi \phi K^+$ with a significance of 5.9σ .

NODE=M260R01;LINKAGE=A

 $T_{c\bar{c}s1}(4220)^+$ REFERENCES

NODE=M260

AAIJ 21E PRL 127 082001 R. Aaij *et al.* (LHCb Collab.) JP

REFID=61150

$T_{c\bar{c}0}(4240)^+$
 $I^G(J^{PC}) = 1^+(0^{--})$
 I, G, C need confirmation.

OMITTED FROM SUMMARY TABLE

was $R_{c0}(4240)$, $X(4240)^\pm$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.Spin and parity assignment $J^P = 0^-$ is favored over 1^- , 2^- , and 2^+ by 8σ and over 1^+ by 1σ , according to the four-dimensional amplitude analysis of AAIJ 14AG.

NODE=M216

NODE=M216

 $T_{c\bar{c}0}(4240)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$4239 \pm 18^{+45}_{-10}$	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

NODE=M216M

NODE=M216M

¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $T_{c\bar{c}1}(4430)$ fit, with significance 6σ including systematic variations.

NODE=M216M;LINKAGE=AA

 $T_{c\bar{c}0}(4240)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$220 \pm 47^{+108}_{-74}$	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$

NODE=M216W

NODE=M216W

¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $T_{c\bar{c}1}(4430)$ fit, with significance 6σ including systematic variations.

NODE=M216W;LINKAGE=AA

 $T_{c\bar{c}0}(4240)^+$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^- \psi(2S)$	seen

NODE=M216215;NODE=M216

DESIG=1

 $T_{c\bar{c}0}(4240)^+$ BRANCHING RATIOS

$\Gamma(\pi^- \psi(2S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+ \pi^- \psi(2S)$	

NODE=M216225

NODE=M216R01
NODE=M216R01¹ From a 4-dimensional analysis when a second, lower mass resonance is allowed in the $T_{c\bar{c}1}(4430)$ fit. No partial branching fraction quoted.

NODE=M216R01;LINKAGE=AA

 $T_{c\bar{c}0}(4240)^+$ REFERENCES

AAIJ	14AG PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.)
------	---------------------	-----------------------	----------------

NODE=M216

REFID=55896

$T_{c\bar{c}}(4250)^+$
 $I^G(J^{PC}) = 1^-(?^{?+})$
 I, G, C need confirmation.

NODE=M192

OMITTED FROM SUMMARY TABLE

was $X(4250)^\pm$

Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

NODE=M192

Observed by MIZUK 08 in the $\pi^+\chi_{c1}(1P)$ invariant mass distribution in $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$ decays. Not seen by LEES 12B in this same mode after accounting for $K\pi$ resonant mass and angular structure.

 $T_{c\bar{c}}(4250)^+$ MASS

NODE=M192M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4250^{+190}_{-50} OUR AVERAGE	[4248 ⁺¹⁹⁰ ₋₅₀ MeV OUR 2023 AVERAGE]		
$4248^{+44}_{-29} + 180_{-35}$	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

NODE=M192M

NEW

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192M;LINKAGE=MI

 $T_{c\bar{c}}(4250)^+$ WIDTH

NODE=M192W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
180^{+320}_{-70} OUR AVERAGE	[177 ⁺³²⁰ ₋₇₀ MeV OUR 2023 AVERAGE]		
$177^{+54}_{-39} + 316_{-61}$	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$

NODE=M192W

NEW

¹ From a Dalitz plot analysis with two Breit-Wigner amplitudes.

NODE=M192W;LINKAGE=MI

 $T_{c\bar{c}}(4250)^+$ DECAY MODES

NODE=M192215;NODE=M192

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^+\chi_{c1}(1P)$	seen

DESIG=1

 $T_{c\bar{c}}(4250)^+$ BRANCHING RATIOS

NODE=M192225

 $\Gamma(\pi^+\chi_{c1}(1P))/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ MIZUK	08	BELL $\bar{B}^0 \rightarrow K^-\pi^+\chi_{c1}(1P)$
not seen	² LEES	12B	BABR $B \rightarrow K\pi\chi_{c1}(1P)$

NODE=M192R01
NODE=M192R01

••• We do not use the following data for averages, fits, limits, etc. •••

¹ With a product branching fraction measurement of $B(\bar{B}^0 \rightarrow K^- T_{c\bar{c}}(4250)^+) \times B(T_{c\bar{c}}(4250)^+ \rightarrow \pi^+\chi_{c1}(1P)) = (4.0^{+2.3+19.7}_{-0.9-0.5}) \times 10^{-5}$.

NODE=M192R01;LINKAGE=MI

² With a product branching fraction limit of $B(\bar{B}^0 \rightarrow T_{c\bar{c}}(4250)^+ K^-) \times B(T_{c\bar{c}}(4250)^+ \rightarrow \chi_{c1}\pi^+) < 4.0 \times 10^{-5}$ at 90% CL.

NODE=M192R01;LINKAGE=LE

 $T_{c\bar{c}}(4250)^+$ REFERENCES

NODE=M192

LEES	12B	PR D85 052003	J.P. Lees <i>et al.</i>	(BABAR Collab.)
MIZUK	08	PR D78 072004	R. Mizuk <i>et al.</i>	(BELLE Collab.)

REFID=54042
REFID=52535

$T_{c\bar{c}1}(4430)^+$
 $I^G(J^{PC}) = 1^+(1^{+-})$
 G, C need confirmation.
was $Z_c(4430), X(4430)^\pm$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.

First seen by CHOI 08 in $B \rightarrow K\pi^+\psi(2S)$ decays, confirmed by AAIJ 14AG, and confirmed in a model-independent way by AAIJ 15BH. Also seen by CHILIKIN 14 in $B \rightarrow K^+\pi J/\psi$ decays. J^P was determined by CHILIKIN 13 and AAIJ 14AG.

NODE=M195

NODE=M195

 $T_{c\bar{c}1}(4430)^+$ MASS

NODE=M195M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M195M

 4478^{+15}_{-18} OUR AVERAGE

$4475 \pm 7^{+15}_{-25}$	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
--------------------------	-------------------	-----------	------------------------------------

$4485 \pm 22^{+28}_{-11}$	¹ CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
---------------------------	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4443^{+15+19}_{-12-13}	² MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
--------------------------	--------------------	---------	--------------------------------

$4433 \pm 4 \pm 2$	³ CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
--------------------	-------------------	---------	--------------------------------

¹ From a four-dimensional amplitude analysis.² From a Dalitz plot analysis. Superseded by CHILIKIN 13.³ Superseded by MIZUK 09 and CHILIKIN 13.

NODE=M195M;LINKAGE=A
 NODE=M195M;LINKAGE=MI
 NODE=M195M;LINKAGE=CH

 $T_{c\bar{c}1}(4430)^+$ WIDTH

NODE=M195W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

NODE=M195W

 181 ± 31 OUR AVERAGE

$172 \pm 13^{+37}_{-34}$	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
--------------------------	-------------------	-----------	------------------------------------

200^{+41+26}_{-46-35}	¹ CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
-------------------------	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

107^{+86+74}_{-43-56}	² MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
-------------------------	--------------------	---------	--------------------------------

45^{+18+30}_{-13-13}	³ CHOI	08 BELL	$B \rightarrow K\pi^+\psi(2S)$
------------------------	-------------------	---------	--------------------------------

¹ From a four-dimensional amplitude analysis.² From a Dalitz plot analysis. Superseded by CHILIKIN 13.³ Superseded by MIZUK 09 and CHILIKIN 13.

NODE=M195W;LINKAGE=A
 NODE=M195W;LINKAGE=MI
 NODE=M195W;LINKAGE=CH

 $T_{c\bar{c}1}(4430)^+$ DECAY MODES

NODE=M195215;NODE=M195

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \pi^+\psi(2S)$	seen
$\Gamma_2 \quad \pi^+J/\psi$	seen

DESIG=1

DESIG=2

 $T_{c\bar{c}1}(4430)^+$ BRANCHING RATIOS

NODE=M195225

$\Gamma(\pi^+\psi(2S))/\Gamma_{\text{total}}$	Γ_1/Γ
---	-------------------

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

seen	¹ AAIJ	14AG LHCB	$B^0 \rightarrow K^+\pi^-\psi(2S)$
------	-------------------	-----------	------------------------------------

seen	² CHILIKIN	13 BELL	$B^0 \rightarrow K^+\pi^-\psi(2S)$
------	-----------------------	---------	------------------------------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	³ AUBERT	09AA BABR	$B \rightarrow K\pi^+\psi(2S)$
----------	---------------------	-----------	--------------------------------

seen	⁴ MIZUK	09 BELL	$B \rightarrow K\pi^+\psi(2S)$
------	--------------------	---------	--------------------------------

NODE=M195R01
 NODE=M195R01

¹ From a four-dimensional amplitude analysis. No product of branching fractions quoted.

² From a four-dimensional amplitude analysis. Measured a product of branching fractions

$$B(B^0 \rightarrow T_{c\bar{c}1}(4430)^- K^+) \times B(T_{c\bar{c}1}(4430)^- \rightarrow \psi(2S)\pi^-) = (6.0_{-2.0}^{+1.7+2.5}) \times 10^{-5}$$

³ AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 T_{c\bar{c}1}(4430)^+) \times B(T_{c\bar{c}1}(4430)^+ \rightarrow \pi^+ \psi(2S)) < 4.7 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- T_{c\bar{c}1}(4430)^+) \times B(T_{c\bar{c}1}(4430)^+ \rightarrow \pi^+ \psi(2S)) < 3.1 \times 10^{-5}$ at 95% CL.

⁴ Measured a product of branching fractions $B(\bar{B}^0 \rightarrow K^- T_{c\bar{c}1}(4430)^+) \times B(T_{c\bar{c}1}(4430)^+ \rightarrow \pi^+ \psi(2S)) = (3.2_{-0.9}^{+1.8+5.3}) \times 10^{-5}$. Superseded by CHILIKIN 13.

NODE=M195R01;LINKAGE=AA
NODE=M195R01;LINKAGE=A

NODE=M195R01;LINKAGE=AU

NODE=M195R01;LINKAGE=MI

$\Gamma(\pi^+ J/\psi)/\Gamma_{\text{total}}$

Γ_2/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	1,2 CHILIKIN	14	BELL $\bar{B}^0 \rightarrow K^- \pi^+ J/\psi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	³ AUBERT	09AA	BABR $B \rightarrow K \pi^+ J/\psi$
¹ CHILIKIN 14 reports $B(\bar{B}^0 \rightarrow T_{c\bar{c}1}(4430)^+ K^-) \times B(T_{c\bar{c}1}(4430)^+ \rightarrow J/\psi \pi^+) = (5.4_{-1.0}^{+4.0+1.1}) \times 10^{-6}$.			
² A broad enhancement seen by AAIJ 19R in the decays $B^0 \rightarrow J/\psi \pi^+ K^-$ at 4600 MeV can be due to an interplay of $T_{c\bar{c}1}(4430)$, $T_{c\bar{c}1}(4200)$ and the fitting polynomials.			
³ AUBERT 09AA quotes $B(B^+ \rightarrow \bar{K}^0 T_{c\bar{c}1}(4430)^+) \times B(T_{c\bar{c}1}(4430)^+ \rightarrow \pi^+ J/\psi) < 1.5 \times 10^{-5}$ and $B(\bar{B}^0 \rightarrow K^- T_{c\bar{c}1}(4430)^+) \times B(T_{c\bar{c}1}(4430)^+ \rightarrow \pi^+ J/\psi) < 0.4 \times 10^{-5}$ at 95% CL.			

NODE=M195R02
NODE=M195R02

NODE=M195R02;LINKAGE=A

NODE=M195R02;LINKAGE=E

NODE=M195R02;LINKAGE=AU

$T_{c\bar{c}1}(4430)^+$ REFERENCES

NODE=M195

AAIJ	19R	PRL 122 152002	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59776
AAIJ	15BH	PR D92 112009	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=57110
AAIJ	14AG	PRL 112 222002	R. Aaij <i>et al.</i>	(LHCb Collab.) JP	REFID=55896
CHILIKIN	14	PR D90 112009	K. Chilikin <i>et al.</i>	(BELLE Collab.)	REFID=56344
CHILIKIN	13	PR D88 074026	K. Chilikin <i>et al.</i>	(BELLE Collab.) JP	REFID=55551
AUBERT	09AA	PR D79 112001	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52940
MIZUK	09	PR D80 031104	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=52960
CHOI	08	PRL 100 142001	S.-K. Choi <i>et al.</i>	(BELLE Collab.)	REFID=52178

NODE=M232

$T_{b\bar{s}}(5568)^+$

$$I(J^P) = 1(?^?)$$

OMITTED FROM SUMMARY TABLE
was $X(5568)^\pm$

Seen as a peak in the $B_s \pi^\pm$ mass spectrum with a significance of more than 3σ by ABAZOV 16E and ABAZOV 18A in inclusive $p\bar{p}$ collisions at 1.96 TeV. Not seen by AAIJ 16AI, AABOUD 18L, AALTONEN 18A, and SIRUNYAN 18J. Needs confirmation.

NODE=M232

$T_{b\bar{s}}(5568)^+$ MASS

NODE=M232M

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$5566.9_{-3.1}^{+3.2+0.6}$	278	¹ ABAZOV	18A	D0 $\rho\bar{p} \rightarrow B_s^0 \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$5567.8 \pm 2.9_{-1.9}^{+0.9}$	133	² ABAZOV	16E	D0 $\rho\bar{p} \rightarrow B_s^0 \pi^\pm X$

NODE=M232M

¹ From the combined analysis of $B_s^0 \rightarrow J/\psi \phi$ and $B_s^0 \rightarrow D_s^\pm \mu^\mp X$ decays.

² Assumes $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ decay. If $T_{b\bar{s}}(5568)^\pm \rightarrow B_s^* \pi^\pm$ decay is assumed, the mass shifts upward by 49 MeV.

NODE=M232M;LINKAGE=B

NODE=M232M;LINKAGE=A

$T_{b\bar{s}}(5568)^+$ WIDTH

NODE=M232W

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$18.6_{-6.1}^{+7.9+3.5}$	278	¹ ABAZOV	18A	D0 $\rho\bar{p} \rightarrow B_s \pi^\pm X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$21.9 \pm 6.4_{-2.5}^{+5.0}$	133	ABAZOV	16E	D0 $\rho\bar{p} \rightarrow B_s \pi^\pm X$

NODE=M232W

¹ From the combined analysis of $B_s^0 \rightarrow J/\psi \phi$ and $B_s^0 \rightarrow D_s^\pm \mu^\mp X$ decays.

NODE=M232W;LINKAGE=B

$T_{b\bar{s}}(5568)^+$ DECAY MODES

NODE=M232215;NODE=M232

Mode	Fraction (Γ_i/Γ)
Γ_1 $B_s \pi^\pm$	seen

DESIG=1

 $T_{b\bar{s}}(5568)^+$ BRANCHING RATIOS

NODE=M232220

$\Gamma(B_s \pi^\pm)/\Gamma_{\text{total}}$					Γ_1/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
seen	145	¹ ABAZOV	18A D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$	
seen	133	² ABAZOV	16E D0	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
not seen		³ AABOUD	18L ATLS	$pp \rightarrow B_s^0 \pi^\pm X$	
not seen		⁴ AALTONEN	18A CDF	$p\bar{p} \rightarrow B_s^0 \pi^\pm X$	OCCUR=2
not seen		⁵ SIRUNYAN	18J CMS	$pp \rightarrow B_s^0 \pi^\pm X$	
not seen		⁶ AAIJ	16AI LHCB	$pp \rightarrow B_s^0 \pi^\pm X$	

NODE=M232R01
NODE=M232R01¹ With B_s mesons reconstructed in decays to $D_s^\pm \mu^\mp X$.NODE=M232R01;LINKAGE=F
NODE=M232R01;LINKAGE=A² Seen in $p\bar{p}$ collisions at 1.96 TeV at a rate of $(8.6 \pm 1.9 \pm 1.4)\%$ relative to inclusive B_s production in the kinematic region $10 < p_T(B_s) < 30$ GeV/c, with B_s mesons reconstructed in decays to $J/\psi\phi$. An alternative possibility, $T_{b\bar{s}}(5568)^\pm \rightarrow B_s^* \pi^\pm$ with a missing γ , could not be ruled out.³ Not seen in 24.4 fb^{-1} of pp collision data at $\sqrt{s} = 7$ and 8 TeV with B_s mesons reconstructed in decays to $J/\psi\phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 1.5% at $p_T(B_s) > 10$ GeV/c and less than 1.6% at $p_T(B_s) > 15$ GeV/c at 95% CL.

NODE=M232R01;LINKAGE=E

⁴ Not seen in 9.6 fb^{-1} of $p\bar{p}$ collision data at $\sqrt{s} = 1.96$ TeV with B_s mesons reconstructed in decays to $J/\psi\phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 6.7% at 95% CL.

NODE=M232R01;LINKAGE=D

⁵ Not seen in 19.7 fb^{-1} of pp collisions data at $\sqrt{s} = 8$ TeV with B_s mesons reconstructed in decays to $J/\psi\phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 1.1% at $p_T(B_s) > 10$ GeV/c and less than 1.0% at $p_T(B_s) > 15$ GeV/c at 95% CL.

NODE=M232R01;LINKAGE=C

⁶ Not seen in 3 fb^{-1} of pp collision data at $\sqrt{s} = 7$ and 8 TeV in a scan over the $T_{b\bar{s}}(5568)$ mass and width, with B_s mesons reconstructed in decays to $D_s^- \pi^+$ or $J/\psi\phi$. An upper limit on the production rate times branching fraction for $T_{b\bar{s}}(5568)^\pm \rightarrow B_s \pi^\pm$ relative to inclusive B_s production is less than 2.1% at $p_T(B_s) > 10$ GeV/c at 90% CL.

NODE=M232R01;LINKAGE=B

 $T_{b\bar{s}}(5568)^+$ REFERENCES

NODE=M232

AABOUD	18L	PRL 120 202007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)
AALTONEN	18A	PRL 120 202006	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	18A	PR D97 092004	V.M. Abazov <i>et al.</i>	(D0 Collab.)
SIRUNYAN	18J	PRL 120 202005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	16AI	PRL 117 152003	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	16E	PRL 117 022003	V.M. Abazov <i>et al.</i>	(D0 Collab.)

REFID=58829
REFID=58828
REFID=58937
REFID=58827
REFID=57549
REFID=57453

$T_{cc\bar{c}\bar{c}}(6900)^0$

$$I^G(J^{PC}) = 0^+(?^{?+})$$

NODE=M268

OMITTED FROM SUMMARY TABLE

was $X(6900)$ State incompatible with a $q\bar{q}$ structure. See the review on "Heavy Non- $q\bar{q}$ Mesons."

NODE=M268

 $T_{cc\bar{c}\bar{c}}(6900)^0$ MASS

NODE=M268M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
6899 ± 12 OUR AVERAGE [2023 AVERAGE]	Error includes scale factor of 1.1. [6886 ± 16 MeV OUR 2023 AVERAGE]		
6910 ± 10 ± 10	¹ AAD	23BL ATLS	$pp \rightarrow J/\psi J/\psi X$
6886 ± 11 ± 11	² AAIJ	20AY LHCB	$pp \rightarrow J/\psi J/\psi X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
6960 ± 50 ± 30	³ AAD	23BL ATLS	$pp \rightarrow J/\psi \psi(2S) X$
¹ In a model with two resonances, one describing the broad structure above threshold (mass $6650 \pm 20^{+30}_{-20}$ MeV, width $440 \pm 50^{+60}_{-50}$ MeV) interfering with single parton scattering, and a non-interfering $T_{cc\bar{c}\bar{c}}(6900)$.			
² In a model where the broad structure above threshold interferes with non-resonant single parton scattering. Without interference the mass is $6905 \pm 11 \pm 7$ MeV.			
³ Assuming a single resonance (could be another state). A 3σ signal is observed for an additional resonance with mass $7220 \pm 30^{+10}_{-40}$ MeV and width $90 \pm 60^{+60}_{-50}$ MeV.			

NODE=M268M
NEW

OCCUR=2

NODE=M268M;LINKAGE=B

NODE=M268M;LINKAGE=A

NODE=M268M;LINKAGE=C

 $T_{cc\bar{c}\bar{c}}(6900)^0$ WIDTH

NODE=M268W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
153 ± 29 OUR AVERAGE [168 ± 80 MeV OUR 2023 AVERAGE]			
150 ± 30 ± 10	¹ AAD	23BL ATLS	$pp \rightarrow J/\psi J/\psi X$
168 ± 33 ± 69	² AAIJ	20AY LHCB	$pp \rightarrow J/\psi J/\psi X$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
510 ± 170 ⁺¹¹⁰ ₋₁₀₀	³ AAD	23BL ATLS	$pp \rightarrow J/\psi \psi(2S) X$
¹ In a model with two resonances, one describing the broad structure above threshold (mass $6650 \pm 20^{+30}_{-20}$ MeV, width $440 \pm 50^{+60}_{-50}$ MeV) interfering with single parton scattering, and a non-interfering $T_{cc\bar{c}\bar{c}}(6900)$.			
² In a model where the broad structure above threshold interferes with non-resonant single parton scattering. Without interference the width is 80 ± 38 MeV.			
³ Assuming a single resonance (could be another state). A 3σ signal is observed for an additional resonance with mass $7220 \pm 30^{+10}_{-40}$ MeV and width $90 \pm 60^{+60}_{-50}$ MeV.			

NODE=M268W
NEW

OCCUR=2

NODE=M268W;LINKAGE=B

NODE=M268W;LINKAGE=C

NODE=M268W;LINKAGE=D

 $T_{cc\bar{c}\bar{c}}(6900)^0$ DECAY MODES

NODE=M268215;NODE=M268

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi J/\psi$	seen

DESIG=1

 $\Gamma(J/\psi J/\psi)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	AAD	23BL ATLS	$pp \rightarrow J/\psi J/\psi X$
seen	AAIJ	20AY LHCB	$pp \rightarrow J/\psi J/\psi X$

NODE=M268R00
NODE=M268R00 $T_{cc\bar{c}\bar{c}}(6900)^0$ REFERENCES

NODE=M268

AAD	23BL PRL 131 151902	G. Aad <i>et al.</i>	(ATLAS)
AAIJ	20AY SCIB 65 1983	R. Aaij <i>et al.</i>	(LHCb Collab.)

REFID=62432
REFID=61631

$T_{bb1}^-(10610)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

was $Z_b(10610)$, $X(10610)$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.Observed by BONDAR 12 in $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

NODE=M207

NODE=M207

 $T_{bb1}^-(10610)^\pm$ MASS

NODE=M207M

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10607.2\pm2.0	¹ BONDAR 12	BELL	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10608.5 \pm 3.4 ^{+3.7} _{-1.4}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10608.1 \pm 1.2 ^{+1.5} _{-0.2}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10607.4 \pm 1.5 ^{+0.8} _{-0.2}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10611 \pm 4 \pm 3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10609 \pm 2 \pm 3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10608 \pm 2 \pm 3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10605 \pm 2 ⁺³ ₋₁	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10599 ⁺⁶ ₋₃ ⁺⁵ ₋₄	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M207M

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.

NODE=M207M;LINKAGE=BO

NODE=M207M;LINKAGE=A

NODE=M207M;LINKAGE=BN

NODE=M207M0

 $T_{bb1}^-(10610)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10609\pm4\pm4	¹ KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$
¹ From a simultaneous fit to the KROKOVNY 13 Dalitz analysis of $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$ decays with fixed width $\Gamma(T_{bb1}^-(10610)^0) = 18.4$ MeV.			

NODE=M207M0

NODE=M207M0;LINKAGE=A

 $T_{bb1}^-(10610)^\pm$ WIDTH

NODE=M207W

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
18.4\pm 2.4	¹ BONDAR 12	BELL	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
18.5 \pm 5.3 ^{+6.1} _{-2.3}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
20.8 \pm 2.5 ^{+0.3} _{-2.1}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
18.7 \pm 3.4 ^{+2.5} _{-1.3}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
22.3 \pm 7.7 ^{+3.0} _{-4.0}	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
24.2 \pm 3.1 ^{+2.0} _{-3.0}	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
17.6 \pm 3.0 \pm 3.0	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
11.4 ⁺ ₋ 4.5 ^{+2.1} _{-3.9} ^{+2.1} _{-1.2}	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
13 ⁺¹⁰ ₋₈ ⁺⁹ ₋₇	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

NODE=M207W

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.

NODE=M207W;LINKAGE=BO

NODE=M207W;LINKAGE=A

NODE=M207W;LINKAGE=BN

$T_{b\bar{b}1}(10610)$ DECAY MODES

NODE=M207215;NODE=M207

Mode	Fraction (Γ_i/Γ)	
Γ_1 $\Upsilon(1S)\pi^+$	$(5.4_{-1.5}^{+1.9}) \times 10^{-3}$	DESIG=1
Γ_2 $\Upsilon(1S)\pi^0$	not seen	DESIG=9
Γ_3 $\Upsilon(2S)\pi^+$	$(3.6_{-0.8}^{+1.1})\%$	DESIG=2
Γ_4 $\Upsilon(2S)\pi^0$	seen	DESIG=10
Γ_5 $\Upsilon(3S)\pi^+$	$(2.1_{-0.6}^{+0.8})\%$	DESIG=3
Γ_6 $\Upsilon(3S)\pi^0$	seen	DESIG=11
Γ_7 $h_b(1P)\pi^+$	$(3.5_{-0.9}^{+1.2})\%$	DESIG=4
Γ_8 $h_b(2P)\pi^+$	$(4.7_{-1.3}^{+1.7})\%$	DESIG=5
Γ_9 $B^+\bar{B}^0$	not seen	DESIG=8
Γ_{10} $B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	$(85.6_{-2.9}^{+2.1})\%$	DESIG=6

 $T_{b\bar{b}1}(10610)$ BRANCHING RATIOS

NODE=M207225

 $\Gamma(\Upsilon(1S)\pi^+)/\Gamma_{\text{total}}$ Γ_1/Γ NODE=M207R01
NODE=M207R01

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
$5.4_{-1.3}^{+1.6+1.1}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0},$ $\pi^- \bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$

¹ Assuming the $T_{b\bar{b}1}(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M207R01;LINKAGE=A

 $\Gamma(\Upsilon(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_2/Γ NODE=M207R09
NODE=M207R09

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	KROKOVNY	13	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^0\pi^0$

 $\Gamma(\Upsilon(2S)\pi^+)/\Gamma_{\text{total}}$ Γ_3/Γ NODE=M207R02
NODE=M207R02

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$3.62_{-0.59}^{+0.76+0.79}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0},$ $\pi^- \bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

¹ Assuming the $T_{b\bar{b}1}(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M207R02;LINKAGE=A

 $\Gamma(\Upsilon(2S)\pi^0)/\Gamma_{\text{total}}$ Γ_4/Γ NODE=M207R10
NODE=M207R10

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ KROKOVNY	13	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^0\pi^0$

¹ Combined significance in $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$, including systematics, of 6.5σ .

NODE=M207R10;LINKAGE=A

 $\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$ Γ_5/Γ NODE=M207R03
NODE=M207R03

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
$2.15_{-0.42}^{+0.55+0.60}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0},$ $\pi^- \bar{B}^0 B^{*+}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

¹ Assuming the $T_{b\bar{b}1}(10610)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M207R03;LINKAGE=A

$\Gamma(\Upsilon(3S)\pi^0)/\Gamma_{\text{total}}$ Γ_6/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	¹ KROKOVNY 13	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^0\pi^0$

¹ Combined significance in $e^+e^- \rightarrow \Upsilon(2S)/\Upsilon(3S)\pi^0\pi^0$, including systematics, of 6.5σ .

NODE=M207R11
NODE=M207R11

NODE=M207R11;LINKAGE=A

 $\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$ Γ_7/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
3.45^{+0.87+0.86}_{-0.71-0.63}	¹ GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$
possibly seen	² MIZUK 16	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
seen	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M207R04
NODE=M207R04

NODE=M207R04;LINKAGE=C

NODE=M207R04;LINKAGE=A

NODE=M207R04;LINKAGE=B

 $\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
4.67^{+1.24+1.18}_{-1.00-0.89}	¹ GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$
possibly seen	² MIZUK 16	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$
seen	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=M207R05
NODE=M207R05

NODE=M207R05;LINKAGE=C

NODE=M207R05;LINKAGE=A

NODE=M207R05;LINKAGE=B

 $\Gamma(B^+ \bar{B}^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^0$

NODE=M207R08
NODE=M207R08 $[\Gamma(B^+ \bar{B}^{*0}) + \Gamma(B^{*+} \bar{B}^0)]/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
85.6^{+1.5+1.5}_{-2.0-2.1}	357	¹ GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- B^{*+} \bar{B}^0$

NODE=M207R00
NODE=M207R00

NODE=M207R00;LINKAGE=A

 $[\Gamma(B^+ \bar{B}^{*0}) + \Gamma(B^{*+} \bar{B}^0)]/[\Gamma(\Upsilon(1S)\pi^+) + \Gamma(\Upsilon(2S)\pi^+) +$ $\Gamma(\Upsilon(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)] \Gamma_{10}/(\Gamma_1 + \Gamma_3 + \Gamma_5 + \Gamma_7 + \Gamma_8)$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
5.93^{+0.99+1.01}_{-0.69-0.73}	357	¹ GARMASH 16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0}, \pi^- \bar{B}^0 B^{*+}$

NODE=M207R07
NODE=M207R07

NODE=M207R07;LINKAGE=A

¹ Combined with the results of BONDAR 12 and MIZUK 16. Not independent from $T_{b\bar{b}1}(10610)$ branching fractions to $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^+ \bar{B}^{*0} + \bar{B}^0 B^{*+}$.

 $T_{b\bar{b}1}(10610)$ REFERENCES

NODE=M207

GARMASH 16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=57446
MIZUK 16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
GARMASH 15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=56811
KROKOVNY 13	PR D88 052016	P. Krokovny <i>et al.</i>	(BELLE Collab.)	REFID=55588
BONDAR 12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)	REFID=53963

$T_{bb\bar{1}}(10650)^+$

 $I^G(J^{PC}) = 1^+(1^{+-})$
 I, G, C need confirmation.
was $Z_b(10650), X(10650)^\pm$ Properties incompatible with a $q\bar{q}$ structure (exotic state). See the review on non- $q\bar{q}$ states.Observed by BONDAR 12 in $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$). $J^P = 1^+$ is favored from angular analyses.

NODE=M208

NODE=M208

$T_{bb\bar{1}}(10650)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10652.2±1.5	¹ BONDAR 12	BELL	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
10656.7±5.0 ^{+1.1} _{-3.1}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10650.7±1.5 ^{+0.5} _{-0.2}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10651.2±1.0 ^{+0.4} _{-0.3}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10657 ±6 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
10651 ±2 ±3	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
10652 ±1 ±2	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
10654 ±3 ⁺¹ ₋₂	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
10651 ⁺² ₋₃ ⁺³ ₋₂	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.

NODE=M208M

NODE=M208M

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

NODE=M208M;LINKAGE=BO

NODE=M208M;LINKAGE=A

NODE=M208M;LINKAGE=BN

$T_{bb\bar{1}}(10650)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
11.5± 2.2	¹ BONDAR 12	BELL	$e^+e^- \rightarrow$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
12.1 ^{+11.3} _{-4.8} ^{+2.7} _{-0.6}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
14.2± 3.7 ^{+0.9} _{-0.4}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
9.3± 2.2 ^{+0.3} _{-0.5}	² GARMASH 15	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
16.3± 9.8 ^{+6.0} _{-2.0}	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
13.3± 3.3 ^{+4.0} _{-3.0}	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
8.4± 2.0± 2.0	³ BONDAR 12	BELL	$e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
20.9 ^{+5.4} _{-4.7} ^{+2.1} _{-5.7}	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
19 ± 7 ⁺¹¹ ₋₇	³ BONDAR 12	BELL	$e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹ Average of the BONDAR 12 measurements in separate channels.² Correlated with the corresponding result from BONDAR 12.³ Superseded by the average measurement of BONDAR 12.

NODE=M208W

NODE=M208W

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

NODE=M208W;LINKAGE=BO

NODE=M208W;LINKAGE=A

NODE=M208W;LINKAGE=BN

$T_{bb\bar{1}}(10650)^+$ DECAY MODES

 $T_{bb\bar{1}}(10650)^-$ decay modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Upsilon(1S)\pi^+$	$(1.7^{+0.8}_{-0.6}) \times 10^{-3}$
Γ_2 $\Upsilon(2S)\pi^+$	$(1.4^{+0.6}_{-0.4}) \%$
Γ_3 $\Upsilon(3S)\pi^+$	$(1.6^{+0.7}_{-0.5}) \%$

NODE=M208215;NODE=M208

NODE=M208

DESIG=1

DESIG=2

DESIG=3

Γ_4	$h_b(1P)\pi^+$	$(8.4^{+2.9}_{-2.4})\%$	DESIG=4
Γ_5	$h_b(2P)\pi^+$	$(15 \pm 4)\%$	DESIG=5
Γ_6	$B^+\bar{B}^0$	not seen	DESIG=8
Γ_7	$B^+\bar{B}^{*0} + B^{*+}\bar{B}^0$	not seen	DESIG=6
Γ_8	$B^{*+}\bar{B}^{*0}$	$(74^{+4}_{-6})\%$	DESIG=7

$T_{bb1}(10650)^+$ BRANCHING RATIOS

$\Gamma(\Upsilon(1S)\pi^+)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.7^{+0.7+0.3}_{-0.6-0.2}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
-----------------------------	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(1S)\pi^+\pi^-$

¹ Assuming the $T_{bb1}(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M208225

NODE=M208R01
NODE=M208R01

NODE=M208R01;LINKAGE=A

$\Gamma(\Upsilon(2S)\pi^+)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.39^{+0.48+0.34}_{-0.38-0.23}$	¹ GARMASH	16	$e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
----------------------------------	----------------------	----	---

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(2S)\pi^+\pi^-$

¹ Assuming the $T_{bb1}(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M208R02
NODE=M208R02

NODE=M208R02;LINKAGE=A

$\Gamma(\Upsilon(3S)\pi^+)/\Gamma_{\text{total}}$ Γ_3/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$1.63^{+0.53+0.39}_{-0.42-0.28}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
----------------------------------	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	GARMASH	15	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$
seen	BONDAR	12	BELL $e^+e^- \rightarrow \Upsilon(3S)\pi^+\pi^-$

¹ Assuming the $T_{bb1}(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

NODE=M208R03
NODE=M208R03

NODE=M208R03;LINKAGE=A

$\Gamma(h_b(1P)\pi^+)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$8.41^{+2.43+1.49}_{-2.12-1.06}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
----------------------------------	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	² MIZUK	16	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$
seen	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(1P)\pi^+\pi^-$

¹ Assuming the $T_{bb1}(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

² Using e^+e^- energies near the $\Upsilon(11020)$.

³ Using e^+e^- energies near the $\Upsilon(10860)$.

NODE=M208R04
NODE=M208R04

NODE=M208R04;LINKAGE=C

NODE=M208R04;LINKAGE=A
NODE=M208R04;LINKAGE=B

$\Gamma(h_b(2P)\pi^+)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$14.7^{+3.2+2.8}_{-2.8-2.3}$	¹ GARMASH	16	BELL $e^+e^- \rightarrow \pi^- B^{*+}\bar{B}^{*0}$
------------------------------	----------------------	----	--

• • • We do not use the following data for averages, fits, limits, etc. • • •

possibly seen	² MIZUK	16	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$
seen	³ BONDAR	12	BELL $e^+e^- \rightarrow h_b(2P)\pi^+\pi^-$

¹ Assuming the $T_{bb1}(10650)$ decay width is saturated by the channels $\pi^+\Upsilon(1S, 2S, 3S)$, $\pi^+h_b(1P, 2P)$, and $B^{*+}\bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16.

² Using e^+e^- energies near the $\Upsilon(11020)$.

³ Using e^+e^- energies near the $\Upsilon(10860)$.

NODE=M208R05
NODE=M208R05

NODE=M208R05;LINKAGE=C

NODE=M208R05;LINKAGE=A
NODE=M208R05;LINKAGE=B

$\Gamma(B^+\bar{B}^0)/\Gamma_{\text{total}}$					Γ_6/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT			NODE=M208R08 NODE=M208R08

not seen	GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^0$		
----------	---------	----	------	--	--	--

$[\Gamma(B^+\bar{B}^{*0}) + \Gamma(B^{*+}\bar{B}^0)]/\Gamma_{\text{total}}$					Γ_7/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT			NODE=M208R00 NODE=M208R00

not seen	GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^+ \bar{B}^{*0},$ $\pi^- \bar{B}^0 B^{*+}$		
----------	---------	----	------	--	--	--

$\Gamma(B^{*+}\bar{B}^{*0})/\Gamma_{\text{total}}$					Γ_8/Γ	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M208R06 NODE=M208R06

$73.7^{+3.4+2.7}_{-4.4-3.5}$	161	¹	GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$
------------------------------	-----	--------------	---------	----	------	--

¹ Assuming the $T_{b\bar{b}1}(10650)$ decay width is saturated by the channels $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+} \bar{B}^{*0}$, and using the results from BONDAR 12 and MIZUK 16. Using the mass and width of the $T_{b\bar{b}1}(10650)$ from BONDAR 12.

NODE=M208R06;LINKAGE=A

$\Gamma(B^{*+}\bar{B}^{*0})/[\Gamma(\Upsilon(1S)\pi^+) + \Gamma(\Upsilon(2S)\pi^+) + \Gamma(\Upsilon(3S)\pi^+) + \Gamma(h_b(1P)\pi^+) + \Gamma(h_b(2P)\pi^+)]$					$\Gamma_8/(\Gamma_1+\Gamma_2+\Gamma_3+\Gamma_4+\Gamma_5)$	
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT		NODE=M208R07 NODE=M208R07

• • • We do not use the following data for averages, fits, limits, etc. • • •

$2.80^{+0.69+0.54}_{-0.40-0.36}$	161	¹	GARMASH	16	BELL	$e^+e^- \rightarrow \pi^- B^{*+} \bar{B}^{*0}$
----------------------------------	-----	--------------	---------	----	------	--

¹ Combined with the results of BONDAR 12 and MIZUK 16. Not independent from $T_{b\bar{b}1}(10650)$ branching fractions to $\pi^+ \Upsilon(1S, 2S, 3S)$, $\pi^+ h_b(1P, 2P)$, and $B^{*+} \bar{B}^{*0}$.

NODE=M208R07;LINKAGE=A

$T_{b\bar{b}1}(10650)^+$ REFERENCES

GARMASH	16	PRL 116 212001	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=57446
MIZUK	16	PRL 117 142001	R. Mizuk <i>et al.</i>	(BELLE Collab.)	REFID=57465
GARMASH	15	PR D91 072003	A. Garmash <i>et al.</i>	(BELLE Collab.)	REFID=56811
BONDAR	12	PRL 108 122001	A. Bondar <i>et al.</i>	(BELLE Collab.)	REFID=53963

NODE=M208

Further States

NODE=M300

OMITTED FROM SUMMARY TABLE

This section contains states observed by a single group or states poorly established that thus need confirmation.

NODE=M300

QUANTUM NUMBERS, MASSES, WIDTHS, AND BRANCHING RATIOS

X(360)	$I^G(J^{PC}) = ?^?(?^?+)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	NODE=M300K08 NODE=M300K08

$360 \pm 7 \pm 9$	64 ± 18	2.3k	¹ ABRAAMYAN	09	CNTR	$2.75 dC \rightarrow \gamma\gamma X$
-------------------	-------------	------	------------------------	----	------	--------------------------------------

¹ Not seen in $pC \rightarrow \gamma\gamma X$ at 5.5 GeV/c.

NODE=M300K08;LINKAGE=AB

X(1070)	$I^G(J^{PC}) = ?^?(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	COMMENT			NODE=M300J07 NODE=M300J07

1072 ± 1	3.5 ± 0.5	¹ VLADIMIRSK...08	40	$\pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$		
--------------	---------------	------------------------------	----	--	--	--

¹ Supersedes GRIGOR'EV 05.

NODE=M300J07;LINKAGE=VL

X(1110)	$I^G(J^{PC}) = 0^+(\text{even}^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		NODE=M300J30 NODE=M300J30

1107 ± 4	$111 \pm 8 \pm 15$	DAFTARI	87	DBC	$0. \bar{p}n \rightarrow \rho^- \pi^+ \pi^-$	
--------------	--------------------	---------	----	-----	--	--

$f_0(1200-1600)$	$I^G(J^{PC}) = 0^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT		NODE=M300J98 NODE=M300J98

1323 ± 8	237 ± 20	VLADIMIRSK...06	SPEC	40	$\pi^- p \rightarrow K_S^0 K_S^0 n$	
--------------	--------------	-----------------	------	----	-------------------------------------	--

1480^{+100}_{-150}	1030^{+80}_{-170}	¹ ANISOVICH	03	SPEC		
----------------------	---------------------	------------------------	----	------	--	--

1530^{+90}_{-250}	560 ± 40	² ANISOVICH	03	SPEC		
---------------------	--------------	------------------------	----	------	--	--

OCCUR=2

¹K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$, $\pi^+ \pi^- \pi^0$, $K^+ K^- \pi^0$, $K_S^0 K_S^0 \pi^0$, $K^+ K_S^0 \pi^-$ at rest, $\bar{p} n \rightarrow \pi^- \pi^- \pi^+$, $K_S^0 K^- \pi^0$, $K_S^0 K_S^0 \pi^-$ at rest.

NODE=M300;LINKAGE=KM

²K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$, $\pi^- p \rightarrow K \bar{K} n$, $\bar{p} p \rightarrow \pi^0 \pi^0 \pi^0$, $\pi^0 \eta \eta$, $\pi^0 \pi^0 \eta$ at rest.

NODE=M300;LINKAGE=MK

X(1420) $I^G(J^{PC}) = 2^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1420±20	160 ± 10	FILIPPI	00	OBLX 0 $\bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$	NODE=M300J61 NODE=M300J61

X(1545) $I^G(J^{PC}) = ??(?^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1545±3	6.0 ± 2.5	¹ VLADIMIRSK...08		40 $\pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$	NODE=M300K07 NODE=M300K07

¹Supersedes VLADIMIRSKII 00.

NODE=M300K07;LINKAGE=VL

X(1575) $I^G(J^{PC}) = ??(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1576 ⁺⁴⁹⁺⁹⁸ ₋₅₅₋₉₁	818 ⁺²²⁺⁶⁴ ₋₂₃₋₁₃₃	¹ ABLIKIM	06S	BES $J/\psi \rightarrow K^+ K^- \pi^0$	NODE=M300J08 NODE=M300J08

¹A broad peak observed at $K^+ K^-$ invariant mass. Mass and width above are its pole position. The observed branching ratio is $B(J/\psi \rightarrow X \pi^0) B(X \rightarrow K^+ K^-) = (8.5 \pm 0.6^{+2.7}_{-3.6}) \times 10^{-4}$.

NODE=M300J08;LINKAGE=AB

X(1600) $I^G(J^{PC}) = 2^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1600±100	400 ± 200	¹ ALBRECHT 91F	ARG	10.2 $e^+ e^- \rightarrow e^+ e^- 2(\pi^+ \pi^-)$	NODE=M300J99 NODE=M300J99

¹Our estimate.

NODE=M300J99;LINKAGE=A

X(1650) $I^G(J^{PC}) = 0^-(?^{?-})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1652±7	<50	100	PROKOSHKIN 96	GAM2	32,38 $\pi p \rightarrow \omega \eta n$

NODE=M300J62
NODE=M300J62

X(1730) $I^G(J^{PC}) = ??(?^{?+})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1731.0±1.2±2.0	3.2 ± 0.8 ± 1.3	58	VLADIMIRSK...07	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 X$

NODE=M300K06
NODE=M300K06

f₂(1750) $I^G(J^{PC}) = 0^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1755±10	67 ± 12	870	¹ SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300JAM
NODE=M300JAM

Γ(K\bar{K})					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
17±5	870	² SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	NODE=M300JA1 NODE=M300JA1

Γ(γγ)					
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.13±0.04	870	² SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	NODE=M300JA2 NODE=M300JA2

Γ(ππ)					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.3±1.0	870	² SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$	NODE=M300JA3 NODE=M300JA3

$\Gamma(\eta\eta)$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2.0±0.5	870	² SCHEGELSKY 06A	RVUE	$\gamma\gamma \rightarrow K_S^0 K_S^0$

¹ From analysis of L3 data at 91 and 183–209 GeV.

² From analysis of L3 data at 91 and 183–209 GeV and using SU(3) relations.

NODE=M300JA4
NODE=M300JA4

NODE=M300JAM;LINKAGE=SC
NODE=M300JA;LINKAGE=SC

X(1775) $I^G(J^{PC}) = 1^-(?^-+)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1763±20	192 ± 60	CONDO 91	SHF	$\gamma p \rightarrow (p\pi^+)(\pi^+\pi^-\pi^-)$
1787±18	118 ± 60	CONDO 91	SHF	$\gamma p \rightarrow n\pi^+\pi^+\pi^-$

NODE=M300J60
NODE=M300J60

OCCUR=2

X(1850 - 3100) $I^G(J^{PC}) = ?^?(1^{--})$		DOCUMENT ID	TECN	COMMENT
$\Gamma(e^+e^-) \cdot B(X \rightarrow \text{hadrons})$ (eV)	CL%			
<120	90	¹ ANASHIN	11	KEDR $e^+e^- \rightarrow \text{hadrons}$

NODE=M300K28
NODE=M300K28

¹ This limit is center-of-mass energy dependent. We quote the most stringent one.

NODE=M300K28;LINKAGE=AN

X(1855) $I^G(J^{PC}) = ?^?(???)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1856.6±5	20 ± 5	BRIDGES	86D	SPEC $0. \bar{p}d \rightarrow \pi\pi N$

NODE=M300J31
NODE=M300J31

X(1870) $I^G(J^{PC}) = ?^?(2??)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1870±40	250 ± 30	ALDE	86D	GAM4 $100 \pi^- p \rightarrow 2\eta X$

NODE=M300J45
NODE=M300J45

a₃(1875) $I^G(J^{PC}) = 1^-(3^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1874±43±96	385 ± 121 ± 114	CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M300J95
NODE=M300J95

B(a₃(1875) → f₂(1270)π)/B(a₃(1875) → ρπ)		DOCUMENT ID	TECN	COMMENT
VALUE				
0.8±0.2		¹ CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M300B7
NODE=M300B7

¹ Using the observable fractions of 50.0% ρπ, 56.5% f₂π, and 11.8% ρ₃π.

NODE=M300B;LINKAGE=C1

B(a₃(1875) → ρ₃(1690)π)/B(a₃(1875) → ρπ)		DOCUMENT ID	TECN	COMMENT
VALUE				
0.9±0.3		¹ CHUNG	02	B852 $18.3 \pi^- p \rightarrow \pi^+\pi^-\pi^- p$

NODE=M300B8
NODE=M300B8

¹ Using the observable fractions of 50.0% ρπ, 56.5% f₂π, and 11.8% ρ₃π.

NODE=M300B8;LINKAGE=C1

a₁(1930) $I^G(J^{PC}) = 1^-(1^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1930 ⁺³⁰ ₋₇₀	155 ± 45	ANISOVICH	01F	SPEC $2.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J92
NODE=M300J92

X(1935) $I^G(J^{PC}) = 1^+(1^{-?})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1935±20	215 ± 30	EVANGELIS...	79	OMEG $10,16 \pi^- p \rightarrow \bar{p}pn$

NODE=M300J33
NODE=M300J33

ρ₂(1940) $I^G(J^{PC}) = 1^+(2^{--})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1940±40	155 ± 40	¹ ANISOVICH	02	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J85
NODE=M300J85

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J85;LINKAGE=AY

ω₃(1945) $I^G(J^{PC}) = 0^-(3^{--})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
1945±20	115 ± 22	¹ ANISOVICH	02B	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J65
NODE=M300J65

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J65;LINKAGE=AZ

$a_2(1950)$ $I^G(J^{PC}) = 1^-(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1950^{+30}_{-70}	180^{+30}_{-70}	¹ ANISOVICH	01F	SPEC	1.96–2.41 $p\bar{p}$

NODE=M300K24
NODE=M300K24

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K24;LINKAGE=AN

$\omega(1960)$ $I^G(J^{PC}) = 0^-(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1960 ± 25	195 ± 60	¹ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J79
NODE=M300J79

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J79;LINKAGE=AZ

$b_1(1960)$ $I^G(J^{PC}) = 1^+(1^{+-})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1960 ± 35	230 ± 50	¹ ANISOVICH	02	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J67
NODE=M300J67

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J67;LINKAGE=AY

$h_1(1965)$ $I^G(J^{PC}) = 0^-(1^{+-})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1965 ± 45	345 ± 75	¹ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J64
NODE=M300J64

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J64;LINKAGE=AZ

$f_1(1970)$ $I^G(J^{PC}) = 0^+(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1971 ± 15	240 ± 45	ANISOVICH	00J	SPEC	

NODE=M300J1
NODE=M300J1

$X(1970)$ $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1970 ± 10	40 ± 20	CHLIAPNIK...	80	HBC	32 $K^+p \rightarrow 2K_S^0 2\pi X$

NODE=M300J46
NODE=M300J46

$X(1975)$ $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1973 ± 15	80	30	CASO	70	HBC 11.2 $\pi^-p \rightarrow \rho 2\pi$

NODE=M300J47
NODE=M300J47

$\omega_2(1975)$ $I^G(J^{PC}) = 0^-(2^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
1975 ± 20	175 ± 25	¹ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J81
NODE=M300J81

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J81;LINKAGE=AZ

$a_2(1990)$ $I^G(J^{PC}) = 1^-(2^{++})$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2050 \pm 10 \pm 40$	$190 \pm 22 \pm 100$	18k	¹ SCHEGELSKY	06	RVUE $\gamma\gamma \rightarrow \pi^+\pi^-\pi^0$
$2003 \pm 10 \pm 19$	$249 \pm 23 \pm 32$		LU	05	B852 $18 \pi^-p \rightarrow \omega\pi^-\pi^0p$

NODE=M300J2
NODE=M300J2

¹ From analysis of L3 data at 183–209 GeV.

NODE=M300J2;LINKAGE=SC

$\Gamma(\gamma\gamma) \Gamma(\pi^+ \pi^- \pi^0) / \Gamma(\text{total})$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.11 \pm 0.04 \pm 0.05$	18k	¹ SCHEGELSKY 06	RVUE	$\gamma\gamma \rightarrow \pi^+ \pi^- \pi^0$

¹ From analysis of L3 data at 183–209 GeV.

NODE=M300J2G
 NODE=M300J2G

NODE=M300J2G;LINKAGE=SC

$\rho(2000)$ $I^G(J^{PC}) = 1^+(1^- -)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2000 ± 30	260 ± 45	¹ BUGG	04C	RVUE Compilation
~ 1988	~ 244	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J77
 NODE=M300J77

NODE=M300;LINKAGE=AY

$f_2(2000)$ $I^G(J^{PC}) = 0^+(2^+ +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2001 ± 10	312 ± 32	ANISOVICH	00J	SPEC
~ 1996	~ 134	HASAN	94	RVUE $\bar{p}p \rightarrow \pi\pi$

NODE=M300J25
 NODE=M300J25

$X(2000)$ $I^G(J^{PC}) = 1^-(?^?+)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1964 ± 35	225 ± 50	¹ ARMSTRONG	93D	E760	$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
~ 2100	~ 500	¹ ANTIPOV	77	CIBS	- $25 \pi^- p \rightarrow p\pi^- \rho_3$
2214 ± 15	355 ± 21	² BALTAY	77	HBC	0 $15 \pi^- p \rightarrow \Delta^{++} 3\pi$
2080 ± 40	340 ± 80	KALELKAR	75	HBC	+ $15 \pi^+ p \rightarrow p\pi^+ \rho_3$

¹ Cannot determine spin to be 3.
² BALTAY 77 favors $J^P = ,3^+$.

NODE=M300K01
 NODE=M300K01

NODE=M300K01;LINKAGE=AA
 NODE=M300K01;LINKAGE=B

$X(2000)$ $I^G(J^{PC}) = ?^?(4^+ +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
$1998 \pm 3 \pm 5$	<15	VLADIMIRSK...03	SPEC	$\pi^- p \rightarrow K_S^0 K_S^0 M M$

NODE=M300J97
 NODE=M300J97

$\eta(2010)$ $I^G(J^{PC}) = 0^+(0^- +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	
2010^{+35}_{-60}	270 ± 60	ANISOVICH	00J	SPEC

NODE=M300J5
 NODE=M300J5

$\pi_1(2015)$ $I^G(J^{PC}) = 1^-(1^- +)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
$2014 \pm 20 \pm 16$	$230 \pm 32 \pm 73$	145k	LU	05	B852 $18 \pi^- p \rightarrow \omega\pi^- \pi^0 p$
$2001 \pm 30 \pm 92$	$333 \pm 52 \pm 49$	69k	KUHN	04	B852 $18 \pi^- p \rightarrow \eta\pi^+ \pi^- \pi^- p$

NODE=M300J05
 NODE=M300J05

$a_0(2020)$ $I^G(J^{PC}) = 1^-(0^+ +)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	
2025 ± 30	330 ± 75	ANISOVICH	99C	SPEC

NODE=M300J6
 NODE=M300J6

$X(2020)$ $I^G(J^{PC}) = ?^?(?^?+)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2015 ± 3	10 ± 4	FERRER	99	RVUE $\pi p \rightarrow p\bar{p}\pi(\pi)$

NODE=M300J34
 NODE=M300J34

$h_3(2025)$ $I^G(J^{PC}) = 0^-(3^+ -)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2025 ± 20	145 ± 30	¹ ANISOVICH	02B	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\eta, \omega\pi^0 \pi^0$

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J78
 NODE=M300J78

NODE=M300J78;LINKAGE=AZ

$b_3(2030)$ $I^G(J^{PC}) = 1^+(3^+ -)$				
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2032 ± 12	117 ± 11	¹ ANISOVICH	02	SPEC $0.6-1.9 p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+ \pi^-$

NODE=M300J69
 NODE=M300J69

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J69;LINKAGE=AY

$a_2(2030)$		$I^G(J^{PC}) = 1^-(2^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2030±20	205 ± 30	¹ ANISOVICH	01F	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K23
NODE=M300K23

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K23;LINKAGE=AN

$a_3(2030)$		$I^G(J^{PC}) = 1^-(3^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2031±12	150 ± 18	¹ ANISOVICH	01F	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K20
NODE=M300K20

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K20;LINKAGE=AN

$\eta_2(2030)$		$I^G(J^{PC}) = 0^+(2^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2030±5±15	205 ± 10 ± 15	ANISOVICH	00E	SPEC	

NODE=M300J8
NODE=M300J8

$B(a_2\pi)_{L=0}/B(a_2\pi)_{L=2}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.05±0.03	¹ ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300B1
NODE=M300B1

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B1;LINKAGE=AN

$B(a_0\pi)/B(a_2\pi)_{L=2}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.10±0.08	¹ ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300B2
NODE=M300B2

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B2;LINKAGE=AN

$B(f_2\eta)/B(a_2\pi)_{L=2}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.13±0.06	¹ ANISOVICH	11	SPEC 0.9–1.94 $p\bar{p}$

NODE=M300B3
NODE=M300B3

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300B3;LINKAGE=AN

$f_3(2050)$		$I^G(J^{PC}) = 0^+(3^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2048±8	213 ± 34	ANISOVICH	00J	SPEC	2.0 $p\bar{p} \rightarrow \eta\pi^0\pi^0$

NODE=M300J7
NODE=M300J7

$f_0(2060)$		$I^G(J^{PC}) = 0^+(0^{++})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
~ 2050	~ 120	¹ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
~ 2060	~ 50	¹ OAKDEN	94	RVUE	0.36–1.55 $\bar{p}p \rightarrow \pi\pi$

NODE=M300J59
NODE=M300J59

¹ See SEMENOV 99 and KLOET 96.

OCCUR=2

NODE=M300J;LINKAGE=A

$\pi(2070)$		$I^G(J^{PC}) = 1^-(0^{-+})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2070±35	310 ⁺¹⁰⁰ ₋₅₀	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J91
NODE=M300J91

$X(2075)$		$I^G(J^{PC}) = ??(1^{+?})$			
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2084 ⁺⁴ ₋₂ ±9	58 ⁺⁴ ₋₃ ±25	^{1,2} ABLIKIM	23BG	BES3	$e^+e^- \rightarrow pK^-\bar{\Lambda}$
2075±12±5	90 ± 35 ± 9	³ ABLIKIM	04J	BES2	$J/\psi \rightarrow K^-\rho\bar{\Lambda}$

NODE=M300J01
NODE=M300J01

- ¹ The reported mass and width are the pole positions in the complex (M, Γ) plane.
² Signal observed with a statistical significance $>20\sigma$ comes from 3883 candidate events. Spin parity determined to be $J^P = 1^+$ with a statistical significance $>5\sigma$ over $0^-, 1^-, 2^+$ hypotheses, and in the range within $3.1-7.5 \sigma$ with respect to 2^- .
³ From a fit in the region $M_{p\bar{\Lambda}} - M_{p\bar{\Lambda}} - M_{\Lambda} < 150$ MeV. S -wave in the $p\bar{\Lambda}$ system preferred. A similar near-threshold enhancement in the $p\bar{\Lambda}$ system is observed in $B^+ \rightarrow p\bar{\Lambda}\bar{D}^0$ by CHEN 11F.

NODE=M300J01;LINKAGE=A
 NODE=M300J01;LINKAGE=B

NODE=M300J01;LINKAGE=AB

X(2080) $I^G(J^{PC}) = ??(???)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2080 ± 10	110 ± 20	KREYMER	80	STRC 13 $\pi^- d \rightarrow p\bar{p}n(n_s)$

NODE=M300J35
 NODE=M300J35

X(2080) $I^G(J^{PC}) = ??(3^-?)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2080 ± 10	190 ± 15	ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J37
 NODE=M300J37

a₁(2095) $I^G(J^{PC}) = 1^-(1^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)	EVTS		
2096 ± 17 ± 121	451 ± 41 ± 81	69k	KUHN	04 B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-p$

NODE=M300J04
 NODE=M300J04

B(a₁(2095) → f₁(1285)π) / B(a₁(2095) → a₁(1260))		DOCUMENT ID	TECN	COMMENT
VALUE	EVTS			
3.18 ± 0.64	69k	KUHN	04	B852 18 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-p$

NODE=M300B03
 NODE=M300B03

η(2100) $I^G(J^{PC}) = 0^+(0^{-+})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)	EVTS		
2050 ⁺³⁰⁺⁷⁵ ₋₂₄₋₂₆	250 ⁺³⁶⁺¹⁸¹ ₋₃₀₋₁₆₄			
		1	ABLIKIM	16N BES3 $J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
2103 ± 50	187 ± 75	586	2	BISELLO 89B DM2 $J/\psi \rightarrow 4\pi\gamma$

NODE=M300J48
 NODE=M300J48

- ¹ From a partial wave analysis of $J/\psi \rightarrow \gamma\phi\phi$, for which the primary signal is $\eta(2225) \rightarrow \phi\phi$, and that also finds significant signals for for 0^{-+} phase space, $f_0(2100)$, $f_2(2010)$, $f_2(2300)$, $f_2(2340)$, and a previously unseen 0^{-+} state $X(2500)$ ($M = 2470^{+15+101}$ ₋₁₉₋₂₃ MeV, $\Gamma = 230^{+64+56}$ ₋₃₅₋₃₃ MeV).

NODE=M300J48;LINKAGE=A

- ² ASTON 81B sees no peak, has 850 events in Ajinenko+Barth bins. ARESTOV 80 sees no peak.

NODE=M300J;LINKAGE=A1

X(2100) $I^G(J^{PC}) = ??(0^{??})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2100 ± 40	250 ± 40	ALDE	86D	GAM4 100 $\pi^- p \rightarrow 2\eta X$

NODE=M300J49
 NODE=M300J49

X(2110) $I^G(J^{PC}) = 1^+(3^-?)$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2110 ± 10	330 ± 20	EVANGELIS...	79	OMEG 10,16 $\pi^- p \rightarrow \bar{p}pn$

NODE=M300J36
 NODE=M300J36

X(2120) $I^G(J^{PC}) = ??(0^{??})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)	EVTS		
2122.4 ± 6.7 ^{+4.7} _{-2.7}	83 ± 16 ⁺³¹ ₋₁₁	647	ABLIKIM	11C BES3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

NODE=M300A07
 NODE=M300A07

f₂(2140) $I^G(J^{PC}) = 0^+(2^{++})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)	EVTS		
2141 ± 12	49 ± 28	389	GREEN	86 MPSF 400 $pA \rightarrow 4KX$

NODE=M300J50
 NODE=M300J50

X(2150) $I^G(J^{PC}) = ??(2^{+?})$		DOCUMENT ID	TECN	COMMENT
MASS (MeV)	WIDTH (MeV)			
2150 ± 10	260 ± 10	ROZANSKA	80	SPRK 18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J38
 NODE=M300J38

$a_2(2175)$ $I^G(J^{PC}) = 1^-(2^{++})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2175±40	310 ⁺⁹⁰ ₋₄₅	ANISOVICH	01F SPEC	2.0 $p\bar{p} \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J88
NODE=M300J88

$\eta(2190)$ $I^G(J^{PC}) = 0^+(0^{-+})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2190±50	850 ± 100	BUGG	99	BES

NODE=M300J13
NODE=M300J13

$\omega_2(2195)$ $I^G(J^{PC}) = 0^-(2^{--})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2195±30	225 ± 40	¹ ANISOVICH	02B SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J82
NODE=M300J82

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J82;LINKAGE=AZ

$X(2210)$ $I^G(J^{PC}) = ??(???)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2210 ⁺⁷⁹ ₋₂₁	203 ⁺⁴³⁷ ₋₈₇	EVANGELIS...	79B OMEG	10 $\pi^- p \rightarrow K^+ K^- n$
2207±22	130	CASO	70 HBC	11.2 $\pi^- p$

NODE=M300J51
NODE=M300J51

$X_2(2210)$ $I^G(J^{PC}) = 0^+(2^{++})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2210±60	360 ± 120	¹ KLEMP	22 RVUE	$J/\psi \rightarrow \gamma\pi^0\pi^0,$ $\gamma K_S^0 K_S^0$

NODE=M300A05
NODE=M300A05

¹ Fit of the tensor partial waves from BES3 in the multipole basis. Might be a cluster of $J^{PC} = 2^{++}$ resonances. The ratio of decay widths $K K^- / \pi\pi$ is 0.23 ± 0.05 .

NODE=M300A05;LINKAGE=A

$h_1(2215)$ $I^G(J^{PC}) = 0^-(1^{+-})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2215±40	325 ± 55	¹ ANISOVICH	02B SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J27
NODE=M300J27

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J27;LINKAGE=AZ

$\rho_2(2225)$ $I^G(J^{PC}) = 1^+(2^{--})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2225±35	335 ⁺¹⁰⁰ ₋₅₀	¹ ANISOVICH	02 SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0,$ $\omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J70
NODE=M300J70

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J70;LINKAGE=AY

$\rho_4(2230)$ $I^G(J^{PC}) = 1^+(4^{--})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2230±25	210 ± 30	¹ ANISOVICH	02 SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0,$ $\omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J74
NODE=M300J74

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J74;LINKAGE=AY

$b_1(2240)$ $I^G(J^{PC}) = 1^+(1^{+-})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2240±35	320 ± 85	¹ ANISOVICH	02 SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0,$ $\omega\eta\pi^0, \pi^+\pi^-$

NODE=M300J87
NODE=M300J87

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J87;LINKAGE=AY

$f_2(2240)$ $I^G(J^{PC}) = 0^+(2^{++})$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2240±15	241 ± 30	¹ ANISOVICH	00J SPEC	1.92-2.41 $p\bar{p}$

NODE=M300K26
NODE=M300K26

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2226 ~ 226 HASAN 94 RVUE $p\bar{p} \rightarrow \pi\pi$

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

NODE=M300K26;LINKAGE=AN

$b_3(2245)$ $I^G(J^{PC}) = 1^+(3^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN
2245 ± 50	320 ± 70	¹ BUGG	04C RVUE

NODE=M300K10
NODE=M300K10

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300K10;LINKAGE=AY

$\eta_2(2250)$ $I^G(J^{PC}) = 0^+(2^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN
2248 ± 20	280 ± 20	ANISOVICH	00I SPEC
2267 ± 14	290 ± 50	ANISOVICH	00J SPEC

NODE=M300J17
NODE=M300J17

$\pi_4(2250)$ $I^G(J^{PC}) = 1^-(4^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 15	215 ± 25	ANISOVICH	01F SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J73
NODE=M300J73

$\omega_4(2250)$ $I^G(J^{PC}) = 0^-(4^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2250 ± 30	150 ± 50	¹ ANISOVICH	02B SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J84
NODE=M300J84

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J84;LINKAGE=AZ

$\omega_5(2250)$ $I^G(J^{PC}) = 0^-(5^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN
2250 ± 70	320 ± 95	¹ BUGG	04 RVUE

NODE=M300K11
NODE=M300K11

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300;LINKAGE=AZ

$\omega_3(2255)$ $I^G(J^{PC}) = 0^-(3^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255 ± 15	175 ± 30	¹ ANISOVICH	02B SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J66
NODE=M300J66

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J66;LINKAGE=AZ

$a_4(2255)$ $I^G(J^{PC}) = 1^-(4^+)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2237 ± 5	291 ± 12	UMAN	06 E835	5.2 $\bar{p}p \rightarrow \eta\eta\pi^0$
2255 ± 40	330 ⁺¹¹⁰ ₋₅₀	¹ ANISOVICH	01F SPEC	1.96-2.41 $\bar{p}p$

NODE=M300K21
NODE=M300K21

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, and ANISOVICH 01F.

NODE=M300K21;LINKAGE=AN

$a_2(2255)$ $I^G(J^{PC}) = 1^-(2^+)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2255 ± 20	230 ± 15	¹ ANISOVICH	01G SPEC	1.96-2.41 $\bar{p}p$

NODE=M300K22
NODE=M300K22

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

NODE=M300K22;LINKAGE=AN

$X(2260)$ $I^G(J^{PC}) = 0^+(4^?)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2260 ± 20	400 ± 100	EVANGELIS...	79 OMEG	10,16 $\pi^- p \rightarrow \bar{p}pn$

NODE=M300J40
NODE=M300J40

$\rho(2270)$ $I^G(J^{PC}) = 1^+(1^-)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
2265 ± 40	325 ± 80	¹ ANISOVICH	02 SPEC	0.6-1.9 $p\bar{p} \rightarrow \omega\pi^0, \omega\eta\pi^0, \pi^+\pi^-$
2280 ± 50	440 ± 110	ATKINSON	85 OMEG	20-70 $\gamma p \rightarrow p\omega\pi^+\pi^-\pi^0$

NODE=M300J86
NODE=M300J86

¹ From the combined analysis of ANISOVICH 00J, ANISOVICH 01D, ANISOVICH 01E, and ANISOVICH 02.

NODE=M300J86;LINKAGE=AY

$a_1(2270)$ $I^G(J^{PC}) = 1^-(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2270 ⁺⁵⁵ ₋₄₀	305 ⁺⁷⁰ ₋₄₀	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J72
NODE=M300J72

$h_3(2275)$ $I^G(J^{PC}) = 0^-(3^{+-})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2275±25	190±45	¹ ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J28
NODE=M300J28

¹ From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J28;LINKAGE=AZ

$a_3(2275)$ $I^G(J^{PC}) = 1^-(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2275±35	350 ⁺¹⁰⁰ ₋₅₀	¹ ANISOVICH	01G	SPEC	1.96–2.41 $\bar{p}p$

NODE=M300K19
NODE=M300K19

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99E, ANISOVICH 01F, and ANISOVICH 01G.

NODE=M300K19;LINKAGE=AN

$\pi_2(2285)$ $I^G(J^{PC}) = 1^-(2^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2285±20±25	250±20±25	¹ ANISOVICH	11	SPEC	0.9–1.94 $p\bar{p}$

NODE=M300K25
NODE=M300K25

¹ Reanalysis of ADOMEIT 96 and ANISOVICH 00E.

NODE=M300K25;LINKAGE=AN

$\omega_3(2285)$ $I^G(J^{PC}) = 0^-(3^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2278±28	224±50	¹ BUGG	04A	RVUE	
2285±60	230±40	² ANISOVICH	02B	SPEC	0.6–1.9 $p\bar{p} \rightarrow \omega\eta, \omega\pi^0\pi^0$

NODE=M300J83
NODE=M300J83

¹ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.

NODE=M300J83;LINKAGE=BU

² From the combined analysis of ANISOVICH 00D, ANISOVICH 01C, and ANISOVICH 02B.

NODE=M300J83;LINKAGE=AZ

$\omega(2290)$ $I^G(J^{PC}) = 0^-(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2290±20	275±35	¹ BUGG	04A	RVUE	

NODE=M300J02
NODE=M300J02

¹ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.

NODE=M300J02;LINKAGE=BU

$f_2(2295)$ $I^G(J^{PC}) = 0^+(2^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2293±13	216±37	¹ ANISOVICH	00J	SPEC	1.92–2.41 $p\bar{p}$

NODE=M300K27
NODE=M300K27

¹ From the combined analysis of ANISOVICH 99C, ANISOVICH 99F, ANISOVICH 99J, ANISOVICH 99K, and ANISOVICH 00B. See also ANISOVICH 12.

NODE=M300K27;LINKAGE=AN

$f_3(2300)$ $I^G(J^{PC}) = 0^+(3^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2334±25	200±20	¹ BUGG	04A	RVUE	

NODE=M300J19
NODE=M300J19

¹ Partial wave analysis of the data on $p\bar{p} \rightarrow \bar{\Lambda}\Lambda$ from BARNES 00.

NODE=M300J19;LINKAGE=BU

$f_1(2310)$ $I^G(J^{PC}) = 0^+(1^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2310±60	255±70	ANISOVICH	00J	SPEC	

NODE=M300J23
NODE=M300J23

$\eta(2320)$ $I^G(J^{PC}) = 0^+(0^{-+})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2320±15	230±35	¹ ANISOVICH	00M	SPEC	

NODE=M300J18
NODE=M300J18

¹ From the combined analysis of $\bar{p}p \rightarrow \eta\eta\eta$ from ANISOVICH 00M and $\bar{p}p \rightarrow \eta\pi^0\pi^0$ from ANISOVICH 00J.

NODE=M300;LINKAGE=B

$\eta_4(2330)$ $I^G(J^{PC}) = 0^+(4^-+)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2328 ± 38	240 ± 90	ANISOVICH	00J	SPEC	2.0 $\bar{p}p \rightarrow \eta\pi^0\pi^0$

NODE=M300J22
NODE=M300J22

$\omega(2330)$ $I^G(J^{PC}) = 0^-(1^{--})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2330 ± 30	435 ± 75	ATKINSON	88	OMEG	25-50 $\gamma p \rightarrow \rho^\pm \rho^0 \pi^\mp$

NODE=M300J53
NODE=M300J53

$X(2340)$ $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2340 ± 20	180 ± 60	126	¹ BALTAY	75	HBC 15 $\pi^+ p \rightarrow p5\pi$

NODE=M300J54
NODE=M300J54

¹ Dominant decay into $\rho^0 \rho^0 \pi^+$. BALTAY 78 finds confirmation in $2\pi^+ \pi^- 2\pi^0$ events which contain $\rho^+ \rho^0 \pi^0$ and $2\rho^+ \pi^-$.

NODE=M300J;LINKAGE=B1

$\pi(2360)$ $I^G(J^{PC}) = 1^-(0^-+)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2360 ± 25	300 ⁺¹⁰⁰ ₋₅₀	ANISOVICH	01F	SPEC	2.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta, \pi^0\eta'$

NODE=M300J90
NODE=M300J90

$X(2360)$ $I^G(J^{PC}) = ??(4^{+?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2360 ± 10	430 ± 30	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J42
NODE=M300J42

• • • We do not use the following data for averages, fits, limits, etc. • • •					
2356 ± 7 ± 15	304 ± 28 ± 54	¹ ABLIKIM	23AY	BES3	$e^+e^- \rightarrow (\Lambda\bar{\Lambda})\eta$

¹ Assuming $J^{PC} = 1^{--}$.

NODE=M300J42;LINKAGE=A

$X(2440)$ $I^G(J^{PC}) = ??(5^{-?})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2440 ± 10	310 ± 20	ROZANSKA	80	SPRK	18 $\pi^- p \rightarrow p\bar{p}n$

NODE=M300J43
NODE=M300J43

$a_6(2450)$ $I^G(J^{PC}) = 1^-(6^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2450 ± 130	400 ± 250	CLELAND	82B	SPEC	50 $\pi p \rightarrow K_S^0 K^\pm p$

NODE=M300K12
NODE=M300K12

$X(2540)$ $I^G(J^{PC}) = 0^+(0^{++})$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2539 ± 14 ⁺³⁸ ₋₁₄	274 ⁺⁷⁷⁺¹²⁶ ₋₆₁₋₁₆₃	UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300K30
NODE=M300K30

$\Gamma(\gamma\gamma) \times B(K\bar{K})$					
VALUE (eV)		DOCUMENT ID	TECN	COMMENT	
40 ⁺⁹⁺¹⁷ ₋₇₋₄₀		UEHARA	13	BELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$

NODE=M300K3G
NODE=M300K3G

$X(2600)$ $I^G(J^{PC}) = ??(???)$					
MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT	
2618.3 ± 2.0 ^{+16.3} _{-1.4}	195 ± 5 ⁺²⁶ ₋₁₇	ABLIKIM	22G	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

NODE=M300A01
NODE=M300A01

$B(J/\psi \rightarrow \gamma X(2600)) \times B(X(2600) \rightarrow f_0(1500)\eta') \times B(f_0(1500) \rightarrow \pi^+\pi^-)$					
VALUE (units 10 ⁻⁵)		DOCUMENT ID	TECN	COMMENT	
3.09 ± 0.21 ^{+1.14} _{-0.77}		¹ ABLIKIM	22G	BES3	$J/\psi \rightarrow \gamma\pi^+\pi^-\eta'$

NODE=M300A02
NODE=M300A02

¹ The $\pi^+\pi^-$ mass spectrum is described by a coherent sum of two Breit-Wigner resonances, $f_0(1500)$ and a new $X(1540)$ with mass 1540.2 ± 7.0 ^{+36.3}_{-6.1} MeV and width 157 ± 19 ⁺¹¹₋₇₇ MeV.

NODE=M300A02;LINKAGE=A

B($J/\psi \rightarrow \gamma X(2600)$) \times B($X(2600) \rightarrow X(1540)\eta'$) \times B($X(1540) \rightarrow \pi^+ \pi^-$)

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

$2.69 \pm 0.19^{+0.38}_{-1.21}$	¹ ABLIKIM	22G BES3	$J/\psi \rightarrow \gamma \pi^+ \pi^- \eta'$
---------------------------------	----------------------	----------	---

¹ The $\pi^+ \pi^-$ mass spectrum is described by a coherent sum of two Breit-Wigner resonances, $f_0(1500)$ and a new $X(1540)$ with mass $1540.2 \pm 7.0^{+36.3}_{-6.1}$ MeV and width $157 \pm 19^{+11}_{-77}$ MeV.

NODE=M300A03
NODE=M300A03

NODE=M300A03;LINKAGE=A

 $K_0^*(2600)$ $I(J^P) = 1/2(0^+)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

$2662 \pm 59 \pm 201$	$480 \pm 47 \pm 72$	¹ AAIJ	23AH LHCB	$B^+ \rightarrow K^+(K_S^0 K \pi)$
-----------------------	---------------------	-------------------	-----------	------------------------------------

¹ From Dalitz plot analyses of $\eta_c(1S, 2S) \rightarrow K_S^0 K^+ \pi^- + c.c.$

NODE=M300A09
NODE=M300A09

NODE=M300A09;LINKAGE=A

 $X(2632)$ $I^G(J^{PC}) = ??(???)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

2635.2 ± 3.3		¹ EVDOKIMOV	04 SELX	$X(2632) \rightarrow D_S^+ \eta$
2631.6 ± 2.1	< 17	² EVDOKIMOV	04 SELX	$X(2632) \rightarrow D_S^0 K^+$

¹ From a mass difference to D_S^+ of 666.9 ± 3.3 MeV.

² From a mass difference to D_S^0 of 767.0 ± 2.0 MeV.

NODE=M300J03
NODE=M300J03

OCCUR=2

NODE=M300J03;LINKAGE=EV
NODE=M300J03;LINKAGE=ED

B($X(2632) \rightarrow D_S^0 K^+$)/B($X(2632) \rightarrow D_S^+ \eta$)

VALUE	DOCUMENT ID	TECN
-------	-------------	------

0.14 ± 0.06	¹ EVDOKIMOV	04 SELX
-----------------	------------------------	---------

¹ Possible interpretation of this decay pattern is discussed by YASUI 07.

NODE=M300B01
NODE=M300B01

NODE=M300B01;LINKAGE=YA

 $X(2680)$ $I^G(J^{PC}) = ??(???)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

2676 ± 27	150	CASO	70 HBC	$11.2 \pi^- p \rightarrow \rho^- \pi^+ \pi^- p$
---------------	-----	------	--------	---

NODE=M300J55
NODE=M300J55

 $X(2710)$ $I^G(J^{PC}) = ??(6^+?)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

2710 ± 20	170 ± 40	ROZANSKA	80 SPRK	$18 \pi^- p \rightarrow p \bar{p} n$
---------------	--------------	----------	---------	--------------------------------------

NODE=M300J44
NODE=M300J44

 $X(2750)$ $I^G(J^{PC}) = ??(7^-?)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

2747 ± 32	195 ± 75	DENNEY	83 LASS	$10 \pi^+ p \rightarrow K^+ K^- \pi^+ p$
---------------	--------------	--------	---------	--

NODE=M300J56
NODE=M300J56

 $f_6(3100)$ $I^G(J^{PC}) = 0^+(6^+ +)$

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

3100 ± 100	700 ± 130	BINON	05 GAMS	$33 \pi^- p \rightarrow \eta \eta n$
----------------	---------------	-------	---------	--------------------------------------

NODE=M300J06
NODE=M300J06

 $X(3250)$ $I^G(J^{PC}) = ??(???)$ 3-Body Decays

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

$3250 \pm 8 \pm 20$	45 ± 18	ALEEV	93 BIS2	$X(3250) \rightarrow \Lambda \bar{p} K^+$
$3265 \pm 7 \pm 20$	40 ± 18	ALEEV	93 BIS2	$X(3250) \rightarrow \bar{\Lambda} p K^-$

NODE=M300J57
NODE=M300J57

OCCUR=2

 $X(3250)$ $I^G(J^{PC}) = ??(???)$ 4-Body Decays

MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	-------------	------	---------

$3245 \pm 8 \pm 20$	25 ± 11	ALEEV	93 BIS2	$X(3250) \rightarrow \Lambda \bar{p} K^+ \pi^\pm$
$3250 \pm 9 \pm 20$	50 ± 20	ALEEV	93 BIS2	$X(3250) \rightarrow \bar{\Lambda} p K^- \pi^\mp$
$3270 \pm 8 \pm 20$	25 ± 11	ALEEV	93 BIS2	$X(3250) \rightarrow K_S^0 p \bar{p} K^\pm$

NODE=M300J58
NODE=M300J58

OCCUR=2

OCCUR=3

 $X(3350)$ $I^G(J^{PC}) = ??(???)$

MASS (MeV)	WIDTH (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	-------------	------	---------

$3350^{+10}_{-20} \pm 20$	$70^{+40}_{-30} \pm 40$	50 ± 10	¹ GABYSHEV	06A BELL	$B^- \rightarrow \Lambda_c^+ \bar{p} \pi^-$
---------------------------	-------------------------	-------------	-----------------------	----------	---

NODE=M300J09
NODE=M300J09

¹A similar enhancement in the $\Lambda_c^+ \bar{p}$ final state is also reported by BABAR collaboration in AUBERT 10H.

NODE=M300J09;LINKAGE=AU

$\psi(4500)$	$I^G(J^{PC}) = 0^-(1^{--})$	MASS (MeV)	WIDTH (MeV)	DOCUMENT ID	TECN	COMMENT
4469.1±26.2±3.6	246.3 ± 36.7 ± 9.4	¹ ABLIKIM	23X	BES3	$e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$	
4484.7±13.3±24.1	111.1 ± 30 ± 15.2	² ABLIKIM	22AU	BES3	$e^+e^- \rightarrow K^+ K^- J/\psi$	

NODE=M300K39
NODE=M300K39

¹From a cross-section measurement of $e^+e^- \rightarrow D^{*0} D^{*-} \pi^+$ between 4.189 and 4.951 GeV, assuming a coherent sum of 3 Breit-Wigner resonances plus a continuum amplitude. $\Gamma(e^+e^-) \cdot B(D^{*0} D^{*-} \pi^+) = 107\text{--}1744$ eV depending on solutions I – VIII with the same fit qualities. The two other resonances have masses (widths) 4209.6 ± 7.5 (81.6 ± 19.9) MeV and 4675.3 ± 29.7 (218.3 ± 73.5) MeV.

NODE=M300K39;LINKAGE=A

²ABLIKIM 22AU cross sections analysis of the process $e^+e^- \rightarrow K^+ K^- J/\psi$ at c.m. energies 4.127–4.600 GeV from 15.6 fb^{-1} of data.

NODE=M300K39;LINKAGE=B

REFERENCES for Further States

NODE=M300

AAIJ	23AH	PR D108 032010	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=62349
ABLIKIM	23AY	PR D107 112001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62066
ABLIKIM	23BG	PRL 131 151901	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62431
ABLIKIM	23X	PRL 130 121901	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=62072
ABLIKIM	22AU	CP C46 111002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61896
ABLIKIM	22G	PRL 129 042001	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61642
KLEMP	22	PL B830 137171	E. Klempt <i>et al.</i>	(BONN)	REFID=61646
ABLIKIM	16N	PR D93 112011	M. Ablikim	(BESIII Collab.)	REFID=57512
UEHARA	13	PTEP 2013 123C01	S. Uehara <i>et al.</i>	(BELLE Collab.)	REFID=55592
ANISOVICH	12	PR D85 014001	A.V. Anisovich <i>et al.</i>		REFID=53961
ABLIKIM	11C	PRL 106 072002	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=53684
ANASHIN	11	PL B703 543	V.V. Anashin <i>et al.</i>	(KEDR Collab.)	REFID=53932
ANISOVICH	11	EPJ C71 1511	A.V. Anisovich <i>et al.</i>	(LOQM, RAL, PNPI)	REFID=53631
CHEN	11F	PR D84 071501	P. Chen <i>et al.</i>	(BELLE Collab.)	REFID=53814
AUBERT	10H	PR D82 031102	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53363
ABRAAMYAN	09	PR C80 034001	Kh.U. Abraamyan <i>et al.</i>		REFID=53100
VLADIMIRSK...	08	PAN 71 2129	V.V. Vladimirovsky <i>et al.</i>	(ITEP)	REFID=52681
VLADIMIRSK...	07	Translated from YAF 71 2166. PAN 70 1706	V. Vladimirovsky <i>et al.</i>		REFID=52058
YASUI	07	PR D76 034009	S. Yasui, M. Oka		REFID=51907
ABLIKIM	06S	PRL 97 142002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=51451
GABYSHEV	06A	PRL 97 242001	N. Gabyshev <i>et al.</i>	(BELLE Collab.)	REFID=51565
SCHEGELSKY	06	EPJ A27 199	V.A. Schegelsky <i>et al.</i>		REFID=51186
SCHEGELSKY	06A	EPJ A27 207	V.A. Schegelsky <i>et al.</i>		REFID=51185
UMAN	06	PR D73 052009	I. Uman <i>et al.</i>	(FNAL E835)	REFID=51063
VLADIMIRSK...	06	PAN 69 493	V.V. Vladimirovsky <i>et al.</i>	(ITEP, Moscow)	REFID=51191
BINON	05	PAN 68 960	F. Binon <i>et al.</i>		REFID=50780
GRIGOR'EV	05	PAN 68 1271	V.K. Grigor'ev <i>et al.</i>	(ITEP)	REFID=50844
LU	05	PRL 94 032002	M. Lu <i>et al.</i>	(BNL E852 Collab.)	REFID=50459
ABLIKIM	04J	PRL 93 112002	M. Ablikim <i>et al.</i>	(BES Collab.)	REFID=50196
BUGG	04	PL B595 556 (errata.)	D.V. Bugg		REFID=49763
BUGG	04A	EPJ C36 161	D.V. Bugg		REFID=50158
BUGG	04C	PRPL 397 257	D.V. Bugg		REFID=50203
EVDOKIMOV	04	PRL 93 242001	A.V. Evdokimov <i>et al.</i>	(SELEX Collab.)	REFID=50337
KUHN	04	PL B595 109	J. Kuhn <i>et al.</i>	(BNL E852 Collab.)	REFID=49773
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>		REFID=49401
VLADIMIRSK...	03	PAN 66 700	V.V. Vladimirovsky <i>et al.</i>		REFID=49419
ANISOVICH	02	PL B542 8	A.V. Anisovich <i>et al.</i>		REFID=48828
ANISOVICH	02B	PL B542 19	A.V. Anisovich <i>et al.</i>		REFID=48829
CHUNG	02	PR D65 072001	S.U. Chung <i>et al.</i>	(BNL E852 Collab.)	REFID=48837
ANISOVICH	01C	PL B507 23	A.V. Anisovich <i>et al.</i>		REFID=48325
ANISOVICH	01D	PL B508 6	A.V. Anisovich <i>et al.</i>		REFID=48327
ANISOVICH	01E	PL B513 281	A.V. Anisovich <i>et al.</i>		REFID=48349
ANISOVICH	01F	PL B517 261	A.V. Anisovich <i>et al.</i>		REFID=48352
ANISOVICH	01G	PL B517 273	A.V. Anisovich <i>et al.</i>		REFID=48353
ANISOVICH	00B	NP A662 319	A.V. Anisovich <i>et al.</i>		REFID=47942
ANISOVICH	00D	PL B476 15	A.V. Anisovich <i>et al.</i>		REFID=47944
ANISOVICH	00E	PL B477 19	A.V. Anisovich <i>et al.</i>		REFID=47945
ANISOVICH	00I	PL B491 40	A.V. Anisovich <i>et al.</i>		REFID=47949
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)	REFID=47950
ANISOVICH	00M	PL B496 145	A.V. Anisovich <i>et al.</i>		REFID=48009
BARNES	00	PR C62 055203	P.D. Barnes <i>et al.</i>		REFID=47965
FILIPPI	00	PL B495 284	A. Filippi <i>et al.</i>	(OBELIX Experiment)	REFID=48006
VLADIMIRSKII	00	JETPL 72 486	V.V. Vladimirovskii <i>et al.</i>		REFID=47997
ANISOVICH	99C	Translated from ZETFP 72 698. PL B452 173	A.V. Anisovich <i>et al.</i>		REFID=46903
ANISOVICH	99E	PL B452 187	A.V. Anisovich <i>et al.</i>		REFID=46902
ANISOVICH	99F	NP A651 253	A.V. Anisovich <i>et al.</i>		REFID=46926
ANISOVICH	99J	PL B471 271	A.V. Anisovich <i>et al.</i>		REFID=47416
ANISOVICH	99K	PL B468 309	A.V. Anisovich <i>et al.</i>		REFID=47472
BUGG	99	PL B458 511	D.V. Bugg <i>et al.</i>		REFID=46938
FERRER	99	EPJ C10 249	A. Ferrer <i>et al.</i>		REFID=47404
SEMENOV	99	SPU 42 847	S.V. Semenov		REFID=47363
ADOMEIT	96	ZPHY C71 227	J. Adomeit <i>et al.</i>	(Crystal Barrel Collab.)	REFID=45202
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)	REFID=45212
PROKOSHKIN	96	PD 41 247	Y.D. Prokoshkin, V.D. Samoilenko	(SERP)	REFID=45182
		Translated from DANS 348 481.			

HASAN	94	PL B334 215	A. Hasan, D.V. Bugg	(LOQM)	REFID=44103
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)	REFID=45210
ALEEV	93	PAN 56 1358	A.N. Aleev <i>et al.</i>	(BIS-2 Collab.)	REFID=43668
		Translated from YAF 56 100.			
ARMSTRONG	93D	PL B307 399	T.A. Armstrong <i>et al.</i>	(FNAL, FERR, GENO+)	REFID=43596
ALBRECHT	91F	ZPHY C50 1	H. Albrecht <i>et al.</i>	(ARGUS Collab.)	REFID=41658
CONDO	91	PR D43 2787	G.T. Condo <i>et al.</i>	(SLAC Hybrid Collab.)	REFID=41588
BISELLO	89B	PR D39 701	G. Busetto <i>et al.</i>	(DM2 Collab.)	REFID=40575
ATKINSON	88	ZPHY C38 535	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=40556
DAFTARI	87	PRL 58 859	I.K. Daftari <i>et al.</i>	(SYRA)	REFID=40412
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)	REFID=20765
BRIDGES	86D	PL B180 313	D.L. Bridges <i>et al.</i>	(SYRA, BNL, CASE+)	REFID=21984
GREEN	86	PRL 56 1639	D.R. Green <i>et al.</i>	(FNAL, ARIZ, FSU+)	REFID=21872
ATKINSON	85	ZPHY C29 333	M. Atkinson <i>et al.</i>	(BONN, CERN, GLAS+)	REFID=22000
DENNEY	83	PR D28 2726	D.L. Denney <i>et al.</i>	(IOWA, MICH)	REFID=20754
CLELAND	82B	NP B208 228	W.E. Cleland <i>et al.</i>	(DURH, GEVA, LAUS+)	REFID=21281
ASTON	81B	NP B189 205	D. Aston <i>et al.</i>	(BONN, CERN, EPOL, GLAS+)	REFID=11553
ARESTOV	80	IHEP 80-165	Y.I. Arestov <i>et al.</i>	(SERP)	REFID=22312
CHLIAPNIK...	80	ZPHY C3 285	P.V. Chliapnikov <i>et al.</i>	(SERP, BRUX, MONS)	REFID=21996
KREYMER	80	PR D22 36	A.E. Kreymer <i>et al.</i>	(IND, PURD, SLAC+)	REFID=21970
ROZANSKA	80	NP B162 505	M. Rozanska <i>et al.</i>	(MPIM, CERN)	REFID=21774
EVANGELIS...	79	NP B153 253	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21966
EVANGELIS...	79B	NP B154 381	C. Evangelista <i>et al.</i>	(BARI, BONN, CERN+)	REFID=21967
BALTAY	78	PR D17 52	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21569
ANTIPOV	77	NP B119 45	Y.M. Antipov <i>et al.</i>	(SERP, GEVA)	REFID=20728
BALTAY	77	PRL 39 591	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU)	REFID=20847
BALTAY	75	PRL 35 891	C. Baltay <i>et al.</i>	(COLU, BING)	REFID=21994
KALELKAR	75	Thesis Nevis 207	M.S. Kalelkar	(COLU)	REFID=21564
CASO	70	LNC 3 707	C. Caso <i>et al.</i>	(GENO, HAMB, MILA, SACL)	REFID=20590
