

# $t'$ ( $4^{th}$ Generation) Quark, Searches for

NODE=Q009

## $t'(2/3)$ -quark/hadron mass limits in $p\bar{p}$ and $pp$ collisions

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt; <math>1.500 \times 10^3</math> (CL = 95%)</b> [ $> 1.280 \times 10^3$ GeV (CL = 95%) OUR 2023 BEST LIMIT]				
<b>&gt;1600</b>	95	<sup>1</sup> AAD	23AV ATLS	$B(t' \rightarrow Z t) = 1$
> 960	95	<sup>2</sup> TUMASYAN	23AX CMS	EW production, $t' \rightarrow H t$ ( $H \rightarrow \gamma\gamma$ )
<b>&gt;1500</b>	95	<sup>3</sup> TUMASYAN	23V CMS	$B(t' \rightarrow h t) = 1$
> 980	95	<sup>4</sup> AABOUD	18CE ATLS	$\geq 2\ell + \cancel{E}_T + \geq 1b_j$
>1030	95	<sup>5,6</sup> AABOUD	18CP ATLS	2,3 $\ell$ , singlet model
>1210	95	<sup>5,7</sup> AABOUD	18CP ATLS	2,3 $\ell$ , doublet model
<b>&gt;1310</b>	95	<sup>8,9</sup> AABOUD	18CR ATLS	singlet $t'$ . ATLAS combination
<b>&gt;1370</b>	95	<sup>8,10</sup> AABOUD	18CR ATLS	$t'$ in a weak isospin doublet ( $t', b'$ ). ATLAS combination.
>1140	95	<sup>11</sup> SIRUNYAN	18BM CMS	$W b, Z t, h t$ modes
> 845	95	<sup>12</sup> SIRUNYAN	18Q CMS	$B(t' \rightarrow W q) = 1$ ( $q=d,s$ )
<b>&gt;1295</b>	95	<sup>13</sup> SIRUNYAN	18W CMS	$B(t' \rightarrow W b) = 1$
> 860	95	<sup>14</sup> SIRUNYAN	17AU CMS	
> 735	95	<sup>15</sup> AAD	14AZ ATLS	$B(b' \rightarrow W t) = 1$
> 350	95	<sup>16</sup> AAD	12BC ATLS	$B(t' \rightarrow W q)=1$ ( $q=d,s,b$ )
> 420	95	<sup>17</sup> AAD	12C ATLS	$t' \rightarrow X t$ ( $m_X < 140$ GeV)
> 685	95	<sup>18</sup> CHATRCHYAN	12BH CMS	$m_{b'} = m_{t'}$
> 557	95	<sup>19</sup> CHATRCHYAN	12P CMS	$t' \bar{t}' \rightarrow W^+ b W^- \bar{b} \rightarrow$ $b \ell^+ \nu \bar{b} \ell^- \bar{\nu}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>1470	95	<sup>20</sup> AAD	23AG ATLS	$B(t' \rightarrow Z t) = 1$
>1280	95	<sup>21</sup> SIRUNYAN	19AQ CMS	$B(t' \rightarrow Z t) = 1$
>1370	95	<sup>22</sup> SIRUNYAN	19BWCMS	$B(t' \rightarrow h t) = 1$
>1010	95	<sup>23</sup> AABOUD	18CL ATLS	$B(t' \rightarrow h t) = 1$
>1160	95	<sup>24</sup> AABOUD	17L ATLS	$B(t' \rightarrow Z t) = 1$
> 770	95	<sup>25</sup> AAD	15AR ATLS	$B(t' \rightarrow W b) = 1$
> 590	95	<sup>26</sup> AAD	15BY ATLS	$W b, Z t, h t$ modes
> 745	95	<sup>27</sup> KHACHATRY...	15AI CMS	$B(t' \rightarrow h t) = 1$
> 700	95	<sup>28</sup> CHATRCHYAN	14A CMS	$B(t' \rightarrow W b) = 1$
> 706	95	<sup>28</sup> CHATRCHYAN	14A CMS	$B(t' \rightarrow Z t) = 1$
> 782	95	<sup>28</sup> CHATRCHYAN	14A CMS	$B(t' \rightarrow h t) = 1$
> 656	95	<sup>29</sup> AAD	13F ATLS	$B(t' \rightarrow W b) = 1$
> 625	95	<sup>30</sup> CHATRCHYAN	13I CMS	$B(t' \rightarrow Z t) = 1$
> 404	95	<sup>31</sup> AAD	12AR ATLS	$B(t' \rightarrow W b) = 1$
> 570	95	<sup>32</sup> CHATRCHYAN	12BC CMS	$t' \bar{t}' \rightarrow W^+ b W^- \bar{b}$
> 400	95	<sup>33</sup> AALTONEN	11AH CDF	$t' \rightarrow X t$ ( $m_X < 70$ GeV)
> 358	95	<sup>34</sup> AALTONEN	11AL CDF	$t' \rightarrow W b$
> 340	95	<sup>34</sup> AALTONEN	11AL CDF	$t' \rightarrow W q$ ( $q=d,s,b$ )
> 360	95	<sup>35</sup> AALTONEN	11O CDF	$t' \rightarrow X t$ ( $m_X < 100$ GeV)
> 285	95	<sup>36</sup> ABAZOV	11Q D0	$t' \rightarrow W q$ ( $q=d,s,b$ )
> 256	95	<sup>37,38</sup> AALTONEN	08H CDF	$t' \rightarrow W q$

<sup>1</sup> AAD 23AV based on  $139 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13$  TeV. Pair production of vector-like  $t'$  is searched for in the mode  $\ell^\pm \ell^\mp + \geq 2j$  ( $\geq 1b$ -tagged) +  $\cancel{E}_T$  or with  $3\ell$ . The data are consistent with the SM background predictions and limits are obtained for different branching ratios.

<sup>2</sup> TUMASYAN 23AX based on  $138 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13$  TeV. A vector-like  $t'$  is searched for in the  $t + H$  ( $H \rightarrow \gamma\gamma$ ) decay channel. EW production via a coupling to third-generation quarks of  $\kappa_T = 0.25$  is assumed. The branching fractions are assumed to be 50, 25, and 25%, respectively, for  $bW$ ,  $tZ$ , and  $tH$  decays.

<sup>3</sup> TUMASYAN 23V based on  $138 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13$  TeV. Pair production of vector-like  $t'$  is searched for in the single-lepton, same-sign charge dilepton and multi-lepton channels. The data are consistent with the SM background predictions and limits are obtained for different branching ratios. Masses below 1.48 TeV are excluded for all decays to third generation quarks.

<sup>4</sup> AABOUD 18CE based on  $36.1 \text{ fb}^{-1}$  of proton-proton data taken at  $\sqrt{s} = 13$  TeV. Events including a same-sign lepton pair are used. The limit is for a singlet model, assuming the branching ratios of  $t'$  into  $Zt$ ,  $Wb$  and  $Ht$  as predicted by the model.

NODE=Q009TPP

NODE=Q009TPP;CHECK LIMITS

OCCUR=2

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

NODE=Q009TPP;LINKAGE=X

NODE=Q009TPP;LINKAGE=Y

NODE=Q009TPP;LINKAGE=V

NODE=Q009TPP;LINKAGE=S

- <sup>5</sup> AABOUD 18CP based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Pair and single production of vector-like  $t'$  are searched for with at least one  $t'$  decaying into  $Zt$ . In the case of  $B(t' \rightarrow Zt) = 1$ , the limit is  $m_{t'} > 1340 \text{ GeV}$ .  
NODE=Q009TPP;LINKAGE=M
- <sup>6</sup> The limit is for the singlet model, assuming that the branching ratios into  $Zt$ ,  $Wb$ , and  $Ht$  add up to one.  
NODE=Q009TPP;LINKAGE=O
- <sup>7</sup> The limit is for the doublet model, assuming that the branching ratios into  $Zt$ ,  $Wb$ , and  $Ht$  add up to one.  
NODE=Q009TPP;LINKAGE=N
- <sup>8</sup> AABOUD 18CR based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . A combination of searches for the pair-produced vector-like  $t'$  in various decay channels ( $t' \rightarrow Wb, Zt, ht$ ). Also a model-independent limit is obtained as  $m_{t'} > 1.31 \text{ TeV}$ , assuming that the branching ratios into  $Zt$ ,  $Wb$  and  $ht$  add up to one.  
NODE=Q009TPP;LINKAGE=P
- <sup>9</sup> The limit is for the singlet  $t'$ .  
NODE=Q009TPP;LINKAGE=R
- <sup>10</sup> The limit is for  $t'$  in a weak isospin doublet ( $t', b'$ ) and  $|V_{t'b}| \ll |V_{tb}|$ .  
NODE=Q009TPP;LINKAGE=Q  
NODE=Q009TPP;LINKAGE=I
- <sup>11</sup> SIRUNYAN 18BM based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $t'$ . Three channels (single lepton, same-charge 2 leptons, or at least 3 leptons) are considered for various branching fraction combinations. Assuming  $B(tH) = 1$ , the limit is  $1270 \text{ GeV}$  and for  $B(tZ) = 1$  it is  $1300 \text{ GeV}$ .
- <sup>12</sup> SIRUNYAN 18Q based on  $19.7 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . The limit is for the pair-produced vector-like  $t'$  that couple only to light quarks. Constraints for other decay channels ( $Zq$  and  $Hq$ ) are also given in the paper.  
NODE=Q009TPP;LINKAGE=J
- <sup>13</sup> SIRUNYAN 18W based on  $35.8 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the vector-like  $t'$  pair-produced by strong interaction using lepton-plus-jets mode and assuming that  $B(t' \rightarrow Wb)$  is 100% of the production cross section and branching fraction to  $Wb$  for any new pair-produced heavy quark decaying to this channel as a narrow resonance.  
NODE=Q009TPP;LINKAGE=H
- <sup>14</sup> SIRUNYAN 17AU based on  $2.3\text{-}2.6 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Limit on pair-produced singlet vector-like  $t'$  using one lepton and several jets. The mass bound is given for a  $t'$  transforming as a singlet under the electroweak symmetry group, assumed to decay through  $W$ ,  $Z$  or Higgs boson (which decays to jets) and to a third generation quark. For a doublet, the limit is  $>830 \text{ GeV}$ . Other limits are also given in the paper.  
NODE=Q009TPP;LINKAGE=G
- <sup>15</sup> Based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . No significant excess over SM expectation is found in the search for pair production or single production of  $t'$  in the events with dilepton from a high  $p_T$   $Z$  and additional jets ( $\geq 1$   $b$ -tag). If instead of  $B(b' \rightarrow Wt) = 1$  an electroweak singlet with  $B(b' \rightarrow Wt) \sim 0.45$  is assumed, the limit reduces to  $685 \text{ GeV}$ .  
NODE=Q009TPP;LINKAGE=B
- <sup>16</sup> Based on  $1.04 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . No signal is found for the search of heavy quark pair production that decay into  $W$  and a quark in the events with dileptons, large  $\cancel{E}_T$ , and  $\geq 2$  jets.  
NODE=Q009TPP;LINKAGE=GA
- <sup>17</sup> Based on  $1.04 \text{ fb}^{-1}$  of data in  $pp$  collisions at  $7 \text{ TeV}$ . AAD 12C looked for  $t'\bar{t}'$  production followed by  $t'$  decaying into a top quark and  $X$ , an invisible particle, in a final state with an isolated high- $p_T$  lepton, four or more jets, and a large missing transverse energy. No excess over the SM  $t\bar{t}$  production gives the upper limit on  $t'\bar{t}'$  production cross section as a function of  $m_{t'}$  and  $m_X$ . The result is obtained for  $B(t' \rightarrow Wt) = 1$ .  
NODE=Q009TPP;LINKAGE=AD
- <sup>18</sup> Based on  $5 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 12BH searched for QCD and EW production of single and pair of degenerate 4<sup>th</sup> generation quarks that decay to  $Wb$  or  $Wt$ . Absence of signal in events with one lepton, same-sign dileptons or tri-leptons gives the bound. With a mass difference of  $25 \text{ GeV}/c^2$  between  $m_{t'}$  and  $m_{b'}$ , the corresponding limit shifts by about  $\pm 20 \text{ GeV}/c^2$ .  
NODE=Q009TPP;LINKAGE=CT
- <sup>19</sup> Based on  $5.0 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 12P looked for  $t'\bar{t}'$  production events with two isolated high  $p_T$  leptons, large  $\cancel{E}_T$ , and 2 high  $p_T$  jets with  $b$ -tag. The absence of signal above the SM background gives the limit for  $B(t' \rightarrow Wb) = 1$ .  
NODE=Q009TPP;LINKAGE=CH
- <sup>20</sup> AAD 23AG based on  $139 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Pair production of vector-like top or  $b_s$  is searched for in the mode  $1\ell + \geq 4j$  ( $\geq 1b$ -tagged) +  $\cancel{E}_T$ . The data are consistent with the SM background predictions and limits are obtained for different branching ratios. Masses below  $1.59 \text{ TeV}$  are excluded assuming a mass-degenerate vector-like doublet ( $t', b'$ ) model.  
NODE=Q009TPP;LINKAGE=W
- <sup>21</sup> SIRUNYAN 19AQ based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Pair production of vector-like  $t'$  is searched for with one  $t'$  decaying into  $Zt$  and the other  $t'$  decaying into  $Wb$ ,  $Zt$ ,  $ht$ . Events with an opposite-sign lepton pair consistent with coming from  $Z$  and jets are used. Mass limits are obtained for a variety of branching ratios of  $t'$ .  
NODE=Q009TPP;LINKAGE=T
- <sup>22</sup> SIRUNYAN 19BW based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $t'$  using all-hadronic final state. The analysis is made for the  $Wb$ ,  $Zt$ ,  $ht$  modes and mass limits are obtained for a variety of branching ratios.  
NODE=Q009TPP;LINKAGE=U
- <sup>23</sup> AABOUD 18CL based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . The limit is for the pair-produced vector-like  $t'$  using all-hadronic final state. The analysis is also made for the  $Wb$ ,  $Zt$ ,  $ht$  modes and mass limits are obtained for a variety of branching ratios.  
NODE=Q009TPP;LINKAGE=L
- <sup>24</sup> AABOUD 17L based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . No signal is found in the search for heavy quark pair production that decay into  $Zt$  followed by  $Z \rightarrow \nu\nu$  in  
NODE=Q009TPP;LINKAGE=F

the events with one lepton, large  $\cancel{E}_T$ , and  $\geq 4$  jets. The lower mass limit 0.87 (1.05) TeV is obtained for the singlet (doublet) model with other possible decay modes.

- 25 AAD 15AR based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Used lepton-plus-jets final state. See Fig. 20 for mass limits in the plane of  $B(t' \rightarrow Ht)$  vs.  $B(t' \rightarrow Wb)$  from a combination of  $t'\bar{t}' \rightarrow Wb + X$  and  $t'\bar{t}' \rightarrow Ht + X$  searches. Any branching ratio scenario is excluded for mass below 715 GeV.
- 26 AAD 15BY based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Limit on pair-produced vector-like  $t'$  assuming the branching fractions to  $W$ ,  $Z$ , and  $h$  modes of the singlet model. Used events containing  $\geq 2\ell + \cancel{E}_T + \geq 2j$  ( $\geq 1b$ ) and including a same-sign lepton pair.
- 27 KHACHATRYAN 15AI based on  $19.7 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . The search exploits all-hadronic final states by tagging boosted Higgs boson using jet substructure and  $b$ -tagging.
- 28 Based on  $19.5 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . The  $t'$  quark is pair produced and is assumed to decay into three different final states of  $bW$ ,  $tZ$ , and  $th$ . The search is carried out using events with at least one isolated lepton.
- 29 Based on  $4.7 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . No signal is found for the search of heavy quark pair production that decay into  $W$  and a  $b$  quark in the events with a high  $p_T$  isolated lepton, large  $\cancel{E}_T$  and at least 3 jets ( $\geq 1b$ -tag). Vector-like quark of charge  $2/3$  with  $400 < m_{t'} < 550 \text{ GeV}$  and  $B(t' \rightarrow Wb) > 0.63$  is excluded at 95% CL.
- 30 Based on  $5.0 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 13I looked for events with one isolated electron or muon, large  $\cancel{E}_T$ , and at least four jets with large transverse momenta, where one jet is likely to originate from the decay of a bottom quark.
- 31 Based on  $1.04 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . No signal is found in the search for pair produced heavy quarks that decay into  $W$  boson and a  $b$  quark in the events with a high  $p_T$  isolated lepton, large  $\cancel{E}_T$  and at least 3 jets ( $\geq 1b$ -tag).
- 32 Based on  $5.0 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 7 \text{ TeV}$ . CHATRCHYAN 12BC looked for  $t'\bar{t}'$  production events with a single isolated high  $p_T$  lepton, large  $\cancel{E}_T$  and at least 4 high  $p_T$  jets with a  $b$ -tag. The absence of signal above the SM background gives the limit for  $B(t' \rightarrow Wb) = 1$ .
- 33 Based on  $5.7 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at 1.96 TeV. AALTONEN 11AH looked for  $t'\bar{t}'$  production followed by  $t'$  decaying into a top quark and  $X$ , an invisible particle, in the all hadronic decay mode of  $t\bar{t}$ . No excess over the SM  $t\bar{t}$  production gives the upper limit on  $t'\bar{t}'$  production cross section as a function of  $m_{t'}$  and  $m_X$ . The result is obtained for  $B(t' \rightarrow Xt) = 1$ .
- 34 Based on  $5.6 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at 1.96 TeV. AALTONEN 11AL looked for  $\ell + \geq 4j$  events and set upper limits on  $\sigma(t'\bar{t}')$  as functions of  $m_{t'}$ .
- 35 Based on  $4.8 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at 1.96 TeV. AALTONEN 11O looked for  $t'\bar{t}'$  production signal when  $t'$  decays into a top quark and  $X$ , an invisible particle, in  $\ell + \cancel{E}_T + \text{jets}$  channel. No excess over the SM  $t\bar{t}$  production gives the upper limit on  $t'\bar{t}'$  production cross section as a function of  $m_{t'}$  and  $m_X$ . The result is obtained for  $B(t' \rightarrow Xt) = 1$ .
- 36 Based on  $5.3 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at 1.96 TeV. ABAZOV 11Q looked for  $\ell + \cancel{E}_T + \geq 4j$  events and set upper limits on  $\sigma(t'\bar{t}')$  as functions of  $m_{t'}$ .
- 37 Searches for pair production of a new heavy top-like quark  $t'$  decaying to a  $W$  boson and another quark by fitting the observed spectrum of total transverse energy and reconstructed  $t'$  mass in the lepton + jets events.
- 38 HUANG 08 reexamined the  $t'$  mass lower bound of 256 GeV obtained in AALTONEN 08H that assumes  $B(b' \rightarrow qZ) = 1$  for  $q = u, c$  which does not hold when  $m_{b'} < m_{t'} - m_W$  or the mixing  $\sin^2(\theta_{bt'})$  is so tiny that the decay occurs outside of the vertex detector. Fig. 1 gives that lower bound on  $m_{t'}$  in the plane of  $\sin^2(\theta_{bt'})$  and  $m_{b'}$ .

NODE=Q009TPP;LINKAGE=C

NODE=Q009TPP;LINKAGE=D

NODE=Q009TPP;LINKAGE=E

NODE=Q009TPP;LINKAGE=HA

NODE=Q009TPP;LINKAGE=AG

NODE=Q009TPP;LINKAGE=A

NODE=Q009TPP;LINKAGE=GD

NODE=Q009TPP;LINKAGE=CA

NODE=Q009TPP;LINKAGE=TN

NODE=Q009TPP;LINKAGE=ON

NODE=Q009TPP;LINKAGE=LT

NODE=Q009TPP;LINKAGE=AB

NODE=Q009TPP;LINKAGE=AA

NODE=Q009TPP;LINKAGE=HU

### $t'(5/3)$ -quark/hadron mass limits in $p\bar{p}$ and $pp$ collisions

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
<b>&gt; 1.460 <math>\times 10^3</math> (CL = 95%)</b> [ <b>&gt;1.350 <math>\times 10^3</math> GeV (CL = 95%) OUR 2023 BEST LIMIT</b> ]				
>1460	95	1 AAD	23AG ATLS	$t'(5/3) \rightarrow tW^+$
>1330	95	2 SIRUNYAN	19T CMS	$t'_R(5/3) \rightarrow tW^+$
>1300	95	2 SIRUNYAN	19T CMS	$t'_L(5/3) \rightarrow tW^+$
>1190	95	3 AABOUD	18CE ATLS	$\geq 2\ell + \cancel{E}_T + \geq 1bj$
>1020	95	4 SIRUNYAN	17J CMS	$t'_R(5/3) \rightarrow tW^+$
> 990	95	4 SIRUNYAN	17J CMS	$t'_L(5/3) \rightarrow tW^+$
> 750	95	5 AAD	15BY ATLS	$t'(5/3) \rightarrow tW^+$
> 840	95	6 AAD	15Z ATLS	$t'(5/3) \rightarrow tW^+$
> 800	95	7 CHATRCHYAN 14T	CMS	$t'(5/3) \rightarrow tW^+$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>1350	95	8 AABOUD	18AW ATLS	$t'(5/3) \rightarrow tW^+$

NODE=Q009TP5  
NODE=Q009TP5

OCCUR=2

OCCUR=2

- <sup>1</sup> AAD 23AG based on  $139 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Pair production of vector-like top or  $b'$  is searched for in the mode  $1\ell + \geq 4j (\geq 1b\text{-tagged}) + \cancel{E}_T$ . The data are consistent with the SM background predictions and limits are obtained for different branching ratios.
- <sup>2</sup> SIRUNYAN 19T based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Signals are searched in the final states of  $t'$  pair production, with same-sign leptons (which come from a  $t'$  decay) or a single lepton (which comes from a  $W$  out of  $4Ws$ ), along with jets, and no excess over the SM expectation is found.
- <sup>3</sup> AABOUD 18CE based on  $36.1 \text{ fb}^{-1}$  of proton-proton data taken at  $\sqrt{s} = 13 \text{ TeV}$ . Events including a same-sign lepton pair are used. The limit is for the pair-produced vector-like  $t'$ . With single  $t'$  production included, assuming  $t'tW$  coupling of one, the limit is  $m_{t'} > 1.6 \text{ TeV}$ .
- <sup>4</sup> SIRUNYAN 17J based on  $2.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Signals are searched in the final states of  $t'$  pair production, with same-sign leptons (which come from a  $t'$  decay) or a single lepton (which comes from a  $W$  out of  $4Ws$ ), along with jets, and no excess over the SM expectation is found.
- <sup>5</sup> AAD 15BY based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Limit on  $t'(5/3)$  in pair and single production assuming its coupling to  $Wt$  is equal to one. Used events containing  $\geq 2\ell + \cancel{E}_T + \geq 2j (\geq 1b)$  and including a same-sign lepton pair.
- <sup>6</sup> AAD 15Z based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Used events with  $\ell + \cancel{E}_T + \geq 6j (\geq 1b)$  and at least one pair of jets from weak boson decay, sensitive to the final state  $b\bar{b}W^+W^-W^+W^-$ .
- <sup>7</sup> CHATRCHYAN 14T based on  $19.5 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . Non-observation of anomaly in  $H_T$  distribution in the same-sign dilepton events leads to the limit when pair produced  $t'(5/3)$  quark decays exclusively into  $t$  and  $W^+$ , resulting in the final state with  $b\bar{b}W^+W^-W^+W^-$ .
- <sup>8</sup> AABOUD 18AW based on  $36.1 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Limit on  $t'(5/3)$  in pair production assuming its coupling to  $Wt$  is equal to one. Lepton-plus-jets final state is used, characterized by  $\ell + \cancel{E}_T + \text{jets} (\geq 1b\text{-tagged})$ .

NODE=Q009TP5;LINKAGE=J

NODE=Q009TP5;LINKAGE=G

NODE=Q009TP5;LINKAGE=F

NODE=Q009TP5;LINKAGE=D

NODE=Q009TP5;LINKAGE=B

NODE=Q009TP5;LINKAGE=C

NODE=Q009TP5;LINKAGE=A

NODE=Q009TP5;LINKAGE=E

### $t'(2/3)$ mass limits from single production in $p\bar{p}$ and $pp$ collisions

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>950	95	<sup>1</sup> AAD	16AV ATLS	$qg \rightarrow q't'b, B(t' \rightarrow Wb)=0.5$
>403	95	<sup>2</sup> ABAZOV	11F D0	$qd \rightarrow q't' \rightarrow q'(Wd)$ $\tilde{\kappa}_{dt'}=1, B(t' \rightarrow Wd)=1$
>551	95	<sup>2</sup> ABAZOV	11F D0	$qu \rightarrow q't' \rightarrow q(Zu)$ $\tilde{\kappa}_{ut'}=\sqrt{2}, B(t' \rightarrow Zu)=1$

NODE=Q009TPS

NODE=Q009TPS;CHECK LIMITS

OCCUR=2

• • • We do not use the following data for averages, fits, limits, etc. • • •

- <sup>3</sup> AAD 22G ATLS  $t' \rightarrow Ht$ , singlet  $t'$
- <sup>4</sup> TUMASYAN 22X CMS  $t' \rightarrow Zt$

- <sup>1</sup> AAD 16AV based on  $20.3 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 8 \text{ TeV}$ . No significant excess over SM expectation is found in the search for a fully reconstructed vector-like  $t'$  in the mode  $\ell + \cancel{E}_T + \geq 2j (\geq 1b)$ . A veto on massive large-radius jets is used to reject the  $t\bar{t}$  background.
- <sup>2</sup> ABAZOV 11F based on  $5.4 \text{ fb}^{-1}$  of data in  $p\bar{p}$  collisions at  $1.96 \text{ TeV}$ . It looked for single production of  $t'$  via the  $Z$  or  $E$  coupling to the first generation up or down quarks, respectively. Model independent cross section limits for the single production processes  $p\bar{p} \rightarrow t'q \rightarrow (Wd)q$ , and  $p\bar{p} \rightarrow t'q \rightarrow (Zd)q$  are given in Figs. 3 and 4, respectively, and the mass limits are obtained for the model of ATRE 09 with degenerate bi-doublets of vector-like quarks.
- <sup>3</sup> AAD 22G based on  $139 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . No significant excess over SM expectation is found in the search for a vector-like  $t'$  in the  $Ht$  decay channel, where  $H$  and  $t$  are reconstructed as single jets. The mass range between 1.0 and 2.3 TeV is targeted and 95% CL limits on the production section times the decay branching fraction are set depending on the coupling and mass of  $t'$ .
- <sup>4</sup> TUMASYAN 22X based on  $137 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . No significant excess over SM expectation is found in the search for a vector-like  $t'$  in the  $Zt$  decay channel, where  $Z$  decays to neutrinos and  $t$  decays hadronically. The 95% CL limits on the production section times the decay branching fraction are set depending on the coupling and mass of  $t'$ .

NODE=Q009TPS;LINKAGE=A

NODE=Q009TPS;LINKAGE=AB

NODE=Q009TPS;LINKAGE=B

NODE=Q009TPS;LINKAGE=C

### $t'(5/3)$ mass limits from single production in $p\bar{p}$ and $pp$ collisions

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

- <sup>1</sup> SIRUNYAN 19AI CMS  $tW \rightarrow t'(5/3) \rightarrow tW$

- <sup>1</sup> SIRUNYAN 19AI based on  $35.9 \text{ fb}^{-1}$  of  $pp$  data at  $\sqrt{s} = 13 \text{ TeV}$ . Exclusion limits are set on the product of the production cross section and branching fraction for the  $b'(-1/3) + t$  and  $t'(5/3) + t$  modes as a function of the vector-like quark mass in Fig. 8 and Tab. 2 for relative vector-like quark widths between 1 and 30% for left- and right-handed vector-like quark couplings. No significant deviation from the SM prediction is observed.

NODE=Q009T5S

NODE=Q009T5S

NODE=Q009T5S;LINKAGE=A

REFERENCES FOR Searches for (Fourth Generation)  $t'$  Quark

NODE=Q009

AAD	23AG	EPJ C83 719	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62172
AAD	23AV	PL B843 138019	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62364
TUMASYAN	23AX	JHEP 2309 057	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62525
TUMASYAN	23V	JHEP 2307 020	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62165
AAD	22G	PR D105 092012	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61744
TUMASYAN	22X	JHEP 2205 093	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61801
SIRUNYAN	19AI	EPJ C79 90	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59702
SIRUNYAN	19AQ	EPJ C79 364	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59721
SIRUNYAN	19BW	PR D100 072001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60010
SIRUNYAN	19T	JHEP 1903 082	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59657
AABOUD	18AW	JHEP 1808 048	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59145
AABOUD	18CE	JHEP 1812 039	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59369
AABOUD	18CL	PR D98 092005	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59472
AABOUD	18CP	PR D98 112010	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59500
AABOUD	18CR	PRL 121 211801	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59532
SIRUNYAN	18BM	JHEP 1808 177	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59137
SIRUNYAN	18Q	PR D97 072008	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58920
SIRUNYAN	18W	PL B779 82	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58999
AABOUD	17L	JHEP 1708 052	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57829
SIRUNYAN	17AU	JHEP 1711 085	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58344
SIRUNYAN	17J	JHEP 1708 073	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=57831
AAD	16AV	EPJ C76 442	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57383
AAD	15AR	JHEP 1508 105	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56648
AAD	15BY	JHEP 1510 150	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56863
AAD	15Z	PR D91 112011	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56592
KHACHATRYAN	15AI	JHEP 1506 080	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56636
AAD	14AZ	JHEP 1411 104	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56201
CHATRCHYAN	14A	PL B729 149	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55674
CHATRCHYAN	14T	PRL 112 171801	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55839
AAD	13F	PL B718 1284	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54843
CHATRCHYAN	13I	JHEP 1301 154	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54941
AAD	12AR	PRL 108 261802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54227
AAD	12BC	PR D86 012007	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54358
AAD	12C	PRL 108 041805	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54080
CHATRCHYAN	12BC	PL B718 307	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54693
CHATRCHYAN	12BH	PR D86 112003	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54772
CHATRCHYAN	12P	PL B716 103	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54183
AALTONEN	11AH	PRL 107 191803	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53835
AALTONEN	11AL	PRL 107 261801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53945
AALTONEN	11O	PRL 106 191801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=16449
ABAZOV	11F	PRL 106 081801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=16469
ABAZOV	11Q	PRL 107 082001	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=53709
ATRE	09	PR D79 054018	A. Atre <i>et al.</i>		REFID=54081
AALTONEN	08H	PRL 100 161803	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52231
HUANG	08	PR D77 037302	P.Q. Hung, M. Sher	(UVA, WILL)	REFID=52505