

# Neutral Higgs Bosons, Searches for

NODE=S055

## CONTENTS:

NODE=S055CNT  
NODE=S055CNT

- Mass Limits for Neutral Higgs Bosons in Supersymmetric Models
  - Mass Limits for heavy neutral Higgs bosons ( $H_2^0$ ,  $A^0$ ) in the MSSM
  - Mass Limits for  $H_1^0$  (Higgs Boson) in Supersymmetric Models
- Mass Limits for Neutral Higgs Bosons in Extended Higgs Models
  - Mass Limits in General two-Higgs-doublet Models
  - Mass Limits for  $H^0$  with Vanishing Yukawa Couplings
  - Mass Limits for  $H^0$  Decaying to Invisible Final States
  - Mass Limits for Light  $A^0$
  - Other Mass Limits
- Electroweak Constraints on the Standard Model Higgs Boson Mass

NODE=S055CNT

## MASS LIMITS FOR NEUTRAL HIGGS BOSONS IN SUPERSYMMETRIC MODELS

NODE=S055250

The minimal supersymmetric model has two complex doublets of Higgs bosons. The resulting physical states are two scalars [ $H_1^0$  and  $H_2^0$ , where we define  $m_{H_1^0} < m_{H_2^0}$ ], a pseudoscalar ( $A^0$ ), and a charged Higgs pair ( $H^\pm$ ).  $H_1^0$  and  $H_2^0$  are also called  $h$  and  $H$  in the literature. There are two free parameters in the Higgs sector which can be chosen to be  $m_{A^0}$  and  $\tan\beta = v_2/v_1$ , the ratio of vacuum expectation values of the two Higgs doublets. Tree-level Higgs masses are constrained by the model to be  $m_{H_1^0} \leq m_Z$ ,  $m_{H_2^0} \geq m_Z$ ,  $m_{A^0} \geq m_{H_1^0}$ , and  $m_{H^\pm} \geq m_W$ . However, as described in the review on "Status of Higgs Boson Physics" in this Volume these relations are violated by radiative corrections.

NODE=S055250

The observed signal at about 125 GeV, see section "H", can be interpreted as one of the neutral Higgs bosons of supersymmetric models. Unless otherwise noted, we identify the lighter scalar  $H_1^0$  with the Higgs discovered at 125 GeV at the LHC (AAD 12AI, CHATRCHYAN 12N).

Unless otherwise noted, the experiments in  $e^+e^-$  collisions search for the processes  $e^+e^- \rightarrow H_1^0 Z^0$  in the channels used for the Standard Model Higgs searches and  $e^+e^- \rightarrow H_1^0 A^0$  in the final states  $b\bar{b}b\bar{b}$  and  $b\bar{b}\tau^+\tau^-$ . Unless otherwise stated, the following results assume no invisible  $H_1^0$  or  $A^0$  decays. Unless otherwise noted, the results are given in the  $m_h^{max}$  scenario, CARENA 13.

In  $p\bar{p}$  and  $pp$  collisions the experiments search for a variety of processes, as explicitly specified for each entry. Limits on the  $A^0$  mass arise from these direct searches, as well as from the relations valid in the minimal supersymmetric model between  $m_{A^0}$  and  $m_{H_1^0}$ . As discussed in the review on "Status of Higgs Boson Physics" in this Volume, these relations depend, via potentially large radiative corrections, on the mass of the  $t$  quark and on the supersymmetric parameters, in particular those of the stop sector. These indirect limits are weaker for larger  $t$  and  $\tilde{t}$  masses. To include the radiative corrections to the Higgs masses, unless otherwise stated, the listed papers use theoretical predictions incorporating two-loop corrections and beyond (SLAVICH 21), and the results are given for the  $M_h^{125}$  benchmark scenario, see BAGNASCHI 19.

## Mass Limits for heavy neutral Higgs bosons ( $H_2^0$ , $A^0$ ) in the MSSM

The limits rely on  $pp \rightarrow H_2^0/A^0 \rightarrow \tau^+\tau^-$  and assume that  $H_2^0$  and  $A^0$  are (sufficiently) mass degenerate. The limits depend on  $\tan\beta$ .

NODE=S055HAD

NODE=S055HAD

NODE=S055HAD

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 835	95	1 TUMASYAN	23S CMS	$\tan\beta = 10$ GeV
>1240	95	1 TUMASYAN	23S CMS	$\tan\beta = 20$ GeV
>1605	95	1 TUMASYAN	23S CMS	$\tan\beta = 30$ GeV
>1820	95	1 TUMASYAN	23S CMS	$\tan\beta = 40$ GeV
>1950	95	1 TUMASYAN	23S CMS	$\tan\beta = 50$ GeV

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

>2062	95	1	TUMASYAN	23S CMS	$\tan\beta = 60$ GeV	OCCUR=6
>1121	95	2	AAD	20AA ATLS	$\tan\beta = 10$ GeV	
>1475	95	2	AAD	20AA ATLS	$\tan\beta = 20$ GeV	OCCUR=2
>1677	95	2	AAD	20AA ATLS	$\tan\beta = 30$ GeV	OCCUR=3
>1826	95	2	AAD	20AA ATLS	$\tan\beta = 40$ GeV	OCCUR=4
>1937	95	2	AAD	20AA ATLS	$\tan\beta = 50$ GeV	OCCUR=5
>2033	95	2	AAD	20AA ATLS	$\tan\beta = 60$ GeV	OCCUR=6

• • • We do not use the following data for averages, fits, limits, etc. • • •

		3	AAD	24AP ATLS	$H_2^0, A^0 \rightarrow t\bar{t}$	
		4	AAD	24CB ATLS	$H$ properties	
		5	AAD	24H ATLS	$H_2^0 \rightarrow HH$	
		6	AAD	20 ATLS	$H$ properties	
		7	AAD	20C ATLS	$H_2^0 \rightarrow HH$	
		8	AAD	20L ATLS	$H_2^0 \rightarrow b\bar{b}$	
		9	SIRUNYAN	20AC CMS	$A^0 \rightarrow ZH$	
		10	SIRUNYAN	20AF CMS	$H_2^0/A^0 \rightarrow t\bar{t}$	
		11	SIRUNYAN	20Y CMS	$H_2^0 \rightarrow W^+W^-$	
		12	SIRUNYAN	19CR CMS	$H_2^0/A^0 \rightarrow \mu^+\mu^-$	
> 377	95	13	AABOUD	18G ATLS	$\tan\beta = 10$ GeV	
> 863	95	13	AABOUD	18G ATLS	$\tan\beta = 20$ GeV	OCCUR=2
>1157	95	13	AABOUD	18G ATLS	$\tan\beta = 30$ GeV	OCCUR=3
>1328	95	13	AABOUD	18G ATLS	$\tan\beta = 40$ GeV	OCCUR=4
>1483	95	13	AABOUD	18G ATLS	$\tan\beta = 50$ GeV	OCCUR=5
>1613	95	13	AABOUD	18G ATLS	$\tan\beta = 60$ GeV	OCCUR=6
		14	SIRUNYAN	18A CMS	$H_2^0 \rightarrow H^0 H^0$	
		15	SIRUNYAN	18BP CMS	$pp \rightarrow H_2^0/A^0 + b + X,$ $H_2^0/A^0 \rightarrow b\bar{b}$	
> 389	95	16	SIRUNYAN	18CX CMS	$\tan\beta = 10$ GeV	
> 832	95	16	SIRUNYAN	18CX CMS	$\tan\beta = 20$ GeV	OCCUR=2
>1148	95	16	SIRUNYAN	18CX CMS	$\tan\beta = 30$ GeV	OCCUR=3
>1341	95	16	SIRUNYAN	18CX CMS	$\tan\beta = 40$ GeV	OCCUR=4
>1496	95	16	SIRUNYAN	18CX CMS	$\tan\beta = 50$ GeV	OCCUR=5
>1613	95	16	SIRUNYAN	18CX CMS	$\tan\beta = 60$ GeV	OCCUR=6
		17	AABOUD	16AA ATLS	$A^0 \rightarrow \tau^+\tau^-$	
		18	KHACHATRY...16A	CMS	$H_{1,2}^0/A^0 \rightarrow \mu^+\mu^-$	
		19	KHACHATRY...16P	CMS	$H_2^0 \rightarrow H^0 H^0, A^0 \rightarrow ZH^0$	
		20	KHACHATRY...15AY	CMS	$pp \rightarrow H_{1,2}^0/A^0 + b + X,$ $H_{1,2}^0/A^0 \rightarrow b\bar{b}$	OCCUR=4
		21	AAD	14AW ATLS	$pp \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau\tau$	
		22	KHACHATRY...14M	CMS	$pp \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau\tau$	
		23	AAD	13O ATLS	$pp \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-,$ $\mu^+\mu^-$	
		24	AAIJ	13T LHCB	$pp \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$	
		25	CHATRCHYAN	13AG CMS	$pp \rightarrow H_{1,2}^0/A^0 + b + X,$ $H_{1,2}^0/A^0 \rightarrow b\bar{b}$	
		26	AALTONEN	12AQ TEVA	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + b + X,$ $H_{1,2}^0/A^0 \rightarrow b\bar{b}$	
		27	AALTONEN	12X CDF	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + b + X,$ $H_{1,2}^0/A^0 \rightarrow b\bar{b}$	
		28	ABAZOV	12G D0	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$	
		29	CHATRCHYAN	12K CMS	$pp \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$	

		30	ABAZOV	11K D0	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + b + X,$ $H_{1,2}^0/A^0 \rightarrow b\bar{b}$
		31	ABAZOV	11W D0	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + b + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$
		32	AALTONEN	09AR CDF	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$
> 90.4		33	ABDALLAH	08B DLPH	$E_{\text{cm}} \leq 209 \text{ GeV}$
> 93.4	95	34	SCHAEEL	06B LEP	$E_{\text{cm}} \leq 209 \text{ GeV}$
		35	ACOSTA	05Q CDF	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + X$
> 85.0	95	36,37	ABBIENDI	04M OPAL	$E_{\text{cm}} \leq 209 \text{ GeV}$
		38	ABBIENDI	03G OPAL	$H_1^0 \rightarrow A^0 A^0$
> 86.5	95	36,39	ACHARD	02H L3	$E_{\text{cm}} \leq 209 \text{ GeV}, \tan\beta > 0.4$
		40	AKERROYD	02 RVUE	
> 90.1	95	36,41	HEISTER	02 ALEP	$E_{\text{cm}} \leq 209 \text{ GeV}, \tan\beta > 0.5$

<sup>1</sup> TUMASYAN 23S search for production of  $H_2^0/A^0 \rightarrow \tau^+\tau^-$  by gluon fusion and  $b$ -associated production using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 13 for excluded regions in the  $m_{A^0}$ - $\tan\beta$  plane in  $M_h^{125}$  and  $M_{h,EFT}^{125}$  MSSM scenarios. In both scenarios  $m_{A^0} < 350 \text{ GeV}$  is excluded at 95% CL.

NODE=S055HAD;LINKAGE=N

<sup>2</sup> AAD 20AA search for  $H_2^0/A^0 \rightarrow \tau^+\tau^-$  produced by gluon fusion or  $b$ -associated production using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 2(c) for excluded region in the  $M_h^{125}$  scenario of MSSM. Values of  $\tan\beta > 8$  (21) are excluded for  $m_{A^0} = 1.0$  (1.5) TeV at 95%CL.

NODE=S055HAD;LINKAGE=G

<sup>3</sup> AAD 24AP search for production of a heavy  $H_2^0$  and  $A^0$  decaying to  $t\bar{t}$  in  $140 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 13(b) for excluded parameter regions in hMSSM.

NODE=S055HAD;LINKAGE=P

<sup>4</sup> AAD 24CB use ATLAS measurements from up to  $139 \text{ fb}^{-1}$  of  $H$  production cross sections and decays to constrain the MSSM Higgs sector. See their Figs. 22 and 23 for excluded regions in various MSSM scenarios.

NODE=S055HAD;LINKAGE=T

<sup>5</sup> AAD 24H combine searches for a scalar resonance decaying to  $HH$  using up to  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  from AAD 22F, AAD 23Z, and AAD 22Y. See their Fig. 3 for the excluded region in the MSSM  $M_{h,EFT}^{125}$  and  $M_{h,EFT}^{125}(\tilde{\chi})$  benchmark scenarios (where only the resonant H-exchange contribution has been taken into account in the signal modelling).

NODE=S055HAD;LINKAGE=O

<sup>6</sup> AAD 20 combine measurements on  $H$  production and decay using data taken in years 2015–2017 (up to  $79.8 \text{ fb}^{-1}$ ) of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 19 for excluded region in the hMSSM parameter space.

NODE=S055HAD;LINKAGE=F

<sup>7</sup> AAD 20C combine searches for a scalar resonance decaying to  $HH$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  from AABOUD 19A, AABOUD 190, AABOUD 18CQ, AABOUD 19T, AABOUD 18CW, and AABOUD 18BU. See their Fig. 7(b) for the excluded region in the hMSSM parameter space.

NODE=S055HAD;LINKAGE=C

<sup>8</sup> AAD 20L search for  $b$ -associated production of  $H_2^0$  decaying to  $b\bar{b}$  in  $27.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for excluded regions in hMSSM,  $m_h^{\text{mod}+}$  and  $m_h^{\text{mod}-}$  scenarios of MSSM.

NODE=S055HAD;LINKAGE=L

<sup>9</sup> SIRUNYAN 20AC search for gluon-fusion and  $b$ -associated production of  $A^0$  decaying to  $ZH$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for excluded regions in the  $M_{h,EFT}^{125}$  and hMSSM scenarios of the MSSM.

NODE=S055HAD;LINKAGE=J

<sup>10</sup> SIRUNYAN 20AF search for  $H_2^0/A^0 \rightarrow t\bar{t}$  with one or two charged leptons in the final state using kinematic variables in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8 for excluded region in the hMSSM scenario of MSSM. Values of  $\tan\beta$  below 1.0–1.5 are excluded for  $m_{A^0} = 0.4$ –0.75 TeV at 95%CL.

NODE=S055HAD;LINKAGE=I

<sup>11</sup> SIRUNYAN 20Y search for gluon-fusion and vector-boson-fusion production of  $H_2^0$  decaying to  $W^+W^-$  in the final states  $\ell\nu\ell\nu$  and  $\ell\nu qq$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 8 and 9 for excluded regions in various MSSM scenarios.

NODE=S055HAD;LINKAGE=K

<sup>12</sup> SIRUNYAN 19CR search for production of  $H_2^0/A^0$  in gluon fusion and in association with a  $b\bar{b}$  pair, decaying to  $\mu^+\mu^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for the excluded region in the MSSM parameter space in the  $m_h^{\text{mod}+}$  and hMSSM scenarios.

NODE=S055HAD;LINKAGE=D

<sup>13</sup> AABOUD 18G search for production of  $H_2^0/A^0 \rightarrow \tau^+\tau^-$  by gluon fusion and  $b$ -associated production in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10 for excluded regions in the  $m_{A^0}$ - $\tan\beta$  plane in several MSSM scenarios.

NODE=S055HAD;LINKAGE=A

<sup>14</sup> SIRUNYAN 18A search for production of a scalar resonance decaying to  $H^0 H^0 \rightarrow b\bar{b}\tau^+\tau^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 (lower) for excluded regions in the  $m_{A^0}$ - $\tan\beta$  plane in the hMSSM scenario.

NODE=S055HAD;LINKAGE=V

- 15 SIRUNYAN 18BP search for production of  $H_2^0/A^0 \rightarrow b\bar{b}$  by  $b$ -associated production in  $35.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for the limits on cross section times branching ratio for  $m_{H_2^0}$ ,  $m_{A^0} = 0.3\text{--}1.3 \text{ TeV}$ , and Fig. 7 for excluded regions in the  $m_{A^0}\text{--}\tan(\beta)$  plane in several MSSM scenarios. NODE=S055HAD;LINKAGE=Y
- 16 SIRUNYAN 18CX search for production of  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  by gluon fusion and  $b$ -associated production in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for excluded regions in the  $m_{A^0}\text{--}\tan(\beta)$  plane in several MSSM scenarios. NODE=S055HAD;LINKAGE=B
- 17 AABOUD 16AA search for production of a Higgs boson in gluon fusion and in association with a  $b\bar{b}$  pair followed by the decay  $A^0 \rightarrow \tau^+\tau^-$  in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5(a, b) for limits on cross section times branching ratio for  $m_{A^0} = 200\text{--}1200 \text{ GeV}$ , and Fig. 5(c, d) for the excluded region in the MSSM parameter space in the  $m_h^{\text{mod+}}$  and hMSSM scenarios. NODE=S055HAD;LINKAGE=U
- 18 KHACHATRYAN 16A search for production of a Higgs boson in gluon fusion and in association with a  $b\bar{b}$  pair followed by the decay  $H_{1,2}^0/A^0 \rightarrow \mu^+\mu^-$  in  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 7 for the excluded region in the MSSM parameter space in the  $m_h^{\text{mod+}}$  benchmark scenario and Fig. 9 for limits on cross section times branching ratio. NODE=S055HAD;LINKAGE=Q
- 19 KHACHATRYAN 16P search for gluon fusion production of an  $H_2^0$  decaying to  $H^0 H^0 \rightarrow b\bar{b}\tau^+\tau^-$  and an  $A^0$  decaying to  $ZH^0 \rightarrow \ell^+\ell^-\tau^+\tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 12 for excluded region in the  $\tan\beta - \cos(\beta - \alpha)$  plane for  $m_{H_2^0} = m_{A^0} = 300 \text{ GeV}$ . NODE=S055HAD;LINKAGE=R
- 20 KHACHATRYAN 15AY search for production of a Higgs boson in association with a  $b$  quark in the decay  $H_{1,2}^0/A^0 \rightarrow b\bar{b}$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  and combine with CHATRCHYAN 13AG 7 TeV data. See their Fig. 6 for the limits on cross section times branching ratio for  $m_{A^0} = 100\text{--}900 \text{ GeV}$  and Figs. 7–9 for the excluded region in the MSSM parameter space in various benchmark scenarios. NODE=S055HAD;LINKAGE=S
- 21 AAD 14AW search for production of a Higgs boson followed by the decay  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  in  $19.5\text{--}20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 11 for the limits on cross section times branching ratio and their Figs. 9 and 10 for the excluded region in the MSSM parameter space. For  $m_{A^0} = 140 \text{ GeV}$ , the region  $\tan\beta > 5.4$  is excluded at 95% CL in the  $m_h^{\text{max}}$  scenario. NODE=S055HAD;LINKAGE=E
- 22 KHACHATRYAN 14M search for production of a Higgs boson in gluon fusion and in association with a  $b$  quark followed by the decay  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  in  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 7 and 8 for one- and two-dimensional limits on cross section times branching ratio and their Figs. 5 and 6 for the excluded region in the MSSM parameter space. For  $m_{A^0} = 140 \text{ GeV}$ , the region  $\tan\beta > 3.8$  is excluded at 95% CL in the  $m_h^{\text{max}}$  scenario. NODE=S055HAD;LINKAGE=H
- 23 AAD 13O search for production of a Higgs boson in the decay  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  and  $\mu^+\mu^-$  with  $4.7\text{--}4.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ . See their Fig. 6 for the excluded region in the MSSM parameter space and their Fig. 7 for the limits on cross section times branching ratio. For  $m_{A^0} = 110\text{--}170 \text{ GeV}$ ,  $\tan\beta \gtrsim 10$  is excluded, and for  $\tan\beta = 50$ ,  $m_{A^0}$  below  $470 \text{ GeV}$  is excluded at 95% CL in the  $m_h^{\text{max}}$  scenario. NODE=S055HAD;LINKAGE=GA
- 24 AAIJ 13T search for production of a Higgs boson in the forward region in the decay  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  in  $1.0 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ . See their Fig. 2 for the limits on cross section times branching ratio and the excluded region in the MSSM parameter space. NODE=S055HAD;LINKAGE=AI
- 25 CHATRCHYAN 13AG search for production of a Higgs boson in association with a  $b$  quark in the decay  $H_{1,2}^0/A^0 \rightarrow b\bar{b}$  in  $2.7\text{--}4.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ . See their Fig. 6 for the excluded region in the MSSM parameter space and Fig. 5 for the limits on cross section times branching ratio. For  $m_{A^0} = 90\text{--}350 \text{ GeV}$ , upper bounds on  $\tan\beta$  of  $18\text{--}42$  at 95% CL are obtained in the  $m_h^{\text{max}}$  scenario with  $\mu = +200 \text{ GeV}$ . NODE=S055HAD;LINKAGE=CR
- 26 AALTONEN 12AQ combine AALTONEN 12X and ABAZOV 11K. See their Table I and Fig. 1 for the limit on cross section times branching ratio and Fig. 2 for the excluded region in the MSSM parameter space. NODE=S055HAD;LINKAGE=OC
- 27 AALTONEN 12X search for associated production of a Higgs boson and a  $b$  quark in the decay  $H_{1,2}^0/A^0 \rightarrow b\bar{b}$ , with  $2.6 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Table III and Fig. 15 for the limit on cross section times branching ratio and Figs. 17, 18 for the excluded region in the MSSM parameter space. NODE=S055HAD;LINKAGE=TA
- 28 ABAZOV 12G search for production of a Higgs boson in the decay  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  with  $7.3 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$  and combine with ABAZOV 11W and ABAZOV 11K. See their Figs. 4, 5, and 6 for the excluded region in the MSSM parameter space. For  $m_{A^0} = 90\text{--}180 \text{ GeV}$ ,  $\tan\beta \gtrsim 30$  is excluded at 95% CL. in the  $m_h^{\text{max}}$  scenario. NODE=S055HAD;LINKAGE=VM

- 29 CHATRCHYAN 12K search for production of a Higgs boson in the decay  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$  with  $4.6 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ . See their Fig. 3 and Table 4 for the excluded region in the MSSM parameter space. For  $m_{A^0} = 160 \text{ GeV}$ , the region  $\tan\beta > 7.1$  is excluded at 95% CL in the  $m_h^{\text{max}}$  scenario. Superseded by KHACHATRYAN 14M. NODE=S055HAD;LINKAGE=CT
- 30 ABAZOV 11K search for associated production of a Higgs boson and a  $b$  quark, followed by the decay  $H_{1,2}^0/A^0 \rightarrow b\bar{b}$ , in  $5.2 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 5/Table 2 for the limit on cross section times branching ratio and Fig. 6 for the excluded region in the MSSM parameter space for  $\mu = -200 \text{ GeV}$ . NODE=S055HAD;LINKAGE=A2
- 31 ABAZOV 11W search for associated production of a Higgs boson and a  $b$  quark, followed by the decay  $H_{1,2}^0/A^0 \rightarrow \tau\tau$ , in  $7.3 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 2 for the limit on cross section times branching ratio and for the excluded region in the MSSM parameter space. NODE=S055HAD;LINKAGE=A1
- 32 AALTONEN 09AR search for Higgs bosons decaying to  $\tau^+\tau^-$  in two doublet models in  $1.8 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 2 for the limit on  $\sigma \cdot \text{B}(H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-)$  for different Higgs masses, and see their Fig. 3 for the excluded region in the MSSM parameter space. NODE=S055HAD;LINKAGE=TN
- 33 ABDALLAH 08B give limits in eight  $CP$ -conserving benchmark scenarios and some  $CP$ -violating scenarios. See paper for excluded regions for each scenario. Supersedes ABDALLAH 04. NODE=S055HAD;LINKAGE=AD
- 34 SCHAEEL 06B make a combined analysis of the LEP data. The quoted limit is for the  $m_h^{\text{max}}$  scenario with  $m_t = 174.3 \text{ GeV}$ . In the  $CP$ -violating CPX scenario no lower bound on  $m_{H_1^0}$  can be set at 95% CL. See paper for excluded regions in various scenarios. See Figs. 2–6 and Tabs. 14–21 for limits on  $\sigma(Z H^0) \cdot \text{B}(H^0 \rightarrow b\bar{b}, \tau^+\tau^-)$  and  $\sigma(H_1^0 H_2^0) \cdot \text{B}(H_1^0, H_2^0 \rightarrow b\bar{b}, \tau^+\tau^-)$ . NODE=S055HAD;LINKAGE=SH
- 35 ACOSTA 05Q search for  $H_{1,2}^0/A^0$  production in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.8 \text{ TeV}$  with  $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$ . At  $m_{A^0} = 100 \text{ GeV}$ , the obtained cross section upper limit is above theoretical expectation. NODE=S055HAD;LINKAGE=AC
- 36 Search for  $e^+e^- \rightarrow H_1^0 A^0$  in the final states  $b\bar{b}b\bar{b}$  and  $b\bar{b}\tau^+\tau^-$ , and  $e^+e^- \rightarrow H_1^0 Z$ . Universal scalar mass of 1 TeV, SU(2) gaugino mass of 200 GeV, and  $\mu = -200 \text{ GeV}$  are assumed, and two-loop radiative corrections incorporated. The limits hold for  $m_t = 175 \text{ GeV}$ , and for the  $m_h^{\text{max}}$  scenario. NODE=S055HAD;LINKAGE=HL
- 37 ABBIENDI 04M exclude  $0.7 < \tan\beta < 1.9$ , assuming  $m_t = 174.3 \text{ GeV}$ . Limits for other MSSM benchmark scenarios, as well as for  $CP$  violating cases, are also given. NODE=S055HAD;LINKAGE=HO
- 38 ABBIENDI 03G search for  $e^+e^- \rightarrow H_1^0 Z$  followed by  $H_1^0 \rightarrow A^0 A^0$ ,  $A^0 \rightarrow c\bar{c}, gg$ , or  $\tau^+\tau^-$ . In the no-mixing scenario, the region  $m_{H_1^0} = 45\text{--}85 \text{ GeV}$  and  $m_{A^0} = 2\text{--}9.5 \text{ GeV}$  is excluded at 95% CL. NODE=S055HAD;LINKAGE=AB
- 39 ACHARD 02H also search for the final state  $H_1^0 Z \rightarrow 2A^0 q\bar{q}$ ,  $A^0 \rightarrow q\bar{q}$ . In addition, the MSSM parameter set in the “large- $\mu$ ” and “no-mixing” scenarios are examined. NODE=S055HAD;LINKAGE=RH
- 40 AKEROYD 02 examine the possibility of a light  $A^0$  with  $\tan\beta < 1$ . Electroweak measurements are found to be inconsistent with such a scenario. NODE=S055HAD;LINKAGE=SY
- 41 HEISTER 02 excludes the range  $0.7 < \tan\beta < 2.3$ . A wider range is excluded with different stop mixing assumptions. Updates BARATE 01C. NODE=S055HAD;LINKAGE=HN

### Mass Limits for $H_1^0$ (Higgs Boson) in Supersymmetric Models

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>89.7		<sup>1</sup> ABDALLAH 08B	DLPH	$E_{\text{cm}} \leq 209 \text{ GeV}$
>92.8	95	<sup>2</sup> SCHAEEL 06B	LEP	$E_{\text{cm}} \leq 209 \text{ GeV}$
>84.5	95	<sup>3,4</sup> ABBIENDI 04M	OPAL	$E_{\text{cm}} \leq 209 \text{ GeV}$
>86.0	95	<sup>3,5</sup> ACHARD 02H	L3	$E_{\text{cm}} \leq 209 \text{ GeV}, \tan\beta > 0.4$
>89.8	95	<sup>3,6</sup> HEISTER 02	ALEP	$E_{\text{cm}} \leq 209 \text{ GeV}, \tan\beta > 0.5$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$$\begin{aligned} &^7 \text{ AALTONEN 12AQ TEVA } p\bar{p} \rightarrow H_{1,2}^0/A^0 + b + X, \\ &H_{1,2}^0/A^0 \rightarrow b\bar{b} \end{aligned}$$

<sup>1</sup> ABDALLAH 08B give limits in eight  $CP$ -conserving benchmark scenarios and some  $CP$ -violating scenarios. See paper for excluded regions for each scenario. Supersedes ABDALLAH 04. NODE=S055HSS;LINKAGE=AD

<sup>2</sup> SCHAEEL 06B make a combined analysis of the LEP data. The quoted limit is for the  $m_h^{\text{max}}$  scenario with  $m_t = 174.3 \text{ GeV}$ . In the  $CP$ -violating CPX scenario no lower bound on  $m_{H_1^0}$  can be set at 95% CL. See paper for excluded regions in various scenarios. See Figs. 2–6 and Tabs. 14–21 for limits on  $\sigma(Z H^0) \cdot \text{B}(H^0 \rightarrow b\bar{b}, \tau^+\tau^-)$  and  $\sigma(H_1^0 H_2^0) \cdot \text{B}(H_1^0, H_2^0 \rightarrow b\bar{b}, \tau^+\tau^-)$ . NODE=S055HSS;LINKAGE=SH

<sup>3</sup> Search for  $e^+e^- \rightarrow H_1^0 A^0$  in the final states  $b\bar{b}b\bar{b}$  and  $b\bar{b}\tau^+\tau^-$ , and  $e^+e^- \rightarrow H_1^0 Z$ . Universal scalar mass of 1 TeV, SU(2) gaugino mass of 200 GeV, and  $\mu = -200$  NODE=S055HSS;LINKAGE=HL

GeV are assumed, and two-loop radiative corrections incorporated. The limits hold for  $m_t=175$  GeV, and for the  $m_h^{\max}$  scenario.

- <sup>4</sup> ABBIENDI 04M exclude  $0.7 < \tan\beta < 1.9$ , assuming  $m_t = 174.3$  GeV. Limits for other MSSM benchmark scenarios, as well as for  $CP$  violating cases, are also given.
- <sup>5</sup> ACHARD 02H also search for the final state  $H_1^0 Z \rightarrow 2A^0 q\bar{q}$ ,  $A^0 \rightarrow q\bar{q}$ . In addition, the MSSM parameter set in the “large- $\mu$ ” and “no-mixing” scenarios are examined.
- <sup>6</sup> HEISTER 02 excludes the range  $0.7 < \tan\beta < 2.3$ . A wider range is excluded with different stop mixing assumptions. Updates BARATE 01C.
- <sup>7</sup> AALTONEN 12AQ combine AALTONEN 12X and ABAZOV 11K. See their Table I and Fig. 1 for the limit on cross section times branching ratio and Fig. 2 for the excluded region in the MSSM parameter space.

NODE=S055HSS;LINKAGE=HO

NODE=S055HSS;LINKAGE=RH

NODE=S055HSS;LINKAGE=HN

NODE=S055HSS;LINKAGE=OC

## MASS LIMITS FOR NEUTRAL HIGGS BOSONS IN EXTENDED HIGGS MODELS

NODE=S055245

NODE=S055245

This Section covers models which do not fit into either the Standard Model or its simplest minimal Supersymmetric extension (MSSM), leading to anomalous production rates, or nonstandard final states and branching ratios. In particular, this Section covers limits which may apply to generic two-Higgs-doublet models (2HDM), or to special regions of the MSSM parameter space where decays to invisible particles or to photon pairs are dominant (see the review on “Status of Higgs Boson Physics”). Concerning the mass limits for  $H^0$  and  $A^0$  listed below, see the footnotes or the comment lines for details on the nature of the models to which the limits apply.

The observed signal at about 125 GeV, see section “H”, can be interpreted as one of the neutral Higgs bosons of an extended Higgs sector.

### Mass Limits in General two-Higgs-doublet Models

NODE=S055H2D  
NODE=S055H2D

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1	AAD	24AB ATLS	$A^0 \rightarrow ZH_2^0$	
2	AAD	24AP ATLS	$H_2^0, A^0 \rightarrow t\bar{t}$	
3	AAD	24CB ATLS	$H$ properties	
4	AAD	24H ATLS	$H_2^0 \rightarrow HH$	
5	HAYRAPETY...24H	CMS	flavor changing $H_2^0, A^0$	
6	AAD	23AD ATLS	$A^0 \rightarrow ZH_2^0, H_2^0 \rightarrow HH$	
7	AAD	23BG ATLS	$t\bar{t}H_2^0/A^0$	
8	AAD	23O ATLS	$A^0 \rightarrow ZH$	
9	AAD	21AF ATLS	$H_2^0 \rightarrow ZZ$	
10	AAD	21AI ATLS	$A^0 \rightarrow ZH_2^0$	
11	AAD	20 ATLS	$H^0$ properties	
12	AAD	20L ATLS	$H_2^0 \rightarrow b\bar{b}$	
13	SIRUNYAN	20AA CMS	$H_2^0 \rightarrow ZA^0$ or $A^0 \rightarrow ZH_2^0$	
14	SIRUNYAN	20Y CMS	$H_2^0 \rightarrow W^+W^-$	
15	SIRUNYAN	19AE CMS	$A^0 \rightarrow \tau^+\tau^-$	
16	SIRUNYAN	19AV CMS	$A^0 \rightarrow ZH^0$	
17	AABOUD	18AH ATLS	$A^0 \rightarrow ZH_2^0$	
18	AABOUD	18AI ATLS	$A^0 \rightarrow ZH^0$	
19	AABOUD	18BF ATLS	$H_2^0 \rightarrow ZZ$	
20	AABOUD	18CE ATLS	$pp \rightarrow H_2^0/A^0 t\bar{t},$ $H_2^0/A^0 \rightarrow t\bar{t}$	
21	HALLER	18 RVUE	global fits	
22	SIRUNYAN	18BP CMS	$pp \rightarrow H_2^0/A^0 + b + X,$ $H_2^0/A^0 \rightarrow b\bar{b}$	
23	SIRUNYAN	18ED CMS	$A^0 \rightarrow ZH^0$	
24	AABOUD	17AN ATLS	$H_2^0, A^0 \rightarrow t\bar{t}$	
25	SIRUNYAN	17AX CMS	$A^0 b\bar{b}, A^0 \rightarrow \mu^+\mu^-$	
26	AAD	16AX ATLS	$H_2^0 \rightarrow ZZ$	
27	KHACHATRY...16P	CMS	$H_2^0 \rightarrow H^0 H^0, A^0 \rightarrow ZH^0$	
28	KHACHATRY...16W	CMS	$A^0 b\bar{b}, A^0 \rightarrow \tau^+\tau^-$	
29	KHACHATRY...16Z	CMS	$H_2^0 \rightarrow ZA^0$ or $A^0 \rightarrow ZH_2^0$	
30	AAD	15BK ATLS	$H_2^0 \rightarrow H^0 H^0$	

		31 AAD	15S ATLS	$A^0 \rightarrow ZH^0$	
		32 KHACHATRY...15BB	CMS	$H_2^0, A^0 \rightarrow \gamma\gamma$	
		33 KHACHATRY...15N	CMS	$A^0 \rightarrow ZH^0$	
		34 AAD	14M ATLS	$H_2^0 \rightarrow H^\pm W^\mp \rightarrow$ $H^0 W^\pm W^\mp, H^0 \rightarrow b\bar{b}$	
		35 KHACHATRY...14Q	CMS	$H_2^0 \rightarrow H^0 H^0, A^0 \rightarrow ZH^0$	
		36 AALTONEN	09AR CDF	$p\bar{p} \rightarrow H_{1,2}^0/A^0 + X,$ $H_{1,2}^0/A^0 \rightarrow \tau^+\tau^-$	
none 1–55	95	37 ABBIENDI	05A OPAL	$H_1^0$ , Type II model	
>110.6	95	38 ABDALLAH	05D DLPH	$H_1^0 \rightarrow 2 \text{ jets}$	
		39 ABDALLAH	04O DLPH	$Z \rightarrow f\bar{f}H$	
		40 ABDALLAH	04O DLPH	$e^+e^- \rightarrow H^0 Z, H^0 A^0$	OCCUR=2
		41 ABBIENDI	02D OPAL	$e^+e^- \rightarrow b\bar{b}H$	
none 1–44	95	42 ABBIENDI	01E OPAL	$H_1^0$ , Type-II model	
> 68.0	95	43 ABBIENDI	99E OPAL	$\tan\beta > 1$	
		44 ABREU	95H DLPH	$Z \rightarrow H^0 Z^*, H^0 A^0$	
		45 PICH	92 RVUE	Very light Higgs	
<p><sup>1</sup> AAD 24AB search for gluon-fusion and <math>b\bar{b}</math> associated production of <math>A^0</math> decaying to <math>A^0 \rightarrow ZH_2^0</math>, in the final states <math>\ell^+\ell^- t\bar{t}</math> and <math>\nu\bar{\nu} b\bar{b}</math> using 140 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 5 and 6 for limits on cross section times branching ratios in terms of two-Higgs-doublet models, and Fig. 7 for excluded model parameter space.</p>					NODE=S055H2D;LINKAGE=JA
<p><sup>2</sup> AAD 24AP search for production of a heavy <math>H_2^0</math> and <math>A^0</math> decaying to <math>t\bar{t}</math> in 140 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Fig. 13(a) for excluded parameter regions in a Type II two-Higgs-doublet model.</p>					NODE=S055H2D;LINKAGE=KA
<p><sup>3</sup> AAD 24CB use ATLAS measurements from up to 139 fb<sup>-1</sup> of <math>H</math> production cross sections and decays to constrain two Higgs doublet models. See their Figs. 19–21 for excluded regions of 2HDM parameter space.</p>					NODE=S055H2D;LINKAGE=LA
<p><sup>4</sup> AAD 24H combine searches for a scalar resonance decaying to <math>HH</math> using up to 139 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV from AAD 22F, AAD 23Z, and AAD 22Y. See their Fig. 2 for the excluded region in the Type-I two-Higgs-doublet model parameter space (where only the resonant H-exchange contribution has been taken into account in the signal modelling).</p>					NODE=S055H2D;LINKAGE=HA
<p><sup>5</sup> HAYRAPETYAN 24H search for <math>H_2^0</math> and <math>A^0</math> having flavor-violating couplings to <math>tc</math> or <math>tu</math>, produced in association with top quark(s), using 138 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. For the interpretation of the data in terms of flavor-violating two-Higgs-doublet models, given in the range <math>m_{A^0} = 0.2\text{--}1.0</math> TeV, see their Figs. 4, 5, and 6.</p>					NODE=S055H2D;LINKAGE=IA
<p><sup>6</sup> AAD 23AD search for associated production of <math>W/ZH_2^0</math> and gluon fusion production of <math>A^0</math> decaying to <math>ZH_2^0</math>, with the decay chain <math>H_2^0 \rightarrow HH \rightarrow b\bar{b}b\bar{b}</math>, using 139 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 12 and 13 for excluded regions in Type-I and lepton-specific 2HDMs.</p>					NODE=S055H2D;LINKAGE=FA
<p><sup>7</sup> AAD 23BG search for production of <math>H_2^0/A^0</math> in association with a <math>t\bar{t}</math> pair, decaying to <math>t\bar{t}</math>, using 139 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Fig. 8 for excluded regions in the parameter space of the type II 2HDM.</p>					NODE=S055H2D;LINKAGE=GA
<p><sup>8</sup> AAD 23O search for production of an <math>A^0</math> in gluon-gluon fusion and in association with a <math>b\bar{b}</math>, decaying to <math>ZH</math> in the final states <math>\nu\bar{\nu} b\bar{b}</math> and <math>\ell^+\ell^- b\bar{b}</math> using 139 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 12 and 13 for excluded regions in the parameter space in various 2HDMs.</p>					NODE=S055H2D;LINKAGE=EA
<p><sup>9</sup> AAD 21AF search for production of a heavy <math>H_2^0</math> state decaying to <math>ZZ</math> in the final states <math>\ell^+\ell^-\ell'^+\ell'^-</math> and <math>\ell^+\ell^-\nu\bar{\nu}</math> in 139 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 6 and 7 for excluded parameter regions of the 2HDM Type I and II.</p>					NODE=S055H2D;LINKAGE=BA
<p><sup>10</sup> AAD 21AI search for production of an <math>A^0</math> in gluon-gluon fusion and in association with a <math>b\bar{b}</math>, decaying to <math>ZH_2^0 \rightarrow \ell^+\ell^- b\bar{b}</math> or <math>\ell^+\ell^- W^+W^-</math> in 139 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 10 and 14 for excluded regions in the parameter space of various 2HDMs.</p>					NODE=S055H2D;LINKAGE=CA
<p><sup>11</sup> AAD 20 combine measurements on <math>H^0</math> production and decay using data taken in years 2015–2017 (up to 79.8 fb<sup>-1</sup>) of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Fig. 18 for excluded regions in various 2HDMs.</p>					NODE=S055H2D;LINKAGE=X
<p><sup>12</sup> AAD 20L search for <math>b</math>-associated production of <math>H_2^0</math> decaying to <math>b\bar{b}</math> in 27.8 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 10 and 11 for excluded regions in the flipped two Higgs doublet model.</p>					NODE=S055H2D;LINKAGE=AA
<p><sup>13</sup> SIRUNYAN 20AA search for <math>H_2^0 \rightarrow ZA^0, A^0 \rightarrow b\bar{b}</math> or <math>A^0 \rightarrow ZH_2^0, H_2^0 \rightarrow b\bar{b}</math> in 35.9 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Figs. 8 and 9 for excluded regions in the parameter space of Type-II two Higgs doublet model.</p>					NODE=S055H2D;LINKAGE=Y
<p><sup>14</sup> SIRUNYAN 20Y search for gluon-fusion and vector-boson-fusion production of <math>H_2^0</math> decaying to <math>W^+W^-</math> in the final states <math>\ell\nu\ell\nu</math> and <math>\ell\nu qq</math> in 35.9 fb<sup>-1</sup> of <math>pp</math> collisions at <math>E_{\text{cm}} = 13</math> TeV. See their Fig. 7 for excluded regions in Type I and II two Higgs doublet models.</p>					NODE=S055H2D;LINKAGE=Z

- 15 SIRUNYAN 19AE search for a pseudoscalar resonance produced in association with a  $b\bar{b}$  pair, decaying to  $\tau^+\tau^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4 for cross section limits for  $m_{A^0} = 25\text{--}70 \text{ GeV}$  and comparison with some representative 2HDMs. NODE=S055H2D;LINKAGE=V
- 16 SIRUNYAN 19AV search for a scalar resonance produced by gluon fusion or  $b$  associated production, decaying to  $ZH^0 \rightarrow \ell^+\ell^-b\bar{b}$  ( $\ell = e, \mu$ ) or  $\nu\bar{\nu}b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 6 and 7 for excluded regions in the parameter space of various 2HDMs. NODE=S055H2D;LINKAGE=W
- 17 AABOUD 18AH search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH^0_2 \rightarrow \ell^+\ell^-b\bar{b}$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for excluded regions in the parameter space of various 2HDMs. NODE=S055H2D;LINKAGE=P
- 18 AABOUD 18AI search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH^0$  in the final states  $\nu\bar{\nu}b\bar{b}$  and  $\ell^+\ell^-b\bar{b}$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 7 and 8 for excluded regions in the parameter space in various 2HDMs. NODE=S055H2D;LINKAGE=O
- 19 AABOUD 18BF search for production of a heavy  $H^0_2$  state decaying to  $ZZ$  in the final states  $\ell^+\ell^-\ell^+\ell^-$  and  $\ell^+\ell^-\nu\bar{\nu}$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 8 and 9 for excluded parameter regions in 2HDM Type I and II. NODE=S055H2D;LINKAGE=R
- 20 AABOUD 18CE search for the process  $pp \rightarrow H^0_2/A^0 t\bar{t}$  followed by the decay  $H^0_2/A^0 \rightarrow t\bar{t}$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 12 for limits on cross section times branching ratio, and for lower limits on  $\tan\beta$  for  $m_{H^0_2}, m_{A^0} = 0.4\text{--}1.0 \text{ TeV}$  in the 2HDM type II. NODE=S055H2D;LINKAGE=U
- 21 HALLER 18 perform global fits in the framework of two-Higgs-doublet models (type I, II, lepton specific, flipped). See their Fig. 8 for allowed parameter regions from fits to LHC  $H^0$  measurements, Fig. 9 bottom and charm decays, Fig. 10 muon anomalous magnetic moment, Fig. 11 electroweak precision data, and Fig. 12 by combination of all data. NODE=S055H2D;LINKAGE=S
- 22 SIRUNYAN 18BP search for production of  $H^0_2/A^0 \rightarrow b\bar{b}$  by  $b$ -associated production in  $35.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for the limits on cross section times branching ratio for  $m_{H^0_2}, m_{A^0} = 0.3\text{--}1.3 \text{ TeV}$ , and Figs. 8 and 9 for excluded regions in the parameter space of type-II and flipped 2HDMs. NODE=S055H2D;LINKAGE=Q
- 23 SIRUNYAN 18ED search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH^0$  in the final states  $\nu\bar{\nu}b\bar{b}$  or  $\ell^+\ell^-b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for excluded regions in the parameter space in Type I and II 2HDMs. NODE=S055H2D;LINKAGE=T
- 24 AABOUD 17AN search for production of a heavy  $H^0_2$  and/or  $A^0$  decaying to  $t\bar{t}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 3 and Table III for excluded parameter regions in Type II Two-Higgs-Doublet-Models. NODE=S055H2D;LINKAGE=M
- 25 SIRUNYAN 17AX search for  $A^0 b\bar{b}$  production followed by the decay  $A^0 \rightarrow \mu^+\mu^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . Limits are set in the range  $m_{A^0} = 25\text{--}60 \text{ GeV}$ . See their Fig. 5 for upper limits on  $\sigma(A^0 b\bar{b}) \cdot \text{B}(A^0 \rightarrow \mu^+\mu^-)$ . NODE=S055H2D;LINKAGE=N
- 26 AAD 16AX search for production of a heavy  $H^0$  state decaying to  $ZZ$  in the final states  $\ell^+\ell^-\ell^+\ell^-$ ,  $\ell^+\ell^-\nu\bar{\nu}$ ,  $\ell^+\ell^-q\bar{q}$ , and  $\nu\bar{\nu}q\bar{q}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 13 and 14 for excluded parameter regions in Type I and II models. NODE=S055H2D;LINKAGE=I
- 27 KHACHATRYAN 16P search for gluon fusion production of an  $H^0_2$  decaying to  $H^0 H^0 \rightarrow b\bar{b}\tau^+\tau^-$  and an  $A^0$  decaying to  $ZH^0 \rightarrow \ell^+\ell^-\tau^+\tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 11 for limits on  $\tan\beta$  for  $m_{A^0} = 230\text{--}350 \text{ GeV}$ . NODE=S055H2D;LINKAGE=K
- 28 KHACHATRYAN 16W search for  $A^0 b\bar{b}$  production followed by the decay  $A^0 \rightarrow \tau^+\tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 3 for upper limits on  $\sigma(A^0 b\bar{b}) \cdot \text{B}(A^0 \rightarrow \tau^+\tau^-)$ . NODE=S055H2D;LINKAGE=J
- 29 KHACHATRYAN 16Z search for  $H^0_2 \rightarrow ZA^0$  followed by  $A^0 \rightarrow b\bar{b}$  or  $\tau^+\tau^-$ , and  $A^0 \rightarrow ZH^0_2$  followed by  $H^0_2 \rightarrow b\bar{b}$  or  $\tau^+\tau^-$ , in  $19.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 4 for cross section limits and Fig. 5 for excluded region in the parameter space. NODE=S055H2D;LINKAGE=L
- 30 AAD 15BK search for production of a heavy  $H^0_2$  decaying to  $H^0 H^0$  in the final state  $b\bar{b}b\bar{b}$  in  $19.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 15–18 for excluded regions in the parameter space. NODE=S055H2D;LINKAGE=H
- 31 AAD 15S search for production of  $A^0$  decaying to  $ZH^0 \rightarrow \ell^+\ell^-b\bar{b}$ ,  $\nu\bar{\nu}b\bar{b}$  and  $\ell^+\ell^-\tau^+\tau^-$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 4 and 5 for excluded regions in the parameter space. NODE=S055H2D;LINKAGE=D
- 32 KHACHATRYAN 15BB search for  $H^0_2, A^0 \rightarrow \gamma\gamma$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 10 for excluded regions in the two-Higgs-doublet model parameter space. NODE=S055H2D;LINKAGE=E
- 33 KHACHATRYAN 15N search for production of  $A^0$  decaying to  $ZH^0 \rightarrow \ell^+\ell^-b\bar{b}$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 5 for excluded regions in the  $\tan\beta - \cos(\beta - \alpha)$  plane for  $m_{A^0} = 300 \text{ GeV}$ . NODE=S055H2D;LINKAGE=C
- 34 AAD 14M search for the decay cascade  $H^0_2 \rightarrow H^\pm W^\mp \rightarrow H^0 W^\pm W^\mp$ ,  $H^0$  decaying to  $b\bar{b}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Table IV for limits in a two-Higgs-doublet model for  $m_{H^0_2} = 325\text{--}1025 \text{ GeV}$  and  $m_{H^\pm} = 225\text{--}825 \text{ GeV}$ . NODE=S055H2D;LINKAGE=A



- 35 KHACHATRYAN 14Q search for  $H_2^0 \rightarrow H^0 H^0$  and  $A^0 \rightarrow Z H^0$  in  $19.5 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 4 and 5 for limits on cross section times branching ratio for  $m_{H_2, A^0} = 260\text{--}360 \text{ GeV}$  and their Figs. 7–9 for limits in two-Higgs-doublet models.
- 36 AALTONEN 09AR search for Higgs bosons decaying to  $\tau^+ \tau^-$  in two doublet models in  $1.8 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 2 for the limit on  $\sigma \cdot \text{B}(H_{1,2}^0/A^0 \rightarrow \tau^+ \tau^-)$  for different Higgs masses, and see their Fig. 3 for the excluded region in the MSSM parameter space.
- 37 ABBIENDI 05A search for  $e^+ e^- \rightarrow H_1^0 A^0$  in general Type-II two-doublet models, with decays  $H_1^0, A^0 \rightarrow q\bar{q}, g\bar{g}, \tau^+ \tau^-$ , and  $H_1^0 \rightarrow A^0 A^0$ .
- 38 ABDALLAH 05D search for  $e^+ e^- \rightarrow H^0 Z$  and  $H^0 A^0$  with  $H^0, A^0$  decaying to two jets of any flavor including  $g\bar{g}$ . The limit is for SM  $H^0 Z$  production cross section with  $\text{B}(H^0 \rightarrow jj) = 1$ .
- 39 ABDALLAH 04o search for  $Z \rightarrow b\bar{b}H^0, b\bar{b}A^0, \tau^+ \tau^- H^0$  and  $\tau^+ \tau^- A^0$  in the final states  $4b, b\bar{b}\tau^+ \tau^-$ , and  $4\tau$ . See paper for limits on Yukawa couplings.
- 40 ABDALLAH 04o search for  $e^+ e^- \rightarrow H^0 Z$  and  $H^0 A^0$ , with  $H^0, A^0$  decaying to  $b\bar{b}, \tau^+ \tau^-$ , or  $H^0 \rightarrow A^0 A^0$  at  $E_{\text{cm}} = 189\text{--}208 \text{ GeV}$ . See paper for limits on couplings.
- 41 ABBIENDI 02D search for  $Z \rightarrow b\bar{b}H_1^0$  and  $b\bar{b}A^0$  with  $H_1^0/A^0 \rightarrow \tau^+ \tau^-$ , in the range  $4 < m_H < 12 \text{ GeV}$ . See their Fig. 8 for limits on the Yukawa coupling.
- 42 ABBIENDI 01E search for neutral Higgs bosons in general Type-II two-doublet models, at  $E_{\text{cm}} \leq 189 \text{ GeV}$ . In addition to usual final states, the decays  $H_1^0, A^0 \rightarrow q\bar{q}, g\bar{g}$  are searched for. See their Figs. 15,16 for excluded regions.
- 43 ABBIENDI 99E search for  $e^+ e^- \rightarrow H^0 A^0$  and  $H^0 Z$  at  $E_{\text{cm}} = 183 \text{ GeV}$ . The limit is with  $m_H = m_A$  in general two Higgs-doublet models. See their Fig. 18 for the exclusion limit in the  $m_H\text{--}m_A$  plane. Updates the results of ACKERSTAFF 98S.
- 44 See Fig. 4 of ABREU 95H for the excluded region in the  $m_{H^0} - m_{A^0}$  plane for general two-doublet models. For  $\tan\beta > 1$ , the region  $m_{H^0} + m_{A^0} \lesssim 87 \text{ GeV}$ ,  $m_{H^0} < 47 \text{ GeV}$  is excluded at 95% CL.
- 45 PICH 92 analyse  $H^0$  with  $m_{H^0} < 2m_\mu$  in general two-doublet models. Excluded regions in the space of mass-mixing angles from LEP, beam dump, and  $\pi^\pm, \eta$  rare decays are shown in Figs. 3,4. The considered mass region is not totally excluded.

NODE=S055H2D;LINKAGE=B

NODE=S055H2D;LINKAGE=TN

NODE=S055H2D;LINKAGE=AN

NODE=S055H2D;LINKAGE=AH

NODE=S055H2D;LINKAGE=AO

NODE=S055H2D;LINKAGE=AP

NODE=S055H2D;LINKAGE=DD

NODE=S055H2D;LINKAGE=EK

NODE=S055H2D;LINKAGE=EB

NODE=S055H2D;LINKAGE=G

NODE=S055H2D;LINKAGE=F

### Mass Limits for $H^0$ with Vanishing Yukawa Couplings

These limits assume that  $H^0$  couples to gauge bosons with the same strength as the Standard Model Higgs boson, but has no coupling to quarks and leptons (this is often referred to as “fermiophobic”).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		1 AALTONEN	13K CDF	$H^0 \rightarrow W W^{(*)}$
none 100–113	95	2 AALTONEN	13L CDF	$H^0 \rightarrow \gamma\gamma, W W^*, Z Z^*$
none 100–116	95	3 AALTONEN	13M TEVA	$H^0 \rightarrow \gamma\gamma, W W^*, Z Z^*$
		4 ABAZOV	13G D0	$H^0 \rightarrow W W^{(*)}$
none 100–113	95	5 ABAZOV	13H D0	$H^0 \rightarrow \gamma\gamma$
		6 ABAZOV	13I D0	$H^0 \rightarrow W W^{(*)}$
		7 ABAZOV	13J D0	$H^0 \rightarrow W W^{(*)}, Z Z^{(*)}$
none 100–114	95	8 ABAZOV	13L D0	$H^0 \rightarrow \gamma\gamma, W W^*, Z Z^*$
none 110–147	95	9 CHATRCHYAN	13AL CMS	$H^0 \rightarrow \gamma\gamma$
none 110–118, 119.5–121	95	10 AAD	12N ATLS	$H^0 \rightarrow \gamma\gamma$
none 100–114	95	11 AALTONEN	12AN CDF	$H^0 \rightarrow \gamma\gamma$
none 110–194	95	12 CHATRCHYAN	12AO CMS	$H^0 \rightarrow \gamma\gamma, W W^{(*)}, Z Z^{(*)}$
none 70–106	95	13 AALTONEN	09AB CDF	$H^0 \rightarrow \gamma\gamma$
none 70–100	95	14 ABAZOV	08U D0	$H^0 \rightarrow \gamma\gamma$
>105.8	95	15 SCHAEEL	07 ALEP	$e^+ e^- \rightarrow H^0 Z, H^0 \rightarrow W W^*$
>104.1	95	16,17 ABDALLAH	04L DLPH	$e^+ e^- \rightarrow H^0 Z, H^0 \rightarrow \gamma\gamma$
>107	95	18 ACHARD	03C L3	$H^0 \rightarrow W W^*, Z Z^*, \gamma\gamma$
>105.5	95	16,19 ABBIENDI	02F OPAL	$H^0 \rightarrow \gamma\gamma$
>105.4	95	20 ACHARD	02C L3	$H^0 \rightarrow \gamma\gamma$
none 60–82	95	21 AFFOLDER	01H CDF	$p\bar{p} \rightarrow H^0 W/Z, H^0 \rightarrow \gamma\gamma$
> 94.9	95	22 ACCIARRI	00S L3	$e^+ e^- \rightarrow H^0 Z, H^0 \rightarrow \gamma\gamma$
>100.7	95	23 BARATE	00L ALEP	$e^+ e^- \rightarrow H^0 Z, H^0 \rightarrow \gamma\gamma$
> 96.2	95	24 ABBIENDI	99O OPAL	$e^+ e^- \rightarrow H^0 Z, H^0 \rightarrow \gamma\gamma$
> 78.5	95	25 ABBOTT	99B D0	$p\bar{p} \rightarrow H^0 W/Z, H^0 \rightarrow \gamma\gamma$
		26 ABREU	99P DLPH	$e^+ e^- \rightarrow H^0 \gamma \text{ and/or } H^0 \rightarrow \gamma\gamma$

NODE=S055H2F

NODE=S055H2F

NODE=S055H2F

- 1 AALTONEN 13K search for  $H^0 \rightarrow WW^{(*)}$  in  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . A limit on cross section times branching ratio which corresponds to (1.3–6.6) times the expected cross section is given in the range  $m_{H^0} = 110\text{--}200 \text{ GeV}$  at 95% CL. NODE=S055H2F;LINKAGE=TT
- 2 AALTONEN 13L combine all CDF searches with  $9.45\text{--}10.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . NODE=S055H2F;LINKAGE=EE
- 3 AALTONEN 13M combine all Tevatron data from the CDF and D0 Collaborations of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . NODE=S055H2F;LINKAGE=LL
- 4 ABAZOV 13G search for  $H^0 \rightarrow WW^{(*)}$  in  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . A limit on cross section times branching ratio which corresponds to (2–9) times the expected cross section is given for  $m_{H^0} = 100\text{--}200 \text{ GeV}$  at 95% CL. NODE=S055H2F;LINKAGE=ZZ
- 5 ABAZOV 13H search for  $H^0 \rightarrow \gamma\gamma$  in  $9.6 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . NODE=S055H2F;LINKAGE=OO
- 6 ABAZOV 13I search for  $H^0$  production in the final state with one lepton and two or more jets plus missing  $E_T$  in  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The search is sensitive to  $WH^0$ ,  $ZH^0$  and vector-boson fusion Higgs production with  $H^0 \rightarrow WW^{(*)}$ . A limit on cross section times branching ratio which corresponds to (8–30) times the expected cross section is given in the range  $m_{H^0} = 100\text{--}200 \text{ GeV}$  at 95% CL. NODE=S055H2F;LINKAGE=MM
- 7 ABAZOV 13J search for  $H^0$  production in the final states  $e\bar{e}\mu$ ,  $e\mu\mu$ ,  $\mu\tau\tau$ , and  $e^\pm\mu^\pm$  in  $8.6\text{--}9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . The search is sensitive to  $WH^0$ ,  $ZH^0$  production with  $H^0 \rightarrow WW^{(*)}$ ,  $ZZ^{(*)}$ , decaying to leptonic final states. A limit on cross section times branching ratio which corresponds to (2.4–13.0) times the expected cross section is given in the range  $m_{H^0} = 100\text{--}200 \text{ GeV}$  at 95% CL. NODE=S055H2F;LINKAGE=D0
- 8 ABAZOV 13L combine all D0 results with up to  $9.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . NODE=S055H2F;LINKAGE=ZC
- 9 CHATRCHYAN 13AL search for  $H^0 \rightarrow \gamma\gamma$  in  $5.1 \text{ fb}^{-1}$  and  $5.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  and  $8 \text{ TeV}$ . NODE=S055H2F;LINKAGE=CY
- 10 AAD 12N search for  $H^0 \rightarrow \gamma\gamma$  with  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  in the mass range  $m_{H^0} = 110\text{--}150 \text{ GeV}$ . NODE=S055H2F;LINKAGE=AA
- 11 AALTONEN 12AN search for  $H^0 \rightarrow \gamma\gamma$  with  $10 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$  in the mass range  $m_{H^0} = 100\text{--}150 \text{ GeV}$ . NODE=S055H2F;LINKAGE=FL
- 12 CHATRCHYAN 12AO use data from CHATRCHYAN 12G, CHATRCHYAN 12E, CHATRCHYAN 12H, CHATRCHYAN 12I, CHATRCHYAN 12D, and CHATRCHYAN 12C. NODE=S055H2F;LINKAGE=CH
- 13 AALTONEN 09AB search for  $H^0 \rightarrow \gamma\gamma$  in  $3.0 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$  in the mass range  $m_{H^0} = 70\text{--}150 \text{ GeV}$ . Associated  $H^0W$ ,  $H^0Z$  production and  $WW$ ,  $ZZ$  fusion are considered. NODE=S055H2F;LINKAGE=TO
- 14 ABAZOV 08U search for  $H^0 \rightarrow \gamma\gamma$  in  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$  in the mass range  $m_{H^0} = 70\text{--}150 \text{ GeV}$ . Associated  $H^0W$ ,  $H^0Z$  production and  $WW$ ,  $ZZ$  fusion are considered. See their Tab. 1 for the limit on  $\sigma \cdot \text{B}(H^0 \rightarrow \gamma\gamma)$ , and see their Fig. 3 for the excluded region in the  $m_{H^0} - \text{B}(H^0 \rightarrow \gamma\gamma)$  plane. NODE=S055H2F;LINKAGE=BA
- 15 SCHAEEL 07 search for Higgs bosons in association with a fermion pair and decaying to  $WW^*$ . The limit is from this search and HEISTER 02L for a  $H^0$  with SM production cross section. NODE=S055H2F;LINKAGE=SA
- 16 Search for associated production of a  $\gamma\gamma$  resonance with a  $Z$  boson, followed by  $Z \rightarrow q\bar{q}$ ,  $\ell^+\ell^-$ , or  $\nu\bar{\nu}$ , at  $E_{\text{cm}} \leq 209 \text{ GeV}$ . The limit is for a  $H^0$  with SM production cross section. NODE=S055H2F;LINKAGE=HA
- 17 Updates ABREU 01F. NODE=S055H2F;LINKAGE=HD
- 18 ACHARD 03C search for  $e^+e^- \rightarrow ZH^0$  followed by  $H^0 \rightarrow WW^*$  or  $ZZ^*$  at  $E_{\text{cm}} = 200\text{--}209 \text{ GeV}$  and combine with the ACHARD 02C result. The limit is for a  $H^0$  with SM production cross section. For  $\text{B}(H^0 \rightarrow WW^*) + \text{B}(H^0 \rightarrow ZZ^*) = 1$ ,  $m_{H^0} > 108.1 \text{ GeV}$  is obtained. See fig. 6 for the limits under different BR assumptions. NODE=S055H2F;LINKAGE=AC
- 19 For  $\text{B}(H^0 \rightarrow \gamma\gamma)=1$ ,  $m_{H^0} > 117 \text{ GeV}$  is obtained. NODE=S055H2F;LINKAGE=LA
- 20 ACHARD 02C search for associated production of a  $\gamma\gamma$  resonance with a  $Z$  boson, followed by  $Z \rightarrow q\bar{q}$ ,  $\ell^+\ell^-$ , or  $\nu\bar{\nu}$ , at  $E_{\text{cm}} \leq 209 \text{ GeV}$ . The limit is for a  $H^0$  with SM production cross section. For  $\text{B}(H^0 \rightarrow \gamma\gamma)=1$ ,  $m_{H^0} > 114 \text{ GeV}$  is obtained. NODE=S055H2F;LINKAGE=HR
- 21 AFFOLDER 01H search for associated production of a  $\gamma\gamma$  resonance and a  $W$  or  $Z$  (tagged by two jets, an isolated lepton, or missing  $E_T$ ). The limit assumes Standard Model values for the production cross section and for the couplings of the  $H^0$  to  $W$  and  $Z$  bosons. See their Fig. 11 for limits with  $\text{B}(H^0 \rightarrow \gamma\gamma) < 1$ . NODE=S055H2F;LINKAGE=AF
- 22 ACCIARRI 00S search for associated production of a  $\gamma\gamma$  resonance with a  $q\bar{q}$ ,  $\nu\bar{\nu}$ , or  $\ell^+\ell^-$  pair in  $e^+e^-$  collisions at  $E_{\text{cm}} = 189 \text{ GeV}$ . The limit is for a  $H^0$  with SM production cross section. For  $\text{B}(H^0 \rightarrow \gamma\gamma)=1$ ,  $m_{H^0} > 98 \text{ GeV}$  is obtained. See their Fig. 5 for limits on  $\text{B}(H \rightarrow \gamma\gamma) \cdot \sigma(e^+e^- \rightarrow Hf\bar{f})/\sigma(e^+e^- \rightarrow Hf\bar{f})$  (SM). NODE=S055H2F;LINKAGE=PC
- 23 BARATE 00L search for associated production of a  $\gamma\gamma$  resonance with a  $q\bar{q}$ ,  $\nu\bar{\nu}$ , or  $\ell^+\ell^-$  pair in  $e^+e^-$  collisions at  $E_{\text{cm}} = 88\text{--}202 \text{ GeV}$ . The limit is for a  $H^0$  with SM production cross section. For  $\text{B}(H^0 \rightarrow \gamma\gamma)=1$ ,  $m_{H^0} > 109 \text{ GeV}$  is obtained. See their Fig. 3 for limits on  $\text{B}(H \rightarrow \gamma\gamma) \cdot \sigma(e^+e^- \rightarrow Hf\bar{f})/\sigma(e^+e^- \rightarrow Hf\bar{f})$  (SM). NODE=S055H2F;LINKAGE=PB

- <sup>24</sup> ABBIENDI 990 search for associated production of a  $\gamma\gamma$  resonance with a  $q\bar{q}$ ,  $\nu\bar{\nu}$ , or  $\ell^+\ell^-$  pair in  $e^+e^-$  collisions at 189 GeV. The limit is for a  $H^0$  with SM production cross section. See their Fig. 4 for limits on  $\sigma(e^+e^- \rightarrow H^0 Z^0) \times B(H^0 \rightarrow \gamma\gamma) \times B(X^0 \rightarrow f\bar{f})$  for various masses. Updates the results of ACKERSTAFF 98Y.
- <sup>25</sup> ABBOTT 99B search for associated production of a  $\gamma\gamma$  resonance and a dijet pair. The limit assumes Standard Model values for the production cross section and for the couplings of the  $H^0$  to  $W$  and  $Z$  bosons. Limits in the range of  $\sigma(H^0 + Z/W) \cdot B(H^0 \rightarrow \gamma\gamma) = 0.80\text{--}0.34$  pb are obtained in the mass range  $m_{H^0} = 65\text{--}150$  GeV.
- <sup>26</sup> ABREU 99P search for  $e^+e^- \rightarrow H^0\gamma$  with  $H^0 \rightarrow b\bar{b}$  or  $\gamma\gamma$ , and  $e^+e^- \rightarrow H^0 q\bar{q}$  with  $H^0 \rightarrow \gamma\gamma$ . See their Fig. 4 for limits on  $\sigma \times B$ . Explicit limits within an effective interaction framework are also given.

NODE=S055H2F;LINKAGE=DI

NODE=S055H2F;LINKAGE=3C

NODE=S055H2F;LINKAGE=PA

### Mass Limits for $H^0$ Decaying to Invisible Final States

These limits are for a neutral scalar  $H^0$  which predominantly decays to invisible final states. Standard Model values are assumed for the couplings of  $H^0$  to ordinary particles unless otherwise stated.

NODE=S055H2I

NODE=S055H2I

NODE=S055H2I

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		1 AABOUD	19AI ATLS	$W W/Z Z$ fusion
		2 AAD	15BD ATLS	$pp \rightarrow H^0 W X, H^0 Z X$
		3 AAD	15BH ATLS	jet + missing $E_T$
		4 AAD	14BA ATLS	secondary vertex
		5 AAD	14O ATLS	$pp \rightarrow H^0 Z X$
		6 CHATRCHYAN	14B CMS	$pp \rightarrow H^0 Z X, qqH^0 X$
		7 AAD	13AG ATLS	secondary vertex
		8 AAD	13AT ATLS	electron jets
		9 CHATRCHYAN	13BJ CMS	
		10 AAD	12AQ ATLS	secondary vertex
		11 AALTONEN	12AB CDF	secondary vertex
		12 AALTONEN	12U CDF	secondary vertex
>108.2	95	13 ABBIENDI	10 OPAL	
		14 ABBIENDI	07 OPAL	large width
>112.3	95	15 ACHARD	05 L3	
>112.1	95	15 ABDALLAH	04B DLPH	
>114.1	95	15 HEISTER	02 ALEP	$E_{\text{cm}} \leq 209$ GeV
>106.4	95	15 BARATE	01C ALEP	$E_{\text{cm}} \leq 202$ GeV
> 89.2	95	16 ACCIARRI	00M L3	

- <sup>1</sup> AABOUD 19AI search for  $H^0_{1,2}$  production by vector boson fusion and decay to invisible final states in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 6(b) for limits on cross section times branching ratios for  $m_{H^0_{1,2}} = 0.1\text{--}3$  TeV.

NODE=S055H2I;LINKAGE=F

- <sup>2</sup> AAD 15BD search for  $pp \rightarrow H^0 W X$  and  $pp \rightarrow H^0 Z X$  with  $W$  or  $Z$  decaying hadronically and  $H^0$  decaying to invisible final states in  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. See their Fig. 6 for a limit on the cross section times branching ratio for  $m_{H^0} = 115\text{--}300$  GeV.

NODE=S055H2I;LINKAGE=D

- <sup>3</sup> AAD 15BH search for events with a jet and missing  $E_T$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. Limits on  $\sigma(H^0) B(H^0 \rightarrow \text{invisible}) < (44\text{--}10) \text{ pb}$  (95%CL) is given for  $m_{H^0} = 115\text{--}300$  GeV.

NODE=S055H2I;LINKAGE=E

- <sup>4</sup> AAD 14BA search for  $H^0$  production in the decay mode  $H^0 \rightarrow X^0 X^0$ , where  $X^0$  is a long-lived particle which decays to collimated pairs of  $e^+e^-$ ,  $\mu^+\mu^-$ , or  $\pi^+\pi^-$  plus invisible particles, in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Figs. 15 and 16 for limits on cross section times branching ratio.

NODE=S055H2I;LINKAGE=C

- <sup>5</sup> AAD 14O search for  $pp \rightarrow H^0 Z X$ ,  $Z \rightarrow \ell\ell$ , with  $H^0$  decaying to invisible final states in  $4.5 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 7$  TeV and  $20.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV. See their Fig. 3 for a limit on the cross section times branching ratio for  $m_{H^0} = 110\text{--}400$  GeV.

NODE=S055H2I;LINKAGE=A

- <sup>6</sup> CHATRCHYAN 14B search for  $pp \rightarrow H^0 Z X$ ,  $Z \rightarrow \ell\ell$  and  $Z \rightarrow b\bar{b}$ , and also  $pp \rightarrow qqH^0 X$  with  $H^0$  decaying to invisible final states using data at  $E_{\text{cm}} = 7$  and 8 TeV. See their Figs. 10, 11 for limits on the cross section times branching ratio for  $m_{H^0} = 100\text{--}400$  GeV.

NODE=S055H2I;LINKAGE=B

- <sup>7</sup> AAD 13AG search for  $H^0$  production in the decay mode  $H^0 \rightarrow X^0 X^0$ , where  $X^0$  is a long-lived particle which decays to  $\mu^+\mu^- X^0$ , in  $1.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV. See their Fig. 7 for limits on cross section times branching ratio.

NODE=S055H2I;LINKAGE=AA

- <sup>8</sup> AAD 13AT search for  $H^0$  production in the decay  $H^0 \rightarrow X^0 X^0$ , where  $X^0$  eventually decays to clusters of collimated  $e^+e^-$  pairs, in  $2.04 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV. See their Fig. 3 for limits on cross section times branching ratio.

NODE=S055H2I;LINKAGE=DA

- <sup>9</sup> CHATRCHYAN 13BJ search for  $H^0$  production in the decay chain  $H^0 \rightarrow X^0 X^0$ ,  $X^0 \rightarrow \mu^+\mu^- X^0$  in  $5.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV. See their Fig. 2 for limits on cross section times branching ratio.

NODE=S055H2I;LINKAGE=AT

- <sup>10</sup> AAD 12AQ search for  $H^0$  production in the decay mode  $H^0 \rightarrow X^0 X^0$ , where  $X^0$  is a long-lived particle which decays mainly to  $b\bar{b}$  in the muon detector, in  $1.94 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ . See their Fig. 3 for limits on cross section times branching ratio for  $m_{H^0} = 120, 140 \text{ GeV}$ ,  $m_{X^0} = 20, 40 \text{ GeV}$  in the  $c\tau$  range of 0.5–35 m.
- <sup>11</sup> AALTONEN 12AB search for  $H^0$  production in the decay mode  $H^0 \rightarrow X^0 X^0$ , where  $X^0$  eventually decays to clusters of collimated  $\ell^+ \ell^-$  pairs, in  $5.1 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . Cross section limits are provided for a benchmark MSSM model incorporating the parameters given in Table VI.
- <sup>12</sup> AALTONEN 12U search for  $H^0$  production in the decay mode  $H^0 \rightarrow X^0 X^0$ , where  $X^0$  is a long-lived particle with  $c\tau \approx 1 \text{ cm}$  which decays mainly to  $b\bar{b}$ , in  $3.2 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Figs. 9 and 10 for limits on cross section times branching ratio for  $m_{H^0} = (130\text{--}170) \text{ GeV}$ ,  $m_{X^0} = 20, 40 \text{ GeV}$ .
- <sup>13</sup> ABBIENDI 10 search for  $e^+ e^- \rightarrow H^0 Z$  with  $H^0$  decaying invisibly. The limit assumes SM production cross section and  $B(H^0 \rightarrow \text{invisible}) = 1$ .
- <sup>14</sup> ABBIENDI 07 search for  $e^+ e^- \rightarrow H^0 Z$  with  $Z \rightarrow q\bar{q}$  and  $H^0$  decaying to invisible final states. The  $H^0$  width is varied between 1 GeV and 3 TeV. A limit  $\sigma \cdot B(H^0 \rightarrow \text{invisible}) < (0.07\text{--}0.57) \text{ pb}$  (95%CL) is obtained at  $E_{\text{cm}} = 206 \text{ GeV}$  for  $m_{H^0} = 60\text{--}114 \text{ GeV}$ .
- <sup>15</sup> Search for  $e^+ e^- \rightarrow H^0 Z$  with  $H^0$  decaying invisibly. The limit assumes SM production cross section and  $B(H^0 \rightarrow \text{invisible}) = 1$ .
- <sup>16</sup> ACCIARRI 00M search for  $e^+ e^- \rightarrow ZH^0$  with  $H^0$  decaying invisibly at  $E_{\text{cm}} = 183\text{--}189 \text{ GeV}$ . The limit assumes SM production cross section and  $B(H^0 \rightarrow \text{invisible}) = 1$ . See their Fig. 6 for limits for smaller branching ratios.

NODE=S055H2I;LINKAGE=AD

NODE=S055H2I;LINKAGE=AN

NODE=S055H2I;LINKAGE=AL

NODE=S055H2I;LINKAGE=BB

NODE=S055H2I;LINKAGE=BI

NODE=S055H2I;LINKAGE=HM

NODE=S055H2I;LINKAGE=PD

### Mass Limits for Light $A^0$

These limits are for a pseudoscalar  $A^0$  in the mass range below  $\mathcal{O}(10) \text{ GeV}$ .

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1	ADACHI	23A BEL2	$\tau \rightarrow eA^0, \tau \rightarrow \mu A^0$
2	TUMASYAN	23AR CMS	$H \rightarrow A^0 A^0 \rightarrow 4\gamma$
3	ABLIKIM	22H BES3	$J/\psi \rightarrow A^0 \gamma$
4	JIA	22 BELL	$\Upsilon(1S) \rightarrow A^0 \gamma$
5	AAD	20AE ATLS	$H^0 \rightarrow Z A^0$
6	AABOUD	18AP ATLS	$H^0 \rightarrow A^0 A^0$
7	KHACHATRY...	17AZ CMS	$H^0 \rightarrow A^0 A^0$
8	ABLIKIM	16E BES3	$J/\psi \rightarrow A^0 \gamma$
9	KHACHATRY...	16F CMS	$H^0 \rightarrow A^0 A^0$
10	LEES	15H BABR	$\Upsilon(1S) \rightarrow A^0 \gamma$
11	LEES	13C BABR	$\Upsilon(1S) \rightarrow A^0 \gamma$
12	LEES	13L BABR	$\Upsilon(1S) \rightarrow A^0 \gamma$
13	LEES	13R BABR	$\Upsilon(1S) \rightarrow A^0 \gamma$
14	ABLIKIM	12 BES3	$J/\psi \rightarrow A^0 \gamma$
15	CHATRCHYAN	12V CMS	$A^0 \rightarrow \mu^+ \mu^-$
16	AALTONEN	11P CDF	$t \rightarrow bH^+, H^+ \rightarrow W^+ A^0$
17,18	ABOUZAID	11A KTEV	$K_L \rightarrow \pi^0 \pi^0 A^0, A^0 \rightarrow \mu^+ \mu^-$
19	DEL-AMO-SA...	11J BABR	$\Upsilon(1S) \rightarrow A^0 \gamma$
20	LEES	11H BABR	$\Upsilon(2S, 3S) \rightarrow A^0 \gamma$
21	ANDREAS	10 RVUE	
18,22	HYUN	10 BELL	$B^0 \rightarrow K^{*0} A^0, A^0 \rightarrow \mu^+ \mu^-$
18,23	HYUN	10 BELL	$B^0 \rightarrow \rho^0 A^0, A^0 \rightarrow \mu^+ \mu^-$
24	AUBERT	09P BABR	$\Upsilon(3S) \rightarrow A^0 \gamma$
25	AUBERT	09Z BABR	$\Upsilon(2S) \rightarrow A^0 \gamma$
26	AUBERT	09Z BABR	$\Upsilon(3S) \rightarrow A^0 \gamma$
18,27	TUNG	09 K391	$K_L \rightarrow \pi^0 \pi^0 A^0, A^0 \rightarrow \gamma \gamma$
28	LOVE	08 CLEO	$\Upsilon(1S) \rightarrow A^0 \gamma$
29	BESSON	07 CLEO	$\Upsilon(1S) \rightarrow \eta_b \gamma$
30	PARK	05 HYCP	$\Sigma^+ \rightarrow p A^0, A^0 \rightarrow \mu^+ \mu^-$
31	BALEST	95 CLE2	$\Upsilon(1S) \rightarrow A^0 \gamma$
32	ANTREASYAN	90C CBAL	$\Upsilon(1S) \rightarrow A^0 \gamma$

NODE=S055H2A

NODE=S055H2A

NODE=S055H2A

OCCUR=2

OCCUR=2

- <sup>1</sup> ADACHI 23A search for flavor-changing  $\tau$  decays  $\tau \rightarrow eA^0$  and  $\tau \rightarrow \mu A^0$ , with  $A^0$  invisible, using  $62.8 \text{ fb}^{-1}$  of  $e^+ e^-$  collisions at  $E_{\text{cm}} = 10.58 \text{ GeV}$ . Limits on  $B(\tau \rightarrow eA^0)/B(\tau \rightarrow e\nu\nu)$  in the range  $1.1 \times 10^{-3}\text{--}9.7 \times 10^{-3}$  (95% CL) and  $B(\tau \rightarrow \mu A^0)/B(\tau \rightarrow \mu\nu\nu)$  in the range  $0.7 \times 10^{-3}\text{--}12.2 \times 10^{-3}$  (95% CL) are given for  $m_{A^0} = 0\text{--}1.6 \text{ GeV}$ . See their Fig. 2.
- <sup>2</sup> TUMASYAN 23AR search for the decay  $H \rightarrow A^0 A^0$  with  $A^0 \rightarrow \gamma\gamma$  (detected as a merged photonlike object) using  $136 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . Limits

NODE=S055H2A;LINKAGE=Q

NODE=S055H2A;LINKAGE=R

- on  $B(H \rightarrow A^0 A^0) \cdot B^2(A^0 \rightarrow \gamma\gamma)$  in the range  $0.9 \times 10^{-3} - 3.3 \times 10^{-3}$  (95% CL) are given for  $m_{A^0} = 0.1 - 1.2$  GeV. See their Fig. 2.
- <sup>3</sup> ABLIKIM 22H search for the process  $J/\psi \rightarrow A^0 \gamma$  with  $A^0$  decaying to  $\mu^+ \mu^-$  in  $9 \times 10^9$   $J/\psi$  events and give limits on  $B(J/\psi \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $1.2 \times 10^{-9} - 7.78 \times 10^{-7}$  (90% CL) for  $0.212 \text{ GeV} \leq m_{A^0} \leq 3.0 \text{ GeV}$ . See their Fig. 4.
- <sup>4</sup> JIA 22 search for the process  $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^- \rightarrow A^0 \gamma \pi^+ \pi^-$  with  $A^0$  decaying to  $\tau^+ \tau^-$  or  $\mu^+ \mu^-$  in  $158 \times 10^6$   $\Upsilon(2S)$  events and give limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \tau^+ \tau^-)$  in the range  $3.8 \times 10^{-6} - 1.5 \times 10^{-4}$  (90% CL) for  $m_{A^0} = 3.6 - 9.2$  GeV, and  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $3.1 \times 10^{-7} - 1.6 \times 10^{-5}$  (90% CL) for  $m_{A^0} = 0.21 - 9.2$  GeV. See their Fig. 4.
- <sup>5</sup> AAD 20AE search for the decay  $H^0 \rightarrow Z A^0$ ,  $Z \rightarrow \ell^+ \ell^-$ ,  $A^0$  decaying hadronically ( $A^0 \rightarrow g g$  or  $s \bar{s}$ ), in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. Limit on the product of production cross section and the  $H^0 \rightarrow Z A^0$  branching ratio in the range 17–340 pb (95% CL) is given for  $m_{A^0} = 0.5 - 4.0$  GeV, see their Table I.
- <sup>6</sup> AABOUD 18AP search for the decay  $H^0 \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- \mu^+ \mu^-$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 10(b) for limits on  $B(H^0 \rightarrow A^0 A^0)$  in the range  $m_{A^0} = 1 - 2.5, 4.5 - 8$  GeV, assuming a type-II two-doublet plus singlet model with  $\tan(\beta) = 5$ .
- <sup>7</sup> KHACHATRYAN 17AZ search for the decay  $H^0 \rightarrow A^0 A^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$ ,  $\mu^+ \mu^- b \bar{b}$ , and  $\mu^+ \mu^- \tau^+ \tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Figs. 4, 5, and 6 for cross section limits in the range  $m_{A^0} = 5 - 62.5$  GeV. See also their Figs. 7, 8, and 9 for interpretation of the data in terms of models with two Higgs doublets and a singlet.
- <sup>8</sup> ABLIKIM 16E search for the process  $J/\psi \rightarrow A^0 \gamma$  with  $A^0$  decaying to  $\mu^+ \mu^-$  and give limits on  $B(J/\psi \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $2.8 \times 10^{-8} - 5.0 \times 10^{-6}$  (90% CL) for  $0.212 \leq m_{A^0} \leq 3.0$  GeV. See their Fig. 5.
- <sup>9</sup> KHACHATRYAN 16F search for the decay  $H^0 \rightarrow A^0 A^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 8 for cross section limits for  $m_{A^0} = 4 - 8$  GeV.
- <sup>10</sup> LEES 15H search for the process  $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^- \rightarrow A^0 \gamma \pi^+ \pi^-$  with  $A^0$  decaying to  $c \bar{c}$  and give limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow c \bar{c})$  in the range  $7.4 \times 10^{-5} - 2.4 \times 10^{-3}$  (90% CL) for  $4.00 \leq m_{A^0} \leq 8.95$  and  $9.10 \leq m_{A^0} \leq 9.25$  GeV. See their Fig. 6.
- <sup>11</sup> LEES 13C search for the process  $\Upsilon(2S, 3S) \rightarrow \Upsilon(1S) \pi^+ \pi^- \rightarrow A^0 \gamma \pi^+ \pi^-$  with  $A^0$  decaying to  $\mu^+ \mu^-$  and give limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $(0.3 - 9.7) \times 10^{-6}$  (90% CL) for  $0.212 \leq m_{A^0} \leq 9.20$  GeV. See their Fig. 5(e) for limits on the  $b - A^0$  Yukawa coupling derived by combining this result with AUBERT 09Z.
- <sup>12</sup> LEES 13L search for the process  $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^- \rightarrow A^0 \gamma \pi^+ \pi^-$  with  $A^0$  decaying to  $g g$  or  $s \bar{s}$  and give limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow g g)$  between  $1 \times 10^{-6}$  and  $2 \times 10^{-2}$  (90% CL) for  $0.5 \leq m_{A^0} \leq 9.0$  GeV, and  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow s \bar{s})$  between  $4 \times 10^{-6}$  and  $1 \times 10^{-3}$  (90% CL) for  $1.5 \leq m_{A^0} \leq 9.0$  GeV. See their Fig. 4.
- <sup>13</sup> LEES 13R search for the process  $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^- \rightarrow A^0 \gamma \pi^+ \pi^-$  with  $A^0$  decaying to  $\tau^+ \tau^-$  and give limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \tau^+ \tau^-)$  in the range  $0.9 - 13 \times 10^{-5}$  (90% CL) for  $3.6 \leq m_{A^0} \leq 9.2$  GeV. See their Fig. 4 for limits on the  $b - A^0$  Yukawa coupling derived by combining this result with AUBERT 09P.
- <sup>14</sup> ABLIKIM 12 searches for the process  $\psi(3686) \rightarrow \pi \pi J/\psi$ ,  $J/\psi \rightarrow A^0 \gamma$  with  $A^0$  decaying to  $\mu^+ \mu^-$ . It gives mass dependent limits on  $B(J/\psi \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $4 \times 10^{-7} - 2.1 \times 10^{-5}$  (90% C.L.) for  $0.212 \leq m_{A^0} \leq 3.0$  GeV. See their Fig. 2.
- <sup>15</sup> CHATRCHYAN 12V search for  $A^0$  production in the decay  $A^0 \rightarrow \mu^+ \mu^-$  with  $1.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV. A limit on  $\sigma(A^0) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range (1.5–7.5) pb is given for  $m_{A^0} = (5.5 - 8.7)$  and (11.5–14) GeV at 95% CL.
- <sup>16</sup> AALTONEN 11P search in  $2.7 \text{ fb}^{-1}$  of  $p \bar{p}$  collisions at  $E_{\text{cm}} = 1.96$  TeV for the decay chain  $t \rightarrow b H^+$ ,  $H^+ \rightarrow W^+ A^0$ ,  $A^0 \rightarrow \tau^+ \tau^-$  with  $m_{A^0}$  between 4 and 9 GeV. See their Fig. 4 for limits on  $B(t \rightarrow b H^+)$  for  $90 < m_{H^+} < 160$  GeV.
- <sup>17</sup> ABOUZAID 11A search for the decay chain  $K_L \rightarrow \pi^0 \pi^0 A^0$ ,  $A^0 \rightarrow \mu^+ \mu^-$  and give a limit  $B(K_L \rightarrow \pi^0 \pi^0 A^0) \cdot B(A^0 \rightarrow \mu^+ \mu^-) < 1.0 \times 10^{-10}$  at 90% CL for  $m_{A^0} = 214.3$  MeV.
- <sup>18</sup> The search was motivated by PARK 05.
- <sup>19</sup> DEL-AMO-SANCHEZ 11J search for the process  $\Upsilon(2S) \rightarrow \Upsilon(1S) \pi^+ \pi^- \rightarrow A^0 \gamma \pi^+ \pi^-$  with  $A^0$  decaying to invisible final states. They give limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \text{invisible})$  in the range  $(1.9 - 4.5) \times 10^{-6}$  (90% CL) for  $0 \leq m_{A^0} \leq 8.0$  GeV, and  $(2.7 - 37) \times 10^{-6}$  for  $8.0 \leq m_{A^0} \leq 9.2$  GeV.

NODE=S055H2A;LINKAGE=O

NODE=S055H2A;LINKAGE=P

NODE=S055H2A;LINKAGE=N

NODE=S055H2A;LINKAGE=M

NODE=S055H2A;LINKAGE=K

NODE=S055H2A;LINKAGE=I

NODE=S055H2A;LINKAGE=H

NODE=S055H2A;LINKAGE=B

NODE=S055H2A;LINKAGE=LE

NODE=S055H2A;LINKAGE=E

NODE=S055H2A;LINKAGE=F

NODE=S055H2A;LINKAGE=J

NODE=S055H2A;LINKAGE=CA

NODE=S055H2A;LINKAGE=A5

NODE=S055H2A;LINKAGE=AB

NODE=S055H2A;LINKAGE=PA

NODE=S055H2A;LINKAGE=D1

- 20 LEES 11H search for the process  $\Upsilon(2S, 3S) \rightarrow A^0 \gamma$  with  $A^0$  decaying hadronically and give limits on  $B(\Upsilon(2S, 3S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \text{hadrons})$  in the range  $1 \times 10^{-6} - 8 \times 10^{-5}$  (90% CL) for  $0.3 < m_{A^0} < 7$  GeV. The decay rates for  $\Upsilon(2S)$  and  $\Upsilon(3S)$  are assumed to be equal up to the phase space factor. See their Fig. 5.
- 21 ANDREAS 10 analyze constraints from rare decays and other processes on a light  $A^0$  with  $m_{A^0} < 2m_\mu$  and give limits on its coupling to fermions at the level of  $10^{-4}$  times the Standard Model value.
- 22 HYUN 10 search for the decay chain  $B^0 \rightarrow K^{*0} A^0$ ,  $A^0 \rightarrow \mu^+ \mu^-$  and give a limit on  $B(B^0 \rightarrow K^{*0} A^0) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $(2.26-5.53) \times 10^{-8}$  at 90%CL for  $m_{A^0} = 212-300$  MeV. The limit for  $m_{A^0} = 214.3$  MeV is  $2.26 \times 10^{-8}$ .
- 23 HYUN 10 search for the decay chain  $B^0 \rightarrow \rho^0 A^0$ ,  $A^0 \rightarrow \mu^+ \mu^-$  and give a limit on  $B(B^0 \rightarrow \rho^0 A^0) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $(1.73-4.51) \times 10^{-8}$  at 90%CL for  $m_{A^0} = 212-300$  MeV. The limit for  $m_{A^0} = 214.3$  MeV is  $1.73 \times 10^{-8}$ .
- 24 AUBERT 09P search for the process  $\Upsilon(3S) \rightarrow A^0 \gamma$  with  $A^0 \rightarrow \tau^+ \tau^-$  for  $4.03 < m_{A^0} < 9.52$  and  $9.61 < m_{A^0} < 10.10$  GeV, and give limits on  $B(\Upsilon(3S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \tau^+ \tau^-)$  in the range  $(1.5-16) \times 10^{-5}$  (90% CL).
- 25 AUBERT 09Z search for the process  $\Upsilon(2S) \rightarrow A^0 \gamma$  with  $A^0 \rightarrow \mu^+ \mu^-$  for  $0.212 < m_{A^0} < 9.3$  GeV and give limits on  $B(\Upsilon(2S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $(0.3-8) \times 10^{-6}$  (90% CL).
- 26 AUBERT 09Z search for the process  $\Upsilon(3S) \rightarrow A^0 \gamma$  with  $A^0 \rightarrow \mu^+ \mu^-$  for  $0.212 < m_{A^0} < 9.3$  GeV and give limits on  $B(\Upsilon(3S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \mu^+ \mu^-)$  in the range  $(0.3-5) \times 10^{-6}$  (90% CL).
- 27 TUNG 09 search for the decay chain  $K_L \rightarrow \pi^0 \pi^0 A^0$ ,  $A^0 \rightarrow \gamma \gamma$  and give a limit on  $B(K_L \rightarrow \pi^0 \pi^0 A^0) \cdot B(A^0 \rightarrow \gamma \gamma)$  in the range  $(2.4-10.7) \times 10^{-7}$  at 90%CL for  $m_{A^0} = 194.3-219.3$  MeV. The limit for  $m_{A^0} = 214.3$  MeV is  $2.4 \times 10^{-7}$ .
- 28 LOVE 08 search for the process  $\Upsilon(1S) \rightarrow A^0 \gamma$  with  $A^0 \rightarrow \mu^+ \mu^-$  (for  $m_{A^0} < 2m_\tau$ ) and  $A^0 \rightarrow \tau^+ \tau^-$ . Limits on  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot B(A^0 \rightarrow \ell^+ \ell^-)$  in the range  $10^{-6}-10^{-4}$  (90% CL) are given.
- 29 BESSON 07 give a limit  $B(\Upsilon(1S) \rightarrow \eta_b \gamma) \cdot B(\eta_b \rightarrow \tau^+ \tau^-) < 0.27\%$  (95% CL), which constrains a possible  $A^0$  exchange contribution to the  $\eta_b$  decay.
- 30 PARK 05 found three candidate events for  $\Sigma^+ \rightarrow p \mu^+ \mu^-$  in the HyperCP experiment. Due to a narrow spread in dimuon mass, they hypothesize the events as a possible signal of a new boson. It can be interpreted as a neutral particle with  $m_{A^0} = 214.3 \pm 0.5$  MeV and the branching fraction  $B(\Sigma^+ \rightarrow p A^0) \cdot B(A^0 \rightarrow \mu^+ \mu^-) = (3.1_{-1.9}^{+2.4} \pm 1.5) \times 10^{-8}$ .
- 31 BALEST 95 give limits  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot 1.5 \times 10^{-5}$  at 90% CL for  $m_{A^0} < 5$  GeV. The limit becomes  $< 10^{-4}$  for  $m_{A^0} < 7.7$  GeV.
- 32 ANTREASYAN 90C give limits  $B(\Upsilon(1S) \rightarrow A^0 \gamma) \cdot 5.6 \times 10^{-5}$  at 90% CL for  $m_{A^0} < 7.2$  GeV.  $A^0$  is assumed not to decay in the detector.

NODE=S055H2A;LINKAGE=L1

NODE=S055H2A;LINKAGE=AN

NODE=S055H2A;LINKAGE=HY

NODE=S055H2A;LINKAGE=HU

NODE=S055H2A;LINKAGE=BR

NODE=S055H2A;LINKAGE=UB

NODE=S055H2A;LINKAGE=AU

NODE=S055H2A;LINKAGE=NG

NODE=S055H2A;LINKAGE=LO

NODE=S055H2A;LINKAGE=BE

NODE=S055H2A;LINKAGE=H5

NODE=S055H2A;LINKAGE=D

NODE=S055H2A;LINKAGE=G

### Other Mass Limits

We use a symbol  $H_1^0$  if mass  $< 125$  GeV or  $H_2^0$  if mass  $> 125$  GeV. The notation  $H$  is reserved for the 125 GeV particle.

NODE=S055H2O

NODE=S055H2O

NODE=S055H2O

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

1	HAYRAPETY...25A	CMS	$H_1^0 \rightarrow \gamma \gamma$
2	AAD	24A ATLS	$H_2^0 \rightarrow Z \gamma$
3	AAD	24AP ATLS	$H_2^0, A^0 \rightarrow t \bar{t}$
4	AAD	24BQ ATLS	$H \rightarrow A^0 A^0$
5	AAD	24BV ATLS	$H_2^0 \rightarrow H H$
6	AAD	24BW ATLS	$H_2^0 \rightarrow H_2^0 H$
7	AAD	24BX ATLS	$H_2^0 \rightarrow H_2^0 H_2^0$
8	AAD	24BX ATLS	$A^0 \rightarrow H_2^0 Z$
9	AAD	24CA ATLS	$H_3^0 \rightarrow H_{1,2}^0 H$
10	AAD	24CD ATLS	$H_2^0 \rightarrow Z A$
11	AAD	24CF ATLS	$A^0 \rightarrow \tau^+ \tau^-$
12	AAD	24CI ATLS	$H_2^0, A^0 \rightarrow \text{invisible}$
13	AAD	24CK ATLS	two doublet + pseudoscalar + Dirac DM
14	AAD	24H ATLS	$H_2^0 \rightarrow H H$
15	AAD	24K ATLS	$H \rightarrow Z A^0$
16	ADACHI	24F BEL2	$H_1^0 \rightarrow \mu^+ \mu^-$
17	HAYRAPETY...24AA	CMS	$H \rightarrow A^0 A^0$

OCCUR=2

18	HAYRAPETY...24AE	CMS	$H_2^0 \rightarrow HH$	
19	HAYRAPETY...24AF	CMS	$H \rightarrow A^0 A^0$	
20	HAYRAPETY...24AI	CMS	$H \rightarrow H_1^0 H_1^0$ , long-lived	
21	HAYRAPETY...24AJ	CMS	$H_2^0 \rightarrow \gamma\gamma$	
22	HAYRAPETY...24AZ	CMS	$A^0 A^0, A^0 \rightarrow \mu^+ \mu^-$	
23	HAYRAPETY...24AZ	CMS	$H, H_{1,2}^0 \rightarrow A^0 A^0, A^0 \rightarrow \mu^+ \mu^-$	OCCUR=2
24	HAYRAPETY...24I	CMS	$H \rightarrow Z A^0$	
25	TUMASYAN 24A	CMS	$H_{1,2}^0 \rightarrow e^+ e^-, \mu^+ \mu^-, \tau^+ \tau^-$	
26	TUMASYAN 24B	CMS	$H_2^0 \rightarrow HH$	
27	TUMASYAN 24B	CMS	$H_3^0 \rightarrow H_{1,2}^0 H$	OCCUR=2
28	AAD 23AD	ATLS	$H_2^0 \rightarrow HH$	
29	AAD 23AD	ATLS	$A^0 \rightarrow Z H_2^0 \rightarrow Z H H$	OCCUR=2
30	AAD 23AJ	ATLS	$H^\pm \rightarrow W^\pm A^0$	
31	AAD 23BD	ATLS	$t \rightarrow q H_{1,2}^0$	
32	AAD 23BE	ATLS	$H_2^0 \rightarrow W^+ W^-$	
33	AAD 23BG	ATLS	$t \bar{t} H_2^0 / A^0$	
34	AAD 23BW	ATLS	$A^0 t \bar{t}, A^0 \rightarrow \mu^+ \mu^-$	
35	AAD 23BX	ATLS	$H + \text{invisible } A^0$	
36	AAD 23CA	ATLS	$H_3^0 \rightarrow H_2^0 H$	
37	AAD 23CR	ATLS	flavor changing $H_2^0$	
38	AAD 23O	ATLS	$A^0 \rightarrow Z H$	
39	AAD 23R	ATLS	$A^0 \rightarrow \gamma\gamma$	
40	AAD 23U	ATLS	$H_2^0 \rightarrow Z \gamma$	
41	AAD 23Z	ATLS	$H_2^0 \rightarrow HH$	
42	HAYRAPETY...23C	CMS	$H_{1,2}^0 \rightarrow e \mu$	
43	HAYRAPETY...23G	CMS	$A^0 \rightarrow \mu^+ \mu^-$	
44	TUMASYAN 23	CMS	$H_3^0 \rightarrow H_{1,2}^0 H$	
45	TUMASYAN 23M	CMS	$H \rightarrow A^0 A^0$	
46	TUMASYAN 23O	CMS	$H_2^0 \rightarrow HH$	
47	TUMASYAN 23S	CMS	$H_{1,2}^0 \rightarrow \tau^+ \tau^-$	
48	AAD 22A	ATLS	$H \rightarrow A^0 A^0$	
49	AAD 22D	ATLS	$Z A^0, A^0 \rightarrow \text{invisible}$	
50	AAD 22F	ATLS	$H_2^0 \rightarrow HH$	
51	AAD 22I	ATLS	$H \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^0, \tilde{\chi}_2^0 \rightarrow A^0 \tilde{\chi}_1^0, A^0 \rightarrow b \bar{b}$	
52	AAD 22J	ATLS	$H \rightarrow Z A^0$	
53	AAD 22J	ATLS	$H \rightarrow A^0 A^0, H_1^0 H_1^0$	OCCUR=2
54	AAD 22P	ATLS	$H_1^0, H_2^0 \rightarrow \text{invisible}$	
55	AAD 22Y	ATLS	$H_2^0 \rightarrow HH$	
56	ABRATENKO 22A	MCBN	$K^+ \rightarrow H_1^0 \pi^+$	
57	TUMASYAN 22AK	CMS	$H_3^0 \rightarrow H_1^0 H_1^0$	
58	TUMASYAN 22D	CMS	$H_2^0 \rightarrow W^+ W^-$	
59	AAD 21AF	ATLS	$H_2^0 \rightarrow ZZ$	
60	AAD 21AI	ATLS	$A^0 \rightarrow Z H_2^0$	
61	AAD 21AY	ATLS	$H_2^0 \rightarrow \gamma\gamma$	
62	AAD 21AZ	ATLS	$A_2^0 \rightarrow H A_1^0$	
63	AAD 21BB	ATLS	$A_2^0 \rightarrow H A_1^0$	
64	AAD 21BE	ATLS	$A_1^0 \rightarrow \text{invisible}$	
65	ABRATENKO 21	MCBN	$K^+ \rightarrow H_1^0 \pi^+$	
66	SIRUNYAN 21A	CMS	$H_2^0 \rightarrow Z A^0, A^0 \rightarrow \text{invisible}$	
67	TUMASYAN 21F	CMS	$H_3^0 \rightarrow H H_{1,2}^0$	
68	AAD 20AA	ATLS	$H_2^0 / A^0 \rightarrow \tau^+ \tau^-$	
69	AAD 20AI	ATLS	$H \rightarrow A^0 A^0$	
70	AAD 20AO	ATLS	$H_2^0 \rightarrow HH$	
71	AAD 20C	ATLS	$H_2^0 \rightarrow HH$	
72	AAD 20L	ATLS	$H_2^0 \rightarrow b \bar{b}$	
73	AAD 20X	ATLS	$H_2^0 \rightarrow HH$	

74	AAIJ	20AL LHCb	$A^0 \rightarrow \mu^+ \mu^-$	
75	SIRUNYAN	20 CMS	$H \rightarrow A^0 A^0$	
76	SIRUNYAN	20AA CMS	$H_2^0 \rightarrow Z A^0$ or $A^0 \rightarrow Z H_2^0$	
77	SIRUNYAN	20AC CMS	$A^0 \rightarrow Z H$	
78	SIRUNYAN	20AD CMS	$H_2^0 \rightarrow \mu \tau, e \tau$	
79	SIRUNYAN	20AF CMS	$H_2^0/A^0 \rightarrow t \bar{t}$	
80	SIRUNYAN	20AP CMS	$H, H_2^0 \rightarrow A^0 A^0$	
81	SIRUNYAN	20Y CMS	$H_2^0 \rightarrow W^+ W^-$	
82	SIRUNYAN	20Z CMS	$t \bar{t} H_{1,2}^0$ or $t \bar{t} A^0, H_{1,2}^0/$ $A^0 \rightarrow e^+ e^-, \mu^+ \mu^-$	
83	AABOUD	19A ATLS	$H_2^0 \rightarrow H H$	
84	AABOUD	19AG ATLS	$H \rightarrow A^0 A^0$	
85	AABOUD	19O ATLS	$H_2^0 \rightarrow H H$	
86	AABOUD	19T ATLS	$H_2^0 \rightarrow H H$	OCCUR=2
87	AABOUD	19V ATLS	two doublet + pseudoscalar model	OCCUR=2
88	AABOUD	19Y ATLS	$H_2^0 \rightarrow \mu^+ \mu^-$	
89	AALTONEN	19 CDF	$H_{1,2}^0 \rightarrow b \bar{b}$	
90	SIRUNYAN	19 CMS	$H_2^0 \rightarrow H H$	
91	SIRUNYAN	19AE CMS	$A^0 \rightarrow \tau^+ \tau^-$	
92	SIRUNYAN	19AN CMS	$A_2^0 \rightarrow H A_1^0$	
93	SIRUNYAN	19AV CMS	$A^0 \rightarrow Z H$	
94	SIRUNYAN	19B CMS	$H_{1,2}^0/A^0 \rightarrow b \bar{b}$	
95	SIRUNYAN	19BB CMS	$H_1^0 \rightarrow \gamma \gamma$	
96	SIRUNYAN	19BD CMS	$H \rightarrow A^0 A^0$	
97	SIRUNYAN	19BE CMS	$H_2^0 \rightarrow H H$	
98	SIRUNYAN	19BQ CMS	$H_{1,2}^0 \rightarrow A^0 A^0$	
99	SIRUNYAN	19CR CMS	$H_2^0/A^0 \rightarrow \mu^+ \mu^-$	
100	SIRUNYAN	19H CMS	$H_2^0 \rightarrow H H$	
101	AABOUD	18AA ATLS	$H_2^0 \rightarrow Z \gamma$	
102	AABOUD	18AG ATLS	$H \rightarrow A^0 A^0$	
103	AABOUD	18AH ATLS	$A^0 \rightarrow Z H_2^0$	
104	AABOUD	18AI ATLS	$A^0 \rightarrow Z H$	
105	AABOUD	18BF ATLS	$H_2^0 \rightarrow Z Z$	
106	AABOUD	18BU ATLS	$H_2^0 \rightarrow H H$	
107	AABOUD	18BX ATLS	$H \rightarrow A^0 A^0$	
108	AABOUD	18CQ ATLS	$H_2^0 \rightarrow H H$	
109	AABOUD	18F ATLS	$H_2^0 \rightarrow W^+ W^-, Z Z$	
110	AAIJ	18AMLHCb	$H_{1,2}^0 \rightarrow \mu \tau$	
111	AAIJ	18AQ LHCb	$A^0 \rightarrow \mu^+ \mu^-$	
112	AAIJ	18AQ LHCb	$H \rightarrow A^0 A^0, A^0 \rightarrow \mu^+ \mu^-$	OCCUR=2
113	SIRUNYAN	18AF CMS	$H_2^0 \rightarrow H H$	
114	SIRUNYAN	18BA CMS	$H_2^0 \rightarrow Z Z$	
115	SIRUNYAN	18CWCMS	$H_2^0 \rightarrow H H$	
116	SIRUNYAN	18DK CMS	$H_2^0 \rightarrow Z \gamma$	
117	SIRUNYAN	18DT CMS	$H \rightarrow A^0 A^0$	
118	SIRUNYAN	18DU CMS	$H_2^0 \rightarrow \gamma \gamma$	
119	SIRUNYAN	18ED CMS	$A^0 \rightarrow Z H$	
120	SIRUNYAN	18EE CMS	$H \rightarrow A^0 A^0$	
121	SIRUNYAN	18F CMS	$pp, 13 \text{ TeV}, H_2^0 \rightarrow H H$	
122	AABOUD	17 ATLS	$H_2^0 \rightarrow Z \gamma$	
123	AABOUD	17AP ATLS	$H_2^0 \rightarrow \gamma \gamma$	
124	AABOUD	17AW ATLS	$H_2^0 \rightarrow Z \gamma$	
125	KHACHATRY...17AZ	CMS	$H \rightarrow A^0 A^0$	
126	KHACHATRY...17D	CMS	$pp, 8, 13 \text{ TeV}, H_2^0 \rightarrow Z \gamma$	
127	KHACHATRY...17R	CMS	$H_2^0 \rightarrow \gamma \gamma$	
128	SIRUNYAN	17CN CMS	$pp, 8 \text{ TeV}, H_2^0 \rightarrow H H$	
129	SIRUNYAN	17Y CMS	$pp, 8, 13 \text{ TeV}, H_2^0 \rightarrow Z \gamma$	
130	AABOUD	16AB ATLS	$H \rightarrow A^0 A^0$	
131	AABOUD	16AE ATLS	$H_2^0 \rightarrow W^+ W^-, Z Z$	





4 AAD 24BQ search for the decay chain $H \rightarrow A^0 A^0 \rightarrow b\bar{b}\tau^+\tau^-$ , produced by gluon fusion, vector boson fusion, or in association with a weak vector boson, in 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 10 for limits on the product of branching ratios in the range $m_{A^0} = 12\text{--}60$ GeV.	NODE=S055H2O;LINKAGE=HF
5 AAD 24BV search for $H_2^0$ produced by vector-boson fusion, decaying to $HH \rightarrow b\bar{b}b\bar{b}$ , using 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 8 for limits on the product of production cross section times branching ratios for $m_{H_2^0} = 1\text{--}5$ TeV.	NODE=S055H2O;LINKAGE=TF
6 AAD 24BW search for production of $H_3^0$ decaying to $H_2^0 H$ , $H_2^0 \rightarrow W^+ W^-$ or $ZZ$ , and $H \rightarrow \gamma\gamma$ using 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Figs. 6–8 for limits on production cross section times branching ratios in the ranges $m_{H_3^0} = 0.3\text{--}1.0$ TeV and $m_{H_2^0} = 0.17\text{--}0.5$ TeV.	NODE=S055H2O;LINKAGE=KF
7 AAD 24BX search for production of $H_4^0$ decaying to $H_2^0 H_3^0$ , $H_2^0 \rightarrow$ invisible, and $H_3^0 \rightarrow ZZ \rightarrow 4\ell$ , using 139 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 8 (a), (b) for limits on production cross section times branching ratios in the ranges $m_{H_4^0} = 0.39\text{--}1.3$ TeV, $m_{H_3^0} = 0.22\text{--}1.0$ TeV, and $m_{H_2^0} = 160$ GeV.	NODE=S055H2O;LINKAGE=LF
8 AAD 24BX search for production of $A^0$ decaying to $H_2^0 Z$ , $H_2^0 \rightarrow ZZ$ , resulting in $4\ell$ final states, using 139 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 8 (c), (d) for limits on production cross section times branching ratios in the ranges $m_{A^0} = 0.32\text{--}1.3$ TeV and $m_{H_2^0} = 0.22\text{--}1.0$ TeV.	NODE=S055H2O;LINKAGE=MF
9 AAD 24CA search for production of $H_3^0$ decaying to $H_{1,2}^0 H$ , $H_{1,2}^0 \rightarrow b\bar{b}$ , and $H \rightarrow \gamma\gamma$ using 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 4 for limits on production cross section times branching ratios in the ranges $m_{H_3^0} = 0.17\text{--}1.0$ TeV and $m_{H_{1,2}^0} = 0.015\text{--}0.5$ TeV.	NODE=S055H2O;LINKAGE=NF
10 AAD 24CD search for production of $H_2^0$ decaying to $ZA$ , $A \rightarrow$ invisible, using 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Figs. 8–12 for excluded parameter regions in two-Higgs-doublet plus singlet pseudoscalar model.	NODE=S055H2O;LINKAGE=OF
11 AAD 24CF search for gluon-fusion production of $A^0$ decaying to $\tau^+\tau^-$ in 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 7 for limits on cross section times branching ratio in the range $m_{A^0} = 20\text{--}90$ GeV, and Fig. 8 for $A^0$ coupling with the top quark.	NODE=S055H2O;LINKAGE=JF
12 AAD 24CI search for production of $H_2^0$ , $A^0$ decaying to invisible final states, in various channels including $t\bar{t} + \text{missing } E_T$ and $t\bar{t}t\bar{t}$ , using up to 140 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 3 for cross section limits in a simplified model with fixed Dark Matter mass and couplings to quarks and Dark Matter.	NODE=S055H2O;LINKAGE=QF
13 AAD 24CK combines published ATLAS data up to 139 fb <sup>-1</sup> to constrain two-Higgs-doublet plus singlet pseudoscalar model with $A_1^0$ decaying to invisibly into Dirac Dark Matter final states. See their Figs. 4–9 for excluded parameter regions.	NODE=S055H2O;LINKAGE=RF
14 AAD 24H combine searches for a scalar resonance decaying to $HH$ using up to 139 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV from AAD 22F, AAD 23Z, and AAD 22Y. See their Fig. 1 for limits on cross section times branching ratio for $m_{H_2^0} = 0.251\text{--}5.0$ TeV.	NODE=S055H2O;LINKAGE=YE
15 AAD 24K search for the decay $H \rightarrow ZA^0$ with $A^0 \rightarrow \gamma\gamma$ and $Z \rightarrow \ell^+\ell^-$ in 139 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. Merged $\gamma\gamma$ events as well as resolved ones are looked for. See their Fig. 3 for limits on the product of the branching ratios for $m_{A^0} = 0.1\text{--}33$ GeV.	NODE=S055H2O;LINKAGE=AF
16 ADACHI 24F search for $H_1^0$ , which couples preferentially to muon pairs, in the process $e^+e^- \rightarrow \mu^+\mu^- H_1^0 \rightarrow \mu^+\mu^-\mu^+\mu^-$ using 178 fb <sup>-1</sup> of $e^+e^-$ collisions at $E_{\text{cm}} = 10.58$ GeV. See their Fig. 13 (lower) for limits on cross section times branching ratio for $m_{H_1^0} = 0.212\text{--}9.0$ GeV. Limits on the model parameter are given in Fig. 14 (lower).	NODE=S055H2O;LINKAGE=CF
17 HAYRAPETYAN 24AA search for the decay chain $H \rightarrow A^0 A^0 \rightarrow b\bar{b}b\bar{b}$ , produced in association with $W^\pm$ or $Z$ , in 138 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 3 for limits on the product of branching ratios in the range $m_{A^0} = 15\text{--}60$ GeV.	NODE=S055H2O;LINKAGE=WE
18 HAYRAPETYAN 24AE search for $H_2^0 \rightarrow HH$ in the final state $b\bar{b}W^+W^-$ using 138 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Fig. 19 (upper) for limit on the product of production cross section times branching ratios for $m_{H_2^0} = 0.25\text{--}0.9$ TeV.	NODE=S055H2O;LINKAGE=VE
19 HAYRAPETYAN 24AF search for $H \rightarrow A^0 A^0$ in the final state $b\bar{b}\mu^+\mu^-$ and $b\bar{b}\tau^+\tau^-$ in 138 fb <sup>-1</sup> of $pp$ collisions at $E_{\text{cm}} = 13$ TeV. See their Figs. 10, 11, and 12 for limits on the product of branching ratios in the range $m_{A^0} = 12\text{--}60$ GeV. Interpretation of the limits in terms of two Higgs doublet plus singlet models is given in their Figs. 13 and 14.	NODE=S055H2O;LINKAGE=BF

- 20 HAYRAPETYAN 24AI search for  $H \rightarrow H_1^0 H_1^0$ , with long-lived  $H_1^0$  decaying in the muon detectors to  $\gamma\gamma$ ,  $e^+e^-$ ,  $\pi^0\pi^0$ ,  $\pi^+\pi^-$ ,  $K^+K^-$ ,  $K^0\bar{K}^0$ ,  $d\bar{d}$ ,  $\tau^+\tau^-$ , or  $b\bar{b}$ , using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 13–17 for limits on  $B(H \rightarrow H_1^0 H_1^0)$  in the mass range between 0.4 and 55 GeV for each of the decay modes. NODE=S055H2O;LINKAGE=EF
- 21 HAYRAPETYAN 24AJ search for production of a scalar resonance decaying to  $\gamma\gamma$  in  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 3 (right) for limits on production cross section times branching ratio for  $m_{H_2^0} = 0.6\text{--}5.0 \text{ TeV}$ , and Fig. 4 (lower) for their dependence on the width. NODE=S055H2O;LINKAGE=XE
- 22 HAYRAPETYAN 24AZ search for pair production of  $A^0$  decaying to  $\mu^+\mu^-$  using  $101\text{--}137 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for limits on cross section times branching ratio for  $m_{A^0} = 0.21\text{--}60 \text{ GeV}$ . NODE=S055H2O;LINKAGE=IF
- 23 HAYRAPETYAN 24AZ search for  $H$  or  $H_{1,2}^0$  decaying to  $A^0 A^0 \rightarrow \mu^+\mu^-\mu^+\mu^-$  using  $101\text{--}137 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8 for limits on cross section times branching ratio for  $m_{A^0} = 0.5\text{--}2.7 \text{ GeV}$  and several choices of  $m_{H_{1,2}^0}$  in a NMSSM scenario. NODE=S055H2O;LINKAGE=SF
- 24 HAYRAPETYAN 24I search for the decay  $H \rightarrow Z A^0$  with  $A^0 \rightarrow \gamma\gamma$  and  $Z \rightarrow \ell^+\ell^-$  in  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on the product of cross section times the branching ratios for  $m_{A^0} = 1\text{--}30 \text{ GeV}$ . NODE=S055H2O;LINKAGE=ZE
- 25 TUMASYAN 24A search for production of  $H_{1,2}^0$  in association with  $W$ ,  $Z$ , or  $t\bar{t}$ , decaying to  $e^+e^-$ ,  $\mu^+\mu^-$ , or  $\tau^+\tau^-$ , using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 13 for limits on cross section times branching ratio for  $m_{H_{1,2}^0} = 15\text{--}350 \text{ GeV}$ . See also Figs. 16 and 17 for limits on the mixing with the Standard Model Higgs boson. NODE=S055H2O;LINKAGE=DF
- 26 TUMASYAN 24B search for  $H_2^0 \rightarrow HH$  in the final state  $\gamma\gamma b\bar{b}$  using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 (upper) for limit on the product of production cross section times branching ratios for  $m_{H_2^0} = 0.26\text{--}1.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=TE
- 27 TUMASYAN 24B search for production of  $H_3^0$  decaying to  $H_{1,2}^0 H \rightarrow b\bar{b}\gamma\gamma$  using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on production cross section times branching ratios for  $m_{H_3^0} = 0.3\text{--}1.0 \text{ TeV}$  and  $m_{H_{1,2}^0} = 0.09\text{--}0.8 \text{ TeV}$ . NODE=S055H2O;LINKAGE=UE
- 28 AAD 23AD search for associated production of  $W/Z H_2^0$  with the decay chain  $H_2^0 \rightarrow HH \rightarrow b\bar{b}b\bar{b}$  using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for limits on cross section times branching ratios for  $m_{H_2^0} = 260\text{--}1000 \text{ GeV}$ . NODE=S055H2O;LINKAGE=BE
- 29 AAD 23AD search for gluon fusion production of  $A^0$  with the decay chain  $A^0 \rightarrow Z H_2^0$ ,  $H_2^0 \rightarrow HH \rightarrow b\bar{b}b\bar{b}$  using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10 for limits on cross section times branching ratios for  $m_{A^0} = 350\text{--}800 \text{ GeV}$  and  $m_{H_2^0} = 260\text{--}400 \text{ GeV}$ . NODE=S055H2O;LINKAGE=CE
- 30 AAD 23AJ search for production of  $H^\pm$  in association with a top quark, followed by  $H^\pm \rightarrow W^\pm A^0$ ,  $A^0 \rightarrow \text{invisible}$ , using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10 for excluded parameter regions of 2HDM +  $CP$ -odd singlet model. NODE=S055H2O;LINKAGE=GE
- 31 AAD 23BD search for a top quark decaying to  $q H_{1,2}^0$  ( $q = u, c$ ),  $H_{1,2}^0 \rightarrow b\bar{b}$ , using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for limits on production cross section times branching ratios for  $m_{H_{1,2}^0} = 20\text{--}160 \text{ GeV}$ . NODE=S055H2O;LINKAGE=YD
- 32 AAD 23BE search for associated production of  $H_2^0 W$  and decay  $H_2^0 \rightarrow W^+ W^-$  assuming the presence of higher dimensional  $H_2^0 W^+ W^-$  interactions, using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for excluded parameter region of higher dimensional operators, and Fig. 7 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.3\text{--}1.5 \text{ TeV}$ . NODE=S055H2O;LINKAGE=FE
- 33 AAD 23BG search for production of  $H_2^0/A^0$  in association with a  $t\bar{t}$  pair, decaying to  $t\bar{t}$ , using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on cross section times branching ratios for  $m_{H_2^0} = m_{A^0} = 0.4\text{--}1.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=EE
- 34 AAD 23BW search for  $A^0$  production in association with a  $t\bar{t}$  pair, decaying to  $\mu^+\mu^-$ , using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5(a) for limits on production cross section times branching ratio for  $m_{A^0} = 15\text{--}72 \text{ GeV}$ . NODE=S055H2O;LINKAGE=RE
- 35 AAD 23BX search for production of  $H \rightarrow \tau^+\tau^-$  with missing transverse momentum using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8 for interpretation of the data in terms of 2HDM + a model. NODE=S055H2O;LINKAGE=LE
- 36 AAD 23CA search for production of  $H_3^0$  decaying to  $H_2^0 H$ ,  $H_2^0 \rightarrow W^+ W^-$  or  $ZZ$ , and  $H \rightarrow \tau^+\tau^-$  using  $140 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 4, 5 for limits on production cross section times branching ratios in the ranges  $m_{H_3^0} = 0.5\text{--}1.5 \text{ TeV}$  and  $m_{H_2^0} = 0.2\text{--}0.5 \text{ TeV}$ . NODE=S055H2O;LINKAGE=KE

- 37 AAD 23CR search for  $H_2^0$  having flavor-violating couplings to  $tc$  or  $tu$ , produced in association with top quark(s), using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 14 for limits on production cross section times branching ratios for  $m_{H_2^0} = 0.2\text{--}1.5 \text{ TeV}$  with various assumptions on the flavor-changing couplings. NODE=S055H2O;LINKAGE=OE
- 38 AAD 23O search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH$  in the final states  $\nu\bar{\nu}b\bar{b}$  and  $\ell^+\ell^-b\bar{b}$  using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for limits on cross section times branching ratio for  $m_{A^0} = 0.22\text{--}2.0 \text{ TeV}$ , and Fig. 11 for limits with both production components. NODE=S055H2O;LINKAGE=XD
- 39 AAD 23R search for the decay  $A^0 \rightarrow \gamma\gamma$  in  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on cross section times branching ratio for  $m_{A^0} = 10\text{--}70 \text{ GeV}$ . NODE=S055H2O;LINKAGE=RD
- 40 AAD 23U search for the decay  $H_2^0 \rightarrow Z\gamma$  with  $Z$  decaying hadronically in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8(a) for limits on production cross section times branching ratios for  $m_{H_2^0} = 1.0\text{--}6.8 \text{ TeV}$ . NODE=S055H2O;LINKAGE=TD
- 41 AAD 23Z search for the decay chain  $H_2^0 \rightarrow HH \rightarrow b\bar{b}\tau^+\tau^-$  using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10 for limits on the product of production cross section times branching ratios for  $m_{H_2^0} = 0.251\text{--}1.6 \text{ TeV}$ . NODE=S055H2O;LINKAGE=WD
- 42 HAYRAPETYAN 23C search for  $H_{1,2}^0 \rightarrow e\mu$  using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on production cross section times branching ratio for  $m_{H_{1,2}^0} = 110\text{--}160 \text{ GeV}$ . NODE=S055H2O;LINKAGE=SE
- 43 HAYRAPETYAN 23G search for dimuon resonance in the mass range  $1.1\text{--}2.6$  or  $4.2\text{--}7.9 \text{ GeV}$  in  $96.6 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ , in inclusive and high  $p_T$  selections. See their Fig. 5 for cross section times branching ratio limits and Fig. 7 for mixing angle limits in two Higgs doublet plus singlet model (at 90% CL). NODE=S055H2O;LINKAGE=JE
- 44 TUMASYAN 23 search for production of  $H_3^0$  decaying to  $H_{1,2}^0 H \rightarrow b\bar{b}b\bar{b}$  using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4 for limits on production cross section times branching ratios for  $m_{H_3^0} = 0.9\text{--}4.0 \text{ TeV}$  and  $m_{H_{1,2}^0} = 60\text{--}600 \text{ GeV}$ , and their interpretation in the NMSSM and the Two Real Singlet Model (TRSM). NODE=S055H2O;LINKAGE=DE
- 45 TUMASYAN 23M search for the decay chain  $H \rightarrow A^0 A^0 \rightarrow \gamma\gamma\gamma\gamma$  in  $132 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for limits on cross section times branching ratio in the range  $m_{A^0} = 15\text{--}62 \text{ GeV}$ . NODE=S055H2O;LINKAGE=SD
- 46 TUMASYAN 23O search for  $H_2^0 \rightarrow HH$ , each  $H$  decaying to either  $WW^*$  or  $\tau^+\tau^-$  using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 14 (upper) for limit on the product of production cross section times branching ratios for  $m_{H_2^0} = 0.25\text{--}1.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=UD
- 47 TUMASYAN 23S search for gluon fusion and  $b$ -associated production of  $H_{1,2}^0$  decaying to  $\tau^+\tau^-$  using  $138 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10 for limits on production cross section times branching ratios for  $m_{H_{1,2}^0} = 0.06\text{--}3.5 \text{ TeV}$ . NODE=S055H2O;LINKAGE=ZD
- 48 AAD 22A search for the decay chain  $H \rightarrow A^0 A^0 \rightarrow \mu^+\mu^-b\bar{b}$  in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 for limits on the overall branching fraction in the range  $m_{A^0} = 16\text{--}62 \text{ GeV}$ . See also Fig. 11 for limits without assuming  $A^0$  is pseudoscalar. NODE=S055H2O;LINKAGE=CD
- 49 AAD 22D search for  $ZA^0$  associate production with  $Z \rightarrow \ell^+\ell^-$ ,  $A^0$  decaying invisibly, in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for excluded regions in the mass parameter space of two Higgs doublet plus singlet (2HDM+ $A^0$ ) model with a certain choice of the model parameters. NODE=S055H2O;LINKAGE=JD
- 50 AAD 22F search for gluon fusion production of  $H_2^0$  decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  using  $126\text{--}139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ .  $B(H \rightarrow b\bar{b}) = 0.582$  is assumed. See their Fig. 14 for limit on the product of production cross section times branching ratios for  $m_{H_2^0} = 0.251\text{--}5.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=ID
- 51 AAD 22I search for  $ZH$  associate production with the decay chain  $H \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^0$ ,  $\tilde{\chi}_2^0 \rightarrow A^0 \tilde{\chi}_1^0$ ,  $A^0 \rightarrow b\bar{b}$ , and  $Z \rightarrow \ell^+\ell^-$ , in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 3 and 4 for limits on the product of cross section times the branching ratios for  $m_{A^0} = 20\text{--}65 \text{ GeV}$  with various choices of NMSSM model parameters. NODE=S055H2O;LINKAGE=KD
- 52 AAD 22J search for the decay  $H \rightarrow ZA^0$  with  $A^0 \rightarrow \mu^+\mu^-$  and  $Z \rightarrow e^+e^-$ ,  $\mu^+\mu^-$  in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  assuming SM gluon-gluon fusion production of the  $H$ . See their Fig. 17(b) for limits on the product of cross section times the branching ratios for  $m_{A^0} = 15\text{--}30 \text{ GeV}$ . NODE=S055H2O;LINKAGE=LD
- 53 AAD 22J search for the decay  $H \rightarrow A^0 A^0$  with  $A^0 \rightarrow \mu^+\mu^-$  in  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  assuming SM gluon-gluon fusion production of the  $H$  in the range of  $m_{A^0} = 1\text{--}60 \text{ GeV}$ . See their Fig. 14(b) for limits on the product of cross section times the branching ratios for  $m_{A^0} = 1.5\text{--}60 \text{ GeV}$  (excluding  $\psi$  and  $\mathcal{T}$  regions). The limit also applies to the decay  $H \rightarrow H_1^0 H_1^0$ . NODE=S055H2O;LINKAGE=MD

- 54 AAD 22P search for invisibly decaying  $H_1^0, H_2^0$  produced by vector boson fusion in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. Limit on the product of cross section times branching ratio in the range 0.1–1 pb (95% CL) is given for the mass range 0.05–2 TeV. See their Fig. 14. NODE=S055H2O;LINKAGE=ND
- 55 AAD 22Y search for gluon fusion production of  $H_2^0$  decaying to  $HH \rightarrow b\bar{b}\gamma\gamma$  in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 15 for limit on the product of production cross section times branching ratios to  $HH$  for  $m_{H_2^0} = 0.251$ –1.0 TeV. NODE=S055H2O;LINKAGE=PD
- 56 ABRATENKO 22A search for a singlet scalar boson  $H_1^0$  having a small mixing with the SM Higgs boson in the decay chain  $K^+ \rightarrow H_1^0 \pi^+, H_1^0 \rightarrow \mu^+ \mu^-$  from data corresponding to  $7.01 \times 10^{20}$  protons on NuMI target. See their Fig. 13 (right) and Table V for limits on the SM Higgs component of  $H_1^0$  for  $m_{H_1^0} = 212$ –279 MeV. NODE=S055H2O;LINKAGE=QD
- 57 TUMASYAN 22AK search for gluon-fusion production of  $H_3^0$  decaying to  $H_1^0 H_1^0 \rightarrow b\bar{b}b\bar{b}$  in 138 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 5 for limits on cross section times branching ratio for  $m_{H_3^0} = 1$ –3 TeV,  $m_{H_1^0} = 25$ –100 GeV. NODE=S055H2O;LINKAGE=OD
- 58 TUMASYAN 22D search for production of an  $H_2^0$  (denoted radion in the paper) in gluon fusion and vector boson fusion, decaying to  $W^+ W^-$  in the final states  $\ell\nu + \text{hadrons}$ , using 137 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 7 for limits on cross section times branching ratio for  $m_{H_2^0} = 1.0$ –4.5 TeV. NODE=S055H2O;LINKAGE=HE
- 59 AAD 21AF search for production of a heavy  $H_2^0$  state decaying to  $ZZ$  in the final states  $\ell^+ \ell^- \ell'^+ \ell'^-$  and  $\ell^+ \ell^- \nu\bar{\nu}$  in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 4 for upper limits on cross section times branching ratio for  $m_{H_2^0} = 0.2$ –2.0 TeV assuming ggF or VBF with narrow width approximation, and Fig. 5 for upper limits on cross section times branching ratio for  $m_{H_2^0} = 0.4$ –2.0 TeV assuming ggF, and with several assumptions on its width. NODE=S055H2O;LINKAGE=XC
- 60 AAD 21AI search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH_2^0 \rightarrow \ell^+ \ell^- b\bar{b}$  or  $\ell^+ \ell^- W^+ W^-$  in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Figs. 9 and 13 for cross section limits for  $m_{A^0} = 230$ –800 GeV and  $m_{H_2^0} = 130$ –700 GeV. NODE=S055H2O;LINKAGE=YC
- 61 AAD 21AY search for production of a scalar resonance decaying to  $\gamma\gamma$  in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 5(a) for limits on fiducial cross section times branching ratio for  $m_{H_2^0} = 0.16$ –3 TeV with narrow width approximation, and Table 2 with several assumptions on the width. NODE=S055H2O;LINKAGE=ZC
- 62 AAD 21AZ search for production of  $A_2^0$  decaying to  $HA_1^0$  followed by  $H \rightarrow \gamma\gamma, A_1^0 \rightarrow$  invisible in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Figs. 10–12 for limits in terms of two-Higgs-doublet model plus singlet pseudoscalar and a fermionic Dark Matter particle. NODE=S055H2O;LINKAGE=FD
- 63 AAD 21BB search for production of  $A_2^0$  by gluon fusion or associated  $A_2^0 b\bar{b}$  production, decaying to  $HA_1^0$  followed by  $H \rightarrow b\bar{b}, A_1^0 \rightarrow$  invisible in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 8 for limits in terms of two-Higgs-doublet plus singlet pseudoscalar model. NODE=S055H2O;LINKAGE=GD
- 64 AAD 21BE search for production of  $A_1^0$  associated with a single top quark and either a light quark or a  $W$  boson, decaying to invisible final states, in 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Figs. 13–15 for limits in terms of two-Higgs-doublet model plus singlet pseudoscalar, which is assumed to decay to a pair of Dark Matter particles. NODE=S055H2O;LINKAGE=HD
- 65 ABRATENKO 21 search for a singlet scalar boson  $H_1^0$  having a small mixing with the SM Higgs boson in the decay chain  $K^+ \rightarrow H_1^0 \pi^+, H_1^0 \rightarrow e^+ e^-$  from data corresponding to  $1.93 \times 10^{20}$  protons on NuMI target. See their Fig. 2 for limits on the SM Higgs component of  $H_1^0$  for  $m_{H_1^0} = 3$ –210 MeV. NODE=S055H2O;LINKAGE=DD
- 66 SIRUNYAN 21A search for  $H_2^0 \rightarrow ZA^0$  with  $Z \rightarrow \ell^+ \ell^-$ ,  $A^0$  decaying invisibly, in 137 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 8 for excluded regions in the mass parameter space of two Higgs doublet plus singlet model with a certain choice of the model parameters. NODE=S055H2O;LINKAGE=VC
- 67 TUMASYAN 21F search for gluon fusion production of  $H_3^0$  decaying to  $HH_{1,2}^0 \rightarrow \tau^+ \tau^- b\bar{b}$  in 137 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Figs. 5 and 6 for limits on cross section times branching ratios for  $m_{H_{1,2}^0} = 0.06$ –2.8 TeV and  $m_{H_3^0} = 0.24$ –3.0 TeV. NODE=S055H2O;LINKAGE=ED
- 68 AAD 20AA search for  $H_2^0/A^0 \rightarrow \tau^+ \tau^-$  produced by gluon fusion or  $b$ -associated production using 139 fb<sup>-1</sup> of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 2(a), 2(b) for limits on the product of cross section and branching ratio for  $m_{H_2^0}, m_{A^0} = 0.2$ –2.5 TeV. NODE=S055H2O;LINKAGE=LC

- 69 AAD 20AI search for  $ZH$  production followed by the decay  $H \rightarrow A^0 A^0 \rightarrow b\bar{b}b\bar{b}$  in  $36 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . The search looks for collimated  $A^0 \rightarrow b\bar{b}$  decays and is complementary to AABOUD 18BX. See their Fig. 10 for limits on the product of production cross section and branching ratios in the range  $m_{A^0} = 15\text{--}30 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=TC
- 70 AAD 20AO search for gluon fusion production of  $H_2^0$  decaying to  $HH \rightarrow \tau^+ \tau^- b\bar{b}$  (with hadronically decaying  $\tau^+ \tau^-$ ) using  $139 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . Limit on the product of production cross section times branching ratios in the range  $28\text{--}817 \text{ fb}$  (95% CL) is given for  $m_{A^0} = 1.0\text{--}3.0 \text{ TeV}$ , see their Fig. 13.  
NODE=S055H2O;LINKAGE=UC
- 71 AAD 20C combine searches for a scalar resonance decaying to  $HH$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  from AABOUD 19A, AABOUD 19O, AABOUD 18CQ, AABOUD 19T, AABOUD 18CW, and AABOUD 18BU. See their Fig. 5(a) for limits on cross section times branching ratio for  $m_{H_2^0} = 0.26\text{--}3 \text{ TeV}$ .  
NODE=S055H2O;LINKAGE=WB
- 72 AAD 20L search for  $b$ -associated production of  $H_2^0$  decaying to  $b\bar{b}$  in  $27.8 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8 for limits on the product of cross section and branching ratio for  $m_{H_2^0} = 0.45\text{--}1.4 \text{ TeV}$ .  
NODE=S055H2O;LINKAGE=RC
- 73 AAD 20X search for vector-boson-fusion production of  $H_2^0$  decaying to  $HH$  using  $126 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for limits on the product of cross section and branching ratio for the assumptions of a narrow- and broad-width resonance.  
NODE=S055H2O;LINKAGE=QC
- 74 AAIJ 20AL search for dimuon resonance in the mass range  $0.2\text{--}60 \text{ GeV}$  in  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ , in inclusive and  $b$  quark associated production. Displaced decays are searched for for masses below  $3 \text{ GeV}$ . See their Figs. 7–9 for cross section limits and Fig. 10 for limits for mixing angle in two Higgs doublet plus singlet model (at 90% CL).  
NODE=S055H2O;LINKAGE=WC
- 75 SIRUNYAN 20 search for the decay  $H \rightarrow A^0 A^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$  or  $\tau^+ \tau^- \mu^+ \mu^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10 for limits on the product of production cross section (normalized to the SM) and branching ratios in the range  $m_{A^0} = 4\text{--}15 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=GC
- 76 SIRUNYAN 20AA search for  $H_2^0 \rightarrow ZA^0$ ,  $A^0 \rightarrow b\bar{b}$  or  $A^0 \rightarrow ZH_2^0$ ,  $H_2^0 \rightarrow b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on the product of cross section and branching ratio for  $m_{H_2^0} = 0.12\text{--}1 \text{ TeV}$  and  $m_{A^0} = 0.03\text{--}1 \text{ TeV}$ .  
NODE=S055H2O;LINKAGE=NC
- 77 SIRUNYAN 20AC search for gluon-fusion production of  $A^0$  decaying to  $ZH$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for limits on the product of cross section and branching ratios for  $m_{A^0} = 220\text{--}400 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=OC
- 78 SIRUNYAN 20AD search for lepton-flavor violating decays  $H_2^0 \rightarrow \mu\tau$ ,  $e\tau$  of gluon-fusion-produced  $H_2^0$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 (9) and Table 5 (6) for limits on production cross section times branching ratio for  $m_{H_2^0} = 0.2\text{--}0.9 \text{ TeV}$  for the  $\mu\tau$  ( $e\tau$ ) final state.  
NODE=S055H2O;LINKAGE=KC
- 79 SIRUNYAN 20AF search for  $H_2^0/A^0 \rightarrow t\bar{t}$  with one or two charged leptons in the final state using kinematic variables in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 5 and 6 for limits on top Yukawa coupling of  $H_2^0$  and  $A^0$  for  $m_{H_2^0}$ ,  $m_{A^0} = 0.4\text{--}0.75 \text{ TeV}$  for various width assumptions.  
NODE=S055H2O;LINKAGE=MC
- 80 SIRUNYAN 20AP search for the decay  $H$  or  $H_2^0 \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- \tau^+ \tau^-$  (for  $m_{H_2^0} = 300 \text{ GeV}$ ) with boosted final-state topology in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on the product of production cross section (normalized to the SM) and branching ratios in the range  $m_{A^0} = 3.6\text{--}21 \text{ GeV}$ , and Figs. 8 and 9 for its interpretation in terms of models with two Higgs doublets plus a singlet.  
NODE=S055H2O;LINKAGE=JC
- 81 SIRUNYAN 20Y search for gluon-fusion and vector-boson-fusion production of  $H_2^0$  decaying to  $W^+ W^-$  in the final states  $\ell\nu\ell\nu$  and  $\ell\nu qq$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for limits on the product of cross section and branching ratio for  $m_{H_2^0} = 0.2\text{--}3 \text{ TeV}$ .  
NODE=S055H2O;LINKAGE=PC
- 82 SIRUNYAN 20Z search for  $H_{1,2}^0$  or  $A^0$  production in association with a  $t\bar{t}$  pair, decaying to  $e^+ e^-$  or  $\mu^+ \mu^-$ , in  $137 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 12 for limits on production cross section times branching ratio for  $m_{H_{1,2}^0}$ ,  $m_{A^0} = 15\text{--}75 \text{ GeV}$  and  $108\text{--}340 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=IC
- 83 AABOUD 19A search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $27.5\text{--}36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9(a) for limits on cross section times branching ratios for  $m_{H_2^0} = 0.26\text{--}3 \text{ TeV}$ .  
NODE=S055H2O;LINKAGE=FB
- 84 AABOUD 19AG search for the decay  $H \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- b\bar{b}$  in  $36.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 (a) for limits on the product of production cross section (normalized to the SM) and branching ratios in the range  $m_{A^0} = 20\text{--}60 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=AC

- 85 AABOUD 19O search for a scalar resonance decaying to  $HH \rightarrow b\bar{b}WW^*$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 12 (left) for limits on cross section times branching ratio for  $m_{H_2^0} = 0.5\text{--}3 \text{ TeV}$ . NODE=S055H2O;LINKAGE=VB
- 86 AABOUD 19T search for a scalar resonance decaying to  $HH \rightarrow WW^*WW^*$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 3 for limits on cross section times branching ratio for  $m_{H_2^0} = 260\text{--}500 \text{ GeV}$ , assuming SM decay rates for the  $H$ . NODE=S055H2O;LINKAGE=FC
- 87 AABOUD 19V combine published ATLAS data to constrain two-Higgs-doublet plus singlet pseudoscalar model with  $A_1^0$  decaying to invisible final states. See their Fig. 19 for excluded parameter regions. NODE=S055H2O;LINKAGE=EC
- 88 AABOUD 19Y search for a narrow scalar resonance produced by gluon fusion or  $b$  associated production, decaying to  $\mu^+\mu^-$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 4 and 5(a) for cross section limits for  $m_{H_2^0} = 0.2\text{--}1.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=SB
- 89 AALTONEN 19 search for  $b$  associated production of a scalar particle decaying to  $b\bar{b}$  in  $5.4 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 3 for limits on cross section times branching ratio for  $m_{H_{1,2}^0} = 100\text{--}300 \text{ GeV}$ . NODE=S055H2O;LINKAGE=XB
- 90 SIRUNYAN 19 search for a narrow scalar resonance decaying to  $HH \rightarrow \gamma\gamma b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 9 (left) for limits on cross section times branching ratios for  $m_{H_2^0} = 260\text{--}900 \text{ GeV}$ . NODE=S055H2O;LINKAGE=CB
- 91 SIRUNYAN 19AE search for a scalar resonance produced in association with a  $b\bar{b}$  pair, decaying to  $\tau^+\tau^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4 for cross section limits for  $m_{A^0} = 25\text{--}70 \text{ GeV}$ . NODE=S055H2O;LINKAGE=TB
- 92 SIRUNYAN 19AN search for production of  $A_2^0$  decaying to  $HA_1^0$  followed by  $H \rightarrow b\bar{b}$ ,  $A_1^0 \rightarrow$  invisible in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ , in the mass range  $m_{A_2^0} = 0.2\text{--}1.6 \text{ TeV}$ ,  $m_{A_1^0} = 0.15\text{--}0.5 \text{ TeV}$ . See their Fig. 6 for limits in terms of two-Higgs-doublet plus singlet pseudoscalar model. NODE=S055H2O;LINKAGE=CC
- 93 SIRUNYAN 19AV search for a scalar resonance produced by gluon fusion or  $b$ -associated production, decaying to  $ZH \rightarrow \ell^+\ell^-b\bar{b}$  ( $\ell = e, \mu$ ) or  $\nu\bar{\nu}b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for cross section limits for  $m_{A^0} = 0.22\text{--}1.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=YB
- 94 SIRUNYAN 19B search for gluon fusion production of narrow scalar resonance with large transverse momentum, decaying to  $b\bar{b}$ , in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 7 and 8 for limits on cross section times branching ratio for the resonance mass of  $50\text{--}350 \text{ GeV}$ . NODE=S055H2O;LINKAGE=QB
- 95 SIRUNYAN 19BB search for the decay  $H_1^0 \rightarrow \gamma\gamma$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  and  $35.9 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 4–6 for limits on cross section times branching ratio for  $m_{H_1^0} = 80\text{--}110 \text{ GeV}$  (some results in Fig. 5 for  $m_{H_1^0} = 70\text{--}110 \text{ GeV}$ ). NODE=S055H2O;LINKAGE=RB
- 96 SIRUNYAN 19BD search for the decay  $H \rightarrow A^0A^0 \rightarrow \mu^+\mu^-b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 5 for limits on the product of cross section times branching ratios in the range  $m_{A^0} = 20\text{--}62.5 \text{ GeV}$ . See also their Figs. 6 and 7 for interpretation of the data in terms of models with two Higgs doublets and a singlet. NODE=S055H2O;LINKAGE=ZB
- 97 SIRUNYAN 19BE combine searches for  $H_2^0 \rightarrow HH$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$  in various  $H$  decay modes, from SIRUNYAN 18A, SIRUNYAN 18AF, SIRUNYAN 18CW, SIRUNYAN 19, and SIRUNYAN 19H. See their Fig. 3 for limits on cross section times branching ratios for  $m_{H_2^0} = 0.25\text{--}3 \text{ TeV}$ . NODE=S055H2O;LINKAGE=UB
- 98 SIRUNYAN 19BQ search for production of  $H_{1,2}^0$  decaying to  $A^0A^0 \rightarrow \mu^+\mu^-\mu^+\mu^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 2 for limits on cross section times branching ratio for  $m_{H_{1,2}^0} = 90\text{--}150 \text{ GeV}$ ,  $m_{A^0} = 0.25\text{--}3.55 \text{ GeV}$ . NODE=S055H2O;LINKAGE=BC
- 99 SIRUNYAN 19CR search for production of  $H_2^0/A^0$  in gluon fusion and in association with a  $b\bar{b}$  pair, decaying to  $\mu^+\mu^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 6 for limits on cross section times branching ratio. NODE=S055H2O;LINKAGE=HC
- 100 SIRUNYAN 19H search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ , where one  $b\bar{b}$  pair is resolved and the other not. Limits on cross section times branching ratios for  $m_{H_2^0} = 0.75\text{--}1.6 \text{ TeV}$  are obtained and combined with data from SIRUNYAN 18AF. See their Fig. 5 (right). NODE=S055H2O;LINKAGE=KB
- 101 AABOUD 18AA search for production of a scalar resonance decaying to  $Z\gamma$ , with  $Z$  decaying hadronically, in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8(a) for limits on cross section times branching ratio for  $m_{H_2^0} = 1.0\text{--}6.8 \text{ TeV}$ . NODE=S055H2O;LINKAGE=TA
- 102 AABOUD 18AG search for the decay  $H \rightarrow A^0A^0 \rightarrow \gamma\gamma gg$  in  $36.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 2 and Table 6 for cross section limits in the range  $m_{A^0} = 20\text{--}60 \text{ GeV}$ . NODE=S055H2O;LINKAGE=YA
- 103 AABOUD 18AH search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH_2^0 \rightarrow \ell^+\ell^-b\bar{b}$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} =$  NODE=S055H2O;LINKAGE=SA

- 13 TeV. See their Fig. 5 for cross section limits for  $m_{A^0} = 230\text{--}800$  GeV and  $m_{H_2^0} = 130\text{--}700$  GeV.
- 104 AABOUD 18AI search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH$  in the final states  $\nu\bar{\nu}b\bar{b}$  and  $\ell^+\ell^-b\bar{b}$  in  $36.1\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 6 for cross section limits for  $m_{A^0} = 0.2\text{--}2$  TeV. See also AABOUD 18CC. NODE=S055H2O;LINKAGE=RA
- 105 AABOUD 18BF search for production of a heavy  $H_2^0$  state decaying to  $ZZ$  in the final states  $\ell^+\ell^-\ell^+\ell^-$  and  $\ell^+\ell^-\nu\bar{\nu}$  in  $36.1\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 6 for upper limits on cross section times branching ratio for  $m_{H_2^0} = 0.2\text{--}1.2$  TeV assuming ggF or VBF with the NWA. See their Fig. 7 for upper limits on cross section times branching ratio for  $m_{H_2^0} = 0.4\text{--}1.0$  TeV assuming ggF, and with several assumptions on its width. NODE=S055H2O;LINKAGE=VA
- 106 AABOUD 18BU search for a narrow scalar resonance decaying to  $HH \rightarrow \gamma\gamma WW^*$  in  $36.1\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 4 for limits on cross section times branching ratios for  $m_{H_2^0} = 260\text{--}500$  GeV. NODE=S055H2O;LINKAGE=EB
- 107 AABOUD 18BX search for associated production of  $WH$  or  $ZH$  followed by the decay  $H \rightarrow A^0 A^0 \rightarrow b\bar{b}b\bar{b}$  in  $36.1\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 9 for limits on cross section times branching ratios for  $m_{A^0} = 20\text{--}60$  GeV. See also their Fig. 10 for the dependence of the limit on  $A^0$  lifetime. NODE=S055H2O;LINKAGE=PB
- 108 AABOUD 18CQ search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}\tau^+\tau^-$  in  $36.1\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 2 (above) for limits on cross section times branching ratios for  $m_{H_2^0} = 260\text{--}1000$  GeV. NODE=S055H2O;LINKAGE=DB
- 109 AABOUD 18F search for production of a narrow scalar resonance decaying to  $W^+W^-$  and  $ZZ$ , followed by hadronic decays of  $W$  and  $Z$ , in  $36.7\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 5(c) for limits on cross section times branching ratio for  $m_{H_2^0} = 1.2\text{--}3.0$  TeV. NODE=S055H2O;LINKAGE=UA
- 110 AAIJ 18AM search for gluon-fusion production of  $H_{1,2}^0$  decaying to  $\mu\tau$  in  $2\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 2 for limits on cross section times branching ratio for  $m_{H_{1,2}^0} = 45\text{--}195$  GeV. NODE=S055H2O;LINKAGE=BB
- 111 AAIJ 18AQ search for gluon-fusion production of a scalar particle  $A^0$  decaying to  $\mu^+\mu^-$  in  $1.99\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV and  $0.98\text{ fb}^{-1}$  at  $E_{\text{cm}} = 7$  TeV. See their Fig. 4 for limits on cross section times branching ratio for  $m_{A^0} = 5.5\text{--}15$  GeV (using the  $E_{\text{cm}} = 8$  TeV data set). NODE=S055H2O;LINKAGE=NB
- 112 AAIJ 18AQ search for the decay  $H \rightarrow A^0 A^0$ , with one of the  $A^0$  decaying to  $\mu^+\mu^-$ , in  $1.99\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV and  $0.98\text{ fb}^{-1}$  at  $E_{\text{cm}} = 7$  TeV. See their Fig. 5 (right) for limits on the product of branching ratios for  $m_{A^0} = 5.5\text{--}15$  GeV (using the  $E_{\text{cm}} = 8$  TeV data set). NODE=S055H2O;LINKAGE=OB
- 113 SIRUNYAN 18AF search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $35.9\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV, where both  $b\bar{b}$  pairs are not resolved. See their Fig. 9 for limits on cross section times branching ratios for  $m_{H_2^0} = 0.75\text{--}3$  TeV. NODE=S055H2O;LINKAGE=WA
- 114 SIRUNYAN 18BA search for production of a heavy  $H_2^0$  state decaying to  $ZZ$  in the final states  $\ell^+\ell^-\ell^+\ell^-$ ,  $\ell^+\ell^-q\bar{q}$ , and  $\ell^+\ell^-\nu\bar{\nu}$  in  $35.9\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Figs. 10 and 11 for upper limits on cross section times branching ratio for  $m_{H_2^0} = 0.13\text{--}3$  TeV with several assumptions on its width and on the fraction of Vector-Boson-Fusion of the total production cross section. NODE=S055H2O;LINKAGE=XA
- 115 SIRUNYAN 18CW search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $35.9\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV, where both  $b\bar{b}$  pairs are resolved. See their Fig. 9 for limits on cross section times branching ratios for  $m_{H_2^0} = 260\text{--}1200$  GeV. NODE=S055H2O;LINKAGE=QA
- 116 SIRUNYAN 18DK search for production of a scalar resonance decaying to  $Z\gamma$ , with  $Z$  decaying to  $\ell^+\ell^-$  or hadronically, in  $35.9\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 7 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.35\text{--}4$  TeV for different assumptions on the width of the resonance. NODE=S055H2O;LINKAGE=JB
- 117 SIRUNYAN 18DT search for the decay  $H \rightarrow A^0 A^0 \rightarrow \tau^+\tau^-b\bar{b}$  in  $35.9\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 7 for limits on the product of branching ratios in the range  $m_{A^0} = 15\text{--}60$  GeV. See also their Fig. 8 for interpretation of the data in terms of models with two Higgs doublets and a singlet. NODE=S055H2O;LINKAGE=LB
- 118 SIRUNYAN 18DU search for production of a narrow scalar resonance decaying to  $\gamma\gamma$  in  $35.9\text{ fb}^{-1}$  (taken in 2016) of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 3 (right) for limits on cross section times branching ratio for  $m_{H_2^0} = 0.5\text{--}5$  TeV for several values of its width-to-mass ratio. NODE=S055H2O;LINKAGE=IB
- 119 SIRUNYAN 18ED search for production of an  $A^0$  in gluon-gluon fusion and in association with a  $b\bar{b}$ , decaying to  $ZH$  in the final states  $\nu\bar{\nu}b\bar{b}$  or  $\ell^+\ell^-b\bar{b}$  in  $35.9\text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13$  TeV. See their Fig. 8 for cross section limits for  $m_{A^0} = 0.8\text{--}2$  TeV. NODE=S055H2O;LINKAGE=AB



- 120 SIRUNYAN 18EE search for the decay  $H \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- \tau^+ \tau^-$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4 for limits on the product of branching ratios in the range  $m_{A^0} = 15\text{--}62.5 \text{ GeV}$ , normalized to the SM production cross section. See also their Fig. 5 for interpretation of the data in terms of models with two Higgs doublets and a singlet. NODE=S055H2O;LINKAGE=MB
- 121 SIRUNYAN 18F search for a narrow scalar resonance decaying to  $HH \rightarrow WWb\bar{b}$  or  $ZZb\bar{b}$  in the final state  $\ell\ell\nu\nu b\bar{b}$  in  $35.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on cross section times branching ratios for  $m_{H_2^0} = 250\text{--}900 \text{ GeV}$ . NODE=S055H2O;LINKAGE=LA
- 122 AABOUD 17 search for production of a scalar resonance decaying to  $Z\gamma$  in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4 for the limits on cross section times branching ratio for  $m_{H_2^0} = 0.25\text{--}3.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=IA
- 123 AABOUD 17AP search for production of a scalar resonance decaying to  $\gamma\gamma$  in  $36.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4(a) for limits on fiducial cross section times branching ratio for  $m_{H_2^0} = 0.2\text{--}2.7 \text{ TeV}$  with narrow width approximation. NODE=S055H2O;LINKAGE=BD
- 124 AABOUD 17AW search for production of a scalar resonance decaying to  $Z\gamma$  in  $36.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 7 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.25\text{--}2.4 \text{ TeV}$ . NODE=S055H2O;LINKAGE=KA
- 125 KHACHATRYAN 17AZ search for the decay  $H \rightarrow A^0 A^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$ ,  $\mu^+ \mu^- b\bar{b}$ , and  $\mu^+ \mu^- \tau^+ \tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 4, 5, and 6 for cross section limits in the range  $m_{A^0} = 5\text{--}62.5 \text{ GeV}$ . See also their Figs. 7, 8, and 9 for interpretation of the data in terms of models with two Higgs doublets and a singlet. NODE=S055H2O;LINKAGE=ZA
- 126 KHACHATRYAN 17D search for production of a scalar resonance decaying to  $Z\gamma$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  and  $2.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 3 and 4 for the limits on cross section times branching ratio for  $m_{H_2^0} = 0.2\text{--}2.0 \text{ TeV}$ . NODE=S055H2O;LINKAGE=OA
- 127 KHACHATRYAN 17R search for production of a narrow scalar resonance decaying to  $\gamma\gamma$  in  $12.9 \text{ fb}^{-1}$  (taken in 2016) of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 2 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.5\text{--}4.5 \text{ TeV}$  for several values of its width-to-mass ratio. Limits from combination with KHACHATRYAN 16M are shown in their Figs. 4 and 6. NODE=S055H2O;LINKAGE=JA
- 128 SIRUNYAN 17CN search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}\tau^+\tau^-$  in  $18.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 5 (above) and Table II for limits on the cross section times branching ratios for  $m_{H_2^0} = 0.3\text{--}1 \text{ TeV}$ , and Fig. 6 (above) and Table III for the corresponding limits by combining with data from KHACHATRYAN 16BQ and KHACHATRYAN 15R. NODE=S055H2O;LINKAGE=NA
- 129 SIRUNYAN 17Y search for production of a scalar resonance decaying to  $Z\gamma$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$  and  $2.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Figs. 3, 4 and Table 3 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.7\text{--}3.0 \text{ TeV}$ , and Fig. 5 for the corresponding limits for  $m_{H_2^0} = 0.2\text{--}3.0 \text{ TeV}$  from combination with KHACHATRYAN 17D data. NODE=S055H2O;LINKAGE=MA
- 130 AABOUD 16AB search for associated production of  $WH$  with the decay  $H \rightarrow A^0 A^0 \rightarrow b\bar{b}b\bar{b}$  in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 8 for limits on cross section times branching ratios for  $m_{A^0} = 20\text{--}60 \text{ GeV}$ . NODE=S055H2O;LINKAGE=FA
- 131 AABOUD 16AE search for production of a narrow scalar resonance decaying to  $W^+W^-$  and  $ZZ$  in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 4 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.5\text{--}3 \text{ TeV}$ . NODE=S055H2O;LINKAGE=EA
- 132 AABOUD 16H search for production of a scalar resonance decaying to  $\gamma\gamma$  in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 12 for limits on cross section times branching ratio for  $m_{H_2^0} = 0.2\text{--}2 \text{ TeV}$  with different assumptions on the width. NODE=S055H2O;LINKAGE=DA
- 133 AABOUD 16I search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 13 \text{ TeV}$ . See their Fig. 10(c) for limits on cross section times branching ratios for  $m_{H_2^0} = 0.5\text{--}3 \text{ TeV}$ . NODE=S055H2O;LINKAGE=T
- 134 AAD 16AX search for production of a heavy  $H$  state decaying to  $ZZ$  in the final states  $\ell^+\ell^-\ell^+\ell^-$ ,  $\ell^+\ell^-\nu\bar{\nu}$ ,  $\ell^+\ell^-q\bar{q}$ , and  $\nu\bar{\nu}q\bar{q}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 12 for upper limits on  $\sigma(H) \text{ B}(H \rightarrow ZZ)$  for  $m_H$  ranging from 140 GeV to 1000 GeV. NODE=S055H2O;LINKAGE=R
- 135 AAD 16C search for production of a heavy  $H$  state decaying to  $W^+W^-$  in the final states  $\ell\nu\ell\nu$  and  $\ell\nu q\bar{q}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 12, 13, and 16 for upper limits on  $\sigma(H) \text{ B}(H \rightarrow W^+W^-)$  for  $m_H$  ranging from 300 GeV to 1000 or 1500 GeV with various assumptions on the total width of  $H$ . NODE=S055H2O;LINKAGE=K
- 136 AAD 16L search for the decay  $H \rightarrow A^0 A^0 \rightarrow \gamma\gamma\gamma\gamma$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 4 (upper right) for limits on cross section times branching ratios (normalized to the SM  $H$  cross section) for  $m_{A^0} = 10\text{--}60 \text{ GeV}$ . NODE=S055H2O;LINKAGE=Z
- 137 AAD 16L search for the decay  $H_2^0 \rightarrow A^0 A^0 \rightarrow \gamma\gamma\gamma\gamma$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 4 (lower right) for limits on cross section times branching NODE=S055H2O;LINKAGE=BA

- ratios for  $m_{H_2^0} = 600$  GeV and  $m_{A^0} = 10\text{--}245$  GeV, and Table 5 for limits for  $m_{H_2^0} = 300$  and  $900$  GeV.
- 138 AALTONEN 16C search for electroweak associated production of  $H_1^0 H^\pm$  followed by the decays  $H^\pm \rightarrow H_1^0 W^*$ ,  $H_1^0 \rightarrow \gamma\gamma$  for  $m_{H_1^0} = 10\text{--}105$  GeV and  $m_{H^\pm} = 30\text{--}300$  GeV. NODE=S055H2O;LINKAGE=S
- See their Fig. 3 for excluded parameter region in a two-doublet model in which  $H_1^0$  has no direct decay to fermions.
- 139 KHACHATRYAN 16BG search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 6 for limits on the cross section times branching ratios for  $m_{H_2^0} = 1.15\text{--}3$  TeV. NODE=S055H2O;LINKAGE=V
- 140 KHACHATRYAN 16BQ search for a resonance decaying to  $HH \rightarrow \gamma\gamma b\bar{b}$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 9 for limits on the cross section times branching ratios for  $m_{H_2^0} = 0.26\text{--}1.1$  TeV. NODE=S055H2O;LINKAGE=PA
- 141 KHACHATRYAN 16F search for the decay  $H \rightarrow H_1^0 H_1^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 8 for cross section limits for  $m_{H_1^0} = 4\text{--}8$  GeV. NODE=S055H2O;LINKAGE=P
- 142 KHACHATRYAN 16M search for production of a narrow resonance decaying to  $\gamma\gamma$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV and  $3.3 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 13$  TeV. See their Fig. 3 (top) for limits on cross section times branching ratio for  $m_{H_2^0} = 0.5\text{--}4$  TeV. NODE=S055H2O;LINKAGE=CA
- 143 KHACHATRYAN 16P search for gluon fusion production of an  $H_2^0$  decaying to  $HH \rightarrow b\bar{b}\tau^+\tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 8 (lower right) for cross section limits for  $m_{H_2^0} = 260\text{--}350$  GeV. NODE=S055H2O;LINKAGE=X
- 144 KHACHATRYAN 16P search for gluon fusion production of an  $A^0$  decaying to  $ZH \rightarrow \ell^+ \ell^- \tau^+ \tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 10 for cross section limits for  $m_{H_2^0} = 220\text{--}350$  GeV. NODE=S055H2O;LINKAGE=Y
- 145 AAD 15BK search for production of a heavy  $H_2^0$  decaying to  $HH$  in the final state  $b\bar{b}b\bar{b}$  in  $19.5 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 14(c) for  $\sigma(H_2^0) \text{B}(H_2^0 \rightarrow HH)$  for  $m_{H_2^0} = 500\text{--}1500$  GeV with  $\Gamma_{H_2^0} = 1$  GeV. NODE=S055H2O;LINKAGE=M
- 146 AAD 15BZ search for the decay  $H \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- \tau^+ \tau^-$  ( $m_H = 125$  GeV) in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 6 for limits on cross section times branching ratio for  $m_{A^0} = 3.7\text{--}50$  GeV. NODE=S055H2O;LINKAGE=H
- 147 AAD 15BZ search for a state  $H_2^0$  via the decay  $H_2^0 \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- \tau^+ \tau^-$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 6 for limits on cross section times branching ratio for  $m_{H_2^0} = 100\text{--}500$  GeV and  $m_{A^0} = 5$  GeV. NODE=S055H2O;LINKAGE=I
- 148 AAD 15CE search for production of a heavy  $H_2^0$  decaying to  $HH$  in the final states  $b\bar{b}\tau^+\tau^-$  and  $\gamma\gamma WW^*$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV and combine with data from AAD 15H and AAD 15BK. A limit  $\sigma(H_2^0) \text{B}(H_2^0 \rightarrow HH) < 2.1\text{--}0.011$  pb (95% CL) is given for  $m_{H_2^0} = 260\text{--}1000$  GeV. See their Fig. 6. NODE=S055H2O;LINKAGE=O
- 149 AAD 15H search for production of a heavy  $H_2^0$  decaying to  $HH$  in the final state  $\gamma\gamma b\bar{b}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. A limit of  $\sigma(H_2^0) \text{B}(H_2^0 \rightarrow HH) < 3.5\text{--}0.7$  pb is given for  $m_{H_2^0} = 260\text{--}500$  GeV at 95% CL. See their Fig. 3. NODE=S055H2O;LINKAGE=N
- 150 AAD 15S search for production of  $A^0$  decaying to  $ZH \rightarrow \ell^+ \ell^- b\bar{b}$ ,  $\nu\bar{\nu}b\bar{b}$  and  $\ell^+ \ell^- \tau^+ \tau^-$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 3 for cross section limits for  $m_{A^0} = 200\text{--}1000$  GeV. NODE=S055H2O;LINKAGE=J
- 151 KHACHATRYAN 15AW search for production of a heavy state  $H_2^0$  of an electroweak singlet extension of the Standard Model via the decays of  $H_2^0$  to  $W^+ W^-$  and  $ZZ$  in up to  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7$  TeV and up to  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8$  TeV in the range  $m_{H_2^0} = 145\text{--}1000$  GeV. See their Figs. 8 and 9 for limits in the parameter space of the model. NODE=S055H2O;LINKAGE=Q
- 152 KHACHATRYAN 15BB search for production of a resonance  $H$  decaying to  $\gamma\gamma$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 7 for limits on cross section times branching ratio for  $m_H = 150\text{--}850$  GeV. NODE=S055H2O;LINKAGE=L
- 153 KHACHATRYAN 15N search for production of  $A^0$  decaying to  $ZH \rightarrow \ell^+ \ell^- b\bar{b}$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 3 for limits on cross section times branching ratios for  $m_{A^0} = 225\text{--}600$  GeV. NODE=S055H2O;LINKAGE=G
- 154 KHACHATRYAN 15O search for production of a high-mass narrow resonance  $A^0$  decaying to  $ZH \rightarrow q\bar{q}\tau^+\tau^-$  in  $19.7 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 6 for limits on cross section times branching ratios for  $m_{A^0} = 800\text{--}2500$  GeV. NODE=S055H2O;LINKAGE=F
- 155 KHACHATRYAN 15R search for a narrow scalar resonance decaying to  $HH \rightarrow b\bar{b}b\bar{b}$  in  $17.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8$  TeV. See their Fig. 5 (top) for limits on cross section times branching ratios for  $m_{H_2^0} = 0.27\text{--}1.1$  TeV. NODE=S055H2O;LINKAGE=GA

- 156 AAD 14AP search for a second  $H$  state decaying to  $\gamma\gamma$  in addition to the state at about 125 GeV in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 4 for limits on cross section times branching ratio for  $m_H = 65\text{--}600 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=C
- 157 AAD 14M search for the decay cascade  $H_2^0 \rightarrow H^\pm W^\mp \rightarrow HW^\pm W^\mp$ ,  $H$  decaying to  $b\bar{b}$  in  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Table III for limits on cross section times branching ratio for  $m_{H_2^0} = 325\text{--}1025 \text{ GeV}$  and  $m_{H^\pm} = 225\text{--}925 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=D
- 158 CHATRCHYAN 14G search for a second  $H$  state decaying to  $WW^{(*)}$  in addition to the observed signal at about 125 GeV using  $4.9 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.4 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Fig. 21 (right) for cross section limits in the mass range 110–600 GeV.  
NODE=S055H2O;LINKAGE=B
- 159 KHACHATRYAN 14P search for a second  $H$  state decaying to  $\gamma\gamma$  in addition to the observed signal at about 125 GeV using  $5.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$  and  $19.7 \text{ fb}^{-1}$  at  $E_{\text{cm}} = 8 \text{ TeV}$ . See their Figs. 27 and 28 for cross section limits in the mass range 110–150 GeV.  
NODE=S055H2O;LINKAGE=A
- 160 AALTONEN 13P search for production of a heavy Higgs boson  $H'^0$  that decays into a charged Higgs boson  $H^\pm$  and a lighter Higgs boson  $H$  via the decay chain  $H'^0 \rightarrow H^\pm W^\mp$ ,  $H^\pm \rightarrow W^\pm H$ ,  $H \rightarrow b\bar{b}$  in the final state  $\ell\nu$  plus 4 jets in  $8.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 4 for limits on cross section times branching ratio in the  $m_{H^\pm} - m_{H'^0}$  plane for  $m_H = 126 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=EN
- 161 CHATRCHYAN 13BJ search for  $H$  production in the decay chain  $H \rightarrow A^0 A^0$ ,  $A^0 \rightarrow \mu^+ \mu^-$  in  $5.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $E_{\text{cm}} = 7 \text{ TeV}$ . See their Fig. 2 for limits on cross section times branching ratio.  
NODE=S055H2O;LINKAGE=AT
- 162 AALTONEN 11P search in  $2.7 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$  for the decay chain  $t \rightarrow bH^+$ ,  $H^+ \rightarrow W^+ A^0$ ,  $A^0 \rightarrow \tau^+ \tau^-$  with  $m_{A^0}$  between 4 and 9 GeV. See their Fig. 4 for limits on  $B(t \rightarrow bH^+)$  for  $90 < m_{H^+} < 160 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=A5
- 163 ABBIENDI 10 search for  $e^+ e^- \rightarrow ZH$  with the decay chain  $H \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0$ ,  $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 + (\gamma \text{ or } Z^*)$ , when  $\tilde{\chi}_1^0$  and  $\tilde{\chi}_2^0$  are nearly degenerate. For a mass difference of 2 (4) GeV, a lower limit on  $m_H$  of 108.4 (107.0) GeV (95% CL) is obtained for SM  $ZH$  cross section and  $B(H \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0) = 1$ .  
NODE=S055H2O;LINKAGE=IE
- 164 SCHAEEL 10 search for the process  $e^+ e^- \rightarrow HZ$  followed by the decay chain  $H \rightarrow A^0 A^0 \rightarrow \tau^+ \tau^- \tau^+ \tau^-$  with  $Z \rightarrow \ell^+ \ell^-$ ,  $\nu\bar{\nu}$  at  $E_{\text{cm}} = 183\text{--}209 \text{ GeV}$ . For a  $HZZ$  coupling equal to the SM value,  $B(H \rightarrow A^0 A^0) = B(A^0 \rightarrow \tau^+ \tau^-) = 1$ , and  $m_{A^0} = 4\text{--}10 \text{ GeV}$ ,  $m_H$  up to 107 GeV is excluded at 95% CL.  
NODE=S055H2O;LINKAGE=SC
- 165 ABAZOV 09V search for  $H$  production followed by the decay chain  $H \rightarrow A^0 A^0 \rightarrow \mu^+ \mu^- \mu^+ \mu^-$  or  $\mu^+ \mu^- \tau^+ \tau^-$  in  $4.2 \text{ fb}^{-1}$  of  $p\bar{p}$  collisions at  $E_{\text{cm}} = 1.96 \text{ TeV}$ . See their Fig. 3 for limits on  $\sigma(H) \cdot B(H \rightarrow A^0 A^0)$  for  $m_{A^0} = 3.6\text{--}19 \text{ GeV}$ .  
NODE=S055H2O;LINKAGE=VO
- 166 ABBIENDI 05A search for  $e^+ e^- \rightarrow H_1^0 A^0$  in general Type-II two-doublet models, with decays  $H_1^0, A^0 \rightarrow q\bar{q}, g\bar{g}, \tau^+ \tau^-$ , and  $H_1^0 \rightarrow A^0 A^0$ .  
NODE=S055H2O;LINKAGE=AN
- 167 ABBIENDI 04K search for  $e^+ e^- \rightarrow HZ$  with  $H$  decaying to two jets of any flavor including  $g\bar{g}$ . The limit is for SM production cross section with  $B(H \rightarrow jj) = 1$ .  
NODE=S055H2O;LINKAGE=AE
- 168 ABDALLAH 04 consider the full combined LEP and LEP2 datasets to set limits on the Higgs coupling to  $W$  or  $Z$  bosons, assuming SM decays of the Higgs. Results in Fig. 26.  
NODE=S055H2O;LINKAGE=AD
- 169 ACHARD 04B search for  $e^+ e^- \rightarrow HZ$  with  $H$  decaying to  $b\bar{b}$ ,  $c\bar{c}$ , or  $g\bar{g}$ . The limit is for SM production cross section with  $B(H \rightarrow jj) = 1$ .  
NODE=S055H2O;LINKAGE=AR
- 170 ACHARD 04F search for  $H$  with anomalous coupling to gauge boson pairs in the processes  $e^+ e^- \rightarrow H\gamma$ ,  $e^+ e^- H$ ,  $HZ$  with decays  $H \rightarrow f\bar{f}, \gamma\gamma, Z\gamma$ , and  $W^* W$  at  $E_{\text{cm}} = 189\text{--}209 \text{ GeV}$ . See paper for limits.  
NODE=S055H2O;LINKAGE=AA
- 171 ABBIENDI 03F search for  $H \rightarrow$  anything in  $e^+ e^- \rightarrow HZ$ , using the recoil mass spectrum of  $Z \rightarrow e^+ e^-$  or  $\mu^+ \mu^-$ . In addition, it searched for  $Z \rightarrow \nu\bar{\nu}$  and  $H \rightarrow e^+ e^-$  or photons. Scenarios with large width or continuum  $H$  mass distribution are considered. See their Figs. 11–14 for the results.  
NODE=S055H2O;LINKAGE=A3
- 172 ABBIENDI 03G search for  $e^+ e^- \rightarrow H_1^0 Z$  followed by  $H_1^0 \rightarrow A^0 A^0$ ,  $A^0 \rightarrow c\bar{c}, g\bar{g}$ , or  $\tau^+ \tau^-$  in the region  $m_{H_1^0} = 45\text{--}86 \text{ GeV}$  and  $m_{A^0} = 2\text{--}11 \text{ GeV}$ . See their Fig. 7 for the limits.  
NODE=S055H2O;LINKAGE=AI
- 173 Search for associated production of a  $\gamma\gamma$  resonance with a  $Z$  boson, followed by  $Z \rightarrow q\bar{q}, \ell^+ \ell^-$ , or  $\nu\bar{\nu}$ , at  $E_{\text{cm}} \leq 209 \text{ GeV}$ . The limit is for a  $H$  with SM production cross section and  $B(H \rightarrow f\bar{f})=0$  for all fermions  $f$ .  
NODE=S055H2O;LINKAGE=HA
- 174 For  $B(H \rightarrow \gamma\gamma)=1$ ,  $m_H > 113.1 \text{ GeV}$  is obtained.  
NODE=S055H2O;LINKAGE=LH
- 175 HEISTER 02M search for  $e^+ e^- \rightarrow HZ$ , assuming that  $H$  decays to  $q\bar{q}, g\bar{g}$ , or  $\tau^+ \tau^-$  only. The limit assumes SM production cross section.  
NODE=S055H2O;LINKAGE=MH
- 176 ABBIENDI 01E search for neutral Higgs bosons in general Type-II two-doublet models, at  $E_{\text{cm}} \leq 189 \text{ GeV}$ . In addition to usual final states, the decays  $H_1^0, A^0 \rightarrow q\bar{q}, g\bar{g}$  are searched for. See their Figs. 15,16 for excluded regions.  
NODE=S055H2O;LINKAGE=EK
- 177 ACCIARRI 00R search for  $e^+ e^- \rightarrow H\gamma$  with  $H \rightarrow b\bar{b}, Z\gamma$ , or  $\gamma\gamma$ . See their Fig. 3 for limits on  $\sigma \cdot B$ . Explicit limits within an effective interaction framework are also given, for which the Standard Model Higgs search results are used in addition.  
NODE=S055H2O;LINKAGE=PE

- 178 ACCIARRI 00R search for the two-photon type processes  $e^+e^- \rightarrow e^+e^-H$  with  $H \rightarrow b\bar{b}$  or  $\gamma\gamma$ . See their Fig. 4 for limits on  $\Gamma(H \rightarrow \gamma\gamma) \cdot B(H \rightarrow \gamma\gamma \text{ or } b\bar{b})$  for  $m_H=70\text{--}170$  GeV.
- 179 GONZALEZ-GARCIA 98B use  $D\bar{D}$  limit for  $\gamma\gamma$  events with missing  $E_T$  in  $p\bar{p}$  collisions (ABBOTT 98) to constrain possible  $ZH$  or  $WH$  production followed by unconventional  $H \rightarrow \gamma\gamma$  decay which is induced by higher-dimensional operators. See their Figs. 1 and 2 for limits on the anomalous couplings.
- 180 KRAWCZYK 97 analyse the muon anomalous magnetic moment in a two-doublet Higgs model (with type II Yukawa couplings) assuming no  $H_1^0 ZZ$  coupling and obtain  $m_{H_1^0} \gtrsim 5$  GeV or  $m_{A^0} \gtrsim 5$  GeV for  $\tan\beta > 50$ . Other Higgs bosons are assumed to be much heavier.
- 181 ALEXANDER 96H give  $B(Z \rightarrow H\gamma) \times B(H \rightarrow q\bar{q}) < 1.4 \times 10^{-5}$  (95%CL) and  $B(Z \rightarrow H\gamma) \times B(H \rightarrow b\bar{b}) < 0.7\text{--}2 \times 10^{-5}$  (95%CL) in the range  $20 < m_H < 80$  GeV.

NODE=S055H2O;LINKAGE=PF

NODE=S055H2O;LINKAGE=W

NODE=S055H2O;LINKAGE=U

NODE=S055H2O;LINKAGE=O2

## Electroweak Constraints on the Standard Model Higgs Boson Mass

Here we list constraints on the mass of the Higgs boson derived from fits to precision electroweak observables, assuming the minimal Standard Model with a doublet Higgs field and three generations of fermions.

NODE=S055HEW

NODE=S055HEW

VALUE (GeV)	DOCUMENT ID	TECN
<b><math>90^{+21}_{-18}</math></b>	<sup>1</sup> HALLER	18 RVUE
• • • We do not use the following data for averages, fits, limits, etc. • • •		
$91^{+30}_{-23}$	<sup>2</sup> BAAK	12 RVUE
$94^{+25}_{-22}$	<sup>3</sup> BAAK	12A RVUE
$91^{+31}_{-24}$	<sup>4</sup> ERLER	10A RVUE
$129^{+74}_{-49}$	<sup>5</sup> LEP-SLC	06 RVUE

NODE=S055HEW

<sup>1</sup> HALLER 18 make Standard Model fits to  $Z$  and neutral current parameters,  $m_t$ ,  $m_W$ , and  $\Gamma_W$  measurements available in 2018. The direct mass measurement at the LHC is not used in the fit.

NODE=S055HEW;LINKAGE=W

<sup>2</sup> BAAK 12 make Standard Model fits to  $Z$  and neutral current parameters,  $m_t$ ,  $m_W$ , and  $\Gamma_W$  measurements available in 2010 (using also preliminary data). The quoted result is obtained from a fit that does not include the limit from the direct Higgs searches. The result including direct search data from LEP2, the Tevatron and the LHC is  $120^{+12}_{-5}$  GeV.

NODE=S055HEW;LINKAGE=BA

<sup>3</sup> BAAK 12A make Standard Model fits to  $Z$  and neutral current parameters,  $m_t$ ,  $m_W$ , and  $\Gamma_W$  measurements available in 2012 (using also preliminary data). The quoted result is obtained from a fit that does not include the measured mass value of the signal observed at the LHC and also no limits from direct Higgs searches.

NODE=S055HEW;LINKAGE=BK

<sup>4</sup> ERLER 10A makes Standard Model fits to  $Z$  and neutral current parameters,  $m_t$ ,  $m_W$  measurements available in 2009 (using also preliminary data). The quoted result is obtained from a fit that does not include the limits from the direct Higgs searches. With direct search data from LEP2 and Tevatron added to the fit, the 90% CL (99% CL) interval is 115–148 (114–197) GeV.

NODE=S055HEW;LINKAGE=ER

<sup>5</sup> LEP-SLC 06 make Standard Model fits to  $Z$  parameters from LEP/SLC and  $m_t$ ,  $m_W$ , and  $\Gamma_W$  measurements available in 2005 with  $\Delta\alpha_{\text{had}}^{(5)}(m_Z) = 0.02758 \pm 0.00035$ . The 95% CL limit is 285 GeV.

NODE=S055HEW;LINKAGE=LE

## SEARCHES FOR NEUTRAL HIGGS BOSONS REFERENCES

NODE=S055

HAYRAPETY...	25A	PL B860 139067	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=63168
AAD	24A	PL B848 138394	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62571
AAD	24AB	JHEP 2402 197	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62838
AAD	24AP	JHEP 2408 013	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62877
AAD	24BQ	PR D110 052013	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63081
AAD	24BV	PL B858 139007	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63118
AAD	24BW	JHEP 2410 104	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63127
AAD	24BX	JHEP 2410 130	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63128
AAD	24CA	JHEP 2411 047	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63135
AAD	24CB	JHEP 2411 097	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63136
AAD	24CD	JHEP 2411 126	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63139
AAD	24CF	JHEP 2412 126	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63143
AAD	24CI	EPJ C84 1102	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63151
AAD	24CK	SCIB 69 3005	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=63161
AAD	24H	PRL 132 231801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62731
AAD	24K	PL B850 138536	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62749
ADACHI	24F	PR D109 112015	I. Adachi <i>et al.</i>	(BELLE II Collab.)	REFID=62814
HAYRAPETY...	24AA	JHEP 2406 097	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62864
HAYRAPETY...	24AE	JHEP 2407 293	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62876
HAYRAPETY...	24AF	EPJ C84 493	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62891
HAYRAPETY...	24AI	PR D110 032007	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62932

HAYRAPETY...	24AJ	JHEP	2408	215	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62948
HAYRAPETY...	24AZ	JHEP	2412	172	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=63144
HAYRAPETY...	24H	PL	B850	138478	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62747
HAYRAPETY...	24I	PL	B852	138582	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62751
TUMASYAN	24A	PR	D110	012013	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62823
TUMASYAN	24B	JHEP	2405	316	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62862
AAD	23AD	EPJ	C83	519	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62169
AAD	23AJ	EPJ	C83	603	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62178
AAD	23BD	JHEP	2307	199	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62391
AAD	23BE	JHEP	2307	200	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62392
AAD	23BG	JHEP	2307	203	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62394
AAD	23BW	PR	D108	092007	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62501
AAD	23BX	JHEP	2309	189	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62532
AAD	23CA	JHEP	2310	009	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62537
AAD	23CR	JHEP	2312	081	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62616
AAD	23O	JHEP	2306	016	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62136
AAD	23R	JHEP	2307	155	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62140
AAD	23U	JHEP	2307	125	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62145
AAD	23Z	JHEP	2307	040	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=62161
ADACHI	23A	PRL	130	181803	I. Adachi <i>et al.</i>	(BELLE II Collab.)	REFID=62218
HAYRAPETY...	23C	PR	D108	072004	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62489
HAYRAPETY...	23G	JHEP	2312	070	A. Hayrapetyan <i>et al.</i>	(CMS Collab.)	REFID=62615
TUMASYAN	23	PL	B842	137392	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62101
TUMASYAN	23AR	PRL	131	101801	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62443
TUMASYAN	23M	JHEP	2307	148	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62141
TUMASYAN	23O	JHEP	2307	095	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62150
TUMASYAN	23S	JHEP	2307	073	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=62157
AAD	22A	PR	D105	012006	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61542
AAD	22D	PL	B829	137066	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61705
AAD	22F	PR	D105	092002	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61741
AAD	22I	JHEP	2201	063	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61774
AAD	22J	JHEP	2203	041	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61779
AAD	22P	JHEP	2208	104	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61818
AAD	22Y	PR	D106	052001	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61938
ABLIKIM	22H	PR	D105	012008	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=61643
ABRATENKO	22A	PR	D106	092006	P. Abratenko <i>et al.</i>	(MicroBooNE Collab.)	REFID=61956
JIA	22	PRL	128	081804	S. Jia <i>et al.</i>	(BELLE Collab.)	REFID=61657
TUMASYAN	22AK	PL	B835	137566	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61936
TUMASYAN	22D	PR	D105	032008	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61720
AAD	21AF	EPJ	C81	332	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61333
AAD	21AI	EPJ	C81	396	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61337
AAD	21AY	PL	B822	136651	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61547
AAD	21AZ	JHEP	2110	013	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61556
AAD	21BB	JHEP	2111	209	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61558
AAD	21BE	EPJ	C81	860	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61569
ABRATENKO	21	PRL	127	151803	P. Abratenko <i>et al.</i>	(MicroBooNE Collab.)	REFID=61500
SIRUNYAN	21A	EPJ	C81	13	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60775
Also		EPJ	C81	333 (errat.)	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=61322
SLAVICH	21	EPJ	C81	450	P. Slavich <i>et al.</i>	(CMS Collab.)	REFID=62641
TUMASYAN	21F	JHEP	2111	057	A. Tumasyan <i>et al.</i>	(CMS Collab.)	REFID=61562
AAD	20	PR	D101	012002	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60046
AAD	20AA	PRL	125	051801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60610
AAD	20AE	PRL	125	221802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60697
AAD	20AI	PR	D102	112006	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60742
AAD	20AO	JHEP	2011	163	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60760
AAD	20C	PL	B800	135103	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60147
AAD	20L	PR	D102	032004	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60347
AAD	20X	JHEP	2007	108	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=60511
Also		JHEP	2101	145 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61275
Also		JHEP	2105	207 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=61301
AAIJ	20AL	JHEP	2010	156	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=60754
SIRUNYAN	20	PL	B800	135087	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60146
SIRUNYAN	20AA	JHEP	2003	055	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60476
SIRUNYAN	20AC	JHEP	2003	065	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60478
SIRUNYAN	20AD	JHEP	2003	103	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60480
SIRUNYAN	20AF	JHEP	2004	171	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60489
SIRUNYAN	20AP	JHEP	2008	139	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60527
SIRUNYAN	20Y	JHEP	2003	034	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60474
SIRUNYAN	20Z	JHEP	2003	051	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60475
AABOUD	19A	JHEP	1901	030	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59370
AABOUD	19AG	PL	B790	1	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59740
AABOUD	19AI	PL	B793	499	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59755
AABOUD	19O	JHEP	1904	092	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59672
AABOUD	19T	JHEP	1905	124	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59684
AABOUD	19V	JHEP	1905	142	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59686
AABOUD	19Y	JHEP	1907	117	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59698
AALTONEN	19	PR	D99	052001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=59838
BAGNASCHI	19	EPJ	C79	617	E. Bagnaschi <i>et al.</i>	(CMS Collab.)	REFID=62642
SIRUNYAN	19	PL	B788	7	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59412
SIRUNYAN	19AE	JHEP	1905	210	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59690
SIRUNYAN	19AN	EPJ	C79	280	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59716
SIRUNYAN	19AV	EPJ	C79	564	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59733
SIRUNYAN	19B	PR	D99	012005	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59516
SIRUNYAN	19BB	PL	B793	320	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59754
SIRUNYAN	19BD	PL	B795	398	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59759
SIRUNYAN	19BE	PRL	122	121803	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59768
SIRUNYAN	19BQ	PL	B796	131	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59930
SIRUNYAN	19CR	PL	B798	134992	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=60137
SIRUNYAN	19H	JHEP	1901	040	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59630
AABOUD	18AA	PR	D98	032015	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58981
AABOUD	18AG	PL	B782	750	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59030
AABOUD	18AH	PL	B783	392	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59037
AABOUD	18AI	JHEP	1803	174	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59080
Also		JHEP	1811	051 (errat.)	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59360
AABOUD	18AP	JHEP	1806	166	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59107
AABOUD	18BF	EPJ	C78	293	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59172
AABOUD	18BU	EPJ	C78	1007	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59333
AABOUD	18BX	JHEP	1810	031	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59349
AABOUD	18CC	JHEP	1811	051 (errat.)	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59360

AABOUD	18CE	JHEP	1812	039	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59369
AABOUD	18CQ	PRL	121	191801	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59529
AABOUD	18CW	JHEP	1811	040	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=59567
AABOUD	18F	PL	B777	91	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58626
AABOUD	18G	JHEP	1801	055	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58788
AAIJ	18AM	EPJ	C78	1008	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59334
AAIJ	18AQ	JHEP	1809	147	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=59345
HALLER	18	EPJ	C78	675	J. Haller <i>et al.</i>	(Gfitter Group)	REFID=59193
SIRUNYAN	18A	PL	B778	101	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58627
SIRUNYAN	18AF	PL	B781	244	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59016
SIRUNYAN	18BA	JHEP	1806	127	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59111
Also		JHEP	1903	128 (errat.)	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59661
SIRUNYAN	18BP	JHEP	1808	113	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59143
SIRUNYAN	18CW	JHEP	1808	152	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59249
SIRUNYAN	18CX	JHEP	1809	007	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59250
SIRUNYAN	18DK	JHEP	1809	148	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59346
SIRUNYAN	18DT	PL	B785	462	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59403
SIRUNYAN	18DU	PR	D98	092001	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59469
SIRUNYAN	18ED	JHEP	1811	172	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59566
SIRUNYAN	18EE	JHEP	1811	018	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=59568
SIRUNYAN	18F	JHEP	1801	054	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58787
AABOUD	17	PL	B764	11	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57699
AABOUD	17AN	PRL	119	191803	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58276
AABOUD	17AP	PL	B775	105	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58296
AABOUD	17AW	JHEP	1710	112	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=58335
KHACHATRYAN...	17AZ	JHEP	1710	076	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=58336
KHACHATRYAN...	17D	JHEP	1701	076	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57750
KHACHATRYAN...	17R	PL	B767	147	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57884
SIRUNYAN	17AX	JHEP	1711	010	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58348
SIRUNYAN	17CN	PR	D96	072004	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58658
SIRUNYAN	17Y	PL	B772	363	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)	REFID=58233
AABOUD	16AA	EPJ	C76	585	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57677
AABOUD	16AB	EPJ	C76	605	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57678
AABOUD	16AE	JHEP	1609	173	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57729
AABOUD	16H	JHEP	1609	001	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57352
AABOUD	16I	PR	D94	052002	M. Aaboud <i>et al.</i>	(ATLAS Collab.)	REFID=57426
AAD	16AX	EPJ	C76	45	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57360
AAD	16C	JHEP	1601	032	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57024
AAD	16L	EPJ	C76	210	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57156
AALTONEN	16C	PR	D93	112010	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=57296
ABLIKIM	16E	PR	D93	052005	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=57265
KHACHATRYAN...	16A	PL	B752	221	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56961
KHACHATRYAN...	16BG	EPJ	C76	371	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57376
KHACHATRYAN...	16BQ	PR	D94	052012	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57433
KHACHATRYAN...	16F	JHEP	1601	079	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57022
KHACHATRYAN...	16M	PRL	117	051802	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57196
KHACHATRYAN...	16P	PL	B755	217	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57209
KHACHATRYAN...	16W	PL	B758	296	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57228
KHACHATRYAN...	16Z	PL	B759	369	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=57236
AAD	15BD	EPJ	C75	337	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56671
AAD	15BH	EPJ	C75	299	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56675
Also		EPJ	C75	408 (errat.)	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=57177
AAD	15BK	EPJ	C75	412	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56678
AAD	15BZ	PR	D92	052002	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56879
AAD	15CE	PR	D92	092004	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56909
AAD	15H	PRL	114	081802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56440
AAD	15S	PL	B744	163	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56504
KHACHATRYAN...	15AW	JHEP	1510	144	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56862
KHACHATRYAN...	15AY	JHEP	1511	071	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56866
KHACHATRYAN...	15BB	PL	B750	494	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56930
KHACHATRYAN...	15N	PL	B748	221	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56527
KHACHATRYAN...	15O	PL	B748	255	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56528
KHACHATRYAN...	15R	PL	B749	560	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56531
LEES	15H	PR	D91	071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=56561
AAD	14AP	PRL	113	171801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56113
AAD	14AW	JHEP	1411	056	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56169
AAD	14BA	JHEP	1411	088	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=56202
AAD	14M	PR	D89	032002	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55784
AAD	14O	PRL	112	201802	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55849
CHATRCHYAN	14B	EPJ	C74	2980	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55703
CHATRCHYAN	14G	JHEP	1401	096	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55713
KHACHATRYAN...	14M	JHEP	1410	160	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56160
KHACHATRYAN...	14P	EPJ	C74	3076	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56183
KHACHATRYAN...	14Q	PR	D90	112013	V. Khachatryan <i>et al.</i>	(CMS Collab.)	REFID=56287
AAD	13AG	PL	B721	32	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55073
AAD	13AT	NJP	15	043009	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=55271
AAD	13O	JHEP	1302	095	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54933
AAIJ	13T	JHEP	1305	132	R. Aaij <i>et al.</i>	(LHCb Collab.)	REFID=55030
AALTONEN	13K	PR	D88	052012	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=55190
AALTONEN	13L	PR	D88	052013	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=55191
AALTONEN	13M	PR	D88	052014	T. Aaltonen <i>et al.</i>	(CDF and D0 Collabs.)	REFID=55192
AALTONEN	13P	PRL	110	121801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=55197
ABAZOV	13G	PR	D88	052006	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55184
ABAZOV	13H	PR	D88	052007	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55185
ABAZOV	13I	PR	D88	052008	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55186
ABAZOV	13J	PR	D88	052009	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55187
ABAZOV	13L	PR	D88	052011	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=55189
CARENA	13	EPJ	C73	2552	M. Carena <i>et al.</i>		REFID=55673
CHATRCHYAN	13AG	PL	B722	207	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55074
CHATRCHYAN	13AL	PL	B725	36	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55091
CHATRCHYAN	13BJ	PL	B726	564	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=55435
LEES	13C	PR	D87	031102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=54949

LEES	13L	PR D88 031701	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55167
LEES	13R	PR D88 071102	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=55451
AAD	12AI	PL B716 1	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54198
AAD	12AQ	PRL 108 251801	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54226
AAD	12N	EPJ C72 2157	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54130
AALTONEN	12AB	PR D85 092001	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=54367
AALTONEN	12AN	PL B717 173	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=54602
AALTONEN	12AQ	PR D86 091101	T. Aaltonen <i>et al.</i>	(CDF and D0 Collabs.)	REFID=54701
AALTONEN	12U	PR D85 012007	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=54360
AALTONEN	12X	PR D85 032005	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=54363
ABAZOV	12G	PL B710 569	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=54162
ABLIKIM	12	PR D85 092012	M. Ablikim <i>et al.</i>	(BESIII Collab.)	REFID=54265
BAAK	12	EPJ C72 2003	M. Baak <i>et al.</i>	(Gitter Group)	REFID=54138
BAAK	12A	EPJ C72 2205	M. Baak <i>et al.</i>	(Gitter Group)	REFID=54615
CHATRCHYAN	12AO	JHEP 1209 111	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54571
CHATRCHYAN	12C	JHEP 1203 081	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54087
CHATRCHYAN	12D	JHEP 1204 036	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54088
CHATRCHYAN	12E	PL B710 91	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54093
CHATRCHYAN	12G	PL B710 403	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54095
CHATRCHYAN	12H	PRL 108 111804	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54096
CHATRCHYAN	12I	JHEP 1203 040	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54097
CHATRCHYAN	12K	PL B713 68	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54178
CHATRCHYAN	12N	PL B716 30	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54181
CHATRCHYAN	12V	PRL 109 121801	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54253
AALTONEN	11P	PRL 107 031801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=16451
ABAZOV	11K	PL B698 97	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=16479
ABAZOV	11W	PRL 107 121801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=53816
ABOUZAID	11A	PRL 107 201803	E. Abouzaid <i>et al.</i>	(KTeV Collab.)	REFID=53842
DEL-AMO-SA...	11J	PRL 107 021804	P. del Amo Sanchez <i>et al.</i>	(BABAR Collab.)	REFID=16495
LEES	11H	PRL 107 021804	J.P. Lees <i>et al.</i>	(BABAR Collab.)	REFID=53877
ABBIENDI	10	PL B682 381	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=53163
ANDREAS	10	JHEP 1008 003	S. Andreas <i>et al.</i>	(DESY)	REFID=53623
ERLER	10A	PR D81 051301	J. Erler	(UNAM)	REFID=53389
HYUN	10	PRL 105 091801	H.J. Hyun <i>et al.</i>	(BELLE Collab.)	REFID=53372
SCHAEI	10	JHEP 1005 049	S. Schael <i>et al.</i>	(ALEPH Collab.)	REFID=53342
AALTONEN	09AB	PRL 103 061803	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=52933
AALTONEN	09AR	PRL 103 201801	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=53081
ABAZOV	09V	PRL 103 061801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52924
AUBERT	09P	PRL 103 181801	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=53062
AUBERT	09Z	PRL 103 081803	B. Aubert <i>et al.</i>	(BABAR Collab.)	REFID=52930
TUNG	09	PRL 102 051802	Y.C. Tung <i>et al.</i>	(KEK E391a Collab.)	REFID=52872
ABAZOV	08U	PRL 101 051801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52399
ABDALLAH	08B	EPJ C54 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=52480
Also		EPJ C56 165 (errat.)	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=52483
LOVE	08	PRL 101 151802	W. Love <i>et al.</i>	(CLEO Collab.)	REFID=52565
ABBIENDI	07	EPJ C49 457	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=51729
BESSON	07	PRL 98 052002	D. Besson <i>et al.</i>	(CLEO Collab.)	REFID=51620
SCHAEI	07	EPJ C49 439	S. Schael <i>et al.</i>	(ALEPH Collab.)	REFID=51728
LEP-SLC	06	PRPL 427 257	ALEPH, DELPHI, L3, OPAL, SLD and working groups		REFID=51219; ERROR=1; ERROR=2
SCHAEI	06B	EPJ C47 547	S. Schael <i>et al.</i>	(LEP Collabs.)	REFID=51390
ABBIENDI	05A	EPJ C40 317	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=50673
ABDALLAH	05D	EPJ C44 147	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=50861
ACHARD	05	PL B609 35	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=50476
ACOSTA	05Q	PR D72 072004	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=50890
PARK	05	PRL 94 021801	H.K. Park <i>et al.</i>	(FNAL HyperCP Collab.)	REFID=50480
ABBIENDI	04K	PL B597 11	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49983
ABBIENDI	04M	EPJ C37 49	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=50150
ABDALLAH	04	EPJ C32 145	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49681
ABDALLAH	04B	EPJ C32 475	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49843
ABDALLAH	04L	EPJ C35 313	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49971
ABDALLAH	04O	EPJ C38 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=50304
ACHARD	04B	PL B583 14	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49819
ACHARD	04F	PL B589 89	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49904
ABBIENDI	03F	EPJ C27 311	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49375
ABBIENDI	03G	EPJ C27 483	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49378
ACHARD	03C	PL B568 191	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49525
ABBIENDI	02D	EPJ C23 397	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48741
ABBIENDI	02F	PL B544 44	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48916
ACHARD	02C	PL B534 28	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48645
ACHARD	02H	PL B545 30	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48962
AKERROYD	02	PR D66 037702	A.G. Akeroyd <i>et al.</i>		REFID=48908
HEISTER	02	PL B526 191	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48539
HEISTER	02L	PL B544 16	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48913
HEISTER	02M	PL B544 25	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48914
ABBIENDI	01E	EPJ C18 425	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48070
ABREU	01F	PL B507 89	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=48071
AFFOLDER	01H	PR D64 092002	T. Affolder <i>et al.</i>	(CDF Collab.)	REFID=48392
BARATE	01C	PL B499 53	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=48046
ACCIARRI	00M	PL B485 85	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47719
ACCIARRI	00R	PL B489 102	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47759
ACCIARRI	00S	PL B489 115	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47760
BARATE	00L	PL B487 241	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47722
ABBIENDI	99E	EPJ C7 407	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=46641
ABBIENDI	99O	PL B464 311	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=47240
ABBOTT	99B	PRL 82 2244	B. Abbott <i>et al.</i>	(D0 Collab.)	REFID=46647
ABREU	99P	PL B458 431	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47040
ABBOTT	98	PRL 80 442	B. Abbott <i>et al.</i>	(D0 Collab.)	REFID=45800
ACKERSTAFF	98S	EPJ C5 19	K. Akerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46153
ACKERSTAFF	98Y	PL B437 218	K. Akerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46254
GONZALEZ...	98B	PR D57 7045	M.C. Gonzalez-Garcia, S.M. Lietti, S.F. Novaes		REFID=46198
KRAWCZYK	97	PR D55 6968	M. Krawczyk, J. Zochowski	(WARS)	REFID=45466
ALEXANDER	96H	ZPHY C71 1	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=44822
ABREU	95H	ZPHY C67 69	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44368
BALEST	95	PR D51 2053	R. Balest <i>et al.</i>	(CLEO Collab.)	REFID=44146
PICH	92	NP B388 31	A. Pich, J. Prades, P. Yepes	(CERN, CPPM)	REFID=43104
ANTREASYAN	90C	PL B251 204	D. Antreasyan <i>et al.</i>	(Crystal Ball Collab.)	REFID=41455