Double- $\beta$ Decay		
OMITTED FROM SUMMARY	TABLE	NODE=5076
For a discussion of the doubl of the review "14. Neutrino See the related review(s	e- $\beta$ decay, see sections 14.9.2 and 14.9.3 Masses, Mixing, and Oscillations." (3):	NODE=S076
Neutrino Masses, Mixing,	and Oscillations	

# Half-life $0\nu$ double- $\beta$ decay

In most cases the transitions (Z,A)  $\rightarrow$  (Z+2,A)  $+2e^-$  to the  $0^+$  ground state of the final nucleus are listed. We also list transitions that decrease the nuclear charge (2 $e^+$ ,  $e^+$  CC and double EC) and transitions to an excited state of the final nucleus (0 $_i^+$ , 2 $^+$ , and 2 $_i^+$ ). In the following Listings only the best or comparable limits for the half-lives of each transition are reported and only those with about  ${\rm T}_{1/2}>10^{23}$  years that are relevant for particle physics.

$t_{1/2}(10^{23} \text{ yr})$	CL%	SOTOPE	TRANSITION	METHOD	DOCUMENT ID		NODE=S076H0N
• • • We do no	ot use	e the follow	ving data for	averages, fits, limi	ts, etc. • • •		
> 3	90	<sup>134</sup> Xe		PandaX-4T	<sup>1</sup> YAN	24	
>2300	90	<sup>136</sup> Xe		KamLAND-Zen	<sup>2</sup> ABE	23	
> 830	90	<sup>76</sup> Ge		MAJORANA	<sup>3</sup> ARNQUIST	23	
> 2.1	90	<sup>100</sup> Mo	g.s. $\rightarrow 2_1^+$	CUPID-Mo	<sup>4</sup> AUGIER	23	
> 1.2	90	<sup>100</sup> Mo	g.s. $\rightarrow 0^{1}_{1}$	CUPID-Mo	<sup>4</sup> AUGIER	23	OCCUR=3
> 13	90	136 <sub>Xe</sub>	5 I	NEXT	<sup>5</sup> NOVELLA	23	
> 220	90	<sup>130</sup> Te		CUORE	<sup>6</sup> ADAMS	22A	
> 36	90	<sup>128</sup> Te		CUORE	<sup>7</sup> ADAMS	22B	
> 12	90	<sup>136</sup> Xe		XENON1T	<sup>8</sup> APRILE	22A	
> 18	90	<sup>100</sup> Mo		CUPID-Mo	<sup>9</sup> AUGIER	22	
> 46	90	<sup>82</sup> Se		CUPID-0	<sup>10</sup> AZZOLINI	22	
> 1.8	90	<sup>82</sup> Se	g.s. $\rightarrow 0^+_1$	CUPID-0	<sup>11</sup> AZZOLINI	22	OCCUR=2
> 3.0	90	<sup>82</sup> Se	g.s. $\rightarrow 2^{+}_{1}$	CUPID-0	<sup>12</sup> AZZOLINI	22	OCCUR=3
> 3.2	90	<sup>82</sup> Se	g.s. $\rightarrow 2^{1}_{2}$	CUPID-0	<sup>13</sup> AZZOLINI	22	OCCUR=4
> 59	90	<sup>130</sup> Te	g.s. $\rightarrow 0_1^2$	CUORE	<sup>14</sup> ADAMS	21A	
> 15	90	100 <sub>Mo</sub>	5 I	CUPID-Mo	<sup>15</sup> ARMENGAUD	21	
> 39.9	90	76 <sub>Ge</sub>	g.s. $\rightarrow 0^+_1$	MAJORANA-Dem	<sup>16</sup> ARNQUIST	21	OCCUR=4
> 21.2	90	76 <sub>Ge</sub>	g.s. $\rightarrow 2^{+}_{1}$	MAJORANA-Dem	<sup>17</sup> ARNQUIST	21	OCCUR=5
> 9.7	90	<sup>76</sup> Ge	g.s. $\rightarrow 2^{+}_{2}$	MAJORANA-Dem	<sup>18</sup> ARNQUIST	21	OCCUR=6
> 320	90	<sup>130</sup> Te	Ζ	CUORE	<sup>19</sup> ADAMS	20A	
>1800	90	<sup>76</sup> Ge		GERDA	<sup>20</sup> AGOSTINI	20B	OCCUR=2
> 14	90	<sup>130</sup> Te	g.s. $\rightarrow 0_1^+$	CUORE-0	<sup>21</sup> ALDUINO	19	
> 0.95	90	<sup>100</sup> Mo	- 1	AMoRE	<sup>22</sup> ALENKOV	19	
> 350	90	<sup>136</sup> Xe		EXO-200	<sup>23</sup> ANTON	19	
> 2.4	90	<sup>136</sup> Xe		PANDAX-II	<sup>24</sup> NI	19	
> 150	90	<sup>130</sup> Te		CUORE	<sup>25</sup> ALDUINO	18	
> 2.5	90	<sup>82</sup> Se		NEMO-3	<sup>26</sup> ARNOLD	18	
> 2.2	90	<sup>116</sup> Cd		AURORA	<sup>27</sup> BARABASH	18	
> 1.1	90	<sup>134</sup> Xe		EXO-200	<sup>28</sup> ALBERT	17C	OCCUR=5
> 1	90	<sup>110</sup> Cd		NEMO-3	<sup>29</sup> ARNOLD	17	OCCUR=2
> 40	90	<sup>130</sup> Te	1	CUORICINO	<sup>30</sup> ALDUINO	16	
> 260	90	<sup>130</sup> Xe	$g.s. \rightarrow 2^+_1$	KamLAND-Zen	<sup>31</sup> ASAKURA	16	
> 260	90	<sup>136</sup> Xe	$g.s. \rightarrow 2^+_2$	KamLAND-Zen	<sup>32</sup> ASAKURA	16	OCCUR=2
> 240	90	<sup>136</sup> Xe	$g.s.\!\to 0_1^+$	KamLAND-Zen	<sup>33</sup> ASAKURA	16	OCCUR=3
> 11	90	<sup>100</sup> Mo		NEMO-3	<sup>34</sup> ARNOLD	15	OCCUR=2
> 9.4	90	<sup>130</sup> Te	g.s. $\rightarrow 0^+_1$	CUORICINO	<sup>35</sup> ANDREOTTI	12	
> 0.58	90	<sup>48</sup> Ca	-	CaF <sub>2</sub> scint.	<sup>36</sup> UMEHARA	08	
> 0.89	90	<sup>100</sup> Mo	g.s. $\rightarrow 0^+_1$	NEMO-3	<sup>37</sup> ARNOLD	07	OCCUR=2
> 1.6	90	100 <sub>Mo</sub>	g.s. $\rightarrow 2^{+}$	NEMO-3	<sup>38</sup> ARNOLD	07	OCCUR=4
> 1.1	90	<sup>128</sup> Te		Cryog. det.	<sup>39</sup> ARNABOLDI	03	OCCUR=4
> 1.7	90	<sup>116</sup> Cd		<sup>116</sup> CdWO <sub>4</sub> scint.	<sup>40</sup> DANEVICH	03	OCCUR=2
> 157	90	<sup>76</sup> Ge		Enriched HPGe	<sup>41</sup> AALSETH	<b>02</b> B	

NODE=S076H0N

NODE=S076H0N

	4/20/2025 20:25 Page 2
<sup>1</sup> YAN 24 make use of 17.9 kg·y of <sup>134</sup> Xe isotope exposure in the PandaX-4T TPC, using natural xenon, to place a limit on the $0\nu\beta\beta$ decay half-life of <sup>134</sup> Xe.	NODE=S076H0N;LINKAGE=JB
<sup>2</sup> ABE 23 use the combined data set of the KamLAND-Zen 400 and 800 experiments, uti- lizing 745 kg of isotopically enriched xenon (90.9% <sup>136</sup> Xe), dissolved in liquid scintillator and an exposure of 970 kg·yr of <sup>136</sup> Xe, to derive this limit on $0\nu\beta\beta$ decay. A half-life sensitivity of $1.5 \times 10^{26}$ yr is reported.	NODE=S076H0N;LINKAGE=CB
<sup>3</sup> ARNQUIST 23 use the final data set of the MAJORANA DEMONSTRATOR experiment, operating enriched in <sup>76</sup> Ge detectors, to set this limit on the $0\nu\beta\beta$ half-life of <sup>76</sup> Ge. The exposure is 64.5 kg yr. A median sensitivity of $8.1 \times 10^{25}$ yr is reported.	NODE=S076H0N;LINKAGE=BB
<sup>4</sup> AUGIER 23 utilize the complete data, set collected by the CUPID-Mo bolometric calorimeter with particle ID and located at the LSM, to study various double beta decays of <sup>100</sup> Mo to excited states of the daughter nucleus. An exposure of 1.47 kg·yr of <sup>100</sup> Mo is available.	NODE=S076H0N;LINKAGE=DB
<sup>5</sup> NOVELLA 23 use data collected by the NEXT-White experiment to limit the $0\nu \beta\beta$ half-life of $^{136}$ Xe. The experiment contains 3.5 kg of enriched Xe and is based on a high-pressure gas TPC. Two different limits are reported, based on different data analysis approaches, $> 5.5 \times 10^{23}$ yr and $> 13 \times 10^{23}$ yr.	NODE=S076H0N;LINKAGE=HB
<sup>6</sup> ADAMS 22A use the CUORE TeO <sub>2</sub> experiment with an exposure of 288.8 kg·yr of <sup>130</sup> Te to place a limit on its $0\nu \beta\beta$ decay. The median sensitivity is reported as $280 \times 10^{23}$ yr. Superseeds ADAMS 20A.	NODE=S076H0N;LINKAGE=MA
<sup>7</sup> ADAMS 22B use the CUORE bolometric calorimeter to place a limit on the $0\nu\beta\beta$ decay	NODE=S076H0N;LINKAGE=PA
<sup>8</sup> APRILE 22A use 36.16 kg.yr of <sup>136</sup> Xe exposure of the XENON1T not enriched detector	NODE=S076H0N;LINKAGE=NA
<sup>9</sup> AUGIER 22 use the final data set of the CUPID-Mo cryogenic calorimeter, utilizing enriched Li <sub>2</sub> <sup>100</sup> MoO <sub>4</sub> and an isotope exposure of 1.47 kg·y, to place a limit on the $0\mu\beta\beta$ docsy balf life	NODE=S076H0N;LINKAGE=AB
$^{10}$ AZZOLINI 22 use the CUPID-0 scintillating cryogenic bolometer to set a limit on the $0\nu\beta\beta$ half-life of <sup>82</sup> Se. The analyzed isotope exposure is 8.82 kg·yr. A median sensitivity of $7 \times 10^{24}$ wire is reported. Supercodes AZZOLINI 10	NODE=S076H0N;LINKAGE=QA
<sup>11</sup> AZZOLINI 22 use CUPID-0 data with an isotope exposure of 8.82 kg·yr to set a limit	NODE=S076H0N;LINKAGE=RA
on the $0\nu\beta\beta$ decay to the first excited 0' state. <sup>12</sup> AZZOLINI 22 use CUPID-0 data with an isotope exposure of 8.82 kg yr to set a limit	NODE=S076H0N;LINKAGE=VA
on the $0\nu\beta\beta$ decay to the first excited 2 <sup>+</sup> state. <sup>13</sup> AZZOLINI 22 use CUPID-0 data with an isotope exposure of 8.82 kg·yr to set a limit	NODE=S076H0N;LINKAGE=XA
on the $0\nu\beta\beta$ decay to the second excited 2 <sup>+</sup> state. <sup>14</sup> ADAMS 21A et al. used 101.76 kg yr of <sup>130</sup> Te exposure of the CUORE (LNGS) bolomet-	NODE=S076H0N:LINKAGE=AA
ric detector to place a limit on the decay to the first excited state of $^{130}$ Xe, superseding ALDUINO 19 as the most restrictive bound on this particular decay.	
<sup>15</sup> ARMENGAUD 21 use the CUPID-Mo 4.2 kg array of enriched Li <sub>2</sub> <sup>100</sup> MoO <sub>4</sub> scintillating bolometers, with 1.17 kg yr exposure, to set this limit.	NODE=S076H0N;LINKAGE=IA
<sup>16</sup> ARNQUIST 21 use the MAJORANA demonstrator to set this limit for the $0\nu \beta\beta$ decay to the first excited $0^+$ state, with a 41.9 kg yr isotopic exposure. The median sensitivity	NODE=S076H0N;LINKAGE=EA
is $39.9 \times 10^{23}$ yr. <sup>17</sup> ARNOUIST 21 use the MAJORANA demonstrator to set this limit for the $0\nu$ $\beta\beta$ decay	
to the first excited $2^+$ state, with a 41.9 kg yr isotopic exposure. The median sensitivity	
<sup>18</sup> ARNQUIST 21 use the MAJORANA demonstrator to set this limit for the 0 $\nu$ $\beta\beta$ decay to	NODE=S076H0N;LINKAGE=HA
the second excited $2^+$ state, with a 41.9 kg yr isotopic exposure. The median sensitivity	
<sup>19</sup> ADAMS 20A use the CUORE detector to search for the $0\nu \beta\beta$ decay of <sup>130</sup> Te. The	NODE=S076H0N;LINKAGE=W
exposure was 372.5 kg·yr of TeO <sub>2</sub> corresponding to 103.6 kg·yr of $^{130}$ Te. The exclusion	
<sup>20</sup> AGOSTINI 20B present the final data set of the GERDA experiment, searching for $0\nu\beta\beta$ decay of <sup>76</sup> Ge with isotopically enriched, high resolution Ge detectors. A final exposure of 127.2 kg·yr is reported. The experiment reports the lowest background and longest half	NODE=S076H0N;LINKAGE=Y
sensitivity equals the limit. Supersedes AGOSTINI 19.	
<sup>21</sup> ALDUINO 19 use the combined data of the CUORICINO and CUORE-0 experiments to place a lower limit on the half life of the $0\nu \beta\beta$ decay of <sup>130</sup> Te to the first excited 0 <sup>+</sup> state of <sup>130</sup> Xe. Supersedes ANDREOTTI 12.	NODE=S076H0N;LINKAGE=S
<sup>22</sup> ALENKOV 19 report the $0\nu \beta\beta$ decay half-life limit based on the 52.1 kg d exposure of <sup>100</sup> Mo, of a a cryogenic dual heat and light detector in the Yangyang underground laboratory. The median sensitivity is $1.1 \times 10^{23}$ years	NODE=S076H0N;LINKAGE=P
<sup>23</sup> ANTON 19 uses he complete dataset of the EXO-200 detector to search for the $0\nu$ $\beta\beta$ docv. The expected is 234.1 kg vs. The median constituities is 5.0 × 10 <sup>25</sup> vs. Supervisites	NODE=S076H0N;LINKAGE=Q
ALBERT 18 and ALBERT 14B. <sup>24</sup> NI 19 use the PandaX-II dual phase TPC at CJPL to search for the $0\nu \beta\beta$ decay of 136Xe. The half-life limit 2.4 × 10 <sup>23</sup> yr is obtained from 22.2 kg yr exposure with a	NODE=S076H0N;LINKAGE=O

 $^{26}$  ARNOLD 18 use the NEMO-3 tracking detector to place a limit on the 0 $\nu\beta\beta$  decay of <sup>82</sup>Se. This is a slightly weaker limit than in BARABASH 11A, using the same detector. Supersedes ARNOLD 05A.

 $^{27}\,\rm BARABASH$  18 use 1.162 kg of  $^{116}\rm CdWO_4$  scintillating crystals to obtain this limit. Supersedes DANEVICH 03 with analogous source and is more sensitive than ARNOLD 17.

 $^{28}\,\text{ALBERT}$  17C uses the EXO-200 detector that contains 19.098  $\pm$  0.014% admixture of  $^{134}{\rm Xe}$  to search for the 0 $\nu$  and 2 $\nu$   $\beta\beta$  decay modes. The exposure is 29.6 kg year. The median sensitivity is  $1.9\times 10^{21}$  years.

 $^{29}$  ARNOLD 17 use the NEMO-3 tracking calorimeter, containing 410 g of enriched  $^{116}$ Cd exposed for 5.26 yr, to determine the half-life limit. Supersedes BARABASH 11A.

 $^{30}\,\text{ALDUINO}$  16 report result obtained with 9.8 kg y of data collected with the CUORE-0 bolometer, combined with data from the CUORICINO. Supersedes ALFONSO 15.

 $^{31}$ ASAKURA 16 use the KamLAND-Zen liquid scintillator calorimeter ( $^{136}$ Xe 89.5 kg yr) to place a limit on the  $0\nu\beta\beta$ -decay into the first excited state of the daughter nuclide.

 $^{32}$ ASAKURA 16 use the KamLAND-Zen liquid scintillator calorimeter ( $^{136}$ Xe 89.5 kg yr) to place a limit on the  $0\nu\beta\beta$ -decay into the second excited state of the daughter nuclide.

 $^{33}$ ASAKURA 16 use the KamLAND-Zen liquid scintillator calorimeter ( $^{136}$ Xe 89.5 kg yr) to place a limit on the  $0
u\beta\beta$ -decay into the third excited state of the daughter nuclide.

 $^{34}$  ARNOLD 15 use the NEMO-3 tracking calorimeter with 34.3 kg yr exposure to determine the limit of  $0
u\beta\beta$ -half life of  $^{100}$ Mo. Supersedes ARNOLD 2005A and BARABASH 11A.

 $^{35}$  ANDREOTTI 12 use high resolution TeO<sub>2</sub> bolometric calorimeter to search for the  $0\nu\beta\beta$ decay of  $^{130}\mathrm{Te}$  leading to the excited  $0^{1}_{+}$  state at 1793.5 keV.

 $^{36}\text{UMEHARA 08}$  use  $\text{CaF}_2$  scintillation calorimeter to search for double beta decay of  $^{48}\text{Ca.}$  Limit is significantly more stringent than quoted sensitivity:  $18\times10^{21}$  years.

 $^{37}$ Limit on 0 $\nu$ -decay to the first excited 0 $^+_1$ -state of daughter nucleus using NEMO-3 tracking calorimeter. Supersedes DASSIE 95.

 $^{38}$  Limit on 0 $\nu$ -decay to the first excited 2+-state of daughter nucleus using NEMO-3 tracking calorimeter.

 $^{39}$  Supersedes ALESSANDRELLO 00. Array of TeO\_2 crystals in high resolution cryogenic calorimeter. Some enriched in <sup>128</sup>Te. Ground state to ground state decay.

 $^{40}$ Limit on 0
uetaeta decay of  $^{116}$ Cd using enriched CdWO<sub>4</sub> scintillators. Supersedes DANEVICH 00.

<sup>41</sup>AALSETH 02B limit is based on 117 mol yr of data using enriched Ge detectors. Background reduction by means of pulse shape analysis is applied to part of the data set. Reported limit is slightly less restrictive than that in KLAPDOR-KLEINGROTHAUS 01 However, it excludes part of the allowed half-life range reported in KLAPDOR-KLEINGROTHAUS 01B for the same nuclide. The analysis has been criticized in KLAPDOR-KLEINGROTHAUS 04B. The criticism was addressed and disputed in AALSETH 04.

# Half-life measurements of the two-neutrino double- $\beta$ decay

The measured half-life values for the transitions (Z,A)  $\rightarrow$  (Z+2,A) + 2 $e^-$  + 2 $\overline{\nu}_{\rho}$  to the  $0^+$  ground state of the final nucleus are listed. We also list the transitions to an excited state of the final nucleus  $(0_i^+,$  etc.). We report only the measuremetnts with the smallest (or comparable) uncertainty for each transition.

$t_{1/2}(10^{20} \text{ yr})$	)			ISOTOPE TRANSITIO	NMETHOD	DOCUMENT ID			NODE=S076H2N
• • • We	do	not use	the followin	g data for averages	s, fits, limits, e	etc. • • •			
109	±	14	$\pm$ 5	$^{124}$ Xe $^{2 u}$ DEC	LZ	<sup>1</sup> AALBERS	24C	1	
20.22	$\pm$	0.18	$\pm$ 0.38	76 <sub>Ge</sub>	GERDA	<sup>2</sup> AGOSTINI	23		
1.11	+	$\begin{array}{c} 0.19\\ 0.14\end{array}$	$^+$ 0.17 $^-$ 0.15	$^{150}\text{Nd}~0^+ \rightarrow 0^+_1$	NEMO-3	<sup>3</sup> AGUERRE	23		
7.5	$\pm$	0.8	$^+$ 0.4 $^-$ 0.3	$^{100}$ Mo 0 $^+  ightarrow$ 0 $^+_1$	CUPID-Mo	<sup>4</sup> AUGIER	23		
0.0707	$7\pm$	0.0002	$2\pm$ 0.0011	100 <sub>Mo</sub>	CUPID-Mo	<sup>5</sup> AUGIER	23A		
0.869	±	0.005	$^{+\ 0.009}_{-\ 0.006}$	<sup>82</sup> Se	CUPID-0	<sup>6</sup> AZZOLINI	23A		OCCUR=2
21.6	+	6.2 4.0	$^{+}$ 4.0 $^{-}$ 2.9	<sup>136</sup> Xe	NEXT	<sup>7</sup> NOVELLA	23		
21900	±	700		<sup>128</sup> Te	CUORE	<sup>8</sup> ADAMS	22в		
110	$\pm$	20	$\pm 10$	<sup>124</sup> Xe	XENON1T	<sup>9</sup> APRILE	22A		
118	$\pm$	13	$\pm 14$	<sup>124</sup> Xe	XENONnT	<sup>10</sup> APRILE	22в		
23.4	+ -	0.8 4.6	$^+$ 3.0 $^-$ 1.7	<sup>136</sup> Xe	NEXT	$^{11}$ NOVELLA	22		
8.76	+	0.09 0.07	$^+$ 0.14 $^-$ 0.17	<sup>130</sup> Te	CUORE	<sup>12</sup> ADAMS	21		
180	+	50	+10	$^{124}$ Xe 2 $\nu$ DEC	XENON1T	<sup>13</sup> APRILE	19E		

NODE=S076H0N;LINKAGE=C

NODE=S076H0N;LINKAGE=J

NODE=S076H0N;LINKAGE=K NODE=S076H0N;LINKAGE=RC

NODE=S076H0N;LINKAGE=YB

NODE=S076H0N;LINKAGE=MB

NODE=S076H0N;LINKAGE=TA

NODE=S076H0N;LINKAGE=UA

NODE=S076H0N;LINKAGE=WA

NODE=S076H0N;LINKAGE=YA

NODE=S076H0N;LINKAGE=ND

NODE=S076H0N;LINKAGE=UM

NODE=S076H0N;LINKAGE=NR

NODE=S076H0N;LINKAGE=NO

NODE=S076H0N;LINKAGE=RO

NODE=S076H0N;LINKAGE=VB

NODE=S076H0N;LINKAGE=SH

NODE=S076H2N NODE=S076H2N

	0.0680	)±	0.0001	+	0.0038 0.0040	100 <sub>Mo</sub>	NEMO-3	<sup>14</sup> ARNOLD	19	
	0.939	±	0.017	±	0.058	<sup>82</sup> Se	NEMO-3	<sup>15</sup> ARNOLD	18	
	0.263	+	$0.011 \\ 0.012$			<sup>116</sup> Cd	AURORA	<sup>16</sup> BARABASH	18	
>	0.87					<sup>134</sup> Xe	EXO-200	<sup>17</sup> ALBERT	17C	OCCUR=2
	8.2	$\pm$	0.2	$\pm$	0.6	<sup>130</sup> Te	CUORE-0	<sup>18</sup> ALDUINO	17	
	0.274	±	0.004	$\pm$	0.018	<sup>116</sup> Cd	NEMO-3	<sup>19</sup> ARNOLD	17	
	0.64	+	0.07 0.06	+	0.12 0.09	<sup>48</sup> Ca	NEMO-3	<sup>20</sup> ARNOLD	16	
	0.0934	÷±	0.0022	2+	0.0062 0.0060	<sup>150</sup> Nd	NEMO-3	<sup>21</sup> ARNOLD	16A	
	19.26	±	0.94			<sup>76</sup> Ge	GERDA	<sup>22</sup> AGOSTINI	15A	
	0.0693	$\pm$	0.0004	ł		<sup>100</sup> Mo	NEMO-3	<sup>23</sup> ARNOLD	15	
	21.65	±	0.16	$\pm$	0.59	<sup>136</sup> Xe	EXO-200	<sup>24</sup> ALBERT	14	
	92	+ 5	55 26	±	13	<sup>78</sup> Kr	BAKSAN	<sup>25</sup> GAVRILYAK	13	
	23.8	±	0.2	$\pm$	1.4	<sup>136</sup> Xe	KamLAND-Z	<sup>26</sup> GANDO	12A	
	7.0	±	0.9	$\pm$	1.1	<sup>130</sup> Te	NEMO-3	<sup>27</sup> ARNOLD	11	
	0.235	±	0.014	$\pm$	0.016	<sup>96</sup> Zr	NEMO-3	<sup>28</sup> ARGYRIADES	10	
	6.9	+ -	1.0 0.8	±	0.7	$^{100}\text{Mo}~\text{O}^+ \rightarrow \text{O}^+_1$	Ge coinc.	<sup>29</sup> BELLI	10	
	5.7	+ -	1.3 0.9	±	0.8	$^{100}\text{Mo}~\text{O}^+ \rightarrow \text{O}^+_1$	NEMO-3	<sup>30</sup> ARNOLD	07	
	0.96	±	0.03	$\pm$	0.10	<sup>82</sup> Se	NEMO-3	<sup>31</sup> ARNOLD	05A	OCCUR=4
	0.29	+ -	0.04 0.03			<sup>116</sup> Cd	$CdWO_4$ sc.	<sup>32</sup> DANEVICH	03	
	1	DС -	046		41		1 DEC 1 20 1			

<sup>1</sup>AALBERS 24C report the observation of <sup>124</sup>Xe  $2\nu$ DEC. 1.39 kg·yr of isotopic exposure of the LZ dark matter experiment, collected during the first science run, were analyzed. The same capture fractions as used in APRILE 22A were assumed.

<sup>2</sup> AGOSTINI 23 report an updated value for the  $2\nu \beta\beta$  half-life of <sup>76</sup>Ge; the final result of the GERDA Phase II experiment. A subset of the data, corresponding to an exposure of exposure is 11.8 kg·yr, is utilized. This is one of the most precise measurements of  $2\nu$  $\beta\beta$  decay reported in the literature. An effective nuclear matrix element of  $0.101 \pm 0.001$ is derived from this result.

<sup>3</sup> AGUERRE 23 report the results of a 5.25 yr search for the  $2\nu \beta\beta$  decay to the exited  $0^+ \rightarrow 0^+_1$  state of the daughter nucleus, using the NEMO-3 tracking calorimeter. 36.6g of <sup>150</sup>Nd isotope were available for the measurement of this decay rate.

<sup>4</sup> AUGIER 23 utilize the complete data, set collected by the CUPID-Mo bolometric calorimeter with particle ID and located at the LSM, to measure the <sup>100</sup>Mo  $2\nu\beta\beta$  half-life to excited  $0^+_1$  state of the daughter nucleus. An exposure of 1.47 kg·yr of <sup>100</sup>Mo is available.

<sup>5</sup> AUGIER 23A use full data set collected by the CUPID-Mo experiment to derive an improved  $2\nu \beta\beta$  g.s. to g.s. half-life of <sup>100</sup>Mo. An exposure of 1.48 kg·yr of <sup>100</sup>Mo is utilized. Supersedes ARMENGAUD 20.

<sup>6</sup>AZZOLINI 23A report an improved measurement of the  $2\nu \beta\beta$  decay with an exposure of 8.82 kg·yr of <sup>82</sup>Se, collected with the CUPID-0 detector. Superseded AZZOLINI 19B.

<sup>7</sup> NOVELLA 23 used the NEXT-White experiment, with a fiducial mass of 3.5 kg of enriched xenon, to measure the  $2\nu \beta\beta$  g.s. to g.s. half-life of <sup>136</sup>Xe. The experiment is based on a high-pressure gas TPC. Supersedes NOVELLA 22.

<sup>8</sup> ADAMS 22B derive the  $2\nu\beta\beta$  half-life of <sup>128</sup>Te from data of the CUORE bolometric calorimeter and the half-live ratio for <sup>130</sup>Te / <sup>128</sup>Te reported in BERNATOWICZ 92.

 $^9$  APRILE 22A report an improved  $^{124}$ Xe  $_{2\nu}$  DEC half-life measurement for  $^{124}$ Xe, using data collected by the XENON1T detector with an isotopically not enriched Xe target. The analyzed  $^{124}$ Xe exposure is 0.87 kg·yr. The statistical significance of the signal is 7.0 sigma. The stated half-life considers captures from the K shell up to the N5 shell. This result supersedes APRILE 19E, which exclusively considered captures from the K shell.

<sup>10</sup> APRILE 22B use data collected by the XENONnT dark matter experiment to derive an improved <sup>124</sup>Xe  $2\nu$ DEC half-life measurement for <sup>124</sup>Xe. This result supersedes APRILE 22A.

<sup>11</sup>NOVELLA 22 report on a high-pressure gas TPC at Canfranc underground laboratory, filled with 3.5 kg (fiducial) xenon gas, used to measure the  $2\nu \beta\beta$  decay of <sup>136</sup>Xe. Topological track reconstruction is utilized in the data analysis. The measurement is based on comparing runs with isotopically enriched and depleted xenon. Other measurements with smaller error exist.

<sup>12</sup> ADAMS 21 use 102.7 kg yr of <sup>130</sup>Te exposure, collected by the CUORE bolometric detector at LNGS, to perform a measurement of the  $2\nu\beta\beta$  decay of <sup>130</sup>Te. The dataset is more than 10-times that collected by the CUORE-0 experiment. The result has been revised in ADAMS 23A. Supersedes ALDUINO 17.

<sup>13</sup> APRILE 19E report first measurement of two-neutrino double electron capture in <sup>124</sup>Xe using the XENON1T detector with a 0.73 t-yr exposure. An excess of 126  $\pm$  29 events

is observed at 64.3  $\pm$  0.6 keV decay energy, corresponding to  $\sqrt{\Delta\chi^2}$  = 4.4 with respect to the background-only hypothesis.

NODE=S076H2N;LINKAGE=HA

NODE=S076H2N;LINKAGE=W

NODE=S076H2N;LINKAGE=X

NODE=S076H2N;LINKAGE=V

NODE=S076H2N;LINKAGE=CA

NODE=S076H2N;LINKAGE=AA

NODE=S076H2N;LINKAGE=DA

NODE=S076H2N;LINKAGE=T

NODE=S076H2N;LINKAGE=Q

NODE=S076H2N;LINKAGE=U

NODE=S076H2N;LINKAGE=P

NODE=S076H2N;LINKAGE=N

NODE=S076H2N;LINKAGE=H

4/20/2025 20:25

 $^{14}$  ARNOLD 19 use the NEMO-3 tracking calorimeter with 34.3 kg y exposure to determine the  $2\nu \beta\beta$  half-life of <sup>100</sup>Mo. Supersedes ARNOLD 15.

 $^{15}$  ARNOLD 18 use the NEMO-3 tracking detector to determine the  $2\nu\beta\beta$  half-life of  $^{82}$  Se. 0.93 kg of  $^{82}$ Se was observed for 5.25 y. The half-life value was obtained based on the single-state-dominance (SSD) hypothesis, preferred in this case by about  $2 \sigma$ . Supersedes ARNOLD 05A.

 $^{16}$  BARABASH 18 use 1.162 kg of  $^{116}$ CdWO<sub>4</sub> scintillating crystals to obtain this value. Supersedes DANEVICH 03 with analogous source and agrees with ARNOLD 17 with the NEMO-3 detector.

 $^{17}$  ALBERT 17C uses the EXO-200 detector that contains 19.098  $\pm$  0.014% admixture of  $^{134}\mathrm{Xe}$  to search for the  $2\nu$   $\beta\beta$  decay mode. The exposure is 29.6 kg year. The median sensitivity is  $1.2\times 10^{21}$  years.

 $^{18}\,\text{ALDUINO}$  17 use the CUORE-0 detector containing 10.8 kg of  $^{130}\text{Te}$  in 52 crystals of TeO<sub>2</sub>. The exposure was 9.3 kg yr of  $^{130}$ Te. This is a more accurate rate determination than in ARNOLD 11 and BARABASH 11A.

19 ARNOLD 17 use the NEMO-3 tracking calorimeter, containing 410 grams of enriched  $^{116}$ Cd exposed for 5.26 years, to determine the half-life value.

 $^{20}$ ARNOLD 16 use the NEMO-3 detector and a source of 6.99 g of  $^{48}$ Ca. The half-life is based on 36.7 g year exposure. It is consistent, although somewhat longer, than the previous determinations of the half-life. Supersedes BARABASH 11A.

 $^{21}$  ARNOLD 16A use the NEMO-3 tracking calorimeter, containing 36.6 g of  $^{150}$ Nd exposed for 1918.5 days, to determine the half-life. Supersedes ARGYRIADES 09.

 $^{22}$  AGOSTINI 15A use 17.9 kg yr exposure of the GERDA calorimeter to derive an improved measurement of the  $2\nu\beta\beta$  decay half life of  $^{76}{\rm Ge}.$ 

 $^{23}$  ARNOLD 15 use the NEMO-3 tracking calorimeter with 34.3 kg yr exposure to determine the  $2\nu\beta\beta$ -half life of <sup>100</sup>Mo. Supersedes ARNOLD 05A and ARNOLD 04.

 $^{24}\text{ALBERT}$  14 use the EXO-200 tracking detector for a re-measurement of the  $2\nu\beta\beta\text{-half}$ life of  $^{136}\text{Xe.}$  A nuclear matrix element of 0.0218  $\pm$  0.0003  $\text{MeV}^{-1}$  is derived from this data. Supersedes ACKERMAN 11.

 $^{25}\,\text{GAVRILYAK}$  13 use a proportional counter filled with Kr gas to search for the  $2\nu2\text{K}$ decay of <sup>78</sup>Kr. Data with the enriched and depleted Kr were used to determine signal and background. A 2.5 $\sigma$  excess of events obtained with the enriched sample is interpreted as an indication for the presence of this decay.

 $^{26}\,{\rm GANDO}$  12A use a modification of the existing KamLAND detector. The  $\beta\beta$  decay source/detector is 13 tons of enriched <sup>136</sup>Xe-loaded scintillator contained in an inner balloon. The  $2\nu\beta\beta$  decay rate is derived from the fit to the spectrum between 0.5 and 4.8 MeV. This result is in agreement with ACKERMAN 11.

<sup>27</sup> ARNOLD 11 use enriched  $1\overline{30}$  Te in the NEMO-3 detector to measure the  $2\nu \beta\beta$  decay

rate. This result is in agreement with, but more accurate than ARNABOLDI 03. <sup>28</sup> ARGYRIADES 10 use 9.4  $\pm$  0.2 g of <sup>96</sup>Zr in NEMO-3 detector and identify its  $2\nu\beta\beta$ decay. The result is in agreement and supersedes ARNOLD 99.

<sup>29</sup> BELLI 10 use enriched <sup>100</sup>Mo with 4 HP Ge detectors to record the 590.8 and 539.5 keV  $\gamma$  rays from the decay of the  $0^+_1$  state in <sup>100</sup>Ru both in singles and coincidences. This result confirms the measurement of KIDD 09 and ARNOLD 07 and supersedes them.

 $^{30}$  First exclusive measurement of  $2\nu$ -decay to the first excited  $0^+_1$ -state of daughter nucleus. ARNOLD 07 use the NEMO-3 tracking calorimeter to detect all particles emitted in decay. Result agrees with the inclusive  $(0\nu + 2\nu)$  measurement of DEBRAECKELEER 01.

 $^{31}\mathrm{ARNOLD}$  05A use the NEMO-3 tracking detector to determine the  $2\nu\beta\beta$  half-life of <sup>82</sup>Se with high statistics and low background (389 days of data taking). Supersedes ARNOLD 04.

 $^{32}$ DANEVICH 03 is calorimetric measurement of 2
uetaeta ground state decay of  $^{116}$ Cd using enrichedCdWO<sub>4</sub> scintillators. Agrees with EJIRI 95 and ARNOLD 96. Supersedes DANEVICH 00.

### $\langle m_{\rm ee} \rangle$ , The Effective Weighted Sum of Majorana Neutrino Masses Contributing to Neutrinoless Double- $\beta$ Decay

 $\langle m_{
m ee} 
angle = |\Sigma U_{ei}^2 m_{
u_i}|$ , i = 1,2,3. It is assumed that  $u_i$  are Majorana particles and that the transition is dominated by the known (light) neutrinos. Note that  $U_{ei}^2$  and not  $|U_{ei}|^2$  occur in the sum, and that consequently cancellations are possible. The experiments obtain the limits on  $\langle m_{
u} 
angle$  from the measured ones on  $T_{1/2}$  using a range of nuclear matrix elements (NME), which is reflected in the spread of  $\langle m_{
u} 
angle$ . Different experiments may choose different NME. All assume  $g_A\,=\,1.27.\,$  In the following Listings, only the best or comparable limits for each isotope are reported. When not mentioned explicitly the transition is between ground states, but transitions between excited states are also reported.

VALUE (eV)	ISOTOPE	METHOD	DOCUMENT ID

• • • We do not use the following data for averages, fits, limits, etc. • • •

NODE=S076H2N;LINKAGE=G

NODE=S076H2N;LINKAGE=E

NODE=S076H2N;LINKAGE=F

NODE=S076H2N;LINKAGE=A

NODE=S076H2N;LINKAGE=UB

NODE=S076H2N;LINKAGE=YC

NODE=S076H2N;LINKAGE=JB

NODE=S076H2N;LINKAGE=XB

NODE=S076H2N;LINKAGE=PA

NODE=S076H2N;LINKAGE=XA

NODE=S076H2N;LINKAGE=Z

NODE=S076H2N;LINKAGE=EA

NODE=S076H2N;LINKAGE=GA

NODE=S076H2N;LINKAGE=AD

NODE=S076H2N;LINKAGE=RG

NODE=S076H2N;LINKAGE=IE

NODE=S076H2N;LINKAGE=NA

NODE=S076H2N;LINKAGE=N4

NODE=S076H2N;LINKAGE=VA

NODE=S076MW NODE=S076MW

NODE=S076MW

Page 5

NODE=S076MW;LINKAGE=OB

NODE=S076MW;LINKAGE=QB

NODE=S076MW;LINKAGE=KB

NODE=S076MW;LINKAGE=LB

NODE=S076MW;LINKAGE=NB

NODE=S076MW;LINKAGE=MB

NODE=S076MW:LINKAGE=JB

NODE=S076MW;LINKAGE=HB

NODE=S076MW;LINKAGE=IB

NODE=S076MW;LINKAGE=EB

NODE=S076MW;LINKAGE=FB

NODE=S076MW;LINKAGE=AB

NODE=S076MW;LINKAGE=VA

NODE=S076MW;LINKAGE=YA

< 0.036-0.156	<sup>136</sup> Xe	KamLAND-Zen	<sup>1</sup> ABE	23	
< 0.113-0.269	<sup>76</sup> Ge	MAJORANA	<sup>2</sup> ARNQUIST	23	
< 0.48-3.19	<sup>136</sup> Xe	NEXT	<sup>3</sup> NOVELLA	23	
< 0.09-0.305	<sup>130</sup> Te	CUORE	<sup>4</sup> ADAMS	22A	
< 0.8–2.5	136 <sub>Xe</sub>	XENON1T	<sup>5</sup> APRILE	22A	
< 0.28–0.49	100 <sub>Mo</sub>	CUPID-Mo	<sup>6</sup> AUGIER	22	
< 0.263–0.545	<sup>82</sup> Se	CUPID-0	<sup>7</sup> AZZOLINI	22	
< 0.31–0.54	100 <sub>Mo</sub>	CUPID-Mo	<sup>8</sup> ARMENGAUD	21	
< 0.075–0.35	130 <sub>Te</sub>	CUORE	<sup>9</sup> ADAMS	20A	
< 0.079–0.180	76 <sub>Ge</sub>	GERDA	<sup>10</sup> AGOSTINI	20в	
< 1.2–2.1	100 <sub>Mo</sub>	AMoRE	<sup>11</sup> ALENKOV	19	
< 0.093–0.286	136 <sub>Xe</sub>	EXO-200	<sup>12</sup> ANTON	19	
< 1.3–3.5	136 <sub>Xe</sub>	PANDAX-II	<sup>13</sup> NI	19	
< 0.11–0.52	130 <sub>Te</sub>	CUORE	<sup>14</sup> ALDUINO	18	
< 1.2–3.0	<sup>82</sup> Se	NEMO-3	<sup>15</sup> ARNOLD	18	OCCUR=6
< 1.0–1.7	$^{116}Cd$	AURORA	<sup>16</sup> BARABASH	18	
< 1.4–2.5	$^{116}Cd$	NEMO-3	<sup>17</sup> ARNOLD	17	
< 0.27–0.76	130 <sub>Te</sub>	CUORICINO	<sup>18</sup> ALDUINO	16	
< 1.6–5.3	<sup>150</sup> Nd	NEMO-3	<sup>19</sup> ARNOLD	16A	
< 0.33–0.62	100 <sub>Mo</sub>	NEMO-3	<sup>20</sup> ARNOLD	15	
< 7.2–19.5	<sup>96</sup> Zr	NEMO-3	<sup>21</sup> ARGYRIADES	10	
< 3.5–22	<sup>48</sup> Ca	CaF <sub>2</sub> scint.	<sup>22</sup> UMEHARA	08	
< 1.5–1.7	$^{116}Cd$	<sup>116</sup> CdWO <sub>4</sub> scint.	<sup>23</sup> DANEVICH	03	
$^1$ ABE 23 utilize 745 k	g of <sup>136</sup> X	e isotope exposure fro	m the combined da	ata set of the	NODE=S076MW;LINKAGE=PB

<sup>1</sup> ABE 23 utilize 745 kg of <sup>130</sup>Xe isotope exposure from the combined data set of the KamLAND-Zen 400 and 800 to derive a limit on  $\langle m_{\beta\beta} \rangle$ . The range reflects the author's assessment of the variability of the theoretically calculated nuclear matrix elements.

 $^2$  ARNQUIST 23 use the final data set of the MAJORANA DEMONSTRATOR experiment, with 64.5 kg yr of isotop exposure, to derive an upper limit for  $\langle m_{\beta\,\beta}\rangle$ . The range reflects the author's assessment of the variability of the theoretically calculated nuclear matrix elements.

- <sup>3</sup> NOVELLA 23 use data collected with the NEXT-White experiment to derive a range of upper limits for  $\langle m_{\beta\beta} \rangle$ . The range reflects the author's assessment of the variability of the theoretically calculated nuclear matrix elements and both half-life limits stated in NOVELLA 23.
- <sup>4</sup> ADAMS 22A use 1038.4 kg-yr of TeO<sub>2</sub> exposure collected by the CUORE experiment to determine this range of limits. The range reflects the uncertainty of nuclear matrix element calculations needed for the conversion of half-life to neutrino mass.
- <sup>5</sup> APRILE 22A use data taken with the XENON1T detector to limit the Majorana neutrino mass. 36.16 kg·yr of <sup>136</sup>Xe exposure were utilized. The reported range of limits is due to uncertainties in the nuclear matrix elements.
- <sup>6</sup> AUGIER 22 use the final data set of the CUPID-Mo cryogenic calorimeter with an isotop exposure of 1.47 kg·y to derive a range of neutrino mass limits. The range reflects the authors' estimate of the spread of nuclear matrix element calculations.
- <sup>7</sup> AZZOLINI 22 use 8.82 kg yr of isotopic exposure of the CPID-0 scintillating cryogenic bolometer to set this range of neutrino mass limits. The range reflects the authors' estimate of the spread of nuclear matrix element calculations.
- <sup>8</sup> ARMENGAUD 21 use the CUPID-Mo demonstrator, with 1.17 kg·yr exposure of <sup>100</sup>Mo, to set this limit. The range reflects the estimated uncertainty of the calculated nuclear matrix elements.
- $^9$  matrix elements.  $^9$  ADAMS 20A use the data of CUORE (372.5 kg·yr exposure of TeO<sub>2</sub>) to obtain this limit.
- <sup>10</sup> AGOSTINI 20B use the final data set of the GERDA experiment, representing an exposure of 127.2 kg·yr to derive an upper limit for  $\langle m_{\beta\beta} \rangle$ . Isotopically enriched Ge detectors were used. The range reflects the variability of the theoretically calculated nuclear matrix elements. Supersedes AGOSTINI 19.
- <sup>11</sup> ALENKOV 19 report the range of the effective masses  $\langle m_{\beta\beta} \rangle$  corresponding to the  $0\nu$  $\beta\beta$  decay half-life limit. It is based on the 52.1 kg d exposure of <sup>100</sup> Mo, in the Yangyang underground laboratory. The median sensitivity is  $1.1 \times 10^{23}$  years. The range of  $\langle m_{\beta\beta} \rangle$  reflects the uncertainty of nuclear matrix elements.
- <sup>12</sup> ANTON 19 uses the complete dataset of the EXO-200 experiment to obtain these limits. The spread reflect the uncertainty in the nuclear matrix elements. Supersedes ALBERT 18 and ALBERT 14B.
- <sup>13</sup> NI 19 use the PandaX-II dual phase TPC at CJPL to search for the  $0\nu \beta\beta$  decay of <sup>136</sup>Xe with 22.2 kg yr exposure. The range in the  $m_{\beta\beta}$  limit of 1.3–3.5 eV reflects the

range of the calculated nuclear matrix elements. The sensitivity is  $1.9 \times 10^{23}$  yr.

- <sup>14</sup> ALDUINO 18 use the combined data of CUORE, CUOREO, and Cuoricino to obtain this limit.
- <sup>15</sup>ARNOLD 18 use the NEMO-3 tracking detector to constrain the  $0\nu\beta\beta$  decay of <sup>82</sup>Se. The limit on  $\langle m_{\beta\beta} \rangle$  is obtained assuming light neutrino exchange; the range reflects different calculations of the nuclear matrix elements. This is a somewhat weaker limit than in BARABASH 11A using the same detector.

NODE=S076MW;LINKAGE=ZA

NODE=S076MW;LINKAGE=IA

- $^{16}$  BARABASH 18 use 1.162 kg of  $^{116} {\rm CdWO}_4$  scintillating crystals to obtain these limits. The spread reflects the estimated uncertainty in the nuclear matrix element. Supersedes DANEVICH 03.
- $17\,\text{ARNOLD}$  17 utilize NEMO-3 data, taken with enriched  $116\,\text{Cd}$  to limit the effective Majorana neutrino mass. The reported range results from the use of different nuclear matrix elements. Supersedes BARABASH 11A.
- <sup>18</sup> ALDUINO 16 place a limit on the effective Majorana neutrino mass using the combined data of the CUORE-0 and CUORICINO experiments. The range reflects the authors' evaluation of the variability of the nuclear matrix elements. Supersededs ALFONSO 15.
- <sup>19</sup> ARNOLD 16A limit is derived from data taken with the NEMO-3 detector and <sup>150</sup>Nd. A range of nuclear matrix elements that include the effect of nuclear deformation have been used. Supersedes ARGYRIADES 09.
- $^{20}$  ARNOLD 15 use the NEMO-3 tracking calorimeter with 34.3 kg yr exposure to determine the neutrino mass limit based on the  $0\nu\beta\beta$ -half life of  $^{100}$ Mo. The spread range reflects different nuclear matrix elements. Supersedes ARNOLD 14 and BARABASH 11A.
- $^{21}$  ARGYRIADES 10 use  $^{96}$  Zr and the NEMO-3 tracking detector to obtain the reported mass limit. The range reflects the fluctuation of the nuclear matrix elements considered.
- $^{22}$  Limit was obtained using CaF $_2$  scintillation calorimeter to search for double beta decay of  $^{48}$ Ca. Reported range of limits reflects spread of QRPA and SM matrix element calculations used. Supersedes OGAWA 04.
- $^{23}$  Limit for  $\langle m_{\nu}\rangle$  is based on the nuclear matrix elements of STAUDT 90 and ARNOLD 96. Supersedes DANEVICH 00.

### Limits on Lepton-Number Violating (V+A) Current Admixture

Considering that a number of experiments earlier than 1989 did not distinguish between  $\lambda$  and  $\eta$ , we list only results from that year on.  $\langle\lambda\rangle = \lambda \sum U_{ej}V_{ej}$  and  $\langle\eta\rangle = \eta \sum U_{ej}V_{ej}$ , where the sum is over the number of neutrino generations. This sum vanishes for massless or unmixed neutrinos. In the following Listings, only best or comparable limits or lifetimes for each isotope are reported.

$\left< \lambda \right>$ (10 <sup>-6</sup> )	CL%	$\left< \eta \right>$ (10 <sup>-8</sup> )	CL%	ISOTOPE	METHOD	DOCUMENT ID		NODE=S076ETA
• • • We d	o not							
< 2.2–2.6	90	< 1.7–2.1	90	<sup>82</sup> Se	NEMO-3	<sup>1</sup> ARNOLD	18	
< 1.8–22	90	< 1.6–21	90	<sup>116</sup> Cd	AURORA	<sup>2</sup> BARABASH	18	OCCUR=2
< 0.9–1.3	90	< 0.5–0.8	90	100 <sub>Mo</sub>	NEMO-3	<sup>3</sup> ARNOLD	14	
<120	90			<sup>100</sup> Mo	$0^+  ightarrow 2^+$	<sup>4</sup> ARNOLD	07	
$0.692 \substack{+0.05 \\ -0.05}$	8 6 <sup>8</sup> 68	$0.305 \substack{+0.02 \\ -0.02}$	6 5 68	76 <sub>Ge</sub>	Enriched HPGe	<sup>5</sup> KLAPDOR-K	06A	
< 2.5	90			<sup>100</sup> Mo	$0\nu$ , NEMO-3	<sup>6</sup> ARNOLD	05A	
< 3.8	90			<sup>82</sup> Se	$0\nu$ , NEMO-3	<sup>7</sup> ARNOLD	05A	OCCUR=2
< 1.5–2.0	90			<sup>100</sup> Mo	$0\nu$ , NEMO-3	<sup>8</sup> ARNOLD	04	
< 3.2–3.8	90			<sup>82</sup> Se	$0\nu$ , NEMO-3	<sup>9</sup> ARNOLD	04	OCCUR=2
< 1.6–2.4	90	< 0.9–5.3	90	<sup>130</sup> Te	Cryog. det.	<sup>10</sup> ARNABOLDI	03	
< 2.2	90	<2.5	90	$^{116}Cd$	<sup>116</sup> CdWO <sub>4</sub> scint.	<sup>11</sup> DANEVICH	03	
< 3.2–4.7	90	< 2.4–2.7	90	100 <sub>Mo</sub>	ELEGANT V	<sup>12</sup> EJIRI	01	
< 1.1	90	<0.64	90	$76_{Ge}$	Enriched HPGe	<sup>13</sup> GUENTHER	97	
< 4.4	90	<2.3	90	<sup>136</sup> Xe	ТРС	<sup>14</sup> VUILLEUMIER	93	
		<5.3		<sup>128</sup> Te	Geochem	<sup>15</sup> BERNATOW	92	

- <sup>1</sup>ARNOLD 18 use the NEMO03 tracking detector, with 0.93 kg of  $^{82}$ Se mass and 5.25 y exposure to obtain the limits for the hypothetical right-handed currents. Supersedes ARNOLD 05A.
- $^{2}$ BARABASH 18 use 1.162 kg of  $^{116}$ CdWO<sub>4</sub> scintillating crystals to obtain this limits for the hypothetical right-handed currents in the 0 $\nu\beta\beta$  decay of  $^{116}$ Cd.
- <sup>3</sup>ARNOLD 14 is based on 34.7 kg yr of exposure of the NEMO-3 tracking calorimeter. The reported range limit on  $\langle \lambda \rangle$  and  $\langle \eta \rangle$  reflects the nuclear matrix element uncertainty in <sup>100</sup>Mo.
- <sup>4</sup> ARNOLD 07 use NEMO-3 half life limit for 0 $\nu$ -decay of <sup>100</sup>Mo to the first excited 2<sup>+</sup>-state of daughter nucleus to limit the right-right handed admixture of weak currents  $\langle \lambda \rangle$ . This limit is not competitive when compared to the decay to the ground state.
- <sup>5</sup> Re-analysis of data originally published in KLAPDOR-KLEINGROTHAUS 04A. Modified pulse shape analysis leads the authors to claim 6 $\sigma$  statistical evidence for observation of 0 $\nu$ -decay. Authors use matrix element of MUTO 89 to determine  $\langle \lambda \rangle$  and  $\langle \eta \rangle$ . Uncertainty of nuclear matrix element is not reflected in stated errors.

<sup>6</sup> ARNOLD 05A derive limit for  $\langle \lambda \rangle$  based on <sup>100</sup>Mo data collected with NEMO-3 detector. No limit for  $\langle \eta \rangle$  is given. Supersedes ARNOLD 04.

<sup>7</sup> ARNOLD 05A derive limit for  $\langle \lambda \rangle$  based on <sup>82</sup>Se data collected with NEMO-3 detector. No limit for  $\langle \eta \rangle$  is given. Supersedes ARNOLD 04.

<sup>8</sup> ARNOLD 04 use the matrix elements of SUHONEN 94 to obtain a limit for  $\langle \lambda \rangle$ , no limit for  $\langle \eta \rangle$  is given. This limit is more stringent than the limit in EJIRI 01 for the same nucleus.

NODE=S076MW;LINKAGE=EA

NODE=S076MW;LINKAGE=FA

NODE=S076MW;LINKAGE=CA

NODE=S076MW;LINKAGE=RG

NODE=S076MW;LINKAGE=UE

NODE=S076MW;LINKAGE=VF

NODE=S076ETA NODE=S076ETA

NODE=S076ETA;LINKAGE=H

NODE=S076ETA;LINKAGE=G

NODE=S076ETA;LINKAGE=E

NODE=S076ETA;LINKAGE=NA

NODE=S076ETA;LINKAGE=KR

NODE=S076ETA;LINKAGE=N1

NODE=S076ETA;LINKAGE=N2

NODE=S076ETA;LINKAGE=A1

- $^9$ ARNOLD 04 use the matrix elements of TOMODA 91 and SUHONEN 91 to obtain a limit for  $\langle\lambda\rangle$ , no limit for  $\langle\eta\rangle$  is given.
- $^{10}\,{\rm Supersedes}$  ALESSANDRELLO 00. Cryogenic calorimeter search. Reported a range reflecting uncertainty in nuclear matrix element calculations.
- $^{11}$  Limits for  $\langle\lambda\rangle$  and  $\langle\eta\rangle$  are based on nuclear matrix elements of STAUDT 90. Supersedes DANEVICH 00.
- <sup>12</sup>The range of the reported  $\langle \lambda \rangle$  and  $\langle \eta \rangle$  values reflects the spread of the nuclear matrix elements. On axis value assuming  $\langle m_{\nu} \rangle = 0$  and  $\langle \lambda \rangle = \langle \eta \rangle = 0$ , respectively.
- $^{13}\,\rm GUENTHER$  97 limits use the matrix elements of STAUDT 90. Supersedes BALYSH 95 and BALYSH 92. <sup>14</sup> VUILLEUMIER 93 uses the matrix elements of MUTO 89. Based on a half-life limit
- $2.6 \times 10^{23}$  y at 90%CL.
- $^{15}\,{\sf BERNATOWICZ}$  92 takes the measured geochemical decay width as a limit on the  $0\nu$ width, and uses the SUHONEN 91 coefficients to obtain the least restrictive limit on  $\eta$ . Further details of the experiment are given in BERNATOWICZ 93.

### Double- $\beta$ Decay REFERENCES

AAI BERS	24C	IP G52 015103		L Aalbers <i>et al</i>		(17	Collab )
VAN	24	DPI 132 152502		X Van at al		(PandaX AT	Collab.)
	24	FRE 132 132302					Collab.
ABE	23	PRL 130 051801	(	S. Abe et al.		(KamLAND-Zen	Collab.
ADAMS	23A	PRL 131 249902	(errat.)	D.Q. Adams et al.		(CUORE	Collab.
AGOSTINI	23	PRL 131 142501		M. Agostini <i>et al.</i>		(GERDA	Collab.)
AGUERRE	23	EPJ C83 1117		X. Aguerre et al.		(NEMO-3	Collab.)
ARNQUIST	23	PRL 130 062501		L.I. Arnquist et al.		(MA)ORANA	Collab.
ALICIER	23	PR (107 025503		C Augier et al		(CUPID-Mo	Collab
	220	DDI 121 162500		C Augier et al		CUPID Me	Collab.)
AUGIER	234	PRL 131 102301		C. Augier et al.			Collab.
AZZOLINI	23A	PRL 151 222501		O. Azzolini et al.		(COPID-0	Collab.
NOVELLA	23	JHEP 2309 190		P. Novella <i>et al.</i>		(NEX I	Collab.
ADAMS	22A	NAT 604 53		D.Q. Adams <i>et al.</i>		(CUORE	Collab.)
ADAMS	22B	PRL 129 222501		D.Q. Adams et al.		(CUORE	Collab.)
APRILE	22A	PR C106 024328		E. Aprile et al.		(XÈNON1T	Collab.
APRILE	22B	PRI 129 161805		E Aprile et al		(XENONnT	Collab
ALICIER	22	FPI C82 1033		C Augier et al		(CUPID-Mo	Collab
	22	DDI 100 111001		O Azzolini ot ol			Collab.
	22	PRC 129 111001		D. Azzolilli et al.			Collab.
NOVELLA	22	PR C105 055501		P. Novella et al.		(NEXT	Collab.
ADAMS	21	PRL 126 171801		D.Q. Adams <i>et al.</i>		(CUORE	Collab.)
Also		PRL 131 249902	(errat.)	D.Q. Adams <i>et al.</i>		(CUORE	Collab.)
ADAMS	21A	EPJ C81 567		D.Q. Adams et al.		(CUORE	Collab.)
ARMENGAUD	21	PRL 126 181802		E. Armengaud et al.		(CUPID-Mo	Collab.)
ARNQUIST	21	PR C103 015501		I.I. Arnquist et al.		(MAJORANA	Collab.
ADAMS	20A	PRI 124 122501		D.O. Adams et al		(CHORE	Collab )
	207	DDI 125 252502		M Agostini at al		(CERDA	Collab.
AGUSTINI	200	FRE 123 232302					Collab.
ARMENGAUD	20	EPJ C80 674		E. Armengaud et al.		(CUPID-Mo	Collab.
AGOSTINI	19	SCI 365 1445		M. Agostini <i>et al.</i>		(GERDA	Collab.)
ALDUINO	19	EPJ C79 795		C. Alduino <i>et al.</i>		(CUORE	Collab.)
ALENKOV	19	EPJ C79 791		V. Alenkov et al.		(AMoRE	Collab.)
ANTON	19	PRL 123 161802		G. Anton <i>et al.</i>		(ÈXO-200	Collab.)
APRILE	19E	NAT 568 532		E. Aprile et al.		(XENON1T	Collab.)
ARNOLD	19	FP1 C79 440		R Arnold et al		(NEMO-3	Collab
A770LINI	19	PRI 123 032501		O Azzolini <i>et al</i>		(CUPID-0	Collab
AZZOLINI	19R	PRI 123 262501		O Azzolini et al		(CUPID-0	Collab.)
NI	10	CP C43 113001		K Ni et al		(PandaX-II	Collab.)
ALBERT	18	PRI 120 072701		I.B. Albert <i>et al</i>		(EXO_200	Collab.)
	10	DRI 120 072701		C Alduino et al			Collab.)
	10	FDL C70 001		C. Aldullo et al.			Collab.
ARNOLD	10	EPJ C/0 021		R. Arnold et al.			Collab.
DARADASH	10	PR D96 092007		A.S. Darabash et al.			Collab.
ALBERT	170	PR D96 092001		J.B. Albert <i>et al.</i>		(EXO-200	Collab.
ALDUINO	17	EPJ C77 13		C. Alduino <i>et al.</i>		(CUORE	Collab.)
ARNOLD	17	PR D95 012007		R. Arnold <i>et al.</i>		(NEMO-3	Collab.)
ALDUINO	16	PR C93 045503		C. Alduino et al.		(CUORE	Collab.)
ARNOLD	16	PR D93 112008		R. Arnold et al.		(NEMO-3	Collab.)
ARNOLD	16A	PR D94 072003		R. Arnold et al.		(NEMO-3	Collab.)
ASAKURA	16	NP A946 171		K. Asakura <i>et al.</i>		(KamLAND-Zen	Collab.
AGOSTINI	15A	FP1 C75 416		M Agostini et al		(GERDA	Collab
ALEONSO	15	PRI 115 102502		K Alfonso et al		CLIORE	Collab )
	15	PR D02 072011		R Arnold at al		(NEMO 3	Collab.)
ALDEDT	14	DD C00 015502		Albert et al		(EVO 200	Collab.
ALDERT	140	NAT E10 220		I.D. Albert et al.		(EXO-200	Callab.)
	14D	NAT 310 229		J.B. Albert et al.		(EAO-200	Collab.
ARNOLD	14	PR D89 111101		R. Arnold et al.		(NEIVIO-3	Collab.)
GAVRILYAK	13	PR C87 035501		Yu.IVI. Gavrilyuk et al.		(	
ANDREOTIT	12	PR C85 045503		E. Andreotti <i>et al.</i>		(CUORICINO	Collab.)
GANDO	12A	PR C85 045504		A. Gando <i>et al.</i>		(KamLAND-Zen	Collab.)
ACKERMAN	11	PRL 107 212501		N. Ackerman <i>et al.</i>		(EXO	Collab.)
ARNOLD	11	PRL 107 062504		R. Arnold et al.		(NEMO-3	Collab.)
BARABASH	11A	PAN 74 312		A.S. Barabash et al.		(NEMO-3	Collab.)
		Translated from Y	AF 74 3	330.		,	,
ARGYRIADES	10	NP A847 168		J. Argyriades et al.		(NEMO-3	Collab.)
BELLI	10	NP A846 143		P. Belli <i>et al.</i>		(DAMA-INR	Collab.)
ARGYRIADES	09	PR C80 032501		J. Argyriades et al.		(NEMO-3	Collab.
KIDD	09	NP A821 251		M Kidd et al		(	,
LIMEHARA	08	PR C78 058501		S Ilmehara et al			
	07	NP A781 200		R Arnold et al		(NEMO 3	Collab )
	064	MDI A01 1547		HV Klandor Kloingrathaus	w	Krivoshoina	conau.)
	064	DDI 05 100200		P Arnold at -/	I. V.		Caller
	03A	FRL 93 102302		R. Annold et al.		(14E1VIO-3	conab.)
AALSEIH	04	FR D/U U/8302		C.E. Aaisetn <i>et al.</i>			C.I
AKNOLD	04	JETPL 80 377		K. Arnold <i>et al.</i>		(NEMO-3	Collab.)
		Translated from Z		0 429.			

NODE=S076ETA;LINKAGE=A2

NODE=S076ETA;LINKAGE=RR

NODE=S076ETA;LINKAGE=VG

NODE=S076ETA;LINKAGE=EJ

NODE=S076ETA;LINKAGE=J1

NODE=S076ETA;LINKAGE=V

NODE=S076ETA;LINKAGE=BB

NODE=S076

ĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸĸ	63173 62725 62010 62586 62255 62209 62469 62533 62469 62469 62533 61928 61928 61928 61928 61928 61976 61915 61838 661976 61181 62586 661976 661838 661259 660704 660167 660167 660167 660167 660167 659810 660167 558306 600647 559810 559810 5598773 557295 557345 556395 557295 556395 557295 556395 556406 555822 554383 556406 558225 554383 556406 558225 554383 556406 558225 554383 556406 558225 554383 556406 558225 554383 556406 558225 554383 556406 558225 554383 556406 55825 554483 556406 55825 554483 556406 55825 554483 556406 55825 554483 556406 55825 554483 556406 55825 554483 556406 55825 554483 556406 55825 556406 55825 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 556406 55826 557275 556406 55826 556406 55826 557275 556406 55826 557275 556406 55826 557275 556406 55826 55826 557275 556406 55826 55826 557275 556406 55826 55856 55826 55856 55826 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856 55856
REF REF REF REF REF REF REF REF REF	53491 53385 53053 52783 52522 51598 51401 50934 50235 50411

DANEVICH03PR D65 092007C.E. Aalseth <i>et al.</i> (IGEX Collab.)DEBRAECKEL01PRL 86 3510L. De Braeckeleer <i>et al.</i> (IGEX Collab.)DEBRAECKEL01PR C63 065501H. Ejiri <i>et al.</i> (IGEX Collab.)KLAPDOR-K01EPJ A12 147H.V. Klapdor-Kleingrothaus <i>et al.</i> KLAPDOR-K01EPJ A12 147H.V. Klapdor-Kleingrothaus <i>et al.</i> ALESSAND00PL B486 13A. Alessandrello <i>et al.</i> ALESSAND00PL B486 13A. Alessandrello <i>et al.</i> ARNOLD90PN C62 045501F.A. Danevich <i>et al.</i> (BUENTHER 97PR D55 54M. Gunther <i>et al.</i> (Heidelberg-Moscow Collab.)GUENTHER 97PR D55 54M. Gunther <i>et al.</i> (BCEN, CAEN, JINR+)BALYSH95PL B356 450A. Balysh <i>et al.</i> (Heidelberg-Moscow Collab.)DASSIE95PR D51 2090D. Dassie <i>et al.</i> (NEMO Collab.)EJIRI95JPSJ 64 339H. Ejiri <i>et al.</i> (OSAK, KIEV)SUHONEN 94PR C49 3055J. Suhonen, O. Civitarese(NEUC, CIT, VILL)BALYSH92PL B283 32A. Balysh <i>et al.</i> (MUSL, TATA)VUILLEUMIER 93PR D48 1009J.C. Vuilleumier <i>et al.</i> (WUSL, TATA)SUHONEN 91NP A535 509J. Suhonen, S.B. Khadkikar, A. Faessler(JYV+)TOMODA91RPP 54 53T. Tomoda(WUSL, TATA)STAUDT90EPL 13 31A. Staudt, K. Muto, H.V. Klapdor-KleingrothausMUIDO80ZPHY X342 187K. Mutt	REFID=48163 REFID=48460 REFID=48628 REFID=47726 REFID=47726 REFID=47325 REFID=45265 REFID=45265 REFID=45277 REFID=44401 REFID=44401 REFID=50527 REFID=50527 REFID=50527 REFID=43219 REFID=43246 REFID=42181 REFID=42181 REFID=45867 REFID=45867 REFID=43301
--	---