See key on page 1343

Meson Full Listings

$K_1(1400)$, $K^*(1410)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 200.0	VERGEEST	79	HBC	–	4.2	$K^- p \to (\overline{K}\pi\pi)^- p$
~ 160	BRANDENB...	76	ASPK	±	13	$K^\pm p \to (K\pi\pi)^\pm p$
80	DAVIS	72	HBC	+	12	$K^+ p$
241 ±30	FIRESTONE	72B	DBC	+	12	$K^+ d$

[4] From partial-wave analysis of $K^0\pi^+\pi^-$ system.

[5] From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.



$K_1(1400)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)
Γ_1	$K^*(892)\pi$	(94 ±6) %
Γ_2	$K\rho$	(3.0±3.0) %
Γ_3	$K f_0(1300)$	(2.0±2.0) %
Γ_4	$K\omega$	(1.0±1.0) %
Γ_5	$K_0^*(1430)\pi$	

$K_1(1400)$ PARTIAL WIDTHS

$\Gamma(K^*(892)\pi)$ Γ_1

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
117.0±10.0	CARNEGIE	77	ASPK	±	13 $K^\pm p \to (K\pi\pi)^\pm p$

$\Gamma(K\rho)$ Γ_2

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
2.0±1.0	CARNEGIE	77	ASPK	±	13 $K^\pm p \to (K\pi\pi)^\pm p$

$\Gamma(K\omega)$ Γ_4

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
23.0±12.0	CARNEGIE	77	ASPK	±	13 $K^\pm p \to (K\pi\pi)^\pm p$

$K_1(1400)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{total}$ Γ_1/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.94±0.06	[6] DAUM	81C	CNTR	63 $K^- p \to K^- 2\pi p$

$\Gamma(K\rho)/\Gamma_{total}$ Γ_2/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.03 ±0.03	[6] DAUM	81C	CNTR	63 $K^- p \to K^- 2\pi p$

$\Gamma(K f_0(1300))/\Gamma_{total}$ Γ_3/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.02 ±0.02	[6] DAUM	81C	CNTR	63 $K^- p \to K^- 2\pi p$

$\Gamma(K\omega)/\Gamma_{total}$ Γ_4/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
0.01 ±0.01	[6] DAUM	81C	CNTR	63 $K^- p \to K^- 2\pi p$

$\Gamma(K_0^*(1430)\pi)/\Gamma_{total}$ Γ_5/Γ

VALUE	DOCUMENT ID		TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 0.00	[6] DAUM	81C	CNTR	63 $K^- p \to K^- 2\pi p$

D-wave/S-wave RATIO FOR $K_1(1400) \to K^*(892)\pi$

VALUE	DOCUMENT ID		TECN	COMMENT
0.04 ±0.01	[6] DAUM	81C	CNTR	63 $K^- p \to K^- 2\pi p$

[6] Average from low and high t data.

$K_1(1400)$ REFERENCES

ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	82B	NP B202 21	+	(BIRM, CERN, GLAS, MSU, CURIN)
TORNQVIST	82B	NP B203 268		(HELS)
DAUM	81C	NP B187 1	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP
VERGEEST	79	NP B158 265	+Jongejans, Dionisi+	(NIJM, AMST, CERN, OXF)
CARNEGIE	77	NP B127 509	+Cashmore, Davier, Dunwoodie, Lasinski+	(SLAC)
BRANDENB...	76	PRL 26 703	Brandenburg, Carnegie, Cashmore+	(SLAC) JP
DAVIS	72	PR D5 2688	+Alston-Garnjost, Barbaro, Flatte, Friedman, Lynch+	(LBL)
FIRESTONE	72B	PR D5 505	+Goldhaber, Lissauer, Trilling	(LBL)

OTHER RELATED PAPERS

SUZUKI	93	PR D47 1252		(LBL)
FERNANDEZ	82	ZPHY C16 95	+Aguilar-Benitez+	(MADR, CERN, CDEF, STOH)
SHEN	66	PRL 17 726	+Butterworth, Fu, Goldhaber, Trilling	(LRL)
Also	66	Private Comm.	Goldhaber	(LRL)
ALMEIDA	65	PL 16 184	+Atherton, Byer, Dornan, Forson+	(CAVE)
ARMENTEROS	64	PL 9 207	+Edwards, D'Andlau+	(CERN, CDEF)
Also	66	PR 145 1095	Barash, Kirsch, Miller, Tan	(COLU)
ARMENTEROS	64B	Dubna Conf. 1 577	+Edwards, D'Andlau+	(CERN, CDEF)
Also	64C	Dubna Conf. 1 617	Armenteros	

$K^*(1410)$

$I(J^P) = \frac{1}{2}(1^-)$

$K^*(1410)$ MASS

All from partial wave amplitude analyses.

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
1412±12 OUR AVERAGE Error includes scale factor of 1.1.					
1367±54	BIRD	89	LASS	–	11 $K^- p \to \overline{K}^0\pi^- p$
1380±21±19	ASTON	88	LASS	0	11 $K^- p \to K^-\pi^+ n$
1420± 7±10	ASTON	87	LASS	0	11 $K^- p \to \overline{K}^0\pi^+\pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1474±25	BAUBILLIER	82B	HBC	0	8.25 $K^- p \to \overline{K}^0 2\pi n$
1500±30	ETKIN	80	MPS	0	6 $K^- p \to \overline{K}^0\pi^+\pi^- n$

$K^*(1410)$ WIDTH

All from partial wave amplitude analyses.

VALUE (MeV)	DOCUMENT ID		TECN	CHG	COMMENT
227± 22 OUR AVERAGE Error includes scale factor of 1.1.					
114±101	BIRD	89	LASS	–	11 $K^- p \to \overline{K}^0\pi^- p$
176± 52±22	ASTON	88	LASS	0	11 $K^- p \to K^-\pi^+ n$
240± 18±12	ASTON	87	LASS	0	11 $K^- p \to \overline{K}^0\pi^+\pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
275± 65	BAUBILLIER	82B	HBC	0	8.25 $K^- p \to \overline{K}^0 2\pi n$
500±100	ETKIN	80	MPS	0	6 $K^- p \to \overline{K}^0\pi^+\pi^- n$

$K^*(1410)$ DECAY MODES

	Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1	$K^*(892)\pi$	> 40 %	95%
Γ_2	$K\pi$	(6.6±1.3) %	
Γ_3	$K\rho$	< 7 %	95%

$K^*(1410)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$ Γ_3/Γ_1

VALUE	CL%	DOCUMENT ID		TECN	CHG	COMMENT
<0.17	95	ASTON	84	LASS	0	11 $K^- p \to \overline{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$ Γ_2/Γ_1

VALUE	CL%	DOCUMENT ID		TECN	CHG	COMMENT
<0.16	95	ASTON	84	LASS	0	11 $K^- p \to \overline{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma_{total}$ Γ_2/Γ

VALUE	DOCUMENT ID		TECN	CHG	COMMENT
0.066±0.010±0.008	ASTON	88	LASS	0	11 $K^- p \to K^-\pi^+ n$

$K^*(1410)$ REFERENCES

BIRD	89	SLAC-332		(SLAC)
ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	+Carnegie, Dunwoodie+	(SLAC, CARL, OTTA) JP
BAUBILLIER	82B	NP B202 21	+	(BIRM, CERN, GLAS, MSU, CURIN)
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP