$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \to \Sigma(2030) \to N\overline{K}^*(892)$, S=3/2, F-wave $\qquad (\Gamma_1\Gamma_{15})^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
$+0.04\pm0.03$	[6] CAMERON	78B	DPWA	$K^- p \to N\overline{K}^*$
-0.12 ± 0.02	CORDEN	77B		$K^- d \to NN\overline{K}^*$

$\Sigma(2030)$ FOOTNOTES

[1] Preferred solution 3; see CORDEN 76 for other possibilities.

[2] The two entries for CORDEN 77C are from two different acceptable solutions.

[3] This coupling is extracted from unnormalized data.

[4] The published sign has been changed to be in accord with the baryon-first convention.

[5] An upper limit.

[6] The upper limit on the G_3 wave is 0.03.

$\Sigma(2030)$ REFERENCES

PDG	84	RMP 56 No. 2 Pt. II	Wohl, Cahn, Rittenberg+	(LBL, CIT, CERN)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
CAMERON	78	NP B143 189	+Franek, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
CAMERON	78B	NP B146 327	+Franek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	+Franek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CORDEN	77B	NP B121 365	+Cox, Kenyon, O'Neale, Stubbs, Sumorok+	(BIRM) IJP
CORDEN	77C	NP B125 61	+Cox, Kenyon, O'Neale, Stubbs, Sumorok+	(BIRM) IJP
DECLAIS	77	CERN 77-16	+Duchon, Louvel, Patry, Seguinot+	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
GOYAL	77	PR D16 2746	+Sodhi	(DELH) IJP
CORDEN	76	NP B104 382	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
CORDEN	75B	NP B92 365	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
HEMINGWAY	75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPIM) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
DEVENISH	74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
KANE	74	LBL-2452		(LBL) IJP
LITCHFIELD	74B	NP B74 19	+Hemingway, Baillon+	(CERN, HEIDH) IJP
LITCHFIELD	74C	NP B74 39	+Hemingway, Baillon+	(CERN, HEIDH) IJP
LITCHFIELD	74D	NP B74 12	+Hemingway, Baillon+	(CERN, HEIDH) IJP
MULLER	69B	UCRL 19372 Thesis		(LRL)
BURGUN	68	NP B8 447	+Meyer, Pauli, Tallini+	(SACL, CDEF, RHEL)
TRIPP	67	NP B3 10	+Leith+	(LRL, SLAC, CERN, HEID, SACL)
COOL	66	PRL 16 1228	+Giacomelli, Kycia, Leontic, Lundby+	(BNL)
WOHL	66	PRL 17 107	+Solmitz, Stevenson	(LRL) IJP

$\Sigma(2070)$ F_{15} $\qquad I(J^P) = 1(\frac{5}{2}^+)$ Status: $*$

OMITTED FROM SUMMARY TABLE

This state suggested by BERTHON 70B finds support in GOPAL 80 with new $K^- p$ polarization and $K^- n$ angular distributions. The very broad state seen in KANE 72 is not required in the later (KANE 74) analysis of $\overline{K}N \to \Sigma\pi$.

$\Sigma(2070)$ MASS

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
≈ 2070 OUR ESTIMATE				
2051 ± 25	GOPAL	80	DPWA	$\overline{K}N \to \overline{K}N$
2057	KANE	72	DPWA	$K^- p \to \Sigma\pi$
2070 ± 10	BERTHON	70B	DPWA	$K^- p \to \Sigma\pi$

$\Sigma(2070)$ WIDTH

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
300 ± 30	GOPAL	80	DPWA	$\overline{K}N \to \overline{K}N$
906	KANE	72	DPWA	$K^- p \to \Sigma\pi$
140 ± 20	BERTHON	70B	DPWA	$K^- p \to \Sigma\pi$

$\Sigma(2070)$ DECAY MODES

	Mode
Γ_1	$N\overline{K}$
Γ_2	$\Sigma\pi$

$\Sigma(2070)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\overline{K})/\Gamma_{total}$ $\qquad \Gamma_1/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
0.08 ± 0.03	GOPAL	80	DPWA	$\overline{K}N \to \overline{K}N$

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \to \Sigma(2070) \to \Sigma\pi$ $\qquad (\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
$+0.104$	KANE	72	DPWA	$K^- p \to \Sigma\pi$
$+0.12\pm0.02$	BERTHON	70B	DPWA	$K^- p \to \Sigma\pi$

$\Sigma(2070)$ REFERENCES

GOPAL	80	Toronto Conf. 159		(RHEL) IJP
KANE	74	LBL-2452		(LBL)
KANE	72	PR D5 1583		(LBL)
BERTHON	70B	NP B24 417	+Vrana, Butterworth+	(CDEF, RHEL, SACL) IJP

$\Sigma(2080)$ P_{13} $\qquad I(J^P) = 1(\frac{3}{2}^+)$ Status: $**$

OMITTED FROM SUMMARY TABLE

Suggested by some but not all partial-wave analyses across this region.

$\Sigma(2080)$ MASS

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
≈ 2080 OUR ESTIMATE				
$2091\pm\ 7$	[1] CORDEN	76	DPWA	$K^- n \to \Lambda\pi^-$
2070 to 2120	DEBELLEFON	76	IPWA	$K^- p \to \Lambda\pi^0$
2120 ± 40	BAILLON	75	IPWA	$\overline{K}N \to \Lambda\pi$ (sol. 1)
2140 ± 40	BAILLON	75	IPWA	$\overline{K}N \to \Lambda\pi$ (sol. 2)
$2082\pm\ 4$	COX	70	DPWA	See CORDEN 76
2070 ± 30	LITCHFIELD	70	DPWA	$K^- N \to \Lambda\pi$

$\Sigma(2080)$ WIDTH

VALUE (MeV)	DOCUMENT ID		TECN	COMMENT
186 ± 48	[1] CORDEN	76	DPWA	$K^- n \to \Lambda\pi^-$
100	DEBELLEFON	76	IPWA	$K^- p \to \Lambda\pi^0$
240 ± 50	BAILLON	75	IPWA	$\overline{K}N \to \Lambda\pi$ (sol. 1)
200 ± 50	BAILLON	75	IPWA	$\overline{K}N \to \Lambda\pi$ (sol. 2)
87 ± 20	COX	70	DPWA	See CORDEN 76
250 ± 40	LITCHFIELD	70	DPWA	$K^- N \to \Lambda\pi$

$\Sigma(2080)$ DECAY MODES

	Mode
Γ_1	$N\overline{K}$
Γ_2	$\Lambda\pi$

$\Sigma(2080)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$(\Gamma_i\Gamma_f)^{1/2}/\Gamma_{total}$ in $N\overline{K} \to \Sigma(2080) \to \Lambda\pi$ $\qquad (\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE	DOCUMENT ID		TECN	COMMENT
-0.10 ± 0.03	[1] CORDEN	76	DPWA	$K^- n \to \Lambda\pi^-$
-0.10	DEBELLEFON	76	IPWA	$K^- p \to \Lambda\pi^0$
-0.13 ± 0.04	BAILLON	75	IPWA	$\overline{K}N \to \Lambda\pi$ (sol. 1 and 2)
-0.16 ± 0.03	COX	70	DPWA	See CORDEN 76
-0.09 ± 0.03	LITCHFIELD	70	DPWA	$K^- N \to \Lambda\pi$

$\Sigma(2080)$ FOOTNOTES

[1] Preferred solution 3; see CORDEN 76 for other possibilities, including a D_{15} at this mass.

$\Sigma(2080)$ REFERENCES

CORDEN	76	NP B104 382	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
Also	75	NP B90 1	De Bellefon, Berthon, Brunet+	(CDEF, SACL) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
COX	70	NP B19 61	+Islam, Colley+	(BIRM, EDIN, GLAS, LOIC) IJP
LITCHFIELD	70	NP B22 269		(RHEL) IJP